



(86) Date de dépôt PCT/PCT Filing Date: 2017/04/27

(87) Date publication PCT/PCT Publication Date: 2017/11/30

(45) Date de délivrance/Issue Date: 2022/12/20

(85) Entrée phase nationale/National Entry: 2018/11/26

(86) N° demande PCT/PCT Application No.: US 2017/029843

(87) N° publication PCT/PCT Publication No.: 2017/204986

(30) Priorités/Priorities: 2016/05/27 (US15/166,476);
2016/08/31 (US15/253,010)

(51) Cl.Int./Int.Cl. *A01N 65/00* (2009.01)

(72) Inventeurs/Inventors:

VANGARA, KIRAN KUMAR, US;

LI, HUAGUANG, US;

YAN, NINGXIN, US;

NGUYEN, HUNG Q., US;

GOSKONDA, VENKAT R., US


(73) Propriétaire/Owner:

RADIUS PHARMACEUTICALS, INC., US

(74) Agent: BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) Titre : FORMULATIONS DE CANNABINOIDES STABLES

(54) Title: STABLE CANNABINOID FORMULATIONS



(57) Abrégé/Abstract:

The present invention is generally directed to substantially pure cannabidiol, stable cannabinoid pharmaceutical formulations, and methods of their use.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
30 November 2017 (30.11.2017)



(10) International Publication Number
WO 2017/204986 A1

(51) International Patent Classification:
A01N 65/00 (2009.01)

1333 South Spectrum Blvd., Suite 100, Chandler, AZ 85286 (US).

(21) International Application Number:
PCT/US2017/029843

(74) Agent: RAINCROW, Jeremy, D. et al.; Wood, Phillips, Katz, Clark, Mortimer, 500 West Madison Street, Suite 1130, Chicago, IL 60661 (US).

(22) International Filing Date:
27 April 2017 (27.04.2017)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
15/166,476 27 May 2016 (27.05.2016) US
15/253,010 31 August 2016 (31.08.2016) US


(71) Applicant: INSYS DEVELOPMENT COMPANY, INC.
[US/US]; 1333 South Spectrum Blvd., Suite 100, Chandler, AZ 85286 (US).

(72) Inventors: VANGARA, Kiran, Kumar; 1333 South Spectrum Blvd, Suite 100, Chandler, AZ 85286 (US). LI, Huaguang; 1333 South Spectrum Blvd., Suite 100, Chandler, AZ 85286 (US). YAN, Ningxin; 1333 South Spectrum Blvd., Suite 100, Chandler, AZ 85286 (US). NGUYEN, Hung, Q.; 1333 South Spectrum Blvd., Suite 100, Chandler, AZ 85286 (US). GOSKONDA, Venkat, R.;

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: STABLE CANNABINOID FORMULATIONS

FIG. 10



(57) Abstract: The present invention is generally directed to substantially pure cannabidiol, stable cannabinoid pharmaceutical formulations, and methods of their use.

[Continued on next page]



WO 2017/204986 A1

WO 2017/204986 A1

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

STABLE CANNABINOID FORMULATIONS

[0001]

Field of the Invention


[0002] The present invention is generally directed to substantially pure cannabidiol, stable cannabinoid pharmaceutical formulations, and methods of their use.

Background

[0003] Cannabinoids are chemicals that are produced by cannabis flowers. Cannabinoids imitate endogenous compounds in humans.

[0004] Cannabinoids include cannabinol, cannabidiol, dronabinol (delta-9-tetrahydrocannabinol), delta-8-tetrahydrocannabinol, 11-hydroxy-tetrahydrocannabinol, 11-hydroxy-delta9-tetrahydrocannabinol, levonantradol, delta-11-tetrahydrocannabinol, tetrahydrocannabivarin, amandamide, nabilone, and acids and analogs thereof. It is now possible to synthesize many cannabinoids in a laboratory thereby eliminating the need to grow cannabis for extraction of the compounds.

[0005] One cannabinoid, cannabidiol, (-)-trans-2-p-mentha-1,8-dien-3-yl-5-pentylresorcinol, is non-psychoactive and has shown promise in treating numerous diseases and disorders. Synthetic cannabidiol has the same structure as naturally occurring cannabidiol.



[0006] Commercially available cannabidiol is usually contaminated with delta 9-tetrahydrocannabinol. The presence of delta-9-tetrahydrocannabinol can be a concern because delta-9-tetrahydrocannabinol is regulated by the United States Drug Enforcement Administration as a Schedule I Drug. Having a higher Schedule number could result in easier access for patients to cannabidiol treatments. Further, delta-9-tetrahydrocannabinol is a hallucinogen and patients receiving cannabidiol wish to avoid this undesirable side effect of the delta-9-tetrahydrocannabinol contaminant. Therefore, there is a need for a substantially pure synthetically synthesized cannabidiol that does not contain delta-9-tetrahydrocannabinol.

[0007] Cannabinoids, including cannabidiol, may be suitable for the treatment of diseases or disorders, or symptoms of diseases or disorders, such as Dravet Syndrome, Lennox Gastaut Syndrome, myoclonic seizures, juvenile myoclonic epilepsy, refractory epilepsy, schizophrenia, juvenile spasms, West syndrome, refractory infantile spasms, infantile spasms, tuberous sclerosis complex, brain tumors, neuropathic pain, cannabis use disorder, post-traumatic stress disorder, anxiety, early psychosis, Alzheimer's Disease, autism, and withdrawal from opioids, cocaine, heroin, amphetamines, and nicotine.

[0008] Accordingly, there is a need for new stable cannabinoid formulations. There is also a need for substantially pure cannabidiol.

Summary of the Invention

[0009] In one aspect, the invention is directed to oral pharmaceutical formulations comprising:

cannabidiol;

a vehicle consisting of a lipid or selected from the group consisting of water, ethanol, glycerin, propylene glycol, polyethylene glycol 400 and a combination thereof,

wherein the formulation provides a greater cannabidiol C_{max} , AUC(last) and/or AUC (0-t) when administered to a subject in a fed condition than when administered to a subject in a fasted condition.

[00010] In another aspect, the cannabidiol is greater than 98% pure.

[00011] In another aspect, the vehicle is a medium chain glyceride, preferably caprylic/capric triglyceride.

[00012] In another aspect, the vehicle is sesame oil.

[00013] In another aspect, the oral pharmaceutical formulations of the present invention further comprise an antioxidant or preservative selected from the group consisting of alpha-tocopherol, ascorbyl palmitate, methyl paraben, propyl paraben and a combination thereof.

[00014] In another aspect, the oral pharmaceutical formulations of the present invention provide wherein the formulation provides a cannabidiol C_{max} of at least 10 times greater, an AUC(0-inf) of at least 3 times greater and/or an AUC(0-t) of at least 6 times greater and/or provides a 7-OH-cannabidiol C_{max} of at least 4 times greater, an AUC(0-inf) of at least 3.5 times greater and/or an AUC(0-t) of at least 3.5 times greater when administered to a subject in a fed condition than when administered to a subject in a fasted condition.

[00015] In another aspect, the invention is directed to oral pharmaceutical formulations comprising:

from about 8% to about 31% w/w cannabidiol; and

a vehicle consisting of from about 60% to about 90% w/w of a lipid or a combination of from about 40% to about 60% w/w ethanol, from about 1% to about 5% w/w polyethylene glycol, from about 5% to about 10% w/w propylene glycol and from about 20% to about 40% w/w water,

wherein the formulation provides a cannabidiol C_{max} of at least 10 times greater, an AUC(last) of at least 7 times greater and/or an AUC(0-t) of at least 6 times greater and/or provides a 7-OH-cannabidiol C_{max} of at least 4.0 times greater, an AUC(last) of at least 3.5 times greater and/or an AUC(0-t) of at least 3.5 times greater when administered to a subject in a fed condition than when administered to a subject in a fasted condition.

[00016] In another aspect, the invention is directed to oral pharmaceutical formulations comprising cannabidiol at a concentration of about 10% w/w and caprylic/capric triglyceride at a concentration of about 89% w/w.

[00017] In another aspect, the invention is directed to oral pharmaceutical formulations comprising cannabidiol at a concentration of about 31% w/w and caprylic/capric triglyceride at a concentration of about 68% w/w.

[00018] In another aspect, the invention is directed to oral pharmaceutical formulations comprising cannabidiol is at a concentration of about 11% w/w and the vehicle is sesame oil at a concentration of about 80% w/w.

[00019] In another aspect, the invention is directed to oral pharmaceutical formulations comprising cannabidiol at a concentration of about 8.8% w/w, and a combination of ethanol at a concentration of about 50% w/w, polyethylene glycol at a concentration of about 3% w/w, propylene glycol at a concentration of about 7.5% w/w and water at a concentration of about 30% w/w.

[00020] In another aspect, the invention is directed to methods of using formulations of the invention to treat or prevent diseases or disorders, or treat symptoms of diseases or disorder selected from the group consisting of Prader-Willi syndrome, obesity, graft versus host disease, gelastic seizures/hypothalamic hamartoma, neonatal seizures, movement disorders including dystonia, central pain syndromes, phantom limb pain, multiple sclerosis, traumatic brain injury, radiation therapy, acute and chronic graft versus host disease, T-cell autoimmune disorders, colitis, Dravet Syndrome, Lennox Gastaut Syndrome, myoclonic seizures, juvenile myoclonic epilepsy, refractory epilepsy, schizophrenia, juvenile spasms, West syndrome, infantile spasms, refractory infantile spasms, tuberous sclerosis complex, brain tumors, neuropathic pain, cannabis use disorder, post-traumatic stress disorder, anxiety, early psychosis, Alzheimer's Disease, autism, acne, Parkinson's disease, social anxiety disorder, depression, diabetic retinopathy, diabetic

nephropathy and diabetic neuropathy, ischemic injury of heart, ischemic injury of brain, chronic pain syndrome, rheumatoid arthritis.

[00021] In another aspect, compositions of the present invention are administered to a patient that is in a fed or fasted condition.

[00022] In another aspect, the invention is directed to methods of using formulations of the invention for assisting with withdrawal from opioids, cocaine, heroin, amphetamines and nicotine; and as an analgesic or to assist with handling of adverse emotional stimuli.

Brief Description of the Figures

[00023] Fig. 1 shows the results from the study detailed in Example 7 and illustrates the advantages of administration of substantially pure, synthetically synthesized, cannabidiol formulations for treatment of neuropathic pain.

[00024] Fig. 2 shows the results from the study detailed in Example 8 and illustrates the advantages of administration of substantially pure, synthetically synthesized, cannabidiol formulations over THC formulations for treatment of neuropathic pain.

[00025] Fig. 3 shows further results from the study detailed in Example 8 and illustrates dose-dependent advantages of administration of substantially pure, synthetically synthesized, cannabidiol formulations over THC formulations for treatment of neuropathic pain.

[00026] Fig. 4 shows further results from the study detailed in Example 8 and illustrates synergistic results of administration of substantially pure, synthetically synthesized, cannabidiol formulations and THC formulations for treatment of neuropathic pain.

[00027] Fig. 5 shows the results from the study detailed in Example 9 and illustrates the advantages administration of higher ratio substantially pure, synthetically synthesized, cannabidiol to THC formulations for treatment of neuropathic pain.

[00028] Fig. 6 shows the results from the study detailed in Example 10 and illustrates the advantages of administration of substantially pure, synthetically synthesized, cannabidiol formulations for treatment of neuropathic pain.

[00029] Fig. 7 shows the results from the study detailed in Example 14 and illustrates the advantages of administration of substantially pure, synthetically synthesized, cannabidiol formulations for treatment of glioblastoma multiforme.

[00030] Fig. 8 shows the results from the study detailed in Example 15 and illustrates the advantages of administration of substantially pure, synthetically synthesized, cannabidiol formulations for treatment of glioblastoma multiforme.

[00031] Fig. 9 shows the results from the study detailed in Example 17 and illustrates the advantages of administration of oral cannabidiol solutions to subjects in the fed condition. Panel A. shows mean cannabidiol concentrations versus time. Panel B. shows the natural log of the mean cannabidiol concentrations versus time.

[00032] Fig. 10 shows the results from the study detailed in Example 17 and illustrates the advantages of administration of oral cannabidiol solutions to subjects in the fed condition. Panel A. shows mean 7-OH-cannabidiol concentrations versus time. Panel B. shows the natural log of the mean 7-OH-cannabidiol concentrations versus time.

Detailed Description

[00033] As indicated above, Applicant created stable formulations with and without alcohol (see Examples 1 and 3). The formulations that do not contain alcohol are especially suitable for administration to children. Further, the alcohol-free formulations are especially suitable for patients in recovery from drug and alcohol addiction.

[00034] In addition, Applicant created stable lipid formulations (see Example 5). These formulations were also unexpectedly stable during storage (see Example 6).

[00035] Further, Applicant unexpectedly found that substantially pure cannabidiol formulations are especially suitable for treatment of neuropathic pain (see Examples 7-10 and Figs. 1-6), epilepsy (see Examples 11-13), and glioblastoma multiforme (see Examples 14 and 15 and Figs. 7 and 8).

Alcohol-Free Formulations

[00036] In one embodiment, the present invention is directed to stable pharmaceutical formulation for oral administration comprising from about 0.1 to about

50 % of a cannabinoid, from about 0.1 to about 40 % of a polyethylene glycol, from about 0.1 to about 50 % of propylene glycol, and from about 0.1 to about 20 % of water, wherein the formulation does not contain alcohol and the formulation has a pH of from about 5 to about 8.

[00037] In a preferred embodiment, the formulations contain from about 1 to about 40 % of a cannabinoid. In more preferred embodiments, the formulations contain from about 5 to about 35 %, from about 20 to about 35 % or from about 30 to 35 % of a cannabinoid.

[00038] In yet another embodiment, the formulations contain a cannabinoid selected from group consisting of cannabinol, cannabidiol, dronabinol (delta-9-tetrahydrocannabinol), delta-8-tetrahydrocannabinol, 11-hydroxy-tetrahydrocannabinol, 11-hydroxy-delta-9-tetrahydrocannabinol, levonantradol, delta-11-tetrahydrocannabinol, tetrahydrocannabivarin, amandamide, nabilone, acids, analogs, and synthetic derivatives thereof. In a preferred embodiment, the cannabinoid is cannabidiol.

[00039] In a preferred embodiment, the formulations contain from about 1 to about 40 % of a cannabidiol. In more preferred embodiments, the formulations contain from about 5 to about 35 %, from about 20 to about 35 % or from about 30 to 35 % of a cannabidiol.

[00040] In yet another embodiment, the formulations contain cannabidiol that is substantially pure and synthetically synthesized which has a purity of greater than 98 %. In a more preferred embodiment, the cannabidiol is greater than 99 % pure. In an even more preferred embodiment, the cannabidiol is greater than 99.5 % pure. In a most preferred embodiment, the cannabidiol formulation contains less than 0.3 % delta-9-tetrahydrocannabinol.

[00041] In another embodiment, the formulations contain from about 0.001 to about 1 % of an antioxidant. In a preferred embodiment, the formulations contain from about 0.01 to about 1 % antioxidant. In a more preferred embodiment, the formulations contain from about 0.02 to about 0.5 % antioxidant.

[00042] Suitable antioxidants include butylated hydroxytoluene ("BHT"), butylated hydroxyl anisole ("BHA"), alpha-tocopherol (Vitamin E), ascorbyl palmitate, ascorbic acid, sodium ascorbate, ethylenediamino tetraacetic acid, cysteine hydrochloride, citric acid, sodium citrate, sodium bisulfate, sodium metabisulfite, lecithin, propyl gallate, sodium sulfate, monothioglycerol tert-butylhydroquinone ("TBHQ") and combinations thereof. In a preferred embodiment, the formulations contain alpha-tocopherol (Vitamin E), ascorbic acid, sodium ascorbate, ascorbyl palmitate or combinations thereof.

[00043] In another embodiment, the formulations contain from about 1 to about 40 % of a polyethylene glycol. In a preferred embodiment, the formulations contain from about 1 to about 35 %, from about 5 to about 35 %, from about 20 to about 30 %, or from about 25 to about 30 % polyethylene glycol.

[00044] Suitable polyethylene glycols include low molecular weight polyethylene glycols with an average molecular weight of between 200 and 10,000. One preferred polyethylene glycol that can be used is polyethylene glycol 400.

[00045] In another embodiment, the formulations contain from about 1 to about 40 % of polyethylene glycol 400. In a preferred embodiment, the formulations contain from about 1 to about 35 %, from about 5 to about 35 %, from about 20 to about 30 %, or from about 25 to about 30 % polyethylene glycol 400.

[00046] In another embodiment, the formulations contain from about 1 to about 50 % of propylene glycol. In a preferred embodiment, the formulations contain from about 1 to about 40 %, from about 5 to about 35 %, from about 20 to about 35 %, or from about 30 to about 35 % propylene glycol.

[00047] In a further embodiment, the formulations contain water. The formulations can contain 0 % water. If the formulations contain water, they can include from about 1 to about 15 % water, from about 1 to about 10 % water, or from about 4 to about 8 % water.

[00048] The pH of the formulations may be modified using any pharmaceutically acceptable means. Preferably the pH of the formulation is from about 5 to about 8. In a more preferred embodiment, the pH of the formulations is from about 6 to about 7. In a most preferred embodiment, the pH of the formulations is from about 6.2 to about 6.7.

[00049] The formulations of the present invention may also contain sweeteners, sweetener enhancers, preservatives, pH modifiers, and flavoring agents.

[00050] Suitable sweeteners include, but are not limited to, sucralose, sucrose, aspartame, saccharin, dextrose, mannitol, xylitol, and combinations thereof.

[00051] If the formulations contain a sweetener, the formulations preferably contain from about 0.001 to about 1 % sweetener.

[00052] If the formulations contain a sweetness enhancer, the formulations preferably contain from about 0.001 to about 1% sweetness enhancer.

[00053] Suitable sweetness enhancers include, but are not limited to, the ammonium salt forms of crude and refined Glycyrrhizic Acid. Magnasweet® products (available from Mafco Worldwide Corporation, Magnasweet is a registered trademark of Mafco Worldwide Corporation) use the ammonium salt forms of crude and refined Glycyrrhizic Acid. Glycyrrhizic Acid is also available as a pure derivative in the sodium and potassium salt forms.

[00054] Suitable pH modifiers include, but are not limited to, hydrochloric acid, ascorbic acid, citric acid, sodium citrate, fumaric acid, sodium hydroxide, sodium bicarbonate, sodium carbonate, ammonium carbonate, and combinations thereof.

[00055] Suitable preservatives include, but are not limited to, methyl paraben, propyl paraben, benzyl alcohol, benzoic acid, sodium benzoate, sorbic acid, and combinations thereof.

[00056] Suitable flavoring agents include, but are not limited to, raspberry, peppermint oil, grape flavor, menthol, spearmint oil, citrus oil, cinnamon oil, strawberry flavor, cherry flavor, raspberry flavor, orange oil, lemon oil, lemon mint flavor, fruit punch flavor, and combinations thereof. In a preferred embodiment, the formulations contain strawberry flavor.

[00057] If the formulations contain a flavoring agent, the formulations preferably contain from about 0.001 to about 1 % flavoring agent. In a more preferred embodiment, the formulations contain from about 0.005 to about 0.5 % of the flavoring agent.

[00058] The formulations are suitable for oral, buccal, sublingual, inhalation or intravenous/intramuscular administration. Preferably, the formulations are liquids administered orally. More preferably, the formulations are simple solutions administered orally.

Formulations Containing Alcohol

[00059] In another embodiment, the invention is directed to stable pharmaceutical formulation for oral administration comprising from about 0.1 to about 40 % of a cannabinoid, from about 0.1 to about 25 % of a polyethylene glycol, from about 0.1 to about 40 % of propylene glycol, optionally from about 0.1 to about 50 % of water, and from about 0.1 to about 70 % of alcohol, wherein the formulation has a pH of from about 5 to about 8.

[00060] In a preferred embodiment, the formulations contain from about 1 to about 35 % of a cannabinoid. In a more preferred embodiment, the formulations contain from about 1 to about 15 %, from about 5 to about 12 % or from about 7 to about 11 % cannabinoid. Alternatively, the formulations may contain from about 20 to about 35 % or from about 30 to about 35 % cannabinoid.

[00061] In yet another embodiment, the formulations contain a cannabinoid selected from group consisting of cannabinol, cannabidiol, dronabinol (delta-9-tetrahydrocannabinol), delta-8-tetrahydrocannabinol, 11-hydroxy-tetrahydrocannabinol, 11-hydroxy-delta-9-tetrahydrocannabinol, levonantradol, delta-11-tetrahydrocannabinol, tetrahydrocannabivarin, amandamide, nabilone, acids, analogs, and synthetic derivatives thereof. In a preferred embodiment, the cannabinoid is cannabidiol.

[00062] In a preferred embodiment, the formulations contain from about 1 to about 35 % of a cannabidiol. In a more preferred embodiment, the formulations contain from about 1 to about 15 %, from about 5 to about 12 % or from about 7 to about 11 % cannabidiol. Alternatively, the formulations may contain from about 20 to about 35 % or from about 30 to about 35 % cannabidiol.

[00063] In yet another embodiment, the formulations contain cannabidiol that is substantially pure and synthetically synthesized which has a purity of greater than 98 %.

In a more preferred embodiment, the cannabidiol is greater than 99 % pure. In an even more preferred embodiment, the cannabidiol is greater than 99.5 % pure. In a most preferred embodiment, the cannabidiol formulation contains less than 0.3 % delta-9-tetrahydrocannabinol.

[00064] In another embodiment, the formulations contain from about 0.001 to about 1 % of an antioxidant. In a preferred embodiment, the formulations contain from about 0.01 to about 1 % antioxidant. In a more preferred embodiment, the formulations contain from about 0.02 to about 0.5 % antioxidant.

[00065] Suitable antioxidants include butylated hydroxytoluene ("BHT"), butylated hydroxyl anisole ("BHA"), alpha-tocopherol (Vitamin E), ascorbyl palmitate, ascorbic acid, sodium ascorbate, ethylenediamino tetraacetic acid, cysteine hydrochloride, citric acid, sodium citrate, sodium bisulfate, sodium metabisulfite, lecithin, propyl gallate, sodium sulfate, tert-butylhydroquinone ("TBHQ") and combinations thereof. In a preferred embodiment, the formulations contain alpha-tocopherol (Vitamin E), ascorbyl palmitate, or combinations thereof.

[00066] In another embodiment, the formulations contain from about 1 to about 20 % of propylene glycol. In a preferred embodiment, the formulations contain from about 1 to about 15 % or from about 5 to about 10 % propylene glycol.

[00067] In an alternative embodiment, the formulations contain from about 20 to about 50 % of propylene glycol. In a preferred embodiment, the formulations contain from about 30 to about 40 % or from about 35 to about 40 % propylene glycol.

[00068] In another embodiment, the formulations contain from about 1 to about 20 % of a polyethylene glycol. In a preferred embodiment, the formulations contain from about 1 to about 10 % or from about 1 to about 5 % polyethylene glycol.

[00069] In an alternative embodiment, the formulations contain from about 10 to about 30 % of a polyethylene glycol. In a preferred alternative embodiment, the formulations contain from about 15 to about 25 % polyethylene glycol.

[00070] Suitable polyethylene glycols include low molecular weight polyethylene glycols with an average molecular weight of between 200 and 10,000. One preferred polyethylene glycol that can be used is polyethylene glycol 400.

[00071] In another embodiment, the formulations contain from about 1 to about 20 % of polyethylene glycol 400. In a preferred embodiment, the formulations contain from about 1 to about 10 % or from about 1 to about 5 % polyethylene glycol 400.

[00072] In an alternative embodiment, the formulations contain from about 1 to about 5 % of polyethylene glycol 400. In a preferred alternative embodiment, the formulations contain from about 15 to about 25 % polyethylene glycol 400.

[00073] In a further embodiment, the formulations contain water. The formulations can contain 0 % water. If the formulations contain water, they can include from about 1 to about 40 % water, from about 5 to about 40 % water, from about 10 to about 35 % water or from about 25 to about 35 % water.

[00074] In yet another embodiment, the formulations contain from about 1 to about 65 % alcohol. In a preferred embodiment, the formulations contain from about 10 to about 65 %, from about 15 to about 60 %, or from about 30 to 55 % alcohol.

[00075] In an alternative embodiment, the formulations contain from about 1 to about 20 % alcohol. In a preferred alternative embodiment, the formulations contain from about 1 to about 10 % or from about 3 to about 7 % alcohol.

[00076] The pH of the formulations may be modified using any pharmaceutically acceptable means. Preferably the pH of the formulations is from about 6 to about 7. In a more preferred embodiment, the pH of the formulations is from about 6.2 to about 6.7.

[00077] The formulations of the present invention may also contain sweeteners, sweetener enhancers, pH modifiers, preservatives, and flavoring agents.

[00078] Suitable sweeteners include, but are not limited to, sucralose, sucrose, aspartame, saccharin, dextrose, mannitol, xylitol, and combinations thereof.

[00079] If the formulations contain a sweetener, the formulations preferably contain from about 0.001 to about 1 % sweetener.

[00080] Suitable sweetness enhancers include, but are not limited to, the ammonium salt forms of crude and refined Glycyrrhizic Acid. Magnasweet® products (available from Mafco Worldwide Corporation, Magnasweet is a registered trademark of Mafco Worldwide Corporation) use the ammonium salt forms of crude and refined Glycyrrhizic Acid. Glycyrrhizic Acid is also available as a pure derivative in the sodium and potassium salt forms.

[00081] If the formulations contain a sweetness enhancer, the formulations preferably contain from about 0.001 to about 1 % sweetness enhancer.

[00082] Suitable pH modifiers include, but are not limited to, hydrochloric acid, ascorbic acid, citric acid, sodium citrate, fumaric acid, sodium hydroxide, sodium bicarbonate, sodium carbonate, ammonium carbonate, and combinations thereof.

[00083] Suitable preservatives include, but are not limited to, methyl paraben, propyl paraben, benzyl alcohol, benzoic acid, sodium benzoate, sorbic acid, and combinations thereof.

[00084] Suitable flavoring agents include, but are not limited to, raspberry, peppermint oil, grape flavor, menthol, spearmint oil, citrus oil, cinnamon oil, strawberry flavor, cherry flavor, raspberry flavor, orange oil, lemon oil, lemon mint flavor, fruit punch flavor, and combinations thereof. In a preferred embodiment, the formulations contain fruit punch flavor, raspberry flavor, grape flavor, or lemon mint flavor.

[00085] If the formulations contain a flavoring agent, the formulations preferably contain from about 0.001 to about 1 % flavoring agent. In a more preferred embodiment, the formulations contain from about 0.005 to about 0.5 % of the flavoring agent.

[00086] The formulations are suitable for oral, buccal, sublingual, inhalation or intravenous/intramuscular administration. Preferably, the formulations are liquids administered orally. More preferably, the formulations are simple solutions administered orally.

Formulations Containing Lipids

[00087] In another embodiment, the invention is directed to stable pharmaceutical formulation for oral administration comprising from about 0.1 to about 40 % of a cannabinoid and from about 10 to about 95 % of a lipid.

[00088] In a preferred embodiment, the lipid is selected from the group consisting of sesame oil, olive oil, corn oil, sunflower oil, safflower oil, flaxseed oil, almond oil, peanut oil, walnut oil, cashew oil, castor oil, coconut oil, palm oil, soybean oil, canola oil, vegetable oil, rice bran oil, fatty acids including caproic acid, enanthic acid, caprylic acid, pelargonic acid, capric acid, undecylenic acid, lauric acid, myristic acid, pentadecylic acid, palmitic acid, margaric acid, oleic acid, stearic acid, nonadecylic acid, linoleic acid, arachidic acid and arachidonic acid, medium chain glycerides, decanoyl glycerides, octanoyl glycerides, caprylic/capric triglyceride, caprylic/capric/linoleic triglyceride, oleoyl polyoxyl-6 glycerides, linoleoyl polyoxyl-6 glycerides, polyglyceryl-3 dioleate, glyceryl monolinoleate, glyceryl monocaprylate, oleic acid, and a combination thereof. In a more preferred embodiment, the lipid is a medium-chain triglyceride whose fatty acids have an aliphatic tail of from 6 to 12 carbon atoms. In a most preferred embodiment, the lipid is caprylic/capric triglyceride.

[00089] Suitable commercial sources for the lipid include Miglyol[®] 812N (caprylic/capric triglyceride) containing a proprietary mixture of decanoyl and octanoyl glycerides (fatty acid esters) (Miglyol is available from and a registered trademark of Cremer Oleo GmbH & Co.) and Miglyol[®] 840 (caprylic/capric/linoleic triglyceride) containing a proprietary mixture of propylene glycol dicaprylate/dicaprate and otherwise known as decanoic acid/octanoic acid/propane-1,2-diol.

[00090] In yet another embodiment, the formulations contain a cannabinoid selected from group consisting of cannabinol, cannabidiol, dronabinol (delta-9-tetrahydrocannabinol), delta-8-tetrahydrocannabinol, 11-hydroxy-tetrahydrocannabinol, 11-hydroxy-delta-9-tetrahydrocannabinol, levonantradol, delta-11-tetrahydrocannabinol, tetrahydrocannabivarin, amandamide, nabilone, acids, analogs, and synthetic derivatives thereof. In a preferred embodiment, the cannabinoid is cannabidiol.

[00091] In yet another embodiment, the formulations contain cannabidiol that is substantially pure and synthetically synthesized which has a purity of greater than 98 %.

In a more preferred embodiment, the cannabidiol is greater than 99 % pure. In an even more preferred embodiment, the cannabidiol is greater than 99.5 % pure. In a most preferred embodiment, the cannabidiol formulation contains less than 0.3 % delta-9-tetrahydrocannabinol.

[00092] In a preferred embodiment, the formulations contain from about 1 to about 35 % of a cannabidiol. In a more preferred embodiment, the formulations contain from about 10 to about 32 % cannabidiol. In a most preferred embodiment, the formulations contain about 31.09% cannabidiol.

[00093] In a preferred embodiment, the formulations contain from about 20 to about 90 % of lipids. In a more preferred embodiment, the formulations contain from about 50 to about 90 % lipids. In a most preferred embodiment, the formulations contain from about 50 to about 74 % lipids.

[00094] In yet another embodiment, the formulations contain alcohol. The formulations can contain 0 % alcohol. If the formulations contain alcohol, they can include from about 0.1 to about 20 % alcohol. In a preferred embodiment, the formulations contain from about 1 to about 15 % alcohol. In a more preferred embodiment, the formulations contain from about 1 to about 10 % alcohol.

[00095] In another embodiment, the formulations contain an antioxidant. The formulations can contain 0 % antioxidant. If the formulations contain antioxidant, they can include from about 0.01 to about 1 % of an antioxidant. In a preferred embodiment, the formulations contain from about 0.02 to about 0.5 % antioxidant.

[00096] Suitable antioxidants include butylated hydroxytoluene, butylated hydroxyl anisole, alpha-tocopherol (Vitamin E), ascorbyl palmitate, ascorbic acid, sodium ascorbate, ethylenediamino tetraacetic acid, cysteine hydrochloride, citric acid, sodium citrate, sodium bisulfate, sodium metabisulfite, lecithin, propyl gallate, sodium sulfate, TBHQ, and combinations thereof. In a preferred embodiment, the formulations contain alpha-tocopherol (Vitamin E), ascorbyl palmitate or combinations thereof.

[00097] Suitable sweeteners include, but are not limited to, sucralose, sucrose, aspartame, saccharin, dextrose, mannitol, xylitol, and combinations thereof. In a preferred embodiment the sweetener is saccharin.

[00098] If the formulations contain a sweetener, the formulations preferably contain from about 0.01 to about 2 % sweetener. In a more preferred embodiment, the formulations contain from about 0.01 to about 0.8 % sweetener. In a most preferred embodiment, the formulations contain from about 0.02 to about 0.05 % sweetener.

[00099] Suitable sweetness enhancers include, but are not limited to, the ammonium salt forms of crude and refined Glycyrrhizic Acid. Magnasweet® products (available from Mafco Worldwide Corporation, Magnasweet is a registered trademark of Mafco Worldwide Corporation) use the ammonium salt forms of crude and refined Glycyrrhizic Acid. Glycyrrhizic Acid is also available as a pure derivative in the sodium and potassium salt forms.

[000100] If the formulations contain a sweetness enhancer, the formulations preferably contain from about 0.001 to about 1 % sweetness enhancer.

[000101] Suitable pH modifiers include, but are not limited to, hydrochloric acid, ascorbic acid, citric acid, sodium citrate, fumaric acid, sodium hydroxide, sodium bicarbonate, sodium carbonate, ammonium carbonate, and combinations thereof.

[000102] Suitable preservatives include, but are not limited to, methyl paraben, propyl paraben, benzyl alcohol, benzoic acid, sodium benzoate, sorbic acid, and combinations thereof.

[000103] Suitable flavoring agents include, but are not limited to, raspberry, peppermint oil, grape flavor, menthol, spearmint oil, citrus oil, cinnamon oil, strawberry flavor, cherry flavor, raspberry flavor, orange oil, lemon oil, lemon mint flavor, fruit punch flavor, and combinations thereof. In a preferred embodiment the flavoring agent is strawberry flavor.

[000104] If the formulations contain a flavoring agent, the formulations preferably contain from about 0.01 to about 1 % flavoring agent. In a more preferred embodiment, the formulations contain from about 0.005 to about 0.5 % of the flavoring agent.

[000105] The formulations are suitable for oral, buccal, sublingual, inhalation or intravenous/intramuscular administration. Preferably, the formulations are liquids administered orally.

Exemplary Uses of Formulations of the Present Invention (Alcohol-Containing, Alcohol-Free, and Lipid) and Synthetically Synthesized, Substantially Pure, Cannabidiol


[000106] The formulations of the present invention are especially suitable for treatment of many diseases or disorders or symptoms of diseases and disorders. Further, cannabidiol which is synthetically synthesized and substantially pure will be even more effective and suitable for the treatment of diseases or symptoms of these diseases.

[000107] The formulations of the present invention may be administered to a patient in a fed condition. As used herein a “fed condition” refers to a patient that consumes food prior to administration of a formulation of the present invention and from which the food has not been cleared from the gastrointestinal tract prior to the administration.

[000108] Disease and disorders or symptoms of these disease or disorders that can be treated or prevented by formulations of the present invention include, but are not limited to, Prader-Willi syndrome, obesity, graft versus host disease, gelastic seizures/hypothalamic hamartoma, neonatal seizures, movement disorders including dystonia, central pain syndromes including but not limited to complex regional pain syndrome, phantom limb pain, multiple sclerosis, traumatic brain injury, radiation therapy, acute and chronic graft versus host disease, T-cell autoimmune disorders, colitis, Dravet Syndrome, Lennox Gastaut Syndrome, myoclonic seizures, juvenile myoclonic epilepsy, refractory epilepsy, schizophrenia, juvenile spasms, West syndrome, infantile spasms, refractory infantile spasms, tuberous sclerosis complex, brain tumors, neuropathic pain, cannabis use disorder, post-traumatic stress disorder, anxiety, early psychosis, Alzheimer’s Disease, autism, acne, Parkinson’s disease, social anxiety disorder, depression, diabetic retinopathy, diabetic nephropathy, diabetic neuropathy, ischemic injury of heart, ischemic injury of brain, chronic pain syndrome, rheumatoid arthritis, patients encountering adverse emotional stimuli, nausea and addiction disorders related to drugs of abuse such as opioid, heroin, cocaine, amphetamine dependence and including acute and long-term treatment of dependence and relapse associated with drugs of abuse.

[000109] As first explained in US Patent Application No. 62/004,495, Applicant unexpectedly created a new synthetic pathway for creating cannabidiol. This new process eliminated the need to grow cannabis in order to extract cannabidiol. Applicant's cannabidiol has a high purity level and is substantially free of Schedule I drugs, including delta-9-tetrahydrocannabinol.

[000110] Applicant chemically synthesized cannabidiol by combining p-menthadienol and olivetol in toluene or dichloromethane or hexane with a p-toluene sulfonic acid catalyst to produce cannabidiol (see diagram below).



[000111] In an embodiment, the present invention is directed to methods for treating Prader-Willi syndrome comprising administering the formulations of the present invention to a patient in need thereof.

[000112] In another embodiment, the present invention is directed to methods for treating a Prader-Willi syndrome comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000113] In an embodiment, the present invention is directed to methods for treating comprising administering the formulations of the present invention to a patient in need thereof.

[000114] In another embodiment, the present invention is directed to methods for treating obesity comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

- [000115] In an embodiment, the present invention is directed to methods for treating graft versus host disease comprising administering the formulations of the present invention to a patient in need thereof.
- [000116] In another embodiment, the present invention is directed to methods for treating graft versus host disease comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.
- [000117] In an embodiment, the present invention is directed to methods for preventing or treating acute and chronic graft versus host disease comprising administering the formulations of the present invention to a patient in need thereof.
- [000118] In another embodiment, the present invention is directed to methods for preventing or treating acute and chronic graft versus host disease comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.
- [000119] In an embodiment, the present invention is directed to methods for treating gelastic seizures/hypothalamic hamartoma comprising administering the formulations of the present invention to a patient in need thereof.
- [000120] In another embodiment, the present invention is directed to methods for treating gelastic seizures/hypothalamic hamartoma comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.
- [000121] In an embodiment, the present invention is directed to methods for treating neonatal seizures comprising administering the formulations of the present invention to a patient in need thereof.
- [000122] In another embodiment, the present invention is directed to methods for treating neonatal seizures comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.
- [000123] In another embodiment, the present invention is directed to methods for treating movement disorders comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof, preferably the movement disorder is dystonia.

[000124] In an embodiment, the present invention is directed to methods for treating movement disorders comprising administering the formulations of the present invention to a patient in need thereof, preferably the movement disorder is dystonia.

[000125] In another embodiment, the present invention is directed to methods for treating central pain syndromes comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof, preferably the central pain syndrome is complex regional pain syndrome.

[000126] In an embodiment, the present invention is directed to methods for treating central pain syndromes comprising administering the formulations of the present invention to a patient in need thereof, preferably the central pain syndrome is complex regional pain syndrome.

[000127] In an embodiment, the present invention is directed to methods for treating phantom limb pain comprising administering the formulations of the present invention to a patient in need thereof.

[000128] In another embodiment, the present invention is directed to methods for treating phantom limb pain comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000129] In an embodiment, the present invention is directed to methods for providing neuroprotection after stroke comprising administering the formulations of the present invention to a patient in need thereof.

[000130] In another embodiment, the present invention is directed to methods for providing neuroprotection after stroke comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000131] In an embodiment, the present invention is directed to methods for treating traumatic brain injury comprising administering the formulations of the present invention to a patient in need thereof.

[000132] In another embodiment, the present invention is directed to methods for treating traumatic brain injury comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000133] In an embodiment, the present invention is directed to methods for treating brain injury due to radiation therapy comprising administering the formulations of the present invention to a patient in need thereof.

[000134] In another embodiment, the present invention is directed to methods for treating brain injury due to radiation therapy comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000135] In an embodiment, the present invention is directed to methods for providing enhancement of neural repair following traumatic brain injury, concussion, cerebral infarction, brain irradiation or encephalomyelitis comprising administering the formulations of the present invention to a patient in need thereof.

[000136] In another embodiment, the present invention is directed to methods for providing enhancement of neural repair following traumatic brain injury, concussion, cerebral infarction, brain irradiation or encephalomyelitis comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000137] In an embodiment, the present invention is directed to methods for providing recovery from myocardial infarction comprising administering the formulations of the present invention to a patient in need thereof.

[000138] In another embodiment, the present invention is directed to methods for providing recovery from myocardial infarction comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000139] In an embodiment, the present invention is directed to methods for providing recovery from radiation injury comprising administering the formulations of the present invention to a patient in need thereof, preferably radiation injury to lung, bowel, kidney, and heart.

[000140] In another embodiment, the present invention is directed to methods for providing recovery from radiation injury comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof, preferably radiation injury to lung, bowel, kidney, and heart.

[000141] In an embodiment, the present invention is directed to methods for treating a brain tumor comprising administering the formulations of the present invention to a patient in need thereof.

[000142] In another embodiment, the present invention is directed to methods for treating a brain tumor comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000143] In an embodiment, the present invention is directed to methods for treating T-cell autoimmune disorders comprising administering the formulations of the present invention to a patient in need thereof.

[000144] In another embodiment, the present invention is directed to methods for treating T-cell autoimmune disorders comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000145] In an embodiment, the present invention is directed to methods for treating colitis comprising administering the formulations of the present invention to a patient in need thereof.

[000146] In another embodiment, the present invention is directed to methods for treating colitis comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000147] In an embodiment, the present invention is directed to methods for treating glioma comprising administering the formulations of the present invention to a patient in need thereof.

[000148] In another embodiment, the present invention is directed to methods for treating glioma comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000149] In an embodiment, the present invention is directed to methods for treating glioblastoma multiforme comprising administering the formulations of the present invention to a patient in need thereof.

[000150] In another embodiment, the present invention is directed to methods for treating glioblastoma multiforme comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000151] In an embodiment, the present invention is directed to methods for treating Dravet Syndrome comprising administering the formulations of the present invention to a patient in need thereof.

[000152] In another embodiment, the present invention is directed to methods for treating Dravet Syndrome comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000153] In yet another embodiment, the present invention is directed to methods for treating Lennox Gastaut Syndrome comprising administering the formulations of the present invention to a patient in need thereof.

[000154] In another embodiment, the present invention is directed to methods for treating Lennox Gastaut Syndrome comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000155] In a further embodiment, the present invention is directed to methods for treating Mycolonic Seizures comprising administering the formulations of the present invention to a patient in need thereof. In a more preferred embodiment, the alcohol-free formulations contain substantially pure cannabidiol.

[000156] In another embodiment, the present invention is directed to methods for treating Mycolonic Seizures comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000157] In a further embodiment, the present invention is directed to methods for treating Juvenile Mycolonic Epilepsy comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000158] In another embodiment, the present invention is directed to methods for treating Juvenile Myoclonic Epilepsy comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000159] In an embodiment, the present invention is directed to methods for treating Refractory Epilepsy comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000160] In another embodiment, the present invention is directed to methods for treating Refractory Epilepsy comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000161] In an embodiment, the present invention is directed to methods for treating juvenile spasms comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000162] In another embodiment, the present invention is directed to methods for treating juvenile spasms comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000163] In an embodiment, the present invention is directed to methods for treating West Syndrome comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000164] In another embodiment, the present invention is directed to methods for treating West Syndrome comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000165] In an embodiment, the present invention is directed to methods for treating infantile spasms comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000166] In another embodiment, the present invention is directed to methods for treating infantile spasms comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000167] In an embodiment, the present invention is directed to methods for treating refractory infantile spasms comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000168] In another embodiment, the present invention is directed to methods for treating refractory infantile spasms comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000169] In an embodiment, the present invention is directed to methods for treating tuberous sclerosis complex comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to young patients in need of treatment.

[000170] In another embodiment, the present invention is directed to methods for treating tuberous sclerosis complex comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000171] In a further embodiment, the present invention is directed to methods for treating neuropathic pain comprising administering the formulations of the present invention to a patient in need thereof. In a further embodiment, the neuropathic pain is caused by neurotoxic chemotherapy agents such as Paclitaxel, Docetaxel, Cisplatin, Oxaliplatin, Carboplatin, Vincristine, Methotrexate, Cytarabine, Fluorouracil, Ifosfamide, Cyclophosphamide, Procarbazine, etoposide, Carmustine, and Lomustine. In yet another embodiment, the neuropathic pain is caused by Paclitaxel and the patient is receiving Paclitaxel due to a diagnosis of breast, cervical, endometrial and/or ovarian cancer. In a further embodiment, the breast, cervical, endometrial and/or ovarian cancer is platinum-resistant. In another embodiment, the breast, cervical, endometrial and/or ovarian cancer is recurrent.

[000172] In another embodiment, the present invention is directed to methods for treating neuropathic pain comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof. In a further embodiment, the neuropathic pain is caused by neurotoxic chemotherapy agents such as Paclitaxel, Docetaxel, Cisplatin, Oxaliplatin, Carboplatin, Vincristine, Methotrexate, Cytarabine, Fluorouracil, Ifosfamide, Cyclophosphamide, Procarbazine, etoposide, Carmustine, and Lomustine. In yet another embodiment, the neuropathic pain is caused by Paclitaxel and the patient is receiving Paclitaxel due to a diagnosis of breast, cervical, endometrial and/or ovarian cancer. In a further embodiment, the breast, cervical, endometrial and/or ovarian cancer is platinum-resistant. In another embodiment, the breast, cervical, endometrial and/or ovarian cancer is recurrent.

[000173] In a further embodiment, the present invention is directed to methods for using cannabidiol as an analgesic comprising administering the formulations of the present invention to a patient in need thereof.

[000174] In another embodiment, the present invention is directed to methods for using cannabidiol as an analgesic comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000175] In a further embodiment, the present invention is directed to methods for treating opioid addiction withdrawal comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000176] In another embodiment, the present invention is directed to methods for treating opioid addiction withdrawal comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000177] In yet another embodiment, the present invention is directed to methods for treating cocaine addiction withdrawal comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000178] In another embodiment, the present invention is directed to methods for treating cocaine addiction withdrawal comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000179] In a further embodiment, the present invention is directed to methods for treating heroin addiction withdrawal comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000180] In another embodiment, the present invention is directed to methods for treating heroin addiction withdrawal comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000181] In a further embodiment, the present invention is directed to methods for treating nicotine addiction withdrawal comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000182] In another embodiment, the present invention is directed to methods for treating nicotine addiction withdrawal comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000183] In a further embodiment, the present invention is directed to methods for treating amphetamine addiction withdrawal comprising administering the formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000184] In another embodiment, the present invention is directed to methods for treating amphetamine addiction withdrawal comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000185] In another embodiment, the present invention is directed to methods for treating drug dependence, wherein treatment is selected from acute and long-term.

[000186] In another embodiment, the present invention is directed to methods for treating relapse associated with drug abuse.

[000187] In an embodiment, the present invention is directed to methods for treating acne comprising administering the formulations of the present invention to a patient in need thereof.

[000188] In another embodiment, the present invention is directed to methods for treating acne comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000189] In an embodiment, the present invention is directed to methods for treating Parkinson's disease comprising administering the formulations of the present invention to a patient in need thereof.

[000190] In another embodiment, the present invention is directed to methods for treating Parkinson's disease comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000191] In an embodiment, the present invention is directed to methods for treating schizophrenia comprising administering the formulations of the present invention to a patient in need thereof.

[000192] In another embodiment, the present invention is directed to methods for treating schizophrenia comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000193] In an embodiment, the present invention is directed to methods for treating social anxiety disorder comprising administering the formulations of the present invention to a patient in need thereof.

[000194] In another embodiment, the present invention is directed to methods for treating social anxiety disorder comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000195] In a further embodiment, the present invention is directed to methods for treating depression comprising administering the formulations of the present invention to a patient in need thereof.

[000196] In another embodiment, the present invention is directed to methods for treating depression comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000197] In a further embodiment, the present invention is directed to methods for treating patients encountering adverse emotional stimuli comprising administering the formulations of the present invention to a patient in need thereof.

[000198] In another embodiment, the present invention is directed to methods for treating patients encountering adverse emotional stimuli comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000199] In an embodiment, the present invention is directed to methods for treating nausea comprising administering the formulations of the present invention to a patient in need thereof.

[000200] In another embodiment, the present invention is directed to methods for treating nausea comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000201] In an embodiment, the present invention is directed to methods for treating multiple sclerosis comprising administering the formulations of the present invention to a patient in need thereof.

[000202] In another embodiment, the present invention is directed to methods for treating multiple sclerosis comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000203] In an embodiment, the invention is directed to methods for treating symptoms of cannabis use disorder comprising administering formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000204] In another embodiment, the present invention is directed to methods for treating symptoms of cannabis use disorder comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000205] In another embodiment, the invention is directed to methods for treating symptoms of early psychosis comprising administering formulations of the present invention to a patient in need thereof.

[000206] In another embodiment, the present invention is directed to methods for treating symptoms of early psychosis comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000207] In another embodiment, the invention is directed to methods for treating symptoms of Alzheimer's Disease comprising administering formulations of the present invention to a patient in need thereof.

[000208] In another embodiment, the present invention is directed to methods for treating symptoms of Alzheimer's Disease comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000209] In yet another embodiment, the invention is directed to methods for treating symptoms of post-traumatic stress disorder ("PTSD") comprising administering formulations of the present invention to a patient in need thereof.

[000210] In another embodiment, the present invention is directed to methods for treating symptoms of post-traumatic stress disorder PTSD comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000211] In an embodiment, the invention is directed to methods for treating symptoms of anxiety comprising administering formulations of the present invention to a patient in need thereof.

[000212] In another embodiment, the present invention is directed to methods for treating anxiety comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

[000213] In a further embodiment, the invention is directed to methods for treating symptoms of autism comprising administering formulations of the present invention to a patient in need thereof. In a preferred embodiment, the alcohol-free formulations of the present invention are administered to the patient in need of treatment.

[000214] In another embodiment, the present invention is directed to methods for treating symptoms of autism comprising administering synthetically synthesized, substantially pure, cannabidiol to a patient in need thereof.

Definitions

[000215] As used herein, a “patient” refers to a single patient and not a patient population.

[000216] As used herein, “synthetic” refers to the chemical synthesis of cannabidiol does not refer to cannabidiol that is extracted from cannabis plant material.

[000217] As used herein, “substantially pure” refers to a preparation having chromatographical purity of cannabidiol of greater than 98 %, preferably greater than 98.5 %, more preferably greater than 99.0 %, and most preferably greater than 99.5 %.

[000218] As used herein, “substantially free of delta-9-tetrahydrocannabinol” refers to a preparation of cannabidiol having less than 0.3 % of delta-9-tetrahydrocannabinol as determined by HPLC. Preferably, the preparation contains less than 0.25 % of delta-9-tetrahydrocannabinol, more preferably 0.2 %, and most preferably less than 0.1 % of delta-9-tetrahydrocannabinol.

[000219] As used herein, all numerical values relating to amounts, weights, and the like, that are defined as “about” each particular value is plus or minus 10 %. For example, the phrase “about 10 % w/w” is to be understood as “9 % w/w to 11 % w/w.” Therefore, amounts within 10 % of the claimed value are encompassed by the scope of the claims.

[000220] As used here, “liquid” refers to a flowable, fluid pharmaceutical formulation. This type of formulation is not a powder or solid.

[000221] All weights herein refer to % w/w or percent weight of the total formulation.

[000222] As used herein the term “effective amount” refers to the amount necessary to treat a patient in need thereof.

[000223] As used herein the term “pharmaceutically acceptable” refers to ingredients that are not biologically or otherwise undesirable in an oral dosage form.

[000224] As used herein, “qs” means a sufficient quantity of that component to reach a desired volume or concentration.

[000225] The disclosed embodiments are simply exemplary embodiments of the inventive concepts disclosed herein and should not be considered as limiting, unless the claims expressly state otherwise.

[000226] The following examples are intended to illustrate the present invention and to teach one of ordinary skill in the art how to use the formulations of the invention. They are not intended to be limiting in any way.

[000227] All claims, aspects and embodiments of the invention, and specific examples thereof, are intended to encompass equivalents thereof.

Examples

Example 1. Alcohol-free formulations

[000228] The formulations in Table 1 below were prepared as follows. All the solvents are purged with nitrogen before using in manufacturing. Vitamin E, methyl paraben, propyl paraben were dissolved in propylene glycol. Polyethylene glycol 400 (PEG400) and a flavoring agent were added to the propylene glycol solution and mixed thoroughly. The water phase was prepared by dissolving sucralose and sodium ascorbate in water. Next, the solutions were combined and pH adjusted using a pH modifier. The cannabinoid was added to the excipient solution and mixed until dissolved.

[000229] Synthetically synthesized, substantially pure, cannabidiol was used as the cannabinoid.

Strawberry flavor was used as the flavoring agent.

Table 1. Alcohol-free Formulations

Formulation	# AF1	# AF2	# AF3	# AF4	# AF5	# AF6	# AF7	# AF8
Cannabinoid	32	32	32	32	32	32	32	32
PEG400	28	28	27.9	27.38			67.95	62.95
Propylene Glycol	34	34	34	34	62.95	67.95		

Water	6	6	6	6	5			5
Vitamin E (Alpha-Tocopherol)				0.05	0.05	0.05	0.05	0.05
Sodium Ascorbate			0.1	0.1				
Methyl Paraben				0.1				
Propyl Paraben				0.02				
Sucralose				0.05				
Flavoring				0.3				
pH adjustment	None	6 to 7	6 to 7	6 to 7				
pH	8.7	6.7	6.6	6.4				

Example 2. Stability of Alcohol-free Formulations

[000230] The formulations listed in Table 1 were subjected to stability at 55 °C ± 2 °C, 40 °C ± 2 °C under 75 % ± 5 % relative humidity, and 25 °C ± 2 °C under 60 % ± 5 % relative humidity. Stability of the formulations was analyzed at specified time points by evaluating for their potency (assay value) and impurity levels. Assay and impurities were detected using high-performance liquid chromatography with an ultraviolet detector. The assay was performed at 228 nm and indicated as a % of initial concentration. For all impurities, analysis was performed at 228 nm and expressed as a % area. Amounts of particular impurities are listed in Tables 2 to 13 as a percentage of area of each formulation along with amount of total impurities. Relative retention time (RRT) is given for each impurity.

Table 2. Stability Data for Cannabidiol Oral Solution Formulation # AF1 stored at 55 °C ± 2 °C

55 °C - Formulation # AF1	RRT	0 Week	1 Week	2 Weeks	3 Weeks	4 Weeks
---------------------------	-----	--------	--------	---------	---------	---------

Assay (% of initial concentration)		100.00	97.11	97.30	94.47	87.91
% Cis-cannabidiol	1.440	0.01	0.02	0.02	0.02	0.02
% Delta-9-tetrahydrocannabinol	1.729	ND	ND	0.01	ND	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.05	0.03	0.03	0.03	0.02
% Unknown Impurity	0.328	ND	BQL	BQL	BQL	0.06
	0.345	ND	BQL	BQL	BQL	0.07
	0.385	ND	BQL	BQL	BQL	0.05
	0.404	ND	0.08	0.13	0.23	0.38
	0.460	ND	0.05	0.07	0.10	0.17
	0.486	ND	0.42	0.65	1.23	2.73
	0.505	BQL	0.22	0.22	0.19	ND
	0.526	ND	0.10	0.14	0.13	0.17
	0.610	ND	ND	BQL	0.05	0.08
	0.702	ND	BQL	BQL	0.07	0.08
	0.742	ND	BQL	BQL	0.05	0.07
	0.774	0.07	0.06	0.06	ND	ND
	0.796	ND	0.58	1.04	2.13	3.80
	0.830	BQL	0.31	0.39	0.59	0.87
	0.933	ND	BQL	0.06	0.17	0.37
	1.881	ND	0.06	0.09	0.06	0.06
	2.025	ND	BQL	BQL	0.34	0.39
	2.291	ND	0.06	ND	ND	ND
Total Impurities (% Area)		0.13	1.99	2.91	5.39	9.41

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 3. Stability Data for Cannabidiol Oral Solution Formulation # AF2 stored at
55 °C ± 2 °C**

55 °C - Formulation # AF2	RRT	0 Week	1 Week	2 Weeks	3 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	100.31	99.90	95.25	96.85
% Cis-cannabidiol	1.440	0.01	0.01	0.01	0.01	0.01
% Delta-9-tetrahydrocannabinol	1.730	ND	ND	0.01	0.03	0.06
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.05	0.07	0.05	0.05	0.04
% Unknown Impurity	0.340	ND	BQL	BQL	0.05	0.07
	0.404	ND	BQL	BQL	BQL	0.08
	0.462	ND	BQL	BQL	BQL	0.05
	0.486	ND	BQL	0.22	0.35	0.94
	0.506	ND	0.07	0.13	0.15	ND
	0.584	ND	BQL	BQL	0.05	0.11
	0.776	0.07	0.07	0.06	0.05	ND
	0.795	ND	BQL	0.30	0.50	1.09
	0.830	BQL	BQL	0.10	0.14	0.22
	0.932	ND	BQL	0.07	0.10	0.18
	2.034	ND	ND	BQL	0.09	BQL
Total Impurities (% Area)		0.13	0.22	0.95	1.57	2.85

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 4. Stability Data for Cannabidiol Oral Solution Formulation # AF3 stored at
55 °C ± 2 °C**

55 °C - Formulation # AF3	RRT	0 Week	1 Week	2 Weeks	3 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	99.25	98.60	98.28	96.12
% Cis-cannabidiol	1.440	0.01	0.01	0.01	0.01	0.01
% Delta-9-tetrahydrocannabinol	1.736	ND	ND	ND	0.01	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.05	0.05	0.05	0.05	0.05
% Unknown Impurity	0.484	ND	ND	ND	BQL	0.14
	0.502	ND	BQL	BQL	0.05	0.09
	0.775	0.06	0.09	0.10	0.06	0.05
	0.793	ND	ND	ND	0.06	0.27
	0.830	BQL	BQL	BQL	BQL	0.06
	0.951	ND	BQL	ND	BQL	0.05
	1.158	ND	0.06	0.08	0.12	0.05
Total Impurities (% Area)		0.12	0.21	0.24	0.36	0.79

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 5. Stability Data for Cannabidiol Oral Solution Formulation # AF4 stored at
55 °C ± 2 °C**

55 °C - Formulation # AF4	RRT	0 Week	1 Week	2 Weeks	3 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	100.92	99.27	100.16	98.10
% Cis-cannabidiol	1.440	0.01	0.01	0.01	0.01	0.01

% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.05	0.05	0.05	0.06	0.07
% Unknown Impurity	0.403	ND	BQL	BQL	BQL	0.06
	0.485	ND	BQL	0.06	0.18	0.38
	0.505	ND	BQL	0.05	0.08	0.12
	0.524	ND	ND	BQL	BQL	0.07
	0.776	0.07	0.08	0.05	0.06	ND
	0.794	ND	ND	0.07	0.31	0.70
	0.822	ND	ND	BQL	0.10	0.15
	0.931	ND	ND	ND	BQL	0.06
	1.159	ND	BQL	0.08	0.10	ND
	1.774	ND	ND	ND	0.05	0.11
Total Impurities (% Area)		0.13	0.14	0.37	0.95	1.73

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 6. Stability Data for Cannabidiol Oral Solution Formulation # AF1 stored at
40 °C ± 2 °C under 75 % ± 5 % relative humidity**

40 °C - Formulation # AF1	RRT	0 Week	2 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	100.18	95.64
% Cis-cannabidiol	1.440	0.01%	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.846	0.05%	0.05%	0.03%
	0.404	ND	BQL	0.12%
	0.460	ND	0.07%	0.08%

% Unknown Impurity	0.486	ND	0.23%	0.87%
	0.505	BQL	0.30%	0.30%
	0.526	ND	0.05%	0.14%
	0.702	ND	BQL	0.06%
	0.774	0.07%	0.07%	ND
	0.796	ND	0.25%	1.31%
	0.830	BQL	0.12%	0.44%
	0.931	ND	ND	0.06%
Total Impurities (% Area)		0.13%	1.15%	3.42%

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

Table 7. Stability Data for Cannabidiol Oral Solution Formulation # AF2 stored at 40 °C ± 2 °C under 75 % ± 5 % relative humidity

40 °C - Formulation # AF2	RRT	0 Week	2 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	100.08	98.77
% Cis-cannabidiol	1.442	0.01%	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.848	0.05%	0.05%	0.04%
% Unknown Impurity	0.484	ND	ND	0.08%
	0.506	ND	BQL	0.11%
	0.776	0.07%	0.07%	0.06%
	0.794	ND	ND	0.09%
	0.830	BQL	BQL	0.05%
Total Impurities (% Area)		0.13%	0.13%	0.44%

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 8. Stability Data for Cannabidiol Oral Solution Formulation #AF3 stored at
40 °C ± 2 °C under 75% ± 5% relative humidity**

40 °C - Formulation # AF3	RRT	0 Week	2 Week	4 Week
Assay (% of initial concentration)		100.00	98.47	96.90
% Cis-cannabidiol	1.442	0.01%	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.846	0.05%	0.05%	0.05%
% Unknown Impurity	0.775	0.06%	0.08%	0.10%
	1.160	ND	ND	0.05%
Total Impurities (% Area)		0.12%	0.14%	0.21%

ND-Not Detected

**Table 9. Stability Data for Cannabidiol Oral Solution Formulation # AF4 stored at
40 °C ± 2 °C under 75 % ± 5 % relative humidity**

40 °C - Formulation # AF4	RRT	0 Week	2 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	99.63	99.50
% Cis-cannabidiol	1.437	0.01%	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.05%	0.05%	0.06%
% Unknown Impurity	0.776	0.07%	0.07%	0.08%
Total Impurities (% Area)		0.13%	0.13%	0.15%

**Table 10. Stability Data for Cannabidiol Oral Solution Formulation # AF1 stored at
25 °C ± 2 °C under 60 % ± 5 % relative humidity**

25 °C - Formulation # AF1	RRT	0 Week	4 Weeks
Assay (% of initial concentration)		100.00	101.24
% Cis-cannabidiol	1.440	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.846	0.05%	0.04%
% Unknown Impurity	0.459	ND	0.09%
	0.483	ND	0.11%
	0.505	BQL	0.27%
	0.774	0.07%	0.06%
	0.796	ND	0.10%
	0.836	BQL	0.06%
Total Impurities (% Area)		0.13%	0.74%

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 11. Stability Data for Cannabidiol Oral Solution Formulation # AF2 stored at
25 °C ± 2 °C under 60 % ± 5 % relative humidity**

25 °C - Formulation # AF2	RRT	0 Week	4 Weeks
Assay (% of initial concentration)		100.00	100.22
% Cis-cannabidiol	1.442	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.848	0.05%	0.05%
% Unknown Impurity	0.776	0.07%	0.07%
Total Impurities (% Area)		0.13%	0.13%

Table 12. Stability Data for Cannabidiol Oral Solution Formulation # AF3 stored at 25 °C ± 2 °C under 60 % ± 5 % relative humidity

25 °C - Formulation # AF3	RRT	0 Week	4 Weeks
Assay (% of initial concentration)		100.00	97.52
% Cis-cannabidiol	1.442	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.846	0.05%	0.05%
% Unknown Impurity	0.775	0.06%	0.08%
Total Impurities (% Area)		0.12%	0.14%

Table 13. Stability Data for Cannabidiol Oral Solution Formulation # AF4 stored at 25 °C ± 2 °C under 60 % ± 5 % relative humidity

25 °C - Formulation # AF4	RRT	T=0	4 Weeks
Assay (% of initial concentration)		100.00	99.26
% Cis-cannabidiol	1.437	0.01%	0.01%
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.05%	0.06%
% Unknown Impurity	0.776	0.07%	0.07%
Total Impurities (% Area)		0.13%	0.14%

[000231] Control formulation (# AF1) showed significant increase in levels of total impurities and decrease in the assay value. Adjusting the pH of formulation (# AF2) in the range of from about 6 to about 7 increased the stability of the formulation in comparison to control formulation. This illustrates the critical role that pH plays in cannabinoid formulations' stability. Applicant determined that the pH should be from about 6 to about 7 for optimal stability. Addition of antioxidants along with pH adjustment further

increased the stability of the cannabinoid formulation. For example, formulations # AF3 and # AF4, containing antioxidant(s) and pH modifiers, showed excellent stability for four weeks regardless of temperature and humidity conditions.

Example 3. Alcohol formulations

[000232] The formulations in Tables 14 and 15 below were prepared as follows. All the solvents were purged with nitrogen before using in manufacturing. Vitamin E, ascorbyl palmitate, methyl paraben, propyl paraben, sucralose were dissolved in ethanol. propylene glycol, polyethylene glycol 400, glycerol, flavoring agent, and water were added to the solution and mixed thoroughly. Then, if applicable, the pH of the solution was adjusted using a pH modifier. The cannabinoid was added to the excipient solution and mixed until completely dissolved.

[000233] Synthetically synthesized, substantially pure, cannabidiol was used as the cannabinoid. Strawberry flavor was used as the flavoring agent.

Table 14. Formulations with Alcohol

Formulation	# A5	# A6	# A7	# A8
Cannabinoid	9.1	9.1	9.1	8.8
Polyethylene glycol 400	3	3	3	3
Propylene Glycol	7.5	7.5	7.5	7.5
Ethanol	50.3	50.2	50.2	49.7
Water	30	30	30	30.5
Vitamin E (Alpha-Tocopherol)		0.05	0.05	0.05
Ascorbyl Palmitate		0.1	0.1	0.1
Sucralose	0.05	0.05	0.05	0.05
Methyl Paraben	0.02	0.02	0.02	0.02
Propyl Paraben	0.02	0.02	0.02	0.02
Flavoring				0.3

pH adjustment	None	None	pH adjusted to 6 to 7	pH adjusted to 6 to 7
Final pH of formulation	6.06	4.9	6.5	6.4

Table 15. Additional Formulations with Alcohol

Formulation	# A9	# A10	# A11	# A12	# A13
Cannabinoid	32	32	8.756	32	32
Polyethylene glycol 400	18.8	23.8	3.0		62.85
Propylene Glycol	39	39	7.5	62.85	
Glycerol	5				
Water			30.204		
Ethanol	5	5	50.0	5	5
Vitamin E (Alpha Tocopherol)	0.05	0.05	0.05	0.05	0.05
Ascorbyl Palmitate	0.1	0.1	0.1	0.1	0.1
Sucralose	0.05	0.05	0.05		
Methyl Paraben	0.02	0.02	0.02		
Propyl Paraben	0.02	0.02	0.02		

Example 4. Stability of Formulations with Alcohol

[000234] The formulations listed in Table 14 and Table 15 were subjected to stability at 25 °C ± 2 °C under 60% ± 5% relative humidity and 40 °C ± 2 °C under 75 % ± 5 % relative humidity. Stability of the formulations was analyzed at specified time points by evaluating for their potency (assay value) and impurity levels. Assay and impurities were detected using high-performance liquid chromatography with an ultraviolet detector. The assay was performed at 228 nm and indicated as a % of initial concentration. For all impurities, analysis was performed at 228 nm and expressed as a % area. Amounts of

particular impurities are listed in Table 16 to 22 as a percentage of area of each formulation along with amount of total impurities. Relative retention time (RRT) is given for each impurity.

Table 16. Stability Data for Cannabidiol Oral Solution Formulation # A5 stored at 25 °C ± 2 °C under 60 % ± 5 % relative humidity

25 °C - Formulation # A5	RRT	0 Month	3 Months	6 Months	9 Months	12 Months
Assay (% of initial concentration)		100.00	92.97	83.87	77.31	68.92
% Cannabinol	1.400	ND	ND	ND	0.01	ND
% Cis-cannabidiol	1.455	0.01	0.01	0.01	0.02	0.02
% Delta-9-tetrahydrocannabinol	1.761	ND	ND	0.01	0.15	0.17
% Unknown Impurity	0.319	ND	0.08	0.18	0.34	0.39
	0.337	ND	BQL	BQL	BQL	0.05
	0.370	ND	BQL	0.07	0.08	0.08
	0.389	ND	0.11	0.24	0.42	0.54
	0.448	ND	0.18	0.23	0.24	0.25
	0.479	ND	0.78	1.65	2.66	3.49
	0.494	ND	0.50	0.72	0.82	0.88
	0.522	ND	0.05	BQL	BQL	BQL
	0.600	ND	BQL	0.05	0.09	0.15
	0.678	ND	BQL	0.10	0.16	0.21
	0.697	ND	BQL	0.08	0.08	0.09
	0.713	ND	ND	ND	0.06	0.10

	0.770	0.05	ND	ND	ND	ND
	0.790	ND	0.99	2.28	4.19	5.55
	0.819	ND	0.39	0.87	1.44	1.97
	0.930	ND	0.05	0.21	0.38	0.56
	1.189	ND	ND	ND	BQL	0.09
	2.053	ND	0.07	ND	BQL	0.14
	3.192	ND	ND	ND	ND	0.09
	3.256	ND	ND	ND	0.08	0.08
	3.650	ND	ND	ND	ND	0.13
Total Impurities (% Area)		0.06	3.21	6.70	11.22	15.03

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 17. Stability Data for Cannabidiol Oral Solution Formulation # A6 stored at
25 °C ± 2 °C under 60 % ± 5 % relative humidity**

25 °C - Formulation # A6	RRT	0 Month	3 Months	6 Months	9 Months	12 Months
Assay (% of initial concentration)		100.00	97.49	94.25	91.14	87.53
% Cannabinol	1.400	ND	ND	ND	0.01	ND
% Cis-cannabidiol	1.455	0.01	0.01	0.01	0.01	ND
% Delta-9-tetrahydrocannabinol	1.761	ND	0.06	0.23	0.68	0.82
	0.390	ND	BQL	0.05	0.10	0.14
	0.479	ND	BQL	0.08	0.17	0.25

% Unknown Impurity	0.496	ND	0.20	0.87	1.80	2.41
	0.577	ND	BQL	BQL	0.08	0.10
	0.721	ND	ND	BQL	BQL	0.05
	0.770	0.05	0.05	BQL	BQL	BQL
	0.790	ND	0.05	0.11	0.25	0.43
	0.834	BQL	BQL	BQL	0.05	0.07
	0.961	ND	0.06	0.33	0.71	0.97
	1.197	ND	ND	ND	ND	0.06
	1.869	BQL	BQL	BQL	0.06	0.27
	2.066	ND	0.07	0.42	0.59	0.86
	3.247	ND	ND	ND	0.07	0.08
	3.655	ND	ND	ND	ND	0.11
Total Impurities (% Area)		0.06	0.50	2.10	4.58	6.62

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 18. Stability Data for Cannabidiol Oral Solution Formulation # A7 stored at
25 °C ± 2 °C under 60 % ± 5 % relative humidity**

25 °C - Formulation # A7	RRT	0 Month	3 Months	6 Months	9 Months	12 Months
Assay (% of initial concentration)		100.00	98.69	96.52	96.30	96.54
% Cis-cannabidiol	1.455	0.01	0.01	0.01	0.01	0.01
% Delta-9-tetrahydrocannabinol	1.761	ND	0.01	0.02	0.03	0.05

% Unknown Impurity	0.479	ND	BQL	BQL	BQL	0.07
	0.495	ND	BQL	0.06	0.14	0.20
	0.770	0.05	0.05	0.05	0.05	BQL
	0.793	ND	BQL	0.06	0.06	0.10
	0.958	ND	ND	ND	BQL	0.06
	1.160	ND	BQL	0.05	BQL	0.05
	1.883	ND	ND	ND	ND	0.06
	2.057	ND	ND	BQL	BQL	0.06
	3.652	ND	ND	ND	ND	0.05
Total Impurities (% Area)		0.06	0.07	0.25	0.29	0.71

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 19. Stability Data for Cannabidiol Oral Solution Formulation # A8 stored at
25 °C ± 2 °C under 60 % ± 5 % relative humidity**

25 °C - Formulation # A8	RRT	0 Month	3 Months	6 Months
Assay (% of initial concentration)		100.00	100.51	100.14
% Cis-cannabidiol	1.454	0.04	0.04	0.04
% Delta-9-tetrahydrocannabinol	1.762	0.03	0.04	0.05
% Unknown Impurity	0.501	BQL	BQL	0.07
	1.162	ND	BQL	0.07
	1.198	ND	ND	0.05
Total Impurities (% Area)		0.07	0.08	0.28

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 20. Stability Data for Cannabidiol Oral Solution Formulation # A5 stored at
40 °C ± 2 °C under 75 % ± 5 % relative humidity**

40 °C - Formulation # A7	RRT	0 Month	3 Months	6 Months
Assay (% of initial concentration)		100.00	95.22	89.72
% Cis-cannabidiol	1.451	0.01	0.01	0.01
% Delta-9-tetrahydrocannabinol	1.753	0.01	0.06	0.16
% Unknown Impurity	0.390	ND	0.05	0.15
	0.450	ND	BQL	0.06
	0.476	BQL	0.23	0.75
	0.501	BQL	0.30	0.80
	0.609	ND	BQL	0.05
	0.675	ND	BQL	0.05
	0.772	0.05	BQL	ND
	0.791	ND	0.36	1.35
	0.830	BQL	0.12	0.37
	0.934	ND	BQL	0.25
	0.958	ND	BQL	0.18
	1.333	ND	ND	0.05
	1.982	ND	ND	0.17
	2.062	BQL	0.05	0.32
	3.253	ND	BQL	0.09
	3.744	ND	ND	0.13

WO 2017/204986

PCT/US2017/029843

Total Impurities (% Area)		0.07	1.18	4.94
---------------------------	--	------	------	------

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

**Table 21. Stability Data for Cannabidiol Oral Solution Formulation # A6 stored at
40 °C ± 2 °C under 75 % ± 5 % relative humidity**

40 °C - Formulation # A8	RRT	0 Month	3 Months	6 Months
Assay (% of initial concentration)		100.00	96.57	92.84
% Cis-cannabidiol	1.454	0.04	0.03	0.03
% Delta-9-tetrahydrocannabinol	1.762	0.03	0.13	0.62
% Unknown Impurity	0.392	ND	0.06	0.14
	0.478	ND	0.22	0.64
	0.501	BQL	0.41	0.84
	0.610	ND	BQL	0.05
	0.670	ND	BQL	0.05
	0.792	ND	0.38	1.15
	0.821	ND	0.12	0.30
	0.931	ND	0.05	0.19
	0.956	ND	0.09	0.21
	2.068	BQL	0.11	0.23
	3.251	ND	BQL	0.09
	3.754	ND	ND	0.13
Total Impurities (% Area)		0.07	1.60	4.67

ND-Not Detected

BQL-Below Quantification Limit, for unknown impurity only

Table 22. Stability Data for Cannabidiol Oral Solution Formulation # A7 stored at 40 °C ± 2 °C under 75 % ± 5 % relative humidity

40 °C - Formulation # A9	RRT	0 Week	2 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	99.77	100.65
% Cis-cannabidiol	1.440	0.01	0.01	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.841	0.05	0.06	0.05
% Unknown Impurity	0.770	0.06	0.07	0.08
Total Impurities (% Area)		0.12	0.14	0.14

Table 23. Stability Data for Cannabidiol Oral Solution Formulation # A8 stored at 40 °C ± 2 °C under 75 % ± 5 % relative humidity

40 °C - Formulation # A10	RRT	0 Week	2 Weeks	4 Weeks
Assay (% of initial concentration)		100.00	101.25	100.78
% Cis-cannabidiol	1.440	0.01	0.01	0.01
% Delta-9-tetrahydrocannabinol	1.723	ND	ND	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.842	0.05	0.05	0.05
% Unknown Impurity	0.770	0.07	0.07	0.06
Total Impurities (% Area)		0.13	0.13	0.13

ND-Not Detected

[000235] Control formulation (# A5) showed significant increase in levels of total impurities and decrease in the assay value. The addition of antioxidants, Vitamin E and ascorbyl palmitate (see # A6) significantly increased the stability of formulation. These results illustrate the critical role of antioxidants in stabilizing cannabinoid formulations. Antioxidants Vitamin E and ascorbic acid (or its salt) show excellent synergism as ascorbic acid (or its salt) strongly inhibits the depletion of Vitamin E by regenerating it. Along with the antioxidants, the addition of pH modifiers to adjust the pH to the range of 6 to 7 resulted

in exceptionally stable formulations (# A7 and # A8). The stability testing data illustrates that the pH range of from about 6 to about 7 is critical. Formulations # A9 and # A10 also showed good stability after four weeks.

Example 5: Lipid Formulations

[000236] The formulations in Table 24 were created by mixing all the solid and liquid excipients in the lipid. Cannabidiol was then dissolved. Synthetically synthesized, substantially pure, cannabidiol used as the source of the cannabinoid. Strawberry was used as the source of flavoring.

Table 24. Formulations with Lipids

Formulation	#LF1	#LF2	#LF3	#LF4	#LF5	#LF6	#LF7	#LF8
Cannabinoid	24.6	19.5	19.5	19.5	19.5	18	28	18
Vitamin E (Alpha Tocopherol)		0.05	0.05	0.05	0.05	0.05	0.05	
Flavor		0.3	0.3	0.3	0.3	0.3		
Sesame oil	75.4	80.15	70.15					
Sunflower oil				80.45				
Soybean oil						81.95		
Corn Oil					80.45			
Olive Oil								82.00
Caprylic/Capric Triglyceride (Miglyol® 812N)							61.95	
Ethanol			10.0				10.0	

Table 25. Additional Formulations with Lipids

Formulation	#LF9	#LF10	#LF11	#LF12	#LF13	#LF14	#LF15	#LF16
Cannabinoid	31.09	31.09	31.09	31.09	31.09	31.09	31.09	31.09
Ascorbyl palmitate					0.1	0.1	0.1	0.1
Vitamin E (Alpha Tocopherol)	0.1	0.2	0.5	1.0		0.1		
Flavor		0.3	0.3		0.3	0.3	0.3	0.3
Saccharin	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
Caprylic/Capric Triglyceride (Miglyol® 812N)	68.785	68.385	68.085	67.885	67.485	67.385	63.485	58.485

Ethanol					1.0	1.0	5.0	10.0
---------	--	--	--	--	-----	-----	-----	------

Formulation	#LF17	#LF18	#LF19	#LF20	#LF21	#LF22	#LF23	#LF24
Cannabinoid	31.09	31.09	31.09	31.09	31.09	31.09	31.09	31.09
BHA	0.05	0.05	0.05		0.05	0.05	0.1	0.01
BHT	0.01	0.01	0.01		0.01	0.01	0.1	0.005
TBHQ				0.02	0.02			
Propyl gallate			0.02					
EDTA						0.05		
Ascorbyl palmitate								
Linoleic acid								
Propylparaben								
Methylparaben								
Vitamin E (Alpha Tocopherol)	0.05							
Flavor							0.3	0.3
Saccharin	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
Caprylic/Capric Triglyceride (Miglyol® 812N)	68.775	68.825	68.805	68.865	68.805	68.775	68.385	68.57
Miglyol® 840								
Olive Oil								
Ethanol								

Formulation	#LF25	#LF26	#LF27	#LF28	#LF29	#LF30	#LF31	#LF32
Cannabinoid	31.09	31.09	31.09	31.09	31.09	31.09	31.09	31.09
BHA		0.01						
BHT	0.005							
TBHQ								
Propyl gallate								
EDTA								
Ascorbyl palmitate				0.05		0.02	0.02	0.02
Linoleic acid			4	4	10			
Propylparaben								
Methylparaben								
Vitamin E (Alpha Tocopherol)								
Flavor	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3

Saccharin	0.025	0.025	0.025	0.025	0.025	0.025	0.025	0.025
Caprylic/Capric Triglyceride (Miglyol® 812N)	68.58	68.575	64.585	64.535	58.585	68.565	63.565	58.565
Miglyol® 840								
Olive Oil								
Ethanol						1	5	10

Formulation	#LF33	#LF34	#LF35	#LF36	#LF37	#LF38	#LF39	#LF40
Cannabinoid	31.09	31.09	31.09	31.09	31.09	31.09	18	18
BHA			0.05	0.05			0.01	
BHT		0.05	0.01	0.01			0.05	0.05
TBHQ								
Propyl gallate								
EDTA								
Ascorbyl palmitate	0.02	0.02						
Linoleic acid								
Propylparaben						0.01		
Methylparaben						0.1		
Vitamin E (Alpha Tocopherol)	0.05				0.2	0.2		
Flavor	0.3	0.3	0.3		0.3	0.3		
Saccharin	0.025	0.025		0.025	0.025	0.025		
Caprylic/Capric Triglyceride (Miglyol® 812N)	63.515	63.515	68.55		68.385	68.275		
Miglyol® 840				68.825				
Olive Oil							81.94	81.95
Ethanol	5	5						

Formulation	#LF41	#LF42
Cannabinoid	10.53	10.98
BHA		
BHT		
TBHQ		
Propyl gallate		
EDTA		
Ascorbyl palmitate		
Linoleic acid		
Propylparaben		

Sucralose		0.05
Vitamin E (Alpha Tocopherol)	0.2	
Flavor	0.3	0.02
Saccharin	0.025	
Caprylic/Capric Triglyceride (Miglyol® 812N)	88.945	
Sesame oil		80.28
Olive Oil		
Ethanol		8.67

Example 6: Stability of a Formulation with Lipids

[000237] Formulation #LF1 was subjected to stability at 25 °C ± 2 °C under 60 % ± 5 % relative humidity and 40 °C ± 2 °C under 75 % ± 5 % relative humidity. Formulations #LF10 and #LF11 were subjected to stability at 55 °C ± 2 °C and 40 °C ± 2 °C under 75 % ± 5 % relative humidity. Formulations #LF8, #LF9 and #LF12-#LF15 were subjected to stability at all 3 storage conditions. The stability of the formulation was analyzed at specified time points by evaluating the potency (assay value) and impurity levels. Assay and impurities were detected using high-performance liquid chromatography with an ultraviolet detector. The assay was performed at 228 nm and indicated as a % of initial concentration. For all impurities, analysis was performed at 228 nm and expressed as a % area. Amounts of particular impurities are listed in Table 25 as a percentage of area of each formulation along with amount of total impurities. Relative retention time (RRT) is given for each impurity.

Table 26. Three Month Stability Data for Cannabidiol Oral Solution

Formulation #LF1 stored at 40 °C ± 2 °C under 75% ± 5% relative humidity and stored at 25 °C ± 2 °C under 60% ± 5% relative humidity

Formulation #LF1	RRT	0 Month	3 Months-40°C	3 Months-25°C
Assay (% of initial concentration)		100.00	100.87	100.72
% Cis-cannabidiol	1.437	0.03	0.04	0.04
% Delta 9-THC	1.736	0.06	0.06	0.08

%Trans-(1R, 6R)-3'-methyl-cannabidiol	1.840	0.02	0.06	0.02
Total Impurities (% Area)		0.11	0.16	0.14

Table 27. Stability Data for Cannabidiol Oral Solution Formulation #LF8

stored at 55°C ± 2 °C

55°C - Formulation #LF8	RRT	T=0	1 Week	2 Week	3 Week	4 Week
Assay (% of Initial Concentration)		100.00	100.25	101.20	100.08	99.41
% Cis-cannabidiol	1.450	0.06	0.05	0.05	0.05	0.05
% Delta 9-THC	1.752	ND	0.02	0.01	0.03	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.07	0.06	0.06
% Total Impurities (% Area)		0.12	0.13	0.13	0.14	0.13

ND-Not Detected

Table 28. Stability Data for Cannabidiol Oral Solution Formulation #LF9

stored at 55°C ± 2 °C

55°C - Formulation #LF9	RRT	T=0	1 Week	2 Week	3 Week	4 Week
Assay (% of Initial Concentration)		100.00	100.69	101.01	98.88	97.63
% Cannabinol	1.395	ND	ND	ND	ND	0.01
% Cis-cannabidiol	1.450	0.01	0.01	0.02	0.02	0.02
% Delta 9-THC	1.749	ND	ND	ND	0.03	0.04
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.05	0.06	0.04	0.04	ND
% Unknown Impurity	0.396	ND	BQL	BQL	0.05	0.06
	0.455	ND	BQL	0.06	0.09	0.11
	0.480	ND	0.11	0.18	0.32	0.39
	0.499	ND	0.07	0.11	0.18	0.23
	0.520	ND	BQL	BQL	0.07	0.08
	0.584	ND	BQL	BQL	0.07	0.09
	0.771	0.07	0.07	0.07	0.05	0.05
	0.796	ND	0.09	0.21	0.40	0.60
	0.824	ND	0.05	0.09	0.10	0.11
	0.853	ND	BQL	BQL	BQL	0.06
	0.920	ND	ND	BQL	BQL	0.05
	1.908	ND	BQL	0.06	0.13	0.22

Total Impurities (% Area)		0.13	0.46	0.84	1.55	2.12
---------------------------	--	------	------	------	------	------

ND-Not Detected

BQL-Below Quantification Limit

**Table 29. Stability Data for Cannabidiol Oral Solution Formulation #LF10 stored at
55°C ± 2 °C**

55°C - Formulation #LF10	RRT	T=0	2 Week	4 Weeks
Assay (% of Initial Concentration)		100.00	98.35	97.12
% Cis-cannabidiol	1.450	0.01	0.01	ND
% Delta 9-THC	1.746	ND	0.04	ND
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.04	0.04	ND
% Unknown Impurity	0.398	ND	BQL	BQL
	0.457	ND	0.09	0.10
	0.483	ND	0.22	0.36
	0.508	ND	0.12	0.17
	0.587	ND	0.05	0.05
	0.771	0.06	0.05	BQL
	0.796	ND	0.29	0.59
	0.823	ND	0.06	0.05
	1.895	ND	0.10	0.07
	18.000	ND	ND	ND
Total Impurities (% Area)		0.11	1.07	1.39

ND-Not Detected

BQL-Below Quantification Limit

**Table 30. Stability Data for Cannabidiol Oral Solution Formulation #LF11
stored at 55°C ± 2 °C**

55°C - Formulation #LF11	RRT	T=0	2 Week	4 Weeks
Assay (% of Initial Concentration)		100.00	98.96	95.50
% Cis-cannabidiol	1.450	0.01	0.01	0.01
% Delta 9-THC	1.751	ND	0.02	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.04	0.05	0.03
	0.397	ND	0.06	0.11
	0.482	ND	0.17	0.35

% Unknown Impurity	0.507	ND	0.13	0.24
	0.771	0.06	0.05	0.05
	0.795	ND	0.32	0.74
	0.823	ND	0.09	0.10
Total Impurities (% Area)		0.11	0.90	1.64

ND-Not Detected

BQL-Below Quantification Limit

Table 31. Stability Data for Cannabidiol Oral Solution Formulation #LF12
stored at 55°C ± 2 °C

55°C - Formulation #LF12	RRT	T=0	1 Week	2 Week	3 Week	4 Week
Assay (% of Initial Concentration)		100.00	99.94	100.87	100.85	99.58
% Cannabinol	1.395	ND	ND	ND	ND	0.01
% Cis-cannabidiol	1.450	0.01	0.01	0.01	0.01	0.01
% Delta 9-THC	1.749	ND	ND	ND	0.05	0.06
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.05	0.07	0.05	0.05	0.05
% Unknown Impurity	0.396	ND	BQL	BQL	0.06	0.08
	0.479	ND	0.06	0.10	0.15	0.23
	0.499	ND	BQL	BQL	0.09	0.11
	0.584	ND	BQL	BQL	0.07	0.10
	0.771	0.07	0.07	0.07	0.06	0.07
	0.796	ND	0.06	0.14	0.21	0.34
	0.824	ND	ND	0.05	BQL	0.06
Total Impurities (% Area)		0.13	0.27	0.42	0.75	1.12

ND-Not Detected

BQL-Below Quantification Limit

Table 32. Stability Data for Cannabidiol Oral Solution Formulation #LF13
stored at 55°C ± 2 °C

55°C - Formulation #LF13	RRT	T=0	1 Week	2 Week	3 Week	4 Week
Assay (% of Initial Concentration)		100.00	99.09	100.73	99.39	99.35
% Cis-cannabidiol	1.450	0.05	0.05	0.05	0.05	0.05
% Delta 9-THC	1.752	0.01	ND	0.02	0.03	0.03

% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.05	0.06	0.06	0.06
% Total Impurities (% Area)		0.12	0.10	0.13	0.14	0.14

ND-Not Detected

Table 33. Stability Data for Cannabidiol Oral Solution Formulation #LF14
stored at 55°C ± 2 °C

55°C - Formulation #LF14	RRT	T=0	1 Week	2 Week	3 Week	4 Week
Assay (% of Initial Concentration)		100.00	100.99	99.20	100.89	100.24
% Cis-cannabidiol	1.450	0.05	0.05	0.05	0.05	0.05
% Delta 9-THC	1.752	0.01	0.01	ND	0.03	0.04
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.08	0.06	0.07
% Total Impurities (% Area)		0.12	0.12	0.13	0.14	0.16

ND-Not Detected

Table 34. Stability Data for Cannabidiol Oral Solution Formulation #LF15
stored at 55°C ± 2 °C

55°C - Formulation #LF15	RRT	T=0	1 Week	2 Week	3 Week	4 Week
Assay (% of Initial Concentration)		100.00	101.11	101.70	100.44	100.70
% Cis-cannabidiol	1.450	0.05	0.05	0.05	0.05	0.05
% Delta 9-THC	1.752	0.01	0.01	0.02	0.03	0.03
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.06	0.06	0.06
% Total Impurities (% Area)		0.12	0.12	0.13	0.14	0.14

Table 35. Stability Data for Cannabidiol Oral Solution Formulation #LF8
stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF8	RRT	T=0	1 Month	2 Months	3 Months
Assay (% of Initial Concentration)		100.00	99.74	99.75	101.93
% Cis-cannabidiol	1.450	0.06	0.05	0.05	0.05
% Delta 9-THC	1.752	ND	ND	ND	0.04

% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.08	0.06
% Unknown Impurity	1.197	ND	BQL	BQL	0.17
% Total Impurities (% Area)		0.12	0.11	0.13	0.32

ND-Not Detected

BQL-Below Quantification Limit

Table 36. Stability Data for Cannabidiol Oral Solution Formulation #LF9
stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF9	RRT	T=0	1 Week	2 Week	4 Week
Assay (% of Initial Concentration)		100.00	100.25	100.92	99.73
% Cis-cannabidiol	1.450	0.01	0.01	0.01	0.01
% Delta 9-THC	1.749	ND	ND	ND	0.04
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.05	0.07	0.06	0.05
% Unknown Impurity	0.455	ND	BQL	BQL	0.05
	0.480	ND	BQL	0.09	0.22
	0.499	ND	BQL	BQL	0.13
	0.771	0.07	0.07	0.07	0.07
	0.796	ND	BQL	BQL	0.19
	0.823	ND	ND	ND	0.08
Total Impurities (% Area)		0.13	0.15	0.23	0.84

ND-Not Detected

BQL-Below Quantification Limit

Table 37. Stability Data for Cannabidiol Oral Solution Formulation #LF10
stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF10	RRT	T=0	2 Week	4 Weeks
Assay (% of Initial Concentration)		100.00	100.33	99.74
% Cis-cannabidiol	1.450	0.01	ND	0.05
% Delta 9-THC	1.746	ND	0.02	ND
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.04	0.04	0.06

% Unknown Impurity	0.483	ND	0.08	0.23
	0.508	ND	0.06	0.16
	0.771	0.06	0.06	0.05
	0.796	ND	0.13	0.43
	0.822	ND	0.05	0.10
Total Impurities (% Area)		0.11	0.44	0.11

ND-Not Detected

BQL-Below Quantification Limit

Table 38. Stability Data for Cannabidiol Oral Solution Formulation #LF11
 stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF11	RRT	T=0	2 Weeks	4 Weeks
Assay (% of Initial Concentration)		100.00	100.99	99.60
% Cis-cannabidiol	1.450	0.01	0.01	0.01
% Delta 9-THC	1.751	ND	0.02	ND
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.04	0.04	ND
% Unknown Impurity	0.482	ND	BQL	0.05
	0.771	0.06	0.05	0.06
	0.795	ND	BQL	0.1
Total Impurities (% Area)		0.11	0.12	0.22

ND-Not Detected

Table 39. Stability Data for Cannabidiol Oral Solution Formulation #LF12
 stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF12	RRT	T=0	1 Week	2 Week	4 Week
Assay (% of Initial Concentration)		100.00	100.04	100.54	100.65
% Cis-cannabidiol	1.450	0.01	0.01	0.01	0.01
% Delta 9-THC	1.749	ND	ND	ND	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.05	0.06	0.05	0.05
% Unknown Impurity	0.771	0.07	0.07	0.07	0.07
% Total Impurities (% Area)		0.13	0.14	0.13	0.15

ND-Not Detected

Table 40. Stability Data for Cannabidiol Oral Solution Formulation #LF13
 stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF13	RRT	T=0	1 Week	2 Week	4 Week
Assay (% of Initial Concentration)		100.00	101.52	101.13	99.79
% Cis-cannabidiol	1.450	0.05	0.06	0.05	0.05
% Delta 9-THC	1.752	0.01	ND	0.02	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.06	0.06
% Total Impurities (% Area)		0.12	0.12	0.13	0.13

ND-Not Detected

Table 41. Stability Data for Cannabidiol Oral Solution Formulation #LF14
 stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF14	RRT	T=0	1 Week	2 Week	4 Week
Assay (% of Initial Concentration)		100.00	101.16	99.75	100.47
% Cis-cannabidiol	1.450	0.05	0.05	0.05	0.05
% Delta 9-THC	1.752	0.01	0.01	ND	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.06	0.06
% Total Impurities (% Area)		0.12	0.12	0.11	0.13

ND-Not Detected

Table 42. Stability Data for Cannabidiol Oral Solution Formulation #LF15
 stored at 40°C ± 2 °C under 75 % ± 5 % relative humidity

40°C - Formulation #LF15	RRT	T=0	1 Week	2 Week	4 Week
Assay (% of Initial Concentration)		100.00	96.78	100.68	100.94
% Cis-cannabidiol	1.450	0.05	0.06	0.05	0.05
% Delta 9-THC	1.752	0.01	ND	0.02	0.02
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06	0.06	0.07
% Total Impurities (% Area)		0.12	0.12	0.13	0.14

ND-Not Detected

Table 43. Stability Data for Cannabidiol Oral Solution Formulation #LF8
 stored at 25°C ± 2 °C under 60 % ± 5 % relative humidity

25°C - Formulation #LF8	RRT	T=0	1 Month	2 Months	3 Months
Assay (% of Initial Concentration)		100.00	100.12	101.11	102.02
% Cis-cannabidiol	1.450	0.06	0.05	0.05	0.05
% Delta 9-THC	1.752	ND	0.01	ND	ND
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.07	0.07	0.06
% Unknown Impurity	1.197	ND	ND	BQL	0.17
% Total Impurities (% Area)		0.12	0.13	0.12	0.22

ND-Not Detected

BQL-Below Quantification Limit

Table 44. Stability Data for Cannabidiol Oral Solution Formulation #LF9
 stored at 25°C ± 2 °C under 60 % ± 5 % relative humidity

25°C - Formulation #LF9	RRT	T=0	4 Week
Assay (% of Initial Concentration)		100.00	100.14
% Cis-cannabidiol	1.450	0.01	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.05	0.05
% Unknown Impurity	0.771	0.07	0.06
Total Impurities (% Area)		0.13	0.12

Table 45. Stability Data for Cannabidiol Oral Solution Formulation #LF12
 stored at 25°C ± 2 °C under 60 % ± 5 % relative humidity

25°C - Formulation #LF12	RRT	T=0	4 Week
Assay (% of Initial Concentration)		100.00	100.69
% Cis-cannabidiol	1.450	0.01	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.05	0.05
% Unknown Impurity	0.771	0.07	0.07
% Total Impurities (% Area)		0.13	0.13

Table 46. Stability Data for Cannabidiol Oral Solution Formulation #LF13
 stored at 25°C ± 2 °C under 60 % ± 5 % relative humidity

25°C - Formulation #LF13	RRT	T=0	4 Week
Assay (% of Initial Concentration)		100.00	99.83
% Cis-cannabidiol	1.450	0.05	0.05
% Delta 9-THC	1.752	0.01	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06
% Total Impurities (% Area)		0.12	0.12

Table 47. Stability Data for Cannabidiol Oral Solution Formulation #LF14
 stored at 25°C ± 2 °C under 60 % ± 5 % relative humidity

25°C - Formulation #LF14	RRT	T=0	4 Week
Assay (% of Initial Concentration)		100.00	100.64
% Cis-cannabidiol	1.450	0.05	0.05
% Delta 9-THC	1.752	0.01	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06
% Total Impurities (% Area)		0.12	0.12

Table 48. Stability Data for Cannabidiol Oral Solution Formulation #LF15
 stored at 25°C ± 2 °C under 60 % ± 5 % relative humidity

25°C - Formulation #LF15	RRT	T=0	4 Week
Assay (% of Initial Concentration)		100.00	100.38
% Cis-cannabidiol	1.450	0.05	0.05
% Delta 9-THC	1.752	0.01	0.01
% Trans-(1R, 6R)-3'-methyl-cannabidiol	1.862	0.06	0.06
% Total Impurities (% Area)		0.12	0.12

[000238] As seen in Table 25 above, formulation # LF1 with sesame oil showed good stability after 3 months at both storage conditions $25^{\circ}\text{C} \pm 2^{\circ}\text{C}/60\% \pm 5\%$ relative humidity and $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \pm 5\%$ relative humidity. Also, formulation #LF8 with olive oil showed good stability after four weeks at storage conditions $55^{\circ}\text{C} \pm 2^{\circ}\text{C}$, after three months at $25^{\circ}\text{C} \pm 2^{\circ}\text{C}/60\% \pm 5\%$ relative humidity and $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \pm 5\%$ relative humidity.

[000239] Formulations #LF9-#LF15 each contain caprylic/capric triglyceride and one of alpha-tocopherol (Vitamin E), ascorbyl palmitate, or a combination thereof as an antioxidant. Formulations #LF13-#LF15 each additionally contain ethanol. Each of formulations #LF9-#LF15 showed good stability after four weeks at storage conditions $55^{\circ}\text{C} \pm 2^{\circ}\text{C}$, $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \pm 5\%$ relative humidity and $25^{\circ}\text{C} \pm 2^{\circ}\text{C}/60\% \pm 5\%$ relative humidity. #LF9-#LF12 demonstrate the ability of alpha-tocopherol (Vitamin E) to surprisingly achieve less than 0.5% total impurities after four weeks at $40^{\circ}\text{C} \pm 2^{\circ}\text{C}/75\% \pm 5\%$ relative humidity. #LF13-#LF15 demonstrate the ability of ascorbyl palmitate to surprisingly achieve less than 0.2% total impurities after four weeks at all 3 storage conditions in formulations containing from 1% to 5% ethanol. #LF14 demonstrates that the addition of alpha tocopherol (Vitamin E) does not improve the surprising stability from the use of ascorbyl palmitate.

Example 7. Paclitaxel Induced Neuropathic Pain Study

[000240] Paclitaxel is an antineoplastic agent that has activity against several types of cancer including ovary, breast, lung and the head and neck. Paclitaxel works by promoting microtubule assembly which results in neuropathy as a toxic side effect. Peripheral sensory neuropathy is the most commonly reported neurotoxic side effect of paclitaxel and it limits treatment with high and cumulative doses of paclitaxel when given alone or in combination with other neurotoxic antineoplastic agents such as cisplatin. Currently there is not a highly effective treatment for this type of pain. Therefore, there is a need for a highly effective treatment to relieve the symptoms of paclitaxel induced neuropathy.

[000241] A mouse study was conducted in order to determine the effects of cannabidiol, delta-9-tetrahydrocannabinol, and cannabidiol plus delta-9-

tetrahydrocannabinol combinations to alleviate neuropathic pain caused by chemotherapy-induced peripheral neuropathy. The cannabidiol administered to the mice was substantially pure, synthetically synthesized, cannabidiol which had a purity greater than 98 %.

[000242] A detailed explanation of Figure 1 is as follows. The Y-axes represent the threshold sensitivity to mechanical stimulation, expressed as a percent of baseline sensitivity. The X-axes represent the dose of drug in milligrams per kilogram ("mg/kg") administered intraperitoneally ("IP"). Whereas the dotted lines represent withdrawal threshold level to mechanical stimulation of saline controls, the dashed lines represent paclitaxel-treated animals. The points along the dashed line indicate neuropathic pain while points along the dotted line represent protection from neuropathic pain. The data shown are mean +SEM sensitivity measured on Day 21 post treatment. * $p < 0.05$ from saline control as determined by one-way ANOVA.

[000243] Specific doses of agents producing similar overt behavioral effects when added to together should produce the additive effect level.

Examples:

- 1) If 1.25 mg/kg cannabidiol produces 100 % alleviation of pain effect and 1.25 mg/kg delta-9-tetrahydrocannabinol produces 0 % effect, then those doses added together should be fully effective (as should the 2.5 mg/kg cannabidiol + 2.5 mg/kg delta-9-tetrahydrocannabinol).
- 2) If 0.625 mg/kg cannabidiol and 0.625 delta-9-tetrahydrocannabinol produce 0% effect, then those doses in combination should be ineffective.

[000244] Applicant found (as illustrated in Fig. 1) that cannabidiol when administered alone provided the most effective level of alleviating chemotherapy-induced neuropathic pain compared to delta-9-tetrahydrocannabinol. The presence of delta-9-tetrahydrocannabinol depending on its concentration can inhibit the ability of cannabidiol to alleviate neuropathic pain. The ability of delta-9-tetrahydrocannabinol to block the pain alleviating activity of cannabidiol is also dependent of the concentration of cannabidiol. This test illustrates that a substantially pure cannabidiol formulation is highly desirable.

Example 8. Additional Paclitaxel Induced Neuropathic Pain Study

Methods

[000245] Paclitaxel was administered on days 1, 3, 5, and 7 following baseline mechanical sensitivity, and cannabinoids were administered 15 minutes prior to each paclitaxel injection. Mechanical sensitivity was then reassessed on days 9, 14, and 21. For mechanical sensitivity testing, mice are placed on a wire mesh surface inside individual clear Plexiglas chambers, and the plantar surface of their hindpaw is touched with increasing thicknesses of Von Frey filaments (0.16 – 2.0 grams of force) until they withdraw their paw from the stimulation. Von Frey hairs are a series of fine, calibrated filaments that are pressed against the plantar surface of the mouse into a bent “C” shape for 6 seconds. For each treatment group the final sample size was 8 animals. Two-way ANOVA were used to determine significant effects of CBD and THC treatment.

[000246] Single agent dose effect curves are shown as percent level of mechanical sensitivity at baseline to normalize data. Dose equivalence analysis was used to determine significant synergistic effects of CBD+THC compared with predicted additive values derived from single agent dose response curves. To obtain predicted and observed effect levels, data were transformed into percent maximal possible effect (MPE) of the cannabinoid to reverse paclitaxel-induced mechanical sensitivity. To determine this value for each animal, the mean sensitivity score of the paclitaxel control group on given test day is set at zero and the animal's baseline score prior to treatment is set at 100. For example, if an animal has a mechanical sensitivity score of 1.0 at baseline and a score of 0.75 on day 9, and the paclitaxel group shows an average score of 0.5 on day 9, then the animal's percent MPE score is 50%. A %MPE score of 0% would indicate that the animal was at least as sensitive as the paclitaxel control group. A %MPE score of 100% would indicate that the animal was as sensitive or less sensitive on test day as it was at baseline. This transformation of the data is necessary to determine effective dose levels (ED50s, ED25s, etc).

Results

[000247] Pretreatment with CBD or THC significantly attenuated paclitaxel-induced mechanical sensitivity, $P < 0.0001$ for each agent. See Figure 2. CBD produced this effect with higher potency, in that the minimal effective dose for CBD was 1.25 mg/kg IP, while

the minimal effective dose for THC was 2.5 mg/kg IP. Two-way ANOVA also revealed a significant difference between the CBD and THC dose response curves, with the 1.25 mg/kg dose of CBD producing a significantly higher % baseline score as compared to 1.25 mg/kg dose of THC. Both drugs appeared to be efficacious.

[000248] Across a wider range of doses, it becomes apparent that both CBD and THC do not produce monotonic dose effects but instead follow an inverted-U or N shaped function. For both CBD and THC at each time point, the curve turns over at between 5.0 and 10 mg/kg, but the treatments regain efficacy at higher doses. See Figure 3, top center and middle center panels. In the combination groups, the data appear more U-shaped, although it is unclear whether the rise would again emerge with larger dose combinations. See Figure 3, bottom middle panel.

[000249] Dose equivalence analysis was used to predict the combined effects of CBD and THC based on their effects alone on the ascending limbs of their dose response curves. The individual dose effect equations are $E=78.47D^{2.5}/D^{2.5}+0.497$ for CBD, and $E=80D^3/D^3+3.44$ for THC. In dose equivalence analysis, for each CBD dose, an effect-equivalent dose of THC is identified. This dose is added to the actual THC dose in each combination so that the sum is the effective dose of the predicted combination. For example, to predict the additive effect of 0.31 mg/kg CBD and 0.31 mg/kg THC, a dose of THC is identified that is equi-effective to 0.31 mg/kg CBD using the determined dose effect equation for CBD. CBD 0.31 mg/kg produces a %MPE of 8.3%. From this, the dose of THC to produce a %MPE of 8.3 is calculated using the determined dose effect equation for THC. The dose of THC required to achieve a %MPE is 0.7 mg/kg, and this represents a dose that is equi-effective to 0.31 mg/kg CBD. The 0.7 mg/kg is added to the 0.31 mg/kg to give 1.01 mg/kg THC, whose effect level will equal the predicted effect level of 0.31 mg/kg CBD + 0.31 mg/kg THC. This predicted effect level is determined to be 13.68 %MPE. When the actual combination experiment was conducted, 0.31 mg/kg CBD + 0.31 mg/kg THC (labeled on the graph as 0.625 mg/kg combination), was actually the ED78 (78% maximum possible effect; Figure 2 bottom panel). Modified t-test statistics are applied and it was determined that the predicted combination dose response curve was statistically significantly different from the observed dose response curve, demonstrating a synergistic effect of CBD+THC combinations. See Figure 4.

Example 9. Additional Paclitaxel Induced Neuropathic Pain Study*Methods*

[000250] A study was designed to test the efficacy of CBD+THC combinations outside of the 1:1 dose ratio. Six additional combinations were tested: 4:1, 3:1, 2:1, 1:2, 1:3, and 1:4. Four doses of each treatment combination were tested in groups of mice treated with paclitaxel. For each treatment group the final sample size was 8 animals.

Results

[000251] The 4:1 combination of CBD to THC produced a similar effect to CBD alone, while 2:1 and 3:1 ratios of CBD to THC were more potent than CBD alone. See Figure 5. Two-way ANOVA revealed an overall effect of treatment and of dose ($p<0.05$) but no significant interaction. Combinations higher in THC than in CBD produced an effect similar to THC alone, with a significant effect of dose ($p<0.05$) but no main effect of treatment and no significant interaction.

Example 10. Oxaliplatin or Vincristine Induced Neuropathic Pain Study*Methods*

[000252] A study was designed to test the efficacy of CBD in preventing oxaliplatin- or vincristine- induced peripheral neuropathy. Two doses of CBD and vehicle were tested against each of these first line chemotherapeutic agents. Oxaliplatin was administered once at a dose of 6 mg/kg. CBD was administered 15 minutes prior to the single oxaliplatin injection. Vincristine was administered once daily for 7 days at a dose of 0.1 mg/kg. CBD was administered 15 minutes prior to each vincristine injection. For each treatment group the final sample size was 8 animals.

Results

[000253] Pretreatment with CBD attenuated oxaliplatin but not vincristine-induced mechanical sensitivity. See Figure 6. Two-way ANOVA for oxaliplatin revealed a significant effect of time and a significant effect of treatment ($p<0.05$) and no significant interaction. Two-way ANOVA for vincristine revealed a significant effect of time ($p<0.05$), but no main effect of treatment and no interaction.

Example 11. Anticonvulsant Study

[000254] This study was conducted as follows according to standard models for anticonvulsant screening including the maximal electroshock test (“MES”), the minimal clonic seizure (“6 Hz”) test and evaluations of toxicity (“TOX”). The data was recorded as number of animals protected (N) out of the number of animals tested (F), see Tables 26 to 29 below. The test was repeated one time. The cannabidiol administered to the mice and rats was substantially pure, synthetically synthesized, cannabidiol which had a purity greater than 98 %. The cannabidiol was dissolved in 0.5% methylcellulose or a 1:1:18 ratio of ethanol:polyethoxylated castor oil:phosphate buffered saline (“PBS”).

[000255] The maximal electroshock test is a model for generalized tonic-clonic seizures and provides an indication of a compound’s ability to prevent seizure spread when all neuronal circuits in the brain are maximally active. These seizures are highly reproducible and are electrophysiologically consistent with human seizures. For all tests based on maximal electroshock convulsions, 60Hz of alternating current (50 mA in mice, 150 in rats) was delivered for 0.2s by corneal electrodes which were primed with an electrolyte solution containing an anesthetic agent (0.5% tetracaine HCl). The mice were tested at various intervals following doses of 10, 30 and 100 mg/kg of cannabidiol given by intraperitoneal injection of a volume of 0.01 mL/g. An animal was considered “protected” from maximal electroshock-induced seizures upon abolition of the hindlimb tonic extensor component of the seizure.

[000256] The minimal motor impairment test was used to determine the compounds’ undesirable side effects or toxicity. During this test, the animals were monitored for overt signs of impaired neurological or muscular function. The rotarod procedure was used to disclose minimal muscular or neurological impairment. When a control mouse is placed on a rod that rotates at a speed of 6 rpm, the animal can maintain its equilibrium for long periods of time. The animal was considered toxic if it fell off this rotating rod three times during a 60 second period. In addition to minimal motor impairment, the animals may have exhibited a circular or zigzag gait, abnormal body posture and spread of the legs, tremors, hyperactivity, lack of exploratory behavior, somnolence, stupor, catalepsy, loss of placing response and changes in muscle tone.

[000257] The third test was the minimal clonic seizure (6Hz) test. Like the maximal electroshock test, the minimal clonic seizure (6Hz) test is used to assess a compound's efficacy against electrically induced seizures but uses a lower frequency (6Hz) and longer duration of stimulation (3s). Cannabidiol was pre-administered to mice via intraperitoneal injection. At varying times, individual mice (four per time point) were challenged with sufficient current delivered through corneal electrodes to elicit a psychomotor seizure in 97 % of animals (32 mA for 3s). Untreated mice will display seizures characterized by a minimal clonic phase followed by stereotyped, automatic behaviors described originally as being similar to the aura of human patients with partial seizures. Animals not displaying this behavior are considered protected.

Table 49. Anticonvulsant Screening, Mice, Methylcellulose

Time (Hours)		0.5	1.0	2.0
Test	Dose	N/F	N/F	N/F
6HZ	10	0/4	0/4	0/4
6HZ	30	0/4	0/4	0/4
6HZ	100	1/4	0/4	0/4
MES	10	0/4	0/4	0/4
MES	30	0/4	0/4	0/4
MES	100	0/4	1/4	2/4
TOX	10	0/8	0/8	0/8
TOX	30	0/8	0/8	0/8
TOX	100	0/8	0/8	0/8

Table 50. Anticonvulsant Screening, Mice, Ethanol:Polyethoxylated castor oil:PBS

Time (Hours)		0.5	1.0	2.0
Test	Dose	N/F	N/F	N/F
6HZ	10	0/4	0/4	0/4
6HZ	30	0/4	0/4	0/4
6HZ	100	2/4	0/4	0/4
MES	10	0/4	0/4	0/4
MES	30	0/4	1/4	0/4
MES	100	0/4	2/4	1/4
TOX	10	0/8	0/8	0/8
TOX	30	0/8	0/8	0/8
TOX	100	0/8	0/8	0/8

Table 51. Anticonvulsant Screening, Rats, Methylcellulose

Time (Hours)		1.0	2.0	4.0
Test	Dose	N/F	N/F	N/F
MES	30	0/4	0/4	0/4
MES	100	0/4	0/4	0/4
TOX	30	0/4	0/4	0/4
TOX	100	0/4	0/4	0/4

Table 52. Anticonvulsant Screening, Rats, Ethanol:Polyethoxylated castor oil:PBS

Time (Hours)		1.0	2.0	4.0
Test	Dose	N/F	N/F	N/F
MES	30	0/4	0/4	0/4
MES	100	1/4	0/4	0/4
TOX	30	0/4	0/4	0/4
TOX	100	0/4	0/4	0/4

[000258] As seen in Tables 49 to 52 above, Applicant found that cannabidiol protected the mice and rats from epilepsy.

Example 12. 6 Hz Psychomotor Seizure Test

[000259] This study was conducted in order to determine the ability of synthetically-synthesized, substantially pure cannabidiol to block a psychomotor seizure induced by long-duration frequency (6 Hz) stimulation. This is a study model for therapy-resistant partial seizures.

[000260] Adult male CFI mice (weighing 18 to 25 g) were pretreated intraperitoneally with the cannabidiol at a dose of 100 mg/kg. The cannabidiol administered to the mice was substantially pure, synthetically synthesized, cannabidiol which had a purity greater than 98 %. The cannabidiol was dissolved in 0.5% methylcellulose or a 1:1:18 ratio of ethanol:polyethoxylated castor oil:PBS.

[000261] Each treatment group (n = 4 mice/group) was examined for anticonvulsive effects at one of five time points (1/4, 1/2, 1, 2, and 4 hours) following treatment with cannabidiol. Following pretreatment, each mouse received a drop of 0.5 % tetracaine hydrochloride applied to each eye. The mouse was then challenged with the low-frequency (6 Hz) stimulation for 3 seconds delivered through corneal electrodes. The low-frequency, long-duration stimuli was initially delivered at 32 mA intensity. Animals were manually

restrained and released immediately following the stimulations and observed for seizure activity. If the test compound was effective in the 32 mA screen, an additional assay wherein the stimulation current is increased to 44 mA is employed using the same protocol as described above. Additionally, a dose response curve can be generated at the time of peak effect (TPE) at the specific stimulation intensity.

[000262] Typically, the 6 Hz stimulation results in a seizure characterized by a minimal clonic phase that is followed by stereotyped, automatistic behaviors, including twitching of the vibrissae, and Straub-tail. Animals not displaying such behaviors were considered protected. Data was analyzed by Mann-Whitney U test, with $p < 0.05$ determined to be statistically significant.

[000263] For each time group, the results are expressed as the total number of animals protected out of the number of animals tested over time (*i.e.*, 2/4 represents 2 out of 4 mice tested were protected).

Table 53. ED50 Biological Response, Methylcellulose

Time (Hours)		0.5
Test	Dose	N/F
6 Hz	30	0/8
6 Hz	65	5/8
6 Hz	130	5/8
6 Hz	160	8/16
6 Hz	190	7/8

Table 54. Time to Peak Effect, Methylcellulose

Time (Hours)		0.25	0.5	1	2	4	6	24
Test	Dose	N/F	N/F	N/F	N/F	N/F	N/F	N/F
6 Hz	300	1/8	0/8	0/8	0/8	0/8	0/8	0/8
6 Hz	500	1/8	0/8	0/8	0/8	0/8	0/8	2/8

Table 55. ED50 Biological Response, Ethanol:Polyethoxylated castor oil:PBS

Test	Dose	Time	N/F
6 Hz	50	0.5	1/8
6 Hz	100	0.5	1/8

6 Hz	130	0.5	4/8
6 Hz	170	0.5	6/8
6 Hz	200	0.5	8/8
TOX	200	2	0/8
TOX	250	2	4/8
TOX	300	2	6/8
TOX	500	2	8/8

Table 56. Time to Peak Effect, Ethanol:Polyethoxylated castor oil:PBS

Time (Hours)		0.25	0.5	1	2	4	6	8	24
Test	Dose	N/F	N/F	N/F	N/F	N/F	N/F	N/F	N/F
TOX	200	-	-	-	0/8	0/8	-	-	-
TOX	250	-	-	-	4/8	3/8	-	-	-
TOX	300	-	-	-	6/8	7/8	4/8	2/8	1/8
TOX	500	0/8	0/8	0/8	8/8	8/8	8/8	-	4/7

[000264] As seen in Tables 53 to 56, cannabidiol in both solvents showed comparable median effective doses that inhibited seizures in 50 % of animals (ED50s) in the 100 mg/kg range. While cannabidiol dissolved in the methylcellulose solvent had an ED50 of 103.75 mg/kg (95 % Confidence Interval of 53.89 mg/kg to 163.84 mg/kg), it showed an ED50 of 121.52 mg/kg when dissolved in the 1:1:18 ethanol:polyethoxylated castor oil:PBS solvent (95 % Confidence Interval of 87.83 mg/kg to 152.96 mg/kg). Based on the toxicity data for the cannabidiol in the methylcellulose solvent, the median toxicity dose where toxicity is observed in 50 % of animals ("TD50") was determined to exceed 500 mg/kg at 0.5 hours post administration. Diarrhea at 24 hours and 1 death was reported at 24 hours at 500 mg/kg, the highest dose tested.

[000265] The TD50 was determined to be 262.37 mg/kg (95 % Confidence Interval of 232.64 to 301.78) with cannabidiol dissolved in the 1:1:18 ethanol:polyethoxylated castor oil:PBS solvent. Death was reported at 24 hours at 300 mg/kg and at 6 and 24 hours for 500 mg/kg with the with the 1:1:18 ethanol:polyethoxylated castor oil:PBS solvent.

[000266] These results further illustrate that cannabidiol is likely to be effective in humans for the treatment of epilepsy and other conditions. Further, synthetically synthesized cannabidiol will likely be less toxic than cannabidiol that is derived from plants and not substantially pure.

Example 13. Maximal Electroshock Seizure and Subcutaneous Metrazol

[000267] The maximal electroshock seizure ("MES") and subcutaneous Metrazol ("sc Met") tests have been the two most widely employed preclinical seizure models for the early identification and high through-put screening of investigational anti-seizure drugs. These tests have been extremely effective in identifying new anti-seizure drugs that may be useful for the treatment of human generalized tonic-clonic seizures and generalized myoclonic seizures. The MES test provides an indication of CBD's ability to prevent seizure spread when all neuronal circuits in the brain are maximally active. The s.c. Met test detects the ability of CBD to raise the chemoconvulsant-induced seizure threshold of an animal and, thus, protect it from exhibiting a clonic, forebrain seizure.

[000268] For the MES test, 60 Hz of alternating current is delivered by corneal electrodes for 0.2 seconds. Supra-maximal seizures are elicited with a current intensity five times that necessary to evoke a threshold tonic extension seizure, *i.e.*, 50 mA in mice and 150 mA in rats. A drop of anesthetic solution, 0.5 % tetracaine hydrochloride, is placed on the eyes of each animal just before the corneal electrodes are applied to the eyes to elicit electrical stimulation. The animals are restrained by hand and released immediately following stimulation to allow observation of the entire seizure. Inhibition of the hind leg tonic extensor component is taken as the endpoint for the MES test.

[000269] A dose of Metrazol (85 mg/kg in mice) will induce convulsions in 97 % of mice (CD97). The CD97 dose of Metrazol is injected into a loose fold of skin in the midline of the neck. The CD97 doses for Metrazol are confirmed annually in mice. It is administered to mice at a volume of 0.01 ml/g body weight. The animals are then placed in isolation cages to minimize stress and continuously monitored for the next 30 min for the presence or absence of a seizure. An episode of clonic spasms, approximately 3 to 5 seconds, of the fore and/or hind limbs, jaws, or vibrissae is taken as the endpoint. Animals not displaying fore and/or hind limb clonus, jaw chomping, or vibrissae twitching are considered protected.

[000270] All quantitative *in vivo* antiseizure/behavioral impairment studies are typically conducted at the previously determined TPE. Groups of at least 8 mice were tested with various doses of cannabidiol until at least two points are established between

the limits of 100 % protection or minimal toxicity and 0 % protection or minimal toxicity. The dose of drug required to produce the desired endpoint in 50 % of animals (ED50 or TD50) in each test, the 95% confidence interval, the slope of the regression line, and the standard error of the mean (S.E.M.) of the slope is then calculated by probit analysis.

[000271] The cannabidiol administered to the mice was substantially pure, synthetically synthesized, cannabidiol which had a purity greater than 98 %. The cannabidiol was dissolved in 0.5% methylcellulose or a 1:1:18 ratio of ethanol:polyethoxylated castor oil:PBS. The maximal electric shock (MES) and subcutaneous Metrazol ("sc MET") are the most widely used preclinical seizure models for the early identification and screening of new antiepileptic drugs.

Table 57. ED50 Biological Response, Methylcellulose

Test	Dose	Time	N/F
MES	200	2	5/8
MES	250	2	4/8
MES	300	2	4/8
MES	350	2	3/8
MES	400	2	3/8
MES	450	2	6/8
MES	500	2	8/8
Sc MET	150	2	1/8
Sc MET	200	2	3/8
Sc MET	300	2	5/8
Sc MET	360	2	7/8
TOX	500	2	0/8

Table 58. Time to Peak Effect, Methylcellulose

Time (Hours)		0.25	0.5	1	2	4
Test	Dose	N/F	N/F	N/F	N/F	N/F
MES	300	0/4	1/4	1/4	4/8	2/4
Sc MET	200	0/4	0/4	2/8	3/8	-
TOX	300	0/4	0/4	0/4	0/4	0/4

Table 59. ED50 Biological Response, Ethanol:Polyethoxylated castor oil:PBS

Test	Dose	Time	N/F
MES	75	2	1/8
MES	95	2	5/8

MES	120	2	7/8
MES	150	2	8/8
Sc MET	120	2	0/8
Sc MET	160	2	2/8
Sc MET	220	2	5/8
Sc MET	260	2	7/8
TOX	175	2	0/8
TOX	250	2	4/8
TOX	325	2	6/8
TOX	500	2	8/8

Table 60. Time to Peak Effect, Ethanol:Polyethoxylated castor oil:PBS

Time (Hours)	0.25	0.5	1	2	4	6	8
Test	Dose	N/F	N/F	N/F	N/F	N/F	N/F
TOX	500	0/8	0/8	0/8	8/8	7/8	7/8

[000272] The ED50 in the MES model for cannabidiol dissolved in the methylcellulose solvent could not be calculated due to a U shaped dose response (1/4 protected at 0.5 hr, 1/4 at 1hr, 4/8 at 2hr and 2/4 at 4hr). However, the ED50 for cannabidiol dissolved in the 1:1:18 ethanol:polyethoxylated castor oil:PBS solvent is 92.21 mg/kg (95 % Confidence Interval of 78.4 mg/kg to 104.63 mg/kg).

[000273] For the MET model, the ED50 was 241.03 mg/kg (95% Confidence Interval of 182.23 to 311.87) for cannabidiol dissolved in the methylcellulose solvent and 198.51 mg/kg (95 % Confidence Interval of 167.76 mg/kg to 232.58 mg/kg) for cannabidiol dissolved in the 1:1:18 ethanol:polyethoxylated castor oil:PBS solvent. Based on the toxicity data for cannabidiol dissolved in the methylcellulose solvent the TD50 was determined to exceed 500 mg/kg, the highest dose tested.

[000274] Myoclonic jerks were reported in at 1 hour with 200 mg/kg dose and at 2 hours with 360 mg/kg dose. The TD50 was determined to be 266.76 mg/kg (95 % Confidence Interval of 222.28 mg/kg to 317.42 mg/kg) with the cannabidiol dissolved in the 1:1:18 ethanol:polyethoxylated castor oil:PBS solvent.

[000275] These results further illustrate that cannabidiol is likely to be effective in humans for the treatment of epilepsy and other conditions. Further, synthetically

synthesized cannabidiol will likely be less toxic than cannabidiol that is derived from plants and not substantially pure.

Example 14. Glioblastoma Multiforme Study

[000276] A study was conducted in order to determine the extent to which systemic administration of cannabidiol or cannabidiol plus delta-9-tetrahydrocannabinol (cannabidiol/delta-9-tetrahydrocannabinol 1:1) can inhibit glioblastoma multiforme progression and enhance the activity of temozolomide, a chemotherapy drug, in an orthotopic mouse model of glioblastoma multiforme utilizing U87 cells. It was previously suggested that the combination of cannabidiol plus delta-9-tetrahydrocannabinol is the most effective treatment for targeting tumors derived from U87 serum-derived glioblastoma multiforme cells.

[000277] The study was conducted as follows. Human U87 luciferase labeled cells were grown in Roswell Park Memorial Institute media with 10 % fetal bovine serum and then harvested from dishes while in their exponential growth phase in culture with 0.1 % trypsin/ethylenediaminetetraacetic acid and washed twice with serum-free Roswell Park Memorial Institute media. For the intracranial model, tumors were generated in female athymic nu/nu mice by the intracranial injection of 0.3×10^6 U87 cells in 4 μ l of Roswell Park Memorial Institute media. Using this model, you can assess drug efficacy (in vivo imaging) as well as survival in the same group of animals. Survival studies were carried out in accordance with the National Institutes of Health's guidelines involving experimental neoplasia and our approved Institutional Animal Care and Use Committees protocol. Animals in all groups are removed from the study when they demonstrate any single sign indicative of significant tumor burden development, including hunched back, sustained decreased general activity, or a significant decrease in weight. In limited cases where tumors were able to escape the intracranial space, the mice were euthanized when the external tumors measured greater than 5mm as assessed by callipers. Additionally, mice with tumors measuring $>500 \times 10^6$ radiance were removed from the study even if symptoms were not observed to assure spontaneous deaths related to seizures did not occur do to the existence of the large intracranial tumor.

[000278] The cannabinoids were dissolved in a mixture of 3 % ethanol, 3 % surfactant and 94 % saline, and temozolomide was dissolved in 30 % dimethyl sulfoxide and 70 % saline. Cannabidiol that was synthetically synthesized and substantially pure was used in this study. The treatments were initiated 9 days after the injection of the tumor cells. Mice were imaged the morning before the first injection to determine initial tumor size and then groups were organized to have equal distribution of tumor size before the initiation of the first injection. Mice were treated once a day for five days with temozolomide. Mice were treated once a day, 5 days a week (Monday through Friday), with the cannabinoids until the completion of the study, except for the first week of the study where mice were injected over the weekend. All mice were administered the treatments via intraperitoneal injection. There were 12 mice per group, for a total of 72 mice. The treatment rates were as follows: cannabidiol (15 mg/kg); cannabidiol/delta-9-tetrahydrocannabinol (1:1, together @ 15mg/kg); and temozolomide (2 mg/kg intraperitoneal injection).

[000279] Significant differences were determined using a one-way ANOVA. Bonferroni-Dunn post-hoc analyses were conducted when appropriate. Survival between groups was compared using a long-rank Mantel-Cox test. P values <0.05 defined statistical significance.

[000280] A detailed explanation of Figure 7 is as follows. The X-axis represents the number of days after treatment and the Y-axis represents the survival rates.

[000281] As seen in Figure 7, while 15 mg/kg of cannabidiol alone or cannabidiol/delta-9-tetrahydrocannabinol (1:1) did not inhibit glioblastoma multiforme progression, it enhanced the antitumor activity of suboptimal doses of temozolomide leading to a significant increase in survival. Further, the substantially pure, synthetically synthesized, cannabidiol produced full regression of 20 % of tumors. This effect was not observed following the 1:1 cannabidiol:delta-9-tetrahydrocannabinol treatments. It was unexpected that substantially pure, synthetically synthesized, cannabidiol would have these effects because previously it was thought that a 1:1 ratio of cannabidiol (that was extracted from cannabis and not substantially pure):delta-9-tetrahydrocannabinol would produce better effects than cannabidiol alone. However, this study again illustrates the superiority of Applicant's substantially pure, synthetically synthesized, cannabidiol.

[000282]

Example 15. Additional Glioblastoma Multiforme Study

[000283] Human U251 luciferase labeled cells were grown in Roswell Park Memorial Institute media with 10% fetal bovine serum and then harvested from dishes while in their exponential growth phase in culture with 0.1% trypsin/ ethylenediaminetetraacetic acid, and washed twice with serum-free Roswell Park Memorial Institute media. For the intracranial model, tumors were generated in female athymic nu/nu mice by the intracranial injection of 0.3×10^6 U251 cells in 4 μ l of Roswell Park Memorial Institute media. Using this model, you can assess drug efficacy (in vivo imaging) as well as survival in the same group of animals. Survival studies were carried out in accordance with the National Institutes of Health's guidelines involving experimental neoplasia and our approved Institutional Animal Care and Use Committees protocol. Animals in all groups are removed from the study when they demonstrate any single sign indicative of significant tumor burden development, including hunched back, sustained decreased general activity, or a significant decrease in weight. In limited cases where tumors were able to escape the intracranial space, the mice were euthanized when the external tumors measured greater than 5mm as assessed by calipers. Additionally, mice with tumors measuring $>500 \times 10^6$ radiance were removed from the study even if symptoms were not observed to assure spontaneous deaths related to seizures did not occur due to the existence of the large intracranial tumor. There were 12 mice per group, for a total of 72 mice. The treatment rates were as follows: cannabidiol (15 mg/kg); temozolomide (1.5 mg/kg intraperitoneal injection); and cannabidiol/temozolomide (10:1, together at 16.5 mg/kg.)

[000284] For drug treatment studies, cannabinoids were dissolved in a mixture of 2.5 % ethanol, 2.5 % Tween[®] 80 and 95% saline, and temozolomide was dissolved in 30% dimethyl sulfoxide and 70% saline. The treatments were initiated 9 days after the injection of the tumor cells. Mice were imaged the morning before the first injection to determine initial tumor size and then groups were organized to have equal distribution of tumor size before the initiation of the first injection. Mice were treated once a day for five days with temozolomide. Mice were treated once a day, 5 days a week (Monday through Friday), with cannabinoids until the completion of the study, except for the first week of the study where mice were injected over the weekend. All mice were injected intraperitoneally.

[000285] Significant differences were determined using a one-way ANOVA. Bonferroni-Dunn post-hoc analyses were conducted when appropriate. Survival between groups was compared using a Kaplan-Meier analysis and long-rank Mantel-Cox test or The Gehan-Breslow-Wilcoxon test. P values <0.05 defined statistical significance.

[000286] A detailed explanation of Figure 8 is as follows. The X-axis represents the number of days after treatment and the Y-axis represents the survival rates.

[000287] One of tumors in the vehicle group fully regressed overtime creating an outlier in the study. Tumor regression in a vehicle treated animal is a rare occurrence but it can occur. Since during the start of the study, the tumor did demonstrate a small increase in growth as assessed by IVIS imaging, it could not be removed from the data set. The data is presented with (Figure 8A) and without (Figure 8B) the outlier for comparison. With the vehicle outlier included, temozolomide alone did not increase survival ($p=0.48$, Figure 8A, $p<0.05$ is considered significant). Cannabidiol alone also did not increase survival. However, the combination of temozolomide + 15 mg/kg of cannabidiol almost reached significance ($p=0.09$) for increasing survival using a Log-rank Mantel-Cox test, $p<0.05$ is considered significant. If this same data set was analyzed with the Gehan-Breslow-Wilcoxon test then the treatment of temozolomide + cannabinoid did produce a significant increase in survival. The Gehan-Breslow-Wilcoxon test however is a less stringent statistical test in comparison to the Log-rank Mantel-Cox test. It should be noted that 2 of the 11 mice are still alive in the temozolomide + cannabinoid group, and in one of the mice the tumor has fully regressed based on in vivo imaging of the tumor.

[000288] If the vehicle outlier was removed from the data set, then treatment with temozolomide significantly increased survival ($p<0.5$, Figure 8B). The combination of temozolomide + 15 mg/kg of cannabidiol, however was highly significant at increasing survival ($p=0.005$). Thus, cannabidiol enhanced the antitumor activity of temozolomide.

Example 16. Pharmacokinetic Study of Multiple Dose Cannabidiol Oral Solution in Pediatric Subjects with Treatment-resistant Seizure Disorders

Protocol

[000289] A Phase 1/2, open label, Multiple Ascending Dose study will be conducted to evaluate the effect of multiple doses of cannabidiol oral solution on pediatric subjects

experiencing treatment-resistant seizures. The study will assess pharmacokinetics, safety, tolerability and preliminary efficacy of 3 doses (10, 20, and 40 milligrams per kilogram per day ("mg/kg/day")) of cannabidiol oral solution administered in a sequential fashion. Specifically, twenty subjects will be enrolled in each dose cohort that A) fit the following criteria: 1. subject and/or parent(s)/caregiver(s) fully comprehend the informed consent form (ICF) and assent form, understand all study procedures, and can communicate satisfactorily with the Investigator and study coordinator; 2. provide informed consent and/or assent (as applicable) of subjects and/or parent(s)/caregiver(s) in accordance with applicable laws, regulations, and local requirements; 3. male or female between 1 and 17 years of age (inclusive) at the time of consent; 4. diagnosed with a treatment-resistant seizure disorder in the opinion of the Investigator and as defined as continued seizures despite: a. adequate trials of ≥ 3 antiepileptic drugs ("AEDs"), and b. ≥ 1 prior adequate treatment course with ≥ 2 AEDs in combination (i.e., concurrently); 5. willingness to remain on established AEDs (stable dosing for ≥ 30 days prior to Day 0 and throughout the duration of the study) a. neither a vagus nerve stimulation (VNS) procedure nor ketogenic diet are considered an AED for the purposes of this study; 6. willingness to not start a ketogenic diet during the Treatment Period or, if already on the diet, to make no changes in the diet during the study; 7. a female subject is eligible to participate in the study if she is: a. premenarchal, or b. of childbearing potential with a negative urine pregnancy test at the Screening Visit and at Day 0. If sexually active, she must agree to fulfill one of the following requirements: i. complete abstinence from intercourse ≥ 4 weeks prior to administration of the first dose of the investigational product, throughout the Treatment Period, and 4 weeks after completion or premature discontinuation from the investigational product, and agreement to use a double-barrier method if she becomes sexually active; ii. use of acceptable methods of contraception throughout the study and 4 weeks after completion or premature discontinuation from investigational product. The acceptable method of contraception is double barrier method (i.e., condom plus spermicide or a condom plus diaphragm); 8. a sexually active male subject must be willing to use acceptable methods of contraception throughout the study and for 4 weeks completion of study participation or premature discontinuation from investigational product. The acceptable methods of birth control are abstinence or double barrier birth control (i.e.,

condom plus spermicide or a condom plus diaphragm); 9. in the opinion of the Investigator, the parent(s)/caregiver(s) are willing and able to comply with the study procedures and visit schedules, including venipuncture, inpatient stay at the study center, dosing at the study center (twice a day as needed while an outpatient), and the Follow-up Visits (if applicable); 10. general good health (defined as the absence of any clinically relevant abnormalities as determined by the Investigator) based on physical and neurological examinations, medical history, and clinical laboratory values (hematology, chemistry, and urinalysis) completed during the Screening Visit; and 11. body weight of ≥ 9 kg; and B) do not meet the following criteria: 1. subject or parent(s)/caregiver(s) have daily commitments during the study duration that would interfere with attending all study visits; 2. currently taking concomitant medications that are strong cytochrome P450 3A4 ("CYP3A4") inhibitors or inducers or CYP3A4 sensitive substrates with a narrow therapeutic index; 3. currently taking any other disallowed medications; 4. currently taking felbamate if they had been receiving it for <6 months prior to the Screening Visit; 5. in the opinion of the Investigator, any clinically significant, unstable medical abnormality, chronic disease, or a history of a clinically significant abnormality of the cardiovascular, gastrointestinal, respiratory, hepatic, or renal systems; 6. any disorder or history of a condition (e.g., malabsorption or gastrointestinal surgery) that may interfere with drug absorption, distribution, metabolism, or excretion; 7. history or presence of abnormal electrocardiograms ("ECGs") that are clinically significant in the opinion of the Investigator; 8. for appropriate subjects, an affirmative answer to queries regarding active suicidal ideation with some intent to act but without a specific plan or active suicidal ideation with specific plan and intent on the Columbia Suicide Severity Rating Scale ("C-SSRS") assessment at the Screening Visit, subjects who have significant findings for suicidal ideation as assessed by the C-SSRS must be referred to the Investigator for follow-up evaluation; 9. any history of attempted suicide; 10. history of poor toleration of venipuncture or poor venous access that would cause difficulty in collecting blood samples; 11. participation in any investigational study currently or within 30 days or 5 half-lives ($t_{1/2}$) of the investigational product (whichever is longer) prior to the Screening Visit; 12. taken any cannabinoids (cannabidiol, Δ^9 -tetrahydrocannabinol [Δ^9 -THC], hemp oil, Realm Oil or marijuana) in the 30 days prior to the Screening Visit; 13. history of an

allergic reaction or a known or suspected sensitivity to any substance that is contained in the investigational product formulation; 14. known infection with hepatitis B, hepatitis C, or human immunodeficiency virus (HIV); 15. In the opinion of the Investigator, the subject is unsuitable in any other way to participate in this study; and 16. body weight of >90 kg.

[000290] Each subject will be enrolled in only one dose cohort. No fewer than 2 subjects between 2 and 12 years of age must be dosed through Day 10 prior to dosing any subject <2 years of age. Each of the 3 planned dose cohorts will include 20 subjects for a study total of 60 subjects: 1 to <2 years of age: 5 subjects; 2 to <12 years of age: 9 subjects with at least 3 under the age of 6; and 12 to ≤17 years of age: 6 subjects with at least 3 subjects under age 16. Each subject will complete a Screening Period of up to 28 days and a Treatment Period of 10 days. Subjects will have a Follow-up Visit on Day 14 and a Follow-up Phone Call on Day 17. On Day 1, the investigational product will be administered once in the morning according to the subject's assigned dose level cohort. The evening dose of the investigational product will not be administered on Day 1. Thus, the subject will receive a half-daily dose only (5, 10, or 20 milligrams/kilogram "mg/kg" total) of the investigational product on Day 1. Subjects will not receive a dose from Day 2 through Day 3 but they will remain in an inpatient setting and complete planned assessments. Subjects will be dosed twice daily (i.e., full daily dose of 10, 20, or 40 mg/kg/day) from Day 4 through Day 10 according to the subject's assigned cohort. Doses will be administered at approximately 12-hour intervals. Doses will be administered to subjects in a fasting state on days on which the serial PK samples will be collected (i.e., Day 1 and Day 10.) Fasting times include 1 hour for ages 1 to less than 2 years and 2 hours for ages 2 to 17 years.

[000291] During the Screening, Treatment, and Follow-up Periods, subjects are not to receive the following: (1) medication(s) that are strong CYP3A4 inhibitors or inducers or CYP3A4-sensitive substrates with a narrow therapeutic index; (2) any cannabinoids (cannabidiol, Δ9-THC, hemp oil, Realm Oil or marijuana); corticosteroids; systemic steroid therapy (excluding inhaled medication for asthma treatment); felbamate (if used for <6 months) or (7) any other investigational drug or investigational device. Subjects will remain on established antiepilepsy therapies (i.e., AEDs for which dosing has been stable ≥30 days prior to Day 0) throughout the duration of the Treatment and Follow-up Periods.

[000292] In summary, for subjects ages 1 to <2 years, serial blood sampling for pharmacokinetic (“PK”) analysis will occur at 2, 4, 8, and 12 hours post the Day 1 dose. Serial blood sampling for PK analysis will also occur at predose, 2, 4, 8, and 12 hours post the Day 10 morning dose. For subjects ages 2 to <6 years, serial blood sampling will occur at predose and at 1, 2, 3, 4, 8, 12, 16, 24 (Day 2), and 48 (Day 3) hours post the Day 1 morning dose. Blood samples for PK trough values for cannabidiol and its 7-hydroxy (“OH”) metabolite will be evaluated on Day 8. Collection will occur prior to the morning dose of the investigational product; no investigational product will be administered on Day 11. Serial blood sampling for PK analysis will also occur at predose, 1, 2, 3, 4, 8, 12, and 24 (Day 11) hours post the Day 10 morning dose. For subjects ages 6 to ≤17 years, serial blood sampling for PK analysis will occur predose and at 1, 2, 3, 4, 6, 8, 12, 16, 24 (Day 2), 36 (Day 2), 48 (Day 3), and 72 (Day 4) hours post the Day 1 morning dose. Blood samples for PK trough values for cannabidiol and its 7-OH metabolite will be evaluated on Day 6 (age ≥12 only), Day 8 and Day 9. Collection will occur prior to the morning dose of the investigational product; no investigational product will be administered on Day 11. Serial blood sampling for PK analysis will also occur at predose and at 1, 2, 3, 4, 6, 8, 12, and 24 (Day 11) hours post the Day 10 morning dose. In addition to the above measurements of cannabidiol and its 7-OH metabolite, levels of clobazam and norclobazam will be measured from the samples taken predose on Day 1 (baseline), Day 8, and Day 10 (predose) for subjects who are aged ≥2 years and are currently taking clobazam.

Endpoints

[000293] Endpoints of the study will include the following: (1) Incidence, type, and severity of adverse events (“AEs”) and serious adverse events (“SAEs”) occurring during the Treatment Period (i.e., treatment-emergent adverse events [“TEAEs”]); (2) Changes from baseline in vital signs; (3) Changes from baseline in ECG findings; (4) Changes from baseline in laboratory values (hematology, chemistry, and urinalysis); (5) Plasma PK variables for cannabidiol (parent compound) and its 7-OH metabolite as appropriate: (a) Maximum plasma concentration (C_{max}) and dose normalized C_{max} (C_{max}/D) (b) time to C_{max} (t_{max}); (c) Half-life ($t_{1/2}$); (d) Elimination rate; (e) Oral clearance (cannabidiol only); (f) Volume of distribution (cannabidiol only); (g) Area under the plasma concentration-

time curve from 0 to 12 hours [$AUC_{(0-12)}$] and dose normalized $AUC_{(0-12)}$ [$AUC_{(0-12)}/D$] on Day 1; (h) Area under curve from time 0 to the last quantifiable concentration [$AUC_{(0-last)}$] on Day 1; (i) Area under the plasma concentration-time curve from 0 to infinity ($AUC_{(0-inf)}$) and dose normalized $AUC_{(0-inf)}$ [$AUC_{(0-inf)}/D$] on Day 1 for study subjects ≥ 2 years of age; (j) Metabolite to parent ratios for C_{max} , $AUC_{(0-inf)}$, $AUC_{(0-12)}$ on Day 1 and Day 10; (k) $AUC_{(0-12)}$ and $AUC_{(0-12)}/D$ on Day 10; (l) Minimum plasma concentration (C_{min}) on Day 10; (m) Average plasma concentration (C_{avg}) on Day 10; (n) Accumulation ratios for C_{max} and $AUC_{(0-12)}$ on Day 10; (o) Time linearity; (6) clinical global impressions of improvement (“CGI-I”) assessment on Day 11; and (7) Change from baseline in clinical global impressions of severity (“CGI-S”) assessment from the Screening Visit to Day 11.

Safety

[000294] Subjects will be assessed by measurement of vital signs and neurological examination daily. A Physical Examination will be completed at the Screening Visit, as well as Day 0, Day 11 and Day 14. A 12-lead ECG, will be completed at the Screening Visit, as well as Day 1, Day 4, Day 8, Day 11, and Day 14 (if clinically indicated). Hematology, chemistry, and urine analysis will be performed at the Screening Visit, as well as Day 1, Day 4, Day 8, and Day 11. Hematology, chemistry, and urine analysis will also be performed on Day 14 if clinically indicated.

Methods

[000295] The PK concentrations and parameters for cannabidiol and its 7-OH metabolite in plasma will be summarized by study day, sampling time (where appropriate) and dose using descriptive statistics and graphic displays as appropriate. These results will be graphically displayed by age and mg/kg dose, as appropriate. Exposure relationship with age and body weight will be evaluated using regression and/or inferential analyses, as appropriate. Dose proportionality of cannabidiol and its 7-OH metabolite exposure will be investigated using graphical methods and statistically using a power model approach, as appropriate. Accumulation of cannabidiol and its 7-OH metabolite will be assessed using an appropriate analysis of variance model for exposure PK parameters. Time linearity will also be assessed. Trough concentrations samples collected prior to dosing at the scheduled time points will be assessed graphically for attainment of steady state. Also, time to reach

steady state will be assessed with a stepwise linear trend analysis. Levels of clobazam and norclobazam will be measured from the samples taken predose on Day 1 (baseline), Day 8, and Day 10 (predose) for subjects who are age ≥ 2 and currently taking clobazam and summarized by time point and treatment. All safety assessments, including AEs, clinical laboratory evaluations, vital signs, 12-lead ECGs, C-SSRS, and physical and neurological examinations will be listed. When appropriate, they will be summarized with descriptive statistics by age and dose cohort. The results of the CGI-I, CGI-S, and daily seizure diary assessments will be summarized by descriptive statistics as appropriate.

Results

[000296] Preliminary results are shown in Table 61 below. Cohort #1 was administered a single dose of 5 mg/kg of an alcohol based formulation and then 5 mg/kg BID (10 mg/kg/day) for 7 days. Cohort #2 was administered a single dose of 10 mg/kg of a lipid based formulation and then 10 mg/kg BID (20 mg/kg/day) for 7 days. For cohort #1 single dosing resulted in a mean C_{max} of 59.029 ng/mL, mean T_{max} of 3.0 hours and an AUC_{inf} of 276.95 h*ng/mL. For cohort #2 single dosing resulted in a mean C_{max} of 110.522 ng/mL, mean T_{max} of 4.45 hours and an AUC_{inf} of 879.273 h*ng/mL. As demonstrated, lipid based formulations at twice the dosage and at a single dosing achieved nearly twice the maximum plasma concentration of the alcohol based formulations in nearly an hour and half longer.

[000297] At repeated BID dosing, administration of alcohol based oral cannabinoid formulations resulted in a mean C_{max} of 119.6 ng/mL, mean T_{max} of 2.75 hours and an AUC_{tau} of 581.744 h*ng/mL and the lipid based or al cannabinoid formulations resulted in a C_{max} of 214.28 ng/mL, mean T_{max} of 2.55 hours and an AUC_{tau} of 1135.345 h*ng/mL. As demonstrated, lipid based formulations at twice the dosage and at administered BID for 7 days achieved less than twice the maximum plasma concentration of the alcohol based formulations twelve minutes faster.

Table 61. Pharmacokinetic Parameters for Oral Cannabinoid Solutions

Group	Single Dosing			Twice-a-day Dosing for 7 days		
	T_{max}	C_{max}	AUC_{inf}	T_{max}	C_{max}	AUC_{tau}
	(h)	(ng/mL)	(h*ng/mL)	(h)	(ng/mL)	(h*ng/mL)

Cohort #1	N	20	20	15	N	20	20	19
	Mean	3	59.029	276.95	Mean	2.75	119.6	581.774
	SD	1.62	99.98	237.749	SD	0.97	105.035	282.096
	Min	1	7.03	84.84	Min	1	11.1	95.71
	Median	2.5	21.1	152.1	Median	3	94.35	528.35
	Max	8	439	839.69	Max	4	508	1107.44
	CV%	54.1	169.4	85.8	CV%	35.1	87.8	48.5
Cohort #2	N	20	20	12	N	20	20	18
	Mean	4.45	110.522	879.273	Mean	2.55	214.28	1135.345
	SD	2.26	142.314	955.01	SD	2.09	279.018	959.48
	Min	1	6.46	139.55	Min	0	16.6	281.96
	Median	4	52.85	587.86	Median	2	107	819.21
	Max	8	462	2919.34	Max	8	1090	4105.46
	CV%	50.8	128.8	108.6	CV%	81.9	130.2	84.5

Example 17. Pharmacokinetic Food-Effect Study of Single Dose Cannabidiol Oral Solution in Healthy Subjects

[000298] An open label, randomized, single-dose, two-period, two-way crossover food-effect study was conducted on healthy subjects. The study assessed pharmacokinetics and safety of single-dose of 20 mg/kg/day cannabidiol (i.e. formulation #LF10 from Table 25, above) administered under fasted or fed conditions. Twenty-four (24) subjects were enrolled in the study and each were subjected to the fasted and fed treatment arms in separate periods followed by a 7-day wash-out period. For pharmacokinetic analysis, nominal time and default lambda-z selections were used. All below quantifiable limit values were set to zero and all subjects were included in the analysis.

Safety

[000299] Safety was assessed using the following parameters: inclusion/exclusion criteria, medical history and demographics, medical history update, continuing eligibility, physical examination, clinical laboratory testing, 12-lead electrocardiogram (ECG), urine drug and alcohol screens, prior medication history, concomitant medication, seated blood pressure, pulse, respiration rate, and oral temperature, and adverse event (AE) assessments.

Statistical Methods

[000300] Data from 13 subjects from the fasted treatment and 24 subject from the fed treatment were included in the pharmacokinetic and statistical analyses.

[000301] Blood samples (1 x 6 mL) for cannabidiol and 7-OH-cannabidiol analysis were collected in Vacutainer tubes containing K₂-EDTA as a preservative at 0 hour (predose), 0.25, 0.5, 0.75, 1.0, 1.5, 2.0, 2.5, 3.0, 4.0, 6.0, 8.0, 12, 16, 24, 36, 48, 72, 96, and 120 hours postdose (20 time points) in each study period. The following pharmacokinetic parameters were calculated: peak concentration in plasma (C_{max}), time to peak concentration (T_{max}), last quantifiable concentration determined directly from individual concentration-time data (C_{last}), time of last quantifiable concentration (T_{last}), elimination rate constant (λ_z), terminal half-life ($T_{1/2}$), area under the concentration-time curve from time-zero to the time of the last quantifiable concentration (AUC_{0-t}), area under the plasma concentration time curve from time-zero extrapolated to infinity (AUC_{inf}), percentage of AUC_{inf} obtained by extrapolation (AUC_{extrap}), calculated as $AUC_{extrap} = [(AUC_{inf} - AUC_{0-t})/AUC_{inf}] * 100$, apparent oral clearance (CL/F), calculated as: $CL/F = \text{Dose}/AUC_{inf}$ for dronabinol only, and volume of distribution in the terminal elimination phase (V_d/F), calculated as $V_d/F = (CL/F)/\lambda_z$ for cannabidiol only.

Results and Conclusions

[000302] Results of the pharmacokinetic and statistical analyses for the oral cannabinoid solutions of the present invention are shown in Tables 62-67. Tables 62-64 shows the pharmacokinetic parameters of cannabidiol comparing administration to subjects in a fasted or fed condition.

Table 62 Summary of PK parameters for cannabidiol with replicates combined after oral administration of single 20 mg/kg dose of cannabidiol solution under fasted or fed conditions to healthy volunteers.

Parameter	Fasted				Fed			
	n	Mean	SD	CV%	n	Mean	SD	CV%
T_{max} (h)	13	12.00 (2.50, 36.00)			24	6.00 (3.00, 12.00)		
C_{max} (ng/mL)	13	27.3	37.7	138.1	24	1560	865	55.3
AUC_{0-t} (h*ng/mL)	13	466.2	354.8	76.1	24	9650	3435	35.6
AUC_{0-inf} (h*ng/mL)	11	360.0	187.3	52.0	21	10090	3725	36.9
AUC_{extrap} (%)	11	3.57	1.16	32.5	21	3.84	1.26	33.0
λ_z (h ⁻¹)	11	0.0257	0.0050	19.7	21	0.0165	0.0017	10.1
$t_{1/2}$ (h)	11	27.89	5.18	18.6	21	42.30	4.08	9.6
T_{last} (h)	13	120.01	0.04	0.0	24	120.00	0.02	0.0
C_{last} (ng/mL)	13	0.867	1.34	154.1	24	6.36	2.24	35.2

CL/F (L/h/kg)	11	68.39	29.42	43.0	21	2.173	0.6205	28.6
Vd/F (L/kg)	11	2680	1156	43.1	21	132.7	41.40	31.2

*Arithmetic mean $\pm$ standard deviation, (n) denotes the number of subjects measured.

T_{max} presented as median (min, max).

Table 63 Statistical analysis of the log-transformed systemic exposure parameters of cannabidiol comparing 20 mg/kg dose of cannabidiol under fed conditions to the same dose under fasted conditions after oral administration to healthy volunteers.

Dependent Variable	n		Geometric Mean ^a		Ratio (%) ^b (Fed/Fasted)	90% CI ^c		Power ANOVA	
	Fed	Fasted	Fed	Fasted		Lower	Upper	CV%	CV%
$\ln(C_{max})$	24	13	1393.0405	44.4612	3133.16	1874.38	5237.32	0.1786	51.41
$\ln(AUC_{0-t})$	24	13	9210.7934	668.6687	1377.48	962.32	1971.75	0.2628	32.78
$\ln(AUC_{0-inf})^d$	21	11	9584.9742	326.2316	2938.09	2122.35	4067.37	0.2930	39.63

^a Geometric mean for Fed and Fasted based on least squares mean of log-transformed parameter values

^b Ratio(%) = geometric mean (fed)/geometric mean (fasted)

^c 90% confidence interval

^d due to the limited number of subjects with data for cannabidiol oral solution, under fasted conditions during period 2, period was removed from the model in order to analyze AUC_{0-inf} [000303] Substantial increases in cannabidiol maximum and total exposure, based on $\ln(C_{max})$, $\ln(AUC_{0-t})$, and $\ln(AUC_{0-inf})$, were observed after administration of 20 mg/kg cannabidiol oral solution with food compared to 20 mg/kg cannabidiol oral solution administered under fasted conditions. Cannabidiol C_{max} was approximately 31-fold higher after administration with food compared to administration under fasted conditions. Cannabidiol AUC_{0-t} and AUC_{0-inf} were approximately 14-fold and 29-fold higher, respectively, after administration with food compared to administration under fasted conditions. Median time to reach maximum concentration of cannabidiol (T_{max}) occurred approximately 6 hours earlier with food (6 hours) compared to that under fasted conditions (12 hours). Inter-subject variability was substantially reduced with food: 55.3% CV from 138.1% for C_{max} and 36.9% CV from 52.0% for AUC_{0-inf} .

Table 64 Summary of AUC for cannabidiol with replicates combined after oral administration of single 20 mg/kg dose of cannabidiol solution under fasted or fed conditions to healthy volunteers.

Parameter*	Fasted	Fed
AUC ₀ (h*ng/mL)	0.325 ± 1.17	0.229 ± 0.636
AUC _{0-0.25} (h*ng/mL)	0.363 ± 1.31	0.498 ± 0.874
AUC _{0-0.5} (h*ng/mL)	0.584 ± 1.33	2.45 ± 3.38
AUC _{0-0.75} (h*ng/mL)	1.25 ± 1.34	7.48 ± 11.2
AUC ₀₋₁ (h*ng/mL)	1.93 ± 1.62	21.8 ± 40.0
AUC _{0-1.5} (h*ng/mL)	2.79 ± 2.14	128 ± 215
AUC ₀₋₂ (h*ng/mL)	4.14 ± 2.75	269 ± 381
AUC _{0-2.5} (h*ng/mL)	5.38 ± 4.09	424 ± 547
AUC ₀₋₃ (h*ng/mL)	5.88 ± 3.99	535 ± 675
AUC ₀₋₄ (h*ng/mL)	6.62 ± 3.85	812 ± 874
AUC ₀₋₆ (h*ng/mL)	12.0 ± 9.29	1090 ± 886
AUC ₀₋₈ (h*ng/mL)	22.9 ± 39.3	819 ± 590
AUC ₀₋₁₂ (h*ng/mL)	14.7 ± 10.2	372 ± 338
AUC ₀₋₁₆ (h*ng/mL)	9.76 ± 4.79	112 ± 133
AUC ₀₋₂₄ (h*ng/mL)	5.45 ± 2.76	34.3 ± 11.7
AUC ₀₋₃₆ (h*ng/mL)	5.00 ± 3.78	20.7 ± 5.11
AUC ₀₋₄₈ (h*ng/mL)	2.66 ± 2.40	15.5 ± 4.31
AUC ₀₋₇₂ (h*ng/mL)	1.73 ± 1.89	10.1 ± 3.42
AUC ₀₋₉₆ (h*ng/mL)	1.35 ± 1.73	7.86 ± 2.44
AUC ₀₋₁₂₀ (h*ng/mL)	0.867 ± 1.34	6.36 ± 2.24

*Arithmetic mean ± standard deviation. Number of subjects tested (“n”) under fasted conditions was 13, n for fed is 24.

[000304] Administration of oral cannabinoid solutions of the present invention under fed conditions resulted in an appreciable AUC difference in cannabidiol over administration under fasted conditions within 30 minutes. See Figure 9. Additionally, administration of oral cannabinoid solutions of the present invention under fed conditions resulted in 90 times greater AUC of cannabidiol over administration under fasted conditions at 6 hours.

[000305] Tables 65-67 shows the pharmacokinetic parameters of 7-OH-cannabidiol, the primary and active metabolite of cannabidiol, comparing oral cannabinoid solutions of the present invention under fasted and fed conditions.

Table 65 Summary of PK parameters for 7-OH-cannabidiol with replicates combined after oral administration of single 20 mg/kg dose of cannabidiol solution under fasted or fed conditions to healthy volunteers.

Parameter	Fasted				Fed			
	n	Mean	SD	CV%	n	Mean	SD	CV%
T _{max} (h)	13	8.00 (2.50, 24.00)			24	6.00 (3.00, 12.00)		
C _{max} (ng/mL)	13	22.8	33.4	146.4	24	474	187	39.4
AUC _{0-t} (h*ng/mL)	13	414.7	291.0	70.2	24	4811	1509	31.4
AUC _{0-inf} (h*ng/mL)	12	425.3	331.4	77.9	24	4902	1520	31.0
AUC _{extrap} (%)	12	2.77	2.10	75.8	24	1.93	1.33	68.7
λ _z (h ⁻¹)	12	0.0348	0.0090	25.8	24	0.0292	0.0056	19.3
t _{1/2} (h)	12	21.36	6.26	29.3	24	24.69	5.36	21.7
T _{last} (h)	13	114.47	10.53	9.2	24	120.00	0.02	0.0
C _{last} (ng/mL)	13	0.561	0.770	137.2	24	2.41	1.26	52.3

*Arithmetic mean ± standard deviation, (n) denotes the number of subjects measured.

Tmax presented as median (min, max).

Table 66 Statistical analysis of the log-transformed systemic exposure parameters of 7-OH-cannabidiol comparing 20 mg/kg dose of cannabidiol under fed conditions to the same dose under fasted conditions after oral administration to healthy volunteers.

Dependent Variable	n		Geometric Mean ^a		Ratio (%) ^b (Fed/Fasted)	90% CI ^c		Power ANOVA	
	Fed	Fasted	Fed	Fasted		Lower	Upper	CV%	
ln(C _{max})	24	13	442.6751	34.8933	1268.65	792.62	2030.58	0.1941	43.92
ln(AUC _{0-t})	24	13	4610.3024	630.3322	731.41	532.52	1004.57	0.3077	31.14
ln(AUC _{0-inf})	24	12	4701.4345	657.5873	714.95	520.18	982.65	0.3060	31.28

^a Geometric mean for Fed and Fasted based on least squares mean of log-transformed parameter values

^b Ratio(%) = geometric mean (fed)/geometric mean (fasted)

^c 90% confidence interval

[000306] Substantial increases in 7-OH-cannabidiol maximum and total exposure, based on ln(C_{max}), ln(AUC_{0-t}), and ln(AUC_{0-inf}), were observed after administration of 20 mg/kg cannabidiol oral solution with food compared to 20 mg/kg cannabidiol oral solution administered under fasted conditions. 7-OH-cannabidiol C_{max} was approximately 13-fold higher after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol AUC_{0-t} and AUC_{0-inf} were both approximately 7-fold higher after

administration with food compared to administration under fasted conditions. Additionally, T_{max} occurs 2 hours sooner under fed conditions.

Table 67 Summary of AUC for 7-OH-cannabidiol with replicates combined after oral administration of single 20 mg/kg dose of cannabidiol solution under fasted or fed conditions to healthy volunteers.

Parameter*	Fasted	Fed
AUC ₀ (h*ng/mL)	0.114 ± 0.410	0.131 ± 0.537
AUC _{0-0.25} (h*ng/mL)	0.101 ± 0.363	0.221 ± 0.676
AUC _{0-0.5} (h*ng/mL)	0.217 ± 0.361	0.980 ± 1.29
AUC _{0-0.75} (h*ng/mL)	0.954 ± 0.843	2.87 ± 4.21
AUC ₀₋₁ (h*ng/mL)	2.29 ± 2.13	7.33 ± 11.7
AUC _{0-1.5} (h*ng/mL)	3.91 ± 3.90	47.5 ± 78.5
AUC ₀₋₂ (h*ng/mL)	5.46 ± 5.23	104 ± 150
AUC _{0-2.5} (h*ng/mL)	6.81 ± 6.85	158 ± 202
AUC ₀₋₃ (h*ng/mL)	7.83 ± 8.27	197 ± 235
AUC ₀₋₄ (h*ng/mL)	8.65 ± 8.64	289 ± 290
AUC ₀₋₆ (h*ng/mL)	9.79 ± 7.71	346 ± 188
AUC ₀₋₈ (h*ng/mL)	18.5 ± 34.2	284 ± 128
AUC ₀₋₁₂ (h*ng/mL)	10.1 ± 8.36	187 ± 121
AUC ₀₋₁₆ (h*ng/mL)	7.23 ± 4.28	85.7 ± 36.9
AUC ₀₋₂₄ (h*ng/mL)	6.09 ± 2.70	47.5 ± 23.0
AUC ₀₋₃₆ (h*ng/mL)	4.39 ± 2.45	26.7 ± 13.1
AUC ₀₋₄₈ (h*ng/mL)	3.00 ± 1.91	16.9 ± 7.73
AUC ₀₋₇₂ (h*ng/mL)	1.65 ± 1.38	7.36 ± 3.10
AUC ₀₋₉₆ (h*ng/mL)	1.04 ± 1.11	4.03 ± 1.94
AUC ₀₋₁₂₀ (h*ng/mL)	0.511 ± 0.799	2.41 ± 1.26

*Arithmetic mean ± standard deviation. Number of subjects tested ("n") under fasted conditions was 13, n for fed is 24.

[000307] Administration of oral cannabinoid solutions of the present invention under fed conditions resulted in an appreciable AUC difference in 7-OH-cannabidiol over administration under fasted conditions within 45 minutes. See Figure 10. Additionally, administration of oral cannabinoid solutions of the present invention under fed conditions resulted in 35 times greater AUC of 7-OH-cannabidiol over administration under fasted conditions at 6 hours.

[000308] In conclusion, oral cannabinoid solutions of the present invention have a substantial food effect resulting in higher peak plasma concentrations in a shorter period of time and also higher overall plasma concentrations after oral administration following food intake as compared to oral administration following fasting.

Example 18. Pharmacokinetic Food-Effect Study of Single Dose Cannabidiol Oral Solution in Healthy Subjects

Method

[000309] An open label, randomized, single-dose, four-treatment four-period, four-way crossover food-effect study of multiple formulations was conducted on healthy subjects. The study assessed pharmacokinetics of a single-dose of 10 mg/kg of 3 separate cannabidiol formulations (#LF41 and #LF42 from Table 26 above and #A11 from Table 15 above) administered under fed conditions and 1 cannabidiol formulation (#LF42 above) under fasted conditions. Subjects were enrolled in the study and each was subjected to the fasted and fed treatment arms in separate periods followed by a 7-day wash-out period. For pharmacokinetic analysis, nominal time and default lambda-z selections were used. All below quantifiable limit values were set to zero and all subjects were included in the analysis.

Results

i. Cannabidiol Oral Solution Sesame Oil Formulation #LF42, fed vs. fasted

[000310] Substantial increases in both cannabidiol and 7-OH-cannabidiol maximum and total exposure, based on $\ln(C_{\max})$, $\ln(AUC_{0-t})$, and $\ln(AUC_{0-\infty})$, were observed after administration of formulation #LF42 at 10 mg/kg with food compared to formulation #LF42 at 10 mg/kg administered under fasted conditions. Cannabidiol $C_{\max}$ was approximately 10.5-fold higher after administration with food compared to administration under fasted conditions. Cannabidiol AUC_{0-t} and $AUC_{0-\infty}$ were approximately 7.5-fold and 3.3-fold higher, respectively, after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol $C_{\max}$ was approximately 4.5-fold higher after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol AUC_{0-t} and $AUC_{0-\infty}$ were approximately 4.3-fold and 3.7-fold higher, respectively, after administration with food compared to administration under fasted

conditions. Thus, there is a substantial food effect when administering lipid formulations of the present invention.

ii. Cannabidiol Oral Solution Medium Chain Triglyceride (MCT) Formulation #LF41 fed vs. #LF42 fasted

[000311] Substantial increases in both cannabidiol and 7-OH-cannabidiol maximum and total exposure, based on $\ln(C_{\max})$, $\ln(AUC_{0-t})$, and $\ln(AUC_{0-\infty})$, were observed after administration of formulation #LF41 at 10 mg/kg with food compared to formulation #LF42 at 10 mg/kg administered under fasted conditions. Cannabidiol $C_{\max}$ was approximately 10.8-fold higher after administration with food compared to administration under fasted conditions. Cannabidiol AUC_{0-t} and $AUC_{0-\infty}$ were approximately 6.8-fold and 3.3-fold higher, respectively, after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol $C_{\max}$ was approximately 4.1-fold higher after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol AUC_{0-t} and $AUC_{0-\infty}$ were approximately 4.0-fold and 3.5-fold higher, respectively, after administration with food compared to administration under fasted conditions. Thus, there is a substantial food effect when administering medium chain glyceride formulations of the present invention.

iii. Cannabidiol Oral Solution Ethanol Formulation #A11 fed vs. #LF42 fasted

[00314] Substantial increases in both cannabidiol and 7-OH-cannabidiol maximum and total exposure, based on $\ln(C_{\max})$, $\ln(AUC_{0-t})$, and $\ln(AUC_{0-\infty})$, were observed after administration of formulation #A11 at 10 mg/kg with food compared to formulation #LF42 at 10 mg/kg administered under fasted conditions. Cannabidiol $C_{\max}$ was approximately 9.5-fold higher after administration with food compared to administration under fasted conditions. Cannabidiol AUC_{0-t} and $AUC_{0-\infty}$ were approximately 6.9-fold and 4.0-fold higher, respectively, after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol $C_{\max}$ was approximately 3.8-fold higher after administration with food compared to administration under fasted conditions. 7-OH-cannabidiol AUC_{0-t} and $AUC_{0-\infty}$ were approximately 4.6-fold and 4.0-fold higher, respectively, after administration with food compared to administration under fasted conditions. Thus, there is a substantial food effect when administering hydro-alcohol formulations of the present invention.

CLAIMS:

1. An oral pharmaceutical formulation comprising:
about 10.53% (w/w) cannabidiol;
about 88.945% (w/w) caprylic/capric triglyceride; and
about 0.2% (w/w) Vitamin E, wherein the cannabidiol is greater than 98% pure, the formulation is a liquid formulation for oral administration, and wherein a 10 mg/kg daily dose of the formulation is for providing a cannabidiol C_{max} of at least 10 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
2. The formulation of claim 1, wherein the cannabidiol is synthetically synthesized.
3. The formulation of claim 1, wherein the formulation is for providing a cannabidiol AUC(0-inf) of at least 3 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
4. The formulation of claim 1, wherein the formulation is for providing a cannabidiol AUC(0-t) of at least 7 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
5. The formulation of claim 1, wherein the formulation is for providing a 7-OH-cannabidiol C_{max} of at least 4 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
6. The formulation of claim 1, wherein the formulation is for providing a 7-OH-cannabidiol AUC(0-inf) of at least 3.5 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
7. The formulation of claim 1, wherein the formulation is for providing a 7-OH-cannabidiol AUC(0-t) of at least 4 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
8. Use of a formulation of claim 1, wherein the formulation is for treating a disease or disorder selected from the group consisting of Prader-Willi syndrome, obesity, graft versus host disease, gelastic seizures/hypothalamic hamartoma, neonatal seizures, movement disorders including dystonia, central pain syndromes, phantom limb pain, multiple sclerosis, traumatic brain injury, radiation therapy, acute and chronic graft versus host disease, T-cell autoimmune disorders, colitis, Dravet Syndrome, Lennox Gastaut Syndrome, myoclonic seizures, juvenile

myoclonic epilepsy, refractory epilepsy, schizophrenia, juvenile spasms, West syndrome, infantile spasms, refractory infantile spasms, tuberous sclerosis complex, brain tumors, neuropathic pain, cannabis use disorder, post-traumatic stress disorder, anxiety, early psychosis, Alzheimer's Disease, autism, acne, Parkinson's disease, social anxiety disorder, depression, diabetic retinopathy, diabetic nephropathy, diabetic neuropathy, ischemic injury of heart, ischemic injury of brain, chronic pain syndrome, and rheumatoid arthritis.

9. Use of a formulation of claim 1, wherein the formulation is for treating withdrawal from opioids, cocaine, heroin, amphetamines or nicotine.

10. The formulation of claim 1, further comprising a flavoring agent.

11. The formulation of claim 10, wherein the flavoring agent is strawberry flavor.

12. The formulation of claim 1, further comprising a sweetener.

13. The formulation of claim 12, wherein the sweetener is saccharin.

14. The formulation of claim 1, wherein the formulation is substantially free of ethanol.

15. An oral pharmaceutical formulation comprising:

about 10.53% w/w cannabidiol;

about 0.2% w/w Vitamin E;

about 0.3 % w/w flavoring agent;

about 0.025% w/w sweetener, and

about 88.945% w/w caprylic/capric triglyceride,

wherein the cannabidiol is greater than 98% pure, the formulation is a liquid formulation for oral administration, and wherein a 10 mg/kg daily dose of the formulation is for providing a cannabidiol C_{max} of at least 10 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.

16. The formulation of claim 15, wherein the cannabidiol is synthetically synthesized.

17. The formulation of claim 15, wherein the formulation is for providing a cannabidiol AUC(0-inf) of at least 3 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.

18. The formulation of claim 15, wherein the formulation is for providing a cannabidiol AUC(0-t) of at least 7 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.

19. The formulation of claim 15, wherein the formulation is for providing a 7-OH-cannabidiol C_{max} of at least 4 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
20. The formulation of claim 15, wherein the formulation is for providing a 7-OH-cannabidiol AUC(0-inf) of at least 3.5 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
21. The formulation of claim 15, wherein the formulation is for providing a 7-OH-cannabidiol AUC(0-t) of at least 4 times greater upon administration to a subject in a fed condition than upon administration to a subject in a fasted condition.
22. The formulation of claim 15, wherein the flavoring agent is strawberry flavor.
23. The formulation of claim 15, wherein the sweetener is saccharin.
24. The formulation of claim 15, wherein the formulation is substantially free of ethanol.

FIG. 1




FIG. 2





FIG. 2 (cont.)




FIG. 3




FIG. 3 (cont.)





FIG. 3 (cont.)






FIG. 3 (cont.)
Day 9 Combo



Day 14 Combo




FIG. 3 (cont.)




FIG. 4




FIG. 5





FIG. 6





FIG. 7




FIG. 8




FIG. 9








FIG. 10



CBD



$\Delta 9$ -THC



CBD+ $\Delta 9$ THC

