

(19)



(11)

EP 2 759 298 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

30.07.2014 Bulletin 2014/31

(51) Int Cl.:

A61K 31/4523 (2006.01)

A61K 31/4748 (2006.01)

A61P 25/00 (2006.01)

A61P 25/02 (2006.01)

A61P 1/00 (2006.01)

(21) Application number: **14165604.1**

(22) Date of filing: **10.10.2012**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

Designated Extension States:

BA ME

(30) Priority: **18.10.2011 US 201161548514 P**

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:

12784677.2 / 2 744 497

(71) Applicant: **Helsinn Healthcare SA
6912 Lugano-Pazzallo (CH)**

(72) Inventors:

- Pietra, Claudio
22100 Como (IT)**
- Cantoreggi, Sergio
6955 Cagiallo (CH)**

(74) Representative: **Zardi, Marco**

M. Zardi & Co. SA

Via Pioda 6

6900 Lugano (CH)

Remarks:

This application was filed on 23-04-2014 as a divisional application to the application mentioned under INID code 62.

(54) Therapeutic combinations of netupitant and palonosetron

(57) Combinations of netupitant and palonosetron, and methods of using such combinations to treat various pain states and irritable bowel syndrome, are provided.

EP 2 759 298 A1

DescriptionFIELD OF THE INVENTION

5 **[0001]** The present invention relates to combinations of netupitant and palonosetron, and to methods of treating pain and irritable bowel syndrome (IBS) using such combinations.


BACKGROUND OF THE INVENTION

10 **[0002]** Palonosetron is a selective 5-HT3 antagonist marketed commercially as Aloxi® for the treatment of emesis. The chemical name of the compound is (3aS)-2-[(S)-1-Azabicyclo [2.2.2]oct-3-yl]- 2,3,3a,4,5,6-hexahydro-1-oxo-1Hberiz[de] isoquinoline, as depicted by the following chemical structure of the hydrochloride salt:



[0003] Methods of synthesizing palonosetron are described in U.S. Patent Nos. 5,202,333 and 5,510,486. Pharmaceutically acceptable dosage forms are described in PCT publications WO 2004/067005 and WO 2008/049552 from 25 Helsinn Healthcare.

[0004] Netupitant is a selective NK₁ receptor antagonist of the formula 2-[3,5-bis(trifluoromethyl)phenyl]-N,2-dimethyl-N-[4-(2-methylphenyl)-6-(4-methylpiperazin-1-yl)pyridin-3-yl]propanamide, or Benzeneacetamide, N,α,α-trimethyl-N-[4-(2-methylphenyl)-6-(4-methyl-1-piperazinyl)-3-pyridinyl]-3,5-bis(trifluoromethyl)-, having the below chemical structure:



[0005] Methods of synthesizing and formulating netupitant and its prodrugs are described in U.S. Patent Nos. 6,297,375, 6,719,996 and 6,593,472 to Hoffmann La Roche.

[0006] Several prior art references disclose the combined use of an NK1 receptor antagonist and a 5-HT3 receptor antagonist for the treatment of emesis. Roila *et al.* report that the co-administration of a NK1 receptor antagonist, such as aprepitant, at the same time as a 5-HT3 antagonist, significantly increases the efficacy of 5-HT3 antagonists in 45 preventing both acute and delayed CINV. See Roila F, Fatigoni S (2006) NEW ANTIEMETIC DRUGS. Ann Oncol 17 Suppl 2: ii96-100. Roche reported in 2006 that "[s]ince 5HT3 and NK1 receptor antagonists exert additive effects [on nausea and vomiting] there is considerable potential for the combined use of Aloxi and netupitant." See NK1 receptor antagonists by Roche, Feb. 23 2006, http://www.hospitalpharma.com/Features/feature.asp?ROW_ID=742.

[0007] Palonosetron and other 5HT3 receptor antagonists were developed originally for the prevention of emesis, but 50 more recently have received attention for their role in pain signaling and transmission in the peripheral and central nervous systems. 5HT3 receptors are known to mediate a descending facilitatory influence on spinal cord activity, a constituent drive that is particularly prominent on mechanical and chemical evoked activity. There is enhancement of this activity following peripheral nerve injury, spinal cord injury and after intense chemical stimulation. A number of studies using opioid-induced-hyperalgesia and activation of ERK support the concept of descending 5HT3 facilitation as a target 55 for pain control. See, e.g., GH McCleane, *et al.*, ANETH ANALG 2003; 97: 1474-8 (reporting that "5HT3 receptors play a pronociceptive role and mediate descending excitatory controls that allow spinal neurons to fully code peripheral stimuli.")

[0008] Spinal NK1 receptors are part of the post-synaptic targets for neurotransmitter released from afferent terminals,

and also have been studied for their role in pain. See De Felipe C, et al. (March 1998) NATURE 392 (6674): 394-7 (reporting that "the peptide neurotransmitter substance P modulates sensitivity to pain by activating the neurokinin-1 (NK-1) receptor, which is expressed by discrete populations of neurons throughout the central nervous system.") LM Thomson et al. (2008 January) J PAIN; 9(1): 11-19 (reporting that substance P systems "may also represent important therapeutic targets for the retention and restoration of pain relief with prolonged morphine treatment.")

5 [0009] Only about 15% of afferents release substance P and this may be reduced after nerve injury although some studies report a phenotypic switch of the transmitter to large fibers. Thus, while several prior art references implicate NK1 receptors in the pain signaling process, considerable doubt has emerged as to the viability of NK1 receptor antagonists to successfully treat pain in human patients. As Hill stated in 2000, "NK1 receptor antagonists have failed to exhibit 10 efficacy in [human] clinical trials of a variety of clinical pain states." See R Hill (July 2000) TRENDS IN PHARMACEUTICAL SCIENCES Vol. 21, pages 244-246. So, the target alone might not be relevant in controlling the pain processing.

10 [0010] Irritable bowel syndrome (IBS), a functional bowel disorder, is a syndrome characterized by abdominal discomfort or pain associated with defecation and abnormal bowel movement in spite of the absence of a detectable intestinal 15 organic disease. The symptoms include diarrhea, abdominal pain, abdominal bloating, and constipation, and are classified into a diarrhea type (IBS-D), a constipation type (IBS-C), and an alternating diarrhea and constipation type (IBS-A). The symptoms may be accompanied by a psychological condition such as anxiety, hypersensitivity, tension, fretfulness, depression, or the like.

20 [0011] The precise pathophysiology of IBS remains to be elucidated. Nevertheless, there is a heightened sensitivity to visceral pain perception, known as peripheral sensitization. This sensitization involves a reduction in the threshold and an increase in the gain of the transduction processes of primary afferent neurons, attributable to a variety of mediators including monoamines (e.g., catecholamines and indoleamines), substance P, and a variety of cytokines and prostanoids such as E-type prostaglandins (see, e.g., Mayer et al., Gastroenterol., 107:271-293 (1994)). Also implicated in the 25 etiopathology of IBS is intestinal motor dysfunction, which leads to abnormal handling of intraluminal contents and/or gas (see, e.g., Kellow et al., Gastroenterol., 92:1885-1893 (1987); Levitt et al., Ann. Int. Med., 124:422-424 (1996)). Psychological factors may also contribute to IBS symptoms appearing in conjunction with, if not triggered by, disturbances 30 including depression and anxiety (see, e.g., Drossman et al., Gastroenterol. Int., 8:47-90 (1995)).

35 [0012] Although no animal models of IBS exist, advances in the understanding of the pathophysiology of IBS have facilitated the development of preclinical rodent models of visceral hypersensitivity. The method of colorectal distention (CRD) can be used to activate nociceptive neuronal pathways from the gastrointestinal (GI) tract, which induces a protective reflex of abdominal muscle contraction (see, e.g., Ness et al., Brain Research, 450: 153-169 (1988)). One method of measuring these contractions is via strain gauges sutured onto the abdominal muscle (see, e.g., Plourde et 40 al., American Journal of Physiology, 273: G191-196 (1997)). Colonic hypersensitivity can be induced experimentally in a rodent model by infusing dilute acetic acid (0.6%) into the colon which causes a transient sensitization of colonic sensory afferents which results in an increase in the VMR to CRD (see e.g., Gaudreau et al., Neuroscience Letters, 351 (2): 59-62 (2003); Venkova et al., Toxicology and Applied Pharmacology, 196: 215-222 (2004)). Although heightened 45 visceral pain perception is well demonstrated in IBS, the existence of abnormalities in somatic pain is controversial with more recent reports suggesting enhanced somatic sensitivity in IBS through viscero-somatic convergence at the level of the spinal cord (see e.g., Chang, Gastroenterology Clinics of North America, 34 (2): 271-2792005; Zhou et al., Pain, 148 (3): 454-461 (2010)).

50 [0013] The gastrointestinal function is highly regulated by the nerves, and a variety of receptors are present. Therefore, an anticholinergic agent, an antidiarrheal, or a laxative is administered according to the gastrointestinal symptoms such as an abdominal pain, diarrhea, and constipation, and an antidepressant or an antianxiety agent is used if needed. Although alosetron HCl, a serotonin 5-HT₃ receptor antagonist, is known as a therapeutic agent for diarrhea-type IBS (IBS-D), this agent is applied only to woman patients with severe symptoms because serious gastrointestinal disorders, particularly ischemic colitis and serious constipation, are observed. Further, although tegaserod maleate, a serotonin 5-HT₄ receptor agonist, is known as a therapeutic agent for constipation-type IBS (IBS-C), this agent is applied only to 55 women patients. In addition, lubiprostone, a ClC-2 chloride channel activator, is applied only to women patients for constipation-type IBS (IBS-C). Further, Cilansetron, another selective 5-HT₃ antagonist, is currently in clinical trials in Europe for the treatment of IBS-D in both men and women; however, in 2005, the sponsor of cilansetron withdrew its application for approval from the US FDA on the basis of a "non-approvable" letter (additional clinical trials were requested). Therefore, the therapeutic options of IBS are limited as the current treatments are gender and IBS subtype specific, and have significant side effects. Therefore, a novel therapeutic agent and method for IBS with no gender or subtype difference and with reduced side effects is strongly desired.

55 SUMMARY OF THE INVENTION

56 [0014] Consistent with other human studies reported in the prior art, applicant has determined that netupitant has a very marginal effect on the treatment of neuropathic pain. In particular, netupitant exhibits zero or minimal effect when

tested in animal models of allodynia / neuropathic pain. However, when the netupitant is added to palonosetron in therapeutically effective amounts, the netupitant acts synergistically with the palonosetron, and produces a much greater reduction in pain than when the palonosetron is administered alone.

[0015] The opposite effect is observed when the compounds are evaluated against various electrophysiological measures of pain control. In the course of these experiments, the inventors have discovered that netupitant has an independent effect, and that palonosetron exhibits zero or minimal effect. However, when the palonosetron is added to netupitant in therapeutically effective amounts, the palonosetron acts synergistically with the netupitant, and induces much greater control over electrophysiological measures of pain than when the netupitant is administered alone.

[0016] Therefore, in one embodiment, the invention provides a synergistic drug combination comprising (a) palonosetron or a pharmaceutically acceptable salt or prodrug thereof; and (b) netupitant or a pharmaceutically acceptable salt or prodrug thereof, in synergistically effective amounts. Various measures can be used to determine whether the amounts in the drug combination are synergistic. In a preferred embodiment, the amounts are synergistically effective when administered to a 70 kg human based on one or more measures selected from (a) the treatment of mechanically evoked allodynia, (b) the modulation of electrically-evoked dorsal horn neural C-fiber responses, and (c) the modulation of electrically-evoked dorsal horn neural AD-fiber responses. An orally-administered combination drug product that contains from 0.01 to 1.0 mg of palonosetron and from 10 to 300 mg of netupitant has been shown to demonstrate synergy under one or more of these criteria.

[0017] In another embodiment, the invention is based on the discovery that netupitant and palonosetron work together to treat chronic neuropathic pain, and that they produce a greater therapeutic effect when used together than when either is administered alone. Therefore, the invention also provides a method of treating chronic neuropathic pain comprising administering to a human patient in need thereof a drug combination comprising palonosetron and netupitant, or their pharmaceutically acceptable salts or prodrugs, in therapeutically effective amounts. The combination is preferably administered orally, and preferably includes from 0.01 to 1.0 mg of palonosetron and from 10 to 300 mg of netupitant administered on a daily basis. In a preferred embodiment, the method is used to treat diabetic peripheral neuropathic pain, fibromyalgia, or post-herpetic neuralgia.

[0018] Further, applicants have determined that palonosetron, netupitant, or a combination of palonosetron and netupitant is effective in treating pain associated with Irritable bowel syndrome (IBS). Specifically, applicants have determined that a combination of a sub-therapeutic dose of palonosetron and a sub-therapeutic dose of netupitant has a synergistic effect in treating a model of Irritable bowel syndrome. Therefore, in one embodiment, the invention provides a synergistic drug combination comprising (a) a sub-therapeutic dose of palonosetron or a pharmaceutically acceptable salt or prodrug thereof; and (b) a sub-therapeutic dose of netupitant or a pharmaceutically acceptable salt or prodrug thereof, in synergistically effective amounts. In another embodiment, the invention provides a method of treating Irritable bowel syndrome comprising administering to a human patient in need thereof a synergistic effective combination comprising (a) a sub-therapeutic dose of palonosetron or a pharmaceutically acceptable salt or prodrug thereof and (b) a sub-therapeutic dose of netupitant or a pharmaceutically acceptable salt or prodrug thereof, in synergistically effective amounts. The drug combinations and methods disclosed in the present invention have advantages over the prior art, e.g., they are not gender specific or IBS-subtype specific, and less side effects are associated with the drug combinations and methods.

BRIEF DESCRIPTION OF THE DRAWINGS

[0019]

FIG. 1 is a series of graphs depicting the temporal development of mechanical and thermal hypersensitivity post nerve injury. Nerve injury induced a significant increase in the number of paw withdrawals to A) vf2g, B) vF 6g and C) acetone stimulation of the ipsilateral hind paw compared with the responses seen in sham controls.

FIG. 2 is a series of graphs depicting the effects of palonosetron on the number of paw withdrawals to A) vf2g, B) vF 6g and C) acetone. A dose related reduction in paw withdrawal frequency is seen in the SNL rats only.

FIG. 3 is a graph comparing the total time spent by SNL and Sham rats on a rotarod. Palonosetron (0.03, 0.3 and 3mg/kg) did not impair motor performance in either group. Data are presented as mean \pm SEM.

FIG. 4 is a series of graphs comparing the effects of Palonosetron (0.03, 0.3 and 3mg/kg s.c.) on the electrical evoked responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 5 is a series of graphs comparing of the effects of Palonosetron (0.03, 0.3 and 3mg/kg s.c.) on the dynamic brush and mechanical punctate evoked responses of spinal dorsal horn neurones in sham and SNL rats. Data are

expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 6 is a series of graphs comparing the effects of Palonosetron (0.03, 0.3 and 3mg/kg s.c.) on the thermal evoked responses of spinal dorsal horn neurones in sham and SNL. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 7 is a series of graphs depicting the effects of netupitant on the number of withdrawal responses to A) vF2g, B) vF6g and C) acetone (cooling stimuli). In SNL rats, 1mg/kg appears to inhibit behavioral hypersensitivities.

FIG. 8 is a series of graphs comparing the effects of 3 doses of netupitant (0.1, 1 and 10mg/kg s.c) on the electrical responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 9 is a series of graphs comparing the effects of 3 doses of netupitant (0.1, 1 and 10mg/kg s.c) on the dynamic brush and mechanical punctate evoked responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 10 is a series of graphs comparing the effects of 3 doses of netupitant (0.1, 1 and 10mg/kg s.c) on the thermal evoked responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 11 is a series of graphs comparing the effects of a combination of palonosetron (0.03 mg/kg s.c) and netupitant (0.1 mg/kg s.c.) on the number of withdrawal responses in the ipsilateral and contralateral paws to A) vF2g, B) vF6g and C) acetone (cooling stimuli) in SNL rats.

FIG. 12 is a series of graphs comparing the effects of a combination of palonosetron (0.03 mg/kg s.c) and netupitant (0.1 mg/kg s.c.) on the electrical responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 13 is a series of graphs comparing the effects of a combination of palonosetron (0.03 mg/kg s.c) and netupitant (0.1 mg/kg s.c.) on the dynamic brush, mechanical punctate and heat evoked responses of spinal dorsal horn neurones in SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.

FIG. 14 shows that Palonosetron and Netupitant dosed in combination at 0.001 mg/kg showed a non-significant trend to inhibit AA-induced somatic hypersensitivity.

FIG. 15 is a series of graphs where visceral sensitivity results presented as individual distension pressure histograms with accompanying scatter-plot comparisons. Palonosetron or Netupitant dosed at 0.001 mg/kg p.o alone or in combination.

DETAILED DESCRIPTION OF THE INVENTION

[0020] The present invention may be understood more readily by reference to the following definitions and detailed description of preferred embodiments of the invention and the non-limiting Examples included therein.

[0021] The pharmaceutically acceptable salts of netupitant and palonosetron, as well as their pharmaceutically acceptable prodrugs can be used in the methods and compositions of the present invention. As used herein, the term "pharmaceutically acceptable salt" refers to a salt of a compound to be administered prepared from pharmaceutically acceptable non-toxic acids. Examples of suitable inorganic acids are hydrochloric, hydrobromic, hydroiodic, nitric, sulfuric, and phosphoric. Suitable organic acids may be selected from aliphatic, aromatic, carboxylic and sulfonic classes of organic acids, examples of which are formic, acetic, propionic, succinic, camphorsulfonic, citric, fumaric, gluconic, isethionic, lactic, malic, mucic, tartaric, para-toluenesulfonic, glycolic, glucuronic, maleic, furoic, glutamic, benzoic, anthranilic, salicylic, phenylacetic, mandelic, embonic (pamoic), methanesulfonic, ethanesulfonic, pantothenic, benzenesulfonic (besylate), stearic, sulfanilic, alginic, galacturonic, and the like.

[0022] The term "prodrug" refers to a chemical derivative of an active ingredient that degrades to the active ingredient in vivo via spontaneous or enzymatic transformation. Prodrugs are designed to overcome pharmaceutically and/or pharmacokinetically based problems associated with the parent drug molecule that might otherwise limit the clinical usefulness of the drug, especially due to drug solubility issues in the case of netupitant. Prodrugs of netupitant are described in U.S. Patent No. 6,593,472, the contents of which are hereby incorporated by reference.

[0023] When dose amounts are expressed herein in reference to a salt or prodrug of an active ingredient, it will be understood that the amount expressed is based on the corresponding amount of the free base of the ingredient. Thus, for a prodrug with a molecular weight of 600, if this document referred to the administration of 100 mg of netupitant or a prodrug thereof, it would be understood that 125.36 mg of the prodrug is administered, since the molecular weight of the prodrug is 125.36% of the molecular weight of the netupitant base.

[0024] As used herein, "therapeutically effective amount" refers to an amount sufficient to elicit the desired biological response. The therapeutically effective amount or dose will depend on the age, sex and weight of the patient, and the current medical condition of the patient. The skilled artisan will be able to determine appropriate dosages depending on these and other factors in addition to the present disclosure.

[0025] The terms "treating" and "treatment," when used herein, refer to the medical management of a patient with the intent to cure, ameliorate, stabilize, or prevent a disease, pathological condition, or disorder. This term includes active treatment, that is, treatment directed specifically toward the improvement of a disease, pathological condition, or disorder, and also includes causal treatment, that is, treatment directed toward removal of the cause of the associated disease, pathological condition, or disorder. In addition, this term includes palliative treatment, that is, treatment designed for the relief of symptoms rather than the curing of the disease, pathological condition, or disorder; preventative treatment, that is, treatment directed to minimizing or partially or completely inhibiting the development of the associated disease, pathological condition, or disorder; and supportive treatment, that is, treatment employed to supplement another specific therapy directed toward the improvement of the associated disease, pathological condition, or disorder.

[0026] As used herein, the term "significantly" refers to a level of statistical significance. The level of statistical significant can be, for example, of at least $p < 0.05$, of at least $p < 0.01$, of at least $p < 0.005$, or of at least $p < 0.001$. When a measurable result or effect is expressed or identified herein, it will be understood that the result or effect can be evaluated based upon its statistical significance relative to a baseline, typically placebo based.

[0027] Throughout the description and claims of this specification, the word "comprise" and variations of the word, such as "comprising" and "comprises," means "including but not limited to," and is not intended to exclude, for example, other additives, components, integers or steps.

[0028] When the singular forms "a," "an" and "the" or like terms are used herein, they will be understood to include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a pharmaceutical carrier" includes mixtures of two or more such carriers, and the like. The word "or" or like terms as used herein means any one member of a particular list and also includes any combination of members of that list.

[0029] When ranges are given by specifying the lower end of a range separately from the upper end of the range, it will be understood that the range can be defined by selectively combining any one of the lower end variables with any one of the upper end variables that is mathematically possible.

[0030] When used herein the term "about" or "ca." will compensate for variability allowed for in the pharmaceutical industry and inherent in pharmaceutical products, such as differences in product strength due to manufacturing variation and time-induced product degradation. The term allows for any variation which in the practice of pharmaceuticals would allow the product being evaluated to be considered bioequivalent to the recited strength of a claimed product.

Methods of Treatment

[0031] As discussed above, it has now been discovered that palonosetron and netupitant are therapeutically effective, and synergistically effective under some circumstances, to treat pain, particularly chronic neuropathic pain. Therefore, in one embodiment, the invention provides a synergistic drug combination comprising (a) palonosetron or a pharmaceutically acceptable salt or prodrug thereof; and (b) netupitant or a pharmaceutically acceptable salt or prodrug thereof, in synergistically effective amounts.

[0032] In another embodiment, the invention is based on the discovery that netupitant and palonosetron work together to treat neuropathic pain, and that they produce a greater therapeutic effect when used together than when either is administered alone. Therefore, the invention also provides a method of treating neuropathic pain comprising administering to a human patient in need thereof a drug combination comprising palonosetron and netupitant, in therapeutically effective amounts.

[0033] While the methods can be used to treat various forms of pain, including nociceptive and neuropathic pain, chronic or acute, in a preferred embodiment the methods are used to treat chronic neuropathic pain (e.g. pain persisting for more than three, six or twelve months). In an even more preferred embodiment, the pain is chronic neuropathic pain characterized by one, two or three or more symptoms selected from burning pain, paresthesia / dysesthesia, shooting / lancinating pain, numbness, hyperalgesia and allodynia.

[0034] The pain can be inherited, or it may be acquired. Acquired peripheral neuropathies are grouped into three broad categories: those caused by systemic disease, those caused by trauma from external agents, and those caused by infections or autoimmune disorders affecting nerve tissue, and each of these categories can be treated using the methods of the current invention.

[0035] Causes of acquired peripheral neuropathy that can be treated using the methods of this invention include physical injury (trauma) to a nerve, tumors, toxins, autoimmune responses, nutritional deficiencies, alcoholism, and vascular and metabolic disorders. Drugs known to induce neuropathies treatable by the current methods include anti-virals such as ddl and ddC, phenytoin, isoniazid, vincristine, high dose vitamins, and folic acid antagonists.

5 [0036] The neuropathic pain treated using the methods of this invention can also be characterized as peripheral or central. Peripheral neuropathies that can be treated by the methods of the current invention include those originating from diabetes, post herpetic neuralgia, fibromyalgia, and physical injuries to the nervous system such as spinal cord injury.

10 [0037] Diabetic peripheral neuropathic pain (DPNP) is a preferred indication for the methods of the current invention. DPNP may result from type I or type II diabetes mellitus, and is preferably defined by a diagnosis of painful distal symmetrical sensorimotor polyneuropathy; and a pain score ≥ 4 on an eleven point scale ranging from zero (no pain) to 10 (worst possible pain)).

15 [0038] Fibromyalgia is another preferred indication for the methods of the current invention. This disease is preferably defined as a history of widespread pain for three months, and pain present at 11 or more of the 18 specific tender point sites on digital palpation with an approximate force of 4 kg., as defined by the American College of Rheumatology criteria for fibromyalgia. See Wolfe F et al. ARTHRITIS RHEUM 1990;33:160-72. The 18 sites include the occiput (bilateral, at the suboccipital muscle insertions), the low cervical (bilateral, at the anterior aspects of the intertransverse spaces at C5-C7), the trapezius (bilateral, at the midpoint of the upper border), the supraspinatus (bilateral, at origins, above the scapula spine near the medial border), the second rib (bilateral, at the second costochondral junctions, just lateral to the junctions on upper surfaces), the lateral epicondyle (bilateral, 2 cm distal to the epicondyles), the gluteal (bilateral, in upper outer quadrants of buttocks in anterior fold of muscle), the greater trochanter (bilateral, posterior to the trochanteric prominence), and the knee (bilateral, at the medial fat pad proximal to the joint line).

20 [0039] In a distinct embodiment, the methods of the present invention are used to treat neuropathic pain characterized by pain at one or a combination of the foregoing specific tender point sites.

25 [0040] Yet another preferred indication for the methods of the current invention is post-herpetic neuralgia, which may be defined as neuralgia persisting for at least three months following healing of herpes zoster rash.

30 [0041] A preferred measure of the success of any treatment is a statistically significant improvement in the number of patients having at least a 50% improvement in pain score from baseline, relative to placebo. The pain score is preferably derived from an eleven point scale ranging from zero (no pain) to 10 (worst possible pain)).

35 [0042] Several other features can also be used to characterize the pain or patients treated by the methods of this invention, regardless of the type or source of pain, or the clinical indication involved. Thus, in one embodiment the pain or patient is unresponsive to one or more analgesic drugs selected from opioid analgesics, non-steroidal anti-inflammatory drugs (NSAIDs), tricyclic antidepressants (TCA) such as amitriptyline, and anticonvulsants such as gabapentin, phenytoin, and carbamazepine. In another embodiment, the pain is accompanied by a comorbid diagnosis of major depressive disorder (MDD), anxiety and/or depression.

40 [0043] Other analgesics can be administered as part of the methods of the current invention, including acetaminophen (up to 4 g/day), an opioid analgesic, or an NSAID.

45 [0044] The dosage can vary within wide limits and will, of course, be fitted to the individual requirements in each particular case. In general, in the case of oral administration a daily dosage of palonosetron of from about 0.01 or 0.05 to about 0.5 or 2.0 mg per person should be appropriate, alternatively from about 0.01 or 0.05 to about 1.0 mg., and alternatively from about 0.025 mg to about 0.5 mg. The dose can be administered in a single daily dosage, due to the excellent pharmacokinetics observed for the molecule, although multiple doses could also be administered (i.e. two or three times daily). With respect to palonosetron, these doses should be substantially the same whether administered orally or via injection.

50 [0045] In the case of oral administration of netupitant, a daily dosage of netupitant of from about 10 to about 300 mg should be appropriate, preferably from about 20 to about 200 mg, or from about 30 to about 150 mg. Once again, it is preferred to administer this dose just once daily, although it is possible to divide the daily dose into two or three doses. In the case of injection, the dose would constitute about 40% of the oral dose, and the foregoing ranges can be adjusted to 40%.

55 [0046] Various measures can be used to determine whether the amounts in the drug combination are synergistic. In a preferred embodiment, the amounts are synergistically effective when administered once daily to a 70 kg human based on one or more measures selected from (a) the treatment of mechanically evoked allodynia, (b) the modulation of electrically-evoked dorsal horn neural C-fiber responses, and (c) the modulation of electrically-evoked dorsal horn neural AD-fiber responses. Combination drug products that contain the amounts of palonosetron and netupitant described herein have been shown to demonstrate synergy under one or more of these criteria.

[0047] Further, it has now been discovered that palonosetron, netupitant or a combination of palonosetron and netupitant are effective to treat irritable bowel syndrome (IBS). In a particularly preferred embodiment, applicants have determined that a combination of a sub-therapeutic dose of palonosetron and a sub-therapeutic dose of netupitant has a synergistic effect in treating Irritable bowel syndrome. Sub-therapeutic dose refers to a dose which is ineffective against

IBS when administered alone. Therefore, in one embodiment, the invention provides a method of treating Irritable bowel syndrome comprising administering to a human patient in need thereof a synergistically-effective combination comprising (a) a sub-therapeutic dose of palonosetron or a pharmaceutically acceptable salt or prodrug thereof; and (b) a sub-therapeutic dose of netupitant or a pharmaceutically acceptable salt or prodrug thereof.

5 [0048] In another embodiment, the above disclosed methods are effective in treating one or more indications selected from the group consisting of alternating IBS (IBS-A), IBS with diarrhea (IBS-D) and IBS with constipation (IBS-C). The primary symptoms of said IBS-C are abdominal pain and constipation. Constipation can be evaluated by accessing stool frequency. The baseline of the abdominal pain can be defined as weekly average of worst abdominal pain in past 24 hours score of equal to or greater than 3.0 in a 0 to 10 point scale. The baseline of stool frequency can be defined as less than three complete spontaneous bowel movements (CSBM) per week. It has been discovered that the above disclosed methods can (1) decrease in weekly average of worst abdominal pain in past 24 hours score of equal to or greater than 30% compared with baseline; and (2) increase of one or more CSBM per week compared with baseline. Therefore, in another embodiment, the invention provides a method of treating Irritable bowel syndrome with constipation (IBS-C) where said method is effective in (1) decreasing intensity of said abdominal pain equal to or greater than 30 percent based on weekly average of worst abdominal pain in past 24 hours as compared with baseline; and (2) increasing at least one complete spontaneous bowel movements (CSBM) per week as compared with baseline.

10 [0049] The primary symptoms of IBS with diarrhea (IBS-D) are abdominal pain and diarrhea. Diarrhea can be evaluated by accessing stool consistency. Stool consistency is described according to the Bristol Stool Scale, which is a medical aid designed to classify the form of human faeces into seven categories. The seven types of stool according to the Bristol 15 Stool Scale are: Type 1: Separate hard lumps, like nuts (hard to pass); Type 2: Sausage-shaped, but lumpy; Type 3: Like a sausage but with cracks on its surface; Type 4: Like a sausage or snake, smooth and soft; Type 5: Soft blobs with clear cut edges (passed easily); Type 6: Fluffy pieces with ragged edges, a mushy stool; and Type 7: Watery, no 20 solid pieces. Entirely liquid. The baseline of the abdominal pain can be defined as weekly average of worst abdominal pain in past 24 hours score of equal to or greater than 3.0 in a 0 to 10 point scale. The baseline of stool consistency can 25 be defined as weekly average equal to or greater than Type 6 Bristol Stool Scale (BSS). It has been discovered that the above disclosed methods can (1) decrease the weekly average of worst abdominal pain in past 24 hours by 30% or more compared with baseline; and (2) achieve a weekly average of equal to or less than Type 5 Bristol Stool Scale (< Type 2 BSS can be considered an adverse event). Therefore, in another embodiment, the invention provides a method 30 of treating Irritable bowel syndrome with diarrhea (IBS-D) where said method is effective in (1) decreasing the intensity of abdominal pain by 30 percent or more based on a weekly average of worst abdominal pain in the previous 24 hours as compared with baseline; and (2) achieving a weekly average of equal to or less than Type 5 Bristol Stool Scale (< Type 2 BSS can be considered an adverse event).

35 [0050] As discussed above, the disclosed methods are non-specific to IBS subtypes. Further, it has been discovered that the above disclosed methods can be applied to both male and female patients. Therefore, in another embodiment, the invention provides a method of treating Irritable bowel syndrome wherein said method is not gender specific. Moreover, since the disclosed methods administer a combination of palonosetron and netupitant to patients where each agent is in sub-therapeutic dose, said methods significantly reduce the potential side effects. Therefore, in another embodiment, the invention provides a method of treating Irritable bowel syndrome wherein associated side effects are significantly reduced.

40 [0051] In another embodiment, the invention provides a method of treating Irritable bowel syndrome where said method is effective to treat other symptoms of IBS-C, e.g., abdominal discomfort, and other symptoms of IBS-D, e.g., abdominal discomfort and fecal incontinence. In still another embodiment, the invention provides a method of treating Irritable bowel syndrome (IBS) wherein said IBS is caused by colonic and/or somatic hypersensitivity. In another embodiment, the invention provides a method of treating Irritable bowel syndrome (IBS) wherein said method is effective in inhibiting said 45 colonic and/or somatic hypersensitivity.

50 [0052] In another embodiment, the invention provides a method of treating Irritable bowel syndrome, comprising administering a palonosetron dose of less than 0.01 mg per kg body weight and a netupitant dose of less than 0.01 mg per kg body weight to a patient in need thereof. In still another embodiment, the invention provides a method of treating Irritable bowel syndrome, comprising administering a palonosetron dose of about 0.001 mg per kg body weight and a netupitant dose of about 0.001 mg per kg body weight to a patient in need thereof.

Pharmaceutical Compositions

55 [0053] Various pharmaceutical compositions can be developed that make use of the combinations described herein. The composition can be administered by any appropriate route, for example, orally, parenterally, or intravenously, in liquid or solid form.

[0054] Preferred modes of administrations of the active compounds are injectable and oral. These compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules (for oral use) or com-

pressed into tablets (for oral or buccal use) or formulated into troches (for buccal use). For these purposes, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition.

[0055] Tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a gliding such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring. When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil. In addition, dosage unit forms can contain various other materials which modify the physical form of the dosage unit, for example, coatings of sugar, shellac, or other enteric agents.

[0056] The compounds can be administered as a component of an elixir, suspension, syrup, wafer, orally disintegrating film, orally disintegrating tablet, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

[0057] Solutions or suspensions used for injection can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride, mannitol and dextrose. An injectable preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

EXAMPLES

[0058] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds claimed herein are made and evaluated, and are intended to be purely exemplary of the invention and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, temperature is in °C or is at room temperature, and pressure is at or near atmospheric.

EXAMPLE 1

[0059] Experiments were performed on male Sprague-Dawley rats, which were housed in cages under a 12-hour alternating light/dark cycle with *ad libitum* access to food and water. All animal experiments were approved by the United Kingdom Home Office and were carried out in accordance with guidelines set by personal and project licenses and thus complied with the UK Animals (Scientific Procedures) Act 1986. All efforts were made to minimise animal suffering and to reduce number of animals used.

SNL Surgery

[0060] A model of experimental neuropathic pain with symptoms of allodynia and hyperalgesia was established by tightly ligating the L5 and L6 spinal nerves as previously described (Kim and Chung, 1992). Within a sterile theatre, rats weighing approximately 130-150g were anaesthetised (1:1 O₂:N₂O, 3% halothane for induction, 1% maintenance) and following a small left-side incision at approximately L4-S2, paraspinal muscle and fat was removed from spinous processes. Part of the L6 traverse process was clipped with rangeurs to expose the parallel-lying L4 and L5 spinal nerves. The L5 nerve was isolated and hooked with a finely-pulled glass rod and tightly tied with non-absorbable 6-0 silk thread distal to the dorsal root ganglion (and proximal to the formation of the sciatic nerve). The L6 nerve was then hooked from under the sacrum and tied in a similar way. Haemostasis was confirmed and the wound was sutured with 3-0 absorbable silk. The surrounding skin was pulled together and secured over the injury with wound clips. Rats recovered in an incubator, and once it had been confirmed that they had no observable motor impairment in their left hindpaw, were re-housed in cages as above. Sham surgery was performed in the same environment and under the same conditions except nerves were not hooked or ligated.

Behavioural Assessment of the Neuropathic State

[0061] On post-operative days 2, 7, 9 and 14, behavioural signs of punctate mechanical and cooling hypersensitivity were assessed in the left ipsilateral hindpaw relative to the right contralateral hindpaw in awake and alert rats. Rats were individually placed in clear acrylic cubes on an elevated floor of wire mesh, and following a period of acclimation (30

minutes), mechanical sensitivities were determined by paw withdrawal to von Frey (vF) filaments with bending forces 2g and 6g (tested in consecutive, ascending force). Each filament was applied to the plantar surface of the paw for ~2-3s with enough force to cause buckling, and for each animal this was repeated 10 times at set positions on each paw. The number of lifts in response to each of the filaments was noted for the contralateral and ipsilateral paw of each rat. Cold hypersensitivity was assessed by applying a drop of acetone to each paw 5 times during each testing period, with each application separated by an interval of at least 5 minutes. Withdrawal frequency was quantified as = (number of foot withdrawals /10 or 5 trials as appropriate).

5 [0062] The rotarod was also used to assess potential effects on motor performance of Netupitant and Palonosetron. Acceleration of the rotarod was set to increase from 0 to 20 revolutions per minute over the period of one minute. Due to initial, non-linear acceleration, rats were placed on the rotarod locked at a speed of 8 revolutions per minute. The rotarod was then set to accelerate and the latency to fall from the rotarod was timed from this point.

Electrophysiology

15 [0063] Electrophysiology experiments were conducted on post-operative days 14-17 in anaesthetised rats.

10 [0064] We recorded the evoked responses of dorsal horn lamina V-VI neurones to stimulation of the peripheral receptive field with:

- 20 ● a train of 16 electrical pulses at 3 x C-fibre threshold
- mechanical von Frey filaments of increasing force - vf2g, 8g, 26g and 60g applied to the centre of the receptive field for 10 seconds
- 25 ● heat - 40, 45 and 48 °C applied to the centre of the receptive field by a constant water jet for 10 seconds

25 [0065] Following 3 stable baseline responses, 0.03 mg/kg Palonosetron and 0.10mg/kg Netupitant were administered into the scruff of the neck by subcutaneous injection. Drug effects were followed for two hours with tests carried out at 10, 30, 50, 70, 90 and 110 minutes post injection.

30 *Explanation of Electrical Measurements*

35 [0066] Input represents the post-synaptic C-fibre-evoked dorsal horn neuronal response following the first of the 16 electrical stimuli in the electrical train. It gives a measure of resting pre-synaptic activity (taken as total neuronal responsiveness upstream of the neurone under study, therefore including afferent excitability, activity at terminals and interneurones) and transmitter release in the absence of potentiation. The baseline 'input' response given numerically and graphically in this and other studies, was calculated as the number of C-fibre-evoked action potentials produced by the first stimulus (i.e. initial baseline response) multiplied by the total number of stimuli. A β -, A δ -, and C-fibre responses denote the number of action potentials respectively generated in the spinal cord as a consequence of A β -, A δ -, and C-fibre activity following electrical stimulation of their peripheral receptive field. Our Spike 2 Data Capturing software 40 separates out which incoming action potentials are carried by which fibre according to their latency to arrive in the spinal cord following generation (given the different conduction velocities of the fibres). Thus, those arriving within 20ms of peripheral stimulation are attributed to A β -fibres, A δ -fibres = 20-90ms, and C-fibre-evoked responses were taken as those recorded 90-300ms after electrical stimulation. Neuronal responses that trail behind between 300-800ms post-stimulus are quantified as post-discharge, and represent spinal cord hyperexcitability that results as a consequence of 45 repetitive stimulation of the neurone. Activity-dependent hyperexcitability can additionally be measured as wind-up. This value is calculated as the difference between the total number of action potentials at C-fibre latency produced by the train of 16 electrical stimuli, and 'input' as defined and calculated above. Therefore, if the first electrical stimulus elicited 15 C-fibre-evoked action potentials, and the *total number* of C-fibre-evoked action potentials recorded after 16 stimuli was 350, then wind up = 350 - (15x10) = 200. If each post-synaptic response was independent of previous activity, then 50 in the example above, the cumulative number of action potentials evoked by C-fibres after the 16 electrical stimuli should theoretically be 150. However, temporal summation of action potentials, and post-synaptic hyperexcitability amplify responses, so wind-up is a measure of the additional action potentials recorded above the predicted baseline level.

55 *Statistical Analysis*

[0067] Analyses were performed using GraphPad Prism version 4 for Apple Macintosh OS 10.4, (GraphPad Software, USA), and for all data a 95% confidence interval was used as a measure of statistical significance. Statistical significance with respect to behavioural scores was calculated using non-parametric Wilcoxon matched pairs tests. For electrophys-

iological data, statistical analyses were performed on raw data using two-way analysis of variance (ANOVA) for responses to mechanical and thermal stimuli, and if significant, Bonferroni post-hoc tests were performed. For responses to electrical stimulation and brush, student's t-tests were used to compare pre-drug baseline values with post-drug values. The data are expressed and presented as mean \pm SEM. Significance was taken as P<0.05 *, P<0.01 **

5 [0068] The data is reproduced in Tables 1-3 herein, and Figures 1-13.

TABLE 1

		BEHAVIORAL ASSESSMENTS: Paw Withdrawal Frequency (PWD) in SNL Rats		
Dose (mg/kg s.c.)		2 g vF Filament	6 g vF Filament	Cooling Acetone
Baseline		5.0 \pm 1.2	6.7 \pm 1.1	2.9 \pm 0.6
0.03 Palo		2.8 \pm 0.8	5.2 \pm 1.5	1.2 \pm 0.4
0.3 Palo		3.1 \pm 0.7	4.1 \pm 1.3	1.2 \pm 0.5
3 Palo		1.4 \pm 0.6	1.1 \pm 0.4*	1.9 \pm 0.3
Baseline		4.4 \pm 1.3	6.7 \pm 1.2	2.8 \pm 0.5
0.1 Netu		4.1 \pm 1.0	6.7 \pm 1.8	1.9 \pm 0.3
1 Netu		3.8 \pm 1.2	6.2 \pm 1.5	2.7 \pm 0.5
10Netu		3.9 \pm 1.1	5.9 \pm 1.0	2.0 \pm 0.4
Baseline		4.1 \pm 0.3	7.8 \pm 0.5	3.6 \pm 0.3
0.1 Netu + 0.03 Palo		0.3 \pm 0.1**	1.8 \pm 0.5**	0.4 \pm 0.2 **
Numbers as Mean \pm S.E. (n=7-9) * p<0.05 ** p<0.01 vs baseline				

TABLE 2

	ELECTROPHYSIOLOGICAL STUDIES: Electrically-Evoked Dorsal Horn Neural Responses in SNL Rats - Number of Action Potentials						
Dose (mg/kg s.c.)		C-Fiber Response	AB-Fiber Response	AD-Fiber Response	Input Response	Post Discharge Response	Wind-up Response
Baseline		408 \pm 25	141 \pm 9	137 \pm 24	419 \pm 70	299 \pm 66	388 \pm 93
0.03 Palo		518 \pm 103	141 \pm 22	192 \pm 51	461 \pm 86	417 \pm 127	559 \pm 192
Baseline		403 \pm 30	140 \pm 7	142 \pm 24	415 \pm 65	225 \pm 48	230 \pm 56
0.1 mNetu		292 \pm 45	135 \pm 10	103 \pm 13	293 \pm 46	195 \pm 29	270 \pm 38
Baseline		301 \pm 43	116:1:13	131 \pm 17	244 \pm 45	182 \pm 31	273 \pm 45
0.1 Netu + 0.03 Palo		124 \pm 29**	104 \pm 15	46 \pm 13*	54 \pm 16**	87 \pm 37**	145 \pm 54
Numbers as Mean \pm S.E. (n=5-7) * p<0.05 ** p<0.01 vs baseline							

50

55

TABLE 3

ELECTROPHYSIOLOGICAL STUDIES: Electrically-Evoked Dorsal Horn Neural Responses in SNL Rats - Number of Action Potentials							
Dose (mg/kg s.c.)	Mechanically-Evoked			Heat-Evoked			Brush-Evoked
	8 g vF Filament	26 g vF Filament	60 g vF Filament	40° Celsius	45° Celsius	48° Celsius	
Baseline	181 ± 53	564 ± 19	772 ± 101	311 ± 138	638 ± 195	915 ± 137	118 ± 43
0.03 Palo	211 ± 42	539 ± 17	799 ± 192	136 ± 58	380 ± 187	816 ± 101	144 ± 59
Baseline	168 ± 51	385 ± 41	671 ± 105	473 ± 125	670 ± 159	971 ± 128	268 ± 35
0.1 Netu	190 ± 42	390 ± 56	486 ± 75	298 ± 72	381 ± 82	495 ± 186	310 ± 40
Baseline	122 ± 54	368 ± 106	578 ± 106	189 ± 67	472 ± 97	659 ± 142	343 ± 78
0.1 Netu + 0.03 Palo	30 ± 20	137 ± 41*	242 ± 54**	175 ± 42	186 ± 49**	396 ± 122**	199 ± 21

Numbers as Mean ± S.E. (n=5-7) * p<0.05 ** p<0.01 vs baseline

EXAMPLE 2

[0069] In a preferred embodiment the combination is administered in a capsule oral dosage form, wherein the capsule houses one or more soft-gel capsules for the palonosetron and one or more hard tablets for the netupitant. Table 4 below describes a representative formulation for a soft-gel capsule containing 0.5 mg. of palonosetron, suitable for inclusion in such a hard shell.

TABLE 4: REPRESENTATIVE SOFT-GEL FORMULATION

Ingredient	Approximate Amount (mg./Capsule)	Function
<i>Fill Solution</i>		
Palonosetron HCl	0.56 ¹	Active
Mono- and di-glycerides of Capryl/Capric Acid (Capmul MCM)	62.19	Solvent vehicle
Glycerin, anhydrous, USP/Ph Eur	3.37	Plasticizer
Polyglyceryl oleate (Plurol Oleique CC 497)	0.87	Surfactant
Purified water, USP/Ph Eur	2.94	Co-solvent
Butylated hydroxyanisole (BHA), NF/Ph Eur	0.07	Antioxidant
Nitrogen	-	
Theoretical fill weight	70.00 mg.	
<i>Gelatine Capsule Shell, 1.5-oval (Catalent Pharma Solutions)²</i>		
Gelatine (type 195), NF/Ph Eur	-	Shell
Sorbitol Special/Glycerin Blend 50/50	-	Plasticizer
Titanium dioxide, USP/Ph Eur	-	Colorant/Opacifier
Purified water, USP/Ph Eur	-	Solvent

1Corresponds to 0.50 mg. free base
2Quantitative composition of capsule shell is proprietary to Catalent Pharma Solutions

[0070] Table 5 below describes a representative formulation for a tablet containing 100 mg. of netupitant, suitable for inclusion in a hard shell.

TABLE 5: REPRESENTATIVE TABLET FORMULATION

5	Ingredient	Approximate Amount (mg./Tablet)	Function
10	Netupitant, milled	100	Active
15	Microcrystalline cellulose pH 101	20.5	Diluent and disintegrant
20	Sucrose Lauric Acid Esters	10.0	Surfactant
25	Polyvinilpyrrolidone K30	7.0	Binder
30	Sodium croscarmellose	3.0	Disintegrant
35	Colloidal Silicon Dioxide	3.0	Glidant
40	Sodium Stearyl Fumarate	1.0	Lubricant
45	Magnesium Stearate	0.5	Lubricant
50	Total weight	145 mg.	

EXAMPLE 3

[0071] Experiments were conducted to investigate the efficacy of Palonosetron (PT) and Netupitant (NT), individually and in combination, to inhibit acetic acid (AA) -induced colonic and somatic hypersensitivity in a rodent model. The overall goal was to determine if synergism would result when sub-therapeutic doses of both compounds were dosed in combination.

Materials and Methods

[0072] Animals: Male Sprague Dawley rats (330-480 g at time of colonic sensitivity assessment) were purchased from Charles River Laboratories. Rats were housed two-per-cage within the University of Oklahoma Health Sciences Center (OUHSC) Department of Comparative Medicine's animal facility under controlled temperature and humidity. All animals had free access to food and water and were acclimated to facility housing for a minimum of one week before experimentation. A total of 85 rats were used to complete this study. The experimental protocol was approved by the University of Oklahoma Health Sciences Center (OUHSC) Institutional Animal Care and Use Committee (IACUC Animal Protocol # 10-077).

[0073] Acclimation: Upon arrival all animals were acclimated to the animal facility for a minimum of one-week. To further acclimate and minimize experimental stress, the rats were brought to the laboratory for an additional week to acclimate to the laboratory environment and animal handlers.

[0074] Induction of Acute Visceral and Somatic Hypersensitivity: Visceral and somatic hypersensitivity were induced by infusing dilute (1.5 ml at 0.6%) acetic acid (AA) into the rat colon via a catheter (Intramedic PE 90 tubing) inserted via the anus to the level of the mid-colon. Within 60 mins colonic and somatic hypersensitivity was evident.

[0075] Colonic Sensitivity Assessment: Visceromotor responses (VMR) to colorectal distension (CRD) were measured by counting the number of abdominal contractions in response to increasing levels of CRD (0-60 mmHg). On the day of the colonic sensitivity assessment, a minor surgical procedure was performed to attach a strain gauge force transducer onto the abdominal oblique muscle and attached via an adapter cable to a Grass Model 7 Polygraph. The cable was connected to a Model 7P1 Low-Level DC Pre-Amp. The Pre-Amp was set at 0.02 mV/cm sensitivity and was connected to a Model 7DA DC Driver Amp with sensitivity set at 5.5 Amps. A 5 cm latex balloon was inserted into the distal colon. The balloon cannula was connected to a Distender Series IIR barostat (G & J Electronics Inc.) for controlled, isobaric inflation of the balloon and CRD using constant pressure (isobaric) tonic distensions were conducted at 0, 20, 40 and 60 mmHg. Each pressure was maintained for a period 10 min. during which time the number of abdominal muscle contractions were counted. A 10 min. recovery period was allowed between the each distension.

[0076] Somatic Sensitivity Assessment: Von Frey filaments were used to determine the level of somatic sensitivity in all animals receiving intracolonic infusion of AA. The Von Frey instrument measures the level of somatic sensitivity by recording the minimal force required to elicit hind paw withdrawal. The animals were placed in a mesh-bottomed Von Frey caging apparatus and a nylon filament was steadily pushed against the rat's foot pad until withdrawal of the hind paw. The force in grams which elicited hind paw withdrawal was then recorded.

[0077] Test Compounds: Following surgery, rats were pre-dosed with Palonosetron (1-hr pre-dose), Netupitant (2-hr

pre-dose) or methyl cellulose vehicle. Palonosetron and Netupitant were supplied by Helsinn Healthcare Inc. and were stored at 4° C until prepared for dosing as a suspension in 1% methyl cellulose solution. The drug suspensions were prepared so that each rat received an oral dosing volume of 0.5 ml per 100 g body weight for doses of 0.001, 0.01, 0.1 or 1 mg/kg. Compounds were prepared fresh on the day of use.

5 [0078] Statistical Analysis: To determine statistical significance between multiple control and treatment groups, data was compared using one-way ANOVA followed by a Bonferroni post-test. Results were deemed significant when p-values were less than 0.05.

Results

10 [0079] Effect of Palonosetron and Netupitant Combination on Somatic Hypersensitivity: As illustrated in figure 14, the oral administration of non-effective doses (0.001 mg/kg) of Palonosetron and Netupitant in combination resulted in a non-significant inhibitory trend toward normalization of somatic hypersensitivity.

15 [0080] The above experiments show that (1) infusion of dilute acetic acid into to the colon results in the development of both visceral and somatic hypersensitivity; (2) The 5-HT₃ antagonist Palonosetron significantly inhibited colonic and somatic hypersensitivity; (3) the NK₁ antagonist Netupitant significantly inhibited colonic sensitivity and somatic hypersensitivity; and (4) when non-effective doses (0.001 mg/kg) of Palonosetron and Netupitant were dosed in combination there was a significant inhibition of colonic sensitivity and a non-significant trend to inhibit somatic sensitivity. The above findings suggest that both Palonosetron and Netupitant could be further developed as possible therapeutic agents for 20 the treatment of abdominal pain and GI dysfunction associated with IBS. Furthermore, the synergic effect shown in the non-effective combination dosing experiments, suggest that combination therapy using these two compounds may result in relief of IBS symptoms with less potential for unfavorable side-effects.

25 [0081] Throughout this application, various publications are referenced. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains. It will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

30

Claims

1. A therapeutically effective amount of palonosetron or a pharmaceutically acceptable salt or prodrug thereof; or a therapeutically effective amount of netupitant or a pharmaceutically acceptable salt or prodrug thereof; or a drug combination comprising: (i) a sub-therapeutic dose of palonosetron or a pharmaceutically acceptable salt or prodrug thereof, wherein said sub-therapeutic dose is ineffective against irritable bowel syndrome when administered alone; and (ii) a sub-therapeutic dose of netupitant or a pharmaceutically acceptable salt or prodrug thereof, wherein said sub-therapeutic dose is ineffective against irritable bowel syndrome when administered alone, for use in a method of treating irritable bowel syndrome in a human patient in need thereof.

35 2. The amount or combination for the use according to claim 1, wherein said administration is oral, and said therapeutically effective amounts comprise, on a daily basis:

40 a) from 0.01 to 1.0 mg of palonosetron or a pharmaceutically acceptable salt or prodrug thereof, based on the weight of the free base; and

b) from 10 to 300 mg of netupitant or a pharmaceutically acceptable salt or prodrug thereof, based on the weight of the free base.

45 3. The amount or combination for the use according to claim 1, wherein said administration is oral, and said sub-therapeutic dose comprises:

50 a) palonosetron or a pharmaceutically acceptable salt thereof in a dose of less than 0.01 mg per kg body weight, based on the weight of the free base; and

b) netupitant or a pharmaceutically acceptable salt thereof in a dose of less than 0.01 mg per kg body weight, based on the weight of the free base.

4. The amount or combination for the use according to claim 1, wherein said administration is oral, and said sub-therapeutic dose comprises:
 - 5 a) palonosetron or a pharmaceutically acceptable salt thereof in a dose of less than 0.001 mg per kg body weight, based on the weight of the free base; and
 - b) netupitant or a pharmaceutically acceptable salt thereof in a dose of less than 0.001 mg per kg body weight, based on the weight of the free base.
- 10 5. The amount or combination for the use according to claims 1-4, wherein said IBS is selected from the group consisting of alternating IBS (IBS-A), IBS with diarrhea (IBS-D), and IBS with constipation (IBS-C).
- 15 6. The amount or combination for the use according to claims 1-5, wherein said method is non-gender specific.
7. The amount or combination for the use according to claims 1-6, wherein said IBS is IBS with constipation (IBS-C), and the method is effective to treat abdominal pain and constipation.
- 15 8. The amount or combination for the use according to claim 7, wherein said constipation is evaluated based on stool frequency.
- 20 9. The amount or combination for the use according to claim 8, wherein said method is effective to:
 - a) decrease the intensity of said abdominal pain by 30 percent or more based on the weekly average of worst abdominal pain in previous 24 hours score as compared with baseline; and
 - 25 b) increasing complete spontaneous bowel movements (CSBM) per week at least once per week as compared with baseline.
10. The amount or combination for the use according to claims 1-6, wherein said IBS is IBS with diarrhea (IBS-D) and said method is effective to treat abdominal pain and diarrhea.
- 30 11. The amount or combination for the use according to claim 10, wherein said diarrhea is evaluated by accessing stool consistency.
12. The amount or combination for the use according to claim 11, wherein said stool consistency is described according to Bristol Stool Scale.
- 35 13. The amount or combination for the use according to claim 12, wherein said method is effective in:
 - a) decreasing intensity of said abdominal pain by 30 percent or more based on weekly average of worst abdominal pain in previous 24 hours score as compared with baseline; and
 - 40 b) achieving a weekly average of equal to or less than type 5 of the Bristol Stool Chart.
14. The amount or combination for the use according to claim 7, wherein other symptoms of said IBS-C include abdominal discomfort.
- 45 15. The amount or combination for the use according to claim 10, wherein other symptoms of said IBS-D include abdominal discomfort and fecal incontinence.
16. The amount or combination for the use according to claims 1-15, wherein said IBS is caused by colonic and/or somatic hypersensitivity.
- 50 17. The amount or combination for the use according to claim 16, wherein said method is effective in inhibiting said colonic and/or somatic hypersensitivity.






Fig. 1. Temporal development of mechanical and thermal hypersensitivity post nerve injury. Nerve injury induced a significant increase in the number of paw withdrawals to A) vf2g, B)vF 6g and C) acetone stimulation of the ipsilateral hind paw compared with the responses seen in sham controls.

A) Effects of Palonosetron on Ipsilateral Behavioural Responses to vF 2g in SNL (n=9) and Sham (n=5) Animals



B) Effects of Palonosetron on Ipsilateral Behavioural Responses to vF 6g in SNL (n=9) and Sham (n=5) Animals



C) Effects of Palonosetron on Ipsilateral Behavioural Responses to Acetone in SNL (n=9) and Sham (n=5) Animals




Fig. 2. Effects of palonosetron on the number of paw withdrawals to A) vF2g, B) vF 6g and C) acetone. A dose related reduction in paw withdrawal frequency is seen in the SNL rats only




Fig.3 A comparison of total time spent by SNL and Sham rats on a rotarod. Palonosetron (0.03, 0.3 and 3mg/kg) did not impair motor performance in either group. Data presented as mean \pm SEM.




Fig.4. A comparison of the effects of Palonosetron (0.03, 0.3 and 3mg/kg s.c.) on the electrical evoked responses of spinal dorsal horn neurones in sham and SNL. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig.5. A comparison of the effects of Palonosetron (0.03, 0.3 and 3mg/kg s.c.) on the dynamic brush and mechanical punctate evoked responses of spinal dorsal horn neurones in sham and SNL. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig.6. Comparison of the effects of Palonosetron (0.03, 0.3 and 3mg/kg s.c.) on the thermal evoked responses of spinal dorsal horn neurones in sham and SNL. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig. 7. Effects of netupitant on the number of withdrawal responses to A) vF2g, B) vF6g and C) acetone (cooling stimuli). In SNL rats, 1mg/kg appears to inhibit behavioral hypersensitivities.




Fig. 8. A comparison of the effects of 3 doses of netupitant (0.1, 1 and 10mg/kg s.c) on the electrical responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig. 9. A comparison of the effects of 3 doses of netupitant (0.1, 1 and 10mg/kg s.c) on the dynamic brush and mechanical punctate evoked responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig. 10. A comparison of the effects of 3 doses of netupitant (0.1, 1 and 10mg/kg s.c) on the thermal evoked responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig. 11. Comparison of the effects of a combination of palonosetron (0.03 mg/kg s.c.) and netupitant (0.1 mg/kg s.c.) on the number of withdrawal responses in the ipsilateral and contralateral paws to A) vF2g, B) vF6g and C) acetone (cooling stimuli) in SNL rats.




Fig. 12. Comparison of the effects of a combination of palonosetron (0.03 mg/kg s.c) and netupitant (0.1 mg/kg s.c.) on the electrical responses of spinal dorsal horn neurones in sham and SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Fig. 13. Comparison of the effects of a combination of palonosetron (0.03 mg/kg s.c) and netupitant (0.1 mg/kg s.c.) on the dynamic brush, mechanical punctate and heat evoked responses of spinal dorsal horn neurones in SNL rats. Data are expressed as the mean percentage of pre-drug control values \pm S.E.M.




Figure 14: Palonosetron and Netupitant dosed in combination at 0.001 mg/kg showed a non-significant trend to inhibit AA-induced somatic hypersensitivity.

*compared to NC or *compared to VC or AA: One-way ANOVA with Bonferroni post-test




Figure 15: Visceral sensitivity results presented as individual distension pressure histograms with accompanying scatter-plot comparisons. Palonosetron or Netupitant dosed at 0.001 mg/kg p.o alone or in combination.

*compared to AA; *compared to NC; Ω compared to NT: One-way ANOVA with Bonferroni post-test



EUROPEAN SEARCH REPORT

Application Number

DOCUMENTS CONSIDERED TO BE RELEVANT					
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)		
Y	WO 2005/097794 A1 (JANSSEN PHARMACEUTICA NV [BE]; JANSSENS FRANS EDUARD [BE]; SCHOENTJES) 20 October 2005 (2005-10-20) * page 7, line 25 - line 2 * * page 8, line 19 - line 29 * -----	1-17	INV. A61K31/4523 A61K31/4748 A61P25/00 A61P25/02 A61P1/00		
Y	WO 99/17755 A2 (GLAXO GROUP LTD [GB]; MANGEL ALLEN WAYNE [US]; NORTHCUTT ALLISON RUTH) 15 April 1999 (1999-04-15) * page 2, line 12 - page 3, line 3; claims 1-10 * -----	1-17			
Y	STEADMAN C J ET AL: "Selective 5-Hydroxytryptamine Type 3 Receptor Antagonism With Ondansetron as Treatment for Diarrhea-Predominant Irritable Bowel Syndrome: A Pilot Study", MAYO CLINIC PROCEEDINGS, DOWDEN HEALTH MEDIA, INC, US, vol. 67, no. 8, 1 August 1992 (1992-08-01), pages 732-738, XP002098638, ISSN: 0025-6196 * the whole document * -----	1-17	TECHNICAL FIELDS SEARCHED (IPC)		
			A61K A61P		
The present search report has been drawn up for all claims					
Place of search	Date of completion of the search	Examiner			
Munich	24 June 2014	Loher, Florian			
CATEGORY OF CITED DOCUMENTS					
X : particularly relevant if taken alone	T : theory or principle underlying the invention				
Y : particularly relevant if combined with another document of the same category	E : earlier patent document, but published on, or after the filing date				
A : technological background	D : document cited in the application				
O : non-written disclosure	L : document cited for other reasons				
P : intermediate document	& : member of the same patent family, corresponding document				

5



Application Number

EP 14 16 5604

10

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing claims for which payment was due.

15

Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for those claims for which no payment was due and for those claims for which claims fees have been paid, namely claim(s):

20

No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for those claims for which no payment was due.

25

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

30

see sheet B

35

All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.

40

As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.

45

Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:

50

None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:

55

The present supplementary European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims (Rule 164 (1) EPC).



5

**LACK OF UNITY OF INVENTION
SHEET B**Application Number
EP 14 16 5604

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

10

1. claims: 1-17

Subject-matter of claims

15 1.1. claims: 1-17(partially)

Use of neptupitant for the treatment of irritable bowel syndrome

20 1.2. claims: 1-17(partially)

Use of palonosetron for the treatment of irritable bowel syndrome

25 1.3. claims: 1-17(partially)

Use of a combination comprising sub-therapeutic doses of palonosetron and netupitant for the treatment of irritable bowel syndrome

30

Please note that all inventions mentioned under item 1, although not necessarily linked by a common inventive concept, could be searched without effort justifying an additional fee.

35

40

45

50

55

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 14 16 5604

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on. The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

24-06-2014

10

		Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		WO 2005097794 A1	20-10-2005	AR 048521 A1 AT 471938 T AU 2005231984 A1 CA 2561975 A1 CN 1946723 A EP 1735313 A1 ES 2346985 T3 JO 2676 B JP 4884371 B2 JP 2007531786 A TW 1360548 B US 2007254904 A1 WO 2005097794 A1	03-05-2006 15-07-2010 20-10-2005 20-10-2005 11-04-2007 27-12-2006 22-10-2010 17-06-2012 29-02-2012 08-11-2007 21-03-2012 01-11-2007 20-10-2005
15					
20					
25					
30		WO 9917755 A2	15-04-1999	AR 017296 A1 AU 750818 B2 AU 9629398 A BR 9812886 A CA 2305751 A1 CN 1281357 A CO 5011096 A1 DZ 2618 A1 EE 2000000214 A EP 1021174 A2 GT 199800157 A HR P200000198 A2 HU 0003750 A2 ID 23874 A IS 5425 A JP 2001518495 A MA 26549 A1 NO 20001776 A NZ 503698 A PA 8461101 A1 PE 120299 A1 PL 340337 A1 SK 4862000 A3 SV 1998000122 A TR 2000000913 T2 US 6284770 B1 US 20010444450 A1 US 2003036549 A1 UY 25200 A1 WO 9917755 A2 YU 20600 A ZA 9809061 A	05-09-2001 25-07-2002 27-04-1999 08-08-2000 15-04-1999 24-01-2001 28-02-2001 01-03-2003 15-06-2001 26-07-2000 30-03-2000 30-04-2001 28-10-2001 25-05-2000 31-03-2000 16-10-2001 20-12-2004 06-06-2000 25-10-2002 24-05-2000 15-12-1999 29-01-2001 18-01-2001 05-07-1999 22-01-2001 04-09-2001 22-11-2001 20-02-2003 31-01-2001 15-04-1999 31-10-2003 05-07-2001
35					
40					
45					
50					
55					

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 5202333 A [0003]
- US 5510486 A [0003]
- WO 2004067005 A [0003]
- WO 2008049552 A [0003]
- US 6297375 B [0005]
- US 6719996 B [0005]
- US 6593472 B, Hoffmann La Roche [0005] [0022]

Non-patent literature cited in the description

- **ROILA F ; FATIGONI S.** NEW ANTIEMETIC DRUGS. *Ann Oncol*, 2006, vol. 17 (2), ii96-100 [0006]
- **ROCHE.** NK1 receptor antagonists, 23 February 2006, http://www.hospitalpharma.com/Features/feature.asp?ROW_ID=742 [0006]
- **GH MCCLEANE et al.** ANETH ANALG, 2003, vol. 97, 1474-8 [0007]
- **DE FELIPE C et al.** NATURE, March 1998, vol. 392 (6674), 394-7 [0008]
- **LM THOMSON et al.** J PAIN, January 2008, vol. 9 (1), 11-19 [0008]
- **R HILL.** TRENDS IN PHARMACEUTICAL SCIENCES, July 2000, vol. 21, 244-246 [0009]
- **MAYER et al.** Gastroenterol., 1994, vol. 107, 271-293 [0011]
- **KELLOW et al.** Gastroenterol., 1987, vol. 92, 1885-1893 [0011]
- **LEVITT et al.** Ann. Int. Med., 1996, vol. 124, 422-424 [0011]
- **DROSSMAN et al.** Gastroenterol. Int., 1995, vol. 8, 47-90 [0011]
- **NESS et al.** Brain Research, 1988, vol. 450, 153-169 [0012]
- **PLOURDE et al.** American Journal of Physiology, 1997, vol. 273, G191-196 [0012]
- **GAUDREAU et al.** Neuroscience Letters, 2003, vol. 351 (2), 59-62 [0012]
- **VENKOVA et al.** Toxicology and Applied Pharmacology, 2004, vol. 196, 215-222 [0012]
- **CHANG.** Gastroenterology Clinics of North America, 2005, vol. 34 (2), 271-279 [0012]
- **ZHOU et al.** Pain, 2010, vol. 148 (3), 454-461 [0012]
- **WOLFE F et al.** ARTHRITIS RHEUM, 1990, vol. 33, 160-72 [0038]

奈妥比坦與帕洛諾司瓊的具療效的組合

本文提供奈妥比坦與帕洛諾司瓊的組合，以及以所述組合治療多種痛楚狀況以及腸易激綜合症。