

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
31 December 2003 (31.12.2003)

PCT

(10) International Publication Number  
**WO 2004/000023 A1**

(51) International Patent Classification<sup>7</sup>: A01N 47/18, 47/20, C07C 271/34, C07D 295/205

(21) International Application Number:  
PCT/CH2003/000403

(22) International Filing Date: 20 June 2003 (20.06.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0214342.8 21 June 2002 (21.06.2002) GB

(71) Applicant (for all designated States except US): GIVAUDAN SA [CH/CH]; Chemin de la Parfumerie 5, CH-1214 Vernier (CH).

(72) Inventor; and

(75) Inventor/Applicant (for US only): GAUTSCHI, Markus [CH/CH]; Am Stutz, CH-4314 Zeiningen (CH).

(74) Agent: MCSTEA, John, Anthony; Global Patents, Überlandstrasse 138, CH-8600 Dübendorf (CH).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/000023 A1

(54) Title: INSECT REPELLENTS

(57) Abstract: Use of an N-substituted p-menthane carboxamide as an active ingredient in an insect repellent composition, which has little or no odour and which is substantially non-toxic for human, pet and livestock.

### Insect repellents

The present invention relates to compounds having insect repellent characteristics and to compositions containing same.

5

Many insects are known as a nuisance and some insect genera even represent a health hazard. Therefore, many efforts have been made to eradicate or at least to control these pests. One method of insect eradication is through the use of synthetically produced insecticides. However, certain insect genera may develop resistance to some whilst others have 10 undesirable effects on human and other animal life, such that their use must be strictly regulated or even forbidden.


Naturally-occurring substances are also known to display insect repellent properties, for example citronella oil, tolu and Peru balsams, camphor and various eucalyptus. However, 15 many possess olfactory properties that makes unacceptable their use in compositions that also contain a perfume, at least in amounts required to have repellent effects.

The prior art has continued to propose highly effective insect repellent compounds that have low, or substantially no odour. Thus, in US 5,182,305 N-aryl and N-cycloalkyl 20 neoalkanamides are described. In US 5,391,578 N-lower alkyl neoalkanamides are described as being superior to DEET in long lasting effectiveness of the insect repellency. In WO 00/16738 menthyl 2-pyrrolidone-5-carboxylate are described to be an effective insect repellent, comparable to DEET. Finally, in WO 02/15692, there are described compositions containing certain menthane carboxamides having excellent insect repellent properties.

25

These prior art efforts reflect the consumer demand for improved insect repellent compounds that are not harmful to the user or the environment, and yet which are highly effective against target insect populations. Furthermore, as the end-uses of such compounds are in high-volume, low-price or commodity products, price sensitivity is an issue that has to be 30 addressed in the development of new compounds.

There has now been found a new category of compounds having interesting insect repellent properties that meet the requirements set forth in the preceding paragraph. Thus the invention provides in a first aspect the use as an insect repellent of a compound of the formula



5 wherein,

R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of H; an aliphatic residue having 1 to 20 carbon atoms, or a cycloaliphatic residue having 5 to 14 carbon atoms, or an aliphatic or cycloaliphatic residue aforementioned containing one or more hetero-atoms

10 selected from O, N or S; an aryl or heteroaryl group having from 6 to 14 carbon atoms and wherein hetero-atoms are selected from O, N or S; or any of the afore-mentioned groups substituted with a group selected from, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>2-4</sub> alkenyl, aryl or heteroaryl as defined above, aryloxy, amino-, amido-, ester, keto-, hydroxyl, and halogen, e.g. Cl, Br or I, or

15

R<sub>1</sub> and R<sub>2</sub> together with the nitrogen atom to which they are attached form a 5- or 6-membered ring that may optionally contain additional hetero-atoms selected from O, N or S.

Preferred groups R<sub>1</sub> and R<sub>2</sub> may be selected from alkyl, e.g. C<sub>1-4</sub> alkyl, more particularly

20 methyl, ethyl, n or iso propyl, or n or sec butyl, cycloalkyl, e.g. having 5- or 6-carbon atoms or phenyl.

Most preferred groups R<sub>1</sub> and R<sub>2</sub> are those groups provided on the compounds of Example 1 hereinunder.


25

The compounds n-butyl-carbamic acid (-)-menthyl ester; iso-butyl-carbamic acid (-)-menthyl ester; diethyl-carbamic acid (-)-menthyl ester; morpholine-4-carboxylic acid (-)-menthyl

ester; and 3-[(-)-menthoxy-carbonylamino]-propionic acid ester have interesting insect repellent properties and these compounds form another aspect of the invention.

The compounds defined hereinabove display good repellent activity against insects, in particular against cockroaches. At the same they are safe to be applied to the human body, pets and livestock, or on or against surfaces which may be contacted by humans, pets or livestock. The compounds also possess good substantivity, thereby providing long-lasting activity against insect infestations on surfaces to which they are applied. Furthermore, the compounds possess little or no odour, which makes them suitable for use in perfumed compositions. It is possible to use more than one such compound in an application.

The compounds of the invention have 3 chiral centres, giving rise to 8 stereoisomers. Thus, all possible stereoisomers are included in the scope of the present invention. However, in general, compounds having the stereo-chemistry set forth below



15

derived from the naturally occurring menthol [(-)-3-p-menthanol] are preferred.

In a preferred embodiment of the present invention, there is provided the use as an insect repellent of a compound or mixture of compounds selected from the compounds set forth in Example 1 below.

Compounds of the present invention may be prepared in a straightforward manner. For example, menthol may be reacted with a chloroformate bearing the groups R<sub>1</sub> and R<sub>2</sub> under alkaline conditions in a manner known in the art. The chloroformates are either commercially available or can be synthesised from known starting materials according to synthetic procedures known to the art. Further and more specific information regarding the syntheses of the compounds is set forth in the Examples.

In another aspect of the present invention there is provided an insect repellent composition comprising at least one of the compounds hereinabove described.

The amount of compound or compounds employed in said composition may vary widely

5 depending on a number of factors including the nature of the insect infestation that is intended to be treated and the presence or absence of other insect repellent agents in the composition. Typically, a compound or mixtures of the compounds is used at levels of from 1ppm to 1000ppm, although levels outside these limits may also be useful in some cases. It is possible to provide the compound or mixture of compounds as a concentrate containing

10 typically from 0.1 to 25% by weight of the composition of the compounds or mixture of compounds. This can be then be added by the user to other ingredients to give an end-product having the desired concentration.

Compositions containing a compound or mixtures of compounds of the present invention

15 may be applied to objects in need of protection against insects, either directly, in liquid solution or dispersion, as aerosols or air-sprays, or dispersed in a powdered carrier or in a suitable composition. Compositions which may be useful to repel insects are, for example, detergent compositions, cleaning compositions, paints, wallpaper, upholstery and/or rug shampoos, liquid soaps, soap bars, floor polishes, floor waxes and furniture polishes.

20 Compositions which are useful to repel insects from the human body are also included in the scope of the present invention and include compositions such as fine fragrances, colognes, skin creams, sun creams, skin lotions, deodorants, talcs, bath oils, soaps, shampoos, hair conditioners and styling agents.

25 The compositions of the present invention may comprise a compound or compounds of the present invention in combination with other known insect repellents, including, but not limited to, N,N-diethyl-m-toluamide (DEET), N,N-diethyl-benzamide, menthyl 2-pyrrolidone-5-carboxylate, N-aryl and N-cycloalkyl neoalkanamides, N-lower alkyl neoalkanamides and nepetalactone. The compositions of the present invention may also

30 comprise natural oils known for their insect repellent characteristics. Examples for such oils include, without limiting, e.g. citronella oil, catnip oil, eucalyptus oil, cypress oil, galbanum oil, tolu and Peru balsams.

A compound of the present invention or mixtures thereof may also be used in conjunction with at least one insecticide in order to repel insects from one area and toward the location, where the insecticide is applied to avoid the action of the insecticide in a special area, for 5 example in areas containing foodstuffs. Alternatively, they may be formulated with insecticides so that after the repellent activity has diminished the treated area will still not be safe for insects.

A compound or mixtures of compounds for use in the present invention may additionally 10 comprise at least one fragrance compound. Such fragrance compounds may be of natural and/or synthetic origin, examples for such natural and synthetic fragrance ingredients can be found e.g. in "Perfume and Flavour Materials of Natural Origin", S. Arctander, Ed., Elizabeth, N.J., 1960 and "Perfume and Flavour Chemicals", S. Arctander, Ed., Vol. I & II, Allured Publishing Corporation, Carol Stream, USA, 1994.

15

Such compositions may additionally contain solvents. Solvents that may be used are known to those skilled in the art and include ethanol, ethylene glycol, propylene glycol, diethyl phthalate and dimethyl phthalate. A preferred solvent is dimethyl phthalate, which is known for its insect repellent characteristics.

20

The compositions of the present invention may comprise other art-recognised ingredients normally used in such formulations. These include antifoaming agents, anti-microbial agents, anti-oxidants, anti-redeposition agents, bleaches, colorants, emulsifiers, enzymes, fats, fluorescent materials, fungicides, hydrotropes, moisturisers, optical brighteners, 25 perfume carriers, perfume, preservatives, proteins, silicones, soil release agents, solubilisers, sugar derivatives, sun screens, surfactants, vitamins and waxes.

Compounds for use in the present invention have useful properties both as contact and vapour repellent. They are superior to various commercial insect repellents in repelling 30 action, especially against German cockroaches, which are considered to be one of the most difficult household pests to control. Due to their low vapour pressure, the compounds are long-lasting on surfaces to which they have been applied. The long-lasting insect repellency

may last up to 2-3 weeks after topical application, depending on the concentration used. Furthermore, the compounds are sufficiently stable in compositions being object of the present invention to maintain their insect repellency.

- 5 Compositions for use in the present invention may also be incorporated in various materials during their manufacturing process. Methods for preparing a product comprising a composition according to the present invention by incorporating said composition into the product during extrusion are preferred.
- 10 In addition to their effectiveness against German cockroaches, compositions formulated according to the present invention are also effective against other insects such as ants, bees, fleas, flies, hornets, mosquitoes, moths, silverfish, and wasps, and against arachnids such as mites, spiders and ticks.
- 15 The effectiveness against mosquitoes is important also for economic reasons, especially against the genera *Anopheles* (which is a known carrier of malaria and transmits also filariasis and encephalitis), *Culex* (which is a carrier of viral encephalitis and filariasis) and *Aedes* (which carries yellow fever, dengue and encephalitis). From the latter genus, the activity against *Aedes aegypti* is especially important.

20

Products that may benefit from the addition or incorporation of compounds and/or compositions provided by this invention include, but are not restricted to, household products, industrial cleansing products, personal care products, and pet or livestock care products. In addition, the invention also provides covering materials especially floor, wall  
25 and furniture coverings, fabrics and plastics materials comprising a composition or compounds as hereinabove defined. The invention therefore also provides a method of repelling insects by the application to a substrate of a preparation comprising at least one compound as hereinabove described.

The invention is now further described, by way of illustration, in the following non-limiting  
30 examples.

Example 1


General procedure for the synthesis of (–)-menthyl carbamates:

5 To a mechanically stirred solution of a corresponding amine (0.4 mol, 5 equiv.) in toluene (100 ml), cooled to 5 °C in an ice-water bath, was added dropwise a solution of (–)-menthyl chloroformate (17.5 g, 0.08 mol, 1 equiv.) in toluene (20 ml) within 0.5 h. The temperature was kept between 5 and 15 °C with cooling. The resulting suspension was allowed to warm to room temperature and stirring continued until the reaction was complete according to TLC  
 10 analysis (0.5 to 3h).

The mixture was poured into ice/water (300 ml), extracted with toluene (2 x 200 ml) and the organic phases were each washed with aqueous hydrochloric acid (5%, 200 ml), water (200 ml) and brine (200 ml). The combined organic layers were dried over sodium sulphate and  
 15 concentrated *in vacuo* to give the crude carbamate. (In the case of methylamine and ethylamine, were ethanolic solutions were used, the reaction mixture remains a solution and ethanol needs to be removed prior to aqueous work-up.) The crudes were either used without purification or purified by re-crystallisation from hexane or 'Kugelrohr' distilled to yield the clean (–)-menthyl carbamates in 75 to 90%.

20

Characterisation of (–)-menthyl carbamates:




25 **Methyl-carbamic acid (–)-menthyl ester**

mp. 108-110 °C (hexane), white solid

<sup>1</sup>H-NMR (400 MHz, CDCl<sub>3</sub>, coupling constants in Hz): 0.79 (3H, d, *J* 7, Me); 0.80-0.98 (2H, m, 2 x CH); 0.89 (6H, d, *J* 7, Me<sub>2</sub>); 1.06 (1H, dq, *J* 13, 4, CH); 1.29 (1H, brt, *J* 11, CH);

1.42-1.55 (1H, m, CH); 1.60-1.72 (2H, m, 2 x CH); 1.86-1.98 (1H, m, CH); 2.01-2.09 (1H, m, CH); 2.79 (3H, d, *J* 5, *MeNH*); 4.28-4.57 (1H, brm, NH); 4.55 (1H, dt, *J* 11, 4, CHOCO).  
**IR** ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ , ATR): 3377w, 2961m, 1688s, 1525s, 1258s, 1139s.  
**MS** [m/z (EI)]: 213 ( $\text{M}^+$ , <1%), 138 (70), 123 (41), 95 (100), 81 (73), 55 (56), 41 (47).

5



### Ethyl-carbamic acid (-)-menthyl ester


10 **mp.** 55-56 °C (hexane), white solid

**$^1\text{H-NMR}$**  (400 MHz,  $\text{CDCl}_3$ , coupling constants in Hz): 0.79 (3H, d, *J* 7, Me); 0.81-0.99 (2H, m, 2 x CH); 0.89 (6H, d, *J* 7,  $\text{Me}_2$ ); 1.05 (1H, dq, *J* 13, 4, CH); 1.13 (3H, t, *J* 7,  $\text{CH}_2\text{Me}$ ); 1.28 (1H, brt, *J* 11, CH); 1.43-1.55 (1H, m, CH); 1.61-1.72 (2H, m, 2 x CH); 1.86-1.99 (1H, m, CH); 2.01-2.09 (1H, m, CH); 3.14-3.27 (2H, brm,  $\text{CH}_2\text{NH}$ ); 4.26-4.65 (1H, brm, NH); 4.48-4.61 (1H, brm, CHOCO).

**IR** ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ , ATR): 3377w, 2960m, 1687s, 1524s, 1249s, 1019m.

**MS** [m/z (EI)]: 227 ( $\text{M}+\text{H}^+$ , <1%), 138 (65), 123 (28), 95 (100), 90 (83), 81 (82), 71 (41), 55 (57), 41 (32).

20



### Butyl-carbamic acid (-)-menthyl ester


**mp.** 60-62 °C (hexane), white solid

25  **$^1\text{H-NMR}$**  (400 MHz,  $\text{CDCl}_3$ , coupling constants in Hz): 0.79 (3H, d, *J* 7, Me); 0.80-0.98 (2H, m, 2 x CH); 0.89 (6H, d, *J* 7,  $\text{Me}_2$ ); 0.92 (3H, t, *J* 7,  $\text{CH}_2\text{Me}$ ); 1.06 (1H, dq, *J* 13, 4, CH); 1.28 (1H, brt, *J* 11, CH); 1.31-1.39 (2H, m,  $\text{CH}_2\text{Me}$ ); 1.41-1.55 (3H, m, CH and  $\text{CH}_2$ );

1.61-1.72 (2H, m, 2 x CH); 1.87-1.98 (1H, m, CH); 2.00-2.09 (1H, m, CH); 3.11-3.23 (2H, brm,  $CH_2NH$ ); 4.32-4.65 (1H, brm, NH); 4.48-4.61 (1H, brm, CHOCO).

**IR** ( $\nu_{max}$ ,  $cm^{-1}$ , ATR): 3369w, 2957m, 1686s, 1525s, 1240s, 1023m.

**MS** [m/z (EI)]: 138 (M-HOCONH $i$ Bu $^+$ , 26%), 123 (20), 118 (18), 95 (59), 81 (48), 71 (26), 56 (61), 41 (100), 27 (34).



#### Isobutyl-carbamic acid (-)-menthyl ester

10


**mp.** 99-102 °C (hexane), white solid

**$^1H$ -NMR** (400 MHz,  $CDCl_3$ , coupling constants in Hz): 0.79 (3H, d,  $J$  7, Me); 0.80-0.98 (2H, m, 2 x CH); 0.89 (6H, d,  $J$  7,  $Me_2$ ); 0.90 (6H, d,  $J$  7,  $Me_2$ ); 1.06 (1H, dq,  $J$  13, 4, CH); 1.28 (1H, brt,  $J$  11, CH); 1.42-1.53 (1H, m, CH); 1.62-1.78 (3H, m, 3 x CH); 1.86-1.97 (1H, m, CH); 2.00-2.08 (1H, m, CH); 2.95-3.03 (2H, brm,  $CH_2NH$ ); 4.45-4.68 (1H, brm, NH); 4.53 (1H, dt,  $J$  11, 4, CHOCO).

**IR** ( $\nu_{max}$ ,  $cm^{-1}$ , ATR): 3373w, 2959m, 1687s, 1528s, 1276m, 1241s, 1144m, 1039m.

**MS** [m/z (EI)]: 138 (M-HOCONH $i$ Bu $^+$ , 28%), 123 (17), 118 (16), 95 (59), 81 (44), 71 (31), 56 (42), 43 (100), 27 (31).

20



#### Diethyl-carbamic acid (-)-menthyl ester

25 **bp.** 90 °C, 0.03 mbar, colourless liquid


**$^1H$ -NMR** (400 MHz,  $CDCl_3$ , coupling constants in Hz): 0.79 (3H, d,  $J$  7, Me); 0.87 (1H, dt,  $J$  11, 3, CH); 0.90 (6H, d,  $J$  7,  $Me_2$ ); 0.95 (1H, q,  $J$  11, CH); 1.07 (1H, dq,  $J$  13, 4, CH); 1.10 (6H, t,  $J$  7, 2 x  $CH_2Me$ ); 1.36 (1H, tt,  $J$  11, 3, CH); 1.42-1.55 (1H, m, CH); 1.62-1.73 (2H, m,

10

2 x CH); 1.94 (1H, ddq, *J* 13, 7, 3, CH); 2.02-2.10 (1H, m, CH); 3.20-3.35 (4H, brm, 2 x CH<sub>2</sub>NH); 4.57 (1H, dt, *J* 11, 4, CHOCO).

**IR** ( $\nu_{\text{max}}$ , cm<sup>-1</sup>, ATR): 2955m, 1695s, 1422m, 1271s, 1172s, 993m.

**MS** [m/z (EI)]: 255 (M<sup>+</sup>, <1%), 138 (42), 118 (91), 95 (29), 83 (100), 69 (34), 55 (48), 41 5 (29).



**Pyrrolidine-1-carboxylic acid (-)-menthyl ester**

10


**crude product:** colourless liquid

**<sup>1</sup>H-NMR** (400 MHz, CDCl<sub>3</sub>, coupling constants in Hz): 0.80 (3H, d, *J* 7, Me); 0.88 (1H, dt, *J* 11, 3, CH); 0.90 (6H, d, *J* 7, Me<sub>2</sub>); 0.95 (1H, q, *J* 11, CH); 1.07 (1H, dq, *J* 13, 4, CH); 1.36 (1H, tt, *J* 11, 3, CH); 1.42-1.56 (1H, m, CH); 1.62-1.72 (2H, m, 2 x CH); 1.81-1.89 (6H, m, CH<sub>2</sub>CH<sub>2</sub>); 1.96 (1H, ddq, *J* 13, 7, 3, CH); 2.05-2.13 (1H, m, CH); 3.28-3.40 (4H, brm, 2 x CH<sub>2</sub>NH); 4.55 (1H, dt, *J* 11, 4, CHOCO).

**IR** ( $\nu_{\text{max}}$ , cm<sup>-1</sup>, ATR): 2952m, 1699s, 1411s, 1100m.

**MS** [m/z (EI)]: 253 (M<sup>+</sup>, <1%), 138 (31), 116 (100), 98 (26), 95 (38), 83 (65), 69 (29), 55 (61), 41 (32).

20



**Piperidine-1-carboxylic acid (-)-menthyl ester**


25 **crude product:** colourless liquid

**<sup>1</sup>H-NMR** (400 MHz, CDCl<sub>3</sub>, coupling constants in Hz): 0.79 (3H, d, *J* 7, Me); 0.88 (1H, dt, *J* 11, 3, CH); 0.91 (6H, d, *J* 7, Me<sub>2</sub>); 0.94 (1H, q, *J* 11, CH); 1.07 (1H, dq, *J* 13, 4, CH); 1.37

(1H, tt, *J* 11, 3, CH); 1.42-1.72 (9H, m, 3 x CH, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>); 1.92 (1H, ddq, *J* 13, 7, 3, CH); 2.03-2.10 (1H, m, CH); 3.40 (4H, t, *J* 6, 2 x CH<sub>2</sub>NH); 4.55 (1H, dt, *J* 11, 4, CHOCO).

**IR** ( $\nu_{\text{max}}$ , cm<sup>-1</sup>, ATR): 2933m, 1695s, 1425m, 1231m.

**MS** [m/z (EI)]: 267 (M<sup>+</sup>, <1%), 138 (41), 130 (100), 95 (43), 83 (66), 69 (41), 55 (47), 41 (46).



#### Morpholine-4-carboxylic acid (-)-menthyl ester

10


**crude product:** colourless liquid

**<sup>1</sup>H-NMR** (400 MHz, CDCl<sub>3</sub>, coupling constants in Hz): 0.79 (3H, d, *J* 7, Me); 0.87 (1H, dt, *J* 11, 3, CH); 0.90 (6H, d, *J* 7, Me<sub>2</sub>); 0.95 (1H, q, *J* 11, CH); 1.08 (1H, dq, *J* 13, 4, CH); 1.36 (1H, tt, *J* 11, 3, CH); 1.42-1.56 (1H, m, CH); 1.63-1.72 (2H, m, 2 x CH); 1.89 (1H, ddq, *J* 13, 7, 3, CH); 2.03-2.11 (1H, m, CH); 3.46 (4H, t, *J* 5, 2 x CH<sub>2</sub>NH); 3.65 (4H, t, *J* 5, 2 x CH<sub>2</sub>O); 4.58 (1H, dt, *J* 11, 4, CHOCO).

**IR** ( $\nu_{\text{max}}$ , cm<sup>-1</sup>, ATR): 2956m, 1699s, 1420m, 1238s, 1117m.

**MS** [m/z (EI)]: 269 (M<sup>+</sup>, <1%), 138 (61), 95 (42), 83 (100), 69 (37), 57 (41), 55 (53), 41 (33).

20



#### Phenyl-carbamic acid (-)-menthyl ester


25 **mp.** 111-113 °C (hexane), white solid

**<sup>1</sup>H-NMR** (400 MHz, CDCl<sub>3</sub>, coupling constants in Hz): 0.81 (3H, d, *J* 7, Me); 0.89 (1H, dt, *J* 11, 3, CH); 0.92 (6H, d, *J* 7, Me<sub>2</sub>); 1.01 (1H, q, *J* 11, CH); 1.10 (1H, dq, *J* 13, 4, CH); 1.37 (1H, tt, *J* 11, 3, CH); 1.45-1.58 (1H, m, CH); 1.66-1.74 (2H, m, 2 x CH); 1.97 (1H, ddq, *J*

13, 7, 3, CH); 2.08-2.15 (1H, m, CH); 4.66 (1H, dt, *J* 11, 4, CHOCO); 6.54 (1H, brs, NH); 7.05 (1H, brt, *J* 7.5, PhH); 7.30 (2H, t, *J* 7.5, 2 x PhH); 7.38 (2H, brd, *J* 7.5, 2 x PhH).

IR ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ , ATR): 3364w, 2955m, 1697s, 1524s, 1443s, 1226s, 1050s.

MS [m/z (EI)]: 275 ( $\text{M}^+$ , 7%), 137 (28), 119 (89), 95 (84), 83 (100), 69 (45), 55 (82), 41 (62).



### 3-[(−)-menthoxy-carbonylamino]-propionic acid ester

10

mp. 49-51 °C (hexane), white solid

<sup>1</sup>H-NMR (400 MHz,  $\text{CDCl}_3$ , coupling constants in Hz): 0.79 (3H, d, *J* 7, Me); 0.80-0.98 (2H, m, 2 x CH); 0.91 (6H, d, *J* 7,  $\text{Me}_2$ ); 1.05 (1H, dq, *J* 13, 4, CH); 1.29 (1H, brt, *J* 11, CH); 1.42-1.55 (1H, m, CH); 1.62-1.73 (2H, m, 2 x CH); 1.85-1.97 (1H, m, CH); 1.99-2.08 (1H, m, CH); 2.55 (2H, t, *J* 6,  $\text{CH}_2\text{CO}$ ); 3.38-3.49 (2H, m,  $\text{CH}_2\text{NH}$ ); 3.70 (3H, s, OMe); 4.53 (1H, dt, *J* 11, 4, CHOCO); 5.07-5.18 (1H, brm, NH).

IR ( $\nu_{\text{max}}$ ,  $\text{cm}^{-1}$ , ATR): 3362w, 2959m, 1734s, 1683s, 1530s, 1256s, 1199m, 1177s, 991m.

MS [m/z (EI)]: 286 ( $\text{M}+\text{H}^+$ , 3%), 148 (69), 138 (100), 123 (45), 116 (53), 104 (92), 95 (91), 81 (60), 55 (37), 41 (24).

20

#### Example 2:

The effectiveness of compounds of formula I as insect repellents is determined using German cockroaches exposed to a melamine-formaldehyde laminate surface partially treated/partially untreated with a test compound of formula I.

The study was conducted in an air-conditioned laboratory at a temperature of  $22^\circ\pm2^\circ\text{C}$  under a normal day/night cycle, using German cockroaches (*Blatella Germanica*) of mixed sex and age. The study was conducted over a 3 day period and the repellent effectiveness was assessed 1, 2 and 3 days after cockroach introduction.

Test method:

A. Treatment

One half of a rectangle of laminate 40cm x 30cm was treated at 20 mg/m<sup>2</sup> of a test compound of formula I. This was achieved by soaking a paper wipe ("Rag on a Roll" ca 5 20cm x 20cm) in a solution of the test compound in ethanol, squeezing out the excess liquid and wiping over the surface to give the required coverage. This was checked by weighing the cloth after application. The surface was allowed to dry. The other half of the rectangle was wiped with ethanol alone (untreated surface).

10 B. Bioassays

The treated laminate rectangle was placed on the bench and a plastic container was placed on the rectangle. This had been treated with "Fluon" (trade mark), a sprayable fluoropolymer, to prevent cockroach escape.

15 Two sheets of laminate (10cm x 10cm) were placed on the laminate, one on the treated surface and one on the untreated surface. This acted as a cockroach harbourage. The laminate sheet to be used in the treated section was treated at the same rate as the treated laminate. The laminate was placed on two 1cm high bottle tops to allow cockroaches access under the laminate.

20 German cockroaches were added (5 adults and 15 nymphs) to the centre of the 20 laminate. No food and water was supplied for these cockroaches.

At 1, 2 and 3 days after cockroach introduction the number of cockroaches under each harbourage was counted. After each count the harbourage position was changed to avoid the possibility of habituation to one location, the harbourages were replaced and new cockroaches were added to replace any dead cockroaches.

25 The above was repeated 3 times to provide a total of 4 replicates.


## C. Repellency results for compound Methyl-carbamic acid (−)-menthyl ester

Time After Cockroach Introduction (Days)	Replicate	Number of Cockroaches present (n=20)			
		Treated Section	Untreated Section	Dead	Total
1	1	0	20	0	20
	2	0	19	1	20
	3	3	17	0	20
	4	1	19	0	20
	<b>Total</b>	<b>4</b>	<b>75</b>	<b>1</b>	<b>80</b>
2	1	1	19	0	20
	2	0	19	1	20
	3	0	20	0	20
	4	1	19	0	20
	<b>Total</b>	<b>2</b>	<b>77</b>	<b>1</b>	<b>80</b>
3	1	0	20	0	20
	2	0	20	0	20
	3	1	19	0	20
	4	0	20	0	20
	<b>Total</b>	<b>1</b>	<b>79</b>	<b>0</b>	<b>80</b>

5 Example 3


The methodology of Example 2 was repeated for all of the compounds set forth in Example 1. The results are presented in the following table.

Structure	Days after treatment	Percentage Present on	
		treated section	untreated section
Compound	Days after treatment		
	1	5.1	94.9
	2	2.5	97.5
	3	1.3	98.7
	Average	3.0	97.0
	1	7.6	92.4
	2	2.5	97.5
	3	2.5	97.5
	Average	4.2	95.8
	1	2.5	97.5
	2	3.7	96.3
	3	2.5	97.5
	Average	2.9	97.1
	1	2.5	97.5
	2	7.5	92.5
	3	12.7	87.3
	Average	7.6	92.4
	1	17.5	82.5
	2	17.5	82.5
	3	25.0	75.0
	Average	20.0	80.0
	1	10.0	90.0
	2	11.3	88.7
	3	3.9	96.1
	Average	8.4	91.6
	1	31.3	68.7
	2	17.5	82.5
	3	16.3	83.7
	Average	21.7	78.3
	1	6.3	93.7
	2	17.5	82.5
	3	7.5	92.5
	Average	10.4	89.6

	1	11.4	88.6
	2	2.5	97.5
	3	10.0	90.0
	Average	8.0	92.0

Claims

1. Use as an insect repellent of a compound of the formula



5 wherein,

R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of H; an aliphatic residue having 1 to 20 carbon atoms, or a cycloaliphatic residue having 5 to 14 carbon atoms, or an aliphatic or cycloaliphatic residue aforementioned containing one or more hetero-atoms selected from O, N or S; an aryl or heteroaryl group having 10 from 6 to 14 carbon atoms and wherein hetero-atoms are selected from O, N or S; or any of the afore-mentioned groups substituted with a group selected from, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>2-4</sub> alkenyl, aryl or heteroaryl as defined above, aryloxy, amino-, amido-, ester, keto-, hydroxyl, and halogen, or

15 R<sub>1</sub> and R<sub>2</sub> together with the nitrogen atom to which they are attached form a 5- or 6-membered ring that may optionally contain additional hetero-atoms selected from O, N or S.

2. Use according to claim 1 wherein the compound is selected from the group consisting 20 of

Methyl-carbamic acid (-)-menthyl ester;  
Ethyl-carbamic acid (-)-menthyl ester;  
Butyl-carbamic acid (-)-menthyl ester;  
Isobutyl-carbamic acid (-)-menthyl ester;  
25 Diethyl-carbamic acid (-)-menthyl ester;  
Pyrrolidine-1-carboxylic acid (-)-menthyl ester;

Piperidine-1-carboxylic acid (-)-menthyl ester;  
 Morpholine-4-carboxylic acid (-)-menthyl ester;  
 Phenyl-carbamic acid (-)-menthyl ester; and  
 3-[(-)-menthoxy-carbonylamino]-propionic acid ester.

5 3. A compound selected from the group consisting of n-butyl-carbamic acid (-)-menthyl ester; iso-butyl-carbamic acid (-)-menthyl ester; diethyl-carbamic acid (-)-menthyl ester; morpholine-4-carboxylic acid (-)-menthyl ester; and 3-[(-)-menthoxy-carbonylamino]-propionic acid ester.


4. A composition comprising a compound as defined in any of the preceding claims in  
 10 an insect-repellent amount.

5. A composition according to claim 4 comprising at least one additional insect repellent.

6. A composition according to claim 4 or claim 5 comprising additionally at least one insecticide.

15 7. A composition according to any one of the claims 4 to 6 comprising additionally at least one fragrance ingredient.

8. A method of repelling insects by applying to a substrate a preparation comprising at least one compound of the formula



20 wherein,

R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of H; an aliphatic residue having 1 to 20 carbon atoms, or a cycloaliphatic residue having 5 to 14 carbon atoms, or an aliphatic or cycloaliphatic residue aforementioned containing one

19

or more hetero-atoms selected from O, N or S; an aryl or heteroaryl group having from 6 to 14 carbon atoms and wherein hetero-atoms are selected from O, N or S; or any of the afore-mentioned groups substituted with a group selected from, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>2-4</sub> alkenyl, aryl or heteroaryl as defined above, aryloxy, amino-, 5 amido-, ester, keto-, hydroxyl, and halogen, or

R<sub>1</sub> and R<sub>2</sub> together with the nitrogen atom to which they are attached form a 5- or 6-membered ring that may optionally contain additional hetero-atoms selected from O, N or S.

10

15

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/CH 03/00403

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 7 A01N47/18 A01N47/20 C07C271/34 C07D295/205

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 02 15692 A (GIVAUDAN SA ;GAUTSCHI MARKUS (CH); BLONDEAU PHILIPPE (FR)) 28 February 2002 (2002-02-28) cited in the application the whole document --- -/-	1-8

 Further documents are listed in the continuation of box C. Patent family members are listed in annex.

## ° Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the international search report

19 September 2003

08/10/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Mue11ners, W

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/CH 03/00403

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE WPI      Section Ch, Week 198547      Derwent Publications Ltd., London, GB;      Class C03, AN 1985-292280      XP002255150      &amp; JP 60 199804 A      abstract      -&amp; JP 60 199804 A (NIPPON KAYAKU KK)      9 October 1985 (1985-10-09)      example 12      -&amp; PATENT ABSTRACTS OF JAPAN      vol. 010, no. 056 (C-331),      6 March 1986 (1986-03-06)      &amp; JP 60 199804 A      abstract</p> <p>---</p>	1-8
A	<p>WO 00 16738 A (HILLS MAXINE JANE ;WATKINS STEPHEN DAVID (GB); BIRCH RICHARD ARTHUR)      30 March 2000 (2000-03-30)      cited in the application      page 1 -page 3, line 14</p> <p>---</p>	1-8
A	<p>KNOWLES, CHARLES O. ET AL: "Biological activity of N-methylcarbamate and dimethyl phosphorothionate esters of various phenols and monoterpenoids"      JOURNAL OF ECONOMIC ENTOMOLOGY (1967),      60(5), 1417-20 ,      1967, XP002255148      page 1417, the abstract      page 1410, right-hand column, paragraph 4      page 1419, table 1, compound VIII</p> <p>---</p>	1-8
A	<p>DATABASE CA 'Online!      CHEMICAL ABSTRACTS SERVICE, COLUMBUS,      OHIO, US;      KYUNG, SÜK-HUN ET AL: "Insecticidal activity of some monoterpenoids"      retrieved from STN      Database accession no. 137:164969 CA      XP002255149      abstract      &amp; HAN'GUK NONGHWA HAKHOECHI (2002), 45(1),      37-41 ,      2002,</p> <p>---</p>	1-8
A	<p>DE 15 18 713 B (DEUTSCHE GOLD- U.      SILBER-SCHEIDEANSTALT)      14 January 1971 (1971-01-14)      column 1, paragraph 1 - paragraph 2;      example 22</p> <p>-----</p>	1-3

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CH 03/00403

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0215692	A	28-02-2002	AU BR WO EP	8769601 A 0113421 A 0215692 A1 1311158 A1	04-03-2002 15-07-2003 28-02-2002 21-05-2003
JP 60199804	A	09-10-1985	JP JP	1711277 C 3080138 B	11-11-1992 24-12-1991
WO 0016738	A	30-03-2000	AU DE EP WO JP US	5634299 A 69903655 D1 1113782 A1 0016738 A1 2002526432 T 6451844 B1	10-04-2000 28-11-2002 11-07-2001 30-03-2000 20-08-2002 17-09-2002
DE 1518713	B	14-01-1971	AT BE CH DE DK FI FR GB NL SE US	280984 B 684038 A 481863 A 1518713 B1 123020 B 44602 B 5603 M 1099558 A 6609768 A 351419 B 3480663 A	11-05-1970 16-12-1966 30-11-1969 14-01-1971 08-05-1972 31-08-1971 11-12-1967 17-01-1968 16-01-1967 27-11-1972 25-11-1969