



(86) **Date de dépôt PCT/PCT Filing Date:** 2014/02/06
(87) **Date publication PCT/PCT Publication Date:** 2014/08/21
(85) **Entrée phase nationale/National Entry:** 2015/08/12
(86) **N° demande PCT/PCT Application No.:** EP 2014/052342
(87) **N° publication PCT/PCT Publication No.:** 2014/124862
(30) **Priorité/Priority:** 2013/02/14 (ITMI2013A000210)

(51) **Cl.Int./Int.Cl. A61K 31/135** (2006.01),
A61K 31/192 (2006.01)

(71) **Demandeur/Applicant:**
LABORATORIOS MENARINI SA, ES

(72) **Inventeurs/Inventors:**
SCHMITZ, REINHARD, DE;
KOHL, TOBIAS, DE

(74) **Agent:** KIRBY EADES GALE BAKER

(54) **Titre : COMPOSITIONS PHARMACEUTIQUES CONTENANT DU DEXKETOPROFENE ET DU TRAMADOL**

(54) **Title: PHARMACEUTICAL COMPOSITIONS CONTAINING DEXKETOPROFEN AND TRAMADOL**

(57) **Abrégé/Abstract:**

A pharmaceutical composition as a solid oral dosage form is described, comprising: i) a combination of two pharmacological active principles, dexketoprofen salt with an organic or inorganic base and tramadol salt with an organic or inorganic acid, wherein: - the organic or inorganic base is selected in the group: trometamol, trimethylamine, dimethylamine, ethylamine, triethylamine, diethylamine, L-lysine, L-arginine, diethanolamine, sodium hydroxide, calcium hydroxide - the organic or inorganic acid is selected in the group: hydrochloric, hydrobromic, phosphoric, sulfuric, methanesulfonic, benzenesulfonic, toluenesulfonic, acetic, propionic, malic, maleic, succinic, citric, L-tartaric, lactic, malonic, aspartic, glutamic; ii) microcrystalline cellulose as a filler; iii) a binder selected in the group: maize starch, pre-gelatinised maize starch, hypromellose or their mixtures; iv) pharmaceutically acceptable excipients.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 August 2014 (21.08.2014)

(10) International Publication Number
WO 2014/124862 A1

- (51) International Patent Classification:
A61K 31/135 (2006.01) *A61K 31/192* (2006.01)
- (21) International Application Number:
PCT/EP2014/052342
- (22) International Filing Date:
6 February 2014 (06.02.2014)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
MI2013A000210 14 February 2013 (14.02.2013) IT
- (71) Applicant: LABORATORIOS MENARINI SA [ES/ES];
Alfonso XII, 587, E-08918 Badalona (ES).
- (72) Inventors: SCHMITZ, Reinhard; Glinicker Weg, 125,
12489 Berlin (DE). KOHL, Tobias; Glinicker Weg, 125,
12489 Berlin (DE).
- (74) Agent: BANFI, Paolo; Bianchetti Bracco Minoja S.r.l.,
Via Plinio, 63, I-20129 Milano (IT).
- (81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,

- AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.
- (84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report (Art. 21(3))

(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING DEXKETOPROFEN AND TRAMADOL

(57) Abstract: A pharmaceutical composition as a solid oral dosage form is described, comprising: i) a combination of two pharma-
cological active principles, dexketoprofen salt with an organic or inorganic base and tramadol salt with an organic or inorganic acid,
wherein: - the organic or inorganic base is selected in the group: trometamol, trimethylamine, dimethylamine, ethylamine, triethyl-
amine, diethylamine, L-lysine, L-arginine, diethanolamine, sodium hydroxide, calcium hydroxide - the organic or inorganic acid is
selected in the group: hydrochloric, hydrobromic, phosphoric, sulfuric, methanesulfonic, benzenesulfonic, toluenesulfonic, acetic,
propionic, malic, maleic, succinic, citric, L-tartaric, lactic, malonic, aspartic, glutamic; ii) microcrystalline cellulose as a filler; iii) a
binder selected in the group: maize starch, pre-gelatinised maize starch, hypromellose or their mixtures; iv) pharmaceutically accept-
able excipients.

WO 2014/124862 A1

**PHARMACEUTICAL COMPOSITIONS CONTAINING DEXKETOPROFEN
AND TRAMADOL**

FIELD OF THE INVENTION

The invention relates to a stable pharmaceutical formulation comprising the combination of the two active pharmaceutical ingredients dexketoprofen and tramadol in a solid oral dosage form with immediate release of both active ingredients.

5 **STATE OF THE ART**

Dexketoprofen is the S-(+) enantiomer of the well known, non-steroidal anti-inflammatory drug (NSAID) and reversible cyclooxygenase (COX) inhibitor, ketoprofen, with the chemical name S-(+)-2-(3-benzoylphenyl)propionic acid. Racemic ketoprofen is used as an analgesic and anti-inflammatory agent and these effects are due to the S-(+) enantiomer (dexketoprofen), while the R-(-) enantiomer does not contribute to these activities. The tromethamine salt of dexketoprofen is registered as 12.5 mg and 25 mg film-coated immediate release oral tablets in EU countries and in several non-European countries for the symptomatic treatment of acute pain of mild to moderate intensity such as musculo-skeletal pain, dysmenorrhoea and dental pain.

15 Tramadol is the centrally acting opioid analgesic agent (1RS,2RS)-2-[(Dimethylamino)methyl]-1-(3-methoxyphenyl)cyclohexanol. Tramadol is in the market since 1977 (Tramal[®], Grünenthal). For oral use, tramadol is available as 50-300 mg capsules, tablets and sustained release formulations. Tramadol hydrochloride is the active ingredient of products with marketing authorizations in many EU countries, such as
20 Contramal[®] (Italy), Adolonta[®] (Spain), Tramal[®] and Tramundin[®] (Germany). In addition, it is marketed in combination with other active ingredients in preparations such as Zaldiar[®] containing 37.5 mg tramadol hydrochloride and 325 mg paracetamol.

The combination of analgesics has been the subject of numerous studies. The combination of a NSAID with an opioid was studied in the case of ibuprofen with
25 oxycodon and ibuprofen with codeine. The combination of 400 mg ibuprofen and 5 mg

oxycodon hydrochloride is marketed in the United States as Combunox[®] (Forest Pharmaceuticals, Inc.). As reported by Raffa et. al. in the US patent 5,516,803, the combination of tramadol and ibuprofen is synergistic for the treatment of pain and tussive conditions. The respective weight ratios were specified in the claims as 1:1 to 1:200 with
5 the preferred ratios of about 1:2 to about 1:20.

In the publication by Tuncer et al. (Eur J Gynaecol Oncol. 2003;24(2):181-4.), it was shown that clinically, ketoprofen seems to reduce morphine requirements by 33 to 40% with ketoprofen supposed central mechanism of analgesia when administered intravenously. According to Siyam et al. in Anesthesiology 2003; 99: A996, the co-
10 administration of tramadol and ketoprofen produced marked antinociceptive synergy and reduced side-effects

The rationale for developing an oral fixed combination of dexketoprofen and tramadol lies in the following considerations:

1. Dexketoprofen and tramadol have different mechanisms of action, the former
15 exerts its anti-nociceptive activity mainly at peripheral level, whereas the latter is a centrally acting analgesic. Therefore their combination is expected to result in an additive or synergistic analgesia, thus allowing a decrease in the required doses of the individual agents, and consequently a reduced risk of adverse events.

2. Dexketoprofen and tramadol have different pharmacokinetic profiles,
20 therefore their combination is expected to be characterised both by quick onset (typical of dexketoprofen) and long duration (peculiar to tramadol) of the analgesic effect.

However in the literature there are a wealth of examples concerning technical hurdles in producing adequate pharmaceutical preparations containing both ketoprofen and tramadol. For example, the common pharmaceutical excipient lactose monohydrate
25 which is used in a variety of pharmaceutical compositions for oral use, e.g. tablets and film-coated tablets, leads to severe discoloration in tablets containing dexketoprofen, thus preventing its use.

Moreover, according to the literature, tramadol hydrochloride injection is found

incompatible with acyclovir and clindamycin, while being stable when mixed with mannitol 20% or lactated Ringer's solution. Furthermore, as assessed for tramadol 50mg/mL solution for injection or infusion, precipitation occurs when tramadol solution is mixed in the same syringe with solutions containing NSAIDs such as diclofenac sodium, indometacin and piroxicam.

In EP546676 a pharmaceutical composition comprising tramadol and a propionic acid derivative NSAID, which may be ketoprofen, is described, the preferred composition consisting of tramadol with ibuprofen. The compositions described in the examples are aqueous solutions and consequently no particular problems with solid compositions were noted.

WO2007008752 describes pharmaceutical compositions comprising two drugs which may be, among others, tramadol and a NSAID; the compositions are in a solid form, but the two drugs are in different layers for separate releases.

US6294195 discloses a sustained release oral analgesic dosage form for once-day administration comprising an opioid analgesic and optionally a non-opioid drug. Tramadol and ketoprofen are included, but shown examples are referred just to morphine.

WO2008/092219 discloses solid compositions with tramadol and ketoprofen. When mixing the solid ketoprofen and tramadol, an interaction between the two active ingredients is reported, as visualized by differential scanning calorimetry (DSC), resulting in a 'thick viscous mass hard to dissolve that could result in an alteration of the bioavailability of the drugs'. The patent application further describes different options to 'vehicle tramadol and ketoprofen in pharmaceutical forms and / or products that prevent any contact between them and / or impede interaction between the two active principles' assuming that the manufacturing following standard techniques and equipment is technically not feasible and will not lead to a stable pharmaceutical form. In this context, it should be noted that the intended field of treatment, namely moderate to severe pain, particularly demands the immediate-release of the active ingredients which might be hindered by the physical interaction of ketoprofen and tramadol.

Additionally, as described by Botha et al. (1989) in Drug Development and Industrial Pharmacy 15 (3), 415-426, Botha et al. (1990) in Drug Development and Industrial Pharmacy 16 (4), 673-683, Tita et al (2011) in Journal of Pharmaceutical and Biomedical Analysis, 56, 221-227, formulation development with ketoprofen and other NSAIDs such as naproxen is not trivial. Incompatibilities were detected for ketoprofen with common pharmaceutical filler materials such as lactose, calcium phosphates, magnesium stearate or Precirol Ato 5 (glyceryl palmitostearate), which might result in increase of disintegration time, impaired dissolution rates and decomposition during stability storage.

In addition, in AU703310B2 (EP7592963) 'A rapidly disintegrating medicinal form of tramadol or a tramadol salt', the preparation of binder-free tablets containing tramadol is described, since the development of binder-containing tablets with tramadol that show fast disintegration and dissolution is considered as not possible. In detail, it was outlined that tramadol hydrochloride closes the capillaries of a tablet during disintegration due to the high water solubility of this compound and leads to the rapid formation of a highly concentrated diffusion boundary layer. Moreover, the disintegration cannot be accelerated by increasing the content of disintegrating agent.

The unfavourable characteristics of tramadol in terms of drug product development become even more pronounced when it is combined with dexketoprofen. In fact, due to its sticky nature and the tendency to agglomerate, the latter should require the further processing in the form of binder-containing granules so precluding any binder-free tablets.

DESCRIPTION OF THE INVENTION

The drawbacks associated with the technology described in WO2008/092219 are overcome by the present invention, which provides the appropriate methodology to prepare pharmaceutically acceptable solid oral formulations of dexketoprofen and tramadol, both as salts, on the basis of standard techniques and equipment without the need of specialized procedures.

From the technological point of view, the basic principles for the manufacturing of these formulations, without the complex methodology which requires the separation of both active ingredients from each other, were identified and consist in:

- The active principles are both used as salts.
- 5 - The use of an appropriate filler material.
- The use of an appropriate binder system for the granulation, inducing a suitable granule size and granule structure.
- The use of an appropriate technology for the manufacturing, particularly in the case of the granulation procedures, i.e. by fluid bed granulation leading to a soft and easily disintegrating /dissolving granule structure which is especially suitable for a solid, immediate-release dosage form.
- 10

In order to avoid the potential physical incompatibility of dexketoprofen and tramadol and the formation of a hard and sticky mass as outlined in WO2008/092219, it is necessary to follow the established principles as described above and in the examples illustrated below. The combination of the identified parameters induces a stable and technologically feasible pharmaceutical dosage form suitable for the treatment of acute pain of moderate to severe intensity originated by different causes and related to different diseases such as: post-operative pain, headache, toothache, backpain, joint pain, inflammations and migraine.

20 According to WO2008/092219 the technological formulation of dexketoprofen and tramadol should necessitate the use of specialised, intricate techniques such as multi-layered tableting, separate granulation of both active ingredients and / or the packaging in separate blister moldings in order to keep the two active ingredients rigorously separated. Surprisingly, however, the present invention shows that the manufacturing of a pharmaceutically stable formulation of these two active ingredients, suitable for the treatment of moderate to severe pain, is feasible without the application of non-standard methods and technologies. The present invention provides a technologically stable formulation containing the active principles dexketoprofen and tramadol together with

25

excipients in pharmaceutically acceptable quality, in particular fillers and binders, wherein the filler is microcrystalline cellulose and the binder is preferably selected in the group of maize starch, pregelatinised maize starch, hypromellose or their mixtures.

For the purposes of the present invention hypromellose indicates
5 hydroxypropylmethylcellulose or HPMC.

In a preferred embodiment, dexketoprofen is used as tromethamine (trometamol) salt and tramadol as the hydrochloride salt. Moreover, the production of the pharmaceutical dosage forms is performed with standard galenical processes such as mixing, sieving, granulating, tableting and film-coating while applying standard
10 technological equipment and machinery. Therefore, both active ingredients are not separated by specialized or unique manufacturing and or / packaging procedures such as tableting in multi-layers, tableting of two different granules / film-coated crystals or packaging in blisters with separated moldings.

According to the invention, dexketoprofen is in the form of a salt with organic or
15 inorganic bases and tramadol as a salt with organic or inorganic acids.

The organic or inorganic base is selected from the group of trometamol, trimethylamine, dimethylamine, ethylamine, triethylamine, diethylamine, L-lysine, L-arginine, diethanolamine, sodium hydroxide, calcium hydroxide; the preferred being trometamol.

20 For the purposes of the present invention trometamol indicates 2-amino-2-hydroxymethyl-1,3-propanediol, also known as tromethamine or TRIS.

The organic or inorganic acid is selected from the group of hydrochloric, hydrobromic, phosphoric, sulfuric, methanesulfonic, benzenesulfonic, toluenesulfonic, acetic, propionic, malic, maleic, succinic, citric, L-tartaric, lactic, malonic, aspartic,
25 glutamic; the preferred being hydrochloric.

The combination is directed to the treatment of acute and postoperative pain of moderate to severe intensity originated by different causes and related to different diseases selected from the group of headache, toothache, inflammations and migraine.

The fixed combination of both compounds is intended to allow optimised pain control with fewer side effects than observed for both compounds alone.

According to the present invention the pharmaceutical forms comprise solid oral dosage forms, preferably immediate-release film-coated tablets.

5 According to the present invention, the pharmaceutical compositions contain a mixture of the two active ingredients, for oral dosage unit, in the following amount:

- Dexketoprofen (mol wt 254.28): between 8 and 50 mg, preferably between 10 and 30 mg, 12.5 mg and 25 mg being the most preferred amounts (equivalent to dexketoprofen trometamol (mol wt 375.42) between 11.8 and 73.8 mg, preferably
10 between 14.8 and 44.3 mg, 18.4 mg and 36.9 mg being the most preferred specific amounts).

- Tramadol (mol wt 263.28): between 17.6 and 105.4 mg, preferably between 26.3 and 87.8 mg, 32.9 mg and 65.9 mg being the most preferred specific amounts [equivalent to tramadol hydrochloride (mol wt 299.84) between 20 and 120 mg, preferably between
15 30 and 100 mg, 37.5 mg and 75 mg being the most preferred amounts].

The pharmaceutical compositions contain a mixture of the two active ingredients for oral dosage unit in a w/w ratio of dexketoprofen / tramadol between 2: 1 to 1: 10, preferably in a w/w ratio between 1: 1 to 1: 6 (equivalent to a w/w ratio of dexketoprofen trometamol / tramadol hydrochloride between 2.59:1 to 1: 7.7, preferably 1.3: 1 to 1:
20 4.63).

The pharmaceutical compositions may contain in addition to the active ingredients one or more pharmaceutically acceptable excipients selected from pharmaceutical fillers, binders, disintegrants, lubricants and glidants.

The filler consists of microcrystalline cellulose and it should be contained in each
25 dosage unit from 45% to 75% w/w of the total amount.

The binder, selected in the group consisting of maize starch, pregelatinised maize starch, hypromellose or mixtures thereof, in amounts from 5% to 20%, preferably from 7% to 12% by the total weight of the composition.

The disintegrant is selected from croscarmellose sodium and sodium starch glycolate or mixtures thereof, in amounts from 4% to 14% by the total composition weight.

The lubricant is selected in the group consisting of glycerol distearate, sodium stearyl fumarate or mixture thereof, sodium stearyl fumarate being the preferred one.

The preferred glidant is anhydrous colloidal silica.

According to the present invention, all the components of the compositions are intimately mixed. With the term intimately mixed a mixing of the components, and in particular of the two active principles, is intended in such a way that a physical contact among the components of the compositions, and in particular between the two active principles, is never precluded.

Oral dosage unit may optionally be coated with pharmaceutical excipients selected from pharmaceutical film formers, pigments, dispersants and polishing agents.

Dosage units are preferably coated with hypromellose or a poly(vinyl) alcohol based outer film coating.

Not all standard pharmaceutical excipients can be used together with dexketoprofen and tramadol. The use of lactose monohydrate which is a standard pharmaceutical excipient commonly used as filler material in combination with microcrystalline cellulose (e.g. MicroceLac[®]) is not possible with dexketoprofen. The resulting tablets show yellow to orange stains indicating a chemical incompatibility and a significantly increased disintegration time after accelerated stability testing.

Also the use of the common binder polyvinylpyrrolidone (PVD) is precluded, as it leads to dull, porous tablet cores which are not suitable for further processing such as film coating, and to a strong increase in disintegration time and resistance to crushing.

Surprisingly pharmaceutical compositions containing i) a salt of dexketoprofen, trometamol being the preferred, ii) a salt of tramadol, hydrochloride being the preferred, iii) microcrystalline cellulose, as a filler, in high percentage w/w of the total amount, iv) a binder, v) one or more pharmaceutically acceptable excipients, wherein all the

constituents are intimately mixed, result to be stable in all the studied conditions.

The pharmaceutical compositions can be manufactured following standard galenical processes in which both ingredients are not separated by specialized or unique manufacturing and / or packaging procedures; they can be manufactured by fluid bed granulation or fluid bed drying in tablets or capsules for the immediate-release of both active principles.

The compositions prepared according to the present invention show a dissolution of the two active ingredients, dexketoprofen and tramadol, in a time range not superior to 15 minutes, more precisely between 3 and 11 minutes.

The compositions show a dissolution rate and a purity profile of the components which remain un-changed during shelf-life of 18 months at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ relative humidity in aluminium-aluminium and PVC/PVDC-aluminium blisters.

EXPERIMENTAL SECTION

Dexketoprofen shows a pronounced physicochemical incompatibility with tramadol. This interaction is elucidated below (Example A).

Example A: Compatibility testing of dexketoprofen trometamol and tramadol hydrochloride as dry powder mixtures

Dexketoprofen trometamol and tramadol hydrochloride were mixed together at different ratios and subjected to accelerated conditions, i.e. $(60 \pm 2)^{\circ}\text{C}$ and 60% relative humidity. The corresponding results are presented in table A1.

Table A - Appearance of dry powder mixtures containing dexketoprofen and tramadol

Composition	Results (after weeks)			
	0	1	2	4
Dexketoprofen trometamol (DKP TRIS)	n.t.	n.t.	loose powder	agglomerated powder
Tramadol hydrochloride (TRA-HCl)	n.t.	n.t.	loose powder	slightly agglutinated powder
DKP TRIS + TRA-HCl (25 mg + 37.5 mg)	n.t.	loose powder	n.t.	agglomerated powder
DKP TRIS + TRA-HCl (12.5 mg + 75 mg)	n.t.	hard and agglutinated mass at the bottom of the vial	n.t.	hard and agglutinated mass at the bottom of the vial

n.t. = not tested

These results demonstrate the physical interaction between dexketoprofen
5 trometamol and tramadol hydrochloride drug substances, especially in the case of high
tramadol doses (e.g. 75 or 100 mg).

The result obtained in EXAMPLE A makes it unlikely for the person skilled in the
art to define an effective and stable pharmaceutical composition of both components in
the form of a binder-containing tablet. As shown in the following EXAMPLES 1 to 4 of
10 the present invention, however, the selection of the right components applied in the
correct ratio and processed in the appropriate manufacturing procedures decide about the
functionality and stability of drug product. The comparative EXAMPLES B and C are
provided to illustrate this finding.

EXAMPLE B: Selection of the appropriate binder system for dexketoprofen +
15 tramadol tablets

For the definition of the appropriate binder system of dexketoprofen + tramadol
drug product, conventional pharmaceutical binders such as polyvinylpyrrolidone,
hypromellose and pregelatinized maize starch were evaluated.

The tablets were manufactured in accordance with the composition presented in
20 table B.

Table B - Composition of tablet cores used for the screening of the appropriate binder system

Ingredient	Reference	Function	Quantity, mg per tablet			
			1	2	3	4
Dexketoprofen trometamol (equivalent to dexketoprofen)	Manufacturers specification	Active ingredient	36.90 (25.0)	36.90 (25.0)	36.90 (25.0)	36.90 (25.0)
Tramadol hydrochloride	Ph. Eur.	Active ingredient	75.00	75.00	75.00	75.00
Microcrystalline cellulose	Ph. Eur.	Filler	239.10	276.54	299.94	309.30
Polyvinylpyrrolidone	Ph. Eur.	Binder	84.24	-	-	-
Maize starch, pregelatinized	Ph. Eur.	Binder	-	46.80	-	-
Hypromellose	Ph. Eur.	Binder	-	-	23.40	14.04
Croscarmellose sodium	Ph. Eur.	Disintegrant	18.72	18.72	18.72	18.72
Glycerol distearate	Ph. Eur.	Lubricant	9.36	9.36	9.36	9.36
Silica colloidal, anhydrous	Ph. Eur.	Glidant	4.68	4.68	4.68	4.68
Purified water*	Ph. Eur.	Solvent for granulation	q.s.	q.s.	q.s.	q.s.
Total tablet core weight			468.00	468.00	468.00	468.00

- 1 = binder system polyvinylpyrrolidone
- 2 = binder system pregelatinized maize starch,
- 3 = binder system hypromellose 5% w/w
- 4 = binder system hypromellose 3% w/w

The amount of polyvinylpyrrolidone was selected in accordance with previous experiences leading to a favourable particle size and granule properties. All the mixtures were sieved, granulated and subjected to tablet compaction. The appearance of the resulting tablet cores is presented in table C.

5 Table C - Appearance of tablet cores manufactured with different binder systems

Parameter	Formulation			
	1	2	3	4
Appearance of the tablet cores	dull, slightly porous	shiny, brilliant	dull, dark-spotted	slightly shining

n.t. = not tested

Subsequently, the tablet cores were subjected to accelerated stability testing at 60°C and tested for disintegration time and tablet hardness (tables E and F).

10 Table D - Disintegration time of dexketoprofen + tramadol formulations in tablets with different binder systems

Formulation	Disintegration time [min.] after weeks at 60°C		
	0	2	4
1 (polyvinylpyrrolidone)	3	16	17
2 (maize starch pregelatinized)	2	7	10
3 (hypromellose 5% w/w)	3	8	8
4 (hypromellose 3% w/w)	2	6	4

Table E - Resistance to crushing of dexketoprofen + tramadol formulations in tablets with different binder systems

Formulation	Resistance to crushing [N] after weeks at 60°C		
	0	2	4
1 (polyvinylpyrrolidone)	90	123	123
2 (maize starch pregelatinized)	134	127	116
3 (hypromellose 5% w/w)	128	120	119
4 (hypromellose 3% w/w)	165	126	122

The obtained results from binder screening underline that the number of suitable binders for dexketoprofen + tramadol containing tablets is limited. The use of the standard binder polyvinylpyrrolidone is precluded, as it leads to dull, porous tablet cores which are not suitable for further processing such as film coating, and to a strong increase of disintegration time and resistance to crushing. The increase of disintegration time was also detected for tablet cores formulated on the basis of pregelatinized maize starch and hypromellose 3% w/w and 5% w/w, but to a far lesser extent and without the increase of tablet hardness. Therefore, any binder system containing the commonly used polyvinylpyrrolidone does not alleviate the incompatibility of dexketoprofen and tramadol, but rather seems to support the negative effects of this interaction. After storage at accelerated conditions, the obtained tablet cores are no longer compliant with the requirements of Ph. Eur., i.e. disintegration time below 15 minutes.

EXAMPLE C illustrates the possible option to overcome the observed increase in disintegration time in tablet cores with maize starch, pregelatinized, by the application of an appropriate lubricant system.

EXAMPLE C: Selection of the appropriate lubricant in combination with appropriate binder system

In order to diminish the observed increase in disintegration time over shelf-life, different types of lubricant were tested in tablet cores formulated with maize starch pregelatinized. As magnesium stearate was described in literature to be incompatible with (dex)ketoprofen (2) and naproxen (3), this excipient was excluded, so that the study was conducted with glycerol distearate and sodium stearyl fumarate. The corresponding formulations and results are summarized in tables F and G.

Table F - Composition of tablet cores used for the lubricant screening

Ingredient	Reference	Function	Quantity, mg per tablet	
			1	2
Dexketoprofen trometamol (equivalent to dexketoprofen)	Manufacturers specification	Active ingredient	36.90 (25.0)	36.90 (25.0)
Tramadol hydrochloride	Ph. Eur.	Active ingredient	75.00	75.00
Microcrystalline cellulose	Ph. Eur.	Filler	253.14	253.14
Maize starch, pregelatinized	Ph. Eur.	Binder	46.80	46.80
Croscarmellose sodium	Ph. Eur.	Disintegrant	46.80	18.72
Glycerol distearate	Ph. Eur.	Lubricant	4.68	-
Sodium stearyl fumarate	Ph. Eur.	Lubricant	-	4.68
Silica colloidal, anhydrous	Ph. Eur.	Glidant	4.68	4.68
Purified water*	Ph. Eur.	Solvent for granulation	q.s.	q.s.
Total tablet core weight			468.00	468.00

Table G - Disintegration time of dexketoprofen + tramadol formulations in tablets with different types of lubricant

Formulation	Disintegration time [min.] after weeks at 60°C		
	0	1	4
1 (glycerol distearate)	2	4	5
2 (sodium stearyl fumarate)	3	2	3

The disintegration time of tablet cores prepared with 1% w/w glycerol distearate significantly increased from 2 min. to 4 min. after 1 week and to 5 min. after 4 weeks at 60°C. In contrast, tablet cores with identical parameters and identical formulation except for the use of 1% sodium stearyl fumarate instead of glycerol distearate showed no increase at all in the disintegration time (3 min. at point zero vs. 2 min. after 1 week and vs. 3 min after 4 weeks at 60°C).

In summary, the experimental EXAMPLES A to C indicate that the formulation development for dexketoprofen + tramadol tablets requires specific strategies and that the number of potential excipients that are appropriate is limited.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1, 2 and 3 are graphs showing the dissolution behaviour of both active ingredients, dexketoprofen TRIS and tramadol hydrochloride, respectively at the initial conditions and after accelerated stability tests (1 and 4 weeks at 60°C). The dissolution tests were performed, according to USP, in pH 6.8 R1 buffer, using a paddle apparatus, at a speed of 50 rpm and in a volume of 1000 ml.

Figure 4: shows the dissolution profile of the film-coated tablets of dexketoprofen TRIS and tramadol hydrochloride derived from a pilot batch production, at the initial conditions.

Figure 5: shows the dissolution profile of the film-coated tablets of dexketoprofen trometamol and tramadol hydrochloride originated from the first manufactured sub-batch

of 100Kg at the initial conditions.

Figure 1: Dissolution profile of active ingredients, initial conditions.

Figure 2: Dissolution profile of active ingredients after 1 week at 60°C.

Figure 3: Dissolution profile of active ingredients after 4 weeks at 60°C.

5 Figure 4: Dissolution profile of film-coated tablets (pilot batch), initial conditions.

Figure 5: Dissolution profile of film-coated tablets (100 kg manufactured sub-batch), initial conditions.

EXAMPLES

The examples comprise solid oral dosage forms in several combinations.

10 However, the dosage strengths which can be prepared with the present invention are not limited to these combinations only.

The detailed description of the examples is intended to illustrate the invention, but not to limit its scope.

EXAMPLE 1: Manufacturing of dexketoprofen trometamol + tramadol
15 hydrochloride tablet or film-coated tablets, with maize starch

Batch size: 4 kg

The composition and batch formulas of the (film-coated) tablets with the combination dosage strengths DKP TRIS 18.4 mg + TRA-HCl 37.5 mg (1), DKP TRIS 18.4 mg + TRA-HCl 75 mg (2), DKP TRIS 36.9 mg + TRA-HCl 37.5 mg (3) and DKP
20 TRIS 36.9 mg + TRA-HCl 75 mg (4) are given in table 1 and table 2, respectively.

Table 1 - Composition of the tablets and film-coated tablets, respectively:

Ingredient	Reference	Function	Quantity, mg per tablet			
			1	2	3	4
Tablet core						
Dexketoprofen trometamol (equivalent to dexketoprofen)	Manufacturers specification	Active ingredient	18.45 (12.5)	18.45 (12.5)	36.90 (25.0)	36.90 (25.0)
Tramadol hydrochloride	Ph. Eur.	Active ingredient	37.50	75.00	37.50	75.00
Microcrystalline cellulose	Ph. Eur.	Filler	378.85	341.35	360.40	322.90
Maize starch	Ph. Eur.	Binder	49.60	49.60	49.60	49.60
Sodium starch glycolate	Ph. Eur.	Disintegrant	27.10	27.10	27.10	27.10
Glycerol distearate	Ph. Eur.	Lubricant	5.20	5.20	5.20	5.20
Silica colloidal, anhydrous	Ph. Eur.	Glidant	3.30	3.30	3.30	3.30
Purified water*	Ph. Eur.	Solvent for granulation	q.s.	q.s.	q.s.	q.s.
Total tablet core weight			520.00	520.00	520.00	520.00
Film-coating (optional)						
Hypromellose	Ph. Eur.	Film former	4.95	4.95	4.95	4.95
Titanium dioxide	Ph. Eur.	Pigment	3.24	3.24	3.24	3.24
Povidone K 30	Ph. Eur.	Dispersant	0.87	0.87	0.87	0.87
Macrogol 4000	Ph. Eur.	Polishing agent	0.95	0.95	0.95	0.95
Purified water*	Ph. Eur.	Solvent for film-coating	q.s.	q.s.	q.s.	q.s.
Total weight of film coated tablet			530.01	530.01	530.01	530.01

* not part of the final drug product

Table 2 - Batch formulas of the tablets and film-coated tablets, respectively:

Ingredient	Reference	Quantity, g per batch			
		1	2	3	4
Tablet core					
Dexketoprofen trometamol	Manufacturers specification	141.923	141.923	283.846	283.846
Tramadol hydrochloride	Ph. Eur.	288.462	576.923	288.462	576.923
Microcrystalline cellulose	Ph. Eur.	2914.231	2625.769	2772.308	2483.846
Maize starch	Ph. Eur.	381.538	381.538	381.538	381.538
Sodium starch glycolate	Ph. Eur.	208.462	208.462	208.462	208.462
Glycerol distearate	Ph. Eur.	40.000	40.000	40.000	40.000
Silica colloidal, anhydrous	Ph. Eur.	25.385	25.385	25.385	25.385
Purified water (solvent for granulation) *	Ph. Eur.	q.s.	q.s.	q.s.	q.s.
Film coating (optional)					
Hypromellose	Ph. Eur.	38.077	38.077	38.077	38.077
Titanium dioxide	Ph. Eur.	24.923	24.923	24.923	24.923
Povidone K 30	Ph. Eur.	6.692	6.692	6.692	6.692
Macrogol 4000	Ph. Eur.	7.308	7.308	7.308	7.308
Purified water (solvent for film coating)*	Ph. Eur.	q.s.	q.s.	q.s.	q.s.

Dexketoprofen trometamol and tramadol hydrochloride are sieved together with microcrystalline cellulose and about ½ of maize starch through 1.0 mm mesh size. The compounds are mixed for 20 min. With the remaining amount of maize starch the granulation fluid is prepared: approximately 1/3 of water is mixed with the second half of

maize starch. The remaining water is heated to near the boiling point and then added to the maize starch slurry. For the wet granulation the binder fluid is stirred to 72-80°C. The granulation is performed applying standard procedures using a fluid bed granulator. It is dried to a loss on drying of 2.5 – 3.5%. Afterwards, the granules are sieved (1.0 mm mesh
5 size) and subsequently mixed with sodium starch glycolate, glycerol distearate and silica colloidal anhydrous for 10 min. The obtained compactable mixture is compressed on a standard rotary die press to tablet cores with the following properties:

- Oblong with dimensions of 17x8 mm
- Tablet mass: $520 \pm 3\%$
- 10 - Height: 5.5 ± 0.2 mm
- Resistance to crushing: ≥ 90 N
- Disintegration time: ≤ 5 min.
- Friability: $\leq 1.0\%$

The resulting tablet cores can be coated with the hypromellose film. The
15 hypromellose is added to purified water heated to near the boiling point. Afterwards, povidone K 30 and macrogol 4000 are added and the mixture is stirred until a clear solution is obtained. Directly before the film-coating, titanium dioxide is added and dispersed with a high frequency stirrer. The film-coating is performed in a standard film-coater.

20 The obtained film-coated tablets show the following properties:

- Tablet mass: $530 \pm 5\%$
- Height: 5.6 ± 0.3 mm
- Resistance to crushing: ≥ 90 N
- Disintegration time: ≤ 7.5 min.

25 Sample batches with different dexketoprofen/tramadol ratios were produced and tested under ICH conditions for 18 months at $(25 \pm 2)^\circ\text{C}$ and $(60 \pm 5)\%$ relative humidity. Results of these tests are presented in tables 3 to 6.

The obtained results indicate excellent chemical and physical stability of the

produced batches (regarding disintegration, dissolution and other tablet parameters).

Table 3 - Stability data of the film-coated tablets DKP TRIS 18.4 mg + TRA-HCl 37.5mg at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ RH in Al-Al blisters:

Test parameters	Shelf-life specification	Results (after months)					
		0	3	6	9	12	18
General and specific characteristics of the dosage form							
Dissolution (after 30 min)							
- Dexketoprofen TRIS [%]	(Q=75)	101	98	99	100	101	101
- Tramadol hydrochloride [%]	(Q=75)	102	102	101	102	102	102
Disintegration time [min]	nmt 15	3	4	5	4	3	4
Assay							
Dexketoprofen TRIS [%]	90.0 to 105.0	99.4	102.1	100.2	100.7	100.4	102.2
Tramadol hydrochloride [%]	90.0 to 105.0	99.7	101.8	102.9	102.0	101.6	105.0
Additional parameter							
Resistance to crushing							
- mean	t: ≥ 90 N	203	252	198	189	189	199

5 nmt = not more than

Table 4 - Stability data of the film-coated tablets DKP TRIS 18.4 mg + TRA-HCl 75mg at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ RH in Al-Al blisters:

Test parameters	Shelf-life specification	Results (after months)					
		0	3	6	9	12	18
General and specific characteristics of the dosage form							
Dissolution (after 30 min)							
- Dexketoprofen TRIS[%]	(Q=75)	102	100	101	101	98	102
- Tramadol hydrochloride [%]	(Q=75)	103	100	102	100	100	102
Disintegration time [min]	nmt 15	6	7	9	9	11	8
Assay							
Dexketoprofen TRIS [%]	90.0 to 105.0	99.1	101.7	101.3	103.3	100.9	101.6
Tramadol hydrochloride [%]	90.0 to 105.0	99.2	101.8	101.3	103.1	101.1	101.2
Additional parameter							
Resistance to crushing							
- mean	t: ≥ 90 N	200	224	189	194	183	192

nmt = not more than

Table 5 - Stability data of the film-coated tablets DKP TRIS 36.9 mg + TRA-HCl 37.5mg at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ RH in Al-Al blisters:

Test parameters	Shelf-life specification	Results (after months)					
		0	3	6	9	12	18
General and specific characteristics of the dosage form							
Dissolution (after 30 min)							
- Dexketoprofen TRIS [%]	(Q=75)	102	101	102	100	101	102
- Tramadol hydrochloride [%]	(Q=75)	101	100	100	100	100	101
Disintegration time [min]	nmt 15	3	5	6	3	1	3
Assay							
Dexketoprofen TRIS [%]	90.0 to 105.0	98.2	102.3	101.6	102.2	101.1	102.4
Tramadol hydrochloride [%]	90.0 to 105.0	96.8	100.5	101.4	100.8	98.7	101.4
Additional parameter							
Resistance to crushing							
- mean	t: ≥ 90 N	207	257	190	205	203	204

nmt = not more than

Table 6 - Stability data of the film-coated tablets DKP TRIS 36.9 mg + TRA-HCl 75mg at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ RH in Al-Al blisters:

Test parameters	Shelf-life specification	Results (after months)					
		0	3	6	9	12	18
General and specific characteristics of the dosage form							
Dissolution (after 30 min)							
- Dexketoprofen TRIS [%]	(Q=75)	104	103	103	n. t.	101	99
- Tramadol hydrochloride [%]	(Q=75) ⁽¹⁾	104	101	102	n. t.	103	99
Disintegration time [min]	nmt 15	3	8	9	8	7	8
Assay							
Dexketoprofen TRIS [%]	90.0 to 105.0	100.5	103.1	103.0	103.2	104.9	104.1
Tramadol hydrochloride [%]	90.0 to 105.0	100.6	102.4	103.1	103.4	102.7	102.6
Additional parameter							
Resistance to crushing							
- mean	t: ≥ 90 N	152	184	170	167	174	183

nmt = not more than

EXAMPLE 2: Manufacturing of dexketoprofen trometamol + tramadol hydrochloride film-coated tablets with pregelatinised maize starch

Batch size: 550g

Composition of the film-coated tablets:

5

Active substances	Amount per single dose unit [mg]	Amount per batch [g]
Dexketoprofen trometamol	36.90	57.99
Tramadol hydrochloride	75.00	117.86
Other ingredients		
<i>Tablet core:</i>		
Microcrystalline cellulose	179.70	282.39
Maize starch pregelatinised	29.20	45.89
Croscarmellose sodium	21.90	34.41
Sodium stearyl fumarate	3.65	5.74
Silica, colloidal anhydrous	3.65	5.74
Purified water*	q.s.	q.s.
<i>Film coating:</i>		
polyvinyl alcohol based coating	7.30	11.47
Purified water*	q.s.	q.s.

* not part of the final product

Dexketoprofen trometamol and tramadol hydrochloride are sieved together with microcrystalline cellulose and ½ of croscarmellose sodium through 1.0 mm mesh size.

The compounds are transferred to the fluid bed granulator. For the preparation of the

10 binder fluid pregelatinised maize starch is added to purified water and stirred for about 60

min. in order to obtain homogenous binder slurry. The granulation is performed applying standard procedures in a fluid bed granulator. It is dried to a loss on drying of 2.5 – 3.5%. Afterwards, the granules are sieved (1.0 mm mesh size) and subsequently mixed with croscarmellose-sodium, silica, colloidal anhydrous for 10 min. After the mixture with sodium stearyl fumarate for 5 min., the obtained compactable mixture is compressed on a standard rotary tablet press to tablet cores with the following properties:

- Oblong with dimensions of 16x7 mm
- Tablet mass: 350 mg $\pm$ 5%
- Height: 4.5 $\pm$ 0.2 mm
- Resistance to crushing: $\geq$ 90 N
- Disintegration time: $\leq$ 4 min.
- Friability: $\leq$ 1.0%

The resulting tablet cores are subsequently coated with an aqueous film coating system based on a polyvinyl alcohol motif, which is prepared according to the manufacturer's recommendation. The film-coating is performed in a standard film-coater. The resulting film-coated tablets show the following properties:

- Tablet mass: 357 mg $\pm$ 5%
- Height: 4.6 $\pm$ 0.2 mm
- Resistance to crushing: $\geq$ 90 N
- Disintegration time: $\leq$ 6 min.

The dissolution profiles of the film-coated tablets in Ph. Eur. buffer pH 6.8 R1 with paddle apparatus, at 50 rpm of stirring speed and 1000 ml of buffer volume are shown in figures 1, 2 and 3. Figure 1 presents the dissolution profiles of both active ingredients at the initial value, whereas figures 2 and 3 displays the dissolution profiles after accelerated stability testing for 1 and 4 weeks, respectively, at 60°C in a dry oven. No significant difference between the dissolution profiles was detectable, so that a physico-chemical interaction of both compounds can be excluded.

EXAMPLE 3: Manufacturing of dexketoprofen + tramadol film-coated tablets with pregelatinised maize starch as binder and croscarmellose sodium as disintegrant in pilot batch scale (12.7 kg)

The identical formulation as presented in EXAMPLE 2 was upscaled to 12.7 kg (pilot scale). The characteristics for the tablet cores are presented below:

- Oblong with dimensions of 14x6 mm
- Tablet mass: 350 mg $\pm$ 5%
- Height: 5.1 $\pm$ 0.5 mm
- Resistance to crushing: 130 $\pm$ 15 N
- Disintegration time: $\leq$ 6 min.
- Friability: $\leq$ 0.5%

The properties of the film-coated tablets are as follows:

- Tablet mass: 357 mg $\pm$ 5%
- Height: 5.2 $\pm$ 0.5 mm
- Resistance to crushing: $\geq$ 90 N
- Disintegration time: $\leq$ 15 min.

The film-coated tablets were tested for the dissolution profile and for stability under accelerated conditions at 40°C / 75% relative humidity. The results are presented in figure 4 and table 7.

Table 7 - Stability data of the film-coated tablets DKP TRIS 36.9 mg + TRA-HCl 75mg at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ RH in Al-Al blisters

Test parameters	Shelf-life specification	Results (after months)		
		0	3	6
General and specific characteristics of the dosage form				
Dissolution (after 30 min)				
- Dexketoprofen TRIS [%]	$(Q=75)^{(1)}$	98	101	98
- Tramadol hydrochloride [%]	$(Q=75)^{(1)}$	99	102	99
Disintegration time [min]	nmt 15	6	7	6
Assay				
Dexketoprofen TRIS [%]	90.0 to 105.0	101.3	99.6	99.2
Tramadol hydrochloride [%]	90.0 to 105.0	102.7	101.4	100.8
Additional parameter				
Resistance to crushing [N] ⁽⁷⁾	t: ≥ 90	191	228	233

EXAMPLE 4: Manufacturing of dexketoprofen + tramadol film-coated tablets with pregelatinised maize starch as binder and croscarmellose sodium as disintegrant in the large scale of 223 kg

The formulation from EXAMPLES 2 and 3 was slightly modified and upscaled to the large-scale batch size of 223 kg. The granulation was performed in two sub-batches.

Composition of the film-coated tablets:

Active substances	Amount per single dose unit [mg]	Amount per granulation sub-batch [kg]	Amount per batch [kg]
Dexketoprofen trometamol	36.90	11.121	22.241
Tramadol hydrochloride	75.00	22.603	45.206
Other ingredients			
<i>Tablet core:</i>			
Microcrystalline cellulose	179.70	54.157	108.314
Maize starch pregelatinised	29.20	8.800	17.600
Croscarmellose sodium (inner phase)	11.02	3.321	6.642
Croscarmellose sodium (outer phase)	22.04	-	13.292
Sodium stearyl fumarate	1.83	-	4.442
Silica, colloidal anhydrous	7.37	-	1.103
Purified water*	q.s.	-	
<i>Film coating:</i>			
polyvinyl alcohol based coating	7.30	-	4.400
Purified water*	q.s.	-	q.s.

For the first sub-batch of 100 kg, dexketoprofen trometamol, tramadol hydrochloride, microcrystalline cellulose and croscarmellose-sodium for the inner phase

5 mixed and sieved together through 1.0 mm mesh size. The compounds are transferred to

the fluid bed granulator. For the preparation of the binder fluid pregelatinised maize starch is added to purified water and stirred for about 60 min. in order to obtain a homogenous binder slurry. The granulation is performed applying standard procedures in a fluid bed granulator. It is dried to a loss on drying of 2.5 – 3.5%. Afterwards, the granules are sieved (1.0 mm mesh size) and subsequently mixed with croscarmellose-sodium and silica, colloidal anhydrous for 10 min. The second sub-batch is prepared following identical procedures and then added to the first sub-batch. Sodium stearyl fumarate is sieved and added to the mixture of both sub-batches. The obtained compactable mixture is compressed on a standard rotary tablet press to tablet cores with the following properties:

- Oblong with dimensions of 14x6 mm
- Tablet mass: 363 mg $\pm$ 2%
- Height: 5.1 $\pm$ 0.2 mm
- Resistance to crushing: 115 - 150 N
- Disintegration time: < 5 min.
- Friability: $\leq$ 0.05%

The properties of the film-coated tablets are as follows:

- Tablet mass: 370 mg $\pm$ 5%
- Height: 5.2 $\pm$ 0.2 mm
- Resistance to crushing: > 90 N
- Disintegration time: $\leq$ 6 min.

Table 8 - Stability data of the film-coated tablets DKP TRIS 36.9 mg + TRA-HCl 75mg at $(40 \pm 2)^{\circ}\text{C}$ and $(75 \pm 5)\%$ RH in Al-Al blisters

Test parameters	Shelf-life specification	Results (after months)		
		0	3	6
General and specific characteristics of the dosage form				
Dissolution (after 30 min)				
- Dexketoprofen TRIS [%]	(Q=75)	100	98	98
- Tramadol hydrochloride [%]	(Q=75)	100	98	100
Disintegration time [min]	nmt 15	8	8	9
Assay				
Dexketoprofen TRIS [%]	90.0 to 105.0	98.7	99.5	98.8
Tramadol hydrochloride [%]	90.0 to 105.0	100.5	100.1	100.0
Additional parameter				
Resistance to crushing [N]	t: ≥ 90	221	255	266

Table 9 - Stability data of the film-coated tablets DKP TRIS 36.9 mg + TRA-HCl 75 mg at $(30 \pm 2)^{\circ}\text{C}$ and $(75 \pm 5)\%$ RH in Al-Al blisters

Test parameters	Shelf-life specification	Results (after months)		
		0	3	6
General and specific characteristics of the dosage form				
Dissolution (after 30 min)				
- Dexketoprofen TRIS [%]	(Q=75)	100	94	96
- Tramadol hydrochloride [%]	(Q=75)	100	95	95
Disintegration time [min]	nmt 15	8	8	8
Assay				
Dexketoprofen TRIS [%]	90.0 to 105.0	98.7	100.1	97.6
Tramadol hydrochloride [%]	90.0 to 105.0	100.5	100.8	99.9
Additional parameter				
Resistance to crushing [N] ⁽⁷⁾	t: ≥ 90	221	232	245

The dissolution profile of the obtained film-coated tablets is shown in figure 5 and is comparable to that of the sample and pilot size batches.

CLAIMS

1. A pharmaceutical composition in a solid oral dosage form, comprising:
 - i) a combination of the two active pharmaceutical ingredients, dexketoprofen as a
5 salt with organic or inorganic bases and tramadol as a salt with organic or inorganic acids, wherein:
 - the organic or inorganic base is selected from the group consisting of trometamol, trimethylamine, dimethylamine, ethylamine, trimethylamine, diethylamine, L-lysine, L-arginine, diethanolamine, sodium hydroxide,
 - 10 - the organic or inorganic acid is selected from the group consisting of hydrochloric, hydrobromic, phosphoric, sulfuric, methanesulfonic, benzenesulfonic, toluenesulfonic, acetic, propionic, malic, maleic, succinic, citric, L-tartaric, lactic, malonic, aspartic, glutamic;
 - ii) a filler which is microcrystalline cellulose;
 - 15 iii) a binder selected from the group consisting of maize starch, maize starch pre-gelatinized and hypromellose, or mixtures thereof;
 - iv) a pharmaceutically acceptable excipient.
2. A pharmaceutical composition according to claim 1, wherein the two active
20 ingredients form a homogeneous mixture in which said active ingredients are intimately mixed.
3. A pharmaceutical composition according to claim 1, wherein the organic base is trometamol and the inorganic acid is hydrochloric acid.
4. A pharmaceutical composition according to claim 1, wherein said pharmaceutical excipient is selected from disintegrant, lubricant and glidant.
- 25 5. A pharmaceutical composition according to claims 1-4, containing a mixture of the two active ingredients in a weight ratio of dexketoprofen to tramadol from 2:1 to 1:10, equivalent to a weight ratio of dexketoprofen trometamol to tramadol hydrochloride from 2.59:1 to 1:7.7.

6. A pharmaceutical composition according to claim 5, containing a mixture of the two active ingredients in a weight ratio of dexketoprofen to tramadol from 1:1 to 1:6, equivalent to a weight ratio of dexketoprofen trometamol to tramadol hydrochloride from 1.3:1 to 1:4.63.

5 7. A pharmaceutical composition according to claims 1-4 containing the two active ingredients, dexketoprofen and tramadol or salts thereof, in the following amounts for each oral dosage unit:

- dexketoprofen from 8 to 50 mg, equivalent to dexketoprofen trometamol from 11.8 to 73.8 mg;

10 - tramadol from 17.6 to 105.4 mg, equivalent to tramadol hydrochloride from 20 to 120 mg.

8. A pharmaceutical composition according to claim 7, containing the two active ingredients, dexketoprofen and tramadol or salts thereof, in the following amounts, for each oral dosage unit:

15 - dexketoprofen from 10 to 30 mg, equivalent to dexketoprofen trometamol from 14.8 to 44.29 mg;

- tramadol from 26.3 to 87.8 mg, equivalent to tramadol hydrochloride from 30 to 100 mg.

20 9. A pharmaceutical composition according to claims 1-8, containing the two active ingredients, dexketoprofen and tramadol or salts thereof, in the following specific amounts for each oral dosage unit:

- dexketoprofen 12.5 mg or 25 mg, equivalent to dexketoprofen trometamol 18.45 mg or 36.95 mg;

25 - tramadol 32.9 mg or 65.9 mg, equivalent to tramadol hydrochloride 37.5 mg or 75 mg.

10. A pharmaceutical composition according to claims 1-9, wherein microcrystalline cellulose is in a concentration range from 45 to 75% by weight of the total amount.

11. A pharmaceutical composition according to claims 1-10, wherein the binder is

selected from the group consisting of maize starch, maize starch pre-gelatinized and hypromellose or mixtures thereof, in amounts from 5% to 20%, preferably from 7% to 12% by the total weight of the composition.

12. A pharmaceutical composition according to claims 1-11, wherein the disintegrant
5 is selected from croscarmellose sodium and sodium starch glycolate or mixtures thereof, in amounts from 4% to 14% by the total composition weight.

13. A pharmaceutical composition according to claims 1-12, wherein the dissolution of the two active ingredients, dexketoprofen and tramadol, occurs in a time range not higher than 15 minutes, preferably between 3 and 11 minutes.

10 14. A pharmaceutical composition according to claims 1 to 13, wherein the dissolution rate and the purity profile of the components remain unchanged during shelf-life of 18 months at $(25 \pm 2)^{\circ}\text{C}$ and $(60 \pm 5)\%$ relative humidity in aluminium - aluminium and PVC/PVDC packages.

15 15. A pharmaceutical composition according to claims 1-14, which is in the form of tablet for immediate release of the active principles.

16. A pharmaceutical composition according to claim 15, wherein said tablet comprises a hypromellose based outer film coating.

17. A pharmaceutical composition according to claim 15, comprising a polyvinyl alcohol based outer film coating.

20 18. A pharmaceutical composition according to claims 1-14, which is in the form of a capsule for the immediate-release of both active principles.


19. A pharmaceutical composition according to claims 1-18, for the treatment of acute pain related to the following conditions: headache, toothache, inflammations and migraine.

25 20. A method of manufacturing a pharmaceutical composition in accordance with claims 1 to 18, said method comprising fluid bed granulation and fluid bed drying of a mixture of active principles and excipients.


Sheet 1/5

Figure 1

A) Dexketoprofen TRIS




B) Tramadol HCl




Sheet 2/5

Figure 2

A) Dexketoprofen TRIS




B) Tramadol HCl




Sheet 3/5

Figure 3

A) Dexketoprofen TRIS




B) Tramadol HCl




Sheet 4/5

Figure 4

A) Dexketoprofen TRIS




B) Tramadol HCl



Sheet 5/5

Figure 5

A) Dexketoprofen



B) Tramadol

