



HU000025361T2



(19) **HU**

MAGYARORSZÁG
Szellemi Tulajdon Nemzeti Hivatala

(11) Lajstromszám: **E 025 361**

(13) **T2**

EURÓPAI SZABADALOM **SZÖVEGÉNEK FORDÍTÁSA**

(21) Magyar ügyszám: **E 12 002626**

(51) Int. Cl.: **A01N 43/56**

(2006.01)

(22) A bejelentés napja: **2005. 02. 21.**

A01N 51/00

(2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20050002626

(97) Az európai bejelentés közzétételi adatai:

EP 2484209 A1 2012. 08. 08.

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2484209 B1 2015. 04. 01.

(30) Elsőbbségi adatai:

2004048376

2004. 02. 24.

JP

2004158349

2004. 05. 27.

JP

(73) Jogosult(ak):

**Sumitomo Chemical Company, Limited, Tokyo
104-8260 (JP)**

(72) Feltaláló(k):

Ohkawara, Yuichi, Hyogo 666-0136 (JP)

(74) Képviselő:

**dr. Fehérvári Flóra, DANUBIA Szabadalmi és
Jogi Iroda Kft., Budapest**

(54)

Rovarirtó készítmények

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



EP 2 484 209 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:

01.04.2015 Bulletin 2015/14

(51) Int Cl.:

A01N 43/56 (2006.01)

A01N 51/00 (2006.01)

(21) Application number: **12002626.5**

(22) Date of filing: **21.02.2005**

(54) Insecticide compositions

Insektizidzusammensetzungen

Compositions d'insecticide

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**

(30) Priority: **24.02.2004 JP 2004048376**

27.05.2004 JP 2004158349

(43) Date of publication of application:

08.08.2012 Bulletin 2012/32

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:

05719327.8 / 1 719 409

(73) Proprietor: **Sumitomo Chemical Company,
Limited**

Tokyo 104-8260 (JP)

(72) Inventor: **Ohkawara, Yuichi
Hyogo 666-0136 (JP)**

(74) Representative: **Witte, Weller & Partner
Patentanwälte mbB
Postfach 10 54 62
70047 Stuttgart (DE)**

(56) References cited:
**WO-A1-03/015519 WO-A1-03/024222
WO-A1-03/027099 WO-A1-2005/048711**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 484 209 B1

Description**TECHNICAL FIELD**

5 [0001] The present invention relates to compositions exhibiting excellent insecticidal action, which comprise the compounds represented by the general formula [Ia] or salts thereof and thiamethoxam and to methods for controlling insect pests by use of a mixture of said compounds.

BACKGROUND ART

10 [0002] The compounds represented by the general formula [Ia] which are usable in the present invention are the known compounds exhibiting insecticidal action (refer to Patent Literature Nos. 1, 2 and 3).

[0003] Also, the neonicotinoid compound thiamethoxam which is utilizable in the present invention, is a compound known to have insecticidal action, and is, thiamethoxam (chemical name: 3-(2-chloro-1,3-thiazol-5-yl-methyl)-5-methyl-1,3,5-oxa-diazinan-4-ylidene(nitro)amine; Non-Patent Literature No. 1, No. 751, page 896; Patent Literature No. 7).

15 [0004] As examples of the compounds which can be incorporated into the compounds represented by the formula [Ia], there is indicated the neonicotinoid compound as described above, together with a large number of other insecticides (Patent Literature Nos. 1 to 3), but no description is given on the working examples in which such compounds are put into practical use as a mixture with the neonicotinoid compound thiamethoxam.

20 Patent Literature No. 1; Pamphlet of WO 01/070671
 Patent Literature No. 2; Pamphlet of WO 03/015519
 Patent Literature No. 3; Pamphlet of WO 03/016284
 Patent Literature No. 4; Gazette of JP-A-Hei 3-157308
 25 Patent Literature No. 5; Gazette of JP-A-Hei 2-000171
 Patent Literature No. 6; Gazette of JP-A-Sho 61-178981
 Patent Literature No. 7; Gazette of JP-A-Hei 6-183918
 Patent Literature No. 8; Gazette of JP-A-Hei 4-154741
 Patent Literature No. 9; Gazette of JP-A-Hei 7-179448
 30 Patent Literature No. 10; Gazette of JP-A-Sho 62-207266
 Non-Patent Literature No. 1; Pesticide Manual, 12th Edition, British Crop Protection Council.

35 [0005] Also, WO 2005/048711 A1 discloses specific combinations of insecticidal active substances, containing compounds which are effective as insecticides on nicotinic acetyl choline receptors as agonists and antagonists, and at least one active substance from the group of antrilic acid amides, which are suitable for protecting plants against pests. Further, WO 03/027099 A1 and WO 03/024222 A1 disclose antranilamides and their use as arthropodicides in agronomic and nonagronomic environments; further, these documents suggest to use such antranilamides in combination with biologically active substances.

40 DISCLOSURE OF THE INVENTION**PROBLEMS TO BE SOLVED BY THE INVENTION**

45 [0006] In recent years, the environmental pollution with a variety of chemical substances have been grasped as a global-scale problem, and the intensified social demand exists for suppressing the release of chemical substances into the environment to a minimum level. In the field of agriculture, also, diverse studies have been conducted to exploit a method for controlling harmful and injurious organisms, or pests, without use of chemical substances, such as the creation or generation of genetically modified crops, the control of pests with their natural enemies and the physical control of pests, etc.

50 [0007] However, such methods for controlling pests without use of chemical substances are encountered with numerous problems, such as their uniquely limited, sole use in controlling specific diseases or insect pests, instability of the effect, and the like, and the need for controlling pests with chemical substances therefore has not yet been reduced.


MEANS FOR SOLVING THE PROBLEM

55 [0008] The present inventors, with a specific view to reduction in application rate of agricultural chemicals or pesticides from the standpoint of prevention of the environmental pollution, etc., conducted repeated intensive research, and as a result, found that a mixture of a compound represented by the general formula [Ia] with thiomethoxam can produce a

greater effect than would be expected when both of the compounds individually are applied solely, thus enabling a reduction in the rate or number of application of agrochemicals or pesticides to be realized. Also, it was found that such mixtures, even when applied to locations other than the sites on which insect pests inflict injuries directly, such as seeds, seed potatoes or soils of nursery beds or farms on which crops are grown, and the like, can control insect pests in an extremely effective manner. Such findings, followed by continued intensive investigation, led to completion of the present invention.

5 [0009] Namely, the present invention relates to

10 [1] an insecticide composition which comprises thiomethoxam and one or not less than two kinds of compounds being selected from a compound represented by the general formula [Ia]:



wherein ;

30 R¹ is a halogen atom or a C₁₋₆ haloalkyl group, R² is a halogen atom, R³ and R⁵ each are a C₁₋₆ alkyl group, R⁴ is a hydrogen or halogen atom, and X is N, or a salt thereof, with the exceptions as indicated in the attached claims, where applicable;

35 [2] an insecticide composition according to the above-mentioned [1], wherein in the compound represented by the general formula [Ia], R¹ is a chlorine or bromine atom or a trifluoromethyl group, R² is a chlorine atom, R³ is a methyl group, R⁵ is an isopropyl group, R⁴ is a hydrogen or chlorine atom, and X is N;

40 [3] an insecticide composition according to the above-mentioned [2], wherein the compound represented by the formula [Ia] is 2-[1-(3-chloropyridin-2-yl)-3-trifluoro-methylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[3-bromo-1-(3-chloropyridin-2-yl)pyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide or 2-[3-bromo-1-(3-chloropyridin-2-yl)pyrazol-5-ylcarbonylamino]-5-chloro-N-isopropyl-3-methylbenzamide, with the exceptions as indicated in the attached claims, where applicable;

45 [4] a method for controlling an insect pest, which comprises applying the insecticide composition as described in any one of the above-mentioned [1] to [3] to locations other than the site where the insect pest inflict injuries directly;

50 [5] a method for controlling an insect pest, characterized in that said method comprises mixing two kinds of the compounds, namely a compound represented by the general formula [Ia] or a salt thereof as described in any one of the above-mentioned [1] to [3] and thiomethoxam followed by drenching onto the soil for raising seedlings in the form of a mixture solution or application on the soil for raising seedlings in the form of a mixture granule, during the period ranging from the sowing time to the seedling-planting time for a crop to be cultivated by the seedling-planting method;

55 [6] a method for controlling an insect pest, characterized in that said method comprises raising seedlings with use of the soil for raising seedlings which has contained therein two kinds of the compounds, namely a compound represented by the general formula [Ia] or a salt thereof as described in any one of the above-mentioned [1] to [3] and thiomethoxam during the period ranging from the sowing time to the seedling-planting time for a crop to be cultivated by the seedling-planting method;

[7] a method for controlling an insect pest, characterized in that said method comprises applying two kinds of the compounds, namely a compound represented by the general formula [Ia] or a salt thereof as described in any one of the above-mentioned [1] to [3] and thiomethoxam, to the soil of a farm field through drenching treatment, planting-

hole treatment, planting-hole treated soil incorporation, plant-root treatment or plant-root treated soil incorporation during the period ranging from the seedling-planting time to the vegetation period for a crop to be cultivated by the seedling- planting method;

[8] a method for controlling an insect pest, characterized in that said method comprises effecting the soaking treatment, dusting treatment or coating treatment of a seed, seed potato or bulb with two kinds of the compounds, namely a compound represented by the general formula [Ia] or a salt thereof as described in any one of the above-mentioned [1] to [3] and thiomethoxam, in the case of a crop to be cultivated by directly sowing or seeding a seed, seed potato or bulb on the farm field; and

[9] a method for controlling an insect pest, characterized in that said method comprises applying two kinds of the compounds, namely a compound represented by the general formula [Ia] or a salt thereof as described in any one of the above-mentioned [1] to [3] and thiomethoxam, to the soil of a farm field through drenching treatment, plant-root treatment or plant-root treated soil incorporation during the vegetation period of a crop to be cultivated by directly sowing or seeding a seed, seed potato or bulb on the farm field.

15 EFFECT OF THE INVENTION

[0010] A combination of a compound represented by the general formula [Ia] with thiomethoxam can result in the development or elicitation of a higher insecticidal effect than would be expected when each of the compounds is applied solely, namely the synergistic effect, thus enabling reductions in the rate or number of application of agrochemicals to be realized.

BEST MODE FOR CARRYING OUT THE INVENTION

[0011] The compounds [Ia] or salts thereof and thiomethoxam in some instances exist in the forms of geometrical isomers and/or stereoisomers, and the present invention include such individual isomers and mixtures thereof.

[0012] Referring to the above-illustrated formula, as the C₁₋₆ alkyl group represented by R³ and R⁵, for example, there is used methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, or tert-butyl, etc.

[0013] As the C₁₋₆ haloalkyl group represented by R¹, for example, there are used C₁₋₆ alkyl groups substituted by 1 to 10 (preferably 1 to 5) of halogen atoms (e.g., fluorine, chlorine, bromine, iodine), such as chloromethyl, fluoromethyl, bromomethyl, 2-chloroethyl, dichloromethyl, trichloromethyl, trifluoromethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, heptafluoropropyl, or nonafluorobutyl, etc.

[0014] As the halogen atom represented by R¹ R² and R⁴, there is used a fluorine, chlorine, bromine or iodine atom.

[0015] R¹ is a halogen atom or C₁₋₆ haloalkyl group, with chlorine and trifluoromethyl being more preferred.

[0016] R² is a halogen atom, with chlorine being particularly preferable. Their position of substitution is the 3-position of the mother cyclic group being a pyridyl group.

[0017] R³ is a C₁₋₆ group which is situated for substitution at the 3-position (the position of substitution refers to a position relative to 2-aminobenzoic acid used as a base nucleus), with a 3-methyl group being particularly preferred.

[0018] R⁴ is a hydrogen atom, or a halogen atom, with a hydrogen atom and a 5-chlorine being particularly preferable.

[0019] R⁵ is a C₁₋₆ alkyl group, with an isopropyl group being particularly preferable.

[0020] X represents N.

[0021] As the compound represented by the general formula [Ia], preferable are 2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamo]N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamo]N-isopropyl-3-methylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamo]N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamo]N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonyl-amino]N-isopropyl-3-methylbenzamide, 2-[3-bromo-1-(3-chloropyridin-2-yl)pyrazol-5-ylcarbonylamo]N-isopropyl-3-methylbenzamide or 2-[3-bromo-1-(3-chloropyridin-2-yl)pyrazol-5-ylcarbonylamo]N-isopropyl-3-methylbenzamide, etc., with the exceptions as indicated in the attached claims, where applicable.

[0022] The salt of the compound [Ia] may be any salts, so long as they are agrochemically acceptable salts. Such salts are exemplified by salts to be formed with inorganic bases (e.g., alkali metals, such as sodium, potassium and lithium, etc., alkaline earth metals, such as calcium and magnesium, etc., ammonia and the like), organic bases (e.g., pyridine, collidine, triethylamine, triethanolamine, etc.), inorganic acids (e.g., hydrochloric acid, hydrobromic acid, hydroiodic acid, phosphoric acid, sulfuric acid, perchloric acid, etc.) or organic acids (e.g., formic acid, acetic acid, tartaric acid, malic acid, citric acid, oxalic acid, succinic acid, benzoic acid, picric acid, methanesulfonic acid, p-toluenesulfonic acid, etc.).

[0023] The compound [Ia] or a salt thereof can be produced, for example, by the methods described in WO01/070671, WO03/015519 and WO03/016284 or methods similar thereto.

[0024] The neonicotinoid compound thiamethoxam can be produced, for example, by the methods as described

individually in JP-A-Hei 3-157308 (Patent Literature No. 4), JP-A-Hei 2-000171 (Patent Literature No. 5), JP-A-Sho 61-178981 (Patent Literature No. 6), JP-A-Hei 6-183918 (Patent Literature No. 7), JP-A-Hei 4-154741 (Patent Literature No. 8), JP-A-Hei 7-179448 (Patent Literature No. 9) and JP-A-Sho 62-207266 (Patent Literature No. 10), or the similar methods thereto.

5 [0025] In utilizing the compositions of the present invention as an agrochemical preparation, such as insecticides, insecticide-acaricide combinations and fungicide-insecticide combinations, etc., one or not less than two kinds (preferably one kind) of the compounds [Ia] or their salts and thiomethoxam, depending upon the object of use, are dissolved or suspended in a suitable liquid carrier, or are admixed with, or adsorbed on, a suitable solid carrier to thereby be used in the forms which the common agrochemicals can take, namely, by way of the formulations, such as wettable powders, aqueous suspensions, emulsions or emulsifiable concentrates, soluble liquids or solutions, ULV preparations, dusts, granules, tablets, jumbo preparations, pastes, foamable preparations, aerosols, microcapsules, coating preparations for seeds, fumigants, smokes, stick preparations for drenching of crops, or oil preparations, etc. If necessary, these agrochemical preparations may suitably be incorporated with ointment bases, emulsifiers, suspending agents, spreaders, penetrants, wetting agents, dispersing agents, stabilizers, binders, fluidizing auxiliary agents, flocculants, antioxidants, floating agents, anti-foaming agents, anti-freezing agents, preservatives, dehydrating agents, UV absorbers, UV scattering agents, coloring agents or suspension stabilizers, etc., and can be prepared by the procedures known per se. Namely, such preparations can be manufactured by mixing uniformly the compound [I] or its salt and the neonicotinoid compound [II] with a liquid or solid carrier as well as the above-described various additives, etc.

10 [0026] For example, an emulsion or emulsifiable concentrate can be manufactured by mixing for dissolution uniformly the compound [Ia] or its salt and thiomethoxam as well as an emulsifier and an organic solvent, etc. For example, a granule, a granular wettable powder and the like can be manufactured by mixing uniformly the compound [Ia] or its salt and thiomethoxam, as well as a dispersing agent (surfactant), binder and extender (or solid carrier), etc., followed by granulation. For example, a dust (e.g. a DL dust, etc.) can be manufactured by mixing for pulverizing uniformly the compound [Ia] or its salt, thiomethoxam and an extender (or solid carrier), etc. For example, a flowable preparation can be manufactured by mixing for dispersion the ingredients, such as the compound [Ia] or its salt and thiomethoxam, a dispersing agent, etc. with use of a stirring machine, followed by wet pulverization with Dyno-Mill, etc. For example, a jumbo preparation can be manufactured by mixing uniformly the compound [Ia] or its salt, thiomethoxam, and a dispersing agent (surfactant), binder, floating agent and extender (or solid carrier), etc., followed by granulation.

15 [0027] Suitable examples of the liquid carrier (e.g., solvents, organic solvents) to be used include solvents, such as water, alcohols (e.g., methyl alcohol, ethyl alcohol, n-propyl alcohol, isopropyl alcohol, ethylene glycol, etc.), ketones (e.g., acetone, methyl ethyl ketone, etc.), ethers (e.g., dioxane, tetrahydrofuran, ethylene glycol monomethyl ether, diethylene glycol monomethyl ether, propylene glycol monomethyl ether, etc.), aliphatic hydrocarbons (e.g., kerosine, paraffin oil, fuel oil, machine oil, etc.), aromatic hydrocarbons (e.g., benzene, toluene, xylene, solvent naphtha, methyl-naphthalene, etc.), halogenated hydrocarbons (e.g., dichloromethane, chloroform, carbon tetrachloride, etc.), acid amides (e.g., N,N-dimethylformamide, N,N-dimethylacetamide, etc.), esters (e.g., ethyl acetate, butyl acetate, fatty acid esters of glycerol, etc.) or nitriles (e.g., acetonitrile, propionitrile, etc.), and the like, and these can suitably be used by mixing one or two or more kinds (preferably not less than one but not more than three kinds) in appropriate ratios.

20 [0028] As the solid carrier (i.e., diluents and extenders), vegetable powders (e.g., soybean meal, tobacco meal, wheat flour, wood powder, etc.), mineral powders (e.g., kaolin, bentonite, sepiolite, clays such as acid clay, etc., talc such as talc venetum and powdered talc, etc., silica such as diatomaceous earth and powdered mica, etc., water-soluble materials such as lactose, ammonium sulfate, urea, sodium hydrogen carbonate, sodium thiosulfate, disodium hydrogenphosphate, sodium acetate and sodium carbonate, etc., and the like), calcium carbonate, alumina, powdered sulfur or activated carbon, etc. and these can suitably be used by mixing one or not less than two (preferably not less than one but not more than three kinds) in appropriate ratios.

25 [0029] As the ointment base, for example, there can be appropriately used one or not less than two kinds (preferably not less than one but not more than three kinds) of polyethylene glycol, pectin, polyhydric alcohol esters of higher fatty acids, such as glycerol monostearate, etc., cellulose derivatives, such as methylcellulose, etc., sodium alginate, bentonite, higher alcohols, polyhydric alcohols, such as glycerol, etc., petrolatum, white petrolatum, liquid paraffin, lard, a variety of vegetable oils, lanolin, dehydrated lanolin, hardened oils, resins and the like, or these ointment bases being additionally incorporated with one or two or more kinds (preferably not less than one but not more than four kinds) of the below-described various surfactants.

30 [0030] Referring particularly to the surfactants which are usable as an emulsifier, spreader, penetrant, wetting agent or dispersing agent, etc., the non-ionic surfactants used include, for example, soaps, polyoxyethylene alkylene ethers (New Calgen D1504, Neugen ET65, Neugen ET83, Neugen ET157, etc.), polyoxyethylene alkylaryl ethers (Neugen EA92, Neugen EA142, etc.), polyoxyethylene alkylphenyl ethers, polyoxyethylene nonylphenyl ethers (Nonipol 20, Nonipol 100, etc.), polyoxyethylene polyoxypropylene ethers, polyoxyethylene distyrenated phenyl ether (Neugen EA87, Neugen EA177, etc.), polyoxyethylene alkyl esters (Ionnet MO20, Ionnet MO600, etc.), sorbitan fatty acid esters (Reodol SP-S10, Reodol TW-S20, etc.), polyoxyethylene sorbitan fatty acid esters, block copolymers of ethylene oxide and

propylene oxide (Newpole PE64), alkanol amides of higher fatty acids, alkyl maleate copolymers (Demol EP), polyhydric alcohol esters (Tween 20, Tween 80, etc.) and the like, and the cationic surfactants used include, for example, alkylamine salts, quaternary ammonium salts, and the like, while the anionic surfactants used include, for example, high-molecular compounds, such as metal salts of naphthalene sulfonate condensates, naphthalene sulfonate formalin condensates (NewCalgen FS4, etc.), alkylnaphthalene sulfonates (Sorpol 5115, etc.), metal ligninsulfonates, alkylallyl sulfonates and alkylallyl sulfonate sulfates, etc., sodium polystyrenesulfonates, metal salts of polycarboxylic acids, ammonium polyoxoethylene histidylphenyl ether sulfates, higher alcohol sulfonates, higher alcohol ether sulfonates, dialkyl sulfosuccinates (NewCalgen EP70P, etc.) or alkali metal salts of higher fatty acids, and the like.

5 [0031] As the stabilizer, use is made of compounds having epoxy groups, antioxidants (e.g., dibutylhydroxytoluene (BHT), butylhydroxyanisole (BHA), tetrakis[3-(3,5-di-tert-butyl-4-hydroxyphenyl)propionyloxymethyl]methane (Irganox 1010), DL-tocopherol, propyl gallate, erysorbic acid, sodium erysorbate, isopropyl citrate, etc.), phosphoric acid, PAP auxiliary agent (isopropyl acid phosphate), cyclodextrin (Toyoderin P), or tall oil fatty acids (Hartall fatty acids), etc., and these can appropriately be used by mixing one or two or more kinds (preferably not less than one but not more than three kinds) of them in suitable ratios.

10 [0032] As the binder, use is made of dextrin, alpha-starch, polyvinyl alcohol, gum arabic, sodium alginate, polyvinylpyrrolidones, glucose, sucrose, mannitol or sorbitol, etc., and these can suitably be used by mixing one or two or more kinds (preferably not less than one but not more than three kinds) of them in appropriate ratios.

15 [0033] As the fluidizing auxiliary agent, use is made of PAP auxiliary agent (e.g., isopropyl acid phosphate), talc, etc., and these can appropriately be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) in suitable ratios.

20 [0034] As the anti-coagulating agent, use is made of white carbon, diatomaceous earth, magnesium stearate, aluminum oxide or titanium dioxide, etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

25 [0035] As the flocculant, use is made of liquid paraffin, ethylene glycol, diethylene glycol, triethylene glycol or isobutylene polymer (e.g., IP Solvent), etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

30 [0036] As the antioxidant, use is made of dibutylhydroxytoluene, 4,4-thiobis-6-tert-butyl-3-methylphenol, butylhydroxyanisole, para-octylphenol, mono(or di- or tri-) (α-methylbenzyl)phenol, 2,6-di-tert-butyl-4-methylphenol or tetrakis[3-(3,5-di-tert-butyl-4-hydroxyphenyl)propionyloxymethyl]methane, etc. and these can appropriately be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in suitable ratios.

35 [0037] The floating agent, which is particularly utilized for the manufacture of jumbo preparations, preferably includes, for example, powdered bases with a specific gravity of not more than 1 (preferably 1 to 0.5), and the like. The said powdered bases are preferably exemplified by those with a particle size of not more than 600 µm, preferably 600 µm to 10 µm; the inorganic ones are naturally-occurring glassy or vitreous materials being provided with one or several independent air bubbles through the burning treatment, and include, for example, perlite composed of perlite and obsidian, Shirasu Balloons (tradename) composed of Shirasu, vermiculite composed of vermiculite rock, etc. as well as Phylite (tradename), or microsized hollow materials based on aluminosilicate as produced through the burning treatment, etc., while the organic ones can be exemplified by higher fatty acids commonly called "waxy substances" which are in the solid state at ambient temperature, such as stearic acid and palmitic acid, as well as higher alcohols, such as stearyl alcohol, paraffin wax and the like. Since such waxy substances are water-repellent and resistant to water-penetration, there develops the likelihood that the active ingredients of agrochemicals tend to be entrapped in such waxy substances over an extended period of time and get difficult to be dispersed into water, and consequently, such active ingredients are preferably used by mixing with the above-described glassy or vitreous hollow materials.

40 [0038] As the anti-foaming agent, use is made of silicone-based anti-foaming agents (e.g., Antifoam E20, etc.) and the like, and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

45 [0039] As the anti-freezing agent, use is made of ethylene glycol, diethylene glycol, polyethylene glycol or glycerol, etc., and these can suitably be utilized by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

50 [0040] As the preservative, use is made of butylparaben or potassium sorbate, etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

55 [0041] As the dehydrating agent, use is made of anhydrous gypsum, silica gel powders, etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

[0042] As the UV absorber, use is made of 2-(2'-hydroxy-5'-methylphenyl)benzotriazole, 2-ethoxy-2'-methyloxalic acid bisanilide or dimethyl succinate/1-(2-hydroxyethyl)-4-hydroxy-2,2,6,6-tetramethylpiperidine polycondensates, etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three

kinds) of them in appropriate ratios.

[0043] As the UV scattering agent, use is made of titanium dioxide, etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

[0044] As the coloring agent, use is made of Cyaningreen G, Eriogreen B400, etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

[0045] As the suspension stabilizer, use is made of polyvinyl alcohols (Gohsenol GH17, etc.), clay minerals (e.g., Kunipia F, VEEGUM R, etc.) or silicon dioxide (Aerosil COK84, etc.), etc., and these can suitably be used by mixing one or two or more kinds (preferably, not less than one but not more than three kinds) of them in appropriate ratios.

[0046] The jumbo preparation, dust, granule, granular wettable powder, wettable powder, etc. may be used after being packed as divided in 20 to 200 g unit portions in bags made of a water-soluble film in order to simplify their application. The said water-soluble film is exemplified by polyvinyl alcohol, carboxymethylcellulose, starch, gelatin, polyvinylpyrrolidone, polyacrylic acid and its salts, Pullulan (tradename: starch-based polysaccharides) or Paogen (tradename: water-soluble thermoplastic polymer), etc.

[0047] In manufacturing the composition preparations according to the present invention, it is possible to control the release of both or either one of the active ingredients, as the case may be, to thereby sustain its or their insecticidal effect(s) over a prolonged period of time.

[0048] The compound [Ia] or its salt and the neonicotinoid compound thiamethoxam as combined are normally contained in the composition of the present invention at the rate of about 0.1 to 80% by weight based on the total weight of the composition. In cases where they are utilized in the form of, for example, emulsifiable concentrate, soluble liquid, wettable powder (e.g., granular wettable powder), aqueous suspension preparation or microemulsion, etc., they are suitably incorporated into such formulations at the rate in the range of normally about 1 to 80% by weight, preferably about 10 to 50% by weight. In cases where they are utilized in the formulations forms, such as oil solution, dust, etc., they are suitably formulated into such formulations at the rate in the range of normally about 0.1 to 50% by weight, preferably about 0.1 to 20% by weight. When they are used in the formulations forms, such as granule, tablet, jumbo preparation, etc., they are incorporated into such formulations at the rate in the range of normally about 0.5 to 50% by weight, preferably about 0.5 to 10% by weight.

[0049] The compound [Ia] or its salt and the neonicotinoid compound thiamethoxam are preferably contained in the composition of the present invention at the rate of 1:0.1 to 1:20 on a weight ratio, more preferably 1:0.2 to 1:10 on a weight ratio.

[0050] The contents of additives other than the above-mentioned active ingredients are usually in the range of about 0.001 to 99.9% by weight, preferably about 1 to 99% by weight, though they vary depending upon the type or content of the active ingredients of agrochemicals, or the formulation of agrochemical preparations, etc. More particularly, it is preferred to add, on the basis of the total weight of the composition, a surfactant at rates in the range of normally about 1 to 30% by weight, preferably about 1 to 15% by weight, a fluidizing auxiliary agent at rates in the range of about 1 to 20% by weight, and a carrier at rates in the range of about 1 to 90% by weight, preferably about 1 to 70% by weight, respectively. Specifically, in the case of manufacture of a liquid preparation, it is preferable to add a surfactant at rates in the range of normally about 1 to 20% by weight, preferably 1 to 10% by weight, and water at rates in the range of about 20 to 90% by weight, respectively. In cases where an emulsifiable concentrate is produced, it is desirable to add a surfactant at rates in the range of normally about 1 to 30% by weight, preferably about 2 to 15% by weight, and an organic solvent. In the case of production of a granular wettable powder, it is desired to add a surfactant at rates in the range of normally about 0.1 to 10% by weight, preferably about 0.5 to 5% by weight, and a binder at rates in the range of about 0.1 to 15% by weight, preferably about 0.5 to 5% by weight, respectively, as well as an extender, such as lactose, ammonium sulfate or clay, etc. In cases where a granule is manufactured, it is desirable to add a surfactant at rates in the range of normally about 0.1 to 10% by weight, preferably about 0.5 to 5% by weight, and a stabilizer at rates in the range of about 0.1 to 10% by weight, preferably about 0.5 to 5% by weight, respectively, as well as an extender, such as clay, etc.

[0051] A wettable powder (e.g., a granular wettable powder) and the like, on the occasion of use, are desirably applied or sprayed after being suitably diluted (e.g., about 100- to 5,000-fold dilution) with water, etc.

[0052] Moreover, the composition of the present invention can suitably be used after being incorporated with the active ingredients of agrochemicals other than the compound [Ia] or its salt and the neonicotinoid compound thiamethoxam, such as other insecticidal active ingredients, acaricidal active ingredients, fungicidal active ingredients, nematicidal active ingredients, herbicidal active ingredients, plant hormone agents, plant growth regulators, synergists (e.g., piperonyl butoxide, sesamex sulfoxide, MGK 264, N-decylimidazole, WARF-antiresistant, TBPT, TPP, IPB, PSCP, CH₃I, t-phenylbutenone, diethyl maleate, DMC, FDMC, ETP, ETN), attractants, repellents or fertilizers, etc.

[0053] Below described are examples of the insecticidal active ingredients, acaricidal active ingredients and bactericidal active ingredients which can be incorporated into the compositions of the present invention:

Insecticidal active ingredients:

5 O-Ethyl O-4-nitrophenyl phosphonothioate (EPN), acephate, isoxathion, isofenfos, isoprocarb, etrimfos, oxy-deprofos, quinalphos, cadusafos, chlorethoxyfos, chlorpyrifos, chlorpyrifos-methyl, chlorofenvinphos, salithion, cyanophos, disulfoton, dimethoate, sulprofos, diazinon, thiometon, tetrachlorvinphos, tebupirimfos, trichlorphon, naled, vamidothion, pyradophos, pyridafenthion, pyrimiphos-methyl, fenitrothion, fenthion, phenthionate, butathiofos, prothiofos, propaphos, profenofos, benclothiaz, phosalone, fosthiazate, marathion, methidathion, metolcarb, monocrotophos, phenobcarb (BPMC), 3,5-xylyl N-methylcarbamate (XMC), alanycarb, ethiofencarb, carbaryl, carbosulfan, carbofuran, xylylcarb, cloethocarb, thiocarbamate, triazamate, pirimicarb, fenoxy carb, fenothiocarb, furathiocarb, propoxur, bendiocarb, benfuracarb, methomyl, acrinathrin, imiprothrin, ethofenprox, cycloprothrin, sigma-cypermethrin, cyhalothrin, cyfluthrin, cypermethrin, silafluofen, tefluthrin, deltamethrin, tralomethrin, fenvalerate, fenpropothrin, flucythrinate, fluvalinate, flufenprox, fluproxyfen, profluthrin, beta-cyfluthrin, benfluthrin, permethrin, cartap, thiocyclam, bensultap, avermectin, emamectin-benzoate, chlorfluazuron, cyromazine, diafenthiuron, dichlorvos, diflubenzuron, spynosyn, spiromesifen, teflubenzuron, tebufenozone, hydroprene, vaniliprole, pymetrozine, pyriproxyfen, fipronil, flufenoxuron, buprofezin, hexaflumuron, milbemycin, lufenuron, chlorphenapyr, pyridalyl, flufendiamide, SI-0009, metofluthrin, noviflumuron, dimefluthrin, cyflumetofen, pyrafluprole and pyriproxyfen.

Acaricidal active ingredients:

25 Clofentezine, dienochlor, tebufenpyrad, pyridaben, hexythiazox, fenazaquin, fenpyroximate, etoxazole, amitraz, bromopropylate, fenbutatin oxide, pyrimidifen, BPPS (propargite), tebufenpyrad and dicofol.

Fungicidal active ingredients:

30 Iprobenphos (IBP), ampropylfos, edifenphos, chlorthiophos, tolclofos-methyl, fosetyl, ipconazole, imazalil, imibenconazole, etaconazole, epoxiconazole, cyproconazole, diniconazole, difenoconazole, tetaconazole, tebuconazole, triadimenol, triadimefon, triticonazole, triforine, bitertanol, viniconazole, fenarimol, fenbuconazole, fluotrimazole, furconazole-cis, flusilazole, flutriafol, bromuconazole, propiconazole, hexaconazole, pefurazole, penconazole, myclobutanil, metconazole, cabendazin, debacarb, prothiocarb, benomyl, maneb, TPN, isoprotiolane, iprodione, iminoctadine-albesil, iminoctadine-triacetate, ethirimol, etridiazole, oxadixyl, oxycarboxin, oxolinic acid, ofurace, kasugamycin, carboxin, captan, clozylacon, chlobenthiazole, cyprodinil, cyprofuram, diethofencarb, dichlofuanid, didomezine, zineb, dimethirimol, dimethomorph, dimefluazole, thiabendazole, thiophanate-methyl, thifluzamide, tecloftalam, triazoxide, triclamide, tricyclazole, tridemorph, triflumizole, validamycin A, hymexazol, pyracarbolid, pyrazophos, pyrifenoxy, pyrimethanil, pyroquilon, ferimzone, fenpiclonil, fenpropidin, fenpropimorph, fthalide, furametpyr, furalaxyd, fluazinam, furcarbanil, fluquinconazole, fludioxonil, flusulfamide, flutolanil, butiobate, prochloraz, procymidone, probenazole, benalaxyl, benodanil, pencycuron, myclozolin, metalaxyl, metsulfovax, methfuroxam, mepanipyrim, mepronil, kresoxim-methyl, azoxystrobin, methominostrobin (SSF-126) carpropamid, acibenzolar-S-methyl, orysastrobin, pyraclostrobin, benthiabarb-carb, t boscalid, metrafenone, fluoxastrobin, proquinazid, flumorph, prothioconazole, penthiopyrad, fluopicolide, amsuldoles, SYP-Z071 and MTF-753.

45 **[0054]** The above-mentioned "other active ingredients of agrochemicals" all are the known active ingredients of agrochemicals. The other active ingredients of agrochemicals may be contained in one or two or more kinds (preferably, not less than one but not more than three kinds) in the compositions.

[0055] The compositions of the present invention exhibit an improved degree of safety to mammals and crops, while 50 they possess enhanced insecticidal action against many species of insect pests (inclusive of arthropods other than the class *Insecta*).

[0056] When not less than two kinds of insecticidal compounds are mixed in aiming at an intensified efficacy or a broadened scope of the targeted insect pests to be controlled, generally, investigation is carried out into a novel application by simply adding the application rates determined for their single applications, whereby there is incurred an increased 55 risk of causing chemical injuries. In the combined treatment according to the present invention, the compounds [Ia] themselves are almost free from their risk of causing chemical injuries, and consequently, the risk of causing chemical injuries to be incurred by such treatment is substantially negligible, in contrast with the treatment through single application of thiamethoxam. Because the compositions can produce the synergistic effect, additionally, the application rates of

either one or both of the compounds [Ia] and thiomethoxam can be reduced from the ones specified for single uses of these compounds, thereby leading to a by far reduced risk of causing chemical injuries.

[0057] The compositions of the present invention are specifically applicable, for example, to the control of the below-described insect pests:

5 **[0058]** Namely, they are especially effective for controlling the insect pests, which are exemplified by: insect pests of the order *Hemiptera*, such as *Eurydema rugosum*, *Scotinophara lurida*, *Riptortus clavatus*, *Stephanitis nashi*, *Laodelphax striatellus*, *Nilaparvata lugens*, *Nephrotettix cincticeps*, *Unaspis yanonensis*, *Aphis glycines*, *Lipaphis erysimi*, *Brevicoryne brassicae*, *Aphis gossypii*, *Myzus persicae*, *Aulacorthum solani*, *Aphis spiraecola*, *Bemisia tabaci*, *Trialeurodes vaporariorum*, *Sogatella furcifera*, *Empoasca onukii*, *Pseudococcus comstocki*, *Planococcus citri*, *Icerya purchasi*, *Plautia stali*, *Eysarcoris parvus* and the like; insect pests of the order *Lepidoptera*, such as *Spodoptera litura*, *Plutella xylostella*, *Pieris rapae*, *crucivora*, *Chilo suppressalis*, *Autographa nigrisigna*, *Helicoverpa assulta*, *Pseudaletia separata*, *Mamestra brassicae*, *Adoxophyes orana fasciata*, *Notarcha derogata*, *Cnaphalocrocis medinalis*, *Phthorimaea operculella*, *Chilo polychrysus*, *Typyra incertulas*, *Spodoptera exigua*, *Agrotis segetum*, *Agrotis ipsilon*, *Heliothis armigera*, *Heliothis virescens*, *Heliothis zea*, *Naranga aenescens*, *Ostrinia nubilalis*, *Ostrinia furnacalis*, *Parnara guttata*, *Adoxophyes* spp., *Caloptilia theivora*, *Phyllonorycter ringoneella*, *Carposina nipponensis*, *Grapholita molesta*, *Cydia pomonella* and the like; insect pests of the order *Coleoptera*, such as *Epilachna vigintioctopunctata*, *Aulacophorafemoralis*, *Phyllotreta striolata*, *Oulema oryzae*, *Echinochenus squameus*, *Lissorhoptrus oryzophilus*, *Anthonomus grandis*, *Callosobruchus chinensis*, *Sphenophorus venatus*, *Popillia japonica*, *Anomala cuprea*, *Diabrotica* spp., *Leptinotarsa decemlineata*, *Agriotes* spp., *Lasioderma serricome*, *Anthrenus verbasci*, *Tribolium castaneum*, *Lyctus brunneus*, *Anoplophora malasiaca*, *Tomicus piniperda* and the like; insect pests of the order *Diptera*, such as *Musca domestica*, *Culex popiens pallens*, *Tabanus trivittatus*, *Delia antique*, *Delia platura*, *Anopheles sinensis*, *Agromyza oryzae*, *Hydrellia griseola*, *Chlorops oryzae*, *Dacus cucurbitae*, *Ceratitis capitata*, *Liriomyza trifolii* and the like; insect pests of the order *Orthoptera*, such as *Locusta migratoria*, *Gryllotalpa Africana*, *Oxya yezoensis*, *Oxya japonica* and the like; insect pests of the order *Thysanoptera*, such as *Thrips tabaci*, *Thrips parmi*, *Frankliniella occidentalis*, *Baliothrips biformis*, *Scirtothrips dorsalis* and the like; insect pests of the order *Hymenoptera*, such as *Athalia rosae*, *Acromyrmex* spp., *Solenopsis* spp. and the like; insect pests of the family *Blattodea*, such as *Blattella germanica*, *Periplaneta fuliginosa*, *Periplaneta japonica*, *Periplaneta Americana* and the like; nematodes, such as *Aphelenchoides besseyi*, *Nothotylenchus acris* and the like; termites, such as *Coptotermes formosanus*, *Reticulitermes speratus*, *Odontotermes formosanus*, *Cryptotermes domesticus* and the like, etc.

25 **[0059]** Also, the compositions of the present invention can find application in the field of treatment of livestock diseases and in the industry of animal husbandry as well as in maintenance of the public hygiene by exterminating arthropods and parasite insects living in and/or on the bodies of the class Vertebrate, such as humans, cattle, sheep, goats, hogs, poultry, dogs, cats and fishes. Examples of the said parasites include *Aedes* spp., *Anopheles* spp., *Culex* spp., *Culicoides* spp., *Musca* spp., *Hypoderma* spp., *Gasterophilus* spp., *Haematobia* spp., *Tabanus* spp., *Simulium* spp., *Triatoma* spp., *Phthiraptera* (e.g., *Damalinia* spp., *Linognathus* spp., *Haematopinus* spp.), fleas (e.g., *Ctenocephalides* spp., *Xenosylla* spp., etc.) or *monomorium pharaonis*, and the like.

30 **[0060]** The compositions of the present invention show an extremely low toxicity and can be used as an excellent agrochemical composition.

35 **[0061]** For example, the compositions of the present invention can be sprayed over paddy fields, fields, fruit tree orchards, non-cultivated waste lands, houses, etc. by use of the per se known methods to thereby exterminate the above-mentioned living insect pests (injurious insects) by allowing such insect pests to contact with or ingest them. In another embodiment of the present invention, for example, the compositions of the present invention can be administered to vertebrates internally (in the body) or externally (on the body surface) to thereby exterminate the arthropods parasitic on such vertebrate or parasites.

40 **[0062]** With reference to the particular application method for the combination preparations comprising the composition of the present invention, use can be made of the same methods as the ordinary agrochemical application methods. Also, the preparations each containing a single different active ingredient can find application through blending on the occasion of use. The application method for such agrochemical preparations can be exemplified by the foliage application, trunk or bark application, ULV application, granule foliar-application, land or soil application, soil drenching, water-surface application, soil incorporation, nursery bed incorporation, seedling-raising box treatment, nursery bed treatment, plant foot treatment, planting furrow treatment, planting row treatment, side row treatment, trunk or bark drenching, trunk or bark spreading, seed coating or dressing, seed immersion or soaking, poison bait, fertilizer incorporation or water blending or mixing for irrigation, etc., but are not understood to be limited to them. The application time for the combination preparations comprising the composition of the present invention or mixtures of the preparations each containing a single different active ingredient may be any arbitrary occasions in advance of planting in the case of treatment of seeds, seed potatoes or bulbs, etc.; and may be the occasions of sowing, seedling-raising or seedling planting for the improved efficacy, although it can be any occasion during the vegetation period after seedling planting, in the case of treatment of the soil; and may be any occasions during the seedling raising period and during the vegetation period on the farm field in the case of foliar application or spraying.

[0063] In cases where sowing is effected on the nursery soil for seedling raising into which the combination preparations comprising the composition of the present invention or mixtures of the formulations each comprising a single different active have been incorporated, when provisional planting is done with use of the said nursery bed soil, or in cases where the soil is treated through solution drenching or granule application during the seedling raising period, all the insect pests appearing during the seedling raising period can also be exterminated.

[0064] As the application method on the occasion of seedling planting, treatment can be conducted through soil incorporation on the whole surface of the field or on furrows in advance of planting, while treatment can also be done by granule application or solution drenching into planting holes. After planting, furthermore, treatment can be effected by immediately applying a granule or drenching a solution at plant foots.

[0065] The crop to be raised by seedling in the farm field is able to be subjected not only to seed treatment, but also soil incorporation treatment on the whole surface of the field or on the furrows in advance of planting.

[0066] The compositions of the present invention, through their mixed use with natural-enemy microbial formulations, their combined utilization with natural-enemy organisms (e.g., natural enemy insects, such as parasitic bees and predacious beetles, predacious acarides, parasitic nematodes, insect-pathogenic microbes, etc.), their combined use with insect pheromones, their combined employment with genetically modified crops, their combined utilization with attractants and repellents, and the like, can contribute to promotion of the IMP (Integrated Pest Management).

[0067] Taking for example the control of diamond-back moths by way of the disturbance in signal communication through use of the pheromones or with utilization of the natural enemies in the cultivation of cabbages, such controlling methods are known to be less effective or entirely ineffective, when the population density of the injurious insect to be controlled becomes high. In the farm field where the population density of diamond-back moths is suppressed to an extremely lower level through soil treatment with the composition of the present invention conducted on the occasion of planting of cabbages, at the point of time when the compound [Ia] or its salt, or thiomethoxam starts to lose its residual effect, the disturbance in signal communication or the utilization of the natural enemies can be employed to thereby secure by far the originally intended effect of such signal communication disturbance or the natural-enemy utilization, thus ensuring that the pest control would keep on working over a prolonged period of time. In controlling insect pests by way of the disturbance in signal communication through use of the pheromones or with utilization of the natural enemies, moreover, the problem arises with proliferation of injurious insects other than the targeted one to be controlled. Under these circumstances, the compositions of the present invention can be applied to thereby suppress the proliferation of injurious insects other than the targeted one, which proliferation has been the problem of great concern in the control of insect pests by way of the disturbance in signal communication or with utilization of natural enemies, and can therefore provide the by far improved integrated pest control.

[0068] The application rate of the compositions of the present invention can be varied over a widened range, depending for example upon the time, location and method of application, etc. but the application is desirably effected at rates of about 0.3 to 3,000 g, preferably about 50 to 1,000 g, of the active ingredient (the total sum of the amounts of the compound [Ia] or its salt and thiomethoxam) per hectare. In cases where the composition of the present invention is formulated into the form of wettable powder, it is desirable to make application after being diluted to a final concentration of the active ingredient (the total sum of the concentrations of the compound [Ia] or its salt and thiomethoxam) of about 0.1 to 1,000 ppm, preferably about 10 to 200 ppm, for foliage application, and about 1 to 10,000 ppm, preferably about 100 to 2,000 ppm, for drenching into the soil.

40 Examples

[0069] Hereunder, the present invention is illustrated in more detail by reference to the following Examples and Test Examples.

[0070] In the Examples described below, the Compounds (I-1), (I-2), (I-3), (I-4), (I-5) and (I-6) are understood to designate 2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloro-pyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[3-bromo-1-(3-chloropyridin-2-yl)pyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide and 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-ylcarbonylamino]-5-chloro-N-isopropyl-3-methylbenzamide, respectively.

Example 1 (comparative example)

[0071] 5 parts of the Compound (I-1), 8 parts of clothianidin, 0.5 part of a non-ionic surfactant (tradename: Noigen EA-177; manufactured by Dai-ichi Kogyo Seiyaku CO., Ltd.), 2 parts of an anionic surfactant (tradename: New Calgen FS-4; manufactured by Takemoto Oils & Fats Co., Ltd.), 2 parts of polyvinyl alcohol (tradename: Gohsenol GH-17; manufactured by Nippon Synthetic Chemical Co., Ltd.), 0.1 part of butylparaben and 82.4 parts of water are mixed and

dispersed sufficiently with a high-speed stirring machine, followed by wet pulverization (1 pass) with a pulverizer, "Dyno-Mill" (constructed by Sinmaru Enterprise Co., 1.0 mm glass beads, 80 % of filling factor, 15 m/sec of peripheral speed), to give a flowable preparation.

5 Example 2 (comparative example)

[0072] 5 parts of the Compound (I-2), 8 parts of clothianidin, 0.5 part of a nonionic surfactant, Noigen EA-177, 1.5 parts of an anionic surfactant, New Calgen FS-4, 2 parts of silicon dioxide (tradename: Aerosil COK84; manufactured by Nippon Aerosil Co., Ltd.), 2 parts of polyvinyl alcohol (tradename: Gohsenol GH-17), 7 parts of ethylene glycol, 0.2 part of a silicone-based anti-foaming agent (tradename: Antifoam E-20; manufactured by Kao Corp.), 0.1 part of butyl-paraben and 73.7 parts of water are mixed and dispersed sufficiently with a high-speed stirring machine, followed by wet pulverization (1 pass) with a pulverizer, "Dyno-Mill" (constructed by Sinmaru Enterprise Co., 1.0 mm glass beads, 80 % of filling factor, 15 m/sec of peripheral speed), to give a flowable preparation.

15 Example 3 (comparative example)

[0073] 1 part of the Compound (I-1), 1 part of clothianidin, 0.5 part of a non-ionic surfactant (tradename: New Pole PE-64; manufactured by Sanyo Chemical Industries Ltd.), 4 parts of alpha-starch and 93.5 parts of clay are mixed uniformly, and the mixture is admixed with 5 to 10 parts of water, kneaded and followed by extrusion through a 0.8mmφ-screen to conduct granulation. The resultant granulated material is dried at 60°C for 1 hour to give a granule preparation.

Example 4 (comparative example)

[0074] A solution of 20 parts of nitenpyram and 80 parts of cyclodextrin (tradename: Toyoderin P; manufactured by JT Foods Co., Ltd.) in 400 parts of water is spray-dried to give a cyclodextrin clathrate A of nitenpyram.

[0075] 1 part of the Compound (I-2), 5 parts of cyclodextrin clathrate A, 2 parts of an anionic surfactant (tradename: New Calgen EP-70P; manufactured by Takemoto Oils & Fats Co., Ltd.), 10 parts of dextrin NDS and 82 parts of clay are mixed uniformly, and the mixture is kneaded with 5 to 10 parts of water, and the same procedure as described in Example 2 is followed to give a granule preparation.

30 Example 5 (comparative example)

[0076] 0.2 part of the Compound (I-1), 0.15 part of clothianidin, 2 parts of an anionic surfactant, New Calgen EP-70P, 0.2 part of a flocculant, IP Solvent, 1 part of white carbon and 96.45 parts of clay are kneaded uniformly, followed by fine pulverization to give a DL dust preparation.

Example 6 (comparative example)

[0077] 0.2 part of the Compound (I-1), 1.25 parts of the cyclodextrin clathrate A prepared in Example 4, 2 parts of an anionic surfactant, New Calgen EP-70P, 0.2 part of a flocculant, IP Solvent, 1.5 parts of white carbon and 94.85 parts of clay are kneaded uniformly, followed by fine pulverization to give a DL dust preparation.

Example 7 (comparative example)

[0078] 0.2 part of the Compound (I-2), 0.15 part of imidacloprid, 1 part of tall oil fatty acid (tradename: Hartall FA-1; manufactured by Harima Chemicals, Inc.), 0.2 part of a flocculant, IP Solvent, 1.5 parts of white carbon and 95.1 parts of clay are kneaded uniformly, followed by fine pulverization to give a DL dust preparation.

50 Example 8 (comparative example)

[0079] 5 parts of the Compound (I-4), 8 parts of clothianidin, 7 parts of an anionic surfactant (tradename: NewCalgen 98147TX; manufactured by Takemoto Oils & Fats Co., Ltd.) and 80 parts of N-(n-dodecyl)pyrrolidone (tradename: AGSOLEX12; manufactured by ISP TECHNOLOGIES INC.) are mixed uniformly for dissolution to give an emulsifiable concentrate.

55 Test Example 1: Insecticidal effect against *Spodoptera litura* by soaking treatment of a bait crop plant:

[0080] A test compound was dissolved by addition of acetone containing 5 % of Tween 20 (tradename) at the rate of

0.1 ml per 1 mg of the test compound, and after the solution is diluted with aqueous 5,000-fold diluted DAIN solution to a predetermined concentration, soybean leaves after complete development of primary leaves were treated through soaking in the insecticide solution for several seconds. After the insecticide solution was dried, four primary leaves were cut off and placed in an ice-cream cup (with a capacity of 180 ml), in which 10 heads of 3-instar larvae of *Spodoptera litura* were released. The ice-cream cup was kept in a breeding room controlled at a constant temperature (25°C), and the number of dead larvae was counted three days later. The death rate of larvae was calculated by the following equation, and the results were shown in Table 1.

10 [Equation 1]

$$\text{Death rate of larvae} = (\text{Number of larvae dead}) / (\text{Number of larvae tested}) \times 100$$

15 Table 1:

Insecticidal effect in soybean plants against <i>Spodoptera litura</i> through soaking treatment of leaves			
Compound	Compound Concentration (ppm)	Death rate, 3 days later, %	
Compound (I-2)	0.03	30	
Clothianidin	0.03	0	
Compound (I-2) + clothianidin (comparative example)	0.03 + 0.03	70*	30**
Notes: *; Activity found by the combination of two kinds of the compounds (actual effect) **; Activity calculated by Colby's equation (expected effect)			

30 [0081] When the effect obtained by combination of two kinds of the active compounds exceeds the expected value E as calculated by the below-described equation of Colby et al., the synergistic effect is deemed to exist.

35 [Equation 2]

$$E = X + Y - X \cdot Y / 100$$

wherein

40 E = the death rate of larvae obtained when the active compounds A and B are used at the dose of m and n, respectively;
X = the death rate of larvae when the active compound A is used at the dose of m;
Y = the death rate of larvae when the active compound B is used at the dose of n.

45 [0082] As is indicated in Table 1, the Compound (I-2), when used as a mixture with clothianidin, was found to produce a greater insecticidal effect than would be expected when individually used alone.

Test Example 2: Insecticidal effect against *Spodoptera litura* through soaking treatment of roots of a bait crop plant

50 [0083] A test compound was dissolved by adding acetone containing 5 % of Tween 20 (tradename) at the rate of 0.1 ml per 1 mg of the test compound, and the insecticide solution prepared by diluting the solution with ion-exchange water to a predetermined concentration was charged into a light-shielded Erlenmeyer flask (100 ml), into which the root portion of a soybean plant in the stage of development of primary leaves was soaked after being washed with running water to remove soil. Five days after root soaking, two primary leaves were cut off and placed in an ice-cream cup (with a capacity of 180 ml), into which 10 heads of 3-instar larvae were released. The ice-cream cup was kept in a breeding room controlled at a constant temperature (25°C), and 5 days later, the number of dead larvae was counted. The death rate of larvae was calculated by the following equation, with the results being tabulated in Tables 2 and 3.

[Equation 3]

Death rate of larvae = (Number of larvae dead)/(Number of larvae tested) x 100

5

Table 2:

Insecticidal effect in soybean plants against <i>Spodoptera litura</i> through soaking treatment of roots			
Compound	Compound Concentration (ppm)	Death rate, 5 days later, %	
Compound (I-1)	0.007	55	
Thiamethoxam	0.007	0	
Compound (I-1) + thiamethoxam	0.007 + 0.007	80*	55**

Notes: * ; Activity found by the combination of two kinds of the compounds (actual effect)
**; Activity calculated by Colby's equation (expected effect)

20

Table 3:

Compound	Compound Concentration (ppm)	Death rate, 5 days later, %	
Compound (I-6)	0.00032	0	0
Clothianidin	0.00032	0	0
Compound (I-6) + clothianidin (comparative example)	0.00032 + 0.00032	65*	0**

30

As is indicated in Table 2, the Compound (I-1), when used as a mixture with thiamethoxam, was found to produce a greater insecticidal effect than would be expected when individually used alone, and the synergistic effect due to the mixing was observed.

35

[0084] As is indicated in Table 3, the Compound (I-6), when used as a mixture with clothianidin, was found to produce a greater insecticidal effect than would be expected when individually used alone, and the synergistic effect due to the mixing was observed. Test Example 3: Insecticidal effect against *Chilo suppressalis* through drenching treatment of the soil with the insecticide solution

40

[0085] A test compound was dissolved by adding acetone containing 5 % of Tween 20 (tradename) at the rate of 0.1 ml per 1 mg of the test compound, and the insecticide solution prepared by diluting the solution with ion-exchange water to a predetermined concentration was drenched onto the surface of the soil for raising rice seedlings (5 to 6 seedlings/plant as planted in a paper pot) in the 2.5- to 3-leaf stage at the rate of 1 ml per plant. Two days later, the stalks about 2 cm higher above the soil surface were cut and placed in a test tube, into which 10 heads of 3-instar larvae of *Chilo suppressalis* were released. The test tube was kept in a breeding chamber controlled at a constant temperature (25°C), and the number of living larvae was counted 4 to 5 days later. The death rate of larvae was calculated by the following equation, and the results were shown in Tables 4 and 5.

45

[Equation 4]

Death rate of larvae = (Number of larvae dead)/(Number of larvae tested) x 100

50

Table 4:

Insecticidal effect in rice plants against <i>Chilo suppressalis</i> through drenching treatment of the soil with the insecticide solution		
Compound	Amount applied mg/plant	Death rate, 4 days later, %
Compound (I-1)	0.01	30

(continued)

5	Insecticidal effect in rice plants against <i>Chilo suppressalis</i> through drenching treatment of the soil with the insecticide solution			
10	Compound	Amount applied mg/plant	Death rate, 4 days later, %	
	Clothianidin	0.001	10	
	Compound (I-1) + clothianidin (comparative example)	0.01 + 0.001	70*	37**
15	Notes: * ; Activity found by the combination of two kinds of the compounds (actual effect) **; Activity calculated by Colby's equation (expected effect)			

Table 5:

15	Compound	Amount applied mg/plant	Death rate, 5 days later, %	
20	Compound (I-5)	0.15	5	
	Dinotefuran	0.44	0	
25	Compound (I-5) + Dinotefuran (comparative example)	0.15 + 0.44	55*	5**
30	Notes: * ; Activity found by the combination of two kinds of the compounds (actual effect) **; Activity calculated by Colby's equation (expected effect)			

[0086] As is indicated in Table 4, the Compound (I-1), when used as a mixture with clothianidin, was found to produce a greater insecticidal effect than would be expected when individually used alone, and the synergistic effect due to the mixing was observed.

[0087] As is indicated in Table 5, the Compound (I-5), when used as a mixture with dinotefuran, was found to produce a greater insecticidal effect than expected when individually used alone, and the synergistic effect due to the mixing was observed. Test Example 4: Insecticidal effect against *Plutella xylostella* through drenching treatment of the soil with the insecticide solution:

[0088] A test compound was dissolved by adding acetone containing 5 % of Tween 20 (tradename) at the rate of 0.1 ml per 1 mg of the test compound, and the solution was diluted with ion-exchange water to a total volume of 3 ml. Each of the insecticide solutions prepared was drenched onto the surface of the plant root soil for a cabbage plant grown in a cell tray (with a soil capacity of 24 ml). Four days later, the above-ground part of the plant was cut and placed in a plastic cup, into which 10 heads of 2-instar larvae of *Plutella xylostella* were released. The cup was kept in a breeding chamber controlled at a constant temperature (25°C), and the number of living larvae was counted 4 days later. The death rate of larvae was calculated by the following equation, and the results were shown in Tables 6 and 7.

[Equation 5]

$$\text{Death rate of larvae} = (\text{Number of larvae dead})/(\text{Number of larvae tested}) \times 100$$

Table 6:

45	Insecticidal effect in cabbage plants against <i>Plutella xylostella</i> through plant-root drenching treatment with the insecticide solution			
50	Compound	Amount applied mg/plant	Death rate, 4 days later, %	
	Compound (I-2)	0.0016	0	
55	Dinotefuran	0.0016	0	

(continued)

5	Insecticidal effect in cabbage plants against <i>Plutella xylostella</i> through plant-root drenching treatment with the insecticide solution			
	Compound	Amount applied mg/plant		Death rate, 4 days later, %
10	Compound (I-2)+ dinotefuran (comparative example)	0.0016 + 0.0016	60*	0**
	Notes: * ; Activity found by the combination of two kinds of the compounds (actual effect) **; Activity calculated by Colby's equation (expected effect)			

Table 7:

15	Compound	Amount applied mg/plant	Death rate, 4 days later, %	
	Compound (I-3)	0.04	50	
20	Dinotefuran	0.04	10	
	Compound (I-3) + dinotefuran (comparative example)	0.04 + 0.04	80*	55**
	Notes: *; Activity found by the combination of two kinds of the compounds (actual effect) **; Activity calculated by Colby's equation (expected effect)			

25 [0089] As is indicated in Table 6, the Compound (I-2), when used as a mixture with dinotefuran, was found to produce a greater insecticidal effect than would be expected when individually used alone, and the synergistic effect due to the mixing was observed.

30 [0090] As is indicated in Table 7, the Compound (I-3), when used as a mixture with dinotefuran, was found to produce a greater insecticidal effect than would be expected when individually used alone, and the synergistic effect due to the mixing was observed. Test Example 5: Insecticidal effect against *Plutella xylostella* through soaking treatment of the bait crop:

35 [0091] A test compound was dissolved by adding acetone containing 5 % of Tween 20 (tradename) at the rate of 0.1 ml per 1 mg of the test compound, and the insecticide solution was prepared by diluting the solution with an aqueous 5,000-fold diluted DAIN solution to a predetermined concentration. A cabbage leaf was cut off at the site of petiole and soaked in the insecticide solution for several seconds. After the solution was dried, the leaf was placed in an ice-cream cup (with a capacity of 180 ml), into which 10 heads of 2-instar larvae of *Plutella xylostella* were released. The cup was kept in a breeding room controlled at a constant temperature (25°C), and the number of dead larvae was counted 4 days later. The death rate of larvae was calculated by the following equation, and the results were shown in Table 8:

40 [Equation 6]

$$\text{Death rate of larvae} = (\text{Number of larvae dead}) / (\text{Number of larvae tested}) \times 100$$

Table 8:

45	Compound	Amount applied mg/plant	Death rate, 4 days later, %	
	Compound (I-4)	0.0064	15	
50	Thiamethoxam	0.0064	10	
	Compound (I-4) + thiamethoxam	0.0064 + 0.0064	50*	23.5**
	Notes: * ; Activity found by the combination of two kinds of the compounds (actual effect) **; Activity calculated by Colby's equation (expected effect)			

55 [0092] As is indicated in Table 8, the Compound (I-4), when used as a mixture with thiamethoxam, was found to produce a greater insecticidal effect than would be expected when individually used alone, and the synergistic effect due to the mixing was observed.

INDUSTRIAL APPLICABILITY

[0093] The compositions of the present invention can be utilized as an insecticide for agricultural and horticultural uses.

5

Claims


10

Claims for the following Contracting State(s): AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, MC, NL, SE

15

1. An insecticide composition which comprises thiamethoxam and one or not less than two kinds of compounds being selected from a compound represented by the formula [Ia]:

20




25

[I a]


30

wherein R1 is a halogen atom or a C1-6 haloalkyl group, R2 is a halogen atom, R3 and R5 each are a C1-6 alkyl group, R4 is a hydrogen or halogen atom, and X is N, or a salt thereof, with the proviso that the following compounds are excluded

35



40



45

50


and

55

5

10

15



2. An insecticide composition as claimed in claim 1, wherein in the compound represented by the general formula [Ia], R1 is a chlorine or bromine atom, or a trifluoromethyl group, R2 is a chlorine atom, R3 is a methyl group, R5 is an isopropyl group, R4 is a hydrogen or chlorine atom, and X is N.

3. An insecticide composition as claimed in claim 1, wherein the compound represented by the formula [Ia] is 2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide.


4. A method for controlling an insect pest, which comprises applying the insecticide composition as claimed in any one of claims 1 to 3 to locations other than the site where the insect pest inflicts injuries directly.

5. A method for controlling an insect pest, **characterized in that** said method comprises mixing two kinds of the compounds, namely a compound represented by the general formula [Ia]:

35

40


45



[I a]

wherein R1 is a halogen atom or a C1-6 haloalkyl group, R2 is a halogen atom, R3 and R5 each are a C1-6 alkyl group, R4 is a hydrogen or halogen atom, and X is N, or a salt thereof and thiamethoxam, followed by drenching onto the soil for raising seedlings in the form of a mixture solution or application on the soil for raising seedlings in the form of a mixture granule, during the period ranging from the sowing time to the seedling-planting time for a crop to be cultivated by the seedling-planting method.


6. A method for controlling an insect pest, **characterized in that** said method comprises growing seedlings with use of the soil for raising seedlings which has contained therein two kinds of the compounds, namely a compound represented by the general formula [Ia]



[I a]

wherein R1 is a halogen atom or a C1-6 haloalkyl group, R2 is a halogen atom, R3 and R5 each are a C1-6 alkyl group, R4 is a hydrogen or halogen atom, and X is N, or a salt thereof and thiamethoxam, during the period ranging from the sowing time to the seedling-planting time for a crop to be cultivated by the seedling-planting method.

20 7. A method for controlling an insect pest, **characterized in that** said method comprises applying two kinds of the compounds, namely a compound represented by the general formula [Ia]



[I a]

or a salt thereof as claimed in any one of claims 1 to 3 and thiamethoxam, to the soil of a farm field through drenching treatment, planting-hole treatment, planting-hole treated soil incorporation, plant-root treatment or plant-root treated soil incorporation during the period ranging from the seedling planting time to the vegetation period for a crop to be cultivated by the seedling-planting method.


40

45

50

8. A method for controlling an insect pest, **characterized in that** said method comprises effecting the immersion treatment, dusting treatment or coating treatment of a seed, seed potato or bulb with two kinds of the compounds, namely a compound represented by the general formula [Ia] or a salt thereof as claimed in any one of claims 1 to 3 and thiamethoxam, in the case of a crop to be grown by directly sowing or seeding a seed, seed potato or bulb on the farm field.

9. A method for controlling an insect pest, **characterized in that** said method comprises treating the soil of a farm field with two kinds of the compounds, namely a compound represented by the general formula [Ia]



[I a]

15


or a salt thereof as claimed in any one of claims 1 to 3 and thiamethoxam, through drenching treatment, plant-root treatment or plant-root treated soil incorporation during the vegetation period for a crop to be cultivated by directly sowing or seeding a seed, seed potato or bulb on the farm field.

20

Claims for the following Contracting State(s): IS, LT

25

1. An insecticide composition which comprises thiamethoxam and one or not less than two kinds of compounds being selected from a compound represented by the formula [Ia]:



[I a]

40

wherein R1 is a halogen atom or a C1-6 haloalkyl group, R2 is a halogen atom, R3 and R5 each are a C1-6 alkyl group, R4 is a hydrogen or halogen atom, and X is N, or a salt thereof.

45


2. An insecticide composition as claimed in claim 1, wherein in the compound represented by the general formula [Ia], R1 is a chlorine or bromine atom, or a trifluoromethyl group, R2 is a chlorine atom, R3 is a methyl group, R5 is an isopropyl group, R4 is a hydrogen or chlorine atom, and X is N.
3. An insecticide composition as claimed in claim 1, wherein the compound represented by the formula [Ia] is 2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-trifluoromethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonyl-amino]-N-isopropyl-3-methylbenzamide, 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-yl-carbonylamino]-N-isopropyl-3-methylbenzamide, or 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-ylcarbonylamino]-5-chloro-N-isopropyl-3-methylbenzamide.
4. A method for controlling an insect pest, which comprises applying the insecticide composition as claimed in any one of claims 1 to 3 to locations other than the site where the insect pest inflicts injuries directly.

5. A method for controlling an insect pest, **characterized in that** said method comprises mixing two kinds of the compounds, namely a compound represented by the general formula [Ia]:

5

10

15



[I a]


20 or a salt thereof as claimed in any one of claims 1 to 3 and thiamethoxam, followed by drenching onto the soil for raising seedlings in the form of a mixture solution or application on the soil for raising seedlings in the form of a mixture granule, during the period ranging from the sowing time to the seedling-planting time for a crop to be cultivated by the seedling-planting method.

25 6. A method for controlling an insect pest, **characterized in that** said method comprises growing seedlings with use of the soil for raising seedlings which has contained therein two kinds of the compounds, namely a compound represented by the general formula [Ia]

30

35

40



[I a]


45 or a salt thereof as claimed in any one of claims 1 to 3 and thiamethoxam, during the period ranging from the sowing time to the seedling-planting time for a crop to be cultivated by the seedling-planting method.


46

7. A method for controlling an insect pest, **characterized in that** said method comprises applying two kinds of the compounds, namely a compound represented by the general formula [Ia]

50


55






15 [I a]

in welcher R1 ein Halogenatom oder eine C1-6 Haloalkylgruppe ist, R2 ein Halogenatom, R3 und R5 jeweils eine C1-6 Alkylgruppe, R4 ein Wasserstoff- oder Halogenatom, und X N bedeutet, oder ein Salz davon, unter der Voraussetzung, dass die folgenden Verbindungen ausgeschlossen sind:



und




2. Insektizidzusammensetzung nach Anspruch 1, wobei in der durch die allgemeine Formel [Ia] dargestellten Verbin-

dung R1 ein Chlor- oder Bromatom ist, oder eine Trifluormethylgruppe, R2 ein Chloratom, R3 eine Methylgruppe, R5 eine Isopropylgruppe, R4 ein Wasserstoff- oder ein Chloratom ist, und X N ist.

5 3. Insektizidzusammensetzung nach Anspruch 1, wobei die durch die allgemeine Formel [Ia] dargestellte Verbindung 2-[1-(3-Chlorpyridin-2-yl)-3-trifluormethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamid, 2-[1-(3-Chlorpyridin-2-yl)-3-chlor-pyrazol-5-yl-carbonyl-amino]-N-isopropyl-3-methylbenzamid, oder 2-[3-Brom-1-(3-chlorpyridin-2-yl)-pyrazol-5-yl-carbonylamino]-N-isopropyl-3-methylbenzamid ist.

10 4. Verfahren zur Bekämpfung eines Schädlings, welches die Anwendung der Insektizidzusammensetzung nach einem der Ansprüche 1 bis 3 an Orten umfasst, die unterschiedlich sind zu derjenigen Stelle, an welcher der Schädling direkt Schädigungen verursacht.


15 5. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Mischung zweier Arten an Verbindungen umfasst, nämlich einer Verbindung, die durch die allgemeine Formel [Ia] dargestellt ist:



(I a)

30 in welcher R1 ein Halogenatom oder eine C1-6 Haloalkylgruppe ist, R2 ein Halogenatom, R3 und R5 jeweils eine C1-6 Alkylgruppe, R4 ein Wasserstoff- oder Halogenatom, und X N bedeutet, oder ein Salz davon, sowie Thiometoxam, gefolgt von dem Tränken eines Bodens für die Aufzucht von Sämlingen in Form einer gemischten Lösung, oder gefolgt von der Anwendung auf einem Boden für die Aufzucht von Sämlingen in Form einer Granulatmischung, und zwar während des Zeitraums, der von der Zeit der Aussaat bis zur Zeit der Sämling-Pflanzung reicht, für Anbaupflanzen, die durch die Sämling-Pflanzungs-Methode kultiviert werden.

35 6. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Züchtung von Sämlingen unter Verwendung eines Bodens zur Aufzucht der Sämlinge umfasst, welcher zwei Arten an Verbindungen beinhaltet, nämlich eine durch die folgende allgemeine Formel [Ia] dargestellte Verbindung:




(I a)

55 in welcher R1 ein Halogenatom oder eine C1-6 Haloalkylgruppe ist, R2 ein Halogenatom, R3 und R5 jeweils eine C1-6 Alkylgruppe, R4 ein Wasserstoff- oder Halogenatom, und X N bedeutet, oder ein Salz davon, sowie Thiometoxam,

thoxam, und zwar während des Zeitraums, der von der Zeit der Aussaat bis zur Zeit der Sämling-Pflanzung reicht, für Anbaupflanzen, die durch die Sämling-Pflanzungs-Methode kultiviert werden.

5 7. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Anwendung zweier Arten an Verbindungen umfasst, nämlich einer Verbindung, die durch die allgemeine Formel $[la]$ dargestellt ist:




[l a]

oder eines Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, und Thiomethoxam, auf einen Boden eines Farmfelds, und zwar mittels Tränk-Behandlung, Pflanzloch-Behandlung, Pflanzloch-Einbau behandelten Bodens, Pflanzwurzel-Behandlung, oder Pflanzwurzel-Einbau behandelten Bodens, und zwar während des Zeitraums, der von der Zeit der Sämling-Pflanzung bis zur Vegetationszeit reicht, für Anbaupflanzen, die durch die Sämling-Pflanzungs-Methode kultiviert werden.

30 8. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Ausführung der Immersions-Behandlung, Bestäubungs-Behandlung oder Beschichtungs-Behandlung einer Saat, Saat-Knolle oder Saat-Zwiebel mit zwei Arten an Verbindungen umfasst, nämlich einer durch die allgemeine Formel [la] dargestellten Verbindung oder einem Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, und Thiomethoxam, für den Fall dass eine Anbaupflanze direkt durch Besäumung oder Aussäen einer Saat, einer Saat-Knolle oder einer Saat-Zwiebel auf dem Farmfeld gezüchtet wird.

35 9. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Behandlung des Bodens eines Farmfelds mit zwei Arten an Verbindungen umfasst, nämlich einer Verbindung, die durch die allgemeine Formel [la] dargestellt ist:




[l a]

oder eines Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, und Thiomethoxam, und zwar mittels Tränk-Behandlung, Pflanzwurzel-Behandlung, oder Pflanzwurzel-Einbau behandelten Bodens, und zwar während der Vegetationszeit für Anbaupflanzen, die direkt durch Besäumung oder Aussäen einer Saat, einer Saat-Knolle oder einer Saat-Zwiebel auf dem Farmfeld gezüchtet werden.

Patentansprüche für folgende(n) Vertragsstaat(en): IS, LT

1. Insektizidzusammensetzung, aufweisend Thiamefoxam und eine oder nicht weniger als zwei Arten an Verbindungen, die ausgewählt sind aus einer Verbindung mit der folgenden Formel [Ia]:

5



[I a]

20

in welcher R1 ein Halogenatom oder eine C1-6 Haloalkylgruppe ist, R2 ein Halogenatom, R3 und R5 jeweils eine C1-6 Alkylgruppe, R4 ein Wasserstoff- oder Halogenatom, und X N bedeutet, oder ein Salz davon.

25

2. Insektizidzusammensetzung nach Anspruch 1, wobei in der durch die allgemeine Formel [Ia] dargestellten Verbindung R1 ein Chlor- oder Bromatom ist, oder eine Trifluormethylgruppe, R2 ein Chloratom, R3 eine Methylgruppe, R5 eine Isopropylgruppe, R4 ein Wasserstoff- oder ein Chloratom ist, und X N ist.

30

3. Insektizidzusammensetzung nach Anspruch 1, wobei die durch die allgemeine Formel [Ia] dargestellte Verbindung 2-[1-(3-Chlorpyridin-2-yl)-3-trifluormethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamid, 5-Chlor-2-[1-(3-chloropyridin-2-yl)-3-trifluormethylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-methylbenzamid, 2-[1-(3-Chlorpyridin-2-yl)-3-chlor-pyrazol-5-yl-carbonyl-amino]-N-isopropyl-3-methylbenzamid, 5-Chlor-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-yl-carbonyl-amino]-N-isopropyl-3-methylbenzamid, 2-[3-Brom-1-(3-chlorpyridin-2-yl)-pyrazol-5-yl-carbonylamino]-N-isopropyl-3-methylbenzamid, oder 2-[3-Brom-1-(3-chloropyridin-2-yl)-pyrazol-5-yl-carbonyl-amino]-5-chloro-N-isopropyl-3-methylbenzamid ist.


35

4. Verfahren zur Bekämpfung eines Schädlings, welches die Anwendung der Insektizidzusammensetzung nach einem der Ansprüche 1 bis 3 an Orten umfasst, die unterschiedlich sind zu derjenigen Stelle, an welcher der Schädling direkt Schädigungen verursacht.

40

5. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Mischung zweier Arten an Verbindungen umfasst, nämlich einer Verbindung, die durch die allgemeine Formel [Ia] dargestellt ist:

45



[I a]


oder ein Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, sowie Thiamefoxam, gefolgt von dem

Tränken eines Bodens für die Aufzucht von Sämlingen in Form einer gemischten Lösung, oder gefolgt von der Anwendung auf einem Boden für die Aufzucht von Sämlingen in Form einer Granulatmischung, und zwar während des Zeitraums, der von der Zeit der Aussaat bis zur Zeit der Sämling-Pflanzung reicht, für Anbaupflanzen, die durch die Sämling-Pflanzungs-Methode kultiviert werden.

5

6. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Züchtung von Sämlingen unter Verwendung eines Bodens zur Aufzucht der Sämlinge umfasst, welcher zwei Arten an Verbindungen beinhaltet, nämlich eine durch die folgende allgemeine Formel [Ia] dargestellte Verbindung:

10



[I a]

15

20

oder ein Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, sowie Thiomethoxam, und zwar während des Zeitraums, der von der Zeit der Aussaat bis zur Zeit der Sämling-Pflanzung reicht, für Anbaupflanzen, die durch die Sämling-Pflanzungs-Methode kultiviert werden.


25

30

7. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Anwendung zweier Arten an Verbindungen umfasst, nämlich einer Verbindung, die durch die allgemeine Formel [Ia] dargestellt ist:

35

40



[I a]

45

50

oder eines Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, und Thiomethoxam, auf einen Boden eines Farmfelds, und zwar mittels Tränk-Behandlung, Pflanzloch-Behandlung, Pflanzloch-Einbau behandelten Bodens, Pflanzwurzel-Behandlung, oder Pflanzwurzel-Einbau behandelten Bodens, und zwar während des Zeitraums, der von der Zeit der Sämling-Pflanzung bis zur Vegetationszeit reicht, für Anbaupflanzen, die durch die Sämling-Pflanzungs-Methode kultiviert werden.

55


8. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Ausführung der Immersions-Behandlung, Bestäubungs-Behandlung oder Beschichtungs-Behandlung einer Saat, Saat-Knolle oder Saat-Zwiebel mit zwei Arten an Verbindungen umfasst, nämlich einer durch die allgemeine Formel [Ia] dargestellten Verbindung oder einem Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, und Thiomethoxam, für den Fall dass eine Anbaupflanze direkt durch Besäumung oder Aussäen einer Saat, einer Saat-Knolle oder einer Saat-Zwiebel auf dem Farmfeld gezüchtet wird.

9. Verfahren zur Bekämpfung eines Schädlings, **dadurch gekennzeichnet, dass** das Verfahren die Behandlung des Bodens eines Farmfelds mit zwei Arten an Verbindungen umfasst, nämlich einer Verbindung, die durch die allgemeine Formel [Ia] dargestellt ist:

5

10

15



[I a]

20 oder eines Salz davon, wie in einem der Ansprüche 1 bis 3 beansprucht, und Thiomethoxam, und zwar mittels Tränk-Behandlung, Pflanzwurzel-Behandlung, oder Pflanzwurzel-Einbau behandelten Bodens, und zwar während der Vegetationszeit für Anbaupflanzen, die direkt durch Besäumung oder Aussäen einer Saat, einer Saat-Knolle oder einer Saat-Zwiebel auf dem Farmfeld gezüchtet werden.

25

Revendications


30 **Revendications pour l'(les) Etat(s) contractant(s) suivant(s): AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, MC, NL, SE**

1. Composition insecticide qui comprend du thiaméthoxam et un ou pas moins de deux types de composés étant choisis parmi un composé représenté par la formule [Ia] :

35

40

45




[I a]


55 dans laquelle R1 est un atome d'halogène ou un groupe halogénoalkyle en C1-6, R2 est un atome d'halogène, R3 et R5 sont chacun un groupe alkyle en C1-6, R4 est un hydrogène ou un atome d'halogène, et X est N, ou un sel de celui-ci, sous réserve que les composés suivants sont exclus:

55

5




10



15

20 et

25



30

35

40 2. Composition insecticide selon la revendication 1, dans laquelle, dans le composé représenté par la formule générale [la], R1 est un atome de chlore ou de bromé, ou un groupe trifluorométhyle, R2 est un atome de chlore, R3 est un groupe méthyle, R5 est un groupe isopropyle, R4 est un atome d'hydrogène ou de chlore, et X est N.

45 3. Composition insecticide selon la revendication 1, dans laquelle le composé représenté par la formule [la] est 2-[1-(3-chloropyridin-2-yl)-3-trifluorométhylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-méthylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-méthylbenzamide, ou 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-ylcarbonylamino]-N-isopropyl-3-méthylbenzamide.


50 4. Procédé de lutte contre un insecte nuisible, qui comprend l'application de la composition insecticide selon l'une quelconque des revendications 1 à 3 à des emplacements autres que le site auquel l'insecte nuisible inflige directement des lésions.

55 5. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend le mélange de deux types des composés, à savoir un composé représenté par la formule générale [la] :

55

5

10



[I a]


15 dans laquelle R1 est un atome d'halogène ou un groupe halogénoalkyle en C1-6, R2 est un atome d'halogène, R3 et R5 sont chacun un groupe alkyle en C1-6, R4 est un hydrogène ou un atome d'halogène, et X est N, ou un sel de celui-ci, et le thiaméthoxam, suivi par une pulvérisation sur la terre pour lever les semis sous la forme d'une solution de mélange ou application sur la terre pour lever les semis sous la forme d'un granule de mélange, pendant la période dans l'intervalle du temps de semis au temps de repiquage pour une culture destinée à être cultivée par
20 le procédé de repiquage.

25

6. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend la croissance de semis avec utilisation de la terre pour lever les semis qui contient dans celle-ci deux types des composés, à savoir un composé représenté par la formule générale [Ia]

30

35



[I a]

40

dans laquelle R1 est un atome d'halogène ou un groupe halogénoalkyle en C1-6, R2 est un atome d'halogène, R3 et R5 sont chacun un groupe alkyle en C1-6, R4 est un hydrogène ou un atome d'halogène, et X est N, ou un sel de celui-ci, et le thiaméthoxam, pendant la période dans l'intervalle du temps de semis au temps de repiquage pour une culture destinée à être cultivée par le procédé de repiquage.

45


7. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend l'application de deux types des composés, à savoir un composé représenté par la formule générale [Ia]

50

55

5

10



[I a]

15 ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, sur la terre d'un champ agricole par traitement par pulvérisation, traitement de trou de plantation, incorporation de terre traitée dans un trou de plantation, traitement des racines de plantes ou incorporation de terre traitée avec les racines des plantes pendant la période dans l'intervalle du temps de repiquage à la période de végétation pour une culture destinée à être cultivée par le procédé de repiquage.

20


8. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend la conduite du traitement d'immersion, traitement de poudrage ou traitement d'enrobage d'une graine, une pomme de terre de semence ou un bulbe avec deux types des composés, à savoir un composé représenté par la formule générale [Ia] ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, dans le cas d'une culture destinée à être cultivée par semis ou ensemencement direct d'une graine, une pomme de terre de semence ou un bulbe sur le champ agricole.

9. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend le traitement de la terre d'un champ agricole avec deux types des composés, à savoir un composé représenté par la formule générale [Ia]

30

35

40



[I a]

45 ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, par traitement de pulvérisation, traitement des racines de plantes ou incorporation de terre traitée avec les racines de plantes pendant la période de végétation pour une culture destinée à être cultivée par semis ou ensemencement direct d'une graine, une pomme de terre de semence ou un bulbe sur le champ agricole.

50


Revendications pour l'(les) Etat(s) contractant(s) suivant(s): IS, LT

1. Composition insecticide qui comprend du thiaméthoxam et un ou pas moins de deux types de composés étant choisis parmi un composé représenté par la formule [Ia] :

55

5

10



(Ia)

15 dans laquelle R1 est un atome d'halogène ou un groupe halogénoalkyle en C1-6, R2 est un atome d'halogène, R3 et R5 sont chacun un groupe alkyle en C1-6, R4 est un hydrogène ou un atome d'halogène, et X est N, ou un sel de celui-ci.

20 2. Composition insecticide selon la revendication 1, dans laquelle, dans le composé représenté par la formule générale [Ia], R1 est un atome de chlore ou de brome, ou un groupe trifluorométhyle, R2 est un atome de chlore, R3 est un groupe méthyle, R5 est un groupe isopropyle, R4 est un atome d'hydrogène ou de chlore, et X est N.

25 3. Composition insecticide selon la revendication 1, dans laquelle le composé représenté par la formule [Ia] est 2-[1-(3-chloropyridin-2-yl)-3-trifluorométhylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-méthylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-trifluorométhylpyrazol-5-ylcarbonylamino]-N-isopropyl-3-méthylbenzamide, 2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonylamino]-N-isopropyl-3-méthylbenzamide, 5-chloro-2-[1-(3-chloropyridin-2-yl)-3-chloropyrazol-5-ylcarbonyl-amino]-N-isopropyl-3-méthylbenzamide, 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-yl-carbonylamino]-N-isopropyl-3-méthylbenzamide, ou 2-[3-bromo-1-(3-chloropyridin-2-yl)-pyrazol-5-ylcarbonylamino]-5-chloro-N-isopropyl-3-méthylbenzamide.


30

4. Procédé de lutte contre un insecte nuisible, qui comprend l'application de la composition insecticide selon l'une quelconque des revendications 1 à 3 à des emplacements autres que le site auquel l'insecte nuisible inflige directement des lésions.

35 5. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend le mélange de deux types des composés, à savoir un composé représenté par la formule générale [Ia] :

40

45



(Ia)

ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, suivi par une pulvérisation sur la terre pour lever les semis sous la forme d'une solution de mélange ou application sur la terre pour lever les semis sous la forme d'un granule de mélange, pendant la période dans l'intervalle du temps de semis au temps de repiquage pour une culture destinée à être cultivée par le procédé de repiquage.


6. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend la croissance de semis

avec utilisation de la terre pour lever les semis qui contient dans celle-ci deux types des composés, à savoir un composé représenté par la formule générale [Ia]

5

10

15



[I a]


ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, pendant la période dans l'intervalle du temps de semis au temps de repiquage pour une culture destinée à être cultivée par le procédé de repiquage.

7. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend l'application de deux types des composés, à savoir un composé représenté par la formule générale [Ia]

25

30

35




[I a]

ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, sur la terre d'un champ agricole par traitement par pulvérisation, traitement de trou de plantation, incorporation de terre traitée dans un trou de plantation, traitement des racines de plantes ou incorporation de terre traitée avec les racines des plantes pendant la période dans l'intervalle du temps de repiquage à la période de végétation pour une culture destinée à être cultivée par le procédé de repiquage.

45 8. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend la conduite du traitement d'immersion, traitement de poudrage ou traitement d'enrobage d'une graine, une pomme de terre de semence ou un bulbe avec deux types des composés, à savoir un composé représenté par la formule générale [Ia] ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, dans le cas d'une culture destinée à être cultivée par semis ou ensemencement direct d'une graine, une pomme de terre de semence ou un bulbe sur le champ agricole.

55 9. Procédé de lutte contre un insecte nuisible, **caractérisé en ce que** ledit procédé comprend le traitement de la terre d'un champ agricole avec deux types des composés, à savoir un composé représenté par la formule générale [Ia]

55



(Ia)

15 ou un sel de celui-ci selon l'une quelconque des revendications 1 à 3 et le thiaméthoxam, par traitement de pulvérisation, traitement des racines de plantes ou incorporation de terre traitée avec les racines de plantes pendant la période de végétation pour une culture destinée à être cultivée par semis ou ensemencement direct d'une graine, une pomme de terre de semence ou un bulbe sur le champ agricole.

20

25

30

35

40

45

50

55


REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.




Patent documents cited in the description

- WO 01070671 A [0004] [0023]
- WO 03015519 A [0004] [0023]
- WO 03016284 A [0004] [0023]
- JP HEI3157308 A [0004] [0024]
- JP HEI2000171 A [0004] [0024]
- JP SHO61178981 A [0004] [0024]
- JP HEI6183918 A [0004] [0024]
- JP HEI4154741 A [0004] [0024]
- JP HEI7179448 A [0004] [0024]
- JP SHO62207266 A [0004] [0024]
- WO 2005048711 A1 [0005]
- WO 03027099 A1 [0005]
- WO 03024222 A1 [0005]

I. Rovarirtó készítmény, amely tartalmaz tiannetoxámot és egy- vagy nem kevesebb mint kétféle vegyületet [la] képletek közül kiválasztva:



ahol R1 jelentése halogénatom vagy C1-6 halogénaikil-csoport, R2 jelentése halogénatom, R3 és R5 mindenügyikének jelentése C1-6 alkillescsoport, R4 jelentése hidrogén- vagy halogénatom, és X jelentése N, vagy ezek szája, azzal a megkötéssel, hogy a következő vegyületek ki vannak zárva:




2. Az 1. igénypontról szerinti rovarirtó készítmény, ahol az [la] általános képletű vegyületben R1 jelentése klór- vagy brómatom, vagy trifluormetilescsoport, R2 jelentése klóratom, R3 jelentése metilescsoport, R5 jelentése izopropilescsoport, R4 jelentése hidrogén- vagy klóratom és X jelentése N.

3. Az 1. igénypontról szerinti rovarirtó készítmény, ahol az [la] képletű vegyület 2-[1-(3-klópiridin-2-il)-3-trifluormetilpirazol-5-ilkarbonilamino]-N-izopropil-3-metilbenzamid, 2-[1-(3-klópiridin-2-il)-3-klópirazol-5-ilkarbonilamino]-N-izopropil-3-metil-benzamid, 2-[3-bróm-1-(3-klópiridin-2-il)-pirazol-5-il-karbonilamino]-N-izopropil-3-metil-benzamid.

4. Eljárás rovarkártevő írtására, amelynek során az 1-3. igénypontról bármelyik szerinti rovarirtó készítményt attól a helytől, ahol a rovarkártevő közvetlenül kifejtő károsító hatását, eltérő helyeken alkalmazzuk.


5. Eljárás rovarkártevő írtására, azzal jellemzve, hogy az eljárás során összekeverünk kétfélé vegyületet, nevezetesen egy [la] általános képletű vegyületet



(Ia)

ahol R1 jelentése halogénatom vagy C1-6 halogénaalkil-csoport, R2 jelentése halogénatom, R3 és R5 mindenek jelentése C1-6 alkikcsoport, R4 jelentése hidrogén- vagy halogénatom és X jelentése N, vagy ennek sóját és tiacetoxámot, majd oldatkeverék formájában ezzel öntözzük a palánták nevelésére alkalmas termőtalajt vagy ezt granulátumkeverék formájában alkalmazzuk a palánták nevelésére alkalmas termőtalajon, a palánta ültetési eljárásai termesztendő növény vetési idejétől a palántaültetés idejéig terjedő időtartam alatt.


6. Eljárás rovarkártevő írtására, azzal jellemzve, hogy az említett eljárás során a palántákat olyan palántanevelő termőtalaj alkalmazásával neveljük, amelyhez kétfélé vegyületet, nevezetesen egy [la] általános képletű vegyületet,



(Ia)

ahol R1 jelentése halogénatom vagy C1-6 halogénaalkil-csoport, R2 jelentése halogénatom, R3 és R5 mindenek jelentése C1-6 alkikcsoport, R4 jelentése hidrogén- vagy halogénatom, és X jelentése N, vagy ennek sóját és tiacetoxámot dolgoztunk be, a palánta ültetési eljárásai termesztett növény vetési idejétől a palántaültetés idejéig terjedő időtartam alatt.

7. Eljárás rovarkártevő írtására, azzal jellemzve, hogy az említett eljárás során kétfélé vegyületet, nevezetesen egy, az 1-3. igénypontok bármelyike szerinti [1a] általános képletű vegyületet




1888.]

vagy ennek sóját és tiame toxámot alkalmazunk egy gazdaság földjének termőtalaján öntözéses kezeléssel, az ültetőlyuk kezelésével, az ültetőlyuk talajába történő bedolgozással, a növény gyökérének kezelésével vagy a növény gyökérének talajába történő bedolgozással, a palánta ültetési eljárással termeszített növény palántaültetési idejétől növekedési idejéig terjedő időtartam alatt.

8. Eljárás rovarkártevő írtására, azzal jellemezve, hogy az említett eljárás során egy vetőmag, vetőburgonya vagy gumó merítéses kezelését, porzó kezelését vagy bevonásos kezelését kétféle vegyülettel, nevezetesen egy, az 1-3. igénypontok bármelyike szerinti [ja] általános képletű vegyülettel vagy sójával és tiame toxámmal végezzük a gazdaság földjén a vetőmag, vetőburgonya vagy gumó közvetlen vetésével vagy ültetéssel termesztiendő növény esetében.

9. Eljárás rovarkártevő írtására, azzal jellemzve, hogy az említett eljárás során egy gazdaság földjének talaját kétfélé vegyűllettet, nevezetesen egy, az 1-3. igénypontok bármelyike szerinti [is] általános képletű vegyűllettel vagy



卷之三

vagy ennek segítsével és tiacetoxámmal kezeljük, öntözéses kezeléssel, a növény gyökérének kezelésével vagy a növény gyökérének talajába történő bedolgozással a gazdaság földjén egy vetőmag, vetőburgonya vagy gumó közvetlen vetésével vagy ültetésével termesztett növény növekedésének időtartama alatt.