

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 199892707 B2**
(10) Patent No. **755745**

(54) Title
Microspore-specific promoter and method for obtaining hybrid plants

(51)⁶ International Patent Classification(s)
C12N 015/82 A01H 005/10
A01H 001/02 C12N 015/29
A01H 005/00

(21) Application No: **199892707** (22) Application Date: **1998.09.23**

(87) WIPO No: **WO99/15678**

(30) Priority Data

(31) Number	(32) Date	(33) Country
97/11812	1997.09.23	FR

(43) Publication Date : **1999.04.12**
(43) Publication Journal Date : **1999.06.10**
(44) Accepted Journal Date : **2002.12.19**

(71) Applicant(s)
Institut National De La Recherche Agronomique

(72) Inventor(s)
Jan Drouaud; Agnes Fourgoux; Georges Pelletier; Philippe Guerche

(74) Agent/Attorney
FREEHILLS CARTER SMITH BEADLE, Level 43, 101 Collins Street, MELBOURNE VIC 3000

(56) Related Art
WO 96/16182
WO 90/08828
EP 790311

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION

International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International patent classification⁶: C12N 15/82, 15/29, A01H 1/02, 5/00, 5/10	A1	(11) International publication number: WO 99/15678 (43) International publication date: 1st April 1999 (01.04.99)
(21) International application number: PCT/FR98/02042 (22) International filing date: 23 September 1998 (23.09.98) (30) Data relating to the priority: 97/11,812 23 September 1997 (23.09.97) DE (71) Applicant (for all designated States except US): INSTITUT NATIONAL DE LA RECHERCHE AGRONOMIQUE [FR/FR]; 147, rue de l'Université, F-75007 Paris (FR). (72) Inventors; and (75) Inventors/Applicants (US only): DROUAUD, Jan [FR/FR]; 7, rue du Général Leclerc, F-78000 Versailles (FR). FOURGOUX, Agnès [FR/FR]; Chemin des Quatre Arpents, F-78330 Fontenay Le Fleury (FR). PELLETIER, Georges [FR/FR]; 28, avenue ed l'Espérance, F-91440 Bures sur Yvette (FR). GUERCHE, Philippe [FR/FR]; 7, rue Marceau, F-92170 Vanves (FR). (74) Representatives: MARTIN, Jean-Jacques etc.; Cabinet Regimbeau, 26, avenue Kléber, F-75116 Paris (FR).		(81) Designated states: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO Patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With the International Search Report. Before expiry of the period provided for amending the claims, will be republished if such amendments are received.
As printed (54) Title: MICROSPORE-SPECIFIC PROMOTER AND METHOD FOR OBTAINING HYBRID PLANTS (54) Titre: PROMOTEUR SPECIFIQUE DES MICROSPORES ET PROCEDE D'OBTENTION DE PLANTES HYBRIDES (57) Abstract The invention concerns a microspore-specific promoter and its use for obtaining plants with gametophytic male sterility with inducible fertility. It also concerns a method for obtaining hybrid plants. (57) Abrégé L'invention concerne un promoteur spécifique des microspores ainsi que sa mise en oeuvre pour obtenir des plantes présentant une stérilité mâle gamétophytique à fertilité inducible. Elle concerne également un procédé d'obtention de plantes hybrides.		

**MICROSPORE-SPECIFIC PROMOTER AND METHOD FOR PRODUCING
HYBRID PLANTS**

5 The present invention concerns in particular a microspore-specific promoter and a method for producing hybrid plants.

The microspore corresponds to a precise stage in the development of the male gamete in higher plants. Male gametogenesis takes place in a specialized organ,
10 the anther, and comprises *sensu stricto* the differentiation of diploid cells into haploid pollen grains. Each diploid cell, called a sporogenic cell, undergoes meiosis to produce four haploid microspores which subsequently differentiate to give mature pollen
15 grains.

Knowing the molecular factors which control the development of the microspore, and how to manipulate them, is a considerable asset not only from a fundamental research point of view but also from a
20 plant-improvement point of view. This is because this knowledge enables the production of pollen grains, and consequently the reproduction of the plant, to be controlled.

Such control proceeds via the production of
25 plants with one of their gametes totally sterile so as to prevent self-fertilization.

So far, male sterility of plants, which is less complex than female sterility, has been widely studied but necessitates the use of genetic systems which are
30 relatively laborious to implement for commercial production of hybrid seeds. One type of male sterility which is highly used is cytoplasmic male sterility which consists in producing:

35 - a female line whose sterile-male characteristic is transmitted through the cytoplasm; such a cytoplasm is called a "male-sterility-inducing cytoplasm"; these "inducing cytoplasm" are, for a given species, in general discovered in the wild, or sometimes



observed in plants which result from interspecific crosses (cross-fertilization, protoplast fusion, etc.),

- a "sterility-maintaining" line whose cytoplasm is normal, and

- a fertility-restoring line if the seeds and/or the fruit of the hybrid plant are harvested.

10 In the female line (carrier of the sterility-inducing cytoplasm) all the pollen grains are killed. To multiply and improve this line it is therefore necessary to have a line which carries neither the inducing cytoplasm (which thus produces pollen grains) nor the restoration gene. This line is termed "sterility-maintaining" because crossing with the female line gives an entirely female lineage.

15 Restoration of the fertility is carried out in the hybrid by crossing the female parent (carrying the sterile male cytoplasm) with the parent comprising a nuclear restoration gene (the restoring line), this cross enabling the production of fertile hybrid plants which will produce seeds by self-fertilization.

20 In the case of sporophytic nuclear sterility, systems have been described, for example, which make it possible to kill the mother cells of the microspores by means of an RNase, and, consequently, to obtain plants lacking in the male gametes. The fertility is restored when the line which no longer produces male gametes is crossed with another line carrying an inhibitor of the RNase, the seeds resulting from this cross comprising both the cytotoxic gene and its inhibitor.

25 As for the present invention, it proposes to produce plants with gametophytic male sterility, which are incapable of producing pollen grains. It consists in using a promoter region which controls the expression, specifically in the microspores, of a gene encoding a cytotoxic molecule, while also having a



means permitting the controlled inhibition of this toxicity, in order to obtain a line of homozygous progenitor plants which are totally sterile as regards their male gametes, and then to obtain fertile hybrid plants (which produce one viable pollen grain in two), thus capable of producing seeds, without
5 having to resort to using a fertility restoration gene.

So far, a single gene which is expressed specifically in the microspore has been described, in tobacco (Oldenhof et al., 1996). This gene does not have any homology with the Brassicaceae as results from a Southern Blot experiment on the genomic DNA of *Brassica oleracea* (data not shown).

10 A subject of the present invention is therefore a nucleotide sequence for which it has been demonstrated that the corresponding gene is expressed specifically in the microspore; this nucleotide sequence corresponds to SEQ ID No. 3.

Consequently, a subject of the present invention is an isolated
15 polynucleotide molecule having the nucleotide sequence corresponding:

- a) the sequence according to SEQ ID. No. 3, or
- b) a sequence which has at least 80% homology with a).

In the context of the present invention, the most valuable part of this nucleotide sequence is the promoter region which is defined as being the
20 sequence preceding (on the 5' side) the translation start codon (ATG). However, at the current stage of knowledge about the nucleotide sequence according to SEQ ID No. 3, three ATGs have been shown: one at position 1965, another at position 2085 and a third at position 2112. It would appear that the functional ATG is the one situated at position 2085. This is not
25 confirmed however; it is the reason why the largest envisagable



promoter region concerning SEQ ID No. 3 stretches from nucleotide 1 to nucleotide 2111, and preferably from nucleotide 1 to nucleotide 2084.

5 This promoter region thus precedes, in the natural state, a coding (orf) sequence which is expressed specifically in the microspores, and in the case where this orf is replaced (by genetic manipulation) by another orf whose product is a cytotoxic molecule, the latter is capable of destroying
10 only said microspores.

A subject of the invention is therefore also cellular expression vectors, comprising a promoter sequence such as that described above, placed upstream of a DNA sequence encoding a cytotoxic product.

15 Advantageously, the cytotoxic product in question is a protease. Specifically, when the protease is expressed specifically in the microspores, it destroys all the proteins thereof, as a result of which the microspore cannot survive. Preferably, the protease
20 is a subtilisin, and in particular the BPN' subtilisin from *Bacillus amyloliquefasciens*. This BPN' subtilisin is part of the family of subtilisins which are found in many organisms and which are proteases known to cleave proteins at the level of serines.

25 It involves, therefore, introducing a vector in accordance with the invention into a bacterial strain capable of carrying out the transformation of plant cells, such as *Agrobacterium tumefaciens*. This may in particular be carried out by the method of infiltration
30 of *Arabidopsis thaliana* plants described by Bechtold et al., 1993. This technique consists in introducing the bacterium into the cells of the floral scapes by infiltration under vacuum. The plants are then bedded out under glass and their seeds harvested. About one
35 seed in a thousand gives rise to plants of which all the cells carry the transgene. The transformation of other plants, and in particular of rape, may be carried out through *Agrobacterium tumefaciens* and/or



Agrobacterium rhizogenes with the aid of various techniques, now conventional (transformation of foliar disks, of hypocotyls, of floral scapes etc) which combine a phase of coculture of the bacterium with the plant tissues, followed by the selection and by the regeneration of the transformed cells into whole plants. Other transformation techniques do not use this bacterium, but make it possible to transfer the cloned gene directly into cells or tissues (electroporation, particle gun etc), and to select and obtain transformed plants (review by Siemens and Schieder).

A subject of the present invention is also the cells of plants transformed with a vector in accordance with the invention and plants comprising said cells.

A subject of the invention is also plants with gametophytic male sterility with inducible fertility, comprising a gene encoding a male-gamete-specific cytotoxic product which is placed downstream from the sequence described above.

As indicated above, the present invention thus enables the production of plants with gametophytic male sterility which inhibits any production of pollen grains. However, these plants, which are homozygous as regards their male sterility, may be obtained only after self-fertilisation of plants which have previously been transformed with a vector in accordance with the invention, i.e. which are hemizygous as regards their male sterility and in which the fertility of the pollen grains carrying the gametophytic sterility has been provisionally restored, so as to allow them to carry out self-fertilisation.

One means of producing plants which are homozygous for this gene would be to use gynogenesis, a technique which consists in regenerating doubled haploid plants from ovule or ovary culture. It involves, in this case, obtaining the formation of a homozygous diploid plant from a female haploid gamete. Gynogenesis is applicable to a certain number of plant species, but production of a large number of plants




which are homozygous for the transgene in question is not envisagable by this technique, because it is tricky to use and its efficacy most often remains very poor.

The present invention also concerns a method for producing plants
5 with gametophytic male sterility with inducible fertility comprising a gene encoding a male-gamete-specific cytotoxic product placed downstream from the sequence according to claims 1 and 2, comprising:

- the insertion into plants of line A of a gene whose expression product is cytotoxic for the microspores, and
- 10 - the production of plants which do not produce male gametes.

More particularly, the method for producing plants with gametophytic male sterility with inducible fertility in accordance with the invention comprises the steps of:

- a) transformation of plants of a line A with a vector in accordance
15 with the invention,
- b) induction of the fertility of the plants obtained in a) by inhibition of the cytotoxicity of the product,
- c) self-fertilisation of the fertile plants obtained in b),
- d) selection of the plants which do not produce male gametes,
20 derived from c).
- e) multiplication of the plants obtained in d) by repeating the steps b) and c).



Thus, in step a) of the method above, a line A is transformed with a vector in accordance with the invention, i.e. comprising a microspore-specific promoter sequence placed upstream of a gene encoding a cytotoxic product. The plants resulting from this transformation all comprise the DNA
25 in question whose gene is expressed only in the microspores. However, at this stage, the plant being diploid at the time of the transformation, it becomes heterozygous as regards its male sterility and is therefore capable, after transformation, of giving rise to microspores of which



only 50% are viable (the other 50% being destroyed following the expression of the transformant).

In step b), the restoration or the induction of the fertility which has been lost by the transformed
5 plant is then carried out by inhibiting the toxicity of the product of the transforming gene.

This may be done in various ways. However, when the cytotoxic product in question is a subtilisin and in particular the BPN' subtilisin from *Bacillus*
10 *amyloliquefasciens*, the inhibition is achieved by the action on the transformed plant of an insecticide molecule of the fluorophosphate family (having a priori no action on the plants). Indeed, this molecule, applied during anthesis, is capable of restoring the
15 total fertility of the hemizygous plants by inhibition of the subtilisin. It may, for example, be applied to the foot of the plant and reach all the tissues. As an insecticide, it should have no effect on the plant. However, it will have its full effect at the level of
20 the microspores, the only organs which express subtilisin.

Next, in step c), the self-fertilization of the plants whose fertility has been restored is carried out, then, in step d), the plants which are homozygous
25 with respect to male sterility, and consequently totally sterile in the absence of treatment i.e. of inhibition of the cytotoxic product, are selected.

The plants thus obtained, which are incapable of producing male gametes but still capable of
30 producing female gametes, i.e. ovules, may be crossed with another line of plants which are totally fertile and have valuable agronomic properties. In this cross, the plant which is homozygous as regards male sterility plays the role of the female parent whilst the other
35 plant plays the role of the male parent. The hybrid resulting from this cross is hemizygous as regards male sterility, and is therefore capable of producing pollen grains of which 50% are viable (the others, carrying



the transgene, are thus destroyed by the cytotoxic product). A 50% production of the pollen is more than enough to give rise to seeds having the qualities of each of the crossed lines which it is specifically
5 desired to combine.

The present invention thus also concerns a method for producing hybrid plants, characterized in that it comprises crossing plants of line A, which have gametophytic male sterility as described above, with
10 plants of line B of agronomic value. It also concerns the seeds derived from the hybrid plants thus obtained.

Advantageously, the plants in accordance with the invention belong to the Brassicacea family; preferably, they are rape.

15 In addition, it should be pointed out that the promoter region in accordance with the invention may also be used in strategies of gene inactivation by utilization of mobile elements such as transposons and retrotransposons.

20 Specifically, this may be carried out with the aim of isolating plants which have a stable mutant genotype, and isolating a very large number of different, independent mutants.

It involves creating a chimeric sequence
25 consisting of a promoter region in accordance with the invention and of the sequence, all or in part, of a mobile element. The expression of this mobile element, which is reduced to the phase of development of the microspore, should make it possible to induce some
30 mutations into the genome of the pollen grains of the transformed plant. It is thus possible, in the lineage obtained from these pollen grains, to isolate individuals which no longer carry the transgene, but merely one or more mutations derived from transposition
35 phenomena. The principle is to bring about, using the abovementioned promoter region, activation of the transposition of these mobile elements for a very short time (microsporogenesis) in a multitude of gametic



cells and to eliminate in the following generation the plants which carry the transgene (i.e. the promoter region + the sequence which allows the activation of the transposition) so that the cycle does not start up again. It then involves investigating, in the lineage, and by various techniques, the plants for which the mobile elements have caused mutations by inserting themselves into genes. The study of these plants would make it possible, in particular, to understand the function of the mutated gene.

Among the mobile elements which can be used in this way, mention may be made of the retrotransposons of the type Tnt1, Ttol, Tnp-2, Tos10-17, Bsl, BARE-1, Ta-1, etc., or the transposons of the type Ac/Ds, Spm, Mu, etc.

Figure 1 illustrates the alignment of the sequences of the two cDNAs M3 (SEQ ID No. 1) and M3.21 (SEQ ID No. 2) derived from the screenings of the *Brassica napus* cv.Brutor microspore cDNA library. The start (ATG) and stop (TGA) codons of the putative coding sequence are underlined.

Figure 2 gives the nucleotide sequence of the clone BnM3.4 (SEQ ID No. 3) from which the M3 cDNA is thought to be derived. The ATG in bold (position 2085) is the one which has the highest probability of being the functional ATG. The ATG underlined in position 2112 is the one present in the M3.21 cDNA sequence. The ATG underlined in position 1965 is the first ATG encountered. The sequence preceding these ATGs is, consequently, taken to be the promoter region of the BnM3.4 gene.

Figure 3 illustrates the Northern Blot hybridization with the P³²-labeled M3 probe on total RNAs (10 µg per well) extracted from different rape tissues. A: buds of 0-2 mm (meiocytes); B: buds of 2-3 mm (mononucleated microspores); C: buds of 3-4 mm (binucleated microspores); D: buds greater than 4 mm (mature pollen grains); E: rape sepals; F: rape



pistils; G: buds of sterile male rape; H: full buds of rape.

Figure 4 illustrates the preparation of the 7152 bp pJD51 plasmid from the 5135 bp pAF1 plasmid (plasmid of origin: pBluescript SK-PROMEGA) and from the 5458 bp pBnB2 plasmid (plasmid of origin pBS SK-PROMEGA).

Figure 5 illustrates the preparation of the 19670 bp pJD101 plasmid from the 15400 bp pEC2 plasmid which is derived from the pDHB 321.1 plasmid (D. Bouchez, personal communication) and from the pJD51 plasmid (cf. Figure 2).

Figure 6 represents a scheme of selection of hybrid varieties of a plant (rape for example) which calls upon a system of gametophytic male sterility with induction of the fertility. SMGfi: gametophytic male sterility with inducible fertility; Induction F: induction of the fertility; AF: self-fertilization.

The invention is not limited to the sole description above; it will be better understood in the light of the following examples, which are given, however, purely as illustrations.

EXAMPLE 1: Demonstration of a microspore-specific promoter

The first step consisted in producing complementary DNA (cDNA) clones expressed specifically in the microspore of rape. For this, cDNAs were synthesized from rape microspore messenger RNAs (mRNA). In parallel, cDNAs were synthesized from floral bud mRNA from sterile male rape. The cDNAs coming from said floral buds were subtracted from the cDNAs derived from the mRNAs expressed in the microspore of rape. The molecules resulting from this subtraction were used in an experiment of differential hybridization of a microspore cDNA library, according to a technique similar to that presented by Atanassov et al. (1996).



One of these isolated clones, the M3 cDNA (SEQ ID No. 1), proved to be the representative of an mRNA which is specifically expressed in the microspore of rape. Another cDNA, named M3.21 (SEQ ID No. 2) was found by screening the library with the M3 cDNA. The sequences of these two cDNAs show 89% identity (Figure 1); they are clearly derived from a family of very close genes, which are expressed specifically in the microspore.

The M3 cDNA clone was used as a probe to screen a rape genomic DNA library sold by CLONTECH Laboratories, Inc., 4030 Fabian Way, Palo Alto, CA 94303-4607, USA; two clones (BnM3.4 and BnM3.2) corresponding to two different genes were isolated. The M3 cDNA is thought to be derived from the BnM3.4 (SEQ ID No. 3) gene, because this gene carries an orf which is identical to the M3 cDNA (Figure 2). This gene has no intron. Sufficient experimental results lead to the thought that the M3.21 cDNA is not derived from the second isolated gene (BnM3.2), which indeed carries a region corresponding to the M3.21 cDNA sequence, but to a third gene, which is very close to the BnM3.2 gene.

The promoter region of this gene is defined as being the sequence immediately upstream of the translation start codon (ATG).

EXAMPLE 2: Verification of the specificity of the promoter of the BnM3.4 gene

A/ Northern Blot

A Northern Blot analysis was carried out with 10 µg of total RNA from sepals, pistil, whole buds, buds from sterile male plants, meiocytes, microspores, binucleated pollen grains and trinucleated pollen grains, hybridized with the M3 cDNA. A band of 1 kb corresponds to the transcript of the BnM3.4 gene, and also to the M3.21 transcript, since they are very close sequences. These transcripts are present uniquely in the first two stages of male gametogenesis, whose



products are difficult to isolate perfectly experimentally (Figure 3).

The proteins deduced from these two cDNA clones are evidently very close and are rich in glycine and proline. They are identical to strictly no other protein in the databanks, but are certainly involved in the formation of the wall.

B/ Transformation with a chimeric gene

Different chimeric genes (i.e. consisting of the sequence encoding a known gene, preceded by the promoter region in accordance with the invention) were constructed in order to study the spatio-temporal specificity of the BnM3.4 promoter.

Figure 4 shows the construction of a bacterial vector pJD51, which combines a fragment of the BnM3.4 promoter with the sequence encoding the β -glucuronidase gene. The pAF1 plasmid containing the sequence encoding β -glucuronidase and the transcription termination sequence of the NOS gene from *Agrobacterium tumefaciens*, was digested with the enzymes BamHI and ClaI. The pBnB2 plasmid contains a 6 kb BamHI-BamHI fragment derived from the BnM3.4 genomic DNA clone, and in which the BnM3.4 gene is present. A fragment corresponding to the largest promoter region possible given the restriction sites (2056 bp) was isolated from the pBnB2 plasmid by a BamHI-NspV digestion, and inserted between the BamHI and ClaI (compatible with NspV) sites of the pAF1 plasmid.

The chimeric gene thus constructed was isolated by a NotI digestion of the pJD51 plasmid, so as to be cloned into a binary plasmid from *Agrobacterium tumefaciens*: pEC2 opened by the enzyme NotI (Figure 5).

The pJD101 binary plasmid containing the chimeric gene was introduced into the C58C1 strain (pMP90) of *Agrobacterium tumefaciens* (Koncz et al. 1986) by electroporation, and the transformants possessing pJD101 were selected on a medium containing kanamycin. One of these *Agrobacterium* transformants was



used to transform *Arabidopsis thaliana* (Wassilevskja ecotype) by the method of infiltration of the floral scapes described by Bechtold et al., 1993. The transformed plants are selected using their resistance
5 to phosphinothrycin, which is conferred by a resistance gene jointly inserted into the T-DNA.

Among these plants, certain show expression of the β -glucuronidase specifically in the microspores (demonstrated by a blue coloration when a
10 β -glucuronidase-specific substrate, X-Glu, is added). No coloration is present in the adjacent tissues of the anther, nor in the somatic tissues of the plant. In a transformed plant which is hemizygous for the chimeric gene, half the microspores produced are blue, because
15 only they contain the chimeric gene.

The specificity of expression conferred by this 2 kb promoter sequence is indeed restricted, within the limits of the sensitivity of the technique, to a single cell type, and from the microspore stage.



REFERENCES

- Atanassov I et al. (1996) Plant Science 118, 185-194
5
- Bechtold N. et al. (1993) Comptes-Rendus de l'Académie
des Sciences 316, 1194-1199
- 10
- Koncz et al. 1986) Molecular General Genetics 204,
383-396
- 15
- Mariani et al. Nature 347 (1990) 737-741
- Oldenholf M.T. et al. (1996) Plant Molecular Biology
31, 213-225
20
- Siemens and Schieder 1996. Plant Tissue Culture and
Biotechnology, 2, 66-75



Editorial Note
92707/98

The following Sequence Listing pages 1 to 4 are part of
the Description.

The Claims pages follow starting from page number 15.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT:
(A) NAME: INRA (INSTITUT NATIONAL DE
LA RECHERCHE AGRONOMIQUE)
(B) STREET: 147 RUE DE L'UNIVERSITE
(C) CITY: PARIS
(E) COUNTRY: FRANCE
(F) POSTAL CODE: 75007

(ii) TITLE OF THE INVENTION: Microspore-specific
promoter and method for producing hybrid plants

(iii) NUMBER OF SEQUENCES: 3

(iv) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version
#1.30 (EPO)

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 497 base pairs
(B) TYPE: nucleotide
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:
(A) NAME/KEY: M3

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TTTGGATCTT TCCATGACCC CTTCTTGACC GGCTATGTCA AGCTACATTG CTCACCCGTT	60
GTGGATCTA CTTACCTCC TCCTCACAG GCTCCTTAC ATGCTCCTTC TTCACAGGCT	120
CCTTCACATG CTCCTTCACA TGCTCCTTCA CAGGCTCCTT TAAATGCTCT TTTAAATGCT	180
CCTTTACATG CTCCTTTACA TGCTCCTTCA CAGGCCCTT CACAGGCCCC TTCACAGGCC	240
CCTTTACATG CTCCTTTACT GCCCCTTCG CAGGCTCCTT CACCGGCTCA GTGATTAGC	300
TATTTGATAG AATTACTCAA GTAATGATGC CCTAGGGACT TTGAGTTTTT CTCGTGTTTT	360



AAAGTTTTGT GTTTATTTTG AGAAAACCGT CTTTGGATTT TAACTTCACT TTGATTTTTT 420
CCCTTATACA ATTTAAATTT AGAGTTTACT TATTAATTTT ATAAATTAGA TGGTACTAAG 480
TTTTTATCAT AATAAAA 497

(2) INFORMATION FOR SEQ ID NO: 2:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 674 base pairs
(B) TYPE: nucleotide
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- (ix) FEATURE:
(A) NAME/KEY: M3.21

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

TCCTGCTATG ATTTTCTTCA TAAGATGTGT CACATCCAAA GTCACAGCAA CAGAACTAGA 60
GTCATCAACT AACCAAGAGC TCTTCCTATC GCGGCACTTG CCTCGCTTTC ACCCCAAGCC 120
ACATTGGCCG TTCTGTGGCT CCGGAAAAGC CTTCCCTGCA GGCCACTTCC GACCAACTCC 180
GTTCCATCTG CCACAGGAAG TCACCAGATG CTTGTCCGAC AAGAAGGAGG TAGGTACATG 240
TTTGTATGAT ATCGTTGAGA CTTTCTTCAC CAGGAAAGCC GTTATTGGAT CGGAATGTTG 300
CGCCGCGATC AAGAAGATGA ACAAAGATTG TGAGAAGACC GTCTTTGGAT CTTTCCATGA 360
CCCCTTCTTG ACAGGCTATG TCAAACTACA TTGCTCCACC GTTGTGGAT CTACTTCACC 420
TCCTCCTTCA CATGCTCCTT CACAGGCTCC TTTACATGCT CCTTCACAGG CTCCTTTACA 480
TGCCCCTTCA CAGGCTCCTT TACTGCCCCC TTCACAGCCT CTCACCACGG CTCAGTGATT 540
TTAGCTATTT GTTAGAATTA TTCAAGTGTG GATGTCCTAG GGAGTTTtag GTTTTTCTTG 600
TTTTAAATTT TTGTGTTTAT TTGAGAAAA CCGTCTTTGG ATCTTAACCT CACTTTGATT 660
TTTTCCTTAT ACAA 674

(2) INFORMATION FOR SEQ ID NO: 3:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 2853 base pairs
(B) TYPE: nucleotide
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA



(ix) FEATURE:

(A) NAME/KEY: BnM3.4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGATCCCA	AAGAAACCG	AAGAAGCAGA	TGTTTCCTAC	CTTCATAAAT	ATATATTGT	60
TTCAGCCTCA	TCAATGTACA	AACAATCCTT	TAGCTCAATG	GTATAAATGT	TGTTGTTTAG	120
ATTTCAATAA	CCCGGGTTCG	AGTCATAGAC	TTGACACTTT	TTACACITTT	TTAAAAGTGG	180
AACGCACATA	TCGCTGACGT	GTGCGATCAG	GAGTGATGCA	ACTGCTCTAT	TATAATGTAG	240
ATTTAAAAGT	GGAACCCACG	TATCGCTGAC	GTGTCGCATC	AGGAGTGATG	CAACTGCCAT	300
ATTATAACGT	AGATTTGACG	TTATTCCTTT	TTAAATCTTA	ATAATAATAC	CAGNGCTTTT	360
ACTTATTAAT	TTTNGCATN	GTTATCATGG	TTTATGCNCT	CTTTTTTTTT	GANCCGTIGA	420
TTGGTTTATG	CTTATTTGAA	TGTNGCCNAC	GTAAGAAATG	AAGAACAATT	TATATTGGGA	480
GAAAATATAA	TTTAATATGT	TCAATATATA	GAGAAAATAT	TATNCCTTGA	TGTTACTGTA	540
TGGATGCGAG	TAGAAGATCT	TTGAATAATA	TTTGAGAACT	TGCCTTTTCT	CAAAAAGTAA	600
AATATTTGAT	ATGTAACCTA	AGTTAACACA	TGAAAATTAA	AAAAAAATTA	AATCAAAATA	660
GAAAAAAGTG	ATAGTGATCT	ACCCCTCAAC	GTTTTGAAGT	TATTCCTGGT	TCACCCCTTA	720
AACCTCTAAG	TTACCCAAAC	AATAAAATTT	CATTATTGCA	TATTCATAT	CTTTTAGAAA	780
GTGAACAAA	ATATTATCAA	GTTATATTAT	GTTTTTCAAA	TAAAAGATA	AAAAATAAAT	840
AAAAAATAAT	AGTAGTTACA	AAAAAATAAT	TTACACGCGT	CANAAAAACAC		900
TAAAACCTAA	ACCCTAAATA	TAAACCTTTT	AGGTAAACCC	TAAACCTTTG	GATAAATCTT	960
AAACATTAAT	CATTAAACAA	CTAAACCTTA	AATCCTAAAC	TCTAAACCTT	TAAGTGTTTA	1020
AATGTTTAGT	GTTTTTGATT	TATAGTTTAG	GATTATCCA	AAGGTTTAAG	GTTTACCCAA	1080
GAGTTTATGG	TTTAGGGATT	ATGACTTAGG	ATTTAGTGTT	TTACTGACGA	CGTTCAAAGT	1140
ATTTTTTAAA	AAATATTTTT	TTTGAACAA	CTACTATTTT	TATTTATTTT	TTTACCTTTT	1200
TATATTAAAA	ACATAATATA	ATTTAATACT	CCATCTGITT	CATATTAAGT	GTCAATTGTAA	1260



CATTATTTTT	TTGTTACAAA	AAAATTGTCA	CTTTAGAATT	CCAATGCAAA	ATTTATTTAT	1320
TTTTCAGCTA	AAATTAAATG	CAAAGTGCAT	TGATCTTATA	AATAATTTTA	TTTATCTCAA	1380
ATGCTATATT	GGTCAAACAT	GTGTAATTAA	TAGAAACTTA	ATTATATTTT	ATTTATTTTT	1440
TCCTAATCTG	TGTA AAAATG	TCAAAGTAAA	ATTTATTTAG	AAACGAATTG	AGTAATATTT	1500
TGTTTCATTT	TTTAAAAGAT	ATCGAATATG	AAATAACACA	ATTTTATTGT	ATGATGAACC	1560
TAAAAATTCA	TCCTAAGAAG	GTGAACGCAA	GAATAAGTCA	ACGTTTTGGG	GAAAGCTAAC	1620
TATGGCCCAA	AGTCATCAAA	ATCTTTCTTG	TATTTATCAA	AATCCTTACA	AATTTAGTTA	1680
GAGTTAATAG	ACCAAACACA	TGATTATCAT	CATATTAGAA	TATTCTAAAA	AATTACTAGC	1740
GAATAATTAA	AATCTTTCTT	TTATTTATCA	AAATCCTTAT	AAAACTTAT	TTATATATAC	1800
TAAAAACAATT	TTAATTAAAA	GAAAATAAGG	GACCATGGAT	ACATAAAAAAT	ATATGTTATT	1860
TCTTAAGATA	GTGATAATAT	TAATATATAC	CAGTCCATAT	ATTTATCAAA	ATAAATAATA	1920
TTTTTCGTAG	TCCGATAATC	ATTACTATAA	ATTCATAAAA	CCACATGTAG	ATGTATATTT	1980
TATTTATATA	TATATATATA	AACCCTAACG	CCTTACCACT	CGATAACCAT	CAAACTTTT	2040
CTTCTCGTTT	CGCTAACTCA	AGGCTTCGAA	AAGTAAAAAA	AACAATGAAG	AATGTCACAC	2100
TTGTTCTTGC	TATGATCCTC	TTCTTAAGCT	GTGTCACATC	CAAAGTTACA	GCAACAGAAC	2160
TAGAGTCATC	AACTARCCAA	GAGCTCTTCC	TATCGGGGCA	CTTACCTCGC	TTTACCCCA	2220
AGCAACATTG	GCCGTTCCGT	GGCTCCGGAA	AAGCCTTCCC	TGCAGGCCAC	TTCCGACTAA	2280
CTCCGTTCCA	TCTGCCACAG	GAAGTCACCA	GATGCTTGAA	CGACAAGAAG	GAGGTAGGTA	2340
CATGTTTTAA	TGATATCGCT	GAGACTTTCT	TCACCAGGAA	AGCCGCTATT	GGATCGGAAT	2400
GTTGCGCCGC	GATCAAGAAG	ATGAACAAAG	ATTGTGAGAA	GACCGTCTTT	GGATCTTTCC	2460
ATGACCCCTT	CTTGACCGGC	TATGTCAAGC	TACATTGCTC	CACCGTTGTT	GGATCTACTT	2520
CACCTCCTCC	TTACAGGGCT	CTTTTACATG	CTCCTTCTTC	ACAGGCTCCT	TCACATGCTC	2580
CTTACATGC	TCCTTCACAG	GCTCCITTTAA	ATGCTCCTTT	AAATGCTCCT	TTACATGCTC	2640
CTTTACATGC	TCCTTCACAG	GCCCTTTCAC	AGGCCCTTTC	ACAGGCCCTT	TTACATGCTC	2700
CTTTACTGCC	CCCTTCGCAG	GCTCCTTCAC	CGGCTCAGTG	ATTTAGCTAT	TTGATAGAAT	2760
TATTCAGTA	TTGATGICCT	AGGGAGTTTT	AGTTTTTTTC	TTGTTTTAAA	ATTTTGTGTT	2820
TATTTTGAGA	AAACCGTCTT	TGGATTTTAA	CTT			2853



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An isolated polynucleotide molecule having the nucleotide sequence corresponding:
 - a) the sequence according to SEQ ID. No. 3, or
 - 5 b) a sequence which has at least 80% homology with a).
2. An isolated polypeptide molecule having the nucleotide sequence according to Claim 1, corresponding:
 - 10 a) the sequence which stretches from nucleotide 1 to nucleotide 2111, preferably from nucleotide 1 to nucleotide 2084 of SEQ ID No. 3, or
 - b) a sequence which has at least 80% homology with a).
3. Cellular expression vector, comprising a sequence according to Claim 2, placed upstream of a DNA sequence encoding a cytotoxic product.
- 15 4. Vector according to Claim 3, characterised in that the cytotoxic product is a protease and preferably a subtilisin.
- 20 5. Plant cells transformed with a vector according to Claim 3 or 4.
6. Plants comprising cells according to Claim 5.
- 25 7. Plants with gametophytic male sterility with inducible fertility, comprising a gene encoding a male-gamete-specific cytotoxic product which is placed downstream from the sequence according to claims 1 and 2.



8. Method for producing plants with gametophytic male sterility with inducible fertility comprising a gene encoding a male-gamete-specific cytotoxic product placed downstream from the sequence according to claims 1 and 2, comprising:

- 5 - the insertion into plants of line A of a gene whose expression product is cytotoxic for the microspores; and
- the production of plants which do not produce male gametes.

- 10 9. Method for producing plants with gametophytic male sterility with inducible fertility according to Claim 8, comprising the steps of:

- a) Transformation of plants of a line A with a vector according to Claim 3 or 4,
- b) Induction of the fertility of the plants obtained in a) by inhibition of the cytotoxicity of the product,
- 15 c) Self-fertilisation of the fertile plants obtained in b),
- d) Selection of the plants which do not produce male gametes, derived from c),
- e) Multiplication of the plants obtained in d) by reproduction of steps b) and c).
- 20

10. Method for producing plants according to Claim 8 or 9, characterised in that, when the cytotoxic product is a subtilisin, the induction of the fertility consists in applying to the plant an insecticide molecule of the fluorophosphate family.
- 25

11. Seeds derived from the hybrid plants obtained by crossing plants of line A, which have gametophytic male sterility with inducible fertility, according to Claim 7, or as obtained by using the method according to one of Claims 8 to 10, with plants of line B of agronomic value.
- 30



12. Plants according to Claim 7, or obtained by using a method according to any one of Claims 8 to 10, characterised in that they belong to the Brassicacea family and preferably in that they are rape.
- 5 13. A method for producing plants with gametophytic male sterility with inducible fertility as defined in Claim 8 substantially as hereinbefore described with reference to the Examples.



DATED:

10

FREEHILLS CARTER SMITH BEADLE

Patent Attorneys for the Applicant

INSTITUTE NATIONAL DE LA RECHERCHE AGRONOMIQUE



```

80      1CTTGGTATG ATTTCTCTCA TAAGATGTGT CACATCCAAA GTCCACAGCA CAGAACTAGA GTCATCAACT AACCAAGAGC
M3      M3.21
160      1CTTCTCTATC GCGGCACTTG CCTCGCTTTC ACCCCAAAGC ACATTGGCCG TTCTGTGGCT CCGGAAAGC CTTCCTCTGA
M3      M3.21
240      GGGCACTTCC GACCAACTCC GTTCCATCTG CCACAGGAAG TCACACAGATG CTGTCCGAC AAGAAGGAGG TAGGTACATG
M3      M3.21
320      TTTTCATGAT ATCGTTGAGA CTTCTCTCAC CAGGAAGCC GTTATTTGGT CGGAATGTTG CCGCCGCGATC AAGAAGATGA
M3      M3.21
400      ACAAGATATG TGAGAAGACC GTTCTTTGGAT CTTTTCATGA CCCCCTCTCTG ACCGGCTATG TCAAGCTACA TTGCTCCACC
M3      M3.21
480      GTTGTGGAT CTACTTCACC TCCTCCTTCA CAGGCTCCTT TACATGCTCC TTCTTCACAG GCTCTCTCAC ATGCTCTCTC
M3      M3.21
560      ACATGCTCCT TCACAGGCTC CTTTAAATGC TCTTTAAAT GCTCTCTTAC ATGCTCTCTT ACATGCTCCT TCACAGGCTC
M3      M3.21
640      CTTTCACAGC CCCCTTCACAG GCCCTCTTAC ATGCTCTCTT ACTGCCCCCT TCGCAGGCTC CTTCACGCGC TCAGTGA-TT
M3      M3.21
720      TAGCTATTGG ATAGAATTAC TCAAGTAATG ATGCCCTAGG CAGTTTGAGT TTCTCTCGTG TTTTAAAGTT TTGCTTTAT
M3      M3.21
800      TTTGACAAA CCGTCTTTGG ATTTTAACCT CACTTTGAT TTTTCCCTTA TACAATTAA ATTTAGAGTT TACTATTAA
M3      M3.21
841      TTTTATAAAT TAGATGGTAC TAGCTTTTAA TCATAATAAA A
M3      M3.21

```

FIGURE 1

FIGURE 2

1 GGATCCCACA AAGAAAACCG AAGAAGCAAA TGTTCCTAC CTTCAATAAT
51 ATATATTTGT TTCAGCCTCA TCAATGTACA AACAACTCTT TAGCTCAATG
101 GTATAAATGT TGTGTTTAG ATTTCAATAA CCCGGGTTCC AGTCATAGAC
151 TTGACACTTT TTCACACTTT TTAAGTGG AACGCACATA TCGCTGACGT
201 GTCGCATCAG GAGTGATGCA ACTGCTCTAT TATAATGTAG ATTTAAAGT
251 GGAACCCACG TATCGCTGAC GTGTCGCATC AGGAGTGATG CAACTGCCAT
301 ATTATAACGT AGATTGACG TTATTCCTTT TTAATCTTA ATAATAATAC
351 CAGNGCTTTT ACTTATTAAT TTGNGCATN GTTATCATGG TTTATGCNCT
401 CTTTTTTTTT GANCCGTTGA TTGGTTTATG CTTATTTGAA TGTNGCCNAC
451 GTAAGAAATG AAGAACAATT TATATTGGA GAAATATAA TTTAATATGT
501 TCAATATATA GAGAAATAT TATNCCTTGA TGTACTGTA TGGATGCGAG
551 TAGAAGATCT TTGAATAATA TTGAGAACT TGCCTTTTCT CAAAAAGTAA
601 AATATTGAT ATGTAACCTTA AGTTAACACA TGAAATTA AAAAAATTA
651 AATCAAAATA GAAAAAATG ATAGTGATCT ACCCTTCAAC GTTTTGAAC
701 TATTCTTGGT TCACCCCTA AACCTCTAAG TTCACCAAAC AATAAATTT
751 CATTATTGCA TATTCTATAT CTTTGTAGAA GTGAAACAAA ATATTATCAA
801 GTTATATTAT GTTTTCAAA TAAAAAGATA AAAAATAAT AAAAAATAAT
851 AGTAGTTACA AAAAAAAAAA ATTAATATTT TTACCAGCGT CANAAACAC
901 TAAACCTAA ACCCTAATA TTAACTTTT AGGTAAACCC TAAACCTTTG
951 GATAAATCTT AAACATTAAA CATTAAACA CTAAACCTA AATCCTAAAC
1001 TCTAAACCT TAAGTGTTA AATGTTTGT GTTTTGATT TATAGTTTAG
1051 GATTATCCA AAGGTTTAA GTTTACCAA GAGTTTATGG TTTAGGGATT
1101 ATGACTTAGG ATTTAGTGT TTAGTGACGA CGTTCAAAGT ATTTTATAA
1151 AAATATTTT TTTGTACAA CTACTATTTT TATTAATTTT TTTACCTTTT
1201 TATATTAAAA ACATAATATA ATTTAATCT CCATCTGTTT CATATTAGT
1251 GTCATGTAA CATTATTTT TTGTTACAA AAAATTGTCA CTTTGAATT

FIGURE 2 (continued)

1301 CCAATGCAAA ATTTATTTAT TTTTCAGCTA AAATTAATTG CAAAGTGCAT
1351 TGATCTTATA AATAATTTA TTTATCTCAA ATGCTATATT GGTCAAACAT
1401 GTGTAATTAA TAGAACTTA ATTATATTTT ATTTATTTTT TCTTAATCTG
1451 TGTAAAAATG TCAAAGTAAA ATTTATTTAG AAACGAATTG AGTAATATTT
1501 TGTTCATTT TTTAAAAGAT ATCGAATATG AAATAACACA ATTTTATTGT
1551 ATGATGAACC TAAAAATTCA TCCTAAGAAG GTGAACGCAA GAATAAGTCA
1601 ACGTTTGGG GAAAGCTAAC TATGGCCCAA AGTCATCAAA ATCTTCTTG
1651 TATTTATCAA AATCCTTACA AATTTAGTTA GAGTTAATAG ACCAAACACA
1701 TGATTATCAT CATATTAGAA TATTCTAAAA AATTACTAGC GAATAATTAA
1751 AATCTTCTT TTATTTATCA AAATCCTTAT AAAAATTAT TTATATATAC
1801 TAAACAATT TTAATTAAAA GAAAATAAGG GACCATGGAT ACATAAAAAAT
1851 ATATGTTATT TCTTAAGATA GTGATAATAT TAATATATAC CAGTCCATAT
1901 ATTTATCAAA ATAAATAATA TTTTTCGTAG TCCGATAATC ATTACTATAA
1951 ATTCATAAAA CCACATGTAG ATGTATATTT TATTATATA TATATATATA
2001 AACCCTAACG CCTTACCACT CGATAACCAT CAAAACTTTT CTTCCTGTTT
2051 CGCTAACTCA AGGCTTCGAA AAGTAAAAA AACAATGAAG AATGTCACAC
2101 TTGTTCTTGC TATGATCCTC TTCTTAAGCT GTGTCACATC CAAAGTTACA
2151 GCAACAGAAC TAGAGTCATC AACTAACCAA GAGCTCTTCC TATCGCGGCA
2201 CTTACCTCGC TTTCACCCCA AGCAACATTG GCCGTTCCGT GGCTCCGGAA
2251 AAGCCTTCCC TGCAGGCCAC TTCCGACTAA CTCGTTCCA TCTGCCACAG
2301 GAAGTCACCA GATGCTTGAA CGACAAGAAG GAGGTAGGTA CATGTTTTAA
2351 TGATATCGCT GAGACTTTCT TCACCAGGAA AGCCGCTATT GGATCGGAAT
2401 GTTGGCCCCG GATCAAGAAG ATGAACAAAG ATTGTGAGAA GACCGTCTTT
M3 TTT
2451 GGATCTTTCC ATGACCCCTT CTTGACCGGC TATGTCAAGC TACATTGCTC
M3 GGATCTTTCC ATGACCCCTT CTTGACCGGC TATGTCAAGC TACATTGCTC
2501 CACCGTTGTT GGATCTACTT CACCTCCTCC TTCACAGGCT CCTTTACATG
M3 CACCGTTGTT GGATCTACTT CACCTCCTCC TTCACAGGCT CCTTTACATG

2551 CTCCTTCCTC ACAGGCTCCT TCACATGCTC CTTACATGC TCCTTCACAG
M3 CTCCTTCCTC ACAGGCTCCT TCACATGCTC CTTACATGC TCCTTCACAG
2601 GCTCCTTTAA ATGCTCCTTT AAATGCTCCT TTACATGCTC CTTTACATGC
M3 GCTCCTTTAA ATGCTCCTTT AAATGCTCCT TTACATGCTC CTTTACATGC
2651 TCCTTCACAG GCCCCTTCAC AGGCCCTTC ACAGGCCCTT TTACATGCTC
M3 TCCTTCACAG GCCCCTTCAC AGGCCCTTC ACAGGCCCTT TTACATGCTC
2701 CTTTACTGCC CCCTTCGCAG GCTCCTTCAC CGGCTCAGTG ATTTAGCTAT
M3 CTTTACTGCC CCCTTCGCAG GCTCCTTCAC CGGCTCAGTG ATTTAGCTAT
2751 TTGATAGAAT TATTCAAGTA TTGATGTCCT AGGGAGTTT AGTTTTTTTC
M3 TTGATAGAAT TACTCAAGTA ATGATGCCCT AGGGAGTTG AGTTTTTCTC
2801 TTGTTTTAAA ATTTTGTGTT TATTTTGAGA AAACCGTCTT TGGATTTTAA
M3 GTGTTTTAAA GTTTTGTGTT TATTTTGAGA AAACCGTCTT TGGATTTTAA
2851 CTT
M3 CTT

FIGURE 2 (continued)




FIGURE 3




FIGURE 4.




FIGURE 4 (continued)




FIGURE 5.




FIGURE 5 (continued)




FIGURE 6