

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2015236340 B2

(54) Title
Lyophilized factor IX formulations

(51) International Patent Classification(s)
F26B 5/06 (2006.01)

(21) Application No: **2015236340** (22) Date of Filing: **2015.03.24**

(87) WIPO No: **WO15/148444**

(30) Priority Data

(31) Number **61/969,801** (32) Date **2014.03.24** (33) Country **US**

(43) Publication Date: **2015.10.01**
(44) Accepted Journal Date: **2020.02.06**

(71) Applicant(s)
Bioverativ Therapeutics Inc.

(72) Inventor(s)
Thome, Brian M.; Parkhurst-Lang, Cherie; Leveille, Brandon W.

(74) Agent / Attorney
Spruson & Ferguson, GPO Box 3898, Sydney, NSW, 2001, AU

(56) Related Art
US 2001/0038859 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2015/148444 A1

(43) International Publication Date
1 October 2015 (01.10.2015)

(51) International Patent Classification:
F26B 5/06 (2006.01)

(74) Agents: KIM, Ji-Eun et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., 1100 New York Avenue, NW, Washington, District of Columbia 20005 (US).

(21) International Application Number:

PCT/US2015/022141

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:

24 March 2015 (24.03.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/969,801 24 March 2014 (24.03.2014) US


(71) Applicant: BIOGEN MA INC. [US/US]; 250 Binney Street, Cambridge, MA 02142 (US).

(72) Inventors: THOME, Brian M.; 81 Dresser Street, Boston, Massachusetts 02127 (US). PARKHURST-LANG, Cherie; 138 Canterbury Lane, Londonderry, New Hampshire 03053 (US). LEVEILLE, Brandon W.; 11 Perry's Way, Berwick, Maine 03901 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]

(54) Title: LYOPHILIZED FACTOR IX FORMULATIONS



Empty Vials
rFIXFc LCM Placebo
= Thermocouple

(57) Abstract: The present invention provides, among other things, pre-lyophilization formulations, reconstituted formulations, and lyophilate powder compositions comprising a Factor IX (FIX) polypeptide. The present invention also provides lyophilization methods for producing lyophilate powder comprising a FIX polypeptide. The present invention is directed to a pre-lyophilization formulation comprising: (a) a Factor IX (FIX) polypeptide having FIX coagulation activity; (b) a buffering agent; (c) a stabilizing agent; (d) a bulking agent; and (e) a surfactant, wherein the formulation has a fill volume of less than about 5 mL, and wherein each of (a) - (e) are at an amount per vial (mg/vial) sufficient to allow (1) improved stability of the FIX polypeptide when lyophilized.

Fig. 1



SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, — *with sequence listing part of description (Rule 5.2(a))*
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

LYOPHILIZED FACTOR IX FORMULATIONS

BACKGROUND

[0001] The present invention relates generally to the field of therapeutics for hemostatic disorders.

[0002] Hemophilia B (also known as Christmas disease) is one of the most common inherited bleeding disorders in the world. It results in decreased *in vivo* and *in vitro* blood clotting activity and requires extensive medical monitoring throughout the life of the affected individual. In the absence of intervention, the afflicted individual will suffer from spontaneous bleeding in the joints, which produces severe pain and debilitating immobility; bleeding into muscles results in the accumulation of blood in those tissues; spontaneous bleeding in the throat and neck can cause asphyxiation if not immediately treated; renal bleeding; and severe bleeding following surgery, minor accidental injuries, or dental extractions also are prevalent.

[0003] Normal *in vivo* blood coagulation at minimum requires the serine proteases Factors II (prothrombin), VII, IX, X and XI (soluble plasma proteins); cofactors including the transmembrane protein tissue factor and the plasma proteins Factors V and VIII; fibrinogen, the transglutaminase Factor XIII, phospholipid (including activated platelets), and calcium. Additional proteins including kallikrein, high molecular weight kininogen, and Factor XII are required for some *in vitro* clotting tests, and can play a role *in vivo* under pathologic conditions.

[0004] In hemophilia, blood clotting is disturbed by a lack of certain plasma blood clotting factors. Hemophilia B is caused by a deficiency in Factor IX that can result from either the decreased synthesis of the Factor IX protein or a defective molecule with reduced activity. The treatment of hemophilia occurs by replacement of the missing clotting factor by exogenous factor concentrates highly enriched in Factor IX. However, generating such a concentrate from blood is fraught with technical difficulties, as is described below.

[0005] Purification of Factor IX from plasma (plasma derived Factor IX; pdFIX) almost exclusively yields active Factor IX. However, such purification of Factor IX from plasma is very difficult because Factor IX is only present in low concentration in plasma (5 μ g/mL. Andersson, Thrombosis Research 7: 451-459 (1975). Further, purification from

- 2 -

blood requires the removal or inactivation of infectious agents such as HIV and HCV. In addition, pdFIX has a short half-life and therefore requires frequent dosing. Recombinant factor IX (rFIX) is also available, but suffers from the same short half-life and need for frequent dosing (e.g., 2-3 times per week for prophylaxis) as pdFIX. rFIX also has a lower incremental recovery (K value) compared to pdFIX, which necessitates the use of higher doses of rFIX than those for pdFIX.

[0006] Reduced mortality, prevention of joint damage and improved quality of life have been important achievements due to the development of plasma-derived and recombinant Factor IX. Prolonged protection from bleeding would represent another key advancement in the treatment of hemophilia B subjects. However, to date, no products that allow for prolonged protection have been developed. Therefore, there remains a need for improved methods of treating hemophilia due to Factor IX deficiency that are more tolerable and more effective than current therapies.

[0007] In particular, there remains a need for improved lyophilized FIX formulations with higher drug product strength, longer shelf life, reduced lyophilization process time, and shorter reconstitution time.

BRIEF SUMMARY OF THE INVENTION

[0008] The present invention is directed to a pre-lyophilization formulation comprising: (a) a Factor IX (FIX) polypeptide having FIX coagulation activity; (b) a buffering agent; (c) a stabilizing agent; (d) a bulking agent; and (e) a surfactant, wherein the formulation has a fill volume of less than about 5 mL, less than about 4mL, or less than about 3 mL and wherein each of (a) – (e) are at an amount per vial (mg/vial) sufficient to allow (1) improved stability of the FIX polypeptide when lyophilized; (2) reduced reconstitution time when lyophilized; (3) reduced splashing onto a stopper comprising the formulation; (4) reduced lyophilization cycle time; (5) increased shelf-life of a lyophilate prepared from the pre-lyophilization formulation at room temperature; or (6) any combinations thereof, compared to a reference pre-lyophilization formulation, wherein the reference formulation comprises (a) – (e) at the amount per vial identical to the pre-lyophilization formulation, but has at least a 5mL fill volume. In a particular embodiment, the fill volume of the formulation is about 2.65mL.

- 3 -

[0009] In some embodiments, the pre-lyophilization formulation comprises at least 100 IU/vial of the FIX polypeptide. In some embodiments, the pre-lyophilization formulation comprises about 200 IU/vial to about 10,000 IU/vial of the FIX polypeptide.

[0010] In some embodiments, the FIX polypeptide comprises wild-type FIX. In some embodiments, the FIX polypeptide further comprises a heterologous moiety fused to wild-type FIX. In one embodiment, the heterologous moiety is a moiety extending half-life of FIX. In another embodiment, the heterologous moiety comprises a polypeptide or a non-polypeptide moiety. In one embodiment, the moiety extending half-life of FIX comprises an FcRn binding partner or an Fc region. In one embodiment, the FIX polypeptide is at least about 80%, at least about 85%, at least about 90%, at least about 95%, or 100% identical to SEQ ID NO: 2.

[0011] In some embodiments, the fill volume is about 4 mL, about 3.5 mL, about 3.0 mL, about 2.9 mL, about 2.8 mL, about 2.7 mL, about 2.65 mL, about 2.6mL, about 2.5 mL, about 2.4 mL, about 2.3 mL, about 2.2 mL, about 2.1 mL, or about 2.0 mL.

[0012] In some embodiments, the reduced reconstitution time is less than 1.5 minute, less than 1 minute, less than 50 seconds, less than 40 seconds, less than 30 seconds, less than 20 seconds, or less than 10 seconds.

[0013] In some embodiments, the buffering agent is L-histidine. In one embodiment, the buffering agent is at a concentration (mg/mL) between about 3 mg/mL and about 15 mg/mL. In another embodiment, the buffering agent is at a concentration between about 8 mg and about 39 mg per vial.

[0014] In some embodiments, the stabilizing agent is sucrose. In one embodiment, the stabilizing agent is at a concentration (mg/mL) between 10 mg/mL and about 50 mg/mL. In another embodiment, the stabilizing agent is at a concentration between about 27 mg and about 132 mg per vial.

[0015] In some embodiments, the bulking agent is mannitol. In one embodiment, the bulking agent is at a concentration (mg/mL) between 20 mg/mL and about 100 mg/mL. In another embodiment, the bulking agent is at a concentration between about 53 mg per vial and about 265 mg per vial.

[0016] In some embodiments, the surfactant is polysorbate 20. In one embodiment, the surfactant is at a concentration (mg/mL) between 0.01 mg/mL and about 5 mg/mL. In

another embodiment, the surfactant is at a concentration between about 0.03 mg and about 13 mg per vial.

[0017] In one aspect, the invention is directed to pre-lyophilization formulation comprising: (a) about 80 to about 2,750 IU/mL of rFIXFc; (b) about 7.76 mg/mL of L-histidine; (c) about 47.6 mg/mL of mannitol; (d) about 23.8 mg/mL of sucrose; and, (e) about 0.2 mg/mL of polysorbate-20.

[0018] The present invention is further directed to a lyophilate powder comprising a FIX polypeptide, a buffering agent, a stabilizing agent, a bulking agent, a surfactant, or any combinations thereof.

[0019] In some embodiment, the residual moisture level of the lyophilate powder is below 1%.

[0020] In one embodiment, the lyophilate powder comprises: (a) a FIX polypeptide at an amount between about 2 mg per vial and about 150 mg per vial; (b) a buffering agent at an amount between 10 mg per vial and about 30 mg per vial; (c) a bulking agent at an amount between 70 mg vial and about 200 mg per vial; (d) a stabilizing agent at an amount between 30 mg per vial and 100 mg per vial; and (e) a surfactant at an amount between 0.05 mg per vial and about 5 mg per vial.

[0021] In another embodiment, the lyophilate powder comprises: (a) the lyophilized FIX polypeptide at an amount between about 2.2 mg per vial and about 125 mg per vial; (b) the buffering agent at an amount between about 12.5 mg per vial and 25 mg per vial; (c) the stabilizing agent at an amount between about 32.5 mg per vial and 80 mg per vial; (d) the bulking agent at an amount between about 75 mg per vial and 150 mg per vial; and (e) the surfactant at an amount between about 0.1 mg/mL and about 2 mg/mL.

[0022] In another embodiment, the lyophilate powder comprises: (a) about 2.2 to about 125 mg/vial of the FIX polypeptide; (b) about 20.6 mg/vial of L-histidine; (c) about 126.1 mg/vial of mannitol; (d) about 63.1 mg/vial of sucrose; and, (e) about 0.53 mg/vial of polysorbate-20;

[0023] The present invention is also directed to a reconstituted formulation comprising the lyophilate powder described herein reconstituted by a reconstitution buffer.

[0024] In one embodiment, the reconstituted formulation comprises: (a) the FIX polypeptide at a concentration between about 0.9 mg/mL and about 50 mg/mL; (b) the buffering agent at a concentration between 1.5 mg/mL and about 7.5 mg/mL; (c) the

bulking agent at a concentration between 10 mg/mL and about 50 mg/mL; (d) the stabilizing agent at a concentration between 5 mg/mL and 25 mg/mL per vial; and (e) the surfactant at a concentration between 0.005 mg/mL and about 2.5 mg/mL.

[0025] In another embodiment, the reconstituted formulation comprises: (a) the FIX polypeptide at a concentration between about 0.9 mg/mL and about 50 mg/mL; (b) the buffering agent at a concentration of about 3.88 mg/mL; (c) the bulking agent at a concentration of about 23.8 mg/mL; (d) the stabilizing agent at a concentration of about 11.9 mg/mL; (e) the surfactant at a concentration of about 0.1 mg/mL; and (f) the reconstitution buffer.

[0026] In another embodiment, the reconstituted formulation comprises: (a) the FIX polypeptide at a concentration between about 80 IU/mL and about 2,750 IU/mL; (b) the buffering agent at a concentration of about 25 mM; (c) the bulking agent at a concentration of about 131 mM; (d) the stabilizing agent at a concentration of about 35 mM; (e) the surfactant at a concentration of 0.01 % (w/v); and (f) the reconstitution buffer.

[0027] The present invention further pertains to a method of administering a FIX polypeptide to a hemophilia B patient in need thereof, or a method of preventing, treating, ameliorating, or managing hemophilia B in a patient in need thereof, comprising administering to the patient the reconstituted formulations described herein.

[0028] The present invention is also directed to a method of producing a lyophilate powder comprising a FIX polypeptide comprising lyophilizing the pre-lyophilization formulations described herein.

[0029] In one aspect, the present invention is directed to a method of lyophilizing a FIX polypeptide comprising: (a) a “freezing step” comprising freezing a pre-lyophilization formulation comprising the FIX polypeptide and an aqueous solvent; (b) a “vacuum step” comprising reducing the pressure of the frozen pre-lyophilization formulation by an amount effective to remove the aqueous solvent from the frozen pre-lyophilization formulation; and, (c) a single “drying step” comprising increasing the temperature of the frozen pre-lyophilization formulation above the collapse temperature, thereby producing a lyophilate powder. In some embodiments, the pre-lyophilization formulation is aseptically filtered and aseptically filled into a vial prior to step (a).

[0030] In another aspect, the present invention is directed to methods of producing a lyophilate powder comprising a FIX polypeptide, comprising: (a) a "freezing step" comprising freezing a pre-lyophilization formulation comprising a FIX polypeptide by ramping down the temperature for about 2 hours to a freezing temperature of about -55°C, and holding the freezing temperature for about 2 hours; (a') an "annealing step" comprising ramping up for about 1.5 hours the temperature of the frozen pre-lyophilization formulation of step (a) to an annealing temperature of about -6°C, holding the annealing temperature for about 3 hours, and ramping down the temperature for about 1.5 hours to about -55°C; (b) a "vacuum step" comprising holding the frozen pre-lyophilization formulation of step (a') at about -55°C for two hours at atmospheric pressure and ramping down the pressure for about 2 hours to about 0.33 mbar; and, (c) a single "drying step" comprising ramping up for 3 hours the temperature of the frozen pre-lyophilization formulation of step (b) to about 40°C, while holding the pressure at about 0.33 mbar, and holding the temperature of the frozen pre-lyophilization formulation at about 40°C for about 25 hours, while holding the pressure at about 0.33 mbar, thereby producing the lyophilate powder.

[0031] In a further aspect, the lyophilate powder has one or more characteristics selected from the group consisting of: (1) improved stability of the FIX polypeptide when lyophilized; (2) reduced reconstitution time when lyophilized; (3) reduced splashing onto a stopper comprising the formulation; (4) reduced lyophilization cycle time; (5) increased shelf-life of a lyophilate prepared from the pre-lyophilization formulation at room temperature; or (6) any combinations thereof,

[0032] In one aspect, the disclosure provides a method of stabilizing a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the lyophilate powder is stabilized as measured by Size Exclusion Chromatography (SEC) with respect to a lyophilate powder prepared by using a lyophilization method comprising more than one drying step.

[0033] In another aspect, the disclosure provides a method of increasing the shelf-life of a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the shelf-life of the lyophilate powder is increased as measured by SEC and/or FIX clotting activity

- 7 -

assay with respect to the shelf-life of a lyophilate powder prepared by using a lyophilization method comprising more than one drying step.

[0034] This disclosure also provides a method to decrease the reconstitution time of a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the reconstitution time of the lyophilate powder is decreased with respect to the reconstitution time of a lyophilate powder prepared by using a lyophilization method comprising more than one drying step.

[0035] This disclosure further provides a method to reduce lyophilization process time of producing a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the lyophilization process time of the pre-lyophilization formulation is reduced with respect to the lyophilization process time of producing a lyophilate powder using a lyophilization method comprising more than one drying step.

BRIEF DESCRIPTION OF THE FIGURES

[0036] Figure 1 shows the vial loading pattern for each lyophilization cycle. The numbers 1, 2, and 3 indicate the thermocouple locations.

[0037] Figure 2 shows the prediction profiler from the DOE analysis of residual moisture as a function of the lyophilization parameters – temperature, vacuum, and time.

[0038] Figure 3 shows the prediction profiler from the DOE analysis of product temperature during sublimation as a function of the lyophilization parameters – temperature and vacuum.

[0039] Figure 4 shows the prediction profiler from the DOE analysis of vial mass flow during sublimation as a function of the lyophilization parameters – temperature and vacuum.

[0040] Figure 5 shows the lyophilization data from DOE Run 8 in Example 2, which is similar to the proposed rFIXFc-2G lyophilization cycle (shelf temperature of 40 °C, a chamber vacuum of 250 mTorr (0.33 mBar) and a drying time of 25 hours).

DETAILED DESCRIPTION

[0041] This disclosure provides, among other things, pre-lyophilization formulations, reconstituted formulations, and lyophilate powder compositions comprising a Factor IX (FIX) polypeptide. The disclosure also provides lyophilization methods for producing lyophilate powder comprising a FIX polypeptide. Also provided are methods of stabilizing a lyophilate powder comprising a FIX polypeptide, method to increase the shelf-life of a lyophilate powder comprising a FIX polypeptide, method to decrease the reconstitute time of a lyophilate powder comprising a FIX polypeptide, and method to reduce lyophilization process time of a pre-lyophilization formulation comprising a FIX polypeptide. In addition, the disclosure provides methods of preventing, treating, ameliorating, or managing hemophilia B in a patient in need of by administering a reconstituted formulation comprising a FIX polypeptide.

Definitions

[0042] Throughout this disclosure, the term "a" or "an" entity refers to one or more of that entity; for example, "a polynucleotide," is understood to represent one or more polynucleotides. As such, the terms "a" (or "an"), "one or more," and "at least one" can be used interchangeably herein.

[0043] Furthermore, "and/or" where used herein is to be taken as specific disclosure of each of the two specified features or components with or without the other. Thus, the term "and/or" as used in a phrase such as "A and/or B" herein is intended to include "A and B," "A or B," "A" (alone), and "B" (alone). Likewise, the term "and/or" as used in a phrase such as "A, B, and/or C" is intended to encompass each of the following aspects: A, B, and C; A, B, or C; A or C; A or B; B or C; A and C; A and B; B and C; A (alone); B (alone); and C (alone).

[0044] It is understood that wherever aspects are described herein with the language "comprising," otherwise analogous aspects described in terms of "consisting of" and/or "consisting essentially of" are also provided.

[0045] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure is related. For example, the Concise Dictionary of Biomedicine and Molecular Biology, Juo, Pei-Show, 2nd ed., 2002, CRC Press; The Dictionary of Cell and Molecular

Biology, 3rd ed., 1999, Academic Press; and the Oxford Dictionary Of Biochemistry And Molecular Biology, Revised, 2000, Oxford University Press, provide one of skill with a general dictionary of many of the terms used in this disclosure.

[0046] Units, prefixes, and symbols are denoted in their Système International de Unites (SI) accepted form. Numeric ranges are inclusive of the numbers defining the range. Unless otherwise indicated, amino acid sequences are written left to right in amino to carboxy orientation. The headings provided herein are not limitations of the various aspects of the disclosure, which can be had by reference to the specification as a whole. Accordingly, the terms defined immediately below are more fully defined by reference to the specification in its entirety.

[0047] The term "about" is used herein to mean approximately, roughly, around, or in the regions of, and the value will depend on how limitations of the measuring system. When the term "about" is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term "about" can modify a numerical value above and below the stated value by a variance of, *e.g.*, 10 percent or 20 percent, up or down (higher or lower). Unless otherwise stated, the meaning of "about" should be assumed to be within an acceptable error range for the particular value for the formulation or composition.

[0048] The term "polypeptide," "peptide" and "protein" are used interchangeably and refer to a polymeric compound comprised of covalently linked amino acid residues.

[0049] The term "polynucleotide" and "nucleic acid" are used interchangeably and refer to a polymeric compound comprised of covalently linked nucleotide residues. Polynucleotides can be DNA, cDNA, RNA, single stranded, or double stranded, vectors, plasmids, phage, or viruses.

[0050] The term "administering," as used herein, means to, *e.g.*, prescribe or give a pharmaceutical composition comprising an FIX polypeptide to a subject. Examples of routes of administration include, but are not limited to, intravenous, *e.g.*, intravenous injection and intravenous infusion, *e.g.*, via central venous access. Additional routes of administration include subcutaneous, intramuscular, oral, nasal, and pulmonary administration. A pharmaceutical composition comprising an FIX polypeptide can comprise one or more excipients, as described herein. Advantages of the methods, compositions, and pharmaceutical kits provided herein include: improved regimen

- 10 -

compliance; reduced break through bleeds; increased protection of joints from bleeds; prevention of joint damage; reduced morbidity; reduced mortality; prolonged protection from bleeding; decreased thrombotic events; and improved quality of life. Administering includes parenteral administration. In some embodiments, the parenteral administration is intravenous or subcutaneous administration.

[0051] The term "treatment" or "treating" as used herein means amelioration or reduction of one or more symptoms of bleeding diseases or disorders including, but not limited to, hemophilia B. In one embodiment, "treatment of" or "treating" a bleeding disease or disorder includes prevention of one or more symptoms of a bleeding disease or disorder. In a bleeding disease or disorder caused by a FIX deficiency (*e.g.*, a low baseline FIX activity), the term "treatment" or "treating" can mean FIX replacement therapy. By administering a FIXFc polypeptide to a subject, the subject can achieve and/or maintain a plasma trough level of a FIX activity at about 1 IU/dl or above 1 IU/dl. In other embodiments, "treatment" or "treating" means reduction of the frequency of one or more symptoms of bleeding diseases or disorders, *e.g.*, spontaneous or uncontrollable bleeding episodes. "Treatment," however, need not be a cure.

[0052] "Patient" as used herein includes an individual who is known to have at least one incidence of uncontrolled bleeding episodes, who has been diagnosed with a disease or disorder associated with uncontrolled bleeding episodes, *e.g.*, a bleeding disease or disorder, *e.g.*, hemophilia B, who are susceptible to uncontrolled bleeding episodes, *e.g.*, hemophilia, or any combinations thereof. Patients can also include an individual who is in danger of one or more uncontrollable bleeding episodes prior to a certain activity, *e.g.*, a surgery, a sport activity, or any strenuous activities. The patient can have a baseline FIX activity less than 1%, less than 0.5%, less than 2%, less than 2.5%, less than 3%, or less than 4%. Patients also include pediatric humans. Pediatric patients are birth to 20 years, preferably birth to 18 years, birth to 16 years, birth to 15 years, birth to 12 years, birth to 11 years, birth to 6 years, birth to 5 years, birth to 2 years, and 2 to 11 years of age.

[0053] "Baseline," as used herein, is the lowest measured plasma Factor IX level in a subject prior to administering a dose. The Factor IX plasma levels can be measured at two time points prior to dosing: at a screening visit and immediately prior to dosing. Alternatively, (a) the baseline in subjects whose pretreatment FIX activity is <1%, who have no detectable FIX antigen, and have nonsense genotypes can be defined as 0%, (b)

the baseline for subjects with pretreatment FIX activity <1% and who have detectable FIX antigen can be set at 0.5%, (c) the baseline for subjects whose pretreatment FIX activity is between 1 – 2% is Cmin (the lowest activity throughout the PK study), and (d) the baseline for subjects whose pretreatment FIX activity is $\geq 2\%$ can be set at 2%. Activity above the baseline pre-dosing can be considered residue drug from prior treatment, and can be decayed to baseline and subtracted from the PK data following rFIXFBP dosing.

[0054] "Trough," as used herein, is the lowest plasma Factor IX activity level reached after administering a dose of chimeric polypeptide of the invention or another Factor IX molecule and before the next dose is administered, if any. Trough is used interchangeably herein with "threshold." Baseline Factor IX levels are subtracted from measured Factor IX levels to calculate the trough level.

[0055] As used herein, the term "half-life" refers to a biological half-life of a particular polypeptide in vivo. Half-life can be represented by the time required for half the quantity administered to a subject to be cleared from the circulation and/or other tissues in the animal.

[0056] The terms "long-acting" and "long-lasting" are used interchangeably herein. In one embodiment, the term "long-acting" or "long-lasting" indicates that a FIX activity as a result of administration of the rFIXFBP polypeptide is longer than the FIX activity of a wild-type FIX (*e.g.*, BENEFIX[®] or plasma-derived FIX ("pdFIX")). The "longer" FIX activity can be measured by any known methods in the art, *e.g.*, aPTT assay, chromogenic assay, ROTEM, TGA, and etc. In one embodiment, the "longer" FIX activity can be shown by the $T_{1/2\beta\alpha}$ (activity). In another embodiment, the "longer" FIX activity can be inferred by the level of FIX antigen present in plasma, *e.g.*, by the $T_{1/2\beta\alpha}$ (antigen).

[0057] The terms "lyophilate," "lyophilate powder," "lyophilized product," or "product cake," as used herein, denote a formulation which is manufactured by freeze-drying methods. The solvent (*e.g.* water) is removed by freezing following sublimation under vacuum and desorption of residual water at elevated temperature. In the pharmaceutical field, the lyophilate is present as a powder or a physical stable cake. The lyophilate is characterized by a fast dissolution after addition of a reconstitution medium.

[0058] The term "pre-lyophilization formulation" or "lyophilization feedstock" as used herein denotes a liquid formulation before the solvent (*e.g.*, water) is removed by a

- 12 -

freeze-drying method. The "fill volume" of a pre-lyophilization formulation is the total volume of the liquid formulation prior to lyophilization.

[0059] "T_{1/2β}," or "T_{1/2 beta}" or "Beta HL," as used herein, is half-life associated with elimination phase, t_{1/2β}=(ln2)/elimination rate constant associated with the terminal phase.

The T_{1/2 beta} can be measured by FIX activity or by FIX antigen level in plasma. The T_{1/2 beta} based on activity is shown as T_{1/2 beta} (activity), and the T_{1/2 beta} based on the FIX antigen level can be shown as T_{1/2 beta} (antigen). Both T_{1/2 beta} (activity) and T_{1/2 beta} (antigen) can be shown as ranges or a geometric mean.

[0060] The term "reconstituted formulation" or "post-reconstitution composition" as used herein denotes a formulation which is lyophilized and re-dissolved by addition of a diluent. The diluent can contain, without limitation, water for injection (WFI), bacteriostatic water for injection (BWFI), sodium chloride solutions (e.g., 0.9% (w/v) NaCl), glucose solutions (e.g., 5% glucose), surfactant containing solutions (e.g., 0.01% polysorbate 20 or polysorbate 80), a pH-buffered solution (e.g. phosphate-buffered solutions) and combinations thereof.

The lyophilization process in general

[0061] Lyophilization, or freeze-drying, is a process widely used in the pharmaceutical industry for the preservation of biological and pharmaceutical materials. The lyophilization process, also known as the lyophilization cycle, is traditionally divided into three distinct stages: freezing, primary drying, and secondary drying. "Lyophilizing," as used herein, refers to the entire process of lyophilization, including both the freezing steps and the drying steps.

[0062] In lyophilization, water present in a material is converted to ice during a freezing step and then removed from the material by direct sublimation under low-pressure conditions during a primary drying step. During freezing, however, not all of the water is transformed to ice. Some portion of the water is trapped in a matrix of solids containing, for example, formulation components and/or the active ingredient. The excess bound water within the matrix can be reduced to a desired level of residual moisture during a secondary drying step. All lyophilization steps, freezing, primary drying and secondary drying, are determinative of the final product properties. The primary drying is typically the longest step in a lyophilization process, therefore, optimization of this portion of the process has significant economic effect.

- 13 -

[0063] In certain aspects of the invention, the lyophilization process only comprises the primary drying step.

[0064] In certain aspects of the invention, the lyophilization process also comprises a separate "vacuum step" between the freezing step and the primary drying step.

[0065] In other aspects of the invention, the lyophilization process further comprises an "annealing step" between the freezing step and the primary drying step.

[0066] The term "annealing step" as used herein refers to a step in the lyophilization process of a polypeptide preparation undergoing lyophilization, prior to the drying step of the preparation, in which the temperature of the preparation is raised from a lower temperature to a higher temperature and then cooled again after a period of time.

[0067] Cycle and formulation optimization has traditionally been performed to assure that the product temperature during primary drying would never exceed the collapse temperature. The term "collapse temperature" as used herein refers to the product temperature during freeze-drying above which product cake begins to lose its original structure. Above the collapse temperature, product could experience slow sporadic bubbling, swelling, foaming, cavitation, fenestration, gross collapse, retraction and beading that may have consequences on the appearance of the product. As a result, Collapse may result in poor product stability, long drying times, uneven drying and loss of texture. *See, e.g.*, US 2010/0041870.

[0068] Lyophilized product in accordance with the present invention can be assessed based on product quality analysis, reconstitution time, quality of reconstitution, high molecular weight, moisture, glass transition temperature (T_g), and biological or biochemical activity. Typically, product quality analysis includes product degradation rate analysis using methods including, but not limited to, size exclusion chromatography (SEC), cation exchange-HPLC (CEX-HPLC), X-ray diffraction (XRD), modulated differential scanning calorimetry (mDSC), reversed phase HPLC (RP-HPLC), multi-angle light scattering detector (MALS), fluorescence, ultraviolet absorption, nephelometry, capillary electrophoresis (CE), SDS-PAGE, and combinations thereof. In some embodiments, evaluation of lyophilized product in accordance with the present invention includes a step of evaluating cake appearance. Additionally, lyophilized product may be assessed based on biological or biochemical activities of the product, typically, after reconstitution.

Lyophilized Factor IX Formulations

[0069] This disclosure provides pre-lyophilization, lyophilized, and post-reconstitution formulations, or pharmaceutical compositions, comprising a FIX polypeptide.

[0070] In certain aspects of the invention, the formulations disclosed herein comprise a FIX polypeptide, a buffering agent, a stabilizing agent, a bulking agent, and a surfactant, or any combinations thereof. The formulation can also contain any other agents that are useful for pharmaceutical formulation.

Factor IX (FIX) polypeptide

[0071] The FIX polypeptide or FIX protein useful for the formulation is a functional Factor FIX protein in its normal role in coagulation, unless otherwise specified. Thus, the FIX polypeptide includes variant polypeptides that are functional and the polynucleotides that encode such functional variant polypeptides. In one embodiment, the FIX polypeptides are the human, bovine, porcine, canine, feline, and murine FIX polypeptides. The full length polypeptide and polynucleotide sequences of FIX are known, as are many functional variants, *e.g.*, fragments, mutants and modified versions. FIX polypeptides include full-length FIX, full-length FIX minus Met at the N-terminus, full-length FIX minus the signal sequence, mature FIX (minus the signal sequence and propeptide), and mature FIX with an additional Met at the N-terminus. FIX can be made by recombinant means ("recombinant Factor IX" or "rFIX"), *i.e.*, it is not naturally occurring or derived from plasma.

[0072] A great many functional FIX variants are known. International publication number WO 02/040544 A3, which is herein incorporated by reference in its entirety, discloses mutants that exhibit increased resistance to inhibition by heparin at page 4, lines 9-30 and page 15, lines 6-31. International publication number WO 03/020764 A2, which is herein incorporated by reference in its entirety, discloses FIX mutants with reduced T cell immunogenicity in Tables 2 and 3 (on pages 14-24), and at page 12, lines 1-27. International publication number WO 2007/149406 A2, which is herein incorporated by reference in its entirety, discloses functional mutant FIX molecules that exhibit increased protein stability, increased *in vivo* and *in vitro* half-life, and increased resistance to proteases at page 4, line 1 to page 19, line 11. WO 2007/149406 A2 also discloses chimeric and other variant FIX molecules at page 19, line 12 to page 20, line 9.

International publication number WO 08/118507 A2, which is herein incorporated by reference in its entirety, discloses FIX mutants that exhibit increased clotting activity at page 5, line 14 to page 6, line 5. International publication number WO 09/051717 A2, which is herein incorporated by reference in its entirety, discloses FIX mutants having an increased number of N-linked and/or O-linked glycosylation sites, which results in an increased half-life and/or recovery at page 9, line 11 to page 20, line 2. International publication number WO 09/137254 A2, which is herein incorporated by reference in its entirety, also discloses Factor IX mutants with increased numbers of glycosylation sites at page 2, paragraph [006] to page 5, paragraph [011] and page 16, paragraph [044] to page 24, paragraph [057]. International publication number WO 09/130198 A2, which is herein incorporated by reference in its entirety, discloses functional mutant FIX molecules that have an increased number of glycosylation sites, which result in an increased half-life, at page 4, line 26 to page 12, line 6. International publication number WO 09/140015 A2, which is herein incorporated by reference in its entirety, discloses functional FIX mutants that have an increased number of Cys residues, which can be used for polymer (*e.g.*, PEG) conjugation, at page 11, paragraph [0043] to page 13, paragraph [0053]. The FIX polypeptides described in International Application No. PCT/US2011/043569 filed July 11, 2011 and published as WO 2012/006624 on January 12, 2012 are also incorporated herein by reference in its entirety.

[0073] In certain embodiments, the FIX polypeptide comprises wild-type FIX. In some embodiments, the FIX polypeptide further comprises a heterologous moiety fused to wild-type FIX. In certain embodiments, the heterologous moiety is a moiety extending half-life of FIX. In certain embodiments, the heterologous moiety comprises a polypeptide or a non-polypeptide moiety.

[0074] In other embodiments, the FIX polypeptide is a long-acting FIX polypeptide. A long-acting FIX polypeptide can comprise a FIX portion and a non FIX portion, *e.g.*, a heterologous moiety that is capable of extending *in vivo* or *in vitro* half-life of the FIX polypeptide. Exemplary non- FIX portions include, *e.g.*, Fc, albumin, a PAS sequence, transferrin, CTP (28 amino acid C-terminal peptide (CTP) of human chorionic gonadotropin (hCG) with its 4 O-glycans), polyethylene glycol (PEG), hydroxyethyl starch (HES), albumin binding polypeptide, albumin-binding small molecules, or any combination thereof. Exemplary long-acting FIX polypeptides of the invention include,

e.g., Factor IX-Fc polypeptides, Factor IX-albumin polypeptides, Factor IX-PAS polypeptides, Factor IX -transferrin polypeptides, Factor IX -CTP polypeptides, Factor IX -PEG polypeptides, Factor IX -HES polypeptides, Factor IX -albumin binding polypeptide polypeptides, or Factor IX -albumin-binding small molecule polypeptides.

[0075] In one embodiment, the FIX polypeptide is rFIXFc, a recombinant fusion protein comprised of human coagulation Factor IX (FIX) and an Fc domain of a human antibody (IgG1 isotype). *See, e.g.*, PCT Application No. PCT/US2011/043569, filed July 11, 2011 and published as WO 2012/006624, which is incorporated herein by reference in its entirety. The rFIXFc polypeptide is a heterodimeric protein with a FIXFc single chain (FIXFc-sc) and an Fc single chain (Fc-sc) bound together through two disulfide bonds in the hinge region of Fc. rFIXFc requires two protein subunits, FIXFc-sc (642 amino acids, SEQ ID NO:2) and Fc-sc (227 amino acids, SEQ ID NO:4), to assemble within a transfected cell line to form the final protein product, rFIXFc. The polynucleotide sequences encoding FIXFc-sc and Fc-sc are presented as SEQ ID NO:1 and SEQ ID NO:3, respectively.

[0076] In certain embodiments, the Factor IX portion of rFIXFc has a primary amino acid sequence that is identical to the Thr148 allelic form of plasma derived Factor IX and has structural and functional characteristics similar to endogenous Factor IX. The Fc domain of rFIXFc contains the hinge, CH2 and CH3 regions of IgG1. The assembled heterodimer mature form of rFIXFc contains 869 amino acids with a molecular weight of approximately 98 kilodaltons. In some embodiments, the rFIXFc polypeptide comprises an amino acid sequence at least 90%, 95%, or 100% identical to amino acids 1 to 642 of SEQ ID NO:2.

[0077] In one embodiment, the second portion fused to FIX is an FcRn binding partner. In another embodiment, an FcRn binding partner fused to FIX is an Fc fragment. An FcRn binding partner is any molecule that can be specifically bound by the FcRn receptor with consequent active transport by the FcRn receptor of the FcRn binding partner. Thus, the term Fc includes any variants of IgG Fc that are functional. The region of the Fc portion of IgG that binds to the FcRn receptor has been described based on X-ray crystallography (Burmeister *et al.*, *Nature* 372:379 (1994), incorporated herein by reference in its entirety). The major contact area of the Fc with the FcRn is near the junction of the CH2 and CH3 domains. Fc-FcRn contacts are all within a single Ig heavy

chain. The FcRn binding partners include, *e.g.*, whole IgG, the Fc fragment of IgG, and other fragments of IgG that include the complete binding region of FcRn. The major contact sites include amino acid residues 248, 250-257, 272, 285, 288, 290-291, 308-311, and 314 of the CH2 domain and amino acid residues 385-387, 428, and 433-436 of the CH3 domain. References made to amino acid numbering of immunoglobulins or immunoglobulin fragments, or regions, are all based on Kabat *et al.* 1991, Sequences of Proteins of Immunological Interest, U. S. Department of Public Health, Bethesda; MD, incorporated herein by reference in its entirety. (The FcRn receptor has been isolated from several mammalian species including humans. The sequences of the human FcRn, rat FcRn, and mouse FcRn are known (Story *et al.*, *J. Exp. Med.* 180: 2377 (1994), incorporated herein by reference in its entirety.) An Fc can comprise the CH2 and CH3 domains of an immunoglobulin with or without the hinge region of the immunoglobulin. Exemplary Fc variants are provided in WO 2004/101740 and WO 2006/074199, incorporated herein by reference in its entirety.

[0078] Fc (or Fc portion of a chimeric polypeptide) can contain one or more mutations, and combinations of mutations.

[0079] Fc (or Fc portion of a chimeric polypeptide) can contain mutations conferring increased half-life such as M252Y, S254T, T256E, and combinations thereof, as disclosed in Oganesyan *et al.*, *Mol. Immunol.* 46:1750 (2009), which is incorporated herein by reference in its entirety; H433K, N434F, and combinations thereof, as disclosed in Vaccaro *et al.*, *Nat. Biotechnol.* 23:1283 (2005), which is incorporated herein by reference in its entirety; the mutants disclosed at pages 1-2, paragraph [0012], and Examples 9 and 10 of US 2009/0264627 A1, which is incorporated herein by reference in its entirety; and the mutants disclosed at page 2, paragraphs [0014] to [0021] of US 20090163699 A1, which is incorporated herein by reference in its entirety.

[0080] Fc (or Fc portion of a chimeric polypeptide) can also include, *e.g.*, the following mutations: The Fc region of IgG can be modified according to well recognized procedures such as site directed mutagenesis and the like to yield modified IgG or Fc fragments or portions thereof that will be bound by FcRn. Such modifications include, *e.g.*, modifications remote from the FcRn contact sites as well as modifications within the contact sites that preserve or even enhance binding to the FcRn. For example the following single amino acid residues in human IgG1 Fc (Fcγ1) can be substituted without

significant loss of Fc binding affinity for FcRn: P238A, S239A, K246A, K248A, D249A, M252A, T256A, E258A, T260A, D265A, S267A, H268A, E269A, D270A, E272A, L274A, N276A, Y278A, D280A, V282A, E283A, H285A, N286A, T289A, K290A, R292A, E293A, E294A, Q295A, Y296F, N297A, S298A, Y300F, R301A, V303A, V305A, T307A, L309A, Q311A, D312A, N315A, K317A, E318A, K320A, K322A, S324A, K326A, A327Q, P329A, A330Q, A330S, P331A, P331S, E333A, K334A, T335A, S337A, K338A, K340A, Q342A, R344A, E345A, Q347A, R355A, E356A, M358A, T359A, K360A, N361A, Q362A, Y373A, S375A, D376A, A378Q, E380A, E382A, S383A, N384A, Q386A, E388A, N389A, N390A, Y391F, K392A, L398A, S400A, D401A, D413A, K414A, R416A, Q418A, Q419A, N421A, V422A, S424A, E430A, N434A, T437A, Q438A, K439A, S440A, S444A, and K447A, where for example P238A represents wildtype proline substituted by alanine at position number 238. In addition to alanine other amino acids can be substituted for the wildtype amino acids at the positions specified above. Mutations can be introduced singly into Fc giving rise to more than one hundred FcRn binding partners distinct from native Fc. Additionally, combinations of two, three, or more of these individual mutations can be introduced together, giving rise to hundreds more FcRn binding partners. Certain of these mutations can confer new functionality upon the FcRn binding partner. For example, one embodiment incorporates N297A, removing a highly conserved N-glycosylation site. The effect of this mutation is to reduce immunogenicity, thereby enhancing circulating half-life of the FcRn binding partner, and to render the FcRn binding partner incapable of binding to FcyRI, FcyRIIA, FcyRIIB, and FcyRIIIA, without compromising affinity for FcRn (Routledge et al. 1995, Transplantation 60:847, which is incorporated herein by reference in its entirety; Friend et al. 1999, Transplantation 68:1632, which is incorporated herein by reference in its entirety; Shields et al. 1995, J. Biol. Chem. 276:6591, which is incorporated herein by reference in its entirety). Additionally, at least three human Fc gamma receptors appear to recognize a binding site on IgG within the lower hinge region, generally amino acids 234-237. Therefore, another example of new functionality and potential decreased immunogenicity can arise from mutations of this region, as for example by replacing amino acids 233-236 of human IgG1 "ELLG" to the corresponding sequence from IgG2 "PVA" (with one amino acid deletion). It has been shown that FcyRI, FcyRII, and FcyRIII which mediate various effector functions will not

bind to IgG1 when such mutations have been introduced (Ward and Ghetie, *Therapeutic Immunology* 2:77 (1995), which is incorporated herein by reference in its entirety; and Armour *et al.*, *Eur. J. Immunol.* 29:2613 (1999), which is incorporated herein by reference in its entirety). As a further example of new functionality arising from mutations described above affinity for FcRn can be increased beyond that of wild type in some instances. This increased affinity can reflect an increased "on" rate, a decreased "off" rate or both an increased "on" rate and a decreased "off" rate. Mutations believed to impart an increased affinity for FcRn include, *e.g.*, T256A, T307A, E380A, and N434A (Shields *et al.*, *J. Biol. Chem.* 276:6591 (2001), which is incorporated herein by reference in its entirety).

[0081] The Fc (or Fc portion of a chimeric polypeptide) can be at least about 60%, about 70%, about 80%, about 90%, about 95%, about 96%, about 97%, about 98%, about 99%, or about 100% identical to the Fc amino acid sequence shown in Table 14 (*e.g.*, amino acids 21 to 247 of SEQ ID NO: 4). The Fc (or Fc portion of a chimeric polypeptide) can be identical to the Fc amino acid sequence shown in Table 14 (*e.g.*, amino acids 21 to 247 of SEQ ID NO: 4).

[0082] As discussed above, exemplary long-acting polypeptides also include FIX fused to one or more albumin polypeptides, albumin binding polypeptides, or albumin-binding small molecules. In one embodiment, the albumin is human albumin. The albumin or albumin binding protein can be fused to either the N-terminal end of FIX or to the C-terminal end of FIX or inserted between two amino acids in FIX. Examples of albumin, *e.g.*, fragments thereof, that can be used in the present invention are known. *e.g.*, U.S. Patent No. 7,592,010; U.S. Patent No. 6,686,179; and Schulte, *Thrombosis Res.* 124 Suppl. 2:S6-S8 (2009), each of which is incorporated herein by reference in its entirety.

[0083] The albumin binding polypeptides can compromise, without limitation, bacterial albumin-binding domains, albumin-binding peptides, or albumin-binding antibody fragments that can bind to albumin. Domain 3 from streptococcal protein G, as disclosed by Kraulis *et al.*, *FEBS Lett.* 378:190-194 (1996) and Linhult *et al.*, *Protein Sci.* 11:206-213 (2002) is an example of a bacterial albumin-binding domain. Examples of albumin-binding peptides include a series of peptides having the core sequence DICLPRWGCLW (SEQ ID NO: 5). See, *e.g.*, Dennis *et al.*, *J. Biol. Chem.* 2002, 277: 35035-35043 (2002). Examples of albumin-binding antibody fragments are disclosed in Muller and

Kontermann, *Curr. Opin. Mol. Ther.* 9:319-326 (2007); Rooverset *et al.*, *Cancer Immunol. Immunother.* 56:303-317 (2007), and Holt *et al.*, *Prot. Eng. Design Sci.*, 21:283-288 (2008), which are incorporated herein by reference in their entireties.

[0084] In certain aspects, a recombinant FIX polypeptide of the invention comprises at least one attachment site for a non-polypeptide small molecule, variant, or derivative that can bind to albumin thereof. An example of such albumin binding moieties is 2-(3-maleimidopropanamido)-6-(4-(4-iodophenyl)butanamido)hexanoate (“Albu” tag) as disclosed by Trusselet *et al.*, *Bioconjugate Chem.* 20:2286-2292 (2009).

[0085] As discussed above, exemplary long-acting polypeptides also include FIX fused to at least one C-terminal peptide (CTP) of the β subunit of human chorionic gonadotropin or fragment, variant, or derivative thereof. The CTP can be fused to FIX either the N-terminal end of FIX or to the C-terminal end of FIX. One or more CTP peptides fused to or inserted into a recombinant protein is known to increase the *in vivo* half-life of that protein. *See, e.g.*, U.S. Patent No. 5,712,122, incorporated by reference herein in its entirety.

Exemplary CTP peptides include

DPRFQDSSSKAPPPSLPSPSRLPGPSDTPIL (SEQ ID NO: 6) or
SSSSKAPPPSLPSPSRLPGPSDTPILPQ. (SEQ ID NO: 7). *See, e.g.*, U.S. Patent Application Publication No. US 2009/0087411 A1, incorporated by reference.

[0086] As discussed above, exemplary long-acting polypeptides also include FIX fused to at least one PAS sequence or fragment, variant, or derivative thereof. The PAS sequence can be fused to either the N-terminal end of FIX or to the C-terminal end of FIX. A PAS peptide or PAS sequence, as used herein, means an amino acid sequence comprising mainly alanine and serine residues or comprising mainly alanine, serine, and proline residues, the amino acid sequence forming random coil conformation under physiological conditions. Accordingly, the PAS sequence is a building block, an amino acid polymer, or a sequence cassette comprising, consisting essentially of, or consisting of alanine, serine, and proline which can be used as a part of the heterologous moiety in the chimeric protein. An amino acid polymer also can form random coil conformation when residues other than alanine, serine, and proline are added as a minor constituent in the PAS sequence. By “minor constituent” is meant that that amino acids other than alanine, serine, and proline can be added in the PAS sequence to a certain degree, *e.g.*, up to about 12%, *i.e.*, about 12 of 100 amino acids of the PAS sequence, up to about 10%, up to about

9%, up to about 8%, about 6%, about 5%, about 4%, about 3%, *i.e.* about 2%, or about 1%, of the amino acids. The amino acids different from alanine, serine and proline can be selected from the group consisting of Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Thr, Trp, Tyr, and Val. Under physiological conditions, a PAS peptide forms a random coil conformation and thereby can mediate an increased *in vivo* and/or *in vitro* stability to a recombinant protein of the invention, and has procoagulant activity.

[0087] Non-limiting examples of the PAS peptides include ASPAAPAPAPASPAAPAPSAPA (SEQ ID NO: 8), AAPASPAPAAAPSAPAPAAPS (SEQ ID NO: 9), APSSPSPSAPSSPSPASPSS (SEQ ID NO: 10), APSSPSPSAPSSPSPASPSP (SEQ ID NO: 11), SSPSAPSPSSPASPSPSSPA (SEQ ID NO: 12), AASPAAPSAPPAAASPAAPSAPPA (SEQ ID NO: 13), ASAAAPAAASAAASAPSAAA (SEQ ID NO: 14) or any variants, derivatives, fragments, or combinations thereof. Additional examples of PAS sequences are known from, *e.g.*, US Pat. Publ. No. 2010/0292130 A1, PCT Appl. Publ. No. WO 2008/155134 A1, and European issued patent EP2173890.

[0088] As discussed above, exemplary long-acting polypeptides also include FIX fused to at least one transferrin peptide or fragment, variant, or derivative thereof. At least one transferrin peptide can be fused to either the N-terminal end of FIX or to the C-terminal end of FIX or inserted between two amino acids in FIX. Any transferrin can be fused to or inserted into a recombinant FIX protein of the invention. As an example, wild-type human Tf (Tf) is a 679 amino acid protein, of approximately 75 KDa (not accounting for glycosylation), with two main domains, N (about 330 amino acids) and C (about 340 amino acids), which appear to originate from a gene duplication. *See* GenBank accession numbers NM001063, XM002793, M12530, XM039845, XM 039847 and S95936 (www.ncbi.nlm.nih.gov), all of which are herein incorporated by reference in their entirety.

[0089] Transferrin transports iron through transferrin receptor (TfR)-mediated endocytosis. After the iron is released into an endosomal compartment and Tf-TfR complex is recycled to cell surface, the Tf is released back extracellular space for next cycle of iron transporting. Tf possesses a long half-life that is in excess of 14-17 days (Li *et al.*, *Trends Pharmacol. Sci.* 23:206-209 (2002)). Transferrin fusion proteins have been studied for half-life extension, targeted delivery for cancer therapies, oral delivery and

sustained activation of proinsulin (Brandsma *et al.*, *Biotechnol. Adv.*, 29: 230-238 (2011); Bai *et al.*, *Proc. Natl. Acad. Sci. USA* 102:7292-7296 (2005); Kim *et al.*, *J. Pharmacol. Exp. Ther.*, 334:682-692 (2010); Wang *et al.*, *J. Controlled Release* 155:386-392 (2011)).

[0090] As discussed above, exemplary long-acting polypeptides also include FIX fused to at least one polyethylene glycol (PEG) moieties.

[0091] PEGylated FIX can refer to a conjugate formed between FIX and at least one polyethylene glycol (PEG) molecule. PEG is commercially available in a large variety of molecular weights and average molecular weight ranges. Typical examples of PEG average molecular weight ranges include, but are not limited to, about 200, about 300, about 400, about 600, about 1000, about 1300-1600, about 1450, about 2000, about 3000, about 3000-3750, about 3350, about 3000-7000, about 3500-4500, about 5000-7000, about 7000-9000, about 8000, about 10000, about 8500-11500, about 16000-24000, about 35000, about 40000, about 60000, and about 80000 daltons. These average molecular weights are provided merely as examples and are not meant to be limiting in any way.

[0092] A recombinant long-acting FIX protein of the invention can be PEGylated to include mono- or poly-(e.g., 2-4) PEG moieties. PEGylation can be carried out by any of the PEGylation reactions known in the art. Methods for preparing a PEGylated protein product will generally include (i) reacting a polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby the peptide of the invention becomes attached to one or more PEG groups; and (ii) obtaining the reaction product(s). In general, the optimal reaction conditions for the reactions will be determined case by case based on known parameters and the desired result.

[0093] There are a number of PEG attachment methods available to those skilled in the art, for example Malik F *et al.*, *Exp. Hematol.* 20:1028-35 (1992); Francis, *Focus on Growth Factors* 3(2):4-10 (1992); European Pat. Pub. Nos. EP0401384, EP0154316, and EP0401384; and International Pat. Appl. Pub. Nos. WO92/16221 and WO95/34326. As a non-limiting example, FIX variants can contain cysteine substitutions in one or more insertion sites in FIX, and the cysteines can be further conjugated to PEG polymer. See Mei *et al.*, *Blood* 116:270-279 (2010) and U.S. Patent No. 7,632,921, which are incorporated herein by reference in their entireties.

[0094] As discussed above, exemplary long-acting polypeptides also include FIX fused to at least one hydroxyethyl starch (HES) polymer. HES is a derivative of naturally

occurring amylopectin and is degraded by alpha-amylase in the body. HES exhibits advantageous biological properties and is used as a blood volume replacement agent and in hemodilution therapy in the clinics. See, e.g., Sommermeyer *et al.*, *Krankenhauspharmazie* 8:271-278 (1987); and Weidler *et al.*, *Arzneim.-Forschung/Drug Res.* 41: 494-498 (1991).

[0095] HES is mainly characterized by the molecular weight distribution and the degree of substitution. HES has a mean molecular weight (weight mean) of from 1 to 300 kD, from 2 to 200kD, from 3 to 100 kD, or from 4 to 70kD. Hydroxyethyl starch can further exhibit a molar degree of substitution of from 0.1 to 3, from 0.1 to 2, from 0.1 to 0.9, or from 0.1 to 0.8, and a ratio between C2:C6 substitution in the range of from 2 to 20 with respect to the hydroxyethyl groups. HES with a mean molecular weight of about 130 kD is VOLUVEN® from Fresenius. VOLUVEN® is an artificial colloid, employed, e.g., for volume replacement used in the therapeutic indication for therapy and prophylaxis of hypovolaemia. There are a number of HES attachment methods available to those skilled in the art, e.g., the same PEG attachment methods described above.

[0096] Factor IX coagulant activity is expresses as International Unit(s) (IU). One IU of Factor IX activity corresponds approximately to the quantity of Factor IX in one milliliter of normal human plasma. Several assays are available for measuring Factor IX activity, including the one stage clotting assay (activated partial thromboplastin time; aPTT), thrombin generation time (TGA) and rotational thromboelastometry (ROTEM®).

Buffering Agent

[0097] Buffering agents useful for the present invention can be a weak acid or base used to maintain the acidity (pH) of a solution near a chosen value after the addition of another acid or base. Suitable buffering agents can maximize the stability of the pharmaceutical formulations by maintaining pH control of the formulation. Suitable buffering agents can also ensure physiological compatibility or optimize solubility. Rheology, viscosity and other properties can also dependent on the pH of the formulation. Common buffering agents include, but are not limited to, histidine, citrate, succinate, acetate and phosphate. In some embodiments, a buffering agent comprises L-histidine or mixtures of L-histidine with L-histidine hydrochloride with isotonicity agents and potentially pH adjustment with an acid or a base known in the art. In certain embodiments, the buffering agent is L-

- 24 -

histidine. In certain embodiments, the pH of the formulation is maintained between about 6 and about 8, or between about 6.5 and about 7.5.

Stabilizing Agent

[0098] Stabilizing agents are added to a pharmaceutical product in order to stabilize that product. Such agents can stabilize proteins in a number of different ways. Common stabilizing agents include, but are not limited to, amino acids such as glycine, alanine, lysine, arginine, or threonine, carbohydrates such as glucose, sucrose, trehalose, raffinose, or maltose, polyols such as glycerol, mannitol, sorbitol, cyclodextrins or destrans of any kind and molecular weight, or PEG. In one aspect of the invention, the stabilizing agent is chosen in order to maximize the stability of FIX polypeptide in lyophilized preparations. In certain embodiments, the stabilizing agent is sucrose.

Bulking Agent

[0099] Bulking agents can be added to a pharmaceutical product in order to add volume and mass to the product, thereby facilitating precise metering and handling thereof. Common bulking agents include, but are not limited to, lactose, sucrose, glucose, mannitol, sorbitol, calcium carbonate, or magnesium stearate. In certain embodiments, the bulking agent is mannitol.

Surfactant

[0100] Surfactants are amphipathic substances with lyophilic and lyophobic groups. A surfactant can be anionic, cationic, zwitterionic, or nonionic. Examples of nonionic surfactants include, but are not limited to, alkyl ethoxylate, nonylphenol ethoxylate, amine ethoxylate, polyethylene oxide, polypropylene oxide, fatty alcohols such as cetyl alcohol or oleyl alcohol, cocamide MEA, cocamide DEA, polysorbates, or dodecyl dimethylamine oxide. In certain embodiments, the surfactant is polysorbate 20 or polysorbate 80.

Pre-Lyophilization Formulation

[0101] In one aspect, the disclosure provides a pre-lyophilization formulation comprising:

- (a) a Factor IX (FIX) polypeptide having FIX coagulation activity;

- 25 -

- (b) a buffering agent;
- (c) a stabilizing agent;
- (d) a bulking agent; and
- (e) a surfactant,

wherein the formulation has a fill volume of less than about 5 mL, less than about 4mL, or less than about 3 mL and wherein each of (a) – (e) are at an amount per vial (mg/vial) sufficient to allow

- (1) improved stability of the FIX polypeptide when lyophilized;
- (2) reduced reconstitution time when lyophilized;
- (3) reduced splashing onto a stopper comprising the formulation;
- (4) reduced lyophilization cycle time;
- (5) increased shelf-life of a lyophilate prepared from the pre-lyophilization formulation at room temperature; or
- (6) any combinations thereof,

compared to a reference pre-lyophilization formulation,

and wherein the reference formulation comprises (a) – (e) at the amount per vial identical to the pre-lyophilization formulation, but has at least 5mL fill volume. In some embodiments, the reference formulation has a 5.3mL fill volume or a 5mL fill volume.

[0102] In other embodiments, the pre-lyophilization formulation allows at least two, at least three, at least four, or at least five properties selected from (1) improved stability of the FIX polypeptide when lyophilized; (2) reduced reconstitution time when lyophilized; (3) reduced splashing onto a stopper comprising the formulation; (4) reduced lyophilization cycle time; and (5) increased shelf-life of a lyophilate prepared from the pre-lyophilization formulation at room temperature. In certain embodiments, the pre-lyophilization formulation allows (1) improved stability of the FIX polypeptide when lyophilized. In certain embodiments, the pre-lyophilization formulation allows (2) reduced reconstitution time when lyophilized. In certain embodiments, the pre-lyophilization formulation allows (3) reduced splashing onto a stopper comprising the formulation. In certain embodiments, the pre-lyophilization formulation allows (4) reduced lyophilization cycle time. In certain embodiments, the pre-lyophilization formulation allows (5) increased shelf-life of a lyophilate prepared from the pre-

lyophilization formulation at room temperature. In certain embodiments, the pre-lyophilization formulation allows (6) any combinations of properties described herein.

[0103] In certain embodiments, the pre-lyophilization formulation comprises at least about 100 IU/vial of the FIX polypeptide. In certain embodiments, the pre-lyophilization formulation comprises at least about 200 IU/vial to about 10,000 IU/vial of the FIX polypeptide, about 200IU/vial to about 6,000 IU/vial, or about 500IU/vial to about 5,000IU/vial. In certain embodiments, the pre-lyophilization formulation comprises about 220 IU/vial, about 250 IU/vial, about 300 IU/vial, about 400 IU/vial, about 500 IU/vial, about 600 IU/vial, about 700 IU/vial, about 800 IU/vial, about 900 IU/vial, about 1,000 IU/vial, about 1,100 IU/vial, about 1,200 IU/vial, about 1,300 IU/vial, about 1,400 IU/vial, about 1,500 IU/vial, about 2,000 IU/vial, about 2,500 IU/vial, about 3,000 IU/vial, about 4,000 IU/vial, about 5,000 IU/vial, about 5,500 IU/vial, about 6,000 IU/vial, about 6,500 IU/vial, about 7,000 IU/vial, about 7,500 IU/vial, about 8,000 IU/vial, about 8,500 IU/vial, about 9,000 IU/vial, about 9,500 IU/vial or about 10,000 IU/vial of the FIX polypeptide.

[0104] In some embodiments, a higher concentration of the pre-lyophilization formulation is achieved by reducing the fill volume. In certain embodiments, the pre-lyophilization formulation has the fill volume of about 4.0 mL, about 3.5 mL, about 3.0 mL, about 2.9 mL, about 2.8 mL, about 2.7 mL, about 2.65 mL, about 2.6mL, about 2.5 mL, about 2.4 mL, about 2.3 mL, about 2.2 mL, about 2.1 mL, or about 2.0 mL. In some embodiments, the fill volume of the pre-lyophilization formulation is about 2.65 mL. In some embodiments, the fill volume of the pre-lyophilization formulation is less than about 5 mL.

[0105] In some embodiments, the FIX polypeptide can be further concentrated by an additional purification step, *e.g.*, a second ultra filtration step.

[0106] In certain embodiments, the reduced reconstitution time is less than 1.5 minute, less than 1 minute, less than 50 seconds, less than 40 seconds, less than 30 seconds, less than 20 seconds, or less than 10 seconds. In specific embodiments, the reduced reconstitution time is less than 30 seconds.

[0107] In certain embodiments, the reduced lyophilization cycle time of the pre-lyophilization formulation is about 4 days or less, about 3 days or less, about 2 days or less, or about a day or less.

[0108] In certain embodiments, the concentration of the buffering agent in the pre-lyophilization formulation is between about 3 mg/mL and about 15 mg/mL, between 4 mg/mL and between 12 mg/mL, between about 5 mg/mL and about 10 mg/mL, or between about 5.82 mg/mL and about 9.7 mg/mL. In one embodiment, the buffering agent is at a concentration of between about 3.88 mg/mL and about 9.7 mg/mL. In one embodiment, the buffering agent is at a concentration of about 7.76 mg/mL. In some embodiments, the pre-lyophilization formulation contains L-histidine at a concentration of about 7.76 mg/mL.

[0109] In certain embodiments, the concentration of the stabilizing agent in the pre-lyophilization formulation is between 10 mg/mL and about 50 mg/mL, between about 13 mg/mL and about 40 mg/mL, between about 15 mg/mL and about 35 mg/mL, or between about 17.85 mg/mL and about 29.95 mg/mL. In one embodiment, the buffering agent is at a concentration of about 23.8 mg/mL. In some embodiments, the pre-lyophilization formulation contains sucrose at a concentration of 23.8 mg/mL.

[0110] In certain embodiments, the concentration of the bulking agent in the pre-lyophilization formulation is between about 20 mg/mL and about 100 mg/mL, between about 30 mg/mL and about 70 mg/mL, between about 30 mg/mL and about 60 mg/mL, or between about 35.7 mg/mL and about 59.5 mg/mL. In one embodiment, the bulking agent is at a concentration of about 47.6 mg/mL. In some embodiments, the pre-lyophilization formulation contains mannitol at a concentration of about 47.6 mg/mL.

[0111] In certain embodiments, the concentration of the surfactant in the pre-lyophilization formulation is between about 0.01 mg/mL and about 5 mg/mL, between about 0.1 mg/mL and about 4 mg/mL, between about 0.1 mg/mL and about 3 mg/mL, between about 0.01 mg/mL and about 2 mg/mL, or between about 0.05 mg/mL and about 1 mg/mL. In one embodiment, the surfactant is at a concentration of about 0.2 mg/mL. In some embodiments, the pre-lyophilization formulation contains polysorbate 20 or polysorbate 80 at a concentration of about 0.2 mg/mL.

[0112] In certain embodiments, the concentration of the FIX polypeptide in the pre-lyophilization formulation is between about 80 IU/mL and about 2,750 IU/mL. In some embodiments, the concentration of the FIX polypeptide in the pre-lyophilization formulation is at least about 100 IU/mL, at least about 200 IU/mL, at least about 300 IU/mL, at least about 400 IU/mL, at least about 500 IU/mL, at least about 600 IU/mL, at

least about 700 IU/mL, at least about 800 IU/mL, at least about 900 IU/mL, at least about 1000 IU/mL, at least about 1500 IU/mL, at least about 2000 IU/mL, or at least about 2500 IU/mL.

[0113] In one aspect, the disclosure further provides a pre-lyophilization formulation comprising:

- (a) about 80 to about 2,750 IU/mL of rFIXFc;
- (b) about 7.76 mg/mL of L-histidine.
- (c) about 47.6 mg/mL of mannitol;
- (d) about 23.8 mg/mL of sucrose; and,
- (e) about 0.2 mg/mL of polysorbate-20.

[0114] In certain embodiments, the fill volume of such pre-lyophilization formulation is about 3 mL, about 2.9 mL, about 2.8 mL, about 2.7 mL, about 2.65 mL, about 2.6 mL, about 2.5 mL, about 2.4 mL, about 2.3 mL, about 2.2 mL, about 2.1 mL, or about 2.0 mL. In one embodiment, the fill volume of such pre-lyophilization formulation is about 2.65 mL.

[0115] In addition, this disclosure provides a lyophilate powder which is lyophilized from any of the above pre-lyophilization formulations. In some embodiments, the pre-lyophilization formulation is any formulation described herein.

Lyophilate Power

[0116] The disclosure also provides a lyophilate powder comprising a FIX polypeptide, a buffering agent, a stabilizing agent, a bulking agent, a surfactant, or any combinations thereof.

[0117] In certain embodiments, the lyophilate powder comprises between about 8 mg and about 39 mg per vial, between about 9mg and about 35 mg per vial, between about 10 mg and about 30 mg per vial, between about 12 mg and about 25 mg per vial, between about 15 mg and about 23 mg per vial of the buffering agent (e.g., L-histidine). In one embodiment, the lyophilate powder comprises about 25 mg per vial, about 24 mg per vial, about 23 mg per vial, about 22 mg per vial, about 21 mg per vial, about 20 mg per vial, about 19 mg per vial, about 18 mg per vial, about 17 mg per vial, about 16 mg per vial, about 15 mg per vial of the buffering agent. In another embodiment, the lyophilate powder comprises about 20.6 mg per vial of the buffering agent. In some embodiments, the buffering agent is L-histidine.

[0118] In certain embodiments, the lyophilate powder comprises between about 27 mg and about 132 mg per vial, between about 30 mg and about 120 mg per vial, between about 40 mg and about 110 mg per vial, between about 50 mg and about 100 mg per vial, between about 60 mg and about 90 mg per vial of the stabilizing agent. In one embodiment, the lyophilate powder comprises about 68 mg per vial, about 67 mg per vial about 66 mg per vial about 65 mg per vial about 64 mg per vial about 63 mg per vial about 62 mg per vial about 61 mg per vial about 60 mg per vial about 59 mg per vial of the stabilizing agent. In another embodiment, the lyophilate power comprises about 63.1 mg per vial of the stabilizing agent. In some embodiments, the stabilizing agent is sucrose.

[0119] In certain embodiments, the lyophilate powder comprises between about 50 mg and about 265 mg per vial, between about 53 mg and about 265 mg per vial, between about 50 mg and about 250 mg per vial, between about 53 mg and about 265 mg per vial, between about 80 mg and about 200 mg per vial, between about 100 mg and about 150 mg per vial, or between about 110 mg and about 140 mg per vial of the bulking agent. In one embodiment, the lyophilate powder comprises about 131 mg per vial, about 130 mg per vial, about 129 mg per vial, about 128 mg per vial, about 127 mg per vial, about 126 mg per vial, about 125 mg per vial, about 124 mg per vial, about 123 mg per vial, or about 122 mg per vial of the bulking agent. In another embodiment, the lyophilate powder comprises about 126.1 mg per vial of the bulking agent. In some embodiments, the bulking agent is mannitol.

[0120] In certain embodiments, the lyophilate powder comprises between about 0.03 mg and about 13 mg per vial, between about 0.05 mg and about 10 mg per vial between about 0.07 mg and about 8 mg per vial between about 0.1 mg and about 2 mg per vial of the surfactant. In one embodiment, the lyophilate powder comprises about 1 mg per vial, about 0.9 mg per vial, about 0.8 mg per vial, about 0.7 mg per vial, about 0.6 mg per vial, about 0.5 mg per vial, about 0.4 mg per vial, about 0.3 mg per vial, about 0.2 mg per vial, or about 0.1 mg per vial of the surfactant. In another embodiment, the lyophilate power comprises about 0.5 mg per vial of the surfactant. In one embodiment, the lyophilate powder comprises about 0.53 mg per vial of the surfactant. In some embodiments, the surfactant is polysorbate 20 or polysorbate 80.

[0121] In certain embodiments, the lyophilate powder comprises:

- 30 -

- (a) a FIX polypeptide at an amount between about 2 mg per vial and about 150 mg per vial;
- (b) a buffering agent at an amount between about 10 mg per vial and about 30 mg per vial;
- (c) a bulking agent at an amount between about 70 mg vial and about 200 mg per vial.
- (d) a stabilizing agent at an amount between about 30 mg per vial and 100 mg per vial; and
- (e) a surfactant at an amount between about 0.05 mg per vial and about 5 mg per vial.

[0122] In some embodiments, the lyophilate powder comprises:

- (a) a FIX polypeptide at an amount between about 2.2 mg per vial and about 125 mg per vial;
- (b) a buffering agent at an amount between about 8 mg per vial and about 39 mg per vial;
- (c) a bulking agent at an amount between about 53 mg vial and about 265 mg per vial.
- (d) a stabilizing agent at an amount between about 27 mg per vial and 132 mg per vial; and
- (e) a surfactant at an amount between about 0.03 mg per vial and about 13 mg per vial.

[0123] In certain embodiments, the lyophilate powder comprises:

- (a) the lyophilized FIX polypeptide at an amount between about 2.2 mg per vial and about 125 mg per vial;
- (b) the buffering agent at an amount between about 12.5 mg per vial and 25 mg per vial;
- (c) the stabilizing agent at an amount between about 32.5 mg per vial and 80 mg per vial;
- (d) the bulking agent at an amount between about 75 mg per vial and 150 mg per vial; and
- (e) the surfactant at an amount between about 0.1 mg/mL and about 2 mg/mL.

[0124] In one embodiment, the lyophilate powder comprises:

- 31 -

- (a) about 2.2 to about 125 mg/vial of the FIX polypeptide;
- (b) about 20.6 mg/vial of L-histidine;
- (c) about 126.1 mg/vial of mannitol;
- (d) about 63.1 mg/vial of sucrose; and
- (e) about 0.53 mg/vial of polysorbate 20.

Reconstituted Formulation

[0125] Furthermore, this disclosure provides a reconstituted formulation comprising any of the above lyophilate powder reconstituted by a reconstitution buffer.

[0126] In certain embodiments, the reconstitution buffer is a NaCl solution. In some embodiments, the reconstitution buffer is 5 mL.

[0127] In certain embodiments, the reconstituted formulation comprises:

- (a) the FIX polypeptide at a concentration between about 0.9 mg/mL and about 50 mg/mL;
- (b) the buffering agent at a concentration between 2 mg/mL and about 5 mg/mL;
- (c) the bulking agent at a concentration between 20 mg/mL and about 30 mg/mL;
- (d) the stabilizing agent at a concentration between 8 mg/mL and 15 mg/mL per vial; and
- (e) the surfactant at a concentration between 0.05 mg/mL and about 0.4 mg/mL.

[0128] In certain embodiments, the reconstituted formulation comprises:

- (a) the FIX polypeptide at a concentration between about 0.9 mg/mL and about 50 mg/mL;
- (b) the buffering agent at a concentration of about 3.88 mg/mL;
- (c) the bulking agent at a concentration of about 23.8 mg/mL;
- (d) the stabilizing agent at a concentration of about 11.9 mg/mL;
- (e) the surfactant at a concentration of about 0.1 mg/mL; and
- (f) the reconstitution buffer comprising about 3.25 mg/mL NaCl.

[0129] In certain embodiments, the reconstituted formulation comprises:

- (a) the FIX polypeptide at a concentration between about 80 IU/mL and about 2,750 IU/mL;

- 32 -

- (b) the buffering agent at a concentration of about 25 mM;
- (c) the bulking agent at a concentration of about 131 mM;
- (d) the stabilizing agent at a concentration of about 35 mM;
- (e) the surfactant at a concentration of 0.01 % (w/v); and
- (f) the reconstitution buffer.

[0130] Examples of the formulation compositions are further provided in Tables 2-4.

[0131] In one aspect, the disclosure further provides a vial comprising the pre-lyophilization formulations, the lyophilate powder, or the reconstituted formulations described herein.

[0132] In another aspect, the disclosure provides a kit comprising a first container comprising the lyophilate powder described herein and a second container comprising a reconstitution buffer at a volume sufficient to produce a reconstituted formulation, when combined with the lyophilate powder of the first container. In certain embodiments, the volume of the reconstitution buffer in the kit is about 5 mL. In some embodiments, the volume is about 5.3 mL. In certain embodiments, the reconstitution buffer of the kit comprises NaCl. In some embodiments, the kit is used to treat hemophilia B.

[0133] In yet another aspect, the disclosure provides a method of administering a FIX polypeptide to a hemophilia B patient in need thereof, comprising administering to the patient the reconstituted formulation described herein, wherein the administration prevents or reduces the frequency or severity of bleeding episodes in the patient.

[0134] The disclosure further provides a method of preventing, treating, ameliorating, or managing hemophilia B in a patient in need thereof by administering the reconstituted formulation described herein.

Methods of producing a lyophilate powder comprising a FIX polypeptide

[0135] This disclosure provides methods of producing a lyophilate powder comprising a FIX polypeptide. In one aspect, the disclosure provides lyophilization methods comprising lyophilizing the pre-lyophilization formulations described herein. In another aspect, the disclosure provides a lyophilization method comprising a single drying step.

[0136] In one aspect, this disclosure provides a method of lyophilizing a FIX polypeptide comprising:

- (a) a "freezing step" comprising freezing a pre-lyophilization formulation comprising the FIX polypeptide and an aqueous solvent;

(b) a "vacuum step" comprising reducing the pressure of the frozen pre-lyophilization formulation by an amount effective to remove the aqueous solvent from the frozen pre-lyophilization formulation; and,

(c) a single "drying step" comprising increasing the temperature of the frozen pre-lyophilization formulation above the collapse temperature, thereby producing a lyophilate powder. In another aspect, the lyophilization process time is reduced compared to a reference method, *e.g.*, lyophilization process having two or more drying steps.

[0137] In other aspects, the lyophilate power produced by the present method has one or more characteristics of the following: (1) improved stability of the FIX polypeptide; (2) reduced reconstitution time; (3) reduced splashing onto a stopper comprising the formulation; or (4) increased shelf-life of the lyophilate powder at room temperature.

[0138] In certain embodiments, the collapse temperature is -1.5 °C.

[0139] In certain embodiments, the pre-lyophilization formulation is frozen to a freezing temperature of about -65 to about -40°C, about -65 to about -45 °C, about -65 to about -55°C, about -60 to about -40 °C, about -60 to about -50°C or about -60 to about -55 °C during the freezing step. In certain embodiments, the pre-lyophilization formulation is frozen to a freezing temperature of about -55°C during the freezing step. In certain embodiments, the freezing temperature is ramped down from about 5°C to about -55°C during the freezing step.

[0140] In certain embodiments, the freezing temperature is held from about 30 minutes to about 5 hours, about 1 hour to about 5 hours, about 1.5 hours to about 5 hours, about 1.5 hours to about 4 hours, about 1.5 hours to about 3 hours, or about 1.5 hours to about 2.5 hours during the freezing step. In certain embodiments, the freezing temperature is held for about 2 hours during the freezing step.

[0141] In certain embodiments, the frozen pre-lyophilization formulation of step (a) is further subject to an "annealing step" (a') prior to the "vacuum step" (b). In certain embodiments, the temperature of the frozen pre-lyophilization formulation of step (a) is ramped up to an annealing temperature of about -15°C to about -2°C during the annealing step. In certain embodiments, the temperature of the frozen pre-lyophilization formulation of step (a) is ramped up to an annealing temperature of about -6°C during the annealing step.

[0142] In certain embodiments, the annealing temperature is held for about 2 hours to about 4 hours during the annealing step. In certain embodiments, the annealing temperature is held for about 30 minutes to about 5 hours, about 1 hour to about 5 hours, about 2 hours to about 5 hours, about 2 hours to about 4 hours or about 2.5 hours to about 3.5 hours during the annealing step. In certain embodiments, the annealing temperature is held for about 3 hours during the annealing step.

[0143] In certain embodiments, the temperature of the frozen pre-lyophilization formulation is ramped down from the annealing temperature to a temperature of about -65°C to about -40°C during the annealing step. In certain embodiments, the temperature of the frozen pre-lyophilization formulation is ramped down from the annealing temperature to a temperature of -55°C during the annealing step.

[0144] In certain embodiments, the "vacuum step" comprises subjecting the frozen pre-lyophilization formulation to a vacuum between about 0.05 and about 1 mbar, between about 0.05 and about 0.50 mbar, between about 0.10 and about 0.50 mbar, between about 0.15 and about 0.50 mbar, between about 0.20 and about 0.50 mbar, or between about 0.25 and about 0.50 mbar. In certain embodiments, the vacuum in the "vacuum step" is about 0.33 mbar.

[0145] In certain embodiments, the vacuum is held in the "vacuum step" for about 5 hours, about 4 hours, about 3 hours, about 2 hours, or about 1 hour. In some embodiments, the vacuum is held in the "vacuum step" for about 2 hours.

[0146] In certain embodiments, the "drying step" comprises ramping up the temperature of the frozen pre-lyophilization formulation from about -55°C to a drying temperature of about 40°C. In certain embodiments, the drying temperature is at least about 30°C, at least about 32°C, at least about 34°C, at least about 35°C, at least about 36°C, at least about 38°C, at least about 39°C, at least about 40°C. In other embodiments, the drying temperature is about 35°C, about 40°C, about 32°C, or about 45 °C.

[0147] In certain embodiments, the drying step further comprises holding the drying temperature for about 10 hours to about 40 hours, about 10 hours to about 30 hours, or about 20 hours to about 30 hours. In certain embodiments, the drying temperature is held for about 25 hours.

[0148] In certain embodiments, the drying step is carried out at a pressure of about 0.05 mbar to about 1 mbar, between about 0.05 and about 0.50 mbar, between about 0.10 and

about 0.50 mbar, between about 0.15 and about 0.50 mbar, between about 0.20 and about 0.50 mbar, or about 0.20 mbar to about 0.45 mbar. In certain embodiments, the pressure is held at about 0.33 mbar during the drying step. The unit of mbar can be converted to Torr or any other units. For example, 1 mbar can be converted to 0.75006375541921 Torr.

[0149] In one aspect, the disclosure provides a method of producing a lyophilate powder comprising a FIX polypeptide, comprising:

(a) a "freezing step" comprising freezing a pre-lyophilization formulation comprising a FIX polypeptide by ramping down the temperature for about 2 hours to a freezing temperature of about -55°C, and holding the freezing temperature for about 2 hours;

(a') an "annealing step" comprising ramping up for about 1.5 hours the temperature of the frozen pre-lyophilization formulation of step (a) to an annealing temperature of about -6°C, holding the annealing temperature for about 3 hours, and ramping down the temperature for about 1.5 hours to about -55°C;

(b) a "vacuum step" comprising holding the frozen pre-lyophilization formulation of step (a') at about -55°C for two hours at atmospheric pressure and ramping down the pressure for about 2 hours to about 0.33 mbar; and,

(c) a single "drying step" comprising ramping up for 3 hours the temperature of the frozen pre-lyophilization formulation of step (b) to about 40°C, while holding the pressure at about 0.33 mbar, and holding the temperature of the frozen pre-lyophilization formulation at about 40°C for about 25 hours, while holding the pressure at about 0.33 mbar,

thereby producing the lyophilate powder. In some embodiments, the lyophilization method takes less cycle time.

[0150] In certain embodiments, the lyophilate powder produced by the methods described herein has the following characteristics:

- (1) improved stability of the FIX polypeptide;
- (2) reduced reconstitution time;
- (3) reduced splashing onto a stopper comprising the formulation;
- (4) increased shelf-life of the lyophilate powder at room temperature; or
- (5) any combinations thereof,

[0151] In some embodiments, the lyophilization cycle period can be less than about 4.5 days, about 4 days, about 3.5 days, about 3 days, about 2.5 days, or about 2 days. In other embodiments, the lyophilization cycle period is about 3 days or less. In certain embodiments, the fill volume of the pre-lyophilization formulation used in the lyophilization method is less than about 5 mL. In certain embodiments, the fill volume is about 4 mL, about 3.5 mL, about 3.0 mL, about 2.9 mL, about 2.8 mL, about 2.7 mL, about 2.65 mL, about 2.6 mL, about 2.5 mL, about 2.4 mL, about 2.3 mL, about 2.2 mL, about 2.1 mL, or about 2.0 mL. In one embodiment, the fill volume is about 2.65 mL.

[0152] In certain embodiments, the reduced reconstitution time of the lyophilate powder produced by the lyophilization method is less than about 1.5 minutes, less than about 1 minute, less than about 50 seconds, less than about 40 seconds, less than about 30 seconds, less than about 20 seconds, or less than about 10 seconds. In certain embodiments, the reduced reconstitution time of the lyophilate powder produced by the lyophilization method is less than about 30 seconds.

[0153] In certain embodiments, the reduced lyophilization cycle time of the pre-lyophilization formulation used in the lyophilization method is about 4 days or less, about 3 days or less, about 2 days or less, or about a day or less.

[0154] In some embodiments, the lyophilate powder is produced from the pre-lyophilization formulation in about 90 hours or less, about 80 hours or less, about 70 hours or less, about 60 hours or less, about 50 hours or less, about 45 hours or less, about 40 hours or less, or about 30 hours or less. In certain embodiments, the lyophilate powder is produced from the pre-lyophilization formulation in about 45 hours or less.

[0155] In certain embodiments, the residual moisture in the lyophilate powder is less than about 1.0%, about 0.7%, about 0.6%, about 0.5%, about 0.4% or about 0.3%. In some embodiments, the residual moisture in the lyophilate powder is less than about 0.5%.

[0156] In one aspect, the disclosure provides a method of stabilizing a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the lyophilate powder is stabilized as measured by Size Exclusion Chromatography (SEC) with respect to a lyophilate powder prepared by using a lyophilization method comprising more than one drying step.

[0157] In another aspect, the disclosure provides a method of increasing the shelf-life of a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the shelf-life of the lyophilate powder is increased as measured by SEC and/or FIX clotting activity assay with respect to the shelf-life of a lyophilate powder prepared by using a lyophilization method comprising more than one drying step.

[0158] This disclosure also provides a method to decrease the reconstitution time of a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the reconstitution time of the lyophilate powder is decreased with respect to the reconstitution time of a lyophilate powder prepared by using a lyophilization method comprising more than one drying step.

[0159] This disclosure further provides a method to reduce lyophilization process time of producing a lyophilate powder comprising a FIX polypeptide, comprising lyophilizing a pre-lyophilization formulation according to the methods described herein, wherein the lyophilization process time of the pre-lyophilization formulation is reduced with respect to the lyophilization process time of producing a lyophilate powder using a lyophilization method comprising more than one drying step.

[0160] Having now described the present invention in detail, the same will be more clearly understood by reference to the following examples, which are included herewith for purposes of illustration only and are not intended to be limiting of the invention. All patents and publications referred to herein are expressly incorporated by reference.

EXAMPLES

Example 1: Factor IX-Fc Drug Substance and Drug Product Compositions

rFIXFc Description

[0161] rFIXFc is a long-acting, fully recombinant fusion protein consisting of human coagulation Factor IX (FIX) covalently linked to the Fc domain of human immunoglobulin G1 (IgG1). The Factor IX portion of rFIXFc has a primary amino acid sequence that is identical to the Thr¹⁴⁸ allelic form of plasma derived Factor IX and has structural and functional characteristics similar to endogenous Factor IX. The Fc domain

- 38 -

of rFIXFc contains the hinge, CH2 and CH3 regions of IgG1. rFIXFc contains 869 amino acids with a molecular weight of approximately 98 kilodaltons.

[0162] rFIXFc is produced by recombinant DNA technology in a human embryonic kidney (HEK) cell line and then purified.

[0163] The rFIXFc drug product formulation of the invention comprising rFIXFc can allow development of high concentration drug products, *e.g.*, 4000+ IU/vial drug product strengths. This requires a higher concentration of rFIXFc protein in the drug substance, as shown in the comparison Table 1 below. The rFIXFc drug product formulation of the invention can increase the shelf life of the 250 & 500 IU/vial drug product strengths to allow increased room temperature stability. Development data suggests that the 250 and 500 IU/vial drug product strengths are significantly more stable under accelerated conditions than the reference drug product.

[0164] The rFIXFc drug product formulation of the invention can also decrease the reconstitution time when lyophilized. Reference drug product reconstitution time varies between 1-2 minutes. Development data suggests that the LCM drug product reduces the reconstitution time to less than 30 seconds.

[0165] The rFIXFc drug product formulation of the invention can also allow reduction of the lyophilization process time. Currently the lyophilization cycle is ~4.5 days. With a lower vial fill volume this can likely be reduced to ~3 days or less for more economical manufacturing.

Drug Substance (DS)

[0166] The drug substance for the rFIXFc formulation of the invention will use the same formulation excipients as the reference drug substance. The higher concentration will be achieved using a second ultrafiltration step during drug substance manufacturing. *See* Table 1.

Table 1: Drug Substance Compositions

Component	Drug Substance Compositions	Reference Drug Product	Drug Product
rFactorIX-Fc		10-13 mg/mL	65-75 mg/mL
L-Histidine		3.88 mg/mL (25 mM)	3.88 mg/mL (25 mM)

- 39 -

Polysorbate 20 0.1 mg/mL (0.01 %) 0.1 mg/mL (0.1 %)

Drug Product (DP)

[0167] In order to accomplish the objectives stated above, a drug product was designed that doubled the concentrations of all components of the reference drug product (protein & excipients), while reducing the fill volume of the vial prior to lyophilization. This ensures that the dose of all components to the patient remains constant, while improving the drug product performance parameters mentioned above.

[0168] Tables 2-4 below detail the composition of the lyophilization feedstock, the contents of the solid product in the vial after lyophilization and the composition after reconstitution. It is important to note the wide variance of the rFIXFc protein concentrations. Due to the fact that each batch of Factor IX is slightly different in its activity in IU/mg, the feedstock is compounded using the assayed activity. This results in a range of protein concentrations, and this variance added to the different strengths of drug products gives the ranges stated below.

Table 2: Drug Product – Pre-lyophilization Formulation Compositions

Drug Product Lyophilization Feedstock Compositions			
Component	Reference Drug Product 250-2000 IU/Vial	Reference Drug Product 3000 IU/Vial	Drug Product 250-5500 IU/Vial
rFactorIX-Fc	0.45-9.5 mg/mL 40-500 IU/mL	5.5-13.7 mg/mL 480-750 IU/mL	0.9-50 mg/mL 80-2750 IU/mL
L-Histidine	3.88 mg/mL (25 mM)	3.88 mg/mL (25 mM)	7.76 mg/mL (50 mM)
Mannitol	23.8 mg/mL (131 mM)	23.8 mg/mL (131 mM)	47.6 mg/mL (261 mM)
Sucrose	11.9 mg/mL (35 mM)	11.9 mg/mL (35 mM)	23.8 mg/mL (70 mM)
Polysorbate 20	0.1 mg/mL (0.01 %)	0.1 mg/mL (0.01 %)	0.2 mg/mL (0.2 %)
Feedstock Fill Volume	5.3 mL	7.4 mL	2.65 mL

Table 3: Drug Product – Lyophilate Compositions

Drug Product Lyophilized Vial Solid Compositions			
Component	Reference Drug Product	Reference Drug Product	Drug Product 250-5500 IU/Vial*

- 40 -

	250-2000 IU/Vial	3000 IU/Vial	
rFactorIX-Fc	2.2-46 mg/vial	28-69 mg/vial	2.2-125 mg/vial
L-Histidine	19.4 mg/vial	27.2 mg/vial	19.4 mg/vial
Mannitol	119 mg/vial	167 mg/vial	119 mg/vial
Sucrose	59.5 mg/vial	83.3 mg/vial	59.5 mg/vial
Polysorbate 20	0.5 mg/vial	0.7 mg/vial	0.5 mg/vial

*The values are the nominal values that do not include the overfill. The values including the overfill are: 20.6 mg/vial (L-histidine), 126.01 mg/vial (mannitol), 63.1 mg/vial (sucrose), and 0.53 mg/vial (polysorbate).

Table 4: Drug Product – Reconstituted Formulation Compositions

Component	Post-Reconstitution Drug Product Vial Compositions		
	Reference Drug Product 250-2000 IU/Vial	Reference Drug Product 3000 IU/Vial	Drug Product 250-5500 IU/Vial
Diluent		5.2 mL of NaCl in Water for Injection	
rFactorIX-Fc	0.45-9.5 mg/mL 40-500 IU/mL	5.5-13.7 mg/mL 480-750 IU/mL	0.9-50.0 mg/mL 80-2750 IU/mL
L-Histidine	3.88 mg/mL (25 mM)	5.43 mg/mL (35 mM)	3.88 mg/mL (25 mM)
Mannitol	23.8 mg/mL (131 mM)	33.3 mg/mL (183 mM)	23.8 mg/mL (131 mM)
Sucrose	11.9 mg/mL (35 mM)	16.7 mg/mL (48 mM)	11.9 mg/mL (35 mM)
Polysorbate 20	0.1 mg/mL (0.01 %)	0.14 mg/mL (0.014 %)	0.1 mg/mL (0.01 %)
Reconstitution buffer	3.25 mg/mL	3.25 mg/mL	3.25 mg/mL
Reconstituted Volume	5.3 mL	5.3 mL	5.3 mL

Example 2: Development of Lyophilization Cycle Parameters for the Second Generation rFIXFc Drug Product

SUMMARY

[0169] The goal of this study was to evaluate ranges around the drying phase process parameters of the lyophilization cycle for the rFIXFc drug product of the invention.

- 41 -

[0170] This report summarizes a statistical design of experiments (DOE) study evaluating the lyophilization process parameters (Drying Shelf Temperature, Chamber Vacuum Level and Drying Time) and their effect on the product temperature during drying, the resulting residual moisture and drying rate of the drug product.

[0171] Preliminary lyophilization cycle design experiments on the Second Generation rFIXFc drug product demonstrated that distinct primary and secondary drying steps were not necessary due to the formulation's high collapse temperature of approximately -1.5 °C. A 12 experiment Design of Experiments (DOE) study was developed to evaluate the effect of the shelf temperature, vacuum level and drying time on the residual moisture levels and product temperatures during the lyophilization process on the placebo. An analysis of the data shows that in order to achieve a residual moisture level below 1%, a minimum shelf temperature of 30 °C during the drying phase is required. The analysis also demonstrated that drying times longer than 25 hours do not further decrease the residual moisture in the vials, and the chamber vacuum level has only a small effect on the residual moisture. The product temperatures during drying are significantly affected by the shelf temperature and chamber vacuum level, but the most aggressive drying conditions in the study (40 °C Shelf Temperature at 1000 mTorr Chamber Vacuum) resulted in a product temperature that was more than 10 °C colder than the collapse temperature. The vial mass flow is predominantly a function of the shelf temperature, and in order to maintain vacuum control, the commercial lyophilizer will need to be able to handle moisture flow rates of 0.7 g/hr/vial. A lyophilization cycle was proposed to achieve product with < 0.5% residual moisture using a shelf temperature of 40 °C, a chamber vacuum of 250 mTorr (0.33 mBar) and a drying time of 25 hours.

INTRODUCTION

[0172] The Second Generation FactorIX-Fc (rFIXFc-2G) Drug Product composition was designed to allow for improved protein stability during accelerated storage, improved reconstitution time, and reduced fill volume to reduce splashing onto the stoppers. This was accomplished by reducing the fill volume from 5.3 mL to 2.65 mL, and doubling the concentrations of protein and excipients in the formulation such that the reconstituted product is the same as the first generation composition. Another benefit to reducing the fill volume was a decrease in the amount of water that needs to be removed during the lyophilization process.

- 42 -

[0173] Since the new drug product design required revalidation of the drug product process, the lyophilization cycle was redeveloped. The collapse temperature of the placebo formulation was measured as approximately -1.5 °C, which would allow a shorter lyophilization cycle to be used than was designed for the reference rFIXFc lyophilization process. It was determined through initial experiments that no separate primary drying step was required, as the product did not undergo collapse even at shelf temperatures in excess of 40 °C. A cycle was designed using the freezing profile from the reference drug product combined with a direct step to the primary drying temperature after vacuum is applied. The placebo is a good surrogate for active rFIXFc-2G vials since the mannitol provides the crystalline structure for the cake making the appearance the same as the active vials. Amorphous sugar is also more difficult to dry than protein, so the resulting residual moisture is slightly higher, providing a worst-case value for the process. Removal of the protein from the vials also reduces the resistance to water vapor, giving a worst case estimate for the vial mass flow rate.

[0174] A statistical design of experiments (DOE) study was carried out to evaluate the lyophilization process parameters (Drying Shelf Temperature, Chamber Vacuum Level and Drying Time) and their effect on the product temperature during the drying process, the resulting residual moisture, and drying rate of the drug product.

MATERIALS AND METHODS

[0175] The goal of this study was to evaluate ranges around the drying phase process parameters of the lyophilization cycle for the second generation rFIXFc drug product. The drying phase parameters and ranges used to design the DOE study in JMP 9 are shown in Table 5. The resulting 12 experiment DOE plan showing the individual run parameter setpoints is shown in Table 6.

Table 5: The Drying Parameter Ranges Used in the DOE Study Design

Drying DOE Parameter	Minimum Value	Maximum Value
Shelf Temperature (°C)	0	40
Chamber Vacuum (mTorr)	100	1000
Drying Time (h)	15	35

Table 6: The Study Design of Experiments Template Output from JMP 9

- 43 -

Run	Temp (C)	Vacuum (mT)	Time (hr)
1	40	100	35
2	0	100	15
3	40	1000	25
4	20	100	28.2
5	20	500	25
6	0	100	25
7	20	550	25
8	40	250	15
9	20	1000	15
10	0	1000	35
11	10	500	25
12	40	500	25

[0176] For each lyophilization cycle in the study, eighty 10 mL Schott vials (P/N: 68000320) were filled with 2.75 mL of Second Generation rFIXFc Placebo as shown in Table 7, which provides a worst case fill volume for residual moisture evaluation. The filled vials were arranged on a single shelf with three thermocouples for each experiment as shown in Figure 1.

Table 7: Second Generation rFIXFc Drug Product Placebo Recipe

Component	FW	g/L	mM	% (w/v)
Histidine	155.15	7.76	50.00	0.776
PS-20	1227.54	0.2	0.1629	0.020
Mannitol	182.172	47.6	261.29	4.760
Sucrose	342.30	23.8	69.53	2.380
	ρ =	1.029	g/mL	

[0177] The lyophilization cycles used were variations of the cycle shown in Table 8. The Lyophilizer Shelf Drying Temperature, Drying Step Time and Drying Vacuum Level were varied based on the Design of Experiments table shown in Table 6. An SP Industries Lyostar II was used for each lyophilization cycle, and the vials were placed on the middle shelf.

- 44 -

Table 8: Lyophilization Cycle Parameters Showing the DOE Inputs of Temperature, Time and Vacuum

Phase	Step	Temp (°C)	Ramp (°C/min)	Soak (min)	Vacuum (mTorr)
Load	Load	25	-	As Needed	-
Freeze	Equilibrate	0	0.5	60	-
	Freeze	-55	0.5	120	-
	Anneal	-6	0.5	180	-
	Freeze	-55	0.5	120	-
Pulldown	Initial Vac	-55	-	-	DOE Vacuum
Drying	1°/2° Drying	DOE Temp	Direct Step	DOE Time	DOE Vacuum
Final	Final	5	5.0	As Needed	DOE Vacuum

[0178] After each lyophilization cycle, five vials were selected from corner and middle positions and measured for residual moisture using procedure TDMP-74, and averaged over the shelf. The product temperature during drying and the vial mass flow rate were measured using thermocouples and by Manometric Temperature Measurement in the Lyostar II software. These outputs were analyzed using JMP 9 software to evaluate the effect of the drying parameters on the second generation rFIXFc lyophilization process. A JMP stepwise analysis was performed to determine the significant variables, and these variables were then analyzed using a standard least squares effect screening algorithm which shows how the process outputs (residual moisture, product temperature and mass flow during drying) respond to the input variables.

RESULTS AND DISCUSSION

[0179] The results of the twelve lyophilization experiments are shown in Table 9.

- 45 -

Table 9: The DOE Results of Residual Moisture, Vial Mass Flow and Product Temperature

Run	Temp (C)	Vacuum (mT)	Time (hr)	NB Ref	% Moisture	dm/dt (g/hr/vial)	T _p (MTM, °C)
1	40	100	35	16322-117	0.33	0.53	-26.5
2	0	100	15	16322-110	2.82	0.28	-32.2
3	40	1000	25	16322-090	0.57	0.45	-15.2
4	20	100	28.2	16322-139	1.28	0.41	-29.5
5	20	500	25	16322-098	1.58	0.57	-20.3
6	0	100	25	16322-151	2.18	0.27	-32.5
7	20	550	25	16322-146	1.49	0.56	-19.7
8	40	250	15	18266-004	0.59	0.70	-21.9
9	20	1000	15	16322-131	1.95	0.38	-15.6
10	0	1000	35	16322-124	2.56	0.17	-15.8
11	10	500	25	16322-104	2.01	0.35	-21.8
12	40	500	25	16322-083	0.49	0.56	-19.8

1. *Analysis of Lyophilization Cycle Parameters on Residual Moisture*

[0180] The rFIXFc-2G placebo was used as a worst-case surrogate for the drug product as it is generally more difficult to remove residual moisture from sugars during secondary drying than protein. The resulting prediction profiler showing the outcome of the DOE analysis is shown in Figure 2. Several observations are apparent: The shelf temperature has the most significant impact on the residual moisture in the drug product. This is expected based on the fact that secondary drying, which removes tightly bound water, is a diffusion and desorption controlled process. The model predicts with high confidence that shelf temperatures higher than 30 °C are required to achieve residual moisture levels lower than 1%. The vacuum level appears to have a small but measurable effect on the resulting residual moisture. The drying time appears to show a point of diminishing returns starting at 25 hours where the addition of further drying time does not continue to decrease the residual moisture level. This type of behavior is consistent with the kinetic approach to an equilibrium boundary determined by the shelf temperature, and the residual moisture approaches an asymptote where further drying is not possible. Based on this residual moisture DOE analysis the drying shelf temperature should be 30 °C or greater and the drying time should be fixed at 25 hours or less.

- 46 -

2. *Analysis of Lyophilization Cycle Parameters on Product Temperature During Sublimation*

[0181] The freeze drying collapse temperature of the rFIXFc-2G Drug Product placebo has been measured as approximately -1.5 °C. Practically, this means that the drug product will maintain an elegant cake structure so long as the product temperature is maintained below this collapse temperature as the bulk water is removed from the vial during lyophilization. The DOE analysis determined that the shelf temperature and chamber vacuum levels both had a significant effect on the product temperature as shown in Figure 3. The chamber vacuum had the largest effect, with higher pressures translating to higher product temperatures during sublimation. Even at 1000 mTorr (1.33 mBar) the highest measured product temperature was -15.2 °C, approximately 13 °C below the product collapse temperature. The shelf temperature also had a modest effect on the product temperature, but the results are less pronounced than the vacuum effect. This analysis shows that there is little risk of collapse even at a shelf temperature of 40 °C and a chamber vacuum level of 1000 mTorr, essentially eliminating the potential of collapse from any practical lyophilization cycle design space.

3. *Analysis of Lyophilization Cycle Parameters on Vial Mass Flow Rate During Sublimation*

[0182] The vial mass flow rate (dm/dt) is a measure of the rate at which water is being removed from the vials during the sublimation process. While faster drying is desirable to reduce the time required for the lyophilization cycle, too much moisture can overwhelm the condensers in manufacturing scale freeze driers and lead to a loss of vacuum control in the product chamber. The placebo represents the worst case vial mass flow conditions. Since there is no protein present in the formulation, the solids percentage in the cake is minimized and this results in a lower resistance to mass flow from the freeze dried cake. The shelf temperature has a significant effect on the vial mass flow as shown in Figure 4 with increasing temperature causing faster sublimation. The chamber vacuum level was included in the DOE analysis model, but the p value is 0.136 which is not significant with 95% confidence. The highest measured dm/dt in the study was 0.7 g/hr/vial.

4. Proposed Second Generation rFIXFc Lyophilization Cycle Based on the Placebo DOE Study

[0183] The data from the Placebo DOE study suggest it is feasible to design a lyophilization cycle to achieve a residual moisture target of < 0.5% while maintaining the product below its collapse temperature using a single drying step. The proposed lyophilization cycle is shown in Table 10, and the data from DOE Run 8 which is under similar conditions to the proposed rFIXFc-2G lyophilization cycle is shown in Figure 5. The residual moisture target of 0.5 % was selected as this is the average value of the first generation rFIXFc drug product strength series. This moisture level provides a cushion so that as the product absorbs moisture during accelerated stability, product quality attributes will not be affected.

[0184] The freezing and annealing portions of the lyophilization cycle were used as developed for the reference rFIXFc drug product lyophilization cycle, and the separate primary and secondary drying steps have been replaced with a single drying step at 40 °C shelf temperature and 250 mTorr vacuum for 25 hours.

Table 10: The Proposed Lyophilization Cycle for rFIXFc-2G

Phase	Step	Temp (°C)	Ramp (°C/min)	Soak (min)	Vacuum (mTorr)
Load	Load	25	-	As Needed	-
Freeze	Equilibrate	0	0.5	60	-
	Freeze	-55	0.5	120	-
	Anneal	-6	0.5	180	-
	Freeze	-55	0.5	120	-
Pulldown	Initial Vac	-55	-	-	250
Drying	1/2° Drying	40	0.5	1500	250
Final	Final	5	5.0	As Needed	250

CONCLUSIONS

[0185] A 12 experiment DOE study evaluating the Second Generation rFIXFc Drug Product lyophilization process parameters on the residual moisture, product temperature and vial mass flow rate of the placebo was completed. An analysis of the data shows that in order to achieve a residual moisture level below 1%, a minimum shelf temperature of 30 °C during the drying phase is required. The analysis also demonstrated that drying times longer than 25 hours do not significantly decrease the residual moisture in the vials, and the vacuum level has only a small effect on the residual moisture. The product temperatures during drying are significantly affected by the shelf temperature and chamber vacuum level. The most aggressive conditions in the

study (40 °C Shelf Temperature at 1000 mTorr Chamber Vacuum) resulted in a product temperature that was more than 10 °C colder than the collapse temperature.

[0186] A lyophilization cycle was proposed to achieve product with < 0.5% residual moisture using a shelf temperature of 40 °C, a chamber vacuum of 250 mTorr (0.33 mBar) and a drying time of 25 hours.

[0187] The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying knowledge within the skill of the art, readily modify and/or adapt for various applications such specific embodiments, without undue experimentation, without departing from the general concept of the present invention. Therefore, such adaptations and modifications are intended to be within the meaning and range of equivalents of the disclosed embodiments, based on the teaching and guidance presented herein. It is to be understood that the phraseology or terminology herein is for the purpose of description and not of limitation, such that the terminology or phraseology of the present specification is to be interpreted by the skilled artisan in light of the teachings and guidance.

[0188] The breadth and scope of the present invention should not be limited by any of the above-described exemplary embodiments, but should be defined only in accordance with the following claims and their equivalents. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein.

[0189] All documents, articles, publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

[0190] The present application claims priority to U.S. Provisional Application No. 61/969,801, filed March 24, 2014, which is incorporated herein by reference in its entirety.

- 49 -

TABLE OF SEQUENCES

Table 11: Polynucleotide Sequences of FIX

FIX-Fc Chain DNA Sequence (FIX signal peptide underlined, FIX sequence double underlined, Fc region in bold) (SEQ ID NO:1, which encodes SEQ ID NO:2)

pSYN-FIX-030 Nucleotide sequence (nt 1 to 7583) :

FIX exon 1 (signal peptide, 1st amino acid propeptide) : nt 690-777

FIX mini intron: nt 778-1076

FIX sequence : nt 1077-2371

Fc : nt 2372-3052

```

1 gcgcgcttg acattgatta ttgacttagtt attaatagta atcaattacg
51 ggttcattag ttcatacgccc atatatggag ttccgcgtta cataacttac
101 ggtaaatggc ccgcctggct gaccgccc aa cgaccggc ccattgacgt
151 caataatgac gtatgttccc atagtaacgc caataggac tttccattga
201 cgtcaatggg tggagtattt acggtaaact gcccacttgg cagtacatca
251 agtgtatcat atgccaagta cgccccctat tgacgtcaat gacggtaaat
301 ggcccgccctg gcattatgcc cagtacatga ccttatggga ctttcctact
351 tggcagtaca tctacgtatt agtcatcgct attaccatgg tgatgcgggt
401 ttggcagtac atcaatgggc gtggatagcg gtttgactca cggggatttc
451 caagtctcca ccccattgac gtcaatggga gtttgggg gcaccaaaat
501 caacgggact ttccaaaatg tcgtaacaac tccgcggcat tgacgcaaat
551 gggcggtagg cgtgtacggt gggaggtcta tataaggcaga gctctctggc
601 taactagaga acccactgct tactggctta tcgaaattaa tacgactcac
651 tatagggaga cccaaagcttc ggcacgtacg gccgcacca tgcagcgcgt
701 gaacatgatc atggcagaat caccaggct catcaccatc tgccttttag
751 gatatctact cagtgctgaa tgtacaggtt tgttccctt tttaaaatac
801 attgagtagt cttgcctttt agatataaaaa atatctgatc ctgtcttctt
851 cactaaattt tgattacatg atttgacagc aatattgaag agtctaacag
901 ccagcacgca gggtggtaag tactgtggga acatcacaga ttttggctcc

```

- 50 -

951 atgccctaaa gagaaattgg ctttcagatt atttggatta aaaacaaaga
1001 ctttcttaag agatgtaaaa ttttcatgat gttttctttt ttgctaaaac
1051 taaagaatta ttctttaca ttcagttt tcttqatcat aaaaacqcca
1101 acaaaattct qaatcgqcca aaqaggtata attcaqgtaa attqqaagag
1151 tttgttcaag qqaatctaga qaqagaatgt atqqaagaaa agtqtagttt
1201 tqaqaqaagca cqqaqaqttt ttqaaaacac tqaaaagaaca actqaatttt
1251 qqaqcaqta tqttqatqqa qatcaqtqtq agtccaatcc atgtttaat
1301 gqcqqcaqtt qcaaggatqa cattaattcc tatqaatqtt qgtqccctt
1351 tggatttgaa qgaaagaact qtgaattaga tqtaacatgt aacattaaga
1401 atqqcaqatq cqagcaqttt tqtaaaaata qtqctqataa caagqtqggtt
1451 tgctcctqta ctqagggata tcqacttqca qaaaaccaga agtccctqtqa
1501 accagcagta ccatttccat qtqqaagadt ttctqttca caaacttcta
1551 agctcacccq tgctqagact qttttcctq atqtggacta tgtaaattct
1601 actqaagctq aaaccatttt qgataacatc actcaaagca ccaaatcatt
1651 taatgacttc actcgqattq ttqgtggaga aqatqccaaa ccaggtcaat
1701 tcccttggca qgttqttttq aatqgtaaag ttqatqcatt ctqtggaggg
1751 tctatcgqta atqaaaaatq qattqtaact qctqcccact qtqttqaaac
1801 tggtgttaaa attacagttq tcqcaaggta acataatatt qagggagacag
1851 acatacaga qcaaaagcga aatqtgattc qaattattcc tcaccacaaac
1901 tacaatgcag ctattaataa qtacaaccat qacattqccc ttctqqaact
1951 gqacqaaccc ttaqtqctaa acagctacgt tacacctatt tqcattgcta
2001 acaaggaata cacqaacatc ttcctcaaatt ttqgatctqg ctatgttaagt
2051 gqctqgggaa gagtcttcca caaagggaga tcaqctttag ttcttcagta
2101 ccttagagtt ccacttgttq accqagccac atgtcttcga tctacaaaagt
2151 tcaccatcta taacaacatq ttctgtqctq qcttccatqa aqqaqqtada
2201 gattcatqtc aaqqaqataq tqqqqqgaccc catqttactq aagtqqaagq
2251 gaccagtttc ttaactggaa ttattagctq qqgtgaaqag tgtgcaatqa
2301 aaggcaaata tqqaatataat accaaqgtgt cccggtatgt caactggatt
2351 aaggaaaaaaa caaagctcac tgacaaaaact cacacatgcc caccgtgccc
2401 agctccggaa ctcctggcg gaccgtcagt cttcctttc cccccaaaac

- 51 -

2451 **ccaaggacac cctcatgatc tcccggaccc ctgaggtcac atgcgtggtg**
2501 **gtggacgtga gccacgaaga ccctgaggtc aagttcaact ggtacgtgga**
2551 **cggcgtggag gtgcataatg ccaagacaaa gccgcgggag gagcagtaca**
2601 **acagcacgta ccgtgtggtc agcgtcctca ccgtcctgca ccaggactgg**
2651 **ctgaatggca aggagtacaa gtgcaaggtc tccaacaaag ccctcccagc**
2701 **ccccatcgag aaaaccatct ccaaagccaa agggcagccc cgagaaccac**
2751 **aggtgtacac cctgccccca tcccggatg agctgaccaa gaaccaggtc**
2801 **agcctgaccc gcctggtcaa aggcttctat cccagcgaca tcgccgtgga**
2851 **gtgggagagc aatgggcagc cggagaacaa ctacaagacc acgcctcccg**
2901 **tgttggactc cgacggctcc ttcttcctct acagcaagct caccgtggac**
2951 **aagagcaggt ggcagcaggg gaacgtcttc tcatgctccg tcatgcatga**
3001 **ggctctgcac aaccactaca cgcagaagag cctctccctg tctccggta**
3051 **aatgagaatt cagacatgat aagatacatt gatgagttt gacaaaccac**
3101 aactagaatg cagtaaaaaa aatgctttat ttgtgaaatt tgtgatgcta
3151 ttgctttatt tgtaaccatt ataagctgca ataaacaagt tgggtgggc
3201 gaagaactcc agcatgagat ccccgcgctg gaggatcatc cagccggcgt
3251 cccggaaaac gattccgaag cccaacctt catagaaggc ggcgggtggaa
3301 tcgaaatctc gtagcacgtg tcagtcctgc tcctcggcca cgaagtgcac
3351 gcagttgccg gccgggtcgc gcagggcgaa ctccggcccc cacggctgct
3401 cgccgatctc ggtcatggcc ggccccggagg cgtcccgaa gttcgtggac
3451 acgacctccg accactcggc gtacagctcg tccaggccgc gcacccacac
3501 ccaggccagg gtgttgcgg accacacccg cgaagtgcgc ctccacgaag
3551 acagggtcac gtcgtcccg accacacccg cgaagtgcgc ctccacgaag
3601 tcccggaga acccgagccg gtcggtccag aactcgaccc ctccggcgcac
3651 gtcgcgcgcg gtgagcaccg gaacggcact ggtcaacttg gccatgggtt
3701 agttcctcac cttgtcgat tataactatgc cgatatacta tgccgatgat
3751 taattgtcaa cacgtgctga tcagatccgaa aatggatata acaagctccc
3801 gggagcttt tgcaaaagcc taggcctcca aaaaagcctc ctcactactt
3851 ctggaatagc tcagaggcag aggcggcctc ggcctctgca taaataaaaa

- 52 -

3901 aaattagtca gccatggggc ggagaatggg cggaactggg cggagtagg
3951 ggcgggatgg gcggagtag gggcgact atgggtgctg actaatttag
4001 atgcatgctt tgccatacttc tgccctgctgg ggagcctggg gactttccac
4051 acctggttgc tgactaattg agatgcatgc tttgcatact tctgcctgct
4101 ggggagcctg gggactttcc acaccctcggt cgagctagct tcgtgaggct
4151 ccggtgcccg tcagtggca gaggcacat cgcccacagt ccccgagaag
4201 ttggggggag gggtcggcaa ttgaaccggt gcctagagaa ggtggcgcgg
4251 ggtaaaactgg gaaagtgatg tcgtgtactg gctccgcctt tttcccgagg
4301 gtgggggaga accgtatata agtgcagtag tcgcccgtgaa cgttttttt
4351 cgcaacgggt ttgcccgg aacacaggt a gtcggctgt gtggcccc
4401 cgggcctggc ctcttacgg gttatggccc ttgcgtgcct tgaattactt
4451 ccacctggct ccagtagctg attcttgcatt cccagctgg a cccaggggcg
4501 ggccttgcgc tttaggagcc ctttcgcctc gtgcttgagt tgaggcctgg
4551 cctggcgct gggccggccg cgtgcgaatc tggtggcacc ttgcgcctg
4601 tctcgctgct ttgcataagt ctctagccat taaaatttt tgatgacctg
4651 ctgcgacgct tttttctgg caagatagtc ttgtaaatgc gggccaggat
4701 ctgcacactg gtatttcggg ttttggggcc gggggcgccg acggggcccg
4751 tgcgtcccg cgcacatgtt cggcgaggcg gggcctgcga ggcggccac
4801 cgagaatcg acggggtag tctcaagctg gccggcctgc tctggcct
4851 ggcctgcgc cggcgtgtat cggccggcc tggcggcaa ggctggcccg
4901 gtcggcacca gttgcgtgag cgaaaagatg gccgcttccc ggccctgctc
4951 cagggggctc aaaatggagg acgcggcgct cggagagcg ggcgggtgag
5001 tcacccacac aaaggaaagg ggctttccg tcctcagccg tcgcttcatg
5051 tgactccacg gagtaccggg cggcgtccag gcacctcgat tagttctgg
5101 gctttggag tacgtcgtct ttaggttggg gggaggggtt ttatgcgtat
5151 gagtttcccc acactgagtg ggtggagact gaagttaggc cagcttggca
5201 cttgatgtaa ttctccttgg aatttgcctt tttttaggttt ggatcttgg
5251 tcattctcaa gcctcagaca gtggttcaaa gttttttct tccatttcag
5301 gtgtcgtgaa cacgtggcgcgccc gccaccatgg agacagacac
5351 actcctgcta tgggtactgc tgctctgggt tccaggtcc actggtgaca

- 53 -

5401 aaactcacac atgcccacccg tgcccagcac ctgaactcct gggaggaccg
5451 tcagtcttcc tcttcccccc aaaacccaag gacaccctca tgatctcccg
5501 gaccctgag gtcacatgcg tgggtggta cgtgagccac gaagaccctg
5551 aggtcaagtt caactggtaac gtggacggcg tggaggtgca taatgccaag
5601 acaaagccgc gggaggagca gtacaacagc acgtaccgtg tggcagcgt
5651 cctcaccgtc ctgcaccagg actggctgaa tggcaaggag tacaagtgca
5701 aggtctccaa caaaggccctc ccagccccca tcgagaaaac catctccaaa
5751 gccaaagggc agccccgaga accacaggtg tacaccctgc ccccatcccg
5801 cgatgagctg accaagaacc aggtcagcct gacctgcctg gtcaaaggct
5851 tctatccag cgacatcgcc gtggagtggg agagcaatgg gcagccggag
5901 aacaactaca agaccacgac tcccgtgtt gactccgacg gtccttctt
5951 cctctacagc aagctcaccg tggacaagag caggtggcag cagggaaacg
6001 tcttctcatg ctccgtgatg catgaggctc tgcacaacca ctacacgcag
6051 aagagcctct ccctgtctcc gggtaaatga ctcgagagat ctggccggct
6101 gggcccggtt cgaaggtaag cctatcccta accctctcct cggtctcgat
6151 tctacgcgta ccggcatca tcaccatcac cattgagttt aaacccgctg
6201 atcagcctcg actgtgcctt ctagttgcca gccatctgtt gtttgcctt
6251 ccccccgtgcc ttccttgacc ctggaaaggtg ccactccac tgcctttcc
6301 taataaaaatg aggaaattgc atcgcattgt ctgagtaggt gtcattctat
6351 tctgggggggt ggggtggggc aggacagcaa gggggaggat tgggaagaca
6401 atagcaggca tgctggggat gcgggtgggt ctatggcttc tgaggcggaa
6451 agaaccagtg gcggtaatac gtttatccac agaattcagg gataacgcag
6501 gaaagaacat gtgagcaaaa ggccagcaaa aggccaggaa ccgtaaaaag
6551 gcccgcgttgc tggcgttttt ccataggctc cggcccccctg acgagcatca
6601 caaaaatcga cgctcaagtc agaggtggcg aaacccgaca ggactataaa
6651 gataccaggc gtttccccc agaagctccc tcgtgcgcctc tcctgttccg
6701 accctgcgc ttaccggata cctgtccgc tttctccctt cgggaagcgt
6751 ggcgcgttct catagctcac gctgttaggtt tctcagttcg gtgttaggtcg
6801 ttgcgtccaa gctgggctgt gtgcacgaac ccccggttca gcccgcaccgc

- 54 -

6851 tgccgccttat ccggtaacta tcgtctttag tccaaacccgg taagacacgaa
 6901 cttatcgcca ctggcagcag ccactggtaa caggattagc agagcgaggt
 6951 atgttaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac
 7001 actagaagaa cagttttgg tatctgcgct ctgctgaagc cagttacctt
 7051 cgaaaaaaaga gttggtagct cttgatccgg caaacaacc accgctggta
 7101 gcggtggttt ttttgggc aaggcagcaga ttacgcgcag aaaaaaaggaa
 7151 tctcaagaag atcctttgat cttttctacg gggctgacg ctcagtggaa
 7201 cgaaaaactca cgttaaggga ttttggtcat gacattaacc tataaaaata
 7251 ggcgtatcac gaggcccttt cgtctcgccg gtttcggta tgacggtgaa
 7301 aacctctgac acatgcagct cccggagacg gtcacagctt gtctgtaaagc
 7351 ggtatgccggg agcagacaag cccgtcaggg cgcgtcagcg ggtgttggcg
 7401 ggtgtcgaaaa ctggcttaac tatgcggcat cagagcagat tgtactgaga
 7451 gtgcaccata tatgcgggtgt gaaataccgc acagatgcgt aaggagaaaa
 7501 taccgcata ggcgcattc gccattcagg ctgcgcaact gttggaaagg
 7551 gcgatcggtg cgggccttt cgctattacg cca

Table 12: Polypeptide Sequences of FIX

FIX-Fc Monomer Hybrid: created by coexpressing FIX-Fc and Fc chains.

A. FIX-Fc chain (SEQ ID NO: 2)

The c-terminal lysine is not present in either subunit; this processing is often observed in recombinant proteins produced in mammalian cell culture, as well as with plasma derived proteins.

FIX-Fc-SC Subunit (the Fc part of FIX-Fc is in bold):

1 YNSGKLEEFV QGNLERECME EKCSFEEARE VFENTERTTE FWKQYVDGDQ
 51 CESNPCLNGG SCKDDINSYE CWCPFGFEGK NCELDVTTCNI KNGRCEQFCK
 101 NSADNKVVCS CTEGYRLAEN QKSCEPAVPF PCGRVSVSQT SKLTRAETVF
 151 PDVDYVNSTE AETILDNITQ STQSFNDFTR VVGGEDAKPG QFPWQVVLNG
 201 KVDAFCGGSI VNEKWIVTAA HCVETGVKIT VVAGEHNIEE TEHTEQKRNW
 251 IRIIPHHHYN AAINKYNHDI ALLELDEPLV LNSYVTPICI ADKEYTNIFL
 301 KFGSGYVSGW GRVFHKGRSA LVLQYLRVPL VDRATCLRST KFTIYNNMFC

- 55 -

351 AGFHEGGRDS CQGDSGGPHV TEVEGTSFLT GIISWGEECA MKGKYGIYTK
401 VSRYVNWIKE KTKLTDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR
451 TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPREEQ YNSTYRVVSV
501 LTVLHQDWLN GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR
551 DELTKNQVSL TCLVKGFYPS DIAVEWESNG QPENNYKTTP PVLDSDGSFF
601 LYSKLTVDKS RWQQGNVFSC SVMHEALHNN YTQKSLSLSP GK

Table 13: Polynucleotide Sequences of Fc

Fc DNA sequence (mouse Ig κ signal peptide underlined) (SEQ ID NO:3, which encodes SEQ ID NO:4)

1 atggagacag acacactcct gctatggta ctgctgctct gggttccagg
51 ttccactggt gacaaaactc acacatgccc accgtgccc gcacctgaac
101 tcctgggagg accgtcagtc ttccctttcc ccccaaaacc caaggacacc
151 ctcatgatct cccggacccc tgaggtcaca tgcgtggtgg tggacgtgag
201 ccacgaagac cctgaggtca agttcaactg gtacgtggac ggcgtggagg
251 tgcataatgc caagacaaag ccgcgggagg agcagtacaa cagcacgtac
301 cgtgtggtca gcgtcctcac cgtcctgcac caggactggc tgaatggcaa
351 ggagtacaag tgcaaggtct ccaacaaagc cctcccgcc cccatcgaga
401 aaaccatctc caaagccaaa gggcagcccc gagaaccaca ggtgtacacc
451 ctgccccat cccgcgatga gctgaccaag aaccaggtca gcctgacctg
501 cctggtcaaa ggcttctatc ccagcgacat cgccgtggag tggagagca
551 atgggcagcc ggagaacaac tacaagacca cgcctccgt gttggactcc
601 gacggctcct tcttcctcta cagcaagctc accgtggaca agagcaggtg
651 gcagcagggg aacgtttctt catgctccgt gatgcattgag gctctgcaca
701 accactacac gcagaagagc ctctccctgt ctccggtaa a

- 56 -

Table 14: Polypeptide Sequences of Fc

Fc chain (SEQ ID NO:4)

1 DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSCHED
51 PEVKFNWYVD GVEVHNNAKTK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK
101 CKVSNKALPA PIEKTISKAK GQPREPQVYT LPPSRDELTK NQVSLTCLVK
151 GFYPSDIAVE WESNGQPENN YKTTPPVLDs DGSFFLYSKL TVDKSRWQQG
201 NVFSCSVMHE ALHNHYTQKS LSLSPGK

[0191] Any reference to publications cited in this specification is not an admission that the disclosures constitute common general knowledge in Australia.

[0192] Definitions of the specific embodiments of the invention as claimed herein follow.

[0193] According to a first embodiment of the invention, there is provided a pre-lyophilization formulation comprising:

- (a) a chimeric Factor IX (FIX) polypeptide comprising FIX and a heterologous moiety that extends the half-life of the FIX, at a concentration between 80 IU/mL and 2,750 IU/mL;
- (b) L-histidine at a concentration between 3 mg/mL and 15 mg/mL;
- (c) sucrose at a concentration between 10 mg/mL and 50 mg/mL;
- (d) mannitol at a concentration between 20 mg/mL and 100 mg/mL; and
- (e) polysorbate 20 at a concentration between 0.01 mg/mL and 5 mg/mL,
wherein the formulation has a fill volume between 2 mL and 3 mL.

[0194] According to a second embodiment of the invention, there is provided a pre-lyophilization formulation comprising:

- (a) a chimeric Factor IX (FIX) polypeptide, comprising the FIXFc-sc polypeptide of SEQ ID NO:2 and the Fc sc polypeptide of SEQ ID NO:4, at a concentration between about 80 IU/mL and about 2,750 IU/mL;
- (b) L-histidine at a concentration of about 7.76 mg/mL;
- (c) sucrose at a concentration of about 23.8 mg/mL;
- (d) mannitol at a concentration of about 47.6 mg/mL; and
- (e) polysorbate 20 at a concentration of about 0.2 mg/mL,
wherein the formulation has a fill volume of about 2.65 mL.

[0195] According to a third embodiment of the invention, there is provided a pre-lyophilization formulation comprising:

- (a) a chimeric Factor IX (FIX) polypeptide, comprising the FIXFc-sc polypeptide of SEQ ID NO: 2, without the C-terminal lysine, and the Fc sc polypeptide of SEQ ID NO:4, without the C-terminal lysine, wherein the FIXFc-sc polypeptide and the Fc sc polypeptide are bound together through two disulfide bonds in the hinge region of Fc, at a concentration between about 80 IU/mL and about 2,750 IU/mL;
- (b) L-histidine at a concentration of about 7.76 mg/mL;

2015236340 20 Jan 2020

- (c) sucrose at a concentration of about 23.8 mg/mL;
- (d) mannitol at a concentration of about 47.6 mg/mL; and
- (e) polysorbate 20 at a concentration of about 0.2 mg/mL,
wherein the formulation has a fill volume of about 2.65 mL.

WHAT IS CLAIMED IS:

1. A pre-lyophilization formulation comprising:
 - (a) a chimeric Factor IX (FIX) polypeptide comprising FIX and a heterologous moiety that extends the half-life of the FIX, at a concentration between 80 IU/mL and 2,750 IU/mL;
 - (b) L-histidine at a concentration between 3 mg/mL and 15 mg/mL;
 - (c) sucrose at a concentration between 10 mg/mL and 50 mg/mL;
 - (d) mannitol at a concentration between 20 mg/mL and 100 mg/mL; and
 - (e) polysorbate 20 at a concentration between 0.01 mg/mL and 5 mg/mL,
wherein the formulation has a fill volume between 2 mL and 3 mL.
2. The pre-lyophilization formulation of claim 1, wherein the fill volume is about 2.65 mL.
3. The pre-lyophilization formulation of claim 1, wherein the fill volume is about 2.5 mL.
4. The pre-lyophilization formulation of any one of claims 1 to 3,
wherein the L-histidine is present at a concentration of about 7.76 mg/mL;
wherein the sucrose is present at a concentration of about 23.8 mg/mL;
wherein the mannitol is present at a concentration of about 47.6 mg/mL; and
wherein the polysorbate 20 is present at a concentration of about 0.2 mg/mL.
5. The pre-lyophilization formulation of any one of claims 1 to 4, wherein the moiety
extending half-life of FIX is Fc.
6. The pre-lyophilization formulation of any one of claim 1 to 5, wherein the chimeric FIX
polypeptide comprises the FIXFc-sc polypeptide of SEQ ID NO: 2 and the Fc sc polypeptide of
SEQ ID NO:4.
7. A pre-lyophilization formulation comprising:
 - (a) a chimeric Factor IX (FIX) polypeptide, comprising the FIXFc-sc polypeptide of
SEQ ID NO:2 and the Fc sc polypeptide of SEQ ID NO:4, at a concentration between about 80
IU/mL and about 2,750 IU/mL;
 - (b) L-histidine at a concentration of about 7.76 mg/mL;
 - (c) sucrose at a concentration of about 23.8 mg/mL;

(d) mannitol at a concentration of about 47.6 mg/mL; and
(e) polysorbate 20 at a concentration of about 0.2 mg/mL,
wherein the formulation has a fill volume of about 2.65 mL.

8. The pre-lyophilization formulation of claim 1, wherein the chimeric FIX polypeptide comprises the FIXFc-sc polypeptide of SEQ ID NO: 2, without the C-terminal lysine, and the Fc sc polypeptide of SEQ ID NO:4, without the C-terminal lysine, wherein the FIXFc-sc polypeptide and the Fc sc polypeptide are bound together through two disulfide bonds in the hinge region of Fc.

9. A pre-lyophilization formulation comprising:

(a) a chimeric Factor IX (FIX) polypeptide, comprising the FIXFc-sc polypeptide of SEQ ID NO: 2, without the C-terminal lysine, and the Fc sc polypeptide of SEQ ID NO:4, without the C-terminal lysine, wherein the FIXFc-sc polypeptide and the Fc sc polypeptide are bound together through two disulfide bonds in the hinge region of Fc, at a concentration between about 80 IU/mL and about 2,750 IU/mL;

(b) L-histidine at a concentration of about 7.76 mg/mL;

(c) sucrose at a concentration of about 23.8 mg/mL;

(d) mannitol at a concentration of about 47.6 mg/mL; and

(e) polysorbate 20 at a concentration of about 0.2 mg/mL,
wherein the formulation has a fill volume of about 2.65 mL.

1/5




Fig. 1

2/5




Fig. 2

3/5




Fig. 3

4/5





Fig. 4

5/5



5

2159397PC01_SequenceListing.txt
SEQUENCE LISTING

<110> BIOGEN IDEC MA INC.
Thome, Brian M.
Parkhurst-Lang, Cherie
Leveille, Brandon W.

<120> LYOPHILIZED FACTOR IX FORMULATIONS

<130> 2159.397PC01/C-K/E-H

<140> To be filed
<141> Herewith

<150> 61/969,801
<151> 2014-03-24

<160> 14

<170> PatentIn version 3.5

<210> 1
<211> 7583
<212> DNA
<213> Artificial Sequence

<220>
<223> FIX-Fc Chain

<400> 1
gcgcgcttg acattgatta ttgactagtt attaatagta atcaattacg gggtcattag 60
ttcatacgccc atatatggag ttccgcgtta cataacttac ggttaatggc ccgcctggct 120
gaccgccccaa cgaccccccgc ccattgacgt caataatgac gtatgttccc atagtaacgc 180
caatagggac tttccattga cgtcaatggg tggagtattt acggtaaact gcccacttgg 240
cagtacatca agtgtatcat atgccaagta cgccccctat tgacgtcaat gacggtaaat 300
ggccgcctg gcattatgcc cagtagatga ccttatggga ctttcctact tggcagttaca 360
tctacgtatt agtcatcgct attaccatgg tgatgcgggtt ttggcagttac atcaatgggc 420
gtggatagcg gtttgactca cggggatttc caagtctcca ccccattgac gtcaatggga 480
gtttgttttg gcacccaaat caacgggact ttccaaaatg tcgtaacaac tccgccttcat 540
tgacgcaaat gggcggtagg cgtgtacggt gggaggtcta tataaggcaga gctctctggc 600
taactagaga acccactgct tactggctta tcgaaattaa tacgactcac tatagggaga 660
cccaagcttc gcgacgtacg gccgcacca tgcagcgcgt gaacatgatc atggcagaat 720
caccaggcct catcaccatc tgccttttag gatatctact cagtgctgaa tgtacagggtt 780
tgtttccttt tttaaaatac attgagttatg cttgcctttt agatataaaaa atatctgtatg 840
ctgtcttctt cactaaattt tgattacatg atttgacagc aatattgaag agtctaacag 900
ccagcacgca ggttggtaag tactgtggga acatcacaga tttggctcc atgccctaaa 960
gagaaattgg ctttcagatt attggatta aaaacaaaga ctttcttaag agatgtaaaa 1020
ttttcatgat gttttctttt ttgctaaaac taaagaatta ttctttaca tttcagtttt 1080
tcttgatcat gaaaacgcca acaaattct gaatcggcca aagaggtata attcaggtaa 1140

2159397PC01_SequenceListing.txt

attggaagag	tttgttcaag	ggaatctaga	gagagaatgt	atggaagaaa	agtgttagtt	1200
tgaagaagca	cgagaagttt	ttgaaaacac	tgaaagaaca	actgaatttt	ggaagcagta	1260
tgttcatgga	gatcagtgtg	agtccaatcc	atgtttaaat	ggcggcagtt	gcaaggatga	1320
cattaattcc	tatgaatgtt	ggtgtccctt	tggattgaa	gaaagaact	gtgaattaga	1380
tgttaacatgt	aacattaaga	atggcagatg	cgagcagttt	tgtaaaaata	gtgctgataa	1440
caagggtggtt	tgctcctgta	ctgagggata	tcgacttgca	gaaaaccaga	agtcctgtga	1500
accagcagt	ccatttccat	gtggaagagt	ttctgttca	caaacttcta	agtcacccg	1560
tgctgagact	gttttcctg	atgtggacta	tgtaaattct	actgaagctg	aaaccatttt	1620
ggataaacatc	actcaaagca	cccaatcatt	taatgacttc	actcgggttt	ttgggtggaga	1680
agatgccaaa	ccaggtcaat	tccctggca	ggttgtttt	aatggtaaag	ttgatgcatt	1740
ctgtggaggc	tctatcgta	atgaaaaatg	gattgtaact	gctgccact	gtgttgaac	1800
tgggtttaaa	attacagttt	tcgcaggtga	acataatatt	gaggagacag	aacatacaga	1860
gcaaaagcga	aatgtgattc	gaattattcc	tcaccacaac	tacaatgcag	ctattaataa	1920
gtacaaccat	gacattgccc	ttctggaact	ggacgaaccc	ttagtgctaa	acagctacgt	1980
tacacctatt	tgcattgctg	acaaggaata	cacgaacatc	ttcctcaa	ttggatctgg	2040
ctatgttgt	ggctggggaa	gagtcttcca	caaaggaga	tcagctttag	ttcttcagta	2100
ccttagagtt	ccacttgg	accgagccac	atgtctcga	tctacaaagt	tcaccatcta	2160
taacaacatg	ttctgtgt	gctccatga	aggaggtaga	gattcatgtc	aaggagatag	2220
tgggggaccc	catgttactg	aagtggagg	gaccagttt	ttaactggaa	ttattagctg	2280
gggtgaagag	tgtcaatga	aaggcaata	tggaatata	accaagggt	cccggtatgt	2340
caactggatt	aaggaaaaaa	caaagctcac	tgacaaaact	cacacatgcc	caccgtgccc	2400
agctccggaa	ctcctggcg	gaccgtcagt	cttccttcc	cccccaaaac	ccaaggacac	2460
cctcatgatc	tcccggaccc	ctgaggtcac	atgcgtgg	gtggacgtga	gccacgaaga	2520
ccctgagg	aagttcaact	ggtacgtgg	cggcgtgg	gtgcataat	ccaagacaaa	2580
gccgcggag	gagcagtaca	acagcacgt	ccgtgtgg	agcgtcctca	ccgtcctgca	2640
ccaggactgg	ctgaatggca	aggagtaca	gtgcaagg	tccaacaa	ccctcccagc	2700
ccccatcgag	aaaaccatct	ccaaagccaa	agggcagccc	cgagaaccac	aggtgtacac	2760
cctgccccca	tcccggatg	agctgacca	gaaccagg	agcctgac	gcctggtaa	2820
aggcttctat	cccagcgaca	tcgcccgtg	gtggagagc	aatgggcagc	cggagaacaa	2880
ctacaagacc	acgcctcccg	tgttggactc	cgacggctcc	ttcttcctct	acagcaagct	2940
caccgtggac	aagagcagg	ggcagcagg	gaacgttcc	tcatgctccg	tgtatgcata	3000
ggctctgcac	aaccactaca	cgcagaagag	cctctccctg	tctccggta	aatgagaatt	3060
cagacatgt	aagatacatt	gatgagttt	gacaaaccac	aactagaatg	cagtaaaaaa	3120
aatgcttat	ttgtgaaatt	tgtatgcata	ttgcttatt	tgtatgcata	ataagctgca	3180

2159397PC01_SequenceListing.txt		
ataaaacaagt	tgggtggc	3240
cagccggcgt	cccggaaaac	3300
tcgaaatctc	gtagcacgtg	3360
gccgggtcgc	gcagggcgaa	3420
ggccgggagg	cgtcccgaa	3480
tccaggccgc	gcacccacac	3540
gcgctgatga	acagggtcac	3600
tcccgggaga	acccgagccg	3660
tgagcaccg	gaacggcact	3720
tatactatgc	cgatatacta	3780
aaatggatat	acaagctccc	3840
ctcactactt	ctggaatagc	3900
aaattagtca	gccatggggc	3960
gcggagttag	gggcgggact	4020
tgcctgctgg	ggagcctggg	4080
tttgactact	tctgctgct	4140
tcgtgaggct	ccggtgcccg	4200
ttgggggag	gggtcgccaa	4260
gaaagtgtat	ttgaaccgg	4320
agtgcagtag	tcgtgtactg	4380
agtgcgtgt	gtggttcccg	4440
tgaattactt	ccacctggct	4500
ggccttgcgc	tttaggagcc	4560
ggggccgccc	cgtgcgaatc	4620
ctctagccat	ttaaaattt	4680
ttgtaaatgc	gggccaggat	4740
acggggcccg	tgcgtccag	4800
cgagaatcgg	cgcacatgtt	4860
cgccgtgtat	ccccccccc	4920
cggaaagatg	gccgcttccc	4980
cgggagagcg	ggcgggtgag	5040
tcgcttcatg	tgactccacg	5100
gctttggag	gagtaccgg	5160
acactgagtg	ggtggagact	5220
gaagtttaggc	cagttggca	
cttgatgtaa		
ttctcccttgg		

2159397PC01_SequenceListing.txt

aatttgcctt	ttttgagttt	ggatcttggt	tcattctcaa	gcctcagaca	gtggttcaaa	5280
gttttttct	tccatttcag	gtgtcgtgaa	cacgtggtcg	cggccgcgccc	gccaccatgg	5340
agacagacac	actcctgcta	tgggtactgc	tgctctgggt	tccaggttcc	actggtgaca	5400
aaactcacac	atgcccaccg	tgcccagcac	ctgaactcct	gggaggaccg	tcagtcttcc	5460
tcttcccccc	aaaacccaag	gacaccctca	tgatctcccg	gaccctgag	gtcacatgcg	5520
tgggtggtgg	cgtgagccac	gaagaccctg	aggtcaagtt	caactggtac	gtggacggcg	5580
tggaggtgca	taatgccaag	acaaagccgc	gggaggagca	gtacaacacgc	acgtaccgtg	5640
tggtcagcgt	cctcaccgtc	ctgcaccagg	actggctgaa	tggcaaggag	tacaagtgca	5700
aggtctccaa	caaagccctc	ccagccccca	tcgagaaaac	catctccaaa	gccaaaggc	5760
agccccgaga	accacaggtg	tacaccctgc	ccccatcccg	cgatgagctg	accaagaacc	5820
aggtcagcct	gacctgcctg	gtcaaaggct	tctatcccag	cgacatcgcc	gtggagtgg	5880
agagcaatgg	gcagccggag	aacaactaca	agaccacgcc	tcccgtgtt	gactccgacg	5940
gctccttctt	cctctacagc	aagctcaccg	tggacaagag	caggtggcag	cagggaaacg	6000
tcttctcatg	ctccgtgatg	catgaggctc	tgcacaacca	ctacacgcag	aagagcctct	6060
ccctgtctcc	ggtaaatga	ctcgagagat	ctggccggct	gggcccgtt	cgaaggtaag	6120
cctatcccta	accctctcct	cggtctcgat	tctacgcgt	ccggcatca	tcaccatcac	6180
cattgagttt	aaaccgcgt	atcagcctcg	actgtgcctt	ctagttgcca	gccatctgtt	6240
gtttgcccct	cccccggtgcc	ttccttgacc	ctggaaaggt	ccactccac	tgtccttcc	6300
taataaaatg	aggaaattgc	atcgattgt	ctgagtaggt	gtcattctat	tctgggggt	6360
ggggtggggc	aggacagcaa	gggggaggat	tggaaagaca	atagcaggca	tgctggggat	6420
gcgggtggct	ctatggcttc	tgaggcggaa	agaaccagt	gcggtaatac	ggttatccac	6480
agaatcaggg	gataacgcag	gaaagaacat	gtgagcaaaa	ggccagcaaa	aggccaggaa	6540
ccgtaaaaag	gccgcgttgc	tggcgcccc	ccataggctc	cgccccctg	acgagcatca	6600
caaaaatcga	cgctcaagtc	agaggtggcg	aaaccgcaca	ggactataaa	gataccaggc	6660
gtttccccct	agaagctccc	tcgtgcgtc	tcctgttccg	accctgccgc	ttaccggata	6720
cctgtccgccc	tttctccctt	cggaaagcgt	ggcgcttct	catagctcac	gctgttaggt	6780
tctcagttcg	gtgttaggtcg	ttcgctccaa	gctgggtgt	gtgcacgaac	cccccggtca	6840
gcccgaccgc	tgcgccttat	ccggtaacta	tcgtcttgag	tccaaccgg	taagacacga	6900
cttatacgcca	ctggcagcag	ccactggtaa	caggattagc	agagcgaggt	atgttaggcgg	6960
tgctacagag	ttcttgaagt	ggtggctaa	ctacggctac	actagaagaa	cagtatttgg	7020
tatctgcgct	ctgctgaagc	cagttacctt	cggaaaaaga	gttggtagct	cttgatccgg	7080
caaacaaacc	accgctggta	gcggtggtt	ttttgttgc	aagcagcaga	ttacgcgcag	7140
aaaaaaaaagga	tctcaagaag	atcctttgat	cttttctacg	gggtctgacg	ctcagtggaa	7200
cgaaaaactca	cgttaaggga	ttttggtcat	gacattaacc	tataaaaata	ggcgtatcac	7260

2159397PC01_SequenceListing.txt

gaggcccttt	cgtctcgcbc	gtttcggta	tgacggtgaa	aacctctgac	acatgcagct	7320
cccggagacg	gtcacagctt	gtctgttaagc	ggatgccggg	agcagacaag	cccgtcaggg	7380
cgcgtcagcg	ggtgttggcg	ggtgtcgggg	ctggcttaac	tatgcggcat	cagagcagat	7440
tgtactgaga	gtgcaccata	tatgcggtgt	gaaataccgc	acagatgcgt	aaggagaaaa	7500
tacccatca	ggcgcattc	gccattcagg	ctgcgcaact	gttggaaagg	gcgatcggtg	7560
cggcccttt	cgctattacg	cca				7583

<210> 2
<211> 642
<212> PRT
<213> Artificial Sequence

<220>
<223> FIX-Fc Chain

<400> 2

Tyr Asn Ser Gly Lys Leu Glu Glu Phe Val Gln Gly Asn Leu Glu Arg
1 5 10 15

Glu Cys Met Glu Glu Lys Cys Ser Phe Glu Glu Ala Arg Glu Val Phe
20 25 30

Glu Asn Thr Glu Arg Thr Thr Glu Phe Trp Lys Gln Tyr Val Asp Gly
35 40 45

Asp Gln Cys Glu Ser Asn Pro Cys Leu Asn Gly Gly Ser Cys Lys Asp
50 55 60

Asp Ile Asn Ser Tyr Glu Cys Trp Cys Pro Phe Gly Phe Glu Gly Lys
65 70 75 80

Asn Cys Glu Leu Asp Val Thr Cys Asn Ile Lys Asn Gly Arg Cys Glu
85 90 95

Gln Phe Cys Lys Asn Ser Ala Asp Asn Lys Val Val Cys Ser Cys Thr
100 105 110

Glu Gly Tyr Arg Leu Ala Glu Asn Gln Lys Ser Cys Glu Pro Ala Val
115 120 125

Pro Phe Pro Cys Gly Arg Val Ser Val Ser Gln Thr Ser Lys Leu Thr
130 135 140

Arg Ala Glu Thr Val Phe Pro Asp Val Asp Tyr Val Asn Ser Thr Glu
145 150 155 160

Ala Glu Thr Ile Leu Asp Asn Ile Thr Gln Ser Thr Gln Ser Phe Asn
165 170 175

Asp Phe Thr Arg Val Val Gly Gly Glu Asp Ala Lys Pro Gly Gln Phe
Page 5

180

185

190

Pro Trp Gln Val Val Leu Asn Gly Lys Val Asp Ala Phe Cys Gly Gly
 195 200 205

Ser Ile Val Asn Glu Lys Trp Ile Val Thr Ala Ala His Cys Val Glu
 210 215 220

Thr Gly Val Lys Ile Thr Val Val Ala Gly Glu His Asn Ile Glu Glu
 225 230 235 240

Thr Glu His Thr Glu Gln Lys Arg Asn Val Ile Arg Ile Ile Pro His
 245 250 255

His Asn Tyr Asn Ala Ala Ile Asn Lys Tyr Asn His Asp Ile Ala Leu
 260 265 270

Leu Glu Leu Asp Glu Pro Leu Val Leu Asn Ser Tyr Val Thr Pro Ile
 275 280 285

Cys Ile Ala Asp Lys Glu Tyr Thr Asn Ile Phe Leu Lys Phe Gly Ser
 290 295 300

Gly Tyr Val Ser Gly Trp Gly Arg Val Phe His Lys Gly Arg Ser Ala
 305 310 315 320

Leu Val Leu Gln Tyr Leu Arg Val Pro Leu Val Asp Arg Ala Thr Cys
 325 330 335

Leu Arg Ser Thr Lys Phe Thr Ile Tyr Asn Asn Met Phe Cys Ala Gly
 340 345 350

Phe His Glu Gly Gly Arg Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro
 355 360 365

His Val Thr Glu Val Glu Gly Thr Ser Phe Leu Thr Gly Ile Ile Ser
 370 375 380

Trp Gly Glu Glu Cys Ala Met Lys Gly Lys Tyr Gly Ile Tyr Thr Lys
 385 390 395 400

Val Ser Arg Tyr Val Asn Trp Ile Lys Glu Lys Thr Lys Leu Thr Asp
 405 410 415

Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly
 420 425 430

Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile
 435 440 445

Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu
 Page 6

2159397PC01_SequenceListing.txt
450 455 460

Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His
465 470 475 480

Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg
485 490 495

Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys
500 505 510

Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu
515 520 525

Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr
530 535 540

Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu
545 550 555 560

Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp
565 570 575

Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val
580 585 590

Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp
595 600 605

Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
610 615 620

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro
625 630 635 640

Gly Lys

<210> 3
<211> 741
<212> DNA
<213> Artificial Sequence

<220>
<223> Fc Portion

<400> 3
atggagacag acacactcct gctatggta ctgctgtct gggttccagg ttccactgg
60
gacaaaactc acacatgccc accgtgccca gcacctgaac tcctgggagg accgtcagtc
120
ttcctcttcc ccccaaaacc caaggacacc ctcatgatct cccggacccc tgaggtcaca
180
tgcgtggagg tggacgtgag ccacgaagac cctgaggta agttcaactg gtacgtggac
240
ggcgtggagg tgcataatgc caagacaaag ccgcgggagg agcagtacaa cagcacgtac
300

2159397PC01_SequenceListing.txt

cgtgtggtca	cggtcctcac	cgtcctgcac	caggactggc	tgaatggcaa	ggagtacaag	360
tgcaaggct	ccaacaaagc	cctcccgacc	cccatcgaga	aaaccatctc	caaagccaaa	420
gggcagcccc	gagaaccaca	ggtgtacacc	ctgccccat	cccgcgatga	gctgaccaag	480
aaccaggta	gcctgacctg	cctggtaaaa	ggcttctatc	ccagcgacat	cgccgtggag	540
tgggagagca	atggcagcc	ggagaacaac	tacaagacca	cgcccccgt	gttggactcc	600
gacggctcct	tcttcctcta	cagcaagctc	accgtggaca	agagcaggtg	gcagcagggg	660
aacgtttct	catgctccgt	gatgcatgag	gctctgcaca	accactacac	gcagaagagc	720
ctctccctgt	ctccggtaa	a				741

<210> 4
<211> 227
<212> PRT
<213> Artificial Sequence

<220>
<223> Fc Portion

<400> 4

Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly
1 5 10 15

Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met
20 25 30

Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His
35 40 45

Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val
50 55 60

His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr
65 70 75 80

Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly
85 90 95

Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile
100 105 110

Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val
115 120 125

Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser
130 135 140

Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu
145 150 155 160

Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro
 165 170 175

Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val
 180 185 190

Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met
 195 200 205

His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
 210 215 220

Pro Gly Lys
 225

<210> 5
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Core sequence of albumin-binding peptides

<400> 5

Asp Ile Cys Leu Pro Arg Trp Gly Cys Leu Trp
 1 5 10

<210> 6
 <211> 32
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CTP peptide

<400> 6

Asp Pro Arg Phe Gln Asp Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser
 1 5 10 15

Leu Pro Ser Pro Ser Arg Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu
 20 25 30

<210> 7
 <211> 28
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> CTP peptide

<400> 7

Ser Ser Ser Ser Lys Ala Pro Pro Pro Ser Leu Pro Ser Pro Ser Arg
 1 5 10 15

Leu Pro Gly Pro Ser Asp Thr Pro Ile Leu Pro Gln
 20 25

2159397PC01_SequenceListing.txt

<210> 8
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 8

Ala Ser Pro Ala Ala Pro Ala Pro Ala Ser Pro Ala Ala Pro Ala Pro
1 5 10 15

Ser Ala Pro Ala
20

<210> 9
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 9

Ala Ala Pro Ala Ser Pro Ala Pro Ala Ala Pro Ser Ala Pro Ala Pro
1 5 10 15

Ala Ala Pro Ser
20

<210> 10
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 10

Ala Pro Ser Ser Pro Ser Pro Ser Ala Pro Ser Ser Pro Ser Pro Ala
1 5 10 15

Ser Pro Ser Ser
20

<210> 11
<211> 19
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 11

Ala Pro Ser Ser Pro Ser Pro Ser Ala Pro Ser Ser Pro Ser Pro Ala
1 5 10 15

2159397PC01_SequenceListing.txt

Ser Pro Ser

<210> 12
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 12

Ser Ser Pro Ser Ala Pro Ser Pro Ser Ser Pro Ala Ser Pro Ser Pro
1 5 10 15

Ser Ser Pro Ala
20

<210> 13
<211> 24
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 13

Ala Ala Ser Pro Ala Ala Pro Ser Ala Pro Pro Ala Ala Ala Ser Pro
1 5 10 15

Ala Ala Pro Ser Ala Pro Pro Ala
20

<210> 14
<211> 20
<212> PRT
<213> Artificial Sequence

<220>
<223> PAS peptide

<400> 14

Ala Ser Ala Ala Ala Pro Ala Ala Ala Ser Ala Ala Ala Ser Ala Pro
1 5 10 15

Ser Ala Ala Ala
20