



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 9/06	A1	(11) International Publication Number: WO 98/10746 (43) International Publication Date: 19 March 1998 (19.03.98)
(21) International Application Number: PCT/US97/15919 (22) International Filing Date: 10 September 1997 (10.09.97) (30) Priority Data: 08/712,454 11 September 1996 (11.09.96) US (71) Applicant: VIROTEX CORPORATION [US/US]; Suite 350, 4200 Research Forest Drive, The Woodlands, TX 77381 (US). (72) Inventor: OSBORNE, David, W.; 19 Quiet Oak Crescent, The Woodlands, TX 77381 (US). (74) Agent: PORTER, Gregory, L.; Arnold, White & Durkee, P.O. Box 4433, Houston, TX 77210 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: COMPOSITIONS AND METHODS FOR TOPICAL APPLICATION OF THERAPEUTIC AGENTS (57) Abstract The present invention generally relates to pharmaceutical compositions that enable control of drug delivery properties and the development of optimal drug delivery strategies customized for particular drugs and particular diseases. The composition includes a dissolved pharmaceutical that has the capacity to permeate the stratum corneum layer of the epidermis and become available systemically, and pharmaceutical in a microparticulate state that does not readily cross the stratum corneum of the epidermis. The dissolved and microparticulate pharmaceuticals may be the same or different pharmaceuticals. Methods for the preparation and use of the compositions are also provided. In a preferred embodiment, the invention finds particular use in a formulation for the topical application of dapsone for the treatment of acne. In another preferred embodiment, the invention finds particular use for the treatment of herpes lesions.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

COMPOSITIONS AND METHODS FOR TOPICAL APPLICATION OF THERAPEUTIC AGENTS

FIELD OF THE INVENTION

5 The present invention relates to novel dermatological compositions that exhibit readily optimized solubility and systemic drug delivery properties for applying drugs and therapeutic agents to the skin of humans and animals and methods for their preparation and use.

BACKGROUND OF THE INVENTION

10 While the skin has long been considered the preferred route of administration for cosmetic applications and dermatological therapies, the introduction of transdermal nitroglycerin patches initiated use of the skin as a route for administering systemic drug therapy. Three types of known product applications which employ the barrier properties of the skin for drug delivery include cosmetic, topical, and transdermal applications. The
15 optimal delivery strategy for administering pharmaceuticals via the skin varies among individual pharmaceuticals and among different disease states.

 Cosmetic applications are limited to negligible drug penetration past the stratum corneum. Thus, any carrier that minimizes penetration or that aids excipient retention within or onto the stratum corneum would be of tremendous advantage. For transdermal
20 applications, steady state drug delivery is preferred. Steady state delivery requires the use of rate-controlling membranes that slow systemic breakthrough of highly permeable drugs such as nitroglycerin. This type of control can be achieved by using matrix type patches that modify delivery rates by using polymer adhesives and solvents. For topical delivery, minimal

- 2 -

systemic breakthrough is always preferred. In order to adequately dose the viable epidermis and dermis, however, large amounts of drug must cross the intact skin barrier, i.e. the stratum corneum, or the lesional delivery barrier, i.e. scab, plaque, etc.

Some dermatological conditions, such as acne, require multiple delivery strategies because they have multiple delivery requirements. Acne is chronic pilosebaceous unit inflammation associated with the face and trunk usually occurring in adolescence due to complex interactions of androgens and bacteria. For the adolescent, circulating androgen results in significantly increased sebum production. The sebaceous glands dramatically enlarge and excrete more sebum than the immature pilosebaceous canals can accommodate. Simultaneously, anaerobic bacteria (*Propionibacterium acnes*) that feed upon the sebum, converting triglycerides to fatty acids, dramatically increase in number due to an increase in volume of the nutrition source. The increase in constricted immature ducts and bacterial waste products results in plugged follicles and typical acne inflammation. Acne severity for a particular anatomical location parallels the number of sebaceous glands per unit of skin.

Acne, which is often treated with antibiotics, is one condition where a highly specialized topical drug delivery is needed. Ideally, a topical antimicrobial would be primarily delivered into the pilosebaceous unit, with only minimal active crossing of the skin barrier. Intact stratum corneum lines the upper third of the pilosebaceous unit, and it is into this upper third of the hair follicle that the sebaceous duct secretes sebum. Thus, a need exists for an acne treatment that maximizes antimicrobial drug levels in the upper third of the pilosebaceous unit.

- 3 -

Additionally, when an anti-inflammatory agent is used to treat acne, it is important to increase the level of drug that will cross the intact stratum corneum lining the upper third of the pilosebaceous unit. By definition, inflammation is the response of the viable epidermis to irritants and sensitizers. In order to reduce the amount of inflammation, the active
5 pharmaceutical must penetrate past the stratum corneum and interfere with the cascade of inflammatory events. Ideally, delivery of an anti-inflammatory for acne requires that steady-state levels be sustained. To date, the ideal delivery system that provides antimicrobial agents above the stratum corneum while providing anti-inflammatory agents below the stratum corneum has not been implemented.

10 Other dermatological conditions, such as herpes lesions, require multiple delivery strategies because the barrier properties of the lesion dramatically change in the course of the disease. Starting with the prodrome and progressing through the formation of vesicles, the lesion has an intact stratum corneum delivery barrier, and thus, maximum penetration of the drug is necessary. While in place, the stratum corneum delays penetration to the target tissue
15 and sustains the time that the dissolved active drug resides in the target tissue. During this stage of the lesion, microparticulate drug will not significantly cross the intact stratum corneum, and thus, has no real effect in treatment of the lesion. Once the herpes lesion vesicles rupture, the stratum corneum is no longer in place, and the dissolved drug rapidly sweeps past the target tissue, providing minimal or insignificant benefit. However, from the
20 time that the vesicle ruptures and through to the complete formation of the scab, the microparticulate drug is deposited directly at the target area, where it can slowly be released for sustained and significant therapeutic benefit. Thus, in order to adequately dose the viable

- 4 -

epidermis from the prodrome through the time of scab formation in a herpes lesion, two distinctly different drug delivery strategies must be implemented.

While the dermatological conditions of acne and herpes lesions serve as conceptual examples of how therapeutic approaches can require dramatically different drug delivery profiles, all skin diseases are best treated by a particular drug delivery strategy tailored specifically to the pharmaceutical and the particular disease. Some diseases are best treated using pulsed or spiked delivery in which high levels of drug are delivered in a short period of time. This type of treatment saturates receptor sites and provides maximum microbial or viral replication inhibition; thus providing optimal therapy for certain diseases. Conversely, a cosmetic, topical, or transdermal product that provides steady state active pharmaceutical delivery while minimizing excipient delivery provides the preferred skin delivery profile for other diseases. Thus, a carrier system that can be adjusted to optimize the delivery profile for the pharmacology of the active drug and the nature of the disease state is needed to advance the effectiveness of pharmaceutical products applied to the skin.

15

SUMMARY OF THE INVENTION

The present invention concerns a pharmaceutical carrier system comprising a dermatological composition that is a semi-solid aqueous gel, wherein a pharmaceutical is dissolved in the gel such that the pharmaceutical has the capacity to cross the stratum corneum layer of the epidermis and become available systemically, and wherein the composition also contains pharmaceutical in a microparticulate state that does not readily cross the stratum corneum of the epidermis. The ratio of microparticulate pharmaceutical to dissolved pharmaceutical is adjustable, but is preferably five or less. The microparticulate

20

- 5 -

pharmaceutical and the dissolved pharmaceutical may be the same drug, or they may be different drugs.

Methods for preparing the compositions of the present invention are also shown. In addition, methods for treating dermatological conditions that include topically applying the dermatological compositions of the invention are shown. More particularly, the invention concerns methods for treating dermatological conditions or diseases such as acne, herpes lesions, and dermatitis. Antimicrobial agents having anti-inflammatory properties such as dapsone are used to treat acne. Antiviral agents or antiviral agents in combination with local anesthetics are used to treat herpes lesions, and anti-inflammatory agents are used to treat dermatitis.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention comprises compositions for application to the skin that can form microparticulate drug precipitates in adjustable ratios of microparticulate drug to dissolved drug, methods for the formation of said compositions, and methods for treatment of skin conditions using said compositions. The advantages of the present invention are appreciated in the treatment of skin conditions or diseases by using cosmetics or topical pharmaceuticals, and in the systemic treatment of illness by using transdermal pharmaceuticals. The present invention is particularly effective in the treatment of acne with antimicrobial actives known to possess anti-inflammatory properties such as dapsone. The invention also finds particular use in the treatment of herpes lesions and dermatitis.

In one embodiment, the present invention is directed to a novel pharmaceutical carrier system comprising a dermatological composition that is a semisolid aqueous gel, wherein the

- 6 -

composition exhibits an optimal balance between a dissolved pharmaceutical that is available to cross through the stratum corneum to become systemically available, and a microparticulate pharmaceutical that is retained in or above the stratum corneum to serve as a reservoir or to provide drug action in the supracorneum zone. The microparticulate pharmaceutical and the dissolved pharmaceutical may be the same or different drugs. The microparticulate pharmaceutical may comprise a crystalline precipitant or an amorphous precipitant.

Optimal balance is accomplished by having a semisolid gel carrier system in which microparticulate pharmaceutical precipitates are formed in reproducible ratios with respect to the dissolved pharmaceutical. For the composition to have a wide range of applicability, the microparticulate to dissolved pharmaceutical ratio preferably should be no greater than five, at therapeutic levels of applied active pharmaceutical.

A composition having a microparticulate to dissolved pharmaceutical ratio of less than two may provide the greatest amount of pharmaceutical available for immediate partition out of the stratum corneum and into the viable epidermis. This should provide minimum reservoir capacity, but may not maintain sustained delivery or provide maximum activity in the supracorneum zone. A composition having a microparticulate to dissolved pharmaceutical ratio of two or greater may have a reduced amount of drug available for immediate partition out of the stratum corneum and into the viable epidermis. This provides maximum reservoir capacity, and maintains sustained delivery, providing maximum activity in the supracorneum zone. For the present invention, the ratio for microparticulate drug to dissolved drug should be no greater than 50, preferably no greater than 10, and most

- 7 -

preferably no greater than 5. Drug delivery from the microparticulate/dissolved pharmaceutical formulation may be optimized to provide higher levels of drug to the supracorneum zone, while maintaining the level of drug partitioning out of the stratum corneum and into the viable epidermis, despite 10-fold increases in the amount of pharmaceutical applied to the skin.

The compositions of the present invention comprise semi-solid and gel-like vehicles that include a polymer thickener, water, preservatives, active surfactants or emulsifiers, antioxidants, sunscreens, and a solvent or mixed solvent system. The solvent or mixed solvent system is important to the formation of the microparticulate to dissolved pharmaceutical ratio. The formation of the microparticulate, however, should not interfere with the ability of the polymer thickener or preservative systems to perform their functions.

Polymer thickeners that may be used include those known to one skilled in the art, such as hydrophilic and hydroalcoholic gelling agents frequently used in the cosmetic and pharmaceutical industries. Preferably, the hydrophilic or hydroalcoholic gelling agent comprises "CARBOPOL®" (B.F. Goodrich, Cleveland, OH), "HYPAN®" (Kingston Technologies, Dayton, NJ), "NATROSOL®" (Aqualon, Wilmington, DE), "KLUCEL®" (Aqualon, Wilmington, DE), or "STABILEZE®" (ISP Technologies, Wayne, NJ). Preferably, the gelling agent comprises between about 0.2% to about 4% by weight of the composition. More particularly, the preferred compositional weight percent range for "CARBOPOL®" is between about 0.5% to about 2%, while the preferred weight percent range for "NATROSOL®" and "KLUCEL®" is between about 0.5% to about 4%. The

- 8 -

preferred compositional weight percent range for both "HYPAN®" and "STABILEZE®" is between about 0.5% to about 4%.

"CARBOPOL®" is one of numerous cross-linked acrylic acid polymers that are given the general adopted name carbomer. These polymers dissolve in water and form a clear or slightly hazy gel upon neutralization with a caustic material such as sodium hydroxide, potassium hydroxide, triethanolamine, or other amine bases. "KLUCCEL®" is a cellulose polymer that is dispersed in water and forms a uniform gel upon complete hydration. Other preferred gelling polymers include hydroxyethylcellulose, cellulose gum, MVE/MA decadiene crosspolymer, PVM/MA copolymer, or a combination thereof.

Preservatives may also be used in this invention and preferably comprise about 0.05% to 0.5% by weight of the total composition. The use of preservatives assures that if the product is microbially contaminated, the formulation will prevent or diminish microorganism growth. Some preservatives useful in this invention include methylparaben, propylparaben, butylparaben, chloroxylenol, sodium benzoate, DMDM Hydantoin, 3-Iodo-2-Propylbutyl carbamate, potassium sorbate, chlorhexidine digluconate, or a combination thereof.

Titanium dioxide may be used as a sunscreen to serve as prophylaxis against photosensitization. Alternative sunscreens include methyl cinnamate. Moreover, BHA may be used as an antioxidant, as well as to protect ethoxydiglycol and/or dapsone from discoloration due to oxidation. An alternate antioxidant is BHT.

Pharmaceuticals for use in all embodiments of the invention include antimicrobial agents, anti-inflammatory agents, antiviral agents, local anesthetic agents, corticosteroids, destructive therapy agents, antifungals, and antiandrogens. In the treatment of acne, active

- 9 -

pharmaceuticals that may be used include antimicrobial agents, especially those having anti-inflammatory properties such as dapsone, erythromycin, minocycline, tetracycline, clindamycin, and other antimicrobials. The preferred weight percentages for the antimicrobials are 0.5% to 10%. In the topical treatment of herpes lesions, active
5 pharmaceuticals that may be used include antiviral or local anesthetic agents. A concentration of about 1.0% to 10% by weight is preferred for nucleoside analogues such as acyclovir, famciclovir, penciclovir, valacyclovir, and ganciclovir.

Local anesthetics include tetracaine, tetracaine hydrochloride, lidocaine, lidocaine hydrochloride, dyclonine, dyclonine hydrochloride, dimethisoquin hydrochloride, dibucaine,
10 dibucaine hydrochloride, butambenpicrate, and pramoxine hydrochloride. A preferred concentration for local anesthetics is about .025% to 5% by weight of the total composition. Anesthetics such as benzocaine may also be used at a preferred concentration of about 2% to 25% by weight.

Corticosteroids that may be used include betamethasone dipropionate, fluocinolone
15 acetonide, betamethasone valerate, triamcinolone acetonide, clobetasol propionate, desoximetasone, diflorasone diacetate, amcinonide, flurandrenolide, hydrocortisone valerate, hydrocortisone butyrate, and desonide are recommended at concentrations of about 0.01% to 1.0% by weight. Preferred concentrations for corticosteroids such as hydrocortisone or methylprednisolone acetate are from about 0.2% to about 5.0% by weight.

20 Destructive therapy agents such as salicylic acid or lactic acid may also be used. A concentration of about 2% to about 40% by weight is preferred. Cantharidin is preferably utilized in a concentration of about 5% to about 30% by weight. Typical antifungals that may

- 10 -

be used in this invention and their preferred weight concentrations include: oxiconazole nitrate (0.1% to 5.0%), ciclopirox olamine (0.1% to 5.0%), ketoconazole (0.1% to 5.0%), miconazole nitrate (0.1% to 5.0%), and butoconazole nitrate (0.1% to 5.0%). For the topical treatment of seborrheic dermatitis, hirsutism, acne, and alopecia, the active pharmaceutical
5 may include an antiandrogen such as flutamide or finasteride in preferred weight percentages of about 0.5% to 10%.

Typically, treatments using a combination of drugs include antibiotics in combination with local anesthetics such as polymycin B sulfate and neomycin sulfate in combination with tetracaine for topical antibiotic gels to provide prophylaxis against infection and relief of
10 pain. Another example is the use of minoxidil in combination with a corticosteroid such as betamethasone dipropionate for the treatment of alopecia areata. The combination of an anti-inflammatory such as cortisone with an antifungal such as ketoconazole for the treatment of tinea infections is also an example.

In one embodiment, the invention comprises a dermatological composition having
15 about 0.5% to 4.0% carbomer and about 0.5% to 10% of a pharmaceutical that exists in both a dissolved state and a microparticulate state. The dissolved pharmaceutical has the capacity to cross the stratum corneum, whereas the microparticulate pharmaceutical does not. Addition of an amine base, potassium hydroxide solution, or sodium hydroxide solution completes the formation of the gel. More particularly, the pharmaceutical may include
20 dapsone, an antimicrobial agent having anti-inflammatory properties. A preferred ratio of microparticulate to dissolved dapsone is five or less.

- 11 -

In another embodiment, the invention comprises about 1% carbomer, about 80-90% water, about 10% ethoxydiglycol, about 0.2% methylparaben, about 0.3% to 3.0% dapsone including both microparticulate dapsone and dissolved dapsone, and about 2% caustic material. More particularly, the carbomer may include "CARBOPOL® 980" and the caustic material may include sodium hydroxide solution.

In a preferred embodiment, the composition comprises dapsone and ethoxydiglycol, which allows for an optimized ratio of microparticulate drug to dissolved drug. This ratio determines the amount of drug delivered, compared to the amount of drug retained in or above the stratum corneum to function in the supracorneum domain. The system of dapsone and ethoxydiglycol may include purified water combined with "CARBOPOL®" gelling polymer, methylparaben, propylparaben, titanium dioxide, BHA, and a caustic material to neutralize the "CARBOPOL®."

Another preferred embodiment of this invention relates to a composition for the treatment of herpes lesions comprising a semisolid aqueous gel; a first pharmaceutical in the gel, partially in a microparticulate form and partially in a dissolved form, where optimized delivery for early state lesions is provided when the pharmaceutical is dissolved and optimized delivery for later state lesions is provided when the pharmaceutical is in a microparticulate form; and a second pharmaceutical dissolved in the gel which provides benefit throughout lesion progression. In a preferred embodiment, the composition comprises acyclovir and 1-methyl-2-pyrrolidone, which allows for an optimized ratio of microparticulate drug to dissolved drug for the treatment of herpes lesions. Acyclovir may be present in dissolved and microparticulate forms. The ratio determines the amount of drug

- 12 -

delivered up to the point of lesion vesicle formation, as compared to the amount of drug available to be deposited into the lesion once the vesicles rupture. The drug delivery system of acyclovir and 1-methyl-2-pyrrolidone may include purified water combined with KLUCEL® hydroxypropyl cellulose gelling polymer, methylparaben, and propylparaben.

5 In another preferred embodiment, a combination drug system of acyclovir and tetracaine HCl may be formulated with 1-methyl-2-pyrrolidone to provide both antiviral and local anesthetic activity. Tetracaine HCl is a local anesthetic that alters membrane function and blocks pain. In a preferred embodiment, acyclovir comprises 5% by weight of the composition. The system of acyclovir, tetracaine HCl, and 1-methyl-2-pyrrolidone can
10 include purified water, sodium lauryl sulfate, KLUCEL® hydroxypropyl cellulose gelling polymer, methylparaben, and propylparaben. The combination of a local anesthetic with sodium lauryl sulfate has been shown to be an effective therapy for herpes lesions. The combination of the nucleoside analogue acyclovir with the anesthetic/late stage antiviral combination tetracaine HCl and sodium lauryl sulfate should provide complete topical
15 therapy for herpes lesions.

The relative percentages for each of the reagents used in the present invention may vary depending upon the desired strength of the target formulation, gel viscosity, and the desired ratio of microparticulate to dissolved pharmaceutical. Unless otherwise designated, all reagents listed above are commonly known by one of ordinary skill in the art and are
20 commercially available from pharmaceutical or cosmetic excipient suppliers.

The present invention also provides methods for preparing the dermatological compositions described above. In a general form, the method for producing a dermatological

- 13 -

gel composition having dissolved drug and microparticulate drug precipitates comprises the steps of completely dissolving a pharmaceutical in a solvent or solvent mixture; adding and adequately dispersing a polymeric thickener in water; and combining the dissolved pharmaceutical with the dispersed polymeric thickener. Alternatively, water may be slowly
5 added to the dissolved pharmaceutical, followed by the addition of a polymeric thickener. Ethoxydiglycol and 1-methyl-2-pyrrolidone are preferred solvents for use in this invention.

In one preferred embodiment, the method for preparing a dermatological composition having dissolved and microparticulate pharmaceutical comprises the steps of forming a homogenous dispersion by stirring purified water vigorously enough to form a vortex and
10 sifting gel polymer into the vortex formed in the water while continuing to stir; forming a pharmaceutical component by dissolving methyl paraben and propylparaben in ethoxydiglycol by mixing to form a solution, and mixing an active pharmaceutical with the solution until the pharmaceutical dissolved; mixing the pharmaceutical component with the homogenous dispersion to form a microparticulate pharmaceutical dispersion; and adding a
15 caustic material. The active pharmaceutical may comprise any of the types mentioned above. In a preferred embodiment, the active pharmaceutical comprises dapsone. In another preferred embodiment, the active pharmaceutical comprises acyclovir or acyclovir in combination with tetracaine or tetracaine HCl.

The order in which reagents are combined may be important, depending on the
20 particular reagents necessary for the target mixture. For example, after a pharmaceutical such as dapsone is dissolved in a solvent such as ethoxydiglycol, water may be slowly added to the dapsone in the ethoxydiglycol solution, or the dapsone in ethoxydiglycol solution may be

- 14 -

added to the water with mixing. Adding the dapsone in ethoxydiglycol solution to water may result in less polydispersity in the size of the microparticulates than adding water to the dapsone in ethoxydiglycol solutions.

The carbomer is generally dispersed in the water component of the formulation, while
5 the remaining ingredients will be dissolved or dispersed in whichever of the two components are best for dissolving or dispersing the ingredient. For example, it is suggested to dissolve methylparaben, propylparaben, and BHA in ethoxydiglycol. After the ethoxydiglycol component and water component are combined, neutralizer is added to formulate the gel.

Finally, in another embodiment of the invention, methods for the treatment of
10 dermatological conditions by topical application of the compositions of this invention are shown. These methods are useful in the treatment of diseases such as acne, herpes lesions, seborrhea dermatitis, hirsutism, and alopecia. In a preferred embodiment, a method for the treatment of dermatological conditions comprises applying topically a gel composition comprising a dissolved pharmaceutical that has the capacity to cross the stratum corneum of
15 the epidermis and become systemically available, and a microparticulate pharmaceutical that has only minimal capacity to cross the stratum corneum in its microparticulate state. In one embodiment, the dissolved pharmaceutical and microparticulate pharmaceutical comprise about 1.0% to 10% antiviral agent. In another embodiment, the dissolved pharmaceutical and microparticulate pharmaceutical comprise about 0.5% to 10% antiandrogen.

20 In a preferred embodiment, a method for the treatment of acne comprises applying topically a gel composition that comprises a dissolved anti-inflammatory pharmaceutical and a microparticulate antimicrobial pharmaceutical, wherein the dissolved anti-inflammatory

- 15 -

pharmaceutical crosses the stratum corneum of the epidermis and is absorbed into the lower two-thirds of the pilosebaceous unit, while the microparticulate antimicrobial pharmaceutical is primarily delivered into the upper third of the pilosebaceous unit, crossing the stratum corneum of the epidermis only minimally. Preferably, the dissolved pharmaceutical and
5 microparticulate pharmaceutical comprise dapsone.

In another preferred embodiment, a method for the treatment of herpes lesions comprises applying topically a semisolid gel composition that comprises a semisolid aqueous gel; a first pharmaceutical in the gel, which exists in a partially microparticulate form and a partially dissolved form, providing for optimized delivery for early state lesions when
10 dissolved and optimized delivery for later state lesions when present as a microparticulate; and a second pharmaceutical dissolved in the gel, providing benefit throughout progression of the lesion. Preferably, the first pharmaceutical comprises a nucleoside analogue, and the second pharmaceutical comprises a local anesthetic. In a preferred embodiment, the nucleoside analogue comprises acyclovir, penciclovir, famciclovir, valacyclovir, or
15 ganciclovir, and the local anesthetic comprises tetracaine, dyclonine, dibucaine, or a salt thereof, such as tetracaine HCl, dyclonine HCl, or dibucaine HCl. More preferably, acyclovir comprises 5% by weight of the composition, and tetracaine HCl comprises 2-5% by weight.

The following examples are provided to enable those of ordinary skill in the art to make and use the methods and compositions of the invention. These examples are not
20 intended to limit the scope of what the inventors regard as their invention. Additional advantages and modifications will be readily apparent to those skilled in the art.

- 16 -

Examples 1 through 6 describe methods for the preparation of compositions of the invention that include microparticulate crystalline dapsone, dissolved dapsone, and combinations of the two. The examples offer illustrations of methods that can be used to control the ratio of dissolved to microparticulate pharmaceuticals in the final product. Since
5 microparticulate pharmaceuticals are retained above the stratum corneum having negligible penetration and dissolved pharmaceuticals penetrate the stratum corneum, controlling the ratios between the two epidermal areas is important in developing a composition having an optimal delivery route for administering pharmaceuticals via the skin.

Example 7 describes a method for the preparation of compositions of this invention
10 using two different pharmaceuticals in combination, resulting in one pharmaceutical dissolved in the composition and the other present in a microparticulate state, such that two epidermal areas may be treated with two different drugs. Example 8 provides a method for preparing a composition having a pharmaceutical partially in a microparticulate state and partially dissolved, combined with a different dissolved pharmaceutical. Examples 9-11
15 provide evaluations of the compositions and methods described herein.

EXAMPLE 1

The following example provides a method for producing a topical therapeutic agent in which the pharmaceutical component is a combination of dissolved and microcrystalline dapsone. Because of the nature of the microcrystalline dapsone in the final product of
20 Example 1, microcrystalline dapsone will be retained in or above the stratum corneum and will therefore serve as a reservoir or provide drug action in the supracorneum zone. The dissolved dapsone will pass through the stratum corneum. The method of Example 1 can also

- 17 -

be used to produce a composition of this invention that includes other pharmaceuticals such as those described above.

A polymer thickener component was prepared by charging 85.7 grams of purified water to a vessel suitable to contain 100 grams of finished semisolid product, and slowly
5 sifting one gram of "CARBOPOL® 980" into a vortex formed by rapidly stirring the purified water. When a homogeneous dispersion of "CARBOPOL® 980" and water was formed, stirring was reduced to minimize air entrapment. Next, an active pharmaceutical component was prepared by charging an appropriately sized container with 10.0 g of ethoxydiglycol. 0.2 g of methylparaben and 0.1 g of propylparaben were added to the ethoxydiglycol and mixed
10 until all of the crystalline solid was dissolved. 1.0 g dapsone was added to the ethoxydiglycol and mixed until the drug was completely dissolved.

The polymer thickener component was added to the pharmaceutical component with mixing, and immediately resulted in the formation of crystalline microparticles. Once the dispersion was homogenous, 2.0 grams of a 10% w/w aqueous sodium hydroxide solution
15 were added to neutralize the CARBOPOL® 980 and form the gel.

EXAMPLE 2

The following example provides another topical therapeutic agent in which the pharmaceutical component is dissolved dapsone. The method of Example 2 can also be used to produce a composition of this invention that includes other pharmaceuticals.

20 To prepare the composition of Example 2, the procedure of Example 1 was followed using the following specific weights of reagents. All dapsone was dissolved in the final product gel, and thus, crystalline microparticles did not form when the polymer thickener

- 18 -

component was added to the pharmaceutical component. All reagent weights are shown per 100 grams of product.

	<u>Component</u>	<u>wt/100 g product</u>
	<u>Polymer Thickener Component</u>	
5	Water	86.67 g
	"CARBOPOL 980"	1.0 g
	<u>Active Pharmaceutical Component</u>	
	Ethoxydiglycol	10.0 g
	Methylparaben	0.2 g
10	Propylparaben	0.1 g
	Dapsone	0.03 g
	<u>Caustic/Amine Component</u>	
	10% w/w Sodium Hydroxide	2.0 g

EXAMPLE 3

15 The following example provides yet another topical therapeutic agent in which the pharmaceutical component is dissolved dapsone. The method of Example 3 can also be used to produce a composition of this invention that includes other pharmaceuticals such as those designated in this application.

The procedure of Example 1 was followed using reagents in the amounts designated
20 below. All of the dapsone was dissolved in the final product gel, thus crystalline microparticles did not form upon adding the polymer thickener component to the pharmaceutical component. All reagent weights are shown per 100 grams of product.

- 19 -

	<u>Component</u>	<u>wt/100 g product</u>
	<u>Polymer Thickener Component</u>	
	Water	86.6 g
	"CARBOPOL 980"	1.0 g
5	<u>Active Pharmaceutical Component</u>	
	Ethoxydiglycol	10.0 g
	Methylparaben	0.2 g
	Propylparaben	0.1 g
	Dapsone	0.1 g
10	<u>Caustic/Amine Component</u>	
	10% w/w Sodium Hydroxide	2.0 g

EXAMPLE 4

The following example provides yet another topical therapeutic agent in which the pharmaceutical component is dissolved dapsone. The method of Example 4 can also be used to produce a composition of this invention that includes other pharmaceuticals such as those designated in this application.

The procedure of Example 1 was followed using reagents in the amounts designated below. All reagent weights are shown per 100 grams of product.

	<u>Component</u>	<u>wt/100 g product</u>
20	<u>Polymer Thickener Component</u>	
	Water	86.4 g
	"CARBOPOL 980"	1.0 g

- 20 -

Active Pharmaceutical

	Ethoxydiglycol	10.0 g
	Methylparaben	0.2 g
	Propylparaben	0.1 g
5	Dapsone	0.3 g

Caustic/Amine Component

	10% w/w Sodium Hydroxide	2.0 g
--	--------------------------	-------

EXAMPLE 5

The following example provides yet another topical therapeutic agent in which the
10 pharmaceutical component is a combination of dissolved and microcrystalline dapsone.
Because of the nature of the microcrystalline dapsone in the final product of Example 5, it
will be primarily retained in or above the stratum corneum and will therefore serve as a
reservoir or provide drug action in the supracorneum zone. The method of Example 5 can
also be used to produce a composition of this invention that includes other pharmaceuticals
15 such as those designated in this application.

The procedure of Example 1 was followed using reagents in the amounts designated
below. All reagent weights are shown per 100 grams of product.

Componentwt/100 g productPolymer Thickener Component

20	Water	86.2 g
	"CARBOPOL 980"	1.0 g

- 21 -

Active Pharmaceutical Component

Ethoxydiglycol	10.0 g
Methylparaben	0.2 g
Propylparaben	0.1 g
5 Dapsone	0.5 g

Caustic/Amine Component

10% w/w Sodium Hydroxide	2.0 g
--------------------------	-------

EXAMPLE 6

The following example provides a method for producing a topical therapeutic agent in
10 which the pharmaceutical component is a combination of dissolved and microcrystalline
dapsones. Because of the nature of the microcrystalline dapsones in the final product of
Example 6, it will be retained in or above the stratum corneum and will therefore serve as a
reservoir or provide drug action in the supracorneum zone. The method of Example 6 can
also be used to produce a composition of this invention that includes other pharmaceuticals
15 such as those designated in this application.

The procedure of Example 1 was followed using reagents in the amounts designated
below. All reagent weights are shown per 100 grams of product.

Componentwt/100 g productPolymer Thickener Component

20 Water	83.7 g
"CARBOPOL 980"	1.0 g

- 22 -

Active Pharmaceutical Component

Ethoxydiglycol	10.0 g
Methylparaben	0.2 g
Propylparaben	0.1 g
5 Dapsone	3.0 g

Caustic/Amine Component

10% w/w Sodium Hydroxide	2.0 g
--------------------------	-------

EXAMPLE 7

Example 7 describes a method for preparing a composition of this invention that
10 includes a microparticulate crystalline pharmaceutical, dapsone, in combination with a
different dissolved pharmaceutical, dyclonine HCl. Since microparticulate pharmaceuticals
are retained above the stratum corneum having negligible penetration and dissolved
pharmaceuticals penetrate the stratum corneum, using different drugs for the two forms
(microparticulate and dissolved) enables treating the two epidermal areas with different
15 drugs. This allows further options to develop optimal delivery routes for administering
pharmaceuticals via the skin.

An active pharmaceutical component was prepared by charging an appropriately sized
container with 15.0 g of ethoxydiglycol. 0.3 g methylparaben and 0.15 g of propylparaben
were added to the ethoxydiglycol and mixed until all of the crystalline solid was dissolved.
20 1.5 g of dapsone were added to the ethoxydiglycol and mixed until the drug was completely
dissolved.

- 23 -

An active pharmaceutical component which would remain dissolved was prepared by charging an appropriately sized container with 127.8 grams of purified water. 1.5 grams of dyclonine HCl was added to the water with mixing until all of the drug was completely dissolved.

5 The solvent phase was added to the aqueous phase and crystalline microparticles of dapsone were immediately formed. 3.75 grams of "NATROSOL"® 250 PHARM were added to form a topical gel containing microcrystalline dapsone and dissolved dyclonine HCl. The presence of microcrystalline dapsone was confirmed by optical microscopy.

EXAMPLE 8

10 Example 8 describes a method for preparing a composition of this invention that includes a pharmaceutical partially in a microparticulate form and partially in a dissolved form in combination with a different dissolved pharmaceutical. The composition finds particular use in the treatment of herpes lesions. The pharmaceutical in both dissolved and microparticulate forms provides optimized delivery for early stage lesions when dissolved
15 and optimized delivery for later stage lesions when present as a microparticulate. The other dissolved pharmaceutical provides benefit throughout the lesion progression.

An active pharmaceutical component was prepared by charging an appropriately sized container with 44.0 g of 1-methyl-2-pyrrolidone. 0.16 g methylparaben and 0.08 g propylparaben as preservatives were added to the 1-methyl-2-pyrrolidone and mixed until all
20 the crystalline solid dissolved. 8.0 g 1 N NaOH was mixed with the 1-methyl-2-pyrrolidone and preservative mixture prior to the addition of 4.0 g acyclovir. Upon heating to

- 24 -

approximately 50°C, all of the added materials dissolved to form a single phase clear solution.

An active pharmaceutical component which would remain dissolved was prepared by charging an appropriately sized container with 17.36 g purified water. 4.0 g tetracaine HCl
5 and 0.8 g of sodium lauryl sulfate was added to the water with mixing until all of the solids were dissolved.

The solvent phase was added to the aqueous phase and crystalline microparticles of acyclovir were immediately formed. 1.60 grams of KLUCEL® HF hydroxypropyl cellulose were added to form a topical gel containing microcrystalline acyclovir, dissolved acyclovir,
10 and dissolved tetracaine HCl and sodium lauryl sulfate. The presence of microcrystalline acyclovir was confirmed by optical microscopy.

EXAMPLE 9--EVALUATION

Example 9 evaluates the compositions of examples 1-6 and demonstrates that increasing the dapsone content of this invention's composition from 0 to 0.3% increases the
15 amount of dapsone that permeates into the skin, whereas further increasing dapsone above 0.3% results in microparticulate dapsone that remains above the stratum corneum or supracorneum zone. Therefore, the amount of dapsone that permeates into the skin does not increase incrementally for compositions having increased dapsone above 0.3%.

In example 9, experiments were performed by loading excised human skin on a
20 standard Franz type vertical diffusion cell having a 15 mm orifice, 7.0 ml volume, and equipped with a Hanson Helix stirrer using a phosphate buffered saline/ethoxydiglycol mixture as the receptor phase. The full thickness of human abdominal skin was removed

- 25 -

from a cadaver within 24 hours of death. The subcutaneous tissue was removed using a #22 scalpel blade. The tissue was cut into 5 X 15 cm sections with care being taken to avoid contamination of the stratum corneum with subcutaneous fat. Each 5 X 15 cm section was placed in a sterile plastic bag and stored on wet ice until being placed in the freezer
5 (maximum transport time 2 hours). A single 5 X 15 cm section of skin was removed from the freezer on the day of an in-vitro skin permeation study. The skin was thawed, rinsed, patted dry with a tissue, and loaded on the Franz diffusion cell. Samples were dosed with 10 mg of the formulation i.e. finite dosing. Receptor solutions were assayed by reverse phase HPLC using UV detection.

10 The cumulative drug concentration in the receptor solution was monitored over a 72 hour period. The data in Table 1 shows the mean quantities of drug transported for quadruplicate *in-vitro* skin permeation experiments in which each formulation was evaluated using the same donor skin on each of four different days. As seen in Table 1, the cumulative amount of dissolved dapsone that was transported across the stratum corneum increased until
15 the dapsone level at which the formation of microparticulate dapsone was reached, i.e. up to 0.3 weight percent dapsone. At concentrations above 0.3 %, the amount of dissolved drug did not increase, and the amount of drug delivered remained the same. Thus, the excess drug (above 0.3 weight percent) was retained in the stratum corneum or supracorneum zone.

For this drug delivery system, the microparticulate to dissolved pharmaceutical ratio
20 of 1.5 to 15 resulted in increasing amounts of drug available for antimicrobial action in the supracorneum zone, while maintaining a set amount of dapsone available for anti-inflammatory activity in the viable epidermis.

Table 1.

**Dapsone Concentration in Receptor Solution Following
in-vitro Dosing of 10 mg of Topically Applied Semisolid**

Dapsone Semisolid Concentration (% w/w)	Example Number	$\mu\text{g Dapsone}/1.77\text{ cm}^2$ by 72 hrs	% of Applied Dose transported by 72 hrs
0.03	2	0.35 ± 0.1	11
0.1	3	0.73 ± 0.3	7
0.3	4	2.6 ± 2.2	8
0.5	5	2.2 ± 0.8	4
1.0	1	2.9 ± 1.7	3
3.0	6	3.3 ± 2.5	1

EXAMPLE 10--EVALUATION

Example 10 demonstrates the importance of using the optimum microparticulate to dissolved pharmaceutical ratio. The results in Table 2 show that for the same amount of applied drug, the amount of drug in the supracorneum zone can be optimized by improving the microparticulate to dissolved pharmaceutical ratio.

In example 10, the procedures of example 9 were used, including Franz diffusion cell and experimental technique, however 1% dapsone formulations were compared. Formulation number 1 was manufactured according to example 1 and contained 1% dapsone and 10% ethoxydiglycol having a microparticulate drug/dissolved drug ratio of 5. Formulation number

- 27 -

2 had the composition 1% dapsone, 25% ethoxydiglycol, 70.7% water, 1% "CARBOPOL 980," 0.2% methylparaben, 0.1% propylparaben, and 2% sodium hydroxide solution (10% w/w). For formulation number 2, all of the dapsone is dissolved with no microparticulate drug present.

5 Approximately 10 mg of product was rubbed into the skin, and after 72 hours, the skin surface was tape stripped to remove all supracorneum drug. The skin was cut into small pieces and extracted while the receptor solution was directly assayed. For formulation number 1 (microparticulate to dissolved drug ratio equal to 5) $68 \pm 4\%$ of the applied dose remained in the supracorneum zone, while only $52 \pm 3\%$ of formulation 2 (particulate to dissolved drug ratio = 0) remained in the supracorneum zone. Each value is the mean of
10 triplicate experiments.

Table 2.

**Distribution of Dapsone in Different Skin Layers
by 72 hrs of *in-vitro* Permeation Study**

Formulation	1% DDS Hydrogel in 10% DGME (microcrystal form)	1% DDS Hydrogel in 25% DGME (soluble form)
Receptor	$0.92\% \pm 0.14$	$1.29\% \pm 0.24$
Surface residual	$68.42\% \pm 3.84$	$52.51\% \pm 2.88$
Stratum corneum	$15.54\% \pm 4.12$	$32.15\% \pm 6.16$
Dermal	$13.59\% \pm 2.15$	$14.06\% \pm 7.31$
Total recovery	$60.7\% \pm 5.3$	$53.7\% \pm 7.7$

- 28 -

EXAMPLE 11--EVALUATION

Example 11 demonstrates that a 2% acyclovir solution in 1-methyl-2 pyrrolidone provides the same level of intact skin delivery as a 5% acyclovir gel formulation that contains approximately 2% dissolved acyclovir and 3% microparticulate acyclovir. These delivery results are compared with the Spruance (1986 Antimicrobial Agents and Chemotherapy, Vol. 29, No. 5, Pgs. 730-732) standard acyclovir intact skin delivery formulation comprised of 5% acyclovir dissolved in 95% dimethyl sulfoxide (DMSO). The 95/5 DMSO/acyclovir standard formulation is known to be 60-fold more permeable than the commercially available ZOVRAX® Ointment (5% acyclovir in a polyethylene glycol base).

In example 11, the procedures of example 9 were used except that 50-500 microliter doses of formulations were applied to dermatomed human skin which was not frozen until after 48 hours post-mortem. The formulation of example 8 which contains 5% acyclovir (of which approximately 40% is dissolved and 60% is in a microparticulate form) was tested along with a 2% acyclovir in 1-methyl-2-pyrrolidone. In direct comparison with the 95/5 DMSO/acyclovir skin delivery standard, both the 5% acyclovir combination gel of example 8 and the 2% acyclovir in 1-methyl-2-pyrrolidone solution delivered 20-25% of the acyclovir delivery standard. The presence of crystalline microparticulate drug does not increase the delivery of acyclovir across intact skin. The delivery of the dissolved acyclovir is considered optimized since it was more than 10-fold greater than skin delivery of acyclovir from the commercialized ZOVRAX® formulation.

- 29 -

Those skilled in the art will recognize that, while specific embodiments have been illustrated and described, various modifications and changes may be made without departing from the spirit and scope of the invention.

- 30 -

CLAIMS:

1. A dermatological composition comprising:

a semisolid aqueous gel;

a pharmaceutical dissolved in said gel, wherein said dissolved pharmaceutical has the
5 capacity to cross the stratum corneum layer of the epidermis and become
available systemically; and

a microparticulate pharmaceutical dispersed in said gel, wherein said microparticulate
pharmaceutical does not cross the stratum corneum of the epidermis in its
microparticulate state.

10

2. The composition of claim 1, wherein said gel comprises carbomer,
hydroxyethylcellulose, hydroxypropylcellulose, cellulose gum, MVE/ME decadiene
crosspolymer, or PVM/MA copolymer.

15 3. The composition of claim 1, wherein said gel comprises a cross-linked acrylic acid
polymer dissolved in an aqueous phase and a caustic material.

4. The composition of claim 1, wherein said microparticulate pharmaceutical is a
crystalline precipitant.

20

5. The composition of claim 1, wherein said microparticulate pharmaceutical is an
amorphous precipitant.

- 31 -

6. The composition of claim 1, wherein said dissolved pharmaceutical and microparticulate pharmaceutical comprise an antimicrobial agent, anti-inflammatory agent, antiviral agent, local anesthetic, corticosteroid, destructive therapy agent, antifungal agent, or antiandrogen agent.

5

7. The composition of claim 1, wherein the ratio of microparticulate pharmaceutical to dissolved pharmaceutical is no greater than 5.

8. The composition of claim 1, wherein said microparticulate pharmaceutical and
10 dissolved pharmaceutical comprise the same pharmaceutical.

9. The composition of claim 1, wherein said microparticulate pharmaceutical and dissolved pharmaceutical comprise different pharmaceuticals.

15 10. A dermatological composition, comprising a semisolid aqueous gel having about 0.5% to 4.0% carbomer; about 0.5% to 10% pharmaceutical, wherein said pharmaceutical comprises a dissolved pharmaceutical having the capacity to cross the stratum corneum of the epidermis and become systemically available and a microparticulate pharmaceutical lacking the ability to cross the stratum corneum of the epidermis in its microparticulate state; and an
20 amine base, sodium hydroxide solution, or potassium hydroxide solution.

- 32 -

11. The dermatological composition of claim 10, wherein said pharmaceutical comprises dapsone.

12. The dermatological composition of claim 11, wherein the microparticulate to
5 dissolved pharmaceutical ratio is no greater than 5.

13. The composition of claim 10, wherein said pharmaceutical comprises an antimicrobial agent, antiviral agent, anti-inflammatory agent, local anesthetic, corticosteroid, destructive therapy agent, antifungal agent, or antiandrogen agent.

10

14. A dermatological gel composition including microparticulate pharmaceutical and dissolved pharmaceutical, which comprises:

about 1 % carbomer;

about 83.7 to 86.4% water;

15 about 10% ethoxydiglycol;

about 0.2% methylparaben;

up to about 3% dapsone in a microparticulate and dissolved state;

and

about 2 % sodium hydroxide solution.

20

15. The dermatological composition of claim 14, wherein the ratio of microparticulate to dissolved dapsone is no greater than 5.

- 33 -

16. A dermatological gel composition for the treatment of herpes lesions comprising:
a semisolid aqueous gel;
a first pharmaceutical in said gel, partially in a microparticulate form and partially in a dissolved form, wherein said first pharmaceutical provides optimized delivery
5 for early state lesions when dissolved and optimized delivery for later state lesions when present as a microparticulate; and
a second pharmaceutical dissolved in said gel, wherein said second pharmaceutical provides benefit throughout lesion progression.
- 10 17. The composition of claim 16, wherein said first pharmaceutical comprises a nucleoside analogue such as acyclovir, penciclovir, famciclovir, valacyclovir, or ganciclovir, and said second pharmaceutical comprises a local anesthetic such as tetracaine, tetracaine HCl, dyclonine, dyclonine HCl, dibucaine, or dibucaine HCl.
- 15 18. The composition of claim 17, wherein said acyclovir comprises 5% by weight of the composition.
19. The composition of claim 17, wherein said tetracaine HCl comprises 5% by weight of the composition.

- 34 -

20. A dermatological composition for the treatment of herpes lesions comprising acyclovir and 1-methyl-2-pyrrolidone in a semisolid aqueous gel, wherein said acyclovir is present in dissolved and microparticulate forms.

5 21. A method for the treatment of dermatological conditions comprising applying topically a semisolid gel composition, wherein said composition comprises:

a semisolid aqueous gel;

a first pharmaceutical in said gel, partially in a microparticulate form and partially in a dissolved form, wherein said first pharmaceutical provides optimized delivery

10 for early state lesions when dissolved and optimized delivery for later state lesions when present as a microparticulate; and

a second pharmaceutical dissolved in said gel, wherein said second pharmaceutical provides benefit throughout lesion progression.

15 22. The method of claim 21, wherein said active pharmaceutical comprises an antimicrobial agent, anti-inflammatory agent, antiviral agent, local anesthetic agent, corticosteroid, destructive therapy agent, antifungal agent, or antiandrogen agent.

23. The method of claim 21, wherein said active pharmaceutical comprises dapsone.

20

24. A method for treating disease comprising applying topically an aqueous gel composition comprising a dissolved pharmaceutical that has the capacity to cross the stratum

- 35 -

corneum of the epidermis and become systemically available, and a microparticulate pharmaceutical that has only minimal capacity to cross the stratum corneum in its microparticulate state.

5 25. The method of claim 24, wherein said dissolved pharmaceutical and said microparticulate pharmaceutical comprise about 1.0% to 10% antiviral agent.

26. The method of claim 24, wherein said dissolved pharmaceutical and said microparticulate pharmaceutical comprise about 0.5 to 10% antiandrogen.

10

27. A method for treating acne, which comprises applying topically a gel composition comprising a dissolved anti-inflammatory pharmaceutical and a microparticulate antimicrobial pharmaceutical, wherein said dissolved anti-inflammatory pharmaceutical crosses the stratum corneum of the epidermis and is absorbed into the lower two-thirds of the pilosebaceous unit, while said microparticulate antimicrobial pharmaceutical is primarily
15 delivered into the upper third of the pilosebaceous unit, crossing the stratum corneum of the epidermis only minimally.

28. The method of claim 27, wherein said anti-inflammatory pharmaceutical and said
20 antimicrobial pharmaceutical comprise dapsone.

- 36 -

29. A method for the preparation of a dermatological gel composition wherein said composition includes dissolved and microparticulate pharmaceutical, comprising the steps of:

forming a homogenous dispersion by stirring purified water vigorously enough to
form a vortex and sifting gel polymer into the vortex formed in the water while
5 continuing to stir;

forming a pharmaceutical component by dissolving methyl paraben and
propylparaben in ethoxydiglycol by mixing to form a solution, and mixing an
active pharmaceutical with said solution until said pharmaceutical dissolves;

mixing said pharmaceutical component with said homogeneous dispersion to form a
10 microparticulate pharmaceutical dispersion;

and

adding a caustic material.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 97/15919

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K9/06

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 679 390 A (ZYMA S.A.) 2 November 1995 see claims 1-5 see column 2, line 7 - line 26 ---	1-6,9, 16,17, 21,22,24
X	GB 2 260 080 A (RHONE-POULENC RORER LIMITED) 7 April 1993 see claims 1,5 see page 2, line 17 - line 24 see page 3, line 4 - line 11 see page 4, line 1 - line 8 -----	1,6,9, 16, 21-24, 27,28

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

16 December 1997

Date of mailing of the international search report

15. 01. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Ventura Amat, A

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 97/15919

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 679390 A	02-11-95	CA 2148034 A GB 2288734 A	30-10-95 01-11-95
GB 2260080 A	07-04-93	NONE	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 15919

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 21-28
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.