(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
19 April 2018 (19.04.2018)

WIPO | PCT

(10) International Publication Number
WO 2018/071606 A1

(51) International Patent Classification:
C07D 233/86 (2006.01) *C07D 471/10* (2006.01)
C07D 401/04 (2006.01) *A61K 31/4164* (2006.01)
C07D 491/107 (2006.01) *A61K 31/4439* (2006.01)

(21) International Application Number:
PCT/US2017/056234

(22) International Filing Date:
11 October 2017 (11.10.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/406,888 11 October 2016 (11.10.2016) US
62/528,385 03 July 2017 (03.07.2017) US

(71) Applicant: **ARVINAS, INC.** [US/US]; 5 Science Park, New Haven, CT 06511 (US).

(72) Inventors: **CREW, Andrew, P.**; 2 Winterberry Road, Guilford, CT 06437 (US). **HORNBERGER, Keith, R.**; 182 Wolf Pit Drive, Southburg, CT 06488 (US). **SNYDER, Lawrence, B.**; 85 Laurel Ridge Trail, Killingworth, CT 06419 (US). **ZIMMERMANN, Kurt**; 5 Skeetfield Point Road, Durham, CT 06422 (US). **WANG, Jing**; 25 Elm Street, Milford, CT 06460 (US). **BERLIN, Michael**; 14 Hendrick Road, Flemington, NJ 08822 (US). **CREWS, Craig, M.**; 286 Livingston Street, New Haven, CT 06511 (US). **DONG, Hanqing**; 248 Opening Hill Rd., Madison, CT 06443 (US).

(74) Agent: **ZERHUSEN, Bryan D.** et al.; Locke Lord LLP, P.O. Box 55874, Boston, MA 02205 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:
— *with international search report (Art. 21(3))*
— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(54) Title: COMPOUNDS AND METHODS FOR THE TARGETED DEGRADATION OF ANDROGEN RECEPTOR

(57) Abstract: The present disclosure relates to bifunctional compounds, which find utility to degrade and (inhibit) Androgen Receptor. In particular, the present disclosure is directed to compounds, which contain on one end a cereblon ligand which binds to the E3 ubiquitin ligase and on the other end a moiety which binds Androgen Receptor, such that Androgen Receptor is placed in proximity to the ubiquitin ligase to effect degradation (and inhibition) of Androgen Receptor. The present disclosure exhibits a broad range of pharmacological activities associated with compounds according to the present disclosure, consistent with the degradation/inhibition of Androgen Receptor.

FIG. 1A



WO 2018/071606 A1

COMPOUNDS AND METHODS FOR THE TARGETED DEGRADATION OF ANDROGEN RECEPTOR

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims priority to United States Provisonal Application No. 62/406,888, filed 11 October 2016, and United States Provisional Application No. 62/528,385, filed 3 July 2017, all of which are incorporated by reference in their entirety for all purposes.

STATEMENT REGARDING FEDERALLY FUNDED RESEARCH

[0002] This invention was made with government support under grant number 1R44CA203199-01 by the National Cancer Institute. The government has certain rights in the invention

INCORPORATION BY REFERENCE

[0003] U.S. Patent Application Serial No. 14/686,640, filed on April 14, 2015, published as U.S. Patent Application Publication No. 2016/0058872; U.S. Patent Application Serial No. 14/792,414, filed on Jul 6, 2015, published as U.S. Patent Application Pulbication No. 2015/0291562; U.S. Patent Application Serial No. 14/371,956, filed on July, 11, 2014, published as U.S. Patent Application Publication No. 2014/0356322; U.S. Patent Application Serial No. 15/074,820, filed on March 18, 2016, published as U.S. Patent Application Publication No. 2016/0272639, are incorporated herein in their entireties. Furthermore, all references cited herein are incorporated by reference herein in their entirety.

FIELD OF THE INVENTION

[0004] The present description provides imide-based compounds, including bifunctional compounds comprising the same, and associated methods of use. The bifunctional compounds are useful as modulators of targeted ubiquitination, especially with respect to a variety of polypeptides and other proteins, which are degraded and/or otherwise inhibited by bifunctional compounds according to the present disclosure. In certain aspects, the bifunctional compounds comprise a Cereblon E3 Ubiquitin Ligase (CRBN) binding moiety, which binds to the cereblon

E3 ubiquitin ligase, a target protein binding moiety, which binds to the target protein (e.g., androgen receptor), and optionally a linker moiety which links the cereblon binding moiety and target protein binding moiety. These compounds work in such a way that the target protein/polypeptide is placed in proximity to the ubiquitin ligase to effect degradation (and inhibition) of that protein (e.g., androgen receptor).

BACKGROUND

[0005] Most small molecule drugs bind enzymes or receptors in tight and well-defined pockets. On the other hand, protein-protein interactions are notoriously difficult to target using small moelcules due to their large contact surfaces and the shallow grooves or flat interfaces involved. E3 ubiquitin ligases (of which hundreds are know in humans) confer substrate specificity for ubiquitination, and therefore, are more attractive therapeutic targets than general proteasome inhibitors due to their specificity for certain protein substrates. The development of ligands of E3 ligases has proven challenging, in part due to the fact that they must disrupt protein-protein interactions. However, recent developments have provided specific ligands wich bind to these ligases. For example, since the discovery of nutlins, the first small molecule E3 ligase inhibitors, additional compounds have been reported that target E3 ligases, but the field remains underdeveloped.

[0006] One E3 ligase with therapeutic potential is Cereblon, a protein that in humans is encoded by the CRBN gene. CRBN orthologs are highly conserved from plants to humans, which underscores its physiological importance. Cereblon forms an E3 ubiquitin ligase complex with damaged DNA binding protein 1 (DDB1), Cullin-4A (CUL4A), and regulator of cullins 1 (ROC1). This complex ubiquitinates a number of other proteins. Through a mechanism which has not been completely elucidated, cereblon ubquitination of target proteins results in increased levels of fibroblast growth factor 8 (FGF8) and fibroblast growth factor 10 (FGF10). FGF8 in turn regulates a number of developmental processes, such as limb and auditory vesicle formation. The net result is that this ubiquitin ligase complex is important for limb outgrowth in embryos. In the absence of cereblon, DDB1 forms a complex with DDB2 that functions as a DNA damage-binding protein.

[0007] Thalidomide, which has been approved for the treatment of a number of immunological indications, has also been approved for the treatment of certain neoplastic diseases, including multiple myeloma. In addition to multiple myeloma, thalidomide and several of its analogs are also currently under investigation for use in treating a variety of other types of cancer. While the precise mechanism of thalidomide's anti-tumor activity is still emerging, it is known to inhibit angiogenesis. Recent literature discussing the biology of the imides includes Lu *et al. Science* 343, 305 (2014) and Krönke *et al. Science* 343, 301 (2014).

[0008] Significantly, thalidomide and its analogs e.g. pomolinamiode and lenalinomide, are known to bind cereblon. These agents bind to cereblon, altering the specificity of the complex to induce the ubiquitination and degradation of Ikaros (IKZF1) and Aiolos (IKZF3), transcription factors essential for multiple myeloma growth. Indeed, higher expression of cereblon has been linked to an increase in efficacy of imide drugs in the treatment of multiple myeloma.

[0009] Androgen Receptor (AR) belongs to a nuclear hormone receptor family that is activated by androgens, such as testosterone and dihydrotestosterone (*Pharmacol. Rev.* 2006, 58(4), 782-97; *Vitam. Horm.* 1999, 55:309-52.). In the absence of androgens, AR is bound by Heat Shock Protein 90 (Hsp90) in the cytosol. When an androgen binds AR, its conformation changes to release AR from Hsp90 and to expose the Nuclear Localization Signal (NLS). The latter enables AR to translocate into the nucleus where AR acts as a transcription factor to promote gene expression responsible for male sexual characteristics (*Endocr. Rev.* 1987, 8(1):1-28; *Mol. Endocrinol.* 2002, 16(10), 2181-7). AR deficiency leads to Androgen Insensitivity Syndrome, formerly termed testicular feminization.

[0010] While AR is responsible for development of male sexual characteristics, it is also a well-documented oncogene in certain forms cancers including prostate cancers (*Endocr. Rev.* 2004, 25(2), 276-308). A commonly measured target gene of AR activity is the secreted Prostate Specific Antigen (PSA) protein. The current treatment regimen for prostate cancer involves inhibiting the androgen-AR axis by two methods. The first approach relies on reduction of androgens, while the second strategy aims to inhibit AR function (Nat. Rev. Drug Discovery, 2013, 12,823–824). Despite the development of effective targeted therapies, most patients develop resistance and the disease progresses. An alternative approach for the treatment of prostate cancer involves eliminating the AR protein. Because AR is a critical driver of

tumorigenesis in many forms of prostate cancers, its elimination should lead to therapeutically beneficial response.

[0011] There exists an ongoing need in the art for effective treatments for diseases, especially cancer, prostate cancer, and Kennedy's Disease. However, non-specific effects, and the inability to target and modulate certain classes of proteins altogether, such as transcription factors, remain as obstacles to the development of effective anti-cancer agents. As such, small molecule therapeutic agents that leverage or potentiate cereblon's substrate specificity and, at the same time, are "tunable" such that a wide range of protein classes can be targeted and modulated with specificity would be very useful as a therapeutic.

BRIEF SUMMARY OF THE INVENTION

[0012] The present disclosure describes bi-functional compounds, which function to recruit endogenous proteins to an E3 Ubiquitin Ligase for degradation, and methods of using the same. In particular, the present disclosure provides bifunctional or proteolysis targeting chimeric (PROTAC) compounds, which find utility as modulators of targeted ubiquitination of a variety of polypeptides and other proteins (such as androgen receptor), which are then degraded and/or otherwise inhibited by the bifunctional compounds as described herein. An advantage of the compounds provided herein is that a broad range of pharmacological activites is possible, consistent with the degradation/inhibition of targeted polypeptides from virtually any protein class or family. In addition, the description provides methods of using an effective amount of the compounds as described herein for the treatment or amelioration of a disease condition including cancer, e.g., prostate cancer, and Kennedy's Disease.

[0013] Thus, in one aspect, the disclosure provides novel imide-based compounds as described herein.

[0014] In an additional aspect, the disclosure provides bifunctional or PROTAC compounds, which comprise an E3 Ubiqutin Ligase binding moiety (i.e. a ligand for an E3 Ubiquitin Ligase or "ULM" group), and a moiety that binds a target protein (i.e. a protein/polypeptide targeting ligand or "PTM" group) such that the target protein/polypeptide is placed in proximity to the ubiquitin ligase to effect degrataion (and inhibition) of that protein. In a preffered embodiment the ULM is a cereblon E3 Ubiquitin Ligase binding moiety (i.e. a "CLM"). For example, the structure of the bifunctional compound can be depicted as:

PTM – CLM.

[0015] The respective positions of the PTM and CLM moieties as well as their number as illustrated herein is provided by way of example only and is not intended to limit the compounds in any way. As would be understood by the skilled artisan, the bifunctional compounds as described herein can be synthesized such that the number and position of the respective functional moieties can be varied as desired.

[0016] In certain embodiments the bifunctional compound may further comprise a chemical linker ("L"). In this example, the structure of the bifunctional compunds can be depicted as:

PTM – L – CLM,


[0017] wherein the PTM is a protein/polypeptide targeting moiety, the L is a chemical linker moiety or bond that links the PTM and the CLM, and the CLM is a cereblon E3 ubiquitin ligase binding moiety.

[0018] In certain preferred embodiments, the E3 Ubiquitin Ligase is cereblon. As such, in certain additional embodiments, the CLM of the bifunctional compound comprises chemistries such as imide, amide, thioamide, thioimide derived moieties. In additional embodiments, the CLM comprises a phthalimido group or an analog or derivative thereof. In still additional embodiments, the CLM comprises a phthalimido-glutarimide group or an analog or derivative thereof. In still other embodiments, the CLM comprises a member of the group consisting of thalidomide, lenalidomide, pomalidomide, and analogs or derivatives thereof.


[0019] In certain embodiments, the compounds as described herein comprise multiple CLMs, multiple PTMs, multiple chemical linkers or a combination thereof.

[0020] It will be understood that the general structures are exemplary and the respective moieties can be arranged spatially in any desired order or configuration, e.g., CLM-L-PTM, and PTM-L-CLM respectively.


[0021] In certain embodiments, the PTM is an AR binding moieties (ABM). In particular embodiments, the ABM is selected from following structures:




ABM-a ,



ABM-b ,



ABM-c , and



ABM-d .

wherein:

W^1 is aryl, heteroaryl, , bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted; for example, optionally substituted by by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, S;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;

Q is a 3-6 membered alicyclic or aromatic ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q,is independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, cyclic, heterocyclic, or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, C_{1-6} alicyclic, heterocyclic, aryl, heteroaryl,biheterocyclic, biaryl, or biheteroaryl, each optionally substituted by 1-10 R^{W2}; each R^{W2} is independently H, halo, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more F), C_{1-6} heteroalkyl (linear, branched, optionally substituted), $-OR^{W2A}$, C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, OC_{1-3}alkyl (optionally substituted; for example, optionally substituted by 1 or more –F), , C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and

R^{W2A} is H, C_{1-6} alkyl (linear, branched), or C_{1-6} heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC_{1-3}alkyl.

[0022] In an additional aspect, the description provides therapeutic compositions comprising an effective amount of a compound as described herein or salt form thereof, and a pharmaceutically acceptable carrier. The therapeutic compositions modulate protein degradation in a patient or subject, for example, an animal such as a human, and can be used for treating or ameliorating disease states or conditions which are modulated through the degraded protein. In certain embodiments, the therapeutic compositions as described herein may be used to effectuate the degradation and/or inhibition of proteins of interest for the treatment or amelioration of a disease, e.g., cancer. In yet another aspect, the present disclosure provides a method of ubiquitinating/ degrading a target protein in a cell. In certain embodiments, the method comprises administering a bifunctional compound as described herein comprising an CLM and a PTM, preferably linked through a linker moiety, as otherwise described herein, wherein the CLM is coupled to the PTM and wherein the CLM recognizes a ubiquitin pathway protein (e.g., a ubiquitin ligase, preferably an E3 ubiquitin ligase such as cereblon) and the PTM recognizes the target protein such that degradation of the target protein will occur when the target protein is placed in proximity to the ubiquitin ligase, thus resulting in degradation/inhibition of the effects of the target protein and the control of protein levels. The control of protein levels afforded by the present disclosure provides treatment of a disease state or condition, which is modulated through the target protein by lowering the level of that protein in the cells of a patient.

[0023] In an additional aspect, the description provides a method for assessing (i.e., determining and/or measuring) a CLM's binding affinity. In certain embodiments, the method comprises

providing a test agent or compound of interest, for example, an agent or compound having an imide moiety, e.g., a phthalimido group, phthalimido-glutarimide group, derivatized thalidomide, derivatized lenalidomide or derivatized pomalidomide, and comparing the cereblon binding affinity and/or inhibitory activity of the test agent or compound as compared to an agent or compound known to bind and/or inhibit the activity of cereblon.

[0024] In still another aspect, the description provides methods for treating or emeliorating a disease, disorder or symptom thereof in a subject or a patient, e.g., an animal such as a human, comprising administering to a subject in need thereof a composition comprising an effective amount, e.g., a therapeutically effective amount, of a compound as described herein or salt form thereof, and a pharmaceutically acceptable carrier, wherein the composition is effective for treating or ameliorating the disease or disorder or symptom thereof in the subject.

[0025] In another aspect, the description provides methods for identifying the effects of the degradation of proteins of interest in a biological system using compounds according to the present disclosure.

[0026] The preceding general areas of utility are given by way of example only and are not intended to be limiting on the scope of the present disclosure and appended claims. Additional objects and advantages associated with the compositions, methods, and processes of the present disclosure will be appreciated by one of ordinary skill in the art in light of the instant claims, description, and examples. For example, the various aspects and embodiments of the invention may be utilized in numerous combinations, all of which are expressly contemplated by the present description. These additional advantages objects and embodiments are expressly included within the scope of the present disclosure. The publications and other materials used herein to illuminate the background of the invention, and in particular cases, to provide additional details respecting the practice, are incorporated by reference.

[0027] Where applicable or not specifically disclaimed, any one of the embodiments described herein are contemplated to be able to combine with any other one or more embodiments, even though the embodiments are described under different aspects of the disclosure. As such, the preceding general areas of utility are given by way of example only and are not intended to be limiting on the scope of the present disclosure and appended claims. Additional objects and advantages associated with the compositions, methods, and processes of the present disclosure will be appreciated by one of ordinary skill in the art in light of the instant

claims, description, and examples. For example, the various aspects and embodiments of the disclosure may be utilized in numerous combinations, all of which are expressly contemplated by the present description. These additional advantages objects and embodiments are expressly included within the scope of the present disclosure. The publications and other materials used herein to illuminate the background of the disclosure, and in particular cases, to provide additional details respecting the practice, are incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0028] The accompanying drawings, which are incorporated into and form a part of the specification, illustrate several embodiments of the present disclosure and, together with the description, serve to explain the principles of the present disclosure. The drawings are only for the purpose of illustrating an embodiment of the disclosure and are not to be construed as limiting the invention. Further objects, features and advantages of the invention will become apparent from the following detailed description taken in conjunction with the accompanying figures showing illustrative embodiments of the disclosure, in which:

[0029] **Figure 1A and Figure 1B.** Illustration of general principle for PROTAC function. **Figure 1A:** Exemplary PROTACs comprise an androgen receptor targeting moiety (ABM; *darkly shaded rectangle*), a E3 ubiquitin Ligase binding moiety, such as a cereblon E3 ubiquitin ligase binding moiety (CLM; *lightly shaded triangle*), and a linker moiety (L; *black line*) coupling or tethering the ABM to the CLM (as described herein, L can be absent or a bond or a chemical linker moiety). **Figure 1B** Illustrates the functional use of the PROTACs as described herein. Briefly, the CLM recognizes and binds to cereblon E3 ubiquitin ligase, and the ABM binds and recruits androgen receptor and brings it into close proximity to the cereblon E3 ubiquitin ligase. Typically, the E3 ubiquitin ligase is complexed with an E2 ubiquitin-conjugating protein, and either alone or via the E2 protein catalyzes attachment of ubiquitin (*dark circles*) to a lysine on the target protein via an isopeptide bond. The poly-ubiquitinated protein (*far right*) is then targeted for degradation by the proteosomal machinery of the cell.

[0030] **Figure 2**. Table 2 that include Exemplary Compounds 1-75, as well as the General Scheme that may be used to produce each of the Exemplary Compounds. Table 2 also includes the DC50, Dmax, M/Z+, and 1H NMR data for each of the Exemplary Compounds. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells): A <1nM; B: 1-

10nM; C: 10-100nM; D: >100nM. Dmax categories (degradation of AR-maximum inhibition (%) AR ELISA in LNCaP and/or VCaP cells): A >50%; B <50%.

[0031] **Figure 3**. Table 3 that include Exemplary Compounds 76-398, as well as the General Scheme that may be used to produce each of the Exemplary Compounds. Table 3 also includes the DC50, M/Z+, and 1H NMR data for each of the Exemplary Compounds. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells): A <1nM; B: 1-10nM; C: 10-100nM; D: >100nM.

[0032] **Figure 4**. Table 4 that include Exemplary Compounds 399-427, as well as the General Scheme that may be used to produce each of the Exemplary Compounds. Table 4 also includes the DC50, Dmax (%), M/Z+, and 1H NMR data for each of the Exemplary Compounds. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells): A <1nM; B: 1-10nM; C: 10-100nM; D: >100nM.

[0033] **Figure 5**. Table 5 that include Exemplary Compounds 428-452. Table 5 also includes the DC50, Dmax (%), M/Z+, and 1H NMR data for each of the Exemplary Compounds. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells): A <1nM; B: 1-10nM; C: 10-100nM; D: >100nM.

[0034] **Figure 6**. Table 6 that include Exemplary Compounds 453-528. Table 6 also includes the DC50, Dmax, M/Z+, and 1H NMR data for each of the Exemplary Compounds. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells): A <1nM; B: 1-10nM; C: 10-100nM; D: >100nM. Dmax categories (degradation of AR-maximum inhibition (%) AR ELISA in LNCaP and/or VCaP cells): A >50%; B <50%.

[0035] **Figure 7**. Table 7 that include Exemplary Compounds 529-625. Table 7 also includes the DC50, M/Z+, and 1H NMR data for each of the Exemplary Compounds. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells): A <1nM; B: 1-10nM; C: 10-100nM; D: >100nM.

[0036]

DETAILED DESCRIPTION

[0037] The following is a detailed description provided to aid those skilled in the art in practicing the present invention. Those of ordinary skill in the art may make modifications and variations in the embodiments described herein without departing from the spirit or scope of the

present disclosure. All publications, patent applications, patents, figures and other references mentioned herein are expressly incorporated by reference in their entirety.

[0038] The present description relates to the surprising and unexpected discovery that an E3 ubiquitin ligase protein can ubiquitinate a target protein, in particular androgen receptor, once the E3 ubiquitin ligase protein and the target protein are brought into proximity by a chimeric construct (e.g., PROTAC) as described herein, in which a moiety that binds the E3 ubiquitin ligase protein is coupled, e.g., covalently, to a moiety that bind the androgen receptor target protein. Accordingly, the present description provides compounds, compositions comprising the same, and associated methods of use for ubiquitination and degradation of a chosen target protein, e.g., androgen receptor (See **Figure 1A** and **Figure 1B**).

[0039] The present description is related in certain aspects to U.S. Patent Publications 2014/0356322A1, 2015/0291562A1, and 2016/0214972A1, all of which are incorporated herein by reference in its entirety for all purposes.

[0040] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. The terminology used in the description is for describing particular embodiments only and is not intended to be limiting of the invention.

[0041] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise (such as in the case of a group containing a number of carbon atoms in which case each carbon atom number falling within the range is provided), between the upper and lower limit of that range and any other stated or intervening value in that stated range is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges is also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either both of those included limits are also included in the invention.

[0042] The following terms are used to describe the present disclosure. In instances where a term is not specifically defined herein, that term is given an art-recognized meaning by those of ordinary skill applying that term in context to its use in describing the present invention.

[0043] The articles "a" and "an" as used herein and in the appended claims are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the

article unless the context clearly indicates otherwise. By way of example, "an element" means one element or more than one element.

[0044] The phrase "and/or," as used herein in the specification and in the claims, should be understood to mean "either or both" of the elements so conjoined, i.e., elements that are conjunctively present in some cases and disjunctively present in other cases. Multiple elements listed with "and/or" should be construed in the same fashion, i.e., "one or more" of the elements so conjoined. Other elements may optionally be present other than the elements specifically identified by the "and/or" clause, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, a reference to "A and/or B", when used in conjunction with open-ended language such as "comprising" can refer, in one embodiment, to A only (optionally including elements other than B); in another embodiment, to B only (optionally including elements other than A); in yet another embodiment, to both A and B (optionally including other elements); etc.

[0045] As used herein in the specification and in the claims, "or" should be understood to have the same meaning as "and/or" as defined above. For example, when separating items in a list, "or" or "and/or" shall be interpreted as being inclusive, i.e., the inclusion of at least one, but also including more than one, of a number or list of elements, and, optionally, additional unlisted items. Only terms clearly indicated to the contrary, such as "only one of' or "exactly one of," or, when used in the claims, "consisting of," will refer to the inclusion of exactly one element of a number or list of elements. In general, the term "or" as used herein shall only be interpreted as indicating exclusive alternatives (i.e., "one or the other but not both") when preceded by terms of exclusivity, such as "either," "one of," "only one of," or "exactly one of."

[0046] The term “about” and the like, as used herein, in association with numeric values or ranges, reflects the fact that there is a certain level of variation that is recognized and tolerated in the art due to practical and/or theoretical limitations. For example, minor variation is tolerated due to inherent variances in the manner in which certain devices operate and/or measurements are taken. In accordance with the above, the phrase “about” is normally used to encompass values within the standard deviation or standard error.

[0047] In the claims, as well as in the specification above, all transitional phrases such as "comprising," "including," "carrying," "having," "containing," "involving," "holding," "composed of," and the like are to be understood to be open-ended, i.e., to mean including but not limited to.

Only the transitional phrases "consisting of' and "consisting essentially of' shall be closed or semi-closed transitional phrases, respectively, as set forth in the United States Patent Office Manual of Patent Examining Procedures, Section 2111.03.

[0048] As used herein in the specification and in the claims, the phrase "at least one," in reference to a list of one or more elements, should be understood to mean at least one element selected from anyone or more of the elements in the list of elements, but not necessarily including at least one of each and every element specifically listed within the list of elements and not excluding any combinations of elements in the list of elements. This definition also allows that elements may optionally be present other than the elements specifically identified within the list of elements to which the phrase "at least one" refers, whether related or unrelated to those elements specifically identified. Thus, as a nonlimiting example, "at least one of A and B" (or, equivalently, "at least one of A or B," or, equivalently "at least one of A and/or B") can refer, in one embodiment, to at least one, optionally including more than one, A, with no B present (and optionally including elements other than B); in another embodiment, to at least one, optionally including more than one, B, with no A present (and optionally including elements other than A); in yet another embodiment, to at least one, optionally including more than one, A, and at least one, optionally including more than one, B (and optionally including other elements); etc.

[0049] It should also be understood that, in certain methods described herein that include more than one step or act, the order of the steps or acts of the method is not necessarily limited to the order in which the steps or acts of the method are recited unless the context indicates otherwise.

[0050] The terms "co-administration" and "co-administering" or “combination therapy” can refer to both concurrent administration (administration of two or more therapeutic agents at the same time) and time varied administration (administration of one or more therapeutic agents at a time different from that of the administration of an additional therapeutic agent or agents), as long as the therapeutic agents are present in the patient to some extent, preferably at effective amounts, at the same time. In certain preferred aspects, one or more of the present compounds described herein, are coadministered in combination with at least one additional bioactive agent, especially including an anticancer agent. In particularly preferred aspects, the co-administration of compounds results in synergistic activity and/or therapy, including anticancer activity.

[0051] The term “effective” can mean, but is in no way limited to, that amount/dose of the active pharmaceutical ingredient, which, when used in the context of its intended use, effectuates or is sufficient to prevent, inhibit the occurrence, ameliorate, delay or treat (alleviate a symptom to some extent, preferably all) the symptoms of a condition, disorder or disease state in a subject in need of such treatment or receiving such treatment. The term effective subsumes all other effective amount or effective concentration terms, e.g., “effective amount/dose,” “pharmaceutically effective amount/dose” or “therapeutically effective amount/dose,” which are otherwise described or used in the present application.

[0052] The effective amount depends on the type and severity of disease, the composition used, the route of administration, the type of mammal being treated, the physical characteristics of the specific mammal under consideration, concurrent medication, and other factors which those skilled in the medical arts will recognize. The exact amount can be ascertainable by one skilled in the art using known techniques (see, e.g., Lieberman, Pharmaceutical Dosage Forms (vols. 1-3, 1992); Lloyd, The Art, Science and Technology of Pharmaceutical Compounding (1999); Pickar, Dosage Calculations (1999); and Remington: The Science and Practice of Pharmacy, 20th Edition, 2003, Gennaro, Ed., Lippincott, Williams & Wilkins).

[0053] The term “pharmacological composition,” “therapeutic composition,” “therapeutic formulation” or “pharmaceutically acceptable formulation” can mean, but is in no way limited to, a composition or formulation that allows for the effective distribution of an agent provided by the present disclosure, which is in a form suitable for administration to the physical location most suitable for their desired activity, e.g., systemic administration.

[0054] The term "pharmaceutically acceptable" or “pharmacologically acceptable” can mean, but is in no way limited to, entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, or a human, as appropriate.

[0055] The term "pharmaceutically acceptable carrier" or “pharmacologically acceptable carrier” can mean, but is in no way limited to, any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Suitable carriers are described in the most recent edition of Remington's Pharmaceutical Sciences, a standard reference text in the field, which is incorporated herein by reference. Preferred examples of such carriers or diluents include, but are

not limited to, water, saline, finger's solutions, dextrose solution, and 5% human serum albumin. Liposomes and non-aqueous vehicles such as fixed oils may also be used. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0056] The term “systemic administration" refers to a route of administration that is, e.g., enteral or parenteral, and results in the systemic distribution of an agent leading to systemic absorption or accumulation of drugs in the blood stream followed by distribution throughout the entire body. Suitable forms, in part, depend upon the use or the route of entry, for example oral, transdermal, or by injection. Such forms should not prevent the composition or formulation from reaching a target cell (i.e., a cell to which the negatively charged polymer is desired to be delivered to). For example, pharmacological compositions injected into the blood stream should be soluble. Other factors are known in the art, and include considerations such as toxicity and forms which prevent the composition or formulation from exerting its effect. Administration routes which lead to systemic absorption include, without limitations: intravenous, subcutaneous, intraperitoneal, inhalation, oral, intrapulmonary and intramuscular. The rate of entry of a drug into the circulation has been shown to be a function of molecular weight or size. The use of a liposome or other drug carrier comprising the compounds of the instant disclosure can potentially localize the drug, for example, in certain tissue types, such as the tissues of the reticular endothelial system (RES). A liposome formulation which can facilitate the association of drug with the surface of cells, such as, lymphocytes and macrophages is also useful.

[0057] The term “local administration" refers to a route of administration in which the agent is delivered to a site that is apposite or proximal, e.g., within about 10 cm, to the site of the lesion or disease.

[0058] The term “compound”, as used herein, unless otherwise indicated, refers to any specific chemical compound disclosed herein and includes tautomers, regioisomers, geometric isomers, and where applicable, stereoisomers, including optical isomers (enantiomers) and other steroisomers (diastereomers) thereof, as well as pharmaceutically acceptable salts and derivatives (including prodrug forms) thereof where applicable, in context. Within its use in context, the term compound generally refers to a single compound, but also may include other compounds such as

stereoisomers, regioisomers and/or optical isomers (including racemic mixtures) as well as specific enantiomers or enantiomerically enriched mixtures of disclosed compounds. The term also refers, in context to prodrug forms of compounds which have been modified to facilitate the administration and delivery of compounds to a site of activity. It is noted that in describing the present compounds, numerous substituents and variables associated with same, among others, are described.

[0059] It is understood by those of ordinary skill that molecules which are described herein are stable compounds as generally described hereunder. When the bond  is shown, both a double bond and single bond are represented or understood within the context of the compound shown and well-known rules for valence interactions.

[0060] As used herein, "derivatives" can mean compositions formed from the native compounds either directly, by modification, or by partial substitution. As used herein, "analogs" can mean compositions that have a structure similar to, but not identical to, the native compound.

[0061] The term “Ubiquitin Ligase” refers to a family of proteins that facilitate the transfer of ubiquitin to a specific substrate protein, targeting the substrate protein for degradation. For example, cereblon is an E3 Ubiquitin Ligase protein that alone or in combination with an E2 ubiquitin-conjugating enzyme causes the attachment of ubiquitin to a lysine on a target protein, and subsequently targets the specific protein substrates for degradation by the proteasome. Thus, E3 ubiquitin ligase alone or in complex with an E2 ubiquitin conjugating enzyme is responsible for the transfer of ubiquitin to targeted proteins. In general, the ubiquitin ligase is involved in polyubiquitination such that a second ubiquitin is attached to the first; a third is attached to the second, and so forth. Polyubiquitination marks proteins for degradation by the proteasome. However, there are some ubiquitination events that are limited to mono-ubiquitination, in which only a single ubiquitin is added by the ubiquitin ligase to a substrate molecule. Mono-ubiquitinated proteins are not targeted to the proteasome for degradation, but may instead be altered in their cellular location or function, for example, via binding other proteins that have domains capable of binding ubiquitin. Further complicating matters, different lysines on ubiquitin can be targeted by an E3 to make chains. The most common lysine is Lys48 on the ubiquitin chain. This is the lysine used to make polyubiquitin, which is recognized by the proteasome.

[0062] The term "patient" or "subject" is used throughout the specification to describe a cell, tissue, or animal, preferably a mammal, e.g., a human or a domesticated animal, to whom treatment, including prophylactic treatment, with the compositions according to the present disclosure is provided. For treatment of those infections, conditions or disease states which are specific for a specific animal such as a human patient, the term patient refers to that specific animal, including a domesticated animal such as a dog or cat or a farm animal such as a horse, cow, sheep, etc. In general, in the present disclosure, the term patient refers to a human patient unless otherwise stated or implied from the context of the use of the term.

[0063] **Compounds and Compositions**

[0064] In one aspect, the present disclosure provides compounds useful for regulating protein activity. The composition comprises a ubiquitin pathway protein binding moiety (preferably for an E3 ubiquitin ligase, alone or in complex with an E2 ubiquitin conjugating enzyme which is responsible for the transfer of ubiquitin to targeted proteins) according to a defined chemical structure and a protein targeting moiety which are linked or coupled together, preferably through a linker, wherein the ubiquitin pathway protein binding moiety recognizes a ubiquitin pathway protein and the targeting moiety recognizes a target protein (e.g., androgen receptor). Such compounds may be referred to herein as PROTAC compounds or PROTACs.

[0065] In one aspect, the description provides compounds comprising an E3 Ubiquitin Ligase binding moiety ("ULM") that is a cereblon E3 Ubiquitin Ligase binding moiety ("CLM"). In one embodiment, the CLM is coupled to a chemical linker group (L) according to the structure:

L-CLM **(I)**

wherein L is a chemical linker group and CLM is a cereblon E3 Ubiquitin Ligase binding moiety. The number and/or relative positions of the moieties in the compounds illustrated herein is provided by way of example only. As would be understood by the skilled artisan, compounds as described herein can be synthesized with any desired number and/or relative position of the respective functional moieties.

[0066] The terms ULM and CLM are used in their inclusive sense unless the context indicates otherwise. For example, the term ULM is inclusive of all ULMs, including those that bind cereblon (i.e., CLMs). Further, the term CLM is inclusive of all possible cereblon E3 Ubiquitin Ligase binding moieties.

[0067] In another embodiment, the description provides a compound which comprises a plurality of CLMs coupled directly or via a chemical linker moiety (L). For example, a compound having two CLMs can be depicted as:

CLM-CLM (**II**) or

CLM-L-CLM (**III**).

[0068] In certain embodiments, where the compound comprises multiple CLMs, the CLMs are identical. In additional embodiments, the compound comprising a plurality of CLMs further comprises at least one PTM coupled to a CLM directly or via a chemical linker (L) or both. In certain additional embodiments, the compound comprising a plurality of CLMs further comprises multiple PTMs. In still additional embodiments, the PTMs are the same or, optionally, different. In still further embodiments, wherein the PTMs are different the respective PTMs may bind the same protein target or bind specifically to a different protein target.

[0069] In additional embodiments, the description provides a compound comprising at least two different CLMs coupled directly or via a chemical linker (L) or both. For example, such a compound having two different CLMs can be depicted as:

CLM-CLM' (**IV**) or

CLM-L-CLM' (**V**)

[0070] wherein CLM' indicates a cereblon E3 Ubiquitin Ligase binding moiety that is structurally different from CLM. In certain embodiments, the compound may comprise a plurality of CLMs and/or a plurality of CLM's. In further embodiments, the compound comprising at least two different CLMs, a plurality of CLMs, and/or a plurality of CLM's further comprises at least one PTM coupled to a CLM or a CLM' directly or via a chemical linker or both. In any of the embodiments described herein, a compound comprising at least two different CLMs can further comprise multiple PTMs. In still additional embodiments, the PTMs are the same or, optionally, different. In still further embodiments, wherein the PTMs are different the respective PTMs may bind the same protein target or bind specifically to a different protein target. In still further embodiments, the PTM itself is a ULM or CLM (or ULM' or CLM')

[0071] In another aspect, the description provides bifunctional or PROTAC compounds, which comprise an E3 Ubiqutin Ligase binding moiety ("ULM") that is a cereblon E3 Ubiquitin Ligase binding moiety ("CLM"), and a moiety that binds a target protein (i.e. a

protein/polypeptide targeting ligand or "PTM" group) that is an AR binding moiety ("ABM"). In one embodiment the structure of the bifunctional compound can be depicted as:

ABM – CLM **(VI),**

[0072] The respective positions of the ABM and CLM moieties as well as their number as illustrated herein is provided by way of example only and is not intended to limit the compounds in any way. As would be understood by the skilled artisan, the bifunctional compounds as described herein can be synthesized such that the number and position of the respective functional moieties can be varied as desired.

[0073] In certain embodiments, the compounds having the following general structure: ABM-L, wherein ABM is an AR binding moiety as described herein, and L is a chemical linker moiety, e.g., a linker as described herein, or optionally a bond.

[0074] In certain embodiments the bifunctional compound further comprises a chemical linker ("L"). In this example, the structure of the bifunctional compunds can be depicted as:

ABM – L - CLM **(VII)**

[0075] where ABM is an AR binding moiety as described herein, CLM is a cereblon E3 ligase binding moiety as described herein, and L is a chemical linker moiety, e.g., a linker as described herein, or optionally a bond, that links the ABM and CLM.

[0076] Further, CLM is inclusive of all possible cereblon E3 Ubiquitin Ligase binding moieties. The CLM group and ABM group may be covalently linked to the linker group through any covalent bond which is appropriate and stable to the chemistry of the linker.

[0077] In certain embodiments, the CLM comprises a moiety that is a ligand of the cereblon E3 Ubiquitin Ligase (CRBN). In certain embodiments, the CLM comprises a chemotype from the "imide" class of molecules. In certain additional embodiments, the CLM comprises a phthalimido group or an analog or derivative thereof. In still additional embodiments, the CLM comprises a phthalimido-glutarimide group or an analog or derivative thereof. In still other embodiments, the CLM comprises a member of the group consisting of thalidomide, lenalidomide, pomalidomide, and analogs or derivatives thereof.

[0078] It will be understood that the general structures are exemplary and the respective moieties can be arranged in any desired order or configuration, e.g., CLM-L-ABM, and CLM-L-ABM respectively. In certain additional embodiments, the compounds comprise a plurality of E3 ligase binding moieties and/or a plurality of ABMs.

[0079] In certain embodiments, the compounds as described herein comprise multiple ABMs (targeting the same or different locations of the AR), multiple CLMs, one or more ULMs (i.e., moieties that bind specifically to another E3 Ubiquitin Ligase, e.g., VHL) or a combination thereof. In any of the aspects of embodiments described herein, the ABMs, CLMs, and ULMs can be coupled directly or via one or more chemical linkers or a combination thereof. In additional embodiments, where a compound has multiple ULMs, the ULMs can be for the same E3 Ubiquitin Ligase or each respective ULM can bind specifically to a different E3 Ubiquitin Ligase. In still further embodiments, where a compound has multiple ABMs, the ABMs are the same or, optionally, different.

[0080] In certain embodiments, where the compound comprises multiple CLMs, the CLMs are identical or, optionally, different. In additional embodiments, the compound comprising a plurality of CLMs further comprises at least one ABM coupled to a CLM directly or via a chemical linker (L) or both. In certain additional embodiments, the compound comprising a plurality of CLMs further comprises multiple ABMs. In still additional embodiments, the ABMs are the same or, optionally, different.

[0081] In certain embodiments, ABM alone, without forming ABM-L-CLM, provides desired properties in regulating protein activity.


[0082] In any of the aspects or embodiments of compounds described herein, unless indicated otherwise, the compounds are intended to encompass pharmaceutically acceptable salts, enantiomers, stereoisomers, solvates or polymorphs thereof.

[0083] In certain embodiments, the compounds as described herein comprise multiple PTMs (targeting the same or different protein targets), multiple ABMs, multiple CLMs, one or more ULMs (i.e., moieties that bind specifically to another E3 Ubiquitin Ligase, e.g., cereblon) or a combination thereof. In any of the aspects of embodiments described herein, the PTMs, ABMs, CLMs, and ULMs can be coupled directly or via one or more chemical linkers or a combination thereof. In additional embodiments, where a compound has multiple ULMs, the ULMs can be for the same E3 Ubiquintin Ligase or each respective ULM can bind specifically to a different E3 Ubiquitin Ligase. In still further embodiments, where a compound has multiple PTMs, the PTMs can bind the same target protein or each respective PTM can bind specifically to a different target protein.


Exemplary CLMs

[0084] Neo-imide Compounds


[0085] In one aspect the description provides compounds useful for binding and/or inhibiting cereblon. In certain embodiments, the compound or CLM is selected from the group consisting of chemical structures:




(a)




(b)




(c)



(d)



(e) and



(f),

wherein:

W is independently selected from the group consisting of CH_2, CHR, C=O, SO_2, NH, and N-alkyl (linear, branched, optionally substituted);

Y is independently selected from the group consisting of CH_2, -C=CR', NH, N-alkyl, N-aryl, N-hetaryl, N-cycloalkyl, N-heterocyclyl, O, and S;

X and Z are each independently O, S, or H_2, except that both X and Z cannot be H_2;

G and G' are independently selected from the group consisting of H, alkyl (linear, branched, optionally substituted with R'), OH, R'OCOOR, R'OCONRR", CH_2-heterocyclyl optionally substituted with R', and benzyl optionally substituted with R';

Q1 – Q4 are each independently C optionally substituted with R', N, or N-oxide;

A is independently selected from the group consisting of alkyl (linear, branched, optionally substituted), cycloalkyl (optionally substituted), Cl, H and F;

R comprises, but is not limited to: -CONR'R", -OR', -NR'R", -SR', -SO_2R', -SO_2NR'R", -CR'R"-, -CR'NR'R"-, -aryl, -hetaryl, -alkyl (linear, branched, optionally substituted), -cycloalkyl, -heterocyclyl, -P(O)(OR')R", -P(O)R'R", -OP(O)(OR')R", -OP(O)R'R", -Cl, -F, -Br, -I, -CF_3, -CN, -NR'SO_2NR'R", -NR'CONR'R", -CONR'COR", -NR'C(=N-CN)NR'R", -C(=N-CN)NR'R", -NR'C(=N-CN)R", -NR'C(=C-NO_2)NR'R", -SO_2NR'COR", -NO_2, -CO_2R', -C(C=N-OR')R", -CR'=CR'R", -CCR', -S(C=O)(C=N-R')R", -SF_5, -R'NR'R", (-R'O)$_n$R", or -OCF_3

R' and R" are each independently a bond, H, alkyl (linear, branched), cycloalkyl, aryl, hetaryl, heterocyclyl, or –C(=O)R, each of which is optionally substituted;


n is an integer from 1-10 (e.g., 1-4, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10);

~~~ represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and


R_n comprises 1-4 independent functional groups or atoms, and optionally, one of which is modified to be covalently joined to an ABM, a chemical linker group (L), a ULM, a CLM (CLM'), or combination thereof.

[0086] Exemplary CLMs


[0087] In any of the compounds described herein, the CLM comprises a chemical structure selected from the group:
~~~




(a)




(b)




(c)



(d)



(e) and



(f),

wherein:

W is CH_2, CHR, C=O, SO_2, NH, or N-alkyl (linear, branched, optionally substituted);

X and Z are each independently O, S or H_2, except that both X and Z cannot be H_2;

Y is independently selected from the group CH_2, -C=CR', NH, N-alkyl, N-aryl, N-hetaryl, N-cycloalkyl, N-heterocyclyl, O, and S;

G is H, alkyl (linear, branched), OH, R'OCOOR, R'OCONRR", CH_2-heterocyclyl, heterocyclic, aryl, or benzyl, each optionally substituted with R';

Q1 – Q4 are each independently C is optionally substituted with R', N, or N-oxide;

A is C_1-C_6 alkyl (linear, branched, optionally substituted), cycloalkyl (optionally substituted), H, Cl or F;

R is -CONR'R", -OR', -NR'R", -SR', -SO2R', -SO2NR'R", -CR'R"-, -CR'NR'R"-, (-CR'O)$_n$R", -aryl, -heteroaryl, -alkyl (linear, branched, optionally substituted), -cycloalkyl, -heterocyclyl, -P(O)(OR')R", -P(O)R'R", -OP(O)(OR')R", -OP(O)R'R", -Cl, -F, -Br, -I, -CF3, -CN, -NR'SO2NR'R", -NR'CONR'R", -CONR'COR", -NR'C(=N-CN)NR'R", -C(=N-CN)NR'R", -NR'C(=N-CN)R", -NR'C(=C-NO2)NR'R", -SO2NR'COR", -NO2, -CO2R', -C(C=N-OR')R", -CR'=CR'R", -CCR', -S(C=O)(C=N-R')R", alicylic, heterocyclic, -SF5, or -OCF3;


R' and R" are each independently a bond, H, N, N-oxide, alkyl (linear, branched), cycloalkyl, aryl, heteroaryl, heterocyclic, -C(=O)R, or, heterocyclyl, each of which is optionally substituted;

n is an integer from 1-10;

〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and

R_n comprises 1-4 independent functional groups or atoms, and optionally, one of which is modified to be covalently joined to an ABM, a chemical linker group (L), a ULM, a CLM (CLM'), or combination thereof.

[0088] In certain embodiments described herein, the CLM or ULM comprises a chemical structure selected from the group:



wherein:

W is independently selected from the group consisting of CH_2, C=O, NH, and N-alkyl;

R is a H, methyl, or C1-C6 alkyl (linear, branched, optionally substituted);

〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and

Rn comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to an ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof.

[0089] The term "independently" is used herein to indicate that the variable, which is independently applied, varies independently from application to application.

[0090] The term "alkyl" shall mean within its context a linear, branch-chained or cyclic fully saturated hydrocarbon radical or alkyl group, preferably a C_1-C_{10}, more preferably a C_1-C_6, alternatively a C_1-C_3 alkyl group, which may be optionally substituted. Examples of alkyl groups are methyl, ethyl, n-butyl, sec-butyl, n-hexyl, n-heptyl, n-octyl, n-nonyl, n-decyl, isopropyl, 2-methylpropyl, cyclopropyl, cyclopropylmethyl, cyclobutyl, cyclopentyl, cyclopentylethyl, cyclohexylethyl and cyclohexyl, among others. In certain embodiments, the alkyl group is end-capped with a halogen group (At, Br, Cl, F, or I). In certain preferred embodiments, compounds according to the present disclosure, which may be used to covalently bind to dehalogenase enzymes. These compounds generally contain a side chain (often linked through a polyethylene glycol group) which terminates in an alkyl group which has a halogen substituent (often chlorine or bromine) on its distal end which results in covalent binding of the compound containing such a moiety to the protein.

[0091] The term "Alkenyl" refers to linear, branch-chained or cyclic C_2-C_{10} (preferably C_2-C_6) hydrocarbon radicals containing at least one C=C bond.

[0092] The term "Alkynyl" refers to linear, branch-chained or cyclic C_2-C_{10} (preferably C_2-C_6) hydrocarbon radicals containing at least one C≡C bond.

[0093] The term "alkylene" when used, refers to a $-(CH_2)_n-$ group (n is an integer generally from 0-6), which may be optionally substituted. When substituted, the alkylene group preferably is substituted on one or more of the methylene groups with a C_1-C_6 alkyl group (including a cyclopropyl group or a t-butyl group), but may also be substituted with one or more halo groups, preferably from 1 to 3 halo groups or one or two hydroxyl groups, O-(C_1-C_6 alkyl) groups or amino acid sidechains as otherwise disclosed herein. In certain embodiments, an alkylene group may be substituted with a urethane or alkoxy group (or other group) which is further substituted with a polyethylene glycol chain (of from 1 to 10, preferably 1 to 6, often 1 to 4 ethylene glycol units) to which is substituted (preferably, but not exclusively on the distal end of the polyethylene glycol chain) an alkyl chain substituted with a single halogen group,

preferably a chlorine group. In still other embodiments, the alkylene (often, a methylene) group, may be substituted with an amino acid sidechain group such as a sidechain group of a natural or unnatural amino acid, for example, alanine, β-alanine, arginine, asparagine, aspartic acid, cysteine, cystine, glutamic acid, glutamine, glycine, phenylalanine, histidine, isoleucine, lysine, leucine, methionine, proline, serine, threonine, valine, tryptophan or tyrosine.

[0094] The term "unsubstituted" shall mean substituted only with hydrogen atoms. A range of carbon atoms which includes C_0 means that carbon is absent and is replaced with H. Thus, a range of carbon atoms which is C_0-C_6 includes carbons atoms of 1, 2, 3, 4, 5 and 6 and for C_0, H stands in place of carbon.

[0095] The term "substituted" or "optionally substituted" shall mean independently (i.e., where more than substituent occurs, each substituent is independent of another substituent) one or more substituents (independently up to five substituents, preferably up to three substituents, often 1 or 2 substituents on a moiety in a compound according to the present disclosure and may include substituents which themselves may be further substituted) at a carbon (or nitrogen) position anywhere on a molecule within context, and includes as substituents hydroxyl, thiol, carboxyl, cyano (C≡N), nitro (NO_2), halogen (preferably, 1, 2 or 3 halogens, especially on an alkyl, especially a methyl group such as a trifluoromethyl), an alkyl group (preferably, C_1-C_{10}, more preferably, C_1-C_6), aryl (especially phenyl and substituted phenyl for example benzyl or benzoyl), alkoxy group (preferably, C_1-C_6 alkyl or aryl, including phenyl and substituted phenyl), thioether (C_1-C_6 alkyl or aryl), acyl (preferably, C_1-C_6 acyl), ester or thioester (preferably, C_1-C_6 alkyl or aryl) including alkylene ester (such that attachment is on the alkylene group, rather than at the ester function which is preferably substituted with a C_1-C_6 alkyl or aryl group), preferably, C_1-C_6 alkyl or aryl, halogen (preferably, F or Cl), amine (including a five- or six-membered cyclic alkylene amine, further including a C_1-C_6 alkyl amine or a C_1-C_6 dialkyl amine which alkyl groups may be substituted with one or two hydroxyl groups) or an optionally substituted –N(C_0-C_6 alkyl)C(O)(O-C_1-C_6 alkyl) group (which may be optionally substituted with a polyethylene glycol chain to which is further bound an alkyl group containing a single halogen, preferably chlorine substituent), hydrazine, amido, which is preferably substituted with one or two C_1-C_6 alkyl groups (including a carboxamide which is optionally substituted with one or two C_1-C_6 alkyl groups), alkanol (preferably, C_1-C_6 alkyl or aryl), or alkanoic acid (preferably, C_1-C_6 alkyl or aryl). Substituents according to the present disclosure may include, for example –

$SiR_1R_2R_3$ groups where each of R_1 and R_2 is as otherwise described herein and R_3 is H or a C_1-C_6 alkyl group, preferably R_1, R_2, R_3 in this context is a C_1-C_3 alkyl group (including an isopropyl or t-butyl group). Each of the above-described groups may be linked directly to the substituted moiety or alternatively, the substituent may be linked to the substituted moiety (preferably in the case of an aryl or heteraryl moiety) through an optionally substituted $-(CH_2)_m-$ or alternatively an optionally substituted $-(OCH_2)_m-$, $-(OCH_2CH_2)_m-$ or $-(CH_2CH_2O)_m-$ group, which may be substituted with any one or more of the above-described substituents. Alkylene groups $-(CH_2)_m-$ or $-(CH_2)_n-$ groups or other chains such as ethylene glycol chains, as identified above, may be substituted anywhere on the chain. Preferred substituents on alkylene groups include halogen or C_1-C_6 (preferably C_1-C_3) alkyl groups, which may be optionally substituted with one or two hydroxyl groups, one or two ether groups (O-C_1-C_6 groups), up to three halo groups (preferably F), or a sideshain of an amino acid as otherwise described herein and optionally substituted amide (preferably carboxamide substituted as described above) or urethane groups (often with one or two C_0-C_6 alkyl substitutents, which group(s) may be further substituted). In certain embodiments, the alkylene group (often a single methylene group) is substituted with one or two optionally substituted C_1-C_6 alkyl groups, preferably C_1-C_4 alkyl group, most often methyl or O-methyl groups or a sidechain of an amino acid as otherwise described herein. In the present disclosure, a moiety in a molecule may be optionally substituted with up to five substituents, preferably up to three substituents. Most often, in the present disclosure moieties which are substituted are substituted with one or two substituents.

[0096] The term "substituted" (each substituent being independent of any other substituent) shall also mean within its context of use C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, amido, carboxamido, sulfone, including sulfonamide, keto, carboxy, C_1-C_6 ester (oxyester or carbonylester), C_1-C_6 keto, urethane $-O-C(O)-NR_1R_2$ or $-N(R_1)-C(O)-O-R_1$, nitro, cyano and amine (especially including a C_1-C_6 alkylene-NR_1R_2, a mono- or di- C_1-C_6 alkyl substituted amines which may be optionally substituted with one or two hydroxyl groups). Each of these groups contain unless otherwise indicated, within context, between 1 and 6 carbon atoms. In certain embodiments, preferred substituents will include for example, -NH-, -NHC(O)-, -O-, =O, $-(CH_2)_m-$ (here, m and n are in context, 1, 2, 3, 4, 5 or 6), -S-, -S(O)-, SO_2- or –NH-C(O)-NH-, $-(CH_2)_nOH$, $-(CH_2)_nSH$, $-(CH_2)_nCOOH$, C_1-C_6 alkyl, $-(CH_2)_nO-(C_1$-C_6 alkyl), $-(CH_2)_nC(O)-(C_1$-C_6 alkyl), $-(CH_2)_nOC(O)-(C_1$-C_6 alkyl), $-(CH_2)_nC(O)O-(C_1$-C_6 alkyl), $-(CH_2)_nNHC(O)-R_1$, -

$(CH_2)_nC(O)$-NR_1R_2, -$(OCH_2)_nOH$, -$(CH_2O)_nCOOH$, C_1-C_6 alkyl, -$(OCH_2)_nO$-(C_1-C_6 alkyl), -$(CH_2O)_nC(O)$-(C_1-C_6 alkyl), -$(OCH_2)_nNHC(O)$-R_1, -$(CH_2O)_nC(O)$-NR_1R_2, -$S(O)_2$-R_S, -$S(O)$-R_S (R_S is C_1-C_6 alkyl or a –$(CH_2)_m$-NR_1R_2 group), NO_2, CN or halogen (F, Cl, Br, I, preferably F or Cl), depending on the context of the use of the substituent. R_1 and R_2 are each, within context, H or a C_1-C_6 alkyl group (which may be optionally substituted with one or two hydroxyl groups or up to three halogen groups, preferably fluorine). The term "substituted" shall also mean, within the chemical context of the compound defined and substituent used, an optionally substituted aryl or heteroaryl group or an optionally substituted heterocyclic group as otherwise described herein. Alkylene groups may also be substituted as otherwise disclosed herein, preferably with optionally substituted C_1-C_6 alkyl groups (methyl, ethyl or hydroxymethyl or hydroxyethyl is preferred, thus providing a chiral center), a sidechain of an amino acid group as otherwise described herein, an amido group as described hereinabove, or a urethane group O-$C(O)$-NR_1R_2 group where R_1 and R_2 are as otherwise described herein, although numerous other groups may also be used as substituents. Various optionally substituted moieties may be substituted with 3 or more substituents, preferably no more than 3 substituents and preferably with 1 or 2 substituents. It is noted that in instances where, in a compound at a particular position of the molecule substitution is required (principally, because of valency), but no substitution is indicated, then that substituent is construed or understood to be H, unless the context of the substitution suggests otherwise.

[0097] The term "aryl" or "aromatic", in context, refers to a substituted (as otherwise described herein) or unsubstituted monovalent aromatic radical having a single ring (e.g., benzene, phenyl, benzyl) or condensed rings (e.g., naphthyl, anthracenyl, phenanthrenyl, etc.) and can be bound to the compound according to the present disclosure at any available stable position on the ring(s) or as otherwise indicated in the chemical structure presented. Other examples of aryl groups, in context, may include heterocyclic aromatic ring systems, "heteroaryl" groups having one or more nitrogen, oxygen, or sulfur atoms in the ring (moncyclic) such as imidazole, furyl, pyrrole, furanyl, thiene, thiazole, pyridine, pyrimidine, pyrazine, triazole, oxazole or fused ring systems such as indole, quinoline, indolizine, azaindolizine, benzofurazan, etc., among others, which may be optionally substituted as described above. Among the heteroaryl groups which may be mentioned include nitrogen-containing heteroaryl groups such as pyrrole, pyridine, pyridone, pyridazine, pyrimidine, pyrazine, pyrazole,


imidazole, triazole, triazine, tetrazole, indole, isoindole, indolizine, azaindolizine, purine, indazole, quinoline, dihydroquinoline, tetrahydroquinoline, isoquinoline, dihydroisoquinoline, tetrahydroisoquinoline, quinolizine, phthalazine, naphthyridine, quinoxaline, quinazoline, cinnoline, pteridine, imidazopyridine, imidazotriazine, pyrazinopyridazine, acridine, phenanthridine, carbazole, carbazoline, pyrimidine, phenanthroline, phenacene, oxadiazole, benzimidazole, pyrrolopyridine, pyrrolopyrimidine and pyridopyrimidine; sulfur-containing aromatic heterocycles such as thiophene and benzothiophene; oxygen-containing aromatic heterocycles such as furan, pyran, cyclopentapyran, benzofuran and isobenzofuran; and aromatic heterocycles comprising 2 or more hetero atoms selected from among nitrogen, sulfur and oxygen, such as thiazole, thiadizole, isothiazole, benzoxazole, benzothiazole, benzothiadiazole, phenothiazine, isoxazole, furazan, phenoxazine, pyrazoloxazole, imidazothiazole, thienofuran, furopyrrole, pyridoxazine, furopyridine, furopyrimidine, thienopyrimidine and oxazole, among others, all of which may be optionally substituted.

[0098] The term "substituted aryl" refers to an aromatic carbocyclic group comprised of at least one aromatic ring or of multiple condensed rings at least one of which being aromatic, wherein the ring(s) are substituted with one or more substituents. For example, an aryl group can comprise a substituent(s) selected from: $-(CH_2)_nOH$, $-(CH_2)_n-O-(C_1-C_6)$alkyl, $-(CH_2)_n-O-(CH_2)_n-(C_1-C_6)$alkyl, $-(CH_2)_n-C(O)(C_0-C_6)$ alkyl, $-(CH_2)_n-C(O)O(C_0-C_6)$alkyl, $-(CH_2)_n-OC(O)(C_0-C_6)$alkyl, amine, mono- or di-(C_1-C_6 alkyl) amine wherein the alkyl group on the amine is optionally substituted with 1 or 2 hydroxyl groups or up to three halo (preferably F, Cl) groups, OH, COOH, C_1-C_6 alkyl, preferably CH_3, CF_3, OMe, OCF_3, NO_2, or CN group (each of which may be substituted in ortho-, meta- and/or para- positions of the phenyl ring, preferably para-), an optionally substituted phenyl group (the phenyl group itself is preferably substituted with a linker group attached to a ABM group, including a ULM group), and/or at least one of F, Cl, OH, COOH, CH_3, CF_3, OMe, OCF_3, NO_2, or CN group (in ortho-, meta- and/or para- positions of the phenyl ring, preferably para-), a naphthyl group, which may be optionally substituted, an optionally substituted heteroaryl, preferably an optionally substituted isoxazole including a methylsubstituted isoxazole, an optionally substituted oxazole including a methylsubstituted oxazole, an optionally substituted thiazole including a methyl substituted thiazole, an optionally substituted isothiazole including a methyl substituted isothiazole, an optionally substituted pyrrole including a methylsubstituted pyrrole, an optionally substituted imidazole including a

methylimidazole, an optionally substituted benzimidazole or methoxybenzylimidazole, an optionally substituted oximidazole or methyloximidazole, an optionally substituted diazole group, including a methyldiazole group, an optionally substituted triazole group, including a methylsubstituted triazole group, an optionally substituted pyridine group, including a halo- (preferably, F) or methylsubstitutedpyridine group or an oxapyridine group (where the pyridine group is linked to the phenyl group by an oxygen), an optionally substituted furan, an optionally substituted benzofuran, an optionally substituted dihydrobenzofuran, an optionally substituted indole, indolizine or azaindolizine (2, 3, or 4-azaindolizine), an optionally substituted quinoline, and combinations thereof.

[0099] "Carboxyl" denotes the group --C(O)OR, where R is hydrogen, alkyl, substituted alkyl, aryl, substituted aryl, heteroaryl or substituted heteroaryl , whereas these generic substituents have meanings which are identical with definitions of the corresponding groups defined herein.

[0100] The term “heteroaryl”or “hetaryl” can mean but is in no way limited to an optionally substituted quinoline (which may be attached to the pharmacophore or substituted on any carbon atom within the quinoline ring), an optionally substituted indole (including dihydroindole), an optionally substituted indolizine, an optionally substituted azaindolizine (2, 3 or 4-azaindolizine) an optionally substituted benzimidazole, benzodiazole, benzoxofuran, an optionally substituted imidazole, an optionally substituted isoxazole, an optionally substituted oxazole (preferably methyl substituted), an optionally substituted diazole, an optionally substituted triazole, a tetrazole, an optionally substituted benzofuran, an optionally substituted thiophene, an optionally substituted thiazole (preferably methyl and/or thiol substituted), an optionally substituted isothiazole, an optionally substituted triazole (preferably a 1,2,3-triazole substituted with a methyl group, a triisopropylsilyl group, an optionally substituted -$(CH_2)_m$-O-C_1-C_6 alkyl group or an optionally substituted -$(CH_2)_m$-C(O)-O-C_1-C_6 alkyl group), an optionally substituted pyridine (2-, 3, or 4-pyridine) or a group according to the chemical structure:



wherein

S^c is CHR^{SS}, NR^{URE}, or O;

R^{HET} is H, CN, NO_2, halo (preferably Cl or F), optionally substituted C_1-C_6 alkyl (preferably substituted with one or two hydroxyl groups or up to three halo groups (e.g. CF_3), optionally substituted O(C_1-C_6 alkyl) (preferably substituted with one or two hydroxyl groups or up to three halo groups) or an optionally substituted acetylenic group –C≡C-R_a where R_a is H or a C_1-C_6 alkyl group (preferably C_1-C_3 alkyl);

R^{SS} is H, CN, NO_2, halo (preferably F or Cl), optionally substituted C_1-C_6 alkyl (preferably substituted with one or two hydroxyl groups or up to three halo groups), optionally substituted O-(C_1-C_6 alkyl) (preferably substituted with one or two hydroxyl groups or up to three halo groups) or an optionally substituted -C(O)(C_1-C_6 alkyl) (preferably substituted with one or two hydroxyl groups or up to three halo groups);

R^{URE} is H, a C_1-C_6 alkyl (preferably H or C_1-C_3 alkyl) or a –C(O)(C_1-C_6 alkyl), each of which groups is optionally substituted with one or two hydroxyl groups or up to three halogen, preferably fluorine groups, or an optionally substituted heterocycle, for example piperidine, morpholine, pyrrolidine, tetrahydrofuran, tetrahydrothiophene, piperidine, piperazine, each of which is optionally substituted, and

Y^C is N or C-R^{YC}, where R^{YC} is H, OH, CN, NO_2, halo (preferably Cl or F), optionally substituted C_1-C_6 alkyl (preferably substituted with one or two hydroxyl groups or up to three halo groups (e.g. CF_3), optionally substituted O(C_1-C_6 alkyl) (preferably substituted

with one or two hydroxyl groups or up to three halo groups) or an optionally substituted acetylenic group –C≡C-R_a where R_a is H or a C_1-C_6 alkyl group (preferably C_1-C_3 alkyl).

[0101] The terms "aralkyl" and "heteroarylalkyl" refer to groups that comprise both aryl or, respectively, heteroaryl as well as alkyl and/or heteroalkyl and/or carbocyclic and/or heterocycloalkyl ring systems according to the above definitions.

[0102] The term "arylalkyl" as used herein refers to an aryl group as defined above appended to an alkyl group defined above. The arylalkyl group is attached to the parent moiety through an alkyl group wherein the alkyl group is one to six carbon atoms. The aryl group in the arylalkyl group may be substituted as defined above.

[0103] The term "Heterocycle" refers to a cyclic group which contains at least one heteroatom, e.g., N, O or S, and may be aromatic (heteroaryl) or non-aromatic. Thus, the heteroaryl moieties are subsumed under the definition of heterocycle, depending on the context of its use. Exemplary heteroaryl groups are described hereinabove.

[0104] Exemplary heterocyclics include: azetidinyl, benzimidazolyl, 1,4- benzodioxanyl, 1,3-benzodioxolyl, benzoxazolyl, benzothiazolyl, benzothienyl, dihydroimidazolyl, dihydropyranyl, dihydrofuranyl, dioxanyl, dioxolanyl, ethyleneurea, 1,3-dioxolane, 1,3-dioxane, 1,4-dioxane, furyl, homopiperidinyl, imidazolyl, imidazolinyl, imidazolidinyl, indolinyl, indolyl, isoquinolinyl, isothiazolidinyl, isothiazolyl, isoxazolidinyl, isoxazolyl, morpholinyl, naphthyridinyl, oxazolidinyl, oxazolyl, pyridone, 2-pyrrolidone, pyridine, piperazinyl, , N-methylpiperazinyl, piperidinyl, phthalimide, succinimide, pyrazinyl, pyrazolinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, pyrrolyl, quinolinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydroquinoline, thiazolidinyl, thiazolyl, thienyl, tetrahydrothiophene, oxane, oxetanyl, oxathiolanyl, thiane among others.

[0105] Heterocyclic groups can be optionally substituted with a member selected from the group consisting of alkoxy, substituted alkoxy, cycloalkyl, substituted cycloalkyl, cycloalkenyl, substituted cycloalkenyl, acyl, acylamino, acyloxy, amino, substituted amino, aminoacyl, aminoacyloxy, oxyaminoacyl, azido, cyano, halogen, hydroxyl, keto, thioketo, carboxy, carboxyalkyl, thioaryloxy, thioheteroaryloxy, thioheterocyclooxy, thiol, thioalkoxy, substituted thioalkoxy, aryl, aryloxy, heteroaryl, heteroaryloxy, heterocyclic, heterocyclooxy, hydroxyamino, alkoxyamino, nitro, —SO-alkyl, —SO-substituted alkyl, —SOaryl, —SO-heteroaryl, —SO2-alkyl, —SO2-substituted alkyl, —SO2-aryl, oxo (=O), and -SO2-heteroaryl.

Such heterocyclic groups can have a single ring or multiple condensed rings. Examples of nitrogen heterocycles and heteroaryls include, but are not limited to, pyrrole, imidazole, pyrazole, pyridine, pyrazine, pyrimidine, pyridazine, indolizine, isoindole, indole, indazole, purine, quinolizine, isoquinoline, quinoline, phthalazine, naphthylpyridine, quinoxaline, quinazoline, cinnoline, pteridine, carbazole, carboline, phenanthridine, acridine, phenanthroline, isothiazole, phenazine, isoxazole, phenoxazine, phenothiazine, imidazolidine, imidazoline, piperidine, piperazine, indoline, morpholino, piperidinyl, tetrahydrofuranyl, and the like as well as N-alkoxy-nitrogen containing heterocycles. The term "heterocyclic" also includes bicyclic groups in which any of the heterocyclic rings is fused to a benzene ring or a cyclohexane ring or another heterocyclic ring (for example, indolyl, quinolyl, isoquinolyl, tetrahydroquinolyl, and the like).


[0106] The term “cycloalkyl” can mean but is in no way limited to univalent groups derived from monocyclic or polycyclic alkyl groups or cycloalkanes, as defnied herein, e.g., saturated monocyclic hydrocarbon groups having from three to twenty carbon atoms in the ring, including, but not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like. The term "substituted cycloalkyl" can mean but is in no way limited to a monocyclic or polycyclic alkyl group and being substituted by one or more substituents, for example, amino, halogen, alkyl, substituted alkyl, carbyloxy, carbylmercapto, aryl, nitro, mercapto or sulfo, whereas these generic substituent groups have meanings which are identical with definitions of the corresponding groups as defined in this legend.


[0107] "Heterocycloalkyl" refers to a monocyclic or polycyclic alkyl group in which at least one ring carbon atom of its cyclic structure being replaced with a heteroatom selected from the group consisting of N, O, S or P. "Substituted heterocycloalkyl" refers to a monocyclic or polycyclic alkyl group in which at least one ring carbon atom of its cyclic structure being replaced with a heteroatom selected from the group consisting of N, O, S or P and the group is containing one or more substituents selected from the group consisting of halogen, alkyl, substituted alkyl, carbyloxy, carbylmercapto, aryl, nitro, mercapto or sulfo, whereas these generic substituent group have meanings which are identical with definitions of the corresponding groups as defined in this legend.


[0108] The term "hydrocarbyl" shall mean a compound which contains carbon and hydrogen and which may be fully saturated, partially unsaturated or aromatic and includes aryl groups, alkyl groups, alkenyl groups and alkynyl groups.


[0109] In any of the embodiments described herein, the W, X, Y, Z, G, G', R, R', R'', Q1-Q4, A, and Rn can independently be covalently coupled to a linker and/or a linker to which is attached one or more ABM, ULM, CLM or CLM' groups.


[0110] More specifically, non-limiting examples of CLMs include those shown below as well as those 'hybrid' molecules that arise from the combination of 1 or more of the different features shown in the molecules below, wherein Rn comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to a ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof..











Exemplary Linkers

[0111] In certain embodiments, the compounds as described herein include one or more ABM chemically linked or coupled to one or more ULMs or CLMs via a chemical linker (L). In certain embodiments, the linker group L is a group comprising one or more covalently connected structural units (e.g., -$A_{1...}A_q$- or –A_q-), wherein A_1 is a group coupled to ABM, and A_q is a group coupled to ULM.

[0112] In certain embodiments, the linker group L is selected from A_q-;

A_q is a group which is connected to a ULM or ABM moiety; and

q is an integer greater than or equal to 1,

wherein A_q is selected from the group consisting of, a bond, $CR^{L1}R^{L2}$, O, S, SO, SO_2, NR^{L3}, SO_2NR^{L3}, $SONR^{L3}$, $CONR^{L3}$, $NR^{L3}CONR^{L4}$, $NR^{L3}SO_2NR^{L4}$, CO, $CR^{L1}=CR^{L2}$, C≡C, $SiR^{L1}R^{L2}$, $P(O)R^{L1}$, $P(O)OR^{L1}$, $NR^{L3}C(=NCN)NR^{L4}$, $NR^{L3}C(=NCN)$, $NR^{L3}C(=CNO_2)NR^{L4}$, C_{3-11}cycloalkyl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, C_{5-13} spirocycloalkyl

optionally substituted with 0-9 R^{L1} and/or R^{L2} groups, C_{3-11}heterocyclyl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, C_{5-13} spiroheterocycloalkyl optionally substituted with 0-8 R^{L1} and/or R^{L2} groups, aryl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, heteroaryl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, where R^{L1} or R^{L2}, each independently are optionally linked to other groups to form cycloalkyl and/or heterocyclyl moiety, optionally substituted with 0-4 R^{L5} groups;

R^{L1}, R^{L2}, R^{L3}, R^{L4} and R^{L5} are, each independently, H, halo, C_{1-8}alkyl, OC_{1-8}alkyl, SC_{1-8}alkyl, NHC_{1-8}alkyl, $N(C_{1-8}alkyl)_2$, C_{3-11}cycloalkyl, aryl, heteroaryl, C_{3-11}heterocyclyl, OC_{1-8}cycloalkyl, SC_{1-8}cycloalkyl, NHC_{1-8}cycloalkyl, $N(C_{1-8}cycloalkyl)_2$, $N(C_{1-8}cycloalkyl)(C_{1-8}alkyl)$, OH, NH_2, SH, SO_2C_{1-8}alkyl, $P(O)(OC_{1-8}alkyl)(C_{1-8}alkyl)$, $P(O)(OC_{1-8}alkyl)_2$, CC-C_{1-8}alkyl, CCH, $CH=CH(C_{1-8}alkyl)$, $C(C_{1-8}alkyl)=CH(C_{1-8}alkyl)$, $C(C_{1-8}alkyl)=C(C_{1-8}alkyl)_2$, $Si(OH)_3$, $Si(C_{1-8}alkyl)_3$, $Si(OH)(C_{1-8}alkyl)_2$, COC_{1-8}alkyl, CO_2H, halogen, CN, CF_3, CHF_2, CH_2F, NO_2, SF_5, SO_2NHC_{1-8}alkyl, $SO_2N(C_{1-8}alkyl)_2$, $SONHC_{1-8}$alkyl, $SON(C_{1-8}alkyl)_2$, $CONHC_{1-8}$alkyl, $CON(C_{1-8}alkyl)_2$, $N(C_{1-8}alkyl)CONH(C_{1-8}alkyl)$, $N(C_{1-8}alkyl)CON(C_{1-8}alkyl)_2$, $NHCONH(C_{1-8}alkyl)$, $NHCON(C_{1-8}alkyl)_2$, $NHCONH_2$, $N(C_{1-8}alkyl)SO_2NH(C_{1-8}alkyl)$, $N(C_{1-8}alkyl)\ SO_2N(C_{1-8}alkyl)_2$, NH $SO_2NH(C_{1-8}alkyl)$, NH $SO_2N(C_{1-8}alkyl)_2$, NH SO_2NH_2.

[0113] In certain embodiments, q is an integer greater than or equal to 0. In certain embodiments, q is an integer greater than or equal to 1.

[0114] In certain embodiments, e.g., where q is greater than 2, A_q is a group which is connected to ULM, and A_1 and A_q are connected via structural units of the linker (L).

[0115] In certain embodiments, e.g., where q is 2, A_q is a group which is connected to A_1 and to a ULM.

[0116] In certain embodiments, e.g., where q is 1, the structure of the linker group L is –A_1–, and A_1 is a group which is connected to a ULM moiety and a ABM moiety.

[0117] In certain embodiments, the linker (L) comprises a group represented by a general structure selected from the group consisting of:

$-NR(CH_2)_n$-(lower alkyl)-, $-NR(CH_2)_n$-(lower alkoxyl)-, $-NR(CH_2)_n$-(lower alkoxyl)-OCH_2-, $-NR(CH_2)_n$-(lower alkoxyl)-(lower alkyl)-OCH_2-, $-NR(CH_2)_n$-(cycloalkyl)-(lower alkyl)-OCH_2-, $-NR(CH_2)_n$-(hetero cycloalkyl)-, $-NR(CH_2CH_2O)_n$-(lower alkyl)-O-CH_2-, $-NR(CH_2CH_2O)_n$-(hetero cycloalkyl)-O-CH_2-, $-NR(CH_2CH_2O)_n$-Aryl-O-CH_2-, $-NR(CH_2CH_2O)_n$-(hetero aryl)-O-CH_2-, $-NR(CH_2CH_2O)_n$-(cyclo alkyl)-O-(hetero aryl)-O-CH_2-, $-NR(CH_2CH_2O)_n$-

(cyclo alkyl)-O-Aryl-O-CH_2-, -NR(CH_2CH_2O)$_n$-(lower alkyl)-NH-Aryl-O-CH_2-, -NR(CH_2CH_2O)$_n$-(lower alkyl)-O-Aryl-CH_2, -NR(CH_2CH_2O)$_n$-cycloalkyl-O-Aryl-, -NR(CH_2CH_2O)$_n$-cycloalkyl-O-(hetero aryl)l-, -NR(CH2CH2)n-(cycloalkyl)-O-(heterocycle)-CH_2, -NR(CH2CH2)n-(heterocycle)-(heterocycle)-CH_2, -N(R1R2)-(heterocycle)-CH2; where

n can be 0 to 10;

R can be H, lower alkyl;

R1 and R2 can form a ring with the connecting N.

[0118] In certain embodiments, the linker (L) comprises a group represented by a general structure selected from the group consisting of:

-N(R)-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-OCH2-,


-O-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-OCH2-,

-O-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-O-;


-N(R)-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-O-;

-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-O-;


-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-OCH2-;




;




;




;




;




;




;



;



; and



; wherein


m, n, o, p, q, and r are each independently 0, 1, 2, 3, 4, 5, 6;

when the number is zero, there is no N-O or O-O bond

R is H, methyl or ethyl; and


X is H or F.


[0119] In certain embodiments, the linker (L) comprises a group represented by a general structure:





wherein m can be 2, 3, 4, 5.


In certain embodiments, the linker (L) comprises a group represented by a general structure selected from the group consisting of:









X = H, F


X = H, F









wherein n and m are each independently 0, 1, 2, 3, 4, 5, 6; and X is H, or F.

[0120] In additional embodiments, the linker group is optionally substituted (poly)ethyleneglycol having between 1 and about 100 ethylene glycol units, between about 1 and about 50 ethylene glycol units, between 1 and about 25 ethylene glycol units, between about 1 and 10 ethylene glycol units, between 1 and about 8 ethylene glycol units and 1 and 6 ethylene glycol units, between 2 and 4 ethylene glycol units,or optionally substituted alkyl groups interdispersed with optionally substituted, O, N, S, P or Si atoms. In certain embodiments, the linker is substituted with an aryl, phenyl, benzyl, alkyl, alkylene, or heterocycle group. In certain embodiments, the linker may be asymmetric or symmetrical.

[0121] In any of the embodiments of the compounds described herein, the linker group may be any suitable moiety as described herein. In one embodiment, the linker is a substituted or unsubstituted polyethylene glycol group ranging in size from about 1 to about 12 ethylene glycol units, between 1 and about 10 ethylene glycol units, about 2 about 6 ethylene glycol units, between about 2 and 5 ethylene glycol units, between about 2 and 4 ethylene glycol units.

[0122] In another embodiment, the present disclosure is directed to a compound which comprises a ABM group as described abovewhich binds to a target protein (e.g., Androgen Receptor) or polypeptide, which is ubiquitinated by an ubiquitin ligase and is chemically linked directly to the ULM group or through a linker moiety L, or ABM is alternatively a ULM' group which is also an ubiquitin ligase binding moiety, which may be the same or different than the ULM group as described above and is linked directly to the ULM group directly or through the linker moiety; and L is a linker moiety as described above which may be present or absent and which chemically (covalently) links ULM to ABM, or a pharmaceutically acceptable salt, enantiomer, stereoisomer, solvate or polymorph thereof.

[0123] In certain embodiments, the ULM shows activity or binds to an E3 ubiquitin ligase with an IC_{50} of less than about 200 μM. The IC_{50} can be determined according to any method known in the art, e.g., a fluorescent polarization assay.

[0124] In certain additional embodiments, the bifunctional compounds described herein demonstrate an activity with an IC_{50} of less than about 100, 50, 10, 1, 0.5, 0.1, 0.05, 0.01, 0.005, 0.001 mM, or less than about 100, 50, 10, 1, 0.5, 0.1, 0.05, 0.01, 0.005, 0.001 μM, or less than about 100, 50, 10, 1, 0.5, 0.1, 0.05, 0.01, 0.005, 0.001 nM.


[0125] Although the ULM group and ABM group may be covalently linked to the linker group through any group which is appropriate and stable to the chemistry of the linker, in preferred aspects of the present disclosure, the linker is independently covalently bonded to the ULM group and the ABM group preferably through an amide, ester, thioester, keto group, carbamate (urethane), carbon or ether, each of which groups may be inserted anywhere on the ULM group and ABM group to provide maximum binding of the ULM group on the ubiquitin ligase and the ABM group on the target protein to be degraded. In certain preferred aspects, the linker may be linked to an optionally substituted alkyl, alkylene, alkene or alkyne group, an aryl group or a heterocyclic group on the ULM and/or ABM groups.

Exemplary Androgen Binding Moieties (ABMs)


[0126] In another aspect, the description provides AR binding moieties (ABM), which in certain aspects and embodiments are coupled to a linker and/or a ULM as described herein.

[0127] In any of the compounds described herein, the ABM comprises a chemical moiety that binds to the androgen receptor (AR). Various androgen receptor binding compounds have been described in literature, including various androgen derivatives such as testosterone, dihydrotestosterone, and metribolone (also known as methyltrienolone or R1881), and non-steroidal compounds such as bicalutamide, enzalutamide. Those of ordinary skill in the art would appreciate that these androgen receptor binding compounds could be potentially used as an ABM moiety in a PROTAC compound. Such literature includes, but not limited to, G. F. Allan et. al, *Nuclear Receptor Signaling*, 2003, 1, e009; R. H. Bradbury et. al, *Bioorganic & Medicinal Chemistry Letters*, 2011 5442–5445; C. Guo et. al, *Bioorganic & Medicinal Chemistry Letters*, 2012 2572-2578; P. K. Poutiainen et. al, *J. Med. Chem.* 2012, 55, 6316 – 6327 A. Pepe et. al, *J. Med. Chem.* 2013, 56, 8280 – 8297; M. E. Jung et al, *J. Med. Chem.* 2010, 53, 2779–2796, which are incorporated by reference herein


[0128] In certain embodiments, the ABM comprises a structure selected from, but not limited to the structures shown below, where a dashed line indicates the attachment point of a linker moiety or a ULM:




ABM-a ;



ABM-b ;



ABM-c ; and



ABM-d ,

wherein:

W^1 is aryl, heteroaryl, bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6}

alkoxyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, S;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;

Q is a 3-6 membered ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q,is independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);



R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, cyclic, heterocyclic , or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, aryl, heteroaryl, alicyclic, heterocyclic, biheterocyclic, biaryl, or biheteroaryl, each optionally substituted by 1-10 R^{W2};

each R^{W2} is independently H, halo, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more F), $-OR^{W2A}$, C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OC_{1-3}alkyl (optionally substituted), OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and

R^{W2A} is H, C_{1-6} alkyl (linear, branched), or C_{1-6} heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC_{1-3}alkyl.


[0129] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.

[0130] In certain embodiments, W^1 is  or  ;


wherein each R_{22} is independently halo, H, optionally substituted alkyl, haloalkyl, cyano, or nitro; and

each R_{23} is independently H, halo, CF_3, optionally substituted alkyl, alkoxy,haloalkyl, cyano, or nitro.


[0131] In certain additional embodiments, W^1 is selected from the group consisting of:




[0132] In particular embodiments, the ABM comprises a structure selected from the following structures shown below, where a  indicates tha attachment point of a linker or a ULM:



,



, and



,

wherein:

R^{Q2} is a H, halogen, CH_3 or CF_3;

R^{Q3} is H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;


Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, heteroaryl, or aryl;

R^{Y1}, R^{Y2} are each independently H, or C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl, cyclic, or heterocyclic); and


R^Q each independently is H, C_1-C_6 alkyl (linear, branched, optionally substituted by 1 or more halo, or C_{1-6} alkoxyl), or two R^Q together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms.

[0133] In a particular embodiment, each R^Q is independently H or CH_3. In another embodiment R^{Q3} is CN.


[0134] In particular embodiments, the ABM comprises a structure selected from the following structures shown below, where a  indicates the attachment point of a linker or a ULM:




,



,



, and



,

wherein:

R^{Q2} is a H, halogen, CN, CH_3 or CF_3; and

R^{Q3} is H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3; .


Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, heteroaryl, or aryl; and

R^{Y1}, R^{Y2} are each independently H or C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl, cyclic, or heterocyclic); and

X is N or C.



[0135] In a particular embodiment R^{Q3} is a CN.

[0136] In certain additional embodiments, the ABM comprises a structure shown below, where a dashed line indicates the attachment point of a linker moiety or a ULM or a CLM:



,

wherein:

W^1 is  or  ;

each R_{22} is independently H or -CN;

each R_{23} is independently H, halo, C_1-C_6 alkyl (linear, branched, optionally substituted), C_1-C_6 alkoxy, or $-CF_3$;

Y^3 is a bond or O;

Y^4 is a bond or NH;

Y^5 is a bond, C=O, C_1-C_6 heteroaryl, or C_1-C_6 aryl;

R^1, R^2, are each independently H, or C_1-C_6 alkyl (linear or branched, optionally substituted; for example, optionally substituted by 1 or more halo, or C_{1-6} alkoxyl);


W^2 is a bond, C_{1-6} aryl, C_{1-6} heteroaryl, C_{1-6} alicyclic, or C_{1-6} heterocyclic, biheterocyclic, biaryl, or biheteroaryl, each optionally substituted by 1-10 R^{W2}; and

each R^{W2} is independently H, or halo; and


[0137] ~~~ represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific.

[0138] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.


[0139] In certain additional embodiments, W^1 is selected from the group consisting of:





[0140] In any aspect or embodiment described herein, W^2 is selected from the group consisting of:



[0141] In certain embodiments, the ABM comprises a structure selected from, but not limited to the structures shown below, where a dashed line indicates the attachment point of a linker moiety or a ULM:



ABM-a ,

wherein W^1 is  or  ;

each R_{22} is independently H or -CN;

each R_{23} is independently H, halo, or $–CF_3$;

Y^1, Y^2 are each independently O or S;






R^1, R^2, are each independently H or a methyl group;

W^2 is a bond, C_{1-6} aryl, or heteroaryl, each optionally substituted by 1, 2 or 3 R^{W2}; and



each R^{W2} is independently H, halo, C_{1-6} alkyl (optionally substituted by 1 or more F), OC_{1-3}alkyl (optionally substituted by 1 or more –F).

[0142] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.


[0143] In certain additional embodiments, W^1 is selected from the group consisting of:



















 ;  ;  ;  ; and  .

[0144] In certain additional embodiments, W2 is selected from the group consisting of:

 and  .

[0145] In certain embodiments, ABM is selected from the group consisting of:

















;

;

;

;

;

;

;

;

;

;

;

;

;

;

;

;

;

;



;


;


.


60


;;

;



;


O
NH
O
N


;



;



[0146] In certain embodiments, the ABM comprises the structure:



ABM-b

wherein W^1 is aryl, or heteroaryl, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;

Q is a 4 membered alicyclic ring with 0-2 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q is independently H, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more

halo, C_{1-6} alkoxyl), or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);


R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, C_{1-6} alicyclic, heterocyclic, aryl, biheterocyclic, biaryl, or biheteroaryl, or heteroaryl, each optionally substituted by 1, 2 or 3 R^{W2}; and

each R^{W2} is independently H, halo, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more F), C_{1-6} heteroalkyl (linear, branched, optionally substituted), -OR^{W2A} OC_{1-3}alkyl (optionally substituted by 1 or more –F), C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl (optionally substituted), C_{1-6} alkyl (optionally substituted), C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), heteroaryl (optionally substituted), bicyclic hereoaryl (optionally substituted), bicyclic aryl, OH, NH_2, $NR^{Y1}R^{Y2}$, or CN; and

R^{W2A} is H, C_{1-6} alkyl (linear, branched), or C_{1-6} heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC_{1-3}alkyl.

[0147] In an additional aspect, the description provides an androgen receptor bindingcompound comprising a structure of:



ABM-e

wherein W^1 is aryl, heteroaryl, , bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, or S;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;

Q is a 3-6 membered alicyclic or aromatic ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q,is independently H, C_{1-6} alkyl (linear, branched, optionally

substituted by 1 or more halo, C_{1-6} alkoxyl), or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);


R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, C_{1-6} alicyclic, heterocyclic, aryl, biheterocyclic, biaryl, or biheteroaryl, or heteroaryl, each optionally substituted by 1, 2 or 3 R^{W2};

each R^{W2} is independently H, halo, C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more F), C_{1-6} heteroalkyl (linear, branched, optionally substituted), -OR^{W2A}, OC_{1-3}alkyl (optionally substituted by 1 or more –F), C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, C_{1-6} alkyl (optionally substituted), C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and


R^{W2A} is H, C_{1-6} alkyl (linear, branched), or C_{1-6} heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC_{1-3}alkyl.


[0148] In certain embodiments, an androgen receptor binding moiety has a structure of:



ABM-8

,

wherein W^1 is 

or 

;

each R_{22} is independently H or -CN;

each R_{23} is independently H, halo, or $–CF_3$;

Y^3 is a bond or O;


Q is a 4 member ring, optionally substituted with 0-4 R^Q, each R^Q is independently H or methyl;

Y4 is a bond or NH;

Y5 is a bond, a C=O, or a C=S;


each W^2 is independently a bond, C1-6 aryl or heteroaryl, each optionally substituted by 1, 2 or 3 R^{W2}, each R^{W2} is independently H, halo, a 6 member alicyclic ring with 1 or 2 heteroatoms or a 5 member aromatic ring with 1 or 2 or 3 heteroatoms.

[0149] In certain additional embodiments, W^2 is selected from the group consisting of:




[0150] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.

[0151] In certain additional embodiments, W^1 is selected from the group consisting of:



[0152] In certain embodiments, an androgen binding moiety has a structure of:



ABM-d

wherein W^1 is aryl, independently substituted by 1 or more halo, CN;

Y^3 are each independently a bond, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O;


Q is a 5 membered aromatic ring with 1 or 2 heteroatoms;

R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched);

W^2 is a bond, aryl, or heteroaryl, each optionally substituted by 1, 2 or 3 R^{W2}; and

each R^{W2} is independently H, halo, C_{1-6} alkyl (optionally substituted by 1 or more F), OC_{1-3}alkyl (optionally substituted by 1 or more –F).





[0153] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.

[0154] In certain embodiments, W^1 is  ;

wherein each R_{22} is independently halo or CN; and


each R_{23} is independently H or halo.

[0155] In certain additional embodiments, W^1 is selected from the group consisting of:


 ;  ;  ; and  .

[0156] In certain additional embodiments, Q is  .

[0157] In certain additional embodiments, W^2 is  .


[0158] In certain additional embodiments, $(Y^3)_{0-5}$ is  .


[0159] In certain embodiments, the ABM comprises a structure selected from, but not limited to the structures shown below, where a dashed line indicates the attachment point of a linker moiety or a ULM:





ABM-a

,





wherein:

W^1 is  or  ;

each R_{22} is independently H or -CN;

each R_{23} is independently H, halo, or $–CF_3$;

Y^1, Y^2 are each independently O or S;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, or SO_2;;


R^1, R^2, are each independently H or a methyl group;

W^2 is a bond, C_{1-6} aryl, or heteroaryl, each optionally substituted by 1, 2 or 3 R^{W2}; and

each R^{W2} is independently H, halo, C_{1-6} alkyl (optionally substituted by 1 or more F), C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, OC_{1-3}alkyl (optionally substituted by 1 or more –F).


[0160] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.

[0161] In certain additional embodiments, W^1 is selected from the group consisting of:




and

[0162] In certain additional embodiments, W2 is selected from the group consisting of:




.

[0163] In certain embodiments, the ABM comprises a structure shown below, where a dashed line indicates the attachment point of a linker moiety or a ULM or a CLM:



,

wherein:

W^1 is  ;

each R_{22} is independently H or -CN;

each R_{23} is independently H, halo, or $–CF_3$;

Y^3 is a bond or O;

Y^4 is a bond or NH;

Y^5 is a bond, C=O, C_1-C_6 heteroaryl, or C_1-C_6 aryl;

R^1, R^2, are each independently H, or C_1-C_6 alkyl (linear or branched, optionally substituted by 1 or more halo, or C_{1-6} alkoxyl);


W^2 is a bond, C_{1-6} aryl, C_{1-6} heteroaryl, C_{1-6} alicyclic, or C_{1-6} heterocyclic, each optionally substituted by 1-10 R^{W2}; and

each R^{W2} is independently H, or halo; and


[0164] ~~~ represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific.








[0165] In any of the embodiments described herein, the W^2 is covalently coupled to one or more ULM or CLM groups, or a linker to which is attached one or more ULM or CLM groups as described herein.

[0166] In certain additional embodiments, W^1 is selected from the group consisting of:



[0167] In certain additional embodiments, W^2 is selected from the group consisting of:




;

;

;

;

;

; and

.

[0168] In certain embodiments, the androgen receptor binding compound of ABM is selected from the group consisting of:

trans-2-Chloro-4-[3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile;

cis-2-Chloro-4-[3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile;

trans 6-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridazine-3-carboxamide;

trans tert-Butyl N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamate;

trans 4-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide;

trans 5-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrazine-2-carboxamide;

trans 2-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide;

4-Methoxy-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide;

trans 1-(2-Hydroxyethyl)-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]-1H-pyrazole-4-carboxamide;

trans 6-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide;

trans 4-[(5-Hydroxypentyl)amino]-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide; and

trans tert-Butyl 2-({5-[(4-{[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoyl}phenyl)aminopentyl}oxy)acetate; and

N-((1r,3r)-3-(4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-methylbenzamide.

[0169] The term "hydrocarbyl" shall mean a compound which contains carbon and hydrogen and which may be fully saturated, partially unsaturated or aromatic and includes aryl groups, alkyl groups, alkenyl groups and alkynyl groups.

[0170] The term "unsubstituted" shall mean substituted only with hydrogen atoms. A range of carbon atoms which includes C_0 means that carbon is absent and is replaced with H. Thus, a range of carbon atoms which is C_0-C_6 includes carbons atoms of 1, 2, 3, 4, 5 and 6 and for C_0, H stands in place of carbon. The term "substituted" or "optionally substituted" shall mean independently (i.e., where more than substituent occurs, each substituent is independent of another substituent) one or more substituents (independently up to five substitutents, preferably up to three substituents, often 1 or 2 substituents on a moiety in a compound according to the present disclosure and may include substituents which themselves may be further substituted) at a carbon (or nitrogen) position anywhere on a molecule within context, and includes as substituents hydroxyl, thiol, carboxyl, cyano (C≡N), nitro (NO_2), halogen (preferably, 1, 2 or 3 halogens, especially on an alkyl, especially a methyl group such as a trifluoromethyl), an alkyl group (preferably, C_1-C_{10}, more preferably, C_1-C_6), aryl (especially phenyl and substituted phenyl for example benzyl or benzoyl), alkoxy group (preferably, C_1-C_6 alkyl or aryl, including phenyl and substituted phenyl), thioether (C_1-C_6 alkyl or aryl), acyl (preferably, C_1-C_6 acyl), ester or thioester (preferably, C_1-C_6 alkyl or aryl) including alkylene ester (such that attachment is on the alkylene group, rather than at the ester function which is preferably substituted with a C_1-C_6 alkyl or aryl group), preferably, C_1-C_6 alkyl or aryl, halogen (preferably, F or Cl), amine (including a five- or six-membered cyclic alkylene amine, further including a C_1-C_6 alkyl amine or a C_1-C_6 dialkyl amine which alkyl groups may be substituted with one or two hydroxyl groups) or an optionally substituted –N(C_0-C_6 alkyl)C(O)(O-C_1-C_6 alkyl) group (which may be optionally substituted with a polyethylene glycol chain to which is further bound an alkyl group containing a single halogen, preferably chlorine substituent), hydrazine, amido, which is preferably substituted with one or two C_1-C_6 alkyl groups (including a carboxamide which is optionally substituted with one or two C_1-C_6 alkyl groups), alkanol (preferably, C_1-C_6 alkyl or aryl), or alkanoic acid (preferably, C_1-C_6 alkyl or aryl). Substituents according to the present disclosure may include, for example – $SiR_1R_2R_3$ groups where each of R_1 and R_2 is as otherwise described herein and R_3 is H or a C_1-C_6 alkyl group, preferably R_1, R_2, R_3 in this context is a C_1-C_3 alkyl group (including an isopropyl or t-butyl group). Each of the above-described groups may be linked directly to the substituted

moiety or alternatively, the substituent may be linked to the substituted moiety (preferably in the case of an aryl or heteraryl moiety) through an optionally substituted $-(CH_2)_m-$ or alternatively an optionally substituted $-(OCH_2)_m-$, $-(OCH_2CH_2)_m-$ or $-(CH_2CH_2O)_m-$ group, which may be substituted with any one or more of the above-described substituents. Alkylene groups $-(CH_2)_m-$ or $-(CH_2)_n-$ groups or other chains such as ethylene glycol chains, as identified above, may be substituted anywhere on the chain. Preferred substituents on alkylene groups include halogen or C_1-C_6 (preferably C_1-C_3) alkyl groups, which may be optionally substituted with one or two hydroxyl groups, one or two ether groups (O-C_1-C_6 groups), up to three halo groups (preferably F), or a sidechain of an amino acid as otherwise described herein and optionally substituted amide (preferably carboxamide substituted as described above) or urethane groups (often with one or two C_0-C_6 alkyl substituents, which group(s) may be further substituted). In certain embodiments, the alkylene group (often a single methylene group) is substituted with one or two optionally substituted C_1-C_6 alkyl groups, preferably C_1-C_4 alkyl group, most often methyl or O-methyl groups or a sidechain of an amino acid as otherwise described herein. In the present disclosure, a moiety in a molecule may be optionally substituted with up to five substituents, preferably up to three substituents. Most often, in the present disclosure moieties that are substituted, are substituted with one or two substituents.

[0171] The term "substituted" (each substituent being independent of any other substituent) shall also mean within its context of use C_1-C_6 alkyl, C_1-C_6 alkoxy, halogen, amido, carboxamido, sulfone, including sulfonamide, keto, carboxy, C_1-C_6 ester (oxyester or carbonylester), C_1-C_6 keto, urethane -O-C(O)-NR_1R_2 or $-N(R_1)$-C(O)-O-R_1, nitro, cyano and amine (especially including a C_1-C_6 alkylene-NR_1R_2, a mono- or di- C_1-C_6 alkyl substituted amines which may be optionally substituted with one or two hydroxyl groups). Each of these groups contain unless otherwise indicated, within context, between 1 and 6 carbon atoms. In certain embodiments, preferred substituents will include for example, –NH-, -NHC(O)-, -O-, =O, $-(CH_2)_m-$ (here, m and n are in context, 1, 2, 3, 4, 5 or 6), -S-, -S(O)-, SO_2- or –NH-C(O)-NH-, $-(CH_2)_nOH$, $-(CH_2)_nSH$, $-(CH_2)_nCOOH$, C_1-C_6 alkyl, $-(CH_2)_nO-(C_1$-C_6 alkyl), $-(CH_2)_nC(O)-(C_1$-C_6 alkyl), $-(CH_2)_nOC(O)-(C_1$-C_6 alkyl), $-(CH_2)_nC(O)O-(C_1$-C_6 alkyl), $-(CH_2)_nNHC(O)-R_1$, $-(CH_2)_nC(O)-NR_1R_2$, $-(OCH_2)_nOH$, $-(CH_2O)_nCOOH$, C_1-C_6 alkyl, $-(OCH_2)_nO-(C_1$-C_6 alkyl), $-(CH_2O)_nC(O)-(C_1$-C_6 alkyl), $-(OCH_2)_nNHC(O)-R_1$, $-(CH_2O)_nC(O)-NR_1R_2$, $-S(O)_2-R_S$, $-S(O)-R_S$ (R_S is C_1-C_6 alkyl or a $-(CH_2)_m-NR_1R_2$ group), NO_2, CN or halogen (F, Cl, Br, I, preferably F or

Cl), depending on the context of the use of the substituent. R_1 and R_2 are each, within context, H or a C_1-C_6 alkyl group (which may be optionally substituted with one or two hydroxyl groups or up to three halogen groups, preferably fluorine). The term "substituted" shall also mean, within the chemical context of the compound defined and substituent used, an optionally substituted aryl or heteroaryl group or an optionally substituted heterocyclic group as otherwise described herein. Alkylene groups may also be substituted as otherwise disclosed herein, preferably with optionally substituted C_1-C_6 alkyl groups (methyl, ethyl or hydroxymethyl or hydroxyethyl is preferred, thus providing a chiral center), a sidechain of an amino acid group as otherwise described herein, an amido group as described hereinabove, or a urethane group O-C(O)-NR_1R_2 group where R_1 and R_2 are as otherwise described herein, although numerous other groups may also be used as substituents. Various optionally substituted moieties may be substituted with 3 or more substituents, preferably no more than 3 substituents and preferably with 1 or 2 substituents. It is noted that in instances where, in a compound at a particular position of the molecule substitution is required (principally, because of valency), but no substitution is indicated, then that substituent is construed or understood to be H, unless the context of the substitution suggests otherwise.

[0172] **Exemplary AR-PROTAC Compounds**

[0173] As described above, in certain aspects, the description provides bifuctional PROTAC compounds comprising at least one ABM group, a linker, and at least one ULM (or CLM) group as described herein.

[0174] In certain embodiments, the compound is selected from the group consisting of compounds 1-625 (i.e., the chemical structures described in Table 2, Table 3, Table 4, Table 5, Table 6, and Table 7 of Figures 2, 3, 4, 5, 6, and 7, respectively), and salts and polymorphs thereof.

[0175] In additional embodiments, the compound is selected from the group consisting of: 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azadodecan-12-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (1); 4-(3-{4-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)ethoxy]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (2); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10,13,16-pentaoxa-1-azaoctadecan-18-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-

(trifluoromethyl)benzonitrile (3); 4-[3-(4-{2-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)ethoxy]ethoxy}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (4); 4-[3-(4-{3-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)ethoxy]propoxy}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (5); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azatetradecan-14-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (6); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azatridecan-13-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (7); 4-(3-{4-[(1-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}-4,7,10-trioxa-1-azadodecan-12-yl)oxy]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (8); 4-(3-{4-[(1-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}-4,7,10-trioxa-1-azadodecan-12-yl)oxy]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (9); 4-[3-(4-{3-[3-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)propoxy]propoxy}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (10); 4-{4,4-dimethyl-3-[4-({1-[2-(3-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azatridecan-13-yl}oxy)phenyl]-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (11); 4-[3-(4-{4-[(5-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}pentyl)oxy]phenyl}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (12); 4-{[5-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)pentyl]oxy}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (13); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-1,4,7,10-tetraoxatridecan-13-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (14); 6-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (15); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-

dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (16); 4-(3-{4-[1-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethyl)-1H-1,3-benzodiazol-5-yl]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (17); 4-{[5-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)pentyl]amino}-3-fluoro-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (18); 6-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (19); 6-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (20); 6-(4-{3-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]propyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (21); 6-{4-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (22); 4-{4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethyl)piperazin-1-yl]butyl}-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (23); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (24); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (25); 4-(6-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}hexyl)-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (26); 4-{3-[4-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propyl)piperazin-1-yl]propyl}-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (27); 4-[5-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)pentyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (28); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-

1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (29); 4-(4-{4-[2-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}oxy)ethyl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (30); 4-(4-{4-[2-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-4-yl}oxy)ethyl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (31); 4-{4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]butyl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (32); 6-{4-[5-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-4-yl}oxy)pentyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (33); 4-(4-{4-[2-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl}oxy)ethyl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (34); 6-{4-[5-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}oxy)pentyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (35); 4-(4-{4-[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (36); 4-{4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]butyl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (37); 6-(4-{6-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]hexyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (38); 4-(4-{4-[2-(2,6-dioxopiperidin-3-yl)-4-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (39); 4-(6-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (40); 6-{4-[5-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl}oxy)pentyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (41); 6-[4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-

yl]oxy}pentyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]pyridine-3-carboxamide (42); 4-[4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (43); 4-(5-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}pentyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (44); 4-(4-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (45); 4-(6-{4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (46); 4-(4-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (47); 4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (48); 4-[3-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)propyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (49); 4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)butyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (50); 4-[5-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)pentyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (51); 4-[6-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)hexyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (52); N-[(2R)-1-[3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl]propan-2-yl]-5-{[4-(4-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}butyl)piperazin-1-yl]methyl}-1H-pyrazole-3-carboxamide (53); 4-(4-{6-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]hexyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (54); 4-(3-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}propyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (55); N-[(2R)-1-[3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl]propan-2-yl]-5-[(3-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}propoxy)methyl]-1H-pyrazole-3-carboxamide (56); 4-[4-(3-{[2-(2,6-

dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}propyl)-1H-1,2,3-triazol-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (57); 6-[4-(6-{[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]amino}hexyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (58); N-[(2R)-1-[3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl]propan-2-yl]-5-({3-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]propoxy}methyl)-1H-pyrazole-3-carboxamide (59); (60); 4-[4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)-1H-1,2,3-triazol-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (61); 6-{4-[2-(4-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}butoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (62); 6-{4-[2-(4-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}butoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (63); 6-(4-{2-[(5-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (64); 6-(4-{2-[(5-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (65); 4-(4-{2-[(5-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (66); 4-(4-{2-[(5-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (67); 4-(4-{6-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-5-yl]hexyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (68); 4-(3-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}propyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (69); 4-[6-(4-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (70); 4-[6-(4-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-

yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (71); 6-[4-(6-{[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}hexyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (72); 6-[4-(6-{[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy}hexyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (73); 4-[6-(4-{2-[(3S)-2,6-dioxopiperidin-3-yl]-3-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (74); and 4-[6-(4-{2-[(3R)-2,6-dioxopiperidin-3-yl]-3-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (75); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(6-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)amino)hexyl)piperazin-1-yl)nicotinamide (76); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperazin-1-yl)nicotinamide (77); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)ethyl)piperazin-1-yl)nicotinamide (78); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-((5-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)pentyl)oxy)ethyl)piperazin-1-yl)nicotinamide (79); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-((5-(2-((R)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)pentyl)oxy)ethyl)piperazin-1-yl)benzamide (80); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)hexyl)benzamide (81); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)hexyl)benzamide (82); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((1R,4R)-5-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)nicotinamide (83); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)benzamide (84); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-1,2,3,6-tetrahydropyridin-4-yl)benzamide (85); N-

((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(5-(4-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)pentyl)benzamide (86); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(5-(4-(2-((R)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)pentyl)benzamide (87); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)benzamide (88); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)benzamide (89); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(6-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)oxy)hexyl)piperazin-1-yl)nicotinamide (90); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)benzamide (91); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)benzamide (92); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperidin-1-yl)hexyl)benzamide (93); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperidin-1-yl)hexyl)nicotinamide (94); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)pentyl)benzamide (95); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-3-oxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (96); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)butyl)benzamide (97); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)pentyl)benzamide (98); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)benzamide (99); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)pentyl)-1H-1,2,3-triazol-1-yl)benzamide (100); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-((1S,4S)-5-(2-((R)-2,6-dioxopiperidin-3-yl)-

1-oxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)hexyl)benzamide (101); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)pentyl)-1H-1,2,3-triazol-1-yl)nicotinamide (102); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-1-yl)hexyl)nicotinamide (103); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)butyl)benzamide (104); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-(4-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)piperidin-1-yl)propyl)-1H-1,2,3-triazol-1-yl)benzamide (105); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-(3-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)azetidin-1-yl)propyl)-1H-1,2,3-triazol-1-yl)benzamide (106); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-1H-1,2,3-triazol-4-yl)ethyl)piperazin-1-yl)nicotinamide (107); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(3-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-1H-1,2,3-triazol-4-yl)propyl)piperazin-1-yl)nicotinamide (108); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (109); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (110); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)nicotinamide (111); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)nicotinamide (112); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(1-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperidin-4-yl)nicotinamide (113); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (114); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)nicotinamide (115); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S)-4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-

dioxoisoindolin-5-yl)oxy)pentyl)-3-(hydroxymethyl)piperazin-1-yl)benzamide (116); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(3-(3-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)azetidin-1-yl)propyl)-1H-1,2,3-triazol-1-yl)nicotinamide (117); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (118); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)propyl)nicotinamide (119); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)nicotinamide (120); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperidin-1-yl)butyl)nicotinamide (121); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)nicotinamide (122); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-1-yl)butyl)nicotinamide (123); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)butyl)nicotinamide (124); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)butyl)nicotinamide (125); N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (126); N-((1r,3r)-3-(3,4-dicyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (127); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(5-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)pentyl)nicotinamide (128); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide (129); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)piperidin-1-yl)butyl)nicotinamide (130); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)nicotinamide (131); N-((1r,3r)-3-(3-

chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)pyrimidine-5-carboxamide (132); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propoxy)ethyl)-3,5-difluorobenzamide (133); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S,5R)-4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-3,5-dimethylpiperazin-1-yl)benzamide (134); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(2-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl)nicotinamide (135); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(1-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)-4,6-dihydropyrrolo[3,4-c]pyrazol-5(1H)-yl)nicotinamide (136); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(5-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)-5,6-dihydropyrrolo[3,4-c]pyrazol-2(4H)-yl)nicotinamide (137); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)nicotinamide (138); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-((S)-2,6-dioxopiperidin-3-yl)-6-fluoro-3-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (139); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-((S)-2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (140); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(9-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-1-oxa-4,9-diazaspiro[5.5]undecan-4-yl)ethyl)benzamide (141); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)pyrimidine-5-carboxamide (142); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrimidine-5-carboxamide (143); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxamide (144); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (145); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-(4-(2-

(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)nicotinamide (146); N-((R)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-((4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)butoxy)methyl)-1H-pyrazole-3-carboxamide (147); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(9-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-1-oxa-4,9-diazaspiro[5.5]undecan-4-yl)propyl)benzamide (148); N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide (149); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrimidine-5-carboxamide (150); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-((S)-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (151); N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)nicotinamide (152); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)nicotinamide (153); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-3-oxopiperazin-1-yl)nicotinamide (154); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)butyl)piperazin-1-yl)pyrimidine-5-carboxamide (155); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(3-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)propyl)piperazin-1-yl)pyrimidine-5-carboxamide (156); 2-chloro-4-(3-(4-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile (157); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (158); 2-chloro-4-(5-(4-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)benzonitrile (159); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)nicotinamide (160); N-((1r,3r)-3-(3,4-

dicyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrimidine-5-carboxamide (161); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(2-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,7-diazaspiro[3.5]nonan-7-yl)nicotinamide (162); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(7-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,7-diazaspiro[3.5]nonan-2-yl)nicotinamide (163); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(2-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-2,6-dihydropyrrolo[3,4-c]pyrazol-5(4H)-yl)nicotinamide (164); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(3-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)azetidin-1-yl)pyrimidine-5-carboxamide (165); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)-2,6-difluorobenzamide (166); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)-1,4-diazepan-1-yl)pyrimidine-5-carboxamide (167); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)-1-oxa-4,9-diazaspiro[5.5]undecan-9-yl)benzamide (168); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-1-oxa-4,9-diazaspiro[5.5]undecan-9-yl)benzamide (169); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)benzamide (170); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-((1R,4R)-5-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethyl)benzamide (171); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazine-1-carbonyl)piperidin-1-yl)benzamide (172); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-3,5-difluorobenzamide (173); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)benzamide (174); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((1R,4R)-5-(2-(2,6-dioxopiperidin-

3-yl)-1,3-dioxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)piperidin-1-yl)benzamide (175); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((1S,4S)-5-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)methyl)piperidin-1-yl)benzamide (176); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-3-fluorobenzamide (177); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(methyl)amino)methyl)piperidin-1-yl)benzamide (178); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-5-fluoronicotinamide (179); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (180); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4,6-difluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (181); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)benzamide (182); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)cyclohexyl)benzamide (183); N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)cyclohexyl)benzamide (184); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (185); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (186); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)benzamide (187); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-oxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)benzamide (188); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-

dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)azetidin-1-yl)benzamide (189); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)ethyl)piperidin-1-yl)benzamide (190); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azepan-1-yl)pyrimidine-5-carboxamide (191); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)cyclohexyl)benzamide (192); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((3S,5R)-4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-3,5-dimethylpiperazin-1-yl)nicotinamide (193); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S,5R)-4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-3,5-dimethylpiperazin-1-yl)benzamide (194); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((3S,5R)-4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-3,5-dimethylpiperazin-1-yl)pyrimidine-5-carboxamide (195); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1s,3S)-3-(((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)methyl)cyclobutyl)piperazin-1-yl)benzamide (196); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)-2-fluorobenzamide (197); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)azetidin-1-yl)piperidin-1-yl)benzamide (198); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-(((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)(methyl)amino)pyrrolidin-1-yl)benzamide (199); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S)-3-(((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)(methyl)amino)pyrrolidin-1-yl)benzamide (200); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-(((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)(methyl)amino)methyl)pyrrolidin-1-yl)benzamide (201); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-((2S,6R)-4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,6-dimethylpiperazin-1-yl)ethyl)benzamide (202); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-

yl)-[1,3'-biazetidin]-1'-yl)benzamide (203); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(methyl)amino)ethyl)piperazin-1-yl)benzamide (204); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(1-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)azetidin-3-yl)piperazin-1-yl)benzamide (205); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)azetidin-3-yl)piperazin-1-yl)ethyl)benzamide (206); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-((2R,6R)-4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,6-dimethylpiperazin-1-yl)ethyl)benzamide (207); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)benzamide (208); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-((2S,6S)-4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,6-dimethylpiperazin-1-yl)ethyl)benzamide (209); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)benzamide (210); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)pyrrolidin-1-yl)benzamide (211); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(4-(6-(2,6-dioxopiperidin-3-yl)-5,7-dioxo-6,7-dihydro-5H-pyrrolo[3,4-b]pyrazin-2-yl)piperazin-1-yl)ethyl)piperidin-1-yl)benzamide (212); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (213); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (214); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((4aR,6R,8aS)-6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)octahydroisoquinolin-2(1H)-yl)nicotinamide (215); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(2-(((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(methyl)amino)ethyl)azetidin-1-yl)benzamide (216); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-4-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)benzamide (217); N-((1r,3R)-3-(3-chloro-4-

cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((1r,3R)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)cyclobutyl)benzamide (218); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((1s,3S)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)cyclobutyl)benzamide (219); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((4aR,6S,8aR)-6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)octahydroisoquinolin-2(1H)-yl)nicotinamide (220); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((4aR,6R,8aR)-6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)octahydroisoquinolin-2(1H)-yl)nicotinamide (221); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)pyridazine-3-carboxamide (222); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pyrrolidin-1-yl)benzamide (223); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)azetidin-3-yl)piperazin-1-yl)benzamide (224); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazine-1-carbonyl)pyrrolidin-1-yl)benzamide (225); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazine-1-carbonyl)pyrrolidin-1-yl)benzamide (226); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)benzamide (227); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((3R)-1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methyl)piperazin-1-yl)benzamide (228); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)-2,6-diazaspiro[3.3]heptan-2-yl)benzamide (229); (3R)-N-(1-(4-(((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)-1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-N-methylpyrrolidine-3-carboxamide (230); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyridazine-3-carboxamide (231); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-

tetramethylcyclobutyl)-2-((2S,4R,6R)-4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2,6-dimethylpiperidin-1-yl)pyrimidine-5-carboxamide (232); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((2S,4S,6R)-4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2,6-dimethylpiperidin-1-yl)pyrimidine-5-carboxamide (233); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((2S,6S)-4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2,6-dimethylpiperidin-1-yl)pyrimidine-5-carboxamide (234); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(3-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)propyl)azetidin-1-yl)benzamide (235); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(7-((3-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-1-yl)methyl)octahydro-2H-pyrido[1,2-a]pyrazin-2-yl)nicotinamide (236); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)methyl)(methyl)amino)piperidin-1-yl)benzamide (237); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((1'-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-[1,4'-bipiperidin]-4-yl)amino)benzamide (238); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperazin-1-yl)benzamide (239); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)benzamide (240); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((1S,4S,5R)-5-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidine-5-carboxamide (241); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((1S,4S,5S)-5-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidine-5-carboxamide (242); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (243); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-((2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)(methyl)amino)pyrrolidin-1-yl)benzamide (244); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-(3-((2-(2,6-dioxopiperidin-3-

yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)pyrrolidin-1-yl)benzamide (245); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(5-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)benzamide (246); N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (247); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)butyl)benzamide (248); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-((R)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)butyl)benzamide (249); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)benzamide (250); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(4-(2-((R)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)benzamide (251); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)benzamide (252); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-((2S,6R)-4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,6-dimethylpiperazin-1-yl)hexyl)benzamide (253); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)ethyl)oxetan-3-yl)methyl)benzamide (254); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)benzamide (255); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-((1S,4S)-5-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)hexyl)benzamide (256); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)benzamide (257); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-((1R,4R)-5-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)hexyl)benzamide (258); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-((1R,4R)-5-(2-((R)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)hexyl)benzamide (259); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-

1-oxoisoindolin-5-yl)piperazin-1-yl)ethyl)oxetan-3-yl)methyl)benzamide (260); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(1-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperidin-4-yl)nicotinamide (261); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(1-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperidin-4-yl)nicotinamide (262); N-((1r,3r)-3-((5-cyano-6-methylpyridin-2-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (263); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)pyrazine-2-carboxamide (264); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)pyridazine-3-carboxamide (265); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)picolinamide (266); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)-1H-pyrazole-4-carboxamide (267); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)-1H-pyrazole-3-carboxamide (268); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)pyrrolidin-3-yl)-1H-pyrazole-4-carboxamide (269); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)benzamide (270); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-((R)-2,6-dioxopiperidin-3-yl)-6-fluoro-3-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (271); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-((R)-2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (272); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)benzamide (273); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-3,6-dihydropyridin-1(2H)-yl)butyl)nicotinamide (274); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1'-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-

yl)oxy)pentyl)-1',2',3',6'-tetrahydro-[3,4'-bipyridine]-6-carboxamide (275); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(5-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-5,6-dihydropyrrolo[3,4-c]pyrazol-2(4H)-yl)nicotinamide (276); N-((S)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-((3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propoxy)methyl)-1H-pyrazole-3-carboxamide (277); N-((R)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-((4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)methyl)-1H-pyrazole-3-carboxamide (278); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((2R)-4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-2-(hydroxymethyl)piperazin-1-yl)nicotinamide (279); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1'-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-1',2',3',6'-tetrahydro-[2,4'-bipyridine]-5-carboxamide (280); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-2-oxopiperazin-1-yl)nicotinamide (281); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(1-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperidin-4-yl)benzamide (282); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(1-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperidin-4-yl)benzamide (283); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)-3-oxopropyl)benzamide (284); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-((R)-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (285); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)azetidin-1-yl)nicotinamide (286); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)azetidin-1-yl)nicotinamide (287); 4-(3-(4-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (288);N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyridazine-3-carboxamide (289); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-

(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrazine-2-carboxamide (290); N-((S)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-(1-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propoxy)ethyl)-1H-pyrazole-3-carboxamide (291); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-((3S)-4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-3-(hydroxymethyl)piperazin-1-yl)picolinamide (292); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (293); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(9-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-3,9-diazaspiro[5.5]undecan-3-yl)nicotinamide (294); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(9-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-3,9-diazaspiro[5.5]undecan-3-yl)pyrimidine-5-carboxamide (295); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-((1R,4R)-5-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethyl)benzamide (296); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)-2,6-difluorobenzamide (297); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)-2,6-difluorobenzamide (298); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)-2-fluorobenzamide (299); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4-(trifluoromethyl)pyrimidine-5-carboxamide (300); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-3'-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-[1,1'-biphenyl]-4-carboxamide (301); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-3-(trifluoromethyl)benzamide (302); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-3'-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)-[1,1'-biphenyl]-4-carboxamide (303); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S,5R)-4-(2-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-

dioxoisoindolin-5-yl)piperidin-4-yl)ethyl)-3,5-dimethylpiperazin-1-yl)benzamide (304); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-2-carboxamide (305); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrimidine-2-carboxamide (306); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperazin-1-yl)benzamide (307); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)azetidin-1-yl)benzamide (308); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)azetidin-1-yl)ethyl)piperazin-1-yl)benzamide (309); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4-methylpyrimidine-5-carboxamide (310); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (311); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)ethyl)benzamide (312); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-3'-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethoxy)-[1,1'-biphenyl]-4-carboxamide (313); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S,5R)-4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)-3,5-dimethylpiperazin-1-yl)benzamide (314); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)azetidin-3-yl)methyl)piperazin-1-yl)benzamide (315); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperazin-1-yl)benzamide (316); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((3S)-4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-3-(hydroxymethyl)piperazin-1-yl)nicotinamide (317); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)-4-fluoronicotinamide

(318); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)-2,6-difluorobenzamide (319); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)azetidin-1-yl)benzamide (320); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S)-3-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methoxy)pyrrolidin-1-yl)benzamide (321); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((3S)-1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methyl)piperazin-1-yl)benzamide (322); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((3R)-1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methyl)piperazin-1-yl)benzamide (323); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-3-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)butyl)benzamide (324); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methoxy)pyrrolidin-1-yl)benzamide (325); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((3R,5R)-4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-3,5-dimethylpiperazin-1-yl)pyrimidine-5-carboxamide (326); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)benzamide (327); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)benzamide (328); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)butyl)piperazin-1-yl)benzamide (329); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)azetidin-3-yl)propyl)benzamide (330); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-4-yl)butyl)piperazin-1-yl)benzamide (331); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-3-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)cyclobutane-1-carboxamide (332); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrazine-2-

carboxamide (333); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3R)-3-((((1r,3R)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(methyl)amino)methyl)pyrrolidin-1-yl)benzamide (334); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((3R)-1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methoxy)piperidin-1-yl)benzamide (335); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((3S)-1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methoxy)piperidin-1-yl)benzamide (336); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(methyl)amino)methyl)azetidin-1-yl)benzamide (337); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((3S)-3-((((1r,3S)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(methyl)amino)methyl)pyrrolidin-1-yl)benzamide (338); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4-fluoronicotinamide (339); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((((3S)-1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methyl)(methyl)amino)piperidin-1-yl)benzamide (340); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((((3R)-1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methyl)(methyl)amino)piperidin-1-yl)benzamide (341); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-((4aR,6S,8aS)-6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)octahydroisoquinolin-2(1H)-yl)nicotinamide (342); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-3-fluorobenzamide (343); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)benzamide (344); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(((3S)-1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)pyrrolidin-3-yl)methyl)piperazin-1-yl)benzamide (345); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (346); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-

tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-3-(2-(2-methoxyethoxy)ethoxy)benzamide (347); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)azetidin-3-yl)methyl)piperidin-1-yl)benzamide (348); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)oxy)piperidin-1-yl)benzamide (349); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperazin-1-yl)-4-fluoronicotinamide (350); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)azetidin-3-yl)methyl)azetidin-1-yl)benzamide (351); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-3,6-dihydropyridin-1(2H)-yl)benzamide (352); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-((1'-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-[1,4'-bipiperidin]-4-yl)oxy)benzamide (353); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (354); N-((1r,3r)-3-(4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (355); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (356); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(6-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)-2-azaspiro[3.3]heptan-2-yl)benzamide (357); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (358); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrazine-2-carboxamide (359); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyridazine-3-carboxamide (360); 2-chloro-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-

1-yl)methyl)piperidin-1-yl)benzamide (361); 2-chloro-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (362); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)pyrimidine-2-carboxamide (363); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)picolinamide (364); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((3S,5S)-4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-3,5-dimethylpiperazin-1-yl)pyrimidine-5-carboxamide (365); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethoxy)propyl)benzamide (366); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl)-4-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)oxy)butyl)-1H-1,2,3-triazol-1-yl)benzamide (367); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-((5-(2-((R)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)pentyl)oxy)ethyl)piperazin-1-yl)nicotinamide (368); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(2-((5-(2-((S)-2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)pentyl)oxy)ethyl)piperazin-1-yl)benzamide (369); (2S)-1-(4-(((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)-N-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)butyl)-N-methylpyrrolidine-2-carboxamide (370); (2R)-1-(4-(((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)benzyl)-N-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)ethyl)-N-methylpyrrolidine-2-carboxamide (371); (2S)-1-(4-(((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)benzyl)-N-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)ethyl)-N-methylpyrrolidine-2-carboxamide (372); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethoxy)ethyl)benzamide (373); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)pyrrolidin-3-yl)-1H-pyrazole-4-carboxamide (374); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(2-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propoxy)ethyl)-1H-pyrazole-4-carboxamide (375); 4-

(3-(4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (376); 2-chloro-4-(3-(4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile (377); 4-(5-(4-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile (378); N-((R)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-((2-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethoxy)ethoxy)methyl)-1H-pyrazole-3-carboxamide (379); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)picolinamide (380); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(3-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)picolinamide (381); 4-(3-(6-(4-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)pyridin-3-yl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (382); 4-(3-(4-(4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazine-1-carbonyl)-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (383); N-((1r,3S)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-((1S,4S)-5-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethyl)benzamide (384); 4-(5-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)pyridin-2-yl)-1-(5-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)-1-methylpiperazin-1-ium (385); 4-(3-(2-((5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)oxy)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (386); 4-(3-(2-((6-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)hexyl)oxy)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (387); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-2-fluorobenzamide (388); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-4-yl)oxy)propyl)piperazin-1-yl)benzamide (389); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)azetidin-3-

yl)ethyl)benzamide (390); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-2,6-difluorobenzamide (391); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-4-yl)oxy)propyl)piperazin-1-yl)benzamide (392); 4-(3-(2-(2-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperazin-1-yl)ethyl)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (393); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-4-yl)azetidin-3-yl)propyl)benzamide (394); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-4-yl)azetidin-3-yl)methyl)piperidin-1-yl)benzamide (395); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperidin-4-yl)-1H-pyrazole-3-carboxamide (396); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-4-yl)-1H-pyrazole-3-carboxamide (397); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(2-(dimethylamino)ethoxy)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide (398); N-((1s,4s)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide (399); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide (400); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (401); N-((1s,4s)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (402); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (403); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrazine-2-carboxamide (404); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-

yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (405); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (406); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (407); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrazine-2-carboxamide (408); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)benzamide (409); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)piperazin-1-yl)benzamide (410); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)piperazin-1-yl)pyridazine-3-carboxamide (411); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)piperazin-1-yl)pyrimidine-5-carboxamide (412); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)pyridazine-3-carboxamide (413); N-((1r,4S)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-((3S)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)pyrimidine-5-carboxamide (414); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyridazine-3-carboxamide (415); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)benzamide (416); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxamide (417); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrimidine-5-carboxamide (418);N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)-4-methylcyclohexyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (419); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyrazine-2-carboxamide (420); N-((1r,4r)-4-(3-chloro-4-

cyanophenoxy)cyclohexyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)picolinamide (421); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)-5-fluoronicotinamide (422); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrazine-2-carboxamide (423); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrimidine-5-carboxamide (424); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)benzamide (425); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrazine-2-carboxamide (426); N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyridazine-3-carboxamide (427); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide (428); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)-1H-pyrazole-4-carboxamide (429); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)-1H-pyrazole-3-carboxamide (430); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(4-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)pyrrolidin-3-yl)-1H-pyrazole-4-carboxamide (431); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)pyrrolidin-3-yl)-1H-pyrazole-4-carboxamide (432); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(2-(3-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propoxy)ethyl)-1H-pyrazole-4-carboxamide (433); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)picolinamide (434); N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)picolinamide

(435); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(3-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)picolinamide (436); 6-(4-(4-((2-(1-butyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)-N,N-dimethylpyridazine-3-carboxamide (437); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(4-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)pyrimidine-5-carboxamide (438); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (439); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (440); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl methyl carbonate (441); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl ethyl carbonate (442); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl isopropyl carbonate (443); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (tetrahydro-2H-pyran-4-yl) carbonate (444); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-acetamidoethyl)carbamate (445); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-(2-aminoacetamido)ethyl)carbamate (446); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-

dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-aminopropanamido)ethyl)carbamate (447); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-amino-3-methylbutanamido)ethyl)carbamate (448); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-((S)-2-amino-3-methylbutanamido)-3-methylbutanamido)ethyl)carbamate (449); (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2,5,8,11-tetraoxatridecan-13-yl) carbonate (450); 2-chloro-4-(3-(3-fluoro-4-(5-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile (451); 2-chloro-4-(5-(3-fluoro-4-(5-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)benzonitrile (452); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azadodecan-12-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (453); 4-(3-{4-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)ethoxy]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (454); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10,13,16-pentaoxa-1-azaoctadecan-18-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (455); 4-[3-(4-{2-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)ethoxy]ethoxy}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (456); 4-[3-(4-{2-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)ethoxy]ethoxy}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (457); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azatetradecan-14-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (458); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azatridecan-13-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-

(trifluoromethyl)benzonitrile (459); 4-(3-{4-[(1-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}-4,7,10-trioxa-1-azadodecan-12-yl)oxy]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (460); 4-(3-{4-[(1-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}-4,7,10-trioxa-1-azadodecan-12-yl)oxy]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (461); 4-[3-(4-{3-[3-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethoxy)propoxy]propoxy}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (462); 4-{4,4-dimethyl-3-[4-({1-[2-(3-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]-4,7,10-trioxa-1-azatridecan-13-yl}oxy)phenyl]-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (463); 4-[3-(4-{4-[(5-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}pentyl)oxy]phenyl}phenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile (464); 4-{[5-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)pentyl]oxy}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (465); 4-{3-[4-({1-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]-1,4,7,10-tetraoxatridecan-13-yl}oxy)phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl}-2-(trifluoromethyl)benzonitrile (466); 6-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (467); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (468); 4-(3-{4-[1-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethyl)-1H-1,3-benzodiazol-5-yl]phenyl}-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (469); 4-{[5-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)pentyl]amino}-3-fluoro-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (470); 6-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (471); 6-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]-N-

[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (472); 6-(4-{3-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]propyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (473); 6-{4-[2-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (474); 4-{4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethyl)piperazin-1-yl]butyl}-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (475); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (476); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (477); 4-(6-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}hexyl)-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (478); 4-{3-[4-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propyl)piperazin-1-yl]propyl}-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (479); 4-[5-(3-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)pentyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (480); 6-{4-[2-(3-{[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}propoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (481); 4-(4-{4-[2-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}oxy)ethyl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (482); 4-(4-{4-[2-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-4-yl}oxy)ethyl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (483); 4-{4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]butyl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (484); 6-{4-[5-({2-[(3S)-2,6-

dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-4-yl}oxy)pentyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (485); 4-(4-{4-[2-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl}oxy)ethyl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (486); 6-{4-[5-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}oxy)pentyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (487); 4-(4-{4-[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (488); 4-{4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]butyl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (489); 6-(4-{6-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]hexyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (490); 4-(4-{4-[2-(2,6-dioxopiperidin-3-yl)-4-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (491); 4-(6-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (492); 6-{4-[5-({2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl}oxy)pentyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (493); 6-[4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]pyridine-3-carboxamide (494); 4-[4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (495); 4-(5-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}pentyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (496); 4-(4-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (497); 4-(6-{4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-

tetramethylcyclobutyl]benzamide (498); 4-(4-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}butyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (499); 4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (500); 4-[3-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)propyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (501); 4-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)butyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (502); 4-[5-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)pentyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (503); 4-[6-(2-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}ethoxy)hexyl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (504); N-[(2R)-1-[3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl]propan-2-yl]-5-{[4-(4-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}butyl)piperazin-1-yl]methyl}-1H-pyrazole-3-carboxamide (505); 4-(4-{6-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]hexyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (506); 4-(3-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}propyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (507); N-[(2R)-1-[3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl]propan-2-yl]-5-[(3-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}propoxy)methyl]-1H-pyrazole-3-carboxamide (508); 4-[4-(3-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}propyl)-1H-1,2,3-triazol-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (509); 6-[4-(6-{[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]amino}hexyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (510); N-[(2R)-1-[3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl]propan-2-yl]-5-({3-[4-(2-{[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl]amino}ethyl)piperazin-1-yl]propoxy}methyl)-1H-pyrazole-3-carboxamide (511); 4-[4-(5-{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}pentyl)-1H-1,2,3-triazol-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (512); 4-[4-(6-

{[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}hexyl)-1H-1,2,3-triazol-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)cyclobutyl]benzamide (513); 6-{4-[2-(4-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}butoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (514); 6-{4-[2-(4-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}butoxy)ethyl]piperazin-1-yl}-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (515); 6-(4-{2-[(5-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (516); 6-(4-{2-[(5-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (517); 4-(4-{2-[(5-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (518); 4-(4-{2-[(5-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1,3-dioxo-2,3-dihydro-1H-isoindol-4-yl}pentyl)oxy]ethyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (519); 4-(4-{6-[2-(2,6-dioxopiperidin-3-yl)-1-oxo-2,3-dihydro-1H-isoindol-5-yl]hexyl}piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (520); 4-(3-{4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl}propyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (521); 4-[6-(4-{2-[(3S)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (522); 4-[6-(4-{2-[(3R)-2,6-dioxopiperidin-3-yl]-1-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (523); 6-[4-(6-{[2-(2,6-dioxopiperidin-3-yl)-5-fluoro-1-oxo-2,3-dihydro-1H-isoindol-4-yl]oxy}hexyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (524); 6-[4-(6-{[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy}hexyl)piperazin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (525); 4-[6-(4-{2-[(3S)-2,6-dioxopiperidin-3-yl]-3-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-

1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (526); 4-[6-(4-{2-[(3R)-2,6-dioxopiperidin-3-yl]-3-oxo-2,3-dihydro-1H-isoindol-5-yl}piperazin-1-yl)hexyl]-N-[(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide (527); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-3-oxo-2,3-dihydro-[1,2,4]triazolo[4,3-a]pyridin-6-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (528); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4-methylene-1-oxo-1,2,3,4-tetrahydroisoquinolin-6-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (529); rac-N-(1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)azetidin-3-yl)-1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-N-methylpiperidine-4-carboxamide (530); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyridazine-3-carboxamide (531); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrazine-2-carboxamide (532); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrimidine-5-carboxamide (533); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrimidine-5-carboxamide (534); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4-(2-(2-methoxyethoxy)ethoxy)nicotinamide (535); rac-N-((1r,4R)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(2-((2R,6R)-4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-2,6-dimethylpiperazin-1-yl)ethyl)benzamide (536); rac-N-((1r,3r)-3-(3,4-dicyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (537); rac-N-((1r,3r)-3-(4-cyano-2-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (538); rac-N-((1r,3r)-3-(2,4-dicyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (539); rac-N-((1r,3r)-3-(4-

cyano-2,6-dimethylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (540); rac-N-((1r,3r)-3-(4-cyano-3-methoxyphenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (541); N-((1r,4r)-4-(4-cyano-3-methylphenoxy)cyclohexyl)-5-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrazine-2-carboxamide (542); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)thio)ethyl)piperazin-1-yl)benzamide (543); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (544);
rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6,7-difluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (545); rac-2-chloro-4-(((1r,4r)-4-(5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-1H-benzo[d]imidazol-2-yl)cyclohexyl)oxy)benzonitrile (546); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)nicotinamide (547); rac-N-((1r,4r)-4-(3-chloro-4-cyano-2-methylphenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (548); rac-N-((1r,4r)-4-(4-cyano-3-methylphenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (549); rac-N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (550); rac-N-((1r,3r)-3-((5-cyano-6-methylpyridin-2-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (551); rac-N-((1r,3r)-3-((5-cyanopyridin-2-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (552); rac-N-((1r,3r)-3-(3,4-dicyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (553); rac-N-((1r,3r)-3-((5-cyano-6-methylpyridin-2-yl)oxy)-

2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (554); rac-N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((1R,4R,5S)-5-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidine-5-carboxamide (555); rac-N-((1r,3r)-3-((5-cyano-3-methylpyridin-2-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (556); rac-N-((1r,3r)-3-((5-cyanopyrimidin-2-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (557); rac-N-((1r,3r)-3-(4-cyano-3,5-dimethylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (558); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)pyrazine-2-carboxamide (559); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)pyrimidine-5-carboxamide (560); rac-N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-((1R,4R,5R)-5-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)-2-azabicyclo[2.2.1]heptan-2-yl)pyrimidine-5-carboxamide (561); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-((2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)(methyl)amino)pyrrolidin-1-yl)benzamide (562); rac-N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1R,3S)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclopentyl)piperazin-1-yl)benzamide (563); rac-N-((1r,3r)-3-((6-cyano-5-methylpyridin-3-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (564); rac-N-((1r,3r)-3-((6-cyano-5-(trifluoromethyl)pyridin-3-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (565); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)pyrazine-2-carboxamide (566); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)pyrazine-2-carboxamide (567); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-

tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (568); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)-4-fluoronicotinamide (569); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (570); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (571); rac-N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (572); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)pyridazine-3-carboxamide (573); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)pyrimidine-5-carboxamide (574); rac-N-((1r,4r)-4-(4-cyano-3-methylphenoxy)cyclohexyl)-2-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)pyrimidine-5-carboxamide (575); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)pyridazine-3-carboxamide (576); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (577); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (578); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrazine-2-carboxamide (579); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (580); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4,6-difluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4,6-dimethylpyrimidine-5-carboxamide (581); rac-N-((1r,3r)-3-(3-

chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-4,6-difluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyridazine-3-carboxamide (582); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-4-(2-morpholinoethoxy)nicotinamide (583); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(3-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)azetidin-1-yl)pyrimidine-5-carboxamide (584); rac-N-((1r,3r)-3-((5-cyano-6-ethylpyridin-2-yl)oxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (585); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(3-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)azetidin-1-yl)pyridazine-3-carboxamide (586); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)benzamide (587); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(1-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperidin-4-yl)benzamide (588); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(3-(((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)oxy)methyl)pyrrolidin-1-yl)benzamide (589); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-oxoisoindolin-5-yl)piperidin-4-yl)methyl)piperazin-1-yl)benzamide (590); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((1-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperidin-4-yl)methyl)piperazin-1-yl)benzamide (591); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4,6-difluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (592); rac-2-chloro-4-(((1r,4r)-4-(5-(4'-(2-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)-1-methyl-1H-imidazol-5-yl)-[1,1'-biphenyl]-4-yl)-1-methyl-1H-imidazol-2-yl)cyclohexyl)oxy)benzonitrile (593); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4,6-difluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (594); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4,6-difluoro-

1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (595); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)benzamide (596); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrazine-2-carboxamide (597); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(1-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperidin-4-yl)pyrimidine-5-carboxamide (598); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)pyridazine-3-carboxamide (599); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyridazine-3-carboxamide (600); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)picolinamide (601); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(6-(4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxoisoindolin-5-yl)piperazin-1-yl)hexyl)nicotinamide (602); rac-2-chloro-4-(((1r,4r)-4-(2-(4-(3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)azetidin-1-yl)phenyl)-1H-imidazol-5-yl)cyclohexyl)oxy)benzonitrile (603); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)cyclohexyl)-1H-pyrazole-3-carboxamide (604); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)-1H-pyrazole-3-carboxamide (605); rac-N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-1-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)cyclohexyl)-1H-pyrazole-3-carboxamide (606); 3-chloro-5-(5-(4-(1-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperidin-4-yl)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile (607); 3-chloro-5-(5-(4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile (608); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-3-methyl-1-oxoisoindolin-5-

yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (609); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1-methyl-3-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (610); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-4-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide (611); 3-chloro-5-(5-(4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile (612); 3-chloro-5-(5-(4-(1-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)methyl)piperidin-4-yl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)picolinonitrile (613); 5-(4-((1-(2-(4-(6,7-dihydro-5H-pyrrolo[1,2-a]imidazol-3-yl)phenoxy)ethyl)piperidin-4-yl)methyl)piperazin-1-yl)-2-(2,6-dioxopiperidin-3-yl)isoindoline-1,3-dione (614); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (615); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(5-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazine-1-carbonyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)benzamide (616); rac-N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-((1r,4R)-4-((1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)(methyl)amino)cyclohexyl)-1H-pyrazole-3-carboxamide (617); 5-(4-((1-(4-(6,7-dihydro-5H-pyrrolo[1,2-a]imidazol-3-yl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-2-(2,6-dioxopiperidin-3-yl)isoindoline-1,3-dione (618); rac-N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(3-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)azetidin-1-yl)pyrazine-2-carboxamide (619); N-((1r,3R)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-((1R,4R)-4-((((1r,3R)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)cyclohexyl)-1H-pyrazole-3-carboxamide (620); N-((1r,3r)-3-(4-cyano-3-methoxyphenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (621); N-((1r,3r)-3-(4-cyano-3-methylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (622); N-

((1r,3r)-3-(4-cyano-3,5-dimethylphenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (623); N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-3-oxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (624); and N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-((((1r,3r)-3-((2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolin-5-yl)oxy)cyclobutyl)(isopropyl)amino)methyl)piperidin-1-yl)pyrimidine-5-carboxamide (625).

[0176] In another embodiment, the present disclosure provides a library of compounds. The library comprises more than one compound wherein each compound has a formula of ABM-L-ULM, wherein ULM is a ubiquitin pathway protein binding moiety (preferably, an E3 ubiquitin ligase moiety as otherwise disclosed herein), e.g., a CLM, and ABM is an AR protein binding moiety, wherein ABM is coupled (preferably, through a linker moiety) to ULM, and wherein the ubiquitin pathway protein binding moiety recognizes an ubiquitin pathway protein, in particular, an E3 ubiquitin ligase.

[0177] The present description includes, where applicable, the compositions comprising the pharmaceutically acceptable salts, in particular, acid or base addition salts of compounds of the present disclosure.

[0178] The term "pharmaceutically acceptable salt" is used throughout the specification to describe, where applicable, a salt form of one or more of the compounds described herein which are presented to increase the solubility of the compound in the gastic juices of the patient's gastrointestinal tract in order to promote dissolution and the bioavailability of the compounds. Pharmaceutically acceptable salts include those derived from pharmaceutically acceptable inorganic or organic bases and acids, where applicable. Suitable salts include those derived from alkali metals such as potassium and sodium, alkaline earth metals such as calcium, magnesium and ammonium salts, among numerous other acids and bases well known in the pharmaceutical art. Sodium and potassium salts are particularly preferred as neutralization salts of the phosphates according to the present disclosure.

[0179] The acids which are used to prepare the pharmaceutically acceptable acid addition salts of the aforementioned base compounds useful in this disclosure are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the

hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3 naphthoate)]salts, among numerous others.

[0180] Pharmaceutically acceptable base addition salts may also be used to produce pharmaceutically acceptable salt forms of the compounds or derivatives according to the present disclosure. The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of the present compounds that are acidic in nature are those that form non-toxic base salts with such compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmacologically acceptable cations such as alkali metal cations (eg., potassium and sodium) and alkaline earth metal cations (eg, calcium, zinc and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines, among others.

[0181] **Compositions**

[0182] In another aspect, the description provides compositions comprising compounds as described herein, including salts thereof, and a pharmaceutically acceptable carrier. In certain embodiments, the compositions are therapeutic or pharmaceutical compositions comprising an effective amount of a compound as described herein and a pharmaceutally acceptable carrier.

[0183] The amount of compound in a pharmaceutical composition of the instant disclosure that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host and disease treated, the particular mode of administration. Generally, an amount between 0.1 mg/kg and 1000 mg/kg body weight/day of active ingredients is administered dependent upon potency of the agent. Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds that exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to

uninfected cells and, thereby, reduce side effects. The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the present disclosure, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0184] The compositions of the present disclosure may be formulated in a conventional manner using one or more pharmaceutically acceptable carriers and may also be administered in controlled-release formulations. Pharmaceutically acceptable carriers that may be used in these pharmaceutical compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as prolamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0185] The active compound is included in the pharmaceutically acceptable carrier or diluent in an amount sufficient to deliver to a patient a therapeutically effective amount for the desired indication, without causing serious toxic effects in the patient treated. A preferred dose of the active compound for all of the herein-mentioned conditions is in the range from about 10 ng/kg to 300 mg/kg, preferably 0.1 to 100 mg/kg per day, more generally 0.5 to about 25 mg per kilogram body weight of the recipient/patient per day. A typical topical dosage will range from 0.01-5% wt/wt in a suitable carrier.

[0186] The compound is conveniently administered in any suitable unit dosage form, including but not limited to one containing less than 1mg, 1 mg to 3000 mg, preferably 5 to 500

mg of active ingredient per unit dosage form. An oral dosage of about 25-250 mg is often convenient.

[0187] The active ingredient is preferably administered to achieve peak plasma concentrations of the active compound of about 0.00001-30 mM, preferably about 0.1-30 μM. This may be achieved, for example, by the intravenous injection of a solution or formulation of the active ingredient, optionally in saline, or an aqueous medium or administered as a bolus of the active ingredient. Oral administration is also appropriate to generate effective plasma concentrations of active agent.

[0188] The concentration of active compound in the drug composition will depend on absorption, distribution, inactivation, and excretion rates of the drug as well as other factors known to those of skill in the art. It is to be noted that dosage values will also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that the concentration ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition. The active ingredient may be administered at once, or may be divided into a number of smaller doses to be administered at varying intervals of time.

[0189] If administered intravenously, preferred carriers are physiological saline or phosphate buffered saline (PBS).

[0190] In one embodiment, the active compounds are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art.

[0191] Liposomal suspensions may also be pharmaceutically acceptable carriers. These may be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811 (which is incorporated herein by reference in its entirety). For example, liposome formulations may be prepared by dissolving appropriate lipid(s) (such as stearoyl phosphatidyl ethanolamine, stearoyl phosphatidyl choline, arachadoyl phosphatidyl

choline, and cholesterol) in an inorganic solvent that is then evaporated, leaving behind a thin film of dried lipid on the surface of the container. An aqueous solution of the active compound are then introduced into the container. The container is then swirled by hand to free lipid material from the sides of the container and to disperse lipid aggregates, thereby forming the liposomal suspension.

[0192] Modes of Adminstration

[0193] In any of the aspects or embodiments described herein, the therapeutic compositions comprising compounds described herein can be in any suitable dosage form configured to be delivered by any suitable route. For example, the compounds can be administered by any appropriate route, for example, orally, parenterally, intravenously, intradermally, subcutaneously, or topically, including transdermally, in liquid, cream, gel, or solid form, rectally, nasally, buccally, vaginally or via an implanted reservoir or by aerosol form.

[0194] The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously.

[0195] The compounds as described herein may be administered in single or divided doses by the oral, parenteral or topical routes. Administration of the active compound may range from continuous (intravenous drip) to several oral administrations per day (for example, Q.I.D.) and may include oral, topical, parenteral, intramuscular, intravenous, sub-cutaneous, transdermal (which may include a penetration enhancement agent), buccal, sublingual and suppository administration, among other routes of administration. Enteric coated oral tablets may also be used to enhance bioavailability of the compounds from an oral route of administration. The most effective dosage form will depend upon the pharmacokinetics of the particular agent chosen as well as the severity of disease in the patient.

[0196] Administration of compounds as sprays, mists, or aerosols for intra-nasal, intra-tracheal or pulmonary administration may also be used. Compounds as described herein may be administered in immediate release, intermediate release or sustained or controlled release forms. Sustained or controlled release forms are preferably administered orally, but also in suppository and transdermal or other topical forms. Intramuscular injections in liposomal form may also be used to control or sustain the release of compound at an injection site.

[0197] Sterile injectable forms of the compositions as described herein may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1, 3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as Ph. Helv or similar alcohol.

[0198] The pharmaceutical compositions as described herein may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added. Oral compositions will generally include an inert diluent or an edible carrier. They may be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound or its prodrug derivative can be incorporated with excipients and used in the form of tablets, troches, or capsules. Pharmaceutically compatible binding agents, and/or adjuvant materials are included as part of the composition.

[0199] The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a dispersing agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring. When the dosage unit form is a capsule, it can contain, in addition to material of the above type, a liquid carrier such as a fatty oil.

In addition, dosage unit forms can contain various other materials which modify the physical form of the dosage unit, for example, coatings of sugar, shellac, or enteric agents.

[0200] The active compound or pharmaceutically acceptable salt thereof can be administered as a component of an elixir, suspension, syrup, wafer, chewing gum or the like. A syrup may contain, in addition to the active compounds, sucrose as a sweetening agent and certain preservatives, dyes and colorings and flavors.

[0201] Alternatively, the pharmaceutical compositions as described herein may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient, which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0202] The pharmaceutical compositions of the disclosure may also be administered topically. Suitable topical formulations are readily prepared for each of these areas or organs. Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-acceptable transdermal patches may also be used. For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this disclosure include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. In certain preferred aspects of the disclosure, the compounds may be coated onto a stent which is to be surgically implanted into a patient in order to inhibit or reduce the likelihood of occlusion occurring in the stent in the patient.

[0203] Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0204] For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with our without a preservative such as benzylalkonium

chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

[0205] The pharmaceutical compositions of this disclosure may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[0206] Solutions or suspensions used for parenteral, intradermal, subcutaneous, or topical application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parental preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0207] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease or condition being treated.

[0208] A patient or subject in need of therapy using compounds as described herein can be treated by administering to the patient (subject) an effective amount of the compound including pharmaceutically acceptable salts, solvates or polymorphs, thereof optionally in a pharmaceutically acceptable carrier or diluent, either alone, or in combination with other known agents.

[0209] **Co-administration**

[0210] Disease states of conditions which may be treated using compounds or compositions according to the present description include, but not limited to, for example, cancer (e.g., prostate cancer), and Kennedy's disease. In certain embodiments, the therapeutic or pharmaceutical compositions comprise an effective amount of an additional biologically or bioactive active agent, e.g., an agent effective for the treatment of cancer, that is co-administered.

[0211] The term "coadministration" or "combination therapy" shall mean that at least two compounds or compositions are administered to the patient at the same time, such that effective amounts or concentrations of each of the two or more compounds may be found in the patient at a given point in time. Although compounds according to the present disclosure may be co-administered to a patient at the same time, the term embraces both administration of two or more agents at the same time or at different times, provided that effective concentrations of all coadministered compounds or compositions are found in the subject at a given time. In certain preferred aspects of the present disclosure, one or more of the present compounds described above, are coadministered in combination with at least one additional bioactive agent, especially including an anticancer agent. In particularly preferred aspects of the disclosure, the co-administration of compounds results in synergistic therapeutic, including anticancer therapy.

[0212] In another aspect, the description provides a composition comprising an effective amount of two or more of the PROTAC compounds as described herein, and a pharmaceutically acceptable carrier. In certain embodiments, the composition further comprises an effective or synergistic amount of another bioactive agent that is not a PROTAC compound.

[0213] Pharmaceutical compositions comprising combinations of an effective amount of at least one bifunctional compound according to the present disclosure, and one or more of the compounds otherwise described herein, all in effective amounts, in combination with a pharmaceutically effective amount of a carrier, additive or excipient, represents a further aspect of the present disclosure.

[0214] The term "bioactive agent" is used to describe an agent, other than the PROTAC compounds described herein, which is used in combination with the present compounds as an agent with biological activity to assist in effecting an intended therapy, inhibition and/or prevention/prophylaxis for which the present compounds are used. Preferred bioactive agents for use herein include those agents which have pharmacological activity similar to that for which the present compounds are used or administered and include for example, anti-cancer agents.

[0215] The term "additional anti-cancer agent" is used to describe an anti-cancer agent, which may be combined with PROTAC compounds according to the present description to treat cancer. These agents include, for example, everolimus, trabectedin, abraxane, TLK 286, AV-299, DN-101, pazopanib, GSK690693, RTA 744, ON 0910.Na, AZD 6244 (ARRY-142886), AMN-107, TKI-258, GSK461364, AZD 1152, enzastaurin, vandetanib, ARQ-197, MK-0457,

MLN8054, PHA-739358, R-763, AT-9263, a FLT-3 inhibitor, an androgen receptor inhibitor, a VEGFR inhibitor, an EGFR TK inhibitor, an aurora kinase inhibitor, a PIK-1 modulator, a Bcl-2 inhibitor, an HDAC inhbitor, a c-MET inhibitor, a PARP inhibitor, a Cdk inhibitor, an EGFR TK inhibitor, an IGFR-TK inhibitor, an anti-HGF antibody, a PI3 kinase inhibitors, an AKT inhibitor, a JAK/STAT inhibitor, a checkpoint-1 or 2 inhibitor, a focal adhesion kinase inhibitor, a Map kinase kinase (mek) inhibitor, a VEGF trap antibody, pemetrexed, erlotinib, dasatanib, nilotinib, decatanib, panitumumab, amrubicin, oregovomab, Lep-etu, nolatrexed, azd2171, batabulin, ofatumumab, zanolimumab, edotecarin, tetrandrine, rubitecan, tesmilifene, oblimersen, ticilimumab, ipilimumab, gossypol, Bio 111, 131-I-TM-601, ALT-110, BIO 140, CC 8490, cilengitide, gimatecan, IL13-PE38QQR, INO 1001, $IPdR_1$ KRX-0402, lucanthone, LY317615, neuradiab, vitespan, Rta 744, Sdx 102, talampanel, atrasentan, Xr 311, romidepsin, ADS-100380, sunitinib, 5-fluorouracil, vorinostat, etoposide, gemcitabine, doxorubicin, liposomal doxorubicin, 5'-deoxy-5-fluorouridine, vincristine, temozolomide, ZK-304709, seliciclib; PD0325901, AZD-6244, capecitabine, L-Glutamic acid, N-[4-[2-(2-amino-4,7-dihydro-4-oxo-1 H - pyrrolo[2,3-d]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate, camptothecin, PEG-labeled irinotecan, tamoxifen, toremifene citrate, anastrazole, exemestane, letrozole, DES(diethylstilbestrol), estradiol, estrogen, conjugated estrogen, bevacizumab, IMC-1C11, CHIR-258); 3-[5-(methylsulfonylpiperadinemethyl)- indolylj-quinolone, vatalanib, AG-013736, AVE-0005, the acetate salt of [D- Ser(Bu t) 6 ,Azgly 10] (pyro-Glu-His-Trp-Ser-Tyr-D-Ser(Bu t)-Leu-Arg-Pro- Azgly-NH_2 acetate [$C_{59}H_{84}N_{18}Oi_4$ -$(C_2H_4O_2)_X$ where x = 1 to 2.4], goserelin acetate, leuprolide acetate, triptorelin pamoate, medroxyprogesterone acetate, hydroxyprogesterone caproate, megestrol acetate, raloxifene, bicalutamide, flutamide, nilutamide, megestrol acetate, CP-724714; TAK-165, HKI-272, erlotinib, lapatanib, canertinib, ABX-EGF antibody, erbitux, EKB-569, PKI-166, GW-572016, Ionafarnib, BMS-214662, tipifarnib; amifostine, NVP-LAQ824, suberoyl analide hydroxamic acid, valproic acid, trichostatin A, FK-228, SU11248, sorafenib, KRN951 , aminoglutethimide, arnsacrine, anagrelide, L-asparaginase, Bacillus Calmette-Guerin (BCG) vaccine, adriamycin, bleomycin, buserelin, busulfan, carboplatin, carmustine, chlorambucil, cisplatin, cladribine, clodronate, cyproterone, cytarabine, dacarbazine, dactinomycin, daunorubicin, diethylstilbestrol, epirubicin, fludarabine, fludrocortisone, fluoxymesterone, flutamide, gleevec, gemcitabine, hydroxyurea, idarubicin, ifosfamide, imatinib, leuprolide, levamisole, lomustine, mechlorethamine, melphalan, 6-

mercaptopurine, mesna, methotrexate, mitomycin, mitotane, mitoxantrone, nilutamide, octreotide, oxaliplatin, pamidronate, pentostatin, plicamycin, porfimer, procarbazine, raltitrexed, rituximab, streptozocin, teniposide, testosterone, thalidomide, thioguanine, thiotepa, tretinoin, vindesine, 13-cis-retinoic acid, phenylalanine mustard, uracil mustard, estramustine, altretamine, floxuridine, 5-deooxyuridine, cytosine arabinoside, 6-mecaptopurine, deoxycoformycin, calcitriol, valrubicin, mithramycin, vinblastine, vinorelbine, topotecan, razoxin, marimastat, COL-3, neovastat, BMS-275291 , squalamine, endostatin, SU5416, SU6668, EMD121974, interleukin-12, IM862, angiostatin, vitaxin, droloxifene, idoxyfene, spironolactone, finasteride, cimitidine, trastuzumab, denileukin diftitox,gefitinib, bortezimib, paclitaxel, cremophor-free paclitaxel, docetaxel, epithilone B, BMS- 247550, BMS-310705, droloxifene, 4-hydroxytamoxifen, pipendoxifene, ERA-923, arzoxifene, fulvestrant, acolbifene, lasofoxifene, idoxifene, TSE-424, HMR- 3339, ZK186619, topotecan, PTK787/ZK 222584, VX-745, PD 184352, rapamycin, 40-O-(2-hydroxyethyl)-rapamycin, temsirolimus, AP-23573, RAD001, ABT-578, BC-210, LY294002, LY292223, LY292696, LY293684, LY293646, wortmannin, ZM336372, L-779,450, PEG-filgrastim, darbepoetin, erythropoietin, granulocyte colony-stimulating factor, zolendronate, prednisone, cetuximab, granulocyte macrophage colony-stimulating factor, histrelin, pegylated interferon alfa-2a, interferon alfa-2a, pegylated interferon alfa-2b, interferon alfa-2b, azacitidine, PEG-L-asparaginase, lenalidomide, gemtuzumab, hydrocortisone, interleukin-11, dexrazoxane, alemtuzumab, all-transretinoic acid, ketoconazole, interleukin-2, megestrol, immune globulin, nitrogen mustard, methylprednisolone, ibritgumomab tiuxetan, androgens, decitabine, hexamethylmelamine, bexarotene, tositumomab, arsenic trioxide, cortisone, editronate, mitotane, cyclosporine, liposomal daunorubicin, Edwina-asparaginase, strontium 89, casopitant, netupitant, an NK-1 receptor antagonist, palonosetron, aprepitant, diphenhydramine, hydroxyzine, metoclopramide, lorazepam, alprazolam, haloperidol, droperidol, dronabinol, dexamethasone, methylprednisolone, prochlorperazine, granisetron, ondansetron, dolasetron, tropisetron, pegfilgrastim, erythropoietin, epoetin alfa, darbepoetin alfa and mixtures thereof.

[0216] Methods of Treatment

[0217] In another aspect, the disclosure provides methods of modulating protein ubiquitination and degradation in a subject, e.g., a cell, a tissue, mammal, or human patient, the method comprising administering an effective amount of a PROTAC compound as described herein or a composition comprising an effective amount of the same to a subject, wherein the

compound or composition comprising the same is effective in modulating protein ubquitination and degradation of the protein in the subject. In certain embodiments, the protein is androgen receptor (AR).

[0218] In certain embodiments, the description provides a method for regulating protein activity of the androgen receptor in a patient in need comprising administering to said patient an amount of a compound as described herein to a patient.

[0219] In still additional embodiments, the description provides a method of treating a disease state or condition in a patient wherein dysregulated protein activity is responsible for said disease state or condition, said method comprising administering to said patient an effective amount of a compound as described herein to said patient in order to regulate said protein activity in said patient. In certain embodiments, the protein is AR.

[0220] The terms "treat", "treating", and "treatment", etc., as used herein, refer to any action providing a benefit to a patient for which the present compounds may be administered, including the treatment of any disease state or condition which is modulated through the protein to which the present compounds bind. Disease states or conditions, including cancer, which may be treated using compounds according to the present disclosure are set forth hereinabove.

[0221] In another aspect, the disclosure provides methods of modulating AR protein ubiquitination and degradation in a subject, e.g., a cell, a tissue, mammal, or human patient, the method comprising administering an effective amount of a compound as described herein or a composition comprising an effective amount of a compound as described herein to a subject, wherein the compound or composition comprising the same is effective in modulating AR protein ubquitination and degradation of the protein in the subject.

[0222] In another aspect, the disclosure provides methods of treating or ameliorating a symptom of a disease related to AR activity in a subject, e.g., a cell, a tissue, mammal, or human patient, the method comprising administering an effective amount of a compound as described herein or a composition comprising an effective amount of the same to a subject in need thereof, wherein the compound or composition comprising the same is effective in treating or ameliorating a symptom of a disease related to AR activity in the subject.

[0223] In certain embodiments, the disease or disorder is asthma, multiple sclerosis, cancer, prostate cancer, Kenney's disease, ciliopathies, cleft palate, diabetes, heart disease, hypertension, inflammatory bowel disease, mental retardation, mood disorder, obesity, refractive

error, infertility, Angelman syndrome, Canavan disease, Coeliac disease, Charcot–Marie–Tooth disease, Cystic fibrosis, Duchenne muscular dystrophy, Haemochromatosis, Haemophilia, Klinefelter's syndrome, Neurofibromatosis, Phenylketonuria, Polycystic kidney disease, (PKD1) or 4 (PKD2) Prader–Willi syndrome, Sickle-cell disease, Tay–Sachs disease, Turner syndrome. The method according to claim 48 wherein said cancer is squamous-cell carcinoma, basal cell carcinoma, adenocarcinoma, hepatocellular carcinomas, and renal cell carcinomas, cancer of the bladder, bowel, breast, cervix, colon, esophagus, head, kidney, liver, lung, neck, ovary, pancreas, prostate, and stomach; leukemias; benign and malignant lymphomas, particularly Burkitt's lymphoma and Non-Hodgkin's lymphoma; benign and malignant melanomas; myeloproliferative diseases; sarcomas, including Ewing's sarcoma, hemangiosarcoma, Kaposi's sarcoma, liposarcoma, myosarcomas, peripheral neuroepithelioma, synovial sarcoma, gliomas, astrocytomas, oligodendrogliomas, ependymomas, gliobastomas, neuroblastomas, ganglioneuromas, gangliogliomas, medulloblastomas, pineal cell tumors, meningiomas, meningeal sarcomas, neurofibromas, and Schwannomas; bowel cancer, breast cancer, prostate cancer, cervical cancer, uterine cancer, lung cancer, ovarian cancer, testicular cancer, thyroid cancer, astrocytoma, esophageal cancer, pancreatic cancer, stomach cancer, liver cancer, colon cancer, melanoma; carcinosarcoma, Hodgkin's disease, Wilms' tumor or teratocarcinomas. In certain embodiments, the disease to be treated is cancer, e.g., prostate cancer, or Kennedy's Disease. In a preferred embodiment, the subject is a human.

[0224] In another aspect, the disclosure provides methods of treating or ameliorating a symptom of a disease related to AR activity in a subject, e.g., a cell, a tissue, mammal, or human patient, the method comprising administering an effective amount of a compound as described herein or a composition comprising an effective amount of the same and an effective or synergistic amount of another bioactive agent to a subject in need thereof, wherein the composition comprising the same is effective in treating or ameliorating a symptom of a disease related to AR activity in the subject. In certain embodiments, the disease to be treated is cancer, e.g., prostate cancer, or Kennedy's Disease. In a preferred embodiment, the subject is a human. In certain additional embodiments, the additional bioactive agent is an anti-cancer agent.

[0225] In alternative aspects, the present disclosure relates to a method for treating a disease state by degrading a protein or polypeptide through which a disease state or condition is modulated comprising administering to said patient or subject an effective amount of at least one

compound as described hereinabove, optionally in combination with an additional bioactive agent. The method according to the present disclosure may be used to treat a large number of disease states or conditions including cancer, by virtue of the administration of effective amounts of at least one compound described herein.

[0226] In another aspect, the disclosure provides methods for identifying the effects of the degradation of proteins of interest in a biological system using compounds according to the present disclosure.

[0227] **Kits**

[0228] In another aspect, the description provides kits comprising compounds or compositions as described herein. The kit may be promoted, distributed, or sold as a unit for performing the methods of the present disclosure. In addition, the kits of the present disclosure may preferably contain instructions, which describe a suitable use. Such kits can be conveniently used, e.g., in clinical settings, to treat patients exhibiting symptoms of, e.g., cancer, prostate cancer or Kennedy's Disease.

[0229] **EXAMPLES**

[0230] **General Chemistry – Analysis and Synthesis**

[0231] Unless otherwise noted, all materials/reagents were obtained from commercial suppliers and used without further purification. Reactions were monitored by LC-MS and/or thin layer chromatography (TLC) on silica gel 60 F_{254} (0.2mm) pre-coated aluminum foil or glass-backed and visualized using UV light. Flash chromatography (alternatively called "ISCO chromatography") was performed using an ISCO CombiFiash RF 75 PSI or equivalent with RediSep normal-phase silica gel cartridges. Preparative TLC was performed on Whatman LK6F Silica Gel 60A size 20x20 cm plates with a thickness of 1000 μm or equivalent.

[0232] ^{1}HNMR (300 or 400 MHz) and ^{13}CNMR (100.6 MHz) spectra were recorded on Bruker spectrometers at room temperature with TMS or the residual solvent peak as the internal standard. The line positions or multiples are given in (δ) and the coupling constants *(J)* are given as absolute values in Hertz (Hz). The multiplicities in ^{1}HNMR spectra are abbreviated as follows: s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), br or broad (broadened).

[0233] Preparative HPLC purifications were performed on a Waters® UV-Directed Purification System equipped with 2545 Binary Gradient Module, 2767 Sample Manager and 2489 UV/Visible Detector, controlled by MassLynx V4.1 software. All purification work was

completed using the following columns: Atlantis Prep T3 OBD Column, SunFire Prep C18 OBD Column and XBridge Prep Phenyl OBD Column. The mobile phases were water (with 0.1%TFA or 0.01% NH_4HCO_3) and acetonitrile; all reagents used were of HPLC grade. The flow rate was 30ml/min. After the columns, a 1:1000 LC packings flow splitter allowed transfer of a small portion of the eluent into the UV detector. The electrospray source was set at 3.0 kV capillary voltage, 30 V conevoltage, 110°C source temperature, 350°C desolvation temperature, 600L/h desolvation gas flow, and 60L/h cone gas flow. For the analyzer, the multiplier was set at 550 for preparative tune method.

[0234] Analytical LC-MS data was collected on a Shimadzu LCMS-2020 with a mobile phase of 0.05% TFA in Acetonitrile (A) and 0.05% TFA in HPLC grade water (B); 0.1% FA in Acetonitrile (A) and 0.1% FA in HPLC grade water (B); Acetonitrile (A) and 5 mM ammonium bicarbonate in HPLC grade water (B).

[0235] Shimadzu LCMS-2020 equipped with LC-20AD or 30AD pumps, SPD-M20A PDA and Alltech 3300 ELSD. The system uses the following conditions for 2.0 min, 2.6 min, 3 min, 3.6 min, 5 min or 5.6 min run time.

[0236] 2.0 minute run: Kinetex XB-C 18 100A column, 2.6 μm, 3.0x 50 mm. The flow rate is 1.5 mL/min, the run time is 2.0 min, and the gradient profiles are 0.01 min 10% A, 1.10 min 100% A, 1.60 min 100% A, 1.70 min 10% A, 2.00 min 10% A.

[0237] 2.6 minute run: Shim-pack VP-ODS column, 2.2 μm, 3.0x 50 mm. The flow rate is 1.5 mL/min, the run time is 2.6 min, and the gradient profiles are 0.01 min 5% A, 1.20 min 100% A, 2.20 min 100% A, 2.30 min 5% A, 2.60 min 5% A.

[0238] 3.0 minute run: ACE UltraCore Super C18 column, 2.5 μm, 3.0x 50 mm. The flow rate is 1.5 mL/min, the run time is 3.0 min, and the gradient profiles are 0.01 min 10% A, 2.00 min 95% A, 2.60 min 95% A, 2.70 min 10% A, 3.00 min 10% A.

[0239] 3.6 minute run: Shim-pack VP-ODS column, 2.2 μm, 3.0x 50 mm. The flow rate is 1.5 mL/min, the run time is 3.6 min, and the gradient profiles are 0.01 min 5% A, 2.20 min 100% A, 3.20 min 100% A, 3.30 min 5% A, 3.60 min 5% A.

[0240] 5.0 minute run: ACE UltraCore Super C18 column, 2.5 μm, 3.0x 50 mm. The flow rate is 1.5 mL/min, the run time is 5.0 min, and the gradient profiles are 0.01 min 10% A, 4.00 min 60% A, 4.70 min 60% A, 4.80 min 10% A, 5.00 min 10% A.

[0241] 5.6 minute run: Shim-pack VP-ODS column, 2.2 µm, 3.0x 50 mm. The flow rate is 1.5 mL/min, the run time is 5.6 min, and the gradient profiles are 0.01 min 5% A, 3.00 min 50% A, 5.00 min 50% A, 5.20 min 5% A, 5.60 min 5% A

[0242] Alternatively, analytical LC-MS data was collected on Agilent infinity 1260 LC, Agilent 6230 TOF mass spectrometer. The analysis is conducted on a Poroshell 120 EC C18 column (50mm x 3.0mm internal diameter 2.7µm packing diameter) at 45°C.

The solvents employed are:

A = 0.1% v/v solution of formic acid in water.

B = 0.1% v/v solution of formic acid in acetonitrile.

The gradient employed are as follows:

Table 1. Exemplary Column Gradients.

Time (minutes)	Flow Rate (mL/min)	% A	% B
0	1	95	5
0.5	1	95	5
3.0	1	1	99
4.0	1	1	99
4.1	1	95	5
4.5	1	95	5


[0243] The UV detection is an averaged signal from wavelength of 210nm to 350nm and mass spectra are recorded on a mass spectrometer using positive mode electrospray ionization.

[0244] Unless otherwise noted, all compounds were prepared with LC-MS purity >95%.


[0245] **Chemical Synthesis**

[0246] A PROTAC of ABM-L-ULM, or their pharmaceutically acceptable salts, polymorphic forms, prodrugs, solvate forms and isotope containing derivatives thereof, may be prepared by the general approaches described below (Schemes 1 and 2, and General Schemes 1-122, 1A-22A, and 1B-25B), together with synthetic methods known in the art of organic chemistry, or modifications and derivatizations that are familiar to those of ordinary skill in the art.


[0247] Scheme 1:




[0248] Scheme 2:




[0249] General Scheme 1:




[0250] General Scheme 2:




[0251] General Scheme 3:




[0252] General Scheme 4:




[0253] General Scheme 5:




[0254] General Scheme 6:




[0255] General Scheme 7:




[0256] General Scheme 8:




[0257] General Scheme 9:




[0258] General Scheme 10:




[0259] General Scheme 11:




[0260] General Scheme 12:




[0261] General Scheme 13:




[0262] General Scheme 14:




[0263] General Scheme 15:




[0264] General Scheme 16:




[0265] General Scheme 17:




[0266] General Scheme 18:




[0267] General Scheme 19:




[0268] General Scheme 20:




[0269] General Scheme 21:




[0270] General Scheme 22:




[0271] General Scheme 23:




[0272] General Scheme 24:




[0273] General Scheme 25:




[0274] General Scheme 26:




[0275] General Scheme 27:




[0276] General Scheme 28:




[0277] General Scheme 29:




[0278] General Scheme 30:




[0279] General Scheme 31:




[0280] General Scheme 32:




[0281] General Scheme 33:




[0282] General Scheme 34:




[0283] General Scheme 35:




[0284] General Scheme 36:




[0285] General Scheme 37:




[0286] General Scheme 38:




[0287] General Scheme 39:




[0288] General Scheme 40:




[0289] General Scheme 41:




[0290] General Scheme 42:




[0291] General Scheme 43:




[0292] General Scheme 44:




[0293] General Scheme 45:




[0294] General Scheme 46:




[0295] General Scheme 47:



[0296] General Scheme 48:



[0297] General Scheme 49:



[0298] To a solution of tert-butyl 4-(4-(hydroxymethyl)piperidin-1-yl)benzoate **3** (15 g, 51.4 mmol) in DCM (100 mL) was added Dess-Martin periodinane (26.1 g, 61.6 mmol) in ice-water bath and the clear solution was stirred at room temperature for 2 hours. The solvent was evaporated and EtOAc was added. The mixture was passed through a pad of silica gel, washed with EtOAc, and the solvent was evaporated to give crude. (It was combined with another batch of crude which started from 10 g starting material for purification.) The crude was dry loaded to ISCO (0-10% MeOH/DCM) for separation. Light yellow solid 16.11 g **4** (yield 65%) was obtained as product.


[0299] To a mixture of tert-butyl 4-(4-formylpiperidin-1-yl)benzoate **4** (15.79 g, 54.2 mmol) and 2-(2,6-dioxopiperidin-3-yl)-5-(piperazin-1-yl)isoindoline-1,3-dione dihydrochloride **5** (22.5 g, 54.2 mmol) in DCM (500 mL) was added sodium acetate (4.44 g, 54.2 mmol). The reaction mixture was allowed to stir for 3 mins and lowered into ice-water bath. $NaBH(OAc)_3$ (20.0 g, 94.8 mmol) was added over 10 mins and the bath was removed after an additional 10 mins. After 21h, water (500 mL) and DCM (150 mL) were added, the organic layer was removed and the aqueous layer was extracted with DCM (150ml), the combined organics were dried with Na_2SO_4, and evaporated the to give crude **6** which was dry-loaded with silica gel to ISCO (0-10% MeOH/DCM, 750 g column) for separation. The fractions were collected and dried, exchanged

with DCM twice, and dried under high vacuum for 26 hours to provide 18.15g **6** (yield 60%) as a yellow solid.


[0300] 4N HCl in dioxane (40 mL, 29.3 mmol) was added to tert-butyl 4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzoate **6** (18.15 g, 29.3 mmol) and the reaction mixture was heated to 50^0C for 7 hours. The solvent was evaporated and DCM exchanged twice, dried under high vacuum for overnight. It was exchanged with DCM again twice, dried under high vacuum for 8 hours to provide 18.77 g (yield 101%) **7** as a light brown solid.

[0301] To a solution of 4-((1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy)-2-chlorobenzonitrile hydrochloride **8** (8.9 g, 28.2 mmol) and 4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzoic acid dihydrochloride **7** (18.7 g, 29.6 mmol) in DCM (300 mL) was added Hunig's base (24.2 mL, 140 mmol), and HATU (11.0 g, 29.0 mmol) was added. The reaction mixture was allowed to stir at ambient temperature for 17h at which time it was washed with sat. $NaHCO_3$ (250 mL) twice, dried with Na_2SO_4, and evaporated to dryness. The crude was dry-loaded with silica gel to column. ISCO (0% 3CV, 0-10% MeOH/DCM 30 CV, 750 g column) separated the mixture. The product was dried in heated vacuum oven at 40 ^{0}C for 16 hours to give **9** 19.6 g (yield 85%, 99.55% pure by 18 mins analytical method) as a yellow solid.


[0302] General Scheme 50:




[0303] General Scheme 52:




[0304] General Scheme 53:




[0305] General Scheme 54:




[0306] General Scheme 55:




[0307] General Scheme 56:




[0308] General Scheme 57:




[0309] General Scheme 58:




[0310] General Scheme 59:




[0311] General Scheme 60:




[0312] General Scheme 61:




[0313] General Scheme 62:




[0314] General Scheme 63:




[0315] General Scheme 64:




[0316] General Scheme 65:




[0317] General Scheme 66:




[0318] General Scheme 67:




[0319] General Scheme 68:




[0320] General Scheme 69:




[0321] General Scheme 70:




[0322] General Scheme 71:




[0323] General Scheme 72:




[0324] General Scheme 73:




[0325] General Scheme 74:




[0326] General Scheme 75:




[0327] General Scheme 76:




[0328] General Scheme 77:




[0329] General Scheme 78:




[0330] General Scheme 79:




[0331] General Scheme 80:




[0332] General Scheme 81:




[0333] General Scheme 82:




[0334] General Scheme 83:




[0335] General Scheme 84:




[0336] General Scheme 85:




[0337] General Scheme 86:




[0338] General Scheme 87:




[0339] General Scheme 88:




[0340] General Scheme 89:




[0341] General Scheme 90:




[0342] General Scheme 91:




[0343] General Scheme 92:




[0344] General Scheme 93:




[0345] General Scheme 94:




[0346] General Scheme 95:




[0347] General Scheme 96:




[0348] General Scheme 97:




[0349] General Scheme 98:



[0350] General Scheme 99:




[0351] General Scheme 100:




Scheme 99


[0352] General Scheme 101:




[0353] General Scheme 102:




[0354] General Scheme 103:




[0355] General Scheme 104:




[0356] General Scheme 105:




[0357] General Scheme 106:




[0358] General Scheme 107:




[0359] General Scheme 108:




[0360] General Scheme 109:




[0361] General Scheme 110:




[0362] General Scheme 111:




[0363] General Scheme 112:




[0364] General Scheme 113:




[0365] General Scheme 114:




[0366] General Scheme 115:




[0367] General Scheme 116:




[0368] General Scheme 117:





[0369] General Scheme 118:




[0370] General Scheme 119:












[0371] General Scheme 120:



[0372] General Scheme 121:




[0373] General Scheme 122:




[0374] General Scheme 1A:

General Scheme 1




[0375] General Scheme 2A:

General Scheme 2




[0376] General Scheme 3A:


General Scheme 3




[0377] General Scheme 4A:




[0378] General Scheme 5A:




[0379] General Scheme 6A:




[0380] General Scheme 7A:




[0381] General Scheme 8A:




[0382] General Scheme 9A:




[0383] General Scheme 10A:




[0384] General Scheme 11A:




[0385] General Scheme 12A:




[0386] General Scheme 13A:




[0387] General Scheme 14A:




[0388] General Scheme 15A:




[0389] General Scheme 16A:




[0390] General Scheme 17A:




[0391] General Scheme 18A:




[0392] General Scheme 19A:




[0393] General Scheme 20A:




[0394] General Scheme 21A:




[0395] General Scheme 22A:




[0396] General Scheme 1B




[0397] General Scheme 2B




[0398] General Scheme 3B




[0399] General Scheme 4B




[0400] General Scheme 5B




[0401] General Scheme 6B




[0402] General Scheme 7B




[0403] General Scheme 8B




[0404] General Scheme 9B




[0405] General Scheme 10B




[0406] General Scheme 11B




[0407] General Scheme 12B




[0408] General Scheme 13B




[0409] General Scheme 14B




[0410] General Scheme 15B




[0411] General Scheme 16B




[0412] General Scheme 17B




[0413] General Scheme 18B




[0414] General Scheme 19B




[0415] General Scheme 20B




[0416] General Scheme 21B




[0417] General Scheme 22B




[0418] General Scheme 23B



[0419] General Scheme 24B



[0420] General Scheme 25B



[0421] **Synthetic Procedure for Compound 406:**

[0422] **Synthesis of tert-butyl N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]carbamate**

[0423] Into a 50.0-mL round-bottom flask, was placed tert-butyl N-[(1r,4r)-4-hydroxycyclohexyl]carbamate (500.0 mg, 2.32 mmol, 1.00 equiv), N,N-dimethylformamide (10.0

mL), sodium hydride (82.8 mg, 3.45 mmol, 1.50 equiv), 2-chloro-4-fluorobenzonitrile (432.6 mg, 2.78 mmol, 1.20 equiv). The resulting solution was stirred for 2 hours at 0°C in a water/ice bath. The reaction was then quenched by the addition of 20.0 mL of water. The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined. The resulting mixture was washed with sodium chloride (40.0 mL). The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/2). The collected fractions were combined and concentrated under vacuum. This resulted in 470.0 mg (58%) of tert-butyl N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]carbamate as yellow oil.

[0424] LC-MS (ES^+): m/z 295.0 [MH^+], t_R = 1.199 min, (1.90 minute run).

[0425] Chemical formula: $C_{18}H_{23}ClN_2O_3$[350.14].

[0426] **Synthesis of 4-(((1r,4r)-4-aminocyclohexyl)oxy)-2-chlorobenzonitrile**

[0427] Into a 50.0-mL round-bottom flask, was placed tert-butyl N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]carbamate (470.0 mg, 1.34 mmol, 1.00 equiv), methanol (5.0 mL), hydrogen chloride. The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was concentrated under vacuum. This resulted in 340.0 mg (88%) of 2-chloro-4-[[(1r,4r)-4-aminocyclohexyl]oxy]benzonitrile hydrochloride as a yellow solid.

[0428] LC-MS (ES^+): m/z 250.90 [MH^+], t_R = 0.537 min, (1.90 minute run).

[0429] Chemical formula: $C_{13}H_{15}ClN_2O$[250.09].

[0430] **Synthesis of 6-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy) cyclohexyl]pyridazine-3-carboxamide):** Into a 100-mL round-bottom flask, was placed 6-[4-(hydroxymethyl)piperidin-1-yl]pyridazine-3-carboxylic acid (1.0 g, 4.21 mmol, 1.00 equiv), 2-chloro-4-[(1r,4r)-4-aminocyclohexyl]oxybenzonitrile hydrochloride (1.2 g, 4.18 mmol, 1.00 equiv), N,N-dimethylformamide (30 mL), N,N,N',N'-Tetramethyl-O-(7-azabenzotriazol-1-yl)uronium hexafluorophospate (2.4 g, 6.31 mmol, 1.50 equiv), N,N-diisopropylethylamine (1.6 g, 12.38 mmol, 3.00 equiv). The resulting solution was stirred for 1 hour at room temperature. The reaction was then quenched by the addition of water (50 mL) and extracted with ethyl acetate (50 mL x 3). The combined organic layers was washed with brine (50 mL), dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (v:v = 12:1). This resulted in 1.1 g (56%) of 6-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide as yellow oil.

[0431] LC-MS (ES^+): *m/z* 470.0 [MH^+], t_R = 0.90 min (1.8 minute run).

[0432] **Synthesis of 6-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl] pyridazine-3-carboxamide:** Into a 100-mL round-bottom flask, was placed 6-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide (700.0 mg, 1.49 mmol, 1.00 equiv), dichloromethane (20 mL), (1,1,1-Triacetoxy)-1,1-dihydro-1,2-benziodoxol-3(1H)-one (947.2 mg, 2.23 mmol, 1.50 equiv). The resulting solution was stirred for 3 hours at room temperature. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (v: v = 1: 3). This resulted in 390.0 mg (56%) of 6-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide as a yellow solid.

[0433] LC-MS (ES^+): *m/z* 468.2 [MH^+], t_R =1.06 min (2.0 minute run).

[0434] **Synthesis of 6-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide:** Into a 100-mL round-bottom flask, was placed 6-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide (180.0 mg, 0.38 mmol, 1.00 equiv), dichloromethane (10 mL), 2-(2,6-dioxopiperidin-3-yl)-5-fluoro-6-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione hydrochloride (152.7 mg, 0.38 mmol, 1.00 equiv), sodium triacetoxyborohydride (244.6 mg, 3.00 equiv). The resulting solution was stirred for 3 hours at room temperature. The reaction was then quenched by water (30 mL), extracted with ethyl acetate (30 mL x 3), washed with brine (30 mL) and concentrated under reduced pressure. The solid was filtered out. The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column, 19*150mm 5 um; mobile phase, water (10 mmol/L ammonium bicarbonate) and acetonitrile (48.0% acetonitrile up to 73.0% in 8 min); Detector, UV 254nm. This resulted in 146.1 mg (47%) of 6-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide as a yellow solid.

[0435] ^1H NMR (400 MHz, DMSO): δ 11.11 (s, 1H), 8.58 (d, J = 8.2 Hz, 1H), 7.86 (d, J = 8.8 Hz, 1H), 7.81 (d, J = 9.5 Hz, 1H), 7.73 (d, J = 11.4 Hz, 1H), 7.46 (d, J = 7.4 Hz, 1H), 7.39 (d, J = 2.4 Hz, 1H), 7.34 (d, J = 9.7 Hz, 1H), 7.15-7.12 (m, 1H), 5.13-5.08 (m, 1H), 4.59-4.45 (m,

3H), 3.90-3.83 (m, 1H), 3.27 (s, 4H), 3.03 (m, 2H), 2.97-2.82 (m, 1H), 2.64-2.53 (m, 5H), 2.46 (m, 1H), 2.23 (m, 2H), 2.14-2.09 (m, 2H), 2.07-2.02 (m, 1H), 1.96-1.79 (m, 5H), 1.65 (m, 2H), 1.52 (m, 2H), 1.19-10.09 (m, 2H); LC-MS (ES^+): *m/z* 812.25 [MH^+], t_R = 1.57 min (3.0 minute run).

[0436] Chemical Formula: C41H43ClFN9O6 [811.30].

[0437] Total H count from HNMR data: 43.

[0438] **Synthetic Procedure for Compound 109:**

[0439] **1. Synthesis of benzyl 6-(6-hydroxyhex-1-yn-1-yl)pyridine-3-carboxylate)**

[0440] Into a 250-mL round-bottom flask, was placed a solution of benzyl 6-chloropyridine-3-carboxylate (2.8 g, 11.31 mmol, 1.00 equiv) in triethylamine (50 mL), Cuprous iodide (1.2 g, 6.30 mmol, 1.50 equiv), four (triphenylphosphine) palladium (2.5 g, 0.50 equiv), hex-5-yn-1-ol (1.3 g, 13.25 mmol, 1.00 equiv). The resulting solution was stirred for 3 hours at 90°C in an oil bath. The resulting solution was extracted with ethyl acetate and the organic layers combined. The resulting mixture was washed with brine. The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). The collected fractions were combined and concentrated under vacuum. This resulted in 1.1 g (31%) of benzyl 6-(6-hydroxyhex-1-yn-1-yl)pyridine-3-carboxylate as yellow oil. LC-MS (ES^+): *m/z* 309.95 [MH^+], t_R =0.873 min (1.90 minute run).

[0441] **2. Synthesis of6-(6-hydroxyhexyl)pyridine-3-carboxylic**

[0442] Into a 250-mL round-bottom flask, was placed a solution of benzyl 6-(6-hydroxyhex-1-yn-1-yl)pyridine-3-carboxylate (1.1 g, 3.56 mmol, 1.00 equiv) in methanol (40 mL), Palladium carbon (2 g, 10.00 equiv), hydrogen (g). The resulting solution was stirred for 48 hours at room temperature. The resulting solution was diluted with 200 mL of methanol. The resulting mixture was concentrated under vacuum. This resulted in 700 mg (88%) of 6-(6-hydroxyhexyl)pyridine-3-carboxylic acid as yellow oil.

[0443] LC-MS (ES^+): *m/z* 223.95 [MH^+], t_R =0.377 min (1.90 minute run).

[0444] **3. Synthesis of6-(6-hydroxyhexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide**

[0445] Into a 100-mL round-bottom flask, was placed a solution of 6-(6-hydroxyhexyl)pyridine-3-carboxylic acid (430 mg, 1.93 mmol, 1.00 equiv) in N,N-dimethylformamide (20 mL), N,N,N',N'-Tetramethyl-O-(7-azabenzotriazol-1-yl)uronium

hexafluorophospate (1.5 g, 3.94 mmol, 2.00 equiv), N,N-diisopropylethylamine (1 mL, 0.30 equiv), 2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile (690 mg, 2.48 mmol, 1.30 equiv). The resulting solution was stirred for 2 hours at room temperature. The reaction was then quenched by the addition of 50 mL of water. The resulting solution was extracted with ethyl acetate and the organic layers combined. The resulting mixture was washed with brine. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/hexane (1/3). The collected fractions were combined and concentrated under vacuum. This resulted in 580 mg (62%) of 6-(6-hydroxyhexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide as a yellow solid.

[0446] LC-MS (ES^+): *m/z* 484.10/486.10 [MH^+], t_R =0.927 min (1.90 minute run).

[0447] **4. Synthesis of6-(5-[[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoyl]pyridin-2-yl)hexyl 4-methylbenzene-1-sulfonate**

[0448] Into a 100-mL round-bottom flask, was placed a solution of 6-(6-hydroxyhexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide (580 mg, 1.20 mmol, 1.00 equiv) in dichloromethane (20 mL), 4-toluene sulfonyl chloride (750 mg, 3.93 mmol, 1.50 equiv), triethylamine (690 mg, 6.82 mmol, 3.00 equiv), 4-dimethylaminopyridine (35 mg, 0.29 mmol, 0.10 equiv). The resulting solution was stirred for 20 hours at room temperature. The reaction was then quenched by the addition of water. The resulting solution was extracted with ethyl acetate and the organic layers combined. The resulting mixture was washed with brine. The solid was dried in an oven under reduced pressure. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 430 mg (56%) of 6-(5-[[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoyl]pyridin-2-yl)hexyl 4-methylbenzene-1-sulfonate as a yellow solid.

[0449] LC-MS (ES^+): *m/z* 638.40/640.40[MH^+], t_R =1.336 min (2.00 minute run).

[0450] **5. Synthesis of6-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide**

[0451] Into a 25-mL round-bottom flask, was placed a solution of 6-(5-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoylpyridin-2-yl)hexyl 4-

methylbenzene-1-sulfonate (220 mg, 0.34 mmol, 1.00 equiv) in acetonitrile (5 mL), 2-(2,6-dioxopiperidin-3-yl)-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione; trifluoroacetic acid (162 mg, 0.35 mmol, 1.50 equiv), potassium carbonate (25 mg, 0.18 mmol, 3.00 equiv), sodium iodide (25 mg, 0.50 equiv). The resulting solution was stirred for 12 hours at 70°C in an oil bath. The resulting mixture was concentrated under vacuum. The crude product (3 mL) was purified by Prep-HPLC with the following conditions: Column, XBridge Shield RP18 OBD Column, 5um, 19*150mm; mobile phase, water (10 mmol/L ammonium bicarbonate) and acetonitrile (56.0% acetonitrile up to 65.0% in 8 min); Detector, UV 254nm. 41.3 mg product was obtained. This resulted in 41.3 mg (15%) of 6-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide as a yellow solid.

[0452] ^{1}HNMR (300 MHz, CD_3OD): δ 8.86 (s, 1H), 8.16-8.09 (m, 1H), 7.72-7.64 (m, 2H), 7.47-7.33 (m, 2H), 7.24-7.17 (m, 1H), 7.12 (s, 1H), 6.98-6.89 (m , 1H), 5.09-5.02 (m, 1H), 4.26 (s, 1H), 4.15(s, 1H), 3.48-3.42 (m, 4H), 2.92-2.49 (m, 9H), 2.46-2.35 (m, 2H), 2.16-2.01 (m, 1H), 1.81-1.71 (m, 2H),1.59-1.47 (m, 2H), 1.42-1.36 (m, 4H), 1.27-1.18 (m, 12H); LC-MS (ES^+): *m/z* 808.85/810.85 [MH^+], t_R =4.79 min (8.0 minute run).

[0453] Chemical formula: $C_{44}H_{50}ClN_7O_6$ [807.35/809.35].

[0454] Total H count from HNMR data: 48.

[0455] **Synthetic Procedure for Compound 158:**

[0456] **1. Synthesis of benzyl 4-fluorobenzoate:** Into a 500-mL round-bottom flask, was placed 4-fluorobenzoic acid (14 g, 99.9 mmol, 1.0 equiv), N,N-dimethylformamide (150.0 mL), (bromomethyl)benzene (18.7 g, 109.3 mmol, 1.1 equiv), dicesium carbonate (27.6 g, 84.7 mmol, 2.0 equiv). The resulting solution was stirred for 4 hours at room temperature. The reaction was then quenched by the addition of 150 mL of water. The resulting solution was extracted with ethyl acetate (100 mL x3) and the organic layers combined and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/6). This resulted in 18.0 g (78%) of benzyl 4-fluorobenzoate as a solid.

[0457] **2. Synthesis of benzyl 4-[4-(hydroxymethyl)piperidin-1-yl]benzoate :** Into a 100-mL round-bottom flask, was placed benzyl 4-fluorobenzoate (5.0 g, 21.7 mmol, 1.1 equiv), N,N-Dimethylformamide (30.0 mL), piperidin-4-ylmethanol (2.3 g, 19.9 mmol, 1.0 equiv), N,N-Diisopropylethylamine (7.6 g, 59.1 mmol, 3.0 equiv). The resulting solution was stirred for 2

hours at 100°C. The reaction was then quenched by the addition of 60.0 mL of water. The resulting solution was extracted with ethyl acetate (60 mL x 3) and the organic layers combined and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). This resulted in 3.6 g (55%) of benzyl 4-[4-(hydroxymethyl)piperidin-1-yl]benzoate as white oil.

[0458] LC-MS (ES^+): 326.30*m/z* [MH^+], t_R = 1.18min, (1.90 minute run).

[0459] **3. Synthesis of benzyl 4-(4-formylpiperidin-1-yl)benzoate:** Into a 100-mL round-bottom flask, was placed benzyl 4-[4-(hydroxymethyl)piperidin-1-yl]benzoate (500 mg, 1.54 mmol, 1.00 equiv), dichloromethane (15.0 mL), (1,1,1-Triacetoxy)-1,1-dihydro-1,2-benziodoxol-3(1H)-one (978 mg, 2.31 mmol, 1.5 equiv). The resulting solution was stirred for 4 hours at room temperature. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). This resulted in 420.0 mg (85%) of benzyl 4-(4-formylpiperidin-1-yl)benzoate as a yellow solid.

[0460] LC-MS (ES^+): 323.95*m/z* [MH^+], t_R = 1.04min, (1.80 minute run).

[0461] **4. Synthesis of benzyl 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]benzoate:** Into a 150-mL round-bottom flask, was placed benzyl 4-(4-formylpiperidin-1-yl)benzoate (500.0 mg, 1.5 mmol, 1.0 equiv), dichloromethane (20.0mL), 2-(2,6-dioxopiperidin-3-yl)-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione, trifluoroacetyl (747 mg, 1.7 mmol, 1.1 equiv), Sodium triacetoxyborohydride (1.39 g, 6.5 mmol, 4.0 equiv). The resulting solution was stirred for 4 hours at room temperature. The reaction was then quenched by the addition of water (100.0 mL). The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 500.0 mg (50%) of benzyl 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]benzoate as a yellow solid.

[0462] LC-MS (ES^+): 650.45*m/z* [MH^+], t_R = 1.37min, (1.90 minute run).

[0463] **5. Synthesis of 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]benzoic acid:** To a solution of benzyl4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]benzoate (500.0 mg, 0.8 mmol, 1.0 equiv) in 20.0 mL methyl alcohol

(30.0 mL) was added Pd/C (10%, 300 mg) under nitrogen atmosphere in a 100.0 mL round bottom flask. The flask was then vacuumed and flushed with hydrogen. The reaction mixture was hydrogenated at room temperature for 12 hours under hydrogen atmosphere using a hydrogen balloon, then filtered through a Celite pad and concentrated under reduced pressure. This resulted in 300.0 mg (69.0%) of 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]benzoic acid as a yellow solid.

[0464] LC-MS (ES^+): 560.35*m/z* $[MH^+]$, t_R = 0.74min, (1.90 minute run).

[0465] **6. Synthesis of 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl] benzamide:** Into a 100-mL round-bottom flask, was placed 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]benzoic acid (300.0 mg, 0.5 mmol, 1.0 equiv), N,N-dimethylformamide (15.0 mL), N,N,N',N'-Tetramethyl-O-(7-azabenzotriazol-1-yl)uronium hexafluorophospate (307.0 mg, 0.8 mmol, 1.5 equiv), 2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile hydrochloride (170.0 mg, 0.5 mmol, 1.0 equiv), N,N-Diisopropylethylamine (207.0 mg, 1.6 mmol, 3.0 equiv). The resulting solution was stirred for 1 hour at room temperature. The reaction was then quenched by the addition of water (80 mL) . The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and concentrated under vacuum. The crude product (5 mL) was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column, 5um,19 *150 mm; mobile phase, Water(10MMOL/L ammonium bicarbonate) and acetonitrile (59.0% acetonitrile up to 80.0% in 8 min); Detector, UV 220nm. 4 mL product was obtained. This resulted in 114 g (25921%) of 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as a yellow solid.

[0466] 1H NMR (300 MHz, DMSO-d6) δ 11.05 (s, 1H), 7.86 (d, J = 8.7 Hz, 1H), 7.67 (dd, J = 17.8, 8.4 Hz, 3H), 7.44 (d, J = 9.2 Hz, 1H), 7.34-7.13 (m, 3H), 7.02-6.87 (m, 3H), 5.04 (dd, J = 12.6, 5.3 Hz, 1H), 4.28 (s, 1H), 4.02 (d, J = 9.0 Hz, 1H), 3.82 (d, J = 12.5 Hz, 2H), 3.58-3.30 (m, 5H), 2.90-2.66 (m, 3H), 2.6-2.47 (m, 3H), 2.50-2.42 (m,4H), 2.17 (d, J = 6.5 Hz, 2H), 1.98 (dt, J = 10.0, 4.5 Hz, 1H), 1.77 (d, J = 11.9 Hz, 3H), 1.18 (s, 6H), 1.09 (s, 6H).

[0467] LC-MS (ES^+): 820.60*m/z* $[MH^+]$, t_R = 2.99min, (4.8 minute run).

[0468] **Synthetic Procedure for Compound 357:**

[0469] **1. Synthesis of tert-butyl 6-(2-ethoxy-2-oxoethylidene)-2-azaspiro[3.3]heptane-2-carboxylate)**

[0470] A mixture of t-BuOK (3.98 g, 35.47 mmol, 2.498 equiv) and ethyl 2-(bromotriphenyl-$l^[5]-phosphanyl)acetate (15.2 g, 35.41 mmol, 2.493 equiv) in THF (90 mL) in a 250-mL round-bottom flask was stirred for 1 hour at room temperature. Then tert-butyl 6-oxo-2-azaspiro[3.3]heptane-2-carboxylate (3 g, 14.20 mmol, 1 equiv) was added and the reaction mixture was stirred for an additional 6 hours at 35℃. Then filtered through Celite and filter cake was washed with THF. The crude was subjected to a silica gel column with ethyl acetate/petroleum ether (1:5). This resulted in 3.3 g (82.60%) of tert-butyl 6-(2-ethoxy-2-oxoethylidene)-2-azaspiro[3.3]heptane-2-carboxylate as a colorless oil.

[0471] LC-MS (ES^+): *m/z* 226.1 [M-(t-Bu)+H^+], t_R = 1.267 min, (2.0 minute run).

[0472] **2. Synthesis of tert-butyl 6-(2-ethoxy-2-oxoethyl)-2-azaspiro[3.3]heptane-2-carboxylate**

[0473] To a solution of tert-butyl 6-(2-ethoxy-2-oxoethylidene)-2-azaspiro[3.3]heptane-2-carboxylate (3.296 g, 11.72 mmol, 1 equiv) in ethanol (330 mL) in a 500-mL 3-necked round-bottom flask was added Pd/C (661.8 mg) under N_2 flow. The reaction flask was vaccumed and flushed with H_2 for three times. The resulting mixture was stirred for 3 hours at room temperature under H_2. Then the mixture was filtered through a celite pad. The collected filtrate was concentrated under reduced pressure and dried under vaccum. This resulted in 3.12 g (93.99%) of tert-butyl 6-(2-ethoxy-2-oxoethyl)-2-azaspiro[3.3]heptane-2-carboxylate as a colorless oil.

[0474] LC-MS (ES^+): *m/z* 228.1 [M-(t-Bu)+H^+], t_R = 1.254 min, (2.0 minute run).

[0475] **3. Synthesis of tert-butyl 6-(2-hydroxyethyl)-2-azaspiro[3.3]heptane-2-carboxylate**

[0476] Into a 250-mL round-bottom flask, was placed tert-butyl 6-(2-ethoxy-2-oxoethyl)-2-azaspiro[3.3]heptane-2-carboxylate (3.08 g, 10.87 mmol, 1 equiv), THF (120 mL), DIBAL-H in THF (1 M) (32.6 mL, 229.28 mmol, 3 equiv). The resulting solution was stirred for 3 hours at room temperature. The reaction was then quenched by the addition of water (60 mL x 1). The resulting solution was extracted with ethyl acetate (120 mL x 2). The resulting mixture was washed with brine (100 mL x 1). The organic layer was dried over anhydrous sodium sulfate,

concentrated and dried under vaccum. This resulted in 2.53 g (96.45%) of tert-butyl 6-(2-hydroxyethyl)-2-azaspiro[3.3]heptane-2-carboxylate as a white solid.

[0477] LC-MS (ES^+): *m/z* 186.1 [M-(t-Bu)+H^+], t_R = 1.061 min, (2.0 minute run).

[0478] **4. Synthesis of 2-[2-azaspiro[3.3]heptan-6-yl]ethan-1-ol; trifluoroacetic acid**

[0479] Into a 500-mL round-bottom flask, was placed a solution of tert-butyl 6-(2-hydroxyethyl)-2-azaspiro[3.3]heptane-2-carboxylate (2.52 g, 10.44 mmol, 1 equiv) in DCM (120 mL), trifluoroacetic acid (7.7 mL). The resulting solution was stirred for 4 hours at room temperature. The resulting mixture was evaporated to dryness and dried under vaccum. This resulted in 2.66 g (99.80%) of 2-[2-azaspiro[3.3]heptan-6-yl]ethan-1-ol; trifluoroacetic acid as light yellow oil.

[0480] **5. Synthesis of tert-butyl 4-[6-(2-hydroxyethyl)-2-azaspiro[3.3]heptan-2-yl]benzoate**

[0481] To a solution of tert-butyl 4-fluorobenzoate (10.25 g, 52.24 mmol, 5.012 equiv) in DMSO (50 mL) in a 250-mL round-bottom flask was added DIEA (14 mL, 84.71 mmol, 8.128 equiv). The mixture was stirred for 5 minutes, then 2-[2-azaspiro[3.3]heptan-6-yl]ethan-1-ol; trifluoroacetic acid (2.66 g, 10.42 mmol, 1 equiv) was added. The resulting solution was stirred for 16 hours at 130 ℃. After cooling to room temperature, the reaction was then quenched by addition of water (100 mL x 1). The resulting solution was extracted with ethyl acetate (100 mL x 3). The combined organic layers were washed with water (50 mL x 2) and brine (50 mL x 1). The organic layer was dried over anhydrous sodium sulfate. The crude was subjected to a silica gel column with ethyl acetate/petroleum ether (1:2). This resulted in 311.5 mg (9.42%) of tert-butyl 4-[6-(2-hydroxyethyl)-2-azaspiro[3.3]heptan-2-yl]benzoate as a light yellow solid.

[0482] LC-MS (ES^+): *m/z* 318.2 [MH^+], t_R = 1.314 min, (2.0 minute run).

[0483] **6. Synthesis of tert-butyl 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]benzoate**

[0484] To a solution of PPh3 (361.7 mg, 1.38 mmol, 1.506 equiv) in THF (15 mL) was added DIAD (280.2 mg, 1.39 mmol, 1.513 equiv) dropwise with stirring at room temperature under N_2. To this was added tert-butyl 4-[6-(2-hydroxyethyl)-2-azaspiro[3.3]heptan-2-yl]benzoate (290.7 mg, 0.92 mmol, 1 equiv), 2-(2,6-dioxopiperidin-3-yl)-5-hydroxy-2,3-dihydro-1H-isoindole-1,3-dione (377.4 mg, 1.38 mmol, 1.503 equiv). The resulting solution was stirred for 4 hours at room temperature. The reaction was then quenched by the addition of water (20 mL).

The resulting solution was extracted with ethyl acetate (30 mL x 3) and the combined organic layers were washed with brine (20 mL x 1). The organic layer was dried over anhydrous sodium sulfate. The crude was subjected to a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 90.8 mg (17.28%) of tert-butyl 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]benzoate as a light brown solid.

[0485] LC-MS (ES^+): *m/z* 574.05 [MH^+], t_R = 1.443 min, (2.0 minute run).

[0486] **7. Synthesis of 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]benzoic acid**

[0487] To a solution of tert-butyl 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]benzoate (70.2 mg, 0.12 mmol, 1 equiv) in DCM (2.0 mL) was added TFA (0.6 mL, 8.08 mmol, 66.008 equiv). The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was evaporated to dryness and dried under vaccum. This resulted in 63.3 mg (99.95%) of 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]benzoic acid as a light brown solid.

[0488] LC-MS (ES^+): *m/z* 518.2 [MH^+], t_R = 1.166 min, (2.0 minute run).

[0489] **8. Synthesis of 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0490] To a solution of 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]benzoic acid (63.3 mg, 0.12 mmol, 1 equiv) in DMF (3.4 mL) was added DIEA (0.32 mL, 1.94 mmol, 15.830 equiv), then HATU (51.5 mg, 0.14 mmol, 1.107 equiv), finally 2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile hydrochloride (43.4 mg, 0.14 mmol, 1.126 equiv). The resulting solution was stirred for 3 hours at room temperature. The reaction was then quenched by the addition of water (10 mL). The resulting solution was extracted with dichloromethane (20 mL x 3); the combined organic layers were washed with water (10 mL x 2) and brine (10 mL x 1). The organic layer was dried over anhydrous sodium sulfate. The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep OBD C18 Column, 19*250mm,5um; mobile phase, Water(10mmol/L NH_4HCO_3) and acetonitrile (20% Phase B up to

60% in 8 min); Detector, UV254/220. This resulted in 57.4 mg (60.30%) of 4-[6-(2-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]ethyl)-2-azaspiro[3.3]heptan-2-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as a white solid.

[0491] ^{1}H NMR (300 MHz, d6-DMSO) δ 1.13 (s, 6H), 1.22 (s, 6H), 1.88-2.01 (m, 5H), 2.29 (d, J = 5.7 Hz, 3H), 2.44-2.56 (m, 2H), 2.79-2.84 (m, 1H), 3.73 (s, 2H), 3.84 (s, 2H), 3.98 (d, J = 9.0 Hz, 1H), 4.07 (t, J = 6.2 Hz, 2H), 4.25 (s, 1H), 5.02-5.08 (m, 1H), 6.34 (d, J = 8.4 Hz, 2H), 6.93 (dd, J1 = 2.1 Hz, J2 = 8.7 Hz, 1H), 7.13 (d, J = 2.4 Hz, 1H), 7.26-7.29 (m, 1H), 7.35-7.39 (m, 2H), 7.63-7.66 (m, 2H), 7.75-7.85 (m, 2H), 11.05 (s, 1H); LC-MS (ES$^+$): *m/z* 778.30 [MH$^+$], t_R = 3.222 min, (4.60 minute run) ; LC-MS (ES$^+$): *m/z* 778.30/780.30 [MH$^+$], t_R = 3.222 min, (4.60 minute run).

[0492] Chemical formula: $C_{43}H_{44}ClN_5O_7$ [777.29/779.29].

[0493] Total H count from HNMR data: 44.

[0494] **Synthetic Procedure for Compound 579:**

[0495] **Synthesis of tert-butyl N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]carbamate**

[0496] Into a 50.0-mL round-bottom flask, was placed tert-butyl N-[(1r,4r)-4-hydroxycyclohexyl]carbamate (500.0 mg, 2.32 mmol, 1.00 equiv), N,N-dimethylformamide (10.0 mL), sodium hydride (82.8 mg, 3.45 mmol, 1.50 equiv), 2-chloro-4-fluorobenzonitrile (432.6 mg, 2.78 mmol, 1.20 equiv). The resulting solution was stirred for 2 hours at 0℃ in a water/ice bath. The reaction was then quenched by the addition of 20.0 mL of water. The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined. The resulting mixture was washed with sodium chloride (40.0 mL). The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/2). The collected fractions were combined and concentrated under vacuum. This resulted in 470.0 mg (58%) of tert-butyl N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]carbamate as yellow oil.

[0497] LC-MS (ES$^+$): m/z 295.0 [MH$^+$], t_R = 1.199 min, (1.90 minute run).

[0498] Chemical formula: $C_{18}H_{23}ClN_2O_3$[350.14]

[0499] **2. Synthesis of 4-(((1r,4r)-4-aminocyclohexyl)oxy)-2-chlorobenzonitrile hydrochloride:**

[0500] Into a 50.0-mL round-bottom flask, was placed tert-butyl N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]carbamate (470.0 mg, 1.34 mmol, 1.00 equiv), methanol (5.0 mL), hydrogen chloride. The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was concentrated under vacuum. This resulted in 340.0 mg (88%) of 4-(((1r,4r)-4-aminocyclohexyl)oxy)-2-chlorobenzonitrile hydrochloride as a yellow solid.

[0501] LC-MS (ES^+): m/z 250.90 [MH^+], t_R = 0.537 min, (1.90 minute run).

[0502] Chemical formula: $C_{13}H_{15}ClN_2O$[250.09].

[0503] **3. Synthesis of methyl 5-(4-(hydroxymethyl)piperidin-1-yl)pyrazine-2-carboxylate**

[0504] Into a 50-mL round-bottom flask, was placed methyl 5-chloropyrazine-2-carboxylate (2 g, 11.59 mmol, 1 equiv), DMSO (15 mL, 0.19 mmol, 0.017 equiv), DIEA (0.2 mL, 0.000 equiv), piperidin-4-ylmethanol (1.3 mg, 0.01 mmol, 1 equiv). The resulting solution was stirred for 16 hours at 120℃ in an oil bath. The resulting solution was extracted ethyl acetate (30 mL x 3) and the organic layers combined.. The resulting mixture was washed with brine (10 mL x 1). The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (1:1). This resulted in 2.21 g (65%) of methyl 5-[4-(hydroxymethyl)piperidin-1-yl]pyrazine-2-carboxylate as a white solid.

[0505] LC-MS (ES+): m/z 251.13 [MH+], t_R = 0.61min, (1.9 minute run).

[0506] **4. Synthesis of 5-[4-(hydroxymethyl)piperidin-1-yl]pyrazine-2-carboxylic acid**

[0507] Into a 100-mL round-bottom flask, was placed methyl 5-[4-(hydroxymethyl)piperidin-1-yl]pyrazine-2-carboxylate (2.21 g, 8.79 mmol, 1 equiv), methanol (40 mL), lithiumol (0.633 mg, 0.03 mmol, 0.003 equiv), water (10 mL). The resulting solution was stirred for 16 hours at room temperature. The resulting mixture was concentrated. The PH was adjusted to 4 with 1M HCl. The resulting solution was extracted dichloromethane (30 mL x 3) and the organic layers combined. The resulting mixture was washed with brine (10 mL x 1). The resulting mixture was concentrated under vacuum. This resulted in 1.7042 g (81.67%) of 5-[4-(hydroxymethyl)piperidin-1-yl]pyrazine-2-carboxylic acid as a white solid.

[0508] LC-MS (ES+): m/z 237.11 [MH+], t_R = 0.75min, (2.0 minute run).

[0509] **5. Synthesis of 5-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1s,4s)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazine-2-carboxamide**

[0510] Into a 100-mL round-bottom flask, was placed 5-[4-(hydroxymethyl)piperidin-1-yl]pyrazine-2-carboxylic acid (310 mg), DMF (15 mL), DIEA (563.69 mg), 2-chloro-4-[[(1s,4s)-4-aminocyclohexyl]oxy]benzonitrile (250 mg), BOP (386.28 mg). The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was extracted with ethyl acetate (50 mL x 3) and the organic layer was washed with brine (30 mL x 1). The organic was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (10:1). The collected fractions were combined and concentrated under vacuum. This resulted in 243.5 mg of 5-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1s,4s)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazine-2-carboxamide as a white solid.

[0511] LC-MS (ES+): m/z 469.19 [MH+], t_R = 0.99min, (1.9 minute run).

[0512] **6. Synthesis of 5-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazine-2-carboxamide**

[0513] Into a 100-mL round-bottom flask, was placed 5-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazine-2-carboxamide (150.1 mg, 0.32 mmol, 1 equiv), dichloromethane (15 mL, 0.18 mmol, 0.553 equiv), Dess-martin (271.37 mg). The resulting mixture was extracted with dichloromethane (50 mL x 3) and the organic layer was washed with brine (30 mL x 1). The organic was dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 200 mg (crude) of 5-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazine-2-carboxamide as a white solid.

[0514] LC-MS (ES+): m/z 467.17 [MH+], t_R = 1.03min, (1.9 minute run).

[0515] **7. Synthesis of 5-(4-[[(propan-2-yl)[(1r,3r)-3-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]cyclobutyl]amino]methyl]piperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazin**

[0516] Into a 100-mL round-bottom flask, was placed 5-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazine-2-carboxamide (220 mg, 0.47 mmol, 1 equiv), dichloromethane (20 mL, 0.24 mmol, 0.501 equiv), Titanium(IV)isopropoxide (2 mL, 0.01 mmol, 0.015 equiv), 2-(2,6-dioxopiperidin-3-yl)-5-[(1r,3r)-3-[(propan-2-yl)amino]cyclobutoxy]-2,3-dihydro-1H-isoindole-1,3-dione (420 mg, 1.09 mmol, 2.318 equiv), $NaH(OAc)_3$ (340 mg). The resulting solution was stirred for 16 hours at 60℃ in an oil bath. The resulting mixture was extracted with dichloromethane (50 mL x 3) and the organic layer was washed with brine (30 mL x 1). The crude product was purified by Prep-HPLC with the

following conditions: Column, XSelect CSH Prep C18 OBD Column, 5um, 19 x 150mm; mobile phase, water(10mmol/L NH_4HCO_3) and acetonitrile (30% Phase B up to 55% in 8.5 min); Detector: UV254/220. This resulted in 26 mg (6.60%) of 5-(4-[[(propan-2-yl)[(1r,3r)-3-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]cyclobutyl]amino]methyl]piperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyrazin as a light yellow solid.


[0517] ^{1}H NMR (300 MHz, DMSO-*d6*) 11.09 (s,1H), 8.56 (s, 1H), 8.23 (s, 1H), 8.12-8.05 (m, 1H), 7.92-7.82 (m, 2H), 7.40-7.36 (s, 1H), 7.26-7.20 (m, 2H), 7.12-7.09 (m, 1H), 5.12-5.10 (m, 1H), 4.90-4.81 (s, 1H), 4.51-4.48 (m, 3H), 3.91-3.80 (s,1H), 3.72-3.62 (m, 1H), 3.00-2.81 (m, 4H), 2.79-2.68 (m, 1H), 2.50-2.42 (m, 3H), 2.30-2.00 (m, 7H), 1.99-1.81 (m, 4H), 1.80-1.36 (m, 5H), 1.21-1.01(m, 2H), 0.91-0.71 (m, 6H); LC-MS (ES^+): *m/z* 837.30/839.30 [MH^+], t_R = 2.934 min, (4.80 minute run).

[0518] Chemical formula: $C_{44}H_{49}ClN_6O_7$ [836.34/838.34].

[0519] Total H count from HNMR data: 49.

[0520] **Synthetic Procedure for Compound 278:**

[0521] Step 1: [(R)-tert-butyl (1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)carbamate].

[0522] 


[0523] A mixture of (R)-2-((tert-butoxycarbonyl)amino)propyl 4-methylbenzenesulfonate (12.6 g, crude), 2-chloro-4-(1H-pyrazol-3-yl)benzonitrile (7.1 g, 34.9 mmol) and cesium carbonate (5.8 g, 48.8 mmol) in acetonitrile (100 ml) was stirred at 70°C for 2 hours. TLC showed the reaction was complete. The reaction mixture was partitioned between water (50 ml) and ethyl acetate (100 ml). The organic layer was collected and the aqueous layer was extracted with ethyl acetate (50 ml x 2). The combined organic layers were washed with brine (100 ml), dried over magnesium sulfate and evaporated under reduced pressure to give a crude residue, which was purified by silica gel flash column chromatography (eluted with 50%-100% ethyl acetate in hexane) to afford tert-butyl 4-(prop-2-yn-1-yl)piperazine-1-carboxylate (7.0 g, yield 58%) as white solid.

[0524] ^{1}H NMR (400 MHz, $CDCl_3$): δ 1.16 (d, J = 6.8 Hz, 3H), 1.42 (s, 9H), 4.02-4.11 (m, 1H), 4.16-4.30 (m, 2H), 6.61 (d, J = 2.4 Hz, 1H), 7.46 (d, J = 2.4 Hz, 1H), 7.67 (d, J = 8.0 Hz, 1H), 7.74-7.80 (m, 1H), 7.96 (s, 1H).

[0525] Chemical Formula: $C_{18}H_{21}ClN_4O_2$; Molecular Weight: 360.84.


[0526] Total H count from HNMR data: 20.

[0527] Step 2: [(R)-4-(1-(2-aminopropyl)-1H-pyrazol-3-yl)-2-chlorobenzonitrile hydrochloride].

[0528] 

[0529] A solution of (R)-tert-butyl (1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)carbamate (7.0 g, 19.4 mmol) in hydrogen chloride in dioxane (4M, 15 ml) was stirred at room temperature for 2 hours. The volatiles were removed under reduced pressure to give a crude residue, which was triturated with dichloromethane (25 ml). The resulting solid was collected by filtration and dried under vacuum to afford (R)-4-(1-(2-aminopropyl)-1H-pyrazol-3-yl)-2-chlorobenzonitrile hydrochloride (4.3 g, yield 85%) as white solid which was used in next step without further purificaiton.

[0530] Step 3: [tert-butyl 4-(prop-2-yn-1-yl)piperazine-1-carboxylate].

[0531] 


[0532] A mixture of tert-butyl piperazine-1-carboxylate (10 g, 53.7 mmol), N-ethyl-N-isopropylpropan-2-amine (12.6 g, 97.6 mmol) and 3-bromoprop-1-yne (5.8 g, 48.8 mmol) in acetonitrile (50 ml) was stirred for 2 hours. TLC showed the reaction was complete. The reaction mixture was partitioned between ethyl acetate (80 ml) and water (80 ml). The organic layer was collected, and the aqueous layer was extracted with ethyl acetate (45 ml x 2). The combined organic layers were washed with brine (100 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash column chromatography (eluted with 50% ethyl acetate in hexane) to afford tert-butyl 4-(prop-2-yn-1-yl)piperazine-1-carboxylate (10.5 g, yield 96%) as pale yellow oil.

[0533] ^{1}H NMR (400 MHz, $CDCl_3$): δ 1.46 (s, 9H), 2.26 (t, J = 2.4 Hz, 1H), 2.51 (t, J = 4.8 Hz, 4H), 3.32 (d, J = 2.4 Hz, 2H), 3.47 (t, J = 4.8 Hz, 4H).

[0534] Chemical Formula: $C_{12}H_{20}N_2O_2$; Molecular Weight: 224.30.

[0535] Total H count from HNMR data: 20.


[0536] Step 2: [tert-butyl 4-((3-(ethoxycarbonyl)-1H-pyrazol-5-yl)methyl)piperazine-1-carboxylate].

[0537] 

[0538] A mixture of tert-butyl 4-(prop-2-yn-1-yl)piperazine-1-carboxylate (3.0 g, 13.4 mmol), ethyl diazoacetate (2.3 g, 20.1 mmol) and zinc trifluoromethanesulfonate (974 mg, 2.68 mmol) in triethylamine (9.3 ml, 67 mmol) was stirred at 100°C overnight. TLC showed the reaction was complete. The reaction mixture was allowed to cool to room temperature and partitioned between ethyl acetate (50 ml) and water (20 ml). The organic layer was collected, and the aqueous layer was extracted with ethyl acetate (30 ml x 2). The combined organic layers were washed with brine (30 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to afford a crude residue, which was purified by silica gel flash chromatography (eluted with 100%-200% ethyl acetate in hexane) to afford tert-butyl 4-((3-(ethoxycarbonyl)-1H-pyrazol-5-yl)methyl)piperazine-1-carboxylate (2.1 g, yield 47%) as yellow oil.

[0539] LC_MS: (ES$^+$): m/z 339.5 $[M+H]^+$. t_R = 1.772 min.

[0540] Step 3: [5-((4-(tert-butoxycarbonyl)piperazin-1-yl)methyl)-1H-pyrazole-3-carboxylic acid].


[0541] 

[0542] A mixture of tert-butyl 4-((3-(ethoxycarbonyl)-1H-pyrazol-5-yl)methyl)piperazine-1-carboxylate (500 mg, 1.48 mmol) and lithium hydroxide monohydrate (250 mg, 5.91 mmol) in tetrahydrofuran (10 ml)-water (2.5 ml)-methanol (2.5 ml) was stirred at 45°C overnight. TLC showed the reaction was complete. The reaction mixture was acidified with diluted hydrochloride acid (1N) till pH 6-7, the resulting mixture was concentrated under reduced pressure to afford 5-((4-(tert-butoxycarbonyl)piperazin-1-yl)methyl)-1H-pyrazole-3-carboxylic acid (1.1 g, crude) as yellow solid.

[0543] LC_MS: (ES$^+$): m/z 311.3 $[M+H]^+$. t_R = 1.378 min.

[0544] Step 4: [(R)-tert-butyl 4-((3-((1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)carbamoyl)-1H-pyrazol-5-yl)methyl)piperazine-1-carboxylate].

[0545]



[0546] A mixture of 5-((4-(tert-butoxycarbonyl)piperazin-1-yl)methyl)-1H-pyrazole-3-carboxylic acid (1.1 g, crude), (R)-4-(1-(2-aminopropyl)-1H-pyrazol-3-yl)-2-chlorobenzonitrile hydrochloride (336 mg, 1.29 mmol), N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride (371 mg, 1.94 mmol), 1-hydroxybenzotriazole (262 mg, 1.94 mmol) and N-ethyl-N-isopropylpropan-2-amine (835 mg, 6.5 mmol) in anhydrous N,N-dimethylformamide (8 ml) was stirred at room temperature for 2 hours, and then 2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (980 mg, 2.58 mmol) was added. The resulting mixture was stirred at room temperature for 1 hour. TLC showed the reaction was complete. The mixture was partitioned between ethyl acetate (20 ml) and water (10 ml). The organic layer was collected, washed with brine (20 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluted with 100%-200% ethyl acetate in hexane) to afford (R)-tert-butyl 4-((3-((1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)carbamoyl)-1H-pyrazol-5-yl)methyl)piperazine-1-carboxylate (300 mg, yield 37% over 2 steps) as white solid.


[0547] ^{1}H NMR (400 MHz, $CDCl_3$): δ 1.21 (d, J = 6.8 Hz, 3H), 1.46 (s, 9H), 2.42 (t, J = 4.8 Hz, 4H), 3.44 (t, J = 4.8 Hz, 4H), 3.62 (s, 2H), 4.24-4.29 (m, 1H), 4.40-4.46 (m, 1H), 6.63 (d, J = 2.4 Hz, 1H), 6.68 (s, 1H), 7.50 (d, J = 2.0 Hz, 1H), 7.68 (d, J = 8.0 Hz, 1H), 7.74-7.78 (m, 1H), 7.97 (d, J = 8.0 Hz, 1H), 8.27 (d, J = 1.2 Hz, 1H).

[0548] Chemical Formula: $C_{27}H_{33}ClN_8O_3$; Molecular Weight: 553.06.

[0549] Total H count from HNMR data: 31.

[0550] LC_MS: (ES$^+$): m/z 553.3 $[M+H]^+$. t_R = 2.181 min.

[0551] Step 5:[(R)-N-(1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-(piperazin-1-ylmethyl)-1H-pyrazole-3-carboxamide dihydrochloride].



[0552]

[0553] A solution of tert-butyl 4-(2-(2-(4-(5-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)pyridin-2-yl)piperazin-1-yl)ethoxy)ethyl)piperazine-1-carboxylate (160 mg, 0.29 mmol) in hydrogen chloride in dioxane (4M, 5 ml) was stirred at room temperature for 2 hours. The volatiles were removed under reduced pressure to give a crude residue which was triturated with dichloromethane (25 ml). The resulting solid was collected by filtration and dried under vacuum to afford (R)-N-(1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-(piperazin-1-ylmethyl)-1H-pyrazole-3-carboxamide dihydrochloride (130 mg, yield 86%) as white solid.


[0554] ^{1}H NMR (400 MHz, CD_3OD): δ 1.31 (d, J = 6.8 Hz, 3H), 3.50-3.62 (m, 8H), 4.30-4.48 (m, 4H), 4.58-4.64 (m, 1H), 6.78 (d, J = 2.0 Hz, 1H), 7.00-7.03 (m, 1H), 7.73 (d, J = 2.0 Hz, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.90 (d, J = 8.0 Hz, 1H), 8.03 (s, 1H).

[0555] Chemical Formula: $C_{22}H_{27}Cl_3N_8O$; Molecular Weight: 525.86.

[0556] Total H count from HNMR data: 23.

[0557] LC_MS: (ES^+): m/z 453.3 $[M+H]^+$. t_R = 1.916 min.

[0558] Step 6: [4-chlorobutyl 4-methylbenzenesulfonate].

[0559] 

[0560] A mixture of 4-chlorobutan-1-ol (5 g, 46.05 mmol), triethylamine (9. 3 g, 92.1 mmol), N,N-dimethylpyridin-4-amine (562 mg, 4.61 mmol) and 4-methylbenzene-1-sulfonyl chloride (9.63 g, 50.66 mmol) in dichloromethane (40 ml) was stirred at room temperature for 1.5 hours. TLC showed the reaction was complete. The reaction mixture was partitioned between water (50 ml) and ethyl acetate (100 ml). The organic layer was collected, and the aqueous layer was extracted with ethyl acetate (100 ml). The combined organic layers were washed with brine (100 ml), dried over sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash column chromatography (eluted with 20% ethyl acetate in hexane) to afford 4-chlorobutyl 4-methylbenzenesulfonate (12.0 g, yield 99%) as colorless oil.


[0561] ^{1}H NMR (400 MHz, $CDCl_3$): δ 1.80-1.84 (m, 4H), 2.46 (s, 3H), 3.49-3.53 (m, 2H), 4.04-4.08 (m, 2H), 7.35 (d, *J* = 8.0 Hz, 2H), 7.79 (d, *J* = 8.0 Hz, 2H).

[0562] Chemical Formula: $C_{11}H_{15}ClO_3S$; Molecular Weight: 262.75.

[0563] Total H count from HNMR data: 15.

[0564] LC_MS: (ES^+): m/z 263.1 $[M+H]^+$. t_R = 2.888 min.


[0565] Step 7: [5-(4-chlorobutoxy)-2-(2,6-dioxopiperidin-3-yl)isoindoline-1,3-dione]

[0566] 

[0567] A mixture of 4-chlorobutyl 4-methylbenzenesulfonate (0.96 g, 3.65 mmol), 2-(2,6-dioxopiperidin-3-yl)-5-hydroxyisoindoline-1,3-dione (1 g, 3.65 mmol) and potassium carbonate (0.76 g, 5.47 mmol) in N,N-dimethylformamide (10 ml) was stirred at 70°C overnight. TLC showed the reaction was complete. The reaction mixture was partitioned between ethyl acetate (30 ml) and water (20 ml). The organic layer was collected, and the aqueous layer was extracted with ethyl acetate (30 ml x 2). The combined organic layers were washed with brine (30 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to afford a crude residue which was purified by silica gel flash chromatography (eluted with 33%-50% ethyl acetate in hexane) to afford 5-(4-chlorobutoxy)-2-(2,6-dioxopiperidin-3-yl)isoindoline-1,3-dione (220 mg, yield 18%) as white solid.

[0568] LC_MS: (ES^+): m/z 365.1 $[M+H]^+$. t_R = 2.547 min.

[0569] Step 8: [N-((R)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-((4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)methyl)-1H-pyrazole-3-carboxamide].

[0570] 

A mixture of (R)-N-(1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-(piperazin-1-ylmethyl)-1H-pyrazole-3-carboxamide dihydrochloride (130 mg, 0.25 mmol), 5-(4-chlorobutoxy)-2-(2,6-dioxopiperidin-3-yl)isoindoline-1,3-dione (90 mg, 0.25 mmol), and N-ethyl-

N-isopropylpropan-2-amine (162 mg, 1.25 mmol) and potassium iodide (124 mg, 0.75 mmol) in acetonitrile (3 ml) was stirred at 100°C in sealed tube overnight. TLC showed the reaction was complete. The mixture was partitioned between ethyl acetate (30 ml) and water (20 ml). The organic layer was collected, washed with brine (30 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by preparative TLC (eluted with 8% methanol in dichloromethane) to afford N-((R)-1-(3-(3-chloro-4-cyanophenyl)-1H-pyrazol-1-yl)propan-2-yl)-5-((4-(4-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)methyl)-1H-pyrazole-3-carboxamide (45 mg, yield 24%) as white solid.

[0571] ^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (d, J = 5.6 Hz, 3H), 1.23 (s, 1H), 1.57 (s, 1H), 1.70-1.80 (m, 2H), 1.98-2.08 (m, 2H), 2.22-2.40 (m, 8H), 2.52-2.68 (m, 2H), 2.84-2.94 (m, 1H), 3.30 (s, 1H), 3.51 (s, 2H), 4.18 (t, J = 6.4 Hz, 2H), 4.42-4.48 (m, 3H), 5.08-5.15 (m, 1H), 6.43 (s, 1H), 6.94 (d, J = 2.4 Hz, 1H), 7.32-7.6 (m, 1H), 7.42 (d, J = 2.0 Hz, 1H), 7.80-7.85 (m, 2H), 7.99 (s, 2H), 8.08 (s, 1H), 8.23 (d, J = 8.0 Hz, 1H), 11.1 (s, 1H), 13.1 (s, 1H).

[0572] Chemical Formula: $C_{39}H_{41}ClN_{10}O_6$; Molecular Weight: 781.26.

[0573] Total H count from HNMR data: 41.

[0574] LC_MS: (ES$^+$): m/z 781.4 $[M+H]^+$. t_R = 2.113 min.

[0575] **Synthetic Procedure for Compound 612:**

[0576] **Synthesis of tert-butyl 4-[[1-(2-fluoro-4-nitrophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate**：Into a 20-30 mL sealed tube, was placed a solution of 1,2-difluoro-4-nitrobenzene (977.5 mg, 6.1 mmol, 1.0 equiv) in dimethylsulfoxide (10 mL), tert-butyl 4-(piperidin-4-ylmethyl)piperazine-1-carboxylate (1.5 g, 5.1 mmol, 0.8 equiv), N,N-Diisopropylethylamine (2.0 g, 15.3 mmol, 2.5 equiv). The resulting solution was stirred overnight at 100°C in an oil bath. The reaction was then quenched by the addition of 10 mL of water/ice. The resulting solution was extracted with ethyl acetate (20 mL x 2) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:0). This resulted in 1.5 g (58%) of tert-butyl 4-[[1-(2-fluoro-4-nitrophenyl) piperidin-4-yl] methyl] piperazine-1-carboxylate as a yellow solid.

[0577] LC-MS (ES$^+$): *m/z* 423.30 $[MH^+]$, t_R =1.58 min (1.9 minute run).

[0578] **Synthesis of tert-butyl 4-[[1-(4-amino-2-fluorophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate:** To a solution of tert-butyl 4-[[1-(2-fluoro-4-nitrophenyl) piperidin-4-yl] methyl] piperazine-1-carboxylate (1.5 g, 3.6 mmol, 1.0 equiv) in 15 mL i-PrOH was added Pd/C (10%, 40.0 mg) under nitrogen atmosphere in a 100 ml round bottom flask. The flask was then vacuumed and flushed with hydrogen. The reaction mixture was hydrogenated at room temperature for 4 hours under hydrogen atmosphere using a hydrogen balloon, then filtered through a Celite pad and concentrated under reduced pressure. The resulting mixture was concentrated under reduced pressure. This resulted in 1.2 g (86%) of tert-butyl 4-[[1-(4-amino-2-fluorophenyl) piperidin-4-yl] methyl] piperazine-1-carboxylate as a pink solid.

[0579] LC-MS (ES^+): *m/z* 393.10 [MH^+], t_R = 0.74 min (1.9 minute run).

[0580] **3. Synthesis of tert-butyl 4-[(1-[4-[(1-cyanocyclobutyl)amino]-2-fluorophenyl]piperidin-4-yl)methyl]piperazine-1- carboxylate:** Into a 100 mL round-bottom flask, was placed a solution of tert-butyl 4-[[1-(4-amino-2-fluorophenyl) piperidin-4-yl] methyl] piperazine-1-carboxylate (1.2 g, 3.1 mmol, 1.0 equiv) in tetrahydrofuran (10 mL), cyclobutanone (428.6 mg, 6.1 mmol, 2.0equiv), $ZnCl_2$ (2.1 g, 15.4 mmol, 5.0 equiv), TMSCN (606.1 mg). The resulting solution was stirred overnight at room temperature. The reaction was then quenched by the addition of 10 mL of 1 mmol/L $FeSO_4$ solvent. The resulting solution was extracted with ethyl acetate (20 mL x 2) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was applied onto a silica gel column with dichloromethane/methanol (5:1). This resulted in 990.0 mg (69%) of tert-butyl 4-[(1-[4-[(1-cyanocyclobutyl) amino]-2-fluorophenyl] piperidin-4-yl) methyl] piperazine-1-carboxylate as a black solid.

[0581] LC-MS (ES^+): *m/z* 472.35 [MH^+], t_R = 1.50 min (1.9 minute run).

[0582] **Synthesis of tert-butyl 4-[(1-[4-[7-(5-chloro-6-cyanopyridin-3-yl)-8-imino-6-sulfanylidene-5,7-diazaspiro[3.4]octan-5-yl]-2-fluorophenyl]piperidin-4-yl)methyl]piperazine-1-carboxylate:** Into a 100-mL round-bottom flask, was placed a solution of tert-butyl 4-[(1-[4-[(1-cyanocyclobutyl)amino]-2-fluorophenyl]piperidin-4-yl)methyl]piperazine-1-carboxylate (402.6 mg, 1.3 mmol, 1.0 equiv) in toluene (10 mL), 3-chloro-5-isothiocyanatopyridine-2-carbonitrile (200.0 mg, 1.5 mmol, 1.2 equiv), 4-dimethylaminopyridine (156.4 mg, 1.9 mmol, 1.5 equiv). The resulting solution was stirred overnight at 100 °C in an oil bath. The reaction was then quenched by the addition of 5 mL of ice

water. The resulting solution was extracted with ethyl acetate (20 mL x 2) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:0). This resulted in 230.0 mg (30.4%) of tert-butyl 4-[(1-[4-[7-(5-chloro-6-cyanopyridin-3-yl)-8-imino-6-sulfanylidene-5, 7- diazaspiro [3.4] octan-5-yl]-2-fluorophenyl] piperidin-4-yl) methyl] piperazine-1-carboxylate as a yellow solid.

[0583] LC-MS (ES^+): *m/z* 667.30 [MH^+], t_R = 1.52 min (1.9 minute run).

[0584] **Synthesis 3-chloro-5-(5-[3-fluoro-4-[4-(piperazin-1-ylmethyl)piperidin-1-yl]phenyl]-8-oxo-6-sulfanylidene-5,7-diazaspiro[3.4]octan-7-yl)pyridine-2-carbonitrile hydrochloride:** Into a 100 mL round-bottom flask, was placed a solution of tert-butyl 4-[(1-[4-[7-(5-chloro-6-cyanopyridin-3-yl)-8-imino-6-sulfanylidene-5,7-diazaspiro[3.4]octan-5-yl]-2-fluorophenyl] piperidin-4-yl)methyl]piperazine-1-carboxylate (230.0 mg, 0.3 mmol, 1.0 equiv) in methanol (10 mL), hydrogen chloride (8 ml). The resulting solution was stirred for 2 hours at 80 °C in an oil bath. The reaction was then quenched by the addition of 5 mL of ice water. The resulting solution was extracted with ethyl acetate (20 mL x 2) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under reduced pressure. This resulted in 200.1 mg (96%) of 3-chloro-5-(5-[3-fluoro-4-[4-(piperazin-1-ylmethyl) piperidin-1-yl] phenyl]-8-oxo-6-sulfanylidene-5, 7- diazaspiro [3.4] octan-7-yl) pyridine-2-carbonitrile hydrochloride as a yellow solid.

[0585] LC-MS (ES^+): *m/z*568.25 [MH^+], t_R = 1.26 min (1.9 minute run).

[0586] **6. Synthesis 3-chloro-5-(5-[4-[4-([4-[2-(2, 6-dioxopiperidin-3-yl)-1,3-dioxo-2, 3-dihydro-1H-isoindol-5-yl] piperazin-1-yl] methyl) piperidin-1-yl]-3-fluorophenyl]-8-oxo-6-sulfanylidene-5, 7-diazaspiro[3.4]octan-7- yl)pyridine-2-carbonitrile:** Into a 20-30 mL sealed tube, was placed a solution of 3-chloro-5-(5-[3-fluoro-4-[4-(piperazin-1-ylmethyl)piperidin-1-yl]phenyl]-8-oxo-6-sulfanylidene-5,7-diazaspiro[3.4]octan-7-yl)pyridine-2-carbonitrile hydrochloride (100.0 mg, 0.2 mmol, 1.0 equiv) in dimethylsulfoxide (10 mL), 2-(2,6-dioxopiperidin-3-yl)-5-fluoro-2,3-dihydro-1H-isoindole-1,3-dione (40.6 mg, 0.2 mmol, 0.9 equiv), N,N-Diisopropylethylamine (56.9 mg, 0.44 mmol, 2.66 equiv). The resulting solution was stirred for 2 hours at 130 °C in an oil bath. The reaction was then quenched by the addition of 5 mL of water/ice. The resulting solution was extracted with of ethyl acetate (20 mL x 2) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under

reduced pressure. The crude product was purified by Prep-HPLC with the following conditions: Mobile Phase A: Water (10MMOL/L ammonium bicarbonate), Mobile Phase B: acetonitrile; Flow rate: 20 mL/min; Gradient: 56% B to 76% B in 8 min; 254 nm; Rt: 7.6 min; 5 mL product was obtained. This resulted in 72.7 mg (60%) of 3-chloro-5-(5-[4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-3-fluorophenyl]-8-oxo-6-sulfanylidene-5,7-diazaspiro[3.4]octan-7-yl)pyridine-2-carbonitrile as a yellow solid.

[0587] ^{1}H NMR (300 MHz, DMSO-d_6) δ 11.10 (s, 1H), 8.90 (d, J = 2.0 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 7.70 (d, J = 8.5 Hz, 1H), 7.37 (s, 1H), 7.33-7.13 (m, 4H), 5.09 (dd, J = 12.6, 5.4 Hz, 1H), 3.50-3.45 (m, 6H), 2.90-2.81 (m, 4H), 2.71-2.2.60 (m, 3H), 2.61-2.45 (m, 5H), 2.38-2.36 (m, 1H), 2.30- 2.28 (m, 2H), 2.05-1.99 (m, 2H), 1.89-1.80 (m, 2H), 1.76-1.72 (m, 1H), 1.61-1.55 (m, 1H), 1.39-1.22 (m, 2H); LC-MS (ES$^+$): m/z 824.35[MH$^+$], t_R = 1.27 min (2.9 minute run).

[0588] Chemical formula: $C_{41}H_{39}ClFN_9O_5S$ [823.25].

[0589] Total H count from HNMR data: 39.

[0590] **Synthetic Procedure for Compound 603:**

[0591] **1. Synthesis of 4-fluoro-N-hydroxybenzene-1-carboximidamide:** Into a 250-mL round-bottom flask, was placed 4-fluorobenzonitrile (5.00 g, 41.28 mmol, 1 equiv), hydroxylamine hydrochloride (14.0 g, 201.47 mmol, 4.880 equiv), ethanol (150 mL) and *N*-ethyl-*N*-isopropylpropan-2-amine (27.0 g, 208.90 mmol, 5.060 equiv). The resulting mixture was stirred for 4 hours at 85 °C in an oil bath. The resulting mixture was concentrated under reduced pressure. The residue was applied onto a silica gel column with methanol: ethyl acetate (1:10). This resulted in 5.70 g (89.57%) of 4-fluoro-N-hydroxybenzene-1-carboximidamide as a white solid.

[0592] LC-MS (ES$^+$): m/z 155.30 [MH]$^+$, t_R = 0.60 min (2.00 minute run).

[0593] **2. Synthesis of 3-(4-fluorophenyl)-5-methyl-1,2,4-oxadiazole:** Into a 100-mL round-bottom flask, was placed 4-fluoro-N-hydroxybenzene-1-carboximidamide (5.70 g, 36.98 mmol, 1.0 equiv), acetyl acetate (54.1 g, 529.93 mmol, 14.3 equiv) and acetic acid (5 mL). The resulting solution was stirred for 16 hours at 120 °C in an oil bath. The reaction was then quenched by the addition of 150 mL of water. The pH value of the solution was adjusted to 7 with $NaHCO_3$ solution. The resulting solution was extracted with ethyl acetate (30 mL x 2) and the organic layers combined. The resulting mixture was washed with brine (30 mL x 2). The mixture

was dried over with anhydrous sodium sulfate and the solid was filtered out. The resulting mixture was concentrated under reduced pressure. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 6.10 g (92.59%) of 3-(4-fluorophenyl)-5-methyl-1,2,4-oxadiazole as a white solid.

[0594] LC-MS (ES^+): *m/z* 178.95 $[MH]^+$, t_R = 1.18 min (2.00 minute run).

[0595] **3. Synthesis of tert-butyl 4-([1-[4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl]azetidin-3-yl]methyl)piperazine-1-carboxylate:** Into a 30-mL sealed tube, was placed 3-(4-fluorophenyl)-5-methyl-1,2,4-oxadiazole (523.0 mg, 2.94 mmol, 1.5 equiv), tert-butyl 4-(azetidin-3-ylmethyl)piperazine-1-carboxylate (500.0 mg, 1.96 mmol, 1 equiv), dimethyl sulphoxide (15 mL, 211.18 mmol, 107.852 equiv) and Cs_2CO_3 (1.91 g, 5.86 mmol, 3.0 equiv). The resulting suspension was stirred for 20 hours at 150 ^oC in an oil bath. The reaction was then quenched by the addition of 150 mL of water. The resulting solution was extracted with ethyl acetate (30 mL x 2) and the organic layers combined. The resulting mixture was washed with brine (30 mL x 2). The mixture was dried over anhydrous sodium sulfate. The solids were filtered out. The resulting mixture was concentrated under reduced pressure. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 380.0 mg (46.93%) of tert-butyl 4-([1-[4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl]azetidin-3-yl]methyl)piperazine-1-carboxylate as a light yellow solid.

[0596] LC-MS (ES^+): *m/z* 414.40 $[MH]^+$, t_R = 0.90 min (2.00 minute run).

[0597] **4. Synthesis of tert-butyl 4-[[1-(4-carbamimidoylphenyl)azetidin-3-yl]methyl]piperazine-1-carboxylate:** Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl 4-([1-[4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl]azetidin-3-yl]methyl)piperazine-1-carboxylate (380.0 mg, 0.92 mmol, 1 equiv), methanol (50 mL), AcOH (2 mL, 34.90 mmol, 37.981 equiv) and 10% Pd/C (300 mg). The flask was rapidly evacuated and then recharged with hydrogen gas for three times, and finally connected with a tyre containing H_2 (10 L). The resulting suspension was stirred for 16 hours at room temperature. The solids were filtered out and further washed successively with 200 mL methanol and 200 ml of acetonitrile. The combined filtrate was concentrated under reduced pressure. This resulted in 100.0 mg (29.14%) of tert-butyl 4-[[1-(4-carbamimidoylphenyl)azetidin-3-yl]methyl]piperazine-1-carboxylate as a white solid.

[0598] LC-MS (ES^+): *m/z* 374.05 $[MH]^+$, t_R = 0.75 min (2.00 minute run).

[0599] **5. Synthesis of 2-chloro-4-[[(1r,4r)-4-(2-bromoacetyl)cyclohexyl]oxy]benzonitrile:** Into a 100-mL three-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed (1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexane-1-carboxylic acid (400.0 mg, 1.43 mmol, 1.0 equiv), dichloromethane (30 mL), oxalyl chloride (220.0 mg, 1.73 mmol, 1.2 equiv) and a drop of *N,N*-dimethylacetamide (30.0 mg, 0.41 mmol, 0.29 equiv). The resulting solution was stirred for 4 hours at room temperature. The reaction mixture was concentrated under reduced pressure, and the residue was dissolved in acetonitrile (20 mL). Into the solution was added (Trimethylsilyl)diazomethane (2.2 mL , 4.4 mmol, 3.0 equiv, 2M in hexane). The resulting solution was stirred for 2 hours at room temperature, and then was cooled with an ice/water bath. Into the cooled mixture was added a solution of HBr in AcOH (40%) (1.05 g, 5.17 mmol, 3.613 equiv), and then the cool bath was removed. The resulting solution was allowed to react with stirring for an additional 2 hours at room temperature. The reaction was then quenched by the addition of 100 mL of water. The resulting solution was extracted with ethyl acetate (30 mL x 2) and the organic layers combined. The resulting mixture was washed with brine (30 mL x 2). The mixture was dried over anhydrous sodium sulfate and the solid was filtered out. The resulting mixture was concentrated under reduced pressure. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:5). This resulted in 380 mg (74.51%) of 2-chloro-4-[[(1r,4r)-4-(2-bromoacetyl)cyclohexyl]oxy]benzonitrile as a white solid.

[0600] LC-MS (ES^+): *m/z* 355.85 $[MH]^+$, t_R = 1.05 min (1.90 minute run).

[0601] **6. Synthesis of tert-butyl 4-[[1-(4-[5-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]-1H-imidazol-2-yl]phenyl)azetidin-3-yl]methyl]piperazine-1-carboxylate:** Into a 50-mL 3-necked round-bottom flask, was placed tert-butyl 4-[[1-(4-carbamimidoylphenyl)azetidin-3-yl]methyl]piperazine-1-carboxylate (100.0 mg, 0.27 mmol, 1.0 equiv), tetrahydrofuran (15 mL) and a solution of $NaHCO_3$ (113.0 mg, 1.35 mmol, 5.0 equiv) in water (3 mL). After stirring the mixture at room temperature for 5 minutes, into the flask, was added a solution of 2-chloro-4-[[(1r,4r)-4-(2-bromoacetyl)cyclohexyl]oxy]benzonitrile (96.0 mg, 0.27 mmol, 1 equiv) in THF (5 mL). The resulting solution was stirred for 24 hours at 50 °C in an oil bath. The reaction was then quenched by the addition of 150 mL of water. The resulting solution was extracted with ethyl acetate (30 mL x 2) and the organic layers combined. The resulting mixture was washed with brine (50 mL x 2). The mixture was dried over anhydrous

sodium sulfate and the solid was filtered out. The resulting mixture was concentrated under reduced pressure. The residue was applied onto a silica gel column with methanol: ethyl acetate (1:10). This resulted in 50.0 mg (29.43%) of tert-butyl 4-[[1-(4-[5-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]-1H-imidazol-2-yl]phenyl)azetidin-3-yl]methyl]piperazine-1-carboxylate as an off-white solid.

[0602] LC-MS (ES^+): *m/z* 631.15 $[MH]^+$, t_R = 0.94 min (2.00 minute run).

[0603] **7. Synthesis of 4-[[1-(4-[5-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]-1H-imidazol-2-yl]phenyl)azetidin-3-yl]methyl]piperazin-1-ium trifluoroacetate:** Into a 100-mL round-bottom flask, was placed tert-butyl 4-[[1-(4-[5-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]-1H-imidazol-2-yl]phenyl)azetidin-3-yl]methyl]piperazine-1-carboxylate (50.0 mg, 0.08 mmol, 1 equiv), dichloromethane (20 mL, 314.60 mmol, 3971.566 equiv) and 2,2,2-trifluoroacetaldehyde (5 mL). The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was concentrated under vacuum. This resulted in 50.0 mg (97.84%) of 4-[[1-(4-[5-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]-1H-imidazol-2-yl]phenyl)azetidin-3-yl]methyl]piperazin-1-ium trifluoroacetate as an off-white solid.

[0604] **8. Synthesis of 2-chloro-4-[[(1r,4r)-4-(2-[4-[3-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)azetidin-1-yl]phenyl]-1H-imidazol-5-yl)cyclohexyl]oxy]benzonitrile:** Into a 8-mL sealed tube, was placed 4-[[1-(4-[5-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]-1H-imidazol-2-yl]phenyl)azetidin-3-yl]methyl]piperazin-1-ium trifluoroacetate (50.0 mg, 0.08 mmol, 1 equiv), 2-(2,6-dioxopiperidin-3-yl)-5-fluoro-2,3-dihydro-1H-isoindole-1,3-dione (54.0 mg, 0.20 mmol, 2.522 equiv), dimethyl sulphoxide (4 mL, 56.31 mmol, 726.583 equiv) and *N*-ethyl-*N*-isopropylpropan-2-amine (2 mL, 12.10 mmol, 156.134 equiv). The resulting solution was stirred for 3 hours at 130 °C in an oil bath. The reaction was then quenched by the addition of 150 mL of water. The resulting solution was extracted with ethyl acetate (30 mL x 2) and the organic layers combined. The resulting mixture was washed with brine (50 mL x 2). The mixture was dried over anhydrous sodium sulfate and the solid was filtered out. The resulting mixture was concentrated under reduced pressure. The residue was applied onto a silica gel column with methanol: ethyl acetate (1:10). The crude product was further purified by Prep-HPLC (Column: XBridge Prep C18 OBD Column, 5um, 19*150mm; Mobile Phase A: Water (10 mmoL/L NH_4HCO_3), Mobile Phase B: ACN; Flow rate: 20 mL/min; Gradient: 45% B to 63% B in 8 min; 254 nm; Rt: 7.8 min). This

resulted in 31.5 mg (51.62%) of 2-chloro-4-[[(1r,4r)-4-(2-[4-[3-([4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)azetidin-1-yl]phenyl]-1H-imidazol-5-yl)cyclohexyl]oxy]benzonitrile as a yellow solid.

[0605] ^{1}H NMR (300 MHz, Me_2SO-d_6): δ 11.08 (s, 1H), 7.86 (d, J = 8.8 Hz, 1H), 7.71-7.67 (m, 3H), 7.39-7.25 (m, 3H), 7.13 (dd, J = 8.8, 2.3 Hz, 1H), 6.74 (s, 1H), 6.45 (d, J = 8.4 Hz, 2H), 5.07 (dd, J = 12.7, 5.5 Hz, 1H), 4.60 (brs, 1H), 3.98 (t, J = 7.6 Hz, 2H), 3.54-3.30 (m, 7H), 3.29-3.120 (m, 5H), 3.11-2.79 (m, 3H), 2.67-2.55 (m, 3H), 2.18-2.01 (m, 5H), 1.75-1.35 (m, 4H); LC-MS (ES$^+$): *m/z* 787.50 [MH$^+$], HPLC: t_R = 5.13 min (10.00 minute run).


[0606] Chemical Formula: $C_{43}H_{43}ClN_8O_5$ [787.31].

[0607] Total H count from HNMR data: 43.

[0608] **Synthetic Procedure for Compound 609:**

[0609] **1. Synthesis of tert-butyl N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamate**

[0610]




[0611] Into a 100-mL 3-necked round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed a solution of tert-butyl N-[(1r,3r)-3-hydroxy-2,2,4,4-tetramethylcyclobutyl]carbamate (600.0 mg, 2.47 mmol, 1.00 equiv) in N,N-dimethylformamide (10.0 mL). This was followed by the addition of sodium hydride (198.0 mg, 8.25 mmol, 2.00 equiv), in portions at 0°C. After 30 minutes, to this was added 2-chloro-4-fluorobenzonitrile (459.0 mg, 2.95 mmol, 1.20 equiv). The resulting solution was stirred for 1 hour at 70°C. The reaction mixture was cooled to room temperature with a water bath. The reaction was then quenched by the addition of water (20 mL). The resulting solution was extracted with ethyl acetate (20 mL x 3) and the organic layers combined. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/5). This resulted in 100.0 mg (11%) of tert-butyl N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamate as colorless oil.

[0612] LC-MS (ES$^+$): *m/z* 279.10 [MH-100]$^+$, t_R= 1.20 min (2.5 minute run).

[0613] 2. Synthesis of 2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile

[0614]



[0615] Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamate (500.0 mg, 1.32 mmol, 1.00 equiv), hydrogen chloride/dioxane (3 mL, 4M), 1,4-dioxane (3 mL). The resulting solution was stirred for 1 hour at room temperature. The resulting mixture was concentrated under vacuum. This resulted in 447.0 mg (87%) of 2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile as a white solid.

[0616] 3. Synthesis of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate

[0617] Into a 100-mL a flask, was placed a solution of 2-(2,6-dioxopiperidin-3-yl)-5-fluoro-2,3-dihydro-1H-isoindole-1,3-dione (5 g, 18.10 mmol, 1.00 equiv) in methyl sulfoxide (30 mL), methyl sulfoxide (30 mL), N,N-diisopropylethylamine (12.5 mL, 2.00 equiv), tert-butyl piperazine-1-carboxylate (3.75 g, 20.13 mmol, 1.10 equiv). The resulting solution was stirred for 16 hours at 110℃ in an oil bath. The reaction was then quenched by the addition of water (100 mL). The resulting solution was extracted with ethyl acetate (100 mL x 3) and the organic layers combined. The resulting mixture was washed with brine (200 mL). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 6.5 g (81%) of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate as a yellow solid.

[0618] LC-MS (ES^+): *m/z* 443.00 [MH^+], t_R = 1.156 min, (2.0 minute run).

[0619] 4. Synthesis of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-3-hydroxy-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate

[0620] Into a 50-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate (2 g, 4.52 mmol, 1.00 equiv), tetrahydrofuran (30 mL). This was followed by the addition of Methylmagnesium bromide (6 mL) dropwise with stirring at -75℃ in 20 minutes. The resulting solution was stirred for 4 hours at 60℃. The reaction was then quenched by the addition of water (50 mL). The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined. The resulting mixture was washed with saturated sodium chloride aqueous solution (50 mL). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:0). This resulted in 1.5 g (72%) of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-3-hydroxy-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate as yellow crude oil.

[0621] **5. Synthesis of 3-[3-methyl-1-oxo-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindol-2-yl]piperidine-2,6-dione**

[0622] Into a 100-mL round-bottom flask, was placed tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-3-hydroxy-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate (1.5 g, 3.27 mmol, 1.00 equiv), dichloromethane (20 mL), triethylsilane (15 mL). This was followed by the addition of Boron trifluoride etherate (15 mL) dropwise with stirring. The resulting solution was stirred for 16 hours at 25℃. The resulting solution was diluted with water (35 mL). The resulting solution was extracted with ethyl acetate (15 mL x 3) and the aqueous layers combined and concentrated under vacuum. This resulted in 1.12 g (100%) of 3-[3-methyl-1-oxo-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindol-2-yl]piperidine-2,6-dione as yellow crude oil.

[0623] LC-MS (ES^+): *m/z* 343.00 [MH^+], t_R = 0.658 min, (2.0 minute run).

[0624] **6. Synthesis of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate**

[0625] Into a 100-mL round-bottom flask, was placed 3-[3-methyl-1-oxo-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindol-2-yl]piperidine-2,6-dione (1.12 g, 3.27 mmol, 1.00 equiv), dichloromethane (50 mL), $(Boc)_2O$ (1.5 g, 6.87 mmol, 2.10 equiv). This was followed by the addition of triethylamine (4 mL) dropwise with stirring. The resulting solution was stirred for 16 hours at 25℃. The resulting solution was diluted with water (50 mL). The resulting solution was

extracted with dichloromethane (20 mL x 3) and the organic layers combined. The resulting mixture was washed with saturated sodium chloride aqueous solution (15 mL x 2). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (8:2). The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column,, 5um,19*150mm; mobile phase, Water(10MMOL/L NH_4HCO_3) and ACN (32.0% ACN up to 41.0% in 9 min); Detector, UV 254nm. This resulted in 435 mg (30%) of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate as yellow oil.

[0626] LCMS42-PH-ARV-LS-046-E-20-3(60861-135Q)1T.

[0627] LC-MS (ES^+): *m/z* 443.40 [MH^+], t_R = 2.034 min, (4.6 minute run).

[0628] **7. Synthesis of 3-[3-methyl-1-oxo-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindol-2-yl]piperidine-2,6-dione; trifluoroacetic acid**

[0629] Into a 50-mL round-bottom flask, was placed tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate (435 mg, 0.98 mmol, 1.00 equiv), dichloromethane (30 mL), trifluoroacetic acid (1 mL). The resulting solution was stirred for 3 hours at 25℃. The resulting mixture was concentrated under vacuum. This resulted in 500 mg (1.11%) of 3-[3-methyl-1-oxo-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindol-2-yl]piperidine-2,6-dione; trifluoroacetic acid as yellow crude oil.

[0630] **8. Synthesis of 4-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0631] Into a 25-mL round-bottom flask, was placed 4-[4-(hydroxymethyl)piperidin-1-yl]benzoic acid (250 mg, 1.06 mmol, 1.00 equiv), N,N-dimethylformamide (6 mL), HATU (485 mg, 1.28 mmol, 1.20 equiv). This was followed by the addition of N,N-diisopropylethylamine (550 mg, 4.26 mmol, 4.01 equiv) in 5 minutes. To this was added 2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile hydrochloride (335 mg, 1.06 mmol, 1.00 equiv). The resulting solution was stirred for 2 hours at 25℃. The resulting solution was diluted with water (25 mL). The resulting solution was extracted with ethyl acetate (15 mL x 3) and the organic layers combined. The resulting mixture was washed with saturated sodium chloride aqueous solution (20 mL). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (10:1).

This resulted in 514 mg (98%) of 4-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as a off-white solid.

[0632] LC-MS (ES^+): *m/z* 496.05 [MH^+], t_R = 1.133 min, (2.0 minute run).

[0633] **9. Synthesis of 4-(4-formylpiperidin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0634] Into a 50-mL round-bottom flask, was placed 4-[4-(hydroxymethyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (600 mg, 1.21 mmol, 1.00 equiv), dichloromethane (20 mL). This was followed by the addition of Dess-Martin (1.02 g), in portions at 0℃. The resulting solution was stirred for 2 hours at 25℃. The reaction was then quenched by the addition of saturated sodium bicarbonate solution (50 mL) and sodium thiosulfate solution (50 mL). The resulting solution was extracted with dichloromethane (30 mL x 3) and the organic layers combined. The resulting mixture was washed with saturated sodium chloride aqueous solution (50 mL). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 450 mg (75%) of 4-(4-formylpiperidin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as a off-white solid.

[0635] LC-MS (ES^+): *m/z* 494.10 [MH^+], t_R = 1.074 min, (1.9 minute run).

[0636] **10. Synthesis of 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0637] Into a 50-mL round-bottom flask, was placed 4-(4-formylpiperidin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (130 mg, 0.26 mmol, 1.00 equiv), 3-[3-methyl-1-oxo-5-(piperazin-1-yl)-2,3-dihydro-1H-isoindol-2-yl]piperidine-2,6-dione; trifluoroacetic acid (110 mg, 0.24 mmol, 0.92 equiv), dichloromethane (20 mL). This was followed by the addition of DIEA (0.3 mL) dropwise with stirring in 16 hours. To this was added sodium triacetoxyborohydride (550 mg), in portions. The resulting solution was stirred for 16 hours at 25℃. The reaction was then quenched by the addition of water (20 mL). The resulting solution was extracted with dichloromethane (15 mL x 3) and the organic layers combined. The resulting mixture was washed with saturated sodium chloride aqueous solution (20 mL). The mixture was dried over anhydrous sodium sulfate and concentrated under

vacuum. The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column,, 5um,19*150mm; mobile phase, Water(10MMOL/L NH_4HCO_3) and ACN (53.0% ACN up to 72.0% in 8 min); Detector, UV 254nm. This resulted in 53.4 mg (25%) of 4-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-3-methyl-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as a white solid.

[0638] H-NMR-PH-ARV-LS-046-E-0: (300 MHz, DMSO, ppm) δ 10.90-10.88 (br, 1H), 7.92-7.89 (d, J = 9Hz, 1H), 7.75-7.72 (d, J = 9Hz, 2H), 7.51-7.44 (m, 2H), 7.21 (br, 1H), 7.10-6.94 (m, 5H), 4.65-4.64 (m, 1H), 4.54-4.52 (m, 1H), 4.32 (s, 1H), 4.06-4.03 (d, J = 9Hz, 1H), 3.88-3.84 (m, 2H), 3.31-3.28 (m, 4H), 2.83-2.75 (m, 3H), 2.63-2.58 (m, 6H), 2.22-2.20 (m, 2H), 1.96-1.94 (m, 1H), 1.83-1.79 (m, 3H), 1.43-1.38 (m, 3H), 1.21 (s, 6H), 1.18-1.17 (m, 2H), 1.12 (s, 6H); LC-MS (ES^+): *m/z* 820.30/822.30 [MH^+], t_R = 2.042 min, (3.0 minute run).

[0639] Chemical Formula: $C_{46}H_{54}ClN_7O_5$ [819.39/821.39].

[0640] Total H count from HNMR data: 54.

[0641] **Synthetic Procedure for Compound 88:**

[0642] **Synthesis of4-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-hydroxy-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide:** Into a 100-mL round-bottom flask, was placed a solution of 4-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (330 mg, 0.40 mmol, 1.00 equiv) in acetic acid (20 mL), Zn (100 mg, 3.00 equiv). The resulting solution was stirred for 30 minutes at room temperature. The resulting solution was diluted with 100 of methyl alcohol. The resulting mixture was concentrated under vacuum. This resulted in 320 mg (97%) of 4-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-hydroxy-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as white oil. *m/z* 827.40 [MH^+], t_R =0.799min (1.90 minute run).

[0643] **2. Synthesis of-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0644] Into a 100-mL round-bottom flask, was placed a solution of 4-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-3-hydroxy-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide (320 g, 386.76 mmol, 1.00 equiv) in trifluoroacetic acid (10 mL), triethylsilane (3 mL, 0.30 equiv). The resulting solution was stirred for 30 minutes at room temperature. The resulting mixture was concentrated under vacuum. The crude product (100 mg) was purified by Column: XBridge Prep OBD C18 Column 19*250mm,5um;Mobile Phase A:Water(0.1%FA), Mobile Phase B: ACN; Flow rate: 20 mL/min; Gradient: 55% B to 75% B in 8 min; 220 nm; Rt: 6.05 min. This resulted in 49.2 mg of 4-(6-[4-[2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1-oxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]hexyl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide as a white solid. [(1r,3r)-3-[4-cyano-3-(trifluoromethyl)phenoxy]-2,2,4,4-tetramethylcyclobutyl]benzamide as a white solid.

[0645] ^{1}H NMR (300 MHz, CD_3OD): δ7.81--7.69 (m, 3H), 7.53--7.46 (m, 1H),7.35--7.27(m,3H), 7.15 (s, 1H),6.97--6.94(m,1H), 5.17--5.04(m, 1H), 4.52--4.39 (m, 2H), 4.27(s , 1H),4.16 (s, 1H), 3.84--3.55 (m, 4H),3.36(s, 2H), 3.27--3.11(m,4H),3.01--2.61(m,4H),2.22--2.07(m,1H),1.87--1.63(m,4H),1.52--1.38(m,4H),1.33--1.16(m,12), *m/z* 811.45[MH^+], t_R =2.415 min (3.20 minute run).

[0646] Chemical formula: C45H52ClFN6O5 [810.37].

[0647] Total H count from HNMR data:52.

[0648] **Synthetic Procedure for Compound 571:**

[0649] **1. Synthesis of 5-bromo-3-fluorobenzene-1,2-dicarboxylic acid.**

[0650] Into a 100-mL round-bottom flask, was placed 4-bromo-2-fluoro-6-methylbenzoic acid (2.5 g, 10.73 mmol, 1.00 equiv), sodium hydroxide (1.72 g, 43.00 mmol, 4.00 equiv), water (20 mL), $KMnO_4$ (3.41 g, 2.00 equiv). The resulting solution was stirred for 16 hours at 100℃ in an oil bath. The solids were filtered out. The pH value of the solution was adjusted to 3 with hydrogen chloride (2 mol/L). The resulting solution was extracted with dichloromethane (10 mL x 1) and the organic layers combined. The resulting solution was extracted with ethyl acetate: methanol = 10:3 (10 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 518 mg (18%) of 5-bromo-3-fluorobenzene-1,2-dicarboxylic acid as a white solid and 1.365 g material as a white solid.

[0651] LC-MS (ES^+): *m/z* 264.75 [MH^+], t_R = 0.675 min (2.0 minute run).

[0652] **2. Synthesis of 1,2-dimethyl 5-bromo-3-fluorobenzene-1,2-dicarboxylate**

[0653] Into a 100-mL round-bottom flask, was placed 5-bromo-3-fluorobenzene-1,2-dicarboxylic acid (1.4 g, 5.32 mmol, 1.00 equiv), methanol (40 mL), sulfuric acid (2 mL). The resulting solution was stirred for 16 hours at 70℃. The reaction was then quenched by the addition of water (10 mL). The pH value of the solution was adjusted to 7 with sodium carbonate. The resulting solution was extracted with ethyl acetate (20 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/hexane (1:10). This resulted in 1.15 g (74%) of 1,2-dimethyl 5-bromo-3-fluorobenzene-1,2-dicarboxylate as colorless oil.

[0654] LC-MS (ES^+): *m/z* 292.80 [MH^+], t_R = 0.939 min (2.0 minute run).

[0655] **3. Synthesis of 1,2-dimethyl 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-fluorobenzene-1,2-dicarboxylate**

[0656] Into a 50-mL round-bottom flask, was placed 1,2-dimethyl 5-bromo-3-fluorobenzene-1,2-dicarboxylate (500 mg, 1.72 mmol, 1.00 equiv), tert-butyl piperazine-1-carboxylate (481 mg, 2.58 mmol, 1.50 equiv), Rouphos Pd (66 mg, 0.09 mmol, 0.05 equiv), Cs_2CO_3 (1.66 g, 5.09 mmol, 3.00 equiv), Toluene (20 mL). The resulting solution was stirred for 12 hours at 100℃ in an oil bath. The resulting solution was diluted with water (40 mL). The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 600 mg (88%) of 1,2-dimethyl 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-fluorobenzene-1,2-dicarboxylate as a solid.

[0657] LC-MS (ES^+): *m/z* 397.25 [MH^+], t_R = 1.238 min (2.0 minute run).

[0658] **4. Synthesis of 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-fluorobenzene-1,2-dicarboxylic acid**

[0659] Into a 100-mL round-bottom flask, was placed 1,2-dimethyl 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-fluorobenzene-1,2-dicarboxylate (800 mg, 2.02 mmol, 1.00 equiv), methanol/water/THF (16 mL), sodiumol (242.4 mg, 6.06 mmol, 3.00 equiv). The resulting solution was stirred for 16 hours at 25℃. The resulting solution was diluted with 16 mL of water (30 mL). The pH value of the solution was adjusted to 8 with hydrogen chloride (2 mol/L). Citric

acid monohydrate was employed to adjust the pH to 3. The resulting solution was extracted with ethyl acetate / methanol =10:1 (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 740 mg (100%) of 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-fluorobenzene-1,2-dicarboxylic acid as a yellow solid.

[0660] LC-MS (ES^+): *m/z* 369.00 [MH^+], t_R = 0.804 min (2.0 minute run).

[0661] **5. Synthesis of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate**

[0662] Into a 100-mL round-bottom flask, was placed 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-fluorobenzene-1,2-dicarboxylic acid (560 mg, 1.52 mmol, 1.00 equiv) in acetonitrile (25mL), CDI (986.6 mg, 6.08 mmol, 4.00 equiv), DIEA (785.6 mg, 6.08 mmol, 4.00 equiv), 3-aminopiperidine-2,6-dione (375.5 mg, 2.93 mmol, 1.50 equiv). The resulting solution was stirred for 5 hours at 70℃ in an oil bath. The resulting solution was diluted with water (40 mL). The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/hexane (1:1). This resulted in 700 mg (100%) of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate as a yellow solid.

[0663] LC-MS (ES^+): *m/z* 483.05 [MNa^+], t_R = 0.884 min (2.0 minute run).

[0664] **6. Synthesis of 2,2,2-trifluoroacetaldehyde-2-(2,6-dioxopiperidin-3-yl)-4-fluoro-6-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione**

[0665] Into a 100-mL round-bottom flask, was placed tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate (710 mg, 1.54 mmol, 1.00 equiv), dichloromethane (10 mL), trifluoroacetic acid (3 mL). The resulting solution was stirred for 2 hours at 25℃. The resulting mixture was concentrated under vacuum. The crude product was purified by re-crystallization from ethanol. This resulted in 320 mg (45%) of 2,2,2-trifluoroacetaldehyde-2-(2,6-dioxopiperidin-3-yl)-4-fluoro-6-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione as a yellow solid.

[0666] LC-MS (ES^+): *m/z* 361.2 [MH^+], t_R = 0.210 min (2.0 minute run).

[0667] **7. Synthesis of 2-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide**

[0668] Into a 100-mL round-bottom flask, was placed 2-(4-formylpiperidin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide (110 mg, 0.22 mmol, 1.00 equiv), 2,2,2-trifluoroacetaldehyde; 2-(2,6-dioxopiperidin-3-yl)-4-fluoro-6-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione (104 mg, 0.23 mmol, 1.30 equiv), dichloromethane (10 mL), acetyl ethaneperoxoate sodioboranyl acetate (70.67 mg, 0.33 mmol, 1.50 equiv). The resulting solution was stirred for 15 minutes at 25℃. The resulting solution was allowed to react, with stirring, for an additional 1 hour at 25℃. The resulting solution was allowed to react, with stirring, for an additional 2 hours at 25℃. The resulting solution was diluted with dichloromethane (30 mL). The resulting mixture was washed with water (30 mL x 1. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/ethyl acetate (3:1). The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column, 5um, 19*150mm; mobile phase, water (10mmol/L NH_4HCO_3) and acetonitrile (32.0% acetonitrile up to 75.0% in 8 min); Detector, UV 254/220 nm. This resulted in 145 mg (78%) of 2-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-7-fluoro-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide as a yellow solid.

[0669] ^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.10 (s, 1H), 8.75 (s, 2H), 7.90 (d, J = 8.7 Hz, 1H), 7.71 (d, J = 9.0 Hz, 1H), 7.24–7.21 (m, 2H), 7.07 – 6.99 (m, 2H), 5.10 (m, 1H), 4.74 (d, J = 13.2 Hz, 2H), 4.29 (s, 1H), 4.04 (d, J = 9.0 Hz, 1H), 3.50 (s, 4H), 3.00 – 2.95 (m, 3H), 2.53 – 2.50 (m, 5H), 2.21 – 2.19 (m, 2H), 2.08 – 1.81 (m, 5H), 1.21 – 1.04 (m, 14H); LC-MS (ES^+): *m/z* 840.30/842.30 [MH^+], t_R = 3.099 min (5.0 minute run).

[0670] Chemical Formula: $C_{43}H_{47}ClFN_9O_6$ [839.33/841.33].

[0671] Total H count from HNMR data: 47.

[0672] **<u>Synthetic Procedure for Compound 358:</u>**

[0673] **1. Synthesis of ethyl 4,6-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate**

[0674] Into a 500-mL round-bottom flask, was placed a solution of ethyl 3-oxobutanoate (10 g, 76.84 mmol, 1.00 equiv) in ethanol (200 mL), urea (6.9 g, 114.89 mmol, 1.50 equiv), acetaldehyde (5 g, 113.50 mmol, 1.50 equiv), hydrogen chloride (2 mL, 0.30 equiv). The resulting

solution was stirred for 5 hours at 85℃ in an oil bath. The resulting mixture was concentrated under vacuum. This resulted in 12 g (79%) of ethyl 4,6-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate as a white solid. *m/z*:199.13[MH^+], t_R =0.865 min (2.00 minute run).

[0675] 2. Synthesis of ethyl 4,6-dimethyl-2-oxo-1,2-dihydropyrimidine-5-carboxylate

[0676] Into a 100-mL round-bottom flask, was placed Nitric acid (20 mL, 5.00 equiv) over a period of 5 minutes. This was followed by the addition of ethyl 4,6-dimethyl-2-oxo-1,2,3,4-tetrahydropyrimidine-5-carboxylate (2 g, 10.09 mmol, 1.00 equiv) dropwise with stirring. The resulting solution was stirred for 3 hours at 0℃ in a water/ice bath. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (1:7). The collected fractions were combined and concentrated under vacuum. This resulted in 1.1 g (56%) of ethyl 4,6-dimethyl-2-oxo-1,2-dihydropyrimidine-5-carboxylate as yellow oil. *m/z*:197.14[MH^+], t_R =0.477 min (1.80 minute run).

[0677] 3. Synthesis of ethyl 2-chloro-4,6-dimethylpyrimidine-5-carboxylate

[0678] Into a 100-mL round-bottom flask, was placed a solution of ethyl 4,6-dimethyl-2-oxo-1,2-dihydropyrimidine-5-carboxylate (300 mg, 1.53 mmol, 1.00 equiv) in phosphoroyl trichloride (10 mL), N,N-Diethylaniline (0.1 mL, 0.10 equiv). The resulting solution was stirred for 3 hours at 105℃ in an oil bath. The reaction was then quenched by the addition of water. The resulting solution was extracted with ethyl acetate and the organic layers combined and concentrated under vacuum. This resulted in 100 mg (30%) of ethyl 2-chloro-4,6-dimethylpyrimidine-5-carboxylate as a yellow solid. *m/z* :214.85[MH^+], t_R =0.842 min (1.90 minute run).

[0679] 4. Synthesis of ethyl 2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-4,6-dimethylpyrimidine-5-carboxylate

[0680] Into a 100-mL round-bottom flask, was placed a solution of ethyl 2-chloro-4,6-dimethylpyrimidine-5-carboxylate (100 mg, 0.47 mmol, 1.00 equiv) in tetrahydrofuran (20 mL), triethylamine (0.3 mL, 0.30 equiv), tert-butyl piperazine-1-carboxylate (104 mg, 0.56 mmol, 1.20 equiv). The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:4). The collected fractions were combined and concentrated under

vacuum. This resulted in 180 mg (94%) of ethyl 2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-4,6-dimethylpyrimidine-5-carboxylate as a yellow solid. *m/z* :365.13[MH^+], t_R =1.418 min (2.00 minute run).

[0681] 5. Synthesis of 2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-4,6-dimethylpyrimidine-5-carboxylic acid

[0682] Into a 100-mL round-bottom flask, was placed a solution of ethyl 2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-4,6-dimethylpyrimidine-5-carboxylate (170 mg, 0.47 mmol, 1.00 equiv) in methanol/water (50/10 mL), sodium hydroxide (93 mg, 2.33 mmol, 5.00 equiv). The resulting solution was stirred for 12 hours at 50℃ in an oil bath. The resulting mixture was concentrated under vacuum. The pH value of the solution was adjusted to 5-6 with hydrogen chloride (1 mol/L). The resulting solution was extracted with ethyl acetate and the organic layers combined. The residue was applied onto a silica gel column with dichloromethane/methanol (4:1). The collected fractions were combined and concentrated under vacuum. This resulted in 150 mg (89%) of 2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-4,6-dimethylpyrimidine-5-carboxylic acid as a white solid. *m/z* 337.24[MH^+], t_R =1.106 min (2.00 minute run).

[0683] 6. Synthesis of tert-butyl 4-(4,6-dimethyl-5-[[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoyl]pyrimidin-2-yl)piperazine-1-carboxylate

[0684] Into a 100-mL round-bottom flask, was placed a solution of 2-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-4,6-dimethylpyrimidine-5-carboxylic acid (130 mg, 0.39 mmol, 1.00 equiv) in N,N-dimethylformamide (30 mL), N,N,N',N'-Tetramethyl-O-(7-azabenzotriazol-1-yl)uronium hexafluorophospate (220 mg, 1.50 equiv), N,N-Diisopropylethylamine(0.3mL,0.30equiv),2-chloro-4-[(1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile hydrochloride (129 mg, 0.41 mmol, 1.20 equiv). The resulting solution was stirred for 30 minutes at room temperature. The resulting solution was extracted with ethyl acetate and the organic layers combined. The resulting mixture was washed with of brine. The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate (100%). The collected fractions were combined and concentrated under vacuum. This resulted in 150 mg (63%) of tert-butyl 4-(4,6-dimethyl-5-[[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-

tetramethylcyclobutyl]carbamoyl]pyrimidin-2-yl)piperazine-1-carboxylate as a yellow solid. *m/z* 597.13[MH^+], t_R =1.428 min (2.00 minute run).

[0685] 7. Synthesis of4,6-dimethyl-2-(piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide hydrochloride

[0686] Into a 100-mL round-bottom flask, was placed tert-butyl 4-(4,6-dimethyl-5-[[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoyl]pyrimidin-2-yl)piperazine-1-carboxylate (140 mg, 0.23 mmol, 1.00 equiv), dioxane/HCl (20 mL, 3.00 equiv). The resulting solution was stirred for 2 hours at room temperature. The resulting mixture was concentrated under vacuum. This resulted in 160 mg (122%) of 4,6-dimethyl-2-(piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide hydrochloride as a yellow solid. *m/z* :497.22[MH^+], t_R =1.046 min (2.00 minute run).

[0687] 8. Synthesis of2-[4-(3-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]propyl)piperazin-1-yl]-4,6-dimethyl-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide

[0688] Into a 25-mL sealed tube, was placed a solution of 4,6-dimethyl-2-(piperazin-1-yl)-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide hydrochloride (140 mg, 0.26 mmol, 1.00 equiv) in acetonitrile (10 mL), potassium carbonate (181 mg, 1.31 mmol, 5.00 equiv), Sodium iodide (51 mg, 1.50 equiv), 5-(3-bromopropoxy)-2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione (125 mg, 0.32 mmol, 1.20 equiv). The resulting solution was stirred for 12 hours at 70℃ in an oil bath. The resulting solution was diluted with 20 mL of acetonitrile. The resulting mixture was concentrated under vacuum. The crude product (50 mL) was purified by Prep-HPLC with the following conditions: Mobile Phase A:Water(0.1% Formic acid), Mobile Phase B: acetonitrile; Flow rate: 20 mL/min; Gradient: 30% B to 48% B in 8 min; 254 nm; Rt: 7.83 min. This resulted in 51 mg (28%) of 2-[4-(3-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]propyl)piperazin-1-yl]-4,6-dimethyl-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamide as a off-white solid.

[0689] 1H NMR (300 MHz, CDCL3) δ 8.29 (s, 1H), 7.81 (d, J = 8.3 Hz, 1H), 7.60 (d, J = 8.6 Hz, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.22 (dd, J = 8.3, 2.3 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.83 (dd, J = 8.8, 2.4 Hz, 1H), 5.96 (d, J = 8.6 Hz, 1H), 4.99 (dd, J = 11.9, 5.5 Hz, 1H), 4.24 (dt, J =

11.9, 7.0 Hz, 7H), 4.08 (s, 1H), 3.09 (s, 9H), 2.98 – 2.77 (m, 5H), 2.40(s, 2H), 2.21- 2.14 (m, 1H), 1.27 (d, J = 3.9 Hz, 12H); LC-MS (ES^+): *m/z* :811.32 [MH^+], t_R =6.278 min (10.00 minute run).

[0690] Chemical formula: $C_{42}H_{47}ClN_8O_7$ [810.33].

[0691] Total H count from HNMR data: 46.

[0692] **Synthetic Procedure for Compound 584:**

[0693] **1. Synthesis of 5-bromo-3-methoxybenzene-1,2-dicarboxylic acid)**

[0694] Into a 100-mL round-bottom flask, was placed 4-bromo-2-methoxy-6-methylbenzonitrile (800 mg, 3.54 mmol, 1.00 equiv), water (10 mL), sodium hydroxide (708 mg, 17.70 mmol, 5.00 equiv), $KMnO_4$ (1.12 g, 7.09 mmol, 2.00 equiv). The resulting solution was stirred for 16 hours at 100℃ in an oil bath. The solids were filtered out. The pH value of the solution was adjusted to 3 with hydrogen chloride (2 mol/L). The resulting solution was extracted with dichloromethane (15 mL x 3) and the aqueous layers combined. The resulting solution was extracted with ethyl acetate / methanol = 10:1 (15 mL x 3) and the organic layers combined and dried in an oven under reduced pressure, concentrated under vacuum. This resulted in 330 mg (34%) of 5-bromo-3-methoxybenzene-1,2-dicarboxylic acid as a white solid.

[0695] **2. Synthesis of 1,2-dimethyl 5-bromo-3-methoxybenzene-1,2-dicarboxylate)**

[0696] Into a 100-mL round-bottom flask, was placed 5-bromo-3-methoxybenzene-1,2-dicarboxylic acid (330 mg, 1.20 mmol, 1.00 equiv), methanol (20 mL), sulfuric acid (5 mL). The resulting solution was stirred for 16 hours at 70℃ in an oil bath. The resulting solution was diluted with water (40 mL). The pH value of the solution was adjusted to 8 with sodium carbonate. The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:10). This resulted in 340 mg (93%) of 1,2-dimethyl 5-bromo-3-methoxybenzene-1,2-dicarboxylate as a white solid.

[0697] LC-MS (ES+): m/z 302.85 [MH+], t_R = 0.906 min (2.0 minute run).

[0698] **3. Synthesis of 1,2-dimethyl 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-methoxybenzene-1,2-dicarboxylate**

[0699] Into a 100-mL round-bottom flask, was placed 1,2-dimethyl 5-bromo-3-methoxybenzene-1,2-dicarboxylate (300 mg, 0.99 mmol, 1.00 equiv), tert-butyl piperazine-1-carboxylate (277 mg, 1.49 mmol, 1.50 equiv), RouphosPd (39 mg, 0.05 mmol, 0.05 equiv), Cs_2CO_3 (978 mg, 3.00 mmol, 3.00 equiv), toluene (15 mL). The resulting solution was stirred for 12 hours at 100℃ in an oil bath. The resulting solution was diluted with water (30 mL). The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/ethyl acetate (10:1). This resulted in 340 mg (84%) of 1,2-dimethyl 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-methoxybenzene-1,2-dicarboxylate as light yellow oil.

[0700] LC-MS (ES+): m/z 409.05 [MH+], t_R = 0.963 min (2.0 minute run).

[0701] **4. Synthesis of 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-methoxybenzene-1,2-dicarboxylic acid**

[0702] Into a 100-mL round-bottom flask, was placed 1,2-dimethyl 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-methoxybenzene-1,2-dicarboxylate (340 mg, 0.83 mmol, 1.00 equiv), methanol/H_2O/THF (8 mL), sodiumol (100 mg, 2.50 mmol, 3.00 equiv). The resulting solution was stirred for 12 hours at 25℃ The resulting solution was diluted with water (30 mL). The pH value of the solution was adjusted to 8 with hydrogen chloride (2 mol/L). citric acid monohydrate was employed to adjust the pH to 3. The resulting solution was extracted with ethyl acetate (30 mL x 3) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 300 mg (95%) of 5-[4-[(tert-butoxy)carbonyl]piperazin-1-yl]-3-methoxybenzene-1,2-dicarboxylic acid as colorless oil.

[0703] LC-MS (ES+): m/z 306.95 [MH+], t_R = 0.853 min (2.0 minute run).

[0704] **5. Synthesis of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-7-methoxy-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate**

[0705] Into a 100-mL round-bottom flask, was placed tert-butyl 4-(7-methoxy-1,3-dioxo-1,3-dihydro-2-benzofuran-5-yl)piperazine-1-carboxylate (260 mg, 0.72 mmol, 1.00 equiv), 3-aminopiperidine-2,6-dione hydrochloride (153.6 mg, 0.93 mmol, 1.30 equiv), pyridine (10 mL). The resulting solution was stirred for 4 hours at 120℃ in an oil bath. The resulting solution was diluted with water (30 mL). The resulting solution was extracted with ethyl acetate (30 mL x 3)

and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/methanol (100:1). This resulted in 280 mg (83%) of tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-7-methoxy-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate as a yellow solid.

[0706] LC-MS (ES+): m/z 417.05 [MH+], t_R = 0.852 min (2.0 minute run).

[0707] **6. Synthesis of 2-(2,6-dioxopiperidin-3-yl)-4-methoxy-6-(piperazin-1-yl)isoindoline-1,3-dione**

[0708] Into a 50-mL round-bottom flask, was placed tert-butyl 4-[2-(2,6-dioxopiperidin-3-yl)-7-methoxy-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazine-1-carboxylate (270 mg, 0.57 mmol, 1 equiv), dichloromethane (6 mL, 0.07 mmol, 0.124 equiv), TFA (2 mL, 0.02 mmol, 0.031 equiv). The resulting solution was stirred for 2 hours at 25 ℃. The resulting mixture was concentrated to give 2-(2,6-dioxopiperidin-3-yl)-4-methoxy-6-(piperazin-1-yl)isoindoline-1,3-dione as a brown oil.

[0709] LC-MS (ES+): m/z 373.05 [MH+], t_R = 0.155 min (2.0 minute run).

[0710] **7. Synthesis of 6-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-7-methoxy-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide**

[0711] Into a 100-mL round-bottom flask, was placed 2,2,2-trifluoroacetaldehyde; 2-(2,6-dioxopiperidin-3-yl)-4-methoxy-6-(piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione (130 mg, 0.28 mmol, 1.078 equiv), dichloromethane (10 mL, 0.12 mmol), 6-(4-formylpiperidin-1-yl)-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide (120 mg, 0.26 mmol, 1 equiv), $NaBH(OAc)_3$ (163.4 mg, 0.77 mmol, 3.006 equiv). The resulting solution was stirred for 2 hours at 25℃. The resulting solution was diluted with dichloromethane (30 mL). The resulting mixture was washed with H_2O (30 mL x 3). The mixture was dried over anhydrous sodium sulfate and concentrated under vacuum. The resulting mixture was concentrated under vacuum. The residue was applied onto a silica gel column with dichloromethane/ethyl acetate (3:1). The crude product was purified by Prep-HPLC with the following conditions: Column, XBridge Prep C18 OBD Column, 5um,19*150mm; mobile phase, Water (10 mmol/L NH_4HCO_3) and acetonitrile (43% Phase B up to 65% in 8 min); Detector, uv. This resulted in 70 mg (33.11%) of 6-[4-([4-[2-(2,6-dioxopiperidin-3-yl)-7-methoxy-1,3-dioxo-2,3-dihydro-1H-isoindol-

5-yl]piperazin-1-yl]methyl)piperidin-1-yl]-N-[(1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl]pyridazine-3-carboxamide as a yellow solid.

[0712] ^{1}H NMR (400 MHz, DMSO-*d6*) δ 11.04 (s, 1H), 8.57 (d, *J* = 8.4Hz, 1H), 7.87–7.79 (m, 2H), 7.39–7.32 (m, 2H), 7.15–7.12 (m, 1H), 6.96 (s, 1H), 6.68 (s, 1H), 5.04–4.98 (m, 1H), 4.50–4.47 (m, 3H), 4.93–3.85 (m, 4H), 3.35-3.33 (m, 5H), 3.07 – 2.81 (m, 3H), 2.51 (s, 3H), 2.27 –22.1 (m, 2H), 2.09 –2.01 (m, 2H), 2.00 – 1.49 (m, 11H), 1.23 – 1.11(m, 3H); LC-MS (ES+): m/z 824.25/826.25 [MH+], t_R = 182 min (3.0 minute run).

[0713] Chemical Formula: $C_{42}H_{46}ClN_9O_7$ [823.32/825.32].

[0714] Total H count from HNMR data: 46.

[0715] **Synthetic Procedure for Compound 618:**

[0716] **1. Synthesis of N-(2,2-dimethoxyethyl)-3,4-dihydro-2H-pyrrol-5-amine**

[0717] Into a 250-mL round-bottom flask, was placed 5-methoxy-3,4-dihydro-2H-pyrrole (4.4 g, 1 equiv), MeOH (50 mL), 2,2-dimethoxyethan-1-amine (4.6 g). The resulting solution was stirred for 12 hours at 60 ℃ in an oil bath. The resulting mixture was concentrated under vacuum. This resulted in 5.7g of N-(2,2-dimethoxyethyl)-3,4-dihydro-2H-pyrrol-5-amine as brown oil.

[0718] LC-MS (ES^+): *m/z* 173.00[MH^+], t_R = 0.17min (1.9inute run).

[0719] **2. Synthesis of 5H,6H,7H-pyrrolo[1,2-a]imidazole**

[0720] Into a 250-mL round-bottom flask, was placed N-(2,2-dimethoxyethyl)-3,4-dihydro-2H-pyrrol-5-amine (5.72 g, 33.21 mmol, 1 equiv), formic acid (100 mL). The resulting solution was stirred for 16 hours at 100 ℃ in an oil bath. The resulting mixture was concentrated under vacuum. This resulted in 3.1 g (86.31%) of 5H,6H,7H-pyrrolo[1,2-a]imidazole as a brown solid.

[0721] LC-MS (ES^+): *m/z* 108.95[MH^+], t_R = 0.15 min (1.9 minute run).

[0722] **3. Synthesis of [5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]boronic acid**

[0723] Into a 100-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed 5H,6H,7H-pyrrolo[1,2-a]imidazole (1.0 g, 9.25 mmol, 1 equiv). This was followed by the addition of THF (20 mL, 246.86 mmol, 26.696 equiv). The resulting solution was turned to -78℃. To this was added n-BuLi (10 mL, 106.16 mmol, 11.480 equiv). The resulting solution was stirred for 10 minutes at -78℃. To the mixture was added

$B(Oi\text{-}Pr)_3$ (5.0 g, 26.59 mmol, 2.875 equiv). The resulting solution was stirred for 6 hours at room temperature. The PH was adjusted to 2 with 2 M HCL. The resulting mixture was concentrated under vacuum. This resulted in 2.1 g (crude) of [5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]boronic acid as a brown solid.

[0724] LC-MS (ES^+): *m/z* 512.4 [MH^+], t_R = 1.27 min (1.9 minute run).

[0725] **4. Synthesis of tert-butyl 4-[[1-(4-nitrophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate**

[0726] Into a 250-mL round-bottom flask, was placed a solution of 1-fluoro-4-nitrobenzene (3 g, 21.26 mmol, 1.00 equiv) in DMSO (80 mL). This was followed by the addition of DIEA (2.7 g, 20.89 mmol, 1.00 equiv) in several batches in 2 minutes. To this was added tert-butyl 4-(piperidin-4-ylmethyl)piperazine-1-carboxylate (6 g, 21.17 mmol, 1.00 equiv). The resulting solution was stirred for 6 hours at 100℃ in an oil bath. The reaction was then quenched by the addition of water (100 mL). The resulting solution was extracted with ethyl acetate (100 mL x 2) and the organic layers combined. The resulting mixture was washed with brine (30 mL x 2). The resulting solution was concentrated under vacuum. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1:1). This resulted in 8.0g (93%) of tert-butyl 4-[[1-(4-nitrophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate as a light yellow solid.

[0727] LC-MS (ES^+): *m/z* 405.10 [MH^+], t_R =0.65min (1.9minute run).

[0728] **5. Synthesis of tert-butyl 4-[[1-(4-aminophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate**

[0729] Into a 250-mL round-bottom flask, was placed a solution of tert-butyl 4-[[1-(4-nitrophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate (4.8 g, 11.87 mmol, 1.00 equiv) in methanol (60 mL) under nitrogen atmosphere. This was followed by the addition of Pd/C (0.96 g, 0.20 equiv). The flask was then vacuumed and flushed with hydrogen. The reaction mixture was hydrogenated at room temperature for 3 hours under hydrogen atmosphere using a hydrogen balloon, then filtered through a Celite pad and concentrated under reduced pressure. This resulted in 2.7 g (61%) of tert-butyl 4-[[1-(4-aminophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate as a light yellow solid.

[0730] LC-MS (ES^+): *m/z* 375.10 [MH^+], t_R =0.73min (2.0minute run).

[0731] **6. Synthesis of tert-butyl 4-[[1-(4-bromophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate**

[0732] Into a 250-mL round-bottom flask purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl 4-[[1-(4-aminophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate (2.0 g), acetonitrile (100 mg). This was followed by the addition of tert-butyl nitrite (770 mg). It was stirred for 30 minutes. To this was added dibromocopper (1.4 g). The resulting solution was stirred for 3 hours at 0℃ in a water/ice bath. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (0:100-1:2). This resulted in 320 mg of tert-butyl 4-[[1-(4-bromophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate as a white solid.

[0733] LC-MS (ES^+): *m/z* 438.15 [MH^+], t_R = 0.69 min (1.9 minute run).

[0734] **7. Synthesis of tert-butyl 4-([1-[4-(1,2-dimethyl-1H-imidazol-5-yl)phenyl]piperidin-4-yl]methyl)piperazine-1-carboxylate**

[0735] Into a 25-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed tert-butyl 4-[[1-(4-bromophenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate (230 mg), [5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]boronic acid (1.0 g), dioxane/H_2O (8 mL /2 mL), Na_2CO_3 (170 mg), Pd(dppf)Cl_2 (50 mg). The resulting solution was stirred for 3 hours at 90℃ in an oil bath. The residue was applied onto a silica gel column with dichloromethane/methanol (10:1). This resulted in 150 mg of tert-butyl 4-([1-[4-(1, 2-dimethyl-1H-imidazol-5-yl)phenyl]piperidin-4-yl]methyl)piperazine-1-carboxylate as brown oil.

[0736] LC-MS (ES^+): *m/z* 466.30 [MH^+], t_R = 0.61 min (1.9 minute run).

[0737] **8. Synthesis of 1-[[1-(4-[5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]phenyl)piperidin-4-yl]methyl]piperazine**

[0738] Into a 100-mL round-bottom flask, was placed tert-butyl 4-[[1-(4-[5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]phenyl)piperidin-4-yl]methyl]piperazine-1-carboxylate (100 mg, 0.21 mmol, 1 equiv), trifluoroacetyl (4 mL), dichloromethane (20 mL). The resulting solution was stirred for 4 hours at room temperature. The resulting mixture was concentrated under reduced pressure. This resulted in 80 mg (90.91%) of 1-[[1-(4-[5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]phenyl)piperidin-4-yl]methyl]piperazine as brown oil.

[0739] LC-MS (ES^+): *m/z* 366.00 [MH^+], t_R = 0.63 min (2.0 minute run).

[0740] **9. Synthesis of 2-(2,6-dioxopiperidin-3-yl)-5-(4-[[1-(4-[5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]phenyl)piperidin-4-yl]methyl]piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione**

[0741] Into a 25-mL sealed tube purged and maintained with an inert atmosphere of nitrogen, was placed 1-[[1-(4-[5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]phenyl)piperidin-4-yl]methyl]piperazine (100 mg, 0.27 mmol, 1 equiv), DMSO (5 mL), DIEA (1.5 m), 2-(2,6-dioxopiperidin-3-yl)-5-fluoro-2,3-dihydro-1H-isoindole-1,3-dione (80 mg, 0.29 mmol, 1.059 equiv). The resulting solution was stirred for 3 hours at 130℃ in an oil bath. The resulting solution was extracted with ethyl acetate (30 mL x 3). The resulting mixture was washed with brine (30 mL x 1). The resulting mixture was concentrated under reduced pressure. The crude product was purified by Prep-HPLC with the following conditions: Column, XB ridge Prep C18 OBD Column, 5um,19*150mm; mobile phase, Water (10mmol/L NH_4HCO_3) and acetonitrile (38% Phase B up to 55% in 8 min); Detector, UV. This resulted in 30.6 mg (17.99%) of 2-(2,6-dioxopiperidin-3-yl)-5-(4-[[1-(4-[5H,6H,7H-pyrrolo[1,2-a]imidazol-3-yl]phenyl)piperidin-4-yl]methyl]piperazin-1-yl)-2,3-dihydro-1H-isoindole-1,3-dione as a yellow solid.


[0742] ^{1}H NMR (300 MHz, DMSO-d_6) δ 11.07 (s, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.37 – 7.25 (m, 4H), 7.04 – 6.89 (m, 3H), 5.06 (dd, J = 12.7, 5.4 Hz, 1H), 4.09 (t, J = 7.0 Hz, 2H), 3.70 (d, J = 12.2 Hz, 2H), 3.43-3.41 (m, 5H), 2.90 – 2.49 (m, 12H), 2.20-2.10 (m, 2H), 2.00-1.90 (m, 1H), 1.80-1.73 (m, 3H), 1.21-1.17 (m, 2H); LC-MS (ES^+): m/z 622.35 [M+ H^+], t_R = 0.68 min, (2.90 minute run).

[0743] Chemical Formula: $C_{35}H_{39}N_7O_4$ [621.31].

[0744] Total H count from HNMR data: 39.

[0745] **<u>Synthetic Procedure for Compound 77:</u>**

[0746] Step 1: [6-(4-(tert-butoxycarbonyl)piperazin-1-yl)nicotinic acid].


[0747] 

[0748] A mixture of tert-butyl 4-(5-((benzyloxy)carbonyl)pyridin-2-yl)piperazine-1-carboxylate (2.0 g, 5.03 mmol) and palladium on carbon (10%, 200 mg) in ethanol (20 ml) was stirred at 30°C overnight under hydrogen atmosphere (hydrogen balloon). TLC showed the reaction was complete. Palladium on carbon was removed through filtration and washed with ethanol (20 ml x 2). The combined filtrates were concentrated under reduced pressure to give 6-

(4-(tert-butoxycarbonyl)piperazin-1-yl)nicotinic acid (1.6 g, crude) as colorless oil which was used in next step without purification.

[0749] Step 2: [tert-butyl 4-(5-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)pyridin-2-yl)piperazine-1-carboxylate].


[0750]



[0751] To a stirred solution of 6-(4-(tert-butoxycarbonyl)piperazin-1-yl)nicotinic acid (300 mg, 0.97 mmol), 4-((1r,3r)-3-amino-2,2,4,4-tetramethylcyclobutoxy)-2-chlorobenzonitrile hydrochloride (306 mg, 0.97 mmol), and N-ethyl-N-isopropylpropan-2-amine (309 mg, 2.4 mmol) in anhydrous N,N-dimethylformamide (8 ml) was added HATU (2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) (684 mg, 1.8 mmol) at 0°C, the resulting mixture was allowed to warm to room temperature and stirred for 20 minutes. TLC showed the reaction was complete. The mixture was partitioned between ethyl acetate (50 ml) and water (80ml). The organic layer was collected, washed with brine (10ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluted with 10 % methanol in dichloromethane) to afford tert-butyl 4-(5-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)pyridin-2-yl)piperazine-1-carboxylate (400 mg, yield 72%) as white solid.

[0752] Step 3 [N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(piperazin-1-yl)nicotinamide]

[0753]



[0754] A mixture of tert-butyl 4-(5-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)pyridin-2-yl)piperazine-1-carboxylate (80 mg, 0.14 mmol) in hydrogen chloride in dioxane solution (4M, 2 ml) was stirred at room temperature for 2 hours.

TLC showed the reaction was complete. The volatiles were evaporated under reduced pressure. The residue was taken up in dichloromethane (20 ml) and washed with aqueous sodium bicarbonate solution (1N, 5 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by pre-TLC (eluted with 10 % methanol in dichloromethane) to afford [N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(piperazin-1-yl)nicotinamide (32 mg, yield 50%) as white solid.


[0755] LC_MS: (ES^+): m/z 468.6 $[M+H]^+$. t_R = 2.285 min.

[0756] ^{1}HNMR (400 MHz, CD_3OD): δ 1.07-1.38 (m, 12H), 3.12-3.40 (m, 4H), 3.51-3.86 (m, 1H), 3.94 (br, 3H), 4.17-4.30 (m, 2H), 6.99-7.15 (m, 2H), 7.74(s, 1H), 8.05(s, 1H), 8.48-8.68 (m, 2H).

[0757] Chemical Formula: $C_{25}H_{30}ClN_5O_2$; Molecular Weight: 467.99.


[0758] Total H count from HNMR data: 28.

[0759] Step 4: [2-(2,6-dioxopiperidin-3-yl)-5-(4-(2-hydroxyethyl)piperazin-1-yl)isoindoline-1,3-dione]

[0760] 

[0761] A mixture of 2-(piperazin-1-yl)ethanol (235.6 mg, 1.8 mmol), 2-(2,6-dioxopiperidin-3-yl)-5-fluoroisoindoline-1,3-dione (500 mg, 1.8 mmol) and N-ethyl-N-isopropylpropan-2-amine (468 mg, 3.6 mmol) in 1-methylpyrrolidin-2-one (5 ml) was stirred at 90°C for 12 hours. TLC showed the reaction was complete. The reaction mixture was partitioned between ethyl acetate (30 ml) and water (50 ml). The organic layer was collected, washed with brine (20 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluted with 2-3% methanol in dichloromethane) to afford 2-(2,6-dioxopiperidin-3-yl)-5-(4-(2-hydroxyethyl)piperazin-1-yl)isoindoline-1,3-dione (370 mg, 62%) as yellow oil.


[0762] Step 2: [2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl 4-methylbenzenesulfonate]

[0763] 

[0764] To a solution of 2-(2,6-dioxopiperidin-3-yl)-5-(4-(2-hydroxyethyl)piperazin-1-yl)isoindoline-1,3-dione (370 mg, 0.96 mmol), N,N-dimethylpyridin-4-amine (12 mg, 0.1 mmol) and triethylamine (291 mg, 2.87 mmol) in dichloromethane (10 ml) was added 4-toluenesulfonyl chloride (201 mg, 1.05 mmol) at 0 °C. The reaction mixture was allowed to warm up to room temperature and stirred at room temperature overnight. TLC showed the reaction was complete. The reaction mixture was diluted with dichloromethane (10 ml), washed with water (10 ml × 2) then brine (10 ml), dried over anhydrous sodium sulfate and concentrated under reduced pressure to give 2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl 4-methylbenzenesulfonate (200 mg, 38%) as yellow solid which was used in next step without further purification.

[0765] LC_MS: (ES$^+$): m/z 541.20 $[M+H]^+$. t_R = 1.965 min.

[0766] Step 3: [N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperazin-1-yl)nicotinamide]

[0767] 

[0768] A mixture of 2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl 4-methylbenzenesulfonate (139 mg, 0.25 mmol), N-ethyl-N-isopropylpropan-2-amine (55.3 mg, 0.43 mmol) and N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(piperazin-1-yl)nicotinamide (100 mg, 0.21 mmol) in dry N,N-dimethylformamide (2 ml) was stirred at 50°C for 16 hours. TLC showed the reaction was complete. The mixture was partitioned between ethyl acetate (20 ml) and water (20 ml). The

organic layer was collected, washed with brine (10 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by pre-TLC (eluted with 10% methanol in dichloromethane) to afford N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperazin-1-yl)nicotinamide (40 mg, 18%) as yellow solid.

[0769] ^{1}H NMR (400 MHz, δ6-DMSO): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.05 (m, 1H), 2.54-2.73 (m, 12H), 2.84-2.92 (m, 1H), 3.41-3.70 (m, 10H), 4.06 (d, J = 9.2 Hz, 1H), 4.30 (s, 1H), 5.05-5.10 (m, 1H), 6.89 (d, J = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, J = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (br, 1H), 7.63 (d, J = 9.2 Hz, 1H), 7.69 (d, J = 8.4 Hz, 1H), 7.90 (d, J = 8.4 Hz, 1H), 7.96-7.99 (m, 1H), 8.63 (d, J = 2.0 Hz, 1H), 11.08 (s, 1H).

[0770] Chemical Formula: $C_{44}H_{50}ClN_9O_6$; Molecular Weight: 836.38.

[0771] Total H count from HNMR data: 50.

[0772] LC_MS: (ES$^+$): m/z 836.40 $[M+H]^+$. t_R = 2.408 min.

[0773] **Synthetic Procedure for Compound 383:**

[0774] **Synthesis of tert-butyl 4-[(4-bromo-2-fluorophenyl)carbonyl]piperazine-1-carboxylate**

[0775] Into a 100.0-mL round-bottom flask, was placed 4-bromo-2-fluorobenzoic acid (5.0 g, 22.83 mmol, 1.00 equiv), N,N-dimethylformamide (20.0 mL), HATU (10.5 g, 27.61 mmol, 1.20 equiv), DIEA (11.9 g, 92.08 mmol, 4.00 equiv), tert-butyl piperazine-1-carboxylate (4.3 g, 23.09 mmol, 1.00 equiv). The resulting solution was stirred overnight at room temperature. The reaction was then quenched by the addition of 20.0 mL of water. The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined. The resulting mixture was washed with sodium chloride (40.0 mL). The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/4). The collected fractions were combined and concentrated under vacuum. This resulted in 8.82 g (100%) of tert-butyl 4-[(4-bromo-2-fluorophenyl)carbonyl]piperazine-1-carboxylate as a yellow solid.

[0776] **2. Synthesis of 2-[[4-([4-[(tert-butoxy)carbonyl]piperazin-1-yl]carbonyl)-3-fluorophenyl]amino]-2-methylpropanoic acid**

[0777] Into a 100.0-mL round-bottom flask, was placed tert-butyl 4-[(4-bromo-2-fluorophenyl)carbonyl]piperazine-1-carboxylate (4.0 g, 10.33 mmol, 1.00 equiv), N,N-dimethylformamide (10.0 mL), potassium carbonate (3.58 g, 25.90 mmol, 2.50 equiv), CuI (393.8 mg, 2.07 mmol, 0.20 equiv), 2-acetylcyclohexan-1-one (261.0 mg, 1.86 mmol, 0.20 equiv), 2-amino-2-methylpropanoic acid (1.6 g, 15.52 mmol, 1.50 equiv). The resulting solution was stirred overnight at 105.0℃ in an oil bath. The reaction was then quenched by the addition of 20.0 mL of water. The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined. The pH value of the solution was adjusted to 8 with hydrogen chloride (1.0 mol/L). The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined and dried over anhydrous sodium sulfate and concentrated under vacuum. This resulted in 3.9 g (92%) of 2-[[4-([4-[(tert-butoxy)carbonyl]piperazin-1-yl]carbonyl)-3-fluorophenyl]amino]-2-methylpropanoic acid as a yellow solid.

[0778] LC-MS (ES^+): m/z 410.2 [MH^+], t_R = 1.049 min, (2.0 minute run).

[0779] Chemical formula: $C_{20}H_{28}FN_3O_5$ [409.20].

[0780] **3. Synthesis of tert-butyl 4-([2-fluoro-4-[(1-methoxy-2-methyl-1-oxopropan-2-yl)amino]phenyl]carbonyl)piperazine-1-carboxylate**

[0781] Into a 250.0-mL round-bottom flask, was placed 2-[[4-([4-[(tert-butoxy)carbonyl]piperazin-1-yl]carbonyl)-3-fluorophenyl]amino]-2-methylpropanoic acid (2.7 g, 6.59 mmol, 1.00 equiv), N,N-dimethylformamide (20.0 mL), potassium carbonate (2.7 g, 19.54 mmol, 3.00 equiv), CH3I (2.8 g, 19.73 mmol, 3.00 equiv). The resulting solution was stirred overnight at room temperature. The reaction was then quenched by the addition of 20.0 mL of water. The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined. The resulting mixture was washed with sodium chloride (40.0 mL). The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). The collected fractions were combined and concentrated under vacuum. This resulted in 2.2 g (79%) of tert-butyl 4-([2-fluoro-4-[(1-methoxy-2-methyl-1-oxopropan-2-yl)amino]phenyl]carbonyl)piperazine-1-carboxylate as a yellow solid.

[0782] LC-MS (ES^+): m/z 424.05 [MH^+], t_R = 0.920 min, (1.90 minute run).

[0783] Chemical formula: $C_{21}H_{30}FN_3O_5$ [423.22].

[0784] 4. Synthesis of tert-butyl 4-[(4-[3-[4-cyano-3-(trifluoromethyl)phenyl]-5,5-dimethyl-4-oxo-2-sulfanylideneimidazolidin-1-yl]-2-fluorophenyl)carbonyl]piperazine-1-carboxylate

[0785] Into a 100.0-mL round-bottom flask, was placed tert-butyl 4-([2-fluoro-4-[(1-methoxy-2-methyl-1-oxopropan-2-yl)amino]phenyl]carbonyl)piperazine-1-carboxylate (500.0 mg, 1.18 mmol, 1.00 equiv), N,N-dimethylformamide (5.0 mL), 4-isothiocyanato-2-(trifluoromethyl)benzonitrile (404.3 mg, 1.77 mmol, 1.50 equiv), t-BuOK (198.24 mg, 1.77 mmol, 1.50 equiv). The resulting solution was stirred overnight at 100.0℃ in an oil bath. The reaction was then quenched by the addition of 20.0 mL of water. The resulting solution was extracted with ethyl acetate (40.0 mL) and the organic layers combined. The resulting mixture was washed with sodium chloride (40.0 mL). The mixture was dried over anhydrous sodium sulfate. The residue was applied onto a silica gel column with ethyl acetate/petroleum ether (1/1). The collected fractions were combined and concentrated under vacuum. This resulted in 500.0 mg (68%) of tert-butyl 4-[(4-[3-[4-cyano-3-(trifluoromethyl)phenyl]-5,5-dimethyl-4-oxo-2-sulfanylideneimidazolidin-1-yl]-2-fluorophenyl)carbonyl]piperazine-1-carboxylate as a yellow solid.

[0786] 5. Synthesis of 4-(3-[3-fluoro-4-[(piperazin-1-yl)carbonyl]phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile hydrochloride

[0787] Into a 50.0-mL round-bottom flask, was placed tert-butyl 4-[(4-[3-[4-cyano-3-(trifluoromethyl)phenyl]-5,5-dimethyl-4-oxo-2-sulfanylideneimidazolidin-1-yl]-2-fluorophenyl)carbonyl]piperazine-1-carboxylate (120.0 mg, 0.19 mmol, 1.00 equiv), methanol (3.0 mL), hydrogen chloride (0 mg). The resulting solution was stirred for 2 hours at room temperature. The crude product (mL) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Prep C18 OBD Column,, 5um,19*150mm; mobile phase, Water(0.1%FA) and acetonitrile (25.0% acetonitrile up to 62.0% in 8 min); This resulted in 89.9 mg (83%) of 4-(3-[3-fluoro-4-[(piperazin-1-yl)carbonyl]phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile hydrochloride as a yellow solid.

[0788] H-NMR (300 MHz, CD3OD) δ 8.17-8.15 (d, J=5.4 Hz, 2H), 8.01-7.98 (d, J=8.4 Hz, 1H), 7.64-7.58 (m, 1H), 7.41-7.37 (m, 2H), 5.49 (s, 1H), 3.83 (s, 2H), 3.46 (s,2H), 3.01-2.92 (d, J=28.5 Hz, 4H), 1.60 (s, 6H), 1.38 (s, 1H).

[0789] LC-MS (ES^+): m/z 520.05 [MH^+], t_R = 1.315 min, (3.0 minute run).

[0790] Chemical formula: $C_{24}H_{21}F_4N_5O_2S$ [519.14].

[0791] Total H count from HNMR data:22.

[0792] **6. Synthesis of 4-[3-(4-[[4-(4-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]butyl)piperazin-1-yl]carbonyl]-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile**

[0793] Into a 50.0-mL round-bottom flask, was placed 4-(3-[3-fluoro-4-[(piperazin-1-yl)carbonyl]phenyl]-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile hydrochloride (150.0 mg, 0.27 mmol, 1.00 equiv), CH3CN (5.0 mL), potassium carbonate (149.1 mg, 1.08 mmol, 4.00 equiv), NaI (40.5 mg, 1.00 equiv), 5-(4-bromobutoxy)-2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione (220.0 mg, 0.54 mmol, 2.00 equiv). The resulting solution was stirred overnight at 70.0℃ in an oil bath. The crude product (mL) was purified by Prep-HPLC with the following conditions (2#-AnalyseHPLC-SHIMADZU(HPLC-10)): Column, XBridge Prep C18 OBD Column,, 5um,19*150mm; mobile phase, Water(0.1%FA) and acetonitrile (25.0% acetonitrile up to 62.0% in 8 min); Detector, UV 254nm. mL product was obtained. This resulted in 27.2 mg (12%) of 4-[3-(4-[[4-(4-[[2-(2,6-dioxopiperidin-3-yl)-1,3-dioxo-2,3-dihydro-1H-isoindol-5-yl]oxy]butyl)piperazin-1-yl]carbonyl]-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-sulfanylideneimidazolidin-1-yl]-2-(trifluoromethyl)benzonitrile as a white solid.


[0794] H-NMR (300 MHz, CDCl3) δ 8.04-7.95 (m, 3H), 7.84-7.77 (m, 2H), 7.60-7.55 (m, 1H), 7.34-7.33 (d, J=1.8 Hz, 1H), 7.21-7.09 (m, 3H), 4.98-4.93 (m, 1H), 4.14-4.10 (m, 2H), 3.91 (s, 2H), 3.49 (s, 2H), 2.94-2.72 (m, 3H), 2.67-2.56 (m, 5H), 2.25-2.13 (m, 2H), 1.92-1.88 (m, 2H), 1.76 (s, 2H), 1.68 (s, 6H).

[0795] LC-MS (ES^+): m/z 848.20 [MH^+], t_R = 3.304 min, (5.0 minute run).

[0796] Chemical formula: $C_{41}H_{37}F_4N_7O_7S$ [847.24].


[0797] Total H count from HNMR data:37.

[0798] **Synthetic Scheme for Compound 450:**



[0799] **Experiments:**

[0800] **Step 1: [chloromethyl 2,5,8,11-tetraoxatridecan-13-yl carbonate]**




[0801] To a stirred solution of chloromethyl carbonochloridate (2.6 g, 20 mmol) in dichloromethane (40 mL) was added a mixture of 2,5,8,11-tetraoxatridecan-13-ol (4.16 g, 20 mmol) and triethylamine (2 g, 20 mmol) in dichloromethane (5 ml) dropwise at 0 °C. The resulting reaction mixture was stirred at 0 °C for 0.5 hour. TLC showed the reaction was completed. The reaction mixture was partitioned between tert-Butyl methyl ether (100 ml) and water (60 ml). The organic layer were collected, washed with brine (50 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluted with 30 % ethyl acetate in hexane) to afford chloromethyl 2,5,8,11-tetraoxatridecan-13-yl carbonate (2.2 g, yield 35.2%) as colorless oil.

[0802] 1H NMR (400 MHz, $CDCl_3$): δ 3.38 (s, 3H), 3.54-3.56 (m, 2H), 3.64-3.66 (m, 10H), 3.74-3.76 (m, 2H), 4.36-4.38 (m, 2H), 5.74 (s, 2H).

[0803] Chemical Formula: $C_{11}H_{21}ClO_7$; Molecular Weight: 300.73;

[0804] Total H count from HNMR data: 21.

[0805] **Step 2: [(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl 2,5,8,11-tetraoxatridecan-13-yl carbonate]**



[0806] A mixture of N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (320 mg, 0.39 mmol), cesium carbonate (190 mg, 0.58mmol) and chloromethyl 2,5,8,11-tetraoxatridecan-13-yl carbonate (140 mg, 0.47mmol) in N,N-dimethylformamide (6 mL) was stirred at room temperature for 2 hours. TLC showed the reaction was completed. The mixture was partitioned between ethyl acetate (50 ml) and water (40 ml). The organic layer were collected, washed with brine (30 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluting with 2.5% methanol in dichloromethane) to afford (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl 2,5,8,11-tetraoxatridecan-13-yl carbonate (160 mg, yield 38%) as yellow solid.


[0807] LC_MS: (ES^+): m/z 1084.7 $[M+H]^+$. t_R = 2.550 min.

[0808] ^1H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.18-1.27 (m, 2H), 1.22 (s, 6H), 1.77-1.83 (m, 3H), 2.04-2.12 (m, 1H), 2.21-2.22 (m, 2H), 2.31-2.45 (m, 1H), 2.58-2.67 (m, 3H), 2.73-2.85 (m, 3H), 3.01-3.12 (m, 1H), 3.23 (s, 3H), 3.31-3.33 (m, 2H), 3.40-3.52 (m, 15H), 3.59-3.62 (m, 2H), 3.86 (d, *J* = 12.4 Hz, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.19-4.21 (m, 2H), 4.32 (s, 1H), 5.25-5.29 (m, 1H), 5.67-5.69 (m, 2H), 6.96 (d, *J* = 8.8 Hz, 2H), 7.00 (dd, *J* = 2.4, 8.8 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.28 (d, *J* = 8.8 Hz, 1H), 7.36 (s, 1H), 7.50 (d, *J* = 9.2 Hz, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.74 (d, *J* = 8.4 Hz, 2H), 7.92 (d, *J* = 8.8 Hz, 1H).

[0809] Chemical Formula: $C_{56}H_{70}ClN_7O_{13}$; Molecular Weight: 1084.65;


[0810] Total H count from HNMR data: 70.

[0811] **Synthetic Scheme for Compound 448:**



[0812] **Experiments:**

[0813] **Step 1: [benzyl (2-aminoethyl)carbamate]**



[0814] To a stirred solution of ethane-1,2-diamine (10 g, 166 mmol) in anhydrous dichloromethane (50 ml) was added benzyl chloroformate (2.83 g, 16.6 mmol) in dichloromethane (10 ml) dropwise at 0°C over 30 min. The reaction mixture was stirred at 0°C for 3 hours. TLC showed the reaction was complete. The mixture solution was acidified with diluted hydrochloride acid (1N) till pH 4-5, and extracted with ethyl acetate (100 ml). The aqueous layer was collected, acidified with saturated sodium hydroxide solution till pH 9-10, and extracted with ethyl acetate (50 ml x 2). The combined organic layers were collected, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give benzyl (2-aminoethyl)carbamate (1.31g, yield 40%) as colorless oil which was used in next step without further purification.


[0815] LC_MS: (ES^+): m/z 195.2 $[M+H]^+$. t_R = 1.198 min.

[0816] ^1H NMR (400 MHz, $CDCl_3$): 2.787 (t, J =6.0 Hz, 2H), 3.14-3.28 (m, 2H), 5.07 (s, 2H), 7.27-7.35 (m, 5H).

[0817] Chemical Formula: $C_{10}H_{14}N_2O_2$; Molecular Weight: 194.23;

[0818] Total H count from HNMR data: 11.

[0819] **Step 2:**




[0820] To a stirred solution of benzyl (2-aminoethyl)carbamate (400 mg, 2.06 mmol), (S)-2-((tert-butoxycarbonyl)amino)-3-methylbutanoic acid (447 mg, 2.06 mmol), and N-ethyl-N-isopropylpropan-2-amine (797mg, 6.18mmol) in anhydrous N,N-dimethylformamide (4 ml) was added HATU (2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) (1.56 g, 4.12 mmol) at room temperature and stirred for 20 min. TLC showed the reaction was complete. The mixture was partitioned between ethyl acetate (40 ml) and water (20 ml). The organic layer was collected, washed with brine (30 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluting with 5% methanol in dichloromethane) to afford Compound 448-2 (670 mg, yield 83%) as white solid.

[0821] ^{1}H NMR (400 MHz, $CDCl_3$): 0.88 (d, J =6.8 Hz, 3H), 0.93 (d, J =6.8 Hz, 3H), 1.43 (s, 9H), 2.07-2.15 (m, 1H), 3.33-3.38 (m, 4H), 3.85 (t, J =1.2 Hz, 3H), 5.08 (s, 2H), 7.30-7.35 (m, 5H).

[0822] Chemical Formula: $C_{20}H_{31}N_3O_5$; Molecular Weight: 393.48;

[0823] Total H count from HNMR data: 28.

[0824] **Step 3: [(S)-tert-butyl (1-((2-aminoethyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate]**




[0825] Palladium on carbon (10%, 200 mg) was added to a stirred solution of Compound 448-2 (670 mg, 1.70 mmol) in methanol (20 ml). The reaction mixture was stirred under hydrogen atmosphere (hydrogen balloon) at room temperature overnight. TLC showed the reaction was complete. Palladium on carbon was removed through filtration and washed with methanol (5 ml x2). The combined filtrates were concentrated under reduced pressure to afford (S)-tert-butyl (1-((2-aminoethyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (470 mg, crude) as colorless oil which was used in next step directly without purification.

[0826] ^{1}H NMR (400 MHz, $CDCl_3$): 0.92 (d, J =6.8 Hz, 3H), 0.96 (d, J =6.8 Hz, 3H), 1.44 (s, 9H), 2.08-2.20 (m, 1H), 2.84 (t, J =5.6 Hz, 2H), 3.27-3.38 (m, 2H), 3.85-3.89 (m, 1H).

[0827] Chemical Formula: $C_{12}H_{25}N_3O_3$; Molecular Weight: 259.35;

[0828] Total H count from HNMR data: 21.


[0829] **Step4: (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-((tert-butoxycarbonyl)amino)-3-methylbutanamido)ethyl)carbamate**



[0830] To a stirred solution of N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide (300 mg, 0.36 mmol), and cesium carbonate (179 mg, 0.55 mmol) in dichloromethane (8 ml) and acetonitrile (1 ml) was added chloromethyl (4-nitrophenyl) carbonate (102 mg, 0.44 mmol) under nitrogen. The reaction mixture was stirred at room temperature under nitrogen for 3 hours. TLC showed the reaction was complete. To the reaction mixture was added (S)-tert-butyl (1-((2-aminoethyl)amino)-3-methyl-1-oxobutan-2-yl)carbamate (93 mg, 0.36 mmol), and the resulting mixture was continued stirring under nitrogen for 2 hours. TLC showed the reaction was complete. The mixture was partitioned between ethyl acetate (30 ml) and water (20 ml). The organic layer was collected, washed with brine (30 ml), dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give a crude residue which was purified by silica gel flash chromatography (eluted with 2.5% methanol in dichloromethane) to afford desired product (240 mg, yield 59%) as yellow solid.

[0831] LC_MS: (ES^+): m/z 1135.7 $[M+H]^+$. t_R = 2.578 min.

[0832] **Step 5: [(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-amino-3-methylbutanamido)ethyl)carbamate]**



[0833] A solution of (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-((tert-butoxycarbonyl)amino)-3-methylbutanamido)ethyl)carbamate (230 mg, 0.20 mmol) in dichloromethane (6 ml) in 4M hydrogen chloride in dioxane (2 ml) was stirred at room temperature for 2 hours. TLC showed the reaction was complete. The mixture solution was acidified with saturated sodium bicarbonate solution till pH 9-10, and extracted with dichloromethane (15 ml x 2). The organic layers were combined, washed with brine (30 ml), dried over sodium sulfate and concentrated under reduced to give a crude residue which was purified by silica gel flash chromatography (eluted with 2.5% methanol in dichloromethane) to afford (3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-amino-3-methylbutanamido)ethyl)carbamate (140 mg, yield 67%) as yellow solid.

[0834] LC_MS: (ES^+): m/z 1035.6 $[M+H]^+$. t_R = 2.215 min.


[0835] ^1H NMR (400 MHz, DMSO-d6): δ 0.76 (d, *J* =6.8 Hz, 3H), 0.84 (d, *J* =6.8 Hz, 3H), 1.12 (s, 6H), 1.18 (s, 2H), 1.22 (s, 6H), 1.80-1.87 (m, 4H), 2.07-2.09 (m, 2H), 2.21 (d, *J* =6.0Hz, 2H), 2.50 (s, 4H), 2.58-2.62 (m, 2H), 2.76-2.85(m, 3H), 2.90 (d, *J* =5.2 Hz, 1H), 3.00-3.06 (m, 3H), 3.11 (t, *J* =6.0 Hz, 2H), 3.45 (s, 4H), 3.86 (d, *J* =12.0 Hz, 2H), 4.05 (d, *J* =9.2 Hz, 1H), 4.32 (s, 1H), 5.20-5.24 (m, 1H), 5.55-5.61 (m, 2H), 6.95 (d, *J* =9.2 Hz, 2H), 6.99-7.02 (m, 1H), 7.20 (d, *J* =2.4 Hz, 1H), 7.27 (d, *J* =8.8 Hz, 1H), 7.32-7.35 (m, 2H), 7.48 (d, *J* =9.2 Hz, 1H), 7.69 (d, *J* =8.4 Hz, 1H), 7.73 (d, *J* =8.8 Hz, 2H), 7.90 (d, *J* =8.8 Hz, 2H).

[0836] Chemical Formula: $C_{54}H_{67}ClN_{10}O_9$; Molecular Weight: 1035.62;

[0837] Total H count from HNMR data: 67.

[0838] **Synthesis of ABM Moieties**

[0839] **ABM-1: 2-chloro-4-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile**



[0840] Step 1: Synthesis of 2-chloro-4-isothiocyanatobenzonitrile (**B**).

[0841] To a stirred solution of 4-amino-2-chlorobenzonitrile (**A**, 1 g, 6.55 mmol) in dichloromethane (9 mL) was added sodium bicarbonate (2.21 g, 26.31 mmol) and water (9 mL). The resulting mixture was cooled to 0 °C, to which thiophosgene (817 mg, 7.11 mmol) was added in drop wise in 30 min at 0 °C. The resulting mixture was then warmed up to room temperature and stirred at room temperature for 1 hour. The reaction mixture was diluted with dichloromethane (200 mL), washed with brine (50 mL x 2), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue. The residue was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v: v = 1: 30)) to give desired product (yield: 71%) ^{1}HNMR (400 MHz, $CDCl_3$): δ 7.69 (d, J = 8.0 Hz, 1H), 7.38 (s, 1H), 7.28 (m, 1H);

[0842] Step 2: Synthesis of 2-chloro-4-[3-(4-hydroxyphenyl)-5-imino-4, 4-dimethyl-2-sulfanylideneimidazolidin-1-yl]benzonitrile (**D**).

[0843] To a stirred solution of 2-chloro-4-isothiocyanatobenzonitrile (**B**, 399 mg, 2.05 mmol) in toluene (5 mL) was added 2-[(4-hydroxyphenyl)amino]-2-methylpropanenitrile (**C**, 300 mg, 1.70 mmol) and 4-dimethylaminopyridine (312 mg, 2.55 mmol). The resulting solution was then heated in an oil bath to 100 °C and stirred at the same temperature for 16h. LC-MS indicated formation of the desired product. The reaction mixture was concentrated under vacuum to give a crude reside which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v: v =1:1)) to give desired product (yield: 48%) as a brown solid. LC-MS (ES^+): m/z 370.95 [MH^+], t_R =0.74 min (2.0 minute run);


[0844] Step 3: Synthesis of 2-chloro-4-[3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo- 2-sulfanylideneimidazolidin-1-yl]benzonitrile (**ABM-1**).

[0845] To a stirred solution of 2-chloro-4-[3-(4-hydroxyphenyl)-5-imino-4, 4-dimethyl-2-sulfanylideneimidazolidin-1-yl]benzonitrile (**D**, 300 mg, 0.81 mmol) in methanol (6 mL) was

added aqueous hydrogen chloride (2N, 3.0 mL). The resulting solution was then heated in an oil bath to 100 °C and stirred at the same temperature for 2h. The reaction mixture was diluted with water (30 mL), extracted with ethyl acetate (60 mL x 3), washed with water (50 mL), dried over anhydrous sodium sulfate and concentrated under vacuum to give titled product (yield: 93%) as a yellow solid, which was used for the next step without any further purifications. LC-MS (ES^+): *m/z* 372.00 [MH^+], t_R =0.97 min (2.0 minute run).


[0846] Unless otherwise noted, the following intermediates and their analogs (for examples, but not limited to, analogs with substitutions such as halogens) were synthesized according to similar procedures described above for the synthesis of **ABM-1**, by utilizing corresponding starting materials and reagents.

[0847] **ABM-2: 2-fluoro-4-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile:**




ABM-2

[0848] **ABM-3: 4-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:**




ABM-3

[0849] **ABM-4: 5-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile:**




ABM-4

[0850] **ABM-5: 4-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-methoxybenzonitrile:**




ABM-5

[0851] ABM-6: 4-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-methylbenzonitrile:




ABM-6

[0852] ABM-7: 3-chloro-5-(3-(4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)picolinonitrile:




ABM-7

[0853] ABM-8: 4-(1-(4-hydroxyphenyl)-4-oxo-2-thioxo-8-oxa-1,3-diazaspiro[4.5]decan-3-yl)-2-(trifluoromethyl)benzonitrile:




ABM-8

[0854] ABM-9: 4-(1-(4-hydroxyphenyl)-8-methyl-4-oxo-2-thioxo-1,3,8-triazaspiro[4.5]decan-3-yl)-2-(trifluoromethyl)benzonitrile:




ABM-9

[0855] ABM-10: 4-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile




ABM-10.

[0856] ABM-11: 5-(5-(4-hydroxyphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-3-(trifluoromethyl)picolinonitrile:




ABM-11.

[0857] ABM-12: 4-(4-(3-(4-cyano-3-(trifluoromethyl)phenyl)-5,5-dimethyl-4-oxo-2-thioxoimidazolidin-1-yl)phenyl)butanoic acid:




ABM-12.

[0858] ABM-13: 2-chloro-4-(3-(4'-hydroxybiphenyl-4-yl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile:




ABM-13.

[0859] ABM-14: 4-(3-(4'-hydroxybiphenyl-4-yl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:




ABM-14.

[0860] ABM-15: 5-(3-(4'-hydroxybiphenyl-4-yl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile:




ABM-15

[0861] **ABM-16: 4-(3-(3-fluoro-4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:**

[0862] 


ABM-16

[0863] **ABM-17: 1-(4-hydroxyphenyl)-5,5-dimethyl-3-(4-nitro-3-(trifluoromethyl)phenyl)-2-thioxoimidazolidin-4-one:**

[0864] 


ABM-17

[0865] **ABM-18: 4-(3-(3,5-difluoro-4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:**

[0866] 


ABM-18

[0867] **ABM-19: 4-(3-(4-hydroxyphenyl)-4,4-dimethyl-2,5-dioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:**

[0868] 

ABM-19


[0869] **ABM-20: 4-(3-(6-hydroxypyridin-3-yl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:**



ABM-20

[0870]


[0871] **ABM-21: 2-chloro-4-(3-(3-fluoro-4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile:**



ABM-21

[0872]


[0873] **ABM-22: 4-(3-(3-fluoro-4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-2-methoxybenzonitrile:**



ABM-22

[0874]


[0875] **ABM-23: 5-(3-(3-fluoro-4-hydroxyphenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile:**



ABM-23

[0876]


[0877] **ABM-24: 5-(3-(2-fluoro-4'-hydroxybiphenyl-4-yl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)-3-(trifluoromethyl)picolinonitrile:**



ABM-24


[0878]

[0879] **ABM-25: 4-(4,4-dimethyl-5-oxo-3-(4-(piperidin-4-yl)phenyl)-2-thioxoimidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile:**


[0880] 

ABM-25.


[0881] **ABM-26: trans-2-Chloro-4-[3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile**.

[0882] 


[0883] **ABM-27: cis-2-Chloro-4-[3-amino-2,2,4,4-tetramethylcyclobutoxy]benzonitrile**

[0884] 


[0885] **ABM-28: trans 6-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridazine-3-carboxamide**

[0886] 

[0887] **ABM-29: trans tert-Butyl N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamate**.

[0888] 

[0889] **ABM-30: trans 4-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0890] 

[0891] Step 1: Synthesis of *tert*-butyl (4-((*trans*-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)carbamate.


[0892] A suspension of 4-((*tert*-butoxycarbonyl)amino)benzoic acid (1.50 g, 6.34 mmol) in methylene dichloride (40 mL) was charged with *N*,*N*-diisopropylethylamine (3.30 mL, 19.0 mmol), followed by 4-(*trans*-3-amino-2,2,4,4-tetramethylcyclobutoxy)-2-chlorobenzonitrile hydrochloride (2.0 g, 6.34 mmol). The mixture was stirred for several minutes and then charged with HATU (2.41 g, 6.34 mmol). The reaction mixture was allowed to stir at room temperature for 2 hours. The mixture was diluted with methylene dichloride (40 mL), washed with aqueous 1N HCl (2 x), saturated aqueous sodium bicarbonate (2 x), brine, and dried over anhydrous Na_2SO_4. The crude product was used in next step;

[0893] Step 2: synthesis of trans 4-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide.


[0894] 4M HCl in Dioxane (1.38 mL, 40.0 mmol) was added to a pre-mixed solution of *tert*-butyl (4-((*trans*-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)carbamate (2.00 g, 4.01 mmol) in MeOH (2 mL) and left to stir at room temperature for 1 hour till completion. The reaction mixture was concentrated in vacuo to a solid, which was dissolved with 5% MeOH in DCM. The organic layer was washed with sodium bicarbonate (2 x), filtered through a Biotage Universal Phase Separator and then concentrated in vacuo to a solid. The crude product was recrystallized from EtOH/Heptanes to afford the desired product as a white solid, 1.2 g, 75% yield. ^{1}H NMR (400 MHz, METHANOL-d4) δ 7.72 (d, J = 8.80 Hz, 1H), 7.61 (d, J = 8.61 Hz, 2H), 7.13 (d, J = 2.35 Hz, 1H), 6.98 (dd, J = 2.45, 8.71 Hz, 1H), 6.69 (d, J = 8.61 Hz, 2H), 4.28 (s, 1H), 4.12 (s, 1H), 1.27 (s, 6H), 1.22 (s, 6H). LC-MS (ES+): *m/z* 398.16/400.15 [MH^+].

[0895] Unless otherwise noted, the following intermediates and their analogs (for examples, but not limited to, analogs with substitutions such as halogens) were synthesized according to similar procedures described above for the synthesis of **ABM-30**, by utilizing corresponding starting materials and reagents.


[0896] **ABM-31: trans 5-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrazine-2-carboxamide**

[0897] 


[0898] **ABM-32: trans 2-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyrimidine-5-carboxamid**

[0899] 


[0900] **ABM-33: 4-Methoxy-N-[(1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0901] 


[0902] **ABM-34: trans 1-(2-Hydroxyethyl)-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]-1H-pyrazole-4-carboxamide**

[0903] 


[0904] **ABM-35: trans 6-Amino-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]pyridine-3-carboxamide**.

[0905] 


[0906] **ABM-36: trans 4-[(5-Hydroxypentyl)amino]-N-[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]benzamide**

[0907] 


[0908] **ABM-37: trans tert-Butyl 2-({5-[(4-{[3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl]carbamoyl}phenyl)aminopentyl}oxy)acetate**

[0909] 


[0910] **ABM-38: N-((1r,3r)-3-(4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-methylbenzamide**



[0911] **ABM-39: N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-methylpyridazine-3-carboxamide**





[0912] **ABM-40: N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-methylpyridazine-3-carboxamide**





[0913] **Synthesis of CLM Moieties**


[0914] Some non-limiting exemplary methods to generate the CLMs as described herein are summarized as shown below.


1 
CO2Me
CO2Me
R
+
H2N
CO2H
CONH2
Py, CDI, HCl
O
NH
N


2 
O
N
CO2H
CONH2
R
condensing agent
NH


3 
O
N
O
R
SiO2 or TFA
NH


4 
O
R
+
NH
H2N
Et3N, DMAP
benzotriazole
CDI
N


5 
O
N
R
NH3, CDI
NH


6 
COCl
COCl
R
+
CO2H
H2N
CONH2
Py, CDI
NH


7 


8 


9 


10 


11 


12 


13 


14 
CO2tBu
CO2H
R
H2N
NH
O
Alk
DIPEA
benzotriazole
carbodiimide
N


15 
HN
NH2
R
carbodiimide
N
H


16 
CO2H
NH2
R
H2N
carbodiimide
N
H


17 
CO2H
OH
R
Cl
OEt
H2N
NH
Et3N
N
H


18 


19 


20 

21 

22 

23 

24 



wherein R comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to a ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof.

[0915] As shown in representative reaction 1, dimethyl phthalate derivatives can be condensed with glutamine (racemate or enantiomer) or glutamine analogs then further reacted with agents such as carbonyl diimidazole to form 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives.

[0916] Alternatively as shown in representative reaction 2, the intermediate phthalimide produced in the initial condensation described above may be separately prepared and/or isolated

and then reacted with dehydrating agents such as trifluoroacetamide, $POCl_3$ or acetic anhydride to form the desired 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives. The same type of intermediate phthalimide can also be reacted with Lawesson's reagent prior to the dehydration step to provide thio analogs such as that shown in representative reactions 8 and 9.

[0917] Protected examples of 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives such as the N^1-BOC species shown in representative example 3 can be deprotected to reveal the target 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives by using, in this case, reagents such as TFA or silica.

[0918] Phthalic anhydrides such as that shown in representative example 4 can be ring-opened by reaction with amines such as 3-aminopiperidine-2,6-dione to form an intermediate carboxylate species, that on treatment with carbonyldiimidazole and benzotriazole will form the target 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives. Alternatively, the two components may be combined in the presence of acetic acid to provide desired product as shown in representative reaction 13.

[0919] In an analogous reaction, anhydride derivatives like those shown in representative reaction 5 may be reacted with amines (ammonia in the example shown) then carbonyldiimidazoleto form the desired 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives.

[0920] Where phthaloyl chlorides are available, direct condensation with glutamine (racemate or enantiomer) or glutamine analogs is possible, followed by further reaction with agents such as carbonyl diimidazole to form 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives as shown in representative reaction 6.

[0921] *o*-Bromobenzamides can be reacted with a source of CO such as the acid chloride shown in representative reaction 7 in the presence of a palladium catalyst and associated phosphine ligand to produce the desired 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives. Alternatively CO gas itself may be used in conjunction with rhodium (II) catalysts and silver carbonate to provide the desired products.

[0922] 2-(2,4-dioxo-1,2,3,4-tetrahydropyrimidin-5-yl)-2,3-dihydro-1H-isoindole-1,3-dione, and 5-(1,3-dioxo-2,3-dihydro-1H-isoindol-2-yl)-1,3-diazinane-2,4,6-trione derivatives can be prepared by analogous means to some of the methods described above for 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives. In representative reactions

20 and 21, a phthalic anhydride can be reacted with 5-amino-1,2,3,4-tetrahydropyrimidine-2,4-dione or 5-amino-1,3-diazinane-2,4,6-trione derivatives, respectively, in the presence of acetic acid to form the desired products.

[0923] Alternatively, 5-(1,3-dioxo-2,3-dihydro-1H-isoindol-2-yl)-1,3-diazinane-2,4,6-trione derivatives can be prepared by reaction of 5-amino-1,3-diazinane-2,4,6-trione derivatives with phthalic acid mono tert-butyl esters in the presence of Hünig's base, a carbodiimide and benzotriazole as shown in representative reaction 12. Similar conditions can be employed for the preparation of 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives from phthalic acid mono tert-butyl esters as shown in representative reaction 14.

[0924] Compounds such as 3-(2,6-dioxopiperidin-3-yl)-1,2,3,4-tetrahydroquinazoline-2,4-dione can be prepared from anthranilic acid derivatives by reaction of 3-aminopiperidine-2,6-diones with a carbodiimide as in representative reaction 16. The intermediate benzamide product may be isolated (or separately produced) and further reacted with a carbodiimide to produce 3-(2,6-dioxopiperidin-3-yl)-1,2,3,4-tetrahydroquinazoline-2,4-dione derivatives as shown in representative reaction 15.

[0925] 3-(2,6-Dioxopiperidin-3-yl)-3,4-dihydro-2H-1,3-benzoxazine-2,4-dione analogs can be prepared by activation of salicylic acids with chloroformates then condensation with 3-aminopiperidine-2,6-diones as shown in representative reaction 17.

[0926] 3,3-Dichloro-2,1λ^6-benzoxathiole-1,1-diones as shown in representative reaction 18 can be prepared by reaction of 2-sulfobenzoic acids with $POCl_3$ and PCl_5. These compounds can be reacted with amino derivatives to produce, for example, desired 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1λ^6,2-benzothiazole-1,1,3-trione derivatives.

[0927] As shown in representative reaction 19, anions of saccharin derivatives can be alkylated with electrophiles such as the 3-bromo-3-methylpiperidin-2-one to produce targeted 2-(3-methyl-2-oxopiperidin-3-yl)-2,3-dihydro-1λ^6,2-benzothiazole-1,1,3-trione derivatives.

[0928] Analogs of 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1λ^6,2-benzothiazole-1,1,3-trione may also be prepared by reaction of methyl 2-[(2,6-dioxopiperidin-3-yl)sulfamoyl]benzoate with strong bases such as sodium hydride (see representative reaction 20).

[0929] Deprotonation of 2-methyl-2,3-dihydro-1H-indene-1,3,dione derivatives with sodium ethoxide then reaction with electrophiles such as 3-bromopiperidin-2,6-dione affords 3-(2-methyl-1,3-dioxo-1H-inden-2-yl)piperidine-2,6-dione as shown in representative reaction 21.

[0930] Preparation of N^1-substituted compounds such as 2-[1-(benzyloxy)-2,6-dioxopiperidin-3-yl]-2,3-dihydro-1H-isoindole-1,4-dione (representative reaction 22) can be achieved by reaction of 2-(1,3-dioxo-2,3-dihydro-1H-isoindol-2-yl)pentanedioic acid with N-benzylhydroxylamine and with trifluoroacetic anhydride.

[0931] In turn, molecules such as 2-[1-(benzyloxy)-2,6-dioxopiperidin-3-yl]-2,3-dihydro-1H-isoindole-1,4-dione (representative reaction 23) may be subject to benzyl removal under hydrogenation conditions to yield N^1-hydroxy analogs such as 2-(1-hydroxy-2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione.

[0932] In representative reaction 24, methyl 1,3-dioxo-2,3-dihydro-1H-isoindole-2-carboxylate (and analogs) is reacted with 3-aminopiperidin-2-one to provide 2-(2-oxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-diones.

[0933] The same amine can also be reacted with phthalic anhydride derivatives in the presence of a Lewis acid such as zinc bromide and trimethylsilyl ether to yield the same type of product as shown in representative reaction 25. Intermediate products from this reaction if isolated or otherwise prepared (representative reaction 26) can be pushed to full cyclization through use of a dehydrating agent.

[0934] The isomeric derivatives such as 2-(6-oxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione shown in representative reaction 27 are attainable through reaction of phthalic acid with 5-aminopiperidin-2-one.


[0935] Preparation of N^1-substituted compounds such as 2-(1-benzyl-2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,4-dione (representative reactions 28 and 29) can be achieved through multiple routes. For example the anhydride (2-(2,6-dioxooxan-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione) can be condensed with 3-aminopiperidine-2,6-dione in the presence of DMAP and carbonyldiimidazole (representative reaction 28), or 2-(2,6-dioxopiperidin-3-yl)-2,3-dihydro-1H-isoindole-1,3-dione derivatives can be alkylated with electrophiles such as benzyl bromide in the presence of base as shown in representative reaction 29.

[0936] In some instances, protecting group strategies and/or functional group interconversions (FGIs) may be required to facilitate the preparation of the desired materials.

Such chemical processes are well known to the synthetic organic chemist and many of these may be found in texts such as "Greene's Protective Groups in Organic Synthesis" Peter G. M. Wuts and Theodora W. Greene (Wiley), and "Organic Synthesis: The Disconnection Approach" Stuart Warren and Paul Wyatt (Wiley).

[0937] **Synthesis of Linker Chemistry, L**

[0938] **L-1: 2-(3-(5-(tosyloxy)pentyloxy)propoxy)acetic acid**



[0939]

[0940] Step 1: Synthesis of ({[5-(prop-2-en-1-yloxy)pentyl]oxy}methyl)benzene

[0941] To a stirred solution of 5-(benzyloxy)pentan-1-ol (**W**, 4.0 g, 20.59 mmol) in N,N-dimethylformamide (50 mL) was added sodium hydride (1.24 g, 51.67 mmol) in portions at 0 °C under an atmosphere of nitrogen. The resulting mixture was then stirred at room temperature for 1 hour. To this mixture was added 3-bromoprop-1-ene (3.71 g, 30.67 mmol), the reaction mixture was stirred overnight at 60 °C in an oil bath. LC-MS indicated formation of the desired product. The reaction mixture was cooled to 0 °C and then quenched by water (100 mL), the resulting mixture was extracted with ethyl acetate (200 mL x 2). The organic layers were combined, washed with saturated aqueous solution of sodium chloride (60 mL), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue. The residue was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:40)) to give 4.57 g of **X**. ^{1}H NMR (300MHz, $CDCl_3$): δ 7.36(s, 4 H), 7.32 (m, 1 H), 5.98 (m, 1 H), 5.33 (m, 1H), 5.21 (m, 1H), 4.53 (s, 2H), 3.99 (m, 2H), 3.53 (m, 4H), 1.72 (m, 4H), 1.52 (m, 2H). LC-MS (ES^+): m/z 235.00 [MH^+], t_R = 1.18 min (2.0 minute run).

[0942] Step 2: Synthesis of 3-{[5-(benzyloxy)pentyl]oxy}propan-1-ol (**Y**)

[0943] To a 250-mL round-bottom flask with 9-BBN (0.5 M in THF, 77 mL) was added a solution of ({[5-(prop-2-en-1-yloxy)pentyl]oxy}methyl)benzene (**X**, 3.0 g, 12.80 mmol) in anhydrous tetrahydrofuran (20 mL) with stirring at 0 °C under an atmosphere of nitrogen. The resulting solution was stirred overnight at room temperature. LC-MS indicated formation of the desired product. Methanol (15 mL, with 30% sodium hydroxide and 30% H_2O_2) was added to the reaction and the resulting mixture was stirred at room temperature for 2 hours. This mixture was then extracted with ethyl acetate (20 mL x 3). The organic layers were combined, washed with saturated aqueous solution of sodium chloride (100 mL), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue. The residue was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v: v = 1:1)) to provide 1.96 g of **Y** as light yellow oil. ^{1}H NMR (300MHz, $CDCl_3$): δ7.34 (m, 5H), 4.49 (s, 2H), 3.75 (m, 2H), 3.59 (m, 2H), 3.49 (m, 4H), 2.65 (bs, 1 H), 1.84 (m, 2H), 1.68 (m, 4H), 1.50 (m, 2H). LC-MS (ES^+): m/z 253.17 [MH^+], t_R = 1.44 min (2.6 minute run).

[0944] Step 3: Synthesis of tert-butyl 2-(3-{[5-(benzyloxy)pentyl]oxy}propoxy)acetate (**Z**)

[0945] To a stirred solution of 3-{[5-(benzyloxy)pentyl]oxy}propan-1-ol (**Y**, 3.7 g, 14.66 mmol) in dichloromethane (30 mL) was added a solution of NaOH in water (37%, 30 mL) followed by tert-butyl 2-bromoacetate (11.39 g, 58.39 mmol) and TBACl (4.17 g). The resulting mixture was stirred at room temperature overnight. LC-MS indicated formation of the desired product. The reaction mixture was then extracted with ethyl acetate (50 mL x 3). The organic layers were combined, washed with saturated aqueous solution of sodium chloride (60 mL), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue. The residue was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:2) to give 3.2g of **Z** as a yellow oil. ^{1}H NMR (400MHz, $CDCl_3$): δ7.34(s, 4 H), 7.29 (m, 1 H), 4.50 (s, 4H), 4.3 (m, 2H), 3.51 (m, 4H), 3.42 (m, 2H), 1.98 (m, 2H), 1.67 (m, 4H), 1.48 (s, 9H), 1.46 (m, 2H). LC-MS (ES^+): m/z 367.25 [MH^+], t_R = 1.28 min (2.0 minute run).

[0946] Step 4: Synthesis of tert-butyl 2-[3-[(5-hydroxypentyl)oxy]propoxy]acetate (**AA**)

[0947] To a stirred solution of tert-butyl 2-(3-{[5-(benzyloxy)pentyl]oxy}propoxy)acetate (**Z**, 3.2 g, 8.73 mmol) in methanol (30 mL) was added AcOH (1.5 mL), palladium on carbon (1.5 g) under an atmosphere of nitrogen. Hydrogen was

then introduced to the reaction mixture via a hydrogen balloon, and the reaction was stirred at room temperature for 3 hours. The solid material was removed by filtration, the solution was concentrated under vacuum to provide 2.3 g of **AA** as light yellow oil, which was used for the next step without any further purifications. LC-MS (ES^+): m/z 277.10 [MH^+], t_R = 0.86 min (2.0 minute run).

[0948] Step 5: Synthesis of tert-butyl 2-[3-({5-[(4-methylbenzenesulfonyl)oxy]pentyl}oxy)propoxy]acetate (**AB**)

[0949] To a stirred solution of tert-butyl 2-[3-[(5-hydroxypentyl)oxy]propoxy]acetate (**AA**, 2.3 g, 8.32 mmol) in dichloromethane (30 mL) was added 4-methylbenzene-1-sulfonyl chloride (3.17 g, 16.63 mmol), triethylamine (2.52 g, 24.90 mmol) and 4-dimethylaminopyridine (203 mg, 1.66 mmol) at room temperature. The resulting mixture was stirred overnight at room temperature. The resulting mixture was concentrated under reduced pressure to give a crude residue, which was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:2) to give 2.6 g of **AB** as a yellow oil. ^1H NMR (300MHz, $CDCl_3$): δ 7.77 (d, J = 8.1 Hz, 2 H), 7.36 (d, J = 8.1 Hz, 2 H), 4.51 (s, 2H), 4.31 (m, 2H), 4.13 (m, 2H), 3.52 (m, 4H), 2.05 (s, 3H), 1.97 (m, 2H), 1.69 (m, 4H), 1.48 (s, 9H), 1.46 (m, 2H). LC-MS (ES^+): m/z 431.20 [MH^+], t_R = 1.21 min (2.0 minute run).

[0950] Step 1: Synthesis of 2-[3-({5-[(4-methylbenzenesulfonyl)oxy]pentyl}oxy)propoxy]acetic acid (**L-1**)


[0951] To a stirred solution of tert-butyl 2-[3-({5-[(4-methylbenzenesulfonyl)oxy]pentyl}oxy)propoxy]acetate (**AB**, 1.3 g, 3.02 mmol) in dichloromethane (10 mL) was added trifluoroacetic acid (10 mL) at room temperature. The resulting solution was stirred at room temperature for 3 hours. The reaction mixture was then concentrated under vacuum to give 1.5 g (crude) of **L-1**, which was used for next step without any further purification. LC-MS (ES^+): *m/z* 375.34 [MH^+], t_R = 1.39 min (2.6 minute run).

[0952]

[0953] The following Linkers (**L**) were prepared in a similar manner as for the preparation of **L-1**.

[0954]


[0955] L-2: 2-(3-(3,3-dimethyl-5-(tosyloxy)pentyloxy)propoxy)acetic acid

[0956] 

L-2 .

[0957]


[0958] **L-3: 2-(3-(3-hydroxy-5-(tosyloxy)pentyloxy)propoxy)acetic acid**

[0959] 

L-3 .

[0960]

[0961] **L-4: 2-(2-(2-(2-(tosyloxy)ethoxy)ethoxy)ethoxy)acetic acid**

[0962] 

.


[0963] To a stirred solution of ethyl 2-[2-(2-{2-[(4-methylbenzenesulfonyl)oxy]ethoxy}ethoxy)ethoxy]acetate (**AC**, 2 g, 5.12 mmol, 1.00 equiv) in methanol (20 mL) was added a solution of NaOH (500 mg, 12.50 mmol) in water (4 mL), and the resulting mixture was stirred at room temperature for 2 hours. Aqueous hydrogen chloride (1 M) was then added to the reaction mixture to adjust pH to ~5. Solids precipitated were collected by filtration to give **L-4** (yield: 98%). Mass (ES+): m/z 363, [MH+].

[0964]

[0965] The following Linkers (**L**) were prepared in a similar manner as for the preparation of **L-4**.

[0966]


[0967] **L-5: 2-(2-((2R,3R)-3-(2-(tosyloxy)ethoxy)butan-2-yloxy)ethoxy)acetic acid**

[0968] 

L-5 .

[0969]

[0970] **L-6: 2-(2-((2S,3S)-3-(2-(tosyloxy)ethoxy)butan-2-yloxy)ethoxy)acetic acid**





L-6

[0971]

[0972]

[0973] **L-7: 2-(4-(4-(tosyloxy)butoxy)butoxy)acetic acid**





[0974]

[0975] Step 1: Synthesis of 4-{4-[(4-methylbenzenesulfonyl)oxy]butoxy}butan-1-ol (**AE**)

[0976] To a stirred solution of 4-(4-hydroxybutoxy)butan-1-ol (**AD**, 2 g, 12.33 mmol) in dichloromethane (20 mL) was added Ag_2O (4.25 g, 18.49 mmol), KI (409 mg, 2.46 mmol) and TsCl (2.345 g, 12.30 mmol). The resulting mixture was stirred at room temperature for 12 hours. The inorganic salt formed was removed by filtration and the organic solution was concentrated under reduced pressure to give a crude residue. The residue was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:1)) to give **AE** (yield: 28%) as a colorless oil.

[0977] Step 2: Synthesis of ethyl 2-(4-{4-[(4-methylbenzenesulfonyl)oxy]butoxy}butoxy)acetate (**AF**)

[0978] To a stirred solution of 4-{4-[(4-methylbenzenesulfonyl)oxy]butoxy}butan-1-ol (**AE**, 1.1 g, 3.48 mmol) in dichloromethane (10 mL) was slowly added $BF_3.Et_2O$ (49.4 mg, 0.35 mmol) followed by ethyl 2-diazoacetate (794 mg, 6.96 mmol) at 0 °C. The resulting mixture was stirred overnight at room temperature. The reaction was then quenched by water (2.0 mL). The resulting mixture was extracted with dichloromethane (50mL x 3), the organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue. The residue was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v: v = 1:4) to give **AF** (yield: 93 as light yellow oil. Mass (ES^+): *m/z* 403.10 [MH^+].


[0979] Step 3: Synthesis of 2-(4-{4-[(4-methylbenzenesulfonyl)oxy]butoxy}butoxy)acetic acid (**L-7**)

[0980] To a stirred solution of ethyl 2-(4-{4-[(4-methylbenzenesulfonyl)oxy]butoxy}butoxy)acetate (**AF**, 1.3 g, 3.23 mmol) in methanol (25mL) was added a solution of NaOH (388 mg, 9.70 mmol) in water (6 mL) at room temperature. The resulting solution was stirred at room temperature for 4 hours. The bulk of organic solvent was removed under reduced pressure, to the resulting mixture was added aqueous hydrogen chloride (1.0 M) to adjust the pH = ~5. The solution was then extracted with ethyl acetate (250 mL x 3), the organic layers were combined and dried over anhydrous sodium sulfate, concentrated under reduced pressure to give **I-7** (yield: 93%) as light yellow oil. Mass (ES^+): *m/z* 375.05 [MH^+].

[0981]

[0982] **L-8: tert-butyl 2-(3-(4-(tosyloxy)butoxy)propoxy)acetate**

[0983]



[0984] Step 1. Synthesis of 3-[4-(benzyloxy)butoxy]propan-1-ol (**AH**)

[0985] To a stirred solution of propane-1, 3-diol (1.52 g, 19.98 mmol) in N, N-dimethylformamide (20 mL) was added sodium hydride (840 mg, 35.00 mmol) at room temperature, the resulting mixture was stirred at room temperature for 30 minutes. Then to the mixture was added 4-(benzyloxy) butyl 4-methylbenzene-1-sulfonate (**AG**, 6.68 g, 19.97 mmol) and the reaction was stirred overnight at 50 °C. TLC indicated formation of the desired product, at this time the reaction was allowed to cool down to room temperature. Water (10 mL) was added slowly to quench the reaction; the resulting mixture was then extracted with ethyl acetate (80 mL x 2). The organic layers were combined, washed with saturated aqueous solution of sodium chloride (20 mL), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue, which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:2)) to give **AH** (yield: 67%) as a light yellow oil. ^1H NMR (300 MHz, $CDCl_3$) δ 7.38-7.29 (m, 5H), 4.52 (m, 2H), 3.80 (m, 2H), 3.61 (m, 2H), 3.49-3.46 (m, 4H), 2.04 (m, 2H), 1.82 (m, 2H), 1.68 (m, 2H); Mass (ES^+): *m/z* 239.05 [MH^+].

[0986] Step 2. Synthesis of tert-butyl 2-[3-[4-(benzyloxy)butoxy]propoxy]acetate (**AI**).

[0987] To a stirred solution of 3-[4-(benzyloxy)butoxy]propan-1-ol (**AH**, 2.38 g, 9.99 mmol) in dichloromethane (15 mL) was added tert-butyl 2-bromoacetate (7.76 g, 39.78 mmol), TBAC (2.78 g, 10.00 mmol) followed by aqueous sodium hydroxide (37 %, 15 mL). The resulting mixture was stirred overnight at room temperature. The reaction mixture was then extracted with dichloromethane (100 mL x 3), the organic layers were combined, washed with saturated aqueous solution of sodium chloride (20 mL), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue. The residue was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1: 5)) to give **AI** (yield 57%) as a yellow oil. Mass (ES^+): *m/z* 353.10 [MH^+].

[0988] Step 3. Synthesis of tert-butyl 2-[3-(4-hydroxybutoxy)propoxy]acetate (**AJ**)


[0989] To a stirred mixture of tert-butyl 2-[3-[4-(benzyloxy)butoxy]propoxy]acetate (**AI**, 1 g, 2.84 mmol), palladium on carbon (10%, 200 mg) in methanol (20 mL) was added acetic acid (0.05 mL) under a nitrogen atmosphere. Hydrogen was then introduced to the reaction mixture via a balloon, the reaction was then stirred overnight at room temperature. The insoluble solids were removed by filtration and the solution phase was concentrated under reduced pressure to give the desired product (yield: 94%) as a yellow oil. Mass (ES^+): *m/z* 263.05 [MH^+].

[0990] Step 4. Synthesis of tert-butyl 2-(3-{4-[(4-methylbenzenesulfonyl)oxy]butoxy}propoxy)acetate (**L-8**)

[0991] To a stirred solution of tert-butyl 2-[3-(4-hydroxybutoxy)propoxy]acetate (**AJ**, 700 mg, 2.67 mmol) in dichloromethane (10 mL) was added 4-methylbenzene-1-sulfonyl chloride (558.4 mg, 2.93 mmol), TEA (539.5 mg, 5.33 mmol) and 4-dimethylaminopyridine (32.6 mg, 0.27 mmol). The resulting mixture was stirred overnight at room temperature. The bulk of solvent was removed under reduced pressure to give a crude residue, which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v= 1: 2)) to give titled product (yield: 52%) as a yellow oil. ^1H NMR (300 MHz, $CDCl_3$) δ7.79 (d, J = 8.4 Hz, 2H), 7.35 (d, J = 8.0 Hz, 2H), 4.05 (m, 2H), 3.95 (s, 2H), 3.59 (m, 2H), 3.48 (m, 2H), 3.38 (m, 2H), 2.46 (s, 3H), 1.82 (m, 2H), 1.70 (m, 2H), 1.57 (m, 2H), 1.50 (s, 9H); Mass (ES^+): *m/z* 417.05 [MH^+].

[0992]

[0993] L-9: tert-butyl 2-(4-(3-(tosyloxy)propoxy)butoxy)acetate




[0994]

[0995] **L-9** was prepared in a similar manner as that used to prepare **L-8**, except that **AK** was used in place of **AG**. Mass (ES^+): *m/z* 439.15 [MNa^+].

[0996]

[0997] **L-10: tert-butyl 2-(6-(tosyloxy)hexa-2,4-diynyloxy)acetate**



[0998]

[0999] Step1: Synthesis of tert-butyl 2-[(6-hydroxyhexa-2,4-diyn-1-yl)oxy]acetate (**AP**)

[1000] To a stirred solution of hexa-2, 4-diyne-1, 6-diol (**AO**, 100 mg, 0.91 mmol) in N, N-dimethylformamide (5 mL) was added sodium hydride (32 mg, 1.33 mmol) at 0 °C. The resulting mixture was then warmed up to room temperature and stirred at room temperature for 30 minutes. The reaction mixture was cooled to 0 °C followed by addition of tert-butyl 2-bromoacetate (176 mg, 0.90 mmol), and the resulting mixture was stirred at 0 °C for 2h. LC-MS indicated formation of the desired product. The reaction was then quenched by water (10 mL, added slowly) at 0 °C, and was extracted with ethyl acetate (20 x 2 mL). The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue, which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:2)) to give **AP** (yield: 49%) as a yellow oil.

[1001] Step 2. Synthesis of tert-butyl 2-({6-[(4-methylbenzenesulfonyl)oxy]hexa-2,4-diyn-1-yl}oxy)acetate (**L-10**)


[1002] To a stirred solution of tert-butyl 2-[(6-hydroxyhexa-2, 4-diyn-1-yl) oxy] acetate (**AP**, 50 mg, 0.22 mmol) in ether (2 mL) was added 4-toluenesulfonyl chloride (51 mg, 0.27 mmol) at 0 °C, followed by potassium hydroxide (125 mg, 2.23 mmol) in several batches at 0 °C. The resulting mixture was stirred at 0 °C for 4 hours. LC-MS indicated formation of the desired product. Water (10 mL) was added to the reaction, and the resulting mixture was extracted with ethyl acetate (20 mL x 2). The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue, which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v = 1:2)) to give **L-10** (yield: 71%) as a yellow oil. ^{1}H NMR (300 MHz, $CDCl_3$): δ 7.83 (d, J = 6.0 Hz, 2H), 7.39 (d, J = 6.0 Hz, 2H), 4.79 (s, 2H), 4.37 (s, 2H), 4.05 (s, 2H), 2.48 (s, 3H), 1.51 (s, 9H); LC-MS (ES$^+$): *m/z* 401.05 [MNa$^+$], t_R = 1.71 min (2.6 minute run).

[1003]

[1004] The following Linkers (**L**) were prepared in a similar manner as for the preparation of **L-10**.

[1005]


[1006] **L-11: tert-butyl 3-(6-(tosyloxy)hexa-2,4-diynyloxy)propanoate**

[1007] 

L-11 .

[1008]


[1009] **L-12: tert-butyl 4-(6-(tosyloxy)hexa-2,4-diynyloxy)butanoate**

[1010] 

L-12 .

[1011]

[1012] **L-13: ethyl 2-(2-(2-aminoethoxy)ethoxy)acetate hydrochloride**

[1013] 

.

[1014] Step 1: Synthesis of tert-butyl N-[2-(2-hydroxyethoxy)ethyl]carbamate (**AR**)

[1015] To a stirred solution of 2-(2-aminoethoxy)ethan-1-ol (**AQ**, 5.25 g, 49.94 mmol) in tetrahydrofuran (100 mL) was added aqueous solution of sodium bicarbonate (20% (w/w), 40 ml) and $(Boc)_2O$ (11.4 g, 52.23 mmol, added in several batches) at 0 °C. The resulting mixture was then warmed up slowly to room temperature and stirred at room temperature for 5 hours. The bulk of organic solvent was removed under reduced pressure and the resulting residue was diluted with water (300 mL), extracted with of ethyl acetate (100 mL x 3). The organic layers were combined, washed with saturated aqueous solution of sodium chloride (20 mL x 2), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give **AR** (yield: 98%) as colorless oil.


[1016] Step 2: Synthesis of ethyl 2-[2-(2-{[(tert-butoxy)carbonyl]amino}ethoxy)ethoxy]acetate (**AS**)

[1017] To a stirred solution of tert-butyl N-[2-(2-hydroxyethoxy)ethyl]carbamate (**AR**, 4.0 g, 19.49 mmol) in dichloromethane (30 mL) was added 1-diazo-3-methoxypropan-2-one (3.34 g, 29.27 mmol) and BF_3-Et_2O (0.2 mL) at room temperature. The resulting solution was stirred at room temperature for 2 hours. Water (20 mL) was added to the reaction mixture, organic layer was separated and washed with brine (20 mL), dried over anhydrous sodium sulfate and concentrated under reduced pressure to give a crude residue. The residue was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v: v = 1:2)) to give **AS** (yield: 18%) as yellow solid. ^{1}H NMR (400MHz, $CDCl_3$): δ 4.25-4.22 (q, J = 7.2 Hz, 2 H), 4.14 (s, 2 H), 3.74 (b, 2 H), 3.72 (b, 1 H), 3.67-3.32 (m, 4 H), 1.414 (s, 9 H), 1.31 (t, J = 7.2 Hz, 3 H).

[1018] Step 3: Synthesis of ethyl 2-[2-(2-aminoethoxy)ethoxy]acetate hydrochloride (**L-13**)

[1019] To a stirred solution of ethyl 2-[2-(2-{[(tert-butoxy)carbonyl]amino}ethoxy)ethoxy]acetate (**AS**, 500 mg, 1.72 mmol) in 1,4-dioxane (10 mL) was introduced hydrogen chloride (gas) via bubbling at room temperature for 2 hours. The solvent was then removed under vacuum to give **L-13** (yield: 99%). LC-MS (ES^+): m/z 192.00 [MH^+], t_R = 0.41 min (2.0 minute run).

[1020] L-14: ethyl 2-(5-aminopentyloxy)acetate

[1021] 

[1022] Step 1: Synthesis of tert-butyl 5-hydroxypentylcarbamate (**AU**)

[1023] To a stirred solution of 5-aminopentan-1-ol (**AT**, 3.1 g, 30.05 mmol) in dichloromethane (30 mL) was added di-tert-butyl dicarbonate (6.56 g, 30.06 mmol) at 0 °C. The resulting mixture was then stirred at room temperature for 4 hours. The solvent was removed under reduced pressure to give a crude residue which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v= 1: 2)) to give **AU** (yield: 98%) as a colorless oil. LC-MS (ES^+): *m/z* 204.00 [MH^+], t_R =1.29 min (2.6 minute run).

[1024] Step 2: Synthesis of ethyl 2-[(5-{[(tert-butoxy)carbonyl]amino}pentyl)oxy]acetate (**AV**)


[1025] To a stirred solution of tert-butyl N-(5-hydroxypentyl)carbamate (**AU**, 1.5 g, 7.38 mmol) in dichloromethane (10 mL) was added $BF_3{\cdot}Et_2O$ (0.1 mL) at 0 °C. To this mixture was then added a solution of ethyl 2-diazoacetate (850 mg, 7.45 mmol) in dichloromethane (2 mL) at 0 °C. The resulting mixture was allowed to warm up to room temperature and stirred at room temperature for 2 hours. Saturated aqueous sodium bicarbonate (30 mL) was added to the reaction, the resulting mixture was extracted with ethyl acetate (150 mL x 3). The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue, which was purified by flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v:v= 1: 7)) to give **AV** (yield: 15%) as a colorless oil. LC-MS (ES^+): *m/z* 290.05 [MH^+], t_R =1.55 min (2.6 minute run).

[1026] Step 3: Synthesis of ethyl 2-(5-aminopentyloxy)acetate (**L-14**)

[1027] To a stirred solution of ethyl ethyl 2-[(5-{[(tert-butoxy)carbonyl]amino}pentyl)oxy]acetate (**AV**, 400 mg, 1.38 mmol) in dichloromethane (5 mL) was added trifluoroacetic acid (5 mL) at room temperature. The resulting solution was stirred at room temperature for 2 hours. The reaction mixture was then concentrated under vacuum to give **L-14** (yield: 84%) as a yellow oil. LC-MS (ES^+): *m/z* 190.00 [MH^+], t_R =1.01 min (2.6 minute run).

[1028]

[1029] **L-15: methyl 2-(2-(2-(methylamino)ethoxy)ethoxy)acetate**



[1030]

[1031] Step 1: Synthesis of 2-[2-(benzylamino)ethoxy]ethan-1-ol (**AX**)

[1032] To a stirred solution of 2-(2-aminoethoxy)ethan-1-ol (**AW**, 5.0 g) and benzaldehyde (5.0 g) in THF (50 mL) was added sodium triacetoxyborohydride (15.8 g, 74.5 mmol) at 0 °C. The resulting solution was then stirred at room temperature for 4 hours. Water (50 mL) was added to the reaction and the resulting mixture was extracted with ethyl acetate (50 mL x 2). The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue, which was purified by flash silica gel chromatography (eluent: dichloromethane/methanol (v:v = 3:1) to give **AX** (yield: 85%) as a white solid. LC-MS (ES^+): *m/z* 195.95[MH^+], t_R = 0.22 min (2.0 minute run).

[1033] Step 2: Synthesis of 2-{2-[benzyl(methyl)amino]ethoxy}ethan-1-ol (**AY**)

[1034] To a stirred solution of of 2-[2-(benzylamino)ethoxy]ethan-1-ol (**AX**, 10.0 g) in methanol (200 mL) was added formaldehyde (38% in water) (4.9 mL) and triacetoxyborohydride (17.0 g) at room temperature. The resulting solution was stirred at room temperature for 2 hours. Saturated aq. sodium bicarbonate (100 mL) was added to the reaction, and bulk of organic solvent was then removed under reduced pressure. The resulting mixture was extracted with ethyl acetate (200 mL x 3). The organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure followed by high vacuum pump to give **AY** (yield: 33%) as a yellow oil. LC-MS (ES^+): *m/z* 210.00 [MH^+], t_R = 0.43 min (2.0 minute run).

[1035] Step 3: Synthesis of methyl 2-(2-{2-[benzyl(methyl)amino]ethoxy}ethoxy)acetate (**AZ**)

[1036] To a stirred solution of 2-{2-[benzyl(methyl)amino]ethoxy}ethan-1-ol (**AY**, 2 g) in dichloromethane (20 mL) was added a solution of sodium hydroxide (37%) in water (20 mL)


followed by tert-butyl 2-bromoacetate (7.76 g) and TBAC (2.78 g) at room temperature. The resulting mixture was stirred at room temperature for 15 hours. The aqueous layer was separated, and to which aq. hydrogen chloride (4N) was added to adjust the pH to ~3 before it was concentrated under reduced pressure to give a crude residue. Methanol (20 mL) was then added to this residue and insoluble salts were filtered out. The solution was concentrated under vacuum to give 2-(2-[2-[benzyl(methyl)amino]ethoxy]ethoxy)acetic acid (yield: 78%) as a yellow oil. To a stirred solution of 2-(2-{2-[benzyl(methyl)amino]ethoxy}ethoxy)acetic acid (2 g, 7.48 mmol, 1.00 equiv) prepare above in methanol (50 mL) was slowly added sulfuric acid (2 mL) at room temperature. The resulting solution was stirred at 70 °C in an oil bath for 3 hours. The bulk of solvent was removed under reduced pressure to give a residue, which was diluted with H_2O (30 mL). Sodium carbonate was then added to the mixture to adjust the pH to ~8. The mixture was then extracted with ethyl acetate (50 mL x 2), the organic layers were combined, dried over anhydrous sodium sulfate and then concentrated under reduced pressure followed by high vacuum pump to give **AZ** (yield: 29%) as a yellow oil. LC-MS (ES^+): *m/z* 281.95 [MH^+], t_R = 0.30 min (2.0 minute run).

[1037] Step 4: Synthesis of methyl 2-{2-[2-(methylamino)ethoxy]ethoxy}acetate (**L-15**)

[1038] To a stirred mixture of methyl 2-(2-{2-[benzyl(methyl)amino]ethoxy}ethoxy)acetate (**AZ**, 600 mg, 2.13 mmol) and palladium on carbon (300 mg) in methanol (30 mL) under a nitrogen atmosphere was charged with hydrogen gas via a balloon. The resulting mixture was stirred at room temperature for 15 hours. The solid material was removed by filtration and the solution was concentrated under vacuum to give **L-15** (400 mg) as yellow oil, which was used for next step without any further purifications. LC-MS (ES^+): *m/z* 191.95 [MH^+], t_R = 0.31 min (2.0 minute run).

[1039] **L-16: ethyl 2-(5-(methylamino)pentyloxy)acetate**

[1040]



[1041] Step 1: Synthesis of ethyl 2-[(5-{[(tert-butoxy)carbonyl](methyl)amino}pentyl)oxy]acetate (**BB**)


[1042] To a stirred solution of ethyl 2-[(5-{[(tert-butoxy)carbonyl]amino}pentyl)oxy]acetate (**BA**, 1.1 g, 3.8 mmol) in N,N-dimethylformamide (10 mL) was added CH_3I (0.71 mL, 11.4 mmol) at 0 °C, followed by sodium hydride (304 mg, 7.60 mmol, 60% in mineral oil) in several portions at 0 °C. The resulting mixture was stirred at room temperature for 16 hours. Water (1.0 mL) was added and the resulting mixture was extracted with ethyl acetate (50 mL x 2). The organic layers were combined, washed with saturated aqueous solution of sodium chloride (100 mL), dried over anhydrous sodium sulfate and then concentrated under reduced pressure to give a crude residue which was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v: v = 1: 10)) to give **BB** (yield: 21%) as a yellow oil. LC-MS (ES^+): m/z 326.20 [MNa^+], t_R = 1.55 min (2.6 minute run).

[1043] Step 2: Synthesis of ethyl 2-{[5-(methylamino)pentyl]oxy}acetate (**L-16**)

[1044] To a stirred solution of ethyl 2-[(5-{[(tert-butoxy)carbonyl](methyl)amino}pentyl)oxy]acetate (**BB**, 240 mg, 0.79 mmol) in dichloromethane (5 mL) was added trifluoroacetic acid (0.5 mL). The resulting solution was stirred at room temperature for 16 hours. The solvents were removed under recued pressure followed by high vacuum pump to give **L-16** (yield: 99%) as a yellow oil. LC-MS (ES^+): m/z 204.20 [MH^+], t_R = 0.56 min (2.0 minute run).

[1045] **L-17: 2-(3-(2-(tosyloxy)ethoxy)propoxy)acetic acid**

[1046]



[1047] Step 1: Synthesis of tert-butyl 2-{3-[2-(benzyloxy)ethoxy]propoxy}acetate (**BD**)

[1048] To a stirred solution of 3-[2-(benzyloxy)ethoxy]propan-1-ol (**BC**, 1.8 g, 8.56 mmol) and tert-butyl 2-bromoacetate (6.6 g, 33.84 mmol, 4.00 equiv) in dichloromethane (40 mL) was added TBAC (2.4 g) and aq. Solution of sodium hydroxide (37%, 40 mL). The resulting mixture was stirred at room temperature overnight. LC-MS indicated formation of the desired product. The

reaction mixture was then extracted with ethyl acetate (150 x 3 mL), the organic layers combined, dried over anhydrous sodium sulfate and concentrated under reduced pressure to give a crude residue, which was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v : v = 1 : 2) to give **BD** (yield: 90%) as a colorless oil. ^{1}H NMR (300 MHz, $CDCl_3$): δ 7.35-7.27 (m, 5H), 4.57 (s, 2H), 3.94 (s, 2H), 3.63-3.57 (m, 8H), 1.96-1.87 (m, 2H), 1.47 (s, 9H); LC-MS (ES^+): *m/z* 347.10 [MNa^+], t_R = 1.72 min (2.6 minute run).

[1049] Step 2: Synthesis of tert-butyl 2-[3-(2-hydroxyethoxy)propoxy]acetate (**BE**)

[1050] To a stirred mixture of tert-butyl 2-{3-[2-(benzyloxy)ethoxy]propoxy}acetate (**BD**, 2.5 g, 7.71 mmol) and palladium on carbon (2.0 g) in methanol (20 mL) under a nitrogen atmosphere was introduced hydrogen gas via a balloon. The resulting mixture was stirred overnight at room temperature under hydrogen gas atmosphere. LC-MS indicated completion of the reaction. The solids were removed by filtration, the solution was concentrated under vacuum to give **BE** (yield: 99%) as a colorless oil. LC-MS (ES^+): m/z 257.10 [MNa^+], t_R = 1.21 min (2.6 minute run).

[1051] Step 3: Synthesis of tert-butyl 2-(3-{2-[(4-methylbenzenesulfonyl)oxy]ethoxy}propoxy)acetate (**BF**)

[1052] To a stirred solution of tert-butyl 2-[3-(2-hydroxyethoxy)propoxy]acetate (**BE**, 1.8 g, 7.68 mmol) in dichloromethane (50 mL) was added 4-toluenesulfonyl chloride (2.2 g, 11.54 mmol), triethylamine (2.33 g, 23.03 mmol) and 4-dimethylaminopyridine (95 mg, 0.78 mmol). The resulting mixture was stirred overnight at room temperature. LC-MS indicated formation of the desired product. The reaction mixture was concentrated under reduced pressure to give a crude residue, which was purified by a flash silica gel chromatography (eluent: ethyl acetate/petroleum ether (v : v = 1 : 2) to give **BF** (yield: 80%) as a yellow oil.^{1}H NMR (400 MHz, $CDCl_3$): δ 7.80 (d, *J* = 8.0 Hz, 2H), 7.34 (d, *J* = 8.4 Hz, 2H), 4.15 (t, *J* = 3.6 Hz, 2H), 3.93 (s, 2H), 3.61 (t, *J* = 3.6 Hz, 2H), 3.55-3.49 (m, 4H), 2.45 (s, 3H), 1.85-1.78 (m, 2H), 1.48 (s, 9H); LC-MS (ES^+): *m/z* 411.00 [MNa^+], t_R = 1.12 min (2.0 minute run).


[1053] Step 4: Synthesis of 2-(3-{2-[(4-methylbenzenesulfonyl)oxy]ethoxy}propoxy)acetic acid (**L-17**)

[1054] To a stirred solution of tert-butyl 2-(3-{2-[(4-methylbenzenesulfonyl)oxy]ethoxy}propoxy)acetate (**BF**, 400 mg, 1.03 mmol) in dichloromethane (3 mL) was added trifluoroacetic acid (1 mL) at room temperature. The resulting solution was stirred at room temperature for 1 hour. LC-MS indicated completion of the reaction.


The reaction mixture was concentrated under reduced pressure to give **L-17** (350 mg) as a yellow oil, which was used for next step without further purifications. LC-MS (ES^+): *m/z* 332.90 [MH^+], t_R = 0.81 min (2.0 minute run).

[1055] Unless otherwise noted, the following intermediates and their analogs (for examples, but not limited to, analogs with substitutions such as halogens) were synthesized according to similar procedures described above for the synthesis of **L-17**, by utilizing corresponding starting materials and reagents.


[1056] L-18: 2-(2-hydroxyethoxy)ethyl 4-methylbenzenesulfonate

[1057] 


[1058] L-19: ethyl 2-(2-(2-(tosyloxy)ethoxy)ethoxy)acetate

[1059] 


[1060] L-20: ethyl 3-(2-(2-(tosyloxy)ethoxy)ethoxy)propanoate

[1061] 


[1062] L-21: ethyl 5-(tosyloxy)pentanoate

[1063] 


[1064] L-22: ethyl 3-(2-(tosyloxy)ethoxy)propanoate

[1065] 


[1066] L-23: ethyl 2-(5-(tosyloxy)pentyloxy)acetate

[1067] 


[1068] L-24: ethyl 3-(5-(tosyloxy)pentyloxy)propanoate

[1069] 


[1070] L-25: 5-hydroxypentyl 4-methylbenzenesulfonate

[1071] 


[1072] L-26: ethyl 2-(5-(tosyloxy)pentyloxy)acetate

[1073] 


[1074] L-27: ethyl 2-(3-(tosyloxy)propoxy)acetate

[1075] 

[1076] **L-28: ethyl 2-(2-(tosyloxy)ethoxy)acetate**


[1077] 

[1078] **L-29: ethyl 2-(4-(2-(tosyloxy)ethoxy)butoxy)acetate**




[1079] **L-29**


[1080] **L-30: 2-(2-(2-hydroxyethoxy)ethoxy)ethyl 4-methylbenzenesulfonate**

[1081] 

[1082] **L-31: 2-((2R,3R)-3-(2-hydroxyethoxy)butan-2-yloxy)ethyl 4-methylbenzenesulfonate**

[1083] 

[1084] **L-32: 1-methyl-4-((1-methylpiperidin-4-yl)methyl)piperazine**

[1085] 

[1086] When referring to the specific exemplary compounds presented herein, the specification uses the terms "example #." For example, compound 1 (Table 2 of Figure 2) is also referred to as Example 1. DC50 (μM) categories (degradation of AR ELISA in LNCaP and/or VCaP cells) of Table 2-7 are as follows: A <1nM; B: 1-10nM; C: 10-100nM; D: >100nM. Dmax categories (degradation of AR-maximum inhibition (%) AR ELISA in LNCaP and/or VCaP cells): A >50%; B <50%.

[1087] **Androgen Receptor ELISA Assay.** Compounds have been evaluated in this assay in LNCaP and/or VCaP cells utilizing similar protocols. The protocols used with VCaP cells are described below. The androgen receptor ELISA assay was performed using PathScan AR Sandwich ELISA (Cell Signaling Catalog#12850) according to the following assay steps:

[1088] VCaP cells are seeded at 40,000 cells/well at a volume of 100 μL/well in VCaP assay medium [Phenol red free RPMI (Gibco Cat#11835-030); 5% Charcoal Stripped (Dextran treated)

FBS (Omega Scientific, Cat#FB-04); 1% penstrep (Life Technologies,Gibco Cat#: 10378-016)] in Corning 3904 plates. The cells are grown for a minimum of 3 days.

[1089] First, cells are dosed with compounds diluted in 0.01% DMSO – use a polypropylene plate avoiding the use of outer columns according to the following protocol: (1)(i) make 1000x stock plate in DMSO; (ii) 20mM stock diluted 1/6.7 with DMSO (5 µL + 28.3 µL DMSO) = 3 mM into row H; (iii) perform serial dilutions in ½ log doses (10 µL of PROTAC + 20 µL DMSO) from row H towards row B. Reserve row A for DMSO; (iv) 7 doses total (final concentration in this 1000x plate will be 3 mM, 1 mM, 333 µM, 111 µM, etc). (2)(i) Make 10x stock plate in media; (ii) transfer 2.5 µL of the 1000x stock to a new 10x stock plate (use 12 channel pipet, start at A (DMSO control) work thru H. When 247.5 µL of media is added to this plate, it will serve as a 10x stock; (iii) make media + 1nM R1881 for making 10x stock plate; (iv) add 247.5 µL of media with 1 nM R1881 to each well of the 10x stock plate, mix.

Then 22 µL of 10x stock is added to cells and incubated for 5 hours. 1x Cell Signaling Cell lysis buffer is made (Catalogue #9803; comes with the kit) - prepare for 50 µL/well. Keep on ice. Media is aspirated, and 100 µL lx cell lysis buffer/well is added. The cells are placed on a shaker located in a cold room for 10 minues and shaken at speek of 7. The lysate mixture is mix and 20µL transferred to 100µl of Diluent in ELISA plate (0.15µg/ml – 0.075µg/ml). The lysate-diluent mixture is store at 4°C overnight on a shaker located in a cold room at speed 5 (gentle swirl).

[1090] The lysate-diluent mixture is shaken for 30 minutes at 37°C. Allow mouse AR antibody, anti-mouse antibody, TMB, and STOP solution to come to room temperature. Make 1x ELISA buffer included in kit, load in reservoir. Run primer program on plate washer. Media from the plates is discarded, the ELISA plate tapped hard on paper towel, and washed 4x 200 µl ELISA wash buffer using a plate washer for the first three washes and an eight channel aspirator for the fourth wash to more thoroughly aspirate the solution.

[1091] Add 100 µL/well mouse AR detection Ab; cover and shake, 37°C for 1 hour; media is discarded from the places, tap the plates on a paper towel, wash 4x 200 µL ELISA wash buffer with a plate washer for the first three washed and an eight channel aspirator for the fourth wash; add 100 µL/well anti-mouse – HRP conjugated Ab (comes with the kit); cover and shake, 37°C for 30 minutes; allow TMB reagent to come to room temperature; discard media from the plate, tap plates on paper towl, wash 4x 200 µL ELISA wash buffer with a plate washer for the first

three washed and an eight channel aspirator for the fourth wash; tap the plates on paper towl; add 100 µL TMB and shake for 2 minutes – while watching color. Add 100 µL Stop solution when light blue color develops. Shake plates and read at 450 nM.

[1092] Progression of prostate cancer in patients treated with anti-androgen therapy usually involves one of several mechanisms of enhanced Androgen Receptor (AR) signaling, including increased intratumoral androgen synthesis, increased AR expression and AR mutations. PROTACs (PROteolysis TArgeting Chimera), which uses bi-functional molecules that simultaneously bind a target of choice and an E3 ligase, cause ubiquitination via induced proximity and degradation of the targeted, pathological protein. As opposed to traditional target inhibition, which is a competitive process, degradation is a progressive process. As such, it is less susceptible to increases in endogenous ligand, target expression, or mutations in the target. Thus this technology seems ideal for addressing the mechanisms of AR resistance in patients with prostate cancer.

[1093] Data is analyzed and plotted using GraphPad Prism software. Compounds described in this application were assayed and c-myc suppression potency is listed in the Table 2-5.

[1094] Apoptosis in VCaP cells. VCaP cells may be cultured in Charcoal Stripped Serum containing media supplemented with 0.1 nM R1881 for 48 hours. The degree of apoptosis can then be ascertained with CaspaseGlo assay (Promega).

[1095] Anti-proliferation in LNCaP F876L. LNCaP cells transduced with an AR construct were cultured in Charcoal Stripped Serum containing media. Indicated doses of enzalutamide or Exemplary Compounds can be added for 7 days. CellTiterGlo reagent (Promega) can be employed to assess proliferation.

[1096] PSA suppression in LNCaP F876L**.** LNCaP F876L cells transduced with an AR construct were cultured in Charcoal Stripped Serum containing media supplemented with 0.1 nM R1881 for 7 days. Secreted PSA in the media was detected by PSA ELISA (Sigma).

[1097] Prostate involution in C57B6 mouse model. 12-week old male C57BL/6 mice are treated with an AR PROTAC and its inactive epimer analog which is unable to bind to cereblon E3 ligase. Compounds are administered for 10 days, upon which the prostates were isolated and weighed.

[1098] Tumor growth inhibition in VCaP xenograft model.

[1099] VcaP Xenograft and Drug Treatment. Three million VCaP cells 75% MatriGel/25% RPMI suspension are implanted into CB17 scid mice subcutaneously. Once the tumors reach about 200-300 mm^3, the mice are surgically castrated, leading to temporary tumor stasis. Allow the mice to recover for at least one week. The mice are then dosed with enzalutamide (PO, QD, 30 mpk) or AR PROTAC (IP, QD, at 30, 10 and 3 mpk) as indicated by oral gavage. Sixteen hours after the last does, the mice are sacrificed and the tumors excised.

[1100] Tissue lysis buffer. Final concentrations: 25 mM HEPES (pH 7.4), 50 mM NaCl, 1% NP-40, 0.1% SDS, protease inhibitor tablets/EDTA free mini (A32955) 1/10ml (if DNA is a problem, protease inhibitor tablets with EDTA is used); if phospho-proteins are observed at end point –a phosphatase inhibitor is added. One µl per mg of tissue is utilized. Protease Inhibitor Cocktail mini is added fresh to the lysis buffer prior to homogenization.

[1101] Tissue lysis. Five mm stainless steel bead in 2 ml tube or 2.5 mm bead for 1.5 ml tubes is added to frozen tumor chunk on dry ice. Bead and tumor is transferred to wet ice. Add 1 µl of tissue lysis buffer per mg of tissue to the tumor and beads mixture. Bead milling of the tissue was performed for for 4 minutes , 25 Hz, 4°C. The beads are removed with a magnet. The lysates are spun for 15 minutes 14,000 x g, 4°C. The total lysate volume is transferred to block plate. The tumor lysate is normalized, usually ~50µg/µl.

[1102] Western blotting. 10 µg/lane are used for LnCAP lysates and 5ug/lane for VCaP lysates. Samples are run at 150 V for 85 minutes. Transfer to nitrocellulose membrane. Block the nitrocellulose membrane for 1 hour at room temprature with 3% BSA. Primary antibody: Cell Signaling antibody AR #5153 1/2000 o.n., mitoC abcam #ab92824 1:1000, ERG antibody Abcam # ab92513 1:1000. Secondary antibody: Cell signaling antibody #7074/7076 1/20000 for 1 hour (Anti-mouse/rabbit HRP). Detect with Femto (THERMO FISHER) or Clarity (BIO-RAD).

[1103] Xenograft data is shown below in Table 8.

Table 8. Xenograft data for select compounds of the present disclosure

Ex. #	PD_AR_castrated_VCaP %AR degraded @10mg_kg @16hr (%)
382	C

378	C
265	C
177	C
124	C
263	C
400	B
134	B
133	B
183	B
293	B
316	B
193	B
377	B
289	B
185	B
41	A
91	A
127	A
178	A
180	A
315	A
359	A
288	A
131	A
179	A
440	A
126	A
310	A
182	A
346	A
132	A
181	A
420	A
270	A
240	A
343	A

426	A
145	A
215	A
114	A
158	A
210	A
234	A
122	A
129	A
247	A
406	A

[1104] AR degradation of PROTAC is E3 ligase dependent. An AR PROTAC is added to LNCaP cells at indicated concentrations for 24 hours in the presence or absence of 10 uM cereblon E3 ligase ligand. (A) AR decradation activity of the AR PROTAC is examined to determine if there is diminished degradation as a result of competition from the cereblon E3 ligase ligand with AR PROTAC in cereblon E3 ligase binding. (B) LNCaP cells are treated with an AR PROTAC and its inactive epimer analog which is unable to bind to cereblon E3 ligase.

[1105] PROTAC prodrug oral pharmacokinetics and PROTAC Subcutaneous phamacokinetics.

[1106] Representative Pharmacokinetic Procedure

[1107] Male CD-1 mice (6-8 weeks old, weighing 20-30 g, 3 per study) with free access to food and water are administered with the test article at 10 mg/kg either by oral gavage or sub-cutaneous injection in the formulation specified in tables 20 and 21, at 10 mL/kg.

[1108] Approximately 0.04 mL blood samples are collected from the dorsal metatarsal vein serially at 0.25, 0.5, 1, 2, 4, 8 and 24 hours timepoints; heparin is used as the anticoagulant. The samples are centrifuged at 4000 g for 5 minutes at 4 °C then stored at -75 °C prior to analysis.

[1109] The plasma samples are analysed via an LC/MS/MS method quantitating for unchanged, administered test article, and/or a derivative species as appropriate. WinNonlin (PhoenixTM) is used for the pharmacokinetic calculations and modeling, to generate parameters such as Cmax and AUC.


[1110] Specific Embodiments

[1111] In certain embodiments, the description provides a compound having a structure selected from the group consisting of Examples 1-625 (see Tables 2-7), a salt, a polymorph, and prodrug thereof. In certain additional embodiments, the description provides a composition comprising at least one of the compounds of Examples 1-625 including a salt, polymorph, and prodrug thereof. In still additional embodiments, the description provides a therapeutic composition comprising at least one of the compounds of Examples 1-625 including a salt, a polymorph, and a prodrug thereof, and a pharmaceutically acceptable carrier.


[1112] An aspect or the present disclosure provides a compound having the structure:

ABM – L – CLM,


wherein ABM is an androgen receptor (AR) binding moiety, L is a chemical linker moiety , CLM is a cereblon E3 ubiquintin ligase binding moiety, wherein the ABM comprises a structure selected from the group consisting of:




ABM-a ;



ABM-b ;



ABM-c ; and



ABM-d ,

wherein:

W^1 is aryl, heteroaryl, bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, S, SO_2, heteroaryl, or aryl;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;


Q is a 3-6 membered ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q is independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);


R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, cyclic, heterocyclic, or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, aryl, heteroaryl, alicyclic, heterocyclic, biheterocyclic, biaryl, or biheteroaryl, each optionally substituted by 1-10 R^{W2};


each R^{W2} is independently H, halo, C_{1-6} alkyl (linear or branched optionally substituted; for example, optionally substituted by 1 or more F), $-OR^{W2A}$, C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OC_{1-3}alkyl (optionally substituted; for example, optionally substituted by 1 or more –F), OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and

R^{W2A} is H, C_{1-6} alkyl (linear, branched), or C_{1-6} heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC_{1-3}alkyl.

[1113] In any aspect or embodiment described herein, W^1 is selected from the group consisting of: 



[1114] In any aspect or embodiment described herein, W^2 is selected from the group consisting of:




[1115] In any aspect or embodiment described herein, the CLM comprises a chemical group derived from an imide, a thioimide, an amide, or a thioamide that binds to the cereblon E3 ubiquitin ligase.


[1116] In any aspect or embodiment described herein, the chemical group is a phthalimido group, or an analog or derivative thereof.

[1117] In any aspect or embodiment described herein, the CLM is thalidomide, lenalidomide, pomalidomide, analogs thereof, isosteres thereof, or derivatives thereof.


[1118] In any aspect or embodiment described herein, the CLM has a chemical structure represented by:




(a) (b)



(c),




(d),



(e),

or



(f),

wherein

W is selected from the group consisting of CH_2, CHR, C=O, SO_2, NH, and N-alkyl;

each X is independently selected from the group consisting of O, S, and H_2;

Y is selected from the group consisting of NH, N-alkyl, N-aryl, N-hetaryl, N-cycloalkyl, N-heterocyclyl, O, and S;

Z is selected from the group consisting of O, S, and H_2;

G and G' are each independently selected from the group consisting of H, alkyl (linear, branched, optionally substituted), OH, R'OCOOR, R'OCONRR", CH_2-heterocyclyl optionally substituted with R', and benzyl optionally substituted with R';

Q_1, Q_2, Q_3, and Q_4 represent a carbon C optionally substituted with a group independently selected from R', N or N-oxide;

A is selected from the group consisting of H, alkyl, cycloalkyl, Cl and F;

R is selected from the group consisting of -CONR'R", -OR', -NR'R", -SR', -SO_2R', -SO_2NR'R", -CR'R"-, -CR'NR'R"-, -aryl, -hetaryl, -alkyl (linear, branched, optionally substituted), -cycloalkyl, -heterocyclyl, -P(O)(OR')R", -P(O)R'R", -OP(O)(OR')R", -OP(O)R'R", -Cl, -F, -Br, -I, -CF_3, -CN, -NR'SO_2NR'R", -NR'CONR'R", -CONR'COR", -NR'C(=N-CN)NR'R", -C(=N-CN)NR'R", -NR'C(=N-CN)R", -NR'C(=C-

NO_2)NR'R", -SO_2NR'COR", -NO_2, -CO_2R', -C(C=N-OR')R", -CR'=CR'R", -CCR', -S(C=O)(C=N-R')R", -SF_5, -R'NR'R", (-R'O)$_n$R", or and -OCF_3;

R' and R" are each independently selected from the group consisting of a bond, H, alkyl (linear, branched), cycloalkyl, aryl, hetaryl, heterocyclyl, or –C(=O)R, each of which is optionally substituted;

〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and


R_n is a functional group or an atom,

wherein n is an integer from 1-10 (e.g., 1-4, 1, 2 , 3, 4, 5, 6, 7, 8, 9, or 10), and wherein

when n is 1, R_n is modified to be covalently joined to the linker group (L), and

when n is 2, 3, or 4, then one R_n is modified to be covalently joined to the linker group (L), and any other R_n is optionally modified to be covalently joined to a ABM, a CLM, a second CLM having the same chemical structure as the CLM, a CLM', a second linker, or any multiple or combination thereof.

[1119] In any aspect or embodiment described herein, the CLM or ULM has a chemical structure represented by:



wherein:

W is independently selected from the group CH2, C=O, NH, and N-alkyl;


R is independently selected from a H, methyl, alkyl;

〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and


Rn comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to a ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof.

[1120] Another aspect of the present disclosure provides a bifunctional compound comprising the chemical structure: ABM-L-CLM, wherein ABM is an androgen receptor (AR) binding moiety, L is absent (a bond) or a chemical linker, and CLM is a cereblon E3 ubiquitin


ligase binding moiety is a chemical group derived from an imide, a thioimide, an amide, or a thioamide,wherein the ABM comprises a structure selected from the group consisting of:




ABM-a ;



ABM-b ;



ABM-c ; and



ABM-d ,

wherein:

W^1 is aryl, heteroaryl, bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, S;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;


Q is a 3-6 membered ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q,is independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, cyclic, heterocyclic , or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);


W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, aryl, heteroaryl, alicyclic, heterocyclic, biheterocyclic, biaryl, or biheteroaryl, each optionally substituted by 1-10 R^{W2};

each R^{W2} is independently H, halo, C_{1-6} alkyl (linear or branched optionally substituted; for example, optionally substituted by 1 or more F), -OR^{W2A} , C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, C_{1-6} alkyl (optionally substituted), C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OC_{1-3}alkyl (optionally substituted; for example, optionally substituted by 1 or more –F) , OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and

RW2A is H, C1-6 alkyl (linear, branched), or C1-6 heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC1-3alkyl..

[1121] In any aspect or embodiment described herein, W^1 is selected from the group consisting of:  .

[1122] In any aspect or embodiment described herein, W^2 is selected from the group consisting of:


 .

[1123] In any aspect or embodiment described herein, CLM comprises a chemical group derived from an imide, a thioimide, an amide, or a thioamide that binds cereblon E3 ubiquitin ligase.


[1124] In any aspect or embodiment described herein, the chemical group is a phthalimido group, or an analog or derivative thereof.

[1125] In any aspect or embodiment described herein, the CLM is thalidomide, lenalidomide, pomalidomide, analogs thereof, isosteres thereof, or derivatives thereof.


[1126] In any aspect or embodiment described herein, the CLM has a chemical structure represented by:




(a),




(b),



(c),




(d),



(e),

or



(f),

wherein

W is selected from the group consisting of CH_2, CHR, C=O, SO_2, NH, and N-alkyl;

each X is independently selected from the group consisting of O, S, and H_2;

Y is selected from the group consisting of CH_2, -C=CR', NH, N-alkyl, N-aryl, N-hetaryl, N-cycloalkyl, N-heterocyclyl, O, and S;

Z is selected from the group consisting of O, S, and H_2;

G and G' are independently selected from the group consisting of H, alkyl (linear or branched optionally substituted; for example optionally substituted with R'), OH, R'OCOOR,

R'OCONRR", CH_2-heterocyclyl optionally substituted with R', and benzyl optionally substituted with R';

Q_1, Q_2, Q_3, and Q_4 are independently a carbon C optionally substituted with a group independently selected from R', N or N-oxide;

A is independently selected from the group H, alkyl, cycloalkyl, Cl and F;

R comprises -CONR'R", -OR', -NR'R", -SR', -SO_2R', -SO_2NR'R", -CR'R"-, -CR'NR'R"-, (-CR'O)$_n$R", -aryl, -hetaryl, -alkyl, -cycloalkyl, -heterocyclyl, -P(O)(OR')R", -P(O)R'R", -OP(O)(OR')R", -OP(O)R'R", -Cl, -F, -Br, -I, -CF_3, -CN, -NR'SO_2NR'R", -NR'CONR'R", -CONR'COR", -NR'C(=N-CN)NR'R", -C(=N-CN)NR'R", -NR'C(=N-CN)R", -NR'C(=C-NO_2)NR'R", -SO_2NR'COR", -NO_2, -CO_2R', -C(C=N-OR')R", -CR'=CR'R", -CCR', -S(C=O)(C=N-R')R", -SF_5 or -OCF_3;

R' and R" are independently selected from the group consisting of a bond, H, alkyl, cycloalkyl, aryl, hetaryl, heterocyclyl, -C(=O)R, each of which is optionally substituted;

〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and


R_n comprises a functional group or an atom,

wherein n is an integer from 1-10 (e.g., 1-4, 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10), and wherein

when n is 1, R_n is modified to be covalently joined to the linker group (L), and

when n is 2, 3, or 4, then one R_n is modified to be covalently joined to the linker group (L), and any other R_n is optionally modified to be covalently joined to a ABM, a CLM, a second CLM having the same chemical structure as the CLM, a CLM', a second linker, or any multiple or combination thereof.

[1127] In any aspect or embodiment described herein, the CLM or ULM has a chemical structure represented by:



wherein:

W is independently selected from the group CH2, C=O, NH, and N-alkyl;

R is independently selected from a H, methyl, alkyl;

~~~ represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and

Rn comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to a ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof.

[1128] In any aspect or embodiment described herein, the linker group L is a group comprises a chemical structural unit represented by the formula:

$$-A_q-$$

wherein A_q is a group coupled to at least one of CLM, ABM, or both; and

q is an integer greater than or equal to 1,

wherein Aq is selected from the group consisting of, a bond, $CR^{L1}R^{L2}$, O, S, SO, SO_2, NR^{L3}, SO_2NR^{L3}, $SONR^{L3}$, $CONR^{L3}$, $NR^{L3}CONR^{L4}$, $NR^{L3}SO_2NR^{L4}$, CO, $CR^{L1}=CR^{L2}$, C≡C, $SiR^{L1}R^{L2}$, $P(O)R^{L1}$, $P(O)OR^{L1}$, $NR^{L3}C(=NCN)NR^{L4}$, $NR^{L3}C(=NCN)$, $NR^{L3}C(=CNO_2)NR^{L4}$, C3-11cycloalkyl optionally substituted with 0-6 RL1 and/or RL2 groups, C5-13 spirocycloalkyl optionally substituted with 0-9 R^{L1} and/or R^{L2} groups, C3-11heterocyclyl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, C5-13 spiroheterocycloalkyl optionally substituted with 0-8 R^{L1} and/or R^{L2} groups, aryl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, heteroaryl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, where R^{L1} or R^{L2}, each independently are optionally linked to other groups to form cycloalkyl and/or heterocyclyl moiety, optionally substituted with 0-4 R^{L5} groups;

R^{L1}, R^{L2}, R^{L3}, R^{L4} and R^{L5} are, each independently, H, halo, C_{1-8}alkyl, OC_{1-8}alkyl, SC_{1-8}alkyl, NHC_{1-8}alkyl, $N(C_{1-8}alkyl)_2$, C_{3-11}cycloalkyl, aryl, heteroaryl, C_{3-11}heterocyclyl, OC_{1-8}cycloalkyl, SC_{1-8}cycloalkyl, NHC_{1-8}cycloalkyl, $N(C_{1-8}cycloalkyl)_2$, $N(C_{1-8}cycloalkyl)(C_{1-8}alkyl)$, OH, NH_2, SH, SO_2C_{1-8}alkyl, $P(O)(OC_{1-8}alkyl)(C_{1-8}alkyl)$, $P(O)(OC_{1-8}alkyl)_2$, CC-C_{1-8}alkyl, CCH, $CH=CH(C_{1-8}alkyl)$, $C(C_{1-8}alkyl)=CH(C_{1-8}alkyl)$, $C(C_{1-8}alkyl)=C(C_{1-8}alkyl)_2$, $Si(OH)_3$, $Si(C_{1-8}alkyl)_3$, $Si(OH)(C_{1-8}alkyl)_2$, COC_{1-8}alkyl, CO_2H, halogen, CN, CF_3, CHF_2, CH_2F, NO_2, SF_5, SO_2NHC_{1-8}alkyl, $SO_2N(C_{1-8}alkyl)_2$, $SONHC_{1-8}$alkyl, $SON(C_{1-8}alkyl)_2$, $CONHC_{1-8}$alkyl, $CON(C_{1-8}alkyl)_2$, $N(C_{1-8}alkyl)CONH(C_{1-8}alkyl)$, $N(C_{1-8}alkyl)CON(C_{1-8}alkyl)_2$, $NHCONH(C_{1-8}alkyl)$, $NHCON(C_{1-8}alkyl)_2$, $NHCONH_2$, $N(C_{1-8}alkyl)SO_2NH(C_{1-8}alkyl)$, $N(C_{1-8}alkyl)\ SO_2N(C_{1-8}alkyl)_2$, NH $SO_2NH(C_{1-8}alkyl)$, NH $SO_2N(C_{1-8}alkyl)_2$, NH SO_2NH_2.
~~~

[1129] In any aspect or embodiment described herein, the linker (L) comprises a group represented by a general structure selected from the group consisting of:

-N(R)-$(CH2)_m$-$O(CH2)_n$-$O(CH2)_o$-$O(CH2)_p$-$O(CH2)_q$-$O(CH2)_r$-OCH2-,


-O-$(CH2)_m$-$O(CH2)_n$-$O(CH2)_o$-$O(CH2)_p$-$O(CH2)_q$-$O(CH2)_r$-OCH2-,


-O-$(CH2)_m$-$O(CH2)_n$-$O(CH2)_o$-$O(CH2)_p$-$O(CH2)_q$-$O(CH2)_r$-O-;


-N(R)-$(CH2)_m$-$O(CH2)_n$-$O(CH2)_o$-$O(CH2)_p$-$O(CH2)_q$-$O(CH2)_r$-O-;


-$(CH2)_m$-$O(CH2)_n$-$O(CH2)_o$-$O(CH2)_p$-$O(CH2)_q$-$O(CH2)_r$-O-;


-$(CH2)_m$-$O(CH2)_n$-$O(CH2)_o$-$O(CH2)_p$-$O(CH2)_q$-$O(CH2)_r$-OCH2-;



;



;



;



;


;


;



;


; and



, wherein m, n, o, p, q, and r are each independently 0, 1, 2, 3, 4, 5, 6, with the provision that when the number is zero, there is no N-O or O-O bond, R is selected from the group H, methyl or ethyl, and X is selected from the group H or F.


[1130] In any aspect or embodiment described herein, the linker (L) comprises a group represented by a general structure:





,


wherein m can be 2, 3, 4, 5.


[1131] In any aspect or embodiment described herein, the linker (L) comprises a group represented by a general structure selected from the group consisting of:









X = H, F


X = H, F












wherein n and m are each independently 0, 1, 2, 3, 4, 5, 6, and X is H, or F.

[1132] In any aspect or embodiment described herein, linker (L) is selected from the group consisting of:








[1133] In any aspect or embodiment described herein, L is a polyethylene group optionally substituted with aryl or phenyl comprising from 1 to 10 ethylene glycol units.

[1134] In any aspect or embodiment described herein, the compound comprises multiple LCMs, multiple ABMs, multiple linkers or any combinations thereof.


[1135] In any aspect or embodiment described herein, the compound is a member selected from the group consisting of Examples 1-625 (e.g., a compound from Tables 2-7), a salt, a polymorph, isotopic derivative, and a prodrug thereof.

[1136] In any aspect or embodiment described herein, the compound is selected from Tables 2-7 (i.e., Exemplary Compounds 1-625).


[1137] In any aspect or embodiment described herein, the ABM is selected from the group consisting of:




,




,




,




,



,



, and



,

wherein:

R^{Q2} is a H, halogen, CH_3 or CF_3;

RQ3 is a H, halo, hydroxyl, nitro, CN, C≡CH, C1-6 alkyl (linear, branched, optionally substituted by 1 or more halo, C1-6 alkoxyl), C1-6 alkoxyl (linear, branched, optionally substituted by 1 or more halo), C2-6 alkenyl, C2-6 alkynyl, or CF_3;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, heteroaryl, or aryl;

R^{Y1}, R^{Y2} are each independently H, or C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl, cyclic, or heterocyclic);


R^Q each independently is H, C_1-C_6 alkyl (linear, branched, optionally substituted by 1 or more halo, or C_{1-6} alkoxyl), or two R^Q together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms; and


X is N or C.


[1138] In any aspect or embodiment described herein, each R^Q is independently H or CH_3.


[1139] In any aspect or embodiment described herein, R^{Q3} is CN;


[1140] In any aspect or embodiment described herein, the ABM is selected from the group consisting of:




























; and










.










;

;


;


;


;

;

;

;

;

;

;






[1141] A further aspect of the present disclosure provides a composition comprising an effective amount of a compound or bifuctional compound of any of claims 1-25, and a pharmaceutically acceptable carrier.

[1142] In any aspect or embodiment described herein, the composition further comprises at least one additional bioactive agent.

[1143] In any aspect or embodiment described herein, the bioactive agent is an anti-cancer agent.

[1144] An additional aspect of the present disclosure provides a therapeutic composition comprising an effective amount of at least two different compounds as described in any of claims 1-25.

[1145] Another aspect of the present disclosure provides a method of treating a disease or disorder in a subject comprising the steps of administering a composition comprising a pharmaceutically acceptable carrier and an effective amount of a compound or bifunctional compound as described in any of claims 1-25 to a subject in need thereof, wherein the compound is effective in treating or ameliorating at least one symptom of the disease or disorder.

[1146] In any aspect or embodiment described herein, the disease or disorder is cancer or Kennedy's Disease or both.

[1147] In any aspect or embodiment described herein, the cancer is prostate cancer.

[1148] In any aspect or embodiment described herein, the composition further comprises an effective amount of at least one additional anti-cancer agent.

[1149] The contents of all references, patents, pending patent applications and published patents, cited throughout this application are hereby expressly incorporated by reference. Except in the Examples, or where otherwise explicitly indicated, all numerical quantities in this description specifying amounts of materials, and the like, are to be understood as modified by the word "about". It is to be understood that the upper and lower amount, range, and ratio limits set forth herein may be independently combined. Similarly, the ranges and amounts for each element of the disclosure can be used together with ranges or amounts for any of the other elements.

[1150] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments and methods described herein. Such equivalents are intended to be encompassed by the scope of the following claims.

[1151] It is understood that the detailed examples and embodiments described herein are given by way of example for illustrative purposes only, and are in no way considered to be limiting to the disclosure. Various modifications or changes in light thereof will be suggested to persons skilled in the art and are included within the spirit and purview of this application and are considered within the scope of the appended claims. For example, the relative quantities of the ingredients may be varied to optimize the desired effects, additional ingredients may be added, and/or similar ingredients may be substituted for one or more of the ingredients described. Additional advantageous features and functionalities associated with the systems, methods, and processes of the present disclosure will be apparent from the appended claims. Moreover, those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the disclosure described herein. Such equivalents are intended to be encompassed by the following claims.


CLAIMS

What Is Claimed Is:


1. A compound having the structure:

ABM – L – CLM,


wherein ABM is an androgen receptor (AR) binding moiety, L is a chemical linker moiety , CLM is a cereblon E3 ubiquintin ligase binding moiety, wherein the ABM comprises a structure selected from the group consisting of:




ABM-a ;



ABM-b ;



ABM-c ; and



ABM-d ,

wherein:

W^1 is aryl, heteroaryl, bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, S, SO2, heteroaryl, or aryl;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;

Q is a 3-6 membered ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q,is independently H, C_{1-6} alkyl (linear, branched, optionally substituted, for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, or 2 R^Q

groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);


R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, cyclic, heterocyclic or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, aryl, heteroaryl, alicyclic, heterocyclic, biheterocyclic, biaryl, or biheteroaryl,each optionally substituted by 1-10 R^{W2};

each R^{W2} is independently H, halo, C_{1-6} alkyl (linear or branched optionally substituted; for example, optionally substituted by 1 or more F), -OR^{W2A} , C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, C_{1-6} alkyl (optionally substituted), , heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OC_{1-3}alkyl (optionally substituted; for example, optionally substituted by 1 or more –F) , OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and

R^{W2A} is H, C_{1-6} alkyl (linear, branched), or C_{1-6} heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC_{1-3}alkyl.





2. The compound of claim 1, wherein W^1 is selected from the group consisting of:



and

.

3. The compound of claim 2, wherein W^2 is selected from the group consisting of:



; 
; 
; and 
.

4. The compound of any of claims 1-3, wherein the CLM comprises a chemical group derived from an imide, a thioimide, an amide, or a thioamide that binds to the cereblon E3 ubiquitin ligase.


5. The compound of claim 4, wherein the chemical group is a phthalimido group, or an analog or derivative thereof.

6. The compound of any of claims 1-3, wherein the CLM is thalidomide, lenalidomide, pomalidomide, analogs thereof, isosteres thereof, or derivatives thereof.


7. The compound of any of claims 1-3, wherein the CLM has a chemical structure represented by:




(a),




(b),



(c),




(d),



(e),

or



(f),

wherein

W is selected from the group consisting of CH_2, CHR, C=O, SO_2, NH, and N-alkyl;

each X is independently selected from the group consisting of O, S, and H_2;

Y is selected from the group consisting of NH, N-alkyl, N-aryl, N-hetaryl, N-cycloalkyl, N-heterocyclyl, O, and S;

Z is selected from the group consisting of O, S, and H_2;

G and G' are each independently selected from the group consisting of H, optionally substituted linear or branched alkyl, OH, R'OCOOR, R'OCONRR", CH_2-heterocyclyl optionally substituted with R', and benzyl optionally substituted with R';

Q_1, Q_2, Q_3, and Q_4 are each independently a carbon C optionally substituted with a group independently selected from R', N or N-oxide;

A is selected from the group consisting of H, alkyl, cycloalkyl, Cl and F;

R is selected from the group consisting of -CONR'R", -OR', -NR'R", -SR', -SO_2R', -SO_2NR'R", -CR'R"-, -CR'NR'R"-, -aryl, -hetaryl, -optionally substituted linear or branched alkyl, -cycloalkyl, -heterocyclyl, -P(O)(OR')R", -P(O)R'R", -OP(O)(OR')R", -OP(O)R'R", -Cl, -

F, -Br, -I, -CF_3, -CN, -NR'SO_2NR'R", -NR'CONR'R", -CONR'COR", -NR'C(=N-CN)NR'R", -C(=N-CN)NR'R", -NR'C(=N-CN)R", -NR'C(=C-NO_2)NR'R", -SO_2NR'COR", -NO_2, -CO_2R', -C(C=N-OR')R", -CR'=CR'R", -CCR', -S(C=O)(C=N-R')R", -SF_5, -R'NR'R", (-R'O)$_n$R", or and -OCF_3;

R' and R" are each independently selected from the group consisting of a bond, H, optionally substituted linear or branched alkyl, cycloalkyl, aryl, hetaryl, heterocyclyl, or –C(=O)R, each of which is optionally substituted;

〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and


R_n is a functional group or an atom,

wherein n is an integer from 1-10, and wherein

when n is 1, R_n is modified to be covalently joined to the linker group (L), and

when n is 2, 3, or 4, then one R_n is modified to be covalently joined to the linker group (L), and any other R_n is optionally modified to be covalently joined to a ABM, a CLM, a second CLM having the same chemical structure as the CLM, a CLM', a second linker, or any multiple or combination thereof.

8. The compound of any of claims 1-3, or 7, wherein the CLM or ULM has a chemical structure represented by:



wherein:


W is independently selected from the group CH2, C=O, NH, and N-alkyl;

R is independently selected from a H, methyl, alkyl;


〰 represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and

Rn comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to a ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof.


9. A bifunctional compound comprising the chemical structure: ABM-L-CLM, wherein ABM is an androgen receptor (AR) binding moiety, L is absent (a bond) or a chemical linker, and CLM is a cereblon E3 ubiquitin ligase binding moiety is a chemical group derived from an imide, a thioimide, an amide, or a thioamide,wherein the ABM comprises a structure selected from the group consisting of:




ABM-a ;



ABM-b ;



ABM-c ; and



ABM-d ,

wherein:

W^1 is aryl, heteroaryl, bicyclic, or biheterocyclic, each independently substituted by 1 or more H, halo, hydroxyl, nitro, CN, C≡CH, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), C_{1-6} alkoxyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo), C_{2-6} alkenyl, C_{2-6} alkynyl, or CF_3;

Y^1, Y^2 are each independently NR^{Y1}, O, S;

Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, C=S, SO, SO_2, heteroaryl, or aryl;


Q is a 3-6 membered ring with 0-4 heteroatoms, optionally substituted with 0-6 R^Q, each R^Q,is independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, or 2 R^Q groups taken together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);


R^1, R^2, R^a, R^b, R^{Y1}, R^{Y2} are each independently H, C_{1-6} alkyl (linear, branched, optionally substituted; for example, optionally substituted by 1 or more halo, C_{1-6} alkoxyl), halogen, C_{1-6} alkoxy, cyclic, heterocyclic, or R^1, R^2 together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms);

W^2 is a bond, C_{1-6} alkyl, C_{1-6} heteroalkyl, O, aryl, heteroaryl, alicyclic, heterocyclic, biheterocyclic, biaryl, or biheteroaryl, each optionally substituted by 1-10 R^{W2};

each R^{W2} is independently H, halo, C_{1-6} alkyl (optionally substituted by 1 or more F), C_{3-6} cycloalkyl, C_{4-6} cycloheteroalkyl, C_{1-6} alkyl (optionally substituted), C_{1-6} alicyclic (optionally substituted), heterocyclic (optionally substituted), aryl (optionally substituted), or heteroaryl (optionally substituted), bicyclic hereoaryl or aryl, OC_{1-3}alkyl (optionally substituted; for example, optionally substituted by 1 or more –F), OH, NH_2, $NR^{Y1}R^{Y2}$, CN; and


R^{W2A} is H, C1-6 alkyl (linear, branched), or C1-6 heteroalkyl (linear, branched), each optionally substituted by a cycloalkyl, cycloheteroalkyl, aryl, heterocyclic, heteroaryl, halo, or OC1-3alkyl.

10. The compound or bifunctional compound of any fo claims 1-9, wherein W^1 is selected from the group consisting of: 

; ; ;


; .

11. The compound or bifunctional compound of any of claims 1-10, wherein W^2 is selected from the group consisting of:




12. The bifunctional compound of any of claims 9-11, wherein CLM comprises a chemical group derived from an imide, a thioimide, an amide, or a thioamide that binds cereblon E3 ubiquitin ligase.


13. The bifunctional compound of claim 12, wherein the chemical group is a phthalimido group, or an analog or derivative thereof.

14. The bifunctional compound of any of claims 9-11, wherein the CLM is thalidomide, lenalidomide, pomalidomide, analogs thereof, isosteres thereof, or derivatives thereof.


15. The bifunctional compound of any of claims 9-11, wherein the CLM has a chemical structure represented by:




(a)




(b)



(c)




(d)



(e)

or



(f),

wherein

W is selected from the group consisting of CH_2, CHR, C=O, SO_2, NH, and N-alkyl;

each X is independently selected from the group consisting of O, S, and H_2;

Y is selected from the group consisting of CH_2, -C=CR', NH, N-alkyl, N-aryl, N-hetaryl, N-cycloalkyl, N-heterocyclyl, O, and S;

Z is selected from the group consisting of O, S, and H_2;

G and G' are independently selected from the group consisting of H, optionally substituted linear or branched alkyl (linear or brached optionally substituted; for example optionally substituted with R'), OH, R'OCOOR, R'OCONRR", CH_2-heterocyclyl optionally substituted with R', and benzyl optionally substituted with R';

Q_1, Q_2, Q_3, and Q_4 are independently a carbon C optionally substituted with a group independently selected from R', N or N-oxide;

A is independently selected from the group H, alkyl, cycloalkyl, Cl and F;

R comprises -CONR'R", -OR', -NR'R", -SR', -SO_2R', -SO_2NR'R", -CR'R"-, -CR'NR'R"-, (-CR'O)$_n$R", -aryl, -hetaryl, -optionally substituted linear or branched alkyl, -cycloalkyl, -heterocyclyl, -P(O)(OR')R", -P(O)R'R", -OP(O)(OR')R", -OP(O)R'R", -Cl, -F, -Br, -I, -CF_3, -CN, -NR'SO_2NR'R", -NR'CONR'R", -CONR'COR", -NR'C(=N-CN)NR'R", -C(=N-CN)NR'R", -NR'C(=N-CN)R", -NR'C(=C-NO_2)NR'R", -SO_2NR'COR", -NO_2, -CO_2R', -C(C=N-OR')R", -CR'=CR'R", -CCR', -S(C=O)(C=N-R')R", -SF_5, -R'NR'R", (-R'O)$_n$R", or -OCF_3;

R' and R" are independently selected from the group consisting of a bond, H, optionally substituted linear or branched alkyl, cycloalkyl, aryl, hetaryl, heterocyclyl, -C(=O)R, each of which is optionally substituted;

~~~ represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and


R_n comprises a functional group or an atom,

wherein n is an integer from 1-10, and wherein

when n is 1, R_n is modified to be covalently joined to the linker group (L), and

when n is 2, 3, or 4, then one R_n is modified to be covalently joined to the linker group (L), and any other R_n is optionally modified to be covalently joined to a ABM, a CLM, a second CLM having the same chemical structure as the CLM, a CLM', a second linker, or any multiple or combination thereof.

16. The compound of any of claims 9-11 or 15, wherein the CLM or ULM has a chemical structure represented by:



wherein:

W is independently selected from the group CH2, C=O, NH, and N-alkyl;

R is independently selected from a H, methyl, alkyl;

~~~ represents a bond that may be stereospecific ((R) or (S)) or non-stereospecific; and

Rn comprises 1-4 independently selected functional groups or atoms, and optionally, one of which is modified to be covalently joined to a ABM, a chemical linker group (L), a ULM, CLM (or CLM') or combination thereof.

17. The compound of any of claims 1-8 or the bifunctional compound of any of claim 9-16, wherein the linker group L is a group comprises a chemical structural unit represented by the formula:

$$-A_q-$$

wherein A_q is a group coupled to at least one of CLM, ABM, or both; and

q is an integer greater than or equal to 1,

wherein Aq is selected from the group consisting of, a bond, $CR^{L1}R^{L2}$, O, S, SO, SO_2, NR^{L3}, SO_2NR^{L3}, $SONR^{L3}$, $CONR^{L3}$, $NR^{L3}CONR^{L4}$, $NR^{L3}SO_2NR^{L4}$, CO, $CR^{L1}=CR^{L2}$, C≡C, $SiR^{L1}R^{L2}$, $P(O)R^{L1}$, $P(O)OR^{L1}$, $NR^{L3}C(=NCN)NR^{L4}$, $NR^{L3}C(=NCN)$, $NR^{L3}C(=CNO_2)NR^{L4}$, C3-11cycloalkyl optionally substituted with 0-6 RL1 and/or RL2 groups, C5-13 spirocycloalkyl optionally substituted with 0-9 R^{L1} and/or R^{L2} groups, C3-11heterocyclyl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, C5-13 spiroheterocycloalkyl optionally substituted with 0-8 R^{L1} and/or R^{L2} groups, aryl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, heteroaryl optionally substituted with 0-6 R^{L1} and/or R^{L2} groups, where R^{L1} or R^{L2}, each independently are optionally linked to other groups to form cycloalkyl and/or heterocyclyl moiety, optionally substituted with 0-4 R^{L5} groups;

R^{L1}, R^{L2}, R^{L3}, R^{L4} and R^{L5} are, each independently, H, halo, C_{1-8}alkyl, OC_{1-8}alkyl, SC_{1-8}alkyl, NHC_{1-8}alkyl, $N(C_{1-8}alkyl)_2$, C_{3-11}cycloalkyl, aryl, heteroaryl, C_{3-11}heterocyclyl, OC_{1-8}cycloalkyl, SC_{1-8}cycloalkyl, NHC_{1-8}cycloalkyl, $N(C_{1-8}cycloalkyl)_2$, $N(C_{1-8}cycloalkyl)(C_{1-8}alkyl)$, OH, NH_2, SH, SO_2C_{1-8}alkyl, $P(O)(OC_{1-8}alkyl)(C_{1-8}alkyl)$, $P(O)(OC_{1-8}alkyl)_2$, CC-C_{1-8}alkyl, CCH, $CH=CH(C_{1-8}alkyl)$, $C(C_{1-8}alkyl)=CH(C_{1-8}alkyl)$, $C(C_{1-8}alkyl)=C(C_{1-8}alkyl)_2$, $Si(OH)_3$, $Si(C_{1-8}alkyl)_3$, $Si(OH)(C_{1-8}alkyl)_2$, COC_{1-8}alkyl, CO_2H, halogen, CN, CF_3, CHF_2, CH_2F, NO_2, SF_5, SO_2NHC_{1-8}alkyl, $SO_2N(C_{1-8}alkyl)_2$, $SONHC_{1-8}$alkyl, $SON(C_{1-8}alkyl)_2$, $CONHC_{1-8}$alkyl, $CON(C_{1-8}alkyl)_2$, $N(C_{1-8}alkyl)CONH(C_{1-8}alkyl)$, $N(C_{1-8}alkyl)CON(C_{1-8}alkyl)_2$, $NHCONH(C_{1-8}alkyl)$, $NHCON(C_{1-8}alkyl)_2$, $NHCONH_2$, $N(C_{1-8}alkyl)SO_2NH(C_{1-8}alkyl)$, $N(C_{1-8}alkyl)\ SO_2N(C_{1-8}alkyl)_2$, NH $SO_2NH(C_{1-8}alkyl)$, NH $SO_2N(C_{1-8}alkyl)_2$, NH SO_2NH_2.

18. The compound or bifunctional compound according to any of claims 1-17, wherein linker (L) comprises a group represented by a general structure selected from the group consisting of:

$-N(R)-(CH2)_m-O(CH2)_n-O(CH2)_o-O(CH2)_p-O(CH2)_q-O(CH2)_r-OCH2-$,


$-O-(CH2)_m-O(CH2)_n-O(CH2)_o-O(CH2)_p-O(CH2)_q-O(CH2)_r-OCH2-$,


$-O-(CH2)_m-O(CH2)_n-O(CH2)_o-O(CH2)_p-O(CH2)_q-O(CH2)_r-O-$;


$-N(R)-(CH2)_m-O(CH2)_n-O(CH2)_o-O(CH2)_p-O(CH2)_q-O(CH2)_r-O-$;


$-(CH2)_m-O(CH2)_n-O(CH2)_o-O(CH2)_p-O(CH2)_q-O(CH2)_r-O-$;


-(CH2)$_m$-O(CH2)$_n$-O(CH2)$_o$-O(CH2)$_p$-O(CH2)$_q$-O(CH2)$_r$-OCH2-;



;



;



;



;



;



;













;



; and



, wherein m, n, o, p, q, and r are each independently 0, 1, 2, 3, 4, 5, 6, with the provision that when the number is zero, there is no N-O or O-O bond, R is selected from the group H, methyl or ethyl, and X is selected from the group H and F;






















X = H, F


X = H, F












wherein n and m are each independently 0, 1, 2, 3, 4, 5, 6, and X is H, or F..


19. The compound or bifunctional compound according to any of claims 1-17, wherein linker (L) is selected from the group consisting of:







20. The compound or bifunctional compound according to any of claims 1-17, wherein the linker (L) comprises a group represented by a general structure:



,

wherein m can be 2, 3, 4, 5.


21. The compound or bifunctional compound according to any of claims 1-17, wherein L is a polyethylene group optionally substituted with aryl or phenyl comprising from 1 to 10 ethylene glycol units.

22. The compound or bifunctional compound according to any of claims 1-21, wherein the compound comprises multiple LCMs, multiple ABMs, multiple linkers or any combinations thereof.


23. The compound or bifunctional compound according to any of claims 1-21, wherein the compound is a member selected from the group consisting of Examples 1-625 (Tables 2-7), a salt, a polymorph, isotopic derivative, and a prodrug thereof.

24. The compound or bifunctional compound according to any of claims 1-21, wherein the compound is selected from Tables 2-7 (i.e., Exemplary Compounds 1-625).


25. The compound or bifunctional compound according to any of claims 1-21, wherein the ABM is selected from the group consisting of:




,




,




,




,



,



, and



,

wherein:

R^{Q2} is a H, halogen, CH_3 or CF_3; and

RQ3 is a H, halo, hydroxyl, nitro, CN, C≡CH, C1-6 alkyl (linear, branched, optionally substituted by 1 or more halo, C1-6 alkoxyl), C1-6 alkoxyl (linear, branched, optionally substituted by 1 or more halo), C2-6 alkenyl, C2-6 alkynyl, or CF_3;


Y^3, Y^4, Y^5 are each independently a bond, O, NR^{Y2}, $CR^{Y1}R^{Y2}$, C=O, heteroaryl, or aryl;













R^{Y1}, R^{Y2} are each independently H, or C_{1-6} alkyl (linear, branched, optionally substituted by 1 or more halo, C_{1-6} alkoxyl, cyclic, or heterocyclic);


R^Q each independently is H, C_1-C_6 alkyl (linear, branched, optionally substituted by 1 or more halo, or C_{1-6} alkoxyl), or two R^Q together with the atom they are attached to, form a 3-8 membered ring system containing 0-2 heteroatoms; and


X is N or C.


26. The compound or bifunctional compound according to any of claims 1-21, wherein the ABM is selected from the group consisting of:












 ;
 ;
 ;
 ;
 ;
 ;
 ;
 ;
 ;
 ;
 ; and
 .









;

; and

,

,

,

;

;






27. A composition comprising an effective amount of a compound or bifuctional compound of any of claims 1-26, and a pharmaceutically acceptable carrier.

28. The composition of claim 27, wherein the composition further comprises at least one additional bioactive agent.

29. The composition of claim 28, wherein the bioactive agent is an anti-cancer agent.

30. A therapeutic composition comprising an effective amount of at least two different compounds as described in any of claims 1-26.

31. A method of treating a disease or disorder in a subject comprising the steps of administering a composition comprising a pharmaceutically acceptable carrier and an effective amount of a compound or bifunctional compound as described in any of claims 1-26 to a subject in need thereof, wherein the compound is effective in treating or ameliorating at least one symptom of the disease or disorder.

32. The method according to claim 31, wherein the disease or disorder is cancer or Kennedy's Disease or both.

33. The method according to claim 32, wherein the cancer is prostate cancer.

34. The method according to any of claims 31-33, wherein the composition further comprises an effective amount of at least one additional anti-cancer agent.

FIG. 1A



ARM
L
CLM

FIG. 1B



E2
E3
Ligase
Androgen
Receptor
Androgen
Receptor
proteasome
Degraded
Target
Protein

FIG. 2.

Table 2. Exemplary Compounds



Ex. No.	Compound Structure	DC50 code	Dmax code	m/z+	1H NMR	Prepared by General Scheme
1		B	B	837.3	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.57 (s, 6H), 2.09-2.14 (m, 1H), 2.70-2.90 (m, 3H), 3.47 (t, *J* = 5.2 Hz, 2H), 3.69-3.74 (m, 10H), 3.88 (t, *J* = 4.6 Hz, 2H), 4.17 (t, *J* = 4.6 Hz, 2H), 4.88-4.93 (m, 1H), 6.50 (br, 1H), 6.92 (d, *J* = 8.8 Hz, 1H), 7.03-7.11 (m, 3H), 7.19 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97 (d, *J* = 8.4 Hz, 2H), 8.06 (s, 1H).	1A
2		D	B	749.3	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.56 (s, 6H), 2.10-2.14 (m, 1H), 2.72-2.90 (m, 3H), 3.50 (t, *J* = 5.4 Hz, 2H), 3.81 (t, *J* = 5.2 Hz, 2H), 3.90 (t, *J* = 4.4 Hz, 2H), 4.20 (t, *J* = 4.2 Hz, 2H), 4.89-4.94 (m, 1H), 6.52 (br, 1H), 6.93 (d, *J* = 8.4 Hz, 1H), 7.05-7.12 (m, 3H), 7.19 (d, *J* = 8.4 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.96-7.99 (m, 2H), 8.10 (br, 1H).	1A

FIG. 2 Continued.



3		C	B	925.3	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.57 (s, 9H), 2.07-2.11 (m, 1H), 2.70-2.82 (m, 3H), 3.44-3.49 (m, 2H), 3.66-3.75 (m, 18H), 3.87-3.89 (m, 2H), 4.16-4.18 (m, 2H), 4.84-4.88 (m, 1H), 6.91 (d, *J* = 8.4 Hz, 1H), 7.03-7.11 (m, 3H), 7.20 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97 (d, *J* = 8.0 Hz, 2H), 8.38 (s, 1H).	1A
4		D	B	793.3	^{1}HNMR (400 MHz, CD_3OD):δ 1.43 (s, 6H), 1.93-1.98(m, 1H), 2.57-2.73 (m, 3H), 3.40 (t, J = 5.2 Hz, 2H), 3.59-3.65 (m, 6H), 3.78 (t, J = 4.4 Hz, 2H), 4.06(t, J = 4.4 Hz, 2H), 4.91-4.95 (m, 1H), 6.93-7.00 (m, 4H), 7.15 (d, J = 8.8 Hz, 2H), 7.41-7.45 (m, 1H), 7.87-7.90 (m, 1H), 8.04-8.06(m, 2H).	1A

FIG. 2 Continued.




5		C	B	807.4	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.56 (s, 6H), 2.08-2.10 (m, 3H), 2.70-2.90 (m, 3H), 3.46 (t, *J* = 5.2 Hz, 2H), 3.65-3.73 (m, 8H), 4.11 (t, *J* = 6 Hz, 2H), 4.88-4.92 (m, 1H), 6.92 (d, *J* = 8.8 Hz, 1H), 7.03 (d, *J* = 8.8 Hz, 2H), 7.10 (d, *J* = 7.2 Hz, 2H), 7.18 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.6 Hz, 1H), 7.84 (d, *J* = 8.0 Hz, 1H), 7.97-7.99 (m, 3H).	1A
6		C	B	865.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.57 (s, 6H), 1.76-1.79 (m, 2H), 1.87-1.90 (m, 2H),2.10-2.14 (m, 1H), 2.71-2.91 (m, 3H), 3.47-3.74 (m, 14H), 4.02 (t, *J* = 6.2 Hz, 2H), 4.88-4.93 (m, 1H), 6.48-6.49 (m, 1H), 6.91-7.02 (m, 3H), 7.10-7.20 (m, 3H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97-8.02 (m, 3H).	1A
7		B	A	851.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.57 (s, 6H), 2.07-2.11 (m, 3H), 2.70-2.90 (m, 3H), 3.46-3.72 (m, 14H), 4.10 (t, *J* = 6.2 Hz, 2H), 4.88-4.92 (m, 1H), 6.48-6.49 (m, 1H), 6.91-7.26 (m, 6H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.85(m, 1H), 7.97-8.02 (m, 3H).	1A

FIG. 2 Continued.



8		C	B	837.2	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.57 (s, 6H), 2.09-2.14 (m, 1H), 2.70-2.90 (m, 3H), 3.47 (t, *J* = 5.2 Hz, 2H), 3.69-3.74 (m, 10H), 3.88 (t, *J* = 4.6 Hz, 2H), 4.17 (t, *J* = 4.6 Hz, 2H), 4.88-4.93 (m, 1H), 6.50 (br, 1H), 6.92 (d, *J* = 8.8 Hz, 1H), 7.03-7.11 (m, 3H), 7.19 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97 (d, *J* = 8.4 Hz, 2H), 8.06 (s, 1H).	1A
9		B	B	837.2	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.57 (s, 6H), 2.09-2.14 (m, 1H), 2.70-2.90 (m, 3H), 3.47 (t, *J* = 5.2 Hz, 2H), 3.69-3.74 (m, 10H), 3.88 (t, *J* = 4.6 Hz, 2H), 4.17 (t, *J* = 4.6 Hz, 2H), 4.88-4.93 (m, 1H), 6.50 (br, 1H), 6.92 (d, *J* = 8.8 Hz, 1H), 7.03-7.11 (m, 3H), 7.19 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97 (d, *J* = 8.4 Hz, 2H), 8.06 (s, 1H).	1A

FIG. 2 Continued.




10		B	B	821.3	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.57 (s, 6H), 1.86-1.89 (m, 2H), 2.04-2.09 (m, 3H), 2.71-2.90 (m, 3H), 3.43 (s, 2H), 3.55-3.63 (m, 8H), 4.10 (t, J = 6.0 Hz, 2H), 4.89-4.93 (m, 1H), 6.91 (d, J=8.8Hz, 1H), 7.02-7.05 (m, 2H), 7.10 (d, J=6.8Hz, 1H), 7.17-7.20 (m, 2H), 7.47-7.51 (m, 1H), 7.83-7.86 (m, 1H), 7.97-7.99(m, 2H), 8.07 (s, 1H).	1A
11		D	B	865.1	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.42 (s, 6H), 1.85 (s, 3H), 1.92-1.96 (m, 3H), 2.44-2.49 (m, 1H), 2.56-2.64 (m, 2H), 3.34 (d, *J* = 5.2 Hz, 2H), 3.50-3.59 (m, 12H), 4.01 (d, *J* = 6.2 Hz, 2H) ,6.86-6.94 (m, 4H), 7.13-7.16 (m, 2H), 7.38-7.42 (m, 1H), 7.86-7.89 (m, 1H), 8.03-8.05 (m, 2H).	1A
12		B	A			1A

FIG. 2 Continued.



13		B	A	798.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.31 (s, 6H), 1.55-1.63 (m, 2H), 1.68-1.74 (m, 2H), 1.80-1.87 (m, 2H), 1.91-1.97 (m, 2H), 2.06-2.11 (m, 1H), 2.67-2.81 (m,3H), 3.43-3.48 (m, 2H), 3.48-3.53 (m, 2H), 3.58-3.61 (m, 2H), 4.04 (t, *J* = 6.4 Hz, 2H), 4.16 (s, 1H), 4.30 (s, 1H), 5.02-5.07 (m, 1H), 6.96-7.08 (m, 5H), 7.14-7.15 (m, 1H), 7.52-7.56 (m, 1H), 7.73-7.80 (m, 3H).	1A
14		D	B		1HNMR (400 MHz, DMSOd6) :δ 10.97 (s, 1H), 8.38 (d, *J* = 8.0 Hz , 1H), 8.29 (s, 1H), 8.07 (d, *J* = 8.4 Hz, 1H), 7.46 (t, *J* = 2.6 Hz, 1H), 7.24-7.33 (m, 4H), 7.08 (d, *J* = 8.4 Hz, 2H), 5.09-5.12 (m, 1H), 4.05-4.39 (m, 9H), 3.77 (s, 2H), 3.51-3.59 (m, 10H), 2.80-3.00 (m, 1H), 2.30-2.70 (m, 2H), 1.90-2.10 (m, 3H), 1.24 (s, 6H).	2A

FIG. 2 Continued.



15		D	B	753.7	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.22 (s, 6H), 1.25 (s, 6H), 2.00-2.03 (m, 2H), 2.22 (t, *J* = 7.4 Hz, 2H), 2.67-2.88 (m, 8H), 3.35 (s, 2H), 3.72 (br, 4H), 4.04 (s, 1H), 4.13-4.22 (m, 2H), 4.32-4.36 (m, 1H), 5.34-5.39 (m, 2H), 6.09 (d, *J* = 8.0 Hz, 1H), 6.67 (d, *J* = 9.2 Hz, 1H), 6.80 (d, *J* = 8.4 Hz, 2H), 6.96 (d, *J* = 1.2 Hz, 1H), 7.29-7.30 (m, 1H), 7.38 (t, *J* = 7.6 Hz, 1H), 7.57 (d, *J* = 8.8 Hz, 1H), 7.93-7.96 (m, 1H), 8.17 (s, 1H), 8.59 (s, 1H).	3A
16		B	A	825.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.31 (s, 6H), 1.96-2.12 (m, 4H), 2.71-2.81 (m, 8H), 3.48 (t, *J* = 6.0 Hz, 2H), 3.64-3.70 (m, 8H), 4.16 (s, 1H), 4.30 (s, 1H), 5.03-5.09 (m, 1H), 6.82 (d, *J* = 8.8 Hz, 1H), 6.99-7.10 (m, 3H), 7.14-7.15 (m, 1H), 7.58 (t, *J* = 7.8 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.95-7.98 (m, 1H), 8.60-8.61 (m, 1H).	7A

FIG. 2 Continued.



17		D	B		^{1}H NMR (400 MHz, CD_3OD): δ 9.64 (s, 1H), 8.19-8.13 (m, 3H), 8.10-8.08 (m, 2H), 8.03-8.00 (m, 1H), 7.94 (d, *J* = 8.8 Hz, 2H), 7.58-7.55 (m, 2H), 7.35-7.33 (m, 1H), 7.29-7.27 (m, 1H), 7.02 (d, *J* = 7.2 Hz, 1H), 4.97-4.93 (m, 1H), 4.86-4.81 (m, 3H), 4.40-4.39 (m, 2H), 4.25 (d, *J* = 16.8 Hz, 1H), 4.09 (d, *J* = 17.2 Hz, 2H), 2.38-2.28 (m, 1H), 2.17-2.12 (m, 1H), 1.63 (s, 6H).	4A
18		B	A	815.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.23 (s, 6H), 1.29 (s, 6H), 1.49-1.56 (m, 2H), 1.66-1.73 (m, 4H), 1.92-1.95 (m, 2H), 2.06-2.12 (m, 1H), 2.69-2.80 (m, 3H), 3.23 (t, *J* = 7.0 Hz, 2H), 3.44 (t, *J* = 6.4 Hz, 2H), 3.50 (t, *J* = 6.2 Hz, 2H), 3.59 (t, *J* = 5.6 Hz, 2H), 4.13-4.15 (m, 1H), 4.30 (s, 1H), 5.03-5.07 (m, 1H), 6.74 (t, *J* = 8.4 Hz, 1H), 6.99-7.08 (m, 3H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.44-7.57 (m, 3H), 7.74 (d, *J* = 8.8 Hz, 1H).	5A

FIG. 2 Continued.



19		D	B	754.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.21 (s, 6H), 1.25 (s, 6H), 2.19-2.24 (m, 1H), 2.32-2.42 (m, 1H), 2.75 (br, 4H), 2.83-2.88 (m, 1H), 2.90-2.95 (m, 3H), 3.73 (br,4H), 4.04 (s, 1H), 4.14 (d, *J* = 8.0 Hz, 1H), 4.28-4.33 (m, 3H), 4.40, 4.44 (two singles, 1H), 5.20-5.25 (m, 1H), 6.09 (d, *J* = 8.0 Hz, 1H), 6.67 (d, J = 8.4 Hz, 1H), 6.80 (dd, *J* = 8.8, 2.4 Hz, 1H), 6.96 (d, *J* = 2.4 Hz, 1H), 7.06 (d, *J* = 8.0 Hz, 1H), 7.43-7.52 (m, 2H), 7.57 (d, *J* = 8.8 Hz, 1H), 7.93 (dd, *J* = 8.8, 2.4 Hz, 1H), 8.20 (br, 1H),8.57-8.58 (m, 1H).	6A
20		D	B	767.4	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.22-1.25 (m, 12H), 2.01-2.02 (m, 2H), 2.59-2.69 (m, 4H), 2.75-2.91 (m, 4H), 3.39-3.40 (m, 2H), 3.73 (s, 4H), 4.04 (s, 1H), 4.14 (d, *J* = 8.0 Hz, 1H), 4.89-4.94 (m, 1H), 6.06 (d, *J* = 7.6 Hz, 1H), 6.67 (d, *J* = 8.8 Hz, 1H), 6.80 (d, *J* = 6.0 Hz, 2H),, 6.89-6.97 (m, 2H), 7.12 (d, *J* = 7.2 Hz, 1H), 7.52-7.58 (m, 2H), 7.92-8.01 (m, 2H), 8.58 (s, 1H).	7A

FIG. 2 Continued.



21		D	B	752.7	^{1}H NMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.18 (s, 6H), 1.83-1.90 (m, 2H), 2.07-2.11 (m, 1H), 2.38-2.45 (m, 3H), 2.50-2.53 (m, 4H), 2.68-2.72 (m, 3H), 2.78-2.86 (m, 1H), 3.57-3.59 (m, 4H), 4.04 (s, 1H), 4.18 (s, 1H), 4.38-4.49 (m, 2H), 5.07-5.12 (m, 1H), 6.74 (d, *J* = 8.8 Hz, 1H), 6.87-6.90 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.38-7.44 (m, 2H), 7.56-7.58 (m, 1H), 7.62 (d, *J* = 8.8 Hz, 1H), 7.85-7.88 (m, 1H), 8.49 (d, *J* = 2.4 Hz, 1H).	8A
22		A	A	798.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.32 (s, 6H), 2.16-2.24 (m, 1H), 2.45-2.56 (m, 1H), 2.74-2.81 (m, 1H), 2.89-2.94 (m, 1H), 3.00-3.06 (m, 6H), 3.73-3.81 (m,4H), 3.85-3.87 (m, 2H), 3.90-3.93 (m, 2H), 4.16-4.18 (m, 1H), 4.30 (s, 1H), 4.35-4.38 (m, 2H), 4.41-4.52 (m, 2H), 5.17-5.21 (m, 1H), 6.87 (d, *J* = 8.8 Hz, 1H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.28 (d, *J* = 8.0 Hz, 1H), 7.42 (d, *J* = 7.2 Hz, 1H), 7.50-7.54 (m, 1H),7.71-7.76 (m, 2H), 7.99 (dd, *J* = 8.8, 2.4 Hz, 1H), 8.62, 8.63 (two singles, 1H).	6A

FIG. 2 Continued.



23		B	A	843.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.18 (s, 6H), 1.24 (s, 6H), 1.50-1.66 (m, 4H), 2.08-2.14 (m, 1H), 2.39-2.75 (m, 13H), 2.81-2.86 (m, 3H), 4.11(s, 1H), 4.23 (t, *J* = 5.4 Hz, 2H), 4.29 (s, 1H), 4.33-4.44 (m, 2H), 5.07-5.11 (m, 1H), 7.16-7.21 (m, 2H), 7.26-7.35 (m, 4H), 7.41-7.45 (m, 1H), 7.69 (d, *J* = 8.0 Hz, 2H), 7.86 (d, *J* = 8.8 Hz, 1H).	8A
24		B	A	825.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.31 (s, 6H), 1.93-1.98 (m, 2H), 2.06-2.12 (m, 1H), 2.67-2.86 (m, 9H), 3.37 (t, *J* = 6.8 Hz, 2H), 3.61-3.72 (m, 8H), 4.15 (s, 1H), 4.31 (s, 1H), 4.63 (s, 1H), 5.03-5.08 (m, 1H), 6.82-6.89 (m, 2H), 6.99-7.04 (m, 2H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.56 (d, *J* = 8.4 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.96-7.99 (m, 1H), 8.62 (d, *J* = 2.4 Hz, 1H).	7A

FIG. 2 Continued.



25		B	A	843.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.32 (d, *J* = 2 Hz, 6H), 1.95-1.99 (m, 2H), 2.07-2.11 (m, 1H), 2.65-2.76 (m, 2H), 2.82-2.94 (m, 7H), 3.37 (s, 1H), 3.46 (t, *J* = 6.2 Hz, 2H), 3.65-3.75 (m, 8H), 4.15 (t, *J* = 4.4 Hz, 1H), 4.31 (s, 1H), 5.04-5.08 (m, 1H), 6.15-6.17 (m, 1H), 6.84 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.12-7.16 (m, 2H), 7.35 (t, *J* = 9.0 Hz, 1H), 7.75 (d, *J* = 8.8 Hz, 1H), 7.96-7.99 (m, 1H), 8.61 (d, *J* = 2.4 Hz, 1H).	7A
26		B	A	841.1	^{1}H NMR (400 MHz, CD_3OD): δ7.94-7.88 (m, 1H), 7.74-7.59 (m, 3H), 7.38-7.19 (m, 6H), 4.35 (s, 1H), 4.14(s, 1H), 3.46 (s, 3H), 2.73-2.54 (m , 7H), 2.41-2.33 (m, 3H), 2.19-2.01 (m, 1H), 1.73--1.58(m, 2H), 1.37(s, 4H), 1.24 (s, 9H), 1.21 (s, 7H), 0.87-0.83 (m, 1H);	10A

FIG. 2 Continued.


27		C	A	856.9	^{1}HNMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.30 (s, 6H), 1.87-2.12 (m, 6H), 2.57-2.85 (m, 16H), 3.41 (t, *J* = 8.0 Hz, 2H), 4.17 (s, 1H), 4.35 (s, 1H), 5.02-5.07 (m, 1H), 7.05 (t, *J* = 8.0 Hz, 2H), 7.24 (d, *J* = 8.0 Hz, 1H), 7.36 (s, 3H), 7.55 (t, *J* = 8.0 Hz, 1H), 7.76 (d, *J* = 4.0 Hz, 2H), 7.92 (d, *J* = 12.0 Hz, 1H).	11A
28		B	B	921.5	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.29 (s, 6H), 1.40-1.44 (m, 2H), 1.62-1.72 (m, 4H), 1.88-1.94 (m, 2H), 2.05-2.12 (m, 1H), 2.67-2.83 (m, 5H), 3.40-3.48 (m, 4H), 3.56 (t, *J* = 5.6 Hz, 2H), 4.15-4.17 (m, 1H), 4.29 (s, 1H), 5.01-5.05 (m, 1H), 6.95-7.02 (m, 1H), 7.01-7.10 (m, 2H), 7.14 (d, *J* = 2.2 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 2H), 7.54 (t, *J* = 8.0 Hz, 1H), 7.71-7.74 (m, 3H).	12A

FIG. 2 Continued.



29		B	A	843.7	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.79-1.82 (m, 2H), 1.99-2.05 (m, 2H), 2.33 (s, 1H), 2.55-2.67 (m, 5H), 2.82-2.91 (m, 2H), 3.48-3.61 (m, 10H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.30 (s, 1H), 5.05-5.09 (m, 1H), 6.54-6.57 (m, 1H), 6.86 (d, *J* = 8.8 Hz, 1H), 6.99-7.02 (m, 1H), 7.07-7.10 (m, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.44-7.49 (m, 1H), 7.62 (d, *J* = 9.2 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 7.95-7.98 (m, 1H), 8.13 (s, 2H), 8.62 (s, 1H), 11.11 (s, 1H).	12A
30		A	A	809.8	^{1}HNMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.18 (s, 6H), 1.52-1.65 (m, 4H), 2.01-2.09 (m, 1H), 2.18-2.54 (m, 2H), 2.61-3.03 (m, 15 H), 4.05-4.07 (m, 1H), 4.13-4.19 (m, 3H), 4.29-4.39 (m, 2H), 4.99-5.04 (m, 1H), 6.87-6.89 (m, 1H), 6.97-7.05 (m, 3H), 7.23-7.25 (m, 2H), 7.60-7.67 (m, 4H).	13A

FIG. 2 Continued.




31		B	A	809.8	^{1}H NMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.71-1.73 (m, 4H), 2.16-2.23 (m, 1H), 2.47-2.74 (m, 3H), 2.78-2.79 (m, 3H), 2.82-3.16 (m, 11H), 4.17-4.19 (m, 1H), 4.31-4.34 (m, 3H), 4.41-4.52 (m, 2H), 5.15-5.20 (m, 1H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.14, 7.15 (two singles, 1H), 7.25-7.27 (m, 1H), 7.35-7.37 (m, 2H), 7.42-7.44 (m, 1H), 7.50-7.54 (m, 1H), 7.71-7.79 (m, 4H).	13A using imide from scheme 6a
32		B	A	840.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.72-1.74 (m, 4H), 2.11-2.16 (m, 1H), 2.70-3.12 (m, 17H), 3.44 (t, *J* = 6.0 Hz, 2H), 4.17 (s, 1H), 4.31 (s, 1H), 4.61 (br, 1H), 5.06-5.11 (m, 1H), 7.00 (dd, *J* = 8.8 Hz, 1H), 7.12-7.15 (m, 2H), 7.33-7.38 (m, 3H), 7.73-7.79 (m, 3H).	11a
33		A	A	796.7	^{1}H NMR (400 MHz, CD_3OD): δ 1.08 (s, 6H), 1.18 (s, 6H), 1.46-1.52 (m, 2H), 1.56-1.64 (m, 2H), 1.76-1.83(m, 2H), 2.05-2.09 (m, 1H), 2.37-2.42 (m, 1H), 2.48-2.53 (m, 2H), 2.58-2.69 (m, 5H), 2.76-2.84 (m, 1H), 3.61-3.67 (m,	6a

FIG. 2 Continued.



					4H),4.04-4.09 (m, 3H), 4.18 (s, 1H), 4.27-4.38 (m, 2H),5.03-5.07 (m, 1H), 6.76 (d, *J* = 8.8 Hz, 1H), 6.88 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.03 (d, *J* = 2.0 Hz, 1H), 7.11 (d, *J* = 8.0 Hz, 1H), 7.28 (d, *J* = 7.6 Hz, 1H), 7.62 (d, *J* = 8.8 Hz, 1H), 7.62-7.64 (m, 1H), 7.88 (dd, *J* = 9.2, 2.4 Hz, 1H), 8.51 (d, *J* = 2.4 Hz, 1H).	
34		A	A	823.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.72-1.73 (m, 4H), 2.12-2.17 (m, 1H), 2.68-3.15 (m, 17H), 4.17-4.19 (m, 1H), 4.31-4.35 (m, 3H), 5.11-5.14 (m, 1H), 7.00 (dd, *J* = 8.6, 2.4 Hz, 1H), 7.14-7.15 (m, 1H), 7.35-7.37 (m, 3H), 7.45-7.46 (m, 1H), 7.73-7.84 (m, 5H).	6a
35		A	A	796.7	^{1}HNMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.45-1.62 (m, 4H), 1.75-1.82 (m, 2H), 1.95-2.03 (m, 1H), 2.33-2.45 (m, 5H), 2.57-2.61 (m, 1H), 2.88-2.95 (m, 1H), 3.59 (br, 4H), 4.05-4.10 (m, 3H), 4.24-4.31 (m, 2H), 4.37-4.41 (m, 1H), 5.05-5.10 (m, 1H), 6.87 (br, 1H), 6.99-7.06 (m, 2H), 7.17 (br, 1H), 7.20-7.21 (m, 1H),	6a

FIG. 2 Continued.



					7.61-7.63 (m, 2H), 7.89-7.91 (m, 1H), 7.96-7.98 (m, 1H), 8.63 (br, 1H), 10.96 (s, 1H).	
36		D	B	797.7	^{1}H NMR (400 MHz, CD_3OD): δ 1.26, 1.31 (two singles, 12H), 1.73 (br, 4H), 2.14 (br, 1H), 2.65-2.76 (m, 5H), 2.79-2.86 (m, 2H),2.93 (br, 4H), 3.56 (br, 4H), 4.18 (br, 1H), 4.32 (s, 1H), 5.11-5.16 (m, 1H), 6.99-7.02 (m, 1H), 7.16 (s, 1H), 7.37-7.50 (m, 4H), 7.74-7.80 (m, 3H), 7.93 (s, 1H).	10a
37		B	A	840.7	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.73-1.75 (m, 4H), 2.11-2.18 (m, 1H), 2.68-2.80 (m, 9H), 2.84-2.90 (m, 2H), 2.92-3.15 (m, 6H), 3.67-3.70 (m, 2H), 4.18 (s, 1H), 4.31 (s, 1H), 5.07-5.12 (m, 1H), 6.99-7.02 (m, 1H), 7.13-7.16 (m, 2H), 7.33-7.39 (m, 3H), 7.72-7.79 (m, 3H).	11a

FIG. 2 Continued.



38		A	A	794.7	^{1}HNMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.30 (s, 6H), 1.44-1.45 (m, 4H), 1.65-1.77 (m, 4H), 2.18-2.23 (m, 1H), 2.49-2.60 (m, 1H), 2.74-2.83 (m, 5H), 2.89-3.03 (m, 5H), 3.85 (br, 4H), 4.16 (t, *J* = 4.4 Hz, 1H), 4.30 (s, 1H), 4.46-4.56 (m, 2H), 5.17-5.22 (m, 1H), 6.92 (d, *J* = 9.2 Hz, 1H), 6.98-7.01 (m, 1H), 7.14 (d, *J* = 2.4 Hz, 1H), 7.48-7.50 (m, 2H), 7.66-7.75 (m, 3H), 8.01-8.04 (m, 1H), 8.65 (d, *J* = 2.4 Hz, 1H).	8a
39		A	A	797.7	^{1}H NMR (400 MHz, CD_3OD): δ 1.14 (s, 6H), 1.23 (s, 6H), 1.43-1.54 (m, 2H), 1.60-1.65 (m, 2H), 1.97-2.06 (m, 2H), 2.33-2.45 (m, 4H), 2.57-2.61 (m, 2H), 2.66-2.69 (m, 2H), 2.84-2.93 (m, 1H), 3.16-3.30 (m, 5H), 4.07 (d, J = 8.8 Hz, 1H), 4.33 (s, 1H), 5.07-5.12 (m, 1H), 6.99-7.02 (m, 1H), 7.21-7.22 (m, 1H), 7.31-7.33 (m, 2H), 7.38 (br, 1H), 7.64-7.66 (m, 1H), 7.75-7.79 (m, 3H), 7.91 (d, J = 8.8 Hz, 1H), 11.11 (s, 1H).	10a

FIG. 2 Continued.




40		B	A	807.4	1H NMR (300 MHz,DMSO): δ11.11 (S, 1H), 7.94--7.92 (d, *J* = 8.4 Hz, 1H), 7.79--7.63 (m, 4H), 7.36--7.18(m, 6H), 7.14--6.96(m, 1H), 5.13--4.98(m , 1H),4.31(s, 1H), 4.09--4.11 (d,J=8.4 Hz, 1H), 3.45 (s,4H),3.01--2.79 (m, 3H), 2.69--2.58 (m, 2H), 2.46(s,3H),2.42--2.27 (m, 2H), 2.08--1.94 (m, 1H),1.66--1.53 (m, 2H),1.48--1.39 (m, 2H),1.31(s,4H),1.21(s,6H),1.15(s,6H)	10a
41		A	A	810.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.46-1.49 (m, 2H), 1.58-1.60 (m, 2H), 1.78-1.82 (m, 2H), 2.03-2.06 (m, 4H), 2.54-2.62 (m, 4H), 2.85-2.93 (m, 1H), 3.17 (d, *J* = 4.8 Hz, 1H), 3.58 (br, 4H), 4.05-4.07 (m, 1H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.31 (s, 1H), 5.10-5.14 (m, 1H), 6.88-6.90 (m, 1H), 7.01 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 8.2, 2.4 Hz, 1H), 7.43-7.44 (m, 1H), 7.62-7.64 (m, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 7.97-7.99 (m, 1H), 8.63-8.64 (m, 1H), 11.11 (s, 1H).	6a

FIG. 2 Continued.

42		D	B	740.3	H-NMR13-PH-ARV-LS-019-B-0: (300 MHz, CD3OD) δ 8.70 (s, 1H), 8.11 – 8.08 (d, *J* = 11.1 MHz, 1H), 7.75 -7.72 (d, *J* = 8.7 MHz, 1H), 7.52 – 7.49 (m, 1H), 7.43 - 7.40 (d, *J* = 7.2 MHz, 1H), 7.26 - 7.23 (d, *J* = 7.8 MHz, 1H), 7.12 (s, 1H), 7.01 - 6.95 (m, 2H), 5.20 - 5.16 (m, 1H), 5.02 - 4.98 (m, 1H), 4.70 - 4.65 (m, 3H), 4.46 - 4.43 (d, *J* = 6.9 MHz, 2H), 4.24 - 4.20 (m, 2H), 3.58 - 3.81 (m, 2H), 3.25 - 3.22 (m, 6H), 3.03 - 2.75 (m, 2H), 2.74 - 2.60 (m, 4H), 2.57 - 2.34 (m, 1H), 2.29 - 2.10 (m, 1H), 1.97 - 1.87 (m, 4H), 1.67 - 1.64 (m, 2H).	14a
43		D	B	739.2	H-NMR- PH-ARV-LS-019-C-0: (400 MHz, CD3OD) δ 7.83 - 7.81 (d, *J* = 8.8 MHz, 2H), 7.71 -7.69 (d, *J* = 8.8 MHz, 1H), 7.51 - 7.47 (m, 1H), 7.40 - 7.38 (d, *J* = 7.9 MHz, 1H), 7.23 - 7.21 (d, *J* = 8 MHz, 1H), 7.09 - 7.06 (m, 3H), 6.96 - 6.93 (d, *J* = 10.8 MHz, 1H), 5.18 - 5.14 (m, 1H), 5.00 - 4.99 (m, 1H), 40.65 - 4.60 (m, 1H), 4.48 - 4.37 (m, 2H), 4.21 - 4.18 (m, 2H), 4.05 - 4.02 (m, 2H), 3.69 - 3.67 (m, 2H), 3.26 - 3.14 (m,	14a

WO 2018/071606 PCT/US2017/056234

FIG. 2 Continued.



					6H), 2.91 - 2.74 (m, 2H), 2.66 - 2.59 (m, 4H), 2.47 - 2.41 (m, 1H), 2.21 - 2.11 (m, 1H), 1.95 - 1.85 (m, 4H), 1.65 - 1.61 (m, 2H)	
44		A	A	793.5	1H NMR (300 MHz, Methanol-d4) δ 7.78- 7.63 (m, 4H), 7.43 (d, J = 2.2 Hz, 1H), 7.36-7.25 (m, 3H), 7.10 (d, J = 2.4 Hz, 1H), 6.95 (dd, J = 8.8, 2.5 Hz, 1H), 5.06 (dd, J = 12.4, 5.4 Hz, 1H), 4.26 (s, 1H), 4.14 (d, J = 5.1 Hz, 1H), 3.65 (s, 4H), 3.23 (d, J = 5.3 Hz, 4H), 2.99 (t, J = 8.2 Hz, 2H), 2.89-2.62 (m, 5H), 2.15- 2.02 (m, 1H), 1.74 (h, J = 7.8 Hz, 4H), 1.43 (q, J = 7.6 Hz, 2H), 1.23 (d, J = 16.2 Hz, 6H), 1.15 (d, J = 16.2 Hz, 6H)	10a
45		A	A	779.5	^{1}H NMR (300 MHz, Chloroform-*d*) δ 8.05 (s, 1H), 7.69 (dd, *J* = 8.2, 5.8 Hz, 3H), 7.55 (d, *J* = 8.7 Hz, 1H), 7.27 (d, *J* = 2.8 Hz, 2H), 7.05 (d, *J* = 8.4 Hz, 1H), 6.90 (m, 1H), 6.75(m, 1H), 6.14 (d, *J* = 2.4 Hz, 1H), 4.79 (dd, *J* = 8.9, 2.4 Hz, 1H), 4.19 (d, *J* = 8.1 Hz, 1H), 3.92 (s, 51H), 3.88-3.39 (m, 4H), 3.00-2.49 (m, 9H), 2.14-2.12 (m, 1H), 1.94-1.68 (m, 6H), 1.32-1.30 (m, 1H), 1.23 (s, 6H),	10a

FIG. 2 Continued.



					1.19 (s, 6H).	
46		A	A	825.4	1H NMR (400 MHz, DMSO): δ 11.11 (s, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 7.78-7.71 (m, 4H), 7.44 (d, *J* = 7.2 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 2H), 7.21 (s, 1H), 7.01 (d, *J* = 8.8 Hz, 1H), 5.13-5.08 (m, 1H), 4.32 (s, 1H), 4.06 (d, *J* = 9.2 Hz, 1H), 3.40-3.26 (m, 5H), 3.25-3.22 (m, 4H), 2.91-2.84 (m, 1H), 2.69-2.59 (m, 3H), 2.34-2.27 (m, 2H), 2.10-2.02 (m, 1H), 1.65-1.58 (m, 2H), 1.48-1.40 (m, 2H), 1.38-1.30 (m, 4H), 1.22 (s, 6H), 1.13 (s, 6H)	10a
47		D	B	697.3	^{1}H NMR (400 MHz, DMSO-d6): δ 1.14 (s, 6H), 1.23 (s, 6H), 1.77 (br, 4H), 1.98-2.00 (m, 1H), 2.43-2.48 (m, 1H), 2.56-2.61 (m, 1H), 2.69-2.73 (m, 2H), 2.87-2.96 (m, 1H), 4.06-4.15 (m, 3H), 4.20-4.25 (m, 1H), 4.33-4.39 (m, 2H), 5.09-5.13 (m, 1H), 7.00-7.02 (m, 1H), 7.21-7.25 (m, 2H), 7.30-7.35 (m, 3H), 7.48 (t, *J* = 7.6 Hz, 1H), 7.75-7.79 (m, 3H), 7.91 (d, *J* = 8.8 Hz, 1H), 10.98 (s, 1H).	15A

FIG. 2 Continued.



48		D	B	711.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.53-1.60 (m, 2H), 1.73-1.80 (m, 2H), 1.84-1.91 (m, 2H), 2.15-2.21 (m, 1H), 2.45-2.56 (m, 1H), 2.75 (t, *J* = 7.6 Hz, 2H), 2.81-2.83 (m, 1H), 2.89-2.93 (m, 1H), 4.13-4.18 (m, 3H), 4.30 (s, 1H), 4.35-4.44 (m, 2H), 5.15 (d, *J* = 13.6 Hz, 1H), 7.00 (d, *J* = 8.8 Hz, 1H), 7.14 (d, *J* = 2.4 Hz, 1H), 7.20 (d, *J* = 8.0 Hz, 1H), 7.33 (d, *J* = 8.4 Hz, 2H), 7.39 (d, *J* = 7.6 Hz, 1H), 7.49 (t, *J* = 8.0 Hz, 1H), 7.73-7.76 (m, 3H).	15a
49		D	B	727.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.90-1.96 (m, 2H), 2.12-2.19 (m, 1H), 2.38-2.49 (m, 1H), 2.74-2.80 (m, 3H), 2.86-2.95 (m, 1H), 3.55-3.58 (m, 2H), 3.85-3.86 (m, 2H), 4.16-4.18 (m, 1H), 4.30-4.33 (m, 3H), 4.44-4.50 (m, 2H),5.13-5.17 (m, 1H), 6.99-7.02 (m, 1H), 7.14-7.15 (m, 1H), 7.26-7.32 (m, 3H), 7.42-7.44 (m, 1H), 7.50-7.54 (m, 1H), 7.71-7.75 (m, 3H).	15a

FIG. 2 Continued.



50		D	B	741.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.14 (s, 6H), 1.23 (s, 6H), 1.49-1.56 (m, 2H), 1.59-1.66 (m, 2H), 1.95-1.99 (m, 1H), 2.35-2.45 (m, 1H), 2.55-2.60 (m, 1H), 2.64 (t, *J* = 7.2 Hz, 2H), 2.86-2.95 (m, 1H), 3.49 (t, *J* = 6.4 Hz, 2H), 3.72-3.74 (m, 2H), 4.07 (d, *J* = 9.2 Hz, 1H), 4.21-4.27 (m, 3H), 4.33-4.38 (m, 2H), 5.08-5.12 (m, 1H), 7.01 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.26-7.28 (m, 3H), 7.31-7.33 (m, 1H), 7.48 (t, *J* = 7.6 Hz, 1H), 7.73-7.76 (m, 3H), 7.91 (d, *J* = 8.8 Hz, 1H), 10.97 (s, 1H).	15a
51		D	B	755.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.38-1.46 (m, 2H), 1.60-1.71 (m, 4H), 2.11-2.23 (m, 1H), 2.41-2.52 (m, 1H), 2.68 (t, *J* = 7.2 Hz, 2H), 2.73-2.79 (m, 1H), 2.86-2.95 (m, 1H), 3.56 (t, *J* = 6.0 Hz, 2H), 3.82 (t, *J* = 4.4 Hz, 2H), 4.15-4.18 (m, 1H), 4.28-4.30 (m, 2H), 4.39-4.49 (m, 2H), 4.59 (s, 1H), 5.14 (d, *J* = 13.2 Hz, 1H), 7.00 (d, *J* = 8.8 Hz, 1H), 7.14 (d, *J* = 2.4 Hz, 1H), 7.23-7.29 (m, 3H), 7.40-7.42 (m, 1H), 7.50 (t, *J*	15a

FIG. 2 Continued.



					= 7.6 Hz, 1H), 7.70-7.75 (m, 3H).	
52		D	B	769.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.31 (s, 6H), 1.36-1.46 (m, 4H), 1.56-1.67 (m, 4H), 2.11-2.17 (m, 1H), 2.41-2.52 (m, 1H), 2.65-2.68 (m, 2H), 2.71-2.78 (m, 1H), 2.83-2.90 (m, 1H), 3.53-3.56 (m, 2H), 3.81-3.83 (m, 2H), 4.16-4.18 (m, 1H), 4.28-4.30 (m, 3H), 4.40-4.50 (m, 2H), 5.11-5.16 (m, 1H), 6.99-7.01 (m, 1H), 7.14-7.15 (m, 1H), 7.24-7.30 (m, 3H), 7.40-7.41 (m, 1H), 7.48-7.52 (m, 1H), 7.73-7.75 (m, 3H).	15a
53		D	B	780.6	^{1}H NMR (400 MHz, CD_3OD): δ 1.26-1.28 (m, 3H), 1.61-1.64 (m, 2H), 1.70-1.71 (m, 4H), 2.03-2.07 (m, 3H), 2.09-2.14 (m, 1H), 2.19-2.23 (t, J = 7.6 Hz, 2H), 2.55-2.64 (m, 4H), 2.70-2.76 (m, 2H), 3.36-3.43 (m, 2H), 3.67 (s, 2H), 4.32-4.37 (m, 1H), 4.40-4.45 (m, 1H), 4.55-4.60 (m, 1H), 5.04-5.09 (m, 1H), 5.35-5.38 (m, 1H), 6.64 (br, 1H), 6.78 (d, J = 2.4 Hz, 1H), 7.04-7.09 (m, 2H), 7.52-7.56 (m, 1H), 7.73 (d, J = 2.4 Hz, 1H), 7.79-7.81 (m, 1H), 7.91-7.93 (m, 1H),	16a

FIG. 2 Continued.




					8.04 (br, 1H).	
54		D	B	807.8	H-NMR (300 MHz, DMSO) δ11.14 (s, 1H), 7.91-7.73 (m, 6H), 7.51-7.47 (m, 1H), 7.19 (s, 1H), 7.01-6.91 (m, 3H), 5.17-5.11 (m, 1H), 4.32 (s, 1H), 4.07-4.04 (d, *J*=9.3 Hz, 7H), 3.33-3.31 (m, 1H), 3.24-3.18 (m, 4H), 2.89-2.73 (m, 3H), 2.61-2.51 (m, 5H), 2.30-2.21 (m, 2H), 2.10-2.00 (m, 1H), 1.67-1.59 (m, 2H), 1.43-1.35 (m, 2H), 1.39-1.25 (m, 4H), 1.19 (s, 6H), 1.09 (m, 6H);	8a
55		D	B	683.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.29 (s, 6H), 2.18-2.21 (m, 3H), 2.46-2.60 (m, 1H), 2.73-2.95 (m, 4H), 4.16 (s, 3H), 4.30 (s, 1H) , 4.42-4.43 (m, 2H), 5.13-5.18 (m, 1H), 6.99-7.01 (m, 1H), 7.14-7.18 (m,2H), 7.37-7.50 (m, 4H), 7.72-7.78 (m, 4H).	15a
56		B	A	767.7	^{1}HNMR (400 MHz, DMSO-d6) : δ 1.12 (br, 3H), 1.72 (br, 2H), 1.99-2.03 (m, 1H), 2.33-2.40 (m, 3H), 2.57-2.67 (m, 2H), 2.84-2.92 (m, 1H), 3.17 (s, 1H), 3.34 (br, 6H), 4.07-4.11 (m, 1H), 4.26-4.49 (m, 5H), 5.06-5.08 (m, 1H), 6.54 (s, 1H), 6.93 (s, 1H), 7.21-7.25 (m, 1H), 7.33 (s, 1H), 7.66-7.68	17a

FIG. 2 Continued.



					(m, 1H), 7.82 (s, 1H), 7.98 (s, 1H), 8.07 (s, 1H), 8.26-8.27 (m, 1H), 11.08 (s, 1H), 13.29 (s, 1H).	
57		D	B	694.2	^{1}H NMR (300 MHz, DMSO-*d6*) 8.63-8.54 (m, 1H), 8.19-7.95 (m, 4H), 7.86-7.83 (m, 1H), 7.52-7.49 (m, 1H), 7.36-7.34 (m, 1H), 7.26-7.20 (m, 2H), 7.04-7.01 (m, 1H), 5.05-4.96 (m, 2H), 4.58-4.49 (m, 1H), 4.41-4.10 (m, 4H), 2.96-2.56 (m, 8H), 2.32-1.89 (m, 4H);	18a
58		B	A	841.3	^{1}H NMR (400 MHz, DMSO-d_6) δ1.12 (6H, s), 1.23 (6H, s), 1.31-1.41 (6H, m), 1.48-1.60 (5H, m), 2.30-2.34 (2H, m), 2.45-2.46 (2H, m), 2.54-2.60 (1H, m), 3.24-3.28 (4H, m), 3.57-3.60 (4H, m), 4.04-4.07 (1H, m), 4.30 (1H, s), 5.04-5.07 (1H, m), 6.88 (1H, d, *J* = 8.8 Hz), 6.94 (1H, brs), 7.00 (1H, dd, *J* = 8.4, 2.4 Hz), 7.10 (1H, d, *J* = 7.2 Hz), 7.21 (1H, d, *J* = 2.4 Hz), 7.57 (1H, d, *J* = 10.4 Hz), 7.62-7.68 (1H, m), 7.91 (1H, d, *J* = 8.8 Hz), 7.99 (1H, dd, *J* = 8.8, 2.4 Hz), 8.64 (1 H, d, *J* = 2.4 Hz), 11.09 (1H, s).	19a

FIG. 2 Continued.



59		D	B	810.3	^{1}HNMR (400 MHz, CD_3OD) : δ 1.27 (t, *J* = 6.8 Hz, 3H), 1.89-1.93 (m, 2H), 2.11-2.13 (m, 1H), 2.70-2.74 (m, 7H), 2.77-2.78 (d, 8H), 3.43 (t, *J* = 6.4 Hz, 2H), 3.59 (t, *J* = 6.0 Hz, 2H), 4.31-4.37 (m, 1H), 4.40-4.44 (m, 1H), 4.55-4.60 (m, 3H), 5.05-5.08 (m, 1H), 6.69 (br, 1H), 6.77 (d, *J* = 2.4 Hz, 1H), 7.05-7.09 (m, 2H), 7.56-7.60 (m, 1H), 7.72 (d, *J* = 2.4 Hz, 1H), 7.78-7.80 (m, 1H), 7.89-7.91 (m, 1H), 8.02-8.03 (m, 1H).	20a
60		D	B	722.2	^{1}H NMR (300 MHz, DMSO) 8.57 (s, 1H), 8.05-7.94 (m, 4H), 7.87-7.84 (m, 1H), 7.53-7.48 (m, 1H), 7.35-7.32 (m, 1H), 7.26-7.20 (m, 2H), 7.04-7.01 (m, 1H), 5.05-4.81 (m, 2H), 4.57-4.50 (m, 1H), 4.41-4.22 (m, 2H), 4.14-4.07 (m, 3H), 2.92-2.77 (m, 4H), 2.65-2.55 (m, 4H) , 2.05-1.90 (m, 1H), 1.81-1.69 (m, 4H), 1.53-1.40 (m, 2H)	18a

FIG. 2 Continued.



61		D	B	736.3	^{1}H NMR (300 MHz, DMSO) 8.501 (s, 1H), 8.01-7.97 (m, 2H), 7.97-7.91 (m, 2H), 7.81-7.78 (m, 1H), 7.47-7.41 (m, 1H), 7.29-7.26 (m, 1H), 7.19-7.14 (m, 2H), 6.98-6.95 (m, 1H), 5.00-4.90 (m, 2H), 4.54-4.45 (m, 1H), 4.37-4.32 (m, 1H), 4.23-4.17 (m, 1H), 4.06-3.92 (m, 2H) , 2.81-2.75 (m, 1H), 2.70-2.68 (m, 2H), 2.66-2.56 (m, 3H), 2.49-2.35 (m, 3H), 2.01-1.90 (m, 1H), 1.70-1.50 (m, 4H), 1.49-1.37 (m, 4H);	18a
62		A	A	824.3	^{1}H NMR (400 MHz, CD_3OD): δ8.64-8.63 (d, *J*=2.0Hz, 1H), 8.18-8.15 (m, 1H), 7.81-7.68 (m, 4H), 7.13-7.11 (m, 2H), 6.99-6.96 (m, 1H), 5.14-5.09 (m, 1H), 4.60-4.51 (m, 2H), 4.29 (s, 1H), 4.15 (s, 1H), 3.83-3.81 (m, 2H), 3.72-3.65 (m, 2H), 3.59-3.56 (m, 2H), 3.46-3.44 (m, 4H), 3.31-3.29 (m, 2H), 2.88-2.82 (m, 3H), 2.77-2.70 (m, 2H), 2.14-2.11 (m, 1H), 1.82-1.65 (m, 4H), 1.29 (s, 6H), 1.21 (s, 6H);	8a

FIG. 2 Continued.



63		A	A	824.3	^{1}H NMR (400 MHz, CD_3OD): δ 8.60-8.59 (d, *J*=2.4Hz, 1H), 7.96-7.93 (m, 1H), 7.79-7.66 (m, 4H), 7.12-7.11 (d, J = 2.4Hz, 2H), 6.98-6.96 (m, 1H), 6.83-6.81 (d, J=9.2Hz, 1H), 5.13-5.09 (m, 1H), 4.27 (s, 1H), 4.13 (s, 1H), 3.67-3.60 (m, 6H), 3.52-3.49 (m, 2H), 2.87-2.82 (m, 3H), 2.76-2.68 (m, 8H), 2.14-2.10 (m, 1H), 1.80-1.76 (m, 2H), 1.67-1.61 (m, 2H), 1.28 (s, 6H), 1.21 (s, 6H);	8a
64		A	A	838.4	H-NMR (300 MHz, CD_3OD) δ 8.61 - 8.60 (d, *J* = 2.1 Hz, 1H)，7.98 - 7.94 (m, 1H), 7.81 - 7.67 (m, 4H), 7.14 - 7.13 (d, *J* = 2.4 Hz, 1H), 7.00 - 6.97 (m, 1H), 6.85 - 6.82 (d, *J* = 9 Hz, 1H), 5.14 - 5.08 (m, 1H), 4.28 (s, 1H), 4.14 (s, 1H), 3.68 - 3.59 (m, 6H), 3.50 - 3.45 (m, 2H), 2.87 - 2.70 (m, 5H), 2.65 - 2.62 (m, 6H), 2.71 - 1.69 (m, 1H), 1.78 - 1.72 (m, 2H), 1.66 - 1.62 (m, 2H), 1.47 - 1.44 (m, 2H), 1.29 (s, 6H), 1.22 (s, 6H).	8a

FIG. 2 Continued.



65		D	B	838.4	H-NMR (300 MHz, CD_3OD) δ 8.61 - 8.60 (d, *J* = 2.4 Hz, 1H), 7.98 - 7.95 (m, 1H), 7.81 - 7.67 (m, 4H), 7.14 - 7.13 (d, *J* = 2.4Hz, 1H), 7.01 - 6.97 (m, 1H), 6.86 - 6.83 (d, *J* = 8.7 Hz, 1H), 5.14 - 5.08 (m, 1H), 4.29 (s, 1H), 4.14 (s, 1H), 3.69 - 3.61 (m, 6H), 3.55 - 3.46 (m, 2H), 2.88 - 2.82 (m, 3H), 2.77 - 2.69 (m, 8H), 2.13 - 2.11 (m, 1H), 1.77 - 1.70 (m, 2H), 1.67 - 1.60 (m, 2H), 1.49 - 1.43 (m, 2H), 1.29 (s, 6H), 1.23 (s, 6H).	8a
66		A	A	837.4	H-NMR (300 MHz, CD_3OD, ppm) δ 7.80 - 7.66 (m, 6H), 7.14 (s, 1H), 7.00 - 6.98 (m, 3H), 5.14 - 5.08 (m, 1H), 4.29 (s, 1H), 4.14 (s, 1H), 3.64 - 3.60 (m, 2H), 3.50 - 3.46 (m, 2H), 3.39 - 3.32 (m, 4H), 2.92 - 2.80 (m, 3H), 2.77 - 2.65 (m, 8H), 2.14 - 2.08 (m, 1H), 1.80 - 1.62 (m, 4H), 1.52 - 1.39 (m, 2H), 1.29 (s, 6H), 1.23 (s, 6H).	8a

FIG. 2 Continued.



67		D	B	837.4	H-NMR (300 MHz, CD_3OD, ppm) δ 7.80 - 7.66 (m, 6H), 7.14 (s, 1H), 7.01 - 6.97 (m, 3H), 5.14 - 5.08 (m, 1H), 4.29 (s, 1H), 4.14 (s, 1H), 3.64 - 3.61 (m, 2H), 3.55 - 3.46 (m, 2H), 3.39 - 3.33 (m, 4H), 2.92 - 2.64 (m, 11H), 2.14 - 2.08 (m, 1H), 1.80 - 1.65 (m, 4H), 1.53 - 1.38 (m, 2H), 1.30 (s, 6H), 1.22 (s, 6H).	8a
68		D	B	793.4	H-NMR (300 MHz, CD_3OD) δ 7.81-7.79 (d, J = 6.6 Hz, 2H), 7.73-7.71 (d, J = 6.6 Hz, 2H), 7.43-7.37 (t, J = 18.9 Hz, 2H), 7.13-7.07 (q, J = 16.8 Hz, 3H), 6.99-6.96 (q, J = 8.4 Hz, 1H), 5.17-5.12 (q, J = 13.8 Hz, 1H), 4.47-4.45 (d, J = 5.7 Hz, 2H), 4.28 (s, 1H), 4.14 (s, 1H), 4.05-4.02 (d, J = 8.1 Hz, 2H), 3.68-3.66 (d, J = 5.7 Hz, 2H), 3.18-3.12 (m, 6H), 2.91-2.86 (t, J = 13.5 Hz, 1H), 2.81-2.77 (t, J = 11.1 Hz, 3H), 2.49-2.47 (d, J = 6.3Hz, 1H), 2.19 (s, 1H), 1.76-1.72 (t, J = 10.8 Hz, 4H), 1.45 (s, 4H) , 1.28 (s, 6H) , 1.22 (s, 6H);	8a

FIG. 2 Continued.



69		A	A	765.3	H-NMR (400 MHz, CD_3OD) δ 7.77-7.66 (m, 4H), 7.37-7.35 (d, *J*=8.4 Hz, 3H), 7.23-7.21 (d, *J*=7.6 Hz, 1H), 7.13-7.12 (d, *J*=2.4 Hz, 1H), 6.99-6.96 (m, 1H), 5.09-5.04 (m, 1H), 4.58 (s, 1H), 4.29 (s, 1H), 4.15 (s, 1H), 3.48-3.46 (m, 4H), 2.87-2.62 (m, 9H), 2.48-2.44 (m, 2H), 2.12-2.09 (m, 2H), 1.31 (s, 6H), 1.26 (s, 6H)	10A
70		A	A	827.4	^{1}H NMR (300 MHz, CD_3OD): δ87.94--7.87 (d, J=7.8Hz, 1H), 7.75--7.68 (d, J=7.8Hz 2H),7.67--7.48 (m,1H), 7.41--7.12 (m, 4H), 7.11--6.95(m, 2H), 5.18--4.99 (m, 1H), 4.58--3.99 (m , 4H),3.76--3.51 (m, 1H), 3.35 (s, 4H), 2.97--2.51(m, 8H), 2.50--2.26(m, 3H), 2.19--2.04 (m, 1H), 1.76--1.44 (m, 4H), 1.43--1.31 (m, 4H),1.28--1.17(m,12H)	21A

FIG. 2 Continued.



71		B	A	827.4	^{1}H NMR (300 MHz, CD_3OD): δδ7.94--7.87 (d, J=7.8Hz, 1H), 7.75--7.68 (d, J=7.8Hz 2H),7.67--7.48 (m,1H), 7.41--7.12 (m, 4H), 7.11--6.95(m, 2H), 5.18--4.99 (m, 1H), 4.58--3.99 (m , 4H),3.76--3.51 (m, 1H), 3.35 (s, 4H), 2.97--2.51(m, 8H), 2.50--2.26(m, 3H), 2.19--2.04 (m, 1H), 1.76--1.44 (m, 4H), 1.43--1.31 (m, 4H),1.28--1.17(m,12H)	21A
72		A	A	828.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.12 (6H, s), 1.22 (6H, s) 1.23-1.26 (2H, m), 1.33-1.40 (2H, m), 1.33-1.49 (4H, m), 1.68-1.73 (2H, m), 1.97-2.01 (1H, m), 2.30-2.47 (6H, m), 2.57-2.61 (1H, m), 2.87-2.93 (1H, m), 3.57 (3H, s), 4.06 (1H, d, J = 8.8 Hz), 4.21 (2H, t, J = 6.0 Hz), 4.30 (1H, s), 4.39 (1H, d, J = 17.2 Hz), 4.56 (1H, d, J = 17.6 Hz), 5.11 (1H, dd, J = 13.2, 4.2 Hz), 6.86 (1H, d, J = 8.8 Hz), 7.00 (1H, dd, J = 8.8, 1.8 Hz), 7.21 (1H, d, J = 2.4 Hz), 7.39-7.44 (2H, m), 7.63 (1H, d, J = 9.2 Hz), 7.91 (1H, d, J = 8.8 Hz), 7.96 (1H, d, J = 8.4 Hz), 8.63 (1H, s), 11.02 (1H, s).	22A

FIG. 2 Continued.



73		B	A	842.3	^{1}H NMR (400 MHz, DMSO-d_6) δ1.12 (6H, s), 1.22 (6H, s), 1.23-1.26 (3H, m),1.30-1.53 (8H, m), 1.78-1.81 (2H, m), 1.99-2.06 (2H, m), 2.18-2.24 (1H, m), 2.54-2.62 (2H, m), 2.87-2.89 (1H, m), 3.51-3.58 (3H, m), 4.06 (1H, d, *J* = 9.2 Hz), 4.27-4.31 (3H, m), 5.13 (1H, dd, *J* = 13.2, 5.6 Hz), 6.88 (1H, d, *J* = 8.8 Hz), 7.00 (1H, dd, *J* = 8.8, 2.4 Hz), 7.21 (1H, d, *J* = 2.4 Hz), 7.63 (1H, d, *J* = 8.8 Hz), 7.72 (1H, d, *J* = 6.8 Hz), 7.87 (1H, d, *J* = 9.6 Hz), 7.91 (1 H, d, *J* = 8.4 Hz), 7.97 (1 H, d, *J* = 9.2 Hz), 8.63 (1 H, d, *J* = 2.0 Hz), 11.13 (1H, s).	19A
74		B	A	827.4	^{1}H NMR (300 MHz, CD_3OD):δ7.94--7.87 (m, 1H), 7.76--7.68 (m, 2H),7.47--7.19 (m,7H), 5.17--5.05 (m, 1H), 4.53--4.06(m, 4H), 3.71 (s, 1H), 3.25(s , 4H),2.94--2.56 (m, 8H), 2.51--2.28 (m, 3H), 2.26--2.07(m, 1H), 1.72--1.48(m, 4H), 1.42(s, 4H), 1.25-1.22 (m, 8H), 1.21--1.18(m,6H),	21A

FIG. 2 Continued.


75		A	A	827.4	. ^{1}H NMR (300 MHz, CD_3OD): δ7.94--7.88 (m, 1H), 7.73--7.67 (m, 2H),7.45--7.17(m,7H), 5.16--4.88 (m, 1H), 4.59--4.09(m, 4H), 3.71--3.52 (m, 1H), 3.04--2.62(m , 7H),2.56--2.01 (m, 5H), 1.76--1.45 (m, 4H), 1.41--1.31(m, 4H), 1.25(s, 7H), 1.21(s, 6H),	21A

FIG. 3

Table 3. Exemplary Compounds


Ex. No.	Compound Structure	DC50 Code	M/Z +	1H NMR	Synthetic Scheme
76		A	827.3	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.21 (6H, s), 1.25 (6H, s), 1.37-1.49 (4H, m), 1.50-1.57 (2H, m), 1.62-1.74 (2H, m), 2.18-2.30 (2H, m), 2.32-2.41 (2H, m), 2.53-2.55 (4H, m), 2.81-2.93 (2H, m), 3.18-3.23 (2H, m), 3.64-3.68 (4H, m), 4.04 (1H, s), 4.13-4.24 (2H, m), 4.34-4.38 (2H, m), 5.15-5.20 (1H, m), 6.07 (1H, d, J = 8.4Hz), 6.64 (2H, m), 6.79-6.82 (1H, m), 6.96-6.97 (1H, d, J = 2.4 Hz), 7.42 (1H, d, J = 8.8 Hz), 7.57 (1H, d, J = 8.8 Hz), 7.91-7.94 (1H, m), 8.04 (1H, brs), 8.74 (1H, d, J = 2.0 Hz).	57

FIG. 3 Continued.


77		A	836.4	^{1}H NMR (400 MHz, δ6-DMSO): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.05 (m, 1H), 2.54-2.73 (m, 12H), 2.84-2.92 (m, 1H), 3.41-3.70 (m, 10H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.30 (s, 1H), 5.05-5.10 (m, 1H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (br, 1H), 7.63 (d, *J* = 9.2 Hz, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 7.96-7.99 (m, 1H), 8.63 (d, *J* = 2.0 Hz, 1H), 11.08 (s, 1H).	38

FIG. 3 Continued.




78		A	835.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.22 (s, 6H), 1.28 (s, 6H), 1.61-1.72 (m, 3H), 1.87-1.90 (m, 2H), 2.04-2.20 (m, 2H), 2.69-2.76 (m, 2H), 2.82-2.87 (m, 1H), 2.93-3.07 (m, 8H), 3.87 (br, 4H), 4.04-4.07 (m, 2H), 4.14-4.16 (m, 1H), 4.28 (s, 1H), 5.04-5.09 (m, 1H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.04-7.06 (m, 1H), 7.13 (d, *J* = 2.4 Hz, 1H), 7.20-7.23 (m, 1H),7.34-7.35 (m, 1H), 7.37-7.39 (m, 1H), 7.51-7.53 (m, 1H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.72 (d, *J* = 8.8 Hz, 1H), 7.87 (d, *J* = 7.8 Hz, 1H), 8.00-8.03 (m, 1H), 8.63 (d, *J* = 2.0 Hz, 1H).	58
79		A	824.6	H-NMR (300 MHz, CD_3OD) δ: (300 MHz, CD_3OD) δ 8.62 - 8.61 (d, *J* = 2.4 Hz, 1H), 8.00 - 7.96 (d, *J* = 11.4 Hz, 1H), 7.74 - 7.71 (d, *J* = 9 Hz, 1H), 7.65 - 7.62 (m, 1H), 7.48 - 7.46 (m, 2H), 7.14 - 7.13 (d, *J* = 2.4 Hz, 1H), 7.00 - 6.97 (d, *J* = 11.1 Hz, 1H), 6.86 - 6.83 (d, *J* = 9Hz, 1H), 5.20 - 5.14 (m, 1H), 4.51 - 4.49 (d,J = 5.4 Hz, 2H), 4.28 (s, 1H), 4.15 (s, 1H), 3.70 - 3.62 (m, 6H), 3.51 - 3.46 (m, 2H), 2.99 - 2.88 (m, 1H), 2.86 - 2.65 (m, 9H), 2.62 - 2.46 (m, 1H), 2.25 - 2.11 (m, 1H), 1.76 - 1.62 (m, 4H), 1.48 - 1.44 (m, 2H), 1.29 (s, 6H), 1.22 (s, 6H)	30

FIG. 3 Continued.

80		A	823.4	H-NMR: (300 MHz, CD_3OD) δ7.77 – 7.71 (m, 3H), 7.65 – 7.62 (m, 1H), 7.50 - 7.46 (m, 2H), 7.14 - 7.13 (d, *J* = 2.4 Hz, 1H), 7.00 – 6.97 (d, *J* = 9 Hz, 3H), 5.19 – 5.13 (m, 1H), 4.50 – 4.49 (d, *J* = 4.8 Hz, 2H), 4.29 (s, 1H), 4.14 (s, 1H), 3.64 - 3.61 (m, 2H), 3.50 - 3.46 (m, 2H), 3.32 - 3.31 (m, 3H), 2.90 - 2.71 (m, 11H), 2.55 - 2.49 (m, 1H), 2.20 - 2.19 (m, 1H), 1.76 - 1.62 (m, 4H), 1.51 – 1.44 (m, 2H), 1.29 (s, 6H), 1.23 (s, 6H);	30


81		A	804.6	^{1}H NMR (300 MHz, CD_3OD): δ 8.10-7.90 (m, 3H), 7.76-7.71 (m, 4H), 7.34-7.31 (m, 2H), 7.13 (s, 1H), 7.00-6.96 (m, 1H), 6.41(m, 1H), 5.19-5.13 (m, 1H), 4.29 (s, 1H), 4.17-4.14 (m, 2H), 3.90-3.60 (m, 3H), 3.26-3.23 (m, 2H), 2.96-2.70 (m, 7H), 2.20-2.10 (m, 1H), 1.81-1.72 (m, 4H), 1.47-1.38 (m, 3H), 1.28-1.23 (d, *J*=16.5Hz, 12H)	42

FIG. 3 Continued.



82		A	790.4	^{1}H NMR (300 MHz, CD_3OD): δ 7.75-7.60 (m, 6H), 7.33-7.30 (m, 2H), 7.13-7.12 (s, 1H), 7.00-6.96 (m, 1H), 6.31 (s, 1H), 5.18-5.12 (m, 1H), 4.49-4.48 (s, 2H), 4.29 (s, 1H), 4.10 (s, 1H), 3.23-3.22 (s, 2H), 2.91-2.46 (m, 11H), 2.19-2.10 (m, 1H), 1.69-1.60 (m, 4H), 1.40-1.30 (s, 4H), 1.28-1.23 (d, *J*=16.5Hz, 12H);	27
83		A	822.7	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.44 (s, 4H), 1.72-1.80 (m, 3H), 1.88-1.90 (m, 1H), 2.03-2.07 (m, 1H), 2.54-2.65 (m, 2H), 2.85-2.95 (m, 2H), 3.32-3.35 (m, 2H), 3.49-3.51 (m, 2H), 3.66 (s, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.16 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 4.69-4.75 (m, 1H), 5.09-5.14 (m, 1H), 6.51-6.54 (m, 1H), 7.00-7.02 (m, 1H), 7.20-7.21 (m, 1H), 7.32-7.34 (m, 1H), 7.41 (s, 1H), 7.55 (d, *J* = 9.2 Hz, 1H), 7.82 (d, *J* = 8.4 Hz, 1H), 7.89-7.93 (m, 2H), 8.16 (s, 1H), 8.59 (s, 1H), 11.11 (s, 1H).	45

FIG. 3 Continued.


84		A	795.4	H-NMR: (300 MHz, DMSO-*d6*, ppm) δ 7.89 - 7.86 (d, *J* = 9 Hz, 1 H), 7.75 7.72 (d, *J* = 9 Hz, 2 H),7.62 - 7.59 (d, *J* = 9 Hz,1 H), 7.52 - 7.49 (d, *J* = 9 Hz, 1 H), 7.19 - 7.14 (m, 2 H), 7.04 - 6.94 (m, 4 H), 5.07 - 5.01 (m, 1 H), 4.40 - 4.21 (m, 3 H), 4.08 - 4.02 (m, 3 H), 3.36 - 3.27 (m, 4 H), 2.88 - 2.79 (m, 1 H), 2.71 - 2.55 (m, 4 H), 2.38 - 2.27 (m, 2 H), 1.99 - 1.94 (m, 1 H), 1.79 - 1.74 (m, 2 H), 1.56 - 1.44 (m, 4 H), 1.21 - 1.19 (m, 7 H), 1.10 (s, 6 H);	31

FIG. 3 Continued.



85		A	806.35	H-NMR01-PH-ARV-LS-024-J-0: (300 MHz, DMSO-*d6*, ppm) δ 8.12 (br, 1 H), 7.89 - 7.79 (m, 5 H), 7.54 -7.51 (d, *J* = 9 Hz, 2 H), 7.47 - 7.41 (m, 1 H), 7.35 - 7.32 (m, 1 H), 7.22 - 7.18 (m, 1 H), 7.10 - 6.97 (m, 1 H), 6.28 (s, 1 H), 5.13 - 5.06 (m, 1 H), 4.31 (s, 1 H), 4.20 - 4.16 (m, 2 H), 4.07 - 4.04 (d, *J* = 9 Hz, 1 H), 3.29 - 3.24 (m, 2 H), 2.93 - 2.83 (m, 3 H), 2.60 - 2.56 (m, 5 H), 2.04 - 1.98 (m, 1 H), 1.81 - 1.76 (m, 2 H), 1.61 - 1.59 (m, 2 H), 1.48 - 1.45 (m, 2 H), 1.46 (s, 6 H), 1.21 (s, 6 H).	45, 50
86		A	779.75	^{1}H NMR (400 MHz, $CDCl_3$): δ 7.71-7.61(m, 4H), 7.58-7.56 (m, 1H), 6.99-6.91 (m, 2H), 6.83-9.79 (m, 2H), 6.22-6.19 (m, 1H), 5.30-5.28 (m, 1H), 4.76-4.74 (m, 1H), 4.17-4.15 (m, 1H), 4.06 (s, 1H), 3.86-3.20 (m, 4H), 2.98-2.89 (m, 2H), 2.72-2.40 (m, 6H), 2.23-2.17 (m, 2H), 1.70-1.62 (m, 3H), 1.44-1.39 (m, 3H),1.27 (s, 7H), 1.23 (s, 6H);	42

FIG. 3 Continued.


87		A	779.75	^{1}H NMR (400 MHz, $CDCl_3$): δ 7.71-7.61(m, 4H), 7.58-7.56 (m, 1H), 6.99-6.91 (m, 2H), 6.83-9.79 (m, 2H), 6.22-6.19 (m, 1H), 5.30-5.28 (m, 1H), 4.76-4.74 (m, 1H), 4.17-4.15 (m, 1H), 4.06 (s, 1H), 3.75-3.20 (m, 4H), 2.98-2.89 (m, 2H), 2.80-2.50 (m, 6H), 2.23-2.17 (m, 2H), 1.70-1.62 (m, 3H), 1.44-1.39 (m, 3H),1.27 (s, 7H), 1.23 (s, 6H);	42

FIG. 3 Continued.


88		A	811.45	^{1}H NMR (400 MHz, $CDCl_3$): δ 7.71-7.61(m, 4H), 7.58-7.56 (m, 1H), 6.99-6.91 (m, 2H), 6.83-9.79 (m, 2H), 6.22-6.19 (m, 1H), 5.30-5.28 (m, 1H), 4.76-4.74 (m, 1H), 4.17-4.15 (m, 1H), 4.06 (s, 1H), 3.75-3.20 (m, 4H), 2.98-2.89 (m, 2H), 2.80-2.50 (m, 6H), 2.23-2.17 (m, 2H), 1.70-1.62 (m, 3H), 1.44-1.39 (m, 3H),1.27 (s, 7H), 1.23 (s, 6H);	59

FIG. 3 Continued.


89		A	811.45	^{1}H NMR (300 MHz, CD_3OD): δ7.81--7.69 (m, 3H), 7.53--7.46 (m, 1H),7.42--7.28(m,3H), 7.15 (s, 1H),7.01--6.88(m,1H), 5.17--5.06(m, 1H), 4.52--4.41 (m, 2H), 4.27(s , 1H),4.16 (s, 1H), 3.75--3.57(m,4H),3.36(s,2H),3.33--3.12(m,4H),2.95--2.69(m,4H),2.59-2.35(m,1H),2.232.01(m,1H),1.89--1.65(m,4H),1.52--1.38(m,4H),1.33--1.16(m,12)	59

FIG. 3 Continued.


90		A	828.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.20 (6H, s), 1.22 (6H, s), 1.37-1.49 (6H, m), 1.79 (2H, t, *J* = 7.2 Hz), 1.98 (1H, t, *J* = 6.0 Hz), 2.29-2.43 (7H, m), 2.56-2.61 (1H, m), 2.87-2.91 (1H, m), 3.59 (4H, s), 4.05 (1H, d, *J* = 9.2 Hz), 4.14 (2H, t, *J* = 6.0 Hz), 4.24-4.40 (3H, m), 5.08 (1H, dd, *J* = 13.2, 4.8 Hz), 6.85 (1H, d, *J* = 8.8 Hz), 7.00 (1H, dd, *J* = 8.8, 2.0 Hz), 7.21 (1H, d, *J* = 2.0 Hz), 7.43 (1H, d, *J* = 6.8 Hz), 7.52 (1H, d, *J* = 10.0 Hz), 7.62 (1H, d, *J* = 9.2 Hz), 7.89-7.97 (2H, m), 8.62 (1H, d, *J* = 2.0 Hz), 11.00 (1H, s).	54, 45

FIG. 3 Continued.



91		A	781.25	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 8.52 -8.20 (brs, 1H), 7.85-7.81 (m, 4H), 7.55-7.46 (m, 3H), 7.40-7.30 (m, 1H), 7.09-6.92 (m, 3H), 5.10 (dd, J = 12.9, 5.4 Hz, 1H), 4.38 (d, J = 7.2 Hz, 1H), 4.28-4.14 (m, 2H), 4.06 (s, 1H), 3.16 (m, 3H), 3.04 (m, 3H), 2.87- 2.42 (m, 4H), 2.10-1.90 (m, 3H), 1.19-1.00 (m, 14H);	45
92		A	795.75	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 7.92 – 7.66 (m, 4H), 7.56 – 7.28 (m, 3H), 7.17 (d, J = 2.4 Hz, 1H), 7.02 – 6.88 (m, 3H), 5.08 (dd, J = 12.9, 5.4 Hz, 1H), 4.28 - 4.19 (t, J = 6.4 Hz, 3H), 4.02 (d, J = 9.1 Hz, 1H), 3.21 (t, J = 5.0 Hz, 4H), 2.90 – 2.76 (m, 1H), 2.62 – 2.46 (m, 6H), 2.37 (t, J = 7.2 Hz, 2H), 2.01 (dd, J = 11.3, 5.6 Hz, 1H), 1.76 (q, J = 9.6, 8.3 Hz, 2H),	45

FIG. 3 Continued.


				1.60 (t, *J* = 7.4 Hz, 2H), 1.18 (s, 6H), 1.09 (s, 6H).	
93		A	792.8	^{1}H NMR (300 MHz, CD_3OD): δ 7.76-7.71 (m, 4H), 7.49-7.31 (m, 4H), 7.13-7.12 (s, 1H), 7.00-6.96 (m, 1H), 6.31 (s, 1H), 5.18-5.11 (m, 1H), 4.48-4.47 (m, 2H), 4.29 (s, 1H), 4.15 (s, 1H), 3.32-3.31 (m, 2H), 2.90-2.36 (m, 10H), 2.20-2.10 (m, 1H), 2.03-1.88 (m, 4H), 1.70-1.65 (m, 4H), 1.40-1.30 (s, 4H), 1.28-1.23 (d, *J*=16.5Hz, 12H)	42

FIG. 3 Continued.



94		A	793.4	^{1}H NMR (300 MHz, CD_3OD): δ7.81--7.69 (m, 3H), 7.53--7.46 (m, 1H),7.42--7.28(m,3H), 7.15 (s, 1H),7.01--6.88(m,1H), 5.17--5.06(m, 1H), 4.52--4.41 (m, 2H), 4.27(s , 1H),4.16 (s, 1H), 3.75--3.57(m,4H),3.36(s,2H),3.33--3.12(m,4H),2.95--2.69(m,4H),2.59-2.35(m,1H),2.232.01(m,1H),1.89--1.65(m,4H),1.52--1.38(m,4H),1.33--1.16(m,12)	27
95		A	790.8	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.14 (m, 6H), 1.23 (s, 6H), 1.34-1.35 (m, 2H), 1.62-1.66 (m, 4H), 2.05-2.09 (m, 1H), 2.65-2.69 (m, 6H), 2.86-2.956 (m, 2H), 3.17 (d, *J* = 4.4 Hz, 1H), 4.07 (d, *J* = 8.8 Hz, 1H), 4.33 (s, 1H), 5.14-5.19 (m, 1H), 6.53 (s, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.32 (d, *J* = 8.4 Hz, 2H), 7.76-7.79 (m, 3H), 7.90-7.97(m, 4H), 11.14 (s, 1H).	1

FIG. 3 Continued.


96		A	796.4	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.21 (s, 6H), 1.25 (s, 6H), 1.50-1.57 (m, 2H), 1.66-1.75 (m, 2H), 1.82-1.89 (m, 2H), 2.31-2.42 (m, 2H), 2.57-2.62 (m, 2H), 2.68-2.97 (m, 6H), 3.79 (s, 4H), 4.02-4.15 (m, 4H), 4.27, 4.31 (two singles, 1H), 4.40, 4.44 (two singles, 1H), 5.18-5.23 (m, 1H), 6.07 (d, J = 8.4 Hz, 1H), 6.67 (d, J = 9.2 Hz, 1H), 6.79-6.82 (m, 1H), 6.93-7.03 (m, 1H), 7.11-7.19 (m, 1H), 7.31-7.43 (m, 2H), 7.57 (d, J = 8.8 Hz, 1H), 7.89-8.00 (m, 1H), 8.01-8.12 (m, 1H), 8.52-8.66 (m, 1H).	31

FIG. 3 Continued.



97		A	776.4	^{1}H NMR (400 MHz, CD_3OD):δ 1.25 (s, 6H), 1.31 (s, 6H), 1.66-1.77 (m, 4H), 2.13-2.18 (m, 1H), 2.61-2.65 (m, 2H), 2.71-2.81 (m, 6H), 2.85-2.91 (m, 3H), 3.28-3.31 (m, 2H), 4.18 (s, 1H), 4.31 (s, 1H), 5.14-5.18 (m, 1H), 6.44-6.46 (m, 1H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.15-7.16 (m, 1H), 7.36-7.38 (m, 2H), 7.73-7.79 (m, 3H), 7.86-7.88 (m, 1H), 7.91-7.93 (m, 1H), 7.97 (br, 1H).	
98		A	776.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.31 (s, 6H), 1.39，1.40 (d, *J* =6.4 Hz, 1H), 1.46-1.52 (m, 2H), 1.74-1.83 (m, 4H), 2.17-2.24 (m, 1H), 2.46-2.57 (m, 1H), 2.75-2.79 (t, *J* =7.6 Hz, 2H), 2.78-2.96 (m, 4H), 3.01-3.09 (m, 2H), 3.78 (s, 2H), 4.16-4.19 (m, 2H), 4.30 (s, 1H), 4.45-4.60 (m, 2H), 5.16-5.19 (m, 1H), 6.32 (s, 1H), 6.99-7.01 (m, 1H), 7.15 (d, *J* =2.4 Hz, 1H), 7.36 (d, *J* =8.4 Hz, 2H), 7.66-7.83 (m, 6H).	27

FIG. 3 Continued.


99		A	808.35	H-NMR01-PH-ARV-LS-024-H-0: (300 MHz, DMSO-*d6*, ppm) δ 8.14 (br, 1 H), 7.89 - 7.74 (m, 5 H), 7.47 7.32 (m, 4 H), 7.18 - 6.96 (m, 2 H), 5.13 - 5.06 (m, 1 H), 4.30 (s, 1 H), 4.20 - 4.16 (m, 2 H), 4.05 - 4.02 (d, *J* = 9 Hz, 1 H), 3.27 - 3.24 (m, 2 H), 2.91 - 2.81 (m, 1 H), 2.77 - 2.64 (m, 3 H), 2.60 - 2.54 (m, 3 H), 2.04 - 1.97 (m, 1 H), 1.87 - 1.74 (m, 6 H), 1.70 - 1.62 (m, 2 H), 1.47 - 1.45 (m, 2 H), 1.20 (s, 6 H), 1.11 (s, 6 H),	45

FIG. 3 Continued.


100		A	778.25	1H NMR (300 MHz, DMSO) 8.53 (s, 1H), 8.02-7.89 (m, 5H), 7.89-7.75 (m, 1H), 7.59-7.57 (m, 1H), 7.16-7.10 (m, 2H), 7.01-6.96 (m, 2H), 5.00-4.90 (m, 1H), 4.32-4.15 (m, 3H), 4.15-3.77 (m, 3H), 2.89-2.48 (m, 4H), 2.38-2.20 (m, 1H), 1.99-1.90 (m, 1H), 1.77-1.68 (m, 4H), 1.48-1.39 (m, 2H), 1.19-1.10 (m, 6H), 1.10-1.01 (m, 6H)	25

FIG. 3 Continued.


101		A	805.4	1H NMR (300 MHz, CDCl3) 7.81-755 (m, 5H), 7.25-7.22 (m, 1H), 6.96-6.90 (m, 1H), 6.82-6.79 (m, 1H), 6.59-6.57 (m, 1H), 6.39-6.30 (m, 1H), 6.19-6.17 (m, 1H), 5.30-5.19 (m, 2H), 4.74-4.70 (m, 1H), 4.28-4.20 (m, 1H), 4.14-4.10 (m, 1H), 4.04-4.00 (m, 1H), 3.67-3.50 (m, 1H), 3.40-3.30 (m, 2H), 3.06-2.90 (m, 2H), 2.63-2.49 (m, 6H), 2.19-1.94 (m, 4H), 1.65-1.44 (m, 4H), 1.40-1.05 (m, 16H)	42, 52

FIG. 3 Continued.


102		A	779.35	^{1}H NMR (300MHz, DMSO-d6): δ 8.97 (s, 1H), 8.69 (s, 1H), 8.51-8.47 (m, 1H), 8.25-8.16 (m, 2H), 7.94-7.88 (m, 1H), 7.63-7.57 (m, 1H), 7.26-6.85 (m , 4H), 5.13-4.97 (m, 1H), 4.51-3.99 (m, 6H), 2.98-2.22 (m, 5H), 2.07-1.89 (m, 1H), 1.87-1.73 (m, 4H), 1.58-1.43 (m, 2H), 1.31-1.08 (m, 13H)	48

FIG. 3 Continued.


103		A	807.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.32 (s, 6H), 1.49-1.51 (m, 4H), 1.78-1.84 (m, 4H), 2.00-2.09 (m, 2H), 2.13-2.23 (m, 2H), 2.73-2.95 (m, 6H), 3.10-3.18 (m, 4H), 3.63-3.71 (m, 2H), 4.20 (s, 1H), 4.31 (s, 1H), 5.16-5.18 (m, 1H), 6.99-7.01 (m, 1H), 7.15 (d, *J* =2.4 Hz, 1H), 7.46 (d, *J* =8.0 Hz, 1H), 7.74 (d, *J* =8.8 Hz, 1H), 7.79 (d, *J* =8.4 Hz, 1H), 7.86 (s, 1H), 7.89 (d, *J* =8.0 Hz, 1H), 8.16-8.18 (m, 1H), 8.89 (d, *J* =2.0 Hz, 1H).	11

FIG. 3 Continued.


104		A	762.5	^{1}H NMR (400 MHz, δ6-DMSO): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.05 (m, 1H), 2.54-2.73 (m, 12H), 2.84-2.92 (m, 1H), 3.41-3.70 (m, 10H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.30 (s, 1H), 5.05-5.10 (m, 1H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (br, 1H), 7.63 (d, *J* = 9.2 Hz, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 7.96-7.99 (m, 1H), 8.63 (d, *J* = 2.0 Hz, 1H), 11.08 (s, 1H).	11

FIG. 3 Continued.


105		A	833.4	H-NMR-PH-ARV-LS-026-D-0: (300 MHz, CD_3OD, ppm) δ 8.47 (s, 1H), 8.06 - 7.98 (m, 4H), 7.75 -7.71 (m, 2H), 7.18 - 7.11 (m, 3H), 7.02 - 6.98 (m, 1H), 5.15 - 5.09 (m, 1H), 4.70 - 4.57 (m, 3H), 4.45 - 4.44 (m, 1H), 4.31 (s, 1H), 4.20 (s, 1H), 3.11 - 3.08 (m, 2H), 2.92 - 2.75 (m, 7H), 2.51 - 2.45 (m, 1H), 2.20 - 2.01 (m, 7H), 1.32 (s, 6H), 1.25 (s, 6H).	10

FIG. 3 Continued.


106		A	806.32	^{1}H NMR (400 MHz, DMSO) 8.610 (s, 1H), 8.10-8.02 (m, 5H), 7.99-7.90 (m, 1H), 7.21-7.19 (m, 1H), 7.07-6.90 (m, 3H), 5.05-5.00 (m, 1H), 4.95-4.85 (m, 1H), 4.45-4.25 (m, 3H), 4.15-4.05 (m, 1H), 3.09-3.00 (m, 2H), 2.90-2.80 (m, 1H), 2.79-2.70 (m, 2H), 2.65-2.60 (m, 4H), 2.05-1.95 (m, 1H), 1.75-1.65 (m, 2H), 1.25-1.12 (m, 14H);	12, 13

FIG. 3 Continued.


107		A	819	H-NMR (400 MHz, CD_3OD) δ 8.69-8.64 (m, 2H), 8.63-8.40 (m, 2H), 8.13-8.11 (d, *J*= 8.0 Hz, 1H), 8.01-7.98 (m, 1H), 7.75-7.73 (d, *J*=8.4Hz, 1H), 7.15-7.15 (m, 1H), 7.01-6.99 (m, 1H), 6.90-6.88 (d, *J*=9.6, 1H), 5.24-5.20 (m, 1H), 4.30 (s, 1H), 4.17-4.16 (d, *J*=4.4z, 1H), 3.79-3.32 (m, 4H), 3.16-3.12 (m, 2H), 2.97-2.93 (m, 2H), 2.89-2.87 (d, *J*=6 Hz, 1H), 2.81-2.77 (m, 6H), 2.21-2.19 (m, 1H), 1.30 (s, 6H), 1.24 (s, 6H)	4

FIG. 3 Continued.





108		A	833.35	H-NMR (400 MHz, CD_3OD) δ8.62-8.60 (m, 2H), 8.41-8.35 (m, 2H), 8.13-8.11 (m, 1H), 8.02-7.99 (m, 1H), 7.71-7.71 (m, 1H), 7.14 (s, 1H), 7.01-6.94 (m, 1H), 6.85-6.83 (m, 1H), 5.25-5.15 (m, 1H), 4.29 (s, 1H), 4.12 (s, 1H), 3.80-3.75 (m, 4H), 2.94-2.72 (m, 11H), 2.20-2.02 (m, 3H), 1.28 (s, 6H), 1.22 (s, 6H);	7
109		A	808.85	^{1}HNMR (300 MHz, CD_3OD): δ 8.86 (s, 1H), 8.16-8.09 (m, 1H), 7.72-7.64 (m, 2H), 7.47-7.33 (m, 2H), 7.24-7.17 (m, 1H), 7.12 (s, 1H), 6.98-6.89 (m , 1H), 5.09-5.02 (m, 1H), 4.26 (s, 1H), 4.15(s, 1H), 3.48-3.42 (m, 4H), 2.92-2.49 (m, 9H), 2.46-2.35 (m, 2H), 2.16-2.01 (m, 1H), 1.81-1.71 (m, 2H),1.59-1.47 (m, 2H), 1.42-1.36 (m, 4H), 1.27-1.18 (m, 12H);	11

FIG. 3 Continued.

110		A	826.4	. ^{1}H NMR (300 MHz, DMSO): δ11.12 (s, 1H), 8.91 (s, 1H),8.18--7.88 (m,3H), 7.76--7.69 (d, J=7.8Hz,1H), 7.49--7.35(m, 2H), 7.22(s, 1H), 7.16--6.96 (m , 1H),5.17--5.06 (m, 1H), 4.32 (s, 1H), 4.11--3.98(m, 1H), 3.37--3.18(m, 4H), 2.97--2.75 (m, 3H), 2.62--2.51 (m, 4H), 2.48--2.37 (m, 2H),2.36--2.28(m,2H),2.12--1.07(m, 1H), 1.78--1.63 (m, 2H), 1.49--1.29 (m, 6H), 1.22 (s, 6H),1.13(s,6H	11
111		A	782.28	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.03 (m, 3H), 2.54-2.62 (m, 6H), 2.85-2.94 (m, 1H), 3.63 (br, 4H), 4.06 (d, *J* = 5.2 Hz ,1H), 4.24-4.31 (m, 3H), 5.10-5.15 (m, 1H), 6.88 (d, *J* = 9.2 Hz ,1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.0 Hz 1H), 7.36-7.38 (m, 1H), 7.45 (d, *J* = 1.6 Hz ,1H), 7.64 (d, *J* = 9.2 Hz,1H), 7.85 (d, *J* = 8.4 Hz ,1H), 7.91 (d, *J* = 8.4 Hz ,1H), 7.98 (d, *J* = 2.4 Hz ,1H), 8.64 (d, *J* = 1.6 Hz ,1H), 11.13 (s, 1H).	45

WO 2018/071606 PCT/US2017/056234

FIG. 3 Continued.



112		A	809.4	^{1}HNMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.19 (s, 6H), 1.46-1.55 (m, 2H), 1.63-1.73 (m, 2H), 1.78-1.86 (m, 2H), 1.88-2.07 (m, 5H), 2.59-2.84 (m, 7H), 2.90-2.95 (m, 1H), 3.31-3.41 (m, 2H), 4.08-4.12 (m, 2H), 4.19 (s, 1H), 4.52 (br, 1H), 4.99-5.03 (m, 1H), 6.89 (d, *J* = 8.8 Hz, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.22 (d, *J* = 8.4 Hz, 1H), 7.31 (d, *J* = 2.0 Hz, 1H), 7.37 (d, *J* = 8.0 Hz, 1H), 7.63 (d, *J* = 8.8 Hz, 1H), 7.71 (d, *J* = 8.4 Hz, 1H), 8.06 (d, *J* = 8.0 Hz, 1H), 8.80-8.81 (m, 1H).	45
113		A	781.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.13(s, 6H), 1.19 (s, 6H), 1.86-1.89 (m, 4H), 2.01-2.05(m, 3H), 2.20-2.26 (m, 2H), 2.60-2.68 (m, 4H), 2.76-2.82 (m, 2H), 3.12 (d, *J* = 11.9 Hz, 2H), 4.07-4.18 (m, 4H), 4.99-5.03 (m,1H), 6.88 (d, *J* = 8.8 Hz, 1H), 7.03 (d, *J* = 2.4 Hz ,1H), 7.23 (d, *J* = 8.4 Hz, 1H), 7.32-7.37 (m, 2H), 7.62 (d, *J* = 8.8 Hz ,1H), 7.71 (d, *J* = 8.0 Hz ,1H), 8.05 (d, *J* = 8.0 Hz ,1H), 8.78 (d, *J* = 2.0 Hz ,1H).	45

FIG. 3 Continued.



114		A	844.32		45
115		A	796.4	^{1}H NMR (400 MHz, CD_3OD): δ 0.88-0.93 (m, 2H), 1.24 (s, 6H), 1.30 (s, 6H), 1.78-1.96 (m, 4H), 2.02-2.18 (m, 2H), 2.63 (t, *J* = 7.2 Hz, 2H), 2.68-2.75 (m, 4H), 2.78-2.82 (m, 1H), 2.83-2.92 (m, 1H), 3.26-3.30 (m, 1H), 3.72-3.77 (m, 4H), 4.16 (s, 1H), 4.24 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 5.10-5.16 (m, 1H), 6.87 (d, *J* = 8.8 Hz, 1H), 6.98-7.02 (m, 1H), 7.15 (d, *J* = 2.4 Hz, 1H),7.32-7.36 (m, 1H), 7.44 (d, *J* = 8.8 Hz, 1H), 7.83 (d, *J* = 8.8 Hz, 1H), 7.97-8.01 (m, 1H),8.63 (d, *J* = 2.4 Hz, 1H).	47

FIG. 3 Continued.


116		A	839.45	H-NMR-PH-ARV-LS-024-F-0: (400 MHz, CD_3OD) δ 7.79 - 7.71 (m, 4H), 7.40 (s, 1H), 7.33 - 7.31 (d, *J* = 8Hz, 1H), 7.13 (s, 1H), 7.04 - 6.96 (m, 3H), 5.11 - 5.08 (m, 1H), 4.28 (s, 1H), 4.21 - 4.18 (m, 2H), 4.14 (s, 1H), 3.81 - 3.67 (m, 4H), 3.30 - 3.22 (m, 1H), 3.13 - 3.09 (m, 3H), 2.87 - 2.73 (m, 6H), 2.15 - 2.08 (m, 1H), 1.93 - 1.90 (m, 2H), 1.80 - 1.58 (m, 4H), 1.28 (s, 6H), 1.22 (s, 6H).	14, 15

FIG. 3 Continued.



117		A	806.25	^{1}H NMR (400 MHz, CDCl3): δ8.93 --8.81 (m, 1H), 8.40 (s, 1H),8.26 (s,2H), 8.18--8.12 (m, 1H), 7.82--7.69(m, 1H), 7.61--7.48 (m, 1H), 6.98 (s , 1H),6.89--6.78 (m,3H), 6.36--6.25 (m, 1H), 5.24--5.15(m, 1H), 4.96--4.82(m, 1H), 4.49--4.18(m, 3H),4.12--3.96(m,3H),3.31(s,1H),2.97--2.77(m,5H),2.43--2.15(m,2H),2.07--1.68(m,4H),1.31(s,6H),1.25(s,6H)	2
118		A	842.85	^{1}H NMR (300 MHz, CD_3OD): δ 8.85 (s, 1H), 8.17-8.08 (m, 1H), 7.93-7.88 (m, 1H), 7.71-7.66 (m, 1H), 7.43-7.29 (m, 3H), 7.26-7.17 (m, 2H), 5.11-5.02 (m, 1H),4.32 (s, 1H), 4.15 (s, 1H), 3.49-3.43 (m, 4H), 2.92-2.58 (m, 9H), 2.46-2.37 (m, 2H), 2.13-2.02 (m, 1H), 1.81-1.72 (m, 2H), 1.58-1.49 (m, 2H), 1.41-1.33 (m, 4H), 1.27-1.18 (m, 12H);	11

FIG. 3 Continued.



119		A	752.3	^{1}H NMR (300 MHz, CD_3OD): δ 8.89 (s, 1H), 8.15 (dd, J = 8.1, 2.3 Hz, 1H), 7.85 (d, J = 8.8 Hz, 1H), 7.70 (d, J = 8.7 Hz, 1H), 7.46 (d, J = 8.1 Hz, 1H), 7.23-7.15 (m, 1H), 7.15- 7.05 (m, 2H) , 6.95 (dd, J = 8.8, 2.4 Hz, 1H), 5.48 (s, 2H), 4.91 - 4.81 (m, 1H), 4.26 (d, J = 0.9 Hz, 1H), 4.16 (d, J = 0.9 Hz, 1H), 3.60 (t, J = 4.9 Hz, 4H), 3.21 (t, J = 5.0 Hz, 4H), 3.01 (dt, J = 17.9, 7.4 Hz, 4H), 2.80-2.69 (m, 2H), 2.29-2.03 (m, 5.8 Hz, 4H), 1.30 (s, 6H), 1.21 (s, 6H).	28
120		A	766.25	^{1}H NMR (300 MHz, CD_3OD): δ 8.92 (m, 1H), 8.14 (dd, J = 8.1, 2.3 Hz, 1H), 7.70 (dd, J = 8.6, 1.9 Hz, 2H), 7.50-7.45 (m, 1H), 7.40 (s, 1H),7.27 (dd, J = 8.6, 2.3 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 6.95 (dd, J = 8.8, 2.4 Hz, 1H), 5.05 (dd, J = 12.3, 5.4 Hz, 1H), 4.26 (s, 1H), 4.15 (s, 1H), 3.59 (t, J = 5.2 Hz, 4H), 3.12- 3.00 (m, 4H), 3.00-2.82 (m, 4H), 2.82-2.62 (m, 3H), 2.21 -2.02 (m, 3H), 1.24 (m, 6H) , 1.20 (s, 6H);	1

FIG. 3 Continued.



121		A	765.8	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.32 (s, 6H), 1.86-1.88 (m, 4H), 2.01-2.08 (m, 2H), 2.06-2.21 (m, 3H), 2.46-2.57 (m, 1H), 2.77-2.83 (m, 1H), 2.89-3.00 (m, 3H), 3.05-3.24 (m, 5H), 3.67-3.70 (m, 2H), 4.20 (s, 1H), 4.31 (s, 1H), 4.46-4.57 (m, 2H), 5.15-5.17 (m, 1H), 6.99-7.01 (m, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.46-7.54 (m, 3H), 7.75 (d, *J* = 8.8 Hz, 1H), 7.79 (d, *J* = 8.0 Hz, 1H), 8.17-8.19 (m, 1H), 8.92 (d, *J* = 1.6 Hz, 1H).	27
122		A	768.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.18 (s, 6H), 1.91-2.05 (m, 2H), 2. 61-2.68 (m, 5H), 2.72-2.78 (m, 1H), 2.84 (s, 2H), 3.60-3.64 (m, 3H), 4.04 (s, 1H), 4.18 (s, 1H), 4.26 (s, 2H), 4.47-4.51 (m, 1H), 4.99-5.04 (m, 1H), 6.72-6.78 (m, 1H), 6.86-6.91 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (d, *J* = 2.0 Hz, 1H), 7.61 (s, 1H), 7.70-7.75 (m, 1H), 7.84-7.89 (m, 1H), 8.48-8.54 (m, 1H).	47

FIG. 3 Continued.



123		A	779.35	(400 MHz, CDCl3) δ 8.94-8.94 (d, J=2.0 Hz, 1H), 8.07-8.04 (m, 1H), 7.86-7.78 (m, 3H), 7.68-7.66 (d, J=7.6 Hz, 1H), 7.61-7.59 (d, J=8.8 Hz, 1H), 7.31-7.29 (d, J=10.4 Hz, 1H), 7.00-6.99 (d, J=2.0 Hz, 1H), 6.85-6.82 (m, 1H), 6.36-6.34 (d, J=8.4 Hz, 1H), 5.02-4.98 (m, 1H), 4.20-4.18 (d, J=8.0 Hz, 1H), 4.10 (s, 1H), 3.41-3.38 (d, J=10.8 Hz, 2H), 2.96-2.74 (m, 8H), 2.43-2.40 (d, J=11.2 Hz, 2H), 2.20-2.15 (m, 3H), 1.98-1.95 (d, J=12.8 Hz, 2H), 1.88-1.77 (m, 4H), 1.31 (s, 6H), 1.26 (s, 6H)	1
124		A	780.4	H-NMR (400 MHz, CD3OD) δ 8.91-8.90 (d, J=2.0 Hz, 1H), 8.19-8.16 (m, 3H), 7.76-7.73 (m, 2H), 7.49-7.43 (m, 2H), 7.31-7.29 (m, 1H), 7.16-7.15 (d, J=2.4 Hz, 1H), 7.02-6.99 (m, 1H), 5.12-5.08 (m, 1H), 4.31 (s, 1H), 4.21 (s, 1H), 3.61-3.58 (m, 4H), 2.98-2.72 (m, 11H), 2.14-2.13 (d, J=5.2 Hz, 1H), 1.88-1.84 (m, 2H), 1.75-1.72 (m, 2H), 1.32 (s, 6H), 1.26 (s, 2H);	1

FIG. 3 Continued.



125		A	766.7	H-NMR (400 MHz, CD3OD) δ 8.91-8.91 (d, J=1.6 Hz, 1H), 8.34-8.17 (m, 3H), 7.86-7.84 (d, J=8.8 Hz, 1H), 7.76-7.73 (d, J=8.8 Hz, 1H), 7.49-7.47 (d, J=8.0 Hz, 1H), 7.18-7.15 (m, 2H), 7.09 (s, 1H), 7.02-6.99 (m, 1H), 5.44 (s, 2H), 4.85-4.81 (m, 1H), 4.31 (s, 1H), 4.21 (s, 1H), 3.62-3.58 (d, J=13.2 Hz, 4H), 3.17 (s, 4H), 3.01-2.95 (m, 4H), 2.79-2.76 (m, 2H), 2.26-2.14 (m, 2H), 1.87-1.80 (m, 4H), 1.32-1.25 (d, J=26 Hz, 12H);	1, 16
126		A	790.05		45

FIG. 3 Continued.


127		A	45

FIG. 3 Continued.



128		A	781.4	^{1}HNMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.27-1.43 (m, 5H), 1.65-1.71 (m, 2H), 2.01-2.07 (m, 1H), 2.53-2.60 (m, 4H), 2.75-2.79 (m, 3H), 2.83-2.94 (m, 2H), 4.05-4.07 (m, 2H), 4.30 (s, 1H), 5.08-5.16 (m, 2H), 6.98-7.01 (m, 1H), 7.19-7.22 (m, 1H), 7.28-7.37 (m, 3H), 7.84-7.90 (m, 2H), 7.99-8.08 (m, 2H), 8.88 (s, 1H), 11.11 (s, 1H).	83
129		A	821.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.04-1.06 (m, 1H), 1.10 (s, 6H), 1.20 (s, 6H), 1.75-1.85 (m, 3H), 1.98-2.02 (m, 1H), 2.18 (d, *J* = 6.4 Hz, 2H), 2.51-2.60 (m, 2H), 2.82-2.92 (m, 4H), 3.30-3.32 (m, 4H), 3.42-3.44 (m, 4H), 4.03 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1H), 4.37-4.43 (m, 2H), 5.03-5.08 (m, 1H), 6.82 (s, 1H), 6.97-7.00 (m, 1H), 7.19 (d, *J* = 2.0 Hz, 1H), 7.22-7.26 (m, 1H), 7.33 (s, 1H), 7.57 (d, *J* = 9.2 Hz, 1H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.87-7.93 (m, 2H), 8.60 (d, *J* = 2.4 Hz, 1H), 11.07 (s, 1H).	49

FIG. 3 Continued.



130		A	795.5	^{1}H NMR (400 MHz, CD_3OD): δ 1.23 (s, 6H), 1.30 (s, 6H), 1.38-1.38 (m, 1H), 1.64-1.69 (m, 2H), 1.77-1.84 (m, 2H), 1.93-1.95 (m, 2H), 2.11-2.15 (m, 3H), 2.68-2.77 (m, 5H), 2.83-2.98 (m, 5H), 4.18 (s, 1H), 4.29 (s, 1H), 4.74 (s, 1H), 5.09, 5.12 (d, *J* = 5.6 Hz, 1H), 6.97-6.99 (m, 1H), 7.13-7.14 (m, 1H), 7.33-7.35 (m, 1H), 7.43-7.47 (m, 2H), 7.71-7.74 (m, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 8.14-8.16 (m, 1H), 8.87 (d, *J* = 2.0 Hz, 1H).	83
131		A	835.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.43-1.44 (m, 2H), 1.62-1.63 (m, 1H), 1.75-1.78 (m, 2H), 2.01-2.03 (m, 1H),2.35-2.43 (m, 3H), 2.59-2.64 (m, 2H), 2.85-2.89 (m, 4H), 3.17-3.45 (m, 8H), 4.05-4.07 (m, 1H), 4.31 (s, 1H), 4.39-4.42 (m, 2H), 5.06-5.10 (m, 1H), 6.85 (d, *J* = 8.8 Hz, 1H), 7.00 (d, *J* = 8.4 Hz, 1H), 7.21-7.27 (m, 2H), 7.35 (s, 1H), 7.59-7.70 (m, 2H), 7.90-7.95 (m, 2H), 8.62 (s, 1H), 11.09 (s, 1H).	23

FIG. 3 Continued.


132		A	811.4	^{1}H-NMR (300 MHz, CD3OD) δ8.78 (s, 2H), 7.85- 7.79(m, 1H), 7.74- 7.68(m, 1H), 7.42- 7.39(m, 1H), 7.35 - 7.28(m, 1H), 7.14 - 7.10 (m, 1H), 7.01 - 6.96 (m, 1H), 5.15- 5.06(m, 1H), 4.29- 4.03(m, 8H), 2.95 - 2.69(m, 9H), 2.17 - 2.12 (m, 1H), 1.96 - 1.87 (m, 2H), 1.81 - 1.72 (m, 2H), 1.64 - 1.54 (m, 2H), 1.28 - 1.21 (d, *J = 20.4Hz* , 12H).	9

FIG. 3 Continued.



133		A	845.3	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.23 (s, 6H), 1.27 (s, 6H), 1.52-1.69 (m, 2H), 1.70-1.86 (m, 2H), 2.12-2.14 (m, 1H), 2.30-2.51 (m, 2H), 2.52-2.66 (m, 3H), 2.72-2.85 (m, 2H), 2.87-2.92 (m, 1H), 3.00 (t, *J* = 6.8 Hz, 2H), 3.31-3.46 (m, 3H), 3.51 (t, *J* = 6.0 Hz, 2H), 3.65 (t, *J* = 6.4 Hz, 2H), 4.07 (s, 1H), 4.13 (d, *J* = 8.0 Hz, 1H), 4.92-4.96 (m, 1H), 6.01-6.21 (m, 1H), 6.80 (dd, *J* = 2.4, 8.8 Hz, 1H), 6.96 (d, *J* = 2.0 Hz, 1H), 7.06 (dd, *J* = 2.0, 8.4 Hz, 1H), 7.28-7.33 (m, 2H), 7.57 (d, *J* = 8.4 Hz, 1H), 7.70 (d, *J* = 8.4 Hz, 1H), 8.02 (s, 1H).	84
134		A	837.3	^{1}H NMR (300 MHz, CD_3OD): δ11.08 (s, 1H), 7.84 (dd, J = 21.4, 8.5 Hz, 2H), 7.71 (d, J = 8.6 Hz, 2H), 7.50-7.39 (m, 2H), 7.33 (dd, J = 8.3, 2.3 Hz, 1H), 7.18 (d, J = 2.4 Hz, 1H), 7.03-6.88 (m, 3H), 5.09 (dd, J = 12.9, 5.4 Hz, 1H), 4.30 (s, 1H), 4.16 (t, J = 6.5 Hz, 2H), 4.02 (d, J = 9.1 Hz, 1H), 3.67 (d, J = 11.5 Hz, 2H), 2.92-2.78 (m, 1H), 2.72-2.50 (m, 5H), 2.42 (s, 1H), 2.34 (d, J = 27.9 Hz, 2H), 2.03 (dq, J = 13.5, 5.6, 4.6 Hz, 1H), 1.76 (dd, J = 8.1, 5.2 Hz, 2H), 1.39 (d, J = 12.5 Hz, 4H), 1.19	21

FIG. 3 Continued.



				(s, 6H), 1.14-0.95 (m, 12H);	
135		A	819.5	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.13 (s, 6H), 1.23 (s, 6H), 1.73-1.80 (m, 2H), 1.93-2.00 (m, 2H), 2.00-2.07 (m, 1H), 2.53-2.67 (m, 2H), 2.85-2.93 (m, 1H), 4.06-4.08 (m, 2H), 4.17-4.22 (m, 4H), 4.31 (s, 1H), 4.51 (br, 2H), 4.73 (s, 2H), 5.09-5.14 (m, 1H), 6.60-6.62 (m, 1H), 7.00-7.03 (m, 1H), 7.21-7.22 (m, 1H), 7.33-7.35 (m, 1H), 7.42-7.43 (m, 1H), 7.65-7.67 (m, 1H), 7.81-7.83 (m, 2H), 7.90-7.92 (m, 1H), 8.03-8.05 (m, 1H), 8.65-8.66 (m, 1H),11.12 (s, 1H).	55
136		A	819.53	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.13 (s, 6H), 1.23 (s, 6H), 1.73-1.80 (m, 2H), 1.93-2.00 (m, 2H), 2.00-2.07 (m, 1H), 2.53-2.67 (m, 2H), 2.85-2.93 (m, 1H), 4.06-4.08 (m, 2H), 4.17-4.22 (m, 4H), 4.31 (s, 1H), 4.51 (br, 2H), 4.73 (s, 2H), 5.09-5.14 (m, 1H), 6.60-6.62 (m, 1H), 7.00-7.03 (m, 1H), 7.21-7.22 (m, 1H), 7.33-7.35 (m, 1H), 7.42-7.43 (m, 1H), 7.65-7.67 (m, 1H), 7.81-7.83 (m, 2H), 7.90-7.92 (m, 1H), 8.03-8.05 (m, 1H), 8.65-8.66 (m, 1H),11.12 (s, 1H).	85

FIG. 3 Continued.



137		A	819.27	^{1}HNMR (400 MHz, CD_3OD): δ 1.14 (s, 6H), 1.21 (s, 6H), 1.72-2.04 (m, 6H), 2.60-2.75 (m, 4H), 2.88 (t, *J* = 7 Hz, 2H), 3.82 (s, 2H), 4.09-4.27 (m, 6H), 4.98-5.03 (m, 1H), 6.89 (d, *J* = 6.8 Hz, 1H), 7.04 (s, 1H), 7.25 (d, *J* = 8.8 Hz, 1H), 7.33 (s, 1H), 7.45 (s, 1H), 7.62 (d, *J* = 4 Hz, 1H), 7.70 (d, *J* = 8 Hz, 1H), 7.88 (d, *J* = 8.4 Hz, 1H), 8.21 (d, *J* = 8.8 Hz, 1H), 8.74 (s, 1H).	86
138		A	794.45	^{1}H NMR (300 MHz, DMSO-d6): *δ* 11.04 (s, 1H), 8.86 (d, J = 2.2 Hz, 1H), 8.10-8.00 (m, 2H), 8.00-7.80 (m, 1H),7.64 (d, J = 8.4 Hz, 1H), 7.39-7.14 (m, 4H), 6.97 (dd, J = 8.7, 2.4 Hz, 1H), 5.03 (dd, J = 12.8, 5.3 Hz, 1H), 4.28 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.38 (t, J = 5.0 Hz, 4H), 3.00-2.73 (m, 3H), 2.73-2.51 (m, 2H), 2.51-2.39 (m, 4H), 2.27 (t, J = 7.3 Hz, 2H), 2.02-1.93 (m, 1H), 1.69-1.63 (m, 2H), 1.47 (p, J = 7.4 Hz, 2H), 1.30 (q, J = 8.0 Hz, 2H), 1.20 (s, 6H), 1.09 (s, 6H)	8

FIG. 3 Continued.



139		A	812.55	^{1}H NMR (400 MHz, CDCl3) 8.845 (s, 1H), 7.97-7.91 (m, 2H), 7.51-7.49 (m, 1H), 7.08-7.05 (m, 1H), 6.90-6.89 (m, 1H), 6.75-6.72 (m, 1H), 6.19-6.17 (m, 1H), 5.13-5.09 (m, 1H), 4.36-4.32 (m, 1H), 4.22-4.18 (m, 1H), 4.10-4.08 (m, 1H), 4.00-3.98 (m, 1H), 3.17-3.10 (m, 4H), 2.83-2.62 (m, 7H), 2.60-2.09 (m, 4H), 2.00-1.76 (m, 2H), 1.75-1.68 (m, 3H), 1.32-1.16 (m, 16H)	20
140		A	812.5	^{1}H NMR (400 MHz, CDCl3) 8.85 (s, 1H), 7.97-7.88 (m, 2H), 7.51-7.42 (m, 2H), 6.93-6.89 (m, 2H), 6.75-6.72 (m, 1H), 6.18-6.15 (m, 1H), 5.12-5.08 (m, 1H), 4.36-4.00 (m, 4H), 3.32-3.20 (m, 4H), 2.87-2.59 (m, 7H), 2.30-2.13 (m, 2H), 1.79-1.50 (m, 7H), 1.41-1.16 (m, 16H)	11, 16

FIG. 3 Continued.



141		A	821.35	^{1}H NMR (400 MHz, CD_3OD): δ 7.77-7.50 (m, 5H), 7.38-7.34 (m, 3H), 7.23-7.13 (m, 2H), 7.00-6.96 (m, 1H), 5.10-5.00 (m, 1H), 4.29 (s, 1H), 4.17-4.14 (m, 1H), 3.76-3.67 (m, 4H), 3.36-3.32 (m, 2H), 2.87-2.42 (m, 11H), 2.10-2.00 (m, 3H), 1.70-1.60 (m, 2H), 1.28-1.23 (m, 12H)	39
142		A	797.4	1H NMR (300 MHz, DMSO-d6) δ 11.08 (s, 1H), 8.73 (s, 2H), 7.83 (dd, J = 18.2, 8.5 Hz, 2H), 7.69 (d, J = 9.2 Hz, 1H), 7.41 (d, J = 2.2 Hz, 1H), 7.33 (dd, J = 8.3, 2.3 Hz, 1H), 7.18 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.5 Hz, 1H), 5.08 (dd, J = 12.8, 5.4 Hz, 1H), 4.29 – 4.13 (m, 3H), 4.00 (d, J = 9.1 Hz, 1H), 3.79 (t, J = 4.7 Hz, 4H), 2.95 – 2.76 (m, 1H), 2.56 (d, J = 16.4 Hz, 2H), 2.45 – 2.30 (m, 6H), 2.00 (dd, J = 12.3, 6.2 Hz, 1H), 1.69 (dq, J = 46.8, 7.0 Hz, 4H), 1.18 (s, 6H), 1.08 (s, 6H).	9

FIG. 3 Continued.




143		A	783.55	1H NMR (300 MHz, DMSO-d6) δ 11.08 (s, 1H), 8.75 (s, 2H), 8.21-7.63 (m, 3H), 7.60- 7.16 (m, 2H), 7.18 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.5 Hz, 1H), 5.09 (dd, J = 12.8, 5.3 Hz, 1H), 4.25 (d, J = 7.4 Hz, 3H), 4.01 (d, J = 9.1 Hz, 1H), 3.28 (s, 4H), 2.89-2.76 (m, 1H), 2.56 (d, J = 19.0 Hz, 2H), 2.24 (s, 6H), 1.99 (s, 3H), 1.13 (d, J = 30.8 Hz, 12H)	9
144		A	769.3	1H NMR (300 MHz, DMSO-d6) δ 11.09 (s, 1H), 10.18 (s, 1H), 8.83 (s, 2H), 7.93-7.74 (m, 3H), 7.54 (d, J = 2.2 Hz, 1H), 7.42 (dd, J = 8.3, 2.3 Hz, 1H), 7.27-7.14 (m, 1H), 7.03-6.86 (m, 1H), 5.10 (dd, J = 12.8, 5.4 Hz, 1H), 4.88-4.47 (m, 4H), 4.27 (s, 1H), 4.02 (d, J = 9.1 Hz, 1H),3.92- 3.62 (s, 4H), 3.30 (d, J = 54.5 Hz, 4H), 2.92-2.77 (m, 1H), 2.57 (d, J = 19.1 Hz, 2H), 2.08-1.97 (m, 1H), 1.18 (s, 6H), 1.08 (s, 6H).	9
145		A	822.57	^{1}H NMR (300 MHz, CD_3OD): δ 8.70 (s, 2H), 7.71-7.66 (m, 2H), 7.36 (m, 1H), 7.24-7.21 (m, 1H), 7.10-7.09 (m, 1H), 6.97-6.93 (m, 1H), 5.08-5.02 (m, 1H), 4.90-4.87 (m, 2H),4.80 (s, 1H), 4.28-4.24 (m, 2H), 4.10 (s, 1H), 3.51 (m, 4H),	49

FIG. 3 Continued.



				3.03-2.95 (m, 2H), 2.78-2.67 (m, 6H), 2.44-2.42 (m, 2H), 2.15-1.80 (m, 4H), 1.75-1.55 (m, 1H), 1.25-1.18 (m, 12H);	
146		A	800.21	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.98-2.07 (m, 2H), 2.11-2.16 (m, 1H), 2.46-2.51(m, 2H), 2.57-2.66 (m, 3H), 2.70-2.87 (m, 3H), 2.88-2.96 (m, 3H), 3.40-3.49 (m, 4H), 4.11 (s, 1H), 4.18 (d, *J* = 8.0 Hz, 1H), 4.92-4.97 (m, 1H), 6.20 (d, *J* = 8.0 Hz, 1H), 7.03-7.08 (m, 2H),7.24 (d, *J* = 2.4 Hz, 1H), 7.28-7.32 (m, 2H), 7.70 (d, *J* = 8.4 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.98-8.11 (m, 2H), 8.92 (d, *J* = 2.0 Hz, 1H).	1
147		A	781.51	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.42-1.58 (m, 4H), 1.98-2.04 (m,1H), 2.26-2.34 (m, 2H), 2.40-2.48 (m, 6H), 2.62-2.70 (m, 3H), 2.82-2.94 (m, 1H), 3.36-3.52 (m, 7H), 4.24-4.50 (m, 5H), 5.02-5.10 (m, 1H), 5.76 (s, 1H), 6.53 (s, 1H), 6.93 (s, 1H), 7.23 (d, *J* = 8.4 Hz, 1H), 7.31 (s, 1H), 7.67 (d, *J* = 8.8 Hz, 1H), 7.98 (s, 2H), 8.07 (s, 1H), 8.26-8.34 (m, 1H), 11.08 (s, 1H), 13.28 (s, 1H).	17a

FIG. 3 Continued.



148		A	835.8	^{1}H NMR (300 MHz, CD3OD) 7.76-7.64 (m, 4H), 7.35-7.32 (m, 3H), 7.23-7.20 (m, 1H), 7.13-7.12 (m, 1H), 7.00-6.96(m, 1H), 5.09-5.03 (m, 1H), 4.28-4.25 (m, 1H), 4.15-4.11 (m, 1H), 3.77-3.68 (m, 4H), 3.38-3.32 (m, 2H), 2.86-2.69 (m, 5H), 2.41-2.29 (m, 6H), 2.11-2.03 (m, 3H), 1.86-1.64 (m, 4H), 1.28-1.23 (m, 12H);	39
149		A	801.5	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.07 (s, 1H), 7.91 (d, J = 8.8 Hz, 1H), 7.71 (dd, J = 19.0, 8.4 Hz, 3H), 7.46 (d, J = 9.2 Hz, 1H), 7.35 (d, J = 2.4 Hz, 1H), 7.27 (dd, J = 8.7, 2.3 Hz, 1H), 7.22 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.4 Hz, 1H), 6.44 (d, J = 8.4 Hz, 2H), 5.08 (dd, J = 12.9, 5.4 Hz, 1H), 4.32 (s, 1H), 4.06 (d, J = 9.1 Hz, 1H), 3.91 (t, J = 7.3 Hz, 2H), 3.64 (ddd, J = 24.3, 9.7, 5.1 Hz, 3H), 3.43 (dt, J = 18.6, 4.7 Hz, 6H), 3.03 – 2.82 (m, 4H), 2.64 – 2.51 (m, 2H),	37

FIG. 3 Continued.


				2.40 (t, *J* = 4.8 Hz, 4H), 2.02 (dt, *J* = 15.2, 5.6 Hz, 1H), 1.22 (s, 6H), 1.13 (s, 6H).	
150		A	836.6	^{1}H-NMR (300 MHz, CDCl3) δ8.69 (s, 2H), 8.01(m, 1H), 7.75-7.69(m, 1H), 7.62- 7.53(m, 1H), 7.29(m, 1H), 7.10 - 7.04(m, 1H), 6.98 - 6.91 (m, 1H), 6.84 - 6.78 (m, 1H), 5.95- 5.88(m, 1H), 4.98- 4.81(m, 3H), 4.15- 4.01(m, 2H), 3.54(m, 4H), 3.11 - 2.60(m, 11H), 2.17 - 2.09 (m, 1H), 1.87 - 1.61(m, 5H), 1.24 - 1.21 (m, 14H)	32

FIG. 3 Continued.



151		A	810.23		14a
152		A	815.32		17

FIG. 3 Continued.



153		A	869.28		17
154		A	824.21	^{1}HNMR (400 MHz, DMSO-d_6) : δ 1.12 (s, 6H), 1.22 (s, 6H), 1.39-1.44 (m, 2H), 1.55-1.61 (m, 2H), 1.75-1.81 (m, 2H), 2.02-2.06 (m, 1H), 2.54-2.69 (m, 2H), 2.85-2.94 (m, 1H), 3.39 (t, *J* = 7.2 Hz, 2H), 3.46 (t, *J* = 5.0 Hz, 2H), 3.88 (t, *J* = 5.2 Hz, 2H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.16-4.19 (m, 4H), 4.31 (s, 1H), 5.10-5.14 (m, 1H), 6.87 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.32-7.35 (m, 1H), 7.43 (s, 1H), 7.66-7.68 (m, 1H), 7.82 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.01-8.03 (m, 1H), 8.66 (d, *J* = 2.0 Hz, 1H), 11.12 (m, 1H),.	87

FIG. 3 Continued.






155		A	852.25	^{1}H NMR (400 MHz, CD_3OD): δ 1.17 (s, 6H), 1.24 (s, 6H), 1.43-1.46 (m, 2H), 1.56-1.59 (m, 2H), 2.07-2.10 (m, 1H), 2.42 (t, *J* = 7.2 Hz, 2H), 2.54 (t, *J* = 4.0 Hz, 4H), 2.62-2.73 (m, 4H), 2.79-2.84 (m, 1H), 3.32-3.35 (m, 2H), 3.86-3.91 (m, 6H), 4.10 (s, 1H), 4.23 (s, 1H), 5.01 (t, *J* = 5.2 Hz, 1H), 5.05-5.10 (m, 1H), 6.93-6.95 (m, 1H), 7.09 (d, *J* = 20.0 Hz, 1H), 7.20-7.25 (m, 2H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.78 (d, *J* = 8.4 Hz, 1H), 8.71 (s, 2H).	73
156		A	838.23	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.11 (s, 6H), 1.21 (s, 6H), 1.49-1.60 (m, 2H), 2.01-2.08 (m, 1H), 2.29-2.49 (m, 7H), 2.53-2.64 (m, 3H), 2.85-2.94 (m, 1H), 3.08-3.26 (m, 2H), 3.85 (s, 6H), 4.04 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1H), 5.06-5.16 (m, 2H), 6.99-7.02 (m, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.26-7.32 (m, 2H), 7.75 (d, *J* = 9.2 Hz, 1H), 7.86 (d, *J* = 9.2 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.77 (s, 2H), 11.14 (s, 1H).	73

FIG. 3 Continued.

157		A	784.1	H-NMR-PH-ARV-LS-031-M-0: (300 MHz, DMSO- *d6*, ppm) δ 11.10 (br, 1H), 8.20 -8.17 (d, *J* = 9Hz, 1H), 8.02 (s, 1H), 7.74 - 7.66 (m, 2H), 7.53 - 7.47 (m, 1H), 7.34 - 7.15 (m, 4H), 5.11 - 5.05 (m, 1H), 3.42 - 3.35 (m, 4H), 2.91 - 2.83 (m, 1H), 2.71 - 2.60 (m, 4H), 2.56 - 2.49 (m, 4H), 2.35 - 2.30 (m, 2H), 2.08 - 2.00 (m, 1H), 1.67 - 1.65 (m, 2H), 1.62 (s, 8H), 1.60 - 1.37 (m, 2H).	43
158		A	820.35	^{1}H NMR (300 MHz, $CDCl_3$) δ 7.95 (s, 1H), 7.70-7.55 (m, 4H), 7.30 (s, 1H), 7.26-6.97 (m, 1H), 6.96-6.79 (m, 4H), 6.12-6.09 (m, 1H), 4.96-4.94 (m, 1H), 4.16-4.04 (m, 2H), 3.89-3.39 (m, 5H), 3.20-2.50 (m, 8H), 2.49-1.60 (m, 6H), 1.26-1.20 (m, 15H);	49
159		A	796.65	H-NMR06-PH-ARV-LS-031-A-0: (300 MHz, CD3OD, ppm) δ 7.96 - 7.93 (d, *J* = 9 Hz, 1 H), 7.83 (s, 1 H), 7.69 - 7.61 (m, 2 H), 7.51 - 7.48 (m, 1 H), 7.35 (s, 1 H), 7.24 – 7.15 (m, 3 H), 5.09 - 5.05 (m, 1 H), 3.48 - 3.45 (m, 4H), 2.82 - 2.42(m, 15 H), 2.12 - 2.09 (m, 2 H), 1.76 - 1.73 (m, 2 H), 1.66 - 1.61 (m, 3 H), 1.48 - 1.46 (m, 2 H).	43

WO 2018/071606 PCT/US2017/056234

FIG. 3 Continued.


160		A	821.35	^{1}H NMR (300 MHz, DMSO-d_6) δ 10.70 (s, 1H), 8.62 (d, J = 2.4 Hz, 1H), 7.92 (dd, J = 10.0, 6.5 Hz, 2H), 7.56 (dd, J = 17.5, 8.9 Hz, 2H), 7.22 (d, J = 2.4 Hz, 1H), 7.03 (ddd, J = 14.6, 8.6, 2.1 Hz, 3H), 6.86 (d, J = 9.1 Hz, 1H), 5.31 (s, 2H), 4.57 (dd, J = 9.1, 5.2 Hz, 1H), 4.41 (d, J = 13.3 Hz, 2H), 4.31 (s, 1H), 4.06 (d, J = 9.2 Hz, 1H), 3.29 (t, J = 4.9 Hz, 4H), 2.96 – 2.81 (m, 2H), 2.61 (t, J = 5.6 Hz, 2H), 2.54 – 2.50 (m, 4H), 2.38 (t, J = 7.4 Hz, 2H), 2.00 (qd, J = 8.2, 3.9 Hz, 2H), 1.77 (d, J = 12.7 Hz, 2H), 1.63 (s, 1H), 1.43 (q, J = 7.2 Hz, 2H), 1.22 (s, 6H), 1.12 (s, 8H).	41

FIG. 3 Continued.



161		A	827.45	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.04 (s, 1H), 8.71 (s, 2H), 8.02 (d, J = 8.8 Hz, 1H), 7.71 – 7.57 (m, 3H), 7.36 – 7.17 (m, 3H), 5.03 (dd, J = 12.7, 5.3 Hz, 1H), 4.69 (d, J = 12.8 Hz, 2H), 4.30 (s, 1H), 4.01 (d, J = 9.0 Hz, 1H), 3.28 (s, 4H), 3.00 – 2.76 (m, 3H), 2.67 (dt, J = 3.7, 1.9 Hz, 4H), 2.41 (t, J = 7.3 Hz, 2H), 2.34 (t, J = 7.3 Hz, 2H), 2.03 – 1.92 (m, 1H), 1.81 – 1.69 (m, 3H), 1.46 – 1.33 (m, 2H), 1.19 (s, 6H), 1.07 (s, 8H)	17
162		A	764.3	H-NMR (300 MHz, DMSO) δ 11.05 (s, 1H), 8.64-8.63 (d, J=2.4 Hz, 1H), 7.98-7.89 (m, 2H), 7.67-7.60 (m, 2H), 7.22-7.21 (d, J=2.4 Hz, 1H), 7.03-6.92 (m, 2H), 6.80-6.79 (d, J=1.8Hz, 1H), 6.69-6.65 (m, 1H), 5.09-5.03 (m, 1H), 4.31 (s, 1H), 4.08-4.05 (d, J=9 Hz, 1H), 3.85 (s, 4H), 3.67 (s, 4H),2.73-2.72 (m, 1H), 2.60 (s, 2H), 2.01-1.99(d, J=5.1 Hz, 1H), 1.82 (s, 4H), 1.22 (s, 6H), 1.12(s, 6H);	44

FIG. 3 Continued.



163		A	764.35	H-NMR (300 MHz, DMSO) δ 11.04 (s, 1H), 8.58 (s, 1H), 7.95-7.86 (m, 2H), 7.66-7.63 (d, J=8.4Hz, 1H), 7.57-7.54 (d, J=9.3Hz, 1H), 7.35-7.18 (m, 3H), 7.00-6.96 (m, 1H), 6.39-6.36 (d, J=9Hz, 1H), 5.08-5.02 (m, 1H), 4.28 (s, 1H), 4.05-4.02 (d, J=9.3Hz, 1H), 3.80 (s, 4H), 3.51 (s, 4H), 2.92-2.81 (m, 1H), 2.70-2.59 (m, 2H), 2.05-1.98 (m, 4H), 1.19 (s, 6H), 1.10 (s, 6H);	19
164		A	833.54	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.24 (s, 6H), 1.38-1.47 (m, 2H), 1.74-1.90 (m, 4H), 1.96-2.10 (m, 2H), 2.54-2.65 (m, 2H), 2.85-2.94 (m, 1H), 3.33 (s, 1H), 4.04-4.11 (m, 1H), 4.14-4.19 (m, 3H), 4.32 (s, 1H), 4.55 (s, 3H), 5.07-5.18 (m, 1H), 6.62 (d, *J* = 8.8 Hz, 1H), 6.96-7.08 (m, 1H), 7.23 (d, *J* = 2.4 Hz, 1H), 7.31-7.38 (m, 1H), 7.43 (d, *J* = 1.6 Hz, 1H), 7.66 (s, 2H), 7.79-7.86 (m, 1H), 7.89-7.95 (m, 1H), 7.98-8.05 (m, 1H), 8.69 (s, 1H), 11.08-11.19 (m, 1H).	85

FIG. 3 Continued.



165		A	740.46	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.21 (s, 6H), 2.03-2.15 (m, 2H), 2.58 (m, 2H), 3.86-3.89 (m, 2H), 4.04 (d, *J* = 8.8 Hz, 1H), 4.23-4.29 (m, 5H), 5.10-5.15 (m, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.35-7.38 (m, 1H), 7.45 (d, *J* = 2.0 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.84-7.91 (m, 2H), 8.74 (s, 2H), 11.12 (s, 1H).	88
166		A	831.53	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.61-1.64 (m, 2H), 1.76-1.82 (m, 2H), 2.01-2.07 (m, 1H), 2.38 (s, 2H), 2.44-2.49 (m, 4H), 2.57 (s, 1H), 2.85-2.94 (m, 1H), 3.24 (s, 4H), 3.32 (s, 1H), 4.00 (d, *J* = 9.2 Hz, 1H), 4.20-4.26 (m, 3H), 5.10-5.15 (m, 1H), 6.66 (d, *J* = 11.6 Hz, 2H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 8.4, 2.4 Hz, 1H), 7.44 (d, *J* = 2.4 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.19 (d, *J* = 9.2 Hz, 1H), 11.13 (s, 1H).	89

FIG. 3 Continued.



167		A	836.5	^{1}HNMR (400 MHz, DMSO-d_6) : δ 1.11 (s, 6H), 1.22 (s, 6H), 1.73-1.83 (m, 5H), 1.99-2.02 (m, 1H), 2.29 (s, 2H), 2.57 (s, 6H), 2.74 (s, 2H), 2.91-2.97 (m, 3H), 3.82-3.86 (m, 4H), 4.01-4.03 (m, 3H), 4.29 (s, 1H), 5.06-5.09 (m, 1H), 7.01 (d, *J* = 8.0 Hz, 1H), 7.22-7.30 (m, 3H), 7.64 (d, *J* = 7.6 Hz, 1H), 7.75 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 8.77 (s, 2H), 11.10 (s, 1H),.	49
168		A	837.75	H-NMR-PH-ARV-LS-032-D-0: (300 MHz, DMSO-d6,ppm) δ 11.11 (s, 1H), 7.86 (dd, *J* = 22.3, 8.5 Hz, 2H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 – 7.43 (m, 2H), 7.36 (dd, *J* = 8.3, 2.3 Hz, 1H), 7.20 (d, *J* = 2.4 Hz, 1H), 7.05 – 6.90 (m, 3H), 5.11 (dd, *J* = 13.0, 5.2 Hz, 1H), 4.31 (s, 3H), 4.04 (d, *J* = 9.1 Hz, 1H), 3.65 (s, 2H), 3.10 (d, *J* = 11.5 Hz, 2H), 2.89 (s, 1H), 2.69 (s, 5H), 2.37 (s, 3H), 2.04 (d, *J* = 11.3 Hz, 1H), 1.85 (s, 2H), 1.58 (s, 2H), 1.21 (s, 6H), 1.12 (s, 6H).	47

FIG. 3 Continued.


169		A	851.25	H-NMR-PH-ARV-LS-032-F-0: (300 MHz, DMSO-d6,ppm) (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 7.84 (dd, J = 19.4, 8.5 Hz, 2H), 7.69 (d, J = 8.6 Hz, 2H), 7.49 – 7.26 (m, 3H), 7.17 (d, J = 2.3 Hz, 1H), 7.02 – 6.87 (m, 3H), 5.09 (dd, J = 12.8, 5.3 Hz, 1H), 4.31 – 4.14 (m, 3H), 4.01 (d, J = 9.1 Hz, 1H), 3.62 (t, J = 4.6 Hz, 2H), 3.44 (d, J = 12.6 Hz, 2H), 3.07 (t, J = 11.3 Hz, 2H), 2.91 – 2.76 (m, 1H), 2.47 (m, 2H), 2.42 – 2.30 (m,4H), 2.23 (s, 2H), 2.01-1.88 (m, 5H), 1.55 (dd, J = 12.6, 8.4 Hz, 2H), 1.18 (s, 6H), 1.09 (s, 6H).	47

FIG. 3 Continued.



170		A	769.3	^{1}H NMR (300 MHz, $CDCl_3$) 8.12 (s, 1H), 7.73-7.71 (m, 2H), 7.60-7.25 (m, 5H), 6.97-6.96 (m, 1H), 6.82-6.79 (m, 1H), 6.21-6.18 (m, 1H), 4.96-4.90 (m, 1H), 4.17-4.14 (m, 1H), 4.05-4.00 (m, 1H), 3.48-3.35 (m, 4H), 2.93-2.68 (m, 10H), 2.16-2.12 (m, 1H) , 1.37-1.23 (m, 13H);	1
171		A	763.6	^{1}H NMR (300 MHz, CD_3OD) δ 7.770-7.66 (m, 3H), 7.60 (d, *J* = 8.5 Hz, 1H), 7.31 (d, *J* = 8.1 Hz, 2H), 7.10 (d, *J* = 2.4 Hz, 1H), 7.04-6.89 (m, 2H), 6.84 (dd, *J* = 8.6, 2.2 Hz, 1H), 5.02 (dd, *J* = 12.3, 5.4 Hz, 1H), 4.54 (s, 1H), 4.25 (s, 1H), 4.11 (s, 1H), 3.75 (s, 1H), 3.46 (d, *J* = 2.5 Hz, 2H), 3.07-2.96 (m, 1H), 2.88-2.61 (m, 8H), 2.13-1.87 (m, 3H), 1.24 (s, 6H), 1.19 (s, 6H);	1

FIG. 3 Continued.



172		A	834.56		
173		A	856.56	^{1}HNMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.21-1.22 (m, 6H), 1.52-1.57 (m, 1H), 1.60-1.69 (m, 2H), 1.98-2.07 (m, 2H), 2.12-2.23 (m, 1H), 2.35-2.43 (m, 2H), 2.55-2.68 (m, 4H), 2.83-2.93 (m, 1H), 3.28-3.34 (m, 3H), 3.40 (s, 4H), 3.52-3.57 (m, 1H), 3.65-3.69 (m, 2H), 4.05-4.08 (m, 1H), 4.31 (s, 1H), 5.06-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.21-7.28 (m, 2H), 7.36 (s, 1H), 7.51-7.54 (m, 2H), 7.63-7.70 (m, 2H), 7.90-7.92 (m, 1H), 11.09 (s, 1H).	49

FIG. 3 Continued.



174		A	834.55	1H NMR (300 MHz, DMSO-d6) δ 11.04 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.67 (dd, J = 15.3, 8.5 Hz, 3H), 7.44 (d, J = 9.2 Hz, 1H), 7.30 (d, J = 2.4 Hz, 1H), 7.27-7.13 (m, 2H), 7.02-6.87 (m, 3H), 5.03 (dd, J = 12.7, 5.3 Hz, 1H), 4.28 (s, 1H), 4.01 (d, J = 9.1 Hz, 1H), 3.81 (d, J = 12.4 Hz, 2H), 3.28 (s, 5H), 2.93-2.69 (m, 4H), 3.0-2.6 (m, 4H), 2.54 (d, J = 16.1 Hz, 5H), 2.35 (t, J = 7.2 Hz, 2H), 2.03-1.93 (m, 1H), 1.73 (d, J = 12.8 Hz, 2H), 1.40 (d, J = 7.3 Hz, 3H), 1.18 (s, 8H), 1.09 (s, 6H).	23
175		A	832.3	^{1}H NMR (400 MHz, CD_3OD) δ 7.74 (dd, *J* = 8.8, 2.7 Hz, 3H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.07-7.04 (m, 1H), 7.00 (td, *J* = 6.3, 3.1 Hz, 3H), 6.93-6.85 (m, 1H), 5.07 (dd, *J* = 12.6, 5.5 Hz, 1H), 4.57 (d, *J* = 2.3 Hz, 1H), 4.30 (s, 1H), 4.15 (s, 1H), 3.90 (d, *J* = 12.2 Hz, 2H), 3.72 (s, 1H), 3.51 (s, 2H), 3.08 (d, *J* = 9.6 Hz, 1H), 2.94-2.70 (m, 5H), 2.66 (d, *J* = 10.0 Hz, 1H), 2.51 (d, *J* = 6.9 Hz, 2H), 2.17-2.03 (m, 2H), 1.99-1.83 (m, 3H), 1.66 (s, 1H), 1.29 (s, 8H), 1.24 (s, 6H)	35

FIG. 3 Continued.



176		A	832.3	^{1}H NMR (400 MHz, CD_3OD) δ 7.73 (dd, *J* = 8.9, 2.8 Hz, 3H), 7.66 (d, *J* = 8.5 Hz, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.05 (s, 1H), 6.99 (td, *J* = 6.3, 3.1 Hz, 3H), 6.89 (d, *J* = 8.5 Hz, 1H), 5.07 (dd, *J* = 12.5, 5.4 Hz, 1H), 4.57 (s, 1H), 4.30 (s, 1H), 4.15 (s, 1H), 3.90 (d, *J* = 12.3 Hz, 2H), 3.71 (s, 1H), 3.51 (s, 2H), 3.07 (d, *J* = 9.7 Hz, 1H), 2.90-2.70 (m, 5H), 2.66 (d, *J* = 10.0 Hz, 1H), 2.50 (d, *J* = 6.9 Hz, 2H), 2.17-2.03 (m, 2H), 1.99-1.83 (m, 3H), 1.66 (s, 1H), 1.29 (s, 8H), 1.24 (s, 6H)	35
177		A	838.57	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.62-1.67 (m, 2H), 1.82-1.88 (m, 2H), 2.00-2.07 (m, 1H), 2.11-2.15 (m, 1H), 2.24-2.28 (m, 1H), 2.56-2.58 (m, 1H), 2.61-2.62 (m, 1H), 2.85-2.93 (m, 2H), 3.14-3.20 (m, 2H), 3.22-3.28 (m, 4H), 3.62-3.65 (m, 4H), 4.06 (d, *J* = 8.8 Hz, 1H), 4.24-4.27 (m, 2H), 4.32 (s, 1H), 5.08-5.13 (m, 1H), 6.72 (t, *J* = 8.8 Hz, 1H), 7.21 (d, *J* = 2.4Hz, 1H), 7.36-7.39 (m ,1H), 7.51-7.55 (m, 2H), 7.60-7.67 (m, 2H), 7.77 (d, *J* = 8.8 Hz ,1H), 7.91 (d, *J* = 8.80 Hz, 1H), 9.65 (br, 1H), 11.11 (s, 1H).	49

FIG. 3 Continued.





178		A	835.2	^{1}H NMR (400 MHz, CD_3OD): δ7.80-7.78 (d, *J*=8.4Hz, 1H), 7.73-7.70 (m, 3H), 7.25-7.19 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.07 (m, 1H), 4.91-4.89 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.92-3.88 (m, 2H), 3.13-3.12 (m, 1H), 2.87-2.65 (m, 6H), 2.47-2.44 (m, 2H), 2.34-2.29 (m, 2H), 2.18-2.13 (m, 6H), 1.92-1.89 (m, 2H), 1.80-1.70 (m, 1H), 1.32-1.31 (m, 1H), 1.28 (s, 6H), 1.21 (s, 6H);	51
179		A	839.57		49
180		A	838.57		49
181		A	856.56		49

FIG. 3 Continued.




182		A	824.45	1H-NMR (300 MHz, DMSO-d6) δ11.10（s, 1H), 7.93- 7.86 (m, 1H), 7.77- 7.69 (m, 3H), 7.50- 7.34 (m, 2H), 7.22- 7.18 (m, 1H), 7.04- 6.96 (m, 1H), 6.58- 6.51 (m , 2H), 5.16 - 5.06 (m, 1H), 4.32(s, 1H), 4.09- 4.02 (m, 1H), 3.55- 3.20 (m, 6H), 3.09- 2.73 (m, 2H), 2.70- 2.38 (m, 10H), 2.20- 2.01 (m, 2H), 1.86- 1.67 (m, 1H), 1.24 - 1.13 (d, *J*=33 Hz, 12H).	46
183		A	819.58	^{1}H NMR (400 MHz, DMSO-d^6) δ 0.97-1.07 (2H, m), 1.13 (6H, s), 1.23 (6H, s), 1.49-1.69 (5H, m), 1.80-2.02 (4H, m), 2.19-2.20 (1H, m), 2.33-2.43(1H, m), 2.50-2.60(2H, m), 2.81-2.91(1H, m), 3.30-3.35 (4H, m), 3.40-3.45 (4H, m), 4.06 (1H, d, *J* = 9.2 Hz), 4.32 (1H, s), 5.03-5.07 (1H, m), 6.99-7.02 (1H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.27 (1H, d, *J* = 8.8 Hz), 7.33-7.39 (3H, m), 7.68 (1H, d, *J* = 8.4 Hz),7.73-7.77 (3H, m), 7.91 (1H, d, *J* = 8.8 Hz), 11.09 (1H, s).	90
184		A	799.63	^{1}H NMR (400 MHz, DMSO-d^6) δ 0.97-1.07 (2H, m), 1.13 (6H, s), 1.23 (6H, s), 1.46-1.69 (5H, m), 1.81-2.02 (4H, m), 2.19-2.20 (1H, m), 2.39-2.41(1H, m), 2.45 (3H, m), 2.53-2.59(2H, m), 2.81-2.89(1H, m), 3.30-3.35 (4H, m), 3.42-3.48 (4H, m), 4.05 (1H, d, *J* = 9.2 Hz), 4.23 (1H, s), 5.02-5.07 (1H, m), 6.80-6.83 (1H, m), 7.25-7.40 (4H, m), 7.67-7.78 (5H, m), 11.10 (1H,	90

FIG. 3 Continued.



				s).	
185		A	824.35	^{1}H NMR (300 MHz, $CDCl_3$) 7.83-7.80 (m, 1H), 7.63-7.39 (m, 4H), 7.19-7.05 (m, 1H), 6.90-6.71 (m, 4H), 6.04-6.01 (m, 1H), 5.14-5.08 (m, 1H), 4.38-4.07 (m, 3H), 3.97-3.95 (m, 1H), 3.81-3.30 (m, 4H), 3.19-2.98 (m, 3H), 2.89-2.55 (m, 6H), 2.31-2.18 (m, 4H), 1.82-1.61 (m, 2H), 1.19-1.15 (m, 15H);	20, 33
186		A	824.35	^{1}H NMR (300 MHz, $CDCl_3$) 7.81-7.80 (m, 1H), 7.62-7.60 (m, 2H), 7.53-7.48 (m, 2H), 6.90-6.71 (m, 5H), 6.04-6.01(m, 1H), 5.13-5.07 (m, 1H), 4.37-4.06 (m, 3H), 3.97-3.41 (m, 5H), 3.12-3.06 (m, 3H), 2.88-2.42 (m, 7H), 2.30-2.15 (m, 4H), 1.82-1.75 (m, 1H), 1.19-1.05 (m, 15H);	41

FIG. 3 Continued.



187		A	838.38	^{1}H NMR (300 MHz, DMSO) 10.96 (s, 1H), 7.91-7.88 (m, 1H), 7.74-7.71 (m, 2H), 7.48-7.41 (m, 2H), 7.24-7.20(m, 2H), 7.01-6.93 (m, 3H), 5.09-5.03 (m, 1H), 4.39-4.21 (m, 3H), 4.06-4.03 (m, 1H), 3.86-3.82 (m, 2H), 3.30-3.28 (m, 1H), 3.11-3.05 (m, 4H), 2.90-2.72 (m, 3H), 2.61-2.50 (m, 4H), 2.42-2.35 (m, 3H), 2.07-1.95 (m, 1H), 1.78-1.75 (m, 2H), 1.51-1.42 (m, 3H), 1.30-1.05 (m, 14H);	33
188		A	838.4	^{1}H NMR (300 MHz, DMSO-*d6*) 10.96 (s, 1H), 7.91-7.88 (m, 1H), 7.74-7.70 (m, 2H), 7.48-7.20 (m, 4H), 7.02-6.93(m, 3H), 5.09-5.05 (m, 1H), 4.39-4.21 (m, 3H), 4.06-4.03 (m, 1H), 3.96-3.82 (m, 2H), 3.30-3.28 (m, 1H), 3.11-3.05 (m, 4H), 2.90-2.72 (m, 3H), 2.70-2.62 (m, 4H), 2.39-2.26 (m, 3H), 2.07-1.79 (m, 1H), 1.75-1.70(m, 2H), 1.52-1.46 (m, 3H), 1.31-1.05 (m, 14H);	33

FIG. 3 Continued.



189		A	806.4	^{1}H NMR (300 MHz, DMSO-d_6): δ11.04 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.67 (dd, J = 13.3, 8.3 Hz, 3H), 7.44-7.13 (m, 4H), 7.02-6.91 (m, 1H), 6.38 (d, J = 8.3 Hz, 2H), 5.04 (dd, J = 12.4, 5.2 Hz, 1H), 4.28 (s, 1H), 3.99 (q, J = 8.1, 7.6 Hz, 3H), 3.50 (t, J = 6.5 Hz, 2H), 3.41 (d, J = 6.4 Hz, 4H), 2.96-2.47 (m, 4H), 2.47 (d, J = 3.4 Hz, 2H), 2.30 (d, J = 7.4 Hz, 2H), 2.09-1.90 (m, 1H), 1.85- 1.71 (m, 2H), 1.18 (s, 6H), 1.09 (s, 6H).	34
190		A	821.35	^{1}H NMR (300 MHz, DMSO): δ 11.11 (s, 1H), 7.91-7.81 (m, 2H), 7.73 (d, J = 9.0 Hz, 2H), 7.48 (d, J = 9.1 Hz, 1H), 7.30-7.27 (m, 2H), 7.20 (m, 1H), 7.02-6.90 (m, 3H), 5.18-5.00 (m, 2H), 4.32 (s, 1H), 4.05 (d, J = 9.0 Hz, 1H), 3.89-3.70 (m, 4H), 3.05-2.98 (m, 2H), 2.98-2.81 (m, 1H), 2.80-2.69 (m, 2H), 2.64-2.56 (m, 2H), 2.49-2.40 (m, 2H), 2.10-1.93 (m, 1H), 1.80-1.79 (m, 2H), 1.60-1.44 (m, 1H), 1.32-1.14 (m, 10H), 1.12 (s, 6H)	23

FIG. 3 Continued.


191		A	836.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.22 (s, 6H), 1.58-1.70 (m, 2H), 1.81-2.05 (m, 6H), 2.10-2.18 (m, 2H), 2.80-2.92 (m, 1H), 3.34-3.76 (m, 7H), 3.53-3.67 (m, 3H), 3.72-3.89 (m, 3H), 3.98-4.08 (m, 2H), 4.27-4.33 (m, 1H), 5.05-5.14 (m, 1H), 7.01 (s, 1H), 7.23-7.33 (m, 3H), 7.67-7.78 (m, 2H), 7.91 (m, 1H), 8.76-8.77 (m, 2H), 11.09 (s, 1H).	49

FIG. 3 Continued.





192		A	805.56	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.13 (6H, s), 1.22 (6H, s), 1.37-1.57 (4H, m), 1.86-2.03 (5H, m), 2.32-2.42 (2H, m), 2.54-2.63 (2H, m), 2.67 (4H, brs), 2.84-2.91 (1H, m), 3.44 (4H, brs), 4.05 (1H, d, J = 9.2 Hz), 4.31 (1H, s), 5.07 (1H, dd, J = 5.6, 9.2 Hz), 7.00 (1H, dd, J = 2.0, 8.8 Hz), 7.21 (1H, d, J = 2.4 Hz), 7.25-7.27 (1H, m), 7.33-7.35 (3H, m), 7.67-7.69 (1H, m), 7.70-7.75 (3H, m), 7.89-7.92 (1H, m), 11.10 (1H, s).	91
193		A	810.35	^{1}H NMR (300 MHz, CD_3OD): δ 8.60 (s, 1H), 7.99-7.95 (m, 1H), 7.83-7.70 (m, 2H), 7.42-7.31 (m, 2H), 7.13-7.12(s, 1H), 7.00-6.86 (m, 2H), 5.09-5.08 (m, 1H), 4.37-4.14 (m, 6H), 3.19-3.13 (m, 2H), 2.89-2.73 (m, 7H), 2.20-2.03 (m, 3H), 1.29-1.22 (m, 18H);	3

FIG. 3 Continued.

194		A	809.2	^{1}H NMR (300 MHz, Methanol-d_4): δ 7.82 (d, J = 8.3 Hz, 1H), 7.79-7.70 (m, 3H), 7.42 (d, J = 2.2 Hz, 1H), 7.33 (dd, J = 8.3, 2.3 Hz, 1H), 7.14 (d, J = 2.4 Hz, 1H), 7.05-6.95 (m, 3H), 5.12 (dd, J = 12.3, 5.4 Hz, 1H), 4.29 (d, J = 0.9 Hz, 1H), 4.21 (t, J = 5.8 Hz, 2H), 4.14 (d, J = 0.8 Hz, 1H), 3.77 (s, 1H), 3.73 (s, 1H), 3.15-3.05 (m, 2H), 2.96-2.69 (m, 5H), 2.60 (t, J = 11.3 Hz, 2H), 2.19-2.08 (m, 1H), 2.03 (d, J = 8.6 Hz, 2H), 1.29 (s, 6H), 1.26 (s, 3H), 1.23 (d, J = 2.1 Hz, 9H).	21
195		A	811.3	^{1}H NMR (400 MHz, DMSO): δ 11.11 (s, 1H), 8.76 (s, 2H), 7.90 (d, J = 8.7 Hz, 1H), 7.84 (d, J = 8.3 Hz, 1H), 7.72 (d, J = 9.2 Hz, 1H), 7.43 (d, J = 2.3 Hz, 1H), 7.36-7.33 (m, 1H), 7.22 (m, 1H), 7.02-6.99 (m, 1H), 5.14-5.09 (m, 1H), 4.53 (d, J = 11.9 Hz, 2H), 4.29 (s, 1H), 4.20 (m, 2H), 4.03 (d, J = 9.1 Hz, 1H), 2.97-2.83 (m, 3H), 2.74 (m, 2H), 2.65-2.52 (m, 3H), 2.49 (m, 1H), 2.11-2.01 (m, 1H), 1.83 (m, 2H), 1.22 (s, 6H), 1.11 (m, 12H)	3

WO 2018/071606 PCT/US2017/056234

FIG. 3 Continued.



196		A	807.54	H-NMR (400 MHz, $CDCl_3$) δ 8.23 (s, 1H), 7.79-7.77 (d, *J*=8.4 Hz, 1H), 7.71-7.68 (d, *J*=9.0 Hz, 2H), 7.58-7.55 (d, *J*=8.7 Hz, 1H), 7.37-7.36 (d, *J*=1.8 Hz, 1H), 7.22-7.19 (d, *J*=8.4 Hz, 1H), 6.97-6.91 (m, 3H), 6.83-6.79 (m, 1H), 6.12-6.10 (d, *J*=8.1 Hz, 1H), 4.98-4.92 (m, 4H), 3.34 (s, 3H), 2.93-2.68 (m, 4H), 2.54 (s, 4H), 2.36 (s, 2H), 2.18-2.13 (m, 1H), 1.62 (s, 4H), 1.26 (s, 6H), 1.22 (s, 6H)	
197		A	769.5		

FIG. 3 Continued.



198		A	861.6		36
199		A	834.59		49

FIG. 3 Continued.




200		A	834.59		49
201		A	834.59		49
202		A	779.35	^{1}H NMR (300 MHz, CD3OD) 7.78-7.68 (m, 4H), 7.37-7.34 (m, 3H), 7.26-7.23 (m, 1H), 7.13-7.11 (m, 1H), 7.00-6.96(m, 1H), 5.08-5.04 (m, 1H), 4.29-4.27 (m, 1H), 4.16-4.14 (m, 1H), 3.92-3.89 (m, 2H), 3.11-2.71 (m, 11H), 2.13-2.11 (m, 1H), 1.31-1.28 (m, 18H);	22

FIG. 3 Continued.



203		A	833.25	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.07 (s, 1H), 7.91 (d, J = 8.8 Hz, 1H), 7.71 (dd, J = 19.0, 8.4 Hz, 3H), 7.46 (d, J = 9.2 Hz, 1H), 7.35 (d, J = 2.4 Hz, 1H), 7.27 (dd, J = 8.7, 2.3 Hz, 1H), 7.22 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.4 Hz, 1H), 6.44 (d, J = 8.4 Hz, 2H), 5.08 (dd, J = 12.9, 5.4 Hz, 1H), 4.32 (s, 1H), 4.06 (d, J = 9.1 Hz, 1H), 3.91 (t, J = 7.3 Hz, 2H), 3.64 (ddd, J = 24.3, 9.7, 5.1 Hz, 3H), 3.43 (dt, J = 18.6, 4.7 Hz, 6H), 3.03 – 2.82 (m, 4H), 2.64 – 2.51 (m, 2H), 2.40 (t, J = 4.8 Hz, 4H), 2.02 (dt, J = 15.2, 5.6 Hz, 1H), 1.22 (s, 6H), 1.13 (s, 6H).	36
204		A	850.25	^{1}H NMR (400 MHz, CD_3OD): δ7.78-7.70 (m, 4H), 7.26-7.20 (m, 2H), 7.13-7.12 (d, J=2.0Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.07 (m, 1H), 4.93-4.89 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.32-3.30 (m, 4H), 3.28-3.26 (m, 1H), 2.87-2.67 (m, 7H), 2.58-2.46 (m, 6H), 2.39-2.34 (m, 2H), 2.25 (s, 3H), 2.14-2.11 (m, 1H), 1.28 (s, 6H), 1.21 (s, 6H);	51

FIG. 3 Continued.



205		A		^{1}H NMR (300 MHz, DMSO-*d6*) 11.12 (s, 1H), 7.92-7.73 (m, 4H), 7.22-7.19 (m, 3H), 7.20-6.95 (m, 3H), 5.20-5.11 (m, 1H), 4.32 (s, 1H), 4.20-4.16 (m, 2H), 4.07-4.04 (m, 1H), 3.47-3.41 (m, 2H), 3.32-3.25 (m, 4H), 2.96-2.94 (m, 6H), 2.61-2.50 (m, 7H), 2.10-2.01 (m, 1H), 1.21 (s, 6H), 1.23 (s, 6H);	93
206		A	806.56		

FIG. 3 Continued.




207		A	779.25	^{1}H NMR (300 MHz, CD_3OD) 7.78-7.64 (m, 4H), 7.39-7.32 (m, 3H), 7.21-7.18 (m, 2H), 7.12-6.95 (m, 1H), 5.09-5.03(m, 1H), 4.28-4.27 (m, 1H), 4.15-4.13 (m, 1H), 3.52-3.47 (m, 2H), 3.29-3.22 (m, 4H), 3.18-2.63 (m, 7H), 2.13-2.05 (m, 1H), 1.28-1.11 (m, 18H);	22
208		A	806.55		49
209		A	779.1	1H NMR (300 MHz, DMSO-d6) δ 11.03 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.73 (dd, J = 8.6, 6.7 Hz, 3H), 7.61 (d, J = 8.5 Hz, 1H), 7.38-7.26 (m, 3H), 7.25-7.14 (m, 2H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 5.03 (dd, J = 12.6, 5.3 Hz, 1H), 4.29 (s, 1H), 4.03 (d, J = 9.1 Hz, 1H), 3.45 (dd, J = 12.6, 3.3 Hz, 2H), 3.19 (dd, J = 12.5, 6.5 Hz, 2H), 3.00 (td, J = 6.4, 3.4 Hz, 2H), 2.91-2.62 (m, 4H), 2.60-2.47 (m, 2H), 1.98 (dd, J = 10.9, 5.6 Hz, 1H), 1.19 (s, 6H), 1.10 (s, 6H), 0.98 (d, J =	22

FIG. 3 Continued.



				6.2 Hz, 6H).	
210		A	792.4	^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.08 (s, 1H), 7.92-7.89 (d, *J* = 9 Hz, 1H), 7.75-7.67 (m, 3H), 7.45-7.42 (d, *J* = 9.0 Hz, 1H), 7.35 (s, 1H), 7.28-7.25 (d, *J* = 9Hz, 1H), 7.21 (s, 1H), 7.02-6.98 (m, 1H), 6.45-6.42 (d, *J* = 9Hz,2H),5.10-5.05 (m, 1H) , 4.32 (s, 1H),4.07-3.99 (m, 3H), 3.59-3.55 (m, 2H), 3.47-3.45 (m,4H), 3.00-2.89 (m, 2H), 2.66-2.50(m, 8H), 2.04-2.00 (m, 1H), 1.21 (s, 6H),1.12 (s, 6H);	40
211		A	820.58		

FIG. 3 Continued.




212		A	836.55	^{1}H-NMR (300 MHz, DMSO-d_6) δ11.08（s, 1H), 8.61 (s, 1H), 7.90-7.84 (m, 1H), 7.74- 7.67 (m, 2H), 7.49-7.41 (m, 1H), 7.20- 7.16 (m, 1H), 7.02-6.89 (m, 3H), 5.17 - 5.07 (m, 1H), 4.29 (s, 1H), 4.06- 3.99 (m, 1H), 3.91- 3.68 (m, 6H), 2.92- 2.50 (m, 7H), 2.42- 2.34 (m, 4H), 2.06- 1.98 (m, 1H), 1.78- 1.60 (m, 2H), 1.54- 1.38 (m, 3H), 1.26 - 1.10 (m, 14H)	5
213		A	838.56		49
214		A	824.59		

FIG. 3 Continued.


215		A		^{1}H NMR (300 MHz, DMSO-d_6): δ 11.07 (s, 1H), 8.60 (s, 1H), 7.95-7.85 (m, 2H), 7.69 (d, J = 8.6 Hz, 1H), 7.57 (d, J = 9.2 Hz, 1H), 7.34 (s, 1H), 7.31(m, 1H) 7.17 (m, 1H), 7.00 (dd, J = 8.8, 2.5 Hz, 1H), 6.85 (d, J = 9.0 Hz, 1H), 5.07 (dd, J = 12.7, 5.6 Hz, 1H), 4.47 (s, 1H), 4.33 (d, J = 19.6 Hz, 2H), 4.05 (d, J = 9.1 Hz, 1H), 3.47 (s, 4H), 2.86 (d, J = 14.6 Hz, 2H), 2.55 (s, 5H), 2.24 (s, 1H),2.02 (m, 3H) 1.67 (m, 2H), 1.58 (m, 3H), 1.20 (m, 7H), 1.16 (m, 10H	94

FIG. 3 Continued.


216		A	821.4	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.12 (s, 1H), 7.91 (d, J = 8.8 Hz, 1H), 7.88-7.82 (m, 1H), 7.74 (d, J = 8.4 Hz, 2H), 7.43 (d, J = 9.2 Hz, 1H), 7.32-7.24 (m, 2H), 7.21 (d, J = 2.3 Hz, 1H), 7.01 (dd, J = 8.8, 2.4 Hz, 1H), 6.43 (d, J = 8.4 Hz, 2H), 5.13 (dd, J = 12.9, 5.3 Hz, 1H), 4.97 (td, J = 6.5, 3.2 Hz, 1H), 4.33 (s, 1H), 4.04-4.02 (m, 3H), 3.54 (t, J = 6.6 Hz, 2H), 3.12-3.10 (m, 1H), 2.93-2.90 (m, 13.9, 5.4 Hz, 1H), 2.79-2.77 (m, 1H), 2.71- 2.52 (m, 2H), 2.40 (dt, J = 12.7, 6.6 Hz, 2H), 2.22 (q, J = 7.4 Hz, 4H), 2.12 - 2.00 (m, 4H), 1.76 (q, J = 7.2 Hz, 2H), 1.22 (s, 6H), 1.13 (s, 6H);	51

FIG. 3 Continued.



217		A	799.52		
218		A	791.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.22 (6H, s), 1.73-1.78 (2H, m), 1.99-2.02 (1H, m), 2.42-2.49 (5H, m), 2.53-2.60 (6H, m), 2.83-2.89 (1H, m), 3.43 (5H, s), 4.06 (1H, d, J = 8.8 Hz), 4.32 (1H, s), 5.07 (1H, dd, J = 5.2, 12.8 Hz), 7.00 (1H, dd, J = 2.4, 8.8 Hz), 7.20-7.26 (2H, m), 7.31-7.33 (3H. m), 7.67 (1H, d, J = 8.8 Hz), 7.73-7.79 (3H, m), 7.90 (1H, d, J = 8.8 Hz), 11.0 (1H, s).	56
219		A		^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.23 (6H, s), 1.98-2.02 (1H, m), 2.15-2.49 (5H, m), 2.52-2.67 (8H, m), 2.83-2.92 (1H, m), 3.44 (4H, s), 3.64-3.68 (1H, m), 4.06 (1H, d, J = 9.2 Hz), 4.32 (1H, s), 5.07 (1H, dd, J = 5.6, 12.8 Hz), 7.00 (1H, dd, J = 2.4, 8.8 Hz), 7.21 (1H, d, J = 2.4 Hz), 7.25 (1H,	56

FIG. 3 Continued.


				d, *J* = 8.8 Hz), 7.34-7.39 (3H, m), 7.68 (1H, d, *J* = 8.4 Hz),7.74-7.80 (3H, m), 7.90 (1H, d, *J* = 8.4 Hz), 11.0 (1H, s).	
220		A		^{1}H NMR (300 MHz, DMSO-d_6): δ 11.06 (s, 1H), 8.57 (d, J = 2.4 Hz, 1H), 7.95-7.83 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H), 7.55 (d, J =9.2 Hz, 1H), 7.34-7.17 (m, 3H), 7.00 (dd, J = 8.8, 2.4 Hz, 1H), 6.83 (d, J = 9.1 Hz, 1H), 5.06 (dd, J = 12.6, 5.3 Hz, 1H), 4.53 (d, J = 13.1 Hz, 1H), 4.31 (d, J = 10.7 Hz, 2H), 4.04 (d, J = 9.2 Hz, 1H), 3.39 (m, 4H), 3.06 (d, J = 12.5 Hz, 1H), 2.86 (d, J = 11.6 Hz, 2H), 2.59 (m, 6H), 2.03 (s, 2H), 1.72 (t, J = 13.2 Hz, 4H), 1.60-1.32 (m, 5H), 1.21 (d, J = 2.0 Hz, 7H), 1.11 (s, 6H);	94

FIG. 3 Continued.



221		A		^{1}H NMR (300 MHz, DMSO-d_6): δ 11.07 (s, 1H), 8.61 (d, J = 2.4 Hz, 1H), 7.96-7.85 (m, 2H), 7.68 (d, J = 8.5 Hz, 1H), 7.56 (d, J = 9.1 Hz, 1H), 7.34-7.17 (m, 3H), 7.00 (dd, J = 8.7, 2.4 Hz, 1H), 6.85 (d, J = 9.0 Hz, 1H), 5.07 (dd, J = 12.7, 5.4 Hz, 1H), 4.30 (s, 1H), 4.20 (d, J = 13.0 Hz, 1H), 4.05 (d, J = 9.2 Hz, 2H), 3.43 (s, 4H), 3.08 (dd, J = 23.5, 12.0 Hz, 2H), 2.88 (t, J = 13.6 Hz, 1H), 2.68 (s, 5H), 2.38 (s, 2H), 2.07-1.96 (m, 1H), 1.87-1.49 (m, 9H), 1.41- 1.28 (m, 1H), 1.21 (s, 6H), 1.11 (s, 6H)	94
222		A	808.3	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.08 (s, 1H), 8.24 (d, J = 9.2 Hz, 1H), 7.90 (d, J = 8.8 Hz, 1H), 7.83 (d, J = 9.5 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.40 (d, J = 9.6 Hz, 1H), 7.33 (d, J = 2.1 Hz, 1H), 7.25 (q, J = 4.0, 3.0 Hz, 2H), 7.03 (dd, J = 8.8, 2.5 Hz, 1H), 5.06 (dd, J = 12.7, 5.3 Hz, 1H), 4.53 (d, J = 13.2 Hz, 2H), 4.46 (s, 1H), 4.00 (d, J = 9.1 Hz, 1H), 3.43 (brs, 4H), 3.05 (t, J = 12.5 Hz, 2H), 2.89 (s, 1H), 2.60 (d, J = 31.4 Hz, 7H), 2.01 (m, 1H), 1.91 (m, 2H), 1.46 (m, 2H), 1.22 (s, 6H), 1.14 (s, 6H).	52

FIG. 3 Continued.



223		A	792.3	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.06 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.69 (dd, J = 16.6, 8.5 Hz, 3H), 7.43 – 7.30 (m, 2H), 7.30 (m, 1H), 7.14 (m, 1H), 6.97 (dd, J = 8.8, 2.5 Hz, 1H), 6.56 (d, J = 8.7 Hz, 2H), 5.05 (dd, J = 12.7, 5.4 Hz, 1H), 4.29 (s, 1H), 4.02 (d, J = 9.3 Hz, 1H), 3.56 (t, J = 8.4 Hz, 1H), 3.39 (s, 5H), 3.13 (t, J = 8.7 Hz, 1H), 2.97 (d, J = 8.7 Hz, 2H), 2.83-2.56 (m, 5H), 2.52-2.04 (m, 2H), 1.88 (m, 3H), , 1.18 (d, J = 27.9 Hz, 6H), 1.04 (d, J = 27.9 Hz, 6H).	26
224		A	778.5	H-NMR-PH-ARV-LS-049-Q-0 (300 MHz, DMSO-d_6, ppm) :δ 11.08 (s, 1H), 7.91 (d, J = 8.7 Hz, 1H), 7.77 (d, J = 8.6 Hz, 2H), 7.67 (d, J = 8.3 Hz, 1H), 7.52 (d, J = 9.2 Hz, 1H), 7.21 (d, J = 2.4 Hz, 1H), 7.06- 6.95 (m, 3H), 6.82 (d, J = 2.0 Hz, 1H), 6.68 (dd, J = 8.4, 2.1 Hz, 1H), 5.07 (dd, J = 12.6, 5.4 Hz, 1H), 4.33 (s, 1H), 4.21-4.12(m, 2H), 4.11-4.01 (m, 1H), 3.93 (dd, J = 8.8, 4.8 Hz, 2H), 3.41 (p, J = 5.7 Hz, 1H), 3.29 (s, 3H), 2.94-2.80 (m, 1H), 2.64-2.51 (m, 4H), 2.50-2.40 (m, 3H), 2.11-1.97 (m, 1H), 1.22 (s, 6H), 1.13 (s, 6H)	19

FIG. 3 Continued.



225		A	820.3	H NMR (400 MHz, DMSO-*d6*) δ11.05 (s, 1H), 7.88 (d, J = 8.7 Hz, 1H), 7.71 (t, J = 8.4 Hz, 3H), 7.37 (d, J = 8.9 Hz, 2H), 7.31- 7.15 (m, 2H), 6.98 (dd, J = 8.7, 2.4 Hz, 1H), 6.56 (d, J = 8.6 Hz, 2H), 5.06 (m, 1H), 4.30 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.75 (m, 2H), 3.70-3.51 (m, 5H), 3.51- 3.32 (m, 5H), 2.85 (m, 1H), 2.57 (m, 2H), 2.37-1.81 (m, 4H), 1.15 (d, 12H)	18
226		A	838.3	^{1}H NMR (400 MHz, DMSO-d_6) δ11.11 (s, 1H), 7.92 (d, J = 8.7 Hz, 1H), 7.78 (m, 3H), 7.53 (d, J = 7.2 Hz, 1H), 7.41 (d, J = 9.2 Hz, 1H), 7.22 (s, 1H), 7.02 (d, J = 8.6 Hz, 1H), 6.60 (d, J = 8.4 Hz, 2H), 5.13 (dd, J = 12.9, 5.3 Hz, 1H), 4.33 (s, 1H), 4.07 (d, J = 9.2 Hz, 1H), 3.80 (s, 2H), 3.61 (m, 2H), 3.44 (m, 2H), 3.36 (d, J = 9.9 Hz, 4H), 3.25 (d, J = 6.1 Hz, 2H), 2.94-2.83 (m, 1H), 2.60 (t, J = 14.3 Hz, 3H), 2.30- 2.11 (m, 2H), 2.06 (m, 1H), 1.23 (s, 6H), 1.14 (s, 6H)	18

FIG. 3 Continued.



227		A	824.3	1H NMR (300 MHz, DMSO-*d6*): δ 11.07 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.70 (m, 3H), 7.38 (dd, J = 27.2, 8.3 Hz, 2H), 7.17 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 6.51 (d, J = 8.7 Hz, 2H), 5.07 (dd, J = 12.9, 5.3 Hz, 1H), 4.28 (s, 1H), 4.02 (d, J = 9.1 Hz, 1H), 3.49-3.30 (m, 2H), 3.24-3.17 (m,, 4H), 3.09-2.98 (m, 1H), 2.91-2.78 (m, 1H), 2.65-2.48 (m, 6H), 2.47-2.42 (m, 2H), 2.41-2.39 (m, 2H), 2.15-1.98 (m, 2H), 1.80-1.60(m, 1H), 1.18 (s, 6H), 1.09 (s, 6H).;	49
228		A	824.2	1H NMR (300 MHz, DMSO-*d6*) δ 11.04 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 (m, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.06 – 6.89 (m, 4H), 5.03 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.28 (s, 1H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.78-3.58 (m, 3H), 3.44 -3.21(m, 4H), 2.94–2.75 (m, 1H), 2.64–2.55 (m, 5H), 2.42 (m, 3H), 2.37 (d, *J* = 7.4 Hz, 2H), 2.12 – 1.92 (m, 2H), 1.68 (m, 1H), 1.18 (s, 6H), 1.09 (s, 6H)	53

FIG. 3 Continued.




229		A	779.3	^{1}H NMR (300 MHz, d6-DMSO) δ 1.13 (s, 6H), 1.22 (s, 6H), 2.07 (s, 1H), 2.52-2.64 (m, 2H), 2.82-2.90 (m, 3H), 3.37-3.44 (m, 4H), 3.96 (s, 4H), 4.04-4.07 (m, 1H), 4.18 (s, 2H), 4.32 (s, 1H), 5.16 (s, 1H), 6.42-6.45 (m, 2H), 7.00-7.03 (m, 1H), 7.21 (s, 1H), 7.35-7.44 (m, 3H), 7.72-7.74 (m, 2H), 7.84-7.93 (m, 2H), 11.13 (s, 1H).	24
230		A	848.4	1H NMR (400 MHz, DMSO-*d6*): δ 11.06 (s, 1H), 7.90 (m, 1H), 7.75 (m, 2H), 7.65 (m, 1H), 7.55 (m, 1H), 7.20 (m, 1H), 7.02 (m, 3H), 6.97 (m, 1H), 6.85 (m, 1H), 5.04 (m, 1H), 4.50 (m, 1H), 4.30 (m, 1H), 4.12 (m, 1H), 4.05 (m, 1H), 3.95 (m, 2H), 3.70 (m, 1H), 3.50 (m, 3H), 3.30 (m, 1H), 3.00 -2.88 (m, 4H), 2.62-2.52 (m, 4H), 2.41-2.02 (m, 3H), 2.01-1.4 (m, 3H), 1.30-1.10 (d, 12H	6
231		A	854.45	1H NMR (300 MHz, $CDCl_3$): δ 8.12 (d, J = 9.1 Hz, 1H), 8.03 (s, 1H), 7.94 (d, J = 9.5 Hz, 1H), 7.54 (d, J = 8.7 Hz, 1H), 7.45 (d, J = 11.0 Hz, 1H), 7.37 (d, J = 7.3 Hz, 1H), 7.01 – 6.91 (m, 2H), 6.78 (dd, J = 8.7, 2.4 Hz, 1H), 4.91 (dd, J = 12.2, 5.3 Hz, 1H), 4.50 (d, J = 13.3 Hz, 2H), 4.17 (d, J = 9.1 Hz, 1H), 4.04 (s, 1H), 3.28 (s, 4H), 3.03 (t, J = 12.3 Hz, 2H), 2.94 – 2.66 (m, 3H), 2.62 (s, 4H), 2.46 (s, 2H), 2.17 – 2.06 (m, 1H), 1.98 (s, 0H), 1.86 (d, J = 12.9 Hz, 2H), 1.69 (s, 1H), 1.26 (s, 9H), 1.19 (s, 6H).	29

FIG. 3 Continued.




232		A	850.5	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.11 (s, 6H), 1.17-1.32 (m, 16H), 1.77-1.82 (m, 1H), 1.95-2.07 (m, 2H), 2.17-2.33 (m, 4H), 2.54-2.61 (m, 4H), 2.83-2.94 (m, 1H), 3.47 (s, 4H), 4.04 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1H), 4.95-5.01 (m, 1H), 5.04-5.11 (m, 1H), 6.98-7.04 (m, 1H), 7.20-7.37 (m, 3H), 7.66-7.73 (m, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.74-8.85 (m, 2H), 11.08 (s, 1H).	95
233		A	850.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.22-1.32 (m, 14H), 1.80 (d, *J* = 12.4 Hz, 2H), 1.99-2.03 (m, 1H), 2.24-2.33 (m, 3H), 2.55-2.61 (m, 6H), 2.84-2.93 (m, 1H), 3.47 (m, 4H), 4.04 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1 H), 4.96-5.01(m, 2H), 5.07 (dd, *J* = 5.2,12.8Hz, 1H), 5.07 (dd, *J* = 2.4,8.8Hz, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.28 (d, *J* = 8.8 Hz, 1H), 7.36 (s, 1H), 7.68-7.74 (m, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.79 (s, 2H), 11.08 (s, 1H).	95
234		A	850.6	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (d, *J* = 7.2 Hz, 6H), 1.26-1.30 (m, 6H), 1.68 (d, *J* = 13.0 Hz, 1H), 1.78-1.88 (m, 2H), 2.00-2.04 (m, 1H), 2.16-2.38 (m, 5H), 2.54-2.68 (m, 5H), 2.84-2.94 (m, 1H), 3.47 (s, 4H), 4.04 (d, *J* = 8.8 Hz, 1H), 4.29 (s, 1H), 4.48-4.56 (m, 2H), 5.08(d, *J* = 12.8 Hz,1H), 7.01 (d, *J* = 8.8 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.27 (d, *J* = 8.4 Hz, 1H), 7.36 (s, 1H), 7.68-7.71 (m, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.78 (s, 2H), 11.09 (s, 1H).	95

FIG. 3 Continued.



235		A	807.54		
236		A	875.65	H-NMR-PH-ARV-LS-035-H-0: (300 MHz, CD_3OD, ppm) δ 8.60-8.59 (d, *J*=2.4Hz, 1H), 7.97-7.94 (m, 1H), 7.87-7.70 (m, 4H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 1H), 6.86-6.83 (m, 1H), 5.17-5.11 (m, 1H), 4.40-4.35 (m, 2H), 4.28 (s, 1H), 4.14 (s, 1H), 3.07-2.77 (m, 10H), 2.28-2.13 (m, 6H), 2.00-1.78 (m, 8H), 1.61-1.58 (m, 1H), 1.42-1.33 (m, 1H), 1.28 (s, 6H), 1.21 (s, 6H), 1.13-1.00 (m, 1H).	96

FIG. 3 Continued.



237		A	835.4	^{1}H NMR (300 MHz, CD_3OD): δ 7.77-7.70 (m, 4H), 7.25-7.19 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 7.01-6.95 (m, 3H), 5.12-5.06 (m, 1H), 4.96-4.90 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 4.01-3.97 (m, 2H), 2.88-2.68 (m, 9H), 2.36-2.31 (m, 7H), 2.10-2.00 (m, 1H), 1.95-1.85 (m, 2H), 1.67-1.60 (m, 2H), 1.24 (s, 6H), 1.18 (s, 6H);	23
238		A	820.35	^{1}H NMR (300 MHz, $CDCl_3$) 7.92 (s, 1H), 7.63-7.47 (m, 4H), 7.22-7.19 (m, 1H), 7.01-6.98 (m, 1H), 6.97-6.89 (m, 1H), 6.88-6.71 (m, 1H), 6.53-6.50 (m, 2H), 6.00 (m, 1H), 4.88-4.84 (m, 1H), 4.08-4.05 (m, 1H), 3.96-3.92 (m, 3H), 3.30 (s, 1H), 2.97-2.80 (m, 5H) , 2.65-2.64 (m, 2H), 2.77-2.64 (m, 1H), 2.33 (s, 1H), 2.06-2.03 (m, 3H), 1.93 (m,2H), 1.51-1.35 (m, 5H), 1.18 (m, 6H), 1.14(m, 6H), 0.81-0.76 (m, 1H);	36

FIG. 3 Continued.



239		A	835.1	^{1}H-NMR (400 MHz, DMSO-*d6*) δ 11.05 (s, 1H), 7.88-7.85 (d, *J*=9.0 Hz, 1H), 7.73-7.63 (m, 3H), 7.50-7.47 (d, *J*=9.0 Hz, 1H), 7.30-7.17 (m, 3H), 6.99-6.92 (m, 3H), 5.06-5.00 (m, 1H), 4.28 (s, 1H), 4.03-4.00 (m, 1H), 3.40-3.22 (m, 8H), 2.86-2.79 (m, 1H), 2.53-2.46 (m, 14H), 2.03-1.98 (m, 1H), 1.18-1.08 (m, 12H);	38
240		A	863.4	H-NMR (300 MHz, DMSO-d*6*) δ 11.09 (s, 1H), 7.90-7.87 (d, *J* = 9Hz, 1H), 7.82-7.80 (d, *J* = 6 Hz, 1H), 7.72-7.69 (d, *J* = 9 Hz, 2H), 7.47-7.44 (d, *J* = 9 Hz, 1H), 7.26-7.18 (m, 3H), 7.00-6.92 (m, 3H), 5.12-5.06 (m, 1H), 4.88 (s, 1H), 4.30 (s, 1H), 4.04-4.01 (d, *J* = 9 Hz, 1H), 3.88-3.84 (d, *J* = 12 Hz, 2H), 3.67-3.30 (m, 1H), 2.91-2.70 (m, 4H), 2.60-2.37 (m, 4H), 2.24-2.13 (m, 4H), 2.05-2.00 (m, 1H), 1.82-1.78 (d, *J* = 12 Hz, 2H), 1.54 (s, 1H), 1.19 (s, 6H), 1.10 (s, 8H), 0.91-0.89 (d, *J*=6 Hz, 6H)	23

FIG. 3 Continued.


241		A	834.5	^{1}H NMR (400 MHz, DMSO): δ 11.08 (s, 1H), 8.77-8.68 (m, 2H), 7.91 (d, J = 8.8 Hz, 1H), 7.68 (d, J = 8.7 Hz, 2H), 7.33 (d, J = 2.3 Hz, 1H), 7.29-7.19 (m, 2H), 7.02-6.99 (m, 1H), 5.09-5.05 (m, 1H), 4.64 (s, 1H), 4.29 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.55 (m, 1H), 3.44-3.42 (m, 4H), 3.35-3.32 (m, 1H), 2.92-2.81 (m, 1H), 2.68-2.53 (m, 5H), 2.48-2.36 (m, 3H), 2.35-2.18 (m, 2H), 2.08-1.91 (m, 2H), 1.81 (m, 1H), 1.69 (m, 1H), 1.22 (s, 6H), 1.12 (s, 6H), 1.06 (m, 1H)	97

FIG. 3 Continued.



242		A	834.1	^{1}H NMR (400 MHz, DMSO): δ 11.08 (s, 1H), 8.73 (d, J = 2.7 Hz, 2H), 7.91 (d, J = 8.8 Hz, 1H), 7.67 (m, 2H), 7.34 (d, J = 2.1 Hz, 1H), 7.29-7.24 (m, 1H), 7.22 (d, J = 2.4 Hz, 1H), 7.01 (m, 1H), 5.07 (m, 1H), 4.68 (s, 1H), 4.30 (s, 1H), 4.04 (d, J = 9.2 Hz, 1H), 3.51-3.39 (m, 5H), 3.34 (m, 1H), 3.26 (m, 1H), 2.96-2.82 (m, 1H), 2.56 (m, 5H), 2.44-2.34 (m, 1H), 2.17 (m, 1H), 2.04-1.96 (m, 2H), 1.80-1.54 (m, 3H), 1.37 (d, J = 11.8 Hz, 1H), 1.22 (m, 7H), 1.12 (s, 6H)	97
243		A	840.35	H-NMR (400 MHz, DMSO-*d6*) δ11.12 (s, 1H), 8.76 (s, 2H), 7.92-7.90 (d, *J* = 8.8Hz, 2H), 7.76-7.70 (m, 2H), 7.48-7.46 (d, *J*=7.2Hz, 1H), 7.23-7.22 (d, *J*=2.4Hz, 1H), 7.03-7.01 (m, 1H), 5.14-5.1 (m, 1H), 4.78-4.74 (d, *J*=12.8Hz, 2H), 4.30 (s, 1H), 4.06-4.04 (d, *J*=9.2Hz, 1H), 3.27 (s, 3H), 3.03-2.97 (m, 3H), 2.63-2.52 (m, 6H), 2.24-2.23 (m, 2H), 2.05-2.04 (m, 1H), 1.86-1.83 (m, 3H), 1.23 (s, 6H), 1.12 (s, 6H), 1.09 -1.06 (m, 2H)	49

FIG. 3 Continued.



244		A	781.3	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.12 (s, 1H), 7.91 (d, J = 8.8 Hz, 1H), 7.84 (d, J = 8.3 Hz, 1H), 7.75 (d, J = 8.6 Hz, 2H), 7.48 (d, J = 2.3 Hz, 1H), 7.45-7.32 (m, 2H), 7.21 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.5 Hz, 1H), 6.57 (d, J = 8.8 Hz, 2H), 5.13 (dd, J = 12.8, 5.2 Hz, 1H), 4.33 (s, 3H), 4.06 (d, J = 9.1 Hz, 1H), 3.55 (t, J = 8.3 Hz, 1H), 3.45 (t, J = 8.7 Hz, 1H), 3.29 (d, J = 8.9 Hz, 2H), 3.20-3.07 (m, 1H), 2.88 (d, J = 12.3 Hz, 3H), 2.60 (d, J = 18.2 Hz, 2H), 2.36 (s, 3H), 2.22 (s, 1H), 2.06 (d, J = 10.6 Hz, 1H), 1.96-1.83 (m, 1H), 1.22 (s, 6H), 1.13 (s, 6H).	93
245		A	779.15	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 7.84 (dd, J = 18.3, 8.7 Hz, 2H), 7.71 (d, J = 8.2 Hz, 2H), 7.35 -7.26 (d, J = 6.5 Hz, 4H), 7.17 (s, 1H), 6.49 (d, J = 8.5 Hz, 2H), 5.15 – 5.02 (m, 2H), 4.29 (s, 1H), 4.02 (d, J = 9.2 Hz, 1H), 3.76 (d, J = 6.9 Hz, 3H), 3.09 (dd, J = 25.1, 8.6 Hz, 4H), 2.98 (s, 2H), 2.54 (s, 2H), 1.97(s,4H),1.18 (s, 6H), 1.09 (s, 6H).	23

FIG. 3 Continued.



246		A	793.25	1H NMR (300 MHz, CDCL3) δ 11.07 (s, 1H), 7.76 (t, J = 8.0 Hz, 3H), 7.58 (d, J = 8.6 Hz, 1H), 7.33 (d, J = 18.6 Hz, 2H), 7.19 (d, J = 8.2 Hz, 1H), 6.99 (d, J = 2.3 Hz, 1H), 6.84 (dd, J = 8.8, 2.4 Hz, 1H), 6.53 (d, J = 8.4 Hz, 2H), 4.22 (d, J = 8.3 Hz, 3H), 4.10 (s, 1H), 3.45 (s, 4H), 3.19 (s, 2H), 3.08 (s, 4H), 2.67 (d, J = 17.7 Hz, 4H), 2.57 (d, J = 13.7 Hz, 1H), 2.28 (s, 1H), 1.77 (s, 1H), 1.37 (s, 3H), 1.28 (d, J = 4.9 Hz, 9H).;	55
247		A	802.5	^{1}H NMR (400 MHz, d-DMSO):δ11.086(s,1H),8.754(s,2H),7.708-7.674(m,3H),7.348(s,1H),7.275-7.253(d,J=8.8Hz,1H),6.938-6.933(d,J=2Hz,1H),6.830-6.803(m,1H),5.098-5.052(m,1H),4.764-4.732(d,J=12.8Hz,2H),4.193(s,1H),4.036-4.013(d,J=9.2Hz,1H),3.450(s,4H),3.020-2.961(m,2H),2.932-2.840(m,1H),2.608-2.537(m,6H),2.507-2.498(m,3H),2.216-2.199(d,J=6.8Hz,2H),2.076-2.018(m,1H),2.005-1.914(m,1H),1.851-1.819(d,J=12.8Hz,2H),1.215(s,6H),1.110(s,6H),1.073-1.018(m,2H)	49

FIG. 3 Continued.



248		B	765.7	^{1}H NMR (400 MHz, d-DMSO): δ 10.69 (s,1H), 7.91-7.89 (d, *J*=8.8Hz, 1H), 7.78-7.74 (m, 3H), 7.53-7.51 (d, *J*=8.4Hz, 1H), 7.32-7.30 (d, *J*=8.0Hz, 2H), 7.21-7.20 (d, *J*=2.4Hz, 1H), 7.05-6.98 (m, 3H), 5.30 (s, 2H), 4.58-4.54 (m, 1H), 4.32 (s,1H), 4.07-4.05 (d, *J*=9.2Hz, 1H), 3.29-3.26 (m, 4H), 2.69-2.58 (m, 4H), 2.50-2.49 (m, 4H), 2.33-2.32 (s, 2H), 2.07-1.95 (m, 2H), 1.64-1.50 (m, 4H), 1.20 (s, 6H), 1.10 (s, 6H);	21A
249		B	765.7	^{1}H NMR (400 MHz, d-DMSO): δ10.69 (s, 1H), 7.91-7.89 (d, *J*=8.8Hz, 1H), 7.78-7.74 (m, 3H), 7.53-7.51 (d, *J*=8.4Hz, 1H), 7.32-7.30 (d, *J*=8.0Hz, 2H), 7.21-7.20 (d, *J*=2.4Hz, 1H), 7.05-6.98 (m, 3H), 5.29 (s, 2H), 4.58-4.54 (m, 1H), 4.32 (s,1H), 4.07-4.05 (d, *J*=9.2Hz, 1H), 3.29-3.27 (m, 4H), 2.69-2.58 (m, 4H), 2.50-2.49 (m, 4H), 2.34-2.32 (s, 2H), 2.01-1.92 (m, 2H), 1.66-1.48 (m, 4H), 1.23 (s, 6H), 1.18 (s, 6H);	21A

FIG. 3 Continued.



250		B	793.7	^{1}H NMR (400 MHz, d-DMSO): δ7.92-7.89 (d, *J*=8.4Hz, 1H), 7.78-7.66 (m, 4H), 7.32-7.30 (d, *J*=8.0Hz, 2H), 7.21-7.10 (m, 3H), 7.01-7.00 (d, J=6.8Hz, 1H), 5.39 (s, 1H), 4.62 (s, 1H), 4.33 (s,1H), 4.07-3.98 (m, 3H), 3.56-3.40 (m, 2H), 3.34-3.08 (m, 6H), 2.67-2.56 (m, 4H), 2.10-1.95 (m, 2H), 1.68-1.55 (m, 4H), 1.33 (m, 5H), 1.23 (s, 6H), 1.13 (s, 6H);	21A
251		B	793.7	^{1}H NMR (400 MHz, *d*-DMSO): δ 10.70 (s, 1H), 7.91-7.89 (d, *J*=8.8Hz, 1H), 7.77-7.74 (m, 3H), 7.55-7.53 (d, *J*=8.4Hz, 1H), 7.31-7.29 (d, *J* = 8.4Hz, 2H), 7.22-7.10 (d, *J*=2.0Hz, 1H), 7.08-6.99 (m, 3H), 5.31 (s, 2H), 4.59-4.55 (s, 1H), 4.32 (s,1H), 4.07-4.05 (m, 1H), 3.34-3.33 (m, 8H), 2.66-2.50 (m, 6H), 2.10-1.90 (m, 2H), 1.60-1.38 (m, 4H), 1.32 (s, 4H), 1.22 (s, 6H), 1.13 (s, 6H);	21A

FIG. 3 Continued.



252		B	843.4	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.11 - 11.04 (m, 1 H), 8.07 (d, J = 8.7 Hz, 1 H), 7.96 (dd, J = 24.7, 8.4 Hz, 1 H), 7.78 (dd, J = 24.7, 8.4 Hz, 2 H), 7.50 (d, J = 9.2 Hz, 1 H), 7.44- 7.21 (m, 4 H), 6.93 (d, J = 8.7 Hz, 2 H), 5.08 (dd, J = 12.8, 5.3 Hz, 1 H), 4.36 (s, 1 H), 4.16 (t, J = 6.3 Hz, 2 H), 4.04 (d, J = 9.3 Hz, 1 H), 3.21 (t, J = 4.8 Hz, 3 H), 2.83 (dd, J = 13.5, 5.3 Hz, 1 H), 2.56 (m, 7 H), 2.31 (t, J = 6.7 Hz, 2 H), 2.00 (s, 1H), 1.96 (t, J = 7.0 Hz, 2 H), 1.47 (s, 4 H), 1.20 (s, 6 H), 1.11 (s, 6 H);	9
253		B	868.4	^{1}H NMR (300 MHz, CD3OD) 7.93-7.90 (m, 1H), 7.77-7.70 (m, 2H), 7.68-7.65 (m, 1H), 7.36-7.29 (m, 4H), 7.29-7.19 (m, 2H), 5.09-5.03 (m, 1H), 4.34-4.31 (m, 1H), 4.17-4.10 (m, 1H), 3.87-3.80 (m, 2H), 2.84-2.67 (m, 10H), 2.11-2.08 (m, 1H), 1.67-1.50 (m, 2H) , 1.50-1.08 (m, 25H);	98

FIG. 3 Continued.



254		B	804.3	^{1}H NMR (300 MHz, CD_3OD): δ 7.84-7.64 (m, 6H), 7.60-7.35 (m, 2H), 7.13-7.12 (s, 1H), 7.00-6.96 (m, 1H), 6.30 (s, 1H), 5.18-5.12 (m, 1H), 4.70-4.43 (m, 6H), 4.29 (s, 1H), 4.16 (s, 1H), 3.31-3.07 (m, 6H), 2.96-2.48 (m, 8H), 2.21-2.17 (m, 1H), 1.97-1.92 (m, 1H), 1.28-1.15 (s, 12H);	99
255		B	801.4	1H NMR (300 MHz, Methanol-d4) δ 7.94 – 7.74 (m, 4H), 7.66 – 7.14 (m, 4H), 7.12 – 7.03 (m, 2H), 5.10 (dd, J = 12.3, 5.4 Hz, 1H), 4.58 (t, J = 4.2 Hz, 2H), 4.32 (s, 1H), 4.14 (t, J = 4.3 Hz, 1H), 3.72 (t, J = 4.6 Hz, 2H), 3.55 (d, J = 8.1 Hz, 7H), 2.75 (dddd, J = 22.4, 17.9, 8.2, 4.6 Hz, 3H), 2.11 (dtd, J = 12.7, 4.8, 2.3 Hz, 1H), 1.94 – 1.33 (m, 2H), 1.24 (d, J = 17.7 Hz, 12H).	9

FIG. 3 Continued.



256		B	805.4	^{1}H NMR (300 MHz, $CDCl_3$) 7.80 (s, 1H), 7.70-7.67 (m, 3H), 7.59-7.56 (m, 1H), 7.27 (s, 1H), 7.01 (s, 1H), 6.83-6.80 (m, 1H), 6.61-6.58 (m, 1H), 6.41 (s, 1H), 6.20-6.18 (m, 1H), 5.32-5.27 (m, 2H), 4.76-4.73 (m, 1H), 4.31 (s, 1H), 4.18-4.15 (m, 1H), 4.06 (s, 1H), 3.93-3.75 (m, 1H), 3.64-3.46 (m, 2H), 2.96-2.88 (m, 1H), 2.59-2.53 (m, 2H), 2.22-2.15 (m, 4H), 1.78-1.35 (m, 6H),1.32-1.23 (m, 18H), 0.88-0.83 (m, 1H);	21A
257		B	767.51	^{1}H NMR (300 MHz, Methanol-d4) δ 7.84-7.64 (m, 4H), 7.44 (d, J = 2.2 Hz, 1H), 7.34 (dd, J = 8.3, 2.3 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 7.03-6.90 (m, 3H), 5.09-5.07 (m, 1H), 4.39- 4.22 (m, 3H), 4.14-4.12 (m, 1H), 3.38-3.32 (m, 4H), 2.89-2.87 (m, 2H), 2.86-2.63 (m, 7H), 2.17 - 2.02 (m, 1H), 1.22 (d, J = 17.9 Hz, 12H) ;	9

FIG. 3 Continued.



258		B	805.4	^{1}H NMR (300 MHz, Methanol-d4) δ 7.84-7.64 (m, 4H), 7.44 (d, J = 2.2 Hz, 1H), 7.34 (dd, J = 8.3, 2.3 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 7.03-6.90 (m, 3H), 5.09-5.07 (m, 1H), 4.39- 4.22 (m, 3H), 4.14-4.12 (m, 1H), 3.38-3.32 (m, 4H), 2.89-2.87 (m, 2H), 2.86-2.63 (m, 7H), 2.17 - 2.02 (m, 1H), 1.22 (d, J = 17.9 Hz, 12H) ;	21A
259		B	805.4	^{1}H NMR (400 MHz, $CDCl_3$) 7.85 (s, 1H), 7.62-7.60 (m, 3H), 7.50-7.48 (m, 1H), 7.19-7.16 (m, 1H), 6.89-6.85 (m, 1H), 6.75-6.72 (m, 1H), 6.52-6.50 (m, 1H), 6.32-6.30 (m, 1H), 6.13-6.11 (m, 1H), 5.22-5.13 (m, 2H), 4.67-4.64 (m, 1H), 4.21-4.15 (m, 1H), 4.10-4.07 (m, 1H), 4.07-3.98 (m, 1H), 3.57-3.50 (m, 1H), 3.32-3.21 (m, 2H), 2.97-2.84 (m, 2H), 2.58-2.40 (m, 6H), 2.12-2.08 (m, 2H), 1.96-1.83 (m, 2H) , 1.54-1.40 (m, 2H), 1.40-1.04 (m, 18H);	21A

FIG. 3 Continued.




260		B	807.4	^{1}H NMR (300 MHz, $CDCl_3$): δ 7.85-7.71 (m, 4H), 7.58-7.56 (m, 1H), 7.26-7.22 (m, 2H), 6.99-6.97 (m, 2H), 6.83-6.79 (m, 2H), 6.25-6.22 (m, 1H), 5.31-5.30 (m, 2H), 4.76-4.65 (m, 3H), 4.45-4.43 (m, 2H), 4.18-4.06 (m, 2H), 3.49-3.30 (s, 4H), 3.07-2.59 (m, 9H), 2.25-2.20 (m, 2H), 2.01-1.94 (m, 2H), 1.28-1.23 (d, *J*=15.0Hz, 13H);	100
261		B	767.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.30 (s, 6H), 1.96-2.01 (m, 4H), 2.11-2.17 (m, 1H), 2.44-2.50 (m, 2H), 2.71-2.78 (m, 2H), 2.83-2.92 (m, 2H), 3.03 (t, *J* = 5.2 Hz, 2H), 3.27 (s, 2H), 4.18 (s, 1H), 4.29 (s, 1H), 4.38 (t, *J* = 5.2 Hz, 2H), 5.10-5.13 (m, 1H), 6.97-6.99 (m, 1H), 7.13-7.14 (m, 1H), 7.37-7.39 (m, 1H), 7.45-7.48 (m, 2H), 7.72-7.74 (m, 1H), 7.82-7.85 (m, 1H), 8.14-8.16 (m, 1H), 8.88 (d, *J* = 2.0 Hz, 1H).	9, 101
262		B	795.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.19 (s, 6H), 1.70-2.05 (m, 10H), 2.35-2.48 (m, 2H), 2.58-2.90 (m, 7H), 4.08 (s, 1H), 4.12 (t, *J* = 5.6 Hz, 2H), 4.19 (s, 1H), 4.98-5.04 (m, 1H), 6.87-6.91 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.22-7.26 (m, 1H), 7.32 (d, J = 2.0 Hz, 1H), 7.35 (d, *J* = 8.4 Hz, 1H), 7.63(d, *J* = 8.8 Hz, 1H), 7.72 (d, *J* = 8.4 Hz, 1H), 8.04-8.08 (m, 1H), 8.79 (d, *J* = 2.0 Hz, 1H).	9, 101

FIG. 3 Continued.



263		B	795.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.19 (s, 6H), 1.70-2.05 (m, 10H), 2.35-2.48 (m, 2H), 2.58-2.90 (m, 7H), 4.08 (s, 1H), 4.12 (t, *J* = 5.6 Hz, 2H), 4.19 (s, 1H), 4.98-5.04 (m, 1H), 6.87-6.91 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.22-7.26 (m, 1H), 7.32 (d, J = 2.0 Hz, 1H), 7.35 (d, *J* = 8.4 Hz, 1H), 7.63(d, *J* = 8.8 Hz, 1H), 7.72 (d, *J* = 8.4 Hz, 1H), 8.04-8.08 (m, 1H), 8.79 (d, *J* = 2.0 Hz, 1H).	9
264		B	811.3	^{1}H-NMR (300 MHz,d-DMSO) δ11.12(s, 1H), 8.65 (s, 1H), 8.38(m, 1H), 7.92- 7.81(m, 3H), 7.50-7.33(m, 2H), 7.25-7.24(m, 1H), 7.14- 7.01(m, 1H), 5.17- 5.07(m, 1H), 4.44(s, 1H), 4.24- 4.17(m, 2H), 3.99- 3.36(m, 5H), 2.91-2.83(m, 2H), 2.64 - 2.54(m, 3H), 2.45- 2.41(m, 2H), 2.28(s, 1H), 2.09- 2.02(m, 1H), 1.85- 1.47(m, 7H), 1.19 - 1.13 (d, *J* = *18Hz* , 12H).	9

FIG. 3 Continued.



265		B	811.7	H-NMR (300 MHz, CD_3OD) δ 7.94 (s, 1H), 7.79-7.68 (m, 2H), 7.38-7.27 (m, 3H), 7.13-7.12 (d, *J*=2.4 Hz, 1H), 6.99-6.96 (m, 1H), 5.11-5.05 (m, 1H), 4.31 (s, 1H), 4.19-4.15 (m, 2H), 4.08 (s, 1H), 3.91 (s, 1H), 3.02-2.99 (m, 4H), 2.85-2.71 (m, 5H), 1.92-1.87 (m, 2H),1.76-1.74 (d, *J*=4.8 Hz, 2H) , 1.60-1.58 (d, *J*=7.2 Hz, 2H), 1.28 (s, 6H), 1.20 (s, 6H);	9
266		B	810.29	H-NMR (400MHz,CD3OD, *ppm*): δ 8.37 (d, J = 2.8 Hz, 1H), 7.97 (d, J = 8.8 Hz, 1H), 7.83 (d, J = 8.3 Hz, 1H), 7.74 (d, J = 8.7 Hz, 1H), 7.51 – 7.40 (m, 2H), 7.34 (dd, J = 8.4, 2.3 Hz, 1H), 7.17 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.4 Hz, 1H), 5.13 (dd, J = 12.6, 5.5 Hz, 1H), 4.34 (s, 1H), 4.22 (t, J = 6.2 Hz, 2H), 4.09 (s, 1H), 3.54 (t, J = 4.9 Hz, 4H), 2.98 (t, J = 4.8 Hz, 5H), 2.95 – 2.67 (m, 4), 2.15 (m, 1H), 2.00 – 1.88 (m, 2H), 1.84 – 1.71 (m, 2H), 1.63 (q, J = 7.9 Hz, 2H), 1.31 (s, 6H), 1.23 (s, 6H).	9

FIG. 3 Continued.



267		B	798.4	^{1}H NMR (300 MHz, CD_3OD): δ 11.11 (s, 1H), 8.36 (s, 1H), 8.16-7.81 (m, 3H), 7.44 (m, 1H), 7.37-7.22 (m, 2H), 7.21 (m, 1H), 7.03-6.99 (m, 1H), 5.15-5.09 (m, 5.3 Hz, 1H), 4.29 (s, 1H), 4.22-4.10 (m, 3H), 4.05 (d, J = 9.3 Hz, 1H), 3.00-2.80 (m, 3H), 2.64-2.52 (m, 2H), 2.40-2.25 (m, 2H), 2.11-1.70 (m, 9H), 1.70-1.39 (m, 4H), 1.20 (s, 6H), 1.09 (s, 6H);	9, 102
268		B	798.3	H-NMR (300 MHz, $CDCl_3$) δ 8.43 (s, 1H), 7.794-7.77 (d, *J*=8.4 Hz, 2H), 7.59-7.52 (m, 2H)7.33-7.32 (d, *J*=2.1 Hz, 1H), 7.21-7.17 (m, 1H), 6.97-6.96 (d, *J*=2.4 Hz, 2H), 6.82-6.78 (m, 1H), 6.54-6.53 (d, *J*=2.1 Hz, 1H), 6.11-6.08 (d, *J*=8.4 Hz, 1H), 5.34 (s, 1H), 4.99-4.93 (m, 1H), 4.12-4.05 (m, 4H), 3.49-3.44 (m, 2H), 2.94-2.72 (m, 7H), 2.48-2.45 (d, *J*=9.9 Hz, 2H), 2.32 (s, 2H), 2.18-2.13 (m, 1H), 1.91-1.85 (m, 4H), 1.60-1.53 (m, 2H), 1.26-1.21 (d, *J*=10.5 Hz, 12H);	9, 103

FIG. 3 Continued.



269		B	770.7	H-NMR-PH-ARV-LS-029-E-0(400MHz,CD3OD, *ppm*): δ 8.30 (d, J = 0.8 Hz, 1H), 8.00 (s, 1H), 7.76 (dd, J = 24.6, 8.5 Hz, 2H), 7.40 (d, J = 2.2 Hz, 1H), 7.32 (dd, J = 8.3, 2.3 Hz, 1H), 7.12 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 5.10 (dt, J = 9.7, 5.2 Hz, 2H), 4.29-4.10 (m, 4H), 3.24 (d, J = 5.7 Hz, 3H), 2.97-2.55 (m, 6H), 2.30 (m, 1H), 2.27 (s, 1H), 2.20-2.05 (m, 1H), 1.88 (dq, J = 27.1, 7.5 Hz, 4H), 1.23 (d, J = 21.8 Hz, 12H).	9, 104
270		B	751.51	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.04 (s, 1H), 7.87 (d, *J* = 8.8 Hz, 1H), 7.79-7.60 (m, 3H), 7.51(m, 1H), 7.38-7.35 (m, 3H), 7.34-7.32 (m, 1H), 7.14 (m, 1H), 6.97 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.04 (dd, *J* = 12.7, 5.4 Hz, 1H), 4.29 (s, 1H), 4.03 (d, *J* = 9.1 Hz, 1H), 3.59 (s, 4H), 3.29 (s, 1H), 2.88-2.76 (m, 3H), 2.58-2.52 (m, 7H), 2.07-1.92 (m, 1H), 1.19 (d, *J* = 28.2 Hz, 12H);	1

FIG. 3 Continued.



271		B	811.4	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.04 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.79-7.60 (m, 3H), 7.51(m, 1H), 7.38-7.35 (m, 3H), 7.34-7.32 (m, 1H), 7.14 (m, 1H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 5.04 (dd, J = 12.7, 5.4 Hz, 1H), 4.29 (s, 1H), 4.03 (d, J = 9.1 Hz, 1H), 3.59 (s, 4H), 3.29 (s, 1H), 2.88-2.76 (m, 3H), 2.58-2.52 (m, 7H), 2.07-1.92 (m, 1H), 1.19 (d, J = 28.2 Hz, 12H);	20
272		B	812.4	^{1}H NMR (300 MHz, Chloroform-d) δ 8.92 (d, J = 2.3 Hz, 1H), 8.22 (s, 1H), 8.04 (dd, J = 8.1, 2.4 Hz, 1H), 7.62 – 7.45 (m, 2H),7.02 – 6.93 (m, 2H), 6.81 (dd, J = 8.8, 2.4 Hz, 1H), 6.24 (d, J = 8.2 Hz, 1H), 5.18 (dd, J = 13.1, 5.2 Hz, 1H), 4.41- 4.07 (m, 4H), 3.28 (s, 4H), 2.99 – 2.74 (m, 8H), 2.53 (s, 2H), 2.33-2.02 (d, J = 6.7 Hz, 3H), 1.78 (s, 3H), 1.40 (s, 4H), 1.26 (d, J = 13.8 Hz, 12H), 0.86 (s, 1H).	20

FIG. 3 Continued.


273		B	799.3	^{1}H NMR (400 MHz, DMSO-d6): δ 1.15 (s, 6H), 1.24 (s, 6H), 1.78 (t, *J* = 7.6 Hz, 2H), 1.99-2.03 (m, 1H), 2.33-2.35 (m, 2H), 2.54-2.60 (m, 2H), 2.69 (t, *J* = 6.8 Hz, 2H), 2.85-2.83 (m, 1H), 3.33 (s, 4H), 3.45 (s, 4H), 4.09 (d, *J* = 8.8 Hz, 1H), 4.10 (s, 1H), 55.05-5.09 (m, 1H), 7.25-7.23 (m, 5H), 7.41 (d, *J* = 2.0 Hz, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.78 (d, *J* = 7.6 Hz, 3H), 8.10 (d, *J* =12.8 Hz, 1H), 11.09 (s, 1H).	1
274		B	777.27	^{1}H NMR (400 MHz, CD_3OD): δ 1.19 (s, 6H), 1.26 (s, 6H), 1.69-1.85 (m, 4H), 2.07-2.13 (m, 1H), 2.65-2.75 (m, 4H), 2.80-2.86 (m, 3H), 2.91 (t, *J* =7.2 Hz, 2H), 3.09 (t, *J* = 5.2 Hz, 2H), 3.52 (s, 2H), 4.14 (s, 1H), 4.25 (s, 1H), 5.09-5.13 (m, 1H), 6.39 (s, 1H), 6.93-6.96 (m, 1H), 7.09 (d, *J* = 2.4 Hz, 1H), 7.43 (d, *J* = 8.0 Hz, 1H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.83 (d, *J* = 8.0 Hz, 1H), 7.87-7.90 (m, 1H), 7.93 (s, 1H), 8.11-8.13 (m, 1H), 8.89 (d, *J* = 2.0 Hz, 1H).	1, 105

FIG. 3 Continued.



275		B	807.2	^{1}HNMR (400 MHz, CD_3OD) : δ 1.25 (s, 6H), 1.32 (s, 6H), 1.61-1.64 (m, 2H), 1.78-1.82 (m, 2H), 1.91-1.98 (m, 2H), 2.13-2.16 (m, 1H), 2.76-2.88 (m, 7H), 3.09 (t, *J* = 5.4 Hz, 2H), 3.53 (s, 2H), 4.10-4.12 (m, 1H), 4.22 (t, *J* = 6.2 Hz, 2H), 4.36 (s, 1H), 5.11-5.15 (m, 1H), 6.43 (s, 1H), 7.00-7.03 (m, 1H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.33-7.35 (m, 1H), 7.42 (d, *J* = 2.0 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.82 (d, *J* = 8.4 Hz, 1H), 8.05-8.11 (m, 2H), 8.78 (s, 1H).	9, 101
276		B	833.28	^{1}HNMR (400 MHz, CD_3OD): δ 1.20 (s, 6H), 1.27 (s, 6H), 1.57-1.65 (m, 2H), 1.74-1.79 (m, 2H), 1.87-1.94 (m, 2H), 2.07-2.11 (m, 1H), 2.67-2.73 (m, 2H), 2.79-2.85 (m, 1H), 3.13 (t, *J* = 7.8 Hz, 2H), 4.13 (s, 2H), 4.17 (t, *J* = 7.0 Hz, 3H), 4.26 (s, 1H), 4.59 (s, 2H), 5.05-5.09 (m, 1H), 6.94-6.97 (m, 1H), 7.10 (d, *J* = 2.4 Hz, 1H), 7.27-7.30 (m, 1H), 7.37 (d, *J* = 2.4 Hz, 1H), 7.57 (s, 1H), 7.69 (d, *J* = 8.8 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 1H), 8.28-8.31 (m, 1H), 8.34 (br, 1H), 8.80 (d, *J* = 2 Hz, 1H).	86

FIG. 3 Continued.



277		B	781.51	^{1}HNMR (400 MHz, CD_3OD): δ 1.20 (s, 6H), 1.27 (s, 6H), 1.57-1.65 (m, 2H), 1.74-1.79 (m, 2H), 1.87-1.94 (m, 2H), 2.07-2.11 (m, 1H), 2.67-2.73 (m, 2H), 2.79-2.85 (m, 1H), 3.13 (t, *J* = 7.8 Hz, 2H), 4.13 (s, 2H), 4.17 (t, *J* = 7.0 Hz, 3H), 4.26 (s, 1H), 4.59 (s, 2H), 5.05-5.09 (m, 1H), 6.94-6.97 (m, 1H), 7.10 (d, *J* = 2.4 Hz, 1H), 7.27-7.30 (m, 1H), 7.37 (d, *J* = 2.4 Hz, 1H), 7.57 (s, 1H), 7.69 (d, *J* = 8.8 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 1H), 8.28-8.31 (m, 1H), 8.34 (br, 1H), 8.80 (d, *J* = 2 Hz, 1H).	17A
278		B	781.25	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (d, *J* = 5.6 Hz, 3H), 1.23 (s, 1H), 1.57 (s, 1H), 1.70-1.80 (m, 2H), 1.98-2.08 (m, 2H), 2.22-2.40 (m, 8H), 2.52-2.68 (m, 2H), 2.84-2.94 (m, 1H), 3.30 (s, 1H), 3.51 (s, 2H), 4.18 (t, *J* = 6.4 Hz, 2H), 4.42-4.48 (m, 3H), 5.08-5.15 (m, 1H), 6.43 (s, 1H), 6.94 (d, *J* = 2.4 Hz, 1H), 7.32-7.6 (m, 1H), 7.42 (d, *J* = 2.0 Hz, 1H), 7.80-7.85 (m, 2H), 7.99 (s, 2H), 8.08 (s, 1H), 8.23 (d, *J* = 8.0 Hz, 1H), 11.1 (s, 1H), 13.1 (s, 1H).	106

FIG. 3 Continued.



279		B	840.3	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.46-1.55 (m, 4H), 1.79-1.82 (m, 2H), 1.91-1.97 (m, 2H), 2.03-2.07 (m, 1H), 2.30 (br, 2H), 2.58-2.69 (m, 2H), 2.84-3.03 (m, 3H), 3.14-3.18 (m, 1H), 3.30 (s, 1H), 3.42 (br, 1H), 3.72-3.76 (m, 1H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.18-4.21 (m, 2H), 4.30-4.34 (m, 2H), 4.72 (br, 1H), 5.10-5.14 (m, 1H), 6.80-6.82 (m, 1H), 7.00-7.02 (m, 1H), 7.21 (d, *J* = 2.0Hz, 1H), 7.35-7.37 (m ,1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.59 (d, *J* = 8.8 Hz ,1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 7.94 (br, 1H), 8.61 (s, 1H), 11.12 (s, 1H).	14
280		B	807.28	^{1}H NMR (400 MHz, DMSO-d6): δ 1.14 (s, 6H), 1.24 (s, 6H), 1.45-1.49 (m, 2H), 1.56-1.60 (m, 2H), 1.78-1.82 (m, 2H), 1.99-2.05 (m, 2H), 2.45-2.47 (m, 3H), 2.58-2.67 (m, 5H), 2.84-2.94 (m, 1H), 3.16 (s, 1H), 4.08 (d, *J* = 9.2 Hz, 1H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.32 (s, 1H), 5.10-5.14 (m, 1H), 6.83 (s, 1H), 7.00-7.03 (m, 1H), 7.22 (d, *J* = 2.4Hz ,1H), 7.34-7.36 (m ,1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.64 (d, *J* = 8.0 Hz ,1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.03 (d, *J* = 9.2 Hz, 1H), 8.13-8.15 (m, 1H), 8.94 (s, 1H), 11.12 (s, 1H).	9, 101

FIG. 3 Continued.



281		B	824.27	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.24 (s, 6H), 1.42-1.64 (m, 4H), 1.72-1.87 (m, 2H), 1.99-2.10 (m, 1H), 2.56-2.70 (m, 3H), 2.77-2.95 (m, 3H), 3.22-3.29 (m, 2H), 3.84-4.00 (m, 2H), 4.05-4.13 (m, 1H), 4.15-4.26 (m, 2H), 4.29-4.37 (m, 1H), 5.07-5.17 (m, 1H), 6.96-7.09 (m, 1H), 7.16-7.29 (m, 1H), 7.33-7.50 (m, 2H), 7.76-7.97 (m, 2H), 8.01-8.22 (m, 3H), 8.81-8.91 (m, 1H), 11.07-11.19 (m, 1H).	87
282		B	794.29	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.24 (s, 6H), 1.42-1.64 (m, 4H), 1.72-1.87 (m, 2H), 1.99-2.10 (m, 1H), 2.56-2.70 (m, 3H), 2.77-2.95 (m, 3H), 3.22-3.29 (m, 2H), 3.84-4.00 (m, 2H), 4.05-4.13 (m, 1H), 4.15-4.26 (m, 2H), 4.29-4.37 (m, 1H), 5.07-5.17 (m, 1H), 6.96-7.09 (m, 1H), 7.16-7.29 (m, 1H), 7.33-7.50 (m, 2H), 7.76-7.97 (m, 2H), 8.01-8.22 (m, 3H), 8.81-8.91 (m, 1H), 11.07-11.19 (m, 1H).	9, 105

FIG. 3 Continued.



283		B	780.27	^{1}H NMR (400 MHz, DMSO-d6): δ 1.14 (s, 6H), 1.23 (s, 6H), 1.75-2.07 (m, 8H), 2.54-2.68 (m, 5H), 2.85-3.17 (m, 4H), 4.06(d, *J* = 9.2 Hz, 1H), 4.27 (br, 2H), 4.32 (s, 1H), 5.11-5.15 (m, 1H), 7.00-7.02 (m, 1H), 7.21 (d, *J* = 2.4Hz ,1H), 7.35-7.39 (m,3H), 7.46 (d, *J* =1.6 Hz, 1H), 7.78 (d, *J* = 8.8 Hz, 3H), 7.86 (d, *J* = 8.0 Hz, 1H), 7.91 (d, *J* = 8.8 Hz, 1H), 11.13 (s.1H).	9, 105
284		B	778.3	H-NMR (300 MHz, CD_3OD, ppm) δ 7.78 - 7.68 (m, 4 H), 7.41 - 7.34 (m, 3 H), 7.22 - 7.18 (m, 1 H), 7.13 (s, 1 H), 6.99 - 6.96 (d, *J* = 9 Hz, 1 H), 5.10 - 5.04 (m, 1 H), 4.27 (s, 1 H), 4.13 (s, 1 H), 3.76 - 3.65 (m, 4 H), 3.46 - 3.43 (m, 2 H), 3.32 - 3.31 (m, 2 H), 3.07 - 3.02 (m, 2 H), 2.87 - 2.68 (m, 5 H), 2.13 - 2.08 (m, 1 H), 1.25 (s, 6 H), 1.21 (s, 6 H).	107

FIG. 3 Continued.



285		B	810.23		9
286		B	739.16	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 2.04-2.16 (m, 3H), 2.56-2.69 (m, 2H), 2.85-2.97 (m, 2H), 3.76-3.79 (m ,2H), 4.05 (d, *J* = 9.2 Hz 1H), 4.16 (t, *J* = 8.4 Hz ,2H), 4.25 (t, *J* = 6.0 Hz ,2H), 4.30 (s, 1H), 5.13 (dd, *J* = 12.8 Hz ,1H), 6.37 (d, *J* = 8.8 Hz ,1H), 7.00 (dd, *J* = 8.8 Hz,1H), 7.20 (d, *J* = 2.4 Hz ,1H), 7.37 (dd, *J* = 8.0 Hz,1H), 7.45 (s, 1H), 7.60 (d, *J* = 9.2 Hz ,1H), 7.84-7.95 (m, 3H), 8.58 (d, *J* = 1.6 Hz ,1H), 11.12 (s, 1H).	9

FIG. 3 Continued.



287		B	767.19	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.42-1.46 (m, 2H), 1.66-1.81 (m, 4H), 2.03-2.07 (m, 1H), 2.50-2.62 (m, 2H), 2.73-2.78 (m, 1H), 2.85-2.87 (m, 1H), 3.64 (dd, *J* = 5.6 Hz, 8.0Hz, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.12 (t, *J* = 8.0 Hz, 2H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 5.12 (dd, *J* = 5.2 Hz, 12.8Hz, 1H), 6.36 (d, *J* = 8.8 Hz, 1H), 7.00 (dd, *J* = 2.4 Hz, 8.8Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 2.0 Hz, 8.4Hz, 1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.59 (d, *J* = 9.2 Hz, 1H), 7.83 (d, *J* = 8.0 Hz, 1H), 7.89-7.95 (m, 2H), 8.57 (d, *J* = 2.0 Hz, 1H), 11.12 (s, 1H).	9
288		B	818.3	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.42-1.46 (m, 2H), 1.66-1.81 (m, 4H), 2.03-2.07 (m, 1H), 2.50-2.62 (m, 2H), 2.73-2.78 (m, 1H), 2.85-2.87 (m, 1H), 3.64 (dd, *J* = 5.6 Hz, 8.0Hz, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.12 (t, *J* = 8.0 Hz, 2H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 5.12 (dd, *J* = 5.2 Hz, 12.8Hz, 1H), 6.36 (d, *J* = 8.8 Hz, 1H), 7.00 (dd, *J* = 2.4 Hz, 8.8Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 2.0 Hz, 8.4Hz, 1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.59 (d, *J* = 9.2 Hz, 1H), 7.83 (d, *J* = 8.0 Hz, 1H), 7.89-7.95 (m, 2H), 8.57 (d, *J* = 2.0 Hz, 1H), 11.12 (s, 1H).	43

FIG. 3 Continued.


289		B	836.35	^{1}H NMR (300 MHz, DMSO): δ 11.10 (s, 1H), 8.25 (d, J = 9.2 Hz, 1H), 7.92 (d, J = 8.8 Hz, 1H), 7.82 (d, J = 9.6 Hz, 1H), 7.69 (d, J = 8.5 Hz, 1H), 7.43-7.32 (m, 2H), 7.28-7.25 (m, 2H), 7.05-7.02 (m, 1H), 5.10-5.04 (m, 1H), 4.53-4.44 (m, 3H), 4.01 (d, J = 9.2 Hz, 1H), 3.45 (m, 4H), 3.14-2.79 (m, 3H), 2.64-2.51 (m, 2H), 2.45 (m, 3H), 2.41-2.34 (m, 2H), 2.08-1.97 (m, 1H), 1.88-1.62 (m, 3H), 1.47-1.43 (m, 2H), 1.22-1.08 (m, 15H);	17

FIG. 3 Continued.



290		B	836.4	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 8.59 (d, J = 1.2 Hz, 1H), 8.32 (d, J = 1.4 Hz, 1H), 7.90 (d, J = 8.7 Hz, 1H), 7.80 (d, J = 8.9 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.34 (s, 1H), 7.30 – 7.20 (m, 2H), 7.02 (dd, J = 8.8, 2.4 Hz, 1H), 5.07 (dd, J = 12.9, 5.4 Hz, 1H), 4.52 – 4.39 (m, 3H), 3.95 (d, J = 9.0 Hz, 1H), 3.43 (s, 4H), 3.06 – 2.79 (m, 5H), 2.36 (d, J = 7.6 Hz, 2H), 2.07 (s, 2H), 1.80 (d, J = 13.0 Hz, 2H), 1.48 – 1.36 (m, 4H), 1.15 (d, J = 18.7 Hz, 16H)	17
291		B	781.51	^{1}H NMR (400 MHz, DMSO-d6): δ 1.41-1.42 (m, 3H), 1.63-1.65 (m, 2H), 1.99-2.03 (m, 1H), 2.30-2.32 (m, 2H), 2.44 (br, 4H), 2.53-2.71 (m, 2H), 2.84-2.92 (m, 1H), 3.10-3.14 (m, 1H), 3.16-3.17 (m, 1H), 3.34-3.38 (m, 4H), 3.57-3.62 (m, 1H), 4.10-4.29 (m, 2H), 4.33-4.47 (m, 2H), 4.56-4.58 (m, 1H), 5.02-5.10 (m, 1H), 6.51 (br, 1H), 6.93-6.94 (m, 1H), 7.22-7.34 (m, 2H), 7.65-7.67 (m, 1H), 7.83 (s, 1H), 7.98 (s, 2H), 8.08 (s, 1H), 8.24-8.26 (m, 1H), 8.75 (br, 1H), 11.10 (s, 1H), 13.30 (br, 1H).	17A, 108

FIG. 3 Continued.



292		B	840.58	^{1}H NMR (400 MHz, CD_3OD): δ 1.23 (s, 6H), 1.31 (s, 6H), 1.53-1.73 (m, 4H), 1.88-1.95 (m, 2H), 2.11-2.17 (m, 1H), 2.55-2.80 (m, 5H), 2.84-2.94 (m, 2H), 2.05-2.09 (m, 1H), 3.12-3.22 (m, 2H), 3.63-3.85 (m, 4H), 4.08 (s, 1H), 4.20 (t, *J* = 6.2 Hz, 2H), 4.34 (s, 1H), 5.10-5.15 (m, 1H), 7.00-7.03 (m, 1H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.32 -7.34 (m, 1H), 7.40-7.43 (m, 2H), 7.74 (d, *J* = 8.4 Hz, 1H), 7.81 (d, *J* = 8.4 Hz, 1H), 7.94 (d, *J* = 8.8 Hz, 1H), 8.34 (d, *J* = 2.4 Hz, 1H).	14
293		B	822.35	^{1}H NMR (400 MHz, DMSO): δ 11.07 (s, 1H), 8.23 (d, J = 9.2 Hz, 1H), 7.89 (d, J = 8.7 Hz, 1H), 7.80 (d, J = 9.6 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.39-7.30 (m, 2H), 7.24 (m, 2H), 7.01 (m, 1H), 5.06 (m, 1H), 4.53-4.42 (m, 3H), 3.99 (m, 1H), 3.43 (m, 4H), 3.08-2.96 (m, 2H), 2.87 (m, 1H), 2.61-2.49 (m, 2H), 2.51 (m, 4H), 2.19 (m, 2H), 2.04-1.87 (m, 2H), 1.83 (m, 2H), 1.16 (m, 14H);	49

FIG. 3 Continued.



294		B	792.4	^{1}H-NMR (300 MHz, DMSO) δ11.07(s, 1H), 8.64- 8.60(m, 1H), 7.98- 7.87(m, 2H), 7.69- 7.55(m, 2H), 7.35- 7.20(m, 3H), 7.04- 6.96(m, 1H), 6.90- 6.83(m , 1H), 5.11 - 5.01 (m, 1H), 4.30(s, 1H), 4.09- 4.02(m, 1H), 3.70- 3.61(m, 4H), 3.58- 3.49(m, 4H), 2.97- 2.79(m, 1H), 2.65- 2.53(m, 2H), 2.05 - 1.97(m, 1H), 1.61- 1.55(m, 8H), 1.21 - 1.12 (d, *J=29.4 Hz,* 12H).	19
295		B	793.35	^{1}H-NMR (300 MHz, DMSO) δ11.07(s, 1H), 8.64- 8.60(m, 1H), 7.98- 7.87(m, 2H), 7.69- 7.55(m, 2H), 7.35- 7.20(m, 3H), 7.04- 6.96(m, 1H), 6.90- 6.83(m , 1H), 5.11 - 5.01 (m, 1H), 4.30(s, 1H), 4.09- 4.02(m, 1H), 3.70- 3.61(m, 4H), 3.58- 3.49(m, 4H), 2.97- 2.79(m, 1H), 2.65- 2.53(m, 2H), 2.05 - 1.97(m, 1H), 1.61- 1.55(m, 8H), 1.21 - 1.12 (d, *J=29.4 Hz,* 12H).	19

FIG. 3 Continued.



296		B	781.3	^{1}H NMR (300 MHz, CD_3OD) 7.75-7.70 (m, 3H), 7.48-7.44 (m, 1H), 7.36-7.33 (m, 2H), 7.16-7.12 (m, 2H), 6.99-6.96 (m, 1H), 5.08-5.02 (m, 1H), 4.63-4.61 (m, 1H), 4.28-4.26 (m, 1H), 4.14-4.11 (m, 1H), 3.73-3.55 (m, 3H), 3.07-3.02 (m, 1H), 2.86-2.64 (m, 8H), 2.11-1.92 (m, 3H), 1.27-1.22 (m, 12H);	1, 16
297		B	817.51	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.92-2.09 (m, 3H), 2.50 (s, 4H), 2.51-2.73 (m, 4H), 2.84-2.93 (m, 1H), 3.28 (br, 4H), 4.00 (d, *J* = 9.6 Hz, 1H), 4.23-4.26 (m, 3H), 5.10-5.15 (m, 1H), 6.67 (d, *J* = 12.0 Hz, 2H), 6.98-7.01 (m, 1H), 7.22 (d, *J* = 2.33 Hz, 1H), 7.35-7.38 (m, 1H), 7.45 (d, *J* = 2.0 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.18 (d, *J* = 9.6 Hz, 1H), 11.12 (s, 1H).	9

FIG. 3 Continued.




298		B	845.55	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.41-1.58 (m, 4H), 1.77-1.81 (m, 2H), 2.01-2.07 (m, 1H), 2.28-2.39 (m, 2H), 2.42-2.49 (m, 4H), 2.54-2.62 (m, 2H), 2.84-2.91 (m, 1H), 3.23 (br, 4H), 4.00 (d, *J* = 9.2 Hz, 1H), 4.19 (t, *J* = 6.4 Hz, 2H), 4.25 (s, 1H), 5.09-5.14 (m, 1H), 6.67 (d, *J* = 11.2 Hz, 2H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.34-7.36 (m, 1H), 7.43 (d, *J* = 2.0 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.18 (d, *J* = 9.2 Hz, 1H), 11.12 (s, 1H).	9
299		B	813.53	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.41-1.58 (m, 4H), 1.77-1.81 (m, 2H), 2.01-2.07 (m, 1H), 2.28-2.39 (m, 2H), 2.42-2.49 (m, 4H), 2.54-2.62 (m, 2H), 2.84-2.91 (m, 1H), 3.23 (br, 4H), 4.00 (d, *J* = 9.2 Hz, 1H), 4.19 (t, *J* = 6.4 Hz, 2H), 4.25 (s, 1H), 5.09-5.14 (m, 1H), 6.67 (d, *J* = 11.2 Hz, 2H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.34-7.36 (m, 1H), 7.43 (d, *J* = 2.0 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.18 (d, *J* = 9.2 Hz, 1H), 11.12 (s, 1H).	9
300		B	890.57	^{1}H NMR (400 MHz, DMSO-d6): δ 1.00-1.07 (m, 2H), 1.09 (s, 6H), 1.18 (s, 6H), 1.82-1.88 (m, 2H), 1.96-2.05 (m, 2H), 2.17-2.25 (m, 2H), 2.53-2.64 (m, 2H), 2.84-2.92 (m, 1H), 2.98-3.06 (m, 2H), 3.30-3.33 (m, 4H), 3.40-3.50 (m, 4H), 3.99 (d, *J* = 9.2 Hz, 1H), 4.26 (s, 1H), 4.63-4.72 (m, 2H), 5.05-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.22-7.28 (m, 2H), 7.33-7.37 (m, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.24 (d, *J* = 9.2 Hz, 1H), 8.60 (s, 1H), 11.09 (s, 1H).	49 starting with ethyl 2-chloro-4-(trifluoromethyl)pyrimidine-5-carboxylate

FIG. 3 Continued.



301		B	813.54		
302		B	888.58	^{1}H NMR (400 MHz, DMSO-d6): δ 1.00-1.07 (m, 2H), 1.09 (s, 6H), 1.18 (s, 6H), 1.82-1.88 (m, 2H), 1.96-2.05 (m, 2H), 2.17-2.25 (m, 2H), 2.53-2.64 (m, 2H), 2.84-2.92 (m, 1H), 2.98-3.06 (m, 2H), 3.30-3.33 (m, 4H), 3.40-3.50 (m, 4H), 3.99 (d, *J* = 9.2 Hz, 1H), 4.26 (s, 1H), 4.63-4.72 (m, 2H), 5.05-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.22-7.28 (m, 2H), 7.33-7.37 (m, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.24 (d, *J* = 9.2 Hz, 1H), 8.60 (s, 1H), 11.09 (s, 1H).	49 starting with 4-fluoro-3-(trifluoromethyl)benzoic acid

FIG. 3 Continued.



303		B	827.56
304		B	862.64

FIG. 3 Continued.



305		B	822.4	H-NMR (300 MHz, DMSO) δ 11.04 (s, 1H), 8.55 (s, 2H), 8.10-8.07 (d, J=9 Hz, 1H), 7.89-7.86 (d, J=9 Hz, 1H), 7.67-7.65 (d, J=8.4 Hz, 1H), 7.32 (s, 1H), 7.26-7.22 (m, 2H), 7.03-6.99 (m, 1H), 5.08-5.02 (m, 1H), 4.40 (s, 1H), 4.01-3.92 (m, 3H), 3.43 (s, 4H), 2.95-2.80 (m, 3H), 2.59 (s, 6H), 2.21-2.19 (d, J=6.3 Hz, 2H), 2.05-1.98 (m, 1H), 1.83-1.79 (d, J=11.1 Hz, 3H), 1.18 (s, 6H), 1.11 (s, 6H);	49 starting with 5-fluoropyrimidine-2-carbonitrile
306		B	836.5	H-NMR (300 MHz, DMSO) δ 11.04 (s, 1H), 8.55 (s, 2H), 8.09-8.06 (d, J=9 Hz, 1H), 7.89-7.86 (d, J=9 Hz, 1H), 7.67-7.64 (d, J=8.4 Hz, 1H), 7.32 (s, 1H), 7.25-7.22 (m, 2H), 7.03-6.99 (m, 1H), 5.08-5.02 (m, 1H), 4.40 (s, 1H), 4.00-3.92 (m, 3H), 3.42 (s, 4H), 2.91-2.83 (m, 3H), 2.59-2.54 (m, 6H), 2.39-2.25 (m, 2H), 2.05-1.98 (m, 1H), 1.80-1.76 (d, J=12 Hz, 2H), 1.58 (s,1H), 1.46-1.39 (m,2H),1.28-1.24 (d, J=11.1 Hz, 8H), 1.18 (s, 6H);	17 starting with 5-fluoropyrimidine-2-carbonitrile

FIG. 3 Continued.



307		B	838.2	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.06 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.69 (dd, J = 15.0, 10.0 Hz, 3H), 7.44 (dd, J = 18.6, 8.3 Hz, 2H), 7.17 (d, J = 2.4 Hz, 1H), 7.02 – 6.89 (m, 3H), 5.07 (dd, J = 12.9, 5.3 Hz, 1H), 4.28 (s, 1H), 4.07 – 3.95 (m, 1H), 3.77 (m, 2H),3.31 – 3.18 (m, 4H), 2.95 – 2.76 (m, 3H), 2.62 – 2.46 (m, 6H), 2.21 (d, J = 6.8 Hz, 2H), 1.99 (d, J = 19.3 Hz, 1H), 1.82 (d, J = 13.2 Hz, 3H), 1.34 (d, J = 27.3 Hz, 8H) , 1.11 (d, J = 27.3 Hz, 6H).	49
308		B	778.3	HNMR (300 MHz, DMSO-*d6*, ppm) δ 11.08 (s, 1 H), 7.92-7.89 (d, J= 8.7Hz, 1H), 7.76-7.68 (m, 3H), 7.44 - 7.213 (m, 4H), 7.02 - 7.01 (d, J= 2.4Hz, 1H), 6.47-6.44 (d, J= 8.7 Hz, 2H), 5.08-5.04 (m, 1H), 4.32(s, 1H), 4.07-4.02 (m, 3H), 3.77-3.73 (m, 2H), 3.49-3.46 (m, 4H), 3.32-3.31 (m, 1H), 2.96-2.82 (m, 1H), 2.60-2.55 (m, 6H), 2.09-1.95 (m, 1H), 1.22 (s, 6H), 1.12 (s, 6H);	19

FIG. 3 Continued.




309		B	822.4	^{1}H NMR (400 MHz, CD_3OD): δ7.82-7.80 (d, *J*=8.0Hz, 1H), 7.75-7.70 (m, 3H), 7.28-7.22 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.02 (m, 2H), 4.28 (s, 1H), 4.13 (s, 1H), 3.94-3.90 (m, 2H), 3.38-3.30(m, 6H), 2.82-2.70 (m, 5H), 2.67-2.64(m, 4H), 2.50-2.47 (m, 2H), 2.13-2.11 (m, 1H), 1.27 (s, 6H), 1.21 (s, 6H);	23
310		B	836.59	^{1}H NMR (400 MHz, CD_3OD): δ7.82-7.80 (d, *J*=8.0Hz, 1H), 7.75-7.70 (m, 3H), 7.28-7.22 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.02 (m, 2H), 4.28 (s, 1H), 4.13 (s, 1H), 3.94-3.90 (m, 2H), 3.38-3.30(m, 6H), 2.82-2.70 (m, 5H), 2.67-2.64(m, 4H), 2.50-2.47 (m, 2H), 2.13-2.11 (m, 1H), 1.27 (s, 6H), 1.21 (s, 6H);	49 starting with ethyl 2-chloro-4-methylpyrimidine-5-carboxylate
311		B	850.6	^{1}HNMR (400 MHz, DMSO-d6): δ 1.10 (s, 6H), 1.18 (s, 6H), 1.25 (d, *J* = 6.4 Hz, 2H), 1.80 (d, *J* = 13.2 Hz, 2H), 1.91 (s, 1H), 2.01-2.04 (m, 1H), 2.18-2.22 (m, 8H), 2.55-2.61 (m, 3H), 2.83-2.89 (m, 3H), 3.34 (s, 2H), 3.45 (br, 4H), 4.08 (d, *J* = 9.6 Hz, 1H), 4.26 (s, 1H), 4.72 (d, *J* = 12.4 Hz, 2H), 5.06-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.22-7.27 (m, 2H), 7.35 (s, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.23 (d, *J* = 9.2 Hz, 1H), 11.10 (s, 1H).	49 starting with ethyl 2-chloro-4-methylpyrimidine-5-carboxylate

FIG. 3 Continued.



312		B	795.55		
313		B	843.55		

FIG. 3 Continued.



314		B	823.6	1H NMR (300 MHz, DMSO-d6) δ 11.07 (s, 1H), 7.83 (dd, J = 20.2, 8.5 Hz, 2H), 7.69 (d, J = 8.6 Hz, 2H), 7.54-7.36 (m, 2H), 7.32 (dd, J = 8.3, 2.3 Hz, 1H), 7.17 (d, J = 2.3 Hz, 1H), 7.02-6.86 (m, 3H), 5.08 (dd, J = 12.8, 5.3 Hz, 1H), 4.28 (s, 1H), 4.16 (t, J = 6.3 Hz, 2H), 4.01 (d, J = 9.1 Hz, 1H), 3.66 (d, J = 11.5 Hz, 2H), 2.94 – 2.83 (m, 1H), 2.65 (dt, J = 29.5, 6.6 Hz, 5H), 2.53 (d, J = 3.1 Hz, 3H), 2.07-1.94 (m, 1H), 1.74-1.63 (m, 2H), 1.48 (p, J = 8.0 Hz, 2H), 1.18 (s, 6H), 1.14-0.98 (m, 12H)..	9
315		B	792.54	H-NMR (300 MHz, DMSO) δ 11.06 (s, 1H), 7.98-7.89 (d, J=9 Hz, 1H), 7.77-7.74 (d, J=8.7 Hz, 2H), 7.65-7.62 (d, J=8.4 Hz, 1H), 7.52-7.49 (d, J=9.3 Hz, 1H), 7.21-7.20 (d, J=2.4 Hz, 1H), 7.02-76.96 (m, 3H), 6.79 (s, 1H), 6.67-6.64 (d, J=8.1 Hz, 1H),5.08-5.02 (m, 1H), 4.32 (s, 1H), 4.18-4.04 (m, 3H), 3.74-3.70 (m, 2H), 3.30-3.26 (d, J=12 Hz, 1H), 3.05 (s, 1H), 2.94-2.82 (m, 1H), 2.73-2.27 (m, 11H), 2.03-1.99 (m,1H), 1.22 (s, 6H), 1.13 (s, 6H);	65

FIG. 3 Continued.



316		B	820.57	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.03 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.72 (d, J = 8.7 Hz, 2H), 7.62 (d, J = 8.5 Hz, 1H), 7.47 (d, J = 9.2 Hz, 1H), 7.31 (m, 1H), 7.20 – 7.13 (m, 2H), 7.02 – 6.89 (m, 3H), 5.23 (dd, J = 12.6, 5.4 Hz, 1H), 4.93 (dd, J = 12.6, 5.4 Hz, 3H), 3.31 – 3.17 (m, 3H), 3.02 – 2.76 (m, 3H), 2.62 – 2.49 (m, 6H), 2.17 (d, J = 6.7 Hz, 2H), 1.98 (dt, J = 10.9, 5.3 Hz, 1H), 1.90 – 1.74 (m, 3H), 1.14 (d, J = 27.4 Hz, 14H).	49, 70
317		B	840.57	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.03 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.72 (d, J = 8.7 Hz, 2H), 7.62 (d, J = 8.5 Hz, 1H), 7.47 (d, J = 9.2 Hz, 1H), 7.31 (m, 1H), 7.20 – 7.13 (m, 2H), 7.02 – 6.89 (m, 3H), 5.23 (dd, J = 12.6, 5.4 Hz, 1H), 4.93 (dd, J = 12.6, 5.4 Hz, 3H), 3.31 – 3.17 (m, 3H), 3.02 – 2.76 (m, 3H), 2.62 – 2.49 (m, 6H), 2.17 (d, J = 6.7 Hz, 2H), 1.98 (dt, J = 10.9, 5.3 Hz, 1H), 1.90 – 1.74 (m, 3H), 1.14 (d, J = 27.4 Hz, 14H).	14

FIG. 3 Continued.



318		B	814.53	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.19 (s, 6H), 1.62-1.67 (m, 2H), 1.77-1.84 (m, 2H), 2.00-2.07 (m, 1H), 2.24-2.44 (m, 6H), 2.54-2.62 (m, 2H), 2.86-2.90 (m, 1H), 3.60 (br, 4H), 3.96 (d, *J* = 9.6 Hz, 1H), 4.20-4.24 (m, 2H), 4.32 (s, 1H), 5.08-5.13 (m, 1H), 6.74-6.77 (m, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36-7.39 (m ,1H), 7.44-7.45 (m, 1H), 7.51-7.55 (m, 1H), 7.84 (d, *J* = 8.4 Hz ,1H), 7.91 (d, *J* = 8.8 Hz, 1H), 8.39 (d, *J* = 10.4 Hz 1H), 11.11 (s, 1H).	109
319		B	787.49	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.11 (s, 6H), 1.18 (s, 6H), 1.98-2.07 (m, 1H), 2.51-2.64 (m, 6H), 2.83-2.94 (m, 3H), 3.45 (s, 4H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.27 (s, 1H), 5.02-5.13 (m, 1H), 6.97-7.05 (m, 1H), 7.12 (d, *J* = 8.8 Hz, 2H), 7.23-7.29 (m, 2H), 7.36 (s, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 8.50 (d, *J* = 9.2 Hz, 1H), 11.10 (s, 1H).	110

FIG. 3 Continued.



320		B	752.49	^{1}HNMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.18 (s, 6H), 1.78-1.79 (m, 4H), 1.92-2.11 (m, 2H), 2.61-2.67 (m, 2H), 2.71-2.77 (m, 2H), 3.50 (t, *J* = 6.0 Hz, 2H), 3.97-4.04 (m, 3H), 4.09 (br, 2H), 4.18 (s, 1H), 4.92-5.03 (m, 1H), 6.35 (d, *J* = 8.4 Hz, 2H), 6.86-6.89 (m, 1H), 7.02 (d, *J* = 2.4 Hz, 1H), 7.21-7.23 (m, 1H), 7.29-7.31 (m, 2H), 7.59-7.63 (m, 3H), 7.70 (d, *J* = 8.0 Hz, 1H).	111
321		B	821.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.73-1.77 (m, 2H), 1.99-2.02 (m, 1H), 2.06-2.10 (m, 2H), 2.54-2.60 (m, 2H), 2.83-2.98 (m, 3H), 3.27-3.34 (m, 5H), 3.45-3.47 (m, 4H), 4.03-4.07 (m, 3H), 4.19 (s, 1H), 4.32 (s, 1H), 5.04-5.08 (m, 1H), 6.55 (d, *J* =8.8 Hz, 2H), 6.99-7.02 (m, 1H),7.20-7.23 (m, 2H), 7.29 (d, *J* =1.6 Hz, 1H), 7.40 (d, *J* = 9.2 Hz, 1H), 7.64 (d, *J* = 8.4 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.08 (s, 1H).	112

FIG. 3 Continued.


322		B	806.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.22 (s, 6H), 1.72-2.17 (m, 4H), 2.34-2.42 (m, 3H), 2.58-2.71 (m, 2H), 2.84-2.93 (m, 1H), 3.14-3.20 (m, 2H), 3.28-3.33 (m, 6H), 3.43-3.60 (m, 3H), 4.05-4.07 (m, 1H), 4.32 (s, 1H), 5.04-5.08 (m, 1H), 6.81-7.01 (m, 5H), 7.21 (s, 1H), 7.53 (d, *J* = 11.6 Hz, 1H), 7.64 (d, *J* = 7.2 Hz, 1H), 7.76 (d, *J* = 6.8 Hz, 2H), 7.90 (d, *J* = 7.2 Hz, 1H), 11.08 (s, 1H).	53

FIG. 3 Continued.



323		B	806.56		53
324		B	779.55		

FIG. 3 Continued.



325		B	821.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 9H), 1.73-1.74 (m, 2H), 1.80-1.84 (m, 1H), 1.99-2.02 (m, 1H), 2.06-2.08 (m, 2H), 2.33-2.39 (m, 3H), 2.54-2.60 (m, 2H), 2.83-2.98 (m, 3H), 3.27-3.27 (m, 2H), 3.33-3.34 (m, 2H), 3.46-3.50 (m, 1H), 4.03-4.07 (m, 3H), 4.19 (s, 1H), 4.32 (s, 1H), 5.04-5.08 (m, 1H), 6.55 (d, *J* = 8.8 Hz, 2H), 6.99-7.02 (m, 1H), 7.20-7.23 (m, 2H), 7.30 (s, 1H), 7.40 (d, *J* = 9.2 Hz, 1H), 7.64 (d, *J* = 8.8 Hz, 1H), 7.75 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.4 Hz, 1H), 11.12 (s, 1H).	113
326		B	811.4	^{1}H-NMR (300 MHz, CD3OD) δ8.71(s, 2H), 7.75- 7.67(m, 2H), 7.21- 7.17(m, 1H), 7.16- 7.08(m, 2H), 7.01- 6.93(m, 1H), 5.16- 5.06 (m, 1H), 4.26(s, 1H), 4.12(s, 1H), 4.02- 3.66(m, 6H), 3.11- 3.3.03(m, 2H), 2.94- 2.77(m, 3H), 2.75- 2.58(m, 1H), 2.48- 2.29(m, 1H), 2.16 - 2.04(m, 1H), 1.82- 1.65(m, 2H), 1.27 - 1.20 (d, *J=20.7 Hz,* 12H), 1.06- 0.09(m, 6H)	9

FIG. 3 Continued.



327		B	806.55	^{1}HNMR (400 MHz, DMSO-d_6) : δ 1.13 (s, 6H), 1.22 (s, 6H), 1.78-1.84 (m, 1H), 2.02-2.05 (m, 1H), 2.25-2.27 (m, 1H), 2.55-2.62 (m, 2H), 2.85-2.93 (m, 2H), 3.08-3.21 (m, 3H), 3.32-3.47 (m, 6H), 3.60-3.73 (m, 3H), 4.07-4.08 (m, 1H), 4.25-4.28 (m, 2H), 4.33 (s, 2H), 5.09-5.13 (m, 1H), 6.57 (d, J = 8.8 Hz, 2H), 7.00-7.02 (m, 1H), 7.20-7.21 (m, 1H), 7.38-7.45 (m, 2H), 7.52 (s, 1H), 7.78 (d, J = 8.4 Hz, 3H), 7.91 (d, J = 8.4 Hz, 1H), 9.97 (br, 1H), 11.12 (s, 1H).	53
328		B	824.55	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.71-1.79 (m, 1H), 2.00-2.05 (m, 1H), 2.12-2.15 (m, 1H), 2.42 (d, J = 6.4 Hz, 2H), 2.57-2.61 (m, 6H), 2.84-2.93 (m, 1H), 3.05 (t, J = 7.2 Hz, 1H), 3.26-3.31 (m, 6H), 3.38-3.40 (m, 1H), 3.46 (t, J = 8.4 Hz, 1H), 4.05 (d, J = 8.8 Hz, 1H), 4.32 (s, 1H), 5.09-5.13 (m, 1H), 6.54 (d, J = 8.8 Hz, 2H), 6.99-7.01 (m, 1H), 7.20 (d, J = 2.0 Hz, 1H), 7.39 (d, J =9.2 Hz, 1H), 7.46 (d, J =3.6 Hz, 1H), 7.72-7.75 (m, 3H), 7.90 (d, J = 8.4 Hz, 1H), 11.12 (s, 1H).	53

FIG. 3 Continued.



329		B	765.56	^{1}H NMR (400 MHz, DMSO-d^6) δ 1.12 (6H, s), 1.22 (6H, s), 1.50-1.53 (2H, m), 1.63-1.66 (2H, m), 1.99-2.02 (1H, m), 2.34-2.47 (7H, m), 2.55-2.59 (1H, m), 2.66-2.70 (2H, m), 2.88-2.97 (1H, m), 3.25-3.31 (4H, m), 4.05 (1H, d, *J* = 9.2 Hz), 4.30-4.34 (2H, m), 4.46-4.50 (1H, m), 5.14 (1H, dd, *J* = 13.2, 4.8 Hz), 6.95-7.02 (3H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.46-7.48 (2H, m), 7.51-7.57 (2H, m), 7.74 (2H, d, *J* = 8.8 Hz), 7.91 (1H, d, *J* = 8.8 Hz), 11.02 (1H, s).	8A
330		B	736.5	^{1}H NMR (400 MHz, DMSO-d^6) δ 1.12 (6H, s), 1.22 (6H, s), 1.50-1.53 (2H, m), 1.63-1.66 (2H, m), 1.99-2.02 (1H, m), 2.34-2.47 (7H, m), 2.55-2.59 (1H, m), 2.66-2.70 (2H, m), 2.88-2.97 (1H, m), 3.25-3.31 (4H, m), 4.05 (1H, d, *J* = 9.2 Hz), 4.30-4.34 (2H, m), 4.46-4.50 (1H, m), 5.14 (1H, dd, *J* = 13.2, 4.8 Hz), 6.95-7.02 (3H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.46-7.48 (2H, m), 7.51-7.57 (2H, m), 7.74 (2H, d, *J* = 8.8 Hz), 7.91 (1H, d, *J* = 8.8 Hz), 11.02 (1H, s).	113

FIG. 3 Continued.



331		B	779.54	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.13 (6H, s), 1.22 (6H, s), 1.50-1.55 (2H, m), 1.62-1.67 (2H, m), 2.04-2.08 (1H, m), 2.33-2.36 (2H, m), 2.47-2.50 (4H, m), 2.55-2.62 (2H, m), 2.86-2.90 (1H, m), 3.07 (2H, t, J = 7.2 Hz), 3.23 (4H, s), 4.05 (1H, d, J = 9.2 Hz), 4.32 (1H, s), 5.14 (1H, dd, J = 12.8, 5.6 Hz), 6.95-7.02 (3H, m), 7.21 (1H, d, J = 2.4 Hz), 7.52 (1H, d, J = 9.2 Hz), 7.71-7.81 (5H, m), 7.91 (1H, d, J = 8.8 Hz), 11.13 (1H, s).	8A
332		B	798.2	^{1}H NMR (400 MHz, CD_3OD): δ 8.51 (s, 1H), 7.71-7.66 (m, 2H), 7.35-7.34 (d, J=1.6Hz, 1H), 7.23-7.21 (d, J=8.4Hz, 1H), 7.10-7.09 (d, J=2.0Hz, 1H), 6.96-6.93 (m, 1H), 5.09-5.04 (m, 1H), 4.19 (s, 1H), 3.98 (s, 1H), 3.52-3.31 (m, 7H), 3.06-3.02 (m, 1H), 2.87-2.81 (m, 1H), 2.76-2.51 (m, 10H), 2.42-2.31 (m, 4H), 2.11-2.10 (m, 3H), 1.90 (m, 1H), 1.45-1.35 (m, 2H), 1.20 (s, 6H), 1.10 (s, 6H);	114

FIG. 3 Continued.



333		B	822.03	^{1}H NMR (400 MHz, CD_3OD): δ 8.51 (s, 1H), 7.71-7.66 (m, 2H), 7.35-7.34 (d, *J*=1.6Hz, 1H), 7.23-7.21 (d, *J*=8.4Hz, 1H), 7.10-7.09 (d, *J*=2.0Hz, 1H), 6.96-6.93 (m, 1H), 5.09-5.04 (m, 1H), 4.19 (s, 1H), 3.98 (s, 1H), 3.52-3.31 (m, 7H), 3.06-3.02 (m, 1H), 2.87-2.81 (m, 1H), 2.76-2.51 (m, 10H), 2.42-2.31 (m, 4H), 2.11-2.10 (m, 3H), 1.90 (m, 1H), 1.45-1.35 (m, 2H), 1.20 (s, 6H), 1.10 (s, 6H);	49
334		B	821.55	^{1}H-NMR (300 MHz, CD_3OD) δ 7.82- 7.68 (m, 4H), 7.28- 7.10 (m, 3H), 7.02- 6.94 (m, 1H), 6.64- 6.57 (m, 2H), 5.14 - 5.05 (m, 1H), 4.94-4.90 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.69- 3.35 (m, 3H), 3.22- 3.06 (m, 2H), 2.96- 2.57 (m, 4H), 2.56-2.24 (m, 6H), 2.24- 2.03 (m, 5H), 1.92- 1.74 (m, 1H), 1.28- 1.22 (d, *J=18 Hz,* 12H);	73

FIG. 3 Continued.



335		B	821.56	^{1}H-NMR (300 MHz, CD_3OD) δ 7.82- 7.68 (m, 4H), 7.28- 7.10 (m, 3H), 7.02- 6.94 (m, 1H), 6.64- 6.57 (m, 2H), 5.14 - 5.05 (m, 1H), 4.94- 4.90 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.69- 3.35 (m, 3H), 3.22- 3.06 (m, 2H), 2.96- 2.57 (m, 4H), 2.56- 2.24 (m, 6H), 2.24- 2.03 (m, 5H), 1.92- 1.74 (m, 1H), 1.28- 1.22 (d, *J=18 Hz*, 12H);	113
336		B	821.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.52-1.54 (m, 2H), 1.81-1.83 (m, 1H), 1.86-1.94 (m, 3H), 1.98-2.03 (m, 1H), 2.16-2.20 (m, 1H), 2.55-2.60 (m, 2H), 2.70 (s, 2H), 2.84-2.93 (m, 1H), 3.03-3.08 (m, 1H), 3.18-3.22 (m, 1H), 3.29-3.32 (m, 2H), 3.48-3.55 (m, 3H), 3.57-3.63 (m, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.32 (s, 1H), 5.05 (d, *J* = 12.8 Hz, 1H), 6.81 (d, *J* = 8.4 Hz, 1H), 6.90 (d, *J* = 1.6 Hz, 1H), 6.96-7.02 (m, 3H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.50 (d, *J* = 9.2 Hz, 1H), 7.63(d, *J* = 8.8 Hz, 1H), 7.73(d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.09 (s, 1H).	113

FIG. 3 Continued.




337		B	807.54	^{1}H NMR (400 MHz, *d6*-DMSO): δ11.11 (s, 1H), 7.91-7.73 (m, 4H), 7.44-7.01 (m, 5H), 6.43 (s, 2H), 5.12-4.96 (m, 2H), 4.31 (s, 1H), 4.04-3.98 (m, 3H), 3.53-3.33 (m, 3H), 3.08 (m, 1H), 3.01-2.92 (m, 2H), 2.61-2.30 (m, 5H), 2.23-2.08 (m, 6H), 1.21 (s, 6H), 1.12 (s, 6H);	73
338		B	821.45	^{1}H-NMR (300 MHz, CD_3OD) δ7.82- 7.68 (m, 4H), 7.28- 7.10 (m, 3H), 7.02- 6.94 (m, 1H), 6.64- 6.57 (m, 2H), 5.14 - 5.05 (m, 1H), 5.01-4.90 (m, 1H), 4.28 (s, 1H), 4.15-4.10 (m, 1H), 3.60- 3.33 (m, 3H), 3.21- 3.09 (m, 2H), 2.92- 2.30 (m, 10H), 2.28- 2.05 (m, 5H), 1.90-1.73 (m, 1H), 1.28- 1.22 (d, *J=18 Hz,* 12H).	73
339		B	839.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.00-1.08 (m, 2H), 1.12 (s, 6H), 1.19 (s, 6H), 1.73-1.94 (m, 3H), 1.99-2.06 (m, 1H), 2.19 (d, J = 6.8 Hz, 2H), 2.46-2.49 (m, 4H), 2.54-2.67 (m, 2H), 2.84-2.96 (m, 3H), 3.38-3.49 (m, 4H), 3.95 (d, J = 8.8 Hz, 1H), 4.31-4.40 (m, 3H), 5.05-5.10 (m, 1H), 6.71, 6.75 (two singles, 1H), 7.00-7.02 (m, 1H), 7.22-7.27 (m, 2H), 7.35 (s, 1H), 7.46-7.49 (m, 1H), 7.68 (d, J = 8.4 Hz, 1H), 7.89 (d, J = 8.8 Hz, 1H), 8.39 (d, J = 11.2 Hz, 1H), 11.10 (s, 1H).	49, 109

FIG. 3 Continued.


340		B	852.59	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.47-1.51 (m, 3H), 1.67-1.78 (m, 3H), 1.99-2.03 (m, 2H), 2.23 (s, 3H), 2.43-2.45 (m, 2H), 2.53-2.60 (m, 3H), 2.73-2.92 (m, 3H), 3.32 (s, 1H), 3.55-3.65 (m, 3H), 3.89-3.92 (m, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.31 (s, 1H), 5.04-5.09 (m, 1H), 6.94-7.04 (m, 4H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.41 (d, *J* = 9.2 Hz, 1H), 7.60 (d, *J* = 12.4 Hz, 1H), 7.73 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.10 (s, 1H).	115

FIG. 3 Continued.


341		B	852.59	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.48-1.49 (m, 3H), 1.67-1.78 (m, 3H), 1.99-2.06 (m, 2H), 2.23 (s, 3H), 2.43-2.45 (m, 2H), 2.53-2.60 (m, 3H), 2.73-2.92 (m, 3H), 3.32 (s, 1H), 3.55-3.65 (m, 3H), 3.89-3.92 (m, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.31 (s, 1H), 5.04-5.09 (m, 1H), 6.94-7.04 (m, 4H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.51 (d, *J* = 9.2 Hz, 1H), 7.60 (d, *J* = 12.8 Hz, 1H), 7.73 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.10 (s, 1H).	115

FIG. 3 Continued.



342		B	861.7	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.08 (s, 1H), 8.61 (d, J = 2.3 Hz, 1H), 7.91 (dd, J = 11.7, 8.8 Hz, 2H), 7.70 (d, J = 8.4 Hz, 1H), 7.60 (d, J = 9.3 Hz, 1H), 7.38 (s, 1H), 7.28 (d, J = 8.8 Hz, 1H), 7.20 (d, J = 2.4 Hz, 1H), 7.00 (dd, J = 8.8, 2.5 Hz, 1H), 6.87 (d, J = 9.1 Hz, 1H), 5.07 (dd, J = 12.8, 5.3 Hz, 1H), 4.43 (dd, J = 25.1, 12.8 Hz, 2H), 4.30 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 2.98 - 2.82 (m, 7H), 2.62 (m, 2H),2.51 (m, 1H), 2.01-1.75 (m, 5H), 1.64 (d, J = 11.7 Hz, 1H), 1.50-1.11 (m, 21H).	94
343		B	856.56	H-NMR-PH-ARV-LS-049-E-0: (300 MHz, DMSO-*d6*, ppm) δ 11.11 (br, 1H), 7.92-7.89 (d, *J*=8.7Hz, 1H), 7.75-7.61 (m, 4H), 7.47-7.45 (d, *J* =7.5Hz, 1H),7.22-7.21 (d, *J* =2.4Hz, 1H), 7.12-7.06 (m, 1H), 7.03-6.99 (m, 1H), 5.14-5.08 (m, 1H), 4.32 (s, 1H), 4.08-4.05 (d, *J* =9.3Hz, 1H), 3.51-3.47 (m, 2H), 3.26-3.24 (m, 4H), 2.89-2.82 (m, 1H), 2.78-2.71 (m, 2H), 2.62-2.55 (m, 6H), 2.27-2.25 (m, 2H), 2.08-2.04 (m, 1H), 1.86-1.73 (m, 3H), 1.31-1.28 (m, 2H), 1.22 (s, 6H), 1.13 (s, 6H);	49

FIG. 3 Continued.



344		B	806.55	H-NMR (300 MHz, DMSO-*d6*, ppm) δ 11.11 (br, 1H), 7.92-7.89 (d, *J*=8.7Hz, 1H), 7.75-7.61 (m, 4H), 7.47-7.45 (d, *J* =7.5Hz, 1H),7.22-7.21 (d, *J* =2.4Hz, 1H), 7.12-7.06 (m, 1H), 7.03-6.99 (m, 1H), 5.14-5.08 (m, 1H), 4.32 (s, 1H), 4.08-4.05 (d, *J* =9.3Hz, 1H), 3.51-3.47 (m, 2H), 3.26-3.24 (m, 4H), 2.89-2.82 (m, 1H), 2.78-2.71 (m, 2H), 2.62-2.55 (m, 6H), 2.27-2.25 (m, 2H), 2.08-2.04 (m, 1H), 1.86-1.73 (m, 3H), 1.31-1.28 (m, 2H), 1.22 (s, 6H), 1.13 (s, 6H);	52
345		B	824.4	1H NMR (300 MHz, DMSO-*d6*) δ 11.04 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 (m, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.06 – 6.89 (m, 4H), 5.03 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.28 (s, 1H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.78-3.56 (m, 3H), 3.35-3.26 (m, 4H), 2.94 – 2.75 (m, 1H), 2.60 – 2.48 (m, 5H), 2.45-2.37 (m, 3H), 2.32 – 2.28 (m, 2H), 2.19-1.98 (m, 2H), 1.88-1.75 (m, 1H), 1.18 (s, 6H), 1.09 (s, 6H).	53

FIG. 3 Continued.



346		B	840.55	1H NMR (300 MHz, DMSO-*d*6) δ 11.04 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 (m, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.06 – 6.89 (m, 4H), 5.03 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.28 (s, 1H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.78-3.56 (m, 3H), 3.35-3.26 (m, 4H), 2.94 – 2.75 (m, 1H), 2.60 – 2.48 (m, 5H), 2.45-2.37 (m, 3H), 2.32 – 2.28 (m, 2H), 2.19-1.98 (m, 2H), 1.88-1.75 (m, 1H), 1.18 (s, 6H), 1.09 (s, 6H).	49
347		B	938.66	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.22 (6H, s), 1.29-1.34 (2H, m), 1.68-1.73 (1H, m), 1.80-1.83 (2H, m), 1.99-2.03 (1H, m), 2.24 (2H, d, *J* = 6.8 Hz), 2.51-2.61 (8H, m), 2.84-2.92 (1H, m), 3.25 (3H, s), 3.46-3.48 (6H, m), 3.55-3.57 (2H, m), 3.60-3.64 (2H, m), 3.78-3.80 (2H, m), 4.05 (1H, d, *J* = 9.6 Hz), 4.14-4.16 (2H, m), 4.32 (1H, s), 5.05-5.10 (1H, m), 6.93 (1H, d, *J* = 8.4 Hz), 6.99-7.02 (1H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.25-7.28 (1H, m), 7.32-7.35 (2H, m), 7.44 (1H, d, *J* = 8.4 Hz), 7.60 (1H, d, *J* = 9.2 Hz), 7.69 (1H, d, *J* = 8.4 Hz), 7.90 (1H, d, *J* = 8.8 Hz), 11.09 (1H, s).	52, 116

FIG. 3 Continued.



348		B	777.56	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.22 (6H, s), 1.29-1.34 (2H, m), 1.68-1.73 (1H, m), 1.80-1.83 (2H, m), 1.99-2.03 (1H, m), 2.24 (2H, d, *J* = 6.8 Hz), 2.51-2.61 (8H, m), 2.84-2.92 (1H, m), 3.25 (3H, s), 3.46-3.48 (6H, m), 3.55-3.57 (2H, m), 3.60-3.64 (2H, m), 3.78-3.80 (2H, m), 4.05 (1H, d, *J* = 9.6 Hz), 4.14-4.16 (2H, m), 4.32 (1H, s), 5.05-5.10 (1H, m), 6.93 (1H, d, *J* = 8.4 Hz), 6.99-7.02 (1H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.25-7.28 (1H, m), 7.32-7.35 (2H, m), 7.44 (1H, d, *J* = 8.4 Hz), 7.60 (1H, d, *J* = 9.2 Hz), 7.69 (1H, d, *J* = 8.4 Hz), 7.90 (1H, d, *J* = 8.8 Hz), 11.09 (1H, s).	117
349		B	821.56	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.12 (6H, s), 1.26 (6H, s), 1.49-1.54 (4H, m), 1.89-1.91 (4H, m), 1.99-2.02 (1H, m), 2.54-2.67 (1H, m), 2.85-2.89 (1H, m), 3.04 (2H, t, *J* = 10.0 Hz), 3.23-3.26 (3H, m), 3.64-3.83 (6H, m), 4.04 (1H, d, *J* = 9.2 Hz), 4.32 (1H, s), 5.04 (1H, dd, *J* = 5.2, 12.8 Hz), 6.96-7.02 (3H, d, m), 7.20 (1H, d, *J* = 2.4 Hz), 7.24 (1H, d, *J* = 10.4 Hz), 7.33 (1H, s), 7.49 (1H, d, *J* = 8.8 Hz), 7.65 (1H, d, *J* = 8.8 Hz), 7.72 (2H, d, *J* = 8.8 Hz), 7.89 (1H, d, *J* = 8.8 Hz), 11.08 (1H, s).	52, 118

FIG. 3 Continued.




350		B	839.35	^{1}H-NMR (400 MHz, CD_3OD) δ8.54（s, 1H), 7.77- 7.64(m, 2H), 7.36(s, 1H), 7.26- 7.12(m, 2H), 7.02- 6.96(m, 1H), 6.65- 6.57(m , 1H), 5.12 - 5.04 (m, 1H), 4.28(s, 1H), 4.12- 4.04(m, 3H), 3.71(m, 4H), 3.08- 2.98(m, 2H), 2.90- 2.55(m, 7H), 2.37- 2.31(m, 2H), 2.15- 1.93(m, 4H), 1.37- 1.30(m, 2H), 1.28 - 1.23 (d, J=21 Hz, 12H).	52, 109
351		B	763.5		
352		B	818.55	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.23 (s, 6H), 1.88-1.92 (m, 1H), 1.99-2.03 (m, 1H), 2.18 (t, J = 8 Hz, 1H), 2.25 (br, 2H), 2.46 (br, 3H), 2.55-2.60 (m, 1H), 2.70 (s, 1H), 2.84-2.89 (m, 1H), 2.96 (s, 2H), 3.46-3.47 (m, 5H), 3.79 (s, 2H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.35 (s, 1H), 5.05-5.11 (m, 1H), 5.76 (s, 1H), 6.96 (d, *J* = 8.8 Hz, 2H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.26 (d, *J* = 8.4 Hz, 1H), 7.34 (s, 1H), 7.53 (d, *J* = 8.4 Hz, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.78 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.08 (s, 1H).	119

FIG. 3 Continued.




353		B	821.2	^{1}H NMR (300 MHz, $CDCl_3$) 7.74-7.67 (m, 3H), 7.58-7.55(m, 1H), 7.29-7.26 (m, 1H), 7.04-7.08 (m, 1H), 6.97-6.93 (m, 3H), 6.83-6.81 (m, 1H), 6.14-6.11 (m, 1H), 4.97-4.91 (m, 1H), 4.42 (s, 1H), 4.16-4.13 (m, 1H), 4.05-3.99 (m, 3H), 3.04-2.92 (m, 4H), 2.88-2.51 (m, 3H) , 2.16-2.10 (m, 3H), 1.96-1.88 (m, 3H), 1.84-1.73(m, 2H) ,1.67-1.65 (m, 4H), 1.46 (m, 6H), 1.27-1.22 (m, 6H);	120
354		B	868.3	^{1}H NMR (400 MHz, $CDCL_3$): δ 8.04 (s, 1H), 7.55 (d, J = 8.7 Hz, 1H), 7.41 (dd, *J* = 25.5, 9.1 Hz, 2H), 6.93 (d, *J* = 2.4 Hz, 1H), 6.77 (dd, *J* = 8.7, 2.4 Hz, 1H), 5.66 (d, *J* = 8.5 Hz, 1H), 4.97-4.78 (m, 3H), 4.14 (d, *J* = 8.5 Hz, 1H), 3.99 (s, 1H), 3.27 (s, 4H), 2.96-2.63 (m, 6H), 2.59 (s, 4H), 2.33 (s, 6H), 2.13 (dd, *J* = 14.4, 6.4 Hz, 1H), 1.83 (d, *J* = 13.2 Hz, 3H), 1.20 (s, 12H), 1.11 (s, 2H).	63
355		B	786.2	^{1}H NMR (400 MHz, $CDCL_3$): δ 8.04 (s, 1H), 7.55 (d, J = 8.7 Hz, 1H), 7.41 (dd, *J* = 25.5, 9.1 Hz, 2H), 6.93 (d, *J* = 2.4 Hz, 1H), 6.77 (dd, *J* = 8.7, 2.4 Hz, 1H), 5.66 (d, *J* = 8.5 Hz, 1H), 4.97-4.78 (m, 3H), 4.14 (d, *J* = 8.5 Hz, 1H), 3.99 (s, 1H), 3.27 (s, 4H), 2.96-2.63 (m, 6H), 2.59 (s, 4H), 2.33 (s, 6H), 2.13 (dd, *J* = 14.4, 6.4 Hz, 1H), 1.83 (d, *J* = 13.2 Hz, 3H), 1.20 (s, 12H), 1.11 (s, 2H).	49

FIG. 3 Continued.



356		B	854.2	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.11 (d, J = 8.6 Hz, 1H), 7.72 (dd, J = 17.8, 8.5 Hz, 3H), 7.52 (d, J = 9.2 Hz, 1H), 7.44-7.22 (m, 4H), 6.96 (d, J = 8.6 Hz, 2H), 5.08 (dd, J = 12.7, 5.4 Hz, 1H), 4.40 (s, 1H), 4.08 (d, J = 9.1 Hz, 1H), 3.95-3.75 (m, 2H), 3.60-3.40 (m, 4H), 3.38-3.20 (m, 4H), 2.95-2.70 (m, 3H), 2.68-2.59 (m, 2H), 2.32 (brs, 2H), 2.04-2.02 (m, 1H), 1.82-1.70 (m, 3H), 1.19 (d, J = 26.9 Hz, 14H).	49
357		B	778.5	^{1}H NMR (300 MHz, *d6*-DMSO) δ 1.13 (s, 6H), 1.22 (s, 6H), 1.89-2.08 (m, 5H), 2.31-2.40 (m, 3H), 2.51-2.64 (m, 2H), 2.82-2.98 (m, 1H), 3.80 (s, 2H), 3.92 (s, 2H), 4.04-4.07 (m, 1H), 4.14 (s, 2H), 4.32 (s, 1H), 5.10-5.16 (m, 1H), 6.40-6.42 (m, 2H), 6.99-7.02 (m, 1H), 7.20 (s, 1H), 7.34-7.36 (m, 1H), 7.43-7.46 (m, 2H), 7.72-7.74 (m, 2H), 7.83-7.92 (m, 2H), 11.1 (s, 1H);	121

FIG. 3 Continued.



358		B	811.32	1H NMR (300 MHz, CDCL3) δ 8.29 (s, 1H), 7.81 (d, J = 8.3 Hz, 1H), 7.60 (d, J = 8.6 Hz, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.22 (dd, J = 8.3, 2.3 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.83 (dd, J = 8.8, 2.4 Hz, 1H), 5.96 (d, J = 8.6 Hz, 1H), 4.99 (dd, J = 11.9, 5.5 Hz, 1H), 4.24 (dt, J = 11.9, 7.0 Hz, 7H), 4.08 (s, 1H), 3.09 (s, 9H), 2.98 – 2.77 (m, 5H), 2.40(s, 2H), 2.21- 2.14 (m, 1H), 1.27 (d, J = 3.9 Hz, 12H);	9, 63
359		B	783.1	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.13 (s, 1H), 8.65 (s, 1H), 8.38 (s, 1H), 8.15 (s, 1H), 7.96-7.81 (m, 3H), 7.47 (d, *J* = 2.2 Hz, 1H), 7.38 (dd, *J* = 8.4, 2.3 Hz, 1H), 7.26 (d, *J* = 2.4 Hz, 1H), 7.04 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.14 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.45 (s, 1H), 4.28 (t, *J* = 6.1 Hz, 2H), 3.98 (d, *J* = 9.0 Hz, 1H), 3.74 (s, 3H), 2.89 (d, *J* = 12.5 Hz, 1H), 2.60 (d, *J* = 15.9 Hz, 4H), 2.04 (s, 4H), 1.17 (d, *J* = 18.4 Hz, 12H), 1.05 (s, 4H);	9, 63

FIG. 3 Continued.




360		B	783.1	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.27 – 8.12 (m, 1H), 7.92 – 7.76 (m, 3H), 7.46 – 7.29 (m, 3H), 7.22 (d, J = 2.4 Hz, 1H), 7.00 (dd, J = 8.8, 2.4 Hz, 1H), 5.09 (dd, J = 12.9, 5.3 Hz, 1H), 4.43 (s, 1H), 4.23 (t, J = 6.3 Hz, 2H), 3.97 (d, J = 9.1 Hz, 1H), 3.70 (t, J = 4.9 Hz, 4H), 2.95 – 2.76 (m, 1H), 2.59-2.50 (s, 6H),2.45(s,2H), 2.07 – 1.89 (m, 3H), 1.19 (s, 6H), 1.11 (s, 6H).	9, 63
361		B	854.3	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.27 – 8.12 (m, 1H), 7.92 – 7.76 (m, 3H), 7.46 – 7.29 (m, 3H), 7.22 (d, J = 2.4 Hz, 1H), 7.00 (dd, J = 8.8, 2.4 Hz, 1H), 5.09 (dd, J = 12.9, 5.3 Hz, 1H), 4.43 (s, 1H), 4.23 (t, J = 6.3 Hz, 2H), 3.97 (d, J = 9.1 Hz, 1H), 3.70 (t, J = 4.9 Hz, 4H), 2.95 – 2.76 (m, 1H), 2.59-2.50 (s, 6H),2.45(s,2H), 2.07 – 1.89 (m, 3H), 1.19 (s, 6H), 1.11 (s, 6H).	49
362		B	872.7	H-NMR (400 MHz, DMSO-*d6*) 11.12 (s, 1H),7.91-7.86 (m, 2H),7.75-7.72 (d, J = 11.6Hz, 1H), 7.47-7.45 (d, J=7.6Hz, 1H), 7.34-7.32 (d, J = 8Hz, 1H), 7.22-7.21 (d, J = 4Hz, 1H), 7.01-6.91 (m, 3H), 5.14 -5.09 (m, 1H), 4.27 (s, 1H), 4.01- 3.98 (d, J = 9.2Hz, 1H), 3.84-3.81 (d, J = 12.8Hz, 2H), 3.26 (s, 4H), 2.94-2.85 (m, 1H), 2.81-2.755 (m,	49

FIG. 3 Continued.


				2H), 2.54-2.48 (m, 6H), 2.22-2.21 (d, *J* = 6Hz, 2H), 2.08-2.03 (m, 1H), 1.81-1.78 (m, 3H), 1.28 (s, 6H), 1.21-1.12 (m, 2H), 1.05 (s, 6H)	
363		C	811.4	^{1}H NMR (300 MHz, CD_3OD): δ 7.79-7.68 (m, 2H), 7.37 (s, 1H),7.31-7.28 (m, 1H), 7.13-7.11 (m, 1H), 6.99-6.94 (m, 1H), 5.10-5.04 (m, 1H), 4.29 (s, 1H), 4.18-4.14 (m, 2H), 4.06 (s, 1H), 3.51-3.45 (m, 4H), 2.80-2.60 (m, 7H), 2.57-2.48 (m, 2H), 2.13-2.02 (m, 1H), 1.92-1.80 (m, 2H), 1.75-1.46 (m, 4H), 1.26 (s, 6H),1.19 (s, 6H);	9

FIG. 3 Continued.



364		C	844.32		
365		C	811.6		

FIG. 3 Continued.




366		D			
367		D			
368		D			

FIG. 3 Continued.



369		D			
370		D			

FIG. 3 Continued.



371		D			
372		D	877.4		

FIG. 3 Continued.



373		D	727.3		
374		D	784.2		

FIG. 3 Continued.



375		D	799.2
376		D	816.2

FIG. 3 Continued.



377		D	782.2
378		D	828.2

FIG. 3 Continued.




379		D	729.18		
380		D	766.27		
381		D	800.3		
382		D	817.2		

FIG. 3 Continued.



383		D	848.2	H-NMR (300 MHz, CDCl3) δ 8.04-7.95 (m, 3H), 7.84-7.77 (m, 2H), 7.60-7.55 (m, 1H), 7.34-7.33 (d, J=1.8 Hz, 1H), 7.21-7.09 (m, 3H), 4.98-4.93 (m, 1H), 4.14-4.10 (m, 2H), 3.91 (s, 2H), 3.49 (s, 2H), 2.94-2.72 (m, 3H), 2.67-2.56 (m, 5H), 2.25-2.13 (m, 2H), 1.92-1.88 (m, 2H), 1.76 (s, 2H), 1.68 (s, 6H);	122
384		D	763.5		

FIG. 3 Continued.




385		D	824.57		
386		D	816.5		
387		D	830.5		

FIG. 3 Continued.



388		D	838.57		
389		D	781.52		

FIG. 3 Continued.



390		D	722.4		
391		D	856.57		

FIG. 3 Continued.



392		D	767.54		
393		D	855.5		

FIG. 3 Continued.


394		D	736.49	
				

FIG. 3 Continued.



395		D	791.54	
396		D	809.6	

FIG. 3 Continued.



397		D	824.6		
398		D	926.7		

FIG. 4

Table 4. Exemplary Compounds



	Compound Structure	MW	Vcap DC50 (μM)	Dmax (%)	Chemical Name	Prepared by General Scheme
399		793.32	0.00235	60.15	N-((1s,4s)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide	1B
400		793.32	0.00066	78.6	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)nicotinamide	1B

FIG. 4 Continued.



401		792.33	0.000625	81.8	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide	2B
402		792.33	0.0019	58.7	N-((1s,4s)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide	2B (using cis aminocyclobutanol)

FIG. 4 Continued.




403		794.31	0.00075	81.15	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide	1B
404		794.31	0.00041	84.65	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrazine-2-carboxamide	1B
405		794.31	0.000165	85.3	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide	1B

FIG. 4 Continued.



406		812.3	0.000242	82.4	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyridazine-3-carboxamide	22B
407		812.3	0.00062	82.05	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide	1B

FIG. 4 Continued.




408		812.3	0.000285	84.05	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrazine-2-carboxamide	22B
409		778.31	0.00035	78.35	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)benzamide	5B
410		778.31	0.00097	68.4	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)piperazin-1-yl)benzamide	8B

FIG. 4 Continued.




411		780.28	0.000195	78.25	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)piperazin-1-yl)pyridazine-3-carboxamide	7B
412		780.28	0.000355	71.05	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(1-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperidin-4-yl)piperazin-1-yl)pyrimidine-5-carboxamide	6B
413		780.28	0.000335	62.85	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)piperidin-1-yl)pyridazine-3-carboxamide	5B

FIG. 4 Continued.




414		780.28	0. 00074	59.85	N-((1r,4S)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-((3S)-3-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)pyrrolidin-1-yl)pyrimidine-5-carboxamide	4B
415		741.2	0.0004	62.1	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyridazine-3-carboxamide	3B
416		739.23	0.000305	44.35	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)benzamide	3B

FIG. 4 Continued.



417		741.2	0.00062	55.62	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyrimidine-5-carboxamide	3B
418		755.23	0.00025	68.4	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrimidine-5-carboxamide	3B

FIG. 4 Continued.






419		806.36	0.0025	38.6	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)-4-methylcyclohexyl)-4-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide	1B, 24B
420		741.2	0.00017	75.4	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)pyrazine-2-carboxamide	14B

FIG. 4 Continued.

421		793.32	0.000575	84.6	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-((4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)picolinamide	1B
422		776.19			N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)ethyl)piperazin-1-yl)-5-fluoronicotinamide	23B
423		773.22			N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-6-fluoro-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrazine-2-carboxamide	3B



424		808.34			N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-2-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyrimidine-5-carboxamide	21B
425		806.36			N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-4-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)benzamide	21B

FIG. 4 Continued.



426		755.23	0.000295	79.6	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-5-(4-(3-((2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)propyl)piperazin-1-yl)pyrazine-2-carboxamide	3B
427		808.34	0. 00014	81.1	N-((1r,4r)-4-(3-chloro-4-cyanophenoxy)cyclohexyl)-6-(4-(2-(4-(2-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)ethyl)piperidin-1-yl)pyridazine-3-carboxamide	21B

FIG. 4 Continued.

FIG. 5

Table 5. Exemplary Compounds


Ex. No.	Compound Structure	MW	Vcap DC50 (μM)	Dmax (%)	
428		824.38	>1	2.1	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-6-(4-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)nicotinamide

FIG. 5 Continued.



429		812.37	>1	1.3	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)-1H-pyrazole-4-carboxamide
430		812.37	>1	8.5	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperidin-4-yl)-1H-pyrazole-3-carboxamide

FIG. 5 Continued.



431		784.31	>1	12.5	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(4-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)pyrrolidin-3-yl)-1H-pyrazole-4-carboxamide
432		798.34	>1	10.1	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(1-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)pyrrolidin-3-yl)-1H-pyrazole-4-carboxamide

FIG. 5 Continued.



433		813.35	>1	15.5	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-1-(2-(3-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propoxy)ethyl)-1H-pyrazole-4-carboxamide
434		824.38	>1	-1	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)picolinamide

FIG. 5 Continued.








435		857.93	>1	0	N-((1r,3r)-3-(4-cyano-3-(trifluoromethyl)phenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(4-(5-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)pentyl)piperazin-1-yl)picolinamide
436		780.32	>1	15.9	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-5-(3-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)propyl)picolinamide
437		619.72	>1	11.9	6-(4-(4-((2-(1-butyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)-N,N-dimethylpyridazine-3-carboxamide

FIG. 5 Continued.

438		811.34	>1	12. 3	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-(4-((2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)oxy)butyl)piperazin-1-yl)pyrimidine-5-carboxamide
439		834.42	0.508++	16.3	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-4-(4-((4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)benzamide
440		836.39	0.667++	31.67	N-((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)-2-(4-((4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)methyl)piperidin-1-yl)pyrimidine-5-carboxamide
441		908.45	0.00057 5	78.8	(3-(5-(4-(((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl methyl carbonate

WO 2018/071606 PCT/US2017/056234

FIG. 5 Continued.





442		922.48	0.00081	77.6	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl ethyl carbonate
443		936.5	0.000755	71.5	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl isopropyl carbonate
444		978.54	0.000635	79.5	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (tetrahydro-2H-pyran-4-yl) carbonate
445		978.55	1++	1.2	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-acetamidoethyl)carbamate

FIG. 5 Continued.



446		993.56	1++	13.55	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-(2-aminoacetamido)ethyl)carbamate
447		1007.59	1++	4.2	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-aminopropanamido)ethyl)carbamate

FIG. 5 Continued.



448		1035.64	1++	11.7	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-amino-3-methylbutanamido)ethyl)carbamate
449		1134.77	0.017	21.7	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2-((S)-2-((S)-2-amino-3-methylbutanamido)-3-methylbutanamido)ethyl)carbamate

FIG. 5 Continued.



450		1084.66	0.00034	77.4	(3-(5-(4-((1-(4-(((1r,3r)-3-(3-chloro-4-cyanophenoxy)-2,2,4,4-tetramethylcyclobutyl)carbamoyl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)-1,3-dioxoisoindolin-2-yl)-2,6-dioxopiperidin-1-yl)methyl (2,5,8,11-tetraoxatridecan-13-yl) carbonate
451		798.33	1++	13.4	2-chloro-4-(3-(3-fluoro-4-(5-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)phenyl)-4,4-dimethyl-5-oxo-2-thioxoimidazolidin-1-yl)benzonitrile

FIG. 5 Continued.


452		810.34	1++	17.1	2-chloro-4-(5-(3-fluoro-4-(5-(4-(2-(1-methyl-2,6-dioxopiperidin-3-yl)-1,3-dioxoisoindolin-5-yl)piperazin-1-yl)pentyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)benzonitrile

FIG. 6

Table 6. Exemplary Compounds


Ex. No.	Compound Structure	Vcap DC50	Dmax	m/z+	1H NMR
453		B	B	837.3	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.57 (s, 6H), 2.09-2.14 (m, 1H), 2.70-2.90 (m, 3H), 3.47 (t, *J* = 5.2 Hz, 2H), 3.69-3.74 (m, 10H), 3.88 (t, *J* = 4.6 Hz, 2H), 4.17 (t, *J* = 4.6 Hz, 2H), 4.88-4.93 (m, 1H), 6.50 (br, 1H), 6.92 (d, *J* = 8.8 Hz, 1H), 7.03-7.11 (m, 3H), 7.19 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97 (d, *J* = 8.4 Hz, 2H), 8.06 (s, 1H).

FIG. 6 Continued.



454		D	B	749.3	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.56 (s, 6H), 2.10-2.14 (m, 1H), 2.72-2.90 (m, 3H), 3.50 (t, *J* = 5.4 Hz, 2H), 3.81 (t, *J* = 5.2 Hz, 2H), 3.90 (t, *J* = 4.4 Hz, 2H), 4.20 (t, *J* = 4.2 Hz, 2H), 4.89-4.94 (m, 1H), 6.52 (br, 1H), 6.93 (d, *J* = 8.4 Hz, 1H), 7.05-7.12 (m, 3H), 7.19 (d, *J* = 8.4 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.96-7.99 (m, 2H), 8.10 (br, 1H).
455		C	B	925.3	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.57 (s, 9H), 2.07-2.11 (m, 1H), 2.70-2.82 (m, 3H), 3.44-3.49 (m, 2H), 3.66-3.75 (m, 18H), 3.87-3.89 (m, 2H), 4.16-4.18 (m, 2H), 4.84-4.88 (m, 1H), 6.91 (d, *J* = 8.4 Hz, 1H), 7.03-7.11 (m, 3H), 7.20 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97 (d, *J* = 8.0 Hz, 2H), 8.38 (s, 1H).

FIG. 6 Continued.



456		D	B	793.3	^{1}HNMR (400 MHz, CD_3OD):δ 1.43 (s, 6H), 1.93-1.98(m, 1H), 2.57-2.73 (m, 3H), 3.40 (t, J = 5.2 Hz, 2H), 3.59-3.65 (m, 6H), 3.78 (t, J = 4.4 Hz, 2H), 4.06(t, J = 4.4 Hz, 2H), 4.91-4.95 (m, 1H), 6.93-7.00 (m, 4H), 7.15 (d, J = 8.8 Hz, 2H), 7.41-7.45 (m, 1H), 7.87-7.90 (m, 1H), 8.04-8.06(m, 2H).
457		C	B	807.4	^{1}HNMR (400 MHz, $CDCl_3$): δ 1.56 (s, 6H), 2.08-2.10 (m, 3H), 2.70-2.90 (m, 3H), 3.46 (t, *J* = 5.2 Hz, 2H), 3.65-3.73 (m, 8H), 4.11 (t, *J* = 6 Hz, 2H), 4.88-4.92 (m, 1H), 6.92 (d, *J* = 8.8 Hz, 1H), 7.03 (d, *J* = 8.8 Hz, 2H), 7.10 (d, *J* = 7.2 Hz, 2H), 7.18 (d, *J* = 8.8 Hz, 2H), 7.49 (t, *J* = 7.6 Hz, 1H), 7.84 (d, *J* = 8.0 Hz, 1H), 7.97-7.99 (m, 3H).

FIG. 6 Continued.




458		C	B	865.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.57 (s, 6H), 1.76-1.79 (m, 2H), 1.87-1.90 (m, 2H),2.10-2.14 (m, 1H), 2.71-2.91 (m, 3H), 3.47-3.74 (m, 14H), 4.02 (t, *J* = 6.2 Hz, 2H), 4.88-4.93 (m, 1H), 6.48-6.49 (m, 1H), 6.91-7.02 (m, 3H), 7.10-7.20 (m, 3H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.86 (m, 1H), 7.97-8.02 (m, 3H).
459		B	A	851.3	^{1}HNMR (400 MHz, CD_3OD): δ 1.57 (s, 6H), 2.07-2.11 (m, 3H), 2.70-2.90 (m, 3H), 3.46-3.72 (m, 14H), 4.10 (t, *J* = 6.2 Hz, 2H), 4.88-4.92 (m, 1H), 6.48-6.49 (m, 1H), 6.91-7.26 (m, 6H), 7.49 (t, *J* = 7.8 Hz, 1H), 7.83-7.85(m, 1H), 7.97-8.02 (m, 3H).
460		C	B	837.2	

FIG. 6 Continued.




461		B	B	837.2	
462		B	B	821.3	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.57 (s, 6H), 1.86-1.89 (m, 2H), 2.04-2.09 (m, 3H), 2.71-2.90 (m, 3H), 3.43 (s, 2H), 3.55-3.63 (m, 8H), 4.10 (t, J = 6.0 Hz, 2H), 4.89-4.93 (m, 1H), 6.91 (d, J=8.8Hz, 1H), 7.02-7.05 (m, 2H), 7.10 (d, J=6.8Hz, 1H), 7.17-7.20 (m, 2H), 7.47-7.51 (m, 1H), 7.83-7.86 (m, 1H), 7.97-7.99(m, 2H), 8.07 (s, 1H).
463		D	B	865.1	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.42 (s, 6H), 1.85 (s, 3H), 1.92-1.96 (m, 3H), 2.44-2.49 (m, 1H), 2.56-2.64 (m, 2H), 3.34 (d, *J* = 5.2 Hz, 2H), 3.50-3.59 (m, 12H), 4.01 (d, *J* = 6.2 Hz, 2H) ,6.86-6.94 (m, 4H), 7.13-7.16 (m, 2H), 7.38-7.42 (m, 1H), 7.86-7.89 (m,

FIG. 6 Continued.



					1H), 8.03-8.05 (m, 2H).
464		B	A		
465		B	A	798.3	

FIG. 6 Continued.



466		D	B		1HNMR (400 MHz, DMSOd6) :δ 10.97 (s, 1H), 8.38 (d, *J* = 8.0 Hz , 1H), 8.29 (s, 1H), 8.07 (d, *J* = 8.4 Hz, 1H), 7.46 (t, *J* = 2.6 Hz, 1H), 7.24-7.33 (m, 4H), 7.08 (d, *J* = 8.4 Hz, 2H), 5.09-5.12 (m, 1H), 4.05-4.39 (m, 9H), 3.77 (s, 2H), 3.51-3.59 (m, 10H), 2.80-3.00 (m, 1H), 2.30-2.70 (m, 2H), 1.90-2.10 (m, 3H), 1.24 (s, 6H).
467		D	B	753.7	

FIG. 6 Continued.



468		B	A	825.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.31 (s, 6H), 1.96-2.12 (m, 4H), 2.71-2.81 (m, 8H), 3.48 (t, *J* = 6.0 Hz, 2H), 3.64-3.70 (m, 8H), 4.16 (s, 1H), 4.30 (s, 1H), 5.03-5.09 (m, 1H), 6.82 (d, *J* = 8.8 Hz, 1H), 6.99-7.10 (m, 3H), 7.14-7.15 (m, 1H), 7.58 (t, *J* = 7.8 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.95-7.98 (m, 1H), 8.60-8.61 (m, 1H).
469		D	B		

FIG. 6 Continued.



470		B	A	815.3	^{1}H NMR (400 MHz, CD_3OD): δ 1.23 (s, 6H), 1.29 (s, 6H), 1.49-1.56 (m, 2H), 1.66-1.73 (m, 4H), 1.92-1.95 (m, 2H), 2.06-2.12 (m, 1H), 2.69-2.80 (m, 3H), 3.23 (t, *J* = 7.0 Hz, 2H), 3.44 (t, *J* = 6.4 Hz, 2H), 3.50 (t, *J* = 6.2 Hz, 2H), 3.59 (t, *J* = 5.6 Hz, 2H), 4.13-4.15 (m, 1H), 4.30 (s, 1H), 5.03-5.07 (m, 1H), 6.74 (t, *J* = 8.4 Hz, 1H), 6.99-7.08 (m, 3H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.44-7.57 (m, 3H), 7.74 (d, *J* = 8.8 Hz, 1H).
471		D	B	754.3	

FIG. 6 Continued.




472		D	B	767.4	
473		D	B	752.7	
474		A	A	798.3	

FIG. 6 Continued.




475		B	A	843.4	
476		B	A	825.3	
477		B	A	843.3	

FIG. 6 Continued.




478		B	A	841.1	
479		C	A	856.9	
480		B	B	921.5	

FIG. 6 Continued.




481		B	A	843.7	
482		A	A	809.8	
483		B	A	809.8	

FIG. 6 Continued.





484		B	A	840.3	
485		A	A	796.7	
486		A	A	823.4	
487		A	A	796.7	

FIG. 6 Continued.




488		D	B	797.7	
489		B	A	840.7	
490		A	A	794.7	

FIG. 6 Continued.




491		A	A	797.7	
492		B	A	807.4	
493		A	A	810.3	

FIG. 6 Continued.




494		D	B	740.3	
495		D	B	739.2	
496		A	A	793.5	

FIG. 6 Continued.



497		A	A	779.5	
498		A	A	825.4	

FIG. 6 Continued.




499		D	B	697.3	
500		D	B	711.3	
501		D	B	727.3	

FIG. 6 Continued.





502		D	B	741.3	
503		D	B	755.3	
504		D	B	769.3	
505		D	B	780.6	

FIG. 6 Continued.




506		D	B	807.8	
507		D	B	683.3	
508		B	A	767.7	

FIG. 6 Continued.





509		D	B	694.2	
510		B	A	841.3	
511		D	B	810.3	
512		D	B	722.2	

FIG. 6 Continued.




513		D	B	736.3	
514		A	A	824.3	
515		A	A	824.3	

FIG. 6 Continued.




516		A	A	838.4	
517		D	B	838.4	
518		A	A	837.4	

FIG. 6 Continued.




519		D	B	837.4	
520		D	B	793.4	
521		A	A	765.3	

FIG. 6 Continued.




522		A	A	827.4	
523		B	A	827.4	
524		A	A	828.3	

FIG. 6 Continued.




525		B	A	842.3	
526		B	A	827.4	
527		A	A	827.4	

FIG. 6 Continued.











528		A	A		




FIG. 7



Table 7. Exemplary Compounds




Ex. #	Chemical Structure	m/z observed	DC50 category	1H NMR	General Scheme
529		832.610	A		
530		834.2	A	^{1}H NMR (300 MHz, *d6*-DMSO) δ 1.13 (s, 6H), 1.22 (s, 6H), 1.59-1.74 (m, 4H), 1.90-2.10 (m, 1H), 2.61-2.69 (m, 2H), 2.89-2.94 (m, 2H), 3.08-3.15 (m, 5H), 3.87-3.90 (m, 1H), 4.04-4.12 (m, 5H), 4.15-4.21 (m, 1H), 4.32 (s, 1H), 5.06-5.10 (m, 1H), 5.15-5.30 (m, 1H), 6.47-6.49 (m, 2H), 6.99-7.02 (m, 1H), 7.20-7.21 (m, 1H), 7.24-7.27 (m, 1H), 7.33 (s, 1H), 7.45-7.48 (m, 1H), 7.65-7.68 (m, 1H), 7.74-7.77 (m, 2H), 7.89-7.92 (m, 1H), 11.07 (s, 1H)	18
531		801.500	B	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.15 (s, 1H), 8.29 (d, J = 9.4 Hz, 1H), 7.92 (dd, J = 9.0, 4.3 Hz, 3H), 7.77 (d, J = 7.1 Hz, 1H), 7.49 (s, 1H), 7.26 (d, J = 2.4 Hz, 1H), 7.05 (dd, J = 8.6, 2.5 Hz, 1H), 5.15 (dd, J = 12.9, 5.4 Hz, 1H), 4.48 -4.41 (s, 3H), 4.03 (d, J = 9.2 Hz, 1H), 3.35 (s, 9H), 2.94 – 2.53 (m, 4H), 2.35-2.08 (s, 3H), 1.24 (s, 6H), 1.16 (s, 6H).	9


532		801.500	B	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.14 (s, 1H), 8.64 (d, J = 1.2 Hz, 1H), 8.36 (s, 1H), 7.96 – 7.80 (m, 3H), 7.76 (d, J = 7.1 Hz, 1H), 7.70 – 7.51 (m, 1H), 7.26 (d, J = 2.4 Hz, 1H), 7.04 (dd, J = 8.8, 2.5 Hz, 1H), 5.14 (dd, J = 12.9, 5.3 Hz, 1H), 4.48 – 4.33 (m, 4H), 3.97 (d, J = 9.0 Hz, 1H), 3.74 (s, 4H), 2.98 – 2.81 (m, 1H), 2.68 – 2.52 (m, 6H), 2.13 – 2.01 (m, 1H), 2.03 (s, 3H), 1.17 (d, J = 18.6 Hz, 12H).	9
533		773.470	A	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.12 (s, 1H), 8.76 (s, 2H), 8.15 (d, J = 7.2 Hz, 1H), 7.86 (d, J = 8.8 Hz, 2H), 7.74 (d, J = 7.2 Hz, 1H), 7.37 (d, J = 2 Hz, 1H), 7.14-7.12 (m, 1H), 5.12 (m, 1H), 4.55 (s, 1H), 4.37-4.34 (m, 2H), 3.82 (s, 6H), 2.94-2.85 (m, 1H), 2.64-2.61(m, 2H), 2.56-2.44 (m, 5H), 2.11-1.91 (m, 7H), 1.57-1.44 (m, 4H)	3 B
534		826.540	A	^{1}H NMR (300 MHz, DMSO-*d6*) 11.12 (s, 1H), 8.74 (s, 2H), 8.12-8.10 (d, *J*= 7.5Hz, 1H), 7.88-7.85 (d, *J*= 8.7Hz, 1H), 7.75-7.71 (d, *J*= 11.4Hz, 1H), 7.47-7.39 (m, 2H), 7.16-7.13 (m, 1H), 5.08-5.04 (m, 1H), 4.74-4.55 (m, 3H), 3.86-3.80 (m, 1H), 3.55-3.51 (m, 6H), 2.96-2.92 (m, 2H), 2.59-2.46 (m, 7H), 2.05-1.80 (m, 5H), 1.79-1.61 (m, 4H), 1.56-1.42 (m, 5H), 1.21-1.14 (m, 2H);	9 B
535		957.650	C	^{1}H-NMR (300 MHz, CD_3OD) δ 8.6（s, 1H), 8.08- 8.01 (m, 1H), 7.76- 7.69 (m, 1H), 7.59- 7.36 (m, 2H), 7.14 (s, 1H), 7.03- 6.96 (m, 1H), 635 (s, 1H), 5.12 - 5.04 (m, 1H), 4.60 (s, 1H), 4.50- 4.39 (m, 4H), 4.30- 4.27 (m, 1H), 4.18- 4.10 (m, 1H), 3.96- 3.90 (m, 2H), 3.73- 3.52 (m, 4H), 3.09- 2.66 (m, 10H), 2.37- 2.31(m, 2H), 2.15- 1.88 (m, 5H), 1.29 - 1.20 (m,	60




				18H)	
536		751.500	A	1H NMR (300 MHz, DMSO-d6) 11.05 (s, 1H), 8.21-8.18 (m, 1H), 7.85-7.82 (m, 1H), 7.76-7.73 (m, 2H), 7.63-7.60 (m, 1H), 7.37-7.30 (m, 4H), 7.22-7.10 (m, 2H), 5.07-5.01 (m, 1H), 4.52-4.45 (m, 1H), 3.80-3.70 (m, 1H), 3.47-3.43 (m, 2H), 3.22-3.16 (m, 2H), 3.00-2.90 (m, 2H), 2.85-2.52 (m, 7H), 2.25-1.87 (m, 5H), 1.54-1.45 (m, 4H), 1.21-0.97 (m, 6H).	10 B
537		811.600	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.05 (s, 1H), 8.03 (d, J = 8.8 Hz, 1H), 7.87-7.59 (m, 4H), 7.45 (d, J = 9.1 Hz, 1H), 7.36-7.16 (m, 3H), 6.92 (d, J = 8.9 Hz, 2H), 5.06-5.03 (m, 1H), 4.33 (s, 1H), 4.03 (d, J = 9.1 Hz, 1H), 3.83 (d, J = 12.6 Hz, 2H), 3.41 (s, 4H), 2.76 (s, 3H), 2.90-2.72 (m, 3H), 2.70-2.65 (m, 2H), 2.28-2.20 (s, 2H), 2.10-1.90 (m, 1H),1.86-1.70 (m, 3H), 1.19 (s, 9H), 1.08 (s, 6H	49
538		800.620	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.06 (s, 1H), 7.75-7.57 (m, 5H), 7.46 (d, J = 9.2 Hz, 1H), 7.36-7.15 (m, 2H), 6.93 (d, J = 8.6 Hz, 2H), 6.80 (d, J = 8.8 Hz, 1H), 5.04 (dd, J = 12.7, 5.3 Hz, 1H), 4.22 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.83 (d, J = 12.5 Hz, 2H), 3.41 (s, 4H), 2.81 (dt, J = 23.9, 12.5 Hz, 3H), 2.55 (d, J = 17.4 Hz, 2H), 2.47 (p, J = 1.8 Hz, 6H), 2.22 (s, 5H), 1.98 (d, J = 11.9 Hz, 1H), 1.78 (d, J = 12.0 Hz, 3H), 1.20 (s, 6H), 1.10 (s, 6H).	49




539		811.600	D	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.08 (s, 1H), 8.43 (d, J = 2.1 Hz, 1H), 8.13 (dd, J = 8.9, 2.2 Hz, 1H), 7.75 (d, J = 8.6 Hz, 3H), 7.70 (d, J = 8.5 Hz, 1H), 7.52 (d, J = 9.0 Hz, 1H), 7.36 (s, 1H), 7.28 (d, J = 7.9 Hz, 1H), 7.17 (d, J = 9.0 Hz, 1H), 6.98 (d, J = 8.7 Hz, 2H), 5.09 (dd, J = 12.6, 5.4 Hz, 1H), 4.45 (s, 1H), 4.11 (d, J = 9.0 Hz, 1H), 3.88 (d, J = 12.3 Hz, 2H), 3.46 (brs, 4H), 2.95-2.75 (m, 3H), 2.66-2.46 (m, 8H), 2.23 (d, J = 6.4 Hz, 2H), 2.02 (brs, 1H), 1.83 (d, J = 12.1 Hz, 3H), 1.26 (s, 6H), 1.17 (s, 6H).	49
540		814.640	D	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.10 (s, 1H), 7.71 (dd, J = 11.8, 8.6 Hz, 3H), 7.6-7.52 (m, 3H), 7.35 (d, J = 2.1 Hz, 1H), 7.27 (d, J = 8.9 Hz, 1H), 6.96 (d, J = 8.6 Hz, 2H), 5.09 (dd, J = 12.9, 5.4 Hz, 1H), 4.33 (s, 1H), 3.99 (d, J = 9.9 Hz, 1H), 3.86 (d, J = 12.6 Hz, 2H), 3.44 (d, J = 5.8 Hz, 4H), 2.95-2.75 (m, 3H), 2.67-2.53 (m, 1H), 2.50 (s, 5H), 2.32 (s, 6H), 2.20 (d, J = 6.6 Hz, 2H), 2.10-2.00 (m, 1H), 1.81 (d, J = 12.3 Hz, 3H), 1.35 (s, 6H), 1.20 (q, J = 12.3, 10.9 Hz, 2H), 1.06 (s, 6H)	49
541		816.620	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.07 (s, 1H), 7.77-7.58 (m, 4H), 7.48 (d, J = 9.1 Hz, 1H), 7.33 (s, 1H), 7.24 (d, J = 8.7 Hz, 1H), 6.94 (d, J = 8.7 Hz, 2H), 6.62 (d, J = 2.1 Hz, 1H), 6.52 (d, J = 8.7 Hz, 1H), 5.05 (dd, J = 12.6, 5.3 Hz, 1H), 4.25 (s, 1H), 4.03 (d, J = 9.2 Hz, 1H), 3.89 (s, 5H), 3.43 (s, 4H), 2.77 (q, J = 13.8, 11.8 Hz, 3H), 2.53 (s, 5H), 2.19 (d, J = 6.4 Hz, 2H), 2.01 (s, 1H), 1.79 (d, J = 12.2 Hz, 3H), 1.16 (d, J = 23.6 Hz, 15H)	49




542		817.620	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.13 (s, 1H), 8.60 (d, *J* = 1.2 Hz, 1H), 8.27 (s, 1H), 8.09 (d, *J* = 8.4 Hz, 1H), 7.85 (d, *J* = 8.0 Hz, 1H), 7.67 (d, *J* = 8.6 Hz, 1H), 7.27 (d, *J* = 8.3 Hz, 2H), 7.05 (d, *J* = 2.3 Hz, 1H), 7.00-6.90 (m, 1H), 5.13 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.91-4.85 (m, 1H), 4.57-4.40 (m, 3H), 3.84-3.73 (m, 1H), 3.68 (t, *J* = 8.1 Hz, 1H), 3.00-2.94 (m, 4H), 2.70-2.61 (m, 1H), 2.60-2.30 (m, 5H), 2.29-2.10 (m, 4H), 2.08-2.01 (m, 3H), 2.00-1.81 (m, 12.9 Hz, 4H),1.76-1.45 (m, 6H), 1.07 (d, *J* = 12.4 Hz, 2H), 0.94 (d, *J* = 6.5 Hz, 6H)	72
543		755.440	B	H-NMR (300 MHz, DMSO-*d6*) δ 11.11 (s, 1H), 7.97 (d, *J*=7.5 Hz, 1H), 7.85-7.70 (m, 6H), 7.37 (d, *J*=2.4 Hz, 1H), 7.14-7.10 (m, 1H), 6.92 (d, *J*=8.7 Hz, 2H), 5.15-5.09 (m, 1H), 4.51 (s, 1H), 3.78 (s, 1H), 3.38-3.36 (m, 2H), 3.22 (s, 4H), 2.87-2.57 (m, 9H), 2.10-2.00 (m, 3H), 1.88-1.86 (m, 2H), 1.53-1.45 (m, 4H)	11 B





544		829.540	B	1H NMR (300 MHz, CDCL3) δ 8.09 (s, 1H), 7.65 – 7.48 (m, 3H), 6.99 (d, J = 2.4 Hz, 1H), 6.83 (dd, J = 8.7, 2.4 Hz, 1H), 5.71 (d, J = 8.5 Hz, 1H), 4.98 (dd, J = 12.2, 5.3 Hz, 1H), 4.31 (t, J = 6.3 Hz, 2H), 4.20 (d, J = 8.5 Hz, 1H), 4.05 (s, 1H), 3.92 (s, 4H), 3.01 – 2.69 (m, 3H), 2.63 (s, 2H), 2.55 (s, 4H), 2.39 (s, 6H), 2.19 (d, J = 14.7 Hz, 4H), 1.26 (s, 12H	62
545		886.590	B	1H NMR (300 MHz, $CDCL_3$): δ 8.05 (s, 1H), 7.60 (d, *J* = 8.7 Hz, 1H), 7.23 (d, *J* = 6.0 Hz, 1H), 6.99 (d, *J* = 2.4 Hz, 1H), 6.83 (dd, *J* = 8.7, 2.4 Hz, 1H), 5.71 (d, *J* = 8.6 Hz, 1H), 5.02 – 4.84 (m, 3H), 4.20 (d, *J* = 8.4 Hz, 1H), 4.05 (s, 1H), 3.39 (s, 3H), 3.00 – 2.73 (m, 5H), 2.64 (s, 4H), 2.38 (s, 5H), 2.31 (s, 2H), 2.23 – 2.12 (m, 1H), 1.89 (d, *J* = 13.3 Hz, 3H), 1.26 (s, 15H), 1.05 - 0.87 (m, 1H);	63
546		789.530	B	^{1}H NMR (300 MHz, DMSO-d_6): *δ* 11.86 (d, *J* = 15.5 Hz, 1H), 11.09 (s, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 7.70 (d, *J* = 8.5 Hz, 1H), 7.41-6.80 (m, 7H), 5.09 (dd, *J* = 12.7, 5.4 Hz, 1H), 4.64 (brs, 1H), 3.58-3.46 (m, 6H), 2.94-2.82 (m, 2H), 2.68-2.55 (m, 4H), 2.55-2.47 (m, 4H), 2.26-1.99 (m, 7H), 1.87-1.51 (m, 7H), 1.31 (q, *J* = 12.4, 11.6 Hz, 2H).	12 B




547		807.550	A	^{1}HNMR (300 MHz, DMSO-d6) 11.07(s, 1H), 8.56 (s, 1H), 8.00-7.98 (m, 1H), 7.98-7.88(m, 2H), 7.85-7.67(m, 1H), 7.37-7.36 (m, 3H), 7.24-7.13 (m, 1H), 6.81-6.78 (m, 1H), 5.08-5.02 (m, 1H), 4.52-4.39 (m, 1H), 4.37-4.31 (m, 2H), 3.80 (s, 1H), 3.77-3.41 (m, 4H), 2.86-2.82 (m, 3H), 2.79(m, 2H), 2.58-2.56 (m, 4H), 2.52-2.47 (m, 2H), 2.32-2.08 (m, 3H), 2.00-1.97 (m, 2H), 1.86-1.74 (m, 2H), 1.71-1.48 (m, 7H), 1.37-1.09 (m, 2H)	9 B
548		826.540	A	1H NMR (300 MHz, DMSO-d6) δ 11.09 (s, 1H), 8.55 (d, J = 8.2 Hz, 1H), 7.74 (td, J = 13.7, 11.7, 7.4 Hz, 3H), 7.42 (d, J = 7.4 Hz, 1H), 7.26 (dd, J = 23.7, 9.3 Hz, 2H), 5.07 (dd, J = 12.8, 5.3 Hz, 1H), 4.45 (d, J = 13.5 Hz, 3H), 3.86 (s, 1H), 3.22 (s, 5H), 3.03-2.95 (m, 2H), 2.87-2.79 (m, 1H), 2.59-2.48 (m, 4H), 2.20-2.18 (m, 5H), 2.10 – 1.95 (m, 3H), 1.90-1.78 (m, 5H), 1.58 (td, J = 21.0, 18.4, 11.8 Hz, 4H), 1.20 (s, 1H), 1.09 (q, J = 11.7 Hz, 2H).	1 B




549		792.570	A	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.11 (s, 1H), 8.58 (d, *J* = 8.2 Hz, 1H), 7.81 (d, *J* = 9.5 Hz, 1H), 7.73 (d, *J* = 11.4 Hz, 1H), 7.66 (d, *J* = 8.6 Hz, 1H), 7.46 (d, *J* = 7.4 Hz, 1H), 7.34 (d, *J* = 9.6 Hz, 1H), 7.06 (d, *J* = 2.5 Hz, 1H), 6.95 (dd, *J* = 8.8, 2.5 Hz, 1H), 5.11 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.47 (t, *J* = 13.0 Hz, 3H), 3.87 (d, *J* = 10.3 Hz, 1H), 3.26 (d, *J* = 5.6 Hz, 4H), 3.03 (t, *J* = 12.5 Hz, 2H), 2.96 – 2.84 (m, 1H), 2.60 (d, *J* = 18.0 Hz, 2H), 2.55 (s, 4H), 2.44 (s, 3H), 2.23 (d, *J* = 7.0 Hz, 2H), 2.15 – 2.00 (m, 3H), 1.95 – 1.84 (m, 4H), 1.83 (s, 1H), 1.57 (dq, *J* = 52.6, 12.2, 11.7 Hz, 4H), 1.14 (q, *J* = 11.5 Hz, 2H);	1 B
550		800.620	A		49
551		801.620	B	^{1}H NMR (400 MHz, DMSO-*d6*): δ 11.09 (s, 1H), 8.07-8.05 (d, *J* = 8.8Hz, 1H), 7.77-7.74 (m, 3H), 7.70-7.68 (d, *J* = 8.4Hz, 1H), 7.56-7.53 (d, *J* = 9.2Hz, 1H), 7.35 (s, 1H), 7.28-7.25 (d, *J* = 8.8Hz, 1H), 6.97 -6.95 (d, *J* = 8.8Hz, 2H), 6.87-6.85 (d, *J* = 8.8Hz, 1H), 5.10-5.05 (m, 1H), 4.75 (s, 1H), 4.04-4.02 (d, *J* = 9.2Hz, 1H), 3.88-3.85 (m, 2H), 3.45 (s, 4H), 2.93-2.76 (m, 3H), 2.61-2.51 (m, 4H), 2.50 (m, 6H), 2.22-2.21(d, *J* = 6.4Hz, 2H), 2.08-1.98 (m, 1H), 1.88-1.68 (m, 3H), 1.21 (s, 6H), 1.12 (s, 6H);	49





552		787.600	D	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.06 (s, 1H), 8.64 (s, 1H), 8.13 (d, *J* = 8.4 Hz, 1H), 7.73-7.65 (m, 3H), 7.55-7.45 (m, 1H), 7.40 (s, 1H), 7.25-7.15 (m, 1H), 7.10-7.00 (m, 1H), 6.95-6.91 (m, 2H), 5.07-5.01 (m, 1H), 4.76 (s, 1H), 4.02 (d, *J* = 9 Hz, 1H), 3.85 (d, *J* = 11.7 Hz, 2H), 3.41 (s, 4H), 2.90-2.72 (m, 3H), 2.58-2.47 (m, 4H), 2.19-2.01(m, 2H), 2.00-1.90 (m, 1H),1.89-1.76 (m, 3H), 1.36-0.98(m, 15H);	49
553		813.600	A	^{1}H NMR (400 MHz, *d6*-DMSO): δ 11.09 (s, 1H), 8.75 (s, 2H), 8.07-8.05 (d, *J*=8.8Hz, 1H), 7.72-7.65 (m, 3H), 7.36-7.34 (m, 2H), 7.28-7.25 (d, *J*=9.2Hz, 1 H), 5.09-5.05 (m, 1H), 4.77-4.73 (m, 2H), 4.34 (s, 1H), 4.06-4.04 (m, 1H), 3.45 (m, 4H), 3.02-2.85 (m, 3H), 2.61-2.54 (m, 2H), 2.51 (m, 4H), 2.22-2.20 (m, 2H), 2.03-2.01 (m, 1H), 1.92 (s, 1H), 1.85-1.82 (m, 2H), 1.23 (s, 6H), 1.10-1.07 (m, 8H);	49
554		803.610	A	^{1}H NMR (400 MHz, d-DMSO):δ11.051(s,1H),8.740(s,2H),8.055-8.027(d,J=11.2Hz,2H),7.731-7.647(m,2H),7.327(s,1H),7.255-7.226(d,J=11.6Hz,1H),6.854-6.825(d,J=11.6Hz,1H),5.082-5.064(m,1H),4.747-4.692(m,3H),4.002-3.971(d,J=12.4Hz,1H),3.427(s,4H),3.305-2.806(m,3H),2.589-2.545(m,9H),2.247-2.173(m,2H),2.052-1.792(m,4H),1.178(s,6H),1.087-1.17(m,8H)	49



555		834.560	A		
556		801.620	C	1H NMR (300 MHz, DMSO-d6,ppm) ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.48 (d, J = 2.3 Hz, 1H), 8.01 (dd, J = 2.3, 1.1 Hz, 1H), 7.71 (dd, J = 17.4, 8.5 Hz, 3H), 7.51 (d, J = 9.3 Hz, 1H), 7.38(m, 1H) 7.21 (m, 1H), 6.95 (d, J = 9.0 Hz, 2H), 5.07 (dd, J = 12.8, 5.4 Hz, 1H), 4.82 (s, 1H), 4.05 (d, J = 9.1 Hz, 1H), 3.86 (d, J = 12.5 Hz, 2H),3.35(m, 8H) ,2.98-2.72 (m, 3H), 2.59 (m, 3H), 2.25 (m, 5H), 2.08-1.97 (m, 1H), 1.81 (d, J = 11.8 Hz, 3H), 1.16 (d, J = 23.5 Hz, 14H)	49
557		788.595	D	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.09 (s, 1H), 9.11 (s, 2H), 7.76 (d, J = 8.7 Hz, 2H), 7.70 (d, J = 8.5 Hz, 1H), 7.57 (d, J = 9.2 Hz, 1H), 7.35 (d, J = 2.1 Hz, 1H), 7.31-7.24 (m, 1H), 6.97 (d, J = 8.8 Hz, 2H), 5.09 (dd, J = 12.9, 5.4 Hz, 1H), 4.79 (s, 1H), 4.07 (d, J = 9.1 Hz, 1H), 3.87 (d, J = 12.4 Hz, 2H), 3.46 (s, 4H), 3.32 (s, 3H), 2.85 (dt, J = 37.2, 12.7 Hz, 3H), 2.64-2.53 (m, 1H), 2.22 (d, J = 6.6 Hz, 2H), 2.03 (d, J = 11.4 Hz, 1H), 1.82 (d, J = 12.3 Hz, 3H), 1.18 (d, J = 26.1 Hz, 16H).	49




558		814.643	A	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.10 (s, 1H), 7.73 (dd, J = 16.0, 8.6 Hz, 3H), 7.49 (d, J = 9.2 Hz, 1H), 7.45 – 7.33 (m, 1H), 7.28 (d, J = 8.7 Hz, 1H), 6.97 (d, J = 8.7 Hz, 2H), 6.75 (s, 2H), 5.09 (dd, J = 12.6, 5.4 Hz, 1H), 4.24 (s, 1H), 4.05 (d, J = 9.1 Hz, 1H), 3.87 (d, J = 12.8 Hz, 2H), 3.47 (s, 6H), 2.83 (q, J = 14.0, 12.0 Hz, 3H), 2.60 (d, J = 16.4 Hz, 2H), 2.44 (s, 8H), 2.24 (s, 2H), 2.03 (d, J = 11.2 Hz, 1H), 1.83 (d, J = 12.0 Hz, 3H), 1.23 (s, 8H), 1.13 (s, 6H).	49
559		780.510	A	^{1}H NMR (400 MHz, DMSO) 11.08 (s, 1H), 8.59 (s, 1H), 8.28 (s, 1H), 8.10 (m, 1H), 7.87-7.85 (m, 1H), 7.68-7.66 (m, 1H), 7.38-7.33 (m, 2H), 7.28-7.22 (m, 1H), 7.14-7.11 (m, 1H), 5.09-5.01 (m, 1H), 4.51-4.45 (m, 3H), 3.90-3.80 (m, 1H), 3.42-3.40 (m, 4H), 3.05-2.80 (m, 3H), 2.64-2.49 (m, 7H), 2.10-1.88 (m, 7H), 1.62-1.38 (m, 6H).	5 B
560		780.510	A	^{1}H NMR (300 MHz, DMSO-*d6*) 11.09 (s, 1H), 8.78-8.72 (m, 2H), 8.29-8.22 (m, 1H), 7.89-7.86 (m, 1H), 7.73-7.70 (m, 1H), 7.40-7.14 (m, 4H), 5.12-5.06 (m, 1H), 4.80-4.56 (m, 3H), 3.81-3.70 (m, 1H), 3.03-2.63 (m, 11H), 2.30-1.92 (m, 8H), 1.55-1.24 (m, 8H);	5 B
561		834.560	A		





562		781.520	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.11 (s, 1H), 7.88 (dd, J=20.8, 8.5 Hz, 2H), 7.75 (d, J=8.6 Hz, 2H), 7.48 (d, J=2.3 Hz, 1H), 7.45 -7.32 (m, 2H), 7.21 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.5 Hz, 1H), 6.57 (d, J = 8.7 Hz, 2H), 5.13 (dd, J = 12.8, 5.4 Hz, 1H), 4.31 (d, J = 7.5 Hz, 3H), 4.06 (d, J = 9.1 Hz, 1H), 3.50 (dt, J = 30.2, 8.2 Hz, 2H), 3.27 (m, 2H), 3.20 - 3.07 (m, 1H), 2.91 (m, 3H), 2.60 (d, J = 19.3 Hz, 2H), 2.36 (s, 3H), 2.21 (s, 1H), 2.04 (s, 1H), 1.96 -1.83 (m, 1H), 1.22 (s, 6H), 1.13 (s, 6H)	46
563		807.540	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.12 (s, 1H), 7.85-7.82 (m, 1H), 7.78-7.75 (m, 1H), 7.75 (d, J = 8.6 Hz, 2H), 7.51 (d, J = 9.2 Hz, 1H), 7.42-7.28 (m, 2H), 7.21 (d, J = 2.4 Hz, 1H), 7.06-6.91 (m, 3H), 5.18-75.03 (m, 2H), 4.32 (s, 1H), 4.05 (d, J = 9.1 Hz, 1H), 3.25-3.43 (m, 4H), 2.99 - 2.80 (m, 1H), 2.68-2.62 (m, 2H), 2.56 (s, 6H), 2.11-1.95 (m, 4H), 1.70-1.51 (m, 2H), 1.22 (s, 6H), 1.13 (s, 6H).	64
564		801.620	B	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.10 (s, 1H), 8.25 (d, J = 2.7 Hz, 1H), 7.73 (dd, J = 17.0, 8.5 Hz, 3H), 7.50 (d, J = 9.1 Hz, 1H), 7.40 – 7.23 (m, 3H), 6.97 (d, J = 8.7 Hz, 2H), 5.09 (dd, J = 12.6, 5.4 Hz, 1H), 4.35 (s, 1H), 4.08 (d, J = 9.0 Hz, 1H), 3.88 (d, J = 12.5 Hz, 2H), 3.00 – 2.73 (m, 5H), 2.66 – 2.54 (m, 6H), 2.50 (s, 4H), 2.23 (d, J = 6.6 Hz, 2H), 2.03 (d, J = 11.5 Hz, 1H), 1.83 (d, J = 12.0 Hz, 3H), 1.20 (d, J = 31.9 Hz, 15H).	49




565		855.600	B	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.09 (s, 1H), 8.70 (d, J = 2.6 Hz, 1H), 7.86-7.65 (m, 4H), 7.52 (d, J = 9.3 Hz, 1H), 7.36-7.28 (m, 2H), 6.98 (d, J = 8.8 Hz, 2H), 5.12-5.08 (m, 1H), 4.60 (s, 1H), 4.11 (d, J = 9.1 Hz, 1H), 3.88 (d, J = 12.4 Hz, 2H), 3.47-3.43 (m, 4H), 2.93-2.70 (m, 3H), 2.63-2.56(m, 5H), 2.55-2.47 (m, 3H), 2.23 (d, J = 6.5 Hz, 2H), 2.11-1.96 (m, 1H), 1.83 (d, J = 12.2 Hz, 3H), 1.25 (s, 6H), 1.17 (s, 6H).	49
566		794.520	B	1H NMR (400 MHz, CDCl3): δ 8.85 (d, J = 1.3 Hz, 1H), 8.06 (s, 1H), 7.76 (m, 8.7 Hz, 2H), 7.67-7.62 (m, 1H), 7.58 (d, J = 8.7 Hz, 1H), 7.32 (d, J = 2.3 Hz, 1H), 7.10 (d, J = 8.5 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.83 (m, 1H), 4.96 (m, 1H), 4.36 (s, 2H), 4.15 (d, J = 8.8 Hz, 1H), 4.08 (s, 1H), 3.95 (s, 2H), 3.50 (s, 4H), 3.15 (s, 1H), 2.96-2.56 (m, 9H), 2.20-2.12 (m, 1H), 1.28 (s, 6H), 1.23 (s, 6H);	65
567		766.490	A	^{1}H NMR (400 MHz,$CDCl_3$): δ 8.85 (d, J = 1.4 Hz, 1H), 8.03 (s, 1H), 7.72 (d, J = 8.5 Hz, 1H), 7.63-7.54 (m, 2H), 7.40 (d, J = 8.2 Hz, 1H), 7.31 (d, J = 2.3 Hz, 1H), 7.08 (m, 1H), 7.01 (d, J = 2.4 Hz, 1H), 6.86 (m, 1H), 4.96 (m, 1H), 4.33 (m, 3H), 4.11-3.99 (m, 1H), 3.91 (m, 2H), 3.45 (s, 4H), 3.11 (s, 1H), 2.96-2.78 (m, 2H), 2.82-2.68 (m, 3H), 2.64 (m, 4H), 2.25-2.10 (m, 5H), 1.78-1.63 (m, 2H), 1.55-1.40 (m, 2H);	7, 16 B




568		806.590	A	^{1}H NMR (300 MHz, DMSO): δ10.68(s, 1H), 7.91-7.87 (m, 1H), 7.72-7.69 (m, 2H), 7.49-7.46(m, 2H), 7.19-7.18 (m, 1H), 7.05-6.89 (m, 5H), 5.28(s, 2H), 4.57-4.52(m, 1H), 4.29 (s, 1H), 4.04-4.01 (m, 1H), 3.86-3.82 (m, 2H), 3.31-3.26(m, 10H), 2.80-2.72 (m, 2H), 2.61-2.57 (m, 2H), 2.19-2.17(m, 2H), 2.10-1.87 (m, 2H), 1.82-1.67 (m, 3H), 1.19 (s, 6H), 1.10 (s, 6H);	13 B
569		776.450	A	H-NMR (300 MHz, DMSO) δ 8.37-8.33 (d, *J*=10.8Hz, 1H), 7.92-7.80 (m, 3H), 7.34 (s, 1H), 7.10-7.08 (d, *J*=8.1Hz, 1H), 6.89-6.84 (d, *J*=14.1Hz, 1H), 5.14-5.08 (m, 1H), 4.67-4.30 (m, 4H), 3.83-3.67 (m, 3H), 3.37-2.93 (m, 6H), 2.91-2.56 (m, 2H), 2.38 (s, 2H), 2.11-1.73 (m, 5H), 1.17 (s, 4H);	14 B
570		812.520	A	^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.10 (s, 1H), 8.60 (d, *J* = 8,4 Hz, 1H), 7.84 (m, 2H), 7.40–7.35 (m, 2H), 7.25 (s, 1H), 7.15 – 6.99 (m, 3H), 5.10 (m, 1H), 4.74 – 4.50 (m, 3H), 3.95 – 3.75 (m, 1H), 3.60 (s, 4H), 3.10 – 2.80 (m, 3H), 2.53 – 2.50 (m, 5H), 2.21 – 2.15 (m, 2H), 2.14 – 1.75 (m, 8H), 1.74 – 1.50 (m, 4H), 1.25 – 1.00 (m, 2H)	15 B
571		840.560	A	^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.10 (s, 1H), 8.75 (s, 2H), 7.90 (d, *J* = 8.7 Hz, 1H), 7.71 (d, *J* = 9.0 Hz, 1H), 7.24–7.21 (m, 2H), 7.07 – 6.99 (m, 2H), 5.10 (m, 1H), 4.74 (d, *J* = 13.2 Hz, 2H), 4.29 (s, 1H), 4.04 (d, *J* = 9.0 Hz, 1H), 3.50 (s, 4H), 3.00 – 2.95 (m, 3H), 2.53 – 2.50 (m, 5H), 2.21 – 2.19 (m, 2H), 2.08 – 1.81 (m, 5H), 1.21 – 1.04 (m, 14H)	66 and 49





572		848.650	B	^{1}H NMR (300 MHz, DMSO-*d6*):δ 11.12 (s, 1H), 8.22 (d, *J* = 9.6 Hz, 1H), 7.80 – 7.65 (m, 2H), 7.47 (d, *J* = 7.4 Hz, 1H), 6.95 (d, *J* = 2.4 Hz, 1H), 6.83 (dd, *J* = 8.6, 2.5 Hz, 1H), 5.12 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.74 (d, J = 12.6 Hz, 2H), 4.17 (s, 1H), 4.08 (d, J = 9.5 Hz, 1H),3.26 - 3.18 (m, 4H), 2.89 (q, *J* = 11.2 Hz, 3H), 2.56 (s, 5H), 2.46 (s, 3H), 2.24 (s, 8H), 2.10 – 1.99 (m, 1H), 1.81 (d, *J* = 13.2 Hz, 3H), 1.26 -1.17 (m, 7H), 1.12 (s, 6H), 1.06 - 0.94 (m,2H);	63
573		794.520	B	^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.08 (s, 1H), 8.23 (d, *J*=9.0 Hz, 1H), 7.92-7.83 (m, 2H), 7.69 (d, *J*=8.7 Hz, 1H), 7.35 (m, 1H), 7.29-7.25 (m, 2H), 7.05-7.01 (m, 1H), 6.89 (d, *J*=9.3 Hz, 1H), 5.10-5.05 (m, 1H), 4.46 (s, 1H), 4.26 (t, *J*=8.3 Hz, 2H), 4.00 (d, *J*=9.0 Hz, 1H), 3.86-3.81 (m, 2H), 3.46 (s, 4H), 3.10 (m, 1H), 2.95-2.83 (m, 1H), 2.69 (d, *J*=7.2 Hz, 2H), 2.62-2.56 (m, 6H), 2.08-2.00 (m, 1H), 1.22-1.14 (m, 12H);	65
574		766.490	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.11 (s, 1H), 8.75 (s, 2H), 8.17 (d, J = 7.3 Hz, 1H), 7.88 (d, J = 8.8 Hz, 1H), 7.70 (d, J = 8.4 Hz, 1H), 7.38-7.36 (m, 2H), 7.28 (d, J = 8.6 Hz, 1H), 7.18-7.16 (m, 1H), 5.12-5.09 (m, 1H), 4.56 (s, 1H), 4.22 (t, J = 8.6 Hz, 2H), 3.82-3.70 (m, 3H), 3.48-3.46 (m, 4H), 3.09-2.75 (m, 2H), 2.67 (d, J = 7.4 Hz, 3H), 2.56 (s, 3H), 2.54-2.52 (m, 2H), 2.20-1.68 (m, 5H), 1.51 (s, 4H);	7, 16 B



575		746.540	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.11 (s, 1H), 8.75 (d, J = 3.1 Hz, 2H), 8.16 (d, J = 7.4 Hz, 1H), 7.70-7.60 (m, 2H), 7.46-7.19 (m, 2H), 7.16-6.89 (m, 2H), 5.11-5.01 (m, 1H), 4.47 (brs, 1H), 4.22 (t, J = 8.6 Hz, 2H), 3.80-3.71 (m, 3H), 3.50-3.42 (m, 4H), 3.10-2.80 (m, 2H), 2.72-2.63 (m, 3H), 2.55-2.52 (m, 5H), 2.45-2.43 (m, 3H), 2.24-1.84 (m, 5H), 1.53-1.48 (m, 4H).	7, 16 B
576		812.520	B	^{1}H NMR (300 MHz, DMSO) δ 11.11 (s, 1H), 8.23 (d, *J*=9.0 Hz, 1H), 7.92-7.83 (m, 2H), 7.74 (d, *J*=11.4 Hz, 1H), 7.47 (d, *J*=7.2 Hz, 1H), 7.25 (d, *J*=2.4 Hz, 1H), 7.05-7.01 (m, 1H), 6.89 (d, *J*=9.3 Hz, 1H), 5.14-5.08 (m, 1H), 4.46 (s, 1H), 4.26 (t, *J*=8.3 Hz, 2H), 4.00 (d, *J*=9.0 Hz, 1H), 3.86-3.81 (m, 2H), 3.27 (s, 4H), 3.11 (m, 1H), 2.96-2.83 (m, 1H), 2.71 (d, *J*=7.2 Hz, 2H), 2.60 (m, 6H), 2.06-2.02 (m, 1H), 1.22 (s, 6H), 1.14 (s, 6H);	65
577		840.560	B	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.24 (d, *J* = 9.2 Hz, 1H), 7.90 (d, *J* = 8.7 Hz, 1H), 7.82 (d, *J* = 9.6 Hz, 1H), 7.37 (d, *J* = 9.7 Hz, 1H), 7.25 (d, *J* = 2.4 Hz, 2H), 7.10-7.00 (m, 2H), 5.07 (dd, *J* = 12.7, 5.4 Hz, 1H), 4.54-4.44 (m, 3H), 4.01 (d, *J* = 9.1 Hz, 1H), 3.51 (s, 4H), 3.05 (t, *J* = 12.2 Hz, 2H), 2.88 (ddd, *J* = 16.6, 13.6, 5.3 Hz, 1H), 2.63-2.51 (m, 3H), 2.24 (s, 2H), 2.20-2.10 (m, 2H), 2.10-2.00(m, 1H), 1.95-1.85(m, 1H), 1.85-1.76(m, 2H), 1.18 (d, *J* = 31.8 Hz, 15H).	49, 66





578		808.580	A	1H NMR (300 MHz, DMSO-*d6*) δ 10.78(s, 1H), 8.77 (s, 2 H), 7.92 (d, *J* = 8.8 Hz, 1H), 7.75 (d, *J* = 9.6 Hz, 2H), 7.34 – 6.92 (m, 4H), 5.43 (s, 2H), 4.77 (d, *J* = 13.2 Hz, 3H), 4.32 (s, 1H), 4.06 (d, *J* = 9.1 Hz, 2H), 3.65 (s, 1H), 3.05 – 3.09 (m, 5H), 2.82 – 2.62 (m, 2H), 1.91 – 2.29 (m, 8H), 1.18 (d, *J* = 31.3 Hz, 16H);	49, 67
579		837.568	A	H-NMR (300 MHz, DMSO-*d6*) δ 11.09 (s, 1H), 8.57 (s, 1H), 8.24-8.04 (m, 2H), 7.86-7.80 (m, 2H), 7.36-7.09 (m, 4H), 5.13-5.07 (m, 1H), 4.88 (s, 1H), 4.50-4.46 (d, *J* = 12 Hz, 3H), 3.82-3.65 (m, 2H), 2.99-2.82 (m, 4H), 2.60 (s, 1H), 2.41-2.40 (d, *J* = 4.8 Hz, 2H), 2.22-2.06 (m, 7H), 1.85 (s, 4H), 1.64-1.51 (m, 5H), 1.46-1.32 (m, 9H);	25B
580		837.567	A	H-NMR-PH-ARV-LS-059-H-0: (300 MHz, DMSO-*d6*, ppm) δ 11.09 (s, 1H), 8.72 (s, 2H), 8.08 (d, *J* = 7.5 Hz, 1H), 7.86-7.82 (m, 2H), 7.37 (s, 1H), 7.26-7.14 (m, 2H), 7.14-7.11 (m, 1H), 5.15-5.09 (m, 1H), 4.91-4.88 (m, 1H), 4.78-4.72 (m, 2H), 4.56-4.53 (m, 1H), 3.77-3.61 (m, 2H), 2.94-2.90 (m, 4H), 2.54-2.51 (m, 1H), 2.48-2.47 (m, 2H), 2.24-2.14 (m, 7H), 1.91-1.80 (m, 4H), 1.71-1.50 (m, 5H), 0.96-0.90 (m, 9H);	25B
581		886.590	A	^{1}H NMR (300 MHz, DMSO): δ 11.15 (s, 1H), 8.23 (d, *J* = 9.6 Hz, 1H), 7.91 (d, *J* = 8.8 Hz, 1H), 7.70 (d, *J* = 9.8 Hz, 1H), 7.24 (d, *J* = 2.4 Hz, 1H), 7.02 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.13 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.74 (d, *J* = 12.6 Hz, 2H), 4.27 (s, 1H), 4.09 (d, *J* = 9.5 Hz, 1H), 3.31 -3.22 (m, 6H),2.89 (q, *J* = 14.1, 11.4 Hz, 3H), 2.61 (d, *J* = 18.0 Hz, 2H), 2.46 – 2.43(m, 2H),	63




				2.24 (s, 8H), 2.04 (d, *J* = 12.2 Hz, 1H), 1.81 (d, *J* = 12.5 Hz, 3H), 1.19 (s, 6H), 1.12 (s, 6H), 1.01 (d, *J* = 12.8 Hz, 2H);	
582		872.570	A	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.14 (s, 1H), 8.30 – 8.14 (m, 1H), 7.92 (d, *J* = 8.7 Hz, 1H), 7.83 (d, *J* = 9.5 Hz, 1H), 7.69 (d, *J* = 9.8 Hz, 1H), 7.38 (d, *J* = 9.7 Hz, 1H), 7.26 (d, *J* = 2.4 Hz, 1H), 7.05 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.13 (dd, *J* = 12.8, 5.3 Hz, 1H), 4.52-4.46 (m, 3H), 4.02-3.99 (d, *J* = 9.1 Hz, 1H), 3.44-3.29 (m, 3H), 3.05-2.93 (m, 4H), 2.62-2.56 (m, 4H), 2.47 – 2.35 (m, 4H), 2.10 – 1.99 (m, 1H), 1.82-1.80 (m, 2H), 1.70-1.69 (m, 1H), 1.54-1.44 (m, 2H), 1.22-1.20 (m, 14H)	49
583		968.670	C	^{1}H NMR (300 MHz, DMSO-d_6) δ11.11 (s, 1H), 8.55 (s, 1H), 7.89 (d, *J* = 9.0 Hz, 1H), 7.75-7.65 (m, 2H), 7.46 (d, *J* = 9.0 Hz, 1H), 7.23 (m, 1H), 7.04-7.00 (m, 1H), 6.38 (s, 1H), 5.13-5.07 (m, 1H), 4.47-4.42 (m, 2H), 4.36 (m, 2H), 4.29 (s, 1H), 4.04 (d, *J* = 9.0 Hz, 1H), 3.60 (m, 4H), 3.31 (m, 1H), 3.26 (m, 4H), 2.94-2.85 (m, 3H), 2.80-2.76 (m, 2H), 2.63 (m, 1H), 2.54 (m, 4H), 2.47 (m, 4H), 2.23 (m, 2H), 2.08 (m, 1H), 1.80 (m, 3H), 1.22 (s, 6H), 1.19 (m, 1H), 1.10 (s, 6H), 1.05 (m, 1H);	60
584		837.570	A		72




585		815.640	D	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.07 (s, 1H), 8.07 (d, J = 8.6 Hz, 1H), 7.75 (d, J = 8.6 Hz, 2H), 7.68 (d, J = 8.5 Hz, 1H), 7.52 (d, J = 9.4 Hz, 1H), 7.35 (d, J = 2.3 Hz, 1H), 7.26 (dd, J = 8.6, 2.1 Hz, 1H), 6.96 (d, J = 8.6 Hz, 2H), 6.87 (d, J = 8.6 Hz, 1H), 5.07 (dd, J = 12.9, 5.4 Hz, 1H), 4.79 (s, 1H), 4.04 (d, J = 9.2 Hz, 1H), 3.86 (d, J = 12.4 Hz, 2H), 3.44 (d, J = 5.2 Hz, 4H), 2.96 -2.74 (m, 5H), 2.63-2.53 (m, 2H), 2.52(s, 1H),2.50(s, 3H), 2.22 (d, J = 6.6 Hz, 2H), 2.02 (d, J = 12.7 Hz, 1H), 1.81 (d, J = 12.2 Hz, 3H), 1.32-1.24 (m, 3H),1.22 (s, 8H), 1.12 (s, 6H);	49
586		837.570	B		72
587		810.530	A	^{1}H NMR (400 MHz, *d6*-DMSO) δ11.13 (s, 1H), 7.93-7.90 (d, *J* = 8.8Hz, 1H), 7.76-7.73 (m, 3H), 7.48-7.44 (m, 2H), 7.22-7.21 (d, *J* = 2.4Hz, 1H), 7.02-7.00 (m, 1H), 6.45-6.43 (d, *J* = 8.8Hz, 2H), 5.14-5.10 (m, 1H), 4.32 (s, 1H), 4.07-4.00 (m, 3H), 3.59-3.56 (m, 2H), 3.26 (s, 4H), 3.00-2.80 (m, 2H), 2.69-2.62 (m, 2H), 2.59-2.51 (m, 6H), 2.08-2.03 (m, 1H), 1.22 (s, 6H), 1.13(m, 6H)	65




588		819.575	A	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.07 (s, 1H), 7.90 (d, J = 8.7 Hz, 1H), 7.77 (dd, J = 8.6, 5.8 Hz, 3H), 7.66 (d, J = 8.6 Hz, 1H), 7.40 – 7.29 (m, 3H), 7.27 – 7.18 (m, 2H), 7.01 (dd, J = 8.8, 2.5 Hz, 1H), 5.07 (dd, J = 12.9, 5.4 Hz, 1H), 4.32 (s, 1H), 4.06 (dd, J = 10.3, 5.2 Hz, 3H), 3.02 – 2.82 (m, 5H), 2.68 – 2.51 (m, 3H), 2.18 (d, J = 6.7 Hz, 2H), 2.05 – 1.94 (m, 3H), 1.82 (d, J = 12. 2 Hz, 3H), 1.79 – 1.61 (m, 4H), 1.18 (d, J = 35.8 Hz, 14H).	69
589		821.560	A		
590		824.580	A	^{1}H NMR (300 MHz, DMSO-*d6*): δ10.96 (s, 1H), 7.90-7.87 (m, 1H), 7.76-7.70 (m, 2H), 7.54-7.48(m, 1H), 7.42-7.38(m, 1H), 7.29-7.27(m, 1H), 7.19-7.18(m, 1H), 7.00-6.94 (m, 3H), 5.09-5.03(m, 1H), 4.37-4.18 (m, 3H), 4.05-4.02 (m, 1H), 3.63-3.45(m, 2H), 3.28-3.16(m, 5H), 2.93-2.59 (m, 5H), 2.48-2.18 (m, 4H), 1.98-1.70(m, 4H), 1.35-0.96 (m, 15H)	70
591		824.580	A	^{1}H NMR (300 MHz, DMSO-*d6*): δ 10.95 (s, 1H), 7.91-7.86 (m, 1H), 7.76-7.69 (m, 2H), 7.56-7.46(m, 1H), 7.42-7.34 (m, 1H), 7.30-7.16 (m, 2H), 7.05-6.89 (m, 3H), 5.08-5.00 (m, 1H), 4.38-4.13 (m, 3H), 4.06-4.01 (m, 1H), 3.49-3.42 (m, 2H), 3.24-3.13 (m, 4H), 2.94-2.58 (m, 5H),	70




				2.36-2.14 (m, 4H), 2.07-1.71 (m, 5H), 1.43-0.96 (m, 15H);	
592		858.550	A	1H NMR (400 MHz, DMSO-d6) δ 11.12 (s, 1H), 8.24 (d, J = 9.3 Hz, 1H), 7.90 (d, J = 8.8 Hz, 1H), 7.82 (d, J = 9.6 Hz, 1H), 7.68 (d, J = 9.8 Hz, 1H), 7.37 (d, J = 9.7 Hz, 1H), 7.25 (d, J = 2.4 Hz, 1H), 7.04 (dd, J = 8.7, 2.4 Hz, 1H), 5.11 (dd, J = 12.9, 5.5 Hz, 1H), 4.55 – 4.44 (m, 3H), 4.01 (d, J = 9.1 Hz, 1H), 3.36-3.29 (m, 4H), 3.04 (t, J = 12.5 Hz, 2H), 2.91 – 2.81 (m, 1H), 2.60-2.51 (m, 3H), 2.48-2.40 (m, 3H)2.23 (d, J = 7.1 Hz, 2H), 2.10 – 1.99 (m, 1H), 1.93 (s, 1H), 1.85 (d, J = 13.5 Hz, 2H), 1.22 (s, 6H), 1.14 (s, 8H).	49
593		804.480	C	^{1}H NMR (300 MHz, DMSO-*d6*) δ11.16 (s, 1H), 8.33-8.29 (m, 2H), 8.06 (d, *J* = 7.8 Hz, 1H), 7.96-7.68 (m, 10H), 7.54 (s, 1H), 7.45 (d, *J* = 2.4 Hz, 1H), 7.19-7.15 (m, 1H), 5.25-5.19 (m, 1H), 4.69 (m, 1H), 3.94 (s, 3H), 3.65 (s, 3H), 2.86 (m, 2H), 2.65-2.54 (m, 2H), 2.20-2.16 (m, 3H), 1.96-1.84 (m, 4H), 1.59-1.55 (m, 2H);	19 B
594		830.510	A	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.12 (s, 1H), 8.58 (d, *J* = 8.3 Hz, 1H), 7.83 (dd, *J* = 20.9, 9.2 Hz, 2H), 7.69 (d, *J* = 9.8 Hz, 1H), 7.39 (d, *J* = 2.4 Hz, 1H), 7.34 (d, *J* = 9.7 Hz, 1H), 7.14 (dd, *J* = 8.7, 2.4 Hz, 1H), 5.13-5.11 (m, 1H), 4.51 (t, *J* = 13.8 Hz, 3H), 3.87 (s, 1H), 3.21 (s, 1H),3.02 (t, *J* = 12.4 Hz, 2H), 2.94-2.81 (m, 1H), 2.59 (d, *J* = 18.6 Hz, 5H), 2.46 (m, 3H), 2.24-2.11(m, 2H), 2.07-1.87 (m, 9H), 1.66-1.62 (m, 2H), 1.55-1.52 (m, 2H), 1.14-1.05 (m, 2H);	5 B



595		858.550	A	^{1}H NMR (300 MHz, Methanol-d4) δ 8.75 (s, 2H), 7.75 (d, J = 8.7 Hz, 1H), 7.49 (d, J = 9.8 Hz, 1H), 7.15 (d, J = 2.4 Hz, 1H), 7.00 (dd, J = 8.8, 2.5 Hz, 1H), 5.12 (dd, J = 12.5, 5.5 Hz, 1H), 4.91 (s, 2H), 4.29 (s, 1H), 4.15 (s, 1H), 3.43 (s, 4H), 3.10-2.70 (m,5H), 2.66 (m, 4H), 2.36 (d, J = 6.7 Hz, 2H), 2.14 (s, 1H), 1.94 (d, J = 13.4 Hz, 3H), 1.30 (s, 6H), 1.23 (s, 8H)	49
596		764.500	A	^{1}H NMR (400 MHz, *d6*-DMSO) δ 11.09 (s, 1H), 7.93-7.91 (d, *J* = 7.6Hz, 1H), 7.87-7.84 (d, J = 8.8Hz, 1H), 7.72-7.67 (m, 3H), 7.39-7.35 (m, 2H), 7.28-7.25 (m, 1H), 7.15-7.12 (m, 1H), 6.41-6.39 (d, J = 8.4Hz, 2H), 5.09-5.05 (m, 1H), 4.54-4.53 (m, 1H), 4.01-3.97 (m, 2H), 3.80(s, 1H), 3.56-3.53 (m, 2H), 3.44 (s, 4H), 2.99-2.80 (m, 2H), 2.65-2.53 (m, 8H), 2.11-2.00 (m, 3H), 1.89-1.87 (m, 2H), 1.53-1.46 (m, 4H);	5, 16 B
597		865.600	A	^{1}H NMR (300 MHz, DMSO-*d6*) 11.09 (s, 1H), 8.59 (s, 1H), 8.31(s, 1H), 7.91-7.73 (m, 3H), 7.25-7.22 (m, 3H), 7.02-6.99 (m, 1H), 5.16-5.03 (m, 1H), 4.88-4.79 (s, 1H), 4.51-4.47 (m, 3H), 3.96-3.90 (m, 1H), 3.72-3.61 (s, 1H), 2.96-2.81 (m, 4H), 2.71-2.69 (m, 1H), 2.68-2.48 (m, 2H), 2.34-2.16 (m, 4H), 2.12-2.00 (s, 1H), 1.91-1.82 (m, 2H), 1.80-1.78 (s, 1H), 1.31-1.06 (m, 15H), 0.96-0.80 (m, 6H);	71
598		821.570	A	^{1}H NMR (400 MHz, DMSO-*d6*) 11.09 (s, 1H), 9.06 (s,2H), 8.24 (m, 1H), 7.92 (m, 1H), 7.67 (m, 1H), 7.31 (s, 1H), 7.24 (m, 2H), 7.03 (m, 1H), 5.21-5.02 (m, 1H), 4.31 (m, 1H), 4.21-4.15 (m, 3H), 3.00-2.81 (m, 6H), 2.71-2.56 (m, 2H), 2.31-2.20 (s, 2H), 2.10-2.02 (m, 3H), 2.00-1.91	70




				(m, 2H), 2.96-2.81 (m, 5H), 1.30-1.28 (m, 6H), 1.21-1.06 (m, 8H);	
599		826.540	A	^{1}H NMR (300 MHz, *d6*-DMSO) δ11.09 (s, 1H), 8.26-8.23 (d, *J* = 9.0Hz, 1H), 7.92-7.89 (d, *J* = 8.7Hz, 1H), 7.85-7.82 (d, *J* = 9.3Hz, 1H), 7.42-7.40 (d, *J* = 9.9Hz, 1H), 7.25-7.23 (m, 2H), 7.07-7.01 (m, 2H), 5.09-5.05 (m, 1H), 4.56-4.51 (m, 2H), 4.46 (s, 1H), 4.02-3.99 (d, *J* = 9.0Hz, 1H), 3.47 (m, 4H), 3.08-3.04 (m, 2H), 3.00-2.80 (m, 1H), 2.63-2.56 (m, 6H), 2.50-2.49 (m, 1H), 2.01-1.89 (m, 3H), 1.47-1.43 (m, 2H), 1.22(s, 6H), 1.14 (s, 6H);	49,, 66
600		865.600	A	H-NMR (300 MHz, DMSO-*d6*) δ 11.09 (s, 1H), 8.23-8.20 (d, *J* = 9.3 Hz, 1H), 7.90-7.78 (m, 3H), 7.36-7.33 (d, *J* = 9.9 Hz, 1H), 7.26-7.23 (m, 3H), 7.03-7.00 (m, 1H), 5.12-5.06 (m, 1H), 4.88 (s, 1H), 4.53-4.44 (m, 3H), 4.00-3.97 (d, *J* = 9.0 Hz, 1H), 3.33-2.87 (m, 4H), 2.81-2.38 (m, 3H), 2.25-2.14 (m, 4H), 2.04-2.00 (m, 1H), 1.86-1.69 (m, 3H), 1.20-1.00 (m, 15H), 0.92-0.89 (d, *J* = 9.9 Hz, 6H);	72
601		836.570	A	^{1}H NMR (300 MHz, DMSO-d_6) δ 8.26 (s, 1H), 7.84 (dd, J = 8.6, 4.9 Hz, 3H), 7.43 (d, J = 8.7 Hz, 1H), 7.38 – 7.26 (m, 3H), 7.10 (d, J = 8.9 Hz, 1H), 5.16 – 5.03 (m, 2H), 4.51 (s, 1H), 4.17 (d, J = 9.2 Hz, 1H), 3.93 (s, 2H), 3.79 (s, 1H), 3.05 (s, 1H), 3.05 (s, 3H), 2.86 (s, 4H), 2.60 (s, 3H), 2.06 (dd, J = 11.1, 12.5 Hz, 8H), 1.57 (dd, J = 25.4, 12.6 Hz, 4H), 1.32 – 1.12 (m, 9H);	72




602		812.590	A	^{1}H NMR (300 MHz, DMSO-d_6): δ 10.96 (s, 1H), 8.88 (s, 1H), 8.15-7.94 (m, 2H), 7.88 (d, J = 8.8 Hz, 1H), 7.45-7.30 (m, 2H), 7.25-7.16 (m, 2H), 7.02-6.99 (m, 1H), 5.10-5.02 (m, 1H), 4.41-4.26 (m, 3H), 4.22 (d, J = 17.1 Hz, 1H), 3.08-2.67 (m, 3H), 2.98-2.60 (m, 6H), 2.53-2.50 (m, 2H), 2.42-2.24 (m, 3H), 2.09-1.90 (m, 1H) 1.72-1.68 (m, 2H), 1.50-1.37 (m, 2H), 1.35-1.30 (m, 4H),1.21 (s, 6H), 1.11 (s, 7H).	20
603		787.520	D	^{1}H NMR (300 MHz, Me_2SO-d_6): δ 11.08 (s, 1H), 7.86 (d, J = 8.8 Hz, 1H), 7.71-7.67 (m, 3H), 7.39-7.25 (m, 3H), 7.13 (dd, J = 8.8, 2.3 Hz, 1H), 6.74 (s, 1H), 6.45 (d, J = 8.4 Hz, 2H), 5.07 (dd, J = 12.7, 5.5 Hz, 1H), 4.60 (brs, 1H), 3.98 (t, J = 7.6 Hz, 2H), 3.54-3.30 (m, 7H), 3.29-3.120 (m, 5H), 3.11-2.79 (m, 3H), 2.67-2.55 (m, 3H), 2.18-2.01 (m, 5H), 1.75-1.35 (m, 4H);	20 B
604		809.570	B		

605		687.420	D	^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.13 (s, 1H), 7.97 – 7.82 (m, 3H), 7.45 (d, *J* = 2.2 Hz, 1H), 7.36 (dd, *J* = 8.3, 2.3 Hz, 1H), 7.27 – 7.17 (m, 2H), 7.01 (dd, *J* = 8.8, 2.4 Hz, 1H), 6.67 (d, *J* = 2.3 Hz, 1H), 5.14 (dd, *J* = 12.8, 5.3 Hz, 1H), 4.47 – 4.30 (m, 3H), 4.19 (t, *J* = 5.8 Hz, 2H), 3.92 (d, *J* = 8.9 Hz, 1H), 2.93 – 2.81 (m, 1H), 2.60 (d, *J* = 19.9 Hz, 2H), 2.31 (d, *J* = 7.4 Hz, 2H), 2.06 (s, 1H), 1.14 – 1.06 (m, 12H);	73
606		781.530	A		
607		823.470	A	^{1}H NMR (400 MHz, DMSO-*d6*) δ 11.08 (s, 1H), 8.89 (s, 1H), 8.53 (s, 1H), 7.69 – 7.56 (m, 2H), 7.35 – 7.21 (m, 4H), 5.10-5.00 (m, 1H), 4.06 (d, *J* = 13.0 Hz, 2H), 3.0-2.90 (m, 4H), 2.88-2.86 (m, 2H), 2.61-2.59 (m, 4H), 2.50-2.40 (m, 2H), 2.20-2.18 (m, 2H), 2.03-1.90 (m, 4H), 1.90-1.80 (m, 7H), 1.60-1.50 (m, 1H), 1.26 – 1.13 (m, 2H);	74



608		806.470	A	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.07 (s, 1H), 8.88 (d, J = 1.9 Hz, 1H), 8.52 (d, J = 1.9 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.33 (d, J = 1.9 Hz, 1H), 7.30 – 7.12 (m, 3H), 7.07 (d, J = 9.0 Hz, 2H), 5.06 (dd, J = 12.6, 5.3 Hz, 1H), 3.81 (d, J = 12.1 Hz, 2H), 3.44 (s, 4H), 2.87 (t, J = 13.4 Hz,4H), 2.76 (t, J = 12.6 Hz, 5H), 2.56 (d, J = 15.3 Hz, 3H), 2.41 (s, 2H), 1.97 -1.75 (s, 6H), 1.55 (s, 1H), 1.23 (d, J = 10.9 Hz, 2H).	75
609		820.610	A	H-NMR-PH-ARV-LS-046-E-0: (300 MHz, DMSO, ppm) δ 10.90-10.88 (br, 1H), 7.92-7.89 (d, J = 9Hz, 1H), 7.75-7.72 (d, J = 9Hz, 2H), 7.51-7.44 (m, 2H), 7.21 (br, 1H), 7.10-6.94 (m, 5H), 4.65-4.64 (m, 1H), 4.54-4.52 (m, 1H), 4.32 (s, 1H), 4.06-4.03 (d, J = 9Hz, 1H), 3.88-3.84 (m, 2H), 3.31-3.28 (m, 4H), 2.83-2.75 (m, 3H), 2.63-2.58 (m, 6H), 2.22-2.20 (m, 2H), 1.96-1.94 (m, 1H), 1.83-1.79 (m, 3H), 1.43-1.38 (m, 3H), 1.21 (s, 6H), 1.18-1.17 (m, 2H), 1.12 (s, 6H);	49, 76
610		820.610	A		49, 76




611		840.560	B	^{1}H NMR (400 MHz, DMSO-d_6): δ 11.12 (s, 1H), 8.25 (d, J = 9.2 Hz, 1H), 7.92 (d, J = 8.7 Hz, 1H), 7.82 (d, J = 9.5 Hz, 1H), 7.65 (d, J = 8.1 Hz, 1H), 7.38 (t, J = 8.2 Hz, 2H), 7.25 (d, J = 2.4 Hz, 1H), 7.03 (dd, J = 8.8, 2.5 Hz, 1H), 5.10 (dd, J = 12.7, 5.4 Hz, 1H), 4.55-4.44 (m, 3H), 4.01 (d, J = 9.2 Hz, 1H), 3.25-3.22 (m, 4H), 3.04 (t, J = 12.4 Hz, 2H), 2.95-2.81 (m, 1H), 2.60-2.56 (m, 5H), 2.23 (d, J = 7.1 Hz, 2H), 2.10-1.99 (m, 1H), 1.94 (s, 1H), 1.84 (d, J = 12.7 Hz, 2H), 1.22 (s, 6H), 1.14 (s, 9H).	49, 77
612		824.470	A	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.06 (s, 1H), 8.87 (d, J = 2.0 Hz, 1H), 8.51 (d, J = 2.0 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.34 – 7.09 (m, 5H), 5.06 (dd, J = 12.7, 5.3 Hz, 1H), 3.47 (d, J = 14.2 Hz, 6H), 3.47 (s, 1H), 2.76 (t, J = 12.5 Hz, 3H), 2.56 (d, J = 14.7 Hz, 6H), 2.25 (d, J = 6.4 Hz, 3H), 2.00 - 1.56 (d, J = 12.1 Hz, 7H), 1.36 – 1.19 (m, 3H).	75

613		805.480	B		74
614		666.520	D	^{1}H NMR (300 MHz, Methanol-d_4) δ 7.69 (d, J = 8.5 Hz, 1H), 7.53 (d, J = 8.8 Hz, 2H), 7.36 (d, J = 2.3 Hz, 1H), 7.29 – 7.15 (m, 2H), 7.14 – 7.05 (m, 2H), 5.07 (dd, J = 12.3, 5.4 Hz, 1H), 4.39 (t, J = 4.9 Hz, 2H), 4.23 (t, J = 7.0 Hz, 2H), 3.62-3.47 (m, 9H), 3.07-2.69 (m, 14H), 2.35 (d, J = 7.1 Hz, 2H), 2.13-2.09 (m, 3H), 1.94-1.90 (m, 1H), 1.51-1.47 (m, 2H);	78
615		865.600	A	1H NMR (300 MHz, DMSO-*d6*) δ 11.10 (s, 1H), 8.75 (s, 2H), 7.90 (d, J = 6.6 Hz, 1H), 7.83 (d, J = 6.3 Hz, 1H), 7.71–7.69 (m, 1H), 7.28 – 7.21 (m, 3H), 7.02 – 7.00 (m, 1H), 5.13 – 5.09 (m, 1H), 4.90 – 4.85 (m, 1H), 4.79 – 4.75 (m, 2H), 4.29 (s, 1H), 4.05 (d, J = 6.9 Hz, 1H), 3.70 – 3.60 (m, 1H), 2.94 – 2.89 (m, 4H), 2.57 – 2.50 (m, 1H), 2.49 – 2.40 (m, 2H), 2.25 – 2.19 (m, 4H), 2.08 – 2.00 (m, 1H), 1.90 – 1.80 (m, 2H), 1.75 – 1.65 (s, 1H), 1.20 (s, 7H), 1.10 (s, 6H), 1.00 – 0.91 (m, 8H);	72

616		861.570	A	1H NMR (300 MHz, DMSO-*d6*): δ 11.06 (s ,1H), 7.92 (d, *J* = 8.7 Hz, 1H), 7.75-7.71 (m, 3H), 7.42 (d, *J* = 9.3 Hz, 1H), 7.34 (s, 1H), 7.29 – 7.18 (m, 2H), 7.01 (dd, *J* = 8.8, 2.5 Hz, 1H), 6.58 (d, *J* = 8.7 Hz, 2H), 5.08 (dd, *J* = 12.7, 5.3 Hz, 1H), 4.33 (s, 1H), 4.07 (d, *J* = 9.1 Hz, 1H), 3.73 - 3.63 (m, 2H), , 3.51- 3.48 (m, 6H), 3.32 – 3.16 (m, 3H), 3.31 - 3.19 (m, 4H), 3.17 - 2.81 (m, 3H), 2.06 (d, *J* = 14.9 Hz, 1H), 1.22 (s, 6H), 1.13 (s, 6H);	49, 79
617		823.590	D	^{1}H NMR (400 MHz, DMSO-*d6*) 11.08 (s, 1H), 7.91-7.89 (m, 2H), 7.66-7.64 (m, 1H), 7.32-7.23 (m, 4H), 7.04-7.01 (m, 1H), 6.67-6.66 (m, 1H), 5.08-5.04 (m, 1H), 4.36-4.30 (m, 2H), 4.12-4.09 (m, 2H), 3.95-3.92 (m, 1H), 3.02-2.88 (m, 4H), 2.67-2.49 (m, 3H), 2.18-1.84 (m, 10H), 1.71-1.46 (m, 6H), 1.20-1.09 (m, 12H).	49, 80
618		622.480	D	^{1}H NMR (300 MHz, DMSO-d_6) δ 8.27 (s, 1H), 7.66 (d, *J* = 8.5 Hz, 1H), 7.41 – 7.28 (m, 3H), 7.24 (dd, *J* = 8.7, 2.3 Hz, 1H), 7.04 – 6.89 (m, 3H), 5.05 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.09 (t, *J* = 7.0 Hz, 2H), 3.70-3.60 (m, 2H), 2.95 – 2.76 (m, 11H), 2.73 – 2.49 (m, 5H), 2.20 (d, *J* = 6.9 Hz, 2H), 1.99 (d, *J* = 12.3 Hz, 1H), 1.79 (d, *J* = 13.1 Hz, 2H), 1.70-1.65 (m, 1H), 1.29-1.10 (m, 3H)	81

619		837.570	B
			72

620		852.605	B		
621		861.650	A	^{1}H-NMR (400 MHz, $CDCl_3$) δ 8.70 (s, 2H), 8.05 (s, 1H), 7.77 (d, J = 8.0 Hz, 1H), 7.45 (d, J = 8.0 Hz, 1H), 7.18 (s, 1H), 7.17-7.0 (m, 1H), 6.45-6.37 (m, 2H), 5.94-5.92 (d, J = 8.0 Hz, 1H), 4.97-4.87 (m, 3H), 4.74 (m, 1H), 4.13-4.11 (d, J = 8.0 Hz, 1H), 4.03 (s, 1H), 3.91(s, 3H), 3.82-3.72 (m, 1H), 2.96-2.74 (m, 6H), 2.40-2.14 (m, 7H), 1.94-1.91 (m, 2H), 1.32-1.24 (m, 14H), 1.22-1.06 (m, 1H), 0.98-0.97 (m, 6H).	72

622		845.650	A	$_1$H-NMR (400 MHz, $CDCl_3$) δ 8.70 (s, 2H), 7.98 (s, 1H), 7.78-7.76 (d, *J* = 8.0 Hz, 1H), 7.52-7.50(d, *J* = 8.0 Hz, 1H), 7.18 (s, 1H), 7.13-7.11 (d, *J* = 8.0 Hz, 1H), 6.77 (s, 1H), 6.69-6.67 (d, J = 8.0 Hz, 1H), 5.94-5.92 (d, *J* = 8.0 Hz, 1H), 4.97-4.87 (m, 3H), 4.74 (m, 1H), 4.12-4.10 (d, *J* = 8.0 Hz, 1H), 4.03 (s, 1H), 3.74-3.70 (m, 1H), 2.96-2.74 (m, 6H), 2.51 (s, 3H), 2.38-2.13 (m, 7H), 1.94-1.91 (m, 2H), 1.67-1.30 (m, 2H), 1.24 (s, 12H), 1.22-0.96 (m, 7H).	72
623		859.670	A		72
624		806.595	A	^{1}H NMR (300 MHz, DMSO): δ10.68(s, 1H), 7.91-7.87 (m, 1H), 7.72-7.69 (m, 2H), 7.49-7.46(m, 2H), 7.19-7.18 (m, 1H), 7.05-6.89 (m, 5H), 5.28(s, 2H), 4.57-4.52(m, 1H), 4.29 (s, 1H), 4.04-4.01 (m, 1H), 3.86-3.82 (m, 2H), 3.31-3.26(m, 10H), 2.80-2.72 (m, 2H),	13B











				2.61-2.57 (m, 2H), 2.19-2.17(m, 2H), 2.10-1.87 (m, 2H), 1.82-1.67 (m, 3H), 1.19 (s, 6H), 1.10 (s, 6H);	
625		823.590	A	^{1}H NMR (300 MHz, DMSO-*d6*) δ 10.96 (s, 1H), 8.72 (s, 2H), 8.09 (d, *J* = 7.4 Hz, 1H), 7.84 (d, *J* = 8.8 Hz, 1H), 7.60 (d, *J* = 8.4 Hz, 1H), 7.37 (d, *J* = 2.4 Hz, 1H), 7.14-7.12 (m, 1H), 7.04 – 7.01 (m, 1H),6.98-6.88(m,1H), 5.13-5.05 (m, 1H), 4.74 (d, *J* = 11.8 Hz,3H), 4.53 (s, 1H), 4.42-4.36 (m, 1H), 4.28-4.22 (m, 1H), 3.77 (s, 1H), 3.62 (t, J = 8.0 Hz, 1H), 2.89 (t, *J* = 12.9 Hz, 4H),2.62- 2.57 (m ,1H), 2.43-2.35 (m, 3H), 2.28- 2.12 (m, 6H), 2.08-1.79 (m, 5H), 1.77-1.38 (m, 5H), 0.90 (d, *J* = 6.5 Hz, 8H).	



FIG. 3



Compound Structure	Potency	M/Z+ found	1H NMR	Synthetic Scheme
	A	827.3	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.21 (6H, s), 1.25 (6H, s), 1.37-1.49 (4H, m), 1.50-1.57 (2H, m), 1.62-1.74 (2H, m), 2.18-2.30 (2H, m), 2.32-2.41 (2H, m), 2.53-2.55 (4H, m), 2.81-2.93 (2H, m), 3.18-3.23 (2H, m), 3.64-3.68 (4H, m), 4.04 (1H, s), 4.13-4.24 (2H, m), 4.34-4.38 (2H, m), 5.15-5.20 (1H, m), 6.07 (1H, d, *J* = 8.4Hz), 6.64 (2H, m), 6.79-6.82 (1H, m), 6.96-6.97 (1H, d, *J* = 2.4 Hz), 7.42 (1H, d, *J* = 8.8 Hz), 7.57 (1H, d, *J* = 8.8 Hz), 7.91-7.94 (1H, m), 8.04 (1H, brs), 8.74 (1H, d, *J* = 2.0 Hz).	57
	A	836.4	^{1}H NMR (400 MHz, δ6-DMSO): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.05 (m, 1H), 2.54-2.73 (m, 12H), 2.84-2.92 (m, 1H), 3.41-3.70 (m, 10H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.30 (s, 1H), 5.05-5.10 (m, 1H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (br, 1H), 7.63 (d, *J* = 9.2 Hz, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 7.96-7.99 (m, 1H), 8.63 (d, *J* = 2.0 Hz, 1H), 11.08 (s, 1H).	38




	A	835.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.22 (s, 6H), 1.28 (s, 6H), 1.61-1.72 (m, 3H), 1.87-1.90 (m, 2H), 2.04-2.20 (m, 2H), 2.69-2.76 (m, 2H), 2.82-2.87 (m, 1H), 2.93-3.07 (m, 8H), 3.87 (br, 4H), 4.04-4.07 (m, 2H), 4.14-4.16 (m, 1H), 4.28 (s, 1H), 5.04-5.09 (m, 1H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.04-7.06 (m, 1H), 7.13 (d, *J* = 2.4 Hz, 1H), 7.20-7.23 (m, 1H),7.34-7.35 (m, 1H), 7.37-7.39 (m, 1H), 7.51-7.53 (m, 1H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.72 (d, *J* = 8.8 Hz, 1H), 7.87 (d, *J* = 7.8 Hz, 1H), 8.00-8.03 (m, 1H), 8.63 (d, *J* = 2.0 Hz, 1H).	58
	A	824.6	H-NMR (300 MHz, CD_3OD) δ: (300 MHz, CD_3OD) δ 8.62 - 8.61 (d, *J* = 2.4 Hz, 1H), 8.00 - 7.96 (d, *J* = 11.4 Hz, 1H), 7.74 - 7.71 (d, *J* = 9 Hz, 1H), 7.65 - 7.62 (m, 1H), 7.48 - 7.46 (m, 2H), 7.14 - 7.13 (d, *J* = 2.4 Hz, 1H), 7.00 - 6.97 (d, *J* = 11.1 Hz, 1H), 6.86 - 6.83 (d, *J* = 9Hz, 1H), 5.20 - 5.14 (m, 1H), 4.51 - 4.49 (d,J = 5.4 Hz, 2H), 4.28 (s, 1H), 4.15 (s, 1H), 3.70 - 3.62 (m, 6H), 3.51 - 3.46 (m, 2H), 2.99 - 2.88 (m, 1H), 2.86 - 2.65 (m, 9H), 2.62 - 2.46 (m, 1H), 2.25 - 2.11 (m, 1H), 1.76 - 1.62 (m, 4H), 1.48 - 1.44 (m, 2H), 1.29 (s, 6H), 1.22 (s, 6H)	30



	A	823.4	H-NMR: (300 MHz, CD_3OD) δ7.77 – 7.71 (m, 3H), 7.65 – 7.62 (m, 1H), 7.50 - 7.46 (m, 2H), 7.14 - 7.13 (d, *J* = 2.4 Hz, 1H), 7.00 – 6.97 (d, *J* = 9 Hz, 3H), 5.19 – 5.13 (m, 1H), 4.50 – 4.49 (d, *J* = 4.8 Hz, 2H), 4.29 (s, 1H), 4.14 (s, 1H), 3.64 - 3.61 (m, 2H), 3.50 - 3.46 (m, 2H), 3.32 - 3.31 (m, 3H), 2.90 - 2.71 (m, 11H), 2.55 - 2.49 (m, 1H), 2.20 - 2.19 (m, 1H), 1.76 - 1.62 (m, 4H), 1.51 – 1.44 (m, 2H), 1.29 (s, 6H), 1.23 (s, 6H);	30
	A	804.6	^{1}H NMR (300 MHz, CD_3OD): δ 8.10-7.90 (m, 3H), 7.76-7.71 (m, 4H), 7.34-7.31 (m, 2H), 7.13 (s, 1H), 7.00-6.96 (m, 1H), 6.41(m, 1H), 5.19-5.13 (m, 1H), 4.29 (s, 1H), 4.17-4.14 (m, 2H), 3.90-3.60 (m, 3H), 3.26-3.23 (m, 2H), 2.96-2.70 (m, 7H), 2.20-2.10 (m, 1H), 1.81-1.72 (m, 4H), 1.47-1.38 (m, 3H), 1.28-1.23 (d, *J*=16.5Hz, 12H)	42
	A	790.4	^{1}H NMR (300 MHz, CD_3OD): δ 7.75-7.60 (m, 6H), 7.33-7.30 (m, 2H), 7.13-7.12 (s, 1H), 7.00-6.96 (m, 1H), 6.31 (s, 1H), 5.18-5.12 (m, 1H), 4.49-4.48 (s, 2H), 4.29 (s, 1H), 4.10 (s, 1H), 3.23-3.22 (s, 2H), 2.91-2.46 (m, 11H), 2.19-2.10 (m, 1H), 1.69-1.60 (m, 4H), 1.40-1.30 (s, 4H), 1.28-1.23 (d, *J*=16.5Hz,12H);	27



	A	822.7	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.44 (s, 4H), 1.72-1.80 (m, 3H), 1.88-1.90 (m, 1H), 2.03-2.07 (m, 1H), 2.54-2.65 (m, 2H), 2.85-2.95 (m, 2H), 3.32-3.35 (m, 2H), 3.49-3.51 (m, 2H), 3.66 (s, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.16 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 4.69-4.75 (m, 1H), 5.09-5.14 (m, 1H), 6.51-6.54 (m, 1H), 7.00-7.02 (m, 1H), 7.20-7.21 (m, 1H), 7.32-7.34 (m, 1H), 7.41 (s, 1H), 7.55 (d, *J* = 9.2 Hz, 1H), 7.82 (d, *J* = 8.4 Hz, 1H), 7.89-7.93 (m, 2H), 8.16 (s, 1H), 8.59 (s, 1H), 11.11 (s, 1H).	45
	A	795.4	H-NMR: (300 MHz, DMSO-*d6*, ppm) δ 7.89 - 7.86 (d, *J* = 9 Hz, 1 H), 7.75 7.72 (d, *J* = 9 Hz, 2 H),7.62 - 7.59 (d, *J* = 9 Hz,1 H), 7.52 - 7.49 (d, *J* = 9 Hz, 1 H), 7.19 - 7.14 (m, 2 H), 7.04 - 6.94 (m, 4 H), 5.07 - 5.01 (m, 1 H), 4.40 - 4.21 (m, 3 H), 4.08 - 4.02 (m, 3 H), 3.36 - 3.27 (m, 4 H), 2.88 - 2.79 (m, 1 H), 2.71 - 2.55 (m, 4 H), 2.38 - 2.27 (m, 2 H), 1.99 - 1.94 (m, 1 H), 1.79 - 1.74 (m, 2 H), 1.56 - 1.44 (m, 4 H), 1.21 - 1.19 (m, 7 H), 1.10 (s, 6 H);	31



	A	806.35	H-NMR01-PH-ARV-LS-024-J-0: (300 MHz, DMSO-*d6*, ppm) δ 8.12 (br, 1 H), 7.89 - 7.79 (m, 5 H), 7.54 -7.51 (d, *J* = 9 Hz, 2 H), 7.47 - 7.41 (m, 1 H), 7.35 - 7.32 (m, 1 H), 7.22 - 7.18 (m, 1 H), 7.10 - 6.97 (m, 1 H), 6.28 (s, 1 H), 5.13 - 5.06 (m, 1 H), 4.31 (s, 1 H), 4.20 - 4.16 (m, 2 H), 4.07 - 4.04 (d, *J* = 9 Hz, 1 H), 3.29 - 3.24 (m, 2 H), 2.93 - 2.83 (m, 3 H), 2.60 - 2.56 (m, 5 H), 2.04 - 1.98 (m, 1 H), 1.81 - 1.76 (m, 2 H), 1.61 - 1.59 (m, 2 H), 1.48 - 1.45 (m, 2 H), 1.46 (s, 6 H), 1.21 (s, 6 H).	45, 50
	A	779.75	^{1}H NMR (400 MHz, $CDCl_3$): δ 7.71-7.61(m, 4H), 7.58-7.56 (m, 1H), 6.99-6.91 (m, 2H), 6.83-9.79 (m, 2H), 6.22-6.19 (m, 1H), 5.30-5.28 (m, 1H), 4.76-4.74 (m, 1H), 4.17-4.15 (m, 1H), 4.06 (s, 1H), 3.86-3.20 (m, 4H), 2.98-2.89 (m, 2H), 2.72-2.40 (m, 6H), 2.23-2.17 (m, 2H), 1.70-1.62 (m, 3H), 1.44-1.39 (m, 3H),1.27 (s, 7H), 1.23 (s, 6H);	42
	A	779.75	^{1}H NMR (400 MHz, $CDCl_3$): δ 7.71-7.61(m, 4H), 7.58-7.56 (m, 1H), 6.99-6.91 (m, 2H), 6.83-9.79 (m, 2H), 6.22-6.19 (m, 1H), 5.30-5.28 (m, 1H), 4.76-4.74 (m, 1H), 4.17-4.15 (m, 1H), 4.06 (s, 1H), 3.75-3.20 (m, 4H), 2.98-2.89 (m, 2H), 2.80-2.50 (m, 6H), 2.23-2.17 (m, 2H), 1.70-1.62 (m, 3H), 1.44-1.39 (m, 3H),1.27 (s, 7H), 1.23 (s, 6H);	42




	A	811.45	^{1}H NMR (400 MHz, $CDCl_3$): δ 7.71-7.61(m, 4H), 7.58-7.56 (m, 1H), 6.99-6.91 (m, 2H), 6.83-9.79 (m, 2H), 6.22-6.19 (m, 1H), 5.30-5.28 (m, 1H), 4.76-4.74 (m, 1H), 4.17-4.15 (m, 1H), 4.06 (s, 1H), 3.75-3.20 (m, 4H), 2.98-2.89 (m, 2H), 2.80-2.50 (m, 6H), 2.23-2.17 (m, 2H), 1.70-1.62 (m, 3H), 1.44-1.39 (m, 3H),1.27 (s, 7H), 1.23 (s, 6H);	59
	A	811.45	^{1}H NMR (300 MHz, CD_3OD): δ7.81--7.69 (m, 3H), 7.53--7.46 (m, 1H),7.42--7.28(m,3H), 7.15 (s, 1H),7.01--6.88(m,1H), 5.17--5.06(m, 1H), 4.52--4.41 (m, 2H), 4.27(s , 1H),4.16 (s, 1H), 3.75--3.57(m,4H),3.36(s,2H),3.33--3.12(m,4H),2.95--2.69(m,4H),2.59-2.35(m,1H),2.232.01(m,1H),1.89--1.65(m,4H),1.52--1.38(m,4H),1.33--1.16(m,12)	59



	A	828.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.20 (6H, s), 1.22 (6H, s), 1.37-1.49 (6H, m), 1.79 (2H, t, *J* = 7.2 Hz), 1.98 (1H, t, *J* = 6.0 Hz), 2.29-2.43 (7H, m), 2.56-2.61 (1H, m), 2.87-2.91 (1H, m), 3.59 (4H, s), 4.05 (1H, d, *J* = 9.2 Hz), 4.14 (2H, t, *J* = 6.0 Hz), 4.24-4.40 (3H, m), 5.08 (1H, dd, *J* = 13.2, 4.8 Hz), 6.85 (1H, d, *J* = 8.8 Hz), 7.00 (1H, dd, *J* = 8.8, 2.0 Hz), 7.21 (1H, d, *J* = 2.0 Hz), 7.43 (1H, d, *J* = 6.8 Hz), 7.52 (1H, d, *J* = 10.0 Hz), 7.62 (1H, d, *J* = 9.2 Hz), 7.89-7.97 (2H, m), 8.62 (1H, d, *J* = 2.0 Hz), 11.00 (1H, s).	54, 45
	A	781.25	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 8.52 - 8.20 (brs, 1H), 7.85-7.81 (m, 4H), 7.55-7.46 (m, 3H), 7.40-7.30 (m, 1H), 7.09-6.92 (m, 3H), 5.10 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.38 (d, *J* = 7.2 Hz, 1H), 4.28-4.14 (m, 2H), 4.06 (s, 1H), 3.16 (m, 3H), 3.04 (m, 3H), 2.87- 2.42 (m, 4H), 2.10-1.90 (m, 3H), 1.19-1.00 (m, 14H);	45
	A	795.75	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 7.92 – 7.66 (m, 4H), 7.56 – 7.28 (m, 3H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.02 – 6.88 (m, 3H), 5.08 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.28 - 4.19 (t, *J* = 6.4 Hz, 3H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.21 (t, *J* = 5.0 Hz, 4H), 2.90 – 2.76 (m, 1H), 2.62 – 2.46 (m, 6H), 2.37 (t, *J* = 7.2 Hz, 2H), 2.01 (dd, *J* = 11.3, 5.6 Hz, 1H), 1.76 (q, *J* = 9.6, 8.3 Hz, 2H), 1.60 (t,	45




			J = 7.4 Hz, 2H), 1.18 (s, 6H), 1.09 (s, 6H).	
	A	792.8	^{1}H NMR (300 MHz, CD_3OD): δ 7.76-7.71 (m, 4H), 7.49-7.31 (m, 4H), 7.13-7.12 (s, 1H), 7.00-6.96 (m, 1H), 6.31 (s, 1H), 5.18-5.11 (m, 1H), 4.48-4.47 (m, 2H), 4.29 (s, 1H), 4.15 (s, 1H), 3.32-3.31 (m, 2H), 2.90-2.36 (m, 10H), 2.20-2.10 (m, 1H), 2.03-1.88 (m, 4H), 1.70-1.65 (m, 4H), 1.40-1.30 (s, 4H), 1.28-1.23 (d, *J*=16.5Hz,12H)	42
	A	793.4	^{1}H NMR (300 MHz, CD_3OD): δ7.81--7.69 (m, 3H), 7.53--7.46 (m, 1H),7.42--7.28(m,3H), 7.15 (s, 1H),7.01--6.88(m,1H), 5.17--5.06(m, 1H), 4.52--4.41 (m, 2H), 4.27(s , 1H),4.16 (s, 1H), 3.75--3.57(m,4H),3.36(s,2H),3.33--3.12(m,4H),2.95--2.69(m,4H),2.59-2.35(m,1H),2.232.01(m,1H),1.89--1.65(m,4H),1.52--1.38(m,4H),1.33--1.16(m,12)	27



	A	790.8	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.14 (m, 6H), 1.23 (s, 6H), 1.34-1.35 (m, 2H), 1.62-1.66 (m, 4H), 2.05-2.09 (m, 1H), 2.65-2.69 (m, 6H), 2.86-2.956 (m, 2H), 3.17 (d, *J* = 4.4 Hz, 1H), 4.07 (d, *J* = 8.8 Hz, 1H), 4.33 (s, 1H), 5.14-5.19 (m, 1H), 6.53 (s, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.32 (d, *J* = 8.4 Hz, 2H), 7.76-7.79 (m, 3H), 7.90-7.97(m, 4H), 11.14 (s, 1H).	1
	A	796.4	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.21 (s, 6H), 1.25 (s, 6H), 1.50-1.57 (m, 2H), 1.66-1.75 (m, 2H), 1.82-1.89 (m, 2H), 2.31-2.42 (m, 2H), 2.57-2.62 (m, 2H), 2.68-2.97 (m, 6H), 3.79 (s, 4H), 4.02-4.15 (m, 4H), 4.27, 4.31 (two singles, 1H), 4.40, 4.44 (two singles, 1H), 5.18-5.23 (m, 1H), 6.07 (d, J = 8.4 Hz, 1H), 6.67 (d, J = 9.2 Hz, 1H), 6.79-6.82 (m, 1H), 6.93-7.03 (m, 1H), 7.11-7.19 (m, 1H), 7.31-7.43 (m, 2H), 7.57 (d, J = 8.8 Hz, 1H), 7.89-8.00 (m, 1H), 8.01-8.12 (m, 1H), 8.52-8.66 (m, 1H).	31
	A	776.4	^{1}H NMR (400 MHz, CD_3OD):δ 1.25 (s, 6H), 1.31 (s, 6H), 1.66-1.77 (m, 4H), 2.13-2.18 (m, 1H), 2.61-2.65 (m, 2H), 2.71-2.81 (m, 6H), 2.85-2.91 (m, 3H), 3.28-3.31 (m, 2H), 4.18 (s, 1H), 4.31 (s, 1H), 5.14-5.18 (m, 1H), 6.44-6.46 (m, 1H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.15-7.16 (m, 1H), 7.36-7.38 (m, 2H), 7.73-7.79 (m, 3H), 7.86-7.88 (m, 1H), 7.91-7.93 (m, 1H), 7.97 (br,	




			1H).	
	A	776.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.31 (s, 6H), 1.39，1.40 (d, *J* =6.4 Hz, 1H), 1.46-1.52 (m, 2H), 1.74-1.83 (m, 4H), 2.17-2.24 (m, 1H), 2.46-2.57 (m, 1H), 2.75-2.79 (t, *J* =7.6 Hz, 2H), 2.78-2.96 (m, 4H), 3.01-3.09 (m, 2H), 3.78 (s, 2H), 4.16-4.19 (m, 2H), 4.30 (s, 1H), 4.45-4.60 (m, 2H), 5.16-5.19 (m, 1H), 6.32 (s, 1H), 6.99-7.01 (m, 1H), 7.15 (d, *J* =2.4 Hz, 1H), 7.36 (d, *J* =8.4 Hz, 2H), 7.66-7.83 (m, 6H).	27
	A	808.35	H-NMR01-PH-ARV-LS-024-H-0: (300 MHz, DMSO-*d6*, ppm) δ 8.14 (br, 1 H), 7.89 - 7.74 (m, 5 H), 7.47 7.32 (m, 4 H), 7.18 - 6.96 (m, 2 H), 5.13 - 5.06 (m, 1 H), 4.30 (s, 1 H), 4.20 - 4.16 (m, 2 H), 4.05 - 4.02 (d, *J* = 9 Hz, 1 H), 3.27 - 3.24 (m, 2 H), 2.91 - 2.81 (m, 1 H), 2.77 - 2.64 (m, 3 H), 2.60 - 2.54 (m, 3 H), 2.04 - 1.97 (m, 1 H), 1.87 - 1.74 (m, 6 H), 1.70 - 1.62 (m, 2 H), 1.47 - 1.45 (m, 2 H), 1.20 (s, 6 H), 1.11 (s, 6 H),	45



	A	778.25	1H NMR (300 MHz, DMSO) 8.53 (s, 1H), 8.02-7.89 (m, 5H), 7.89-7.75 (m, 1H), 7.59-7.57 (m, 1H), 7.16-7.10 (m, 2H), 7.01-6.96 (m, 2H), 5.00-4.90 (m, 1H), 4.32-4.15 (m, 3H), 4.15-3.77 (m, 3H), 2.89-2.48 (m, 4H), 2.38-2.20 (m, 1H), 1.99-1.90 (m, 1H) , 1.77-1.68 (m, 4H), 1.48-1.39 (m, 2H), 1.19-1.10 (m, 6H), 1.10-1.01 (m, 6H)	25
	A	805.4	1H NMR (300 MHz, CDCl3) 7.81-755 (m, 5H), 7.25-7.22 (m, 1H), 6.96-6.90 (m, 1H), 6.82-6.79 (m, 1H), 6.59-6.57 (m, 1H), 6.39-6.30 (m, 1H), 6.19-6.17 (m, 1H), 5.30-5.19 (m, 2H), 4.74-4.70 (m, 1H), 4.28-4.20 (m, 1H), 4.14-4.10 (m, 1H), 4.04-4.00 (m, 1H), 3.67-3.50 (m, 1H), 3.40-3.30 (m, 2H), 3.06-2.90 (m, 2H), 2.63-2.49 (m, 6H), 2.19-1.94 (m, 4H), 1.65-1.44 (m, 4H), 1.40-1.05 (m, 16H)	42, 52
	A	779.35	^{1}H NMR (300MHz, DMSO-d6): δ 8.97 (s, 1H), 8.69 (s, 1H), 8.51-8.47 (m, 1H), 8.25-8.16 (m, 2H), 7.94-7.88 (m, 1H), 7.63-7.57 (m, 1H), 7.26-6.85 (m , 4H), 5.13-4.97 (m, 1H), 4.51-3.99 (m, 6H), 2.98-2.22 (m, 5H), 2.07-1.89 (m, 1H), 1.87-1.73 (m, 4H), 1.58-1.43 (m, 2H), 1.31-1.08 (m, 13H)	48




	A	807.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.32 (s, 6H), 1.49-1.51 (m, 4H), 1.78-1.84 (m, 4H), 2.00-2.09 (m, 2H), 2.13-2.23 (m, 2H), 2.73-2.95 (m, 6H), 3.10-3.18 (m, 4H), 3.63-3.71 (m, 2H), 4.20 (s, 1H), 4.31 (s, 1H), 5.16-5.18 (m, 1H), 6.99-7.01 (m, 1H), 7.15 (d, *J* =2.4 Hz, 1H), 7.46 (d, *J* =8.0 Hz, 1H), 7.74 (d, *J* =8.8 Hz, 1H), 7.79 (d, *J* =8.4 Hz, 1H), 7.86 (s, 1H), 7.89 (d, *J* =8.0 Hz, 1H), 8.16-8.18 (m, 1H), 8.89 (d, *J* =2.0 Hz, 1H).	11
	A	762.5	^{1}H NMR (400 MHz, δ6-DMSO): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.05 (m, 1H), 2.54-2.73 (m, 12H), 2.84-2.92 (m, 1H), 3.41-3.70 (m, 10H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.30 (s, 1H), 5.05-5.10 (m, 1H), 6.89 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (br, 1H), 7.63 (d, *J* = 9.2 Hz, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 7.96-7.99 (m, 1H), 8.63 (d, *J* = 2.0 Hz, 1H), 11.08 (s, 1H).	11



	A	833.4	H-NMR-PH-ARV-LS-026-D-0: (300 MHz, CD_3OD, ppm) δ 8.47 (s, 1H), 8.06 - 7.98 (m, 4H), 7.75 -7.71 (m, 2H), 7.18 - 7.11 (m, 3H), 7.02 - 6.98 (m, 1H), 5.15 - 5.09 (m, 1H), 4.70 - 4.57 (m, 3H), 4.45 - 4.44 (m, 1H), 4.31 (s, 1H), 4.20 (s, 1H), 3.11 - 3.08 (m, 2H), 2.92 - 2.75 (m, 7H), 2.51 - 2.45 (m, 1H), 2.20 - 2.01 (m, 7H), 1.32 (s, 6H), 1.25 (s, 6H).	10
	A	806.32	^{1}H NMR (400 MHz, DMSO) 8.610 (s, 1H), 8.10-8.02 (m, 5H), 7.99-7.90 (m, 1H), 7.21-7.19 (m, 1H), 7.07-6.90 (m, 3H), 5.05-5.00 (m, 1H), 4.95-4.85 (m, 1H), 4.45-4.25 (m, 3H), 4.15-4.05 (m, 1H), 3.09-3.00 (m, 2H), 2.90-2.80 (m, 1H), 2.79-2.70 (m, 2H), 2.65-2.60 (m, 4H), 2.05-1.95 (m, 1H), 1.75-1.65 (m, 2H), 1.25-1.12 (m, 14H);	12, 13
	A	819	H-NMR (400 MHz, CD_3OD) δ 8.69-8.64 (m, 2H), 8.63-8.40 (m, 2H), 8.13-8.11 (d, *J*= 8.0 Hz, 1H), 8.01-7.98 (m, 1H), 7.75-7.73 (d, *J*=8.4Hz, 1H), 7.15-7.15 (m, 1H), 7.01-6.99 (m, 1H), 6.90-6.88 (d, *J*=9.6, 1H), 5.24-5.20 (m, 1H), 4.30 (s, 1H), 4.17-4.16 (d, *J*=4.4z, 1H), 3.79-3.32 (m, 4H), 3.16-3.12 (m, 2H), 2.97-2.93 (m, 2H), 2.89-2.87 (d, *J*=6 Hz, 1H), 2.81-2.77 (m, 6H), 2.21-2.19 (m, 1H), 1.30 (s, 6H), 1.24 (s, 6H)	4




	A	833.35	H-NMR (400 MHz, CD_3OD) δ8.62-8.60 (m, 2H), 8.41-8.35 (m, 2H), 8.13-8.11 (m, 1H), 8.02-7.99 (m, 1H), 7.71-7.71 (m, 1H), 7.14 (s, 1H), 7.01-6.94 (m, 1H), 6.85-6.83 (m, 1H), 5.25-5.15 (m, 1H), 4.29 (s, 1H), 4.12 (s, 1H), 3.80-3.75 (m, 4H), 2.94-2.72 (m, 11H), 2.20-2.02 (m, 3H), 1.28 (s, 6H), 1.22 (s, 6H);	7
	A	808.85	^{1}HNMR (300 MHz, CD_3OD): δ 8.86 (s, 1H), 8.16-8.09 (m, 1H), 7.72-7.64 (m, 2H), 7.47-7.33 (m, 2H), 7.24-7.17 (m, 1H), 7.12 (s, 1H), 6.98-6.89 (m , 1H), 5.09-5.02 (m, 1H), 4.26 (s, 1H), 4.15(s, 1H), 3.48-3.42 (m, 4H), 2.92-2.49 (m, 9H), 2.46-2.35 (m, 2H), 2.16-2.01 (m, 1H), 1.81-1.71 (m, 2H),1.59-1.47 (m, 2H), 1.42-1.36 (m, 4H), 1.27-1.18 (m, 12H);	11



	A	826.4	. ^{1}H NMR (300 MHz, DMSO): δ11.12 (s, 1H), 8.91 (s, 1H),8.18--7.88 (m,3H), 7.76--7.69 (d, J=7.8Hz,1H), 7.49--7.35(m, 2H), 7.22(s, 1H), 7.16--6.96 (m , 1H),5.17--5.06 (m, 1H), 4.32 (s, 1H), 4.11--3.98(m, 1H), 3.37--3.18(m, 4H), 2.97--2.75 (m, 3H), 2.62--2.51 (m, 4H), 2.48--2.37 (m, 2H),2.36--2.28(m,2H),2.12--1.07(m, 1H), 1.78--1.63 (m, 2H), 1.49--1.29 (m, 6H), 1.22 (s, 6H),1.13(s,6H	11
	A	782.28	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.99-2.03 (m, 3H), 2.54-2.62 (m, 6H), 2.85-2.94 (m, 1H), 3.63 (br, 4H), 4.06 (d, *J* = 5.2 Hz ,1H), 4.24-4.31 (m, 3H), 5.10-5.15 (m, 1H), 6.88 (d, *J* = 9.2 Hz ,1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.0 Hz 1H), 7.36-7.38 (m, 1H), 7.45 (d, *J* = 1.6 Hz ,1H), 7.64 (d, *J* = 9.2 Hz,1H), 7.85 (d, *J* = 8.4 Hz ,1H), 7.91 (d, *J* = 8.4 Hz ,1H), 7.98 (d, *J* = 2.4 Hz ,1H), 8.64 (d, *J* = 1.6 Hz ,1H), 11.13 (s, 1H).	45
	A	809.4	^{1}HNMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.19 (s, 6H), 1.46-1.55 (m, 2H), 1.63-1.73 (m, 2H), 1.78-1.86 (m, 2H), 1.88-2.07 (m, 5H), 2.59-2.84 (m, 7H), 2.90-2.95 (m, 1H), 3.31-3.41 (m, 2H), 4.08-4.12 (m, 2H), 4.19 (s, 1H), 4.52 (br, 1H), 4.99-5.03 (m, 1H), 6.89 (d, *J* = 8.8 Hz, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.22 (d, *J* = 8.4 Hz, 1H), 7.31 (d, *J* = 2.0 Hz, 1H), 7.37 (d, *J* = 8.0 Hz, 1H), 7.63 (d, *J* = 8.8 Hz, 1H), 7.71 (d, *J* = 8.4 Hz,	45




			1H), 8.06 (d, *J* = 8.0 Hz, 1H), 8.80-8.81 (m, 1H).	
	A	781.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.13(s, 6H), 1.19 (s, 6H), 1.86-1.89 (m, 4H), 2.01-2.05(m, 3H), 2.20-2.26 (m, 2H), 2.60-2.68 (m, 4H), 2.76-2.82 (m, 2H), 3.12 (d, *J* = 11.9 Hz, 2H), 4.07-4.18 (m, 4H), 4.99-5.03 (m,1H), 6.88 (d, *J* = 8.8 Hz, 1H), 7.03 (d, *J* = 2.4 Hz ,1H), 7.23 (d, *J* = 8.4 Hz, 1H), 7.32-7.37 (m, 2H), 7.62 (d, *J* = 8.8 Hz ,1H), 7.71 (d, *J* = 8.0 Hz ,1H), 8.05 (d, *J* = 8.0 Hz ,1H), 8.78 (d, *J* = 2.0 Hz ,1H).	45
	A	844.32		45



	A	796.4	^{1}H NMR (400 MHz, CD_3OD): δ 0.88-0.93 (m, 2H), 1.24 (s, 6H), 1.30 (s, 6H), 1.78-1.96 (m, 4H), 2.02-2.18 (m, 2H), 2.63 (t, *J* = 7.2 Hz, 2H), 2.68-2.75 (m, 4H), 2.78-2.82 (m, 1H), 2.83-2.92 (m, 1H), 3.26-3.30 (m, 1H), 3.72-3.77 (m, 4H), 4.16 (s, 1H), 4.24 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 5.10-5.16 (m, 1H), 6.87 (d, *J* = 8.8 Hz, 1H), 6.98-7.02 (m, 1H), 7.15 (d, *J* = 2.4 Hz, 1H),7.32-7.36 (m, 1H), 7.44 (d, *J* = 8.8 Hz, 1H), 7.83 (d, *J* = 8.8 Hz, 1H), 7.97-8.01 (m, 1H),8.63 (d, *J* = 2.4 Hz, 1H).	47
	A	839.45	H-NMR-PH-ARV-LS-024-F-0: (400 MHz, CD_3OD) δ 7.79 - 7.71 (m, 4H), 7.40 (s, 1H), 7.33 -7.31 (d, *J* = 8Hz, 1H), 7.13 (s, 1H), 7.04 - 6.96 (m, 3H), 5.11 - 5.08 (m, 1H), 4.28 (s, 1H), 4.21 - 4.18 (m, 2H), 4.14 (s, 1H), 3.81 - 3.67 (m, 4H), 3.30 - 3.22 (m, 1H), 3.13 - 3.09 (m, 3H), 2.87 - 2.73 (m, 6H), 2.15 - 2.08 (m, 1H), 1.93 - 1.90 (m, 2H), 1.80 - 1.58 (m, 4H), 1.28 (s, 6H), 1.22 (s, 6H).	14, 15
	A	806.25	^{1}H NMR (400 MHz, CDCl3): δ8.93 -- 8.81 (m, 1H), 8.40 (s, 1H),8.26 (s,2H), 8.18--8.12 (m, 1H), 7.82--7.69(m, 1H), 7.61--7.48 (m, 1H), 6.98 (s , 1H),6.89--6.78 (m,3H), 6.36--6.25 (m, 1H), 5.24--5.15(m, 1H), 4.96--4.82(m, 1H), 4.49--4.18(m, 3H),4.12--3.96(m,3H),3.31(s,1H),2.97--2.77(m,5H),2.43--2.15(m,2H),2.07--1.68(m,4H),1.31(s,6H),1.25(s,6H)	2




	A	842.85	^{1}H NMR (300 MHz, CD_3OD): δ 8.85 (s, 1H), 8.17-8.08 (m, 1H), 7.93-7.88 (m, 1H), 7.71-7.66 (m, 1H), 7.43-7.29 (m, 3H), 7.26-7.17 (m, 2H), 5.11-5.02 (m, 1H),4.32 (s, 1H), 4.15 (s, 1H), 3.49-3.43 (m, 4H), 2.92-2.58 (m, 9H), 2.46-2.37 (m, 2H), 2.13-2.02 (m, 1H), 1.81-1.72 (m, 2H), 1.58-1.49 (m, 2H), 1.41-1.33 (m, 4H), 1.27-1.18 (m, 12H);	11
	A	752.3	^{1}H NMR (300 MHz, CD_3OD): δ 8.89 (s, 1H), 8.15 (dd, J = 8.1, 2.3 Hz, 1H), 7.85 (d, J = 8.8 Hz, 1H), 7.70 (d, J = 8.7 Hz, 1H), 7.46 (d, J = 8.1 Hz, 1H), 7.23-7.15 (m, 1H), 7.15- 7.05 (m, 2H)，6.95 (dd, J = 8.8, 2.4 Hz, 1H), 5.48 (s, 2H), 4.91 - 4.81 (m, 1H), 4.26 (d, J = 0.9 Hz, 1H), 4.16 (d, J = 0.9 Hz, 1H), 3.60 (t, J = 4.9 Hz, 4H), 3.21 (t, J = 5.0 Hz, 4H), 3.01 (dt, J = 17.9, 7.4 Hz, 4H), 2.80-2.69 (m, 2H), 2.29-2.03 (m, 5.8 Hz, 4H), 1.30 (s, 6H), 1.21 (s, 6H).	28



	A	766.25	^{1}H NMR (300 MHz, CD_3OD): δ 8.92 (m, 1H), 8.14 (dd, J = 8.1, 2.3 Hz, 1H), 7.70 (dd, J = 8.6, 1.9 Hz, 2H), 7.50-7.45 (m, 1H), 7.40 (s, 1H),7.27 (dd, J = 8.6, 2.3 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 6.95 (dd, J = 8.8, 2.4 Hz, 1H), 5.05 (dd, J = 12.3, 5.4 Hz, 1H), 4.26 (s, 1H), 4.15 (s, 1H), 3.59 (t, J = 5.2 Hz, 4H), 3.12- 3.00 (m, 4H), 3.00-2.82 (m, 4H), 2.82-2.62 (m, 3H), 2.21 -2.02 (m, 3H), 1.24 (m, 6H) , 1.20 (s, 6H);	1
	A	765.8	^{1}HNMR (400 MHz, CD_3OD): δ 1.25 (s, 6H), 1.32 (s, 6H), 1.86-1.88 (m, 4H), 2.01-2.08 (m, 2H), 2.06-2.21 (m, 3H), 2.46-2.57 (m, 1H), 2.77-2.83 (m, 1H), 2.89-3.00 (m, 3H), 3.05-3.24 (m, 5H), 3.67-3.70 (m, 2H), 4.20 (s, 1H), 4.31 (s, 1H), 4.46-4.57 (m, 2H), 5.15-5.17 (m, 1H), 6.99-7.01 (m, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.46-7.54 (m, 3H), 7.75 (d, *J* = 8.8 Hz, 1H), 7.79 (d, *J* = 8.0 Hz, 1H), 8.17-8.19 (m, 1H), 8.92 (d, *J* = 1.6 Hz, 1H).	27
	A	768.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.18 (s, 6H), 1.91-2.05 (m, 2H), 2. 61-2.68 (m, 5H), 2.72-2.78 (m, 1H), 2.84 (s, 2H), 3.60-3.64 (m, 3H), 4.04 (s, 1H), 4.18 (s, 1H), 4.26 (s, 2H), 4.47-4.51 (m, 1H), 4.99-5.04 (m, 1H), 6.72-6.78 (m, 1H), 6.86-6.91 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.25-7.30 (m, 1H), 7.37 (d, *J* = 2.0 Hz, 1H), 7.61 (s, 1H), 7.70-7.75 (m, 1H), 7.84-7.89 (m, 1H), 8.48-8.54 (m, 1H).	47



	A	779.35	(400 MHz, CDCl3) δ 8.94-8.94 (d, J=2.0 Hz, 1H), 8.07-8.04 (m, 1H), 7.86-7.78 (m, 3H), 7.68-7.66 (d, J=7.6 Hz, 1H), 7.61-7.59 (d, J=8.8 Hz, 1H), 7.31-7.29 (d, J=10.4 Hz, 1H), 7.00-6.99 (d, J=2.0 Hz, 1H), 6.85-6.82 (m, 1H), 6.36-6.34 (d, J=8.4 Hz, 1H), 5.02-4.98 (m, 1H), 4.20-4.18 (d, J=8.0 Hz, 1H), 4.10 (s, 1H), 3.41-3.38 (d, J=10.8 Hz, 2H), 2.96-2.74 (m, 8H), 2.43-2.40 (d, J=11.2 Hz, 2H), 2.20-2.15 (m, 3H), 1.98-1.95 (d, J=12.8 Hz, 2H), 1.88-1.77 (m, 4H), 1.31 (s, 6H), 1.26 (s, 6H)	1
	A	780.4	H-NMR (400 MHz, CD3OD) δ 8.91-8.90 (d, J=2.0 Hz, 1H), 8.19-8.16 (m, 3H), 7.76-7.73 (m, 2H), 7.49-7.43 (m, 2H), 7.31-7.29 (m, 1H), 7.16-7.15 (d, J=2.4 Hz, 1H), 7.02-6.99 (m, 1H), 5.12-5.08 (m, 1H), 4.31 (s, 1H), 4.21 (s, 1H), 3.61-3.58 (m, 4H), 2.98-2.72 (m, 11H), 2.14-2.13 (d, J=5.2 Hz, 1H), 1.88-1.84 (m, 2H), 1.75-1.72 (m, 2H), 1.32 (s, 6H), 1.26 (s, 2H);	1




	A	766.7	H-NMR (400 MHz, CD3OD) δ 8.91-8.91 (d, J=1.6 Hz, 1H), 8.34-8.17 (m, 3H), 7.86-7.84 (d, J=8.8 Hz, 1H), 7.76-7.73 (d, J=8.8 Hz, 1H), 7.49-7.47 (d, J=8.0 Hz, 1H), 7.18-7.15 (m, 2H), 7.09 (s, 1H), 7.02-6.99 (m, 1H), 5.44 (s, 2H), 4.85-4.81 (m, 1H), 4.31 (s, 1H), 4.21 (s, 1H), 3.62-3.58 (d, J=13.2 Hz, 4H), 3.17 (s, 4H), 3.01-2.95 (m, 4H), 2.79-2.76 (m, 2H), 2.26-2.14 (m, 2H), 1.87-1.80 (m, 4H), 1.32-1.25 (d, J=26 Hz, 12H);	1, 16
	A	790.05		45
	A			45



	A	781.4	^{1}HNMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.27-1.43 (m, 5H), 1.65-1.71 (m, 2H), 2.01-2.07 (m, 1H), 2.53-2.60 (m, 4H), 2.75-2.79 (m, 3H), 2.83-2.94 (m, 2H), 4.05-4.07 (m, 2H), 4.30 (s, 1H), 5.08-5.16 (m, 2H), 6.98-7.01 (m, 1H), 7.19-7.22 (m, 1H), 7.28-7.37 (m, 3H), 7.84-7.90 (m, 2H), 7.99-8.08 (m, 2H), 8.88 (s, 1H), 11.11 (s, 1H).	83
	A	821.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.04-1.06 (m, 1H), 1.10 (s, 6H), 1.20 (s, 6H), 1.75-1.85 (m, 3H), 1.98-2.02 (m, 1H), 2.18 (d, *J* = 6.4 Hz, 2H), 2.51-2.60 (m, 2H), 2.82-2.92 (m, 4H), 3.30-3.32 (m, 4H), 3.42-3.44 (m, 4H), 4.03 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1H), 4.37-4.43 (m, 2H), 5.03-5.08 (m, 1H), 6.82 (s, 1H), 6.97-7.00 (m, 1H), 7.19 (d, *J* = 2.0 Hz, 1H), 7.22-7.26 (m, 1H), 7.33 (s, 1H), 7.57 (d, *J* = 9.2 Hz, 1H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.87-7.93 (m, 2H), 8.60 (d, *J* = 2.4 Hz, 1H), 11.07 (s, 1H).	49




	A	795.5	^{1}H NMR (400 MHz, CD_3OD): δ 1.23 (s, 6H), 1.30 (s, 6H), 1.38-1.38 (m, 1H), 1.64-1.69 (m, 2H), 1.77-1.84 (m, 2H), 1.93-1.95 (m, 2H), 2.11-2.15 (m, 3H), 2.68-2.77 (m, 5H), 2.83-2.98 (m, 5H), 4.18 (s, 1H), 4.29 (s, 1H), 4.74 (s, 1H), 5.09, 5.12 (d, *J* = 5.6 Hz, 1H), 6.97-6.99 (m, 1H), 7.13-7.14 (m, 1H), 7.33-7.35 (m, 1H), 7.43-7.47 (m, 2H), 7.71-7.74 (m, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 8.14-8.16 (m, 1H), 8.87 (d, *J* = 2.0 Hz, 1H).	83
	A	835.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.43-1.44 (m, 2H), 1.62-1.63 (m, 1H), 1.75-1.78 (m, 2H), 2.01-2.03 (m, 1H),2.35-2.43 (m, 3H), 2.59-2.64 (m, 2H), 2.85-2.89 (m, 4H), 3.17-3.45 (m, 8H), 4.05-4.07 (m, 1H), 4.31 (s, 1H), 4.39-4.42 (m, 2H), 5.06-5.10 (m, 1H), 6.85 (d, *J* = 8.8 Hz, 1H), 7.00 (d, *J* = 8.4 Hz, 1H), 7.21-7.27 (m, 2H), 7.35 (s, 1H), 7.59-7.70 (m, 2H), 7.90-7.95 (m, 2H), 8.62 (s, 1H), 11.09 (s, 1H).	23
	A	811.4	^{1}H-NMR (300 MHz, CD3OD) δ8.78 (s, 2H), 7.85-7.79(m, 1H), 7.74- 7.68(m, 1H), 7.42- 7.39(m, 1H), 7.35 - 7.28(m, 1H), 7.14 - 7.10 (m, 1H), 7.01 - 6.96 (m, 1H), 5.15- 5.06(m, 1H), 4.29- 4.03(m, 8H), 2.95 - 2.69(m, 9H), 2.17 - 2.12 (m, 1H), 1.96 - 1.87 (m, 2H), 1.81 - 1.72 (m, 2H), 1.64 - 1.54 (m, 2H), 1.28 - 1.21 (d, *J = 20.4Hz* , 12H).	9



	A	845.3	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.23 (s, 6H), 1.27 (s, 6H), 1.52-1.69 (m, 2H), 1.70-1.86 (m, 2H), 2.12-2.14 (m, 1H), 2.30-2.51 (m, 2H), 2.52-2.66 (m, 3H), 2.72-2.85 (m, 2H), 2.87-2.92 (m, 1H), 3.00 (t, *J* = 6.8 Hz, 2H), 3.31-3.46 (m, 3H), 3.51 (t, *J* = 6.0 Hz, 2H), 3.65 (t, *J* = 6.4 Hz, 2H), 4.07 (s, 1H), 4.13 (d, *J* = 8.0 Hz, 1H), 4.92-4.96 (m, 1H), 6.01-6.21 (m, 1H), 6.80 (dd, *J* = 2.4, 8.8 Hz, 1H), 6.96 (d, *J* = 2.0 Hz, 1H), 7.06 (dd, *J* = 2.0, 8.4 Hz, 1H), 7.28-7.33 (m, 2H), 7.57 (d, *J* = 8.4 Hz, 1H), 7.70 (d, *J* = 8.4 Hz, 1H), 8.02 (s, 1H).	84
	A	837.3	^{1}H NMR (300 MHz, CD_3OD): δ11.08 (s, 1H), 7.84 (dd, J = 21.4, 8.5 Hz, 2H), 7.71 (d, J = 8.6 Hz, 2H), 7.50-7.39 (m, 2H), 7.33 (dd, J = 8.3, 2.3 Hz, 1H), 7.18 (d, J = 2.4 Hz, 1H), 7.03-6.88 (m, 3H), 5.09 (dd, J = 12.9, 5.4 Hz, 1H), 4.30 (s, 1H), 4.16 (t, J = 6.5 Hz, 2H), 4.02 (d, J = 9.1 Hz, 1H), 3.67 (d, J = 11.5 Hz, 2H), 2.92-2.78 (m, 1H), 2.72-2.50 (m, 5H), 2.42 (s, 1H), 2.34 (d, J = 27.9 Hz, 2H), 2.03 (dq, J = 13.5, 5.6, 4.6 Hz, 1H), 1.76 (dd, J = 8.1, 5.2 Hz, 2H), 1.39 (d, J = 12.5 Hz, 4H),	21




			1.19 (s, 6H), 1.14-0.95 (m, 12H);	
	A	819.5	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.13 (s, 6H), 1.23 (s, 6H), 1.73-1.80 (m, 2H), 1.93-2.00 (m, 2H), 2.00-2.07 (m, 1H), 2.53-2.67 (m, 2H), 2.85-2.93 (m, 1H), 4.06-4.08 (m, 2H), 4.17-4.22 (m, 4H), 4.31 (s, 1H), 4.51 (br, 2H), 4.73 (s, 2H), 5.09-5.14 (m, 1H), 6.60-6.62 (m, 1H), 7.00-7.03 (m, 1H), 7.21-7.22 (m, 1H), 7.33-7.35 (m, 1H), 7.42-7.43 (m, 1H), 7.65-7.67 (m, 1H), 7.81-7.83 (m, 2H), 7.90-7.92 (m, 1H), 8.03-8.05 (m, 1H), 8.65-8.66 (m, 1H),11.12 (s, 1H).	55
	A	819.53	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.13 (s, 6H), 1.23 (s, 6H), 1.73-1.80 (m, 2H), 1.93-2.00 (m, 2H), 2.00-2.07 (m, 1H), 2.53-2.67 (m, 2H), 2.85-2.93 (m, 1H), 4.06-4.08 (m, 2H), 4.17-4.22 (m, 4H), 4.31 (s, 1H), 4.51 (br, 2H), 4.73 (s, 2H), 5.09-5.14 (m, 1H), 6.60-6.62 (m, 1H), 7.00-7.03 (m, 1H), 7.21-7.22 (m, 1H), 7.33-7.35 (m, 1H), 7.42-7.43 (m, 1H), 7.65-7.67 (m, 1H), 7.81-7.83 (m, 2H), 7.90-7.92 (m, 1H), 8.03-8.05 (m, 1H), 8.65-8.66 (m, 1H),11.12 (s, 1H).	85



	A	819.27	^{1}HNMR (400 MHz, CD_3OD): δ 1.14 (s, 6H), 1.21 (s, 6H), 1.72-2.04 (m, 6H), 2.60-2.75 (m, 4H), 2.88 (t, *J* = 7 Hz, 2H), 3.82 (s, 2H), 4.09-4.27 (m, 6H), 4.98-5.03 (m, 1H), 6.89 (d, *J* = 6.8 Hz, 1H), 7.04 (s, 1H), 7.25 (d, *J* = 8.8 Hz, 1H), 7.33 (s, 1H), 7.45 (s, 1H), 7.62 (d, *J* = 4 Hz, 1H), 7.70 (d, *J* = 8 Hz, 1H), 7.88 (d, *J* = 8.4 Hz, 1H), 8.21 (d, *J* = 8.8 Hz, 1H), 8.74 (s, 1H).	86
	A	794.45	^{1}H NMR (300 MHz, DMSO-d6): *δ* 11.04 (s, 1H), 8.86 (d, J = 2.2 Hz, 1H), 8.10-8.00 (m, 2H), 8.00-7.80 (m, 1H),7.64 (d, J = 8.4 Hz, 1H), 7.39-7.14 (m, 4H), 6.97 (dd, J = 8.7, 2.4 Hz, 1H), 5.03 (dd, J = 12.8, 5.3 Hz, 1H), 4.28 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.38 (t, J = 5.0 Hz, 4H), 3.00-2.73 (m, 3H), 2.73-2.51 (m, 2H), 2.51-2.39 (m, 4H), 2.27 (t, J = 7.3 Hz, 2H), 2.02-1.93 (m, 1H), 1.69-1.63 (m, 2H), 1.47 (p, J = 7.4 Hz, 2H), 1.30 (q, J = 8.0 Hz, 2H), 1.20 (s, 6H), 1.09 (s, 6H)	8
	A	812.55	^{1}H NMR (400 MHz, CDCl3) 8.845 (s, 1H), 7.97-7.91 (m, 2H), 7.51-7.49 (m, 1H), 7.08-7.05 (m, 1H), 6.90-6.89 (m, 1H), 6.75-6.72 (m, 1H), 6.19-6.17 (m, 1H), 5.13-5.09 (m, 1H), 4.36-4.32 (m, 1H), 4.22-4.18 (m, 1H), 4.10-4.08 (m, 1H), 4.00-3.98 (m, 1H), 3.17-3.10 (m, 4H), 2.83-2.62 (m, 7H), 2.60-2.09 (m, 4H), 2.00-1.76 (m, 2H), 1.75-1.68 (m, 3H), 1.32-1.16 (m, 16H)	20




	A	812.5	^{1}H NMR (400 MHz, CDCl3) 8.85 (s, 1H), 7.97-7.88 (m, 2H), 7.51-7.42 (m, 2H), 6.93-6.89 (m, 2H), 6.75-6.72 (m, 1H), 6.18-6.15 (m, 1H), 5.12-5.08 (m, 1H), 4.36-4.00 (m, 4H), 3.32-3.20 (m, 4H), 2.87-2.59 (m, 7H), 2.30-2.13 (m, 2H), 1.79-1.50 (m, 7H), 1.41-1.16 (m, 16H)	11, 16
	A	821.35	**^{1}H NMR (400 MHz, CD_3OD): δ 7.77-7.50 (m, 5H), 7.38-7.34 (m, 3H), 7.23-7.13 (m, 2H), 7.00-6.96 (m, 1H), 5.10-5.00 (m, 1H), 4.29 (s, 1H), 4.17-4.14 (m, 1H), 3.76-3.67 (m, 4H), 3.36-3.32 (m, 2H), 2.87-2.42 (m, 11H), 2.10-2.00 (m, 3H), 1.70-1.60 (m, 2H), 1.28-1.23 (m, 12H)**	39



	A	797.4	1H NMR (300 MHz, DMSO-d6) δ 11.08 (s, 1H), 8.73 (s, 2H), 7.83 (dd, J = 18.2, 8.5 Hz, 2H), 7.69 (d, J = 9.2 Hz, 1H), 7.41 (d, J = 2.2 Hz, 1H), 7.33 (dd, J = 8.3, 2.3 Hz, 1H), 7.18 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.5 Hz, 1H), 5.08 (dd, J = 12.8, 5.4 Hz, 1H), 4.29 – 4.13 (m, 3H), 4.00 (d, J = 9.1 Hz, 1H), 3.79 (t, J = 4.7 Hz, 4H), 2.95 – 2.76 (m, 1H), 2.56 (d, J = 16.4 Hz, 2H), 2.45 – 2.30 (m, 6H), 2.00 (dd, J = 12.3, 6.2 Hz, 1H), 1.69 (dq, J = 46.8, 7.0 Hz, 4H), 1.18 (s, 6H), 1.08 (s, 6H).	9
	A	783.55	1H NMR (300 MHz, DMSO-d6) δ 11.08 (s, 1H), 8.75 (s, 2H), 8.21-7.63 (m, 3H), 7.60- 7.16 (m, 2H), 7.18 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.5 Hz, 1H), 5.09 (dd, J = 12.8, 5.3 Hz, 1H), 4.25 (d, J = 7.4 Hz, 3H), 4.01 (d, J = 9.1 Hz, 1H), 3.28 (s, 4H), 2.89-2.76 (m, 1H), 2.56 (d, J = 19.0 Hz, 2H), 2.24 (s, 6H), 1.99 (s, 3H), 1.13 (d, J = 30.8 Hz, 12H)	9
	A	769.3	1H NMR (300 MHz, DMSO-d6) δ 11.09 (s, 1H), 10.18 (s, 1H), 8.83 (s, 2H), 7.93-7.74 (m, 3H), 7.54 (d, J = 2.2 Hz, 1H), 7.42 (dd, J = 8.3, 2.3 Hz, 1H), 7.27-7.14 (m, 1H), 7.03-6.86 (m, 1H), 5.10 (dd, J = 12.8, 5.4 Hz, 1H), 4.88-4.47 (m, 4H), 4.27 (s, 1H), 4.02 (d, J = 9.1 Hz, 1H),3.92- 3.62 (s, 4H), 3.30 (d, J = 54.5 Hz, 4H), 2.92-2.77 (m, 1H), 2.57 (d, J = 19.1 Hz, 2H), 2.08-	9




			1.97 (m, 1H), 1.18 (s, 6H), 1.08 (s, 6H).	
	A	822.57	^{1}H NMR (300 MHz, CD_3OD): δ 8.70 (s, 2H), 7.71-7.66 (m, 2H), 7.36 (m, 1H), 7.24-7.21 (m, 1H), 7.10-7.09 (m, 1H), 6.97-6.93 (m, 1H), 5.08-5.02 (m, 1H), 4.90-4.87 (m, 2H),4.80 (s, 1H), 4.28-4.24 (m, 2H), 4.10 (s, 1H), 3.51 (m, 4H), 3.03-2.95 (m, 2H), 2.78-2.67 (m, 6H), 2.44-2.42 (m, 2H), 2.15-1.80 (m, 4H), 1.75-1.55 (m, 1H), 1.25-1.18 (m, 12H);	49
	A	800.21	^{1}H NMR (400 MHz, $CDCl_3$): δ 1.25 (s, 6H), 1.30 (s, 6H), 1.98-2.07 (m, 2H), 2.11-2.16 (m, 1H), 2.46-2.51(m, 2H), 2.57-2.66 (m, 3H), 2.70-2.87 (m, 3H), 2.88-2.96 (m, 3H), 3.40-3.49 (m, 4H), 4.11 (s, 1H), 4.18 (d, *J* = 8.0 Hz, 1H), 4.92-4.97 (m, 1H), 6.20 (d, *J* = 8.0 Hz, 1H), 7.03-7.08 (m, 2H),7.24 (d, *J* = 2.4 Hz, 1H), 7.28-7.32 (m, 2H), 7.70 (d, *J* = 8.4 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.98-8.11 (m, 2H), 8.92 (d, *J* = 2.0 Hz, 1H).	1



	A	781.51	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.42-1.58 (m, 4H), 1.98-2.04 (m,1H), 2.26-2.34 (m, 2H), 2.40-2.48 (m, 6H), 2.62-2.70 (m, 3H), 2.82-2.94 (m, 1H), 3.36-3.52 (m, 7H), 4.24-4.50 (m, 5H), 5.02-5.10 (m, 1H), 5.76 (s, 1H), 6.53 (s, 1H), 6.93 (s, 1H), 7.23 (d, *J* = 8.4 Hz, 1H), 7.31 (s, 1H), 7.67 (d, *J* = 8.8 Hz, 1H), 7.98 (s, 2H), 8.07 (s, 1H), 8.26-8.34 (m, 1H), 11.08 (s, 1H), 13.28 (s, 1H).	17a
	A	835.8	^{1}H NMR (300 MHz, CD3OD) 7.76-7.64 (m, 4H), 7.35-7.32 (m, 3H), 7.23-7.20 (m, 1H), 7.13-7.12 (m, 1H), 7.00-6.96(m, 1H), 5.09-5.03 (m, 1H), 4.28-4.25 (m, 1H), 4.15-4.11 (m, 1H), 3.77-3.68 (m, 4H), 3.38-3.32 (m, 2H), 2.86-2.69 (m, 5H), 2.41-2.29 (m, 6H), 2.11-2.03 (m, 3H), 1.86-1.64 (m, 4H), 1.28-1.23 (m, 12H);	39
	A	801.5	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.07 (s, 1H), 7.91 (d, *J* = 8.8 Hz, 1H), 7.71 (dd, *J* = 19.0, 8.4 Hz, 3H), 7.46 (d, *J* = 9.2 Hz, 1H), 7.35 (d, *J* = 2.4 Hz, 1H), 7.27 (dd, *J* = 8.7, 2.3 Hz, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.01 (dd, *J* = 8.8, 2.4 Hz, 1H), 6.44 (d, *J* = 8.4 Hz, 2H), 5.08 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.32 (s, 1H), 4.06 (d, *J* = 9.1 Hz, 1H), 3.91 (t, *J* = 7.3 Hz, 2H), 3.64 (ddd, *J* = 24.3, 9.7, 5.1 Hz, 3H), 3.43 (dt, *J* = 18.6, 4.7 Hz, 6H), 3.03 – 2.82 (m, 4H), 2.64 – 2.51 (m, 2H), 2.40 (t, *J* = 4.8 Hz,	37



			4H), 2.02 (dt, *J* = 15.2, 5.6 Hz, 1H), 1.22 (s, 6H), 1.13 (s, 6H).	
	A	836.6	^{1}H-NMR (300 MHz, CDCl3) δ8.69 (s, 2H), 8.01(m, 1H), 7.75- 7.69(m, 1H), 7.62- 7.53(m, 1H), 7.29(m, 1H), 7.10 - 7.04(m, 1H), 6.98 - 6.91 (m, 1H), 6.84 - 6.78 (m, 1H), 5.95- 5.88(m, 1H), 4.98- 4.81(m, 3H), 4.15- 4.01(m, 2H), 3.54(m, 4H), 3.11 - 2.60(m, 11H), 2.17 - 2.09 (m, 1H), 1.87 - 1.61(m, 5H), 1.24 - 1.21 (m, 14H)	32
	A	810.23		14a




	A	815.32		17
	A	869.28		17
	A	824.21	^{1}HNMR (400 MHz, DMSO-d_6) : δ 1.12 (s, 6H), 1.22 (s, 6H), 1.39-1.44 (m, 2H), 1.55-1.61 (m, 2H), 1.75-1.81 (m, 2H), 2.02-2.06 (m, 1H), 2.54-2.69 (m, 2H), 2.85-2.94 (m, 1H), 3.39 (t, *J* = 7.2 Hz, 2H), 3.46 (t, *J* = 5.0 Hz, 2H), 3.88 (t, *J* = 5.2 Hz, 2H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.16-4.19 (m, 4H), 4.31 (s, 1H), 5.10-5.14 (m, 1H), 6.87 (d, *J* = 9.2 Hz, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.32-7.35 (m, 1H), 7.43 (s, 1H), 7.66-7.68 (m, 1H), 7.82 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J*	87



			= 8.8 Hz, 1H), 8.01-8.03 (m, 1H), 8.66 (d, *J* = 2.0 Hz, 1H), 11.12 (m, 1H),.	
	A	852.25	^{1}H NMR (400 MHz, CD_3OD): δ 1.17 (s, 6H), 1.24 (s, 6H), 1.43-1.46 (m, 2H), 1.56-1.59 (m, 2H), 2.07-2.10 (m, 1H), 2.42 (t, *J* = 7.2 Hz, 2H), 2.54 (t, *J* = 4.0 Hz, 4H), 2.62-2.73 (m, 4H), 2.79-2.84 (m, 1H), 3.32-3.35 (m, 2H), 3.86-3.91 (m, 6H), 4.10 (s, 1H), 4.23 (s, 1H), 5.01 (t, *J* = 5.2 Hz, 1H), 5.05-5.10 (m, 1H), 6.93-6.95 (m, 1H), 7.09 (d, *J* = 20.0 Hz, 1H), 7.20-7.25 (m, 2H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.78 (d, *J* = 8.4 Hz, 1H), 8.71 (s, 2H).	73
	A	838.23	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.11 (s, 6H), 1.21 (s, 6H), 1.49-1.60 (m, 2H), 2.01-2.08 (m, 1H), 2.29-2.49 (m, 7H), 2.53-2.64 (m, 3H), 2.85-2.94 (m, 1H), 3.08-3.26 (m, 2H), 3.85 (s, 6H), 4.04 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1H), 5.06-5.16 (m, 2H), 6.99-7.02 (m, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.26-7.32 (m, 2H), 7.75 (d, *J* = 9.2 Hz, 1H), 7.86 (d, *J* = 9.2 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.77 (s, 2H), 11.14 (s, 1H).	73




	A	784.1	H-NMR-PH-ARV-LS-031-M-0: (300 MHz, DMSO- *d6*, ppm) δ 11.10 (br, 1H), 8.20 -8.17 (d, *J* = 9Hz, 1H), 8.02 (s, 1H), 7.74 - 7.66 (m, 2H), 7.53 - 7.47 (m, 1H), 7.34 - 7.15 (m, 4H), 5.11 - 5.05 (m, 1H), 3.42 - 3.35 (m, 4H), 2.91 - 2.83 (m, 1H), 2.71 - 2.60 (m, 4H), 2.56 - 2.49 (m, 4H), 2.35 - 2.30 (m, 2H), 2.08 - 2.00 (m, 1H), 1.67 - 1.65 (m, 2H), 1.62 (s, 8H), 1.60 - 1.37 (m, 2H).	43
	A	820.35	^{1}H NMR (300 MHz, $CDCl_3$) δ 7.95 (s, 1H), 7.70-7.55 (m, 4H), 7.30 (s, 1H), 7.26-6.97 (m, 1H), 6.96-6.79 (m, 4H), 6.12-6.09 (m, 1H), 4.96-4.94 (m, 1H), 4.16-4.04 (m, 2H), 3.89-3.39 (m, 5H), 3.20-2.50 (m, 8H), 2.49-1.60 (m, 6H), 1.26-1.20 (m, 15H);	49
	A	796.65	H-NMR06-PH-ARV-LS-031-A-0: (300 MHz, CD3OD, ppm) δ 7.96 - 7.93 (d, *J* = 9 Hz, 1 H), 7.83 (s, 1 H), 7.69 - 7.61 (m, 2 H), 7.51 - 7.48 (m, 1 H), 7.35 (s, 1 H), 7.24 – 7.15 (m, 3 H), 5.09 - 5.05 (m, 1 H), 3.48 - 3.45 (m, 4H), 2.82 - 2.42(m, 15 H), 2.12 - 2.09 (m, 2 H), 1.76 - 1.73 (m, 2 H), 1.66 - 1.61 (m, 3 H), 1.48 - 1.46 (m, 2 H).	43



	A	821.35	^{1}H NMR (300 MHz, DMSO-d_6) δ 10.70 (s, 1H), 8.62 (d, *J* = 2.4 Hz, 1H), 7.92 (dd, *J* = 10.0, 6.5 Hz, 2H), 7.56 (dd, *J* = 17.5, 8.9 Hz, 2H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.03 (ddd, *J* = 14.6, 8.6, 2.1 Hz, 3H), 6.86 (d, *J* = 9.1 Hz, 1H), 5.31 (s, 2H), 4.57 (dd, *J* = 9.1, 5.2 Hz, 1H), 4.41 (d, *J* = 13.3 Hz, 2H), 4.31 (s, 1H), 4.06 (d, *J* = 9.2 Hz, 1H), 3.29 (t, *J* = 4.9 Hz, 4H), 2.96 – 2.81 (m, 2H), 2.61 (t, *J* = 5.6 Hz, 2H), 2.54 – 2.50 (m, 4H), 2.38 (t, *J* = 7.4 Hz, 2H), 2.00 (qd, *J* = 8.2, 3.9 Hz, 2H), 1.77 (d, *J* = 12.7 Hz, 2H), 1.63 (s, 1H), 1.43 (q, *J* = 7.2 Hz, 2H), 1.22 (s, 6H), 1.12 (s, 8H).	41
	A	827.45	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.04 (s, 1H), 8.71 (s, 2H), 8.02 (d, *J* = 8.8 Hz, 1H), 7.71 – 7.57 (m, 3H), 7.36 – 7.17 (m, 3H), 5.03 (dd, *J* = 12.7, 5.3 Hz, 1H), 4.69 (d, *J* = 12.8 Hz, 2H), 4.30 (s, 1H), 4.01 (d, *J* = 9.0 Hz, 1H), 3.28 (s, 4H), 3.00 – 2.76 (m, 3H), 2.67 (dt, *J* = 3.7, 1.9 Hz, 4H), 2.41 (t, *J* = 7.3 Hz, 2H), 2.34 (t, *J* = 7.3 Hz, 2H), 2.03 – 1.92 (m, 1H), 1.81 – 1.69 (m, 3H), 1.46 – 1.33 (m, 2H), 1.19 (s, 6H), 1.07 (s, 8H)	17


	A	764.3	H-NMR (300 MHz, DMSO) δ 11.05 (s, 1H), 8.64-8.63 (d, J=2.4 Hz, 1H), 7.98-7.89 (m, 2H), 7.67-7.60 (m, 2H), 7.22-7.21 (d, J=2.4 Hz, 1H), 7.03-6.92 (m, 2H), 6.80-6.79 (d, J=1.8Hz, 1H), 6.69-6.65 (m, 1H), 5.09-5.03 (m, 1H), 4.31 (s, 1H), 4.08-4.05 (d, J=9 Hz, 1H), 3.85 (s, 4H), 3.67 (s, 4H),2.73-2.72 (m, 1H), 2.60 (s, 2H), 2.01-1.99(d, J=5.1 Hz, 1H), 1.82 (s, 4H), 1.22 (s, 6H), 1.12(s, 6H);	44
	A	764.35	H-NMR (300 MHz, DMSO) δ 11.04 (s, 1H), 8.58 (s, 1H), 7.95-7.86 (m, 2H), 7.66-7.63 (d, J=8.4Hz, 1H), 7.57-7.54 (d, J=9.3Hz, 1H), 7.35-7.18 (m, 3H), 7.00-6.96 (m, 1H), 6.39-6.36 (d, J=9Hz, 1H), 5.08-5.02 (m, 1H), 4.28 (s, 1H), 4.05-4.02 (d, J=9.3Hz, 1H), 3.80 (s, 4H), 3.51 (s, 4H), 2.92-2.81 (m, 1H), 2.70-2.59 (m, 2H), 2.05-1.98 (m, 4H), 1.19 (s, 6H), 1.10 (s, 6H);	19



	A	833.54	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.24 (s, 6H), 1.38-1.47 (m, 2H), 1.74-1.90 (m, 4H), 1.96-2.10 (m, 2H), 2.54-2.65 (m, 2H), 2.85-2.94 (m, 1H), 3.33 (s, 1H), 4.04-4.11 (m, 1H), 4.14-4.19 (m, 3H), 4.32 (s, 1H), 4.55 (s, 3H), 5.07-5.18 (m, 1H), 6.62 (d, *J* = 8.8 Hz, 1H), 6.96-7.08 (m, 1H), 7.23 (d, *J* = 2.4 Hz, 1H), 7.31-7.38 (m, 1H), 7.43 (d, *J* = 1.6 Hz, 1H), 7.66 (s, 2H), 7.79-7.86 (m, 1H), 7.89-7.95 (m, 1H), 7.98-8.05 (m, 1H), 8.69 (s, 1H), 11.08-11.19 (m, 1H).	85
	A	740.46	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.21 (s, 6H), 2.03-2.15 (m, 2H), 2.58 (m, 2H), 3.86-3.89 (m, 2H), 4.04 (d, *J* = 8.8 Hz, 1H), 4.23-4.29 (m, 5H), 5.10-5.15 (m, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.35-7.38 (m, 1H), 7.45 (d, *J* = 2.0 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.84-7.91 (m, 2H), 8.74 (s, 2H), 11.12 (s, 1H).	88
	A	831.53	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.61-1.64 (m, 2H), 1.76-1.82 (m, 2H), 2.01-2.07 (m, 1H), 2.38 (s, 2H), 2.44-2.49 (m, 4H), 2.57 (s, 1H), 2.85-2.94 (m, 1H), 3.24 (s, 4H), 3.32 (s, 1H), 4.00 (d, *J* = 9.2 Hz, 1H), 4.20-4.26 (m, 3H), 5.10-5.15 (m, 1H), 6.66 (d, *J* = 11.6 Hz, 2H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 8.4, 2.4 Hz, 1H), 7.44 (d, *J* = 2.4 Hz, 1H), 7.84 (d, *J* = 8.4	89



			Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.19 (d, *J* = 9.2 Hz, 1H), 11.13 (s, 1H).	
	A	836.5	^{1}HNMR (400 MHz, DMSO-d_6) : δ 1.11 (s, 6H), 1.22 (s, 6H), 1.73-1.83 (m, 5H), 1.99-2.02 (m, 1H), 2.29 (s, 2H), 2.57 (s, 6H), 2.74 (s, 2H), 2.91-2.97 (m, 3H), 3.82-3.86 (m, 4H), 4.01-4.03 (m, 3H), 4.29 (s, 1H), 5.06-5.09 (m, 1H), 7.01 (d, *J* = 8.0 Hz, 1H), 7.22-7.30 (m, 3H), 7.64 (d, *J* = 7.6 Hz, 1H), 7.75 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 8.77 (s, 2H), 11.10 (s, 1H),.	49
	A	837.75	H-NMR-PH-ARV-LS-032-D-0: (300 MHz, DMSO-d6,ppm) δ 11.11 (s, 1H), 7.86 (dd, *J* = 22.3, 8.5 Hz, 2H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 – 7.43 (m, 2H), 7.36 (dd, *J* = 8.3, 2.3 Hz, 1H), 7.20 (d, *J* = 2.4 Hz, 1H), 7.05 – 6.90 (m, 3H), 5.11 (dd, *J* = 13.0, 5.2 Hz, 1H), 4.31 (s, 3H), 4.04 (d, *J* = 9.1 Hz, 1H), 3.65 (s, 2H), 3.10 (d, *J* = 11.5 Hz, 2H), 2.89 (s, 1H), 2.69 (s, 5H), 2.37 (s, 3H), 2.04 (d, *J* = 11.3 Hz, 1H), 1.85 (s, 2H), 1.58 (s, 2H), 1.21 (s, 6H), 1.12 (s, 6H).	47




	A	851.25	H-NMR-PH-ARV-LS-032-F-0: (300 MHz, DMSO-d6,ppm) (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 7.84 (dd, *J* = 19.4, 8.5 Hz, 2H), 7.69 (d, *J* = 8.6 Hz, 2H), 7.49 – 7.26 (m, 3H), 7.17 (d, *J* = 2.3 Hz, 1H), 7.02 – 6.87 (m, 3H), 5.09 (dd, *J* = 12.8, 5.3 Hz, 1H), 4.31 – 4.14 (m, 3H), 4.01 (d, *J* = 9.1 Hz, 1H), 3.62 (t, *J* = 4.6 Hz, 2H), 3.44 (d, *J* = 12.6 Hz, 2H), 3.07 (t, *J* = 11.3 Hz, 2H), 2.91 – 2.76 (m, 1H), 2.47 (m, 2H), 2.42 – 2.30 (m,4H), 2.23 (s, 2H), 2.01-1.88 (m, 5H), 1.55 (dd, *J* = 12.6, 8.4 Hz, 2H), 1.18 (s, 6H), 1.09 (s, 6H).	47
	A	769.3	^{1}H NMR (300 MHz, $CDCl_3$) 8.12 (s, 1H), 7.73-7.71 (m, 2H), 7.60-7.25 (m, 5H), 6.97-6.96 (m, 1H), 6.82-6.79 (m, 1H), 6.21-6.18 (m, 1H), 4.96-4.90 (m, 1H), 4.17-4.14 (m, 1H), 4.05-4.00 (m, 1H), 3.48-3.35 (m, 4H), 2.93-2.68 (m, 10H), 2.16-2.12 (m, 1H) , 1.37-1.23 (m, 13H);	1
	A	763.6	^{1}H NMR (300 MHz, CD_3OD) δ 7.770-7.66 (m, 3H), 7.60 (d, *J* = 8.5 Hz, 1H), 7.31 (d, *J* = 8.1 Hz, 2H), 7.10 (d, *J* = 2.4 Hz, 1H), 7.04-6.89 (m, 2H), 6.84 (dd, *J* = 8.6, 2.2 Hz, 1H), 5.02 (dd, *J* = 12.3, 5.4 Hz, 1H), 4.54 (s, 1H), 4.25 (s, 1H), 4.11 (s, 1H), 3.75 (s, 1H), 3.46 (d, *J* = 2.5 Hz, 2H), 3.07-2.96 (m, 1H), 2.88-2.61 (m, 8H), 2.13-1.87 (m, 3H), 1.24 (s, 6H), 1.19 (s, 6H);	1



	A	834.56		
	A	856.56	^{1}HNMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.21-1.22 (m, 6H), 1.52-1.57 (m, 1H), 1.60-1.69 (m, 2H), 1.98-2.07 (m, 2H), 2.12-2.23 (m, 1H), 2.35-2.43 (m, 2H), 2.55-2.68 (m, 4H), 2.83-2.93 (m, 1H), 3.28-3.34 (m, 3H), 3.40 (s, 4H), 3.52-3.57 (m, 1H), 3.65-3.69 (m, 2H), 4.05-4.08 (m, 1H), 4.31 (s, 1H), 5.06-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.21-7.28 (m, 2H), 7.36 (s, 1H), 7.51-7.54 (m, 2H), 7.63-7.70 (m, 2H), 7.90-7.92 (m, 1H), 11.09 (s, 1H).	49




	A	834.55	1H NMR (300 MHz, DMSO-d6) δ 11.04 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.67 (dd, J = 15.3, 8.5 Hz, 3H), 7.44 (d, J = 9.2 Hz, 1H), 7.30 (d, J = 2.4 Hz, 1H), 7.27-7.13 (m, 2H), 7.02-6.87 (m, 3H), 5.03 (dd, J = 12.7, 5.3 Hz, 1H), 4.28 (s, 1H), 4.01 (d, J = 9.1 Hz, 1H), 3.81 (d, J = 12.4 Hz, 2H), 3.28 (s, 5H), 2.93-2.69 (m, 4H), 3.0-2.6 (m, 4H), 2.54 (d, J = 16.1 Hz, 5H), 2.35 (t, J = 7.2 Hz, 2H), 2.03-1.93 (m, 1H), 1.73 (d, J = 12.8 Hz, 2H), 1.40 (d, J = 7.3 Hz, 3H), 1.18 (s, 8H), 1.09 (s, 6H).	23
	A	832.3	^{1}H NMR (400 MHz, CD_3OD) δ 7.74 (dd, *J* = 8.8, 2.7 Hz, 3H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.07-7.04 (m, 1H), 7.00 (td, *J* = 6.3, 3.1 Hz, 3H), 6.93-6.85 (m, 1H), 5.07 (dd, *J* = 12.6, 5.5 Hz, 1H), 4.57 (d, *J* = 2.3 Hz, 1H), 4.30 (s, 1H), 4.15 (s, 1H), 3.90 (d, *J* = 12.2 Hz, 2H), 3.72 (s, 1H), 3.51 (s, 2H), 3.08 (d, *J* = 9.6 Hz, 1H), 2.94-2.70 (m, 5H), 2.66 (d, *J* = 10.0 Hz, 1H), 2.51 (d, *J* = 6.9 Hz, 2H), 2.17-2.03 (m, 2H), 1.99-1.83 (m, 3H), 1.66 (s, 1H), 1.29 (s, 8H), 1.24 (s, 6H)	35



	A	832.3	^{1}H NMR (400 MHz, CD_3OD) δ 7.73 (dd, *J* = 8.9, 2.8 Hz, 3H), 7.66 (d, *J* = 8.5 Hz, 1H), 7.15 (d, *J* = 2.4 Hz, 1H), 7.05 (s, 1H), 6.99 (td, *J* = 6.3, 3.1 Hz, 3H), 6.89 (d, *J* = 8.5 Hz, 1H), 5.07 (dd, *J* = 12.5, 5.4 Hz, 1H), 4.57 (s, 1H), 4.30 (s, 1H), 4.15 (s, 1H), 3.90 (d, *J* = 12.3 Hz, 2H), 3.71 (s, 1H), 3.51 (s, 2H), 3.07 (d, *J* = 9.7 Hz, 1H), 2.90-2.70 (m, 5H), 2.66 (d, *J* = 10.0 Hz, 1H), 2.50 (d, *J* = 6.9 Hz, 2H), 2.17-2.03 (m, 2H), 1.99-1.83 (m, 3H), 1.66 (s, 1H), 1.29 (s, 8H), 1.24 (s, 6H)	35
	A	838.57	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.62-1.67 (m, 2H), 1.82-1.88 (m, 2H), 2.00-2.07 (m, 1H), 2.11-2.15 (m, 1H), 2.24-2.28 (m, 1H), 2.56-2.58 (m, 1H), 2.61-2.62 (m, 1H), 2.85-2.93 (m, 2H), 3.14-3.20 (m, 2H), 3.22-3.28 (m, 4H), 3.62-3.65 (m, 4H), 4.06 (d, *J* = 8.8 Hz, 1H), 4.24-4.27 (m, 2H), 4.32 (s, 1H), 5.08-5.13 (m, 1H), 6.72 (t, *J* = 8.8 Hz, 1H), 7.21 (d, *J* = 2.4Hz, 1H), 7.36-7.39 (m ,1H), 7.51-7.55 (m, 2H), 7.60-7.67 (m, 2H), 7.77 (d, *J* = 8.8 Hz ,1H), 7.91 (d, *J* = 8.80 Hz, 1H), 9.65 (br, 1H), 11.11 (s, 1H).	49




	A	835.2	^{1}H NMR (400 MHz, CD_3OD): δ7.80-7.78 (d, *J*=8.4Hz, 1H), 7.73-7.70 (m, 3H), 7.25-7.19 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.07 (m, 1H), 4.91-4.89 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.92-3.88 (m, 2H), 3.13-3.12 (m, 1H), 2.87-2.65 (m, 6H), 2.47-2.44 (m, 2H), 2.34-2.29 (m, 2H), 2.18-2.13 (m, 6H), 1.92-1.89 (m, 2H), 1.80-1.70 (m, 1H), 1.32-1.31 (m, 1H), 1.28 (s, 6H), 1.21 (s, 6H);	51
	A	839.57		49



	A	838.57		49
	A	856.56		49
	A	824.45	1H-NMR (300 MHz, DMSO-d*6*) δ11.10（s, 1H), 7.93- 7.86 (m, 1H), 7.77- 7.69 (m, 3H), 7.50- 7.34 (m, 2H), 7.22- 7.18 (m, 1H), 7.04- 6.96 (m, 1H), 6.58- 6.51 (m , 2H), 5.16 - 5.06 (m, 1H), 4.32(s, 1H), 4.09- 4.02 (m, 1H), 3.55- 3.20 (m, 6H), 3.09- 2.73 (m, 2H), 2.70- 2.38 (m, 10H), 2.20- 2.01 (m, 2H), 1.86- 1.67 (m, 1H), 1.24 - 1.13 (d, *J*=33 Hz, 12H).	46




	A	819.58	^{1}H NMR (400 MHz, DMSO-d^6) δ 0.97-1.07 (2H, m), 1.13 (6H, s), 1.23 (6H, s), 1.49-1.69 (5H, m), 1.80-2.02 (4H, m), 2.19-2.20 (1H, m), 2.33-2.43(1H, m), 2.50-2.60(2H, m), 2.81-2.91(1H, m), 3.30-3.35 (4H, m), 3.40-3.45 (4H, m), 4.06 (1H, d, *J* = 9.2 Hz), 4.32 (1H, s), 5.03-5.07 (1H, m), 6.99-7.02 (1H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.27 (1H, d, *J* = 8.8 Hz), 7.33-7.39 (3H, m), 7.68 (1H, d, *J* = 8.4 Hz),7.73-7.77 (3H, m), 7.91 (1H, d, *J* = 8.8 Hz), 11.09 (1H, s).	90
	A	799.63	^{1}H NMR (400 MHz, DMSO-d^6) δ 0.97-1.07 (2H, m), 1.13 (6H, s), 1.23 (6H, s), 1.46-1.69 (5H, m), 1.81-2.02 (4H, m), 2.19-2.20 (1H, m), 2.39-2.41(1H, m), 2.45 (3H, m), 2.53-2.59(2H, m), 2.81-2.89(1H, m), 3.30-3.35 (4H, m), 3.42-3.48 (4H, m), 4.05 (1H, d, *J* = 9.2 Hz), 4.23 (1H, s), 5.02-5.07 (1H, m), 6.80-6.83 (1H, m), 7.25-7.40 (4H, m), 7.67-7.78 (5H, m), 11.10 (1H, s).	90



	A	824.35	^{1}H NMR (300 MHz, $CDCl_3$) 7.83-7.80 (m, 1H), 7.63-7.39 (m, 4H), 7.19-7.05 (m, 1H), 6.90-6.71 (m, 4H), 6.04-6.01 (m, 1H), 5.14-5.08 (m, 1H), 4.38-4.07 (m, 3H), 3.97-3.95 (m, 1H), 3.81-3.30 (m, 4H), 3.19-2.98 (m, 3H), 2.89-2.55 (m, 6H), 2.31-2.18 (m, 4H), 1.82-1.61 (m, 2H), 1.19-1.15 (m, 15H);	20, 33
	A	824.35	^{1}H NMR (300 MHz, $CDCl_3$) 7.81-7.80 (m, 1H), 7.62-7.60 (m, 2H), 7.53-7.48 (m, 2H), 6.90-6.71 (m, 5H), 6.04-6.01(m, 1H), 5.13-5.07 (m, 1H), 4.37-4.06 (m, 3H), 3.97-3.41 (m, 5H), 3.12-3.06 (m, 3H), 2.88-2.42 (m, 7H), 2.30-2.15 (m, 4H), 1.82-1.75 (m, 1H), 1.19-1.05 (m, 15H);	41
	A	838.38	^{1}H NMR (300 MHz, DMSO) 10.96 (s, 1H), 7.91-7.88 (m, 1H), 7.74-7.71 (m, 2H), 7.48-7.41 (m, 2H), 7.24-7.20(m, 2H), 7.01-6.93 (m, 3H), 5.09-5.03 (m, 1H), 4.39-4.21 (m, 3H), 4.06-4.03 (m, 1H), 3.86-3.82 (m, 2H), 3.30-3.28 (m, 1H), 3.11-3.05 (m, 4H), 2.90-2.72 (m, 3H), 2.61-2.50 (m, 4H), 2.42-2.35 (m, 3H), 2.07-1.95 (m, 1H), 1.78-1.75 (m, 2H), 1.51-1.42 (m, 3H), 1.30-1.05 (m, 14H);	33




	A	838.4	^{1}H NMR (300 MHz, DMSO-*d6*) 10.96 (s, 1H), 7.91-7.88 (m, 1H), 7.74-7.70 (m, 2H), 7.48-7.20 (m, 4H), 7.02-6.93(m, 3H), 5.09-5.05 (m, 1H), 4.39-4.21 (m, 3H), 4.06-4.03 (m, 1H), 3.96-3.82 (m, 2H), 3.30-3.28 (m, 1H), 3.11-3.05 (m, 4H), 2.90-2.72 (m, 3H), 2.70-2.62 (m, 4H), 2.39-2.26 (m, 3H), 2.07-1.79 (m, 1H), 1.75-1.70(m, 2H), 1.52-1.46 (m, 3H), 1.31-1.05 (m, 14H);	33
	A	806.4	^{1}H NMR (300 MHz, DMSO-d_6): δ11.04 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.67 (dd, J = 13.3, 8.3 Hz, 3H), 7.44-7.13 (m, 4H), 7.02-6.91 (m, 1H), 6.38 (d, J = 8.3 Hz, 2H), 5.04 (dd, J = 12.4, 5.2 Hz, 1H), 4.28 (s, 1H), 3.99 (q, J = 8.1, 7.6 Hz, 3H), 3.50 (t, J = 6.5 Hz, 2H), 3.41 (d, J = 6.4 Hz, 4H), 2.96-2.47 (m, 4H), 2.47 (d, J = 3.4 Hz, 2H), 2.30 (d, J = 7.4 Hz, 2H), 2.09-1.90 (m, 1H), 1.85- 1.71 (m, 2H), 1.18 (s, 6H), 1.09 (s, 6H).	34



	A	821.35	^{1}H NMR (300 MHz, DMSO): δ 11.11 (s, 1H), 7.91-7.81 (m, 2H), 7.73 (d, J = 9.0 Hz, 2H), 7.48 (d, J = 9.1 Hz, 1H), 7.30-7.27 (m, 2H), 7.20 (m, 1H), 7.02-6.90 (m, 3H), 5.18-5.00 (m, 2H), 4.32 (s, 1H), 4.05 (d, J = 9.0 Hz, 1H), 3.89-3.70 (m, 4H), 3.05-2.98 (m, 2H), 2.98-2.81 (m, 1H), 2.80-2.69 (m, 2H), 2.64-2.56 (m, 2H), 2.49-2.40 (m, 2H), 2.10-1.93 (m, 1H), 1.80-1.79 (m, 2H), 1.60-1.44 (m, 1H), 1.32-1.14 (m, 10H), 1.12 (s, 6H)	23
	A	836.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.22 (s, 6H), 1.58-1.70 (m, 2H), 1.81-2.05 (m, 6H), 2.10-2.18 (m, 2H), 2.80-2.92 (m, 1H), 3.34-3.76 (m, 7H), 3.53-3.67 (m, 3H), 3.72-3.89 (m, 3H), 3.98-4.08 (m, 2H), 4.27-4.33 (m, 1H), 5.05-5.14 (m, 1H), 7.01 (s, 1H), 7.23-7.33 (m, 3H), 7.67-7.78 (m, 2H), 7.91 (m, 1H), 8.76-8.77 (m, 2H), 11.09 (s, 1H).	49
	A	805.56	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.13 (6H, s), 1.22 (6H, s), 1.37-1.57 (4H, m), 1.86-2.03 (5H, m), 2.32-2.42 (2H, m), 2.54-2.63 (2H, m), 2.67 (4H, brs), 2.84-2.91 (1H, m), 3.44 (4H, brs), 4.05 (1H, d, J = 9.2 Hz), 4.31 (1H, s), 5.07 (1H, dd, J = 5.6, 9.2 Hz), 7.00 (1H, dd, J = 2.0, 8.8 Hz), 7.21 (1H, d, J = 2.4 Hz), 7.25-7.27 (1H, m), 7.33-7.35 (3H, m), 7.67-7.69 (1H, m), 7.70-7.75 (3H, m), 7.89-7.92 (1H, m), 11.10 (1H, s).	91



	A	810.35	^{1}H NMR (300 MHz, CD_3OD): δ 8.60 (s, 1H), 7.99-7.95 (m, 1H), 7.83-7.70 (m, 2H), 7.42-7.31 (m, 2H), 7.13-7.12(s, 1H), 7.00-6.86 (m, 2H), 5.09-5.08 (m, 1H), 4.37-4.14 (m, 6H), 3.19-3.13 (m, 2H), 2.89-2.73 (m, 7H), 2.20-2.03 (m, 3H), 1.29-1.22 (m, 18H);	3
	A	809.2	^{1}H NMR (300 MHz, Methanol-d_4): δ 7.82 (d, J = 8.3 Hz, 1H), 7.79-7.70 (m, 3H), 7.42 (d, J = 2.2 Hz, 1H), 7.33 (dd, J = 8.3, 2.3 Hz, 1H), 7.14 (d, J = 2.4 Hz, 1H), 7.05-6.95 (m, 3H), 5.12 (dd, J = 12.3, 5.4 Hz, 1H), 4.29 (d, J = 0.9 Hz, 1H), 4.21 (t, J = 5.8 Hz, 2H), 4.14 (d, J = 0.8 Hz, 1H), 3.77 (s, 1H), 3.73 (s, 1H), 3.15-3.05 (m, 2H), 2.96-2.69 (m, 5H), 2.60 (t, J = 11.3 Hz, 2H), 2.19-2.08 (m, 1H), 2.03 (d, J = 8.6 Hz, 2H), 1.29 (s, 6H), 1.26 (s, 3H), 1.23 (d, J = 2.1 Hz, 9H).	21




	A	811.3	^{1}H NMR (400 MHz, DMSO): δ 11.11 (s, 1H), 8.76 (s, 2H), 7.90 (d, J = 8.7 Hz, 1H), 7.84 (d, J = 8.3 Hz, 1H), 7.72 (d, J = 9.2 Hz, 1H), 7.43 (d, J = 2.3 Hz, 1H), 7.36-7.33 (m, 1H), 7.22 (m, 1H), 7.02-6.99 (m, 1H), 5.14-5.09 (m, 1H), 4.53 (d, J = 11.9 Hz, 2H), 4.29 (s, 1H), 4.20 (m, 2H), 4.03 (d, J = 9.1 Hz, 1H), 2.97-2.83 (m, 3H), 2.74 (m, 2H), 2.65-2.52 (m, 3H), 2.49 (m, 1H), 2.11-2.01 (m, 1H), 1.83 (m, 2H), 1.22 (s, 6H), 1.11 (m, 12H)	3
	A	807.54	H-NMR (400 MHz, $CDCl_3$) δ 8.23 (s, 1H), 7.79-7.77 (d, *J*=8.4 Hz, 1H), 7.71-7.68 (d, *J*=9.0 Hz, 2H), 7.58-7.55 (d, *J*=8.7 Hz, 1H), 7.37-7.36 (d, *J*=1.8 Hz, 1H), 7.22-7.19 (d, *J*=8.4 Hz, 1H), 6.97-6.91 (m, 3H), 6.83-6.79 (m, 1H), 6.12-6.10 (d, *J*=8.1 Hz, 1H), 4.98-4.92 (m, 4H), 3.34 (s, 3H), 2.93-2.68 (m, 4H), 2.54 (s, 4H), 2.36 (s, 2H), 2.18-2.13 (m, 1H), 1.62 (s, 4H), 1.26 (s, 6H), 1.22 (s, 6H)	
	A	769.5		


	A	861.6		36
	A	834.59		49




	A	834.59		49
	A	834.59		49
	A	779.35	^{1}H NMR (300 MHz, CD3OD) 7.78-7.68 (m, 4H), 7.37-7.34 (m, 3H), 7.26-7.23 (m, 1H), 7.13-7.11 (m, 1H), 7.00-6.96(m, 1H), 5.08-5.04 (m, 1H), 4.29-4.27 (m, 1H), 4.16-4.14 (m, 1H), 3.92-3.89 (m, 2H), 3.11-2.71 (m, 11H), 2.13-2.11 (m, 1H), 1.31-1.28 (m, 18H);	22



	A	833.25	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.07 (s, 1H), 7.91 (d, *J* = 8.8 Hz, 1H), 7.71 (dd, *J* = 19.0, 8.4 Hz, 3H), 7.46 (d, *J* = 9.2 Hz, 1H), 7.35 (d, *J* = 2.4 Hz, 1H), 7.27 (dd, *J* = 8.7, 2.3 Hz, 1H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.01 (dd, *J* = 8.8, 2.4 Hz, 1H), 6.44 (d, *J* = 8.4 Hz, 2H), 5.08 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.32 (s, 1H), 4.06 (d, *J* = 9.1 Hz, 1H), 3.91 (t, *J* = 7.3 Hz, 2H), 3.64 (ddd, *J* = 24.3, 9.7, 5.1 Hz, 3H), 3.43 (dt, *J* = 18.6, 4.7 Hz, 6H), 3.03 – 2.82 (m, 4H), 2.64 – 2.51 (m, 2H), 2.40 (t, *J* = 4.8 Hz, 4H), 2.02 (dt, *J* = 15.2, 5.6 Hz, 1H), 1.22 (s, 6H), 1.13 (s, 6H).	36
	A	850.25	^{1}H NMR (400 MHz, CD_3OD): δ7.78-7.70 (m, 4H), 7.26-7.20 (m, 2H), 7.13-7.12 (d, J=2.0Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.07 (m, 1H), 4.93-4.89 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.32-3.30 (m, 4H), 3.28-3.26 (m, 1H), 2.87-2.67 (m, 7H), 2.58-2.46 (m, 6H), 2.39-2.34 (m, 2H), 2.25 (s, 3H), 2.14-2.11 (m, 1H), 1.28 (s, 6H), 1.21 (s, 6H);	51



	A		^{1}H NMR (300 MHz, DMSO-*d6*) 11.12 (s, 1H), 7.92-7.73 (m, 4H), 7.22-7.19 (m, 3H), 7.20-6.95 (m, 3H), 5.20-5.11 (m, 1H), 4.32 (s, 1H), 4.20-4.16 (m, 2H), 4.07-4.04 (m, 1H), 3.47-3.41 (m, 2H), 3.32-3.25 (m, 4H), 2.96-2.94 (m, 6H), 2.61-2.50 (m, 7H), 2.10-2.01 (m, 1H), 1.21 (s, 6H), 1.23 (s, 6H);	93
	A	806.56		




	A	779.25	^{1}H NMR (300 MHz, CD_3OD) 7.78-7.64 (m, 4H), 7.39-7.32 (m, 3H), 7.21-7.18 (m, 2H), 7.12-6.95 (m, 1H), 5.09-5.03(m, 1H), 4.28-4.27 (m, 1H), 4.15-4.13 (m, 1H), 3.52-3.47 (m, 2H), 3.29-3.22 (m, 4H), 3.18-2.63 (m, 7H), 2.13-2.05 (m, 1H), 1.28-1.11 (m, 18H);	22
	A	806.55		49
	A	779.1	1H NMR (300 MHz, DMSO-d6) δ 11.03 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.73 (dd, J = 8.6, 6.7 Hz, 3H), 7.61 (d, J = 8.5 Hz, 1H), 7.38-7.26 (m, 3H), 7.25-7.14 (m, 2H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 5.03 (dd, J = 12.6, 5.3 Hz, 1H), 4.29 (s, 1H), 4.03 (d, J = 9.1 Hz, 1H), 3.45 (dd, J = 12.6, 3.3 Hz, 2H), 3.19 (dd, J = 12.5, 6.5 Hz, 2H), 3.00 (td, J = 6.4, 3.4 Hz, 2H), 2.91-2.62 (m, 4H), 2.60-2.47 (m, 2H), 1.98 (dd, J = 10.9, 5.6 Hz, 1H), 1.19 (s, 6H), 1.10 (s, 6H), 0.98 (d, J = 6.2 Hz, 6H).	22



	A	792.4	^{1}H NMR (300 MHz, DMSO-*d6*) δ 11.08 (s, 1H), 7.92-7.89 (d, *J* = 9 Hz, 1H), 7.75-7.67 (m, 3H), 7.45-7.42 (d, *J* = 9.0 Hz, 1H), 7.35 (s, 1H), 7.28-7.25 (d, *J* = 9Hz, 1H), 7.21 (s, 1H), 7.02-6.98 (m, 1H), 6.45-6.42 (d, *J* = 9Hz,2H),5.10-5.05 (m, 1H) , 4.32 (s, 1H),4.07-3.99 (m, 3H), 3.59-3.55 (m, 2H), 3.47-3.45 (m,4H), 3.00-2.89 (m, 2H), 2.66-2.50(m, 8H), 2.04-2.00 (m, 1H), 1.21 (s, 6H),1.12 (s, 6H);	40
	A	820.58		




	A	836.55	^{1}H-NMR (300 MHz, DMSO-d_6) δ11.08（s, 1H), 8.61 (s, 1H), 7.90- 7.84 (m, 1H), 7.74- 7.67 (m, 2H), 7.49- 7.41 (m, 1H), 7.20- 7.16 (m, 1H), 7.02- 6.89 (m, 3H), 5.17 - 5.07 (m, 1H), 4.29 (s, 1H), 4.06- 3.99 (m, 1H), 3.91- 3.68 (m, 6H), 2.92- 2.50 (m, 7H), 2.42- 2.34 (m, 4H), 2.06- 1.98 (m, 1H), 1.78- 1.60 (m, 2H), 1.54- 1.38 (m, 3H), 1.26 - 1.10 (m, 14H)	5
	A	838.56		49
	A	824.59		



	A		^{1}H NMR (300 MHz, DMSO-d_6): δ 11.07 (s, 1H), 8.60 (s, 1H), 7.95-7.85 (m, 2H), 7.69 (d, J = 8.6 Hz, 1H), 7.57 (d, J = 9.2 Hz, 1H), 7.34 (s, 1H), 7.31(m, 1H) 7.17 (m, 1H), 7.00 (dd, J = 8.8, 2.5 Hz, 1H), 6.85 (d, J = 9.0 Hz, 1H), 5.07 (dd, J = 12.7, 5.6 Hz, 1H), 4.47 (s, 1H), 4.33 (d, J = 19.6 Hz, 2H), 4.05 (d, J = 9.1 Hz, 1H), 3.47 (s, 4H), 2.86 (d, J = 14.6 Hz, 2H), 2.55 (s, 5H), 2.24 (s, 1H),2.02 (m, 3H) 1.67 (m, 2H), 1.58 (m, 3H), 1.20 (m, 7H), 1.16 (m, 10H	94
	A	821.4	^{1}H NMR (400 MHz, DMSO-d_6) δ 11.12 (s, 1H), 7.91 (d, J = 8.8 Hz, 1H), 7.88-7.82 (m, 1H), 7.74 (d, J = 8.4 Hz, 2H), 7.43 (d, J = 9.2 Hz, 1H), 7.32-7.24 (m, 2H), 7.21 (d, J = 2.3 Hz, 1H), 7.01 (dd, J = 8.8, 2.4 Hz, 1H), 6.43 (d, J = 8.4 Hz, 2H), 5.13 (dd, J = 12.9, 5.3 Hz, 1H), 4.97 (td, J = 6.5, 3.2 Hz, 1H), 4.33 (s, 1H), 4.04-4.02 (m, 3H), 3.54 (t, J = 6.6 Hz, 2H), 3.12-3.10 (m, 1H), 2.93-2.90 (m, 13.9, 5.4 Hz, 1H), 2.79-2.77 (m, 1H), 2.71- 2.52 (m, 2H), 2.40 (dt, J = 12.7, 6.6 Hz, 2H), 2.22 (q, J = 7.4 Hz, 4H), 2.12 - 2.00 (m, 4H), 1.76	51



			(q, J = 7.2 Hz, 2H), 1.22 (s, 6H), 1.13 (s, 6H);	
	A	799.52		
	A	791.3	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.22 (6H, s), 1.73-1.78 (2H, m), 1.99-2.02 (1H, m), 2.42-2.49 (5H, m), 2.53-2.60 (6H, m), 2.83-2.89 (1H, m), 3.43 (5H, s), 4.06 (1H, d, *J* = 8.8 Hz), 4.32 (1H, s), 5.07 (1H, dd, *J* = 5.2, 12.8 Hz), 7.00 (1H, dd, *J* = 2.4, 8.8 Hz), 7.20-7.26 (2H, m), 7.31-7.33 (3H. m), 7.67 (1H, d, *J* = 8.8 Hz), 7.73-7.79 (3H, m), 7.90 (1H, d, *J* = 8.8 Hz), 11.0 (1H, s).	56



	A		^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.23 (6H, s), 1.98-2.02 (1H, m), 2.15-2.49 (5H, m), 2.52-2.67 (8H, m), 2.83-2.92 (1H, m), 3.44 (4H, s), 3.64-3.68 (1H, m), 4.06 (1H, d, *J* = 9.2 Hz), 4.32 (1H, s), 5.07 (1H, dd, *J* = 5.6, 12.8 Hz), 7.00 (1H, dd, *J* = 2.4, 8.8 Hz), 7.21 (1H, d, *J* = 2.4 Hz), 7.25 (1H, d, *J* = 8.8 Hz), 7.34-7.39 (3H, m), 7.68 (1H, d, *J* = 8.4 Hz),7.74-7.80 (3H, m), 7.90 (1H, d, *J* = 8.4 Hz), 11.0 (1H, s).	56
	A		^{1}H NMR (300 MHz, DMSO-d_6): δ 11.06 (s, 1H), 8.57 (d, J = 2.4 Hz, 1H), 7.95-7.83 (m, 2H), 7.66 (d, J = 8.5 Hz, 1H), 7.55 (d, J =9.2 Hz, 1H), 7.34-7.17 (m, 3H), 7.00 (dd, J = 8.8, 2.4 Hz, 1H), 6.83 (d, J = 9.1 Hz, 1H), 5.06 (dd, J = 12.6, 5.3 Hz, 1H), 4.53 (d, J = 13.1 Hz, 1H), 4.31 (d, J = 10.7 Hz, 2H), 4.04 (d, J = 9.2 Hz, 1H), 3.39 (m, 4H), 3.06 (d, J = 12.5 Hz, 1H), 2.86 (d, J = 11.6 Hz, 2H), 2.59 (m, 6H), 2.03 (s, 2H), 1.72 (t, J = 13.2 Hz, 4H), 1.60-1.32 (m, 5H), 1.21 (d, J = 2.0 Hz, 7H), 1.11 (s, 6H);	94
	A		^{1}H NMR (300 MHz, DMSO-d_6): δ 11.07 (s, 1H), 8.61 (d, J = 2.4 Hz, 1H), 7.96-7.85 (m, 2H), 7.68 (d, J = 8.5 Hz, 1H), 7.56 (d, J = 9.1 Hz, 1H), 7.34-7.17 (m, 3H), 7.00 (dd, J = 8.7, 2.4 Hz, 1H), 6.85 (d, J = 9.0 Hz, 1H), 5.07 (dd, J = 12.7, 5.4 Hz, 1H), 4.30 (s, 1H), 4.20 (d, J = 13.0 Hz, 1H), 4.05 (d, J = 9.2 Hz, 2H), 3.43 (s, 4H), 3.08 (dd, J = 23.5, 12.0 Hz, 2H), 2.88 (t, J = 13.6 Hz,	94




			1H), 2.68 (s, 5H), 2.38 (s, 2H), 2.07-1.96 (m, 1H), 1.87-1.49 (m, 9H), 1.41- 1.28 (m, 1H), 1.21 (s, 6H), 1.11 (s, 6H)	
	A	808.3	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.08 (s, 1H), 8.24 (d, J = 9.2 Hz, 1H), 7.90 (d, J = 8.8 Hz, 1H), 7.83 (d, J = 9.5 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.40 (d, J = 9.6 Hz, 1H), 7.33 (d, J = 2.1 Hz, 1H), 7.25 (q, J = 4.0, 3.0 Hz, 2H), 7.03 (dd, J = 8.8, 2.5 Hz, 1H), 5.06 (dd, J = 12.7, 5.3 Hz, 1H), 4.53 (d, J = 13.2 Hz, 2H), 4.46 (s, 1H), 4.00 (d, J = 9.1 Hz, 1H), 3.43 (brs, 4H), 3.05 (t, J = 12.5 Hz, 2H), 2.89 (s, 1H), 2.60 (d, J = 31.4 Hz, 7H), 2.01 (m, 1H), 1.91 (m, 2H), 1.46 (m, 2H), 1.22 (s, 6H), 1.14 (s, 6H).	52
	A	792.3	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.06 (s, 1H), 7.87 (d, J = 8.8 Hz, 1H), 7.69 (dd, J = 16.6, 8.5 Hz, 3H), 7.43 – 7.30 (m, 2H), 7.30 (m, 1H), 7.14 (m, 1H), 6.97 (dd, J = 8.8, 2.5 Hz, 1H), 6.56 (d, J = 8.7 Hz, 2H), 5.05 (dd, J = 12.7, 5.4 Hz, 1H), 4.29 (s, 1H), 4.02 (d, J = 9.3 Hz, 1H), 3.56 (t, J = 8.4 Hz, 1H), 3.39 (s, 5H), 3.13 (t, J = 8.7 Hz, 1H), 2.97 (d, J = 8.7 Hz, 2H), 2.83-2.56 (m, 5H), 2.52-2.04 (m, 2H), 1.88 (m, 3H), , 1.18 (d, J = 27.9 Hz, 6H), 1.04 (d, J = 27.9 Hz, 6H).	26



	A	778.5	H-NMR-PH-ARV-LS-049-Q-0 (300 MHz, DMSO-d_6, ppm) :δ 11.08 (s, 1H), 7.91 (d, *J* = 8.7 Hz, 1H), 7.77 (d, *J* = 8.6 Hz, 2H), 7.67 (d, *J* = 8.3 Hz, 1H), 7.52 (d, *J* = 9.2 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.06- 6.95 (m, 3H), 6.82 (d, *J* = 2.0 Hz, 1H), 6.68 (dd, *J* = 8.4, 2.1 Hz, 1H), 5.07 (dd, *J* = 12.6, 5.4 Hz, 1H), 4.33 (s, 1H), 4.21-4.12(m, 2H), 4.11-4.01 (m, 1H), 3.93 (dd, *J* = 8.8, 4.8 Hz, 2H), 3.41 (p, *J* = 5.7 Hz, 1H), 3.29 (s, 3H), 2.94-2.80 (m, 1H), 2.64-2.51 (m, 4H), 2.50-2.40 (m, 3H), 2.11-1.97 (m, 1H), 1.22 (s, 6H), 1.13 (s, 6H)	19
	A	820.3	H NMR (400 MHz, DMSO-*d6*) δ11.05 (s, 1H), 7.88 (d, J = 8.7 Hz, 1H), 7.71 (t, J = 8.4 Hz, 3H), 7.37 (d, J = 8.9 Hz, 2H), 7.31- 7.15 (m, 2H), 6.98 (dd, J = 8.7, 2.4 Hz, 1H), 6.56 (d, J = 8.6 Hz, 2H), 5.06 (m, 1H), 4.30 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.75 (m, 2H), 3.70-3.51 (m, 5H), 3.51- 3.32 (m, 5H), 2.85 (m, 1H), 2.57 (m, 2H), 2.37-1.81 (m, 4H), 1.15 (d, 12H)	18
	A	838.3	^{1}H NMR (400 MHz, DMSO-d_6) δ11.11 (s, 1H), 7.92 (d, J = 8.7 Hz, 1H), 7.78 (m, 3H), 7.53 (d, J = 7.2 Hz, 1H), 7.41 (d, J = 9.2 Hz, 1H), 7.22 (s, 1H), 7.02 (d, J = 8.6 Hz, 1H), 6.60 (d, J = 8.4 Hz, 2H), 5.13 (dd, J = 12.9, 5.3 Hz, 1H), 4.33 (s, 1H), 4.07 (d, J = 9.2 Hz, 1H), 3.80 (s, 2H), 3.61 (m, 2H), 3.44 (m, 2H), 3.36 (d, J = 9.9 Hz, 4H), 3.25 (d, J = 6.1 Hz, 2H), 2.94-2.83 (m, 1H),	18




			2.60 (t, J = 14.3 Hz, 3H), 2.30- 2.11 (m, 2H), 2.06 (m, 1H), 1.23 (s, 6H), 1.14 (s, 6H)	
	A	824.3	1H NMR (300 MHz, DMSO-*d6*): δ 11.07 (s, 1H), 7.87 (d, J = 8.7 Hz, 1H), 7.70 (m, 3H), 7.38 (dd, J = 27.2, 8.3 Hz, 2H), 7.17 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 6.51 (d, J = 8.7 Hz, 2H), 5.07 (dd, J = 12.9, 5.3 Hz, 1H), 4.28 (s, 1H), 4.02 (d, J = 9.1 Hz, 1H), 3.49-3.30 (m, 2H), 3.24-3.17 (m,, 4H), 3.09-2.98 (m, 1H), 2.91-2.78 (m, 1H), 2.65-2.48 (m, 6H), 2.47-2.42 (m, 2H), 2.41-2.39 (m, 2H), 2.15-1.98 (m, 2H), 1.80-1.60(m, 1H), 1.18 (s, 6H), 1.09 (s, 6H).;	49
	A	824.2	1H NMR (300 MHz, DMSO-*d*6) δ 11.04 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 (m, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.06 – 6.89 (m, 4H), 5.03 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.28 (s, 1H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.78-3.58 (m, 3H), 3.44 -3.21(m, 4H), 2.94–2.75 (m, 1H), 2.64–2.55 (m, 5H), 2.42 (m, 3H), 2.37 (d, *J* = 7.4 Hz, 2H), 2.12 – 1.92 (m, 2H), 1.68 (m, 1H), 1.18 (s, 6H), 1.09 (s, 6H)	53



	A	779.3	^{1}H NMR (300 MHz, d6-DMSO) δ 1.13 (s, 6H), 1.22 (s, 6H), 2.07 (s, 1H), 2.52-2.64 (m, 2H), 2.82-2.90 (m, 3H), 3.37-3.44 (m, 4H), 3.96 (s, 4H), 4.04-4.07 (m, 1H), 4.18 (s, 2H), 4.32 (s, 1H), 5.16 (s, 1H), 6.42-6.45 (m, 2H), 7.00-7.03 (m, 1H), 7.21 (s, 1H), 7.35-7.44 (m, 3H), 7.72-7.74 (m, 2H), 7.84-7.93 (m, 2H), 11.13 (s, 1H).	24
	A	848.4	1H NMR (400 MHz, DMSO-*d*6): δ 11.06 (s, 1H), 7.90 (m, 1H), 7.75 (m, 2H), 7.65 (m, 1H), 7.55 (m, 1H), 7.20 (m, 1H), 7.02 (m, 3H), 6.97 (m, 1H), 6.85 (m, 1H), 5.04 (m, 1H), 4.50 (m, 1H), 4.30 (m, 1H), 4.12 (m, 1H), 4.05 (m, 1H), 3.95 (m, 2H), 3.70 (m, 1H), 3.50 (m, 3H), 3.30 (m, 1H), 3.00 -2.88 (m, 4H), 2.62-2.52 (m, 4H), 2.41-2.02 (m, 3H), 2.01-1.4 (m, 3H), 1.30-1.10 (d, 12H	6
	A	854.45	1H NMR (300 MHz, $CDCl_3$): δ 8.12 (d, J = 9.1 Hz, 1H), 8.03 (s, 1H), 7.94 (d, J = 9.5 Hz, 1H), 7.54 (d, J = 8.7 Hz, 1H), 7.45 (d, J = 11.0 Hz, 1H), 7.37 (d, J = 7.3 Hz, 1H), 7.01 – 6.91 (m, 2H), 6.78 (dd, J = 8.7, 2.4 Hz, 1H), 4.91 (dd, J = 12.2, 5.3 Hz, 1H), 4.50 (d, J = 13.3 Hz, 2H), 4.17 (d, J = 9.1 Hz, 1H), 4.04 (s, 1H), 3.28 (s, 4H), 3.03 (t, J = 12.3 Hz, 2H), 2.94 – 2.66 (m, 3H), 2.62 (s, 4H), 2.46 (s, 2H), 2.17 – 2.06 (m, 1H), 1.98 (s, 0H), 1.86 (d, J = 12.9 Hz, 2H), 1.69 (s, 1H), 1.26 (s,	29



			9H), 1.19 (s, 6H).	
	A	850.5	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.11 (s, 6H), 1.17-1.32 (m, 16H), 1.77-1.82 (m, 1H), 1.95-2.07 (m, 2H), 2.17-2.33 (m, 4H), 2.54-2.61 (m, 4H), 2.83-2.94 (m, 1H), 3.47 (s, 4H), 4.04 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1H), 4.95-5.01 (m, 1H), 5.04-5.11 (m, 1H), 6.98-7.04 (m, 1H), 7.20-7.37 (m, 3H), 7.66-7.73 (m, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.74-8.85 (m, 2H), 11.08 (s, 1H).	95
	A	850.5	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (s, 6H), 1.22-1.32 (m, 14H), 1.80 (d, *J* = 12.4 Hz, 2H), 1.99-2.03 (m, 1H), 2.24-2.33 (m, 3H), 2.55-2.61 (m, 6H), 2.84-2.93 (m, 1H), 3.47 (m, 4H), 4.04 (d, *J* = 9.2 Hz, 1H), 4.29 (s, 1 H), 4.96-5.01(m, 2H), 5.07 (dd, *J* = 5.2,12.8Hz, 1H), 5.07 (dd, *J* = 2.4,8.8Hz, 1H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.28 (d, *J* = 8.8 Hz, 1H), 7.36 (s, 1H), 7.68-7.74 (m, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.79 (s, 2H), 11.08 (s, 1H).	95



	A	850.6	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (d, *J* = 7.2 Hz, 6H), 1.26-1.30 (m, 6H), 1.68 (d, *J* = 13.0 Hz, 1H), 1.78-1.88 (m, 2H), 2.00-2.04 (m, 1H), 2.16-2.38 (m, 5H), 2.54-2.68 (m, 5H), 2.84-2.94 (m, 1H), 3.47 (s, 4H), 4.04 (d, *J* = 8.8 Hz, 1H), 4.29 (s, 1H), 4.48-4.56 (m, 2H), 5.08(d, *J* = 12.8 Hz,1H), 7.01 (d, *J* = 8.8 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.27 (d, *J* = 8.4 Hz, 1H), 7.36 (s, 1H), 7.68-7.71 (m, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.78 (s, 2H), 11.09 (s, 1H).	95
	A	807.54		
	A	875.65	H-NMR-PH-ARV-LS-035-H-0: (300 MHz, CD_3OD, ppm) δ 8.60-8.59 (d, *J*=2.4Hz, 1H), 7.97-7.94 (m, 1H), 7.87-7.70 (m, 4H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 1H), 6.86-6.83 (m, 1H), 5.17-5.11 (m, 1H), 4.40-4.35 (m, 2H), 4.28 (s, 1H), 4.14 (s, 1H), 3.07-2.77 (m, 10H), 2.28-2.13 (m, 6H), 2.00-1.78 (m, 8H), 1.61-1.58 (m, 1H), 1.42-1.33 (m, 1H), 1.28 (s, 6H), 1.21 (s, 6H), 1.13-1.00 (m, 1H).	96




	A	835.4	^{1}H NMR (300 MHz, CD_3OD): δ 7.77-7.70 (m, 4H), 7.25-7.19 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 7.01-6.95 (m, 3H), 5.12-5.06 (m, 1H), 4.96-4.90 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 4.01-3.97 (m, 2H), 2.88-2.68 (m, 9H), 2.36-2.31 (m, 7H), 2.10-2.00 (m, 1H), 1.95-1.85 (m, 2H), 1.67-1.60 (m, 2H), 1.24 (s, 6H), 1.18 (s, 6H);	23
	A	820.35	^{1}H NMR (300 MHz, $CDCl_3$) 7.92 (s, 1H), 7.63-7.47 (m, 4H), 7.22-7.19 (m, 1H), 7.01-6.98 (m, 1H), 6.97-6.89 (m, 1H), 6.88-6.71 (m, 1H), 6.53-6.50 (m, 2H), 6.00 (m, 1H), 4.88-4.84 (m, 1H), 4.08-4.05 (m, 1H), 3.96-3.92 (m, 3H), 3.30 (s, 1H), 2.97-2.80 (m, 5H) , 2.65-2.64 (m, 2H), 2.77-2.64 (m, 1H), 2.33 (s, 1H), 2.06-2.03 (m, 3H), 1.93 (m,2H), 1.51-1.35 (m, 5H), 1.18 (m, 6H), 1.14(m, 6H), 0.81-0.76 (m, 1H);	36



	A	835.1	^{1}H-NMR (400 MHz, DMSO-*d6*) δ 11.05 (s, 1H), 7.88-7.85 (d, *J*=9.0 Hz, 1H), 7.73-7.63 (m, 3H), 7.50-7.47 (d, *J*=9.0 Hz, 1H), 7.30-7.17 (m, 3H), 6.99-6.92 (m, 3H), 5.06-5.00 (m, 1H), 4.28 (s, 1H), 4.03-4.00 (m, 1H), 3.40-3.22 (m, 8H), 2.86-2.79 (m, 1H), 2.53-2.46 (m, 14H), 2.03-1.98 (m, 1H), 1.18-1.08 (m, 12H);	38
	A	863.4	H-NMR (300 MHz, DMSO-d*6*) δ 11.09 (s, 1H), 7.90-7.87 (d, *J* = 9Hz, 1H), 7.82-7.80 (d, *J* = 6 Hz, 1H), 7.72-7.69 (d, *J* = 9 Hz, 2H), 7.47-7.44 (d, *J* = 9 Hz, 1H), 7.26-7.18 (m, 3H), 7.00-6.92 (m, 3H), 5.12-5.06 (m, 1H), 4.88 (s, 1H), 4.30 (s, 1H), 4.04-4.01 (d, *J* = 9 Hz, 1H), 3.88-3.84 (d, *J* = 12 Hz, 2H), 3.67-3.30 (m, 1H), 2.91-2.70 (m, 4H), 2.60-2.37 (m, 4H), 2.24-2.13 (m, 4H), 2.05-2.00 (m, 1H), 1.82-1.78 (d, *J* = 12 Hz, 2H), 1.54 (s, 1H), 1.19 (s, 6H), 1.10 (s, 8H), 0.91-0.89 (d, *J*=6 Hz, 6H)	23
	A	834.5	^{1}H NMR (400 MHz, DMSO): δ 11.08 (s, 1H), 8.77-8.68 (m, 2H), 7.91 (d, J = 8.8 Hz, 1H), 7.68 (d, J = 8.7 Hz, 2H), 7.33 (d, J = 2.3 Hz, 1H), 7.29-7.19 (m, 2H), 7.02-6.99 (m, 1H), 5.09-5.05 (m, 1H), 4.64 (s, 1H), 4.29 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 3.55 (m, 1H), 3.44-3.42 (m, 4H), 3.35-3.32 (m, 1H), 2.92-2.81 (m, 1H), 2.68-2.53 (m, 5H), 2.48-2.36 (m, 3H), 2.35-2.18	97




			(m, 2H), 2.08-1.91 (m, 2H), 1.81 (m, 1H), 1.69 (m, 1H), 1.22 (s, 6H), 1.12 (s, 6H), 1.06 (m, 1H)	
	A	834.1	^{1}H NMR (400 MHz, DMSO): δ 11.08 (s, 1H), 8.73 (d, J = 2.7 Hz, 2H), 7.91 (d, J = 8.8 Hz, 1H), 7.67 (m, 2H), 7.34 (d, J = 2.1 Hz, 1H), 7.29-7.24 (m, 1H), 7.22 (d, J = 2.4 Hz, 1H), 7.01 (m, 1H), 5.07 (m, 1H), 4.68 (s, 1H), 4.30 (s, 1H), 4.04 (d, J = 9.2 Hz, 1H), 3.51-3.39 (m, 5H), 3.34 (m, 1H), 3.26 (m, 1H), 2.96-2.82 (m, 1H), 2.56 (m, 5H), 2.44-2.34 (m, 1H), 2.17 (m, 1H), 2.04-1.96 (m, 2H), 1.80-1.54 (m, 3H), 1.37 (d, J = 11.8 Hz, 1H), 1.22 (m, 7H), 1.12 (s, 6H)	97
	A	840.35	H-NMR (400 MHz, DMSO-*d6*) δ11.12 (s, 1H), 8.76 (s, 2H), 7.92-7.90 (d, *J* = 8.8Hz, 2H), 7.76-7.70 (m, 2H), 7.48-7.46 (d, *J*=7.2Hz, 1H), 7.23-7.22 (d, *J*=2.4Hz, 1H), 7.03-7.01 (m, 1H), 5.14-5.1 (m, 1H), 4.78-4.74 (d, *J*=12.8Hz, 2H), 4.30 (s, 1H), 4.06-4.04 (d, *J*=9.2Hz, 1H), 3.27 (s, 3H), 3.03-2.97 (m, 3H), 2.63-2.52 (m, 6H), 2.24-2.23 (m, 2H), 2.05-2.04 (m, 1H), 1.86-1.83 (m, 3H), 1.23 (s, 6H), 1.12 (s, 6H), 1.09 -1.06 (m, 2H)	49



	A	781.3	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.12 (s, 1H), 7.91 (d, J = 8.8 Hz, 1H), 7.84 (d, J = 8.3 Hz, 1H), 7.75 (d, J = 8.6 Hz, 2H), 7.48 (d, J = 2.3 Hz, 1H), 7.45-7.32 (m, 2H), 7.21 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.5 Hz, 1H), 6.57 (d, J = 8.8 Hz, 2H), 5.13 (dd, J = 12.8, 5.2 Hz, 1H), 4.33 (s, 3H), 4.06 (d, J = 9.1 Hz, 1H), 3.55 (t, J = 8.3 Hz, 1H), 3.45 (t, J = 8.7 Hz, 1H), 3.29 (d, J = 8.9 Hz, 2H), 3.20-3.07 (m, 1H), 2.88 (d, J = 12.3 Hz, 3H), 2.60 (d, J = 18.2 Hz, 2H), 2.36 (s, 3H), 2.22 (s, 1H), 2.06 (d, J = 10.6 Hz, 1H), 1.96-1.83 (m, 1H), 1.22 (s, 6H), 1.13 (s, 6H).	93
	A	779.15	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 7.84 (dd, J = 18.3, 8.7 Hz, 2H), 7.71 (d, J = 8.2 Hz, 2H), 7.35 -7.26 (d, J = 6.5 Hz, 4H), 7.17 (s, 1H), 6.49 (d, J = 8.5 Hz, 2H), 5.15 – 5.02 (m, 2H), 4.29 (s, 1H), 4.02 (d, J = 9.2 Hz, 1H), 3.76 (d, J = 6.9 Hz, 3H), 3.09 (dd, J = 25.1, 8.6 Hz, 4H), 2.98 (s, 2H), 2.54 (s, 2H), 1.97(s,4H),1.18 (s, 6H), 1.09 (s, 6H).	23




	A	793.25	1H NMR (300 MHz, CDCL3) δ 11.07 (s, 1H), 7.76 (t, J = 8.0 Hz, 3H), 7.58 (d, J = 8.6 Hz, 1H), 7.33 (d, J = 18.6 Hz, 2H), 7.19 (d, J = 8.2 Hz, 1H), 6.99 (d, J = 2.3 Hz, 1H), 6.84 (dd, J = 8.8, 2.4 Hz, 1H), 6.53 (d, J = 8.4 Hz, 2H), 4.22 (d, J = 8.3 Hz, 3H), 4.10 (s, 1H), 3.45 (s, 4H), 3.19 (s, 2H), 3.08 (s, 4H), 2.67 (d, J = 17.7 Hz, 4H), 2.57 (d, J = 13.7 Hz, 1H), 2.28 (s, 1H), 1.77 (s, 1H), 1.37 (s, 3H), 1.28 (d, J = 4.9 Hz, 9H).;	55
	A	802.5	^{1}H NMR (400 MHz, d-DMSO):δ11.086(s,1H),8.754(s,2H),7.708-7.674(m,3H),7.348(s,1H),7.275-7.253(d,J=8.8Hz,1H),6.938-6.933(d,J=2Hz,1H),6.830-6.803(m,1H),5.098-5.052(m,1H),4.764-4.732(d,J=12.8Hz,2H),4.193(s,1H),4.036-4.013(d,J=9.2Hz,1H),3.450(s,4H),3.020-2.961(m,2H),2.932-2.840(m,1H),2.608-2.537(m,6H),2.507-2.498(m,3H),2.216-2.199(d,J=6.8Hz,2H),2.076-2.018(m,1H),2.005-1.914(m,1H),1.851-1.819(d,J=12.8Hz,2H),1.215(s,6H),1.110(s,6H),1.073-1.018(m,2H)	49



	B	765.7	^{1}H NMR (400 MHz, d-DMSO): δ 10.69 (s,1H), 7.91-7.89 (d, *J*=8.8Hz, 1H), 7.78-7.74 (m, 3H), 7.53-7.51 (d, *J*=8.4Hz, 1H), 7.32-7.30 (d, *J*=8.0Hz, 2H), 7.21-7.20 (d, *J*=2.4Hz, 1H), 7.05-6.98 (m, 3H), 5.30 (s, 2H), 4.58-4.54 (m, 1H), 4.32 (s,1H), 4.07-4.05 (d, *J*=9.2Hz, 1H), 3.29-3.26 (m, 4H), 2.69-2.58 (m, 4H), 2.50-2.49 (m, 4H), 2.33-2.32 (s, 2H), 2.07-1.95 (m, 2H), 1.64-1.50 (m, 4H), 1.20 (s, 6H), 1.10 (s, 6H);	21A
	B	765.7	^{1}H NMR (400 MHz, d-DMSO): δ10.69 (s, 1H), 7.91-7.89 (d, *J*=8.8Hz, 1H), 7.78-7.74 (m, 3H), 7.53-7.51 (d, *J*=8.4Hz, 1H), 7.32-7.30 (d, *J*=8.0Hz, 2H), 7.21-7.20 (d, *J*=2.4Hz, 1H), 7.05-6.98 (m, 3H), 5.29 (s, 2H), 4.58-4.54 (m, 1H), 4.32 (s,1H), 4.07-4.05 (d, *J*=9.2Hz, 1H), 3.29-3.27 (m, 4H), 2.69-2.58 (m, 4H), 2.50-2.49 (m, 4H), 2.34-2.32 (s, 2H), 2.01-1.92 (m, 2H), 1.66-1.48 (m, 4H), 1.23 (s, 6H), 1.18 (s, 6H);	21A
	B	793.7	^{1}H NMR (400 MHz, d-DMSO): δ7.92-7.89 (d, *J*=8.4Hz, 1H), 7.78-7.66 (m, 4H), 7.32-7.30 (d, *J*=8.0Hz, 2H), 7.21-7.10 (m, 3H), 7.01-7.00 (d, J=6.8Hz, 1H), 5.39 (s, 1H), 4.62 (s, 1H), 4.33 (s,1H), 4.07-3.98 (m, 3H), 3.56-3.40 (m, 2H), 3.34-3.08 (m, 6H), 2.67-2.56 (m, 4H), 2.10-1.95 (m, 2H), 1.68-1.55 (m, 4H), 1.33 (m, 5H), 1.23 (s, 6H), 1.13 (s, 6H);	21A




	B	793.7	^{1}H NMR (400 MHz, *d*-DMSO): δ 10.70 (s, 1H), 7.91-7.89 (d, *J*=8.8Hz, 1H), 7.77-7.74 (m, 3H), 7.55-7.53 (d, *J*=8.4Hz, 1H), 7.31-7.29 (d, *J* = 8.4Hz, 2H), 7.22-7.10 (d, *J*=2.0Hz, 1H), 7.08-6.99 (m, 3H), 5.31 (s, 2H), 4.59-4.55 (s, 1H), 4.32 (s,1H), 4.07-4.05 (m, 1H), 3.34-3.33 (m, 8H), 2.66-2.50 (m, 6H), 2.10-1.90 (m, 2H), 1.60-1.38 (m, 4H), 1.32 (s, 4H), 1.22 (s, 6H), 1.13 (s, 6H);	21A
	B	843.4	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.11 -11.04 (m, 1 H), 8.07 (d, *J* = 8.7 Hz, 1 H), 7.96 (dd, *J* = 24.7, 8.4 Hz, 1 H), 7.78 (dd, *J* = 24.7, 8.4 Hz, 2 H), 7.50 (d, *J* = 9.2 Hz, 1 H), 7.44- 7.21 (m, 4 H), 6.93 (d, *J* = 8.7 Hz, 2 H), 5.08 (dd, *J* = 12.8, 5.3 Hz, 1 H), 4.36 (s, 1 H), 4.16 (t, *J* = 6.3 Hz, 2 H), 4.04 (d, *J* = 9.3 Hz, 1 H), 3.21 (t, *J* = 4.8 Hz, 3 H), 2.83 (dd, *J* = 13.5, 5.3 Hz, 1 H), 2.56 (m, 7 H), 2.31 (t, *J* = 6.7 Hz, 2 H), 2.00 (s, 1H), 1.96 (t, *J* = 7.0 Hz, 2 H), 1.47 (s, 4 H), 1.20 (s, 6 H), 1.11 (s, 6 H);	9



	B	868.4	^{1}H NMR (300 MHz, CD3OD) 7.93-7.90 (m, 1H), 7.77-7.70 (m, 2H), 7.68-7.65 (m, 1H), 7.36-7.29 (m, 4H), 7.29-7.19 (m, 2H), 5.09-5.03 (m, 1H), 4.34-4.31 (m, 1H), 4.17-4.10 (m, 1H), 3.87-3.80 (m, 2H), 2.84-2.67 (m, 10H), 2.11-2.08 (m, 1H), 1.67-1.50 (m, 2H) , 1.50-1.08 (m, 25H);	98
	B	804.3	^{1}H NMR (300 MHz, CD_3OD): δ 7.84-7.64 (m, 6H), 7.60-7.35 (m, 2H), 7.13-7.12 (s, 1H), 7.00-6.96 (m, 1H), 6.30 (s, 1H), 5.18-5.12 (m, 1H), 4.70-4.43 (m, 6H), 4.29 (s, 1H), 4.16 (s, 1H), 3.31-3.07 (m, 6H), 2.96-2.48 (m, 8H), 2.21-2.17 (m, 1H), 1.97-1.92 (m, 1H), 1.28-1.15 (s, 12H);	99
	B	801.4	1H NMR (300 MHz, Methanol-d4) δ 7.94 – 7.74 (m, 4H), 7.66 – 7.14 (m, 4H), 7.12 – 7.03 (m, 2H), 5.10 (dd, J = 12.3, 5.4 Hz, 1H), 4.58 (t, J = 4.2 Hz, 2H), 4.32 (s, 1H), 4.14 (t, J = 4.3 Hz, 1H), 3.72 (t, J = 4.6 Hz, 2H), 3.55 (d, J = 8.1 Hz, 7H), 2.75 (dddd, J = 22.4, 17.9, 8.2, 4.6 Hz, 3H), 2.11 (dtd, J = 12.7, 4.8, 2.3 Hz, 1H), 1.94 – 1.33 (m, 2H), 1.24 (d, J = 17.7 Hz, 12H).	9




	B	805.4	^{1}H NMR (300 MHz, $CDCl_3$) 7.80 (s, 1H), 7.70-7.67 (m, 3H), 7.59-7.56 (m, 1H), 7.27 (s, 1H), 7.01 (s, 1H), 6.83-6.80 (m, 1H), 6.61-6.58 (m, 1H), 6.41 (s, 1H), 6.20-6.18 (m, 1H), 5.32-5.27 (m, 2H), 4.76-4.73 (m, 1H), 4.31 (s, 1H), 4.18-4.15 (m, 1H), 4.06 (s, 1H), 3.93-3.75 (m, 1H), 3.64-3.46 (m, 2H), 2.96-2.88 (m, 1H), 2.59-2.53 (m, 2H), 2.22-2.15 (m, 4H), 1.78-1.35 (m, 6H),1.32-1.23 (m, 18H), 0.88-0.83 (m, 1H);	21A
	B	767.51	^{1}H NMR (300 MHz, Methanol-d4) δ 7.84-7.64 (m, 4H), 7.44 (d, J = 2.2 Hz, 1H), 7.34 (dd, J = 8.3, 2.3 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 7.03-6.90 (m, 3H), 5.09-5.07 (m, 1H), 4.39- 4.22 (m, 3H), 4.14-4.12 (m, 1H), 3.38-3.32 (m, 4H), 2.89-2.87 (m, 2H), 2.86-2.63 (m, 7H), 2.17 - 2.02 (m, 1H), 1.22 (d, J = 17.9 Hz, 12H) ;	9



	B	805.4	^{1}H NMR (300 MHz, Methanol-d4) δ 7.84-7.64 (m, 4H), 7.44 (d, J = 2.2 Hz, 1H), 7.34 (dd, J = 8.3, 2.3 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 7.03-6.90 (m, 3H), 5.09-5.07 (m, 1H), 4.39- 4.22 (m, 3H), 4.14-4.12 (m, 1H), 3.38-3.32 (m, 4H), 2.89-2.87 (m, 2H), 2.86-2.63 (m, 7H), 2.17 - 2.02 (m, 1H), 1.22 (d, J = 17.9 Hz, 12H) ;	21A
	B	805.4	^{1}H NMR (400 MHz, $CDCl_3$) 7.85 (s, 1H), 7.62-7.60 (m, 3H), 7.50-7.48 (m, 1H), 7.19-7.16 (m, 1H), 6.89-6.85 (m, 1H), 6.75-6.72 (m, 1H), 6.52-6.50 (m, 1H), 6.32-6.30 (m, 1H), 6.13-6.11 (m, 1H), 5.22-5.13 (m, 2H), 4.67-4.64 (m, 1H), 4.21-4.15 (m, 1H), 4.10-4.07 (m, 1H), 4.07-3.98 (m, 1H), 3.57-3.50 (m, 1H), 3.32-3.21 (m, 2H), 2.97-2.84 (m, 2H), 2.58-2.40 (m, 6H), 2.12-2.08 (m, 2H), 1.96-1.83 (m, 2H) , 1.54-1.40 (m, 2H), 1.40-1.04 (m, 18H);	21A
	B	807.4	^{1}H NMR (300 MHz, $CDCl_3$): δ 7.85-7.71 (m, 4H), 7.58-7.56 (m, 1H), 7.26-7.22 (m, 2H), 6.99-6.97 (m, 2H), 6.83-6.79 (m, 2H), 6.25-6.22 (m, 1H), 5.31-5.30 (m, 2H), 4.76-4.65 (m, 3H), 4.45-4.43 (m, 2H), 4.18-4.06 (m, 2H), 3.49-3.30 (s, 4H), 3.07-2.59 (m, 9H), 2.25-2.20 (m, 2H), 2.01-1.94 (m, 2H), 1.28-1.23 (d, *J*=15.0Hz,13H);	100




	B	767.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.24 (s, 6H), 1.30 (s, 6H), 1.96-2.01 (m, 4H), 2.11-2.17 (m, 1H), 2.44-2.50 (m, 2H), 2.71-2.78 (m, 2H), 2.83-2.92 (m, 2H), 3.03 (t, *J* = 5.2 Hz, 2H), 3.27 (s, 2H), 4.18 (s, 1H), 4.29 (s, 1H), 4.38 (t, *J* = 5.2 Hz, 2H), 5.10-5.13 (m, 1H), 6.97-6.99 (m, 1H), 7.13-7.14 (m, 1H), 7.37-7.39 (m, 1H), 7.45-7.48 (m, 2H), 7.72-7.74 (m, 1H), 7.82-7.85 (m, 1H), 8.14-8.16 (m, 1H), 8.88 (d, *J* = 2.0 Hz, 1H).	9, 101
	B	795.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.19 (s, 6H), 1.70-2.05 (m, 10H), 2.35-2.48 (m, 2H), 2.58-2.90 (m, 7H), 4.08 (s, 1H), 4.12 (t, *J* = 5.6 Hz, 2H), 4.19 (s, 1H), 4.98-5.04 (m, 1H), 6.87-6.91 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.22-7.26 (m, 1H), 7.32 (d, J = 2.0 Hz, 1H), 7.35 (d, *J* = 8.4 Hz, 1H), 7.63(d, *J* = 8.8 Hz, 1H), 7.72 (d, *J* = 8.4 Hz, 1H), 8.04-8.08 (m, 1H), 8.79 (d, *J* = 2.0 Hz, 1H).	9, 101



	B	795.4	^{1}H NMR (400 MHz, CD_3OD): δ 1.13 (s, 6H), 1.19 (s, 6H), 1.70-2.05 (m, 10H), 2.35-2.48 (m, 2H), 2.58-2.90 (m, 7H), 4.08 (s, 1H), 4.12 (t, *J* = 5.6 Hz, 2H), 4.19 (s, 1H), 4.98-5.04 (m, 1H), 6.87-6.91 (m, 1H), 7.03 (d, *J* = 2.4 Hz, 1H), 7.22-7.26 (m, 1H), 7.32 (d, J = 2.0 Hz, 1H), 7.35 (d, *J* = 8.4 Hz, 1H), 7.63(d, *J* = 8.8 Hz, 1H), 7.72 (d, *J* = 8.4 Hz, 1H), 8.04-8.08 (m, 1H), 8.79 (d, *J* = 2.0 Hz, 1H).	9
	B	811.3	^{1}H-NMR (300 MHz,d-DMSO) δ11.12(s, 1H), 8.65 (s, 1H), 8.38(m, 1H), 7.92- 7.81(m, 3H), 7.50- 7.33(m, 2H), 7.25-7.24(m, 1H), 7.14- 7.01(m, 1H), 5.17-5.07(m, 1H), 4.44(s, 1H), 4.24- 4.17(m, 2H), 3.99-3.36(m, 5H), 2.91- 2.83(m, 2H), 2.64 - 2.54(m, 3H), 2.45- 2.41(m, 2H), 2.28(s, 1H), 2.09- 2.02(m, 1H), 1.85- 1.47(m, 7H), 1.19 - 1.13 (d, *J = 18Hz* , 12H).	9
	B	811.7	H-NMR (300 MHz, CD_3OD) δ 7.94 (s, 1H), 7.79-7.68 (m, 2H), 7.38-7.27 (m, 3H), 7.13-7.12 (d, *J*=2.4 Hz, 1H), 6.99-6.96 (m, 1H), 5.11-5.05 (m, 1H), 4.31 (s, 1H), 4.19-4.15 (m, 2H), 4.08 (s, 1H), 3.91 (s, 1H), 3.02-2.99 (m, 4H), 2.85-2.71 (m, 5H), 1.92-1.87 (m, 2H),1.76-1.74 (d, *J*=4.8 Hz, 2H) , 1.60-1.58 (d, *J*=7.2 Hz, 2H), 1.28 (s, 6H), 1.20 (s, 6H);	9



	B	810.29	H-NMR (400MHz,CD3OD, *ppm*): δ 8.37 (d, J = 2.8 Hz, 1H), 7.97 (d, J = 8.8 Hz, 1H), 7.83 (d, J = 8.3 Hz, 1H), 7.74 (d, J = 8.7 Hz, 1H), 7.51 – 7.40 (m, 2H), 7.34 (dd, J = 8.4, 2.3 Hz, 1H), 7.17 (d, J = 2.4 Hz, 1H), 7.01 (dd, J = 8.8, 2.4 Hz, 1H), 5.13 (dd, J = 12.6, 5.5 Hz, 1H), 4.34 (s, 1H), 4.22 (t, J = 6.2 Hz, 2H), 4.09 (s, 1H), 3.54 (t, J = 4.9 Hz, 4H), 2.98 (t, J = 4.8 Hz, 5H), 2.95 – 2.67 (m, 4), 2.15 (m, 1H), 2.00 – 1.88 (m, 2H), 1.84 – 1.71 (m, 2H), 1.63 (q, J = 7.9 Hz, 2H), 1.31 (s, 6H), 1.23 (s, 6H).	9
	B	798.4	^{1}H NMR (300 MHz, CD_3OD): δ 11.11 (s, 1H), 8.36 (s, 1H), 8.16-7.81 (m, 3H), 7.44 (m, 1H), 7.37-7.22 (m, 2H), 7.21 (m, 1H), 7.03-6.99 (m, 1H), 5.15-5.09 (m, 5.3 Hz, 1H), 4.29 (s, 1H), 4.22-4.10 (m, 3H), 4.05 (d, J = 9.3 Hz, 1H), 3.00-2.80 (m, 3H), 2.64-2.52 (m, 2H), 2.40-2.25 (m, 2H), 2.11-1.70 (m, 9H), 1.70-1.39 (m, 4H), 1.20 (s, 6H), 1.09 (s, 6H);	9, 102



	B	798.3	H-NMR (300 MHz, $CDCl_3$) δ 8.43 (s, 1H), 7.794-7.77 (d, *J*=8.4 Hz, 2H), 7.59-7.52 (m, 2H)7.33-7.32 (d, *J*=2.1 Hz, 1H), 7.21-7.17 (m, 1H), 6.97-6.96 (d, *J*=2.4 Hz, 2H), 6.82-6.78 (m, 1H), 6.54-6.53 (d, *J*=2.1 Hz, 1H), 6.11-6.08 (d, *J*=8.4 Hz, 1H), 5.34 (s, 1H), 4.99-4.93 (m, 1H), 4.12-4.05 (m, 4H), 3.49-3.44 (m, 2H), 2.94-2.72 (m, 7H), 2.48-2.45 (d, *J*=9.9 Hz, 2H), 2.32 (s, 2H), 2.18-2.13 (m, 1H), 1.91-1.85 (m, 4H), 1.60-1.53 (m, 2H), 1.26-1.21 (d, *J*=10.5 Hz, 12H);	9, 103
	B	770.7	H-NMR-PH-ARV-LS-029-E-0(400MHz,CD3OD, *ppm*): δ 8.30 (d, J = 0.8 Hz, 1H), 8.00 (s, 1H), 7.76 (dd, J = 24.6, 8.5 Hz, 2H), 7.40 (d, J = 2.2 Hz, 1H), 7.32 (dd, J = 8.3, 2.3 Hz, 1H), 7.12 (d, J = 2.4 Hz, 1H), 6.97 (dd, J = 8.8, 2.4 Hz, 1H), 5.10 (dt, J = 9.7, 5.2 Hz, 2H), 4.29-4.10 (m, 4H), 3.24 (d, J = 5.7 Hz, 3H), 2.97-2.55 (m, 6H), 2.30 (m, 1H), 2.27 (s, 1H), 2.20-2.05 (m, 1H), 1.88 (dq, J = 27.1, 7.5 Hz, 4H), 1.23 (d, J = 21.8 Hz, 12H).	9, 104
	B	751.51	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.04 (s, 1H), 7.87 (d, *J* = 8.8 Hz, 1H), 7.79-7.60 (m, 3H), 7.51(m, 1H), 7.38-7.35 (m, 3H), 7.34-7.32 (m, 1H), 7.14 (m, 1H), 6.97 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.04 (dd, *J* = 12.7, 5.4 Hz, 1H), 4.29 (s, 1H), 4.03 (d, *J* = 9.1 Hz, 1H), 3.59 (s, 4H), 3.29 (s, 1H), 2.88-2.76 (m, 3H), 2.58-2.52 (m, 7H), 2.07-1.92 (m, 1H), 1.19 (d, *J* = 28.2 Hz, 12H);	1



	B	811.4	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.04 (s, 1H), 7.87 (d, *J* = 8.8 Hz, 1H), 7.79-7.60 (m, 3H), 7.51(m, 1H), 7.38-7.35 (m, 3H), 7.34-7.32 (m, 1H), 7.14 (m, 1H), 6.97 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.04 (dd, *J* = 12.7, 5.4 Hz, 1H), 4.29 (s, 1H), 4.03 (d, *J* = 9.1 Hz, 1H), 3.59 (s, 4H), 3.29 (s, 1H), 2.88-2.76 (m, 3H), 2.58-2.52 (m, 7H), 2.07-1.92 (m, 1H), 1.19 (d, *J* = 28.2 Hz, 12H);	20
	B	812.4	^{1}H NMR (300 MHz, Chloroform-*d*) δ 8.92 (d, *J* = 2.3 Hz, 1H), 8.22 (s, 1H), 8.04 (dd, *J* = 8.1, 2.4 Hz, 1H), 7.62 – 7.45 (m, 2H),7.02 – 6.93 (m, 2H), 6.81 (dd, *J* = 8.8, 2.4 Hz, 1H), 6.24 (d, *J* = 8.2 Hz, 1H), 5.18 (dd, *J* = 13.1, 5.2 Hz, 1H), 4.41- 4.07 (m, 4H), 3.28 (s, 4H), 2.99 – 2.74 (m, 8H), 2.53 (s, 2H), 2.33-2.02 (d, *J* = 6.7 Hz, 3H), 1.78 (s, 3H), 1.40 (s, 4H), 1.26 (d, *J* = 13.8 Hz, 12H), 0.86 (s, 1H).	20




	B	799.3	^{1}H NMR (400 MHz, DMSO-d6): δ 1.15 (s, 6H), 1.24 (s, 6H), 1.78 (t, *J* = 7.6 Hz, 2H), 1.99-2.03 (m, 1H), 2.33-2.35 (m, 2H), 2.54-2.60 (m, 2H), 2.69 (t, *J* = 6.8 Hz, 2H), 2.85-2.83 (m, 1H), 3.33 (s, 4H), 3.45 (s, 4H), 4.09 (d, *J* = 8.8 Hz, 1H), 4.10 (s, 1H), 55.05-5.09 (m, 1H), 7.25-7.23 (m, 5H), 7.41 (d, *J* = 2.0 Hz, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.78 (d, *J* = 7.6 Hz, 3H), 8.10 (d, *J* =12.8 Hz, 1H), 11.09 (s, 1H).	1
	B	777.27	^{1}H NMR (400 MHz, CD_3OD): δ 1.19 (s, 6H), 1.26 (s, 6H), 1.69-1.85 (m, 4H), 2.07-2.13 (m, 1H), 2.65-2.75 (m, 4H), 2.80-2.86 (m, 3H), 2.91 (t, *J* =7.2 Hz, 2H), 3.09 (t, *J* = 5.2 Hz, 2H), 3.52 (s, 2H), 4.14 (s, 1H), 4.25 (s, 1H), 5.09-5.13 (m, 1H), 6.39 (s, 1H), 6.93-6.96 (m, 1H), 7.09 (d, *J* = 2.4 Hz, 1H), 7.43 (d, *J* = 8.0 Hz, 1H), 7.68 (d, *J* = 8.8 Hz, 1H), 7.83 (d, *J* = 8.0 Hz, 1H), 7.87-7.90 (m, 1H), 7.93 (s, 1H), 8.11-8.13 (m, 1H), 8.89 (d, *J* = 2.0 Hz, 1H).	1, 105
	B	807.2	^{1}HNMR (400 MHz, CD_3OD) : δ 1.25 (s, 6H), 1.32 (s, 6H), 1.61-1.64 (m, 2H), 1.78-1.82 (m, 2H), 1.91-1.98 (m, 2H), 2.13-2.16 (m, 1H), 2.76-2.88 (m, 7H), 3.09 (t, *J* = 5.4 Hz, 2H), 3.53 (s, 2H), 4.10-4.12 (m, 1H), 4.22 (t, *J* = 6.2 Hz, 2H), 4.36 (s, 1H), 5.11-5.15 (m, 1H), 6.43 (s, 1H), 7.00-7.03 (m, 1H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.33-7.35 (m, 1H), 7.42 (d, *J* = 2.0 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 1H), 7.82 (d, *J* = 8.4 Hz, 1H), 8.05-	9, 101



			8.11 (m, 2H), 8.78 (s, 1H).	
	B	833.28	^{1}HNMR (400 MHz, CD_3OD): δ 1.20 (s, 6H), 1.27 (s, 6H), 1.57-1.65 (m, 2H), 1.74-1.79 (m, 2H), 1.87-1.94 (m, 2H), 2.07-2.11 (m, 1H), 2.67-2.73 (m, 2H), 2.79-2.85 (m, 1H), 3.13 (t, *J* = 7.8 Hz, 2H), 4.13 (s, 2H), 4.17 (t, *J* = 7.0 Hz, 3H), 4.26 (s, 1H), 4.59 (s, 2H), 5.05-5.09 (m, 1H), 6.94-6.97 (m, 1H), 7.10 (d, *J* = 2.4 Hz, 1H), 7.27-7.30 (m, 1H), 7.37 (d, *J* = 2.4 Hz, 1H), 7.57 (s, 1H), 7.69 (d, *J* = 8.8 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 1H), 8.28-8.31 (m, 1H), 8.34 (br, 1H), 8.80 (d, *J* = 2 Hz, 1H).	86
	B	781.51	^{1}HNMR (400 MHz, CD_3OD): δ 1.20 (s, 6H), 1.27 (s, 6H), 1.57-1.65 (m, 2H), 1.74-1.79 (m, 2H), 1.87-1.94 (m, 2H), 2.07-2.11 (m, 1H), 2.67-2.73 (m, 2H), 2.79-2.85 (m, 1H), 3.13 (t, *J* = 7.8 Hz, 2H), 4.13 (s, 2H), 4.17 (t, *J* = 7.0 Hz, 3H), 4.26 (s, 1H), 4.59 (s, 2H), 5.05-5.09 (m, 1H), 6.94-6.97 (m, 1H), 7.10 (d, *J* = 2.4 Hz, 1H), 7.27-7.30 (m, 1H), 7.37 (d, *J* = 2.4 Hz, 1H), 7.57 (s, 1H), 7.69 (d, *J* = 8.8 Hz, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.97 (d, *J* = 8.8 Hz, 1H), 8.28-8.31 (m, 1H),	17A



			8.34 (br, 1H), 8.80 (d, *J* = 2 Hz, 1H).	
	B	781.25	^{1}H NMR (400 MHz, DMSO-d6): δ 1.11 (d, *J* = 5.6 Hz, 3H), 1.23 (s, 1H), 1.57 (s, 1H), 1.70-1.80 (m, 2H), 1.98-2.08 (m, 2H), 2.22-2.40 (m, 8H), 2.52-2.68 (m, 2H), 2.84-2.94 (m, 1H), 3.30 (s, 1H), 3.51 (s, 2H), 4.18 (t, *J* = 6.4 Hz, 2H), 4.42-4.48 (m, 3H), 5.08-5.15 (m, 1H), 6.43 (s, 1H), 6.94 (d, *J* = 2.4 Hz, 1H), 7.32-7.6 (m, 1H), 7.42 (d, *J* = 2.0 Hz, 1H), 7.80-7.85 (m, 2H), 7.99 (s, 2H), 8.08 (s, 1H), 8.23 (d, *J* = 8.0 Hz, 1H), 11.1 (s, 1H), 13.1 (s, 1H).	106
	B	840.3	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.46-1.55 (m, 4H), 1.79-1.82 (m, 2H), 1.91-1.97 (m, 2H), 2.03-2.07 (m, 1H), 2.30 (br, 2H), 2.58-2.69 (m, 2H), 2.84-3.03 (m, 3H), 3.14-3.18 (m, 1H), 3.30 (s, 1H), 3.42 (br, 1H), 3.72-3.76 (m, 1H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.18-4.21 (m, 2H), 4.30-4.34 (m, 2H), 4.72 (br, 1H), 5.10-5.14 (m, 1H), 6.80-6.82 (m, 1H), 7.00-7.02 (m, 1H), 7.21 (d, *J* = 2.0Hz, 1H), 7.35-7.37 (m ,1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.59 (d, *J* = 8.8 Hz ,1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz,	14



			1H), 7.94 (br, 1H), 8.61 (s, 1H), 11.12 (s, 1H).	
	B	807.28	^{1}H NMR (400 MHz, DMSO-d6): δ 1.14 (s, 6H), 1.24 (s, 6H), 1.45-1.49 (m, 2H), 1.56-1.60 (m, 2H), 1.78-1.82 (m, 2H), 1.99-2.05 (m, 2H), 2.45-2.47 (m, 3H), 2.58-2.67 (m, 5H), 2.84-2.94 (m, 1H), 3.16 (s, 1H), 4.08 (d, *J* = 9.2 Hz, 1H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.32 (s, 1H), 5.10-5.14 (m, 1H), 6.83 (s, 1H), 7.00-7.03 (m, 1H), 7.22 (d, *J* = 2.4Hz ,1H), 7.34-7.36 (m ,1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.64 (d, *J* = 8.0 Hz ,1H), 7.83 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.03 (d, *J* = 9.2 Hz, 1H), 8.13-8.15 (m, 1H), 8.94 (s, 1H), 11.12 (s, 1H).	9, 101
	B	824.27	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.24 (s, 6H), 1.42-1.64 (m, 4H), 1.72-1.87 (m, 2H), 1.99-2.10 (m, 1H), 2.56-2.70 (m, 3H), 2.77-2.95 (m, 3H), 3.22-3.29 (m, 2H), 3.84-4.00 (m, 2H), 4.05-4.13 (m, 1H), 4.15-4.26 (m, 2H), 4.29-4.37 (m, 1H), 5.07-5.17 (m, 1H), 6.96-7.09 (m, 1H), 7.16-7.29 (m, 1H), 7.33-7.50 (m, 2H), 7.76-7.97 (m, 2H), 8.01-8.22 (m, 3H), 8.81-8.91 (m, 1H), 11.07-11.19 (m, 1H).	87




	B	794.29	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.24 (s, 6H), 1.42-1.64 (m, 4H), 1.72-1.87 (m, 2H), 1.99-2.10 (m, 1H), 2.56-2.70 (m, 3H), 2.77-2.95 (m, 3H), 3.22-3.29 (m, 2H), 3.84-4.00 (m, 2H), 4.05-4.13 (m, 1H), 4.15-4.26 (m, 2H), 4.29-4.37 (m, 1H), 5.07-5.17 (m, 1H), 6.96-7.09 (m, 1H), 7.16-7.29 (m, 1H), 7.33-7.50 (m, 2H), 7.76-7.97 (m, 2H), 8.01-8.22 (m, 3H), 8.81-8.91 (m, 1H), 11.07-11.19 (m, 1H).	9, 105
	B	780.27	^{1}H NMR (400 MHz, DMSO-d6): δ 1.14 (s, 6H), 1.23 (s, 6H), 1.75-2.07 (m, 8H), 2.54-2.68 (m, 5H), 2.85-3.17 (m, 4H), 4.06(d, *J* = 9.2 Hz, 1H), 4.27 (br, 2H), 4.32 (s, 1H), 5.11-5.15 (m, 1H), 7.00-7.02 (m, 1H), 7.21 (d, *J* = 2.4Hz ,1H), 7.35-7.39 (m,3H), 7.46 (d, *J* =1.6 Hz, 1H), 7.78 (d, *J* = 8.8 Hz, 3H), 7.86 (d, *J* = 8.0 Hz, 1H), 7.91 (d, *J* = 8.8 Hz, 1H), 11.13 (s.1H).	9, 105



	B	778.3	H-NMR (300 MHz, CD_3OD, ppm) δ 7.78 - 7.68 (m, 4 H), 7.41 - 7.34 (m, 3 H), 7.22 - 7.18 (m, 1 H), 7.13 (s, 1 H), 6.99 - 6.96 (d, *J* = 9 Hz, 1 H), 5.10 - 5.04 (m, 1 H), 4.27 (s, 1 H), 4.13 (s, 1 H), 3.76 - 3.65 (m, 4 H), 3.46 - 3.43 (m, 2 H), 3.32 - 3.31 (m, 2 H), 3.07 - 3.02 (m, 2 H), 2.87 - 2.68 (m, 5 H), 2.13 - 2.08 (m, 1 H), 1.25 (s, 6 H), 1.21 (s, 6 H).	107
	B	810.23		9
	B	739.16	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 2.04-2.16 (m, 3H), 2.56-2.69 (m, 2H), 2.85-2.97 (m, 2H), 3.76-3.79 (m ,2H), 4.05 (d, *J* = 9.2 Hz 1H), 4.16 (t, *J* = 8.4 Hz ,2H), 4.25 (t, *J* = 6.0 Hz ,2H), 4.30 (s, 1H), 5.13 (dd, *J* = 12.8 Hz ,1H), 6.37 (d, *J* = 8.8 Hz ,1H), 7.00 (dd, *J* = 8.8 Hz,1H), 7.20 (d, *J* = 2.4 Hz ,1H), 7.37 (dd, *J* = 8.0 Hz,1H), 7.45 (s, 1H), 7.60 (d, *J* = 9.2 Hz ,1H), 7.84-7.95 (m, 3H), 8.58 (d, *J* = 1.6 Hz ,1H), 11.12 (s, 1H).	9



	B	767.19	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.42-1.46 (m, 2H), 1.66-1.81 (m, 4H), 2.03-2.07 (m, 1H), 2.50-2.62 (m, 2H), 2.73-2.78 (m, 1H), 2.85-2.87 (m, 1H), 3.64 (dd, *J* = 5.6 Hz, 8.0Hz, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.12 (t, *J* = 8.0 Hz, 2H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 5.12 (dd, *J* = 5.2 Hz, 12.8Hz, 1H), 6.36 (d, *J* = 8.8 Hz, 1H), 7.00 (dd, *J* = 2.4 Hz, 8.8Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 2.0 Hz, 8.4Hz, 1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.59 (d, *J* = 9.2 Hz, 1H), 7.83 (d, *J* = 8.0 Hz, 1H), 7.89-7.95 (m, 2H), 8.57 (d, *J* = 2.0 Hz, 1H), 11.12 (s, 1H).	9
	B	818.3	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.42-1.46 (m, 2H), 1.66-1.81 (m, 4H), 2.03-2.07 (m, 1H), 2.50-2.62 (m, 2H), 2.73-2.78 (m, 1H), 2.85-2.87 (m, 1H), 3.64 (dd, *J* = 5.6 Hz, 8.0Hz, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.12 (t, *J* = 8.0 Hz, 2H), 4.20 (t, *J* = 6.4 Hz, 2H), 4.30 (s, 1H), 5.12 (dd, *J* = 5.2 Hz, 12.8Hz, 1H), 6.36 (d, *J* = 8.8 Hz, 1H), 7.00 (dd, *J* = 2.4 Hz, 8.8Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36 (dd, *J* = 2.0	43




			Hz, 8.4Hz, 1H), 7.44 (d, *J* = 2.0 Hz, 1H), 7.59 (d, *J* = 9.2 Hz, 1H), 7.83 (d, *J* = 8.0 Hz, 1H), 7.89-7.95 (m, 2H), 8.57 (d, *J* = 2.0 Hz, 1H), 11.12 (s, 1H).	
	B	836.35	^{1}H NMR (300 MHz, DMSO): δ 11.10 (s, 1H), 8.25 (d, J = 9.2 Hz, 1H), 7.92 (d, J = 8.8 Hz, 1H), 7.82 (d, J = 9.6 Hz, 1H), 7.69 (d, J = 8.5 Hz, 1H), 7.43-7.32 (m, 2H), 7.28-7.25 (m, 2H), 7.05-7.02 (m, 1H), 5.10-5.04 (m, 1H), 4.53-4.44 (m, 3H), 4.01 (d, J = 9.2 Hz, 1H), 3.45 (m, 4H), 3.14-2.79 (m, 3H), 2.64-2.51 (m, 2H), 2.45 (m, 3H), 2.41-2.34 (m, 2H), 2.08-1.97 (m, 1H), 1.88-1.62 (m, 3H), 1.47-1.43 (m, 2H), 1.22-1.08 (m, 15H);	17
	B	836.4	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.08 (s, 1H), 8.59 (d, *J* = 1.2 Hz, 1H), 8.32 (d, *J* = 1.4 Hz, 1H), 7.90 (d, *J* = 8.7 Hz, 1H), 7.80 (d, *J* = 8.9 Hz, 1H), 7.67 (d, *J* = 8.5 Hz, 1H), 7.34 (s, 1H), 7.30 – 7.20 (m, 2H), 7.02 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.07 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.52 – 4.39 (m, 3H), 3.95 (d, *J* = 9.0 Hz, 1H), 3.43 (s, 4H), 3.06 – 2.79 (m, 5H), 2.36 (d, *J* = 7.6 Hz, 2H), 2.07 (s, 2H), 1.80 (d, *J* = 13.0 Hz, 2H), 1.48 – 1.36 (m, 4H), 1.15 (d, *J* = 18.7 Hz, 16H)	17



	B	781.51	^{1}H NMR (400 MHz, DMSO-d6): δ 1.41-1.42 (m, 3H), 1.63-1.65 (m, 2H), 1.99-2.03 (m, 1H), 2.30-2.32 (m, 2H), 2.44 (br, 4H), 2.53-2.71 (m, 2H), 2.84-2.92 (m, 1H), 3.10-3.14 (m, 1H), 3.16-3.17 (m, 1H), 3.34-3.38 (m, 4H), 3.57-3.62 (m, 1H), 4.10-4.29 (m, 2H), 4.33-4.47 (m, 2H), 4.56-4.58 (m, 1H), 5.02-5.10 (m, 1H), 6.51 (br, 1H), 6.93-6.94 (m, 1H), 7.22-7.34 (m, 2H), 7.65-7.67 (m, 1H), 7.83 (s, 1H), 7.98 (s, 2H), 8.08 (s, 1H), 8.24-8.26 (m, 1H), 8.75 (br, 1H), 11.10 (s, 1H), 13.30 (br, 1H).	17A, 108
	B	840.58	^{1}H NMR (400 MHz, CD_3OD): δ 1.23 (s, 6H), 1.31 (s, 6H), 1.53-1.73 (m, 4H), 1.88-1.95 (m, 2H), 2.11-2.17 (m, 1H), 2.55-2.80 (m, 5H), 2.84-2.94 (m, 2H), 2.05-2.09 (m, 1H), 3.12-3.22 (m, 2H), 3.63-3.85 (m, 4H), 4.08 (s, 1H), 4.20 (t, *J* = 6.2 Hz, 2H), 4.34 (s, 1H), 5.10-5.15 (m, 1H), 7.00-7.03 (m, 1H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.32 -7.34 (m, 1H), 7.40-7.43 (m, 2H), 7.74 (d, *J* = 8.4 Hz, 1H), 7.81 (d, *J* = 8.4 Hz, 1H), 7.94 (d, *J* = 8.8 Hz, 1H), 8.34 (d, *J* = 2.4 Hz, 1H).	14




	B	822.35	^{1}H NMR (400 MHz, DMSO): δ 11.07 (s, 1H), 8.23 (d, J = 9.2 Hz, 1H), 7.89 (d, J = 8.7 Hz, 1H), 7.80 (d, J = 9.6 Hz, 1H), 7.67 (d, J = 8.5 Hz, 1H), 7.39-7.30 (m, 2H), 7.24 (m, 2H), 7.01 (m, 1H), 5.06 (m, 1H), 4.53-4.42 (m, 3H), 3.99 (m, 1H), 3.43 (m, 4H), 3.08-2.96 (m, 2H), 2.87 (m, 1H), 2.61-2.49 (m, 2H), 2.51 (m, 4H), 2.19 (m, 2H), 2.04-1.87 (m, 2H), 1.83 (m, 2H), 1.16 (m, 14H);	49
	B	792.4	^{1}H-NMR (300 MHz, DMSO) δ11.07(s, 1H), 8.64-8.60(m, 1H), 7.98- 7.87(m, 2H), 7.69- 7.55(m, 2H), 7.35- 7.20(m, 3H), 7.04- 6.96(m, 1H), 6.90- 6.83(m , 1H), 5.11 - 5.01 (m, 1H), 4.30(s, 1H), 4.09- 4.02(m, 1H), 3.70- 3.61(m, 4H), 3.58- 3.49(m, 4H), 2.97-2.79(m, 1H), 2.65- 2.53(m, 2H), 2.05 - 1.97(m, 1H), 1.61- 1.55(m, 8H), 1.21 - 1.12 (d, *J=29.4 Hz,* 12H).	19
	B	793.35	^{1}H-NMR (300 MHz, DMSO) δ11.07(s, 1H), 8.64-8.60(m, 1H), 7.98- 7.87(m, 2H), 7.69- 7.55(m, 2H), 7.35- 7.20(m, 3H), 7.04- 6.96(m, 1H), 6.90- 6.83(m , 1H), 5.11 - 5.01 (m, 1H), 4.30(s, 1H), 4.09- 4.02(m, 1H), 3.70- 3.61(m, 4H), 3.58- 3.49(m, 4H), 2.97-2.79(m, 1H), 2.65- 2.53(m, 2H), 2.05 - 1.97(m, 1H), 1.61- 1.55(m, 8H), 1.21 - 1.12 (d, *J=29.4 Hz,* 12H).	19



	B	781.3	^{1}H NMR (300 MHz, CD_3OD) 7.75-7.70 (m, 3H), 7.48-7.44 (m, 1H), 7.36-7.33 (m, 2H), 7.16-7.12 (m, 2H), 6.99-6.96 (m, 1H), 5.08-5.02 (m, 1H), 4.63-4.61 (m, 1H), 4.28-4.26 (m, 1H), 4.14-4.11 (m, 1H), 3.73-3.55 (m, 3H), 3.07-3.02 (m, 1H), 2.86-2.64 (m, 8H), 2.11-1.92 (m, 3H), 1.27-1.22 (m, 12H);	1, 16
	B	817.51	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.92-2.09 (m, 3H), 2.50 (s, 4H), 2.51-2.73 (m, 4H), 2.84-2.93 (m, 1H), 3.28 (br, 4H), 4.00 (d, *J* = 9.6 Hz, 1H), 4.23-4.26 (m, 3H), 5.10-5.15 (m, 1H), 6.67 (d, *J* = 12.0 Hz, 2H), 6.98-7.01 (m, 1H), 7.22 (d, *J* = 2.33 Hz, 1H), 7.35-7.38 (m, 1H), 7.45 (d, *J* = 2.0 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.18 (d, *J* = 9.6 Hz, 1H), 11.12 (s, 1H).	9




	B	845.55	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.41-1.58 (m, 4H), 1.77-1.81 (m, 2H), 2.01-2.07 (m, 1H), 2.28-2.39 (m, 2H), 2.42-2.49 (m, 4H), 2.54-2.62 (m, 2H), 2.84-2.91 (m, 1H), 3.23 (br, 4H), 4.00 (d, *J* = 9.2 Hz, 1H), 4.19 (t, *J* = 6.4 Hz, 2H), 4.25 (s, 1H), 5.09-5.14 (m, 1H), 6.67 (d, *J* = 11.2 Hz, 2H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.34-7.36 (m, 1H), 7.43 (d, *J* = 2.0 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.18 (d, *J* = 9.2 Hz, 1H), 11.12 (s, 1H).	9
	B	813.53	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.10 (s, 6H), 1.17 (s, 6H), 1.41-1.58 (m, 4H), 1.77-1.81 (m, 2H), 2.01-2.07 (m, 1H), 2.28-2.39 (m, 2H), 2.42-2.49 (m, 4H), 2.54-2.62 (m, 2H), 2.84-2.91 (m, 1H), 3.23 (br, 4H), 4.00 (d, *J* = 9.2 Hz, 1H), 4.19 (t, *J* = 6.4 Hz, 2H), 4.25 (s, 1H), 5.09-5.14 (m, 1H), 6.67 (d, *J* = 11.2 Hz, 2H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.34-7.36 (m, 1H), 7.43 (d, *J* = 2.0 Hz, 1H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.18 (d, *J* = 9.2 Hz, 1H), 11.12 (s, 1H).	9



	B	890.57	^{1}H NMR (400 MHz, DMSO-d6): δ 1.00-1.07 (m, 2H), 1.09 (s, 6H), 1.18 (s, 6H), 1.82-1.88 (m, 2H), 1.96-2.05 (m, 2H), 2.17-2.25 (m, 2H), 2.53-2.64 (m, 2H), 2.84-2.92 (m, 1H), 2.98-3.06 (m, 2H), 3.30-3.33 (m, 4H), 3.40-3.50 (m, 4H), 3.99 (d, *J* = 9.2 Hz, 1H), 4.26 (s, 1H), 4.63-4.72 (m, 2H), 5.05-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.22-7.28 (m, 2H), 7.33-7.37 (m, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.24 (d, *J* = 9.2 Hz, 1H), 8.60 (s, 1H), 11.09 (s, 1H).	49 starting with ethyl 2-chloro-4-(trifluoromethyl)pyrimidine-5-carboxylate
	B	813.54		
	B	888.58	^{1}H NMR (400 MHz, DMSO-d6): δ 1.00-1.07 (m, 2H), 1.09 (s, 6H), 1.18 (s, 6H), 1.82-1.88 (m, 2H), 1.96-2.05 (m, 2H), 2.17-2.25 (m, 2H), 2.53-2.64 (m, 2H), 2.84-2.92 (m, 1H), 2.98-3.06 (m, 2H), 3.30-3.33 (m, 4H), 3.40-3.50 (m, 4H), 3.99 (d, *J* = 9.2 Hz, 1H), 4.26 (s, 1H), 4.63-4.72 (m, 2H), 5.05-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.22-7.28 (m, 2H), 7.33-7.37 (m, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.89 (d, *J* = 8.8 Hz, 1H), 8.24	49 starting with 4-fluoro-3-(trifluoromethyl)benzoic acid




			(d, *J* = 9.2 Hz, 1H), 8.60 (s, 1H), 11.09 (s, 1H).	
	B	827.56		
	B	862.64		



	B	822.4	H-NMR (300 MHz, DMSO) δ 11.04 (s, 1H), 8.55 (s, 2H), 8.10-8.07 (d, J=9 Hz, 1H), 7.89-7.86 (d, J=9 Hz, 1H), 7.67-7.65 (d, J=8.4 Hz, 1H), 7.32 (s, 1H), 7.26-7.22 (m, 2H), 7.03-6.99 (m, 1H), 5.08-5.02 (m, 1H), 4.40 (s, 1H), 4.01-3.92 (m, 3H), 3.43 (s, 4H), 2.95-2.80 (m, 3H), 2.59 (s, 6H), 2.21-2.19 (d, J=6.3 Hz, 2H), 2.05-1.98 (m, 1H), 1.83-1.79 (d, J=11.1 Hz, 3H), 1.18 (s, 6H), 1.11 (s, 6H);	49 starting with 5-fluoropyrimidine-2-carbonitrile
	B	836.5	H-NMR (300 MHz, DMSO) δ 11.04 (s, 1H), 8.55 (s, 2H), 8.09-8.06 (d, J=9 Hz, 1H), 7.89-7.86 (d, J=9 Hz, 1H), 7.67-7.64 (d, J=8.4 Hz, 1H), 7.32 (s, 1H), 7.25-7.22 (m, 2H), 7.03-6.99 (m, 1H), 5.08-5.02 (m, 1H), 4.40 (s, 1H), 4.00-3.92 (m, 3H), 3.42 (s, 4H), 2.91-2.83 (m, 3H), 2.59-2.54 (m, 6H), 2.39-2.25 (m, 2H), 2.05-1.98 (m, 1H), 1.80-1.76 (d, J=12 Hz, 2H), 1.58 (s,1H), 1.46-1.39 (m,2H),1.28-1.24 (d, J=11.1 Hz, 8H), 1.18 (s, 6H);	17 starting with 5-fluoropyrimidine-2-carbonitrile
	B	838.2	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.06 (s, 1H), 7.87 (d, *J* = 8.8 Hz, 1H), 7.69 (dd, *J* = 15.0, 10.0 Hz, 3H), 7.44 (dd, *J* = 18.6, 8.3 Hz, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.02 – 6.89 (m, 3H), 5.07 (dd, *J* = 12.9, 5.3 Hz, 1H), 4.28 (s, 1H), 4.07 – 3.95 (m, 1H), 3.77 (m, 2H),3.31 – 3.18 (m, 4H), 2.95 – 2.76 (m, 3H), 2.62 – 2.46 (m, 6H), 2.21 (d, *J* = 6.8 Hz, 2H), 1.99 (d, *J* = 19.3 Hz, 1H), 1.82 (d, *J* = 13.2 Hz, 3H), 1.34 (d, *J* = 27.3 Hz,	49




			8H) , 1.11 (d, *J* = 27.3 Hz, 6H).	
	B	778.3	HNMR (300 MHz, DMSO-*d6*, ppm) δ 11.08 (s, 1 H), 7.92-7.89 (d, *J*= 8.7Hz, 1H), 7.76-7.68 (m, 3H), 7.44 - 7.213 (m, 4H), 7.02 - 7.01 (d, *J*= 2.4Hz, 1H), 6.47-6.44 (d, *J*= 8.7 Hz, 2H), 5.08-5.04 (m, 1H), 4.32(s, 1H), 4.07-4.02 (m, 3H), 3.77-3.73 (m, 2H), 3.49-3.46 (m, 4H), 3.32-3.31 (m, 1H), 2.96-2.82 (m, 1H), 2.60-2.55 (m, 6H), 2.09-1.95 (m, 1H), 1.22 (s, 6H), 1.12 (s, 6H);	19
	B	822.4	^{1}H NMR (400 MHz, CD_3OD): δ7.82-7.80 (d, *J*=8.0Hz, 1H), 7.75-7.70 (m, 3H), 7.28-7.22 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.02 (m, 2H), 4.28 (s, 1H), 4.13 (s, 1H), 3.94-3.90 (m, 2H), 3.38-3.30(m, 6H), 2.82-2.70 (m, 5H), 2.67-2.64(m, 4H), 2.50-2.47 (m, 2H), 2.13-2.11 (m, 1H), 1.27 (s, 6H), 1.21 (s, 6H);	23



	B	836.59	^{1}H NMR (400 MHz, CD_3OD): δ7.82-7.80 (d, *J*=8.0Hz, 1H), 7.75-7.70 (m, 3H), 7.28-7.22 (m, 2H), 7.13-7.12 (d, *J*=2.4Hz, 1H), 6.99-6.96 (m, 3H), 5.12-5.02 (m, 2H), 4.28 (s, 1H), 4.13 (s, 1H), 3.94-3.90 (m, 2H), 3.38-3.30(m, 6H), 2.82-2.70 (m, 5H), 2.67-2.64(m, 4H), 2.50-2.47 (m, 2H), 2.13-2.11 (m, 1H), 1.27 (s, 6H), 1.21 (s, 6H);	49 starting with ethyl 2-chloro-4-methylpyrimidine-5-carboxylate
	B	850.6	^{1}HNMR (400 MHz, DMSO-d6): δ 1.10 (s, 6H), 1.18 (s, 6H), 1.25 (d, *J* = 6.4 Hz, 2H), 1.80 (d, *J* = 13.2 Hz, 2H), 1.91 (s, 1H), 2.01-2.04 (m, 1H), 2.18-2.22 (m, 8H), 2.55-2.61 (m, 3H), 2.83-2.89 (m, 3H), 3.34 (s, 2H), 3.45 (br, 4H), 4.08 (d, *J* = 9.6 Hz, 1H), 4.26 (s, 1H), 4.72 (d, *J* = 12.4 Hz, 2H), 5.06-5.10 (m, 1H), 6.99-7.02 (m, 1H), 7.22-7.27 (m, 2H), 7.35 (s, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.8 Hz, 1H), 8.23 (d, *J* = 9.2 Hz, 1H), 11.10 (s, 1H).	49 starting with ethyl 2-chloro-4-methylpyrimidine-5-carboxylate
	B	795.55		



	B	843.55		
	B	823.6	1H NMR (300 MHz, DMSO-d6) δ 11.07 (s, 1H), 7.83 (dd, J = 20.2, 8.5 Hz, 2H), 7.69 (d, J = 8.6 Hz, 2H), 7.54-7.36 (m, 2H), 7.32 (dd, J = 8.3, 2.3 Hz, 1H), 7.17 (d, J = 2.3 Hz, 1H), 7.02-6.86 (m, 3H), 5.08 (dd, J = 12.8, 5.3 Hz, 1H), 4.28 (s, 1H), 4.16 (t, J = 6.3 Hz, 2H), 4.01 (d, J = 9.1 Hz, 1H), 3.66 (d, J = 11.5 Hz, 2H), 2.94 – 2.83 (m, 1H), 2.65 (dt, J = 29.5, 6.6 Hz, 5H), 2.53 (d, J = 3.1 Hz, 3H), 2.07-1.94 (m, 1H), 1.74-1.63 (m, 2H), 1.48 (p, J = 8.0 Hz, 2H), 1.18 (s, 6H), 1.14-0.98 (m, 12H)..	9



	B	792.54	H-NMR (300 MHz, DMSO) δ 11.06 (s, 1H), 7.98-7.89 (d, J=9 Hz, 1H), 7.77-7.74 (d, J=8.7 Hz, 2H), 7.65-7.62 (d, J=8.4 Hz, 1H), 7.52-7.49 (d, J=9.3 Hz, 1H), 7.21-7.20 (d, J=2.4 Hz, 1H), 7.02-76.96 (m, 3H), 6.79 (s, 1H), 6.67-6.64 (d, J=8.1 Hz, 1H),5.08-5.02 (m, 1H), 4.32 (s, 1H), 4.18-4.04 (m, 3H), 3.74-3.70 (m, 2H), 3.30-3.26 (d, J=12 Hz, 1H), 3.05 (s, 1H), 2.94-2.82 (m, 1H), 2.73-2.27 (m, 11H), 2.03-1.99 (m,1H), 1.22 (s, 6H), 1.13 (s, 6H);	65
	B	820.57	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.03 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.7 Hz, 2H), 7.62 (d, *J* = 8.5 Hz, 1H), 7.47 (d, *J* = 9.2 Hz, 1H), 7.31 (m, 1H), 7.20 – 7.13 (m, 2H), 7.02 – 6.89 (m, 3H), 5.23 (dd, *J* = 12.6, 5.4 Hz, 1H), 4.93 (dd, *J* = 12.6, 5.4 Hz, 3H), 3.31 – 3.17 (m, 3H), 3.02 – 2.76 (m, 3H), 2.62 – 2.49 (m, 6H), 2.17 (d, *J* = 6.7 Hz, 2H), 1.98 (dt, *J* = 10.9, 5.3 Hz, 1H), 1.90 – 1.74 (m, 3H), 1.14 (d, *J* = 27.4 Hz, 14H).	49, 70
	B	840.57	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.03 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.7 Hz, 2H), 7.62 (d, *J* = 8.5 Hz, 1H), 7.47 (d, *J* = 9.2 Hz, 1H), 7.31 (m, 1H), 7.20 – 7.13 (m, 2H), 7.02 – 6.89 (m, 3H), 5.23 (dd, *J* = 12.6, 5.4 Hz, 1H), 4.93 (dd, *J* = 12.6, 5.4 Hz, 3H), 3.31 – 3.17 (m, 3H), 3.02 – 2.76 (m, 3H), 2.62 – 2.49 (m, 6H), 2.17 (d, *J* = 6.7 Hz, 2H), 1.98 (dt, *J* = 10.9, 5.3 Hz, 1H), 1.90 – 1.74 (m, 3H), 1.14 (d, *J* = 27.4 Hz, 14H).	14




	B	814.53	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.19 (s, 6H), 1.62-1.67 (m, 2H), 1.77-1.84 (m, 2H), 2.00-2.07 (m, 1H), 2.24-2.44 (m, 6H), 2.54-2.62 (m, 2H), 2.86-2.90 (m, 1H), 3.60 (br, 4H), 3.96 (d, *J* = 9.6 Hz, 1H), 4.20-4.24 (m, 2H), 4.32 (s, 1H), 5.08-5.13 (m, 1H), 6.74-6.77 (m, 1H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.36-7.39 (m ,1H), 7.44-7.45 (m, 1H), 7.51-7.55 (m, 1H), 7.84 (d, *J* = 8.4 Hz ,1H), 7.91 (d, *J* = 8.8 Hz, 1H), 8.39 (d, *J* = 10.4 Hz 1H), 11.11 (s, 1H).	109
	B	787.49	^{1}H NMR (400 MHz, DMSO-*d6*): δ 1.11 (s, 6H), 1.18 (s, 6H), 1.98-2.07 (m, 1H), 2.51-2.64 (m, 6H), 2.83-2.94 (m, 3H), 3.45 (s, 4H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.27 (s, 1H), 5.02-5.13 (m, 1H), 6.97-7.05 (m, 1H), 7.12 (d, *J* = 8.8 Hz, 2H), 7.23-7.29 (m, 2H), 7.36 (s, 1H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.90 (d, *J* = 8.4 Hz, 1H), 8.50 (d, *J* = 9.2 Hz, 1H), 11.10 (s, 1H).	110



	B	752.49	^{1}HNMR (400 MHz, CD_3OD): δ 1.12 (s, 6H), 1.18 (s, 6H), 1.78-1.79 (m, 4H), 1.92-2.11 (m, 2H), 2.61-2.67 (m, 2H), 2.71-2.77 (m, 2H), 3.50 (t, *J* = 6.0 Hz, 2H), 3.97-4.04 (m, 3H), 4.09 (br, 2H), 4.18 (s, 1H), 4.92-5.03 (m, 1H), 6.35 (d, *J* = 8.4 Hz, 2H), 6.86-6.89 (m, 1H), 7.02 (d, *J* = 2.4 Hz, 1H), 7.21-7.23 (m, 1H), 7.29-7.31 (m, 2H), 7.59-7.63 (m, 3H), 7.70 (d, *J* = 8.0 Hz, 1H).	111
	B	821.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.73-1.77 (m, 2H), 1.99-2.02 (m, 1H), 2.06-2.10 (m, 2H), 2.54-2.60 (m, 2H), 2.83-2.98 (m, 3H), 3.27-3.34 (m, 5H), 3.45-3.47 (m, 4H), 4.03-4.07 (m, 3H), 4.19 (s, 1H), 4.32 (s, 1H), 5.04-5.08 (m, 1H), 6.55 (d, *J* =8.8 Hz, 2H), 6.99-7.02 (m, 1H),7.20-7.23 (m, 2H), 7.29 (d, *J* =1.6 Hz, 1H), 7.40 (d, *J* = 9.2 Hz, 1H), 7.64 (d, *J* = 8.4 Hz, 1H), 7.74 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.08 (s, 1H).	112
	B	806.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.22 (s, 6H), 1.72-2.17 (m, 4H), 2.34-2.42 (m, 3H), 2.58-2.71 (m, 2H), 2.84-2.93 (m, 1H), 3.14-3.20 (m, 2H), 3.28-3.33 (m, 6H), 3.43-3.60 (m, 3H), 4.05-4.07 (m, 1H), 4.32 (s, 1H), 5.04-5.08 (m, 1H), 6.81-7.01 (m, 5H), 7.21 (s, 1H), 7.53 (d, *J* = 11.6 Hz, 1H), 7.64 (d, *J* = 7.2 Hz, 1H), 7.76 (d, *J* = 6.8 Hz, 2H), 7.90 (d, *J* = 7.2 Hz, 1H), 11.08 (s, 1H).	53




	B	806.56		53
	B	779.55		



	B	821.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 9H), 1.73-1.74 (m, 2H), 1.80-1.84 (m, 1H), 1.99-2.02 (m, 1H), 2.06-2.08 (m, 2H), 2.33-2.39 (m, 3H), 2.54-2.60 (m, 2H), 2.83-2.98 (m, 3H), 3.27-3.27 (m, 2H), 3.33-3.34 (m, 2H), 3.46-3.50 (m, 1H), 4.03-4.07 (m, 3H), 4.19 (s, 1H), 4.32 (s, 1H), 5.04-5.08 (m, 1H), 6.55 (d, *J* = 8.8 Hz, 2H), 6.99-7.02 (m, 1H), 7.20-7.23 (m, 2H), 7.30 (s, 1H), 7.40 (d, *J* = 9.2 Hz, 1H), 7.64 (d, *J* = 8.8 Hz, 1H), 7.75 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.4 Hz, 1H), 11.12 (s, 1H).	113
	B	811.4	^{1}H-NMR (300 MHz, CD3OD) δ8.71(s, 2H), 7.75-7.67(m, 2H), 7.21- 7.17(m, 1H), 7.16- 7.08(m, 2H), 7.01- 6.93(m, 1H), 5.16- 5.06 (m, 1H), 4.26(s, 1H), 4.12(s, 1H), 4.02- 3.66(m, 6H), 3.11- 3.3.03(m, 2H), 2.94- 2.77(m, 3H), 2.75- 2.58(m, 1H), 2.48- 2.29(m, 1H), 2.16 - 2.04(m, 1H), 1.82- 1.65(m, 2H), 1.27 - 1.20 (d, *J=20.7 Hz,* 12H), 1.06- 0.09(m, 6H)	9




	B	806.55	^{1}HNMR (400 MHz, DMSO-d_6) : δ 1.13 (s, 6H), 1.22 (s, 6H), 1.78-1.84 (m, 1H), 2.02-2.05 (m, 1H), 2.25-2.27 (m, 1H), 2.55-2.62 (m, 2H), 2.85-2.93 (m, 2H), 3.08-3.21 (m, 3H), 3.32-3.47 (m, 6H), 3.60-3.73 (m, 3H), 4.07-4.08 (m, 1H), 4.25-4.28 (m, 2H), 4.33 (s, 2H), 5.09-5.13 (m, 1H), 6.57 (d, *J* = 8.8 Hz, 2H), 7.00-7.02 (m, 1H), 7.20-7.21 (m, 1H), 7.38-7.45 (m, 2H), 7.52 (s, 1H), 7.78 (d, *J* = 8.4 Hz, 3H), 7.91 (d, *J* = 8.4 Hz, 1H), 9.97 (br, 1H), 11.12 (s, 1H).	53
	B	824.55	^{1}H NMR (400 MHz, DMSO-d_6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.71-1.79 (m, 1H), 2.00-2.05 (m, 1H), 2.12-2.15 (m, 1H), 2.42 (d, *J* = 6.4 Hz, 2H), 2.57-2.61 (m, 6H), 2.84-2.93 (m, 1H), 3.05 (t, *J* = 7.2 Hz, 1H), 3.26-3.31 (m, 6H), 3.38-3.40 (m, 1H), 3.46 (t, *J* = 8.4 Hz, 1H), 4.05 (d, *J* = 8.8 Hz, 1H), 4.32 (s, 1H), 5.09-5.13 (m, 1H), 6.54 (d, *J* = 8.8 Hz, 2H), 6.99-7.01 (m, 1H), 7.20 (d, *J* = 2.0 Hz, 1H), 7.39 (d, *J* =9.2 Hz, 1H), 7.46 (d, *J* =3.6 Hz, 1H), 7.72-7.75 (m, 3H), 7.90 (d, *J* = 8.4 Hz, 1H), 11.12 (s, 1H).	53



	B	765.56	^{1}H NMR (400 MHz, DMSO-d^6) δ 1.12 (6H, s), 1.22 (6H, s), 1.50-1.53 (2H, m), 1.63-1.66 (2H, m), 1.99-2.02 (1H, m), 2.34-2.47 (7H, m), 2.55-2.59 (1H, m), 2.66-2.70 (2H, m), 2.88-2.97 (1H, m), 3.25-3.31 (4H, m), 4.05 (1H, d, *J* = 9.2 Hz), 4.30-4.34 (2H, m), 4.46-4.50 (1H, m), 5.14 (1H, dd, *J* = 13.2, 4.8 Hz), 6.95-7.02 (3H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.46-7.48 (2H, m), 7.51-7.57 (2H, m), 7.74 (2H, d, *J* = 8.8 Hz), 7.91 (1H, d, *J* = 8.8 Hz), 11.02 (1H, s).	8A
	B	736.5	^{1}H NMR (400 MHz, DMSO-d^6) δ 1.12 (6H, s), 1.22 (6H, s), 1.50-1.53 (2H, m), 1.63-1.66 (2H, m), 1.99-2.02 (1H, m), 2.34-2.47 (7H, m), 2.55-2.59 (1H, m), 2.66-2.70 (2H, m), 2.88-2.97 (1H, m), 3.25-3.31 (4H, m), 4.05 (1H, d, *J* = 9.2 Hz), 4.30-4.34 (2H, m), 4.46-4.50 (1H, m), 5.14 (1H, dd, *J* = 13.2, 4.8 Hz), 6.95-7.02 (3H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.46-7.48 (2H, m), 7.51-7.57 (2H, m), 7.74 (2H, d, *J* = 8.8 Hz), 7.91 (1H, d, *J* = 8.8 Hz), 11.02 (1H, s).	113
	B	779.54	^{1}H NMR (400 MHz, $CDCl_3$) *δ* 1.13 (6H, s), 1.22 (6H, s), 1.50-1.55 (2H, m), 1.62-1.67 (2H, m), 2.04-2.08 (1H, m), 2.33-2.36 (2H, m), 2.47-2.50 (4H, m), 2.55-2.62 (2H, m), 2.86-2.90 (1H, m), 3.07 (2H, t, *J* = 7.2 Hz), 3.23 (4H, s), 4.05 (1H, d, *J* = 9.2 Hz), 4.32 (1H, s), 5.14 (1H, dd, *J* = 12.8, 5.6 Hz), 6.95-7.02 (3H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.52 (1H, d, *J* = 9.2 Hz), 7.71-7.81 (5H, m), 7.91 (1H, d, *J* = 8.8 Hz), 11.13 (1H, s).	8A



	B	798.2	^{1}H NMR (400 MHz, CD_3OD): δ 8.51 (s, 1H), 7.71-7.66 (m, 2H), 7.35-7.34 (d, *J*=1.6Hz, 1H), 7.23-7.21 (d, *J*=8.4Hz, 1H), 7.10-7.09 (d, *J*=2.0Hz, 1H), 6.96-6.93 (m, 1H), 5.09-5.04 (m, 1H), 4.19 (s, 1H), 3.98 (s, 1H), 3.52-3.31 (m, 7H), 3.06-3.02 (m, 1H), 2.87-2.81 (m, 1H), 2.76-2.51 (m, 10H), 2.42-2.31 (m, 4H), 2.11-2.10 (m, 3H), 1.90 (m, 1H), 1.45-1.35 (m, 2H), 1.20 (s, 6H), 1.10 (s, 6H);	114
	B	822.03	^{1}H NMR (400 MHz, CD_3OD): δ 8.51 (s, 1H), 7.71-7.66 (m, 2H), 7.35-7.34 (d, *J*=1.6Hz, 1H), 7.23-7.21 (d, *J*=8.4Hz, 1H), 7.10-7.09 (d, *J*=2.0Hz, 1H), 6.96-6.93 (m, 1H), 5.09-5.04 (m, 1H), 4.19 (s, 1H), 3.98 (s, 1H), 3.52-3.31 (m, 7H), 3.06-3.02 (m, 1H), 2.87-2.81 (m, 1H), 2.76-2.51 (m, 10H), 2.42-2.31 (m, 4H), 2.11-2.10 (m, 3H), 1.90 (m, 1H), 1.45-1.35 (m, 2H), 1.20 (s, 6H), 1.10 (s, 6H);	49



	B	821.55	^{1}H-NMR (300 MHz, CD_3OD) δ 7.82- 7.68 (m, 4H), 7.28- 7.10 (m, 3H), 7.02- 6.94 (m, 1H), 6.64- 6.57 (m, 2H), 5.14 - 5.05 (m, 1H), 4.94- 4.90 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.69- 3.35 (m, 3H), 3.22- 3.06 (m, 2H), 2.96- 2.57 (m, 4H), 2.56- 2.24 (m, 6H), 2.24- 2.03 (m, 5H), 1.92- 1.74 (m, 1H), 1.28- 1.22 (d, *J=18 Hz*, 12H);	73
	B	821.56	^{1}H-NMR (300 MHz, CD_3OD) δ 7.82- 7.68 (m, 4H), 7.28- 7.10 (m, 3H), 7.02- 6.94 (m, 1H), 6.64- 6.57 (m, 2H), 5.14 - 5.05 (m, 1H), 4.94- 4.90 (m, 1H), 4.28 (s, 1H), 4.13 (s, 1H), 3.69- 3.35 (m, 3H), 3.22- 3.06 (m, 2H), 2.96- 2.57 (m, 4H), 2.56- 2.24 (m, 6H), 2.24- 2.03 (m, 5H), 1.92- 1.74 (m, 1H), 1.28- 1.22 (d, *J=18 Hz*, 12H);	113
	B	821.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.22 (s, 6H), 1.52-1.54 (m, 2H), 1.81-1.83 (m, 1H), 1.86-1.94 (m, 3H), 1.98-2.03 (m, 1H), 2.16-2.20 (m, 1H), 2.55-2.60 (m, 2H), 2.70 (s, 2H), 2.84-2.93 (m, 1H), 3.03-3.08 (m, 1H), 3.18-3.22 (m, 1H), 3.29-3.32 (m, 2H), 3.48-3.55 (m, 3H), 3.57-3.63 (m, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.32 (s, 1H), 5.05 (d, *J* = 12.8 Hz, 1H), 6.81 (d, *J* = 8.4 Hz, 1H), 6.90 (d, *J* = 1.6 Hz, 1H), 6.96-7.02 (m, 3H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.50 (d, *J* = 9.2	113



			Hz, 1H), 7.63(d, *J* = 8.8 Hz, 1H), 7.73(d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.09 (s, 1H).	
	B	807.54	^{1}H NMR (400 MHz, *d6*-DMSO): δ11.11 (s, 1H), 7.91-7.73 (m, 4H), 7.44-7.01 (m, 5H), 6.43 (s, 2H), 5.12-4.96 (m, 2H), 4.31 (s, 1H), 4.04-3.98 (m, 3H), 3.53-3.33 (m, 3H), 3.08 (m, 1H), 3.01-2.92 (m, 2H), 2.61-2.30 (m, 5H), 2.23-2.08 (m, 6H), 1.21 (s, 6H), 1.12 (s, 6H);	73
	B	821.45	^{1}H-NMR (300 MHz, CD_3OD) δ7.82- 7.68 (m, 4H), 7.28- 7.10 (m, 3H), 7.02- 6.94 (m, 1H), 6.64- 6.57 (m, 2H), 5.14 - 5.05 (m, 1H), 5.01- 4.90 (m, 1H), 4.28 (s, 1H), 4.15- 4.10 (m, 1H), 3.60- 3.33 (m, 3H), 3.21-3.09 (m, 2H), 2.92- 2.30 (m, 10H), 2.28- 2.05 (m, 5H), 1.90-1.73 (m, 1H), 1.28- 1.22 (d, *J=18 Hz,* 12H).	73




	B	839.56	^{1}H NMR (400 MHz, DMSO-d6): δ 1.00-1.08 (m, 2H), 1.12 (s, 6H), 1.19 (s, 6H), 1.73-1.94 (m, 3H), 1.99-2.06 (m, 1H), 2.19 (d, J = 6.8 Hz, 2H), 2.46-2.49 (m, 4H), 2.54-2.67 (m, 2H), 2.84-2.96 (m, 3H), 3.38-3.49 (m, 4H), 3.95 (d, J = 8.8 Hz, 1H), 4.31-4.40 (m, 3H), 5.05-5.10 (m, 1H), 6.71, 6.75 (two singles, 1H), 7.00-7.02 (m, 1H), 7.22-7.27 (m, 2H), 7.35 (s, 1H), 7.46-7.49 (m, 1H), 7.68 (d, J = 8.4 Hz, 1H), 7.89 (d, J = 8.8 Hz, 1H), 8.39 (d, J = 11.2 Hz, 1H), 11.10 (s, 1H).	49, 109
	B	852.59	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.47-1.51 (m, 3H), 1.67-1.78 (m, 3H), 1.99-2.03 (m, 2H), 2.23 (s, 3H), 2.43-2.45 (m, 2H), 2.53-2.60 (m, 3H), 2.73-2.92 (m, 3H), 3.32 (s, 1H), 3.55-3.65 (m, 3H), 3.89-3.92 (m, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.31 (s, 1H), 5.04-5.09 (m, 1H), 6.94-7.04 (m, 4H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.41 (d, *J* = 9.2 Hz, 1H), 7.60 (d, *J* = 12.4 Hz, 1H), 7.73 (d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.10 (s, 1H).	115
	B	852.59	^{1}H NMR (400 MHz, DMSO-d6): δ 1.12 (s, 6H), 1.21 (s, 6H), 1.48-1.49 (m, 3H), 1.67-1.78 (m, 3H), 1.99-2.06 (m, 2H), 2.23 (s, 3H), 2.43-2.45 (m, 2H), 2.53-2.60 (m, 3H), 2.73-2.92 (m, 3H), 3.32 (s, 1H), 3.55-3.65 (m, 3H), 3.89-3.92 (m, 2H), 4.05 (d, *J* = 9.2 Hz, 1H), 4.31 (s, 1H), 5.04-5.09 (m, 1H), 6.94-7.04 (m, 4H), 7.21 (d, *J* = 2.0 Hz, 1H), 7.51 (d, *J* = 9.2 Hz, 1H), 7.60 (d, *J* = 12.8 Hz, 1H), 7.73 (d, *J* = 8.8 Hz, 2H), 7.90	115



			(d, *J* = 8.8 Hz, 1H), 11.10 (s, 1H).	
	B	861.7	^{1}H NMR (300 MHz, DMSO-d_6): δ 11.08 (s, 1H), 8.61 (d, J = 2.3 Hz, 1H), 7.91 (dd, J = 11.7, 8.8 Hz, 2H), 7.70 (d, J = 8.4 Hz, 1H), 7.60 (d, J = 9.3 Hz, 1H), 7.38 (s, 1H), 7.28 (d, J = 8.8 Hz, 1H), 7.20 (d, J = 2.4 Hz, 1H), 7.00 (dd, J = 8.8, 2.5 Hz, 1H), 6.87 (d, J = 9.1 Hz, 1H), 5.07 (dd, J = 12.8, 5.3 Hz, 1H), 4.43 (dd, J = 25.1, 12.8 Hz, 2H), 4.30 (s, 1H), 4.04 (d, J = 9.1 Hz, 1H), 2.98 - 2.82 (m, 7H), 2.62 (m, 2H),2.51 (m, 1H), 2.01-1.75 (m, 5H), 1.64 (d, J = 11.7 Hz, 1H), 1.50-1.11 (m, 21H).	94
	B	856.56	H-NMR-PH-ARV-LS-049-E-0: (300 MHz, DMSO-*d6*, ppm) δ 11.11 (br, 1H), 7.92-7.89 (d, *J*=8.7Hz, 1H), 7.75-7.61 (m, 4H), 7.47-7.45 (d, *J* =7.5Hz, 1H),7.22-7.21 (d, *J* =2.4Hz, 1H), 7.12-7.06 (m, 1H), 7.03-6.99 (m, 1H), 5.14-5.08 (m, 1H), 4.32 (s, 1H), 4.08-4.05 (d, *J* =9.3Hz, 1H), 3.51-3.47 (m, 2H), 3.26-3.24 (m, 4H), 2.89-2.82 (m, 1H), 2.78-2.71 (m, 2H), 2.62-2.55 (m, 6H), 2.27-2.25 (m, 2H), 2.08-2.04 (m, 1H), 1.86-1.73 (m, 3H), 1.31-1.28 (m, 2H), 1.22 (s, 6H), 1.13 (s, 6H);	49



	B	806.55	H-NMR (300 MHz, DMSO-*d6*, ppm) δ 11.11 (br, 1H), 7.92-7.89 (d, *J*=8.7Hz, 1H), 7.75-7.61 (m, 4H), 7.47-7.45 (d, *J* =7.5Hz, 1H),7.22-7.21 (d, *J* =2.4Hz, 1H), 7.12-7.06 (m, 1H), 7.03-6.99 (m, 1H), 5.14-5.08 (m, 1H), 4.32 (s, 1H), 4.08-4.05 (d, *J* =9.3Hz, 1H), 3.51-3.47 (m, 2H), 3.26-3.24 (m, 4H), 2.89-2.82 (m, 1H), 2.78-2.71 (m, 2H), 2.62-2.55 (m, 6H), 2.27-2.25 (m, 2H), 2.08-2.04 (m, 1H), 1.86-1.73 (m, 3H), 1.31-1.28 (m, 2H), 1.22 (s, 6H), 1.13 (s, 6H);	52
	B	824.4	1H NMR (300 MHz, DMSO-*d*6) δ 11.04 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 (m, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.06 – 6.89 (m, 4H), 5.03 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.28 (s, 1H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.78-3.56 (m, 3H), 3.35-3.26 (m, 4H), 2.94 – 2.75 (m, 1H), 2.60 – 2.48 (m, 5H), 2.45-2.37 (m, 3H), 2.32 – 2.28 (m, 2H), 2.19-1.98 (m, 2H), 1.88-1.75 (m, 1H), 1.18 (s, 6H), 1.09 (s, 6H).	53



	B	840.55	1H NMR (300 MHz, DMSO-*d*6) δ 11.04 (s, 1H), 7.87 (d, *J* = 8.7 Hz, 1H), 7.72 (d, *J* = 8.6 Hz, 2H), 7.52 (m, 2H), 7.17 (d, *J* = 2.4 Hz, 1H), 7.06 – 6.89 (m, 4H), 5.03 (dd, *J* = 12.8, 5.4 Hz, 1H), 4.28 (s, 1H), 4.02 (d, *J* = 9.1 Hz, 1H), 3.78-3.56 (m, 3H), 3.35-3.26 (m, 4H), 2.94 – 2.75 (m, 1H), 2.60 – 2.48 (m, 5H), 2.45-2.37 (m, 3H), 2.32 – 2.28 (m, 2H), 2.19-1.98 (m, 2H), 1.88-1.75 (m, 1H), 1.18 (s, 6H), 1.09 (s, 6H).	49
	B	938.66	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.22 (6H, s), 1.29-1.34 (2H, m), 1.68-1.73 (1H, m), 1.80-1.83 (2H, m), 1.99-2.03 (1H, m), 2.24 (2H, d, *J* = 6.8 Hz), 2.51-2.61 (8H, m), 2.84-2.92 (1H, m), 3.25 (3H, s), 3.46-3.48 (6H, m), 3.55-3.57 (2H, m), 3.60-3.64 (2H, m), 3.78-3.80 (2H, m), 4.05 (1H, d, *J* = 9.6 Hz), 4.14-4.16 (2H, m), 4.32 (1H, s), 5.05-5.10 (1H, m), 6.93 (1H, d, *J* = 8.4 Hz), 6.99-7.02 (1H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.25-7.28 (1H, m), 7.32-7.35 (2H, m), 7.44 (1H, d, *J* = 8.4 Hz), 7.60 (1H, d, *J* = 9.2 Hz), 7.69 (1H, d, *J* = 8.4 Hz), 7.90 (1H, d, *J* = 8.8 Hz), 11.09 (1H, s).	52, 116




	B	777.56	^{1}H NMR (400 MHz, DMSO-d_6) δ 1.13 (6H, s), 1.22 (6H, s), 1.29-1.34 (2H, m), 1.68-1.73 (1H, m), 1.80-1.83 (2H, m), 1.99-2.03 (1H, m), 2.24 (2H, d, *J* = 6.8 Hz), 2.51-2.61 (8H, m), 2.84-2.92 (1H, m), 3.25 (3H, s), 3.46-3.48 (6H, m), 3.55-3.57 (2H, m), 3.60-3.64 (2H, m), 3.78-3.80 (2H, m), 4.05 (1H, d, *J* = 9.6 Hz), 4.14-4.16 (2H, m), 4.32 (1H, s), 5.05-5.10 (1H, m), 6.93 (1H, d, *J* = 8.4 Hz), 6.99-7.02 (1H, m), 7.21 (1H, d, *J* = 2.4 Hz), 7.25-7.28 (1H, m), 7.32-7.35 (2H, m), 7.44 (1H, d, *J* = 8.4 Hz), 7.60 (1H, d, *J* = 9.2 Hz), 7.69 (1H, d, *J* = 8.4 Hz), 7.90 (1H, d, *J* = 8.8 Hz), 11.09 (1H, s).	117
	B	821.56	^{1}H NMR (400 MHz, $CDCl_3$) δ 1.12 (6H, s), 1.26 (6H, s), 1.49-1.54 (4H, m), 1.89-1.91 (4H, m), 1.99-2.02 (1H, m), 2.54-2.67 (1H, m), 2.85-2.89 (1H, m), 3.04 (2H, t, *J* = 10.0 Hz), 3.23-3.26 (3H, m), 3.64-3.83 (6H, m), 4.04 (1H, d, *J* = 9.2 Hz), 4.32 (1H, s), 5.04 (1H, dd, *J* = 5.2, 12.8 Hz), 6.96-7.02 (3H, d, m), 7.20 (1H, d, *J* = 2.4 Hz), 7.24 (1H, d, *J* = 10.4 Hz), 7.33 (1H, s), 7.49 (1H, d, *J* = 8.8 Hz), 7.65 (1H, d, *J* = 8.8 Hz), 7.72 (2H, d, *J* = 8.8 Hz), 7.89 (1H, d, *J* = 8.8 Hz), 11.08 (1H, s).	52, 118



	B	839.35	^{1}H-NMR (400 MHz, CD_3OD) δ8.54（s, 1H), 7.77-7.64(m, 2H), 7.36(s, 1H), 7.26- 7.12(m, 2H), 7.02-6.96(m, 1H), 6.65- 6.57(m , 1H), 5.12 - 5.04 (m, 1H), 4.28(s, 1H), 4.12- 4.04(m, 3H), 3.71(m, 4H), 3.08-2.98(m, 2H), 2.90- 2.55(m, 7H), 2.37- 2.31(m, 2H), 2.15- 1.93(m, 4H), 1.37- 1.30(m, 2H), 1.28 - 1.23 (d, J=21 Hz, 12H).	52, 109
	B	763.5		
	B	818.55	^{1}H NMR (400 MHz, DMSO-d6): δ 1.13 (s, 6H), 1.23 (s, 6H), 1.88-1.92 (m, 1H), 1.99-2.03 (m, 1H), 2.18 (t, J = 8 Hz, 1H), 2.25 (br, 2H), 2.46 (br, 3H), 2.55-2.60 (m, 1H), 2.70 (s, 1H), 2.84-2.89 (m, 1H), 2.96 (s, 2H), 3.46-3.47 (m, 5H), 3.79 (s, 2H), 4.06 (d, *J* = 9.2 Hz, 1H), 4.35 (s, 1H), 5.05-5.11 (m, 1H), 5.76 (s, 1H), 6.96 (d, *J* = 8.8 Hz, 2H), 6.99-7.02 (m, 1H), 7.21 (d, *J* = 2.4 Hz, 1H), 7.26 (d, *J* = 8.4 Hz, 1H), 7.34 (s, 1H), 7.53 (d, *J* = 8.4 Hz, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.78	119

			(d, *J* = 8.8 Hz, 2H), 7.90 (d, *J* = 8.8 Hz, 1H), 11.08 (s, 1H).	
	B	821.2	^{1}H NMR (300 MHz, $CDCl_3$) 7.74-7.67 (m, 3H), 7.58-7.55(m, 1H), 7.29-7.26 (m, 1H), 7.04-7.08 (m, 1H), 6.97-6.93 (m, 3H), 6.83-6.81 (m, 1H), 6.14-6.11 (m, 1H), 4.97-4.91 (m, 1H), 4.42 (s, 1H), 4.16-4.13 (m, 1H), 4.05-3.99 (m, 3H), 3.04-2.92 (m, 4H), 2.88-2.51 (m, 3H) , 2.16-2.10 (m, 3H), 1.96-1.88 (m, 3H), 1.84-1.73(m, 2H) ,1.67-1.65 (m, 4H), 1.46 (m, 6H), 1.27-1.22 (m, 6H);	120
	B	868.3	^{1}H NMR (400 MHz, $CDCL_3$): δ 8.04 (s, 1H), 7.55 (d, J = 8.7 Hz, 1H), 7.41 (dd, *J* = 25.5, 9.1 Hz, 2H), 6.93 (d, *J* = 2.4 Hz, 1H), 6.77 (dd, *J* = 8.7, 2.4 Hz, 1H), 5.66 (d, *J* = 8.5 Hz, 1H), 4.97- 4.78 (m, 3H), 4.14 (d, *J* = 8.5 Hz, 1H), 3.99 (s, 1H), 3.27 (s, 4H), 2.96-2.63 (m, 6H), 2.59 (s, 4H), 2.33 (s, 6H), 2.13 (dd, *J* = 14.4, 6.4 Hz, 1H), 1.83 (d, *J* = 13.2 Hz, 3H), 1.20 (s, 12H), 1.11 (s, 2H).	63

	B	786.2	^{1}H NMR (400 MHz, $CDCL_3$): δ 8.04 (s, 1H), 7.55 (d, J = 8.7 Hz, 1H), 7.41 (dd, *J* = 25.5, 9.1 Hz, 2H), 6.93 (d, *J* = 2.4 Hz, 1H), 6.77 (dd, *J* = 8.7, 2.4 Hz, 1H), 5.66 (d, *J* = 8.5 Hz, 1H), 4.97- 4.78 (m, 3H), 4.14 (d, *J* = 8.5 Hz, 1H), 3.99 (s, 1H), 3.27 (s, 4H), 2.96-2.63 (m, 6H), 2.59 (s, 4H), 2.33 (s, 6H), 2.13 (dd, *J* = 14.4, 6.4 Hz, 1H), 1.83 (d, *J* = 13.2 Hz, 3H), 1.20 (s, 12H), 1.11 (s, 2H).	49
	B	854.2	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.11 (d, *J* = 8.6 Hz, 1H), 7.72 (dd, *J* = 17.8, 8.5 Hz, 3H), 7.52 (d, *J* = 9.2 Hz, 1H), 7.44-7.22 (m, 4H), 6.96 (d, *J* = 8.6 Hz, 2H), 5.08 (dd, *J* = 12.7, 5.4 Hz, 1H), 4.40 (s, 1H), 4.08 (d, *J* = 9.1 Hz, 1H), 3.95-3.75 (m, 2H), 3.60-3.40 (m, 4H), 3.38-3.20 (m, 4H), 2.95-2.70 (m, 3H), 2.68-2.59 (m, 2H), 2.32 (brs, 2H), 2.04-2.02 (m, 1H), 1.82-1.70 (m, 3H), 1.19 (d, *J* = 26.9 Hz, 14H).	49
	B	778.5	^{1}H NMR (300 MHz, *d6*-DMSO) *δ* 1.13 (s, 6H), 1.22 (s, 6H), 1.89-2.08 (m, 5H), 2.31-2.40 (m, 3H), 2.51-2.64 (m, 2H), 2.82-2.98 (m, 1H), 3.80 (s, 2H), 3.92 (s, 2H), 4.04-4.07 (m, 1H), 4.14 (s, 2H), 4.32 (s, 1H), 5.10-5.16 (m, 1H), 6.40-6.42 (m, 2H), 6.99-7.02 (m, 1H), 7.20 (s, 1H), 7.34-7.36 (m, 1H), 7.43-7.46 (m, 2H), 7.72-7.74 (m, 2H), 7.83-7.92 (m, 2H), 11.1 (s, 1H);	121




	B	811.32	1H NMR (300 MHz, CDCL3) δ 8.29 (s, 1H), 7.81 (d, J = 8.3 Hz, 1H), 7.60 (d, J = 8.6 Hz, 1H), 7.36 (d, J = 2.2 Hz, 1H), 7.22 (dd, J = 8.3, 2.3 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.83 (dd, J = 8.8, 2.4 Hz, 1H), 5.96 (d, J = 8.6 Hz, 1H), 4.99 (dd, J = 11.9, 5.5 Hz, 1H), 4.24 (dt, J = 11.9, 7.0 Hz, 7H), 4.08 (s, 1H), 3.09 (s, 9H), 2.98 – 2.77 (m, 5H), 2.40(s, 2H), 2.21- 2.14 (m, 1H), 1.27 (d, J = 3.9 Hz, 12H);	9, 63
	B	783.1	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.13 (s, 1H), 8.65 (s, 1H), 8.38 (s, 1H), 8.15 (s, 1H), 7.96-7.81 (m, 3H), 7.47 (d, *J* = 2.2 Hz, 1H), 7.38 (dd, *J* = 8.4, 2.3 Hz, 1H), 7.26 (d, *J* = 2.4 Hz, 1H), 7.04 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.14 (dd, *J* = 12.9, 5.4 Hz, 1H), 4.45 (s, 1H), 4.28 (t, *J* = 6.1 Hz, 2H), 3.98 (d, *J* = 9.0 Hz, 1H), 3.74 (s, 3H), 2.89 (d, *J* = 12.5 Hz, 1H), 2.60 (d, *J* = 15.9 Hz, 4H), 2.04 (s, 4H), 1.17 (d, *J* = 18.4 Hz, 12H), 1.05 (s, 4H);	9, 63

	B	783.1	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.27 – 8.12 (m, 1H), 7.92 – 7.76 (m, 3H), 7.46 – 7.29 (m, 3H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.09 (dd, *J* = 12.9, 5.3 Hz, 1H), 4.43 (s, 1H), 4.23 (t, *J* = 6.3 Hz, 2H), 3.97 (d, *J* = 9.1 Hz, 1H), 3.70 (t, *J* = 4.9 Hz, 4H), 2.95 – 2.76 (m, 1H), 2.59-2.50 (s, 6H),2.45(s,2H), 2.07 – 1.89 (m, 3H), 1.19 (s, 6H), 1.11 (s, 6H).	9, 63
	B	854.3	^{1}H NMR (300 MHz, DMSO-d_6) δ 11.09 (s, 1H), 8.27 – 8.12 (m, 1H), 7.92 – 7.76 (m, 3H), 7.46 – 7.29 (m, 3H), 7.22 (d, *J* = 2.4 Hz, 1H), 7.00 (dd, *J* = 8.8, 2.4 Hz, 1H), 5.09 (dd, *J* = 12.9, 5.3 Hz, 1H), 4.43 (s, 1H), 4.23 (t, *J* = 6.3 Hz, 2H), 3.97 (d, *J* = 9.1 Hz, 1H), 3.70 (t, *J* = 4.9 Hz, 4H), 2.95 – 2.76 (m, 1H), 2.59-2.50 (s, 6H),2.45(s,2H), 2.07 – 1.89 (m, 3H), 1.19 (s, 6H), 1.11 (s, 6H).	49
	B	872.7	H-NMR (400 MHz, DMSO-*d6*) 11.12 (s, 1H),7.91-7.86 (m, 2H),7.75-7.72 (d, *J* = 11.6Hz, 1H), 7.47-7.45 (d, *J*=7.6Hz, 1H), 7.34-7.32 (d, *J* = 8Hz, 1H), 7.22-7.21 (d, *J* = 4Hz, 1H), 7.01-6.91 (m, 3H), 5.14 -5.09 (m, 1H), 4.27 (s, 1H), 4.01- 3.98 (d, *J* = 9.2Hz, 1H), 3.84-3.81 (d, *J* = 12.8Hz, 2H), 3.26 (s, 4H), 2.94-2.85 (m, 1H), 2.81-2.755 (m, 2H), 2.54-2.48 (m, 6H), 2.22-2.21 (d, *J* = 6Hz, 2H), 2.08-2.03 (m, 1H), 1.81-1.78 (m, 3H), 1.28 (s, 6H), 1.21-1.12 (m, 2H), 1.05 (s, 6H)	49

	C	811.4	^{1}H NMR (300 MHz, CD_3OD): δ 7.79-7.68 (m, 2H), 7.37 (s, 1H),7.31-7.28 (m, 1H), 7.13-7.11 (m, 1H), 6.99-6.94 (m, 1H), 5.10-5.04 (m, 1H), 4.29 (s, 1H), 4.18-4.14 (m, 2H), 4.06 (s, 1H), 3.51-3.45 (m, 4H), 2.80-2.60 (m, 7H), 2.57-2.48 (m, 2H), 2.13-2.02 (m, 1H), 1.92-1.80 (m, 2H), 1.75-1.46 (m, 4H), 1.26 (s, 6H),1.19 (s, 6H);	9
	C	844.32		

C	D	D

	D	D



D	D	D
		

	D	D

	D			
	D			
	D			

		729.18
	D	D

	D	766.27	
	D	800.3	
	D		

	D	848.2	H-NMR (300 MHz, CDCl3) δ 8.04-7.95 (m, 3H), 7.84-7.77 (m, 2H), 7.60-7.55 (m, 1H), 7.34-7.33 (d, J=1.8 Hz, 1H), 7.21-7.09 (m, 3H), 4.98-4.93 (m, 1H), 4.14-4.10 (m, 2H), 3.91 (s, 2H), 3.49 (s, 2H), 2.94-2.72 (m, 3H), 2.67-2.56 (m, 5H), 2.25-2.13 (m, 2H), 1.92-1.88 (m, 2H), 1.76 (s, 2H), 1.68 (s, 6H);	122
	D			

	D	824.57	
	D		
	D		

	838.57	781.52
	D	D

	856.57	
D	D	D

		736.49
	D	D

	D	791.54	
	D		
	D		

	D

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2017/056234

A. CLASSIFICATION OF SUBJECT MATTER

C07D 233/86(2006.01)i, C07D 401/04(2006.01)i, C07D 491/107(2006.01)i, C07D 471/10(2006.01)i, A61K 31/4164(2006.01)i, A61K 31/4439(2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D 233/86; C07D 495/14; C07D 403/06; C07D 413/14; A61K 47/48; A61K 38/05; C07D 401/04; C07D 491/107; C07D 471/10; A61K 31/4164; A61K 31/4439

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
Korean utility models and applications for utility models
Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
eKOMPASS(KIPO internal), STN (Registry, Caplus), Google & Keywords: androgen receptor, ubiquintination, linker, protein, polypeptide, E3 ubiquitin ligase, cereblon, thalidomide, lenalidomide, pomalidomide, degradation, inhibition

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2016-0058872 A1 (ARVINAS, INC.) 3 March 2016 See claims 1-4; and pages 92, 94.	1-7,9
X	WO 2016-118666 A1 (ARVINAS, INC.) 28 July 2016 See claims 1-7.	1-3
A	WINTER, G. E. et al., "Phthalimide conjugation as a strategy for in vivo target protein degradation", SCIENCE, 19 June 2015, Vol. 348, Issue 6241, pages 1376-1381 See abstract; and figures 1-2.	1-7,9
A	US 2014-0356322 A1 (YALE UNIVERSITY et al.) 4 December 2014 See claim 147; and claim 172, lines 8-9.	1-7,9
A	WO 02-00617 A2 (BRISTOL-MYERS SQUIBB COMPANY) 3 January 2002 See pages 34-44, table 1.	1-7,9

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:
"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier application or patent but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed
"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents,such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
12 February 2018 (12.02.2018)	**12 February 2018 (12.02.2018)**

Name and mailing address of the ISA/KR
International Application Division
Korean Intellectual Property Office
189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea
Facsimile No. +82-42-481-8578

Authorized officer
KIM, Sun Hee
Telephone No. +82-42-481-5405

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2017/056234

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 31-34
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 31-34 pertain to methods for treatment of the human body by therapy and thus relate to a subject matter which this International Searching Authority is not required to search under PCT Article 17(2)(a)(i) and Rule 39.1(iv).

2. ☒ Claims Nos.: 13,28-29,32-33
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims 13,28-29 and 32-33 refer to claims which are not drafted in accordance with the third sentence of Rule 6.4(a).

3. ☒ Claims Nos.: 8,10-12,14-27,30-31,34
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required addtional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/056234

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2016-0058872 A1	03/03/2016	AU 2015-247817 A1	27/10/2016
		AU 2016-270442 A1	07/12/2017
		CA 2945975 A1	22/10/2015
		CN 106458993 A	22/02/2017
		EP 3131588 A2	22/02/2017
		EP 3131588 A4	10/01/2018
		JP 2017-513862 A	01/06/2017
		KR 10-2017-0002446 A	06/01/2017
		MX 2016013563 A	09/05/2017
		US 2015-0291562 A1	15/10/2015
		WO 2015-160845 A2	22/10/2015
		WO 2015-160845 A3	10/12/2015
		WO 2016-197032 A1	08/12/2016
WO 2016-118666 A1	28/07/2016	AU 2016-209349 A1	10/08/2017
		CA 2974367 A1	28/07/2016
		CN 107428734 A	01/12/2017
		EP 3247708 A1	29/11/2017
		KR 10-2017-0103977 A	13/09/2017
		MX 2017009454 A	20/10/2017
		US 2016-0214972 A1	28/07/2016
		US 2017-0327469 A1	16/11/2017
US 2014-0356322 A1	04/12/2014	AU 2013-207900 A1	24/07/2014
		AU 2013-207900 A8	31/07/2014
		AU 2013-207900 B2	07/12/2017
		CA 2861066 A1	18/07/2013
		CN 104736569 A	24/06/2015
		EP 2802608 A2	19/11/2014
		EP 2802608 A4	05/08/2015
		HK 1208875 A1	18/03/2016
		JP 2015-508414 A	19/03/2015
		KR 10-2015-0010935 A	29/01/2015
		MX 2014008520 A	14/08/2015
		RU 2014133019 A	27/02/2016
		WO 2013-106643 A2	18/07/2013
		WO 2013-106643 A3	06/09/2013
WO 02-00617 A2	03/01/2002	AR 031709 A1	01/10/2003
		AR 035340 A1	12/05/2004
		AR 037950 A1	22/12/2004
		AT 318822 T	15/03/2006
		AT 411319 T	15/10/2008
		AT 420880 T	15/01/2009
		AU 1560902 A	08/01/2002
		AU 2001-269943 B2	09/11/2006
		AU 2001-288213 B2	14/04/2005
		AU 2002-359727 A1	09/07/2003
		AU 2002-359727 A8	09/07/2003

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/056234

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		AU 2002-361785 B2	22/05/2008
		AU 6700801 A	08/01/2002
		AU 6994301 A	02/04/2002
		AU 8821301 A	08/01/2002
		BG 107675 A	31/12/2003
		BR 0111298 A	10/05/2005
		BR 0111869 A	23/09/2003
		BR 0113980 A	24/06/2003
		BR 0215281 A	19/10/2004
		CA 2413417 A1	03/01/2002
		CA 2413596 A1	03/01/2002
		CA 2413683 A1	03/01/2002
		CA 2423071 A1	28/03/2002
		CA 2471342 A1	31/07/2003
		CN 1307181 C	28/03/2007
		CN 1443187 A	17/09/2003
		CN 1454083 A	05/11/2003
		CN 1589271 A	02/03/2005
		CN 1608069 A	20/04/2005
		CN 1995039 A	11/07/2007
		CO 5640082 A2	31/05/2006
		CZ 20024214 A3	16/04/2003
		CZ 20024250 A3	15/10/2003
		CZ 20030780 A3	14/01/2004
		DE 60117551 T2	18/01/2007
		DK 1319007 T3	19/06/2006
		DK 1458723 T3	09/02/2009
		EC SP024410 A	31/03/2003
		EE 200300108 A	15/02/2005
		EP 1297108 A2	02/04/2003
		EP 1299094 A2	09/04/2003
		EP 1299385 A2	09/04/2003
		EP 1319007 A1	18/06/2003
		EP 1319007 B1	01/03/2006
		EP 1319007 B9	10/10/2007
		EP 1458723 A1	22/09/2004
		EP 1458723 A4	21/12/2005
		EP 1458723 B1	15/10/2008
		EP 1463728 A2	06/10/2004
		EP 1463728 A4	21/12/2005
		EP 1463728 B1	14/01/2009
		EP 1854798 A2	14/11/2007
		EP 1854798 A3	28/11/2007
		ES 2260244 T3	01/11/2006
		ES 2314127 T3	16/03/2009
		ES 2319511 T3	08/05/2009
		GE P20063817 B	10/05/2006
		GE P20074144 B	10/07/2007
		HK 1054230 A1	28/07/2006
		HR P20030305 A2	30/04/2005

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/056234

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		HR P20030305 B1	31/12/2007
		HR P20030305 B9	31/03/2008
		HR P20040567 A2	30/06/2005
		HU 0303165 A2	29/12/2003
		HU 0303165 A3	29/05/2006
		HU 0303172 A2	29/12/2003
		HU 0303172 A3	29/05/2006
		HU 0400455 A2	28/06/2004
		HU 0400455 A3	29/10/2007
		HU 0402554 A2	29/03/2005
		HU 0402554 A3	28/01/2009
		HU 0402672 A2	28/04/2005
		HU 0402672 A3	29/05/2006
		IS 6750 A	18/03/2003
		IS 7324 A	18/06/2004
		JP 2004-509072 A	25/03/2004
		JP 2004-509895 A	02/04/2004
		JP 2004-515462 A	27/05/2004
		JP 2004-517606 A	17/06/2004
		JP 2005-523257 A	04/08/2005
		JP 2005-524610 A	18/08/2005
		JP 4966477 B2	04/07/2012
		KR 10-0765670 B1	10/10/2007
		KR 10-2003-0014407 A	17/02/2003
		KR 10-2003-0016310 A	26/02/2003
		KR 10-2003-0028847 A	10/04/2003
		KR 10-2004-0086248 A	08/10/2004
		MX PA02012563 A	14/05/2003
		MX PA02012573 A	14/05/2003
		MX PA02012605 A	14/05/2003
		MX PA03002412 A	05/05/2004
		MX PA04005876 A	16/05/2005
		MY 138531 A	30/06/2009
		NO 20026167 A	20/12/2002
		NO 20026194 A	26/02/2003
		NO 20031266 A	13/05/2003
		NO 20043068 A	08/09/2004
		NO 329931 B1	24/01/2011
		NZ 523802 A	25/11/2005
		NZ 524803 A	24/09/2004
		NZ 533471 A	31/05/2007
		PE 01862003 A1	23/03/2003
		PE 07292002 A1	11/08/2002
		PL 361707 A1	04/10/2004
		PL 366565 A1	07/02/2005
		PL 370904 A1	30/05/2005
		PL 371685 A1	27/06/2005
		PT 1319007 E	31/05/2006
		PT 1458723 E	26/12/2008
		RU 2003101918 A	27/08/2004

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		RU 2003102390 A	27/05/2004
		RU 2004122403 A	27/05/2005
		RU 2298554 C2	10/05/2007
		RU 2330038 C2	27/07/2008
		SI 1319007 T1	31/08/2006
		SI 1458723 T1	28/02/2009
		SK 4982003 A3	04/05/2004
		TR 200202702 T2	21/12/2004
		TW 200304810 A	16/10/2003
		TW I263640 B	11/10/2006
		TW I305208 B	11/01/2009
		UA 78265 C2	15/03/2007
		UA 78686 C2	25/04/2007
		US 2002-0058290 A1	16/05/2002
		US 2002-0173445 A1	21/11/2002
		US 2003-0114420 A1	19/06/2003
		US 2004-0077605 A1	22/04/2004
		US 2004-0077606 A1	22/04/2004
		US 2004-0176324 A1	09/09/2004
		US 2005-0119228 A1	02/06/2005
		US 2005-0192253 A1	01/09/2005
		US 2005-0256048 A1	17/11/2005
		US 2005-0272799 A1	08/12/2005
		US 2005-0282813 A1	22/12/2005
		US 2006-0223832 A1	05/10/2006
		US 2006-0264459 A1	23/11/2006
		US 2008-0214643 A1	04/09/2008
		US 6953679 B2	11/10/2005
		US 6960474 B2	01/11/2005
		US 7001911 B2	21/02/2006
		US 7141578 B2	28/11/2006
		US 7432267 B2	07/10/2008
		US 7470797 B2	30/12/2008
		US 7517904 B2	14/04/2009
		US 7655689 B2	02/02/2010
		UY 26808 A1	31/01/2002
		UY 26809 A1	31/01/2002
		UY 27595 A1	31/07/2003
		WO 02-00617 A3	30/01/2003
		WO 02-00653 A2	03/01/2002
		WO 02-00653 A3	25/07/2002
		WO 02-00716 A2	03/01/2002
		WO 02-00716 A3	13/06/2002
		WO 02-24702 A1	28/03/2002
		WO 03-053354 A2	03/07/2003
		WO 03-053354 A3	27/11/2003
		WO 03-062241 A1	31/07/2003
		WO 03-062241 A8	21/10/2004
		YU 20003 A	03/03/2006
		YU 53604 A	17/08/2006

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/056234

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		ZA 200209530 B	23/02/2004
		ZA 200210313 B	19/03/2004
		ZA 200302963 B	15/07/2004
		ZA 200404812 B	26/08/2005