

[19] Patents Registry
The Hong Kong Special Administrative Region
香港特別行政區
專利註冊處

[11] 1151036 B
EP 2261249 B1

[12]

STANDARD PATENT SPECIFICATION
標準專利說明書

[21] Application No. 申請編號
11104992.7

[51] Int.C1.⁸ C07K C12N A61K A61P

[22] Date of filing 提交日期
15.05.2009

[22] Date Lodged 交件日期
20.05.2011

[54] KDR PEPTIDES AND VACCINES COMPRISING THE SAME KDR 肽及含有 KDR 肽的疫苗

[30] Priority 優先權

12.09.2002 JP 2002267285

07.03.2003 JP 2003062003

11.06.2003 JP 2003167042

[43] Date of publication of application 申請發表日期
20.01.2012

[45] Publication of the grant of the patent 批予專利的發表日期
07.08.2015

EP Application No. & Date 歐洲專利申請編號及日期

EP 10010173.2 12.09.2003

EP Publication No. & Date 歐洲專利申請發表編號及日期

EP 2261249 15.12.2010

Date of Grant in Designated Patent Office 指定專利當局批予專利日期
11.02.2015

[62] Derived from Application No. 09104475.7 under section 22
of the Patents Ordinance (Chapter 514)
根據專利條例 (第 514 章) 第 22 條, 此為申請編號
09104475.7 的分開申請

[73] Proprietor 專利所有人
OncoTherapy Science, Inc.
2-1, Sakado 3-chome Takatsu-ku
Kawasaki-shi, Kanagawa 213-0012
JAPAN

[72] Inventor 發明人
Tahara, Hideaki
Wada, Satoshi
Tsunoda, Takuya

[74] Agent and / or address for service 代理人及/或送達地址
Ella Cheong (Hong Kong) Limited
5001 Hopewell Centre
183 Queen's Road East
HONG KONG



(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
11.02.2015 Bulletin 2015/07

(21) Application number: **10010173.2**

(22) Date of filing: **12.09.2003**

(51) Int Cl.:
C07K 14/71 ^(2006.01) **C12N 15/12** ^(2006.01)
C12N 5/00 ^(2006.01) **A61K 38/17** ^(2006.01)
A61K 39/00 ^(2006.01) **A61P 3/10** ^(2006.01)
A61P 9/10 ^(2006.01) **A61P 17/06** ^(2006.01)
A61P 27/02 ^(2006.01) **A61P 29/00** ^(2006.01)
A61P 35/00 ^(2006.01) **C07K 7/06** ^(2006.01)

(54) **KDR peptides and vaccines comprising the same**

KDR-Peptide und Impfstoffe damit

Peptides KDR et vaccins les comportant

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**

(30) Priority: **12.09.2002 JP 2002267285**
07.03.2003 JP 2003062003
11.06.2003 JP 2003167042

(43) Date of publication of application:
15.12.2010 Bulletin 2010/50

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
08018875.8 / 2 014 678
03795432.8 / 1 548 032

(73) Proprietor: **Oncotherapy Science, Inc.**
Kawasaki-shi
Kanagawa 213-0012 (JP)

(72) Inventors:
• **Tahara, Hideaki**
Tokyo 113-8654 (JP)
• **Wada, Satoshi**
Saitama-shi
Saitama 338-8553 (JP)
• **Tsunoda, Takuya**
Tokyo 108-0071 (JP)

(74) Representative: **Vossius & Partner**
Siebertstrasse 4
81675 München (DE)

(56) References cited:

EP-A1- 1 055 684 WO-A-98/11223
WO-A-99/43801

- **NORGREN R B ET AL:** "Kinase insert domain receptor (fragment)", EMBL, 1 June 2002 (2002-06-01), XP002348370,
- **HIDA N ET AL:** "A simple culture protocol to detect peptide-specific cytotoxic T lymphocyte precursors in the circulation", **CANCER IMMUNOLOGY, IMMUNOTHERAPY**, vol. 51, 6 April 2002 (2002-04-06), pages 219-228, XP002615531, DOI: 10.1007/s00262-002-0273-7
- **SADOVNIKOVA E ET AL:** "Generation of human tumor-reactive cytotoxic T cells against peptides presented by non-self HLA class I molecules", **EUROPEAN JOURNAL OF IMMUNOLOGY**, **WILEY - V C H VERLAG GMBH & CO. KGAA, DE**, vol. 28, no. 1, 1 January 1998 (1998-01-01), pages 193-200, XP000864767, ISSN: 0014-2980, DOI: 10.1002/(SICI)1521-4141(199801)28:01<193::AID-IMMU193>3.3.CO;2-B
- **BINETRUY-TOURNAIRE ROSELYNE ET AL:** "Identification of a peptide blocking vascular endothelial growth factor (VEGF)-mediated angiogenesis", **EMBO JOURNAL, OXFORD UNIVERSITY PRESS, SURREY, GB**, vol. 19, no. 7, 3 April 2000 (2000-04-03), pages 1525-1533, XP002179245, ISSN: 0261-4189
- **WADA S ET AL.:** "Rationale for Antiangiogenic Cancer Therapy with Vaccination Using Epitope Peptides Derived from Human Vascular Endothelial Growth Factor Receptor 2", **CANCER RESEARCH**, vol. 65, no. 11, 1 June 2005 (2005-06-01), pages 4939-4946, XP002348368,

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionTechnical Field

5 **[0001]** The present invention relates to novel peptides that are extremely useful as cancer vaccines, and pharmaceuticals that comprise these peptides, for treating and preventing tumors.

Background Art

10 **[0002]** Tumor-rejection antigen genes have mainly been identified for malignant melanomas, and accordingly, cancer immunotherapies utilizing these genes are developed. Specifically, with recognition of the importance of CD8-positive T cells in anti-tumor immune responses, cancer vaccine therapy that induces tumor-specific CD8-positive T cells *in vivo* has received attention, and is being applied to various clinical applications. In addition, the mechanism by which peptides consisting of approximately ten amino acid residues activate T cells via the Class I pathway, by the assistance of various
15 costimulatory molecules, to induce tumor-specific cytotoxic T cells (CTLs) was also elucidated. Furthermore, peptides restricted to individual HLA molecules are being actively identified.

[0003] However, complete control of tumors is currently impossible. This may be due to tumor cell heterogeneity, a reduction or disappearance of MHC-Class I expression, in tumor cells, and the absence of target molecules in the tumor cells. Furthermore, the currently identified tumor antigen peptides exist in some types of tumors, but can not exist in all
20 tumor types. Thus, to resolve these problems, the present inventors did not use tumor cells as target cells, but rather focused on the endothelial cells of tumor vessels. More specifically, endothelial cells have hardly any problems involving a decrease or disappearance of MHC-Class I expression, or heterogeneity. Thus, if CTLs that target tumor vessels can be induced, problems in conventional cancer vaccine therapy, such as the disappearance of Class I and absence of a target molecule, may be overcome, regardless of the type of tumor, and excellent therapeutic effects can be anticipated.
25 Studies on tumor angiogenesis were started from a pioneering hypothesis proposed by Folkman *et al.* in the 1970's, and have been conducted from various angles. Many studies have been carried out on vascular endothelial growth factor (VEGF)-VEGF receptors (VEGFRs) to evaluate their significance in tumor angiogenesis. Angiogenesis inhibitors have been vigorously developed as target-oriented drugs, particularly in cancer therapy, and are already being clinically tested. However, therapies that use this concept for cancer vaccine treatments are not yet in use. One of the reasons
30 may be immunotolerance to VEGFR, which is expressed in normal cells. However, in the 1990's, Plate, Millauer, and Risau *et al.* confirmed that VEGFRs were strongly expressed in the endothelial cells of tumoral tissues. In addition, the immune response to autoantigens such as CEA and HER/neu, which are also expressed in normal cells, is not necessarily one of immunotolerance. Thus, the present inventors reasoned that VEGFRs may be used as targets for cancer vaccine therapy.

35 **[0004]** Recently, it was reported that active immunization against VEGFRs can inhibit angiogenesis in tumors, and metastasis (The Journal of Experimental Medicine, 2002, 195:12, 1575-1584). However, this literature merely used soluble VEGFR proteins, and made no investigation of the amino acid sequences of effective peptides.

[0005] WO99/4380 describes the identification of stimulating epitopes from angiogenic targets including the Tek protein or the KDR protein for the design of anti-angiogenic vaccines.
40

Disclosure of the Invention

[0006] The present inventors focused on possible cancer vaccine therapies that target VEGFR2 (KDR/flk-1; referred to below as KDR), where VEGFR2 is strongly expressed in tumoral tissue endothelial cells, and is thought to be involved
45 in the proliferation of endothelial cells on the VEGF signal. Furthermore, the present inventors screened for peptides that can be effectively used as vaccines, examining their specificity, and completing this invention.

[0007] More specifically, the present invention provides [1] to [15] as follows.

50 [1] A nonapeptide consisting of the amino acid sequence of SEQ ID NO: 2.

[2] A peptide with cytotoxic T cell inducibility, wherein one or two amino acids are substituted or added to the amino acid sequence of SEQ ID NO: 2, wherein the second amino acid from the N-terminus is phenylalanine, tyrosine, methionine, or tryptophan.

55 [3] A peptide with cytotoxic T cell inducibility, wherein one or two amino acids are substituted or added to the amino acid sequence of SEQ ID NO: 2, wherein the C-terminal amino acid is phenylalanine, leucine, isoleucine, tryptophan, or methionine.

[4] The peptide of claim 2, wherein the C-terminal amino acid is phenylalanine, leucine, isoleucine, tryptophan, or methionine.

[5] A pharmaceutical composition comprising one or more peptides of any one of claims 1 to 4.

[6] A vaccine comprising a peptide of any one of claims 1 to 4 as an active ingredient.

[7] A peptide for use in treating and/or preventing tumors, wherein the peptide is selected from the group consisting of the peptides of claims 2 to 4.

[8] A peptide for use in treating diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, wherein the peptide is selected from the group consisting of the peptides of claims 1 to 4.

[9] A pharmaceutical composition for use in treating and/or preventing tumors, wherein the pharmaceutical composition comprises one or more peptides of any one of claims 2 to 4.

[10] A pharmaceutical composition for use in treating diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, wherein the pharmaceutical composition comprises one or more peptides of any one of claims 1 to 4.

[11] An exosome that presents on its surface a complex consisting of a peptide of any one of claims 1 to 4, and an HLA-A2402 antigen.

[12] A vaccine for use in treating and/or preventing tumors, wherein the vaccine comprises a peptide of any one of claims 2 to 4 as an active ingredient.

[13] A vaccine for use in treating diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, wherein the vaccine comprises a peptide of any one of claims 1 to 4 as an active ingredient.

[14] The peptide of claim 7 or 8, the pharmaceutical composition of claim 9 or 10, or the vaccine of claim 12 or 13, which is used for administration to a subject whose HLA antigen is HLA-A24.

[15] The peptide, the pharmaceutical composition, or the vaccine of claim 14, which is used for administration to a subject whose HLA antigen is HLA-A2402.

Brief Description of the Drawings

[0008]

Fig. 1 shows the cytotoxicity of CTL clones against the target cells.

Fig. 2 shows the results of HLA-tetramer analyses of established CTL clones: (a) CTL clones induced by the peptides of this invention, and feeder cells; and (b) feeder cells (control).

Fig. 3 shows the effect of vaccination using the peptides of this invention on the survival rate of Colon 26-inoculated BALB/c mice.

Fig. 4 shows the effect of vaccination using the peptides of this invention on the growth of Colon 26-derived tumors.

Fig. 5 shows the effect of vaccination using the peptides of this invention on the survival rate of B16-inoculated mice.

Fig. 6 shows the effect of vaccination using the peptides of this invention on the growth of B 16 melanoma-derived tumors. Before challenge with a subcutaneous injection of B 16 tumor cells, A2/Kb TGM were vaccinated twice, with a one-week interval, with DC pulsed with KDR773 (black circles), DC alone (white triangles), or HBSS (white squares). The figure shows mean tumor growth ($P < 0.01$).

Fig. 7 shows the cytotoxic effect of CTL clone C29-169 on HLA-A24-positive cells presenting peptides of this invention.

Fig. 8 shows the cytotoxic effect of CTL clone C29-169 on HLA-A24-positive cells presenting peptides of this invention.

Fig. 9 shows the cytotoxic effect of CTL clone KDR C85-775 (KWC85-775) on HLA-A0201-positive cells presenting peptides of this invention.

Fig. 10 shows the cytotoxic effect of CTL clone KDR C85-775 (KWC85-775) on HLA-A0201-positive cells presenting KDR peptides.

Fig. 11 shows the cytotoxic effect of CTL clone KDR C51-1328 (KWC51) on HLA-A0201-positive cells presenting

peptides of this invention.

Fig. 12 shows the cytotoxic effect of CTL clone KDR C51-1328 (KWC51) on HLA-A0201-positive cells presenting KDR peptides.

Fig. 13 shows the cytotoxic effect of CTL clone KDR C44-773-2L (KWC44-773-2L) on HLA-A0201-positive cells presenting peptides of this invention.

Fig. 14 shows the cytotoxic effect of CTL clone KDR C44-773-2L (KWC44-773-2L) on HLA-A0201-positive cells presenting peptides of this invention.

Fig. 15 shows the cytotoxic effect of CTL clone KDR C44-773-2L (KWC44-773-2L) on HLA-A0201-positive cells presenting KDR peptides.

Fig. 16 shows the inhibitory effect of each type of antibody on the cytotoxic effect of the CTL clones.

Fig. 17 shows the inhibitory activity of the peptides of this invention on the B 16 cell-induced angiogenic response in A2/Kb TGM. Mice were vaccinated twice with HBSS, unpulsed DC, or DC pulsed with the epitope peptides of this invention (KDR190, KDR772, KDR773, KDR773-2L, KDR775, or KDR1084). The arrows indicate newly formed blood vessels, which run in a characteristic zigzag pattern.

Fig. 18 shows the suppressive effect of the peptides of this invention on the increase in volume of human tumors in transgenic mice.

Fig. 19 shows the cytotoxic effect of CTL clone KWC65-190 on HLA-A0201-positive cells presenting peptides of this invention, in the presence (black boxes) or absence (white boxes) of the peptides.

Fig. 20 shows the cytotoxic effect of CTL clone KWC72-772 on HLA-A0201-positive cells presenting peptides of this invention, in the presence (black boxes) or absence (white boxes) of the peptides.

Fig. 21 shows the cytotoxic effect of CTL clone KWC72-772 on HLA-A0201-positive cells presenting KDR peptides. HePG2-VEGFR2: black boxes; HePG2-EGFP: white boxes.

Fig. 22 shows the cytotoxic effect of CTL clone C7-1318 on HLA-A24-positive cells presenting peptides of this invention, in the presence (black boxes) or absence (white boxes) of the peptides.

Fig. 23 shows the cytotoxic effect of CTL clone KWC46 on HLA-A24-positive cells presenting peptides of this invention, in the presence (black boxes) or absence (white boxes) of the peptides.

Fig. 24 shows the cytotoxic effect of CTL clone C18-189 on HLA-A24-positive cells presenting peptides of this invention, in the presence (black boxes) or absence (white boxes) of the peptides.

Fig. 25 shows the cytotoxic effect of CTL clone C65-826 on HLA-A24-positive cells presenting peptides of this invention, in the presence (black boxes) or absence (white boxes) of the peptides.

Fig. 26 shows the cytotoxic effect of CTL clone C7-1318 on HLA-A24-positive cells presenting peptides of this invention. HT29-VEGFR2: black boxes; HT29-EGFP: white boxes.

Fig. 27 shows the cytotoxic effect of CTL clone KWC46 on HLA-A24-positive cells presenting peptides of this invention. HT29-VEGFR2: black boxes; HT29-EGFP: white boxes.

Fig. 28 shows the cytotoxic effect of CTL clone C18-189 on HLA-A24-positive cells presenting peptides of this invention. HT29-VEGFR2: black boxes; HT29-EGFP: white boxes.

Fig. 29 shows the cytotoxic effect of CTL clone C65-826 on HLA-A24-positive cells presenting peptides of this invention. HT29-VEGFR2: black boxes; HT29-EGFP: white boxes.

Fig. 30 shows the inhibitory activity of the peptides of this invention on the Colon 26 cell-induced angiogenic response in BALB/c mice. The mice were vaccinated twice with

[0009] HBSS, unpulsed DC, or DC pulsed with an epitope peptide of this invention (KDR189 or KDR826). The arrows indicate newly formed blood vessels, which run in a characteristic zigzag pattern.

Fig. 31 shows the inhibition of Colon 26 cell-induced angiogenic response in BALB/c by the peptides of this invention. The bars indicate the average, and the vertical error bars indicate standard error.

Fig. 32 shows the cytotoxic effect of CTL, derived from patients stimulated with the peptides of this invention, on HLA-A24-positive cells, in the presence (black diamonds) or absence (white squares) of the peptides.

(A) The cytotoxic effect of colon cancer patient-derived CTL obtained by stimulation with SEQ ID NO: 8 (in which the amino acid initiation position is KDR169);

(B) The cytotoxic effect of colon cancer patient-derived CTL obtained by stimulation with SEQ ID NO: 5 (in which the amino acid initiation position is KDR189); and

(C) The cytotoxic effect of colon cancer patient-derived CTL obtained by stimulation with SEQ ID NO: 3 (in which the amino acid initiation position is KDR220).

Mode for Carrying Out the Invention

[0010] The present inventors first considered that various proteins are presented *in vivo* on antigen presenting cells after being degraded to 9-mer peptides (nonapeptides), and examined the binding affinity of 9-mer or 10-mer partial peptides of KDR proteins to HLA antigens, which are human major histocompatibility antigens (MHC antigens). The lower case letters in the peptide sequences shown on the right of Table 1 indicate the fifth amino acids.

[0011] The amino acid sequence of human KDR protein is well known, disclosed by US Patent No. 5,861,301 for example, and can be easily obtained by one skilled in the art. The 9-mer and 10-mer peptides can be obtained by synthesizing peptides initiated from any position, based on the full-length amino acid sequence of the obtained KDR protein. Peptides can be synthesized according to conventional methods used in peptide chemistry. Commonly used synthesis methods are described, for example, in literatures such as Peptide Synthesis, Interscience, New York, 1966; The Proteins, vol. 2, Academic Press Inc., New York, 1976; Peptide Synthesis (Peputido Gousei), Maruzen Co., Ltd., 1975; Foundations and Experiments in Peptide Synthesis (Peputido Gousei no Kiso to Jikken), Maruzen Co., Ltd., 1985; and Development of Pharmaceuticals, New Series (Iyaku hin no Kaihatsu, Zoku), Volume 14, Peptide Synthesis (Peputido Gousei), Hirokawa Shoten, 1991, and in publications such as International Publication No. WO 99/67288. Binding with HLA antigens can be measured by isolating cells that comprise HLA antigens on their cell surface, such as dendritic cells, and using conventional methods to measure the binding of peptides to cells.

[0012] Alternatively, software programs now available on the Internet, such as those described in Parker K. C., J. Immunol. 152, 1994, may be used to calculate the binding affinities between various peptides and HLA antigens *in silico*. Binding affinity with HLA antigens can be measured as described, for example, in Parker, K. C., J. Immunol., 152, 1994; and Nukaya, I., Int. J. Cancer, 80, 1999.

[0013] To obtain sufficient results, A-24 type and A-02 type antigens, said to be highly expressed among the Japanese, are preferably used as the HLA antigens. More preferably, subtypes such as A-2402 and A-0201 are used. However, clinically, by predetermining the HLA antigen type of a patient in need of treatment, a peptide can be appropriately selected to have a high level of binding affinity to that antigen, or level of cytotoxic T cell (CTL) inducibility upon antigen presentation. Furthermore, to obtain peptides with high levels of binding affinity and CTL inducibility, substitution or addition of one, two, or several amino acids may be performed based on the amino acid sequence of the naturally occurring KDR partial peptide. Herein, the term "several" means five or less, or preferably three or less. In addition to peptides in nature, the sequence regularity of the peptides displayed by binding to HLA antigens is already known (J. Immunol., 152, 3913, 1994; Immunogenetics. 41:178, 1995; J. Immunol. 155:4307, 1994). Thus, modifications based on this regularity can also be performed on the obtained peptides. For example, peptides in which the second amino acid from the N terminus is substituted with phenylalanine, tyrosine, methionine, or tryptophan, and in which the C-terminal amino acid is substituted with phenylalanine, leucine, isoleucine, tryptophan, or methionine may also be preferably used as peptides with high HLA-24 binding affinity. On the other hand, peptides in which the second amino acid from the N terminus is substituted with leucine or methionine, and in which C-terminal amino acid is substituted with valine or leucine may be preferably used as peptides with high HLA-0201 binding affinity. Furthermore, one to two amino acids may also be added to the N and/or C terminus of the peptides.

[0014] However, when a peptide sequence is identical to a portion of an amino acid sequence of an endogenous or exogenous protein with a different function, it may cause side effects such as autoimmune diseases or allergic symptoms against specific substances. Therefore, it is preferable to use available databases to carry out homology searches, to avoid situations in which a sequence matches the amino acid sequence of another protein. Furthermore, if homology searches reveal that no peptides comprising even a one- or two- amino acid difference exist, there is no risk of such problems caused by modifications of the above-mentioned amino acid sequences for increasing the binding affinity with HLA antigens and/or the CTL inducibility.

[0015] As described above, peptides with high binding affinity to the HLA antigens are expected to be highly effective as cancer vaccines. However, it is necessary to determine whether the candidate peptides, which are selected using high binding affinity as an indicator, actually have CTL inducibility or not. Confirmation of CTL inducibility is carried out by inducing antigen-presenting cells comprising human MHC antigens (such as B-lymphocytes, macrophages, and dendritic cells), more specifically dendritic cells derived from human peripheral blood mononuclear leukocytes; stimulating the cells with the peptides; mixing the cells with CD8-positive cells; and then measuring cytotoxicity against the target cells. As the reaction system, transgenic animals generated to express a human HLA antigen may be used (for example, those described in Hum. Immunol. 2000 Aug.; 61(8):764-79 Related Articles, Books, Linkout Induction of CTL response by a minimal epitope vaccine in HLA A*0201/DR1 transgenic mice: dependence on HLA class II restricted T(H) response., BenMohamed L., Krishnan R., Longmate J., Auge C., Low L., Primus J., and Diamond D.J.). For example, by radiolabeling the target cells with ⁵¹Cr or such, cytotoxicity can be calculated from the radioactivity released from the target cells. Alternatively, the activity can be examined by measuring the IFN- γ produced and released by CTL in the presence of antigen-presenting cells that have immobilized peptides, and visualizing the inhibition zone on the media using anti-IFN- γ monoclonal antibodies.

[0016] The results of examining the CTL inducibility of peptides as described above revealed that those with high binding affinity to HLA antigens do not necessarily have high inducibility. Furthermore, nonapeptides selected from peptides comprising the amino acid sequences of VYSSEEAEI (SEQ ID NO: 2), GYRIYDVVL (SEQ ID NO: 3), SYMISY-AGM (SEQ ID NO: 5), RFVPDGNRI (SEQ ID NO: 8), KWEFPRDRL (SEQ ID NO: 11), or DFLTLEHLI (SEQ ID NO: 12), and nonapeptides and decapeptides selected from peptides comprising the amino acid sequence of AMFFWLLL (SEQ ID NO: 29), VIAMFFWLL (SEQ ID NO: 30), AVIAMFFWL (SEQ ID NO: 33), KLIEIGVQT (SEQ ID NO: 34), YMISYAGMV (SEQ ID NO: 40), or IQSDVWSFGV (SEQ ID NO: 46), showed particularly high CTL inducibility.

[0017] The present disclosure further provides peptides with cytotoxic T cell inducibility, wherein one, two, or several amino acids are substituted with or added to the amino acid sequence of SEQ ID NO: 2, 3, 5, 8, 11, or 12. One, two, or several amino acids can be substituted or added to the amino acid sequences of SEQ ID Nos: 2, 3, 5, 8, 11, and 12, consisting of nine amino acids, as long as there is CTL inducibility and the sequences do not match the amino acid sequence of another protein. In particular, for example, the second amino acid from the N terminus is preferably substituted to phenylalanine, tyrosine, methionine, or tryptophan, or the C-terminal amino acid is preferably substituted to phenylalanine, leucine; isoleucine, tryptophan, or methionine; or one or two amino acids are added to the N terminus and/or C terminus.

[0018] The present disclosure also provides peptides with cytotoxic T cell inducibility, wherein one, two, or several amino acids are substituted or added to the amino acid sequence of SEQ ID NOs: 29, 30, 33, 34, 40, or 46. One, two, or several amino acids can be substituted or added to the amino acid sequences of SEQ ID Nos: 29, 30, 33, 34, 40, and 46, consisting of nine or ten amino acids, as long as there is CTL inducibility and the sequences do not match the amino acid sequence of another protein.

[0019] In particular, for example, the second amino acid from the N terminus is preferably substituted to leucine or methionine, or the C-terminal amino acid is preferably substituted to valine or leucine; or one or two amino acids are added to the N terminus and/or C terminus. An example of such a modified peptide is the peptide of SEQ ID NO: 30, in which the second amino acid from the N terminus is substituted to leucine (SEQ ID NO: 54), but this example is not limiting. CTL clones obtained on stimulation with these modified peptides can recognize the original peptides, and cause damage.

[0020] The peptides of this invention can be used alone or in combinations of two or more, as cancer vaccines capable of inducing CTL *in vivo*. By administering the peptides of this invention, the peptides are presented at a high density on the HLA antigens of antigen-presenting cells. CTLs that specifically react with the complexes formed between the displayed peptides and the HLA antigens are induced, increasing aggression against vascular endothelial cells in the tumor cells to be targeted. Alternatively, antigen-presenting cells on whose cell surface the peptides of this invention are immobilized can be obtained by deriving dendritic cells from a subject, and stimulating them with the peptides of this invention. The obtained antigen-presenting cells are re-administered to the subjects to induce CTL in the subjects. As a result, aggression towards the target cells can be increased.

[0021] More specifically, the present invention provides pharmaceuticals for use in treating tumors or preventing proliferation, metastasis, and such of tumors, where the pharmaceuticals comprise one or more peptides of this invention. Angiogenesis at pathologic sites is closely associated with not only tumors, but also with diseases such as diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, and the metastasis of solid tumors (Folkman, J., Nature Med. 1:27-31 (1995); Bicknell, R., Harris, A.L., Curr. Opin. Oncol. 8:60-65 (1996)). Therefore, the peptides of this invention can be used to treat tumors, diseases such as diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, atherosclerosis, and metastasis of solid tumors.

[0022] The peptides of this invention were confirmed to inhibit the formation of tortuous blood vessels, which are morphologically different from normal blood vessels, and which are formed in malignant tumoral tissues. The results of analyzing wound healing and fertility in vaccinated mice also confirmed that the peptides do not have an adverse effect on normal physiological angiogenesis. Furthermore, by using CTL clones that recognize the peptides of this invention, cytotoxicity against non-proliferative or proliferative endothelial cells was tested *in vitro*. These clones showed stronger activity towards proliferative endothelial cells than non-proliferative endothelial cells. More specifically, they can function very specifically for disorders involving proliferative endothelial cells, and particularly cancer.

[0023] *In vivo* and *in vitro* stimulation of dendritic cells by the peptides of this invention can be easily performed by exposing the cells to a high concentration of the peptides, which causes these peptides to replace the peptides originally immobilized on the cells. Therefore, the peptides used in this invention must have at least a certain level of binding affinity to HLA antigens.

[0024] The pharmaceuticals of this invention may be directly administered as the peptides of this invention themselves, or may be administered as pharmaceutical compositions that have been formulated by conventional formulation methods. In such cases, the pharmaceuticals of this invention can appropriately include, in addition to the peptides of this invention, carriers, excipients, and such that are ordinarily used for pharmaceuticals, without particular limitations. The pharmaceuticals of this invention can be used for treatment and prevention of various tumors, such as gastric cancer, duodenal cancer, colon cancer, lung cancer, breast cancer, prostate cancer, and brain tumors. The peptides of this invention do

not target the tumor cells themselves, but target the endothelial cells of blood vessels that are newly formed in tumoral tissues. Therefore, a wide variety of tumors can be treatment targets, and the pharmaceuticals of this invention are not particularly limited in their use.

[0025] Pharmaceuticals for use in treating and/or preventing tumors, which comprise a peptide of this invention as an active ingredient, can be administered with adjuvants so that cellular immunity is induced effectively; can be administered with other active ingredients such as antitumor agents; and can be administered in granular forms. Those adjuvants described in the literature (Clin. Microbiol. Rev., 7:277-289, 1994) or such are applicable. Furthermore, the pharmaceuticals of this invention can be administered as liposome formulations, as granular formulations bound to beads of a few micrometers in diameter, and as formulations to which lipids are bound. Administration methods may be carried out, for example, orally, intradermally, or subcutaneously, or through intravenous injection, or such. Systemic administration or local administration to the vicinity of the target tumor may be applicable. Doses of the peptides of this invention can be adjusted appropriately, depending on the disease to be treated, age and weight of the patients, administration methods, and such. Ordinarily, 0.001 mg to 1,000 mg, preferably 0.01 mg to 100 mg, more preferably 0.1 mg to 10 mg, of the peptides of this invention are preferably administered once in a few days to a few months.

[0026] One skilled in the art can appropriately select suitable doses.

[0027] Alternatively, the present invention provides intracellular vesicles which present complexes formed between the peptides of this invention and HLA antigens on their surface. These intracellular vesicles are called exosomes. Exosomes can be prepared, for example, according to the methods specifically described in Published Japanese Translation of International Publication Nos. Hei 11-510507 and 2000-512161. Exosomes can preferably be prepared using antigen-presenting cells obtained from subjects who are to be the target of therapy or prophylaxis. The exosomes of this invention can be inoculated as cancer vaccines, as for the peptides of this invention.

[0028] The type of HLA antigens to be used must match that of the subject in need of therapy and/or prophylaxis. For example, for Japanese people, HLA-A24 or HLA-A02, particularly HLA-A2402 or HLA-0201, is often appropriate.

[0029] The present disclosure also provides methods for inducing antigen-presenting cells using the peptides of this invention. The antigen-presenting cells can be induced by inducing dendritic cells from peripheral blood monocytes; and then contacting (stimulating) them with the peptides of this invention, *in vitro* or *in vivo*. Administering the peptides of this invention to subjects induces in the body of the subject antigen-presenting cells to which the peptides of this invention are immobilized. Alternatively, the peptides of this invention can be immobilized to the antigen-presenting cells to be administered to the subject as a vaccine.

[0030] The present disclosure also provides methods for inducing antigen-presenting cells with a high level of cytotoxic T cell inducibility, wherein said methods comprise the *in vitro* introduction to an antigen-presenting cell of a gene that comprises a polynucleotide encoding a peptide of this invention. The genes to be introduced may be in the form of DNA or RNA. The introduction methods may be various methods commonly used in the art, such as lipofection, electroporation, and calcium phosphate methods, without particular limitations. More specifically, the methods may be performed as described in, for example, Cancer Res., 56:5672, 1996; J. Immunol., 161:5607, 1998; J. Exp. Med., 184:465, 1996; or Published Japanese Translation of International Publication No. 2000-509281. By introducing the genes into antigen-presenting cells, the genes undergo transcription, translation, and such in the cells. The obtained proteins are then subjected to Class I or II MHC processing and a presentation pathway to present partial peptides.

[0031] Furthermore, the present disclosure provides methods for inducing CTLs using the peptides of this invention. By administering the peptides of this invention to a subject, CTLs are induced in the body of the subject, enhancing immunity against the angiogenic endothelial cells in the tumoral tissues. Alternatively, the methods may be used for *ex vivo* therapeutic methods, in which subject-derived antigen-presenting cells, CD8-positive cells, or peripheral blood mononuclear leukocytes are contacted (stimulated) *in vitro* with the peptides of this invention to induce CTLs, and then the cells are returned to the subject.

[0032] In addition, the present disclosure provides cytotoxic T cells, that are induced using the peptides of this invention and then isolated. The cytotoxic T cells, which have been induced by stimulation with antigen-presenting cells that present the peptides of this invention, are preferably derived from subjects to be the target of therapy and/or prophylaxis. The cytotoxic T cells can be administered alone or, for the purpose of antitumor effect, in combination with other drugs, including the peptides, exosomes and so on of this invention. The obtained cytotoxic T cells act specifically against target cells presenting the peptides of this invention, or preferably, against target cells presenting the same peptides used for induction. The target cells may be cells that endogenously express KDR, or cells forced to express KDR. Furthermore, cells that present the peptides of this invention on their cell surface due to stimulation by these peptides can also be targeted.

[0033] The present disclosure also provides antigen-presenting cells that present complexes formed between HLA antigens and the peptides of this invention. The antigen-presenting cells that are obtained by contact with the peptides of this invention, or with nucleotides encoding the peptides of this invention, are preferably derived from subjects to be targeted for therapy and/or prophylaxis. The antigen-presenting cells can be administered as vaccines alone, or in combination with other drugs such as the peptides of this invention, exosomes, and cytotoxic T cells.

Best Mode for Carrying Out the Invention

[0034] Herein below, the present invention will be specifically described using Examples, but it is not to be construed as being limited thereto.

[0035] Peptides that are not selected from any one of the peptides according to appended claims 1 to 4 are disclosed for comparison only.

[Example 1] Prediction of VEGFR-2 (KDR)-derived peptides -- 1 (HLA-A*2402)

[0036] Based on the entire amino acid sequence of KDR protein, twelve types of 9-mer peptides (SEQ ID NOs: 1 to 12), and twelve types of 10-mer peptides (SEQ ID NOs: 13 to 24) were predicted in order from that with the highest binding affinity to HLA-A*2402, using Bioinformatics & Molecular Analysis Section (BIMAS) HLA Peptide Binding Prediction software (http://bimas.dcrf.nih.gov/cgi-bin/molbio/ken_parker_comboform) (Table 1). Table 1 shows the binding affinity of each of the 9 mers and 10 mers in order from the highest value, together with the position of their N termini in the amino acid sequence of the KDR protein. In the table, CE652 (SEQ ID NO: 25) refers to one of the epitopes of tumor antigen CEA (carcinoembryonic antigen) that has been reported by Nukaya, I. (Int. J. Cancer 80, 1999). HIV peptides (ILKEPVHGV (SEQ ID NO: 55) and RYLRLDQQLL (SEQ ID NO: 56)) were used as the negative control peptides.

[0037] The peptides were synthesized according to standard solid-phase synthesis methods using Cleaved PepSet (Mimotope, San Diego, CA), and purified by reverse phase HPLC. The purity (>95%) and the type of peptide were individually determined using HPLC and mass spectrometry.

Table 1 Binding of HLA-A*2402 to KDR-derived peptides

SEQ ID NO:	INITIATION POSITION	AA SEQUENCE (9mers)	BINDING AFFINITY	SEQ ID NO:	INITIATION POSITION	AA SEQUENCE (10mers)	BINDING AFFINITY*1
1	KDR583	WYKLG PQPL	240	13	KDR950	DYVGaIPVDL	420
2	KDR1318	VYSSEEAEEL	220	14	KDR434	SYQYgTTQTL	360
3	KDR220	GYRIYDVVL	200	15	KDR1058	DYVRkGDARL	300
4	KDR920	KFGNLSTYL	48	16	KDR304	LYTCaASSGL	200
5	KDR189	SYMLSYAGM	38	17	KDR1318	VYSSeEAELL	200
6	KDR828	EFPRDRKLK	33	18	KDR734	LYTCqACSVL	200
7	KDR1152	TFSELVEHL	29	19	KDR926	TYLRsKRNEF	198
8	KDR169	RFVPDGNRI	22	20	KDR353	KYLGyPPPEI	165
9	KDR331	AFGSG MESL	20	21	KDR116	VYVQdYRSPF	150
10	KDR604	KNLDTLWKL	16	22	KDR395	NYTViLTNPI	72
11	KDR826	KWEFPRDRL	12	23	KDR777	FFWLILVIL	24
12	KDR998	DFLTLEHLI	9	24	KDR193	SYAGmVFCEA	9.2
25	CE652 *2	TYACFVSNL	200				

*1 Parker KO:1994 J.Immunol 152

*2 Nukaya I:1999 Int.J.Cancer 80

[Example 2] Establishment of CTL lines using the predicted peptides

[0038] Epstein-Barr virus (EBV)-immortalized B cell lines, TISI (HLA-A24/24) and EHM (HLA-A3/3), were provided by Takara Shuzo Biotechnology Research Laboratory.

[0039] HLA serotyping was performed using 7 mL of peripheral blood collected from a healthy subject. Peripheral blood mononuclear lymphocytes (PBMCs) were isolated from HLA-A24-positive peripheral blood using Ficoll Paque (Pharmacia) specific gravity centrifugation.

[0040] The obtained PBMCs were left to stand for ten hours in a culture flask (Corning, 43072). After removing the suspended cells, the cells that adhered to the flask were cultured in AIM-V medium (Invitrogen) supplemented with 2% autoserum by adding 1,000 U/mL of human GM-CSF (provided by Kirin Brewery), and 1,000 U/mL of human IL-4 (Genzyme). Five days later, the cells were cultured for another 48 hours in 10 µg/mL of OK-432 (provided by Chugai Pharmaceutical Co., Ltd.), and were used as antigen-presenting cells for CTL induction. Dendritic cells (DCs) were confirmed to be DCs by reaction with FITC-labeled anti-ClassII, CD80, and CD86 antibodies, and with PE-labeled anti-ClassI, CD11c, CD40 (all from Beckton-Dickinson), and CD83 (Immunotech) antibodies; followed by analysis of surface

antigens by FACS-Calibur (Beckton-Dickinson) using Cell Quest software.

[0041] Pre-induced DCs were pulsed for four hours at 20°C with twelve kinds of 9-mer peptides (SEQ ID NOs: 1 to 12) (20 µg/mL), that were predicted in Example 1 to have high binding affinity, in the presence of 3 µg/mL of β2-microglobulin. The resulting DCs were mixed with CD8-positive cells selected by magnetic beads (Dynabeads M-450 and Detachabeads) from PBMC, at a ratio of 1:20 or 1:2, and then cultured in a 48-well plate (Corning), in the presence of 10 ng/mL of human IL-7 (Genzyme). Three days later, the final concentration of 10 U/mL of IL-2 (Sigma) was added to the culture. And seven and 14 days later, the same DCs were stimulated to induce CTLs, and 20 days later, cytotoxicity was measured for each well using peptide-pulsed TISI cells as targets. Only the positive wells were cultured by a stimulation method with allo-PBMC, EBV-immortalized B cell lines (EHM), and 30 ng/mL of anti-CD3 antibody. CTLs were functionally analyzed 14 days later.

[0042] Cytotoxic activity was evaluated by a four-hour ⁵¹Cr release assay. A concentration of 20 µg/mL of the target cells were pulsed with peptides overnight. The target cells were labeled with 100 µCi of Na₂⁵¹CrO₄ at 37°C for one hour, and then washed three times with RPMI1640. The target cells (1x 10⁴/100 µL) and 100 µL of effector cells at various concentrations with a total volume of 200 µL were placed into a U-shaped 96-well microtiter plate (Corning), and cultured at 37°C in a CO₂ incubator for four hours. After culturing, 100 µL of supernatant was collected from each well, and measured using a γ counter. Natural decay was the radioactivity of the target cells and the medium, and maximum decay was the radioactivity of the target cells and 1 M HCl. Cytotoxic activity (percentage of specific lysis) was calculated by the following formula:

$$\text{Cytotoxic activity (\% Cytotoxicity)} = \frac{(\text{Experimental decay} - \text{natural decay})}{(\text{maximum decay} - \text{natural decay})} \times 100$$

[0043] Six types of CTL lines were established as a result, as shown in Table 2. In particular, nonapeptides shown by SEQ ID NO: 2 (amino acid initiation position KDR1318), SEQ ID NO: 3 (KDR220), SEQ ID NO: 5 (KDR189), SEQ ID NO: 8 (KDR169), SEQ ID NO: 11 (KDR826), and SEQ ID NO: 12 (KDR998) were shown to be effective as epitope peptides.

Table 2

				CYTOTOXIC ACTIVITY			
				× 20		× 2	
SEQ ID NO:	INITIATION POSITION	AA SEQUENCE	BINDING *1 AFFINITY	Pep(+)	Pep(--)	Pep(+)	Pep(--)
1	KDR583	WYKLG PQPL	240	6%	4%	0%	0%
2	KDR1318	VYSSEEAE L	220	39%	21%	8%	3%
3	KDR220	GYRIYDVVL	200	57%	2%	20%	0%
4	KDR920	KFGNLSTYL	48	3%	4%	0%	2%
5	KDR189	SYMISYAGM	38	41%	3%	20%	0%
6	KDR828	EFPRDLKL	33	3%	2%	1%	0%
7	KDR1152	TFSELVEHL	29	0%	0%	0%	0%
8	KDR169	RFVPDGNRI	22	98%	2%	53%	0%
9	KDR331	AFGSGMESL	20	6%	6%	0%	0%
10	KDR604	KNLDTLWKL	16	3%	2%	1%	1%
11	KDR826	KWEFPRDRL	12	23%	0%	11%	0%
12	KDR998	DFLTLEHLI	9	13%	4%	8%	0%
25	CE652*2	TYACFVSNL	200	29%	16%	7%	1%

*1 Parker.KC:1994.J.Immunol 152

*2 Nukaya I:1999 Int.J.Cancer 80

[Example 3] Establishment of CTL clones derived from predicted peptides

[0044] The six types of CTL lines established in Example 2 were diluted to 0.3, 1, or 3 cells/well in a U-shaped 96-well plate. 7x 10⁴ cells/well of allo-PBMC, 1x10⁴ cells/well of EHM, 30 ng/mL of anti-CD3 antibody, and 125 U/mL of IL-2 were added to each well, and AIM-V supplemented with 5% autoserum was added thereto, so that total concentration was 150 µL/well. Ten days later, 50 µL/well of IL-2-supplemented medium adjusted was added to a final IL-2 concentration

EP 2 261 249 B1

of 125 U/mL. CTLs were functionally analyzed on the 14th day, and CTLs with activity were cultured on a large scale.

[0045] As a result, from each of the five types of nonapeptides shown by SEQ ID NO: 8 (amino acid initiation position KDR169), SEQ ID NO: 5 (KDR189), SEQ ID NO: 3 (KDR220), SEQ ID NO: 2 (KDR1318), and SEQ ID NO: 11 (KDR826), eight types (C13-169, C19-169, C29-169, C53-169, C72-169, C61-169, K5-169, and K25-169), two types (C12-189 and C18-189), eleven types (KWC3, 23, 25, 26, 36, 42, 46, 58, 61, 70, and 77), one type (C7-1318), and one type (C65-826) of CTL clones were established, respectively (Table 3).

Table 3

CYTOTOXIC ACTIVITY					
CLONE NAME	× 20		× 2		
	Pep(+)	Pep--)	Pep(+)	Pep(--)	
C13-169	98%	2%	66%	2%	
C19-169	94%	2%	50%	0%	
C29-169	100%	1%	70%	0%	
C53-169	100%	2%	91%	2%	
C72-169	87%	1%	39%	0%	
C61-169	88%	0%	38%	0%	
K5-169	36%	0%	6%	0%	
K25-169	27%	0%	5%	0%	
C12-189	27%	18%	7%	3%	
C18-189	67%	18%	28%	7%	
KWC3	54%	2%	11%	1%	
KWC23	95%	1%	49%	1%	
KWC25	93%	1%	70%	1%	
KWC26	87%	1%	34%	0%	
KWC36	62%	2%	11%	2%	
KWC42			67%	0%	
KWC46	94%	1%	51%	0%	
KWC58			76%	0%	
KWC61	70%	2%	19%	1%	
KWC70	92%	3%	64%	2%	
KWC77	99%	3%	70%	2%	
C65-826	38%	6%	8%	3%	
C7-1318	84%	1%	62%	1%	

[Example 4] Measurement of the cytotoxicity of established CTL clones

[0046] The degrees of cytotoxicity of the ten types of CTL clones established in Example 3 were examined by varying their ratio with the target cells (E:T ratio).

[0047] As a result, C53-169 derived from the nonapeptide of SEQ ID NO: 8 (amino acid initiation position KDR169) showed the strongest cytotoxicity.

[Example 5] HLA-tetramer analysis of established CTL clones

[0048] An HLA-tetramer was synthesized from HLA-A*2402 and the peptide of SEQ ID NO: 8 (amino acid initiation position KDR169). More specifically, plasmid vectors expressing the H chain (approximately 35 kDa) and L chain (approximately 11 kDa) were produced individually, and their recombinant protein was expressed using *E. coli*. The H-chain plasmid vector consists only of the extracellular domain, in which the carboxyl terminus was substituted with a biotinylation enzyme recognition sequence. The H chain and L chain form an HLA peptide complex by refolding with an antigenic peptide. The HLA peptide complexes were biotinylated by a biotinylation enzyme, excess biotin was removed by gel filtration chromatography and ion exchange chromatography, and the biotinylated HLA peptide complexes were isolated and purified. HLA-tetramer was synthesized by reacting the obtained biotinylated HLA peptide complex with streptavidin (SA) at a mole ratio of 4: to form a tetramer. When analyzing the samples, SA fluorescently labeled with phycoerythrin

(PE) (SA-PE) was used to form the tetramer. CD8 molecule is known to bind to the 245th amino acid in the $\alpha 3$ domain, which is alanine, with the exception of portions of HLA-A, B, and C. Therefore, the present inventors mutagenized the alanine in the HLA-tetramers to valine, to produce a mutant HLA-tetramer capable of selectively detecting CTLs with high avidity towards HLA peptide complexes, independent of the binding between the CD8 molecule and the $\alpha 3$ domain.

[0049] Established CTL clones were detected by the synthesized HLA-A24 KDR169-tetramer mutant. As a result, as shown in Fig. 2, the established CTL clones were found to be strongly stained by the tetramer-positive and CD8-positive fractions, and to specifically recognize complexes formed between the peptides of this invention and HLA antigens. The signals in the tetramer-negative and CD8-negative fractions are due to feeder cells.

[Example 6] Prediction of VEGFR-2 (KDR)-derived peptides -- 2 (HLA-A*0201)

[0050] In a similar manner to Example 1, 15 types of 9-mer peptides (SEQ ID NOs: 26 to 40), and 12 types of 10-mer peptides (SEQ ID NOs: 41 to 52) were predicted from the entire amino acid sequence of the KDR protein, in order from highest binding affinity with HLA-A*0201, using HLA Peptide Binding Prediction software (http://bimas.dcrt.nih.gov/cgi-bin/molbio/ken_parker_comboform) (Table 4). Table 4 shows the binding affinity of each of the 9 mers and 10 mers in order from the highest value, together with the position of their N terminal in the amino acid sequence of the KDR protein. In the table, CEA588 (SEQ ID NO: 53) refers to one of the epitopes of tumor antigen CEA (carcinoembryonic antigen) reported by Tanaka, H. et al. (poster presentation, AACR #3669, vol. 42, p681-682, March 2001).

Table 4 Binding of HLA-A*0201 to KDR-derived peptides

SEQ ID NO:	INITIATION POSITION	AA SEQUENCE (9mers)	BINDING AFFINITY	SEQ ID NO:	INITIATION POSITION	AA SEQUENCE (10mers)	BINDING AFFINITY
26	KDR1093	VLLWEIFSL	1792	41	KDR1094	LLWEIFSLGA	1092
27	KDR 633	SLQDQGDYV	769	42	KDR 5	VLLAVALWLC	539
28	KDR 5	VLLAVALWL	739	43	KDR 313	LMTKkNSTFV	470
29	KDR 775	AMFFWLLL	427	44	KDR 779	WLLLvILRT	292
30	KDR 773	VIAMFFWLL	270	45	KDR 505	ALIEgKNKTV	285
31	KDR1034	ILLSEKNVV	179	46	KDR1084	IQSDvWSFGV	285
32	KDR 604	KNLDTLWKL	128	47	KDR 733	GLYTcQACSV	223
33	KDR 772	AVIAMFFWL	113	48	KDR 780	LLLVIILRTV	201
34	KDR1328	KLIEIGVQT	107	49	KDR 6	LLAVaLWLCV	201
35	KDR 698	IMWFKDNET	76	50	KDR 137	YITEnKNKTV	180
36	KDR 608	TLWKLNATM	41	51	KDR1035	LLSEkNVVKI	167
37	KDR 491	FQGGNKIEV	32	52	KDR 4	KVLLaVALWL	133
38	KDR 435	YQYGTTQTL	32				
39	KDR 505	ALIEGKNKT	31				
40	KDR 190	YMISYAGMV	30	53	CEA 588*1	DVLYGPDTPi	0.25

*1 Tanaka H et al.:Poster presented at AACR 2001

[Example 7] Determination of HLA-A*0201 immobilizing epitope peptides

[0051] CTLs were induced by a vaccination method, using the 16 types of 9-mer and 10-mer peptides (SEQ ID NO: 26 to 41) predicted in Example 6 on transgenic mice that express human HLA (A0201) (Hum. Immunol. 2000 Aug.;61(8):764-79 Related Articles, Books, Linkout Induction of CTL response by a minimal epitope vaccine in HLAA*0201/DR1 transgenic mice: dependence on HLA class II restricted T(H) response., BenMohamed L., Krishnan R., Longmate J., Auge C., Low L., Primus J., and Diamond DJ.).

[0052] Six- to eight-week old BALB/C(H-2^d) mice were obtained from CLEA Japan. A2/Kb transgenic mice (TGM) were provided by F. Jim Primus, Ph. D. at Vanderbilt-Ingram Cancer Center. T2 cells (TAP-deficient and HLA-A*0201-positive cells) were provided by Prof. H. Shiku at Mie University's Third Department of Internal Medicine.

[0053] Bone marrow was collected from mouse femur and tibia, and lymphocytes and granulocytes were removed using anti-CD4, CD8, Gr-1 antibodies (Beckton-Dickinson), anti-B220 antibody (Bioscience) and rabbit complement (PeL-Freez). The obtained cells were cultured in a 6-well plate. Suspended cells were collected the following day to culture in RPMI1640 (Invitrogen) medium supplemented with 10% FBS in another 6-well plate, to which 1,000 U/mL of mouse GM-CSF (provided by Kirin Brewery) and 1,000 U/mL of mouse IL-4 (PEPRO TECH) were added. Three days

later, half of the above-mentioned medium was changed, and another two days later, the cells were cultured for 20 hours in 10 µg/mL of OK-432. The resulting suspended cells were used as DCs. DCs were reacted with FITC-labeled anti-ClassII and CD40 antibodies, and PE-labeled anti-ClassI, CD11c, CD80, and CD86 antibodies (all from Beckton-Dickinson), and confirmed to be DCs by analysis of the surface antigens by FACS-Calibur (Beckton-Dickinson) using Cell Quest software.

[0054] Next, 100 µg of each peptide, 140 µg of HbcAg120-140 helper peptide, and 100 µL of IFA were mixed (200 µL in total). The mixture was administered subcutaneously to the right abdomen on day 0, and to the left abdomen on day 11. The spleens of the vaccinated mice were collected on day 21, erythrocytes were hemolytically removed using red lysis buffer (Sigma), and a portion of those cells were used as responder cells for IFN-γ ELISPOT assay. The remaining cells were placed into a 24-well plate (Corning) at 6x 10⁶ cells/well with feeder cells at a ratio of 3:1, and stimulated again. Their cytotoxicity was measured five days later. Erythrocyte-free splenocytes, which were derived from syngeneic mice, were cultured for three days in the presence of 25 µg/mL of lipopolysaccharide (LPS) to use as feeder cells.

[0055] In this Example, the IFN-γ ELISPOT method was used to evaluate CTL inducibility according to SPOTs produced by IFN-γ producing cells. Flat-bottomed 96-well multiscreen plate MAHA S45 (Millipore) was treated at 4°C overnight with anti-mouse IFN-γ monoclonal antibody (Pharmingen). On the following day, the plate washed with PBS containing 0.05% Tween 20, and then reacted with a blocking buffer at room temperature for two hours. Thereafter, peptide-pulsed T2 cells and unpulsed T2 cells were added to each well at 10⁵ cells/100 µL. The splenocytes of vaccinated mice were further added to each well at a maximum of 4x 10⁶ cells/100 µL (200 µL in total), and cultured overnight at 37°C. The next day, each well was washed, and reacted for two hours with biotinylated rat anti-mouse IFN-γ antibodies (Pharmingen). After washing the plate thoroughly, Extravidin was added to each well and reacted at room temperature for two hours. After washing, alkaline phosphatase conjugate substrate (BIO-RAD) was added to the wells, which were left to stand at room temperature for five minutes. The blue spots resulting from IFN-γ production were measured using KS ELISPOT compact release (Carl Zeiss).

[0056] Specific blue spots were determined as:

$$\text{Specific blue spot} = (\text{TISI}(+)\text{SPOT} - \text{TISI}(-)\text{SPOT})$$

[0057] Those satisfying the following condition were considered effective:

$$\text{TISI}(+)\text{SPOT} / \text{TISI}(-)\text{SPOT} \geq 2$$

[0058] As a result, five types of epitope peptides, SEQ ID NO: 29 (amino acid initiation position KDR775), SEQ ID NO: 30 (KDR773), SEQ ID NO: 33 (KDR772), SEQ ID NO: 34 (KDR1328), and SEQ ID NO: 40 (KDR190) were obtained by IFN-γ ELISPOT assay (Table 5).

Table 5 ELISPOT Analysis

SEQ ID NO:	INITIATION POSITION	AA SEQUENCE	BINDING AFFINITY	ELISPOT ANALYSIS X 40 (SFC)			
				MOUSE1		MOUSE2	
				pep(+)	pep(-)	pep(+)	pep(-)
26	KDR1093	VLLWEIFSL	1792	5	8	7	1
41	KDR1094	LLWEIFSLGA	1092	0	0	0	0
27	KDR 633	SLQDQGDYV	769	0	0	0	0
28	KDR 5	VLLAVALWL	739	6	0	7	3
29	KDR 775	AMFFWLLL	427	95	32	322	17
30	KDR 773	VIAMFFWLL	270	171	6	193	1
31	EDR1034	ILLSEKNVV	179	0	0	0	0
32	KDR 604	KNLDTLWKL	128	0	7	0	0
33	KDR 772	AVIAMFFWL	113	88	19	143	31
34	KDR1328	KLIEIGVQT	107	0	0	81	5
35	KDR 698	IMWFKDNET	76	0	0	0	0
36	KDR 608	TLWFLNATM	41	0	0	0	0
37	KDR 491	FQGGNKIEV	32	0	0	0	0

(continued)

SEQ ID NO:	INITIATION POSITION	AA SEQUENCE	BINDING AFFINITY	ELISPOT ANALYSIS X 40 (SFC)			
				MOUSE1		MOUSE2	
				pep(+)	pep(-)	pep(+)	pep(-)
38	KDR 435	YQYGTI Q TL	32	0	0	0	0
39	KDR 505	ALIEGKNKT	31	0	0	0	0
40	KDR 190	YMISYAGMV	30	122	6	33	11
53	CEA 588*1	DVLYGPDTP	0.25	229	2	230	6

*1 Tanaka H et al.:Poster presented at AACR 2001

[Example 8] Confirmation of *in vivo* antitumor effect -- I

[0059] Bone marrow was collected from six- to eight-week old male BALB/c mice (CLEA Japan, Inc.) with that same anchor motif as human HLA-A2402. As in Example 7, GM-CSF (Kirin Brewery) (1,000 U/mL) and IL-4 (Genzyme) (1,000 U/mL) were added to the bone marrow and cultured to prepare dendritic cells. The dendritic cells were pulsed with the peptide of SEQ ID NO: 5 (amino acid initiation position KDR189) and then administered (1x 10⁶ cells/mouse) twice to nine BALB/c mice at the right lower quadrant, with an interval of seven days. Seven days later, colon cancer cell line (5x 10⁵ cells), Colon 26 (Taiho Pharmaceutical), was inoculated to the right abdomens of the mice, and mice survival and tumor growth were examined.

[0060] First, KDR expression was analyzed by RT-PCR, as indicated in Fig. 3, to confirm that KDR was not expressed in Colon 26 cells alone, but expressed in the tumoral tissues of mice inoculated with Colon 26. More specifically, tumor vessels were confirmed to appear in tumoral tissues, and to form the tumor.

[0061] Fig. 3 shows the survival curve of Colon 26-inoculated mice. 35 days after tumor inoculation, two out of nine mice survived when HBSS (Hank's Balanced Salt Solution) was administered alone (100 μ L) as a control; five out of nine mice survived when inoculated with dendritic cells that were not pulsed with peptides; and all of the nine mice survived when inoculated with dendritic cells pulsed with the peptide of SEQ ID NO: 5.

[0062] Fig. 4 shows the effect on the growth of Colon 26-derived tumors in mice. Compared to the control, suppressive effects on tumor growth were observed when inoculating dendritic cells, and remarkable suppression effects were observed when inoculating dendritic cells pulsed with the peptide of SEQ ID NO: 5.

[Example 9] Homology searches for the peptides of this invention

[0063] Homology searches for the peptides of this invention (SEQ ID NO: 2 (amino acid initiation position KDR1318), SEQ ID NO: 3 (KDR220), SEQ ID NO: 5 (KDR189), SEQ ID NO: 8 (KDR169), SEQ ID NO: 11 (KDR826), SEQ ID NO: 30 (KDR773), and SEQ ID NO: 40 (KDR190)) were carried out by using the BLAST program (<http://www.ncbi.nlm.nih.gov/blast/blast.cgi>). No peptides with a sequence completely identical to any one of these peptides could be found (Table 6). The 9-mer peptide (KDR169) of SEQ ID NO: 8 capable of inducing strong CTL activity in Example 4 had only one sequence containing two mismatches (77.8% homology), and two sequences containing three mismatches (66.7% homology). The peptide of SEQ ID NO: 5 (KDR189), whose remarkable *in vivo* anti-tumor effect was observed in Example 8, had only one sequence containing three mismatches (66.7% homology).

Table 6 Homology analysis using the BLAST program (<http://www.ncbi.nlm.nih.gov/blast/blast.cgi>)

	KDR169	KDR189	KDR826	KDR220	KDR1318	KDR773	KDR190
IDENTITY (9/9)	0	0	0	0	0	0	0
IDENTITY (8/9)	0	0	3	0	0	0	0
IDENTITY (7/9)	1	0	2	0	2	2	0
IDENTITY (6/9)	2	1	-	0	-	2	0

[Example 10] Confirmation of *in vivo* anti-tumor effect -- 2

[0064] Bone marrow was collected from six- to eight-week old male A2/Kb transgenic mice (TGM) expressing human HLA-A0201. As in Example 8, GM-CSF (Kirin Brewery) (1,000 U/mL) and IL-4 (Genzyme) (1,000 U/mL) were added to the bone marrow and then cultured to prepare dendritic cells. The dendritic cells were pulsed with the peptide of SEQ ID NO: 30 (amino acid initiation position KDR773), and administered (1x 10⁶ cells/mouse) twice to the left lower quadrant

of twelve A2/Kb transgenic mice, with an interval of seven days. Seven days later, B16 melanoma cells (ATCC) were inoculated (1×10^6 cells) to the right abdomen of the mice, and mice survival and tumor growth were examined.

[0065] Fig. 5 shows the survival rate of B16-inoculated mice. 35 days after tumor inoculation, eight of twelve mice survived when HBSS was administered alone (100 μ L) as a control; eleven of twelve mice survived when inoculated with dendritic cells that were not pulsed with peptides; and all twelve mice survived when inoculated with dendritic cells pulsed with the peptide of SEQ ID NO: 30.

[0066] Fig. 6 shows the effect on the growth of B16 mice-derived tumors in mice. Compared to the control, suppressive effects were observed against tumor growth when inoculating with dendritic cells, and remarkable suppressive effects were observed on inoculating dendritic cells pulsed with the peptide of SEQ ID NO: 30.

[Example 11] Cytotoxic activity against cells forcibly expressing KDR

[0067] Using adenoviruses, KDR (see, J. Biol. Chem. 1994 Oct 28; 269(43):26988-95) was forcibly expressed in HLA-A24-positive human colon cancer HT29 strain (ATCC), according to the method described by Miyake, S. et al., Proc. Natl. Acad. Sci. USA. 93, 1320-1324 (1996). More specifically, based on the titer determined by plaque assay of human 293 cells, cells were infected at a specific multiplicity. HT29 cells were plated at a cell density of 10^5 cells in a 60-mm dish, and incubated for 24 hours. On the following day, the medium was removed and exchanged with 200 μ L of viruses continuously diluted with fresh medium. The cells were incubated at 37°C for one hour, and then growth medium was added to the cells, which were cultured for 48 hours to be used as target cells. HLA-A24-positive HT29 strain, forced by adenoviruses to express EGFP (see, Leukemia 1999 Apr; 13(4):605-13), was used as a control. HT29 strains forcibly expressing KDR and EGFP, respectively, can potentially present partial peptides of each protein on their respective cell surface.

[0068] Using CTL clone C29-169, induced from the peptide of SEQ ID NO: 8 (amino acid initiation position KDR169), as described in Example 3, which binds to HLA-A24, the cytotoxic effect on HLA-A24-positive HT29 cells forced to express KDR was measured by chromium release assay. The results are shown in Fig. 7.

[0069] Cytotoxic activity was measured by ^{51}Cr release assay for four hours. The target cells were cultured in a 6-well plate at a concentration of 1×10^6 cells/well for 24 hours, infected with adenovirus to which KDR or EGFP has been inserted (Ad-KDR or Ad-EGFP) at MOI 50, and used after 48 hours. The target cells were labeled with 100 μCi of $\text{Na}_2^{51}\text{CrO}_4$ at 37°C for one hour, and washed three times with RPMI1640. The target cells (1×10^4 cells/100 μL) and 100 μL of effector cells at various concentrations were added (200 μL in total) to a U-shaped 96-well microtiter plate (Coming), and cultured in a CO_2 incubator at 37°C for four hours. After culturing, 100 μL of supernatant was collected from each well and measured using a γ counter to calculate cytotoxicity, in the same manner as in Example 2.

[0070] As a result, as shown in Fig. 7, CTL clone C29-169 showed remarkably high cytotoxic effect towards HLA-A24-positive HT29 forcibly expressing KDR, compared to that towards cells forcibly expressing EGFP.

[Example 12] Cytotoxic activity against cells endogenously expressing KDR

[0071] The cytotoxic effect of the CTL clone C29-169 described in Example 3, against HUVEC KT5 strain, derived from HLA-A24-positive human umbilical cord vascular endothelial cells that endogenously express KDR, was examined using KDR-positive and HLA-A24-negative HUVEC strain, P8, as a control. These cells present partial peptides of KDR on their cell surface.

[0072] Cytotoxic activity was measured by a four hour ^{51}Cr release assay. The target cells were labeled with 100 μCi of $\text{Na}_2^{51}\text{CrO}_4$ at 37°C for one hour, then washed three times with RPMI1640. Target cells (1×10^4 cells/100 μL) and 100 μL of effector cells at various concentrations were added (200 μL in total) to a U-shaped 96-well microtiter plate (Corning), and cultured in a CO_2 incubator at 37°C for four hours. After culturing, 100 μL of supernatant was collected from each well and measured using a γ counter to calculate the cytotoxicity in the same manner as in Example 2.

[0073] As apparent from the results of Fig. 8, CTL clone C29-169 showed cytotoxic effect against HLA-A24-positive HUVEC endogenously expressing KDR, and showed weak cytotoxic effect against HLA-A24-negative HUVEC.

[Example 13] Establishment of CTL clones using HLA-A0201-binding peptides

[0074] By using the peptide of SEQ ID NO: 40 (amino acid initiation position KDR190), KDR773-2L peptide (SEQ ID NO: 54) in which the second amino acid from the N terminal in the peptide of SEQ ID NO: 30 (KDR773) has been converted to leucine, the peptide of SEQ ID NO: 29 (KDR775), the peptide of SEQ ID NO: 34 (KDR1328), and the peptide of SEQ ID NO: 33 (KDR772), which bind to HLA-A0201 and which were determined to be effective according to the ELISPOT analysis in Example 7, two types (KWC14-190 and KWC65-190), four types (KWC44-773-2L, KWC76-773-2L, KWC129-773-2L, and KWC134-773-2L), two types (KWC81-775 and KWC85-775), twelve types (KWC16, KWC21, KWC22, KWC47, KWC51, KWC108, KWC117, KWC132, KWC151, KWC153, KWC156, and

KWC159), and one type (KWC72-772) of CTL clones, respectively, were obtained as in Example 3. The respective cytotoxicities are shown in Table 7.

Table 7
CYTOTOXIC ACTIVITY

CLONE NAME	×20		×2	
	Pep(+)	Pep(--)	Pep(+)	Pep(--)
KWC14-190	43%	0%	9%	0%
KWC65 -190	83%	0%	27%	0%
KWC44 -773 -2L	87%	0%	64%	0%
KWC76 -773 -2L	83%	2%	68%	1%
KWC129 -773 -2L	89%	0%	82%	1%
KWC134-773-2L	89%	2%	54%	2%
kWC81 -775	86%	0%	64%	0%
KWC85 -775	90%	0%	63%	0%
KWC16	95%	0%	85%	0%
KWC21	100%	0%	91%	0%
KWC22	93%	0%	77%	0%
KWC47	109%	2%	87%	1%
KWC51	101%	0%	93%	1%
KWC108	71%	2%	23%	2%
KWC117	83%	0%	42%	0%
KWC132	102%	0%	55%	0%
KWC151	102%	0%	101%	0%
KWC153	47%	0%	18%	0%
KWC156	64%	0%	23%	0%
KWC159	93%	0%	86%	0%
KWC72-772	101%	0%	67%	0%

[0075] As apparent from the results in Table 7, the peptides of SEQ ID NOs: 29, 33, 34, and 40, as well as the peptide of SEQ ID NO: 54, in which the second amino acid from the N terminus in the peptide of SEQ ID NO: 30 is leucine, can all induce CTLs having remarkable cytotoxicity, and were thus shown to be effective as epitope peptides.

[Example 14]

[0076] The peptide of SEQ ID NO: 29 (amino acid initiation position KDR775) was added to HLA-A0201-positive T2 strain to produce a target cell. By using the cell, the cytotoxic effect of the CTL clone (KDR C85-775 (KWC85-775)) obtained from the peptide of SEQ ID NO: 29 in Example 13 was examined by chromium release assay.

[0077] In the results, shown in Fig. 9, CTL clone C85-775 (KWC85-775) clearly showed a cytotoxic effect against HLA-A0201-positive cells presenting the peptide of SEQ ID NO: 29. In contrast, absolutely no cytotoxic effect was observed against control cells not presenting the peptide of SEQ ID NO: 29.

[Example 15]

[0078] The cytotoxic effect of the CTL clone (KDR C85-775 (KWC85-775)) against hepatocellular carcinoma cell line HLA-A0201-positive HePG2, forced by adenoviruses to express KDR, was examined by chromium release assay. HLA-A0201-positive HePG2 forced to express EGFP was used as control.

[0079] In the results, shown in Fig. 10, CTL clone KDR C85-775 (KWC85-775) showed remarkably high cytotoxic effect against HLA-A0201-positive cells presenting KDR peptides on the surface, compared to activity against cells presenting the EGFP peptide.

[Example 16]

[0080] The peptide of SEQ ID NO: 34 (amino acid initiation position KDR1328) was added to HLA-A0201-positive T2

strain to produce target cells. Using these cells, the cytotoxic effect of the CTL clone (KDR C51-1328 (KWC51)) obtained from the peptide of SEQ ID NO: 34 in Example 13 was examined by chromium release assay.

[0081] In the results, shown in Fig. 11, CTL clone KDR C51-1328 (KWC51) clearly showed a cytotoxic effect against HLA-A0201-positive cells presenting the peptide of SEQ ID NO: 34. In contrast, absolutely no cytotoxic effect was observed against control cells not presenting the peptide of SEQ ID NO: 34.

[Example 17]

[0082] The cytotoxic effect of the CTL clone (KDR C51-1328 (KWC51)) against hepatocellular carcinoma cell line HLA-A0201-positive HePG2, forced by adenoviruses to express KDR, was examined by chromium release assay. HLA-A0201-positive HePG2 forced to expressing EGFP was used as control.

[0083] In the results, shown in Fig. 12, CTL clone KDR C51-1328 (KWC51) showed remarkably high cytotoxicity against HLA-A0201-positive cells presenting KDR peptides on the surface, compared to activity against cells presenting the EGFP peptide.

[Example 18]

[0084] The peptide of SEQ ID NO: 54 (KDR773-2L), in which the second peptide of SEQ ID NO: 30 is converted to leucine, was added to HLA-A0201-positive T2 strain to produce target cells. Using these cells, the cytotoxic effect of the CTL clone (KDR C44-773-2L (KWC44-773-2L)) obtained from the peptide of SEQ ID NO: 54 in Example 13 was examined by chromium release assay. Fig. 13 shows the results.

[0085] In the results, shown in Fig. 13, CTL clone KDR C44-773-2L (KWC44-773-2L) clearly showed a cytotoxic effect against HLA-A0201-positive cells presenting KDR773-2L. In contrast, absolutely no cytotoxic effect was observed against control cells not presenting KDR773-2L.

[Example 19]

[0086] The cytotoxic effect of the CTL clone (KDR C44-773-2L (KWC44-773-2L)) obtained from the peptide of SEQ ID NO: 54 (KDR773-2L), in which the second peptide of SEQ ID NO: 30 has been converted to leucine, was examined by chromium release assay using HLA-A0201-positive T2 strain pulsed with the unmodified peptide of SEQ ID NO: 30 (amino acid initiation position KDR773) as the target cell.

[0087] In the results, shown in Fig. 14, CTL clones obtained by stimulation with the modified peptide could recognize and damage the post-modification peptide.

[Example 20]

[0088] The cytotoxic effect of the CTL clone (KDR C44-773-2L (KWC44-773-2L)) against hepatocellular carcinoma cell line HLA-A0201-positive HePG2, forced by adenoviruses to express KDR, was examined by chromium release assay. HLA-A0201-positive HePG2 forced to express EGFP was used as control.

[0089] In the results, shown in Fig. 15, CTL clone KDR C44-773-2L (KWC44-773-2L) showed remarkably high cytotoxic effect against HLA-A0201-positive cells presenting KDR peptides on the surface, compared to activity against cells presenting the EGFP peptide.

[Example 21]

[0090] The present inventors determined whether various antibodies can block the cytotoxic effect of the CTL clone (KDR C51-1328 (KWC51)), obtained from the peptide of SEQ ID NO: 34, against hepatocellular carcinoma cell line HLA-A0201-positive HePG2, forced by adenoviruses to express KDR.

[0091] In the results, shown in Fig. 16, cytotoxic function was inhibited by anti-HLA Class I antibody against HLA antigens displayed by target cells, and anti-CD8 antibody against CD8 used as a marker for cytotoxic T cells.

[Example 22] Examination of angiogenesis-inhibiting activity by dorsal air sac (DAS) assay

[0092] The actual angiogenesis-inhibiting activity of peptide epitopes was evaluated in mice using a DAS assay (see, Clinical Cancer Research, Vol. 5, 2185-2191, August 1999; Cancer Research 62, 6116-6123, November 1, 2002).

[0093] Specifically, a membrane filter was fixed to both sides of a chamber ring, and sterilized with ethylene oxide gas. B16 melanoma cells were prepared to 5×10^5 cells/0.15 mL, and injected into the chamber. Mice (A2/kb transgenic mice (derived from B 16)) were immobilized face down, and an air sac was produced by injecting 5 to 10 mL of air in to

the murine dorsal hypoderm using a syringe. An approximately 1-cm incision was made to the lower part of the sac. The chamber ring, filled with cell suspension solution, was implanted in the dorsal hypoderm, and the skin was closed using a skin stapler. Six days after implantation, the mouse skin was decorticated together with the chamber, fixed in an extended form, and marked with a black rubber ring to observe the newly formed blood vessels using a microscope. Blood vessels that are newly formed by angiogenic factors released from malignant tumor cells characteristically run in a zigzag manner, and are morphologically different from naturally occurring background blood vessels. Therefore, blood vessels of 3-mm or longer within the rubber ring and running in a zigzag manner were determined to be newly formed blood vessels, and were evaluated using an angiogenesis index (AI). AIs was evaluated into six levels, from 0 to 5 according to the number of tortuous blood vessels. All those with five or more tortuous blood vessels were determined to be 5.

[0094] Vaccination was performed 14 days prior to chamber implantation, and repeated after one week (a total of two times) by respectively injecting 5×10^4 cells of HBSS, DC alone, or peptide-pulsed DC, to the tail vein. The results are shown in Fig. 17 and in Table 8, as shown below.

Table 8 Effect of DAS assay

INOCULATION	PEPTIDE STIMULATION	ANGIOGENESIS INDEX (AI)	P
VEHICLE	-	(0.7 $\pm$ 0.2)	
HBSS	-	(4.5 $\pm$ 0.2)	
DC	NONE	(4.2 $\pm$ 0.4)	
DC	KDR190 (SEQ ID NO:40)	(3.6 $\pm$ 0.5)	0.3453
DC	KDR772 (SEQ ID NO:33)	(2.8 $\pm$ 0.5)	<u>0.0346</u>
DC	KDR773 (SEQ ID NO:30)	(2.2 $\pm$ 0.7)	<u>0.0034</u>
DC	KDR773-2L (SEQ ID NO:54)	(1.3 $\pm$ 0.8)	<u>0.0001</u>
DC	(KDR775 (SEQ ID NO:29)	(2.2 $\pm$ 0.5)	<u>0.0007</u>
DC	KDR1084 (SEQ ID NO:46)	(1.7 $\pm$ 0.3)	<u><0.0001</u>

[0095] Fig. 17 and Table 8 showed that the formation of tortuous blood vessels was clearly suppressed in the groups vaccinated with DCs pulsed with the peptides of SEQ ID NO: 40 (amino acid initiation position KDR190), SEQ ID NO: 33 (KDR772), SEQ ID NO: 30 (KDR773), SEQ ID NO: 54 (KDR773-2L), SEQ ID NO: 29 (KDR775), and SEQ ID NO: 46 (KDR1084), respectively, as compared to the group to which DC was administered alone. This indicated a statistically significant angiogenesis inhibitory effect.

[0096] Wound healing and fertility in vaccinated mice were analyzed in order to observe the adverse effects of vaccination using these epitope peptides on normal physiological angiogenesis. However, no significant adverse effects were found in the vaccinated mice. Furthermore, using CTL clones that recognize KDR peptides, cytotoxicity against non-proliferative or proliferative endothelial cells was tested *in vitro* to examine adverse effects in humans. As a result, these clones showed stronger activity against proliferative endothelial cells than against non-proliferative endothelial cells.

[Example 23]

[0097] Using A2/Kb transgenic mice expressing human HLA-A0201, the peptides of SEQ ID NO: 30 (amino acid initiation position KDR773) and SEQ ID NO: 34 (KDR1084) were each mixed with incomplete Freund's adjuvant, then administered on day 3 and 13 to colon cancer strain MC38-implanted tumor-bearing mice so as to indicate changes in tumor volume.

[0098] The *in vivo* anti-tumor effect in a therapeutic model of HLA-A* 0201-immobilized epitope peptide (KDR773) was evaluated using the area of the tumor, using the same vaccination method as in Example 22. The result, as apparent from Fig. 18, confirm a significant tumor growth suppression effect by vaccination with the peptides of SEQ ID NO: 30 and SEQ ID NO: 34.

[Example 24]

[0099] The peptide of SEQ ID NO: 40 (amino acid initiation position KDR190) was added to HLA-A0201-positive T2 strain to produce target cells. Using these cells, the cytotoxic effect of the CTL clone (KWC65-190) obtained from the peptide of SEQ ID NO: 40 in Example 13 was examined using chromium release assay.

[0100] In the results, shown in Fig. 19, CTL clone KWC65-190 clearly showed a cytotoxic effect against HLA-A0201-positive cells presenting the peptide of SEQ ID NO: 40. In contrast, absolutely no cytotoxic effect was observed against control cells not presenting the peptide of SEQ ID NO: 40.

[Example 25]

[0101] The peptide of SEQ ID NO: 33 (amino acid initiation position KDR772) was added to HLA-A0201-positive T2 strain to produce target cells. Using these cells, the cytotoxic effect of the CTL clone (KWC72-772) obtained from the peptide of SEQ ID NO: 33 in Example 13 was examined using a chromium release assay.

[0102] In the results, shown in Fig. 20, CTL clone KWC72-772 clearly showed a cytotoxic effect against HLA-A0201-positive cells presenting the peptide of SEQ ID NO: 33. In contrast, absolutely no cytotoxic effect was observed against control cells that did not present the peptide of SEQ ID NO: 33.

[Example 26]

[0103] The cytotoxic effect of the CTL clone (KWC72-772) against hepatocellular carcinoma cell line HLA-A0201-positive HePG2, forced by adenoviruses to express KDR, was examined by chromium release assay. HLA-A0201-positive HePG2 forced to express EGFP was used as control.

[0104] In the results, as shown in Fig. 21, CTL clone KWC72-772 showed remarkably high cytotoxic effect against HLA-A0201-positive cells presenting KDR peptides on their surface, as compared to activity against cells presenting the EGFP peptide.

[Example 27]

[0105] The peptides of SEQ ID NO: 2 (amino acid initiation position KDR1318), SEQ ID NO: 3 (KDR220), SEQ ID NO: 5 (KDR189), and SEQ ID NO: 11 (KDR826) were added to HLA-A24-positive human B-lymphocyte strain A24-LCL (HLA-A24/24, Takara Shuzo) to produce target cells. Using these cells, the cytotoxic effects of CTL clones, C7-1318, KWC46, C18-189, and C65-826, described in Example 3, were examined by chromium release assay.

[0106] In the results, shown in Figs. 22 to 25, each of the CTL clones, C7-1318, KWC46, C18-189, and C65-826, clearly showed a cytotoxic effect against each of the HLA-A24-positive cells presenting the peptides of SEQ ID NOs: 2, 3, 5, and 11, respectively. In contrast, hardly any or absolutely no cytotoxic effect was observed against each of the control cells, which did not present the peptides.

[Example 28]

[0107] As for Example 11, the cytotoxic effects of the CTL clones, C7-1318, KWC46, C18-189, and C65-826, described in Example 3, against human colon cancer HT29 strain (ATCC), forced by adenoviruses to express KDR, were examined by chromium release assay. HLA-A24-positive HT29 strain forcedly expressing EGFP was used as a control.

[0108] In the results, shown in Figs. 26 to 29, all CTL clones, C7-1318, KWC46, C18-189, and C65-826, showed remarkably high cytotoxic effect against HLA-A24-positive HT29 forcedly expressing KDR, compared to activity against cells forcedly expressing EGFP.

[Example 29]

[0109] As for Example 22, the angiogenesis inhibitory activity of peptide epitopes against the angiogenic response induced by Colon 26 cells in BALB/c mice was evaluated by DAS assay.

[0110] As apparent from the results of Figs. 30 and 31, in the groups vaccinated with DCs pulsed with the peptides of SEQ ID NO: 5 (amino acid initiation position KDR189) and SEQ ID NO: 11 (KDR826), respectively, the formation of tortuous blood vessels was clearly suppressed as compared to the group administered with DC alone. This indicated a statistically significant angiogenesis suppression effect.

[Example 30]

[0111] The CTL response against each type of following epitope peptides was investigated using PBMC of cancer patients after stimulation with each peptide:

Peptides of SEQ ID NO: 8 (amino acid initiation position KDR169), SEQ ID NO: 5 (KDR189), SEQ ID NO: 3 (KDR220), SEQ ID NO: 11 (KDR826), and SEQ ID NO: 17

EP 2 261 249 B1

(KDR1318), with high HLA-24 binding affinity; and peptides of SEQ ID NO: 40 (KDR190), SEQ ID NO: 33 (KDR772), SEQ ID NO: 30 (KDR773), SEQ ID NO: 29 (KDR775), and SEQ ID NO: 34 (KDR1328), with high HLA-A0201 binding affinity.

[0112] Using the method reported by Maeda, Y. et al. in Br. J. Cancer 87, 796-804 (2002), peptide-specific CTLs were detected in PBMC. First, PBMC collected from patients (1×10^5 cells) were incubated with 10 μ M of each peptide in a U-shaped bottom 96-well plate comprising 200 μ L medium. The medium consisted of 45% RPMI1640 medium, 45% AIM-V medium, 10% FBS, 100 U/mL interleukin-2 (IL-2), and 0.1 μ M MEM non-essential amino acid solution. Every three days, half of the medium was removed and replaced with fresh medium containing the corresponding peptide (20 μ M). After incubation for twelve days, cells were collected to test IFN- γ production ability in response to each peptide. When the amount of IFN- γ produced by peptide-stimulated PBMC in response to a corresponding peptide was two times or more than the amount produced in response to the control HIV peptide, that test peptide was deemed positive.

[0113] These results showed that a CTL precursor is produced in HLA-A2- and HLA-A24-type cancer patients, as in a healthy donor (Table 9). In the table, patients A, B, and C are HLA-A24-type patients; and patient D is an HLA-A02-type patient. Patients A and D are colon cancer patients; and patients B and C are melanoma patients. CMVs refer to the cytomegalovirus-derived peptides used as positive controls. The peptides with high A-24 binding affinity are described in Kuzusima, K. et al., Blood 2001, 98(6):p1872-1881, and such; and the peptides with high A-2 binding affinity are described in Solache, A. et al., J. of Immunol. 1999, 163(10); p5512-5518, and such.

Table 9 Cancer patient-derived CTL precursor

	PATIENT A			PATIENT B			PATIENT C			PATIENT D		
	POSITIVE WELLS	TOTAL WELLS		POSITIVE WELLS	TOTAL WELLS		POSITIVE WELLS	TOTAL WELLS		POSITIVE WELLS	TOTAL WELLS	
CMV	1	32		1	24		0	16	CMV	5	20	
KDR 169	12	32		4	32		1	24	KDR 190	3	24	
KDR 189	3	32		1	32		0	24	KDR 772	3	24	
KDR 220	0	32		3	32		0	24	KDR 773	1	24	
KDR 826	11	32		3	32		0	24	KDR 775	2	24	
KDR	0	32		9	32		1	24	KDR	0	24	
1318									1328			

[0114] As for Example 14 and the like, a standard ⁵¹Cr release assay was performed for four hours to establish CTL lines from the CTL precursors (Fig. 32).

Industrial Applicability

[0115] The present invention provides novel peptides, which induce cytotoxic T cells by targeting the endothelial cells formed in a wide range of tumoral tissues, and which are extremely effective as cancer vaccines. The present invention also provides pharmaceuticals for treating and preventing tumors, where the pharmaceuticals comprise these peptides.

SEQUENCE LISTING

[0116]

<110> ONCOTHERAPY SCIENCE, INC.

<120> KDR PEPTIDES AND VACCINES COMPRISING THE SAME

<130> L1360 EP/2 S3

<150> EP 03 79 5432.8

<151> 2003-09-12

<150> JP 2002-267285

<151> 2002-09-12

<150> JP 2003-62003

<151> 2003-03-07

<150> JP 2003-167042

<151> 2003-06-11

<160> 56

<170> PatentIn Ver. 2.0

<210> 1

<211> 9

<212> PRT

<213> Homo sapiens

<400> 1

Trp Tyr Lys Leu Gly Pro Gln Pro Leu
1 5

<210> 2

<211> 9

<212> PRT

<213> Homo sapiens

<400> 2

Val Tyr Ser Ser Glu Glu Ala Glu Leu
1 5

<210> 3

<211> 9

EP 2 261 249 B1

<212> PRT
 <213> Homo sapiens
 <400> 3

5

Gly Tyr Arg Ile Tyr Asp Val Val Leu
 1 5

10 <210> 4
 <211> 9
 <212> PRT
 <213> Homo sapiens

15 <400> 4

Lys Phe Gly Asn Leu Ser Thr Tyr Leu
 1 5

20 <210> 5
 <211> 9
 <212> PRT
 <213> Homo sapiens
 25 <400> 5

Ser Tyr Met Ile Ser Tyr Ala Gly Met
 1 5

30 <210> 6
 <211> 9
 <212> PRT
 <213> Homo sapiens
 35 <400> 6

Glu Phe Pro Arg Asp Arg Leu Lys Leu
 1 5

40 <210> 7
 <211> 9
 <212> PRT
 <213> Homo sapiens
 45 <400> 7

Thr Phe Ser Glu Leu Val Glu His Leu
 1 5

50 <210> 8
 <211> 9
 <212> PRT
 <213> Homo sapiens

EP 2 261 249 B1

<400> 8

5 Arg Phe Val Pro Asp Gly Asn Arg Ile
1 5

<210> 9

<211> 9

<212> PRT

10 <213> Homo sapiens

<400> 9

15 Ala Phe Gly Ser Gly Met Glu Ser Leu
1 5

<210> 10

<211> 9

20 <212> PRT

<213> Homo sapiens

<400> 10

25 Lys Asn Leu Asp Thr Leu Trp Lys Leu
1 5

<210> 11

<211> 9

30 <212> PRT

<213> Homo sapiens

<400> 11

35 Lys Trp Glu Phe Pro Arg Asp Arg Leu
1 5

<210> 12

40 <211> 9

<212> PRT

<213> Homo sapiens

<400> 12

45 Asp Phe Leu Thr Leu Glu His Leu Ile

50 1 5

<210> 13

<211> 10

<212> PRT

55 <213> Artificial Sequence

<220>

<221> Source

EP 2 261 249 B1

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 13

5
Asp Tyr Val Gly Ala Ile Pro Val Asp Leu
1 5 10

<210> 14

10 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

15 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 14

20
Ser Tyr Gln Tyr Gly Thr Thr Gln Thr Leu
1 5 10

<210> 15

25 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

30 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 15

35
Asp Tyr Val Arg Lys Gly Asp Ala Arg Leu
1 5 10

<210> 16

40 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

45 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 16

50
Leu Tyr Thr Cys Ala Ala Ser Ser Gly Leu
1 5 10

<210> 17

55 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

EP 2 261 249 B1

<221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 17

5

Val Tyr Ser Ser Glu Glu Ala Glu Leu Leu

10

1

5

10

<210> 18

<211> 10

<212> PRT

15

<213> Artificial Sequence

<220>

<221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

20

<400> 18

Leu Tyr Thr Cys Gln Ala Cys Ser Val Leu

25

1

5

10

<210> 19

<211> 10

<212> PRT

30

<213> Artificial Sequence

<220>

<221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

35

<400> 19

Thr Tyr Leu Arg Ser Lys Arg Asn Glu Phe

40

1

5

10

<210> 20

<211> 10

<212> PRT

45

<213> Artificial Sequence

<220>

<221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

50

<400> 20

Lys Tyr Leu Gly Tyr Pro Pro Pro Glu Ile

55

1

5

10

<210> 21

<211> 10

EP 2 261 249 B1

<212> PRT

<213> Artificial Sequence

<220>

5 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 21

10

Val	Tyr	Val	Gln	Asp	Tyr	Arg	Ser	Pro	Phe
1				5					10

<210> 22

15 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

20 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 22

25

Asn	Tyr	Thr	Val	Ile	Leu	Thr	Asn	Pro	Ile
1				5					10

<210> 23

30 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

35 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 23

40

Phe	Phe	Trp	Leu	Leu	Leu	Val	Ile	Ile	Leu
1				5					10

<210> 24

45 <211> 10

<212> PRT

<213> Artificial Sequence

<220>

50 <221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 24

55

Ser	Tyr	Ala	Gly	Met	Val	Phe	Cys	Glu	Ala
1				5					10

EP 2 261 249 B1

<210> 25
 <211> 9
 <212> PRT
 <213> Homo sapiens
 5
 <400> 25
 Thr Tyr Ala Cys Phe Val Ser Asn Leu
 1 5
 10
 <210> 26
 <211> 9
 <212> PRT
 <213> Homo sapiens
 15
 <400> 26
 Val Leu Leu Trp Glu Ile Phe Ser Leu
 1 5
 20
 <210> 27
 <211> 9
 <212> PRT
 <213> Homo sapiens
 25
 <400> 27
 Ser Leu Gln Asp Gln Gly Asp Tyr Val
 1 5
 30
 <210> 28
 <211> 9
 <212> PRT
 <213> Homo sapiens
 35
 <400> 28
 Val Leu Leu Ala Val Ala Leu Trp Leu
 1 5
 40
 <210> 29
 <211> 9
 <212> PRT
 <213> Homo sapiens
 45
 <400> 29
 Ala Met Phe Phe Trp Leu Leu Leu Val
 1 5
 50
 <210> 30
 <211> 9
 <212> PRT
 <213> Homo sapiens
 55

EP 2 261 249 B1

<400> 30

5 Val Ile Ala Met Phe Phe Trp Leu Leu
1 5

<210> 31

<211> 9

<212> PRT

10 <213> Homo sapiens

<400> 31

15 Ile Leu Leu Ser Glu Lys Asn Val Val
1 5

<210> 32

<211> 9

20 <212> PRT

<213> Homo sapiens

<400> 32

25 Lys Asn Leu Asp Thr Leu Trp Lys Leu
1 5

<210> 33

30 <211> 9

<212> PRT

<213> Homo sapiens

<400> 33

35 Ala Val Ile Ala Met Phe Phe Trp Leu
1 5

<210> 34

<211> 9

<212> PRT

<213> Homo sapiens

<400> 34

45 Lys Leu Ile Glu Ile Gly Val Gln Thr
1 5

50 <210> 35

<211> 9

<212> PRT

<213> Homo sapiens

55 <400> 35

EP 2 261 249 B1

Ile Met Trp Phe Lys Asp Asn Glu Thr
1 5

5 <210> 36
<211> 9
<212> PRT
<213> Homo sapiens

10 <400> 36

Thr Leu Trp Lys Leu Asn Ala Thr Met
1 5

15 <210> 37
<211> 9
<212> PRT
<213> Homo sapiens

20 <400> 37

Phe Gln Gly Gly Asn Lys Ile Glu Val
1 5

25 <210> 38
<211> 9
<212> PRT
<213> Homo sapiens

30 <400> 38

Tyr Gln Tyr Gly Thr Thr Gln Thr Leu
1 5

35 <210> 39
<211> 9
<212> PRT
<213> Homo sapiens

40 <400> 39

Ala Leu Ile Glu Gly Lys Asn Lys Thr
1 5

45 <210> 40
<211> 9
<212> PRT
<213> Homo sapiens

50 <400> 40

Tyr Met Ile Ser Tyr Ala Gly Met Val
1 5

55

EP 2 261 249 B1

<210> 41
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 5
 <220>
 <221> Source
 <223> /note ="Description of Artificial Sequence: Synthetic peptide"
 10
 <400> 41

 Leu Leu Trp Glu Ile Phe Ser Leu Gly Ala
 1 5 10
 15
 <210> 42
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 20
 <220>
 <221> Source
 <223> /note ="Description of Artificial Sequence: Synthetic peptide"
 25
 <400> 42

 Val Leu Leu Ala Val Ala Leu Trp Leu Cys
 1 5 10
 30
 <210> 43
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 35
 <220>
 <221> Source
 <223> /note ="Description of Artificial Sequence: Synthetic peptide"
 40
 <400> 43

 Leu Met Thr Lys Lys Asn Ser Thr Phe Val
 1 5 10
 45
 <210> 44
 <211> 10
 <212> PRT
 <213> Artificial Sequence
 50
 <220>
 <221> Source
 <223> /note ="Description of Artificial Sequence: Synthetic peptide"
 55
 <400> 44

EP 2 261 249 B1

Trp Leu Leu Leu Val Ile Ile Leu Arg Thr
1 5 10

5 <210> 45
<211> 10
<212> PRT
<213> Artificial Sequence

10 <220>
<221> Source
<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 45

Ala Leu Ile Glu Gly Lys Asn Lys Thr Val
1 5 10

20 <210> 46
<211> 10
<212> PRT
<213> Artificial Sequence

25 <220>
<221> Source
<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 46

Ile Gln Ser Asp Val Trp Ser Phe Gly Val
1 5 10

35 <210> 47
<211> 10
<212> PRT
<213> Artificial Sequence

40 <220>
<221> Source
<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 47

Gly Leu Tyr Thr Cys Gln Ala Cys Ser Val
1 5 10

50 <210> 48
<211> 10
<212> PRT
<213> Artificial Sequence

55 <220>
<221> Source
<223> /note ="Description of Artificial Sequence: Synthetic peptide"

EP 2 261 249 B1

<400> 48

5 Leu Leu Leu Val Ile Ile Leu Arg Thr Val
 1 5 10

<210> 49

<211> 10

<212> PRT

10 <213> Artificial Sequence

<220>

<221> Source

15 <223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 49

20 Leu Leu Ala Val Ala Leu Trp Leu Cys Val
 1 5 10

<210> 50

<211> 10

<212> PRT

25 <213> Artificial Sequence

<220>

<221> Source

30 <223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 50

35 Tyr Ile Thr Glu Asn Lys Asn Lys Thr Val
 1 5 10

<210> 51

<211> 10

<212> PRT

40 <213> Artificial Sequence

<220>

<221> Source

45 <223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 51

50 Leu Leu Ser Glu Lys Asn Val Val Lys Ile
 1 5 10

<210> 52

<211> 10

<212> PRT

55 <213> Artificial Sequence

<220>

<221> Source

EP 2 261 249 B1

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 52

5
Lys Val Leu Leu Ala Val Ala Leu Trp Leu
1 5 10

<210> 53

10 <211> 10

<212> PRT

<213> Homo sapiens

<400> 53

15
Asp Val Leu Tyr Gly Pro Asp Thr Pro Ile
1 5 10

20 <210> 54

<211> 9

<212> PRT

<213> Artificial Sequence

25 <220>

<221> Source

<223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 54

30
Val Leu Ala Met Phe Phe Trp Leu Leu
1 5

<210> 55

35 <211> 9

<212> PRT

<213> Artificial Sequence

<220>

40 <223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 55

45
Ile Leu Lys Glu Pro Val His Gly Val
1 5

<210> 56

<211> 9

50 <212> PRT

<213> Artificial Sequence

<220>

<221> Source

55 <223> /note ="Description of Artificial Sequence: Synthetic peptide"

<400> 56

Arg Tyr Leu Arg Asp Gln Gln Leu Leu
1 5

Claims

1. A nonapeptide consisting of the amino acid sequence of SEQ ID NO: 2.
2. A peptide with cytotoxic T cell inducibility, wherein one or two amino acids are substituted or added to the amino acid sequence of SEQ ID NO: 2, wherein the second amino acid from the N-terminus is phenylalanine, tyrosine, methionine, or tryptophan.
3. A peptide with cytotoxic T cell inducibility, wherein one or two amino acids are substituted or added to the amino acid sequence of SEQ ID NO: 2, wherein the C-terminal amino acid is phenylalanine, leucine, isoleucine, tryptophan, or methionine.
4. The peptide of claim 2, wherein the C-terminal amino acid is phenylalanine, leucine, isoleucine, tryptophan, or methionine.
5. A pharmaceutical composition comprising one or more peptides of any one of claims 1 to 4.
6. A vaccine comprising a peptide of any one of claims 1 to 4 as an active ingredient.
7. A peptide for use in treating and/or preventing tumors, wherein the peptide is selected from the group consisting of the peptides of claims 2 to 4.
8. A peptide for use in treating diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, wherein the peptide is selected from the group consisting of the peptides of claims 1 to 4.
9. A pharmaceutical composition for use in treating and/or preventing tumors, wherein the pharmaceutical composition comprises one or more peptides of any one of claims 2 to 4.
10. A pharmaceutical composition for use in treating diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, wherein the pharmaceutical composition comprises one or more peptides of any one of claims 1 to 4.
11. An exosome that presents on its surface a complex consisting of a peptide of any one of claims 1 to 4, and an HLA-A2402 antigen.
12. A vaccine for use in treating and/or preventing tumors, wherein the vaccine comprises a peptide of any one of claims 2 to 4 as an active ingredient.
13. A vaccine for use in treating diabetic retinopathy, chronic rheumatoid arthritis, psoriasis, and atherosclerosis, wherein the vaccine comprises a peptide of any one of claims 1 to 4 as an active ingredient.
14. The peptide of claim 7 or 8, the pharmaceutical composition of claim 9 or 10, or the vaccine of claim 12 or 13, which is used for administration to a subject whose HLA antigen is HLA-A24.
15. The peptide, the pharmaceutical composition, or the vaccine of claim 14, which is used for administration to a subject whose HLA antigen is HLA-A2402.

Patentansprüche

1. Nonapeptid, bestehend aus der Aminosäuresequenz der SEQ ID NO:2.
2. Peptid mit cytotoxischer T-Zell-Induzierbarkeit, wobei eine oder zwei Aminosäuren in der Aminosäuresequenz der SEQ ID NO:2 ersetzt oder hinzugefügt sind, wobei die zweite Aminosäure vom N-Terminus Phenylalanin, Tyrosin, Methionin oder Tryptophan ist.

3. Peptid mit cytotoxischer T-Zell-Induzierbarkeit, wobei eine oder zwei Aminosäuren in der Aminosäuresequenz der SEQ ID NO:2 ersetzt oder hinzugefügt sind, wobei die C-terminale Aminosäure Phenylalanin, Leucin, Isoleucin, Tryptophan oder Methionin ist.
- 5 4. Peptid nach Anspruch 2, wobei die C-terminale Aminosäure Phenylalanin, Leucin, Isoleucin, Tryptophan oder Methionin ist.
5. Arzneimittel, umfassend ein oder mehrere Peptide nach einem der Ansprüche 1 bis 4.
- 10 6. Impfstoff, umfassend ein Peptid nach einem der Ansprüche 1 bis 4 als Wirkstoff.
7. Peptid zur Verwendung bei der Behandlung und/oder Vorbeugung von Tumoren, wobei das Peptid ausgewählt ist aus der Gruppe bestehend aus den Peptiden nach den Ansprüchen 2 bis 4.
- 15 8. Peptid zur Verwendung bei der Behandlung von diabetischer Retinopathie, chronischer rheumatoider Arthritis, Psoriasis und Atherosklerose, wobei das Peptid ausgewählt ist aus der Gruppe bestehend aus den Peptiden nach den Ansprüchen 1 bis 4.
- 20 9. Arzneimittel zur Verwendung bei der Behandlung und/oder Vorbeugung von Tumoren, wobei das Arzneimittel ein oder mehrere Peptide nach einem der Ansprüche 2 bis 4 umfasst.
10. Arzneimittel zur Verwendung bei der Behandlung von diabetischer Retinopathie, chronischer rheumatoider Arthritis, Psoriasis und Atherosklerose, wobei das Arzneimittel ein oder mehrere Peptide nach einem der Ansprüche 1 bis 4 umfasst.
- 25 11. Exosom, das auf seiner Oberfläche einen Komplex präsentiert, der aus einem Peptid nach einem der Ansprüche 1 bis 4 und einem HLA-A2402-Antigen besteht.
12. Impfstoff zur Verwendung bei der Behandlung und/oder Vorbeugung von Tumoren, wobei der Impfstoff ein Peptid nach einem der Ansprüche 2 bis 4 als Wirkstoff umfasst.
- 30 13. Impfstoff zur Verwendung bei der Behandlung von diabetischer Retinopathie, chronischer rheumatoider Arthritis, Psoriasis und Atherosklerose, wobei der Impfstoff ein Peptid nach einem der Ansprüche 1 bis 4 als Wirkstoff umfasst.
- 35 14. Peptid nach Anspruch 7 oder 8, Arzneimittel nach Anspruch 9 oder 10 oder Impfstoff nach Anspruch 12 oder 13, das/der zur Verabreichung an ein Individuum verwendet wird, dessen HLA-Antigen HLA-A24 ist.
- 40 15. Peptid, Arzneimittel oder Impfstoff nach Anspruch 14, das/der zur Verabreichung an ein Individuum verwendet wird, dessen HLA-Antigen HLA-A2402 ist.

Revendications

- 45 1. Nonapeptide constitué de la séquence d'acides aminés de SEQ ID NO : 2.
2. Peptide possédant la capacité d'induire les lymphocytes T cytotoxiques, dans lequel un ou deux acides aminés sont substitués ou ajoutés à la séquence d'acides aminés de SEQ ID NO : 2, où le deuxième acide aminé à partir de l'extrémité N-terminale est la phénylalanine, la tyrosine, la méthionine ou le tryptophane.
- 50 3. Peptide possédant la capacité d'induire les lymphocytes T cytotoxiques, dans lequel un ou deux acides aminés sont substitués ou ajoutés à la séquence d'acides aminés de SEQ ID NO : 2, où l'acide aminé C-terminal est la phénylalanine, la leucine, l'isoleucine, le tryptophane ou la méthionine.
- 55 4. Peptide selon la revendication 2, dans lequel l'acide aminé C-terminal est la phénylalanine, la leucine, l'isoleucine, le tryptophane ou la méthionine.
5. Composition pharmaceutique comprenant un ou plusieurs peptides selon l'une quelconque des revendications 1 à 4.

6. Vaccin comprenant un peptide selon l'une quelconque des revendications 1 à 4 en tant que principe actif.
7. Peptide pour une utilisation dans le traitement et/ou la prévention des tumeurs, où le peptide est choisi dans le groupe constitué des peptides selon les revendications 2 à 4.
8. Peptide pour une utilisation dans le traitement de la rétinopathie diabétique, de la polyarthrite rhumatoïde chronique, du psoriasis et de l'athérosclérose, où le peptide est choisi dans le groupe constitué des peptides selon les revendications 1 à 4.
9. Composition pharmaceutique pour une utilisation dans le traitement et/ou la prévention des tumeurs, où la composition pharmaceutique comprend un ou plusieurs peptides selon l'une quelconque des revendications 2 à 4.
10. Composition pharmaceutique pour une utilisation dans le traitement de la rétinopathie diabétique, de la polyarthrite rhumatoïde chronique, du psoriasis et de l'athérosclérose, où la composition pharmaceutique comprend un ou plusieurs peptides selon l'une quelconque des revendications 1 à 4.
11. Exosome qui présente sur sa surface un complexe constitué d'un peptide selon l'une quelconque des revendications 1 à 4, et un antigène HLA-A2402.
12. Vaccin pour une utilisation dans le traitement et/ou la prévention des tumeurs, où le vaccin comprend un peptide selon l'une quelconque des revendications 2 à 4 en tant que principe actif.
13. Vaccin pour une utilisation dans le traitement de la rétinopathie diabétique, de la polyarthrite rhumatoïde chronique, du psoriasis et de l'athérosclérose, où le vaccin comprend un peptide selon l'une quelconque des revendications 1 à 4 en tant que principe actif.
14. Peptide selon la revendication 7 ou 8, composition pharmaceutique selon la revendication 9 ou 10, ou vaccin selon la revendication 12 ou 13, qui est utilisé pour une administration à un sujet dont l'antigène HLA est HLA-A24.
15. Peptide, composition pharmaceutique, ou vaccin selon la revendication 14, qui est utilisé pour une administration à un sujet dont l'antigène HLA est HLA-A2402.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- WO 994380 A [0005]
- US 5861301 A [0011]
- WO 9967288 A [0011]
- JP HE111510507 B [0027]
- JP 2000512161 A [0027]
- JP 2000509281 A [0030]
- EP 03795432 A [0116]
- JP 2002267285 A [0116]
- JP 2003062003 A [0116]
- JP 2003167042 A [0116]

Non-patent literature cited in the description

- *The Journal of Experimental Medicine*, 2002, vol. 195 (12), 1575-1584 [0004]
- Peptide Synthesis. Interscience, 1966 [0011]
- The Proteins. Academic Press Inc, 1976, vol. 2 [0011]
- Peptide Synthesis. Maruzen Co., Ltd, 1975 [0011]
- Foundations and Experiments in Peptide Synthesis. Maruzen Co., Ltd, 1985 [0011]
- Peptide Synthesis. Development of Pharmaceuticals, New Series. Hirokawa Shoten, 1991, vol. 14 [0011]
- PARKER K. C. *J. Immunol.*, 1994, 152 [0012]
- PARKER, K. C. *J. Immunol.*, 1994, 152 [0012]
- NUKAYA, I. *Int. J. Cancer*, 1999, 80 [0012]
- *J. Immunol.*, 1994, vol. 152, 3913 [0013]
- *Immunogenetics*, 1995, vol. 41, 178 [0013]
- *J. Immunol.*, 1994, vol. 155, 4307 [0013]
- BENMOHAMED L. ; KRISHNAN R. ; LONGMATE J. ; AUGE C. ; LOW L. ; PRIMUS J. ; DIAMOND DJ. Related Articles, Books, Linkout Induction of CTL response by a minimal epitope vaccine in HLA A*0201/DR1 transgenic mice: dependence on HLA class II restricted T(H) response. *Hum. Immunol.*, 2000, vol. 61 (8), 764-79 [0015]
- FOLKMAN, J. *Nature Med*, 1995, vol. 1, 27-31 [0021]
- BICKNELL, R. ; HARRIS, A.L. *Curr. Opin. Oncol.*, 1996, vol. 8, 60-65 [0021]
- *Clin. Microbiol. Rev.*, 1994, vol. 7, 277-289 [0025]
- *Cancer Res.*, 1996, vol. 56, 5672 [0030]
- *J. Immunol.*, 1998, vol. 161, 5607 [0030]
- *J. Exp. Med.*, 1996, vol. 184, 465 [0030]
- *Int. J. Cancer*, 1999, 80 [0036]
- PARKER KO. *J. Immunol*, 1994, 152 [0037]
- NUKAYA I. *Int. J. Cancer*, 1999, 80 [0037] [0043]
- PARKER.KC. *J. Immunol*, 1994, 152 [0043]
- TANAKA, H. et al. *AACR #3669*, March 2001, vol. 42, 681-682 [0050]
- TANAKA H et al. *AACR*, 2001 [0050] [0058]
- BENMOHAMED L. ; KRISHNAN R. ; LONGMATE J. ; AUGE C. ; LOW L. ; PRIMUS J. ; DIAMOND DJ. Related Articles, Books, Linkout Induction of CTL response by a minimal epitope vaccine in HLA A*0201/DR1 transgenic mice: dependence on HLA class II restricted T(H) response. *Hum. Immunol.*, August 2000, vol. 61 (8), 764-79 [0051]
- *J. Biol. Chem.*, 28 October 1994, vol. 269 (43), 26988-95 [0067]
- MIYAKE, S. et al. *Proc. Natl. Acad. Sci. USA.*, 1996, vol. 93, 1320-1324 [0067]
- *Leukemia*, April 1999, vol. 13 (4), 605-13 [0067]
- *Clinical Cancer Research*, August 1999, vol. 5, 2185-2191 [0092]
- *Cancer Research*, 01 November 2002, vol. 62, 6116-6123 [0092]
- MAEDA, Y. et al. *Br. J. Cancer*, 2002, vol. 87, 796-804 [0112]
- KUZUSLIMA, K. et al. *Blood*, 2001, vol. 98 (6), 1872-1881 [0113]
- SOLACHE, A. et al. *J. of Immunol.*, 1999, vol. 163 (10), 5512-5518 [0113]