



(86) Date de dépôt PCT/PCT Filing Date: 1995/10/03
(87) Date publication PCT/PCT Publication Date: 1996/04/11
(45) Date de délivrance/Issue Date: 2009/07/21
(85) Entrée phase nationale/National Entry: 1997/03/27
(86) N° demande PCT/PCT Application No.: US 1995/012963
(87) N° publication PCT/PCT Publication No.: 1996/010404
(30) Priorité/Priority: 1994/10/03 (US317,226)

(51) Cl.Int./Int.Cl. *A61K 31/353* (2006.01),
A61K 36/185 (2006.01), *A61P 35/00* (2006.01),
C07D 311/62 (2006.01), *C09K 15/34* (2006.01)

(72) Inventeurs/Inventors:
ROMANCZYK, LEO J., JR., US;
HAMMERSTONE, JOHN F., JR., US;
BUCK, MARGARET M., US

(73) Propriétaire/Owner:
MARS, INCORPORATED, US

(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : EXTRAITS ANTINEOPLASIQUES DE CACAO ET LEURS PROCEDES DE PREPARATION ET
D'UTILISATION

(54) Title: ANTINEOPLASTIC COCOA EXTRACTS AND METHODS FOR MAKING AND USING THE SAME

(57) **Abrégé/Abstract:**

Disclosed and claimed are cocoa extracts such as polyphenols or procyanidins, methods for preparing such extracts, as well as uses for them, especially as antineoplastic agents and antioxidants. Disclosed and claimed are antineoplastic compositions containing cocoa polyphenols or procyanidins and methods for treating patients employing the compositions. Additionally disclosed and claimed is a kit for treating a patient in need of treatment with an antineoplastic agent containing cocoa polyphenols or procyanidins as well as a lyophilized antineoplastic composition containing cocoa polyphenols or procyanidins. Further, disclosed and claimed is the use of the invention in antioxidant, preservative and topoisomerase-inhibiting compositions and methods.



ABSTRACT

Disclosed and claimed are cocoa extracts such as polyphenols or procyanidins, methods for preparing such extracts, as well as uses for them, especially as
5 antineoplastic agents and antioxidants. Disclosed and claimed are antineoplastic compositions containing cocoa polyphenols or procyanidins and methods for treating patients employing the compositions. Additionally disclosed and claimed is a kit for treating a patient in
10 need of treatment with an antineoplastic agent containing cocoa polyphenols or procyanidins as well as a lyophilized antineoplastic composition containing cocoa polyphenols or procyanidins. Further, disclosed and claimed is the use of the invention in antioxidant,
15 preservative and topiosomerase-inhibiting compositions and methods.

ANTINEOPLASTIC COCOA EXTRACTS AND METHODS FOR MAKING AND
USING THE SAME

FIELD OF THE INVENTION

This invention relates to cocoa extracts such
5 as polyphenols preferably polyphenols enriched with
procyanidins. This invention also relates to methods for
preparing such extracts, as well as to uses for them; for
instance, as antineoplastic agents and antioxidants.

Documents are referred to in this disclosure
10 with full details for each appearing in a References
section at the end of the specification, preceding the
claims. These documents pertain to the field of this
invention.

15 BACKGROUND OF THE INVENTION

Polyphenols are an incredibly diverse group of
compounds (Ferreira *et al.*, 1992) which widely occur in a
variety of plants, some of which enter into the food
chain. In some cases they represent an important class
20 of compounds for the human diet. Although some of the
polyphenols are considered to be nonnutrative, interest
in these compounds has arisen because of their possible
beneficial effects on health. For instance, quercetin (a
flavonoid) has been shown to possess anticarcinogenic
25 activity in experimental animal studies (Deshner *et al.*,
1991 and Kato *et al.*, 1983). (+)-Catechin and (-)-
epicatechin (flavan-3-ols) have been shown to inhibit
Leukemia virus reverse transcriptase activity (Chu *et al.*,
1992). Nobotanin (an oligomeric hydrolyzable
30 tannin) has also been shown to possess anti-tumor
activity (Okuda *et al.*, 1992). Statistical reports have
also shown that stomach cancer mortality is significantly
lower in the tea producing districts of Japan.
Epigallocatechin gallate has been reported to be the
35 pharmacologically active material in green tea that
inhibits mouse skin tumors (Okuda *et al.*, 1992). Ellagic
acid has also been shown to possess anticarcinogen
activity in various animal tumor models (Bukharta *et al.*,

1992). Lastly, proanthocyanidin oligomers have been patented by the Kikkoman Corporation for use as antimutagens. Indeed, the area of phenolic compounds in foods and their modulation of tumor development in experimental animal models has been recently presented at the 202nd National Meeting of The American Chemical Society (Ho et al., 1992; Huang et al., 1992).

However, none of these reports teaches or suggests cocoa extracts, any methods for preparing such extracts, or, any uses as antineoplastic agents for cocoa extracts.

Since unfermented cocoa beans contain substantial levels of polyphenols, the present inventors considered it possible that similar activities of and uses for cocoa extracts, e.g., compounds within cocoa, could be revealed by extracting such compounds from cocoa and screening the extracts for activity. The National Cancer Institute has screened various *Theobroma* and *Herrania* species for anti-cancer activity as part of their massive natural product selection program. Low levels of activity were reported in some extracts of cocoa tissues, and the work was not pursued. Thus, in the antineoplastic or anti-cancer art, cocoa and its extracts were not deemed to be useful; i.e., the teachings in the antineoplastic or anti-cancer art lead the skilled artisan away from employing cocoa and its extracts as cancer therapy. Since a number of analytical procedures were developed to study the contributions of cocoa polyphenols to flavor development (Clapperton et al., 1992), the present inventors decided to apply analogous methods to prepare samples for anti-cancer screening, contrary to the knowledge in the antineoplastic or anti-cancer art. Surprisingly, and contrary to the knowledge in the art, e.g., the National Cancer Institute screening, the present inventors discovered that cocoa polyphenol extracts which contain procyanidins, have significant utility as anti-cancer or

antineoplastic agents. Additionally, the inventors demonstrate that cocoa extracts containing procyanidins have utility as antioxidants.

OBJECTS AND SUMMARY OF THE INVENTION

5 It is an object of the present invention to provide a method for producing cocoa extract.

It is another object of the invention to provide a cocoa extract.

10 It is another object of the invention to provide an antioxidant composition.

It is another object of the invention to demonstrate inhibition of DNA topoisomerase II enzyme activity.

15 It is yet another object of the present invention to provide a method for treating tumors or cancer.

It is still another object of the invention to provide an anti-cancer, anti-tumor or antineoplastic composition.

20 It is a further object of the invention to provide a method for making an anti-cancer, anti-tumor or antineoplastic composition.

And, it is an object of the invention to provide a kit for use in treating tumors or cancer.

25 It has been surprisingly discovered that cocoa extract has anti-tumor, anti-cancer or antineoplastic activity; or, is an antioxidant composition or, inhibits DNA topoisomerase II enzyme activity. Accordingly, the present invention provides a substantially pure cocoa
30 extract. The extract preferably comprises polyphenol(s) such as polyphenol(s) enriched with cocoa procyanidin(s), such as polyphenols of at least one cocoa procyanidin selected from (-) epicatechin, procyanidin B-2, procyanidin oligomers 2 through 12, preferably 2 through
35 5 or 4 through 12, procyanidin B-5, procyanidin A-2 and procyanidin C-1. The present invention also provides an anti-tumor, anti-cancer or antineoplastic or antioxidant

or DNA topoisomerase II inhibitor composition comprising a substantially pure cocoa extract or synthetic cocoa polyphenol(s) such as polyphenol(s) enriched with procyanidin(s) and a suitable carrier. The extract
5 preferably comprises cocoa procyanidin(s). The cocoa extract is preferably obtained by a process comprising reducing cocoa beans to powder, defatting the powder and, extracting active compound(s) from the powder.

The present invention further comprehends a method
10 for treating a patient in need of treatment with an anti-tumor, anti-cancer, or antineoplastic agent or an antioxidant or a DNA topoisomerase II inhibitor comprising administering to the patient a composition comprising an effective quantity of a substantially pure
15 cocoa extract or synthetic cocoa polyphenol(s) or procyanidin(s) and a carrier. The cocoa extract can be cocoa procyanidin(s); and, is preferably obtained by reducing cocoa beans to powder, defatting the powder and, extracting active compound(s) from the powder.

20 Additionally, the present invention provides a kit for treating a patient in need of treatment with an anti-tumor, anti-cancer, or antineoplastic agent or antioxidant or DNA topoisomerase II inhibitor comprising a substantially pure cocoa extract or synthetic cocoa
25 polyphenol(s) or procyanidin(s) and a suitable carrier for admixture with the extract or synthetic polyphenol(s) or procyanidin(s).

In one embodiment there is preferably provided a solid or liquid composition suitable for oral

4a

administration which comprises a suitable carrier and a cocoa extract comprising at least one cocoa procyanidin oligomer 4 to 12.

5 In another preferred embodiment there is provided a cocoa extract comprising cocoa polyphenols, which extract is obtainable by a process which comprises: reducing unfermented or partially fermented cocoa beans to a powder; defatting the powder; extracting cocoa polyphenols from the powder by solvent extraction
10 techniques; and purifying the cocoa extract thus obtained.

These and other objects and embodiments are disclosed or will be obvious from the following Detailed Description.

15 **BRIEF DESCRIPTION OF THE DRAWINGS**

The following Detailed Description will be better understood by reference to the accompanying drawings wherein:

Fig. 1 shows a representative gel permeation
20 chromatogram from the fractionation of crude cocoa procyanidins;

Fig. 2A shows a representative reverse-phase HPLC chromatogram showing the separation (elution profile) of cocoa procyanidins extracted from unfermented cocoa;

5 Fig. 2B shows a representative normal phase HPLC separation of cocoa procyanidins extracted from unfermented cocoa;

Fig. 3 shows several representative procyanidin structures;

10 Figs. 4A-4E show representative HPLC chromatograms of five fractions employed in screening for anti-cancer or antineoplastic activity;

Figs. 5 and 6A-6D show the dose-response relationship between cocoa extracts and cancer cells ACHN (Fig. 5) and PC-3 (Figs. 6A-6D) (fractional survival vs. dose, $\mu\text{g/ml}$); M&M2 F4/92, M&MA+E U12P1, M&MB+E Y192P1, M&MC+E U12P2, M&MD+E U12P2;

20 Figs. 7A to 7H show the typical dose response relationships between cocoa procyanidin fractions A, B, C, D, E, A+B, A+E, and A+D, and the PC-3 cell line (fractional survival vs. dose, $\mu\text{g/ml}$); MM-1A 0212P3, MM-1 B 0162P1, MM-1 C 0122P3, MM-1 D 0122P3, MM-1 E 0292P8, MM-1 A/B 0292P6, MM-1 A/E 0292P6, MM-1 A/D 0292P6;

25 Figs. 8A to 8H show the typical dose response relationships between cocoa procyanidin fractions A, B, C, D, E, A+B, B+E, and D+E and the KB Nasopharyngeal/HeLa cell line (fractional survival vs. dose, $\mu\text{g/ml}$); MM-1A092K3, MM-1 B 0212K5, MM-1 C 0162K3, MM-1 D 0212K5, MM-1 E 0292K5, MM-1 A/B 0292K3, MM-1 B/E 0292K4, MM-1 D/E 30 0292K5;

Figs. 9A to 9H show the typical dose response relationship between cocoa procyanidin fractions A, B, C, D, E, B+D, A+E and D+E and the HCT-116 cell line (fractional survival vs. dose, $\mu\text{g/ml}$); MM-1 C 0192H5, D 35 0192H5, E 0192H5, MM-1 B&D 0262H2, A/E 0262H3, MM-1 D&E 0262H1;

Figs. 10A to 10H show typical dose response relationships between cocoa procyanidin fractions A, B, C, D, E, B+D, C+D and A+E and the ACHN renal cell line (fractional survival vs. dose, $\mu\text{g/ml}$); MM-1 A 092A5, MM-1
5 B 092A5, MM-1 C 0192A7, MM-1 D 0192A7, M&M1 E 0192A7, MM-1 B&D 0302A6, MM-1 C&D 0302A6, MM-1 A&E 0262A6;

Figs. 11A to 11H show typical dose response relationships between cocoa procyanidin fractions A, B, C, D, E, A+E, B+E and C+E and the A-549 lung cell line
10 (fractional survival vs. dose, $\mu\text{g/ml}$); MM-1 A 019258, MM-1 B 09256, MM-1 C 019259, MM-1 D 019258, MM-1 E 019258, A/E 026254, MM-1 B&E 030255, MM-1 C&E N6255;

Figs. 12A to 12H show typical dose response relationships between cocoa procyanidin fractions A, B,
15 C, D, E, B+C, C+D and D+E and the SK-5 melanoma cell line (fractional survival vs. dose $\mu\text{g/ml}$); MM-1 A 0212S4, MM-1 B 0212S4, MM-1 C 0212S4, MM-1 D 0212S4, MM-1 E N32S1, MM-1 B&C N32S2, MM-1 C&D N32S3, MM-1 D&E N32S3;

Figs. 13A to 13H show typical dose response
20 relationships between cocoa procyanidin fractions A, B, C, D, E, B+C, C+E, and D+E and the MCF-7 breast cell line (fractional survival vs. dose, $\mu\text{g/ml}$); MM-1 A N22M4, MM-1 B N22M4, MM-1 C N22M4, MM-1 D N22M3, MM-1 E 0302M2, MM-1 B/C 0302M4, MM-1 C&E N22M3, MM-1 D&E N22M3;

25 Fig. 14 shows typical dose response relationships for cocoa procyanidin (particularly fraction D) and the CCRF-CEM T-cell leukemia cell line (cells/ml vs. days of growth; open circle is control, darkened circle is 125 μg fraction D, open inverted
30 triangle is 250 μg fraction D, darkened inverted triangle is 500 μg fraction D);

Fig. 15 A shows a comparison of the XTT and Crystal Violet cytotoxicity assays against MCF-7 p168 breast cancer cells treated with fraction D+E (open
35 circle is XTT and darkened circle is Crystal Violet);

Fig. 15 B shows a typical dose response curve obtained from MDA MB231 breast cell line treated with

varying levels of crude polyphenols obtained from UIT-1 cocoa genotype (absorbance (540 nm) vs. Days; open circle is control, darkened circle is vehicle, open inverted triangle is 250 $\mu\text{g/ml}$, darkened inverted triangle is 100 $\mu\text{g/ml}$, open square is 10 $\mu\text{g/ml}$; absorbance of 2.0 is maximum of plate reader and may not be necessarily representative of cell number);

Fig. 15 C shows a typical dose response curve obtained from PC-3 prostate cancer cell line treated with varying levels of crude polyphenols obtained from UIT-1 cocoa genotype (absorbance (540nm) vs. Days; open circle is control, darkened circle is vehicle, open inverted triangle is 250 $\mu\text{g/ml}$, darkened inverted triangle is 100 $\mu\text{g/ml}$ and open square is 10 $\mu\text{g/ml}$);

Fig. 15 D shows a typical dose-response curve obtained from MCF-7 p168 breast cancer cell line treated with varying levels of crude polyphenols obtained from UIT-1 cocoa genotype (absorbance (540 nm) vs. Days; open circle is control, darkened circle is vehicle, open inverted triangle is 250 $\mu\text{g/ml}$, darkened inverted triangle is 100 $\mu\text{g/ml}$, open square is 10 $\mu\text{g/ml}$, darkened square is 1 $\mu\text{g/ml}$; absorbance of 2.0 is maximum of plate reader and may not be necessarily representative of cell number);

Fig. 15 E shows a typical dose response curve obtained from Hela cervical cancer cell line treated with varying levels of crude polyphenols obtained from UIT-1 cocoa genotype (absorbance (540 nm) vs. Days; open circle is control, darkened circle is vehicle, open inverted triangle is 250 $\mu\text{g/ml}$, darkened inverted triangle is 100 $\mu\text{g/ml}$, open square is 10 $\mu\text{g/ml}$; absorbance of 2.0 is maximum of plate reader and may not be necessarily representative of cell number);

Fig. 15 F shows cytotoxic effects against Hela cervical cancer cell line treated with different cocoa polyphenol fractions (absorbance (540nm) vs. Days; open circle is 100 $\mu\text{g/ml}$ fractions A-E, darkened circle is 100

$\mu\text{g/ml}$ fractions A-C, open inverted triangle is 100 $\mu\text{g/ml}$ fractions D&E; absorbance of 2.0 is maximum of plate reader and not representative of cell number);

Fig. 15 G shows cytotoxic effects at 100 $\mu\text{l/ml}$ against SKBR-3 breast cancer cell line treated with different cocoa polyphenol fractions (absorbance (540nm) vs. Days; open circle is fractions A-E, darkened circle is fractions A-C, open inverted triangle is fractions D&E);

Fig. 15 H shows typical dose-response relationships between cocoa procyanidin fraction D+E on Hela cells (absorbance (540nm) vs. Days; open circle is control, darkened circle is 100 $\mu\text{g/ml}$, open inverted triangle is 75 $\mu\text{g/ml}$, darkened inverted triangle is 50 $\mu\text{g/ml}$, open square is 25 $\mu\text{g/ml}$, darkened square is 10 $\mu\text{g/ml}$; absorbance of 2.0 is maximum of plate reader and is not representative of cell number);

Fig. 15 I shows typical dose-response relationship between cocoa procyanidin fraction D+E on SKBR-3 cells (absorbance (540nm) vs. Days; open circle is control, darkened circle is 100 $\mu\text{g/ml}$, open inverted triangle is 75 $\mu\text{g/ml}$, darkened inverted triangle is 50 $\mu\text{g/ml}$, open square is 25 $\mu\text{g/ml}$, darkened square is 10 $\mu\text{g/ml}$);

Fig. 15 J shows typical dose-response relationships between cocoa procyanidin fraction D+E on Hela cells using the Soft Agar Cloning assay (bar chart; number of colonies vs. control, 1, 10, 50, and 100 $\mu\text{g/ml}$);

Fig. 15 K shows the growth inhibition of Hela cells when treated with crude polyphenol extracts obtained from eight different cocoa genotypes (% control vs. concentration, $\mu\text{g/ml}$; open circle is C-1, darkened circle is C-2, open inverted triangle is C-3, darkened inverted triangle is C-4, open square is C-5, darkened square is C-6, open triangle is C-7, darkened triangle is C-8; C-1 = UF-12: horti race = Criollo and description is

crude extracts of UF-12 (Brazil) cocoa polyphenols
 (decaffeinated/detheobrominated); C-2 = NA-33: horti
 race = Forastero and description is crude extracts of NA-
 33 (Brazil) cocoa polyphenols (decaffeinated/
 5 detheobrominated); C-3 = EEG-48: horti race = Forastero
 and description is crude extracts of EEG-48 (Brazil)
 cocoa polyphenols (decaffeinated/detheobrominated); C-4 =
 unknown: horti race = Forastero and description is crude
 extracts of unknown (W. African) cocoa polyphenols
 10 (decaffeinated/detheobrominated); C-5 = UF-613: horti
 race = Trinitario and description is crude extracts of
 UF-613 (Brazil) cocoa polyphenols (decaffeinated/
 detheobrominated); C-6 = ICS-100: horti race = Trinitario
 and description is crude extracts of ICS-100 (Brazil)
 15 cocoa polyphenols (decaffeinated/detheobrominated); C-7 =
 ICS-139: horti race = Trinitario and description is crude
 extracts of ICS-139 (Brazil) cocoa polyphenols
 (decaffeinated/detheobrominated); C-8 = UIT-1: horti race
 = Trinitario and description is crude extracts of UIT-1
 20 (Malaysia) cocoa polyphenols (decaffeinated/
 detheobrominated));

Fig. 15 L shows the growth inhibition of Hela
 cells when treated with crude polyphenol extracts
 obtained from fermented cocoa beans and dried cocoa beans
 25 (stages throughout fermentation and sun drying; % control
 vs. concentration, $\mu\text{g/ml}$; open circle is day zero
 fraction, darkened circle is day 1 fraction, open
 inverted triangle is day 2 fraction, darkened inverted
 triangle is day 3 fraction, open square is day 4 fraction
 30 and darkened square is day 9 fraction);

Fig. 15 M shows the effect of enzymically
 oxidized cocoa procyanidins against Hela cells (dose
 response for polyphenol oxidase treated crude cocoa
 polyphenol; % control vs. concentration, $\mu\text{g/ml}$; darkened
 35 square is crude UIT-1 (with caffeine and theobromine),
 open circle crude UIT-1 (without caffeine and

theobromine) and darkened circle is crude UIT-1 (polyphenol oxidase catalyzed));

Fig. 15 N shows a representative semi-preparative reverse phase HPLC separation for combined
5 cocoa procyanidin fractions D and E;

Fig. 15 O shows a representative normal phase semi-preparative HPLC separation of a crude cocoa polyphenol extract;

Fig. 16 shows typical Rancimat Oxidation curves
10 for cocoa procyanidin extract and fractions in comparison to the synthetic antioxidants BHA and BHT (arbitrary units vs. time; dotted line and cross (+) is BHA and BHT; * is D-E; x is crude; open square is A-C; and open diamond is control);

Fig. 17 shows a typical Agarose Gel indicating inhibition of topoisomerase II catalyzed decatenation of
15 kinetoplast DNA by cocoa procyanidin fractions (Lane 1 contains 0.5 μ g of marker (M) monomer-length kinetoplast DNA circles; Lanes 2 and 20 contain kinetoplast DNA that
20 was incubated with Topoisomerase II in the presence of 4% DMSO, but in the absence of any cocoa procyanidins. (Control -C); Lanes 3 and 4 contain kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.5 and 5.0 μ g/mL cocoa procyanidin fraction A; Lanes 5
25 and 6 contain kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.5 and 5.0 μ g/mL cocoa procyanidin fraction B; Lanes 7, 8, 9, 13, 14 and 15 are replicates of kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.05, 0.5 and
30 5.0 μ g/mL cocoa procyanidin fraction D; Lanes 10, 11, 12, 16, 17 and 18 are replicates of kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.05, 0.5, and 5.0 μ g/mL cocoa procyanidin fraction E; Lane 19 is a replicate of kinetoplast DNA that was incubated with
35 Topoisomerase II in the presence of 5.0 μ g/mL cocoa procyanidin fraction E);

Fig. 18 shows dose response relationships of cocoa procyanidin fraction D against DNA repair competent and deficient cell lines (fractional survival vs. $\mu\text{g/ml}$; left side xrs-6 DNA Deficient Repair Cell Line, MM-1 D D282X1; right side BR1 Competent DNA Repair Cell Line, MM-1 D D282B1);

Fig. 19 shows the dose-response curves for Adriamycin resistant MCF-7 cells in comparison to a MCF-7 p168 parental cell line when treated with cocoa fraction D+E (% control vs. concentration, $\mu\text{g/ml}$; open circle is MCF-7 p168; darkened circle is MCF-7 ADR); and

Fig. 20 shows the dose-response effects on Hela cells when treated at 100 $\mu\text{g/mL}$ and 25 $\mu\text{g/mL}$ levels of twelve fractions prepared by Normal phase semi-preparative HPLC (bar chart, % control vs. control and fractions 1-12).

DETAILED DESCRIPTION

As discussed above, it has now been surprisingly found that cocoa extracts exhibit anti-cancer, anti-tumor or antineoplastic activity, antioxidant activity and, inhibit DNA topoisomerase II enzyme. The extracts are generally prepared by reducing cocoa beans to a powder, defatting the powder, and extracting the active compound(s) from the defatted powder. The powder can be prepared by freeze-drying the cocoa beans and pulp, depulping the cocoa beans and pulp, dehulling the freeze-dried cocoa beans, and grinding the dehulled beans. The extraction of active compound(s) can be by solvent extraction techniques. The extracts can be purified; for instance, by gel permeation chromatography or by preparative High Performance Liquid Chromatography (HPLC) techniques or by a combination of such techniques. The extracts having activity, without wishing to necessarily be bound by any particular theory, have been identified as cocoa polyphenol(s) such as procyanidins. These cocoa procyanidins have significant anti-cancer,

anti-tumor or antineoplastic activity; antioxidant activity; and inhibit DNA topoisomerase II enzyme.

Anti-cancer, anti-tumor or antineoplastic or, antioxidant or DNA topoisomerase II enzyme inhibiting
5 compositions containing the inventive cocoa polyphenols or procyanidins can be prepared in accordance with standard techniques well known to those skilled in the pharmaceutical art. Such compositions can be administered to a patient in need of such administration
10 in dosages and by techniques well known to those skilled in the medical arts taking into consideration such factors as the age, sex, weight, and condition of the particular patient, and the route of administration. The compositions can be co-administered or sequentially
15 administered with other antineoplastic, anti-tumor or anti-cancer agents or antioxidant or DNA topoisomerase II enzyme inhibiting agents and/or with agents which reduce or alleviate ill effects of antineoplastic, anti-tumor or anti-cancer agents or antioxidant or DNA topoisomerase II
20 enzyme inhibiting agents; again, taking into consideration such factors as the age, sex, weight, and condition of the particular patient, and, the route of administration.

Examples of compositions of the invention
25 include solid compositions for oral administration such as capsules, tablets, pills and the like, as well as chewable solid formulations, to which the present invention may be well-suited since it is from an edible source (e.g., cocoa or chocolate flavored solid
30 compositions); liquid preparations for orifice, e.g., oral, nasal, anal, vaginal etc., administration such as suspensions, syrups or elixirs; and, preparations for parental, subcutaneous, intradermal, intramuscular or intravenous administration (e.g., injectable
35 administration) such as sterile suspensions or emulsions. However, the active ingredient in the compositions may complex with proteins such that when administered into

the bloodstream, clotting may occur due to precipitation of blood proteins; and, the skilled artisan should take this into account. In such compositions the active cocoa extract may be in admixture with a suitable carrier, diluent, or excipient such as sterile water, physiological saline, glucose or the like. The active cocoa extract of the invention can be provided in lyophilized form for reconstituting, for instance, in isotonic aqueous, saline buffer.

Further, the invention also comprehends a kit wherein the active cocoa extract is provided. The kit can include a separate container containing a suitable carrier, diluent or excipient. The kit can also include an additional anti-cancer, anti-tumor or antineoplastic agent or antioxidant or DNA topoisomerase II enzyme inhibiting agent and/or an agent which reduces or alleviates ill effects of antineoplastic, anti-tumor or anti-cancer agents or antioxidant or DNA topoisomerase II enzyme inhibiting agents for co- or sequential-administration. The additional agent(s) can be provided in separate container(s) or in admixture with the active cocoa extract. Additionally, the kit can include instructions for mixing or combining ingredients and/or administration.

Furthermore, while the invention is described with respect to cocoa extracts preferably comprising cocoa procyanidins, from this disclosure the skilled organic chemist will appreciate and envision synthetic routes to obtain the active compounds. Accordingly, the invention comprehends synthetic cocoa polyphenols or procyanidins or their derivatives which include, but are not limited to glycosides, gallates, esters, etc. and the like.

The following non-limiting Examples are given by way of illustration only and are not to be considered a limitation of this invention, many apparent variations

of which are possible without departing from the spirit or scope thereof.

EXAMPLES

Example 1: Cocoa Source and Method of Preparation

5 Several *Theobroma cacao* genotypes which
represent the three recognized horticultural races of
cocoa (Enriquez, 1967; Engels, 1981) were obtained from
the three major cocoa producing origins of the world. A
list of those genotypes used in this study are shown in
10 Table 1. Harvested cocoa pods were opened and the beans
with pulp were removed for freeze drying. The pulp was
manually removed from the freeze dried mass and the beans
were subjected to analysis as follows. The unfermented,
freeze dried cocoa beans were first manually dehulled,
15 and ground to a fine powdery mass with a TEKMAR Mill.TM
The resultant mass was then defatted overnight by Soxhlet
extraction using redistilled hexane as the solvent.
Residual solvent was removed from the defatted mass by
vacuum at ambient temperature.

Table 1: Description of *Theobroma cacao* Source Material

GENOTYPE	ORIGIN	HORTICULTURAL RACE
UIT-1	Malaysia	Trinitario
Unknown	West Africa	Forastero
ICS-100	Brazil	Trinitario
ICS-39	Brazil	Trinitario
UF-613	Brazil	Trinitario
EEG-48	Brazil	Forastero
UF-12	Brazil	Criollo
NA-33	Brazil	Forastero

Example 2: Procyanidin Extraction Procedures**A. Method 1**

Procyanidins were extracted from the defatted, unfermented, freeze dried cocoa beans of Example 1 using a modification of the method described by Jalal and Collin (1977). Procyanidins were extracted from 50 gram batches of the defatted cocoa mass with 2X 400 mL 70% acetone/deionized water followed by 400mL 70% methanol/deionized water. The extracts were pooled and the solvents removed by evaporation at 45°C with a rotary evaporator held under partial vacuum. The resultant aqueous phase was diluted to 1L with deionized water and extracted 2X with 400mL CHCl₃. The solvent phase was discarded. The aqueous phase was then extracted 4X with 500mL ethyl acetate. Any resultant emulsions were broken by centrifugation on a Sorvall RC 28S centrifuge operated at 2,000 xg for 30 min. at 10°C. To the combined ethyl acetate extracts, 100-200mL deionized water was added. The solvent was removed by evaporation at 45°C with a rotary evaporator held under partial vacuum. The resultant aqueous phase was frozen in liquid N₂ followed by freeze drying on a LABCONCO Freeze Dry System. The yields of crude procyanidins that were obtained from the different cocoa genotypes are listed in Table 2.

Table 2: Crude Procyanidin Yields

GENOTYPE	ORIGIN	YIELDS (g)
UIT-1	Malaysia	3.81
Unknown	West Africa	2.55
ICS-100	Brazil	3.42
ICS-39	Brazil	3.45
UF-613	Brazil	2.98
EEG-48	Brazil	3.15
UF-12	Brazil	1.21
NA-33	Brazil	2.23

B. Method 2

Alternatively, procyanidins are extracted from defatted, unfermented, freeze dried cocoa beans of Example 1 with 70% aqueous acetone. Ten grams of defatted material was slurried with 100 mL solvent for 5-10 min. The slurry was centrifuged for 15 min. at 4°C at 3000 xg and the supernatant passed through glass wool. The filtrate was subjected to distillation under partial vacuum and the resultant aqueous phase frozen in liquid N₂, followed by freeze drying on a LABCONCO Freeze Dry System. The yields of crude procyanidins ranged from 15-20%.

Without wishing to be bound by any particular theory, it is believed that the differences in crude yields reflected variations encountered with different genotypes, geographical origin, horticultural race, and method of preparation.

Example 3: Partial Purification of Cocoa Procyanidins**30 A. Gel Permeation Chromatography**

Procyanidins obtained from Example 2 were partially purified by liquid chromatography on SephadexTM LH-20 (28 x 2.5 cm). Separations were aided by a step gradient from deionized water into methanol. The initial gradient composition started with 15% methanol in deionized water which was followed step wise every 30

min. with 25% methanol in deionized water, 35% methanol in deionized water, 70% methanol in deionized water, and finally 100% methanol. The effluent following the elution of the xanthine alkaloids (caffeine and theobromine) was collected as a single fraction. The fraction yielded a xanthine alkaloid free subfraction which was submitted to further subfractionation to yield five subfractions designated MM2A through MM2E. The solvent was removed from each subfraction by evaporation at 45°C with a rotary evaporator held under partial vacuum. The resultant aqueous phase was frozen in liquid N₂ and freeze dried overnight on a LABCONCO Freeze Dry System. A representative gel permeation chromatogram showing the fractionation is shown in Figure 1. Approximately, 100mg of material was subfractionated in this manner.

Figure 1: Gel Permeation Chromatogram of Crude Procyanidins on Sephadex LH-20

Chromatographic Conditions: Column; 28 x 2.5 cm Sephadex LH-20, Mobile Phase: Methanol/Water Step Gradient, 15:85, 25:75, 35:65, 70:30, 100:0 Stepped at 1/2 Hour Intervals, Flow Rate; 1.5ml/min, Detector; UV @ $\lambda_1=254$ nm and $\lambda_2=365$ nm, Chart Speed: 0.5 mm/min, Column Load; 120 mg.

B. Semi-preparative High Performance Liquid Chromatography (HPLC)

Method 1: Reverse Phase Separation

Procyanidins obtained from Example 2 and/or 3A were partially purified by semi-preparative HPLC. A Hewlett PackardTM 1050 HPLC System equipped with a variable wavelength detector, RheodyneTM 7010 injection valve with 1 mL injection loop was assembled with a PharmaciaTM FRAC-100 Fraction Collector. Separations were effected on a Phenomenex UltracarbTM 10 μ ODS column (250 x 22.5 mm) connected with a Phenomenex 10 μ ODS Ultracarb (60 x 10 mm) guard column. The mobile phase composition was A =

water; B = methanol used under the following linear gradient conditions: [Time, %A]; (0,85), (60,50), (90,0), and (110,0) at a flow rate of 5 mL/min.

A representative Semi-preparative HPLC trace is shown in Figure 15N for the separation of procyanidins present in fraction D + E. Individual peaks or select chromatographic regions were collected on timed intervals or manually by fraction collection for further purification and subsequent evaluation. Injection loads ranged from 25-100 mg of material.

Method 2. Normal Phase Separation

Procyanidin extracts obtained from Examples 2 and/or 3A were partially purified by semi-preparative HPLC. A Hewlett Packard 1050 HPLC system, MilliporeTM Waters Model 480 LC detector set at 254 nm was assembled with a Pharmacia Frac-100 Fraction Collector set in peak mode. Separations were effected on a SupelcoTM 5 μ Supelcosil LC-Si column (250 x 10mm) connected with a Supelco 5 μ SupelguardTM LC-Si guard column (20 x 4.6mm). Procyanidins were eluted by a linear gradient under the following conditions: (Time, %A, %B); (0,82,14), (30, 67.6, 28.4), (60, 46, 50), (65, 10, 86), (70, 10, 86) followed by a 10 min. re-equilibration. Mobile phase composition was A=dichloromethane; B=methanol; and C=acetic acid: water (1:1). A flow rate of 3 mL/min was used. Components were detected by UV at 254 nm, and recorded on a Kipp & ZonanTM BD41 recorder. Injection volumes ranged from 100-250 μ L of 10mg of procyanidin extracts dissolved in 0.25 mL 70% aqueous acetone. A representative semi-preparative HPLC trace is shown in Figure 15 O. Individual peaks or select chromatographic regions were collected on timed intervals or manually by fraction collection for further purification and subsequent evaluation.

HPLC Conditions: 250 x 10mm Supelco Supelcosil LC-Si
(5 μ m) Semipreparative Column
20 x 4.6mm Supelco Supelcosil LC-Si
(5 μ m) Guard Column

Detector: Waters LC
Spectrophotometer Model
480 @ 254nm

Flow rate: 3mL/min,
Column Temperature: ambient,
Injection: 250 μ L of 70% aqueous
acetone extract.

Gradient: Time (min)	CH ₂ Cl ₂	Methanol	Acetic Acid/H ₂ O (1:1)
0	82	14	4
30	67.6	28.4	4
60	46	50	4
65	10	86	4
70	10	86	4

The fractions obtained were as follows:

FRACTION	TYPE
1	dimers
2	trimers
3	tetramers
4	pentamers
5	hexamers
6	heptamers
7	octamers
8	nonamers
9	decamers
10	undecamers
11	dodecamers
12	higher oligomers

Example 4: Analytical HPLC Analysis of Procyanidin Extracts

Method 1: Reverse Phase Separation

5 Procyanidin extracts obtained from Example 3 were filtered through a 0.45μ filter and analyzed by a Hewlett Packard 1090 ternary HPLC system equipped with a Diode Array detector and a HP model 1046A Programmable Fluorescence Detector. Separations were effected at 45°C
10 on a Hewlett-Packard 5μ HypersilTM ODS column (200 x 2.1mm). The flavanols and procyanidins were eluted with a linear gradient of 60% B into A followed by a column wash with B at a flow rate of 0.3mL/min. The mobile phase composition was B= 0.5% acetic acid in methanol and
15 A= 0.5% acetic acid in deionized water. Acetic acid levels in A and B mobile phases can be increased to 2%. Components were detected by fluorescence, where $\lambda_{\text{ex}} = 276\text{nm}$ and $\lambda_{\text{em}} = 316\text{nm}$. Concentrations of (+)-catechin and (-)-epicatechin were determined relative to reference
20 standard solutions. Procyanidin levels were estimated by using the response factor for (-)-epicatechin. A representative HPLC chromatogram showing the separation of the various components is shown in Figure 2A for one cocoa genotype. Similar HPLC profiles were obtained from
25 the other cocoa genotypes.

21

HPLC Conditions: Column: 200 x 2.1mm Hewlett Packard
Hypersil ODS (5 μ)
Guard column: 20 x 2.1mm Hewlett
Packard Hypersil ODS (5 μ)
Detectors: Diode Array @ 280nm
Fluorescence λ_{ex} = 276nm;
 λ_{em} = 316nm.
Flow rate: 0.3mL/min.
Column Temperature: 45°C

Gradient: Time (min)	0.5% Acetic Acid in deionized water	0.5% Acetic acid in methanol
0	100	0
50	40	60
60	0	100

Method 2: Normal Phase Separation

Procyanidin extracts obtained from Examples 2 and/or 3 were filtered through a 0.45 μ filter and analyzed by a Hewlett Packard 1090 Series II HPLC system equipped with a HP model 1046A Programmable Fluorescence detector and Diode Array detector. Separations were effected at 37°C on a 5 μ Phenomenex LichrosphereTM Silica 100 column (250 x 3.2mm) connected to a Supelco Supelguard LC-Si 5 μ guard column (20 x 4.6mm). Procyanidins were eluted by linear gradient under the following conditions: (Time, %A, %B); (0, 82, 14), (30, 67.6, 28.4), (60, 46, 50), (65, 10, 86), (70, 10, 86) followed by an 8 min. re-equilibration. Mobile phase composition was A=dichloromethane, B=methanol, and C=acetic acid: water at a volume ratio of 1:1. A flow rate of 0.5 mL/min. was used. Components were detected by fluorescence, where λ_{ex} = 276nm and λ_{em} = 316nm or by UV at 280 nm. A representative HPLC chromatogram showing the separation of the various procyanidins is shown in Figure 2B for one genotype. Similar HPLC profiles were obtained from other cocoa genotypes.

HPLC Conditions:

250 x 3.2mm Phenomenex Lichrosphere Silica 100
 column (5 μ) 20 x 4.6mm Supelco Supelguard LC-Si
 (5 μ) guard column
 Detectors: Photodiode Array @ 280nm
 Fluorescence λ_{ex} = 276nm;
 λ_{em} = 316nm.
 Flow rate: 0.5 mL/min.
 Column Temperature: 37°C

Gradient: Time (min.)	CH ₂ -Cl ₂	Methanol	Acetic Acid/Water (1:1)
0	82	14	4
30	67.6	28.4	4
60	46	50	4
65	10	86	4
70	10	86	4

Example 5: Identification of Procyanidins

Procyanidins were purified by liquid chromatography on Sephadex LH-20 (28 x 2.5cm) columns followed by semi-preparative HPLC using a 10 μ μ BondapakTM C18 (100 x 8mm) column or by semi-preparative HPLC using a 5 μ Supelcosil LC-Si (250 x 10mm) column.

Partially purified isolates were analyzed by Fast Atom Bombardment - Mass Spectrometry (FAB-MS) on a VG ZAB-TTM high resolution MS system using a Liquid Secondary Ion Mass Spectrometry (LSIMS) technique in positive and negative ion modes. A cesium ion gun was used as the ionizing source at 30kV and a "Magic Bullet Matrix" (1:1 dithiothreitol/dithioerythritol) was used as the proton donor.

Analytical investigations of these fractions by LSIMS revealed the presence of a number of flavan-3-ol oligomers as shown in Table 3.

Table 3: LSIMS (Positive Ion) Data from Cocoa Procyanidin Fractions

Oligomer	(M + 1) ⁺ m/z	(M + Na) ⁺ m/z	Mol. Wt.
Monomers (catechins)	291	313	290
Dimer(s)	577/579	599/601	576/578
Trimer(s)	865/867	887/889	884/866
Tetramer(s)	1155	1177	1154
Pentamer(s)	1443	1465	1442
Hexamer(s)	1731	1753	1730
Heptamer(s)	---	2041	2018
Octamer(s)	---	2329	2306
Nonamer(s)	---	2617	2594
Decamer(s)	---	2905	2882
Undecamer(s)	---	---	3170
Dodecamer(s)	---	---	3458

The major mass fragment ions were consistent with work previously reported for both positive and negative ion FAB-MS analysis of procyanidins (Self et al., 1986 and Porter et al., 1991). The ion corresponding to m/z 577 ($M+H$)⁺ and its sodium adduct at m/z 599 ($M+Na$)⁺ suggested the presence of doubly linked procyanidin dimers in the isolates. It was interesting to note that the higher oligomers were more likely to form sodium adducts ($M+Na$)⁺ than their protonated molecular ions ($M+H$)⁺. The procyanidin isomers B-2, B-5 and C-1 were tentatively identified based on the work reported by Revilla et al. (1991), Self et al. (1986) and Porter et al. (1991). Procyanidins up to both the octamer and decamer were verified by FAB-MS in the partially purified fractions. Additionally, evidence for procyanidins up to the dodecamer were observed from normal phase HPLC analysis (see Figure 2B). Without wishing to be bound by any particular theory, it is

believed that the dodecamer is the limit of solubility in the solvents used in the extraction and purification schemes. Table 4 lists the relative concentrations of the procyanidins found in xanthine alkaloid free isolates based on reverse phase HPLC analysis. Table 5 lists the relative concentrations of the procyanidins based on normal phase HPLC analysis.

Table 4: Relative Concentrations of Procyanidins in the Xanthine Alkaloid Free Isolates

Component	Molecular Weight	Amount
(+)-catechin	290	1.6%
(-)-epicatechin	290	38.2%
B-2 Dimer	578	11.0%
B-5 Dimer	578	5.3%
C-1 Trimer	866	9.3%
Doubly linked dimers	576	3.0%
Tetramer(s)	1154	4.5%
Pentamer-Octamer	1442-2306	24.5%
Unknowns and higher oligomers	---	2.6%

Table 5: Relative Concentrations of Procyanidins in Aqueous Acetone Extracts

5	Component	Molecular Weight	Amount
	(+)-catechin and (-) -epicatechin	290 (same for each)	41.9%
	B-2 and B-5 Dimers	578	13.9%
	Trimers	884/866	11.3%
10	Tetramers	1154	9.9%
	Pentamers	1442	7.8%
	Hexamers	1730	5.1%
	Heptamers	2018	4.2%
	Octamers	2306	2.8%
15	Nonamers	2594	1.6%
	Decamers	2882	0.7%
	Undecamers	3170	0.2%
	Dodecamers	3458	<0.1%

20 Figure 3 shows several procyanidin structures
and Figures 4A-4E show the representative HPLC
chromatograms of the five fractions employed in the
following screening for anti-cancer or antineoplastic
activity. The HPLC conditions for Figs. 4A-4E were as
25 follows:

HPLC Conditions: Hewlett Packard 1090 ternary
HPLC System equipped with HP Model 1046A
Programmable Fluorescence Detector.

30 Column: Hewlett Packard 5 μ Hypersil ODS (200 x
2.1 mm) Linear Gradient of 60% B into A at a
flow rate of 0.3 ml/min. B = 0.5% acetic acid
in methanol; A = 0.5% acetic acid in deionized
water. λ_{ex} = 280nm; λ_{em} = 316nm.

Figure 15 O shows a representative semi-prep
35 HPLC chromatogram of an additional 12 fractions employed
in the screening for anticancer or antineoplastic
activity (HPLC conditions stated above).

**Example 6: Anti-Cancer, Anti-Tumor or Antineoplastic
 Activity of Cocoa Extracts (Procyanidins)**

 The MTT (3-[4,5-dimethyl thiazol-2yl]-2,5-
5 diphenyltetrazolium bromide) - microtiter plate
 tetrazolium cytotoxicity assay originally developed by
 Mosmann (1983) was used to screen test samples from
 Example 5. Test samples, standards (cisplatin and
 chlorambucil) and MTT reagent were dissolved in 100% DMSO
10 (dimethyl sulfoxide) at a 10 mg/mL concentration. Serial
 dilutions were prepared from the stock solutions. In the
 case of the test samples, dilutions ranging from 0.01
 through 100 µg/mL were prepared in 0.5% DMSO.

 All human tumor cell lines were obtained from
15 the American Type Culture Collection. Cells were grown
 as mono layers in alpha-MEM containing 10% fetal bovine
 serum, 100 units/mL penicillin, 100 µg/mL streptomycin
 and 240 units/mL nystatin. The cells were maintained in
 a humidified, 5% CO₂ atmosphere at 37°C.

20 After trypsinization, the cells are counted and
 adjusted to a concentration of 50×10^5 cells/mL (varied
 according to cancer cell line). 200µL of the cell
 suspension was plated into wells of 4 rows of a 96-well
 microtiter plate. After the cells were allowed to attach
25 for four hours, 2µL of DMSO containing test sample
 solutions were added to quadruplicate wells. Initial
 dose-response finding experiments, using order of
 magnitude test sample dilutions were used to determine
 the range of doses to be examined. Well absorbencies at
30 540nm were then measured on a BIO RAD MP450TM plate reader.
 The mean absorbance of quadruplicate test sample treated
 wells was compared to the control, and the results
 expressed as the percentage of control absorbance
 plus/minus the standard deviation. The reduction of MTT
35 to a purple formazan product correlates in a linear
 manner with the number of living cells in the well.
 Thus, by measuring the absorbance of the reduction
 product, a quantitation of the percent of cell survival

at a given dose of test sample can be obtained. Control wells contained a final concentration of 1% DMSO.

Two of the samples were first tested by this protocol. Sample MM1 represented a very crude isolate of
5 cocoa procyanidins and contained appreciable quantities of caffeine and theobromine. Sample MM2 represented a cocoa procyanidin isolate partially purified by gel permeation chromatography. Caffeine and theobromine were absent in MM2. Both samples were screened for activity
10 against the following cancer cell lines using the procedures previously described:

HCT 116 colon cancer
ACHN renal adenocarcinoma
SK-5 melanoma
15 A498 renal adenocarcinoma
MCF-7 breast cancer
PC-3 prostate cancer
CAPAN-2 pancreatic cancer

Little or no activity was observed with MM1 on
20 any of the cancer cell lines investigated. MM2 was found to have activity against HCT-116, PC-3 and ACHN cancer cell lines. However, both MM1 and MM2 were found to interfere with MTT such that it obscured the decrease in absorbance that would have reflected a decrease in viable
25 cell number. This interference also contributed to large error bars, because the chemical reaction appeared to go more quickly in the wells along the perimeter of the plate. A typical example of these effects is shown in Figure 5. At the high concentrations of test material,
30 one would have expected to observe a large decrease in survivors rather than the high survivor levels shown. Nevertheless, microscopic examinations revealed that cytotoxic effects occurred, despite the MTT interference effects. For instance, an IC_{50} value of $0.5 \mu\text{g/mL}$ for the
35 effect of MM2 on the ACHN cell line was obtained in this manner.

These preliminary results, in the inventors' view, required amendment of the assay procedures to preclude the interference with MTT. This was accomplished as follows. After incubation of the plates at 37°C in a humidified, 5% CO₂ atmosphere for 18 hours, the medium was carefully aspirated and replaced with fresh alpha-MEM media. This media was again aspirated from the wells on the third day of the assay and replaced with 100µL of freshly prepared McCoy's medium. 11µL of a 5mg/mL stock solution of MTT in PBS (Phosphate Buffered Saline) were then added to the wells of each plate. After incubation for 4 hours in a humidified, 5% CO₂ atmosphere at 37°C, 100µL of 0.04 N HCl in isopropanol was added to all wells of the plate, followed by thorough mixing to solubilize the formazan produced by any viable cells. Additionally, it was decided to subfractionate the procyanidins to determine the specific components responsible for activity.

The subfractionation procedures previously described were used to prepare samples for further screening. Five fractions representing the areas shown in Figure 1 and component(s) distribution shown in Figures 4A - 4E were prepared. The samples were coded MM2A through MM2E to reflect these analytical characterizations and to designate the absence of caffeine and theobromine.

Each fraction was individually screened against the HCT-116, PC-3 and ACHN cancer cell lines. The results indicated that the activity did not concentrate to any one specific fraction. This type of result was not considered unusual, since the components in "active" natural product isolates can behave synergistically. In the case of the cocoa procyanidin isolate (MM2), over twenty detectable components comprised the isolate. It was considered possible that the activity was related to a combination of components present in the different

fractions, rather than the activity being related to an individual component(s).

On the basis of these results, it was decided to combine the fractions and repeat the assays against the same cancer cell lines. Several fraction combinations produced cytotoxic effects against the PC-3 cancer cell lines. Specifically, IC_{50} values of 40 $\mu\text{g/mL}$ each for MM2A and MM2E combination, and of 20 $\mu\text{g/mL}$ each for MM2C and MM2E combination, were obtained. Activity was also reported against the HCT-116 and ACHN cell lines, but as before, interference with the MTT indicator precluded precise observations. Replicate experiments were repeatedly performed on the HCT-116 and ACHN lines to improve the data. However, these results were inconclusive due to bacterial contamination and exhaustion of the test sample material. Figures 6A-6D show the dose-response relationship between combinations of the cocoa extracts and PC-3 cancer cells.

Nonetheless, from this data, it is clear that cocoa extracts, especially cocoa polyphenols or procyanidins, have significant anti-tumor; anti-cancer or antineoplastic activity, especially with respect to human PC-3 (prostate), HCT-116 (colon) and ACHN (renal) cancer cell lines. In addition, those results suggest that specific procyanidin fractions may be responsible for the activity against the PC-3 cell line.

Example 7: Anti-Cancer, Anti-Tumor or Antineoplastic Activity of Cocoa Extracts (Procyanidins)

To confirm the above findings and further study fraction combinations, another comprehensive screening was performed.

All prepared materials and procedures were identical to those reported above, except that the standard 4-replicates per test dose was increased to 8 or 12-replicates per test dose. For this study, individual and combinations of five cocoa procyanidin fractions were screened against the following cancer cell lines.

PC-3 Prostate
KB Nasopharyngeal/HeLa
HCT-116 Colon
ACHN Renal
5 MCF-7 Breast
SK-5 Melanoma
A-549 Lung
CCRF-CEM T-cell leukemia

Individual screenings consisted of assaying
10 different dose levels (0.01-100 μ g/mL) of fractions A, B,
C, D, and E (See Figs. 4A-4E and discussion thereof,
supra) against each cell line. Combination screenings
consisted of combining equal dose levels of fractions
A+B, A+C, A+D, A+E, B+C, B+D, B+E, C+D, C+E, and D+E
15 against each cell line. The results from these assays
are individually discussed, followed by an overall
summary.

A. PC-3 Prostate Cell Line

Figures 7A - 7H show the typical dose response
20 relationship between cocoa procyanidin fractions and the
PC-3 cell line. Figures 7D and 7E demonstrate that
fractions D and E were active at an IC₅₀ value of 75
 μ g/mL. The IC₅₀ values that were obtained from dose-
response curves of the other procyanidin fraction
25 combinations ranged between 60 - 80 μ g/mL when fractions
D or E were present. The individual IC₅₀ values are
listed in Table 6.

B. KB Nasopharyngeal/HeLa Cell Line

Figures 8A - 8H show the typical dose response
30 relationship between cocoa procyanidin fractions and the
KB Nasopharyngeal/HeLa cell line. Figures 8D and 8E
demonstrate that fractions D and E were active at an IC₅₀
value of 75 μ g/mL. Figures 8F - 8H depict representative
results obtained from the fraction combination study. In
35 this case, procyanidin fraction combination A+B had no
effect, whereas fraction combinations B+E and D+E were
active at an IC₅₀ value of 60 μ g/mL. The IC₅₀ values that

were obtained from other dose response curves from other fraction combinations ranged from 60 - 80 $\mu\text{g/mL}$ when fractions D or E were present. The individual IC_{50} values are listed in Table 6. These results were essentially
5 the same as those obtained against the PC-3 cell line.

C. HCT-116 Colon Cell Line

Figure 9A - 9H show the typical dose response relationships between cocoa procyanidin fractions and the HCT-116 colon cell line. Figures 9D and 9E demonstrate
10 that fraction E was active at an IC_{50} value of approximately 400 $\mu\text{g/mL}$. This value was obtained by extrapolation of the existing curve. Note that the slope of the dose response curve for fraction D also indicated activity. However, no IC_{50} value was determined from this
15 plot, since the slope of the curve was too shallow to obtain a reliable value. Figures 9F - 9H depict representative results obtained from the fraction combination study. In this case, procyanidin fraction combination B+D did not show appreciable activity,
20 whereas fraction combinations A+E and D+E were active at IC_{50} values of 500 $\mu\text{g/mL}$ and 85 $\mu\text{g/mL}$, respectively. The IC_{50} values that were obtained from dose response curves of other fraction combinations averaged about 250 $\mu\text{g/mL}$ when fraction E was present. The extrapolated IC_{50} values
25 are listed in Table 6.

D. ACHN Renal Cell Line

Figure 10A - 10H show the typical dose response relationships between cocoa procyanidin fractions and the ACHN renal cell line. Figures 10A - 10E indicated that
30 no individual fraction was active against this cell line. Figures 10F - 10H depict representative results obtained from the fraction combination study. In this case, procyanidin fraction combination B+C was inactive, whereas the fraction combination A+E resulted in an
35 extrapolated IC_{50} value of approximately 500 $\mu\text{g/mL}$. Dose response curves similar to the C+D combination were considered inactive, since their slopes were too shallow.

Extrapolated IC_{50} values for other fraction combinations are listed in Table 6.

E. A-549 Lung Cell Line

Figures 11A - 11H show the typical dose response relationships between cocoa procyanidin fractions and the A-549 lung cell line. No activity could be detected from any individual fraction or combination of fractions at the doses used in the assay. However, procyanidin fractions may nonetheless have utility with respect to this cell line.

F. SK-5 Melanoma Cell Line

Figure 12A - 12H show the typical dose response relationships between cocoa procyanidin fractions and the SK-5 melanoma cell line. No activity could be detected from any individual fraction or combination of fractions at the doses used in the assay. However, procyanidin fractions may nonetheless have utility with respect to this cell line.

G. MCF-7 Breast Cell Line

Figures 13A - 13H show the typical dose response relationships between cocoa procyanidin fractions and the MCF-7 breast cell line. No activity could be detected from any individual fraction or combination of fractions at the doses used in the assay. However, procyanidin fractions may nonetheless have utility with respect to this cell line.

H. CCRF-CEM T-Cell Leukemia Line

Atypical dose response curves were originally obtained against the CCRF-CEM T-cell leukemia line. However, microscopic counts of cell number versus time at different fraction concentrations indicated that 500 μ g of fractions A, B and D effected an 80% growth reduction over a four day period. A representative dose response relationship is shown in Figure 14.

I. Summary

The IC_{50} values obtained from these assays are collectively listed in Table 6 for all the cell lines

except for CCRF-CEM T-cell leukemia. The T-cell leukemia data was intentionally omitted from the Table, since a different assay procedure was used. A general summary of these results indicated that the most activity was
5 associated with fractions D and E. These fractions were most active against the PC-3 (prostate) and KB (nasopharyngeal/HeLa) cell lines. These fractions also evidenced activity against the HCT-116 (colon) and ACHN (renal) cell lines, albeit but only at much higher doses.
10 No activity was detected against the MCF-7 (breast), SK-5 (melanoma) and A-549 (lung) cell lines. However, procyanidin fractions may nonetheless have utility with respect to these cell lines. Activity was also shown against the CCRF-CEM (T-cell leukemia) cell line. It
15 should also be noted that fractions D and E are the most complex compositionally. Nonetheless, from this data it is clear that cocoa extracts, especially cocoa procyanidins, have significant anti-tumor, anti-cancer or antineoplastic activity.

Table 6: IC₅₀ Values for Cocoa Procyanidin Fractions Against Various Cell Lines

(IC₅₀ values in $\mu\text{g/mL}$)

5		FRACTION	PC-3	KB	HCT-116	ACHN	MCF-7	SK-5	A-549
		A							
		B							
		C							
10		D	90	80					
		E	75	75	400				
		A+B							
		A+C	125	100					
		A+D	75	75					
15		A+E	80	75	500	500			
		B+C							
		B+D	75	80					
		B+E	60	65	200				
		C+D	80	75		1000			
20		C+E	80	70	250				
		D+E	80	60	85				

Values above 100 $\mu\text{g/mL}$ were extrapolated from dose response curves

25 Example 8. Anti-Cancer, Anti-Tumor or Antineoplastic Activity of Cocoa Extracts (Procyanidins)

Several additional in vitro assay procedures were used to complement and extend the results presented in Examples 6 and 7.

Method A. Crystal Violet Staining Assay

All human tumor cell lines were obtained from the American Type Culture Collection. Cells were grown as monolayers in IMEM containing 10% fetal bovine serum without antibiotics. The cells were maintained in a humidified, 5% CO₂ atmosphere at 37°C.

After trypsinization, the cells were counted and adjusted to a concentration of 1,000-2,000 cells per

100 μ L. Cell proliferation was determined by plating the cells (1,000-2,000 cells/well) in a 96 well microtiter plate. After addition of 100 μ L cells per well, the cells were allowed to attach for 24 hours. At the end of
5 the 24 hour period, various cocoa fractions were added at different concentrations to obtain dose response results. The cocoa fractions were dissolved in media at a 2 fold concentration and 100 μ L of each solution was added in triplicate wells. On consecutive days, the plates were
10 stained with 50 μ L crystal violet (2.5g crystal violet dissolved in 125 mL methanol, 375 mL water), for 15 min. The stain was removed and the plate was gently immersed into cold water to remove excess stain. The washings were repeated two more times, and the plates allowed to
15 dry. The remaining stain was solubilized by adding 100 μ L of 0.1 M sodium citrate/50% ethanol to each well. After solubilization, the number of cells were quantitated on an ELISA plate reader at 540nm (reference filter at 410nm). The results from the ELISA reader were
20 graphed with absorbance on the y-axis and days growth on the x-axis.

Method B. Soft Agar Cloning Assay

Cells were cloned in soft agar according to the method described by Nawata et al. (1981). Single cell
25 suspensions were made in media containing 0.8% agar with various concentrations of cocoa fractions. The suspensions were aliquoted into 35mm dishes coated with media containing 1.0% agar. After 10 days incubation, the number of colonies greater than 60 μ m in diameter
30 were determined on an Ominicron 3600 Image Analysis System. The results were plotted with number of colonies on the y-axis and the concentrations of a cocoa fraction on the x-axis.

Method C. XTT-Microculture Tetrazolium Assay

35 The XTT assay procedure described by Scudiero et al. (1988) was used to screen various cocoa fractions. The XTT assay was essentially the same as that described

using the MTT procedure (Example 6) except for the following modifications. XTT ((2,3-bis(2-methoxy-4-nitro-5-sulfophenyl)-5-[(phenylamino)carbonyl]-2H-tetrazolium hydroxide) was prepared at 1mg/mL medium
5 without serum, prewarmed to 37°C. PMS was prepared at 5mM PBS. XTT and PMS were mixed together; 10 μ L of PMS per mL XTT and 50 μ L PMS-XTT were added to each well. After an incubation at 37°C for 4 hr, the plates were mixed 30 min. on a mechanical shaker and the absorbance
10 measured at 450-600nm. The results were plotted with the absorbance on the y-axis and days growth or concentration on the x-axis.

For methods A and C, the results were also plotted as the percent control as the y-axis and days
15 growth or concentration on the x-axis.

A comparison of the XTT and Crystal Violet Assay procedures was made with cocoa fraction D & E (Example 3B) against the breast cancer cell line MCF-7 p168 to determine which assay was most sensitive. As
20 shown in Figure 15A, both assays showed the same dose-response effects for concentrations >75 μ g/mL. At concentrations below this value, the crystal violet assay showed higher standard deviations than the XTT assay results. However, since the crystal violet assay was
25 easier to use, all subsequent assays, unless otherwise specified, were performed by this procedure.

Crystal violet assay results are presented (Figures 15B-15E) to demonstrate the effect of a crude polyphenol extract (Example 2) on the breast cancer cell
30 line MDA MB231, prostate cancer cell line PC-3, breast cancer cell line MCF-7 p163, and cervical cancer cell line Hela, respectively. In all cases a dose of 250 μ g/mL completely inhibited all cancer cell growth over a period of 5-7 days. The Hela cell line appeared to be more
35 sensitive to the extract, since a 100 μ g/mL dose also inhibited growth. Cocoa fractions from Example 3B were also assayed against Hela and another breast cancer cell

line SKBR-3. The results (Figures 15F and 15G) showed that fraction D & E has the highest activity. As shown in Figures 15H and 15I, IC_{50} values of about $40\mu\text{g/mL}$ D & E were obtained from both cancer cell lines.

5 The cocoa fraction D & E was also tested in the soft agar cloning assay which determines the ability of a test compound(s) to inhibit anchorage independent growth. As shown in Figure 15J, a concentration of $100\mu\text{g/mL}$ completely inhibited colony formation of Hela cells.

10 Crude polyphenol extracts obtained from eight different cocoa genotypes representing the three horticultural races of cocoa were also assayed against the Hela cell line. As shown in Figure 15K all cocoa varieties showed similar dose-response effects. The
15 UIT-1 variety exhibited the most activity against the Hela cell line. These results demonstrated that all cocoa genotypes possess a polyphenol fraction that elicits activity against at least one human cancer cell line that is independent of geographical origin,
20 horticultural race, and genotype.

 Another series of assays were performed on crude polyphenol extracts prepared on a daily basis from a one ton scale traditional 5-day fermentation of Brazilian cocoa beans, followed by a 4-day sun drying
25 stage. The results shown in Figure 15L showed no obvious effect of these early processing stages, suggesting little change in the composition of the polyphenols. However, it is known (Lehrian and Patterson, 1983) that polyphenol oxidase (PPO) will oxidize polyphenols during
30 the fermentation stage. To determine what effect enzymically oxidized polyphenols would have on activity, another experiment was performed. Crude PPO was prepared by extracting finely ground, unfermented, freeze dried, defatted Brazilian cocoa beans with acetone at a ratio of
35 1 gm powder to 10 mL acetone. The slurry was centrifuged at 3,000 rpm for 15 min. This was repeated three times, discarding the supernatant each time with the fourth

extraction being poured through a Buchner filtering funnel. The acetone powder was allowed to air dry, followed by assay according to the procedures described by McLord and Kilara, (1983). To a solution of crude
5 polyphenols (100mg/10mL Citrate-Phosphate buffer, 0.02M, pH 5.5) 100mg of acetone powder (4,000 μ /mg protein) was added and allowed to stir for 30 min. with a stream of air bubbled through the slurry. The sample was centrifuged at 5,000 xg for 15 min. and the supernatant
10 extracted 3X with 20 mL ethyl acetate. The ethyl acetate extracts were combined, taken to dryness by distillation under partial vacuum and 5mL water added, followed by lyophilization. The material was then assayed against Hela cells and the dose-response compared to crude
15 polyphenol extracts that were not enzymically treated. The results (Figure 15M) showed a significant shift in the dose-response curve for the enzymically oxidized extract, showing that the oxidized products were more inhibitory than their native forms.

20 **Example 9: Antioxidant Activity of Cocoa Extracts
 Containing Procyanidins**

Evidence in the literature suggests a relationship between the consumption of naturally
25 occurring antioxidants (Vitamins C, E and B-carotene) and a lowered incidence of disease, including cancer (Designing Foods, 1993; Caragay, 1992). It is generally thought that these antioxidants affect certain oxidative and free radical processes involved with some types of
30 tumor promotion. Additionally, some plant polyphenolic compounds that have been shown to be anticarcinogenic, also possess substantial antioxidant activity (Ho et al., 1992; Huang et al., 1992).

To determine whether cocoa extracts containing
35 procyanidins possessed antioxidant properties, a standard Rancimat method was employed. The procedures described in Examples 1, 2 and 3 were used to prepare cocoa extracts which were manipulated further to produce two

fractions from gel permeation chromatography. These two fractions are actually combined fractions A through C, and D and E (See Figure 1) whose antioxidant properties were compared against the synthetic antioxidants BHA and BHT.

Peanut Oil was pressed from unroasted peanuts after the skins were removed. Each test compound was spiked into the oil at two levels, ~ 100 ppm and ~ 20 ppm, with the actual levels given in Table 7. 50 μ L of methanol solubilized antioxidant was added to each sample to aid in dispersion of the antioxidant. A control sample was prepared with 50 μ L of methanol containing no antioxidant.

The samples were evaluated in duplicate, for oxidative stability using the Rancimat stability test at 100°C and 20 cc/min of air. Experimental parameters were chosen to match those used with the Active Oxygen Method (AOM) or Swift Stability Test (Van Oosten et al., 1981). A typical Rancimat trace is shown in Figure 16. Results are reported in Table 8 as hours required to reach a peroxide level of 100 meq.

Table 7: Concentrations of Antioxidants

SAMPLE	LEVEL 1	LEVEL 2
		ppm
Butylated Hydroxytoluene (BHT)	24	120
Butylated Hydroxyanisole (BHA)	24	120
Crude Ethyl Acetate Fraction of Cocoa	22	110
Fraction A-C	20	100
Fraction D-E	20	100

**Table 8: Oxidative Stability of Peanut Oil
with Various Antioxidants**

5	SAMPLE	20 ppm	100 ppm
		average	
10	Control	10.5 $\pm$ 0.7	
	BHT	16.5 $\pm$ 2.1	12.5 $\pm$ 2.1
	BHA	13.5 $\pm$ 2.1	14.0 $\pm$ 1.4
	Crude Cocoa Fraction	18.0 $\pm$ 0.0	19.0 $\pm$ 1.4
	Fraction A-C	16.0 $\pm$ 6.4	17.5 $\pm$ 0.0
	Fraction D-E	14.0 $\pm$ 1.4	12.5 $\pm$ 0.7

These results demonstrated increased oxidative stability of peanut oil with all of the additives tested. The highest increase in oxidative stability was realized by the sample spiked with the crude ethyl acetate extract of cocoa. These results demonstrated that cocoa extracts containing procyanidins have antioxidant potential equal to or greater than equal amounts of synthetic BHA and BHT. Accordingly, the invention may be employed in place of BHT or BHA in known utilities of BHA or BHT, such as for instance as an antioxidant and/or food additive. And, in this regard, it is noted too that the invention is from an edible source. Given these results, the skilled artisan can also readily determine a suitable amount of the invention to employ in such "BHA or BHT" utilities, e.g., the quantity to add to food, without undue experimentation.

Example 10: Topoisomerase II Inhibition Study

DNA topoisomerase I and II are enzymes that catalyze the breaking and rejoining of DNA strands, thereby controlling the topological states of DNA (Wang, 1985). In addition to the study of the intracellular function of topoisomerase, one of the most significant findings has been the identification of topoisomerase II as the primary cellular target for a number of clinically important antitumor compounds (Yamashita et al., 1990)

which include intercalating agents (m-AMSA, Adriamycin® and ellipticine) as well as nonintercalating epipodophyllotoxins. Several lines of evidence indicate that some antitumor drugs have the common property of stabilizing the DNA - topoisomerase II complex ("cleavable complex") which upon exposure to denaturing agents results in the induction of DNA cleavage (Muller et al., 1989). It has been suggested that the cleavable complex formation by antitumor drugs produces bulky DNA adducts that can lead to cell death.

According to this attractive model, a specific new inducer of DNA topoisomerase II cleavable complex is useful as an anti-cancer, anti-tumor or antineoplastic agent. In an attempt to identify cytotoxic compounds with activities that target DNA, the cocoa procyanidins were screened for enhanced cytotoxic activity against several DNA - damage sensitive cell lines and enzyme assay with human topoisomerase II obtained from lymphoma.

A. Decatenation of Kinetoplast DNA by Topoisomerase II

The in vitro inhibition of topoisomerase II decatenation of kinetoplast DNA, as described by Muller et al. (1989), was performed as follows. Nuclear extracts containing topoisomerase II activity were prepared from human lymphoma by modifications of the methods of Miller et al. (1981) and Danks et al. (1988). One unit of purified enzyme was enough to decatenate 0.25 µg of kinetoplast DNA in 30 min. at 34°C. Kinetoplast DNA was obtained from the trypanosome *Crithidia fasciculata*. Each reaction was carried out in a 0.5 mL microcentrifuge tube containing 19.5 µL H₂O, 2.5 µL 10X buffer (1X buffer contains 50 mM tris-HCl, pH 8.0, 120 mM KCl, 10mM MgCl₂, 0.5 mM ATP, 0.5 mM dithiothreitol and 30 µg BSA/mL), 1 µL kinetoplast DNA (0.2 µg), and 1 µL DMSO-containing cocoa procyanidin test fractions at various concentrations. This combination was mixed thoroughly and kept on ice. One unit of topoisomerase was added

immediately before incubation in a waterbath at 34°C for 30 min.

Following incubation, the decatenation assay was stopped by the addition of 5 µL stop buffer (5% sarkosyl, 0.0025% bromophenol blue, 25% glycerol) and placed on ice. DNA was electrophoresed on a 1% agarose gel in TAE buffer containing ethidium bromide (0.5 µg/mL). Ultraviolet illumination at 310 nm wavelength allowed the visualization of DNA. The gels were photographed using a PolaroidTM Land camera.

Figure 17 shows the results of these experiments. Fully catenated kinetoplast DNA does not migrate into a 1% agarose gel. Decatenation of kinetoplast DNA by topoisomerase II generates bands of monomeric DNA (monomer circle, forms I and II) which do migrate into the gel. Inhibition of the enzyme by addition of cocoa procyanidins is apparent by the progressive disappearance of the monomer bands as a function of increasing concentration. Based on these results, cocoa procyanidin fractions A, B, D, and E were shown to inhibit topoisomerase II at concentrations ranging from 0.5 to 5.0 µg/mL. These inhibitor concentrations were very similar to those obtained for mitoxanthrone and m-AMSA [4'-(9-acridinylamino) methanesulfon-m-anisidide].

B. Drug Sensitive Cell Lines

Cocoa procyanidins were screened for cytotoxicity against several DNA-damage sensitive cell lines. One of the cell lines was the xrs-6 DNA double strand break repair mutant developed by P. Jeggo (Kemp et al., 1984). The DNA repair deficiency of the xrs-6 cell line renders them particularly sensitive to x-irradiation, to compounds that produce DNA double strand breaks directly, such as bleomycin, and to compounds that inhibit topoisomerase II, and thus may indirectly induce double strand breaks as suggested by Warters et al. (1991). The cytotoxicity toward the repair deficient

line was compared to the cytotoxicity against a DNA repair proficient CHO line, BR1. Enhanced cytotoxicity towards the repair deficient (xrs-6) line was interpreted as evidence for DNA cleavable double strand break
5 formation.

The DNA repair competent CHO line, BR1, was developed by Barrows et al. (1987) and expresses O⁶-alkylguanine - DNA - alkyltransferase in addition to normal CHO DNA repair enzymes. The CHO double strand
10 break repair deficient line (xrs-6) was a generous gift from Dr. P. Jeggo and co-workers (Jeggo et al., 1989). Both of these lines were grown as monolayers in alpha-MEM containing serum and antibiotics as described in Example 6. Cells were maintained at 37°C in a humidified
15 5% CO₂ atmosphere. Before treatment with cocoa procyanidins, cells grown as monolayers were detached with trypsin treatment. Assays were performed using the MTT assay procedure described in Example 6.

The results (Figure 18) indicated no enhanced
20 cytotoxicity towards the xrs-6 cells suggesting that the cocoa procyanidins inhibited topoisomerase II in a manner different from cleavable double strand break formation. That is, the cocoa procyanidins interact with topoisomerase II before it has interacted with the DNA to
25 form a noncleavable complex.

Noncleavable complex forming compounds are relatively new discoveries. Members of the anthracyclines, podophyllin alkaloids, anthracenediones, acridines, and ellipticines are all approved for clinical
30 anti-cancer, anti-tumor or antineoplastic use, and they produce cleavable complexes (Liu, 1989). Several new classes of topoisomerase II inhibitors have recently been identified which do not appear to produce cleavable complexes. These include amonafide (Hsiang et al.,
35 1989), distamycin (Fesen et al., 1989), flavanoids (Yamashita et al., 1990), saintopin (Yamashita et al., 1991), membranone (Drake et al., 1989), terpenoids

(Kawada *et al.*, 1991), anthrapyrazoles (Fry *et al.*, 1985), dioxopiperazines (Tanabe *et al.*, 1991), and the marine acridine - dercitin (Burres *et al.*, 1989).

Since the cocoa procyanidins inactivate
5 topoisomerase II before cleavable complexes are formed,
they have chemotherapy value either alone or in
combination with other known and mechanistically defined
topoisomerase II inhibitors. Additionally, cocoa
procyanidins also appear to be a novel class of
10 topoisomerase II inhibitors, (Kashiwada *et al.*, 1993) and
may thus be less toxic to cells than other known
inhibitors, thereby enhancing their utility in
chemotherapy.

The human breast cancer cell line MCF-7 (ADR)
15 which expresses a membrane bound glycoprotein (gp170) to
confer multi-drug resistance (Leonessa *et al.*, 1994) and
its parental line MCF-7 p168 were used to assay the
effects of cocoa fraction D & E. As shown in Figure 19,
the parental line was inhibited at increasing dose levels
20 of fraction D & E, whereas the adriamycin (ADR) resistant
line was less effected at the higher doses. These
results show that cocoa fraction D & E has an effect on
multi-drug resistant cell lines.

Example 11: Synthesis of Procyanidins

25 The synthesis of procyanidins was performed
according to the procedures developed by Delcour *et al.*
(1983), with modification. In addition to condensing
(+)-catechin with dihydroquercetin under reducing
conditions, (-)-epicatechin was also used to reflect the
30 high concentrations of (-)-epicatechin that naturally
occur in unfermented cocoa beans. The synthesis products
were isolated, purified, analyzed, and identified by the
procedures described in Examples 3, 4 and 5. In this
manner, the biflavanoids, triflavanoids and
35 tetraflavanoids are prepared and used as analytical
standards and, in the manner described above with respect
to cocoa extracts.

Example 12: Assay of Normal Phase Semi-Preparative Fractions

Since the polyphenol extracts are compositionally complex, it was necessary to determine which components were active against cancer cell lines for further purification, dose-response assays and comprehensive structural identification. A normal phase semi preparative HPLC separation (Example 3B) was used to separate cocoa procyanidins on the basis of oligomeric size. In addition to the original extract, twelve fractions were prepared (Figures 2B and 15 O) and assayed at 100 μ g/mL and 25 μ g/mL doses against Hela to determine which oligomer possessed the greatest activity. As shown in Figure 20, fractions 4-11 (pentamer-dodecamer) demonstrated IC₅₀ values of approximately 25 μ g/mL. These results indicated that these specific oligomers had the greatest activity against Hela cells. Additionally, normal phase HPLC analysis of cocoa fraction D & E indicated that this fraction was enriched with these oligomers.

From the foregoing, it is clear that the extract and cocoa polyphenols, as well as the compositions method and kit, of the invention have utility. In this regard, it is mentioned that the invention is from an edible source and, that the activity in vitro can demonstrate at least some activity in vivo, especially considering the doses discussed above.

Additionally, the above description shows that the extract and cocoa polyphenols, as well as the compositions, method and kit have antioxidant activity like that of BHT and BHA, as well as oxidative stability. Thus, the invention can be employed in place of BHT or BHA in known utilities of BHA and BHT, such as an antioxidant, for instance, an antioxidant food additive. The invention can also be employed in place of topoisomerase-inhibitors in the presently known utilities therefor. Accordingly, there are many compositions and

methods envisioned by the invention; for instance, antioxidant or preservative compositions, topoisomerase-inhibiting compositions, methods for preserving food or any desired item such as from oxidation, and methods for
5 inhibiting topoisomerase which comprise either the extract and/or cocoa polyphenol(s) or which comprise contacting the food, item or topoisomerase with the respective composition or with the extract and/or cocoa polyphenol(s).

10 Having thus described in detail the preferred embodiments of the present invention, it is to be understood that the invention defined by the appended claims is not to be limited by particular details set forth in the above descriptions as many apparent
15 variations thereof are possible without departing from the spirit or scope of the present invention.

REFERENCES

1. Barrows, L.R., Borchers, A.H., and Paxton, M.B.,
Transfectant CHO Cells Expressing O⁶ - alkylguanine
- DNA-alkyltransferase Display Increased Resistance
to DNA Damage Other than O⁶-guanine Alkylation,
Carcinogenesis, 8:1853 (1987).
2. Boukharta, M., Jalbert, G. and Castonguay, A.,
Efficacy of Ellagitannins and Ellagic Acid as Cancer
Chemopreventive Agents - Presented at the XVIth
International Conference of the Groupe Polyphenols,
Lisbon, Portugal, July 13-16, 1992.
3. Burres, N.S., Sazesh, J., Gunawardana, G.P., and
Clement, J.J., Antitumor Activity and Nucleic Acid
Binding Properties of Dercitin, a New Acridine
Alkaloid Isolated from a Marine Dercitus species
Sponge, Cancer Research, 49, 5267-5274 (1989).
4. Caragay, A.B., Cancer Preventive Foods and
Ingredients, Food Technology, 46:4, 65-79 (1992).
5. Chu, S.-C., Hsieh, Y.-S. and Lim, J.-Y., Inhibitory
Effects of Flavonoids on Maloney Murine Leukemia
Virus Reverse Transcriptase Activity, J. of Natural
Products, 55:2, 179-183 (1992).
6. Clapperton, J., Hammerstone, J.F. Jr., Romanczyk,
L.J. Jr., Chan, J., Yow, S., Lim, D. and Lockwood,
R., Polyphenols and Cocoa Flavor - Presented at the
XVIth International Conference of the Groupe
Polyphenols, Lisbon, Portugal, July 13-16, 1992.
7. Danks, M.K., Schmidt, C.A., Cirtain, M.C., Suttle,
D.P., and Beck, W.T., Altered Catalytic Activity of
and DNA Cleavage by DNA Topoisomerase II from Human
Leukemic Cells Selected for Resistance to VM-26,
Biochem., 27:8861 (1988).
8. Delcour, J.A., Ferreira, D. and Roux, D.G.,
Synthesis of Condensed Tannins, Part 9, The
Condensation Sequence of Leucocyanidin with (+)-
Catechin and with the Resultant Procyanidins, J.
Chem. Soc. Perkin Trans. I, 1711-1717 (1983).
9. Deschner, E.E., Ruperto, J., Wong, G. and Newmark,
H.L., Quercetin and Rutin as Inhibitors of
Azoxymethanol - Induced Colonic Neoplasia,
Carcinogenesis, 7, 1193-1196 (1991).
10. Designing Foods, Manipulating Foods to Promote
Health, Inform, 4:4, 344-369 (1993).
11. Drake, F.H., Hofmann, G.A., Mong., S.-M., Bartus,
J.O., Hertzberg, R.P., Johnson, R.K., Mattern, M.R.,

- and Mirabelli, C.K., In vitro and Intercellular Inhibition of Topoisomerase II by the Antitumor Agent Membranone, *Cancer Research*, **49**, 2578-2583 (1989).
- 5
12. Engels J.M.M., Genetic Resources of Cacao: A Catalogue of the CATIE Collection, Tech. Bull. 7, Turrialba, Costa Rica (1981).
- 10
13. Enriquez G.A. and Soria J.V., Cocoa Cultivars Register IICA, Turrialba, Cost Rica (1967).
14. Ferreira, D., Steynberg, J.P., Roux, D.G. and Brandt, E.V., Diversity of Structure and Function in Oligomeric Flavanoids, *Tetrahedron*, **48:10**, 1743-1803 (1992).
- 15
15. Fesen, M. and Pommier, Y., Mammalian Topoisomerase II Activity is Modulated by the DNA Minor Groove Binder - Distainycin in Simian Virus 40 DNA, *J. Biol. Chem.*, **264**, 11354-11359 (1989).
- 20
16. Fry, D.W., Boritzki, T.J., Besserer, J.A., and Jackson, R.C., In vitro Strand Scission and Inhibition of Nucleic Acid Synthesis on L1210 Leukemia Cells by a New Class of DNA Complexes, the anthra [1, 9-CD]pyrazol-6(2H)-ones (anthrapyrazoles), *Biochem. Pharmacol.*, **34**, 3499-3508 (1985).
- 25
17. Hsiang, Y.-H., Jiang, J.B., and Liu, L.F., Topoisomerase II Mediated DNA Cleavage by Amonafide and Its Structural Analogs, *Mol. Pharmacol.*, **36**, 371-376 (1989).
- 30
18. Jalal, M.A.F. and Collin, H.A., Polyphenols of Mature Plant, Seedling and Tissue Cultures of *Theobroma Cacao*, *Phytochemistry*, **6**, 1377-1380 (1978).
- 35
19. Jeggo, P.A., Caldecott, K., Pidsley, S., and Banks, G.R., Sensitivity of Chinese Hamster Ovary Mutants Defective in DNA Double Strand Break Repair to Topoisomerase II Inhibitors, *Cancer Res.*, **49:7057** (1989).
- 40
20. Kashiwada, Y., Nonaka, G.-I., Nishioka, I., Lee, K.J.-H., Bori, I., Fukushima, Y., Bastow, K.F., and Lee, K.-H., Tannin as Potent Inhibitors of DNA Topoisomerase II in vitro, *J. Pharm. Sci.*, **82:5**, 487-492 (1993).
- 45
21. Kato, R., Nakadate, T., Yamamoto, S. and Sugimura, T., Inhibition of 12-O-tetradecanoylphorbol-13-acetate Induced Tumor Promotion and Ornithine Decarboxylase Activity by Quercitin: Possible
- 50
- 55

Involvement of Lipoxygenase Inhibition,
Carcinogenesis, 4, 1301-1305 (1983).

- 5 22. Kawada, S.-Z., Yamashita, Y., Fujii, N. and Nakano, H., Induction of Heat Stable Topoisomerase II-DNA Cleavable Complex by Nonintercalative Terpenoids, Terpentecin and Clerocidin, Cancer Research, 51, 2922-2929 (1991).
- 10 23. Kemp, L.M., Sedgwick, S.G. and Jeggo, P.A., X-ray Sensitive Mutants of Chinese Hamster Ovary Cells Defective in Double Strand Break Rejoining, Mutat. Res., 132:189 (1984).
- 15 24. Kikkoman Corporation, Antimutagenic Agent Containing Proanthocyanidin Oligomer Preferably Having Flavan-3-ol-Diol Structure, JP 04190774-A, July 7, 1992.
- 20 25. Lehrian, D.W.; Patterson, G.R. In Biotechnology; Reed, G., Ed.; Verlag Chemie: Weinheim, 1983, Vol.5, Chapter 12.
- 25 26. Leonessa, F., Jacobson, M., Boyle, B., Lippman, J., McGarvey, M., and Clarke, R. Effect of Tamoxifen on the Multidrug-Resistant Phenotype in Human Breast Cancer Cells: Isobolograms, Drug Accumulation, and M_r 170,000 Glycoprotein (gp 170) Binding Studies, Cancer Research, 54, 441-447 (1994).
- 30 27. Liu, L.F., DNA Topoisomerase Poisons as Antitumor Drugs, Ann. Rev. Biochem., 58, 351-375 (1989).
- 35 28. McCord, J.D. and Kilara A. Control of Enzymatic Browning in Processed Mushrooms (*Agaricus bisporus*). J. Food Sci., 48:1479 (1983).
- 40 29. Miller, K.G., Liu, L.F. and Englund, P.A., Homogeneous Type II DNA Topoisomerase from Hela Cell Nuclei, J. Biol. Chem., 256:9334 (1981).
- 45 30. Mosmann, T., Rapid Colorimetric Assay for Cellular Growth and Survival: Application to Proliferation and Cytotoxicity Assays, J. Immunol. Methods, 65, 55 (1983).
- 50 31. Muller, M.T., Helal, K., Soisson, S. and Spitzer, J.R., A Rapid and Quantitative Microtiter Assay for Eukaryotic Topoisomerase II, Nuc. Acid Res., 17:9499 (1989).
- 55 32. Nawata, H., Chong, M.T., Bronzert, D. and Lippman, M.E. Estradiol-Independent growth of a Subline of MCF-7 Human Breast Cancer Cells in Culture, J. Biol. Chem., 256:13, 6895-6902 (1981).

33. Okuda, T., Yoshida, T., and Hatano, T., Molecular Structures and Pharmacological Activities of Polyphenols - Oligomeric Hydrolyzable Tannins and Others - Presented at the XVIth International Conference of the Groupe Polyphenols, Lisbon, Portugal, July 13-16, 1992.
34. Phenolic Compounds in Foods and Their Effects on Health II. Antioxidants & Cancer Prevention, Huang, M.-T., Ho, C.-T., and Lee, C.Y. editors, ACS Symposium Series 507, American Chemical Society, Washington, D.C. (1992).
35. Phenolic Compounds in Foods and Their Effects on Health I, Analysis, Occurrence & Chemistry, Ho, C.-T., Lee, C.Y., and Huang, M.-T editors, ACS Symposium Series 506, American Chemical Society, Washington, D.C. (1992).
36. Porter, L.J., Ma, Z. and Chan, B.G., Cocoa Procyanidins: Major Flavanoids and Identification of Some Minor Metabolites, *Phytochemistry*, 30, 1657-1663 (1991).
37. Revilla, E., Bourzeix, M. and Alonso, E., Analysis of Catechins and Procyanidins in Grape Seeds by HPLC with Photodiode Array Detection, *Chromatographia*, 31, 465-468 (1991).
38. Scudiero, D.A., Shoemaker, R.H., Paull, K.D., Monks, A., Tierney, S., Nofziger, T.H., Currens, M.J., Seniff, D., and Boyd, M.R. Evaluation of a Soluble Tetrazolium/Formazan Assay for Cell Growth and Drug Sensitivity in Culture Using Human and Other Tumor Cell Lines, *Canur Research*, 48, 4827-4833 (1988).
39. Self, R., Eagles, J., Galletti, G.C., Mueller-Harvey, I., Hartley, R.D., Lee, A.G.H., Magnolato, D., Richli, U., Gujur, R. and Haslam, E., Fast Atom Bombardment Mass Spectrometry of Polyphenols (syn. Vegetable Tannins), *Biomed Environ. Mass Spec.* 13, 449-468 (1986).
40. Tanabe, K., Ikegami, Y., Ishida, R. and Andoh, T., Inhibition of Topoisomerase II by Antitumor Agents bis(2,6-dioxopiperazine) Derivatives, *Cancer Research*, 51, 4903-4908 (1991).
41. Van Oosten, C.W., Poot, C. and A.C. Hensen, The Precision of the Swift Stability Test, *Fette, Seifen, Anstrichmittel*, 83:4, 133-135 (1981).
42. Wang, J.C., DNA Topoisomerases, *Ann. Rev. Biochem.*, 54, 665-697 (1985).

43. Wartens, R.L., Lyons, B.W., Li, T.M. and Chen, D.J.,
Topoisomerase II Activity in a DNA Double-Strand
Break Repair Deficient Chinese Hamster Ovary Cell
Line, Mutat. Res., 254:167 (1991).
5
44. Yamashita, Y., Kawada, S.-Z. and Nakano, H.,
Induction of Mammalian Topoisomerase II Dependent DNA
Cleavage by Nonintercalative Flavanoids, Genistein
and Orbol., Biochem Pharm, 39:4, 737-744 (1990).
10
45. Yamashita, Y., Kawada, S.-Z., Fujii, N. and Nakano,
H., Induction of Mammalian DNA Topoisomerase I and
II Mediated DNA Cleavage by Saintopin, a New
Antitumor Agent from Fungus, Biochem., 30, 5838-5845
15 (1991).


CLAIMS

1. A solid composition suitable for oral administration which comprises a suitable carrier and a cocoa extract comprising at least one cocoa procyanidin oligomer 4 to 12.
2. A solid composition suitable for oral administration which comprises a suitable carrier and an extract of unfermented or partially fermented cocoa beans, the extract comprising at least one cocoa procyanidin selected from epicatechin, procyanidin B-2, procyanidin oligomers 2 to 12, procyanidin B-5, procyanidin A-2 and procyanidin C-1.
3. A solid composition suitable for oral administration which comprises a suitable carrier and a synthetic procyanidin oligomer 2 to 12 or a gallate ester thereof.
4. A solid composition according to any one of claims 1 to 3 which is a chewable solid composition.
5. A solid composition according to any one of claims 1 to 4 which is a cocoa- or chocolate-flavoured solid composition.
6. A liquid preparation suitable for oral administration which comprises a suitable carrier and a cocoa extract comprising at least one cocoa procyanidin oligomer 4 to 12.
7. A liquid preparation suitable for oral administration which comprises a suitable carrier and an extract of unfermented or partially fermented cocoa beans, the extract comprising at least one cocoa procyanidin selected from epicatechin, procyanidin B-2, procyanidin oligomers 2 to 12, procyanidin B-5, procyanidin A-2 and procyanidin C-1.

8. A liquid preparation suitable for oral administration which comprises a suitable carrier and a synthetic procyanidin oligomer 2 to 12 or a gallate ester thereof.
9. A liquid preparation according to claim 8 which comprises a synthetic procyanidin oligomer 2 to 5 or a gallate ester thereof.
10. A liquid preparation according to any one of claims 6 to 9 which is a suspension, syrup or elixir.
11. A tablet, capsule or pill which comprises a suitable carrier and a cocoa extract comprising a cocoa procyanidin oligomer 2 to 12.
12. A tablet, capsule or pill which comprises a suitable carrier and a synthetic procyanidin oligomer 2 to 12 or a gallate ester thereof.
13. A cocoa extract comprising cocoa polyphenols, which extract is obtained by a process which comprises:
 - reducing unfermented or partially fermented cocoa beans to a powder;
 - defatting the powder;
 - extracting cocoa polyphenols from the powder by solvent extraction techniques; and
 - purifying the cocoa extract thus obtained.
14. A cocoa extract according to claim 13 wherein the process further comprises purifying the extract by gel permeation chromatography and/or by high performance liquid chromatography (HPLC).

15. A cocoa extract according to claim 13 or 14 which comprises at least one cocoa procyanidin selected from (-)-epicatechin, procyanidin B-2, procyanidin oligomers 4 –12, procyanidin B-5, procyanidin A-2 and procyanidin C-1.
16. A cocoa extract comprising cocoa polyphenols, which comprises at least one each of a cocoa procyanidin dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer, decamer, undecamer and dodecamer.
17. A cocoa extract comprising cocoa polyphenols, which extract is free of xanthine alkaloids and is obtained by a process which comprises:
 - reducing cocoa beans to powder;
 - defatting the powder;
 - extracting cocoa polyphenols from the powder by solvent extraction techniques; and
 - removing xanthine alkaloids from the extract to yield a xanthine alkaloid-free subfraction.
18. A cocoa extract according to any one of claims 13 to 15 or claim 17 wherein the process of reducing cocoa beans to powder comprises:
 - freeze-drying beans and pulp;
 - depulping the freeze-dried mass;
 - dehulling the freeze-dried cocoa beans; and
 - grinding the dehulled beans.
19. A cocoa extract according to claim any one of claims 13 to 18 which is in lyophilised form.

Figure 1: Gel Permeation Chromatogram of Crude Procyanidins on Sephadex LH-20



Chromatographic Conditions: Column; 28 x 2.5 cm Sephadex LH-20, Mobile Phase: Methanol/Water Step Gradient, 15:85, 25:75, 35:65, 70:30, 100:0 Stepped at 1/2 Hour Intervals, Flow Rate; 1.5 ml/min, Detector; UV @ $\lambda_1=254$ nm and $\lambda_2=365$ nm, Chart Speed; 0.5 mm/min, Column Load; 120 mg.

Figure 2 A: Elution Profile of Cocoa Procyanidins Extracted from UIT-1 Unfermented Cocoa





Figure 2B. Analytical Normal Phase HPLC Separation of Cocoa Procyanidins



HPLC Conditions:

250 x 3.2mm Lichrosphere 5 Silica column (5 μ)20 x 4.6mm Supelguard LC-Si (5 μ) guard column


Detector: Photodiode Array @ 280nm

Gradient: Time (min.)	CH ₂ Cl ₂	Methanol	Acetic Acid/Water (1:1)
0	82	14	4
30	67.6	28.4	4
60	46	50	4
65	10	86	4
70	10	86	4


Flow rate: 0.5 mL/min

Column Temperature: 37°C


Figure 3: Representative Structures of Cocoa Procyanidins




(-) Epicatechin




Procyanidin B-2




Procyanidin Oligomers n = 2 through 5



Procyanidin B-5



Procyanidin A-2




Procyanidin C-1

HPLC Conditions: Hewlett Packard 1090 ternary HPLC System equipped with HP Model 1046A Programmable Fluorescence Detector.

Column: 5 μ Hypersil ODS (200 x 2.1 mm)

Linear gradient of 60% B into A at a flow rate of 0.3 ml/min. B=0.5% acetic acid in methanol; A=0.5% acetic acid in deionized water. λ_{ex} =280nm; λ_{em} =316nm

Figure 4A Representative HPLC Traces of Procyanidin Fractions




5/91

HPLC Conditions: Hewlett Packard 1090 ternary HPLC System equipped with HP Model 1046A Programmable Fluorescence Detector.

Column: 5 μ Hypersil ODS (200 x 2.1 mm)

Linear gradient of 60% B into A at a flow rate of 0.3 ml/min. B=0.5% acetic acid in methanol; A=0.5% acetic acid in deionized water. λ_{ex} =280nm; λ_{em} =316nm

Figure 4B Representative HPLC Traces of Procyanidin Fractions




HPLC Conditions: Hewlett Packard 1090 ternary HPLC System equipped with HP Model 1046A Programmable Fluorescence Detector.

Column: 5 μ Hypersil ODS (200 x 2.1 mm)

Linear gradient of 60% B into A at a flow rate of 0.3 ml/min. B=0.5% acetic acid in methanol; A=0.5% acetic acid in deionized water. λ_{ex} =280nm; λ_{em} =316nm

Figure 4C Representative HPLC Traces of Procyanidin Fractions



HPLC Conditions: Hewlett Packard 1090 ternary HPLC System equipped with HP Model 1046A Programmable Fluorescence Detector.

Column: 5 μ Hypersil ODS (200 x 2.1 mm)

Linear gradient of 60% B into A at a flow rate of 0.3 ml/min. B=0.5% acetic acid in methanol; A=0.5% acetic acid in deionized water. λ_{ex} =280nm; λ_{em} =316nm




Figure 4D Representative HPLC Traces of Procyanidin Fractions

HPLC Conditions: Hewlett Packard 1090 ternary HPLC System equipped with HP Model 1046A Programmable Fluorescence Detector.

Column: 5 μ Hypersil ODS (200 x 2.1 mm)

Linear gradient of 60% B into A at a flow rate of 0.3 ml/min. B=0.5% acetic acid in methanol; A=0.5% acetic acid in deionized water. λ_{ex} =280nm; λ_{em} =316nm




Figure 4E Representative HPLC Traces of Procyanidin Fractions

Figure 5: Dose - Response Relationship Between Amount MM2 and ACHN Survival






Figure 6A Dose-Response Relationships Between Combinations of Procyanidin Fractions and PC-3 Cancer Cell Line




11/91

Figure 6B Dose-Response Relationships Between Combinations of Procyanidin Fractions and PC-3 Cancer Cell Line




12/91

Figure 6C Dose-Response Relationships Between Combinations
of Procyanidin Fractions and PC-3 Cancer Cell Line



13/91

Figure 6D Dose-Response Relationships Between Combinations of Procyanidin Fractions and PC-3 Cancer Cell Line



14/91

Figure 7A

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Individual Procyanidin Fractions




Figure 7B

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Individual Procyanidin Fractions




Figure 7C

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Individual Procyanidin Fractions





Figure 7D

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Individual Procyanidin Fractions



18/91

Figure 7E

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Individual Procyanidin Fractions




Figure 7F

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Representative Procyanidin Fraction Combinations




Figure 7G

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Representative Procyanidin Fraction Combinations




Figure 7H

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the PC-3 Prostate Cell Line

Representative Procyanidin Fraction Combinations




Figure 8A

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Individual Procyanidin Fractions




Figure 8B

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Individual Procyanidin Fractions




Figure 8C

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Individual Procyanidin Fractions




Figure 8D

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Individual Procyanidin Fractions




Figure 8E

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Individual Procyanidin Fractions




Figure 8F

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Representative Procyanidin Fraction Combinations




Figure 8G

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Representative Procyanidin Fraction Combinations





Figure 8H

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the KB Nasopharyngeal/HeLa Cell Line

Representative Procyanidin Fraction Combinations



30/91

Figure 9A

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Individual Procyanidin Fractions




Figure 9B

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Individual Procyanidin Fractions




Figure 9C

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Individual Procyanidin Fractions




Figure 9D

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Individual Procyanidin Fractions




Figure 9E

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
HCT-116 Cell Line

Individual Procyanidin Fractions




Figure 9F

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Representative Procyanidin Fraction Combinations




Figure 9G

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Representative Procyanidin Fraction Combinations




Figure 9H

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the HCT-116 Cell Line

Representative Procyanidin Fraction Combinations




Figure 10A

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the ACHN Renal Cell Line

Individual Procyanidin Fractions




Figure 10B

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
ACHN Renal Cell Line

Individual Procyanidin Fractions




Figure 10C

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the ACHN Renal Cell Line

Individual Procyanidin Fractions




Figure 10D

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
ACHN Renal Cell Line

Individual Procyanidin Fractions




Figure 10E

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
ACHN Renal Cell Line

Individual Procyanidin Fractions




Figure 10F Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
ACHN Renal Cell Line

Representative Procyanidin Fraction Combinations




Figure 10G

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the ACHN Renal Cell Line

Representative Procyanidin Fraction Combinations




Figure 10H

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the ACHN Renal Cell Line

Representative Procyanidin Fraction Combinations





Figure 11A

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Individual Procyanidin Fractions



47/91

Figure 11B

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Individual Procyanidin Fractions




Figure 11C

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Individual Procyanidin Fractions




Figure 11D

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Individual Procyanidin Fractions




Figure 11E

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
A-549 Lung Cell Line

Individual Procyanidin Fractions




Figure 11F

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Representative Procyanidin Fraction Combinations




Figure 11G

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Representative Procyanidin Fraction Combinations




Figure 11H

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the A-549 Lung Cell Line

Representative Procyanidin Fraction Combinations




Figure 12A

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
SK-5 Melanoma Cell Line

Individual Procyanidin Fractions




Figure 12B

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
SK-5 Melanoma Cell Line

Individual Procyanidin Fractions




Figure 12C

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
SK-5 Melanoma Cell Line

Individual Procyanidin Fractions




Figure 12D

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the SK-5 Melanoma Cell Line

Individual Procyanidin Fractions




Figure 12E

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
SK-5 Melanoma Cell Line

Individual Procyanidin Fractions




Figure 12F

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the SK-5 Melanoma Cell Line

Representative Procyanidin Fraction Combinations




Figure 12G

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the SK-5 Melanoma Cell Line

Representative Procyanidin Fraction Combinations




Figure 12H Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
SK-5 Melanoma Cell Line

† Representative Procyanidin Fraction Combinations




Figure 13A

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7 Breast Cell Line

Individual Procyanidin Fractions




Figure 13B

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7 Breast Cell Line

Individual Procyanidin Fractions





Figure 13C

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7 Breast Cell Line

Individual Procyanidin Fractions




65/91

Figure 13D

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7 Breast Cell Line

Individual Procyanidin Fractions



66/91

Figure 13E

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7 Breast Cell Line

Individual Procyanidin Fractions




Figure 13F

Dose-Response Relationships Between Cocoa Procyanidin Fractions and the MCF-7 Breast Cell Line

Representative Procyanidin Fraction Combinations





Figure 13G

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7 Breast Cell Line

Representative Procyanidin Fraction Combinations




69/91

Figure 13H

Dose-Response Relationships Between
Cocoa Procyanidin Fractions and the
MCF-7-Breast Cell Line

Representative Procyanidin Fraction Combinations



70/91

FIGURE 14

Dose-Response Relationships Between Cocoa Procyanidin
Fraction D and the CCRF-CEM T-Cell Leukemia Cell Line




Figure 15 A: Comparison of XTT and Crystal Violet Cytotoxicity Assays on MCF - 7 p168 cells Treated with Fraction D + E






Figure 15 B: Dose Response for UTT-1 Crude Polyphenol Extract on MDA MB 231 Cells




* NOTE: ABSORBANCE OF 2.0 INDICATES THE MAXIMUM ABSORBANCE OF THE PLATE READER. IT IS NOT REPRESENTATIVE OF CELL NUMBER.

Figure 15 C: Dose Response for UIT-1 Crude Polyphenol Extract on PC-3 Cells




74/91

Figure 15 D: Dose Response for UIT-1 Crude Polyphenol Extract on MCF-7 p168 Cells




* NOTE: ABSORBANCE OF 2.0 INDICATES THE MAXIMUM ABSORBANCE OF THE PLATE READER. IT IS NOT REPRESENTATIVE OF CELL NUMBER.

Figure 15 E: Dose Response for UIT-1 Crude Polyphenol Extract on Hela Cells



* NOTE: ABSORBANCE OF 2.0 INDICATES THE MAXIMUM ABSORBANCE OF THE PLATE READER. IT IS NOT REPRESENTATIVE OF CELL NUMBER.

Figure 15 F: Cytotoxicity of Cocoa Fractions at 100 μ L/mL on Hela Cells



* NOTE: ABSORBANCE OF 2.0 INDICATES THE MAXIMUM ABSORBANCE OF THE PLATE READER. IT IS NOT REPRESENTATIVE OF CELL NUMBER.

Figure 15 G: Cytotoxicity Of Cocoa Fractions at 100 μ L/mL on SKBR - 3 Cells





Figure 15 H: Dose Response for Cocoa Fraction D + E on Hela Cells

* NOTE: ABSORBANCE OF 2.0 INDICATES THE MAXIMUM ABSORBANCE OF THE PLATE READER. IT IS NOT REPRESENTATIVE OF CELL NUMBER.


Figure 15 I: Dose Response for Cocoa Fraction D + E on SKBR - 3 Cells

Figure 15 J: Dose Response for Cocoa Fraction D + E on Hela Cells by Soft Agar Cloning Assay





Figure 15 K: Growth Inhibition of Hela Cells by Crude Cocoa Extracts Prepared from Different Cocoa Genotypes



#	GENOTYPE	HORTL. RACE	DESCRIPTION
C-1	UF-12	CRIOLO	CRUDE EXTRACTS OF UF-12 (BRAZIL) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-2	NA-33	FORASTERO	CRUDE EXTRACTS OF NA-33 (BRAZIL) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-3	EEG-48	FORASTERO	CRUDE EXTRACTS OF EEG-48 (BRAZIL) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-4	UNKNOWN	FORASTERO	CRUDE EXTRACTS OF UNKNOWN (W. AFRICAN) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-5	UF-613	TRINITARIO	CRUDE EXTRACTS OF UF-613 (BRAZIL) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-6	ICS-100	TRINITARIO	CRUDE EXTRACTS OF ICS-100 (BRAZIL) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-7	ICS-139	TRINITARIO	CRUDE EXTRACTS OF ICS-139 (BRAZIL) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)
C-8	UIT-1	TRINITARIO	CRUDE EXTRACTS OF UIT-1 (MALAYSIA) COCOA POLYPHENOLS (DECAFFEINATED/DETHREOBROMINATED)

Figure 15 L: Growth Inhibition of Hela Cells by Cocoa Polyphenol Extracts Taken at Different Time Stages Throughout a Fermentation and Sun Drying Stage




Figure 15 M: Dose Response for Polyphenol Oxidase Treated Crude Cocoa Polyphenol Extract




Figure 15 N: Reverse Phase Semi-Preparative HPLC Separation of Fraction D + E





Figure 15 O: Normal Phase Semi-Preparative HPLC Separation of a Crude Cocoa Polyphenol Extract



HPLC Conditions: 250 x 10mm Supelcosil LC-Si (5 μ m) Semipreparative Column
 20 x 4.6mm Supelcosil LC-Si (5 μ m) Guard Column
 Gradient: Time (min) CH₂Cl₂ Methanol Acetic Acid/H₂O (1:1)

0	82	14	4
30	67.6	28.4	4
60	46	50	4
65	10	86	4
70	10	86	4

Detector: Waters LC Spectrophotometer Model 480 @ 254nm
 Flow rate: 3mL/min, ambient temperature
 250 μ L of 70% aqueous acetone extract injected

Figure 16. Rancimat Oxidation Curves for Cocoa Procyanidins and Synthetic Antioxidants

20 ppm Sample Set





Figure 17. Inhibition of Topoisomerase II Catalyzed Decatenation of Kinetoplast DNA by Cocoa Procyanidin Fractions



Lane 1 contains 0.5 μg of marker (M) monomer-length kinetoplast DNA circles

Lanes 2 and 20 contain kinetoplast DNA that was incubated with Topoisomerase II in the presence of 4% DMSO, but in the absence of any cocoa procyanidins. (Control -C)

Lanes 3 and 4 contain kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.5 and 5.0 $\mu\text{g/mL}$ cocoa procyanidin fraction A.

Lanes 5 and 6 contain kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.5 and 5.0 $\mu\text{g/mL}$ cocoa procyanidin fraction B.

Lanes 7,8,9,13,14, and 15 are replicates of kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.05,0.5 and 5.0 $\mu\text{g/mL}$ cocoa procyanidin fraction D.

Lanes 10,11,12,16,17, and 18 are replicates of kinetoplast DNA that was incubated with Topoisomerase II in the presence of 0.05,0.5 and 5.0 $\mu\text{g/mL}$ cocoa procyanidin fraction E.

Lane 19 is a replicate of kinetoplast DNA that was incubated with Topoisomerase II in the presence of 5.0 $\mu\text{g/mL}$ cocoa procyanidin fraction E.

Figure 18. Dose Response Relationships Between Cocoa Procyanidin Fraction D and DNA Repair Competent and Deficient Cell Lines




Figure 19: Dose Response Curve for Adriamycin Resistant MCF-7 Cells in Comparison to MCF-7 p168 Parental Cell Line with Cocoa Fraction D + E





Figure 20: Dose Response Effect on Hela by Normal Phase Fractions

9/1/91