



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12N 15/54, 9/12, C12Q 1/68, A61K 38/45, C07R 16/40</p>	<p>A2</p>	<p>(11) International Publication Number: WO 00/22143</p> <p>(43) International Publication Date: 20 April 2000 (20.04.00)</p>
<p>(21) International Application Number: PCT/US99/24202</p> <p>(22) International Filing Date: 15 October 1999 (15.10.99)</p> <p>(30) Priority Data: 09/173,581 15 October 1998 (15.10.98) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/173,581 (CIP) Filed on 15 October 1998 (15.10.98)</p> <p>(71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): BANDMAN, Olga [US/US]; 366 Anna Avenue, Mountain View, CA 94043 (US). TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). HILLMAN, Jennifer, L. [US/US]; 230 Monroe Drive, #12, Mountain View, CA 94040 (US). YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale, CA 94087 (US). GUEGLER, Karl, J. [CH/US]; 1048 Oakland Avenue, Menlo Park, CA 94025 (US). CORLEY, Neil, C.</p>	<p>[US/US]; 1240 Dale Avenue, #30, Mountain View, CA 94040 (US). GORGONE, Gina, A. [US/US]; 1253 Pinecrest Drive, Boulder Creek, CA 95006 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). LU, Dyung, Aina, M. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US).</p> <p>(74) Agents: BILLINGS, Lucy, J. et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>	
<p>(54) Title: PROTEIN KINASE HOMOLOGS</p> <p>(57) Abstract</p> <p>The invention provides human protein kinase homologs (PKH) and polynucleotides which identify and encode PKH. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing disorders associated with expression of PKH.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PROTEIN KINASE HOMOLOGS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of protein kinase homologs and to
5 the use of these sequences in the diagnosis, treatment, and prevention of cancer,
autoimmune/inflammatory disorders, and reproductive disorders.

BACKGROUND OF THE INVENTION

Kinases and phosphatases are critical components of intracellular signal transduction
10 mechanisms. Kinases catalyze the transfer of high energy phosphate groups from adenosine triphosphate
(ATP) to hydroxyamino acids of various target proteins. Phosphatases, in contrast, remove phosphate
groups from proteins. Reversible protein phosphorylation is the main strategy for regulating protein
activity in eukaryotic cells. In general, proteins are activated by phosphorylation in response to
extracellular signals such as hormones, neurotransmitters, and growth and differentiation factors. Protein
15 dephosphorylation occurs when down-regulation of a signaling pathway is required. The combined
activities of kinases and phosphatases regulate key cellular processes such as proliferation, differentiation,
and cell cycle progression.

Kinases comprise the largest known enzyme superfamily and vary widely in their target proteins.
Kinases may be categorized as protein tyrosine kinases (PTKs), which phosphorylate tyrosine residues,
20 and protein serine/threonine kinases (STKs), which phosphorylate serine and/or threonine residues. Some
kinases have dual specificity for both serine/threonine and tyrosine residues. Almost all kinases contain a
conserved 250-300 amino acid catalytic domain. This domain can be further divided into 11 subdomains.
N-terminal subdomains I-IV fold into a two-lobed structure which binds and orients the ATP donor
molecule, and subdomain V spans the two lobes. C-terminal subdomains VI-XI bind the protein substrate
25 and transfer the gamma phosphate from ATP to the hydroxyl group of a serine, threonine, or tyrosine
residue. Each of the 11 subdomains contains specific catalytic residues or amino acid motifs
characteristic of that subdomain. For example, subdomain I contains an 8-amino acid glycine-rich ATP
binding consensus motif, subdomain II contains a critical lysine residue required for maximal catalytic
activity, and subdomains VI through IX comprise the highly conserved catalytic core. STKs and PTKs
30 also contain distinct sequence motifs in subdomains VI and VIII which may confer hydroxyamino acid
specificity. Some STKs and PTKs possess structural characteristics of both families. In addition, kinases
may also be classified by additional amino acid sequences, generally between 5 and 100 residues, which
either flank or occur within the kinase domain. These additional amino acid sequences regulate kinase
activity and determine substrate specificity. (Reviewed in Hardie, G. and S. Hanks (1995) The Protein
35 Kinase Facts Book, Vol I:7-20 Academic Press, San Diego CA.)

PTKs may be classified as either transmembrane or non-transmembrane proteins.

Transmembrane tyrosine kinases function as receptors for most growth factors. Binding of growth factor to the receptor activates the transfer of a phosphate group from ATP to selected tyrosine residues in the receptor itself and in specific second messenger proteins. Growth factors (GF) that associate with
5 receptor PTKs include epidermal GF, platelet-derived GF, fibroblast GF, hepatocyte GF, insulin and insulin-like GFs, nerve GF, vascular endothelial GF, and macrophage colony stimulating factor.

Non-transmembrane PTKs form signaling complexes with the cytosolic domains of plasma membrane receptors. Receptors that signal through non-transmembrane PTKs include cytokine, hormone, and antigen-specific lymphocytic receptors. Many PTKs were first identified as oncogene
10 products in cancer cells in which PTK activation was no longer subject to normal cellular controls. In fact, about one third of the known oncogenes encode PTKs. Furthermore, cellular transformation (oncogenesis) is often accompanied by increased tyrosine phosphorylation activity (Carbonneau, H. and N.K. Tonks (1992) *Annu. Rev. Cell Biol.* 8:463-93). Regulation of PTK activity may therefore be an important strategy in controlling some types of cancer.

15 The discovery of new protein kinase homologs and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of cancer, autoimmune/inflammatory disorders, and reproductive disorders.

SUMMARY OF THE INVENTION

20 The invention features substantially purified polypeptides, protein kinase homologs, referred to collectively as "PKH" and individually as "PKH-1", "PKH-2", "PKH-3", "PKH-4", "PKH-5", "PKH-6", "PKH-7", "PKH-8", and "PKH-9". In one aspect, the invention provides a substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof.

25 The invention further provides a substantially purified variant having at least 90% amino acid identity to at least one of the amino acid sequences selected from the group consisting of SEQ ID NO:1-9, and fragments thereof. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof. The invention also includes an isolated and purified polynucleotide variant having
30 at least 70% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof.

Additionally, the invention provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof. The invention
35 also provides an isolated and purified polynucleotide having a sequence which is complementary to the

polynucleotide encoding the polypeptide comprising the amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof.

The invention also provides an isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, and fragments thereof. The invention
5 further provides an isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, and fragments thereof. The invention also provides an isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, and fragments thereof.

10 The invention also provides a method for detecting a polynucleotide in a sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide sequence to at least one of the polynucleotides of the sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide in the sample. In one aspect, the method further comprises
15 amplifying the polynucleotide prior to hybridization.

The invention further provides an expression vector containing at least a fragment of the polynucleotide encoding the polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof. In another aspect, the expression vector is contained within a host cell.

20 The invention also provides a method for producing a polypeptide, the method comprising the steps of: (a) culturing the host cell containing an expression vector containing at least a fragment of a polynucleotide under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified
25 polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide selected from the group consisting of SEQ ID NO:1-9, and fragments thereof. The invention also provides a purified agonist and a purified antagonist to the polypeptide.

30 The invention also provides a method for treating or preventing a disorder of cell proliferation associated with decreased expression or activity of PKH, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising a substantially purified polypeptide having the amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof, in conjunction with a suitable pharmaceutical carrier.

35 The invention also provides a method for treating or preventing a disorder of cell proliferation

associated with increased expression or activity of PKH, the method comprising administering to a subject in need of such treatment an effective amount of an antagonist of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof.

5

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows nucleotide and polypeptide sequence identification numbers (SEQ ID NO), clone identification numbers (clone ID), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding PKH.

Table 2 shows features of each polypeptide sequence including potential motifs, homologous
10 sequences, and methods and algorithms used for characterization of PKH.

Table 3 shows the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis, diseases, disorders, or conditions associated with these tissues, and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which Incyte cDNA
15 clones encoding PKH were isolated.

Table 5 shows the programs, their descriptions, references, and threshold parameters used to analyze PKH.

DESCRIPTION OF THE INVENTION

20 Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

25 It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings
30 as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in
35 connection with the invention. Nothing herein is to be construed as an admission that the invention is not

entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

“PKH” refers to the amino acid sequences of substantially purified PKH obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and
5 preferably the human species, from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term “agonist” refers to a molecule which, when bound to PKH, increases or prolongs the duration of the effect of PKH. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of PKH.

10 An “allelic variant” is an alternative form of the gene encoding PKH. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of
15 these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding PKH include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same as PKH or a polypeptide with at least one functional characteristic of PKH. Included within this definition are
20 polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding PKH, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding PKH. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent PKH. Deliberate
25 amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of PKH is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine,
30 isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine.

The terms “amino acid” or “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, “fragments,” “immunogenic fragments,” or “antigenic fragments” refer to fragments of PKH
35 which are preferably at least 5 to about 15 amino acids in length, most preferably at least 14 amino acids,

and which retain some biological activity or immunological activity of PKH. Where “amino acid sequence” is recited to refer to an amino acid sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

5 “Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which, when bound to PKH, decreases the amount or the duration of the effect of the biological or immunological activity of PKH. Antagonists may include
10 proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of PKH.

The term “antibody” refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind PKH polypeptides can be prepared using intact polypeptides or using fragments containing small peptides
15 of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

20 The term “antigenic determinant” refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune
25 response) for binding to an antibody.

The term “antisense” refers to any composition containing a nucleic acid sequence which is complementary to the “sense” strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to
30 block either transcription or translation. The designation “negative” can refer to the antisense strand, and the designation “positive” can refer to the sense strand.

The term “biologically active,” refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, “immunologically active” refers to the capability of the natural, recombinant, or synthetic PKH, or of any oligopeptide thereof, to induce a specific immune
35 response in appropriate animals or cells and to bind with specific antibodies.

The terms “complementary” or “complementarity” refer to the natural binding of polynucleotides by base pairing. For example, the sequence “5' A-G-T 3'” bonds to the complementary sequence “3' T-C-A 5'.” Complementarity between two single-stranded molecules may be “partial,” such that only some of the nucleic acids bind, or it may be “complete,” such that total complementarity exists between the single
5 stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A “composition comprising a given polynucleotide sequence” or a “composition comprising a
10 given amino acid sequence” refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding PKH or fragments of PKH may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an
15 aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

“Consensus sequence” refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte
20 Clone using a computer program for fragment assembly, such as the GELVIEW Fragment Assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

The term “correlates with expression of a polynucleotide” indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding PKH, by northern
25 analysis is indicative of the presence of nucleic acids encoding PKH in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding PKH.

A “deletion” refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term “derivative” refers to the chemical modification of a polypeptide sequence, or a
30 polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was
35 derived.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" or "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MEGALIGN program (DNASTAR, Madison WI) which creates alignments between two or more sequences according to methods selected by the user, e.g., the clustal method. (See, e.g., Higgins, D.G. and P.M. Sharp (1988) *Gene* 73:237-244.) The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no similarity between the two amino acid sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by varying hybridization conditions.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization” refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

The term “hybridization complex” refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization
5 complex may be formed in solution (e.g., C₀t or R₀t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words “insertion” or “addition” refer to changes in an amino acid or nucleotide sequence
10 resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

“Immune response” can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and
15 systemic defense systems.

The term “microarray” refers to an arrangement of distinct polynucleotides on a substrate.

The terms “element” or “array element,” in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term “modulate” refers to a change in the activity of PKH. For example, modulation may
20 cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of PKH.

The phrases “nucleic acid” or “nucleic acid sequence,” as used herein, refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense
25 or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, “fragments” refers to those nucleic acid sequences which comprise a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:10-18, for example, as distinct from any other sequence in the same genome. For example, a fragment of SEQ ID NO:10-18 is useful in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:10-
30 18 from related polynucleotide sequences. A fragment of SEQ ID NO:10-18 is at least about 15-20 nucleotides in length. The precise length of the fragment of SEQ ID NO:10-18 and the region of SEQ ID NO:10-18 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment. In some cases, a fragment, when translated, would produce polypeptides retaining some functional characteristic, e.g., antigenicity, or structural domain
35 characteristic, e.g., ATP-binding site, of the full-length polypeptide.

The terms “operably associated” or “operably linked” refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the translation of the encoded polypeptide. While operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements, e.g., repressor
5 genes, are not contiguously linked to the sequence encoding the polypeptide but still bind to operator sequences that control expression of the polypeptide.

The term “oligonucleotide” refers to a nucleic acid sequence of at least about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or microarray. “Oligonucleotide” is
10 substantially equivalent to the terms “amplimer,” “primer,” “oligomer,” and “probe,” as these terms are commonly defined in the art.

“Peptide nucleic acid” (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially
15 bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

The term “sample” is used in its broadest sense. A sample suspected of containing nucleic acids encoding PKH, or fragments thereof, or PKH itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in
20 solution or bound to a substrate; a tissue; a tissue print; etc.

The terms “specific binding” or “specifically binding” refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope “A,” the presence of a polypeptide
25 containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term “stringent conditions” refers to conditions which permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt concentration, the concentration of organic solvent, e.g., formamide, temperature, and other conditions
30 well known in the art. In particular, stringency can be increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

The term “substantially purified” refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from other components with which they are naturally
35 associated.

A “substitution” refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

“Substrate” refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

“Transformation” describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term “transformed” cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A “variant” of PKH polypeptides refers to an amino acid sequence that is altered by one or more amino acid residues. The variant may have “conservative” changes, wherein a substituted amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have “nonconservative” changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE software (DNASTAR).

The term “variant,” when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to PKH. This definition may also include, for example, “allelic” (as defined above), “splice,” “species,” or “polymorphic” variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass “single nucleotide polymorphisms” (SNPs) in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

THE INVENTION

The invention is based on the discovery of new human protein kinase homologs (PKH), the polynucleotides encoding PKH, and the use of these compositions for the diagnosis, treatment, or prevention of cancer, autoimmune/inflammatory disorders, and reproductive disorders.

5 Table 1 lists the Incyte Clones used to derive full length nucleotide sequences encoding PKH. Columns 1 and 2 show the sequence identification numbers (SEQ ID NO) of the amino acid and nucleic acid sequences, respectively. Column 3 shows the Clone ID of the Incyte Clone in which nucleic acids encoding each PKH were identified, and column 4, the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones, their corresponding cDNA libraries, and shotgun sequences.
10 The clones and shotgun sequences are part of the consensus nucleotide sequence of each PKH and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of the polypeptides of the invention: column 1 references the amino acid SEQ ID NO; column 2 shows the number of amino acid residues in each polypeptide; column 3, potential phosphorylation sites; column 4, the amino acid residues comprising
15 signature sequences and motifs; column 5, the identity of each protein; and column 6, analytical methods used to characterize and identify each protein through sequence homology and protein motifs.

The columns of Table 3 show the tissue specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding PKH. The first column of Table 3 lists the nucleotide SEQ ID NO; the second column lists tissue categories which express PKH as a fraction of total tissue
20 categories expressing PKH. The third column lists the diseases, disorders, or conditions associated with those tissues expressing PKH. The fourth column lists the vectors used to subclone the cDNA library.

The following fragments of the nucleotide sequences encoding PKH are useful in hybridization or amplification technologies to identify SEQ ID NO:10-18 and to distinguish between SEQ ID NO:10-18 and related polynucleotide sequences. The useful fragments are the fragment of SEQ ID NO:10 from
25 about nucleotide 473 to about nucleotide 532; the fragment of SEQ ID NO:11 from about nucleotide 65 to about nucleotide 125; the fragment of SEQ ID NO:12 from about nucleotide 96 to about nucleotide 155; the fragment of SEQ ID NO:13 from about nucleotide 805 to about nucleotide 864; the fragment of SEQ ID NO:14 from about nucleotide 230 to about nucleotide 289; the fragment of SEQ ID NO:15 from about nucleotide 154 to about nucleotide 213; the fragment of SEQ ID NO:16 from about nucleotide 110 to
30 about nucleotide 169; the fragment of SEQ ID NO:17 from about nucleotide 482 to about nucleotide 541; and the fragment of SEQ ID NO:18 from about nucleotide 115 to about nucleotide 174.

The invention also encompasses PKH variants. A preferred PKH variant is one which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% amino acid sequence identity to the PKH amino acid sequence, and which contains at least one functional or
35 structural characteristic of PKH.

The invention also encompasses polynucleotides which encode PKH. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:10-18, which encodes PKH.

The invention also encompasses a variant of a polynucleotide sequence encoding PKH. In particular, such a variant polynucleotide sequence will have at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding PKH. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:10-18 which has at least about 70%, more preferably at least about 85%, and most preferably at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:10-18. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of PKH.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding PKH, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring PKH, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode PKH and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring PKH under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding PKH or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding PKH and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode PKH and PKH derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding PKH or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing

to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:10-18 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and most preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed. In a preferred embodiment, hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 $\mu\text{g/ml}$ denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50 % formamide, and 200 $\mu\text{g/ml}$ ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

The washing steps which follow hybridization can also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include temperature of at least about 25°C, more preferably of at least about 42°C, and most preferably of at least about 68°C. In a preferred embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 42°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a most preferred embodiment, wash steps will occur at 68°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations on these conditions will be readily apparent to those skilled in the art.

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of

polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the Hamilton MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), Peltier Thermal Cycler 200 (PTC200; MJ Research, Watertown MA) and the ABI CATALYST 800 (Perkin-Elmer).

5 Sequencing is then carried out using either ABI 373 or 377 DNA sequencing systems (Perkin-Elmer) or the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA). The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

10 The nucleic acid sequences encoding PKH may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse
15 PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) *PCR Methods Applic.*
20 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids
25 the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

30 When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

35 Capillary electrophoresis systems which are commercially available may be used to analyze the

size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode PKH may be cloned in recombinant DNA molecules that direct expression of PKH, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express PKH.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter PKH-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding PKH may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.) Alternatively, PKH itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A Peptide Synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of PKH, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active PKH, the nucleotide sequences encoding PKH or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the

necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding PKH. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding PKH. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding PKH and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding PKH and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding PKH. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding PKH. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding PKH can be achieved using a multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or pSPORT1 plasmid (Life Technologies). Ligation of sequences encoding PKH into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When

large quantities of PKH are needed, e.g. for the production of antibodies, vectors which direct high level expression of PKH may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of PKH. A number of vectors containing
5 constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

10 Plant systems may also be used for expression of PKH. Transcription of sequences encoding PKH may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science*
15 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where
20 an adenovirus is used as an expression vector, sequences encoding PKH may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses PKH in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus
25 (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or
30 vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of PKH in cell lines is preferred. For example, sequences encoding PKH can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of
35 the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to

selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

5 Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr*^r cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers
10 resistance to the aminoglycosides, neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA
15 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

20 Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding PKH is inserted within a marker gene sequence, transformed cells containing sequences encoding PKH can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding PKH under the control of a single
25 promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding PKH and that express PKH may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay
30 or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of PKH using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated
35 cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies

reactive to two non-interfering epitopes on PKH is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and
5 Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding PKH include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences
10 encoding PKH, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable
15 reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding PKH may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced
20 by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode PKH may be designed to contain signal sequences which direct secretion of PKH through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted
25 sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa,
30 MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding PKH may be ligated to a heterologous sequence resulting in translation of a fusion protein in
35 any of the aforementioned host systems. For example, a chimeric PKH protein containing a heterologous

moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of PKH activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP),
5 thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion
10 protein may also be engineered to contain a proteolytic cleavage site located between the PKH encoding sequence and the heterologous protein sequence, so that PKH may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

15 In a further embodiment of the invention, synthesis of radiolabeled PKH may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract systems (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, preferably ³⁵S-methionine.

20 Fragments of PKH may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A Peptide Synthesizer (Perkin-Elmer). Various fragments of PKH may be synthesized separately and then combined to produce the full length molecule.

25 THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of PKH and various protein kinase homologs. In addition, the expression of PKH is closely associated with cancer, reproductive tissues and hematopoietic/immune tissues. Therefore, PKH appears to play a role in cancer, autoimmune/inflammatory disorders, and reproductive disorders. In the treatment
30 of cancer, autoimmune/inflammatory disorders, and reproductive disorders associated with increased PKH expression or activity, it is desirable to decrease the expression or activity of PKH. In the treatment of the above conditions associated with decreased PKH expression or activity, it is desirable to increase the expression or activity of PKH.

Therefore, in one embodiment, PKH or a fragment or derivative thereof may be administered to a
35 subject to treat or prevent a disorder associated with decreased expression or activity of PKH. Examples

of such disorders include, but are not limited to, a cancer, such as adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.; and a reproductive disorder, such as disorders of prolactin production; infertility, including tubal disease, ovulatory defects, and endometriosis; disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia.

25 In another embodiment, a vector capable of expressing PKH or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PKH including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified PKH in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PKH including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of PKH may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of PKH including, but not limited to, those listed above.

35 In a further embodiment, an antagonist of PKH may be administered to a subject to treat or

prevent a disorder associated with increased expression or activity of PKH. Examples of such disorders include, but are not limited to, those described above. In one aspect, an antibody which specifically binds PKH may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissue which express PKH.

5 In an additional embodiment, a vector expressing the complement of the polynucleotide encoding PKH may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of PKH including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate
10 therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

15 An antagonist of PKH may be produced using methods which are generally known in the art. In particular, purified PKH may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind PKH. Antibodies to PKH may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab
20 expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with PKH or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase
25 immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to PKH
30 have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of PKH amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

35 Monoclonal antibodies to PKH may be prepared using any technique which provides for the

production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce PKH-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

Antibodies may also be produced by inducing *in vivo* production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; Winter, G. et al. (1991) *Nature* 349:293-299.)

Antibody fragments which contain specific binding sites for PKH may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between PKH and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering PKH epitopes is preferred, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for PKH. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of PKH-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a

preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple PKH epitopes, represents the average affinity, or avidity, of the antibodies for PKH. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular PKH epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the PKH-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of PKH, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is preferred for use in procedures requiring precipitation of PKH-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra; Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding PKH, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding PKH may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding PKH. Thus, complementary molecules or fragments may be used to modulate PKH activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding PKH.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding PKH. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

Genes encoding PKH can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding PKH. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating

vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding PKH. Oligonucleotides derived from the transcription initiation site, e.g.,
5 between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and
10 Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme
15 molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding PKH.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU,
20 and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

25 Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding PKH. Such DNA sequences may be incorporated into a wide variety of vectors with suitable
30 RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within
35 the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in

all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of PKH, antibodies to PKH, and mimetics, agonists, antagonists, or inhibitors of PKH. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain

tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and 5 tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene 10 glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit 15 capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous 20 solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic 25 fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

30 The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts 35 tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms.

In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate
5 container and labeled for treatment of an indicated condition. For administration of PKH, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

10 For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

15 A therapeutically effective dose refers to that amount of active ingredient, for example PKH or fragments thereof, antibodies of PKH, and agonists, antagonists or inhibitors of PKH, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose lethal to 50% of the
20 population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the LD₅₀/ED₅₀ ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED₅₀ with little or no toxicity. The dosage varies within this
25 range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of
30 the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1
35 gram, depending upon the route of administration. Guidance as to particular dosages and methods of

delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

5 In another embodiment, antibodies which specifically bind PKH may be used for the diagnosis of disorders characterized by expression of PKH, or in assays to monitor patients being treated with PKH or agonists, antagonists, or inhibitors of PKH. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for PKH include methods which utilize the antibody and a label to detect PKH in human body fluids or in extracts of cells or tissues. The
10 antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring PKH, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of PKH expression. Normal or standard
15 values for PKH expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to PKH under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities of PKH expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject
20 values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding PKH may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and
quantitate gene expression in biopsied tissues in which expression of PKH may be correlated with
25 disease. The diagnostic assay may be used to determine absence, presence, and excess expression of PKH, and to monitor regulation of PKH levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding PKH or closely related molecules may be used to identify nucleic acid sequences which encode PKH. The specificity of the probe, whether it is made from
30 a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding PKH, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at
35 least 50% sequence identity to any of the PKH encoding sequences. The hybridization probes of the

subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:10-18 or from genomic sequences including promoters, enhancers, and introns of the PKH gene.

Means for producing specific hybridization probes for DNAs encoding PKH include the cloning of polynucleotide sequences encoding PKH or PKH derivatives into vectors for the production of mRNA
5 probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes *in vitro* by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ³²P or ³⁵S, or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

10 Polynucleotide sequences encoding PKH may be used for the diagnosis of disorders associated with expression of PKH. Examples of such disorders include, but are not limited to, cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis,
15 prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; autoimmune/inflammatory disorders such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia
20 with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic
25 sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and reproductive disorders including disorders of prolactin production; infertility, including tubal disease, ovulatory defects, and endometriosis; disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome,
30 endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia. The polynucleotide sequences encoding PKH may be used in Southern or northern analysis, dot blot, or
35 other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like

assays; and in microarrays utilizing fluids or tissues from patients to detect altered PKH expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding PKH may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences
5 encoding PKH may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding PKH in the sample indicates the presence of
10 the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of PKH, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment
15 thereof, encoding PKH, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a
20 disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

25 With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further
30 progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding PKH may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding PKH, or a fragment of a polynucleotide complementary to the polynucleotide encoding PKH, and will be employed
35 under optimized conditions for identification of a specific gene or condition. Oligomers may also be

employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of PKH include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard
5 curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the
10 polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

15 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

20 In another embodiment of the invention, nucleic acid sequences encoding PKH may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single
25 chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online
30 Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding PKH on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as
35 linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often

the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, PKH, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between PKH and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate. The test compounds are reacted with PKH, or fragments thereof, and washed. Bound PKH is then detected by methods well known in the art. Purified PKH can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding PKH specifically compete with a test compound for binding PKH. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PKH.

In additional embodiments, the nucleotide sequences which encode PKH may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in

particular U.S. Ser. No. 09/173,581, are hereby expressly incorporated by reference.

EXAMPLES

I. Construction of cDNA Libraries

5 RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or
10 sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A⁺) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Valencia CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue
15 lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERSCRIPT plasmid system (Life Technologies), using the
20 recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6). Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography
25 (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), pSPORT1 plasmid (Life Technologies), or pINCY (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent *E. coli* cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

30 II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision, using the UNIZAP vector system (Stratagene) or cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8
35 Ultra Plasmid purification systems or the R.E.A.L. Prep 96 plasmid kit from QIAGEN. Following

precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) *Anal. Biochem.* 216:1-14). Host cell lysis and thermal cycling steps 5 were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a Fluoroskan II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

10 The cDNAs were prepared for sequencing using the ABI CATALYST 800 (Perkin-Elmer) or the HYDRA microdispenser (Robbins Scientific) or MICROLAB 2200 (Hamilton) systems in combination with the PTC-200 thermal cyclers (MJ Research). The cDNAs were sequenced using the ABI PRISM 373 or 377 sequencing systems (Perkin-Elmer) and standard ABI protocols, base calling software, and kits. In one alternative, cDNAs were sequenced using the MEGABACE 1000 DNA sequencing system 15 (Molecular Dynamics). In another alternative, the cDNAs were amplified and sequenced using the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). In yet another alternative, cDNAs were sequenced using solutions and dyes from Amersham Pharmacia Biotech. Reading frames for the ESTs were determined using standard methods (reviewed in Ausubel, 1997, *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in 20 Example V.

The polynucleotide sequences derived from cDNA, extension, and shotgun sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the software programs, descriptions, references, and threshold parameters used. The first column of Table 5 shows the tools, programs, and algorithms used, 25 the second column provides a brief description thereof, the third column presents the references which are incorporated by reference herein, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the probability the greater the homology). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software 30 (DNASTAR).

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as GenBank primate, rodent, mammalian, vertebrate, and eukaryote 35 databases, and BLOCKS to acquire annotation, using programs based on BLAST, FASTA, and BLIMPS.

The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently
5 analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, PFAM, and Prosite.

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:10-18. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and
10 amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra,
15 ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ database (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or
20 similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to
25 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding PKH occurred. Analysis involved the categorization of cDNA libraries by
30 organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation/trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories.
35 Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of PKH Encoding Polynucleotides

The full length nucleic acid sequence of SEQ ID NO:10-18 was produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other
5 primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

10 Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -
15 mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step
20 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton
25 MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates,
30 digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia
35 Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and

transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulphoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequence of SEQ ID NO:10-18 is used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:10-18 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT-AR film (Eastman Kodak, Rochester NY) is exposed to the blots for several hours, hybridization patterns are compared visually.

VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on

the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized
5 probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well
10 known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470;
15 Shalon, D. et al. (1996) *Genome Res.* 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the PKH-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring PKH. Although use of oligonucleotides
20 comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of PKH. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is
25 designed to prevent ribosomal binding to the PKH-encoding transcript.

IX. Expression of PKH

Expression and purification of PKH is achieved using bacterial or virus-based expression systems. For expression of PKH in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription.
30 Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express PKH upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of PKH in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant *Autographica*
35 *californica* nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential

polyhedrin gene of baculovirus is replaced with cDNA encoding PKH by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or
5 human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, PKH is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-
10 based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from PKH at specifically
15 engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified PKH obtained by these methods can be used directly in the following activity assay.

X. Demonstration of PKH Activity

20 An assay for PKH activity measures the phosphorylation of a substrate in the presence of gamma-labeled ^{32}P -ATP. PKH is incubated with an appropriate substrate and ^{32}P -ATP in a buffered solution. ^{32}P -labeled product is separated from free ^{32}P -ATP by gel electrophoresis or chromatographic procedures, and the incorporated ^{32}P is quantified by phosphoimage analysis or scintillation counter. The amount of ^{32}P detected is proportional to the activity of PKH in this assay. The specific amino acid
25 residue phosphorylated by PKH may be determined by phosphoamino acid analysis of the labeled, hydrolyzed protein.

XI. Functional Assays

PKH function is assessed by expressing the sequences encoding PKH at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector
30 containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT plasmid (Life Technologies) and pCR3.1 plasmid (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μg of recombinant vector are transiently transfected into a human cell line, preferably of endothelial or hematopoietic origin, using either liposome formulations or electroporation. 1-2 μg of an additional plasmid containing sequences encoding a marker
35 protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected

cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP, and to evaluate properties, for example, their
5 apoptotic state. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins
10 as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of PKH on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding PKH and either CD64 or CD64-GFP. CD64 and CD64-GFP
15 are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding PKH and other genes of interest can be analyzed by northern analysis or microarray techniques.

20 XII. Production of PKH Specific Antibodies

PKH substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the PKH amino acid sequence is analyzed using LASERGENE software
25 (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides 15 residues in length are synthesized using an ABI 431A Peptide
30 Synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with
35 radio-iodinated goat anti-rabbit IgG.

XIII. Purification of Naturally Occurring PKH Using Specific Antibodies

Naturally occurring or recombinant PKH is substantially purified by immunoaffinity chromatography using antibodies specific for PKH. An immunoaffinity column is constructed by covalently coupling anti-PKH antibody to an activated chromatographic resin, such as CNBr-activated
5 SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing PKH are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of PKH (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/PKH binding (e.g., a
10 buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and PKH is collected.

XIV. Identification of Molecules Which Interact with PKH

PKH, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (Bolton, A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539). Candidate molecules previously
15 arrayed in the wells of a multi-well plate are incubated with the labeled PKH, washed, and any wells with labeled PKH complex are assayed. Data obtained using different concentrations of PKH are used to calculate values for the number, affinity, and association of PKH with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.
20 Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

Table 1

Amino Acid SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	10	119819	MUSCNOT01	119819H1 (MUSCNOT01), 1476434H1 (CORPNOT02), 1719355H1 (BLADNOT06), 3016364F6 (MUSCNOT07), 3373249F6 (CONNTUT05)
2	11	132750	BMARNOT02	132750H1 (BMARNOT02), 132750X335D1 (BMARNOT02), 287300R1 (EOSIHET02), 1271292F1 (TESTTUT02), 3343321H1 (SPLNNOT09)
3	12	507669	TMLR3DT01	101010R6 (ADRENOT01), 507669H1 (TMLR3DT01), 507669R6 (TMLR3DT01), 697674H1 (SYNORAT03), 851624T1 (NGANNOT01), 1270195F7 (BRAINOT09)
4	13	1439938	THYRNOT03	814363X11 (OVARTUT01), 822246X13 (KERANOT02), 1439938H1 (THYRNOT03), 3365318F6 (PROSBPT02)
5	14	1447427	PLACNOT02	712366R6 (SYNORAT04), 1447427H1 (PLACNOT02), 2179842F6 (SININOT01), 2697460F6 (UTRSNOT12)
6	15	1567782	UTRSNOT05	1567782H1 (UTRSNOT05), 1567786F6 (UTRSNOT05), 2289257X42F1 (BRAINON01), 2839231T6 (DRGLNOT01), SAFB00259F1
7	16	2295842	BRSTNOT05	606173H1 (BRSTTUT01), 865560R1 (BRAITUT03), 1441018F6 (THYRNOT03), 1706132T6 (DUODNOT02), 1811824F6 (PROSTUT12), 2223155H1 (LUNGNOT18), 2295842H1 (BRSTNOT05)
8	17	2605059	LUNGTUT07	381188R6 (HYPONOB01), 381188T6 (HYPONOB01), 691185T6 (LUNGTUT02), 1824201F6 (GBLATUT01), 2605059F6 (LUNGTUT07), 2605059H1 (LUNGTUT07), 2605059X304F1 (LUNGTUT07)
9	18	3000825	TLYMNOT06	2015630F6 (ADRENOT07), 3000825F6 (TLYMNOT06), 3000825H1 (TLYMNOT06)

Table 2

Amino Acid SEQ ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Signature Sequence	Identification	Analytical Methods
1	297	S285 T29 S197 S202 S225	A117 - I295	Muscle-specific serine kinase	PFAM, BLAST, BLOCKS, MOTIFS
2	287	S283 T106 T179 S194 S274 T69 T184 S233 S270	S31 - T284	Protein kinase homolog	PFAM, BLOCKS, PRINTS, BLAST, MOTIFS
3	346	S7 T105 S160 T232 T244 T282 S329 T336 S214 T255 Y231	E125 - D333	Tyrosine kinase	PFAM, BLAST, BLOCKS, PRINTS, PROFILESCAN, MOTIFS
4	90	S60 S9 S62	F28 - R78	Protein kinase homolog	PFAM, BLOCKS, BLAST, MOTIFS
5	327	S96 T210 S277 T40 S137 S179 T273 Y178 Y310	Y11 - N235	Protein kinase homolog	PFAM, BLOCKS, PRINTS, BLAST, MOTIFS
6	345	S327 S23 S41 S48 S123 S219 T319 T166 S175 Y30	Y4 - I226	Serine/threonine- and tyrosine-specific protein kinase, Nek1	PFAM, BLAST, BLOCKS, PRINTS, PROFILESCAN, MOTIFS
7	424	T97 S218 S298 T389 S413 T54 T62 T89 T109 S112 S151 S223 S229 S286 S318	D202 - V412	Protein kinase homolog	PFAM, BLOCKS, PRINTS, BLAST, MOTIFS
8	99			Cell cycle related protein kinase	BLAST
9	138	T91 T24 T57 T14	R72 - V101	Serine/threonine protein kinase	PFAM, BLAST, MOTIFS

Table 3

Nucleotide SEQ ID NO:	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
10	Nervous (0.400) Musculoskeletal (0.200) Cardiovascular (0.100)	Cancer (0.500) Neurological (0.300)	PBLUESCRIPT
11	Reproductive (0.316) Hematopoietic/Immune (0.211) Gastrointestinal (0.158)	Inflammation (0.421) Cancer (0.368)	PBLUESCRIPT
12	Hematopoietic/Immune (0.514) Gastrointestinal (0.189) Reproductive (0.081)	Inflammation (0.595) Cancer (0.243)	PBLUESCRIPT
13	Reproductive (0.375) Developmental (0.125) Endocrine (0.125)	Cancer (0.375) Inflammation (0.250)	pINCY
14	Reproductive (0.346) Nervous (0.269) Hematopoietic/Immune (0.231)	Cancer (0.462) Inflammation (0.385)	pINCY
15	Nervous (0.500) Developmental (0.167) Musculoskeletal (0.167)	Cancer (0.833) Inflammation (0.333)	pINCY
16	Reproductive (0.290) Gastrointestinal (0.145) Nervous (0.130)	Cancer (0.420) Inflammation (0.362)	PSPORT1
17	Nervous (0.250) Gastrointestinal (0.167) Hematopoietic/Immune (0.167)	Cancer (0.500) Inflammation (0.250)	pINCY
18	Endocrine (0.333) Hematopoietic/Immune (0.333) Reproductive (0.333)	Inflammation (0.667) Cancer (0.333)	pINCY

Table 4

Nucleotide SEQ ID NO:	Library	Library Comment
10	MUSCNOT01	Library was constructed at Stratagene (STR937209), using RNA isolated from the skeletal muscle tissue of a patient with malignant hyperthermia.
11	BMARNOT02	Library was constructed using RNA isolated from the bone marrow of 24 male and female Caucasian donors, 16 to 70 years old. (RNA came from Clontech.)
12	TMLR3DT02	Library was constructed using RNA isolated from non-adherent peripheral blood mononuclear cells collected from a pool of male and female donors. Cells from each donor were purified on Ficoll Hypaque, then co-cultured for 72 hours. The cells were pooled, washed once in PBS, lysed in a buffer containing GuSCN, and spun through CsCl to obtain RNA. PolyA RNA was isolated using a Qiagen Oligotex kit.
13	THYRNOT03	Library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
14	PLACNOT02	Library was constructed using RNA isolated from the placental tissue of a Hispanic female fetus, who was prematurely delivered at 21 weeks' gestation. Serologies of the mother's blood were positive for cytomegalovirus.
15	UTRSNOT05	Library was constructed using RNA isolated from the uterine tissue of a 45-year-old Caucasian female during a total abdominal hysterectomy and total colectomy. Pathology for the associated tumor tissue indicated multiple leiomyomas of the myometrium.

Table 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
16	BRSTNOT05	Library was constructed using RNA isolated from breast tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated multicentric invasive grade 4 lobular carcinoma. Family history included breast and prostate cancer.
17	LUNGTUT07	Library was constructed using RNA isolated from lung tumor tissue removed from the upper lobe of a 50-year-old Caucasian male during segmental lung resection. Pathology indicated an invasive grade 4 squamous cell adenocarcinoma. Patient history included tobacco use. Family history included skin cancer.
18	TLYMNOT06	Library was constructed using RNA isolated from activated Th2 cells. These cells were differentiated from umbilical cord CD4 T cells with IL-4 in the presence of anti-IL-12 antibodies and B7-transfected COS cells, and then activated for six hours with anti-CD3 and anti-CD28 antibodies.

TABLE 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLocks IMProved Searcher that matches a sequence against those in BLOCKS and PRINTS databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S and J.G. Henikoff, Nucl. Acid Res., 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and Probability value= 1.0E-3 or less
PFAM	A Hidden Markov Models-based application useful for protein family search.	Krogh, A. et al. (1994) J. Mol. Biol., 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) Nucleic Acids Res. 26:320-322.	Score=10-50 bits, depending on individual protein families

TABLE 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Score= 4.0 or greater
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. A substantially purified polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and fragments thereof.
- 5 2. A substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of claim 1.
3. An isolated and purified polynucleotide encoding the polypeptide of claim 1.
4. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 3.
- 10 5. An isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide of claim 3.
6. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 3.
7. A method for detecting a polynucleotide, the method comprising the steps of:
 - 15 (a) hybridizing the polynucleotide of claim 6 to at least one nucleic acid in a sample, thereby forming a hybridization complex; and
 - (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of the polynucleotide in the sample.
8. The method of claim 7 further comprising amplifying the polynucleotide prior to
20 hybridization.
9. An isolated and purified polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, and fragments thereof.
10. An isolated and purified polynucleotide variant having at least 70% polynucleotide sequence identity to the polynucleotide of claim 9.
- 25 11. An isolated and purified polynucleotide having a sequence which is complementary to the polynucleotide of claim 9.
12. An expression vector comprising at least a fragment of the polynucleotide of claim 3.
13. A host cell comprising the expression vector of claim 12.
14. A method for producing a polypeptide, the method comprising the steps of:
 - 30 a) culturing the host cell of claim 13 under conditions suitable for the expression of the polypeptide; and
 - b) recovering the polypeptide from the host cell culture.
15. A pharmaceutical composition comprising the polypeptide of claim 1 in conjunction with a suitable pharmaceutical carrier.
- 35 16. A purified antibody which specifically binds to the polypeptide of claim 1.

17. A purified agonist of the polypeptide of claim 1.
18. A purified antagonist of the polypeptide of claim 1.
19. A method for treating or preventing a disorder associated with decreased expression or activity of PKH, the method comprising administering to a subject in need of such treatment an effective
5 amount of the pharmaceutical composition of claim 15.
20. A method for treating or preventing a disorder associated with increased expression or activity of PKH, the method comprising administering to a subject in need of such treatment an effective amount of the antagonist of claim 18.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.
 BANDMAN, Olga
 TANG, Y. Tom
 HILLMAN, Jennifer L.
 YUE, Henry
 GUEGLER, Karl J.
 CORLEY, Neil C.
 GORGONE, Gina A.
 AZIMZAI, Yalda
 LU, Dyung Aina M.

<120> PROTEIN KINASE HOMOLOGS

<130> PF-0614 PCT

<140> To Be Assigned

<141> Herewith

<150> 09/173,581

<151> 1998-10-15

<160> 18

<170> PERL Program

<210> 1

<211> 297

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 119819CD1

<400> 1

Met	Arg	Arg	Lys	Arg	Lys	Gln	Gln	Lys	Arg	Leu	Leu	Glu	Glu	Arg
1			5						10					15
Leu	Arg	Asp	Leu	Gln	Arg	Leu	Glu	Ala	Met	Glu	Ala	Ala	Thr	Gln
			20						25					30
Ala	Glu	Asp	Ser	Gly	Leu	Arg	Leu	Asp	Gly	Gly	Ser	Gly	Ser	Thr
			35						40					45
Ser	Ser	Ser	Gly	Cys	His	Pro	Gly	Gly	Ala	Arg	Ala	Gly	Pro	Ser
			50						55					60
Pro	Ala	Ser	Ser	Ser	Pro	Ala	Pro	Gly	Gly	Gly	Arg	Ser	Leu	Ser
			65						70					75
Ala	Gly	Ser	Gln	Thr	Ser	Gly	Phe	Ser	Gly	Ser	Leu	Phe	Ser	Pro
			80						85					90
Ala	Ser	Cys	Ser	Ile	Leu	Ser	Gly	Ser	Ser	Asn	Gln	Arg	Glu	Thr
			95						100					105
Gly	Gly	Leu	Leu	Ser	Pro	Ser	Thr	Pro	Phe	Gly	Ala	Ser	Asn	Leu
			110						115					120
Leu	Val	Asn	Pro	Leu	Glu	Pro	Gln	Asn	Ala	Asp	Lys	Ile	Lys	Ile
			125						130					135

Lys Ile Ala Asp Leu Gly Asn Ala Cys Trp Val His Lys His Phe
 140 145 150
 Thr Glu Asp Ile Gln Thr Arg Gln Tyr Arg Ala Val Glu Val Leu
 155 160 165
 Ile Gly Ala Glu Tyr Gly Pro Pro Ala Asp Ile Trp Ser Thr Ala
 170 175 180
 Cys Met Ala Phe Glu Leu Ala Thr Gly Asp Tyr Leu Phe Glu Pro
 185 190 195
 His Ser Gly Glu Asp Tyr Ser Arg Asp Glu Asp His Ile Ala His
 200 205 210
 Ile Val Glu Leu Leu Gly Asp Ile Pro Pro Ala Phe Ala Leu Ser
 215 220 225
 Gly Arg Tyr Ser Arg Glu Phe Phe Asn Arg Arg Gly Glu Leu Arg
 230 235 240
 His Ile His Asn Leu Lys His Trp Gly Leu Tyr Glu Val Leu Met
 245 250 255
 Glu Lys Tyr Glu Trp Pro Leu Glu Gln Ala Thr Gln Phe Ser Ala
 260 265 270
 Phe Leu Leu Pro Met Asn Glu Tyr Ile Pro Glu Lys Arg Ala Ser
 275 280 285
 Ala Arg Asp Cys Leu Gln His Pro Trp Leu Gln Pro
 290 295

<210> 2
 <211> 287
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 132750CD1

<400> 2
 Met Gln Glu Ile Pro Gln Glu Gln Ile Lys Glu Ile Lys Lys Glu
 1 5 10 15
 Gln Leu Ser Gly Ser Pro Trp Ile Leu Leu Arg Glu Asn Glu Val
 20 25 30
 Ser Thr Leu Tyr Lys Gly Glu Tyr His Arg Ala Pro Val Ala Ile
 35 40 45
 Lys Val Phe Lys Lys Leu Gln Ala Gly Ser Ile Ala Ile Val Arg
 50 55 60
 Gln Thr Phe Asn Lys Glu Ile Lys Thr Met Lys Lys Phe Glu Ser
 65 70 75
 Pro Asn Ile Leu Arg Ile Phe Gly Ile Cys Ile Asp Glu Thr Val
 80 85 90
 Thr Pro Pro Gln Phe Ser Ile Val Met Glu Tyr Cys Glu Leu Gly
 95 100 105
 Thr Leu Arg Glu Leu Leu Asp Arg Glu Lys Asp Leu Thr Leu Gly
 110 115 120
 Lys Arg Met Val Leu Val Leu Gly Ala Ala Arg Gly Leu Tyr Arg
 125 130 135
 Leu His His Ser Glu Ala Pro Glu Leu His Gly Lys Ile Arg Ser
 140 145 150
 Ser Asn Phe Leu Val Thr Gln Gly Tyr Gln Val Lys Leu Ala Gly
 155 160 165

Phe Glu Leu Arg Lys Thr Gln Thr Ser Met Ser Leu Gly Thr Thr
 170 175 180
 Arg Glu Lys Thr Asp Arg Val Lys Ser Thr Ala Tyr Leu Ser Pro
 185 190 195
 Gln Glu Leu Glu Asp Val Phe Tyr Gln Tyr Asp Val Lys Ser Glu
 200 205 210
 Ile Tyr Ser Phe Gly Ile Val Leu Trp Glu Ile Ala Thr Gly Asp
 215 220 225
 Ile Pro Phe Gln Gly Cys Asn Ser Glu Lys Ile Arg Lys Leu Val
 230 235 240
 Ala Val Lys Arg Gln Gln Glu Pro Leu Gly Glu Asp Cys Pro Ser
 245 250 255
 Glu Leu Arg Glu Ile Ile Asp Glu Cys Arg Ala His Asp Pro Ser
 260 265 270
 Val Arg Pro Ser Val Asp Glu Ile Leu Lys Lys Leu Ser Thr Phe
 275 280 285
 Ser Lys

<210> 3
 <211> 346
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 507669CD1

<400> 3
 Met Gly Cys Gly Cys Ser Ser His Pro Glu Asp Asp Trp Met Glu
 1 5 10 15
 Asn Ile Asp Val Cys Glu Asn Cys His Tyr Pro Ile Val Pro Leu
 20 25 30
 Asp Gly Lys Gly Thr Leu Leu Ile Arg Asn Gly Ser Glu Val Arg
 35 40 45
 Asp Pro Leu Val Thr Tyr Glu Gly Ser Asn Pro Pro Ala Ser Pro
 50 55 60
 Leu Gln Asp Asn Leu Val Ile Ala Leu His Ser Tyr Glu Pro Ser
 65 70 75
 His Asp Gly Asp Leu Gly Phe Glu Lys Gly Glu Gln Leu Arg Ile
 80 85 90
 Leu Glu Gln Ser Gly Glu Trp Trp Lys Ala Gln Ser Leu Thr Thr
 95 100 105
 Gly Gln Glu Gly Phe Ile Pro Phe Asn Phe Val Ala Lys Ala Asn
 110 115 120
 Ser Leu Glu Pro Glu Ala Asn Leu Met Lys Gln Leu Gln His Gln
 125 130 135
 Arg Leu Val Arg Leu Tyr Ala Val Val Thr Gln Glu Pro Ile Tyr
 140 145 150
 Ile Ile Thr Glu Tyr Met Glu Asn Gly Ser Leu Val Asp Phe Leu
 155 160 165
 Lys Thr Pro Ser Gly Ile Lys Leu Thr Ile Asn Lys Leu Leu Asp
 170 175 180
 Met Ala Ala Gln Ile Ala Glu Gly Met Ala Phe Ile Glu Glu Arg
 185 190 195
 Asn Tyr Ile His Arg Asp Leu Arg Ala Ala Asn Ile Leu Val Ser

	200		205		210
Asp Thr Leu Ser Cys Lys Ile Ala Asp Phe Gly Leu Ala Arg Leu					
	215		220		225
Ile Glu Asp Asn Glu Tyr Thr Ala Arg Glu Gly Ala Lys Phe Pro					
	230		235		240
Ile Lys Trp Thr Ala Pro Glu Ala Ile Asn Tyr Gly Thr Phe Thr					
	245		250		255
Ile Lys Ser Asp Val Trp Ser Phe Gly Ile Leu Leu Thr Glu Ile					
	260		265		270
Val Thr His Gly Arg Ile Pro Tyr Pro Gly Met Thr Asn Pro Glu					
	275		280		285
Val Ile Gln Asn Leu Glu Arg Gly Tyr Arg Met Val Arg Pro Asp					
	290		295		300
Asn Cys Pro Glu Glu Leu Tyr Gln Leu Met Arg Leu Cys Trp Lys					
	305		310		315
Glu Arg Pro Glu Asp Arg Pro Thr Phe Asp Tyr Leu Arg Ser Val					
	320		325		330
Leu Glu Asp Phe Phe Thr Ala Thr Glu Gly Gln Tyr Gln Pro Gln					
	335		340		345

Pro

<210> 4
 <211> 90
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1439938CD1

<400> 4

Met Pro Ala Gly Gly Arg Ala Gly Ser Leu Lys Asp Pro Asp Val					
1	5		10		15
Ala Glu Leu Phe Phe Lys Asp Asp Pro Glu Lys Leu Phe Ser Asp					
	20		25		30
Leu Arg Glu Ile Gly His Gly Ser Phe Gly Ala Val Tyr Phe Ala					
	35		40		45
Arg Asp Val Arg Asn Ser Glu Val Val Ala Ile Lys Lys Met Ser					
	50		55		60
Tyr Ser Gly Lys Gln Ser Asn Glu Lys Trp Gln Asp Ile Ile Lys					
	65		70		75
Glu Val Arg Arg Arg Arg Arg Val Gly Arg Glu Asp Glu Glu Arg					
	80		85		90

<210> 5
 <211> 327
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1447427CD1

<400> 5

Met Ser Ser Phe Leu Pro Glu Gly Gly Cys Tyr Glu Leu Leu Thr
 1 5 10 15
 Val Ile Gly Lys Gly Phe Glu Asp Leu Met Thr Val Asn Leu Ala
 20 25 30
 Arg Tyr Lys Pro Thr Gly Glu Tyr Val Thr Val Arg Arg Ile Asn
 35 40 45
 Leu Glu Ala Cys Ser Asn Glu Met Val Thr Phe Leu Gln Gly Glu
 50 55 60
 Leu His Val Ser Lys Leu Phe Asn His Pro Asn Ile Val Pro Tyr
 65 70 75
 Arg Ala Thr Phe Ile Ala Asp Asn Glu Leu Trp Val Val Thr Ser
 80 85 90
 Phe Met Ala Tyr Gly Ser Ala Lys Asp Leu Ile Cys Thr His Phe
 95 100 105
 Met Asp Gly Met Asn Glu Leu Ala Ile Ala Tyr Ile Leu Gln Gly
 110 115 120
 Val Leu Lys Ala Leu Asp Tyr Ile His His Met Gly Tyr Val His
 125 130 135
 Arg Ser Val Lys Ala Ser His Ile Leu Ile Ser Val Asp Gly Lys
 140 145 150
 Val Tyr Leu Ser Gly Leu Arg Thr Thr Leu Ser Met Ile Ser His
 155 160 165
 Gly Gln Arg Gln Arg Val Val His Asp Phe Pro Lys Tyr Ser Val
 170 175 180
 Lys Val Leu Pro Trp Leu Ser Pro Glu Val Leu Gln Gln Asn Leu
 185 190 195
 Gln Gly Tyr Asp Ala Lys Ser Asp Ile Tyr Ser Val Gly Ile Thr
 200 205 210
 Ala Cys Glu Leu Ala Asn Gly His Val Pro Phe Lys Asp Met Pro
 215 220 225
 Ala Thr Gln Met Leu Leu Glu Lys Leu Asn Gly Thr Val Pro Cys
 230 235 240
 Leu Leu Asp Thr Ser Thr Ile Pro Ala Glu Glu Leu Thr Met Ser
 245 250 255
 Pro Ser Arg Ser Val Ala Asn Ser Gly Leu Ser Asp Ser Leu Thr
 260 265 270
 Thr Ser Thr Pro Arg Pro Ser Asn Gly Asp Ser Pro Ser His Pro
 275 280 285
 Tyr His Arg Thr Phe Ser Pro His Phe His His Phe Val Glu Gln
 290 295 300
 Cys Leu Gln Arg Asn Pro Asp Ala Arg Tyr Pro Cys Trp Pro Gly
 305 310 315
 Pro Gly Leu Arg Glu Ser Arg Gly Cys Ser Gly Gly
 320 325

<210> 6
 <211> 345
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1567782CD1

<400> 6

Met Glu Lys Tyr Val Arg Leu Gln Lys Ile Gly Glu Gly Ser Phe
 1 5 10 15
 Gly Lys Ala Ile Leu Val Lys Ser Thr Glu Asp Gly Arg Gln Tyr
 20 25 30
 Val Ile Lys Glu Ile Asn Ile Ser Arg Met Ser Ser Lys Glu Arg
 35 40 45
 Glu Glu Ser Arg Arg Glu Val Ala Val Leu Ala Asn Met Lys His
 50 55 60
 Pro Asn Ile Val Gln Tyr Arg Glu Ser Phe Glu Gly Ile Leu Asp
 65 70 75
 Trp Phe Val Gln Ile Cys Leu Ala Leu Lys His Val His Asp Arg
 80 85 90
 Lys Ile Leu His Arg Asp Ile Lys Ser Gln Asn Ile Phe Leu Thr
 95 100 105
 Lys Asp Gly Thr Val Gln Leu Gly Asp Phe Gly Ile Ala Arg Val
 110 115 120
 Leu Asn Ser Thr Val Glu Leu Ala Arg Thr Cys Ile Gly Thr Pro
 125 130 135
 Tyr Tyr Leu Ser Pro Glu Ile Cys Glu Asn Lys Pro Tyr Asn Asn
 140 145 150
 Lys Ser Asp Ile Trp Ala Leu Gly Cys Val Leu Tyr Glu Leu Cys
 155 160 165
 Thr Leu Lys His Ala Phe Glu Ala Gly Ser Met Lys Asn Leu Val
 170 175 180
 Leu Lys Ile Ile Ser Gly Ser Phe Pro Pro Val Ser Leu His Tyr
 185 190 195
 Ser Tyr Asp Leu Arg Ser Leu Val Ser Gln Leu Phe Lys Arg Asn
 200 205 210
 Pro Arg Asp Arg Pro Ser Val Asn Ser Ile Leu Glu Lys Gly Phe
 215 220 225
 Ile Ala Lys Arg Ile Glu Lys Phe Leu Ser Pro Gln Leu Ile Ala
 230 235 240
 Glu Glu Phe Cys Leu Lys Thr Phe Ser Lys Phe Gly Ser Gln Pro
 245 250 255
 Ile Pro Ala Lys Arg Pro Ala Ser Gly Gln Asn Ser Ile Ser Val
 260 265 270
 Met Pro Ala Gln Lys Ile Thr Lys Pro Ala Ala Lys Tyr Gly Ile
 275 280 285
 Pro Leu Ala Tyr Lys Lys Tyr Gly Asp Lys Lys Leu His Glu Lys
 290 295 300
 Lys Pro Leu Gln Lys His Lys Gln Ala His Gln Thr Pro Glu Lys
 305 310 315
 Arg Val Asn Thr Gly Glu Glu Arg Arg Lys Ile Ser Glu Glu Ala
 320 325 330
 Ala Arg Lys Arg Arg Leu Glu Phe Ile Glu Lys Asp Lys Glu Arg
 335 340 345

<210> 7
 <211> 424
 <212> PRT
 <213> Homo sapiens

 <220>
 <221> misc_feature
 <223> Incyte ID No: 2295842CD1

<400> 7
 Met Ile Ser Phe Cys Pro Asp Cys Gly Lys Ser Ile Gln Ala Ala
 1 5 10 15
 Phe Lys Phe Cys Pro Tyr Cys Gly Asn Ser Leu Pro Val Glu Glu
 20 25 30
 His Val Gly Ser Gln Thr Phe Val Asn Pro His Val Ser Ser Phe
 35 40 45
 Gln Gly Ser Gly Ser Arg Pro Pro Thr Pro Lys Ser Ser Pro Gln
 50 55 60
 Lys Thr Arg Lys Ser Pro Gln Val Thr Arg Gly Ser Pro Gln Lys
 65 70 75
 Thr Ser Cys Ser Pro Gln Lys Thr Arg Gln Ser Pro Gln Thr Leu
 80 85 90
 Lys Arg Ser Arg Val Thr Thr Ser Leu Glu Ala Leu Pro Thr Gly
 95 100 105
 Thr Val Leu Thr Asp Lys Ser Gly Arg Gln Trp Lys Leu Lys Ser
 110 115 120
 Phe Gln Thr Arg Asp Asn Gln Gly Ile Leu Tyr Glu Ala Ala Pro
 125 130 135
 Thr Ser Thr Leu Thr Cys Asp Ser Gly Pro Gln Lys Gln Lys Phe
 140 145 150
 Ser Leu Lys Leu Asp Ala Lys Asp Gly Arg Leu Phe Asn Glu Gln
 155 160 165
 Asn Phe Phe Gln Arg Ala Ala Lys Pro Leu Gln Val Asn Lys Trp
 170 175 180
 Lys Lys Leu Tyr Ser Thr Pro Leu Leu Ala Ile Pro Thr Cys Met
 185 190 195
 Gly Phe Gly Val His Gln Asp Lys Tyr Arg Phe Leu Val Leu Pro
 200 205 210
 Ser Leu Gly Arg Ser Leu Gln Ser Ala Leu Asp Val Ser Pro Lys
 215 220 225
 His Val Leu Ser Glu Arg Ser Val Leu Gln Val Ala Cys Arg Leu
 230 235 240
 Leu Asp Ala Leu Glu Phe Leu His Glu Asn Glu Tyr Val His Gly
 245 250 255
 Asn Val Thr Ala Glu Asn Ile Phe Val Asp Pro Glu Asp Gln Ser
 260 265 270
 Gln Val Thr Leu Ala Gly Tyr Gly Phe Ala Phe Arg Tyr Cys Pro
 275 280 285
 Ser Gly Lys His Val Ala Tyr Val Glu Gly Ser Arg Ser Pro His
 290 295 300
 Glu Gly Asp Leu Glu Phe Ile Ser Met Asp Leu His Lys Gly Cys
 305 310 315
 Gly Pro Ser Arg Arg Ser Asp Leu Gln Ser Leu Gly Tyr Cys Met
 320 325 330
 Leu Lys Trp Leu Tyr Gly Phe Leu Pro Trp Thr Asn Cys Leu Pro
 335 340 345
 Asn Thr Glu Asp Ile Met Lys Gln Lys Gln Lys Phe Val Asp Lys
 350 355 360
 Pro Gly Pro Phe Val Gly Pro Cys Gly His Trp Ile Arg Pro Ser
 365 370 375
 Glu Thr Leu Gln Lys Tyr Leu Lys Val Val Met Ala Leu Thr Tyr
 380 385 390
 Glu Glu Lys Pro Pro Tyr Ala Met Leu Arg Asn Asn Leu Glu Ala
 395 400 405
 Leu Leu Gln Asp Leu Arg Val Ser Pro Tyr Asp Pro Ile Gly Leu

410 415 420
 Pro Met Val Pro

<210> 8
 <211> 99
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2605059CD1

<400> 8
 Met Pro Leu Glu Glu Val Leu Pro Asp Val Ser Pro Gln Ala Leu
 1 5 10 15
 Asp Leu Leu Gly Gln Phe Leu Leu Tyr Pro Pro His Gln Arg Ile
 20 25 30
 Ala Ala Ser Lys Ala Leu Leu His Gln Tyr Phe Phe Thr Ala Pro
 35 40 45
 Leu Pro Ala His Pro Ser Glu Leu Pro Ile Pro Gln Arg Leu Gly
 50 55 60
 Gly Pro Ala Pro Lys Ala His Pro Gly Pro Pro His Ile His Asp
 65 70 75
 Phe His Val Asp Arg Pro Leu Glu Glu Ser Leu Leu Asn Ser Glu
 80 85 90
 Leu Ile Arg Pro Phe Ile Leu Glu Gly
 95

<210> 9
 <211> 138
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3000825CD1

<400> 9
 Met Trp Val Val Pro Pro Ile Gly Ala Glu Phe Leu Gly Thr Glu
 1 5 10 15
 Lys Gly Gly Leu Arg Asp Gln Lys Thr Pro Asp Asp His Glu Ala
 20 25 30
 Glu Thr Gly Ile Lys Ser Lys Glu Ala Arg Lys Tyr Ile Phe Asn
 35 40 45
 Cys Leu Asp Ala Cys Val Gln Val Asn Met Thr Thr Asp Leu Glu
 50 55 60
 Gly Ser Asp Met Leu Val Glu Lys Ala Asp Arg Arg Glu Phe Ile
 65 70 75
 Asp Leu Leu Lys Lys Met Leu Thr Ile Asp Ala Asp Lys Arg Ile
 80 85 90
 Thr Pro Ile Glu Thr Leu Asn His Pro Phe Val Thr Met Thr His
 95 100 105
 Leu Leu Asp Phe Pro His Ser Thr His Val Lys Ser Cys Phe Gln
 110 115 120

Asn Met Glu Ile Cys Lys Arg Arg Val Asn Met Tyr Asp Thr Val
 125 130 135
 Asn Gln Ser

<210> 10
 <211> 1427
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 119819CB1

<400> 10
 cggagccaca gtggetccac cccccacctt cacgcactcc cacggtggta atccccaaag 60
 gctgggtggc tgggctgacg gtaattcccc ggggggggtca agtgcccaa actgctcttg 120
 gtgaaaggat gctgtcttcc ccgaatggcc acttccgcct gccttagctt gggctgagag 180
 gggacagaga gcaacctgag gggggccggc caggtcttcc cactcctaata ggagctgtgg 240
 ggagtggggc cacaggcggg gaggcagggg gagtagtgag tagctgggtg caaggggcgc 300
 tggcgccaca ttctgggtgtc catgggagcc ctggggcccg gagaggcctc ttccctggcg 360
 gctgtgcagg gaaacctcca ctcatgctg actggggcgg gcgacaggaa ccctggggtg 420
 acctggctc tgacagcaga ccgtaagct gtccaaaaac aagaggaaga agatgaggcg 480
 caaacggaaa cagcagaagc ggctgctgga ggagcggctg cgggacctgc agaggctgga 540
 ggcatggag gctgccaccc aggctgagga ctctggcttg agactagacg gggcagcgg 600
 ctccacatcc tcttcaggct gtcaccccg gggcgccaga gcaggccct cccagcctc 660
 ttctcccccc gccccagggg gcggcgtag cctcagcgcg ggctcacaga cctcaggctt 720
 ctccggctcc ctcttctctc ctgcctcctg ctccatcctc tccggctcgt ccaatcagcg 780
 agagaccggg ggctcctctg cgcttagcac accattcggg gcctcgaacc tctggtgaa 840
 cccctggag ccccaaatg cagataagat caagatcaag atcgagacc tgggcaacgc 900
 ctgctgggtg cacaagcact tcacggaaga catccagact cggcagtagc gggcctcga 960
 ggtgctgatc ggcgcgcaat acggcccccc ggcagacatc tggagcacag cctgcatggc 1020
 cttcgagctg gccactgggt actacctgtt cgagccgcat tctggagaag actacagtgc 1080
 tgatgaggac cacatcgctc acatagtgga gcttctgggg gacatcccc cagccttcgc 1140
 cctctcaggc cgctattccc gggagttctt caaccggaga ggagagctgc ggcacatcca 1200
 caatctcaag cactggggcc tgtacgaggt actcatggaa aagtacgagt ggcccctaga 1260
 gcaggccaca cagttcagcg ctttctgtct gccatgaat gagtacatcc ccgaaaagcg 1320
 ggccagtgcc cgtgactgcc tccagcacc ctggctccaa ccctagggcc cggctgtggc 1380
 tccacctcca gctctcctg ctttaaggg aaaagcggga cagctcc 1427

<210> 11
 <211> 1586
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 132750CB1

<400> 11
 gctcattgac tcttttgtct tctttctctc cgggggtgag gtcagattta ccacaaaat 60
 gcatgcagga gatcccgcaa gagcaaatca aggagatcaa gaaggagcag ctttcaggat 120
 ccccggtgat tctgctaagg gaaaatgaag tcagcacact ttataaagga gaataccaca 180
 gagctccagt ggccataaaa gtattcaaaa aactccaggc tggcagcatt gcaatagtga 240
 ggcagacttt caataaggag atcaaaaacca tgaagaaatt cgaatctccc aacatcctgc 300
 gtatatttgg gatttgcatt gatgaaacag tgactccgcc tcaattctcc attgtcatgg 360

```

agtactgtga actcgggacc ctgagggagc tgttggatag ggaaaaagac ctcacacttg 420
gcaagcgcac ggtcctagtc ctggggggcag cccgaggcct ataccggcta caccattcag 480
aagcacctga actccacgga aaaatcagaa gctcaaactt cctggtaact caaggctacc 540
aagtgaagct tgcaggattt gagttgagga aaacacagac ttccatgagt ttgggaacta 600
cgagagaaaa gacagacaga gtcaaatcta cagcatatct ctcacctcag gaactggaag 660
atgtatttta tcaatatgat gtaaagtctg aaatatacag ctttggaaatc gtcctctggg 720
aaatcggcac tggagatatc ccgtttcaag gctgtaattc tgagaagatc cgcaagctgg 780
tggctgtgaa gcggcagcag gagccactgg gtgaagactg cccttcagag ctgicgggaga 840
tcattgatga gtgccgggccc catgatccct ctgtgicggcc ctctgtggat gaaatcttaa 900
agaaactctc caccttttct aagtagtgta tcaaaatcta aaccaaggag tctctggaca 960
agaagctggg agaggcacia actggacatc tctctctctc atatccttcg gcattgggtt 1020
atctatggga gcaaggagtg ggcacgcttc tctgttacia atagaaaacg attccagtca 1080
tacaggacac atcccactcc aatgatatt tccaaaaaca tacctctgac agtaactttg 1140
atagatgggt tgtcaaatgt atctttctgg gtatccacac ctcttgcaa tgaaatttgc 1200
agctcctccc ttccataaat gaagtctctt tccccacct ttgaaatctgg gctggcactg 1260
tgacttgatt tgatcaatag aatgtggaag aagtgactgt atgccagttc caagcctagg 1320
tttcaagagg cttataaat gtctgttggg accttaccga gccatgaaca tgttgagtga 1380
gcatgctgga gaatgagaga ccacatgaag cagaaacatg ctttcttagc tgaagtcata 1440
ctagcccaac caacatggca gctaacacat gaatgaggcc aatcaagacc agaagaacca 1500
ctcaagcaga tcccagccca aattgcccac tcacacaatc aggagctaaa taaattactg 1560
ttgtcttaac actaaaaaaaa aaaaaa

```

<210> 12

<211> 1574

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 507669CB1

<400> 12

```

cgacggcgaa gggagctgag actgtccagg cagccagggt aggccaggag gaccatgtga 60
atggggccag aaggctcccg ggctgggagc ggaccatggg ctgtggctgc agctcacacc 120
cggaagatga ctggatggaa aacatcgatg tgtgtgagaa ctgccattat cccatagtcc 180
cactggatgg caagggcacg ctgctcatcc gaaatggctc tgagggtgagg gaccactgag 240
ttacctacga aggtcccaat ccgcccggct ccccactgca agacaacctg gttatcgtc 300
tgacacgcta tgagccctct cacgacggag atctgggctt tgagaagggg gaacagctcc 360
gcatcctgga gcagagcggc gagtgggtgga aggcgcagtc cctgaccacg ggccagggaag 420
gcttcatccc cttcaatttt gtggccaaaag cgaacagcct ggagcccagc gccaacctca 480
tgaagcagct gcaacaccag cggctgggtc ggctctacgc tgtggtcacc caggagccca 540
tctacatcat cactgaatac atggagaatg ggagtctagt ggattttctc aagaccctt 600
caggcatcaa gttgaccatc acaaaactcc tggacatggc agcccaaatt gcagaaggca 660
tggcattcat tgaagagcgg aattatattc atcgtgacct tggggctgcc aacattctgg 720
tgtctgacac cctgagctgc aagattgcag actttggcct agcacgcctc attgaggaca 780
acgagtacac agccagggag ggggccaagt tcccattaa gtggacagcg ccagaagcca 840
ttaactacgg gacattcacc atcaagtcag atgtgtggtc ttttgggatc ctgctgacgg 900
aaattgtcac ccacggccgc atcccctacc cagggatgac caaccgggag gtgattcaga 960
acctggagcg aggctaccgc atgggtgcgc ctgacaactg tccagaggag ctgtaccaac 1020
tcatgaggct gtgctggaag gagcgcaccg aggaccggcc cacctttgac tacctgcgca 1080
gtgtgctgga ggaactcttc acggccacag agggccagta ccagcctcag ccttgagagg 1140
ccttgagagg ccctgggggt ctcccccttt ctctccagcc tgacttgggg agatggagtt 1200
cttgtgccat agtcacatgg cctatgcaca tatggactct gcacatgaat cccaccaca 1260
tgtgacacat atgcaccttg tgtctgtaca cgtgtcctgt agttgcgtgg actctgcaca 1320
tgtcttgtag atgtgtagcc tgtgcatgta tgtcttgagc actgtacaag gtacccttt 1380
ctggctctcc catttcttga gaccacagag agaggggaga agcctgggat tgacagaagc 1440

```

ttctgcccac ctacttttct ttctctcagat catccagaag ttcttcaagg gccaggactt 1500
 tatctaatac ctctgtgtgc tcctccttgg tgccctggcct ggcacacatc aggagttcaa 1560
 taaatgtctg ttga 1574

<210> 13
 <211> 1866
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1439938CB1

<400> 13
 cgggaggaag agggagaggg agaccgggac gagaccgggg ctgtggtgcg gagagaggct 60
 gagacggaga agaggagagg cagagagggc gcggggaccg tcagcagcac cttagctaca 120
 atcgttcagc tattctcggga agagagaagg gagagggagg aggccggggc gggagtgggg 180
 gctgtcacc cgggacccc gcgtgagagg ggcctgtcgg ccggacgtcc tcgggggtggg 240
 cccccagtcg gtggccgaag acctacagct caggccccctg ggtcccaaat ttccaggctt 300
 tgccccctct cttttctcag ataccgggtt aacagtcctc atagtccaga tatccgggac 360
 tcgggtccca acctctctaa acctgggtct ctgtttcata gaatttcaa tatcagggtt 420
 aggcccttgc gtgcaccagt atccgggggtt cattcccccg gcgttcagat atcggattca 480
 gtctccatcc cgttcagata ttcgggggtt agaccccaca atcagaaatc cgaattcgg 540
 cagctgtcgc cctcgacgag ggggaggact ggaccgag gtcagattag gttgtcacc 600
 cctcccctcc aggggaggct tcccgggccc gccctcagg aagggcgaaa gccgaggaag 660
 aggtggcaag gggaaaggct tccttgcccc tctccctgct tggcagagcc gctggaggac 720
 cccaggcgga agcggaggcg ctggggcacc atagtgacct ctaccaggcc aggccccact 780
 ctcaggggcc ccaggggcca ccatgccagc tgggggcccg gccgggagcc tgaaggacct 840
 agatgtggct gagctcttct tcaaggatga cccagaaaag ctcttctctg acctccggga 900
 aattggccat ggcagctttg gagccgtata ctttgcccgg gatgtccgga atagtggagt 960
 ggtggccatc aagaagatgt cctacagtgg gaagcagtcc aatgagaaat ggcaagacat 1020
 catcaaggag gtgcgggagc gaaggagagt agggaggag gatgaagaga gataaggggg 1080
 agaaaagaga ggggcatgag agtggagcgg agctaagaag gggtagaaga gagagtgggt 1140
 gaaggggaag agacgtagag aaagtgtgga gagaggaaag gcatagcgag agaacgagg 1200
 agagagaagt ggaaggggga agtaagagag gataagagga acgagaggag ggaaggggtg 1260
 gggacgagaa cgaagagcat gatggagagg aaagatagag aagagaggaa gtggaggcag 1320
 ttagggggca tggaggagag agagagatga gggagagtgg gagcacgggg cggatggacg 1380
 ggggtggagaa gaagagaggg aggagatgag aggggaaga ggtgggagaa ccgagcgagg 1440
 gaaaagatgg aggaggcagt agagaggggtg tgcaaggggt gaaaagaaa agaagggaaa 1500
 aggatggagg gagtgaaggt aggagacgag gaggaggat gggagagaaat ggagggtagc 1560
 gtgtggatgg tgagtggtag agaatagtga gatggtgaga agcggagaaa ggcagcagag 1620
 gatgggggtg aagcgggaag caaagacaat aggggatgga ggaggagagg agcaggagga 1680
 agacgaagag cgaagggtt gaaagagggga gaagagagta gtaaggggta ggtatgtaga 1740
 tgcgagtagg agaggaagag aaggaatgaa tgagagagag tagagagtag agagagaacg 1800
 aaggaacggg gcagagggag aggaaggaca gaaggagaag agaacaatcg aagaatgaga 1860
 gtgttt 1866

<210> 14
 <211> 1498
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 1350, 1355, 1372, 1444
 <223> a or g or c or t, unknown, or other

<220>
 <221> misc_feature
 <223> Incyte ID No: 1447427CB1

<400> 14
 ctcccctccc agcaaccggt ctggcgggcg cgcggcagta aaactgagga ggcggagcaa 60
 gacggtcggg gctgcttgct aactccagga acaggtttaa gtttttgaaa ctgaagtagg 120
 tctacacagt aggaactcat gtcatttctt gtaagtaaac ccagagcgaa tccaggacca 180
 atgatgagag ctcagagtca atagcatcct tctctaaaca ggaggtcag agtagctttc 240
 tgccagaggg aggggtgttac gagctgctca ctgtgatagg caaaggattt gaggacctga 300
 tgactgtgaa tctagcaagg tacaaccaa caggagagta cgtgactgta cggaggatta 360
 acctagaagc ttgttccaat gagatggtaa cattcttgca gggcgagctg catgtctcca 420
 aactcttcaa ccatcccaat atcgtgccat atcgagccac ttttattgca gacaatgagc 480
 tgtgggttgt cacatcattc atggcatacg gttctgcaaa agatctcatc tgtacacact 540
 tcatggatgg catgaatgag ctggcgattg cttacatcct gcaggggggtg ctgaaggccc 600
 tcgactacat ccaccacatg ggatatgtac acaggagtgt caaagccagc cacatcctga 660
 tctctgtgga tgggaagggtc tacctgtctg gtttgccgac aacgctcagc atgataagcc 720
 atgggcagcg gcagcgagtg gtccacgatt tcccaagta cagtgtcaag gttctgccgt 780
 ggctcagccc cgaggtcctc cagcagaatc tccagggtta tgatgccaag tctgacatct 840
 acagtgtggg aatcacagcc tgtgaactgg ccaacggcca tgtccccttt aaggatatgc 900
 ctgccacca gatgctgcta gagaaactga acggcacagt gccctgcctg ttggatacca 960
 gcaccatccc cgctgaggag ctgacctga gcccttcgcg ctcagtggcc aactctggcc 1020
 tgagtgcag cctgaccacc agcacccccc ggccctccaa cggtgactcg cctcccacc 1080
 cctaccaccg aaccttctcc ccccacttcc accactttgt ggagcagtgc cttcagcgca 1140
 acccgatgc caggtatccc tgctggcctg ggccctgggtc tcgggagagc agaggggtgt 1200
 caggagggta aggccagggt gtgaaggac ttacctcca aaggttctgc aggggaatct 1260
 ggagctacac acaggagggg tcagctcctg ggtgtgtcag aggccagcct ggggagctct 1320
 ggccactgct tcccatgagc tgagggagan ggagnaggga cccgagcctg angcataagt 1380
 ggcaggattt tcggaagctg gggacacggc agtgatgctg cggctctctc ctcccctacc 1440
 tcangctcag tgcagcacc tctgaacact ctttctcagc agtatcgtag ccttcggt 1498

<210> 15
 <211> 1846
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1567782CB1

<400> 15
 taggaattcg tcgaccacg cgatccgccc tcagaagact gccacaccta gactgatgct 60
 tattagtcac caccggtatt cctactaacg tcctgtgtca ctgagttttt taaatgtcta 120
 gcatatctgt aaagatgcct tagaaaaaga atcatggaga agtatgttag actacagaag 180
 attggagaag gttcatttgg aaaagccatt cttgttaaata ctacagaaga tggcagacag 240
 tatgttatca aggaaattaa catctcaaga atgtccagta aagaaagaga agaatcaagg 300
 agagaagttg cagtattggc aaacatgaag catccaaata ttgtccagta tagagaatca 360
 tttgaaggaa ttttggactg gtttgtacag atatgtttgg ccctgaaaca tgtacatgat 420
 agaaaaattc ttcacgaga cattaatct cagaacatat ttttaactaa agatggaaca 480
 gtacaacttg gagatttttg aattgctaga gttcttaata gtactgtaga gctggctcga 540
 acttgcatag ggaccccata ctacttgtca cctgaaatct gtgaaaaca accttacaat 600
 aataaaagt acatttgggc tctgggggtg gtcctttatg agctgtgtac acttaaacat 660
 gcttttgaag ctggcagtat gaaaaacctg gtactgaaga taatatctgg atcttttcca 720
 cctgtgtctt tgcattattc ctatgatctc cgcagtttgg tgtctcagtt atttaaaaga 780
 aatcctaggg atagaccatc agtcaactcc atattggaga aaggttttat agccaaaccg 840
 attgaaaagt ttctctctcc tcagcttatt gcagaagaat tttgtctaaa aacattttcg 900

```

aagtttgat cacagcctat accagctaaa agaccagctt caggacaaaa ctcgatttct 960
ggtatgcttg ctcagaaaaat tacaaaagcct gccgctaaat atggaatacc tttagcatat 1020
aagaaatag gagataaaaa attacacgaa aagaaaccac tgcaaaaaaca taaacaggcc 1080
catcaaactc cagagaagag agtgaatact ggagaagaaa ggaggaaaaat atctgaggaa 1140
gcagcaagaa agagaaggct ggaatattt gaaaaagata aggaacggta ggatcagatt 1200
attagtttaa tgaaggctga acaaatgaaa aggcaagaca aggaagggtt ggaaagaata 1260
aataggcca gggacaagg atggagaaat gtgctaagtg ctggtggaag tgggtaagta 1320
aaggtaggca ttttatacca atatggttat actaccattt tcccctccag tccaccttg 1380
ttctataaaa tgcattgact tgggattttc tttctttctt tagtgtacaa ttaattttta 1440
cctagaattc ttttaacattt attatgaata cttagctttc ctgcatgtat ctgatatgta 1500
acttggttg ctgttatgtg actatactca aaattgcttt aaaagttttt tgtgaagact 1560
atgataacat tattcctgtc aggaatattt aaaaattatg tacaattcat gacactgcag 1620
cctaaaatcg ttctgtaatt tcatgtagcc ttgaagatta agttctcaga agatgcttct 1680
taaatccgat ccctgttgct tctccaattt catcaccatt cattccccta ccacatactg 1740
ggaaggcct attccatggc ggaatgaag ggccataatt tgtaggtttt ccattaccaa 1800
taatgggggg ttggcccaaa tcctactttt gggcctttgg aacctt 1846

```

```

<210> 16
<211> 1721
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc_feature
<223> Incyte ID No: 2295842CB1

```

```

<400> 16
agttggacga ggctcagtga aagttttcgc tgggcaactg agaaggcgc tgcaaatg 60
gagtttccaa cccagtaaat ccaaggcca gaccgtgacc tcataaagca tgatctcctt 120
ctgtccagac tgtggcaaaa gtatccaagc ggcatccaaa ttctgcccct actgtggaaa 180
ttctttgcct gtagaggagc atgtagggtc ccagaccttt gtcaatccac atgtgtcatc 240
cttccaaggc tccgggagca gacccccaac ccccaaaagc agccctcaga agaccaggaa 300
gagccctcag gtgaccaggg gtagccctca gaagaccagc tgtagccctc agaagaccag 360
gcagagccct cagacgctga agcggagccg agtgaccacc tcacttgaag ctttgcccac 420
agggacagtg ctgacagaca agagtggcg acagtggaag ctgaagtcct tccagaccag 480
ggacaaccag ggcattctct atgaagctgc acccacctcc accctcacct gtgactcagg 540
accacagaag caaaagttct cactcaaact ggatgccaaag gatgggcgct tgttcaatga 600
gcagaacttc ttccagcggg ccgccaaagc tctgcaagtc aacaagtgga agaagctgta 660
ctcgaccca ctgctggcca tccctacctg catgggtttc ggtgttcacc aggacaaata 720
caggttcttg gtgttaccca gcctggggag gagccttcag tccgcccctgg atgtcagccc 780
aaagcatgtg ctgtcagaga ggtctgtgct gcagggtggc tgccggctgc tggatgcct 840
ggagttcctc catgagaatg agtatgttca tggaaatgtg acagctgaaa atatctttgt 900
ggatccagag gaccagagtc aggtgacttt ggcaggctat ggcttcgctt tccgctattg 960
cccaagtggc aaacacgtgg cctacgtgga aggcagcagg agccctcacg agggggacct 1020
tgagttcatt agcatggacc tgcacaaggg atgcggggcc tcccgcgca gcgacctcca 1080
gagcctgggc tactgcatgc tgaagtggct ctacgggttt ctgccatgga caaattgcct 1140
tcccacact gaggacatca tgaagcaaaa acagaagttt gttgataagc cggggccctt 1200
cgtgggacct tgcggtcact ggatcaggcc ctccagagacc ctgcagaagt acctgaagg 1260
ggtgatggcc ctccagctat aggagaagcc gccctacgcc atgctgagga acaacctaga 1320
agctttgctg caggatctgc gtgtgtctcc atatgacccc attggcctcc cgatggtgcc 1380
ctaggtgaa tccagaactt tccatttgca gtgtgcaaca gaaaaaaaaa aatgaagtaa 1440
tgtgactcaa ggctgctgt ttaatcacag ataagcttct agaacaagcc ctggaatgtg 1500
cattcctgcc actggtttca ggatactcat cagtcctgat tagcctcccg gaggggccca 1560
gtttccctcc cgtgaatgtg aagttccca tcttggtggc ctgcccttca gccagtgtcc 1620
tagcaaagct ggatggggtt gggcggccc acagggggga cccctctac ccttgacacc 1680
tctgtgcttt ggtaataaat tgttttacca gaaaaaaaaa a 1721

```

<210> 17
 <211> 1985
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 2605059CB1

<400> 17
 ttgcacatctt gggcacccca aaccctcaag tctggccggt ttgtaggggc ccttgggtgag 60
 gtgggtgtgg ggcaggttta ctccactccc aacagcaagt aaccactccc tcccctgaac 120
 cttctctctc ctggcccca ccccccttga tggacagggg cactgtcctt ggcccaactc 180
 agggcttctt ccttcctgct gtcatttggg ttggggtaga tctgtcctt tgtecccttt 240
 caccctagta cacacatgtg cagtgtctca gcaagctgtg cacagagtgc tcatctgaga 300
 gggcaagggg atggatgaag gaatacaggg gtgggtgagt gaatgaatga tgggtcaggg 360
 agacacatgg atgggagagc acccccatg tgagtgtgtg ttaggggctg agagttgaca 420
 gcagagagca tggcaagggg cgggaactac tctcattgta cctgttctt tctccctggc 480
 ccaggagctc actgagctgc cggactaaa caagatctcc ttaaggagc aggtgcccat 540
 gccctgggag gaggtgctgc ctgacgtctc tcccaggca ttggatctgc tgggtcaatt 600
 cttctctac cctcctcacc agegcatcgc agcttccaag gctctcctcc atcagtactt 660
 cttcacagct cccctgcctg cccatccatc tgagctgccg attcctcagc gtctaggggg 720
 acctgcccc aaggcccatc cagggcccc ccacatccat gacttccacg tggaccggcc 780
 tcttgaggag tcgctgttga actcagagct gattcggccc ttcactcctgg aggggtgaga 840
 agttggcctt ggtcccgtct gcctgtcctt caggaccact cagtccacct gttcctctgc 900
 cacctgcctg gcttcaccct ccaaggcctc cccatggcca cagtgggcc acaccacacc 960
 ctgcccctta gcccttgca gggttggtct cgaggcagag gtcattgtcc cagccaagag 1020
 tatgagaaca tccagtgcag cagaggagat tcatggcctg tgctcgggta gccttacctt 1080
 ctgtgtgcta ctgacgtacc catcaggaca gtgagctctg ctgccagtca aggcctgcat 1140
 atgcagaatg acgatgcctg ccttgggtgct gcttccccga gtgctgctc ctgggtcaagg 1200
 agaagtgcag agagtaaggt gtccttatgt tggaaactca agtgggaagga agatttgggt 1260
 tgggtttatt ctccagagcca ttaaacacta gttcagtatg tgagatatag attctaaaaa 1320
 cctcaggtgg ctctgcctta tgtctgttcc tcttcattt ctctcaaggg aaatggctaa 1380
 ggtggcattg tctcatggct ctctgttttg gggctatggg gagggtagca ccaggcatag 1440
 ccacttttgc cctgagggac tctgtgtgct ttcacatcac tgagcactca ttagaagtgc 1500
 agggagacag aagtctaggg ccagggatgg ctccagttgg ggatccagca ggagaccctc 1560
 tgcacatgag gctggtttac caacatctac tccctcagga tgagcgtgag ccagaagcag 1620
 ctgtgtatct aaggaaacaa gcgttcctgg aattaattta taaatttaaat aaatcccaat 1680
 ataatacccag ctagtgtctt ttccctatta taatttgata aggtgattat aaaagataca 1740
 tgggaaggaag tgggaaccaga tgcagaagag gaaatgatgg aaggacttat ggtatcagat 1800
 accaatatctt aaaagtttgt ataataataa agagtatgat tgtggttcaa ggataaaaaac 1860
 agacttagaga aacttattct tagccatcct ttatttttat tttattttatt ttttgatgga 1920
 gtcttgctct gttgccact gcaattcaag ccttgggtgac agactctggt ctcaaaaaaa 1980
 aaaaa 1985

<210> 18
 <211> 661
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 3000825CB1

<400> 18
 tgaggagtga tgaagctgc atttcaactt aactgatgaa agcaggagca gtttacatcc 60

tgtcattcag atatatttgc aggtcccagc agcagccctc tccccttctt ggggcacagc 120
ccctctctgc ctttcctgca gagagaaaag ccacatcctg tgggcaatga caacatgtgg 180
gtgggtgcctc ccataggggc agagtccctg ggaactgaga aagggggctt gagagatcag 240
aagacaccag atgaccatga agcagagaca gggattaagt caaaagaagc aagaaagtac 300
atthttcaact gtttagatgc ttgcgtccag gtgaacatga cgacagattt ggaagggagc 360
gacatggttg tagaaaaggc tgaccggcgg gagttcattg acctggtgaa gaagatgctg 420
accattgatg ctgacaagag aatcactcca atcgaaacct tgaaccatcc ctttgtcacc 480
atgacacact tactcgattt tccccacagc acacacgtca aatcatgttt ccagaacatg 540
gagatctgca agcgtcgggt gaatatgtat gacacgggtga accagagcta aacctagccc 600
caaaccctc tgccgaatat cctcgtcga gggccaaatt ccctatagtg gtcgtattac 660
g 661