

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 199934426 B2
(10) Patent No. 742013

(54) Title
Composition comprising ketanserin and L-carnitine or an alkanoyl L-carnitine for the treatment of CRPS

(51)⁶ International Patent Classification(s)
A61K 031/00

(21) Application No: **199934426** (22) Application Date: **1999 .04 .08**

(87) WIPO No: **WO99/52517**

(30) Priority Data

(31) Number	(32) Date	(33) Country
MI98A0774	1998 .04 .10	IT

(43) Publication Date : **1999 .11 .01**

(43) Publication Journal Date : **2000 .01 .06**

(44) Accepted Journal Date : **2001 .12 .13**

(71) Applicant(s)
Sigma-Tau Industrie Farmaceutiche Riunite S.p.A.

(72) Inventor(s)
Claudio Cavazza; Menotti Calvani

(74) Agent/Attorney
LORD and COMPANY, 4 Douro Place, WEST PERTH WA 6005

34426/99

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00	A2	(11) International Publication Number: WO 99/52517 (43) International Publication Date: 21 October 1999 (21.10.99)
(21) International Application Number: PCT/IT99/00084 (22) International Filing Date: 8 April 1999 (08.04.99) (30) Priority Data: MI98A000774 10 April 1998 (10.04.98) IT (71) Applicant (for all designated States except US): SIGMA-TAU INDUSTRIE FARMACEUTICHE RIUNITE S.P.A. [IT/IT]; Viale Shakespeare, 47, I-00144 Rome (IT). (72) Inventors; and (75) Inventors/Applicants (for US only): CAVAZZA, Claudio [IT/IT]; Piazza Campitelli, 2, I-00186 Rome (IT). CALVANI, Menotti [IT/IT]; Via Longanesi, 14, I-00146 Rome (IT). (74) Agent: SPADARO, Marco; Sigma-Tau Industrie Farmaceutiche Riunite S.p.A., Viale Shakespeare, 47, I-00144 Rome (IT).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published Without international search report and to be republished upon receipt of that report. IP AUSTRALIA 1 NOV 1999 RECEIVED
(54) Title: COMPOSITION COMPRISING KETANSERIN AND L-CARNITINE OR AN ALKANOYL L-CARNITINE FOR THE TREATMENT OF CRPS (57) Abstract The use of ketanserin in combination with L-carnitine, some alkanoyl L-carnitines and the pharmacologically acceptable salts thereof is disclosed for the treatment of CRPS.		

**Composition comprising ketanserin and L-carnitine or an alkanoyl
L-carnitine for the treatment of CRPS**

Pharmaceutical composition for the treatment of Chronic
Regional Pain Syndrome

5 The present invention relates to L-carnitine and lower alkanoyl
L-carnitines or the pharmacologically acceptable salts thereof in
combination with ketanserin, for the therapeutic treatment of Chronic
Regional Pain Syndrome (CRPS).

CRPS is a pain syndrome, which affects several subjects after a
10 trauma, even a mild entity trauma.

To CRPS are associated disturbs of blood circulation with
ischemia and pain, wherein, ischemia gives pain and pain causes
ischemia.

Other important symptoms shown by patients affected by CRPS
15 are hyperpathia and allodinia.

CRPS origin is still not clear; for several years it has been
considered that the sympathetic nervous system is involved.

For this reason the most effective therapy was effected by
blocking the sympathetic innervation with phenol, thermolesion or
20 treating with guanethidine.

The circulation restoration with ketanserin, a serotonin
antagonist, constituted an improvement in the treatment of such
disease.

With this drug it is possible to treat most of symptoms, but not
25 those caused by hyperpathia and allodinia [A. Moesker et al., in


./.

Konservative Therapie Arterieller Durchblutungsstörungen, Georg Thieme Verlag (Stuttgart, New York), 148-152, 1986; A. Moesker et al., The pain Clinic vol. 8, n°1, 31-37 (1995); A. Moesker et al., ibid.vol 12, 269-302 (1991)].

5 More recently, following to the identification of the oxygen free radical during CRPS, dimethyl sulfoxide has been utilised (R. J. A. Goris et al. Free Rad. Res. Comm. 1987, 13-18; W. W. Zuurmond et al., Acta Anaesthesiol. Scan.) without any therapeutic result for the symptoms caused by hyperpathia and allodinia.

10 The therapeutic treatment with dimethyl sulfoxide proved to be efficacious only during acute CRPS.

Ketanserin is a well known synthetic drug (The Merck Index 11th Ed., pag. 834), having formula



and is a specific S₂-receptor antagonist with hypotensive properties, first described in EP application n°13.612.

Ketanserin has been utilised for evaluating its effect on central
20 haemodynamics and coronary circulation [J. Cardiovasc Pharmacol 1998 Dec; 32(6):983-7]; and
in the treatment of intermittent claudicatio [J. De Cree et al., Lancet 2, 775 (1984)].

./.

L-carnitine and alkanoyl L-carnitines are well known compounds. US Patent 4,255,449 and US Patent 4,268,524 describe the use of L-carnitine and alkanoyl L-carnitines, respectively, for normalising abnormally high ratios of low-density lipoproteins (LDL) +
5 very low- density lipoproteins (VLDL) to high-density lipoproteins (HDL), which constitute an etiological factor in various cardiovascular diseases. Through beta-oxidation of fatty acids, L-carnitine is capable of preventing their accumulation and of supplying the cell energy requirement (Bremner Y, *TIBS* 2, 207, 1977) via modulation of extra-
10 and intra-mitochondrial CoA.

L-carnitine and particularly propionyl L-carnitine or acetyl L-carnitine can act by varying the lipid substrate from which the various vasoconstrictor and aggregation-promoting factors derive as a result of the effects of cyclo-oxygenase and lipo-oxygenase, by reducing their
15 formation and by promoting the synthesis of antiaggregant and vasodilators factors.

Carnitine contain a single centre of asymmetry and therefore may exist as two enantiomers, designated D(+)-carnitine and L(-)-carnitine and, obviously in form of racemate. Of these only L(-)-
20 carnitine is found in living organism, were it functions as a vehicle for transporting fatty acids across mitochondrial membranes. Moreover, L-carnitine may be in form of inner salt or in form of pharmacologically acceptable salt.

./.

For the sake of simplicity in the following reference will be made only to L-carnitine or alkanoyl L-carnitine, it should be understood that the compositions described herein apply to L-carnitine or alkanoyl L-carnitines inner salt, or pharmacological acceptable salts thereof.

5 To date, the combined use of L-carnitine and ketanserin is not known for any therapeutic indication.

It has now unexpectedly been found that the co-ordinated use, a term which will be precisely defined here below, of L-carnitine or of an alkanoyl L-carnitine in which the linear or branched-chain alkanoyl
10 has 2-6 carbon atoms, or one of their pharmacologically acceptable salts, in combination with ketanserin show a potent synergistic effect in the treatment of CRPS.

With the composition of the invention patients are able either to recover from blood circulation symptoms or from hyperpathia and
15 allodinia symptoms.

This pharmacological activity is very important because for the first time it is possible to cure these symptoms in patients affected by CRPS.

The well known lack of toxic and side effects of L-carnitine or of
20 the alkanoyl L-carnitines and ketanserin makes their co-ordinated use, according to the invention, particularly useful and safe for the treatment of CRPS.

./.

In the context of the invention described herein, what is meant by "co-ordinated use" of the afore-mentioned compounds is either their co-administration, i.e. the substantially simultaneous administration of L-carnitine or one of the alkanoyl L-carnitines, or one of their
5 pharmacologically acceptable salts, and ketanserin or, indifferently, the administration of a composition containing a combination or mixture of the aforesaid active ingredients, in addition to any excipient included.

The scope of the present invention therefore encompasses both
10 the co-administration of L-carnitine or of an alkanoyl L-carnitine, or one of their pharmacologically acceptable salts, together with ketanserin and pharmaceutical compositions, which can be administered orally, parenterally or per intravenous infusion, containing a mixture of the two active ingredients.

15 In a preferred embodiment of the invention, the alkanoyl L-carnitine will be selected from the group consisting of acetyl, propionyl, butyryl, valeryl and isovaleryl L-carnitine or one of their pharmacologically acceptable salts.

What is meant by pharmacologically acceptable salt of L-
20 carnitine or of an alkanoyl L-carnitine is any salt with an acid that does not give rise to unwanted toxic or side effects.

These acids are well known to pharmacologists and to experts in pharmacy.

./.

Examples of pharmacologically acceptable salts of L-carnitine or alkanoyl L-carnitine, though not exclusively these, are chloride, bromide, orotate, aspartate, acid aspartate, acid citrate, acid phosphate, fumarate and acid fumarate, lactate, maleate and acid
5 maleate, mucate, acid oxalate, acid sulphate, glucose phosphate, tartrate and acid tartrate.

One preferred composition, in unit dosage form, is a composition containing 5-100 mg of ketanserin and 500-3000 mg of L-carnitine or an equivalent amount of alkanoyl L-carnitine.

10 This pharmaceutical composition is useful for the treatment of symptoms related to CRPS.

Here below are given, by way of example, the pharmacological results of experimental studies aimed at providing evidence of the surprising and unexpected synergistic effect achieved with the
15 combination of the invention.

In the following description, reference will be made only to L-carnitine, it being understood that the compositions described also apply to the above-mentioned alkanoyl L-carnitines and to the pharmacologically acceptable salts of both L-carnitine and the above-
20 mentioned alkanoyl L-carnitines.

Method of diagnosis of CRPS

The diagnosis of chronic regional pain syndrome was made in the presence of at least five of the following symptoms:

Persistent pain at rest;

5 Increasing pain during exercise;

Abnormal feeling of pain like hyperpathia or allodynia ;

Cold skin;

Glancing skin;

Hyperhidrosis;

10 Oedema;

Impaired mobility.

These symptoms were scored at the start of the treatment and after 3 months of oral treatment.

All patients gave their informed consent. Patients data are listed
15 in the following table 1.

20

TAB. 1

patient	M/F	Age (Years)	Delay (months)	Skin Tempera ture	Area
A	F	37	13	28.1	Foot
B	F	38	6	29.1	Foot
C	F	39	3	33.6	Hand
D	F	40	30	31.4	Hand
E	F	24	3	28.5	Foot
F	M	48	108	25.9	Foot
G	M	37	192	23.1	Foot
H	M	42	6	29.2	Knee
I	F	55	6	33.0	Hand
J	F	50	3	31.7	Hand
K	F	36	16	25.3	Hand
L	F	42	9	32.5	Foot
	Mean	40.7	32.9		
	S.D.	7.5	55.6		

All patient had a story of surgery on the affected area of hand,
 5 foot, or knee.

In spite of the fact that this group comprised only 12 patients,
 they matched with the general picture of CRPS patients found in
 greater studies, like H. J. M. Veldman et al., The Lancet, 342, 1012-
 1016 (1993).

./.

In a group of 829 CRPS patients Veldman had 76 % females and 24 % males. In the group treated according to the present invention, there were 67 % females and 33 % males.

The mean age of the Valdeman's study showed 42 years, in the
5 study according to the example of the present invention, it was 40.7 years. Veldman found 59 % CRPS of the upper extremity and 41 % lower extremity.

In our group were 42 % CRPS of the upper extremity and 58 % CRPS of the lower extremity.

10 The delay time between start of the symptoms and start of the treatment was in our case of 33 months.

This extremely long time interval is the consequence of two patients with a delay of 108 and 192 months.

When we eliminate these two patients, delay time is reduced to
15 9.5 months, and for six patients delay time is six months or less.

Looking at the phenomenon of a warm or a cold CRPS the objective measurement of the skin temperature was used.

Taking into account that normal skin temperature is about 32 °C, 7 cold extremities and five normal or warm extremities were
20 observed.

Special attention was given to the symptom of abnormal pain feeling, hyperpathia and allodinia.

This kind of symptom was found in 6 patients.

./.

Treatment

All i.v. treatments were performed after at least 15 minutes of acclimatisation at constant temperature (20-23 °C).

The temperature of the affected limb was recorded with Hewlett
5 Packard electro-skin probe between digit 4 and 5. In addition
continuous photoplethysmography of the affected hand (digit 2) or foot
(digit 1), blood pressure and heart rate were recorded.

All patients received a bolus of 10 mg of ketanserin i.v. in a
running saline infusion.

10 Ketanserin administration was continued by rate of 4 mg per
hour.

After one hour, a bolus injection of 1000 mg of L-carnitine was
given.

After this infusion start, oral maintenance therapy was started
15 with ketanserin three times daily 20 mg, and L-carnitine three times
daily 990 mg.

Results

To compare the clinical manifestation of the CRPS before and
after the infusion treatment followed by three months oral therapy an
20 evaluation of seven symptoms was made.

The seven symptoms were: pain at rest, pain during exercise,
impaired mobility, hyperhidrosis, oedema, small veins and
allodynia/hyperpathia.

./.

A scale with five degree was used. The nurse recorded the degree of the symptoms of the patient according to the following scale: absent, mild, reasonable, moderate and severe.

Fig. 1 shows the inventorisation at starting time; special
5 attention was given to the symptom allodinia, which is pointed on the first line.

This symptom was present in six out of these twelve patients. The results after three months are shown in Fig. 2.

After the intravenous treatment with ketanserin all patients had
10 peripheral circulation of the affected limb normalised, as proven in the past by Moesker (above cited) and by M. H. Hanna et al. [Pain, 38, 145-150 (1989)].

The symptom of small vein disappeared and the skin temperature was normalised.

15 After three months treatment with the two compounds according to the invention, patient "F" which was a severe case with a delay time of 108 months, had still the worst symptomatology, with pain at rest, pain during exercise and impaired mobility.

Patient "C" was a remarkably case, who had good recovery, but
20 still oedema and Hyperhidrosis.

Patient "B" was fully cured of any complaint of CRPS, despite she had a three months old full blow CRPS, with a skin temperature of 29,1 °C and seriously debilitating allodinia of the affected foot. Is

./.

noteworthy that the allodynia symptom disappeared in all patients that showed this symptom.

Impaired mobility and pain during exercise did not disappear completely probably because the treatment time with the combination
5 according to the present invention was too short.

Other positive results were obtained changing the therapeutic treatment protocol above cited.

In fact, it is possible to increase or decrease the amount of the two compounds either during the initial intravenous infusion or during
10 the maintenance therapeutic treatment (os).

In mild or recent CRPS it is possible to start directly with the above cited maintenance therapeutic treatment (os) with ketanserin and L-carnitine, three times daily.

It is an object of the present invention an orally or parenterally
15 administrable composition comprising ketanserin and L-carnitine or an alkanoyl L-carnitine or one of its pharmacologically acceptable salts.

Object of the present invention is also the co-ordinated use of the above-mentioned compounds i.e. the substantially simultaneous
20 administration of L-carnitine or one of the alkanoyl L-carnitines, or one of their pharmacologically acceptable salts, and ketanserin or, indifferently, the administration of a composition containing a

./.

combination or mixture of the aforesaid active ingredients, in addition to any excipient included.

The composition of the invention can be orally, parenterally or intravenously per infusion administered.

5 The composition according to the invention can be in the form of tablets, capsules, effervescent sachets, suppositories or vials.

Further object of the present invention is to provide a therapeutical kit comprising in the same package:

a) a first set of vials for intravenous infusion, said first set of vials
10 comprising 5-50 mg of ketanserin in admixture with pharmacologically acceptable vehicle and/or excipient, a second set of vials for intravenous infusion, said second set of vials comprising 500-2000 mg of L-carnitine or an equivalent amount of an alkanoyl L-carnitine or one pharmacologically acceptable salts
15 pharmaceutical composition comprising 100-3000 thereof, in admixture with pharmacologically acceptable vehicle and/or excipient;

and further comprising in the same package,

b) a first orally administrable pharmaceutical composition
20 comprising 5-100 mg of Ketanserin and a second orally administrable mg of L-carnitine or of an alkanoyl L-carnitine or one pharmacologically acceptable salts thereof, in admixture with pharmacologically acceptable vehicle and/or excipient.

./.

CLAIMS

1. A combination consisting of ketanserin and L-carnitine or an alkanoyl L-carnitine or one pharmacologically acceptable salts thereof.
- 5 2. Pharmaceutical composition comprising as active ingredients ketanserin and L-carnitine or an alkanoyl L-carnitine wherein the linear or branched-chain alkanoyl group has 2-6 carbon atoms or one pharmacologically acceptable salts thereof, in admixture with pharmacologically acceptable vehicle and/or excipient.
- 10 3. The composition according to claim 2, for oral, parenteral or intravenous infusion administration.
4. The composition according to claim 2, wherein the alkanoyl L-carnitine is selected from the group consisting of acetyl L-carnitine, propionyl L-carnitine, butyryl L-carnitine, valeryl L-carnitine and isovaleryl L-carnitine.
- 15 5. The composition according to claim 2, wherein the pharmacologically acceptable salt of L-carnitine or of the alkanoyl L-carnitine is selected from the group consisting of chloride, bromide, orotate, aspartate, acid aspartate, acid citrate, acid phosphate, fumarate and acid fumarate, lactate, maleate and acid maleate, mucate, acid oxalate, acid sulphate, glucose phosphate, tartrate and acid tartrate.
- 20

./.

6. The composition according to claim 2, in unit dosage form, comprising 5-100 mg of Ketanserin and 100-3000 mg of L-carnitine or of an alkanoyl L-carnitine.
7. The composition according to claim 6, suitable for oral administration.
8. The composition according to any of the preceding claims for the therapeutic treatment of CRPS.
9. Co-ordinated use of ketanserin and L-carnitine or of an alkanoyl L-carnitine or one of their pharmacologically acceptable salts for the treatment of CRPS.
10. A kit comprising in the same package a first set of vials for intravenous infusion, said first set of vials comprising 5-50 mg of ketanserin in admixture with pharmacologically acceptable vehicle and/or excipient, a second set of vials for intravenous infusion, said second set of vials comprising 500-2000 mg of L-carnitine or an equivalent amount of an alkanoyl L-carnitine wherein the linear or branched-chain alkanoyl group has 2-6 carbon atoms or one pharmacologically acceptable salts thereof, in admixture with pharmacologically acceptable vehicle and/or excipient.
11. A kit according to claim 10, further comprising in the same package a composition of claim 7.


./.

12. A kit according to claim 10 , further comprising in the same package a first orally administrable pharmaceutical composition comprising 5-100 mg of Ketanserin and a second orally administrable pharmaceutical composition comprising 100-3000 mg of L-carnitine or of an alkanoyl L-carnitine.
13. Use of ketanserin and L-carnitine or of an alkanoyl L-carnitine wherein the linear or branched-chain alkanoyl group has 2-6 carbon atoms or one of their pharmacologically acceptable salts for the preparation of a medicament for the treatment of CRPS.
14. Use according to claim 13, wherein the alkanoyl L-carnitine is selected from the group consisting of acetyl L-carnitine, propionyl L-carnitine, butyryl L-carnitine, valeryl L-carnitine and isovaleryl L-carnitine.
15. Use according to claim 13, wherein the pharmacologically acceptable salt of L-carnitine or of the alkanoyl L-carnitine is selected from the group consisting of chloride, bromide, orotate, aspartate, acid aspartate, acid citrate, acid phosphate, fumarate and acid fumarate, lactate, maleate and acid maleate, mucate, acid oxalate, acid sulphate, glucose phosphate, tartrate and acid tartrate.

16. Use according to claim 13, wherein said medicament comprises,
5-100 mg of Ketanserin and 100-3000 mg of L-carnitine or of an
alkanoyl L-carnitine, in a single unit dosage form.

17. Use according to claim 13, wherein said medicament comprises,
5-100 mg of Ketanserin in a first unit dosage form and 100-3000
mg of L-carnitine or of an alkanoyl L-carnitine in a second unit
dosage form, said first and second unit dosage forms being
physically separated between them.

CRPS Symptomatology



CRPS Symptomatology after 3 month

