

REPUBLIC OF SOUTH AFRICA
PATENTS ACT, 1978**PUBLICATION PARTICULARS AND ABSTRACT**

(Section 32(3)(a) - Regulation 22(1)(g) and 31)

OFFICIAL APPLICATION NO.

LODGING DATE

ACCEPTANCE DATE

21	01	20010240
----	----	----------

22	9 JAN 2001
----	------------

43	13 - 9 - 2001
----	---------------

INTERNATIONAL CLASSIFICATION

NOT FOR PUBLICATION

51	A61L; A61P
----	------------

CLASSIFIED BY: WIPO

FULL NAMES OF APPLICANT

71	LABORATORIOS DEL DR. ESTEVE, S.A.
----	-----------------------------------

FULL NAMES OF INVENTORS

72	1. MERCE-VIDAL, RAMON 2. FRIGOLA-CONSTANSA, JORDI
----	--

EARLIEST PRIORITY CLAIMED

COUNTRY

NUMBER

DATE

33	ES
----	----

31	P 9801467
----	-----------

32	10 JUL 1998
----	-------------

TITLE OF INVENTION

54	UTILIZATION OF DERIVATIVES OF TETRAHYDROPYRIDINES (OR 4-HYDROXYPIPERIDINES)- BUTYLAZOLS IN THE PREPARATION OF A MEDICAMENT FOR THE TREATMENT OF PAIN
----	---

57	ABSTRACT (NOT MORE THAT 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	50
------------------	----

If no classification is finished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.

Abstract


Derivatives of tetrahydropyridines (or 4-hydroxypiperidines)-butylazols of formula (I), wherein R1, R2 and R3, which are similar or different, represent each of them hydrogen, halogen, alkyl C1-C4, trifluoromethyl, hydroxyl, alkoxy, or two adjacent radicals can form a ring; A is a C atom and the dotted line represents an additional bond, or A is a C atom joined to a hydroxyl group and the dotted line represents absence of additional bond; Z1 is N or CR4; Z2 is N or CR5; Z4 is N or CR7; and R4, R5, R6 and R7, which are similar or different, represent hydrogen, halogen, alkyl C1-C4, aryl or substituted aryl, or two adjacent radicals can form part of another ring. These derivatives are useful for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including human beings.

5 Field of the Invention

The present invention relates to the use of derivatives of tetrahydropyridines (or 4-hydroxypyperidines)butylazoles of general formula (I), as well as their physiologically acceptable salts, in the preparation of medicaments useful in human and/or veterinary therapy for the treatment of acute pain, neuropathic pain and nociceptive pain, either alone or in combination with other analgesics, producing in this case a synergy.

Background of the invention

15 In our patent application WO 96/04287 compounds of general formula (I) are disclosed




20 that have an affinity for the σ and 5HT_{1A} receptors, and which are claimed as medicaments useful for the treatment of anxiety, psychosis, epilepsy, convulsion, amnesia, cerebro-vascular diseases and senile dementia.

We have now discovered that the compounds of general formula (I), as well as their pharmaceutically acceptable salts, are especially useful for the preparation of medicaments, useful in human and/or veterinary therapy for the prophylaxis, alleviation or curing of acute pain, neuropathic pain and nociceptive pain, either alone or in combination with other analgesics, giving rise in this case to a synergy.

Detailed description of the invention

The present invention relates to the use of derivatives of tetrahydropyridines (or 4-hydroxypyperidine)butylazoles of general formula



where

R_1 , R_2 and R_3 are either identical or different and represent a hydrogen atom, a halogen atom, a C_1 - C_4 alkyl group, a trifluoromethyl radical, a hydroxyl or alkoxy radical, and furthermore, two adjacent radicals can form part of an six-member aromatic ring;

A represents a carbon atom and the dotted line represents an additional bond, or A represents a carbon atom bound to a hydroxyl group (C-OH) and the dotted line represents the lack of an additional bond;

Z_1 represents a nitrogen atom or a substituted carbon atom that can be represented by C- R_4 ;

Z_2 represents a nitrogen atom or a substituted carbon atom that can be represented by C- R_5 ;

Z_4 represents a nitrogen atom or a substituted carbon atom that can be represented by C- R_7 ;

with the condition that Z_1 , Z_2 and Z_4 taken together can represent, at most, two nitrogen atoms; and

R_4 , R_5 , R_6 and R_7 , are identical or different and represent a hydrogen atom, a halogen atom, a C_1 - C_4 alkyl group, an aryl or substituted aryl group, or two adjacent radicals can form part of a six-member aromatic ring;

or one of their physiologically acceptable salts,

in the elaboration of a medicament for the treatment of acute pain, neuropatic pain or nocipeptive pain in mammals, including man.

The term "a halogen atom" represents a fluorine, chlorine or bromine atom.

5 The term "aryl or substituted aryl" represents a phenyl radical or a phenyl radical substituted by halogen.

The term "alkoxyl" represents a methoxyl or ethoxyl radical.

10 The term " C₁-C₄ alkyl" represents a straight chain or branched radical that is based on a saturated hydrocarbon of 1 to 4 atoms of carbon, such as methyl, ethyl, propyl, , isopropyl, butyl, isobutyl, sec-butyl y *terc*-butyl for example.

15 Physiologically acceptable salts of the compounds of general formula (I) refer both to salts formed with inorganic acids and organic acids, in particular, to salts of hydrochloric acid, hydrobromic acid, sulphuric acid, phosphoric acid, acetic acid, lactic acid, malonic acid, succinic acid, glutaric acid, fumaric acid, malic acid, tartaric acid, citric acid, ascorbic acid, maleic acid, benzoic acid, phenylacetic acid, cinamic acid, salicylic acid and alkyl, cycloalkyl or arylsulphonic acids.

20 The use of derivatives of general formula (I) for the treatment of pain refers to the use of analgesics in clinical practice. The term acute pain includes, but is not limited to, headache, arthritis, muscular tension or dysmenorrhea. The term neuropathic pain includes, but is not limited to, chronic back pains, pain associated with arthritis, herpes, pain associated with cancer, pain of a phantom limb, pain during childbirth or neuropathic pain resistant to opioids. The term nociceptive pain includes, but is not limited to, post-operation pain, dental pain, pain arising from surgery, pain caused by serious burns, post-natal pain or pain related with the genitourinary tract.

The derivatives of general formula (I) can be prepared according to the procedures disclosed in our patent application WO 96/04287.

30 In human therapy, the dosage administered of the compounds of the present invention varies as a function of the seriousness of the affliction to be treated. Normally the dosage will lie between 1 and 100 mg/day. The compounds of the invention can be administered as the only active ingredient


or in conjunction with another analgesic in a proportion of one part of compound of general formula (I) with around one to ten parts of the other analgesic, with the aim of provoking a synergy. Other analgesics include, but are not limited to, non-steroid anti-inflammatory compounds such as aspirin or indomethacine, other analgesics such as paracetamol, narcotic analgesics or related compounds such as morphine, meperidine or pentazocine. The compounds of the invention, with a suitable pharmaceutical formulation, are administered by different routes, such as orally, transdermically, parenterally, subcutaneously, intranasally, intramuscularly or intravenously.

5

10 Pharmaceutical compositions that contain compounds of general formula (I) are disclosed in our patent application WO 96/04287.

Illustrative examples of compounds included in the scope of the present invention include compounds that are characterised by the data indicated in tables 1 and 2.

TABLE 1



Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
1	H	H	H	H	H	N	CH	Cl	CH	102-103°C	3364 (b.a., OH), 2950, 2810, 1375, 1130, 991, 969, 760, 696, 605 KBr	1.56 (quin, J=7.1Hz, 2H); 1.65 (b.a., 1H); 1.76 (d, J=12.4Hz, 2H); 1.90 (quin, J=7.6Hz, 2H); 2.20 (m, 2H); 2.40-2.55 (a.c., 4H); 2.83 (d, J=9.5Hz, 2H); 4.11 (t, J=7Hz, 2H); 7.21-7.42 (a.c., 5H); 7.52 (d, J=8.5Hz, 2H) (CDCl ₃)
2	H	H	H	H	H	C-CH ₃	N	Cl	CCl	86-89°C	3196 (b.a., OH), 2951, 2924, 2824, 1406, 1247, 1146, 762, 703 KBr	1.59 (m, J=5.3 J'=6.6, 2H); 1.70-1.32 (a.c., 4H); 2.16 (d, J=13.0Hz J'=4.4Hz, 2H); 2.37 (s, 3H); 2.41-2.55 (a.c., 5H); 2.79 (d, J=11.3Hz, 2H); 3.88 (t, J=7.5Hz, 2H); 7.27 (t, J=7.2Hz, 1H); 7.36 (t, J=7.6Hz, 2H); 7.51 (d, J=7.3Hz, 2H) (CDCl ₃)
3	H	H	H	H	H	CH	N	CH=CH-CH=CH-C	CH=CH-CH=CH-C	122-123°C	3180 (b.a., OH), 2929, 2818, 1496, 1467, 1459, 1445, 1286, 1219, 1143, KBr	1.51 (quin, J=7.4Hz, 2H); 1.73 (d, J=12.7Hz, 2H); 1.87 (quin, J=7.6Hz, 2H); 2.10 (dt, J=12.9Hz J'=4.1Hz, 2H); 2.36-2.50 (a.c., 4H); 2.70 (d, J=11.2Hz, 2H); 3.25 (b.a., 1H); 4.12 (t, J=7.1Hz, 2H); 7.21-7.40 (a.c., 6H); 7.51

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R _G	Z ₄	m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											769, 743, 707 KBr	(d, J=8.3Hz, 2H); 7.70-7.75 (a.c., 2H) (CDCl ₃)
4	H	H	H	H	H	CH	N	H	N	123°C	3180 (b.a., OH), 2949, 2919, 2838, 1276, 1145, 1135, 1006, 770, 707, 676 KBr	1.45 (quin, J=7.5Hz, 2H); 1.69 (d, J=12.9Hz, 2H); 1.85 (quin, J=7.5Hz, 2H); 2.07 (dt, J=13.0Hz, J'=4.1Hz, 2H); 2.33-2.45 (a.c., 4H); 2.69 (d, J=11.2Hz, 2H); 2.93 (b.a., 1H); 4.10 (t, J=6.9Hz, 2H); 7.18 (t, J=7Hz, 1H); 7.27 (t, J=7.8Hz, 2H); 7.46 (d, J=8.3Hz, 2H); 7.80 (s, 1H); 7.91 (s, 1H) (CDCl ₃)
5	H	H	Cl	H	H	N	CH	Cl	CH	106°C	3145 (b.a., OH), 2947, 2918, 2834, 1318, 1147, 1083, 1112, 990, 817, 612 KBr	1.47 (quin, J=7.5Hz, 2H); 1.69 (d, J=11.9Hz, 2H); 1.84 (quin, J=7.6Hz, 2H); 2.05 (dt, J=13Hz, J'=4.4Hz, 2H); 2.34-2.50 (a.c., 5H); 2.72 (d, J=11.2Hz, 2H); 4.05 (t, J=7.0Hz, 2H); 7.29 (AB system, J=8.6Hz, 2H); 7.36 (s, 2H); 7.42 (AB system, J=8.6Hz, 2H) (CDCl ₃)
6	H	H	Cl	H	H	C-CH ₃	N	Cl	CCl	oil	3340 (b.a., OH), 2946, 2820, 1537, 1492, 1471, 1406, 1376, 1247, 1135, 1094, 1013, 828, 755 film	1.54 (m, 2H); 1.67-1.78 (a.c., 4H); 2.06 (dt, J=13Hz, J'=4.2Hz, 2H); 2.32 (s, 3H); 2.38-2.45 (a.c., 5H); 2.73 (d, J=11.2Hz, 2H); 3.86 (t, J=7.3Hz, 2H); 7.28 (AB system, J=8.6Hz, 2H); 7.43 (AB system, J=8.6Hz, 2H) (CDCl ₃)
7	H	CF ₃	H	H	H	N	CH	Cl	CH	oil	3360 (b.a., OH), 2948, 2823, 1438, 1378, 1330, 1212, film	1.48 (quin, J=7.6Hz, 2H); 1.71 (d, J=12.5Hz, 2H); 1.85 (quin, J=7.6Hz, 2H); 2.06-2.21 (a.c., 3H); 2.36-2.43 (a.c., 4H); 2.76 (d, J=11.5Hz, 2H); 4.06 (t, J=7.1Hz,

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											1165, 1124, 1047, 972, 804, 704 film	2H); 7.35 (s, 2H); 7.43-7.51 (a.c., 2H); 7.66 (d, J=7.5Hz, 1H); 7.79 (s, 1H) (CDCl ₃)
8	H	CF ₃	H	H	H	C-CH ₃	N	Cl	CCl	oil	3340 (b.a., OH), 2948, 2823, 1408, 1330, 1165, 1126, 1075, 789, 763, 704 film	1.57 (quin, J=7.5Hz, 2H); 1.70-1.80 (a.c., 4H); 2.15 (dt, J=12.9Hz J'=3.6Hz, 2H); 2.35 (s, 3H); 2.40-2.52 (a.c., 4H); 2.80 (d, J=11.7Hz, 2H); 3.88 (t, J=7.0Hz, 2H); 7.42-7.57 (a.c., 2H); 7.69 (d, J=7.5Hz, 1H); 7.82 (s, 1H) (CDCl ₃)
9	H	H	F	H	H	C-CH ₃	N	Cl	CCl	oil	3330 (b.a., OH), 2946, 2818, 1509, 1406, 1247, 1222, 1160, 835 film	1.58 (m, 2H); 1.64-1.81 (a.c., 4H); 2.14 (dt, J=12.9Hz J'=3.6Hz, 2H); 2.32 (s, 3H); 2.43-2.60 (a.c., 4H); 2.84 (d, J=11Hz, 2H); 3.87 (t, J=7.1Hz, 2H); 4.18 (b.a., 1H); 7.01 (t, J=8.8Hz, 2H); 7.46 (dd, J=8.8Hz J'=5.2Hz, 2H) (CDCl ₃)
10	H	H	H	H	H	CH	CH	CH=CH-CH=CH-C		109-111°C	3190 (b.a., OH), 2956, 2823, 1461, 1446, 1319, 1303, 1218, 1142, 738, 703 KBr	1.57 (m, 2H); 1.73 (d, J=14Hz, 2H); 1.80 (b.a., 1H); 1.90 (m, 2H); 2.13 (dt, J=13Hz J'=4Hz, 2H); 2.32-2.46 (a.c., 4H); 2.76 (d, J=11.3Hz, 2H); 4.16 (t, J=7.1Hz, 2H); 6.50 (d, J=3.1Hz, 1H); 7.05-7.14 (a.c., 2H); 7.18-7.40 (a.c., 5H); 7.50 (d, J=7.8Hz, 2H); 7.00 (d, J=7.3Hz, 1H) (CDCl ₃)
11	H	H	CH ₃	H	H	C-CH ₃	N	Cl	CCl	oil	3360 (b.a., OH), 2946, 2818, 1535, 1471, 1406, 1376, 1247, 1134, 817, 755	1.53 (m, 2H); 1.66-1.84 (a.c., 4H); 2.09 (dt, J=12.9Hz, J'=3.6Hz, 2H); 2.33 (s, 3H); 2.36 (s, 3H); 2.39-2.50 (a.c., 4H); 2.77 (d, J=11.2Hz, 2H); 3.87 (t, J=7.0Hz, 2H); 7.15 (AB system, J=7.8Hz, 2H); 7.33 (AB system, J=7.8Hz, 2H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											film	
12	H	H	H	H	H	N	CH	H	CH	89-91°C	3137 (b.a., OH) 2947, 2532, 1396, 1378, 1119, 1046, 756, 697 KBr	1.51 (quin, J=7.6Hz, 2H); 1.73 (d, J=12.3Hz, 2H); 1.89 (quin, J=7.6Hz, 2H); 2.00-2.20 (a.c., 3H); 2.35-2.45 (a.c., 4H); 2.76 (d, J=10.2Hz, 2H); 4.13 (t, J=7.1Hz, 2H); 6.21 (s, 1H); 7.21 (m, 1H); 7.30-7.37 (a.c., 3H); 7.44-7.52 (a.c., 3H) (CDCl ₃)
13	H	H	H	H	H	N	CH	CH=CH-CH=CH-C		107-109°C	3311 (b.a., OH) 2953, 2803, 1465, 1375, 1133, 1117, 1043, 1017, 761, 744, 704 KBr	1.53 (m, 2H); 1.71 (d, J=12.2Hz, 2H); 1.95 (m, 2H); 2.10 (m, 2H); 2.29 (b.a., 1H); 2.35-2.47 (a.c., 4H); 2.71 (d, 2H); 4.39 (t, J=7.1Hz, 2H); 7.13 (t, 1H); 7.22-7.44 (a.c., 5H); 7.50 (d, J=8Hz, 2H); 7.71 (d, J=8.3Hz, 1H); 7.95 (s, 1H) (CDCl ₃)
14	H	H	H	H	H	N	C-CH=CH- CH=CH	CH	CH	120-122°C	3295 (b.a., OH) 2946, 2817, 1377, 1126, 786, 735, 700 KBr	1.58 (m, 2H); 1.73 (d, J=13.5Hz, 2H); 1.90-2.20 (a.c., 5H); 2.38-2.47 (a.c., 4H); 2.75 (d, J=10.5Hz, 2H); 4.42 (t, J=6.6Hz, 2H); 7.06 (t, J=7.5Hz, 1H); 7.22-7.37 (a.c., 4H); 7.49 (d, J=7.8Hz, 2H); 7.61-7.71 (a.c., 2H); 7.90 (s, 1H) (CDCl ₃)
15	H	H	CH ₃	H	H	N	CH	Cl	CH	81-82°C	3122 (b.a., OH) 2936, 1475, 1434, 1378, 1319, 989, 973, 814 KBr	1.51 (quin, J=7.6Hz, 2H); 1.73 (d, J=11.7Hz, 2H); 1.87 (quin, J=7.6Hz, 2H); 2.12 (dt, J=12.8Hz J'=4.4Hz, 2H); 2.33 (s, 3H); 2.35-2.48 (a.c., 5H); 2.74 (d, J=11.2Hz, 2H); 4.07 (t, J=7.1Hz, 2H); 7.15 (d, J=8Hz, 2H); 7.25-7.40 (a.c., 4H) (CDCl ₃)
16	H	H	CH ₃ O	H	H	N	CH	Cl	CH	122-123°C	3190 (b.a., OH) 2954, 2923, 2827, 1509, 1314, 1243, KBr	1.49 (quin, J=7.6Hz, 2H); 1.72 (d, J=11.8Hz, 2H); 1.84 (quin, J=7.4Hz, 2H); 2.00-2.14 (a.c. (dt+b.a.), 3H); 2.34-2.47 (a.c., 4H); 2.72 (d, J=11Hz, 2H); 3.77 (s,


Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
17	H	H	H	H	H	CPh	N	H	CH	108-110°C	1178, 971 KBr 3220 (b.a., OH) 2944, 2817, 1473, 1446, 1421, 1136, 1046, 787, 773, 761, 700 KBr	3H); 4.05 (t, J=7.1Hz, 2H); 6.85 (d, J=9Hz, 2H), 7.24-7.42 (a.c., 4H) (CDCl ₃) 1.45 (quin, J=7.6Hz, 2H); 1.68-1.82 (a.c., 4H); 2.08 (dt, J=13.0Hz, J'=4.1Hz, 2H); 2.29-2.42 (a.c., 4H); 2.5 (b.a., 1H); 2.67 (d, J=11.2Hz, 2H); 4.01 m (t, J=7.3Hz, 2H); 7.01 (s, 1H); 7.08 (s, 1H); 7.20-7.56 (a.c., 10H) (CDCl ₃)
18	H	H	CH ₃	H	H	CH	N	CH=CH-CH=CH-C		oil	3260 (b.a., OH) 2944, 2817, 1497, 1459, 1381, 1287, 1135, 1046, 817, 745 film	1.58 (quin, J=7.6Hz, 2H); 1.74 (d, J=12Hz, 2H); 1.82 (b.a., 1H); 1.95 (quin, J=7.6Hz, 2H); 2.11 (dt, 2H); 2.33 (s, 3H); 2.40-2.50 (a.c., 4H); 2.74 (d, J=11.5Hz, 2H); 4.20 (t, J=7.1Hz, 2H); 7.15 (d, J=8.3Hz, 2H); 7.22-7.35 (a.c., 3H); 7.37-7.43 (a.c., 2H); 7.79 (m, 1H); 7.87 (s, 1H) (CDCl ₃)
19	H	H	H	H	H	CH	N	Ph	CPh	138-139°C	3194 (b.a., OH) 2939, 2806, 1509, 1446, 773, 766, 758, 696 KBr	1.38 (m, 2H); 1.56 (m, 2H); 1.72 (d, J=12.4Hz, 2H); 2.09 (dt, 2H); 2.25 (t, J=7.4Hz, 2H); 2.39 (m, 2H); 2.66 (m, 2H); 3.10 (b.a., 1H); 3.78 (t, J=7.2Hz, 2H); 7.10-7.52 (a.c., 16H);

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
20	CH=CH-CH=CH	CH=CH=CH	H	H	H	N	CH	Cl	CH	oil	3357 (b.a., OH), 2946, 2833, 1434,	1.44 (quin, J=7.3Hz, 2H); 1.77 (quin, J=7.5Hz, 2H); 2.15-2.30 (a.c., 5H); 2.34 (t, J=7.5Hz, 2H); 2.57 (m, 2H);

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											KBr	(DMSO-d ₆)
25	H	H	F	H	H	N	CH	CH=CH-CH=CH-C		136-137°C	3303, 2951, 2805, 1506, 1464, 1376, 1218, 1162, 1118, 832, 741	1.54 (m, 2H); 1.60-1.80 (a.c., 3H); 1.97 (m, 2H); 2.06 (dt, J=13.0Hz, J'=4.3Hz, 2H); 2.30-2.43 (a.c. 4H, 2.72 (m, 2H); 4.40 (t, J=7.0Hz, 2H); 6.99 (t, J=8.8Hz, 2H); 7.12 (m, 1H); 7.32-7.47 (a.c., 4H); 7.71 (d, J=8.1Hz, 1H); 7.96 (s, 1H) (CDCl ₃ -CD ₃ OD [1:1])
26	H	H	F	H	H	N	C-CH=CH-CH=CH	CH		148-150°C	3325, 2950, 2923, 2812, 1509, 1377, 1218, 1131, 834, 758	1.57 (m, 2H); 1.70-1.77 (a.c., 3H); 1.98-2.19 (a.c., 4H); 2.35-2.49 (a.c., 4H); 2.77 (d, J=11.2Hz, 2H); 4.45 (t, J=7.0Hz, 2H); 6.98-7.15 (a.c., 3H); 7.25-7.49 (a.c., 3H); 7.63 (d, J=8.3Hz, 1H); 7.69 (d, J=7.8Hz, 1H); 7.91 (s, 1H) (CDCl ₃ -CD ₃ OD [1:1])
27	H	H	F	H	H	N	C-CH=CH-CH=CH	N		109-110°C	3400, 2931, 2812, 1509, 1229, 1101, 831, 745	1.47-1.80 (a.c., 4H); 1.90-2.25 (a.c., 5H); 2.30-2.55 (a.c., 4H); 2.70 (m, 2H); 4.78 (t, J=6.9Hz, 2H); 7.01 (t, J=8.7Hz, 2H); 7.26-7.54 (a.c., 4H); 7.85 (dd, J=6.7Hz, J'=3.0Hz, 2H) (CDCl ₃ -CD ₃ OD [1:1])
28	H	H	F	H	H	N	CH=CH-CH=CH-C			102-103°C	3430, 2952, 2925, 1508, 1223, 1140, 833, 744	1.45-1.80 (a.c., 4H); 1.85-2.25 (a.c., 5H); 2.25-2.55 (a.c., 4H); 2.77 (m, 2H); 4.69 (t, J=6.9Hz, 2H); 7.01 (t, J=8.7Hz, 2H); 7.26-7.53 (a.c., 5H); 8.06 (d, J=7.3Hz, 1H) (CDCl ₃ -CD ₃ OD [1:1])
29	H	H	F	H	H	CH	N	H	N	oil	3350 (b.a., OH), 2947, 2818, 1509, 1222, 1138, 836, 681	1.55 (m, 2H); 1.74 (d, J=12.6 Hz, 2H); 1.94 (m, 2H); 2.13 (m, 2H); 2.40-2.55 (a.c.; 4H); 2.79 (m, 2H); 4.20 (t, J=6.9 Hz, 2H); 7.02 (t, J=8.4 Hz, 2H); 7.46 (m, 2H); 7.91 (s, 1H); 8.04 (s, 1H) (CDCl ₃)
30	H	H	Cl	H	H	CH	N	H	N	89-91°C	3119 (b.a., OH), 2956, 2829, 1509,	1.46 (m, 2H); 1.71 (m, 2H); 1.90 (quin, J=7.4Hz, 2H); 2.05 (m, 2H); 2.33-2.50 (a.c., 4H); 2.54 (b.a., 1H); 2.72

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											1379, 1277, 1145, 1007, 824, 685 KBr	(m, 2H); 4.16 (t, J=7.1Hz, 2H); 7.28 (m, 2H); 7.42 (m, 2H); 7.86 (s, 1H); 7.99 (s, 1H) (CDCl ₃)

TABLE 2



5

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
1a	H	H	H	H	H	N	CH	Cl	CH	62-64°C	3113, 2920, 2745, 1375, 1325, 1138, 965, 837, 742, 688 KBr	1.56 (quin, J=7.6Hz, 2H); 1.91 (quin, J=7.6Hz, 2H); 2.47 (t, J=7.4Hz, 2H); 2.58 (m, 2H); 2.65 (t, J=5.6Hz, 2H); 3.14 (m, 2H); 4.11 (t, J=7.1Hz, 2H); 6.06 (m, 1H); 7.23-7.42 (a.c., 7H) (CDCl ₃)
2a	H	H	H	H	H	CH	N	CH=CH-CH=CH-C		66-69°C	2933, 1495, 745, 694, 665 film	1.55 (quin, J=7.6Hz, 2H); 1.92 (quin, J=7.6Hz, 2H); 2.43 (t, J=7.3Hz, 2H); 2.52 (m, 2H); 2.61 (t, J=5.6Hz, 2H); 3.07 (m, 2H); 4.14 (t, J=7.1Hz, 2H); 6.02 (m, 1H); 7.20-7.40 (a.c., 8H); 7.80 (m, 1H); 7.86 (s, 1H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
3a	H	H	H	H	H	CH	N	H	N	63-64°C	2942, 1438, 1381, 1271, 1142, 1006, 753, 697, 681, KBr	1.56 (m, 2H); 1.95 (m, 2H); 2.47 (t, J=7.1Hz, 2H); 2.56 (m, 2H); 2.66 (t, J=5.3Hz, 2H); 3.11 (m, 2H); 4.19 (t, J=7.0Hz, 2H); 6.05 (s, 1H); 7.21 (m, 1H); 7.30 (t, J=7.6Hz, 2H); 7.36 (d, J=7.8Hz, 2H); 7.94 (s, 1H); 8.06 (s, 1H) (CDCl ₃)
4a	H	H	Cl	H	H	N	CH	Cl	CH	103-104°C	2939, 1493, 1436, 1381, 1306, 1122, 1097, 973, 843, 824, 730 KBr	1.54 (m, 2H); 1.90 (m, 2H); 2.45 (t, J=7.4Hz, 2H); 2.51 (m, 2H); 2.65 (t, J=5.6Hz, 2H); 3.10 (m, 2H); 4.10 (t, J=7.0Hz, 2H); 6.03 (m, 1H); 7.26 (AB system, J=8.6Hz, 2H); 7.29 (AB system, J=8.6Hz, 2H); 7.37 (s, 1H); 7.41 (s, 1H) (CDCl ₃)
5a	H	H	Cl	H	H	C-CH ₃	N	Cl	CCl	119-120°C	2922, 1531, 1494, 1469, 1403, 1380, 1366, 1245, 1094, 1010 KBr	1.59 (m, 2H); 1.76 (m, 2H); 2.36 (s, 3H); 2.42-2.53 (a.c., 4H); 2.67 (t, J=5.3Hz, 2H); 3.12 (m, 2H); 3.88 (t, J=7.4Hz, 2H); 6.04 (m, 1H); 7.27 (AB system, J=9.1Hz, 2H); 7.30 (AB system, J=9.1Hz, 2H) (CDCl ₃)
6a	H	CF ₃	H	H	H	N	CH	Cl	CH	oil	2944, 1434, 1375, 1331, 1247, 1165, 1126, 1076, 972, 800, 698 film	1.53 (quin, J=7.5Hz, 2H); 1.89 (quin, J=7.7Hz, 2H); 2.45 (t, J=7.3Hz, 2H); 2.54 (m, 2H); 2.66 (t, J=5.5Hz, 2H); 3.10 (m, 2H); 4.08 (t, J=7.1Hz, 2H); 6.10 (m, 1H); 7.35-7.56 (a.c., 5H); 7.59 (s, 1H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
7a	H	CF ₃	H	H	H	C-CH ₃	N	Cl	CCl	oil	2931, 2815, 1533, 1405, 1331, 1246, 1165, 1125, 1076, 797, 699 film	1.62 (quin, J=6.6Hz, 2H); 1.77 (quin, J=7.6Hz, 2H); 2.37 (s, 3H); 2.51 (t, J=7.2Hz, 2H); 2.60 (m, 2H); 2.71 (t, J=5.6Hz, 2H); 3.17 (m, 2H); 3.89 (t, J=7.3Hz, 2H); 6.14 (m, 1H); 7.40-7.50 (a.c., 2H); 7.55 (d, J=7.5Hz, 1H); 7.62 (s, 1H) (CDCl ₃)
8a	H	H	F	H	H	N	CH	Cl	CH	86-87°C	2936, 1512, 1378, 1326, 1229, 988, 967 KBr	1.60 (quin, J=7.5Hz, 2H); 1.91 (quin, J=7.5Hz, 2H); 2.50-2.82 (a.c., 4H); 2.76 (t, J=5.6Hz, 2H); 3.19 (m, 2H); 4.11 (t, J=6.9Hz, 2H); 5.97 (s, 1H); 6.99 (t, J=8.8Hz, 2H); 7.32 (dd, J=8.8Hz, J'=5.4Hz, 2H); 7.38 (s, 1H); 7.40 (s, 1H) (CDCl ₃)
9a	H	H	F	H	H	C-CH ₃	N	Cl	CCl	79-82°C	2934, 1531, 1512, 1408, 1247, 1225, 1167, 818 KBr	1.74 (m, 4H); 2.35 (s, 3H); 2.60-2.72 (a.c., 4H); 2.90 (m, 2H); 3.33 (m, 2H); 3.88 (m, 2H); 5.95 (s, 1H); 6.99 (t, J=8.6Hz, 2H); 7.31 (a.c., 2H) (CDCl ₃)
10a	H	H	H	H	H	C-CH ₃	N	Cl	CCl	oil	2929, 1533, 1405, 1246, 748 film	1.59 (m, 2H); 1.76 (m, 2H); 2.37 (s, 3H); 2.49 (t, J=7.3Hz, 2H); 2.58 (m, 2H); 2.69 (t, J=5.4Hz, 2H); 3.14 (m, 2H); 3.89 (t, J=7.4Hz, 2H); 6.06 (m, 1H); 7.22-7.40 (a.c., 5H) (CDCl ₃)
11a	H	H	H	H	H	C-CH ₃	N	Cl	CCl	• HCl 203-204°C	2930, 2576, 1407, 1376, 1245, 750 KBr	1.69 (m, 2H); 1.81 (m, 2H); 2.35 (s, 3H); 2.71 (d, J=7.2Hz, 1H); 2.91 (m, 1H); 3.17 (a.c., 3H); 3.56 (m, 1H); 3.75 (m, 1H); 3.90-3.97 (a.c., 3H); 6.17 (s, 1H); 7.25-7.40 (a.c., 3H); 7.47 (d, J=7.6Hz, 2H); 11.30 (b.a., 1H) (DMSO-d ₆)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
12a	H	H	H	H	H	C-CH ₃	N	Cl	CCl	• 2HCl 192-194°C	3569, 2941, 2692, 2556, 1601, 1446, 769, 753, 698 KBr	1.67 (m, 2H); 1.79 (m, 2H); 2.36 (s, 3H); 2.69 (d, J=18.0Hz, 1H); 2.88 (m, 1H); 3.15 (a.c., 3H); 3.54 (m, 1H); 3.72 (m, 1H); 3.85-3.98 (a.c., 3H); 6.15 (s, 1H); 7.22-7.38 (a.c., 3H); 7.45 (d, J=7.3Hz, 2H); 9.93 (b.a., 1H); 11.36 (b.a., 1H) (DMSO-d ₆)
13a	H	H	F	H	H	CH	CH	CH=CH-CH=CH-C	CH=CH-CH=CH-C	oil	2937, 1510, 1464, 1230, 1161, 816, 742 film	1.61 (quin, J=7.7Hz, 2H); 1.93 (quin, J=7.6Hz, 2H); 2.42-2.58 (a.c., 4H); 2.66 (t, J=5.6Hz, 2H); 3.11 (m, 2H); 4.17 (t, J=7.0Hz, 2H); 5.98 (m, 1H); 6.51 (d, J=3.9Hz, 1H); 6.95-7.39 (a.c., 8H); 7.65 (d, J=7.8Hz, 1H) (CDCl ₃)
14a	H	H	H	H	H	CH	CH	CH=CH-CH=CH-C	CH=CH-CH=CH-C	oil	2938, 1510, 1485, 1463, 1446, 1376, 1336, 1315, 763, 740, 695 film	1.63 (quin, J=7.4Hz, 2H); 1.94 (quin, J=7.4Hz, 2H); 2.49 (t, J=7.6Hz, 2H); 2.60 (m, 2H); 2.69 (t, J=5.3Hz, 2H); 3.14 (m, 2H); 4.19 (t, J=7.1Hz, 2H); 6.08 (m, 1H); 6.53 (m, 1H); 7.08-7.44 (a.c., 9H); 7.67 (d, J=8.1Hz, 1H) (CDCl ₃)
15a	H	H	CH ₃	H	H	C-CH ₃	N	Cl	CCl	87-88°C	2939, 2916, 1529, 1404, 1378, 1243, 1166, 1131, 1016 film	1.59 (m, 2H); 1.75 (m, 2H); 2.32 (s, 3H); 2.36 (s, 3H); 2.47 (t, J=7.2Hz, 2H); 2.54 (m, 2H); 2.67 (t, J=5.2Hz, 2H); 3.11 (m, 2H); 3.87 (t, J=7.3Hz, 2H); 6.01 (s, 1H); 7.11 (AB system, J=8.1Hz, 2H); 7.27 (AB system, J=8.1Hz, 2H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
16a	H	H	H	H	H	N	CH	H	CH	36-38°C	2941, 1396, 748, 695 film	1.54 (quin, J=7.6Hz, 2H); 1.91 (quin, J=7.6Hz, 2H); 2.45 (t, J=7.6Hz, 2H); 2.55 (m, 2H); 2.65 (t, J=5.6Hz, 2H); 3.11 (m, 2H); 4.14 (t, J=7.1Hz, 2H); 6.03 (m, 1H); 6.21 (m, 1H); 7.20-7.39 (a.c., 6H); 7.49 (m, 1H) (CDCl ₃)
17a	H	H	H	H	H	N	CH	CH=CH-CH=CH-C		50-52°C	2942, 1465, 1158, 832, 740, 691 KBr	1.61 (quin, 2H); 2.00 (quin, J=7.5Hz, 2H); 2.43-2.58 (a.c., 4H); 2.68 (m, 2H); 3.14 (s, 2H); 4.43 (t, J=6.6Hz, 2H); 6.02 (s, 1H); 7.13 (t, J=7.3Hz, 1H); 7.20-7.51 (a.c., 7H); 7.73 (d, J=7.9Hz, 1H); 7.99 (s, 1H) (CDCl ₃)
18a	H	H	H	H	H	N	C-CH=CH- CH=CH	CH=CH- CH=CH	CH	73-75°C	3049, 2940, 2778, 1467, 1371, 1158, 1143, 1131, 757, 742, 692 KBr	1.60 (quin, J=7.6Hz, 2H); 2.09 (quin, J=7.4Hz, 2H); 2.48 (t, J=7.4Hz, 2H); 2.55 (m, 2H); 2.66 (t, J=5.6Hz, 2H); 3.11 (d, J=2.9Hz, 2H); 4.45 (t, J=7.1Hz, 2H); 6.03 (s, 1H); 7.07 (t, J=7.5Hz, 1H); 7.20-7.39 (a.c., 6H); 7.63 (d, J=4.3Hz, 1H); 7.70 (d, J=8Hz, 1H); 7.91 (s, 1H) (CDCl ₃)
19a	H	H	CH ₃	H	H	N	CH	Cl	CH	72-73°C	3115, 2938, 2740, 1376, 1328, 1137, 986, 966, 844, 824, 797 KBr	1.55 (quin, 2H); 1.90 (quin, J=7.5Hz, 2H); 2.33 (s, 3H); 2.46 (t, J=7.5Hz, 2H); 2.55 (m, 2H); 2.66 (t, J=6.4Hz, 2H); 3.11 (m, 2H); 4.10 (t, J=7.0Hz, 2H); 6.01 (s, 1H); 7.12 (AB system, J=8Hz, 2H); 7.27 (AB system, J=8Hz, 2H); 7.37 (s, 1H); 7.41 (s, 1H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
20a	H	H	CH ₃ O	H	H	N	CH	Cl	CH	104-105°C	2923, 1533, 1405, 1379, 1246, 749 KBr	1.54 (quin, 2H); 1.89 (quin, J=7.6Hz, 2H); 2.44 (t, J=7.4Hz, 2H); 2.52 (m, 2H); 2.65 (t, J=5.3Hz, 2H); 3.10 (m, 2H); 3.78 (s, 3H); 4.09 (t, J=7.0Hz, 2H); 5.95 (s, 1H); 6.84 (AB system, J=8.5Hz, 2H); 7.31 (AB system, J=8.5Hz, 2H); 7.36 (s, 1H); 7.40 (s, 1H) (CDCl ₃)
21a	H	H	H	H	H	N	CH	Cl	CH	oil	2948, 2923, 2811, 2774, 1446, 1382, 1316, 971, 748, 695 film	2.08 (quin, J=7.0Hz, 2H); 2.42 (t, J=7.0Hz, 2H); 2.58 (m, 2H); 2.67 (t, J=5.6Hz, 2H); 3.13 (m, 2H); 4.17 (t, J=6.9Hz, 2H); 6.07 (m, 1H); 7.23-7.45 (a.c., 7H) (CDCl ₃)
22a	H	H	H	H	H	CCH ₃	N	Cl	CCl	oil	2923, 1533, 1405, 1379, 1246, 749 film	1.95 (quin, J=7.2Hz, 2H); 2.39 (s, 3H); 2.46 (t, J=7.0Hz, 2H); 2.58 (m, 2H); 2.69 (t, J=4.9Hz, 2H); 3.13 (m, 2H); 3.96 (t, J=7.3Hz, 2H); 6.07 (m, 1H); 7.20-7.41 (a.c., 5H) (CDCl ₃)
23a	H	H	H	H	H	CPh	N	H	CH	oil	2940, 1496, 1474, 1445, 1379, 1275, 774, 698 film	1.51 (m, 2H); 1.81 (m, 2H); 2.40 (t, J=7.4Hz, 2H); 2.56 (m, 2H); 2.63 (t, J=4.9Hz, 2H); 3.09 (m, 2H); 4.04 (t, J=7.2Hz, 2H); 6.03 (m, 1H); 7.03 (m, 1H); 7.13 (m, 1H); 7.22-7.48 (a.c., 8H); 7.58 (m, 2H) (CDCl ₃)

Ex	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
24a	H	H	CH ₃	H	H	CH	N	CH=CH-CH=CH-C		90-91°C	2939, 2915, 1500, 1461, 1377, 1365, 750 KBr	1.59 (m, 2H); 1.95 (m, 2H); 2.32 (s, 3H); 2.46 (t, J=7.3Hz, 2H); 2.53 (m, 2H); 2.63 (t, J=5.5Hz, 2H); 3.08 (m, 2H); 4.20 (t, J=6.95Hz, 2H); 6.00 (s, 1H); 7.11 (d, J=7.8Hz, 2H); 7.27 (a.c., 4H); 7.40 (m, 1H); 7.80 (m, 1H); 7.89 (s, 1H) (CDCl ₃)
25a	H	H	H	H	H	CH	N	Ph	CPh	100-101°C	3130, 2939, 2770, 1600, 1506, 1443, 1259, 954, 780, 774, 750, 696, 649 KBr	1.46 (quin, J=7.5Hz, 2H); 1.65 (quin, J=7.6Hz, 2H); 2.33 (t, J=7.3Hz, 2H); 2.53 (m, 2H); 2.60 (m, 2H); 3.05 (m, 2H); 3.84 (t, J=7.2Hz, 2H); 6.02 (m, 1H); 7.05-7.50 (a.c., 15H); 7.61 (s, 1H) (CDCl ₃)

Ex	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
26a	CH=CH-CH=CH		H	H	H	N	CH	Cl	CH	oil	3057, 3043, 2942, 2806, 2768, 1378, 1365, 971, 801, 778 film	1.61 (quin, J=7.5Hz, 2H); 1.95 (quin, J=7.6Hz, 2H); 2.51-2.57 (a.c., 4H); 2.76 (t, J=5.6Hz, 2H); 3.20 (m, 2H); 4.14 (t, J=7.1Hz, 2H); 5.74 (m, 1H); 7.26-7.50 (a.c., 6H); 7.75 (d, J=8Hz, 1H); 7.84 (m, 1H); 8.02 (m, 1H); (CDCl ₃)
27a	H	CH=CH-CH=CH		H	H	N	CH	Cl	CH	95-96°C	3111, 2920, 2806, 1374, 1326, 966, 826, 749, 612, KBr	1.57 (m, 2H); 1.92 (m, 2H); 2.48 (m, 2H); 2.71 (a.c., 4H); 3.18 (m, 2H); 4.11 (m, 2H); 6.22 (m, 1H); 7.38-7.50 (a.c., 4H); 7.61 (m, 1H); 7.75-7.84 (a.c., 4H) (CDCl ₃)

Ex	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
28a	H	H	F	H	H	CH	N	CH=CH-CH=CH-C		135-136°C	3050, 2920, KBr	2.54 (m, 2H); 2.74 (t, J=5.6Hz, 2H); 2.92 (t, J=6.7Hz, 2H); (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											2780, 2760, 1510, 1492, 1459, 1224, 1202, 1161, 771, 751 KBr	2H); 3.24 (m, 2H); 4.35 (t, J=6.7Hz, 2H); 5.98 (m, 1H); 7.00 (t, J=8.7Hz, 2H); 7.26-7.40 (a.c., 4H); 7.42 (m, 1H); 7.81 (m, 1H); 8.01 (s, 1H) (CDCl ₃)
29a	H	H	H	H	H	CH	N	CH=CH-CH=CH-C		HCl 177-178°C	2940, 2488, 1500, 1420, 1390, 742 KBr	1.70-1.90 (a.c., 4H); 2.78 (m, 2H); 3.17 (m, 2H); 3.20-3.50 (b.a., 2H); 3.79 (m, 2H); 4.30 (t, J=6.6Hz, 2H); 6.15 (s, 1H); 7.17-7.40 (a.c., 5H); 7.45 (d, J=7.3Hz, 2H); 7.65 (m, 2H); 8.35 (s, 1H) (DMSO-d ₆)
30a	H	H	F	H	H	CH	N	CH=CH-CH=CH-C		106-108°C	2942, 1512, 1498, 1460, 1376, 1221, 756 KBr	1.59 (quin, J=7.5Hz, 2H); 1.96 (quin, J=7.5Hz, 2H); 2.40- 2.50 (a.c., 4H); 2.63 (t, J=5.5Hz, 2H); 3.09 (m, 2H); 4.21 (t, J=7.1Hz, 2H); 5.97 (m, 1H); 6.98 (t, J=8.1Hz, 2H); 7.20-7.35 (a.c., 4H); 7.40 (m, 1H); 7.80 (m, 1H); 7.89 (s, 1H) (CDCl ₃)
31a	H	H	F	H	H	CH	N	CH=CH-CH=CH-C		HCl	2930, 1600, 1510, 1275 KBr	1.70-2.00 (a.c., 4H); 2.78 (m, 2H); 3.20 (m, 2H); 3.20- 3.60 (b.a., 2H); 3.81 (m, 2H); 4.38 (t, J=6.6Hz, 2H); 6.13 (s, 1H); 7.19 (t, J=8.7Hz, 2H); 7.33 (m, 2H); 7.49 (m, 2H); 7.71 (d, J=7.8Hz, 1H); 7.77 (d, J=7.6Hz, 1H); 8.79 (s, 1H); 11.20 (b.a., 1H) (DMSO-d ₆)
32a	H	CF ₃	H	H	H	CCH ₃	N	Cl	CCl	HCl 205-206°C	2930, 2490, 1330, 1243, 1164, KBr	1.67 (m, 2H); 1.79 (m, 2H); 2.33 (s, 3H); 2.79 (m, 1H); 2.91 (m, 1H); 3.10-3.20 (a.c., 3H); 3.55 (m, 1H); 3.77 (m, 1H); 3.91-4.00 (a.c., 3H); 6.33 (s, 1H); 7.58-7.80

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											1119, 1076 KBr	(a.c., 4H); 11.32 (b.a., 1H) (DMSO-d ₆)
33a	H	H	F	H	H	N	CH	Cl	CH	HCl 191-192°C	2543, 1512, 1232, 967, 807 KBr	1.71-1.85 (a.c., 4H); 2.68 (m, 1H); 2.86 (m, 1H); 3.10-3.20 (a.c., 3H); 3.55 (m, 1H); 3.72 (m, 1H); 3.90 (m, 1H); 4.12 (t, J=6.5Hz, 2H); 6.14 (s, 1H); 7.20 (t, J=8.7Hz, 2H); 7.40-7.55 (a.c., 3H); 8.06 (s, 1H); 11.20 (b.a., 1H) (DMSO-d ₆)
34a	H	H	H	H	H	N	CH	CH=CH-CH=CH-C		HCl 193-194°C	2931, 2566, 742 KBr	1.80 (m, 2H); 1.91 (m, 2H); 2.67 (m, 1H); 2.88 (m, 1H); 3.10-3.20 (a.c., 3H); 3.52 (m, 1H); 3.71 (m, 1H); 3.90 (m, 1H); 4.46 (t, J=6.7Hz, 2H); 6.15 (s, 1H); 7.14 (t, J=7.5Hz, 1H); 7.25-7.41 (a.c., 4H); 7.46 (d, J=8.6Hz, 2H); 7.71 (d, J=8.6Hz, 1H); 7.75 (d, J=8.3Hz, 1H); 8.08 (s, 1H); 11.18 (b.a., 1H) (DMSO-d ₆)
35a	H	H	F	H	H	CCH ₃	N	Cl	CCl	HCl 160-161°C	2930, 2590, 1512, 1409, 1241, 827 KBr	1.67 (m, 2H); 1.79 (m, 2H); 2.33 (s, 3H); 2.67 (m, 1H); 2.90 (m, 1H); 3.10-3.25 (a.c., 3H); 3.54 (m, 1H); 3.72 (m, 1H); 3.85-3.98 (a.c., 3H); 6.13 (s, 1H); 7.19 (m, 2H); 7.50 (m, 2H); 11.28 (b.a., 1H) (DMSO-d ₆)
36a	H	H	H	H	H	N	CH	4-CIPh	CH	HCl 198-199°C	2472, 1560, 1450, 1095, 955, 810, 745 KBr	1.77 (m, 2H); 1.87 (m, 2H); 2.70 (m, 1H); 2.86 (m, 1H); 3.16 (a.c., 3H); 3.55 (m, 1H); 3.73 (m, 1H); 3.90 (m, 1H); 4.17 (t, J=6.6Hz, 2H); 6.15 (m, 1H); 7.25-7.47 (a.c., 7H); 7.59 (m, 2H); 7.90 (s, 1H); 8.27 (s, 1H); 10.91 (b.a., 1H) (DMSO-d ₆)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
37a	H	H	H	H	H	N	CH	4-CipH	CH	126-127°C	2935, 1570, 1493, 1455, 1379, 1091, 953, 815, 746 KBr	1.60 (m, 2H); 1.97 (m, 2H); 2.48 (t, J=7.3Hz, 2H); 2.56 (m, 2H); 2.67 (t, J=5.1Hz, 2H); 3.13 (m, 2H); 4.18 (t, J=7.1Hz, 2H); 6.05 (m, 1H); 7.23-7.40 (a.c., 9H); 7.61 (s, 1H); 7.74 (s, 1H) (CDCl ₃)
38a	H	H	F	H	H	CH	N	H	N	HCl 166-168°C	3450, 2429, 2707, 2593, 1512, 1437, 1230, 816, 626 KBr	1.74 (m, 2H); 1.86 (m, 2H); 2.68 (m, 1H); 2.84 (m, 1H); 3.16 (a.c., 3H); 3.53 (m, 1H); 3.70 (m, 1H); 3.91 (m, 1H); 4.27 (t, J=6.7Hz, 2H); 6.12 (s, 1H); 7.19 (t, J=8.9Hz, 2H); 7.50 (dd, J=8.9Hz, J'=5.5Hz, 2H); 8.23 (s, 1H); 8.93 (s, 1H); 11.02 (b a., 1H) (DMSO-d ₆)
39a	H	H	F	H	H	CH	N	H	N	oil	2944, 2808, 2773, 1602, 1510, 1273, 1227, 1161, 1140, 846, 824, 681 film	1.60 (m, 2H); 1.97 (m, 2H); 2.40-2.70 (a.c., 6H); 3.12 (m, 2H); 4.22 (t, J=6.9Hz, 2H); 5.99 (m, 1H); 6.98 (m, 2H); 7.35 (m, 2H); 7.95 (s, 1H); 8.07 (s, 1H) (CDCl ₃)
40a	H	H	F	H	H	CCH ₃	N	CH=CH-CH=CH-C	CH=CH-CH=CH-C	oil	2932, 1512, 1456, 1404, 1231, 744 film	1.63 (m, 2H); 1.88 (m, 2H); 2.42-2.55 (a.c., 4H); 2.61 (s, 3H); 2.65 (t, J=5.5Hz, 2H); 3.09 (m, 2H); 4.14 (t, J=7.3Hz, 2H); 5.97 (m, 1H); 6.99 (m, 2H); 7.19-7.35 (a.c., 5H); 7.68 (m, 1H) (CDCl ₃)
41a	H	H	F	H	H	N	CH	CH=CH-CH=CH-C	CH=CH-CH=CH-C	oil	2932, 2805,	1.57 (m, 2H); 1.99 (m, 2H); 2.42-2.50 (a.c., 4H); 2.62

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											1511, 1465, 1230, 1160, 825, 752, 741 film	(t,J=5.6Hz, 2H); 3.06 (m, 2H); 4.42 (t,J=6.9Hz, 2H); 5.95 (m, 1H); 6.97 (t,J=8.8Hz, 2H); 7.12 (m, 1H); 7.25- 7.41 (a.c., 4H); 7.71 (d,J=8Hz, 1H); 7.99 (s, 1H) (CDCl ₃)
42a	H	H	F	H	H	N	C-CH=CH-CH=CH	CH		102-103°C	2941, 1510, 1374, 1226, 1162, 806, 759, 741 KBr	1.59 (quin., J=7.0Hz, 2H); 2.09 (quin., J=7.5Hz, 2H); 2.40-2.50 (a.c., 4H); 2.64 (t,J=6.2Hz, 2H); 3.10 (m, 2H); 4.45 (t,J=7.1Hz, 2H); 5.96 (m, 1H); 6.98 (t,J=8.8Hz, 2H); 7.07 (t,J=7.6Hz, 1H); 7.20-7.35 (a.c., 3H); 7.63 (d,J=8.5Hz, 1H); 7.71 (d,J=8.6Hz, 1H); 7.90 (s, 1H) (CDCl ₃)
43a	H	H	F	H	H	N	C-CH=CH-CH=CH	N		HCl 208-209°C	2574, 2482, 1510, 1231, 745 KBr	1.80 (m, 2H); 2.11 (quin., J=7.2Hz, 2H); 2.69 (m, 1H); 2.83 (m, 1H); 3.10-3.20 (a.c., 3H); 3.52 (m, 1H); 3.71 (m, 1H); 3.88 (m, 1H); 4.80 (t,J=6.3Hz, 2H); 6.11 (s, 1H); 7.19 (m, 2H); 7.41 (m, 2H); 7.50 (m, 2H); 7.91 (m, 2H); 11.07 (b a, 1H) (DMSO-d ₆)
44a	H	H	F	H	H	N	C-CH=CH-CH=CH	N		76-77°C	2913, 1511, 1470, 1380, 1327, 1224, 1172, 1132, 851, 826, 757 KBr	1.60 (quin., J=7.5Hz, 2H); 2.19 (quin., J=8.2Hz, 2H); 2.41-2.59 (a.c., 4H); 2.64 (t,J=5.7Hz, 2H); 3.08 (m, 2H); 4.77 (t,J=7.0Hz, 2H); 5.95 (m, 1H); 6.97 (t,J=8.8Hz, 2H); 7.25-7.40 (a.c., 4H); 7.85 (m, 2H) (CDCl ₃)
45a	H	H	F	H	H	N	CH=CH-CH=CH-C			HCl 204-205°C	2928, 2680, 2573, 2559, 1515, KBr	1.81 (m, 2H); 1.99 (m, 2H); 2.67 (m, 1H); 2.84 (m, 1H); 3.10-3.20 (a.c., 3H); 3.53 (m, 1H); 3.72 (m, 1H); 3.90 (m, 1H); 4.76 (t,J=6.9Hz, 2H); 6.12 (s, 1H); 7.19

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											1454, 1272, 1242, 1224, 1166, 819, 745 KBr	(t,J=8.8Hz, 2H); 7.39 (t,J=7.6Hz, 1H); 7.45-7.60 (a.c., 3H); 7.94 (d,J=8.3Hz, 2H); 8.03 (d,J=8.3Hz, 2H); 11.04 (b.a., 1H) (DMSO-d ₆)
46a	H	H	F	H	H	N	N	CH=CH-CH=CH-C		88-90°C	2939, 1510, 1229, 1209, 1164, 744 KBr	1.58 (quin., J=7.5Hz, 2H); 2.07 (quin, J=7.5Hz, 2H); 2.40-2.50 (a.c., 4H); 2.61 (m, 2H); 3.05 (m, 2H); 4.66 (t,J=7.0Hz, 2H); 5.95 (m, 1H); 6.96 (t,J=8.8Hz, 2H); 7.23-7.38 (a.c., 3H); 7.44 (m, 1H); 7.52 (m, 1H); 8.04 (d,J=8.3Hz, 1H) (CDCl ₃)
47a	H	H	Cl	H	H	N	CH	Cl	CH	HCl 172-173°C	3068, 2948, 1491, 1445, 1320, 1308, 1096, 968, 809, 799 KBr	1.71 (m, 2H); 1.80 (m, 2H); 2.70 (m, 1H); 2.83 (m, 1H); 3.15-3.30 (a.c., 3H); 3.44 (m, 1H); 3.72 (m, 1H); 3.89 (m, 1H); 4.11 (t,J=6.5Hz, 2H); 6.20 (s, 1H); 7.41 (Syst. AB, J _{AB} =8.8Hz, 2H); 7.48 (Syst. AB, J _{AB} =8.8Hz, 2H); 7.52 (s, 1H); 8.04 (s, 1H); 10.98 (b.a., 1H) (DMSO-d ₆)
48a	H	H	H	H	H	N	CH	H	CH	HCl 180-181°C	2955, 2929, 2530, 1445, 965, 761, 745 KBr	1.70-1.90 (a.c., 4H); 2.69 (m, 1H); 2.89 (m, 1H); 3.10-3.20 (a.c., 3H); 3.53 (m, 1H); 3.70 (m, 1H); 3.91 (m, 1H); 4.15 (t,J=6.5Hz, 2H); 6.16 (m, 1H); 6.23 (m, 1H); 7.28-7.50 (a.c., 6H); 7.78 (m, 1H); 11.26 (b.a., 1H) (DMSO-d ₆)
49a	H	H	H	H	H	CH	N	H	N	HCl 122-123°C	2937, 2370, 1503, 1276, 1142, 774, 755	1.74 (m, 2H); 1.84 (m, 2H); 2.72 (m, 1H); 2.87 (m, 1H); 3.10-3.20 (a.c., 3H); 3.54 (m, 1H); 3.73 (m, 1H); 3.88 (m, 1H); 4.22 (t,J=6.6Hz, 2H); 6.15 (s, 1H); 7.27-7.70 (a.c., 3H); 7.47 (m, 2H); 7.97 (s, 1H); 8.59 (s, 1H);

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											KBr	11.20 (b.a., 1H) (DMSO-d ₆)
50a	H	H	H	H	H	CPh	N	H	CH	HCl 170-171°C	2930, 2554, 1469, 1459, 1444, 1278, 1075, 774, 762, 749, 732, 711, 702, 690	1.62-1.78 (a.c., 4H); 2.75 (m, 2H); 3.00 (m, 2H); 3.25 (m, 2H); 3.69 (m, 2H); 4.08 (t, J=6.7Hz, 2H); 6.13 (s, 1H); 7.07 (s, 1H); 7.24-7.40 (a.c., 3H); 7.42-7.52 (a.c., 6H); 7.62 (Syst. AB, J _{AB} =7.6Hz, 2H) (DMSO-d ₆)
51a	H	H	H	H	H	CH	CH	H	CH	HCl 197-199°C	2930, 2482, 1448, 1280, 1090, 732	1.60-1.80 (a.c., 4H); 2.70 (m, 1H); 2.84 (m, 1H); 3.08-3.22 (a.c., 3H); 3.50 (m, 1H); 3.71 (m, 1H); 3.86-3.96 (a.c., 3H); 5.97 (t, J=2.1Hz, 2H); 6.16 (m, 1H); 6.76 (t, J=2.1Hz, 2H); 7.25-7.50 (a.c., 5H); 10.74 (b.a., 1H) (DMSO-d ₆)
52a	H	H	H	H	H	CH	CH	H	CH	58-60°C	2928, 1498, 1280, 1262, 1137, 1087, 1060, 747, 723, 691	1.58 (m, 2H); 1.84 (m, 2H); 2.47 (t, J=7.5Hz, 2H); 2.58 (m, 2H); 2.68 (m, 2H); 3.13 (m, 2H); 3.92 (t, J=7.1Hz, 2H); 6.06 (m, 1H); 6.15 (t, J=2.2Hz, 2H); 6.67 (t, J=2.2Hz, 2H); 7.24-7.42 (a.c., 5H) (CDCl ₃)
53a	H	H	H	H	H	N	CCl	CH=CH-CH=CH-C	oil		2939, 1495, 1467, 1338, 745	1.58 (quin, J=7.6Hz, 2H); 1.99 (quin, J=7.6Hz, 2H); 2.47 (m, 2H); 2.55 (m, 2H); 2.65 (m, 2H); 3.10 (m, 2H); 4.36 (t, J=7.1Hz, 2H); 6.04 (m, 1H); 7.18-7.42 (a.c., 8H); 7.67 (d, J=7.6Hz, 1H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
54a	H	H	H	H	H	N	CCl	CH=CH-CH=CH-C		HCl 164-165°C	3460, 2940, 2550, 1338, 743 KBr	1.80 (m, 2H); 1.90 (m, 2H); 2.70 (m, 1H); 2.87 (m, 1H); 3.07-3.22 (a.c., 3H); 3.52 (m, 1H); 3.71 (m, 1H); 3.87 (m, 1H); 4.43 (t, J=6.6Hz, 2H); 6.14 (s, 1H); 7.20-7.52 (a.c., 7H); 7.65 (m, 1H); 7.79 (m, 1H); 11.16 (b.a., 1H) (DMSO-d ₆)
55a	H	H	OH	H	H	CCH ₃	N	Cl	CCl	HCl 216-217°C	3062, 2561, 1516, 1248 KBr	1.69 (m, 2H); 1.75 (m, 2H); 2.33 (s, 3H); 2.68 (m, 1H); 2.79 (m, 1H); 3.14 (a.c., 3H); 3.55 (m, 1H); 3.68 (m, 1H); 3.87-4.00 (a.c., 3H); 5.97 (s, 1H); 6.77 (Syst. AB, J=8.8Hz, 2H); 7.28 (Syst. AB, J=8.8Hz, 2H); 9.62 (s, 1H); 10.82 (b.a., 1H) (DMSO-d ₆)
56a	H	H	H	H	H	CH	N	Cl	CCl	HCl 166-167°C	2336, 1254 KBr	1.75 (a.c., 4H); 2.70 (m, 1H); 2.87 (m, 1H); 3.17 (a.c., 3H); 3.56 (m, 1H); 3.74 (m, 1H); 3.87-4.15 (a.c., 3H), 6.17 (s, 1H); 7.27-7.40 (a.c., 3H); 7.47 (m, 2H); 7.91 (s, 1H); 11.02 (b.a., 1H) (DMSO-d ₆)
57a	H	H	F	H	H	CH	N	H	N	Citrate 132-133°C	1720, 1709, 1513, 1225, 1193, 1166, 1133 KBr	1.90 (m, 2H); 2.08 (quint., J=7.5Hz, 2H); 2.86 (AB, J=15.5 Hz, 4H); 2.93 (b.a., 2H); 3.29 (m, 2H); 3.54 (t, J=5.9 Hz, 2H) 3.93 (b.a., 2H); 4.43 (t, J=6.6 Hz, 2H); 6.17 (b.a., 1H); 7.19 (m, 2H); 7.59 (m, 2H); 8.10 (s, 1H), 8.60 (s, 1H) (MeOH-d ₄)
58a	H	H	Br	H	H	CH	N	H	N		2939, 2773, 2736, 1509, 1490, 1380, 1271, 1140, 1071, KBr	1.55 (m, 2H); 1.95 (m, 2H); 2.40-2.55 (a.c., 4H); 2.64 (m, 2H); 3.08 (m, 2H); 4.20 (t, J=7.1 Hz, 2H); 6.03 (m, 1H); 7.22 (AB, J=8.5 Hz, 2H); 7.40 (AB, J=8.5 Hz, 2H); 7.93 (s, 1H); 8.05 (s, 1H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
59a	H	H	Br	H	H	CH	N	H	N	HCl 162-164°C	1006, 961, 844, 827, 800, 680 KBr 3066, 2937, 2479(b.a.) 1514, 1146, 1012, 802 KBr	1.76 (m, 2H); 1.84 (m, 2H); 2.71 (m, 1H); 2.85 (m, 1H); 3.17 (a.c., 3H); 3.55 (m, 1H); 3.74 (m, 1H); 3.80 (m, 1H); 4.23 (t, J=6.6 Hz, 2H); 6.22 (s, 1H); 7.42 (Syst. AB, J=8.1 Hz, 2H); 7.56 (Syst. AB, J=8.1 Hz, 2H); 7.98 (s, 1H); 8.60 (s, 1H) (DMSO-d ₆)
60a	H	H	Cl	H	H	CH	N	H	N	101-103°C	2930, 2775, 2737, 1509, 1493, 1381, 1271, 1141.1091, 1010, 961, 847, 828, 680 KBr	1.56 (quint, J=7.5 Hz, 2H); 1.97 (quint, J=7.5 Hz, 2H); 2.40-2.70 (a.c., 4H); 2.66 (t, J=5.7 Hz, 2H); 3.10 (d, J=3 Hz, 2H); 4.21 (t, J=7.0 Hz, 2H); 6.04 (s, 1H); 7.20-7.35 (m, 4H); 7.94 (s, 1H); 8.06 (s, 1H) (CDCl ₃)
61a	H	H	Cl	H	H	CH	N	H	N	HCl 165-166°C	2951, 2505 (b.a.) 1502, 1494, 1275, 1136, 1098, 1013, 810, 686 KBr	1.73 (m, 2H); 1.83 (m, 2H); 2.70 (m, 1H); 2.85 (m, 1H); 3.10-3.20 (a.c., 3H); 3.54 (m, 1H); 3.73 (m, 1H); 3.88 (m, 1H); 4.22 (t, J=6.6 Hz, 2H); 6.20 (s, 1H); 7.42 (Syst. AB, J=8.6 Hz, 2H); 7.49 (Syst. AB, J=8.6 Hz, 2H); 7.97 (s, 1H); 8.59 (s, 1H); 11.17 (b.a., 1H) (DMSO-d ₆)
62a	H	H	Cl	H	H	CPh	N	H	CH	oil	1445, 1379, 1271, 774, 681 film	1.48 (m, 2H); 1.80 (m, 2H); 2.36 (t, J=7.4 Hz, 2H); 2.47 (m, 2H); 2.59 (m, 2H); 3.04 (d, J=3 Hz, 2H); 4.03 (t, J=7.4 Hz, 2H); 6.01 (s, 1H); 7.01 (d, J=1.2 Hz, 1 H); 7.11 (d, J=1.2 Hz, 1H); 7.27 (m., 4H); 7.35-7.60 (a.c., 5H) (CDCl ₃)

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
63a	H	H	Cl	H	H	CPh	N	H	CH	HCl 70°C (hygrosc.)	2935, 2695, 2591, 1493, 1094, 777, 702 KBr	1.65 (m, 2H); 1.80 (m, 2H); 2.67 (m, 1H); 2.82 (m, 1H); 3.05-3.21 (a.c., 3H); 3.55 (m, 1H); 3.69 (m, 1H); 3.88 (m, 1H); 4.20 (t, J=6.6 Hz, 2H); 6.18 (s, 1H); 7.40 (Syst. AB, J=8.7 Hz, 2H); 7.47 (Syst. AB, J=8.7 Hz, 2H); 7.60- 7.80 (a.c., 5H); 7.83 (s, 1H); 7.96 (s, 1H) (DMSO-d ₆ + TFA)
64a	H	H	H	H	H	CCH ₃	CH	H	CCH ₃	oil	2929, 1408, 1299, 746, 693 film	1.65 (a.c., 4H); 2.23 (s, 6H); 2.48 (m, 2H); 2.58 (m, 2H); 2.69 (m, 2H); 3.15 (m, 2H); 3.76 (t, J=7.2 Hz, 2H); 5.76 (s, 2H); 6.06 (s, 1H); 7.20-7.40 (a.c., 5H) (CDCl ₃)
65a	H	H	H	H	H	CCH ₃	CH	H	CCH ₃	HCl 178-180°C	3434 (b.a.), 2935, 2560, 1443, 1405, 1298, 748, 692 KBr	1.56 (m, 2H); 1.77 (m, 2H); 2.15 (s, 6H); 2.70 (m, 1H); 2.84 (m, 1H); 3.08-3.22 (a.c., 3H); 3.59 (m, 1H); 3.70- 3.80 (a.c., 3H); 3.93 (m, 1H); 5.59 (s, 2H); 6.17 (s, 1H); 7.25-7.50 (a.c., 5H); 10.72 (b.a., 1H) (DMSO-d ₆)
66a	H	H	Cl	H	H	CCH ₃	CH	H	CCH ₃	86-88°C	2933, 1493, 1413, 1376, 1300, 750 film	1.65 (a.c., 4H); 2.24 (s, 6H); 2.48 (m, 2H); 2.54 (m, 2H); 2.69 (m, 2H); 3.15 (m, 2H); 3.77 (t, J=7.1 Hz, 2H); 5.77 (s, 2H); 6.06 (s, 1H); 7.30 (m, 4H) (CDCl ₃)
67a	H	H	Cl	H	H	CCH ₃	CH	H	CCH ₃	HCl 182-184°C	3432 (b.a.), 2936, 2570, 1495, 1410, 1298, 1097, 804, 752 KBr	1.56 (m, 2H); 1.76 (m, 2H); 2.14 (s, 6H); 2.70 (m, 1H); 2.84 (m, 1H); 3.00-3.28 (a.c., 3H); 3.58 (m, 1H); 3.69- 3.77 (a.c., 3H); 3.92 (m, 1H); 5.58 (s, 2H); 6.22 (s, 1H); 7.42 (AB, J=8.6, 2H); 7.50 (AB, J=8.6, 2H); 10.65 (b.a., 1H) (DMSO-d ₆)
68a	H	H	Cl	H	H	CH	CH	H	CH	102-104°C	2931, 1492,	1.56 (t, J=7.5 Hz, 2H); 1.84 (t, J=7.4 Hz, 2H); 2.46

Ex.	R ₁	R ₂	R ₃	R ₄	R ₅	Z ₁	Z ₂	R ₆	Z ₄	Salt / m.p.	IR cm ⁻¹	¹ H-RMN (300 MHz), δ (solvent)
											1280, 1090, 967, 828, 727 KBr	(t, J=7.5 Hz, 2H); 2.53 (m, 2H); 2.66 (t, J=5.6 Hz, 2H); 3.12 (m, 2H); 3.92 (t, J=7.1 Hz, 2H); 6.05 (m, 1H); 6.15 (d, J=1.8 Hz, 2H); 6.66 (d, J=1.8 Hz, 2H); 7.26 (AB, J=8.4, 2H); 7.30 (AB, J=8.4, 2H) (CDCl ₃)
69a	H	H	Cl	H	H	CH	CH	H	CH	HCl 194-195°C	2937, 2479, 1492, 1282, 1096, 810, 737 KBr	1.72 (m, 4H); 2.65 (m, 1H); 2.87 (m, 1H); 3.08-3.22 (a.c., 3H); 3.52 (m, 1H); 3.70 (m, 1H); 3.80-4.00 (a.c., 3H); 5.96 (t, J=2.1 Hz, 2H); 6.19 (s, 1H); 6.76 (t, J=2.1 Hz, 2H); 7.42 (AB, J=8.6 Hz, 2H); 7.48 (AB, J=8.6 Hz, 2H); 11.12 (b.a., 1H) (DMSO-d ₆)
70 ^a	H	H	Cl	H	H	CH	N	H	N	Citrate 133°C	3384 (b.a.), 3200-2200 (b.a.) 1726, 1702, 1594, 1432, 1221, 1131, 802 KBr	1.54 (m, 2H); 1.83 (m, 2H); 2.54 (Syst. AB, J=15 Hz, 2H); 2.63 (Syst. AB, J=15 Hz, 2H); 2.82 (m, 2H); 3.03 (m, 2H); 3.20-3.50 (a.c., 4H); 4.21 (t, J=6.8 Hz, 2H); 6.20 (s, 1H); 7.40 (Syst. AB, J=8.8 Hz, 2H); 7.48 (Syst. AB, J=8.8 Hz, 2H); 7.97 (s, 1H); 8.52 (s, 1H) (DMSO-d ₆)

EBiological assays

The analgesic activity of the products object of the invention have been studied in several assays using the Swiss albino mice as the experimental animal. The assay of contortions induced by phenylbenzoquinone, the hot-plate assay and the hot-point assay are now described. The examples that are presented by way of illustration describe some of the pharmacological assays and should not limit the scope of the invention in any way.

10 The assay of contortions induced by phenylbenzoquinone was carried out following the method described by E. Siegmund *et al.* (*Proc. Soc. Exp. Biol. Med.* 95 : 729-731, 1957). In this assay the mice received the product orally or sub-cutaneously (s.c.) and after 60 minutes (after oral administration) or after 30 minutes (after s.c. administration) they received
15 an intraperitoneal (i.p.) injection of an 0.02% aqueous solution of phenylbenzoquinone, at a dosage of 10 ml/kg. The degree of analgesic was expressed as a percentage of the contortions with respect to the control group at each one of the dosages assayed. Using the results obtained the effective dose-50 (ED-50) was calculated, that is to say the
20 dose able to inhibit by 50% the contortions induced by phenylbenzoquinone.

The hot-plate assay was carried out following the method described by M. Ocaña *et al.* (*Europ.J.Pharmacol.* 186 : 377-378, 1990). The product under study was administered s.c. or i.p. and 30 minutes later the
25 analgesic effect was registered. For this the animals were placed on a metallic surface kept at 50° C or 55° C and the time registered (latency) until the licked their hind legs and a jump. The analgesic activity was calculated at each dose, comparing the potency of the treated group with the control group. Using the results obtained the ED-50 was calculated.

30 The assay of withdrawal of the tail from a hot spot (tail flick) was carried out following the method described by M. Ocaña *et al.* (*Br. J. Pharmacol.* 110 : 1049-1054, 1993). The mice were introduced into an immobiliser and placed on the tail-flick apparatus (LI7100, Letica, S.A). A beam of light was focussed on the tail, at 4 cm from the tip, and the latency

for withdrawal of the tail automatically registered. Ten minutes before administering the product of the study the basal latency was registered. After the product had been administered s.c. the tail withdrawal latencies at 10, 20, 30, 40, 45, 60, 90 and 120 minutes were registered. For each
5 animal the area below curve of the latency was calculated during the time period following the method described by R.J. Tallarida and R.B. Murray (Manual of pharmacologic calculations with computer programs, Springer-Verlag, Berlin, p. 297, 1987). The degree of analgesic of each dosage was calculated comparing the area under curve of latency of the group treated
10 with the medicament with the control group. Using these data the ED-50 was calculated.

The products object of the invention have a notable analgesic activity in the assay of contortions induced by phenylbenzoquinone. Several products have activity of the same order as morphine and clearly
15 better than the products that inhibit biosynthesis of prostaglandins, such as aspirin and dipirone (see table 3).

The analgesic activity has also been demonstrated in the hot-plate assay, considered as a demonstration of analgesic action at the central nervous system level (see table 4).

20 The analgesic activity has also been demonstrated in the hot-beam assay applied to the mouse's tail, finding a good correspondence between the results obtained in the assay of the hot-plate and the assay of the calorific beam (see table 5).

Furthermore, the capacity of the products object of the invention for
25 exhibiting synergistic analgesic activity with other analgesics, for example, pentazocine, has also been demonstrated. This has been shown for the compound of example 47a (see table 6). Effectively, the latency time in responding licking of the paws when the mice were placed on the hot-plate at 55° C, is much greater after the combined treatment with the compound
30 of example 47a and pentazocine than the sum of the latencies of each one of the treatments carried out separately.

In summary, the products object of the invention have shown a clear analgesic activity in different assays, such as phenylbenzoquinone, hot-

plate and calorific beam applied to the tail of the mouse. The activity of these products has been clearly superior to that of the inhibitors of the biosynthesis of prostaglandins such as aspirin and dipirone, and the activity has been shown to of the order of that of morphine. Furthermore, the capacity for forming synergistic combinations with other analgesics has been demonstrated, as can be seen for the case of the compound of example 47a administered along with pentazocine in the hot-plate assay for mice.

10 Table 3. Analgesic activity in the assay of contortions induced by phenylbenzoquinone in mice

PRODUCT	ED-50 (mg/kg)	
	Oral route	S.C. route
Example 5	20	28
Example 6	80	34
Example 33a	30	2
Example 35a	37	1
Example 38a	5	1
Example 41a	58	6
Example 47a	19	26
Example 48a	38	1
Example 49a	2	1
Example 50a	10	2
Example 51a	9	3
Example 59a	13	2
Example 61a	22	2
Example 63a	44	33
Morfine	4	1
Dipirone	223	24
Aspirin	100	80

Table 4.- Analgesic activity in the hot-plate assay (55°C) in mice

PRODUCT	ED-50 (mg/kg, sc)
Example 38a	7
Example 47a	89
Example 48a	5
Example 49a	4
Example 50a	58
Example 51a	2
Example 59a	43
Example 61a	48
Morfina	2

Table 5.- Analgesic activity in the calorific beam assay in mouse's tail.

PRODUCT	ED-50 (mg/kg, sc)
Example 51a	5
Example 59a	60
Example 63a	70
Morfine	4


5

Table 6.- Analgesic activity in the hot plate assay in mouse.

PRODUCT	DOSE (mg/kg, ip)	Δ Latency (Seconds)
Example 47a	40	5
Pentazocine	10	6
Example 47a	40	
+	+	20
Pentazocine	10	

CLAIMS

1. Use of a derivative of tetrahydropyridine (or 4-hydroxypiperidine)butylazole of general formula (I)



where

R₁, R₂ and R₃ are either identical or different and represent a hydrogen atom, a halogen atom, a C₁-C₄ alkyl group, a trifluoromethyl radical, a hydroxyl or alkoxy radical, and furthermore, two adjacent radicals can form part of an six-member aromatic ring;

A represents a carbon atom and the dotted line represents an additional bond, or A represents a carbon atom bound to a hydroxyl group (C-OH) and the dotted line represents the lack of an additional bond;

Z₁ represents a nitrogen atom or a substituted carbon atom that can be represented by C-R₄;

Z₂ represents a nitrogen atom or a substituted carbon atom that can be represented by C-R₅;

Z₄ represents a nitrogen atom or a substituted carbon atom that can be represented by C-R₇;

with the condition that Z₁, Z₂ and Z₄ taken together can represent, at most, two nitrogen atoms; and

R₄, R₅, R₆ and R₇, are identical or different and represent a hydrogen atom, a halogen atom, a C₁-C₄ alkyl group, an aryl or substituted aryl group, or two adjacent radicals can form part of a six-member aromatic ring;

or physiologically acceptable salts thereof,

in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.

2. Use according to claim 1, of a compound of general formula (I), wherein R_1 , R_2 and R_3 are chosen from a hydrogen atom, a fluorine atom, a chlorine atom, a bromine atom, a trifluoromethyl radical, a hydroxyl radical, a methoxyl radical, a methyl radical, an ethyl radical, a propyl radical, an isopropyl radical, a sec-butyl radical and a terc-butyl radical, or a physiologically acceptable salt thereof, in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.

3. Use according to claim 1, of a compound of general formula (I), wherein R_1 , R_2 and R_3 are chosen in such a way that two adjacent radicals can form part of a six-member aromatic ring, or a physiologically acceptable salt thereof, in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.

4. Use according to claim 1, of a compound of general formula (I), wherein R_4 , R_5 , R_6 and R_7 are chosen from a hydrogen atom, a fluorine atom, a chlorine atom, a bromine atom, a methyl radical, an ethyl radical, a propyl radical, an isopropyl radical, a butyl radical, an iso-butyl radical, a sec-butyl radical and a terc-butyl radical, a phenyl radical, a phenyl radical substituted by an atom of fluorine, chlorine or bromine, or a physiologically acceptable salt thereof, in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.

5. Use according to claim 1, of a compound of general formula (I), wherein R_4 , R_5 , R_6 and R_7 are chosen in such a way that two adjacent radicals can form part of a six-member aromatic ring, or a physiologically acceptable salt thereof, in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.

6. Use according to claim 1, of a compound of general formula (I), chosen from the group formed by:

- [1] 4-chloro-1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-pirazole
- [2] 4,5-dichloro-1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-2-methyl-
5 1*H*-imidazole
- [3] 1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-benzimidazole
- [4] 1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-1,2,4-triazole
- [5] 4-chloro-1-{4-[4-(4-chlorophenyl)-4-hydroxy-1-piperdiny]]butyl}-1*H*-
pirazole
- 10 [6] 4,5-dichloro-1-{4-[4-hydroxy-4-(4-chlorophenyl)-1-piperdiny]]butyl}-2-
methyl-1*H*-imidazole
- [7] 4-chloro-1-{4-[4-hydroxy-4-(3-trifluoromethylphenyl)-1-piperdiny]]-
butyl}-1*H*-pirazole
- [8] 4,5-dichloro-1-{4-[4-hydroxy-4-(3-trifluoromethylphenyl)-1-piperdiny]]-
15 butyl}-2-methyl-1*H*-imidazole
- [9] 4,5-dichloro-1-{4-[4-(4-fluorophenyl)-4-hydroxy-1-piperdiny]]butyl}-
2-methyl-1*H*-imidazole
- [10] 1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-indole
- [11] 4,5-dichloro-1-{4-[4-hydroxy-4-(4-methylphenyl)-1-piperdiny]]-
20 butyl}-2-methyl-1*H*-imidazole
- [12] 1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-pirazole
- [13] 1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-indazole
- [14] 2-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-2*H*-indazole
- [15] 4-chloro-1-{4-[4-hydroxy-4-(4-methylphenyl)-1-piperdiny]]-
25 butyl}-1*H*-pirazole
- [16] 4-chloro-1-{4-[4-hydroxy-4-(4-methoxyphenyl)-1-piperdiny]]-
butyl}-1*H*-pirazole
- [17] 1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-2-phenyl-1*H*-imidazole
- [18] 1-{4-[4-hydroxy-4-(4-methylphenyl)-1-piperdiny]]butyl}-1*H*-
30 benzimidazole
- [19] 4,5-diphenyl-1-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl]-1*H*-imidazole
- [20] 4-chloro-1-{4-[4-hydroxy-4-(1-naphthyl)-1-piperdiny]]butyl}-1*H*-pirazole
- [21] 4-chloro-1-{4-[4-hydroxy-4-(2-naphthyl)-1-piperdiny]]butyl}-1*H*-pirazole

- [22] 4-chloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-pirazole
- [23] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-benzimidazole
- 5 [24] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-1,2,4-triazole
- [25] 4-chloro-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pirazole
- [26] 4,5-dichloro-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-2-methyl-1*H*-imidazole
- 10 [27] 4-chloro-1-{4-[4-(3-trifluoromethylphenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-pirazole
- [28] 4,5-dichloro-2-methyl-1-{4-[4-(3-trifluoromethylphenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-imidazole
- [29] 4-chloro-1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-pirazole
- 15 [30] 4,5-dichloro-1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-2-methyl-1*H*-imidazole
- [31] 4,5-dichloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-2-methyl-1*H*-imidazole
- 20 [32] 4,5-dichloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2-methyl-1*H*-imidazole hydrochloride.
- [33] 4,5-dichloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2-methyl-1*H*-imidazole dihydrochloride.
- [34] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-indole
- 25 [35] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-indole
- [36] 4,5-dichloro-2-methyl-1-{4-[4-(4-methylphenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-imidazole
- [37] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pirazole
- [38] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-indazole
- 30 [39] 2-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2*H*-indazole
- [40] 4-chloro-1-{4-[4-(4-methylphenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-pirazole

- [41] 4-chloro-1-{4-[4-(4-methoxyphenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-pirazole
- [42] 4-chloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]propyl}-1*H*-pirazole
- [43] 4,5-dichloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]propyl}-2-methyl-
5 1*H*-imidazole
- [44] 2-phenyl-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-imidazole
- [45] 1-{4-[4-(4-methylphenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-
benzimidazole
- [46] 4,5-diphenyl-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-
10 imidazole
- [47] 4-chloro-1-{4-[4-(1-naphthyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-
pirazole
- [48] 4-chloro-1-{4-[4-(2-naphthyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-
pirazole
- [49] 1-{2-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]ethyl}-1*H*-
15 benzimidazole.
- [50] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-benzimidazole
hydrochloride.
- [51] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-
20 benzimidazole.
- [52] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-ben-
zimidazole hydrochloride.
- [53] 4,5-dichloro-2-methyl-1-{4-[4-(3-trifluoromethylphenyl)-1-(1,2,3,6-tetra-
hydropyridinyl)]butyl}-1*H*-imidazole hydrochloride.
- [54] 4-chloro-1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-
25 pirazole hydrochloride.
- [55] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-indazole
hydrochloride.
- [56] 4,5-dichloro-1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-2-
30 methyl-1*H*-imidazole hydrochloride.
- [57] 4-(4-chlorophenyl)-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]-butyl}-1*H*-
pirazole hydrochloride.

- [58] 4-(4-chlorophenyl)-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pirazole.
- [59] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-triazole hydrochloride.
- 5 [60] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-triazole.
- [61] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2-methyl-1*H*-benzimidazole.
- [62] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-indazole.
- [63] 2-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2*H*-indazole.
- 10 [64] 2-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2*H*-benzotriazole hydrochloride.
- [65] 2-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-2*H*-benzotriazole.
- [66] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-benzotriazole hydrochloride.
- 15 [67] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-benzotriazole.
- [68] 4-chloro-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pirazole hydrochloride.
- 20 [69] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pirazole hydrochloride.
- [70] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-triazole hydrochloride.
- [71] 2-phenyl-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-imidazole hydrochloride.
- 25 [72] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pyrrol hydrochloride.
- [73] 1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1*H*-pyrrol.
- [74] 4-(4-chlorophenyl)-1-{4-[4-(4-hydroxy-4-phenyl-1-piperdiny)]butyl}-1*H*-pirazole.
- [75] 1-{4-[4-(4-fluorophenyl)-4-hydroxy-1-piperdiny)]butyl}-1*H*-benzimidazole.
- 30 [76] 4-chloro-1-{4-[4-(4-hydroxy-4-(3-trifluoromethylphenyl)-1-piperdiny)]butyl}-1*H*-pirazole.
- [77] 1-{4-[4-(4-fluorophenyl)-4-hydroxy-1-piperdiny)]butyl}-1*H*-indazole.
- [78] 2-{4-[4-(4-fluorophenyl)-4-hydroxy-1-piperdiny)]butyl}-2*H*-indazole.

- [79] 2-{4-[4-(4-fluorophenyl)-4-hydroxy-1-piperdiny]]butyl}-2*H*-benzotriazole.
- [80] 1-{4-[4-(4-fluorophenyl)-4-hydroxy-1-piperdiny]]butyl}-1*H*-benzotriazole.
- [81] 3-chloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-indazole.
- [82] 3-chloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-indazole
5 hydrochloride.
- [83] 1-{4-[4-hydroxy-4-(4-fluorophenyl)-1-piperdiny]]butyl}-1*H*-triazole.
- [84] 1-{4-[4-hydroxy-4-(4-chlorophenyl)-1-piperdiny]]butyl}-1*H*-triazole.
- [85] 4,5-dichloro-1-{4-[4-(4-hydroxyphenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-
2-methyl-1*H*-imidazole hydrochloride.
- 10 [86] 4,5-dichloro-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-
imidazole hydrochloride.
- [87] 1-{4-[4-(4-fluorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-triazole
citrate.
- [88] 1-{4-[4-(4-bromophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-triazole.
- 15 [89] 1-{4-[4-(4-bromophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-triazole
hydrochloride.
- [90] 1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-triazole.
- [91] 1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-triazole
hydrochloride.
- 20 [92] 2-phenyl-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-
imidazole.
- [93] 2-phenyl-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-
imidazole hydrochloride.
- [94] 2,5-dimethyl-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-pyrrol.
- 25 [95] 2,5-dimethyl-1-{4-[4-phenyl-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-pyrrol
hydrochloride.
- [96] 2,5-dimethyl-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-
1*H*-pyrrol.
- [97] 2,5-dimethyl-1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-
30 1*H*-pyrrol hydrochloride.
- [98] 1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-pyrrol.
- [99] 1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridiny)]butyl}-1*H*-pyrrol
hydrochloride.

[100] 1-{4-[4-(4-chlorophenyl)-1-(1,2,3,6-tetrahydropyridinyl)]butyl}-1H-triazole
citrate

in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.

7. Use according to claim 1, of a compound of general formula (I), or a physiologically acceptable salt thereof, along with another analgesic of those normally used in the treatment of pain, in the preparation of a medicament for the treatment of acute pain, neuropathic pain or nociceptive pain in mammals, including man.