

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
11 March 2004 (11.03.2004)

PCT

(10) International Publication Number
WO 2004/019940 A1

(51) International Patent Classification⁷: **A61K 31/4375**,
31/502, A61P 9/10, 31/12, 31/20

W. [US/US]; 306 Briar Bluff, Thousand Oaks, CA 91360-2734 (US). **WATHEN, Lynne, K.** [US/US]; 306 Briar Bluff, Thousand Oaks, CA 91360-2734 (US).

(21) International Application Number:
PCT/US2003/026838

(74) Agents: **WILLIAMS, JR., Sidney, B.** et al.; Flynn, Thiel, Boutell & Tanis, P.C., 2026 Rambling Road, Kalamazoo, MI 49008-1631 (US).

(22) International Filing Date: 28 August 2003 (28.08.2003)

(25) Filing Language: English

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(26) Publication Language: English

(30) Priority Data:
60/407,563 30 August 2002 (30.08.2002) US
60/469,629 9 May 2003 (09.05.2003) US

(71) Applicant (*for all designated States except US*): **PHARMACIA & UPJOHN COMPANY** [US/US]; 301 Henrietta Street, Kalamazoo, MI 49001 (US).





(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **WATHEN, Michael**,

[Continued on next page]

(54) Title: METHOD OF PREVENTING OR TREATING ATHEROSCLEROSIS OR RESTENOSIS



(57) Abstract: The present invention provides a method of preventing or treating atherosclerosis or restenosis in mammals, which comprises administering an effective amount of a compound selected from the group consisting of structures of Formulas VI, VII, VIII and IX: wherein the various groups in the structure are as defined herein.

WO 2004/019940 A1



Published:

- *with international search report*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

HETEROCYCLIC CARBOXAMIDES FOR THE TREATMENT OF ATHEROSCLEROSIS AND RESTENOSIS

BACKGROUND OF THE INVENTION

This invention relates to a method of preventing or treating atherosclerosis and restenosis in mammals.

Atherosclerosis is characterized by the deposition of fatty substances in and fibrosis of the inner layer of the arteries. Restenosis is an accelerated form of atherosclerosis that commonly occurs after angioplastic surgery and atherectomy.

Cardiovascular diseases (CVD) contribute substantially to illness and death worldwide and ranks second only to infectious and parasitic diseases as human affliction. Atherosclerosis, a major component of CVD, has properly been considered a public health problem of industrialized countries, accounting for an estimated one third of deaths overall. It has been reported that in the United States alone, atherosclerosis affects one in four persons, causing approximately 42% of all deaths. O'connor et al, "Potential Infectious Etiologies of Atherosclerosis: A Multifactorial Perspective", Emerging Infectious Disease, Vol. 7, No. 5, September-October 2001.

It has been suggested that the number of chronic infective pathogens which an individual has been exposed independently contribute to the long-term prognosis in patients with documented coronary artery disease. HJ Rupprecht et al, "Impact of Viral and Bacterial Infective Burden on Long-term Prognosis in Patients with Coronary Artery Disease. (Circulation (2001) 104:25-31. Seropositivity to multiple herpesviruses is an independent risk factor for death from cardiovascular disease and risk is proportional to the number of different herpesviruses that have infected an individual.

Other investigators that have suggested a connection between infectious pathogens and atherosclerosis include Espinola-Klein et al, "Impact of Infectious Burden on Extent and Long-Term Prognosis of Atherosclerosis", Circulation (2002) 105:15-21; O'Connor et al, Supra: and Zhou et al, "Association Between Prior Cytomegalovirus Infection and the Risk of Restenosis after Coronary Atherectomy", The New England Journal of Medicine (1996). An antiviral drug, Ganciclovir, has been shown to prevent atherosclerosis resulting from CMV infection of rats (K.B. Lemstrom et al. Cytomegalovirus infection-enhanced allograft arteriosclerosis is prevented by DHPG prophylaxis in the rat. Circulation, 1994,90:1969-1978).

Herpesviruses are believed to be a particular problem in atherosclerosis because they reside latently in an infected individual and can reactivate causing a chronic inflammatory response.

The herpesvirus family contains eight known human viruses; herpes simplex virus type 1 (HSV-1), herpes simplex virus type 2 (HSV-2), varicella zoster virus (VZV), human cytomegalovirus (HCMV), human herpes virus 6 (HHV-6), human herpes virus 7 (HHV-7), Epstein-barr virus (EBV) and human herpes virus 8 (HHV-8). One of the hallmarks of herpesviruses is their ability to establish latent infections in their host and to recur during times of stress or immunosuppression. The human herpesviruses are associated with a diverse set of diseases ranging in severity from mild cold sores to life-threatening illness in immunocompromised patients (Table 1).

Table 1. Herpesvirus diseases and treatment

| Virus | Associated Diseases | | Marketed Antivirals |
|-------|------------------------------|---|--|
| | Normal Host | Immunocompromised Host | |
| HSV-1 | Herpes labialis (cold sores) | Disseminated herpes | <ul style="list-style-type: none"> • Acyclovir • Penciclovir |
| HSV-2 | Genital herpes | Disseminated herpes | <ul style="list-style-type: none"> • Acyclovir • Valaciclovir • Famciclovir |
| VZV | Chicken pox Herpes zoster | Herpes zoster | <ul style="list-style-type: none"> • Acyclovir • Valaciclovir • Famciclovir |
| CMV | Congenital CMV disease | Retinitis Pneumonia GI disease Graft rejection | <ul style="list-style-type: none"> • Ganciclovir • Valganciclovir • Foscarnet • Cidofovir • Formivirsen |
| EBV | Infectious mononucleosis | Lymphomas (PTLD) | <ul style="list-style-type: none"> • None |
| HHV-6 | Exanthem subitum | Graft rejection | <ul style="list-style-type: none"> • None |
| HHV-7 | Exanthem subitum | Graft rejection | <ul style="list-style-type: none"> • None |
| HHV-8 | Kaposi's sarcoma | Kaposi's sarcoma | <ul style="list-style-type: none"> • None |

HSV-1, HCMV, VZV and EBV are ubiquitous viruses with seroprevalence rates in adults of 70-80% for HSV-1 and 90-100% for HCMV, VZV and EBV. Seroprevalence of HSV-2 increases from about 10% in young adults to 35% by age 60. Antibodies to HHV-8 are also found in about 33% of adults in the United States. The high seroprevalence of multiple viruses and their ability to reactivate from latent infections, make these herpesviruses prime candidates for causing chronic inflammatory responses leading to atherosclerosis.

Numerous studies and articles on the epidemiology of the herpesvirus family are in the prior art. Wathen, Michael W., "Non-nucleoside inhibitor of herpesviruses",

Rev. Med. Virol, 2002; 12: 167-178; Whitley et al, "Herpes Simplex Viruses", Clinical Infection Diseases, 1998; 26: 541-55, Cohen, Jeffrey I., "Epstein-Barr Virus Infection", Medical Progress, Volume 343, Number 7, The New England Journal of Medicine, August 17, 2000, pp. 481-492; Blouvelt et al; "Human Herpes Virus 8 Infection Occurs Following Adolescence in the United States", The Journal of Infectious Disease, 1997, 176: 771-4; Field, A. Kirk, "Human Cytomegalovirus: challenge opportunities and new drug development", Antiviral Chemistry and Chemotherapy 10: 219-232.

INFORMATION DISCLOSURE

U.S. Patent 6 239 142 discloses 4-oxo-4,7-dihydro-thieno[2,3-b]pyridine-5-carboxamide derivatives, compounds of Formula I and I' that are useful as antiviral agents. These compounds have now been found to be useful in the method of this invention.

U.S. Patent 6 291 437 describes a method for preventing or retarding the development of atherosclerotic lesions or restenosis comprising administering to a subject, preferably a human, an effective amount of an anti-viral composition directed against CMV, and optionally anti-microbial composition directed against *C. pneumoniae*.

WO 02/48148 A2 discloses anti-viral compounds and a method of using them for the prophylaxis or treatment of atherosclerosis, coronary artery disease or restenosis.

An antiviral drug, Ganciclovir, has been shown to prevent atherosclerosis resulting from CMV infection of rats (K.B. Lemstrom et al. Cytomegalovirus infection-enhanced allograft arteriosclerosis is prevented by DHPG prophylaxis in the rat. Circulation, 1994, 90:1969-1978).

U.S. Patent 6 239 142 disclosed compounds and their use to treat herpesvirus infections.

WO 02/06513 disclosed method of screening 4-hydroxyquinoline, 4-oxo-dihydroquinoline, and 4-oxo-dihydrothienopyridine derivatives as non-nucleoside herpesvirus DNA polymerase inhibitors.

EP 443568 disclosed fused thiophene derivatives, their production and use.

WO 02/04445 disclosed a variety of tricyclic core structures which have antiviral activity against herpesviruses.

WO 02/04444, WO 02/04443, and WO 02/04422 disclosed a variety of bicyclic core structures which have antiviral activity against herpesviruses.

U.S. Patent 6 248 739 disclosed compounds in which the core structure is a quinoline and useful as antivirals against herpesviruses.

OBJECT OF THE INVENTION


It is the object of this invention to provide a method for preventing or treating atherosclerosis or restenosis in mammals.

It is a further objective of this invention to provide a method for prophylaxis of atherosclerosis and treat patients who have atherosclerosis.

It is still a further objective of the invention to provide a method that prevents or ameliorates the occurrence of restenosis in patients anticipating coronary atherosclerosis or angioplasty.

SUMMARY OF THE INVENTION

A method of preventing or treating atherosclerosis or restenosis in a mammal, comprising administering to said mammal an effective amount of a compound selected from the group consisting of structures Formula VI, Formula VII, Formula VIII and Formula IX, wherein Formula VI is:



or a pharmaceutically acceptable salt thereof wherein,
 A^{VI} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO_2 , or
- e) F;

R^{VI-1} is

- a) R^{VI-5} , or
- b) SO_2R^{VI-9}

R^{VI-2} , R^{VI-3} and R^{VI-4} may be the same or different and are selected from the group consisting of:

- a) H,
- b) $halo^{VI}$,
- c) $aryl^{VI}$,
- d) $S(O)_mR^{VI-6}$,
- e) $(C=O)R^{VI-6}$,
- f) $(C=O)OR^{VI-9}$,
- g) cyano,
- h) het^{VI} , wherein said het^{VI} is bound via a carbon atom,
- i) OR^{VI-10} ,
- j) $Ohet^{VI}$,
- k) $NR^{VI-7}R^{VI-8}$
- l) SR^{VI-10} ,
- m) $Shet^{VI}$,
- n) $NHCOR^{VI-12}$,
- o) $NHSO_2R^{VI-12}$,

- p) C₁₋₇alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{VI-11}, OR^{VI-13}, SR^{VI-10}, SR^{VI-13}, NR^{VI-7}R^{VI-8}, halo, (C=O)C₁₋₇alkyl, or SO_mR^{VI-9}, and
- q) R^{VI-3} together with R^{VI-2} or R^{VI-4} form a carbocyclic or ^{VI}het which may be optionally substituted by NR^{VI-7}R^{VI-8}, or C₁₋₇alkyl which may be optionally substituted by OR^{VI-14};

R^{VI-5} is

- a) (CH₂CH₂O)_iR^{VI-10},
- b) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of NR^{VI-7}R^{VI-8}, R^{VI-11}, SO_mR^{VI-9}, or OC₂₋₄alkyl which may be further substituted by het^{VI}, OR^{VI-10}, or NR^{VI-7}R^{VI-8}, or
- c) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11}, NR^{VI-7}R^{VI-8}, SO_mR^{VI-9}, or C₁₋₇alkyl optionally substituted by R^{VI-11}, NR^{VI-7}R^{VI-8}, or SO_mR^{VI-9};

R^{VI-6} is

- a) C₁₋₇alkyl,
- b) NR^{VI-7}R^{VI-8},
- c) aryl^{VI}, or
- d) het^{VI}, wherein said het^{VI} is bound via a carbon atom;

R^{VI-7} and R^{VI-8} are independently

- a) H,
- b) aryl^{VI},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting

of aryl^{VI}, NR^{VI-10}R^{VI-10}, R^{VI-11}, SO_mR^{VI-9},
CONR^{VI-10}R^{VI-10}, or halo, or;

- d) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11}, NR^{VI-7}R^{VI-8}, SO_m^{VI}R^{VI-9}, or C₁₋₇alkyl optionally substituted by R^{VI-11}, NR^{VI-7}R^{VI-8}, or SO_m^{VI}R^{VI-9}, or
- e) R^{VI-7} and R^{VI-8} together with the nitrogen to which they are attached form a het^{VI};

R^{VI-9} is

- a) aryl^{VI},
- b) het^{VI},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of NR^{VI-10}R^{VI-10}, R^{VI-11}, SH, CONR^{VI-10}R^{VI-10}, or halo;

R^{VI-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VI-11} is

- a) OR^{VI-10},
- b) Ohet^{VI},
- c) Oaryl^{VI},
- d) CO₂R^{VI-10},
- e) het^{VI},
- f) ^{VI}-aryl^{VI},
- g) CN, or
- h) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11}, NR^{VI-7}R^{VI-8}, SO_m^{IV}R^{VI-9}, or

C₁₋₇alkyl optionally substituted by R^{VI-11}, NR^{VI-7}
R^{VI-8}, or SO_mR^{VI-9};

R^{VI-12} is

- a) H,
- b) het^{VI},
- c) aryl^{VI},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VI-7}R^{VI-8}
or R^{VI-11};

R^{VI-13} is

- a) (P=O)(OR^{VI-14})₂,
- b) CO(CH₂)_n^{IV}CON(CH₃)-(CH₂)_nSO₃⁻M^{VI+},
- c) an amino^{VI} acid,
- d) C(=O)aryl^{VI},
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VI-7}
R^{VI-8}, aryl^{VI}, het^{VI}, CO₂H, or O(CH₂)_nCO₂R^{VI-14}, or
- f) C(=O)NR^{VI-7}R^{VI-8}

R^{VI-14} is

- a) H, or
- b) C₁₋₇alkyl;

each i^{VI} is independently 2, 3, or 4;

each n^{VI} is independently 1, 2, 3, 4 or 5;

each m^{VI} is independently 0, 1, or 2;

M^{VI} is sodium, potassium, or lithium;

aryl^{VI} is a phenyl radical or an ortho-fused bicyclic
carbocyclic radical wherein at least one ring is
aromatic;


wherein any aryl^{VI} is optionally substituted with one or
more substituents selected from the group consisting
of halo, OH, cyano, CO₂R^{VI-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆
alkyl which maybe further substituted by one to
three SR^{VI-14}, NR^{VI-14}R^{VI-14}, OR^{VI-14}, or CO₂R^{VI-14};

het^{VI} is a four- (4), five- (5), six- (6), or seven- (7)

membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VI} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VI-14}}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which maybe further substituted by one to three $\text{SR}^{\text{VI-14}}$, $\text{NR}^{\text{VI-14}}\text{R}^{\text{VI-14}}$, $\text{OR}^{\text{VI-14}}$, or $\text{CO}_2\text{R}^{\text{VI-14}}$;

wherein Formula VII is



VII

or a pharmaceutically acceptable salt thereof,

wherein

A^{VII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VII-1} is

- a) aryl^{VII},
- b) $\text{S}(\text{O})_m^{\text{VII}}\text{R}^{\text{VII-6}}$,
- c) $(\text{C}=\text{O})\text{R}^{\text{VII-6}}$, with the proviso that if R^{VII-6} is $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, then R^{VII-7} and R^{VII-8} do not both equal H,
- d) $(\text{C}=\text{O})\text{OR}^{\text{VII-9}}$,
- e) cyano,

- f) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- g) Ohet^{VII} ,
- h) $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$ with the proviso that $\text{R}^{\text{VII-7}}$ and $\text{R}^{\text{VII-8}}$ do not both equal H,
- i) $\text{SR}^{\text{VII-10}}$,
- j) Shet^{VII} ,
- k) $\text{NHCOR}^{\text{VII-12}}$,
- l) $\text{NHSO}_2\text{R}^{\text{VII-12}}$,
- m) C_{1-7} alkyl which is partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VII-11}}$, $\text{OR}^{\text{VII-13}}$, $\text{SR}^{\text{VII-10}}$, $\text{SR}^{\text{VII-13}}$, $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m\text{R}^{\text{VII-9}}$, or
- n) C_{1-7} alkyl which is substituted by one or more substituents of the group $\text{R}^{\text{VII-11}}$, $\text{OR}^{\text{VII-13}}$, $\text{SR}^{\text{VII-10}}$, $\text{SR}^{\text{VII-13}}$, $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m\text{R}^{\text{VII-9}}$;

$\text{R}^{\text{VII-2}}$ is

- a) H,
- b) halo,
- c) aryl^{VII},
- d) $\text{S}(\text{O})_m\text{R}^{\text{VII-6}}$,
- e) $(\text{C}=\text{O})\text{R}^{\text{VII-6}}$,
- f) $(\text{C}=\text{O})\text{OR}^{\text{VII-9}}$,
- g) cyano,
- h) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- i) $\text{OR}^{\text{VII-10}}$,
- j) Ohet^{VII} ,
- k) $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$,
- l) $\text{SR}^{\text{VII-10}}$,
- m) Shet^{VII} ,
- n) $\text{NHCOR}^{\text{VII-12}}$,
- o) $\text{NHSO}_2\text{R}^{\text{VII-12}}$, or

- p) C₁₋₇alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{VII-11}, OR^{VII-13}, SR^{VII-10}, SR^{VII-13}, NR^{VII-7}R^{VII-8}, halo, (C=O)C₁₋₇alkyl, or SO_m^{VII}R^{VII-9}, or
- q) R^{VII-1} together with R^{VII-2} form a carbocyclic or het^{VII} which may be optionally substituted by NR^{VII-7}R^{VII-8}, or C₁₋₇alkyl which may be optionally substituted by OR^{VII-14};

R^{VII-6} is

- a) C₁₋₇alkyl,
- b) NR^{VII-7}R^{VII-8}
- c) aryl^{VII}, or
- d) het^{VII}, wherein said het^{VII} is bound via a carbon atom;

R^{VII-7} and R^{VII-8} are independently

- a) H,
- b) aryl^{VII},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VII-10}R^{VII-10}, R^{VII-11}, SO_mR^{VII-9}, CONR^{VII-10}R^{VII-10}, or halo, or,
- d) R^{VII-7} and R^{VII-8} together with the nitrogen to which they are attached form a het^{VII};

R^{VII-9} is

- a) aryl^{VII},
- b) het^{VII},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VII-10}R^{VII-10}, R^{VII-11}, SH, CONR^{VII-10}R^{VII-10}, or halo;

R^{VII-10} is

- a) H,

- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VII-11} is

- a) OR^{VII-10},
- b) Ohet^{VII},
- c) Oaryl^{VII},
- d) CO₂R^{VII-10},
- e) het^{VII},
- f) aryl^{VII},
- g) CN, or
- h) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VII-11}, NR^{VII-7}R^{VII-8}, SO_m^{VII}R^{VII-9}, or C₁₋₇alkyl optionally substituted by R^{VII-11}, NR^{VII-7}R^{VII-8}, or SO_mR^{VII-9};

R^{VII-12} is

- a) H,
- b) het^{VII},
- c) aryl^{VII},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VII-7}R^{VII-8} or R^{VII-11};

R^{VII-13} is

- a) (P=O)(OR^{VII-14})₂,
- b) CO(CH₂)_nCON(CH₃)-(CH₂)_nSO₃⁻M⁺,
- c) an amino acid,
- d) C(=O)aryl^{VII}, or
- e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VII-7}R^{VII-8}, aryl^{VII}, het^{VII}, CO₂H, or O(CH₂)_n^{VII}CO₂R^{VII-14};

R^{VII-14} is

- a) H, or
- b) C₁₋₇alkyl;

each n^{VII} is independently 1, 2, 3, 4 or 5;

each m^{VII} is independently 0, 1, or 2;

M^{VII} is sodium, potassium, or lithium;


aryl^{VII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, $\text{CO}_2\text{R}^{\text{VII-14}}$, CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VII-14}}$, $\text{NR}^{\text{VII-14}}\text{R}^{\text{VII-14}}$, $\text{OR}^{\text{VII-14}}$, or $\text{CO}_2\text{R}^{\text{VII-14}}$ groups;

het^{VII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VII-14}}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three $\text{SR}^{\text{VII-14}}$, $\text{NR}^{\text{VII-14}}\text{R}^{\text{VII-14}}$, $\text{OR}^{\text{VII-14}}$, or $\text{CO}_2\text{R}^{\text{VII-14}}$ groups;

wherein Formula VIII is



and pharmaceutically acceptable salts thereof,

wherein

A^{VIII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VIII-1} is

- a) R^{VIII-5},
- b) NR^{VIII-7}R^{VIII-8}, or
- c) SO₂R^{VIII-9};

R^{VIII-2} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) SO_m^{VIII}R^{VIII-6},
- d) OC₂₋₇ alkyl substituted by OH,
- e) SC₂₋₇ alkyl substituted by OH, or
- f) C₂₋₈ alkyl which is partially unsaturated and is optionally substituted by one or more substituents selected from R^{VIII-11}, OR^{VIII-13}, SR^{VIII-13}, NR^{VIII-7}R^{VIII-8}, halo, (C=O)C₁₋₇ alkyl or SO_m^{VIII}R^{VIII-9};

with the proviso that when R^{VIII-1} = R^{VIII-5} =

(CH₂CH₂O)_i^{VIII}R^{VIII-10}, then R^{VIII-2} may additionally represent

- a) H,
- b) halo,
- c) (C=O)R^{VIII-6},
- d) (C=O)OR^{VIII-9},
- e) cyano,
- f) OR^{VIII-10},
- g) het^{VIII},
- h) NR^{VIII-7}R^{VIII-8},
- i) SR^{VIII-10},
- j) het^{VIII},
- k) NHCOR^{VIII-12},
- l) NHSO₂R^{VIII-12}, or

- m) R^{VIII-2} together with R^{VIII-3} or R^{VIII-4} form a carbocyclic or het^{VIII} which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-3} and R^{VIII-4} are independently:

- a) H,
- b) halo,
- c) aryl^{VIII},
- d) $S(O)_m^{VIII}R^{VIII-6}$,
- e) $(C=O)R^{VIII-6}$,
- f) $(C=O)OR^{VIII-9}$,
- g) cyano,
- h) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom,
- i) $OR^{VIII-10}$,
- j) Ohet^{VIII},
- k) $NR^{VIII-7}R^{VIII-8}$,
- l) $SR^{VIII-10}$,
- m) Shet^{VIII},
- n) $NHCOR^{VIII-12}$,
- o) $NHSO_2R^{VIII-12}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $R^{VIII-11}$, $OR^{VIII-13}$, $SR^{VIII-10}$, $SR^{VIII-13}$, $NR^{VIII-7}R^{VIII-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{VIII}R^{VIII-9}$, or
- q) R^{VIII-4} together with R^{VIII-3} form a carbocyclic or het which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-5} is

- a) $(CH_2CH_2O)_i^{VIII}R^{VIII-10}$,
- b) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom,
- c) aryl^{VIII},

- d) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VIII-7}R^{VIII-8}, R^{VIII-11}, SO_mR^{VIII-9}, or OC₂₋₄alkyl which may be further substituted by het^{VIII}, OR^{VIII-10}, or NR^{VIII-7}R^{VIII-8}, or
- e) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from R^{VIII-11}, NR^{VIII-7}R^{VIII-8}, SO_mR^{VIII-9}, or C₁₋₇alkyl optionally substituted by R^{VIII-11}, NR^{VIII-7}R^{VIII-8}, or SO_mR^{VIII-9};

R^{VIII-6} is

- a) C₁₋₇alkyl,
 b) NR^{VIII-7}R^{VIII-8},
 c) aryl^{VIII}, or
 d) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom;

R^{VIII-7} and R^{VIII-8} are independently

- a) H,
 b) aryl^{VIII},
 c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VIII-10}R^{VIII-10}, R^{VIII-11}, SO_mR^{VIII-9}, CONR^{VIII-10}R^{VIII-10}, or halo, or,
 d) R^{VIII-7} and R^{VIII-8} together with the nitrogen to which they are attached form a het^{VIII};

R^{VIII-9} is

- a) aryl^{VIII},
 b) het^{VIII},
 c) C₃₋₈cycloalkyl,
 d) methyl, or
 e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more

substituents selected from $\text{NR}^{\text{VIII-10}}\text{R}^{\text{VIII-10}}$,
 $\text{R}^{\text{VIII-11}}$, SH, $\text{CONR}^{\text{VIII-10}}\text{R}^{\text{VIII-10}}$, or halo;

$\text{R}^{\text{VIII-10}}$ is

- a) H,
- b) methyl, or
- c) C_{2-7} alkyl optionally substituted by OH;

$\text{R}^{\text{VIII-11}}$ is

- a) $\text{OR}^{\text{VIII-10}}$,
- b) $\text{Ohet}^{\text{VIII}}$,
- c) $\text{Oaryl}^{\text{VIII}}$,
- d) $\text{CO}_2\text{R}^{\text{VIII-10}}$,
- e) het^{VIII} ,
- f) $\text{aryl}^{\text{VIII}}$, or
- g) CN;

$\text{R}^{\text{VIII-12}}$ is

- a) H,
- b) het^{VIII} ,
- c) $\text{aryl}^{\text{VIII}}$,
- d) C_{3-8} cycloalkyl,
- e) methyl, or
- f) C_{2-7} alkyl optionally substituted by $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$
or $\text{R}^{\text{VIII-11}}$;

$\text{R}^{\text{VIII-13}}$ is

- a) $(\text{P}=\text{O})(\text{OR}^{14})_2$,
- b) $\text{CO}(\text{CH}_2)_n^{\text{VIII}}\text{CON}(\text{CH}_3) - (\text{CH}_2)_n^{\text{VIII}}\text{SO}_3^-\text{M}^+$,
- c) an amino acid,
- d) $\text{C}(=\text{O})\text{aryl}^{\text{VIII}}$, or
- e) $\text{C}(=\text{O})\text{C}_{1-7}$ alkyl optionally substituted by
 $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, $\text{aryl}^{\text{VIII}}$, het^{VIII} , CO_2H , or
 $\text{O}(\text{CH}_2)_n^{\text{VIII}}\text{CO}_2\text{R}^{\text{VIII-14}}$;

$\text{R}^{\text{VIII-14}}$ is


- a) H, or
- b) C_{1-7} alkyl;

each i^{VIII} is independently 2, 3, or 4;

each n^{VIII} is independently 1, 2, 3, 4 or 5;

each m^{VIII} is independently 0, 1, or 2;
 M^{VIII} is sodium, potassium, or lithium;
 $aryl^{VIII}$ is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;
 wherein any $aryl^{VIII}$ is optionally substituted with one or more substituents selected from halo, OH, cyano, $CO_2R^{VIII-14}$, CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three $SR^{VIII-14}$, $NR^{VIII-14}R^{VIII-14}$, $OR^{VIII-14}$, or $CO_2R^{VIII-14}$ groups;
 het^{VIII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;
 wherein any het^{VIII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $CO_2R^{VIII-14}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three $SR^{VIII-14}$, $NR^{VIII-14}R^{VIII-14}$, $OR^{VIII-14}$, or $CO_2R^{VIII-14}$ groups;

wherein Formula IX is



IX

and pharmaceutically acceptable salts thereof,
 wherein,
 R^{IX-1} is

- a) Cl,
- b) Br,
- c) CN,

d) NO_2 , or

e) F;

$\text{R}^{\text{IX}-2}$, $\text{R}^{\text{IX}-3}$ and $\text{R}^{\text{IX}-4}$ are independently selected from:

a) H,

b) halo,

c) aryl^{IX},

d) $\text{S}(\text{O})_m^{\text{IX}}\text{R}^{\text{IX}-6}$,

e) $(\text{C}=\text{O})\text{R}^{\text{IX}-6}$,

f) $(\text{C}=\text{O})\text{OR}^{\text{IX}-9}$,

g) cyano,

h) het^{IX}, wherein said ^{IX}het is bound via a carbon atom,

i) $\text{OR}^{\text{IX}-10}$,

j) Ohet^{IX},

k) $\text{NR}^{\text{IX}-7}\text{R}^{\text{IX}-8}$

l) $\text{SR}^{\text{IX}-10}$,

m) Shet^{IX},

n) $\text{NHCOR}^{\text{IX}-12}$,

o) $\text{NHSO}_2\text{R}^{\text{IX}-12}$, or

p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{IX}-11}$, $\text{OR}^{\text{IX}-13}$, $\text{SR}^{\text{IX}-10}$, $\text{SR}^{\text{IX}-13}$, $\text{NR}^{\text{IX}-7}\text{R}^{\text{IX}-8}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m^{\text{IX}}\text{R}^{\text{IX}-9}$;

$\text{R}^{\text{IX}-6}$ is

a) C_{1-7} alkyl,

b) $\text{NR}^{\text{IX}-7}\text{R}^{\text{IX}-8}$,

c) aryl^{IX}, or

d) het^{IX}, wherein said het^{IX} is bound via a carbon atom;

$\text{R}^{\text{IX}-7}$ and $\text{R}^{\text{IX}-8}$ are independently

a) H,

b) aryl^{IX},

c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more

substituents selected from $\text{NR}^{\text{IX-10}}\text{R}^{\text{IX-10}}$, $\text{R}^{\text{IX-11}}$, $\text{SO}_m\text{R}^{\text{IX-9}}$, $\text{CONR}^{\text{IX-10}}\text{R}^{\text{IX-10}}$, or halo, or,

- d) $\text{R}^{\text{IX-7}}$ and $\text{R}^{\text{IX-8}}$ together with the nitrogen to which they are attached form a IX-het ;

$\text{R}^{\text{IX-9}}$ is

- a) aryl^{IX},
 b) het^{IX},
 c) C_{3-8} cycloalkyl,
 d) methyl, or
 e) C_{2-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $\text{NR}^{\text{IX-10}}\text{R}^{\text{IX-10}}$, $\text{R}^{\text{IX-11}}$, SH, $\text{CONR}^{\text{IX-10}}\text{R}^{\text{IX-10}}$, or halo;

$\text{R}^{\text{IX-10}}$ is

- a) H,
 b) methyl, or
 c) C_{2-7} alkyl optionally substituted by OH;

$\text{R}^{\text{IX-11}}$ is

- a) $\text{OR}^{\text{IX-10}}$,
 b) Ohet^{IX} ,
 c) Oaryl^{IX} ,
 d) $\text{CO}_2\text{R}^{\text{IX-10}}$,
 e) het^{IX},
 f) aryl^{IX}, or
 g) CN;

$\text{R}^{\text{IX-12}}$ is

- a) H,
 b) het^{IX},
 c) aryl^{IX},
 d) C_{3-8} cycloalkyl,
 e) methyl, or
 f) C_{2-7} alkyl optionally substituted by $\text{NR}^{\text{IX-7}}\text{R}^{\text{IX-8}}$ or $\text{R}^{\text{IX-11}}$;

$\text{R}^{\text{IX-13}}$ is

- a) $(\text{P}=\text{O})(\text{OR}^{\text{IX-14}})_2$,

- b) $\text{CO}(\text{CH}_2)_n^{\text{IX}}\text{CON}(\text{CH}_3) - (\text{CH}_2)_n^{\text{IX}}\text{SO}_3^- \text{M}^{\text{IX}+}$,
- c) an amino acid,
- d) $\text{C}(=\text{O})\text{aryl}^{\text{IX}}$, or
- e) $\text{C}(=\text{O})\text{C}_{1-7}\text{alkyl}$ optionally substituted by $\text{NR}^{\text{IX}-7}\text{R}^{\text{IX}-8}$, aryl^{IX} , het^{IX} , CO_2H , or $\text{O}(\text{CH}_2)_n\text{CO}_2\text{R}^{\text{IX}-14}$;

$\text{R}^{\text{IX}-14}$ is

- a) H, or
- b) $\text{C}_{1-7}\text{alkyl}$;

each n^{IX} is independently 1, 2, 3, 4 or 5;

each m^{IX} is independently 0, 1, or 2;

M^{IX} is sodium, potassium, or lithium;

aryl^{IX} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, $\text{CO}_2\text{R}^{\text{IX}-14}$, CF_3 , $\text{C}_{1-6}\text{alkoxy}$, and $\text{C}_{1-6}\text{alkyl}$ which may be further substituted by one to three $\text{SR}^{\text{IX}-14}$, $\text{NR}^{\text{IX}-14}\text{R}^{\text{IX}-14}$, $\text{OR}^{\text{IX}-14}$, or $\text{CO}_2\text{R}^{\text{IX}-14}$ groups;

het^{IX} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{IX}-14}$, CF_3 , $\text{C}_{1-6}\text{alkoxy}$, oxo, oxime, and $\text{C}_{1-6}\text{alkyl}$ which may be further substituted by one to three $\text{SR}^{\text{IX}-14}$, $\text{NR}^{\text{IX}-14}\text{R}^{\text{IX}-14}$, $\text{OR}^{\text{IX}-14}$, or $\text{CO}_2\text{R}^{\text{IX}-14}$ groups.

Also provided is the use of compounds of Formulas VI, VII, VIII and IX to prepare medicaments for preventing or treating atherosclerosis or reestenosis in mammals.

The advantage of using compounds of Formulas VI, VII, VIII and IX in the method of our invention is their extensive activity against herpesviruses since atherosclerosis is related to the number of herpesvirus infections. Drugs containing compounds of Formulae VI-IX could prevent the inflammatory response resulting from reactivation of HCMV, EBV, HSV-1, HSV-2, HHV-8 and VZV.

DETAILED DESCRIPTION OF THE INVENTION

The compounds of Formula VI, their method of preparation and formulation into pharmaceutical dosage forms are described in U. S. Patent Application Serial No. 09/808 836, filed March 15, 2001. The disclosure of U. S. Patent Application Serial No. 09/808 836 is herein incorporated in its entirety by reference.

The compounds of Formula VII, their method of preparation and formulation into pharmaceutical dosage forms are disclosed in U. S. Patent Application Serial No. 09/808 902, filed March 15, 2001. The disclosure of U.S. Patent Application Serial No. 09/808 902 is herein incorporated in its entirety by reference.

The compounds of Formula VIII, their method of preparation and formulation into pharmaceutical dosage forms are described in U.S. Patent Application Serial No. 09/808 757, filed March 15, 2001. The disclosure of U.S. Patent Application Serial No. 09/808 757 is herein incorporated in its entirety by reference.

The compounds of Formula IX, their method of preparation and formulation into pharmaceutical dosage forms are described in U.S. Patent No. 6 413 958. U.S. Patent No. 6 413 958 is herein incorporated in its entirety by reference.

The correspondence between the compounds utilized in the method of the invention and the compounds incorporated by reference is as follows:

Formula VI corresponds to Formula I of U. S. Patent Application Serial No. 09/808 836.

Formula VII corresponds to Formula I of U. S. Patent Application Serial No. 09/808 902.

Formula VIII corresponds to Formula I of U.S. Patent Application Serial No. 09/808 757.

Formula IX corresponds to Formula I of U.S. Patent No. 6 413 958.

The invention further provides:

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI and wherein A^{VI} is Cl.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI and wherein R^{VI-1} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, carboxymethyl, (C1-7 alkoxy)carbonylmethyl, 2-hydroxyethyl, 2-(2-methoxyethoxy)ethyl, 3-(2-tetrahydropyranyloxy)propyl, 2-morpholinoethyl, 2-(diethylamino)ethyl, 2-(dimethylamino)ethyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(1-methylpyrrolidin-2-yl)ethyl, 2-(diisopropylamino)ethyl, 2-pyrrolidin-1-ylethyl, 3-(dimethylamino)propyl, and vinyl.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI and wherein R^{VI-2} is selected from the group consisting of CH_2 -morpholine, alkylnl- CH_2OH , CH_2 -(tetrahydro-2H-pyran-4-yl), and $(CH_2)_3OH$.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI and wherein the compound administered is selected from the group consisting of

N-(4-chlorobenzyl)-6-iodo-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(hydroxymethyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-{[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethynyl}-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(1-hydroxycyclohexyl)ethynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3S)-3-hydroxy-1-butynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butynyl}-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butynyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxy-1-butynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(5-hydroxy-1-pentynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[1R,2S]-2-hydroxycyclopentyl]ethynyl]-1-methyl-6-(4-

morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-methyl-1-butynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-(phenylethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-phenyl-1-propynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[3-(methylsulfonyl)propyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[3-(methylsulfonyl)propyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[(2-hydroxyethoxy)methyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-3-furanyl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-(1,2-diethyl-4-pyrazolidinyl)-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-1-(3-oxetanyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-{3-[3-(3-hydroxypropyl)sulfonyl]propyl}-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[2-(2-ethoxyethoxy)ethyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-[(phenylsulfinyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-[(phenylsulfonyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-[(phenylsulfanyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-2H-pyran-3-yl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[(methylsulfanyl)methyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[[(4-chlorophenyl) sulfinyl]methyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-2H-pyran-4-yl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-(4-thiomorpholinylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(4-hydroxy-1-piperidinyl)methyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxypyrrolidinylmethyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3-hydroxy-1-piperidinyl)methyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

[3-[(4-chlorobenzyl)amino]carbonyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-8-cinnolinylmethyl 4-morpholinecarboxylate

N-(4-chlorobenzyl)-8-(hydroxymethyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3-cyanobenzyl)amino]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6,8-bis(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-[(1-acetyl-4-piperidinyl)amino]-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-([1-methyl-2-(phenylsulfonyl)ethyl]amino)-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-{[3-(4-methoxyphenyl)-1-methylpropyl]amino}-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-amino-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-8-[(3-nitrobenzyl)amino]-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-(tetrahydro-2H-pyran-4-ylamino)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-hydroxy-1-butyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-([(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(1-hydroxycyclohexyl)ethyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3S)-3-hydroxy-1-butyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butyl}-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxy-1-butyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(5-hydroxy-1-pentyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-{[(1R,2S)-2-hydroxycyclopentyl]ethyl}-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-methyl-1-butyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1H-imidazol-1-yl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1H-imidazol-1-yl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-(phenylethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-phenyl-1-propyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-{[methyl(tetrahydro-2-furanylmethyl)amino]methyl}-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

and pharmaceutically acceptable salts thereof.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VII and wherein A^{VII} is Cl.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VII and wherein R^{VII-1} is selected from the group consisting of CH₂-morpholine, alkynyl-CH₂OH, CH₂-(tetrahydro-2H-pyran-4-yl) and (CH₂)₃OH.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VII and wherein the compound administered is selected from the group consisting of

N-(4-chlorobenzyl)-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

Methyl 3-[(4-chlorobenzyl)amino]carbonyl-4-hydroxy-6-cinnolinecarboxylate;

N-(4-chlorobenzyl)-4-hydroxy-6-(hydroxymethyl)-3-cinnolinecarboxamide

N-(4-chlorobenzyl)-8-(cyclopropylethynyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1-hydroxycyclohexyl)ethynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propynyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3S)-3-hydroxy-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butynyl}-N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3R)-3-hydroxy-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(5-hydroxy-1-pentynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[[(1R,2S)-2-hydroxycyclopentyl]ethynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-methyl-1-butynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1-hydroxycyclohexyl)ethyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3S)-3-hydroxy-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butyl}-N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3R)-3-hydroxy-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(5-hydroxy-1-pentyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[[(1R,2S)-2-hydroxycyclopentyl]ethyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-methyl-1-butyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(phenylethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-phenyl-1-propynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-([(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(phenylethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-phenyl-1-propyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-([(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

and pharmaceutically acceptable salts thereof.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VIII and wherein A^{VIII} is Cl.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VIII and wherein R^{VIII-1} is selected from the group consisting of methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, carboxymethyl, (C1-7 alkoxy)carbonylmethyl, 2-hydroxyethyl, 2-(2-methoxyethoxy)ethyl, 3-(2-tetrahydropyranyloxy)propyl, 2-morpholinoethyl, 2-(diethylamino)ethyl, 2-(dimethyl-

amino)ethyl, 2-piperidinoethyl, 3-piperidinopropyl, 2-(1-methylpyrrolidin-2-yl)ethyl, 2-(diisopropylamino)ethyl, 2-pyrrolidin-1-ylethyl, 3-(dimethylamino)propyl, and vinyl.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VIII and wherein R^{VIII-2} is alkynyl-CH₂OH.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VIII and wherein the compound administered is N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide, or N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VIII and wherein the compound administered is

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-Chlorobenzyl)-6-iodo-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-1,7-dimethyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-1-methyl-4,7-dioxo-1,4,7,8-tetrahydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

ethyl 6-([(4-chlorobenzyl)amino]carbonyl)-2-methoxy-8-methyl-5-oxo-5,8-dihydro[1,8]naphthyridine-3-carboxylate;

and pharmaceutically acceptable salts thereof.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula IX and wherein R^{IX-1} is Cl.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula IX and wherein R^{IX-3} is selected from the group consisting of CH₂-morpholine, alkynyl-CH₂OH, CH₂-(tetrahydro-2H-pyran-4-yl) and (CH₂)₃OH.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula IX and is selected from a group consisting of

N-(4-chlorobenzyl)-4-hydroxy-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-7-methyl-6-(tetrahydro-2H-pyran-4-ylmethyl)[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-7-methyl-6-(4-morpholinylmethyl)[1,8]naphthyridine-3-carboxamide;

6-bromo-N-(4-chlorobenzyl)-4-hydroxy-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(3-hydroxy-1-propynyl)-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-iodo-7-methyl[1,8]naphthyridine-3-carboxamide; and
Methyl 6-{{(4-chlorobenzyl) amino}carbonyl}-5-hydroxy-2-methyl[1,8]naphthyridine-3-carboxylate.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formulae VI, VII, VIII or IX and wherein said mammal is a human.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI, VII, VIII or IX and wherein said mammal is a livestock or companion animal.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI, VII, VIII or IX and wherein the amount administered is from about 0.1 to about 300 mg/kg of body weight.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI, VII, VIII or IX and wherein the amount administered is from about 1 to about 30 mg/kg of body weight.

A method for preventing or treating atherosclerosis or restenosis in mammals, wherein the compound administered has the Formula VI, VII, VIII or IX and wherein the compound is administered parenterally, intravaginally, intranasally, topically, orally, or rectally.

Use of a compound of Formula VI, VII, VIII or IX for the manufacture of a medicament useful for preventing or treating atherosclerosis or restenosis in a mammal.

Dosages and Dosage Forms

By the term "effective amount" of a compound as provided herein is meant a nontoxic but sufficient amount of one or more anti-atherosclerosis or anti-restenosis agents to provide the desired effect. The desired effect may be to prevent, give relief from, or ameliorate atherosclerosis or restenosis.

As pointed out below, the exact amount of the anti-atherosclerosis or anti-restenosis agent required to treat atherosclerosis or restenosis will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the disease that is being treated, the particular compound(s) used, the mode of administration, such as the route and frequency of administration, and the particular compound(s) employed, and the like. Thus, it is not possible to specify an exact "effective amount." However, an appropriate effective amount may be determined by one of ordinary skill in the art using only routine experimentation.

Pharmaceutical compositions including one or more anti-atherosclerosis or anti-restenosis agents can be administered orally or parenterally at dose levels, calculated as the free base, of each of the anti-atherosclerosis or anti-restenosis agent at 0.1 to 300 mg/kg, preferably 1.0 to 30 mg/kg or mammal body weight, and can be used in a human in a unit dosage form, administered one to four times daily in the amount of 1 to 1000 mg per unit dose.

Initial treatment of a patient suffering from atherosclerosis or restenosis can begin with a dosage

regimen as indicated above. Treatment is generally continued as necessary over a period of several weeks to several months or years until the condition or disorder has been controlled or eliminated. Patients undergoing treatment with a composition of the invention can be routinely monitored by any of the methods well known in the art to determine the effectiveness of therapy. Continuous analysis of data from such monitoring permits modification of the treatment regimen during therapy so that optimally effective amounts of drug are administered at any point in time, and so that the duration of treatment can be determined. In this way, the treatment regimen and dosing schedule can be rationally modified over the course of therapy so that the lowest amount of the compounds of this invention exhibiting satisfactory effectiveness is administered, and so that administration is continued only for so long as is necessary to successfully treat the condition or disorder.

Also, it is to be understood that the initial dosage administered may be increased beyond the above upper level in order to rapidly achieve the desired plasma concentration. On the other hand, the initial dosage may be smaller than the optimum and the daily dosage may be progressively increased during the course of treatment depending on the particular situation.

In a combination therapy, the anti-atherosclerosis or anti-restenosis agent compound(s) and other inhibitor compound(s) can be administered simultaneously or at separate intervals. When administered simultaneously the anti-atherosclerosis or anti-restenosis agent compound(s) and the other inhibitor compound(s) can be incorporated into a single pharmaceutical composition or into separate compositions, e.g., anti-atherosclerosis or anti-restenosis agent compound(s) in one composition and the other inhibitor compound(s) in another composition. For

instance the combination therapy, the anti-atherosclerosis or anti-restenosis agent compound(s) may be administered concurrently or concomitantly with the other inhibitor compound(s). The term "concurrently" means the subject being treated takes one drug within about 5 minutes of taking the other drug. The term "concomitantly" means the subject being treated takes one drug within the same treatment period of taking the other drug. The same treatment period is preferably within twelve hours and up to forty-eight hours.

When separately administered, therapeutically effective amounts of anti-atherosclerosis or anti-restenosis agent compound(s) and the other inhibitor compound(s) are administered on a different schedule. One may be administered before the other as long as the time between the two administrations falls within a therapeutically effective interval. A therapeutically effective interval is a period of time beginning when one of either (a) the anti-atherosclerosis or anti-restenosis agent compound(s), or (b) the other inhibitor compound(s) is administered to a mammal and ending at the limit of the beneficial effect in the treatment of atherosclerosis or restenosis of the combination of (a) and (b). The methods of administration of the anti-atherosclerosis or anti-restenosis agent compound(s) and the other inhibitor compound(s) may vary. Thus, one agent may be administered orally, while the other is administered by injection.

A specific active agent may have more than one recommended dosage range, particularly for different routes of administration. Generally, an effective amount of dosage of anti-atherosclerosis or anti-restenosis agent compound(s), either administered individually or in combination with other inhibitor compound(s), will be in the range of about 0.1 to about 300 mg/kg of body

weight/day, preferably about 1 to about 30 mg/kg of body weight/day. It is to be understood that the dosages of active component(s) may vary depending upon the requirements of each subject being treated and the severity of the atherosclerosis or restenosis.

In addition to the anti-atherosclerosis or anti-restenosis agents, the composition for therapeutic use may also comprise one or more non-toxic, pharmaceutically acceptable carrier materials or excipients. The term "carrier" material or "excipient" herein means any substance, not itself a therapeutic agent, used as a carrier and/or diluent and/or adjuvant, or vehicle for delivery of a therapeutic agent to a subject or added to a pharmaceutical composition to improve its handling or storage properties or to permit or facilitate formation of a dose unit of the composition into a discrete article such as a capsule or tablet suitable for oral administration. Excipients can include, by way of illustration and not limitation, diluents, disintegrants, binding agents, adhesives, wetting agents, polymers, lubricants, glidants, substances added to mask or counteract a disagreeable taste or odor, flavors, dyes, fragrances, and substances added to improve appearance of the composition. Acceptable excipients include lactose, sucrose, starch powder, cellulose esters of alkanolic acids, cellulose alkyl esters, talc, stearic acid, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulfuric acids, gelatin, acacia gum, sodium alginate, polyvinyl-pyrrolidone, and/or polyvinyl alcohol, and then tableted or encapsulated for convenient administration. Such capsules or tablets may contain a controlled-release formulation as may be provided in a dispersion or active compound in hydroxypropyl-methyl cellulose, or other methods known to those skilled in the art. For oral administration, the

pharmaceutical composition may be in the form of, for example, a tablet, capsule, suspension or liquid. If desired, other active ingredients may be included in the composition.

In addition to the oral dosing, noted above, the compositions of the present invention may be administered by any suitable route, in the form of a pharmaceutical composition adapted to such a route, and in a dose effective for the treatment intended. The compositions may, for example, be administered parenterally, e.g., intravascularly, intraperitoneally, subcutaneously, or intramuscularly. For parenteral administration, saline solution, dextrose solution, or water may be used as a suitable carrier. Formulations for parenteral administration may be in the form of aqueous or non-aqueous isotonic sterile injection solutions or suspensions. These solutions and suspensions may be prepared from sterile powders or granules having one or more of the carriers or diluents mentioned for use in the formulations for oral administration. The compounds may be dissolved in water, polyethylene glycol, propylene glycol, ethanol, corn oil, cottonseed oil, peanut oil, sesame oil, benzyl alcohol, sodium chloride, and/or various buffers. Other adjuvants and modes of administration are well and widely known in the pharmaceutical art.

Generally, the concentration of each of the anti-atherosclerosis or anti-restenosis agents in a liquid composition, such as a lotion, will be from about 0.1 wt.% to about 20 wt.%, preferably from about 0.5 wt.% to about 10 wt.%. The solution may contain other ingredients, such as emulsifiers, antioxidants or buffers. The concentration in a semi-solid or solid composition, such as a gel or a powder, will be about 0.1 wt.% to about 5 wt.%, preferably about 0.5 wt.% to about

2.5 wt.%. When the topically deliverable, pharmaceutical composition of the present invention is utilized to effect targeted treatment of a specific internal site, each of the anti-atherosclerosis or anti-restenosis agent is preferably contained in the composition in an amount of from 0.05-10 wt.%, more preferably 0.5-5 wt.%.

Routes of Administration

In therapeutic use for treating, or preventing atherosclerosis or restenosis in a mammal (i.e., human and animals) the pharmaceutical composition including the anti-atherosclerosis or anti-restenosis agent(s) of Formulae VI, VII, VIII and IX can be administered orally, parenterally, topically, rectally, or intranasally.

Parenteral administrations include injections to generate a systemic effect or injections directly to the afflicted area. Examples to parenteral administrations are subcutaneous, intravenous, intramuscular, intradermal, intrathecal, intraocular, intraventricular, and general infusion techniques.

Topical administrations include the treatment of infectious areas or organs readily accessibly by local application, such as, for example, eyes, ears including external and middle ear infections, vaginal, open and sutured or closed wounds and skin. It also includes transdermal delivery to generate a system effect.

The rectal administration includes the form of suppositories.

The intranasally administration includes nasal aerosol or inhalation applications.

Pharmaceutical compositions including the anti-atherosclerosis or anti-restenosis agent(s) may be prepared by methods well known in the art, e.g., by means of conventional mixing, dissolving, granulation, dragee-

making, levigating, emulsifying, encapsulating, entrapping, lyophilizing processes or spray drying.

Pharmaceutical compositions for use in accordance with the present invention may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For oral administration, the anti-atherosclerosis or anti-restenosis agent(s) can be formulated by combining the active compounds with pharmaceutically acceptable carriers well known in the art. Such carriers enable the compounds of the invention to be formulated as tablets, pills, lozenges, dragees, capsules, liquids, solutions, emulsions, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient. A carrier can be at least one substance which may also function as a diluent, flavoring agent, solubilizer, lubricant, suspending agent, binder, tablet disintegrating agent, and encapsulating agent. Examples of such carriers or excipients include, but are not limited to, magnesium carbonate, magnesium stearate, talc, sugar, lactose, sucrose, pectin, dextrin, mannitol, sorbitol, starches, gelatin, cellulosic materials, low melting wax, cocoa butter or powder, polymers such as polyethylene glycols and other pharmaceutical acceptable materials.

Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings

for identification or to characterize different combinations of active compound doses.

Pharmaceutical compositions which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with a filler such as lactose, a binder such as starch, and/or a lubricant such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, liquid polyethylene glycols, cremophor, capmul, medium or long chain mono-, di- or triglycerides. Stabilizers may be added in these formulations, also.

Liquid form compositions include solutions, suspensions and emulsions. For example, there may be provided solutions of pharmaceutical compositions with the anti-atherosclerosis or anti-restenosis agent(s) dissolved in water and water-propylene glycol and water-polyethylene glycol systems, optionally containing suitable conventional coloring agents, flavoring agents, stabilizers and thickening agents.

The anti-atherosclerosis or anti-restenosis agent(s) may also be formulated for parenteral administration, e.g., by injections, bolus injection or continuous infusion. Formulations for parenteral administration may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oil or aqueous vehicles, and may contain formulating materials such as suspending, stabilizing and/or dispersing agents.

For injection, the anti-atherosclerosis or anti-restenosis agent(s) may be formulated in aqueous

solution, preferably in physiologically compatible buffers or physiological saline buffer. Suitable buffering agents include tri-sodium orthophosphate, sodium bicarbonate, sodium citrate, N-methyl-glucamine, L(+)-lysine and L(+)-arginine.

The compositions can also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the active compound or its salts can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

Pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the active ingredient which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and anti-fungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal; and the like. In many cases, it will

be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

Other parenteral administrations also include aqueous solutions of a water soluble form, such as, without limitation, a salt, of the anti-atherosclerosis or anti-restenosis agent(s). Additionally, suspensions of the active compounds may be prepared in a lipophilic vehicle. Suitable lipophilic vehicles include fatty oils such as sesame oil, synthetic fatty acid esters such as ethyl oleate and triglycerides, or materials such as liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers and/or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Alternatively, the anti-atherosclerosis or anti-restenosis agent(s) may be in a powder form for constitution with a suitable vehicle, e.g., sterile, pyrogen-free water, before use.

For suppository administration, the pharmaceutical compositions may also be formulated by mixing the anti-atherosclerosis or anti-restenosis agent(s) with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and other glycerides.

For administration by inhalation, the anti-atherosclerosis or anti-restenosis agent(s) can be conveniently delivered through an aerosol spray in the form of solution, dry powder, or cream. The aerosol may use a pressurized pack or a nebulizer and a suitable propellant. In the case of a pressurized aerosol, the dosage unit may be controlled by providing a valve to deliver a metered amount. Capsules and cartridges of, for example, gelatin for use in an inhaler may be formulated containing a powder base such as lactose or starch.

For ophthalmic and otitis uses, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative, such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment, such as petrolatum.

In addition to the formulations described previously, the anti-atherosclerosis or anti-restenosis agent(s) may also be formulated as depot preparations. Such long acting formulations may be in the form of implants. The anti-atherosclerosis or anti-restenosis agent(s) may be formulated for this route of administration with suitable polymers, hydrophobic

materials, or as a sparingly soluble derivative such as, without limitation, a sparingly soluble salt.

Additionally, the anti-atherosclerosis or anti-restenosis agent(s) may be delivered using a sustained-release system. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the compounds for 24 hours up to several days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

In certain embodiments, the anti-atherosclerosis or anti-restenosis agent(s) are applied topically. For topical applications, the pharmaceutical composition may be formulated in a suitable ointment containing the anti-atherosclerosis or anti-restenosis agent(s) suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion such as suspensions, emulsion, or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

Several different animal models are available to evaluate reduction of atherosclerosis or restenosis by antiviral drug treatment. In these models histological changes in the atherosclerotic lesions of aortic arteries

are measured in animals infected with a herpesvirus and treated or untreated with an antiviral drug capable of inhibiting replication of the herpesvirus. The models include murine CMV infection of apoE deficient mice and rat CMV infection of rats. These models would mimic the effects of human CMV infection. MHV-68 is a murine gammaherpesvirus related to EBV. Antiviral treatment has been shown to reduce atherosclerosis caused by HMV-68 infection in apoE deficient mice. Drugs containing compounds of Formula I and II inhibit replication of these animal viruses so the models could be used to show an effect of drugs containing compounds of Formula I and II on development of atherosclerosis. Lemstrom, et al, "Cytomegalovirus Infection-Enhanced Allograft Atherosclerosis is prevented by DHPG Prophylaxis in the Rat", Circulation Vol. 90, No. 4, October 1994, pp 1969-1978; Burnell et al, "Atherosclerosis in a poE Knockout Mice Infected with Multiple Pathogens". Both of these references are herein incorporated by reference.


The terms and expressions which have been employed in the foregoing specification are used therein as terms of description and not of limitation, and there is no intention, in the use of such terms and expressions, of excluding equivalents of the features shown and described or portions thereof, it being recognized that the scope of the invention is defined and limited by the claims which follow.

All published documents are incorporated by reference herein.

What is claimed is:

1. Use of a compound selected from the group consisting of structures of Formula VI, VII, VIII or IX to manufacture a medicament useful for preventing or treating atherosclerosis or restenosis in a mammal,

wherein Formula VI is:



or a pharmaceutically acceptable salt thereof wherein, A^{VI} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VI-1} is

- a) R^{VI-5}, or
- b) SO₂R^{VI-9}

R^{VI-2}, R^{VI-3} and R^{VI-4} may be the same or different and are selected from the group consisting of:

- a) H,
- b) halo^{VI},
- c) aryl^{VI},
- d) S(O)_mR^{VI-6},
- e) (C=O)R^{VI-6},
- f) (C=O)OR^{VI-9},
- g) cyano,
- h) het^{VI}, wherein said het^{VI} is bound via a carbon atom,

- i) $\text{OR}^{\text{VI}-10}$,
- j) Ohet^{VI} ,
- k) $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$
- l) $\text{SR}^{\text{VI}-10}$,
- m) Shet^{VI} ,
- n) $\text{NHCOR}^{\text{VI}-12}$,
- o) $\text{NHSO}_2\text{R}^{\text{VI}-12}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VI}-11}$, $\text{OR}^{\text{VI}-13}$, $\text{SR}^{\text{VI}-10}$, $\text{SR}^{\text{VI}-13}$, $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m\text{R}^{\text{VI}-9}$, and
- q) $\text{R}^{\text{VI}-3}$ together with $\text{R}^{\text{VI}-2}$ or $\text{R}^{\text{VI}-4}$ form a carbocyclic or $^{\text{VI}}$ -het which may be optionally substituted by $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, or C_{1-7} alkyl which may be optionally substituted by $\text{OR}^{\text{VI}-14}$;

$\text{R}^{\text{VI}-5}$ is

- a) $(\text{CH}_2\text{CH}_2\text{O})_i\text{R}^{\text{VI}-10}$,
- b) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, $\text{R}^{\text{VI}-11}$, $\text{SO}_m\text{R}^{\text{VI}-9}$, or OC_{2-4} alkyl which may be further substituted by het^{VI} , $\text{OR}^{\text{VI}-10}$, or $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, or
- c) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of $\text{R}^{\text{VI}-11}$, $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, $\text{SO}_m\text{R}^{\text{VI}-9}$, or C_{1-7} alkyl optionally substituted by $\text{R}^{\text{VI}-11}$, $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, or $\text{SO}_m\text{R}^{\text{VI}-9}$;

$\text{R}^{\text{VI}-6}$ is

- a) C_{1-7} alkyl,
- b) $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$,
- c) aryl $^{\text{VI}}$, or

- d) het^{VI} , wherein said het^{VI} is bound via a carbon atom;

$\text{R}^{\text{VI-7}}$ and $\text{R}^{\text{VI-8}}$ are independently

- a) H,
 b) aryl^{VI} ,
 c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of aryl^{VI} , $\text{NR}^{\text{VI-10}}\text{R}^{\text{VI-10}}$, $\text{R}^{\text{VI-11}}$, $\text{SO}_m\text{R}^{\text{VI-9}}$, $\text{CONR}^{\text{VI-10}}\text{R}^{\text{VI-10}}$, or halo, or;
 d) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of $\text{R}^{\text{VI-11}}$, $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, $\text{SO}_m^{\text{VI}}\text{R}^{\text{VI-9}}$, or C_{1-7} alkyl optionally substituted by $\text{R}^{\text{VI-11}}$, $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, or $\text{SO}_m^{\text{VI}}\text{R}^{\text{VI-9}}$, or
 e) $\text{R}^{\text{VI-7}}$ and $\text{R}^{\text{VI-8}}$ together with the nitrogen to which they are attached form a het^{VI} ;

$\text{R}^{\text{VI-9}}$ is

- a) aryl^{VI} ,
 b) het^{VI} ,
 c) C_{3-8} cycloalkyl,
 d) methyl, or
 e) C_{2-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of $\text{NR}^{\text{VI-10}}\text{R}^{\text{VI-10}}$, $\text{R}^{\text{VI-11}}$, SH, $\text{CONR}^{\text{VI-10}}\text{R}^{\text{VI-10}}$, or halo;

$\text{R}^{\text{VI-10}}$ is

- a) H,
 b) methyl, or
 c) C_{2-7} alkyl optionally substituted by OH;

$\text{R}^{\text{VI-11}}$ is

- a) $\text{OR}^{\text{VI-10}}$,
 b) Ohet^{VI} ,
 c) Oaryl^{VI} ,

- d) $\text{CO}_2\text{R}^{\text{VI}-10}$,
- e) het^{VI} ,
- f) $^{\text{VI}}\text{-aryl}^{\text{VI}}$,
- g) CN, or
- h) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of $\text{R}^{\text{VI}-11}$, $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, $\text{SO}_m^{\text{IV}}\text{R}^{\text{VI}-9}$, or C_{1-7} alkyl optionally substituted by $\text{R}^{\text{VI}-11}$, $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, or $\text{SO}_m\text{R}^{\text{VI}-9}$;

$\text{R}^{\text{VI}-12}$ is

- a) H,
- b) het^{VI} ,
- c) aryl^{VI} ,
- d) C_{3-8} cycloalkyl,
- e) methyl, or
- f) C_{2-7} alkyl optionally substituted by $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$ or $\text{R}^{\text{VI}-11}$;

$\text{R}^{\text{VI}-13}$ is

- a) $(\text{P}=\text{O})(\text{OR}^{\text{VI}-14})_2$,
- b) $\text{CO}(\text{CH}_2)_n^{\text{IV}}\text{CON}(\text{CH}_3) - (\text{CH}_2)_n\text{SO}_3^-\text{M}^{\text{VI}+}$,
- c) an amino $^{\text{VI}}$ acid,
- d) $\text{C}(=\text{O})\text{aryl}^{\text{VI}}$,
- e) $\text{C}(=\text{O})\text{C}_{1-7}$ alkyl optionally substituted by $\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$, aryl^{VI} , het^{VI} , CO_2H , or $\text{O}(\text{CH}_2)_n\text{CO}_2\text{R}^{\text{VI}-14}$, or
- f) $\text{C}(=\text{O})\text{NR}^{\text{VI}-7}\text{R}^{\text{VI}-8}$

$\text{R}^{\text{VI}-14}$ is

- a) H, or
- b) C_{1-7} alkyl;

each i^{VI} is independently 2, 3, or 4;

each n^{VI} is independently 1, 2, 3, 4 or 5;

each m^{VI} is independently 0, 1, or 2;

M^{VI} is sodium, potassium, or lithium;

aryl^{VI} is a phenyl radical or an ortho-fused bicyclic


carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VI} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO₂R^{VI-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆ alkyl which maybe further substituted by one to three SR^{VI-14}, NR^{VI-14}R^{VI-14}, OR^{VI-14}, or CO₂R^{VI-14};

het^{VI} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VI} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO₂R^{VI-14}, CF₃, C₁₋₆alkoxy, oxo, oxime, and C₁₋₆ alkyl which maybe further substituted by one to three SR^{VI-14}, NR^{VI-14}R^{VI-14}, OR^{VI-14}, or CO₂R^{VI-14};

wherein Formula VII is



VII

or a pharmaceutically acceptable salt thereof,

wherein

A^{VII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VII-1} is

- a) aryl^{VII},
- b) S(O)_m^{VII}R^{VII-6},
- c) (C=O)R^{VII-6}, with the proviso that if R^{VII-6} is NR^{VII-7}R^{VII-8}, then R^{VII-7} and R^{VII-8} do not both equal H,
- d) (C=O)OR^{VII-9},
- e) cyano,
- f) het^{VII}, wherein said het^{VII} is bound via a carbon atom,
- g) Ohet^{VII},
- h) NR^{VII-7}R^{VII-8} with the proviso that R^{VII-7} and R^{VII-8} do not both equal H,
- i) SR^{VII-10},
- j) Shet^{VII},
- k) NHCOR^{VII-12},
- l) NHSO₂R^{VII-12},
- m) C₁₋₇alkyl which is partially unsaturated and optionally substituted by one or more substituents of the group R^{VII-11}, OR^{VII-13}, SR^{VII-10}, SR^{VII-13}, NR^{VII-7}R^{VII-8}, halo, (C=O)C₁₋₇alkyl, or SO_mR^{VII-9}, or
- n) C₁₋₇alkyl which is substituted by one or more substituents of the group R^{VII-11}, OR^{VII-13}, SR^{VII-10}, SR^{VII-13}, NR^{VII-7}R^{VII-8}, halo, (C=O)C₁₋₇alkyl, or SO_mR^{VII-9};

R^{VII-2} is

- a) H,
- b) halo,
- c) aryl^{VII},
- d) S(O)_m^{VII}R^{VII-6},
- e) (C=O)R^{VII-6},
- f) (C=O)OR^{VII-9},
- g) cyano,

- h) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- i) $\text{OR}^{\text{VII-10}}$,
- j) Ohet^{VII} ,
- k) $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$,
- l) $\text{SR}^{\text{VII-10}}$,
- m) Shet^{VII} ,
- n) $\text{NHCOR}^{\text{VII-12}}$,
- o) $\text{NHSO}_2\text{R}^{\text{VII-12}}$, or
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VII-11}}$, $\text{OR}^{\text{VII-13}}$, $\text{SR}^{\text{VII-10}}$, $\text{SR}^{\text{VII-13}}$, $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m^{\text{VII}}\text{R}^{\text{VII-9}}$, or
- q) $\text{R}^{\text{VII-1}}$ together with $\text{R}^{\text{VII-2}}$ form a carbocyclic or het^{VII} which may be optionally substituted by $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$, or C_{1-7} alkyl which may be optionally substituted by $\text{OR}^{\text{VII-14}}$;

$\text{R}^{\text{VII-6}}$ is

- a) C_{1-7} alkyl,
- b) $\text{NR}^{\text{VII-7}}\text{R}^{\text{VII-8}}$
- c) aryl^{VII}, or
- d) het^{VII} , wherein said het^{VII} is bound via a carbon atom;

$\text{R}^{\text{VII-7}}$ and $\text{R}^{\text{VII-8}}$ are independently

- a) H,
- b) aryl^{VII},
- c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $\text{NR}^{\text{VII-10}}\text{R}^{\text{VII-10}}$, $\text{R}^{\text{VII-11}}$, $\text{SO}_m^{\text{VII-9}}$, $\text{CONR}^{\text{VII-10}}\text{R}^{\text{VII-10}}$, or halo, or,
- d) $\text{R}^{\text{VII-7}}$ and $\text{R}^{\text{VII-8}}$ together with the nitrogen to which they are attached form a het^{VII} ;

$\text{R}^{\text{VII-9}}$ is

- a) aryl^{VII},

- b) het^{VII},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VII-10}R^{VII-10}, R^{VII-11}, SH, CONR^{VII-10}R^{VII-10}, or halo;

R^{VII-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VII-11} is

- a) OR^{VII-10},
- b) Ohet^{VII},
- c) Oaryl^{VII},
- d) CO₂R^{VII-10},
- e) het^{VII},
- f) aryl^{VII},
- g) CN, or
- h) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VII-11}, NR^{VII-7}R^{VII-8}, SO_m^{VII}R^{VII-9}, or C₁₋₇alkyl optionally substituted by R^{VII-11}, NR^{VII-7}R^{VII-8}, or SO_mR^{VII-9};

R^{VII-12} is

- a) H,
- b) het^{VII},
- c) aryl^{VII},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{VII-7}R^{VII-8} or R^{VII-11};

R^{VII-13} is

- a) (P=O)(OR^{VII-14})₂,

- b) $\text{CO}(\text{CH}_2)_n\text{CON}(\text{CH}_3) - (\text{CH}_2)_n\text{SO}_3^- \text{M}^+$,
- c) an amino acid,
- d) $\text{C}(=\text{O})\text{aryl}^{\text{VII}}$, or
- e) $\text{C}(=\text{O})\text{C}_{1-7}\text{alkyl}$ optionally substituted by $\text{NR}^{\text{VII}-7}\text{R}^{\text{VII}-8}$, aryl^{VII} , het^{VII} , CO_2H , or $\text{O}(\text{CH}_2)_n^{\text{VII}}\text{CO}_2\text{R}^{\text{VII}-14}$;

$\text{R}^{\text{VII}-14}$ is

- a) H, or
- b) $\text{C}_{1-7}\text{alkyl}$;

each n^{VII} is independently 1, 2, 3, 4 or 5;

each m^{VII} is independently 0, 1, or 2;

M^{VII} is sodium, potassium, or lithium;


aryl^{VII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, $\text{CO}_2\text{R}^{\text{VII}-14}$, CF_3 , $\text{C}_{1-6}\text{alkoxy}$, and $\text{C}_{1-6}\text{alkyl}$ which may be further substituted by one to three $\text{SR}^{\text{VII}-14}$, $\text{NR}^{\text{VII}-14}\text{R}^{\text{VII}-14}$, $\text{OR}^{\text{VII}-14}$, or $\text{CO}_2\text{R}^{\text{VII}-14}$ groups;

het^{VII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VII}-14}$, CF_3 , $\text{C}_{1-6}\text{alkoxy}$, oxo, oxime, and $\text{C}_{1-6}\text{alkyl}$ which may be further substituted by one to three $\text{SR}^{\text{VII}-14}$, $\text{NR}^{\text{VII}-14}\text{R}^{\text{VII}-14}$, $\text{OR}^{\text{VII}-14}$, or $\text{CO}_2\text{R}^{\text{VII}-14}$ groups;

wherein Formula VIII is



and pharmaceutically acceptable salts thereof,
wherein

A^{VIII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VIII-1} is

- a) R^{VIII-5},
- b) NR^{VIII-7}R^{VIII-8}, or
- c) SO₂R^{VIII-9};

R^{VIII-2} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) SO_mR^{VIII-6},
- d) OC₂₋₇ alkyl substituted by OH,
- e) SC₂₋₇ alkyl substituted by OH, or
- f) C₂₋₈ alkyl which is partially unsaturated and is optionally substituted by one or more substituents selected from R^{VIII-11}, OR^{VIII-13}, SR^{VIII-13}, NR^{VIII-7}R^{VIII-8}, halo, (C=O)C₁₋₇ alkyl or SO_mR^{VIII-9};

with the proviso that when R^{VIII-1} = R^{VIII-5} =

(CH₂CH₂O)_i^{VIII}R^{VIII-10}, then R^{VIII-2} may additionally represent

- a) H,
- b) halo,
- c) (C=O)R^{VIII-6},

- d) $(C=O)OR^{VIII-9}$,
- e) cyano,
- f) $OR^{VIII-10}$,
- g) het^{VIII} ,
- h) $NR^{VIII-7}R^{VIII-8}$,
- i) $SR^{VIII-10}$,
- j) het^{VIII} ,
- k) $NHCOR^{VIII-12}$,
- l) $NHSO_2R^{VIII-12}$, or
- m) R^{VIII-2} together with R^{VIII-3} or R^{VIII-4} form a carbocyclic or het^{VIII} which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-3} and R^{VIII-4} are independently:

- a) H,
- b) halo,
- c) $aryl^{VIII}$,
- d) $S(O)_m R^{VIII-6}$,
- e) $(C=O)R^{VIII-6}$,
- f) $(C=O)OR^{VIII-9}$,
- g) cyano,
- h) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
- i) $OR^{VIII-10}$,
- j) $Ohet^{VIII}$,
- k) $NR^{VIII-7}R^{VIII-8}$,
- l) $SR^{VIII-10}$,
- m) $Shet^{VIII}$,
- n) $NHCOR^{VIII-12}$,
- o) $NHSO_2R^{VIII-12}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more

substituents of the group $R^{VIII-11}$, $OR^{VIII-13}$, $SR^{VIII-10}$, $SR^{VIII-13}$, $NR^{VIII-7}R^{VIII-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{VIII}R^{VIII-9}$, or

- q) R^{VIII-4} together with R^{VIII-3} form a carbocyclic or het which may be optionally substituted by $NR^{VIII-7}R^{VIII-8}$, or C_{1-7} alkyl which may be optionally substituted by $OR^{VIII-14}$;

R^{VIII-5} is

- a) $(CH_2CH_2O)_iR^{VIII-10}$,
- b) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom,
- c) aryl^{VIII},
- d) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VIII-7}R^{VIII-8}$, $R^{VIII-11}$, SO_mR^{VIII-9} , or OC_{2-4} alkyl which may be further substituted by het^{VIII}, $OR^{VIII-10}$, or $NR^{VIII-7}R^{VIII-8}$, or
- e) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from $R^{VIII-11}$, $NR^{VIII-7}R^{VIII-8}$, $SO_m^{VIII}R^{VIII-9}$, or C_{1-7} alkyl optionally substituted by $R^{VIII-11}$, $NR^{VIII-7}R^{VIII-8}$, or $SO_m^{VIII}R^{VIII-9}$;

R^{VIII-6} is

- a) C_{1-7} alkyl,
- b) $NR^{VIII-7}R^{VIII-8}$,
- c) aryl^{VIII}, or
- d) het^{VIII}, wherein said het^{VIII} is bound via a carbon atom;

R^{VIII-7} and R^{VIII-8} are independently

- a) H,

- b) aryl^{VIII},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VIII-10}R^{VIII-10}, R^{VIII-11}, SO_m^{VIII}R^{VIII-9}, CONR^{VIII-10}R^{VIII-10}, or halo, or,
- d) R^{VIII-7} and R^{VIII-8} together with the nitrogen to which they are attached form a het^{VIII};

R^{VIII-9} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{VIII-10}R^{VIII-10}, R^{VIII-11}, SH, CONR^{VIII-10}R^{VIII-10}, or halo;

R^{VIII-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VIII-11} is

- a) OR^{VIII-10},
- b) Ohet^{VIII},
- c) Oaryl^{VIII},
- d) CO₂R^{VIII-10},
- e) het^{VIII},
- f) aryl^{VIII}, or
- g) CN;

R^{VIII-12} is

- a) H,

- b) het^{VIII} ,
- c) $\text{aryl}^{\text{VIII}}$,
- d) $\text{C}_{3-8}\text{cycloalkyl}$,
- e) methyl, or
- f) $\text{C}_{2-7}\text{alkyl}$ optionally substituted by $\text{NR}^{\text{VIII}-7}\text{R}^{\text{VIII}-8}$ or $\text{R}^{\text{VIII}-11}$;

$\text{R}^{\text{VIII}-13}$ is

- a) $(\text{P}=\text{O})(\text{OR}^{14})_2$,
- b) $\text{CO}(\text{CH}_2)_n^{\text{VIII}}\text{CON}(\text{CH}_3)-(\text{CH}_2)_n^{\text{VIII}}\text{SO}_3^-\text{M}^+$,
- c) an amino acid,
- d) $\text{C}(=\text{O})\text{aryl}^{\text{VIII}}$, or
- e) $\text{C}(=\text{O})\text{C}_{1-7}\text{alkyl}$ optionally substituted by $\text{NR}^{\text{VIII}-7}\text{R}^{\text{VIII}-8}$, $\text{aryl}^{\text{VIII}}$, het^{VIII} , CO_2H , or $\text{O}(\text{CH}_2)_n^{\text{VIII}}\text{CO}_2\text{R}^{\text{VIII}-14}$;

$\text{R}^{\text{VIII}-14}$ is

- a) H, or
- b) $\text{C}_{1-7}\text{alkyl}$;

each i^{VIII} is independently 2, 3, or 4;

each n^{VIII} is independently 1, 2, 3, 4 or 5;

each m^{VIII} is independently 0, 1, or 2;

M^{VIII} is sodium, potassium, or lithium;

$\text{aryl}^{\text{VIII}}$ is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;


wherein any $\text{aryl}^{\text{VIII}}$ is optionally substituted with one or more substituents selected from halo, OH, cyano, $\text{CO}_2\text{R}^{\text{VIII}-14}$, CF_3 , $\text{C}_{1-6}\text{alkoxy}$, and $\text{C}_{1-6}\text{alkyl}$ which may be further substituted by one to three $\text{SR}^{\text{VIII}-14}$, $\text{NR}^{\text{VIII}-14}\text{R}^{\text{VIII}-14}$, $\text{OR}^{\text{VIII}-14}$, or $\text{CO}_2\text{R}^{\text{VIII}-14}$ groups;

het^{VIII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the

group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VIII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO₂R^{VIII-14}, CF₃, C₁₋₆alkoxy, oxo, oxime, and C₁₋₆ alkyl which may be further substituted by one to three SR^{VIII-14}, NR^{VIII-14}R^{VIII-14}, OR^{VIII-14}, or CO₂R^{VIII-14} groups;

wherein Formula IX is



IX

and pharmaceutically acceptable salts thereof, wherein,

R^{IX-1} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{IX-2}, R^{IX-3} and R^{IX-4} are independently selected from:

- a) H,
- b) halo,
- c) aryl^{IX},
- d) S(O)_mR^{IX-6},
- e) (C=O)R^{IX-6},
- f) (C=O)OR^{IX-9},
- g) cyano,
- h) het^{IX}, wherein said ^{IX}-het is bound via a carbon atom,

- i) OR^{IX-10} ,
- j) $Ohet^{IX}$,
- k) $NR^{IX-7}R^{IX-8}$
- l) SR^{IX-10} ,
- m) $Shet^{IX}$,
- n) $NHCOR^{IX-12}$,
- o) $NHSO_2R^{IX-12}$, or
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{IX-11} , OR^{IX-13} , SR^{IX-10} , SR^{IX-13} , $NR^{IX-7}R^{IX-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{IX}R^{IX-9}$;

R^{IX-6} is

- a) C_{1-7} alkyl,
- b) $NR^{IX-7}R^{IX-8}$,
- c) aryl^{IX}, or
- d) het^{IX}, wherein said het^{IX} is bound via a carbon atom;

R^{IX-7} and R^{IX-8} are independently

- a) H,
- b) aryl^{IX},
- c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{IX-10}R^{IX-10}$, R^{IX-11} , SO_mR^{IX-9} , $CONR^{IX-10}R^{IX-10}$, or halo, or,
- d) R^{IX-7} and R^{IX-8} together with the nitrogen to which they are attached form a ^{IX}-het;

R^{IX-9} is

- a) aryl^{IX},
- b) het^{IX},
- c) C_{3-8} cycloalkyl,
- d) methyl, or

- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{IX-10}R^{IX-10}, R^{IX-11}, SH, CONR^{IX-10}R^{IX-10}, or halo;

R^{IX-10} is

- a) H,
 b) methyl, or
 c) C₂₋₇alkyl optionally substituted by OH;

R^{IX-11} is

- a) OR^{IX-10},
 b) Ohet^{IX},
 c) Oaryl^{IX},
 d) CO₂R^{IX-10},
 e) het^{IX},
 f) aryl^{IX}, or
 g) CN;

R^{IX-12} is

- a) H,
 b) het^{IX},
 c) aryl^{IX},
 d) C₃₋₈cycloalkyl,
 e) methyl, or
 f) C₂₋₇alkyl optionally substituted by NR^{IX-7}R^{IX-8} or R^{IX-11};

R^{IX-13} is

- a) (P=O) (OR^{IX-14})₂,
 b) CO(CH₂)_n^{IX}CON(CH₃) - (CH₂)_n^{IX}SO₃⁻M^{IX+},
 c) an amino acid,
 d) C(=O)aryl^{IX}, or
 e) C(=O)C₁₋₇alkyl optionally substituted by NR^{IX-7}R^{IX-8}, aryl^{IX}, het^{IX}, CO₂H, or O(CH₂)_nCO₂R^{IX-14};

R^{IX-14} is

a) H, or

b) C₁₋₇alkyl;

each n^{IX} is independently 1, 2, 3, 4 or 5;

each m^{IX} is independently 0, 1, or 2;

M^{IX} is sodium, potassium, or lithium;


aryl^{IX} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO₂R^{IX-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆alkyl which may be further substituted by one to three SR^{IX-14}, NR^{IX-14}R^{IX-14}, OR^{IX-14}, or CO₂R^{IX-14} groups;

het^{IX} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO₂R^{IX-14}, CF₃, C₁₋₆alkoxy, oxo, oxime, and C₁₋₆alkyl which may be further substituted by one to three SR^{IX-14}, NR^{IX-14}R^{IX-14}, OR^{IX-14}, or CO₂R^{IX-14} groups.

2. The use of claim 1, wherein the compound has the Formula



or a pharmaceutically acceptable salt thereof,
wherein,

A^{VI} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VI-1} is

- a) R^{VI-5}, or
- b) SO₂R^{VI-9}

R^{VI-2}, R^{VI-3} and R^{VI-4} may be the same or different and are selected from the group consisting of:

- a) H,
- b) halo,
- c) aryl^{VI},
- d) S(O)_mR^{VI-6},
- e) (C=O)R^{VI-6},
- f) (C=O)OR^{VI-9},
- g) cyano,
- h) het^{VI}, wherein said het^{VI} is bound via a carbon atom,
- i) OR^{VI-10},
- j) Ohet^{VI},
- k) NR^{VI-7}R^{VI-8}
- l) SR^{VI-10},

- m) Shet^{VI} ,
- n) $\text{NHCOR}^{\text{VI-12}}$,
- o) $\text{NHSO}_2\text{R}^{\text{VI-12}}$,
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VI-11}}$, $\text{OR}^{\text{VI-13}}$, $\text{SR}^{\text{VI-10}}$, $\text{SR}^{\text{VI-13}}$, $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m^{\text{VI}}\text{R}^{\text{VI-9}}$, and
- q) $\text{R}^{\text{VI-3}}$ together with $\text{R}^{\text{VI-2}}$ or $\text{R}^{\text{VI-4}}$ form a carbocyclic or het^{VI} which may be optionally substituted by $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, or C_{1-7} alkyl which may be optionally substituted by $\text{OR}^{\text{VI-14}}$;

$\text{R}^{\text{VI-5}}$ is

- a) $(\text{CH}_2\text{CH}_2\text{O})_i^{\text{VI}}\text{R}^{\text{VI-10}}$,
- b) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, $\text{R}^{\text{VI-11}}$, $\text{SO}_m^{\text{VI}}\text{R}^{\text{VI-9}}$, or OC_{2-4} alkyl which may be further substituted by het^{VI} , $\text{OR}^{\text{VI-10}}$, or $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, or
- c) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of $\text{R}^{\text{VI-11}}$, $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, $\text{SO}_m^{\text{VI}}\text{R}^{\text{VI-9}}$, or C_{1-7} alkyl optionally substituted by $\text{R}^{\text{VI-11}}$, $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$, or $\text{SO}_m^{\text{VI}}\text{R}^{\text{VI-9}}$;

$\text{R}^{\text{VI-6}}$ is

- a) C_{1-7} alkyl,
- b) $\text{NR}^{\text{VI-7}}\text{R}^{\text{VI-8}}$,
- c) aryl^{VI}, or
- d) het^{VI} , wherein said het^{VI} is bound via a carbon atom;

$\text{R}^{\text{VI-7}}$ and $\text{R}^{\text{VI-8}}$ are independently

- a) H,
- b) aryl^{VI},

- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of aryl^{VI}, NR^{VI-10}R^{VI-10}, R^{VI-11}, SO_m^{VI}R^{VI-9}, CONR^{VI-10}R^{VI-10}, or halo, or;
- d) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11}, NR^{VI-7}R^{VI-8}, SO_m^{VI}R^{VI-9}, or C₁₋₇alkyl optionally substituted by R^{VI-11}, NR^{VI-7}R^{VI-8}, or SO_m^{VI}R^{VI-9}, or
- e) R^{VI-7} and R^{VI-8} together with the nitrogen to which they are attached form a het^{VI};

R^{VI-9} is

- a) aryl^{VI},
- b) het^{VI},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from a group consisting of NR^{VI-10}R^{VI-10}, R^{VI-11}, SH, CONR^{VI-10}R^{VI-10}, or halo;

R^{VI-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VI-11} is

- a) OR¹⁰,
- b) Ohet^{VI},
- c) Oaryl^{VI},
- d) CO₂R¹⁰,
- e) het^{VI},
- f) aryl^{VI},
- g) CN, or

- h) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VI-11}, NR^{VI-7}R^{VI-8}, SO_m^{VI}R^{VI-9}, or C₁₋₇alkyl optionally substituted by R^{VI-11}, NR^{VI-7}R^{VI-8}, or SO_m^{VI}R^{VI-9};

R^{VI-12} is

- a) H,
 b) het^{VI},
 c) aryl^{VI},
 d) C₃₋₈cycloalkyl,
 e) methyl, or
 f) C₂₋₇alkyl optionally substituted by NR^{VI-7}R^{VI-8} or R^{VI-11};

R^{VI-13} is

- a) (P=O)(OR^{VI-14})₂,
 b) CO(CH₂)_n^{VI}CON(CH₃)-(CH₂)_nSO₃^{-VI+},
 c) an amino acid,
 d) C(=O)aryl^{VI},
 e) C(=O)C₁₋₇alkyl optionally substituted by NR^{VI-7}R^{VI-8}, aryl^{VI}, het^{VI}, CO₂H, or O(CH₂)_n^{VI}CO₂R^{VI-14}, or
 f) C(=O)NR^{VI-7}R^{VI-8}

R^{VI-14} is

- a) H, or
 b) C₁₋₇alkyl;

each i^{VI} is independently 2, 3, or 4;

each n^{VI} is independently 1, 2, 3, 4 or 5;

each m^{VI} is independently 0, 1, or 2;

M^{VI} is sodium, potassium, or lithium;

aryl^{VI} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VI} is optionally substituted with one or

more substituents selected from the group consisting of halo, OH, cyano, $\text{CO}_2\text{R}^{\text{VI}-14}$, CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which maybe further substituted by one to three $\text{SR}^{\text{VI}-14}$, $\text{NR}^{\text{VI}-14}\text{R}^{\text{VI}-14}$, $\text{OR}^{\text{VI}-14}$, or $\text{CO}_2\text{R}^{\text{VI}-14}$;

het^{VI} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VI} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, $\text{CO}_2\text{R}^{\text{VI}-14}$, CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which maybe further substituted by one to three $\text{SR}^{\text{VI}-14}$, $\text{NR}^{\text{VI}-14}\text{R}^{\text{VI}-14}$, $\text{OR}^{\text{VI}-14}$, or $\text{CO}_2\text{R}^{\text{VI}-14}$.

3. The use of Claim 2, wherein A^{VI} is Cl.

4. The use of Claim 2, wherein the compound administered is selected from the group consisting of

N-(4-chlorobenzyl)-6-iodo-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide; N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(hydroxymethyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide; N-(4-

chlorobenzyl)-8- {[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethynyl}-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(1-hydroxycyclohexyl)ethynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3S)-3-hydroxy-1-butynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butynyl}-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butynyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxy-1-butynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(5-hydroxy-1-pentynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[1R,2S]-2-hydroxycyclopentyl]ethynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-methyl-1-butynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-(phenylethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-phenyl-1-propynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butynyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[3-(methylsulfonyl)propyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[3-(methylsulfonyl)propyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[(2-hydroxyethoxy)methyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-tetrahydro-3-furanyl-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-(1,2-diethyl-4-pyrazolidinyl)-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-1-(3-oxetanyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-{3-[(3-hydroxypropyl)sulfonyl]propyl}-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[2-(2-ethoxyethoxy)ethyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-[(phenylsulfinyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-[(phenylsulfonyl)methyl]-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-
[(phenylsulfonyl)methyl]-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-
tetrahydro-2H-pyran-3-yl-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[(methylsulfonyl)methyl]-6-(4-
morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-[[(4-chlorophenyl) sulfinyl]methyl]-
6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-morpholinylmethyl)-4-oxo-1-
tetrahydro-2H-pyran-4-yl-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-
oxo-8-(4-thiomorpholinylmethyl)-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(4-hydroxy-1-piperidinyl)methyl]-1-
methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[(3R)-3-
hydroxypyrrolidinyl]methyl]-1-methyl-6-(4-
morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3-hydroxy-1-piperidinyl)methyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

[3-[(4-chlorobenzyl)amino]carbonyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-8-cinnolinyl]methyl 4-morpholinecarboxylate;

N-(4-chlorobenzyl)-8-(hydroxymethyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3-cyanobenzyl)amino]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6,8-bis(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-[(1-acetyl-4-piperidinyl)amino]-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[[1-methyl-2-(phenylsulfonyl)ethyl]amino]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[[3-(4-methoxyphenyl)-1-methylpropyl]amino]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-amino-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-8-
[(3-nitrobenzyl)amino]-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-
oxo-8-(tetrahydro-2H-pyran-4-ylamino)-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propyl)-1-methyl-4-oxo-
1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-6-(4-hydroxy-1-butyl)-1-methyl-4-oxo-
1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-{[(1R,2R)-1-hydroxy-2-
methylcyclohexyl]ethyl}-1-methyl-4-oxo-6-(tetrahydro-2H-
pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethyl)-1-methyl-6-(4-
morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-1-
methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-{4-[(4R)-2-oxo-1,3-
oxazolidin-4-yl]-1-butyl}-6-(tetrahydro-2H-pyran-4-
ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-6-(4-
morpholinylmethyl)-4-oxo-1,4-dihydro-3-
cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(1-hydroxycyclohexyl)ethyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3S)-3-hydroxy-1-butyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butyl}-N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butyl]-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[(3R)-3-hydroxy-1-butyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(5-hydroxy-1-pentyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-{{[(1R,2S)-2-hydroxycyclopentyl]ethyl}-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-methyl-1-butyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1H-imidazol-1-yl)-1-propyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1H-imidazol-1-yl)-1-propynyl]-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-4-oxo-8-(phenylethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-3-phenyl-1-propyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3-hydroxy-1-propyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;


N-(4-chlorobenzyl)-8-(4-hydroxy-1-butyl)-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-1-methyl-4-oxo-6-(tetrahydro-2H-pyran-4-ylmethyl)-1,4-dihydro-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-1-methyl-8-{[methyl(tetrahydro-2-furanylmethyl)amino]methyl}-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro-3-cinnolinecarboxamide;

and pharmaceutically acceptable salts thereof.

5. The use of Claim 1, wherein the compound administered has the Formula VII



VII

or a pharmaceutically acceptable salt thereof,
wherein,

A^{VII} is

a) Cl,

- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VII-1} is

- a) aryl^{VII},
- b) S(O)_m^{VII}R^{VII-6},
- c) (C=O)R^{VII-6}, with the proviso that if R^{VII-6} is NR^{VII-7}R^{VII-8}, then R^{VII-7} and R^{VII-8} do not both equal H
- d) (C=O)OR^{VII-9},
- e) cyano,
- f) het^{VII}, wherein said het^{VII} is bound via a carbon atom,
- g) Ohet^{VII},
- h) NR^{VII-7}R^{VII-8} with the proviso that R^{VII-7} and R^{VII-8} do not both equal H
- i) SR^{VII-10},
- j) Shet^{VII},
- k) NHCOR^{VII-12},
- l) NHSO₂R^{VII-12},
- m) C₁₋₇alkyl which is partially unsaturated and optionally substituted by one or more substituents of the group R^{VII-11}, OR^{VII-13}, SR^{VII-10}, SR^{VII-13}, NR^{VII-7}R^{VII-8}, halo, (C=O)C₁₋₇alkyl, or SO_mR^{VII-9}, or
- n) C₁₋₇alkyl which is substituted by one or more substituents of the group R^{VII-11}, OR^{VII-13}, SR^{VII-10}, SR^{VII-13}, NR^{VII-7}R^{VII-8}, halo, (C=O)C₁₋₇alkyl, or SO_mR^{VII-9};

R^{VII-2} is

- a) H,
- b) halo,
- c) aryl^{VII},
- d) S(O)_m^{VII}R^{VII-6},

- e) $(C=O)R^{VII-6}$,
- f) $(C=O)OR^{VII-9}$,
- g) cyano,
- h) het^{VII} , wherein said het^{VII} is bound via a carbon atom,
- i) OR^{VII-10} ,
- j) $Ohet^{VII}$,
- k) $NR^{VII-7}R^{VII-8}$,
- l) SR^{VII-10} ,
- m) $Shet^{VII}$,
- n) $NHCOR^{VII-12}$,
- o) $NHSO_2R^{VII-12}$, or
- p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{VII-11} , OR^{VII-13} , SR^{VII-10} , SR^{VII-13} , $NR^{VII-7}R^{VII-8}$, halo, $(C=O)C_{1-7}$ alkyl, or $SO_m^{VII}R^{VII-9}$, or
- q) R^{VII-1} together with R^{VII-2} form a carbocyclic or het^{VII} which may be optionally substituted by $NR^{VII-7}R^{VII-8}$, or C_{1-7} alkyl which may be optionally substituted by OR^{VII-14} ;

R^{VII-6} is

- a) C_{1-7} alkyl,
- b) $NR^{VII-7}R^{VII-8}$
- c) aryl^{VII}, or
- d) het^{VII} , wherein said het^{VII} is bound via a carbon atom;

R^{VII-7} and R^{VII-8} are independently

- a) H,
- b) aryl^{VII},
- c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VII-10}R^{VII-10}$, R^{VII-11} , SO_mR^{VII-9} , $CONR^{VII-10}R^{VII-10}$, or halo, or,

d) R^{VII-7} and R^{VII-8} together with the nitrogen to which they are attached form a het^{VII} ;

R^{VII-9} is

- a) aryl^{VII},
- b) het^{VII} ,
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $NR^{VII-10}R^{VII-10}$, R^{VII-11} , SH, $CONR^{VII-10}R^{VII-10}$, or halo;

R^{VII-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{VII-11} is

- a) OR^{VII-10} ,
- b) $Ohet^{VII}$,
- c) $Oaryl^{VII}$,
- d) CO_2R^{VII-10} ,
- e) het^{VII} ,
- f) aryl^{VII},
- g) CN, or
- h) C₃₋₈cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from a group consisting of R^{VII-11} , $NR^{VII-7}R^{VII-8}$, $SO_m^{VII}R^{VII-9}$, or C₁₋₇alkyl optionally substituted by R^{VII-11} , $NR^{VII-7}R^{VII-8}$, or $SO_m^{VII}R^{VII-9}$;

R^{VII-12} is

- a) H,
- b) het^{VII} ,
- c) aryl^{VII},
- d) C₃₋₈cycloalkyl,
- e) methyl, or

- f) C₂₋₇alkyl optionally substituted by NR^{VII-7}R^{VII-8}
or R^{VII-11};

R^{VII-13} is

- a) (P=O)(OR^{VII-14})₂,
b) CO(CH₂)_n^{VII}CON(CH₃)-(CH₂)_nSO₃⁻M^{VII+},
c) an amino acid,
d) C(=O)aryl^{VII}, or
e) C(=O)C₁₋₇alkyl optionally substituted by
NR^{VII-7}R^{VII-8}, aryl^{VII}, het^{VII}, CO₂H, or
O(CH₂)_n^{VII}CO₂R^{VII-14};

R^{VII-14} is

- a) H, or
b) C₁₋₇alkyl;

each n^{VII} is independently 1, 2, 3, 4 or 5;

each m^{VII} is independently 0, 1, or 2;

M^{VII} is sodium, potassium, or lithium;

aryl^{VII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO₂R^{VII-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆alkyl which may be further substituted by one to three SR^{VII-14}, NR^{VII-14}R^{VII-14}, OR^{VII-14}, or CO₂R^{VII-14} groups;

het^{VII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO₂R^{VII-14}, CF₃, C₁₋₆alkoxy, oxo, oxime, and C₁₋₆alkyl which may be further

substituted by one to three $\text{SR}^{\text{VII-14}}$, $\text{NR}^{\text{VII-14}}\text{R}^{\text{VII-14}}$, $\text{OR}^{\text{VII-14}}$, or $\text{CO}_2\text{R}^{\text{VII-14}}$ groups.

6. The use of Claim 5, wherein A^{VII} is Cl.

7. The use of Claim 6, wherein $\text{R}^{\text{VII-1}}$ is selected from the group consisting of CH_2 -morpholine, alkynyl- CH_2OH , CH_2 -(tetrahydro-2H-pyran-4-yl) and $(\text{CH}_2)_3\text{OH}$.

8. The use of Claim 6, wherein the compound administered is selected from the group consisting of

N-(4-chlorobenzyl)-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

Methyl 3-{[(4-chlorobenzyl)amino]carbonyl}-4-hydroxy-6-cinnolinecarboxylate;

N-(4-chlorobenzyl)-4-hydroxy-6-(hydroxymethyl)-3-cinnolinecarboxamide N-(4-chlorobenzyl)-8-(cyclopropylethynyl)-4-hydroxy-6-

(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1-hydroxycyclohexyl)ethynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propynyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3S)-3-hydroxy-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butynyl}-N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3R)-3-hydroxy-1-butynyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(5-hydroxy-1-pentynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{[(1R,2S)-2-hydroxycyclopentyl]ethynyl}-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-methyl-1-butynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propynyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(cyclopropylethyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1-hydroxycyclohexyl)ethyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-(3,3-dicyclopropyl-3-hydroxy-1-propyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3S)-3-hydroxy-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

8-{3-[(aminocarbonyl)amino]-3-methyl-1-butyl}-N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[3-methyl-3-(4-thioxo-1,3,5-triazinan-1-yl)-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(3R)-3-hydroxy-1-butyl]-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(4-morpholinylmethyl)-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(1,1-dioxido-4-thiomorpholinyl)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(5-hydroxy-1-pentyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[[1R,2S]-2-hydroxycyclopentyl]ethyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-methyl-1-butyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(4,5-dichloro-1H-imidazol-1-yl)-1-propyl]-4-hydroxy-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propyl)-6-(4-morpholinylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(phenylethynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-phenyl-1-propynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butynyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propynyl]-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethynyl]-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butynyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-1-propyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(phenylethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(3-hydroxy-3-phenyl-1-propyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-(4-hydroxy-1-butyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;


N-(4-chlorobenzyl)-8-[3-(dimethylamino)-1-propyl]-4-hydroxy-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-[(1R,2R)-1-hydroxy-2-methylcyclohexyl]ethyl)-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

N-(4-chlorobenzyl)-4-hydroxy-8-{4-[(4R)-2-oxo-1,3-oxazolidin-4-yl]-1-butyl}-6-(tetrahydro-2H-pyran-4-ylmethyl)-3-cinnolinecarboxamide;

and pharmaceutically acceptable salts thereof.

9. A use of Claim 1, wherein the compound is Formula VIII



and pharmaceutically acceptable salts thereof, wherein A^{VIII} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{VIII-1} is

- a) R^{VIII-5},
- b) NR^{VIII-7}R^{VIII-8}, or
- c) SO₂R^{VIII-9};

R^{VIII-2} is

- a) aryl^{VIII},
- b) het^{VIII},
- c) SO_m^{VIII}R^{VIII-6},
- d) OC₂₋₇ alkyl substituted by OH,
- e) SC₂₋₇ alkyl substituted by OH, or
- f) C₂₋₈ alkyl which is partially unsaturated and is optionally substituted by one or more substituents selected from R^{VIII-11}, OR^{VIII-13}, SR^{VIII-13}, NR^{VIII-7}R^{VIII-8}, halo, (C=O)C₁₋₇ alkyl or SO_m^{VIII}R^{VIII-9};

with the proviso that when R^{VIII-1} = R^{VIII-5} =

(CH₂CH₂O)_iR^{VIII-10}, then R^{VIII-2} may additionally represent

- a) H,
- b) halo,
- c) (C=O)R^{VIII-6},
- d) (C=O)OR^{VIII-9},
- e) cyano,
- f) OR^{VIII-10},
- g) Ohet^{VIII},
- h) NR^{VIII-7}R^{VIII-8},
- i) SR^{VIII-10},
- j) Shet^{VIII},
- k) NHCOR^{VIII-12},

- l) $\text{NHSO}_2\text{R}^{\text{VIII-12}}$, or
 m) $\text{R}^{\text{VIII-2}}$ together with $\text{R}^{\text{VIII-3}}$ or $\text{R}^{\text{VIII-4}}$ form a carbocyclic or het^{VIII} which may be optionally substituted by $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, or C_{1-7} alkyl which may be optionally substituted by $\text{OR}^{\text{VIII-14}}$;

$\text{R}^{\text{VIII-3}}$ and $\text{R}^{\text{VIII-4}}$ are independently:

- a) H,
 b) halo,
 c) aryl^{VIII},
 d) $\text{S}(\text{O})_m^{\text{VIII}}\text{R}^{\text{VIII-6}}$,
 e) $(\text{C}=\text{O})\text{R}^{\text{VIII-6}}$,
 f) $(\text{C}=\text{O})\text{OR}^{\text{VIII-9}}$,
 g) cyano,
 h) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
 i) $\text{OR}^{\text{VIII-10}}$,
 j) $\text{Ohet}^{\text{VIII}}$,
 k) $\text{R}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$,
 l) $\text{SR}^{\text{VIII-10}}$,
 m) $\text{Shet}^{\text{VIII}}$,
 n) $\text{NHCOR}^{\text{VIII-12}}$,
 o) $\text{NHSO}_2\text{R}^{\text{VIII-12}}$,
 p) C_{1-7} alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group $\text{R}^{\text{VIII-11}}$, $\text{OR}^{\text{VIII-13}}$, $\text{SR}^{\text{VIII-10}}$, $\text{SR}^{\text{VIII-13}}$, $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, halo, $(\text{C}=\text{O})\text{C}_{1-7}$ alkyl, or $\text{SO}_m^{\text{VIII}}\text{R}^{\text{VIII-9}}$, or
 q) $\text{R}^{\text{VIII-4}}$ together with $\text{R}^{\text{VIII-3}}$ form a carbocyclic or het^{VIII} which may be optionally substituted by $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, or C_{1-7} alkyl which may be optionally substituted by $\text{OR}^{\text{VIII-14}}$;

$\text{R}^{\text{VIII-5}}$ is

- a) $(\text{CH}_2\text{CH}_2\text{O})_i \text{R}^{\text{VIII-10}}$,
- b) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom,
- c) $\text{aryl}^{\text{VIII}}$,
- d) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, $\text{R}^{\text{VIII-11}}$, $\text{SO}_m \text{R}^{\text{VIII-9}}$, or OC_{2-4} alkyl which may be further substituted by het^{VIII} , $\text{OR}^{\text{VIII-10}}$, or $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, or
- e) C_{3-8} cycloalkyl which may be partially unsaturated and optionally substituted by one or more substituents selected from $\text{R}^{\text{VIII-11}}$, $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, $\text{SO}_m \text{R}^{\text{VIII-9}}$, or C_{1-7} alkyl optionally substituted by $\text{R}^{\text{VIII-11}}$, $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$, or $\text{SO}_m \text{R}^{\text{VIII-9}}$;

$\text{R}^{\text{VIII-6}}$ is

- a) C_{1-7} alkyl,
- b) $\text{NR}^{\text{VIII-7}}\text{R}^{\text{VIII-8}}$,
- c) $\text{aryl}^{\text{VIII}}$, or
- d) het^{VIII} , wherein said het^{VIII} is bound via a carbon atom;

$\text{R}^{\text{VIII-7}}$ and $\text{R}^{\text{VIII-8}}$ are independently

- a) H,
- b) $\text{aryl}^{\text{VIII}}$,
- c) C_{1-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $\text{NR}^{\text{VIII-10}}\text{R}^{\text{VIII-10}}$, $\text{R}^{\text{VIII-11}}$, $\text{SO}_m \text{R}^{\text{VIII-9}}$, $\text{CONR}^{\text{VIII-10}}\text{R}^{\text{VIII-10}}$, or halo, or,
- d) $\text{R}^{\text{VIII-7}}$ and $\text{R}^{\text{VIII-8}}$ together with the nitrogen to which they are attached form a het^{VIII} ;

$\text{R}^{\text{VIII-9}}$ is

- a) $\text{aryl}^{\text{VIII}}$,

- b) het^{VIII} ,
- c) C_{3-8} cycloalkyl,
- d) methyl, or
- e) C_{2-7} alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from $\text{NR}^{\text{VIII}-10}\text{R}^{\text{VIII}-10}$, $\text{R}^{\text{VIII}-11}$, SH , $\text{CONR}^{\text{VIII}-10}\text{R}^{\text{VIII}-10}$, or halo;

$\text{R}^{\text{VIII}-10}$ is

- a) H,
- b) methyl, or
- c) C_{2-7} alkyl optionally substituted by OH;

$\text{R}^{\text{VIII}-11}$ is

- a) $\text{OR}^{\text{VIII}-10}$,
- b) $\text{Ohet}^{\text{VIII}}$,
- c) $\text{Oaryl}^{\text{VIII}}$,
- d) $\text{CO}_2\text{R}^{\text{VIII}-10}$,
- e) het^{VIII} ,
- f) $\text{aryl}^{\text{VIII}}$, or
- g) CN;

$\text{R}^{\text{VIII}-12}$ is

- a) H,
- b) het^{VIII} ,
- c) $\text{aryl}^{\text{VIII}}$,
- d) C_{3-8} cycloalkyl,
- e) methyl, or
- f) C_{2-7} alkyl optionally substituted by $\text{NR}^{\text{VIII}-7}\text{R}^{\text{VIII}-8}$ or $\text{R}^{\text{VIII}-11}$;

$\text{R}^{\text{VIII}-13}$ is

- a) $(\text{P}=\text{O})(\text{OR}^{14})_2$,
- b) $\text{CO}(\text{CH}_2)_n^{\text{VIII}}\text{CON}(\text{CH}_3) - (\text{CH}_2)_n\text{SO}_3^-\text{M}^{\text{VIII}+}$,
- c) an amino acid,
- d) $\text{C}(=\text{O})\text{aryl}^{\text{VIII}}$, or

- e) C(=O)C₁₋₇alkyl optionally substituted by
 NR^{VIII-7}R^{VIII-8}, aryl^{VIII}, het^{VIII}, CO₂H, or
 O(CH₂)_n^{VIII}CO₂R^{VIII-14};

R^{VIII-14} is

- a) H, or
 b) C₁₋₇alkyl;

each i^{VIII} is independently 2, 3, or 4;

each n^{VIII} is independently 1, 2, 3, 4 or 5;

each m^{VIII} is independently 0, 1, or 2;

M^{VIII} is sodium, potassium, or lithium;

aryl^{VIII} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{VIII} is optionally substituted with one or more substituents selected from halo, OH, cyano, CO₂R^{VIII-14}, CF₃, C₁₋₆alkoxy, and C₁₋₆ alkyl which may be further substituted by one to three SR^{VIII-14}, NR^{VIII-14}R^{VIII-14}, OR^{VIII-14}, or CO₂R^{VIII-14} groups;

het^{VIII} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{VIII} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO₂R^{VIII-14}, CF₃, C₁₋₆alkoxy, oxo, oxime, and C₁₋₆ alkyl which may be further substituted by one to three SR^{VIII-14}, NR^{VIII-14}R^{VIII-14}, OR^{VIII-14}, or CO₂R^{VIII-14} groups.

10. The use of Claim 9, wherein A^{VIII} is Cl.

11. The use of Claim 9, wherein R^{VIII-2} is alkynyl-CH₂OH.

12. The use of Claim 9, wherein the compound administered is N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide, N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide; or a pharmaceutically acceptable salt thereof.

13. The use of Claim 11, wherein the compound administered is:

N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-1,7-dimethyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-Chlorobenzyl)-6-iodo-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-1,7-dimethyl-6-(4-morpholinylmethyl)-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-1-methyl-4,7-dioxo-1,4,7,8-tetrahydro[1,8]naphthyridine-3-carboxamide;


N-(4-chlorobenzyl)-6-(3-hydroxy-1-propynyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-6-(3-hydroxypropyl)-7-methoxy-1-methyl-4-oxo-1,4-dihydro[1,8]naphthyridine-3-carboxamide;

ethyl 6-([(4-chlorobenzyl)amino]carbonyl)-2-methoxy-8-methyl-5-oxo-5,8-dihydro[1,8]naphthyridine-3-carboxylate;

and pharmaceutically acceptable salts thereof.

14. A use of Claim 1, wherein the compound has the Formula IX



IX

and pharmaceutically acceptable salts thereof, wherein, R^{IX-1} is

- a) Cl,
- b) Br,
- c) CN,
- d) NO₂, or
- e) F;

R^{IX-2}, R^{IX-3} and R^{IX-4} are independently selected from:

- a) H,
- b) halo,
- c) aryl^{IX},
- d) S(O)_m^{IX}R^{IX-6},
- e) (C=O)R^{IX-6},
- f) (C=O)OR^{IX-9},

- g) cyano,
- h) het^{IX}, wherein said het^{IX} is bound via a carbon atom,
- i) OR^{IX-10},
- j) Ohet^{IX},
- k) NR^{IX-7}R^{IX-8}
- l) SR^{IX-10},
- m) S^{IX}-het,
- n) NHCOR^{IX-12},
- o) NHSO₂R^{IX-12}, or
- p) C₁₋₇alkyl which may be partially unsaturated and optionally substituted by one or more substituents of the group R^{IX-11}, OR^{IX-13}, SR^{IX-10}, SR^{IX-13}, NR^{IX-7}R^{IX-8}, halo, (C=O)C₁₋₇alkyl, or SO_mR^{IX-9};

R^{IX-6} is

- a) C₁₋₇alkyl,
- b) NR^{IX-7}R^{IX-8},
- c) aryl^{IX}, or
- d) het^{IX}, wherein said het^{IX} is bound via a carbon atom;

R^{IX-7} and R^{IX-8} are independently

- a) H,
- b) aryl^{IX},
- c) C₁₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{IX-10}R^{IX-10}, R^{IX-11}, SO_mR^{IX-9}, CONR^{IX-10}R^{IX-10}, or halo, or,
- d) R^{IX-7} and R^{IX-8} together with the nitrogen to which they are attached form a het^{IX};

R^{IX-9} is

- a) aryl^{IX},

- b) het^{IX},
- c) C₃₋₈cycloalkyl,
- d) methyl, or
- e) C₂₋₇alkyl which may be partially unsaturated and is optionally substituted by one or more substituents selected from NR^{IX-10}R^{IX-10}, R^{IX-11}, SH, CONR^{IX-10}R^{IX-10}, or halo;

R^{IX-10} is

- a) H,
- b) methyl, or
- c) C₂₋₇alkyl optionally substituted by OH;

R^{IX-11} is

- a) OR^{IX-10},
- b) Ohet^{IX},
- c) Oaryl^{IX},
- d) CO₂R^{IX-10},
- e) het^{IX},
- f) aryl^{IX}, or
- g) CN;

R^{IX-12} is

- a) H,
- b) het^{IX},
- c) aryl^{IX},
- d) C₃₋₈cycloalkyl,
- e) methyl, or
- f) C₂₋₇alkyl optionally substituted by NR^{IX-7}R^{IX-8} or R^{IX-11};

R^{IX-13} is

- a) (P=O)(OR^{IX-14})₂,
- b) CO(CH₂)_n^{IX}CON(CH₃)-(CH₂)_n^{IX}SO₃⁻M^{IX+},
- c) an amino acid,
- d) C(=O)aryl, or

- e) $C(=O)C_{1-7}$ alkyl optionally substituted by $NR^{IX-7}R^{IX-8}$, aryl^{IX}, het^{IX}, CO_2H , or $O(CH_2)_nCO_2R^{IX-14}$;

R^{IX-14} is

- a) H, or
b) C_{1-7} alkyl;

each n^{IX} is independently 1, 2, 3, 4 or 5;

each m^{IX} is independently 0, 1, or 2;

M^{IX} is sodium, potassium, or lithium;

aryl^{IX} is a phenyl radical or an ortho-fused bicyclic carbocyclic radical wherein at least one ring is aromatic;

wherein any aryl^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, CO_2R^{IX-14} , CF_3 , C_{1-6} alkoxy, and C_{1-6} alkyl which may be further substituted by one to three SR^{IX-14} , $NR^{IX-14}R^{IX-14}$, OR^{IX-14} , or CO_2R^{IX-14} groups;

het^{IX} is a four- (4), five- (5), six- (6), or seven- (7) membered saturated or unsaturated heterocyclic ring having 1, 2, or 3 heteroatoms selected from the group consisting of oxygen, sulfur, and nitrogen, which is optionally fused to a benzene ring, or any bicyclic heterocycle group;

wherein any het^{IX} is optionally substituted with one or more substituents selected from the group consisting of halo, OH, cyano, phenyl, CO_2R^{IX-14} , CF_3 , C_{1-6} alkoxy, oxo, oxime, and C_{1-6} alkyl which may be further substituted by one to three SR^{IX-14} , $NR^{IX-14}R^{IX-14}$, OR^{IX-14} , or CO_2R^{IX-14} groups.

15. The use of Claim 14, wherein R^{IX-1} is Cl.

16. The use of Claim 14, wherein the compound administered is selected from a group consisting of

N-(4-chlorobenzyl)-4-hydroxy-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-7-methyl-6-(tetrahydro-2H-pyran-4-ylmethyl)[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-7-methyl-6-(4-morpholinylmethyl)[1,8]naphthyridine-3-carboxamide;

6-bromo-N-(4-chlorobenzyl)-4-hydroxy-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-(3-hydroxy-1-propynyl)-7-methyl[1,8]naphthyridine-3-carboxamide;

N-(4-chlorobenzyl)-4-hydroxy-6-iodo-7-methyl[1,8]naphthyridine-3-carboxamide; and

Methyl 6-([(4-chlorobenzyl)amino]carbonyl)-5-hydroxy-2-methyl[1,8]naphthyridine-3-carboxylate.

17. The use according to any one of Claims 1 to 16, wherein said mammal is a human.

18. The use according to any one of Claims 1 to 16, wherein said mammal is a livestock or companion animal.

19. The use according to any one of Claims 1 to 16, wherein the amount administered is from about 0.1 to about 300 mg/kg of body weight.

20. The use according to any one of Claims 1 to 16, wherein the amount of compound administered is from about 1 to about 30 mg/kg of body weight.

21. The use according to any one of Claims 1 to 20, wherein the compound is administered parenterally, intravaginally, intranasally, topically, orally, or rectally.

INTERNATIONAL SEARCH REPORT

International Classification No
PCT/US 03/26838

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/4375 A61K31/502 A61P9/10 A61P31/12 A61P31/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, PAJ, WPI Data, BIOSIS, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category ° | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| Y | ZHOU ET AL: "Association between prior cytomegalovirus infection and the risk of restenosis after coronary atherectomy" NEW ENGLAND JOURNAL OF MEDICINE, THE, MASSACHUSETTS MEDICAL SOCIETY, WALTHAM, MA, US, no. 335, 29 August 1996 (1996-08-29), pages 624-630, XP002076129 ISSN: 0028-4793 page 624, right-hand column page 630, left-hand column, paragraph 3 --- | 1,9-21 |
| Y | WO 02 04444 A (WICNIENSKI NANCY ANNE ;PALMER JOHN R (US); UPJOHN CO (US); WILHITE) 17 January 2002 (2002-01-17) claims 1-114 ----- | 1,9-21 |

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

26 January 2004

Date of mailing of the international search report

05/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Albayrak, T

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 03/26838

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|--|------------------|-------------------------|------------------|
| WO 0204444 | A | AU 6970001 A | 21-01-2002 |
| | | EP 1301493 A2 | 16-04-2003 |
| | | WO 0204444 A2 | 17-01-2002 |
| | | US 2003207880 A1 | 06-11-2003 |
| | | US 2002025960 A1 | 28-02-2002 |
| ----- | | | |