



(11)

EP 3 567 106 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
13.11.2019 Bulletin 2019/46

(51) Int Cl.:
C12N 15/00 (2006.01) **C12N 15/86** (2006.01)
A61K 48/00 (2006.01)

(21) Application number: 18248250.5

(22) Date of filing: 22.04.2011

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: 23.04.2010 US 32762710 P

(62) Document number(s) of the earlier application(s) in accordance with Art. 76 EPC:
18190083.8 / 3 536 781
14181861.7 / 2 826 860
11772784.2 / 2 561 073

(71) Applicant: **University of Massachusetts**
Boston, MA 02110 (US)

(72) Inventors:
• **Gao, Guangping**
Westborough, MA 01581 (US)

- **Zhang, Hongwei**
Worcester, MA 01655 (US)
- **Wang, Hongyan**
Worcester, MA 01655 (US)
- **Xu, Zuoshang**
Wellesley, MA 02482 (US)

(74) Representative: **Potter Clarkson**
The Belgrave Centre
Talbot Street
Nottingham NG1 5GG (GB)

Remarks:

This application was filed on 28-12-2018 as a divisional application to the application mentioned under INID code 62.

(54) CNS TARGETING AAV VECTORS AND METHODS OF USE THEREOF

(57) The invention in some aspects relates to recombinant adeno-associated viruses useful for targeting transgenes to CNS tissue, and compositions comprising the same, and methods of use thereof. In some aspects, the invention provides methods and compositions for treating CNS-related disorders.




Fig. 1-1

Description

RELATED APPLICATIONS

5 [0001] This application claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional application USSN 61/327,627, filed April 23, 2010, and entitled "CNS Targeting AAV Vectors and Methods of Use Thereof," the entire contents of which are incorporated herein by reference.

FIELD OF THE INVENTION

10 [0002] The invention in some aspects relates to recombinant adeno-associated viruses useful for targeting transgenes to CNS tissue, and compositions comprising the same, and methods of use thereof.

BACKGROUND OF THE INVENTION

15 [0003] Gene therapy has been investigated for delivery of therapeutic genes to the CNS cells for treatment of various CNS disease, e.g., Canavan disease, ALS, Parkinson disease (PD), etc. In some limited cases, therapeutic benefits have been observed using certain viruses, e.g., recombinant adenovirus (rAd), lentivirus (LV) and adeno-associated virus (AAV) to express a variety of therapeutic genes. AAV2 has been used in clinical trials for treatment of PD and 20 Leber congenital amaurosis (an eye disease) and preliminary findings suggest symptomatic improvements without noticeable toxicity [2-4].

25 [0004] However, AAV-based gene therapy to treat CNS disease has still faced major obstacle. Many CNS diseases including, for example, ALS affect both cortical and spinal motor neurons that are distributed in a very broad area in the CNS. It has frequently been the case that viral vectors injected into CNS tissue transduce cells only in the vicinity of the injection site, have a very limited spread and generally have not impacted the lifespan in CNS disease animal models [See, e.g., Ref. 5]. Still, a variety of other viral administration methods have been tested. One example, involves injecting the viral particles into skeletal muscle and allowing the nerve terminals to internalize the viral genome, which is then retrogradely transported back to the spinal motor neurons. This approach has shown some positive results in certain mouse models [68]. However, to apply this method in larger mammals, like adult humans, would be impractical. Overall, 30 the transduction efficiency observed with muscle injection is relatively low. Some investigators have tried to improve this efficiency by modifying viral capsid proteins with the nerve binding domains of tetanus toxin or botulinum toxin. These efforts have not been fruitful due to various technical difficulties. Another problem with muscle injection in larger mammals, is a need for large doses, which is technically challenging, expensive, and carries a high risk for adverse effects, ranging from immune reaction to transduction of unintended cells (e.g., germ cells).

35 [0005] Another method that has been evaluated for delivering transgenes into motor neurons is to inject the virus into large nerves, which maximizes that exposure of the virus to motor axons, allowing the motor neurons to internalize the viral genome and retrogradely transport them back to the cell body. This method has been demonstrated to be more efficient in transducing motor neurons than muscle injection [9]. Still, to implement a method such as this in larger mammals would be challenging.

40 SUMMARY OF THE INVENTION

45 [0006] Aspects of the invention, are based on the discovery of recombinant AAVs that achieve wide-spread distribution throughout CNS tissue of a subject. In some embodiments, the rAAVs spread throughout CNS tissue following direct administration into the cerebrospinal fluid (CSF), e.g., via intrathecal and/or intracerebral injection. In other embodiments, the rAAVs cross the blood-brain-barrier and achieve wide-spread distribution throughout CNS tissue of a subject following intravenous administration. In some aspects the invention relates to rAAVs having distinct central nervous system tissue targeting capabilities (e.g., CNS tissue tropisms), which achieve stable and nontoxic gene transfer at high efficiencies. 50 Methods involving co-administration via intrathecal and intracerebral (e.g., intraventricular) injection of rAAVs are provided in some aspects. For example, it has been discovered that rAAVs having a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 achieves wide-spread distribution following intrathecal injection throughout the CNS, and thus, are particularly useful for treating CNS-associated disorders such as, for example, ALS. In still further aspects of the invention methods are provided for treating Canavan disease.

55 [0007] According to some aspects of the invention, methods for delivering a transgene to CNS tissue in a subject are provided. In some embodiments, the methods comprise administering an effective amount of a rAAV by intrathecal administration, wherein the rAAV comprises (i) a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 and (ii) a nucleic acid comprising a promoter operably linked with a transgene. In some embodiments, the methods further comprise administering an effective amount of the rAAV by intracerebral administration. In some embodiments,

the methods comprise administering an effective amount of a rAAV by intrathecal administration and by intracerebral administration, wherein the rAAV infects cells of CNS tissue in the subject and comprises a nucleic acid comprising a promoter operably linked with a transgene. In certain embodiments, the intracerebral administration is an intraventricular administration. In one embodiment, the intraventricular administration is an administration into a ventricular region of the forebrain of the subject. In certain embodiments, the intrathecal administration is in the lumbar region of the subject. In some embodiments, the dose of the rAAV for intrathecal administration is in a range of 10^{10} genome copies/subject to 10^{11} genome copies/subject. In some embodiments, the dose of the rAAV for intrathecal administration is in a range of 10^{11} genome copies/subject to 10^{12} genome copies/subject. In some embodiments, the dose of the rAAV for intrathecal administration is in a range of 10^{12} genome copies/subject to 10^{13} genome copies/subject. In some embodiments, the dose of the rAAV for intrathecal administration is in a range of 10^{13} genome copies/subject to 10^{14} genome copies/subject. In some embodiments, the dose of the rAAV for intracerebral administration is in a range of 10^{10} genome copies/subject to 10^{11} genome copies/subject. In some embodiments, the dose of the rAAV for intracerebral administration is in a range of 10^{11} genome copies/subject to 10^{12} genome copies/subject. In some embodiments, the dose of the rAAV for intracerebral administration is in a range of 10^{12} genome copies/subject to 10^{13} genome copies/subject. In some embodiments, the dose of the rAAV for intracerebral administration is in a range of 10^{13} genome copies/subject to 10^{14} genome copies/subject. In some embodiments, the dose of the rAAV for intracerebral or intrathecal administration is formulated for injection of a volume in a range of 1 μ l to 10 μ l. In some embodiments, the dose of the rAAV for intracerebral or intrathecal administration is formulated for injection of a volume in a range of 10 μ l to 100 μ l. In some embodiments, the rAAV for the intracerebral or intrathecal administration is formulated for injection of a volume in a range of 100 μ l to 1 ml. In some embodiments, the rAAV for the intracerebral or intrathecal administration is formulated for injection of a volume of 1 ml or more. In some embodiments, the transgene encodes a reporter protein. In certain embodiments, the reporter protein is a fluorescent protein, an enzyme that catalyzes a reaction yielding a detectable product, or a cell surface antigen. In certain embodiments, the enzyme is a luciferase, a beta-glucuronidase, a chloramphenicol acetyl-transferase, an aminoglycoside phosphotransferase, an aminocyclitol phosphotransferase, or a Puromycin N-acetyl-transferase. In some embodiments, the transgene is a CNS-associated gene. In some embodiments, the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA). In some embodiments, the transgene encodes an inhibitory RNA that binds specifically to SOD1 mRNA and inhibits expression of SOD1 in the subject. In some embodiments, the inhibitory RNA is an antisense RNA, a shRNA or a miRNA. In some embodiments, the inhibitory RNA has a sequence as set forth in SEQ ID NO: 26. Thus, according to some aspects of the invention a nucleic acid comprising a sequence as set forth in SEQ ID NO: 26 is provided. In some embodiments, a nucleic acid comprising a promoter operably linked with a region having a sequence as set forth in SEQ ID NO: 26 is provided.

[0008] In further aspects of the invention a recombinant AAV comprising a nucleic acid comprising a sequence as set forth in SEQ ID NO: 26 is provided. In some aspects of the invention a recombinant AAV comprising a nucleic acid comprising a promoter operably linked with a region having a sequence as set forth in SEQ ID NO: 26 is provided. In some embodiments the recombinant AAV further comprises a capsid protein comprising a sequence as set forth in SEQ ID NO: 9.

[0009] According to some aspects of the invention, methods for treating amyotrophic lateral sclerosis (ALS) in a subject in need thereof are provided. In some embodiments, the methods comprise administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises (i) a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 and (ii) a nucleic acid comprising a promoter operably linked with a region encoding an inhibitory RNA that binds specifically to SOD1 mRNA and inhibits expression of SOD1 in the subject. In some embodiments, the inhibitory RNA is an antisense RNA, a shRNA or a miRNA. In some embodiments, the inhibitory RNA has a sequence as set forth in SEQ ID NO: 26. In some embodiments, the methods comprise administering an effective amount of a rAAV to the subject, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding a sequence as set forth in SEQ ID NO: 26 and wherein the rAAV infects cells of CNS tissue in the subject.

[0010] According to some aspects of the invention, methods for delivering a transgene to a CNS tissue in a subject are provided that comprise administering an effective amount of a rAAV by intravenous administration, wherein the rAAV infects cells of CNS tissue in the subject and comprises a nucleic acid comprising a promoter operably linked with a transgene. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{10} genome copies/subject to 10^{11} genome copies/subject. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{11} genome copies/subject to 10^{12} genome copies/subject. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{12} genome copies/subject to 10^{13} genome copies/subject. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{13} genome copies/subject to 10^{14} genome copies/subject. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{14} genome copies/subject to 10^{15} genome copies/subject. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{10} genome copies/kg to 10^{11} genome copies/kg. In some embodiments, the dose of

the rAAV for intravenous administration is in a range of 10^{11} genome copies/kg to 10^{12} genome copies/kg. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{12} copies/kg to 10^{13} genome copies/kg. In some embodiments, the dose of the rAAV for intravenous administration is in a range of 10^{13} genome copies/kg to 10^{14} genome copies/kg.

5 [0011] According to some aspects of the invention, methods for delivering a transgene to a CNS tissue in a subject are provided that comprise administering to the subject an effective amount of a rAAV that comprises (i) a capsid protein having a sequence as set forth in any one of SEQ ID NO: 10 to 12 and (ii) a nucleic acid comprising a promoter operably linked with a transgene. In some embodiments, the methods comprise administering to the subject an effective amount of a rAAV comprising a transgene to a subject, wherein the rAAV comprises a capsid protein of a AAV serotype, or serotype variant, selected from the group consisting of: AAV1, AAV2, AAV5, AAV6, AV6.2, AAV7, AAV8, AAV9, rh.10, rh.39, rh.43 and CSp3, and wherein: (a) if the AAV serotype is AAV1, the administration route is not intracerebral, intramuscular, intranerve, or intraventricular and/or the subject is not a mouse, rat or feline; (b) if the AAV serotype is AAV2, the administration route is not intracerebral or intraventricular administration and/or the subject is not a rat, mouse, feline, marmoset, or macaque; (c) if the AAV serotype is AAV5, the administration route is not intracerebral or intraventricular administration and/or the subject is not a rat, mouse, or marmoset; (d) if the AAV serotype is AAV6, the subject is not a mouse; (e) if the AAV serotype is AAV7, the administration route is not intracerebral administration and/or the subject is not a mouse or macaque; (f) if the AAV serotype is AAV8, the administration route is not intracerebral, intraperitoneal, or intravascular administration and/or the subject is not a mouse or macaque; (g) if the AAV serotype is AAV9, the administration route is not intracerebral or intravascular administration and/or the subject is not a rat or mouse; and (h) if the AAV serotype is AAVrh.10, the administration route is not intracerebral or intravascular administration and/or the subject is not a rat or mouse. In some embodiments, the AAV serotype, or serotype variant, is selected from AAV1, AAV6, AAV7, rh.39, rh.43, and CSp3, and the administration route is intravascular administration. In some embodiments, the AAV serotype is AAV7 and the administration route is intravascular administration. In some embodiments, the CNS tissue is selected from cortex, hippocampus, thalamus, hypothalamus, cerebellum, brain stem, cervical spinal cord, thoracic spinal cord, and lumbar spinal cord. In some embodiments, the transgene encodes a reporter protein. In certain embodiments, the reporter protein is a fluorescent protein, an enzyme that catalyzes a reaction yielding a detectable product, or a cell surface antigen. In certain embodiments, the enzyme is a luciferase, a beta-glucuronidase, a chloramphenicol acetyltransferase, an aminoglycoside phosphotransferase, an aminocyclitol phosphotransferase, or a Puromycin N-acetyl-transferase. In some embodiments, the transgene is a CNS-associated gene. In certain embodiments, the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA). In some embodiments, the rAAV is administered by intravenous injection.

35 [0012] According to some aspects of the invention a rAAV that comprises (i) a capsid protein having a sequence as set forth in any one of SEQ ID NO: 10 to 12 and (ii) a nucleic acid comprising a promoter operably linked with a CNS-associated gene is provided. In certain embodiments, the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA). In some embodiments, mRNA expressed from the CNS-associated gene comprises a miRNA binding site of a miRNA that is preferentially expressed in non-CNS tissue. In certain embodiments, the miRNA binding site is a binding site for miR-122. In certain embodiments, the miRNA binding site is a binding site for miR-1. In some embodiments, mRNA expressed from the CNS-associated gene does not comprise a miRNA binding site of a miRNA that is preferentially expressed in CNS tissue. In some embodiments, the promoter is a CNS tissue specific promoter. In certain embodiments, the promoter is a promoter of a gene selected from: neuronal nuclei (NeuN), glial fibrillary acidic protein (GFAP), adenomatous polyposis coli (APC), and ionized calcium-binding adapter molecule 1 (Iba-1).

40 [0013] According to some aspects of the invention, a composition comprising a rAAV that comprises (i) a capsid protein having a sequence as set forth in SEQ ID NO: 10 to 12 and (ii) a nucleic acid comprising a promoter operably linked with a CNS-associated gene is provided. In certain embodiments the composition further comprises a pharmaceutically acceptable carrier. According to some aspects of the invention, a kit comprising a container housing the composition is provided. In some embodiments, the container is a sealed vial or ampule. In some embodiments, the container is a syringe.

45 [0014] According to some aspects of the invention, an isolated mammalian cell is provided that comprises a nucleic acid encoding a capsid protein having a sequence as set forth in any one of SEQ ID NO: 10 to 12 and a rAAV vector comprising a nucleic acid encoding a CNS-disease associated gene. In some embodiments, the isolated mammalian cell further comprises an AAV helper function vector. In some embodiments, isolated mammalian cell further comprises an accessory function vector. In certain embodiments, the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA).

[0015] According to further aspects of the invention, a method for treating Canavan disease in a subject in need thereof is provided. In some embodiments, the methods comprise administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises (i) a capsid protein other than a capsid protein of AAV serotype 2 and (ii) a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA). Any of the rAAV serotypes disclosed herein may be used in the methods for treating Canavan disease. In some embodiments, the rAAV has a capsid protein having an amino acid sequence as set forth in SEQ ID NO: 8 or 9 or a variant thereof. In some embodiments, administering is performed intrathecally or intracerebrally. In some embodiments, administering is performed intravascularly.

[0016] In some embodiments, the methods comprise administering an effective amount of a rAAV to CNS tissue of the subject by a route other than intracerebral administration, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA). In some embodiments, the methods comprise administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA); and evaluating kidney function in the subject at least once after the administration. Any suitable method known in the art may be used to evaluate a subject's kidney function. The evaluation may involve, for example, an examination of blood or urine urea nitrogen levels, an examination of blood or urine creatinine levels, a creatinine clearance rate examination, a glomerular filtration rate examination, a filtration fraction examination, a renal plasma flow examination, an ultrasound examination, a microscopic examination of a kidney tissue biopsy or any other suitable kidney function test. It should be appreciated that in some embodiments an improvement in a subject's kidney function following treatment with an rAAV-mediated gene therapy is indicative of efficacy of the gene therapy for treating Canavan disease.

[0017] In some embodiments, the methods comprise administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA); and evaluating vision of the subject at least once after the administration. Any suitable method known in the art may be used to evaluate a subject's vision. The evaluation may involve, for example, an external examination of the eye, a visual acuity examination, an examination of pupil function, a retinal examination, an ocular motility examination, an intraocular pressure test, or an ophthalmoscopic examination. The evaluation may involve a determination regarding a subject's ability to discriminate colors, objects or shapes or the ability of a subject to discern colors, objects or shapes from a particular distance. It should be appreciated that in some embodiments an improvement in a subject's vision following treatment with an rAAV-mediated gene therapy is indicative of efficacy of the gene therapy for treating Canavan disease.

[0018] In some embodiments, the nucleic acid expresses an aspartoacylase (ASPA) mRNA comprising one or more miRNA binds sites for one or more miRNAs that are more abundant in one or more non-CNS tissues in comparison to CNS tissue. Accordingly, in some embodiments, the mRNA is targeted for degradation by an miRNA in one or more non-CNS tissues. In some embodiments, the one or more non-CNS tissue is not kidney tissue or retinal tissue. In some embodiments, the one or more miRNAs that are more abundant in non-CNS tissues in comparison to CNS tissue are at least two-fold, at least three-fold, at least four-fold, at least five-fold, or at least ten-fold more abundant. MiRNAs that are more abundant in non-CNS tissue versus CNS tissue are known in the art. For example, one study discloses the expression levels of more than three-hundred different human miRNAs in 40 different tissues, including CNS tissue, kidney tissue. (See Liang Y, et al., Characterization of microRNA expression profiles in normal human tissues. *BMC Genomics*. 2007 Jun 12;8:166, the contents of which relating to miRNAs are incorporated herein by reference). Thus, in some embodiments, the skilled artisan could readily select (e.g., based on data such as are disclosed in Liang et al.) a suitable miRNA that is more abundant in non-CNS tissue and incorporate a binding site for that miRNA into the encoded mRNA.

[0019] Each of the limitations of the invention can encompass various embodiments of the invention. It is, therefore, anticipated that each of the limitations of the invention involving any one element or combinations of elements can be included in each aspect of the invention. This invention is not limited in its application to the details of construction and the arrangement of components set forth in the following description or illustrated in the drawings. The invention is capable of other embodiments and of being practiced or of being carried out in various ways.

50 BRIEF DESCRIPTION OF THE DRAWINGS

[0020]

Figure 1 depicts quantitative results of EGFP intensities from fluorescence microscopic images of a panel of CNS tissue sections from neonatal mice infected with various rAAVs harboring EGFP expression vectors. Neonatal mice were administered the rAAVs by intravenous administration (superficial temporal vein injection).

Figure 2 depicts quantitative results of EGFP intensities from fluorescence microscopic images of a panel of CNS tissue sections from adult mice infected with various rAAVs harboring EGFP expression vectors. Adult mice were

administered the rAAVs by intravenous administration (tail vein injection).

Figure 3 depicts quantitation of EGFP expression in neonatal mice spinal cord (cervical, thoracic and lumbar regions) 21 days post IV injection (5 mice per group). Neonatal mice were administered the rAAVs by intravenous administration (superficial temporal vein injection).

Figure 4A depicts results showing that direct CSF injection of AAVrh.10 harboring a EGFP gene leads to EGFP expression in broad areas of the CNS. Tissue sections, prepared 60 days post virus injection, from brainstem, cervical spinal cord, thoracic spinal cord and lumbar spinal cord are shown. Gray/black pixels correspond with EGFP expression.

Figure 4B depicts results showing that direct CSF injection of AAVrh.10 harboring a EGFP gene leads to EGFP expression in astrocytes. Gray/black pixels correspond with EGFP expression.

Figure 5A depicts a rAAVrh.10 vector that expresses a microRNA targeting SOD1. The construct employs CAG (chicken β -actin promoter with a CMV enhancer) to drive the expression of EGFP and miR-SOD1 that is located in an intron in the 3'-UTR. pA stands for poly A signal. ITRs mark the inverted repeats of the AAV.

Figure 5B depicts results of experiments that test the silencing potency of 9 different miRNA constructs, miR-SOD1#5 was found to silence SOD1 expression most potently.

Figure 5C depicts results of experiments in which miR-SOD1#5 was packaged into AAVrh. 10 and used to infect HEK293 cells. Total cellular protein was extracted 43 hours after the infection and blotted to detect SOD1. Scr stands for scrambled miRNA; Sod stands for miR-SOD1#5; and C stands for a control that expresses EGFP only.

Figure 5D depicts a plasmid map of pAAVscCB6 EGFPmir SOD5 (5243bp) (SEQ ID NO: 21).

Figure 6A depicts results of gene transfer studies in SOD1 (G93A) mutant mice showing that rAAV rh.10-SOD1 miRNA knockdowns levels of mutant SOD1 in astrocytes. Staining in motor neurons was also observed.

Figure 6B depicts results of gene transfer studies in SOD1 (G93A) mutant mice showing that rAAV rh.10-SOD1 shRNA increases live span, compared with a rAAV rh.10-scrambled miRNA.

Figure 7A depicts quantitation of EGFP expression in cervical, thoracic, and lumbar spinal cord tissue compared with life spans of individual mice infected with rAAV rh.10-SOD1 miRNA; rAAV rh.10-SOD1 was administered directly to the CSF.

Figure 7B depicts quantitation of EGFP expression in cervical, thoracic, and lumbar spinal cord tissue compared with life spans of individual mice infected with rAAV rh.10-scrambled miRNA; rAAV rh.10-scrambled miRNA was administered directly to the CSF.

Figure 8 depicts fluorescence microscopy analysis of mice that have been administered intrathecal injections of various AAVs. In this experiment, both AAV9 and AAVrh10 transduce cells along the full length of the spinal cord after a single injection into the CSF in lumbar subarachnoid space.

Figure 9 depicts the effects of AAV10-miR-SOD1 treatment. AAV10-miR-SOD1 treatment slows disease progression as indicated by the slower loss of body weight in treated compared with the control G93A mice.

Figure 10 depicts fluorescence microscopy analysis of mice that have been administered intrathecal injections of various AAVs. In this experiment, AAV9 and AAVrh10 can transduce cells in the broad forebrain areas after a single injection into the CSF in the third ventricle.

Figure 11 depicts fluorescence microscopy analysis of tissue sections from AAV9-injected mice. A single injection of AAV9 and AAVrh10 into the third ventricle can transduce cells in the broad forebrain areas, including cortex, hippocampus, striatum, thalamus, cerebellum and some scattered cells in the spinal cord. The same general pattern is also observed in AAV10-injected mice.

Figure 12 depicts an *in vitro* validation of artificial miRNA-binding sites for reporter silencing. Plasmids harboring the rAAVCBnLacZ genome with or without miR-1 or miR-122-binding sites were transfected into human hepatoma (HuH7) cells (a) which express miR-122 or cotransfected into 293 cells, together with a plasmid expressing either pri-miR-122 (b) or pri-miR-1 (c) at molar ratios of 1:3 (low) or 1:10 (high). 0X: no miRNA-binding site; 1X: one miRNA-binding site; 3X: three miRNA-binding sites. The cells were fixed and stained histochemically with X-gal 48 hours after transfection and blue cells counted. The percentage of nLacZ-positive cells in each transfection were compared to transfection of the control plasmid (prAAVCBnLacZ). CB, chicken β -actin; miR, microRNA; *n*LacZ, β -galactosidase reporter transgene; rAAV, recombinant adeno-associated viruses.

Figure 13 depicts an *in vivo* evaluation of endogenous miRNA-mediated transgene silencing in an rAAV9 transduction. (a-c) Adult male C588L/6 mice were injected intravenously with 5×10^{13} genome copies per kg (GC/kg) each of rAAV9CBnLacZ (no binding site), (a) rAAV9CBnLacZmiR-122BS (one miR-122-binding site) and rAAV9C8nLacZ-(miR-122BS)₃ (three miR-122-binding sites), (b) rAAV9CBnLacZ-miR-1 BS (one miR-1 binding site) and rAAV9CBnLacZ-(miR-1BS)₃ (three miR-1-binding sites), (c) rAAV9CBnLacZ-miR-1BS-miR-122BS (IX each binding site) and rAAV9CBnLacZ-(miR-1BS)₃-(miR-122BS)₃ (three miR-1 and three miR-122-binding sites). The animals were necropsied 4 weeks after vector administration, and appropriate tissues were harvested for cryosectioning and X-gal histochemical staining. miR, microRNA; *n*LacZ, β -galactosidase reporter transgene; rAAV, recombinant adeno-associated viruses, and (d) quantification of β -galactosidase activities in liver tissue from animals that

received rAAV*nLacZ* vectors with and without miRNA-binding sites.

Figure 14 depicts an analysis of expression levels of cognate miRNA, mRNA, and protein of endogenous miRNA target genes in mice transduced with rAAV9CB*nLacZ* with or without miRNA-binding sites. Total cellular RNA or protein was prepared from (a-c) liver or (d) heart. (a) Northern blot detection of miRNAs. U6 small nuclear RNA provided a loading control. (b) Quantitative reverse-transcription PCR measuring cyclin G1 mRNA. The data are presented as relative *cyclin G1* mRNA levels normalized to β -actin. (c,d) Western blot analyses of protein levels of endogenous targets of miR-122 and miR-1. Total cellular protein prepared from (c) liver or (d) heart was analyzed for cyclin G1 and calmodulin. (e) Serum cholesterol levels. Serum samples from mice that received rAAV9 with or without miRNA-binding sites were collected after 4 weeks and measured for total cholesterol, high-density lipoprotein (HDL) and low-density lipoprotein (LDL). miR, microRNA; *nLacZ*, β -galactosidase reporter transgene; rAAV, recombinant adeno-associated viruses.

Figure 15 depicts a molecular characterization of transgene mRNAs with or without miRNA binding sites. (a) Locations of the probes and primers, the sequences of mature miR-122 and its perfectly complementary binding site in the transgene mRNA are presented. (b) Total cellular RNA from liver was analyzed either by conventional reverse-transcription PCR (RT-PCR) by using primers that span a region between the 3' end of *nLacZ* and the 5' end of poly(A) signal (c) or by quantitative RT-PCR; data are presented as relative *nLacZ* mRNA levels normalized to β -actin. (d) For the northern blot analysis of *nLacZ* mRNA, 18S RNA served as a loading control, and the blots were hybridized with either a transgene DNA (e) or RNA probe. (f) In addition, poly(A) bearing mRNA from the liver of an animal received rAAV containing three miR-1- and three miR-122-binding sites was analyzed by 5' RACE; the PCR product was resolved on an ethidium bromide-stained agarose gel. miR, microRNA; *nLacZ*, β -galactosidase reporter transgene; rAAV, recombinant adeno-associated viruses.

Figure 16 depicts an alignment of sequences spanning the miRNA-binding sites and poly(A) signal regions recovered by 5 RACE. Poly(A)-containing mRNA was isolated from the (a) liver and (b) heart of an animal injected with rAAV9CB*nLacZ*-(miR-1BS)₃-(miR-122BS)₃. Twenty-one liver-derived and twenty-two heart-derived clones were sequenced. The putative cleavage sites in each clone are identified by arrows; the frequencies of miRNA-directed, site-specific cleavage for each miRNA-binding site are reported; triangles point to the positions of the expected miRNA-directed cleavage sites (a,b). miRNA, microRNA, *nLacZ*, β -galactosidase reporter transgene; rAAV, recombinant adeno-associated viruses.

Figure 17 depicts an endogenous miRNA-repressed, CNS-directed EGFP gene transfer by systemically delivered rAAV9. Ten-week-old male C57BL/6 mice were injected intravenously with scAAV9CBEGFP or scAAV9CB*nLacZ*(miR-1BS)₃-(miR-122BS)₃ at a dose of 2×10^{14} genome copies per kg (GC/ kg) body weight. The animals were necropsied 3 weeks later for whole body fixation by transcardiac perfusion. (a) Brain, spinal cord, liver, heart, and muscle were harvested for cryosectioning, immunofluorescent staining for EGFP (brain and cervical spinal cord), and fluorescence microscopy to detect EGFP. Total cellular DNA and RNA were extracted from brain, liver, heart and muscle to measure the amount of persistent vector genome by qPCR and EGFP mRNA by qRT-PCR. (b) For each tissue, the relative abundance of the EGFP mRNA containing miRNA-binding sites was compared to that of the EGFP mRNA lacking miRNA-binding sites. For each sample, mRNA abundance was normalized to the amount of vector genome detected in the tissue. EGFP, enhanced green fluorescent protein; miRNA, microRNA; *nLacZ*, β -galactosidase reporter transgene; qRT-PCR, quantitative reverse-transcription PCR; rAAV, recombinant adeno-associated viruses.

Figure 18 depicts a molecular model for endogenous miRNA-regulated rAAV expression. miRNA, microRNA; rAAV, recombinant adeno-associated viruses.

Figure 19 depicts a quantification of GFP intensity levels in the brain and spinal cord of neonatal mice transduced with various AAV vectors. 4×10^{11} genome copies (GCs) of ten different AAV vectors were injected into neonatal P1 pups through superficial vein. The mice were sacrificed 21 days after injection. The brain tissues were extracted and 40 μ m thick cryosections were prepared. The sections were stained against anti-EGFP antibody. The images were analyzed and the intensity/pixel values of all AAV serotypes in various regions in brain and spinal cord (A) were calculated by using Nikon NIS elements AR software version 3.2. Average intensities of the brain and spinal cord regions for different rAAVs were also presented (B). Region of interest (ROI) of each anatomical structure was fixed for all vectors to ensure the parallel comparison.

Figure 20 depicts a strong and widespread EGFP expression in neonatal mouse brain after intravenous injection of rAAVs. 4×10^{11} genome copies (GCs) of rAAVs 7, 9, rh.10, rh.39 and rh.43 were injected into neonatal P1 pups through superficial vein. The mice were sacrificed 21 days after injection. The brain tissues were extracted and 40 μ m thick cryosections were prepared. The sections were stained against anti-EGFP antibody. Bars represent 100 μ m. The regions shown are: olfactory bulb, striatum, hippocampus, cortex, hypothalamus, cerebellum and medulla.

Figure 21 depicts EGFP expression in neonatal mouse spinal cord after intravenous injection of rAAVs. 4×10^{11} GCs of rAAVs 7, 9, rh.10, rh.39 and rh.43 were injected into neonatal P1 pups through superficial vein. The mice were sacrificed 21 days after injection. The spinal cord tissues were extracted and 40 μ m thick cryosections were

prepared. The sections from cervical, thoracic and lumbar regions were stained against anti-EGFP antibody. Bars represent 100 μ m.

Figure 22 depicts EGFP expression in dorsal root ganglia transduced by intravascularly delivered rAAVsl, 2, 6, 6.2, 7, 9, rh.10 and rh.39. Neonatal pups received 4×10^{11} GCs of rAAVs at PI and were necropsied 21 days after injection. Forty μ m thick cryosections were processed for double immunohistochemical staining for EGFP (green) and Neurons (NeuN, red). Bars represent 75 μ m.

Figure 23 depicts confocal microscopic analysis of the transduced cell types in mouse CNS after systemic delivery of rAAVs to PI neonates. The 40 μ m thick brain and spinal cord sections of the animals treated with different rAAVs were co-stained against anti-EGFP antibody and cell-type specific markers. Anti-NeuN was used to stain neuronal cells; anti-GFAP was used to stain astrocytes; anti-Calbindin was used to stain Purkinje cells; anti-ChAT was used to stain motor neurons; anti-DARPP was used to stain dopaminergic neurons in the substantia nigra. All rAAVs were examined, but for each cell type, only one representative picture was shown here.

Figure 24 depicts a transduction of the brain ventricular structures by intravascularly delivered rAAVs. Neonatal pups received 4×10^{11} GCs of rAAVs at PI and were necropsied 21 days after injection. The choroid plexuses in different ventricles were well preserved during tissue process. Forty μ m thick cryosections were stained against anti-EGFP antibody. Bars represent 100 μ m.

Figure 25 depicts an analysis of purity and morphological integrity of rAAV vectors. A. Silver stained SDS-Page analysis of CsCl gradient purified rAAVCBEGFP vectors used in this study. Approximately 1.5×10^{10} virus particles each of rAAVs 1, 2, 5, 6, 6.2, 7, 9, rh10, rh39 and rh43 were loaded in the corresponding lane. B. Transmission electron microscopy of negative stained recombinant AAV virions. rAAV virions were spread on a freshly prepared carbon coated- Formvar support film and stained with 1% uranyl acetate for transmission microscopy. The images of virus particles from representative vector lots were taken at 92,000X and presented.

Figure 26 depicts a transduction of neonatal mouse dorsal root ganglia by systemically delivered rAAVs 1, 6, 6.2 and rh43. Neonatal pups received 4×10^{11} GCs of rAAVs at PI were necropsied 21 days after injection. Forty μ m thick cryosections were stained against anti-EGFP antibody. Bars represent 75 μ m.

Figure 27 depicts a transduction of the brain capillary vessels by intravascularly delivered rAAVs. Neonatal pups received 4×10^{11} GCs of rAAVs at PI were necropsied 21 days after injection. Forty μ m thick cryosections of the brains were stained against: (a) anti-EGFP antibody (AAV1, AAV6, AAV6.2, AAV7, AAV9, AAVrh.10, AAVrh.39 and AAVrh.43); (b) anti-EGFP and anti-CD34 antibodies (rh.10 only). Bars represent 100 μ m.

Figure 28 depicts an evaluation of microgliosis in mice brain after systemic delivery of rAAVs to PI neonates. The 40 μ m thick brain sections of the animals treated with different rAAVs were co-stained against anti-EGFP antibody and anti-IBa-1. Only the staining result of rAAVrh.10 was shown.

Figure 29 depicts native EGFP expression in mice CNS after systemic delivery of rAAVs to PI neonates. Neonatal pups received 4×10^{11} GCs of rAAVs at PI were necropsied 21 days after injection. Forty μ m thick cryosections were mounted and observed under microscope without immunostaining. The exposure times for each image were indicated.

Figure 30 depicts results showing the effects of rAAV based gene therapy in the treatment of Canavan disease. Figure 30A shows that treatment corrected gait and motor function of the CD mice. Figure 30B shows that treatment mitigated retinopathy and restored vision in CD mice. Figure 30C shows that NAA levels in the treated CD mice approach those in the normal mice. Figure 30D indicates that APSA activity is detected in the brains of CD mice. Figure 30E indicates APSA expression is detected in the brains of CD mice.

Figure 31A depicts that vacuolation in both brain and spinal cord of the treated mice is more patchy and variable with generally smaller-sized vacuoles and that some areas of the cerebral cortex show almost no vacuolation. Figure 31B shows ASPA expression in the cerebral cortex *in situ*.

Figure 32 depicts results of a quantitative analysis of vacuolation in various brain regions. Figure 32A shows that olfactory bulb had a dramatic mitigation in the white matter degeneration after gene therapy and that the large vacuoles were essentially eliminated in other tissues. Figure 32B shows results from a similar analysis on spinal cord sections.

Figure 33 depicts results from a histopathological evaluation of kidneys in the CD mice. Figure 33A shows that the renal tubular epithelium of the kidney was diffusely attenuated and exhibited enlargement of the tubular lumens in untreated CD mice. Figure 33B shows that treated CD mouse had normal glomeruli. Figure 33C and 33D depict results of an analysis of two lead candidate vectors, rAAV9 and rh.10, respectively, for efficiency of kidney transduction after IV delivery.

55 DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS OF THE INVENTION

[0021] Adeno-associated virus (AAV) is a small (26 nm) replication-defective, nonenveloped virus, that depends on the presence of a second virus, such as adenovirus or herpes virus, for its growth in cells. AAV is not known to cause

disease and induces a very mild immune response. AAV can infect both dividing and non-dividing cells and may incorporate its genome into that of the host cell. Aspects of the invention provide methods for delivering a transgene to a CNS tissue in a subject using recombinant AAV-based gene transfer. Accordingly, methods and compositions for treating CNS-related disorders are provided herein. Further aspects of the invention, are based on the discovery of rAAVs that achieve wide-spread distribution throughout CNS tissue. In some embodiments, the rAAVs spread throughout CNS tissue following direct administration into the cerebrospinal fluid (CSF), e.g., via intrathecal and/or intracerebral injection. In other embodiments, the rAAVs cross the blood-brain-barrier and achieve wide-spread distribution throughout CNS tissue of a subject following intravenous administration. Such rAAVs are useful for the treatment of CNS-related disorders, including, for example, amyotrophic lateral sclerosis (ALS) and Canavan disease (CD).

Methods and Compositions for Targeting CNS tissue

[0022] Methods for delivering a transgene to central nervous system (CNS) tissue in a subject are provided herein. The methods typically involve administering to a subject an effective amount of a rAAV comprising a nucleic acid vector for expressing a transgene in the subject. An "effective amount" of a rAAV is an amount sufficient to infect a sufficient number of cells of a target tissue in a subject. An effective amount of a rAAV may be an amount sufficient to have a therapeutic benefit in a subject, e.g., to extend the lifespan of a subject, to improve in the subject one or more symptoms of disease, e.g., a symptom of ALS, a symptom of Canavan disease, etc. In some cases, an effective amount of a rAAV may be an amount sufficient to produce a stable somatic transgenic animal model. The effective amount will depend on a variety of factors such as, for example, the species, age, weight, health of the subject, and the CNS tissue to be targeted, and may thus vary among subject and tissue. An effective amount may also depend on the mode of administration. For example, targeting a CNS tissue by intravascular injection may require different (e.g., higher) doses, in some cases, than targeting CNS tissue by intrathecal or intracerebral injection. In some cases, multiple doses of a rAAV are administered. An effective amount may also depend on the rAAV used. For example, dosages for targeting a CNS tissue may depend on the serotype (e.g., the capsid protein) of the rAAV. For example, the rAAV may have a capsid protein of a AAV serotype selected from the group consisting of: AAV1, AAV2, AAV5, AAV6, AAV6.2, AAV7, AAV8, AAV9, rh.10, rh.39, rh.43 and CSp3. In certain embodiments, the effective amount of rAAV is 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} genome copies per kg. In certain embodiments, the effective amount of rAAV is 10^{10} , 10^{11} , 10^{12} , 10^{13} , 10^{14} , or 10^{15} genome copies per subject.

[0023] A method for delivering a transgene to CNS tissue in a subject may comprise administering a rAAV by a single route or by multiple routes. For example, delivering a transgene to CNS tissue in a subject may comprise administering to the subject, by intravenous administration, an effective amount of a rAAV that crosses the blood-brain-barrier. Delivering a transgene to CNS tissue in a subject may comprise administering to the subject an effective amount of a rAAV by intrathecal administration or intracerebral administration, e.g., by intraventricular injection. A method for delivering a transgene to CNS tissue in a subject may comprise co-administering of an effective amount of a rAAV by two different administration routes, e.g., by intrathecal administration and by intracerebral administration. Co-administration may be performed at approximately the same time, or different times.

[0024] The CNS tissue to be targeted may be selected from cortex, hippocampus, thalamus, hypothalamus, cerebellum, brain stem, cervical spinal cord, thoracic spinal cord, and lumbar spinal cord, for example. The administration route for targeting CNS tissue typically depends on the AAV serotype. For example, in certain instances where the AAV serotype is selected from AAV1, AAV6, AAV6.2, AAV7, AAV8, AAV9, rh.10, rh.39, rh.43 and CSp3, the administration route may be intravascular injection. In some instances, for example where the AAV serotype is selected from AAV1, AAV2, AAV5, AAV6, AAV6.2, AAV7, AAV8, AAV9, rh.10, rh.39, rh.43 and CSp3, the administration route may be intrathecal and/or intracerebral injection.

Intravascular Administration

[0025] As used herein the term "intravascular administration" refers to the administration of an agent, e.g., a composition comprising a rAAV, into the vasculature of a subject, including the venous and arterial circulatory systems of the subject. Typically, rAAVs that cross the blood-brain-barrier may be delivered by intravascular administration for targeting CNS tissue. In some cases, intravascular (e.g., intravenous) administration facilitates the use of larger volumes than other forms of administration (e.g., intrathecal, intracerebral). Thus, large doses of rAAVs (e.g., up to 10^{15} GC/subject) can be delivered at one time by intravascular (e.g., intravenous) administration. Methods for intravascular administration are well known in the art and include for example, use of a hypodermic needle, peripheral cannula, central venous line, etc.

Intrathecal and/or Intracerebral Administration

[0026] As used herein the term "intrathecal administration" refers to the administration of an agent, e.g., a composition

comprising a rAAV, into the spinal canal. For example, intrathecal administration may comprise injection in the cervical region of the spinal canal, in the thoracic region of the spinal canal, or in the lumbar region of the spinal canal. Typically, intrathecal administration is performed by injecting an agent, e.g., a composition comprising a rAAV, into the subarachnoid cavity (subarachnoid space) of the spinal canal, which is the region between the arachnoid membrane and pia mater of the spinal canal. The subarachnoid space is occupied by spongy tissue consisting of trabeculæ (delicate connective tissue filaments that extend from the arachnoid mater and blend into the pia mater) and intercommunicating channels in which the cerebrospinal fluid is contained. In some embodiments, intrathecal administration is not administration into the spinal vasculature.

[0027] As used herein, the term "intracerebral administration" refers to administration of an agent into and/or around the brain. Intracerebral administration includes, but is not limited to, administration of an agent into the cerebrum, medulla, pons, cerebellum, intracranial cavity, and meninges surrounding the brain. Intracerebral administration may include administration into the dura mater, arachnoid mater, and pia mater of the brain. Intracerebral administration may include, in some embodiments, administration of an agent into the cerebrospinal fluid (CSF) of the subarachnoid space surrounding the brain. Intracerebral administration may include, in some embodiments, administration of an agent into ventricles of the brain, e.g., the right lateral ventricle, the left lateral ventricle, the third ventricle, the fourth ventricle. In some embodiments, intracerebral administration is not administration into the brain vasculature.

[0028] Intracerebral administration may involve direct injection into and/or around the brain. In some embodiments, intracerebral administration involves injection using stereotaxic procedures. Stereotaxic procedures are well known in the art and typically involve the use of a computer and a 3-dimensional scanning device that are used together to guide injection to a particular intracerebral region, e.g., a ventricular region. Micro-injection pumps (e.g., from World Precision Instruments) may also be used. In some embodiments, a microinjection pump is used to deliver a composition comprising a rAAV. In some embodiments, the infusion rate of the composition is in a range of 1 μ l / minute to 100 μ l / minute. As will be appreciated by the skilled artisan, infusion rates will depend on a variety of factors, including, for example, species of the subject, age of the subject, weight/size of the subject, serotype of the AAV, dosage required, intracerebral region targeted, etc. Thus, other infusion rates may be deemed by a skilled artisan to be appropriate in certain circumstances.

Methods and Compositions for Treating CNS-related Disorders

[0029] Methods and compositions for treating CNS-related disorders are also provided herein. As used herein, a "CNS-related disorder" is a disease or condition of the central nervous system. A CNS-related disorder may affect the spinal cord (e.g., a myelopathy), brain (e.g., a encephalopathy) or tissues surrounding the brain and spinal cord. A CNS-related disorder may be of a genetic origin, either inherited or acquired through a somatic mutation. A CNS-related disorder may be a psychological condition or disorder, e.g., Attention Deficient Hyperactivity Disorder, Autism Spectrum Disorder, Mood Disorder, Schizophrenia, Depression, Rett Syndrome, etc. A CNS-related disorder may be an autoimmune disorder. A CNS-related disorder may also be a cancer of the CNS, e.g., brain cancer. A CNS-related disorder that is a cancer may be a primary cancer of the CNS, e.g., an astrocytoma, glioblastomas, etc., or may be a cancer that has metastasized to CNS tissue, e.g., a lung cancer that has metastasized to the brain. Further non-limiting examples of CNS-related disorders, include Parkinson's Disease, Lysosomal Storage Disease, Ischemia, Neuropathic Pain, Amyotrophic lateral sclerosis (ALS), Multiple Sclerosis (MS), and Canavan disease (CD).

[0030] Methods for treating amyotrophic lateral sclerosis (ALS) in a subject in need thereof are provided herein. A subject in need of a treatment for ALS is a subject having or suspected of having ALS. In some cases, ALS has been linked to a mutation in the gene coding for superoxide dismutase (SOD1). Elevated levels of SOD1 appear to be associated with ALS in some instances. It has been shown that transgenic expression of shRNA against SOD1 can knockdown mutant SOD1 expression, delay disease onset and extend survival (Xia et al. 2006, *Neurobiol Dis* 23: 578). Intrathecal infusion of siRNA against SOD1 at disease onset has also been found to knockdown mutant SOD1 expression and extend survival (Wang et al. 2008, *JBC* 283: 15845). Furthermore, nerve injection of adenovirus expressing shRNA against SOD1 at the disease onset can knockdown mutant SOD1 expression and extend survival (Wu et al. 2009, *Antiox Redox Sig* 11: 1523).

[0031] Aspects of the invention, are based on the discovery of AAV-based therapies that achieve, with low-toxicity, long-term inhibition of SOD1 expression that is wide-spread throughout CNS tissue of the subject. Methods for treating ALS that are provided herein, typically involve administering to CNS tissue of a subject an effective amount of a rAAV that harbors a nucleic acid comprising a promoter operably linked with a region encoding an inhibitory RNA that binds specifically to SOD1 mRNA (e.g., that hybridizes specifically to a nucleic acid having a sequence as set forth in SEQ ID NO 17 or 19) and inhibits expression of SOD1 in the subject. It has been discovered that rAAVs having a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 achieve wide-spread distribution throughout the CNS following intrathecal injection and/or intracerebral injection, and thus, are particularly useful for treating ALS. This result is surprising in light of certain other rAAVs that infect cells only within the immediate vicinity of the injection site, or that achieve only a limited distribution, following intrathecal injection. Thus, rAAVs that achieve wide-spread distribution throughout the

CNS are particularly useful as gene transfer vectors for treating ALS.

[0032] In some embodiments, it has been discovered that co-administration by intrathecal injection and intracerebral injection, e.g., intraventricular injection, of rAAVs having a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 and a nucleic acid comprising a promoter operably linked with a region encoding an inhibitory RNA that binds specifically to SOD1 mRNA and inhibits expression of SOD1, achieves long-term inhibition of SOD1 and improves outcome (e.g., lifespan) in an animal model of ALS (See, e.g., Figure 6A). In some embodiments, the inhibitory RNA is an antisense RNA, a shRNA or a miRNA. The inhibitory RNA may have a sequence as set forth in SEQ ID NO: 26. The inhibitory RNA may have a sequence as set forth in any one of SEQ ID NO: 22 to 30. Thus, in some embodiments, a nucleic acid comprising a promoter operably linked with a nucleic acid having a sequence as set forth in any one of SEQ ID NO: 22 to 30 is provided. In some embodiments, a recombinant AAV that harbors a nucleic acid comprising a sequence as set forth in any one of SEQ ID NO: 22 to 30 is provided. The recombinant AAV may have a capsid protein comprising a sequence as set forth in SEQ ID NO: 9. The recombinant AAV may have a capsid protein comprising a sequence as set forth in any one of SEQ ID NO: 1 to 12.

[0033] Methods for treating Canavan disease (CD) in a subject in need thereof are provided herein. A subject in need of a treatment for CD is a subject having or suspected of having CD. Canavan disease is caused by a defective ASPA gene which is responsible for the production of the enzyme aspartoacylase. This enzyme normally breaks down the concentrated brain molecule N-acetyl aspartate. Decreased aspartoacylase activity in subjects with CD prevents the normal breakdown of N-acetyl aspartate, and the lack of breakdown appears to interfere with growth of the myelin sheath of the nerve fibers in the brain. Symptoms of Canavan disease, which may appear in early infancy and progress rapidly, may include mental retardation, loss of previously acquired motor skills, feeding difficulties, abnormal muscle tone (i.e., floppiness or stiffness), poor head control, and megalcephaly (abnormally enlarged head). Paralysis, blindness, or seizures may also occur. Aspects of the invention improve one or more symptoms of CD in a subject by administering to the subject a recombinant AAV harboring a nucleic acid that expresses aspartoacylase (ASPA). For example, a method for treating Canavan disease in a subject in need thereof may comprise administering an effective amount of a rAAV to CNS tissue of the subject by intravascular administration, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding ASPA (e.g., a region having a sequence as set forth in SEQ ID NO: 14 or 16). A method for treating Canavan disease in a subject in need thereof may comprise administering an effective amount of a rAAV to CNS tissue of the subject by intrathecal administration, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding ASPA. In some cases, methods for treating CD involve administering, to CNS tissue of the subject, an effective amount of a rAAV that comprises a capsid protein other than a capsid protein of AAV serotype 2 (e.g., other than a protein having an amino acid sequence as set forth in SEQ ID NO: 2) and a nucleic acid comprising a promoter operably linked with a region encoding ASPA. In another example, a method for treating Canavan disease in a subject in need thereof comprises administering an effective amount of a rAAV to CNS tissue of the subject by a route other than intracerebral administration, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding ASPA. In some embodiments, ASPA expressed in CNS tissue following administration of the rAAV results in a decrease in aspartoacylase activity and breakdown of N-acetyl aspartate in the CNS tissue. Thus, in some embodiments, a recombinant AAV vector is provided that comprises a nucleic acid encoding a sequence as set forth in SEQ ID NO: 14 or 16. In some embodiments, a recombinant AAV is provided that harbors a nucleic acid comprising a promoter operably linked with a region having a sequence as set forth in SEQ ID NO: 14 or 16. In some embodiments, a recombinant AAV is provided that harbors a nucleic acid comprising a promoter operably linked with a region encoding a protein having a sequence as set forth in SEQ ID NO: 13 or 15. The recombinant AAV may have a capsid protein comprising an amino acid sequence as set forth in any one of SEQ ID NO: 1 to 12. The recombinant AAV may have a capsid protein comprising a sequence as set forth in any one of SEQ ID NO: 1 and 3 to 12.

45 *Recombinant AAVs*

[0034] In some aspects, the invention provides isolated AAVs. As used herein with respect to AAVs, the term "isolated" refers to an AAV that has been isolated from its natural environment (e.g., from a host cell, tissue, or subject) or artificially produced. Isolated AAVs may be produced using recombinant methods. Such AAVs are referred to herein as "recombinant AAVs". Recombinant AAVs (rAAVs) preferably have tissue-specific targeting capabilities, such that a transgene of the rAAV will be delivered specifically to one or more predetermined tissue(s). The AAV capsid is an important element in determining these tissue-specific targeting capabilities. Thus, a rAAV having a capsid appropriate for the tissue being targeted can be selected. In some embodiments, the rAAV comprises a capsid protein having an amino acid sequence as set forth in any one of SEQ ID NOs 1 to 12, or a protein having substantial homology thereto.

[0035] Methods for obtaining recombinant AAVs having a desired capsid protein are well known in the art (See, for example, US 2003/0138772, the contents of which are incorporated herein by reference in their entirety). AAVs capsid protein that may be used in the rAAVs of the invention include, for example, those disclosed in G. Gao, et al., J. Virol., 78(12):6381-6388 (June 2004); G. Gao, et al, Proc Natl Acad Sci USA, 100(10):6081-6086 (May 13, 2003); US

2003-0138772, US 2007/0036760, US 2009/0197338, and U.S. provisional application serial number 61/182,084, filed May 28, 2009, the contents of which relating to AAVs capsid proteins and associated nucleotide and amino acid sequences are incorporated herein by reference. Typically the methods involve culturing a host cell which contains a nucleic acid sequence encoding an AAV capsid protein (e.g., a nucleic acid encoding a protein having a sequence as set forth in any one of SEQ ID NOs 1-12) or fragment thereof; a functional *rep* gene; a recombinant AAV vector composed of, AAV inverted terminal repeats (ITRs) and a transgene; and sufficient helper functions to permit packaging of the recombinant AAV vector into the AAV capsid proteins.

[0036] The components to be cultured in the host cell to package a rAAV vector in an AAV capsid may be provided to the host cell in *trans*. Alternatively, any one or more of the required components (e.g., recombinant AAV vector, *rep* sequences, cap sequences, and/or helper functions) may be provided by a stable host cell which has been engineered to contain one or more of the required components using methods known to those of skill in the art. Most suitably, such a stable host cell will contain the required component(s) under the control of an inducible promoter. However, the required component(s) may be under the control of a constitutive promoter. Examples of suitable inducible and constitutive promoters are provided herein, in the discussion of regulatory elements suitable for use with the transgene. In still another alternative, a selected stable host cell may contain selected component(s) under the control of a constitutive promoter and other selected component(s) under the control of one or more inducible promoters. For example, a stable host cell may be generated which is derived from 293 cells (which contain E1 helper functions under the control of a constitutive promoter), but which contain the *rep* and/or cap proteins under the control of inducible promoters. Still other stable host cells may be generated by one of skill in the art.

[0037] The recombinant AAV vector, *rep* sequences, cap sequences, and helper functions required for producing the rAAV of the invention may be delivered to the packaging host cell using any appropriate genetic element (vector). The selected genetic element may be delivered by any suitable method, including those described herein. The methods used to construct any embodiment of this invention are known to those with skill in nucleic acid manipulation and include genetic engineering, recombinant engineering, and synthetic techniques. See, e.g., Sambrook et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. Similarly, methods of generating rAAV virions are well known and the selection of a suitable method is not a limitation on the present invention. See, e.g., K. Fisher et al, *J. Virol.*, 70:520-532 (1993) and U.S. Pat. No. 5,478,745.

[0038] In some embodiments, recombinant AAVs may be produced using the triple transfection method (e.g., as described in detail in U.S. Pat. No. 6,001,650, the contents of which relating to the triple transfection method are incorporated herein by reference). Typically, the recombinant AAVs are produced by transfecting a host cell with a recombinant AAV vector (comprising a transgene) to be packaged into AAV particles, an AAV helper function vector, and an accessory function vector. An AAV helper function vector encodes the "AAV helper function" sequences (i.e., *rep* and cap), which function in *trans* for productive AAV replication and encapsidation. Preferably, the AAV helper function vector supports efficient AAV vector production without generating any detectable wild-type AAV virions (i.e., AAV virions containing functional *rep* and cap genes). Non-limiting examples of vectors suitable for use with the present invention include pHLP19, described in U.S. Pat. No. 6,001,650 and pRep6cap6 vector, described in U.S. Pat. No. 6,156,303, the entirety of both incorporated by reference herein. The accessory function vector encodes nucleotide sequences for non-AAV derived viral and/or cellular functions upon which AAV is dependent for replication (i.e., "accessory functions"). The accessory functions include those functions required for AAV replication, including, without limitation, those moieties involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of cap expression products, and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses such as adenovirus, herpesvirus (other than herpes simplex virus type-1), and vaccinia virus.

[0039] In some aspects, the invention provides transfected host cells. The term "transfection" is used to refer to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art. See, e.g., Graham et al. (1973) *Virology*, 52:456, Sambrook et al. (1989) *Molecular Cloning*, a laboratory manual, Cold Spring Harbor Laboratories, New York, Davis et al. (1986) *Basic Methods in Molecular Biology*, Elsevier, and Chu et al. (1981) *Gene* 13:197. Such techniques can be used to introduce one or more exogenous nucleic acids, such as a nucleotide integration vector and other nucleic acid molecules, into suitable host cells.

[0040] A "host cell" refers to any cell that harbors, or is capable of harboring, a substance of interest. Often a host cell is a mammalian cell. A host cell may be used as a recipient of an AAV helper construct, an AAV minigene plasmid, an accessory function vector, or other transfer DNA associated with the production of recombinant AAVs. The term includes the progeny of the original cell which has been transfected. Thus, a "host cell" as used herein may refer to a cell which has been transfected with an exogenous DNA sequence. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

[0041] In some aspects, the invention provides isolated cells. As used herein with respect to cell, the term "isolated" refers to a cell that has been isolated from its natural environment (e.g., from a tissue or subject). As used herein, the

term "cell line" refers to a population of cells capable of continuous or prolonged growth and division *in vitro*. Often, cell lines are clonal populations derived from a single progenitor cell. It is further known in the art that spontaneous or induced changes can occur in karyotype during storage or transfer of such clonal populations. Therefore, cells derived from the cell line referred to may not be precisely identical to the ancestral cells or cultures, and the cell line referred to includes such variants. As used herein, the terms "recombinant cell" refers to a cell into which an exogenous DNA segment, such as DNA segment that leads to the transcription of a biologically-active polypeptide or production of a biologically active nucleic acid such as an RNA, has been introduced.

[0042] As used herein, the term "vector" includes any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, artificial chromosome, virus, virion, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors. In some embodiments, useful vectors are contemplated to be those vectors in which the nucleic acid segment to be transcribed is positioned under the transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrases "operatively positioned," "under control" or "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene. The term "expression vector or construct" means any type of genetic construct containing a nucleic acid in which part or all of the nucleic acid encoding sequence is capable of being transcribed. In some embodiments, expression includes transcription of the nucleic acid, for example, to generate a biologically-active polypeptide product or inhibitory RNA (e.g., shRNA, miRNA) from a transcribed gene.

[0043] The foregoing methods for packaging recombinant vectors in desired AAV capsids to produce the rAAVs of the invention are not meant to be limiting and other suitable methods will be apparent to the skilled artisan.

Recombinant AAV vectors

[0044] "Recombinant AAV (rAAV) vectors" of the invention are typically composed of, at a minimum, a transgene and its regulatory sequences, and 5' and 3' AAV inverted terminal repeats (ITRs). It is this recombinant AAV vector which is packaged into a capsid protein and delivered to a selected target cell. In some embodiments, the transgene is a nucleic acid sequence, heterologous to the vector sequences, which encodes a polypeptide, protein, functional RNA molecule (e.g., miRNA, miRNA inhibitor) or other gene product, of interest. The nucleic acid coding sequence is operatively linked to regulatory components in a manner which permits transgene transcription, translation, and/or expression in a cell of a target tissue.

[0045] The AAV sequences of the vector typically comprise the cis-acting 5' and 3' inverted terminal repeat sequences (See, e.g., B. J. Carter, in "Handbook of Parvoviruses", ed., P. Tijsser, CRC Press, pp. 155 168 (1990)). The ITR sequences are about 145 bp in length. Preferably, substantially the entire sequences encoding the ITRs are used in the molecule, although some degree of minor modification of these sequences is permissible. The ability to modify these ITR sequences is within the skill of the art. (See, e.g., texts such as Sambrook et al, "Molecular Cloning. A Laboratory Manual", 2d ed., Cold Spring Harbor Laboratory, New York (1989); and K. Fisher et al., J Virol., 70:520 532 (1996)). An example of such a molecule employed in the present invention is a "cis-acting" plasmid containing the transgene, in which the selected transgene sequence and associated regulatory elements are flanked by the 5' and 3' AAV ITR sequences. The AAV ITR sequences may be obtained from any known AAV, including presently identified mammalian AAV types.

[0046] In addition to the major elements identified above for the recombinant AAV vector, the vector also includes conventional control elements which are operably linked to the transgene in a manner which permits its transcription, translation and/or expression in a cell transfected with the plasmid vector or infected with the virus produced by the invention. As used herein, "operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. A great number of expression control sequences, including promoters which are native, constitutive, inducible and/or tissue-specific, are known in the art and may be utilized.

[0047] As used herein, a nucleic acid sequence (e.g., coding sequence) and regulatory sequences are said to be operably linked when they are covalently linked in such a way as to place the expression or transcription of the nucleic acid sequence under the influence or control of the regulatory sequences. If it is desired that the nucleic acid sequences be translated into a functional protein, two DNA sequences are said to be operably linked if induction of a promoter in the 5' regulatory sequences results in the transcription of the coding sequence and if the nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of

the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably linked to a nucleic acid sequence if the promoter region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide. Similarly two or more coding regions are operably linked when they are linked in such a way that their transcription from a common promoter results in the expression of two or more proteins having been translated in frame. In some embodiments, operably linked coding sequences yield a fusion protein. In some embodiments, operably linked coding sequences yield a functional RNA (e.g., shRNA, miRNA).

[0048] For nucleic acids encoding proteins, a polyadenylation sequence generally is inserted following the transgene sequences and before the 3' AAV ITR sequence. A rAAV construct useful in the present invention may also contain an intron, desirably located between the promoter/enhancer sequence and the transgene. One possible intron sequence is derived from SV-40, and is referred to as the SV-40 T intron sequence. Another vector element that may be used is an internal ribosome entry site (IRES). An IRES sequence is used to produce more than one polypeptide from a single gene transcript. An IRES sequence would be used to produce a protein that contain more than one polypeptide chains. Selection of these and other common vector elements are conventional and many such sequences are available [see, e.g., Sambrook et al, and references cited therein at, for example, pages 3.18 3.26 and 16.17 16.27 and Ausubel et al., Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1989]. In some embodiments, a Foot and Mouth Disease Virus 2A sequence is included in polyprotein; this is a small peptide (approximately 18 amino acids in length) that has been shown to mediate the cleavage of polyproteins (Ryan, M D et al., EMBO, 1994; 4: 928-933; Mattion, N M et al., J Virology, November 1996; p. 8124-8127; Furler, S et al., Gene Therapy, 2001; 8: 864-873; and Halpin, C et al., The Plant Journal, 1999; 4: 453-459). The cleavage activity of the 2A sequence has previously been demonstrated in artificial systems including plasmids and gene therapy vectors (AAV and retroviruses) (Ryan, M D et al., EMBO, 1994; 4: 928-933; Mattion, N M et al., J Virology, November 1996; p. 8124-8127; Furler, S et al., Gene Therapy, 2001; 8: 864-873; and Halpin, C et al., The Plant Journal, 1999; 4: 453-459; de Felipe, P et al., Gene Therapy, 1999; 6: 198-208; de Felipe, P et al., Human Gene Therapy, 2000; 11: 1921-1931.; and Klump, H et al., Gene Therapy, 2001; 8: 811-817).

[0049] The precise nature of the regulatory sequences needed for gene expression in host cells may vary between species, tissues or cell types, but shall in general include, as necessary, 5' non-transcribed and 5' non-translated sequences involved with the initiation of transcription and translation respectively, such as a TATA box, capping sequence, CAAT sequence, enhancer elements, and the like. Especially, such 5' non-transcribed regulatory sequences will include a promoter region that includes a promoter sequence for transcriptional control of the operably joined gene. Regulatory sequences may also include enhancer sequences or upstream activator sequences as desired. The vectors of the invention may optionally include 5' leader or signal sequences. The choice and design of an appropriate vector is within the ability and discretion of one of ordinary skill in the art.

[0050] Examples of constitutive promoters include, without limitation, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter [Invitrogen].

[0051] Inducible promoters allow regulation of gene expression and can be regulated by exogenously supplied compounds, environmental factors such as temperature, or the presence of a specific physiological state, e.g., acute phase, a particular differentiation state of the cell, or in replicating cells only. Inducible promoters and inducible systems are available from a variety of commercial sources, including, without limitation, Invitrogen, Clontech and Ariad. Many other systems have been described and can be readily selected by one of skill in the art. Examples of inducible promoters regulated by exogenously supplied promoters include the zinc-inducible sheep metallothioneine (MT) promoter, the dexamethasone (Dex)-inducible mouse mammary tumor virus (MMTV) promoter, the T7 polymerase promoter system (WO 98/10088); the ecdysone insect promoter (No et al, Proc. Natl. Acad. Sci. USA, 93:3346-3351 (1996)), the tetracycline-repressible system (Gossen et al, Proc. Natl. Acad. Sci. USA, 89:5547-5551 (1992)), the tetracycline-inducible system (Gossen et al, Science, 268:1766-1769 (1995), see also Harvey et al, Curr. Opin. Chem. Biol., 2:512-518 (1998)), the RU486-inducible system (Wang et al, Nat. Biotech., 15:239-243 (1997) and Wang et al, Gene Ther., 4:432-441 (1997)) and the rapamycin-inducible system (Magari et al, J. Clin. Invest., 100:2865-2872 (1997)). Still other types of inducible promoters which may be useful in this context are those which are regulated by a specific physiological state, e.g., temperature, acute phase, a particular differentiation state of the cell, or in replicating cells only.

[0052] In another embodiment, the native promoter, or fragment thereof, for the transgene will be used. The native promoter may be preferred when it is desired that expression of the transgene should mimic the native expression. The native promoter may be used when expression of the transgene must be regulated temporally or developmentally, or in a tissue-specific manner, or in response to specific transcriptional stimuli. In a further embodiment, other native expression control elements, such as enhancer elements, polyadenylation sites or Kozak consensus sequences may also be used to mimic the native expression.

[0053] In some embodiments, the regulatory sequences impart tissue-specific gene expression capabilities. In some cases, the tissue-specific regulatory sequences bind tissue-specific transcription factors that induce transcription in a

tissue specific manner. Such tissue-specific regulatory sequences (e.g., promoters, enhancers, etc.) are well known in the art. Exemplary tissue-specific regulatory sequences include, but are not limited to the following tissue specific promoters: neuronal such as neuron-specific enolase (NSE) promoter (Andersen et al., *Cell. Mol. Neurobiol.*, 13:503-15 (1993)), neurofilament light-chain gene promoter (Piccioli et al., *Proc. Natl. Acad. Sci. USA*, 88:5611-5 (1991)), and the neuron-specific vgf gene promoter (Piccioli et al., *Neuron*, 15:373-84 (1995)). In some embodiments, the tissue-specific promoter is a promoter of a gene selected from: neuronal nuclei (NeuN), glial fibrillary acidic protein (GFAP), adenomatous polyposis coli (APC), and ionized calcium-binding adapter molecule 1 (Iba-1). Other appropriate tissue specific promoters will be apparent to the skilled artisan. In some embodiments, the promoter is a chicken Beta-actin promoter.

[0054] In some embodiments, one or more binding sites for one or more of miRNAs are incorporated in a transgene of a rAAV vector, to inhibit the expression of the transgene in one or more tissues of a subject harboring the transgenes, e.g., non-CNS tissues. The skilled artisan will appreciate that binding sites may be selected to control the expression of a transgene in a tissue specific manner. For example, expression of a transgene in the liver may be inhibited by incorporating a binding site for miR-122 such that mRNA expressed from the transgene binds to and is inhibited by miR-122 in the liver. Expression of a transgene in the heart may be inhibited by incorporating a binding site for miR-133a or miR-1, such that mRNA expressed from the transgene binds to and is inhibited by miR-133a or miR-1 in the heart. The miRNA target sites in the mRNA may be in the 5' UTR, the 3' UTR or in the coding region. Typically, the target site is in the 3' UTR of the mRNA. Furthermore, the transgene may be designed such that multiple miRNAs regulate the mRNA by recognizing the same or multiple sites. The presence of multiple miRNA binding sites may result in the cooperative action of multiple RISCs and provide highly efficient inhibition of expression. The target site sequence may comprise a total of 5-100, 10-60, or more nucleotides. The target site sequence may comprise at least 5 nucleotides of the sequence of a target gene binding site.

Transgene Coding Sequences: CNS-Related Genes

[0055] The composition of the transgene sequence of a rAAV vector will depend upon the use to which the resulting vector will be put. For example, one type of transgene sequence includes a reporter sequence, which upon expression produces a detectable signal. In another example, the transgene encodes a therapeutic protein or therapeutic functional RNA. In another example, the transgene encodes a protein or functional RNA that is intended to be used for research purposes, e.g., to create a somatic transgenic animal model harboring the transgene, e.g., to study the function of the transgene product. In another example, the transgene encodes a protein or functional RNA that is intended to be used to create an animal model of disease. Appropriate transgene coding sequences will be apparent to the skilled artisan.

[0056] In some aspects, the invention provides rAAV vectors for use in methods of preventing or treating one or more gene defects (e.g., heritable gene defects, somatic gene alterations) in a mammal, such as for example, a gene defect that results in a polypeptide deficiency or polypeptide excess in a subject, and particularly for treating or reducing the severity or extent of deficiency in a subject manifesting a CNS-associated disorder linked to a deficiency in such polypeptides in cells and tissues. In some embodiments, methods involve administration of a rAAV vector that encodes one or more therapeutic peptides, polypeptides, shRNAs, microRNAs, antisense nucleotides, etc. in a pharmaceutically-acceptable carrier to the subject in an amount and for a period of time sufficient to treat the CNS-associated disorder in the subject having or suspected of having such a disorder.

[0057] A rAAV vector may comprise as a transgene, a nucleic acid encoding a protein or functional RNA that modulates or treats a CNS-associated disorder. The following is a non-limiting list of genes associated with CNS-associated disorders: neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), aspartoacylase (ASPA), superoxide dismutase (SOD1) and amino acid decarboxylase (AADC). For example, a useful transgene in the treatment of Parkinson's disease encodes TH, which is a rate limiting enzyme in the synthesis of dopamine. A transgene encoding GTPCH, which generates the TH cofactor tetrahydrobiopterin, may also be used in the treatment of Parkinson's disease. A transgene encoding GDNF or BDNF, or AADC, which facilitates conversion of L-Dopa to DA, may also be used for the treatment of Parkinson's disease. For the treatment of ALS, a useful transgene may encode: GDNF, BDNF or CNTF. Also for the treatment of ALS, a useful transgene may encode a functional RNA, e.g., shRNA, miRNA, that inhibits the expression of SOD1. For the treatment of ischemia a useful transgene may encode NAIP or NGF. A transgene encoding Beta-glucuronidase (GUS) may be useful for the treatment of certain lysosomal storage diseases (e.g., Mucopolysaccharidosis type VII (MPS VII)). A transgene encoding a prodrug activation gene, e.g., HSV-Thymidine kinase which converts ganciclovir to a toxic nucleotide which disrupts DNA synthesis and leads to cell death, may be useful for treating certain cancers, e.g., when administered in combination with the prodrug. A transgene encoding an endogenous opioid, such a β-endorphin may be useful for treating pain. Other examples of transgenes that may be used in the rAAV vectors of the invention will be apparent to the skilled artisan (See, e.g., Costantini LC, et al., *Gene Therapy* (2000) 7, 93-109).

[0058] In some embodiments, the cloning capacity of the recombinant RNA vector may be limited and a desired coding

sequence may involve the complete replacement of the virus's 4.8 kilobase genome. Large genes may, therefore, not be suitable for use in a standard recombinant AAV vector, in some cases. The skilled artisan will appreciate that options are available in the art for overcoming a limited coding capacity. For example, the AAV ITRs of two genomes can anneal to form head to tail concatamers, almost doubling the capacity of the vector. Insertion of splice sites allows for the removal of the ITRs from the transcript. Other options for overcoming a limited cloning capacity will be apparent to the skilled artisan.

5 *Recombinant AAV Administration*

10 [0059] rAAVS are administered in sufficient amounts to transfect the cells of a desired tissue and to provide sufficient levels of gene transfer and expression without undue adverse effects. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the selected tissue (e.g., intracerebral administration, intrathecal administration), intravenous, oral, inhalation (including intranasal and intratracheal delivery), intraocular, intravenous, intramuscular, subcutaneous, intradermal, intratumoral, and other parental routes of administration. Routes of administration may be combined, if desired.

15 [0060] Delivery of certain rAAVs to a subject may be, for example, by administration into the bloodstream of the subject. Administration into the bloodstream may be by injection into a vein, an artery, or any other vascular conduit. Moreover, in certain instances, it may be desirable to deliver the rAAVs to brain tissue, meninges, neuronal cells, glial cells, astrocytes, oligodendrocytes, cerebrospinal fluid (CSF), interstitial spaces and the like. In some embodiments, recombinant AAVs may be delivered directly to the spinal cord or brain by injection into the ventricular region, as well as to 20 the striatum (e.g., the caudate nucleus or putamen of the striatum), and neuromuscular junction, or cerebellar lobule, with a needle, catheter or related device, using neurosurgical techniques known in the art, such as by stereotactic injection (see, e.g., Stein et al., J Virol 73:3424-3429, 1999; Davidson et al., PNAS 97:3428-3432, 2000; Davidson et al., Nat. Genet. 3:219-223, 1993; and Alisky and Davidson, Hum. Gene Ther. 11:2315-2329, 2000). In certain circumstances it will be desirable to deliver the rAAV-based therapeutic constructs in suitably formulated pharmaceutical 25 compositions disclosed herein either subcutaneously, intrapancreatically, intranasally, parenterally, intravenously, intramuscularly, intracerebrally, intrathecally, intracerebrally, orally, intraperitoneally, or by inhalation. In some embodiments, the administration modalities as described in U.S. Pat. Nos. 5,543,158; 5,641,515 and 5,399,363 (each specifically incorporated herein by reference in its entirety) may be used to deliver rAAVs.

30 *Recombinant AAV Compositions*

35 [0061] The rAAVs may be delivered to a subject in compositions according to any appropriate methods known in the art. The rAAV, preferably suspended in a physiologically compatible carrier (e.g., in a composition), may be administered to a subject, e.g., a human, mouse, rat, cat, dog, sheep, rabbit, horse, cow, goat, pig, guinea pig, hamster, chicken, turkey, or a non-human primate (e.g., Macaque). The compositions of the invention may comprise a rAAV alone, or in combination with one or more other viruses (e.g., a second rAAV encoding having one or more different transgenes). In some embodiments, a compositions comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more different rAAVs each having one or more different transgenes.

40 [0062] Suitable carriers may be readily selected by one of skill in the art in view of the indication for which the rAAV is directed. For example, one suitable carrier includes saline, which may be formulated with a variety of buffering solutions (e.g., phosphate buffered saline). Other exemplary carriers include sterile saline, lactose, sucrose, calcium phosphate, gelatin, dextran, agar, pectin, peanut oil, sesame oil, and water. The selection of the carrier is not a limitation of the present invention.

45 [0063] Optionally, the compositions of the invention may contain, in addition to the rAAV and carrier(s), other conventional pharmaceutical ingredients, such as preservatives, or chemical stabilizers. Suitable exemplary preservatives include chlorobutanol, potassium sorbate, sorbic acid, sulfur dioxide, propyl gallate, the parabens, ethyl vanillin, glycerin, phenol, and parachlorophenol. Suitable chemical stabilizers include gelatin and albumin.

50 [0064] The dose of rAAV virions required to achieve a desired effect or "therapeutic effect," e.g., the units of dose in vector genomes/per kilogram of body weight (vg/kg), will vary based on several factors including, but not limited to: the route of rAAV administration, the level of gene or RNA expression required to achieve a therapeutic effect, the specific disease or disorder being treated, and the stability of the gene or RNA product. One of skill in the art can readily determine a rAAV virion dose range to treat a subject having a particular disease or disorder based on the aforementioned factors, as well as other factors that are well known in the art. An effective amount of the rAAV is generally in the range of from about 10 μ l to about 100 ml of solution containing from about 10^9 to 10^{16} genome copies per subject. Other volumes of solution may be used. The volume used will typically depend, among other things, on the size of the subject, the dose of the rAAV, and the route of administration. For example, for intrathecal or intracerebral administration a volume in range of 1 μ l to 10 μ l or 10 μ l to 100 μ l may be used. For intravenous administration a volume in range of 10 μ l to 100 μ l, 100 μ l to 1 ml, 1 ml to 10 ml, or more may be used. In some cases, a dosage between about 10^{10} to 10^{12} rAAV

genome copies per subject is appropriate. In certain embodiments, 10^{12} rAAV genome copies per subject is effective to target CNS tissues. In some embodiments the rAAV is administered at a dose of 10^{10} , 10^{11} , 10^{12} , 10^{13} , 10^{14} , or 10^{15} genome copies per subject. In some embodiments the rAAV is administered at a dose of 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} genome copies per kg.

5 [0065] In some embodiments, rAAV compositions are formulated to reduce aggregation of AAV particles in the composition, particularly where high rAAV concentrations are present (e.g., $\sim 10^{13}$ GC/ml or more). Methods for reducing aggregation of rAAVs are well known in the art and, include, for example, addition of surfactants, pH adjustment, salt concentration adjustment, etc. (See, e.g., Wright FR, et al., Molecular Therapy (2005) 12, 171-178, the contents of which are incorporated herein by reference.)

10 [0066] Formulation of pharmaceutically-acceptable excipients and carrier solutions is well-known to those of skill in the art, as is the development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens. Typically, these formulations may contain at least about 0.1% of the active ingredient or more, although the percentage of the active ingredient(s) may, of course, be varied and may conveniently be between about 1 or 2% and about 70% or 80% or more of the weight or volume of the total formulation. Naturally, 15 the amount of active ingredient in each therapeutically-useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

20 [0067] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In many cases the form is sterile and fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and 25 storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size 30 in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents 35 delaying absorption, for example, aluminum monostearate and gelatin.

40 [0068] For administration of an injectable aqueous solution, for example, the solution may be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion, (see for example, "Remington's Pharmaceutical Sciences" 15th Edition, pages 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the host. The person responsible for administration will, in any event, determine the appropriate dose for the individual host.

45 [0069] Sterile injectable solutions are prepared by incorporating the active rAAV in the required amount in the appropriate solvent with various of the other ingredients enumerated herein, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

50 [0070] The rAAV compositions disclosed herein may also be formulated in a neutral or salt form. Pharmaceutically-acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms such as injectable solutions, drug-release capsules, and the like.

55 [0071] As used herein, "carrier" includes any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids,

and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Supplementary active ingredients can also be incorporated into the compositions. The phrase "pharmaceutically-acceptable" refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a host.

5 [0072] Delivery vehicles such as liposomes, nanocapsules, microparticles, microspheres, lipid particles, vesicles, and the like, may be used for the introduction of the compositions of the present invention into suitable host cells. In particular, the rAAV vector delivered *transgenes* may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, or a nanoparticle or the like.

10 [0073] Such formulations may be preferred for the introduction of pharmaceutically acceptable formulations of the nucleic acids or the rAAV constructs disclosed herein. The formation and use of liposomes is generally known to those of skill in the art. Recently, liposomes were developed with improved serum stability and circulation half-times (U.S. Pat. No. 5,741,516). Further, various methods of liposome and liposome like preparations as potential drug carriers have been described (; U.S. Pat. Nos. 5,567,434; 5,552,157; 5,565,213; 5,738,868 and 5,795,587).

15 [0074] Liposomes have been used successfully with a number of cell types that are normally resistant to transfection by other procedures. In addition, liposomes are free of the DNA length constraints that are typical of viral-based delivery systems. Liposomes have been used effectively to introduce genes, drugs, radiotherapeutic agents, viruses, transcription factors and allosteric effectors into a variety of cultured cell lines and animals. In addition, several successful clinical trials examining the effectiveness of liposome-mediated drug delivery have been completed.

20 [0075] Liposomes are formed from phospholipids that are dispersed in an aqueous medium and spontaneously form multilamellar concentric bilayer vesicles (also termed multilamellar vesicles (MLVs)). MLVs generally have diameters of from 25 nm to 4 μ m. Sonication of MLVs results in the formation of small unilamellar vesicles (SUVs) with diameters in the range of 200 to 500 ANG. , containing an aqueous solution in the core.

25 [0076] Alternatively, nanocapsule formulations of the rAAV may be used. Nanocapsules can generally entrap substances in a stable and reproducible way. To avoid side effects due to intracellular polymeric overloading, such ultrafine particles (sized around 0.1 μ m) should be designed using polymers able to be degraded *in vivo*. Biodegradable polyalkyl-cyanoacrylate nanoparticles that meet these requirements are contemplated for use.

30 [0077] In addition to the methods of delivery described above, the following techniques are also contemplated as alternative methods of delivering the rAAV compositions to a host. Sonophoresis (i.e., ultrasound) has been used and described in U.S. Pat. No. 5,656,016 as a device for enhancing the rate and efficacy of drug permeation into and through the circulatory system. Other drug delivery alternatives contemplated are intraosseous injection (U.S. Pat. No. 5,779,708), microchip devices (U.S. Pat. No. 5,797,898), ophthalmic formulations (Bourlaiss et al., 1998), transdermal matrices (U.S. Pat. Nos. 5,770,219 and 5,783,208) and feedback-controlled delivery (U.S. Pat. No. 5,697,899).

Kits and Related Compositions

35 [0078] The agents described herein may, in some embodiments, be assembled into pharmaceutical or diagnostic or research kits to facilitate their use in therapeutic, diagnostic or research applications. A kit may include one or more containers housing the components of the invention and instructions for use. Specifically, such kits may include one or more agents described herein, along with instructions describing the intended application and the proper use of these agents. In certain embodiments agents in a kit may be in a pharmaceutical formulation and dosage suitable for a particular application and for a method of administration of the agents. Kits for research purposes may contain the components in appropriate concentrations or quantities for running various experiments.

40 [0079] The kit may be designed to facilitate use of the methods described herein by researchers and can take many forms. Each of the compositions of the kit, where applicable, may be provided in liquid form (e.g., in solution), or in solid form, (e.g., a dry powder). In certain cases, some of the compositions may be constitutable or otherwise processable (e.g., to an active form), for example, by the addition of a suitable solvent or other species (for example, water or a cell culture medium), which may or may not be provided with the kit. As used herein, "instructions" can define a component of instruction and/or promotion, and typically involve written instructions on or associated with packaging of the invention. Instructions also can include any oral or electronic instructions provided in any manner such that a user will clearly recognize that the instructions are to be associated with the kit, for example, audiovisual (e.g., videotape, DVD, etc.), Internet, and/or web-based communications, etc. The written instructions may be in a form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which instructions can also reflects approval by the agency of manufacture, use or sale for animal administration.

45 [0080] The kit may contain any one or more of the components described herein in one or more containers. As an example, in one embodiment, the kit may include instructions for mixing one or more components of the kit and/or isolating and mixing a sample and applying to a subject. The kit may include a container housing agents described herein. The agents may be in the form of a liquid, gel or solid (powder). The agents may be prepared sterilely, packaged in syringe and shipped refrigerated. Alternatively it may be housed in a vial or other container for storage. A second

container may have other agents prepared sterilely. Alternatively the kit may include the active agents premixed and shipped in a syringe, vial, tube, or other container. The kit may have one or more or all of the components required to administer the agents to a subject, such as a syringe, topical application devices, or IV needle tubing and bag.

5 EXAMPLES

EXAMPLE 1: Characterization of 12 AAV vectors for intravascular delivery to target CNS and detarget non-CNS tissues by miRNA regulation

10 [0081] The CNS gene transfer properties of 12 scAAVEGFP vectors of different serotypes, or natural variants were evaluated. RAAVs that cross the blood-brain-barrier (BBB) and target oligodendrocytes were discovered. Experiments were performed in neonatal mice (1 day old) and in adult mice (10 week old) (C57BL/6). The following AAV serotypes were tested: AAV1, AAV2, AAV5, AAV6, AAV6.2, AAV7, AAV8, AAV9, rh.10 (also referred to herein as AAVrh.10), rh.39, rh.43, CSp3.

15 [0082] The recombinant AAV vectors expressed an enhanced GFP reporter gene under the CMV-enhanced chicken β -actin hybrid promoter and were produced by transient transfection in 293 cells. The neonatal day 1 pups were anesthetized with isoflurane. Then 100 μ L of rAAV vectors (4×10^{11} GC per mouse) was injected to the pups via superficial temporal vein under a dissection microscope. In adult mice, rAAV was administered by tail vein injection (two different doses were evaluated 4×10^{11} GC per mouse or 4×10^{12} GC per mouse). Twenty-one days post injection, the treated animals were anesthetized and transcardially perfused with cold PBS and 4% (v/v) paraformaldehyde. Brains were extracted, immersed in 20% sucrose, and embedded in Tissue-Tek OCT. 40 μ m thick sections were cut and stained in 12-well plate with primary antibodies, e.g., anti-NeuN, anti-EGFP and anti-GFAP, overnight at 4°C, then with secondary antibodies for 2 h at room temperature. Control mice received PBS injections.

20 [0083] In the neonatal study, the distribution of EGFP (+) cells throughout the brain at 3 wks post-infusion was observed. Large numbers of EGFP (+) cells with variable intensities were visible in different regions of the brains from the animals treated with 10 out of 12 vectors. In many instances the choroid plexus showed very strong EGFP expression, and transduced brain parenchyma cells appeared predominantly in periventricular regions. This indicates that a fraction of IV delivered vectors may enter the CNS via the choroid plexus-blood interface. In adults, substantial staining of brain vasculature was observed. Overall targeting efficiencies by AAVs to different regions of the brain was ranked as hypothalamus > medulla > cortex > hippocampus > cerebellum > thalamus. EGFP expression was not detected at high levels in neonatal mice that were administered rAAV2 or rAAV5 harboring the EGFP reporter gene by injection of 4×10^{11} GC per mouse in the superficial temporal vein. (See Table 1 and Figures 1 and 2 for summary data).

25 [0084] Tissue sections were also immunofluorescently stained with anti-EGFP and -cell type specific marker antibodies to classify EGFP (+) cell types in the CNS. Detection sensitivity for EGFP (+) cells, particularly neurons and oligodendrocytes, was improved dramatically. Although different vectors transduced neurons at variable efficiencies, all 10 vectors (including AAV9) exhibited stronger tropisms to non-neuronal cells, especially astrocytes. One vector (AAV7) targeted oligodendrocytes more efficiently than the other 9 vectors. Several rAAVs transduced both neurons and/or astrocytes at higher efficiencies as compared to rAAV9 (AAVrh.10, rh.34, and rh.43). Extensive astrocyte transduction was observed in hypothalamus and medulla. Injection of certain vectors resulted in substantial neuron transduction in different regions of the brain, including neocortex, hippocampus, and hypothalamus. Some vectors appeared to transduce Purkinje cells in cerebella cortex (e.g., CSp3), while others effectively transduced blood vessel in neocortex, thalamus and hypothalamus. In addition, choroid plexuses in 3rd ventricle, lateral ventricle and 4th ventricle showed strong EGFP expression. EGFP expression was also evaluated in different spinal cord regions of neonatal and adult mice (results for neonatal studies are shown in Figure 3).

30 [0085] Transduction of non-CNS tissues such as heart and skeletal muscle was observed (e.g., for AAV9, AAV8, and CSp3). In some cases, this may lead to some undesirable side effects. To address this issue, miRNA binding sites were incorporated into the 3' UTR of the transgene cassette and achieved highly specific and effective detargeting of AAV transduction from non-CNS tissues. To inhibit expression in liver, miRNA binding(s) for mR-122 were used. To inhibit expression in skeletal muscle and heart, miRNA binding(s) for mR-1 were used.

45

50

55

Table 1: AAV CNS TROPISMS

		AAV1	AAV2	AAV5	AAV6	AAV6.2	AAV7	AAV8	AAV9	rh.10	rh.39	rh.43	CSp3	
5	Adult	Cortex	+			+	+	++	++	+++	++	+	-	+
		Hippocampus	+			+	+	++	++	+++	++	+	-	+
		Thalamus	+			+	+	++	++	++++	+++	++	+	+
		Hypothalamus	+			++	+	+++	++	++	+++	+++	+	++
		Cerebellum	+			++	+	++	+++	+++	++++	+	+	+
		Brain Stem	+			++	+	++	++	++++	+++	++	-	+
		Cervical	+++			+	+	+++	+++	++++	+++	+++	-	+
		Thoracic	+++			+	+	+++	+++	++++	+++	++	-	+
		Lumbar	+++			+	+	+++	+++	++++++	+++	++	-	+
10	Neo-Natal	Cortex	++	+	-	++	+	+	+++	++	++	++	++	++
		Hippocampus	+	+	-	-	-	+++	++	+	+	++	++	+
		Thalamus	+	+	-	-	+	++	+	+	+	++	+	+
		Hypothalamus	++	-	-	+	+	+++++	+++++	+	+	+++++	+++	-
		Cerebellum	++	-	-	+	-	+	+	+	+	++	+	+
		Brain Stem	++	-	-	+	-	++	+	+	+	++++	+++	++
		Cervical	-	-	-	+	++	++	+++	++	++++	++++	+++	++
		Thoracic	+	-	-	+	++	+++	++	++	++	++++	+++	+
		Lumbar	+	-	-	+	++	++	+	+	++	+++	+++	+

Extent of Tissue Tropism (- no tropism; +++++ high tropism) Based on Data in Figures 1 and 2.

Example 2: Construction and evaluation of a recombinant AAVrh.10 vector to treat CD

[0086] Canavan disease (CD) is an inherited neurodegenerative disorder caused by mutations in the aspartoacylase gene (ASPA), leading to accumulation of N-acetyl-aspartic acid (NAA) in oligodendrocytes with resultant spongy degeneration of white matter in the brain. An initial clinical study on rAAV2-based ASPA gene therapy for CD achieved very limited success. It is believed, without wishing to be bound by theory, that an effective CD gene therapy will transduce oligodendrocytes throughout the CNS.

[0087] A rAAV vector is constructed that comprises a promoter operably linked with a region encoding ASPA protein (SEQ ID NO: 13 or 15) as a gene therapy vector for CD. The construct employs CAG (chicken β -actin promoter with CMV enhancer) to drive the expression of ASPA having a coding sequence as set forth in SEQ ID NO: 14 or 16. The rAAV vector is package into rAAV particles using the triple transfection method. To evaluate its effectiveness, rAAV-ASPA is examined in an ASAP knock-out mouse model of CD for its ability to eliminate or attenuate the CD-like phenotypic of homozygous ASPA knock-out mice (Matalon R et al. The Journal of Gene Medicine, Volume 2 Issue 3, Pages 165 - 175). Homozygous ASPA knock-out mice exhibit neurological impairment, macrocephaly, generalized white matter disease, deficient ASPA activity and high levels of NAA in urine. Magnetic resonance imaging (MRI) and spectroscopy (MRS) of the brain of the homozygous mice show white matter changes characteristic of Canavan disease and elevated NAA levels. Heterozygous ASPA knock-out mice, which have no overt phenotype at birth, serve as controls.

Example 3: Therapeutic Efficacy and Safety Evaluation of an AAV vector to treat CD

[0088] The mouse model of CD is a C57BL/6 derived ASPA gene KO strain. The homozygous KO animals present biochemical and neurological defects similar to those observed in CD patients. CD mice provide an animal model for evaluating gene therapy and other therapeutics for the treatment of CD. CD mice are used to study the efficacy and safety of the novel gene therapy strategies for the treatment of CD.

Experiment design

[0089] To examine therapeutic efficacy and safety, scAAV vectors (e.g., AAV7, AAV8, CSp3 and AAV9) carrying an optimized ASPA expression cassette are investigated in a preclinical gene therapy trial of CD. The vectors include miRNA binding site(s) to inhibit ASPA expression in non-CNS tissues. Both postnatal day-1 and 3-month-old adult animals are treated with each vector at two doses, 1 and 3×10^{14} GC/kg by intravenous administration. For the neonatal CD mice, two litters of animals receive each vector at each dose via temporal vein injections for necropsy of one litter each at 1- and 3-month time points. For the 3-month-old adult CD mice, 12 male animals are treated with each vector at each dose via tail vein injections. Six each of the treated animals are necropsied 1 and 3 months later. In further experiments, both postnatal day-1 and 3-month-old adult animals are treated with vectors at a dose in a range of 10^{11}

to 10^{12} GC/subject by direct intraventricular administration.

Functional and neurological measurements during the live phase of the study

5 [0090]

- 1). NAA metabolism. Urine samples are collected from the treated, untreated control, and wild type animals at days 14, 30, 45, 60, 75, and 90. The samples are analyzed by HPLC to determine the NAA levels.
- 2). NAA accumulation and NAA-induced water retention in brain. MRI/MRS-based neuroimaging studies are performed on the live animals in all study groups at 1, 2, and 3 months after the vector treatment to measure spectral peak integrals for creatine/phosphocreatine and NAA as well as abnormal hyperintense areas in the brain.
- 3). Liver function tests. Serum samples are collected from the animals in all study groups at days 14, 30, 60, and 90 to measure the levels of alanine transaminase (ALT) and aspartate aminotransferase (AST) as indicators of vector-related liver toxicity.
- 4). Neurological tests. Tremors, walking with splayed legs at a slow and shaky pace, and ataxia are among the prominent neurological features of the CD mice. At 1, 2, and 3 months after the gene therapy treatment, the animals in all study groups are subjected to a walking-pattern analysis by staining their feet with color ink and then recording their walking patterns as footprints on white paper. The animals also are tested and scored on a rotarod test for their ability to maintain balance.

20 *Enzymatic and histopathological analyses at the endpoints of the study*

[0091]

- 1). ASPA activities in the brain and non-CNS tissues. On-target and off-target expression of ASPA are analyzed by collecting brain, liver, heart and pancreatic tissues at necropsy to measure ASAP activities in the respective tissue homogenates.
- 2). Brain white matter and liver pathologies. To examine potential improvement in brain white-matter pathology and vector-related liver toxicity resulting from the gene therapy, brain and liver tissues are harvested and fixed, paraffin-embedded and sectioned, and stained with hematoxylin and eosin. Histopathological examination is performed by a pathologist.

Example 4: Delivery of therapeutic genes to the CNS cells by AAVrh.10

35 [0092] A screen of different AAV serotypes, was developed to identify candidates for a therapeutic gene transfer to the CNS. A recombinant AAV vector was constructed that expresses EGFP. The rAAV vector was packaged into four different AAVs: AAV1, 8, 9 and 10. Adult mice were injected with the AAVs into the CSF in the lumbar position. AAV1, 8 and 9 transduced cells only in the vicinity of the injection site at the lumbar region of the spinal cord following administration of $\sim 4.8 \times 10^{10}$ particles. Surprisingly, AAVrh.10 transduced cells in the gray matter along the entire spinal cord and brainstem following the same injection protocol and dosage as AAV1, AAV8 and AAV9 (Fig. 4A). Recently, AAV9 has been shown to cross the blood brain barrier (BBB) and transduce spinal cord cells after intravenous injection. A weak signal was observed in the cerebellum and strong signals in the brainstem and spinal cord. A weak signal (similar to the cerebellum) in the forebrain was also observed. Without wishing to be bound by theory, it is believed that CSF flow and diffusion allows the virus spread along the entire spinal cord, but that the ability of a virus to flow and diffuse depends on the structure of the viral capsid. The transduced cell types include neurons and oligodendrocytes. But the majority appears to be astrocytes (Fig. 4B), as indicated by overlap of EGFP with GFAP-positive cells. Substantial overlap with the microglia marker, Iba-1 was not observed. A number of motor neurons were transduced as indicated by overlap of EGFP expression and NeuN staining. It was surprising that among the astrocytes, only those situated in the gray matter were transduced and those that were situated in the white matter and beneath the pia matter were not transduced. This was striking because the virus is likely to be exposed to astrocytes in these areas since it was administered in the subarachnoid space.

Example 5: Construction of a recombinant AAVrh.10 vector to treat ALS

55 [0093] An recombinant AAV system was developed as a treatment for ALS. A rAAVrh. 10 vector was constructed that expresses a microRNA targeting SOD1 (Figure 5A). This microRNA was identified as miR-SOD1. The construct employed CAG (chicken β -actin promoter with CMV enhancer) to drive the expression of EGFP and miR-SOD1 that was located in an intron in the 3'-UTR.

[0094] The silencing potency of 9 miRNA constructs was evaluated. The constructs were transfected into HEK293 cells. After 48 hours, RNA was isolated and Northern blot was carried out to detect SOD1 mRNA (Figure 5B). MiR-SOD1#5 (SEQ ID NO: 26) silenced SOD1 expression most potently. Next, miR-SOD1#5 was packaged into AAVrh.10 (Figure 5D), which was used to infect HEK293 cells. Total cellular protein was extracted 43 hours after the infection and blotted to detect SOD1 (Figure 5C). Inhibition of expression of SOD1 at the protein level was observed.

5 *Example 6: Delivery of therapeutic genes to the CNS cells to treat ALS*

[0095] Large batches of AAVrh.10-miR-SOD1 and AAVrh.10-miR-Scr (scrambled miRNA) were produced using standard techniques. Self-complementary AAV (scAAV) was made because it mediates transduction with higher efficiency than conventional single stranded AAV [14]. A scAAVrh.10 was tested and found to express EGFP more rapidly (within 1 week) and stronger than a single stranded AAV.

[0096] AAVrh.10-miR-SOD1 was administered to one group of G93A mice (high SOD1 expressers) and AAVrh.10-miR-Scr to another group of G93A mice (n=15). The AAVrh.10 was injected intrathecally into the CSF in the lumbar area and injected intraventricularly into the forebrain in mice of 60 days of age ($\sim 4.8 \times 10^{10}$ particles in 8ul).

[0097] The animals were allowed to live their natural lifespan before succumbing to ALS. The lifespan was compared between the two groups. It was found that mice receiving the AAVrh.10-miR-SOD1 virus, which expresses a SOD1miR5 (SEQ ID NO: 26), lived on average 135 days (± 14 days), whereas mice receiving the AAVrh.10-miR-Scr, which expresses a scrambled miRNA (SEQ ID NO: 31), lived on average 122 days (± 6 days) (Figure 6B). Moreover, by examining the extent of EGFP expression in cervical, thoracic, and lumbar spinal cord tissue, a correlation in the levels of expression in these tissues, particularly with cervical tissue, and lifespan was observed in AAVrh.10-miR-SOD1 treated mice (Figure 7A), but not AAVrh.10-miR-Scr treated mice (Figure 7B). These results suggest that silencing mutant SOD1 expression in the cervical spinal cord is particularly beneficial in extending survival. A subset of the animals from each group were perfused with fixative, sectioned and stained for SOD1 in the spinal cord. SOD1 was detected using standard techniques [9]. SOD1 staining intensity in EGFP expressing cells was reduced compared with the non-EGFP cells that are transduced with AAVrh.10-miR-SOD1 (Figure 6A, showing knockdown of SOD1 expression in astrocytes). Reduction of expression of SOD1 was not observed in cells transduced with AAVrh.10-miR-Scr.

[0098] Tissues from another subset of animals in both groups were dissected to estimate transduction levels. The levels of transduction were estimated by determining the viral genome content using PCR on DNA samples obtained from different CNS and non-CNS regions. Measurements in non-CNS tissues (e.g. liver) provided an indication of whether virus had leaked to the periphery. Northern and Western analysis was performed to measure the SOD1 levels in the spinal cord. The antibody used for SOD1 detection was polyclonal, sheep anti-human SOD1, by Biodesign International, catalog#K90077C.

35 *Example 7: Combined Intrathecal/Intraventricular Administration Protocol*

[0099] AAV viruses were injected into mouse CSF by lumbar intrathecal injection and/or brain third ventricle injection. Injection into mice lumbar subarachnoid space was carried out using a method modified from Wu et al. [22]. A thin catheter (about 5cm) was made by stretching PE10 tube to the inner diameter 0.12 mm. The stretched section was cut to 1.7 to 1.9 mm, and two beads (1 mm apart) were made between the thin and the thick sections by heating and pressing the tube. To implant the catheter, the mouse was anesthetized by injection of Avertin (1.2% 2,2,2-tribromoethanol in 2% tert-amyl alcohol and PBS) intraperitoneally at 0.23 ml/10 g of body weight [23]. The catheter was then implanted between the L5 and L6 vertebra. The catheter was stitched to the surface muscle at the beaded area. Viruses of dose from 4.80E+10 Genome Copy (for virus screening, in 6ul) to 2.40E+10 Genome copy (for therapy, in 8ul) were injected via the catheter by a Hamilton syringe at a speed of 2 ul/minute. The catheter was sealed at the end by heat and left in place for one day. Wound was closed by clips. Injection into brain third ventricle was carried out using a Stoelting Stereotaxic Instrument and micro-injection pumps from World Precision Instruments following standard stereotaxic procedure. Same doses of virus were injected into the third ventricle at a rate of 1 ul/minutes.

[0100] Estimated doses for human and monkeys and comparison with IV injection are shown below. The two types of monkey are similar in size.

Table 2 - Estimated Doses for Human and Monkeys

Species	Avg CSF ml	Estimated CSF production rate ml/hour	Estimate dose (GC)	particles/g of body weight
mouse	0.035	0.018	2.40E+10	1.2E+09

(continued)

Species	Avg CSF ml	Estimated CSF production rate ml/hour	Estimate dose (GC)	particles/g of body weight
human	140	21	9.6E+13	1.3E+09
Macaca mulatta (rhesus monkeys)	14	2.5	9.6E+12	1.7E+09
Macaca fascicularis (cynomolgus macaque)	See, Foust KD, et al., <i>Nature Biotechnology</i> , Volume 28, Number 3, March 2010, 271-274		1.00E+14	2.20E+11

References for Background and Examples 1-7**[0101]**

1. Daya S, Berns KI: Gene Therapy Using Adeno-Associated Virus Vectors. *Clin Microbiol Rev* 2008, 21:583-593.
2. Eberling JL, Jagust WJ, Christine CW, Starr P, Larson P, Bankiewicz KS, Aminoff MJ: Results from a phase I safety trial of hAAADC gene therapy for Parkinson disease. *Neurology* 2008, 70:1980-1983.
3. Feigin A, Kaplitt MG, Tang C, Lin T, Mattis P, Dhawan V, During MJ, Eidelberg D: Modulation of metabolic brain networks after subthalamic gene therapy for Parkinson's disease. *Proceedings of the National Academy of Sciences* 2007, 104:19559-19564.
4. Cideciyan AV, Hauswirth WW, Aleman TS, Kaushal S, Schwartz SB, Boye SL, Windsor EAM, Conlon TJ, Sumaroka A, Pang J-j, et al: Human RPE65 Gene Therapy for Leber Congenital Amaurosis: Persistence of Early Visual Improvements and Safety at 1 Year. *Hum Gen Ther*, 0.
5. Raoul C, Abbas-Terki T, Bensadoun J-C, Guillot S, Haase G, Szulc J, Henderson CE, Aebischer P: Lentiviral-mediated silencing of SOD1 through RNA interference retards disease onset and progression in a mouse model of ALS. *Nat Med* 2005, 11:423-428.
6. Kaspar BK, Llado J, Sherkat N, Rothstein JD, Gage FH: Retrograde Viral Delivery of IGF-1 Prolongs Survival in a Mouse ALS Model. *Science* 2003, 301:839-842.
7. Azzouz M, Ralph GS, Storkebaum E, Walmsley LE, Mitrophanous KA, Kingsman SM, Carmeliet P, Mazarakis ND: VEGF delivery with retrogradely transported lentivector prolongs survival in a mouse ALS model. *Nature* 2004, 429:413-417.
8. Ralph GS, Radcliffe PA, Day DM, Carthy JM, Leroux MA, Lee DCP, Wong L-F, Bilsland LG, Greensmith L, Kingsman SM, et al: Silencing mutant SOD1 using RNAi protects against neurodegeneration and extends survival in an ALS model. *Nat Med* 2005, 11:429-433.
9. Wu R, Wang H, Xia X, Zhou H, Liu C, Castro M, Xu Z: Nerve Injection of Viral Vectors Efficiently Transfers Transgenes into Motor Neurons and Delivers RNAi Therapy Against ALS. *Antioxidants & Redox Signaling* 2009, 11:1523-1534.
10. Foust KD, Nurre E, Montgomery CL, Hernandez A, Chan CM, Kaspar BK: Intravascular AAV9 preferentially targets neonatal neurons and adult astrocytes. *Nat Biotechnol* 2009, 27:59-65.
11. Duque S, Joussemet B, Riviere C, Marais T, Dubreil L, Douar AM, Fyfe J, Moullier P, Colle MA, Barkats M: Intravenous administration of self-complementary AAV9 enables transgene delivery to adult motor neurons. *Mol Ther* 2009, 17:1187-1196.
12. Vandenberghe LH, Wilson JM, Gao G: Tailoring the AAV vector capsid for gene therapy. *Gene Ther* 2009, 16:311-319.
13. McBride JL, Boudreau RL, Harper SQ, Staber PD, Monteys AM, Martins I, Gilmore BL, Burstein H, Peluso RW, Polisky B, et al: Artificial miRNAs mitigate shRNA-mediated toxicity in the brain: Implications for the therapeutic development of RNAi. *Proceedings of the National Academy of Sciences* 2008, 105:5868-5873.
14. McCarty DM: Self-complementary AAV Vectors; Advances and Applications. *Mol Ther* 2008, 16:1648-1656.
15. Boilée S, Yamanaka K, Lobsiger CS, Copeland NG, Jenkins NA, Kassiotis G, Kollias G, Cleveland DW: Onset and progression in inherited ALS determined by motor neurons and microglia. *Science* 2006, 312:1389-1392.
16. Xia X, Zhou H, Huang Y, Xu Z: Allele-specific RNAi selectively silences mutant SOD1 and achieves significant therapeutic benefit in vivo. *Neurobiol Dis* 2006, 23:578-586.
17. Yamanaka K, Chun SJ, Boilée S, Fujimori-Tonou N, Yamashita H, Gutmann DH, Takahashi R, Misawa H, Cleveland DW: Astrocytes as determinants of disease progression in inherited amyotrophic lateral sclerosis. *Nat*

Neurosci 2008, 11:251253.

18. Di Giorgio FP, Carrasco MA, Siao MC, Maniatis T, Eggan K: Non-cell autonomous effect of glia on motor neurons in an embryonic stem cell-based ALS model. *Nat Neurosci* 2007, 10:608-614.

19. Nagai M, Re DB, Nagata T, Chalazonitis A, Jessell TM, Wichterle H, Przedborski S: Astrocytes expressing ALS-linked mutated SOD1 release factors selectively toxic to motor neurons. *Nat Neurosci* 2007, 10:615-622.

5 20. Wang Y, Ou Mao X, Xie L, Banwait S, Marti HH, Greenberg DA, Jin K: Vascular Endothelial Growth Factor Overexpression Delays Neurodegeneration and Prolongs Survival in Amyotrophic Lateral Sclerosis Mice. *J Neurosci* 2007, 27:304-307.

10 21. Storkebaum E, Lambrechts D, Dewerchin M, Moreno-Murciano M-P, Appelmans S, Oh H, Van Damme P, Rutten B, Man WY, De Mol M, et al: Treatment of motoneuron degeneration by intracerebroventricular delivery of VEGF in a rat model of ALS. *Nat Neurosci* 2005, 8:85-92.

22. Wu, W. P., Xu, X. J., and Hao, J. X. (2004) *J. Neurosci. Methods* 133, 65-69

23. Papaioannou, V. E., and Fox, J. G. (1993) *Lab. Anim. Sci.* 43, 189-192

15 *Example 8. MicroRNA-regulated, Systemically Delivered rAAV9*

Introduction to the Example

[0102] This example involves the use of tissue-specific, endogenous microRNAs (miRNAs) to repress rAAV expression outside the CNS, by engineering perfectly complementary miRNA-binding sites into the rAAV9 genome. The example describes recombinant adeno-associated viruses (rAAVs) that can cross the blood-brain-barrier and achieve efficient and stable transvascular gene transfer to the central nervous system (CNS), while de-targeting certain other tissues (e.g., liver, heart, skeletal muscle and other tissues). The approaches described in this example allowed simultaneous multi-tissue regulation and CNS-directed stable transgene expression without detectably perturbing the endogenous miRNA pathway. Regulation of rAAV expression by miRNA was primarily via site-specific cleavage of the transgene mRNA, generating specific 5' and 3' mRNA fragments.

[0103] Gene transfer mediated by recombinant adeno-associated virus (rAAV), as disclosed herein, is useful for treatment of a large number of neurological disorders. It has been found that rAAV vectors disclosed herein cross the blood-brain barrier and are specifically expressed in the CNS. Thus, the vectors may be used for intravascular delivery of rAAV for gene therapy of CNS diseases, including those that affect large areas of the brain and spinal cord.

[0104] This example describes the use of endogenous microRNAs (miRNAs) to suppress transgene expression outside the CNS. miRNAs are small, noncoding RNAs that regulate gene expression by post-transcriptional silencing. In general, miRNAs may silence genes by two mechanisms. When partially complementary to mRNA sequences, they typically reduce target mRNA stability and protein expression (e.g., by two- to fourfold or less), a mode of regulation thought to tune mRNA expression. In contrast, when miRNAs are nearly perfectly complementary to their mRNA targets, they typically bring about cleavage of the mRNA, triggering its wholesale destruction.

[0105] In particular, this example describes the use of miRNAs to detarget rAAV9 expression both separately and concurrently in the liver, heart, and skeletal muscle, the three tissues that are most efficiently targeted by intravenously delivered rAAV9. Silencing of transgene expression in liver, heart, and muscle exploited the natural expression of the abundant (\geq 60,000 copies/cell) miRNAs, miR-122, which is expressed in hepatocytes, and miR-1, a miRNA found in the heart and skeletal muscle of virtually all animals. miR-122-binding sites have been successfully used to prevent hepatotoxicity of a transgene from an adenovirus vector. Perfectly complementary sites for miR-1, miR-122, or both were engineered into the 3' untranslated region (UTR) of a nuclear-targeted, β -galactosidase (*nLacZ*) reporter transgene whose expression was driven by a cytomegalovirus-enhancer, chicken β -actin (CB) promoter. This example presents multiple independent results indicating that the miRNAs repress *nLacZ* expression by cleaving the transgene mRNA at exactly the same site as by all Argonaute-bound small RNAs in eukaryotic cells. When delivered systemically *in vivo*, the miRNA-detargeted rAAV9 vector successfully expressed the reporter transgene in the CNS, but not the liver or heart or skeletal muscle.

50 *Results*

miRNAs efficiently repress reporter gene expression in cultured cells

[0106] To evaluate a strategy for rAAV-mediated transduction, one or three tandem copies of a perfectly complementary binding site for miR-1 or miR-122 were introduced into the 3' UTR of *nLacZ* in a rAAV plasmid vector. The constructs were transfected into HuH7 cells, a human hepatoma cell line expressing ~16,000 copies of miR-122 per cell, and measured the number of *nLacZ*-positive cells. The number of *nLacZ*-expressing HuH7 cells for the one-site plasmid was about half that of the no site control; three sites reduced the number of *nLacZ*-expressing cells more than sevenfold

(Figure 12a).

[0107] Next, expression of the *nLacZ* constructs was analyzed in human embryonic kidney 293 cells, which naturally express low levels of both miR-122 and miR-1, when miR-1 or miR122 was introduced as a pri-miRNA from a second plasmid. 293 cells were transfected with the *nLacZ* reporter plasmids carrying 0, 1, or 3 miR-122 or miR-1-binding sites, together with a plasmid expressing either pri-miR-122 (Figure 12b) or pri-miR-1 (Figure 12c). To vary the concentration of the miRNA, either a low (1:3) or a high (1:10) molar ratio of the *nLacZ*-binding site plasmid to the miRNA expression plasmid was used. When miR-122 or miR-1 was introduced into the cells, *nLacZ* expression was repressed only when the *nLacZ* reporter mRNA contained the corresponding miRNA-binding sites; there was no reduction of nLacZ-positive cells when miR-1 was coexpressed with *nLacZ* containing miR-122-binding sites or when miR-122 was coexpressed with *nLacZ* containing miR-1-binding sites (Figure 12b,c).

Tissue-specific endogenous miRNAs regulate expression of rAAV9 delivered systemically in adult mice

[0108] To evaluate miRNA regulation of systemically delivered AAV9CBnLacZ vectors *in vivo*, AAV9CBnLacZ vectors carrying 0, 1, or 3 miRNA-binding sites perfectly complementary to either miR-122 or miR-1 were produced. The vectors were administered by tail vein injection to adult male C56BL/6 mice at a dose of 5×10^{13} genome copies per kg (GC/kg) body weight. Four weeks later, the liver and heart of the transduced animals were examined. LacZ staining revealed that the *nLacZ* transgene was silenced by the endogenous miRNAs in the cell type and organ in which they are predominantly expressed: the transgene was specifically silenced by miR-122 in the liver and by miR-1 in the heart (Figure 13a,b). While nLacZ positive cells were reduced in the livers of the animals treated with rAAV9CBnLacZ bearing one or three miR-122-binding sites, nLacZ expression levels in the hearts of the same animals were similar to those in the animals treated with AAV9CBnLacZ bearing no sites (Figure 13a). Similarly, nLacZ expression was not detected in the hearts of the animals that received AAV9CBnLacZ containing one or three miR-1-binding sites, but nLacZ expression in the livers of the same animals was not affected as compared to that in the control animal (Figure 13b). These data suggest that the greater the number of sites for a miRNA in rAAV, the lower the nLacZ expression in the tissue where the corresponding miRNA was expressed (Figure 13a,b).

[0109] Next, to evaluate whether transgene silencing could be achieved simultaneously in multiple tissues, different numbers of both miR-122- and miR-1-binding sites were inserted in the 3' UTR of the rAAV9CBnLacZ genome and examined for their expression in rAAV9 transduced mice. Histochemical staining of tissue sections showed that nLacZ expression was suppressed in both heart and liver for rAAV9CBnLac containing one or three copies each of the miR-1- and miR-122-binding sites, but nLacZ was readily detectable in pancreas, where expression of both miR-122 and miR-1 was low (Figure 13c). Quantitative, β -galactosidase assays of homogenized liver tissue similarly showed that nLacZ expression was significantly lower when the transgene contained the miRNA-binding sites (one miR-122-binding site: $7.8 \pm 7.4\%$, *P* value = 0.005; three miR-122-binding sites: $1.6 \pm 1.0\%$, *P* value = 0.005; one miR-1- plus one miR-122-binding site: $8.6 \pm 5.7\%$, *P* value = 0.005; three miR-1-plus three miR-122- binding sites: $3.1 \pm 1.2\%$, *P* value = 0.005; three miR-1-binding sites: $105.7 \pm 11.6\%$) (Figure 13d).

miRNA repression of rAAV expression does not perturb endogenous miRNA pathways

[0110] Highly expressed transgenes bearing miRNA-complementary sites have been reported to promote degradation of the corresponding miRNA. The levels of miR- 122, miR-22, miR-26a, and *let-7* were determined in rAAV transduced liver. No difference in abundance of the four miRNAs was detected among the three study groups (Figure 14a). Moreover, data from high throughput sequencing analyses of small RNA from the livers of one animal each from the three study groups show no change in miRNA levels.

[0111] In order to determine whether the miRNA-binding sites in the transgene transcripts would deregulate the expression of the known endogenous target mRNAs of miR-122 or miR-1, the expression of cyclin G1, a miR-122 target in liver (Figure 14b,c) and calmodulin, a miR-1 target in heart (Figure 14d) were analyzed. No significant alteration in cyclin G1 or calmodulin expression was detected. miR-122 regulates cholesterol biosynthesis in the liver, and agents that block miR-122 function may produce readily detectable changes in serum cholesterol levels. No change in total cholesterol, high-density lipoprotein, or low-density lipoprotein levels was detected in mice 4 weeks after transduction with either control rAAV9 or rAAV9 expressing a transgene bearing miR-122-binding sites (Figure 14e). It was concluded that in this example miRNA-mediated detargeting of rAAV expression had no detectable effect on endogenous miRNA expression or function.

Endogenous miRNAs silence rAAV transduction by site-specific cleavage of transgene mRNA

[0112] To determine how miRNAs suppress expression of transgenes delivered by rAAV *in vivo*, the transgene mRNA in liver was characterized by conventional PCR (Figure 15b), quantitative reverse transcription PCR (qRT-PCR) (Figure

15c), Northern hybridization (Figure 15d,e), and rapid amplification of 5' complimentary DNA (cDNA) ends (5' RACE; Figure 15f). When primers were used that amplify the region between the 3' end of *nLacZ* (A⁺F primer) and the 5' end of the poly(A) signal (A⁺R primer), an amplicon that spans the miRNA-biding sites, a 145 basepair (bp) product was detected after 26 cycles of amplification for the samples that received control rAAV. An additional six cycles of amplification 5 were required to detect a weak 220bp band for the samples transduced by rAAV containing three miR-122-binding sites. These data are consistent with low levels of intact *nLacZ* mRNA (Figure 15a,b).

[0113] To quantitatively assess the extent of the miRNA-directed repression of the transgene transcripts, qRT-PCR was performed using either oligo(dT) or random hexamer primers for reverse-transcription and PCR primer pairs that span either a 5' (nLacZ5'F/5'R), or 3' (nLacZ 3'F13'R) region of the *nLacZ* coding sequence (Figure 15a). The levels of 10 *nLacZ* mRNA were examined with intact 5' and 3' ends in total liver RNA extracted from four animals that received the control rAAV9CB*nLacZ* and four that received rAAV9CB*nLacZ* containing three miR-122-binding sites in the 3' UTR. Reductions ranging from 3 ± 1 (random hexamer) to 7 ± 1 (oligo[dT])-fold in *nLacZ* mRNA with an intact 3' end were 15 observed in the animals that had received rAAV9 containing miR-122-binding sites, relative to the control. In contrast, little or no decrease in *nLacZ* mRNA with an intact 5' end were detected for the same samples using the 5'F/5'R primer pair (Figure 15c). These results indicate that the primary mode of turnover of the mRNA that has been cleaved by a miRNA was 3'-to-5' exonucleolytic degradation.

[0114] To further characterize the fate of the transgene mRNA targeted by miR-1 or miR-122, Northern blot analyses 20 was performed. A transgene probe binding to the 5' end of *nLacZ* mRNA detected a -3.4 kb RNA in an animal injected with control rAAV9C*b**nLacZ*, the expected size of the of the full-length *nLacZ* transcript; a slightly larger band was detected in the liver sample from a mouse treated with rAAV9CB*nLacZ* bearing three miR1-binding sites (Figure 15a,d). In contrast to the single transcript detected for the rAAV9 expressing *nLacZ* bearing three miR-1- binding sites, two 25 RNAs of different sizes were detected for the rAAV expressing *nLacZ* bearing three miR-122 sites (Figure 15d).

[0115] The lengths of these transcripts indicate that the longer transcript likely represents the full-length mRNA, whereas 25 the shorter, more abundant transcript corresponds to 5' fragments of *nLacZ* RNA cleaved by miR-122 at the corresponding miR-122-binding sites in the 3' UTR (Figure 15d).

[0116] To confirm this observation, the Northern analysis was repeated using an RNA probe spanning a portion of 3' 30 UTR of the transgene mRNA. In addition to detecting full-length *nLacZ* transcripts in the samples transduced by rAAV9 lacking miRNA-binding sites, two closely migrating species smaller than the 281 nucleotide RNA marker were detected. The size of these fragments was consistent with miRNA-directed 3' cleavage products of the *nLacZ* mRNA (Figure 15e). These two 3' cleavage products were also detected by gel electrophoresis of the product from the 5' RACE experiment 35 described below (Figure 15f).

[0117] To determine whether such target cleavage occurs *in vivo* when the *nLacZ* transcript contained miR-1 or miR-122-binding sites, rapid amplification of 5' cDNA ends (5' RACE) was performed. Figure 16 presents the sequences of 40 21 clones recovered using 5' RACE from liver RNA (Figure 16a) and 22 clones isolated from heart RNA (Figure 16b) from the animals injected with rAAV9 in which the *nLacZ* 3' UTR contained three miR-1 and three miR-122-binding sites. In liver, the sequence signatures for miR-122- directed cleavage of the transgene mRNA were detected at each miR-122-binding site: 5% for the first binding site, 48% for the second binding site, and 43% for the third binding site. All 5' ends mapped to the phosphate that lies between the target nucleotides that pair with positions 10 and 11 of the sequence perfectly complementary to miR-122, the precise site cleaved by small RNAs bound to Argonaute proteins in all eukaryotes 45 (Figure 17a). Similar results were obtained in the heart for the rniR-1 sites (Figure 17b).

[0118] Table 3 presents an expanded 5' RACE analysis for additional vector groups. It was noted that none of the 5' RACE products sequenced corresponded to miR-1-directed site-specific cleavage in liver or miR-122-directed site-specific cleavage in heart (Table 3). Although no cleavage was detected within miR-1-binding sites in the liver, some clones from heart were cleaved within the miR-122-binding sites, but not at the hallmark position for miRNA-directed 45 cleavage.

Intravascularly delivered rAAV9 can be efficiently controlled by endogenous miRNAs

[0119] MiRNA-1 and miRNA-122-binding sites were added into the scAAV9CB enhanced GFP (EGFP) vector genome 50 and injected 10-week-old C57BL/6 male mice with 2 x 10¹⁴ GC/kg. After 3 weeks, 40 µm sections of brain and spinal cord and 8 µm sections of liver, heart, and skeletal muscle were prepared and examined for EGFP protein expression. It was found that intravenously delivered scAAV9CBEGFP efficiently transduced the CNS; EGFP was readily detectable 55 in the thalamus region of the brain and the cervical region of the spinal cord, but also in non-CNS tissues such as liver, heart, and muscle (Figure 17a). In contrast, transgene expression in those non-CNS tissues was reduced when miR-1 and miR122-binding sites were included in the transgene; EGFP expression was unaltered in the CNS, where miR-1 and miR-122 were not present (Figure 17a). Quantitative RT-PCR was used to measure the differential expression of the miRNA-repressed EGFP transgene in brain (41.2 ± 7.7%), liver (3.0 ± 0.5%), heart (0.4 ± 0.1%), and muscle (1.3 ± 0.4%), relative to the EGFP transgene lacking miRNA-binding sites (Figure 17b). To eliminate changes associated 55

with transduction efficiency between experiments, the data were normalized to the number of vector genomes detected in the experimental and control samples. Similar to the microscopic analyses of native EGFP expression, the qRT-PCR data show that the presence of miR-122- or miR-1-binding sites reduced transgene expression in liver (20-fold), heart (100-fold), and muscle (50-fold), but did not detectably alter transgene expression in brain.

5

Discussion Of Results

[0120] This example shows that rAAV9 can be engineered so that endogenous miRNAs repress transgene expression outside the CNS. The results indicate that such engineered rAAV9s may be used in therapies for the degenerating neurons associated with Parkinson's disease, Alzheimer's disease and amyotrophic lateral sclerosis, by expressing neurotrophic growth factors such as insulin-like growth factor, brain-derived neurotrophic factor or glial-derived neurotrophic factor in the transduced astrocytes. This approaches eliminates or lessens non-CNS expression derived from the peripheral tissues transduced by systemically delivered rAAV9.

[0121] Achieving transgene expression in primarily only the target tissues is a consideration for the clinical development of safe CNS gene delivery. The results in this example indicate that endogenous miRNAs can be harnessed to restrict the tissue- and cell-type specificity of rAAV expression, as was initially shown for lentiviral vectors. The data demonstrate that endogenous miRNAs can effectively repress transgene expression from rAAV. In both heart and liver, the miRNAs repressed transgene expression by directing endonucleolytic cleavage of the transgene mRNA (Figure 18). MiRNA regulation of rAAV expression did not perturb the expression or function of the corresponding endogenous miRNA, allowing transgene expression to be restricted to the CNS in mice. The example indicates that a strategy that combines multiple binding sites for miRNAs expressed in the periphery but not the CNS is useful for the development of safer, CNS-specific gene therapy vectors.

Materials And Methods

25

[0122] *Vector design, construction, and production.* Perfectly complementary miRNA-binding sites were designed based on the annotated miR-1 and miR-122 sequences in miRBase and inserted into the *BstBI* restriction site in the 3' UTR of the nLacZ expression cassette of the ubiquitously expressed pAAVCB nuclear-targeted β -galactosidase (nLacZ) plasmid using synthetic oligonucleotides (Figure 15a and Table 3). This vector uses a hybrid cytomegalovirus enhancer/CB promoter cassette that is active in most cells and tissues. To express miR-122 and miR-1, pri-miR-122 and pri-miR-1 fragments were amplified by PCR from C57/B6 mouse genomic DNA (Table 4) and inserted into the *XbaI* restriction site 3' to a firefly luciferase cDNA in the pAAVCBELuc plasmid. The identity of each pri-miRNA was verified by sequencing. AAV9 vectors used in this study were generated, purified, and tittered.

[0123] *Cell culture and transfection.* HEK-293 and HuH7 cells were cultured in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum and 100 mg/l of penicillin-streptomycin (Hyclone, South Logan, UT). Cells were maintained in a humidified incubator at 37°C and 5% CO₂. Plasmids were transiently transfected using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions.

[0124] *Mouse studies.* Male C57BL/6 mice (Charles River Laboratories, Wilmington, MA) were obtained and maintained. To monitor lipid profiles of the study animals, serum samples were collected 4 weeks after rAAV9 injection and analyzed for total cholesterol, high-density lipoprotein and low-density lipoprotein on a COBAS C 111 analyzer (Roche Diagnostics, Lewes, UK). To evaluate endogenous miRNA-mediated, CNS-restricted EGFP gene transfer, 10-week-old male C57BL/6 mice were injected intravenously (tail vein) with AAV9CBnLacZ-[miR-122-binding site (BS)₁]. AAV9CBnLacZ-(miR-122BS)₃. AAV9CBnLacZ-(miR-1BS)₁. AAV9CBnLacZ-(miR-1BS)₃. AAV9CBnLacZ-(miR-1BS)₁-(miR-122BS)₁, and AAV9CBnLacZ-(miR-1BS)₃-(miR-122BS)₃, respectively, at 5 x 10¹³ GC/kg body weight or scAAV9CBEGFP at 2 x 10¹⁴ GC/kg body weight). Animals receiving nLacZ vectors were necropsied 4 weeks later; 8 μ m cryosections of liver, heart, and pancreas tissues were prepared for X-gal-histochemical staining. Animals that received EGFP vectors were necropsied 3 weeks later and fixed by transcardial perfusion with 4% (wt/vol) paraformaldehyde. Brain, spinal cord, liver, heart, and muscle were harvested for cryosectioning. Brain and cervical spinal cord tissue were stained as floating sections in a 12-well plate using rabbit anti-EGFP antibody (Invitrogen) diluted 1:500, followed by goat anti-rabbit secondary antibody (Invitrogen) diluted 1:400. Outside the CNS, EGFP expression was detected directly by fluorescence. EGFP and antibody fluorescence was recorded using a Nikon TE-2000S inverted microscope at x10 magnification and an exposure time of 3 seconds for liver, heart, and muscle, and 5 seconds for thalamus (brain) and cervical spinal cord.

[0125] *Vector genome quantification by qPCR.* Genome DNA was extracted from the selected tissues using QIAamp DNA Mini Kit (Qiagen, West Sussex, UK), according to the manufacturer's instructions. Quantitative PCR were carried out in triplicate using Ring DNA and 0.3 μ mol/l EGFP-specific primers (EGFP-F and EGFP-R) using GoTaq qPCR master mix (Promega, Madison, WI) in a StepOne Plus real-time PCR instrument (Applied Biosystems, Foster City, CA).

[0126] *qRT-PCR analysis.* RNA was extracted using Trizol (Invitrogen), according to the manufacturer's instructions.

Total RNA (0.5-1.0 µg) was primed with random hexamers or oligo(dT) and reverse-transcribed with MultiScribe Reverse Transcriptase (Applied Biosystems). Quantitative PCR were performed in triplicate with 0.3 µmol/l gene-specific primer pairs (nLacZ5'F/5'R, nLacZ 3'F/3'R, cyclinG1F/R and EGFP-F/EGFP-R) using the GoTaq qPCR master mix in a StepOne Plus Real-time PCR device. The specificity of qRT-PCR products derived from the 5' and 3' ends of *nLacZ* mRNA was confirmed by gel electrophoresis.

[0127] *Northern blot analysis.* Total RNA was extracted from mouse liver and analyzed by Northern hybridization. To detect *nLacZ* mRNA, a 618 bp fragment of *nLacZ* cDNA was isolated by Ncol and Pcil digestion of pAAVCBnLacZ and labeled with α -³²P dCTP by random priming (Takara, Shiga, Japan). To detect 3' fragments of the cleaved *nLacZ* mRNA, an 111 bp fragment of the poly(A) sequence in the vector genome was cloned into pCR4-TOPO (Invitrogen) for preparation of antisense RNA probe labeled with α -³²P CTP during *in vitro* transcription using the Riboprobe System T7 kit (Promega). To detect miR-122, miR-26a, miR-22, and let-7 or U6 in total liver RNA, small RNAs were resolved by denaturing 15% polyacrylamide gels, transferred to Hybond N+ membrane (Amersham BioSciences, Pittsburgh, PA), and crosslinked with 254 nm light (Stratagene, La Jolla, CA). Synthetic oligonucleotides, 5' end-labeled with γ -³²P ATP using T4 polynucleotide kinase (New England Biolabs, Beverly, MA), were used as DNA probes (Table 4) and hybridized in Church buffer (0.5 mol/l NaHPO₄, pH 7.2, 1 mmol/l EDTA, 7% (w/v) sodium dodecyl sulphate) at 37°C. Membranes were washed using 1 x SSC (150 mM sodium chloride, 15 mM sodium citrate), 0.1% sodium dodecyl sulphate buffer, and then visualized using an FLA-5100 Imager (Fujifilm, Tokyo, Japan).

[0128] *Western blot analysis.* Proteins were extracted with radioimmunoprecipitation assay buffer [25 mmol/l Tris-HCl, pH 7.6, 150 mmol/l NaCl, 1% (vol/vol) NP-40, 1% (wt/vol) sodium deoxycholate, 0.1% (w/v) sodium dodecyl sulphate] containing a protease inhibitor mixture (Boston BP, Boston, MA). Protein concentration was determined using the Bradford method (Bio-Rad, Melville, NY). Protein samples, 50 µg each, were loaded onto 12% polyacrylamide gels, electrophoresed, and transferred to nitrocellulose membrane (Amersham BioSciences). Briefly, membranes were blocked with blocking buffer (LI-COR Biosciences, Lincoln, NE) at room temperature for 2 hours, followed by incubation with either anti-GAPDH (Millipore, Billerica, MA), anti-cyclin G1 (Santa Cruz Biotechnology, Santa Cruz, CA) or anti-calmodulin (Millipore) for 2 hours at room temperature. After three washes with PBS containing 0.1% (vol/vol) Tween-20, membranes were incubated with secondary antibodies conjugated to LI-COR IRDye for 1 hour at room temperature, and then antibodies detected using the Odyssey Imager (LI-COR).

[0129] β -*Galactosidase assay.* Proteins were extracted with radioimmunoprecipitation assay buffer and quantified as described above. Fifty micrograms of protein was used for each β -galactosidase assay using the Galacto-Star System (Applied Biosystems), according to the manufacturer's instructions.

[0130] *5'RACE.* 5' RACE was performed as described. The 5' RACE Outer Primer and the *nLacZ* gene-specific primer bGHpolyAR (Table 4) were used for the first round of nested PCR. The 5' RACE Inner Primer and the *nLacZ* gene-specific primer nLacZpolyR, which is located near the stop codon of *nLacZ* cDNA, were used for the second round of nested PCR (Table 4). PCR products were TOPO-cloned into pCR-4.0 (Invitrogen) and sequenced.

[0131] *Statistical analysis.* All results are reported as mean \pm SD and compared between groups using the two-tailed Student's t-test.

Table 3 Summary of microRNA-guided transgene mRNA cleavage in mouse liver and heart

		Cleavage site						
miR BS cleavage	Position	Between 10 and 11 nt		Between 17 and 18 nt		Between 18 and 19 nt		Random site
Liver 1 Copy of miR-122 BS (21 clones)	1	17/21	81%	ND		ND		19%
3 Copies of miR-122 BS (11 clones)	1	ND	100%	ND		ND		0%
	2	4/11						
	3	7/11						
3 Copies each of miR-1 and miR 122 miR 13x BS	1	ND	ND	ND		ND		0%
BS in a single vector (21 clones)	2	ND						
	3	ND						

(continued)

		Cleavage site					
5	miR BS cleavage	Position	Between 10 and 11 nt	Between 17 and 18 nt	Between 18 and 19 nt	Random site	
10	miR-122 3x BS	1	1/21	95%	ND	ND	5%
		2	10/21				
		3	9/21				
15	Heart 1 Copy of miR-IBS (12 clones)	1	12/12	100%	ND	ND	0%
	3 Copies of miR 1BS (21 clones)	1	ND	80%	4/21	20%	ND
		2	16/21		ND		
20		3	1/21		ND		
	3 Copies each of miR 1 and miR 122 miR-122 3x BS	1	ND	ND	ND	1/22	14%
		2	ND			1/22	
25	BS in a single vector (22 clones)	2	ND			ND	
		3	ND				
	miR 1 3x BS	1	1/22	73%	ND	9%	ND
30		2	7/22		1/22		
		3	8/22		1/22		

Table 4. Oligonucleotide primers and probes used in Example 8.

Oligo nucleotides	Sequence	SEQ ID NO
(miR-1) ₁ sense	[PHOS]CGAAATACATACTTCTTACATTCC ATT	SEQ ID NO: 32
(miR-1) ₁ anti-sense	[PHOS]CGAATGGAATGTAAAGAAGTATGT ATTT	SEQ ID NO: 33
(miR-122) ₁ sense	[PHOS]CGAAACAAACACCATTGTCACACT CCATT	SEQ ID NO: 34
(miR-122) ₁ anti-sense	[PHOS]CGAATGGAGTGTGACAATGGTGT TGT	SEQ ID NO: 35
(miR-1) ₃ sense	[PHOS]CGAAATACATACTTCTTACATTCC AATACATACTTCTTACATTCCAACATA CTTCTTACATTCCATT	SEQ ID NO: 36
(miR-1) ₃ anti-sense	[PHOS]CGAATGGAATGTAAAGAAGTATGT ATTGGAATGTAAAGAAGTATGTATTGGAA TGTAAAGAAGTATGTATT	SEQ ID NO: 37
(miR-122) ₃ sense	[PHOS]CGAAACAAACACCATTGTCACACT CCAACAAACACCATTGTCACACTCCAACA AACACCATTGTCACACTCCATT	SEQ ID NO: 38

(continued)

Oligo nucleotides	Sequence	SEQ ID NO
5 (miR-122) ₃ anti-sense	[PHOS]CGAATGGAGTGTGACAATGGTGT TGTGGAGTGTGACAATGGTGTGGTGG AGTGTGACAATGGTGTGGT	SEQ ID NO: 39
10 (miR-1) ₁ -(miR-122) ₁ sense	[PHOS]CGAAATACATACTTCTTACATTCC AACAAACACCATTGTCACACTCCATT	SEQ ID NO: 40
15 (miR-1) ₁ -(miR-122) ₁ anti-sense	[PHOS]CGAATGGAGTGTGACAATGGTGT TGTGGAAATGAAAGAAGTATGTATT	SEQ ID NO: 41
20 Synthesized (miR-1) ₃ - (miR-122) ₃ fragment	TTCGAACTCGAGATACATACTTCTTACAT TCCAATACATACTTCTTACATTCCAATAC ATACTTCTTACATTCCACCATGGACTAGT ACAAACACCATTGTCACACTCCAACAAAC ACCATTGTCAACTCCAACAAACACCATT GTCACACTCCAGCGGCCGTCGAA	SEQ ID NO: 42
25 Pri-miR-122F	ATCGGGCCCGACTGCAGTTCAGCGTTG	SEQ ID NO: 43
Pri-miR-122R	CGCGGGCCCGACTTACATTACACACAAT	SEQ ID NO: 44
Pri-miR-1F	CGCGGGCCCGACTGATGTGTGAGAGAGAC	SEQ ID NO: 45
Pri-miR-1R	CGCGGGCCCGACTTCGGCCTCCGAGGC	SEQ ID NO: 46
30 nLacZS¢F(5¢F)	TGAAGCTGAAGCCTGTGATG	SEQ ID NO: 47
nLacZ 5¢R(5¢R)	GAGCACCTGACAGCATTGAA	SEQ ID NO: 48
nLacZ3¢F(3¢F)	CTCAGCAACAGCTCATGGAA	SEQ ID NO: 49
nLacZ3¢R(3¢R)	TTACTTCTGGCACCAACACCA	SEQ ID NO: 50
nLacZpolyF(A ⁺ F)	TGGTGTGGTGCCAGAAGTAA	SEQ ID NO: 51
nLacZpolyR(A ⁺ R)	CAACAGATGGCTGGCAACTA	SEQ ID NO: 52
35 bGHpolyAR(bGH ⁺ AR)	TGGGAGTGGCACCTTCCA	SEQ ID NO: 53
EGFP-F	CGACCACTACCAGCAGAACAA	SEQ ID NO: 54
EGFP-R	CTTGTACAGCTCGTCCATGC	SEQ ID NO: 55
40 CyclinG1F	AATGGCCTCAGAATGACTGC	SEQ ID NO: 56
CyclinG1R	AGTCGCTTCACAGCCAAAT	SEQ ID NO: 57
MM-ActinF	ATGCCAACACAGTGTCTGG	SEQ ID NO: 58
MM-ActinR	TGCTTGCTGATCCACATCTGCT	SEQ ID NO: 59
45 miR-122 probe	TGGAGTGTGACAATGGTGTGG	SEQ ID NO: 60
Let-7 probe	AACTATACAACCTACTACCTCA	SEQ ID NO: 61
miR-26a probe	AGCCTATCCTGGATTACTTGAA	SEQ ID NO: 62
50 miR-22 Probe	ACAGTTCTCAACTGGCAGCT	SEQ ID NO: 63
U6 probe	CTCTGTATCGTCCAATTTAGTATA	SEQ ID NO: 64

Example 9: Intravenous injection of rAAVs mediated widespread transduction in neonatal mouse CNS

55
Introduction to the Example

[0132] This example describes an analysis of nine scAAV vectors for CNS gene transfer properties after systemic administration. This study involved identifying more effective vectors for the CNS gene transfer, In some aspects the

study examined serotypes or natural variants of rAAVs for enhanced-permeation of the BBB. In some cases, the study sought to identify rAAV vectors with improved delivery of enhanced green fluorescent protein (EGFP) to the CNS following facial vein injection on postnatal day 1 (P1). AAV9 was included in the study. Except for rAAV2 and rAAV5, all other 7 vectors crossed the BBB with varied transduction efficiency, among which rAAVrh.10, rAAVrh.39, rAAVrh.43, rAAV9 and rhAAV7 rank in the top 5, mediating robust EGFP expression in both neuronal and glial cells throughout the CNS in this study. The performance of rAAVrh. 10 was comparable to that of rAAV9 and in some case better. Several rAAVs efficiently transduce neurons, motor neurons, astrocytes and Purkinje cells; among them, rAAVrh.10 is at least as efficient as rAAV9 in many of the regions examined. Intravenously delivered rAAVs did not cause abnormal microgliosis in the CNS. The rAAVs that achieve stable widespread gene transfer in the CNS are useful as therapeutic vectors for neurological disorders affecting large regions of the CNS as well as convenient biological tools for neuroscience research.

Results

[0133] Twenty one days after vector administration in P1 mice, the CNS transduction profiles of the following recombinant AAV vectors encoding EGFP: rAAV1, rAAV2, rAAV5, rAAV6, rAAV6.2, rAAV7, rAAV9, rAAVrh.10, rAAVrh.39 and rhAAVrh.43 were compared. The vectors used in this study were comparable in purity and morphological integrity (Figure 19). As assessed by the scoring system described in the methods, rAAV9 was among the top performers; most other rAAVs tested (rAAV1, rAAV6, rAAV6.2, rAAV7, rAAVrh.10, rAAVrh.39 and rAAVrh.43) also gave rise to EGFP expression throughout the CNS (Table 2). The number of apparent EGFP positive cells (Table 5) among sub-anatomical structures was influenced by the particular vector used. For these seven rAAVs, and rAAV9 (total of eight rAAVs), that permeated the BBB and accomplished CNS transduction after i.v. delivery, EGFP positive cells were found in hypothalamus followed by medulla, striatum, hippocampus, cortex and cerebellum. In contrast, the transduction efficiency in olfactory bulb and thalamus was relatively low (Table 5). A quantitative assessment of EGFP gene transfer efficiency was made of each rAAV. 12 sub-anatomically and functionally important regions in the brain were selected for quantitative analysis of the mean EGFP intensity/pixel in each region for each rAAV by using Nikon NIS elements AR software V. 3.2 (Figure 19a) (see Methods). For the eight vectors that achieved CNS transduction after i.v. injection, the mean EGFP intensity/pixel was relatively low in cortex, habenular nucleus, cornu ammonis, dentate gyrus, thalamus, cerebellum and olfactory bulb, moderate in choroid plexus and caudate-putamen, but high in hypothalamus, medulla and amygdale (Figure 19a). The average EGFP intensities of all 12 regions for different rAAVs were compared in Figure 19b. AAVrh.10, AAVrh.39 and AAVrh.43 were noted for gene transduction efficiency in brain, followed by AAV7, AAV9, and AAV1 (Figures 19a and 19b). Those eight effective serotypes also mediated EGFP expression throughout the spinal cord, to different degrees. The same quantitative analysis was performed for each rAAV in the cervical, thoracic and lumbar sections of the spinal cord (Figure 19a); the average EGFP intensities of the three sections for different rAAVs were also compared (Figure 19b). AAV1, AAV9, AAVrh10, AAV.rh39 and AAV.rh43 displayed strong transduction in the spinal cord with the high EGFP intensity observed in the cervix, followed by thoracic and lumbar sections of the spinal cord (Figures 19a and 19b). For rAAV2 there were few EGFP-positive cells in hippocampus, cortex and hypothalamus. EGFP-positive cells were observed in the hypothalamus in AAV5-injected mice. A description of the observations made in different CNS structures is provided below. The subanatomic CNS structures may serve as a target for CNS gene therapy. In some cases, the subanatomic CNS structures are associated with pathological changes in one or more neurological disorders. In some cases, the subanatomic CNS structure have distinct transduction profiles for one or more rAAVs.

[0134] *Striatum.* Pathology of the striatum is associated with Huntington's disease, choreas, choreoathetosis, and dyskinesias. Addiction may involve plasticity at striatal synapses. Systemic injection of rAAV9 in neonatal mice transduces striatal tissue. In this study, a large number of cells with neuronal morphology in this region were also transduced by rAAVrh.10 (Figure 20), which was confirmed by co-staining with a neuronal marker as described below. Other vectors, including rAAVrh.39 and rAAV7, also mediated moderate transduction in striatum (Figure 20). In contrast, rAAV6, rAAV6.2, and rAAV1 resulted in relatively lower EGFP expression in this structure (Figure 20).

[0135] *Hippocampus.* The hippocampus is a region associated with long-term memory and spatial navigation, which is usually damaged by stress and pathogenesis of diseases such as epilepsy and Schizophrenia. Large numbers of EGFP-positive neurons were observed bilaterally in all regions of the hippocampus, namely dentate gyrus, hilus, CA1, CA2 and CA3 for the mice received intravenous rAAVrh.10, rAAV9, rAAV7, rAAVrh.39, and rAAVrh.43 (ranked by transduction efficiency in this structure, Table 5 and Figures 19 and 20). In addition to the neuronal transduction pattern, EGFP-positive cells had morphologic appearance of astrocytes (Figure 20). This was further confirmed by double staining with antibodies against EGFP and astrocytic marker as described below. For intravenously delivered rAAV1, rAAV6 and rAAV6.2 vectors there were small numbers of EGFP-positive cells in the hippocampus (Figure 20).

[0136] *Cortex.* Pathological changes in the cortex have been implicated in Alzheimer's and Parkinson's diseases. AAV7, AAV9, AAVrh.10, AAVrh.39 and AAVrh.43 vectors achieved moderate EGFP transduction in cortex (Table 5 and Figures 19 and 20). The morphology of transduced cells was consistent with both neurons and astrocytes as further

confirmed by cellular marker staining and confocal microscopic analysis described below. Prominent EGFP-positive cells were typically observed in the ventrolateral regions of the cortex, including posterior agranular insular cortex, piriform cortex, lateral entorhinal cortex, posterolateral cortical amygdaloid nucleus and posteromedial cortical amygdaloid nucleus (Figure 20). Strong EGFP signals spread from +1.5 to -3.3 mm in relation to the Bregma (0.0 mm). The cortical transduction efficiency of rAAVrh.10, rAAV9, rAAVrh.39 and rAAVrh.43 was comparable (Table 5 and Figures 19 and 20). AAV1, AAV6 and AAV6.2 vectors also transduced cells in the cortex (Figure 20).

[0137] Hypothalamus. A role for the hypothalamus is to secret neurohormones to control certain metabolic processes. The hypothalamus is also indicated in the etiology of diabetes. EGFP signal was observed in the hypothalamus for eight vectors. Intravenous administration of rAAVrh.10 resulted in the highest EGFP expression in the entire hypothalamus, followed by rAAVrh.39, rAAV7, rAAV6.2, rAAVrh.43, rAAV9, rAAV1 and rAAV6 (Figures 19 and 20 and Table 5). Interestingly most EGFP-positive cells in this structure have an astrocytic morphology which was ascertained by immunostaining for an astrocytic cell type specific marker as described below. The astrocytic EGFP signal tended to obscure direct examination of morphological details of other transduced cells. However, this was clarified by double immunofluorescent staining of tissue sections with antibodies for EGFP and neuronal cell markers as described below.

[0138] Cerebellum. The pathological lesions in cerebellum are often found in diseases such as cerebellar-cognitive affective syndrome, developmental coordination disorder, posterior fossa syndrome, linguistic deficits, aging, attention deficit hyperreactivity disorder, autism, dementia and schizophrenia. EGFP-positive cells and fibers were detected in cerebellum for most rAAV vectors (Table 5 and Figures 19 and 20). A large number of EGFP-expressing cells were found in the Purkinje and granule cell layers for rAAV7, rAAV9, rAAVrh.10, rAAVrh.39 and rAAVrh.43 (Figure 20). The transduction profile of rAAV1 vector indicated expression in cells in the granule cell layer, while rAAV6 and rAAV6.2 were localized in cells in the Purkinje cell layer (Figure 20).

[0139] Medulla. The medulla is a potential gene therapy target for treating chronic pain. Most rAAVs mediated moderate to robust EGFP expression in medulla with most green cells being present in the outer rim (Figure 20). Transduction efficiencies of these rAAV in this region are ranked in the following order: rAAVrh.39 = rAAVrh.43 > rAAV.rh10 > rAAV1 > rAAV9 > rAAV7 > rAAV6.2 > rAAV6 (Table 5 and Figure 19a). The morphology of most EGFP-transduced cells was consistent with the cells being astrocytes.

[0140] Spinal cord. The spinal cord is involved with motor neurons diseases. rAAVrh.10, rAAV9, rAAVrh.39 and rAAVrh.43 gave rise to very robust EGFP expression in cervical gray and white matter, while rAAV1, rAAV6.2 and rAAV7 showed moderate EGFP intensity (Table 5 and Figures 19 and 21). For rAAV1 the EGFP signal was observed in white matter. The transduction ability of all effective rAAVs decreased from cervical to lumbar spinal cord. EGFP-positive cells were visible in the latter region. Large populations of EGFP-positive cells with astrocytic morphology were observed throughout the spinal cord (Figure 21). In addition, rAAVrh.10, rAAV9, rAAVrh.39, rAAVrh.43 and rAAV7 also transduced cells with motor neuron morphology in the ventral regions of spinal cord (Figure 21). Ascending dorsal column fibers showed clear EGFP signal. In addition, dorsal root ganglia (DRG) displayed remarkable transduction with strong EGFP expression in DRG neurons (Figure 22 and Figure 26). The identities of rAAV transduced cell types in the spinal cord were characterized by co-immunofluorescence staining with antibodies against EGFP and cell type specific markers as described below.

IV administration of AAV vectors leads to transduction of different cell types in the CNS

[0141] To confirm the identity of transduced cells in different regions of the CNS, double immunofluorescent staining was performed with antibodies for EGFP and NeuN (generic neuronal marker), glial fibrillary acid protein (GFAP; astrocyte marker), calbindin-D28K (Purkinje cell marker), and choline acetyl transferase (ChAT; motor neuron marker) (Figure 23). The immunostaining results showed that a large number of NeuN positive cells expressed EGFP throughout the mouse brain, which indicated widespread neuronal transduction. The regions with high density of transduced neurons included striatum, hippocampus, cortex and hypothalamus. rAAVrh.10, rAAV9, rAAV7 and rAAVrh.39 vectors were efficient in mediating neuronal transduction, followed by AAV6.2, AAV1 and AAV6 (Figures 19 and 23). In addition, dopaminergic neurons in substantia nigra were transduced by AAV.rh10 (Figure 23). Transduced cells in the CNS included GFAP-positive astrocytes with small cell bodies and highly ramified processes (Figure 23). The calbindin-D28K immunostaining confirmed the identity of a number of transduced cells in the cerebellum as Purkinje cells, with EGFP expression in both cell body and their tree-like processes (Figure 23). The rAAVs proficient in transducing Purkinje cells include: rAAVrh.10, rAAV9, rAAVrh.39, rAAV7, rAAV6.2 and rAAVrh.43. rAAV1 and rAAV6 transduced a portion of Purkinje cells with relatively low EGFP intensity (Figure 19). Transduction of motor neurons was confirmed by the presence of large EGFP+/ChAT+ cells in the ventral spinal cord for several rAAV vectors (Figure 23). rAAVrh.10, rAAV9, rAAV7, rAAVrh.39 showed comparable efficiency transduction of motor neurons (Figure 21).

IV administration of AAV vectors mediated robust transduction in ventricles and brain blood vessels

[0142] EGFP expression was observed in the choroid plexus cells in lateral, 3rd and 4th ventricles of the animals infused with rAAVrh.39, rAAVrh.10, rAAVrh.43, rAAV7 and rAAV9 (ranked by transduction efficiency, Table 5 and Figures 19 and 24). EGFP expression in different ventricles of the same mouse brain was similar (Figure 24). Ependymal cells lining the ventricles were also transduced. An observation regarding the distribution of EGFP-positive cells was the apparent gradient with the highest number of transduced cells in peri-ventricular regions and progressively lower numbers with increasing distance to the ventricles. This was apparent in areas around the 3rd and 4th ventricles than the lateral ventricles (Figure 24). Extensive EGFP signal was also found with blood vessels throughout mouse brain and spinal cord. This was verified by dual immunofluorescent staining with antibodies directed to EGFP and a blood vessel endothelium specific marker, CD34 (Figures 27a and 27b). Unlike the rAAV transduction profiles in different regions of the brain parenchyma, the EGFP transduction of the blood vessels throughout the CNS was relatively uniform for any given vector. However, transduction of blood vessels was influenced by the particular rAAV used. A majority of rAAVs mediated moderate (e.g., rAAV6) to highly efficient (e.g. rAAVrh.10) blood vessel transduction in the CNS.

IV injection of AAV vectors did not cause microgliosis

[0143] Brain sections were also stained with antibody against Iba-1 to label microglial cells. The Iba-1-positive cells in the sections from mice received rAAVrh.10 was no more than those in naive or PBS-injected mice (Figure 28). This result indicated that intravascularly delivered rAAVs do not cause sustained inflammation in the CNS of mice 3 weeks after the injection of PI neonates.

Discussion of Results

[0144] In this study, the CNS transduction profile was evaluated for 10 different rAAV vectors delivered by intravascular infusion in neonatal mice. Most of the rAAVs can cross the BBB and mediate gene transfer to the neonatal mouse CNS with varying degrees of efficiency (Figures 19-21 and Table 5). After systemic administration, rAAVrh.10, rAAVrh.39, rAAVrh.43, and rAAV9 are the effective rAAVs with similar transduction capabilities and cellular tropism, as assessed by overall EGFP expression in the CNS. Specifically, a number of regions in the mouse CNS, including striatum, hippocampus, cortex, hypothalamus, cerebellum, medulla, and cervical spinal cord, revealed substantial EGFP expression. In addition, rAAV6.2 and rAAV7 were also effective. AAV1 and AAV6, achieved CNS transduction (Table 5). Native EGFP expression was detectable in brain and spinal cord sections for most of the rAAVs without immunostaining (Figure 29).

[0145] This example has clinical significance for gene therapy of CNS-related disorders, including for young patients. For a variety of neurological diseases, early treatment during infancy may be necessary to prevent irreversible CNS injury. The capacity of rAAVs to transduce large numbers of neuronal cells in different regions is relevant for treating neurological diseases such as spinal muscular atrophies, neuronal ceroid lipofuscinoses, and spinocerebellar degenerations. The efficiency of some rAAV vectors in transducing Purkinje and granule layer cells indicates that the vectors may be used for treating spinocerebellar ataxias. Transduction of astrocytes by rAAVs expressing secreted neurotrophic factors may be also beneficial for a number of neurodegenerative diseases such as Canavan's disease and amyotrophic lateral sclerosis. The vascular transduction in the CNS may be relevant for treating brain ischemia and stroke. The clinical application of intravascular rAAV-mediated gene delivery may also extend to the peripheral nervous system (PNS). Efficient transduction of DRG provides new therapeutic strategies for patients suffering from chronic pain.

[0146] Systemic gene delivery to the CNS is also useful as a method to manipulate gene expression in research. Effective and stable transgene expression in the CNS by intravenous administration of rAAVs may be applied to establish somatic transgenic animal models, which is a potentially cheaper, faster and simpler method than conventional transgenesis. Somatic CNS gene knock-down animal models may also be created using the method described herein.

[0147] Some rAAVs indeed demonstrated unique transduction profiles in the CNS. For instance, rAAV1 displayed transduced granule cells in the cerebellum, while rAAV6 and rAAV6.2 transduced Purkinje cells, and others transduced both types of cells (Figure 9). This indicates that once across the BBB, the rAAVs have distinct tropisms, which can be attributed to the capsid because that the vector genome used in all vectors was the same.

[0148] AAV serotypes disclosed herein can efficiently transduce brain capillary endothelial cells, neurons and astrocytes. This indicates that these vectors may extravasate from the circulation and reach the CNS parenchyma, possibly by crossing the BBB. AAV may cross the endothelial barrier by a transcytosis pathway. In this study, choroid plexuses and their surrounding parenchymal tissue were efficiently transduced. In addition, there was an apparent gradient of EGFP intensity from peri-ventricular (higher) to deep parenchymal (lower) tissue. These observations indicate that AAV may enter the neonatal mouse CNS through the choroid plexus, followed by widespread distribution via CSF and/or interstitial fluid flow to transduce neuronal and glial cells.

[0149] Neuronal- or glial-specific promoters, such as synapsin-1, and GFAP promoters may be used to restrict gene expression to a specific cell type. A further method to achieve targeted CNS gene delivery is to utilize RNA interference to detarget the peripheral tissues by post-transcriptional regulatory mechanisms. By adding microRNA binding sites into the 3' end of the transgene cassettes, transgene expression after systemic administration of AAV vectors may be reduced or eliminated in tissues such as liver, heart and skeletal muscle, while maintaining CNS transduction.

Materials And Methods

AAV production

[0150] ScAAV vectors were produced by trans-encapsidation of rAAV vector genome flanking by inverted terminal repeats (ITRs) from AAV2 with the capsids of different AAVs using the method transient transfection of 293 cells and CsCl gradient sedimentation as previously described. Vector preparations were titered by quantitative PCR. Purity of vectors was assessed by 4-12% SDS-acrylamide gel electrophoresis and silver staining (Invitrogen, Carlsbad, CA). Morphological integrity of each vector used in the study was examined by transmission electron microscopy of negative stained recombinant AAV virions. The expression of EGFP in the scAAV vector genome is directed by a hybrid CMV enhancer/chicken β -actin promoter.

Neonatal mouse injections

[0151] Wild-type C57BL/6 mice littermates were used. Mice breeding were conducted using programmatic timing method. Pregnant mice were monitored daily from embryonic day 17 to 21 to ensure the newborn pups could be dosed with vectors on P1. The mother (singly housed) of each litter to be injected was removed from the cage. Vectors were diluted to concentration of 4×10^{12} GCs/mL in PBS and 100 μ L of solution was subsequently drawn into 31G insulin syringes (BD Ultra-Fine II U-100 Insulin Syringes). P1 pups of C57BL/6 mice were anesthetized using isoflurane and rested on ice. For intravenous injections, a dissection microscope was used to visualize the temporal vein (located just anterior to the ear). The needle was inserted into the vein and the plunger was manually depressed. Correct injection was verified by noting blanching of the vein. Each pup received 4×10^{11} GCs of different scrAAVCBEGFP vectors (rAAV1, rAAV2, rAAV5, rAAV6, rAAV6.2, rAAV7, rAAV9, rAAVrh.10, rAAVrh.39, rAAVrh.43; n=6-8 mice per group) via the superficial temporal vein. After the injection pups were carefully cleaned, rubbed with their original bedding, and then returned to their original cage. The mother was then reintroduced to the cage after brief nose numbing using ethanol pads.

Histological processing

[0152] The study animals were anesthetized 21 days post-injection, then transcardially perfused with 15 mL of cold PBS followed by 15 mL of fixation solution containing 4% paraformaldehyde (v/v) with 0.2% of glutaraldehyde (v/v) in PBS. Then the whole carcasses were post-fixed in fixation solution for 5 days. Spinal cords and brains were extracted under a bright-field dissecting microscope, rinsed in PBS, and then cryoprotected in 30% sucrose (w/v) in PBS at 4 °C. Once the tissues sank to the bottom of the sucrose solution, they were embedded in Tissue-Tek OCT compound (Sakura Finetek, Torrance, CA) and frozen in a dry ice/ethanol bath. The tissue blocks were stored at -80 °C until sectioning. Serial 40 μ m floating sections of the entire brain were cut in a Cryostat (Thermo Microm HM 550). For the spinal cord, 3 mm length sections were taken from cervical, thoracic and lumbar regions, and then serial 40 μ m transverse sections prepared as above.

Immunostaining and microscopy imaging analysis

[0153] Brain and spinal cord sections were stained as floating sections in 12-well plates. Sections were washed 3 times in PBS for 5 min each time, and then incubated in blocking solution containing 1% Triton-X100 (v/v) (Fisher, Pittsburgh, PA), 5% dry-milk (w/v) and 10% goat serum (v/v) (Invitrogen) for 2h at room temperature. Then the sections were incubated with primary antibodies diluted in blocking solution at 4°C overnight. The following day tissue sections were washed twice in 0.05% Tween-20 (v/v) in PBS (PBST) and once with PBS, with each washing step lasting 10 min. Afterwards sections were incubated with appropriate secondary antibodies in blocking solution at room temperature for 2 h. Sections were washed again as above before mounting on glass slides. Vectashield with DAPI (Vector Laboratories, Burlingame, CA) was used to coverslip all slides, and then they were analyzed using a fluorescent inverted microscope (Nikon Eclipse Ti) or a Leica TSC-SP2 AOBS confocal microscope equipped with a 63 \times oil lens and a DM-IRE2 inverted microscope. The primary antibodies used in this study were as follows: rabbit anti-GFP (Invitrogen), goat anti-ChAT and mouse anti-NeuN (both from Millipore, Billerica, MA), mouse anti-GFAP (Cell signaling, Danvers, MA), rat anti-CD34 (Abcam, Cambridge, MA), mouse anti-Calbindin D-28k (Sigma, St Louis, MO) and rabbit anti-DARPP (Abcam, Cam-

bridge, MA). The secondary antibodies used in the study included: DyLight 488 AffiniPure Donkey Anti-rabbit IgG (Jackson ImmunoResearch, West Grove, PA); DyLight 549 AffiniPure Donkey Anti-Goat IgG (Jackson ImmunoResearch); DyLight 549 Affinipure Goat Anti-Rat IgG (Jackson ImmunoResearch); DyLight 594 AffiniPure Goat Anti-Mouse IgG (Jackson ImmunoResearch); goat anti-rabbit IgG-Alexa fluro 488 (Invitrogen) and goat anti-mouse IgG-Alexa fluro 568 (Invitrogen).

5

Semi-quantitative and quantitative comparison of EGFP transduction by different vectors

[0154] To generate a quantifiable and comparable data format, a semi-quantitative scoring system was developed to estimate transduction efficiency of different rAAV vectors in different regions of the mouse CNS. Briefly, regions with no EGFP positive cells were marked as (-). Regions with very few EGFP positive cells were scored (+), regions with some EGFP positive cells were ranked as (++)+, regions with many EGFP positive cells were marked as (+++). Finally, regions filled with EGFP positive cells were marked as (++++).

[0155] Next, 12 sub-anatomically and functionally important regions in the brain as well as cervical, thoracic and lumbar sections of the spinal cord were selected for quantitative analysis of images that were taken on a Nikon Eclipse Ti inverted microscope equipped with a Retiga 2000-RV CCD cooled camera. Nikon NIS elements AR software v. 3.2 was used for intensity quantification. Prior to quantification, optimal light source intensity and exposure times were obtained by plotting an intensity/exposure time curve using fluorescence reference slides (Ted Pella, prod. 2273). It was found that the intensity and exposure times had linear correlation. In addition, overexposure and extreme underexposure distorts the linear correlation. The maximum intensity (ND1) and a 20ms exposure were used for all sections to avoid overexposure. For quantification, fixed region of interest (ROI) was used to quantify the brightest area of any given brain region. A mean intensity (total intensity/size of ROI) was obtained for each region of all serotypes.

Table 5. Transduction characteristics of AAV serotypes following intravascular injections into neonatal mouse brain

	Olfactory Bulb		Striatum		Hippocampus		Cortex		Thalamus		Hypothalamus		Cerebellum		Medulla		Cervical		Thoracic		Lumber		Choroid Plexus	
	score	n	score	n	score	n	score	n	score	n	score	n	score	n	score	n	score	n	score	n	score	n	score	n
AAV1	+	3	++	3	++	3	++	3	+	3	+++	3	+++	3	+++	3	+++	3	+++	3	+	3	+++	3
AAV2	-	3	-	3	+	3	+	3	+	3	+	3	+	3	+	3	+	3	-	3	-	3	-	3
AAV5	-	3	-	3	-	3	+	3	-	3	+	3	-	3	-	3	-	3	-	3	-	3	-	3
AAV6	+	3	+	3	++	3	++	3	+	3	+++	2	++	3	++	3	++	3	+	3	+	3	++	3
AAV6.2	-	3	+++	2	++	3	++	3	+	3	++++	3	++	3	++	3	++	3	++	3	+	3	++++	3
AAV7	++	1	+++	3	+++	2	++	3	+	3	++++	3	+++	1	++	3	++	3	+	3	+	3	+++	3
AAV9	++	2	++	3	++	3	+++	1	+	3	++++	1	+++	1	++	3	+++	1	++	3	+	3	+++	3
rh10	++	1	+++	1	++	2	++	2	++	2	++++	3	+++	1	+++	3	++++	1	++	3	+	3	+++	3
rh39	++	1	++++	2	+++	3	+++	1	+	3	++++	3	+++	1	++++	1	++++	1	+++	3	+	3	+++	3

rh43	++	2	+++	1			++	2																
	++	3	+++	3	+++	3	+++	3	+	3	+++	1	++	3	+++	1	+++	2	+++	3	+	3	+++	3

Scoring: (-) no transduction, (+) very few positive cells, (++) some positive cells, (+++) many positive cells, and (++++) region is almost saturated with EGFP-positive cells.
The number of animals (n) with the particular score is given to the right of the score.

55

Example 10: Evaluation of an rAAV Based Treatment in a Canavan Disease Model Introduction to the Example

[0156] CD is a rare and fatal childhood leukodystrophy caused by autosomal recessive mutations in the aspartoacylase

gene (ASPA) [as established by G.G.'s graduate work (12)]. ASPA deficiency in CD patients leads to elevated N-Acetyl-Aspartic Acid (NAA) in urine (a hallmark of CD) and spongy degeneration of white matter throughout the CNS, producing severe psychomotor retardation and early death. An *ASPA*^{-/-} mouse model mimics the neuropathology and clinical manifestations seen in CD patients, i.e., spongy degeneration of white matter, motor deficits, developmental delays, and early death (within 3 weeks after birth).

[0157] In this study, i.v. deliverable rAAVs were used to target the CNS globally to treat diffused WM degeneration in CD mice. Single i.v. injections of *ASPA* vector to the neonatal CD mice corrected metabolic defect, psychomotor malfunction and other disease phenotypes, and prolonged survival. While untreated CD mice started showing growth retardation, psychomotor malfunction in the 2nd wk after birth and uniformly died soon after weaning, the treated mice began to gain weight 2 wks after vector injection and nearly caught up with their heterozygous littermates within 7 -8 weeks. Unlike CD mice, the mobility of the treated animals was similar to Wt littermates. Data from rotarod test on the treated mice showed no significant differences in the latency time among the treated CD mice and their age-matched Wt littermates, indicating that gene therapy corrected the ataxia, a typical neuromuscular symptom of CD. Biochemical characterization indicated reduction of NAA levels in the urine samples and restoration of ASPA activity in their brain and kidney tissues. Mitigation of the biochemical and clinical phenotypes was well correlated with globally ameliorated histopathology in not only the brain, spinal cord but also in the peripheral tissues such as kidney, indicating that CD is not just a CNS disorder.

Results

[0158] In CD mice were dosed at P1 (facial vein, 4x10¹¹ GCs) with AAV9ASPA. The mice were monitored for growth, gait, motor function on rotarod, NAA levels in urine and ASPA activities in brain. The results showed that i) Untreated CD mice started losing weight at the 2nd week and died in the 3rd week after birth; ii) The treated animals recovered their capacity to grow in the 5th week and caught up with *ASPA*^{+/+} animal by the 10th week; iii). Gene therapy completely corrected gait of CD mice as well as motor function of the CD mice treated at P1 (Figure 30A) as measured by rotarod test; iv). Gene therapy restored the vision of CD mice. The electroretinography (ERG) tests on the eyes of the CD mice showed non-recordable responses to light, while well-defined ERG responses were readily detectable in the treated CD mice (Figure 30B). These data indicate a severer retinopathy and loss of vision in CD mice and gene therapy can mitigate the retinopathy and restore the vision of CD mice; v). Gene therapy clearly improved metabolic defects of NAA as the NAA levels in the treated CD mice approach those in the control mice (Figure 30C); and vi) correction of NAA metabolism is well correlated with restoration of ASPA expression (Figure 30E) and activities (Figure 30D) in the brain of the treated CD mice.

[0159] To determine if the phenotypic corrections are correlated with alleviated neuropathology as well as *in situ* expression of ASPA in the brain sections of the treated CD mice, brain sections were analyzed at 3 months after gene therapy for neuropathology and ASPA immunohistochemistry. While the untreated mouse brain shows marked vacuolation that diffusely involves all regions of the brain and spinal cord, the vacuolation in both brain and spinal cord of the treated animal appears more patchy and variable with generally smaller-sized vacuoles. Some areas of the cerebral cortex show almost no vacuolation (Figure 31A). In addition, avidin-biotin complex (ABC) system was used to stain brain sections to detect ASPA expression in the cerebral cortex *in situ* (Figure 31B). To generate quantitative measurements on the improvement of neuropathology in the treated CD mice, the "vacuoles" in the brain and spinal cord sections caused by the white matter degeneration in the CD mice were quantified before and after gene therapy treatment. For this quantitative analysis, a Nikon Eclipse Ti inverted microscope and Nikon NIS elements AR software V.3.2 were used. Vacuoles that were > 3,000 pixels, 1,000 - 3,000 pixels and 100- 1,000 pixels were defined as large, medium and small vacuoles respectively. Among the 5 brain regions evaluated in this experiment, the olfactory bulb had the most dramatic mitigation in the white matter degeneration after gene therapy (Figure 32A). For the other 3 regions, while the large vacuoles were completely eliminated and the numbers of medium vacuoles were remarkably reduced, the reduction in the numbers of small vacuoles (<100 um) was not as significant in this experiment (Figure 32A). The same analysis on the spinal cord sections revealed a similar trend (Figure 32B).

[0160] Histopathology of the kidneys in the CD mice were evaluated. The glomeruli showed normal structure but were associated with dilation of Bowman's spaces. The renal tubular epithelium was diffusely attenuated (or atrophic) in association with enlargement of the tubular lumens (Figure 33A). In contrast, the treated CD mouse had normal glomeruli. Renal tubular epithelial cells were well-stained and normal in volume (Figure 33B). These results indicate the involvement of kidney in the pathophysiology of CD and kidney as a peripheral target for CD gene therapy. This result also indicates renal tropism of AAV vectors as a consideration for selection of a vector for CD gene therapy. Two vectors, rAAV9 and rh.10 were evaluated for efficiency of kidney transduction after IV delivery to 10 week old C57BL/6 mice. The results indicate the use of rAAVrh.10 (Figure 33D) as a useful vector for CD gene therapy because it transduces kidney efficiently in addition to its efficient CNS transduction (Figure 33C).

NUCLEIC ACID AND AMINO ACID SEQUENCES

[0161]

5 >gi|9632548|ref|NP_049542.1| capsid protein [Adeno-associated virus - 1]
 (SEQ ID NO: 1)

10 MAADGYLPDWLEDNLSEGIREWWDLKPGAPPKKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAALEHDKAYDQQLKAGDNPYLRYNHADAEGQERLQEDTS
 FGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKRPVEQSPQEPDSSSGIGKTGQQP
 AKKRLNFGQTGDSEVPDPQPLGEPPATPAAVGPTTMASGGGAPMADNNNEGADGV
 GNASGNWHCDSTWLGRVITTSTRTWALPTYNNHLYKQISSASTGASNDNHYFGYS
 TPWGYFDFNRFHCHFSRWDQRLINNNWGRPKRLNFKNLQVKEVTTNDGVTTIA
 NNLTSTVQVFSQMLRTGNNFTFSYTFEEVPFHSSYAHQSQSLRMLNPLIDQYLYLNRT
 15 SFYCLEYFPSQMLRTGNNFTFSYTFEEVPFHSSYAHQSQSLRMLNPLIDQYLYLNRT
 QNQSGSAQNKDLLSRGSPAGMSVQPKNWLPGPCYRQQRVSKTKTDNNNSNFTWT
 GASKYNLNGRESIINPGTAMASHKDDEDKFFPMMSGVMIFGKESAGASNTALDNVMIT
 20 DEEEIKAATNPVATERFGTVAVNFQSSSTDATGDVHAMGALPGMVWQDRDVYLQG
 PIWAKIPHTDGHFHPSPLMGGFGLKNPPPQILIKNTPVPANPPAEGSATKFASFITQYST
 GQVSVEIEWELQKENSKRWNPEVQYTSNYAKSANVDFTVDNNGLYTEPRPIGTRYL
 TRPL

25 >gi|110645923|ref|YP-680426.1| major coat protein VP1 [Adeno-associated virus - 2]
 (SEQ ID NO: 2)

30 MAADGYLPDWLEDTLSEGIRQWWKLKPGPPPKPAERHKDDSRGLVLPGYKYLGP
 NGLDKGEPVNEADAALEHDKAYDRQLSGDNPYLKYHADAEGQERLKEDTSFG
 GNLGRAVFQAKKRVLEPLGLVEEPVKTAPGKRPVEHSPVEPDSSSGTGKAGQQPA
 RKRLNFGQTGDADSPDPQPLGQPPAAPSGLGTNTMATGSGAPMADNNNEGADGV
 NSSGNWHCDSTWMGDRVITTSTRTWALPTYNNHLYKQISSQSGASNDNHYFGYSTP
 35 WGYFDFNRFHCHFSRWDQRLINNNWGRPKRLNFKNLQVKEVTTNDGVTTIAN
 NLTSTVQVFTDSEYQLPYVLGSAHQGCLPPFPADVFMVPQYGYLTNNGSQAVGRS
 SFYCLEYFPSQMLRTGNNFTFSYTFEDVPFHSSYAHQSQSLRMLNPLIDQYLYLSRT
 NTPSGTTTQSRLQFSQAGASDIRDQSRNWLPGPCYRQQRVSKTSADNNNSEYSWTG
 40 ATKYHLNGRDSLVPGPAMASHKDDDEKFFPQSGVLIFGKQGSEKTNVDIEKVMITD
 EEEIRTTNPVATEQYGSVSTNLQRGNRQAATADVNTQGVLPGMVWQDRDVYLQGPI
 WAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVPANPSTTFSAAKFASFITQYSTG
 QVSVEIEWELQKENSKRWNPEIQYTSNYNKSVDFTVDTNGVYSEPRPIGTRYLTR
 NL

45 >gi|51593838|ref|YP_068409.1| capsid protein [Adeno-associated virus - 5]
 (SEQ ID NO: 3)

50

55

5 MSFVDHPPDWLEEVGEGLREFLGLEAGPPKPKPNQQHQDQARGLVLPGYNYLPGN
 GLDRGEPVNRADEVAREHDISYNEQLEAGDNPYLKYNHADADEFQEKLADDSFGGN
 LGKAVFQAKKRVLEPFGLEEGAKTAPGTGKRIDDHFHKRKARTEEDSKPSTSSDAE
 10 AGPSGSQQLQIPAQPASSLGADTMSAGGGPLGDNNQGADGVGNASGDWHCDSTW
 MGDRVVTKSTRTWLPSYNHQQREIKSGVDGSNANA YFGYSTPWGYFDFNRFHS
 HWSPRDWQRLLINNYWGFRPRSLRKIFNIQVKETVQDSTTIANLTSTVQVFTDD
 DYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERSSFCLEYFPSK
 MLRTGNNFEFTYNFEEVPFHSSFAPSQNLFKLANPLVDQYLYRFVSTNNTGGVQFNK
 15 NLAGRYANTYKNWFGPMRTQGWNLGSGVNRASVAFATTNRMELEGASYQVPP
 QPNGMTNQLQGSNTYALENTMIFNSQPANPGTTATYLEGNMLITSESETQPVNRVAY
 NVGGQMATNNQSSSTTAPATGTYNLQEIVPGSVWMERDVYLQGPIWAKIPETGAHFH
 PSPAMGGFGLKHPPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEMEWEKK
 ENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTTRPIGTRYLTRPL

15 >gi|2766607|gb|AAB95450.1| capsid protein VP1 [Adeno-associated virus - 6]
 (SEQ ID NO: 4)

20 MAADGYLPDWLEDNLSEGIREWWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLRYNHADADEFQERLQEDTS
 FGGNLGRAVFQAKKRVLEPFGLEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQP
 25 AKKRLNFGQTGDSESVPDPQPLGEPPATPAAVGPTTMASGGGAPMADNNEGADGV
 GNASGNWHDSTWLGRVRTTSTRWALPTYNHLYKQISSASTGASNDNHYFGYS
 TPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLNFKNLQVKEVTTNDGVTTIA
 NNLTSTVQVFSDEYQLPYVLGSAHQGCLPPFPADVFMPQYGYTLNNGSQAVGRS
 SFYCLEYFPSQMLRTGNNFTSYTFEDVPFHSSYAHSQLDRLMNPLIDQYLYLNRT
 30 QNQSGSAQNKDLLFSRGSPAGMSVQPKNWLPGPCYRQQRVSKTKTDDNNNSNFTWT
 GASKYNLNGRESIINPGTAMASHKDDDKFPPMSGVMIFGKESAGASNTALDNVMIT
 DEEEIKATNPVATERFGTVAVNLQSSSTDATGDVHVMGALPGMVWQDRDVYLQG
 PIWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVANPPAERSATKFASFITQYST
 GQVSVEIEWELQKENSKRWNPEVQYTSNYAKSANVDFTVDNNGLYTEPRPIGTRYL
 TRPL

35 >gi|171850125|gb|ACB55302.1| capsid protein VP1 [Adeno-associated virus - 6.2]
 (SEQ ID NO: 5)

40 MAADGYLPDWLEDNLSEGIREWWDLKPGAPKPKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLRYNHADADEFQERLQEDTS
 FGGNLGRAVFQAKKRVLEPGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKTGQQP
 45 AKKRLNFGQTGDSESVPDPQPLGEPPATPAAVGPTTMASGGGAPMADNNEGADGV
 GNASGNWHDSTWLGRVRTTSTRWALPTYNHLYKQISSASTGASNDNHYFGYS
 TPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLNFKNLQVKEVTTNDGVTTIA
 NNLTSTVQVFSDEYQLPYVLGSAHQGCLPPFPADVFMPQYGYTLNNGSQAVGRS
 SFYCLEYFPSQMLRTGNNFTSYTFEDVPFHSSYAHSQLDRLMNPLIDQYLYLNRT
 50 QNQSGSAQNKDLLFSRGSPAGMSVQPKNWLPGPCYRQQRVSKTKTDDNNNSNFTWT
 GASKYNLNGRESIINPGTAMASHKDDDKFPPMSGVMIFGKESAGASNTALDNVMIT
 DEEEIKATNPVATERFGTVAVNLQSSSTDATGDVHVMGALPGMVWQDRDVYLQG
 PIWAKIPHTDGHFHPSPLMGGFGLKHPPPQILIKNTPVANPPAERSATKFASFITQYST
 GQVSVEIEWELQKENSKRWNPEVQYTSNYAKSANVDFTVDNNGLYTEPRPIGTRYL
 TRPL

55 >gi|22652861|gb|AAN03855.1|AF513851_2 capsid protein [Adeno-associated virus - 7]
 (SEQ ID NO: 6)

5 MAADGYLPDWLEDNLSEGIREWWDLKPGAPPKKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLRYNHADAECERLQEDTS
 FGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPAKKRPVEPSPQRSPDSSTGIGKKGQQ
 PARKRLNFGQTGDSEVPDPQPLGEPPAAPSSVSGTVAAAGGGAPMADNNEGADGV
 GNASGNWHCDSTWLGRVITTSTRTWALPTYNHLYKQISSETAGSTNDNTYFGYS
 TPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKLRFKLFNIQVKEVTNDGTTIA
 NNLTSTIQVFSQMLRTGNNFEFSYSFEDVPFHSSYAHQSLSLRLMNPLIDQYLYLARTQ
 10 SNPGGTAGNRELQFYQGGPSTMAEQAKNWLPGPCFRQQRVSKTLDQNNNSNFAWT
 GATKYHLNGRNSLVNPGVAMATHKDDEDRFPSSGVLIFGKTGATNKTTLENVLMT
 NEEEIRPTNPVATEEYGIVSSNLQAANTAAQTQVVNNQGALPGMVWQNRDVYLQGP
 15 IWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTAFNPKLNSFITQYSTG
 QVSVEIEWELQKENSKRWNPEIQYTSNFEKQTGVDFAVDSQGVYSEPRPIGTRYLTR
 NL

20 >gi|22652864|gb|AAN03857.1|AF513852_2 capsid protein [Adeno-associated virus - 8]
 (SEQ ID NO: 7)

25 MAADGYLPDWLEDNLSEGIREWWALKPGAPPKKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLQAGDNPYLRYNHADAECERLQEDTS
 FGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKRPVEPSPQRSPDSSTGIGKKGQQ
 PARKRLNFGQTGDSEVPDPQPLGEPPAAPSGVGPNMAAGGGAPMADNNEGADG
 VGSSSGNWHCDSTWLGRVITTSTRTWALPTYNHLYKQISNGTSGGATNDNTYFG
 YSTPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLSFKLFNIQVKEVTQNEGKT
 IANNLTSTIQVFTDSEYQLPYVLGSAHQGCLPPFPADVFMPQYGYLTNNNGSQAVGR
 30 SSFYCLEYFPSQMLRTGNNFQFTYTFEDVPFHSSYAHQSLSLRLMNPLIDQYLYLRS
 TQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGCYRQQRVSTTGQNNNSNFAW
 TAGTKYHLNGRNSLANPGIAMATHKDDERFPNSNGILIFGKQNAARDNADYSDVM
 LTSEEEIKTTNPVATEEYGIVADNLQQQNTAPQIGTVNSQGALPGMVWQNRDVYLQ
 35 GPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTFNQSKLNSFITQYS

40 TGQVSVEIEWELQKENSKRWNPEIQYTSNYYKSTSVDFAVNTEGVYSEPRPIGTRYL
 TRNL

45 >gi|46487805|gb|AAS99264.1| capsid protein VP1 [Adeno-associated virus 9]
 (SEQ ID NO: 8)

50 MAADGYLPDWLEDNLSEGIREWWALKPGAPQPKANQQHQDNARGLVLPGYKYLG
 PGNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLYKYNHADAECERLQEDTS
 FGGNLGRAVFQAKKRLLEPLGLVEEAAKTAPGKRPVEQSPQEPDSSAGIGKSGAQ
 AKKRLNFGQTGDTESVPDPQPIGEPPAAPSGVGSLTMASGGAPVADNNEGADGV
 SSSGNWHCDSQWLGRVITTSTRTWALPTYNHLYKQISNSTSGGSSNDNAYFGYST
 PWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLNFKLFNIQVKEVTDNNGVKTIA
 NNLTSTVQVFTSDYQLPYVLGSAHEGCLPPFPADVFMPQYGYLTNDGSQAVGRS
 SFYCLEYFPSQMLRTGNNFQFSYEFENVPFHSSYAHQSLSLRLMNPLIDQYLYLRS
 55 INGSGQNQQTLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTQNNNSEFAWPGA
 SSWALNGRNSLMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITN
 EEEIKTTNPVATESYGVATNHQSAQAQATGWWQNQGILPGMVWQDRDVYLQGP
 IWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTAFNPKLNSFITQYST
 GQVSVEIEWELQKENSKRWNPEIQYTSNYYKSNNEFAVNTEGVYSEPRPIGTRYLT
 RNL

>gi|29650526|gb|AAO88201.1| capsid protein [Non-human primate Adeno-associated virus]

(SEQ ID NO: 9 rh-10

5 MAADGYLPDWLEDNLSEGIREWWDLKPGAPPKKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLRYNHADAEGQERLQEDTS
 FGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEPSPQRSPDSSTGIGKKGQQ
 PAKKRLNFGQTGDSEVPDPQPIGEPPAGPSGLSGTMAAGGGAPMADNNEGADGV
 GSSSGNWHCDSTWLGDRVITTSTRTWALPTYNHLYKQISNGTSGGSTDNTYFGY
 STPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLNFKLFNQIYKVEVTQNEGKTI
 10 ANNLTSTIQVFTDSEYQLPYVLGSAHQGCLPPFPADVFMPQYGYLTNNNGSQAVGR
 SSFYCLEYFPSQMLRTGNNFEFSYQFEDVPFHSSYAHSQLDRLMNPLIDQYLYLSR
 TQSTGGTAGTQQQLFSQAGPNNMSAQAKNWLPGPCYRQQRVSTTLSQNNNSNFAW
 TGATKYHLNGRDSLNVPGVAMATHKDDEERFPSSGVLMFGKQGAGRDNVDYSSV
 15 MLTSEEEIKTTNPVATEQYGVVADNLQQQNAAPIVGAVNSQGALPGMVWQNRDVY
 LQGPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTFSQAKLASFIT
 QYSTGQVSVEIEWELQKENSKRWNPEIYQTSNYYKSTNVDAVNTDGTYSEPRPIGT
 RYLTRNL

>gi|171850147|gb|ACB55313.1| capsid protein VP1 [Adeno-associated virus - rh.39]

(SEQ ID NO: 10)

20 MAADGYLPDWLEDNLSEGIREWWALKPGAPPKKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLRYNHADAEGQERLQEDTS
 FGGNLGRAVFQAKKRVLEPLGLVEEAAKTAPGKKRPVEPSPQRSPDSSTGIGKKGQQ
 PAKKRLNFGQTGDSEVPDPQPIGEPPAGPSGLSGTMAAGGGAPMADNNEGADGV
 GSSSGNWHCDSTWLGDRVITTSTRTWALPTYNHLYKQISNGTSGGSTDNTYFGY
 STPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLSFKLFNQIYKVEVTQNEGKTI
 25 ANNLTSTIQVFTDSEYQLPYVLGSAHQGCLPPFPADVFMPQYGYLTNNNGSQAVGR
 SSFYCLEYFPSQMLRTGNNFEFSYTFEDVPFHSSYAHSQLDRLMNPLIDQYLYLSR
 TQSTGGTQGTQQQLFSQAGPANMSAQAKNWLPGPCYRQQRVSTTLSQNNNSNFAW

30 TGATKYHLNGRDSLNVPGVAMATHKDDEERFPSSGVLMFGKQGAGRDNVDYSSV
 MLTSEEEIKTTNPVATEQYGVVADNLQQNTGPIVGNVNSQGALPGMVWQNRDVY
 LQGPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTFSQAKLASFIT
 35 QYSTGQVSVEIEWELQKENSKRWNPEIYQTSNYYKSTNVDAVNTEGTYSEPRPIGT
 RYLTRNL

>gi|46487767|gb|AAS99245.1| capsid protein VP1 [Adeno-associated virus rh.43]

(SEQ ID NO: 11)

40 MAADGYLPDWLEDNLSEGIREWWDLKPGAPPKKANQQKQDDGRGLVLPGYKYLG
 PFNGLDKGEPVNAADAAALEHDKAYDQQLEAGDNPYLRYNHADAEGQERLQEDTS
 FGGNLGRAVFQAKKRVLEPLGLVEEGAKTAPGKKRPVEQSPQEPDSSSGIGKKGQQP
 ARKRLNFGQTGDSEVPDPQPLGEPPAAPSGVGPNTMAAGGGAPMADNNEGADGV
 GSSSGNWHCDSTWLGDRVITTSTRTWALPTYNHLYKQISNGTSGGATNDNTYFGY
 STPWGYFDFNRFHCHFSPRDWQRLINNNWGRPKRLSFKLFNQIYKVEVTQNEGKTI
 45 ANNLTSTIQVFTDSEYQLPYVLGSAHQGCLPPFPADVFMPQYGYLTNNNGSQAVGR
 SSFYCLEYFPSQMLRTGNNFQFTYTFEDVPFHSSYAHSQLDRLMNPLIDQYLYLSR
 TQTTGGTANTQTLGFSQGGPNTMANQAKNWLPGPCYRQQRVSTTGQNNNSNFAW
 TAGTKYHLNGRNSLANPGIAMIATHKDDEERFPVTGSCFWQQNAARDNADYSDVM
 50 LTSEEEIKTTNPVATEEYGIVADNLQQQNTAPQIGTVNSQGALPGMVWQNRDVYLQ
 GPIWAKIPHTDGNFHPSPLMGGFGLKHPPPQILIKNTPVPADPPTFNQSKLNSFITQYS
 TGQVSVEIEWELQKENSKRWNPEIYQTSNYYKSTSVDFAVNTEGVYSEPRPIGTRYL
 55 TRNL

>capsid protein VP1 [Adeno-associated virus] CSp3
(SEQ ID NO: 12)

5 MAADGYLPDWLEDNLSEGIREWWALKPGAPQPKANQQHQDNARGLVLPGYKYLG
PGNGLDKGEPVNAADAAALEHDKAYDQQLKAGDNPYLKYNHADAEGQERLKEDTS
FGGNLGRAVFQAKKRLLEPLGLVEEAAKTAPGKKRPVEQSPQEPDSSAGIGKSGAQP
AKKRLNFGQTGDTESVPDPQPIGEPPAAPSGVGSLTIASGGGAPVADNNEGADGVGS
SSGNWHCDSQWLGDRTTSTRTWALPTYNHLYKRISNSTSGSSNDNAYFGYSTP
WGYFDFNRFHCHFSRWDQRLINNNWGRPKRLNFKLFNIRVKEVTDNNGVKTITN
10 NLTSTVQVFTDSDYQLPYVLGSAHEGCLPPFPADVFMPQYGYLTLDGSQAVGRSS
FYCLEYFPSPQMLRTGNNFQFSYEFENVPFHSSYAHQSQDRLMNPLIDQYLYYLSKTI
NGSGQNQQLKFSVAGPSNMAVQGRNYIPGPSYRQQRVSTTVTRNNNSEFAWPGAS
SWALNGRNSLMNPGPAMASHKEGEDRFFPLSGSLIFGKQGTGRDNVDADKVMITNE
15 EEEIKTTNPVATESYQGVATNHQSAQAQATGWVQNQGILPGMVWQDRDVYLQGPI
WAKIPHTDGNFHPSPLMGGFGVKHPPPQILIKNTPVPADPPTAFNKKDKLNSFITQYST
GQVSVEIEWELQKENSKRWNPEIYQYTSNYYKSNNVEFAVNTEGVYSEPRPIGTRYLT
RNL

20 >gi|189339202|ref|NP_001121557.1| aspartoacylase [Homo sapiens]
(SEQ ID NO: 13)

25 MT\$CHIAEEHIQKVAIFGGTHGNELTGVFLVKHWLENGAEIQRTGLEVKPFITNPRAV
KKCTRYIDCDLNRFIDLENLGKKMSEDLPYEVRRQAQEINHLFGPKDSEDSYDIIFDLH
NTTSNMGCTLILEDSRNNFLIQMFHYIKTSLAPLPCYVYIEHPSLKYATTRSIAKYPV
GIEVGPQPQGVLRADILDQMRKMIKHALDFIHFFNEGKEFPPCAIEVYKIEKVDYPR
DENGIAAIHPNLQDQDWKPLHPGDPMLTLDGKTIPLGGDCTVYPVFVNEAAYYE
30 KKEAFAKTTKLTNAKSIRCCLH

>gi|189339201:92-1033 Homo sapiens aspartoacylase (Canavan disease) (ASPA), transcript variant 2, mRNA
(SEQ ID NO: 14)

35 ATGACTTCTTGTACATTGCTGAAGAACATATACAAAAGGTTGCTATCTTGGAG
GAACCCATGGGAATGAGCTAACCGGAGTATTCTGGTTAACGATTGGCTAGAGA
ATGGCGCTGAGATTAGAGAACAGGGCTGGAGGTAAAACCATTATTACTAACCC
CCAGAGCAGTGAAGAACAGTGTACCAAGATATTGACTGTGACCTGAATCGCATT
TGACCTTGAAAATCTTGGCAAAAAAAATGTCAGAACAGATTTGCCATATGAAGTGAG
AAGGGCTCAAGAAATAATCATTATTGGTCCAAAAGACAGTGAAGATTCTAT
GACATTATTTGACCTTCACAACACCACCTCTAACATGGGTCACACTTTATTCT
TGAGGATTCCAGGAATAACTTTAATTCAAGATGTTCAATTACATTAAAGACTTCTC
40 TGGCTCCACTACCCTGCTACGTTATCTGATTGAGCATCCTCCCTCAAATATGCG
ACCACTCGTCCATAGCCAAGTATCCTGTGGGTATAGAACAGTTGGCCTCAGCCTC
AAGGGGTTCTGAGAGCTGATATCTGGATCAAATGAGAAAAATGATTAAACATG
CTCTGATTATACATCATTCAATGAAGGAAAAGAATTCCCTCCCTGCCATT
GAGGTCTATAAAATTATAGAGAAAGTTGATTACCCCCGGATGAAATGGAGAA
ATTGCTGCTATCATCCATCTAACATCTGCAGGATCAAGACTGGAAACCAC
45 CTGGGGATCCCATGTTTAACCTCTGATGGGAAGACGATCCCACGGCGGAGA
CTGTACCGTGTACCCGTGTTGTGAATGAGGCCGATATTACGAAAAGAAGA
AGCTTTGCAAAGACAACAACTAACGCTCAATGCAAAAGTATTGCTGCTGT
50 TTACATTAG

55 >gi|31560279|ref|NP_075602.2| aspartoacylase [Mus Musculus]
(SEQ ID NO: 15)

5 MTSCVAKEPIKKIAIFGGTHGNELTVFVTHWLRNGTEVHRAGLDVKPFITNPRAV
 EKCTRYIDCDLNRFVFDLENLSKEMSEDLPYEVRRRAQEINHFGPKNSDDAYDLVFDL
 HNTTSNMGCTLILEDSRNDFLIQMFHYIKTCMAPLPCSVYLIEHPSLKYATTRSIAKYP
 VGIEVGPQPHGVLRADILDQMRKMIKHALDFIQHFNEGKEFPPCSIDVYKIMEKVDYP
 RNESGDMAAVIHPNLQDQDWKPLHPGDPVFVSLDGKVIPLGGDCTVYPVFVNEAAY
 YEKKEAFAKTTKLTLSAKSIRSTLH

10 >gi|142354273:148-1086 Mus musculus aspartoacylase (Aspa), mRNA
 (SEQ ID NO: 16)

15 ATGACCTCTTGTGCTAAAGAACCTATTAAGAAGATTGCCATCTTGGAGGGAA
 CTCATGGAAATGAAC TGACCGGAGTGTCTAGTTACTCACTGGCTAAGGAATGG
 CACTGAAGTTCACAGAGCAGGGCTGGACGTGAAGCCATTCAATTACCAATCCAAG
 GGC GGTTGGAGAAGTGCACCAAGATACTGACTGTGACCTGAATCGTGTGTTTGAC
 CTTGAAAATCTTAGCAAAGAGATGTCTGAAGACTTGCATATGAAGTGAGAAGG
 GCTCAAGAAATAAATCATTATTGGTCCAAAAAATAGTGTATGCCTATGACC
 TTGTTTTGACCTTCACAACACCACTCTAACATGGGTGCACTCTTATTCTTGAG
 GATTCCAGGAATGACTTTAATTCAAGATGTTCACTATATTAAAGACTTGATGGC
 TCCATTACCCCTGCTCTGTTATCTCATTGAGCATCCTCACTCAAATATGCAACCA
 CTCGTTCCATTGCCAAGTATCCTGTTGGTATAGAAGTTGGCCTCAGCCTCACGGT
 GTCCTTAGAGCTGATATTAGACCAATGAGAAAAATGATAAAACATGCTCTG
 ATTTATACAGCATTCATGAAGGAAAAGAATTCCCTCCCTGTTCTATTGACGTC
 TATAAAATAATGGAGAAAGTTGATTATCCAAGGAATGAAAGTGGAGACATGGCT
 GCTGTTATTCACTCTAACAGACTGGAAACCATTGCACCCCTGGAG

30 >gi|48762945:149-613 Homo sapiens superoxide dismutase 1, soluble (SOD1), mRNA
 (SEQ ID NO: 17)

40 ATGGCGACGAAGGCCGTGCGTGCTGAAGGGCGACGGCCAGTGCAGGGCATIC
 ATCAATTTCGAGCAGAAGGAAAGTAATGGACCAGTGAAAGGTGTGGGGAAAGCATT
 AAAGGACTGACTGAAGGCCTGCATGGATTCCATGTTATGAGTTGGAGATAATA
 CAGCAGGCTGTACCAAGTGCAGGTCTCACTTTAACCTCTATCCAGAAAACACGG
 TGGGCCAAAGGATGAAGAGAGGCATGTTGGAGACTTGGCAATGTGACTGCTGA
 CAAAGATGGTGTGCCGATGTGTCTATTGAAGATTCTGTGATCTCACTCTCAGGA
 GACCATTGCATCATTGGCCGACACTGGTGGTCCATGAAAAGCAGATGACTTG
 GGCAAAGGTGAAATGAAGAAAGTACAAAGACAGGAAACGCTGGAAGTCGTTT
 GGCTTGTGGTGTATTGGATCGCCCAATAA

50 >gi|4507149|ref|NP_000445.1| superoxide dismutase [Homo sapiens]
 (SEQ ID NO: 18)

55 MATKAVCVLKGDGPVQGIINFEQKESNGPVKVVGSIKGLTEGLHGFHVHEFDNTA
 GCTSAGPHFNPLSRKHGGPKDEERHVGDLGNTADKDGVADVSIEDSVISLSDHCII
 GRTLVVHEKADDLGKGGNEESTKTGNAGSRLACGVIGIAQ

>gi|45597446:117-581 Mus musculus superoxide dismutase 1, soluble (Sod1), mRNA
 (SEQ ID NO: 19)

ATGGCGATGAAAGCGGTGTGCGTGCTGAAGGGCGACGGTCCGGTGCAGGGAACC
 5 ATCCACTCGAGCAGAAGGCAAGCGGTGAACCAGTTGTTGTCAGGACAAATT
 ACAGGATTAACGTGAAAGGCCAGCATGGGTTCCACGTCCATCAGTATGGGGACAAT
 ACACAAGGCTGTACCAAGTGCAGGACCTCATTTAACCTCACTCTAAGAAACATG
 GTGGCCCGGCGGATGAAGAGAGGCATGTTGGAGACCTGGGCAATGTGACTGCTG
 10 GAAAGGACGGTGTGGCAATGTGTCATTGAAGATCGTGTGATCTCACTCTCAGG
 AGAGCATTCCATCATTGGCCGTACAATGGTGGTCCATGAGAAACAAGATGACTT
 GGGCAAAGGTGAAATGAAGAAAGTACAAAGACTGGAAATGCTGGAGCCGCT
 TGGCCTGTGGAGTGATTGGGATTGCGCAGTAA

>gi|45597447|ref|NP_035564.1| superoxide dismutase [Mus musculus]
 (SEQ ID NO: 20)

15 MAMKAVCVLKGDPVQGTIHFEQKASGEPVVLSGQITGLTEGQHGFHVHQYGDNT
 QGCTSAGPHFNPHSKKHGGPADEERHVGDLGNVTAGKDGVANVSIEDRVISLSGEH
 SIIGRTMVVHEKQDDLGKGGNEESTKTGNAGSRLACGVIGIAQ

20 >pAAVscCB6 EGFPmir SOD5 (direct) 5243bp (SEQ ID NO: 21)

25 CTGCGCGCTCGCTCGACTGAGGCCGCCGGCAAAGCCCGGGCGTCGGCG
 ACCTTGGTCGCCCGCCTCAGTGAGCGAGCGAGCGCAGAGAGGGAGTGTAG
 CCATGCTCTAGGAAGATCAATTCAATTACCGGGTCATTAGTTCATAGCCCATA
 TATGGAGTTCC
 TATTAATAGTAATCAATTACGGTAAATGGCCCGCTGGCTGACCGCCAACGACCCCCG
 GCGTTACATAACTACGGTAAATGGCCCGCTGGCTGACCGCCAACGACCCCCG
 CCCATTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCAATAGGGACTTTC

30

35

40

45

50

55

CATTGACGTCAATGGGTGGATATTACGGTAAACTGCCCACTGGCAGTACATCA
 5 AGTGTATCATATGCCAAGTACGCCCTATTGACGTCAATGACGGTAAATGGCC
 GCCTGGCATTATGCCAGTACATGACCTTATGGGACTTCTACTTGGCAGTACA
 TCTACGTATTAGTCATCGCTATTACCATGTCGAGGCCACGTTCTGCTTCACTCTCC
 CCATCTCCCCCCCCTCCCCACCCCCAATTGTATTATTATTATTAAATTATT
 GTGCAGCGATGGGGCGGGGGGGGGCGCGCAGGCAGGGCGGGCGGGCGGG
 GCGAGGGCGGGCGGGCGAGGCAGGGTGCAGGCAGCCAATCAGAGC
 GGCAGCGCTCCGAAAGTTCTTTATGGCAGGGCGGGCGGGCGGCCCTATA
 10 AAAAGCGAAGCGCGCGGGCGGGAGCAAGCTCTAGCCTCGAGAATTACCGCG
 TGGTACCTCTAGAGCAGAGCTCGTTAGTGAACCGTCAGTCGAATGCCACCA
 TGGTGAGCAAGGGCGAGGAGCTGTTACCGGGTGGTGCCTCCTGGTCAGC
 TGGACGGCGACGTAAACGCCACAAGTTCAGCGTGTCCGGCAGGGCGAGGGCG
 ATGCCACCTACGGCAAGCTGACCTGAAGTTCATCTGACCAACCGCAAGCTGCC
 15 CGTGCCTGGCCACCTCGTACCGACCCCTGACCTACGGCGTGCAGTGCTTCAGC
 CGCTACCCCGACCACATGAAGCAGCACGACTTCTCAAGTCCGCATGCCGAAG
 GCTACGTCCAGGAGCGCACCATCTTCAAGGACGACGGCAACTACAAGACCC
 GCGCCGAGGTGAAGTTCGAGGGGACACCCCTGGTGAACCGCATCGAGCTGAAGG
 GCATCGACTTCAAGGAGGACGGCAACATCCTGGGCACAAGCTGGAGTACAAC
 20 ACAACAGCCACACGTCTATATCATGGCGACAAGCAGAAGAACGGCATCAAGG
 TGAACCTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGTCGCCGACC
 ACTACCAGCAGAACACCCCCATCGCGACGGCCCGTGCTGCTGCCGACAACC
 ACTACCTGAGCACCCAGTCCGCCCTGAGCAAAGACCCAACGAGAACGGCGATC
 ACATGGTCTGCTGGAGITCGTACCGCCGGGATCACTCTGGCATGGACGA
 25 GCTGTACAAGTAAGTAACAGGTAAAGTGCATCGTAATGCGGGAAAGCTTTAT
 TCGGGTGAGATGGGCTGGGACCATCTGGGGACCCCTGACGTGAAGTTGTCACT
 GACTGGAGAACTCGGTTGTCGTCTGTTGCGGGGGCGGCAGTTATGGCGGTGCC
 TTGGGCAGTGCACCGTACCTTGGAGCGCGCAGCCCTCGTGTGTCGTGACGTC
 30 ACCCGTTCTGTTGGTACCTGCTGTTGACAGTGAAGCGACGCAATGTGACTTCGCTG
 ACAAAAGCTGTGAAGCCACAGATGGGCTTGTCAAGCAGTCACATTGCGCTGCCTAC
 TGCCTCGGACTTCAAGGGCTCGAGAAATTCAAGGGTGGGCCACCTGCCGGTAGGT
 GTGCGGTAGGCTTTCTCCGTCGAGGACGCAAGGTTGGCCTAGGGTAGGCTC
 35 TCCTGAATCGACAGGCGCCGGACCTCTGGCGGCCAACAACACGCGTCTGACC
 ATTCACTCTCTTCTTCTGCAGGTTGTGGAAGAAATGGGATCCGATCTTT
 TCCCTCTGCCAAAAATTATGGGACATCATGAAGCCCTTGAGCATTGACTTCT
 GGCTAATAAAAGGAAATTATTTCAATTGCAATAGTGTGTTGGAATTTTGTGTCT
 CTCACTCGGCCTAGGTAGATAAGTAGCATGGCGGTTAATCAATTAACTACAAGG
 40 AACCCCTAGTGAATGGAGTTGCCACTCCCTCTGCGCGCTCGCTCGACTGA
 GGCGGGCGACCAAAGGTCGCCGACGCCGGCTTGCCCCGGCGCCCTCAGT
 GAGCGAGCGAGCGCGCAGCCTTAATTAACTTAATTCACTGGCCGTTACAA
 CGTCGTGACTGGAAAACCCCTGGCGTACCCAACCTTAATGCCCTGCAGCACATC
 CCCCTTCGCCAGCTGGCTAATAGCGAAGAGGCCGACCGATGCCCTCCCA
 ACAGTTGCGCAGCCTGAATGGCAATGGACGCGCCCTGTAGCGCGCATTAAG
 45 CGCGGGGGTGTGGTTACGCGCAGCGTACCGCTACACTTGCCAGCGCCCTA
 GCGCCCGCTCTTCGCTTCTTCCCTTCTGCCACGTTGCCGGCTTCC
 CGTCAAGCTCTAAATGGGGGCTCCCTTAGGGTCCGATTAGTGTCTTACGGC
 ACCTCGACCCAAAAACTGATTAGGGTGTGGTACGTTACGTAAGTGGGCCATGCC
 CTGATAGACGGTTTCGCCCTTGACGTTGGAGTCCACGTTCTTAATAGTGGAC
 50 TCTTGTCTCAAACGGAAACAACACTCAACCTATCTCGGTCTATTCTTTGATTAA

TAAGGGATTTGCCGATTCGGCCTATTGGTTAAAAAATGAGCTGATTAAACAAA
 5 AATTAAACGCGAATTAAACAAATATTAACGCTACAATTAGGTGGCACTTT
 CGGGGAATGTGCGCGAACCCCTATTGTTATTCTAAATACATTCAAATA
 TGTATCCGCTCATGAGACAATAACCCGTATAAAATGCTCAATAATATTGAAAAAG
 GAAGAGTATGAGTATTCAACATTCCGTGTCGCCCTATTCCCTTTGCGGCAT
 10 TTTGCCTTCTGTTTGCTCACCCAGAACGCTGGTAAAGTAAAAGATGCTGA
 AGATCAGTGGGTGCACGAGTGGTTACATCGAACTGGATCTCAACAGCGGTAA
 GATCCTGAGAGTTCGCCCCGAAGAACGTTCCAATGATGAGCAGCTTTAAA
 GTTCTGCTATGTGGCGCGGTATTATCCGTATTGACGCCGGCAAGAGCAACTCG
 GTCGCCGCATAACACTATTCTCAGAACATGACTGGTGTGAGTACTCACCAGTCACAGA
 15 AAAGCATCTTACGGATGGCATGACAGTAAGAGAATTATGCAGTGCTGCCATAAC
 CATGAGTGATAACACTGCCAACTTACTTGACAAACGATCGAGGACCGAA
 GGAGCTAACCGCTTTTGACAACATGGGGGATCATGTAACTCGCCTGATCGT
 TGGGAACCGGAGCTGAATGAAGCCATACCAAACGACGAGCGTGACACCACGATG
 CCTGTAGCAATGGCAACAAACGTTGCGCAAACATATTAACTGGCGAACTACTACTC
 TAGCTTCCCAGCAACAATTAAATAGACTGGATGGAGGCCATAAGTTGAGGAC
 20 CACTTCTGCGCTCGGCCCTCCGGCTGGCTGGTTATTGCTGATAAATCTGGAGCC
 GGTGAGCGTGGGTCTCGCGGTATCATTGCAACTGGGCCAGATGGTAAGCCC
 TCCCGTATCGTAGTTATCACACGACGGGAGTCAGGCAACTATGGATGAACGA
 AATAGACAGATCGCTGAGATAGGTGCTCACTGATTAAGCATGGTAACTGTCAG
 25 ACCAAGTTACTCATATAACTTAGTGGATTAAACTTCATTAAATTTAA
 AGGATCTAGGTGAAGATCCCTTTGATAATCTCATGACCAAATCCCTAACGTG
 AGTTTCGTTCACTGAGCGTCAGACCCGTAGAAAAGATCAAAGGATCTTCTTG
 AGATCCTTTCTGCGCTAATCTGCTGCTGCAAACACAAAAACCCACCGCTA
 CCAGCGGTGGTTGTTGCCGATCAAGAGCTACCAACTCTTCCGAAGGTAA
 30 CTGGCTTCAGCAGAGCGCAGAACGACGGGAGTCAGGCTAGTT
 AGGCCACCACTCAAGAAACTCTGTAGCACCCTACATACCTCGCTGCTAATC
 CTGTTACCACTGGCTGCTGCCAGTGGCGATAAGTCGTGCTTACCGGGTTGGACT
 CAAGACGATAGTACCGGATAAGGCGCAGCGCTGGCTGAACGGGGGTTCGT
 GCACACAGCCCAGCTGGAGCGAACGACCTACACCGAACTGAGATACCTACAGC
 35 GTGAGCTATGAGAAAGCGCCACGCTTCCGAAGGGAGAAAGGCAGGACAGGTATC
 CGGTAAAGCGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTCCAGGGGAA
 AACGCCTGGTATCTTATAGCTCTGCGGTTGCCACCTCTGACTTGAGCGTCG
 ATTTTGATGCTCGTCAGGGGGCGGAGCCTATGGAAAAACGCCAGCAACGC
 40 GGCCTTCTGCGCTGCTGGCTTTGCTGGCTTTGCTCACATGTTCTTCCCTGC
 GTTATCCCCGATTCGTGATAACCGTATTACGCCCTTGAGTGAGCTGATAACC
 GCTCGCCGAGCCAAACGACCGAGCGCAGCGAGTCAGTGAGCGAGGAAGCGGA
 AGAGCGCCAATACGCAAACCGCCTCTCCCGCGCTGGCCATTACATTAATGC
 AGCTGGCACGACAGGTTCCGACTGGAAAGCGGGAGTGAGCGAACGCAATT
 45 AATGTGAGTTAGCTCACTCATTAGGCACCCAGGCTTACACTTATGCTCCGG
 CTCGTATGTTGTGGAATTGTGAGCGGATAACAATTACACAGGAAACAGCTA
 TGACCATGATTAGCCAGATTAATTAAGGCCTTAATTAGG

>sod1mir1 (direct) 108bp
 (SEQ ID NO: 22)

50 TGCTGTTGACAGTGAGCGACATCATCAATTCCGAGCAGAACTGTGAAGCCACA
 GATGGGTTCTGCTCGAAATTGATGATGCTGCTACTGCCCTGGACTTCAAGGG

>sod1mir2 (direct) 106bp
 (SEQ ID NO: 23)

55 TGCTGTTGACAGTGAGCGACGCATTAAGGATCCTGACTGACTGTGAAGCCACA
 GATGGGTCAGTCAGTCCTTAATGCGCTGCCACTGCCCTGGACTTCAAGGG

5 >sodlmir3 (direct) 108bp
(SEQ ID NO: 24)

TGCTGTTGACAGTGAGCGACTGCATGGATTCTCCATGTCATCTGTGAAGCCACA
GATGGGATGAACATGGAATCCATGCAGCTGCCTACTGCCTCGGACTTCAAGGG

10 >sodlmir4 (direct) 106bp
(SEQ ID NO: 25)

TGCTGTTGACAGTGAGCGACAAGGATGAAGATCGAGGCATGCTGTGAAGCCACA
GATGGGCATGCCTCTCTTCATCCTGCTGCCTACTGCCTCGGACTTCAAGGG

15 >sodlmir5 (direct) 110bp
(SEQ ID NO: 26)

TGCTGTTGACAGTGAGCGACGCAATGTGACTTCGCTGACAAAGCTGTGAAGCCA
CAGATGGGCTTGTCAGCAGTCACATTGCGCTGCCTACTGCCTCGGACTTCAAGG
G

20 >sodlmir6 (direct) 108bp
(SEQ ID NO: 27)

TGCTGTTGACAGTGAGCGACCGATGTGTCTATCTGAAGATTCTGTGAAGCCACA
GATGGGAATCTCAATAGACACATCGGCTGCCTACTGCCTCGGACTTCAAGGG

25 >sodlmir7 (direct) 106bp
(SEQ ID NO: 28)

TGCTGTTGACAGTGAGCGACGGTGAAATGATCAGAAAGTACTGTGAAGCCACA
GATGGGTACTTCTCATTCCACCGCTGCCTACTGCCTCGGACTTCAAGGG

30 >sodlmir8 (direct) 110bp
(SEQ ID NO: 29)

TGCTGTTGACAGTGAGCGACGCTGTAGAAATTCGTATCCTGATCTGTGAAGCCAC
AGATGGGATCAGGATACATTCTACAGCGCTGCCTACTGCCTCGGACTTCAAGGG

35 >sodlmir9 (direct) 106bp
(SEQ ID NO: 30)

TGCTGTTGACAGTGAGCGAGGTATTAACATTGTCAGAATTAGTGAAGCCACAGA
TGTAAATTCTGACAAGTTAATACCTGCCTACTGCCTCGGACTTCAAGGG

40 >pAAVscCB6 EGFPmir scr (1820bp - 1925bp, direct) 106bp
(SEQ ID NO: 31)

TGCTGTTGACAGTGAGCGACGATGCTCTAACCGTTCTATCAAGTGAAGCCACAG
ATGTTGATAGAACCTTAGAGCATCGCTGCCTACTGCCTCGGACTTCAAGGG

[0162] This invention is not limited in its application to the details of construction and the arrangement of components

set forth in this description or illustrated in the drawings. The invention is capable of other embodiments and of being practiced or of being carried out in various ways. Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of "including," "comprising," or "having," "containing," "involving," and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

[0163] Having thus described several aspects of at least one embodiment of this invention, it is to be appreciated various alterations, modifications, and improvements will readily occur to those skilled in the art. Such alterations, modifications, and improvements are intended to be part of this disclosure, and are intended to be within the spirit and scope of the invention. Accordingly, the foregoing description and drawings are by way of example only.

10 **Embodiments of the invention will now be described in the following numbered paragraphs:**

1. A method for delivering a transgene to CNS tissue in a subject, the method comprising: administering an effective amount of a rAAV by intrathecal administration, wherein the rAAV comprises (i) a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 and (ii) a nucleic acid comprising a promoter operably linked with a transgene.
- 15 2. The method of paragraph 1 further comprising administering an effective amount of the rAAV by intracerebral administration.
- 20 3. A method for delivering a transgene to central nervous system (CNS) tissue in a subject, the method comprising: administering an effective amount of a rAAV by intrathecal administration and by intracerebral administration, wherein the rAAV infects cells of CNS tissue in the subject and comprises a nucleic acid comprising a promoter operably linked with a transgene.
- 25 4. The method of paragraph 2 or 3, wherein the intracerebral administration is an intraventricular administration.
- 5 5. The method of any one of paragraphs 1 to 4, wherein the intrathecal administration is in the lumbar region of the subject
- 30 6. The method of paragraph 4 or 5, wherein the intraventricular administration is an administration into a ventricular region of the forebrain of the subject.
- 35 7. The method of any one of paragraphs 1 to 6, wherein the dose of the rAAV for intrathecal administration is in a range of 10^{10} genome copies to 10^{11} genome copies.
8. The method of any one of paragraphs 2 to 6, wherein the dose of the rAAV for intracerebral administration is in a range of 10^{10} genome copies to 10^{11} genome copies.
- 40 9. The method of any one of paragraphs 1 to 8, wherein the transgene is a CNS-associated gene.
10. The method of paragraph 9, wherein the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AACD) or aspartoacylase (ASPA).
- 45 11. The method of any one of paragraphs 1 to 8, wherein the transgene encodes an inhibitory RNA that binds specifically to SOD1 mRNA and inhibits expression of SOD1 in the subject.
12. The method of paragraph 11, wherein the inhibitory RNA is an antisense RNA, a shRNA or a miRNA.
- 50 13. The method of paragraph 11, wherein the inhibitory RNA has a sequence as set forth in SEQ ID NO: 26.
14. A nucleic acid comprising a sequence as set forth in SEQ ID NO: 26.
- 55 15. A recombinant AAV comprising the nucleic acid of paragraph 14.
16. The recombinant AAV of paragraph 15 further comprising a capsid protein comprising a sequence as set forth in SEQ ID NO: 9.

17. A method for treating amyotrophic lateral sclerosis (ALS) in a subject in need thereof, the method comprising: administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises (i) a capsid protein comprising a sequence as set forth in SEQ ID NO: 9 and (ii) a nucleic acid comprising a promoter operably linked with a region encoding an inhibitory RNA that binds specifically to SOD1 mRNA and inhibits expression of SOD1 in the subject.

18. The method of paragraph 17, wherein the inhibitory RNA is an antisense RNA, a shRNA or a miRNA.

19. The method of paragraph 18, wherein the inhibitory RNA has a sequence as set forth in SEQ ID NO: 26.

20. A method for treating amyotrophic lateral sclerosis (ALS) in a subject in need thereof, the method comprising: administering an effective amount of a rAAV to the subject, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding a sequence as set forth in SEQ ID NO: 26 and wherein the rAAV infects cells of CNS tissue in the subject.

21. A method for delivering a transgene to a CNS tissue in a subject, the method comprising: administering to a subject an effective amount of a rAAV that comprises (i) a capsid protein having a sequence as set forth in any one of SEQ ID NO: 10 to 12 and (ii) a nucleic acid comprising a promoter operably linked with a transgene.

22. The method of paragraph 21, wherein the transgene encodes a reporter protein.

23. The method of paragraph 22, wherein the reporter protein is a fluorescent protein, an enzyme that catalyzes a reaction yielding a detectable product, or a cell surface antigen.

24. The method of paragraph 23, wherein the enzyme is a luciferase, a beta-glucuronidase, a chloramphenicol acetyltransferase, an aminoglycoside phosphotransferase, an aminocyclitol phosphotransferase, or a Puromycin N-acetyltranferase.

25. The method of paragraph 21, wherein the transgene is a CNS-associated gene.

26. The method of paragraph 25, wherein the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA).

27. A rAAV that comprises (i) a capsid protein having a sequence as set forth in any one of SEQ ID NO: 10 to 12 and (ii) a nucleic acid comprising a promoter operably linked with a CNS-associated gene.

28. The rAAV of paragraph 25 or 27, wherein mRNA expressed from the CNS- associated gene comprises a miRNA binding site of a miRNA that is preferentially expressed in non-CNS tissue.

29. The rAAV of paragraph 27 or 28, wherein mRNA expressed from the CNS- associated gene does not comprise a miRNA binding site of a miRNA that is preferentially expressed in CNS tissue.

30. The rAAV of any one of paragraphs 27 to 28, wherein the promoter is a CNS tissue specific promoter.

31. The rAAV of any one of paragraphs 27 to 30, wherein the promoter is a promoter of a gene selected from: neuronal nuclei (NeuN), glial fibrillary acidic protein (GFAP), adenomatous polyposis coli (APC), and ionized calcium-binding adapter molecule 1 (Iba-1).

32. The rAAV of any one of paragraphs 27 to 31, wherein the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA).

33. A composition comprising the rAAV of any one of paragraphs 27 to 32.

34. The composition of paragraph 33 further comprising a pharmaceutically acceptable carrier.

35. A kit comprising a container housing the composition of paragraph 33 or 34.

5 36. The kit of paragraph 35, wherein the container is a sealed vial or ampule.

37. The kit of paragraph 35, wherein the container is a syringe.

10 38. An isolated mammalian cell comprising a nucleic acid encoding a capsid protein having a sequence as set forth in SEQ ID NO: 10 to 12 and a rAAV vector comprising a nucleic acid encoding a CNS-disease associated gene.

15 39. The isolated mammalian cell of paragraph 38 further comprising an AAV helper function vector.

40. The isolated mammalian cell of paragraph 38 or 39 further comprising an accessory function vector.

15 41. The isolated mammalian cell of any one of paragraphs 38 to 40, wherein the CNS-associated gene is neuronal apoptosis inhibitory protein (NAIP), nerve growth factor (NGF), glial-derived growth factor (GDNF), brain-derived growth factor (BDNF), ciliary neurotrophic factor (CNTF), tyrosine hydroxylase (TH), GTP-cyclohydrolase (GTPCH), amino acid decarboxylase (AADC) or aspartoacylase (ASPA).

20 42. A method for treating Canavan disease in a subject in need thereof, the method comprising: administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises (i) a capsid protein other than a capsid protein of AAV serotype 2 and (ii) a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA).

25 43. A method for treating Canavan disease in a subject in need thereof, the method comprising: administering an effective amount of a rAAV to CNS tissue of the subject by a route other than intracerebral administration, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA).

30 44. A method for treating Canavan disease in a subject in need thereof, the method comprising: administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA), and

35 evaluating kidney function in the subject at least once after the administration.

45. A method for treating Canavan disease in a subject in need thereof, the method comprising: administering an effective amount of a rAAV to CNS tissue of the subject, wherein the rAAV comprises a nucleic acid comprising a promoter operably linked with a region encoding aspartoacylase (ASPA), and

40 evaluating vision of the subject at least once after the administration.

45 46. The method of any one of paragraphs 42 to 45, wherein the rAAV has a capsid protein having an amino acid sequence as set forth in SEQ ID NO: 8 or 9.

47. The method of any one of paragraphs 44 to 46, wherein administering is performed intrathecally or intracerebrally.

50 48. The method of any one of paragraphs 42 to 46, wherein administering is performed intravascularly.

49. The method of paragraph 48, wherein the nucleic acid expresses an aspartoacylase (ASPA) mRNA comprising one or more miRNA binds sites for one or more miRNAs that are more abundant in one or more non-CNS tissues in comparison to a CNS tissue.

55 50. The method of paragraph 49, wherein the one or more miRNAs that are more abundant in one or more non-CNS tissues in comparison to the CNS tissue are at least twofold more abundant.

51. The method of paragraph 49 or 50, wherein the one or more non-CNS tissue is not kidney tissue or retinal tissue.

52. A method for delivering a transgene to central nervous system (CNS) tissue in a subject, the method comprising:

5 administering an effective amount of a rAAV comprising a transgene to a subject, wherein the rAAV comprises a capsid protein of a AAV serotype selected from the group consisting of: AAV1, AAV2, AAV5, AAV6, AAV6.2, AAV7, AAV8, AAV9, rh.10, rh.39, rh.43 and CSp3, and

10 wherein:

15 (a) if the AAV serotype is AAV1, the administration route is not intracerebral, intramuscular, intranerve, or intraventricular and/or the subject is not a mouse, rat or feline;

20 (b) if the AAV serotype is AAV2, the administration route is not intracerebral or intraventricular administration and/or the subject is not a rat, mouse, feline, marmoset, or macaque;

25 (c) if the AAV serotype is AAV5, the administration route is not intracerebral or intraventricular administration and/or the subject is not a rat, mouse, or marmoset;

30 (d) if the AAV serotype is AAV6, the subject is not a mouse;

35 (e) if the AAV serotype is AAV7, the administration route is not intracerebral administration and/or the subject is not a mouse or macaque;

40 (f) if the AAV serotype is AAV8, the administration route is not intracerebral, intraperitoneal, or intravascular administration and/or the subject is not a mouse or macaque;

45 (g) if the AAV serotype is AAV9, the administration route is not intracerebral or intravascular administration and/or the subject is not a rat or mouse; and (h) if the AAV serotype is AAVrh.10, the administration route is not intracerebral or intravascular administration and/or the subject is not a rat or mouse.

53. The method of paragraph 52, wherein the AAV serotype is selected from AAV1, AAV6, AAV7, rh.39, rh.43, and CSp3, and the administration route is intravascular administration.

54. The method of paragraph 52, wherein the AAV serotype is AAV7 and the administration route is intravascular administration.

55. The method of paragraph 52, wherein the CNS tissue is selected from cortex, hippocampus, thalamus, hypothalamus, cerebellum, brain stem, cervical spinal cord, thoracic spinal cord, and lumbar spinal cord.

40 [0164] It should be understood that various changes and modifications to the presently preferred embodiments described herein will be apparent to those skilled in the art. Such changes and modifications can be made without departing from the spirit and scope of the present invention and without diminishing its intended advantages. It is therefore intended that such changes and modifications be covered by the following paragraphs.

45

50

55

SEQUENCE LISTING

<110> University of Massachusetts
5 <120> CNS TARGETING AAV VECTORS AND METHODS OF USE THEREOF
<130> WOLBR/P52496EPdiv3
10 <150> 61/327,627
<151> 2010-04-23
<160> 64
<170> PatentIn version 3.5
15 <210> 1
<211> 736
<212> PRT
<213> Adeno associated virus 1
<400> 1
20 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15
25 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
20 25 30
30 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro
35 40 45
35 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60
40 Val Asn Ala Ala Asp Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80
45 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
85 90 95
50 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110
55 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
115 120 125
Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140
Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly
145 150 155 160
55 Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr

EP 3 567 106 A1

	165	170	175
5	Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190		
10	Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205		
15	Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220		
20	Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240		
25	Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255		
30	Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270		
35	Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285		
40	His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300		
45	Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315 320		
50	Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335		
55	Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350		
60	Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365		
65	Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380		
70	Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400		
75	Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe 405 410 415		

EP 3 567 106 A1

Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp
420 425 430

5 Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg
435 440 445

10 Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser
450 455 460

Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro
465 470 475 480

15 Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn
485 490 495

20 Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn
500 505 510

25 Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys
515 520 525

Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly
530 535 540

30 Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile
545 550 555 560

35 Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg
565 570 575

40 Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala
580 585 590

45 Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln
595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

50 Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
625 630 635 640

Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

55 Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr
660 665 670

EP 3 567 106 A1

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685

5 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn
 690 695 700

10 Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu
 705 710 715 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
 725 730 735

15 <210> 2
 <211> 735
 <212> PRT
 <213> Adeno associated virus 2
 20 <400> 2

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
 1 5 10 15

25 Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
 20 25 30

30 Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
 35 40 45

35 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60

40 Val Asn Glu Ala Asp Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65 70 75 80

45 Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95

50 Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
 100 105 110

55 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
 115 120 125

Leu Gly Leu Val Glu Glu Pro Val Lys Thr Ala Pro Gly Lys Lys Arg
 130 135 140

55 Pro Val Glu His Ser Pro Val Glu Pro Asp Ser Ser Ser Gly Thr Gly
 145 150 155 160

EP 3 567 106 A1

Lys Ala Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr
 165 170 175

5 Gly Asp Ala Asp Ser Val Pro Asp Pro Gln Pro Leu Gly Gln Pro Pro
 180 185 190

10 Ala Ala Pro Ser Gly Leu Gly Thr Asn Thr Met Ala Thr Gly Ser Gly
 195 200 205

Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser
 210 215 220

15 Ser Gly Asn Trp His Cys Asp Ser Thr Trp Met Gly Asp Arg Val Ile
 225 230 235 240

20 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
 245 250 255

25 Tyr Lys Gln Ile Ser Ser Gln Ser Gly Ala Ser Asn Asp Asn His Tyr
 260 265 270

Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe His
 275 280 285

30 Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn Trp
 290 295 300

35 Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln Val
 305 310 315 320

40 Lys Glu Val Thr Gln Asn Asp Gly Thr Thr Thr Ile Ala Asn Asn Leu
 325 330 335

45 Thr Ser Thr Val Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu Pro Tyr
 340 345 350

50 Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala Asp
 355 360 365

55 Val Phe Met Val Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly Ser
 370 375 380

Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro Ser
 385 390 395 400

Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe Glu
 405 410 415

EP 3 567 106 A1

Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg
420 425 430

5 Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser Arg Thr
435 440 445

10 Asn Thr Pro Ser Gly Thr Thr Gln Ser Arg Leu Gln Phe Ser Gln
450 455 460

15 Ala Gly Ala Ser Asp Ile Arg Asp Gln Ser Arg Asn Trp Leu Pro Gly
465 470 475 480

Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Ser Ala Asp Asn Asn
485 490 495

20 Asn Ser Glu Tyr Ser Trp Thr Gly Ala Thr Lys Tyr His Leu Asn Gly
500 505 510

25 Arg Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His Lys Asp
515 520 525

Asp Glu Glu Lys Phe Pro Gln Ser Gly Val Leu Ile Phe Gly Lys
530 535 540

30 Gln Gly Ser Glu Lys Thr Asn Val Asp Ile Glu Lys Val Met Ile Thr
545 550 555 560

35 Asp Glu Glu Glu Ile Arg Thr Thr Asn Pro Val Ala Thr Glu Gln Tyr
565 570 575

40 Gly Ser Val Ser Thr Asn Leu Gln Arg Gly Asn Arg Gln Ala Ala Thr
580 585 590

Ala Asp Val Asn Thr Gln Gly Val Leu Pro Gly Met Val Trp Gln Asp
595 600 605

45 Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His Thr
610 615 620

50 Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu Lys
625 630 635 640

His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala Asn
645 650 655

55 Pro Ser Thr Thr Phe Ser Ala Ala Lys Phe Ala Ser Phe Ile Thr Gln

EP 3 567 106 A1

5	660	665	670
	Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln Lys		
	675	680	685
10	Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn Tyr		
	690	695	700
15	Asn Lys Ser Val Asn Val Asp Phe Thr Val Asp Thr Asn Gly Val Tyr		
	705	710	715
	Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu		
	725	730	735
20	<210> 3		
	<211> 724		
	<212> PRT		
	<213> Adeno associated virus 5		
	<400> 3		
25	Met Ser Phe Val Asp His Pro Pro Asp Trp Leu Glu Glu Val Gly Glu		
	1	5	10
	Gly Leu Arg Glu Phe Leu Gly Leu Glu Ala Gly Pro Pro Lys Pro Lys		
	20	25	30
30	Pro Asn Gln Gln His Gln Asp Gln Ala Arg Gly Leu Val Leu Pro Gly		
	35	40	45
35	Tyr Asn Tyr Leu Gly Pro Gly Asn Gly Leu Asp Arg Gly Glu Pro Val		
	50	55	60
40	Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp Ile Ser Tyr Asn Glu		
	65	70	75
	Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala Asp		
	85	90	95
45	Ala Glu Phe Gln Glu Lys Leu Ala Asp Asp Thr Ser Phe Gly Gly Asn		
	100	105	110
50	Leu Gly Lys Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro Phe		
	115	120	125
55	Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Thr Gly Lys Arg Ile		
	130	135	140
	Asp Asp His Phe Pro Lys Arg Lys Lys Ala Arg Thr Glu Glu Asp Ser		

EP 3 567 106 A1

145	150	155	160
5			
Lys Pro Ser Thr Ser Ser Asp Ala Glu Ala Gly Pro Ser Gly Ser Gln 165 170 175			
10			
Gln Leu Gln Ile Pro Ala Gln Pro Ala Ser Ser Leu Gly Ala Asp Thr 180 185 190			
15			
Met Ser Ala Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly Ala 195 200 205			
20			
Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr Trp 210 215 220			
25			
Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu Pro 225 230 235 240			
30			
Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val Asp 245 250 255			
35			
Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr 260 265 270			
40			
Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp Gln 275 280 285			
45			
Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg Val 290 295 300			
50			
Lys Ile Phe Asn Ile Gln Val Lys Glu Val Thr Val Gln Asp Ser Thr 305 310 315 320			
55			
Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp 325 330 335			
Leu Pro Ala Phe Pro Pro Gln Val Phe Thr Leu Pro Gln Tyr Gly Tyr 355 360 365			
Ala Thr Leu Asn Arg Asp Asn Thr Glu Asn Pro Thr Glu Arg Ser Ser 370 375 380 385			
Phe Phe Cys Leu Glu Tyr Phe Pro Ser Lys Met Leu Arg Thr Gly Asn 390 395 400			

EP 3 567 106 A1

Asn Phe Glu Phe Thr Tyr Asn Phe Glu Glu Val Pro Phe His Ser Ser
405 410 415

5 Phe Ala Pro Ser Gln Asn Leu Phe Lys Leu Ala Asn Pro Leu Val Asp
420 425 430

10 Gln Tyr Leu Tyr Arg Phe Val Ser Thr Asn Asn Thr Gly Gly Val Gln
435 440 445

Phe Asn Lys Asn Leu Ala Gly Arg Tyr Ala Asn Thr Tyr Lys Asn Trp
450 455 460

15 Phe Pro Gly Pro Met Gly Arg Thr Gln Gly Trp Asn Leu Gly Ser Gly
465 470 475 480

20 Val Asn Arg Ala Ser Val Ser Ala Phe Ala Thr Thr Asn Arg Met Glu
485 490 495

25 Leu Glu Gly Ala Ser Tyr Gln Val Pro Pro Gln Pro Asn Gly Met Thr
500 505 510

Asn Asn Leu Gln Gly Ser Asn Thr Tyr Ala Leu Glu Asn Thr Met Ile
515 520 525

30 Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu Glu
530 535 540

35 Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn Arg
545 550 555 560

Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser Ser
565 570 575

40 Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val Pro
580 585 590

45 Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp
595 600 605

50 Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala Met
610 615 620

Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys Asn
625 630 635 640

55 Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val Ser
645 650 655

EP 3 567 106 A1

Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met Glu
660 665 670

5 Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln
675 680 685

10 Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro Asp
690 695 700

Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr Leu
705 710 715 720

15 Thr Arg Pro Leu

20 <210> 4
<211> 736
<212> PRT
<213> Adeno associated virus 6

25 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

30 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
20 25 30

35 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro
35 40 45

40 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

45 Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

50 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
85 90 95

55 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
115 120 125

55 Phe Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

EP 3 567 106 A1

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly
145 150 155 160

5 Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

10 Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro
180 185 190

Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly
195 200 205

15 Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala
210 215 220

20 Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile
225 230 235 240

25 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245 250 255

Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His
260 265 270

30 Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe
275 280 285

35 His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn
290 295 300

Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln
305 310 315 320

40 Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn
325 330 335

45 Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro
340 345 350

50 Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala
355 360 365

Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly
370 375 380

55 Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro
385 390 395 400

EP 3 567 106 A1

Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe
405 410 415

5 Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp
420 425 430

10 Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg
435 440 445

15 Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser
450 455 460

20 Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn
485 490 495

25 Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn
500 505 510

30 Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys
515 520 525

35 Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile
545 550 555 560

40 Thr Asp Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg
565 570 575

45 Phe Gly Thr Val Ala Val Asn Leu Gln Ser Ser Ser Thr Asp Pro Ala
580 585 590

50 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

55 Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala

EP 3 567 106 A1

645

650

655

5 Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr
660 665 670

10 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

15 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn
690 695 700

20 Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu
705 710 715 720

25 Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

30 <210> 5
<211> 736
<212> PRT
<213> Adeno associated virus 6.2

35 <400> 5

40 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

45 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
20 25 30

50 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro
35 40 45

55 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

60 Val Asn Ala Ala Asp Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

65 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
85 90 95

70 Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

75 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
115 120 125

80 Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Arg

EP 3 567 106 A1

	130	135	140
5	Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly 145 150 155 160		
	Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175		
10	Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190		
15	Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205		
20	Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220		
	Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240		
25	Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255		
30	Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270		
	Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285		
35	His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300		
40	Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315 320		
45	Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335		
	Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350		
50	Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365		
55	Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380		

EP 3 567 106 A1

Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro
385 390 395 400

5 Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe
405 410 415

10 Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp
420 425 430

Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg
435 440 445

15 Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser
450 455 460

20 Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro
465 470 475 480

25 Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn
485 490 495

Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn
500 505 510

30 Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys
515 520 525

35 Asp Asp Lys Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly
530 535 540

40 Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile
545 550 555 560

45 Phe Gly Thr Val Ala Val Asn Leu Gln Ser Ser Thr Asp Pro Ala
580 585 590

50 Thr Gly Asp Val His Val Met Gly Ala Leu Pro Gly Met Val Trp Gln
595 600 605

55 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
625 630 635 640

EP 3 567 106 A1

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

5 Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr
660 665 670

10 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn
690 695 700

15 Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu
705 710 715 720

20 Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

25 <210> 6

<211> 737

<212> PRT

<213> Adeno associated virus 7

30 <400> 6

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

35 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro
20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asn Gly Arg Gly Leu Val Leu Pro
35 40 45

40 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

45 Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

50 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

55 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
115 120 125

EP 3 567 106 A1

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Ala Lys Lys Arg
 130 135 140

5 Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile
 145 150 155 160

10 Gly Lys Lys Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln
 165 170 175

15 Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro
 180 185 190

20 Pro Ala Ala Pro Ser Ser Val Gly Ser Gly Thr Val Ala Ala Gly Gly
 195 200 205

25 Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn
 210 215 220

30 Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val
 225 230 235 240

35 Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His
 245 250 255

40 Leu Tyr Lys Gln Ile Ser Ser Glu Thr Ala Gly Ser Thr Asn Asp Asn
 260 265 270

45 Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285

50 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300

Asn Trp Gly Phe Arg Pro Lys Lys Leu Arg Phe Lys Leu Phe Asn Ile
 305 310 315 320

Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn
 325 330 335

Asn Leu Thr Ser Thr Ile Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu
 340 345 350

55 Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro
 355 360 365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn
 370 375 380

EP 3 567 106 A1

Gly Ser Gln Ser Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400

5 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Glu Phe Ser Tyr Ser
 405 410 415

10 Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430

15 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ala
 435 440 445

20 Arg Thr Gln Ser Asn Pro Gly Gly Thr Ala Gly Asn Arg Glu Leu Gln
 450 455 460

25 Phe Tyr Gln Gly Gly Pro Ser Thr Met Ala Glu Gln Ala Lys Asn Trp
 465 470 475 480

30 Leu Pro Gly Pro Cys Phe Arg Gln Gln Arg Val Ser Lys Thr Leu Asp
 485 490 495

35 Gln Asn Asn Asn Ser Asn Phe Ala Trp Thr Gly Ala Thr Lys Tyr His
 500 505 510

40 Leu Asn Gly Arg Asn Ser Leu Val Asn Pro Gly Val Ala Met Ala Thr
 515 520 525

45 His Lys Asp Asp Glu Asp Arg Phe Phe Pro Ser Ser Gly Val Leu Ile
 530 535 540

50 Phe Gly Lys Thr Gly Ala Thr Asn Lys Thr Thr Leu Glu Asn Val Leu
 545 550 555 560

55 Met Thr Asn Glu Glu Glu Ile Arg Pro Thr Asn Pro Val Ala Thr Glu
 565 570 575

Glu Tyr Gly Ile Val Ser Ser Asn Leu Gln Ala Ala Asn Thr Ala Ala
 580 585 590

60 Gln Thr Gln Val Val Asn Asn Gln Gly Ala Leu Pro Gly Met Val Trp
 595 600 605

65 Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro
 610 615 620

70 His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly

EP 3 567 106 A1

625	630	635	640
Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro			
5	645	650	655
Ala Asn Pro Pro Glu Val Phe Thr Pro Ala Lys Phe Ala Ser Phe Ile			
	660	665	670
10	Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu		
	675	680	685
15	Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser		
	690	695	700
20	Asn Phe Glu Lys Gln Thr Gly Val Asp Phe Ala Val Asp Ser Gln Gly		
	705	710	715
	720		
	Val Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn		
	725	730	735
25	Leu		
	<210> 7		
30	<211> 738		
	<212> PRT		
	<213> Adeno associated virus 8		
	<400> 7		
35	Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser		
	1	5	10
	15		
40	Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Lys Pro		
	20	25	30
	35		
	40		
45	Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro		
	35	40	45
	50		
	55		
50	Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro		
	50	55	60
	65		
	70		
	75		
	80		
55	Val Asn Ala Ala Asp Ala Ala Leu Glu His Asp Lys Ala Tyr Asp		
	85	90	95
	Gln Gln Leu Gln Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala		
	Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly		

EP 3 567 106 A1

5	Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125	100 105 110
10	Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140	
15	Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile 145 150 155 160	
20	Gly Lys Lys Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln 165 170 175	
25	Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro 180 185 190	
30	Pro Ala Ala Pro Ser Gly Val Gly Pro Asn Thr Met Ala Ala Gly Gly 195 200 205	
35	Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser 210 215 220	
40	Ser Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val 225 230 235 240	
45	Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His 245 250 255	
50	Leu Tyr Lys Gln Ile Ser Asn Gly Thr Ser Gly Gly Ala Thr Asn Asp 260 265 270	
55	Asn Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn 275 280 285	
60	Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn 290 295 300	
65	Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Ser Phe Lys Leu Phe Asn 305 310 315 320	
70	Ile Gln Val Lys Glu Val Thr Gln Asn Glu Gly Thr Lys Thr Ile Ala 325 330 335	
75	Asn Asn Leu Thr Ser Thr Ile Gln Val Phe Thr Asp Ser Glu Tyr Gln 340 345 350	

EP 3 567 106 A1

Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe
 355 360 365

5 Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn
 370 375 380

10 Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr
 385 390 395 400

Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Thr Tyr
 405 410 415

15 Thr Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser
 420 425 430

20 Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu
 435 440 445

Ser Arg Thr Gln Thr Thr Gly Gly Thr Ala Asn Thr Gln Thr Leu Gly
 450 455 460

25 Phe Ser Gln Gly Gly Pro Asn Thr Met Ala Asn Gln Ala Lys Asn Trp
 465 470 475 480

30 Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Thr Thr Thr Gly
 485 490 495

35 Gln Asn Asn Asn Ser Asn Phe Ala Trp Thr Ala Gly Thr Lys Tyr His
 500 505 510

40 Leu Asn Gly Arg Asn Ser Leu Ala Asn Pro Gly Ile Ala Met Ala Thr
 515 520 525

45 His Lys Asp Asp Glu Glu Arg Phe Phe Pro Ser Asn Gly Ile Leu Ile
 530 535 540

50 Phe Gly Lys Gln Asn Ala Ala Arg Asp Asn Ala Asp Tyr Ser Asp Val
 545 550 555 560

55 Met Leu Thr Ser Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr
 565 570 575

Glu Glu Tyr Gly Ile Val Ala Asp Asn Leu Gln Gln Asn Thr Ala
 580 585 590

55 Pro Gln Ile Gly Thr Val Asn Ser Gln Gly Ala Leu Pro Gly Met Val
 595 600 605

EP 3 567 106 A1

Trp Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile
610 615 620

5 Pro His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe
625 630 635 640

10 Gly Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val
645 650 655

15 Pro Ala Asp Pro Pro Thr Thr Phe Asn Gln Ser Lys Leu Asn Ser Phe
660 665 670

20 Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu
675 680 685

25 Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr
690 695 700

30 Ser Asn Tyr Tyr Lys Ser Thr Ser Val Asp Phe Ala Val Asn Thr Glu
705 710 715 720

35 Asn Leu

<210> 8
<211> 736
<212> PRT
<213> Adeno associated virus 9

<400> 8

40 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

45 Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

50 Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

55 Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

60 Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

EP 3 567 106 A1

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

5 Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

10 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

15 Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

20 Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

25 Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Met Ala Ser Gly Gly Gly
195 200 205

30 Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210 215 220

35 Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
225 230 235 240

40 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245 250 255

45 Tyr Lys Gln Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260 265 270

50 Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275 280 285

Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290 295 300

55 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
305 310 315 320

Gln Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Ala Asn
325 330 335

EP 3 567 106 A1

Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
340 345 350

Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
355 360 365

Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
370 375 380

Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385 390 395 400

Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
405 410 415

Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
420 425 430

Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
435 440 445

Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
450 455 460

Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
465 470 475 480

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Gln Asn
485 490 495

Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
500 505 510

Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys
515 520 525

Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
530 535 540

Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
545 550 555 560

Thr Asn Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
565 570 575

Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln Ala Gln

EP 3 567 106 A1

	580	585	590
5	Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln 595 600 605		
10	Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 615 620		
15	Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Met 625 630 635 640		
20	Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655		
25	Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr 660 665 670		
30	Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685		
35	Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn 690 695 700		
40	Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val 705 710 715 720		
45	Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu 725 730 735		
50	<210> 9 <211> 738 <212> PRT <213> Adeno associated virus rh.10		
55	<400> 9 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser 1 5 10 15 Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60 Val Asn Ala Ala Asp Ala Ala Leu Glu His Asp Lys Ala Tyr Asp		

EP 3 567 106 A1

65	70	75	80
Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala			
85		90	95
Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly			
100		105	110
Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro			
115		120	125
Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg			
130		135	140
Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile			
145		150	155
Gly Lys Lys Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln			
165		170	175
Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro			
180		185	190
Pro Ala Gly Pro Ser Gly Leu Gly Ser Gly Thr Met Ala Ala Gly Gly			
195		200	205
Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser			
210		215	220
Ser Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val			
225		230	235
Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His			
245		250	255
Leu Tyr Lys Gln Ile Ser Asn Gly Thr Ser Gly Gly Ser Thr Asn Asp			
260		265	270
Asn Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn			
275		280	285
Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn			
290		295	300
Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn			
305		310	315

EP 3 567 106 A1

Ile Gln Val Lys Glu Val Thr Gln Asn Glu Gly Thr Lys Thr Ile Ala
325 330 335

5 Asn Asn Leu Thr Ser Thr Ile Gln Val Phe Thr Asp Ser Glu Tyr Gln
340 345 350

10 Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe
355 360 365

15 Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn
370 375 380

20 Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr
385 390 395 400

25 Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Glu Phe Ser Tyr
405 410 415

30 Gln Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser
420 425 430

35 Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu
435 440 445

40 Ser Arg Thr Gln Ser Thr Gly Gly Thr Ala Gly Thr Gln Gln Leu Leu
450 455 460

45 Phe Ser Gln Ala Gly Pro Asn Asn Met Ser Ala Gln Ala Lys Asn Trp
465 470 475 480

50 Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Thr Thr Leu Ser
485 490 495

55 Gln Asn Asn Asn Ser Asn Phe Ala Trp Thr Gly Ala Thr Lys Tyr His
500 505 510

60 Leu Asn Gly Arg Asp Ser Leu Val Asn Pro Gly Val Ala Met Ala Thr
515 520 525

65 His Lys Asp Asp Glu Glu Arg Phe Phe Pro Ser Ser Gly Val Leu Met
530 535 540

70 Phe Gly Lys Gln Gly Ala Gly Lys Asp Asn Val Asp Tyr Ser Ser Val
545 550 555 560

75 Met Leu Thr Ser Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr
565 570 575

EP 3 567 106 A1

Glu Gln Tyr Gly Val Val Ala Asp Asn Leu Gln Gln Asn Ala Ala
580 585 590

5 Pro Ile Val Gly Ala Val Asn Ser Gln Gly Ala Leu Pro Gly Met Val
595 600 605

10 Trp Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile
610 615 620

15 Pro His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe
625 630 635 640

20 Gly Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val
645 650 655

25 Pro Ala Asp Pro Pro Thr Thr Phe Ser Gln Ala Lys Leu Ala Ser Phe
660 665 670

30 Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu
675 680 685

35 Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr
690 695 700

40 Ser Asn Tyr Tyr Lys Ser Thr Asn Val Asp Phe Ala Val Asn Thr Asp
705 710 715 720

45 Gly Thr Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg
725 730 735

Asn Leu

50

<210> 10
<211> 738
<212> PRT
<213> Adeno associated virus rh.39
<400> 10

55 Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Lys Pro
20 25 30

55 Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro
35 40 45

EP 3 567 106 A1

5	Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro	50	55	60	
	Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp	65	70	75	80
10	Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala	85	90	95	
	Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly	100	105	110	
15	Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro	115	120	125	
20	Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg	130	135	140	
25	Pro Val Glu Pro Ser Pro Gln Arg Ser Pro Asp Ser Ser Thr Gly Ile	145	150	155	160
	Gly Lys Lys Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln	165	170	175	
30	Thr Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro	180	185	190	
35	Pro Ala Gly Pro Ser Gly Leu Gly Ser Gly Thr Met Ala Ala Gly Gly	195	200	205	
	Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser	210	215	220	
40	Ser Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val	225	230	235	240
45	Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His	245	250	255	
50	Leu Tyr Lys Gln Ile Ser Asn Gly Thr Ser Gly Ser Thr Asn Asp	260	265	270	
	Asn Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn	275	280	285	
55	Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn	290	295	300	

EP 3 567 106 A1

Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Ser Phe Lys Leu Phe Asn
305 310 315 320

Ile Gln Val Lys Glu Val Thr Gln Asn Glu Gly Thr Lys Thr Ile Ala
5 325 330 335

Asn Asn Leu Thr Ser Thr Ile Gln Val Phe Thr Asp Ser Glu Tyr Gln
10 340 345 350

Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe
355 360 365

Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn
15 370 375 380

Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr
20 385 390 395 400

Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Glu Phe Ser Tyr
25 405 410 415

Thr Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser
420 425 430

Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu
30 435 440 445

Ser Arg Thr Gln Ser Thr Gly Gly Thr Gln Gly Thr Gln Gln Leu Leu
35 450 455 460

Phe Ser Gln Ala Gly Pro Ala Asn Met Ser Ala Gln Ala Lys Asn Trp
40 465 470 475 480

Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Thr Thr Leu Ser
485 490 495

Gln Asn Asn Asn Ser Asn Phe Ala Trp Thr Gly Ala Thr Lys Tyr His
45 500 505 510

Leu Asn Gly Arg Asp Ser Leu Val Asn Pro Gly Val Ala Met Ala Thr
50 515 520 525

His Lys Asp Asp Glu Glu Arg Phe Phe Pro Ser Ser Gly Val Leu Met
55 530 535 540

Phe Gly Lys Gln Gly Ala Gly Arg Asp Asn Val Asp Tyr Ser Ser Val

EP 3 567 106 A1

545	550	555	560
Met Leu Thr Ser Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr			
565	570	575	
Glu Gln Tyr Gly Val Val Ala Asp Asn Leu Gln Gln Thr Asn Thr Gly			
580	585	590	
Pro Ile Val Gly Asn Val Asn Ser Gln Gly Ala Leu Pro Gly Met Val			
595	600	605	
Trp Gln Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile			
610	615	620	
Pro His Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe			
625	630	635	640
Gly Leu Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val			
645	650	655	
Pro Ala Asp Pro Pro Thr Thr Phe Ser Gln Ala Lys Leu Ala Ser Phe			
660	665	670	
Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu			
675	680	685	
Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr			
690	695	700	
Ser Asn Tyr Tyr Lys Ser Thr Asn Val Asp Phe Ala Val Asn Thr Glu			
705	710	715	720
Gly Thr Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg			
725	730	735	
Asn Leu			
<210> 11			
<211> 736			
<212> PRT			
<213> Adeno associated virus rh.43			
<400> 11			
Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser			
1	5	10	15
Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro			

EP 3 567 106 A1

20	25	30		
5	Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro			
	35	40	45	
10	Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro			
	50	55	60	
15	Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp			
	65	70	75	80
20	Gln Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala			
	85	90	95	
25	Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly			
	100	105	110	
30	Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro			
	115	120	125	
35	Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg			
	130	135	140	
40	Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly			
	145	150	155	160
45	Lys Lys Gly Gln Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr			
	165	170	175	
50	Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro			
	180	185	190	
55	Ala Ala Pro Ser Gly Val Gly Pro Asn Thr Met Ala Ala Gly Gly Gly			
	195	200	205	
60	Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser			
	210	215	220	
65	Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile			
	225	230	235	240
70	Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu			
	245	250	255	
75	Tyr Lys Gln Ile Ser Asn Gly Thr Ser Gly Gly Ala Thr Asn Asp Asn			
	260	265	270	

EP 3 567 106 A1

Thr Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
275 280 285

5 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
290 295 300

10 Asn Trp Gly Phe Arg Pro Lys Arg Leu Ser Phe Lys Leu Phe Asn Ile
305 310 315 320

15 Gln Val Lys Glu Val Thr Gln Asn Glu Gly Thr Lys Thr Ile Ala Asn
325 330 335

20 Asn Leu Thr Ser Thr Ile Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu
340 345 350

25 Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro
355 360 365

30 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn
370 375 380

35 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
385 390 395 400

40 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Thr Tyr Thr
405 410 415

45 Phe Glu Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
420 425 430

50 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
435 440 445

55 Arg Thr Gln Thr Thr Gly Gly Thr Ala Asn Thr Gln Thr Leu Gly Phe
450 455 460

60 Ser Gln Gly Gly Pro Asn Thr Met Ala Asn Gln Ala Lys Asn Trp Leu
465 470 475 480

65 Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Thr Thr Thr Gly Gln
485 490 495

70 Asn Asn Asn Ser Asn Phe Ala Trp Thr Ala Gly Thr Lys Tyr His Leu
500 505 510

75 Asn Gly Arg Asn Ser Leu Ala Asn Pro Gly Ile Ala Met Ala Thr His
515 520 525

EP 3 567 106 A1

Lys Asp Asp Glu Glu Arg Phe Phe Pro Val Thr Gly Ser Cys Phe Trp
 530 535 540

5 Gln Gln Asn Ala Ala Arg Asp Asn Ala Asp Tyr Ser Asp Val Met Leu
 545 550 555 560

10 Thr Ser Glu Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Glu
 565 570 575

Tyr Gly Ile Val Ala Asp Asn Leu Gln Gln Asn Thr Ala Pro Gln
 580 585 590

15 Ile Gly Thr Val Asn Ser Gln Gly Ala Leu Pro Gly Met Val Trp Gln
 595 600 605

20 Asn Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620

25 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
 625 630 635 640

Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655

30 Asp Pro Pro Thr Thr Phe Asn Gln Ser Lys Leu Asn Ser Phe Ile Thr
 660 665 670

35 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
 675 680 685

40 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
 690 695 700

Tyr Tyr Lys Ser Thr Ser Val Asp Phe Ala Val Asn Thr Glu Gly Val
 705 710 715 720

45 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
 725 730 735

50 <210> 12
 <211> 736
 <212> PRT
 <213> Adeno associated virus CSp3

55 <400> 12

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1 5 10 15

EP 3 567 106 A1

Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Gln Pro
20 25 30

5 Lys Ala Asn Gln Gln His Gln Asp Asn Ala Arg Gly Leu Val Leu Pro
35 40 45

10 Gly Tyr Lys Tyr Leu Gly Pro Gly Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

15 Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

20 Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

25 Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

30 Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Leu Leu Glu Pro
115 120 125

35 Leu Gly Leu Val Glu Glu Ala Ala Lys Thr Ala Pro Gly Lys Lys Arg
130 135 140

40 Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ala Gly Ile Gly
145 150 155 160

45 Lys Ser Gly Ala Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr
165 170 175

50 Gly Asp Thr Glu Ser Val Pro Asp Pro Gln Pro Ile Gly Glu Pro Pro
180 185 190

55 Ala Ala Pro Ser Gly Val Gly Ser Leu Thr Ile Ala Ser Gly Gly Gly
195 200 205

60 Ala Pro Val Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Ser Ser
210 215 220

65 Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
225 230 235 240

70 Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
245 250 255

75 Tyr Lys Arg Ile Ser Asn Ser Thr Ser Gly Gly Ser Ser Asn Asp Asn
260 265 270

EP 3 567 106 A1

Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg
 275 280 285

5 Phe His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn
 290 295 300

10 Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile
 305 310 315 320

15 Arg Val Lys Glu Val Thr Asp Asn Asn Gly Val Lys Thr Ile Thr Asn
 325 330 335

20 Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp Ser Asp Tyr Gln Leu
 340 345 350

25 Pro Tyr Val Leu Gly Ser Ala His Glu Gly Cys Leu Pro Pro Phe Pro
 355 360 365

30 Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asp
 370 375 380

35 Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe
 385 390 395 400

40 Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Glu
 405 410 415

45 Phe Glu Asn Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu
 420 425 430

50 Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Ser
 435 440 445

55 Lys Thr Ile Asn Gly Ser Gly Gln Asn Gln Gln Thr Leu Lys Phe Ser
 450 455 460

Val Ala Gly Pro Ser Asn Met Ala Val Gln Gly Arg Asn Tyr Ile Pro
 465 470 475 480

Gly Pro Ser Tyr Arg Gln Gln Arg Val Ser Thr Thr Val Thr Arg Asn
 485 490 495

Asn Asn Ser Glu Phe Ala Trp Pro Gly Ala Ser Ser Trp Ala Leu Asn
 500 505 510

Gly Arg Asn Ser Leu Met Asn Pro Gly Pro Ala Met Ala Ser His Lys

EP 3 567 106 A1

515

520

525

5 Glu Gly Glu Asp Arg Phe Phe Pro Leu Ser Gly Ser Leu Ile Phe Gly
530 535 540

10 Lys Gln Gly Thr Gly Arg Asp Asn Val Asp Ala Asp Lys Val Met Ile
545 550 555 560

15 Thr Asn Glu Glu Ile Lys Thr Thr Asn Pro Val Ala Thr Glu Ser
565 570 575

20 Tyr Gly Gln Val Ala Thr Asn His Gln Ser Ala Gln Ala Gln
580 585 590

25 Thr Gly Trp Val Gln Asn Gln Gly Ile Leu Pro Gly Met Val Trp Gln
595 600 605

30 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
610 615 620

35 Thr Asp Gly Asn Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Val
625 630 635 640

40 Lys His Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala
645 650 655

45 Asp Pro Pro Thr Ala Phe Asn Lys Asp Lys Leu Asn Ser Phe Ile Thr
660 665 670

50 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln
675 680 685

55 Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn
690 695 700

60 Tyr Tyr Lys Ser Asn Asn Val Glu Phe Ala Val Asn Thr Glu Gly Val
705 710 715 720

65 Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu
725 730 735

70 <210> 13
<211> 313
<212> PRT
<213> Homo sapiens

75 <400> 13

80 Met Thr Ser Cys His Ile Ala Glu Glu His Ile Gln Lys Val Ala Ile

EP 3 567 106 A1

1	5	10	15
Phe Gly Gly Thr His Gly Asn Glu Leu Thr Gly Val Phe Leu Val Lys			
5	20	25	30
His Trp Leu Glu Asn Gly Ala Glu Ile Gln Arg Thr Gly Leu Glu Val			
10	35	40	45
Lys Pro Phe Ile Thr Asn Pro Arg Ala Val Lys Lys Cys Thr Arg Tyr			
15	50	55	60
Ile Asp Cys Asp Leu Asn Arg Ile Phe Asp Leu Glu Asn Leu Gly Lys			
20	65	70	75
Lys Met Ser Glu Asp Leu Pro Tyr Glu Val Arg Arg Ala Gln Glu Ile			
25	85	90	95
Asn His Leu Phe Gly Pro Lys Asp Ser Glu Asp Ser Tyr Asp Ile Ile			
30	100	105	110
Phe Asp Leu His Asn Thr Thr Ser Asn Met Gly Cys Thr Leu Ile Leu			
35	115	120	125
Glu Asp Ser Arg Asn Asn Phe Leu Ile Gln Met Phe His Tyr Ile Lys			
40	130	135	140
Thr Ser Leu Ala Pro Leu Pro Cys Tyr Val Tyr Leu Ile Glu His Pro			
45	145	150	155
160			
Ser Leu Lys Tyr Ala Thr Thr Arg Ser Ile Ala Lys Tyr Pro Val Gly			
50	165	170	175
Ile Glu Val Gly Pro Gln Pro Gln Gly Val Leu Arg Ala Asp Ile Leu			
55	180	185	190
Asp Gln Met Arg Lys Met Ile Lys His Ala Leu Asp Phe Ile His His			
60	195	200	
205			
Phe Asn Glu Gly Lys Glu Phe Pro Pro Cys Ala Ile Glu Val Tyr Lys			
65	210	215	220
Ile Ile Glu Lys Val Asp Tyr Pro Arg Asp Glu Asn Gly Glu Ile Ala			
70	225	230	235
240			
Ala Ile Ile His Pro Asn Leu Gln Asp Gln Asp Trp Lys Pro Leu His			
75	245	250	255

EP 3 567 106 A1

Pro Gly Asp Pro Met Phe Leu Thr Leu Asp Gly Lys Thr Ile Pro Leu
 260 265 270

5 Gly Gly Asp Cys Thr Val Tyr Pro Val Phe Val Asn Glu Ala Ala Tyr
 275 280 285

10 Tyr Glu Lys Lys Glu Ala Phe Ala Lys Thr Thr Lys Leu Thr Leu Asn
 290 295 300

Ala Lys Ser Ile Arg Cys Cys Leu His
 305 310

15 <210> 14
 <211> 942
 <212> DNA
 <213> Homo sapiens

20 <400> 14
 atgacttctt gtcacattgc tgaagaacat atacaaaagg ttgctatctt tggaggaacc 60
 catggaaatg agctaaccgg agtatttctg gttaagcatt ggctagagaa tggcgctgag 120
 25 attcagagaa cagggctgga ggtaaaacca tttattacta accccagagc agtgaagaag 180
 tgtaccagat atattgactg tgacctgaat cgcattttg accttgaaaa tcttggcaaa 240
 aaaaatgtcag aagatttgcc atatgaagtg agaagggctc aagaaataaa tcatttattt 300
 30 ggtccaaaag acagtgaaga ttcctatgac attattttg accttcacaa caccacctct 360
 aacatgggtt gcactcttat tcttgaggat tccaggaata acttttaat tcagatgttt 420
 cattacatta agacttctct ggctccacta ccctgctacg tttatctgat tgagcatcct 480
 35 tccctcaaatt atgcgaccac tcgttccata gccaagtatc ctgtgggtat agaagttgg 540
 cctcagccctc aaggggttct gagagctgat atcttggatc aatgagaaa aatgattaaa 600
 catgctcttg attttataca tcatttcaat gaaggaaaag aatttcctcc ctgcgccatt 660
 40 gaggtctata aaattataga gaaagttgat taccgggggg atgaaaatgg agaaattgct 720
 gctatcatcc atcctaattct gcaggatcaa gactggaaac cactgcatcc tggggatccc 780
 atgtttttaa ctcttgatgg gaagacgatc ccactggcg gagactgtac cgtgtacccc 840
 45 gtgtttgtga atgaggccgc atattacgaa aagaaagaag ctttgcaaa gacaactaaa 900
 ctaacgctca atgcaaaaag tattcgctgc tgtttacatt ag 942

50 <210> 15
 <211> 312
 <212> PRT
 <213> Mus Musculus

55 <400> 15

Met Thr Ser Cys Val Ala Lys Glu Pro Ile Lys Lys Ile Ala Ile Phe

EP 3 567 106 A1

1	5	10	15
Gly Gly Thr His Gly Asn Glu Leu Thr Gly Val Phe Leu Val Thr His			
5	20	25	30
Trp Leu Arg Asn Gly Thr Glu Val His Arg Ala Gly Leu Asp Val Lys			
	35	40	45
10	15	20	25
Pro Phe Ile Thr Asn Pro Arg Ala Val Glu Lys Cys Thr Arg Tyr Ile			
	50	55	60
Asp Cys Asp Leu Asn Arg Val Phe Asp Leu Glu Asn Leu Ser Lys Glu			
15	65	70	75
	80		
Met Ser Glu Asp Leu Pro Tyr Glu Val Arg Arg Ala Gln Glu Ile Asn			
20	85	90	95
His Leu Phe Gly Pro Lys Asn Ser Asp Asp Ala Tyr Asp Leu Val Phe			
	100	105	110
25	30	35	40
Asp Leu His Asn Thr Thr Ser Asn Met Gly Cys Thr Leu Ile Leu Glu			
	115	120	125
Asp Ser Arg Asn Asp Phe Leu Ile Gln Met Phe His Tyr Ile Lys Thr			
	130	135	140
Cys Met Ala Pro Leu Pro Cys Ser Val Tyr Leu Ile Glu His Pro Ser			
35	145	150	155
	160		
Leu Lys Tyr Ala Thr Thr Arg Ser Ile Ala Lys Tyr Pro Val Gly Ile			
	165	170	175
40	45	50	55
Glu Val Gly Pro Gln Pro His Gly Val Leu Arg Ala Asp Ile Leu Asp			
	180	185	190
Gln Met Arg Lys Met Ile Lys His Ala Leu Asp Phe Ile Gln His Phe			
	195	200	205
Asn Glu Gly Lys Glu Phe Pro Pro Cys Ser Ile Asp Val Tyr Lys Ile			
	210	215	220
Met Glu Lys Val Asp Tyr Pro Arg Asn Glu Ser Gly Asp Met Ala Ala			
	225	230	240
Val Ile His Pro Asn Leu Gln Asp Gln Asp Trp Lys Pro Leu His Pro			
	245	250	255

EP 3 567 106 A1

Gly Asp Pro Val Phe Val Ser Leu Asp Gly Lys Val Ile Pro Leu Gly
 260 265 270

5 Gly Asp Cys Thr Val Tyr Pro Val Phe Val Asn Glu Ala Ala Tyr Tyr
 275 280 285

10 Glu Lys Lys Glu Ala Phe Ala Lys Thr Thr Lys Leu Thr Leu Ser Ala
 290 295 300

Lys Ser Ile Arg Ser Thr Leu His
 305 310

15 <210> 16
 <211> 939
 <212> DNA
 <213> Mus Musculus

20 <400> 16
 atgacctctt gtgttgctaa agaacctatt aagaagattt ccatcttgg agggactcat 60
 ggaaatgaac tgaccggagt gtttctagtt actcactggc taaggaatgg cactgaagtt 120
 25 cacagagcag ggctggacgt gaagccattc attaccaatc caagggcggt ggagaagtgc 180
 accagataca ttgactgtga cctgaatcgt gttttgacc ttgaaaatct tagcaaagag 240
 atgtctgaag acttgcata tgaagtgaga agggctcaag aaataaatca tttatgggt 300
 30 ccaaaaaata gtgatgatgc ctatgacattt gttttgacc ttcacaacac cacttcta 360
 atgggttgca ctcttattct tgaggattcc aggaatgact ttttaattca gatgtttcac 420
 tataattaaga cttgcattggc tccattaccc tgctctgttt atctcattga gcattttca 480
 35 ctcaaatatg caaccactcg ttccatttgcc aagtatcctg ttggataga agttggtcct 540
 cagcctcactg gtgtccttag agctgatatt tttagacaaa tgagaaaaat gataaaacat 600
 gctcttgatt ttatacagca tttcaatgaa ggaaaagaat ttcctccctg ttctattgac 660
 40 gttattcattc ctaatctgca ggttcaagac tggaaaccat tgcaccctgg agatcctgtg 720
 tttgtgtctc ttgatggaaa agttattcca ctgggtggag actgtaccgt gtacccagtg 780
 45 tttgtgaatg aagctgcata ttatgaaaaaa aaagaagcat ttgcaaagac aacaaaacta 840
 acactcagcg caaaaagcat ccgctccact ttgcactaa 900
 50
 <210> 17
 <211> 465
 <212> DNA
 <213> Homo sapiens

55 <400> 17
 atggcgacga aggccgtgtg cgtgctgaag ggcgacggcc cagtgcaggg catcatcaat 60

ttcgagcaga aggaaagtaa tggaccagtg aaggtgtggg gaagcattaa aggactgact 120
 gaaggcctgc atggattcca tttcatgag tttggagata atacagcagg ctgtaccagt 180
 5 gcaggtcctc actttaatcc tctatccaga aaacacggtg ggccaaagga tgaagagagg 240
 catgttggag acttgggcaa tgtgactgct gacaaagatg gtgtggccga tgtgtctatt 300
 gaagattctg tgatctcact ctcaggagac cattgcatca ttggccgcac actgggtggc 360
 10 catgaaaaag cagatgactt gggcaaaggt ggaaatgaag aaagtacaaa gacaggaaac 420
 gctggaagtc gtttggcttg tgggttaatt gggatcgccc aataa 465

15 <210> 18
 <211> 154
 <212> PRT
 <213> Homo sapiens

20 <400> 18

25 Met Ala Thr Lys Ala Val Cys Val Leu Lys Gly Asp Gly Pro Val Gln
 1 5 10 15

Gly Ile Ile Asn Phe Glu Gln Lys Glu Ser Asn Gly Pro Val Lys Val
 20 25 30

30 Trp Gly Ser Ile Lys Gly Leu Thr Glu Gly Leu His Gly Phe His Val
 35 40 45

His Glu Phe Gly Asp Asn Thr Ala Gly Cys Thr Ser Ala Gly Pro His
 50 55 60

35 Phe Asn Pro Leu Ser Arg Lys His Gly Gly Pro Lys Asp Glu Glu Arg
 65 70 75 80

40 His Val Gly Asp Leu Gly Asn Val Thr Ala Asp Lys Asp Gly Val Ala
 85 90 95

Asp Val Ser Ile Glu Asp Ser Val Ile Ser Leu Ser Gly Asp His Cys
 100 105 110

45 Ile Ile Gly Arg Thr Leu Val Val His Glu Lys Ala Asp Asp Leu Gly
 115 120 125

50 Lys Gly Gly Asn Glu Glu Ser Thr Lys Thr Gly Asn Ala Gly Ser Arg
 130 135 140

Leu Ala Cys Gly Val Ile Gly Ile Ala Gln
 145 150

55 <210> 19

<211> 465
 <212> DNA
 <213> Mus Musculus

5 <400> 19
 atggcgatga aagcggtgtg cgtgctgaag ggcgacggtc cggtgcaggg aaccatccac 60
 ttcgagcaga aggcaagcgg tgaaccagtt gtgttgcag gacaaattac aggattaact 120
 10 gaaggccagc atgggttcca cgtccatca gatggggaca atacacaagg ctgtaccagt 180
 gcaggacctc attttaatcc tcactctaag aaacatggtg gcccggcgg a tgaagagagg 240
 catgttggag acctgggcaa tgtgactgct ggaaaggacg gtgtggccaa tgtgtccatt 300
 15 gaagatcgtg tcatctcaact ctcaggagag cattccatca ttggccgtac aatggtggc 360
 catgagaaac aagatgactt gggcaaagggt ggaaatgaag aaagtacaaa gactggaaat 420
 gctggagcc gcttggcctg tggagtgatt gggattgcgc agtaa 465

20 <210> 20
 <211> 154
 <212> PRT
 <213> Mus Musculus

25 <400> 20

Met Ala Met Lys Ala Val Cys Val Leu Lys Gly Asp Gly Pro Val Gln
 1 5 10 15

30 Gly Thr Ile His Phe Glu Gln Lys Ala Ser Gly Glu Pro Val Val Leu
 20 25 30

35 Ser Gly Gln Ile Thr Gly Leu Thr Glu Gly Gln His Gly Phe His Val
 35 40 45

40 His Gln Tyr Gly Asp Asn Thr Gln Gly Cys Thr Ser Ala Gly Pro His
 50 55 60

45 Phe Asn Pro His Ser Lys Lys His Gly Gly Pro Ala Asp Glu Glu Arg
 65 70 75 80

50 His Val Gly Asp Leu Gly Asn Val Thr Ala Gly Lys Asp Gly Val Ala
 85 90 95

55 Asn Val Ser Ile Glu Asp Arg Val Ile Ser Leu Ser Gly Glu His Ser
 100 105 110

Ile Ile Gly Arg Thr Met Val Val His Glu Lys Gln Asp Asp Leu Gly
 115 120 125

55 Lys Gly Gly Asn Glu Glu Ser Thr Lys Thr Gly Asn Ala Gly Ser Arg
 130 135 140

Leu Ala Cys Gly Val Ile Gly Ile Ala Gln
145 150

5 <210> 21
<211> 5243
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Synthetic Oligonucleotide

<400> 21 ctgcgcgctc gctcgctcac tgaggccgccc cgggcaaagc ccgggcgtcg ggacaccc 60
ggtcgccccgg cctcagttag cgagcgagcg cgccagagagg gaggtagcc atgctctagg 120
aagatcaatt caattcacgc gtcgacattt attattgact agttattaaat agtaatcaat 180
tacggggtca ttagttcata gcccataatgg gtagttccgc gttacataac ttacggtaaa 240
tggcccgccct ggctgaccgc ccaacgaccc ccgcattt acgtcaataa tgacgtatgt 300
tcccatagta acgccaatag ggactttcca ttgacgtcaa tgggtggata tttacggtaa 360
actgcccact tggcagtaca tcaagtgtat catatgccaa gtacgcccc tattgacgtc 420
aatgacggta aatggccgcg ctggcattat gcccagtaca tgaccttatgg gactttccct 480
acttggcagt acatctacgt attagtcata gctattacca tgctgaggcc acgttctgct 540
tcactctccc catctcccccc ccctcccccac ccccaatttt gtatttattt attttttaat 600
tattttgc agcgatgggg gcgggggggg ggggcgcgcg ccaggcgggg cggggcgggg 660
cgaggggcg ggccggcgaa ggcggagagg tgccggcgca gccaatcaga gccggcgct 720
ccgaaagttt cttttatgg cgaggcgccg gcggcgccgg ccctataaaa agcgaagcg 780
gcggcgccg ggagcaagct ctagcctcga gaattcacgc gtggtagctc tagagcagag 840
ctcgtttagt gaaccgtcag ttgcggatcg ccaccatggt gagcaaggc gaggagctgt 900
tcaccgggtt ggtgccatc ctggcgagc tggacggcgaa cgtaaacggc cacaagttca 960
gcgtgtccgg cgaggcgag ggcgatgcca cttacggcaa gctgaccctg aagttcatct 1020
gcaccaccgg caagctgccc gtgcctggc ccaccctcgat gaccaccctg acctacggcg 1080
tgcagtgtttt cagccgctac cccgaccaca tgaagcagca cgacttcttca aagtccggca 1140
tgcccgaaagg ctacgtccag gagcgcacca tcttcttcaa ggacgacggc aactacaaga 1200
cccgccgcca ggtgaagtgc gagggcgaca ccctggtaa ccgcacatcgag ctgaaggcg 1260
tcgacttcaa ggaggacggc aacatcctgg ggcacaaagct ggagtacaac tacaacagcc 1320
acaacgtcta tatcatggcc gacaaggcaga agaacggcat caaggtaac ttcaagatcc 1380
gccacaaacat cgaggacggc aqcgatgcgac tcgcccqacca ctaccacqaa aacacccca 1440

	tcggcgacgg ccccggtgctg ctgcccaca accactacct gagcacccag tccgcccgtga	1500
5	gcaaagaccc caacgagaag cgcgatcaca tggtcctgct ggagttcgta accgcccgcg	1560
	ggatcaactct cggcatggac gagctgtaca agtaagtaac aggttaagtgc gatcgctaat	1620
	gcgggaaagc tcttattcgg gtgagatggg ctggggcacc atctgggac cctgacgtga	1680
10	agtttgcac tgactggaga actcggttg tcgtctgttg cggggcggc agttatggcg	1740
	gtgcgcgttgg gcagtgcacc cgtacccttg ggagcgcgcg ccctcgctgt gtcgtgacgt	1800
	caccgcgttct gttggtacct gctgttgaca gtgagcgcacg caatgtgact tcgctgacaa	1860
15	agctgtgaag ccacagatgg gctttgtcag cagtcacatt gcgctgccta ctgcctcgga	1920
	cttcaagggc tcgagaattc aggggtgggc cacctgcccgg taggtgtgcg gtaggccttt	1980
	ctccgtcgca ggacgcaggg ttcgggccta gggtaggctc tcctgaatcg acaggcgcgcg	2040
20	gacctctggc ggccgcaaca acgcgttctt gaccattcat cctatcttctt tttcctgcag	2100
	gcttgtggaa gaaatggat ccgatcttt tccctctgccc aaaaattatg gggacatcat	2160
	gaagccccctt gagcatctga cttctggcta ataaaggaaa tttatttca ttgcaatagt	2220
25	gtgttggaaat tttttgtgtc ttcactcggt cctaggtaga taagtagcat ggcgggttaa	2280
	tcattaacta caaggaaccc ctatgtatgg agttggccac tccctctctg cgcgctcgat	2340
	cgctcaactga ggccggggcga ccaaaggatcg cccgacgcgc gggatttgcgc cggggcggcct	2400
30	cagtgagcga gcgagcgcgc agccttaatt aacctaattc actggccgtc gttttacaac	2460
	gtcgtgactg ggaaaaccct ggcgttaccc aacttaatcg cttgcagca catccccctt	2520
	tccgcagctg gctaatagc gaagaggccc gcaccgatcg ccattccaa cagttgcgcga	2580
35	gcctgaatgg cgaatgggac ggcgcgtgtc gcgccgcatt aagcgccggc ggtgtgggtgg	2640
	ttacgcgcag cgtgaccgct acacttgcca ggcgcctagc gcccgcctt ttcgcctttat	2700
	tcccttcctt ttcgcgcacg ttccgcggct ttccccgtca agctctaaat cgggggctcc	2760
40	ctttagggtt ccgatttatgt gcttacggc acctcgaccc caaaaaactt gattagggtg	2820
	atggttcacg tagtgggcca tccgcctgat agacggtttt tccgcctttg acgttggagt	2880
	ccacgttctt taatagtggc ctcttgcgttcc aaactggaaac aacactcaac cctatctcg	2940
45	tctattcttt tgatttataa gggatttgc cgatttcggc ctattggta aaaaatgagc	3000
	tgatTTTACA aaaaatTTAA gcgaatTTTA aaaaaatTTTAA aacgcttaca atttaggtgg	3060
	cacttttcgg ggaaatgtgc gcgaaacccc tatttggtaa tttttctaaa tacattcaaa	3120
50	tatgtatccg ctcatgagac aataaccctg ataaatgctt caataatatt gaaaaaggaa	3180
	gagtagttagt attcaacatt tccgtgtcgc ccttattccc tttttgcgg cattttgcct	3240
55	tcctgttttt gctcacccag aaacgctggtaaa gaaagtaaaa gatgctgaag atcagttggg	3300
	tgcacgagtg ggttacatcg aactggatct caacagcggtaa aagatccttg agagttttag	3360

ccccgaagaa	cgttttccaa	tgatgagcac	ttttaaagtt	ctgctatgtg	gcgcggtatt	3420	
atcccgtatt	gacgcccggc	aagagcaact	cggtcgccgc	atacactatt	ctcagaatga	3480	
5	cttggttgag	tactcaccag	tcacagaaaa	gcatcttacg	gatggcatga	cagtaagaga	3540
attatgcagt	gctgccataa	ccatgagtga	taacactgcg	gccaacttac	ttctgacaac	3600	
10	gatcgaggaga	ccgaaggago	taaccgctt	tttgcacaac	atggggatc	atgttaactcg	3660
ccttgatcgt	tgggaaccgg	agctgaatga	agccatacca	aacgacgagc	gtgacaccac	3720	
15	gatgcctgta	gcaatggcaa	caacgttgcg	caaactatta	actggcgaac	tacttactct	3780
agcttcccg	caacaattaa	tagactggat	ggaggcggat	aaagttgcag	gaccacttct	3840	
20	gcgctaggcc	cttccggctg	gctggttat	tgctgataaa	tctggagccg	gtgagcgtgg	3900
gtctcgcggt	atcattgcag	cactgggccc	agatggtaag	ccctcccgta	tcgtagttat	3960	
25	ctacacgacg	gggagtcagg	caactatgga	tgaacgaaat	agacagatcg	ctgagatagg	4020
tgcctcaactg	attaaggcatt	ggtaactgtc	agaccaagtt	tactcatata	tacttttagat	4080	
tgataaaaa	cttcattttt	aattttaaag	gatctaggtg	aagatccctt	ttgataatct	4140	
30	catgaccaaa	atcccttaac	gtgagtttc	gttccactga	gcgtcagacc	ccgtagaaaa	4200
gatcaaagga	tcttctttag	atcctttttt	tctgcgcgta	atctgctgct	tgcaaacaaa	4260	
35	aaaaccaccc	ctaccagcg	tggttgttt	gccggatcaa	gagctaccaa	ctcttttcc	4320
gaaggttaact	ggcttcagca	gagcgcagat	accaaatact	gttctctag	tgtagccgta	4380	
gttaggccac	cacttcaaga	actctgttagc	accgcctaca	tacctcgctc	tgctaattcct	4440	
40	gttaccagt	gctgctgcca	gtggcgataa	gtcgtgtctt	accgggttgg	actcaagacg	4500
atagttaccg	gataaggcgc	agcggtcggg	ctgaacgggg	ggttcgtgca	cacagcccag	4560	
45	cttggagcga	acgacacctaca	ccgaactgag	atacctacag	cgtgagctat	gagaaagcgc	4620
cacgcttccc	gaagggagaa	aggcggacag	gtatccggta	agcggcaggg	tcggaacagg	4680	
50	agagcgcacg	aggagacttc	cagggggaaa	cgcctggtat	ctttatagtc	ctgtcggggtt	4740
tcgccacctc	tgacttgagc	gtcgattttt	gtgatgctcg	tcaggggggc	ggagcctatg	4800	
55	aaaaaacgcc	agcaacgcgg	ccttttacg	gttcctggcc	ttttgctggc	cttttgctca	4860
catgttcttt	cctgcgttat	cccctgattc	tgtggataac	cgtattaccg	cctttgagtg	4920	
agctgataacc	gctcgccgca	gccgaacgac	cgagcgcagc	gagtcagtga	gcgaggaagc	4980	
55	ggaagagcgc	ccaatacgca	aaccgcctct	ccccgcgcgt	tggccgattc	attaatgcag	5040
ctggcacgac	aggtttcccg	actggaaagc	gggcagttag	cgcaacgcaa	ttaatgtgag	5100	
ttagctcaact	cattaggcac	cccaggctt	acactttatg	cttccggctc	gtatgttg	5160	
tggaatttgt	agcggataac	aatttcacac	aggaaacagc	tatgaccatg	attacgcccag	5220	

	attnaattaa ggccttaatt agg	5243
5	<210> 22	
	<211> 108	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 22	
	tgctgttgac agtgagcgac atcatcaatt ttccgagcag aactgtgaag ccacagatgg	60
	gttctgctcg aaattgatga tgctgcctac tgccctggac ttcaaggg	108
15	<210> 23	
	<211> 106	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 23	
	tgctgttgac agtgagcgac gcattaaagg atcctgactg actgtgaagc cacagatggg	60
25	tcagtcagtc cttaatgcg ctgcctactg cctcggactt caaggg	106
30	<210> 24	
	<211> 108	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 24	
	tgctgttgac agtgagcgac tgcattggatt ctccatgttc atctgtgaag ccacagatgg	60
	gatgaacatg gaatccatgc agctgcctac tgccctggac ttcaaggg	108
40	<210> 25	
	<211> 106	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 25	
	tgctgttgac agtgagcgac aaggatgaag atcgaggcat gctgtgaagc cacagatggg	60
50	catgcctctc ttcatccttg ctgcctactg cctcggactt caaggg	106
55	<210> 26	
	<211> 110	
	<212> DNA	
	<213> Artificial Sequence	

<220>
 <223> Synthetic Oligonucleotide
 <400> 26
 5 tgctgttgac agtgagcgac gcaatgtgac ttcgctgaca aagctgtgaa gccacagatg 60
 ggctttgtca gcagtcacat tgcgctgcct actgcctcgg acttcaaggg 110

10 <210> 27
 <211> 108
 <212> DNA
 <213> Artificial Sequence
 <220>
 15 <223> Synthetic Oligonucleotide
 <400> 27
 tgctgttgac agtgagcgac cgatgtgtct atcttgaaga ttctgtgaag ccacagatgg 60
 20 gaatcttcaa tagacacatc ggctgcctac tgccctcggac ttcaaggg 108

25 <210> 28
 <211> 106
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Synthetic Oligonucleotide
 30 <400> 28
 tgctgttgac agtgagcgac ggtggaaatg atcagaaaatg actgtgaagc cacagatgg 60
 tactttcttc atttccaccg ctgcctactg cctcggactt caaggg 106

35 <210> 29
 <211> 110
 <212> DNA
 <213> Artificial Sequence
 <220>
 40 <223> Synthetic Oligonucleotide
 <400> 29
 tgctgttgac agtgagcgac gctgttagaaa ttctgtatcct gatctgtgaa gccacagatg 60
 45 ggatcaggat acatttctac agcgctgcct actgcctcgg acttcaaggg 110

50 <210> 30
 <211> 106
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Synthetic Oligonucleotide
 55 <400> 30
 tgctgttgac agtgagcgag gtattaaact tgtcagaatt tagtgaagcc acagatgtaa 60

	attctgacaa gttaataacc ctgcctactg cctcggactt caaggg	106
5	<210> 31	
	<211> 106	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 31	
	tgctgttgac agtgagcgac gatgctctaa tcggttctat caagtgaagc cacagatgtt	60
	gatagaacct tagagcatcg ctgcctactg cctcggactt caaggg	106
15	<210> 32	
	<211> 28	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 32	
	cgaaatacat acttctttac attccatt	28
25	<210> 33	
	<211> 28	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 33	
	cgaatggaat gtaaagaagt atgtattt	28
35	<210> 34	
	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
40	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 34	
	cgaaacaaac accattgtca cactccatt	29
45	<210> 35	
	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 35	
	cgaatggagt gtgacaatgg tggttgttt	29

5 <210> 36
<211> 72
<212> DNA
<213> Artificial Sequence

10 <220>
<223> Synthetic Oligonucleotide

15 <400> 36
cgaaatacat acttctttac attccaatac atacttcttt acattccaat acataacttct 60
ttacattcca tt 72

20 <210> 37
<211> 72
<212> DNA
<213> Artificial Sequence

25 <220>
<223> Synthetic Oligonucleotide

30 <400> 37
cgaatggaat gtaaagaagt atgtattgga atgtaaagaa gtatgtattg gaatgtaaaag 60
aagtatgtat tt 72

35 <210> 38
<211> 75
<212> DNA
<213> Artificial Sequence

40 <220>
<223> Synthetic Oligonucleotide

45 <400> 38
cgaaacaaac accattgtca cactccaaca aacaccattg tcacactcca acaaacacca 60
ttgtcacact ccatt 75

50 <210> 39
<211> 75
<212> DNA
<213> Artificial Sequence

55 <220>
<223> Synthetic Oligonucleotide

60 <400> 39
cgaatggagt gtgacaatgg tgtttggatgg agtgtgacaa tggtgtttgt tggagtgtga 60
caatgggtt tt 75

65 <210> 40
<211> 51
<212> DNA
<213> Artificial Sequence

```

<220>
<223> Synthetic Oligonucleotide

5 <400> 40
    cgaatacacat acttctttac attccaacaa acaccattgt cacactccat t      51

    <210> 41
    <211> 51
    <212> DNA
10  <213> Artificial Sequence

    <220>
    <223> Synthetic Oligonucleotide

15 <400> 41
    cgaatggagt gtgacaatgg tggatgttgg aatgtaaaga agtatgtatt t      51

    <210> 42
    <211> 173
    <212> DNA
20  <213> Artificial Sequence

    <220>
    <223> Synthetic Oligonucleotide

25 <400> 42
    ttcaactcg agatacatac ttctttacat tccaaatcacat acttctttac attccaatac      60
        atacttcttt acattccacc atggactagt acaaacacca ttgtcacact ccaacaaaca      120
        ccattgtcac actccaacaa acaccattgt cacactccag cggccgcttc gaa      173

    <210> 43
    <211> 29
    <212> DNA
35  <213> Artificial Sequence

    <220>
    <223> Synthetic Oligonucleotide

40 <400> 43
    atcgggccccg actgcagttt cagcgttt      29

    <210> 44
    <211> 29
    <212> DNA
45  <213> Artificial Sequence

    <220>
    <223> Synthetic Oligonucleotide

50 <400> 44
    cgcgggccccg actttacatt acacacaat      29

    <210> 45
    <211> 29
    <212> DNA
55  <213> Artificial Sequence

```

<220>		
<223>	Synthetic Oligonucleotide	
5	<400> 45	29
	cgcgcccccg actgatgtgt gagagagac	
10	<210> 46	
	<211> 29	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 46	29
	cgcgcccccg actttcggcc tcccgaggc	
20	<210> 47	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
25	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 47	20
	tgaagctgaa gcctgtgatg	
30	<210> 48	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 48	20
	gagcacctga cagcattgaa	
40	<210> 49	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 49	20
	ctcagcaaca gctcatggaa	
50	<210> 50	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
55	<220>	
	<223> Synthetic Oligonucleotide	

	<400> 50	
	ttacttctgg caccacacca	20
5	<210> 51	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
10	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 51	
	tggtgtggtg ccagaagtaa	20
15	<210> 52	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 52	
	caacagatgg ctggcaacta	20
25	<210> 53	
	<211> 18	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
	<223> Synthetic Oligonucleotide	
	<400> 53	
35	tgggagtggc accttcca	18
	<210> 54	
	<211> 20	
	<212> DNA	
40	<213> Artificial Sequence	
	<220>	
	<223> Synthetic Oligonucleotide	
45	<400> 54	
	cgaccactac cagcagaaca	20
50	<210> 55	
	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
	<220>	
	<223> Synthetic Oligonucleotide	
55	<400> 55	
	cttgtacagc tcgtccatgc	20

```

<210> 56
<211> 20
<212> DNA
<213> Artificial Sequence
5
<220>
<223> Synthetic Oligonucleotide

<400> 56
aatggcctca gaatgactgc
10
20

<210> 57
<211> 20
<212> DNA
15
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 57
agtgcgttcc acagccaaat
20
20

<210> 58
<211> 22
25
<212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide
30
<400> 58
atgccaacac agtgctgtct gg
22

<210> 59
35
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
40
<223> Synthetic Oligonucleotide

<400> 59
tgcttgctga tccacatctg ct
22

<210> 60
45
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
50
<223> Synthetic Oligonucleotide

<400> 60
tggagtgtga caatggtgtt tg
22
55
<210> 61

```

```

<211> 22
<212> DNA
<213> Artificial Sequence
5
<220>
<223> Synthetic Oligonucleotide

<400> 61
aactatacaa cctactacct ca 22
10

<210> 62
<211> 22
<212> DNA
15 <213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide

<400> 62
20 agcctatcct ggattacttg aa 22

<210> 63
<211> 22
25 <212> DNA
<213> Artificial Sequence

<220>
<223> Synthetic Oligonucleotide
30 <400> 63
acagttcttc aactggcagc tt 22

<210> 64
35 <211> 26
<212> DNA
<213> Artificial Sequence

<220>
40 <223> Synthetic Oligonucleotide

<400> 64
ctctgtatcg ttccaaattt agtata 26
45

```

Claims

1. A recombinant adeno-associated virus (rAAV) comprising (i) a capsid protein of a AAV serotype and (ii) a nucleic acid comprising a transgene, wherein the transgene encodes glial-derived growth factor (GDNF), for use as a medicament; wherein the rAAV is to be delivered in an effective amount to central nervous system (CNS) tissue of a subject by intracerebral injection into the putamen of the subject.
2. The rAAV for the use of claim 1 wherein the AAV serotype is AAV2.
3. The rAAV for the use of claim 1 or claim 2 wherein the intracerebral injection includes the use of a stereotactic device to guide the injection.
4. The rAAV for the use of any one of claims 1 to 3 wherein the intracerebral injection includes the use of a catheter

to deliver the rAAV into the putamen of the subject.

5 5. The rAAV for the use of any one of claims 1 to 4 wherein the rAAV is formulated in a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

6 6. The rAAV for the use of claim 5, wherein between 100 µl to 1 ml of the pharmaceutical composition is injected into the putamen of the subject.

10 7. The rAAV for the use of any one of claims 1 to 6, wherein the effective amount of the rAAV for intracerebral injection to the subject is in a range of 1×10^{10} genome copies to 1×10^{11} genome copies.

8. The rAAV for the use of any one of claims 1 to 6, wherein the effective amount of the rAAV for intracerebral injection to the subject is in a range of 1×10^{11} genome copies to 1×10^{12} genome copies.

15 9. The rAAV for the use of any one of claims 1 to 6, wherein the effective amount of the rAAV for intracerebral injection to the subject is in a range of 1×10^{12} genome copies to 1×10^{13} genome copies.

10 10. The rAAV for the use of any one of claims 1 to 6, wherein the effective amount of the rAAV for intracerebral injection to the subject is in a range of 1×10^{13} genome copies to 1×10^{14} genome copies.

20 11. The rAAV for the use of any one of claims 1 to 10, wherein the rAAV is for use in treating Parkinson's Disease and is to be delivered in an effective amount to central nervous system (CNS) tissue of a subject afflicted with Parkinson's Disease.

25

30

35

40

45

50

55





Fig. 1-1







Fig. 2-1








Fig. 3






Fig. 5A




Fig. 5B




Fig. 5C




Fig. 5D




Fig. 6A




Fig. 6B




Fig. 7A




Fig. 7B




Fig. 8





Fig. 9






Fig. 11




Fig. 12A




Fig. 12B




Fig. 12C




Fig. 13A




Fig. 13B




Fig. 13C




Fig. 13D




Fig. 14A




Fig. 14B




Fig. 14C




Fig. 14D




Fig. 14E





Fig. 15B




Fig. 15C




Fig. 15D




Fig. 15E




Fig. 15F




Fig. 16A




Fig. 16B




Fig. 17A




Fig. 17B




Fig. 18




Fig. 19A-1




Fig. 19A-2




Fig. 19A-3




Fig. 19B




Fig. 20




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25A




Fig. 25B




Fig. 26




Fig. 27A




Fig. 27B




Fig. 28




Fig. 29




Fig. 30A




Fig. 30B




Fig. 30C




Fig. 30D





Fig. 30E






Fig. 32A




Fig. 32B




Fig. 33A




Fig. 33B




Fig. 33C




Fig. 33D



EUROPEAN SEARCH REPORT

 Application Number
 EP 18 24 8250

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
	Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim
10	Y, P	WO 2010/129021 A1 (GENZYME CORP [US]; PASSINI MARCO A [US]; SHIHABUDDIN LAMYA [US]; CHENG) 11 November 2010 (2010-11-11) * page 4, line 29 - line 31 * * page 5, line 8 - line 17; figure 2 * -----	1-11
15	Y	DOLAN SONDHI ET AL: "Enhanced Survival of the LINCL Mouse Following CLN2 Gene Transfer Using the rh.10 Rhesus Macaque-derived Adeno-associated Virus Vector", MOLECULAR THERAPY, vol. 15, no. 3, 19 December 2006 (2006-12-19), pages 481-491, XP055084809, ISSN: 1525-0016, DOI: 10.1038/sj.mt.6300049 * abstract * * figure 1 * * page 484, right-hand column, paragraph 2 * * page 488, left-hand column, paragraphs 2, 3 * * page 488, right-hand column, paragraphs 2, last * -----	1-11
20			
25			
30	Y	A S BEUTLER ET AL: "AAV for pain: steps towards clinical translation", GENE THERAPY, vol. 16, no. 4, 1 April 2009 (2009-04-01), pages 461-469, XP055070356, ISSN: 0969-7128, DOI: 10.1038/gt.2009.23 * page 464, right-hand column, last paragraph - page 466, left-hand column, paragraph 2 * * page 462, right-hand column, paragraph 3 * -----	1-11
35			
40			
45			
1	The present search report has been drawn up for all claims		
50	Place of search Munich		Date of completion of the search 13 September 2019 Examiner Rojo Romeo, Elena
	CATEGORY OF CITED DOCUMENTS		
	X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document

EPO FORM 1503 03.82 (P04C01)

55

page 1 of 3



EUROPEAN SEARCH REPORT

Application Number
EP 18 24 8250

5

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
10	Y TENENBAUM L ET AL: "Recombinant AAV-mediated gene delivery to the central nervous system.", THE JOURNAL OF GENE MEDICINE FEB 2004, vol. 6 Suppl 1, February 2004 (2004-02), pages S212-S222, XP002715192, ISSN: 1099-498X * page S216, paragraph "Neuroprotection" - page S217 * * figure 2 *	1-11	
15			
20	A MARCO A. PASSINI ET AL: "CNS-targeted gene therapy improves survival and motor function in a mouse model of spinal muscular atrophy", JOURNAL OF CLINICAL INVESTIGATION, vol. 120, no. 4, 1 April 2010 (2010-04-01), pages 1253-1264, XP055040159, ISSN: 0021-9738, DOI: 10.1172/JCI41615 * abstract * * page 1254, left-hand column, paragraph 1 * * page 1254, right-hand column, paragraph 2 *	1-11	
25			TECHNICAL FIELDS SEARCHED (IPC)
30			
35	A CEARLEY ET AL: "Transduction characteristics of adeno-associated virus vectors expressing cap serotypes 7, 8, 9, and Rh10 in the mouse brain", MOLECULAR THERAPY, NATURE PUBLISHING GROUP, GB, vol. 13, no. 3, 1 March 2006 (2006-03-01), pages 528-537, XP005326765, ISSN: 1525-0016, DOI: 10.1016/J.YMTH.2005.11.015 * Results *	1-11	
40			
45			
50	1 The present search report has been drawn up for all claims		
55			
EPO FORM 1503 03/82 (P04C01)	Place of search	Date of completion of the search	Examiner
	Munich	13 September 2019	Rojo Romeo, Elena
CATEGORY OF CITED DOCUMENTS		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document			



EUROPEAN SEARCH REPORT

Application Number
EP 18 24 8250

5

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (IPC)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
10	A CASSIA N CEARLEY ET AL: "Expanded Repertoire of AAV Vector Serotypes Mediate Unique Patterns of Transduction in Mouse Brain", MOLECULAR THERAPY, vol. 16, no. 10, 19 August 2008 (2008-08-19), pages 1710-1718, XP055022703, ISSN: 1525-0016, DOI: 10.1038/mt.2008.166 * tables 1, 2 *	1-11	
15			
20	T BROOKE R. SNYDER ET AL: "Comparison of Adeno-Associated Viral Vector Serotypes for Spinal Cord and Motor Neuron Gene Delivery", HUMAN GENE THERAPY, vol. 22, no. 9, 1 September 2011 (2011-09-01), pages 1129-1135, XP055099540, ISSN: 1043-0342, DOI: 10.1089/hum.2011.008 * the whole document *		TECHNICAL FIELDS SEARCHED (IPC)
25			
30			
35			
40			
45			
50	1 The present search report has been drawn up for all claims		
55	1 Place of search Munich	Date of completion of the search 13 September 2019	Examiner Rojo Romeo, Elena
	CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document	T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/82 (P04C01)

ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.

EP 18 24 8250

5 This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

13-09-2019

10	Patent document cited in search report	Publication date		Patent family member(s)	Publication date
15	WO 2010129021 A1 11-11-2010	BR CA CN CN DK EP EP ES HR HU JP JP JP JP JP JP JP KR KR LT MX MX PL RU RU SG SG SI US US US WO		PI1010868 A2 2759801 A1 102459611 A 107083400 A 2424991 T3 2424991 A1 3421603 A1 2686504 T3 P20181423 T1 E039345 T2 5879256 B2 6397391 B2 2012526046 A 2016052309 A 2018148927 A 20120006073 A 20180027619 A 2424991 T 341073 B 356669 B 2424991 T3 2011149094 A 2016140850 A 175409 A1 10201404528S A 2424991 T1 2013287736 A1 2016051699 A1 2017087212 A1 2010129021 A1	12-06-2018 11-11-2010 16-05-2012 22-08-2017 17-09-2018 07-03-2012 02-01-2019 18-10-2018 30-11-2018 28-12-2018 08-03-2016 26-09-2018 25-10-2012 14-04-2016 27-09-2018 17-01-2012 14-03-2018 25-09-2018 05-08-2016 08-06-2018 30-11-2018 10-06-2013 14-12-2018 29-12-2011 30-10-2014 30-11-2018 31-10-2013 25-02-2016 30-03-2017 11-11-2010
20	-----				
25					
30					
35					
40	-----				
45					
50					
55					

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61327627 B [0001]
- US 20030138772 A [0035]
- US 20070036760 A [0035]
- US 20090197338 A [0035]
- US 61182084 A [0035]
- US 5478745 A [0037]
- US 6001650 A [0038]
- US 6156303 A [0038]
- WO 9810088 A [0051]
- US 5543158 A [0060]
- US 5641515 A [0060]
- US 5399363 A [0060]
- US 5741516 A [0073]
- US 5567434 A [0073]
- US 5552157 A [0073]
- US 5565213 A [0073]
- US 5738868 A [0073]
- US 5795587 A [0073]
- US 5656016 A [0077]
- US 5779708 A [0077]
- US 5797898 A [0077]
- US 5770219 A [0077]
- US 5783208 A [0077]
- US 5697899 A [0077]

Non-patent literature cited in the description

- **LIANG Y et al.** Characterization of microRNA expression profiles in normal human tissues. *BMC Genomics*, 12 June 2007, vol. 8, 166 [0018]
- **XIA et al.** *Neurobiol Dis*, 2006, vol. 23, 578 [0030]
- **WANG et al.** *JBC*, 2008, vol. 283, 15845 [0030]
- **WU et al.** *Antiox Redox Sig*, 2009, vol. 11, 1523 [0030]
- **G. GAO et al.** *J. Virol*, June 2004, vol. 78 (12), 6381-6388 [0035]
- **G. GAO et al.** *Proc Natl Acad Sci USA*, 13 May 2003, vol. 100 (10), 6081-6086 [0035]
- **SAMBROOK et al.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Press [0037]
- **K. FISHER et al.** *J. Virol.*, 1993, vol. 70, 520-532 [0037]
- **GRAHAM et al.** *Virology*, 1973, vol. 52, 456 [0039]
- **SAMBROOK et al.** Molecular Cloning, a laboratory manual. Cold Spring Harbor Laboratories, 1989 [0039]
- **DAVIS et al.** Basic Methods in Molecular Biology. Elsevier, 1986 [0039]
- **CHU et al.** *Gene*, 1981, vol. 13, 197 [0039]
- **B. J. CARTER.** Handbook of Parvoviruses. CRC Press, 1990, 155-168 [0045]
- **SAMBROOK et al.** Molecular Cloning. A Laboratory Manual. Cold Spring Harbor Laboratory, 1989 [0045]
- **K. FISHER et al.** *J Virol.*, 1996, vol. 70, 520-532 [0045]
- **AUSUBEL et al.** Current Protocols in Molecular Biology. John Wiley & Sons, 1989 [0048]
- **RYAN, M D et al.** *EMBO*, 1994, vol. 4, 928-933 [0048]
- **MATTION, N M et al.** *J Virology*, November 1996, 8124-8127 [0048]
- **FURLER, S et al.** *Gene Therapy*, 2001, vol. 8, 864-873 [0048]
- **HALPIN, C et al.** *The Plant Journal*, 1999, vol. 4, 453-459 [0048]
- **DE FELIPE, P et al.** *Gene Therapy*, 1999, vol. 6, 198-208 [0048]
- **DE FELIPE, P et al.** *Human Gene Therapy*, 2000, vol. 11, 1921-1931 [0048]
- **KLUMP, H et al.** *Gene Therapy*, 2001, vol. 8, 811-817 [0048]
- **BOSHART et al.** *Cell*, 1985, vol. 41, 521-530 [0050]
- **NO et al.** *Proc. Natl. Acad. Sci. USA*, 1996, vol. 93, 3346-3351 [0051]
- **GOSSEN et al.** *Proc. Natl. Acad. Sci. USA*, 1992, vol. 89, 5547-5551 [0051]
- **GOSSEN et al.** *Science*, 1995, vol. 268, 1766-1769 [0051]
- **HARVEY et al.** *Curr. Opin. Chem. Biol.*, 1998, vol. 2, 512-518 [0051]
- **WANG et al.** *Nat. Biotech.*, 1997, vol. 15, 239-243 [0051]
- **WANG et al.** *Gene Ther.*, 1997, vol. 4, 432-441 [0051]
- **MAGARI et al.** *J. Clin. Invest.*, 1997, vol. 100, 2865-2872 [0051]
- **ANDERSEN et al.** *Cell. Mol. Neurobiol.*, 1993, vol. 13, 503-15 [0053]
- **PICCIOLI et al.** *Proc. Natl. Acad. Sci. USA*, 1991, vol. 88, 5611-5 [0053]
- **PICCIOLI et al.** *Neuron*, 1995, vol. 15, 373-84 [0053]
- **COSTANTINI LC et al.** *Gene Therapy*, 2000, vol. 7, 93-109 [0057]
- **STEIN et al.** *J Virol*, 1999, vol. 73, 3424-3429 [0060]

- DAVIDSON et al. *PNAS*, 2000, vol. 97, 3428-3432 [0060]
- DAVIDSON et al. *Nat. Genet.*, 1993, vol. 3, 219-223 [0060]
- ALISKY ; DAVIDSON. *Hum. Gene Ther.*, 2000, vol. 11, 2315-2329 [0060]
- WRIGHT FR et al. *Molecular Therapy*, 2005, vol. 12, 171-178 [0065]
- Remington's Pharmaceutical Sciences. 1035-1038, 1570-1580 [0068]
- MATALON R et al. *The Journal of Gene Medicine*, vol. 2 (3), 165-175 [0087]
- FOUST KD et al. *Nature Biotechnology*, March 2010, vol. 28 (3), 271-274 [0100]
- DAYA S ; BERNS KI. Gene Therapy Using Adeno-Associated Virus Vectors. *Clin Microbiol Rev*, 2008, vol. 21, 583-593 [0101]
- EBERLING JL ; JAGUST WJ ; CHRISTINE CW ; STARR P ; LARSON P ; BANKIEWICZ KS ; AMINOFF MJ. Results from a phase I safety trial of hAA� gene therapy for Parkinson disease. *Neurology*, 2008, vol. 70, 1980-1983 [0101]
- FEIGIN A ; KAPLITTMG ; TANG C ; LINT ; MATTIS P ; DHAWAN V ; DURING MJ ; EIDELBERG D. Modulation of metabolic brain networks after subthalamic gene therapy for Parkinson's disease. *Proceedings of the National Academy of Sciences*, 2007, vol. 104, 19559-19564 [0101]
- CIDECIYAN AV ; HAUSWIRTH WW ; ALEMAN TS ; KAUSHAL S ; SCHWARTZ SB ; BOYE SL ; WINDSOR EAM ; CONLON TJ ; SUMAROKA A ; PANG J-J et al. Human RPE65 Gene Therapy for Leber Congenital Amaurosis: Persistence of Early Visual Improvements and Safety at 1 Year. *Hum Gen Ther*, vol. 0 [0101]
- RAOUL C ; ABBAS-TERKI T ; BENSADOUN J-C ; GUILLOT S ; HAASE G ; SZULC J ; HENDERSON CE ; AEBISCHER P. Lentiviral-mediated silencing of SOD1 through RNA interference retards disease onset and progression in a mouse model of ALS. *Nat Med*, 2005, vol. 11, 423-428 [0101]
- KASPAR BK ; LLADO J ; SHERKAT N ; ROTHSTEIN JD ; GAGE FH. Retrograde Viral Delivery of IGF-1 Prolongs Survival in a Mouse ALS Model. *Science*, 2003, vol. 301, 839-842 [0101]
- AZZOUZ M ; RALPH GS ; STORKEBAUM E ; WALMSLEY LE ; MITROPHANOUS KA ; KINGSMAN SM ; CARMELIET P. Mazarakis ND: VEGF delivery with retrogradely transported lentivector prolongs survival in a mouse ALS model. *Nature*, 2004, vol. 429, 413-417 [0101]
- RALPH GS ; RADCLIFFE PA ; DAY DM ; CARTHY JM ; LEROUX MA ; LEE DCP ; WONG L-F ; BILSLAND LG ; GREENSMITH L ; KINGSMAN SM et al. Silencing mutant SOD1 using RNAi protects against neurodegeneration and extends survival in an ALS model. *Nat Med*, 2005, vol. 11, 429-433 [0101]
- WU R ; WANG H ; XIA X ; ZHOU H ; LIU C ; CASTRO M ; XU Z. Nerve Injection of Viral Vectors Efficiently Transfers Transgenes into Motor Neurons and Delivers RNAi Therapy Against ALS. *Antioxidants & Redox Signaling*, 2009, vol. 11, 1523-1534 [0101]
- FOUST KD ; NURRE E ; MONTGOMERY CL ; HERNANDEZ A ; CHAN CM ; KASPAR BK. Intravascular AAV9 preferentially targets neonatal neurons and adult astrocytes. *Nat Biotechnol*, 2009, vol. 27, 59-65 [0101]
- DUQUE S ; JOUSSEMET B ; RIVIERE C ; MARAIS T ; DUBREUIL ; DOUAR AM ; FYFE J ; MOULLIER P ; COLLE MA ; BARKATS M. Intravenous administration of self-complementary AAV9 enables transgene delivery to adult motor neurons. *Mol Ther*, 2009, vol. 17, 1187-1196 [0101]
- VANDENBERGHE LH ; WILSON JM ; GAO G. Tailoring the AAV vector capsid for gene therapy. *Gene Ther*, 2009, vol. 16, 311-319 [0101]
- MCBRIDE JL ; BOUDREAU RL ; HARPER SQ ; STABER PD ; MONTEYS AM ; MARTINS I ; GILMORE BL ; BURSTEIN H ; PELUSO RW ; POLISKY B et al. Artificial miRNAs mitigate shRNA-mediated toxicity in the brain: Implications for the therapeutic development of RNAi. *Proceedings of the National Academy of Sciences*, 2008, vol. 105, 5868-5873 [0101]
- MCCARTY DM. Self-complementary AAV Vectors: Advances and Applications. *Mol Ther*, 2008, vol. 16, 1648-1656 [0101]
- BOILLEE S ; YAMANAKA K ; LOBSIGER CS ; COPELAND NG ; JENKINS NA ; KASSIOTIS G ; KOLLIAS G ; CLEVELAND DW. Onset and progression in inherited ALS determined by motor neurons and microglia. *Science*, 2006, vol. 312, 1389-1392 [0101]
- XIA X ; ZHOU H ; HUANG Y ; XU Z. Allele-specific RNAi selectively silences mutant SOD1 and achieves significant therapeutic benefit in vivo. *Neurobiol Dis*, 2006, vol. 23, 578-586 [0101]
- YAMANAKA K ; CHUN SJ ; BOILLEE S ; FUJIMORI-TONOU N ; YAMASHITA H ; GUTMANN DH ; TAKAHASHI R ; MISAWA H ; CLEVELAND DW. Astrocytes as determinants of disease progression in inherited amyotrophic lateral sclerosis. *Nat Neurosci*, 2008, vol. 11, 251253 [0101]
- DIGIORGIO FP ; CARRASCO MA ; SIAO MC ; MANIATIS T ; EGGAN K. Non-cell autonomous effect of glia on motor neurons in an embryonic stem cell-based ALS model. *Nat Neurosci*, 2007, vol. 10, 608-614 [0101]
- NAGAI M ; RE DB ; NAGATA T ; CHALAZONITIS A ; JESSELL TM ; WICHTERLE H ; PRZEDBORSKI S. Astrocytes expressing ALS-linked mutated SOD1 release factors selectively toxic to motor neurons. *Nat Neurosci*, 2007, vol. 10, 615-622 [0101]

- **WANG Y ; OU MAO X ; XIE L ; BANWAITS ; MARTI HH ; GREENBERG DA ; JIN K.** Vascular Endothelial Growth Factor Overexpression Delays Neurodegeneration and Prolongs Survival in Amyotrophic Lateral Sclerosis Mice. *J Neurosci*, 2007, vol. 27, 304-307 [0101]
- **STORKEBAUM E ; LAMBRECHTS D ; DEW-ERCHIN M ; MORENO-MURCIANO M-P ; APPELMANS S ; OH H ; VAN DAMME P ; RUTTEN B ; MAN WY ; DE MOL M et al.** Treatment of motoneuron degeneration by intracerebroventricular delivery of VEGF in a rat model of ALS. *Nat Neurosci*, 2005, vol. 8, 85-92 [0101]
- **WU, W. P. ; XU, X. J. ; HAO, J. X.** *J. Neurosci. Methods*, 2004, vol. 133, 65-69 [0101]
- **PAPAIOANNOU, V. E. ; FOX, J. G.** *Lab. Anim. Sci.*, 1993, vol. 43, 189-192 [0101]