

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 January 2010 (07.01.2010)

PCT

(10) International Publication Number
WO 2010/002655 A2

(51) International Patent Classification:

C07D 401/14 (2006.01) **C07D 409/14** (2006.01)
C07D 403/12 (2006.01) **C07D 413/14** (2006.01)
C07D 403/14 (2006.01) **A61K 31/506** (2006.01)

(21) International Application Number:

PCT/US2009/048428

(22) International Filing Date:

24 June 2009 (24.06.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/075,583 25 June 2008 (25.06.2008) US
61/155,434 25 February 2009 (25.02.2009) US

(71) Applicant (for all designated States except US): **IRM LLC** [US/—]; 131 Front Street, P.O. Box 2899, Hamilton, HM LX (BM).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MARSILJE III, Thomas H.** [US/US]; 1844 Narragansett Court, San Diego, California 92107 (US). **LU, Wenshuo** [CA/US]; 7594 Wallingford Court, San Diego, California 92126 (US). **CHEN, Bei** [CN/US]; 11048 Creekbridge Place, San Diego, California 92128 (US). **HE, Xiaohui** [CN/US]; 12461 Dormouse Road, San Diego, California 92129 (US). **LEE, Christian Cho-Hua** [US/US]; 11115 Ice Skate Place, San Diego, California 92126 (US). **JIANG, Songchun** [CN/US]; 10512 Rancho Carmel Drive, San Diego, California 92128 (US). **YANG, Kunyong** [CN/US]; 11861 Spruce Run Drive, #C, San Diego, California 92131 (US).

(74) Agents: **TONGCO WU, Emily** et al.; Genomics Institute of the Novartis Research Foundation, 10675 John Jay Hopkins Drive, San Diego, California 92121 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: COMPOUNDS AND COMPOSITIONS AS KINASE INHIBITORS

(57) Abstract: The invention provides novel pyrimidine derivatives and pharmaceutical compositions thereof, and methods for using such compounds. For example, the pyrimidine derivatives of the invention may be used to treat, ameliorate or prevent a condition which responds to inhibition of insulin-like growth factor (IGF-1R) or analplastic lymphoma kinase (ALK).



WO 2010/002655 A2

COMPOUNDS AND COMPOSITIONS AS KINASE INHIBITORS

Cross-Reference to Related Applications

[0001] This application claims the benefit of U.S. provisional application serial numbers 61/075,583, filed 25 June 2008, and 61/155,434, filed 25 February 2009, each of which is incorporated herein by reference in its entirety.

Technical Field

[0002] The invention relates to protein kinase inhibitors, more particularly novel pyrimidine derivatives and pharmaceutical compositions thereof, and their use as pharmaceuticals.

Background Art

[0003] Insulin-like growth factor (IGF-1) signaling is highly implicated in cancer, with the IGF-1 receptor (IGF-1R) as the predominating factor. IGF-1R is important for tumor transformation and survival of malignant cells, but is only partially involved in normal cell growth. Targeting of IGF-1R has been suggested to be a promising option for cancer therapy. (Larsson et al., Br. J. Cancer 92:2097-2101 (2005)).


[0004] Anaplastic lymphoma kinase (ALK), a member of the insulin receptor superfamily of receptor tyrosine kinases, has been implicated in oncogenesis in hematopoietic and non-hematopoietic tumors. The aberrant expression of full-length ALK receptor proteins has been reported in neuroblastomas and glioblastomas; and ALK fusion proteins have occurred in anaplastic large cell lymphoma. The study of ALK fusion proteins has also raised the possibility of new therapeutic treatments for patients with ALK-positive malignancies. (Pulford et al., Cell. Mol. Life Sci. 61:2939-2953 (2004)).

[0005] Because of the emerging disease-related roles of IGF-1R and ALK, there is a continuing need for compounds which may be useful for treating and preventing a disease which responds to inhibition of IGF-1R and ALK.

Disclosure of the Invention

[0006] The invention relates to novel pyrimidine derivatives and pharmaceutical compositions thereof, and their use as pharmaceuticals.

[0007] In one aspect, the invention provides a compound of Formula (1):



or a physiologically acceptable salt thereof;

wherein ring E may optionally contain a double bond;

one of Z^1 , Z^2 and Z^3 is NR^6 , $N(R^6)^+-O^-$ or $S(O)_{1-2}$ and the others are CR_2 ;

R^1 is halo or an optionally halogenated C_{1-6} alkyl;


R^2 is pyridine-2-onyl, azepan-2-onyl or a monocyclic 5-6 membered heteroaryl having 1-3 heteroatoms selected from N, O and S; each of which is optionally substituted substituted with R^9 wherein R^9 is C_{1-6} alkyl, C_{1-6} haloalkyl or C_{3-7} cycloalkyl;

R^3 and R^4 are each H;

R^5 is halo, hydroxyl, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkoxy, cyano or $C(O)O_{0-1}R^8$;

R^6 is H; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $(CR_2)_pNR(CR_2)_pOR^7$, $(CR_2)_pNR-L-C(O)R^8$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $-C(O)-(CR_2)_p-OR^7$, $L-Y$, $-L-C(O)R^7$, $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)NR(CR_2)_pOR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-C(O)NR(CR_2)_pSR^7$, $-L-C(O)NR(CR_2)_pS(O)_{1-2}R^8$, $-L-S(O)_2R^8$, $-L-S(O)_2-(CR_2)_q-NRR^7$, $-L-S(O)_2NR(CR_2)_pNR(R^7)$ or $-L-S(O)_2NR(CR_2)_pOR^7$;

alternatively, R^6 is a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} and R^{16} are independently selected from H; C_{1-6} alkoxy; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R^{11} and R^{12} , R^{12} and R^{15} , R^{15} and R^{16} , R^{13} and R^{14} , or R^{13} and R^{15} together with the atoms to which they are attached may form a 3-7 membered saturated, unsaturated or partially unsaturated ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;

L is $(CR_2)_{1-4}$ or a bond;

Y is C_{3-7} carbocyclic ring, C_{6-10} aryl, or a 5-10 membered heteroaryl or 4-10 membered heterocyclic ring, each of which is optionally substituted with 1-3 R^5 groups;

R^7 , R^8 and R^{8a} are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or cyano; $(CR_2)_q Y$ or C_{1-6} alkoxy; or R^7 is H;

each R is independently H or C_{1-6} alkyl;

R and R^7 together with N in each NRR^7 may form a 5-6 membered ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;


m is 2-4;

n is 1-3;

p is 1-4; and

q is 0-4.


[0008] In one embodiment, the invention provides a compound is of Formula (2):



wherein one of R^{5a} , R^{5b} and R^{5c} is H and the others are independently halo, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, cyano or $C(O)O_{0-1}R^8$ wherein R^8 is C_{1-6} alkyl; and

R^1 , R^2 , R^3 , R^4 , E, Z^1 , Z^2 and Z^3 are as defined in Formula (1).

[0009] In some examples, Z^3 in the above Formula (2) is NR^6 or $N(R^6)^+-O^-$; and Z^1 and Z^2 are CH_2 . In other examples, R^6 is H, C_{1-6} alkyl optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, L-Y, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $-L-C(O)R^7$, $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-S(O)_2R^8$, $-L-S(O)_2-(CR_2)_q-NRR^7$, or a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;


R^{8a} is C_{1-6} alkyl; and

R^7 , R^8 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R, L, Y, p and q are as defined in

Formula (1).

[0010] In the above Formula (1) and (2), R^2 may be pyrazolyl, isoxazolyl, pyridine-2-onyl or azepan-2-onyl, each of which is substituted with C_{1-6} alkyl, C_{1-6} haloalkyl or C_{3-7} cycloalkyl.

[0011] In another embodiment, the invention provides a compound of Formula (3):




or a tautomer thereof;

wherein R^{5b} is H; and R^{5a} and R^{5c} are independently halo, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, cyano or $C(O)O_{0-1}R^8$ wherein R^8 is C_{1-6} alkyl;

Z^1 and Z^2 are CH_2 ;

Z^3 is NR^6 or $N(R^6)^+-O^-$;

R^6 is H, C_{1-6} alkyl optionally substituted with halo and/or hydroxyl groups;
 $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$,
 $-(CR_2)_p-C(O)OR^7$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $L-Y$, $-L-C(O)R^7$,
 $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-S(O)_2R^8$,
 $-L-S(O)_2-(CR_2)_q-NRR^7$, or a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;


R^{8a} is C_{1-6} alkyl; and

$R^1, R^3, R^4, R^7, R^8, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R, L, Y, p, q$ and E are as defined in Formula (1).

[0012] In the above Formula (3), R^{5a} may be halo and R^{5c} is C_{1-6} alkyl. In some examples, R^6 is C_{1-6} alkyl or a radical of formula (a) or (c); and R^{10} is O.

[0013] In the above Formula (1), (2) and (3), R^1 may be halo. In particular examples, R^1 is chloro. In other examples, R^3 and R^4 are H.

[0014] In another aspect, the present invention provides a compound of Formula (4):



or a physiologically acceptable salt thereof;

wherein X is a monocyclic 5-6 membered heterocyclic ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted on a substitutable ring carbon with oxo and at any substitutable ring nitrogen by R^6 ;


R^1 is halo;

R^3 and R^4 are each H;

R^5 is halo, hydroxyl, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkoxy, cyano or $C(O)O_{0-1}R^8$;

R^6 is H; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $(CR_2)_pNR(CR_2)_pOR^7$, $(CR_2)_pNR-L-C(O)R^8$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $-C(O)-(CR_2)_p-OR^7$, $L-Y$, $-L-C(O)R^7$, $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)NR(CR_2)_pOR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-C(O)NR(CR_2)_pSR^7$, $-L-C(O)NR(CR_2)_pS(O)_{1-2}R^8$, $-L-S(O)_2R^8$, $-L-S(O)_2-(CR_2)_q-NRR^7$, $-L-S(O)_2NR(CR_2)_pNR(R^7)$ or $-L-S(O)_2NR(CR_2)_pOR^7$;

alternatively, R^6 is a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} and R^{16} are independently selected from H; C_{1-6} alkoxy; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R^{11} and R^{12} , R^{12} and R^{15} , R^{15} and R^{16} , R^{13} and R^{14} , or R^{13} and R^{15} together with the atoms to which they are attached may form a 3-7 membered saturated, unsaturated or partially unsaturated ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;

L is $(CR_2)_{1-4}$ or a bond;

Y is C_{3-7} carbocyclic ring, C_{6-10} aryl, or a 5-10 membered heteroaryl or 4-10 membered heterocyclic ring, each of which is optionally substituted with 1-3 R^5 groups;

R^7 , R^8 and R^{8a} are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or cyano; $(CR_2)_qY$ or C_{1-6} alkoxy; or R^7 is H;

each R is independently H or C₁₋₆ alkyl;

R and R⁷ together with N in each NRR⁷ may form a 5-6 membered ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R⁵ groups;

m is 2-4;

n is 1-3;

p is 1-4; and

q is 0-4.

[0015] In the above Formula (4), X may be morpholinyl, piperazinyl, piperazin-2-onyl, oxazolidin-2-onyl or piperidin-2-onyl each of which is substituted at any substitutable ring nitrogen by C₁₋₆ alkyl; or X is tetrahydro-2H-pyran-2-onyl; and R¹ is chloro.

[0016] In another aspect, the present invention provides pharmaceutical compositions comprising a compound having Formula (1), (2), (3) and (4), and a physiologically acceptable excipient.

[0017] In yet another aspect, the invention provides methods for inhibiting IGF-1R in a cell, comprising contacting the cell with an effective amount of a compound having Formula (1), (2), (3) and (4) or a pharmaceutical composition thereof.

[0018] The invention also provides methods to treat, ameliorate or prevent a condition which responds to inhibition of IGF-1R or anaplastic lymphoma kinase (ALK) in a mammal suffering from said condition, comprising administering to the mammal a therapeutically effective amount of a compound having Formula (1), (2), (3) and (4) or a pharmaceutical composition thereof, and optionally in combination with a second therapeutic agent. Alternatively, the present invention provides for the use of a compound having Formula (1), (2), (3) and (4), and optionally in combination with a second therapeutic agent, in the manufacture of a medicament for treating a condition mediated by IGF-1R or ALK. The compounds of the invention may be administered, for example, to a mammal suffering from an autoimmune disease, a transplantation disease, an infectious disease or a cell proliferative disorder. In particular examples, the compounds of the invention may be used alone or in combination with a chemotherapeutic agent to treat a cell proliferative disorder, including but not limited to, multiple myeloma, neuroblastoma, synovial, hepatocellular, Ewing's Sarcoma or a solid tumor selected from a osteosarcoma,

melanoma, and tumor of breast, renal, prostate, colorectal, thyroid, ovarian, pancreatic, lung, uterine or gastrointestinal tumor.

Definitions

[0019] “Alkyl” refers to a moiety and as a structural element of other groups, for example halo-substituted-alkyl and alkoxy, and may be straight-chained or branched. An optionally substituted alkyl, alkenyl or alkynyl as used herein may be optionally halogenated (e.g., CF₃), or may have one or more carbons that is substituted or replaced with a heteroatom, such as NR, O or S (e.g., –OCH₂CH₂O–, alkylthiols, thioalkoxy, alkylamines, etc).

[0020] “Aryl” refers to a monocyclic or fused bicyclic aromatic ring containing carbon atoms. “Arylene” means a divalent radical derived from an aryl group. For example, an aryl group may be phenyl, indenyl, indanyl, naphthyl, or 1,2,3,4-tetrahydronaphthalenyl, which may be optionally substituted in the ortho, meta or para position.

[0021] “Heteroaryl” as used herein is as defined for aryl above, where one or more of the ring members is a heteroatom. For example, a heteroaryl substituent for use in the compounds of the invention may be a monocyclic or bicyclic 5-10 membered heteroaryl containing 1-4 heteroatoms selected from N, O, and S. Examples of heteroaryls include but are not limited to pyridyl, pyrazinyl, indolyl, indazolyl, quinoxaliny, quinoliny, benzofuranyl, benzopyranyl, benzothiopyranyl, benzo[1,3]dioxole, imidazolyl, benzo-imidazolyl, pyrimidinyl, furanyl, oxazolyl, isoxazolyl, triazolyl, benzotriazolyl, tetrazolyl, pyrazolyl, thienyl, pyrrolyl, isoquinoliny, purinyl, thiazolyl, tetrazinyl, benzothiazolyl, oxadiazolyl, benzoxadiazolyl, etc.

[0022] A “carbocyclic ring” as used herein refers to a saturated or partially unsaturated, monocyclic, fused bicyclic or bridged polycyclic ring containing carbon atoms, which may optionally be substituted, for example, with =O. Examples of carbocyclic rings include but are not limited to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylene, cyclohexanone, etc.

[0023] A “heterocyclic ring” as used herein is as defined for a carbocyclic ring above, wherein one or more ring carbons is a heteroatom. For example, a heterocyclic ring for use in the compounds of the invention may be a 4-7 membered heterocyclic ring containing 1-3 heteroatoms selected from N, O and S, or a combination thereof

such as -S(O) or -S(O)₂-. Examples of heterocyclic rings include but are not limited to azetidiny, morpholino, pyrrolidinyl, pyrrolidinyl-2-one, piperaziny, piperidinyl, piperidinylone, 1,4-dioxa-8-aza-spiro[4.5]dec-8-yl, 1,2,3,4-tetrahydroquinolinyl, etc. Heterocyclic rings as used herein may encompass bicyclic amines and bicyclic diamines.

[0024] As used herein, an H atom in any substituent groups (e.g., CH₂) encompasses all suitable isotopic variations, e.g., H, ²H and ³H.

[0025] The term “pharmaceutical combination” as used herein refers to a product obtained from mixing or combining active ingredients, and includes both fixed and non-fixed combinations of the active ingredients. The term “fixed combination” means that the active ingredients, e.g. a compound of Formula (1) and a co-agent, are both administered to a patient simultaneously in the form of a single entity or dosage. The term “non-fixed combination” means that the active ingredients, e.g. a compound of Formula (1) and a co-agent, are both administered to a patient as separate entities either simultaneously, concurrently or sequentially with no specific time limits, wherein such administration provides therapeutically effective levels of the active ingredients in the body of the patient. The latter also applies to cocktail therapy, e.g. the administration of three or more active ingredients.

[0026] “Mammal” refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. In particular examples, the mammal is human.

[0027] The term "administration" or "administering" of the subject compound means providing a compound of the invention and prodrugs thereof to a subject in need of treatment. Administration “in combination with” one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order, and in any route of administration.

[0028] An “effective amount” of a compound is an amount sufficient to carry out a specifically stated purpose. An “effective amount” may be determined empirically and in a routine manner, in relation to the stated purpose.

[0029] The term “therapeutically effective amount” refers to an amount of a compound (e.g., an IGF-1R antagonist) effective to “treat” an IGF-1R-mediated disorder in a subject or mammal. In the case of cancer, the therapeutically effective amount of the drug may reduce the number of cancer cells; reduce the tumor size;

inhibit (i.e., slow to some extent and preferably stop) cancer cell infiltration into peripheral organs; inhibit (i.e., slow to some extent and preferably stop) tumor metastasis; inhibit, to some extent, tumor growth; and/or relieve to some extent one or more of the symptoms associated with the cancer. See the definition herein of “treating”. To the extent the drug may prevent growth and/or kill existing cancer cells, it may be cytostatic and/or cytotoxic.

[0030] The term “cancer” refers to the physiological condition in mammals that is typically characterized by unregulated cell growth/proliferation. Examples of cancer include, but are not limited to: carcinoma, lymphoma, blastoma, and leukemia. More particular examples of cancers include, but are not limited to: chronic lymphocytic leukemia (CLL), lung, including non small cell (NSCLC), breast, ovarian, cervical, endometrial, prostate, colorectal, intestinal carcinoid, bladder, gastric, pancreatic, hepatic (hepatocellular), hepatoblastoma, esophageal, pulmonary adenocarcinoma, mesothelioma, synovial sarcoma, osteosarcoma, head and neck squamous cell carcinoma, juvenile nasopharyngeal angiofibromas, liposarcoma, thyroid, melanoma, basal cell carcinoma (BCC), medulloblastoma and desmoid.

[0031] “Treating” or “treatment” or “alleviation” refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic disease or condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to having the disorder or those in whom the disorder is to be prevented (prophylaxis). When the IGF-1R-mediated disorder is cancer, a subject or mammal is successfully “treated” or shows a reduced tumor burden if, after receiving a therapeutic amount of an IGF-1R antagonist according to the methods of the present invention, the patient shows observable and/or measurable reduction in or absence of one or more of the following: reduction in the number of cancer cells or absence of the cancer cells; reduction in the tumor size; inhibition (i.e., slow to some extent and preferably stop) of cancer cell infiltration into peripheral organs including the spread of cancer into soft tissue and bone; inhibition (i.e., slow to some extent and preferably stop) of tumor metastasis; inhibition, to some extent, of tumor growth; and/or relief to some extent, one or more of the symptoms associated with the specific cancer; reduced morbidity and mortality, and improvement in quality of life issues. To the extent the IGF-1R antagonist may prevent growth and/or kill existing cancer cells, it may be cytostatic

and/or cytotoxic. Reduction of these signs or symptoms may also be felt by the patient.

[0032] “Carriers” as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN®, polyethylene glycol (PEG), and PLURONICS®.

[0033] A “chemotherapeutic agent” is a chemical compound useful in the treatment of cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); delta-9-tetrahydrocannabinol (dronabinol, MARINOL®); beta-lapachone; lapachol; colchicines; betulinic acid; a camptothecin (including the synthetic analogue topotecan (HYCAMTIN®), CPT-11 (irinotecan, CAMPTOSAR®), acetylcamptothecin, scoplectin, and 9-aminocamptothecin); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); podophyllotoxin; podophyllinic acid; teniposide; cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratiastatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine,

trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gamma1I and calicheamicin omega1I (see, e.g., Agnew, Chem Intl. Ed. Engl., 33: 183-186 (1994)); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antiobiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, caminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitio stanol, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elfomithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; niraerine; pentostatin; phenamet; pirarubicin; losoxantrone; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, Oreg.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine (ELDISINE®, FILDESIN®); dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); thiotepa; taxoids, e.g., TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE™ Cremophor-free,

albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Ill.), and TAXOTERE® doxorubicin (Rhône-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine (GEMZAR®); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine (VELBAN®); platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine (ONCOVIN®); oxaliplatin; leucovorin; vinorelbine (NAVELBINE®); novantrone; edatrexate; daunomycin; aminopterin; ibandronate; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine (XELODA®); pharmaceutically acceptable salts, acids or derivatives of any of the above; as well as combinations of two or more of the above such as CHOP, an abbreviation for a combined therapy of cyclophosphamide, doxorubicin, vincristine, and prednisolone, and FOLFOX, an abbreviation for a treatment regimen with oxaliplatin (ELOXATIN™) combined with 5-FU and leucovorin.


[0034] Furthermore, a “chemotherapeutic agent” may include anti-hormonal agents that act to regulate, reduce, block, or inhibit the effects of hormones that can promote the growth of cancer, and are often in the form of systemic, or whole-body treatment. They may be hormones themselves. Examples include anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including NOLVADEX® tamoxifen), EVISTA® raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and FARESTON® toremifene; anti-progesterones; estrogen receptor down-regulators (ERDs); agents that function to suppress or shut down the ovaries, for example, leutinizing hormone-releasing hormone (LHRH) agonists such as LUPRON® and ELIGARD® leuprolide acetate, goserelin acetate, buserelin acetate and triptorelin; other anti-androgens such as flutamide, nilutamide and bicalutamide; and aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example, 4(5)-imidazoles, aminoglutethimide, MEGASE® megestrol acetate, AROMASIN® exemestane, formestane, fadrozole, RIVISOR® vorozole, FEMARA® letrozole, and ARIMIDEX® anastrozole. In addition, such definition of chemotherapeutic agents includes bisphosphonates such as clodronate (for example, BONEFOS® or OSTAC®), DIDROCAL® etidronate, NE-58095, ZOMETA® zoledronic acid/zoledronate, FOSAMAX® alendronate, AREDIA® pamidronate, SKELID® tiludronate, or ACTONEL® risedronate; as well as troxacitabine (a 1,3-

dioxolane nucleoside cytosine analog); antisense oligonucleotides, particularly those that inhibit expression of genes in signaling pathways implicated in aberrant cell proliferation, such as, for example, PKC- α , Raf, H-Ras, and epidermal growth factor receptor (EGF-R); vaccines such as THERATOPE® vaccine and gene therapy vaccines, for example, ALLOVECTIN® vaccine, LEUVECTIN® vaccine, and VAXID® vaccine; LURTOTECAN® topoisomerase 1 inhibitor; ABARELIX® rmRH; lapatinib ditosylate (an ErbB-2 and EGFR dual tyrosine kinase small-molecule inhibitor also known as GW572016); and pharmaceutically acceptable salts, acids or derivatives of any of the above.

Modes of Carrying Out the Invention

[0035] The invention provides novel pyrimidine derivatives and pharmaceutical compositions thereof, and methods for using such compounds.

[0036] In one aspect, the invention provides a compound of Formula (1):



or a physiologically acceptable salt thereof;

wherein ring E may optionally contain a double bond;

one of Z^1 , Z^2 and Z^3 is NR^6 , $N(R^6)^+-O^-$ or $S(O)_{1-2}$ and the others are CR_2 ;

R^1 is halo or an optionally halogenated C_{1-6} alkyl;

R^2 is pyridine-2-onyl, azepan-2-onyl or a monocyclic 5-6 membered heteroaryl having 1-3 heteroatoms selected from N, O and S; each of which is optionally substituted with R^9 wherein R^9 is C_{1-6} alkyl, C_{1-6} haloalkyl or C_{3-7} cycloalkyl;


R^3 and R^4 are each H;

R^5 is halo, hydroxyl, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkoxy, cyano or $C(O)O_{0-1}R^8$;

R^6 is H; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $(CR_2)_pNR(CR_2)_pOR^7$,

$(\text{CR}_2)_p\text{NR-L-C(O)R}^8$, $\text{C(O)(CR}_2)_q\text{OR}^8$, $-\text{C(O)O-(CR}_2)_p\text{-NRR}^7$, $-\text{C(O)-(CR}_2)_p\text{-OR}^7$, L-Y,
 $-\text{L-C(O)R}^7$, $-\text{L-C(O)-NRR}^7$, $-\text{L-C(O)-NR-(CR}_2)_p\text{-NRR}^7$, $-\text{L-C(O)NR(CR}_2)_p\text{OR}^7$,
 $-\text{L-C(O)-(CR}_2)_q\text{-NR-C(O)-R}^8$, $-\text{L-C(O)NR(CR}_2)_p\text{SR}^7$, $-\text{L-C(O)NR(CR}_2)_p\text{S(O)}_{1-2}\text{R}^8$,
 $-\text{L-S(O)}_2\text{R}^8$, $-\text{L-S(O)}_2\text{-(CR}_2)_q\text{-NRR}^7$, $-\text{L-S(O)}_2\text{NR(CR}_2)_p\text{NR(R}^7)$ or
 $-\text{L-S(O)}_2\text{NR(CR}_2)_p\text{OR}^7$;

alternatively, R^6 is a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} and R^{16} are independently selected from H; C_{1-6} alkoxy;
 C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted
 with halo, amino or hydroxyl groups; or R^{11} and R^{12} , R^{12} and R^{15} , R^{15} and R^{16} , R^{13}
 and R^{14} , or R^{13} and R^{15} together with the atoms to which they are attached may form a
 3-7 membered saturated, unsaturated or partially unsaturated ring containing 1-3
 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5
 groups;

L is $(\text{CR}_2)_{1-4}$ or a bond;

Y is C_{3-7} carbocyclic ring, C_{6-10} aryl, or a 5-10 membered heteroaryl or 4-10
 membered heterocyclic ring, each of which is optionally substituted with 1-3 R^5
 groups;

R^7 , R^8 and R^{8a} are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each
 of which may be optionally substituted with halo, amino, hydroxyl or cyano; $(\text{CR}_2)_q\text{Y}$
 or C_{1-6} alkoxy; or R^7 is H;

each R is independently H or C_{1-6} alkyl;

R and R^7 together with N in each NRR^7 may form a 5-6 membered ring
 containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with
 oxo and 1-3 R^5 groups;


m is 2-4;

n is 1-3;

p is 1-4; and

q is 0-4.


[0037] In one embodiment, the invention provides a compound is of Formula (2):



wherein one of R^{5a} , R^{5b} and R^{5c} is H and the others are independently halo, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, cyano or $C(O)O_{0-1}R^8$ wherein R^8 is C_{1-6} alkyl; and

R^1 , R^2 , R^3 , R^4 , E, Z^1 , Z^2 and Z^3 are as defined in Formula (1).

[0038] In another embodiment, the invention provides a compound of Formula (3):




or a tautomer thereof;

wherein R^{5b} is H; and R^{5a} and R^{5c} are independently halo, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, cyano or $C(O)O_{0-1}R^8$ wherein R^8 is C_{1-6} alkyl;

Z^1 and Z^2 are CH^2 ;

Z^3 is NR^6 or $N(R^6)^+-O^-$;

R^6 is H, C_{1-6} alkyl optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, L-Y, -L-C(O) R^7 , -L-C(O)-NRR⁷, -L-C(O)-NR-(CR₂)_p-NRR⁷, -L-C(O)-(CR₂)_q-NR-C(O)-R⁸, -L-S(O)₂R⁸, -L-S(O)₂-(CR₂)_q-NRR⁷, or a radical selected from formula (a), (b), (c) or (d):




R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;

R^{8a} is C_{1-6} alkyl; and

$R^1, R^3, R^4, R^7, R^8, R^{10}, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R, L, Y, p, q$ and E are as defined in Formula (1).

[0039] In another aspect, the present invention provides a compound of Formula (4):



or a physiologically acceptable salt thereof;

wherein X is a monocyclic 5-6 membered heterocyclic ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted on a substitutable ring carbon with oxo and at any substitutable ring nitrogen by R^6 ;


R^1 is halo;

R^3 and R^4 are each H;

R^5 is halo, hydroxyl, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkoxy, cyano or $C(O)O_{0-1}R^8$;

R^6 is H; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $(CR_2)_pNR(CR_2)_pOR^7$, $(CR_2)_pNR-L-C(O)R^8$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $-C(O)-(CR_2)_p-OR^7$, $L-Y$, $-L-C(O)R^7$, $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)NR(CR_2)_pOR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-C(O)NR(CR_2)_pSR^7$, $-L-C(O)NR(CR_2)_pS(O)_{1-2}R^8$, $-L-S(O)_2R^8$, $-L-S(O)_2-(CR_2)_q-NRR^7$, $-L-S(O)_2NR(CR_2)_pNR(R^7)$ or $-L-S(O)_2NR(CR_2)_pOR^7$;

alternatively, R^6 is a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} and R^{16} are independently selected from H; C_{1-6} alkoxy; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R^{11} and R^{12} , R^{12} and R^{15} , R^{15} and R^{16} , R^{13} and R^{14} , or R^{13} and R^{15} together with the atoms to which they are attached may form a 3-7 membered saturated, unsaturated or partially unsaturated ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;

L is $(CR_2)_{1-4}$ or a bond;

Y is C_{3-7} carbocyclic ring, C_{6-10} aryl, or a 5-10 membered heteroaryl or 4-10 membered heterocyclic ring, each of which is optionally substituted with 1-3 R^5 groups;

R^7 , R^8 and R^{8a} are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or cyano; $(CR_2)_qY$ or C_{1-6} alkoxy; or R^7 is H;

each R is independently H or C_{1-6} alkyl;

R and R^7 together with N in each NRR^7 may form a 5-6 membered ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;


m is 2-4;

n is 1-3;

p is 1-4; and

q is 0-4.

[0040] In yet another aspect, the invention provides compounds of Formula (5):



or physiologically acceptable salts thereof;

wherein X is a 4-7 membered heterocyclic ring containing NR^6 , $\text{—N}^+\text{R}^6$, O or S, and optionally substituted with 1-3 R^5 groups;

wherein R^1 is halo, C_{1-6} alkyl, or a halo-substituted C_{1-6} alkyl;


R^2 is an optionally substituted 5-6 membered heteroaryl or 5-7 membered heterocyclic ring, each having 1-3 heteroatoms selected from N, O and S and optionally substituted with 1-3 R^9 groups;

R^3 and R^4 are independently H, $\text{C}(\text{O})\text{R}^8$, C_{1-6} alkyl or halo-substituted C_{1-6} alkyl;

R^5 and R^9 are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R^5 and R^9 are independently C_{1-6} alkoxy, halo-substituted C_{1-6} alkoxy, halo, nitro, cyano, $\text{CR}(\text{OR}^7)\text{R}^7$, OR^7 , $\text{O}(\text{CR}_2)_p\text{—OR}^7$, $\text{NR}(\text{R}^7)$, $\text{CR}(\text{R}^7)\text{NRR}^7$, $(\text{CR}_2)_{1-6}\text{NR}(\text{CR}_2)_p\text{OR}^7$, $(\text{CR}_2)_{1-6}\text{NR}(\text{CR}_2)_q\text{C}(\text{O})\text{R}^8$, $\text{—C}(\text{O})\text{—}(\text{CR}_2)_q\text{—NR—C}(\text{O})\text{—R}^8$, $(\text{CR}_2)_q\text{Y}$, $\text{C}(\text{O})\text{O}_{0-1}\text{R}^7$, $\text{C}(\text{O})\text{NR}(\text{R}^7)$, $\text{C}(\text{O})\text{CRR}^7\text{—NR}(\text{R}^7)$, $\text{C}(\text{O})\text{NR}(\text{CR}_2)_p\text{NR}(\text{R}^7)$, $\text{C}(\text{O})\text{NR}(\text{CR}_2)_p\text{OR}^7$, $\text{C}(\text{O})\text{NR}(\text{CR}_2)_p\text{SR}^7$, $\text{C}(\text{O})\text{NR}(\text{CR}_2)_p\text{S}(\text{O})_{1-2}\text{R}^8$, $\text{S}(\text{O})_{0-2}\text{R}^8$, $\text{S}(\text{O})_2\text{NRR}^7$, $\text{S}(\text{O})_2\text{NR}(\text{CR}_2)_p\text{NR}(\text{R}^7)$, or $\text{S}(\text{O})_2\text{NR}(\text{CR}_2)_p\text{OR}^7$;

R^6 is H; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or alkoxy; $\text{—}(\text{CR}_2)_p\text{—CN}$; $\text{C}(\text{O})\text{H}$, $\text{CR}(\text{OR}^7)\text{R}^7$, $(\text{CR}_2)_p\text{—OR}^7$, $(\text{CR}_2)_p\text{—NR}(\text{R}^7)$, $\text{CR}(\text{R}^7)\text{NRR}^7$, —L—Y , $\text{—L—C}(\text{O})\text{O}_{0-1}\text{—}(\text{CR}_2)_q\text{—R}^8$, $\text{—}(\text{CR}_2)_p\text{—C}(\text{O})\text{O}_{0-1}\text{—R}^7$, $\text{—L—C}(\text{O})\text{—NRR}^7$, $\text{—L—C}(\text{O})\text{O}_{0-1}\text{—CR}(\text{R}^7)\text{—NRR}^7$, $\text{—L—C}(\text{O})\text{—NR—}(\text{CR}_2)_p\text{—NRR}^7$, $\text{—L—C}(\text{O})\text{NR}(\text{CR}_2)_p\text{OR}^7$, $\text{—L—C}(\text{O})\text{—}(\text{CR}_2)_q\text{—NR—C}(\text{O})\text{—R}^8$, $\text{—L—C}(\text{O})\text{NR}(\text{CR}_2)_p\text{SR}^7$, $\text{—L—C}(\text{O})\text{NR}(\text{CR}_2)_p\text{S}(\text{O})_{1-2}\text{R}^8$, $\text{—L—S}(\text{O})_2\text{R}^8$, $(\text{CR}_2)_p\text{NR}(\text{CR}_2)_p\text{OR}^7$, $(\text{CR}_2)_p\text{NR—L—C}(\text{O})\text{R}^8$, $\text{—L—S}(\text{O})_2\text{NRR}^7$, $\text{—L—S}(\text{O})_2\text{NR}(\text{CR}_2)_p\text{NR}(\text{R}^7)$ or $\text{—L—S}(\text{O})_2\text{NR}(\text{CR}_2)_p\text{OR}^7$;

alternatively, R^6 is a radical selected from formula (a), (b), (c) or (d):



wherein R^{11} , R^{12} , R^{13} , R^{14} , R^{15} and R^{16} are independently selected from H or C_{1-6} alkyl, C_{1-6} alkoxy, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R^{11} and R^{12} , R^{12} and R^{15} , R^{15} and R^{16} , R^{13} and R^{14} , or R^{13} and R^{15} together with the carbon and/or nitrogen atoms to which they are attached may form a 3-7 membered saturated, unsaturated or partially unsaturated ring optionally containing up to 3 atoms or groups selected from C(O), N, O and S(O)₀₋₂;

R^{17} is O or SO₂R⁸;

L is (CR₂)₁₋₄ or a bond;

Y is C₃₋₇ carbocyclic ring, C₆₋₁₀ aryl, or a 5-10 membered heteroaryl or 4-10 membered heterocyclic ring, each of which is optionally substituted with 1-3 R⁵ groups;

R^7 and R^8 are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or cyano; (CR₂)_qY or C_{1-6} alkoxy; or R^7 is H;

each R is H or C_{1-6} alkyl;

m is 2-4;

n is 1-3;

p is 1-4; and

q is 0-4.

[0041] In the above Formula (5), examples of 5-6 membered heteroaryl or 5-7 membered heterocyclic ring R² groups include but are not limited to pyrazolyl, pyrrolyl, thiophenyl, pyrimidinyl, isoxazolyl, pyridyl, azepan-2-onyl, 2H-thiopyran, 3H-thiopyran, 4H-thiopyran, tetrahydrothiopyran, 2H-pyran, 4H-pyran, tetrahydropyran, piperidine, 1,2-dithiin, 1,2-dithiane, 1,3-dithiin, 1,3-dithiane, 1,4-dithiin, 1,4-dithiane, 1,2-dioxin, 1,2-dioxane, 1,3-dioxin, 1,3-dioxane, 1,4-dioxin, 1,4-dioxane, piperazine, 1,2-oxathiin, 1,2-oxathiane, 4H-1,3-oxathiin, 1,3-oxathiane, 1,4-oxathiin, 1,4-oxathiane, 2H-1,2-thiazine, tetrahydro-1,2-thiazine, 2H-1,3-thiazine,

4H-1, 3-thiazine, 5, 6-dihydro-4H-thiazine, 4H-1, 4-thiazine, tetrahydro-1, 4-thiazine, 2H-1, 2-oxazine, 4H-1, 2-oxazine, 6H-1, 2-oxazine, 2H-1,3-oxazine, 4H-1,3-oxazine, 4H-1, 4-oxazine, morpholine, trioxane, 4H-1, 2,3-trithiin, 1,2, 3-trithiane, 1,3, 5-trithiane, hexahydro-1,3, 5-triazine, tetrahydrothiophene, tetrahydrofuran, pyrroline, pyrrolidine, pyrrolidone, pyrrolidione, pyrazoline, pyrazolidine, imidazoline, imidazolidine, 1,2-dioxole, 1,2-dioxolane, 1,3-dioxole, 1, 3-dioxolane, 3H-1,2-dithiole, 1, 2-dithiolane, 1,3-dithiole, 1,3-dithiolane, isoxazoline, isoxazolidine, oxazoline, oxazolidine, thiazoline, thiozolidine, 3H-1, 2-oxathiole, 1,2-oxathiolane, 5H-1,2-oxathiole, 1,3-oxathiole, 1,3-oxathiolane, 1,2, 3-trithiole, 1,2, 3-trithiolane, 1,2, 4-trithiolane, 1,2, 3- trioxole, 1,2, 3-trioxolane. 1,2, 4-trioxolane, 1,2, 3-triazoline and 1,2, 3-triazolidine.

[0042] In each of the above formula, any asymmetric carbon atoms may be present in the (R)-, (S)-or (R,S)-configuration. The compounds may thus be present as mixtures of isomers or as pure isomers, for example, as pure enantiomers or diastereomers. The invention further encompasses possible tautomers of the inventive compounds.

[0043] Any formula given herein is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, and chlorine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}F , ^{31}P , ^{32}P , ^{35}S , ^{36}Cl , ^{125}I respectively.

[0044] The invention includes various isotopically labeled compounds as defined herein, for example, those into which radioactive isotopes such as ^3H , ^{13}C , and ^{14}C , are present. Such isotopically labelled compounds are useful in metabolic studies (with, for example, ^{14}C), reaction kinetic studies (with, for example ^2H or ^3H), detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In other examples, an ^{18}F or labeled compound may be used for PET or SPECT studies. Isotopic variations of the compounds have the potential to change a compound's metabolic fate and/or create small changes in physical properties such as hydrophobicity, and the like.

Isotopic variations also have the potential to enhance efficacy and safety, enhance bioavailability and half-life, alter protein binding, change biodistribution, increase the proportion of active metabolites and/or decrease the formation of reactive or toxic metabolites. Isotopically labeled compounds of this invention and prodrugs thereof can generally be prepared by carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

[0045] In each of the above formula, each optionally substituted moiety may be substituted with C₁₋₆ alkyl, C₂₋₆ alkenyl or C₃₋₆ alkynyl, each of which may be optionally halogenated or optionally having a carbon that may be replaced or substituted with N, S, O, or a combination thereof (for example, hydroxylC₁₋₈alkyl, C₁₋₈alkoxyC₁₋₈alkyl); halo, amino, amidino, C₁₋₆ alkoxy; hydroxyl, methylenedioxy, carboxy; C₁₋₈ alkylcarbonyl, C₁₋₈ alkoxy carbonyl, carbamoyl, C₁₋₈ alkylcarbamoyl, sulfamoyl, cyano, oxo, nitro, or an optionally substituted carbocyclic ring, heterocyclic ring, aryl or heteroaryl as previously described.

Pharmacology and Utility

[0046] The compounds of the invention and their pharmaceutically acceptable salts exhibit valuable pharmacological properties when tested in vitro in cell-free kinase assays and in cellular assays, and are therefore useful as pharmaceuticals.

[0047] In one aspect, the compounds of the invention may inhibit insulin like growth-factor receptor 1 (IGF-1R), and may be useful in the treatment of IGF-1 R mediated diseases. Examples of IGF-1R mediated diseases include but are not limited to proliferative diseases, such as tumors, for example breast, renal, prostate, colorectal, thyroid, ovarian, pancreas, neuronal, lung, uterine and gastro intestinal tumors, as well as osteosarcomas and melanomas. The efficacy of the compounds of the invention as inhibitors of IGF-1R tyrosine kinase activity may be demonstrated using a cellular capture ELISA. In this assay, the activity of the compounds of the invention against (IGF-1)-induced autophosphorylation of the IGF-1R is determined.

[0048] In another aspect, the compounds of the invention may inhibit the tyrosine kinase activity of anaplastic lymphoma kinase (ALK) and the fusion protein of NPM-ALK. This protein tyrosine kinase results from a gene fusion of nucleophosmin (NPM) and ALK, rendering the protein tyrosine kinase activity of ALK ligand independent. NPM-ALK plays a key role in signal transmission in a number of

hematopoietic and other human cells leading to hematological and neoplastic diseases, for example in anaplastic large-cell lymphoma (ALCL) and non-Hodgkin's lymphomas (NHL), specifically in ALK+NHL or Alkomas, in inflammatory myofibroblastic tumors (IMT) and neuroblastomas. (Duyster et al. 2001 *Oncogene* 20, 5623-5637). In addition to NPM-ALK, other gene fusions have been identified in human hematological and neoplastic diseases; for example, TPM3-ALK (a fusion of nonmuscle tropomyosin with ALK).

[0049] The inhibition of ALK tyrosine kinase activity may be demonstrated using known methods, for example using the recombinant kinase domain of the ALK in analogy to the VEGF-R kinase assay described in J. Wood et al. *Cancer Res.* 60, 2178-2189 (2000). In general, in vitro enzyme assays using GST-ALK protein tyrosine kinase are performed in 96-well plates as a filter binding assay in 20 mM Tris HCl, pH = 7.5, 3 mM MgCl₂, 10 mM MnCl₂, 1 mM DTT, 0.1 µCi/assay (=30 µl) [γ -³³P]-ATP, 2 µM ATP, 3 µg/mL poly (Glu, Tyr 4:1) Poly-EY (Sigma P-0275), 1 % DMSO, 25 ng ALK enzyme. Assays are incubated for 10 min at ambient temperature. Reactions are terminated by adding 50 µl of 125 mM EDTA, and the reaction mixture is transferred onto a MAIP Multiscreen plate (Millipore, Bedford, MA, USA), previously wet with methanol, and rehydrated for 5 min with H₂O. Following washing (0.5 % H₃PO₄), plates are counted in a liquid scintillation counter. IC₅₀ values are calculated by linear regression analysis of the percentage inhibition.

[0050] The compounds of the invention may potently inhibit the growth of human NPM-ALK overexpressing murine BaF3 cells (DSMZ Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Germany). The expression of NPM-ALK may be achieved by transfecting the BaF3 cell line with an expression vector pCIneoTM (Promega Corp., Madison WI, USA) coding for NPM-ALK and subsequent selection of G418 resistant cells. Non-transfected BaF3 cells depend on IL-3 for cell survival. In contrast, NPM-ALK expressing BaF3 cells (named BaF3-NPM-ALK hereinafter) can proliferate in the absence of IL-3 because they obtain proliferative signal through NPM-ALK kinase. Putative inhibitors of the NPM-ALK kinase therefore abolish the growth signal and may result in antiproliferative activity. The antiproliferative activity of putative inhibitors of the NPM-ALK kinase can however be overcome by addition of IL-3, which provides growth signals through an NPM-

ALK independent mechanism. An analogous cell system using FLT3 kinase has also been described (see, E Weisberg et al. Cancer Cell; 1, 433-443 (2002)).

[0051] The inhibitory activity of the compounds of the invention may be determined as follows. In general, BaF3-NPM-ALK cells (15,000/microtitre plate well) are transferred to 96-well microtitre plates. Test compounds dissolved in dimethyl sulfoxide (DMSO) are added in a series of concentrations (dilution series) in such a manner that the final concentration of DMSO is not greater than 1 % (v/v). After the addition, the plates are incubated for two days during which the control cultures without test compound are able to undergo two cell-division cycles. The growth of the BaF3-NPM-ALK cells is measured by means of YOPROTM staining [T Idziorek et al. J. Immunol. Methods; 185: 249-258 (1995)]: 25 µl of lysis buffer comprising 20 mM sodium citrate, pH 4.0, 26.8 mM sodium chloride, 0.4 % NP40 and 20 mM EDTA is added to each well. Cell lysis is completed within 60 min at room temperature and total amount of YOPROTM bound to DNA is determined by measurement using the Cytofluor II 96-well reader (PerSeptive Biosystems) with the following settings: Excitation (nm) 485/20 and Emission (nm) 530/25.

[0052] The compounds of the invention may also be useful in the treatment and/or prevention of acute or chronic inflammatory diseases or disorders or autoimmune diseases e.g. rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, diabetes (type I and II) and the disorders associated therewith, respiratory diseases such as asthma or inflammatory liver injury, inflammatory glomerular injury, cutaneous manifestations of immunologically-mediated disorders or illnesses, inflammatory and hyperproliferative skin diseases (such as psoriasis, atopic dermatitis, allergic contact dermatitis, irritant contact dermatitis and further eczematous dermatitis, seborrhoeic dermatitis), inflammatory eye diseases, e.g. Sjogren's syndrome, keratoconjunctivitis or uveitis, inflammatory bowel disease, Crohn's disease or ulcerative colitis.

[0053] In accordance with the foregoing, the present invention provides:

- (1) a compound of the invention for use as a pharmaceutical;
- (2) a compound of the invention for use as an IGF-1R inhibitor, for example for use in any of the particular indications hereinbefore set forth;
- (3) a pharmaceutical composition, e.g. for use in any of the indications herein before set forth, comprising a compound of the invention as active ingredient together

with one or more pharmaceutically acceptable diluents or carriers;

(4) a method for the treatment of any particular indication set forth hereinbefore in a subject in need thereof which comprises administering an effective amount of a compound of the invention or a pharmaceutical composition comprising same;

(5) the use of a compound of the invention for the manufacture of a medicament for the treatment or prevention of a disease or condition in which IGF-1R activation plays a role or is implicated;

(6) the method as defined above under (4) comprising co-administration, e.g. concomitantly or in sequence, of a therapeutically effective amount of a compound of the invention and one or more further drug substances, said further drug substance being useful in any of the particular indications set forth hereinbefore;

(7) a combination comprising a therapeutically effective amount of a compound of the invention and one or more further drug substances, said further drug substance being useful in any of the particular indications set forth hereinbefore;

(8) use of a compound of the invention for the manufacture of a medicament for the treatment or prevention of a disease which responds to inhibition of the anaplastic lymphoma kinase;

(9) the use according to (8), wherein the disease to be treated is selected from anaplastic large cell lymphoma, non-Hodgkin's lymphomas, inflammatory myofibroblastic tumors, neuroblastomas and neoplastic diseases;

(10) the use according to (8) or (9), wherein the compound is any one of the examples, or a pharmaceutically acceptable salt thereof;

(11) a method for the treatment of a disease which responds to inhibition of the anaplastic lymphoma kinase, especially a disease selected from anaplastic large-cell lymphoma, non Hodgkin's lymphomas, inflammatory myofibroblastic tumors, neuroblastomas and neoplastic diseases, comprising administering an effective amount of a compound of the invention or a pharmaceutically acceptable salt thereof.

Administration and Pharmaceutical Compositions

[0054] In general, compounds of the invention will be administered in therapeutically effective amounts via any of the usual and acceptable modes known in the art, either singly or in combination with one or more therapeutic agents. A therapeutically effective amount may vary widely depending on the severity of the

disease, the age and relative health of the subject, the potency of the compound used and other factors known to those of ordinary skill in the art. For example, for the treatment of neoplastic diseases and immune system disorders, the required dosage will also vary depending on the mode of administration, the particular condition to be treated and the effect desired.

[0055] In general, satisfactory results are indicated to be obtained systemically at daily dosages of from about 0.01 to about 100 mg/kg per body weight, or particularly, from about 0.03 to 2.5 mg/kg per body weight. An indicated daily dosage in the larger mammal, e.g. humans, may be in the range from about 0.5 mg to about 2000 mg, or more particularly, from about 0.5 mg to about 100 mg, conveniently administered, for example, in divided doses up to four times a day or in retard form. Suitable unit dosage forms for oral administration comprise from ca. 1 to 50 mg active ingredient.

[0056] Compounds of the invention may be administered as pharmaceutical compositions by any conventional route; for example, enterally, e.g., orally, e.g., in the form of tablets or capsules; parenterally, e.g., in the form of injectable solutions or suspensions; or topically, e.g., in the form of lotions, gels, ointments or creams, or in a nasal or suppository form.

[0057] Pharmaceutical compositions comprising a compound of the present invention in free form or in a pharmaceutically acceptable salt form in association with at least one pharmaceutically acceptable carrier or diluent may be manufactured in a conventional manner by mixing, granulating, coating, dissolving or lyophilizing processes. For example, pharmaceutical compositions comprising a compound of the invention in association with at least one pharmaceutical acceptable carrier or diluent may be manufactured in conventional manner by mixing with a pharmaceutically acceptable carrier or diluent. Unit dosage forms for oral administration contain, for example, from about 0.1 mg to about 500 mg of active substance.

[0058] In one embodiment, the pharmaceutical compositions are solutions of the active ingredient, including suspensions or dispersions, such as isotonic aqueous solutions. In the case of lyophilized compositions comprising the active ingredient alone or together with a carrier such as mannitol, dispersions or suspensions can be made up before use. The pharmaceutical compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents,

solution promoters, salts for regulating the osmotic pressure and/or buffers. Suitable preservatives include but are not limited to antioxidants such as ascorbic acid, or microbicides, such as sorbic acid or benzoic acid. The solutions or suspensions may further comprise viscosity-increasing agents, including but not limited to, sodium carboxymethylcellulose, carboxymethylcellulose, dextran, polyvinylpyrrolidone, gelatins, or solubilizers, e.g. Tween 80 (polyoxyethylene(20)sorbitan mono-oleate).

[0059] Suspensions in oil may comprise as the oil component the vegetable, synthetic, or semi-synthetic oils customary for injection purposes. Examples include liquid fatty acid esters that contain as the acid component a long-chained fatty acid having from 8 to 22 carbon atoms, or in some embodiments, from 12 to 22 carbon atoms. Suitable liquid fatty acid esters include but are not limited to lauric acid, tridecylic acid, myristic acid, pentadecylic acid, palmitic acid, margaric acid, stearic acid, arachidic acid, behenic acid or corresponding unsaturated acids, for example oleic acid, elaidic acid, erucic acid, brassidic acid and linoleic acid, and if desired, may contain antioxidants, for example vitamin E, 3-carotene or 3,5-di-tert-butyl-hydroxytoluene. The alcohol component of these fatty acid esters may have six carbon atoms and may be monovalent or polyvalent, for example a mono-, di- or trivalent, alcohol. Suitable alcohol components include but are not limited to methanol, ethanol, propanol, butanol or pentanol or isomers thereof; glycol and glycerol.

[0060] Other suitable fatty acid esters include but are not limited ethyl-oleate, isopropyl myristate, isopropyl palmitate, LABRAFIL® M 2375, (polyoxyethylene glycerol), LABRAFIL® M 1944 CS (unsaturated polyglycolized glycerides prepared by alcoholysis of apricot kernel oil and comprising glycerides and polyethylene glycol ester), LABRASOL™ (saturated polyglycolized glycerides prepared by alcoholysis of TCM and comprising glycerides and polyethylene glycol ester; all available from GaKefosse, France), and/or MIGLYOL® 812 (triglyceride of saturated fatty acids of chain length C₈ to C₁₂ from Hüls AG, Germany), and vegetable oils such as cottonseed oil, almond oil, olive oil, castor oil, sesame oil, soybean oil, or groundnut oil.

[0061] Pharmaceutical compositions for oral administration may be obtained, for example, by combining the active ingredient with one or more solid carriers, and if

desired, granulating a resulting mixture, and processing the mixture or granules by the inclusion of additional excipients, to form tablets or tablet cores.

[0062] Suitable carriers include but are not limited to fillers, such as sugars, for example lactose, saccharose, mannitol or sorbitol, cellulose preparations, and/or calcium phosphates, for example tricalcium phosphate or calcium hydrogen phosphate, and also binders, such as starches, for example corn, wheat, rice or potato starch, methylcellulose, hydroxypropyl methylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone, and/or, if desired, disintegrators, such as the above-mentioned starches, carboxymethyl starch, crosslinked polyvinylpyrrolidone, alginic acid or a salt thereof, such as sodium alginate. Additional excipients include flow conditioners and lubricants, for example silicic acid, talc, stearic acid or salts thereof, such as magnesium or calcium stearate, and/or polyethylene glycol, or derivatives thereof.

[0063] Tablet cores may be provided with suitable, optionally enteric, coatings through the use of, inter alia, concentrated sugar solutions which may comprise gum arabic, talc, polyvinylpyrrolidone, polyethylene glycol and/or titanium dioxide, or coating solutions in suitable organic solvents or solvent mixtures, or, for the preparation of enteric coatings, solutions of suitable cellulose preparations, such as acetylcellulose phthalate or hydroxypropylmethylcellulose phthalate. Dyes or pigments may be added to the tablets or tablet coatings, for example for identification purposes or to indicate different doses of active ingredient.

[0064] Pharmaceutical compositions for oral administration may also include hard capsules comprising gelatin or soft-sealed capsules comprising gelatin and a plasticizer, such as glycerol or sorbitol. The hard capsules may contain the active ingredient in the form of granules, for example in admixture with fillers, such as corn starch, binders, and/or glidants, such as talc or magnesium stearate, and optionally stabilizers. In soft capsules, the active ingredient may be dissolved or suspended in suitable liquid excipients, such as fatty oils, paraffin oil or liquid polyethylene glycols or fatty acid esters of ethylene or propylene glycol, to which stabilizers and detergents, for example of the polyoxyethylene sorbitan fatty acid ester type, may also be added.

[0065] Pharmaceutical compositions suitable for rectal administration are, for example, suppositories comprising a combination of the active ingredient and a

suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, paraffin hydrocarbons, polyethylene glycols or higher alkanols.

[0066] Pharmaceutical compositions suitable for parenteral administration may comprise aqueous solutions of an active ingredient in water-soluble form, for example of a water-soluble salt, or aqueous injection suspensions that contain viscosity-increasing substances, for example sodium carboxymethylcellulose, sorbitol and/or dextran, and, if desired, stabilizers. The active ingredient, optionally together with excipients, can also be in the form of a lyophilizate and can be made into a solution before parenteral administration by the addition of suitable solvents. Solutions such as are used, for example, for parenteral administration can also be employed as infusion solutions. The manufacture of injectable preparations is usually carried out under sterile conditions, as is the filling, for example, into ampoules or vials, and the sealing of the containers.

[0067] The compounds of the invention may be administered as the sole active ingredient, or together with other drugs useful against neoplastic diseases or useful in immunomodulating regimens. For example, the compounds of the invention may be used in accordance with the invention in combination with pharmaceutical compositions effective in various diseases as described above, e.g. with cyclophosphamide, 5-fluorouracil, fludarabine, gemcitabine, cisplatin, carboplatin, vincristine, vinblastine, etoposide, irinotecan, paclitaxel, docetaxel, rituxan, doxorubicine, gefitinib, or imatinib; or also with cyclosporins, rapamycins, ascomycins or their immunosuppressive analogs, e.g. cyclosporin A, cyclosporin G, FK-506, sirolimus or everolimus, corticosteroids, e.g. prednisone, cyclophosphamide, azathioprene, methotrexate, gold salts, sulfasalazine, antimalarials, brequinar, leflunomide, mizoribine, mycophenolic acid, mycophenolate, mofetil, 15-deoxyspergualine, immuno-suppressive monoclonal antibodies, e.g. monoclonal antibodies to leukocyte receptors, e.g. MHC, CD2, CD3, CD4, CD7, CD25, CD28, I CD40, CD45, CD58, CD80, CD86, CD152, CD137, CD154, ICOS, LFA-1, VLA-4 or their ligands, or other immunomodulatory compounds, e.g. CTLA41g.

[0068] The invention also provides for a pharmaceutical combinations, e.g. a kit, comprising a) a first agent which is a compound of the invention as disclosed herein, in free form or in pharmaceutically acceptable salt form, and b) at least one co-agent. The kit can comprise instructions for its administration.

Processes for Making Compounds of the Invention

[0069] General procedures for preparing compounds of the invention are described in the Examples, *infra*. In the reactions described, reactive functional groups, for example hydroxy, amino, imino, thio or carboxy groups, where these are desired in the final product, may be protected to avoid their unwanted participation in the reactions. Conventional protecting groups may be used in accordance with standard practice (see e.g., T.W. Greene and P. G. M. Wuts in “Protective Groups in Organic Chemistry”, John Wiley and Sons, 1991).

[0070] The compounds of the invention, including their salts, are also obtainable in the form of hydrates, or their crystals may include for example the solvent used for crystallization (present as solvates). Salts can usually be converted to compounds in free form, e.g., by treating with suitable basic agents, for example with alkali metal carbonates, alkali metal hydrogen carbonates, or alkali metal hydroxides, such as potassium carbonate or sodium hydroxide. A compound of the invention in a base addition salt form may be converted to the corresponding free acid by treating with a suitable acid (e.g., hydrochloric acid, etc.). In view of the close relationship between the novel compounds in free form and those in the form of their salts, including those salts that may be used as intermediates, for example in the purification or identification of the novel compounds, any reference to the free compounds is to be understood as referring also to the corresponding salts, as appropriate.

[0071] Salts of the inventive compounds with a salt-forming group may be prepared in a manner known per se. Acid addition salts of compounds of Formula (1), (2), (3) and (4), may thus be obtained by treatment with an acid or with a suitable anion exchange reagent. Pharmaceutically acceptable salts of the compounds of the invention may be formed, for example, as acid addition salts, with organic or inorganic acids, from compounds of Formula (1), (2), (3) and (4), with a basic nitrogen atom.

[0072] Suitable inorganic acids include, but are not limited to, halogen acids, such as hydrochloric acid, sulfuric acid, or phosphoric acid. Suitable organic acids include, but are not limited to, carboxylic, phosphoric, sulfonic or sulfamic acids, for example acetic acid, propionic acid, octanoic acid, decanoic acid, dodecanoic acid, glycolic acid, lactic acid, fumaric acid, succinic acid, adipic acid, pimelic acid, suberic acid, azelaic acid, -malic acid, tartaric acid, citric acid, amino acids, such as glutamic acid or

aspartic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, cyclohexanecarboxylic acid, adamantanecarboxylic acid, benzoic acid, salicylic acid, 4 aminosalicylic acid, phthalic acid, phenylacetic acid, mandelic acid, cinnamic acid, methane-or ethane-sulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 1,5-naphthalene-disulfonic acid, 2-, 3-or 4 methylbenzenesulfonic acid, methylsulfuric acid, ethylsulfuric acid, dodecylsulfuric acid, N cyclohexylsulfamic acid, N-methyl-, N-ethyl-or N-propyl-sulfamic acid, or other organic protonic acids, such as ascorbic acid. For isolation or purification purposes, it is also possible to use pharmaceutically unacceptable salts, for example picrates or perchlorates. For therapeutic use, only pharmaceutically acceptable salts or free compounds are employed (where applicable in the form of pharmaceutical preparations).

[0073] Compounds of the invention in unoxidized form may be prepared from N-oxides of compounds of the invention by treating with a reducing agent (e.g., sulfur, sulfur dioxide, triphenyl phosphine, lithium borohydride, sodium borohydride, phosphorus trichloride, tribromide, or the like) in a suitable inert organic solvent (e.g. acetonitrile, ethanol, aqueous dioxane, or the like) at 0 to 80°C.

[0074] Prodrug derivatives of the compounds of the invention may be prepared by methods known to those of ordinary skill in the art (e.g., for further details see Saulnier et al., (1994), Bioorganic and Medicinal Chemistry Letters, Vol. 4, p. 1985). For example, appropriate prodrugs may be prepared by reacting a non-derivatized compound of the invention with a suitable carbamylating agent (e.g., 1,1-acyloxyalkylcarbanochloridate, para-nitrophenyl carbonate, or the like).

[0075] Protected derivatives of the compounds of the invention may be made by means known to those of ordinary skill in the art. A detailed description of techniques applicable to the creation of protecting groups and their removal may be found in T. W. Greene, "Protecting Groups in Organic Chemistry", 3rd edition, John Wiley and Sons, Inc., 1999.

[0076] Compounds of the invention may be prepared as their individual stereoisomers by reacting a racemic mixture of the compound with an optically active resolving agent to form a pair of diastereoisomeric compounds, separating the diastereomers and recovering the optically pure enantiomers. Resolution of enantiomers may be carried out using covalent diastereomeric derivatives of the

compounds of the invention, or by using dissociable complexes (e.g., crystalline diastereomeric salts). Diastereomers have distinct physical properties (e.g., melting points, boiling points, solubilities, reactivity, etc.) and may be readily separated by taking advantage of these dissimilarities. The diastereomers may be separated by fractionated crystallization, chromatography, or by separation/resolution techniques based upon differences in solubility. The optically pure enantiomer is then recovered, along with the resolving agent, by any practical means that would not result in racemization. A more detailed description of the techniques applicable to the resolution of stereoisomers of compounds from their racemic mixture may be found in Jean Jacques, Andre Collet, Samuel H. Wilen, "Enantiomers, Racemates and Resolutions", John Wiley And Sons, Inc., 1981.

[0077] In summary, the compounds of the invention may be made by a process as described in the Examples; and


- (a) optionally converting a compound of the invention into a pharmaceutically acceptable salt;
- (b) optionally converting a salt form of a compound of the invention to a non-salt form;
- (c) optionally converting an unoxidized form of a compound of the invention into a pharmaceutically acceptable N-oxide;
- (d) optionally converting an N-oxide form of a compound of the invention to its unoxidized form;
- (e) optionally resolving an individual isomer of a compound of the invention from a mixture of isomers;
- (f) optionally converting a non-derivatized compound of the invention into a pharmaceutically acceptable prodrug derivative; and
- (g) optionally converting a prodrug derivative of a compound of the invention to its non-derivatized form.

[0078] Insofar as the production of the starting materials is not particularly described, the compounds are known or can be prepared analogously to methods known in the art or as disclosed in the Examples hereinafter. One of skill in the art will appreciate that the above transformations are only representative of methods for preparation of the compounds of the present invention, and that other well known methods can similarly be used. The present invention is further exemplified, but not

limited, by the following and Examples that illustrate the preparation of the compounds of the invention.

Intermediate 1


2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine




[0079] A mixture of 5-methyl-1H-pyrazol-3-amine (3.00 g, 30.9 mmol), 2,4,5-trichloropyrimidine (5.67 g, 30.9 mmol, 1 equiv.) and Na₂CO₃ (3.60 g, 34.0 mmol, 1.1 equiv.) in EtOH (100 mL) was heated at 40 °C for 24 h. The solvent was removed *in vacuo*. The resulting residue was partitioned between EtOAc (350 mL) and water (100 mL). The EtOAc layer was washed with water (3x), saturated aqueous NaCl (1x) and dried over Na₂SO₄. The resulting EtOAc solution was concentrated *in vacuo*, providing the product 2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine; ESMS *m/z* 244.0 (M + H⁺).

Intermediate 2


2-chloro-N-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-4-amine



[0080] A mixture of 2,4-dichloro-5-(trifluoromethyl)pyrimidine (1.06 g, 4.86 mmol), 5-methyl-1H-pyrazol-3-amine (472.2 mg, 4.86 mmol) and sodium carbonate (2.06 g, 19.4 mmol) in 100 mL of EtOH was stirred at room temperature overnight. The reaction mixture was concentrated *in vacuo*. The crude solid was partitioned between EtOAc and water. The combined organic extracts were dried (Na₂SO₄), concentrated *in vacuo*, and purified by silica chromatography (EtOAc/ hexanes: 1/1) to afford 2-chloro-N-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-4-amine; ESMS *m/z* 278.0 (M + H⁺).


Intermediate 32-chloro-5-methyl-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine

[0081] A mixture of 5-methyl-1H-pyrazol-3-amine (3.00 g, 30.9 mmol), 2,4-dichloro-5-methylpyrimidine (5.03 g, 30.9 mmol, 1 equiv.) and Na₂CO₃ (3.60 g, 34.0 mmol, 1.1 equiv.) in EtOH (100 mL) was heated at 40 °C for 24 h. The solvent was removed *in vacuo*. The resulting residue was partitioned between EtOAc (350 mL) and water (100 mL). The EtOAc layer was washed with water (3x), saturated aqueous NaCl (1x), dried over Na₂SO₄, and concentrated *in vacuo*. The resulting crude product was sonicated in Et₂O (200 mL) and the resulting precipitate collected by filtration. This powder was further washed with Et₂O, providing the product 2-chloro-5-methyl-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine; ESMS *m/z* 224.1 (M + H⁺).

Intermediate 4N-(2,5-dichloropyrimidin-4-yl)-5-methylisoxazol-3-amine

[0082] The mixture 5-methylisoxazol-3-amine (98 mg, 1.0 mmol), 2,4,5-trichloropyrimidine (344 µL, 3.0 mmol), and sodium carbonate (106 mg, 1.0 mmol) in 3 mL of EtOH was heated at 60°C overnight. The reaction mixture was concentrated and then partitioned between EtOAc and brine. The collected organic extracts were dried (Na₂SO₄), concentrated *in vacuo*, and purified with silica gel chromatography (MeOH/DCM: 1/9) to afford N-(2,5-dichloropyrimidin-4-yl)-5-methylisoxazol-3-amine; ESMS *m/z* 245.0 (M + H⁺).

Intermediate 52,5-dichloro-N-(5-cyclopropyl-1H-pyrazol-3-yl)pyrimidin-4-amine



[0083] A mixture of 5-cyclopropyl-1H-pyrazol-3-amine (246 mg, 2.00 mmol), 2,4,5-trichloropyrimidine (367 mg, 2.00 mmol, 1 equiv.) and Na_2CO_3 (233 mg, 2.20 mmol, 1.1 equiv.) in EtOH (10 mL) was heated at 40 °C for 16 h. The crude reaction mixture was diluted with EtOAc and sequentially washed with: water (3x) and saturated aqueous NaCl (1x). The resulting EtOAc layer was dried over Na_2SO_4 and then concentrated *in vacuo*, providing 2,5-dichloro-N-(5-cyclopropyl-1H-pyrazol-3-yl)pyrimidin-4-amine; ESMS m/z 270.0 ($\text{M} + \text{H}^+$).

Intermediate 6


3-(2,5-dichloropyrimidin-4-ylamino)azepan-2-one



[0084] A mixture of ($\pm$)- α -amino- ϵ -caprolactam (256 mg, 2.0 mmol), 2,4,5-trichloropyrimidine (366 mg, 2.0 mmol, 1 equiv.) and NaHCO_3 (168 mg, 2 mmol, 1 equiv.) in a mixture of MeOH (12 mL) and H_2O (6 mL) was stirred at room temperature for 15 h. The resulting precipitate was collected by vacuum filtration and washed with small amounts of MeOH and water, providing the product 3-(2,5-dichloropyrimidin-4-ylamino)azepan-2-one; ESMS m/z 275.0 ($\text{M} + \text{H}^+$).

Intermediate 7

3-(2,5-dichloropyrimidin-4-ylamino)pyridin-2(1H)-one




[0085] A mixture of 3-aminopyridin-2(1H)-one (99 mg, 0.90 mmol), 2,4,5-trichloropyrimidine (165 mg, 0.90 mmol, 1 equiv.) and NaHCO_3 (76 mg, 0.90 mmol, 1 equiv.) in a mixture of MeOH (6 mL) and H_2O (3 mL) was stirred at room

temperature for 24 h. The resulting precipitate was collected by vacuum filtration and washed with small amounts of MeOH and water, providing the product 3-(2,5-dichloropyrimidin-4-ylamino)pyridin-2(1H)-one; ESMS m/z 257.0 ($M + H^+$).

Intermediate 8


2,5-Dichloro-N-(5-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-3-yl)pyrimidin-4-amine



[0086] To a mixture of 2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine (2.44 g, 10 mmol) and 4-toluenesulfonic acid monohydrate (1.9g, 10 mmol) in THF (200 mL), was added DHP (4.25g, 50 mmol). After stirring overnight, the reaction turned clear. After concentration, the residue was dissolved in ethyl acetate and washed with Na_2CO_3 saturated aqueous solution. The organic layer was then washed with brine, dried over sodium sulfate and concentrated *in vacuo* to afford 2,5-Dichloro-N-(5-methyl-1-(tetrahydro-2H-pyran-2-yl)-1H-pyrazol-3-yl)pyrimidin-4-amine as light yellow solid, which was used in subsequent reactions without further purification; ESMS m/z 244 ($M\text{-THP}+H^+$); TLC $R_f=0.6$ (Silica; 1:1 ethyl acetate/hexanes; starting material $R_f=0.2$).

Intermediate 9

2,5-Dichloro-N-(5-(trifluoromethyl)-1H-pyrazol-3-yl)pyrimidin-4-amine



Step 1: 5-(Trifluoromethyl)-1H-pyrazol-3-amine

[0087] A mixture of (*E*)-4-amino-4-ethoxy-1,1,1-trifluorobut-3-en-2-one (5.00 g, 27 mmol) and hydrazine (1.05 g) in anhydrous EtOH (20 mL) was stirred at 80 °C in a sealed vial overnight. The reaction was quenched with water (50 mL), and extracted with EtOAc (100ml, then 2x50 mL). The EtOAc layers were combined and


sequentially washed with water (25 mL), brine (25 mL), dried over Na₂SO₄ and evaporated to give a light brown oily residue. The crude product was purified by silica chromatography (30-100% EtOAc in hexanes gradient) to give 5-(trifluoromethyl)-1*H*-pyrazol-3-amine as an off white solid; ESMS *m/z* 152.0 (M+H⁺).

Step 2: 2,5-Dichloro-*N*-(5-(trifluoromethyl)-1*H*-pyrazol-3-yl)pyrimidin-4-amine

[0088] A mixture of 5-(trifluoromethyl)-1*H*-pyrazol-3-amine (500 mg, 3.3 mmol), 2,4,5-trichloropyrimidine (607 mg, 3.3 mmol) and sodium carbonate (525 mg, 5 mmol) in anhydrous EtOH (10 mL) was stirred at rt. After one day, LCMS showed the reaction was not complete. Additional 2,4,5-trichloropyrimidine (0.15 mL) and sodium carbonate (525 mg) were added and the reaction continued for two more days. The solvent was evaporated, and the residue was extracted with DCM (3x20 mL). The combined DCM layers were washed with brine (10 mL), dried over Na₂SO₄ and evaporated. The crude product was purified by silica chromatography (0-50% EtOAc in hexanes gradient) to give 2,5-Dichloro-*N*-(5-(trifluoromethyl)-1*H*-pyrazol-3-yl)pyrimidin-4-amine as a light tan colored solid; ESMS *m/z* 297.9 (M+H⁺).

Intermediate 10


2,5-Dichloro-*N*-(5-ethyl-1*H*-pyrazol-3-yl)pyrimidin-4-amine



[0089] A mixture of 5-ethyl-1*H*-pyrazol-3-amine (1.00 g, 6.77 mmol), 2,4,5-trichloropyrimidine (1.24 g, 6.77 mmol) and Na₂CO₃ (1.79 g, 16.9 mmol) in *iso*-propanol (15 mL) was stirred in a sealed vial at rt for two days. Water (10 mL) was added. The solid was collected by filtration, washed with water (10 mL) and dried to give 2,5-Dichloro-*N*-(5-ethyl-1*H*-pyrazol-3-yl)pyrimidin-4-amine as a light yellow solid; ESMS *m/z* 258.0 (M+H⁺).

Intermediate 11


1-chloro-5-methyl-4-nitro-2-(trifluoromethyl)benzene



[0090] To a solution of 2-chloro-4-methyl-1-(trifluoromethyl)benzene (6.65g, 34.1 mmol) in conc. sulfuric acid (25 mL) at 0 °C was added potassium nitrate (3.46 g, 34.1 mmol) in sulfuric acid (15 mL). After stirring at 0 °C for 1.5 h, the reaction mixture was poured into 500 mL of ice water. The precipitate was collected by filtration and washed with copious amounts of water, yielding 1-chloro-5-fluoro-4-nitro-2-(trifluoromethyl)benzene as an orange solid; ¹H NMR (400 MHz, chloroform-d) δ 8.356 (s, 1H), 7.543 (s, 1H), 2.660 (s, 3H).

Intermediate 12


2-Bromo-4-methyl-5-nitrobenzonitrile



[0091] Following the same procedure as described in Intermediate 11 starting with 2-bromo-4-methylbenzonitrile, 2-bromo-4-methyl-5-nitrobenzonitrile was obtained as a yellow solid; ¹H NMR (400 MHz, chloroform-d) δ 8.290 (s, 1H), 7.742 (s, 1H), 2.683 (s, 3H).

Intermediate 13

5-Amino-2-bromo-4-methylbenzonitrile




[0092] To a solution of 2-bromo-4-methyl-5-nitrobenzonitrile (500mg, 2.07 mmol) in ethanol (20mL) was added tin chloride (785mg, 4.14 mmol). The mixture was refluxed for 5 h. After cooling to room temperature, triethylamine (1.01g, 10 mmol) was added. The mixture was concentrated and purified by silica

chromatography (30-50% ethyl acetate in hexanes gradient) to afford 5-Amino-2-bromo-4-methylbenzonitrile a as beige solid; ESMS m/z 211 ($M + H^+$).

Intermediate 14


Methyl 2-bromo-4-methyl-5-nitrobenzoate



[0093] Following the same procedure as described in Intermediate 11 starting with methyl 2-bromo-4-methylbenzoate, the title compound was obtained as a white solid; ^1H NMR (400 MHz, chloroform- d) δ 8.505 (s, 1H), 7.711 (s, 1H), 3.967 (s, 3H), 2.640 (s, 3H).

Intermediate 15


Methyl 5-amino-2-bromo-4-methylbenzoate



[0094] Following the same procedure as described in Intermediate 13, the title compound was obtained as a beige solid; ESMS m/z 244 ($M + H^+$).

Intermediate 16

4-Bromo-5-methoxy-2-methylaniline



Step 1: N-(5-methoxy-2-methylphenyl)acetamide

[0095] To a solution of triethylamine (2.9 g, 14.4 mmol) and 5-methoxy-2-methylaniline (1.0 g, 7.2 mmol) in DCM (30 mL) at 0 °C, was added acetyl chloride

(0.57 g, 7.2 mmol) dropwise. The mixture was stirred at room temperature for 1 hour. The reaction was quenched with saturated ammonium chloride aqueous solution, and extracted with DCM. The organic layer was dried over Na₂SO₄ and concentrated to give N-(5-methoxy-2-methylphenyl)acetamide as needle-shaped crystals. The crude product was used in the next step without further purification.

Step 2: N-(4-bromo-5-methoxy-2-methylphenyl)acetamide


[0096] To a solution of N-(5-methoxy-2-methylphenyl)acetamide in acetic acid (40 mL) was added Br₂ (3 g, 19 mmol) slowly. The mixture was capped and stirred at 50 °C for 5 hours. The reaction was cooled to room temperature, then quenched with sodium sulfite aqueous solution, and extracted with ethyl acetate (3 x 50 mL). The combined organic layer was dried over Na₂SO₄ and concentrated to give crude N-(4-bromo-5-methoxy-2-methylphenyl)acetamide as a brown oil, which was used in the next step without further purification.

Step 3: 4-bromo-5-methoxy-2-methylaniline

[0097] N-(4-bromo-5-methoxy-2-methylphenyl)acetamide was dissolved in methanol (15 mL) and concentrated HCl (30 mL). The mixture was refluxed at 95 °C overnight. After cooling to room temperature, the mixture was poured into ice water, and basified to pH = 12 with conc. NaOH aqueous solution. The mixture was then extracted with ethyl acetate (3 x 50 mL). The combined organic layer was dried over Na₂SO₄ and concentrated. The crude product was purified by silica chromatography (15% ethyl acetate in hexanes) to give 4-bromo-5-methoxy-2-methylaniline as a beige solid; ¹H NMR (400 MHz, CDCl₃) 7.20 (s, 1H), 6.37 (s, 1H), 4.6 (br, 2H), 3.82 (s, 3H), 2.12 (s, 3H).

Intermediate 17

2-Fluoro-5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline




[0098] A mixture of 4-bromo-2-fluoro-5-methylaniline (2.0 g, 9.8 mmol),

bis(pinacolato)diboron (2.74 g, 10.8 mmol), tricyclohexylphosphine (274.8 mg, 0.98 mmol), $\text{Pd}_2(\text{dba})_3$ (448.9 mg, 0.49 mmol), and potassium acetate (1.92 g, 19.6 mmol) in 60 mL of 1,4-dioxane was degassed and purged with nitrogen. The reaction mixture was heated at 80°C overnight and then cooled to room temperature. The mixture was partitioned between EtOAc and water and the collected organic extracts were dried (Na_2SO_4), concentrated *in vacuo*, and purified by silica chromatography (EtOAc / Hexanes: 20/ 80) to afford 2-fluoro-5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline; ESMS m/z 252.2 ($\text{M} + \text{H}^+$).

Intermediate 18

2-Isopropoxy-5-methyl-4-(1-methyl-piperidin-4-yl)-phenylamine



Step 1: 1-bromo-5-fluoro-2-methyl-4-nitrobenzene

[0099] To a solution of 2-bromo-4-fluoro-1-methylbenzene (5 g, 26.4 mmol) dissolved in conc. sulfuric acid (20 mL) was added potassium nitrate (2.67 g, 26.4 mmol) in sulfuric acid (5 mL) at 0 °C. After stirring at 0 °C for 1hr, the reaction was diluted with water, extracted with ethyl acetate (3 x 40 mL). The combined organic phases were washed with saturated sodium bicarbonate solution and brine sequentially, and then dried over Na_2SO_4 . The crude product obtained after concentration was purified by silica gel column chromatography with mixed solvent hexanes/EtOAc (95/5) to afford 1-bromo-5-fluoro-2-methyl-4-nitrobenzene as an orange solid. ^1H NMR (400 MHz, chloroform- d) δ 7.93 (d, 1H, $J = 7.6$ Hz), 7.50 (d, 1H, $J = 10$ Hz), 2.43 (s, 3H).

Step 2: 1-bromo-5-isopropoxy-2-methyl-4-nitrobenzene

[0100] To a solution of 1-bromo-5-fluoro-2-methyl-4-nitrobenzene (Step 1, 1.5 g, 6.4 mmol) in 2-propanol (30 mL) was added cesium carbonate (5.3 g, 16 mmol). The mixture was heated at 50 °C overnight. The mixture was concentrated and partitioned between ethyl acetate and water. The organic layer was concentrated and the residue

was purified with silica gel column chromatography with 5% ethyl acetate in hexanes to afford 1-bromo-5-isopropoxy-2-methyl-4-nitrobenzene as a yellow solid.

Step 3: 4-(5-Isopropoxy-2-methyl-4-nitrophenyl)pyridine

[0101] To a mixture of 1-bromo-5-isopropoxy-2-methyl-4-nitrobenzene (Step 2, 1g, 3.65 mmol), pyridine-4-boronic acid (490 mg, 4 mmol), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (300 mg, 0.73 mmol) and potassium phosphate (1.55g, 7.3 mmol) in mixed solvent of dioxane (15 mL) and water (7.5 mL) was added tris(dibenzylidene-acetone)dipalladium (0) (334mg, 0.36 mmol). This mixture was sealed and purged with nitrogen for 3 minutes and then heated at 120 °C for 5 hours. The mixture was cooled to room temperature, filtered, and the filtrate was concentrated. After concentration, the crude product was purified with silica gel flash column chromatography (60% ethyl acetate in hexanes) to afford 4-(5-isopropoxy-2-methyl-4-nitrophenyl)pyridine as a yellow solid: ESMS m/z 273.2 ($M + H^+$).

Step 4: 4-(5-Isopropoxy-2-methyl-4-nitro-phenyl)-1-methyl-pyridinium iodide

[0102] 4-(5-Isopropoxy-2-methyl-4-nitro-phenyl)-pyridine (Step 3, 217 mg, 0.797 mmol) was dissolved in anhydrous THF (9 mL). Iodomethane (0.10 mL, 1.61 mmol, 2 equiv.) was added, and the reaction was stirred at 40 °C in a sealed tube for 2 days. The volatiles were removed under vacuum, generating 4-(5-Isopropoxy-2-methyl-4-nitro-phenyl)-1-methyl-pyridinium iodide as a brown solid: ESMS m/z 287.1 (M^+).


Steps 5 and 6: 2-Isopropoxy-5-methyl-4-(1-methyl-piperidin-4-yl)-phenylamine

[0103] 4-(5-Isopropoxy-2-methyl-4-nitro-phenyl)-1-methyl-pyridinium iodide (Step 4, 0.697 mmol) was dissolved in CH_3OH (20 mL) and cooled to 0 °C. $NaBH_4$ (264 mg, 6.97 mmol, 10 equiv.) was slowly added. After this addition was complete, the cooling bath was removed and the reaction was stirred at room temperature for 1 h. The reaction was quenched by the slow addition of 1N aqueous HCl (14 mL). The CH_3OH was partially removed by vacuum. The resulting residue was partitioned between EtOAc and 1 N aqueous NaOH. Additional 50 % aqueous NaOH was added until the aqueous layer had a pH >12. The EtOAc layer was washed with 1 N aqueous NaOH (2x), the organic layer was then dried over sodium sulfate, filtered,

and concentrated under vacuum. After concentration, the crude product (175 mg) was dissolved in acetic acid (10 mL). TFA (0.15 mL, 3 equiv.) and PtO₂ (53 mg, 30% w/w) were added and the reaction was placed under 50 psi H₂ gas in a Parr Shaker for 14 h. The reaction mixture was filtered and the filtrate was concentrated under vacuum. The resulting residue was partitioned between EtOAc and 1 N aqueous NaOH. Additional 50% aqueous NaOH was added until the aqueous layer has a pH >12. The EtOAc layer was washed with 1 N aqueous NaOH (2x), the organic layer was then dried over sodium sulfate, filtered, and concentrated under vacuum to give 2-isopropoxy-5-methyl-4-(1-methyl-piperidin-4-yl)-phenylamine, which was used in subsequent steps without further purification: ESMS m/z 263.2 (M + H⁺).

Intermediate 19

tert-butyl 4-(4-amino-5-fluoro-2-methylphenyl)piperidine-1-carboxylate



Step 1: 4-Trifluoromethanesulfonyloxy-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl ester

[0104] A solution of *N*-tert-Butoxycarbonyl-4-piperidone (10.17 g, 0.05 mol) in THF (100 mL) was added dropwise into a cooled (-78 °C), vigorously stirring solution of LDA (40 mL of 1.5 M solution in cyclohexanes, 0.06 mol) in THF (100 mL), under N₂. The reaction mixture was left at -78 °C for 30 min before adding a solution of phenyl trifluorosulfonimide (19.85 g, 0.055 mol) in THF (50 mL). Then the reaction mixture was warmed to room temperature and stirred for 3 h. The reaction was quenched at 0 °C with 100 mL of saturated aqueous NH₄Cl and filtered through Celite. The filtrate was added to 100 mL of EtOAc, and the layers were separated. The organic layer was washed with H₂O, dried over MgSO₄ and concentrated. The crude product was purified by silica gel chromatography (0-30% EtOAc in Hexanes as eluent and checked by TLC stained with 2% of KMnO₄ in EtOH) to afford 4-Trifluoromethanesulfonyloxy-3,6-dihydro-2H-pyridine-1-carboxylic acid tert-butyl

ester as a yellow oil.

Step 2: *Tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0105] A solution of *tert*-butyl 4-(trifluoromethylsulfonyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (16.0 g, 48.2 mmol) in DMSO (200 mL) was treated with bis(pinacolato)diboron (12.6 g, 49.6 mmol), potassium acetate (14.65 g, 149 mmol) and Pd(dppf)Cl₂ (790 mg, 0.96 mmol). The solution was purged with N₂ (g) for 5 minutes, and then sealed and heated at 80 °C for 15 h. The reaction was cooled to room temperature and poured into ice water. The mixture was extracted with ethyl acetate (3 x 100 mL). The combined organic phase was washed with brine, dried over sodium sulfate, and concentrated in vacuo. The crude product was purified with silica gel chromatography (15% ethyl acetate in hexanes) to afford *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate as a white solid.


Steps 3 and 4: *Tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)piperidine-1-carboxylate

[0106] To a mixture of 5'-chloro-N,2'-dimethyl-4'-nitrobiphenyl-4-carboxamide (204 mg, 1 mmol), *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (370 mg, 1.2 mmol) and sodium carbonate (742 mg, 7 mmol) in DMF/H₂O (12/3 mL) was added tetrakis(triphenylphosphine)palladium (0) (58 mg, 5% mmol). The reaction tube was sealed, the mixture was purged with N₂ for 3 min and then heated at 90°C under N₂ for overnight. The reaction was cooled to room temperature, and poured into saturated aqueous ammonia chloride solution. The crude reaction mixture was extracted with ethyl acetate (3 x 15 mL). The organic extracts were combined, washed with brine and concentrated. The crude product was purified with silica gel chromatography (20% ethyl acetate in hexanes) to afford *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate as a yellow oil. The obtained oil was dissolved in methanol (20 mL). To the solution was added Pd/C (10%). The reaction mixture was degassed and purged with H₂ for several times and stirred under H₂ (1 atm) overnight. The mixture was filtered and concentrated to afford the *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)piperidine-1-carboxylate as a white solid. ESMS *m/z* 207 (M -Boc+

H⁺).

Intermediate 20


2,5-Dimethyl-4-(piperidin-2-yl)aniline



[0107] To a mixture of 1-bromo-2,5-dimethyl-4-nitrobenzene (185 mg, 1 mmol) and 2-(tributylstannyl)pyridine (202 mg, 1.1 mmol) in DMF (4 mL) was added tetrakis(triphenylphosphine) palladium (0) (58 mg, 5% mmol). The reaction tube was sealed, the mixture was purged with N₂ for 3 min and then heated at 120 °C under N₂ for overnight. The reaction was cooled to room temperature and poured into saturated aqueous ammonia chloride solution. The crude reaction mixture was extracted with ethyl acetate (3 x 15 mL). The organic extracts were combined, washed with brine and concentrated. The crude product was purified with silica gel column chromatography (60% ethyl acetate in hexanes) to afford 2-(2,5-dimethyl-4-nitrophenyl)pyridine as a white solid. The obtained solid was dissolved in acetic acid/TFA (15 mL/200 uL). To this solution was added PtO₂ (10% w/w). The reaction mixture was degassed and purged with H₂ for several times and stirred under 1 atm. hydrogen gas overnight. The mixture was filtered and concentrated to afford 2,5-Dimethyl-4-(piperidin-2-yl)aniline as a yellow oil. ESMS m/z 205 (M + H⁺).

Intermediate 21

tert-butyl 3-(4-amino-2,5-dimethylphenyl)azetidine-1-carboxylate



Step 1: *tert*-Butyl 3-(2,5-dimethyl-4-nitrophenyl)azetidine-1-carboxylate


[0108] Prepared from *tert*-butyl 3-iodoazetidine-1-carboxylate and 1-bromo-2,5-dimethyl-4-nitrobenzene by following the general protocol as described in Billotte, S. *Synlett* 1998, 379.

Step 2: *tert*-butyl 3-(4-amino-2,5-dimethylphenyl)azetidine-1-carboxylate

[0109] *tert*-Butyl 3-(2,5-dimethyl-4-nitrophenyl)azetidine-1-carboxylate was reduced to *tert*-butyl 3-(4-amino-2,5-dimethylphenyl)azetidine-1-carboxylate by standard Raney Ni hydrogenation at room temperature using MeOH as the solvent.

Intermediate 22

5-Ethyl-2-methyl-4-(piperidin-4-yl)aniline



Step 1: *tert*-butyl 4-(4-amino-5-methyl-2-vinylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate

[0110] To a mixture of 4-bromo-5-chloro-2-methylaniline (500 mg, 2.27mmol), *tert*-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (840 mg, 3.73 mmol) and sodium carbonate (2.52 g, 15.9 mmol) in DMF/H₂O (20/5 mL) was added tetrakis(triphenylphosphine) palladium (0) (131 mg, 5 mol%). The reaction tube was sealed, the mixture was purged with N₂ for 3 min and then heated at 100°C under N₂ for overnight. The reaction was cooled to room temperature and poured into saturated aqueous ammonia chloride solution. The crude reaction mixture was extracted with ethyl acetate (3 x 15 mL). The organic extracts were combined, washed with brine and concentrated. The crude product was purified with silica chromatography (80% ethyl acetate in hexanes) to afford *tert*-butyl 4-(4-amino-2-chloro-5-methylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate as a yellow solid. This obtained product (322 mg, 1 mmol) was dissolved in dioxane/NMP (anhydrous, 4 mL, 3/1). To this solution was added tributyl(vinyl)stannane (380 mg, 1.2 mmol), cesium fluoride (304 mg, 2 mmol) and palladium bis(tri-*tert*-butyl-


phosphine) (51 mg, 10 mol%). This mixture was purged with N₂ for 3 minutes and then heated in a sealed tube at 120 °C for overnight. The mixture was cooled to room temperature and diluted with ethyl acetate. The resulting mixture was sequentially washed with saturated aqueous ammonium chloride and brine, and finally dried over sodium sulfate. After concentration, the crude product was purified via silica chromatography (10% ethyl acetate in hexanes) to afford *tert*-butyl 4-(4-amino-5-methyl-2-vinylphenyl)-5,6-dihydropyridine-1(2H)-carboxylate as a light yellow oil; ESMS *m/z* 215.2 (M -Boc + H⁺).

Step 2: 5-Ethyl-2-methyl-4-(piperidin-4-yl)aniline

[0111] The product obtained from the previous step was dissolved in methanol (20 mL). To this solution was added conc. aqueous HCl (200 uL) and platinum oxide (23 mg, 0.1 mmol). The reaction mixture was degassed and purged with H₂ for several times and vigorously stirred under 1 atm. H₂ for 3 h. The mixture was filtered and concentrated to afford 5-Ethyl-2-methyl-4-(piperidin-4-yl)aniline as a yellow solid. ESMS *m/z* 219.2 (M + H⁺).

Intermediate 23

2,5-Dimethyl-4-(piperidin-3-yl)aniline



Step 1: 2,5-Dimethyl-4-(pyridin-3-yl)aniline

[0112] A suspension of 4-bromo-2,5-dimethylaniline (4.00 g, 20 mmol), pyridin-3-ylboronic acid (2.70 g, 11 mmol), Pd₂(dba)₃ (0.55 g, 0.6 mmol), 2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl (0.98 g, 1.2 mmol) and Na₂CO₃ (10.6 g, 100 mmol) in n-BuOH (50 mL) was degassed by a stream of argon gas for 15 min. The reaction flask was sealed and placed in a pre-heated oil bath (115 °C). After stirring overnight, the reaction was cooled and filtered. The filter cake was washed with DCM and the filtrate was concentrated *in vacuo*. The resulting residue was dissolved in EtOAc (150 mL). EtOAc was sequentially washed with water (20 mL),


brine (20 mL), dried over Na₂SO₄ and evaporated. The crude product was purified by silica chromatography (0-50% EtOAc in hexanes gradient) to give 2,5-dimethyl-4-(pyridin-3-yl)aniline as a yellow solid; ESMS *m/z* 199.1 (M+H⁺).

Step 2: 2,5-Dimethyl-4-(piperidin-3-yl)aniline

[0113] 2,5-Dimethyl-4-(pyridin-3-yl)aniline (403 mg, 2.03 mmol) was dissolved in MeOH (5 mL) and conc. aqueous HCl (1 mL), followed by addition of PtO₂ (40 mg). The flask was purged with H₂ and the reaction was vigorously stirred at rt under H₂ (1 atm). After two days, LCMS shows the completion of the reaction. The catalyst was removed by filtration and the remaining solution was concentrated *in vacuo* to give 2,5-Dimethyl-4-(piperidin-3-yl)aniline as a white solid, which can be used in subsequent reactions without further purification; ESMS *m/z* 205.2 (M+H⁺).

Intermediate 24


1-(tert-butoxycarbonyl(methyl)amino)cyclopropanecarboxylic acid



[0114] To a solution of 1-(tert-butoxycarbonylamino)cyclopropanecarboxylic acid (201mg, 1.0 mmol) in DMF (2 mL) was added NaH (120 mg, 3.0 mmol) at 0 °C. The resulting mixture was stirred at 0°C for 10 min. before the addition of MeI (568 mg, 4.0 mmol). The mixture was then warmed up to room temperature and stirred for 2 hr. Saturated NH₄Cl aqueous solution (20 mL) was added into the mixture, and the solution was extracted with EtOAc (3 x 10 mL). The combined organic layers were concentrated and the residue was dissolved in MeOH (3 ml) with NaOH (3 N, 1mL). This mixture was stirred at 80 °C for 1 hr and cooled down to room temperature. The clear solution was purified directly on preparative RP-HPLC to provide 1-(tert-butoxycarbonyl(methyl)amino)cyclopropanecarboxylic acid. ESMS *m/z* 238.2 (M + Na⁺).

Intermediate 25

(R)-2-bromopropanamide



Step 1: (R)-2-bromopropanoyl chloride


[0115] Under nitrogen, to the solution of (R)-2-bromopropanoic acid (5.0 g, 32.68 mmol) in 100 mL of DCM was added thionyl chloride (7.1 mL, 98.04 mmol) and 1 mL of DMF sequentially at 0°C. The reaction was stirred at room temperature overnight, and concentrated *in vacuo*. The resulting crude product was directly used in next step without further purification.

Step 2: (R)-2-bromopropanamide

[0116] (R)-2-bromopropanoyl chloride was slowly added to a 37% ammonia hydroxide water solution cooled to 0°C. The reaction was allowed to warm to room temperature and was stirred for 2 h. The product was extracted with EtOAc and the collected organic extracts were dried (Na₂SO₄) and concentrated *in vacuo* to afford (R)-2-bromopropanamide.

Intermediate 26


(R)-2-bromo-N-methylpropanamide



[0117] (R)-2-bromopropanoyl chloride was slowly added to a methylamine water solution cooled to 0°C. The reaction was allowed to warm to room temperature and was stirred for 2 h. The product was extracted with EtOAc and the collected organic extracts were dried (Na₂SO₄) and concentrated *in vacuo* to afford (R)-2-bromo-N-methylpropanamide.

Intermediate 27

1-(tert-butoxycarbonyl(ethyl)amino)cyclopropanecarboxylic acid




Step 1: Ethyl 1-(tert butoxycarbonyl(ethyl)amino)cyclopropanecarboxylate

[0118] Under nitrogen, to the solution of 1-(*tert*-butoxycarbonylamino)cyclopropanecarboxylic acid (201.2 mg, 1.0 mmol) in 4 mL of DMF was added NaH (120.0 mg, 3.0 mmol) at 0°C. After 30 min stirring, iodoethane (0.4 mL, 5.0 mmol) was added to the reaction. The reaction was warmed gradually to room temperature and stirred overnight. The reaction mixture was partitioned between EtOAc and water. The combined organic extracts were dried (Na₂SO₄), and concentrated *in vacuo* to afford crude ethyl 1-(*tert* butoxycarbonyl(ethyl)amino)cyclopropanecarboxylate, which was used in the next step without further purification.

Step 2: 1-(tert-butoxycarbonyl(ethyl)amino)cyclopropanecarboxylic acid

[0119] The crude product from Step 1 was dissolved in a mixture of EtOH (4.0 mL) and water (1.0 mL). LiOH (82.0 mg, 2.0 mmol) was added to the reaction. The reaction was heated at 120°C in a microwave reactor for 10 min. The reaction was concentrated *in vacuo*, and the crude product was partitioned between EtOAc and water. The water layer was neutralized with 4N aq. HCl to pH 4 and then extracted with EtOAc. The combined organic extracts were dried (Na₂SO₄), and concentrated *in vacuo* to afford 1-(*tert* butoxycarbonyl(ethyl)amino)cyclopropanecarboxylic acid.

Intermediate 281-(tert-butoxycarbonyl(ethyl)amino)cyclobutanecarboxylic acidStep 1: Ethyl 1-(tert-butoxycarbonyl(ethyl)amino)cyclobutanecarboxylate

[0120] Under nitrogen, to the solution of 1-(*tert*-butoxycarbonylamino)cyclobutanecarboxylic acid (430.0 mg, 2.0 mmol) in 4 mL of DMF was added NaH (240.0 mg, 6.0 mmol) at 0°C. After stirring for 30 min., iodoethane (0.8 mL, 10.0 mmol) was added to the reaction. The reaction was warmed to room temperature gradually and stirred overnight. The reaction mixture was partitioned between EtOAc and water. The combined organic extracts were dried


(Na₂SO₄), and concentrated *in vacuo* to afford crude ethyl 1-(*tert*-butoxycarbonyl(ethyl)amino) cyclobutanecarboxylate.

Step 2: 1-(*tert*-butoxycarbonyl(ethyl)amino)cyclobutanecarboxylic acid

[0121] The crude product from Step 1 was dissolved in EtOH (4.0 mL) and water (1.0 mL). LiOH (164.0 mg, 4.0 mmol) was added to the reaction. The reaction was heated at 120°C in a microwave reactor for 10 min. The reaction was concentrated *in vacuo*, and the crude was partitioned between EtOAc and water. The water layer was neutralized with 4N aq. HCl to pH 4 and then extracted with EtOAc. The combined organic extracts were dried (Na₂SO₄), and concentrated *in vacuo* to afford 1-(*tert*-butoxycarbonyl(ethyl)amino) cyclobutanecarboxylic acid.

Intermediate 29

1-(*tert*-butoxycarbonyl(methyl)amino)cyclobutanecarboxylic acid



Step 1: Methyl 1-(*tert*-butoxycarbonyl(ethyl)amino)cyclobutanecarboxylate

[0122] Under nitrogen, to the solution of 1-(*tert*-butoxycarbonylamino) cyclobutanecarboxylic acid (430.0 mg, 2.0 mmol) in 4 mL of DMF was added NaH (240.0 mg, 6.0 mmol) at 0°C. After stirring for 30 min., iodomethane (0.62 mL, 10.0 mmol) was added to the reaction. The reaction was warmed to room temperature gradually and stirred overnight. The reaction mixture was partitioned between EtOAc and water. The combined organic extracts were dried (Na₂SO₄), and concentrated *in vacuo* to afford crude methyl 1-(*tert*-butoxycarbonyl(ethyl)amino) cyclobutanecarboxylate.


Step 2: 1-(*tert*-butoxycarbonyl(methyl)amino)cyclobutanecarboxylic acid

[0123] The crude product from Step 1 was dissolved in MeOH (4.0 mL) and water (1.0 mL). LiOH (164.0 mg, 4.0 mmol) was added to the reaction. The reaction was heated at 120°C in a microwave reactor for 10 min. The reaction was concentrated *in vacuo*, and the crude was partitioned between EtOAc and water. The water layer was neutralized with 4N aq. HCl to pH 4, and then extracted with EtOAc.

The combined organic extracts were dried (Na_2SO_4), and concentrated *in vacuo* to afford 1-(*tert*-butoxycarbonyl(methyl)amino) cyclobutanecarboxylic acid.

Intermediate 30


2-bromobutanamide



[0124] Under nitrogen, to a solution of 2-bromobutanoic acid (2.0 g, 12.0 mmol) in DCM (50 mL) at 0°C were sequentially added thionyl chloride (2.6 mL, 35.9 mmol) and 0.5 mL of DMF. The reaction was allowed to warm to room temperature and was stirred overnight. The reaction mixture was concentrated *in vacuo* and then cooled to 0°C. Thirty (30) mL of 37% aqueous ammonia hydroxide was added slowly. The reaction was allowed to warm to room temperature and was stirred for 2 h followed by extraction with EtOAc. The combined organic extracts were dried (Na_2SO_4), and concentrated *in vacuo* to afford 2-bromobutanamide.

Intermediate 31


2-bromo-N-methylbutanamide



[0125] Under nitrogen, to a solution of 2-bromobutanoic acid (2.0 g, 12.0 mmol) in DCM (50 mL) at 0°C were sequentially added thionyl chloride (2.6 mL, 35.9 mmol) and 0.5 mL of DMF. The reaction was allowed to warm to room temperature and was stirred overnight. The reaction mixture was concentrated *in vacuo* and then cooled to 0°C. Thirty (30) mL of 40% methylamine water solution was added slowly. The reaction was allowed to warm to room temperature and was stirred for 2 h followed by extraction with EtOAc. The combined organic extracts were dried (Na_2SO_4), and concentrated *in vacuo* to afford 2-bromo-N-methylbutanamide.

Intermediate 32


2-bromo-N-ethylbutanamide



[0126] Under nitrogen, to a solution of 2-bromobutanoic acid (2.0 g, 12.0 mmol) in DCM (50 mL) at 0°C were sequentially added thionyl chloride (2.6 mL, 35.9 mmol) and 0.5 mL of DMF. The reaction was allowed to warm to room temperature and was stirred overnight. The reaction mixture was concentrated *in vacuo* and then cooled to 0°C. Thirty (30) mL of 40% ethylamine water solution was added slowly. The reaction was allowed to warm to room temperature and was stirred for 2 h, followed by extraction with EtOAc. The combined organic extracts were dried (Na₂SO₄), and concentrated *in vacuo* to afford 2-bromo-*N*-ethylbutanamide.

Intermediate 33


1-(Methylsulfonyl)azetidin-3-one



[0127] To a mixture of 3-azetidinone hydrochloride (471 mg, 3.28 mmol) and K₂CO₃ (1.50 g, 11 mmol) in CHCl₃/H₂O (5/5 mL) was added methanesulfonic anhydride (1.14 g, 6.57 mmol) in one portion at 0 °C. After stirring for 2 h, the reaction was diluted with saturated aqueous NaHCO₃ (5 mL) and extracted with DCM (3x25 mL). The combined DCM layers were washed with brine (5 mL), dried over Na₂SO₄ and evaporated to afford 1-(Methylsulfonyl)azetidin-3-one as an off white solid, which was used without further purification; ¹H NMR (400 MHz, CDCl₃) δ 4.82 (s, 4H), 3.05 (s, 3H).

Example 1

5-Chloro-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (1)




[0128] Step 1: A mixture of 2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine (Intermediate 1, 132 mg, 0.54 mmol) and *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)piperidine-1-carboxylate (Intermediate 19, 166 mg, 0.54 mmol) in 2-propanol (15 mL) was treated with conc. aqueous HCl (14 drops). The mixture was sealed and heated in a microwave at 130 °C for 60 min. The mixture was concentrated to afford 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a yellow solid. A portion of the crude product was purified with preparative RP-HPLC to afford 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ESMS m/z 416.1 ($M + H^+$). The remainder of the crude product was used directly for the next step without further purification.

[0129] Step 2: To a solution of the crude product from previous step in THF (5 mL) and methanol (5 mL) was added formaldehyde (100 μ L, 1.3 mmol) and 10 drops of AcOH sequentially. The reaction mixture was stirred at room temperature for 1 h, then sodium cyanoborohydride (175 mg, 2.78 mmol) was added in one portion, and the reaction was stirred for an additional 30 min. The reaction was quenched by saturated aqueous NH_4Cl and concentrated *in vacuo* to give an oily residue. The residue was purified with preparative RP-HPLC to afford 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 10.33 (br, 1H), 9.90 (br, 1H), 9.60 (br, 1H), 8.22 (s, 1H), 7.56 (d, 1H), 7.00 (d, 1H), 6.20 (s, 1H), 3.53-3.50 (m, 2H), 3.16-3.10 (m, 2H), 2.97-2.92 (m, 1H), 2.81 (d, 3H), 2.26 (s, 3H), 2.16 (s, 3H), 1.94-1.91 (m, 2H), 1.83-1.73 (m, 2H); ESMS m/z 430.1 ($M + H^+$).

Example 2


5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (2)




[0130] A mixture of 2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine (Intermediate 1, 132 mg, 0.54 mmol) and *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)piperidine-1-carboxylate (Intermediate 19, 166 mg, 0.54 mmol) in 2-propanol (15 mL) was treated with conc. aqueous HCl (14 drops). The mixture was sealed and heated in a microwave at 130 °C for 60 min. The mixture was concentrated to afford 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as yellow solid. A portion of the crude product was purified with preparative RP-HPLC to afford 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ESMS m/z 416.1 ($M + H^+$).

Examples 3 and 4

(*S*)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one (10)




(*R*)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one (11)




[0131] ($\pm$)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one was synthesized by the same series of procedures as described in Example 1, utilizing Intermediate 6 and Intermediate 19. Chiral separation of the racemic mixture was conducted with normal phase HPLC using ChiralPaK AD column using the following solvent system: hexanes (80%) and iPrOH (20%) modified with 0.1% Diethylamine. The two purified enantiomer peaks were collected separately: (*S*)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one and (*R*)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one; both peaks: ESMS m/z 461.2 ($M + H^+$). The earlier eluting peak ($RT = 7.26$ min.) and the later eluting peak ($RT = 10.17$ min.) were arbitrarily assigned as the (*S*) and (*R*) enantiomer, respectively.

Example 5


5-Chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (19)




[0132] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (83 μ L, 0.6 mmol) in DMF (1.5 mL), was added 4-(chloromethyl)-3,5-dimethylisoxazole (35 mg, 0.24 mmol). The mixture was stirred at room temperature for 3 hours. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to afford 5-Chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid. ^1H NMR (400 MHz, MeOD- d_4) 8.12 (s, 1H), 7.75 (d, 1H), 7.03 (s, 1H), 6.27 (s, 1H), 4.25 (s, 2H), 3.69-3.67 (m, 2H), 3.27-3.20 (m, 3H), 2.54 (s, 3H), 2.37 (s, 3H), 2.34 (s, 3H), 2.30 (s, 3H), 2.10-1.95 (m, 4H); ESMS m/z 525.1 ($M + H^+$).

Example 62-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanol (22)


[0133] 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) was dissolved in anhydrous DMF (1 mL). TEA (50 uL, 0.36 mmol, 3 equiv.) was added followed by 2-bromoethanol (0.018 mL, 0.24 mmol, 2 equiv.) dissolved in anhydrous DMF (0.7 mL). The reaction vessel was sealed and heated in a microwave at 100 °C for 20 min. After cooling to room temperature, the reaction was concentrated and the crude product was purified using preparative RP-HPLC to give 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanol as a white solid; ESMS *m/z* 460.2 (M + H⁺).

Example 75-chloro-N²-(2-fluoro-5-methyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (24)


[0134] The mixture of 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol), 3-bromo-1,1,1-trifluoropropane (85.0 uL, 0.60 mmol) and triethylamine (102.0 uL, 0.60 mmol) in 2 mL of DMF was stirred overnight. The reaction mixture was purified by preparative RP-HPLC to afford 5-chloro-N²-(2-fluoro-5-methyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine. ESMS *m/z* 512.2 (M + H⁺).

Example 83-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol (30)


[0135] The mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (50.0 mg, 0.12 mmol) and 2-(trifluoromethyl)oxirane (67.8 mg, 0.60 mmol) in 2 mL of DMF was stirred overnight. The reaction mixture was purified by preparative RP-HPLC to afford 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol. ESMS m/z 528.2 ($M + H^+$).

Example 92-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide (39)


[0136] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (50 μ L, 0.36 mmol) in DMF (1.5 mL), was added 2-bromo-acetamide (33 mg, 0.24 mmol). The mixture was stirred at room temperature for 2 hours. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to give 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl) piperidin-1-yl)acetamide as a white solid: ^1H NMR (400 MHz, MeOD- d_4) δ 8.11 (s, 1H), 7.76 (d 1H), 7.09 (d, 1H), 6.27 (s, 1H), 4.0 (s, 2H), 3.75-3.73 (m, 2H), 3.22-3.13 (m, 3H), 2.34 (s, 3H), 2.30 (s, 3H), 2.04 -2.00 (m, 4H); ESMS m/z 473.2 ($M + H^+$).

Example 105-Chloro-N2-(2-fluoro-5-methyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (55)


[0137] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (83 μ L, 0.6 mmol) in DMF (1.5 mL), was added methyl vinyl sulfone (38 mg, 0.36 mmol). The mixture was stirred at room temperature for 1 hour. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to afford 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid. ESMS m/z 522.1 ($M + H^+$).

Example 115-Chloro-N2-(2-fluoro-5-methyl-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (62)

[0138] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (50 μ L, 0.36 mmol) in DMF (1.5 mL) was added methanesulfonyl chloride (18 μ L, 0.24 mmol). The mixture was stirred at room temperature for 2 hours. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to give 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid: ESMS m/z 494.2 ($M + H^+$).

Example 12Ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate (66)

[0139] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (50 μ L, 0.36 mmol) in DMF (1.5 mL) was added ethyl chloroformate (26 mg, 0.24 mmol). The mixture was stirred at room temperature for 2 hours. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to give Ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate as a white solid: ^1H NMR (400 MHz, $\text{MeOD-}d_4$) δ 8.13 (s, 1H), 7.54 (d, 1H), 7.10 (d, 1H), 6.28 (s, 1H), 4.30-4.26 (m, 2H), 4.14 (q, 2H), 2.99-2.96 (m, 3H), 2.33 (s, 3H), 2.29 (s, 3H), 1.8-1.77 (m, 2H), 1.64-1.55 (m, 2H), 1.28 (t, 3H); ESMS m/z 488.2 ($\text{M} + \text{H}^+$).


Example 134-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-N,N-dimethylpiperidine-1-carboxamide (69)

[0140] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (83 μ L, 0.6 mmol) in DMF (1.5 mL) was added dimethylcarbamic chloride (39 mg, 0.36 mmol). The mixture was stirred at room temperature for 1 hour. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to afford 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-N,N-dimethylpiperidine-1-carboxamide as a white solid.

ESMS m/z 487.2 ($M + H^+$).

Example 14


1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone (73)



[0141] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (50 μ L, 0.36 mmol) in DMF (1.5 mL) was added 2-(dimethylamino)acetyl chloride (38 mg, 0.24 mmol). The mixture was stirred at room temperature for 1 hour. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to give 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone as a white solid: ^1H NMR (400 MHz, $\text{MeOD-}d_4$) δ 8.08 (s, 1H), 7.71 (d, 1H), 7.03 (d, 1H), 6.26 (s, 1H), 4.71-4.67 (m, 2H), 3.81-3.77 (m, 2H), 3.14-3.12 (m, 2H), 2.98 (s, 3H), 2.96 (s, 3H), 2.90-2.83 (m, 1H), 2.34 (s, 3H), 2.28 (s, 3H), 1.87 (d, 2H), 1.76-1.55 (m, 2H); ESMS m/z 501.2 ($M + H^+$).

Example 15

1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(4-methylpiperazin-1-yl)ethanone (105)




[0142] To the solution of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (100.0 mg, 0.24 mmol) in 4 mL of DCM was added 2-chloroacetyl chloride (23.0 μ L, 0.29 mmol) and triethylamine (67.0 μ L, 0.48 mmol) sequentially. The reaction was stirred at room

temperature for 1 h then washed by brine. The organic extract was dried over Na_2SO_4 , followed by concentration in vacuo to afford a crude product. The crude product was mixed with 1-methylpiperazine (116.0 mg, 1.16 mmol) in 3 mL of DMF and the reaction was stirred overnight at room temperature. The reaction mixture was purified by preparative HPLC to afford 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(4-methylpiperazin-1-yl)ethanone. ESMS m/z 556.3 ($M + H^+$).

Example 16


Azetidin-3-yl(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone (131)



[0143] To a mixture of 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol), diisopropylethylamine (50 μL , 0.36 mmol) and HATU (55 mg, 0.14 mmol) in DMF (1.5 mL) was added 1-(tert-butoxycarbonyl)azetidine-3-carboxylic acid (29 mg, 0.14 mmol). After stirring at room temperature for 4 hours, the mixture was diluted with methanol (1 mL) and conc. aqueous HCl (1 mL). The mixture was stirred at 50 °C for 30 min. The mixture was filtered and purified by preparative RP-HPLC to give azetidin-3-yl(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone as white solid. ^1H NMR (400 MHz, $\text{MeOD}-d_4$) δ 8.08 (s, 1H), 7.68 (d, 2H), 7.04 (d, 2H), 6.26 (s, 1H), 4.74-4.68 (m, 2H), 4.39-4.35 (m, 1H), 4.29-4.24 (m, 2H), 4.11-4.07 (m, 1H), 3.74-3.71 (m, 1H), 3.25-3.24 (m, 1H), 3.07-3.04 (m, 1H), 2.87-2.80 (m, 1H), 2.33 (s, 3H), 2.28 (s, 3H), 1.87-1.84 (m, 2H), 1.66-1.56 (m, 2H); ESMS m/z 499.2 ($M + H^+$).

Example 17


5-chloro-N²-(2-fluoro-5-methyl-4-(1-(tetrahydro-1,1-dioxido-3-thienyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (145)



[0144] To 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (30 mg, 0.072 mmol) in EtOH (0.5 mL) was added thiophene-2,3-dihydro-1,1-dioxide (17 mg, 0.144 mmol). The resulting mixture was then heated to 130 °C for 2 hr.. After cooling down to room temperature, the mixture was concentrated and purified by preparative RP-HPLC to provide 5-chloro-N²-(2-fluoro-5-methyl-4-(1-(tetrahydro-1,1-dioxido-3-thienyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine. ESMS m/z 534.2 (M + H⁺).

Example 18

5-chloro-N²-(2,5-dimethyl-4-(1-(tetrahydro-1,1-dioxido-2H-thiopyran-4-yl))piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
(146)




[0145] Step 1: To a solution of 5-chloro-N²-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (30 mg, .073

mmol) in acetonitrile (1 mL) was added Cs_2CO_3 (47 mg, 0.15 mmol) and 4-iodotetrahydro-2H-thiopyran (48 mg, 0.22 mmol). This mixture was then stirred at 80 °C for 16 hr. After cooling down to room temperature, the mixture was treated with saturated aqueous NH_4Cl solution (3 mL) and extracted with EtOAc (3x3 mL). The organic layers were combined and concentrated. The residue was purified by silica gel chromatography (Gradient 0~8% $\text{MeOH}/\text{CH}_2\text{Cl}_2$ with NH_3) to provide 5-chloro-N²-(2,5-dimethyl-4-(1-(tetrahydro-2H-thiopyran-4-yl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid. ESMS m/z 512.2 ($\text{M} + \text{H}^+$).

[0146] Step 2: To 5-chloro-N²-(2,5-dimethyl-4-(1-(tetrahydro-2H-thiopyran-4-yl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (18 mg, .035 mmol) in CH_2Cl_2 (1 mL) was added *m*-CPBA (16 mg, .071 mmol) at 0 °C. The mixture was then warmed up to room temperature and stirred for 30 min; saturated aqueous NaHCO_3 solution (3 mL) was then added and the crude product extracted with CH_2Cl_2 (3 x3 mL). The combined organic layers were concentrated and purified by preparative thin layer chromatography (silica gel, 12 % $\text{MeOH}/\text{CH}_2\text{Cl}_2$ with NH_3) to provide 5-chloro-N²-(2,5-dimethyl-4-(1-(1,1-dioxido-2H-thiopyran-4-yl))piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine. ESMS m/z 544.2 ($\text{M} + \text{H}^+$).

Example 19

5-chloro-N²-(2-fluoro-5-methyl-4-(1-(1,1-dioxido-3-thietanyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (148)




[0147] To 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (30 mg, 0.072 mmol) in MeOH (1

mL) was added 3-bromothietane 1,1-dioxide (15mg, 0.079 mmol) followed by TEA (15mg). The resulting mixture was stirred at room temperature for 5 hr and then concentrated. The resulting residue was purified by preparative RP-HPLC to provide 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1,1-dioxido-3-thietanyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine. ¹H NMR (400 MHz, DMSO-d₆ + 1 drop D₂O) δ 8.02 (s, 1H), 7.37 (s, 1H), 7.02 (s, 1H), 6.23 (s, 1H), 4.28-4.22 (m, 2H), 4.12-4.07 (m, 2H), 3.22-3.18 (m, 1H), 2.95-2.92 (m, 2H), 2.68-2.62 (m, 1H), 2.22 (s, 3H), 2.13 (s, 3H), 2.07-2.01 (m, 2H), 1.71-1.55 (m, 4H). ESMS m/z 520.2 (M + H⁺).

Example 20


5-Chloro-N2-(4-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (149)



[0148] To a mixture of 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.10 mmol) and triethylamine (83 μ L, 0.6 mmol) in DMF (1.5 mL), was added 2-chloro-5-ethylpyrimidine (27 mg, 0.20 mmol). The mixture was heated in a microwave at 120 °C for 10 min. The reaction was filtered and the filtrate was purified by preparative RP-HPLC to afford 5-chloro-N2-(4-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid. ESMS m/z 522.2 (M + H⁺).

Example 21

4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-2-one (151)



[0149] Step 1: In a 5 mL microwave reaction tube was added 2-(2,5-dimethyl-4-nitrophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (143 mg, 0.516 mmol, prepared from 1-bromo-2,5-dimethyl-4-nitrobenzene by standard protocol), 1-(4-methoxybenzyl)-5,6-dihydropyridin-2(1H)-one (56 mg, 0.26 mmol, prepared by following a similar procedure as reported by Lerchner etc. in *Chem. Eur. J.* 2006, 12, 8208), chloro(1,5-cyclooctadiene)rhodium (I) dimer (13 mg, .026 mmol), KOH (0.13 mL 1N aqueous solution) and dioxane (1.2 mL). The tube was degassed, filled with N₂ and sealed. The reaction tube was then heated to 100 °C in a microwave reactor for 10 min. After the reaction tube was opened, the mixture was treated with saturated NH₄Cl aqueous solution (3 mL) and extracted with EtOAc (3 x 4 mL). The organic layers were combined and concentrated. The residue was purified by flash column chromatography (silica gel, gradient 0%~70% EtOAc/hexane), to provide 4-(2,5-dimethyl-4-nitrophenyl)-1-(4-methoxybenzyl)piperidin-2-one. ESMS m/z 369.2 (M + H⁺).

[0150] Step 2: To 4-(2,5-dimethyl-4-nitrophenyl)-1-(4-methoxybenzyl)piperidin-2-one (63 mg, 0.17 mmol) in MeOH (10 mL) was added 10% w/w Pd/C (6 mg), the mixture was degassed and stirred under H₂ at room temperature for 14 hr. After removing the catalyst by filtration, the filtrate was concentrated to provide 4-(4-amino-2,5-dimethylphenyl)-1-(4-methoxybenzyl)piperidin-2-one as a pale yellowish oil. ESMS m/z 339.2 (M + H⁺).


[0151] Step 3: To a mixture of 4-(4-amino-2,5-dimethylphenyl)-1-(4-methoxybenzyl)piperidin-2-one (23 mg, 0.094 mmol) and 2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine (29 mg, 0.086 mmol) in iPrOH (1 mL) was added HCl (60 uL, 4N in dioxane). The reaction vessel was then sealed and heated to 130 °C for 4 hr. After cooling to room temperature, the mixture was treated with saturated NaHCO₃ aqueous solution (3 mL) and extracted with EtOAc (3 x 4mL). The organic layers were combined and concentrated. The residue was purified by flash column

chromatography (silica gel, gradient 0~10% MeOH/CH₂Cl₂) to provide 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)-1-(4-methoxybenzyl)piperidin-2-one. ESMS m/z 546.2 (M + H⁺).

[0152] Step 4: A solution of 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)-1-(4-methoxybenzyl)piperidin-2-one (45 mg) in TFA (0.5 mL) was heated to 100 °C for 24 hr. After cooling down to room temperature, the mixture was concentrated and purified by preparative RP-HPLC to provide 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-2-one. ESMS m/z 426.2 (M + H⁺).

Example 22

5-chloro-N²-(2-fluoro-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (157)




[0153] Step 1: To a microwave reaction tube was added 4-bromo-2-fluoro-5-(trifluoromethyl)aniline (256 mg, 1.0 mmol), 1-methylpiperazine (300mg, 3 mmol) Pd₂(dba)₃ (91.5 mg, 0.1 mmol), di-tert-butyl(2',4',6'-triisopropylbiphenyl-2-yl)phosphine (60 mg, 0.2 mmol), NaOtBu (144mg, 1.5 mmol) and THF (3 mL). After the reaction tube was degassed and filled with N₂, the tube was heated to 120 °C in a microwave reactor for 40 min. The mixture was then poured into saturated NH₄Cl aqueous solution (10 mL) and extracted with EtOAc (3 x10 mL). The organic layers were combined and concentrated. The residue was purified by flash column chromatography (silica gel, gradient MeOH/CH₂Cl₂, 0~10%) to provide 2-fluoro-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)aniline as a white solid. ESMS m/z 278.1 (M + H⁺).

[0154] Step 2: To a mixture of 2-fluoro-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)aniline (37 mg, 0.133 mmol) and 2,5-dichloro-N-(5-methyl-1H-

pyrazol-3-yl)pyrimidin-4-amine (36 mg, 0.147 mmol) in iPrOH (1 mL) was added HCl (100 μ L, 4N in dioxane). The reaction vessel was then sealed and heated to 130 °C for 4 hr. After cooling to room temperature, the mixture was treated with saturated NaHCO₃ aqueous solution (5 mL) and extracted with EtOAc (3 x 5mL). The organic layers were combined and concentrated. The residue was purified by preparative thin layer chromatography (silica gel, 8% MeOH/CH₂Cl₂), to provide 5-chloro-N²-(2-fluoro-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine. ESMS m/z 485.2 (M + H⁺).

Example 23


(S)-2-(4-(4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide (163)



[0155] A mixture of 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (50.0 mg, 0.12 mmol), (*R*)-2-bromopropanamide (90.6 mg, 0.60 mmol), and triethylamine (102.0 μ L, 0.60 mmol) in 2 mL of DMF was heated at 150°C in a microwave reactor for 30 min. The reaction mixture was directly purified by preparative RP-HPLC to afford (*S*)-2-(4-(4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide; ESMS m/z 487.2 (M + H⁺).

Example 24


(S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylpropanamide (164)



[0156] A mixture of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (50.0 mg, 0.12 mmol), (*R*)-2-bromo-*N*-methylpropanamide (100.0 mg, 0.60 mmol), and triethylamine (102.0 μ L, 0.60 mmol) in 2 mL of DMF was heated at 150°C in a microwave reactor for 30 min. The reaction mixture was directly purified by preparative RP-HPLC to afford (*S*)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-*N*-methylpropanamide; ESMS m/z 501.2 ($M + H^+$).

Example 25


(*S*)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide (166)



[0157] A mixture of 5-chloro- N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (50.0 mg, 0.12 mmol), (*R*)-2-bromopropanamide (90.6 mg, 0.60 mmol), and triethylamine (102.0 μ L, 0.60 mmol) in 2 mL of DMF was heated at 150°C in a microwave reactor for 30 min. The reaction mixture was directly purified by preparative RP-HPLC to afford (*S*)-2-(4-(4-(5-chloro-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide; ESMS m/z 483.2 ($M + H^+$).

Example 26

(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclopropyl)methanone (169)




[0158] A mixture of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-

N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (67.3 mg, 0.17 mmol), 1-(*tert*-butoxycarbonyl(ethyl)amino)cyclopropanecarboxylic acid (38.0 mg, 0.17 mmol), HATU (63.0 mg, 0.17 mmol) and diisopropylethylamine (57 μ L, 0.34 mmol) in 1 mL of DMF was stirred at room temperature overnight. The reaction mixture was partitioned between EtOAc and water, the combined organic extracts were dried (Na_2SO_4) and then concentrated *in vacuo*. The resulting crude mixture was dissolved in 5 mL of DCM and 4 mL of TFA and was then stirred for 2 h followed by concentration *in vacuo*. The crude product was purified by RP-HPLC to afford (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclopropyl)methanone; ESMS m/z 527.2 ($M + H^+$).

Example 27


1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(ethylamino)-2-methylpropan-1-one (175)



[0159] To a solution of 2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methylpropan-1-one (6mg, 0.012 mmol) in acetone (0.5 mL) was added K_2CO_3 (11 mg, 0.079 mmol). The mixture was stirred at room temperature for 10 min. before the addition of EtI (6 mg, 0.038 mmol). The resulting mixture was then stirred at room temperature for 14 h before it was treated with saturated aqueous NH_4Cl (1 mL) and extracted with EtOAc (3 x 2mL). The organic layers were combined, concentrated, and the residue was purified by preparative thin layer chromatography (silica gel, 8% MeOH/DCM/ NH_3) followed by further purification with preparative RP-HPLC to provide 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(ethylamino)-2-methylpropan-1-one; ESMS m/z 525.3 ($M + H^+$).

Example 28


(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclobutyl)methanone (176)



[0160] A mixture of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (50.0 mg, 0.12 mmol), 1-(*tert*-butoxycarbonyl(ethyl)amino)cyclobutanecarboxylic acid (29.2 mg, 0.12 mmol), HATU (45.8 mg, 0.12 mmol) and di-isopropylethylamine (20 μ L, 0.12 mmol) in 1 mL of DMF was stirred at room temperature overnight.. The reaction mixture was partitioned between EtOAc and water. The combined organic extracts were dried (Na_2SO_4), and concentrated *in vacuo*. The resulting crude mixture was dissolved in 5 mL of DCM and 4 mL of TFA and was then stirred for 2 h followed by concentration *in vacuo*. The crude product was purified by RP-HPLC to afford (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclobutyl)methanone; ESMS m/z 541.3 ($M + H^+$).

Example 29

(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(methylamino)cyclobutyl)methanone (177)




[0161] A mixture of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (50.0 mg, 0.12 mmol), 1-(*tert*-butoxycarbonyl(methyl)amino)cyclobutanecarboxylic acid (27.5 0.12 mmol), HATU

(45.8 mg, 0.12 mmol) and di-isopropylethylamine (20 μ L, 0.12 mmol) in 1 mL of DMF was stirred at room temperature overnight. The reaction mixture was partitioned between EtOAc and water. The combined organic extracts were dried (Na_2SO_4), and concentrated *in vacuo*. The resulting crude mixture was dissolved in 5 mL of DCM and 4 mL of TFA, and was stirred for 2 h followed by concentration *in vacuo*. The crude product was purified by RP-HPLC to afford (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(methylamino)cyclobutyl)methanone; ESMS m/z 527.2 ($\text{M} + \text{H}^+$).

Example 30


(S)-3-(4-(2,5-dimethyl-4-(5-methyl-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol (179)



[0162] To a solution of N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-5-methyl- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine hydrochloride (36.0 mg, 0.084 mmol) in DMF (0.5 mL) were added TEA (23.4 μ L, 0.168 mmol) and (S)-2-(trifluoromethyl) oxirane (72.8 μ L, 0.84 mmol). The reaction mixture was stirred at room temperature overnight, and purified by preparative RP-HPLC to provide (S)-3-(4-(2,5-dimethyl-4-(5-methyl-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol; ESMS m/z 504.3 ($\text{M} + \text{H}^+$).

Example 31


2-(2-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide (181)




[0163] To a mixture of 5-Chloro-N2-(2,5-dimethyl-4-(piperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (0.12 mmol) and triethylamine (83 μ L, 0.6 mmol) in DMF (1.5 mL), was added 2-bromoacetamide (35 mg, 0.24 mmol). The mixture was stirred at room temperature for 3 h. The reaction was filtered and the filtrate was purified by RP-HPLC to afford 2-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide as a white solid; ^1H NMR (400 MHz, MeOD- d_4) 8.09 (s, 1H), 7.65 (d, 1H), 7.43 (s, 1H), 6.41 (s, 1H), 4.79-4.74 (m, 2H), 3.83-3.48 (m, 2H), 2.71 (s, 3H), 2.42 (s, 3H), 2.34 (s, 6H), 2.15-1.84 (m, 7H); ESMS m/z 469.2 ($M + H^+$).

Examples 32 and 33

(S) 5-Chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (183)



(R) 5-Chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (184)



[0164] Step 1: To a mixture of 1-bromo-2,5-dimethyl-4-nitrobenzene (185 mg, 1 mmol) and 2-(tributylstannyl)pyridine (202 mg, 1.1 mmol) in DMF (4 mL) was added tetrakis(triphenylphosphine) palladium (0) (58 mg, 0.05 mmol). The reaction tube was sealed, the mixture was purged with N_2 for 3 min and then heated at 120 $^\circ\text{C}$ under

N₂ for overnight. The reaction was cooled to room temperature and poured into saturated aqueous ammonia chloride solution. The crude reaction mixture was extracted with ethyl acetate (3 x 15 mL). The organic extracts were combined, washed with brine and concentrated. The crude product was purified with silica chromatography (60% ethyl acetate in hexanes) to afford 2-(2,5-dimethyl-4-nitrophenyl)pyridine as a white solid. The obtained solid was dissolved in acetic acid/TFA (15 mL/200 μ L). To this solution was added PtO₂ (10% w/w). The reaction mixture was degassed and purged with H₂ for several times and then vigorously stirred under 1 atm. H₂ overnight. The mixture was filtered and the filtrate concentrated to afford 2,5-Dimethyl-4-(piperidin-2-yl)aniline as a yellow oil; ESMS m/z 205 (M + H⁺).

[0165] Step 2: A mixture of 2,5-dichloro-N-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine (120 mg, 0.49 mmol) and 2,5-dimethyl-4-(piperidin-2-yl)aniline (100 mg, 0.49 mmol) in 2-propanol (10 mL) was treated with conc. aqueous HCl (7 drops). The mixture was sealed and heated in a microwave at 130 °C for 45 min. The mixture was concentrated to afford 5-Chloro-N2-(2,5-dimethyl-4-(piperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine; ESMS m/z 412.1 (M + H⁺). The crude product was used directly for the next step without further purification.


[0166] Step 3: To a solution of the crude product from the previous step in THF (1 mL) and methanol (1 mL) was sequentially added formaldehyde (100 μ L, 1.3 mmol) and 5 drops of AcOH. The reaction mixture was stirred at room temperature for 1 h, then sodium cyanoborohydride (160 mg, 2.45 mmol) was added in one portion, and the reaction was stirred for an additional 30 min. The reaction was quenched by saturated aqueous NH₄Cl and concentrated *in vacuo* to give an oily residue. The residue was purified with RP-HPLC to afford ($\pm$)-5-Chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ESMS m/z 426.2 (M + H⁺).

[0167] Step 4: Chiral separation of the racemic mixture was conducted with normal phase HPLC using ChiralPaK AD-H column using the following solvent system: hexanes (95%), EtOH (2.5%), MeOH (2.5%). The two purified enantiomer peaks were collected separately: (*R*)-5-Chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

and (*S*)-5-Chloro-N²-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine; both peaks: ESMS *m/z* 426.2 (*M* + *H*⁺). The earlier eluting peak has been arbitrarily assigned as the (*R*) enantiomer.

Example 34

5-Chloro-N²-(2,5-dimethyl-4-(1-(3-morpholinopropylsulfonyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (194)




[0168] Step 1: To a solution of 5-chloro-N²-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (78 mg, 0.19 mmol) and TEA (0.52 mL, 3.78 mmol) in DCM (10 mL) was added 3-chloropropylsulfonyl chloride (63 mg, 0.36 mmol, in 1 mL DCM). After stirring at rt for one hour, EtOAc (100 mL) was added. The mixture was sequentially washed with water (10 mL), brine (10 mL), dried over Na₂SO₄ and concentrated *in vacuo*. The crude product was purified by silica chromatography (0-100% EtOAc in hexanes gradient) to give 5-chloro-N²-(4-(1-(3-chloropropylsulfonyl)piperidin-4-yl)-2,5-dimethylphenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as an off white solid; ESMS *m/z* 552.2 (*M*+*H*⁺).

[0169] Step 2: The product from Step 1 was stirred in neat morpholine (0.5 mL) at 100 °C in a sealed vial for one hour. The reaction was purified by RP-HPLC to give 5-Chloro-N²-(2,5-dimethyl-4-(1-(3-morpholinopropylsulfonyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white powder; ESMS *m/z* 603.2 (*M*+*H*⁺).

Example 35


1-((4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methyl)cyclopropanecarbonitrile (198)



[0170] To a solution of 5-chloro- N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (30 mg, 0.078 mmol) in acetonitrile (1.0 mL) was added (1-cyanocyclopropyl)methyl 4-methylbenzenesulfonate (16 mg, 0.10 mmol, prepared according to the procedure described in patent WO2005063247) followed by DIEA (30mg, 0.23 mmol) and KI (catalytic amount). The resulting mixture was heated to 70 °C for 14 h and then cooled down to room temperature. The resulting mixture was directly purified by preparative RP-HPLC to provide 1-((4-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5dimethylphenyl)-piperidin-1-yl)methyl)cyclopropanecarbonitrile; ESMS m/z 491.2 ($M + H^+$).

Example 36

2-(4-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetonitrile (199)




[0171] To a solution of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (20 mg, 0.048 mmol) in acetonitrile (0.5 mL) was added Cs_2CO_3 (31 mg, 0.096 mmol) and chloroacetonitrile (7 mg, 0.096 mmol). The resulting mixture was stirred at room temperature for 14 h before it was treated with saturated aqueous NH_4Cl (1 mL) and extracted with EtOAc (3 x 2mL). The organic layers were combined, concentrated, and the residue was purified by preparative RP-HPLC to provide 2-(4-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-

yl)acetonitrile; ESMS m/z 455.2 ($M + H^+$).

Example 37


3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanenitrile (200)




[0172] To a solution of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (20 mg, 0.048 mmol) in MeOH (0.5 mL) was added acrylonitrile (5 mg, 0.096 mmol). The resulting mixture was stirred at room temperature for 14 h and then purified directly by preparative RP-HPLC to provide 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanenitrile; ESMS m/z 469.2 ($M + H^+$).

Examples 38 and 39

(R)-2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide (207)




(S)-2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide (208)



[0173] A solution of 5-chloro-*N*²-(2,5-dimethyl-4-(piperidin-3-yl)phenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (700 mg, 1.70 mmol), 2-bromo-*N*-methylacetamide (258mg, 1.70 mmol) and TEA (1.2 mL, 8.5 mmol) in DMF (4 mL) was stirred at rt for 30 min. The reaction was purified by RP-HPLC to give the product as a racemate; ESMS *m/z* 483.2 (*M*+1). Chiral separation of the racemic mixture was conducted with chiral HPLC (ChiralCel OD-H, Hex:EtOH:MeOH/80:10:10, 15 min run time, 1 mL/min). The two purified enantiomer peaks were collected separately, both as white solids: (*R*)-2-(3-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-*N*-methylacetamide and (*S*)-2-(3-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-*N*-methylacetamide; both peaks: ESMS *m/z* 483.2 (*M*+*H*⁺). The earlier eluting peak (*rt* = 5.63 min. vs. *rt* = 7.62 min.) was arbitrarily assigned as the (*R*) enantiomer.

Example 40

5-chloro-*N*²-(2-fluoro-5-methyl-4-(1-(tetrahydro-2*H*-pyran-4-yl)piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (209)




[0174] To a suspension of 5-chloro-*N*²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (2.50 g, 6.0 mmol) in a mixed solvent of MeOH (42 mL) and DCM (10 mL) was added dihydro-2*H*-pyran-4(3*H*)-one (3.33 mL, 36.0 mmol) and TEA (8.36 mL, 60.0 mmol). The reaction mixture was heated at 60 °C for 1 h before NaCNBH₃ (1.66 g, 26.4 mmol) was added. The mixture was heated at 60 °C for another 2 h, cooled down to room temperature, diluted with DCM (60 mL), dry loaded onto silica gel (12.5 g), and purified by silica gel chromatography (DCM with 1% NH₃ to 10% MeOH in DCM with 1% NH₃ gradient) to provide 5-chloro-*N*²-(2-fluoro-5-methyl-4-(1-(tetrahydro-2*H*-pyran-4-yl)piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ¹H NMR (DMSO-*d*₆, 400 MHz) δ 12.08 (br, 1H), 9.58 (br, 1H), 8.65 (d, *J*

= 28.4 Hz, 1H), 8.02 (s, 1H), 7.36 (d, $J = 8.4$ Hz, 1H), 7.03 (d, $J = 12.4$ Hz, 1H), 6.08 (br, 1H), 3.89 (dd, $J = 10.8, 3.6$ Hz, 2H), 3.28 (t, $J = 10.4$ Hz, 2H), 2.99 (d, $J = 10.8$ Hz, 2H), 2.64-2.43 (m, 2H), 2.24-2.15 (m, 8H), 1.71-1.40 (m, 8H); ESMS m/z 500.2 ($M + H^+$).

Example 41


5-chloro- N^2 -(4-(1-cyclopropylpiperidin-4-yl)-2-fluoro-5-methylphenyl)- N^4 -(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (210)



[0175] To a solution of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (30 mg, 0.072 mmol) in MeOH (0.7 mL) was added AcOH (35 mg, 0.58 mmol), (1-ethoxycyclopropoxy)trimethylsilane (37 mg, 0.216 mmol) and a small amount of 4Å molecular sieves. The mixture was stirred at room temperature for 1 h before the addition of Na(CN)BH₃ (14 mg, 0.216 mmol). The resulting mixture was then stirred at 60 °C for 14 h and cooled to room temperature. Saturated aqueous NH₄Cl (2 mL) was added and the mixture was extracted with EtOAc (3 x 3 mL). The combined organic layers were concentrated and purified by preparative RP-HPLC to provide 5-chloro- N^2 -(4-(1-cyclopropylpiperidin-4-yl)-2-fluoro-5-methylphenyl)- N^4 -(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine; ¹H NMR (400 MHz, MeOD-*d*₄) δ 7.97 (s, 1H), 7.84 (d, 1H), 7.50 (s, 1H), 6.93 (d, 1H), 6.29 (s, 1H), 5.33 (s, 1H), 2.80 (m, 1H), 2.58 (s, 2H), 2.28 (s, 3H), 2.24 (s, 3H), 1.96 (m, 1H), 1.80-1.85(m, 2H), 1.65-1.75 (m, 2H), 0.2-0.7 (m, 6H); ESMS m/z 456.2 ($M + H^+$).

Example 42

N^2 -(4-(1-(3-(azetidin-1-ylsulfonyl)propyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro- N^4 -(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (214)




[0176] Step 1: To a solution of azetidine (12 mg, 0.21 mmol) and TEA (0.1 mL, 0.71 mmol) in DCM (5 mL) was slowly added a solution of 3-chloropropylsulfonyl chloride (34 mg, 0.19 mmol) in DCM (1 mL). After stirring at rt overnight, the solvent was removed *in vacuo*. The resulting 1-(3-chloropropylsulfonyl)azetidine crude product was dissolved in NMP (1.9 mL, 0.1 mmol/mL) and used in the next step without further purification; ESMS m/z 198.0 ($M+H^+$).

[0177] Step 2: A mixture of 5-chloro- N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (42 mg, 0.1 mmol), 1-(3-chloropropylsulfonyl) azetidine (Step 1, 0.1 mmol), TEA (70 μ L) and sodium iodide (150 mg) in NMP (1 mL) was placed in a sealed vial which was heated in a microwave for 30 min at 150 °C. The reaction was purified by RP-HPLC to give N^2 -(4-(1-(3-(azetidin-1-ylsulfonyl)propyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white powder; ESMS m/z 573.3 ($M+H^+$).

Example 43

1-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-4-morpholinobutan-1-one (224)




[0178] Step 1: To a solution of HATU (95 mg, 0.255 mmol) and 4-chlorobutanoic acid (36 mg, 0.25 mmol) in DMF (1 mL) was added DIEA (0.1 mL). After stirring at rt for 3 min, 5-chloro- N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (100 mg, 0.24 mmol) in DMF (2 mL) was added. The reaction was stirred at rt for 90 min. at which point LCMS indicates the reaction was complete and that 4-chloro-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-

ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)butan-1-one has been generated; ESMS m/z 516.1 ($M+H^+$).

[0179] Step 2: Morpholine (0.5 mL) was added directly to the crude reaction mixture from Step 1 (1 mL). The resulting solution was stirred at 100 °C in a sealed vial for 1.5 h. The crude product was purified by RP-HPLC to give 1-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-4-morpholinobutan-1-one as a white powder; ESMS m/z 567.3 ($M+H^+$).

Example 44

1-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-morpholinopropan-1-one (225)




[0180] Step 1: A solution of HATU (108 mg, 0.28 mmol), 3-chloropropanoic acid (30 mg, 0.28 mmol) and DIEA (0.1 mL) in DMF (1 mL) was stirred at rt. After 3 min, this solution was transferred to a solution of 5-chloro- N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine solution (111 mg, 0.27 mmol) in DMF (10 mL). The resulting mixture was stirred at rt overnight. The reaction was quenched with water (20 mL) and extracted with EtOAc (3x30 mL). The combined EtOAc layers were washed with brine (10 mL), dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by silica chromatography (0-10% MeOH in DCM gradient with 1% NH_3 additive) to give 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)prop-2-en-1-one as a light brown oil; ESMS m/z 466.1 ($M+H^+$).

[0181] Step 2: A solution of 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)prop-2-en-1-one (41 mg, 0.88 mmol) and morpholine (0.1 mL) in DMF (1 mL) was stirred at rt overnight. The crude product was purified by RP-HPLC to give 1-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-

dimethylphenyl)piperidin-1-yl)-3-morpholinopropan-1-one as a white powder; ESMS m/z 553.2 ($M+H^+$).

Example 45

(S)-4-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)oxazolidin-2-one (230)



[0182] Step 1: To a microwave reaction vessel was added a mixture of 1-bromo-2,5-dimethyl-4-nitrobenzene (1.15 g, 5.0 mmol), dibutyl vinylboronate (1.16 g, 6.3 mmol), $Pd(PPh_3)_4$ (289 mg, 0.25 mmol), CsF (2.28g, 15 mmol) and a mixed solvent (1,2-dimethoxyethane/MeOH=2:1, 15 mL). The mixture was degassed and sealed with N_2 . The reaction vessel was then heated to 130 °C in a microwave reactor for 15 min. After opening the reaction vessel, the mixture was poured into saturated NH_4Cl aqueous solution (30 mL) and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with brine and dried ($MgSO_4$). After removing the drying agent by filtration, the filtrate was concentrated and purified by flash column chromatography (silica gel, 0-30% EtOAc in hexanes gradient) to provide 1,4-dimethyl-2-nitro-5-vinylbenzene as needlelike crystals; ESMS m/z 178.1 ($M + H^+$).

[0183] Step 2: 1,4-dimethyl-2-nitro-5-vinylbenzene was reacted via the asymmetric synthetic method reported by N. Barta *et al.* (*Org. Lett.* **2000**, 2, 2821) followed by purification with preparative RP-HPLC to provide (S)-4-(2,5-dimethyl-4-nitrophenyl)oxazolidin-2-one; ESMS m/z 237.1 ($M + H^+$).


[0184] Step 3: To a solution of (S)-4-(2,5-dimethyl-4-nitrophenyl)oxazolidin-2-one (55 mg, 0.23 mmol) in MeOH (5 mL) was added Pd/C (5mg, 10% w/w). The mixture was degassed and stirred at room temperature under 1 atm. H_2 for 14 h. The Pd/C was removed by filtering through Celite. The filtrate was concentrated to provide (S)-4-(4-amino-2,5-dimethylphenyl)-oxazolidin-2-one which was used in the next step without further purification; ESMS m/z 207.1 ($M + H^+$).

[0185] Step 4: To a reaction tube was added (S)-4-(4-amino-2,5-

dimethylphenyl)oxazolidin-2-one (48 mg, 0.23 mmol), 2,5-dichloro-N-(5-methyl-1*H*-pyrazol-3-yl)pyrimidin-4-amine (77 mg, 0.32 mmol), ¹PrOH (2 mL) and HCl (0.13 mL, 0.52 mmol, 4N in dioxane). The tube was then sealed and heated at 130 °C for 6 h. After cooling down to room temperature, the mixture was concentrated and purified by preparative RP-HPLC to provide (*S*)-4-(4-(5-chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethyl-phenyl)oxazolidin-2-one; ESMS *m/z* 414.1 (*M* + *H*⁺).

Example 46


5-chloro-N²-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (232)



[0186] A mixture of 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (400 mg, 0.96 mmol), iodoethane (92.5 μ L, 1.2 mmol) and triethylamine (201.5 μ L, 1.4 mmol) in 4 mL of DMF was heated at 60°C for 2 h. The reaction mixture was partitioned between EtOAc and water. The combined organic extracts were dried (Na₂SO₄) and concentrated *in vacuo*. The crude product was purified by silica gel chromatography (MeOH/DCM: 1/9) to afford 5-chloro-N²-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine; ¹HNMR (DMSO-*d*₆) δ 8.01 (s, 1H), 7.35 (m, 1H), 7.02 (d, 1H), 6.22 (s, 1H), 2.97 (d, 2H), 2.62 (m, 1H), 2.34 (q, 2H), 2.21 (s, 3H), 2.13 (s, 3H), 1.98 (t, 2H), 1.64 (m, 4H), 1.01 (t, 3H); ESMS *m/z* 444.2 (*M* + *H*⁺).

Example 47


5-chloro-N²-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (235)




[0187] A mixture of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (40 mg, 0.096 mmol), 2-(bromomethyl)-1,1-difluorocyclopropane (32.8 mg, 0.19 mmol) and triethylamine (20.2 μ L, 0.14 mmol) in 1 mL of DMF was heated at 150°C in a microwave reactor for 30 min. The crude product was purified by RP-HPLC to afford 5-chloro- N^2 -(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine; ESMS m/z 506.2 ($M + H^+$).

Example 48


3-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide (236)



[0188] A solution of 5-chloro- N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (41 mg, 0.1 mmol), acrylamide (24 mg, 0.33 mmol) and DIEA (87 μ L, 0.5 mmol) in NMP (1 mL) was stirred at 80 °C for 10 h. LCMS at this point shows the reaction was not complete, so additional acrylamide (2x36mg) was added and the reaction continued until LCMS shows the reaction was complete. The crude product was purified by RP-HPLC to give 3-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide as a white powder; ESMS m/z 483.2 ($M+1$).

Example 494-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1-morpholinobutan-1-one (246)

[0189] A mixture of 5-chloro-*N*²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (41 mg, 0.1 mmol), 4-chloro-1-morpholinobutan-1-one (29 mg, 0.15 mmol) and DIEA (0.1 mL, 0.58 mmol) in NMP (1 mL) was stirred at 80 °C. After 10 h, additional 4-chloro-1-morpholinobutan-1-one (51 mg) was added, and the reaction continued overnight. The crude product was purified by RP-HPLC to give 4-(4-(4-(5-Chloro-4-(5-methyl-1*H*-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1-morpholinobutan-1-one as a white powder; ESMS *m/z* 571.3.3 (*M*+*H*⁺).


Example 50*N*²-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine (251)

[0190] A mixture of *N*²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine (100 mg, 0.22 mmol), 2-(bromomethyl)-1,1-difluorocyclopropane (76.4 mg, 0.45 mmol) and triethylamine (60.6 μL, 0.42 mmol) in 1 mL of DMF was heated at 60°C in a microwave reactor for 2 h. The crude product was purified by RP-HPLC to afford *N*²-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-*N*⁴-(5-methyl-1*H*-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine; ESMS

m/z 540.2 ($M + H^+$).

Example 51


N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine (257)



[0191] A mixture of 2-chloro- N -(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidin-4-amine (513.0 mg, 0.9 mmol), tert-butyl-4-(4-amino-2,5-dimethylphenyl)piperidine-1-carboxylate (282.7 mg, 0.9 mmol), and concentrated aqueous HCl (10 drops) in *i*-PrOH (12 mL) was heated at 150°C in a microwave reactor for 30 min. The crude product was purified by RP-HPLC to afford N^2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine; ESMS m/z 446.2 ($M + H^+$).

Example 52

5-Chloro- N^2 -(2-fluoro-4-(1-((3-isopropyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-5-methylphenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (259)




[0192] A mixture of 5-chloro- N^2 -(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (42 mg, 0.1 mmol), 5-(chloromethyl)-3-isopropyl-1,2,4-oxadiazole (17 mg, 0.10 mmol) and DIEA (86 μ L, 0.5 mmol) in DMF (1 mL) was stirred at rt overnight. The reaction was diluted with EtOAc (50 mL) and washed with water (2x5mL). The EtOAc layer was dried over Na_2SO_4 and concentrated *in vacuo*. The crude product was purified by silica

chromatography (50-100% EtOAc in hexanes gradient) to give 5-Chloro-N²-(2-fluoro-4-(1-((3-isopropyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-5-methylphenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ESMS *m/z* 540.2 (*M*+H⁺).

Example 53

(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(3-isopropyl-1,2,4-oxadiazol-5-yl)methanone (265)



[0193] Step 1: A solution of isobutyronitrile (4.49 mL, 50 mmol) and hydroxylamine (12.3 mL, 200 mmol) in anhydrous ethanol (20 mL) was stirred at 60 °C for two days. The solvent was evaporated. The residue was coevaporated repeatedly with toluene to give (*E*)-*N'*-hydroxyisobutyrimidamide as a light yellow liquid; ¹H NMR (400 MHz, CDCl₃) δ 7.50 (br s, 1H), 4.52 (br s, 2H), 2.43 (septet, *J* = 6.8 Hz, 1H), 1.15 (d, *J* = 6.8 Hz, 6H).


[0194] Step 2: A solution of (*E*)-*N'*-hydroxyisobutyrimidamide (0.51 g, 5 mmol) and trichloroacetic anhydride (2.31 g, 7.5 mmol) in toluene (20 mL) was stirred at 80 °C. After 4 h, the reaction was diluted with EtOAc (50 mL), washed with saturated aqueous NaHCO₃ (2x10 mL), dried over Na₂SO₄ and evaporated to give 3-Isopropyl-5-(trichloromethyl)-1,2,4-oxadiazole as a clear liquid which was used without further purification; ¹H NMR (400 MHz, CDCl₃) δ 3.16 (septet, *J* = 7.2 Hz, 1H), 1.39 (d, *J* = 6.8 Hz, 6H).

[0195] Step 3: A mixture of 5-chloro-*N*²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (42 mg, 0.1 mmol), 3-isopropyl-5-(trichloromethyl)-1,2,4-oxadiazole (46 mg, 0.2 mmol) and DIEA (40 μL, 0.23 mmol) in MeOH (1 mL) was stirred at 70 °C overnight. The crude reaction mixture was purified by RP-HPLC to give (4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(3-

isopropyl-1,2,4-oxadiazol-5-yl)methanone as a white solid; ^1H NMR (400 MHz, CDCl_3) δ 9.67 (br s, 2H), 8.50 (s, 1H), 8.07 (s, 1H), 7.57 (d, $J = 8.4$ Hz, 1H), 7.05 (d, $J = 12.4$ Hz, 1H), 6.16 (s, 1H), 5.53 (s, 1H), 3.98 (d, $J = 13.2$ Hz, 1H), 3.34 (dt, $J = 2.0, 2.8$ Hz, 1H), 3.15 (m, 1H), 2.82 (m, 2H), 2.62 (t, $J = 4.8$ Hz, 2H), 2.34 (s, 3H), 2.25 (s, 3H), 1.90 (m, 2H), 1.76 (2 H); ESMS m/z 554.2 ($\text{M} + \text{H}^+$).

Example 54

5-Chloro- N_2 -(4-(1-(2-(3-isopropyl-1,2,4-oxadiazol-5-yl)ethyl)piperidin-4-yl)-2,5-dimethylphenyl)- N_4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (269)




[0196] Step 1: A mixture of (*E*)-*N'*-hydroxyisobutyrimidamide (510 mg, 5 mmol), 3-chloropropanoic acid (542 mg, 5 mmol) and dicyclohexylcarbodiimide (1.24 g, 6 mmol) in anhydrous dioxane (20 mL) was stirred at 0 °C for 1 h and then at room temperature for an additional 1 h. The reaction was then heated at 80 °C for an additional 18 h. The reaction was filtered and the filtrate was evaporated. The resulting residue was purified by silica chromatography (0-100% EtOAc in hexanes gradient) to give 5-(2-chloroethyl)-3-isopropyl-1,2,4-oxadiazole; ^1H NMR (400 MHz, CDCl_3) δ 3.85 (t, $J = 7.2$ Hz, 2H), 3.27 (t, $J = 7.2$ Hz, 2H), 3.03 (septet, $J = 7.2$ Hz, 1H), 1.13 (d, $J = 7.2$ Hz, 6H); ESMS m/z 175.1 ($\text{M} + \text{H}^+$).

[0197] Step 2: A mixture of 5-chloro- N_2 -(2,5-dimethyl-4-(piperidin-4-yl)phenyl)- N_4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (14 mg, 0.033 mmol), 5-(2-chloroethyl)-3-isopropyl-1,2,4-oxadiazole (9 mg, 0.053 mmol) and DIEA (29 μL , 0.17 mmol) in NMP (1 mL) was stirred at 50 °C for 3 days. The crude reaction mixture was purified by RP-HPLC to give 5-Chloro- N_2 -(4-(1-(2-(3-isopropyl-1,2,4-oxadiazol-5-yl)ethyl)piperidin-4-yl)-2,5-dimethylphenyl)- N_4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ESMS m/z 550.3 ($\text{M} + \text{H}^+$).

Example 55


5-Chloro-*N*₂-(2-fluoro-5-methyl-4-(1-(6-methylpyridazin-3-yl)piperidin-4-yl)phenyl)-*N*₄-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (277)




[0198] A suspension of 5-chloro-*N*₂-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-*N*₄-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (20 mg, 0.05 mmol), 3-chloro-6-methylpyridazine (13 mg, 0.1 mmol) and Cs₂CO₃ (33 mg, 0.1 mmol) in dioxane (1 mL) was stirred at 150 °C in a sealed vial. After 15 h, additional 3-chloro-6-methylpyridazine (26 mg) and Cs₂CO₃ (66 mg) were added. The reaction was continued for an additional 3 h. The crude reaction mixture was purified by RP-HPLC to give 5-Chloro-*N*₂-(2-fluoro-5-methyl-4-(1-(6-methylpyridazin-3-yl)piperidin-4-yl)phenyl)-*N*₄-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid; ESMS *m/z* 508.2 (M+H⁺).

Examples 56 and 57

5-chloro-*N*₂-(2-fluoro-5-methyl-4-((*trans*)-2-methylpiperidin-4-yl)phenyl)-*N*₄-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (279)



5-chloro-*N*₂-(2-fluoro-5-methyl-4-((*cis*)-2-methylpiperidin-4-yl)phenyl)-*N*₄-(5-methyl-1*H*-pyrazol-3-yl)pyrimidine-2,4-diamine (280)



[0199] Step 1: A solution of *tert*-butyl-2-methyl-4-oxopiperidine-1-carboxylate (1 g, 4.69 mmol) in THF (20 mL) was added dropwise into a cooled (-78 °C), vigorously stirring solution of LDA (3.75 mL of 1.5 M solution in cyclohexanes, 5.63 mmol) in THF (20 mL), under N₂. The reaction mixture was stirred at -78 °C for 30 min before adding a solution of phenyl trifluorosulfonimide (1.84 g, 5.16 mmol) in THF (20 mL). Following this addition, the reaction mixture was warmed to room temperature and stirred for 3 h. The reaction was cooled to 0 °C and quenched with 100 mL of saturated aqueous NH₄Cl, followed by filtration through Celite. The filtrate was added to 100 mL of EtOAc and the layers were separated. The organic layer was washed with water, dried over MgSO₄ and concentrated. The crude product was purified by silica chromatography (0-30% EtOAc in hexanes gradient and checked by TLC stained with 2% of KMnO₄ in EtOH) to afford *tert*-butyl 2-methyl-4-(trifluoromethylsulfonyloxy)-5,6-dihydropyridine-1(2H)-carboxylate as a yellow solid.

[0200] Step 2: To a mixture of 2-fluoro-5-methyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (587 mg, 2.33 mmol), *tert*-butyl 2-methyl-4-(trifluoromethylsulfonyloxy)-5,6-dihydropyridine-1(2H)-carboxylate (968 mg, 2.80 mmol) and sodium carbonate (1.73 g, 16.35 mmol) in DMF/H₂O (20/5 mL) was added tetrakis(triphenylphosphine) palladium (0) (135 mg, 5% mmol). The reaction tube was sealed, the mixture was purged with N₂ for 3 min and then heated at 100°C under N₂ for 8 h. The reaction was cooled to room temperature and poured into saturated aqueous ammonia chloride solution. The crude reaction mixture was extracted with ethyl acetate (3 x 15 mL). The organic extracts were combined, washed with brine and concentrated. The crude product was purified with silica chromatography (50% ethyl acetate in hexanes) to afford *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)-6-methyl-5,6-dihydropyridine-1(2H)-carboxylate as a yellow oil. The obtained oil was dissolved in methanol (20 mL). To the solution was added Pd/C (10%). The reaction mixture was degassed and purged with H₂ for several times and stirred vigorously under 1 atm. H₂ overnight. The mixture was filtered and the filtrate concentrated to afford *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)-2-methylpiperidine-1-carboxylate as a white solid. ESMS *m/z* 267 (M -56+ H⁺).


[0201] Step 3: A mixture of 2,5-dichloro-*N*-(5-methyl-1H-pyrazol-3-yl)pyrimidin-4-amine (346 mg, 1.42 mmol) and *tert*-butyl 4-(4-amino-5-fluoro-2-methylphenyl)-2-

methylpiperidine-1-carboxylate (352 mg, 1.09 mmol) in 2-propanol (10 mL) was treated with conc. HCl (12N, 0.43 mL). The mixture was sealed and heated in a microwave at 130 °C for 45 min. The mixture was concentrated. The residue was purified by RP-HPLC to afford 5-Chloro-N²-(2-fluoro-5-methyl-4-(2-methylpiperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as a white solid. ESMS m/z 430.1 ($M + H^+$).

[0202] Step 4: 5-Chloro-N²-(2-fluoro-5-methyl-4-(2-methylpiperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (20 mg, 0.046 mmol) was dissolved in a solvent mixture of DCM and methanol (0.5 mL, 10/1 v/v + 0.175N NH₃). The mixture was applied to preparative TLC in order to separate the *cis* and *trans* stereoisomers (eluent: 6% MeOH in DCM with 0.1N NH₃) of 5-Chloro-N²-(2-fluoro-5-methyl-4-(2-methylpiperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as white solids. The *trans* isomer was the preparative TLC top band and the *cis* isomer was the preparative TLC bottom band; ESMS m/z 430.1 ($M + H^+$); *cis* stereoisomer ¹H NMR (400 MHz, MeOD-d₄): δ 8.03 (s, 1H), 7.84 (d, 1H), 7.02 (d, 1H), 6.24 (d, 1H), 3.90-3.85 (m, 1H), 2.32 (s, 3H), 2.27 (s, 3H), 2.03-1.84 (m, 2H), 1.65-1.54 (m, 1H), 1.55 (d, 3H), 1.38-1.19 (m, 4H).

Example 58

3-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclopent-2-en-one (282)




[0203] A mixture of 5-chloro-N²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (43 mg, 0.1 mmol), cyclopentane-1,3-dione (89 mg, 9.1 mmol), sodium cyanoborohydride (19 mg, 0.3 mmol) and DIEA (0.2 mL, 11 mmol) in MeOH (1 mL) was stirred in a sealed vial at 60 °C. After 2.5 h, additional sodium cyanoborohydride (19 mg, 0.3 mmol) was added and the reaction continued to stir at 60 °C overnight. The mixture was purified by RP-HPLC to give 3-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-

ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclopent-2-en-one as an off white solid; ^1H NMR (400 MHz, DMSO- d_6) δ 9.97 (br s, 1H), 9.83 (br s, 1H), 8.90 (s, 1H), 8.32 (d, $J = 6.8$ Hz, 1H), 7.88 (d, $J = 12.8$ Hz, 1H), 6.97 (s, 1H), 6.04 (s, 1H), 5.30 (br s, 1H), 4.71 (d, $J = 12.0$ Hz, 2H), 4.04 (t, $J = 12.4$ Hz, 2H), 3.84 (m, 1H), 3.53 (m, 2H), 3.13 (m, 2H), 3.08 (s, 3H), 2.96 (s, 3H), 2.59-2.40 (m, 4H); ESMS m/z 496.2 ($\text{M} + \text{H}^+$).

Example 59


4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-1-ethylpiperidine-1-oxide (289)



[0204] To a solution of 5-chloro- N^2 -(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (22.2 mg, 0.05 mmol) in DCM (10 mL) and MeOH (1 mL) was added MCPBA (11.2 mg, 0.07 mmol). The reaction was stirred at room temperature overnight. The reaction mixture was partitioned between DCM and saturated aqueous NaHCO_3 , the combined organic extracts were dried (Na_2SO_4), and concentrated *in vacuo*. The crude product was purified by RP-HPLC to afford 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-1-ethylpiperidine-1-oxide; ^1H NMR (MeOD- d_4) δ 7.91 (s, 1H), 7.69 (d, 1H), 7.00 (d, 1H), 6.15 (s, 1H), 3.28 (m, 4H), 2.91 (m, 1H), 2.32 (m, 2H), 2.20 (s, 3H), 2.16 (s, 3H), 1.83 (s, 2H), 1.64 (d, 2H), 1.31 (t, 3H).; ESMS m/z 460.2 ($\text{M} + \text{H}^+$).

Example 60

5-Chloro- N^2 -(2-fluoro-5-methyl-4-(1-(pyrimidin-5-ylmethyl)piperidin-4-yl)phenyl)- N^4 -(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (314)











[0205] To a solution of 5-chloro-*N*²-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine (45 mg, 0.1 mmol) and pyrimidine-5-carbaldehyde (32 mg, 0.3 mmol) in DCM (1.5 mL) was added NaBH(OAc)₃ (64 mg, 0.3 mmol), followed by addition of AcOH (12 uL). After stirring at room temperature in a sealed vial for 16 h, the reaction was quenched with EtOAc (50 mL). The reaction mixture was sequentially washed with water (10 mL), brine (5 mL), dried over Na₂SO₄ and concentrated *in vacuo*. The resulting residue was purified by silica chromatography (0-10% MeOH in EtOAc gradient with 1% NH₃ in MeOH additive) to give 5-Chloro-*N*²-(2-fluoro-5-methyl-4-(1-(pyrimidin-5-ylmethyl)piperidin-4-yl)phenyl)-*N*⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine as an off white solid; ¹H NMR (400 MHz, CD₃OD) δ 9.09 (s, 1H), 8.81 (s, 2H), 8.00 (s, 1H), 7.73 (d, *J* = 8.4 Hz, 1H), 6.99 (d, *J* = 13.2 Hz, 1H), 6.23 (br s, 1H), 3.65 (s, 2H), 3.03 (m, 2H), 2.76 (m, 1H), 2.27 (s, 3H), 2.25 (s, 3H), 2.25 (m, 2H), 1.80-1.68 (m, 4H); ESMS *m/z* 508.2 (M+H⁺).





[0206] The following compounds in Table 1 are obtained by repeating the procedures described in examples above and using appropriate starting materials.





Table 1





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
1	5-chloro-<i>N</i>²-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-<i>N</i>⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	¹ H NMR (400 MHz, DMSO-d ₆) δ 10.33 (br, 1H), 9.90 (br, 1H), 9.60 (br, 1H), 8.22 (s, 1H), 7.56 (d, 1H), 7.00 (d, 1H), 6.20 (s, 1H), 3.53-3.50 (m, 2H), 3.16-3.10 (m, 2H), 2.97-2.92 (m, 1H), 2.81 (d, 3H), 2.26 (s, 3H), 2.16 (s, 3H), 1.94-1.91 (m, 2H), 1.83-1.73 (m, 2H); ESMS <i>m/z</i> 430.1 (M + H ⁺).	0.011





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
2	 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 416.1 (M + H ⁺).	0.050
3	 5-chloro-N2-(2-isopropoxy-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 470.2 (M + H ⁺).	0.104
4	 N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine	ESMS m/z 450.2 (M + H ⁺).	0.150
5	 N2-(2-isopropoxy-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-5-methyl-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 450.3 (M + H ⁺).	0.805





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
6	 5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 426.2 (M + H ⁺).	0.015
7	 5-chloro-N2-(5-chloro-2-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 446.1 (M + H ⁺).	0.015
8	 5-chloro-N2-(4-fluoro-2-methyl-5-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 461.2 (M + H ⁺).	1.93
9	 3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one	ESMS m/z 461.2 (M + H ⁺).	0.087





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
10	 (S)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one	ESMS m/z 461.2 (M + H ⁺).	0.869
11	 (R)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one	ESMS m/z 461.2 (M + H ⁺).	0.065
12	 5-chloro-N2-(4-(1-isopropylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 454.2 (M + H ⁺).	0.010
13	 5-chloro-N2-(4-fluoro-5-(1-isopropylpiperidin-4-yl)-2-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 458.2 (M + H ⁺).	





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
14	 5-chloro-N2-(2,5-dimethyl-4-(piperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 412.2 ($M + H^+$).	0.016
15	 5-chloro-N2-(2,5-dimethyl-4-(piperidin-3-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 412.2 ($M + H^+$).	0.050
16	 5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 426.2 ($M + H^+$).	0.019
17	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)pyrimidine-2,4-diamine	ESMS m/z 442.2 ($M + H^+$).	0.160





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
18	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 438.2 (M + H ⁺).	0.095
19	 5-Chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	¹ H NMR (400 MHz, MeOD-d ₄) 8.12 (s, 1H), 7.75 (d, 1H), 7.03 (s, 1H), 6.27 (s, 1H), 4.25 (s, 2H), 3.69-3.67 (m, 2H), 3.27-3.20 (m, 3H), 2.54 (s, 3H), 2.37 (s, 3H), 2.34 (s, 3H), 2.30 (s, 3H), 2.10-1.95 (m, 4H); ESMS <i>m/z</i> 525.1 (M + H ⁺).	0.056
20	 5-chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 521.2 (M + H ⁺).	0.011
21	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyridazin-4-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 504.2 (M + H ⁺)	0.002





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
22	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanol	ESMS m/z 460.2 (M + H ⁺).	0.059
23	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)ethanol	ESMS m/z 456.21 (M + H ⁺).	0.006
24	 5-chloro-N²-(2-fluoro-5-methyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 512.2 (M + H ⁺).	0.085
25	 5-chloro-N²-(2,5-dimethyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 508.2 (M + H ⁺).	0.031





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
26	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 498.2 (M + H ⁺).	0.142
27	 5-chloro-N2-(5-chloro-2-methyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 528.2 (M + H ⁺).	0.044
28	 5-chloro-N2-(4-fluoro-2-methyl-5-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 512.2 (M + H ⁺).	5.46
29	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl) piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 524.2 (M + H ⁺).	0.023





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
30	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 528.2 (M + H ⁺).	0.071
31	 3-(4-(5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-fluoro-4-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 528.2 (M + H ⁺).	6.47
32	 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 513.2 (M + H ⁺).	0.042
33	 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 509.2 (M + H ⁺).	0.006





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
34	 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide	ESMS m/z 499.2 (M + H ⁺).	0.049
35	 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	ESMS m/z 495.2 (M + H ⁺).	0.011
36	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-(trifluoromethyl)phenyl)piperidin-1-yl)acetamide	ESMS m/z 527.2 (M + H ⁺).	0.019
37	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	ESMS m/z 469.2 (M + H ⁺).	0.013




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
38	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 483.2 ($M + H^+$).	0.010
39	 2-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide	¹ H NMR (400 MHz, MeOD- <i>d</i> ₄) δ 8.11 (s, 1H), 7.76 (d 1H), 7.09 (d, 1H), 6.27 (s, 1H), 4.0 (s, 2H), 3.75-3.73 (m, 2H), 3.22-3.13 (m, 3H), 2.34 (s, 3H), 2.30 (s, 3H), 2.04 - 2.00 (m, 4H); ESMS m/z 473.2 ($M + H^+$).	0.019
40	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	ESMS m/z 469.2 ($M + H^+$).	
41	 2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	ESMS m/z 469.2 ($M + H^+$).	0.026





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
42	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 487.2 (M + H ⁺).	0.019
43	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 483.2 (M + H ⁺).	
44	 2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 483.2 (M + H ⁺).	0.015
45	 2-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 503.2 (M + H ⁺).	0.018





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
46	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 499.2 (M + H ⁺).	0.013
47	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methyl-2-(trifluoromethyl)phenyl)piperidin-1-yl)acetamide	ESMS m/z 523.2 (M + H ⁺).	0.022
48	 2-(4-(5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-fluoro-4-methylphenyl)piperidin-1-yl)propanamide	ESMS m/z 487.2 (M + H ⁺).	3.68
49	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide	ESMS m/z 487.2 (M + H ⁺).	0.015





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
50	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylpropanamide	ESMS m/z 501.2 (M + H ⁺).	0.022
51	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide	ESMS m/z 483.2 (M + H ⁺).	0.009
52	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylpropanamide	ESMS m/z 497.2 (M + H ⁺).	0.006
53	 2-(4-(5-fluoro-2-methyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)propanamide	ESMS m/z 521.2 (M + H ⁺).	0.078





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
54	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanoic acid	ESMS m/z 488.2 (M + H ⁺).	1.53
55	 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 522.1 (M + H ⁺).	0.024
56	 5-chloro-N2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 518.2 (M + H ⁺).	0.006
57	 5-chloro-N2-(4-(1-(2-(ethylsulfonyl)ethyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 536.2 (M + H ⁺).	0.021




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
58	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N,N-dimethylethanesulfonamide	ESMS m/z 551.2 (M + H ⁺).	0.036
59	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanesulfonamide	ESMS m/z 523.2 (M + H ⁺).	0.038
60	 N2-(4-(1-(2-(azetidin-1-ylsulfonyl)ethyl)piperidin-4-yl)-5-chloro-2-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 579.2 (M + H ⁺).	0.011




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
61	 N2-(4-(1-(2-(azetidin-1-ylsulfonyl)ethyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 563.2 (M + H ⁺).	0.056
62	 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 494.2 (M + H ⁺).	0.005
63	 5-chloro-N2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 490.2 (M + H ⁺).	0.002
64	 5-chloro-N2-(4-(1-(cyclopropylsulfonyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 520.2 (M + H ⁺).	0.059




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
65	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-3-ylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 557.2 ($M + H^+$).	0.076
66	 Ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate	^1H NMR (400 MHz, MeOD- d_4) δ 8.13 (s, 1H), 7.54 (d, 1H), 7.10 (d, 1H), 6.28 (s, 1H), 4.30-4.26 (m, 2H), 4.14 (q, 2H), 2.99-2.96 (m, 3H), 2.33 (s, 3H), 2.29 (s, 3H), 1.8-1.77 (m, 2H), 1.64-1.55 (m, 2H), 1.28 (t, 3H); ESMS m/z 488.2 ($M + H^+$).	0.064
67	 ethyl 4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidine-1-carboxylate	ESMS m/z 504.2 ($M + H^+$).	0.182
68	 isopropyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate	ESMS m/z 502.2 ($M + H^+$).	0.080




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
69	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-N,N-dimethylpiperidine-1-carboxamide	ESMS m/z 487.2 (M + H ⁺).	0.011
70	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)-N,N-dimethylpiperidine-1-carboxamide	ESMS m/z 483.2 (M + H ⁺).	0.006
71	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)-N-ethylpiperidine-1-carboxamide	ESMS m/z 483.2 (M + H ⁺).	0.002
72	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidine-1-carboxamide	ESMS m/z 455.2 (M + H ⁺).	0.006




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
73	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	¹ H NMR (400 MHz, MeOD- <i>d</i> ₄) δ 8.08 (s, 1H), 7.71 (d, 1H), 7.03 (d, 1H), 6.26 (s, 1H), 4.71-4.67 (m, 2H), 3.81-3.77 (m, 2H), 3.14-3.12 (m, 2H), 2.98 (s, 3H), 2.96 (s, 3H), 2.90-2.83 (m, 1H), 2.34 (s, 3H), 2.28 (s, 3H), 1.87 (d, 2H), 1.76-1.55 (m, 2H); ESMS <i>m/z</i> 501.2 (M + H ⁺).	0.019
74	 1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS <i>m/z</i> 517.2 (M + H ⁺).	0.007
75	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS <i>m/z</i> 497.2 (M + H ⁺).	0.011
76	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS <i>m/z</i> 513.2 (M + H ⁺).	0.009




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
77	 1-(4-(2,5-dichloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS m/z 537.1 (M + H ⁺).	0.007
78	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-3-(dimethylamino)propan-1-one	ESMS m/z 515.2 (M + H ⁺).	0.138
79	 2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanone	ESMS m/z 473.2 (M + H ⁺).	0.075




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
80	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(methylamino)ethanone	ESMS m/z 487.2 (M + H ⁺).	0.103
81	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(diethylamino)ethanone	ESMS m/z 529.3 (M + H ⁺).	0.030
82	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(methylamino)ethanone	ESMS m/z 487.2 (M + H ⁺).	




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
83	 1-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS m/z 497.2 (M + H ⁺).	0.013
84	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(diethylamino)ethanone	ESMS m/z 529.3 (M + H ⁺).	
85	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(methylamino)propan-1-one	ESMS m/z 501.2 (M + H ⁺).	0.042




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
86	 2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propan-1-one	ESMS m/z 487.2 (M + H ⁺).	0.141
87	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-methyl-2-(methylamino)propan-1-one	ESMS m/z 515.2 (M + H ⁺).	0.017
88	 1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-2-methyl-2-(methylamino)propan-1-one	ESMS m/z 531.2 (M + H ⁺).	0.007




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
89	 2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methylpropan-1-one	ESMS m/z 497.2 (M + H ⁺).	0.023
90	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-(methylamino)cyclopropyl)methanone	ESMS m/z 509.2 (M + H ⁺).	0.005
91	 (1-aminocyclobutyl)(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methanone	ESMS m/z 509.2 (M + H ⁺).	0.052





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
92	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-(dimethylamino)cyclobutyl)methanone	ESMS m/z 537.3 (M + H ⁺).	0.015
93	 (1-aminocyclopropyl)(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone	ESMS m/z 499.2 (M + H ⁺).	0.033
94	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(dimethylamino)cyclopropyl)methanone	ESMS m/z 527.2 (M + H ⁺).	0.013





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
95	 1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-2-morpholinoethanone	ESMS m/z 559.2 (M + H ⁺).	0.006
96	 2-(azetidin-1-yl)-1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)ethanone	ESMS m/z 529.2 (M + H ⁺).	0.024
97	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-morpholinoethanone	ESMS m/z 543.2 (M + H ⁺).	0.020




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
98	 2-(azetidin-1-yl)-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanone	ESMS m/z 513.2 (M + H ⁺).	0.064
99	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(pyrrolidin-1-yl)ethanone	ESMS m/z 527.2 (M + H ⁺).	0.038
100	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(1H-pyrrol-1-yl)ethanone	ESMS m/z 523.2 (M + H ⁺).	0.025





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
101	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(piperidin-1-yl)ethanone	ESMS m/z 537.3 (M + H ⁺).	0.007
102	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-morpholinoethanone	ESMS m/z 539.3 (M + H ⁺).	0.009
103	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(2,6-dimethylpiperidin-1-yl)ethanone	ESMS m/z 565.3 (M + H ⁺).	0.009




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
104	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(2,2,6,6-tetramethylpiperidin-1-yl)ethanone	ESMS m/z 593.3 (M + H ⁺).	0.018
105	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(4-methylpiperazin-1-yl)ethanone	ESMS m/z 556.3 (M + H ⁺).	0.051
106	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methoxyethanone	ESMS m/z 484.2 (M + H ⁺).	0.002




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
107	 N-(2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-oxoethyl)acetamide	ESMS m/z 515.2 (M + H ⁺).	0.057
108	 2-(dimethylamino)ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate	ESMS m/z 531.2 (M + H ⁺).	0.053
109	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-N-(2-(dimethylamino)ethyl)piperidine-1-carboxamide	ESMS m/z 530.2 (M + H ⁺).	0.060
110	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)ethanone	ESMS m/z 454.2 (M + H ⁺).	0.013




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
111	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(morpholino)methanone	ESMS m/z 529.2 (M + H ⁺).	0.008
112	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(4-methylpiperazin-1-yl)methanone	ESMS m/z 542.2 (M + H ⁺).	0.021
113	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidine-1-carbonyl)imidazolidin-2-one	ESMS m/z 524.2 (M + H ⁺).	0.013
114	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methoxyethanone	ESMS m/z 569.2 (M + H ⁺).	0.048




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
115	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(piperidin-4-yl)methanone	ESMS m/z 527.2 (M + H ⁺).	1.57
116	 (S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(pyrrolidin-2-yl)methanone	ESMS m/z 513.2 (M + H ⁺).	0.137
117	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methylazetidin-3-yl)methanone	ESMS m/z 513.2(M + H ⁺).	0.099




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
118	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methylpiperidin-4-yl)methanone	ESMS m/z 541.2 (M + H ⁺).	0.081
119	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carbonyl)azetidin-2-one	ESMS m/z 513.2 (M + H ⁺).	2.99
120	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylazetidin-3-yl)methanone	ESMS m/z 509.2 (M + H ⁺).	0.025
121	 (S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylpyrrolidin-2-yl)methanone	ESMS m/z 523.2 (M + H ⁺).	0.017




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
122	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylpiperidin-4-yl)methanone	ESMS m/z 537.2 ($M + H^+$).	
123	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)((2R,4R)-4-hydroxy-1-methylpyrrolidin-2-yl)methanone	ESMS m/z 539.3 ($M + H^+$).	0.048
124	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1H-pyrrol-2-yl)methanone	ESMS m/z 509.2 ($M + H^+$).	0.145




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
125	 (S)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl(1-isopropylpiperidin-2-yl)methanone	ESMS m/z 569.3 (M + H ⁺).	0.012
126	 (S)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl(4-isopropylmorpholin-3-yl)methanone	ESMS m/z 571.3 (M + H ⁺).	0.029
127	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl(2,4-dimethyloxazol-5-yl)methanone	ESMS m/z 539.2 (M + H ⁺).	0.027




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
128	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)((2S)-4-hydroxy-1-methylpyrrolidin-2-yl)methanone	ESMS m/z 543.2 (M + H ⁺).	0.175
129	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1,3,5-trimethyl-1H-pyrazol-4-yl)methanone	ESMS m/z 552.2 (M + H ⁺).	0.025
130	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methyl-5-(trifluoromethyl)-1H-pyrazol-4-yl)methanone	ESMS m/z 592.2 (M + H ⁺).	0.061





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
131	 azetidin-3-yl(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone	¹ H NMR (400 MHz, MeOD- <i>d</i> ₄) δ 8.08 (s, 1H), 7.68 (d, 2H), 7.04 (d, 2H), 6.26 (s, 1H), 4.74-4.68 (m, 2H), 4.39-4.35 (m, 1H), 4.29-4.24 (m, 2H), 4.11 -4.07 (m, 1H), 3.74-3.71 (m, 1H), 3.25 -3.24 (m, 1H), 3.07-3.04 (m, 1H), 2.87-2.80 (m, 1H), 2.33 (s, 3H), 2.28 (s, 3H), 1.87-1.84 (m, 2H), 1.66-1.56 (m, 2H); ESMS <i>m/z</i> 499.2 (M + H ⁺).	1.25
132	 (S)-azetidin-2-yl(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone	ESMS <i>m/z</i> 499.2 (M + H ⁺).	0.161
133	 (R)-4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylpyrrolidin-2-yl)methanone	ESMS <i>m/z</i> 523.3 (M + H ⁺).	0.013





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
134	 (R)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl) piperidin-1-yl)(1-methylpyrrolidin-2-yl)methanone	ESMS m/z 527.2 (M + H ⁺).	0.021
135	 (R)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-isopropylpyrrolidin-2-yl)methanone	ESMS m/z 555.3 (M + H ⁺).	0.048
136	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(4-methyloxazol-5-yl)methanone	ESMS m/z 525.2 (M + H ⁺).	0.035





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
137	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(3,5-dimethylisoxazol-4-yl)methanone	ESMS m/z 539.2 (M + H ⁺).	0.018
138	 (S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(2-methylpyrrolidin-2-yl)methanone	ESMS m/z 527.2 (M + H ⁺).	0.091
139	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)((2S,3R)-3-hydroxy-1-methylpyrrolidin-2-yl)methanone	ESMS m/z 543.2 (M + H ⁺).	0.175





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
140	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1H-tetrazol-1-yl)methanone	ESMS m/z 512.2 (M + H ⁺).	0.139
141	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(2-methyl-5-(trifluoromethyl)oxazol-4-yl)methanone	ESMS m/z 593.2 (M + H ⁺).	0.069
142	 6-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carbonyl)piperidin-2-one	ESMS m/z 541.2 (M + H ⁺).	0.282




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
143	 (S)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl(morpholin-3-yl)methanone	ESMS m/z 529.2 (M + H ⁺).	0.071
144	 5-chloro-N²-(2-methyl-5-chloro-4-(1-(tetrahydro-1,1-dioxido-3-thienyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 550.1 (M + H ⁺).	0.007
145	 5-chloro-N²-(2-fluoro-5-methyl-4-(1-(tetrahydro-1,1-dioxido-3-thienyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 534.2 (M + H ⁺).	0.018




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
146	 5-chloro-N2-(2,5-dimethyl-4-(1-(tetrahydro-1,1-dioxido-2H-thiopyran-4-yl))piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 544.2 (<i>M</i> + <i>H</i> ⁺).	0.258
147	 5-chloro-N2-(2-chloro-5-methyl-4-(1-(1,1-dioxido-3-thietanyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 536.1 (<i>M</i> + <i>H</i> ⁺).	0.016
148	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1,1-dioxido-3-thietanyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 520.2 (<i>M</i> + <i>H</i> ⁺). ¹ H NMR (400 MHz, DMSO- <i>d</i> ₆ + 1 drop D ₂ O) δ 8.02 (s, 1H), 7.37 (s, 1H), 7.02 (s, 1H), 6.23 (s, 1H), 4.28-4.22 (m, 2H), 4.12-4.07 (m, 2H), 3.22-3.18 (m, 1H), 2.95-2.92 (m, 2H), 2.68-2.62 (m, 1H), 2.22 (s, 3H), 2.13 (s, 3H), 2.07-2.01 (m, 2H), 1.71-1.55 (m, 4H).	0.018
149	 5-chloro-N2-(4-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 522.2 (<i>M</i> + <i>H</i> ⁺).	0.955




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
150	 5-chloro-N2-(2-fluoro-5-methyl-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 414.1 (M + H ⁺).	0.025
151	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-2-one	ESMS m/z 426.2 (M + H ⁺).	0.009
152	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)tetrahydro-2H-pyran-2-one	ESMS m/z 427.2 (M + H ⁺).	0.155
153	 5-chloro-N2-(2-fluoro-4-morpholino-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 472.1 (M + H ⁺).	0.091




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
154	 5-Chloro-<i>N</i>²-[4-(1,1-dioxo-tetrahydro-1λ⁶-thiophen-3-yl)-2,5-dimethyl-phenyl]-<i>N</i>⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)-pyrimidine-2,4-diamine	ESMS <i>m/z</i> 447.1 (M + H ⁺).	0.012
155	 1-(4-(5-chloro-4-(5-methyl-1<i>H</i>-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperazin-2-one	ESMS <i>m/z</i> 427.2 (M + H ⁺).	1.65
156	 5-chloro-<i>N</i>⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)-<i>N</i>²-(2-methyl-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)phenyl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 481.2 (M + H ⁺)	0.040
157	 5-chloro-<i>N</i>²-(2-fluoro-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)phenyl)-<i>N</i>⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 485.2 (M + H ⁺).	0.075




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
158	 3-(5-chloro-2-(2,5-dimethyl-4-(piperidin-4-yl)phenylamino)pyrimidin-4-ylamino)pyridin-2(1H)-one	ESMS m/z 425.2 (M + H ⁺).	0.062
159	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methylisoxazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 431.2 (M + H ⁺).	0.073
160	 3-(5-chloro-2-(2,5-dimethyl-4-(piperidin-4-yl)phenylamino)pyrimidin-4-ylamino)pyridin-2(1H)-one	ESMS m/z 425.2 (M + H ⁺).	
161	 5-chloro-N2-(2-isopropoxy-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 456.2 (M + H ⁺).	0.332




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
162	 ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidine-1-carboxylate	ESMS m/z 438.9 ($M + H^+$).	0.013
163	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide	ESMS m/z 487.2 ($M + H^+$)	0.056
164	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylpropanamide	ESMS m/z 501.2 ($M + H^+$); ^1H NMR (MeOD- d_4) δ 8.25 (s, 1H), 7.56 (m, 1H), 7.20 (d, 1H), 6.46 (s, 1H), 3.97 (m, 1H), 3.79 (m, 1H), 3.62 (m, 1H), 3.25 (m, 3H), 2.84 (s, 3H), 2.40 (d, 6H), 2.08 (m, 4H), 1.62 (m, 3H).	0.033




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
165	 (R)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide	ESMS m/z 487.2 ($M + H^+$); ^1H NMR (MeOD- d_4) δ 8.22 (s, 1H), 7.61 (t, 1H), 7.18 (d, 1H), 6.40 (s, 1H), 3.79 (m, 1H), 3.65 (m, 1H), 3.54 (m, 1H), 3.22 (m, 3H), 2.39 (d, 6H), 2.08 (m, 4H), 1.66 (d, 3H)	0.038
166	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide	ESMS m/z 483.2 ($M + H^+$)	0.007
167	 (R)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide	ESMS m/z 483.2 ($M + H^+$)	0.006




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
168	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylpropanamide	ESMS m/z 497.2 (M + H ⁺)	0.01
169	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclopropyl)methanone	ESMS m/z 527.2 (M + H ⁺); ¹ HNMR (MeOD-d ₄) δ 8.24 (s, 1H), 7.49 (d, 1H), 7.23 (d, 1H), 6.47 (s, 1H), 4.52 (d, 2H), 3.26 (q, 2H), 3.18 (m, 3H), 2.40 (s, 6H), 1.91(d, 2 H), 1.71(m, 2H), 1.49 (m, 3H), 1.39 (t, 3H).	0.004
170	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-(ethylamino)cyclopropyl)methanone	ESMS m/z 523.3 (M + H ⁺)	0.003




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
171	 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-(trifluoromethyl)-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 470.1 (M + H ⁺)	0.067
172	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 541.2 (M + H ⁺)	0.069
173	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide	ESMS m/z 527.2 (M + H ⁺)	0.049




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
174	 5-chloro-N2-(2,5-dimethyl-4-(1-(tetrahydro-2H-thiopyran-4-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 512.2 (M + H ⁺)	0.008
175	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(ethylamino)-2-methylpropan-1-one	ESMS <i>m/z</i> 525.3 (M + H ⁺)	0.005
176	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclobutyl)methanone	ESMS <i>m/z</i> 541.3 (M + H ⁺)	0.084




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
177	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(methylamino)cyclobutyl)methanone	ESMS m/z 527.2 (M + H ⁺)	0.08
178	 N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-5-methyl-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 392.2 (M + H ⁺)	3.229
179	 (S)-3-(4-(2,5-dimethyl-4-(5-methyl-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 504.3 (M + H ⁺)	0.12




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
180	 1-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS m/z 497.2 (M + H ⁺)	0.025
181	 2-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	ESMS m/z 469.2 (M + H ⁺)	0.024
182	 2-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 483.2 (M + H ⁺)	0.019




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
183	 (S)-5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 426.2 (M + H ⁺)	
184	 (R)-5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 426.2 (M + H ⁺)	
185	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)butanamide	ESMS m/z 497.2 (M + H ⁺)	0.009





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
186	 1-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS m/z 523.3 (M + H ⁺)	0.012
187	 (1-aminocyclopropyl)(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methanone	ESMS m/z 495.2 (M + H ⁺)	0.095
188	 (1-aminocyclopropyl)(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methanone	ESMS m/z 521.2 (M + H ⁺)	0.019





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
189	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylbutanamide	ESMS m/z 511.3 (M + H ⁺)	0.007
190	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-ethylbutanamide	ESMS m/z 525.3 (M + H ⁺)	0.007
191	 5-chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-2-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 521.2 (M + H ⁺)	0.113




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
192	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-hydroxycyclopropyl)methanone	ESMS m/z 500.2 (M + H ⁺)	0.021
193	 (2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-hydroxycyclopropyl)methanone	ESMS m/z 496.2 (M + H ⁺)	0.076
194	 5-chloro-N2-(2,5-dimethyl-4-(1-(3-morpholinopropylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 603.3 (M + H ⁺)	0.013




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
195	 N2-(4-(1-(3-(azetidin-1-yl)propylsulfonyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 573.2 (M + H ⁺)	0.027
196	 (3R,4R)-1-(3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-ylsulfonyl)propyl)pyrrolidine-3,4-diol	ESMS m/z 619.3 (M + H ⁺)	0.37
197	 (2S)-3-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 524.2 (M + H ⁺)	0.21





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
198	 1-((4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methyl)cyclopropanecarbonitrile	ESMS m/z 491.2 (M + H ⁺)	0.022
199	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetonitrile	ESMS m/z 455.2 (M + H ⁺)	0.077
200	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanenitrile	ESMS m/z 469.2 (M + H ⁺)	0.036




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
201	 (2S)-3-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 524.2 (M + H ⁺)	0.046
202	 2-(4-(2,5-dimethyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)acetamide	ESMS m/z 503.2 (M + H ⁺); ¹ HNMR (MeOD-d ₄) δ 8.23 (s, 1H), 7.26 (s, 1H), 7.20 (s, 1H), 6.20 (s, 1H), 4.00 (s, 2H), 3.75 (d, 2H), 3.55 (m, 2H), 3.14 (m, 1H), 2.36 (s, 3H), 2.26 (s, 3H), 2.24 (s, 3H), 2.08 (m, 4H).	0.02
203	 2-(4-(2,5-dimethyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 517.3 (M + H ⁺)	0.015
204	 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)ethanol	ESMS m/z 482.2 (M + H ⁺)	0.016




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
205	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(4-(1-ethylpiperidin-4-yl)-2,5-dimethylphenyl)pyrimidine-2,4-diamine	ESMS m/z 466.2 (M + H ⁺)	0.019
206	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carbaldehyde	ESMS m/z 444.2 (M + H ⁺)	0.03
207	 (R)-2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 483.2 (M + H ⁺)	0.026
208	 (S)-2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 483.2 (M + H ⁺)	0.01




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
209	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 500.2 (M + H ⁺); ¹ H NMR (DMSO- <i>d</i> ₆ , 400 MHz) δ 12.08 (br, 1H), 9.58 (br, 1H), 8.65 (d, <i>J</i> = 28.4 Hz, 1H), 8.02 (s, 1H), 7.36 (d, <i>J</i> = 8.4 Hz, 1H), 7.03 (d, <i>J</i> = 12.4 Hz, 1H), 6.08 (br, 1H), 3.89 (dd, <i>J</i> = 10.8, 3.6 Hz, 2H), 3.28 (t, <i>J</i> = 10.4 Hz, 2H), 2.99 (d, <i>J</i> = 10.8 Hz, 2H), 2.64-2.43 (m, 2H), 2.24-2.15 (m, 8H), 1.71-1.40 (m, 8H)	0.027
210	 5-chloro-N2-(4-(1-cyclopropylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 456.2 (M + H ⁺)	0.032
211	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 440.2 (M + H ⁺)	0.009





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
212	 5-chloro-N2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 470.2 (M + H ⁺)	0.013
213	 (S)-3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS <i>m/z</i> 540.2 (M + H ⁺)	0.012
214	 N2-(4-(1-(3-(azetidin-1-ylsulfonyl)propyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 573.2 (M + H ⁺)	0.04




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
215	 5-chloro-N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N4-(5-(trifluoromethyl)-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 466.2 (M + H ⁺)	0.047
216	 5-chloro-N2-(2-fluoro-4-(1-methylpiperidin-4-yl)-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 484.2 (M + H ⁺)	0.047
217	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	ESMS m/z 523.2 (M + H ⁺)	0.034
218	 5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)-N2-(2-methyl-4-(1-methylpiperidin-4-yl)-5-(trifluoromethyl)phenyl)pyrimidine-2,4-diamine	ESMS m/z 480.2 (M + H ⁺)	0.026





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
219	 5-chloro-N2-(5-methoxy-2-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 442.2 (M + H ⁺)	0.009
220	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	ESMS m/z 537.2 (M + H ⁺)	0.037
221	 N2-(4-(1-ethylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine	ESMS m/z 474.3 (M + H ⁺)	0.034




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
222	 2-(4-(2,5-dimethyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)ethanol	ESMS m/z 490.2 (M + H ⁺)	0.038
223	 1-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	ESMS m/z 551.2 (M + H ⁺)	0.036
224	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-4-morpholinobutan-1-one	ESMS m/z 567.3 (M + H ⁺)	0.02




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
225	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-morpholinopropan-1-one	ESMS m/z 553.3 (M + H ⁺)	0.017
226	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-(pyrrolidin-1-yl)propan-1-one	ESMS m/z 537.3 (M + H ⁺)	0.061
227	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-4-(pyrrolidin-1-yl)butan-1-one	ESMS m/z 551.3 (M + H ⁺)	0.123




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
228	 (R)-3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 540.2 (M + H ⁺)	0.023
229	 (2R)-3-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol	ESMS m/z 524.2 (M + H ⁺)	0.211
230	 (S)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)oxazolidin-2-one	ESMS m/z 414.1 (M + H ⁺)	0.04
231	 (R)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)oxazolidin-2-one	ESMS m/z 414.1 (M + H ⁺)	0.05




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
232	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 444.2 (M + H ⁺); ¹ HNMR (DMSO-d ₆) δ 8.01 (s, 1H), 7.35 (m, 1H), 7.02 (d, 1H), 6.22 (s, 1H), 2.97 (d, 2H), 2.62 (m, 1H), 2.34 (q, 2H), 2.21 (s, 3H), 2.13 (s, 3H), 1.98 (t, 2H), 1.64 (m, 4H), 1.01 (t, 3H).	0.020
233	 5-chloro-N2-(2,5-dimethyl-4-(1-(3-(pyrrolidin-1-yl)propylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 587.3 (M + H ⁺)	0.054
234	 (R)-1-(3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-ylsulfonyl)propyl)pyrrolidin-3-ol	ESMS m/z 603.3 (M + H ⁺)	0.073





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
235	 5-chloro-N2-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 506.2 (M + H ⁺)	0.063
236	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide	ESMS <i>m/z</i> 483.2 (M + H ⁺)	0.059
237	 N2-(4-(azetidin-3-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 384.2 (M + H ⁺)	0.41
238	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1-(pyrrolidin-1-yl)propan-1-one	ESMS <i>m/z</i> 537.3 (M + H ⁺)	0.039





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
239	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-(dimethylamino)propan-1-one	ESMS m/z 511.3 (M + H ⁺)	0.073
240	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 498.2 (M + H ⁺)	0.102
241	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2-methyl-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 494.2 (M + H ⁺)	0.068





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
242	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1-(pyrrolidin-1-yl)propan-1-one	ESMS m/z 541.3 (M + H ⁺)	0.06
243	 N2-(4-(1-(3-(azetidin-1-ylsulfonyl)propyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 577.2 (M + H ⁺)	0.076
244	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(3-(pyrrolidin-1-ylsulfonyl)propyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 591.2 (M + H ⁺)	0.084




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
245	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(3-(morpholinosulfonyl)propyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 607.2 (M + H ⁺)	0.077
246	 4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1-morpholinobutan-1-one	ESMS m/z 571.3 (M + H ⁺)	0.058
247	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-cyclopropylpropanamide	ESMS m/z 527.2 (M + H ⁺)	0.062




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
248	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide	ESMS m/z 487.2 (M + H ⁺)	0.14
249	 5-chloro-N2-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 502.2 (M + H ⁺)	0.032
250	 5-chloro-N2-(4-(1-cyclopropylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 452.2 (M + H ⁺)	0.025




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
251	 N2-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 540.2 (M + H ⁺)	0.052
252	 5-chloro-N2-(2,5-dimethyl-4-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 496.3 (M + H ⁺)	0.014
253	 5-chloro-N2-(2-fluoro-4-(piperidin-4-yl)-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 470.1 (M + H ⁺)	0.075
254	 5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)-N2-(2-methyl-4-(piperidin-4-yl)-5-(trifluoromethyl)phenyl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 466.2 (M + H ⁺)	0.07




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
255	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(4-(1-cyclopropylpiperidin-4-yl)-2-fluoro-5-methylphenyl)pyrimidine-2,4-diamine	ESMS m/z 482.2 (M + H ⁺)	0.228
256	 5-chloro-N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 412.2 (M + H ⁺)	0.044
257	 N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine	ESMS m/z 446.2 (M + H ⁺)	0.064
258	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(4-(1-cyclopropylpiperidin-4-yl)-2,5-dimethylphenyl)pyrimidine-2,4-diamine	ESMS m/z 478.2 (M + H ⁺)	0.026




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
259	 5-chloro-N2-(2-fluoro-4-(1-((3-isopropyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 540.2 (M + H ⁺)	0.064
260	 5-chloro-N2-(4-(1-((3-isopropyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 536.3 (M + H ⁺)	0.018
261	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-ethyl-5-methylphenyl)piperidin-1-yl)acetamide	ESMS m/z 483.2 (M + H ⁺)	0.027
262	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)acetamide	ESMS m/z 485.2 (M + H ⁺)	0.019




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
263	 2-(4-(4-(5-chloro-4-(5-ethyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide	ESMS m/z 487.2 (M + H ⁺)	0.033
264	 5-Chloro-N2-{4-[1-(1,1-dioxo-1λ6-thietan-3-yl)-piperidin-4-yl]-2-fluoro-5-methyl-phenyl}-N4-(5-ethyl-1H-pyrazol-3-yl)-pyrimidine-2,4-diamine	ESMS m/z 534.2 (M + H ⁺)	0.021
265	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(3-isopropyl-1,2,4-oxadiazol-5-yl)methanone	ESMS m/z 554.2 (M + H ⁺); ¹ H NMR (400 MHz, CDCl ₃) δ 9.67 (br s, 2H), 8.50 (s, 1H), 8.07 (s, 1H), 7.57 (d, J = 8.4 Hz, 1H), 7.05 (d, J = 12.4 Hz, 1H), 6.16 (s, 1H), 5.53 (s, 1H), 3.98 (d, J = 13.2 Hz, 1H), 3.34 (dt, J = 2.0, 2.8 Hz, 1H), 3.15 (m, 1H), 2.82 (m, 2H), 2.62 (t, J = 4.8 Hz, 2H), 2.34 (s, 3H), 2.25 (s, 3H), 1.90 (m, 2H), 1.76 (2 H)	0.091




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
266	 N2-(4-(1-((5-tert-butyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 550.3 (M + H ⁺)	0.025
267	 N2-(4-(1-((5-tert-butyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 554.3 (M + H ⁺)	0.077
268	 5-chloro-N4-(5-ethyl-1H-pyrazol-3-yl)-N2-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)pyrimidine-2,4-diamine	ESMS m/z 458.2 (M + H ⁺)	0.022





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
269	 5-chloro-N2-(4-(1-(2-(3-isopropyl-1,2,4-oxadiazol-5-yl)ethyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 550.3 (M + H ⁺)	0.039
270	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(3-isopropyl-1,2,4-oxadiazol-5-yl)methanone	ESMS m/z 550.2 (M + H ⁺)	0.018
271	 5-chloro-N4-(5-ethyl-1H-pyrazol-3-yl)-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)pyrimidine-2,4-diamine	ESMS m/z 444.2 (M + H ⁺)	0.013




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
272	 5-chloro-N2-(4-(1-cyclobutylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 470.2 (M + H ⁺)	0.050
273	 5-chloro-N2-(2-fluoro-5-methyl-4-(2-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 430.2 (M + H ⁺)	0.018
274	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-2-methylpiperidin-1-yl)acetamide	ESMS m/z 487.2 (M + H ⁺)	0.006




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
275	 5-chloro-N2-(4-(1,2-dimethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 444.2 (M + H ⁺)	0.009
276	 N2-(4-(1,4'-bipiperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 499.2 (M + H ⁺)	0.437
277	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(6-methylpyridazin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 508.2 (M + H ⁺)	0.097




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
278	 5-chloro-N2-(2-fluoro-5-methyl-4-(1'-(methylsulfonyl)-1,4'-bipiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 577.2 (M + H ⁺)	0.009
279	 5-chloro-N2-(2-fluoro-5-methyl-4-((trans)-2-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 430.2 (M + H ⁺)	0.064
280	 5-chloro-N2-(2-fluoro-5-methyl-4-((cis)-2-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 430.2 (M + H ⁺)	0.163




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
281	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 507.2 (M + H ⁺)	0.087
282	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclopent-2-enone	ESMS m/z 496.2 (M + H ⁺); ¹ H NMR (400 MHz, DMSO-d ₆) δ 9.97 (br s, 1H), 9.83 (br s, 1H), 8.90 (s, 1H), 8.32 (d, J = 6.8 Hz, 1H), 7.88 (d, J = 12.8 Hz, 1H), 6.97 (s, 1H), 6.04 (s, 1H), 5.30 (br s, 1H), 4.71 (d, J = 12.0 Hz, 2H), 4.04 (t, J = 12.4 Hz, 2H), 3.84 (m, 1H), 3.53 (m, 2H), 3.13 (m, 2H), 3.08 (s, 3H), 2.96 (s, 3H), 2.59-2.40 (m, 4H)	0.048
283	 N2-(4-(1-(1,4-dioxaspiro[4.5]decan-8-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 556.3 (M + H ⁺)	0.016




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
284	 5-chloro-N2-(4-(1-((3-ethyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 526.2 (M + H ⁺)	0.033
285	 5-chloro-N2-(2-fluoro-4-(1-((5-isopropyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 540.2 (M + H ⁺)	0.068
286	 5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-4-methyl-2-(1-methylpiperidin-4-yl)benzonitrile	ESMS <i>m/z</i> 437.2 (M + H ⁺)	0.037
287	 methyl 5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-4-methyl-2-(1-methylpiperidin-4-yl)benzoate	ESMS <i>m/z</i> 470.2 (M + H ⁺)	0.027




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
288	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-4-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 507.2 ($M + H^+$)	0.093
289	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-1-ethylpiperidine 1-oxide	ESMS m/z 460.2 ($M + H^+$); ^1H NMR (MeOD- d_4) δ 7.91 (s, 1H), 7.69 (d, 1H), 7.00 (d, 1H), 6.15 (s, 1H), 3.28 (m, 4H), 2.91 (m, 1H), 2.32 (m, 2H), 2.20 (s, 3H), 2.16 (s, 3H), 1.83 (s, 2H), 1.64 (d, 2H), 1.31 (t, 3H).	0.37
290	 4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanone	ESMS m/z 512.2 ($M + H^+$)	0.023




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
291	 tert-butyl 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)pyrrolidine-1-carboxylate	ESMS m/z 585.3 (M + H ⁺)	0.067
292	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-((5-methyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 512.2 (M + H ⁺)	0.023
293	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-((3-methyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 512.2 (M + H ⁺)	0.026





	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
294	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-1-(tetrahydro-2H-pyran-4-yl)piperidine 1-oxide	ESMS m/z 516.2 (M + H ⁺)	0.064
295	 5-Chloro-N2-{4-[1-(1,1-dioxo-1λ6-thietan-3-yl)-1-oxide-piperidin-4-yl]-2-fluoro-5-methylphenyl}-N4-(5-ethyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 536.2 (M + H ⁺)	0.039
296	 N2-(4-(1-(azetidin-3-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 471.2 (M + H ⁺)	0.101




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
297	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-3-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 507.2 (M + H ⁺)	0.079
298	 4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanol	ESMS m/z 514.2 (M + H ⁺)	0.026
299	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 485.2 (M + H ⁺)	0.101




	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
300	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-methylsulfonyl)azetidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 549.2 (M + H ⁺)	0.032
301	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-methylsulfonyl)pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 563.2 (M + H ⁺)	0.038
302	 5-chloro-N2-(2,5-dimethyl-4-(1-((3-methyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 508.2 (M + H ⁺)	0.006

	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
303	 5-chloro-N2-(4-(1-((3-ethyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 522.2 (M + H ⁺)	0.006
304	 5-chloro-N2-(2,5-dimethyl-4-(1-((5-methyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 508.2 (M + H ⁺)	0.005
305	 5-chloro-N2-(4-(1-((5-isopropyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS <i>m/z</i> 536.3 (M + H ⁺)	0.025

	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
306	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(oxetan-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 472.2 (M + H ⁺)	0.015
307	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyridin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 503.2 (M + H ⁺)	0.007
308	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyridin-3-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 503.2 (M + H ⁺)	0.006

	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
309	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyridin-4-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 503.2 (M + H ⁺)	0.008
310	 Trans-4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanol	ESMS m/z 514.2 (M + H ⁺)	0.027
311	 Cis-4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanol	ESMS m/z 514.2 (M + H ⁺)	0.021
312	 (R)-5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-(methylsulfonyl)pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 563.2 (M + H ⁺)	0.025

	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC50 (uM)
313	 (S)-5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-(methylsulfonyl)pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 563.2 (M + H ⁺)	0.030
314	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyrimidin-5-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 508.2 (M + H ⁺); ¹ H NMR (400 MHz, CD ₃ OD) δ 9.09 (s, 1H), 8.81 (s, 2H), 8.00 (s, 1H), 7.73 (d, J = 8.4 Hz, 1H), 6.99 (d, J = 13.2 Hz, 1H), 6.23 (br s, 1H), 3.65 (s, 2H), 3.03 (m, 2H), 2.76 (m, 1H), 2.27 (s, 3H), 2.25 (s, 3H), 2.25 (m, 2H), 1.80-1.68 (m, 4H)	0.037
315	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyrazin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 508.2 (M + H ⁺)	0.036

	STRUCTURE	NMR or ESMS	IGF-1R Ba/F3 IC ₅₀ (uM)
316	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridazin-4-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 508.2 (M + H ⁺)	0.027
317	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyrimidin-5-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 504.2 (M + H ⁺)	0.004
318	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyrazin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	ESMS m/z 504.2 (M + H ⁺)	0.010

Assays

[0207] The IC₅₀ of a drug may be determined constructing a dose-response curve and examining the effect of different concentrations of antagonist on reversing agonist activity. IC₅₀ values may be calculated for a given antagonist by determining the concentration needed to inhibit half of the maximum biological response of the agonist. To calculate IC₅₀ values, a series of dose-response data (e.g., drug concentrations x₁, x₂, ..., x_n and growth inhibition y₁, y₂, ..., y_n, the values of y are in the range of 0-1) is generated. IC₅₀ values may be determined by a computer-aided system using the formula:

$$y = D + ((A - D) / (1 + 10^{(x - \log(\text{IC}_{50}))B}))$$

where A is the ratio of growth inhibition between lowest drug concentration and control; B is the slope of sigmoidal curve; and D is the ratio of growth inhibition between highest drug concentration and control.

[0208] The IC₅₀ value is given as that concentration of the test compound that results in growth inhibition that is 50 % lower than that obtained using the control without inhibitor. The compounds of the invention in free form or in pharmaceutically acceptable salt form may exhibit valuable pharmacological properties, for example, as indicated by the *in vitro* tests described in this application. In general, compounds of the invention have IC₅₀ values from 1 nM to 10 μM. In some examples, compounds of the invention have IC₅₀ values from 0.01 μM to 5 μM. In other examples, compounds of the invention have IC₅₀ values from 0.01 μM to 1 μM, or more particularly from 1 nM to 1 μM. In yet other examples, compounds of the invention have IC₅₀ values of less than 1 nM or more than 10 μM. The compounds of the invention may exhibit a percentage inhibition of greater than 50%, or in other embodiments, may exhibit a percentage inhibition greater than about 70%, against IGF-1R at 10 μM.

Ba/F3 cell line panel and reagents

[0209] Ba/F3 is a murine IL-3-dependent pro-B lymphoma cell line. Parental Ba/F3 cells are used to generate a panel of sublines whose proliferation and survival is rendered IL-3-independent by stable transduction with individual tyrosine kinases activated by fusion with the amino-terminal portion of TEL (amino acid 1-375) or BCR. In order to generate Ba/F3 cell lines transformed by Tel-Tyrosine Kinase (TK)

fusions, parental Ba/F3 cells are infected with a retrovirus harboring each TEL-fusion kinase and subjected to puromycin selection and IL-3 withdrawal to obtain IL-3-independent, transformed Ba/F3 cells.

[0210] Each transformed Ba/F3 cells are cultured in RPMI-1640 media (Gibco Cat #11875093, Carlsbad, CA) supplemented with 10% FBS (Hyclone Cat #SV30014.03, Logan, UT), 4.5 g/L glucose (Sigma #G5400, St.Louis, MO), 1.5 g/L sodium bicarbonate (Biowhittaker #17-613E, Walkersville, MD) and Pen/Strep (Gibco #10378-016, Carlsbad, CA). Cells are splitted twice weekly.

Ba/F3 cell viability inhibition assay

[0211] The potency of test compounds against various Tel-TK transformed Ba/F3 lines is determined as follows. Exponentially growing BaF3 Tel-TK cells are diluted in fresh medium to 75,000 cells/mL and seeded into 384-well plates (3750 cells/well) at 50 μ L/well using a μ Fill liquid dispenser (BioTek, Winooski, VT, USA). Duplicate plates are run for each cell line. Test and control compounds are serially diluted with DMSO and arrayed in a polypropylene 384-well plate. 50 nL of compound is transferred into the assay plates using a pin-transfer device, and the plates are incubated at 37 °C (5% CO₂) for 48 hours. 25 μ L Britelite (Perkin Elmer) is added and luminescence is quantified using Analyst GT (Molecular Devices). Custom curve-fitting software is used to produce a logistic fit of percent cell viability as a function of the logarithm of inhibitor concentration. The IC₅₀ is interpolated as the concentration of compound needed to reduce cell viability to 50% of a DMSO control. Parental Ba/F3 cells that are maintained and cultured in presence of IL-3 (1 ng/ml in final) are diluted in fresh medium containing IL-3 (1 ng/ml in final) to 75,000 cells/mL following the same procedure as described above.

Enzymatic HTRF assay

[0212] IGF-1R and INSR (insulin receptor) are purchased from Upstate. Following reagents are prepared in-house; 10 x kinase buffer (KB) (200 mM Tris (pH 7.0), 100 mM MgCl₂, 30 mM MnCl₂, 50 nM NaVO₄), 10 mM ATP, 100 mg/ml BSA, 0.5 M EDTA, 4 M KF. Proxiplate-384 from Perkin-Elmer is used for set up assay. All the HTRF reagents including substrate (Biotin-poly-GT (61GT0BLB), Mab PT66-K, (61T66KLB), Streptavidin-XL^{ent} (611SAXLB)) are purchased from CIS-US, Inc.

[0213] The substrate/ATP mix is prepared by adding ATP (final concentration, 3

μM) and biotinylated poly-GT (final concentration, 10 ng/ μl) into 1x KB, and dispensed into Proxiplate-384 at 5 μl /well using μFill (Bio-TEK). Serially diluted compounds (in DMSO) are transferred into plate using 50 nL pinhead. 5 μL of prepared Enzyme mix (enzyme (final concentration, 5 ng/ μl), mixed with BSA and DTT in 1x KB) is added to initiate kinase reaction using μFill (Bio-TEK). Assay plate is incubated at room temperature for 2 hours. Detection mix is prepared by adding both Mab PT66-K and Streptavidin-XL^{ent} into 0.5 x KB solution containing KF (final concentration, 125 mM), EDTA (final concentration, 50 mM) and BSA (final concentration, 100 $\mu\text{g/ml}$) in. At the end of reaction, 10 μL of detection mix is added and incubated for 30 minutes at room temperature before measurement. HTRF signal is detected using Analyst-GT (molecular Devices).

Cancer cell proliferation inhibition assay

[0214] For luciferizing cancer cell line, each cell line is transduced by ampholytic retrovirus carrying both luciferase gene and puromycin-resistant gene whose expression is driven by LTR. Briefly, the retroviral vector pMSCV-Puro-Luc is transfected into Phoenix cell line using Eugene6 (Roche) according to manufacturer's instruction. Two days after transfection, supernatant containing virus is harvested and filtered with 0.2 μm filter. Harvested virus is used immediately or stored at -80°C . For infection, cultured cancer cells are harvested and plated (5×10^5 cells/well in 1 ml medium) on 6-well tissue culture plate. For each well, 3ml virus supernatant is added together with 400 μl FBS, 40 μl 1 M HEPES (pH8.0) and 4 μl of polybrene (10 $\mu\text{g/ml}$, Specialty media). The plate is centrifuged down for 90 minutes at 2500 rpm for spin-infection and is transferred into an incubator for overnight infection. Next day, infected cell line is transferred into T-75 flask containing fresh medium and incubated for one day. Two days after infection, puromycin is added at the final concentration of 1 $\mu\text{g/ml}$ to begin selection. Within 1-2 weeks, puromycin-resistant cell line is established after at least two subsequent splits and is preserved as luciferized stock.

[0215] Each cell line is harvested while in log phase growth by trypsinization and diluted in respective media to appropriate density prior to plating. Cells are dispensed using μFill (BioTeK) at 50 μl /well into white walled clear bottom plates (Greiner – custom for GNF). Cells are then placed in 37 $^{\circ}\text{C}$ incubator supplying 5% CO_2 overnight. Compounds are transferred using 50nL/well Pintool technology via

Platmate (Matrix). Assay plates are then placed back into the incubator for 3 days. On the third day following compound transfer, BRITELITE® (Perkin Elmer, diluted according to manufacturer's suggestion) is added to assay plates and read on Analyst GT (Molecular Devices) or Envision (Perkin Elmer). Raw data is generated in RLU.

Tumor Activity of Exemplary Compounds

[0216] Three to 4×10^6 human neuroblastoma SK-N-MC cells resuspended in HBSS and mixed in 50% matrigel were injected subcutaneously (0.05 ml/mouse) into female nude (HsdNpa:athymic/nu) mice 10-12 weeks of age. Treatments were initiated when the mean tumor volumes were approximately 150-200 mm³. Body weights and tumor volumes were recorded three times a week. Tumor volumes were measured with calipers and determined according to the formula $\text{length} \times \text{diameter}^2 \times \pi/6$. In addition to presenting fractional changes of tumor volumes over the course of treatments, antitumor activity is expressed as T/C % (mean change of tumor volume of treated animals / mean change of tumor volume of control animals) $\times 100$. Efficacy of test compounds was determined by initiating oral dosing on day 19 post-cell injection following randomization of the mice so that each group has similar mean tumor size. Dosing with an appropriate schedule continued for 7 days based on the general health condition of the animals. All test compounds were formulated in NMP/PEG300 (10:90) and applied daily by gavage. Vehicle consisted of NMP/PEG300 (10:90). All application volumes were 5 ml/kg. The activity of exemplary compounds of the invention on tumor growth is shown in Tables 2-4.

Table 2

Treatment	Tumor response			
	T/C (%)	Regr. (%)	Δ tumor volume (mean mm ³ $\pm$ SEM)	Mean fold change in tumor growth
NMP/PEG300 (10/90) 5 ml/kg po q8/16h	100	0	334 $\pm$ 59	3.1 $\pm$ 0.3
Test Compound 1 20 mg/kg, q8/16h, po	53	-	231 $\pm$ 145	2.1 $\pm$ 0.4
Test Compound 1 40 mg/kg, q8/16h, po	35	-	107 $\pm$ 48*	1.7 $\pm$ 0.2*

Table 3

Treatment	Tumor response			
	T/C (%)	Regr. (%)	Δ tumor volume (mean mm ³ $\pm$ SEM)	Mean fold change in tumor growth
NMP/PEG300 (10/90) 5 ml/kg po q8/16h	100	0	334 $\pm$ 59	3.1 $\pm$ 0.3
Test Compound 2 12.5 mg/kg, q8/16h, po	69	-	232 $\pm$ 103	2.5 $\pm$ 0.4
Test Compound 2 25 mg/kg, q8/16h, po	22	-	42 $\pm$ 10*	1.5 $\pm$ 0.2*

Table 4


Treatment	Tumor response			
	T/C (%)	Regr. (%)	Δ tumor volume (mean mm³ ± SEM)	Mean fold change in tumor growth
NMP/PEG300 (10/90) 5 ml/kg po q8/16h	100	0	334 ± 59	3.1 ± 0.3
Test Compound 3 12.5 mg/kg, q8/16h, po	58	-	220 ± 67	2.2 ± 0.3
Test Compound 3 25 mg/kg, q8/16h, po	26	-	102 ± 40*	1.6 ± 0.3*

* p < 0.05 vs. Vehicle controls – ANOVA on ranks and post hoc Dunnett's test.

[0217] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference for all purposes.

Claims

1. A compound of Formula (1):



or a physiologically acceptable salt thereof;

wherein ring E may optionally contain a double bond;

one of Z^1 , Z^2 and Z^3 is NR^6 , $N(R^6)^+-O^-$ or $S(O)_{1-2}$ and the others are CR_2 ;

R^1 is halo or an optionally halogenated C_{1-6} alkyl;


R^2 is pyridine-2-onyl, azepan-2-onyl or a monocyclic 5-6 membered heteroaryl having 1-3 heteroatoms selected from N, O and S; each of which is optionally substituted with R^9 wherein R^9 is C_{1-6} alkyl, C_{1-6} haloalkyl or C_{3-7} cycloalkyl;

R^3 and R^4 are each H;

R^5 is halo, hydroxyl, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkoxy, cyano or $C(O)O_{0-1}R^8$;

R^6 is H; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $(CR_2)_pNR(CR_2)_pOR^7$, $(CR_2)_pNR-L-C(O)R^8$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $-C(O)-(CR_2)_p-OR^7$, L-Y, $-L-C(O)R^7$, $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)NR(CR_2)_pOR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-C(O)NR(CR_2)_pSR^7$, $-L-C(O)NR(CR_2)_pS(O)_{1-2}R^8$, $-L-S(O)_2R^8$, $-L-S(O)_2-(CR_2)_q-NRR^7$, $-L-S(O)_2NR(CR_2)_pNR(R^7)$ or $-L-S(O)_2NR(CR_2)_pOR^7$;

alternatively, R^6 is a radical selected from formula (a), (b), (c) or (d):



R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} ;

R^{11} , R^{12} , R^{13} , R^{14} , R^{15} and R^{16} are independently selected from H; C_{1-6} alkoxy; C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R^{11} and R^{12} , R^{12} and R^{15} , R^{15} and R^{16} , R^{13} and R^{14} , or R^{13} and R^{15} together with the atoms to which they are attached may form a 3-7 membered saturated, unsaturated or partially unsaturated ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;

L is $(CR_2)_{1-4}$ or a bond;

Y is C_{3-7} carbocyclic ring, C_{6-10} aryl, or a 5-10 membered heteroaryl or 4-10 membered heterocyclic ring, each of which is optionally substituted with 1-3 R^5 groups;

R^7 , R^8 and R^{8a} are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or cyano; $(CR_2)_qY$ or C_{1-6} alkoxy; or R^7 is H;

each R is independently H or C_{1-6} alkyl;

R and R^7 together with N in each NRR^7 may form a 5-6 membered ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;


m is 2-4;

n is 1-3;

p is 1-4; and

q is 0-4.

2. The compound of claim 1, wherein said compound is of Formula (2):




wherein one of R^{5a} , R^{5b} and R^{5c} is H and the others are independently halo, C_{1-6} alkyl, C_{1-6} alkoxy, halo-substituted C_{1-6} alkyl, cyano or $C(O)O_{0-1}R^8$ wherein R^8 is C_{1-6} alkyl; and

R^1 , R^2 , R^3 , R^4 , E, Z^1 , Z^2 and Z^3 are as defined in claim 1.

3. The compound of claim 2, wherein Z^3 is NR^6 or $N(R^6)^+-O^-$; and Z^1 and Z^2 are CH_2 .

4. The compound of claim 3, wherein R^6 is H, C_{1-6} alkyl optionally substituted with halo and/or hydroxyl groups; $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$, $-(CR_2)_p-C(O)OR^7$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, L-Y, $-L-C(O)R^7$, $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-S(O)_2R^8$, $-L-S(O)_2-(CR_2)_q-NRR^7$, or a radical selected from formula (a), (b), (c) or (d):




R^{10} is O, S, NR^{17} wherein R^{17} is H, C_{1-6} alkyl, SO_2R^{8a} or CO_2R^{8a} and R^{8a} is C_{1-6} alkyl; and

R^7 , R^8 , R^{10} , R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R, L, Y, p and q are as defined in claim 1.

5. The compound of any one of claims 1-4, wherein R^2 is pyrazolyl, isoxazolyl, pyridine-2-onyl or azepan-2-onyl, each of which is substituted with C_{1-6} alkyl, C_{1-6} haloalkyl or C_{3-7} cycloalkyl.

6. The compound of claim 1, wherein said compound is of Formula (3):




or a tautomer thereof;

wherein R^{5b} is H; and R^{5a} and R^{5c} are independently halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, halo-substituted C₁₋₆ alkyl, cyano or C(O)O₀₋₁R⁸ wherein R⁸ is C₁₋₆ alkyl;

Z^1 and Z^2 are CH^2 ;

$$\text{Z}^3 \text{ is } \text{NR}^6 \text{ or } \text{N}(\text{R}^6)^+ \text{-O}^-;$$

R^6 is H, C_{1-6} alkyl optionally substituted with halo and/or hydroxyl groups;
 $-(CR_2)_p-OR^7$, $-(CR_2)_p-CH(OH)C_tF_{2t+1}$ wherein t is 1-3, $(CR_2)_p-CN$; $(CR_2)_p-NR(R^7)$,
 $-(CR_2)_p-C(O)OR^7$, $C(O)(CR_2)_qOR^8$, $-C(O)O-(CR_2)_p-NRR^7$, $L-Y$, $-L-C(O)R^7$,
 $-L-C(O)-NRR^7$, $-L-C(O)-NR-(CR_2)_p-NRR^7$, $-L-C(O)-(CR_2)_q-NR-C(O)-R^8$, $-L-$
 $S(O)_2R^8$,
 $-L-S(O)_2-(CR_2)_q-NRR^7$, or a radical selected from formula (a), (b), (c) or (d):

















R¹⁰ is O, S, NR¹⁷ wherein R¹⁷ is H, C₁₋₆ alkyl, SO₂R^{8a} or CO₂R^{8a} and R^{8a} is C₁₋₆ alkyl; and









$R^1, R^3, R^4, R^7, R^8, R^{11}, R^{12}, R^{13}, R^{14}, R^{15}, R^{16}, R, L, Y, p, q$ and E are as defined in claim 1.









7. The compound of claim 6, wherein R^{5a} is halo and R^{5c} is C₁₋₆ alkyl.
8. The compound of claim 6, wherein R⁶ is C₁₋₆ alkyl or a radical of formula (a) or (c); and R¹⁰ is O.







9. The compound of any one of claims 1-8, wherein R¹ is halo.
10. The compound of claim 1, wherein said compound is selected from the group


 5-chloro-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidine-1-carboxylate
 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide
 5-chloro-N2-(2-isopropoxy-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylpropanamide

 N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine	 (R)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide
 N2-(2-isopropoxy-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-5-methyl-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide
 5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (R)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide
 5-chloro-N2-(5-chloro-2-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (S)-2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylpropanamide


 5-chloro-N2-(4-fluoro-2-methyl-5-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclopropyl)methanone
 3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-(ethylamino)cyclopropyl)methanone
 (S)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one	 5-chloro-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)-N4-(5-(trifluoromethyl)-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (R)-3-(5-chloro-2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenylamino)pyrimidin-4-ylamino)azepan-2-one	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylacetamide

 5-chloro-N2-(4-(1-isopropylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide
 5-chloro-N2-(4-fluoro-5-(1-isopropylpiperidin-4-yl)-2-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N2-(2,5-dimethyl-4-(1-(tetrahydro-2H-thiopyran-4-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N2-(2,5-dimethyl-4-(piperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(ethylamino)-2-methylpropan-1-one
 5-chloro-N2-(2,5-dimethyl-4-(piperidin-3-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(ethylamino)cyclobutyl)methanone


 5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(methylamino)cyclobutyl)methanone
 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(2-fluoro-5-methyl-4-(piperidin-4-yl)phenyl)pyrimidine-2,4-diamine	 N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-5-methyl-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)pyrimidine-2,4-diamine	 (S)-3-(4-(2,5-dimethyl-4-(5-methyl-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol




5-Chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




1-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone




5-chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine











2-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide










5-chloro-N2-(2,5-dimethyl-4-(1-(pyridazin-4-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




2-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide

 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanol	 (S)-5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)ethanol	 (R)-5-chloro-N2-(2,5-dimethyl-4-(1-methylpiperidin-2-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N²-(2-fluoro-5-methyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)butanamide
 5-chloro-N2-(2,5-dimethyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 1-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone


 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (1-aminocyclopropyl)(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methanone
 5-chloro-N2-(5-chloro-2-methyl-4-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (1-aminocyclopropyl)(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methanone
 5-chloro-N2-(4-fluoro-2-methyl-5-(1-(3,3,3-trifluoropropyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylbutanamide




3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl) piperidin-1-yl)-1,1,1-trifluoropropan-2-ol




2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-ethylbutanamide




3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino) pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl) piperidin-1-yl)-1,1,1-trifluoropropan-2-ol









5-chloro-N2-(4-(1-((3,5-dimethylisoxazol-4-yl)methyl)piperidin-2-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




3-(4-(5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-fluoro-4-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol




(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-hydroxycyclopropyl)methanone


 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylacetamide	 (2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-hydroxycyclopropyl)methanone
 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	 5-chloro-N2-(2,5-dimethyl-4-(1-(3-morpholinopropyl)sulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide	 N2-(4-(1-(3-(azetidin-1-yl)propyl)sulfonyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide




(3R,4R)-1-(3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-ylsulfonyl)propyl)pyrrolidine-3,4-diol




2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-(trifluoromethyl)phenyl)piperidin-1-yl)acetamide









(2S)-3-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol

















2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide















1-((4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methyl)cyclopropanecarbonitrile









 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetonitrile
 2-(4-(4-(5-Chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanenitrile
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	 (2S)-3-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol


 2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide	 2-(4-(2,5-dimethyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)acetamide
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylacetamide	 2-(4-(2,5-dimethyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-N-methylacetamide
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	 2-(4-(4-(5-chloro-4-(5-cyclopropyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)ethanol

 2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(4-(1-ethylpiperidin-4-yl)-2,5-dimethylphenyl)pyrimidine-2,4-diamine
 2-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-N-methylacetamide	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carbaldehyde
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-N-methylacetamide	 (R)-2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methyl-2-(trifluoromethyl)phenyl)piperidin-1-yl)acetamide	 (S)-2-(3-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide


 2-(4-(5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-fluoro-4-methylphenyl)piperidin-1-yl)propanamide	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide	 5-chloro-N2-(4-(1-cyclopropylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-methylpropanamide	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide	 5-chloro-N2-(4-(1-(2-methoxyethyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylpropanamide	 (S)-3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol
 2-(4-(5-fluoro-2-methyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)propanamide	 N2-(4-(1-(3-(azetidin-1-yl)sulfonyl)propyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanoic acid	 5-chloro-N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N4-(5-(trifluoromethyl)-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N2-(2-fluoro-4-(1-methylpiperidin-4-yl)-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N2-(2,5-dimethyl-4-(1-(2-(methylsulfonyl)ethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)acetamide
 5-chloro-N2-(4-(1-(2-(ethylsulfonyl)ethyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)-N2-(2-methyl-4-(1-methylpiperidin-4-yl)-5-(trifluoromethyl)phenyl)pyrimidine-2,4-diamine




2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N,N-dimethylethanesulfonamide




5-chloro-N2-(5-methoxy-2-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanesulfonamide









2-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-N-methylacetamide




N2-(4-(1-(2-(azetidin-1-ylsulfonyl)ethyl)piperidin-4-yl)-5-chloro-2-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




N2-(4-(1-ethylpiperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine


 N2-(4-(1-(2-(azetidin-1-ylsulfonyl)ethyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 2-(4-(2,5-dimethyl-4-(4-(5-methyl-1H-pyrazol-3-ylamino)-5-(trifluoromethyl)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)ethanol
 5-Chloro-N2-(2-fluoro-5-methyl-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 1-(4-(4-(5-chloro-4-(5-(trifluoromethyl)-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone
 5-chloro-N2-(2,5-dimethyl-4-(1-(methylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-4-morpholinobutan-1-one




5-chloro-N2-(4-(1-(cyclopropylsulfonyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-morpholinopropan-1-one




5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-3-ylsulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine











1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-(pyrrolidin-1-yl)propan-1-one



















Ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate















1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-4-(pyrrolidin-1-yl)butan-1-one


 ethyl 4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidine-1-carboxylate	 (R)-3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol
 isopropyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate	 (2R)-3-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1,1,1-trifluoropropan-2-ol
 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-N,N-dimethylpiperidine-1-carboxamide	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)-N,N-dimethylpiperidine-1-carboxamide	 5-chloro-N2-(2,5-dimethyl-4-(1-(3-(pyrrolidin-1-yl)propyl)sulfonyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)-N-ethylpiperidine-1-carboxamide	 (R)-1-(3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-ylsulfonyl)propyl)pyrrolidin-3-ol
 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidine-1-carboxamide	 5-chloro-N2-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)propanamide
 1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	 N2-(4-(azetidin-3-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-1-(pyrrolidin-1-yl)propan-1-one
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-3-(dimethylamino)propan-1-one
 1-(4-(2,5-dichloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)phenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-3-(dimethylamino)propan-1-one	 5-chloro-N2-(4-(1-ethylpiperidin-4-yl)-2-methyl-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanone	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1-(pyrrolidin-1-yl)propan-1-one
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(methylamino)ethanone	 N2-(4-(1-(3-(azetidin-1-yl)sulfonyl)propyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(diethylamino)ethanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(3-(pyrrolidin-1-yl)sulfonyl)propyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(methylamino)ethanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(3-(morpholinosulfonyl)propyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 1-(2-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(dimethylamino)ethanone	 4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-1-morpholinobutan-1-one
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(diethylamino)ethanone	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-N-cyclopropylpropanamide




1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(methylamino)propan-1-one




3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propanamide




2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)propan-1-one




5-chloro-N2-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-methyl-2-(methylamino)propan-1-one




5-chloro-N2-(4-(1-cyclopropyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-2-methyl-2-(methylamino)propan-1-one




N2-(4-(1-((2,2-difluorocyclopropyl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine




2-amino-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methylpropan-1-one











5-chloro-N2-(2,5-dimethyl-4-(1-(tetrahydro-2H-pyran-4-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine















(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-(methylamino)cyclopropyl)methanone




5-chloro-N2-(2-fluoro-4-(piperidin-4-yl)-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 (1-aminocyclobutyl)(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)methanone	 5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)-N2-(2-methyl-4-(piperidin-4-yl)-5-(trifluoromethyl)phenyl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-(dimethylamino)cyclobutyl)methanone	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(4-(1-cyclopropylpiperidin-4-yl)-2-fluoro-5-methylphenyl)pyrimidine-2,4-diamine
 (1-aminocyclopropyl)(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone	 5-chloro-N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-(dimethylamino)cyclopropyl)methanone	 N2-(2,5-dimethyl-4-(piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-5-(trifluoromethyl)pyrimidine-2,4-diamine

 1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)-2-morpholinoethanone	 5-chloro-N4-(5-cyclopropyl-1H-pyrazol-3-yl)-N2-(4-(1-cyclopropylpiperidin-4-yl)-2,5-dimethylphenyl)pyrimidine-2,4-diamine
 2-(azetidin-1-yl)-1-(4-(2-chloro-4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-methylphenyl)piperidin-1-yl)ethanone	 5-chloro-N2-(2-fluoro-4-(1-((3-isopropyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-morpholinoethanone	 5-chloro-N2-(4-(1-((3-isopropyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 2-(azetidin-1-yl)-1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)ethanone	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-ethyl-5-methylphenyl)piperidin-1-yl)acetamide
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(pyrrolidin-1-yl)ethanone	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2-methoxy-5-methylphenyl)piperidin-1-yl)acetamide
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(1H-pyrrol-1-yl)ethanone	 2-(4-(4-(5-chloro-4-(5-ethyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)acetamide




1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(piperidin-1-yl)ethanone




5-Chloro-N2-{4-[1-(1,1-dioxo-1λ6-thietan-3-yl)-piperidin-4-yl]-2-fluoro-5-methylphenyl}-N4-(5-ethyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-morpholinoethanone









(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(3-isopropyl-1,2,4-oxadiazol-5-yl)methanone















1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(2,6-dimethylpiperidin-1-yl)ethanone

















N2-(4-(1-((5-tert-butyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine







 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-(2,2,6,6-tetramethylpiperidin-1-yl)ethanone	 N2-(4-(1-((5-tert-butyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-(4-methylpiperazin-1-yl)ethanone	 5-chloro-N4-(5-ethyl-1H-pyrazol-3-yl)-N2-(4-(1-ethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)pyrimidine-2,4-diamine
 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methoxyethanone	 5-chloro-N2-(4-(1-(2-(3-isopropyl-1,2,4-oxadiazol-5-yl)ethyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 N-(2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)-2-oxoethyl)acetamide	 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(3-isopropyl-1,2,4-oxadiazol-5-yl)methanone
 2-(dimethylamino)ethyl 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carboxylate	 5-chloro-N4-(5-ethyl-1H-pyrazol-3-yl)-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)pyrimidine-2,4-diamine
 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-N-(2-(dimethylamino)ethyl)piperidine-1-carboxamide	 5-chloro-N2-(4-(1-cyclobutylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)ethanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(2-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(morpholino)methanone	 2-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-2-methylpiperidin-1-yl)acetamide
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(4-methylpiperazin-1-yl)methanone	 5-chloro-N2-(4-(1,2-dimethylpiperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 1-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidine-1-carbonyl)imidazolidin-2-one	 N2-(4-(1,4'-bipiperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)-2-methoxyethanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(6-methylpyridazin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(piperidin-4-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1'-(methylsulfonyl)-1,4'-bipiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 (S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(pyrrolidin-2-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-((trans)-2-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methylazetidin-3-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-((cis)-2-methylpiperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methylpiperidin-4-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carbonyl)azetidin-2-one	 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclopent-2-enone


 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylazetidin-3-yl)methanone	 N2-(4-(1-(1,4-dioxaspiro[4.5]decan-8-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylpyrrolidin-2-yl)methanone	 5-chloro-N2-(4-(1-((3-ethyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylpiperidin-4-yl)methanone	 5-chloro-N2-(2-fluoro-4-(1-((5-isopropyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)((2R,4R)-4-hydroxy-1-methylpyrrolidin-2-yl)methanone




5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-4-methyl-2-(1-methylpiperidin-4-yl)benzonitrile




(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1H-pyrrol-2-yl)methanone




methyl 5-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-4-methyl-2-(1-methylpiperidin-4-yl)benzoate




(S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-isopropylpiperidin-2-yl)methanone




5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-4-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




(S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(4-isopropylmorpholin-3-yl)methanone




4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-1-ethylpiperidine 1-oxide




(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(2,4-dimethyloxazol-5-yl)methanone









4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanone




(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)((2S)-4-hydroxy-1-methylpyrrolidin-2-yl)methanone




tert-butyl 3-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)pyrrolidine-1-carboxylate


 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1,3,5-trimethyl-1H-pyrazol-4-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-((5-methyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methyl-5-(trifluoromethyl)-1H-pyrazol-4-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-((3-methyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 azetidin-3-yl(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone	 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)-1-(tetrahydro-2H-pyran-4-yl)piperidine 1-oxide




(S)-azetidin-2-yl(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)methanone




5-Chloro-N2-{4-[1-(1,1-dioxo-1λ6-thietan-3-yl)-1-oxide-piperidin-4-yl]-2-fluoro-5-methyl-phenyl}-N4-(5-ethyl-1H-pyrazol-3-yl)-pyrimidine-2,4-diamine




(R)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-1-yl)(1-methylpyrrolidin-2-yl)methanone









N2-(4-(1-(azetidin-3-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)-pyrimidine-2,4-diamine




(R)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-methylpyrrolidin-2-yl)methanone




5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridin-3-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)-pyrimidine-2,4-diamine


 (R)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1-isopropylpyrrolidin-2-yl)methanone	 4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanol
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(4-methyloxazol-5-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(3,5-dimethylisoxazol-4-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-(methylsulfonyl)azetidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




(S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(2-methylpyrrolidin-2-yl)methanone




5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-(methylsulfonyl)pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine




(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)((2S,3R)-3-hydroxy-1-methylpyrrolidin-2-yl)methanone









5-chloro-N2-(2,5-dimethyl-4-(1-((3-methyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine



















(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(1H-tetrazol-1-yl)methanone











5-chloro-N2-(4-(1-((3-ethyl-1,2,4-oxadiazol-5-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine


 (4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(2-methyl-5-(trifluoromethyl)oxazol-4-yl)methanone	 5-chloro-N2-(2,5-dimethyl-4-(1-((5-methyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 6-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidine-1-carbonyl)piperidin-2-one	 5-chloro-N2-(4-(1-((5-isopropyl-1,2,4-oxadiazol-3-yl)methyl)piperidin-4-yl)-2,5-dimethylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 (S)-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)(morpholin-3-yl)methanone	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(oxetan-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 5-chloro-N²-(2-methyl-5-chloro-4-(1-(tetrahydro-1,1-dioxido-3-thienyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N²-(2,5-dimethyl-4-(1-(pyridin-2-ylmethyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N²-(2-fluoro-5-methyl-4-(1-(tetrahydro-1,1-dioxido-3-thienyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N²-(2,5-dimethyl-4-(1-(pyridin-3-ylmethyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N²-(2,5-dimethyl-4-(1-(tetrahydro-1,1-dioxido-2<i>H</i>-thiopyran-4-yl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N²-(2,5-dimethyl-4-(1-(pyridin-4-ylmethyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N²-(2-chloro-5-methyl-4-(1-(1,1-dioxido-3-thietanyl)piperidin-4-yl)phenyl)-N⁴-(5-methyl-1<i>H</i>-pyrazol-3-yl)pyrimidine-2,4-diamine	 Trans-4-(4-(4-(5-chloro-4-(5-methyl-1<i>H</i>-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanol

 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1,1-dioxido-3-thietanyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 Cis-4-(4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-5-fluoro-2-methylphenyl)piperidin-1-yl)cyclohexanol
 5-chloro-N2-(4-(1-(5-ethylpyrimidin-2-yl)piperidin-4-yl)-2-fluoro-5-methylphenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (R)-5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-(methylsulfonyl)pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N2-(2-fluoro-5-methyl-4-(1,2,3,6-tetrahydropyridin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (S)-5-chloro-N2-(2-fluoro-5-methyl-4-(1-(1-(methylsulfonyl)pyrrolidin-3-yl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-Chloro-N²-[4-(1,1-dioxo-tetrahydro-1λ⁶-thiophen-3-yl)-2,5-dimethyl-phenyl]-N⁴-(5-methyl-1H-pyrazol-3-yl)-pyrimidine-2,4-diamine	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyrimidin-5-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

 3-(5-chloro-2-(2,5-dimethyl-4-(piperidin-4-yl)phenylamino)pyrimidin-4-ylamino)pyridin-2(1H)-one	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyrazin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N2-(2-fluoro-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(5-methylisoxazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N2-(2-fluoro-5-methyl-4-(1-(pyridazin-4-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 3-(5-chloro-2-(2,5-dimethyl-4-(piperidin-4-yl)phenylamino)pyrimidin-4-ylamino)pyridin-2(1H)-one	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyrimidin-5-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N2-(2-isopropoxy-5-methyl-4-(1-methylpiperidin-4-yl)phenyl)-N4-(1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 5-chloro-N2-(2,5-dimethyl-4-(1-(pyrazin-2-ylmethyl)piperidin-4-yl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine

11. A compound of Formula (4):



or a physiologically acceptable salt thereof;

wherein X is a monocyclic 5-6 membered heterocyclic ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted on a substitutable ring carbon with oxo and at any substitutable ring nitrogen by R⁶;


R¹ is halo;

R³ and R⁴ are each H;

R⁵ is halo, hydroxyl, C₁₋₆ alkyl, C₁₋₆ alkoxy, halo-substituted C₁₋₆ alkyl, halo-substituted C₁₋₆ alkoxy, cyano or C(O)O₀₋₁R⁸;

R⁶ is H; C₁₋₆ alkyl, C₂₋₆ alkenyl or C₂₋₆ alkynyl, each of which may be optionally substituted with halo and/or hydroxyl groups; -(CR₂)_p-OR⁷, -(CR₂)_p-CH(OH)C_tF_{2t+1} wherein t is 1-3, (CR₂)_p-CN; (CR₂)_p-NR(R⁷), -(CR₂)_p-C(O)OR⁷, (CR₂)_pNR(CR₂)_pOR⁷, (CR₂)_pNR-L-C(O)R⁸, C(O)(CR₂)_qOR⁸, -C(O)O-(CR₂)_p-NRR⁷, -C(O)-(CR₂)_p-OR⁷, L-Y, -L-C(O)R⁷, -L-C(O)-NRR⁷, -L-C(O)-NR-(CR₂)_p-NRR⁷, -L-C(O)NR(CR₂)_pOR⁷, -L-C(O)-(CR₂)_q-NR-C(O)-R⁸, -L-C(O)NR(CR₂)_pSR⁷, -L-C(O)NR(CR₂)_pS(O)₁₋₂R⁸, -L-S(O)₂R⁸, -L-S(O)₂-(CR₂)_q-NRR⁷, -L-S(O)₂NR(CR₂)_pNR(R⁷) or -L-S(O)₂NR(CR₂)_pOR⁷;

alternatively, R⁶ is a radical selected from formula (a), (b), (c) or (d):



R¹⁰ is O, S, NR¹⁷ wherein R¹⁷ is H, C₁₋₆ alkyl, SO₂R^{8a} or CO₂R^{8a};

R¹¹, R¹², R¹³, R¹⁴, R¹⁵ and R¹⁶ are independently selected from H; C₁₋₆ alkoxy; C₁₋₆ alkyl, C₂₋₆ alkenyl or C₂₋₆ alkynyl, each of which may be optionally substituted with halo, amino or hydroxyl groups; or R¹¹ and R¹², R¹² and R¹⁵, R¹⁵ and R¹⁶, R¹³

and R^{14} , or R^{13} and R^{15} together with the atoms to which they are attached may form a 3-7 membered saturated, unsaturated or partially unsaturated ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;

L is $(CR_2)_{1-4}$ or a bond;

Y is C_{3-7} carbocyclic ring, C_{6-10} aryl, or a 5-10 membered heteroaryl or 4-10 membered heterocyclic ring, each of which is optionally substituted with 1-3 R^5 groups;

R^7 , R^8 and R^{8a} are independently C_{1-6} alkyl, C_{2-6} alkenyl or C_{2-6} alkynyl, each of which may be optionally substituted with halo, amino, hydroxyl or cyano; $(CR_2)_q Y$ or C_{1-6} alkoxy; or R^7 is H;

each R is independently H or C_{1-6} alkyl;

R and R^7 together with N in each NRR^7 may form a 5-6 membered ring containing 1-3 heteroatoms selected from N, O and S, and optionally substituted with oxo and 1-3 R^5 groups;

m is 2-4;







n is 1-3;



p is 1-4; and

q is 0-4.

12. The compound of claim 11, wherein X is morpholinyl, piperazinyl, piperazin-2-onyl, oxazolidin-2-onyl or piperidin-2-onyl each of which is substituted at any substitutable ring nitrogen by C_{1-6} alkyl; or X is tetrahydro-2H-pyran-2-onyl; and R^1 is chloro.

13. The compound of claim 11, wherein said compound is selected from the group consisting of

 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperidin-2-one	 5-chloro-N4-(5-methyl-1H-pyrazol-3-yl)-N2-(2-methyl-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)phenyl)pyrimidine-2,4-diamine
 4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)tetrahydro-2H-pyran-2-one	 5-chloro-N2-(2-fluoro-4-(4-methylpiperazin-1-yl)-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine
 5-chloro-N2-(2-fluoro-4-morpholino-5-(trifluoromethyl)phenyl)-N4-(5-methyl-1H-pyrazol-3-yl)pyrimidine-2,4-diamine	 (S)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)oxazolidin-2-one

 1-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)piperazin-2-one	 (R)-4-(4-(5-chloro-4-(5-methyl-1H-pyrazol-3-ylamino)pyrimidin-2-ylamino)-2,5-dimethylphenyl)oxazolidin-2-one
--	--

14. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of claims 1-13 and a physiologically acceptable carrier.

15. A method for inhibiting IGF-1R in a cell, comprising contacting the cell with an effective amount of a compound of any one of claims 1-13 or a pharmaceutical composition thereof.

16. A method for treating an IGF-1R-mediated condition in a mammal suffering therefrom, comprising administering to the mammal a therapeutically effective amount of a compound of any one of claims 1-13 or a pharmaceutical composition thereof, and optionally in combination with a second therapeutic agent; wherein said condition is an autoimmune disease, a transplantation disease, an infectious disease or a cell proliferative disorder.

17. The method of claim 16, wherein said condition is a cell proliferative disorder.

18. The method of claim 17, wherein said cell proliferative disorder is multiple myeloma, neuroblastoma, synovial, hepatocellular, Ewing's Sarcoma or a solid tumor selected from a osteosarcoma, melanoma, and tumor of breast, renal, prostate, colorectal, thyroid, ovarian, pancreatic, lung, uterine or gastrointestinal tumor.

19. The method of claim 16, wherein said second therapeutic agent is a

chemotherapeutic agent.

20. The use of a compound of any one of claims 1-13, or a pharmaceutical composition thereof, for the manufacture of a medicament for treating a condition mediated by IGF-1R or anaplastic lymphoma kinase, and optionally in combination with a second therapeutic agent, wherein said condition is an autoimmune disease, a transplantation disease, an infectious disease or a cell proliferative disorder.