



(43) International Publication Date  
19 March 2015 (19.03.2015)

(10) International Publication Number  
**WO 2015/036379 A1**

(51) International Patent Classification:

*A01N 43/80* (2006.01) *A01N 63/04* (2006.01)  
*A01N 63/00* (2006.01) *A01P 3/00* (2006.01)  
*A01N 63/02* (2006.01) *A01P 21/00* (2006.01)

(21) International Application Number:

PCT/EP2014/069132

(22) International Filing Date:

9 September 2014 (09.09.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

13184283.3 13 September 2013 (13.09.2013) EP

(71) Applicant: **BAYER CROPSCIENCE AG** [DE/DE]; Alfred-Nobel-Str. 50, 40789 Monheim (DE).

(72) Inventors: **WACHENDORFF-NEUMANN, Ulrike**; Oberer Markenweg 85, 56566 Neuwied (DE). **HILLEBRAND, Stefan**; Lothringer Str. 22, 41462 Neuss (DE).

(74) Agent: **BIP PATENTS**; c/o Bayer Intellectual Property GmbH, Alfred-Nobel-Str. 10, 40789 Monheim am Rhein (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,

DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: FUNGICIDAL COMPOSITIONS CONTAINING THIAZOLYLISOXAZOLINE FUNGICIDE AND BIOLOGICAL FUNGICIDE

(57) Abstract: The application relates to fungicidal compositions comprising (A) a thiazolylisoxazoline fungicide (I) and at least one biological control agent (B) particularly selected from: (B1) bacteria, (B2) fungi, (B3) protozoas, (B4) viruses, (B5) entomopathogenic nematode, (B6) proteins or secondary metabolites and further to processes for preparing these compositions and to the use thereof as biologically active compounds, for control of phytopathogenic fungi in crop protection, in the protection of materials and as plant growth regulators.



WO 2015/036379 A1

## FUNGICIDAL COMPOSITIONS CONTAINING THIAZOLYLISOXAZOLINE FUNGICIDE AND BIOLOGICAL FUNGICIDE

**Field of invention**

The present invention relates to novel compositions, to a process for preparing these compositions and to the use thereof as biologically active compositions, especially for the control of phytopathogenic fungi in plants, in the protection of materials and as plant growth regulators.

**Background**

It is already known that certain thiazolylisoxazolines can be used as fungicides (see WO 2008/013925, WO 2008/013622, WO 2009/094407, WO 2009/094445, WO 2010/065579, WO 2011/85170, WO 2011/076699, WO 2012/020060, WO 2012/025557, WO 2012/055837, WO 2012/082580, WO 2012/104273, WO 2013/037768, and WO 2013/098229).

Moreover, it is known that these compounds can be mixed with different fungicidal compounds: WO 2009/055514, WO 2012/168188.

However the ecological and economic demands made on modern active ingredients, for example fungicides, are increasing constantly, for example with respect to activity spectrum, toxicity, selectivity, application rate, formation of residues and favourable manufacture. A further problem arising with the use of synthetic fungicides is that the repeated and exclusive application of a fungicide often leads to the selection of resistant microorganisms. Normally, such are also cross-resistant to other active ingredients having the same mode of action. An effective control of the pathogens with said active compounds is then no longer possible. However, active ingredients having new mechanisms of action are difficult to find and expensive to develop.

The risk of resistance development in pathogen populations as well as environmental and human health concerns have fostered interest in identifying alternatives to synthetic fungicides for managing plant diseases. The use of biological control agents (BCAs) is one alternative. However, the effectiveness of most BCAs is not at the same level as for conventional fungicides, especially in case of severe infection pressure. Consequently, known biological control agents, their mutants and metabolites produced by them are, in particular in low application rates, not entirely satisfactory.

Thus, there is a constant need for developing new, alternative plant protection products which in some areas at least help to fulfill the above-mentioned requirements. One way of fulfilling such need can be the development of novel compositions comprising of fungicides and biological control agents which have advantages over the known compositions at least in some areas.


In view of this, it was in particular an object of the present invention to provide compositions which exhibit activity against phytopathogenic fungi in plants, in the protection of materials and as plant growth regulators. Moreover, it was a further particular object of the present invention, to reduce the application rates and broaden the activity spectrum of the fungicides and the biological control agents, and thereby to provide a composition which, prefera-

bly at a reduced total amount of active compounds applied, has improved activity against phytopathogenic fungi. In particular, it was a further object of the present invention to provide a composition which, when applied to a crop, results in a decreased amount of residues in the crop, and nevertheless provides efficient disease control.

## 5 Summary of the invention

It has now surprisingly been found that compositions comprising

(A) at least one thiazolyloxazoline of formula (I)



10 in which

R<sup>1</sup> represents phenyl, which is at least substituted with one methylsulfonyloxy and optionally be additionally substituted by one substituent selected from the group consisting of methyl, methoxy, fluoro or chloro,

or an agrochemically acceptable salt thereof,

15 (B) at least one biological control agent,

and wherein the biological control agent comprises bacteria, fungi, proteins or secondary metabolites, act in a fungicidal fashion.

In some embodiments, such compositions act in a synergistic fashion.

20 The invention also comprises a method for preparing an agricultural composition comprising adding agriculturally suitable components such as suitable extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost protectants, thickeners, adjuvants or the like to the composition according to the invention. Furthermore the invention comprises a method for reducing damage of plants and plant parts or losses in harvested fruits or vegetables caused by phytopathogenic fungi by controlling such phytopathogenic fungi, comprising applying the composition to the plant or the phytopathogenic fungi or the habitat of the plant or the habitat of the phytopathogenic fungi.

25

In view of this, the problem underlying the present invention has been solved by providing novel compositions which exhibit fungicidal and/ or synergistic activity against phytopathogenic fungi in plants, in the protection of materials and as plant growth regulators. Moreover, the novel compositions according to the invention enable reduced application rates and broaden the activity spectrum of the fungicides and the biological control agents. Final-

ly the novel compositions provide improved activity against phytopathogenic fungi and consequently provide efficient disease control for reducing damage of plants and plant parts or losses in harvested fruits or vegetables.

### **Description of the invention**

5 The problem underlying the present invention has been solved by providing novel compositions comprising

(A) at least one compound of the general formula (I) and

(B) at least one biological control agent,

which act in a fungicidal and / or synergistic fashion for efficient disease control comprising reducing damage of plants and plant parts or losses in harvested fruits or vegetables.

10 Preference is given to combinations comprising at least one compound of the formula (I) selected from the group consisting of

(I-1) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}phenyl methanesulfonate,

15 (I-2) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-6-chlorophenyl methanesulfonate,

(I-3) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate,

(I-4) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-5-methylphenyl methanesulfonate,

20 (I-5) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-5-chlorophenyl methanesulfonate,

(I-6) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-4-methylphenyl methanesulfonate,

25 (I-7) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-4-chlorophenyl methanesulfonate,

(I-8) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-6-fluorophenyl methanesulfonate.

In general “pesticidal” means the ability of a substance to increase mortality or inhibit the growth rate of phytopathogenic fungi. The definition also comprises the ability of a substance to increase mortality or inhibit the growth rate of phytopathogenic fungi and / or plant pests. The term is used herein, to describe the property of a substance to exhibit activity against phytopathogenic fungi, insects, mites and / or nematodes.

30

“Fungicides” as well as the terms “fungicidal” and “acting in a fungicidal fashion” refer to the ability of a substance to increase mortality or inhibit growth rates of phytopathogenic fungi. Fungicides can be used in crop protection for control of phytopathogenic fungi. They are characterized by an outstanding efficacy against a broad spectrum of phytopathogenic fungi. As used herein, the term “phytopathogenic fungi” comprises all organisms of the kingdom of fungi which can cause damage of plants and / or damage of plant parts and/or losses in harvested fruits or vegetables. Specific phytopathogenic fungi are described later.

“Insecticides” as well as the term “insecticidal” refers to the ability of a substance to increase mortality or inhibit growth rate of insects. As used herein, the term “insects” comprises all organisms in the class “Insecta”.

“Nematicide” and “nematicidal” refers to the ability of a substance to increase mortality or inhibit the growth rate of nematodes. In general, the term “nematode” comprises eggs, larvae, juvenile and mature forms of said organism.

“Acaricide” and “acaricidal” refers to the ability of a substance to increase mortality or inhibit growth rate of ectoparasites belonging to the class Arachnida, sub-class Acari.

#### **Biological control agents**

As used herein, the term “biological control” is defined as control of harmful organisms such as a phytopathogenic fungi and/or insects and/or acarids and/or nematodes by the use or employment of a biological control agent.

As used herein, the term “biological control agent” is defined as a second organism and / or proteins or secondary metabolites produced by such a second organism for the purpose of biological control. Mutants of the second organism shall be included within the definition of the biological control agent. The term “mutant” refers to a variant of the parental strain as well as methods for obtaining a mutant or variant in which the pesticidal activity is greater than that expressed by the parental strain. The “parent strain” is defined herein as the original strain before mutagenesis. To obtain such mutants the parental strain may be treated with a chemical such as N-methyl-N'-nitro-N-nitrosoguanidine, ethylmethanesulfone, or by irradiation using gamma, x-ray, or UV-irradiation, or by other means well known to those skilled in the art. Known mechanisms of biological control agents comprise enteric bacteria that control root rot by out-competing fungi for space on the surface of the root. Bacterial toxins, such as antibiotics, have been used to control pathogens. The toxin can be isolated and applied directly to the plant or the bacterial species may be administered so it produces the toxin *in situ*.

A “variant” is a strain having all the identifying characteristics of the NRRL or ATCC Accession Numbers as indicated in this text and can be identified as having a genome that hybridizes under conditions of high stringency to the genome of the NRRL or ATCC Accession Numbers.

“Hybridization” refers to a reaction in which one or more polynucleotides react to form a complex that is stabilized via hydrogen bonding between the bases of the nucleotide residues. The hydrogen bonding may occur by Watson-Crick base pairing, Hoogsteen binding, or in any other sequence-specific manner. The complex may comprise two strands forming a duplex structure, three or more strands forming a multi-stranded complex, a

single self-hybridizing strand, or any combination of these. Hybridization reactions can be performed under conditions of different “stringency”. In general, a low stringency hybridization reaction is carried out at about 40 °C in 10 X SSC or a solution of equivalent ionic strength/temperature. A moderate stringency hybridization is typically performed at about 50 °C in 6 X SSC, and a high stringency hybridization reaction is generally performed at about 60 °C in 1 X SSC.

A variant of the indicated NRRL or ATCC Accession Number may also be defined as a strain having a genomic sequence that is greater than 85%, more preferably greater than 90% or more preferably greater than 95% sequence identity to the genome of the indicated NRRL or ATCC Accession Number. A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) has a certain percentage (for example, 80%, 85%, 90%, or 95%) of “sequence identity” to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example, those described in Current Protocols in Molecular Biology (F. M. Ausubel et al., eds., 1987).

NRRL is the abbreviation for the Agricultural Research Service Culture Collection, an international depositary authority for the purposes of depositing microorganism strains under the Budapest treaty on the international recognition of the deposit of microorganisms for the purposes of patent procedure, having the address National Center for Agricultural Utilization Research, Agricultural Research service, U.S. Department of Agriculture, 1815 North university Street, Peroira, Illinois 61604 USA.

ATCC is the abbreviation for the American Type Culture Collection, an international depositary authority for the purposes of depositing microorganism strains under the Budapest treaty on the international recognition of the deposit of microorganisms for the purposes of patent procedure, having the address ATCC Patent Depository, 10801 University Blvd., Manassas, VA 10110 USA.

According to the present invention the biological control agents (B) are particularly selected from the group consisting of:

B1. bacteria,

B2. fungi,

B3. protozoas,

B4. viruses,

B5. entomopathogenic nematode,

B6. proteins or secondary metabolites.

According to the invention the term “bacteria” include spore-forming, root-colonizing bacteria, or bacteria and their metabolites useful as biological fungicides, biological insecticides, biological nematocides or biological miticides, or soil amendments improving plant health and growth. Examples of such bacteria to be used or employed according to the invention are (the numbering is used throughout the complete following description of the invention): B1.1 *Bacillus subtilis*, in particular strain QST713/AQ713 (e.g. SERENADE MAX from Bayer CropScience

LP, US, having NRRL Accession No. B21661 and described in U.S. Patent No. 6,060,051); B1.2 *Bacillus subtilis* strain AQ153 (having NRRL Accession No. 55614 and described in U.S. Patent No. 5,753,222); B1.3 *Bacillus sp.* strain AQ175 (having ATCC Accession No. 55608 and described in U.S. Patent No. 5,869,042); B1.4 *Bacillus sp.* strain AQ177 (having ATCC Accession No. 55609 and described in U.S. Patent No. 5,869,042); B1.5 *Bacillus sp.* strain AQ178 (having ATCC Accession No. 53522 and described in U.S. Patent No. 5,869,042); B1.6 *Bacillus pumilus*, in particular strain QST2808 (e.g. Sonata® from Bayer CropScience LP, US, having Accession No. NRRL B-30087 and described in U.S. Patent No. 6,245,551); B1.7 *Bacillus pumilus*, in particular strain GB34 (e.g. Yield Shield® from Bayer CropScience AG, DE); B1.8 *Bacillus thuringiensis* strain AQ52 (having Accession No. NRRL B-21619 and described in U.S. Patent No. 5,919,447); B1.9 *Streptomyces sp.* strains, in particular the strain having Accession No. NRRL B-30145 and described in U.S. Patent No. 6,524,577; B1.10 *Streptomyces galbus* (= *Streptomyces griseoviridis*), in particular *Streptomyces galbus* strain QST 6047 and mutants thereof (having Accession No. NRRL B-30232 and described in US Patent Nos. 6,682,925); B1.11 *Bacillus chitinosporus*, in particular strain AQ746 (having Accession No. NRRL B-21618 and described in U.S. Patent No. 5,733,544); B1.12 *Bacillus mycoides*, strain AQ726 (having Accession No. NRRL B-21664 and described in U.S. Patent Nos. 5,906,818 and 6,210,665); B1.13 *Bacillus pumilus*, in particular strain AQ717 (having Accession No. NRRL B21662 and described in U.S. Patent No. 6,001,637); B1.14 *Bacillus subtilis*, in particular strain AQ743 (having Accession No. NRRL B-21665 and described in U.S. Patent No. 6,015,553); B1.15 *Rhodococcus globerulus* strain AQ719 (having Accession No. NRRL B21663); B1.16 *Bacillus thuringiensis subsp. aizawai*, in particular strain ABTS-1857 (SD-1372; e.g. XenTari® from Bayer Crop Science AG, DE); B1.17 *Bacillus firmus*, in particular, strain CNMC I-1582 (e.g. VOTIVO from Bayer CropScience); B1.18 *Bacillus subtilis*, in particular strain AQ30002, (having Accession No. NRRL B-50421 and described in U.S. Patent Application No. 13/330,576); B1.19 *Bacillus subtilis*, in particular strain AQ30004 (having Accession No. NRRL B-50455 and described in U.S. Patent Application No. 13/330,576); B1.20 *Bacillus amyloliquefaciens*, strain D747 (e.g. Bacstar® from Etec Crop Solutions, NZ and also e.g. Double Nickel™ from Certis, US); B1.21 *Bacillus pumilus*, in particular strain BU F-33 (e.g. Integral F-33 from Becker Underwood, US); B1.22 *B. subtilis var. amyloliquefaciens* strain FZB24 (e.g. Taegro® from Novozymes, US); B1.23 *Paenibacillus polymyxa*, in particular strain AC-1 (e.g. Topseed from Green Biotech Company Ltd.); B1.24 *Pseudomonas proradix* (e.g. Proradix® from Sourcon Padena); B1.25 *Bacillus amyloliquefaciens* strain MBI 600 (e.g. Subtilex from Becker Underwood, US); B1.26 *Bacillus amyloliquefaciens* strain GB03 (e.g. Kodiak® from Bayer Crop Science AG, DE); B1.27 *Bacillus amyloliquefaciens* strain DB 101 (e.g. Shelter from Dagut Bio lab, ZA); B1.28 *Bacillus amyloliquefaciens* strain DB 102 (e.g. Artemis from Dagut Bio lab, ZA); B1.29 *Bacillus amyloliquefaciens* isolate B246 (e.g. Avogreen from University of Pretoria); B1.30 *Bacillus licheniformis*, in particular strain SB3086 (e.g. EcoGuard™ Biofungicide and Green Releaf from Novozymes); B1.31 *Pseudomonas syringae*, in particular strain MA-4 (e.g. Biosave from EcoScience, US); B1.32 *Pseudomonas syringae* strain 742RS (e.g. Frostban C from Frost Technology Corp); B1.33 *Pseudomonas fluorescens*, in particular strain 1629RS (e.g. Frostban D from Frost Technology Corp); B1.34 *Streptomyces galbus* (*Streptomyces griseoviridis*), in particular strain K61 (Accession No. DSM 7206; e.g. Mycostop® from Verdera, cf. Crop Protection 2006, 25, 468-475); B1.35 *Streptomyces lydicus*, in particular strain WYEC108 (e.g. Actinovate from Natural Industries, US); B1.36 *Agrobacterium radiobacter*, in particular strain K84 (e.g. Galltrol-A from AgBioChem, CA); B1.37 *Agrobacterium radiobacter* strain K1026 (e.g. Nogall from Becker Underwood,

US); B1.38 *Bacillus lentimorbus*; B1.39 *Bacillus mycoides*, isolate J. (e.g. BmJ from Certis USA); B1.40 *Bacillus sphaericus*, in particular Serotype H5a5b strain 2362 (e.g. VectoLex® from Valent BioSciences, US); B1.41 *Bacillus thuringiensis subsp. kurstaki* strain BMP 123 from Becker Microbial Products, IL; B1.42 *Bacillus thuringiensis subsp. aizawai*, in particular serotype H-7 (e.g. Florbac WG from Valent BioSciences, US); B1.43 *Bacillus thuringiensis subsp. kurstaki* strain HD-1 (e.g. Dipel® ES from Valent BioSciences, US); B1.44 *Bacillus thuringiensis subsp. tenebrionis* strain NB 176 (SD-5428; e.g. Novodor® FC from BioFa DE); B1.45 *Bacillus thuringiensis var. japonensis* strain Buibui; B1.46 *Burkholderia spp.*, in particular strain A396 (Accession No. NRRL B-50319; e.g. MBI-206 TGAI from Marrone Bio Innovations); B1.47 *Chromobacterium subtsugae*, in particular strain PRAA4-1T (MBI-203; e.g. Grandevo from Marrone Bio Innovations); B1.48 *Paenibacillus popilliae* (formerly *Bacillus popilliae*; e.g. Milky spore disease from St. Gabriel Laboratories); B1.49 *Xenorhabdus luminescens*; B1.50 *Xenorhabdus nematophila*; B1.51 *Bacillus thuringiensis*, in particular *Bacillus thuringiensis* subspecies *israelensis* (serotype H-14); B1.52 *Bacillus amyloliquefaciens*, in particular strain FZB42 (e.g. RhizoVital® from ABiTEP, DE); B1.53 *Bacillus cereus*; B1.54 spores of *Bacillus cereus* strain CNCM I-1562 (cf. US 6,406,690); B1.55 *Bacillus laterosporus* (also known as *Brevibacillus laterosporus*; e.g. Bio-Tode from Agro-Organics, SA); B1.56 *Bacillus megaterium*, strain YFM3.25 (e.g. Bioarc from BioArc); B1.57 *Bacillus mojavenensis*, strain SR11 (CECT-7666) by Probelte, Sa; B1.58 *Bacillus nematocida*; B1.59 *Pasteuria nishizawae* (e.g. oyacyst LF/ST from Pasteuria Bioscience); B1.60 *Pasteuria penetrans* (formerly *Bacillus penetrans*, e.g. Pasteuria wettable powder from Pasteuria Bioscience); B1.61 *Pasteuria usgae* (e.g. Econem™ from Pasteuria Bioscience); B1.62 compositions comprising nematicidal *Streptomyces sp.*, such as *Streptomyces lydicus* (commercially e.g. ACTINOVATE®); B1.63 compositions comprising nematicidal *Streptomyces sp.*, such as *Streptomyces saraceticus* (e.g. Clanda from A & A Group (Agro Chemical Corp.); B1.64 *Bacillus amyloliquefaciens*, in particular strain IN937a; B1.65 *Bacillus cereus*, in particular strain BP01 (ATCC 55675, e.g. Mepichlor from Arysta, US and also e.g. Mepplus from Micro-Flo Company LLC, US); B1.66 *Bradyrhizobium japonicum* (e.g. Optimize from Novozymes); B1.67 *Delftia acidovorans*, in particular strain RAY209 (e.g. BioBoost® from Brett Young Seeds); B1.68 *Lactobacillus sp.* (e.g. Lactoplant from LactoPAFI); B1.69 *Pseudomonas aeruginosa*, in particular strain PN1; B1.70 *Rhizobium leguminosarum*, in particular *bv. viceae* strain Z25 (Accession No. CECT 4585); B1.71 *Streptomyces acidiscabies*, in particular strain RL-110T (e.g. MBI-005EP from Marrone Bioinnovations, CA); B1.72 *Bacillus azotoformans*; B1.73 *Bacillus smithii*; B1.74 *Bacillus subtilis*, in particular strain DB 101; B1.75 *Lysobacter antibioticus*, in particular strain 13-1 (cf. Biological Control 2008, 45, 288-296); B1.76 *Pantoea agglomerans*, in particular strain E325 (Accession No. NRRL B-21856); B1.77 *Bacillus coagulans*, in particular strain TQ33; B1.78 *Bacillus popilliae* (e.g. Cronox by Bio-Crop, CO); B1.79 *Bacillus cepacia* (e.g. Deny Stine by Microbial Products); B1.80 *Lactobacillus acidophilus* (e.g. Fruitsan by Inagrosa-Industrias Agrobiológicas, S.A); B1.81 *Lysobacter enzymogenes*, in particular strain C3 (cf. J Nematol. 2006 June; 38(2): 233-239); B1.82 *Pseudomonas aeruginosa*, in particular strain WS-1; B1.83 *Pseudomonas aureofaciens*, in particular strain TX-1 (e.g. Spot-Less Biofungicide by Eco Soils Systems, CA); B1.84 *Pseudomonas cepacia* (formerly known as *Burkholderia cepacia*), in particular type Wisconsin, strain M54; B1.85 *Pseudomonas cepacia* (formerly known as *Burkholderia cepacia*), in particular type Wisconsin, strain J82; B1.86 *Pseudomonas chlororaphis*, in particular strain 63-28 (e.g. ATEze by EcoSoil Systems); B1.87 *Pseudomonas fluorescens*, in particular strain A506 (e.g. Blightban by NuFarm and also e.g. Frostban B by Frost Technology Corp); B1.88 *Pseudomonas resinovorans* (e.g. Solanacure



by Agricultural Research Council, SA); B1.89 *Streptomyces* sp., in particular strain WYE 324 (KCTC0342BP); B1.90 *Streptomyces candidus*, in particular strain Y21007-2 (e.g. BioBac by Biontech, TW); B1.91 *Brevibacillus brevis* (formerly *Bacillus brevis*; e.g. Brevisin); B1.92 *Pectobacterium carotovorum* (formerly *Erwinia carotovora*; e.g. BioKeeper by Nissan, JP); B1.93 *Pseudomonas chlororaphis*, in particular strain MA 342 (e.g. Cedomon by Bioagri, S); B1.94 *Bacillus thuringiensis* subsp. *kurstaki* strain ABTS 351; B1.95 *Bacillus thuringiensis* subsp. *kurstaki* strain PB 54; B1.96 *Bacillus thuringiensis* subsp. *kurstaki* strain SA 11; B1.97 *Bacillus thuringiensis* subsp. *kurstaki* strain SA 12; B1.98 *Bacillus thuringiensis* subsp. *kurstaki* strain EG 2348; B1.99 *Bacillus thuringiensis* var. *Colmeri* (e.g. TianBaoBTc by Changzhou Jianghai Chemical Factory); B1.100 *Bacillus thuringiensis* subsp. *aizawai* strain GC-91; B1.101 *Rhizobium meliloti*; B1.102 *Serratia entomophila* (e.g. Invade® by Wrightson Seeds); B1.103 *Serratia marcescens*, in particular strain SRM (Accession No. MTCC 8708); B1.104 *Streptomyces prasinus* (cf. "Prasinons A and B: potent insecticides from *Streptomyces prasinus*" Applied microbiology 1973 Nov); B1.105 *Bacillus kurstaki*; B1.106 *Bacillus aizawai*; B1.107 *Bacillus albolactis*; B1.108 *Bacillus thuringiensis* strain CR-371 (Accession No. ATCC 55273); B1.109 *Pasteuria thornei*; B1.110 *Azorhizobium caulinodans*, preferably strain ZB-SK-5; B1.111 *Azospirillum amazonense*; B1.112 *Azospirillum brasilense*; B1.113 *Azospirillum halopraeferense*; B1.114 *Azospirillum irakense*; B1.115 *Azospirillum lipoferum*; B1.116 *Azotobacter chroococcum*, preferably strain H 23; B1.117 *Azotobacter vinelandii*, preferably strain ATCC 12837; B1.118 *Bacillus lacticola* by Micro-Flo Company; B1.119 *Bacillus lactimorbis* by Micro-Flo Company; B1.120 *Bacillus lactis* from Micro Flo Company; B1.121 *Bacillus maroccanus* from Micro Flo Company; B1.122 *Bacillus metiens* from Micro Flo Company; B1.123 *Bacillus nigrificans* from Micro Flo Company; B1.124 *Bacillus siamensis*, in particular strain KCTC 13613T; B1.125 *Bacillus tequilensis*, in particular strain NII-0943; B1.126 *Gluconacetobacter diazotrophicus*; B1.127 *Rhizobium fredii*; B1.128 *Thiobacillus* sp. (e.g. Cropaid from Cropaid Ltd UK); B1.129 *Xanthomonas campestris* (herbicidal activity), in particular pv poae (e.g. Camperico); B1.130 *Agrobacterium vitis*, in particular the non-pathogenic strain VAR03-1; B1.131 *Bacillus acidocaldarius*; B1.132 *Bacillus acidoterrestris*; B1.133 *Bacillus alcalophilus*; B1.134 *Bacillus alvei*; B1.135 *Bacillus aminoglucosidicus*; B1.136 *Bacillus aminovorans*; B1.137 *Bacillus amylolyticus*; B1.138 *Bacillus amyloliquefaciens*, strain B3; B1.139 *Bacillus aneurinolyticus*; B1.140 *Bacillus atrophaeus*; B1.141 *Bacillus badius*; B1.142 *Bacillus circulans*; B1.143 *Bacillus fastidiosus*; B1.144 *Bacillus lautus*; B1.145 *Bacillus lentus*; B1.146 *Bacillus medusa*; B1.147 *Bacillus psychrosaccharolyticus*; B1.148 *Bacillus subtilis* subspecies *natto* (formerly *Bacillus natto*); B1.149 *Bacillus thuringiensis*, in particular strain AM65-52 (e.g. VectoBac® by Valent BioSciences, US); B1.150 *Bacillus thuringiensis israelensis* strain BMP 144 (e.g. Aquabac by Becker Microbial Products IL); B1.151 *Bacillus thuringiensis* subspecies. *Aegypti* (e.g. Agerin); B1.152 *Bacillus thuringiensis* var. *darmstadiensis* strain 24-91 (e.g. Baciturin); B1.153 *Bacillus thuringiensis* var. *dendrolimus* (e.g. Dendrobacillin); B1.154 *Bacillus thuringiensis* var. *galleriae*; B1.155 *Bacillus thuringiensis* subsp. *Morrisoni*; B1.156 *Bacillus thuringiensis* var. *san diego* (e.g. M-One® by Mycogen Corporation, US); B1.157 *Bacillus thuringiensis* subsp. *thuringiensis* serotype 1, strain MPPL002; B1.158 *Bacillus thuringiensis* var. *thuringiensis*; B1.159 *Bacillus thuringiensis* var. 7216 (e.g. Amactic and Pethian); B1.160 *Bacillus thuringiensis* var. *T36* (e.g. Cahat); B1.161 *Bacillus uniflagellatus*; B1.162 *Brevibacillus brevis* (formerly *Bacillus brevis*), in particular strain SS86-3; B1.163 *Brevibacillus brevis* (formerly *Bacillus brevis*), in particular strain SS86-4; B1.164 *Brevibacillus brevis* (formerly *Bacillus brevis*), in particular strain SS86-5; B1.165 *Brevibacillus brevis* (formerly *Bacillus brevis*), in particular strain 2904; B1.166 *Brevibacillus laterosporus* (formerly *Ba-*

*cillus laterosporus*); B1.167 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain ATCC 64; B1.168 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain NRS 1111; B1.169 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain NRS 1645; B1.170 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain NRS 1647; B1.171 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain BPM3; B1.172 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain G4; B1.173 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain NCIMB; B1.174 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), in particular strain 41419; B1.175 *Corynebacterium paurometabolum*; B1.176 *Herbaspirillum rubrisubalbicans*; B1.177 *Herbaspirillum seropedicae*; B1.178 *Paenibacillus alvei*, in particular strain III3DT-1A; B1.179 *Paenibacillus alvei*, in particular strain III2E; B1.180 *Paenibacillus alvei*, in particular strain 46C3; B1.181 *Paenibacillus alvei*, in particular strain 2771; B1.182 *Paenibacillus macerans*; B1.183 *Pasteuria ramose*; B1.184 *Pasteuria reniformis*; B1.185 *Pseudomonas putida*; B1.186 *Rhizobium loti*; B1.187 *Rhizobium trifolii*; B1.188 *Rhizobium tropici*; B1.189 *Serratia marcescens*, in particular strain R35; B1.190 *Streptomyces colombiensis*; B1.191 *Streptomyces goshikiensis*; B1.192 *Streptomyces lavendulae*; B1.193 *Streptomyces lydicus*, in particular strain WYCD108US; B1.194 *Streptomyces rimosus*; B1.195 *Streptomyces venezuelae*; B1.196 *Virgibacillus pantothenicus* (formerly *Bacillus pantothenicus*), in particular strain ATCC 14576 / DSM 491; B1.197 *Bacillus thuringiensis* strain BD#32 (Accession No. NRRL B-21530 and described in US 5,645,831); B1.198 *Streptomyces* sp. strain WYE 20 (KCTC 0341BP) and B1.199 *Bacillus agri*.

In some embodiments of the invention such biological control agents which are summarized under the term bacteria according to the invention include: B1.1.1 *Bacillus subtilis*, strain QST713/AQ713 (Accession No. NRRL B21661); B1.1.6 *Bacillus pumilus*, strain QST2808 (Accession No. NRRL B-30087); B1.1.7 *Bacillus pumilus*, strain GB34; B1.1.9 *Streptomyces* sp., strain having Accession No. NRRL B-30145; B1.1.10 *Streptomyces galbus* (= *Streptomyces griseoviridis*), strain QST 6047 (Accession No. NRRL B-30232); B1.1.11 *Bacillus chitinosporus*, strain AQ746 (Accession No. NRRL B-21618); B1.1.13 *Bacillus pumilus*, strain AQ717 (having Accession No. NRRL B21662); B1.1.14 *Bacillus subtilis*, strain AQ743 (having Accession No. NRRL B-21665); B1.1.16 *Bacillus thuringiensis* subsp. *aizawai*, strain ABTS-1857 (SD-1372); B1.1.17 *Bacillus firmus*, strain CNMC I-1582; B1.1.18 *Bacillus subtilis*, strain AQ30002, (Accession No. NRRL B-50421); B1.1.19 *Bacillus subtilis*, strain AQ30004 (Accession No. NRRL B-50455); B1.1.21 *Bacillus pumilus*, strain BU F-33; B1.1.23 *Paenibacillus polymyxa*, strain AC-1; B1.1.30 *Bacillus licheniformis*, strain SB3086; B1.1.31 *Pseudomonas syringae*, strain MA-4; B1.1.33 *Pseudomonas fluorescens*, strain 1629RS; B1.1.34 *Streptomyces galbus* (*Streptomyces griseoviridis*), strain K61 (Accession No. DSM 7206); B1.1.35 *Streptomyces lydicus*, strain WYEC108; B1.1.36 *Agrobacterium radiobacter*, strain K84; B1.1.40 *Bacillus sphaericus*, Serotype H5a5b strain 2362; B1.1.42 *Bacillus thuringiensis* subsp. *aizawai*, serotype H-7; B1.1.46 *Burkholderia* spp., strain A396 (Accession No. NRRL B-50319); B1.1.47 *Chromobacterium subtsugae*, strain PRAA4-1T (MBI-203); B1.1.51 *Bacillus thuringiensis* subspecies *israelensis* (serotype H-14); B1.1.52 *Bacillus amyloliquefaciens*, strain FZB42; B1.1.64 *Bacillus amyloliquefaciens*, strain IN937a; B1.1.65 *Bacillus cereus*, strain BP01 (ATCC Accession No. 55675); B1.1.67 *Delftia acidovorans*, strain RAY209; B1.1.69 *Pseudomonas aeruginosa*, strain PN1; B1.1.70 *Rhizobium leguminosarum*; Bv. *viceae* strain Z25 (Accession No. CECT 4585); B1.1.71 *Streptomyces acidiscabies*, strain RL-110T; B1.1.74 *Bacillus subtilis*, strain DB 101; B1.1.75 *Lysobacter antibioticus*,

- strain 13-1; B1.1.76 *Pantoea agglomerans*, strain E325 (Accession No. NRRL B-21856); B1.1.77 *Bacillus coagulans*, strain TQ33; B1.1.81 *Lysobacter enzymogenes*, strain C3; B1.1.82 *Pseudomonas aeruginosa*, strain WS-1; B1.1.83 *Pseudomonas aureofaciens*, strain TX-1; B1.1.84 *Pseudomonas cepacia* (formerly known as *Burkholderia cepacia*); type Wisconsin, strain M54; B1.1.85 *Pseudomonas cepacia* (formerly known as *Burkholderia cepacia*); type Wisconsin, strain J82; B1.1.86 *Pseudomonas chlororaphis*, strain 63-28; B1.1.87 *Pseudomonas fluorescens*, strain A506; B1.1.89 *Streptomyces* sp., strain WYE 324 (KCTC0342BP); B1.1.90 *Streptomyces candidus*, strain Y21007-2; B1.1.93 *Pseudomonas chlororaphis*, strain MA 342; B1.1.103 *Serratia marcescens*, strain SRM; B1.1.124 *Bacillus siamensis*, strain KCTC 13613T; B1.1.125 *Bacillus tequilensis*, strain NII-0943; B1.1.129 *Xanthomonas campestris* (herbicidal activity); pv poae; B1.1.130 *Agrobacterium vitis*, the non-pathogenic strain VAR03-1; B1.1.149 *Bacillus thuringiensis*, strain AM65-52; B1.1.162 *Brevibacillus brevis* (formerly *Bacillus brevis*), strain SS86-3; B1.1.163 *Brevibacillus brevis* (formerly *Bacillus brevis*), strain SS86-4; B1.1.164 *Brevibacillus brevis* (formerly *Bacillus brevis*), strain SS86-5; B1.1.165 *Brevibacillus brevis* (formerly *Bacillus brevis*), strain 2904; B1.1.167 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain ATCC 64; B1.1.168 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain NRS 1111; B1.1.169 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain NRS 1645; B1.1.170 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain NRS 1647; B1.1.171 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain BPM3; B1.1.172 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain G4; B1.1.173 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain NCIMB; B1.1.174 *Brevibacillus laterosporus* (formerly *Bacillus laterosporus*), strain 41419; B1.1.178 *Paenibacillus alvei*, strain III3DT-1A; B1.1.179 *Paenibacillus alvei*, strain III2E; B1.1.180 *Paenibacillus alvei*, strain 46C3; B1.1.181 *Paenibacillus alvei*, strain 2771; B1.1.188 *Rhizobium tropici*; B1.1.189 *Serratia marcescens*, strain R35; B1.1.193 *Streptomyces lydicus*, strain WYCD108US and B1.1.196 *Virgibacillus pantothenicus* (formerly *Bacillus pantothenicus*), strain ATCC 14576 / DSM 491.
- 25 According to the invention the term “fungi” comprises all organisms belonging to the kingdom of fungi, including yeast, useful as biological fungicides, biological insecticides, biological nematocides or biological miticides, or soil amendments improving plant health and growth. Examples of such fungi to be used or employed according to the invention are (the numbering is used throughout the complete following description of the invention): B2.1 *Coniothyrium minitans*, in particular strain CON/M/91-8 (Accession No. DSM-9660; e.g. Contans® from Encore Technologies, LLC); B2.2 *Metschnikowia fructicola*, in particular strain NRRL Y-30752 (e.g. Shemer® from Bayer CropScience); B2.3 *Microsphaeropsis ochracea* (e.g. Microx® from Prophyta); B2.4 *Muscodor albus*, in particular strain QST 20799 (Accession No. NRRL 30547); B2.5 *Trichoderma* spp., including *Trichoderma atroviride*, strain SC1 (described in International Application No. PCT/IT2008/000196); B2.6 *Trichoderma harzianum* rifai strain KRL-AG2 (also known as strain T-22, /ATCC 208479, e.g. PLANTSHIELD T-22G, Rootshield®, and TurfShield from BioWorks, US); B2.7 *Muscodor roseus* strain A3-5 (Accession No. NRRL 30548); B2.8 *Paecilomyces lilacinus*, in particular spores of *P. lilacinus* strain 251 (AGAL 89/030550; e.g. BioAct from Prophyta); B2.9 *Trichoderma koningii*; B2.10 *Talaromyces flavus*, strain V117b (e.g. PROTUS® WG by Prophyta, DE); B2.11 *Trichoderma atroviride*, strain no. V08/002387; B2.12 *Trichoderma atroviride*, strain no. NMI No.

- V08/002388; B2.13 *Trichoderma atroviride*, strain no. NMI No. V08/002389; B2.14 *Trichoderma atroviride*, strain no. NMI No. V08/002390; B2.15 *Trichoderma harzianum*, strain ITEM 908 (e.g. Triatum-P from Koppert); B2.16 *Pseudozyma aphidis*; B2.17 *Pseudozyma aphidis* (from Yissum Research Development Company of the Hebrew University of Jerusalem); B2.18 *Arthrobotrys dactyloides*; B2.19 *Arthrobotrys oligospora*; B2.20 *Arthrobotrys superba*; B2.21 *Aspergillus flavus*, strain NRRL 21882 (e.g. Afla-Guard® from Syngenta); B2.22 *Aspergillus flavus*, strain AF36 (e.g. AF36 from Arizona Cotton Research and Protection Council, US); B2.23 *Cryptococcus albidus* (e.g. YieldPlus® from Anchor Bio-Technologies, ZA); B2.24 *Cryptococcus flavescentis*, strain 4C (NRRL Y-50379); B2.25 *Gliocladium roseum*, strain 321U from W.F. Stoneman Company LLC; B2.26 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain VRA 1835 (ATCC 90304); B2.27 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain VRA 1984 (DSM16201); B2.28 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain VRA 1985 (DSM16202); B2.29 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain VRA 1986 (DSM16203); B2.30 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG B20/5 (IMI390096); B2.31 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG SP log6 (IMI390097); B2.32 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG SP log5 (IMI390098); B2.33 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG BU3 (IMI390099); B2.34 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG BU4 (IMI390100); B2.35 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG 410.3 (IMI390101); B2.36 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG 97/1062/116/1.1 (IMI390102); B2.37 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG B22/SP1287/3.1 (IMI390103); B2.38 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG SH1 (IMI390104); B2.39 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, in particular strain FOC PG B22/SP1190/3.2 (IMI390105; e.g. Rotstop® from Verdara and FIN, PG-Agromaster®, PG-Fungler®, PG-IBL®, PG-Poszwald®, and Rotex® from e-nema, DE); B2.40 *Pythium oligandrum*, strain DV74 or M1 (ATCC 38472; e.g. Polyversum from Biopreparaty, CZ); B2.41 *Saccharomyces cerevisiae*, strain CNCM No. I-3936 (from Lesaffre et Compagnie, FR); B2.42 *Saccharomyces cerevisiae*, strain CNCM No. I-3937 (from Lesaffre et Compagnie, FR); B2.43 *Saccharomyces cerevisiae*, strain CNCM No. I-3938 (from Lesaffre et Compagnie, FR); B2.44 *Saccharomyces cerevisiae*, strain CNCM No. I-3939 (from Lesaffre et Compagnie, FR); B2.45 *Sclerotinia citrinum*; B2.46 *Trichoderma asperellum*, strain ICC 012 from Isagro; B2.47 *Trichoderma asperellum*, strain SKT-1 (e.g. ECO-HOPE® from Kumiai Chemical Industry); B2.48 *Trichoderma atroviride*, strain CNCM I-1237 (e.g. Esquive® WP from Agrauxine, FR); B2.49 *Trichoderma atroviride*, strain LC52 (e.g. Tenet by Agrimm Technologies Limited); B2.50 *Trichoderma atroviride*, strain ATCC 20476 (IMI 206040); B2.51 *Trichoderma atroviride*, strain T11 (IMI352941/ CECT20498); B2.52 *Trichoderma harmatum*; B2.53 *Trichoderma harzianum*; B2.54 *Trichoderma harzianum rifai T39* (e.g. Trichodex® from Makhteshim, US); B2.55 *Trichoderma harzianum*, in particular, strain KD (e.g. Trichoplus from Biological Control Products, SA (acquired by Becker Underwood)); B2.56 *Trichoderma harzianum*, strain KD (e.g. Eco-T from Plant Health Products, SZ); B2.57 *Trichoderma harzianum*, strain TH35 (e.g. Root-Pro by Mycontrol); B2.58 *Trichoderma virens* (also known as *Gliocladium virens*), in particular strain GL-21 (e.g. SoilGard 12G by Certis, US); B2.59 *Trichoderma viride*, strain TV1 (e.g. Triatum-P by Koppert); B2.60 *Beauveria bassiana*, strain ATCC 74040 (e.g. Naturalis® from Intrachem Bio Italia); B2.61 *Beauveria bassiana* strain GHA (Accession No. ATCC 74250; e.g. BotaniGuard Es and Mycontrol-O from Laverlam Interna-

tional Corporation); B2.62 *Beauveria bassiana* strain ATP02 (Accession No. DSM 24665); B2.63 *Beauveria bassiana* strain CG 716 (e.g. BoveMax® from Novozymes); B2.64 *Hirsutella citriformis*; B2.65 *Hirsutella thompsonii* (with some strains e.g. Mycohit and ABTEC from Agro Bio-tech Research Centre, IN); B2.66 *Lecanicillium lecanii* (formerly known as *Verticillium lecanii*) conidia of strain KV01 (e.g. Mycotal® and Vertalec® from Koppert/Arysta); B2.67 *Lecanicillium lecanii* (formerly known as *Verticillium lecanii*) conidia of strain DA-OM198499; B2.68 *Lecanicillium lecanii* (formerly known as *Verticillium lecanii*) conidia of strain DA-OM216596; B2.69 *Lecanicillium muscarium* (formerly *Verticillium lecanii*), strain VE 6 / CABI(=IMI) 268317/ CBS102071/ 2.70 ARSEF5128; B2.71 *Metarhizium anisopliae*, strain F52 (DSM3884/ ATCC 90448; e.g. BIO 1020 by Bayer CropScience and also e.g. Met52 by Novozymes); B2.72 *Metarhizium anisopliae* var. *acridum* (e.g. GreenGuard by Becker Underwood, US); B2.73 *Metarhizium anisopliae* var. *acridum* isolate IMI 330189 (ARSEF7486; e.g. Green Muscle by Biological Control Products); B2.74 *Nomuraea rileyi*; B2.75 *Paecilomyces fumosoroseus* (new: *Isaria fumosorosea*), strain apopka 97 (e.g. PreFeRal® WG from Biobest); B2.76 *Paecilomyces fumosoroseus* (new: *Isaria fumosorosea*) strain FE 9901 (e.g. NoFly® from Natural Industries Inc., a Novozymes company); B2.77 *Harposporium anguillulae*; B2.78 *Hirsutella minnesotensis*; B2.79 *Monacrosporium cionopagum*; B2.80 *Monacrosporium psychrophilum*; B2.81 *Myrothecium verrucaria*, strain AARC-0255 (e.g. DiTera™ by Valent Biosciences); B2.82 compositions comprising the fungus *Paecilomyces lilacinus* (commercially available as e.g. MELOCON® or BIOACT); B2.83 *Paecilomyces variotii*, strain Q-09 (e.g. Nemaquim® from Quimia, MX); B2.84 compositions comprising the bacterium *Pasteuria* including *Pasteuria usgae* (commercially available as e.g. ECONEM®); B2.85 *Stagonospora phaseoli* (commercially available e.g. from Syngenta); B2.86 *Trichoderma lignorum*, in particular strain TL-0601 (e.g. Mycotric from Futureco Bioscience, ES); B2.87 *Penicillium bilaii*, strain ATCC 22348 (e.g. JumpStart® from Novozymes); B2.88 *Penicillium bilaii*, in particular strain ATCC 22348 (e.g. PB-50 PROVIDE from Philom Bios Inc., Saskatoon, Saskatchewan); B2.89 *Rhizopogon amylopogon* (e.g. Myco-Sol from Helena Chemical Company); B2.90 *Rhizopogon fulvigleba* (e.g. Myco-Sol from Helena Chemical Company); B2.91 *Trichoderma harzianum*, strain TSTh20; B2.92 *Phoma macrostroma*, strain 94-44B (e.g. Phoma H and Phoma P by Scotts, US); B2.93 *Sclerotinia minor*, strain IMI 344141 (e.g. Sarritor by Agrium Advanced Technologies); B2.94 *Ampelomyces quisqualis*, in particular strain AQ 10 (e.g. AQ 10® by IntrachemBio Italia); B2.95 Arkansas fungus 18, ARF; B2.96 *Aureobasidium pullulans*, in particular blastospores of strain DSM14940; B2.97 *Aureobasidium pullulans*, in particular blastospores of strain DSM14941; B2.98 *Aureobasidium pullulans*, in particular mixtures of blastospores of strains DSM14940 and DSM14941 (e.g. Botector® by bio-ferm, CH); B2.99 *Candida oleophila*, strain O (e.g. Nexy® by BioNext); B2.100 *Candida oleophila*, isolate I-182 (e.g. Aspire® by Ecogen, US); B2.101 *Candida saitoana*, strain NRRL Y-21022 (by Biotechnology Research And Development Corporation); B2.102 *Chaetomium cupreum* (e.g. BIODUPRUM™ by AgriLife); B2.103 *Chaetomium globosum* (e.g. Rivadiom by Rivale); B2.104 *Cladosporium cladosporioides*, strain H39 (by Stichting Dienst Landbouwkundig Onderzoek); B2.105 *Cryptococcus flaveszens*, strain 3C (NRRL Y-50378); B2.106 *Dactylaria candida*; B2.107 *Dilophosphora alopecuri* (e.g. Twist Fungus); B2.108 *Fusarium oxysporum*, strain Fo47 (e.g. Fusaclean by Natural Plant Protection); B2.109 *Gliocladium catenulatum* (Synonym: *Clonostachys rosea* f. *catenulate*) strain J1446 (e.g. Prestop® by AgBio Inc. and also e.g. Primastop® by Kemira Agro Oy); B2.110 *Penicillium vermiculatum*; B2.111 *Pichia anomala*, strain WRL-076 (NRRL Y-30842); B2.112 *Pseudozyma flocculosa*, strain PF-A22 UL (e.g. Sporodex® L by Plant Products Co., CA); B2.113 *Trichoderma*

- atroviride*, strain SKT-1 (FERM P-16510); B2.114 *Trichoderma atroviride*, strain SKT-2 (FERM P-16511); B2.115 *Trichoderma atroviride*, strain SKT-3 (FERM P-17021); B2.116 *Trichoderma gamsii* (formerly *T. viride*), strain ICC080 (IMI CC 392151 CABI, e.g. BioDerma by AGROBIOSOL DE MEXICO, S.A. DE C.V.); B2.117 *Trichoderma harzianum*, strain DB 103 (e.g. T-Gro 7456 by Dagutat Biolab); B2.118 *Trichoderma polysporum*, strain IMI 206039 (e.g. Binab TF WP by BINAB Bio-Innovation AB, Sweden); B2.119 *Trichoderma stromaticum* (e.g. Tricovab by Ceplac; Brazil); B2.120 *Tsukamurella paurometabola*, strain C-924 (e.g. HeberNem®); B2.121 *Ulocladium oudemansii*, in particular strain HRU3 (e.g. Botry-Zen® by Botry-Zen Ltd, NZ); B2.122 *Verticillium albo-atrum* (formerly *V. dahliae*), strain WCS850 (CBS 276.92; e.g. Dutch Trig by Tree Care Innovations); B2.123 *Aschersonia aleyrodis*; B2.124 *Beauveria brongniartii* (e.g. Beaupro from Andermatt Biocontrol AG); B2.125 *Conidiobolus obscurus*; B2.126 *Entomophthora virulenta* (e.g. Vektor from Ecomic); B2.127 *Lagenidium giganteum*; B2.128 *Metarhizium flavoviride*; B2.129 *Mucor haemelis* (e.g. BioAvard from Indore Biotech Inputs & Research); B2.130 *Pandora delphacis*; B2.131 *Sporothrix insectorum* (e.g. Sporothrix Es from Biocerto; BR); B2.132 *Zoophtora radicans*; B2.133 *Fusarium solani*, strain Fs5; B2.134 *Hirsutella rhossiliensis*; B2.135 *Monacrosporium drechsleri*; B2.136 *Monacrosporium gephyropagum*; B2.137 *Nematoctonus geogenius*; B2.138 *Nematoctonus leiosporus*; B2.139 *Neocosmospora vasinfecta*; B2.140 *Paraglomus sp.*, in particular *Paraglomus brasili-anum*; B2.141 *Pochonia chlamydosporia* (also known as *Vercillium chlamydosporium*), in particular *var catenula-ta* (IMI SD 187; e.g. KlamiC from The National Center of Animal and Plant Health (CENSA); CU); B2.142 *Stagonospora heteroderae*; B2.143 *Glomus aggregatum*; B2.144 *Glomus clarum*; B2.145 *Glomus deserticola*; B2.146 *Glomus etunicatum*; B2.147 *Glomus intraradices*; B2.148 *Glomus monosporum*; B2.149 *Glomus mosseae*; B2.150 *Laccaria bicolor*; B2.151 *Rhizopogon luteolus*; B2.152 *Rhizopogon tinctorus*; B2.153 *Rhizopogon vil-lolulus*; B2.154 *Scleroderma cepa*; B2.155 *Suillus granulatus*; B2.156 *Suillus punctatapius*; B2.157 *Trichoderma harzianum*, strain 1295-22; B2.158 *Colletotrichum gloeosporioides*, strain ATCC 20358 (e.g. Collego (aka Lock-Down by Agriultural Research Initiatives); B2.159 *Stagonospora atriplici*; B2.160 *Cylindrocarpon heteronema*; B2.161 *Exophiala jeanselmei*; B2.162 *Exophiala pisciphila*; B2.163 *Fusarium aspergilus*; B2.164 *Gigaspora mar-garita*; B2.165 *Gigaspora monosporum*; B2.166 *Glomus brasilianum*; B2.167 *Laccaria laccata*; B2.168 *Ophi-ostoma piliferum*, strain D97 (e.g. Sylvanex); B2.169 *Sarcocystis singaporensis*; B2.170 *Trichoderma asperellum*, strain T34 (e.g. T34 Biocontrol by Bioncontrol Technologies, ES); B2.171 *Meristacrum asterospermum*; B2.172 *Muscodor albus*, in particular strain 620 (Accession No. NRRL 30547; described in US 2012/0114610); B2.173 *Nomuraea rileyi*, in particular strain SA86101 (cf. Braz. Arch. Biol. Technol. Vol. 46, No.1, pp 13-18); B2.174 *Nomuraea rileyi*, in particular strain GU87401 (cf. Braz. Arch. Biol. Technol. Vol. 46, No.1, pp 13-18); B2.175 *Nomuraea rileyi*, in particular strain SR86151 (cf. Braz. Arch. Biol. Technol. Vol. 46, No.1, pp 13-18); B2.176 *Nomuraea rileyi*, in particular strain CG128 (cf. Braz. Arch. Biol. Technol. Vol. 46, No.1, pp 13-18); B2.177 *Nomuraea rileyi*, in particular strain VA9101 (cf. Braz. Arch. Biol. Technol. Vol. 46, No.1, pp 13-18); B2.178 *Trichoderma album* (product known as e.g. Bio-Zeid) and B2.179 mixtures of *Trichoderma asperellum* strain ICC 012 and *Trichoderma gamsii* strain ICC 080 (product known as e.g. BIO-TAM™ from Bayer CropScience LP, US).

In some embodiments of the invention such biological control agents which are summarized under the term fungi according to the invention include: B2.2.1 *Coniothyrium minitans*, strain CON/M/91-8 (Accession No. DSM-9660); B2.2.2 *Metschnikowia fructicola*, strain NRRL Y-30752; B2.2.4 *Muscodor albus*, strain QST 20799 (Ac-

cession No. NRRL 30547); B2.2.8 *Paecilomyces lilacinus*, spores of *P. lilacinus* strain 251 (AGAL 89/030550); B2.2.26 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1835 (ATCC Accession No. 90304); B2.2.27 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1984 (DSM16201); B2.2.28 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1985 (DSM16202); B2.2.29 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1986 (DSM16203); B2.2.30 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG B20/5 (IMI390096); B2.2.31 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG SP log6 (IMI390097); B2.2.32 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG SP log5 (IMI390098); B2.2.33 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG BU3 (IMI390099); B2.2.34 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG BU4 (IMI390100); B2.2.35 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG 410.3 (IMI390101); B2.2.36 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG 97/1062/116/1.1 (IMI390102); B2.2.37 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG B22/SP1287/3.1 (IMI390103); B2.2.38 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG SH1 (IMI390104); B2.2.39 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG B22/SP1190/3.2 (IMI390105); B2.2.55 *Trichoderma harzianum*, strain KD; B2.2.58 *Trichoderma virens* (also known as *Gliocladium virens*), strain GL-21; B2.2.86 *Trichoderma lignorum*, strain TL-0601; B2.2.88 *Penicillium bilaii*, strain ATCC 22348; B2.2.96 *Aureobasidium pullulans*; Blastospores of strain DSM14940; B2.2.97 *Aureobasidium pullulans*; Blastospores of strain DSM14941; B2.2.98 *Aureobasidium pullulans*, mixtures of blastospores of strains DSM14940 and DSM14941; B2.2.121 *Ulocladium oudemansii*, strain HRU3; B2.2.140 *Paraglomus sp*, *Paraglomus brasilianum*; B2.2.141 *Pochonia chlamydosporia* (also known as *Vercillium chlamydosporium*); var. *catenulata* (IMI SD 187); B2.2.172 *Muscodor albus*, strain 620 (Accession No. NRRL 30547); B2.2.173 *Nomuraea rileyi*, strain SA86101; B2.2.174 *Nomuraea rileyi*, strain GU87401; B2.2.175 *Nomuraea rileyi*, strain SR86151; B2.2.176 *Nomuraea rileyi*, strain CG128 and B2.2.177 *Nomuraea rileyi*, strain VA9101.

According to the invention examples of protozoas to be used or employed according to the invention are (the numbering is used throughout the complete following description of the invention): B3.1 *Nosema locustae* (product known as e.g. NoloBait); B3.2 *Thelohania solenopsis* and B3.3 *Vairimorpha spp.*.

According to the invention examples of viruses to be used or employed according to the invention are (the numbering is used throughout the complete following description of the invention): B4.1 *Adoxophyes orana* (summer fruit tortrix) granulosis virus (GV); (product known as e.g. BIOFA - Capex®); B4.2 *Agrotis segetum* (turnip moth) nuclear polyhedrosis virus (NPV); B4.3 *Anticarsia gemmatalis* (Woolly pyrol moth) mNPV (product known as e.g. Polygen); B4.4 *Autographa californica* (Alfalfa Looper ) mNPV (product known as e.g. VPN80 from Agricola El Sol); B4.5 *Biston suppressaria* (tea looper) NPV; B4.6 *Bombyx mori* (silkworm) NPV; B4.7 *Cryptophlebia leucotreta* (false codling moth) GV (product known as e.g. Cryptex); B4.8 *Cydia pomonella* (Codling moth) granulosis virus (GV) (product known as e.g. Madex Plus); B4.9 *Dendrolimus punctatus* (Mas-son pine moth) CPV; B4.10 *Helicoverpa armigera* NPV (product known as e.g. AgBiTech - ViVUS Max); B4.11 *Helicoverpa* (previously *Heliothis*) *zea* (corn earworm) NPV (product known as e.g. Elcar); B4.12 *Leucoma salicis* (satin moth) NPV; B4.13 *Lymantria dispar* (gypsy moth) NPV (product known as e.g. Gypcheck); B4.14 *Neodiprion abietis* (balsam-fir sawfly) NPV (product known as e.g. Abietiv); B4.15 *Neodiprion lecontei*

(red-headed pinesawfly) NPV (product known as e.g. Lecontvirus); B4.16 *Neodiprion sertifer* (Pine sawfly) NPV (product known as e.g. Neocheck-S); B4.17 *Orgyia pseudotsugata* (Douglas-fir tussock moth) NPV (product known as e.g. Virtuss); B4.18 *Phthorimaea operculella* (tobacco leaf miner) GV (product known as e.g. Matapol); B4.19 *Pieris rapae* (small white) GV; B4.20 *Plutella xylostella* (diamondback moth) GV (product known as e.g. Plutec); B4.21 *Spodoptera albula* (gray-streaked armyworm moth) mNPV (product known as e.g. VPN 82); B4.22 *Spodoptera exempta* (true armyworm) mNPV (product known as e.g. Spodec); B4.23 *Spodoptera exigua* (sugarbeet armyworm) mNPV (product known as e.g. Spexit from Andermatt Biocontrol); B4.24 *Spodoptera frugiperda* (fall armyworm) mNPV (product known as e.g. Baculovirus VPN); B4.25 *Spodoptera littoralis* (tobacco cutworm) NPV (products known as Spodoptrin from NPP Calliope France); and B4.26 *Spodoptera litura* (oriental leafworm moth) NPV (product known as e.g. Littovir).

According to the invention examples of nematodes to be used or employed according to the invention are (the numbering is used throughout the complete following description of the invention): B5.1 *Abbreviata caucasica*; B5.2 *Acuaria spp.*; B5.3 *Agamermis decaudata*; B5.4 *Allantonema spp.*; B5.5 *Amphimermis spp.*; B5.6 *Beddingia* (= *Deladenus*) *siridicola*; B5.7 *Bovienema spp.*; B5.7a *Cameronia spp.*; B5.8 *Chitwoodiella ovofilamentata*; B5.9 *Contortylenchus spp.*; B5.10 *Culicimermis spp.*; B5.11 *Diplotrinae spp.*; B5.12 *Empidomeremis spp.*; B5.13 *Filipjevimermis leipsandra*; B5.14 *Gastromermis spp.*; B5.15 *Gongylonema spp.*; B5.16 *Gynopocilia pseudovipara*; B5.17 *Heterorhabditis spp.*, in particular *Heterorhabditis bacteriophora* (product known as e.g. B-Green); or *Heterorhabditis baujardi*, or *Heterorhabditis heliothidis* (product known as e.g. Nematon); or *Heterorhabditis indica*, *Heterorhabditis marelatus*, *Heterorhabditis megidis*, *Heterorhabditis zealandica*; B5.18 *Hexameris spp.*; B5.19 *Hydromermis spp.*; B5.20 *Isomeris spp.*; B5.21 *Limnomermis spp.*; B5.22 *Maupasiana weissii*; B5.23 *Mermis nigrescens*; B5.24 *Mesomeris spp.*; B5.25 *Neomesomeris spp.*; B5.26 *Neoparasitylenchus rugulosi*; B5.27 *Octomyomeris spp.*; B5.28 *Parasitaphelenchus spp.*; B5.29 *Parasitorhabditis spp.*; B5.30 *Parasitylenchus spp.*; B5.31 *Perutilimermis culicis*; B5.32 *Phasmarhabditis hermaphrodita*; B5.33 *Phy-saloptera spp.*; B5.34 *Protrellatus spp.*; B5.35 *Pterygodermatites spp.*; B5.36 *Romanomeris spp.*; B5.37 *Seuratum cadarachense*; B5.38 *Sphaerulariopsis spp.*; B5.39 *Spirura guianensis*; B5.40 *Steinernema spp.* (= *Neoaplectana spp.*), in particular *Steinernema carpocapsae* (product known as e.g. Biocontrol); or *Steinernema feltiae* (= *Neoaplectana carpocapsae*); (product known as e.g. Nemasys®); or *Steinernema glaseri* (products known as Biotopia); or *Steinernema kraussei* (product known as e.g. Larvesure); or *Steinernema riobrave* (product known as e.g. Biovector); or *Steinernema scapterisci* (product known as e.g. Nematac S); or *Steinernema scarabaei*, or *Steinernema siamkayai*; B5.41 *Strelkovimermis peterseni*; B5.42 *Subulura spp.*; B5.43 *Sulphuretylenchus elongatus* and B5.44 *Tetrameres spp.*

The term “proteins or secondary metabolite” refers to any compound, substance or byproduct of a fermentation of a microorganism that has pesticidal activity. The definition comprises any compound, substance or byproduct of a fermentation of a microorganism that has fungicidal activity. Examples of such proteins or secondary metabolites to be used or employed according to the invention are (the numbering is used throughout the complete following description of the invention): B6.1 Harpin (isolated by *Erwinia amylovora*, product known as e.g. Harp-N-Tek™, Messenger®, Employ™, ProAct™); B6.2 terpene constituents of extract of *Chenopodium am-*



*brotsioides* near *ambrosioides* as synthetically manufactured (product known as e.g. Requiem® from Bayer CropScience LP, US).

In some embodiments of the invention such biological control agents which are summarized under the term proteins or secondary metabolites according to the invention comprise B6.6.2 terpene constituents of extract of *Che-*  
 5 *nopodium ambrosioides* near *ambrosioides* as synthetically manufactured.

Unless indicated otherwise, the embodiments described below for the compositions disclosed herein are also applicable to respective embodiments of other aspects disclosed herein.

Preferred are biological control agents selected from the group of bacteria consisting of

- 10 B1.1.1 *Bacillus subtilis*, strain QST713/AQ713 (Accession No. NRRL B21661); B1.2 *Bacillus subtilis* strain AQ153 (Accession No. NRRL 55614); B1.3 *Bacillus sp.* strain AQ175 (ATCC Accession No. 55608); B1.4 *Bacillus sp.* strain AQ177 (ATCC Accession No. 55609); B1.5 *Bacillus sp.* strain AQ178 (ATCC Accession No. 53522); B1.1.6 *Bacillus pumilus*, strain QST2808 (Accession No. NRRL B-30087); B1.1.7 *Bacillus pu-*  
 15 *milus*, strain GB34; B1.8 *Bacillus thuringiensis* strain AQ52 (Accession No. NRRL B-21619); B1.1.9 *Streptomyces sp.*, strain having Accession No. NRRL B-30145; B1.1.10 *Streptomyces galbus* (= *Streptomyces gris-*  
*eoviridis*), strain QST 6047 (Accession No. NRRL B-30232); B1.1.11 *Bacillus chitinosporus*, strain AQ746 (Accession No. NRRL B-21618); B1.12 *Bacillus mycoides*, strain AQ726 (having Accession No. NRRL B-21664); B1.1.13 *Bacillus pumilus*, strain AQ717 (having Accession No. NRRL B21662); B1.1.14 *Bacillus*  
 20 *subtilis*, strain AQ743 (having Accession No. NRRL B-21665); B1.15 *Rhodococcus globerulus* strain AQ719 (Accession No. NRRL B21663); B1.1.16 *Bacillus thuringiensis subsp. aizawai*, strain ABTS-1857 (SD-1372); B1.1.17 *Bacillus firmus*, strain CNMC I-1582; B1.1.18 *Bacillus subtilis*, strain AQ30002, (Accession No. NRRL B-50421); B1.1.19 *Bacillus subtilis*, strain AQ30004 (Accession No. NRRL B-50455); B1.20 *Ba-*  
*cillus amyloliquefaciens*, strain D747; B1.1.21 *Bacillus pumilus*, strain BU F-33; B1.22 *B. subtilis var. amy-*  
*loliquefaciens* strain FZB24; B1.1.23 *Paenibacillus polymyxa*, strain AC-1; B1.24 *Pseudomonas prora-*  
 25 *dix*, B1.25 *Bacillus amyloliquefaciens* strain MBI 600; B1.26 *Bacillus amyloliquefaciens* strain GB03; B1.27 *Bacillus amyloliquefaciens* strain DB 101; B1.28 *Bacillus amyloliquefaciens* strain DB 102; B1.29 *Bacillus amyloliquefaciens* isolate B246 ; B1.1.30 *Bacillus licheniformis*, strain SB3086; B1.1.31 *Pseudomonas syrin-*  
*gae*, strain MA-4, B1.32 *Pseudomonas syringae* strain 742RS; B1.1.33 *Pseudomonas fluorescens*, strain 1629RS; B1.1.34 *Streptomyces galbus* (*Streptomyces griseoviridis*), strain K61 (Accession No. DSM 7206);  
 30 B1.1.35 *Streptomyces lydicus*, strain WYEC108; B1.1.36 *Agrobacterium radiobacter*, strain K84; B1.37 *Agrobacterium radiobacter* strain K1026; B1.38 *Bacillus lentimorbus*; B1.39 *Bacillus mycoides*, isolate J.; B1.1.40 *Bacillus sphaericus*, Serotype H5a5b strain 2362; B1.41 *Bacillus thuringiensis subsp. kurstaki* strain BMP 123 from Becker Microbial Products, IL; B1.1.42 *Bacillus thuringiensis subsp. aizawai*, serotype H-7; B1.43 *Bacillus thuringiensis subsp. kurstaki* strain HD-1; B1.44 *Bacillus thuringiensis subsp. tenebrionis*  
 35 strain NB 176; B1.45 *Bacillus thuringiensis var. japonensis* strain Buibui; B1.1.46 *Burkholderia spp.*, strain A396 (Accession No. NRRL B-50319); B1.1.47 *Chromobacterium subtsugae*, strain PRAA4-1T (MBI-203); B1.48 *Paenibacillus popilliae* (formerly *Bacillus popilliae*); B1.49 *Xenorhabdus luminescens*; B1.50

*Xenorhabdus nematophila*; B1.1.51 *Bacillus thuringiensis* subspecies *israelensis* (serotype H-14); B1.1.52 *Bacillus amyloliquefaciens*, strain FZB42; B1.53 *Bacillus cereus*; B1.54 spores of *Bacillus cereus* strain CNCM I-1562; B1.55 *Bacillus laterosporus* (also known as *Brevibacillus laterosporus*); B1.56 *Bacillus megaterium*, strain YFM3.25; B1.57 *Bacillus mojavenensis*, strain SR11 (CECT-7666); B1.58 *Bacillus nemato-*  
 5 *cida*; B1.59 *Pasteuria nishizawae*; B1.60 *Pasteuria penetrans* (formerly *Bacillus penetrans*); B1.61 *Pasteuria usgae*; B1.62 compositions comprising nematicidal *Streptomyces* sp., such as *Streptomyces lydicus*; B1.63 compositions comprising nematicidal *Streptomyces* sp., such as *Streptomyces saraceticus*; B1.1.64 *Bacillus amyloliquefaciens*, strain IN937a; B1.1.65 *Bacillus cereus*, strain BP01 (ATCC Accession No. 55675); B1.66 *Bradyrhizobium japonicum*; B1.1.67 *Delftia acidovorans*, strain RAY209; B1.68 *Lactobacillus* sp.; B1.1.69  
 10 *Pseudomonas aeruginosa*, strain PN1; B1.1.70 *Rhizobium leguminosarum*; Bv. *viceae* strain Z25 (Accession No. CECT 4585); B1.1.71 *Streptomyces acidiscabies*, strain RL-110T.

Preferred are biological control agents selected from the group of fungi consisting of

B2.2.1 *Coniothyrium minitans*, strain CON/M/91-8 (Accession No. DSM-9660); B2.2.2 *Metschnikowia fruc-*  
 15 *ticola*, strain NRRL Y-30752; B2.3 *Microsphaeropsis ochracea*; B2.2.4 *Muscodor albus*, strain QST 20799 (Accession No. NRRL 30547); B2.5 *Trichoderma atroviride*, strain SC1; B2.6 *Trichoderma harzianum* rifai strain KRL-AG2 (also known as strain T-22, /ATCC Accession No. 208479); B2.7 *Muscodor roseus* strain A3-5 (Accession No. NRRL 30548); B2.2.8 *Paecilomyces lilacinus*, spores of *P. lilacinus* strain 251 (AGAL 89/030550); B2.9 *Trichoderma koningii*; B2.10 *Talaromyces flavus*, strain V117b; B2.11 *Trichoderma atro-*  
 20 *viride*, strain no. V08/002387; B2.12 *Trichoderma atroviride*, strain no. NMI No. V08/002388; B2.13 *Trichoderma atroviride*, strain no. NMI No. V08/002389; B2.14 *Trichoderma atroviride*, strain no. NMI No. V08/002390; B2.15 *Trichoderma harzianum*, strain ITEM 908; B2.16 *Pseudozyma aphidis*; B2.17 *Pseudozyma aphidis* (from Yissum Research Development Company of the Hebrew University of Jerusa-  
 lem); B2.18 *Arthrobotrys dactyloides*; B2.19 *Arthrobotrys oligospora*; B2.20 *Arthrobotrys superba*; B2.21  
 25 *Aspergillus flavus*, strain NRRL 21882; B2.22 *Aspergillus flavus*, strain AF36; B2.23 *Cryptococcus albidus*; B2.24 *Cryptococcus flavescens*, strain 4C (Accession No. NRRL Y-50379); B2.25 *Gliocladium roseum*, strain 321U from W.F. Stoneman Company LLC; B2.2.26 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1835 (ATCC Accession No. 90304); B2.2.27 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1984 (DSM16201); B2.2.28 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1985  
 30 (DSM16202); B2.2.29 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain VRA 1986 (DSM16203); B2.2.30 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG B20/5 (IMI390096); B2.2.31 *Phle-*  
*biopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG SP log6 (IMI390097); B2.2.32 *Phlebiopsis* (or  
*Phlebia* or *Peniophora*) *gigantea*, strain FOC PG SP log5 (IMI390098); B2.2.33 *Phlebiopsis* (or *Phlebia* or  
*Peniophora*) *gigantea*, strain FOC PG BU3 (IMI390099); B2.2.34 *Phlebiopsis* (or *Phlebia* or *Peniophora*)  
 35 *gigantea*, strain FOC PG BU4 (IMI390100); B2.2.35 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG 410.3 (IMI390101); B2.2.36 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG 97/1062/116/1.1 (IMI390102); B2.2.37 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG B22/SP1287/3.1 (IMI390103); B2.2.38 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG

- SH1 (IMI390104); B2.2.39 *Phlebiopsis* (or *Phlebia* or *Peniophora*) *gigantea*, strain FOC PG B22/SP1190/3.2 (IMI390105); B2.40 *Pythium oligandrum*, strain DV74 or M1 (ATCC 38472); B2.41 *Saccharomyces cerevisiae*, strain CNCM No. I-3936 (from Lesaffre et Compagnie, FR); B2.42 *Saccharomyces cerevisiae*, strain CNCM No. I-3937 (from Lesaffre et Compagnie, FR); B2.43 *Saccharomyces cerevisiae*, strain CNCM No. I-3938 (from Lesaffre et Compagnie, FR); B2.44 *Saccharomyces cerevisiae*, strain CNCM No. I-3939 (from Lesaffre et Compagnie, FR); B2.45 *Scleroderma citrinum*; B2.46 *Trichoderma asperellum*, strain ICC 012 from Isagro; B2.47 *Trichoderma asperellum*, strain SKT-1; B2.48 *Trichoderma atroviride*, strain CNCM I-1237; B2.49 *Trichoderma atroviride*, strain LC52; B2.50 *Trichoderma atroviride*, strain ATCC 20476 (IMI 206040); B2.51 *Trichoderma atroviride*, strain T11 (IMI352941/ CECT20498); B2.52 *Trichoderma harmatum*; B2.53 *Trichoderma harzianum*; B2.54 *Trichoderma harzianum* rifai T39; B2.2.55 *Trichoderma harzianum*, strain KD; B2.56 *Trichoderma harzianum*, strain KD; B2.57 *Trichoderma harzianum*, strain TH35; B2.2.58 *Trichoderma virens* (also known as *Gliocladium virens*), strain GL-21; B2.59 *Trichoderma viride*, strain TV1(); B2.60 *Beauveria bassiana*, strain ATCC 74040; B2.61 *Beauveria bassiana* strain GHA (ATCC Accession No. 74250); B2.62 *Beauveria bassiana* strain ATP02 (Accession No. DSM 24665); B2.63 *Beauveria bassiana* strain CG 716; B2.64 *Hirsutella citriformis*; B2.65 *Hirsutella thompsonii*; B2.66 *Lecanicillium lecanii* (formerly known as *Verticillium lecanii*) conidia of strain KV01; B2.67 *Lecanicillium lecanii* (formerly known as *Verticillium lecanii*) conidia of strain DAOM198499; B2.68 *Lecanicillium lecanii* (formerly known as *Verticillium lecanii*) conidia of strain DAOM216596; B2.69 *Lecanicillium muscarium* (formerly *Verticillium lecanii*), strain VE 6 / CABI(=IMI) 268317/ CBS102071/ ARSEF5128; B2.71 *Metarhizium anisopliae*, strain F52 (DSM3884/ ATCC 90448); B2.72 *Metarhizium anisopliae* var. *acridum*; B2.73 *Metarhizium anisopliae* var. *acridum* isolate IMI 330189 (ARSEF7486); B2.74 *Nomuraea rileyi*; B2.75 *Paecilomyces fumosoroseus* (new: *Isaria fumosorosea*), strain apopka 97; B2.76 *Paecilomyces fumosoroseus* (new: *Isaria fumosorosea*) strain FE 9901; B2.77 *Harposporium anguillulae*; B2.78 *Hirsutella minnesotensis*; B2.79 *Monacrosporium cionopagum*; B2.80 *Monacrosporium psychrophilum*; B2.81 *Myrothecium verrucaria*, strain AARC-0255; B2.82 compositions comprising the fungus *Paecilomyces lilacinus*; B2.83 *Paecilomyces variotii*, strain Q-09; B2.84 compositions comprising the bacterium *Pasteuria* including *Pasteuria usgae*; B2.85 *Stagonospora phaseoli*; B2.2.86 *Trichoderma lignorum*, strain TL-0601; B2.87 *Penicillium bilaii*, strain ATCC 22348; B2.2.88 *Penicillium bilaii*, strain ATCC 22348; B2.89 *Rhizopogon amylopogon*; B2.90 *Rhizopogon fulvigleba*; B2.91 *Trichoderma harzianum*, strain TSTh20; B2.92 *Phoma macrostroma*, strain 94-44B; B2.93 *Sclerotinia minor*, strain IMI 344141.

Preferred are biological control agents selected from the group of proteins or secondary metabolites consisting of: B6.6.2 terpene constituents of extract of *Chenopodium ambrosioides* near *ambrosioides* as synthetically manufactured.

35

Particularly preferred are biological control agents selected from the group of bacteria consisting of B1.1.1 *Bacillus subtilis*, strain QST713/AQ713 (Accession No. NRRL B21661); B1.2 *Bacillus subtilis* strain AQ153 (Accession No. NRRL 55614); B1.3 *Bacillus* sp. strain AQ175 (ATCC Accession No. 55608); B1.4 *Bacillus* sp.

strain AQ177 (ATCC Accession No. 55609); B1.5 *Bacillus* sp. strain AQ178 (ATCC Accession No. 53522); B1.1.6 *Bacillus pumilus*, strain QST2808 (Accession No. NRRL B-30087); B1.1.7 *Bacillus pumilus*, strain GB34; B1.8 *Bacillus thuringiensis* strain AQ52 (Accession No. NRRL B-21619); B1.1.9 *Streptomyces* sp., strain having Accession No. NRRL B-30145; B1.1.10 *Streptomyces galbus* (= *Streptomyces griseoviridis*), strain  
 5 QST 6047 (Accession No. NRRL B-30232); B1.1.11 *Bacillus chitinosporus*, strain AQ746 (Accession No. NRRL B-21618); B1.12 *Bacillus mycoides*, strain AQ726 (having Accession No. NRRL B-21664); B1.1.13 *Bacillus pumilus*, strain AQ717 (having Accession No. NRRL B21662); B1.1.14 *Bacillus subtilis*, strain AQ743 (having Accession No. NRRL B-21665); B1.15 *Rhodococcus globerulus* strain AQ719 (Accession No. NRRL B21663); B1.1.16 *Bacillus thuringiensis* subsp. *aizawai*, strain ABTS-1857 (SD-1372); B1.1.17 *Bacillus firmus*,  
 10 strain CNMC I-1582; B1.1.18 *Bacillus subtilis*, strain AQ30002, (Accession No. NRRL B-50421); B1.1.19 *Bacillus subtilis*, strain AQ30004 (Accession No. NRRL B-50455).

Particularly preferred are biological control agents selected from the group of fungi consisting of B2.2.1 *Coniothyrium minitans*, strain CON/M/91-8 (Accession No. DSM-9660); B2.2.2 *Metschnikowia fructicola*, strain NRRL Y-30752; B2.3 *Microsphaeropsis ochracea*; B2.2.4 *Muscodor albus*, strain QST 20799 (Accession No.  
 15 NRRL 30547); B2.5 *Trichoderma atroviride*, strain SC1; B2.6 *Trichoderma harzianum* rifai strain KRL-AG2 (also known as strain T-22, /ATCC Accession No. 208479); B2.7 *Muscodor roseus* strain A3-5 (Accession No. NRRL 30548); B2.2.8 *Paecilomyces lilacinus*, spores of *P. lilacinus* strain 251 (AGAL 89/030550); B2.9 *Trichoderma koningii*; B2.10 *Talaromyces flavus*, strain V117b; B2.11 *Trichoderma atroviride*, strain no. V08/002387; B2.12 *Trichoderma atroviride*, strain no. NMI No. V08/002388; B2.13 *Trichoderma atroviride*,  
 20 strain no. NMI No. V08/002389; B2.14 *Trichoderma atroviride*, strain no. NMI No. V08/002390; B2.15 *Trichoderma harzianum*, strain ITEM 908.

Most preferred are biological control agents selected from the group of bacteria consisting of B1.1 *Bacillus subtilis*, in particular strain QST713/AQ713 (available as SERENADE MAX from Bayer CropScience LP, US,  
 25 having NRRL Accession No. B21661 and described in U.S. Patent No. 6,060,051); B1.6 *Bacillus pumilus*, in particular strain QST2808 (available as Sonata® from Bayer CropScience LP, US, having Accession No. NRRL B-30087 and described in U.S. Patent No. 6,245,551) and B1.18 *Bacillus subtilis*, in particular strain AQ30002, (having Accession No. NRRL B-50421 and described in U.S. Patent Application No. 13/330,576).

30 Furthermore preferred are biological control agents selected from the group of bacteria consisting of B1.1.1 *Bacillus subtilis*, strain QST713/AQ713 (Accession No. NRRL B21661); B1.1.6 *Bacillus pumilus*, strain QST2808 (Accession No. NRRL B-30087) and B1.1.18 *Bacillus subtilis*, strain AQ30002, (Accession No. NRRL B-50421).

35 The following combinations exemplify specific embodiments of the composition according to the present invention:

[illegible]

[illegible]

[illegible]

[illegible]



[illegible]

(B1.141), (I-3) + (B1.142), (I-3) + (B1.143), (I-3) + (B1.144), (I-3) + (B1.145), (I-3) + (B1.146), (I-3) +  
(B1.147), (I-3) + (B1.148), (I-3) + (B1.149), (I-3) + (B1.150), (I-3) + (B1.151), (I-3) + (B1.152), (I-3) +  
(B1.153), (I-3) + (B1.154), (I-3) + (B1.155), (I-3) + (B1.156), (I-3) + (B1.157), (I-3) + (B1.158), (I-3) +  
(B1.159), (I-3) + (B1.160), (I-3) + (B1.161), (I-3) + (B1.162), (I-3) + (B1.163), (I-3) + (B1.164), (I-3) +  
5 (B1.165), (I-3) + (B1.166), (I-3) + (B1.167), (I-3) + (B1.168), (I-3) + (B1.169), (I-3) + (B1.170), (I-3) +  
(B1.171), (I-3) + (B1.172), (I-3) + (B1.173), (I-3) + (B1.174), (I-3) + (B1.175), (I-3) + (B1.176), (I-3) +  
(B1.177), (I-3) + (B1.178), (I-3) + (B1.179), (I-3) + (B1.180), (I-3) + (B1.181), (I-3) + (B1.182), (I-3) +  
(B1.183), (I-3) + (B1.184), (I-3) + (B1.185), (I-3) + (B1.186), (I-3) + (B1.187), (I-3) + (B1.188), (I-3) +  
(B1.189), (I-3) + (B1.190), (I-3) + (B1.191), (I-3) + (B1.192), (I-3) + (B1.193), (I-3) + (B1.194), (I-3) +  
10 (B1.195), (I-3) + (B1.196), (I-3) + (B1.197), (I-3) + (B1.198), (I-3) + (B1.199), (I-3) + (B1.1.1), (I-3) +  
(B1.1.6), (I-3) + (B1.1.7), (I-3) + (B1.1.9), (I-3) + (B1.1.10), (I-3) + (B1.1.11), (I-3) + (B1.1.13), (I-3) +  
(B1.1.14), (I-3) + (B1.1.16), (I-3) + (B1.1.17), (I-3) + (B1.1.18), (I-3) + (B1.1.19), (I-3) + (B1.1.21), (I-3) +  
(B1.1.23), (I-3) + (B1.1.30), (I-3) + (B1.1.31), (I-3) + (B1.1.33), (I-3) + (B1.1.34), (I-3) + (B1.1.35), (I-3) +  
(B1.1.36), (I-3) + (B1.1.40), (I-3) + (B1.1.42), (I-3) + (B1.1.46), (I-3) + (B1.1.47), (I-3) + (B1.1.51), (I-3) +  
15 (B1.1.52), (I-3) + (B1.1.64), (I-3) + (B1.1.65), (I-3) + (B1.1.67), (I-3) + (B1.1.69), (I-3) + (B1.1.70), (I-3) +  
(B1.1.71), (I-3) + (B1.1.74), (I-3) + (B1.1.75), (I-3) + (B1.1.76), (I-3) + (B1.1.77), (I-3) + (B1.1.81), (I-3) +  
(B1.1.82), (I-3) + (B1.1.83), (I-3) + (B1.1.84), (I-3) + (B1.1.85), (I-3) + (B1.1.86), (I-3) + (B1.1.87), (I-3) +  
(B1.1.89), (I-3) + (B1.1.90), (I-3) + (B1.1.93), (I-3) + (B1.1.103), (I-3) + (B1.1.124), (I-3) + (B1.1.125), (I-3) +  
(B1.1.129), (I-3) + (B1.1.130), (I-3) + (B1.1.149), (I-3) + (B1.1.162), (I-3) + (B1.1.163), (I-3) + (B1.1.164), (I-  
20 3) + (B1.1.165), (I-3) + (B1.1.167), (I-3) + (B1.1.168), (I-3) + (B1.1.169), (I-3) + (B1.1.170), (I-3) +  
(B1.1.171), (I-3) + (B1.1.172), (I-3) + (B1.1.173), (I-3) + (B1.1.174), (I-3) + (B1.1.178), (I-3) + (B1.1.179), (I-  
3) + (B1.1.180), (I-3) + (B1.1.181), (I-3) + (B1.1.189), (I-3) + (B1.1.193), (I-3) + (B1.1.196), (I-3) + (B2.1), (I-  
3) + (B2.2), (I-3) + (B2.3), (I-3) + (B2.4), (I-3) + (B2.5), (I-3) + (B2.6), (I-3) + (B2.7), (I-3) + (B2.8), (I-3) +  
(B2.9), (I-3) + (B2.10), (I-3) + (B2.11), (I-3) + (B2.12), (I-3) + (B2.13), (I-3) + (B2.14), (I-3) + (B2.15), (I-3) +  
25 (B2.16), (I-3) + (B2.17), (I-3) + (B2.18), (I-3) + (B2.19), (I-3) + (B2.20), (I-3) + (B2.21), (I-3) + (B2.22), (I-3) +  
(B2.23), (I-3) + (B2.24), (I-3) + (B2.25), (I-3) + (B2.26), (I-3) + (B2.27), (I-3) + (B2.28), (I-3) + (B2.29), (I-3) +  
(B2.30), (I-3) + (B2.31), (I-3) + (B2.32), (I-3) + (B2.33), (I-3) + (B2.34), (I-3) + (B2.35), (I-3) + (B2.36), (I-3) +  
(B2.37), (I-3) + (B2.38), (I-3) + (B2.39), (I-3) + (B2.40), (I-3) + (B2.41), (I-3) + (B2.42), (I-3) + (B2.43), (I-3) +  
(B2.44), (I-3) + (B2.45), (I-3) + (B2.46), (I-3) + (B2.47), (I-3) + (B2.48), (I-3) + (B2.49), (I-3) + (B2.50), (I-3) +  
30 (B2.51), (I-3) + (B2.52), (I-3) + (B2.53), (I-3) + (B2.54), (I-3) + (B2.55), (I-3) + (B2.56), (I-3) + (B2.57), (I-3) +  
(B2.58), (I-3) + (B2.59), (I-3) + (B2.60), (I-3) + (B2.61), (I-3) + (B2.62), (I-3) + (B2.63), (I-3) + (B2.64), (I-3) +  
(B2.65), (I-3) + (B2.66), (I-3) + (B2.67), (I-3) + (B2.68), (I-3) + (B2.69), (I-3) + (B2.70), (I-3) + (B2.71), (I-3) +  
(B2.72), (I-3) + (B2.73), (I-3) + (B2.74), (I-3) + (B2.75), (I-3) + (B2.76), (I-3) + (B2.77), (I-3) + (B2.78), (I-3) +  
(B2.79), (I-3) + (B2.80), (I-3) + (B2.81), (I-3) + (B2.82), (I-3) + (B2.83), (I-3) + (B2.84), (I-3) + (B2.85), (I-3) +  
35 (B2.86), (I-3) + (B2.87), (I-3) + (B2.88), (I-3) + (B2.89), (I-3) + (B2.90), (I-3) + (B2.91), (I-3) + (B2.92), (I-3) +  
(B2.93), (I-3) + (B2.94), (I-3) + (B2.95), (I-3) + (B2.96), (I-3) + (B2.97), (I-3) + (B2.98), (I-3) + (B2.99), (I-3) +  
(B2.100), (I-3) + (B2.101), (I-3) + (B2.102), (I-3) + (B2.103), (I-3) + (B2.104), (I-3) + (B2.105), (I-3) +  
(B2.106), (I-3) + (B2.107), (I-3) + (B2.108), (I-3) + (B2.109), (I-3) + (B2.110), (I-3) + (B2.111), (I-3) +  
(B2.112), (I-3) + (B2.113), (I-3) + (B2.114), (I-3) + (B2.115), (I-3) + (B2.116), (I-3) + (B2.117), (I-3) +

(B2.118), (I-3) + (B2.119), (I-3) + (B2.120), (I-3) + (B2.121), (I-3) + (B2.122), (I-3) + (B2.123), (I-3) + (B2.124), (I-3) + (B2.125), (I-3) + (B2.126), (I-3) + (B2.127), (I-3) + (B2.128), (I-3) + (B2.129), (I-3) + (B2.130), (I-3) + (B2.131), (I-3) + (B2.132), (I-3) + (B2.133), (I-3) + (B2.134), (I-3) + (B2.135), (I-3) + (B2.136), (I-3) + (B2.137), (I-3) + (B2.138), (I-3) + (B2.139), (I-3) + (B2.140), (I-3) + (B2.141), (I-3) + (B2.142), (I-3) + (B2.143), (I-3) + (B2.144), (I-3) + (B2.145), (I-3) + (B2.146), (I-3) + (B2.147), (I-3) + (B2.148), (I-3) + (B2.149), (I-3) + (B2.150), (I-3) + (B2.151), (I-3) + (B2.152), (I-3) + (B2.153), (I-3) + (B2.154), (I-3) + (B2.155), (I-3) + (B2.156), (I-3) + (B2.157), (I-3) + (B2.158), (I-3) + (B2.159), (I-3) + (B2.160), (I-3) + (B2.161), (I-3) + (B2.162), (I-3) + (B2.163), (I-3) + (B2.164), (I-3) + (B2.165), (I-3) + (B2.166), (I-3) + (B2.167), (I-3) + (B2.168), (I-3) + (B2.169), (I-3) + (B2.170), (I-3) + (B2.171), (I-3) + (B2.172), (I-3) + (B2.173), (I-3) + (B2.174), (I-3) + (B2.175), (I-3) + (B2.176), (I-3) + (B2.177), (I-3) + (B2.178), (I-3) + (B2.179), (I-3) + (B2.2.1), (I-3) + (B2.2.2), (I-3) + (B2.2.4), (I-3) + (B2.2.8), (I-3) + (B2.2.26), (I-3) + (B2.2.27), (I-3) + (B2.2.28), (I-3) + (B2.2.29), (I-3) + (B2.2.30), (I-3) + (B2.2.31), (I-3) + (B2.2.32), (I-3) + (B2.2.33), (I-3) + (B2.2.34), (I-3) + (B2.2.35), (I-3) + (B2.2.36), (I-3) + (B2.2.37), (I-3) + (B2.2.38), (I-3) + (B2.2.39), (I-3) + (B2.2.55), (I-3) + (B2.2.58), (I-3) + (B2.2.86), (I-3) + (B2.2.88), (I-3) + (B2.2.94), (I-3) + (B2.2.96), (I-3) + (B2.2.97), (I-3) + (B2.2.98), (I-3) + (B2.2.121), (I-3) + (B2.2.140), (I-3) + (B2.2.141), (I-3) + (B2.2.172), (I-3) + (B2.2.173), (I-3) + (B2.2.174), (I-3) + (B2.2.175), (I-3) + (B2.2.176), (I-3) + (B2.2.177), (I-3) + (B3.1), (I-3) + (B3.2), (I-3) + (B3.3), (I-3) + (B4.1), (I-3) + (B4.2), (I-3) + (B4.3), (I-3) + (B4.4), (I-3) + (B4.5), (I-3) + (B4.6), (I-3) + (B4.7), (I-3) + (B4.8), (I-3) + (B4.9), (I-3) + (B4.10), (I-3) + (B4.11), (I-3) + (B4.12), (I-3) + (B4.13), (I-3) + (B4.14), (I-3) + (B4.15), (I-3) + (B4.16), (I-3) + (B4.17), (I-3) + (B4.18), (I-3) + (B4.19), (I-3) + (B4.20), (I-3) + (B4.21), (I-3) + (B4.22), (I-3) + (B4.23), (I-3) + (B4.24), (I-3) + (B4.25), (I-3) + (B4.26), (I-3) + (B5.1), (I-3) + (B5.2), (I-3) + (B5.3), (I-3) + (B5.4), (I-3) + (B5.5), (I-3) + (B5.6), (I-3) + (B5.7), (I-3) + (B5.8), (I-3) + (B5.9), (I-3) + (B5.10), (I-3) + (B5.11), (I-3) + (B5.12), (I-3) + (B5.13), (I-3) + (B5.14), (I-3) + (B5.15), (I-3) + (B5.16), (I-3) + (B5.17), (I-3) + (B5.18), (I-3) + (B5.19), (I-3) + (B5.20), (I-3) + (B5.21), (I-3) + (B5.22), (I-3) + (B5.23), (I-3) + (B5.24), (I-3) + (B5.25), (I-3) + (B5.26), (I-3) + (B5.27), (I-3) + (B5.28), (I-3) + (B5.29), (I-3) + (B5.30), (I-3) + (B5.31), (I-3) + (B5.32), (I-3) + (B5.33), (I-3) + (B5.34), (I-3) + (B5.35), (I-3) + (B5.36), (I-3) + (B5.37), (I-3) + (B5.38), (I-3) + (B5.39), (I-3) + (B5.40), (I-3) + (B5.41), (I-3) + (B5.42), (I-3) + (B5.43), (I-3) + (B5.44), (I-3) + (B6.1), (I-3) + (B6.2), (I-3) + (B6.6.2).

The following combinations are particularly preferred:

(I-1) + (B1.1), (I-1) + (B1.6), (I-1) + (B1.18), (I-2) + (B1.1), (I-2) + (B1.6), (I-2) + (B1.18), (I-3) + (B1.1), (I-3) + (B1.6), (I-3) + (B1.18), (I-1) + (B1.1.1), (I-1) + (B1.1.6), (I-1) + (B1.1.18), (I-2) + (B1.1.1), (I-2) + (B1.1.6), (I-2) + (B1.1.18), (I-3) + (B1.1.1), (I-3) + (B1.1.6), (I-3) + (B1.1.18).

In the context of the present invention the term “composition” shall mean a physical mixture comprising compounds of the formula (I) and at least one biological control agent.

As used herein, the term “comprising” is to be interpreted as specifying the presence of the stated features, integers, steps, components or compounds as referred to, but does not preclude the presence or addition of one or more features, integers, steps, components or compounds, or groups thereof. Thus, e.g. a composition comprising a

compound of the formula (I) and at least one biological control agent may comprise more compounds or more biological control agents than the actually cited ones, i.e. the composition may further comprise at least one compound (C) selected from the group consisting of fungicides. However, in context with the present disclosure, the term “comprising” also encloses “consisting of” and “including”.

- 5 As used herein the term “at least one” shall refer to either one compound or biological control agent or the like, but also encompasses the presence of at least two, at least three or at least four compounds or biological control agents or the like.

10 The compositions may be used alone or in combination with other active ingredients such as fungicides. The active ingredients specified herein by their “common name” are known and described, for example, in the Pesticide Manual. Where a compound can be present in tautomeric form, such a compound is understood herein before mentioned and herein below also to include, where applicable, corresponding tautomeric forms, even when these are not specifically mentioned in each case.

15 Examples of such fungicides to be used in combination with the inventive composition are (the numbering is used throughout the complete following description of the invention): 1) Inhibitors of the ergosterol biosynthesis, for example (1.1) aldimorph, (1.2) azaconazole, (1.3) bitertanol, (1.4) bromuconazole, (1.5) cyproconazole, (1.6) diclobutrazole, (1.7) difenoconazole, (1.8) diniconazole, (1.9) diniconazole-M, (1.10) dodemorph, (1.11) dodemorph acetate, (1.12) epoxiconazole, (1.13) etaconazole, (1.14) fenarimol, (1.15) fenbuconazole, (1.16) fenhexamid, (1.17) fenpropidin, (1.18) fenpropimorph, (1.19) fluquinconazole, (1.20) flurprimidol, (1.21) flusilazole, (1.22) flutriafol, (1.23) furconazole, (1.24) furconazole-cis, (1.25) hexaconazole, (1.26) imazalil, (1.27) imazalil sulfate, (1.28) imibenconazole, (1.29) ipconazole, (1.30) metconazole, (1.31) myclobutanil, (1.32) naptifine, (1.33) nuarimol, (1.34) oxpoconazole, (1.35) paclobutrazol, (1.36) pefurazoate, (1.37) penconazole, (1.38) piperalin, (1.39) prochloraz, (1.40) propiconazole, (1.41) prothioconazole, (1.42) pyributicarb, (1.43) pyrifenoxy, (1.44) quinconazole, (1.45) simeconazole, (1.46) spiroxamine, (1.47) tebuconazole, (1.48) terbinafine, (1.49) tetraconazole, (1.50) triadimefon, (1.51) triadimenol, (1.52) tridemorph, (1.53) triflumizole, (1.54) triforine, (1.55) triticonazole, (1.56) uniconazole, (1.57) uniconazole-p, (1.58) viniconazole, (1.59) voriconazole, (1.60) 1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol, (1.61) methyl 1-(2,2-dimethyl-2,3-dihydro-1H-inden-1-yl)-1H-imidazole-5-carboxylate, (1.62) N'-{5-(difluoromethyl)-2-methyl-4-[3-(trimethylsilyl)propoxy]phenyl}-N-ethyl-N-methylimidoforamide, (1.63) N-ethyl-N-methyl-N'-{2-methyl-5-(trifluoromethyl)-4-[3-(trimethylsilyl)propoxy]phenyl}imidoforamide, (1.64) O-[1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl] 1H-imidazole-1-carbothioate, (1.65) Pyrisoxazole.

2) Inhibitors of the respiratory chain at complex I or II, for example (2.1) bixafen, (2.2) boscalid, (2.3) carboxin, (2.4) diflumerimol, (2.5) fenfuram, (2.6) fluopyram, (2.7) flutolanil, (2.8) fluxapyroxad, (2.9) furametpyr, (2.10) furmecyclox, (2.11) isopyrazam (mixture of syn-epimeric racemate 1RS,4SR,9RS and anti-epimeric racemate 1RS,4SR,9SR), (2.12) isopyrazam (anti-epimeric racemate 1RS,4SR,9SR), (2.13) isopyrazam (anti-epimeric enantiomer 1R,4S,9S), (2.14) isopyrazam (anti-epimeric enantiomer 1S,4R,9R), (2.15) isopyrazam (syn epimeric racemate 1RS,4SR,9RS), (2.16) isopyrazam (syn-epimeric enantiomer 1R,4S,9R), (2.17) isopyrazam (syn-

epimeric enantiomer 1S,4R,9S), (2.18) mepronil, (2.19) oxycarboxin, (2.20) penflufen, (2.21) penthiopyrad, (2.22) sedaxane, (2.23) thifluzamide, (2.24) 1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (2.25) 3-(difluoromethyl)-1-methyl-N-[2-(1,1,2,2-tetrafluoroethoxy)phenyl]-1H-pyrazole-4-carboxamide, (2.26) 3-(difluoromethyl)-N-[4-fluoro-2-(1,1,2,3,3,3-hexafluoropropoxy)phenyl]-1-methyl-1H-pyrazole-4-carboxamide, (2.27) N-[1-(2,4-dichlorophenyl)-1-methoxypropan-2-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.28) 5,8-difluoro-N-[2-(2-fluoro-4-{[4-(trifluoromethyl)pyridin-2-yl]oxy}phenyl)ethyl]quinazolin-4-amine, (2.29) benzovindiflupyr, (2.30) N-[(1S,4R)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.31) N-[(1R,4S)-9-(dichloromethylene)-1,2,3,4-tetrahydro-1,4-methanonaphthalen-5-yl]-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (2.32) 3-(difluoromethyl)-1-methyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.33) 1,3,5-trimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.34) 1-methyl-3-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (2.35) 1-methyl-3-(trifluoromethyl)-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.36) 1-methyl-3-(trifluoromethyl)-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.37) 3-(difluoromethyl)-1-methyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.38) 3-(difluoromethyl)-1-methyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.39) 1,3,5-trimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.40) 1,3,5-trimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (2.41) benodanil, (2.42) 2-chloro-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)pyridine-3-carboxamide, (2.43) Isofetamid.

3) Inhibitors of the respiratory chain at complex III, for example (3.1) ametocetradin, (3.2) amisulbrom, (3.3) azoxystrobin, (3.4) cyazofamid, (3.5) coumethoxystrobin, (3.6) coumoxystrobin, (3.7) dimoxystrobin, (3.8) enoxastrobin, (3.9) famoxadone, (3.10) fenamidone, (3.11) flufenoxystrobin, (3.12) fluoxastrobin, (3.13) kresoxim-methyl, (3.14) metominostrobin, (3.15) orysastrobin, (3.16) picoxystrobin, (3.17) pyraclostrobin, (3.18) pyrametostrobin, (3.19) pyraoxystrobin, (3.20) pyribencarb, (3.21) triclopyricarb, (3.22) trifloxystrobin, (3.23) (2E)-2-(2-{[6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl]oxy}phenyl)-2-(methoxyimino)-N-methylacetamide, (3.24) (2E)-2-(methoxyimino)-N-methyl-2-(2-{[6-(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy]methyl}phenyl)acetamide, (3.25) (2E)-2-(methoxyimino)-N-methyl-2-{2-[(E)-({1-[3-(trifluoromethyl)phenyl]ethoxy}imino)methyl]phenyl}acetamide, (3.26) (2E)-2-{2-[(E)-1-(3-{[(E)-1-fluoro-2-phenylvinyl]oxy}phenyl)ethylidene]amino}oxy)methyl}phenyl}-2-(methoxyimino)-N-methylacetamide, (3.27) Fenaminostrobin, (3.28) 5-methoxy-2-methyl-4-(2-{[6-(1E)-1-[3-(trifluoromethyl)phenyl]ethylidene}amino)oxy]methyl}phenyl)-2,4-dihydro-3H-1,2,4-triazol-3-one, (3.29) methyl (2E)-2-{2-[(cyclopropyl[(4-methoxyphenyl)imino]methyl)sulfanyl]methyl}phenyl}-3-methoxyacrylate, (3.30) N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-formamido-2-hydroxybenzamide, (3.31) 2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide, (3.32) 2-{2-[(2,5-dimethylphenoxy)methyl]phenyl}-2-methoxy-N-methylacetamide.

- 4) Inhibitors of the mitosis and cell division, for example (4.1) benomyl, (4.2) carbendazim, (4.3) chlorfenazole, (4.4) diethofencarb, (4.5) ethaboxam, (4.6) fluopicolide, (4.7) fuberidazole, (4.8) pencycuron, (4.9) thiabendazole, (4.10) thiophanate-methyl, (4.11) thiophanate, (4.12) zoxamide, (4.13) 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine, (4.14) 3-chloro-5-(6-chloropyridin-3-yl)-6-methyl-4-(2,4,6-trifluorophenyl)pyridazine.
- 5) Compounds capable to have a multisite action, for example (5.1) bordeaux mixture, (5.2) captafol, (5.3) captan, (5.4) chlorothalonil, (5.5) copper hydroxide, (5.6) copper naphthenate, (5.7) copper oxide, (5.8) copper oxychloride, (5.9) copper(2+) sulfate, (5.10) dichlofluanid, (5.11) dithianon, (5.12) dodine, (5.13) dodine free base, (5.14) ferbam, (5.15) fluorofolpet, (5.16) folpet, (5.17) guazatine, (5.18) guazatine acetate, (5.19) iminocadine, (5.20) iminocadine albesilate, (5.21) iminocadine triacetate, (5.22) mancozeb, (5.23) mancozeb, (5.24) maneb, (5.25) metiram, (5.26) metiram zinc, (5.27) oxine-copper, (5.28) propamidine, (5.29) propineb, (5.30) sulfur and sulfur preparations including calcium polysulfide, (5.31) thiram, (5.32) tolylfluanid, (5.33) zineb, (5.34) ziram, (5.35) anilazine.
- 6) Compounds capable to induce a host defence, for example (6.1) acibenzolar-S-methyl, (6.2) isotianil, (6.3) probenazole, (6.4) tiadinil, (6.5) laminarin.
- 7) Inhibitors of the amino acid and/or protein biosynthesis, for example (7.1) andoprim, (7.2) blasticidin-S, (7.3) cyprodinil, (7.4) kasugamycin, (7.5) kasugamycin hydrochloride hydrate, (7.6) mepanipyrim, (7.7) pyrimethanil, (7.8) 3-(5-fluoro-3,3,4,4-tetramethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (7.9) oxytetracycline, (7.10) streptomycin.
- 8) Inhibitors of the ATP production, for example (8.1) fentin acetate, (8.2) fentin chloride, (8.3) fentin hydroxide, (8.4) silthiofam.
- 9) Inhibitors of the cell wall synthesis, for example (9.1) benthiavalicarb, (9.2) dimethomorph, (9.3) flumorph, (9.4) iprovalicarb, (9.5) mandipropamid, (9.6) polyoxins, (9.7) polyoxorim, (9.8) validamycin A, (9.9) valifenalate, (9.10) polyoxin B.
- 10) Inhibitors of the lipid and membrane synthesis, for example (10.1) biphenyl, (10.2) chloroneb, (10.3) dicloran, (10.4) edifenphos, (10.5) etridiazole, (10.6) iodocarb, (10.7) iprobenfos, (10.8) isoprothiolane, (10.9) propamocarb, (10.10) propamocarb hydrochloride, (10.11) prothiocarb, (10.12) pyrazophos, (10.13) quintozone, (10.14) tecnazene, (10.15) tolclofos-methyl.
- 11) Inhibitors of the melanin biosynthesis, for example (11.1) carpropamid, (11.2) diclocymet, (11.3) fenoxanil, (11.4) phthalide, (11.5) pyroquilon, (11.6) tricyclazole, (11.7) 2,2,2-trifluoroethyl {3-methyl-1-[(4-methylbenzoyl)amino]butan-2-yl} carbamate.
- 12) Inhibitors of the nucleic acid synthesis, for example (12.1) benalaxyl, (12.2) benalaxyl-M (kiralaxyl), (12.3) bupirimate, (12.4) clozylacon, (12.5) dimethirimol, (12.6) ethirimol, (12.7) furalaxyl, (12.8) hymexazol, (12.9)

metalaxyl, (12.10) metalaxyl-M (mefenoxam), (12.11) ofurace, (12.12) oxadixyl, (12.13) oxolinic acid, (12.14) octhilinone.

13) Inhibitors of the signal transduction, for example (13.1) chlozolate, (13.2) fenpiclonil, (13.3) fludioxonil, (13.4) iprodione, (13.5) procymidone, (13.6) quinoxifen, (13.7) vinclozolin, (13.8) proquinazid.

5 14) Compounds capable to act as an uncoupler, for example (14.1) binapacryl, (14.2) dinocap, (14.3) ferimzone, (14.4) fluazinam, (14.5) meptyldinocap.

15) Further compounds, for example (15.1) benthiazole, (15.2) bethoxazin, (15.3) capsimycin, (15.4) carvone, (15.5) chinomethionat, (15.6) pyriofenone (chlazafenone), (15.7) cufraneb, (15.8) cyflufenamid, (15.9) cy-moxanil, (15.10) cyprosulfamide, (15.11) dazomet, (15.12) debacarb, (15.13) dichlorophen, (15.14) di-  
 10 clomezine, (15.15) difenzoquat, (15.16) difenzoquat metilsulfate, (15.17) diphenylamine, (15.18) ecomate, (15.19) fenpyrazamine, (15.20) flumetover, (15.21) fluoroimide, (15.22) flusulfamide, (15.23) flutianil, (15.24) fosetyl-aluminium, (15.25) fosetyl-calcium, (15.26) fosetyl-sodium, (15.27) hexachlorobenzene, (15.28) iru-mamycin, (15.29) methasulfocarb, (15.30) methyl isothiocyanate, (15.31) metrafenone, (15.32) mildiomyacin, (15.33) natamycin, (15.34) nickel dimethyldithiocarbamate, (15.35) nitrothal-isopropyl, (15.37) oxamocarb,  
 15 (15.38) oxyfenthiiin, (15.39) pentachlorophenol and salts, (15.40) phenothrin, (15.41) phosphorous acid and its salts, (15.42) propamocarb-fosetilate, (15.43) propanosine-sodium, (15.44) pyrimorph, (15.45) (2E)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, (15.46) (2Z)-3-(4-tert-butylphenyl)-3-(2-chloropyridin-4-yl)-1-(morpholin-4-yl)prop-2-en-1-one, (15.47) pyrrolnitrine, (15.48) tebufloquin, (15.49) tecloftalam, (15.50) tolnifanide, (15.51) triazoxide, (15.52) trichlamide, (15.53) zarilamid, (15.54)  
 20 (3S,6S,7R,8R)-8-benzyl-3-[(3-[(isobutyryloxy)methoxy]-4-methoxypyridin-2-yl)carbonylamino]-6-methyl-4,9-dioxo-1,5-dioxonan-7-yl 2-methylpropanoate, (15.55) 1-(4-{4-[(5R)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.56) 1-(4-{4-[(5S)-5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone, (15.57) 1-(4-{4-[5-(2,6-difluorophenyl)-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)-2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]ethanone,  
 25 (15.58) 1-(4-methoxyphenoxy)-3,3-dimethylbutan-2-yl 1H-imidazole-1-carboxylate, (15.59) 2,3,5,6-tetrachloro-4-(methylsulfonyl)pyridine, (15.60) 2,3-dibutyl-6-chlorothieno[2,3-d]pyrimidin-4(3H)-one, (15.61) 2,6-dimethyl-1H,5H-[1,4]dithiino[2,3-c:5,6-c']dipyrrole-1,3,5,7(2H,6H)-tetrone, (15.62) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5R)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.63) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-(4-{4-[(5S)-5-phenyl-4,5-dihydro-1,2-oxazol-3-yl]-1,3-thiazol-2-yl}piperidin-1-yl)ethanone, (15.64) 2-[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]-1-{4-[4-(5-phenyl-4,5-dihydro-1,2-oxazol-3-yl)-1,3-thiazol-2-yl]piperidin-1-yl}ethanone,  
 30 (15.65) 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, (15.66) 2-chloro-5-[2-chloro-1-(2,6-difluoro-4-methoxyphenyl)-4-methyl-1H-imidazol-5-yl]pyridine, (15.67) 2-phenylphenol and salts, (15.68) 3-(4,4,5-trifluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (15.69) 3,4,5-trichloropyridine-2,6-dicarbonitrile, (15.70) 3-chloro-5-(4-chlorophenyl)-4-(2,6-difluorophenyl)-6-methylpyridazine, (15.71) 4-(4-chlorophenyl)-5-(2,6-difluorophenyl)-3,6-dimethylpyridazine, (15.72) 5-amino-1,3,4-thiadiazole-2-thiol, (15.73) 5-chloro-N'-

- phenyl-N'-(prop-2-yn-1-yl)thiophene-2-sulfonohydrazide, (15.74) 5-fluoro-2-[(4-fluorobenzyl)oxy]pyrimidin-4-amine, (15.75) 5-fluoro-2-[(4-methylbenzyl)oxy]pyrimidin-4-amine, (15.76) 5-methyl-6-octyl[1,2,4]triazolo[1,5-a]pyrimidin-7-amine, (15.77) ethyl (2Z)-3-amino-2-cyano-3-phenylacrylate, (15.78) N'-(4-{[3-(4-chlorobenzyl)-1,2,4-thiadiazol-5-yl]oxy}-2,5-dimethylphenyl)-N-ethyl-N-methylimidoforamide, (15.79) N-(4-chlorobenzyl)-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (15.80) N-[(4-chlorophenyl)(cyano)methyl]-3-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]propanamide, (15.81) N-[(5-bromo-3-chloropyridin-2-yl)methyl]-2,4-dichloronicotinamide, (15.82) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloronicotinamide, (15.83) N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-iodonicotinamide, (15.84) N-{(E)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide, (15.85) N-{(Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide, (15.86) N'-{4-[(3-tert-butyl-4-cyano-1,2-thiazol-5-yl)oxy]-2-chloro-5-methylphenyl}-N-ethyl-N-methylimidoforamide, (15.87) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-(1,2,3,4-tetrahydronaphthalen-1-yl)-1,3-thiazole-4-carboxamide, (15.88) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1R)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide, (15.89) N-methyl-2-(1-{[5-methyl-3-(trifluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-N-[(1S)-1,2,3,4-tetrahydronaphthalen-1-yl]-1,3-thiazole-4-carboxamide, (15.90) pentyl {6-[[[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino]oxy)methyl]pyridin-2-yl} carbamate, (15.91) phenazine-1-carboxylic acid, (15.92) quinolin-8-ol, (15.93) quinolin-8-ol sulfate (2:1), (15.94) tert-butyl {6-[[[(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino]oxy)methyl]pyridin-2-yl} carbamate, (15.95) 1-methyl-3-(trifluoromethyl)-N-[2'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (15.96) N-(4'-chlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (15.97) N-(2',4'-dichlorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, (15.98) 3-(difluoromethyl)-1-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (15.99) N-(2',5'-difluorobiphenyl-2-yl)-1-methyl-3-(trifluoromethyl)-1H-pyrazole-4-carboxamide, (15.100) 3-(difluoromethyl)-1-methyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (15.101) 5-fluoro-1,3-dimethyl-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]-1H-pyrazole-4-carboxamide, (15.102) 2-chloro-N-[4'-(prop-1-yn-1-yl)biphenyl-2-yl]nicotinamide, (15.103) 3-(difluoromethyl)-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, (15.104) N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, (15.105) 3-(difluoromethyl)-N-(4'-ethynylbiphenyl-2-yl)-1-methyl-1H-pyrazole-4-carboxamide, (15.106) N-(4'-ethynylbiphenyl-2-yl)-5-fluoro-1,3-dimethyl-1H-pyrazole-4-carboxamide, (15.107) 2-chloro-N-(4'-ethynylbiphenyl-2-yl)nicotinamide, (15.108) 2-chloro-N-[4'-(3,3-dimethylbut-1-yn-1-yl)biphenyl-2-yl]nicotinamide, (15.109) 4-(difluoromethyl)-2-methyl-N-[4'-(trifluoromethyl)biphenyl-2-yl]-1,3-thiazole-5-carboxamide, (15.110) 5-fluoro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, (15.111) 2-chloro-N-[4'-(3-hydroxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]nicotinamide, (15.112) 3-(difluoromethyl)-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1-methyl-1H-pyrazole-4-carboxamide, (15.113) 5-fluoro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]-1,3-dimethyl-1H-pyrazole-4-carboxamide, (15.114) 2-chloro-N-[4'-(3-methoxy-3-methylbut-1-yn-1-yl)biphenyl-2-yl]nicotinamide, (15.115) (5-bromo-2-methoxy-4-methylpyridin-3-yl)(2,3,4-trimethoxy-6-methylphenyl)methanone, (15.116) N-[2-(4-{[3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-



- methoxyphenyl)ethyl]-N2-(methylsulfonyl)valinamide, (15.117) 4-oxo-4-[(2-phenylethyl)amino]butanoic acid, (15.118) but-3-yn-1-yl {6-[(Z)-(1-methyl-1H-tetrazol-5-yl)(phenyl)methylene]amino}oxy)methyl]pyridin-2-yl} carbamate, (15.119) 4-amino-5-fluoropyrimidin-2-ol (tautomeric form: 4-amino-5-fluoropyrimidin-2(1H)-one), (15.120) propyl 3,4,5-trihydroxybenzoate, (15.121) 1,3-dimethyl-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1H-pyrazole-4-carboxamide, (15.122) 1,3-dimethyl-N-[(3R)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (15.123) 1,3-dimethyl-N-[(3S)-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1H-pyrazole-4-carboxamide, (15.124) [3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (15.125) (S)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (15.126) (R)-[3-(4-chloro-2-fluorophenyl)-5-(2,4-difluorophenyl)-1,2-oxazol-4-yl](pyridin-3-yl)methanol, (15.127) 2-{[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.128) 1-{[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-1H-1,2,4-triazol-5-yl thiocyanate, (15.129) 5-(allylsulfanyl)-1-{[3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-1H-1,2,4-triazole, (15.130) 2-[1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.131) 2-{[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.132) 2-{[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.133) 1-{[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-1H-1,2,4-triazol-5-yl thiocyanate, (15.134) 1-{[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-1H-1,2,4-triazol-5-yl thiocyanate, (15.135) 5-(allylsulfanyl)-1-{[rel(2R,3S)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-1H-1,2,4-triazole, (15.136) 5-(allylsulfanyl)-1-{[rel(2R,3R)-3-(2-chlorophenyl)-2-(2,4-difluorophenyl)oxiran-2-yl]methyl}-1H-1,2,4-triazole, (15.137) 2-[(2S,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.138) 2-[(2R,4S,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.139) 2-[(2R,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.140) 2-[(2S,4R,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.141) 2-[(2S,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.142) 2-[(2R,4S,5R)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.143) 2-[(2R,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.144) 2-[(2S,4R,5S)-1-(2,4-dichlorophenyl)-5-hydroxy-2,6,6-trimethylheptan-4-yl]-2,4-dihydro-3H-1,2,4-triazole-3-thione, (15.145) 2-fluoro-6-(trifluoromethyl)-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)benzamide, (15.146) 2-(6-benzylpyridin-2-yl)quinazoline, (15.147) 2-[6-(3-fluoro-4-methoxyphenyl)-5-methylpyridin-2-yl]quinazoline, (15.148) 3-(4,4-difluoro-3,3-dimethyl-3,4-dihydroisoquinolin-1-yl)quinoline, (15.149) Absciscic acid, (15.150) 3-(difluoromethyl)-N-methoxy-1-methyl-N-[1-(2,4,6-trichlorophenyl)propan-2-yl]-1H-pyrazole-4-carboxamide, (15.151) N'-[5-bromo-6-(2,3-dihydro-1H-inden-2-yloxy)-2-methylpyridin-3-yl]-N-ethyl-N-methylimidofornamide, (15.152) N'-{5-bromo-6-[1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidofornamide, (15.153) N'-{5-bromo-6-[(1R)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidofornamide, (15.154) N'-{5-bromo-6-[(1S)-1-(3,5-difluorophenyl)ethoxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidofornamide, (15.155) N'-{5-bromo-6-

[(cis-4-isopropylcyclohexyl)oxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide, (15.156) N'-{5-bromo-6-[(trans-4-isopropylcyclohexyl)oxy]-2-methylpyridin-3-yl}-N-ethyl-N-methylimidoforamide, (15.157) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (15.158) N-cyclopropyl-N-(2-cyclopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.159) N-(2-tert-butylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.160) N-(5-chloro-2-ethylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.161) N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.162) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-fluorobenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.163) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(5-fluoro-2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (15.164) N-cyclopropyl-N-(2-cyclopropyl-5-fluorobenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.165) N-(2-cyclopentyl-5-fluorobenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.166) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-fluoro-6-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (15.167) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-5-methylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.168) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropyl-5-methylbenzyl)-1-methyl-1H-pyrazole-4-carboxamide, (15.169) N-cyclopropyl-N-(2-cyclopropyl-5-methylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.170) N-(2-tert-butyl-5-methylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.171) N-[5-chloro-2-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.172) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[5-methyl-2-(trifluoromethyl)benzyl]-1H-pyrazole-4-carboxamide, (15.173) N-[2-chloro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.174) N-[3-chloro-2-fluoro-6-(trifluoromethyl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.175) N-cyclopropyl-3-(difluoromethyl)-N-(2-ethyl-4,5-dimethylbenzyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide, (15.176) N-cyclopropyl-3-(difluoromethyl)-5-fluoro-N-(2-isopropylbenzyl)-1-methyl-1H-pyrazole-4-carbothioamide, (15.177) 3-(difluoromethyl)-N-(7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)-1-methyl-1H-pyrazole-4-carboxamide, (15.178) 3-(difluoromethyl)-N-[(3R)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide, (15.179) 3-(difluoromethyl)-N-[(3S)-7-fluoro-1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl]-1-methyl-1H-pyrazole-4-carboxamide.

All named mixing partners of the classes (1) to (15) can, if their functional groups enable this, optionally form salts with suitable bases or acids.

Although the composition according to the present invention comprises compounds of the formula (I) and at least a biological control agent, the final used composition is usually formulated to be suitable for agrochemical application e.g. by mixing the compounds of the formula (I) with the biological control agent and an additional component such as suitable extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost protectants, thickeners, adjuvants or the like. Those compositions are referred to as formulations.

Accordingly, in one aspect of the present invention such formulations, and application forms prepared from them, are provided as crop protection agents and/or pesticidal agents, such as drench, drip and spray liquors, comprising the composition of the invention. The application forms may comprise activity-enhancing adjuvants such as penetrants, examples being vegetable oils such as, for example, rapeseed oil, sunflower oil, mineral oils such as, for example, liquid paraffins, alkyl esters of vegetable fatty acids, such as rapeseed oil or soybean oil methyl esters, or alkanol alkoxylates, and/or spreaders such as, for example, alkylsiloxanes and/or salts, examples being organic or inorganic ammonium or phosphonium salts, examples being ammonium sulphate or diammonium hydrogen phosphate, and/or retention promoters such as dioctyl sulphosuccinate or hydroxypropylguar polymers and/or humectants such as glycerol and/or fertilizers such as ammonium, potassium or phosphorous fertilizers, for example.

- 5 Examples of typical formulations include water-soluble liquids (SL), emulsifiable concentrates (EC), emulsions in water (EW), suspension concentrates (SC, SE, FS, OD), water-dispersible granules (WG), granules (GR) and capsule concentrates (CS); these and other possible types of formulation are described, for example, by Crop Life International and in Pesticide Specifications, Manual on development and use of FAO and WHO specifications for pesticides, FAO Plant Production and Protection Papers – 173, prepared by the FAO/WHO Joint Meeting on Pesticide Specifications, 2004, ISBN: 9251048576. The formulations may comprise active agrochemical compounds other than one or more active compounds of the invention.

- The formulations or application forms in question preferably comprise auxiliaries, such as extenders, solvents, spontaneity promoters, carriers, emulsifiers, dispersants, frost protectants, biocides, thickeners and/or other auxiliaries, such as adjuvants, for example. An adjuvant in this context is a component which enhances the biological effect of the formulation, without the component itself having a biological effect. Examples of adjuvants are agents which promote the retention, spreading, attachment to the leaf surface, or penetration.

These formulations are produced in a known manner, for example by mixing the active compounds with auxiliaries such as, for example, extenders, solvents and/or solid carriers and/or further auxiliaries, such as, for example, surfactants. The formulations are prepared either in suitable plants or else before or during the application.

- 25 Suitable for use as auxiliaries are substances which are suitable for imparting to the formulation of the active compound or the application forms prepared from these formulations (such as, e.g., usable crop protection agents, such as spray liquors or seed dressings) particular properties such as certain physical, technical and/or biological properties.

- Suitable extenders are, for example, water, polar and nonpolar organic chemical liquids, for example from the classes of the aromatic and non-aromatic hydrocarbons (such as paraffins, alkylbenzenes, alkylnaphthalenes, chlorobenzenes), the alcohols and polyols (which, if appropriate, may also be substituted, etherified and/or esterified), the ketones (such as acetone, cyclohexanone), esters (including fats and oils) and (poly)ethers, the unsubstituted and substituted amines, amides, lactams (such as N-alkylpyrrolidones) and lactones, the sulphones and sulfoxides (such as dimethyl sulfoxide).

If the extender used is water, it is also possible to employ, for example, organic solvents as auxiliary solvents. Essentially, suitable liquid solvents are: aromatics such as xylene, toluene or alkylnaphthalenes, chlorinated aromatics and chlorinated aliphatic hydrocarbons such as chlorobenzenes, chloroethylenes or methylene chloride, aliphatic hydrocarbons such as cyclohexane or paraffins, for example petroleum fractions, mineral and vegetable oils, alcohols such as butanol or glycol and also their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents such as dimethylformamide and dimethyl sulphoxide, and also water.

In principle it is possible to use all suitable solvents. Suitable solvents are, for example, aromatic hydrocarbons, such as xylene, toluene or alkylnaphthalenes, for example, chlorinated aromatic or aliphatic hydrocarbons, such as chlorobenzene, chloroethylene or methylene chloride, for example, aliphatic hydrocarbons, such as cyclohexane, for example, paraffins, petroleum fractions, mineral and vegetable oils, alcohols, such as methanol, ethanol, isopropanol, butanol or glycol, for example, and also their ethers and esters, ketones such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, for example, strongly polar solvents, such as dimethyl sulphoxide, and water.

All suitable carriers may in principle be used. Suitable carriers are in particular for example, ammonium salts and ground natural minerals such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as finely divided silica, alumina and natural or synthetic silicates, resins, waxes and/or solid fertilizers. Mixtures of such carriers may likewise be used. Carriers suitable for granules include the following: for example, crushed and fractionated natural minerals such as calcite, marble, pumice, sepiolite, dolomite, and also synthetic granules of inorganic and organic meals, and also granules of organic material such as sawdust, paper, coconut shells, maize cobs and tobacco stalks.

Liquefied gaseous extenders or solvents may also be used. Particularly suitable are those extenders or carriers which at standard temperature and under standard pressure are gaseous, examples being aerosol propellants, such as halogenated hydrocarbons, and also butane, propane, nitrogen and carbon dioxide.

Examples of emulsifiers and/or foam-formers, dispersants or wetting agents having ionic or nonionic properties, or mixtures of these surface-active substances, are salts of polyacrylic acid, salts of lignosulphonic acid, salts of phenolsulphonic acid or naphthalenesulphonic acid, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, with substituted phenols (preferably alkylphenols or arylphenols), salts of sulphosuccinic esters, taurine derivatives (preferably alkyltaurates), phosphoric esters of polyethoxylated alcohols or phenols, fatty acid esters of polyols, and derivatives of the compounds containing sulphates, sulphonates and phosphates, examples being alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates, protein hydrolysates, lignin-sulphite waste liquors and methylcellulose. The presence of a surface-active substance is advantageous if one of the active compounds and/or one of the inert carriers is not soluble in water and if application takes place in water.

Further auxiliaries that may be present in the formulations and in the application forms derived from them include colorants such as inorganic pigments, examples being iron oxide, titanium oxide, Prussian Blue, and organic dyes,

such as alizarin dyes, azo dyes and metal phthalocyanine dyes, and nutrients and trace nutrients, such as salts of iron, manganese, boron, copper, cobalt, molybdenum and zinc.

Stabilizers, such as low-temperature stabilizers, preservatives, antioxidants, light stabilizers or other agents which improve chemical and/or physical stability may also be present. Additionally present may be foam-formers or defoamers.

Furthermore, the formulations and application forms derived from them may also comprise, as additional auxiliaries, stickers such as carboxymethylcellulose, natural and synthetic polymers in powder, granule or latex form, such as gum arabic, polyvinyl alcohol, polyvinyl acetate, and also natural phospholipids, such as cephalins and lecithins, and synthetic phospholipids. Further possible auxiliaries include mineral and vegetable oils.

There may possibly be further auxiliaries present in the formulations and the application forms derived from them. Examples of such additives include fragrances, protective colloids, binders, adhesives, thickeners, thixotropic substances, penetrants, retention promoters, stabilizers, sequestrants, complexing agents, humectants and spreaders. Generally speaking, the active compounds may be combined with any solid or liquid additive commonly used for formulation purposes.

Suitable retention promoters include all those substances which reduce the dynamic surface tension, such as dioctyl sulphosuccinate, or increase the viscoelasticity, such as hydroxypropylguar polymers, for example.

Suitable penetrants in the present context include all those substances which are typically used in order to enhance the penetration of active agrochemical compounds into plants. Penetrants in this context are defined in that, from the (generally aqueous) application liquor and/or from the spray coating, they are able to penetrate the cuticle of the plant and thereby increase the mobility of the active compounds in the cuticle. This property can be determined using the method described in the literature (Baur et al., 1997, Pesticide Science 51, 131-152). Examples include alcohol alkoxylates such as coconut fatty ethoxylate (10) or isotridecyl ethoxylate (12), fatty acid esters such as rapeseed or soybean oil methyl esters, fatty amine alkoxylates such as tallowamine ethoxylate (15), or ammonium and/or phosphonium salts such as ammonium sulphate or diammonium hydrogen phosphate, for example.

The formulations preferably comprise between 0.00000001% and 98% by weight of active compound or, with particular preference, between 0.01% and 95% by weight of active compound, more preferably between 0.5% and 90% by weight of active compound, based on the weight of the formulation. The content of the active compound is defined as the sum of the compound of formula (I) and the at least one biological control agent.

The active compound content of the application forms (crop protection products) prepared from the formulations may vary within wide ranges. The active compound concentration of the application forms may be situated typically between 0.00000001% and 95% by weight of active compound, preferably between 0.00001% and 1% by weight, based on the weight of the application form. Application takes place in a customary manner adapted to the application forms.

The inventive compositions may be present as such or in their (commercial) formulations and in the use forms prepared from these formulations as a mixture with other (known) active ingredients, such as attractants, sterilants, bactericides, acaricides, nematocides, fungicides, insecticides, growth regulators, herbicides, fertilizers, safeners and/or semiochemicals. The (commercial) formulation comprising a compound of the formula (I) and a biological control agent and the agriculturally suitable components as described above, can be used as a pesticide. In a specific embodiment of the invention the (commercial) formulation as defined above can be used as a fungicide. In another specific embodiment of the invention the (commercial) formulation as defined above acts in a synergistic fashion.

According to the present invention, in the composition according to the invention, the compound ratio (I)/B may be advantageously chosen so as to produce a synergistic effect. The term “synergistic effect” as well as the term “acting in a synergistic fashion” is understood to mean in particular that defined by Colby in an article entitled "Calculation of the synergistic and antagonistic responses of herbicide combinations" Weeds, (1967), 15, pages 20-22.

The latter article mentions the formula:

$$E = X + Y - \frac{XY}{100}$$

wherein E represents the expected percentage of inhibition of the pest for the combination of the two compounds at defined doses (for example equal to x and y respectively), X is the percentage of inhibition observed for the pest by compound (A) at a defined dose (equal to x), Y is the percentage of inhibition observed for the pest by compound (B) at a defined dose (equal to y). When the percentage of inhibition observed for the combination is greater than E, there is a synergistic effect.

The term “synergistic effect” also means the effect defined by application of the Tammes method, “Isoboles, a graphic representation of synergism in pesticides”, Netherlands Journal of Plant Pathology, 70(1964), pages 73-80.

A synergistic effect in fungicides is always present when the fungicidal action of the composition exceeds the expected action of the active compounds applied individually.

The expected fungicidal action for a given combination of two or three active compounds can be calculated as follows, according to S.R. Colby (“Calculating Synergistic and Antagonistic Responses of Herbicide Combinations”, Weeds 1967, 15, 20-22):

If

X is the *efficacy* when employing active compound A at an application rate of  $\underline{m}$  g/ha,

Y is the *efficacy* when employing active compound B at an application rate of  $\underline{n}$  g/ha and

E is the *efficacy* when employing active compounds A and B at application rates of  $\underline{m}$  and  $\underline{n}$  g/ha,

then  $E = X + Y - (X*Y)/100$  .

Here, the efficacy is determined in %. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed. If the actual fungicidal action exceeds the calculated value, the action of the combination is superadditive, i.e. a synergistic effect is present. In this case, the actually observed efficacy must exceed the value calculated using the above formula for the expected efficacy (E).

The composition according to the present invention can be combined in any specific ratio between the two mandatory components. In the compositions according to the invention the compounds of the general formula (I) and the biological control agent (B) as defined above are present in a synergistically effective weight ratio of (I) : (B) in a range of 1:5000 to 1:1, preferably in a weight ratio of 1:1000 to 1:1, most preferably in a weight ratio of 1:500 to 1:100. Further ratios of (I) : (B) which can be used according to the present invention with increasing preferences the order given are: 1:5000 to 1:1, 1:4000 to 1:1, 1:3000 to 1:1, 1:2000 to 1:1, 1:1000 to 1:1, 1:900 to 1:1, 1:800 to 1:1, 1:700 to 1:1, 1:600 to 1:1, 1:500 to 1:1, 1:400 to 1:1, 1:300 to 1:1, 1:200 to 1:1, 1:100 to 1:1, 1:50 to 1:1, 1:1000 to 1:2, 1:1000 to 1:5, 1:500 to 1:2, 1:500 to 1:5, 1:500 to 1:10, 1:500 to 1:50, 1:500 to 1:100, 1:500 to 1:250, 1:500 to 1:300, 1:500 to 1:350, 1:500 to 1:400, 1:500 to 1:450.

In a preferred embodiment the biological control agent e.g. their spores are present in a solo-formulation or the combined formulation in a concentration of at least  $10^5$  colony forming units per gram preparation (e. g. cells/g preparation, spores/g preparation), such as  $10^5$  -  $10^{12}$  cfu/g, preferably  $10^6$  -  $10^{11}$  cfu/g, more preferably  $10^7$  -  $10^{10}$  cfu/g and most preferably  $10^9$  -  $10^{10}$  cfu/g at the time point of applying biological control agents on a plant or plant parts such as seeds, fruits or vegetables. Also references to the concentration of biological control agents in form of, e.g., spores or cells - when discussing ratios between the amount of a preparation of at least one biological control agent and the amount of the compound of the formula (I)- are made in view of the time point when the biological control agent is applied on a plant or plant parts such as seeds, fruits or vegetables.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and / or a widening of the activity spectrum and / or an increase in the activity of the compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, bigger fruits, larger plant height, greener leaf color, earlier flowering, higher quality and / or a higher nutritional value of the harvested products, higher sugar concentration within the fruits, better storage stability and / or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

At certain application rates, the compositions according to the invention may also have a strengthening effect in plants. Accordingly, they are also suitable for mobilizing the defense system of the plant against attack by phytopathogenic fungi. This may, if appropriate, be one of the reasons of the enhanced activity of the compositions

according to the invention, for example against fungi, in particular phytopathogenic fungi. "Plant-strengthening" (resistance-inducing) substances are to be understood as meaning, in the present context, those substances or combinations of substances which are capable of stimulating the defense system of plants in such a way that, when subsequently inoculated with harmful microorganisms, the treated plants display a substantial degree of resistance to these microorganisms. In a specific embodiment of the invention, harmful microorganisms are phytopathogenic fungi. Thus, the compositions according to the invention can be employed for protecting plants against attack by the abovementioned pathogens within a certain period of time after the treatment. The period of time within which protection is effected generally extends from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the inventive composition.

In a specific embodiment, the invention comprises a method for reducing damage of plants and plant parts or losses in harvested fruits or vegetables caused by phytopathogenic fungi by controlling such phytopathogenic fungi, comprising applying the composition to the plant or the phytopathogenic fungi or the habitat of the plant or the habitat of the phytopathogenic fungi. Accordingly, the present invention also relates to compositions for controlling phytopathogenic fungi, comprising an effective but non-phytotoxic amount of the inventive compositions. These are preferably fungicidal compositions which comprise agriculturally suitable components as described above.

By "plants" is meant all plants and plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering methods. Plants also comprise plant parts.

By "plant parts" is meant all above ground and below ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed as well as roots, corms and rhizomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners and seeds also belong to plant parts.

In this context the term "damage of plants" or "damage of plant parts" comprises e.g. the decrease of plant growth or yield or the decrease of plant vigor, comprising plant health, plant quality and seed vigor, a decreased resistance to harmful microorganisms such as phytopathogenic fungi, a decreased abiotic stress tolerance, comprising temperature tolerance, drought tolerance and recovery after drought stress, water use efficiency (correlating to reduced water consumption), flood tolerance, ozone stress and UV tolerance, tolerance towards chemicals like heavy metals, salts, pesticides (safener) or the like, an increased stand failure, decreased recovery, impaired greening effect and decreased photosynthetic efficiency. Reducing the damage of plants and plant parts often results in healthier plants and/or in an increase in plant vigor and yield. The compositions according to the invention, in combination with good plant tolerance and favourable toxicity to warm-blooded animals and being tolerated well by the environment, are suitable for protecting plants and plant parts, for increasing harvest yields and for improving the quality of the harvested material.



An “effective but non-phytotoxic amount” means an amount of the inventive composition which is sufficient to control the fungal disease of the plant in a satisfactory manner or to eradicate the fungal disease completely, and which, at the same time, does not cause any significant symptoms of phytotoxicity. In general, this application rate may vary within a relatively wide range. It depends on several factors, for example on the fungus to be controlled, the plant, the climatic conditions and the ingredients of the inventive compositions.

In the context of the present invention, “control of phytopathogenic fungi” means a reduction in infestation by phytopathogenic fungi, compared with the untreated plant measured as fungicidal efficacy, preferably a reduction by 25-50 %, compared with the untreated plant (100 %), more preferably a reduction by 40-79 %, compared with the untreated plant (100 %); even more preferably, the infection by phytopathogenic fungi is entirely suppressed (by 70-100 %). The control may be curative, i.e. for treatment of already infected plants, or protective, for protection of plants which have not yet been infected. In a specific embodiment the compositions according to the invention can be used for curative or protective or preventive control of phytopathogenic fungi. The invention therefore also relates to curative and protective methods for controlling phytopathogenic fungi by the use of the inventive compositions. The compositions according to the invention are active against normally sensitive and resistant species of phytopathogenic fungi and against all or some stages of development. The term “preventive control of phytopathogenic fungi” comprises the use of compositions according to the invention for the prevention of diseases. Thus, a further aspect of the present invention is the use of compositions according to the invention for controlling diseases such as phytopathogenic fungi, e.g., in agriculture, in horticulture, in forests, in gardens and leisure facilities as well as in the protection of stored products and materials.

The term “applying the composition to the plant or the phytopathogenic fungi or the habitat of the plant or the habitat of the phytopathogenic fungi” refers to the treatment of a plant and / or phytopathogenic fungi and / or the habitat of the plant or the phytopathogenic fungi with the compositions according to the invention. The treatment is effected directly or by action on their surroundings, habitat or storage space by the customary treatment methods, for example by dipping, spraying, atomizing, irrigating, evaporating, dusting, fogging, broadcasting, foaming, painting, spreading-on, watering (drenching) and drip irrigating. It is also possible to deploy the compositions by the ultra-low volume method or to inject the composition preparation or the compositions itself into the soil.

When using the inventive compositions or the (commercial) formulation comprising the inventive composition and the agriculturally suitable components as described above as fungicides, the application rates can be varied within a relatively wide range, depending on the kind of application. The application rate of the compositions is

- in the case of treatment of a plant or plant parts, for example leaves: from 0.1 to 10 000 g/ha, preferably from 10 to 1000 g/ha, more preferably from 10 to 800 g/ha, even more preferably from 50 to 300 g/ha (in the case of application by watering or dripping, it is even possible to reduce the application rate, especially when inert substrates such as rockwool or perlite are used);
- in the case of soil treatment: from 0.1 to 10 000 g/ha, preferably from 1 to 5000 g/ha.

These application rates are merely by way of example and are not limiting for the purposes of the invention.

The inventive compositions can thus be used to protect plants from attack by the pathogens mentioned below for a certain period of time after treatment. The period for which protection is provided extends generally for 1 to 28 days, preferably for 1 to 14 days, more preferably for 1 to 10 days, most preferably for 1 to 7 days, after the treatment of the plant or the plant parts with the inventive compositions.

The method of treatment according to the invention also provides the use or application of compounds according to formula (I) and one biological control agent as defined above in a simultaneous, separate or sequential manner. If the single active ingredients are applied in a sequential manner, i.e. at different times, they are applied one after the other within a reasonably short period, such as a few hours or days. Preferably the order of applying the compounds according to formula (I) and the biological control agent as defined above is not essential for working the present invention.

The plants listed further down in the description can particularly advantageously be treated in accordance with the compositions according to the invention. The preferred ranges stated above for the compositions also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the compositions specifically mentioned in the present text.

The fact that the compositions are well tolerated by plants at the concentrations required for controlling phytopathogenic fungi allows the treatment of above-ground parts of plants, of propagation stock and seeds, and of the soil.

The "habitat of the plant" comprises the environment where the plant is growing, e.g. the soil or nutrition medium - which is in a radius of at least 10 cm, 20 cm, 30 cm around the caulis or bole of a plant to be treated or which is at least 10 cm, 20 cm, 30 cm around the root system of said plant to be treated, respectively. The term "habitat of the phytopathogenic fungi" as used herein is defined as the seed, the plant or plant parts or the fruit. It also comprises the soil or the nutrition medium in which the plant grows - which is in a radius of at least 10 cm, 20 cm, 30 cm around the caulis or bole of a plant to be treated or which is at least 10 cm, 20 cm, 30 cm around the root system of said plant to be treated, respectively.

## **Plants**

According to the invention all plants and plant parts can be treated. By plants is meant all plants and plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering methods. By plant parts is meant all above ground and below ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed as well as roots, corms and rhi-

zomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners and seeds also belong to plant parts.

The inventive compositions, when they are well tolerated by plants, have favourable homeotherm toxicity and are well tolerated by the environment, are suitable for protecting plants and plant organs, for enhancing harvest yields, for improving the quality of the harvested material. They can preferably be used as crop protection compositions. They are active against normally sensitive and resistant species and against all or some stages of development.

Plants which can be treated in accordance with the invention include the following main crop plants including fruits and vegetables: maize, soya bean, alfalfa, cotton, sunflower, *Brassica* oil seeds such as *Brassica napus* (e.g. canola, rapeseed), *Brassica rapa*, *B. juncea* (e.g. (field) mustard) and *Brassica carinata*, *Arecaceae* sp. (e.g. oil-palm, coconut), rice, wheat, sugar beet, sugar cane, oats, rye, barley, millet and sorghum, triticale, flax, nuts, grapes and vine and various fruit and vegetables from various botanic taxa, e.g. *Rosaceae* sp. (e.g. pome fruits such as apples and pears, but also stone fruits such as apricots, cherries, almonds, plums and peaches, and berry fruits such as strawberries, raspberries, red and black currant and gooseberry), *Ribesiodae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp. (e.g. olive tree), *Actinidaceae* sp., *Lauraceae* sp. (e.g. avocado, cinnamon, camphor), *Musaceae* sp. (e.g. banana trees and plantations), *Rubiaceae* sp. (e.g. coffee), *Theaceae* sp. (e.g. tea), *Sterculiaceae* sp., *Rutaceae* sp. (e.g. lemons, oranges, mandarins and grapefruit); *Solanaceae* sp. (e.g. tomatoes, potatoes, peppers, capsicum, aubergines, tobacco), *Liliaceae* sp., *Compositae* sp. (e.g. lettuce, artichokes and chicory – including root chicory, endive or common chicory), *Umbelliferae* sp. (e.g. carrots, parsley, celery and celeriac), *Cucurbitaceae* sp. (e.g. cucumbers – including gherkins, pumpkins, watermelons, calabashes and melons), *Alliaceae* sp. (e.g. leeks and onions), *Cruciferae* sp. (e.g. white cabbage, red cabbage, broccoli, cauliflower, Brussels sprouts, pak choi, kohlrabi, radishes, horseradish, cress and chinese cabbage), *Leguminosae* sp. (e.g. peanuts, peas, lentils and beans – e.g. common beans and broad beans), *Chenopodiaceae* sp. (e.g. Swiss chard, fodder beet, spinach, beetroot), *Linaceae* sp. (e.g. hemp), *Cannabaceae* sp. (e.g. cannabis), *Malvaceae* sp. (e.g. okra, cocoa), *Papaveraceae* (e.g. poppy), *Asparagaceae* (e.g. asparagus); useful plants and ornamental plants in the garden and woods including turf, lawn, grass and *Stevia rebaudiana*; and in each case genetically modified types of these plants.

### **Phytopathogenic fungi**

The inventive compositions comprise a method for reducing damage of plants and plant parts or losses in harvested fruits or vegetables caused by phytopathogenic fungi by controlling such phytopathogenic fungi. Accordingly the inventive compositions have potent microbicidal activity and can be used for control of phytopathogenic fungi in crop protection and in the protection of materials.

Such phytopathogenic fungi include soilborne pathogens, which are in particular members of the classes *Plasmodiophoromycetes*, *Peronosporomycetes* (Syn. *Oomycetes*), *Chytridiomycetes*, *Zygomycetes*, *Ascomycetes*, *Basid-*

*oomycetes* and *Deuteromycetes* (Syn. *Fungi imperfecti*). Some fungicides are systemically active and can be used in plant protection as foliar, seed dressing or soil fungicide. Furthermore, they are suitable for combating fungi, which inter alia infest wood or roots of plant.

Examples of phytopathogenic fungi which can be treated in accordance with the invention comprise:

- 5 diseases caused by powdery mildew pathogens, for example *Blumeria* species, for example *Blumeria graminis*; *Podosphaera* species, for example *Podosphaera leucotricha*; *Sphaerotheca* species, for example *Sphaerotheca fuliginea*; *Uncinula* species, for example *Uncinula necator*;

- diseases caused by rust disease pathogens, for example *Gymnosporangium* species, for example *Gymnosporangium sabinae*; *Hemileia* species, for example *Hemileia vastatrix*; *Phakopsora* species, for example *Phakopsora pachyrhizi* and *Phakopsora meibomia*; *Puccinia* species, for example *Puccinia recondite*, *P. trititica*, *P. graminis* or *P. striiformis*; *Uromyces* species, for example *Uromyces appendiculatus*;

- diseases caused by pathogens from the group of the *Oomycetes*, for example *Albugo* species, for example *Albugo candida*; *Bremia* species, for example *Bremia lactucae*; *Peronospora* species, for example *Peronospora pisi* or *P. brassicae*; *Phytophthora* species, for example *Phytophthora infestans*; *Plasmopara* species, for example *Plasmopara viticola*; *Pseudoperonospora* species, for example *Pseudoperonospora humuli* or *Pseudoperonospora cubensis*; *Pythium* species, for example *Pythium ultimum*;

- leaf blotch diseases and leaf wilt diseases caused, for example, by *Alternaria* species, for example *Alternaria solani*; *Cercospora* species, for example *Cercospora beticola*; *Cladosporium* species, for example *Cladosporium cucumerinum*; *Cochliobolus* species, for example *Cochliobolus sativus* (conidia form: *Drechslera*, Syn: *Helminthosporium*), *Cochliobolus miyabeanus*; *Colletotrichum* species, for example *Colletotrichum lindemuthianum*; *Cyloconium* species, for example *Cyloconium oleaginum*; *Diaporthe* species, for example *Diaporthe citri*; *Elsinoe* species, for example *Elsinoe fawcettii*; *Gloeosporium* species, for example *Gloeosporium laeticolor*; *Glomerella* species, for example *Glomerella cingulata*; *Guignardia* species, for example *Guignardia bidwelli*; *Leptosphaeria* species, for example *Leptosphaeria maculans*, *Leptosphaeria nodorum*; *Magnaporthe* species, for example *Magnaporthe grisea*; *Microdochium* species, for example *Microdochium nivale*; *Mycosphaerella* species, for example *Mycosphaerella graminicola*, *M. arachidicola* and *M. fijiensis*; *Phaeosphaeria* species, for example *Phaeosphaeria nodorum*; *Pyrenophora* species, for example *Pyrenophora teres*, *Pyrenophora tritici repentis*; *Ramularia* species, for example *Ramularia collo-cygni*, *Ramularia areola*; *Rhynchosporium* species, for example *Rhynchosporium secalis*; *Septoria* species, for example *Septoria apii*, *Septoria lycopersii*; *Typhula* species, for example *Typhula incarnata*; *Venturia* species, for example *Venturia inaequalis*;

- root and stem diseases caused, for example, by *Corticium* species, for example *Corticium graminearum*; *Fusarium* species, for example *Fusarium oxysporum*; *Gaeumannomyces* species, for example *Gaeumannomyces graminis*; *Rhizoctonia* species, such as, for example *Rhizoctonia solani*; *Sarocladium* diseases caused for example by *Sarocladium oryzae*; *Sclerotium* diseases caused for example by *Sclerotium oryzae*; *Tapesia* species, for example *Tapesia acutiformis*; *Thielaviopsis* species, for example *Thielaviopsis basicola*;

ear and panicle diseases (including corn cobs) caused, for example, by *Alternaria* species, for example *Alternaria* spp.; *Aspergillus* species, for example *Aspergillus flavus*; *Cladosporium* species, for example *Cladosporium cladosporioides*; *Claviceps* species, for example *Claviceps purpurea*; *Fusarium* species, for example *Fusarium culmorum*; *Gibberella* species, for example *Gibberella zeae*; *Monographella* species, for example *Monographella nivalis*; *Septoria* species, for example *Septoria nodorum*;

diseases caused by smut fungi, for example *Sphacelotheca* species, for example *Sphacelotheca reiliana*; *Tilletia* species, for example *Tilletia caries*, *T. controversa*; *Urocystis* species, for example *Urocystis occulta*; *Ustilago* species, for example *Ustilago nuda*, *U. nuda tritici*;

fruit rot caused, for example, by *Aspergillus* species, for example *Aspergillus flavus*; *Botrytis* species, for example *Botrytis cinerea*; *Penicillium* species, for example *Penicillium expansum* and *P. purpurogenum*; *Sclerotinia* species, for example *Sclerotinia sclerotiorum*; *Verticillium* species, for example *Verticillium alboatrum*;

seed and soilborne decay, mould, wilt, rot and damping-off diseases caused, for example, by *Alternaria* species, caused for example by *Alternaria brassicicola*; *Aphanomyces* species, caused for example by *Aphanomyces eutiches*; *Ascochyta* species, caused for example by *Ascochyta lentis*; *Aspergillus* species, caused for example by *Aspergillus flavus*; *Cladosporium* species, caused for example by *Cladosporium herbarum*; *Cochliobolus* species, caused for example by *Cochliobolus sativus*; (Conidiaform: Drechslera, Bipolaris Syn: Helminthosporium); *Colletotrichum* species, caused for example by *Colletotrichum coccodes*; *Fusarium* species, caused for example by *Fusarium culmorum*; *Gibberella* species, caused for example by *Gibberella zeae*; *Macrophomina* species, caused for example by *Macrophomina phaseolina*; *Monographella* species, caused for example by *Monographella nivalis*; *Penicillium* species, caused for example by *Penicillium expansum*; *Phoma* species, caused for example by *Phoma lingam*; *Phomopsis* species, caused for example by *Phomopsis sojae*; *Phytophthora* species, caused for example by *Phytophthora cactorum*; *Pyrenophora* species, caused for example by *Pyrenophora graminea*; *Pyricularia* species, caused for example by *Pyricularia oryzae*; *Pythium* species, caused for example by *Pythium ultimum*; *Rhizoctonia* species, caused for example by *Rhizoctonia solani*; *Rhizopus* species, caused for example by *Rhizopus oryzae*; *Sclerotium* species, caused for example by *Sclerotium rolfsii*; *Septoria* species, caused for example by *Septoria nodorum*; *Typhula* species, caused for example by *Typhula incarnata*; *Verticillium* species, caused for example by *Verticillium dahliae*;

cancers, galls and witches' broom caused, for example, by *Nectria* species, for example *Nectria galligena*;

wilt diseases caused, for example, by *Monilinia* species, for example *Monilinia laxa*;

leaf blister or leaf curl diseases caused, for example, by *Exobasidium* species, for example *Exobasidium vexans*;

*Taphrina* species, for example *Taphrina deformans*;

decline diseases of wooden plants caused, for example, by Esca disease, caused for example by *Phaemoniella clamydospora*, *Phaeoacremonium aleophilum* and *Fomitiporia mediterranea*; *Eutypa* dyeback, caused for exam-

ple by *Eutypa lata* ; Ganoderma diseases caused for example by *Ganoderma boninense*; Rigidoporus diseases caused for example by *Rigidoporus lignosus*;

diseases of flowers and seeds caused, for example, by *Botrytis* species, for example *Botrytis cinerea*;

diseases of plant tubers caused, for example, by *Rhizoctonia* species, for example *Rhizoctonia solani*; *Helminthosporium* species, for example *Helminthosporium solani*;

Club root caused, for example, by *Plasmodiophora* species, for example *Plasmodiophora brassicae*;

diseases caused by bacterial pathogens, for example *Xanthomonas* species, for example *Xanthomonas campestris* pv. *oryzae*; *Pseudomonas* species, for example *Pseudomonas syringae* pv. *lachrymans*; *Erwinia* species, for example *Erwinia amylovora*.

- 10 Fungal diseases on roots and the stem base caused, for example, by black root rot (*Calonectria crotalariae*), charcoal rot (*Macrophomina phaseolina*), fusarium blight or wilt, root rot, and pod and collar rot (*Fusarium oxysporum*, *Fusarium orthoceras*, *Fusarium semitectum*, *Fusarium equiseti*), mycoleptodiscus root rot (*Mycoleptodiscus terrestris*), neocosmospora (*Neocosmospora vasinfecta*), pod and stem blight (*Diaporthe phaseolorum*), stem canker (*Diaporthe phaseolorum* var. *caulivora*), phytophthora rot (*Phytophthora megasperma*), brown stem
- 15 rot (*Phialophora gregata*), pythium rot (*Pythium aphanidermatum*, *Pythium irregulare*, *Pythium debaryanum*, *Pythium myriotylum*, *Pythium ultimum*), rhizoctonia root rot, stem decay, and damping-off (*Rhizoctonia solani*), sclerotinia stem decay (*Sclerotinia sclerotiorum*), sclerotinia southern blight (*Sclerotinia rolfsii*), thielaviopsis root rot (*Thielaviopsis basicola*).

## 20 **Plant Growth Regulation**

- In some cases, the inventive compositions can, at particular concentrations or application rates, also be used as herbicides, safeners, growth regulators or agents to improve plant properties, or as microbicides, for example as fungicides, antimycotics, bactericides, viricides (including compositions against viroids) or as compositions against MLO (Mycoplasma-like organisms) and RLO (Rickettsia-like organisms). The active ingredients of the
- 25 inventive composition intervene in the metabolism of the plants and can therefore also be used as growth regulators.

- Plant growth regulators may exert various effects on plants. The effect of the substances depends essentially on the time of application in relation to the developmental stage of the plant, and also on the amounts of active ingredient applied to the plants or their environment and on the type of application. In each case, growth regulators
- 30 should have a particular desired effect on the crop plants.

Plant growth-regulating compounds can be used, for example, to inhibit the vegetative growth of the plants. Such inhibition of growth is of economic interest, for example, in the case of grasses, since it is thus possible to

reduce the frequency of grass cutting in ornamental gardens, parks and sport facilities, on roadsides, at airports or in fruit crops. Also of significance is the inhibition of the growth of herbaceous and woody plants on roadsides and in the vicinity of pipelines or overhead cables, or quite generally in areas where vigorous plant growth is unwanted.

- 5 Also important is the use of growth regulators for inhibition of the longitudinal growth of cereal. This reduces or completely eliminates the risk of lodging of the plants prior to harvest. In addition, growth regulators in the case of cereals can strengthen the culm, which also counteracts lodging. The employment of growth regulators for shortening and strengthening culms allows the deployment of higher fertilizer volumes to increase the yield, without any risk of lodging of the cereal crop.
- 10 In many crop plants, inhibition of vegetative growth allows denser planting, and it is thus possible to achieve higher yields based on the soil surface. Another advantage of the smaller plants obtained in this way is that the crop is easier to cultivate and harvest.

Inhibition of the vegetative plant growth may also lead to enhanced yields because the nutrients and assimilates are of more benefit to flower and fruit formation than to the vegetative parts of the plants.

- 15 Frequently, growth regulators can also be used to promote vegetative growth. This is of great benefit when harvesting the vegetative plant parts. However, promoting vegetative growth may also promote generative growth in that more assimilates are formed, resulting in more or larger fruits.

- In some cases, yield increases may be achieved by manipulating the metabolism of the plant, without any detectable changes in vegetative growth. In addition, growth regulators can be used to alter the composition of the plants, which in turn may result in an improvement in quality of the harvested products. For example, it is possible to increase the sugar content in sugar beet, sugar cane, pineapples and in citrus fruit, or to increase the protein content in soya or cereals. It is also possible, for example, to use growth regulators to inhibit the degradation of desirable ingredients, for example sugar in sugar beet or sugar cane, before or after harvest. It is also possible to positively influence the production or the elimination of secondary plant ingredients. One example is the stimulation of the flow of latex in rubber trees.
- 20
- 25

Under the influence of growth regulators, parthenocarpic fruits may be formed. In addition, it is possible to influence the sex of the flowers. It is also possible to produce sterile pollen, which is of great importance in the breeding and production of hybrid seed.

- Use of growth regulators can control the branching of the plants. On the one hand, by breaking apical dominance, it is possible to promote the development of side shoots, which may be highly desirable particularly in the cultivation of ornamental plants, also in combination with an inhibition of growth. On the other hand, however, it is also possible to inhibit the growth of the side shoots. This effect is of particular interest, for example, in the cultivation of tobacco or in the cultivation of tomatoes.
- 30

Under the influence of growth regulators, the amount of leaves on the plants can be controlled such that defoliation of the plants is achieved at a desired time. Such defoliation plays a major role in the mechanical harvesting of cotton, but is also of interest for facilitating harvesting in other crops, for example in viticulture. Defoliation of the plants can also be undertaken to lower the transpiration of the plants before they are transplanted.

- 5 Growth regulators can likewise be used to regulate fruit dehiscence. On the one hand, it is possible to prevent premature fruit dehiscence. On the other hand, it is also possible to promote fruit dehiscence or even flower abortion to achieve a desired mass ("thinning"), in order to eliminate alternation. Alternation is understood to mean the characteristic of some fruit species, for endogenous reasons, to deliver very different yields from year to year. Finally, it is possible to use growth regulators at the time of harvest to reduce the forces required to detach the fruits, in order to allow mechanical harvesting or to facilitate manual harvesting.

- 15 Growth regulators can also be used to achieve faster or else delayed ripening of the harvested material before or after harvest. This is particularly advantageous as it allows optimal adjustment to the requirements of the market. Moreover, growth regulators in some cases can improve the fruit colour. In addition, growth regulators can also be used to concentrate maturation within a certain period of time. This establishes the prerequisites for complete mechanical or manual harvesting in a single operation, for example in the case of tobacco, tomatoes or coffee.

- 20 By using growth regulators, it is additionally possible to influence the resting of seed or buds of the plants, such that plants such as pineapple or ornamental plants in nurseries, for example, germinate, sprout or flower at a time when they are normally not inclined to do so. In areas where there is a risk of frost, it may be desirable to delay budding or germination of seeds with the aid of growth regulators, in order to avoid damage resulting from late frosts.

Finally, growth regulators can induce resistance of the plants to frost, drought or high salinity of the soil. This allows the cultivation of plants in regions which are normally unsuitable for this purpose.

### **Resistance Induction / Plant Health and other effects**

- 25 The compositions according to the invention also exhibit a potent strengthening effect in plants. Accordingly, they can be used for mobilizing the defences of the plant against attack by undesirable microorganisms.

- 30 Plant-strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances which are capable of stimulating the defence system of plants in such a way that the treated plants, when subsequently inoculated with undesirable microorganisms, develop a high degree of resistance to these microorganisms.

The active compounds according to the invention are also suitable for increasing the yield of crops. In addition, they show reduced toxicity and are well tolerated by plants.



Further, in context with the present invention plant physiology effects comprise the following:

Abiotic stress tolerance, comprising temperature tolerance, drought tolerance and recovery after drought stress, water use efficiency (correlating to reduced water consumption), flood tolerance, ozone stress and UV tolerance, tolerance towards chemicals like heavy metals, salts, pesticides (safener) etc..

- 5 Biotic stress tolerance, comprising increased fungal resistance and increased resistance against nematodes, viruses and bacteria. In context with the present invention, biotic stress tolerance preferably comprises increased fungal resistance and increased resistance against nematodes

Increased plant vigor, comprising plant health / plant quality and seed vigor, reduced stand failure, improved appearance, increased recovery, improved greening effect and improved photosynthetic efficiency.

- 10 Effects on plant hormones and/or functional enzymes.

Effects on growth regulators (promoters), comprising earlier germination, better emergence, more developed root system and/or improved root growth, increased ability of tillering, more productive tillers, earlier flowering, increased plant height and/or biomass, shorting of stems, improvements in shoot growth, number of kernels/ear, number of ears/m<sup>2</sup>, number of stolons and/or number of flowers, enhanced harvest index, bigger leaves, less  
 15 dead basal leaves, improved phyllotaxy, earlier maturation / earlier fruit finish, homogenous riping, increased duration of grain filling, better fruit finish, bigger fruit/vegetable size, sprouting resistance and reduced lodging.

Increased yield, referring to total biomass per hectare, yield per hectare, kernel/fruit weight, seed size and/or hectolitre weight as well as to increased product quality, comprising:

- 20 improved processability relating to size distribution (kernel, fruit, etc.), homogenous riping, grain moisture, better milling, better vinification, better brewing, increased juice yield, harvestability, digestibility, sedimentation value, falling number, pod stability, storage stability, improved fiber length/strength/uniformity, increase of milk and/or meat quality of silage fed animals, adaption to cooking and frying;

- 25 further comprising improved marketability relating to improved fruit/grain quality, size distribution (kernel, fruit, etc.), increased storage / shelf-life, firmness / softness, taste (aroma, texture, etc.), grade (size, shape, number of berries, etc.), number of berries/fruits per bunch, crispness, freshness, coverage with wax, frequency of physiological disorders, colour, etc.;

further comprising increased desired ingredients such as e.g. protein content, fatty acids, oil content, oil quality, aminoacid composition, sugar content, acid content (pH), sugar/acid ratio (Brix), polyphenols, starch content, nutritional quality, gluten content/index, energy content, taste, etc.;

- 30 and further comprising decreased undesired ingredients such as e.g. less mycotoxines, less aflatoxines, geosmin level, phenolic aromas, lacchase, polyphenol oxidases and peroxidases, nitrate content etc.

Sustainable agriculture, comprising nutrient use efficiency, especially nitrogen (N)-use efficiency, phosphorus (P)-use efficiency, water use efficiency, improved transpiration, respiration and/or CO<sub>2</sub> assimilation rate, better nodulation, improved Ca-metabolism etc..

5 Delayed senescence, comprising improvement of plant physiology which is manifested, for example, in a longer grain filling phase, leading to higher yield, a longer duration of green leaf colouration of the plant and thus comprising colour (greening), water content, dryness etc.. Accordingly, in the context of the present invention, it has been found that the specific inventive application of the inventive compositions makes it possible to prolong the green leaf area duration, which delays the maturation (senescence) of the plant. The main advantage to the farmer is a longer grain filling phase leading to higher yield. There is also an advantage to the farmer on the basis of greater flexibility in the harvesting time.

15 Therein “sedimentation value” is a measure for protein quality and describes according to Zeleny (Zeleny value) the degree of sedimentation of flour suspended in a lactic acid solution during a standard time interval. This is taken as a measure of the baking quality. Swelling of the gluten fraction of flour in lactic acid solution affects the rate of sedimentation of a flour suspension. Both a higher gluten content and a better gluten quality give rise to slower sedimentation and higher Zeleny test values. The sedimentation value of flour depends on the wheat protein composition and is mostly correlated to the protein content, the wheat hardness, and the volume of pan and hearth loaves. A stronger correlation between loaf volume and Zeleny sedimentation volume compared to SDS sedimentation volume could be due to the protein content influencing both the volume and Zeleny value ( *Czech J. Food Sci. Vol. 21, No. 3: 91–96, 2000*).

20 Further the “falling number” as mentioned herein is a measure for the baking quality of cereals, especially of wheat. The falling number test indicates that sprout damage may have occurred. It means that changes to the physical properties of the starch portion of the wheat kernel has already happened. Therein, the falling number instrument analyzes viscosity by measuring the resistance of a flour and water paste to a falling plunger. The time (in seconds) for this to happen is known as the falling number. The falling number results are recorded as an index of enzyme activity in a wheat or flour sample and results are expressed in time as seconds. A high falling number (for example, above 300 seconds) indicates minimal enzyme activity and sound quality wheat or flour. A low falling number (for example, below 250 seconds) indicates substantial enzyme activity and sprout-damaged wheat or flour.

30 The term “more developed root system” / “improved root growth” refers to longer root system, deeper root growth, faster root growth, higher root dry/fresh weight, higher root volume, larger root surface area, bigger root diameter, higher root stability, more root branching, higher number of root hairs, and/or more root tips and can be measured by analyzing the root architecture with suitable methodologies and Image analysis programmes (e.g. WinRhizo).

35 The term “crop water use efficiency” refers technically to the mass of agriculture produce per unit water consumed and economically to the value of product(s) produced per unit water volume consumed and can e.g. be measured in terms of yield per ha, biomass of the plants, thousand-kernel mass, and the number of ears per m<sup>2</sup>.

The term “nitrogen-use efficiency” refers technically to the mass of agriculture produce per unit nitrogen consumed and economically to the value of product(s) produced per unit nitrogen consumed, reflecting uptake and utilization efficiency.

Improvement in greening / improved colour and improved photosynthetic efficiency as well as the delay of senescence can be measured with well-known techniques such as a HandyPea system (Hansatech). Fv/Fm is a parameter widely used to indicate the maximum quantum efficiency of photosystem II (PSII). This parameter is widely considered to be a selective indication of plant photosynthetic performance with healthy samples typically achieving a maximum Fv/Fm value of approx. 0.85. Values lower than this will be observed if a sample has been exposed to some type of biotic or abiotic stress factor which has reduced the capacity for photochemical quenching of energy within PSII. Fv/Fm is presented as a ratio of variable fluorescence (Fv) over the maximum fluorescence value (Fm). The Performance Index is essentially an indicator of sample vitality. (See e.g. *Advanced Techniques in Soil Microbiology*, 2007, 11, 319-341; *Applied Soil Ecology*, 2000, 15, 169-182.)

The improvement in greening / improved colour and improved photosynthetic efficiency as well as the delay of senescence can also be assessed by measurement of the net photosynthetic rate (Pn), measurement of the chlorophyll content, e.g. by the pigment extraction method of Ziegler and Ehle, measurement of the photochemical efficiency (Fv/Fm ratio), determination of shoot growth and final root and/or canopy biomass, determination of tiller density as well as of root mortality.

Within the context of the present invention preference is given to improving plant physiology effects which are selected from the group comprising: enhanced root growth / more developed root system, improved greening, improved water use efficiency (correlating to reduced water consumption), improved nutrient use efficiency, comprising especially improved nitrogen (N)-use efficiency, delayed senescence and enhanced yield.

Within the enhancement of yield preference is given as to an improvement in the sedimentation value and the falling number as well as to the improvement of the protein and sugar content – especially with plants selected from the group of cereals (preferably wheat).

Preferably the novel use of the fungicidal compositions of the present invention relates to a combined use of a) preventively and/or curatively controlling pathogenic fungi, with or without resistance management, and b) at least one of enhanced root growth, improved greening, improved water use efficiency, delayed senescence and enhanced yield. From group b) enhancement of root system, water use efficiency and N-use efficiency is particularly preferred.

### Mycotoxins

In addition, the inventive treatment can reduce the mycotoxin content in the harvested material and the foods and feeds prepared therefrom. Mycotoxins include particularly, but not exclusively, the following: deoxynivalenol (DON), nivalenol, 15-Ac-DON, 3-Ac-DON, T2- and HT2-toxin, fumonisins, zearalenon, moniliformin, fusarin, diacetoxyscirpenol (DAS), beauvericin, enniatin, fusaroproliferin, fusarenol, ochratoxins, patulin, ergot alkaloids and aflatoxins which can be produced, for example, by the following fungi: *Fusarium* spec., such as *F. acuminatum*, *F. asiaticum*, *F. avenaceum*, *F. crookwellense*, *F. culmorum*, *F. graminearum* (*Gibberella zae*), *F. equiseti*, *F. fujikoro*i, *F. musarum*, *F. oxysporum*, *F. proliferatum*, *F. poae*, *F. pseudograminearum*, *F. sambucinum*, *F. scirpi*, *F. semitectum*, *F. solani*, *F. sporotrichoides*, *F. langsethiae*, *F. subglutinans*, *F. tricinctum*, *F. verticillioides* etc., and also by *Aspergillus* spec., such as *A. flavus*, *A. parasiticus*, *A. nomius*, *A. ochraceus*, *A. clavatus*, *A. terreus*, *A. versicolor*, *Penicillium* spec., such as *P. verrucosum*, *P. viridicatum*, *P. citrinum*, *P. expansum*, *P. claviforme*, *P. roqueforti*, *Claviceps* spec., such as *C. purpurea*, *C. fusiformis*, *C. paspali*, *C. africana*, *Stachybotrys* spec. and others.

### Material Protection

The inventive compositions or compositions can also be used in the protection of materials, for protection of industrial materials against attack and destruction by phytopathogenic fungi.

In addition, the inventive compositions can be used as antifouling compositions, alone or in combinations with other active ingredients.

Industrial materials in the present context are understood to mean inanimate materials which have been prepared for use in industry. For example, industrial materials which are to be protected by inventive compositions from microbial alteration or destruction may be adhesives, glues, paper, wallpaper and board/cardboard, textiles, carpets, leather, wood, fibers and tissues, paints and plastic articles, cooling lubricants and other materials which can be infected with or destroyed by microorganisms. Parts of production plants and buildings, for example cooling-water circuits, cooling and heating systems and ventilation and air-conditioning units, which may be impaired by the proliferation of microorganisms may also be mentioned within the scope of the materials to be protected. Industrial materials within the scope of the present invention preferably include adhesives, sizes, paper and card, leather, wood, paints, cooling lubricants and heat transfer fluids, more preferably wood.

The inventive compositions may prevent adverse effects, such as rotting, decay, discoloration, decoloration or formation of mould.

In the case of treatment of wood the compositions according to the invention may also be used against fungal diseases liable to grow on or inside timber. The term "timber" means all types of species of wood, and all types of working of this wood intended for construction, for example solid wood, high-density wood, laminated wood, and plywood. The method for treating timber according to the invention mainly consists in contacting a

composition according to the invention; this includes for example direct application, spraying, dipping, injection or any other suitable means.

In addition, the inventive compositions can be used to protect objects which come into contact with saltwater or brackish water, especially hulls, screens, nets, buildings, moorings and signalling systems, from fouling.


- 5 The inventive method for controlling phytopathogenic fungi can also be employed for protecting storage goods. Storage goods are understood to mean natural substances of vegetable or animal origin or processed products thereof which are of natural origin, and for which long-term protection is desired. Storage goods of vegetable origin, for example plants or plant parts, such as stems, leaves, tubers, seeds, fruits, grains, can be protected freshly harvested or after processing by (pre)drying, moistening, comminuting, grinding, pressing or roasting. Storage  
10 goods also include timber, both unprocessed, such as construction timber, electricity poles and barriers, or in the form of finished products, such as furniture. Storage goods of animal origin are, for example, hides, leather, furs and hairs. The inventive compositions may prevent adverse effects, such as rotting, decay, discoloration, decoloration or formation of mould.

- Microorganisms capable of degrading or altering the industrial materials include, for example, bacteria, fungi,  
15 yeasts, algae and slime organisms. The inventive compositions preferably act against fungi, especially moulds, wood-discoloring and wood-destroying fungi (*Ascomycetes*, *Basidiomycetes*, *Deuteromycetes* and *Zygomycetes*), and against slime organisms and algae. Examples include microorganisms of the following genera: *Alternaria*, such as *Alternaria tenuis*; *Aspergillus*, such as *Aspergillus niger*; *Chaetomium*, such as *Chaetomium globosum*; *Coniophora*, such as *Coniophora puetana*; *Lentinus*, such as *Lentinus tigrinus*; *Penicillium*, such as *Penicillium*  
20 *glaucum*; *Polyporus*, such as *Polyporus versicolor*; *Aureobasidium*, such as *Aureobasidium pullulans*; *Sclerophoma*, such as *Sclerophoma pityophila*; *Trichoderma*, such as *Trichoderma viride*; *Ophiostoma spp.*, *Ceratocystis spp.*, *Humicola spp.*, *Petriella spp.*, *Trichurus spp.*, *Coriolus spp.*, *Gloeophyllum spp.*, *Pleurotus spp.*, *Poria spp.*, *Serpula spp.* and *Tyromyces spp.*, *Cladosporium spp.*, *Paecilomyces spp.*, *Mucor spp.*, *Escherichia*, such as *Escherichia coli*; *Pseudomonas*, such as *Pseudomonas aeruginosa*; *Staphylococcus*, such as *Staphylococcus aureus*,  
25 *Candida spp.* and *Saccharomyces spp.*, such as *Saccharomyces cerevisiae*.

**Claims:**

1. A composition, comprising

(A) at least one thiazolyloxazoline of formula (I)



5 in which

R<sup>1</sup> represents phenyl, which is at least substituted with one methylsulfonyloxy and optionally be additionally substituted by one substituent selected from the group consisting of methyl, methoxy, fluoro or chloro,

or an agrochemically acceptable salt thereof,

10 (B) at least one biological control agent.

2. A composition according to claim 1, whereby the compound of the general formula (I) is represented by one of the compounds (I-1) to (I-8):

15 (I-1) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl} phenyl methanesulfonate,

(I-2) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-6-chlorophenyl methanesulfonate,

(I-3) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-3-chlorophenyl methanesulfonate,

20 (I-4) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-5-methylphenyl methanesulfonate,

(I-5) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-5-chlorophenyl methanesulfonate,

25 (I-6) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl} piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-4-methylphenyl methanesulfonate,

(I-7) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-4-chlorophenyl methanesulfonate,

(I-8) 2-{3-[2-(1-{[3,5-bis(difluoromethyl)-1H-pyrazol-1-yl]acetyl}piperidin-4-yl)-1,3-thiazol-4-yl]-4,5-dihydro-1,2-oxazol-5-yl}-6-fluorophenyl methanesulfonate.

5

3. The composition according to claim 1 or 2, whereby the biological control agent (B) comprises bacteria, fungi, proteins or secondary metabolites.

4. The composition according to any one of claims 1 to 3, wherein said composition acts in a fungicidal fashion.

10 5. The composition according to any one of claims 1 to 4, wherein said composition acts in a synergistic fashion.

6. The composition according to any one of claims 1 to 5, whereby the biological control agent is selected from B1 bacteria consisting of B1.1.1 *Bacillus subtilis*, strain QST713/AQ713, Accession No. NRRL B21661; B1.2 *Bacillus subtilis* strain AQ153, Accession No. NRRL 55614; B1.3 *Bacillus sp.* strain AQ175, ATCC Accession No. 55608; B1.4 *Bacillus sp.* strain AQ177, ATCC Accession No. 55609; B1.5 *Bacillus sp.* strain AQ178, ATCC Accession No. 53522; B1.1.6 *Bacillus pumilus*, strain QST2808, Accession No. NRRL B-30087; B1.1.7 *Bacillus pumilus*, strain GB34; B1.8 *Bacillus thuringiensis* strain AQ52, Accession No. NRRL B-21619; B1.1.9 *Streptomyces sp.*, strain, Accession No. NRRL B-30145; B1.1.10 *Streptomyces galbus*, syn. *Streptomyces griseoviridis*, strain QST 6047, Accession No. NRRL B-30232; B1.1.11 *Bacillus chitinosporus*, strain AQ746, Accession No. NRRL B-21618; B1.12 *Bacillus mycoides*, strain AQ726, Accession No. NRRL B-21664; B1.1.13 *Bacillus pumilus*, strain AQ717, Accession No. NRRL B21662; B1.1.14 *Bacillus subtilis*, strain AQ743, Accession No. NRRL B-21665; B1.15 *Rhodococcus globerulus* strain AQ719, Accession No. NRRL B21663; B1.1.16 *Bacillus thuringiensis subsp. aizawai*, strain ABTS-1857, SD-1372; B1.1.17 *Bacillus firmus*, strain CNMC I-1582; B1.1.18 *Bacillus subtilis*, strain AQ30002, Accession No. NRRL B-50421; B1.1.19 *Bacillus subtilis*, strain AQ30004, Accession No. NRRL B-50455) or variants or mutants thereof.

7. The composition according to any one of claims 1 to 5, whereby the biological control agent is selected from B2 fungi consisting of B2.2.1 *Coniothyrium minitans*, strain CON/M/91-8, Accession No. DSM-9660; B2.2.2 *Metschnikowia fructicola*, strain NRRL Y-30752; B2.3 *Microsphaeropsis ochracea*; B2.2.4 *Muscodora albus*, strain QST 20799, Accession No. NRRL 30547; B2.5 *Trichoderma atroviride*, strain SC1; B2.6 *Trichoderma harzianum rifai* strain KRL-AG2, also known as strain T-22, /ATCC Accession No. 208479; B2.7 *Muscodora roseus* strain A3-5, Accession No. NRRL 30548; B2.2.8 *Paecilomyces lilacinus*, spores of *P. lilacinus* strain 251, AGAL 89/030550; B2.9 *Trichoderma koningii*; B2.10 *Talaromyces flavus*, strain V117b; B2.11 *Trichoderma atroviride*, strain no. V08/002387; B2.12

35

*Trichoderma atroviride*, strain no. NMI No. V08/002388; B2.13 *Trichoderma atroviride*, strain no. NMI No. V08/002389; B2.14 *Trichoderma atroviride*, strain no. NMI No. V08/002390; B2.15 *Trichoderma harzianum*, strain ITEM 908) or variants or mutants thereof.

8. The composition according to any one of claims 1 to 5, whereby the biological control agent is selected from B6 proteins or secondary metabolites consisting of B6.6.2 terpene constituents of extract of *Chenopodium ambrosioides* near *ambrosioides* as synthetically manufactured containing a mixture of three terpenes, i.e.  $\alpha$ -terpinene, p-cymene and limonene, as pesticidally active ingredients.

9. The composition according to any one of claims 1 to 5, whereby the biological control agent is selected from

B1 bacteria consisting of B1.1.1 *Bacillus subtilis*, strain QST713/AQ713, Accession No. NRRL B21661; B1.1.6 *Bacillus pumilus*, strain QST2808, Accession No. NRRL B-30087 and B1.1.18 *Bacillus subtilis* strain AQ30002, Accession No. NRRL B-50421 or variants or mutants thereof.

10. Method for preparing an agricultural composition comprising adding at least one agriculturally suitable extender, solvent, spontaneity promoter, carrier, emulsifier, dispersant, frost protectant, biocides, thickener or adjuvant to the composition according to any one of claims 1 to 9.

11. Method for reducing damage of plants and plant parts or losses in harvested fruits or vegetables caused by phytopathogenic fungi by controlling such phytopathogenic fungi, comprising applying the composition according to any one of claims 1 to 9 to the plant or the phytopathogenic fungi or the habitat of the plant or the habitat of the phytopathogenic fungi.

12. Use of the composition according to any one of claims 1 to 9 for reducing damage of plants and plant parts or losses in harvested fruits or vegetables caused by phytopathogenic fungi by controlling such phytopathogenic fungi.

13. The method or the use according to any one of the claims 11 to 12, wherein the plants are genetically modified.



## INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2014/069132

A. CLASSIFICATION OF SUBJECT MATTER		
INV.	A01N43/80	A01N63/00 A01N63/02 A01N63/04 A01P3/00
ADD.	A01P21/00	
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) A01N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/127704 A1 (BAYER IP GMBH [DE]) 6 September 2013 (2013-09-06) pages 2, 3 page 6, line 29 page 7, lines 1, 25, 33-35 page 35, lines 34-37 page 42, lines 26-27 page 45, lines 10-11	1-6,9-13
Y	US 2012/070421 A1 (DIETZ JOCHEN [DE] ET AL) 22 March 2012 (2012-03-22) cited in the application paragraphs [0001], [0013], [0045] ----- -/-	1-6,9-13
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search  6 October 2014		Date of mailing of the international search report  22/01/2015
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer  Sawicki, Marcin

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2014/069132

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2012/025557 A1 (BAYER CROPSCIENCE AG [DE]; TSUCHIYA TOMOKI [DE]; WASNAIRE PIERRE [DE];) 1 March 2012 (2012-03-01) cited in the application pages 1, 73-77; table 1 page 55, line 25 - page 56, line 8 -----	1-6,9-13
Y	US 2012/034315 A1 (HANAGAN MARY ANN [US] ET AL OBERHOLZER MATTHEW RICHARD [US] ET AL) 9 February 2012 (2012-02-09) cited in the application pages 33-35; table 7 page 37; table 8 paragraph [0345] -----	1-6,9-13
Y	US 2012/003199 A1 (SCHERER MARIA [DE] ET AL) 5 January 2012 (2012-01-05) paragraphs [0027] - [0039], [0059] -----	1-6,9-13
A,P	WO 2013/178656 A1 (BAYER CROPSCIENCE AG [DE]) 5 December 2013 (2013-12-05) page 11; claim 1; compounds F313-F316, F320-F322 -----	1-6,9-13

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP2014/069132

## Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

6, 9(completely); 1-5, 10-13(partially)

### Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 6, 9(completely); 1-5, 10-13(partially)

relating to a composition comprising (A) at least one thiazolyloxazoline of formula (1) and (B1) bacteria.

---

2. claims: 7(completely); 1-5, 10-13(partially)

relating to a composition comprising (A) at least one thiazolyloxazoline of formula (1) and (B2) fungi.

---

3. claims: 1, 2, 4, 5, 10-13(all partially)

relating to a composition comprising (A) at least one thiazolyloxazoline of formula (1) and (B3) protozoas.

---

4. claims: 1, 2, 4, 5, 10-13(all partially)

relating to a composition comprising (A) at least one thiazolyloxazoline of formula (1) and (B4) viruses.

---

5. claims: 1, 2, 4, 5, 10-13(all partially)

relating to a composition comprising (A) at least one thiazolyloxazoline of formula (1) and (B5) entomopathogenic nematode.

---

6. claims: 8(completely); 1-5, 10-13(partially)

relating to a composition comprising (A) at least one thiazolyloxazoline of formula (1) and (B6) proteins and secondary metabolites.

---

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2014/069132

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013127704 A1	06-09-2013	AR 090157 A1 CA 2865300 A1 CN 104244716 A CO 7061086 A2 EP 2819518 A1 PE 17352014 A1 WO 2013127704 A1	22-10-2014 06-09-2013 24-12-2014 19-09-2014 07-01-2015 12-11-2014 06-09-2013
US 2012070421 A1	22-03-2012	AU 2010255831 A1 CA 2762268 A1 CN 102458119 A CR 20110613 A EA 201101699 A1 EP 2437604 A2 JP 2012528820 A KR 20120046149 A US 2012070421 A1 WO 2010139656 A2	12-01-2012 09-12-2010 16-05-2012 06-01-2012 30-07-2012 11-04-2012 15-11-2012 09-05-2012 22-03-2012 09-12-2010
WO 2012025557 A1	01-03-2012	AU 2011295045 A1 CA 2809211 A1 CN 103180317 A EA 201390240 A1 EP 2609094 A1 JP 2013539464 A KR 20130138751 A PE 01852014 A1 US 2012122929 A1 US 2013296272 A1 US 2014206646 A1 WO 2012025557 A1	21-03-2013 01-03-2012 26-06-2013 30-08-2013 03-07-2013 24-10-2013 19-12-2013 16-02-2014 17-05-2012 07-11-2013 24-07-2014 01-03-2012
US 2012034315 A1	09-02-2012	AR 076384 A1 AU 2010239456 A1 CA 2757055 A1 CN 102459256 A CO 6400143 A2 DK 2421861 T3 EP 2421861 A1 ES 2402113 T3 JP 2012524784 A KR 20120007056 A NZ 595278 A PE 08792012 A1 PT 2421861 E RU 2011147184 A TW 201038568 A US 2012034315 A1 US 2014127322 A1 UY 32567 A WO 2010123791 A1	08-06-2011 13-10-2011 28-10-2010 16-05-2012 15-03-2012 15-04-2013 29-02-2012 29-04-2013 18-10-2012 19-01-2012 22-02-2013 16-08-2012 18-03-2013 27-05-2013 01-11-2010 09-02-2012 08-05-2014 30-11-2010 28-10-2010
US 2012003199 A1	05-01-2012	AR 076150 A1 AU 2010227493 A1 CA 2753150 A1 CN 102361551 A CR 20110460 A EA 201101333 A1	18-05-2011 20-10-2011 30-09-2010 22-02-2012 24-10-2011 30-05-2012

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2014/069132

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		EC SP11011349 A	31-10-2011
		EP 2410840 A2	01-02-2012
		JP 2012521390 A	13-09-2012
		KR 20110132461 A	07-12-2011
		NZ 594887 A	29-11-2013
		PE 05362012 A1	05-05-2012
		US 2012003199 A1	05-01-2012
		US 2014212401 A1	31-07-2014
		WO 2010108973 A2	30-09-2010
-----			
WO 2013178656	A1	05-12-2013	
		AU 2013269723 A1	20-11-2014
		CA 2874916 A1	05-12-2013
		WO 2013178656 A1	05-12-2013
-----			