



(86) Date de dépôt PCT/PCT Filing Date: 2016/02/19
(87) Date publication PCT/PCT Publication Date: 2016/08/25
(45) Date de délivrance/Issue Date: 2019/12/17
(85) Entrée phase nationale/National Entry: 2017/08/15
(86) N° demande PCT/PCT Application No.: JP 2016/054822
(87) N° publication PCT/PCT Publication No.: 2016/133194
(30) Priorité/Priority: 2015/02/20 (JP2015-032201)

(51) Cl.Int./Int.Cl. *A61K 31/4439* (2006.01),
A61K 31/5377 (2006.01), *A61P 35/00* (2006.01),
A61P 35/02 (2006.01)
(72) Inventeur/Inventor:
SEKI, TAKAHIKO, JP
(73) Propriétaire/Owner:
DAIICHI SANKYO COMPANY, LIMITED, JP
(74) Agent: MARKS & CLERK

(54) Titre : POLYTHERAPIE POUR TRAITER UN CANCER
(54) Title: METHOD FOR TREATING CANCER BY COMBINED USE

(57) **Abrégé/Abstract:**

It is intended to provide a medicament and a method for treating cancer comprising a compound having MDM2 inhibiting activity and a compound having FLT3 inhibiting activity in combination. The present invention provides a medicament comprising (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6''-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2''-oxo-1'',2''-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3''-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof in combination, and a treatment method using these compounds or salts in combination.

Abstract

It is intended to provide a medicament and a method for treating cancer comprising a compound having MDM2 inhibiting activity and a compound having FLT3 inhibiting activity in combination. The present invention provides a medicament comprising (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl)urea or a pharmaceutically acceptable salt thereof in combination, and a treatment method using these compounds or salts in combination.

Description

Title of Invention: METHOD FOR TREATING CANCER BY
COMBINED USE

Technical Field

[0001]

The present invention relates to a medicament and a method for treating cancer comprising a compound having murine double minute 2 (MDM2) inhibiting activity and a compound having Fms-like tyrosine kinase 3 (FLT3) inhibiting activity in combination.

Background Art

[0002]

p53 is known as an important factor for inhibiting canceration of cells. p53 is a transcription factor that induces the expression of genes involved in the cell cycle and cellular apoptosis in response to various stresses. p53 is thought to inhibit canceration of cells by a transcription regulating function thereof. In fact, deletion or mutation of the p53 gene is observed in about half of human cancer cases.

[0003]

Meanwhile, overexpression of murine double minute 2 (MDM2), a type of E3 ubiquitin ligase, is known as a factor for canceration of cells that are cancerated in

spite of the presence of normal p53. MDM2 is a protein whose expression is induced by p53. MDM2 negatively regulates p53 by binding to the transcription activity domain of p53 to decrease the transcription activity of p53, exporting p53 out of the nucleus, and mediating degradation of p53 by acting as an ubiquitination ligase against p53. Therefore, it is thought that inactivation of functions of and degradation of p53 are promoted in cells in which MDM2 is overexpressed, resulting in canceration (Non Patent Document 1).

[0004]

Paying attention to such functions of MDM2, many approaches have been attempted using substances that inhibit the suppression of p53 functions by MDM2 as candidate anti-tumor agents. Examples of MDM2 inhibitors targeting the MDM2-p53 binding site have been reported, which include spirooxindole derivatives (Patent Documents 1 to 15 and Non Patent Documents 1 to 3), indole derivatives (Patent Document 16), pyrrolidine-2-carboxamide derivatives (Patent Document 17), pyrrolidinone derivatives (Patent Document 18), isoindolinone derivatives (Patent Document 19 and Non Patent Document 4) and dispiropyrrrolidine compounds (Patent Document 20).

[0005]

FLT3 is a protein belonging to receptor tyrosine kinase class III together with KIT, FMS and PDGFR, etc.,

and is thought to be involved in the hematopoietic system (Non Patent Documents 5 to 8). Its structure has an extracellular region composed of five immunoglobulin-like domains, one juxtamembrane region (JM domain), two tyrosine kinase domains (TK1 and TK2) divided by a kinase insert domain (KI domain), and a C-terminal domain. FLT3 is highly expressed in brain, placenta, liver and hematopoietic stem cells (Non Patent Documents 6 to 9).

A ligand of FLT3 (FL) is expressed in stromal cells of bone marrow and stimulates stem cells, either alone or in collaboration with other cytokines (Non Patent Documents 10 to 13). The ligand-receptor interaction between FL and FLT3 is considered to have important functions in the hematopoietic system.

Meanwhile, high expression of FLT3 is observed in most cases in samples of acute myeloid leukemia (AML) or acute lymphatic leukemia (ALL) patients, and high expression of FLT3 is also seen in chronic myeloid leukemia (CML). It is also known that growth of AML cells is more remarkably enhanced than that of ALL cells by stimulation of FL (Non Patent Documents 14 to 18). The FLT3 gene is the gene that is most frequently mutated in acute myeloid leukemia (AML) cases, and either of internal tandem duplications (ITDs) in the juxtamembrane region (Non Patent Document 19) or a mutation in the FLT3 activation loop region (Non Patent Document 20) is confirmed in approximately 30% to 35% of patients. The

mutation of FLT3-ITD or the activation loop region is associated with constitutive activation of tyrosine kinase activity.

[0006]

N-(5-tert-Butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea (quizartinib) having FLT3 inhibiting activity is known to have anti-tumor activity. Treatment of various cancers using quizartinib has been proposed in documents. Various dosing regimens have been reported. See, for example, Patent Documents 21 to 23. Also, the effects of combined use of quizartinib and anthracycline, a topoisomerase inhibitor or a tumor cell metabolic antagonist have been reported (Patent Document 24).

[0007]

As for the relation between an MDM2 inhibitor and a FLT3 inhibitor, it has been reported that administration of an MDM2 inhibitor is preferred for patients whose cells contain FLT3 having an activating mutation (Patent Document 25). This document also states that combined administration of a FLT3 inhibitor and an MDM2 inhibitor is preferred for patients whose cells contain FLT3 having an activating mutation, but it does not disclose specific effects of the combined use of specific drugs.

[0008]

There are various reports on the effects of combined use of various MDM2 inhibitors and various anti-tumor agents (Patent Documents 26 to 29).

Citation List

Patent Documents

[0009]

- Patent Document 1: WO2006/091646
- Patent Document 2: WO2006/136606
- Patent Document 3: WO2007/104664
- Patent Document 4: WO2007/104714
- Patent Document 5: WO2008/034736
- Patent Document 6: WO2008/036168
- Patent Document 7: WO2008/055812
- Patent Document 8: WO2008/141917
- Patent Document 9: WO2008/141975
- Patent Document 10: WO2009/077357
- Patent Document 11: WO2009/080488
- Patent Document 12: WO2010/084097
- Patent Document 13: WO2010/091979
- Patent Document 14: WO2010/094622
- Patent Document 15: WO2010/121995
- Patent Document 16: WO2008/119741
- Patent Document 17: WO2010/031713
- Patent Document 18: WO2010/028862
- Patent Document 19: WO2006/024837
- Patent Document 20: WO2012/121361

Patent Document 21: U. S. Patent Application Publication
US 2007/0232604

Patent Document 22: U. S. Patent Application Publication
US 2009/0123418

Patent Document 23: U. S. Patent Application Publication
US 2009/0131426

Patent Document 24: WO2010/111172

Patent Document 25: WO2011/127058

Patent Document 26: EP1712235

Patent Document 27: WO2007/115289

Patent Document 28: WO2013/139724

Patent Document 29: WO2014/107713

Non Patent Documents

[0010]

Non Patent Document 1: Ding K. et al., "Structure-Based
Design of Potent Non-Peptide MDM2 Inhibitors",
J.Am.Chem.Soc., 27 July 2005, 127(29), 10130-10131

Non Patent Document 2: Ding K. et al., "Structure-Based
Design of Spiro-oxindoles as Potent, Specific Small-
Molecule Inhibitors of the MDM2-p53 Interaction",
J.Med.Chem., 15 June 2006, 49(12), 3432-3435

Non Patent Document 3: Yu S. et al., "Potent and Orally
Active Small-Molecule Inhibitors of the MDM2-p53
Interaction", J.Med.Chem., 24 December 2009, 52(24),
7970-7973

Non Patent Document 4: Hardcastle I. R. et al., "Small-Molecule Inhibitors of the MDM2-p53 Protein-Protein Interaction Based on an Isoindolinone Scaffold", J.Med.Chem., 19 October 2006, 49(21), 6209-6221

Non Patent Document 5: Rosnet O. et al., "Isolation and chromosomal localization of a novel FMS-like tyrosine kinase gene", Genomics, February 1991, 9(2), 380-385

Non Patent Document 6: Rosnet O. et al., "Murine Flt 3, a gene encoding a novel tyrosine kinase receptor of the PDGFR/CSF1R family", Oncogene, September 1991, 6(9), 1641-1650

Non Patent Document 7: Matthews W. et al., "A receptor tyrosine kinase specific to hematopoietic stem and progenitor cell-enriched populations", Cell, 28 June 1991, 65(7), 1143-1152

Non Patent Document 8: Rosnet O. et al., "Human FLT3/FLK2 gene: cDNA cloning and expression in hematopoietic cells", Blood, 15 August 1993, 82(4), 1110-1119

Non Patent Document 9: Rusten L. S. et al., "The FLT3 ligand is a direct and potent stimulator of the growth of primitive and committed human CD34+ bone marrow progenitor cells in vitro", Blood, 15 February 1996, 87(4), 1317-1325

Non Patent Document 10: Hannum C. et al., "Ligand for FLT3/FLK2 receptor tyrosine kinase regulates growth of haematopoietic stem cells and is encoded by variant RNAs", Nature, 14 April 1994, 368(6472), 643-648

Non Patent Document 11: McKenna H. J. et al., "Effect of flt3 ligand on the ex vivo expansion of human CD34+ hematopoietic progenitor cells", Blood, 1 November 1995, 86(9), 3413-3420

Non Patent Document 12: Hirayama F. et al., "The flt3 ligand supports proliferation of lymphohematopoietic progenitors and early B-lymphoid progenitors", Blood, 1 April 1995, 85(7), 1762-1768

Non Patent Document 13: Lisovsky M. et al., "Flt3-ligand production by human bone marrow stromal cells", Leukemia, June 1996, 10(6), 1012-1018

Non Patent Document 14: Piacibello W. et al., "Effects of human FLT3 ligand on myeloid leukemia cell growth: heterogeneity in response and synergy with other hematopoietic growth factors", Blood, 1 November 1995, 86(11), 4105-4114

Non Patent Document 15: Stacchini A. et al., "Expression of type III receptor tyrosine kinases FLT3 and KIT and responses to their ligands by acute myeloid leukemia blasts", Leukemia, October 1996, 10(10), 1584-1591

Non Patent Document 16: Lisovsky M. et al., "Flt3 ligand stimulates proliferation and inhibits apoptosis of acute myeloid leukemia cells: regulation of Bcl-2 and Bax", Blood, 15 November 1996, 88(10), 3987-3997

Non Patent Document 17: Birg F. et al., "Expression of the FMS/KIT-like gene FLT3 in human acute leukemias of

the myeloid and lymphoid lineages", Blood, 15 November 1992, 80(10), 2584-2593

Non Patent Document 18: Dehmel U. et al., "Effects of FLT3 ligand on human leukemia cells. I. Proliferative response of myeloid leukemia cells", Leukemia, February 1996, 10(2), 261-270

Non Patent Document 19: Nakao M. et al., "Internal tandem duplication of the flt3 gene found in acute myeloid leukemia", Leukemia, December 1996, 10(12), 1911-1918

Non Patent Document 20: Yamamoto Y. et al., "Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies", Blood, 15 April 2001, 97(8), 2434-2439.

Summary of Invention

Technical Problem

[0011]

An object of the present invention is to provide a medicament and a method for treating cancer comprising a compound having MDM2 inhibiting activity and a compound having FLT3 inhibiting activity in combination.

Solution to the Problem

[0012]

As a result of extensive studies, the present inventors have found that use of (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-

4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide, which is a compound having MDM2 inhibiting activity, or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl)urea, which is a compound having FLT3 inhibiting activity, or a pharmaceutically acceptable salt thereof in combination particularly produces an excellent anti-tumor effect while keeping adverse reaction (e.g. weight loss) low, and accomplished the present invention.

[0013]

Specifically, the present invention relates to the following [1] to [21]:

[1] A medicament for cancer treatment comprising (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl)urea or a pharmaceutically acceptable salt thereof which are administered in combination.

[2] A medicament according to [1], wherein the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-

3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-ylethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof are separately contained as active ingredients in different formulations and administered at the same time or different times.

[3] A medicament according to [1], wherein the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-ylethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof are contained in a single formulation.

[4] A medicament according to [1], wherein the medicament is a kit formulation comprising the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-

tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof.

[5] A method for treating cancer comprising administering (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof in combination.

[6] A medicament according to any one of [1] to [4], wherein the respective salts of the compounds are (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide p-toluenesulfonate and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea dihydrochloride.

[7] A treatment method according to [5], wherein the respective salts of the compounds are (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-

indole]-5'-carboxamide p-toluenesulfonate and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl)urea dihydrochloride.

[8] A medicament according to any one of [1] to [4] and [6], wherein the cancer is blood cancer (leukemia, lymphoma or multiple myeloma), brain tumor, head and neck cancer, esophageal cancer, stomach cancer, appendix cancer, colon cancer, anus cancer, gallbladder cancer, cholangiocarcinoma, pancreatic cancer, gastrointestinal stromal tumor, lung cancer, liver cancer, mesothelioma, thyroid cancer, renal cancer, prostate cancer, neuroendocrine tumor, melanoma, breast cancer, endometrial cancer, cervical cancer, ovarian cancer, osteosarcoma, soft tissue sarcoma, Kaposi's sarcoma, myosarcoma, bladder cancer or testicular cancer.

[9] A treatment method according to [5] or [7], wherein the cancer is blood cancer (leukemia, lymphoma or multiple myeloma), brain tumor, head and neck cancer, esophageal cancer, stomach cancer, appendix cancer, colon cancer, anus cancer, gallbladder cancer, cholangiocarcinoma cancer, pancreatic cancer, gastrointestinal stromal tumor, lung cancer, liver cancer, mesothelioma, thyroid cancer, renal cancer, prostate cancer, neuroendocrine tumor, melanoma, breast cancer, endometrial cancer, cervical cancer, ovarian cancer,

osteosarcoma, soft tissue sarcoma, Kaposi's sarcoma, myosarcoma, bladder cancer or testicular cancer.

[10] A medicament according to any one of [1] to [4] and [6], wherein the cancer is leukemia.

[11] A treatment method according to [5] or [7], wherein the cancer is leukemia.

[12] A medicament according to any one of [1] to [4] and [6], wherein the cancer is leukemia having an activating mutation of FLT3.

[13] A treatment method according to [5] or [7], wherein the cancer is leukemia having an activating mutation of FLT3.

[14] A medicament according to any one of [1] to [4] and [6], wherein the cancer is acute myeloid leukemia (AML).

[15] A treatment method according to [5] or [7], wherein the cancer is acute myeloid leukemia (AML).

[16] A medicament according to any one of [1] to [4] and [6], wherein the cancer is acute myeloid leukemia (AML) having a FLT3-ITD mutation.

[17] A treatment method according to [5] or [7], wherein the cancer is acute myeloid leukemia (AML) having a FLT3-ITD mutation.

[18] A medicament according to any one of [1] to [4] and [6], wherein the cancer has wild-type TP53.

[19] A treatment method according to [5] or [7], wherein the cancer is cancer having wild-type TP53.

[20] A medicament according to any one of [1] to [4] and [6], wherein the cancer is cancer confirmed to be MDM2 inhibitor-sensitive using a gene signature.

[21] A treatment method according to [5] or [7], wherein the cancer is confirmed to be MDM2 inhibitor-sensitive using a gene signature.

[22] Use of (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for use in the treatment of cancer in combination with N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof.

[23] Use of N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for use in the treatment of cancer in combination with (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-

pyrrolidine-3',3''-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof.

[24] A use according to [22] or [23] which results in the manufacture of a medicament as defined above in any one of [1], [2], [3], [4], [6], [8], [10], [12], [14], [16], [18], and [20], or as otherwise described herein.

Advantageous Effects of Invention

[0014]

The present invention is useful as a method for treating cancer and/or an anti-cancer agent.

Brief Description of Drawings

[0015]

[Figure 1-1] Figure 1-1 is a diagram showing *in vivo* effects of combined use of (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6''-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2''-oxo-1'',2''-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3''-indole]-5'-carboxamide tosylate (Compound A) and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea (quizartinib) on subcutaneously transplanted tumors of human acute myeloid leukemia cell line MOLM-13 having a FLT3-ITD mutation and wild-type TP53 in mice, and body weight change caused by combined administration thereof.

The symbol x depicts an untreated control group, the symbol open circle depicts 0.5 mg/kg quizartinib, the symbol filled circle depicts 1 mg/kg quizartinib, the symbol open triangle depicts 25 mg/kg Compound A, the symbol open square depicts 25 mg/kg Compound A + 0.5 mg/kg quizartinib, and the symbol filled square depicts 25 mg/kg Compound A + 1 mg/kg quizartinib. The horizontal axis shows the number of days after tumor inoculation. The vertical axis of the upper panel shows estimated tumor volume calculated from tumor size. The vertical axis of the lower panel shows body weight change % relative to body weight on the first day of administration. The symbol filled triangle on the horizontal axis depicts the administration day of each compound. The error bar represents SE for the upper panel and SD for the lower panel.

[Figure 1-2] Figure 1-2 is a diagram showing *in vivo* effects of combined use of (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide tosylate (Compound A) and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea (quizartinib) on subcutaneously transplanted tumors of human acute myeloid leukemia cell line MOLM-13 having a FLT3-ITD mutation and wild-type TP53 in mice, and body

weight change caused by combined administration thereof. The symbol x depicts an untreated control group, the symbol open circle depicts 0.5 mg/kg quizartinib, the symbol filled circle depicts 1 mg/kg quizartinib, the symbol filled triangle depicts 50 mg/kg Compound A, the symbol open square depicts 50 mg/kg Compound A + 0.5 mg/kg quizartinib, and the symbol filled square depicts 50 mg/kg Compound A + 1 mg/kg quizartinib. The abscissa shows the number of days after tumor inoculation. The ordinate of the upper panel shows estimated tumor volume calculated from tumor size. The ordinate of the lower panel shows body weight change % relative to body weight on the first day of administration. The symbol filled triangle on the abscissa depicts the administration day of each compound. The error bar represents SE for the upper panel and SD for the lower panel.

Description of Embodiments

[0016]


In the present invention, the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide is a compound of Example 70 in WO2012/121361. This compound can be produced by a method described in WO2012/121361.

[0017]

In the present invention, the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea is also referred to as 1-(5-tert-butyl-1,2-oxazol-3-yl)-3-(4-{7-[2-(morpholin-4-yl)ethoxy]imidazo[2,1-b][1,3]benzothiazol-2-yl}phenyl)urea or also referred to as quizartinib or AC220. This compound is represented by the following formula:

[0018]

[Formula 1]



[0019]

This compound can be produced by a method described in WO2007/109120.

[0020]

In the present invention, the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea may be various pharmaceutically acceptable salts.

Examples of the salts can include: hydrohalides such as hydrochloride and hydroiodide; inorganic acid salts such as nitrate, perchlorate, sulfate and phosphate; lower alkanesulfonates such as methanesulfonate, trifluoromethanesulfonate and ethanesulfonate; arylsulfonates such as benzenesulfonate and p-toluenesulfonate; organic acid salts such as formate, acetate, malate, fumarate, succinate, citrate, tartrate, oxalate and maleate; amino acid salts such as ornithine salt, glutamate and aspartate; alkali metal salts such as sodium salt, potassium salt and lithium salt; alkaline earth metal salts such as calcium salt and magnesium salt; inorganic salts such as ammonium salt; and organic amine salts such as dibenzylamine salt, morpholine salt, phenylglycine alkyl ester salt, ethylenediamine salt, N-methylglucamine salt, diethylamine salt, triethylamine salt, cyclohexylamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, diethanolamine salt, N-benzyl-N-(2-phenylethoxy)amine salt, piperazine salt, tetramethylammonium salt and tris(hydroxymethyl)aminomethane salt.

The salt of the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide is preferably p-toluenesulfonate. The salt of the N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-

(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea is preferably hydrochloride, particularly, dihydrochloride.

[0021]

In the present invention, the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or their pharmaceutically acceptable salts may each be present in a free or solvate form. This compound or the salt thereof may be present in a hydrate form, for example, by absorbing moisture in the air. The solvate is not particularly limited so long as it is pharmaceutically acceptable. Specifically, the solvate is preferably a hydrate, an ethanol solvate or the like. Moreover, this compound may be in an N-oxide form when containing a nitrogen atom. These solvate and N-oxide forms are also included in the present invention.

[0022]

The (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-

ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or their pharmaceutically acceptable salts may have stereoisomers depending on their structures. The compounds or the salts also encompass all these stereoisomers and mixtures of these stereoisomers in any ratio. The stereoisomers are as defined in 1996 IUPC, Pure and Applied Chemistry 68, 2193-2222. When the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or their pharmaceutically acceptable salts are each present as tautomers, these tautomers may be present in equilibrium or a certain form may be dominantly present. All these cases are included in the scope of the present invention. The tautomers refer to isomers resulting from the shift of a proton of one atom of the molecule to another atom.

[0023]

The (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea

or their pharmaceutically acceptable salts may each be a "pharmaceutically acceptable prodrug compound" that is converted to the desired compound through enzymatic oxidation, reduction, hydrolysis or the like or through hydrolysis or the like induced by gastric acid or the like, due to a reaction induced by an enzyme, gastric acid or the like under physiological conditions *in vivo*.
[0024]

Examples of the prodrug include compounds obtained by acylation, alkylation or phosphorylation.
[0025]

Prodrugs of the compounds can be produced from the compound according to a method known in the art. Moreover, prodrugs of the compounds also include those converted to the desired compounds under physiological conditions as described in "Development of Pharmaceutical Products", vol. 7, Molecule Design, p. 163-198, Hirokawa-Shoten Ltd. (1990).
[0026]

In the present invention, the terms "tumor" and "cancer" are used interchangeably. Furthermore, in the present invention, tumor, malignant tumor, cancer, malignant neoplasm, carcinoma, sarcoma, and the like may be collectively referred to as "tumor" or "cancer".
[0027]

In the present invention, "FLT3" means Fms-like tyrosine kinase 3 (FLT3) and has the same meaning as FLK2,

STK1, CD135 and FLK-2. FLT3 also includes homologs derived from various animal species. Human FLT3 is a molecule registered in NCBI under RefSeq: accession NM_004119.2 (protein: RefSeq NP_004110.2).

FLT3 mRNA has a sequence as given below. However, it should be understood that even FLT3 having no mutation may differ in sequence among individuals due to polymorphism, etc.

[0028]

```
acctgcagcgcgagggcgcgccgctccaggcggcatcgcagggctgggccggcgcggc
ctggggaccccgggctccggaggccatgccggcggttggcgcgcgacggcgggccagct
gccgctgctcgttgttttttctgcaatgatatttgggactattacaaatcaagatct
gcctgtgatcaagtgtgttttaataatcataagaacaatgattcatcagtggggaa
gtcatcatcatatcccatgggtatcagaatccccggaagacctcgggtgtgcgttgag
accccagagctcagggacagtgtacgaagctgccgctgtggaagtggatgtatctgc
ttccatcacactgcaagtgtgtggcgacgccccaggggaacatttcctgtctctgggt
ctttaagcacagctccctgaattgccagccacattttgatttacaaaacagaggagt
tgtttccatgggtcattttgaaaatgacagaaaccaagctggagaatacctactttt
tattcagagtgaagctaccaattacacaatatgttttacagtgagtataagaaatac
cctgctttacacattaagaagaccttacttttagaaaaatggaaaaccaggacgcctt
ggctctgcatactctgagagcggttcagagccgatcgtggaatgggtgctttgcgattc
acaggggggaaagctgtaaagaagaaagtccagctgttggttaaaaaggaggaaaaagt
gcttcatgaattatttgggacggacataaggtgctgtgccagaaatgaactgggcag
ggaatgcaccaggctgttcacaatagatctaaatcaaactcctcagaccacattgcc
acaattatttcttaaagtaggggaacccttatggataaggtgcaaagctgttcatgt
gaaccatggattcgggctcacctgggaattagaaaacaaagcactcgaggaggggcaa
ctactttgagatgagtacctattcaacaaacagaactatgatacggattctgtttgc
```

ttttgtatcatcagtggaagaaacgacaccggatactacacttgttcctcttcaaa
gcatcccagtcfaatcagctttgggtaccatcgtagaaaagggatttataaatgctac
caattcaagtgaagattatgaaattgaccaatatgaagagttttgtttttctgtcag
gtttaaagcctacccacaaatcagatgtacgtggaccttctctcgaaaatcatttcc
ttgtgagcaaaaagggcttgataacggatacagcatatccaagttttgcaatcataa
gcaccagccaggagaatatattccatgcagaaaatgatgatgcccaatttaccaa
aatgttcacgctgaatataagaaggaaacctcaagtgctcgcagaagcatcggcaag
tcaggcgtcctgtttctcggtatggataccattaccatcttggacctggaagaagt
ttcagacaagtctcccaactgcacagaagagatcacagaaggagctctggaatagaaa
ggctaacagaaaaagtgtttggacagtggtgtcgagcagtactctaaacatgagtga
agccataaaaagggttcctggtcaagtgtgtgcatacaattcccttggcacatcttg
tgagacgatccttttaactctccaggcccccttccctttcatccaagacaacatctc
attctatgcaacaattggtgtttgtctcctcttcattgtcgttttaaccctgcta
ttgtcacaagtacaaaaagcaatttaggtatgaaagccagctacagatggtacaggt
gaccggctcctcagataatgagtacttctacgttgatttcagagaatatgaatatga
tctcaaatgggagtttccaagagaaaaatttagagtttgggaaggtagtaggatcagg
tgcttttggaaaagtgatgaacgcaacagcttatggaattagcaaaacaggagctc
aatccagggttgccgtcaaaatgctgaaagaaaaagcagacagctctgaaagagaggc
actcatgtcagaactcaagatgatgaccagctgggaagccacgagaatatgtgaa
cctgctgggggctgcacactgtcaggaccaatttacttgatttttgaatactgttg
ctatggtgatcttctcaactatctaagaagtaaaagagaaaaatttcacaggacttg
gacagagattttcaaggaaacacaatttcagtttttacccttccaatcacatcc
aaattccagcatgcctggttcaagagaagttcagatacaccggactcggatcaaat
ctcagggttcatgggaattcatttcactctgaagatgaaattgaatatgaaaacca
aaaaaggctggaagaagaggaggacttgaatgtgcttacatttgaagatcttctttg
ctttgcataatcaagttgccaagggaatggaatttctggaatttaagtcgtgtgttca
cagagacctggccgcccaggaaacgtgcttgtcaccacgggaaagtgggtgaagatatg

tgactttggattggctcgagatatcatgagtgattccaactatgttgtcaggggcaa
tgcccgctctgcctgtaaaatggatggcccccgaaagcctgtttgaaggcatctacac
cattaagagtgatgtctgggtcatatggaatattactgtgggaaatcttctcacttgg
tgtgaatccttaccctggcattccggttgatgctaacttctacaaactgattcaaaa
tggatttaaaatggatcagccattttatgctacagaagaaatatacattataatgca
atcctgctggggcttttgactcaaggaaacggccatccttccctaatttgacttcgtt
tttaggatgtcagctggcagatgcagaagaagcgatgtatcagaatgtggatggccg
tgtttcggaatgtcctcacacctacccaaaacaggcgacctttcagcagagagatgga
tttggggctactctctccgcaggctcaggctcgaagattcgtagaggaacaatttagt
tttaaggacttcatccctccacctatccctaacaggctgtagattaccaaaacaaga
ttaatttcatcactaaaagaaaatctattatcaactgctgcttcaccagacttttct
ctagaagctgtctgcgtttactcttgttttcaaagggacttttgtaaaatcaaataca
tcctgtcacaaaggcaggaggagctgataatgaactttattggagcattgatctgcat
ccaaggccttctcaggctggcttgagtgaattgtgtacctgaagtacagtatattct
tgtaaatacataaaaacaaaagcatttttgctaaggagaagctaatatgattttttaag
tctatgtttttaaataatatgtaaatttttcagctatttagtgatatattttatggg
tggaataaaaatttctactacagaattgccattattgaattatttacatgggtataa
ttagggcaagtcttaactggagttcacgaacccccctgaaattgtgcacccatagcca
cctacacattccttccagagcacgtgtgcttttaccccaagatacaaggaatgtgta
ggcagctatggttgtcacagcctaagatttctgcaacaacaggggttgattggggg
aagtttataatgaataggtgttctaccataaagagtaatacatcacctagacacttt
ggcggccttcccagactcagggccagtcagaagtaacatggaggattagtagttttca
ataaagttactcttgtccccacaaaaaaa.

[0029]

FLT3 protein has an amino acid sequence as described below. However, it should be understood that even FLT3

having no mutation may differ in sequence among individuals due to polymorphism, etc.

mpalardggqlpllvvfsamifgtitnqdlpvikcvlinhknndssvgksssympvs
espedlgcalrpqssgtvyaaaavevdsasitlqvlvdapgnisclwvfkhslncl
qphfdlqnrgvvsmvilkmtdetqageyllfiqseatnytilftvsirntllytlrrp
yfrkmenqdalvcisesvpepivewvlcdsqqgesckeespavvkkeekvlhelgtd
irccarnelgrectrlftidlnqtpqttlpqlflkvgeplwirckavhvnhgfgltw
elenkaleegnyfemstystnrtmirilfafvssvarndtgyytcssskhpsqsalv
tivekgfinatnssedyeidqyeefcfsvrfkaypqiactwtfsrksfpceqkgldn
gysiskfcnhkhqpgeyifhaenddaqftkmftlnirrkpqvlaeasasqascfsdg
yplpswtwkkcsdkspncteeitegvwnrkanrkvfqgqvwssstlnmseaikgflvk
ccaynslgtscetillnspgpfpiqdnisfyatigvcllfivvltllichkykkqf
ryesqlqmvqvtgssdneyfyvdfreyeydlkwefprenlefgkvlgsqafgkvmna
taygisktgvsigvavkmlkekadsserealmseikmmtqlgshenivnllgactls
gpiylifeycygdllnylrskrekfhrtwteifkehnfsfyptfqshpnssmpgsr
evqihpdsdqisglhgnsfhsedeieyenqkrleeeedlnvltfedllcfayqvakg
meflefkscvhrdlaarnvltvthgkvvkicdfglardimsdsnyvvrgnarlpvkwm
apeslfegiytiksdvwsygyllweifslgvnpypgipvdanfykliqngfkmdqpf
yateeyiimqscwafdsrkrsfpnltsflgcqladaeeamyqnvldgrvsecphty
qnrrpfsremdlglspqaqveds.

[0030]

In the present invention, "activating mutation of FLT3" means a mutation that causes ligand-independent activation of FLT3. Examples thereof include, but are not particularly limited to, internal tandem duplications (ITDs) in the juxtamembrane region (JM region), and point mutations D835V, D835E, D835N, D835Y and D835H that occur

in the activation loop region of FLT3. The FLT3-ITD mutation occurs mainly in exon 14 of the JM region and is also found in exon 15.

[0031]

In the present invention, "wild-type TP53" means that a gene TP53 encoding p53 protein is a gene having a sequence registered in NCBI under RefSEQ: accession NM_000546 (protein: RefSeq NP_000537).

Wild-type TP53 mRNA has a sequence given below. However, it should be understood that even TP53 having no mutation may differ in sequence among individuals due to polymorphism, etc.

```
gatgggattgggggttttcccctcccatgtgctcaagactggcgctaaaagttttgag
cttctcaaaagtctagagccaccgtccaggaggagcaggtagctgctgggctccgggga
cactttgcgttcgggctgggagcgtgctttccacgacggtgacacgcttccttgat
tggcagccagactgccttcgggtcactgccatggaggagccgcagtcagatcctag
cgtcgagccccctctgagtcaggaaacattttcagacctatggaaactacttcctga
aaacaacgttctgtcccccttgccgtcccaagcaatggatgatttgatgctgtcccc
ggacgatattgaacaatggttcactgaagaccaggtccagatgaagctcccagaat
gccagaggctgctccccccgtggccccctgcaccagcagctcctacaccggcgggccc
tgcaccagccccctcctggccccctgtcatctttctgtcccttcccagaaaacctacca
gggcagctacggttttccgtctgggcttcttgcatctctgggacagccaagtctgtgac
ttgcacgtactccccctgccctcaacaagatgttttgccaactggccaagacctgccc
tgtgcagctgtggggttgattccacacccccgcccggcaccgcgtccgcgcatggc
catctacaagcagtcacagcacatgacggaggttgtagggcgctgccccaccatga
gcgctgctcagatagcgatgggtctggccccctcctcagcatcttatccgagtggaagg
aaatttgctgtgtggagtatttgatgacagaaacacttttcgacatagtgtggtggt
```

gccctatgagccgcctgaggttggctctgactgtaccaccatccactacaactacat
gtgtaacagttcctgcatgggcggcatgaaccggaggcccatcctcaccatcatcac
actggaagactccagtggtaatctactgggacggaacagctttgaggtgcgtgtttg
tgctgtcctgggagagaccggcgccacagaggaagagaatctccgcaagaaagggga
gcctcaccacgagctgccccaggaggagcactaagcgagcactgcccacaacaccag
ctcctctccccagccaaagaagaaccactggatggagaatatctcacccttcagat
ccgtgggcgtgagcgccttcgagatggtccgagagctgaatgaggccttggaactcaa
ggatgccagggctgggaaggagccaggggggagcagggctcactccagccacctgaa
gtccaaaaagggtcagctctacctcccgccataaaaaactcatgttcaagacagaagg
gcctgactcagactgacattctccacttcttgttccccactgacagcctcccacccc
catctctccctcccctgccattttgggttttgggtctttgaacccttgcttgcaata
gggtgtgcgtcagaagcaccaggaacttccatttgctttgtcccggggctccactgaa
caagttggcctgcactgggtgttttgttgtggggaggaggatggggagtaggacatac
cagcttagattttaagggtttttactgtgagggatgtttgggagatgtaagaaatgtt
cttgacgttaagggtagttttacaatcagccacattctaggtagggggcccacttcac
cgtactaaccagggaagctgtccctcactgttgaattttctctaacttcaaggccca
tatctgtgaaatgctggcatttgcacctacctcacagagtgcattgtgagggttaat
gaaataatgtacatctggccttgaaaccaccttttattacatgggggtctagaacttg
acccccttgaggggtgcttgttccctctccctgttgggtcgggtgggttggtagtttcta
cagttgggcagctggtaggttagaggagttgtcaagtctctgctggcccagccaaa
ccctgtctgacaacctcttgggtgaaccttagtacctaaaaggaaatctcaccctatc
ccacacctggaggatttcatctcttgtatatgatgatctggatccaccaagacttg
ttttatgctcaggggtcaatttctttttctttttttttttttttttttttttttttt
gagactgggtctcgctttgttggccaggctggagtggagtggcgtgatcttggctta
ctgcagcctttgcctccccggctcgagcagtcctgcctcagcctccggagtagctgg
gaccacaggttcatgccaccatggccagccaacttttgcattgtttttagagatggg
gtctcacagtgttggccaggctgggtctcaaactcctggggtcaggcgatccacctgt

ctcagcctcccagagtgctgggattacaattgtgagccaccacgtccagctggaagg
gtcaacatctttttacattctgcaagcacatctgcattttcaccacccttcccctc
cttctcccttttttatatcccatttttatatcgatctcttattttacaataaaaacttt
gctgccacctgtgtgtctgaggggtg.

[0032]

TP53 protein has an amino acid sequence as described below. However, it should be understood that even TP53 having no mutation may differ in sequence among individuals due to polymorphism, etc.

meepqsdpsvepplsgetfsdlwklpennvlspplsqamddmlspddieqwfte
pgpdeaprmpeaappvapapaaptaapapapswplsssvpsqktyqgsygfrlgfl
hsgtaksvtctyspalnkmfcqlaktcpvqlwvdstpppgtrvramaiykqsqhmte
vvrircphhercsdsdglappqhlirvegnlrveylddrntfrhsvvvpyppevg
cttihynymcnsscmgmnrrpiltiitledssgnllgrnsfevrvcacpgrdrte
eenlrkkgephhelppgstkralpnntssspqpkkkpldgeyftlqirgrerfemfr
elnealelkdaqagkepghsrahsshlkskkgqstsrhkkmlfktegpdsd.

[0033]

In the present invention, "gene signature" means a single gene or a gene group consisting of a plurality of genes, a plurality of genes whose expression pattern is characteristic of a biological phenotype or a medical condition, such as morbidity of a certain disease, response to a certain medicament, or prognosis of a certain disease.

In the present invention, "biological sample" refers to tissues, liquids or cells isolated from an individual, or a mixture thereof. Examples thereof can include, but

are not limited to, tumor biopsy, spinal fluid, pleural fluid, intra-abdominal fluid, lymph, skin sections, blood, urine, feces, sputum, respiratory organ, intestinal tract, genitourinary tract, saliva, milk, digestive organ, and cells collected therefrom. Preferred examples of a "biological sample" can include a portion of test subject-derived resected tissues obtained during surgery performed for the purpose of treating a cancer disease, a portion of tissues collected by biopsy or the like from a test subject suspected of having a cancer disease, and cells derived from pleural fluid or intra-abdominal fluid.
[0034]

The biological sample may be protein extracts or nucleic acid extracts prepared from tissues, liquids or cells isolated from an individual, or a mixture thereof, etc. The protein extracts or the nucleic acid extracts can be prepared by use of a protein preparation method or a nucleic acid preparation method known per se in the art.
[0035]

One aspect of the present invention relates to a medicament for cancer treatment comprising (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-

b) [1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof which are administered in combination.

[0036]

In the present invention, a "medicament" comprising (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof "which are administered in combination" is a medicament based on the assumption that both the drugs are administered in combination.

[0037]

In the present invention, the "administration in combination" of (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof means that both the drugs are incorporated into

the body of a recipient in a given period. A formulation containing both the drugs in a single formulation may be administered, or the drugs may be prepared into separate formulations and separately administered. In the case of preparing separate formulations, the timing of their administration is not particularly limited. The separate formulations may be administered at the same time or may be administered at different times or on different days in a staggered manner. In the case of administering the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof at different times or on different days, the order of their administration is not particularly limited. Usually, these formulations are administered according to their respective administration methods. Therefore, these formulations may be administered in the same number of doses or may be administered in a different number of doses. Also, in the case of preparing separate formulations, the respective administration methods (administration routes) of the formulations may be the same as each other, or these formulations may be

administered by different administration methods (administration routes). Both the drugs do not have to exist at the same time in the body and may be incorporated into the body over a given period (e.g., 1 month, preferably 1 week, more preferably a few days, even more preferably 1 day). One of the active ingredients may have disappeared from the body at the time of administration of the other active ingredient.

[0038]

Examples of a dosage form of the medicament of the present invention include 1) administration of a single formulation comprising (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof, 2) concurrent administration through the same administration route of two formulations separately prepared from (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-

(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof, 3) administration in a staggered manner through the same administration route of two formulations separately prepared from (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof, 4) concurrent administration through different administration routes of two formulations separately prepared from (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof, and 5) administration in a staggered manner through different administration routes of two formulations separately prepared from (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-

1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof.

[0039]

In the present invention, the two different formulations may be in the form of a kit comprising these formulations.

[0040]

A medicament according to the present invention can contain (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and/or N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier and can be administered as various injections such as intravenous injection, intramuscular injection, and subcutaneous injection or by various methods such as oral administration or percutaneous administration. A pharmaceutically acceptable carrier means a pharmaceutically acceptable material that is involved in

transport of the compound of the present invention or a composition containing the compound of the present invention (e.g., an excipient, a diluent, an additive and a solvent) from a given organ to another organ.

[0041]

A formulation can be prepared by selecting a suitable formulation form (e.g., oral formulation or injection) depending on the administration method and using various methods conventionally used for preparing a formulation. Examples of oral formulations can include tablets, powders, granules, capsules, pills, lozenges, solutions, syrups, elixirs, emulsions and oily or aqueous suspensions. In oral administration, the free compound or a salt form may be used. An aqueous formulation can be prepared by forming an acid adduct with a pharmaceutically acceptable acid or by forming an alkali metal salt such as sodium. As an injection, a stabilizer, a preservative, a dissolving aid, and the like can be used in the formulation. After filling a solution that may contain these aids and the like in a vessel, a formulation for use may be prepared as a solid formulation by lyophilization or the like. Furthermore, one dose may be filled in one vessel, or two or more doses may be filled in a vessel.

[0042]

Examples of solid formulations include tablets, powders, granules, capsules, pills and lozenges. These

solid formulations may contain pharmaceutically acceptable additives together with a compound of the present invention. Examples of additives include fillers, extenders, binders, disintegrating agents, dissolution promoting agents, skin wetting agents and lubricants. These additives can be selected and mixed as required to prepare a formulation.

[0043]

Examples of liquid formulations include solutions, syrups, elixirs, emulsions and suspensions. Examples of additives include suspending agents and emulsifiers. These additives can be selected and mixed as required to prepare a formulation.

[0044]

Examples of pharmaceutical materials can include, but are not limited to: amino acids such as glycine, alanine, glutamine, asparagine, arginine and lysine; antimicrobial agents; antioxidants such as ascorbic acid, sodium sulfate and sodium bisulfite; buffers such as phosphate, citrate or borate buffers, sodium bicarbonate and Tris-HCl solutions; fillers such as mannitol and glycine; chelating agents such as ethylenediaminetetraacetic acid (EDTA); complexing agents such as caffeine, polyvinylpyrrolidone, β -cyclodextrin and hydroxypropyl- β -cyclodextrin; bulking agents such as glucose, mannose and dextrin; other carbohydrates such as monosaccharides and disaccharides; coloring agents;

corrigents; diluents; emulsifiers; hydrophilic polymers such as polyvinylpyrrolidone; low-molecular-weight polypeptides; salt-forming counterions; antiseptics such as benzalkonium chloride, benzoic acid, salicylic acid, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid and hydrogen peroxide; solvents such as glycerin, propylene glycol and polyethylene glycol; sugar alcohols such as mannitol and sorbitol; suspending agents; surfactants such as sorbitan ester, polysorbates such as polysorbate 20 and polysorbate 80, triton™, tromethamine, lecithin and cholesterol; stability enhancers such as sucrose and sorbitol; elasticity enhancers such as sodium chloride, potassium chloride, mannitol and sorbitol; transport agents; excipients; and/or pharmaceutical additives. The amount of these pharmaceutical materials added is preferably 0.01 to 100 times, particularly, 0.1 to 10 times the weight of the drug. The recipe of a preferred pharmaceutical composition in a formulation can be appropriately determined by those skilled in the art according to an applicable disease, an applicable administration route, etc.

[0045]

An excipient or a carrier in a pharmaceutical composition may be liquid or solid. Appropriate excipients or carriers may be other materials usually used in injectable water, physiological saline,

artificial cerebrospinal fluid, and parenteral administration. Neutral physiological saline or physiological saline containing serum albumin may be used as a carrier. The pharmaceutical composition can contain a Tris buffer of pH 7.0 to 8.5, an acetate buffer of pH 4.0 to 5.5, or a citrate buffer of pH 3.0 to 6.2. These buffers can also contain sorbitol or other compounds.

Preferred examples of the formulation of the N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-ylethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl)urea or the pharmaceutically acceptable salt thereof include formulations described in WO2014/055397.

[0046]

The medicament of the present invention can be used in cancer treatment of mammals, particularly, humans. The dose and the administration interval of the medicament of the present invention can be suitably selected depending on the site of the disease, the patient's height, body weight, sex, or medical history, according to a physician's discretion. When the medicament of the present invention is administered to a human, the dose range is approximately 0.01 to 500 mg/kg body weight, preferably, approximately 0.1 to 100 mg/kg body weight, per day with respect to one type of active ingredient. Preferably, the active ingredient of the present invention is administered to a human once a day, or the dose is divided two to four times, and

administration is repeated at an appropriate interval. Furthermore, the daily dose may exceed the above-mentioned dose at a physician's discretion, if necessary.
[0047]

For examples of the administration method of the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-ylethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof, see a method described in WO2009/061446, a method described in WO2010/132787 and a method described in US8357690. This active ingredient may be administered once a day for 1 week, 2 weeks, 3 weeks, 4 weeks or 5 weeks. Preferred examples thereof include a method of continuously administering 12 to 450 mg, for example, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 90 mg, 135 mg, 200 mg, 300 mg or 450 mg, of this agent for 28 days, a method of continuously administering the above-mentioned dose for 8 to 21 days together with an additional anticancer agent, and a method of continuously administering the above-mentioned dose for 4 to 17 days together with an additional anticancer agent.

[0048]

The type of cancer to be treated is not particularly limited as long as the cancer is confirmed to be sensitive to treatment by combined use of the present invention. Examples thereof include blood cancer (leukemia, lymphoma or multiple myeloma), brain tumor,

head and neck cancer, esophageal cancer, stomach cancer, appendix cancer, colon cancer, anus cancer, gallbladder cancer, cholangiocarcinoma cancer, pancreatic cancer, gastrointestinal stromal tumor, lung cancer, liver cancer, mesothelioma, thyroid cancer, renal cancer, prostate cancer, neuroendocrine tumor, melanoma, breast cancer, endometrial cancer, cervical cancer, ovarian cancer, osteosarcoma, soft tissue sarcoma, Kaposi's sarcoma, myosarcoma, bladder cancer and testicular cancer. Among them, leukemia, particularly, acute myeloid leukemia (AML) is preferred. Leukemia having an activating mutation of FLT3 is more preferred, and acute myeloid leukemia having a FLT3-ITD mutation is particularly preferred.

[0049]

In the present invention, a method for detecting the "activating mutation of FLT3" includes a method of detecting a mutation on genomic DNA as well as, when the mutation on the genomic DNA is reflected to base change in a transcription product or amino acid change in a translation product, a method of detecting this change in the transcription product or translation product (i.e., indirect detection), and a method based on detection of phosphorylated FLT3 because activation of FLT3 involves a rise in phosphorylation level.

[0050]

In a preferred embodiment, a method for detecting a mutation includes a method of directly determining the nucleotide sequence of a gene region in a test subject-derived biological sample to thereby detect a mutation. In the present invention, the "FLT3 gene region" means a given region on genomic DNA containing the FLT3 gene. The region also contains the expression control regions (e.g., a promoter region and an enhancer region) of the FLT3 gene, the 3'-terminal untranslated region of the FLT3 gene, and the like. A mutation in these regions can influence, for example, the transcription activity of the FLT3 gene.

[0051]

In this method, first, a DNA sample is prepared from a test subject-derived biological sample. Examples of the DNA sample include a genomic DNA sample and a cDNA sample prepared from RNA by reverse transcription.

[0052]

The method for extracting genomic DNA or RNA from the biological sample is not particularly limited, and an approach known in the art can be appropriately selected for use. Examples of the method for extracting genomic DNA include a SDS phenol method (method which involves denaturing proteins in tissues preserved in a urea-containing solution or ethanol, using a proteolytic enzyme (proteinase K), a surfactant (SDS) and phenol, and extracting DNA by precipitation from the tissues using

ethanol), and DNA extraction methods using Clean Columns(R) (manufactured by Nexttec Biotechnologie GmbH), AquaPure(R) (manufactured by Bio-Rad Laboratories, Inc.), ZR Plant/Seed DNA Kit (manufactured by Zymo Research Corp.), Aqua Genomic Solution(R) (manufactured by MoBiTec GmbH), prepGEM(R) (manufactured by ZyGEM NZ Ltd.) or BuccalQuick(R) (manufactured by TrimGen Corp.). Examples of the method for extracting RNA include extraction methods using phenol and a chaotropic salt (more specifically, extraction methods using a commercially available kit such as TRIzol (manufactured by Invitrogen Corp.) or ISOGEN (manufactured by Wako Pure Chemical Industries, Ltd.)), and methods using other commercially available kits (RNAPrep Total RNA Extraction Kit (manufactured by Beckman Coulter, Inc.), RNeasy Mini (manufactured by Qiagen N.V.), RNA Extraction Kit (manufactured by Pharmacia Biotech Inc.), etc.). Examples of reverse transcriptase for use in the preparation of cDNA from the extracted RNA include, but are not particularly limited to, reverse transcriptase derived from retrovirus such as RAV (Rous associated virus) or AMV (avian myeloblastosis virus), and reverse transcriptase derived from mouse retrovirus such as MMLV (Moloney murine leukemia virus).

[0053]

In this embodiment, DNA containing a mutation site in the FLT3 gene region is subsequently isolated, and the

nucleotide sequence of the isolated DNA is determined. The isolation of the DNA can be performed by, for example, PCR using a pair of oligonucleotide primers designed to flank the mutation in the FLT3 gene region, and the genomic DNA or the RNA as a template. The determination of the nucleotide sequence of the isolated DNA can be performed by a method generally known to those skilled in the art, such as the Maxam-Gilbert method or Sanger method.

[0054]

The determined nucleotide sequence of the DNA or the cDNA can be compared with a control (e.g., the nucleotide sequence of DNA or cDNA derived from non-cancer tissues of the same test subject) to thereby determine the presence or absence of the mutation in the FLT3 gene region in the cancer cells of the test subject.

[0055]

The method for detecting a mutation in the FLT3 gene region can be performed by various methods capable of detecting a mutation, in addition to the method of directly determining the nucleotide sequence of DNA or cDNA.

[0056]

For example, in one of the methods, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, a reporter fluorescent dye- and quencher fluorescent dye-labeled oligonucleotide probe having a

nucleotide sequence complementary to a nucleotide sequence containing the mutation in the FLT3 gene region is prepared. Then, the oligonucleotide probe is hybridized to the DNA sample. Then, the nucleotide sequence containing the mutation in the FLT3 gene region is amplified using the DNA sample hybridized with the oligonucleotide probe as a template. Then, fluorescence emitted by the reporter fluorescent dye through the decomposition of the oligonucleotide probe associated with the amplification is detected. Subsequently, the detected fluorescence is compared with a control. Examples of such a method include a double die probe method, so-called TaqMan(R) probe method.

[0057]

In an alternative method, a DNA or cDNA sample is prepared from the biological sample. Subsequently, the nucleotide sequence containing the mutation in the FLT3 gene region is amplified using the DNA sample as a template in a reaction system containing an intercalator that emits fluorescence upon insertion between two strands of DNA. Then, the temperature of the reaction system is changed, and variation in the intensity of the fluorescence emitted by the intercalator is detected. The detected variation in the intensity of the fluorescence caused by the change in the temperature is compared with a control. Examples of such a method include a high resolution melting (HRM) method.

[0058]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, DNA containing a mutation site in the FLT3 gene region is amplified. Then, the amplified DNA is cleaved with restriction enzymes. Subsequently, the DNA fragments are separated according to their sizes. Subsequently, the detected sizes of the DNA fragments are compared with a control. Examples of such a method include a method using restriction fragment length polymorphism (RFLP) and PCR-RFLP.

[0059]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, DNA containing a mutation site in the FLT3 gene region is amplified. Then, the amplified DNA is dissociated into single-stranded DNA. Subsequently, the single-stranded DNA thus obtained by dissociation is separated on a non-denaturing gel. The mobility of the separated single-stranded DNA on the gel is compared with a control. Examples of such a method include PCR-SSCP (single-strand conformation polymorphism).

[0060]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, DNA containing a mutation site in the FLT3 gene region is amplified. Then, the amplified DNA is

separated on a gel in which the concentration of a DNA denaturant is gradually elevated. Subsequently, the mobility of the separated DNA on the gel is compared with a control. Examples of such a method include denaturant gradient gel electrophoresis (DGGE).

[0061]

A further alternative method is a method using DNA containing a mutation site in the FLT3 gene region prepared from the biological sample, and a substrate with immobilized oligonucleotide probes hybridizing to the DNA. Examples of such a method include a DNA array method.

[0062]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Also, an "oligonucleotide primer having a nucleotide sequence complementary to the base downstream by one base from the base at the mutation site in the FLT3 gene region, and a downstream nucleotide sequence thereof" is prepared. Subsequently, a ddNTP primer extension reaction is performed using the DNA as a template and the primer. Subsequently, the primer extension reaction product is applied to a mass spectrometer to conduct mass spectrometry. Subsequently, the genotype is determined from the mass spectrometry results. Then, the determined genotype is compared with a control. Examples of such a method include MALDI-TOF/MS.

[0063]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, an oligonucleotide probe consisting of 5' - "a nucleotide sequence complementary to the base at the mutation site in the FLT3 gene region, and an upstream nucleotide sequence thereof" - "a nucleotide sequence hybridizing neither to the base downstream by one base from the mutation site in the FLT3 gene region, nor to a downstream nucleotide sequence thereof" - 3' (flap) is prepared. Also, an "oligonucleotide probe having a nucleotide sequence complementary to the base at the mutation site in the FLT3 gene region, and a downstream nucleotide sequence thereof" is prepared. Subsequently, the prepared DNA is hybridized to these two types of oligonucleotide probes. Subsequently, the hybridized DNA is cleaved with a single-stranded DNA-cleaving enzyme to release the flap. Examples of the single-stranded DNA-cleaving enzyme include, but are not particularly limited to, cleavase. In this method, a fluorescent reporter- and fluorescent quencher-labeled oligonucleotide probe having a sequence complementary to the flap is then hybridized to the flap. Subsequently, the intensity of the generated fluorescence is measured. Then, the measured intensity of the fluorescence is compared with a control. Examples of such a method include an invader method.

[0064]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, DNA containing a mutation site in the FLT3 gene region is amplified. Then, the amplified DNA is dissociated into single strands, and only one of the single strands of the dissociated DNA is separated. Subsequently, an extension reaction is performed one by one from a base close to the base at the mutation site in the FLT3 gene region. Pyrophosphoric acid generated during this reaction is enzymatically allowed to develop light. The intensity of the light is measured. The measured intensity of the fluorescence is compared with a control. Examples of such a method include pyrosequencing.

[0065]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, DNA containing a mutation site in the FLT3 gene region is amplified. Subsequently, an "oligonucleotide primer having a nucleotide sequence complementary to the base downstream by one base from the base at the mutation site in the FLT3 gene region, and a downstream nucleotide sequence thereof" is prepared. Subsequently, a single-base extension reaction is performed using the amplified DNA as a template and the prepared primer in the presence of fluorescently labeled nucleotides. Then, the degree of polarization of

fluorescence is measured. Subsequently, the measured degree of polarization of fluorescence is compared with a control. Examples of such a method include the AcycloPrime method.

[0066]

In a further alternative method, a DNA or cDNA sample is first prepared from the biological sample. Subsequently, DNA containing a mutation site in the FLT3 gene region is amplified. Subsequently, an "oligonucleotide primer having a nucleotide sequence complementary to the base downstream by one base from the base at the mutation site in the FLT3 gene region, and a downstream nucleotide sequence thereof" is prepared. Subsequently, a single-base extension reaction is performed using the amplified DNA as a template and the prepared primer in the presence of fluorescently labeled nucleotides. Subsequently, the base species used in the single-base extension reaction are determined. Then, the determined base species are compared with a control. Examples of such a method include the SNuPE method.

[0067]

Provided that the mutation results in an amino acid change (e.g., substitution, deletion or insertion) in the FLT3 protein, a sample prepared from the biological sample may be a protein. In such a case, a method using a molecule (e.g., an antibody) specifically binding to a

site having the amino acid change ascribable to the mutation can be used for detecting the mutation.

Since the activating mutation of FLT3 elevates the phosphorylation level of FLT3, the activating mutation of FLT3 can also be detected by the quantification of phosphorylated FLT3. A phosphorylated protein measurement method known in the art can be used as a method for quantitatively measuring the phosphorylated FLT3 protein. For example, various methods using antibodies against the phosphorylated FLT3 protein can be utilized. Specific examples thereof can include Western blotting, immunoprecipitation, enzyme-linked immunosorbent assay (ELISA) and radioimmunoassay (RIA).
[0068]

A humanized antibody, a mouse antibody, a rat antibody, a rabbit antibody, a sheep antibody or the like can be appropriately used as an antibody against the mutated FLT3 protein or the phosphorylated FLT3 protein as long as the antibody is directed to the mutated FLT3 protein or the phosphorylated FLT3 protein as an antigen and specifically binds to the antigen. The antibody may be a polyclonal antibody or may be a monoclonal antibody. A monoclonal antibody is preferred from the viewpoint that homogeneous antibodies can be stably produced. The polyclonal antibody and the monoclonal antibody can be prepared by methods well known to those skilled in the

art. A desired antibody can also be selected, for use, from commercially available antibodies.

[0069]

A hybridoma producing the monoclonal antibody can basically be prepared by use of a technique known in the art as follows: the antigen of interest or cells expressing the antigen of interest is used as a sensitizing antigen, and a desired animal is immunized with this sensitizing antigen according to a conventional immunization method. The obtained immunocytes are fused with known parent cells by a conventional cell fusion method. Then, cells producing the desired monoclonal antibody (hybridoma cells) can be selected by a conventional screening method. The preparation of the hybridoma can be carried out according to, for example, the method of Millstein ("Methods of Enzymology", 1981, Vol. 73, p. 3-46).

[0070]

In this context, the phosphorylated FLT3 protein or a fragment thereof can be used as the antigen for preparing the monoclonal antibody. Those skilled in the art can easily obtain the phosphorylated FLT3 protein or the fragment thereof according to a method described in a book, for example, Sambrook ed., "Molecular Cloning: A Laboratory Manual", the 2nd edition, Vol. 1-3, Cold Spring Harbor Laboratory Press, NY, 1989.

[0071]

The protein or the fragment thereof and the antibody may be immobilized on a support and used for quantifying the phosphorylated FLT3 protein. The support is not limited as long as the support permits immobilization of proteins. General examples thereof can include: inorganic materials such as glass plates, silicon wafers and resins; natural polymer materials including nitrocellulose; and synthetic polymer materials including nylon and polystyrene.

[0072]

More specific examples of the method for detecting the activating mutation of FLT3 include a method for detecting a FLT3-ITD mutation described in WO9817808 and its corresponding U.S. patent US6846630. This method can be performed by using a detection kit commercially available from Takara Bio Inc., etc.

[0073]

A method of performing RT-PCR using mRNA obtained from the test subject-derived biological sample, followed by capillary electrophoresis can also be used as another similar method (Leukemia, 2005, 19, 1479-1482).

Specific examples of the method for detecting the phosphorylated FLT3 protein include a method described in WO2010/054185.

From another viewpoint, cancer sensitive to an MDM2 inhibitor and that having wild-type TP53 are preferable as the types of the cancer to be treated.

Various approaches mentioned above as methods for confirming a mutation in FLT3 can be similarly utilized as methods for confirming TP53 to be wild-type. More specific examples thereof include a microarray method using a probe specific for a mutated DNA sequence (AmpliChip p53, Roche Molecular Systems, Inc., etc.), PCR using a probe specific for a mutated DNA sequence (qBiomarker Somatic Mutation PCR Arrays, Qiagen N.V., etc.), a method of reading the p53 gene sequence using a Sanger sequencer, and a method of reading the p53 gene sequence using a next-generation sequencer (TruSeq Amplicon - Cancer Panel, Illumina, Oncomine(R) Cancer Research Panel, Life Technologies Corp., etc.).

[0074]

A method using a gene signature can also be preferably used as a method for predicting sensitivity to an MDM2 inhibitor. Examples of the gene signature for predicting sensitivity to an MDM2 inhibitor include, but are not particularly limited to, a gene group described in WO2014/020502. More specifically, a gene group comprising at least one gene selected from the group consisting of MDM2, CDKN1A, ZMAT3, DDB2, FDXR, RPS27L, BAX, RPM2B, SESN1, CCNG1, XPC, TNFSF10B and AEN (the gene group may comprise all of these genes) can be preferably used. Other examples thereof include a gene group described in WO2015/000945. More specifically, a gene group comprising at least one gene selected from the

group consisting of BAX, RPS27L, EDA2R, XPC, DDB2, FDXR, MDM2, CDKN1A, TRIAP1, BBC3, CCNG1, TNFRSF10B and CDKN2A (the gene group may comprise all of these genes) can be preferably used. The number of genes contained in the gene group is not limited. A sensitive signature that allows the cancer to be confirmed as sensitive to an MDM2 inhibitor when the gene contained in the gene signature is highly expressed can be preferably used.

[0075]

The medicament according to the present invention may be used in combination with an additional anti-tumor agent. Examples thereof include anti-tumor antibiotics, anti-tumor plant constituents, BRMs (biological response modifiers), hormones, vitamins, anti-tumor antibodies, molecular target drugs, alkylating agents, metabolic antagonists and other anti-tumor agents.

[0076]

More specifically, examples of alkylating agents include: alkylating agents such as nitrogen mustard, nitrogen mustard N-oxide, bendamustine and chlorambucil; aziridine alkylating agents such as carboquone and thiotepa; epoxide alkylating agents such as dibromomannitol and dibromodulcitol; nitrosourea alkylating agents such as carmustine, lomustine, semustine, nimustine hydrochloride, streptozocin, chlorozotocin and ranimustine; and busulfan, improsulfan tosylate, temozolomide and dacarbazine.

[0077]

Various examples of metabolic antagonists include: purine metabolic antagonists such as 6-mercaptopurine, 6-thioguanine and thioinosine; pyrimidine metabolic antagonists such as fluorouracil, tegafur, tegafur-uracil, carmofur, doxifluridine, broxuridine, cytarabine and enocitabine; and folic acid metabolic antagonists such as methotrexate and trimetrexate.

[0078]

Examples of anti-tumor antibiotics include: mitomycin C, bleomycin, peplomycin, daunorubicin, aclarubicin, doxorubicin, idarubicin, pirarubicin, THP-adriamycin, 4'-epidoxorubicin and epirubicin; and chromomycin A3 and actinomycin D.

[0079]

Examples of anti-tumor plant constituents and their derivatives include: vinca alkaloids such as vindesine, vincristine and vinblastine; taxanes such as paclitaxel, docetaxel and cabazitaxel; and epipodophyllotoxins such as etoposide and teniposide.

[0080]

Examples of BRMs include tumor necrosis factors and indomethacin.

[0081]

Examples of hormones include hydrocortisone, dexamethasone, methylprednisolone, prednisolone, prasterone, betamethasone, triamcinolone, oxymetholone,

nandrolone, metenolone, fosfestrol, ethinylestradiol, chlormadinone, mepitiostane and medroxyprogesterone.

[0082]

Examples of vitamins include vitamin C and vitamin A.

[0083]

Examples of anti-tumor antibodies and molecular target drugs include trastuzumab, rituximab, cetuximab, nimotuzumab, denosumab, bevacizumab, infliximab, ipilimumab, nivolumab, pembrolizumab, avelumab, pidilizumab, atezolizumab, ramucirumab, imatinib mesylate, dasatinib, gefitinib, erlotinib, osimertinib, sunitinib, lapatinib, dabrafenib, trametinib, cobimetinib, pazopanib, palbociclib, panobinostat, sorafenib, crizotinib, vemurafenib, ibrutinib, bortezomib, carfilzomib, ixazomib and gilteritinib.

[0084]

Examples of other anti-tumor agents include cisplatin, carboplatin, oxaliplatin, tamoxifen, letrozole, anastrozole, exemestane, toremifene citrate, fulvestrant, bicalutamide, flutamide, mitotane, leuprorelin, goserelin acetate, camptothecin, ifosfamide, cyclophosphamide, melphalan, L-asparaginase, aceglatone, sizofiran, picibanil, procarbazine, pipobroman, neocarzinostatin, hydroxyurea, ubenimex, azacitidine, decitabine, thalidomide, lenalidomide, pomalidomide, eribulin, tretinoin and krestin.

[0085]

Another aspect of the present invention relates to a method for predicting responsiveness to treatment of cancer with (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof in combination, comprising using a test subject-derived biological sample, detecting the presence or absence of an activating mutation of FLT3 contained in the biological sample, and confirming the test subject having the detected activating mutation of FLT3 to be responsive to the treatment of cancer with the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof in combination.

An alternative aspect of the present invention relates to a method for selecting a subject for treatment

of cancer with (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof and N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof in combination, comprising using a test subject-derived biological sample, detecting the presence or absence of an activating mutation of FLT3 in the biological sample, and selecting the test subject having the detected activating mutation of FLT3 as the subject for the treatment of cancer with the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-(4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof in combination.

[0086]

The present invention further relates to a method for predicting responsiveness to treatment of cancer or a method for selecting a subject for treatment of cancer,

wherein the activating mutation of FLT3 is a FLT3-ITD mutation.

[0087]

A method for collecting the test subject-derived biological sample and a method for detecting the activating mutation of FLT3 or the FLT3-ITD mutation in the biological sample are as mentioned above.

Examples

[0088]

Hereinafter, the present invention will be specifically explained with reference to the Examples given below. However, the present invention is not limited to these examples, and they should not be construed in any limitative way.

(Test Example 1 Study on *in vivo* effect of combined use of Compound A and quizartinib)

Human acute myeloid leukemia cell line MOLM-13 cells having a FLT3-ITD mutation and wild-type TP53 were suspended to 5×10^7 cells/mL using phosphate-buffered saline. 0.1 mL of the prepared cell suspension was subcutaneously transplanted to each NOD-SCID mouse (male, 5 to 7 weeks old). On 6 days after the tumor inoculation, after confirmation that the average tumor volume exceeded 100 mm³, the mice were randomized (N = 6/group) on the basis of their tumor volume values. 25 mg/kg or 50 mg/kg Compound A or 0.5 mg/kg or 1 mg/kg quizartinib (LC

Laboratories) was administered by oral gavage to the mice. For a combined use group, 25 mg/kg or 50 mg/kg Compound A and 0.5 mg/kg or 1 mg/kg quizartinib were orally administered sequentially by forced administration. The administration was performed once a day for 5 consecutive days (6 to 10 days after the tumor inoculation) from the date of randomization (6 days after the tumor inoculation), and after a 2-day drug holiday, performed once a day for 4 consecutive days (13 to 16 days after the tumor inoculation). The length (mm) and width (mm) of tumor were measured over time using an electronic digital caliper. Tumor growth inhibition % (TGI%) on the date of assessment (17 days after the tumor inoculation) calculated according to calculation formula (4) shown below was used in evaluation. Also, the body weights were measured over time using an automatic balance for small animals, and body weight change % was calculated according to calculation formula (5) shown below to assess the influence of drug administration on the body weights. In addition, the results of the last body weight measurement were used in dose calculation.

[0089]

$$\text{TGI (\%)} = (1 - A / B) \times 100 \dots (4)$$

A: Average tumor volume of the compound-administered group on the date of assessment (*)

B: Average tumor volume of the untreated control group on the date of assessment (*)

*: The tumor volume was calculated according to $1/2 \times [\text{Major axis of tumor}] \times [\text{Minor axis of tumor}] \times [\text{Minor axis of tumor}]$.

[0090]

Body weight change (%) = Average body weight change % of the individuals ... (5)

Body weight change % of each individual = $(1 - \text{BWn} / \text{BW}_s) \times 100$

BWn: Body weight on day n

BW_s: Body weight on the start day of administration

The results are shown in Figure 1 and Tables 1 to 3.

[0091]

[Table 1]

Group	TGI (%)
Compound A 25 mg/kg	38
Compound A 50 mg/kg	82
Compound A 25 mg/kg + Quizartinib 0.5 mg/kg	57
Compound A 50 mg/kg + Quizartinib 0.5 mg/kg	85
Compound A 25 mg/kg + Quizartinib 1 mg/kg	85
Compound A 50 mg/kg + Quizartinib 1 mg/kg	97
Quizartinib 0.5 mg/kg	29
Quizartinib 1 mg/kg	61

[0092]

[Table 2]

Estimated tumor volume (mm ³)							
Group (N=6)		Days after tumor inoculation					
		6	8	10	13	15	17
1.Untreated	average	163	302	625	1227	1667	2667
	SE	12	39	64	136	152	195
2.Compound A 25 mg/kg	average	169	266	425	812	983	1661
	SE	14	24	41	78	63	104
3.Compound A 50 mg/kg	average	162	195	210	325	375	467
	SE	12	15	23	37	49	64
4.Compound A 25 mg/kg + Quizartinib 0.5 mg/kg	average	169	168	220	673	753	1149
	SE	14	10	21	50	66	83
5.Compound A 50 mg/kg + Quizartinib 0.5 mg/kg	average	164	134	106	290	297	387
	SE	13	12	11	37	47	81
6.Compound A 25 mg/kg + Quizartinib 1 mg/kg	average	165	132	98	314	318	387
	SE	12	12	11	45	40	43
7.Compound A 50 mg/kg + Quizartinib 1 mg/kg	average	171	111	62	116	96	67
	SE	15	8	7	18	17	14
8.Quizartinib 0.5 mg/kg	average	167	239	477	897	1212	1893
	SE	13	28	36	61	96	154
9.Quizartinib 1 mg/kg	average	163	168	203	593	663	1039
	SE	13	11	18	39	57	67

[0093]

[Table 3]

Body weight change (%)								
Group (N=6)		Days after tumor inoculation						
		6	8	10	13	15	17	
1.Untreated	average	0.0	2.6	6.0	10.8	15.2	19.2	
	SD	0.0	1.2	1.3	1.0	2.0	2.8	
2.Compound A 25 mg/kg	average	0.0	2.1	3.1	6.4	8.0	10.6	
	SD	0.0	1.1	1.9	1.9	2.5	2.7	
3.Compound A 50 mg/kg	average	0.0	1.5	2.7	5.5	6.8	5.8	
	SD	0.0	1.4	2.3	2.5	3.5	3.4	
4.Compound A 25 mg/kg + Quizartinib 0.5 mg/kg	average	0.0	0.3	1.4	5.1	5.5	6.1	
	SD	0.0	2.1	2.9	3.1	4.1	4.2	
5.Compound A 50 mg/kg + Quizartinib 0.5 mg/kg	average	0.0	1.0	0.3	2.9	4.2	3.6	
	SD	0.0	2.0	3.1	3.8	4.6	4.2	
6.Compound A 25 mg/kg + Quizartinib 1 mg/kg	average	0.0	1.1	0.7	4.7	2.8	3.3	
	SD	0.0	2.4	3.7	2.8	4.1	4.5	
7.Compound A 50 mg/kg + Quizartinib 1 mg/kg	average	0.0	-0.6	-0.3	2.5	2.6	3.2	
	SD	0.0	2.6	4.8	3.3	5.3	5.1	
8.Quizartinib 0.5 mg/kg	average	0.0	4.2	4.2	8.5	9.6	10.7	
	SD	0.0	2.9	2.6	2.1	1.6	2.8	
9.Quizartinib 1 mg/kg	average	0.0	0.2	1.8	4.6	5.1	5.8	
	SD	0.0	1.7	1.9	3.2	3.4	3.0	

Free Text of Sequence Listing

[0094]

SEQ ID NO: 1: FLT3 mRNA encoding FLT3 protein (SEQ ID NO: 2).

SEQ ID NO: 2: Amino acid sequence of the FLT3 protein.

SEQ ID NO: 3: TP53 mRNA encoding TP53 protein (SEQ ID NO: 4).

SEQ ID NO: 4: Amino acid sequence of the TP53 protein.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. Use of (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for use in the treatment of cancer in combination with N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof.

2. Use of N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or a pharmaceutically acceptable salt thereof for the manufacture of a medicament for use in the treatment of cancer in combination with (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or a pharmaceutically acceptable salt thereof.

3. Use according to claim 1 or claim 2, wherein the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-ylethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof are separately contained as active ingredients in different formulations for administration at the same time or different times.

4. Use according to claim 1 or claim 2, wherein the (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide or the pharmaceutically acceptable salt thereof and the N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-ylethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea or the pharmaceutically acceptable salt thereof are contained in a single formulation.

5. Use according to any one of claims 1 to 4, wherein the respective salts of the compounds are (3'R,4'S,5'R)-N-[(3R,6S)-6-carbamoyltetrahydro-2H-pyran-3-yl]-6"-

chloro-4'-(2-chloro-3-fluoropyridin-4-yl)-4,4-dimethyl-2"-oxo-1",2"-dihydrodispiro[cyclohexane-1,2'-pyrrolidine-3',3"-indole]-5'-carboxamide p-toluenesulfonate and N-(5-tert-butyl-isoxazol-3-yl)-N'-{4-[7-(2-morpholin-4-yl-ethoxy)imidazo[2,1-b][1,3]benzothiazol-2-yl]phenyl}urea dihydrochloride.

6. Use according to any one of claims 1 to 5, wherein the cancer is blood cancer, leukemia, lymphoma, multiple myeloma, brain tumor, head and neck cancer, esophageal cancer, stomach cancer, appendix cancer, colon cancer, anus cancer, gallbladder cancer, cholangiocarcinoma cancer, pancreatic cancer, gastrointestinal stromal tumor, lung cancer, liver cancer, mesothelioma, thyroid cancer, renal cancer, prostate cancer, neuroendocrine tumor, melanoma, breast cancer, endometrial cancer, cervical cancer, ovarian cancer, osteosarcoma, soft tissue sarcoma, Kaposi's sarcoma, myosarcoma, bladder cancer or testicular cancer.

7. Use according to any one of claims 1 to 5, wherein the cancer is leukemia.

8. Use according to any one of claims 1 to 5, wherein the cancer is leukemia having an activating mutation of FLT3.

9. Use according to any one of claims 1 to 5, wherein the cancer is acute myeloid leukemia (AML).

10. Use according to any one of claims 1 to 5, wherein the cancer is acute myeloid leukemia (AML) having a FLT3-ITD mutation.



11. Use according to any one of claims 1 to 5, wherein the cancer is cancer having wild-type TP53.

12. Use according to any one of claims 1 to 5, wherein the cancer is cancer confirmed to be MDM2 inhibitor-sensitive using a gene signature.

13. Use according to any one of claims 1 to 5, wherein the cancer is cancer having an activating mutation of FLT3 and having wild-type TP53.

14. Use according to any one of claims 1 to 5, wherein the cancer is cancer having an activating mutation of FLT3 and determined to be MDM2 inhibitor-sensitive using a gene signature.

[Figure 1-1]



[Figure 1-2]

