



(51) International Patent Classification:

C07D 223/20 (2006.01) A61P 25/00 (2006.01)
C07D 223/22 (2006.01) A61P 25/18 (2006.01)
A61K 31/55 (2006.01) A61P 25/28 (2006.01)

(21) International Application Number:

PCT/IB2013/050711

(22) International Filing Date:

28 January 2013 (28.01.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1782/CHE/2012 7 May 2012 (07.05.2012) IN

(72) Inventor; and

(71) Applicant : KANDULA, Mahesh [IN/IN]; Door No. 11-61, Samalkot Mandal, East Godavari Dist, G.Medapadu 533 434, Andhra Pradesh (IN).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))



WO 2013/167985 A1

(54) Title: COMPOSITIONS AND METHODS FOR THE TREATMENT OF NEUROLOGICAL DISORDERS

(57) Abstract: The invention relates to the compounds of formula (I) and formula (1A) or its pharmaceutical acceptable salts, as well as polymorphs, solvates, enantiomers, stereoisomers and hydrates thereof. The pharmaceutical compositions comprising an effective amount of compounds of formula (I) and formula (IA); and methods for treating or preventing neurological diseases may be formulated for oral, buccal, rectal, topical, transdermal, transmucosal, intravenous, parenteral administration, syrup, or injection. Such compositions may be used to treatment, of epilepsy, bipolar disorder, trigeminal neuralgia, attention-deficit, hyperactivity disorder (ADHD), schizophrenia, neuropathic pain, seizures, bipolar disorder, mania, phantom limb syndrome, complex regional pain syndrome, paroxysmal extreme pain disorder, neuromyotonia, intermittent explosive disorder, borderline personality disorder. Myotonia congenita and post-traumatic stress disorder.

**COMPOSITIONS AND METHODS FOR THE TREATMENT OF
NEUROLOGICAL DISORDERS**

PRIORITY

[0001] The present application claims the benefit of Indian Provisional Patent Application No. 1782/CHE/2012 filed on 07-May-2012, the entire disclosure of which is relied on for all purposes and is incorporated into this application by reference.

FIELD OF THE INVENTION

[0002] This disclosure generally relates to compounds and compositions for the treatment of neurological disorders. More particularly, this invention relates to treating subjects with a pharmaceutically acceptable dose of compounds, crystals, esters, salts, enantiomers, stereoisomers, polymorphs, hydrates, prodrugs, or mixtures thereof.

BACKGROUND OF THE INVENTION

[0003] Disorders such as the periodic paralyses, nondystrophic myotonias, episodic ataxias, paroxysmal dyskinesias, long QT syndrome, migraine headache, and epilepsy all share the feature of being episodic in nature. Affected individuals are often completely healthy between attacks. Stress and fatigue precipitate attacks in all of these diseases, and various dietary factors can also contribute to attack onset. The drugs used to treat these disorders overlap significantly.

[0004] Carbonic anhydrase inhibitors are effective for many patients with periodic paralysis, episodic ataxia, and migraine headache. Mexiletine hydrochloride, an effective antiarrhythmic medication, can be beneficial in treating myotonia in patients with paramyotonia congenita. The anticonvulsant carbamazepine is an extremely efficacious drug for treating the episodic movements of paroxysmal kinesigenic dyskinesia. All of these disorders have a tendency to begin in infancy or childhood and to worsen through adolescence and young adult life. In some cases, they decrease in severity and frequency in middle to late adult life.

[0005] Epilepsy and stroke are the 2 most common neurological disorders: at any one time 7 in 1000 people in the general population have epilepsy. Epilepsy usually begins in childhood, potentially impeding education, employment, social relationships and development of a sense of self-worth. Prompt, accurate diagnosis with appropriate social and medical management will optimize the situation. A family physician, in conjunction with a neurologist, can ascertain (a) if the episodes represent epileptic seizures and (b) if so, which epileptic syndrome they represent.

[0006] A harmonized partnership between family physician and neurologist will facilitate the recognition and care of epileptic disorders. As the role of the family physician in the care of patients with epilepsy increases, the principles delineated in this article will be ever more utilized. Trigeminal neuralgia is defined as sudden, usually unilateral, severe, brief, stabbing recurrent episodes of pain within the distribution of one or more branches of the trigeminal nerve, which has a profound effect on quality of life. The diagnosis is made on history alone, and time needs to be taken to elicit the key features and differentiate from toothache or one of the trigeminal autonomic cephalalgias. Most trigeminal neuralgia is idiopathic, but a small percentage is due to secondary causesd for example, tumours or multiple sclerosis which can be picked up on CT or MRI.


[0007] Recently published international guidelines suggest that carbamazepine and oxcarbazepine are the first-line drugs. There is limited evidence for the use of lamotrigine and baclofen. If there is a decrease in efficacy or tolerability of medication, surgery needs to be considered.

[0008] Managing acute pathology of often relies on the addressing underlying pathology and symptoms of the disease. There is currently a need in the art for new compositions to treatment of Neurological disorders.

SUMMARY OF THE INVENTION


[0009] The present invention provides compounds, compositions containing these compounds and methods for using the same to treat, prevent and/or ameliorate the effects of the conditions such as Neurological disorders.

[0010] The invention herein provides compositions comprising of formula I or pharmaceutical acceptable salts thereof. The invention also provides pharmaceutical compositions comprising one or more compounds of formula I or enantiomers of formula I or intermediates thereof and one or more of pharmaceutically acceptable carriers, vehicles or diluents. These compositions may be used in the treatment of Neurological disorders and its associated complications.



Formula I

[0011] In certain embodiments, the present invention relates to the compounds and compositions of formula I, or pharmaceutically acceptable salts thereof,





Formula I

[0012] Wherein,


R¹ each independently represents Oxygen (O), D, Hydrogen (H) or Flourine (F);







R³ independently represents H, D,






R^2 independently represents








a is independently 2,3 or 7;

each b is independently 3, 5 or 6;

e is independently 1, 2 or 6;


c and d are each independently H, D, -OH, -OD, C₁-C₆-alkyl, -NH₂ or -COCH₃.

[0013] In the illustrative embodiments, examples of compounds of formula I are as set forth below:



(1-1)

[0014] In certain embodiments, the present invention relates to the compounds and compositions of formula IA in S- enantiomer or R- enantiomer form or pharmaceutically acceptable salts thereof,





Active Stereoisomer of Formula IA

[0015] Wherein,


R¹ each independently represents Oxygen (O), D, Hydrogen (H) or Flourine (F),


R³ each independently represents D,






R² each independently represents Acetyl (CH₃CO-),








a is independently 2,3 or 7;

each b is independently 3, 5 or 6;

e is independently 1, 2 or 6;

c and d are each independently H, D, -OH, -OD, C₁-C₆-alkyl, -NH₂ or -COCH₃.

[0016] In the illustrative embodiments, examples of compounds of active stereoisomer form of formula IA are as set forth below:



Active stereoisomer form of formula IA (1-2)

[0017] Herein the application also provides a kit comprising any of the pharmaceutical compositions disclosed herein. The kit may comprise instructions for use in the treatment of Neurological disorders or its related complications.

[0018] The application also discloses a pharmaceutical composition comprising a pharmaceutically acceptable carrier and any of the compositions herein. In some aspects, the pharmaceutical composition is formulated for systemic administration, oral administration, sustained release, parenteral administration, injection, subdermal administration, or transdermal administration.

[0019] Herein, the application additionally provides kits comprising the pharmaceutical compositions described herein. The kits may further comprise instructions for use in the treatment of Neurological disorders or its related complications.

[0020] The compositions described herein have several uses. The present application provides, for example, methods of treating a patient suffering from Neurological disorders or its related complications manifested from metabolic conditions, chronic diseases or disorders; Hepatology, Cancer, Neurological, Hematological, Orthopedic, Cardiovascular, Renal, Skin, Vascular or Ocular complications.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0021] As used herein, the following terms and phrases shall have the meanings set forth below. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art.

[0022] The compounds of the present invention can be present in the form of pharmaceutically acceptable salts. The compounds of the present invention can also be present in the form of pharmaceutically acceptable esters (i.e., the methyl and ethyl esters of the acids of formula I and formula IA to be used as prodrugs). The compounds of the present invention can also be solvated, i.e. hydrated. The solvation can be affected in the course of the manufacturing process or can take place i.e. as a consequence of hygroscopic properties of an initially anhydrous compound of formula I and formula IA(hydration).

[0023] Compounds that have the same molecular formula but differ in the nature or sequence of bonding of their atoms or the arrangement of their atoms in space are termed "isomers." Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers." Diastereomers are stereoisomers with opposite configuration at one or more chiral centers which are not enantiomers. Stereoisomers bearing one or more asymmetric centers that are non- superimposable mirror images of each other are termed "enantiomers." When a compound has an asymmetric center, for example, if a carbon atom is bonded to four different groups, a pair of enantiomers is possible. An enantiomer can be characterized by the absolute configuration of its asymmetric center or centers and is described by the R- and S-sequencing rules of Cahn, Ingold and Prelog, or by the manner in which the molecule rotates the plane of polarized light and designated as dextrorotatory or levorotatory (i.e., as (+) or (-)-isomers respectively). A chiral compound can exist as either individual enantiomer or as a mixture thereof. A mixture containing equal proportions of the enantiomers is called a "racemic mixture".

[0024] As used herein, the term “metabolic condition” refers to an Inborn errors of metabolism (or genetic metabolic conditions) are genetic disorders that result from a defect in one or more metabolic pathways; specifically, the function of an enzyme is affected and is either deficient or completely absent.

[0025] In some embodiments, a molecular conjugate comprises of compounds selected from the group consisting of R-lipoic acid (CAS No. 1200-22-2), salsalate (CAS No. 552-94-3), acetylcysteine (CAS No. 616-91-1), Eicosapentaenoic acid (CAS No. 10417-94-4), Docosahexaenoic acid (CAS No. 6217-54-5).

[0026] The term "polymorph" as used herein is art-recognized and refers to one crystal structure of a given compound.

[0027] The phrases “parenteral administration” and “administered parenterally” as used herein refer to modes of administration other than enteral and topical administration, such as injections, and include without limitation intravenous, intramuscular, intrapleural, intravascular, intrapericardial, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradental, intraperitoneal, transtracheal, subcutaneous, subcuticular, intra-articular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0028] A “patient,” “subject,” or “host” to be treated by the subject method may mean either a human or non-human animal, such as primates, mammals, and vertebrates.

[0029] The phrase “pharmaceutically acceptable” is art-recognized. In certain embodiments, the term includes compositions, polymers and other materials and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of mammals, human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0030] The phrase “pharmaceutically acceptable carrier” is art-recognized, and includes, for example, pharmaceutically acceptable materials, compositions or vehicles, such as a liquid or solid filler, diluent, solvent or encapsulating material involved in carrying or transporting any subject composition, from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of a subject composition and not injurious to the patient. In certain embodiments, a pharmaceutically acceptable carrier is non-pyrogenic. Some examples of materials which may serve as pharmaceutically acceptable carriers include: (1) sugars, such as lactose, glucose and sucrose; (2) starches, such as corn starch and potato starch; (3) cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; (4) powdered tragacanth; (5) malt; (6) gelatin; (7) talc; (8) cocoa butter and suppository waxes; (9) oils, such as peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil and soybean oil; (10) glycols, such as propylene glycol; (11) polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; (12) esters, such as ethyl oleate and ethyl laurate; (13) agar; (14) buffering agents, such as magnesium hydroxide and aluminum hydroxide; (15) alginic acid; (16) pyrogen-free water; (17) isotonic saline; (18) Ringer’s solution; (19) ethyl alcohol; (20) phosphate buffer solutions; and (21) other non-toxic compatible substances employed in pharmaceutical formulations.

[0031] The term “prodrug” is intended to encompass compounds that, under physiological conditions, are converted into the therapeutically active agents of the present invention. A common method for making a prodrug is to include selected moieties that are hydrolyzed under physiological conditions to reveal the desired molecule. In other embodiments, the prodrug is converted by an enzymatic activity of the host animal.

[0032] The term “prophylactic or therapeutic” treatment is art-recognized and includes administration to the host of one or more of the subject compositions. If it is administered prior to clinical manifestation of the unwanted condition (e.g., disease or

other unwanted state of the host animal) then the treatment is prophylactic, i.e., it protects the host against developing the unwanted condition, whereas if it is administered after manifestation of the unwanted condition, the treatment is therapeutic, (i.e., it is intended to diminish, ameliorate, or stabilize the existing unwanted condition or side effects thereof).

[0033] The term "predicting" as used herein refers to assessing the probability related diseases patient will suffer from abnormalities or complication and/or terminal platelet aggregation or failure and/or death (i.e. mortality) within a defined time window (predictive window) in the future. The mortality may be caused by the central nervous system or complication. The predictive window is an interval in which the subject will develop one or more of the said complications according to the predicted probability. The predictive window may be the entire remaining lifespan of the subject upon analysis by the method of the present invention.

[0034] The term "treating" is art -recognized and includes preventing a disease, disorder or condition from occurring in an animal which may be predisposed to the disease, disorder and/or condition but has not yet been diagnosed as having it; inhibiting the disease, disorder or condition, e.g., impeding its progress; and relieving the disease, disorder, or condition, e.g., causing regression of the disease, disorder and/or condition. Treating the disease or condition includes ameliorating at least one symptom of the particular disease or condition, even if the underlying pathophysiology is not affected, such as treating the neurological condition such as epilepsy, bipolar disorder, trigeminal neuralgia, attention-deficit hyperactivity disorder (ADHD), schizophrenia, neuropathic pain, seizures, bipolar disorder, mania, phantom limb syndrome, complex regional pain syndrome, paroxysmal extreme pain disorder, neuromyotonia, intermittent explosive disorder, borderline personality disorder, Myotonia congenita and post-traumatic stress disorder of a subject by administration of an agent even though such agent does not treat the cause of the condition. The term "treating", "treat" or "treatment" as used herein includes curative, preventative (e.g., prophylactic), adjunct and palliative treatment.

[0035] The phrase "therapeutically effective amount" is an art-recognized term. In certain embodiments, the term refers to an amount of a salt or composition disclosed herein that produces some desired effect at a reasonable benefit/risk ratio applicable to any medical treatment. In certain embodiments, the term refers to that amount necessary or sufficient to eliminate or reduce medical symptoms for a period of time. The effective amount may vary depending on such factors as the disease or condition being treated, the particular targeted constructs being administered, the size of the subject, or the severity of the disease or condition. One of ordinary skill in the art may empirically determine the effective amount of a particular composition without necessitating undue experimentation.

[0036] In certain embodiments, the pharmaceutical compositions described herein are formulated in a manner such that said compositions will be delivered to a patient in a therapeutically effective amount, as part of a prophylactic or therapeutic treatment. The desired amount of the composition to be administered to a patient will depend on absorption, inactivation, and excretion rates of the drug as well as the delivery rate of the salts and compositions from the subject compositions. It is to be noted that dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions. Typically, dosing will be determined using techniques known to one skilled in the art.

[0037] Additionally, the optimal concentration and/or quantities or amounts of any particular salt or composition may be adjusted to accommodate variations in the treatment parameters. Such treatment parameters include the clinical use to which the preparation is put, e.g., the site treated, the type of patient, e.g., human or non-human, adult or child, and the nature of the disease or condition.

[0038] In certain embodiments, the dosage of the subject compositions provided herein may be determined by reference to the plasma concentrations of the therapeutic composition or other encapsulated materials. For example, the maximum plasma concentration (C_{max}) and the area under the plasma concentration-time curve from time 0 to infinity may be used.

[0039] When used with respect to a pharmaceutical composition or other material, the term "sustained release" is art-recognized. For example, a subject composition which releases a substance over time may exhibit sustained release characteristics, in contrast to a bolus type administration in which the entire amount of the substance is made biologically available at one time. For example, in particular embodiments, upon contact with body fluids including blood, spinal fluid, mucus secretions, lymph or the like, one or more of the pharmaceutically acceptable excipients may undergo gradual or delayed degradation (e.g., through hydrolysis) with concomitant release of any material incorporated therein, e.g., an therapeutic and/or biologically active salt and/or composition, for a sustained or extended period (as compared to the release from a bolus). This release may result in prolonged delivery of therapeutically effective amounts of any of the therapeutic agents disclosed herein.

[0040] The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" are art-recognized, and include the administration of a subject composition, therapeutic or other material at a site remote from the disease being treated. Administration of an agent for the disease being treated, even if the agent is subsequently distributed systemically, may be termed "local" or "topical" or "regional" administration, other than directly into the central nervous system, e.g., by subcutaneous administration, such that it enters the patient's system and, thus, is subject to metabolism and other like processes.

[0041] The phrase "therapeutically effective amount" is an art-recognized term. In certain embodiments, the term refers to an amount of a salt or composition disclosed

herein that produces some desired effect at a reasonable benefit/risk ratio applicable to any medical treatment. In certain embodiments, the term refers to that amount necessary or sufficient to eliminate or reduce medical symptoms for a period of time. The effective amount may vary depending on such factors as the disease or condition being treated, the particular targeted constructs being administered, the size of the subject, or the severity of the disease or condition. One of ordinary skill in the art may empirically determine the effective amount of a particular composition without necessitating undue experimentation.

[0042] The present disclosure also contemplates prodrugs of the compositions disclosed herein, as well as pharmaceutically acceptable salts of said prodrugs.

[0043] This application also discloses a pharmaceutical composition comprising a pharmaceutically acceptable carrier and the composition of a compound of Formula I and formula IA may be formulated for systemic or topical or oral administration. The pharmaceutical composition may be also formulated for oral administration, oral solution, injection, subdermal administration, or transdermal administration. The pharmaceutical composition may further comprise at least one of a pharmaceutically acceptable stabilizer, diluent, surfactant, filler, binder, and lubricant.

[0044] In many embodiments, the pharmaceutical compositions described herein will incorporate the disclosed compounds and compositions (Formula I and IA) to be delivered in an amount sufficient to deliver to a patient a therapeutically effective amount of a compound of formula I and formula IA or composition as part of a prophylactic or therapeutic treatment. The desired concentration of formula I and formula IA or its pharmaceutically acceptable salts will depend on absorption, inactivation, and excretion rates of the drug as well as the delivery rate of the salts and compositions from the subject compositions. It is to be noted that dosage values may also vary with the severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the

individual need and the professional judgment of the person administering or supervising the administration of the compositions. Typically, dosing will be determined using techniques known to one skilled in the art.

[0045] Additionally, the optimal concentration and/or quantities or amounts of any particular compound of formula I and formula IA may be adjusted to accommodate variations in the treatment parameters. Such treatment parameters include the clinical use to which the preparation is put, e.g., the site treated, the type of patient, e.g., human or non-human, adult or child, and the nature of the disease or condition.

[0046] The concentration and/or amount of any compound of formula I and formula IA may be readily identified by routine screening in animals, e.g., rats, by screening a range of concentration and/or amounts of the material in question using appropriate assays. Known methods are also available to assay local tissue concentrations, diffusion rates of the salts or compositions, and local blood flow before and after administration of therapeutic formulations disclosed herein. One such method is microdialysis, as reviewed by T. E. Robinson et al., 1991, *microdialysis in the neurosciences, Techniques*, volume 7, Chapter 1. The methods reviewed by Robinson may be applied, in brief, as follows. A microdialysis loop is placed in situ in a test animal. Dialysis fluid is pumped through the loop. When compounds with formula I and formula IA such as those disclosed herein are injected adjacent to the loop, released drugs are collected in the dialysate in proportion to their local tissue concentrations. The progress of diffusion of the salts or compositions may be determined thereby with suitable calibration procedures using known concentrations of salts or compositions.

[0047] In certain embodiments, the dosage of the subject compounds of formula I and formula IA provided herein may be determined by reference to the plasma concentrations of the therapeutic composition or other encapsulated materials. For example, the maximum plasma concentration (C_{max}) and the area under the plasma concentration-time curve from time 0 to infinity may be used.

[0048] Generally, in carrying out the methods detailed in this application, an effective dosage for the compounds of Formulas I is in the range of about 0.01 mg/kg/day to about 100 mg/kg/day in single or divided doses, for instance 0.01 mg/kg/day to about 50 mg/kg/day in single or divided doses. The compounds of Formulas I may be administered at a dose of, for example, less than 0.2 mg/kg/day, 0.5 mg/kg/day, 1.0 mg/kg/day, 5 mg/kg/day, 10 mg/kg/day, 20 mg/kg/day, 30 mg/kg/day, or 40 mg/kg/day. Compounds of Formula I and formula IA may also be administered to a human patient at a dose of, for example, between 0.1 mg and 1000 mg, between 5 mg and 80 mg, or less than 1.0, 9.0, 12.0, 20.0, 50.0, 75.0, 100, 300, 400, 500, 800, 1000, 2000, 5000 mg per day. In certain embodiments, the compositions herein are administered at an amount that is less than 95%, 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, or 10% of the compound of formula I and formula IA required for the same therapeutic benefit.

[0049] An effective amount of the compounds of formula I and formula IA described herein refers to the amount of one of said salts or compositions which is capable of inhibiting or preventing a disease.

[0050] An effective amount may be sufficient to prohibit, treat, alleviate, ameliorate, halt, restrain, slow or reverse the progression, or reduce the severity of a complication resulting from nerve damage or demyelization and/or elevated reactive oxidative-nitrosative species and/or abnormalities in neurotransmitter homeostasis's, in patients who are at risk for such complications. As such, these methods include both medical therapeutic (acute) and/or prophylactic (prevention) administration as appropriate. The amount and timing of compositions administered will, of course, be dependent on the subject being treated, on the severity of the affliction, on the manner of administration and on the judgment of the prescribing physician. Thus, because of patient-to-patient variability, the dosages given above are a guideline and the physician may titrate doses of the drug to achieve the treatment that the physician considers appropriate for the patient. In considering the degree of treatment desired, the physician must balance a variety of

factors such as age of the patient, presence of preexisting disease, as well as presence of other diseases.

[0051] The compositions provided by this application may be administered to a subject in need of treatment by a variety of conventional routes of administration, including orally, topically, parenterally, e.g., intravenously, subcutaneously or intramedullary. Further, the compositions may be administered intranasally, as a rectal suppository, or using a "flash" formulation, i.e., allowing the medication to dissolve in the mouth without the need to use water. Furthermore, the compositions may be administered to a subject in need of treatment by controlled release dosage forms, site specific drug delivery, transdermal drug delivery, patch (active/passive) mediated drug delivery, by stereotactic injection, or in nanoparticles.

[0052] The compositions may be administered alone or in combination with pharmaceutically acceptable carriers, vehicles or diluents, in either single or multiple doses. Suitable pharmaceutical carriers, vehicles and diluents include inert solid diluents or fillers, sterile aqueous solutions and various organic solvents. The pharmaceutical compositions formed by combining the compositions and the pharmaceutically acceptable carriers, vehicles or diluents are then readily administered in a variety of dosage forms such as tablets, powders, lozenges, syrups, injectable solutions and the like. These pharmaceutical compositions can, if desired, contain additional ingredients such as flavorings, binders, excipients and the like. Thus, for purposes of oral administration, tablets containing various excipients such as L-arginine, sodium citrate, calcium carbonate and calcium phosphate may be employed along with various disintegrates such as starch, alginic acid and certain complex silicates, together with binding agents such as polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules. Appropriate materials for this include lactose or milk sugar and high molecular weight polyethylene glycols. When aqueous suspensions or elixirs are

desired for oral administration, the essential active ingredient therein may be combined with various sweetening or flavoring agents, coloring matter or dyes and, if desired, emulsifying or suspending agents, together with diluents such as water, ethanol, propylene glycol, glycerin and combinations thereof. The compounds of formula I and formula IA may also comprise enterically coated comprising of various excipients, as is well known in the pharmaceutical art.

[0053] For parenteral administration, solutions of the compositions may be prepared in (for example) sesame or peanut oil, aqueous propylene glycol, or in sterile aqueous solutions may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, the sterile aqueous media employed are all readily available by standard techniques known to those skilled in the art.

[0054] The formulations, for instance tablets, may contain e.g. 10 to 100, 50 to 250, 150 to 500 mg, or 350 to 800 mg e.g. 10, 50, 100, 300, 500, 700, 800 mg of the compounds of formula I and formula IA disclosed herein, for instance, compounds of formula I and formula IA or pharmaceutical acceptable salts of a compounds of Formula I.

[0055] Generally, a composition as described herein may be administered orally, or parenterally (e.g., intravenous, intramuscular, subcutaneous or intramedullary). Topical administration may also be indicated, for example, where the patient is suffering from gastrointestinal disorder that prevent oral administration, or whenever the medication is best applied to the surface of a tissue or organ as determined by the attending physician. Localized administration may also be indicated, for example, when a high dose is desired at the target tissue or organ. For buccal administration the active composition may take the form of tablets or lozenges formulated in a conventional manner.

[0056] The dosage administered will be dependent upon the identity of the neurological disease; the type of host involved, including its age, health and weight; the kind of concurrent treatment, if any; the frequency of treatment and therapeutic ratio.

[0057] Illustratively, dosage levels of the administered active ingredients are: intravenous, 0.1 to about 200 mg/kg; intramuscular, 1 to about 500 mg/kg; orally, 5 to about 1000 mg/kg; intranasal instillation, 5 to about 1000 mg/kg; and aerosol, 5 to about 1000 mg/kg of host body weight.

[0058] Expressed in terms of concentration, an active ingredient can be present in the compositions of the present invention for localized use about the cutis, intranasally, pharyngolaryngeally, bronchially, intravaginally, rectally, or ocularly in a concentration of from about 0.01 to about 50% w/w of the composition; preferably about 1 to about 20% w/w of the composition; and for parenteral use in a concentration of from about 0.05 to about 50% w/v of the composition and preferably from about 5 to about 20% w/v.

[0059] The compositions of the present invention are preferably presented for administration to humans and animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, suppositories, sterile parenteral solutions or suspensions, sterile non-parenteral solutions of suspensions, and oral solutions or suspensions and the like, containing suitable quantities of an active ingredient. For oral administration either solid or fluid unit dosage forms can be prepared.

[0060] As discussed above, the tablet core contains one or more hydrophilic polymers. Suitable hydrophilic polymers include, but are not limited to, water swellable cellulose derivatives, polyalkylene glycols, thermoplastic polyalkylene oxides, acrylic polymers, hydrocolloids, clays, gelling starches, swelling cross-linked polymers, and mixtures thereof. Examples of suitable water swellable cellulose derivatives include, but are not limited to, sodium carboxymethylcellulose, cross-linked hydroxypropylcellulose, hydroxypropyl cellulose (HPC), hydroxypropylmethylcellulose (HPMC), hydroxyisopropylcellulose, hydroxybutylcellulose, hydroxyphenylcellulose,

hydroxyethylcellulose (HEC), hydroxypentylcellulose, hydroxypropylethylcellulose, hydroxypropylbutylcellulose, and hydroxypropylethylcellulose, and mixtures thereof. Examples of suitable polyalkylene glycols include, but are not limited to, polyethylene glycol. Examples of suitable thermoplastic polyalkylene oxides include, but are not limited to, poly(ethylene oxide). Examples of suitable acrylic polymers include, but are not limited to, potassium methacrylatedivinylbenzene copolymer, polymethylmethacrylate, high-molecular weight crosslinked acrylic acid homopolymers and copolymers such as those commercially available from Noveon Chemicals under the tradename CARBOPOL™. Examples of suitable hydrocolloids include, but are not limited to, alginates, agar, guar gum, locust bean gum, kappa carrageenan, iota carrageenan, tara, gum arabic, tragacanth, pectin, xanthan gum, gellan gum, maltodextrin, galactomannan, pustulan, laminarin, scleroglucan, gum arabic, inulin, pectin, gelatin, whelan, rhamnan, zooglan, methylan, chitin, cyclodextrin, chitosan, and mixtures thereof. Examples of suitable clays include, but are not limited to, smectites such as bentonite, kaolin, and laponite; magnesium trisilicate; magnesium aluminum silicate; and mixtures thereof. Examples of suitable gelling starches include, but are not limited to, acid hydrolyzed starches, swelling starches such as sodium starch glycolate and derivatives thereof, and mixtures thereof. Examples of suitable swelling cross-linked polymers include, but are not limited to, cross-linked polyvinyl pyrrolidone, cross-linked agar, and cross-linked carboxymethylcellulose sodium, and mixtures thereof.

[0061] The carrier may contain one or more suitable excipients for the formulation of tablets. Examples of suitable excipients include, but are not limited to, fillers, adsorbents, binders, disintegrants, lubricants, glidants, release-modifying excipients, superdisintegrants, antioxidants, and mixtures thereof.

[0062] Suitable binders include, but are not limited to, dry binders such as polyvinyl pyrrolidone and hydroxypropylmethylcellulose; wet binders such as water-soluble polymers, including hydrocolloids such as acacia, alginates, agar, guar gum, locust bean, carrageenan, carboxymethylcellulose, tara, gum arabic, tragacanth, pectin, xanthan,

gellan, gelatin, maltodextrin, galactomannan, pustulan, laminarin, scleroglucan, inulin, wheilan, rhamnan, zooglan, methylan, chitin, cyclodextrin, chitosan, polyvinyl pyrrolidone, celluloses, sucrose, and starches; and mixtures thereof. Suitable disintegrants include, but are not limited to, sodium starch glycolate, cross-linked polyvinylpyrrolidone, cross-linked carboxymethylcellulose, starches, microcrystalline cellulose, and mixtures thereof.

[0063] Suitable lubricants include, but are not limited to, long chain fatty acids and their salts, such as magnesium stearate and stearic acid, talc, glycerides waxes, and mixtures thereof. Suitable glidants include, but are not limited to, colloidal silicon dioxide. Suitable release-modifying excipients include, but are not limited to, insoluble edible materials, pH-dependent polymers, and mixtures thereof.

[0064] Suitable insoluble edible materials for use as release-modifying excipients include, but are not limited to, water-insoluble polymers and low-melting hydrophobic materials, copolymers thereof, and mixtures thereof. Examples of suitable water-insoluble polymers include, but are not limited to, ethylcellulose, polyvinyl alcohols, polyvinyl acetate, polycaprolactones, cellulose acetate and its derivatives, acrylates, methacrylates, acrylic acid copolymers, copolymers thereof, and mixtures thereof. Suitable low-melting hydrophobic materials include, but are not limited to, fats, fatty acid esters, phospholipids, waxes, and mixtures thereof. Examples of suitable fats include, but are not limited to, hydrogenated vegetable oils such as for example cocoa butter, hydrogenated palm kernel oil, hydrogenated cottonseed oil, hydrogenated sunflower oil, and hydrogenated soybean oil, free fatty acids and their salts, and mixtures thereof. Examples of suitable fatty acid esters include, but are not limited to, sucrose fatty acid esters, mono-, di-, and triglycerides, glyceryl behenate, glyceryl palmitostearate, glyceryl monostearate, glyceryl tristearate, glyceryl triaurylate, glyceryl myristate, GlycoWax-932, lauroyl macrogol-32 glycerides, stearyl macrogol-32 glycerides, and mixtures thereof. Examples of suitable phospholipids include phosphotidyl choline, phosphotidyl serene, phosphotidyl enositol, phosphotidic acid, and mixtures thereof. Examples of

suitable waxes include, but are not limited to, carnauba wax, spermaceti wax, beeswax, candelilla wax, shellac wax, microcrystalline wax, and paraffin wax; fat-containing mixtures such as chocolate, and mixtures thereof. Examples of super disintegrants include, but are not limited to, croscarmellose sodium, sodium starch glycolate and cross-linked povidone (crospovidone). In one embodiment the tablet core contains up to about 5 percent by weight of such super disintegrant.

[0065] Examples of antioxidants include, but are not limited to, tocopherols, ascorbic acid, sodium pyrosulfite, butylhydroxytoluene, butylated hydroxyanisole, edetic acid, and edetate salts, and mixtures thereof. Examples of preservatives include, but are not limited to, citric acid, tartaric acid, lactic acid, malic acid, acetic acid, benzoic acid, and sorbic acid, and mixtures thereof.

[0066] In one embodiment, the immediate release coating has an average thickness of at least 50 microns, such as from about 50 microns to about 2500 microns; e.g., from about 250 microns to about 1000 microns. In embodiment, the immediate release coating is typically compressed at a density of more than about 0.9 g/cc, as measured by the weight and volume of that specific layer.

[0067] In one embodiment, the immediate release coating contains a first portion and a second portion, wherein at least one of the portions contains the second pharmaceutically active agent. In one embodiment, the portions contact each other at a center axis of the tablet. In one embodiment, the first portion includes the first pharmaceutically active agent and the second portion includes the second pharmaceutically active agent.

[0068] In one embodiment, the first portion contains the first pharmaceutically active agent and the second portion contains the second pharmaceutically active agent. In one embodiment, one of the portions contains a third pharmaceutically active agent. In one embodiment one of the portions contains a second immediate release portion of the same pharmaceutically active agent as that contained in the tablet core.

[0069] In one embodiment, the outer coating portion is prepared as a dry blend of materials prior to addition to the coated tablet core. In another embodiment the outer coating portion is included of a dried granulation including the pharmaceutically active agent.

[0070] Formulations with different drug release mechanisms described above could be combined in a final dosage form containing single or multiple units. Examples of multiple units include multilayer tablets, capsules containing tablets, beads, or granules in a solid or liquid form. Typical, immediate release formulations include compressed tablets, gels, films, coatings, liquids and particles that can be encapsulated, for example, in a gelatin capsule. Many methods for preparing coatings, covering or incorporating drugs, are known in the art.

[0071] The immediate release dosage, unit of the dosage form, i.e., a tablet, a plurality of drug-containing beads, granules or particles, or an outer layer of a coated core dosage form, contains a therapeutically effective quantity of the active agent with conventional pharmaceutical excipients. The immediate release dosage unit may or may not be coated, and may or may not be admixed with the delayed release dosage unit or units (as in an encapsulated mixture of immediate release drug-containing granules, particles or beads and delayed release drug-containing granules or beads).

[0072] Extended release formulations are generally prepared as diffusion or osmotic systems, for example, as described in "Remington—The Science and Practice of Pharmacy", 20th. Ed., Lippincott Williams & Wilkins, Baltimore, Md., 2000). A diffusion system typically consists of one of two types of devices, reservoir and matrix, which are wellknown and described in die art. The matrix devices are generally prepared by compressing the drug with a slowly dissolving polymer carrier into a tablet form.

[0073] An immediate release portion can be added to the extended release system by means of either applying an immediate release layer on top of the extended release core; using coating or compression processes or in a multiple unit system such as a capsule containing extended and immediate release beads.

[0074] Delayed release dosage formulations are created by coating a solid dosage form with a film of a polymer which is insoluble in the acid environment of the stomach, but soluble in the neutral environment of small intestines. The delayed release dosage units can be prepared, for example, by coating a drug or a drug-containing composition with a selected coating material. The drug-containing composition may be a tablet for incorporation into a capsule, a tablet for use as an inner core in a "coated core" dosage form, or a plurality of drug-containing beads, particles or granules, for incorporation into either a tablet or capsule.

[0075] A pulsed release dosage form is one that mimics a multiple dosing profile without repeated dosing and typically allows at least a twofold reduction in dosing frequency as compared to the drug presented as a conventional dosage form (e.g., as a solution or prompt drug-releasing, conventional solid dosage form). A pulsed release profile is characterized by a time period of no release (lag time) or reduced release followed by rapid drug release.

[0076] Each dosage form contains a therapeutically effective amount of active agent. In one embodiment of dosage forms that mimic a twice daily dosing profile, approximately 30 wt. % to 70 wt. %, preferably 40 wt. % to 60 wt. %, of the total amount of active agent in the dosage form is released in the initial pulse, and, correspondingly approximately 70 wt. % to 3.0 wt. %, preferably 60 wt. % to 40 wt. %, of the total amount of active agent in the dosage form is released in the second pulse. For dosage forms mimicking the twice daily dosing profile, the second pulse is preferably released approximately 3 hours to less than 14 hours, and more preferably approximately 5 hours to 12 hours, following administration.

[0077] Another dosage form contains a compressed tablet or a capsule having a drug-containing immediate release dosage unit, a delayed release dosage unit and an optional second delayed release dosage unit. In this dosage form, the immediate release dosage unit contains a plurality of beads, granules particles that release drug substantially immediately following oral administration to provide an initial dose. The delayed release dosage unit contains a plurality of coated beads or granules, which release drug approximately 3 hours to 14 hours following oral administration to provide a second dose.

[0078] For purposes of transdermal (e.g., topical) administration, dilute sterile, aqueous or partially aqueous solutions (usually in about 0.1% to 5% concentration), otherwise similar to the above parenteral solutions, may be prepared.

[0079] Methods of preparing various pharmaceutical compositions with a certain amount of one or more compounds of formula I and formula IA or other active agents are known, or will be apparent in light of this disclosure, to those skilled in this art. For examples of methods of preparing pharmaceutical compositions, see Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 19th Edition (1995).

[0080] In addition, in certain embodiments, subject compositions of the present application maybe lyophilized or subjected to another appropriate drying technique such as spray drying. The subject compositions may be administered once, or may be divided into a number of smaller doses to be administered at varying intervals of time, depending in part on the release rate of the compositions and the desired dosage.

[0081] Formulations useful in the methods provided herein include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal, aerosol and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of a subject composition which may be combined with a carrier material to

produce a single dose may vary depending upon the subject being treated, and the particular mode of administration.

[0082] Methods of preparing these formulations or compositions include the step of bringing into association subject compositions with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a subject composition with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

[0083] The compounds of formula I and formula IA described herein may be administered in inhalant or aerosol formulations. The inhalant or aerosol formulations may comprise one or more agents, such as adjuvants, diagnostic agents, imaging agents, or therapeutic agents useful in inhalation therapy. The final aerosol formulation may for example contain 0.005-90% w/w, for instance 0.005-50%, 0.005-5% w/w, or 0.01-1.0% w/w, of medicament relative to the total weight of the formulation.

[0084] In solid dosage forms for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the subject composition is mixed with one or more pharmaceutically acceptable carriers and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, acetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be

employed as fillers in soft and hard-filled gelatin capsules using lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

[0085] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the subject compositions, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, corn, peanut, sunflower, soybean, olive, castor, and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

[0086] Suspensions, in addition to the subject compositions, may contain suspending agents such as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol, and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

[0087] Formulations for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing a subject composition with one or more suitable non-irritating carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax, or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the appropriate body cavity and release the encapsulated compound(s) and composition(s). Formulations which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams, or spray formulations containing such carriers as are known in the art to be appropriate.

[0088] Dosage forms for transdermal administration include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches, and inhalants. A subject

composition may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants that may be required. For transdermal administration, the complexes may include lipophilic and hydrophilic groups to achieve the desired water solubility and transport properties.

[0089] The ointments, pastes, creams and gels may contain, in addition to subject compositions, other carriers, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof. Powders and sprays may contain, in addition to a subject composition, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of such substances. Sprays may additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

[0090] Methods of delivering a composition or compositions via a transdermal patch are known in the art. Exemplary patches and methods of patch delivery are described in US Patent Nos. 6,974,588, 6,564,093, 6,312,716, 6,440,454, 6,267,983, 6,239,180, and 6,103,275.

[0091] In another embodiment, a transdermal patch may comprise: a substrate sheet comprising a composite film formed of a resin composition comprising 100 parts by weight of a polyvinyl chloride-polyurethane composite and 2-10 parts by weight of a styrene-ethylene-butylene-styrene copolymer, a first adhesive layer on the one side of the composite film, and a polyalkylene terephthalate film adhered to the one side of the composite film by means of the first adhesive layer, a primer layer which comprises a saturated polyester resin and is formed on the surface of the polyalkylene terephthalate film; and a second adhesive layer comprising a styrene-diene-styrene block copolymer containing a pharmaceutical agent layered on the primer layer. A method for the manufacture of the above-mentioned substrate sheet comprises preparing the above resin

composition molding the resin composition into a composite film by a calendar process, and then adhering a polyalkylene terephthalate film on one side of the composite film by means of an adhesive layer thereby forming the substrate sheet, and forming a primer layer comprising a saturated polyester resin on the outer surface of the polyalkylene terephthalate film.

[0092] Another type of patch comprises incorporating the drug directly in a pharmaceutically acceptable adhesive and laminating the drug-containing adhesive onto a suitable backing member, e.g. a polyester backing membrane. The drug should be present at a concentration which will not affect the adhesive properties, and at the same time deliver the required clinical dose.

[0093] Transdermal patches may be passive or active. Passive transdermal drug delivery systems currently available, such as the nicotine, estrogen and nitroglycerine patches, deliver small-molecule drugs. Many of the newly developed proteins and peptide drugs are too large to be delivered through passive transdermal patches and may be delivered using technology such as electrical assist (iontophoresis) for large-molecule drugs.


[0094] Iontophoresis is a technique employed for enhancing the flux of ionized substances through membranes by application of electric current. One example of an iontophoretic membrane is given in U.S. Pat. No. 5,080,646 to Theeuwes. The principal mechanisms by which iontophoresis enhances molecular transport across the skin are (a) repelling a charged ion from an electrode of the same charge, (b) electroosmosis, the convective movement of solvent that occurs through a charged pore in response the preferential passage of counter-ions when an electric field is applied or (c) increase skin permeability due to application of electrical current.

[0095] In some cases, it may be desirable to administer in the form of a kit, it may comprise a container for containing the separate compositions such as a divided bottle or

a divided foil packet. Typically the kit comprises directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral), are administered at different dosage intervals, or when titration of the individual components of the combination is desired by the prescribing physician.

[0096] An example of such a kit is a so-called blister pack. Blister packs are well known in the packaging industry and are widely used for the packaging of pharmaceutical unit dosage forms (tablets, capsules, and the like). Blister packs generally consist of a sheet of relatively stiff material covered with a foil of a plastic material that may be transparent.

[0097] Methods and compositions for the treatment of Neurological disorders. Among other things, herein is provided a method of treating Neurological disorders, comprising administering to a patient in need thereof a therapeutically effective amount of compound of Formula I:




Formula I


Wherein,













R¹ each independently represents Oxygen (O), D, Hydrogen (H) or Flourine (F),


R³ each independently represents



R² independently represents Acetyl (CH₃CO-),







a is independently 2,3 or 7;

each b is independently 3, 5 or 6;

e is independently 1, 2 or 6;

c and d are each independently H, D, -OH, -OD, C₁-C₆-alkyl, -NH₂ or -COCH₃.


Methods for using compounds of formula I and formula IA:

[0098] The invention also includes methods for treating neurological disorders such as convulsions and mood-stabilizing agent and, also used primarily in the treatment of epilepsy and bipolar disorder, as well as trigeminal neuralgia. It is also used for a variety of indications, including attention-deficit hyperactivity disorder (ADHD), schizophrenia, neuropathic pain, seizures, bipolar disorder, mania, phantom limb syndrome, complex regional pain syndrome, paroxysmal extreme pain disorder, neuromyotonia, intermittent explosive disorder, borderline personality disorder, Myotonia congenita and post-traumatic stress disorder.


METHODS OF MAKING

[0099] Examples of synthetic pathways useful for making compounds of formula IA are set forth in example below and generalized in scheme 1.

Scheme – 1:




[00100] Step-1: Synthesis of compound 2:




[00101] To a stirred suspension of compound 1 (200 g, 847.5 mmol) and sodium carbonate (287.4 g, 2711 mmol) in dichloromethane (1000 ml) were added tablets of potassium permanganate supported on alumina (3.5% w/w, 3.46 g, 0.77 mmol). Thereafter, peroxyacetic acid (39% solution in acetic acid, 432 ml, 2538 mmol) was added dropwise over one hour, causing a gradual rise in temperature until gentle reflux of the solvent. The mixture was stirred for twenty minutes and then allowed to stand for twenty minutes. The sodium carbonate and supported catalyst were then removed by filtration and washed by dichloromethane (200 ml); the alumina beads were separated from sodium carbonate by screening through a sieve. The combined filtrate was then stirred with an aqueous solution of sodium sulphite (20 g) and sodium bicarbonate (20 g) in water (250 ml) for one hour. The phases were then separated and the aqueous phase extracted by dichloromethane (50 ml). The combined organic layers were washed by water (100 ml), saturated aqueous sodium bicarbonate (100 ml), water again (100 ml) and brine, then dried over anhydrous sodium sulphate and filtered. Evaporation of the solvent (rotary evaporator, water aspirator pressure, 40 degree. C.) gave the crude epoxide 2 as a beige solid which was crystallised from ethyl acetate (100 ml) to give the product as an off-white solid, 194.2 g, (91% yield).

[00102] Step-2: Synthesis of compound 3:




[00103] To a solution of the epoxide 2 (5.03 g, 20 mmol) in methanol (100 ml), dichloromethane (50 ml) and water (5 ml) at room temperature under nitrogen was added ammonium formate (3.78 g, 60 mmol) followed by 10% palladium on charcoal (540 mg, 0.51 mmol Pd). The resulting mixture was stirred at room temperature for one hour and then the catalyst was recovered by filtration through celite. The filter pad was washed with dichloromethane (20 ml), and the organic phase of the combined filtrate was separated and dried over anhydrous sodium sulphate. Filtration and evaporation of the solvent (rotary evaporator, water aspirator pressure, 40.degree. C.) gave the crude alcohol (3) which was crystallized from ethyl acetate (20 ml) to afford white crystals, 4.7 g, (93% yield).

[00104] Step-4: Synthesis of compound 4:




[00105] Dichloromethane 2160 ml, 164.96 g (R)-(-)-acetyl mandelic acid, 180 g racemic 3, 86.48 g 4-dimethylamino pyrimidine and 175.28g dicyclohexylcarbodiimide were mixed in a round bottom flask and the resultant reaction mixture was refluxed for one hour. After completion of reaction the reaction mixture was filtered and the filtrate was washed with aqueous acetic acid and aqueous sodium bicarbonate solution. The organic layer was collected and it was concentrated. To the residue thus obtained 1620ml of methyl isobutylketone was added. The resultant reaction mixture was heated at 60 - 65 °C for one hour and filtered at the same temperature. The solid thus obtained was again purified by using methyl isobutylketone at 60 - 65 °C to get title compound with 97.6% of optical purity.

[00106] Step-5: Synthesis of compound 5:



[00107] 250 ml water, 50 g compound 4 and 50 g sodium hydroxide were mixed in a round bottom flask. The resultant reaction mixture was heated at 80 -85 °C for 1 hour. After completion of reaction the reaction mixture was cooled to ambient temperature and the pH of the reaction mixture was adjusted to acidic to get 26 g of compound 5.

[00108] Step-6: Synthesis of compound 6:



[00109] 240 ml dichloromethane, 5.29 g, acetic acid, 20 g compound 5, 9.6 g 4-dimethylamino pyridine and 19.47 g dicyclohexylcarbodiimide were charged in a round bottom flask. The reaction mixture was refluxed for 1 hour, after completion of reaction the reaction mixture was filtered, and the filtrate was washed with aqueous acetic acid and aqueous sodium bicarbonate solution. The organic layer was separated and it was concentrated, to the residue thus obtained isopropyl alcohol (140 ml) was added. The resultant mixture was heated to reflux for 30 minutes followed by cooling to 25 - 30 °C. The product thus separated and filtered to obtain 17g of compound 6 with 99.6% optical purity.

combined organic layers were washed with water (2 x 5mL) followed by brine solution (10 mL), dried over anhydrous Na_2SO_4 and evaporated under reduced pressure. The crude was purified by column chromatography over 100-200 mesh silica gel to get the product **10**.

EXAMPLES

[00114] Toxicokinetic Effects of Carbamazepine and Formula (1-2) in Diabetic Rats

Carbamazepine (CBZ) is one of the most widely used antiepileptic drugs. It is classified as a narrow therapeutic index drug and is belonging to a group of drugs that are highly susceptible to drug-drug interactions.

Methods:

Diabetes was induced in adult male Sprague Dawely rats by streptozotocin (60 mg/kg, ip). Animals were divided into three sets, normal and diabetic sets; each set was divided into control and three-treated groups where rats received orally carbamazepine and formula (1-2) in dose levels of 100 and 200 mg/kg once daily for 28 days. The pharmacokinetic profiles were carried out over a period of 24 hours. The liver functions were evaluated by the determination of albumin, ALT and AST plasma levels and kidney functions were determined by creatinine and urea. Histopathology of rat's livers and kidneys were also investigated. Carbamazepine plasma levels were determined by a validated HPLC method.

Results:

The obtained results showed that Liver enzymes, urea and creatinine levels were significantly elevated in carbamazepine-treated groups ($P < 0.05$). These effects were more pronounced in diabetic rats. Diabetes significantly altered the rate and extent of carbamazepine absorption ($P < 0.05$).

Carbamazepine pharmacokinetic parameters showed that diabetes reduced C_{max} , AUC_{0-24} and $AUC_{0-\infty}$ by 6.8%, 13.6 % and 22.8%, respectively following administration of

carbamazepine 100 mg/kg and by 20.3%, 12.2% and 32.8%, respectively following administration of carbamazepine 200 mg/kg.

Formula (1-2) pharmacokinetic parameters showed better C_{max}, AUC₀₋₂₄ and AUC_{0-inf} by 8.1%, 26.4 % and 31.3%, respectively following administration of formula (1-2) 100 mg/kg and by 32.9%, 19.8% and 39.8%, respectively following administration of formula (1-2) 200 mg/kg.

[00115] The term "sample" refers to a sample of a body fluid, to a sample of separated cells or to a sample from a tissue or an organ. Samples of body fluids can be obtained by well known techniques and include, preferably, samples of blood, plasma, serum, or urine, more preferably, samples of blood, plasma or serum.

EQUIVALENTS


[00116] The present disclosure provides among other things compositions and methods for treating Neurological disorders and their complications. While specific embodiments of the subject disclosure have been discussed, the above specification is illustrative and not restrictive. Many variations of the systems and methods herein will become apparent to those skilled in the art upon review of this specification. The full scope of the claimed systems and methods should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

INCORPORATION BY REFERENCE

[00117] All publications and patents mentioned herein, including those items listed above, are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually indicated to be incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.

CLAIMS:

1. A compound of formula I:




Formula I


or a pharmaceutically acceptable salt, hydrate, polymorph, solvate, prodrug, enantiomer, or stereoisomer thereof;

Wherein,


R¹ each independently represents Oxygen (O), D, Hydrogen (H) or Flourine (F);









R³ each independently represents H, D,






R² independently represents Acetyl (CH₃CO-),








a is independently 2,3 or 7;

each b is independently 3, 5 or 6;

e is independently 1, 2 or 6;

c and d are each independently H, D, -OH, -OD, C₁-C₆-alkyl, -NH₂ or -COCH₃.

2. Compounds and compositions of formula IA in a R-enantiomer or stereoisomeric form




Formula IA













or a pharmaceutically acceptable salt, hydrate, polymorph, solvate, prodrug, enantiomer, or stereoisomer thereof;


Wherein,

R¹ each independently represents Oxygen (O), D, Hydrogen (H) or Flourine (F);

R³ each independently represents H, D,







a is independently 2,3 or 7;

each b is independently 3, 5 or 6;

e is independently 1, 2 or 6;

c and d are each independently H, D, -OH, -OD, C₁-C₆-alkyl, -NH₂ or -COCH₃.

3. A Pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.
4. A Pharmaceutical composition comprising a compound of claim 2 and a pharmaceutically acceptable carrier.
5. The pharmaceutical composition of claim 3, which is formulated to treat the underlying etiology with an effective amount administering the patient in need by oral administration, delayed release or sustained release, transmucosal, syrup, topical, parenteral administration, injection, subdermal, oral solution, rectal administration, buccal administration or transdermal administration.
6. The pharmaceutical composition of claim 4, which is formulated to treat the underlying etiology with an effective amount administering the patient in need by

oral administration, delayed release or sustained release, transmucosal, syrup, topical, parenteral administration, injection, subdermal, oral solution, rectal administration, buccal administration or transdermal administration.

7. Compounds and compositions of claim 5 are formulated for the treatment of epilepsy, bipolar disorder, trigeminal neuralgia, attention-deficit hyperactivity disorder (ADHD), schizophrenia, neuropathic pain, seizures, bipolar disorder, mania, phantom limb syndrome, complex regional pain syndrome, paroxysmal extreme pain disorder, neuromyotonia, intermittent explosive disorder, borderline personality disorder, Myotonia congenita and post-traumatic stress disorder.
8. Compounds and compositions of claim 6 are formulated for the treatment of epilepsy, bipolar disorder, trigeminal neuralgia, attention-deficit hyperactivity disorder (ADHD), schizophrenia, neuropathic pain, seizures, bipolar disorder, mania, phantom limb syndrome, complex regional pain syndrome, paroxysmal extreme pain disorder, neuromyotonia, intermittent explosive disorder, borderline personality disorder, Myotonia congenita and post-traumatic stress disorder.
9. A molecular conjugate of 5H-dibenzo[b,f]azepine-5-carboxamide and R-Lipoic acid.
10. A molecular conjugate of 5H-dibenzo[b,f]azepine-5-carboxamide and eicosapentaenoic acid.
11. A molecular conjugate of 5H-dibenzo[b,f]azepine-5-carboxamide and docosahexaenoic acid.
12. A molecular conjugate of 10-oxo-10,11-dihydro-5H-dibenzo[b,f]azepine-5-carboxamide and R-Lipoic acid.
13. A molecular conjugate of 10-oxo-10,11-dihydro-5H-dibenzo[b,f]azepine-5-carboxamide and eicosapentaenoic acid.
14. A molecular conjugate of 10-oxo-10,11-dihydro-5H-dibenzo[b,f]azepine-5-carboxamide and docosahexaenoic acid.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2013/050711

A. CLASSIFICATION OF SUBJECT MATTER

See the extra sheet

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CA, EPOQUE, WPI, Google, CNTXT: STN structure search, CAS number 85756-57-6, 298-46-4, 28721-07-5, carmazepine, oxcarbazepine, conjugate, conjugates, lipoic acid, eicosapentaenoic acid, docosahexaenoic acid

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94/13298 A1 (RHONEPOULENC RORER S.A.), 23 June 1994 (23.06.1994), see claim 1	1,3,5,7
X	WO 2004/031155 A1 (NOVARTIS AG), 15 April 2004 (15.04.2004), see claims 1-14	2,4,6,8
A	US 4404366 A (MILES LABORATORIES, INC.), 13 September 1983 (13.09.1983), see column 14 lines 23-59	9-14
A	EP 0141393 A2 (ABBOTT LABORATORIES), 15 May 1985 (15.05.1985), see claims 1-8	9-14

Further documents are listed in the continuation of Box C.

See patent family annex.

<p>* Special categories of cited documents:</p> <p>“A” document defining the general state of the art which is not considered to be of particular relevance</p> <p>“E” earlier application or patent but published on or after the international filing date</p> <p>“L” document which may throw doubts on priority claim (S) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>“O” document referring to an oral disclosure, use, exhibition or other means</p> <p>“P” document published prior to the international filing date but later than the priority date claimed</p>	<p>“T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>“X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>“Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>“&” document member of the same patent family</p>
--	---

Date of the actual completion of the international search
01 July 2013 (01.07.2013)

Date of mailing of the international search report
25 Jul. 2013 (25.07.2013)

Name and mailing address of the ISA/CN
The State Intellectual Property Office, the P.R.China
6 Xitucheng Rd., Jimen Bridge, Haidian District, Beijing, China
100088
Facsimile No. 86-10-62019451

Authorized officer
WANG Jingjing
Telephone No. (86-10)62411133

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2013/050711

A. CLASSIFICATION OF SUBJECT MATTER

C07D 223/20 (2006.01)i

C07D 223/22 (2006.01)i

A61K 31/55 (2006.01)i

A61P 25/00 (2006.01)i

A61P 25/18 (2006.01)i

A61P 25/28 (2006.01)i

According to International Patent Classification (IPC) or to both national classification and IPC

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/IB2013/050711

Patent Documents referred in the Report	Publication Date	Patent Family	Publication Date
WO 94/13298 A1	23.06.1994	AU 5653994 A	04.07.1997
		NO 952229 A	06.06.1995
		SK 78595 A3	08.05.1996
		JPH 08504429 A	14.05.1996
		HUT 71839 A	28.02.19996
WO 2004/031155 A1	15.04.2004	AU 2003276055 A1	23.04.2004
		EP 1551808 A1	13.07.2005
		NO 20052244 A1	07.07.2005
		BR 0315113 A	23.08.2005
		TW 200413324 A	01.08.2004
		JP 2006504710 A	09.02.2006
		ZA 200502561 A	22.02.2006
		MXPA 05003737 A	01.07.2005
		CN 1703404 A	30.11.2005
		KR 20050071549 A	07.07.2005
		US 2006142566 A1	29.06.2006
		INCHENP 200500552 E	22.06.02007
		CN 101062932 A	31.10.2007
		AU 2003276055 B	03.01.2008
INCHENP 200702174 E	07.09.2007		
US 4404366 A	13.09.1983	none	
EP 0141393 A2	15.05.1985	AU 3471284 A	09.05.1985
		JPS 60120892 A	28.06.1985
		ZA 8408272 A	23.04.1985
		DK 8405228 A	05.05.1985
		US 4543412 A	24.09.1985
		US 4559173 A	17.12.1985

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/IB2013/050711

Patent Documents referred in the Report	Publication Date	Patent Family	Publication Date
		ES 8700672 A	16.01.1987
		CA 1224783 A	28.07.1987
		IL 73293 A	29.02.1988
		EP 0141393 B	17.08.1988
		DE 3473456 G	22.09.1988
		KR 900001620 B	17.03.1990
		JPH 0455198 B	02.09.1992