

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 December 2010 (02.12.2010)

(10) International Publication Number
WO 2010/138868 A2

(51) International Patent Classification:

A61K 9/12 (2006.01)

(74) Agent: **WEBB, Samuel E.**; STOEL RIVES LLP, 600 University Street, Suite 3600, Seattle, Washington 98101 (US).

(21) International Application Number:

PCT/US2010/036659

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:

28 May 2010 (28.05.2010)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/182,565	29 May 2009 (29.05.2009)	US
61/258,172	4 November 2009 (04.11.2009)	US
61/309,365	1 March 2010 (01.03.2010)	US
61/345,536	17 May 2010 (17.05.2010)	US

(71) Applicant (for all designated States except US): **PEARL THERAPEUTICS, INC.** [US/US]; 200 Saginaw Drive, Redwood City, California 94063 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VEHRING, Reinhard** [US/US]; 200 Saginaw Drive, Redwood City, California 94063 (US). **HARTMAN, Michael Steven** [US/US]; 3820 Park Blvd., Apt. #23, Palo Alto, California 94306 (US). **SMITH, Adrian Edward** [US/US]; 2 Wilmington Acres Court, Emerald Hills, California 94062 (US). **JOSHI, Vidya B.** [US/US]; 812 Seal Pointe Drive, Redwood City, California 94065 (US). **DWIVEDI, Sarvajna Kumar** [IN/US]; 409 Whisper Lane, Redwood City, California 94065 (US).

Published:

— without international search report and to be republished upon receipt of that report (Rule 48.2(g))

(54) Title: COMPOSITIONS FOR PULMONARY DELIVERY OF LONG-ACTING MUSCARINIC ANTAGONISTS AND LONG-ACTING β_2 ADRENERGIC RECEPTOR AGONISTS AND ASSOCIATED METHODS AND SYSTEMS




FIG. 8

(57) Abstract: Compositions, methods and systems are provided for pulmonary delivery of long-acting muscarinic antagonists and long-acting β_2 adrenergic receptor agonists via a metered dose inhaler. In particular embodiments, the compositions include a suspension medium, active agent particles, and suspending particles, in which the active agent particles and suspending particles form a co-suspension within the suspension medium.

WO 2010/138868 A2

COMPOSITIONS FOR PULMONARY DELIVERY OF LONG-ACTING MUSCARINIC ANTAGONISTS AND LONG-ACTING B₂ADRENERGIC RECEPTOR AGONISTS AND ASSOCIATED METHODS AND SYSTEMS

Technical Field

[0001] The present disclosure relates generally to pharmaceutical formulations and methods for delivery of active agents via the respiratory tract. In certain aspects, the present disclosure relates to compositions, methods, and systems for pulmonary delivery of long-acting muscarinic antagonists and long-acting β₂ adrenergic receptor agonists via a metered dose inhaler.

Background

[0002] Methods of targeted drug delivery that deliver an active agent at the site of action are often desirable. For example, targeted delivery of active agents can reduce undesirable side effects, lower dosing requirements and decrease therapeutic costs. In the context of respiratory delivery, inhalers are well known devices for administering an active agent to a subject's respiratory tract, and several different inhaler systems are currently commercially available. Three common inhaler systems include dry powder inhalers, nebulizers and metered dose inhalers (MDIs).

[0003] MDIs may be used to deliver medicaments in a solubilized form or as a suspension. Typically, MDIs use a relatively high vapor pressure propellant to expel aerosolized droplets containing an active agent into the respiratory tract when the MDI is activated. Dry powder inhalers generally rely on the patient's inspiratory efforts to introduce a medicament in a dry powder form to the respiratory tract. On the other hand, nebulizers form a medicament aerosol to be inhaled by imparting energy to a liquid solution or suspension.

[0004] MDIs are active delivery devices that utilize the pressure generated by a propellant. Conventionally, chlorofluorocarbons (CFCs) have been used as propellants in MDI systems because of their low toxicity, desirable vapor pressure and suitability for formulation of stable suspensions. However, traditional CFC propellants are understood to have a negative environmental impact, which has led to the development of alternative propellants that are believed to be more

environmentally-friendly, such as perfluorinated compounds (PFCs) and hydrofluoroalkanes (HFAs).

[0005] The active agent to be delivered by a suspension MDI is typically provided as a fine particulate dispersed within a propellant or combination of two or more propellants (i.e., a propellant “system”). In order to form the fine particulates, the active agent is typically micronized. Fine particles of active agent suspended in a propellant or propellant system tend to aggregate or flocculate rapidly. This is particularly true of active agents present in micronized form. In turn, aggregation or flocculation of these fine particles may complicate the delivery of the active agent. For example, aggregation or flocculation can lead to mechanical failures, such as those that might be caused by obstruction of the valve orifice of the aerosol container. Unwanted aggregation or flocculation of drug particles may also lead to rapid sedimentation or creaming of drug particles, and such behavior may result in inconsistent dose delivery, which can be particularly troublesome with highly potent, low dose medicaments. Another problem associated with such suspension MDI formulations relates to crystal growth of the drug during storage, resulting in a decrease over time of aerosol properties and delivered dose uniformity of such MDIs. More recently, solution approaches, such as those disclosed in U.S. Patent No. 6,964,759, have been proposed for MDI formulations containing anticholinergics.

[0006] One approach to improve aerosol performance in dry powder inhalers has been to incorporate fine particle carrier particles, such as lactose. Use of such fine excipients has not been investigated to any great extent for MDIs. A recent report by Young et al., “The influence of micronized particulates on the aerosolization properties of pressurized metered dose inhalers”; *Aerosol Science* 40, pgs. 324-337 (2009), suggests that the use of such fine particle carriers in MDIs actually result in a decrease in aerosol performance.

[0007] In traditional CFC systems, when the active agent present in an MDI formulation is suspended in the propellant or propellant system, surfactants are often used to coat the surfaces of the active agent in order to minimize or prevent the problem of aggregation and maintain a substantially uniform dispersion. The use of surfactants in this manner is sometimes referred to as “stabilizing” the suspension. However, many surfactants that are soluble and thus effective in CFC systems are

not effective in HFA and PFC propellant systems because such surfactants exhibit different solubility characteristics in non-CFC propellants.

Brief Description of the Drawings

[0008] FIG. 1 is a graph, which depicts the particle size distribution exhibited by an exemplary co-suspension composition according to the present description, which included glycopyrrolate, a long-acting muscarinic antagonist, as the active agent. Co-suspension MDIs were subjected to temperature cycling conditions (alternating 6h hold time at -5 or 40 °C) for 12 weeks.

[0009] FIG. 2 is a graph, which depicts the particle size distribution exhibited by an exemplary co-suspension composition according to the present description, which included glycopyrrolate, a long-acting muscarinic antagonist, as the active agent. Co-suspension MDIs were subjected to temperature cycling conditions (alternating 6h hold time at -5 or 40 °C) for 24 weeks.

[0010] FIG. 3 provides a micrograph illustrating the morphologies of a variety of suspending particles prepared according to Example 5.

[0011] FIG. 4 is a photograph of two vials that allows visualization of a co-suspension formed using active agent particles formed using glycopyrrolate and suspending particles formed using a saccharide.

[0012] FIG. 5 is a graph, which depicts the serum glycopyrrolate concentration level achieved over a period of 24 hours after a single administration of four different doses of glycopyrrolate delivered from a co-suspension composition as described herein.

[0013] FIG. 6 is a graph, which depicts the mean change in FEV₁ from baseline (in liters) experienced in patients over a period of 24 hours after receiving a single administration of the indicated dose of glycopyrrolate formulated in a co-suspension as described herein. In this study, Spiriva (18 µg Tiotropium) was included as an active control, and the mean change in FEV₁ from baseline (in liters) experienced in patients receiving a single administration of Spiriva is also depicted.

[0014] FIG. 7 is a bar graph, which depicts the peak change in FEV₁ from baseline (in liters) experienced in patients after receiving a single administration of the indicated dose of glycopyrrolate formulated in a co-suspension as described herein relative to placebo, the area under the curve of the FEV₁ over 12 hours after

dosing, and the area under the curve of the FEV₁ over 24 hours after dosing relative to placebo across the four doses evaluated. In this study, Spiriva (18 µg Tiotropium) was included as an active control and the results following single administration of Spiriva for the above parameters are also depicted in this figure.

[0015] FIG. 8 is a graph, which depicts the proportion of patients which achieved a greater than 12% change in FEV₁ from baseline and an improvement of 150 mL change from baseline or an absolute improvement of 200 mL from baseline regardless of % change in FEV₁ from baseline, after receiving a single administration of the indicated doses of a glycopyrrolate co-suspension as described herein. In this study, Spiriva (18 µg Tiotropium) was included as an active control and the results following single administration of Spiriva for the above parameter are also depicted on this figure.

[0016] FIG. 9 is a bar graph, which depicts the peak change in inspiratory capacity experienced in patients after receiving a single administration of the indicated doses of a glycopyrrolate co-suspension as described herein. In this study, Spiriva (18 µg Tiotropium) was included as an active control and the results following single administration of Spiriva for the above parameter are also depicted on this figure.

[0017] FIG. 10 is a bar graph providing the change in FEV₁ AUC achieved in patients after receiving a single administration of the indicated doses of a glycopyrrolate co-suspension as described herein. The results achieved by the glycopyrrolate co-suspension according to the present description are shown in comparison with the change in FEV₁ AUC reported in a published study in patients who received a powder formulation of glycopyrrolate not prepared according to the teachings provided herein.

[0018] FIG. 11 is a graph, which depicts the particle size distribution of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 4.5 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles and subjected to temperature cycling conditions (alternating 6h hold time at -5 or 40 °C).

[0019] FIG. 12 is a graph, which depicts the particle size distribution of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 36 µg/actuation delivered dose of glycopyrrolate and 6mg/mL

suspending particles and subjected to temperature cycling conditions (alternating 6h hold time at -5 or 40 °C).

[0020] FIG. 13 is a graph, which depicts the delivered dose through canister life of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 4.5 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles.

[0021] FIG. 14 is a graph, which depicts the delivered dose through canister life of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 36 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles.

[0022] FIG. 15 is a graph, which depicts the particle size distribution of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 36 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles and subjected to 12 months storage at 25°C/60% RH unprotected.

[0023] FIG. 16 is a graph, which depicts the mean delivered dose through canister life of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 32 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles and subjected to temperature cycling conditions (alternating 6h hold time at -5 or 40 °C).

[0024] FIG. 17 is a graph, which depicts the particle size distribution of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 32 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles and subjected to temperature cycling conditions (alternating 6h hold time at -5 or 40 °C)

[0025] FIG. 18 is a graph, which depicts the particle size distribution of an exemplary glycopyrrolate co-suspension prepared according to the present description, containing 24 µg/actuation delivered dose of glycopyrrolate and 6mg/mL suspending particles and subjected to 6 weeks storage at 50°C/ambient relative humidity and 12 weeks at 40°C

[0026] FIG. 19 is a photograph that allows visualization of co-suspension compositions prepared according to the present description which include formoterol fumarate active agent particles.

[0027] FIG. 20 is a graph, which depicts the delivered dose uniformity achieved by formoterol fumarate co-suspension compositions prepared according to the present description.

[0028] FIG. 21 is a graph, which depicts the aerodynamic particle size distribution determined by cascade impaction of exemplary formoterol fumarate co-suspension compositions prepared according to the present description and stored for three months at 25 °C/ 75% RH, without protective overwrap, or at 40 °C / 75%RH with protective overwrap.

[0029] FIG. 22 is a graph, which depicts the chemical stability of exemplary co-suspension compositions including formoterol fumarate as the active agent. The results depicted in this figure allow comparison of the chemical stability of formoterol fumarate achieved in a co-suspension composition formulated using crystalline formoterol fumarate with the chemical stability of suspension formulations prepared using spray dried formoterol fumarate.

[0030] FIG. 23 through FIG. 26 are electron micrographs of suspending particles prepared from various different materials, with Figure 23 providing a micrograph of trehalose suspending particles, Figure 24 providing a micrograph of HP- β -cyclodextrin suspending particles, Figure 25 providing a micrograph of Ficoll MP 70 suspending particles, and Figure 26 providing a micrograph of inulin suspending particles.

[0031] FIG. 27 provides a graph that depicts the aerodynamic particle size distribution determined by cascade impaction of exemplary co-suspension compositions prepared according to the present description and including glycopyrrolate active agent particles.

[0032] FIG. 28 provides a graph that depicts the aerodynamic particle size distribution determined by cascade impaction of exemplary co-suspension compositions prepared according to the present description and including formoterol fumarate active agent particles.

[0033] FIG. 29 provides a graph that depicts the delivered dose uniformity achieved by ultra low-dose formoterol fumarate co-suspension compositions prepared according to the present description.

[0034] FIG. 30 provides graphs illustrating the particle size distribution of glycopyrrolate (top) and formoterol (bottom) achieved by an exemplary co-

suspension compared to particle size distributions achieved by formulations including either glycopyrrolate or formoterol fumarate alone.

Detailed Description

[0035] The present disclosure provides compositions, methods, and systems for respiratory delivery of active agents via an MDI. In particular embodiments, the compositions, methods and systems described herein are adapted for respiratory delivery of active agents selected from a long-acting muscarinic antagonist ("LAMA") and a long-acting β_2 adrenergic receptor agonist ("LABA"). In certain embodiments, the LAMA or LABA active agent may be potent or highly potent and, therefore, formulated at low concentrations and delivered in low doses. The pharmaceutical compositions described herein may be formulated for pulmonary or nasal delivery via an MDI. The methods described herein include methods of stabilizing formulations including LAMA or LABA active agents for respiratory delivery, as well as methods for pulmonary delivery of LAMA and LABA active agents via an MDI. Also described herein are methods for preparing an MDI for delivery of a LAMA or LABA active agent.

[0036] In specific embodiments, the methods described herein include methods for treating a pulmonary disease or disorder amenable to treatment by delivery of a LAMA or LABA active agent through an MDI. For example, and the compositions, methods and systems described herein can be used to treat inflammatory or obstructive pulmonary diseases or conditions. In certain embodiments, the compositions, methods and systems described herein can be used to treat patients suffering from a disease or disorder selected from asthma, chronic obstructive pulmonary disease (COPD), exacerbation of airways hyper reactivity consequent to other drug therapy, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, pulmonary hypertension, pulmonary vasoconstriction, and any other respiratory disease, condition, trait, genotype or phenotype that can respond to the administration of a LAMA or LABA, alone or in combination with other therapies. In certain embodiments, the compositions, systems and methods described herein can be used to treat pulmonary inflammation and obstruction associated with cystic fibrosis. As used herein, the terms "COPD" and "chronic obstructive pulmonary

disease" encompass chronic obstructive lung disease (COLD), chronic obstructive airway disease (COAD), chronic airflow limitation (CAL) and chronic obstructive respiratory disease (CORD) and include chronic bronchitis, bronchiectasis, and emphysema. As used herein, the term "asthma" refers to asthma of whatever type or genesis, including both intrinsic (non-allergic) asthma and extrinsic (allergic) asthma, mild asthma, moderate asthma, severe asthma, bronchitic asthma, exercise-induced asthma, occupational asthma and asthma induced following bacterial infection. Asthma is also to be understood as embracing wheezy-infant syndrome.

[0037] It will be readily understood that the embodiments described herein are exemplary. The following more detailed description of various embodiments is not intended to limit the scope of the present disclosure, but is merely representative of various embodiments. Moreover, the order of the steps or actions of the methods described in connection with the embodiments disclosed herein may be changed by those skilled in the art without departing from the scope of the present disclosure. In other words, unless a specific order of steps or actions is required for proper operation of the embodiment, the order or use of specific steps or actions may be modified.

I. Definitions

[0038] Unless specifically defined otherwise, the technical terms, as used herein, have their normal meaning as understood in the art. The following terms are specifically defined for the sake of clarity.

[0039] The term "active agent" is used herein to include any agent, drug, compound, composition or other substance that may be used on, or administered to a human or animal and is a LAMA or LABA. The term "active agent" may be used interchangeably with the terms, "drug," "pharmaceutical," "medicament," "drug substance," or "therapeutic."

[0040] The terms "associate," "associate with" or "association" refers to an interaction or relationship between a chemical entity, composition, or structure in a condition of proximity to a surface, such as the surface of another chemical entity, composition, or structure. The association includes, for example, adsorption, adhesion, covalent bonding, hydrogen bonding, ionic bonding and electrostatic

attraction, Lifshitz-van der Waals interactions and polar interactions. The term “adhere” or “adhesion” is a form of association and is used as a generic term for all forces tending to cause a particle or mass to be attracted to a surface. “Adhere” also refers to bringing and keeping particles in contact with each other, such that there is substantially no visible separation between particles due to their different buoyancies in a propellant under normal conditions. In one embodiment, a particle that attaches to or binds to a surface is encompassed by the term “adhere.” Normal conditions may include storage at room temperature or under an accelerative force due to gravity. As described herein, active agent particles may associate with suspending particles to form a co-suspension, where there is substantially no visible separation between the suspending particles and the active agent particles or flocculates thereof due to differences in buoyancy within a propellant.

[0041] “Suspending particles” refer to a material or combination of materials that is acceptable for respiratory delivery, and acts as a vehicle for active agent particles. Suspending particles interact with the active agent particles to facilitate repeatable dosing, delivery or transport of active agent to the target site of delivery, i.e., the respiratory tract. The suspending particles described herein are dispersed within a suspension medium including a propellant or propellant system, and can be configured according to any shape, size or surface characteristic suited to achieving a desired suspension stability or active agent delivery performance. Exemplary suspending particles include particles that exhibit a particle size that facilitates respiratory delivery of active agent and have physical configurations suited to formulation and delivery of the stabilized suspensions as described herein.

[0042] The term “co-suspension” refers to a suspension of two or more types of particles having different compositions within a suspension medium, wherein one type of particle associates at least partially with one or more of the other particle types. The association leads to an observable change in one or more characteristics of at least one of the individual particle types suspended in the suspension medium. Characteristics modified by the association may include, for example, one or more of the rate of aggregation or flocculation, the rate and nature of separation, i.e. sedimentation or creaming, density of a cream or sediment layer, adhesion to container walls, adhesion to valve components, and rate and the level of dispersion upon agitation.

[0043] Exemplary methods for assessing whether a co-suspension is present can include the following: If one particle type has a pycnometric density greater than the propellant and another particle type has a pycnometric density lower than the propellant, a visual observation of the creaming or sedimentation behavior can be employed to determine the presence of a co-suspension. The term “pycnometric density” refers to the density of a material that makes up a particle, excluding voids within the particle. In one embodiment, the materials can be formulated or transferred into a transparent vial, typically a glass vial, for visual observation. After initial agitation the vial is left undisturbed for a sufficient time for formation of a sediment or cream layer, typically 24 hours. If the sediment or cream layer is observed to be completely or mostly a uniform single layer, a co-suspension is present. The term “co-suspension” includes partial co-suspensions, where a majority of the at least two particle types associate with each other, however, some separation (i.e., less than a majority) of the at least two particle types may be observed.

[0044] The exemplary co-suspension test may be performed at different propellant temperatures to accentuate the sedimentation or creaming behavior of particle types with a density close to the propellant density at room temperature. If the different particle types have the same nature of separation, i.e. all sediment or all cream, the presence of a co-suspension can be determined by measuring other characteristics of the suspension, such as rate of aggregation or flocculation, rate of separation, density of cream or sediment layer, adhesion to container walls, adhesion to valve components, and rate and level of dispersion upon agitation, and comparing them to the respective characteristics of the similarly suspended individual particle types. Various analytical methods generally known to those skilled in the art can be employed to measure these characteristics.

[0045] In the context of a composition containing or providing respirable aggregates, particles, drops, etc., such as compositions described herein, the term “fine particle dose” or “FPD” refers to the dose, either in total mass or fraction of the nominal dose or metered dose, that is within a respirable range. The dose that is within the respirable range is measured *in vitro* to be the dose that deposits beyond the throat stage of a cascade impactor, i.e., the sum of dose delivered at stages 3 through filter in a Next Generation Impactor operated at a flow rate of 30 l/min.

[0046] In the context of a composition containing or providing respirable aggregates, particles, drops, etc., such as compositions described herein, the term “fine particle fraction” or “FPF” refers to the proportion of the delivered material relative to the delivered dose (i.e., the amount that exits the actuator of a delivery device, such as an MDI) that is within a respirable range. The amount of delivered material within the respirable range is measured *in vitro* as the amount of material that deposits beyond the throat stage of a cascade impactor, e.g., the sum of the material delivered at stages 3 through filter in a Next Generation Impactor operated at a flow rate of 30 l/min.

[0047] As used herein, the term “inhibit” refers to a measurable lessening of the tendency of a phenomenon, symptom or condition to occur or the degree to which that phenomenon, symptom or condition occurs. The term “inhibit” or any form thereof, is used in its broadest sense and includes minimize, prevent, reduce, repress, suppress, curb, constrain, restrict, slow progress of and the like.

[0048] “Mass median aerodynamic diameter” or “MMAD” as used herein refers to the aerodynamic diameter of an aerosol below which 50% of the mass of the aerosol consists of particles with an aerodynamic diameter smaller than the MMAD, with the MMAD being calculated according to monograph 601 of the United States Pharmacopeia (“USP”).

[0049] When referred to herein, the term “optical diameter” indicates the size of a particle as measured by the Fraunhofer diffraction mode using a laser diffraction particle size analyzer equipped with a dry powder dispenser (e.g., Sympatec GmbH, Clausthal-Zellerfeld, Germany).

[0050] The term solution mediated transformation refers to the phenomenon in which a more soluble form of a solid material (i.e. particles with small radius of curvature (a driving force for Ostwald ripening), or amorphous material) dissolves and recrystallizes into the more stable crystal form that can coexist in equilibrium with its saturated propellant solution.

[0051] A “patient” refers to an animal in which LAMA or LABA active agents will have a therapeutic effect. In one embodiment, the patient is a human being.

[0052] “Perforated microstructures” refer to suspending particles that include a structural matrix that exhibits, defines or comprises voids, pores, defects, hollows, spaces, interstitial spaces, apertures, perforations or holes that allow the surrounding

suspension medium to permeate, fill or pervade the microstructure, such as those materials and preparations described in U.S. Patent No. 6,309,623 to Weers, et al. The primary form of the perforated microstructure is, generally, not essential, and any overall configuration that provides the desired formulation characteristics is contemplated herein. Accordingly, in one embodiment, the perforated microstructures may comprise approximately spherical shapes, such as hollow, suspending, spray-dried microspheres. However, collapsed, corrugated, deformed or fractured particulates of any primary form or aspect ratio may also be compatible.

[0053] As is true of suspending particles described herein, perforated microstructures may be formed of any biocompatible material that does not substantially degrade or dissolve in the selected suspension medium. While a wide variety of materials may be used to form the particles, in some embodiments, the structural matrix is associated with, or includes, a surfactant such as, a phospholipid or fluorinated surfactant. Although not required, the incorporation of a compatible surfactant in the perforated microstructure or, more generally, the suspending particles, can improve the stability of the respiratory dispersions, increase pulmonary deposition and facilitate the preparation of the suspension.

[0054] The term “suspension medium” as uses herein refers to a substance providing a continuous phase within which active agent particles and suspending particles can be dispersed to provide a co-suspension formulation. The suspension medium used in co-suspension formulations described herein includes propellant. As used herein, the term “propellant” refers to one or more pharmacologically inert substances which exert a sufficiently high vapor pressure at normal room temperature to propel a medicament from the canister of an MDI to a patient on actuation of the MDI’s metering valve. Therefore, the term “propellant” refers to both a single propellant and to a combination of two or more different propellants forming a “propellant system.”

[0055] The term “respirable” generally refers to particles, aggregates, drops, etc. sized such that they can be inhaled and reach the airways of the lung.

[0056] When used to refer to co-suspension compositions described herein, the terms “physical stability” and “physically stable” refer to a composition that is resistant to one or more of aggregation, flocculation, and particle size changes due to solution mediated transformations and is capable of substantially maintaining the

MMAD of suspending particles and the fine particle dose. In one embodiment, physical stability may be evaluated through subjecting compositions to accelerated degradation conditions, such as by temperature cycling as described herein.

[0057] When referring to active agents, the term "potent" indicates active agents that are therapeutically effective at or below doses ranging from about 0.01 mg/kg to about 1 mg/kg. Typical doses of potent active agents generally range from about 100 µg to about 100 mg.

[0058] When referring to active agents, the term "highly potent" indicates active agents that are therapeutically effective at or below doses of about 10 µg/kg. Typical doses of highly potent active agents generally range up to about 100 µg.

[0059] The terms "suspension stability" and "stable suspension" refer to suspension formulations capable of maintaining the properties of a co-suspension of active agent particles and suspending particles over a period of time. In one embodiment, suspension stability may be measured through delivered dose uniformity achieved by co-suspension compositions described herein.

[0060] The term "substantially insoluble" means that a composition is either totally insoluble in a particular solvent or it is poorly soluble in that particular solvent. The term "substantially insoluble" means that a particular solute has a solubility of less than one part per 100 parts solvent. The term "substantially insoluble" includes the definitions of "slightly soluble" (from 100 to 1000 parts solvent per 1 part solute), "very slightly soluble" (from 1000 to 10,000 parts solvent per 1 part solute) and "practically insoluble" (more than 10,000 parts solvent per 1 part solute) as given in Table 16-1 of Remington: The Science and Practice of Pharmacy, 21st ed. Lippincott, Williams & Wilkins, 2006, p. 212.

[0061] The term "surfactant," as used herein, refers to any agent which preferentially adsorbs to an interface between two immiscible phases, such as the interface between water and an organic polymer solution, a water/air interface or organic solvent/air interface. Surfactants generally possess a hydrophilic moiety and a lipophilic moiety, such that, upon adsorbing to microparticles, they tend to present moieties to the continuous phase that do not attract similarly-coated particles, thus reducing particle agglomeration. In some embodiments, surfactants may also promote adsorption of a drug and increase bioavailability of the drug.

[0062] A “therapeutically effective amount” is the amount of compound which achieves a therapeutic effect by inhibiting a disease or disorder in a patient or by prophylactically inhibiting or preventing the onset of a disease or disorder. A therapeutically effective amount may be an amount which relieves to some extent one or more symptoms of a disease or disorder in a patient; returns to normal either partially or completely one or more physiological or biochemical parameters associated with or causative of the disease or disorder; and/or reduces the likelihood of the onset of the disease or disorder.

[0063] The terms “chemically stable” and “chemical stability” refer to co-suspension formulations wherein the individual degradation products of active agent remain below the limits specified by regulatory requirements during the shelf life of the product for human use (e.g., 1% of total chromatographic peak area per ICH guidance Q3B(R2)) and there is acceptable mass balance (e.g., as defined in ICH guidance Q1E) between active agent assay and total degradation products.

II. Pharmaceutical Compositions

[0064] The compositions described herein are co-suspensions that include a suspension medium including a propellant, LAMA or LABA active agent particles, and suspending particles. Of course, if desired, the compositions described herein may include one or more additional constituents. Moreover, variations and combinations of components of the compositions described herein may be used. For example, two or more species of suspending particles may be used in compositions for the formulation and delivery of a selected LAMA or LABA active agent. Alternatively, for example, the compositions described herein may include two or more species of active agent particles. In certain such embodiments, the compositions may include LAMA or LABA active agent particles co-suspended with suspending particles, wherein, in addition to the active agent material included in the active agent particles, at least some of the suspending particles incorporate the selected LAMA or LABA active agent. Even further, if desired, the compositions described herein may include two or more different species of particles containing the selected LAMA or LABA active agent in combination with two or more different species of suspending particles.

[0065] It has been found that, in formulations according to the present description, active agent particles exhibit an association with the suspending particles such that separation of the active agent particles from the suspending particles is substantially prevented, resulting in co-location of active agent particles and suspending particles within the suspension medium. Generally, due to density differences between distinct species of particles and the medium within which they are suspended (e.g., a propellant or propellant system), buoyancy forces cause creaming of particles with lower density than the propellant and sedimentation of particles with higher density than the propellant. Therefore, in suspensions that include a mixture of particles that vary in their densities, the sedimentation or creaming behavior of each type of particle may vary and may lead to separation of the different particle types within the propellant.

[0066] However, the combinations of propellant, active agent particles and suspending particles described herein provide co-suspensions wherein the active agent particles and suspending particles co-locate within the propellant (i.e., the active agent particles associate with the suspending particles such that suspending particles and active agent particles do not exhibit substantial separation relative to each other, such as by differential sedimentation or creaming, even after a time sufficient for the formation of a cream or sediment layer). In particular embodiments, for example, the compositions described herein form co-suspensions wherein the suspending particles remain associated with active agent particles when subjected to buoyancy forces amplified by temperature fluctuations and/or centrifugation at accelerations up to an over, for example, 1 g, 10 g, 35 g, 50 g, and 100 g. However, the co-suspensions described herein need not be defined by or limited to a specific threshold force of association. For example, a co-suspension as contemplated herein may be successfully achieved where the active agent particles associate with the suspending particles such that there is no substantial separation of active agent particles and suspending particles within the continuous phase formed by the suspension medium under typical patient use conditions.

[0067] Co-suspension compositions according to the present description provide desirable formulation and delivery characteristics for LAMA and LABA active agents. For example, in certain embodiments, when present within an MDI canister, co-suspensions as described herein can inhibit or reduce one or more of the following:

flocculation of active agent material; differential sedimentation or creaming of active agent particles and suspending particles; solution mediated transformation of active agent material; and loss of active agent to the surfaces of the container closure system, in particular the metering valve components. In addition, compositions as described herein provide chemical stability for the active agents contained therein. Such qualities work to achieve and preserve aerosol performance as the co-suspension is delivered from an MDI such that desirable fine particle fraction, fine particle dose and delivered dose uniformity characteristics are achieved and substantially maintained throughout emptying of an MDI canister within which the co-suspension composition is contained. Additionally, as illustrated by embodiments detailed herein, co-suspensions according to the present description can provide a stable formulation that provides consistent dosing and respiratory delivery characteristics for LAMA and LABA active agents, while utilizing a relatively simple HFA suspension medium that does not require modification by the addition of, for example, cosolvents, antisolvents, solubilizing agents or adjuvants.

[0068] Providing a co-suspension according to the present description may also simplify formulation, delivery and dosing of LAMA and LABA active agents. Without being bound by a particular theory, it is thought that by achieving a co-suspension of active agent particles and suspending particles, the delivery and dosing of active agent contained within such a dispersion may be substantially controlled through control of the size, composition, morphology and relative amount of the suspending particles, and less dependent upon the size and morphology of the active agent particles.

[0069] Accordingly, the pharmaceutical compositions disclosed herein provide for delivery of LAMA and LABA active agents from an MDI. Delivery of the co-suspension compositions described herein provides desirable pharmacokinetic and pharmacodynamic characteristics, and MDI delivery of the pharmaceutical compositions described herein is suitable for treating patients suffering from an inflammatory or obstructive pulmonary disease or condition that responds to the administration of a LAMA or LABA active agent. In particular embodiments, the pharmaceutical compositions described herein may be used in treating a disease or condition selected from asthma, COPD, exacerbation of airways hyper reactivity consequent to other drug therapy, allergic rhinitis, sinusitis, pulmonary

vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, pulmonary hypertension, pulmonary vasoconstriction, emphysema, and any other respiratory disease, condition, trait, genotype or phenotype that can respond to the administration of a LAMA or LABA, alone or in combination with other therapies. In certain embodiments, the compositions, systems and methods described herein can be used to treat pulmonary inflammation and obstruction associated with cystic fibrosis.

(i) Suspension Medium

[0070] The suspension medium included in a composition described herein includes one or more propellants. In general, suitable propellants for use as suspension mediums are those propellant gases that can be liquefied under pressure at room temperature, and upon inhalation or topical use, are safe and toxicologically innocuous. Additionally, it is desirable that the selected propellant be relatively non-reactive with the suspending particles or active agent particles. Exemplary compatible propellants include hydrofluoroalkanes (HFAs), perfluorinated compounds (PFCs), and chlorofluorocarbons (CFCs).

[0071] Specific examples of propellants that may be used to form the suspension medium of the co-suspensions disclosed herein include 1,1,1,2-tetrafluoroethane ($\text{CF}_3\text{CH}_2\text{F}$) (HFA-134a), 1,1,1,2,3,3-heptafluoro-n-propane ($\text{CF}_3\text{CHFCF}_3$) (HFA-227), perfluoroethane, monochloro-fluoromethane, 1,1 difluoroethane, and combinations thereof. Even further, suitable propellants include, for example: short chain hydrocarbons; C_{1-4} hydrogen-containing chlorofluorocarbons such as CH_2ClF , $\text{CCl}_2\text{FCHClF}$, CF_3CHClF , $\text{CHF}_2\text{CClF}_2$, CHClFCHF_2 , $\text{CF}_3\text{CH}_2\text{Cl}$, and CClF_2CH_3 ; C_{1-4} hydrogen-containing fluorocarbons (e.g., HFAs) such as CHF_2CHF_2 , $\text{CF}_3\text{CH}_2\text{F}$, CHF_2CH_3 , and $\text{CF}_3\text{CHFCF}_3$; and perfluorocarbons such as CF_3CF_3 and $\text{CF}_3\text{CF}_2\text{CF}_3$.

[0072] Specific fluorocarbons, or classes of fluorinated compounds, that may be used as suspension media include, but are not limited to, fluoroheptane, fluorocycloheptane, fluoromethylcycloheptane, fluorohexane, fluorocyclohexane, fluoropentane, fluorocyclopentane, fluoromethylcyclopentane, fluorodimethylcyclopentanes, fluoromethylcyclobutane, fluorodimethylcyclobutane, fluorotrimethylcyclobutane, fluorobutane, fluorocyclobutane, fluoropropane, fluoroethers,

fluoropolyethers and fluorotriethylamines. These compounds may be used alone or in combination with more volatile propellants.

[0073] In addition to the aforementioned fluorocarbons and hydrofluoroalkanes, various exemplary chlorofluorocarbons and substituted fluorinated compounds may also be used as suspension media. In this respect, FC-11 (CCl₃F), FC-11B1 (CBrCl₂F), FC-11B2 (CBr₂ClF), FC12B2 (CF₂Br₂), FC21 (CHCl₂F), FC21B1 (CHBrClF), FC-21B2 (CHBr₂F), FC-31B1 (CH₂BrF), FC113A (CCl₃CF₃), FC-122 (CClF₂CHCl₂), FC-123 (CF₃CHCl₂), FC-132 (CHClFCHClF), FC-133 (CHClFCHF₂), FC-141 (CH₂ClCHClF), FC-141B (CCl₂FCH₃), FC-142 (CHF₂CH₂Cl), FC-151 (CH₂FCH₂Cl), FC-152 (CH₂FCH₂F), FC-1112 (CClF=CClF), FC-1121 (CHCl=CCl) and FC-1131 (CHCl=CHF) may also be used, while recognizing the possible attendant environmental concerns. As such, each of these compounds may be used, alone or in combination with other compounds (i.e., less volatile fluorocarbons) to form the stabilized suspensions disclosed herein.

[0074] In some embodiments, the suspension medium may be formed of a single propellant. In other embodiments, a combination of propellants (a "propellant system") may be used to form the suspension medium. In some embodiments, relatively volatile compounds may be mixed with lower vapor pressure components to provide suspension media having specified physical characteristics selected to improve stability or enhance the bioavailability of the dispersed active agent. In some embodiments, the lower vapor pressure compounds will comprise fluorinated compounds (e.g. fluorocarbons) having a boiling point greater than about 25°C. In some embodiments, lower vapor pressure fluorinated compounds for use in the suspension medium may include perfluoroctylbromide C₈F₁₇Br (PFOB or perflubron), dichlorofluorooctane C₈F₁₆Cl₂, perfluoroctylethane C₈F₁₇C₂H₅ (PFOE), perfluorodecylbromide C₁₀F₂₁Br (PFDB) or perfluorobutylethane C₄F₉C₂H₅. In certain embodiments, these lower vapor pressure compounds are present in a relatively low level. Such compounds may be added directly to the suspension medium or may be associated with the suspending particles.

[0075] The suspension medium included in compositions as described herein may be formed of a propellant or propellant system that is substantially free of additional materials, including, for example, antisolvents, solubilizing agents, cosolvents or adjuvants. For example, in some embodiments, the suspension

medium may be formed of a non-CFC propellant or propellant system, such as an HFA propellant or propellant system, that is substantially free of additional materials. Such embodiments simplify the formulation and manufacture of pharmaceutical compositions suited for respiratory delivery of a LAMA or LABA active agent.

[0076] However, in other embodiments, depending on the selection of propellant, the properties of the suspending particles, or the nature of active agent to be delivered, the suspension medium utilized may include materials in addition to the propellant or propellant system. Such additional materials may include, for example, one or more of an appropriate antisolvent, solubilizing agent, cosolvent or adjuvant to adjust, for example, the vapor pressure of the formulation or the stability, or solubility of suspended particles. For example, propane, ethanol, isopropyl alcohol, butane, isobutane, pentane, isopentane or a dialkyl ether, such as dimethyl ether, may be incorporated with the propellant in the suspension medium. Similarly, the suspension medium may contain a volatile fluorocarbon. In other embodiments, one or both of polyvinylpyrrolidone ("PVP") or polyethylene glycol ("PEG") may be added to the suspension medium. Adding PVP or PEG to the suspension medium may achieve one or more desired functional characteristics, and in one example, PVP or PEG may be added to the suspension medium as a crystal growth inhibitor. In general, where a volatile cosolvent or adjuvant is used, such an adjuvant or cosolvent may be selected from known hydrocarbon or fluorocarbon materials and may account for up to about 1% w/w of the suspension medium. For example, where a cosolvent or adjuvant is incorporated in the suspension medium, the cosolvent or adjuvant may comprise less than about 0.01%, 0.1%, or 0.5% w/w of the suspension medium. Where PVP or PEG are included in the suspension medium, such constituents may be included at up to about 1% w/w, or they may comprise less than about 0.01%, 0.1%, or 0.5% w/w of the suspension medium.

(ii) Active Agent Particles

[0077] The active agent particles included in the co-suspensions described herein are formed to be capable of being dispersed and suspended within the suspension medium and are sized to facilitate delivery of respirable particles from the co-suspension. In one embodiment, therefore, the active agent particles are provided as a micronized material wherein at least 90% of the active agent particle material by

volume exhibits an optical diameter of about 7 μm or less. In other embodiments, the active agent particles are provided as a micronized material wherein at least 90% of the active agent particles by volume exhibit an optical diameter selected from a range of about 6 μm to about 1 μm , about 5 μm to about 2 μm , and about 4 μm to about 3 μm . In further embodiments, the active agent particles are provided as a micronized material wherein at least 90% of the active agent particles by volume exhibit an optical diameter selected from 6 μm or less, 5 μm or less, and 4 μm or less. In another embodiment, the active agent particles are provided as a micronized material wherein at least 50% of the active agent particle material by volume exhibits an optical diameter of about 5 μm or less. In other embodiments, the active agent particles are provided as a micronized material wherein at least 50% of the active agent particles by volume exhibit an optical diameter selected from a range of about 4 μm to about 1 μm , about 3 μm to about 1 μm , and about 2.5 μm to about 1 μm . In another embodiment, the active agent particles are provided as a micronized material wherein at least 50% of the active agent particles by volume exhibit an optical diameter selected from 4 μm or less, 3 μm or less, and 2 μm or less.

[0078] In specific embodiments, the active agent material used as or to form the active agent particles may be entirely or substantially crystalline, i.e., a majority of the active agent molecules are arranged in a regularly repeating pattern, over a long range or external face planes. In another embodiment, the active agent particles may be present in both crystal and amorphous states. In yet another embodiment, the active agent particles may be present in substantially an amorphous state, i.e., the active agent particles are overall noncrystalline in nature and do not have a regularly repeating arrangement of molecules maintained over along range. Suitable excipients for formulation of active agent particles include those described herein in association with the suspending particles. In specific embodiments, for example, active agent particles may be formulated with one or more of the lipid, phospholipid, carbohydrate, amino acid, organic salt, peptide, protein, alditols, synthetic or natural polymer, or surfactant materials as described, for example, in association with the suspending particles. In other embodiments, the active agent particles are formed solely from micronized active agent material.

[0079] Because the compositions disclosed enable the formulation and reproducible delivery of very low doses of active agents, in certain embodiments, the active agents included in the compositions described herein may be selected from one or more potent or highly potent active agents. For example, in certain embodiments, the compositions described herein may include a potent active agent that is delivered at a single administration dose selected from between about 100 µg and about 100 mg per dose, about 100 µg and about 10 mg per dose, and about 100 µg and 1 mg per dose. In other embodiments, the compositions described herein may include a potent or highly potent active agent that is delivered at a dose selected from up to about 80 µg per single administration dose, up to about 40 µg per single administration dose, up to about 20 µg per single administration dose, up to about 10 µg per single administration dose or between about 10 µg and about 100 µg per single administration dose. Additionally, in certain embodiments, the compositions described herein may include a highly potent active agent delivered at a dose selected from between about 0.1 and about 2 µg per single administration dose, about 0.1 and about 1 µg per single administration dose, and about 0.1 and about 0.5 µg per single administration dose.

[0080] In certain embodiments, the active agent included in the compositions described herein is a LAMA active agent. Where the compositions include a LAMA active agent, in particular embodiments, the LAMA active agent may be selected from, for example, glycopyrrolate, dexipirronium, tiotropium, trospium, aclidinium, darotropium, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

[0081] Glycopyrrolate can be used to treat inflammatory or obstructive pulmonary diseases and disorders such as, for example, those described herein. As an anticholinergic, glycopyrrolate acts as a bronchodilator and provides an antisecretory effect, which is a benefit for use in the therapy of pulmonary diseases and disorders characterized by increased mucus secretions. Glycopyrrolate is a quaternary ammonium salt. Where appropriate, glycopyrrolate may be used in the form of salts (e.g. alkali metal or amine salts, or as acid addition salts) or as esters or as solvates (hydrates). Additionally, the glycopyrrolate may be in any crystalline form or isomeric form or mixture of isomeric forms, for example a pure enantiomer, a mixture of enantiomers, a racemate or a mixture thereof. In this regard, the form of

glycopyrrolate may be selected to optimize the activity and/or stability of glycopyrrolate and/or to minimize the solubility of glycopyrrolate in the suspension medium. Suitable counter ions are pharmaceutically acceptable counter ions including, for example, fluoride, chloride, bromide, iodide, nitrate, sulfate, phosphate, formate, acetate, trifluoroacetate, propionate, butyrate, lactate, citrate, tartrate, malate, maleate, succinate, benzoate, p-chlorobenzoate, diphenyl-acetate or triphenylacetate, o-hydroxybenzoate, p-hydroxybenzoate, 1-hydroxynaphthalene-2-carboxylate, 3-hydroxynaphthalene-2-carboxylate, methanesulfonate and benzenesulfonate. In particular embodiments of the compositions described herein, the bromide salt of glycopyrrolate, namely 3-[(cyclopentyl-hydroxyphenylacetyl)oxy]-1,1-dimethylpyrrolidinium bromide, is used and can be prepared according to the procedures set out in U.S. Pat. No. 2,956,062.

[0082] Where the compositions described herein include glycopyrrolate, in certain embodiments, the compositions may include sufficient glycopyrrolate to provide a target delivered dose selected from between about 10 µg and about 200 µg per actuation of an MDI, about 15 µg and about 150 µg per actuation of an MDI, and about 18 µg and 144 µg per actuation of an MDI. In other such embodiments, the formulations include sufficient glycopyrrolate to provide a dose selected from up to about 200 µg, up to about 150 µg, up to about 75 µg, up to about 40 µg, or up to about 20 µg per actuation. In yet further embodiments, the formulations include sufficient glycopyrrolate to provide a dose selected from about 18 µg per actuation, 36 µg per actuation, or about 72 µg per actuation. In order to achieve targeted delivered doses as described herein, where compositions described herein include glycopyrrolate as the active agent, in specific embodiments, the amount of glycopyrrolate included in the compositions may be selected from, for example, between about 0.04 mg/ml and about 2.25 mg/ml.

[0083] In other embodiments, tiotropium, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, may be selected as a LAMA active agent for inclusion in a composition as described herein. Tiotropium is a known, long-acting anticholinergic drug suitable for use in treating diseases or disorders associated with pulmonary inflammation or obstruction, such as those described herein. Tiotropium, including crystal and pharmaceutically acceptable salt forms of tiotropium, is described, for example, in U.S. Patent No. 5,610,163, U.S.

Patent No. RE39820, U.S. Patent No. 6,777,423, and U.S. Patent No. 6,908,928. Where the compositions described herein include tiotropium, in certain embodiments, the compositions may include sufficient tiotropium to provide a delivered dose selected from between about 2.5 µg and about 50 µg, about 4 µg and about 25 µg, about 2.5 µg and about 20 µg, about 10 µg and about 20 µg, and about 2.5 µg and about 10 µg per actuation of an MDI. In other such embodiments, the formulations include sufficient tiotropium to provide a delivered dose selected from up to about 50 µg, up to about 20 µg, up to about 10 µg, up to about 5 µg, or up to about 2.5 µg per actuation of an MDI. In yet further embodiments, the formulations include sufficient tiotropium to provide a delivered dose selected from about 3 µg, 6 µg, 9 µg, 18 µg, and 36 µg per actuation of the MDI. In order to achieve delivered doses as described herein, where compositions described herein include tiotropium as the active agent, in specific embodiments, the amount of tiotropium included in the compositions may be selected from, for example, between about 0.01 mg/ml and about 0.5 mg/ml.

[0084] In certain embodiments, the compositions described herein include a LABA active agent. In such embodiments, a LABA active agent can be selected from, for example, bambuterol, clenbuterol, formoterol, salmeterol, carmoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantyl-derived β_2 agonists, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof. In certain such embodiments, formoterol is selected as the LABA active agent. Formoterol can be used to treat inflammatory or obstructive pulmonary diseases and disorders such as, for example, those described herein. Formoterol has the chemical name (\pm) -2-hydroxy-5-[(1RS)-1-hydroxy-2-[(1RS)-2-(4-methoxyphenyl)-1-methylethyl]-amino]ethyl] formanilide, and is commonly used in pharmaceutical compositions as the racemic fumarate dihydrate salt. Where appropriate, formoterol may be used in the form of salts (e.g. alkali metal or amine salts or as acid addition salts) or as esters or as solvates (hydrates). Additionally, the formoterol may be in any crystalline form or isomeric form or mixture of isomeric forms, for example a pure enantiomer, a mixture of enantiomers, a racemate or a mixture thereof. In this regard, the form of formoterol may be selected to optimize the activity and/or stability of formoterol and/or to minimize the solubility of formoterol in the suspension medium. Pharmaceutically acceptable salts of formoterol include,

for example, salts of inorganic acids such as hydrochloric, hydrobromic, sulfuric and phosphoric acids, and organic acids such as fumaric, maleic, acetic, lactic, citric, tartaric, ascorbic, succinic, glutaric, gluconic, tricarballylic, oleic, benzoic, p-methoxybenzoic, salicylic, o- and p-hydroxybenzoic, p-chlorobenzoic, methanesulfonic, p-toluenesulfonic and 3-hydroxy-2-naphthalene carboxylic acids. Hydrates of formoterol are described, for example, in U.S. Pat. No. 3,994,974 and U.S. Pat. No. 5,684,199. Specific crystalline forms of formoterol and other β_2 adrenergic receptor agonists are described, for example, in WO95/05805, and specific isomers of formoterol are described in U.S. Patent No. 6,040,344.

[0085] In specific embodiments, the formoterol material utilized to form the formoterol particles is formoterol fumarate, and in one such embodiment, the formoterol fumarate is present in the dihydrate form. Where the compositions described herein include formoterol, in certain embodiments, the compositions described herein may include formoterol at a concentration that achieves a targeted delivered dose selected from between about 1 μ g and about 30 μ g, about 1 μ g and about 10 μ g, about 2 μ g and 5 μ g, about 2 μ g and about 10 μ g, about 5 μ g and about 10 μ g, and 3 μ g and about 30 μ g per actuation of an MDI. In other embodiments, the compositions described herein may include formoterol in an amount sufficient to provide a targeted delivered dose selected from up to about 30 μ g, up to about 10 μ g, up to about 5 μ g, up to about 2.5 μ g, up to about 2 μ g, or up to about 1.5 μ g per actuation. In order to achieve targeted delivered doses as described herein, where compositions described herein include formoterol as the active agent, in specific embodiments, the amount of formoterol included in the compositions may be selected from, for example, between about 0.01 mg/ml and about 1 mg/ml, between about 0.01 mg/ml and about 0.5 mg/ml, and between about 0.03 mg/ml and about 0.4 mg/ml.

[0086] Where the pharmaceutical co-suspension compositions described herein include a LABA active agent, in certain embodiments, the active agent may be salmeterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof. Salmeterol can be used to treat inflammatory or obstructive pulmonary diseases and disorders such as, for example, those described herein. Salmeterol, pharmaceutically acceptable salts of salmeterol, and methods for

producing the same are described, for example, in U.S. Patent No. 4,992,474, U.S. Patent No. 5,126,375, and U.S. patent 5,225,445.

[0087] Where salmeterol is included as a LABA active agent, in certain embodiments, the compositions described herein may include salmeterol at a concentration that achieves a delivered dose selected from between about 2 μ g and about 120 μ g, about 4 μ g and about 40 μ g, about 8 μ g and 20 μ g, about 8 μ g and about 40 μ g, about 20 μ g and about 40 μ g, and 12 μ g and about 120 μ g per actuation of an MDI. In other embodiments, the compositions described herein may include salmeterol in an amount sufficient to provide a delivered dose selected from up to about 120 μ g, up to about 40 μ g, up to about 20 μ g, up to about 10 μ g, up to about 8 μ g, or up to about 6 μ g per actuation of an MDI. In order to achieve targeted delivered doses as described herein, where compositions described herein include salmeterol as the active agent, in specific embodiments, the amount of salmeterol included in the compositions may be selected from, for example, between about 0.04 mg/ml and about 4 mg/ml, between about 0.04 mg/ml and about 2.0 mg/ml, and between about 0.12 mg/ml and about 0.8 mg/ml. For example, the compositions described herein may include sufficient salmeterol to provide a target delivered dose selected from between about 4 μ g and about 120 μ g, about 20 μ g and about 100 μ g, and between about 40 μ g and about 120 μ g per actuation of an MDI. In still other embodiments, the compositions described herein may include sufficient salmeterol to provide a targeted delivered dose selected from up to about 100 μ g, up to about 40 μ g, or up to about 15 μ g per actuation of an MDI.

[0088] Though the active agent material included in the compositions described herein may be amorphous or substantially amorphous, in specific embodiments, the active agent material used as or in the formation of the active agent particles included in the compositions described herein is substantially or entirely crystalline. Active agent material that is substantially or entirely crystalline may be selected to improve the chemical stability of the LAMA or LABA active agent when formulated in the compositions described herein. Therefore, in specific embodiments, the active agent material included in the compositions described herein is a micronized, crystalline LAMA material. In one such embodiment, the active agent particles are formed solely of micronized, crystalline LAMA material, such as a micronized crystalline material selected from glycopyrrolate, dexipirronium, tiotropium, trospium,

aclidinium, darotropium, and any pharmaceutically acceptable salts, esters or solvates thereof. In other specific embodiments, the active agent material included in the compositions described herein is a micronized, crystalline LABA material. In one such embodiment, the active agent particles are formed solely of micronized, crystalline LABA material, such as a micronized crystalline material selected from bambuterol, clenbuterol, formoterol, salmeterol, caramoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantly-derived β_2 agonists, and any pharmaceutically acceptable salts, esters or solvates thereof.

[0089] Any suitable process may be employed to achieve micronized active agent material as or in the formulation of the active agent particles included in the compositions described herein. A variety of processes may be used to create active agent particles suitable for use in the co-suspension formulations described herein, including, but not limited to micronization by milling or grinding processes, crystallization or recrystallization processes, and processes using precipitation from supercritical or near-supercritical solvents, spray drying, spray freeze drying, or lyophilization. Patent references teaching suitable methods for obtaining micronized active agent particles are described, for example, in U.S. Patent No. 6,063,138, U.S. Patent No. 5,858,410, U.S. Patent No. 5,851,453, U.S. Patent No. 5,833,891, U.S. Patent No. 5, 707,634, and International Patent Publication No. WO 2007/009164. Where the active agent particles include active agent material formulated with one or more excipient or adjuvant, micronized active agent particles can be formed using one or more of the preceding processes and such processes can be utilized to achieve active agent particles having a desired size distribution and particle configuration.

(iii) Suspending Particles

[0090] The suspending particles included in the co-suspension compositions described herein work to facilitate stabilization and delivery of the active agent included in the compositions. Though various forms of suspending particles may be used, the suspending particles are typically formed from pharmacologically inert material that is acceptable for inhalation and is substantially insoluble in the propellant selected. Generally, the majority of suspending particles are sized within a respirable range. In particular embodiments, therefore, the MMAD of the

suspending particles will not exceed about 10 μm but is not lower than about 500 nm. In an alternative embodiment, the MMAD of the suspending particles is between about 5 μm and about 750 nm. In yet another embodiment, the MMAD of the suspending particles is between about 1 μm and about 3 μm . When used in an embodiment for nasal delivery from an MDI, the MMAD of the suspending particles is between 10 μm and 50 μm .

[0091] In order to achieve respirable suspending particles within the MMAD ranges described, the suspending particles will typically exhibit a volume median optical diameter between about 0.2 μm and about 50 μm . In one embodiment, the suspending particles exhibit a volume median optical diameter that does not exceed about 25 μm . In another embodiment, the suspending particles exhibit a volume median optical diameter selected from between about 0.5 μm and about 15 μm , between about 1.5 μm and about 10 μm , and between about 2 μm and about 5 μm .

[0092] The concentration of suspending particles included in a composition according to the present description can be adjusted, depending on, for example, the amount of active agent particles and suspension medium used. In one embodiment, the suspending particles are included in the suspension medium at a concentration selected from about 1 mg/ml to about 15 mg/ml, about 3 mg/ml to about 10 mg/ml, 5 mg/ml to about 8 mg/ml, and about 6 mg/ml. In another embodiment, the suspending particles are included in the suspension medium at a concentration of up to about 30 mg/ml. In yet another embodiment, the suspending particles are included in the suspension medium at a concentration of up to about 25 mg/ml.

[0093] The relative amount of suspending particles to active agent particles is selected to achieve a co-suspension as contemplated herein. A co-suspension composition may be achieved where the amount of suspending particles, as measured by mass, exceeds that of the active agent particles. For example, in specific embodiments, the ratio of the total mass of the suspending particles to the total mass of active agent particles may be between about 3:1 and about 15:1, or alternatively from about 2:1 and 8:1. Alternatively, the ratio of the total mass of the suspending particles to the total mass of active agent particles may be above about 1, such as up to about 1.5, up to about 5, up to about 10, up to about 15, up to about 17, up to about 20, up to about 30, up to about 40, up to about 50, up to about 60, up to about 75, up to about 100, up to about 150, and up to about 200, depending on

the nature of the suspending particles and active agent particles used. In further embodiments, the ratio of the total mass of the suspending particles to the total mass of the active agent particles may be selected from between about 10 and about 200, between about 60 and about 200, between about 15 and about 60, between about 15 and about 170, between about 15 and about 60, about 16, about 60, and about 170.

[0094] In other embodiments, the amount of suspending particles, as measured by mass, is less than that of the active agent particles. For example, in particular embodiments, the mass of the suspending particles may be as low as 20% of the total mass of the active agent particles. However, in some embodiments, the total mass of the suspending particles may also approximate or equal the total mass of the active agent particles.

[0095] Suspending particles suitable for use in the compositions described herein may be formed of one or more pharmaceutically acceptable materials or excipients that are suitable for inhaled delivery and do not substantially degrade or dissolve in the suspension medium. In one embodiment, perforated microstructures, as defined herein, may be used as the suspending particles. Exemplary excipients that may be used in the formulation of suspending particles described herein include but are not limited to (a) carbohydrates, e.g., monosaccharides such as fructose, galactose, glucose, D-mannose, sorbose, and the like; disaccharides, such as sucrose, lactose, trehalose, cellobiose, and the like; cyclodextrins, such as 2-hydroxypropyl- β -cyclodextrin; and polysaccharides, such as raffinose, maltodextrins, dextrans, starches, chitin, chitosan, inulin, and the like; (b) amino acids, such as alanine, glycine, arginine, aspartic acid, glutamic acid, cysteine, lysine, leucine, isoleucine, valine, and the like; (c) metal and organic salts prepared from organic acids and bases, such as sodium citrate, sodium ascorbate, magnesium gluconate, sodium gluconate, tromethamin hydrochloride, and the like; (d) peptides and proteins such as aspartame, trileucine, human serum albumin, collagen, gelatin, and the like; (e) alditols, such as mannitol, xylitol, and the like; (f) synthetic or natural polymers or combinations thereof, such as polylactides, polylactide-glycolides, cyclodextrins, polyacrylates, methylcellulose, carboxymethylcellulose, polyvinyl alcohols, polyanhydrides, polylactams, polyvinyl pyrrolidones, hyaluronic acid, polyethylene glycols; and (g) surfactants including fluorinated and nonfluorinated compounds such

as saturated and unsaturated lipids, nonionic detergents, nonionic block copolymers, ionic surfactants and combinations thereof. In particular embodiments, suspending particles may include a calcium salt, such as calcium chloride, as described, for example, in U.S. Patent No. 7,442,388.

[0096] Additionally, phospholipids from both natural and synthetic sources may be used in preparing suspending particles suitable for use in the compositions described herein. In particular embodiments, the phospholipid chosen will have a gel to liquid crystal phase transition of greater than about 40°C. Exemplary phospholipids are relatively long chain (i.e., C₁₆-C₂₂) saturated lipids and may comprise saturated phospholipids, such as saturated phosphatidylcholines having acyl chain lengths of 16 C or 18 C (palmitoyl and stearoyl). Exemplary phospholipids include phosphoglycerides such as dipalmitoylphosphatidylcholine, distearylphosphatidylcholine, diarachidoylphosphatidylcholine, dibehenoylphosphatidylcholine, diphosphatidyl glycerol, short-chain phosphatidylcholines, long-chain saturated phosphatidylethanolamines, long-chain saturated phosphatidylserines, long-chain saturated phosphatidylglycerols, and long-chain saturated phosphatidylinositols. Additional excipients are disclosed in International Patent Publication No. WO 96/32149 and U.S. Patent Nos. 6,358,530, 6,372,258 and 6,518,239.

[0097] In particular embodiments, the suspending particles may be formed using one or more lipids, phospholipids or saccharides, as described herein. In some embodiments, suspending particles include one or more surfactants. The use of suspending particles formed of or incorporating one or more surfactants may promote absorption of the selected active agent, thereby increasing bioavailability. The suspending particles described herein, such as, for example, suspending particles formed using one or more lipids, can be formed to exhibit a desired surface rugosity (roughness), which can further reduce inter-particle interactions and improve aerosolization by reducing the surface area available for particle-particle interaction. In further embodiments, if suitable, a lipid that is naturally occurring in the lung could be used in forming the suspending particles, as such suspending particles that have the potential to reduce opsonization (and thereby reducing phagocytosis by alveolar macrophages), thus providing a longer-lived controlled release particle in the lung.

[0098] In another aspect, the suspending particles utilized in the compositions described herein may be selected to increase storage stability of the selected active agent, similar to that disclosed in International Patent Publication No WO 2005/000267. For example, in one embodiment, the suspending particles may include pharmaceutically acceptable glass stabilization excipients having a Tg of at least 55 °C, at least 75 °C, or at least 100 °C. Glass formers suitable for use in compositions described herein include, but are not limited to, one or more of trileucine, sodium citrate, sodium phosphate, ascorbic acid, inulin, cyclodextrin, polyvinyl pyrrolidone, mannitol, sucrose, trehalose, lactose, and, proline. Examples of additional glass-forming excipients are disclosed in U. S. Patent Nos. RE 37,872, 5,928,469, 6,258,341.

[0099] The suspending particles may be designed, sized and shaped as desired to provide desirable stability and active agent delivery characteristics. In one exemplary embodiment, the suspending particles comprise perforated microstructures as described herein. Where perforated microstructures are used as suspending particles in the compositions described herein, they may be formed using one or more excipients as described herein. For example, in particular embodiments, perforated microstructures may include at least one of the following: lipids, phospholipids, nonionic detergents, nonionic block copolymers, ionic surfactants, biocompatible fluorinated surfactants and combinations thereof, particularly those approved for pulmonary use. Specific surfactants that may be used in the preparation of perforated microstructures include poloxamer 188, poloxamer 407 and poloxamer 338. Other specific surfactants include oleic acid or its alkali salts. In one embodiment, the perforated microstructures include greater than about 10% w/w surfactant.

[0100] In some embodiments, suspending particles may be prepared by forming an oil-in-water emulsion, using a fluorocarbon oil (e.g., perfluoroctyl bromide, perfluorodecalin) which may be emulsified using a surfactant such as a long chain saturated phospholipid. The resulting perfluorocarbon in water emulsion may be then processed using a high pressure homogenizer to reduce the oil droplet size. The perfluorocarbon emulsion may be fed into a spray dryer, optionally with an active agent solution, if it is desirable to include active agent within the matrix of the perforated microstructures. As is well known, spray drying is a one-step process that

converts a liquid feed to a dried particulate form. Spray drying has been used to provide powdered pharmaceutical material for various administrative routes, including inhalation. Operating conditions of the spray dryer (such as inlet and outlet temperature, feed rate, atomization pressure, flow rate of the drying air and nozzle configuration) can be adjusted to produce the desired particle size producing a yield of the resulting dry microstructures. Such methods of producing exemplary perforated microstructures are disclosed in U.S. Patent No. 6,309,623 to Weers et al.

[0101] Perforated microstructures as described herein may also be formed through lyophilization and subsequent milling or micronization. Lyophilization is a freeze-drying process in which water is sublimed from the composition after it is frozen. This process allows drying without elevated temperatures. In yet further embodiments, the suspending particles may be produced using a spray freeze drying process, such as is disclosed in U.S. Patent 5,727,333.

[0102] Furthermore, suspending particles as described herein may include bulking agents, such as polymeric particles. Polymeric polymers may be formed from biocompatible and/or biodegradable polymers, copolymers or blends. In one embodiment, polymers capable of forming aerodynamically light particles may be used, such as functionalized polyester graft copolymers and biodegradable polyanhydrides. For example, bulk eroding polymers based on polyesters including poly(hydroxy acids) can be used. Polyglycolic acid (PGA), polyactic acid (PLA) or copolymers thereof may be used to form suspending particles. The polyester may include a charged or functionalizable group, such as an amino acid. For example, suspending particles may be formed of poly(D,L-lactic acid) and/or poly(D,L-lactic-co-glycolic acid) (PLGA), which incorporate a surfactant such as DPPC.

[0103] Other potential polymer candidates for use in suspending particles may include polyamides, polycarbonates, polyalkylenes such as polyethylene, polypropylene, poly(ethylene glycol), poly(ethylene oxide), poly(ethylene terephthalate), poly vinyl compounds such as polyvinyl alcohols, polyvinyl ethers, and polyvinyl esters, polymers of acrylic and methacrylic acids, celluloses and other polysaccharides, and peptides or proteins, or copolymers or blends thereof. Polymers may be selected with or modified to have the appropriate stability and degradation rates in vivo for different controlled drug delivery applications.

[0104] The compositions described herein may include two or more species of suspending particles. Even further, compositions according to the present description can include suspending particles that include glycopyrrolate incorporated into the suspending particles. Where active agent is incorporated into suspending particles, the suspending particles will be of a respirable size and can be formulated and produced using, for example, the methods and materials described herein.

[0105] Compositions formulated according to the present teachings can inhibit degradation of active agent included therein. For example, in specific embodiments, the compositions described herein inhibit one or more of flocculation, aggregation and the solution mediated transformation of active agent material included in the compositions. The pharmaceutical compositions described herein are suited for respiratory delivery via and MDI in a manner that achieves desirable delivered dose uniformity (“DDU”) of LABA and LAMA active agents, including potent and highly potent LABA and LAMA agents throughout emptying of an MDI canister. As is described in detail in the Examples included herein, even when delivering very low doses of LAMA or LABA active agents, compositions described herein can achieve a DDU for the active agent of \pm 30%, or better throughout emptying of an MDI canister. In one such embodiment, compositions described herein achieve a DDU for the active agent of \pm 25%, or better throughout emptying of an MDI canister. In yet another such embodiment, compositions described herein achieve a DDU for the active agent of \pm 20%, or better throughout emptying of an MDI canister.

[0106] Pharmaceutical compositions described herein also serve to substantially preserve FPF and FPD performance throughout emptying of an MDI canister, even after being subjected to accelerated degradation conditions. For instance, compositions according to the present description maintain as much as 80%, 90%, 95%, or more, of the original FPF and FPD performance throughout emptying of an MDI canister, even after being subjected to accelerated degradation conditions. Compositions described herein provide the added benefit of achieving such performance while being formulated using non-CFC propellants. In specific embodiments, the compositions described herein achieve desired one or all of a targeted DDU, FPF and FPD performance while being formulated with suspension medium including only one or more non-CFC propellants and without the need to modify the characteristics of the non-CFC propellant, such as by the addition of, for

example, one or more cosolvent, antisolvent, solubilizing agent, adjuvant or other propellant modifying material.

[0107] In one embodiment, a co-suspension composition as described herein includes: a suspension medium comprising a pharmaceutically acceptable HFA propellant; a plurality of active agent particles comprising glycopyrrolate, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, suspended in the suspension medium at a concentration sufficient to provide a delivered dose of glycopyrrolate of between about 20 μg and about 150 μg per actuation of the metered dose inhaler; and a plurality of respirable suspending particles comprising perforated microstructures as described herein exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , wherein perforated microstructures associate with the plurality of active agent particles to form a co-suspension. In one such embodiment, the glycopyrrolate active agent particles are formed of crystalline glycopyrrolate material. In another such embodiment, the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In yet another such embodiment, the glycopyrrolate active agent particles are formed of crystalline glycopyrrolate material and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In still another such embodiment, the glycopyrrolate active agent particles are formed of crystalline glycopyrrolate material, at least 90% of the glycopyrrolate active agent particles by volume exhibit an optical diameter of less than 7 μm , and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

[0108] In another embodiment, a co-suspension composition as described herein includes: a suspension medium comprising a pharmaceutically acceptable HFA propellant; a plurality of active agent particles comprising tiotropium, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, suspended in the suspension medium at a concentration sufficient to provide a delivered dose of glycopyrrolate of between about 5 μg and about 40 μg per actuation of the metered dose inhaler; and a plurality of respirable suspending particles comprising perforated

microstructures as described herein exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , wherein perforated microstructures associate with the plurality of active agent particles to form a co-suspension. In one such embodiment, the tiotropium active agent particles are formed of crystalline tiotropium material. In another such embodiment, the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In yet another such embodiment, the tiotropium active agent particles are formed of crystalline tiotropium material and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In still another such embodiment, the tiotropium active agent particles are formed of crystalline tiotropium material, at least 90% of the tiotropium active agent particles by volume exhibit an optical diameter of less than 7 μm , and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

[0109] In another embodiment, a co-suspension composition as described herein includes: a suspension medium comprising a pharmaceutically acceptable HFA propellant; a plurality of active agent particles comprising formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, suspended in the suspension medium at a concentration sufficient to provide a delivered dose of formoterol of between about 0.5 μg and about 10 μg per actuation of the metered dose inhaler; and a plurality of respirable suspending particles comprising perforated microstructures as described herein exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , wherein perforated microstructures associate with the plurality of active agent particles to form a co-suspension. In one such embodiment, the formoterol active agent particles are formed of crystalline formoterol material. In another such embodiment, the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In yet another such embodiment, the formoterol active agent particles are formed of crystalline formoterol material and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about

15:1 and between about 2:1 and 8:1. In still another such embodiment, the formoterol active agent particles are formed of crystalline formoterol material, at least 90% of the formoterol active agent particles by volume exhibit an optical diameter of less than 7 μm , and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

[0110] In one embodiment, a co-suspension composition as described herein includes: a suspension medium comprising a pharmaceutically acceptable HFA propellant; a plurality of active agent particles comprising formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, suspended in the suspension medium at a concentration sufficient to provide a delivered dose of formoterol of between about 2 μg and about 10 μg per actuation of the metered dose inhaler; and a plurality of respirable suspending particles comprising perforated microstructures as described herein exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , wherein perforated microstructures associate with the plurality of active agent particles to form a co-suspension. In one such embodiment, the formoterol active agent particles are formed of crystalline formoterol material. In another such embodiment, the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In yet another such embodiment, the formoterol active agent particles are formed of crystalline formoterol material and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In still another such embodiment, the formoterol active agent particles are formed of crystalline formoterol material, at least 90% of the formoterol active agent particles by volume exhibit an optical diameter of less than 7 μm , and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

[0111] In another embodiment, a co-suspension composition as described herein includes: a suspension medium comprising a pharmaceutically acceptable HFA propellant; a plurality of active agent particles comprising salmeterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, suspended in

the suspension medium at a concentration sufficient to provide a delivered dose of salmeterol of between about 8 μg and about 40 μg per actuation of the metered dose inhaler; and a plurality of respirable suspending particles comprising perforated microstructures as described herein exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , wherein perforated microstructures associate with the plurality of active agent particles to form a co-suspension. In one such embodiment, the salmeterol active agent particles are formed of crystalline salmeterol material. In another such embodiment, the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In yet another such embodiment, the salmeterol active agent particles are formed of crystalline salmeterol material and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1. In still another such embodiment, the salmeterol active agent particles are formed of crystalline salmeterol material, at least 90% of the salmeterol active agent particles by volume exhibit an optical diameter of less than 7 μm , and the ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

III. Metered Dose Inhaler Systems

[0112] As described in relation to the methods provided herein, the compositions disclosed herein may be used in an MDI system. MDIs are configured to deliver a specific amount of a medicament in aerosol form. In one embodiment, an MDI system includes a pressurized, liquid phase formulation-filled canister disposed in an actuator formed with a mouthpiece. The MDI system may include the formulations described herein, which include a suspension medium, glycopyrrolate and at least one species of suspending particles. The canister used in the MDI be any of any suitable configuration, and in one exemplary embodiment, the canister may have a volume ranging from about 5 mL to about 25 mL, such as, for example a canister having a 19 mL volume. After shaking the device, the mouthpiece is inserted into a patient's mouth between the lips and teeth. The patient typically exhales deeply to empty the lungs and then takes a slow deep breath while actuating the cartridge.

[0113] Inside an exemplary cartridge is a metering valve including a metering chamber capable of holding a defined volume of the formulation (e.g., 63 μ l or any other suitable volume available in commercially available metering valves), which is released into an expansion chamber at the distal end of the valve stem when actuated. The actuator retains the canister and may also include a port with an actuator nozzle for receiving the valve stem of the metering valve. When actuated, the specified volume of formulation travels to the expansion chamber, out the actuator nozzle and into a high-velocity spray that is drawn into the lungs of a patient.

IV. Methods

[0114] Methods for formulating pharmaceutical compositions for respiratory delivery of LAMA and LAMA active agents are provided herein. In particular embodiments, such methods involve the steps of providing a suspension medium, active agent particles selected from active agent particles comprising a LAMA and active agent particles comprising a LABA, and one or more species of suspending particles, as described herein, and combining such constituents to form a formulation wherein the active agent particles associate with the suspending particles and co-locate with the suspending particles within the suspension medium such that a co-suspension is formed. In one such embodiment, the association of the glycopyrrolate particles and the suspending particles is such that they do not separate due to their different buoyancies in a propellant. As will be appreciated, the method may include providing two or more species of suspending particles in combination with active agent particles. In another embodiment, the method may include providing two or more species of active agent particles and combining the two or more species of active agent particles with one or more species of suspending particles in a manner that results in a co-suspension. In certain embodiments, the active agent particles consist essentially of a LAMA or LABA active agent as described herein.

[0115] In specific embodiments of methods for providing a stabilized composition of a LAMA or LABA active agent for pulmonary delivery, the present disclosure provides methods for inhibiting the solution mediated transformation of the LAMA or LABA active agent in a pharmaceutical composition for pulmonary delivery. In one

embodiment, a suspension medium as described herein, such as a suspension medium formed by an HFA propellant, is obtained. Suspending particles are also obtained or prepared as described herein. Active agent particles are also obtained, and the suspension medium, suspending particles and active agent particles are combined to form a co-suspension wherein the active agent particles associate with suspending particles and co-locate with the suspending particles within the continuous phase formed by the suspension medium. When compared to active agent particles contained in the same suspension medium in the absence of suspending particles, co-suspensions according to the present description have been found to exhibit a higher tolerance to solution mediated phase transformation that leads to irreversible crystal aggregation, and thus may lead to improved stability and dosing uniformity.

[0116] In further embodiments, methods for forming stabilized compositions of LAMA and LABA active agents for pulmonary delivery include for preserving the FPF and/or FPD of the composition throughout emptying of an MDI canister. In specific embodiments of methods for preserving the FPF and/or FPD provided by a pharmaceutical composition for pulmonary delivery, a respirable co-suspension as described herein is provided which is capable of maintaining the FPD and/or the FPF to within \pm 20%, \pm 10%, or even \pm 5% the initial FPD and/or FPF, respectively, throughout emptying of an MDI canister. Such performance can be achieved even after the co-suspension is subjected to accelerated degradation conditions. In one embodiment, a suspension medium as described herein, such as a suspension medium formed by an HFA propellant, is obtained. Suspending particles are also obtained or prepared as described herein. Active agent particles are also obtained, and the suspension medium, suspending particles and active agent particles are combined to form a co-suspension wherein the glycopyrrolate particles associate with suspending particles and co-locate with the suspending particles within the suspension medium. Even after exposure of such composition to one or more temperature cycling events, the co-suspension maintains an FPD or FPF within \pm 20%, \pm 10%, or even \pm 5% of the respective values measured prior to exposure of the composition to the one or more temperature cycling events.

[0117] Methods for preparing an MDI for pulmonary delivery of LAMA or LABA active agent are disclosed. The method of preparing the MDI may include loading a

canister, as described herein, with active agent particles and suspending particles. An actuator valve can be attached to an end of the canister and the canister sealed. The actuator valve may be adapted for dispensing a metered amount of the glycopyrrolate pharmaceutical formulation per actuation. The canister can be charged with a pharmaceutically acceptable suspension medium, such as a propellant as described herein. Whereupon the active agent particles and suspending particles yield a stable co-suspension in the suspension medium.

[0118] In methods involving pulmonary delivery of a LAMA or LABA active agent using compositions described herein, the compositions may be delivered by an MDI. Therefore, in particular embodiments of such methods, an MDI loaded with a composition described herein is obtained, and a LAMA or LABA active agent is administered to a patient through pulmonary delivery through actuation of the MDI. For example, in one embodiment, after shaking the MDI device, the mouthpiece is inserted into a patient's mouth between the lips and teeth. The patient typically exhales deeply to empty the lungs and then takes a slow deep breath while actuating the cartridge of the MDI. When actuated, the specified volume of formulation travels to the expansion chamber, out the actuator nozzle and into a high-velocity spray that is drawn into the lungs of a patient. In one embodiment the dose of active agent delivered throughout emptying of an MDI canister is not more than 30% greater than the mean delivered dose and is not less than 30% less than the mean delivered dose. Therefore, methods of achieving a desired DDU of glycopyrrolate delivered from an MDI are also provided. In such embodiments, the method may include achieving a DDU for glycopyrrolate delivered from an MDI selected from, for example, a DDU of \pm 30%, or better, a DDU of \pm 25%, or better, and a DDU of \pm 20%, or better.

[0119] Methods for treating patients suffering from an inflammatory or obstructive pulmonary disease or condition are provided herein. In specific embodiments, such methods include pulmonary delivery of a pharmaceutical composition described herein, and in certain such embodiments, pulmonary administration of the pharmaceutical composition is accomplished by delivering the composition using an MDI. The disease or condition to be treated can be selected from any inflammatory or obstructive pulmonary disease or condition that responds to the administration of a LAMA or LABA agent. In particular embodiments, the pharmaceutical

compositions described herein may be used in treating a disease or disorder selected from asthma, COPD, exacerbation of airways hyper reactivity consequent to other drug therapy, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, pulmonary hypertension, pulmonary vasoconstriction, emphysema, and any other respiratory disease, condition, trait, genotype or phenotype that can respond to the administration of a LAMA or LABA, alone or in combination with other therapies. In certain embodiments, the pharmaceutical compositions described herein may be used in treating pulmonary inflammation and obstruction associated with cystic fibrosis.

[0120] Additionally, pharmaceutical compositions according to the present description delivered from an MDI provide desirable pharmacodynamic (PD) performance. In particular embodiments, pulmonary delivery of the pharmaceutical compositions described herein results in rapid, significant improvement in the lung capacity, which can be characterized by an improvement in the patient's forced expiratory volume in one second (FEV₁). For example, in particular embodiments, methods for achieving a clinically relevant increase in FEV₁ are provided, wherein such methods include providing a co-suspension composition comprising a LABA or LAMA active agent as described herein and administering such composition to a patient experiencing pulmonary inflammation or obstruction via an MDI. For purposes of the present disclosure, a clinically relevant increase in FEV₁ is any increase of 100 ml or greater, and in certain embodiments of the methods described herein, administration of compositions according to the present description to patient results in a clinically significant increase in FEV₁ within 1 hour or less. In other such embodiments, methods for administering a composition as described herein to a patient via an MDI result in a clinically significant increase in FEV₁ within 0.5 hours or less. The compositions provided and delivered in such embodiments may include a composition including a LAMA active agent or a composition including a LABA active agent as described herein.

[0121] In further embodiments, methods are provided for achieving an increase in FEV₁ greater than 100 ml. For example, in certain embodiments, the methods described herein include methods for achieving an FEV₁ of 150 ml or greater within a period of time selected from 0.5 hours or less, 1 hour or less, and 1.5 hours or less.

In other embodiments, the methods described herein include methods for achieving an FEV₁ of 200 ml or greater within a period of time selected from 0.5 hours or less, 1 hour or less, and 1.5 hours or less, and 2 hours or less. In certain such embodiments, a composition comprising a LABA or LAMA active agent as described herein is provided and administered to a patient experiencing pulmonary inflammation or obstruction via an MDI.

[0122] In still further embodiments, methods for achieving and maintaining a clinically significantly increase in FEV₁ are provided. In particular embodiments, upon administration of a single dose of a LABA or LAMA active agent formulated in a composition as described herein to a patient via an MDI, a clinically significant increase in FEV₁ is achieved in a period of time selected from 0.5 hours or less, 1 hour or less, and 1.5 hours or less, and the clinically significant increase in FEV₁ is maintained for up 12 hours or more. In certain such embodiments, the increase in FEV₁ may be selected from an increase of 150 ml or greater, 200 ml or greater and 250 ml or greater, and the increase in FEV₁ remains clinically significant for a time period selected from up to 4 hours, up to 6 hours, up to 8 hours, up to 10 hours, and up to 12 hours, or more. In certain such embodiments, a composition comprising a LABA or LAMA active agent as described herein is provided and administered a patient experiencing pulmonary inflammation or obstruction via an MDI.

[0123] Compositions, systems and methods described herein are not only suited to achieving desirable pharmacodynamic performance in short periods of time, but will achieve such results in a high percentage of patients. For example, methods are provided herein for achieving a 10% or greater increase in FEV₁ in 50% or more of patients experiencing pulmonary inflammation or obstruction. For example, in particular embodiments, methods for achieving a 10% or greater increase in FEV₁ in a patient include providing a co-suspension composition comprising a LABA or LAMA active agent as described herein and administering such composition via an MDI to a patient experiencing pulmonary inflammation or obstruction. In certain such embodiments, administration of the composition results in 10% or greater increase in FEV₁ within a period of time selected from 0.5 hours or less, 1 hour or less, 1.5 hours or less, and 2 hours in 50% or more of patients. In other such embodiments, administration of the composition results in 10% or greater increase in FEV₁ within a period of time selected from 0.5 hours or less, 1 hour or less, 1.5 hours or less, and

2 or less hours in 60% or more of patients. In still other such embodiments, administration of the composition results in 10% or greater increase in FEV₁ within a period of time selected from 0.5 hours or less, 1 hour or less, 1.5 hours or less, and 2 hours or less in 70% or more of patients. In yet other such embodiments, administration of the composition results in 10% or greater increase in FEV₁ within a period of time selected from 0.5 hours or less, 1 hour or less, 1.5 hours or less, and 2 or less hours in 80% or more of patients

[0124] In specific embodiments, the methods described herein facilitate treatment of patients experiencing pulmonary inflammation or obstruction, wherein such methods include providing a co-suspension composition comprising a LABA or LAMA active agent as described herein and administering such composition to a patient experiencing pulmonary inflammation or obstruction via an MDI and result in a high proportion of such patients experiencing either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml. In certain such embodiments, administration of the composition results in either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1 hour or less, 1.5 hours or less, 2 hours or less, and 2.5 hours or less in 50% or more of patients. In other such embodiments, administration of the composition results in an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1 hour or less, 1.5 hours or less, 2 hours or less, and 2.5 hours or less in 60% or more of patients. In still other such embodiments, administration of the composition results in either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1.5 hours or less, 2 hours or less, 2.5 hours or less, and 3 hours or less in 70% or more of patients. In yet other such embodiments, administration of the composition results in either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1.5 hours or less, 2 hours or less, 2.5 hours or less, and 3 hours or less in 80% or more of patients.

[0125] In some embodiments, pharmaceutical compositions according to the present description delivered from an MDI provide improvement in the lung capacity, which can be characterized by an improvement inspiratory capacity (IC), which is defined as the maximal volume of gas that can be taken into the lungs in a full inhalation following a normal expiration. For example, in particular embodiments, methods for achieving a clinically relevant increase in IC are provided, wherein such methods include providing a co-suspension composition comprising a LABA or LAMA active agent as described herein and administering such composition to a patient experiencing pulmonary inflammation or obstruction via an MDI. For purposes of the present disclosure, a clinically relevant increase in IC is any increase of 70 ml or greater, and in certain embodiments of the methods described herein, administration of compositions according to the present description to patient results in a clinically significant increase in IC within 2 hours or less. In other such embodiments, methods for administering a composition as described herein to a patient via an MDI result in a clinically significant increase in IC within 1 hour or less. In other such embodiments, administration of compositions according to the present description to patient results in an increase in IC of 100 ml or greater within a period of time selected from 1 hour or less and 2 hours or less. In still other such embodiments, administration of compositions according to the present description to patient results in an increase in IC of 150 ml or greater within a period of time selected from 1 hour or less and 2 hours or less. In even further such embodiments, administration of compositions according to the present description to patient results in an increase in IC of 300 ml or greater within a period of time selected from 1 hour or less and 2 hours or less. The compositions provided and delivered in such embodiments may include a composition including a LAMA active agent or a composition including a LABA active agent as described herein.

[0126] In particular embodiments of the methods described herein, the compositions provided include a LAMA active agent. In such embodiments, the LAMA active agent can be selected from, for example, glycopyrrolate, dexipirronium, tiotropium, trospium, aclidinium, and darotropium, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof. In specific embodiments of the methods described herein, the composition is a co-suspension composition as described herein that includes glycopyrrolate or any pharmaceutically acceptable

salt, ester, isomer or solvate thereof. In other specific embodiments of the methods described herein, the composition is a co-suspension composition as described herein that includes tiotropium or any pharmaceutically acceptable salt, ester, isomer or solvate thereof. Where glycopyrrolate or tiotropium is selected as the active agent for use in the compositions produced or administered as part of the methods described herein, the amount of glycopyrrolate or tiotropium included in the composition may be selected from, for example, those amounts specifically disclosed with respect to the pharmaceutical compositions described herein.

[0127] In further embodiments of the methods described herein, the compositions provided include a LABA active agent. In such embodiments, the LABA active agent can be selected from, for example, bambuterol, clenbuterol, formoterol, salmeterol, carmoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantyl-derived β_2 agonists, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof. In specific embodiments of the methods described herein that includes formoterol or any pharmaceutically acceptable salt, ester, isomer or solvate thereof. In other specific embodiments of the methods described herein, the composition is a co-suspension composition as described herein that includes salmeterol or any pharmaceutically acceptable salt, ester, isomer or solvate thereof. Where formoterol or salmeterol is selected as the active agent for use in the compositions produced or administered as part of the methods described herein, the amount of formoterol or salmeterol included in the composition may be selected from, for example, those amounts specifically disclosed with respect to the pharmaceutical compositions described herein.

[0128] Compositions, methods and systems described herein provide desirable dose efficiency and dose response for LAMA or LABA active agents formulated for pulmonary delivery. For example, pulmonary delivery of glycopyrrolate for treatment of conditions such as COPD has been previously suggested or reported by Schroeckenstein et al., J. Allergy Clin. Immunol., 1988; 82(1): 115-119, Leckie et al., Exp. Opin. Invest. Drugs, 2000; 9(1): 3-23, Skorodin, Arch. Intern. Med., 1993; 153: 814-828, Walker et al., Chest, 1987; 91(1): 49-51, and International Patent Publication WO/1997/039758. These references report a minimum effective dose for glycopyrrolate of 200 μ g – 1,000 μ g. Such dosing requirements are in line with

human clinical results reported by Bannister et al. in U.S. Patent No. 7,229,607, wherein subjects were given a 480 μ g dose of glycopyrrolate. As is described in Example 6 provided herein, compositions of glycopyrrolate prepared according to the present description and delivered to human subjects via an MDI achieved quick onset of action and clinically relevant improvements in FEV₁ and IC in accordance with the methods detailed herein, even when delivering significantly smaller doses of glycopyrrolate (the largest single dose delivered in the study was 144 μ g).

[0129] Singh et al. [D Singh, P A Corris, and S D Snape. "NVA237, a once-daily inhaled antimuscarinic, provides 24-hour bronchodilator efficacy in patients with moderate to-severe COPD" Poster presented at the American Thoracic Society International Conference, San Diego, California, May 19-24, 2006] reported clinical work wherein glycopyrrolate was administered to human subjects via pulmonary delivery at doses of 20 μ g, 125 μ g, 250 μ g, and 400 μ g. Though such doses ranged below the 200 μ g threshold previously reported, as is also detailed in Example 6, compositions of glycopyrrolate formulated and delivered as described herein still achieved a relatively improved dose efficiency. For example, changes in FEV₁ AUC achieved by glycopyrrolate co-suspensions as described and evaluated in the clinical trial described in Example 6 are compared to those achieved by the compositions of Singh et al. in Figure 10. The 18 μ g glycopyrrolate dose from Example 6 provided significantly better bronchodilator response than the 20 μ g dose reported by Singh et al., and the 36 μ g and 144 μ g glycopyrrolate doses from Example 6 providing comparable bronchodilator response to the 125 μ g and 250 μ g doses, respectively, reported by Singh et al.

[0130] In particular embodiments, methods for achieving desired pharmacodynamic effects are provided, wherein the methods include administering a co-suspension composition as described herein to a patient via a metered dose inhaler, wherein the co-suspension includes glycopyrrolate active agent particles as described herein to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is administered to the patient. In one embodiment, a method for achieving a clinically significant increase in FEV₁ is provided, wherein the method includes administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is

administered to the patient. In one such embodiment, a delivered dose of no more than 100 μ g glycopyrrolate is administered to the patient, and in another embodiment, a delivered dose of no more than 80 μ g glycopyrrolate is administered to the patient. Even where doses of no more than 80 μ g, no more than 100 μ g glycopyrrolate, or no more than 150 μ g glycopyrrolate are administered to the patient, in particular embodiments, the clinically significant increase in FEV₁ is achieved in 1 hour or less. In some such embodiments, the clinically significant increase in FEV₁ is achieved in 0.5 hours or less.

[0131] In further embodiments, methods are provided for achieving an increase in FEV₁ greater than 100 ml, wherein the methods include administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is administered to the patient. For example, in certain embodiments, methods for achieving an FEV₁ of 150 ml or greater within a period of time selected from 0.5 hours or less, 1 hour or less, and 1.5 hours or less, are provided, wherein the methods include administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is administered to the patient. In other embodiments, the methods described herein include methods for achieving an FEV₁ of 200 ml or greater within a period of time selected from 0.5 hours or less, 1 hour or less, and 1.5 hours or less, and 2 hours or less, wherein the methods include administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is administered to the patient.

[0132] In still further embodiments, methods for achieving and maintaining a clinically significantly increase in FEV₁ are provided, wherein the methods include administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is administered to the patient. In certain such embodiments, upon administration of a single delivered dose of glycopyrrolate of no more than 150 μ g, a clinically significant increase in FEV₁ is achieved in a period of time selected from 0.5 hours or less, 1 hour or less, and 1.5 hours or less, and the

clinically significant increase in FEV₁ is maintained for up 12 hours or more. For example, in particular embodiments, the increase in FEV₁ may be selected from an increase of 150 ml or greater, 200 ml or greater and 250 ml or greater, and the increase in FEV₁ remains clinically significant for a time period selected from up to 4 hours, up to 6 hours, up to 8 hours, up to 10 hours, and up to 12 hours, or more.

[0133] Methods for achieving an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml are also provided, wherein the methods include administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 µg glycopyrrolate is administered to the patient. In certain such embodiments, administration of a delivered dose of no more than 150 µg glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1 hour or less, 1.5 hours or less, 2 hours or less, and 2.5 hours or less in 50% or more of patients. In other such embodiments, administration of a delivered dose of no more than 150 µg glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1 hour or less, 1.5 hours or less, 2 hours or less, and 2.5 hours or less in 60% or more of patients. In still other such embodiments, administration of a delivered dose of no more than 150 µg glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml within a period of time selected from 1.5 hours or less, 2 hours or less, 2.5 hours or less, and 3 hours or less in 70% or more of patients. In yet other such embodiments, administration of a delivered dose of no more than 150 µg glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in either an increase from baseline in FEV₁ of at least 200 ml or a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at

least 150 ml within a period of time selected from 1.5 hours or less, 2 hours or less, 2.5 hours or less, and 3 hours or less in 80% or more of patients.

[0134] Methods for achieving a clinically significant increase in IC are provided, wherein the methods include administering a co-suspension as described herein comprising glycopyrrolate active agent particles to a patient via a metered dose inhaler such that a delivered dose of no more than 150 μ g glycopyrrolate is administered to the patient. In certain such embodiments, administration of a delivered dose of no more than 150 μ g glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in a clinically significant increase in IC within 1 hour or less. In other such embodiments, administration of a delivered dose of no more than 150 μ g glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in an increase in IC of 100 ml or greater within a period of time selected from 1 hour or less and 2 hours or less. In still other such embodiments, administration of a delivered dose of no more than 150 μ g glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in an increase in IC of 150 ml or greater within a period of time selected from 1 hour or less and 2 hours or less. In even further such embodiments, administration of a delivered dose of no more than 150 μ g glycopyrrolate from a co-suspension as described herein via a metered dose inhaler results in an increase in IC of 300 ml or greater within a period of time selected from 1 hour or less and 2 hours or less.

[0135] The specific examples included herein are for illustrative purposes only and are not to be considered as limiting to this disclosure. Moreover, the compositions, systems and methods disclosed herein have been described in relation to certain embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled in the art that the invention is susceptible to additional embodiments and that certain of the details described herein may be varied without departing from the basic principles of the invention. Any active agents and reagents used in the following examples are either commercially available or can be prepared according to standard literature procedures by those skilled in the art of organic synthesis. The entire contents of all publications, patents, and patent applications referenced herein are hereby incorporated herein by reference.

Example 1

[0136] Active agent particles formed of glycopyrrolate (Pyrrolidinium, 3-((cyclopentylhydroxyphenylacetyl)oxy)-1,1-dimethyl-, bromide) were formed by micronizing glycopyrrolate using a jet mill. The particle size distribution of the micronized glycopyrrolate (GP) was determined by laser diffraction. 50% by volume of the micronized particles exhibited an optical diameter smaller than 2.1 μm , 90% by volume were smaller than 5 μm .

[0137] Suspending particles were manufactured as follows: 500 mL of a fluorocarbon-in water emulsion of PFOB (perfluorooctyl bromide) stabilized by a phospholipid was prepared. 18.7 g of the phospholipid, DSPC (1,2-Distearoyl-sn-Glycero-3-Phosphocholine), and 1.3 g of calcium chloride were homogenized in 400 mL of hot water (75 °C) using a high shear mixer. 100 mL of PFOB were added slowly during homogenization. The resulting coarse emulsion was then further homogenized using a high pressure homogenizer (Model C3, Avestin, Ottawa, CA) at pressures of up to 170 MPa for 5 passes.

[0138] The emulsion was spray dried in nitrogen using the following spray drying conditions: Inlet temperature 95 °C, outlet temperature 72 °C, emulsion feed rate 2.4 mL/min, total gas flow 525 L/min. The particle size distribution of the suspending particles was determined by laser diffraction. 50% by volume of the suspending particles were smaller than 2.9 μm , the Geometric Standard Deviation of the distribution was 1.8.

[0139] Metered dose inhalers were prepared by weighing the target masses of micronized GP particles and suspending particles into fluorinated ethylene polymer (FEP) coated aluminum canisters (Presspart, Blackburn, UK) with 19 mL volume. The target masses and the target delivered dose assuming 20% actuator deposition are given in Table 1 for five different configurations (configurations 1A through 1C representing different suspensions of GP particles and suspending particles; configuration 1D representing GP particles alone; configuration 1E representing suspending particles alone). The canisters were crimp sealed with 63 μl valves (# BK 357, Bespak, King's Lynn, UK) and filled with 12.4 g of HFA 134a (1,1,1,2-tetrafluoroethane) (Ineos Fluor, Lyndhurst, UK) by overpressure through the valve stem. After injecting the propellant, the canisters were sonicated for 15 seconds and agitated on a wrist action shaker for 30 minutes. The canisters were fitted with

polypropylene actuators with a 0.3 mm orifice (# BK 636, Bespak, King's Lynn, UK). Additional inhalers for visual observation of suspension quality were prepared using glass vials.

Table 1: Results for Glycopyrrolate Co-suspensions of Example 1

Configuration ID	GP (mg/can)	Suspending particles (mg/can)	Target delivered dose (µg)	Delivered Dose (µg)	FPF (%)	MMAD (µm)
1A	3.4	61	16.5	17.8	41.3	3.7
1B	4.1	61	20	19.4	42.0	3.9
1C	4.1	15	20	19.2	42.7	3.2
1D	4.1	0	20	11.1-15.3	27.0	3.3
1E	0	61	-	-	53.6 *	3.2

* Based on DSPC assay.

[0140] Aerosol performance was assessed shortly after manufacturing in accordance with USP <601> (United States Pharmacopeia Monograph 601). A Next Generation Impactor (NGI) operated at a flow rate of 30 L/min was used for determination of particle size distribution. Sample canisters were seated into an actuator with two waste actuations and two additional waste priming actuations. Five actuations were collected in the NGI with a USP throat attached. The valve, actuator, throat, NGI cups, stages, and filter were rinsed with volumetrically dispensed solvent. The sample solutions were assayed using a drug specific chromatographic method. The fine particle fraction was defined using the sum of stages 3 through filter. Delivered dose uniformity through use testing was performed using a Dose Uniformity Sampling Apparatus as described in USP <601>. Inhalers were seated and primed as described before. Two actuations were collected and assayed at beginning, middle and end of use.

[0141] Visual observation of the co-suspended configurations (1A, 1B, 1C) showed no sedimentation of drug crystals. The suspension flocculated slowly and formed a homogeneous, single cream layer similar to the comparator configuration 1E, which included suspending particles suspended alone. In contrast, the micronized GP particles alone (configuration 1D) flocculated and sedimented quickly. Configuration 1B showed no indication of separation of GP particles from the suspending particles even after centrifugation at 35g for 20 minutes. The same result was observed (i.e., lack of GP particle separation) when centrifuged up to

200g. Configuration 1C (low suspending concentration) showed a small amount of GP crystals settling out after centrifugation at 35g for 20 minutes.

[0142] While the co-suspended configurations achieved a delivered dose within 10 % of target, the GP particles suspended alone showed much higher variability in delivered dose in a range significantly below target. The fine particle fraction relative to configuration 1D was improved by more than 50%. The MMADs of the co-suspended configurations were acceptable and depended on the suspension concentration of the suspending particles. The delivered dose uniformity through use was tested for configurations 1B and 1C. All individual delivered doses were within $\pm 20\%$ of mean. The results showed that the drug crystals forming the GP particles associate to the suspending particles, a co-suspension was formed, and the aerosol performance of the co-suspension was mostly determined by the suspending particles.

[0143] The association between GP crystals and suspending particles was strong enough to overcome buoyancy forces, as it was observed that GP crystals do not separate from the perforated microstructures and settling of the crystals is inhibited.

Example 2

[0144] Glycopyrrolate (GP) particles were formed by micronization using a jet mill. Suspending particles were manufactured as described in Example 1. The particle size distribution of the micronized GP was determined by laser diffraction. 50% by volume of the micronized particles exhibited an optical diameter smaller than 1.7 μm , 90% by volume exhibited an optical diameter smaller than 4.1 μm . Five different lots of metered dose inhalers were different lots were made. For configurations 2A, 2B and 2C the total concentration of DSPC, CaCl_2 , and GP in the feedstock was 40 mg/mL, for configuration 2D and 2E this concentration was doubled.

[0145] Metered dose inhalers were prepared by weighing the target masses of GP particles and suspending particles into canisters as described in Example 1. No further excipients were used. The target masses were 4 mg / canister for GP particles and 60 mg / canister for the suspending particles, resulting in a suspending particle to GP particle ratio of 15 for configurations 2A and 2D. The target masses were 5.1 mg / canister for GP particles and 51 mg / canister for the suspending particles, resulting in a suspending particle to GP particle ratio of 10 for configuration

2B. The target masses were 8 mg / canister for GP particles and 60 mg / canister for the suspending particles, resulting in a suspending particle to GP particle ratio of 7.5 for configurations 2C and 2E. Propellant and container closure system were as described in Example 1.

[0146] The GP crystals were placed in HFA 134a in a canister under pressure and were equilibrated for 3 weeks at room temperature to determine their solubility in the propellant. The samples were filtered under pressure at ambient temperature through filters with a pore width of 0.22 µm. The filtrate was evaporated and the GP dissolved in methanol and chromatographically analyzed. A solubility of 0.17 ± 0.07 µg/g was found. Using this value it was determined that 2.1 µg or 0.05% of GP present in the canister dissolved in the propellant. Previous articles teach that microcrystalline material with a measurable solubility in the propellant will not be physically stable due to solution mediated transformation [N. C. Miller, The Effects of Water in Inhalation Suspension Aerosol Formulations, in: P. A. Byron, Ed. , Respiratory Drug Delivery, CRC Press, 1990, p 250], or that actives with solubility's above 0.1 µg/g should be formulated with an adjuvant to prevent a solution mediated transformation [P. Rogueda, Novel Hydrofluoroalkane Suspension Formulations for Respiratory Drug Delivery, Expert Opin. Drug Deliv. 2, 625-638, 2005].

[0147] The filled metered dose inhalers were stored valve down without overwrap at two different conditions: 1) refrigerated at 5°C; and 2) room temperature at 25°C / 60% RH. Aerosol performance and delivered dose uniformity tests as described in Example 1 were carried out at different time points. The results, which are summarized in Table 2, show a stable fine particle fraction at refrigerated and room temperature conditions.

Table 2: Fine particle fraction of configurations in Example 2

#	Storage	FPF in %			
		Initial	2 months	3 months	6 months
2A	5°C	49	51	52	-
	25 °C/60 % RH		48	51	-
2B	25 °C/60 % RH	50	46	49	48
2D	5 °C	51	54	54	-
	25 °C/60 % RH		46	49	49

[0148] Configurations 2C and 2E were subjected to a temperature cycling test. The canisters were subjected to -5 °C and 40 °C alternating between temperatures every 6 hours for a total duration of twelve weeks. Fine particle fraction was 53% for both configurations at the beginning of the study. After twelve weeks of cycling the FPF was unchanged, i.e. at 55% for configuration 2C and at 53% for configuration 2E.

[0149] The delivered dose uniformity through use was tested at the 1, 2 and 6 month time points. All individual delivered doses were within $\pm 20\%$ of mean. Figures 1 and 2 show the aerosol particle size distributions as measured by the NGI for configurations 2A and 2B, respectively. Also shown are the amounts of drug recovered from actuator, and from the induction port (throat) and its mouth piece adaptor. Recovered masses are expressed as percent of nominal dose. For configuration 2A, aerodynamic particle size distribution individual replicates are shown at 4, 8 and 12 weeks and at 8, 12 and 24 week for configuration 2B. Though there is a measureable fraction of the suspended GP dissolved in the propellant, there is no evidence of a coarsening of the size distributions. Moreover, as evidenced by these Examples, the aerosol performance of a co-suspension at suitable suspending particle to GP ratios is determined largely by the suspending particles.

Example 3

[0150] Several similar batches of suspending particles were made as described in Example 1. The suspending particles were combined with glycopyrrolate (GP) particles that were micronized to different extents, using two different types of jet mills with various milling parameters. The optical diameter and particle size distribution of the micronized GP particles was determined by laser diffraction. Table 3 lists the d_{50} and d_{90} values for the different lots of micronized material used. d_{50} and d_{90} denote the particle size at which the cumulative volume distribution reported by the particle sizing instrument reaches 50% and 90% respectively.

[0151] Twelve different lots of metered dose inhalers were prepared as described in Example 1. In all cases the suspension concentration of GP particles in HFA 134a was in the range of 0.32 – 0.45 mg/mL and the suspension concentration of the suspending particles was in the range of 5.8 – 6.1 mg/mL. The configurations were

deemed similar enough to pool the data for a meta-analysis presented in this Example.

[0152] The filled metered dose inhalers were stored valve down without overwrap at two different conditions: refrigerated at 5 °C and controlled room temperature at 25 °C / 60% RH. Aerosol performance tests as described in Example 1 were carried out at different time points. The results did not show any statistically significant trend as a function of time up to twelve weeks of storage. No difference between room temperature storage and refrigerated storage was discernible. Hence, results from different stress conditions and time points were pooled to determine how the particle size distribution of the micronized material affects aerosol performance.

[0153] Table 3 summarizes the MMAD results of the meta-analysis. The first column describes the six different configurations. The second column identifies how many individual lots were used in the compilation of the data for the respective configuration. The third column lists the number of individual MMAD determinations used to calculate the average MMAD for the respective configuration. Columns four and five show the d_{90} and d_{50} of the micronized material used to manufacture the co-suspensions. The results are sorted by d_{90} value from coarse to fine. The last two columns display the average MMAD and standard deviation.

Table 3: Pooled MMAD results for 12 glycopyrrolate co-suspensions, sorted by the d_{90} of the micronized glycopyrrolate particles.

Lot ID	number of lots	Number of MMAD measurements	d_{90} (μm)	d_{50} (μm)	Average MMAD (μm)	SD
3A	3	21	5.0	1.8	4.0	0.28
3B	2	9	4.9	2.1	4.1	0.37
3C	1	6	4.8	1.8	3.6	0.12
3D	1	4	4.3	1.7	3.5	0.22
3E	3	20	4.1	1.6	3.7	0.28
3F	2	10	3.5	1.7	3.6	0.10

[0154] These results show a weak dependence of MMAD on the d_{90} of the micronized material. A similar analysis for the d_{50} showed no statistically significant trend. It can be concluded that changes in the size distribution of the micronized material (e.g., different micronized material lots, or induced by solution mediated transformations) lead to only minor differences in the size distribution of the aerosol emitted from the metered dose inhaler.

Example 4

[0155] Micronized glycopyrrolate (GP) particles were formed tested as described in Example 1. The optical diameter of the micronized GP particles was determined and 50% by volume of the micronized GP particles were smaller than 1.7 μm , 90% by volume were smaller than 3.8 μm .

[0156] Five batches of suspending particles were made as described in Example 1. The batches differed in concentration, C_F , and volume fraction of PFOB, V_{PFOB} , of the feed emulsion prior to spray drying, ranging from 20 mg/mL to 160 mg/mL and 20% to 40%, respectively. The different configurations are described in Table 4.

[0157] Metered dose inhalers were prepared by weighing the target masses of micronized GP and suspending particles into coated glass vials with 15 mL volume. The target suspension concentrations and suspending particle to GP ratios are given in Table 4 for the 26 different vials tested. The canisters were crimp sealed with 63 μl valves (Valois, Les Vaudreuil, France) and filled with 10 g or 12 g of HFA 134a (1,1,1,2-tetrafluoroethane) (Ineos Fluor, Lyndhurst, UK) by overpressure through the valve stem. After injecting the propellant, the canisters were sonicated for 15 seconds and agitated on a wrist action shaker for 30 minutes.

[0158] As described in Example 1, micronized GP particles formulated alone flocculated and sedimented quickly. The glass vials in this example were left to settle for at least 24 h without agitation and then it was tested by visual observation whether the crystal, GP particles were co-suspended completely. For the vials marked with "Yes" in Table 4, no GP particles were observed at the bottom of the vials, except for very few foreign particulates in some vials. Occasional foreign particles were also visible in a similar very low amount in vials filled with suspending particles only. For the vials marked "Partial," a fraction of the GP particles was visible at the bottom of the vial.

Table 4: Co-suspension observations for glycopyrrolate configurations with various suspending particle to glycopyrrolate particle ratios.

	C_F in mg/mL	V_{PFOB} (%)	C_S (mg/mL)	Suspending particle to glycopyrrolate particle ratio	Co-suspension
#			Suspending particle		
4A	20	40	1.8	3.8	Partial
	20	40	7.2	15	Yes
4B	40	40	3.0	1.9	Partial
	40	40	1.8	3.8	Partial
4B	40	40	3.0	3.8	Yes
	40	40	6.0	3.8	Yes
4B	40	40	9.0	5.6	Yes
	40	40	3.0	7.5	Yes
4B	40	40	6.0	7.5	Yes
	40	40	9.0	11.3	Yes
4B	40	40	6.0	15	Yes
	40	40	7.2	15	Yes
4B	40	40	9.0	22.5	Yes
	80	20	3.0	1.9	Partial
4C	80	20	3.0	3.8	Partial
	80	20	6.0	3.8	Yes
4C	80	20	9.0	5.6	Yes
	80	20	3.0	7.5	Yes
4C	80	20	6.0	7.5	Yes
	80	20	9.0	11.3	Yes
4C	80	20	6.0	15	Yes
	80	20	9.0	22.5	Yes
4D	80	40	1.8	3.8	Partial
	80	40	7.2	15	Yes
4E	160	40	1.8	3.8	Partial
	160	40	7.2	15	Yes

Example 5

[0159] Glycopyrrolate (GP) particles were micronized with a jetmill and tested as described in Example 1. 50% by volume of the micronized particles exhibited an optical diameter smaller than 1.7 μm , 90% by volume exhibited an optical diameter smaller than 4.4 μm .

[0160] Six batches of suspending particles were made by spray drying as described in Example 1. Configuration 5A was spray dried from an emulsion.

Configuration 5B was manufactured in a similar fashion but using dipalmitoylphosphatidylcholine (DPPC) instead of DSPC. Configuration 5C was spray dried from an ethanolic solution. For configurations 5D, 5E, and 5F, saccharides were spray dried from aqueous solution. The spray drying parameters for all configurations are given in Table 5a.

Table 5a: Suspending particle configurations used in Example 5.

Lot #	Powder composition (% w/w)	Feed composition (% v/v)	C_F (mg/mL)	Spray Drying Parameters			
				Feed rate (mL/min)	T_{in} (°C)	T_{out} (°C)	Total Gas Flow (L/min)
5A	93.5 % DSPC 6.5 % CaCl_2	80 % H_2O 20 % PFOB	40	2.4	95	72	526
5B	92.9 % DPPC 7.1 % CaCl_2	70 % H_2O 30 % PFOB	60	2.4	95	67	525
5C	100 % DSPC	95 % Ethanol 5 % PFOB	100	5	95	70	520
5D	100 % Lactose	100 % H_2O	100	4	95	70	668
5E	100 % Trehalose	100 % H_2O	10	2.4	100	68	527
5F	100 % Trehalose	100 % H_2O	89	4	100	71	670

[0161] The particle size distribution of the suspending particles was determined by laser diffraction. The volume median optical diameter, VMD, and geometric standard deviation, GSD, for the different configurations are given in Table 5b.

Table 5b: Characteristics of suspending particle configurations used in Example 5.

Lot #	VMD (μ m)	GSD	Separation	Co-suspension	Comment
5A	3.6	1.8	Creams	Yes	No or few crystals visible on bottom of vials
5B	3.6	1.8	Creams	Yes	
5C	1.2	1.9	Creams	Partial	
5D	1.7	2.3	Sediments	Yes	Causes GP crystals to sediment with the suspending particles
5E	0.9	1.7	Sediments	Yes	
5F	1.7	2.4	Sediments	Yes	

[0162] Electron micrographs of the suspending particles showed a variety of morphologies, summarized in Figure 3. The particles that were spray dried from emulsion, 5A and 5B, had high porosity and low density. The DSPC particle spray dried from an ethanolic solution, 5C, showed a much smaller particle size with no noticeable porosity, indicating a high density. All saccharides produced smooth particles with no visible porosity. Configuration 5E had the smallest particles, as expected due to its low feed concentration.

[0163] Metered dose inhalers were prepared by weighing the 4 mg of micronized GP particles and 60 mg of suspending particles into coated glass vials with 15 mL volume. The canisters were crimp sealed with 63 μ L valves (Valois DF30/63 RCU, Les Vaudreuil, France) and filled with 9.5 mL of HFA 134a (Ineos Fluor, Lyndhurst, UK) by overpressure through the valve stem. After injecting the propellant, the canisters were sonicated for 15 seconds and agitated on a wrist action shaker for 30 minutes. Additional inhalers with suspending particles only were manufactured as control for each configuration.

[0164] The suspending particles in Examples 5A, 5B, and 5C, have true densities lower than the propellant. They formed a cream layer and were tested for the presence of a co-suspension as described in Example 4. No GP particles were visible at the bottom of the vials for configuration 5A and 5B. Configuration 5C formed a partial co-suspension.

[0165] The saccharide particles sediment because they have a higher true density than the propellant. However, all control vials for the saccharide configurations showed a significantly faster sedimentation rate than micronized GP

particles alone. In configurations 5D, 5E, and 5F, the sedimentation rate was similar to that of the control vials with the suspending particles alone and faster than the micronized GP particles alone, demonstrating the association of the GP crystals with the suspending particles. A co-suspension was formed in these cases. Figure 4 shows an example of this behavior for configuration 5D. The glass vial was observed one minute after agitation. The co-suspension has already settled leaving a clear propellant layer, while in the control containing GP particles alone, most of the crystals are still suspended in the propellant.

Example 6

[0166] Pharmaceutical compositions according to the present description were evaluated in a multi-center clinical trial. MDI devices containing a pharmaceutical composition of glycopyrrolate prepared according to the present description were provided.

[0167] Suspending particles used were prepared in a similar manner described in Example 1. MDI manufacturing was accomplished using a drug addition vessel (DAV) by first adding half of Suspending particle quantity, next filling the microcrystalline GP, and lastly adding the remaining half of suspending particles to the top. Materials were added to the vessel in a humidity controlled environment of <10% RH. The DAV was then connected to a 4 L suspension vessel and flushed with HFA 134a propellant and then mixed. The temperature inside the vessel was maintained at 21-23 °C throughout the entire batch production. After recirculation of the batch for 30 min canisters were filled with the suspension mixture through 50 µL EPDM valves. Sample canisters were then selected at random for total canister assay to ensure correct formulation quantities. The freshly manufactured co-suspension MDI batch was then placed on one week quarantine before initial product performance analysis.

[0168] The composition was formulated and the MDI devices configured to provide a dose of 18 µg glycopyrrolate per MDI actuation.

[0169] The study was a randomized, double-blind, four-period, six-treatment, placebo and active-controlled crossover study which evaluated single administration of 4 ascending doses of glycopyrrolate in patients with mild to moderate COPD compared to placebo and open label tiotropium (18 µg via the Spiriva Handihaler) as

an active control. The six study treatments were Glycopyrrolate MDI at doses of 18, 36, 72 and 144 µg were achieved by one, two, four or eight consecutive actuations of the 18 µg per actuation Glycopyrrolate MDI. Tiotropium Handihaler at 18 µg, and Placebo MDI, which was identical to the Glycopyrrolate MDI but without glycopyrrolate. Each patient was randomized to one of six possible sequences that included four of the study treatments. Each sequence included two or three Glycopyrrolate MDI doses, which were administered in ascending order to each patient. Glycopyrrolate MDI and Placebo MDI treatments were blinded and tiotropium was open label. Thirty-three patients were enrolled and analyzed for safety; thirty patients were analyzed for efficacy. Peak improvement in FEV₁ relative to test day baseline (FEV₁ is the maximum volume of air exhaled during the first second of maximum effort from a maximum inhalation), time to onset of action, time to peak FEV₁, FEV₁ AUC₀₋₁₂, FEV₁ AUC₀₋₂₄, FEV₁ AUC₁₂₋₂₄, 12 and 24-hour trough FEV₁, and similar analyses for peak expiratory flow rate (PEFR) and FVC, as well as peak improvement in inspiratory capacity (IC) were evaluated. Blood samples were collected pre-dose and 2, 6, 20 minutes, and 1, 2, 4, 8, 12, and 24 hours post-dose for determining plasma concentrations used to calculate PK parameters. The ratios of clinical spirometry outcomes (FEV₁) to glycopyrrolate PK outcomes (AUC₀₋₁₂ and C_{max}) were determined.

[0170] All doses of Glycopyrrolate MDI were safe and well tolerated, and the mean plasma glycopyrrolate concentration-time profiles were well characterized with rapidly occurring peak plasma concentrations, generally within 20 minutes. Plasma glycopyrrolate increased with dose level. Figure 5 shows the serum glycopyrrolate concentration (in pg/mL) compared to placebo over a 24 hour period experienced in the study subjects.

[0171] Glycopyrrolate MDI showed statistically significant and clinically relevant superior efficacy compared to Placebo MDI (p<0.001 for all four glycopyrrolate doses) with a clear dose response relationship. The efficacy of Glycopyrrolate MDI 144 µg and Glycopyrrolate 72 µg bracketed that of tiotropium 18 µg in terms of peak improvement in FEV₁ over time. For improvement in secondary FEV₁ endpoints relative to test day baseline, including trough FEV₁ at 12 hours, FEV₁ AUC₀₋₁₂, FEV₁ AUC₀₋₂₄, FEV₁ AUC₁₂₋₂₄, and 12 and 24-hour trough FEV₁, all doses of Glycopyrrolate MDI demonstrated clinically relevant and statistical superiority

compared to Placebo MDI ($p \leq 0.049$ for all four dose levels), with the exception of improvement in trough FEV₁ at 24 hours following administration of Glycopyrrolate MDI 36 μ g (difference compared to placebo = 0.073L; $p=0.059$). Similar to the clear dose-response relationship observed for improvement in peak FEV₁, dose ordering across all four doses of Glycopyrrolate MDI evaluated was also observed for improvements in FEV₁ AUC₀₋₁₂, FEV₁ AUC₀₋₂₄, and FEV₁ AUC₁₂₋₂₄.

[0172] The Glycopyrrolate MDI 144 μ g and 72 μ g doses were shown to be statistically non-inferior to tiotropium 18 μ g in terms of peak change in FEV₁, FEV₁ AUC₀₋₁₂, and FEV₁ AUC₀₋₂₄, with the *a priori* defined non-inferiority bound of 100 mL. The Glycopyrrolate 144 μ g dose was also non-inferior to tiotropium for 12-hour trough and FEV₁ AUC₁₂₋₂₄. Point-estimates for the majority of the FEV₁ parameters for the 72 and 144 μ g doses were within ± 50 mL compared to tiotropium. In general, the secondary endpoints (time to onset of effect, peak and trough FEV₁, FVC, PEFR, and peak IC) confirmed the findings of the primary endpoint. Glycopyrrolate MDI demonstrated a more rapid onset of action compared to tiotropium 18 μ g, with mean time to $\geq 10\%$ improvement in FEV₁ of 1 hour or less for all doses of Glycopyrrolate MDI evaluated, compared to approximately 3 hours for tiotropium 18 μ g.

[0173] Figure 6 plots the mean change in FEV₁ from baseline (in liters) experienced by the study subjects over a period of 24 hours. Figure 7 depicts the change in FEV₁ from baseline (in liters) for patients at different glycopyrrolate dosing levels compared to the results obtained for tiotropium. Specifically, Figure 7 compares the peak change from baseline over the placebo value for different glycopyrrolate concentrations and the area under the curve over a 12 hour and 24 hour period. Figure 8 depicts the proportion of patients which experienced either 1) an increase from baseline in FEV₁ of at least 200 mL or 2) a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 mL or greater. Figure 9 shows the peak improvement in IC experienced by patients administered the various doses of Glycopyrrolate, as well as the peak improvement in IC for patients receiving tiotropium. Figure 10 shows change in FEV₁ cumulatively over a 24 hour period in patients receiving glycopyrrolate, compared with the results obtained from another clinical study where NVA237 (a powder formulation of glycopyrrolate) was given at various doses by Singh *et al* (D Singh, P A Corris, and

S D Snape. "NVA237, a once-daily inhaled antimuscarinic, provides 24-hour bronchodilator efficacy in patients with moderate to-severe COPD" Poster presented at the American Thoracic Society International Conference, San Diego, California, May 19-24, 2006).

Example 7

[0174] Glycopyrrolate (GP) was micronized using a jet mill to a volume median optical diameter (d_{50}) of 1.4 μm with 90% of the cumulative distribution (d_{90}) having a volume optical diameter below 3.0 μm . Suspending particles were manufactured similarly to those in Example 1. MDI canisters were manufactured using FEP coated Presspart cans to provide products with metered dose of 5.5 $\mu\text{g}/\text{actuation}$ GP and 44 $\mu\text{g}/\text{actuation}$ GP which correlates to approximately 4.5 $\mu\text{g}/\text{actuation}$ and 36 $\mu\text{g}/\text{actuation}$ GP delivered dose from a 50 μl volume metering chamber from commercially available Bespak valves. The formulations contained 6mg/mL of suspending particles. The MDI canisters were manufactured using standard pressure filling process where drug substance and the suspending were mixed with HFA 134a in a suspension vessel and filled into canisters through a commercially available filling head.

[0175] Each lot was tested for delivered dose uniformity through can life and aerodynamic particle size distribution by Next Generation Impactor after manufacture. The aerodynamic particle size distributions as measured by the NGI are shown in Figures 11 and 12. Also shown are the amounts of drug recovered from valve stem and actuator, and from the induction port (throat) and its mouth piece adaptor. Recovered masses are expressed as percent of nominal dose. The fine particle fraction remained unchanged over 168 cycles, illustrating the stability of the GP co-suspensions disclosed herein across a GP dose range. The delivered dose through life of the MDI canisters is shown in Figures 13 and 14. No change in delivered dose from beginning to middle of can is observed and a ~10% increase from middle to end of canister. The change from middle to end is anticipated based upon evaporative losses of propellant as the can is emptied. The compositions described in this example demonstrate desirable delivered dose uniformity for MDI for doses as low as 4.5 $\mu\text{g}/\text{actuation}$.

[0176] In addition, canisters from each lot were subjected to a temperature cycling stability study. The canisters were subjected to -5 °C and 40 °C alternating between temperatures every 6 hours for a total duration of 84 cycles (3 weeks) and 168 cycles (6 weeks). After 184 cycles, the % FPF (ex-actuator) is not significantly different from initial. A summary of the stability of the fine particle fraction is shown in Table 6.

Table 6. Temperature Cycling Stability of the Fine Particle Fraction of crystalline GP co suspended with suspending particles at two doses in MDI containing HFA 134a

Timepoint	4.5 µg/actuation (%FPF ex-actuator)	36 µg/actuation (%FPF ex-actuator)
Initial	60.9	57.4
3 Weeks (84 cycles)	61.9	58.0
6 Weeks (168 cycles)	60.6	59.0

Example 8

[0177] MDI Canisters were manufactured to contain 6mg/mL suspending particle concentration and to provide a metered dose of 36 µg/actuation with a 50µl valve volume according to Example 7. Micronized GP had a d_{50} and d_{90} of 1.6µm and 4.1µm respectively and suspending particles were manufactured similarly to the process described in Example 1. The canisters were placed on stability without protective packaging at 25°C/60% RH and stored for duration of 12 months. Aerodynamic particle size distribution was determined by next generation impaction at 2 weeks, 1, 2, 3, 6 or 12 months. The fine particle fraction, as a percentage of GP ex-actuator, at initial sampling was 50.2%. No significant change in the fine particle fraction was noted at any of the timepoints out to 12 months, with FPF of 47.7% after 12 months. Figure 15 provides a view of the entire aerodynamic size distribution for each of the timepoints demonstrating desirable consistency on aerosol delivery. A summary of the fine particle fraction is shown in Table 7.

Table 7. Stability of the Fine Particle Fraction of crystalline GP co suspended with suspending particles in MDI containing HFA 134a at 25°C and 60%RH with no protective packaging

Time-Point	% FPF (ex actuator)
Initial	50.2
2 Week	46.1
1 Month	42.0
2 Month	46.0
3 Month	48.9
6 Month	47.7
12 Month	47.7

Example 9

[0178] MDI Canisters were manufactured to contain 6mg/mL suspending particle concentration and to provide a metered dose of 36 μg /actuation as described in Example 7. These canisters were packaged in a heat sealed aluminum foil overwrap containing desiccant, and cycled for 6 weeks (6 hours at -5 °C and 6 hours at 40 °C). The delivered dose uniformity through use was tested at the 0, 2, 4 and 6 weeks time points. The mean glycopyrrolate delivered dose of each lot each time period was within $\pm 15\%$ of the mean, with one exception, as demonstrated in Figure 16. The aerodynamic particle size distribution as measured by NGI remain unchanged after 168 temperature cycles as shown in Figure 17.

Example 10

[0179] MDI Canisters were manufactured to contain 6mg/mL suspending particle concentration and to provide a metered dose of 24 μg per actuation according to Example 7. These canisters were stored for six weeks at 50 °C under ambient humidity. Another lot was stored for 8 weeks at 40 °C and 75% relative humidity. Yet another lot was stored for 12 weeks at 40 °C and 75% relative humidity. The fine particle fraction was 59.3% initially. The canister stored for 6 weeks at 50 °C had an FPF that was unchanged compared to the initial lot, i.e. at 58.4 %. The lot stored at 40 °C for 8 and 12 weeks had an FPF that was also unchanged compared to the initial, i.e. at 56.8 % and 57.6% respectively. The aerodynamic particle size distributions as measured by the NGI are shown in Figure 18. The MMAD remains relatively unchanged after 6 weeks at 50 °C, 3.94 μm , and up to 12 weeks at 40 °C, 3.84 μm , compared to the initial at 3.54 μm . In addition, the FPF and the amounts of

glycopyrrolate recovered from valve stem and actuator, and from the induction port (throat) and its mouth piece adaptor, remained relatively unchanged over 3 months at elevated temperatures.

Example 11

[0180] Metered dose inhalers including pharmaceutical compositions of formoterol fumarate as described herein were prepared. Formoterol fumarate, (\pm)-2-hydroxy-5-[(1RS)-1-hydroxy-2-[[[(1RS)-2-(4-methoxyphenyl)-1-methylethyl]-amino]ethyl] formanilide fumarate, also known as (\pm)-2'-hydroxy-5'-[(RS)-1-hydroxy-2-[[[(RS)-p-methoxy- α -methylphenethyl]-amine]ethyl] formanilide fumarate, dihydrate was micronized to form active agent particles. The particle size distribution of the micronized formoterol fumarate (FF) was determined by laser diffraction. 50% by volume of the micronized particles exhibited an optical diameter smaller than 1.6 μ m, and 90% by volume exhibited an optical diameter smaller than 3.9 μ m.

[0181] Suspending particles were manufactured as follows: 503 mL of a fluorocarbon-in-water emulsion of PFOB (perfluorooctyl bromide) stabilized by a phospholipid was prepared. 20.6 g of the phospholipid, DSPC (1,2-disteroyl-sn-glycero-3-phosphocholine), and 1.9 g of calcium chloride were homogenized in 403 mL of hot water (75°C) using a high shear mixer. 100 mL of PFOB were added slowly during homogenization. The resulting coarse emulsion was then further homogenized using a high pressure homogenizer (Model C3, Avestin, Ottawa, CA) at pressures of up to 170 MPa for 5 passes.

[0182] The emulsion was spray dried in nitrogen using the following spray drying conditions: Inlet temperature 95°C, outlet temperature 71°C, emulsion feed rate 2.4 mL/min, total gas flow 498 L/min. The particle size distribution of the suspending particles was determined by laser diffraction. 50% by volume of the suspending particles were smaller than 3 μ m, the geometric standard deviation of the distribution was 1.9.

[0183] Metered dose inhalers were prepared by weighing the target masses of micronized active agent particles and suspending particles into coated glass vials with 15 mL volume. The target masses and the target delivered dose assuming 20% actuator deposition are given in Table 8 for three different configurations. For each configuration, additional glass bottles were filled with the respective amount of FF

active agent particles without any suspending particles. The canisters were crimp sealed with 63 μ l valves (Valois, Les Vaudreuil, France) and filled with 11 g (9.1 mL at 25 °C) of HFA 134a (1,1,1,2-tetrafluoroethane) (Ineos Fluor, Lyndhurst, UK) by overpressure through the valve stem. After injecting the propellant, the canisters were sonicated for 15 seconds and agitated on a wrist action shaker for 30 minutes.

Table 8: Target doses for formoterol fumarate co-suspensions of Example 10

Configuration #	FF Active Agent Particles μ g/can	Suspending Particles mg/can	Target delivered dose μ g	Suspending Particle to active particle ratio
6A	300	50	1.7	167
6B	860		4.6	58
6C	3010		16.5	16.6

[0184] Visual observation of the co-suspended configurations (6A, 6B, 6C) showed no sedimentation of the crystalline FF forming the active agent particles. The suspension flocculated slowly and formed a homogeneous, single cream layer. For all concentrations tested the micronized active agent particles alone sedimented quickly. Pictures of the co-suspension and the traditional comparator suspensions, indicated by an asterisk, are shown in Figure 19. The vials were left to settle for 24 h without agitation. No FF crystals were visible at the bottom of any of the co-suspension vials.

[0185] The results showed that the FF crystals associated with the suspending particles. The association between FF particles and suspending particles was strong enough to overcome buoyancy forces, as FF particles did not separate from the suspending particles and settling of the active agent particles was successfully inhibited in each of the three different formulation configurations.

Example 12

[0186] Formoterol fumarate MDI compositions were prepared according to the present invention. Micronized formoterol fumarate was commercially obtained and its particle size distribution measured as described in Example 1 was characterized by a d_{10} , d_{50} , d_{90} of 0.6, 1.9 and 4.4 μ m respectively and a Span of 2.0. Suspending

particles used were prepared in a similar manner described in Example 1. MDI manufacturing was accomplished using a drug addition vessel (DAV) by first adding half of suspending particle quantity, next filling the microcrystalline FF, and lastly adding the remaining half of suspending particles to the top. Materials were added to the DAV in a humidity controlled environment of <10% RH. The DAV was then connected to a 4 L suspension vessel. A slurry was then formed by adding a known amount of HFA-134a propellant (Ineos Fluor, Lyndhurst, UK) into the DAV, which is then removed from the suspension vessel and gently swirled. The slurry is then transferred back to the suspension mixing vessel and diluted with additional HFA-134a to form the final suspension at target concentration stirring gently with an impeller. The temperature inside the vessel was maintained at 21-23 °C throughout the entire batch production. After recirculation of the batch for 30 min, 14-mL fluorinated ethylene polymer (FEP) coated aluminum canisters (Presspart, Blackburn, UK) were filled with the suspension mixture through 50 µL EPDM valves (Bespak, King's Lynn, UK). Sample canisters were then selected at random for total canister assay to ensure correct formulation quantities.

[0187] The freshly manufactured co-suspension MDI batch was then placed on one week quarantine before initial performance analysis. Aerosol performance was assessed in accordance with USP <601> (United States Pharmacopeia monograph 601). A Next Generation Impactor (NGI) operated at a flow rate of 30 L/min was used for determination of particle size distribution. Sample canisters were seated into an actuator with two waste actuations and two additional waste priming actuations. Five actuations were collected in the NGI with a USP throat attached. The valve, actuator, throat, NGI cups, stages, and filter were rinsed with volumetrically dispensed solvent. The sample solutions were assayed using a drug specific chromatographic method. The fine particle fraction was defined using the sum of stages 3 through filter. Delivered dose uniformity through use testing was performed using a Dose Uniformity Sampling Apparatus as described by USP <601>. Two actuations were collected and assayed at beginning, middle and end of use.

[0188] Figure 20 shows the delivered dose uniformity for a co-suspension of FF at a 4.8 µg target dose per actuation. The individual delivered dose per actuation for

beginning, middle and end of actuations was within $\pm 25\%$ of the mean delivered dose.

Example 13

[0189] Formoterol Fumarate MDI compositions were prepared according to the present invention. Micronized formoterol fumarate was commercially obtained and its particle size distribution measured as described in Example 1 was characterized by a d_{10} , d_{50} , d_{90} of 0.6, 1.9 and 4.4 μm respectively and a Span of 2.0. Suspending particles used were prepared in a similar manner described in Example 1. MDI manufacturing was accomplished as described in Example 12.

Aerosol performance was assessed in accordance with USP <601>. A Next Generation Impactor (NGI) operated at a flow rate of 30 L/min was used for determination of particle size distribution. Sample canisters were seated into an actuator with two waste actuations and two additional waste priming actuations. Five actuations were collected in the NGI with a USP throat attached. The valve, actuator, throat, NGI cups, stages, and filter were rinsed with volumetrically dispensed solvent. The sample solutions were assayed using a drug specific chromatographic method. The fine particle fraction was defined using the sum of stages 3 through filter. The aerodynamic particle size distribution of a FF co-suspension formulation was evaluated after manufacture and after three months of storage at 25 °C and 75%RH (unprotected canisters) and 40 °C and 75 %RH (protected canisters wrapped in aluminum foil pouch). The aerodynamic particle size distributions shown in Figure 21 demonstrate that the compositions described in the present invention display desirable stability characteristics even at accelerated conditions.

Example 14

[0190] The chemical stability of formoterol fumarate (FF) included in a co-suspension formulation prepared according Example 11 was evaluated. FF MDI canisters containing HFA 134a were overwrapped with an aluminum foil pouch and stored at 25 °C and 60% relative humidity and 40 °C and 75% relative humidity for thirteen and six months, respectively. Likewise FF MDI canisters containing HFA

227ea were overwrapped with an aluminum foil pouch and stored at 25 °C and 60% relative humidity and 40 °C and 75% relative humidity for six months. The amount of impurity F, a characteristic degradation product of FF, and total impurities were determined by reverse phase HPLC assay as follows: each canister is chilled, cut open, and the can contents are transferred to a centrifuge tube; the contents were dissolved in organic solvent, followed by the addition of an aqueous solvent to precipitate excipient (DSPC) from the solution; the solution was centrifuged to produce a clear supernatant solution; and each sample solution was analyzed using a C18 column, 4.6 x 150 mm and 3.0 µm particle size. The column temperature was kept at 30 °C. The injection volume was 20 µl, and flow rate was set at 1 mL/min and detected by determining the UV absorption at 214 nm. A gradient was used mixing pH 3.1 aqueous phosphate buffer and acetonitrile, 17% acetonitrile first 27 minutes, then 50% acetonitrile for 30 seconds followed by 6.5 minutes at 75% acetonitrile and 17% acetonitrile for 8 minutes. Impurities were reported as area percent of formoterol peak area (corrected for relative response factors, where available). As shown in Figure 22 (or Table 9 and 10), a co-suspension prepared using crystalline FF active agent particles suspended in HFA 134a with suspending particles was chemically stable for 18 months at a temperature of 25°C and 60% relative humidity, in contrast a spray dried, non co-suspended formoterol formulation showed a faster degradation rate under the same storage conditions. Likewise crystalline FF active agent particles formed a chemically stable co-suspension in HFA 227a, as shown in Table 11.

Table 9. Chemical Stability of Spray Dried FF Suspending Particles in FF MDI Containing HFA 134a at 25°C/60%RH, Overwrapped in Aluminum Foil Pouches

Time (months)	0	2	3	12	18
Impurity F (%)	ND	0.12%	0.04%	1.16%	2.77%
Total Impurities (%)	0.62%	1.42%	1.75%	2.33%	4.39%

ND= Not detected

Table 10. Chemical Stability of Crystalline FF Co-suspended with Suspending Particles in FF MDI Containing HFA 134a at 25°C/60%RH, Overwrapped in Aluminum Foil Pouches

Time (months)	0	1	2	3	6	10	13
Impurity F (%)	0.05%	0.08%	0.08%	0.14%	0.06%	0.22%	0.35%
Total Impurities (%)	0.44%	0.32%	0.32%	0.37%	0.18%	0.45%	0.64%
at 40°C/75%RH, Overwrapped in Aluminum Foil Pouches							
Time (months)	0	1	2	3	6		
Impurity F (%)	0.05%	0.11%	0.31%	1.18%	1.74%		
Total Impurities (%)	0.44%	0.41%	0.75%	1.58%	2.54%		

Table 11. Chemical Stability of Crystalline FF Co-suspended with Suspending Particles in FF MDI Containing HFA 227ea at 25°C/60%RH, Overwrapped in Aluminum Foil Pouches

Time (months)	0	1	2	3	6
Impurity F (%)	0.04	0.06	0.07	0.13	0.05
Total Impurities (%)	0.4	0.3	0.3	0.4	0.1
at 40°C/75%RH, Overwrapped in aluminum foil pouches					
Time (months)	0	1	2	3	6
Impurity F (%)	0.04	0.08	0.18	0.80	1.14
Total Impurities (%)	0.40	0.39	0.53	1.13	1.56

Example 15

[0191] Micronized formoterol fumarate dihydrate (FF) (Inke, S.A., Barcelona, Spain) used in the present example had with particle size distribution by laser diffraction of 50% by volume of the micronized particles exhibited an optical diameter smaller than 1.9 μm , 90% by volume exhibited an optical diameter smaller than 4.1 μm . Four batches of suspending particles were manufactured by spray drying as described in Example 1. All four batches were spray-dried from aqueous solution; solution concentration and spray drying parameters are given in Table 12.

Table 12: Suspending particle configurations used in Example 15

#	Powder Composition	C_f in mg/mL	Spray Drying Parameters				Particle Size Distribution	
			Feed rate	T_{in} in	T_{out}	Total Gas	VMD in	GSD

			in mL/min	° C	in ° C	Flow in std L/min	μm	
XA	100 % trehalose	80	10	150	82	385	1.62	2.20
XB	100 % HP- β -cyclodextrin	80	10	100	68	885	1.61	2.21
XC	100% Ficoll PM 70	80	10	100	70	885	1.19	2.27
XD	100% Inulin	80	10	100	70	885	1.23	2.20

[0192] Electron micrographs of the suspending particles showed a variety of morphologies, and are shown in Figure 23 through Figure 26, with Figure 23 providing a micrograph of trehalose suspending particles, Figure 24 providing a micrograph of HP- β -cyclodextrin suspending particles, Figure 25 providing a micrograph of Ficoll MP 70 suspending particles, and Figure 26 providing a micrograph of inulin suspending particles. Trehalose particles appear to be spherical, with a smooth surface. HP- β -cyclodextrin particles show extensive wrinkling of the surface, suggesting a partially buckled exterior with a hollow core. Ficoll MP 70 and Inulin particles display some surface rugosity but are generally spheroidal.

[0193] Metered dose inhalers were prepared by weighing 0.9 mg of the micronized FF active agent particles and 60 mg of suspending particles into coated glass vials with 15 mL volume. FF was combined with each type of the four suspending particle species of Table 11. The canisters were crimp sealed with 50 μ L valves (Valois DF31/50 RCU, Les Vaudreuil, France) and filled with 10 mL of HFA propellant 134a (Ineos Fluor, Lyndhurst, UK) by overpressure through the valve stem. After injecting the propellant, the canisters were sonicated for 30 seconds and agitated on a wrist action shaker for 30 minutes. Additional inhalers containing suspending particles only and active agent particles only were filled as a control for each configuration.

[0194] Crystalline FF has a greater density than propellant 134a at room temperature, as do all four species of suspending particles in the present example. Consequently both FF and suspending particles settled to the bottom of the inhalers at room temperature. To test these inhalers for active-suspending agent particle interactions indicating a co-suspension, the inhalers were immersed in an ethanol

bath at $\leq -10^{\circ}\text{C}$ (resulting in increased propellant density) and allowed to equilibrate for a minimum of 30 minutes. At this temperature, the FF active agent particles are less dense than the propellant and consequently cream to the top of the propellant volume, while all four species of suspending agent particles remain settled at the bottom of the propellant volume.

[0195] The tested configurations and the results of the observations are presented in Table 13. FF active agent particles alone formed a cream layer atop the propellant volume, and trehalose, HP- β -cyclodextrin, inulin, and Ficoll PM70 particles alone all settled to the bottom of the glass vial. FF active agent particles in combination with trehalose suspending particles formed a single sediment layer, with no particles creamed or afloat in the propellant, indicating that the FF particles interact with the trehalose suspending particles, and a co-suspension is formed. In the case of FF particles in combination with HP- β -cyclodextrin suspending particles, some turbidity was present in the propellant, similar to that observed in the suspending particle only control vial. Additionally, some floating flocs were observed, which may have been FF particles; however, such flocs accounted for a small amount of solid mass relative to the control vial, indicating that some if not all FF particles were interacting with the suspending agent particles. Thus, this configuration is an example of a partial co-suspension. FF particles in combination with inulin suspending particles formed a single sediment layer, indicating a co-suspension was formed. Though some turbidity was present in this configuration, similar cloudiness was observed in the inulin-only control vial. FF active agent particles in combination with Ficoll PM70 suspending particles formed a sediment layer at the bottom of the vial, indicating that a co-suspension was formed. While some turbidity and floating flocs were observed in this configuration, similar turbidity, and floc frequency were observed in the Ficoll-only control vial.

Table 13: Summary of tested configurations and results of observations

Container ID	Contents in 10 mL p134a	Suspending Particle to Active Particle Ratio	Observational Notes, $\leq -10^{\circ}\text{C}$	Co-suspension
0-FF	0.9 mg FF	n/a	Creamed to top	n/a
T	60 mg trehalose	n/a	Settled to bottom	n/a

Container ID	Contents in 10 mL p134a	Suspending Particle to Active Particle Ratio	Observational Notes, $\leq -10^{\circ}\text{C}$	Co-suspension
T-FF	60 mg trehalose, 0.9 mg FF	67	Sediment layer; no particles creamed	Yes
C	60 mg HP- β -cyclodextrin	n/a	Settled to bottom; some turbidity	n/a
C-FF	60 mg HP- β -cyclodextrin, 0.9 mg FF	67	Solids mostly in sediment layer at bottom; some turbidity; some floating flocs present	partial
I	60 mg Inulin	n/a	Settled to bottom; some turbidity	n/a
I-FF	60 mg Inulin, 0.9 mg FF	67	Sediment layer; no particles creamed; some turbidity	Yes
F	60 mg Ficoll PM70	n/a	Settled to bottom, with some floating flocs	n/a
F-FF	60 mg Ficoll PM70, 0.9 mg FF	67	Sediment layer; very few floating flocs	Yes

Example 16

[0196] Co-suspension compositions including glycopyrrolate (GP) and formoterol fumarate (FF) active agent particles were produced and MDIs incorporating the co-suspension compositions were prepared. The co-suspension compositions produced included GP active agent particles, FF active agent particles or a combination of both GP and FF active agent particles. The GP and FF material was supplied as micronized, crystalline material with particle size distribution as shown in Table 14.

[0197] Suspending particles were manufactured via spray dried emulsion at a feed stock concentration of 80 mg/mL with a composition of 93.44% DSPC (1,2-Distearoyl-sn-Glycero-3-Phosphocholine) and 6.56% anhydrous calcium chloride (equivalent to a 2:1 DSPC:CaCl₂ mole/mole ratio). During the emulsion preparation, DSPC and CaCl₂ was dispersed with a high shear mixer at 8000-10000 rpm in a vessel containing heated water ($80 \pm 3^{\circ}\text{C}$) with PFOB slowly added during the process. The emulsion was then processed with 6 passes in a high pressure homogenizer (10000-25000 psi). The emulsion was then spray dried via a spray drier fitted with a 0.42" atomizer nozzle with a set atomizer gas flow of 18 SCFM.

The drying gas flow rate was set to 72 SCFM with an inlet temperature of 135 °C, outlet temperature 70 °C, and an emulsion flow rate of 58 mL/min.

[0198] The co-suspensions were prepared by first dispensing the appropriate quantities of micronized GP and FF active agent particles and suspending particles into a drug addition vessel (DAV) inside a humidity controlled chamber (RH < 5%). In the present Example, the suspending particles were added in three equal portions intercalating the addition of GP and FF after the first and second addition respectively. The DAV is then sealed under a nitrogen atmosphere and connected to the suspension vessel containing 12 kg of HFA-134a (Ineos Fluor, Lyndhurst, UK). A slurry was then formed by adding 0.5-1 kg of HFA-134a into the DAV, which is then removed from the suspension vessel and gently swirled. The slurry is then transferred back to the suspension mixing vessel and diluted with additional HFA-134a to form the final suspension at target concentration stirring gently with an impeller. The suspension is then recirculated via a pump to the filling system for a minimum time prior to initiation of filling. Mixing and recirculation continue throughout the filling process. 50 µL valves (Bespak, King's Lynn, UK) are placed onto 14-mL fluorinated ethylene polymer (FEP) coated aluminum canisters (Presspart, Blackburn, UK) canisters and then purged of air either by a vacuum crimping process, or an HFA-134a purging process followed by valve crimping. The crimped canisters are then filled through-the-valve with the appropriate quantity of suspension, adjusted by the metering cylinder.

Table 14: Glycopyrrolate and Formoterol Fumarate particle size distributions.

Designation	d ₁₀ (µm)	d ₅₀ (µm)	d ₉₀ (µm)	Span
FF API	0.6	1.9	4.1	1.8
GP API	0.5	1.3	3.0	1.9

[0199] MDIs containing the dual co-suspensions described in this Example were prepared to contain two different doses GP and FF. Specifically, a first run of dual co-suspension compositions were prepared to provide 18 µg per actuation GP and 4.8 µg per actuation FF ("low dose"), and a second run of dual co-suspension compositions were prepared to provide 36 µg per actuation GP and 4.8 µg per actuation FF ("high dose"). In addition to the dual co-suspensions compositions, co-

suspensions including a single species of active agent particle were prepared. These compositions included either GP active agent particles or FF active agent particles and were referred to as "mono" or "monotherapy" co-suspensions. The monotherapy co-suspension compositions were prepared as described for the dual co-suspensions, except that they included only one species of active agent particles (either GP or FF). The monotherapy co-suspensions were formulated and monotherapy MDIs prepared to provide the following targeted delivered doses: 18 μ g per actuation of GP, and 0.5, 1.0, 3.6 or 4.8 μ g per actuation of FF. The compositions and MDIs providing 0.5 μ g FF and 1 μ g FF per actuation are referred to as "ultra low" dose and were manufactured in a similar manner at a 4L scale.

[0200] The drug specific aerodynamic size distributions achieved with MDIs containing the co-suspension compositions prepared according to this Example were determined as described in Example 1. The proportionality of the aerodynamic size distributions of GP obtained from the low and high dose dual co-suspensions as well as the equivalency between the dual and monotherapy co-suspensions is demonstrated in Figure 27. In the same manner, the proportionality of the aerodynamic size distributions of FF obtained from the dual and monotherapy co-suspensions, including the ultralow, low, and high dose compositions is demonstrated in Figure 28.

[0201] The delivered dose uniformity of the ultra low dose FF monotherapy MDIs was also measured as described in Example 1. The DDU for the FF MDI containing 0.5 μ g per actuation and 1.0 μ g per actuation are shown in Figure 29. Desirable dose delivery uniformity is achieved demonstrating the utility of the present invention to consistently deliver ultra low doses. In order to evaluate whether the combination of GP and FF within a single formulation would result in the degradation of the aerosol properties relative to compositions including a single active agent, the aerosol properties of co-suspension compositions were assessed relative to suspension compositions including only a single active agent. As can be seen in Figure 30, the aerosol performance of the combination co-suspension composition including both GP and FF active agent was no different than the aerosol performance achieved by suspension compositions including either GP or FF alone. Therefore, there were no combination effects observed.

Example 17

[0202] Micronized salmeterol xinafoate (4-hydroxy- α 1-[[6-(4-phenylbutoxy)hexyl]amino] methyl]-1,3-benzenedimethanol, 1-hydroxy-2-naphthalenecarboxylate) was received by the manufacturer (Inke SA, Germany) and used as active agent particles. The particle size distribution of the salmeterol xinafoate (SX) was determined by laser diffraction. 50% by volume of the micronized particles exhibited an optical diameter smaller than 2 μ m, 90% by volume exhibited an optical diameter smaller than 3.9 μ m.

[0203] Suspending particles were manufactured as follows: 150 mL of a fluorocarbon-in water emulsion of PFOB (perfluorooctyl bromide) stabilized by a phospholipid was prepared. 12.3 g of the phospholipid, DSPC (1,2-Distearoyl-sn-Glycero-3-Phosphocholine), and 1.2 g of calcium chloride were homogenized in 100 mL of hot water (70 °C) using a high shear mixer. 65 mL of PFOB were added slowly during homogenization. The resulting coarse emulsion was then further homogenized using a high pressure homogenizer (Model C3, Avestin, Ottawa, CA) at pressures of up to 140 MPa for 3 passes

[0204] The emulsion was spray dried in nitrogen using the following spray drying conditions: Inlet temperature 90 °C, outlet temperature 69 °C, emulsion feed rate 2.4 mL/min, total gas flow 498 l/min. The particle size distribution of the suspending particles, VMD, was determined by laser diffraction. 50% by volume of the suspending particles were smaller than 2.7 μ m, the Geometric Standard Deviation of the distribution was 2.0. Additionally, the aerodynamic particle size distribution of the suspending particles was determined with a time-of-flight particle sizer. 50% by volume of the suspending particles had an aerodynamic particle diameter smaller than 1.6 μ m. The large difference between aerodynamic particle diameter and optical particle diameter indicates that the suspending particles had a low particle density < 0.5 kg/L.

[0205] Metered dose inhalers were prepared by weighing 2 mg of SX active agent particles and 60 mg of suspending particles into fluorinated ethylene polymer (FEP) coated aluminum canisters (Presspart, Blackburn, UK) with 19 mL volume. The suspending particle to active particle ratio was 30. The target delivered dose assuming 20% actuator deposition was 10 μ g. The canisters were crimp sealed with 63 μ l valves (# BK 357, Bespak, King's Lynn, UK) and filled with 10 mL of HFA 134a

(1,1,1,2-tetrafluoroethane) by overpressure through the valve stem. After injecting the propellant, the canisters were sonicated for 15 seconds and agitated on a wrist action shaker for 30 minutes. The canisters were fitted with polypropylene actuators with a 0.3 mm orifice (# BK 636, Bespak, King's Lynn, UK). Additional inhalers for visual observation of suspension quality were prepared using 15 mL glass vials including a comparator filled with micronized SX only. Aerosol performance was assessed as described in Example 1. The MMAD was 3.7 μm and the fine particle fraction was 48%. Because the SX crystals forming the active agent particles and the propellant were nearly density matched at 15 °C – 20 °C, the visual observation was conducted on glass vials that were heated up to 30 °C – 35 °C in a water bath. Under these conditions the SX active agent particles formulated alone sedimented rapidly, but no SX crystals were visible at the bottom of the co-suspension vial.

[0206] Micronized salmeterol xinafoate active agent particles were co-suspended through association with suspending particles of low density that were formulated according to the disclosure provided herein. The association between salmeterol crystals and the suspending particles was strong enough to overcome buoyancy forces as it was observed that settling of the crystals is inhibited.

CLAIMS

1. A pharmaceutical composition deliverable from a metered dose inhaler, comprising:
 - a suspension medium comprising a pharmaceutically acceptable propellant;
 - a plurality of active agent particles comprising an active agent selected from a long-acting muscarinic antagonist (LAMA) active agent and a long-acting β_2 adrenergic receptor agonist (LABA) active agent; and
 - a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles to form a co-suspension.
2. A pharmaceutical composition according to claim 1, wherein the active agent included in the active agent particles is a LAMA active agent selected from glycopyrrolate, dexipirronium, tiotropium, trospium, aclidinium, darotropium, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.
3. The pharmaceutical composition according to claim 2, wherein the active agent particles comprise glycopyrrolate, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof.
4. The pharmaceutical composition according to claim 3, wherein the active agent particles comprise crystalline glycopyrrolate.
5. The pharmaceutical composition according to claim 3, wherein the glycopyrrolate active agent particles are included in the suspension medium at a concentration sufficient to provide a delivered dose of glycopyrrolate per actuation of the metered dose inhaler selected from between about 2 μ g and about 200 μ g per actuation, between about 10 μ g and about 150 μ g per actuation, and between about 15 μ g and about 80 μ g per actuation.
6. The pharmaceutical composition according to claim 3, wherein the concentration of glycopyrrolate included in the co-suspension is between about 0.04 mg/ml and about 2.25 mg/ml.
7. The pharmaceutical composition according to claim 3, wherein at least 90% of the glycopyrrolate active agent particles by volume exhibit an optical diameter of 7 μ m or less.

8. The pharmaceutical composition according to claim 3, wherein at least 50% of the glycopyrrolate active agent particles by volume exhibit an optical diameter of 5 μm or less.

9. The pharmaceutical composition according to claim 2, wherein the suspending particles comprise perforated microstructures.

10. The pharmaceutical composition according to claim 9, wherein the perforated microstructures are prepared using a spray drying process.

11. The pharmaceutical composition according to claim 10, wherein the perforated microstructures comprise a spray dried emulsion of perfluoroctyl bromide, DSPC and calcium chloride in water.

12. The pharmaceutical composition according to claim 2, wherein the suspending particles comprise an excipient selected from at least one of lipids, phospholipids, nonionic detergents, polymers, nonionic block copolymers, surfactants, non-ionic surfactants, biocompatible fluorinated surfactants, carbohydrates, amino acids, organic salts, peptides, proteins, alditols, and combinations thereof.

13. The pharmaceutical composition according to claim 2, wherein the suspending particles exhibit an MMAD selected from between about 10 μm and about 500 nm, between about 5 μm and about 750 nm, between about and 1 μm and about 3 μm .

14. The pharmaceutical composition according to claim 2, wherein the suspending particles exhibit a volume median optical diameter selected from between about 0.2 μm and about 50 μm , between about 0.5 μm and about 15 μm , between about 1.5 μm and about 10 μm , and between about 2 μm and about 5 μm .

15. The pharmaceutical composition according to claim 2, wherein the propellant comprises a propellant selected from an HFA propellant, a PFC propellant and combinations thereof, and wherein the propellant is substantially free of additional constituents.

16. The pharmaceutical composition according to claim 2, wherein a total mass of the suspending particles exceeds a total mass of the active agent particles.

17. The pharmaceutical composition according to claim 16, wherein a ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from above about 1.5, up to about 5, up to about 10, up to about

15, up to about 17, up to about 20, up to about 30, up to about 40, up to about 50, up to about 60, up to about 75, up to about 100, up to about 150, and up to about 200.

18. The pharmaceutical composition according to claim 16, wherein a ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

19. The pharmaceutical composition according to claim 1, wherein the suspending particles remain associated with the active agent particles even when subjected to buoyancy forces amplified by centrifugation at an acceleration selected from accelerations of at least 1 g, at least 10 g, at least 50 g, and at least 100 g.

20. A method for treating a pulmonary disease or disorder in a patient, the method comprising:

providing metered dose inhaler comprising a pharmaceutically acceptable co-suspension, the co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable propellant;

a plurality of active agent particles comprising an active agent selected from a LAMA active agent and a LABA active agent; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and administering the co-suspension to the patient by actuating the metered dose inhaler, wherein said administering of the co-suspension composition comprises delivering a therapeutically effective amount of the LAMA or LABA active agent to the patient.

21. The method of claim 20, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising a LAMA active agent.

22. The method of claim 21, wherein the LAMA active agent is selected from glycopyrrolate, dexipirronium, tiotropium, trospium, aclidinium, darotropium, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

23. The method of claim 22, wherein the pulmonary disease or disorder is selected from at least one of asthma, COPD, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress

syndrome, pulmonary hypertension, pulmonary vasoconstriction, pulmonary inflammation associated with cystic fibrosis, and pulmonary obstruction associated with cystic fibrosis.

24. The method of claim 23, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising glycopyrrolate, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

25. The method of claim 24, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising glycopyrrolate, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, and the concentration of glycopyrrolate included in the co-suspension is between about 0.04 mg/ml and about 2.25 mg/ml

26. The method of claim 22, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in FEV₁ in the patient within 1 hour, or less.

27. The method of claim 22, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in FEV₁ in the patient within 0.5 hours, or less.

28. The method of claim 22, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in an increase of FEV₁ of 150 ml or greater within a period of time selected from 0.5 hours, or less, 1 hour, or less, and 1.5 hours, or less.

29. The method of claim 22, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in FEV₁ in the patient within 0.5 hours, or less, and providing a clinically significant increase in FEV₁ for a time period selected from up to 4 hours, up to 6 hours, up to 8 hours, up to 10 hours, and up to 12 hours, or more.

30. The method of claim 23, wherein administering the pharmaceutically acceptable co-suspension comprises administering to the patient a dose of LAMA

active agent selected from 200 µg, or less, 150 µg, or less, 100 µg, or less, 75 µg, or less, 50 µg, or less, and 25 µg, or less, per actuation of the metered dose inhaler.

31. The method of claim 23, wherein administering the pharmaceutically acceptable co-suspension comprises administering to the patient a dose of glycopyrrolate or a pharmaceutically acceptable salt, ester, isomer or solvate thereof, selected from about 150 µg, or less, about 80 µg, or less, about 40 µg, about 20 µg, or less, and about 10 µg, or less, per actuation of the metered dose inhaler.

32. The method of claim 26, wherein administering the pharmaceutically acceptable co-suspension comprises administering to the patient a dose of LAMA active agent selected from 200 µg, or less, 150 µg, or less, 100 µg, or less, 75 µg, or less, 50 µg, or less, and 25 µg, or less, per actuation of the metered dose inhaler.

33. The method of claim 26, wherein administering the pharmaceutically acceptable co-suspension comprises administering to the patient a dose of glycopyrrolate or a pharmaceutically acceptable salt, ester, isomer or solvate thereof, selected from about 150 µg, or less, about 80 µg, or less, about 40 µg, about 20 µg, or less, and about 10 µg, or less, per actuation of the metered dose inhaler.

34. The method of claim 22, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in inspiratory capacity.

35. A method for respiratory delivery of a LAMA or LABA active agent to a patient, the method comprising;

providing metered dose inhaler comprising a canister containing a pharmaceutically acceptable co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable propellant;

a plurality of active agent particles comprising an active agent selected from a LAMA active agent and a LABA active agent; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and actuating the metered dose inhaler to provide respiratory delivery of the LAMA or LABA active agent to the patient.

36. The method of claim 35, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising a LAMA active agent.

37. The method of claim 36, wherein the LAMA active agent is selected from glycopyrrolate, dexipirronium, tiotropium, trospium, aclidinium, darotropium, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

38. The method of claim 37, wherein the LAMA active agent is glycopyrrolate, including pharmaceutically acceptable salts, esters, isomers or solvates thereof.

39. The method of claim 36, wherein actuating the metered dose inhaler to provide respiratory delivery of the LAMA active agent comprises delivering the LAMA active agent to the patient at a DDU selected from a DDU of $\pm 30\%$, or better, a DDU of $\pm 25\%$, or better, and a DDU of $\pm 20\%$, or better, throughout emptying of the canister.

40. The method of claim 36, wherein actuating the metered dose inhaler to provide respiratory delivery of the LAMA active agent comprises delivering the LAMA active agent at an initial fine particle fraction and the initial fine particle fraction delivered from the metered dose inhaler is substantially maintained, such that, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 80% of the initial fine particle fraction.

41. The method of claim 40, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 90% of the initial fine particle fraction.

42. The method of claim 40, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 95% of the initial fine particle fraction.

43. A pharmaceutical composition according to claim 1, wherein the active agent included in the active agent particles is a LABA active agent selected from bambuterol, clenbuterol, formoterol, salmeterol, carmoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantyl-derived β_2 agonists, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

44. The pharmaceutical composition according to claim 43, wherein the active agent particles comprise formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

45. The pharmaceutical composition according to claim 44, wherein the active agent particles comprise crystalline formoterol.

46. The pharmaceutical composition according to claim 44, wherein the formoterol active agent particles are included in the composition at a concentration sufficient to provide a delivered dose of formoterol selected from between about 1 μ g and about 30 μ g, between about 0.5 μ g and about 10 μ g, between about 2 μ g and 5 μ g, between about 2 μ g and about 10 μ g, between about 5 μ g and about 10 μ g, and between 3 μ g and about 30 μ g per actuation of the metered dose inhaler.

47. The pharmaceutical composition according to claim 44, wherein the formoterol active agent particles are included in the composition at a concentration sufficient to provide a delivered dose of formoterol selected from up to about 30 μ g, up to about 10 μ g, up to about 5 μ g, up to about 2.5 μ g, up to about 2 μ g, or up to about 1.5 μ g per actuation of the metered dose inhaler.

48. The pharmaceutical composition according to claim 44, wherein the concentration of formoterol included in the co-suspension is selected from between about 0.01 mg/ml and about 1 mg/ml, between about 0.01 mg/ml and about 0.5 mg/ml, and between about 0.03 mg/ml and about 0.4 mg/ml.

49. The pharmaceutical composition according to claim 44, wherein at least 90% of the formoterol active agent particles by volume exhibit an optical diameter of 5 μ m or less.

50. The pharmaceutical composition according to claim 44, wherein at least 50% of the formoterol active agent particles by volume exhibit an optical diameter of 2 μ m or less.

51. The pharmaceutical composition according to claim 43, wherein the suspending particles comprise perforated microstructures.

52. The pharmaceutical composition according to claim 51, wherein the perforated microstructures are prepared using a spray drying process.

53. The pharmaceutical composition according to claim 52, wherein the perforated microstructures comprise a spray dried emulsion of perfluoroctyl bromide, DSPC and calcium chloride in water.

54. The pharmaceutical composition according to claim 43, wherein the suspending particles comprise an excipient selected from at least one of lipids, phospholipids, nonionic detergents, polymers, nonionic block copolymers, surfactants, non-ionic surfactants, biocompatible fluorinated surfactants, carbohydrates, amino acids, organic salts, peptides, proteins, alditols, and combinations thereof.

55. The pharmaceutical composition according to claim 43, wherein the suspending particles exhibit an MMAD selected from between about 10 μm and about 500 nm, between about 5 μm and about 750 nm, and 1 μm and about 3 μm .

56. The pharmaceutical composition according to claim 43, wherein the suspending particles exhibit a volume median optical diameter selected from between about 0.2 μm and about 50 μm , between about 0.5 μm and about 15 μm , between about 1.5 μm and about 10 μm , and between about 2 μm and about 5 μm .

57. The pharmaceutical composition according to claim 43, wherein the propellant comprises a propellant selected from an HFA propellant, a PFC propellant and combinations thereof, and wherein the propellant is substantially free of additional constituents.

58. The pharmaceutical composition according to claim 43, wherein a total mass of the suspending particles exceeds a total mass of the active agent particles.

59. The pharmaceutical composition according to claim 58, wherein a ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from above about 1.5, up to about 5, up to about 10, up to about 15, up to about 17, up to about 20, up to about 30, up to about 40, up to about 50, up to about 60, up to about 75, up to about 100, up to about 150, and up to about 200.

60. The pharmaceutical composition according to claim 58, wherein a ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

61. The pharmaceutical composition according to claim 43, wherein the suspending particles remain associated with the active agent particles even when subjected to buoyancy forces amplified by centrifugation at an acceleration selected from accelerations of at least 1 g, at least 10 g, at least 50 g, and at least 100 g.

62. The method of claim 20, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising a LABA active agent.

63. The method of claim 62, wherein the LABA active agent is selected from bambuterol, clenbuterol, formoterol, salmeterol, carmoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantly-derived β_2 agonists, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

64. The method of claim 63, wherein the pulmonary disease or disorder is selected from at least one of asthma, COPD, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, pulmonary hypertension, pulmonary vasoconstriction, pulmonary inflammation associated with cystic fibrosis, and pulmonary obstruction associated with cystic fibrosis.

65. The method of claim 63, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

66. The method of claim 65, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, and the concentration of formoterol included in the co-suspension is selected from between about 0.01 mg/ml and about 1 mg/ml, between about 0.01 mg/ml and about 0.5 mg/ml, and between about 0.03 mg/ml and about 0.4 mg/ml.

67. The method of claim 63, wherein administering the pharmaceutical composition comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in FEV₁ in the patient within 1 hour, or less.

68. The method of claim 63, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in FEV₁ in the patient within 0.5 hours, or less.

69. The method of claim 63, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in an increase of FEV₁ of 150 ml or greater within a period of time selected from 0.5 hours, or less, 1 hour, or less, and 1.5 hours, or less.

70. The method of claim 63, wherein administering the pharmaceutically acceptable co-suspension comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in FEV₁ in the patient within 0.5 hours, or less, and providing a clinically significant increase in FEV₁ for a time period selected from up to 4 hours, up to 6 hours, and up to 8 hours, or more.

71. The method of claim 67, wherein administering the pharmaceutically acceptable co-suspension comprises administering to the patient a dose of LABA active agent selected from between about 1 µg and about 50 µg, between about 1 µg and about 30 µg, between about 2 µg and 5 µg, between about 2 µg and about 10 µg, between about 5 µg and about 10 µg, and between 3 µg and about 30 µg per actuation.

72. The method of claim 67, wherein administering the pharmaceutically acceptable co-suspension comprises administering to the patient a dose of formoterol or a pharmaceutically acceptable salt, ester, isomer or solvate thereof, selected from up to about 30 µg, up to about 10 µg, up to about 5 µg, up to about 2.5 µg, up to about 2 µg, and up to about 1.5 µg per actuation.

73. The method of claim 63, wherein administering the pharmaceutical composition comprises administering the pharmaceutical composition in an amount resulting in a clinically significant increase in inspiratory capacity.

74. The method of claim 35, wherein providing a pharmaceutically acceptable co-suspension comprises providing a co-suspension comprising a plurality of active agent particles comprising a LABA active agent.

75. The method of claim 74, wherein the LABA active agent is selected from bambuterol, clenbuterol, formoterol, salmeterol, carmoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantly-derived β₂ agonists, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

76. The method of claim 75, wherein the LABA active agent is formoterol, including pharmaceutically acceptable salts, esters, isomers or solvates thereof.

77. The method of claim 75, wherein actuating the metered dose inhaler to provide respiratory delivery of the LABA active agent comprises delivering the LABA active agent to the patient at a DDU selected from a DDU of $\pm 30\%$, or better, a DDU of $\pm 25\%$, or better, and a DDU of $\pm 20\%$, or better, throughout emptying of the canister.

78. The method of claim 75, wherein actuating the metered dose inhaler to provide respiratory delivery of the LABA active agent comprises delivering the LABA active agent at an initial fine particle fraction and the initial fine particle fraction delivered from the metered dose inhaler is substantially maintained, such that, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 80% of the initial fine particle fraction.

79. The method of claim 78, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 90% of the initial fine particle fraction.

80. The method of claim 79, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 95% of the initial fine particle fraction.

81. A metered dose inhaler for delivery of a LAMA or a LABA active agent, the metered dose inhaler comprising:

a canister containing a pharmaceutically acceptable co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable propellant;

a plurality of active agent particles comprising an active agent selected from a LAMA active agent and a LABA active agent; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles;

wherein the LAMA or LABA active agent included in the pharmaceutically acceptable co-suspension is chemically stable over a period of at least 18 months when stored at 5°C.

82. The metered dose inhaler according to claim 81, wherein the LAMA or LABA active agent included in the pharmaceutically acceptable co-suspension is chemically stable over a period of at least 18 months when stored at 25°C.

83. The metered dose inhaler according to claim 81, wherein the LAMA or LABA active agent included in the pharmaceutically acceptable co-suspension is a LABA active agent selected from bambuterol, clenbuterol, formoterol, salmeterol, carmoterol, milveterol, indacaterol, and saligenin- or indole- containing and adamantyl-derived β_2 agonists, and any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

84. The metered dose inhaler according to claim 83 wherein the LABA active agent is formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof.

85. The metered dose inhaler according to claim 84, wherein a rate of formation of N-(2-hydroxy-5-(1-(2-hydroxy-5-(1-hydroxy-2-(1-(4-methoxyphenyl)propan-2-ylamino)ethyl)phenylamino)-2-(1-(4-methoxyphenyl)propan-2-ylamino)ethyl)phenyl)acetamide within the pharmaceutically acceptable co-suspension is not greater than about 0.15%, after the canister is subjected to a temperature of 40°C and a relative humidity of 75% for a period of one month.

86. The metered dose inhaler according to claim 84, wherein a rate of formation of N-(2-hydroxy-5-(1-(2-hydroxy-5-(1-hydroxy-2-(1-(4-methoxyphenyl)propan-2-ylamino)ethyl)phenylamino)-2-(1-(4-methoxyphenyl)propan-2-ylamino)ethyl)phenyl)acetamide within the pharmaceutically acceptable co-suspension is not greater than about 0.5%, after the canister is subjected to a temperature of 40°C and a relative humidity of 75% for a period of one month.

87. A pharmaceutical composition deliverable from a metered dose inhaler, comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising glycopyrrolate, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, said active agent particles are included in the suspension medium at a concentration sufficient to provide a delivered dose of glycopyrrolate of between about 15 μ g and about 80 μ g per actuation of the metered dose inhaler; and

a plurality of respirable suspending particles comprising perforated microstructures exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , said perforated microstructures associate with the plurality of active agent particles to form a co-suspension.

88. The pharmaceutical composition according to claim 87, wherein a ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

89. A pharmaceutical composition deliverable from a metered dose inhaler, comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising formoterol, including any pharmaceutically acceptable salts, esters, isomers or solvates thereof, said active agent particles are included in the suspension medium at a concentration sufficient to provide a delivered dose of formoterol of between about 2 μg and about 10 μg per actuation of the metered dose inhaler; and

a plurality of respirable suspending particles comprising perforated microstructures exhibiting a volume median optical diameter of between about 1.5 μm and about 10 μm , said perforated microstructures associate with the plurality of active agent particles to form a co-suspension.

90. The pharmaceutical composition according to claim 89, wherein a ratio of the total mass of the suspending particles to the total mass of the active agent particles is selected from between about 3:1 and about 15:1 and between about 2:1 and 8:1.

91. A method for respiratory delivery of LAMA active agent to a patient, the method comprising;

providing a metered dose inhaler comprising a canister containing a pharmaceutically acceptable co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising glycopyrrolate, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and actuating the metered dose inhaler to provide respiratory delivery of glycopyrrolate to the patient at a DDU of \pm 20%, or better, throughout emptying of the canister.

92. The method of claim 91, wherein actuating the metered dose inhaler to provide respiratory delivery of glycopyrrolate to the patient comprises delivering the glycopyrrolate at an initial fine particle fraction and the initial fine particle fraction delivered from the metered dose inhaler is substantially maintained, such that, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 80% of the initial fine particle fraction.

93. The method of claim 92, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 90% of the initial fine particle fraction.

94. The method of claim 93, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 95% of the initial fine particle fraction.

95. A method for respiratory delivery of LABA active agent to a patient, the method comprising;

providing a metered dose inhaler comprising a canister containing a pharmaceutically acceptable co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising formoterol, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and

actuating the metered dose inhaler to provide respiratory delivery of formoterol to the patient at a DDU of \pm 20%, or better, throughout emptying of the canister.

96. The method of claim 95, wherein actuating the metered dose inhaler to provide respiratory delivery of formoterol to the patient comprises delivering the formoterol at an initial fine particle fraction and the initial fine particle fraction

delivered from the metered dose inhaler is substantially maintained, such that, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 80% of the initial fine particle fraction.

97. The method of claim 96, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 90% of the initial fine particle fraction.

98. The method of claim 97, wherein, throughout emptying of the canister, the fine particle fraction delivered from the metered dose inhaler is maintained within 95% of the initial fine particle fraction.

99. A method for treating a pulmonary disease or disorder in a patient, the method comprising:

providing metered dose inhaler comprising a pharmaceutically acceptable co-suspension, the co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising glycopyrrolate, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and

administering the co-suspension to the patient by actuating the metered dose inhaler, wherein said administering of the co-suspension composition comprises delivering a dose of 150 μ g, or less, of glycopyrrolate per actuation of the metered dose inhaler and results in a clinically significant increase in FEV₁ in the patient.

100. The method of claim 99, wherein the pulmonary disease or disorder is selected from at least one of asthma, COPD, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, pulmonary hypertension, pulmonary vasoconstriction, pulmonary inflammation associated with cystic fibrosis, and pulmonary obstruction associated with cystic fibrosis.

101. A method for treating a pulmonary disease or disorder in a patient, the method comprising:

providing metered dose inhaler comprising a pharmaceutically acceptable co-suspension, the co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising formoterol, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and administering the co-suspension to the patient by actuating the metered dose inhaler, wherein said administering of the co-suspension composition comprises delivering a dose of 10 μ g, or less, of formoterol per actuation of the metered dose inhaler and results in a clinically significant increase in FEV₁ in the patient.

102. The method of claim 101, wherein the pulmonary disease or disorder is selected from at least one of asthma, COPD, allergic rhinitis, sinusitis, pulmonary vasoconstriction, inflammation, allergies, impeded respiration, respiratory distress syndrome, pulmonary hypertension, pulmonary vasoconstriction, pulmonary inflammation associated with cystic fibrosis, and pulmonary obstruction associated with cystic fibrosis.

103. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 μ g.

104. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 μ g.

105. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 μ g.

106. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 μ g, and said administration results in an increase in FEV₁ of at least 150 mL within 0.5 hours, or less.

107. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 μ g, and said administration results in an increase in FEV₁ of at least 150 mL within 0.5 hours, or less.

108. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg, and said administration results in an increase in FEV₁ of at least 150 mL within 0.5 hours, or less

109. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg, and said administration results in an increase in FEV₁ of at least 200 mL within 1.0 hour, or less.

110. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 µg, and said administration results in an increase in FEV₁ of at least 200 mL within 1.0 hour, or less.

111. The method of claim 99, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg, and said administration results in an increase in FEV₁ of at least 200 mL within 1.0 hour, or less.

112. A method for treating a pulmonary disease or disorder in a patient, the method comprising:

providing metered dose inhaler comprising a pharmaceutically acceptable co-suspension, the co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising glycopyrrolate, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and administering the co-suspension to the patient by actuating the metered dose inhaler, wherein said administering of the co-suspension composition comprises delivering a dose of 150 µg, or less, of glycopyrrolate per actuation of the metered dose inhaler and results clinically significant increase in FEV₁ in 0.5 hours, or less, and a clinically significant increase in FEV₁ is maintained for up to 12 hours.

113. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg.

114. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 µg.

115. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg.

116. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg, and said administration results in an increase in FEV₁ of at least 150 mL within 0.5 hours, or less, and a clinically significant increase in FEV₁ is maintained for at least 12 hours.

117. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 µg, and said administration results in an increase in FEV₁ of at least 150 mL within 0.5 hours, or less, and a clinically significant increase in FEV₁ is maintained for at least 10 hours.

118. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg, and said administration results in an increase in FEV₁ of at least 150 mL within 0.5 hours, or less, and a clinically significant increase in FEV₁ is maintained for at least 8 hours.

119. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg, and said administration results in an increase in FEV₁ of at least 200 mL within 1.0 hour, or less, and a clinically significant increase in FEV₁ is maintained for at least 12 hours.

120. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 µg, and said administration results in an increase in FEV₁ of at

least 200 mL within 1.0 hour, or less, and a clinically significant increase in FEV₁ is maintained for at least 10 hours.

121. The method of claim 112, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg, and said administration results in an increase in FEV₁ of at least 200 mL within 1.0 hour, or less, and a clinically significant increase in FEV₁ is maintained for at least 8 hours.

122. A method for treating a pulmonary disease or disorder in a patient population, the method comprising:

providing metered dose inhalers comprising a pharmaceutically acceptable co-suspension, the co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising glycopyrrolate, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and

administering the co-suspension to the patient population by actuating the metered dose inhaler, wherein said administering of the co-suspension composition comprises delivering a dose of 150 µg, or less, of glycopyrrolate per actuation of the metered dose inhaler and, in at least 50% of the patient population, results in an increase from baseline in FEV₁ selected from (i) an increase in baseline FEV₁ of at least 200 ml and (ii) a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml.

123. The method of claim 122, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg.

124. The method of claim 122, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 µg.

125. The method of claim 122, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg.

126. The method of claim 122, wherein, for at least 60% of the patient population, said administering of the co-suspension composition results in an increase from baseline in FEV₁ selected from either (i) an increase in baseline FEV₁ of at least 200 ml and (ii) a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml.

127. The method of claim 122, wherein, for at least 80% of the patient population, said administering of the co-suspension composition results in an increase from baseline in FEV₁ selected from either (i) an increase in baseline FEV₁ of at least 200 ml and (ii) a 12%, or greater, increase from baseline in FEV₁ coupled with total increase in FEV₁ of at least 150 ml.

128. A method for treating a pulmonary disease or disorder in a patient, the method comprising:

providing metered dose inhaler comprising a pharmaceutically acceptable co-suspension, the co-suspension comprising:

a suspension medium comprising a pharmaceutically acceptable HFA propellant;

a plurality of active agent particles comprising glycopyrrolate, including pharmaceutically acceptable salts, esters, isomers or solvates thereof; and

a plurality of respirable suspending particles, wherein the plurality of active agent particles associate with the plurality of suspending particles; and

administering the co-suspension to the patient by actuating the metered dose inhaler, wherein said administering of the co-suspension composition comprises delivering a dose of 150 µg, or less, of glycopyrrolate per actuation of the metered dose inhaler and results in a clinically significant increase in inspiratory capacity (IC) in the patient.

129. The method of claim 128, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg.

130. The method of claim 128, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 100 µg.

131. The method of claim 128, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 80 µg.

132. The method of claim 128, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg, and said administration results in an increase in IC of at least 300 ml within a period of time selected from 1 hour or less and 2 hours or less.

133. The method of claim 128, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg, and said administration results in an increase in IC of at least 150 ml within a period of time selected from 1 hour or less and 2 hours or less.

134. The method of claim 128, wherein said administering of the co-suspension composition comprises administering a delivered dose of glycopyrrolate of no more than 150 µg, and said administration results in an increase in IC of at least 100 ml within a period of time selected from 1 hour or less and 2 hours or less.

1/30




FIG. 1

2/30




FIG. 2

3/30




FIG. 3

4/30




FIG. 4

5/30




FIG. 5

6/30




FIG. 6

7/30




FIG. 7

8/30




FIG. 8

9/30




FIG. 9

10/30




FIG. 10

11/30




FIG. 11

12/30




FIG. 12

13/30




FIG. 13

14/30




FIG. 14

15/30




FIG. 15

16/30




FIG. 16

17/30




FIG. 17

18/30




FIG. 18

19/30




FIG. 19

20/30




FIG. 20

21/30




FIG. 21

22/30




FIG. 22

23/30




FIG. 23

24/30




FIG. 24

25/30




FIG. 25

26/30




FIG. 26

27/30




FIG. 27

28/30




FIG. 28

29/30




FIG 29

30/30




FIG. 30