



HU000030311T2

(19) **HU**(11) Lajstromszám: **E 030 311**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 12 772808**
(22) A bejelentés napja: **2012. 08. 24.**

(51) Int. Cl.: **C07D 471/08** (2006.01)
A61K 314/39 (2006.01)
A61P 31/04 (2006.01)

(96) Az európai bejelentés bejelentési száma:
EP 20120772808

(97) Az európai bejelentés közzétételi adatai:
EP 2748165 A1 **2013. 03. 07.**

(86) A nemzetközi (PCT) bejelentési szám:
PCT/IB 12/054290

(97) Az európai szabadalom megadásának meghirdetési adatai:
EP 2748165 B1 **2016. 11. 02.**

(87) A nemzetközi közzétételi szám:
WO 13030733

(30) Elsőbbségi adatok:
MM24122011 **2011. 08. 27.** **IN**

(73) Jogosult(ak):
Wockhardt Limited, Aurangabad 431210 (IN)

(72) Feltaláló(k):
PATEL, Mahesh Vithalbhai, Aurangabad 431 003 (IN)
DESHPANDE, Prasad Keshav, Aurangabad 431003 (IN)
BHAWSAR, Satish, Aurangabad 431005 (IN)
BHAGWAT, Sachin, Aurangabad (IN)
JAFRI, Mohammad Alam, Aurangabad 431001 (IN)
MISHRA, Amit, Lucknow Uttar Pradesh 226010 (IN)
PAVASE, Laxmikant, Ahmednagar 414005 (IN)
GUPTA, Sunil, Kota 324002 (IN)
KALE, Rajesh, Amravati 444606 (IN)
JOSHI, Sanjeev, Aurangabad (IN)

(74) Képviselő:
Danubia Szabadalmi és Jogi Iroda Kft.,
Budapest

(54) **1,6-Diazabiciklo[3.2.1]oktán-7-on-származékok és alkalmazásuk bakteriális fertőzések kezelésére**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 748 165 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
02.11.2016 Bulletin 2016/44

(51) Int Cl.:
C07D 471/08 ^(2006.01) **A61K 31/439** ^(2006.01)
A61P 31/04 ^(2006.01)

(21) Application number: **12772808.7**

(86) International application number:
PCT/IB2012/054290

(22) Date of filing: **24.08.2012**

(87) International publication number:
WO 2013/030733 (07.03.2013 Gazette 2013/10)

(54) **1,6-DIAZABICYCLO[3,2,1]OCTAN-7-ONE DERIVATIVES AND THEIR USE IN THE TREATMENT OF BACTERIAL INFECTIONS.**

1,6-DIAZABICYCLO-[3,2,1]-OCTAN-7-ON-DERIVATE UND IHRE VERWENDUNG ZUR
BEHANDLUNG BAKTERIELLER INFEKTIONEN

DÉRIVÉS 1,6-DIAZABICYCLO[3,2,1]OCTANE-7-ONE ET LEUR UTILISATION DANS LE
TRAITEMENT D'INFECTIONS BACTÉRIENNES

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **27.08.2011 IN MM24122011**

(43) Date of publication of application:
02.07.2014 Bulletin 2014/27

(73) Proprietor: **Wockhardt Limited
Aurangabad 431210 (IN)**

(72) Inventors:
• **PATEL, Mahesh Vithalbhai
Aurangabad 431 003 (IN)**
• **DESHPANDE, Prasad Keshav
Aurangabad 431003 (IN)**
• **BHAWSAR, Satish
Aurangabad 431005 (IN)**
• **BHAGWAT, Sachin
Aurangabad (IN)**
• **JAFRI, Mohammad Alam
Aurangabad 431001 (IN)**

- **MISHRA, Amit
Lucknow
Uttar Pradesh 226010 (IN)**
- **PAVASE, Laxmikant
Ahmednagar 414005 (IN)**
- **GUPTA, Sunil
Kota 324002 (IN)**
- **KALE, Rajesh
Amravati 444606 (IN)**
- **JOSHI, Sanjeev
Aurangabad (IN)**

(74) Representative: **Nederlandsch Octrooibureau
P.O. Box 29720
2502 LS The Hague (NL)**

(56) References cited:
WO-A2-2009/091856 US-A1- 2010 093 784

Remarks:

The file contains technical information submitted after
the application was filed and not included in this
specification

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 748 165 B1

Description**FIELD OF THE INVENTION**

[0001] The invention relates to nitrogen containing compounds, their preparation and their use in preventing and/or treating bacterial infections.

BACKGROUND OF THE INVENTION

[0002] Emergence of bacterial resistance to known antibacterial agents is becoming a major challenge in treating bacterial infections. One way forward to treat bacterial infections, and especially those caused by resistant bacteria, is to develop newer antibacterial agents that can overcome the bacterial resistance. Coates et al. (Br. J. Pharmacol. 2007; 152(8), 1147-1154.) have reviewed novel approaches to developing new antibiotics. However, the development of new antibacterial agents is a challenging task. For example, Gwynn et al. (Annals of the New York Academy of Sciences, 2010, 1213: 5-19) have reviewed the challenges in the discovery of antibacterial agents.


[0003] Several antibacterial agents have been described in the prior art (for example, see PCT International Application Nos. PCT/US2010/060923, PCT/EP2010/067647, PCT/US2010/052109, PCT/US2010/048109, PCT/GB2009/050609, PCT/EP2009/056178 and PCT/US2009/041200). However, there remains a need for potent antibacterial agents for preventing and/or treating bacterial infections, including those caused by bacteria that are resistant to known antibacterial agents. WO2009/091856 and US2010/0093784 disclose bicyclic heterocycles for the treatment of bacterial infections.

[0004] The inventors have surprisingly discovered nitrogen containing compounds with antibacterial properties.

SUMMARY OF THE INVENTION

[0005] Accordingly there are provided nitrogen containing compounds, methods for preparation of these compounds, pharmaceutical compositions comprising these compounds, and also disclosed is a method for preventing or treating bacterial infection in a subject using these compounds.

[0006] In one general aspect, there are provided compounds of Formula (I):



or a stereoisomer or a pharmaceutically acceptable salt thereof; wherein:

R₁ is:

- (a) hydrogen,
- (b) (CO)_n-R₃,
- (c) COOR₄, or
- (d) COCH₂COR₃

n is 0, 1 or 2;

R₂ is:

- (a) SO₃M,
- (b) SO₂NH₂,
- (c) PO₃M,
- (d) CH₂COOM,
- (e) CF₂COOM,
- (f) CHFCOOM, or
- (g) CF₃;

M is hydrogen or a cation;

R₃ is:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, NR₅CONR₆R₇, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) CN,
- (d) NR₆R₇,
- (e) CONR₆R₇,
- (f) NHCONR₆R₇,
- (g) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (h) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, NHC(NH)NR₆R₇, or NHCONR₆R₇,
- (i) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (j) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (k) cycloalkyl substituted with C₁-C₆ alkyl wherein C₁-C₆ alkyl is further substituted with one or more substituents independently selected from OR₅, NR₆R₇, halogen, CN, or CONR₆R₇, or (1) OR₈;

R₄ is:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (d) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (e) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇, or
- (f) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇;

R₅ and R₈ are each independently:

- (a) hydrogen, or
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, CN, CONR₆R₇, NR₆R₇, heterocyclyl, heteroaryl, cycloalkyl or aryl;

R₆ and R₇ are each independently:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅, CN, COOR₅, CONR₅R₈, NR₅R₈, NR₅COR₈, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (d) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (e) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (f) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (g) R₆ and R₇ are joined together to form a four to seven member ring.

[0007] In another general aspect, there are provided pharmaceutical compositions comprising a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0022] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable derivative thereof.

[0023] Also disclosed is a method for preventing or treating bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) at least one antibacterial agent or a pharmaceutically acceptable derivative thereof.

[0024] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) at least one antibacterial agent or a pharmaceutically acceptable derivative thereof.

[0025] Also disclosed is a method for preventing or treating bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, (b) at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable derivative thereof, and (c) at least one antibacterial agent or a pharmaceutically acceptable derivative thereof.

[0026] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, (b) at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable derivative thereof, and (c) at least one antibacterial agent or a pharmaceutically acceptable derivative thereof.

[0027] Also disclosed is methods for increasing antibacterial effectiveness of a antibacterial agent in a subject, said method comprising co-administering said antibacterial agent or a pharmaceutically acceptable derivative thereof with a pharmaceutically effective amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0028] The details of one or more embodiments of the invention are set forth in the description below.

DETAILED DESCRIPTION OF THE INVENTION

[0029] It must be noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content clearly dictates otherwise.

[0030] The inventors have surprisingly discovered novel nitrogen containing compounds having antibacterial properties.

[0031] The term " C_1 - C_6 alkyl" as used herein refers to branched or unbranched acyclic hydrocarbon radical with 1 to 6 carbon atoms. Typical, non-limiting examples of " C_1 - C_6 alkyl" include methyl, ethyl, *n*-propyl, *iso*-propyl, *n*-butyl, *iso*-butyl, *tert*-butyl, *n*-pentyl, *iso*-pentyl, *n*-hexyl and the like. The " C_1 - C_6 alkyl" may be unsubstituted, or substituted with one or more substituents. Typical, non-limiting examples of such substituents include halogen, alkoxy, CN, COOH, CONH₂, OH, -NH₂, -NHCOCH₃, cycloalkyl, heterocyclyl, heteroaryl, aryl and the like.

[0032] The term "cycloalkyl" as used herein refers to three to seven member cyclic hydrocarbon radicals. The cycloalkyl group optionally incorporates one or more double or triple bonds, or a combination of double bonds and triple bonds, but which is not aromatic. Typical, non-limiting examples of cycloalkyl groups include cyclopropane, cyclobutane, cyclopentane, cyclohexane, and cycloheptane. The cycloalkyl may be unsubstituted, or substituted with one or more substituents. Typical, non-limiting examples of such substituents include C_1 - C_6 alkyl, halogen, alkoxy, CN, COOH, CONH₂, OH, NH₂, NHCOCH₃, heterocyclyl, heteroaryl, aryl, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, -OSO₂-aryl and the like.

[0033] The term "heterocyclyl" as used herein refers to four to seven member cycloalkyl group containing one or more heteroatoms selected from nitrogen, oxygen or sulfur. The heterocycloalkyl group optionally incorporates one or more double or triple bonds, or a combination of double bonds and triple bonds, but which is not aromatic. Typical, non-limiting examples of heterocycloalkyl groups include azetidine, pyrrolidine, 2-oxo-pyrrolidine, imidazolidin-2-one, piperidine, oxazine, thiazine, piperazine, piperazin-2,3-dione, morpholine, thiamorpholine, azapane, and the like. The heterocycloalkyl may be unsubstituted, or substituted with one or more substituents. Typical, non-limiting examples of such substituents include C_1 - C_6 alkyl, halogen, alkoxy, CN, COOH, CONH₂, OH, NH₂, NHCOCH₃, heterocyclyl, heteroaryl, aryl, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl and the like.

[0034] The term "aryl" as used herein refers to a monocyclic or polycyclic aromatic hydrocarbon. Typical, non-limiting examples of aryl groups include phenyl, naphthyl, anthracenyl, fluorenyl, phenanthrenyl, and the like. The aryl group may be unsubstituted, or substituted with one or more substituents. Typical, non-limiting examples of such substituents

include C₁-C₆ alkyl, halogen, alkoxy, CN, COOH, CONH₂, OH, NH₂, NHCOCH₃, heterocyclyl, heteroaryl, aryl, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl and the like.

[0035] The term "heteroaryl" as used herein refers to a monocyclic or polycyclic aromatic hydrocarbon group wherein one or more carbon atoms have been replaced with heteroatoms selected from nitrogen, oxygen, and sulfur. If the heteroaryl group contains more than one heteroatom, the heteroatoms may be the same or different. Typical, non-limiting example of heteroaryl groups include 1,2,4-oxadiazol, 1,3,4-oxadiazol, 1,3,4-thiadiazol, 1,2,3,4-tetrazol, 1,3-oxazol, 1,3-thiazole, pyridine, pyrimidine, pyrazine, pyridazine, furan, pyrrol, thiophene, imidazole, pyrazole, benzofuran, benzothio-
 5 phenene, benzimidazole, benzoxazole, benzothiazole, thiazole, and the like. The heteroaryl group may be unsubstituted, or substituted with one or more substituents. Typical, non-limiting examples of such substituents include C₁-C₆ alkyl,
 10 halogen, alkoxy, CN, COOH, CONH₂, OH, NH₂, NHCOCH₃, heterocyclyl, heteroaryl, aryl, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl and the like.

[0036] The term "stereoisomers" as used herein refers to compounds that have identical chemical constitution, but differ with regard to the arrangement of their atoms or groups in space. The compounds of Formula (I) may contain asymmetric or chiral centers and, therefore, exist in different stereoisomeric forms. It is intended, unless specified
 15 otherwise, that all stereoisomeric forms of the compounds of Formula (I) as well as mixtures thereof, including racemic mixtures, form part of the present invention. In addition, all geometric and positional isomers (including *cis* and *trans*-forms) as well as mixtures thereof, are also embraced within the scope of the invention. In general, a reference to a compound is intended to cover its stereoisomers and mixture of various stereoisomers.

[0037] The term "optionally substituted" as used herein means that substitution is optional and therefore includes both unsubstituted and substituted atoms and moieties. A "substituted" atom or moiety indicates that any hydrogen on the
 20 designated atom or moiety can be replaced with a selection from the indicated substituent group, provided that the normal valency of the designated atom or moiety is not exceeded, and that the substitution results in a stable compound.

[0038] The term "pharmaceutically acceptable salt" as used herein refers to one or more salts of a given compound which possesses the desired pharmacological activity of the free compound and which are neither biologically nor
 25 otherwise undesirable. In general, the "pharmaceutically acceptable salts" refer to salts that are suitable for use in contact with the tissues of human and animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, et al. (J. Pharmaceutical Sciences, 66: 1-19 (1977)), incorporated herein by reference in its entirety, describes various pharmaceutically acceptable salts in details.

[0039] In general, the compounds according to the invention contain basic (e.g. nitrogen atoms) as well as acid moieties (e.g. compounds of Formula (I) wherein M is hydrogen). A person of skills in the art would appreciate that such compounds, therefore, can form acidic salts (formed with inorganic and/or organic acids), as well as basic salts (formed with inorganic
 30 and/or organic bases). Such salts can be prepared using procedures described in the art. For example, the basic moiety can be converted to its salt by treating a compound with a suitable amount of acid. Typical, non-limiting examples of such suitable acids include hydrochloric acid, trifluoroacetic acid, methanesulphonic acid, or the like. Alternatively, the acid moiety may be converted into its salt by treating with a suitable base. Typical non-limiting examples of such bases
 35 include sodium carbonate, sodium bicarbonate, potassium carbonate, potassium bicarbonate or the like. In case of compounds containing more than one functional groups capable of being converted into salt, each such functional group may be converted to salt independently. For example, in case of compounds containing two basic nitrogen atoms, one basic nitrogen can form salt with one acid while the other basic nitrogen can form salt with another acid. Some compounds
 40 according to the invention contain both, acidic as well as basic moieties, and thus can form inner salts or corresponding zwitterions. In general, all pharmaceutically acceptable salt forms of compounds of Formula (I) according to invention including acid addition salts, base addition salts, zwitterions or the like are contemplated to be within the scope of the present invention and are generically referred to as pharmaceutically acceptable salts.

[0040] The term "halogen" or "halo" as used herein refers to chlorine, bromine, fluorine, or iodine.

[0041] The term "infection" or "bacterial infection" as used herein includes presence of bacteria, in or on a subject, which, if its growth were inhibited, would result in a benefit to the subject. As such, the term "infection" in addition to referring to the presence of bacteria also refers to normal flora, which is not desirable. The term "infection" includes
 45 infection caused by bacteria.

[0042] The term "treat", "treating" or "treatment" as used herein refers to administering a medicament, including a pharmaceutical composition, or one or more pharmaceutically active ingredients, for prophylactic and/or therapeutic
 50 purposes. The term "prophylactic treatment" refers to treating a subject who is not yet infected, but who is susceptible to, or otherwise at a risk of infection (preventing the bacterial infection). The term "therapeutic treatment" refers to administering treatment to a subject already suffering from infection. The terms "treat", "treating" or "treatment" as used
 55 herein also refer to administering compositions or one or more of pharmaceutically active ingredients discussed herein, with or without additional pharmaceutically active or inert ingredients, in order to: (i) reduce or eliminate either a bacterial infection or one or more symptoms of the bacterial infection, or (ii) retard the progression of a bacterial infection or of one or more symptoms of the bacterial infection, or (iii) reduce the severity of a bacterial infection or of one or more

symptoms of the bacterial infection, or (iv) suppress the clinical manifestation of a bacterial infection, or (v) suppress the manifestation of adverse symptoms of the bacterial infection.

[0043] The term "pharmaceutically effective amount" or "therapeutically effective amount" or "effective amount" as used herein refers to an amount, which has a therapeutic effect or is the amount required to produce a therapeutic effect in a subject. For example, a therapeutically or pharmaceutically effective amount of an antibacterial agent or a pharmaceutical composition is the amount of the antibacterial agent or the pharmaceutical composition required to produce a desired therapeutic effect as may be judged by clinical trial results, model animal infection studies, and/or in vitro studies (e.g. in agar or broth media). The pharmaceutically effective amount depends on several factors, including but not limited to, the microorganism (e.g. bacteria) involved, characteristics of the subject (for example height, weight, sex, age and medical history), severity of infection and the particular type of the antibacterial agent used. For prophylactic treatments, a therapeutically or prophylactically effective amount is that amount which would be effective in preventing a microbial (e.g. bacterial) infection.

[0044] The term "administration" or "administering" includes delivery of a composition or one or more pharmaceutically active ingredients to a subject, including for example, by any appropriate methods, which serves to deliver the composition or its active ingredients or other pharmaceutically active ingredients to the site of the infection. The method of administration may vary depending on various factors, such as for example, the components of the pharmaceutical composition or the nature of the pharmaceutically active or inert ingredients, the site of the potential or actual infection, the microorganism involved, severity of the infection, age and physical condition of the subject and a like. Some non-limiting examples of ways to administer a composition or a pharmaceutically active ingredient to a subject according to this invention includes oral, intravenous, topical, intratracheal, intraperitoneal, intramuscular, parenteral, sublingual, transdermal, intranasal, aerosol, intraocular, intratracheal, intrarectal, vaginal, gene gun, dermal patch, eye drop, ear drop or mouthwash. In case of a pharmaceutical composition comprising more than one ingredient (active or inert), one of way of administering such composition is by admixing the ingredients (e.g. in the form of a suitable unit dosage form such as tablet, capsule, solution, powder and a like) and then administering the dosage form. Alternatively, the ingredients may also be administered separately (simultaneously or one after the other) as long as these ingredients reach beneficial therapeutic levels such that the composition as a whole provides a synergistic and/or desired effect.

[0045] The term "growth" as used herein refers to a growth of one or more microorganisms and includes reproduction or population expansion of the microorganism (e.g. bacteria). The term also includes maintenance of on-going metabolic processes of a microorganism, including processes that keep the microorganism alive.

[0046] The term, "effectiveness" as used herein refers to ability of a treatment or a composition or one or more pharmaceutically active ingredients to produce a desired biological effect in a subject. For example, the term "antibacterial effectiveness" of a composition or an antibacterial agent refers to the ability of the composition or the antibacterial agent to prevent or treat the microbial (e.g. bacterial) infection in a subject.

[0047] The term "synergistic" or "synergy" as used herein refers to the interaction of two or more agents so that their combined effect is greater than their individual effects.

[0048] The term "antibacterial agent" as used herein refers to any substance, compound or a combination of substances or a combination compounds capable of: (i) inhibiting, reducing or preventing growth of bacteria; (ii) inhibiting or reducing ability of a bacteria to produce infection in a subject; or (iii) inhibiting or reducing ability of bacteria to multiply or remain infective in the environment. The term "antibacterial agent" also refers to compounds capable of decreasing infectivity or virulence of bacteria.

[0049] The term "beta-lactam antibacterial agent" as used herein refers to compounds with antibacterial properties and containing a beta-lactam nucleus in their molecular structure.

[0050] The term "beta-lactamase" as used herein refers to any enzyme or protein or any other substance that breaks down a beta-lactam ring. The term "beta-lactamase" includes enzymes that are produced by bacteria and have the ability to hydrolyze the beta-lactam ring in a beta-lactam compound, either partially or completely.

[0051] The term "beta-lactamase inhibitor" as used herein refers to a compound capable of inhibiting activity of one or more beta-lactamase enzymes, either partially or completely.

[0052] The term "pharmaceutically inert ingredient" or "carrier" or "excipient" refers to a compound or material used to facilitate administration of a compound, including for example, to increase the solubility of the compound. Typical, non-limiting examples of solid carriers include, starch, lactose, dicalcium phosphate, sucrose, and kaolin and so on. Typical, non-limiting examples of liquid carriers include, sterile water, saline, buffers, non-ionic surfactants, and edible oils such as oil, peanut and sesame oils and so on. In addition, various adjuvants commonly used in the art may be included. These and other such compounds are described in the literature, for example, in the Merck Index (Merck & Company, Rahway, N.J.). Considerations for inclusion of various components in pharmaceutical compositions are described, for example, in Gilman et al. (Eds.) (1990); Goodman and Gilman's: The Pharmacological Basis of Therapeutics, 8th Ed., Pergamon Press., which is incorporated herein by reference in its entirety.


[0053] The term "subject" as used herein refers to vertebrate or invertebrate, including a mammal. The term "subject" includes human, animal, a bird, a fish, or an amphibian. Typical, non-limiting examples of a "subject" includes humans,

cats, dogs, horses, sheep, bovine cows, pigs, lambs, rats, mice and guinea pigs.

[0054] The term "pharmaceutically acceptable derivative" as used herein refers to and includes any pharmaceutically acceptable salt, pro-drugs, metabolites, esters, ethers, hydrates, polymorphs, solvates, complexes, enantiomers or adducts of a compound described herein which, upon administration to a subject, is capable of providing (directly or indirectly) the parent compound. For example, the term "antibacterial agent or a pharmaceutically acceptable derivative thereof" includes all derivatives of the antibacterial agent (such as salt, pro-drugs, metabolites, esters, ethers, hydrates, polymorphs, solvates, complexes, enantiomers or adducts) which, upon administration to a subject, is capable of providing (directly or indirectly) the antibacterial compound.

[0055] In general, the term "cation" includes Na, K, Mg, Ca, NCH_4^+ , $(\text{CH}_3\text{CH}_2)_3\text{N}^+$ etc.

[0056] In one general aspect, there are provided compounds of Formula (I):



or a stereoisomer or a pharmaceutically acceptable salt thereof; wherein:

R_1 is:

- (a) hydrogen,
- (b) $(\text{CO})_n\text{-R}_3$,
- (c) COOR_4 , or
- (d) $\text{COCH}_2\text{COR}_3$

n is 0, 1 or 2;

R_2 is:

- (a) SO_3M ,
- (b) SO_2NH_2 ,
- (c) PO_3M ,
- (d) CH_2COOM ,
- (e) CF_2COOM ,
- (f) CHF_2COOM , or
- (g) CF_3 ;

M is hydrogen or a cation;

R_3 is:

- (a) hydrogen,
- (b) $\text{C}_1\text{-C}_6$ alkyl optionally substituted with one or more substituents independently selected from halogen, OR_5 , CN, COOR_5 , CONR_6R_7 , NR_6R_7 , NR_5COR_8 , $\text{NR}_5\text{CONR}_6\text{R}_7$, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) CN,
- (d) NR_6R_7 ,
- (e) CONR_6R_7 ,
- (f) $\text{NHCONR}_6\text{R}_7$,
- (g) aryl optionally substituted with one or more substituents independently selected from $\text{C}_1\text{-C}_6$ alkyl, OR_5 , NR_6R_7 , halogen, CN, CONR_6R_7 , $\text{SO}_2\text{-alkyl}$, $\text{SO}_2\text{-aryl}$, $\text{OSO}_2\text{-alkyl}$, $\text{OSO}_2\text{-aryl}$, or $\text{NHCONR}_6\text{R}_7$,
- (h) heterocyclyl optionally substituted with one or more substituents independently selected from $\text{C}_1\text{-C}_6$ alkyl, OR_5 , NR_6R_7 , halogen, CN, CONR_6R_7 , $\text{SO}_2\text{-alkyl}$, $\text{SO}_2\text{-aryl}$, $\text{OSO}_2\text{-alkyl}$, $\text{OSO}_2\text{-aryl}$, $\text{NHC(NH)NR}_6\text{R}_7$, or $\text{NHCONR}_6\text{R}_7$,
- (i) heteroaryl optionally substituted with one or more substituents independently selected from $\text{C}_1\text{-C}_6$ alkyl, OR_5 , NR_6R_7 , halogen, CN, CONR_6R_7 , $\text{SO}_2\text{-alkyl}$, $\text{SO}_2\text{-aryl}$, $\text{OSO}_2\text{-alkyl}$, $\text{OSO}_2\text{-aryl}$, or $\text{NHCONR}_6\text{R}_7$,
- (j) cycloalkyl optionally substituted with one or more substituents independently selected from $\text{C}_1\text{-C}_6$ alkyl, OR_5 ,

NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
 (k) cycloalkyl substituted with C₁-C₆ alkyl wherein C₁-C₆ alkyl is further substituted with one or more substituents
 independently selected from OR₅, NR₆R₇, halogen, CN, or CONR₆R₇, or
 (l) OR₈;

R₄ is:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅,
 CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅,
 NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (d) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl,
 OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (e) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl,
 OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇, or
- (f) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅,
 NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇;

R₅ and R₈ are each independently:

- (a) hydrogen, or
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, CN,
 CONR₆R₇, NR₆R₇, heterocyclyl, heteroaryl, cycloalkyl or aryl;

R₆ and R₇ are each independently:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅,
 CN, COOR₅, CONR₅R₈, NR₅R₈, NR₅COR₈, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅,
 NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (d) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl,
 OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (e) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl,
 OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (f) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅,
 NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈, or
- (g) R₆ and R₇ are joined together to form a four to seven member ring.

[0057] Typical non-limiting examples of compounds according to the invention include:

trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-{7-oxo-2-[N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-acetic acid;

trans-difluoro-{7-oxo-2-[N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-acetic acid;

trans-sulfuric acid mono-[2-hydrazinocarbonyl-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

EP 2 748 165 B1

trans-sulfuric acid mono-[2-(N'-(amino-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(3-amino-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

5 trans-sulfuric acid mono-[2-(N'-(4-amino-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-((2S)-2-amino-3-hydroxy-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

10 trans-sulfuric acid mono-[2-(N'-[(2S,4S)-4-fluoro-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S,4R)-4-methoxy-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

15 trans-sulfuric acid mono-[2-(N'-(piperidin-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

20 trans-sulfuric acid mono-[2-(N'-((RS)-piperidin-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-((S)-piperidin-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

25 trans-sulfuric acid mono-[2-(N'-((RS)-piperidin-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-((S)-piperidin-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

30 trans-sulfuric acid mono-[2-(N'-((R)-piperidin-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

35 trans-sulfuric acid mono-[2-(N'-(piperazin-4-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-((RS)-1-amino-1-phenyl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

40 trans-sulfuric acid mono-[2-(N'-((RS)-3-amino-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(3-amino-2,2-dimethyl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

45 trans-sulfuric acid mono-[2-(N'-(1-aminomethyl-cyclopropan-1-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

50 trans-sulfuric acid mono-[2-(N'-(2-amino-4-carboxamido-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(5-amino-pentanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

55 trans-sulfuric acid mono-[2-(N'-((2S)-2,6-diamino-hexanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-((2-aminoethoxy)-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-

yl]ester;

trans-sulfuric acid mono-[2-(N'-[azetidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[pyrrolidin-1-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(S)-3-pyrrolidin-2-yl]-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S,4R)-4-hydroxy-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S,4S)-4-amino-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S,4S)-4-guanidino-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-3-piperidin-2-yl-propionyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-piperazin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(S)-morpholin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(3-oxo-3-piperazin-1-yl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-1-amino-1-pyridin-2-yl-acetyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2-amino-thiazol-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-[(cyano-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-N'-(7-oxo-6-sulfooxy-1,6-diaza-bicyclo[3.2.1]octane-2-carboxylic acid)-hydrazinocarboxylic acid tert-butyl ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-[(morpholin-4-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-[(6-carboxamido-pyridin-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-(morpholin-4-oxo-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-[(S)-1-carbamoyl-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-[(2S,4S)-1-carbamoyl-4-fluoro-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-[(S)-1-methanesulfonyl-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

Sodium salt of trans-sulfuric acid mono-[2-(N'-(cyano-dimethyl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;


Sodium salt of trans-sulfuric acid mono-[2-(N'-[(S)-5-oxo-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0058] In general, the compounds of the invention can be prepared according to the following procedures (Scheme 1). A person of skills in the art would appreciate that the described methods can be varied or optimized further to provide the desired and related compounds. In the following procedures, all variables are as defined above.

[0059] As described in Scheme-1, trans-6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carboxylic acid (**1a**), which is described in PCT International Publication No. WO 2009/091856, was reacted with corresponding acid hydrazides in presence of a suitable coupling agent such as EDC hydrochloride, or dicyclohexylcarbodiimide (DCC), in a solvent such as N,N dimethyl formamide; N,N dimethyl acetamide; 1,4 dioxane; chloroform; dichloromethane; or dichloroethane at a temperature ranging from -15°C to 60°C for about 1 to 24 hours to obtain intermediate compound (**1b**).

[0060] The intermediate compound (**1b**) was subjected for hydrogenolysis in presence of a suitable catalyst (e.g. 5% or 10% palladium on carbon, or 20% palladium hydroxide on carbon) in presence of a suitable hydrogen source (such as hydrogen gas, ammonium formate, cyclohexene) in a suitable solvent (such as methanol, ethanol, methanol-dichloromethane mixture, or N,N dimethyl formamide-dichloromethane mixture) at a temperature ranging from about 25°C to 60°C for about 1 to 14 hours to obtain intermediate compound (**1c**).




Scheme-1

[0061] The intermediate compound (**1c**) was sulfonated by reacting it with a sulfonating reagent (such as sulfur trioxide-pyridine complex, or sulfur trioxide-N,N-dimethyl formamide complex) in a suitable solvent (such as pyridine, N,N-dimethyl formamide) at a temperature ranging from about 25°C to 90°C for about 1 to 24 hours to obtain pyridine salt of sulfonic acid which when treated with tetrabutyl ammonium sulfate provided tetrabutylammonium salt of sulfonic acid as an intermediate compound (**1d**).

[0062] Some compounds according to the invention were isolated as zwitterions, by treating intermediate compound

(1d) with trifluoroacetic acid, in a suitable solvent (such as dichloromethane, chloroform, or acetonitrile) at a temperature ranging from about -10°C to 40°C for about 1 to 14 hours, especially when R in intermediate compound (1d) contained tert-butoxycarbonyl protected amine function.

[0063] Some other compounds according to the invention were isolated as a corresponding sodium salt, by passing intermediate compound (1d) through sodium form of Aberlite 200C resin in a tetrahydrofuran-water mixture followed by evaporation of the solvent under vacuum.



[0064] As described in Scheme-2, the hydroxyl intermediate (1c) obtained as per Scheme-1, was subjected for alkylation with alkylating agent (such as ethyl-bromoacetate, ethyl-fluoroacetate or ethyl-difluoroacetate) in presence of a base (such as potassium carbonate, diisopropylethylamine or triethylamine) in a suitable solvent (such as N,N dimethyl formamide, N,N dimethyl acetamide or N-methyl pyrrolidine) to provide O-alkylated compound (2d).

[0065] The compound (2d) was subjected for hydrolysis in presence of a base (such as lithium hydroxide or potassium hydroxide) in a suitable solvent (such as aqueous tetrahydrofuran, aqueous dioxane) to provide compound of Formula (I) after pH adjustment.

[0066] Optionally, if R₁ bears amine function protected with Boc group, then it was removed in an additional step of deprotection by using a suitable deprotecting agent (such as trifluoroacetic acid or HF pyridine) in a solvent (such as dichloromethane, chloroform or acetonitrile) to provide compound of Formula (I).

[0067] In some embodiments, there are provided pharmaceutical compositions comprising a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0068] Also disclosed is a method for preventing or treating bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0069] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0070] Also disclosed is a method for preventing or treating bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of a pharmaceutical composition comprising a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0071] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of a pharmaceutical composition comprising a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0072] In some embodiments, there are provided pharmaceutical compositions comprising: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable derivative thereof.

[0073] In some other embodiments, there are provided pharmaceutical compositions comprising: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) at least one antibacterial agent or a pharmaceutically acceptable derivative thereof.

[0074] In some other embodiments, there are provided pharmaceutical compositions comprising: (a) a compound of Formula (I), or a stereoisomer or a pharmaceutically acceptable salt thereof, (b) at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable derivative thereof, and (c) at

acceptable derivative thereof.

[0087] In some embodiments, there are provided pharmaceutical compositions comprising: (a) trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0088] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of: (a) trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0089] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, said method comprising administering to said subject a pharmaceutically effective amount of: (a) trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0090] Also disclosed is a method for preventing or treating bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of a pharmaceutical composition comprising: (a) trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0091] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of a pharmaceutical composition comprising: (a) trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0092] In some embodiments, there are provided pharmaceutical compositions comprising: (a) trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0093] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of: (a) trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0094] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, said method comprising administering to said subject a pharmaceutically effective amount of: (a) trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0095] Also disclosed is a method for preventing or treating bacterial infection in a subject, said method comprising administering to said subject a pharmaceutically effective amount of a pharmaceutical composition comprising: (a) trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0096] Also disclosed is a method for preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes, wherein the method comprises administering to said subject a pharmaceutically effective amount of a pharmaceutical composition comprising: (a) trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable derivative thereof.

[0097] Also disclosed is methods for increasing antibacterial effectiveness of a antibacterial agent in a subject, said method comprising co-administering said antibacterial agent or a pharmaceutically acceptable derivative thereof with a pharmaceutically effective amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0098] In some embodiments, the compositions and methods according to the invention use compounds of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof in combination with at least one antibacterial agent or a pharmaceutically acceptable derivative thereof. A wide variety of antibacterial agents can be used. Typical, non-limiting examples of antibacterial agents include one or more of antibacterial compounds generally classified as aminoglycosides, Ansamycins, Carbacephems, Cephalosporins, Cephamycins, Lincosamides, Lipopeptides, Macrolides, Mono-

bactams, Nitrofurans, Penicillins, Polypeptides, Quinolones, Sulfonamides, Tetracyclines, Oxazolidinone and the like.

[0099] Typical, non-limiting examples of Aminoglycoside antibacterial agents include Amikacin, Gentamicin, Kanamycin, Neomycin, Netilmicin, Tobramycin, Paromomycin, Arbekacin, Streptomycin, Apramycin and the like.

[0100] Typical, non-limiting examples of Ansamycin antibacterial agents include Geldanamycin, Herbimycin and the like.

[0101] Typical, non-limiting examples of Carbacephem antibacterial agents include Loracarbef and the like.

[0102] Typical, non-limiting examples of Carbapenem antibacterial agents include Ertapenem, Doripenem, Imipenem, Meropenem and the like.

[0103] Typical, non-limiting examples of Cephalosporin and Cepharmycin antibacterial agents include Cefazolin, Cefacetrile, Cefadroxil, Cefalexin, Cefaloglycin, Cefalonium, Cefaloridine, Cefalotin, Cefapirin, Cefatrizine, Cefazedone, Cefazaflur, Cefradine, Cefroxadine, Ceftezole, Cefaclor, Cefamandole, Cefminox, Cefonicid, Ceforanide, Cefotiam, Cefprozil, Cefbuperazone, Cefuroxime, Cefuzonam, Cepharmycin, Cefoxitin, Cefotetan, Cefmetazole, Carbacephem, Cefixime, Ceftazidime, Ceftriaxone, Cefcapene, Cefdaloxime, Cefdinir, Cefditoren, Cefetamet, Cefmenoxime, Cefodizime, Cefoperazone, Cefotaxime, Cefpimizole, Cefpiramide, Cefpodoxime, Cefsulodin, Cefteram, Cefibuten, Ceftiolene, Ceftizoxime, Oxacephem, Cefepime, Cefozopran, Cefpirome, Cefquinome, Ceftobiprole, Ceftiofur, Cefquinome, Cefovecin, CXA-101, Ceftaroline, Ceftobiprole etc.

[0104] Typical, non-limiting examples of Lincosamide antibacterial agents include Clindamycin, Lincomycin and the like.

[0105] Typical, non-limiting examples of Macrolide antibacterial agents include Azithromycin, Clarithromycin, Dirithromycin, Erythromycin, Roxithromycin, Troleandomycin, Telithromycin, Spectinomycin, Solithromycin and the like.

[0106] Typical, non-limiting examples of Monobactam antibacterial agents include Aztreonam and the like.

[0107] Typical, non-limiting examples of Nitrofurantoin antibacterial agents include Furazolidone, Nitrofurantoin and the like.

[0108] Typical, non-limiting examples of Penicillin antibacterial agents include Amoxicillin, Ampicillin, Azlocillin, Carbenicillin, Cloxacillin, Dicloxacillin, Flucloxacillin, Mezlocillin, Methicillin, Nafcillin, Oxacillin, Penicillin G, Penicillin V, Piperacillin, Temocillin, Ticarcillin and the like.

[0109] Typical, non-limiting examples of Polypeptide antibacterial agents include Bacitracin, Colistin, Polymyxin B and the like.

[0110] Typical, non-limiting examples of Quinolone antibacterial agents include Ciprofloxacin, Enoxacin, Gatifloxacin, Levofloxacin, Lomefloxacin, Moxifloxacin, Nalidixic acid, Levonadifloxacin, Norfloxacin, Ofloxacin, Trovafloxacin, Grepafloxacin, Sparfloxacin, Temafloxacin and the like.

[0111] Typical, non-limiting examples of Sulfonamide antibacterial agents include Mafenide, Sulfonamidochrysoidine, Sulfacetamide, Sulfadiazine, Sulfamethizole, Sulfamethoxazole, Sulfasalazine, Sulfisoxazole, Trimethoprim and the like.

[0112] Typical, non-limiting examples of Tetracycline antibacterial agents include Demeclocycline, Doxycycline, Minocycline, Oxytetracycline, Tetracycline, Tigecycline and the like.

[0113] Typical, non-limiting examples of Oxazolidinone antibacterial agents include Tedizolid, Linezolid, Ranbezolid, Torezolid, Radezolid etc.

[0114] The pharmaceutical compositions according to the invention may include one or more pharmaceutically acceptable carriers or excipients or the like. Typical, non-limiting examples of such carriers or excipient include mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, sodium crosscarmellose, glucose, gelatin, sucrose, magnesium carbonate, wetting agents, emulsifying agents, solubilizing agents, pH buffering agents, lubricants, stabilizing agents, binding agents etc.

[0115] The pharmaceutical compositions according to this invention can exist in various forms. In some embodiments, the pharmaceutical composition is in the form of a powder or a solution. In some other embodiments, the pharmaceutical compositions according to the invention are in the form of a powder that can be reconstituted by addition of a compatible reconstitution diluent prior to parenteral administration. Non-limiting example of such a compatible reconstitution diluent includes water.

[0116] In some other embodiments, the pharmaceutical compositions according to the invention are in the form of a frozen composition that can be diluted with a compatible diluent prior to parenteral administration.

[0117] In some other embodiments, the pharmaceutical compositions according to the invention are in the form ready to use for parenteral administration.

[0118] In the methods according to the invention, the pharmaceutical composition and/or other pharmaceutically active ingredients disclosed herein may be administered by any appropriate method, which serves to deliver the composition or its constituents or the active ingredients to the desired site. The method of administration can vary depending on various factors, such as for example, the components of the pharmaceutical composition and nature of the active ingredients, the site of the potential or actual infection, the microorganism (e.g. bacteria) involved, severity of infection, age and physical condition of the subject. Some non-limiting examples of administering the composition to a subject according to this invention include oral, intravenous, topical, intraspiratory, intraperitoneal, intramuscular, parenteral,

sublingual, transdermal, intranasal, aerosol, intraocular, intratracheal, intrarectal, vaginal, gene gun, dermal patch, eye drop, ear drop or mouthwash.

[0119] The compositions according to the invention can be formulated into various dosage forms wherein the active ingredients and/or excipients may be present either together (e.g. as an admixture) or as separate components. When the various ingredients in the composition are formulated as a mixture, such composition can be delivered by administering such a mixture. The composition or dosage form wherein the ingredients do not come as a mixture, but come as separate components, such composition/dosage form may be administered in several ways. In one possible way, the ingredients may be mixed in the desired proportions and the mixture is then administered as required. Alternatively, the components or the ingredients (active or inert) may be separately administered (simultaneously or one after the other) in appropriate proportion so as to achieve the same or equivalent therapeutic level or effect as would have been achieved by administration of the equivalent mixture.

[0120] Similarly, in the methods according to the invention, the active ingredients disclosed herein may be administered to a subject in several ways depending on the requirements. In some embodiments, the active ingredients are admixed in appropriate amounts and then the admixture is administered to a subject. In some other embodiments, the active ingredients are administered separately. Since the invention contemplates that the active ingredients agents may be administered separately, the invention further provides for combining separate pharmaceutical compositions in kit form. The kit may comprise one or more separate pharmaceutical compositions, each comprising one or more active ingredients. Each of such separate compositions may be present in a separate container such as a bottle, vial, syringes, boxes, bags, and the like. Typically, the kit comprises directions for the administration of the separate components. The kit form is particularly advantageous when the separate components are preferably administered in different dosage forms (e.g., oral and parenteral) or are administered at different dosage intervals. When the active ingredients are administered separately, they may be administered simultaneously or sequentially.

[0121] The pharmaceutical composition or the active ingredients according to the present invention may be formulated into a variety of dosage forms. Typical, non-limiting examples of dosage forms include solid, semi-solid, liquid and aerosol dosage forms; such as tablets, capsules, powders, solutions, suspensions, suppositories, aerosols, granules, emulsions, syrups, elixirs and a like.

[0122] In general, the pharmaceutical compositions and method disclosed herein are useful in preventing or treating bacterial infections. Advantageously, the compositions and methods disclosed herein are also effective in preventing or treating infections caused by bacteria that are considered be less or not susceptible to one or more of known antibacterial agents or their known compositions. Some non-limiting examples of such bacteria known to have developed resistance to various antibacterial agents include *Acinetobacter*, *E. coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Enterobacter*, *Klebsiella*, *Citrobacter* and a like. Other non-limiting examples of infections that may be prevented or treated using the compositions and/or methods of the invention include: skin and soft tissue infections, febrile neutropenia, urinary tract infection, intraabdominal infections, respiratory tract infections, pneumonia (nosocomial), bacteremia meningitis, surgical, infections etc.

[0123] Surprisingly, the compounds, compositions and methods according to the invention are also effective in preventing or treating bacterial infections that are caused by bacteria producing one or more beta-lactamase enzymes. The ability of compositions and methods according to the present invention to treat such resistant bacteria with typical beta-lactam antibiotics represents a significant improvement in the art.

[0124] In general, the compounds of Formula (I) or a stereoisomer or pharmaceutically acceptable salt thereof according to invention are also useful in increasing antibacterial effectiveness of an antibacterial agent in a subject. The antibacterial effectiveness of one or more antibacterial agents may increased, for example, by co-administering said antibacterial agent or a pharmaceutically acceptable derivative thereof with a pharmaceutically effective amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof according to the invention. For example, those skilled in the art will recognize that the invention may be practiced using a variety of different compounds within the described generic descriptions.


EXAMPLES

[0125] The following examples illustrate the embodiments of the invention that are presently best known. However, it is to be understood that the following are only exemplary or illustrative of the application of the principles of the present invention. Thus, while the present invention has been described above with particularity, the following examples provide further detail in connection with what are presently deemed to be the most practical and preferred embodiments of the invention.

Example-1

trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester

[0126]



Step-1: Preparation of trans-2-[N'-(6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(S)-pyrrolidin-1-carboxylic acid tert-butyl ester:

[0127] To a clear solution of trans-6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carboxylic acid (15 gm, 0.054 mol) in N,N-dimethyl formamide (150 ml), was added EDC hydrochloride (15.57 gm, 0.082 mol) followed by HOBt (11.0 gm, 0.082 mol) at about 25°C to 35°C under stirring. The reaction mixture was stirred for 15 minutes and a solution of (S)-N-tert-butoxycarbonyl-pyrrolidin-2-carboxylic acid hydrazide (14.93 gm, 0.065 mol) dissolved in N,N-dimethyl formamide (75 ml), followed by N,N-di-isopropyl ethylamine (28.4 ml, 0.163 mol) were added. The reaction mixture was stirred at a temperature between 25°C to 35°C for 16 hours. The reaction mixture was poured under stirring into 10% aqueous citric acid solution (2250 ml). The resulting mixture was extracted with diethyl ether (1000 ml x 3). Combined organic layer was washed with water (1000 ml) followed by brine solution (500 ml) and dried over sodium sulfate. Concentration of organic layer under vacuum afforded the crude residue in 13 gm quantity. The residue was purified using silica gel column chromatography to provide the product (trans-2-[N'-(6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(S)-pyrrolidin-1-carboxylic acid tert-butyl ester) in 6.3 gm quantity as a white powder.

Analysis: MS (ES+) $C_{24}H_{33}N_5O_6 = 488.1$ (M+1);

1H NMR (DMSO- d_6) = 9.86 (br d, 1H), 9.75 (br d, 1H), 7.34-7.44 (m, 5H), 4.92 (dd, 2H), 4.07-4.10 (m, 1H), 3.78-3.82 (m, 1H), 3.68 (br d, 1H), 3.20-3.25 (m, 3H), 2.87 (br d, 1H), 1.62-2.10 (m, 8H), 1.34 (s, 9H).

Step-2: Preparation of trans-2-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(S)-pyrrolidin-1-carboxylic acid tert-butyl ester:

[0128] To a clear solution of step-1 product (3.0 gm, 6.15 mmol) in methanol (30 ml) was added 10% palladium on carbon (300 mg). The suspension was stirred under atmospheric hydrogen pressure at a temperature of about 30°C for 2 hours. The catalyst was filtered over a celite bed and catalyst-containing bed was washed with additional methanol (10 ml) and dichloromethane (10 ml). The filtrate was concentrated in vacuum to provide a white powder, which was triturated with diethyl ether to provide trans-2-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(S)-pyrrolidin-1-carboxylic acid tert-butyl ester as a white powder in 2.00 gm quantity in 82% yield.

Analysis: MS (ES+) $C_{17}H_{27}N_5O_6 = 398.0$ (M+1);

1H NMR (DMSO- d_6) = 9.82 (d, 1H), 9.70-9.80 (m, 2H), 4.08-4.15 (m, 1H), 3.4.0-3.78 (m, 1H), 3.59 (br s, 1H), 3.17-3.40 (m, 3H), 2.97 (br d, 1H), 1.55-2.15 (m, 8H), 1.35 (s, 9H).

Step-3: Preparation of tetrabutylammonium salt of trans-2-[N'-(6-sulfooxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(S)-pyrrolidin-1-carboxylic acid tetrabutyl ester:

[0129] The product obtained in step-2 (2.00 gm, 5.03 mmol) was dissolved in pyridine (40 ml) and to the clear solution was added pyridine sulfur trioxide complex (4.03 gm, 25.18 mmol). The suspension was stirred at a temperature 25°C to 35°C for overnight. The suspension was filtered and the solids were washed with dichloromethane (25 ml x 2). The filtrate was evaporated under vacuum and the residue was stirred in 0.5 N aqueous potassium dihydrogen phosphate

solution (200 ml) for 0.5 hour. The solution was washed with ethyl acetate (100 ml x 4) and layers were separated. To the aqueous layer was added tetrabutylammonium sulphate (1.71 gm, 5.03 mmol) and stirred for four hours at about 25°C. The mixture was extracted with dichloromethane (100 ml x 2). The combined organic extract was washed with brine (50 ml) and dried on sodium sulfate and evaporated under vacuum to provide solid, that was triturated with diethyl ether and filtered to provide white powder as a step-3 product (tetrabutylammonium salt of trans-2-[N'-(6-sulfooxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(S)-pyrrolidin-1-carboxylic acid tetr-butyl ester), in 3.0 gm quantity (83% yield).

Analysis: MS (ES-) $C_{17}H_{26}N_5O_9S.N(C_4H_9)_4$ as a salt = 476.0 (M-1) as a free sulfonic acid;

1H NMR ($CDCl_3$) = 9.13 (br s, 1H), 8.49 (br s, 1H), 4.35 (br s, 2H), 3.98 (d, 1H), 3.24-3.50 (m, 10H), 3.13 (br d, 1H), 2.35 (dd, 2H), 2.16 (br s, 2H), 1.91-2.01 (m, 4H), 1.61-1.70 (m, 10H), 1.40-1.48 (m, 17H), 0.98-1.02 (m, 12H).

Step-4: trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester:

[0130] To the powder obtained in step-3 (3.0 gm, 4.17 mmol) was added, a solution of trifluoroacetic acid (7 ml) in dichloromethane (7 ml) slowly by syringe at -5°C over a period of 5 minutes. The mixture was maintained under stirring for 1 hr. Solvents were removed below 40°C under high vacuum to provide a residue, which was triturated with diethyl ether (50 ml x 5) and each time diethyl ether was decanted. The obtained white solid was further triturated with acetonitrile (100 ml x 2). The resultant solid was stirred in dichloromethane (100 ml) and the suspension was filtered. The solid was dried under vacuum to provide title compound of the invention (trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester) in 1.2 gm quantity (59% yield).


Analysis: MS (ES-) $C_{12}H_{19}N_5O_7S$ = 376.2 (M-1) as a free sulfonic acid;

1H NMR ($DMSO-d_6$) = 10.39 (br s, 1H), 10.15 (s, 1H), 8.96 (br s, 2H), 4.19 (t, 1H), 4.03 (br s, 1H), 3.86 (d, 1H), 3.16-3.25 (m, 3H), 3.02 (br d, 1H), 2.27-2.33 (m, 1H), 1.92-2.23 (m, 1H), 1.84-1.90 (m, 4H), 1.69-1.75 (m, 1H), 1.54-1.62 (m, 1H).

Example-2

trans-sulfuric acid mono-r2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester

[0131]



Step-1: Preparation of trans-3-[N'-(6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester:

[0132] By using the procedure described in Step-1 of Example-1 above, and by using trans-6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carboxylic acid (25 gm, 0.084 mol), N,N-dimethyl formamide (625 ml), EDC hydrochloride (24 gm, 0.126 mol), HOBT (16.96 gm, 0.126 mol), (R)-N-tert-butoxycarbonyl-piperidin-3-carboxylic acid hydrazide (21.40 gm, 0.088 mol) to provide the title compound in 17.0 gm quantity, 41% yield as a white solid.

Analysis: MS (ES+) $C_{25}H_{35}N_5O_6$ = 502.1 (M+1);

1H NMR ($CDCl_3$) = 8.40 (br s, 1H), 7.34-7.44 (m, 5H), 5.05 (d, 1H), 4.90 (d, 1H), 4.00 (br d, 1H), 3.82 (br s, 1H), 3.30 (br s, 1H), 3.16-3.21 (m, 1H), 3.06 (br d, 1H), 2.42 (br s, 1H), 2.29-2.34 (m, 1H), 1.18-2.02 (m, 4H), 1.60-1.75 (m, 4H), 1.45-1.55 (m, 2H), 1.44 (s, 9H).

Step-2: Preparation of trans-3-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester:

[0133] By using the procedure described in Step-2 of Example-1 above, and by using trans-3-[N'-(6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester (16.5 gm , 0.033 mol), methanol (170 ml) and 10% palladium on carbon (3.5 gm) to provide the title compound in 13.5 gm quantity as a pale pink solid and it was used for the next reaction immediately.

Analysis: MS (ES+) $C_{18}H_{29}N_5O_6 = 411.1$ (M+1);

Step-3: Preparation of tetrabutylammonium salt of trans-3-[N'-(6-sulfooxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester:

[0134] By using the procedure described in Step-3 of Example-1 above, and by using trans-3-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester (13.5 gm , 0.033 mol), pyridine (70 ml) and pyridine sulfur trioxide complex (26.11 gm, 0.164 mol), 0.5 N aqueous potassium dihydrogen phosphate solution (400 ml) and tetrabutylammonium sulphate (9.74 gm, 0.033 mol) to provide the title compound in 25 gm quantity as a yellowish solid, in quantitative yield.

Analysis: MS (ES-) $C_{18}H_{28}N_5O_9S.N(C_4H_9)_4$ as a salt = 490.0 (M-1) as a free sulfonic acid;

Step-4: trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester:

[0135] By using the procedure described in Step-4 of Example-1 above, and by using tetrabutylammonium salt of trans-3-[N'-(6-sulfooxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester (24 gm , 0.032 mmol), dichloromethane (60 ml) and trifluoroacetic acid (60 ml) to provide the title compound in 10 gm quantity as a white solid, in 79% yield.

Analysis: MS (ES-)= $C_{13}H_{21}N_5O_7S = 390.2$ (M-1) as a free sulfonic acid;


1H NMR (DMSO- d_6) = 9.97 (d, 2H), 8.32 (br s, 2H), 4.00 (br s, 1H), 3.81 (d, 1H), 3.10-3.22 (m, 3H), 2.97-3.02 (m, 2H), 2.86-2.91 (m, 1H), 2.65-2.66 (m, 1H), 1.97-2.03 (m, 1H), 1.57-1.88 (m, 7H).

$[\alpha]_D^{25} = -32.6^\circ$, (c 0.5, water).

Example-3

trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester

[0136]



Step-1: Preparation of trans-3-[N'-(6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-pyrrolidin-1-carboxylic acid tert-butyl ester:

[0137] By using the procedure described in Step-1 of Example-1, and by using trans-6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carboxylic acid (15.7 gm, 0.057 mol), N,N-dimethyl formamide (390 ml), EDC hydrochloride (16.24 gm, 0.085 mol), HOBT (11.48 gm, 0.085 mol), (R)-N-tert-butoxycarbonyl-pyrrolidin-3-carboxylic acid hydrazide (13.7 gm , 0.06 mol) to provide the title compound in 11.94 gm quantity, 43% yield as a white solid.

EP 2 748 165 B1

Analysis: MS (ES+) $C_{24}H_{33}N_5O_6 = 488.2$ (M+1);

1H NMR ($CDCl_3$, D_2O exchange) = 7.30-7.39 (m, 5H), 4.85 (s, 2H), 3.77 (d, 1H), 3.68 (br s, 1H), 3.39-3.41 (m, 1H), 3.17-3.26 (m, 3H), 3.01 (d, 1H), 2.90-2.92 (m, 2H), 1.97-2.03 (m, 2H), 1.79-1.89 (m, 2H), 1.66-1.70 (m, 1H), 1.55-1.57 (m, 1H), 1.32 (s, 9H).

Step-2: Preparation of trans-3-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-pyrrolidin-1-carboxylic acid tert-butyl ester:

[0138] By using the procedure described in Step-2 of Example-1, and by using trans-3-[N'-(6-benzyloxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-pyrrolidin-1-carboxylic acid tert-butyl ester (11.5 gm, 0.024 mol), methanol (115 ml) and 10% palladium on carbon (3.0 gm) to provide the title compound in 9.5 gm quantity as a pale brown solid and it was used for the next reaction immediately.

Analysis: MS (ES+) $C_{17}H_{27}N_5O_6 = 398.2$ (M+1);

Step-3: Preparation of tetrabutylammonium salt of trans-3-[N'-(6-sulfooxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-pyrrolidin-1-carboxylic acid tert-butyl ester:

[0139] By using the procedure described in Step-3 of Example-1, and by using trans-3-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-pyrrolidin-1-carboxylic acid tert-butyl ester (9.5 gm, 0.024 mol), pyridine (95 ml) and pyridine sulfur trioxide complex (19.08 gm, 0.12 mol), 0.5 N aqueous potassium dihydrogen phosphate solution (300 ml) and tetrabutylammonium sulphate (8.15 gm, 0.024 mol) to provide the title compound in 15.3 gm quantity as a yellowish solid, in 87% yield.

Analysis: MS (ES-) $C_{17}H_{26}N_5O_9S.N(C_4H_9)_4$ as a salt = 476.1 (M-1) as a free sulfonic acid;

1H NMR ($DMSO-d_6$) = 9.82 (d, 2H), 3.97 (br s, 1H), 3.79 (d, 1H), 3.42-3.44 (m, 1H), 3.00-3.18 (m, 10H), 2.65-2.97 (m, 2H), 1.98-2.01 (m, 2H), 1.74-1.83 (m, 2H), 1.63-1.72 (m, 1H), 1.38-1.55 (m, 9H), 1.33 (s, 9H), 1.24-1.28 (m, 8H), 0.91-0.99 (m, 12H).

Step-4: trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester:

[0140] By using the procedure described in Step-4 of Example-1, and by using tetrabutylammonium salt of trans-3-[N'-(6-sulfooxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-pyrrolidin-1-carboxylic acid tert-butyl ester (15 gm, 0.021 mmol), dichloromethane (37 ml) and trifluoroacetic acid (37 ml) to provide the title compound in 7.7 gm quantity as a white solid.


Analysis: MS (ES-) = $C_{12}H_{19}N_5O_7S = 376.1$ (M-1) as a free sulfonic acid;

1H NMR ($DMSO-d_6$) = 10.04 (s, 1H), 9.96 (s, 1H), 8.79 (br s, 1H), 8.68 (br s, 1H), 4.00 (br s, 1H), 3.82 (d, 1H), 3.18-3.32 (m, 4H), 3.08-3.12 (m, 1H), 3.00 (br d, 1H), 2.05-2.29 (m, 1H), 1.96-2.05 (m, 2H), 1.84-1.87 (m, 1H), 1.69-1.73 (m, 1H), 1.56-1.67 (m, 1H). $[\alpha]_D^{25} = -44.2^\circ$, (c 0.5, water).

Example-4

trans-{7-oxo-2-[N'-((R)-piperidin-3-carbonyl)-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-acetic acid

[0141]



Step-1: Preparation of trans-3-[N'-(6-ethoxycarbonylmethoxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester:

[0142] The intermediate compound trans-3-[N'-(6-hydroxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester (4.0 gm, 9.73 mmol), obtained in Step-2 of Example-2) was dissolved in DMF (12 ml) and to the clear solution was added potassium carbonate (1.61 gm, 11.6 mmol) followed by ethyl bromo acetate (1.2 ml, 10.0 mmol) under stirring and the suspension was stirred for 18 hours at about 25°C. The reaction was monitored by TLC. DMF was evaporated under vacuum to provide a residue. The residue was purified by silica gel column chromatography to provide titled Step-1 intermediate compound in 2.6 gm quantity as a solid in 53.7% yield.

Analysis: MS (+) = $C_{22}H_{35}N_5O_8$ = 498.1 (M+1);

1H NMR ($CDCl_3$) = 8.45 (br s, 2H), 4.58 (s, 2H), 4.19-4.27 (m, 2H), 4.02-4.12 (m, 2H), 3.25 (br d, 1H), 3.15 (br d, 1H), 2.38 (br s, 1H), 2.35 (dd, 1H), 2.15-2.20 (m, 1H), 1.79-2.02 (m, 4H), 1.67-1.77 (m, 4H), 1.44-1.51 (m, 11H), 1.28 (t, 3H).

Step-2: Preparation of trans-3-[N'-(6-carboxymethoxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester:

[0143] To a clear solution of trans-3-[N'-(6-ethoxycarbonylmethoxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester (600 mg, 1.20 mmol) in tetrahydrofuran (32 ml) and water (12 ml) was added lithium hydroxide (43.2 mg, 1.8 mmol) at 0°C. The reaction mixture was stirred for 3 hours and was neutralized to pH 6 by addition of aqueous 1 N potassium hydrogen sulfate. It was extracted with ethyl acetate (3 X 25 ml). Layers were separated and aqueous layer was acidified with 1 N potassium hydrogen sulfate to pH 1 and extracted with ethyl acetate (3 X 25 ml). The Organic layer was dried over sodium sulfate and evaporated to dryness under vacuum to provide 160 mg of titled intermediate as a solid in 27% yield.

Analysis: MS (ES+) = $C_{20}H_{31}N_5O_8$ = 470.1 (M+1)

1H NMR ($CDCl_3$) = 8.40 (br s, 2H), 4.67 (d, 1H), 4.52 (d, 1H), 4.07-4.14 (m, 2H), 3.95 (br s, 1H), 3.43 (br d, 1H), 3.19 (brd, 1H), 2.47 (brs, 1H), 2.39 (dd, 1H), 2.09-2.13 (m, 2H), 1.77-2.00 (m, 4H), 1.68-1.77 (m, 2H), 1.45-1.51 (m, 11H)..

Step-3: trans- {7-oxo-2-[N'-((R)-piperidin-3-carbonyl)-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]-oct-6-yloxy}-acetic acid:

[0144] To a clear solution of trans-3-[N'-(6-carboxymethoxy-7-oxo-1,6-diaza-bicyclo[3.2.1]octane-2-carbonyl)-hydrazinocarbonyl]-(R)-piperidin-1-carboxylic acid tert-butyl ester (150 mg, 0.32 mmol) in dichloromethane (2 ml) was added trifluoroacetic acid under stirring at -10°C. The reaction mixture was stirred for about 1 hour at -10°C and the solvents were evaporated under vacuum to provide a residue. The residue was triturated successively with diethyl ether (25 ml) and acetonitrile (25 ml) and solvents were decanted to provide solid that was dried under vacuum to provide 59 mg of titled compound in 50 % yield.


Analysis: MS (ES-) = $C_{15}H_{23}N_5O_6$ = 368.0 (M-1)

1H NMR ($DMSO-d_6$) = 9.97 (br s, 2H), 4.48 (d, 1H), 4.29 (d, 1H), 3.91 (s, 1H), 3.83 (d, 1H), 3.36 (q, 1H), 3.11-3.21 (m, 4H), 2.84-3.01 (m, 3H), 2.66 (br s, 1H), 1.90-2.05 (m, 3H), 1.69-1.76 (m, 2H), 1.59-1.66 (m, 3H).

Example-5

trans-difluoro-{7-oxo-2-[N'-((R)-peridin-3-carbonyl)-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]-oct-6-yloxy}-acetic acid

[0145]



[0146] By using procedure the described in Example 4 and by using ethyl-bromo difluoroacetate (2.0 gm, 10.0 mmol) in the place of ethyl bromo acetate, the titled compound was prepared in 30 mg quantity as a solid.

Analysis: MS (ES⁺) = C₁₅H₂₁F₂N₅O₆ = 406.2 (M+1)

¹H NMR (DMSO-d₆) = 10.99 (d, 2H), 8.59 (br s, 2H), 3.89-4.00 (m, 2H), 3.13-3.31 (m, 4H), 2.95-3.07 (m, 2H), 2.81-2.88 (m, 1H), 2.62-2.78 (m, 1H), 1.97-2.05 (m, 1H), 1.84-1.95 (m, 1H), 1.72-1.79 (m, 2H), 1.59-1.64 (m, 3H).

[0147] Compounds **6 to 42** (Table 1) were prepared using the procedure described as in Example-1 and using corresponding R₁CONHNH₂, in place of (S)-N-tert-butoxycarbonylpyrrolidin-2-carboxylic acid hydrazide.


























Table 1.








Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	¹ H NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
6.	t-Boc-NHNH ₂	H	11.05 (br s, 1H), 9.39 (br s, 2H), 4.04 (d, 1H), 3.92 (d, 1H), 3.04 (br d, 1H), 2.48 (d, 1H), 1.99-2.05 (m, 1H), 1.86-1.90 (m, 1H), 1.63-1.77 (m, 2H).	279.1 (C ₇ H ₁₂ N ₄ O ₆ S)
7.	t-Boc-HNCH ₂ -CO NHNH ₂	H ₂ NCH ₂ -CO-	10.20 (bs, 1H), 8.20 (br s, 3H), 4.01 (br s, 1H), 3.86 (br d, 1H), 3.61 (s, 2H), 2.99 (d, 1H), 1.97-2.05 (m, 1H), 1.78-1.88 (m, 1H), 1.68-1.72 (m, 1H), 1.54-1.67 (m, 2H).	336.2 (C ₉ H ₁₅ N ₅ O ₇ S)
8.	t-Boc-HNCH ₂ CH ₂ -CONHNH ₂	H ₂ NCH ₂ CH ₂ -CO-	9.34 (d, 2H), 7.66 (br s, 2H), 4.00 (br d, 1H), 3.84 (d, 1H), 3.16 (d, 1H), 2.92-3.00 (m, 3H), 2.43-2.53 (m, 2H), 1.85-2.05 (m, 1H), 1.74-1.75 (m, 1H), 1.51-1.73 (m, 2H).	350.2 (C ₁₀ H ₁₆ N ₅ O ₇ S)
9.	t-Boc-HN-(CH ₂) ₃ -CONHNH ₂	H ₂ N(CH ₂) ₃ -CO-	9.87 (s, 1H), 9.77 (s, 1H), 7.63 (br s, 3H), 4.00 (br s, 1H), 3.86 (d, 1H), 3.16 (d, 1H), 2.76-2.73 (m, 2H), 2.20 (t, 2H), 1.97-2.01 (m, 1H), 1.51-1.84 (m, 6H).	363.9 (C ₁₁ H ₁₉ N ₅ O ₇ S)
10.			10.33 (s, 1H), 10.09 (s, 1H), 8.19 (br s, 3H), 5.48 (br s, 1H), 4.02 (br s, 1H), 3.86 (d, 1H), 3.79 (d, 1H), 3.64-3.68 (m, 1H), 3.21 (d, 1H), 3.02 (d, 1H), 1.98-2.05 (m, 1H), 1.82-1.88 (m, 1H), 1.68-1.75 (m, 1H), 1.54-1.62 (m, 1H).	366.2 (C ₁₀ H ₁₇ N ₅ O ₈ S)
11.			10.46 (s, 1H), 10.20 (s, 1H), 9.73 (br s, 1H), 9.15 (br s, 1H), 5.44 (br s, 1H), 5.31-4.45 (d, 1H), 3.99 (s, 1H), 3.87 (d, 1H), 3.61 (dd, 2H), 3.28-3.42 (m, 1H), 3.02-3.12 (m, 2H), 2.60-2.82 (m, 1H), 2.30-2.42 (m, 1H), 1.99-2.04 (m, 1H), 1.80-1.90 (m, 1H), 1.58-1.74 (m, 2H).	393.9 (C ₁₂ H ₁₈ N ₅ FO ₇ S)
12.			10.41 (s, 1H), 10.19 (s, 1H), 9.59 (br s, 1H), 8.97 (br s, 1H), 4.19-4.24 (m, 1H), 4.13 (br s, 1H), 4.02 (br s, 1H), 3.87 (d, 1H), 3.38 (s, 3H), 3.16-3.33 (m, 3H), 3.02 (br d, 1H), 1.57-2.05 (m, 6 H).	406.2 (C ₁₃ H ₂₁ N ₅ O ₈ S)
13.			9.82 (d, 1H), 8.41 (br d, 1H), 8.19 (br d, 1H), 3.99 (br s, 1H), 3.80 (d, 1H), 3.18-3.39 (m, 5H), 2.90-3.00 (m, 3H), 1.97-2.03 (m, 1H), 1.75-1.84 (m, 2H), 1.67-1.71 (m, 3H), 1.57-1.62 (m, 2H).	390.3 (C ₁₃ H ₂₁ N ₅ O ₇ S)

(continued)












Table 1.

Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	¹ H NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
14.			9.99 (d, 2H), 8.40 (br s, 2H), 4.00 (br s, 1H), 3.81 (d, 1H), 3.10-3.21 (m, 4H), 2.91-3.01 (m, 3H), 2.65-2.66 (m, 1H), 1.97-2.05 (m, 1H), 1.54-1.88 (m, 7H).	389.9 (C ₁₃ H ₂₁ N ₅ O ₇ S)
15.			9.96 (d, 2H), 8.39 (br s, 2H), 4.00 (br s, 1H), 3.81 (d, 1H), 3.09-3.17 (m, 3H), 2.97-3.02 (m, 1H), 2.87-2.92 (m, 1H), 2.60-2.65 (m, 1H), 1.97-2.02 (m, 1H), 1.58-1.84 (m, 8H).	390.2 (C ₁₃ H ₂₁ N ₅ O ₇ S)
16.			10.29 (d, 1H), 10.16 (d, 1H), 8.90-9.03 (m, 1H), 8.70-8.78 (m, 1H), 4.02 (br s, 1H), 3.82-3.87 (m, 2H), 3.12-3.22 (m, 2H), 2.93-3.03 (m, 2H), 1.86-2.10 (m, 2H), 1.24-1.72 (m, 8H).	390.2 (C ₁₃ H ₂₁ N ₅ O ₇ S)
17.			10.30 (s, 1H), 10.14 (s, 1H), 9.00 (br d, 1H), 8.70-8.75 (m, 1H), 4.02 (br s, 1H), 3.81-3.87 (m, 2H), 3.17-3.21 (m, 2H), 2.93-3.03 (m, 2H), 2.10 (br d, 1H), 1.98-2.05 (m, 1H), 1.85-1.92 (m, 1H), 1.69-1.77 (m, 3H), 1.46-1.66 (m, 4H).	390.3 (C ₁₃ H ₂₁ N ₅ O ₇ S)
18.			10.27 (s, 1H), 10.17 (s, 1H), 8.93 (br d, 1H), 8.76-8.78 (m, 1H), 4.02 (br s, 1H), 3.79-3.86 (m, 2H), 3.18-3.23 (m, 2H), 2.93-3.02 (m, 2H), 2.14 (br d, 1H), 1.97-2.05 (m, 1H), 1.83-1.93 (m, 1H), 1.46-1.77 (m, 7H).	390.3 (C ₁₃ H ₂₁ N ₅ O ₇ S)
19.			10.00 (d, 1H), 9.68 (br s, 1H), 4.01 (br s, 1H), 3.83 (d, 1H), 3.42 (s, 1H), 3.12-3.21 (m, 6H), 2.93-3.02 (m, 4H), 1.98-2.05 (m, 1H), 1.84-1.92 (m, 1H), 1.68-1.72 (m, 1H), 1.54-1.62 (m, 1H).	404.9 (C ₁₃ H ₂₂ N ₆ O ₇ S)
20.			10.44 (d, 1H), 10.18 (d, 1H), 8.67 (br d, 3H), 7.43-7.55 (m, 5H), 4.98 (br s, 1H), 4.01 (s, 1H), 3.83 (br d, 1H), 3.19 (d, 1H), 3.00 (br d, 1H), 1.85-2.05 (m, 2H), 1.59-1.72 (m, 2H).	411.9 (C ₁₅ H ₁₉ N ₅ O ₇ S)

(continued)

Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	H ¹ NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
21.	(RS)-t-Boc-HNCH(CH ₃)CH ₂ -CONHNH ₂	(RS)-H ₂ NCH(CH ₃)CH ₂ -CO-	10.00 (s, 1H), 9.95 (d, 1H), 7.74 (br s, 3H), 4.01 (br s, 1H), 3.83 (d, 1H), 3.53-3.48 (m, 1H), 3.12-3.19 (m, 1H), 3.00 (br d, 1H), 2.36-2.42 (m, 1H), 1.98-2.06 (m, 1H), 1.78-1.87 (m, 1H), 1.66-1.76 (m, 1H), 1.54-1.62 (m, 2H), 1.19 (d, 3H)	364.1 (C ₁₁ H ₁₉ N ₅ O ₇ S)
22.			9.92 (s, 1H), 9.82 (s, 1H), 7.71 (br s, 3H), 4.01 (br s, 1H), 3.84 (d, 1H), 3.21 (d, 1H), 3.01 (br d, 1H), 2.85-2.92 (m, 2H), 1.98-2.04 (m, 1H), 1.85-1.87 (m, 1H), 1.69-1.73 (m, 1H), 1.57-1.60 (m, 1H), 1.22 (d, 6 H).	378.2 (C ₁₂ H ₂₁ N ₅ O ₇ S)
23.			9.82 (s, 1H), 7.76 (br s, 3H), 4.00 (br s, 1H), 3.81 (d, 1H), 3.22 (d, 1H), 2.00 (d, 2H), 2.88 (d, 1H), 1.97-2.05 (m, 1H), 1.74-1.85 (m, 1H), 1.16-1.73 (m, 1H), 1.54-1.61 (m, 1H), 1.27 (t, 2H) 1.01-1.08 (m, 2H).	376.2 (C ₁₂ H ₁₉ N ₅ O ₇ S)
24.			10.37 (s, 1H), 10.16 (s, 1H), 8.24 (br s, 3H), 7.39 (br s, 1H), 6.95 (br s, 1H), 4.02 (br s, 1H), 3.86 (d, 1H), 3.19 (d, 1H), 3.02 (br d, 1H), 2.20-2.34 (m, 2H), 1.92-2.05 (m, 4H), 1.67-1.79 (m, 1H), 1.55-1.63 (m, 1H).	407.2 (C ₁₂ H ₂₀ N ₆ O ₈ S)
25.	t-Boc-HN-(CH ₂) ₄ -CONHNH ₂	H ₂ N(CH ₂) ₄ -CO-	9.18 (s, 1H), 9.70 (s, 1H), 7.58 (br s, 3H), 3.99 (br s, 1H), 3.80 (d, 1H), 3.21 (d, 1H), 3.99 (br d, 1H), 2.76-(br s, 2H), 2.15-2.20 (m, 1H), 1.97-2.01 (m, 1H), 1.82-1.85 (m, 1H), 1.67-1.73 (m, 1H), 1.55-1.62 (m, 6H).	378.2 (C ₁₂ H ₂₁ N ₅ O ₇ S)
26.			10.16 (s, 1H), 7.86 (br s, 6H), 4.03 (br s, 1H), 3.88 (d, 1H), 3.77 (t, 1H), 3.16 (d, 1H), 3.02 (br d, 1H), 2.73 (t, 2H), 1.99-2.05 (m, 1H), 1.72-1.76 (m, 1H), 1.62-1.71 (m, 3H), 1.13-1.60 (m, 6H).	407.3 (C ₁₃ H ₂₄ N ₆ O ₇ S.CF ₃ COOH)

(continued)

Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	H ¹ NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
27.			9.97 (s, 1H), 9.71 (s, 1H), 7.74 (br s, 3H), 4.04 (s, 2H), 4.01 (br s, 1H), 3.83 (d, 1H), 3.64 (t, 2H), 3.20 (d, 1H), 3.00-3.05 (m, 3H), 1.98-2.05 (m, 1H), 1.82-1.86 (m, 1H), 1.68-1.73 (m, 1H), 1.58-1.63 (m, 1H).	380.2 (C ₁₁ H ₁₉ N ₅ O ₈ S)
28.			10.03 (br s, 1H), 8.66 (br s, 1H), 4.05-4.12 (m, 1H), 3.96-4.01 (m, 4H), 3.84 (d, 1H), 3.53-3.58 (m, 1H), 3.21 (d, 1H), 3.01 (br d, 1H), 1.98-2.03 (m, 1H), 1.72-1.85 (m, 1H), 1.68-1.71 (m, 1H), 1.57-1.62 (m, 1H).	362.2 (C ₁₁ H ₁₇ N ₅ O ₇ S)
29.			9.49 (s, 1H), 8.08 (s, 1H), 3.98 (br s, 1H), 3.76 (d, 1H), 3.35 (d, 2H), 3.20-3.25 (m, 4H), 2.96 (br d, 1H), 1.86-2.06 (m, 1H), 1.57-1.80 (m, 7H).	376.1 (C ₁₂ H ₁₉ N ₅ O ₇ S)
30.			10.35 (br s, 1H), 10.18 (s, 1H), 8.96 (br s, 2H), 4.18 (t, 1H), 4.01 (br s, 1H), 3.86 (d, 1H), 3.17-3.25 (m, 3H), 3.01 (br d, 1H), 2.31-2.35 (m, 1H), 1.98-2.03 (m, 1H), 1.76-1.91 (m, 4H), 1.66-1.74 (m, 1H), 1.57-1.62 (m, 1H).	376.2 (C ₁₂ H ₁₉ N ₅ O ₇ S)
31.			9.88 (s, 1H), 9.81 (s, 1H), 8.77 (br s, 1H), 8.29 (br s 1H), 4.00 (br s, 1H), 3.81 (d, 1H), 3.37-3.43 (m, 1H), 3.12-3.19 (m, 4H), 2.63 (br d, 1H), 2.08-2.24 (m, 2H), 1.69-2.06 (m, 6H), 1.47-1.61 (m, 3H).	404.2 (C ₁₄ H ₂₃ N ₅ O ₇ S)
32.			10.02 (d, 1H), 9.96 (s, 1H), 8.80 (br s, 1H), 8.68 (br s, 1H), 4.00 (br s, 1H), 3.83 (d, 1H), 3.08-3.17 (m, 6H), 3.00 (br d, 1H), 2.06-2.21 (m, 1H), 1.98-2.06 (m, 2H), 1.78-1.82 (m, 1H), 1.67-1.75 (m, 1H), 1.53-1.65 (m, 1H).	376.2 (C ₁₂ H ₁₉ N ₅ O ₇ S)
33.			10.03 (s, 1H), 9.96 (s, 1H), 8.80 (br s, 1H), 8.68 (br s, 1H), 4.00 (br s, 1H), 3.82 (d, 1H), 3.26-3.37 (m, 4H), 3.17-3.25 (m, 1H), 3.06-3.10 (m, 1H), 3.00 (br d, 1H), 2.16-2.23 (m, 1H), 2.00-2.05 (m, 1H), 1.86-1.85 (m, 1H), 1.65-1.75 (m, 1H), 1.53-1.61 (m, 1H).	376.1 (C ₁₂ H ₁₉ N ₅ O ₇ S)

(continued)

Table 1.

Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	¹ H NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
34.			10.44 (s, 1H), 10.17 (s, 1H), 9.62 (br s, 1H), 8.92 (br s, 1H), 5.48-5.52 (m, 1H), 4.44 (br s, 1H), 4.33 (t, 1H), 4.01 (s, 1H), 3.86 (d, 1H), 3.19 (d, 1H), 3.00-3.10 (m, 2H), 2.26-2.31 (m, 1H), 1.87-2.06 (m, 4H), 1.66-1.75 (m, 1H), 1.54-1.62 (m, 1H).	392.3 (C ₁₂ H ₁₉ N ₅ O ₈ S)
35.			10.28 (s, 1H), 9.11 (br s, 1H), 8.05 (br s, 2H), 4.29 (t, 1H), 4.03 (br s, 1H), 3.87-3.95 (m, 2H), 3.48-3.53 (m, 2H), 3.04-3.24 (m, 3H), 3.02 (br d, 1H), 2.72-2.79 (m, 1H), 1.99-2.04 (m, 1H), 1.88-1.93 (m, 2H), 1.69-1.78 (m, 1H), 1.54-1.63 (m, 1H).	391.2 (C ₁₀ H ₂₀ N ₆ O ₇ S)
36.			10.48 (br s, 1H), 10.20 (s, 1H), 7.73 (d, 1H), 7.32 (br s, 4H), 4.16-4.26 (m, 2H), 4.03 (br s, 1H), 3.87 (d, 1H), 3.51-3.56 (m, 1H), 3.01-3.17 (m, 3H), 2.74-2.86 (m, 1H), 1.99-2.05 (m, 1H), 1.55-1.86 (m, 5H).	433.2 (C ₁₃ H ₂₂ N ₈ O ₇ S)
37.			D ₂ O exchange: 3.98 (br s, 1H), 3.85 (d, 1H), 3.17 (br d, 1H), 3.02-3.11 (m, 2H), 2.92-2.98 (m, 1H), 2.77 (t, 1H), 2.25-2.28 (m, 2H), 2.00-2.04 (m, 1H), 1.75-1.86 (m, 3H), 1.60-1.71 (m, 4H), 1.23-1.47 (m, 4H).	418.0 (C ₁₅ H ₂₅ N ₅ O ₇ S)
38.			10.54 (br s, 1H), 10.22 (d, 1H), 8.96 (br s, 2H), 4.09 (d, 1H), 4.02 (br s, 1H), 3.87 (d, 1H), 3.62 (t, 2H), 3.38 (t, 3H), 3.01-3.17 (m, 4H), 1.98-2.05 (m, 1H), 1.86-1.88 (m, 1H), 1.68-1.75 (m, 1H), 1.55-1.62 (m, 1H).	391.3 (C ₁₂ H ₂₀ N ₆ O ₇ S.CF ₃ COOH)
39.			10.46 (s, 1H), 10.18 (s, 1H), 9.37 (br d, 2H), 4.13 (br d, 2H), 4.02 (br s, 1H), 3.86 (br d, 2H), 3.57-3.62 (m, 2H), 3.17 (br d, 3H), 3.02 (br d, 1H), 1.98-2.04 (m, 1H), 1.85-1.94 (m, 1H), 1.66-1.76 (m, 1H), 1.56-1.61 (m, 1H).	392.2 (C ₁₂ H ₁₉ N ₅ O ₈ S)


(continued)

Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	H ¹ NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
40.			D ₂ O exchange: 3.97 (br s, 1H), 3.83 (d, 1H), 3.59-3.68 (m, 4H), 3.40 (s, 2H), 3.03-3.06 (m, 6H), 1.98-2.04 (m, 1H), 1.75-1.84 (m, 1H), 1.65-1.73 (m, 1H), 1.57-1.61 (m, 1H).	433.3 (C ₁₄ H ₂₂ N ₆ O ₈ S)
41.			10.62 (s, 1H), 10.23 (d, 1H), 8.72 (br s, 3H), 8.62 (br s, 1H), 7.94 (t, 1H), 7.76 (dd, 1H), 7.47 (t, 1H), 5.09-5.10 (m, 1H), 4.00 (br s, 1H), 3.84 (br d, 1H), 3.14-3.21 (m, 1H), 2.99 (br d, 1H), 1.94-2.02 (m, 1H), 1.82-1.91 (m, 1H), 1.66-1.76 (m, 1H), 1.57-1.62 (m, 1H).	413.2 (C ₁₉ H ₁₈ N ₆ O ₇ S.CF ₃ COOH)
42.			D ₂ O exchange: 7.33 (s, 1H), 4.00 (br s, 1H), 3.83 (d, 1H), 3.28 (d, 1H), 3.00 (br d, 1H), 2.00-2.05 (m, 1H), 1.76-1.86 (m, 1H), 1.67-1.75 (m, 1H), 1.59-1.64 (m, 1H).	405.1 (C ₁₁ H ₁₄ N ₆ O ₇ S ₂)

Example- 43

Sodium salt of trans-sulfuric acid mono-[2-(N'-(cyano-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester

[0148]



[0149] Tetrabutylammonium salt of trans-sulfuric acid mono-[2-(N'-(cyano-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester (600 mg, obtained using procedure described in Step-1 to Step-3 of Example 1 (using cyano acetic acid hydrazide in the place of (S)-N-tert-butoxycarbonyl-pyrrolidine-2-carboxylic acid hydrazide) was loaded in tetrahydrofuran and water 1:9 mixture (10 ml) and passed slowly through freshly activated Amberlite 200C resin in sodium form (100 gm). The column was eluted with 10% tetrahydrofuran in water mixture. The fractions were analyzed on TLC and desired fractions were evaporated to remove volatile solvent under vacuum below 40°C. The aqueous layer was then washed with dichloromethane (25 ml x 2) and layers were separated. The aqueous layer was concentrated under vacuum below 40°C to provide residue, which was azeotroped with acetone and triturated with diethyl ether to provide a suspension. The suspension was filtered to provide title compound (Sodium salt of trans-sulfuric acid mono-[2-(N'-(2-cyano-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester) in 300 mg quantity and in 81% yield.

Analysis: MS (ES-) $C_{10}H_{12}N_5O_7SNa$, 346.2 (M-1) as a free sulfonic acid;

1H NMR (DMSO- d_6) = 10.2 (s, 1H), 10.05 (s, 1H), 3.99 (s, 1H), 3.82 (d, 1H), 3.72 (s, 1H), 3.36 (s, 1H), 3.14 (br d, 1H), 2.99 (d, 1H), 1.98-2.03 (m, 1H), 1.75-1.84 (m, 1H), 1.56-1.72 (m, 2H).

[0150] Compounds **44 to 52** (Table 2) were using a procedure described in Example-16 and using corresponding $R_1CONHNH_2$, in the place of cyano acetic acid hydrazide.
















Table 2





Example No.	Acid hydrazide ($R_1CONHNH_2$)	R_1	1H NMR (DMSO- d_6)	Mass (ES-1) as free acid (MF)
44.	t-Boc-NHNNH ₂	t-C ₄ H ₉ -O-CO-	9.70 (s, 1H), 8.71 (s, 1H), 3.98 (s, 1H), 3.73 (d, 1H), 3.13 (d, 1H), 2.97 (br d, 1H), 1.97-2.01 (m, 1H), 1.73-1.83 (m, 1H), 1.55-2.72 (m, 2H), 1.38 (s, 9H).	379.3 (C ₁₂ H ₁₉ N ₄ O ₈ SNa)

(continued)

Table 2

Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	¹ H NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
45.			10.02 (br s, 2H), 3.98 (br s, 1H), 3.85 (d, 1H), 3.66-2.70 (m, 4H), 3.37 (br s, 2H), 3.05 (s, 2H), 2.76 (br s, 4H), 2.00-2.04 (m, 1H), 1.73-1.82 (m, 1H), 1.58-1.70 (m, 2H).	406.3 (C ₁₃ H ₂₀ N ₅ O ₈ SNa)
46.			11.00 (s, 1H), 10.17 (s, 1H), 8.91 (s, 1H), 8.17-8.20 (m, 3H), 7.74 (s, 1H), 4.04 (br s, 1H), 3.91 (d, 1H), 3.06 (br d, 1H), 1.90-2.06 (m, 2H), 1.77-1.87 (m, 1H), 1.61-1.77 (m, 2H).	426.9 (C ₁₄ H ₁₅ N ₆ O ₈ SNa)
47.			10.45 (br s, 1H), 10.08 (br s, 1H), 3.99 (br s, 1H), 3.81 (br d, 1H), 3.56-3.58 (m, 4H), 3.41-3.51 (m, 2H), 2.99-3.10 (m, 2H), 2.06 (br s, 2H), 2.01-2.03 (m, 1H), 2.82-1.83 (m, 1H), 1.59-1.72 (m, 2H).	419.9 (C ₁₃ H ₁₈ N ₅ O ₉ SNa)
48.			9.81 (br s, 1H), 9.69 (br s, 1H), 5.77 (s, 2H), 4.21 (t, 1H), 3.98 (br s, 1H), 3.77 (d, 1H), 3.34-3.38 (m, 1H), 3.16-3.29 (m, 2H), 2.96 (br d, 1H), 1.73-1.98 (m, 6H), 1.58-1.72 (m, 2H).	419.2 (C ₁₃ H ₁₉ N ₆ O ₈ S.Na)
49.			9.77 (br s, 2H), 5.95 (s, 2H), 5.32 (s, 0.5H), 5.18 (br s 0.5H), 4.38 (dd, 1H), 3.98 (br s, 1H), 3.81 (d, 1H), 3.35-3.59 (m, 2H), 2.97-3.25 (m, 1H), 2.96 (br d, 1H), 1.99-2.39 (m, 2H), 1.82-1.95 (m, 1H), 1.75-1.82 (m, 1H), 1.59-1.69 (m, 2H).	437.2 (C ₁₃ H ₁₈ N ₆ O ₈ FS.Na)
50.			9.83 (br s, 2H), 7.32 (s, 1H), 4.21 (dd, 1H), 3.98 (br s, 1H), 3.77 (d, 1H), 3.38-3.44 (m, 1H), 3.21-3.25 (m, 1H), 2.90-2.98 (m, 4H), 1.56-2.12 (m, 8H).	454.2 (C ₁₃ H ₂₀ N ₅ O ₉ S ₂ .Na)

(continued)

Table 2				
Example No.	Acid hydrazide (R ₁ CONHNH ₂)	R ₁	H ¹ NMR (DMSO-d ₆)	Mass (ES-1) as free acid (MF)
51.			D ₂ O exchange: 3.95-4.05 (m, 2H), 3.10-3.19 (m, 1H), 2.95-3.05 (m, 1H), 1.68-1.98 (m, 4H), 1.29 (s, 3H), 1.21 (s, 3H).	374.2 (C ₁₂ H ₁₆ N ₅ O ₇ S.Na)
52.			9.95 (s, 2H), 7.93 (s, 1H), 4.03-4.05 (m, 1H), 3.99 (br s, 1H), 3.81 (d, 1H), 3.19 (d, 1H), 2.98 (br d, 1H), 2.28-2.48 (m, 1H), 1.84-2.25 (m, 5H), 1.56-1.75 (m, 2H).	

[0151] Compounds of the invention from Example 1 to 52 were prepared using (S)-pyrroglutamic acid as a starting compound. The absolute stereochemistry is therefore (2S, 5R) 7-oxo-1,6-diaza-bicyclo[3.2.1]octane ring. Thus, the compound of Example-2, trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester has the absolute stereochemistry as (2S,5R)-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester. Alternatively, if the starting compound used is (R)-pyrroglutamic acid the resulting compounds will have (2R, 5S) stereochemistry in 7-oxo-1,6-diaza-bicyclo[3.2.1]octane ring. A reference to a compound according to the invention also includes corresponding compounds having (2S, 5R) and (2R, 5S) stereochemistry.

Biological Activity

[0152] The biological activity of representative compounds according to the invention against various bacterial strains was investigated. In a typical study, overnight grown bacterial cultures were diluted appropriately and inoculated on the agar media containing doubling dilutions of the test compounds. Observation for growth or no growth was performed after 16-20 hours of incubation at 35 ± 2°C in ambient air. The overall procedure was performed as per Clinical and Laboratory Standards Institute (CLSI) recommendations (Clinical and Laboratory Standards Institute (CLSI), Performance Standards for Antimicrobial Susceptibility Testing, 20th Informational Supplement, M 100 - S20, Volume 30, No. 1, 2010). The results of these studies are summarized in Tables 3 to 8.

[0153] Table 3 details antibacterial activity of representative compounds according to the invention against various E. coli strains (NCTC 13351, M 50 and 7 MP) expressing ESBL (Extended Spectrum Beta Lactamases).

Table 3. Antibacterial activity of representative compounds according to the invention (MIC expressed in mcg/ml)				
Sr.	Compound of Example No.	Bacterial Strain		
		E. coli NCTC 13351	E. coli M50	E. coli 7 MP
1.	1	0.025	0.25	0.25
2.	2	0.25	0.25	1
3.	3	0.25	0.25	1
4.	4	8	8	16
5.	5	0.25	0.25	0.5
6.	6	32	64	128
7.	7	2	2	4
8.	8	1	1	2
9.	9	1	2	4

(continued)

Table 3. Antibacterial activity of representative compounds according to the invention (MIC expressed in mcg/ml)

Sr.	Compound of Example No.	Bacterial Strain		
		E. coli NCTC 13351	E. coli M50	E. coli 7 MP
10.	10	1	8	32
11.	11	2	1	4
12.	12	1	8	32
13.	13	1	1	2
14.	14	0.5	0.25	2
15.	15	8	4	16
16.	16	1	1	2
17.	17	0.5	0.5	2
18.	18	8	4	32
19.	19	1	1	2
20.	20	2	2	8
21.	21	1	1	4
22.	22	4	4	8
23.	23	4	4	8
24.	24	1	1	4
25.	25	1	1	4
26.	26	1	1	2
27.	27	4	2	8
28.	28	1	1	4
29.	29	32	16	128
30.	30	16	8	32
31.	31	1	1	4
32.	32	0.5	0.5	1
33.	33	1	0.5	2
34.	34	2	2	8
35.	35	1	1	4
36.	36	1	1	2
37.	37	2	2	4
38.	38	4	2	8
39.	39	4	2	8
40.	40	1	1	4
41.	41	4	4	16
42.	42	32	32	64
43.	43	1	2	8
44.	44	32	32	64
45.	45	4	8	8

(continued)

Table 3. Antibacterial activity of representative compounds according to the invention (MIC expressed in mcg/ml)

Sr.	Compound of Example No.	Bacterial Strain		
		E. coli NCTC 13351	E. coli M50	E. coli 7 MP
46.	46	32	64	128
47.	47	32	64	128
48.	48	16	8	32
49.	49	32	16	16
50.	50	16	16	32
51.	51	16	16	32
52.	52	0.5	0.5	2

[0154] Tables 4 and 5 provide antibacterial activity of representative compounds according to the invention against various Multi Drug Resistant (MDR) Gram-negative bacterial strains expressing various ESBLs. The activities are expressed as MICs (mcg/ml). For comparison, the activity of several known antibacterial agents (for example, Ceftazidime, Aztreonam, Imipenem, Ciprofloxacin and Tigecycline) are also included. As can be seen, the representative compounds according to the invention exhibit antibacterial activity against various MDR strains. The data in Table 4 and 5 also indicates that the compounds according to the invention exhibit potent activity against a wide variety of bacteria, even against those producing different types of beta-lactamase enzymes. In general, the activity of the compounds according to the invention against various beta-lactamase producing bacterial strains is even better than the other antibacterial agents currently employed in the clinical practice to treat such infections.

[0155] The antibacterial activity of representative compounds according to the invention was also investigated in combination with at least one antibacterial agent using the above study protocol and the results are given Table 6. As can be seen, the use of compounds according to the invention significantly lowered MIC values of the antibacterial agent (e.g. in this case Ceftazidime). The results also suggest the compounds according to the invention increase antibacterial effectiveness of the antibacterial agent when said antibacterial agent is co-administered with a pharmaceutically effective amount of a compound of Formula (I) or a stereoisomer or a pharmaceutically acceptable salt thereof.

[0156] The antibacterial activity of representative compounds according to the invention was also investigated in combination with a beta-lactamase inhibitor using the above study protocol and the results are given in Table 7. As can be seen, the compounds according to the invention in combination with a beta lactam inhibitor exhibited excellent antibacterial activity against various bacterial strains. For example, a combination comprising compound of Example 2 (and also Example 3) according to the invention in combination with sulbactam exhibited much superior MIC values compared with when these were used alone. The results also suggest that compounds according to the invention in combination with a beta-lactam inhibitor can be effectively used in preventing or treating a bacterial infection in a subject, including those infections caused by bacteria producing one or more beta-lactamase enzymes.

[0157] The antibacterial activity of representative compounds according to the invention was also investigated in combination with a beta-lactamase inhibitor and an antibacterial agent using the above study protocol and the results are given in Table 8. As can be seen, the compounds according to the invention in combination with at least one beta lactam inhibitor and at least one antibacterial agent exhibited excellent antibacterial activity against various bacterial strains. For example, a combination comprising compound of Example 2 (and also Example 3) according to the invention in combination with sulbactam and Cefepime exhibited better MIC values compared with when these were used alone. The results also suggest that compounds according to the invention in combination with at least one beta-lactam inhibitor and at least one antibacterial agent can be effectively used in preventing or treating a bacterial infection in a subject, including those infections caused by bacteria producing one or more beta-lactamase enzymes.

Sr.	Compound	Class A ESBL				Class C ESBL				KPC ESBL			
		E. Coli W 13353	E. Coli W 13351	E. Coli W 13352	E. Coli M 50	E. Coli H 484	E. Coll B 89	K pneumoniae H 521	K pneumoniae H 523	K pneumoniae H 525			
1.	Ceftazidime	32	32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	
2.	Aztreonam	> 32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	> 32	
3.	Imipenem	0.25	0.25	0.25	0.5	4	0.5	16	16	16	32	16	
4.	Ciprofloxacin	> 32	0.5	0.12	> 32	> 32	> 32	32	8	32	32	32	
5.	Tigecyclin	1	1	0.25	0.5	0.25	0.5	2	8	2	2	2	
6.	Example 1	0.25	0.5	0.5	0.5	2	0.5	0.5	0.5	0.5	0.5	0.5	
7.	Example 2	0.12	0.25	0.25	0.25	0.12	0.25	0.5	1	0.5	0.5	0.5	
8.	Example 3	0.25	0.5	0.5	0.5	0.5	0.25	0.5	0.5	0.5	0.5	0.5	
9.	Example 5	0.5	0.25	0.5	1	1	0.5	1	1	1	2	2	
10.	Example 7	0.5	2	1	2	4	4	2	1	2	2	2	
11.	Example 8	1	1	1	1	2	1	1	0.5	1	1	1	
12.	Example 9	1	1	1	1	2	1	1	0.5	0.5	0.5	0.5	
13.	Example 11	2	2	2	2	16	2	4	2	2	2	2	
14.	Example 13	1	1	2	1	4	1	1	0.5	1	1	1	
15.	Example 14	0.5	0.5	0.5	0.5	2	0.5	0.5	0.25	0.25	0.25	0.25	
16.	Example 17	0.25	0.5	0.5	0.5	4	0.5	0.5	0.5	0.5	0.5	0.5	
17.	Example 19	2	2	2	2	4	1	1	1	1	1	1	
18.	Example 30	8	16	16	8	8	8	8	8	> 32	> 32	> 32	
19.	Example 35	1	1	1	1	1	1	2	1	1	1	1	
20.	Example 38	2	4	4	2	8	2	2	2	2	2	2	
21.	Example 43	1	1	1	1	8	1	2	1	1	1	1	

Table 5. Comparative antibacterial activity of representative compounds according to the invention against various Multi Drug Resistant (MDR) Gram negative strains (expressed as MICs (mcg/ml)).

Sr.	Compound	Class B ESBL			P. aeruginosa		
		K. pneumoniae S 48	E. Coli M 3	E. Coli M 44	ATCC 27853	Ps 21	Ps 32
1.	Ceftazidime	> 32	> 32	> 32	1	> 32	> 32
2.	Aztreonam	> 32	> 32	> 32	2	8	8
3.	Imipenem	16	8	32	4	> 32	> 32
4.	Ciprofloxacin	> 32	> 32	> 32	0.5	32	0.12
5.	Tigecyclin	1	4	0.25	16	16	16
6.	Example 1	2	1	0.25	16	16	32
7.	Example 2	0.5	2	0.5	8	8	8
8.	Example 3	0.5	0.5	0.12	2	4	4
9.	Example 5	2	1	0.5	> 32	> 32	> 32
10.	Example 7	4	4	2	> 32	> 32	> 32
11.	Example 8	1	4	0.5	8	16	8
12.	Example 9	1	2	0.5	8	8	16
13.	Example 11	2	8	1	> 32	> 32	> 32
14.	Example 13	1	1	1	32	32	16
15.	Example 14	4	0.5	0.25	8	16	8
16.	Example 17	0.5	0.5	0.5	> 32	> 32	> 32
17.	Example 19	4	1	1	32	> 32	> 32
18.	Example 30	> 32	8	16	> 32	> 32	> 32
19.	Example 35	2	1	2	> 32	> 32	32
20.	Example 38	4	1	4	> 32	> 32	> 32
21.	Example 43	1	4	0.5	> 32	> 32	> 32

Table 6. Antibacterial activity of Ceftazidime in presence of representative compounds of the invention against various Multi Drug Resistant (MDR) Gram negative strains.

Sr.	Composition	Ceftazidime MIC (expressed in mcg/ml)	
		K. pneumoniae ATCC 700603 (ESBL type: Class A)	P. vulgaris S-137B (ESBL type: Class C)
1.	Ceftazidime alone	> 32	> 32
2.	Ceftazidime + Compound of Example 1 (4 mcg/ml)	0.06	1
3.	Ceftazidime + Compound of Example 2 (4 mcg/ml)	0.06	1
4.	Ceftazidime + Compound of Example 3 (4 mcg/ml)	0.06	1
5.	Ceftazidime + Compound of Example 5 (4 mcg/ml)	0.25	4

(continued)

Table 6. Antibacterial activity of Ceftazidime in presence of representative compounds of the invention against various Multi Drug Resistant (MDR) Gram negative strains.

Sr.	Composition	Ceftazidime MIC (expressed in mcg/ml)	
		K. pneumoniae ATCC 700603 (ESBL type: Class A)	P. vulgaris S-137B (ESBL type: Class C)
6.	Ceftazidime + Compound of Example 8 (4 mcg/ml)	0.06	0.5
7.	Ceftazidime + Compound of Example 9 (4 mcg/ml)	0.12	1
8.	Ceftazidime + Compound of Example 14 (4 mcg/ml)	0.03	0.5

Note: The MICs of each of compounds of Example 1, 2, 3, 5, 8, 9 and 14 when used alone (in the absence of Ceftazidime) is > 32 mcg/ml.

Table 7. Antibacterial activity of sulbactam in combination with a compound according to the invention.

Sr.	Composition	MIC of sulbactam (expressed in mcg/ml)			
		A. baumannii J-143	A. baumannii 1460648	A. baumannii S-334	A. baumannii G-165
1.	Sulbactam alone	32	32	32	32
2.	Sulbactam + compound of Example 2 (4 mcg/ml)	4	4	4	2
3.	Sulbactam + compound of Example 2 (8 mcg/ml)	4	4	2	2
4.	Sulbactam + compound of Example 3 (4 mcg/ml)	8	4	8	2
5.	Sulbactam + compound of Example 3 (8 mcg/ml)	1	2	4	2

Standalone MIC of compound of Example 2 and Example 3 for each of the strains was > 32 mcg/ml.

Table 8. Antibacterial activity of an antibacterial agent in combination with sulbactam and a compound according to the invention.

Sr.	Composition	MIC of Cefepime (expressed in mcg/ml)			
		A. baumannii J-143	A. baumannii 1460648	A. baumannii S-334	A. baumannii G-165
1.	Cefepime alone	> 32	> 32	32	> 32
2.	Cefepime + compound of Example 2 (4 mcg/ml)	> 32	32	32	> 32
3.	Cefepime + compound of Example 2 (8 mcg/ml)	32	32	32	32
4.	Cefepime + Sulbactam (4 mcg/ml)	16	32	16	16
5.	Cefepime + Sulbactam (8 mcg/ml)	16	16	8	8


(continued)

Table 8. Antibacterial activity of an antibacterial agent in combination with sulbactam and a compound according to the invention.

Sr.	Composition	MIC of Cefepime (expressed in mcg/ml)			
		A. baumannii J-143	A. baumannii 1460648	A. baumannii S-334	A. baumannii G-165
6.	Cefepime + Sulbactam (8 mcg/ml) + compound of Example 2 (4 mcg/ml)	0.25	0.25	0.25	0.25
7.	Cefepime + Sulbactam (8 mcg/ml) + compound of Example 2 (8 mcg/ml)	0.25	0.25	0.25	0.25
8.	Cefepime + compound of Example 3 (4 mcg/ml)	> 32	32	32	> 32
9.	Cefepime + compound of Example 3 (8 mcg/ml)	32	32	32	32
10.	Cefepime + Sulbactam (8 mcg/ml) + compound of Example 3 (4 mcg/ml)	0.12	0.12	0.12	0.12
11.	Cefepime + Sulbactam (8 mcg/ml) + compound of Example 3 (8 mcg/ml)	0.06	0.06	0.06	0.06
Standalone MIC of sulbactam for each of the strains was 32 mcg/ml. Standalone MIC of compound of Example 2 and Example 3 for each of the strains was > 32 mcg/ml.					

Claims

1. A compound of Formula (I)



or a stereoisomer or a pharmaceutically acceptable salt thereof; wherein:

R₁ is:

- (a) hydrogen,
- (b) (CO)_n-R₃,
- (c) COOR₄, or
- (d) COCH₂COR₃

n is 0, 1 or 2;

R₂ is:

- (a) SO₃M,
- (b) SO₂NH₂,
- (c) PO₃M,
- (d) CH₂COOM,
- (e) CF₂COOM,
- (f) CHF₂COOM, or
- (g) CF₃.

M is hydrogen or a cation;

R₃ is:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₆, NR₅CONR₆R₇, heterocyclyl, heteroaryl, Cycloalkyl or aryl,
- (c) CN,
- (d) NR₆R₇,
- (e) CONR₆R₇,
- (f) NHCONR₆R₇,
- (g) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (h) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, NHC(NH)NR₆R₇, or NHCONR₆R₇,
- (i) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (j) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (k) cycloalkyl substituted with C₁-C₆ alkyl wherein C₁-C₆ alkyl is further substituted with one or more substituents independently selected from OR₅, NR₆R₇, halogen, CN, or CONR₆R₇, or
- (l) OR₆;

R₄ is:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (d) heterocyclyl optionally substituted with one or more substituents independently selected from C₅-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇,
- (e) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇, or
- (f) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₆, NR₆R₇, halogen, CN, CONR₆R₇, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₆R₇;

R₅ and R₈ are each independently:

- (a) hydrogen, or
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, CN, CONR₆R₇, NR₆R₇, heterocyclyl, heteroaryl, cycloalkyl or aryl;

R₆ and R₇ are each independently:

- (a) hydrogen,
- (b) C₁-C₆ alkyl optionally substituted with one or more substituents independently selected from halogen, OR₅, CN, COOR₅, CONR₅R₈, NR₅R₆, NR₅COR₈, heterocyclyl, heteroaryl, cycloalkyl or aryl,
- (c) aryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (d) heterocyclyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (e) heteroaryl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈,
- (f) cycloalkyl optionally substituted with one or more substituents independently selected from C₁-C₆ alkyl, OR₅, NR₅R₈, halogen, CN, CONR₅R₈, SO₂-alkyl, SO₂-aryl, OSO₂-alkyl, OSO₂-aryl, or NHCONR₅R₈, or
- (g) R₆ and R₇ are joined together to form a four to seven member ring.

2. A compound according to Claim 1, selected from:

trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-{7-oxo-2-[N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl]-1,8-diaza-bicyclo[3.2.1]oct-6-yloxy}-acetic acid;

trans-difluoro-{7-oxo-2-[N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-acetic acid;

trans-sulfuric acid mono-[2-hydrazinocarbonyl-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(amino-acetyl)hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(3-amino-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(4-amino-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S)-2-amino-3-hydroxy-propionyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S,4S)-4-fluoro-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S,4R)-4-methoxy-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(piperidin-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(S)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-piperidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(S)-piperidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(piperazin-4-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-1-amino-1-phenyl-acetyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(RS)-3-amino-butanoyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(3-amino-2,2-dimethyl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(1-aminomethyl-cyclopropan-1-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(2-amino-4-carboxamido-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-(5-amino-pentanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2S)-2,6-diamino-hexanoyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(2-aminoethoxy)-acetyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[azetidin-3-carbonyl]-hydrazinocarbonyl)-1-oxa-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[pyrrolidin-1-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
trans-sulfuric acid mono-[2-(N'-[(S)-3-pyrrolidin-2-yl]-propionyl)-hydrazinocarbonyl]-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;
5 trans-sulfuric acid mono-[2-(N'-[(RS)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
trans-sulfuric acid mono-[2-(N'-[(S)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
10 trans-sulfuric acid mono-[2-(N'-[(2S,4R)-4-hydroxy-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
trans-sulfuric acid mono-[2-(N'-[(2S,4S)-4-amino-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
trans-sulfuric acid mono-[2-(N'-[(2S,4S)-4-guanidino-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
15 trans-sulfuric acid mono-[2-(N'-[(RS)-3-piperidin-2-yl-propionyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;
trans-sulfuric acid mono-[2-(N'-[(RS)-piperazin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
trans-sulfuric acid mono-[2-(N'-[(S)-morpholin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
20 trans-sulfuric acid mono-[2-(N'-[(3-oxo-3-piperazin-1-yl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
trans-sulfuric acid mono-[2-(N'-[(RS)-1-amino-1-pyridin-2-yl-acetyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
25 trans-sulfuric acid mono-[2-(N'-[(2-amino-thiazol-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
or a stereoisomer or a pharmaceutically acceptable salt thereof.

3. A compound according to Claim 1, selected from:

30 Sodium salt of trans-sulfuric acid mono-[2-(N'-(cyano-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
Sodium salt of trans-N'-(7-oxo-6-sulfooxy-1,6-diaza-bicyclo[3.2.1]octane-2-carboxylic acid)-hydrazinecarboxylic acid tert-butyl ester;
35 Sodium salt of trans-sulfuric acid mono-[2-(N'-(morpholin-4-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
Sodium salt of trans-sulfuric acid mono-[2-(N'-(6-carboxamido-pyridin-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
Sodium salt of trans-sulfuric acid mono-[2-(N'-(morpholin-4-oxo-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
40 Sodium salt of trans-sulfuric acid mono-[2-(N'-[(S)-1-carbamoyl-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
Sodium salt of trans-sulfuric acid mono-[2-(N'-[(2S,4S)-1-carbamoyl-4-fluoro-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
45 Sodium salt of trans-sulfuric acid mono-[2-(N'-[(S)-1-methanesulfonyl-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
Sodium salt of trans-sulfuric acid mono-[2-(N'-(cyano-dimethyl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
50 Sodium salt of trans-sulfuric acid mono-[2-(N'-[(S)-5-oxo-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

or a stereoisomer thereof.

4. A pharmaceutical composition comprising a compound according to any of the Claims 1 to 3.

5. A pharmaceutical composition according to Claim 4, further comprising at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof.


6. A pharmaceutical composition according to Claim 4 or 5, further comprising at least one antibacterial agent or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof.
- 5 7. A pharmaceutical composition according to Claim 6, wherein the antibacterial agent is selected from a group consisting of aminoglycosides, ansamycins, carbacephems, cephalosporins, cephamycins, lincosamides, lipopeptides, macrolides, monobactams, nitrofurans, penicillins, polypeptides, quinolones, sulfonamides, tetracyclines, or oxazolidinone antibacterial agents.
- 10 8. A pharmaceutical composition according to Claim 6, wherein the antibacterial agent is a beta-lactam antibacterial agent.
9. A pharmaceutical composition according to Claim 6, wherein said antibacterial agent is selected from a group consisting of penicillins, penems, carbapenems, cephalosporins, and monobactams.
- 15 10. A pharmaceutical composition according to Claim 6, wherein the antibacterial agent is a cephalosporin antibiotic selected from a group consisting of cephalothin, cephaloridine, cefaclor, cefadroxil, cefamandole, cefazolin, cephalixin, cephadrine, ceftizoxime, cefoxitin, cephacetrile, cefotiam, cefotaxime, cefsulodin, cefoperazone, ceftizoxime, cefmenoxime, cefmetazole, cephaloglycin, cefonicid, cefodizime, cefpirome, ceftazidime, ceftiraxone, cefpiramide, cefbuparazone, cefozopran, cefepime, cefoselis, ceftuprenam, cefuzonam, cefpimizole, cefclidin, cefixime, ceftibuten, cefdinir, cefpodoxime axetil, cefpodoxime proxetil, ceftaram pivoxil, cefetamet pivoxil, cefcapene pivoxil or cefditoren pivoxil, cefuroxime, cefuroxime axetil, loracarbacef, ceftaroline and latamoxef.
- 20 11. A pharmaceutical composition according to Claim 6, wherein the antibacterial agent is selected from a group consisting of ceftazidime, cefepime, cefpirome, piperacillin doripenem, meropenem, imipenem, ceftaroline and ceftolozane.
- 25 12. A pharmaceutical composition according to Claim 6, wherein the antibacterial agent is selected from a group consisting of aminoglycosides, ansamycins, carbacephems, cephalosporins, cephamycins, lincosamides, lipopeptides, macrolides, monobactams, nitrofurans, penicillins, polypeptides, quinolones, sulfonamides, tetracyclines, or oxazolidinone antibacterial agents.
- 30 13. A pharmaceutical composition according to Claim 5, comprising (a) trans-sulfuric acid mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof.
- 35 14. A pharmaceutical composition according to Claim 5, comprising (a) trans-sulfuric acid mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) sulbactam or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof.
- 40 15. A compound according to any of the Claims 1 to 3 or a pharmaceutical composition for use according to any of Claims 4 to 14 in preventing or treating bacterial infection in a subject.
- 45 16. A compound according to any of the Claims 1 to 3 or a pharmaceutical composition according to any of Claim 4 to 14 for use in preventing or treating a bacterial infection in a subject according to claim 15, said infection being caused by bacteria producing one or more beta-lactamase enzymes.
- 50 17. A compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof for use in preventing or treating a bacterial infection in a subject, said use comprising administering to said subject a pharmaceutically effective amount of (a) a compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) at least one antibacterial agent or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof.
- 55 18. A compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof for use in preventing or treating a bacterial infection in a subject, said use comprising administering to said subject a pharmaceutically effective amount of, according to claim 17: (a) a compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof, and (b) an antibacterial agent, which is at least one

beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof.

19. A compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof for use in preventing or treating a bacterial infection in a subject, said use comprising administering to said subject a pharmaceutically effective amount of, according to claim 18: (a) a compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof; (b) an antibacterial agent, which is at least one beta-lactamase inhibitor selected from sulbactam, tazobactam, clavulanic acid, or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof, and (c) at least one antibacterial agent or a pharmaceutically acceptable salt, hydrate, solvate, polymorph or enantiomer thereof for use in preventing or treating bacterial infection in a subject.
20. A compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof for use according to claim 17, claim 18 or claim 19 in preventing or treating a bacterial infection in a subject, said infection being caused by bacteria producing one or more beta-lactamase enzymes.
21. A compound of Formula (I) according to Claim 1 or a stereoisomer or a pharmaceutically acceptable salt thereof for use in increasing antibacterial effectiveness of an antibacterial agent in a subject.

Patentansprüche

1. Verbindung der Formel (I)



Formel (I)

oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon, worin:

R₁ ist:

- (a) Wasserstoff,
- (b) (CO)_n-R₃,
- (c) COOR₄ oder
- (d) COCH₂COR₃,

wobei n 0, 1 oder 2 ist,

R₂ ist:

- (a) SO₃M,
- (b) SO₂NH₂,
- (c) PO₃M,
- (d) CH₂COOM,
- (e) CF₂COOM,
- (f) CHF₂COOM oder
- (g) CF₃,

worin M Wasserstoff oder ein Kation ist,

R₃ ist:

- (a) Wasserstoff,

(b) C₁-C₆-Alkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus Halogen, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, NR₅CONR₆R₇, Heterocyclyl, Heteroaryl, Cycloalkyl oder Aryl,

(c) CN,

(d) NR₆R₇,

(e) CONR₆R₇,

(f) NHCONR₆R₇,

(g) Aryl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇,

(h) Heterocyclyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl, NHC(NH)NR₆R₇ oder NHCONR₆R₇,

(i) Heteroaryl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇,

(j) Cycloalkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇,

(k) Cycloalkyl, substituiert mit C₁-C₆-Alkyl, worin C₁-C₆-Alkyl des Weiteren substituiert ist mit einem oder mehreren Substituenten, unabhängig ausgewählt aus OR₅, NR₆R₇, Halogen, CN oder CONR₆R₇ oder

(l) OR₈;

R₄ ist:

(a) Wasserstoff,

(b) C₁-C₆-Alkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus Halogen, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, Heterocyclyl, Heteroaryl, Cycloalkyl oder Aryl,

(c) Aryl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇,

(d) Heterocyclyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇,

(e) Heteroaryl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇, oder

(f) Cycloalkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₆R₇, Halogen, CN, CONR₆R₇, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₆R₇,

R₅ und R₈ jeweils unabhängig voneinander, sind:

(a) Wasserstoff oder

(b) C₁-C₆-Alkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus Halogen, CN, CONR₆R₇, NR₆R₇, Heterocyclyl, Heteroaryl, Cycloalkyl oder Aryl,

R₆ und R₇, jeweils unabhängig voneinander, sind:

(a) Wasserstoff,

(b) C₁-C₆-Alkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus Halogen, OR₅, CN, COOR₅, CONR₅R₈, NR₅R₈, NR₅COR₈, Heterocyclyl, Heteroaryl, Cycloalkyl oder Aryl,

(c) Aryl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₅R₈, Halogen, CN, CONR₅R₈, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₅R₈,

(d) Heterocyclyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausge-

wählt aus C₁-C₆-Alkyl, OR₅, NR₅R₈, Halogen, CN, CONR₅R₈, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₅R₈,

(e) Heteroaryl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₅R₈, Halogen, CN, CONR₅R₈, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₅R₈,

(f) Cycloalkyl, gegebenenfalls substituiert mit einem oder mehreren Substituenten, unabhängig ausgewählt aus C₁-C₆-Alkyl, OR₅, NR₅R₈, Halogen, CN, CONR₅R₈, SO₂-Alkyl, SO₂-Aryl, OSO₂-Alkyl, OSO₂-Aryl oder NHCONR₅R₈ oder

(g) R₆ und R₇ miteinander verbunden sind, um einen vier- bis siebengliedrigen Ring zu bilden.

2. Verbindung nach Anspruch 1, ausgewählt unter:

trans-Schwefelsäure-mono-[2-(N'-[(S)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-{7-Oxo-2-[N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-Essigsäure;

trans-Difluor-{7-oxo-2-[N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-Essigsäure;

trans-Schwefelsäure-mono-[2-hydrazinocarbonyl-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(amino-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(3-amino-propioyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(4-amino-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(2S)-2-amino-3-hydroxy-propioyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(2S,4S)-4-fluor-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(2S,4R)-4-methoxy-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(piperidin-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(RS)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(S)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(RS)-piperidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(S)-piperidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(R)-piperidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(piperazin-4-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(RS)-1-amino-1-phenyl-acetyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-[(RS)-3-amino-butanoyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(3-amino-2,2-dimethyl-propioyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(1-aminomethyl-cyclopropan-1-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;

trans-Schwefelsäure-mono-[2-(N'-(2-amino-4-carboxamido-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bi-

cyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-(5-amino-pentanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;
 trans-Schwefelsäure-mono-[2-(N'-((2S)-2,6-diamino-hexanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-((2-aminoethoxy)-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[azetidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;
 trans-Schwefelsäure-mono-[2-(N'-[pyrrolidin-1-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(R)-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(S)-3-pyrrolidin-2-yl]-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(RS)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(S)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(2S,4R)-4-hydroxy-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(2S,4S)-4-amino-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(2S,4S)-4-guanidino-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-[(RS)-3-piperidin-2-yl-propionyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl] ester;
 trans-Schwefelsäure-mono-[2-(N'-[(RS)-piperazin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-((S)-morpholin-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-(3-oxo-3-piperazin-1-yl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-((RS)-1-amino-1-pyridin-2-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 trans-Schwefelsäure-mono-[2-(N'-(2-amino-thiazol-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester;
 oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon.

3. Verbindung nach Anspruch 1, ausgewählt aus:

Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-(cyano-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-N'-(7-oxo-6-sulfooxy-1,6-diaza-bicyclo[3.2.1]octan-2-carbonsäure)-hydrazinocarbonsäure-tert-butyl-esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-(morpholin-4-yl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-(6-carboxamido-pyridin-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-(morpholin-4-oxo-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-[(S)-1-carbamoyl-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-[(2S,4S)-1-carbamoyl-4-fluor-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-[(S)-1-methansulfonyl-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters;
 Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-(cyano-dimethyl-acetyl)-hydrazinocarbonyl)-7-oxo-1,6-dia-

za-bicyclo[3.2.1]oct-6-yl]esters;

Natriumsalz des trans-Schwefelsäure-mono-[2-(N'-[(S)-5-oxo-pyrrolidin-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]esters oder ein Stereoisomer davon.


- 5 **4.** Pharmazeutische Zusammensetzung, umfassend eine Verbindung nach irgendeinem der Ansprüche 1 bis 3.
- 5.** Pharmazeutische Zusammensetzung nach Anspruch 4, des Weiteren umfassend mindestens einen beta-Lactamase-Inhibitor, ausgewählt aus Sulbactam, Tazobactam, Clavulansäure, oder ein pharmazeutisch annehmbares Salz, Hydrat, Solvat, Polymorph oder Enantiomer davon.
- 10 **6.** Pharmazeutische Zusammensetzung nach Anspruch 4 oder 5, des Weiteren umfassend mindestens ein antibakterielles Mittel oder ein pharmazeutisch annehmbares Salz, Hydrat, Solvat, Polymorph oder Enantiomer davon.
- 7.** Pharmazeutische Zusammensetzung nach Anspruch 6, wobei das antibakterielle Mittel ausgewählt ist aus der Gruppe, bestehend aus den antibakteriellen Mitteln Aminoglykosiden, Ansamycinen, Carbacephemem, Cephalosporinen, Cephamycinen, Lincosamiden, Lipopeptiden, Macroliden, Monobactamen, Nitrofuranen, Penicillinen, Polypeptiden, Chinolonen, Sulfonamiden, Tetracyclinen oder Oxazolidinon antibakterielle Mittel.
- 15 **8.** Pharmazeutische Zusammensetzung nach Anspruch 6, wobei das antibakterielle Mittel ein beta-Lactam-antibakterielles Mittel ist.
- 9.** Pharmazeutische Zusammensetzung nach Anspruch 6, wobei das antibakterielle Mittel ausgewählt ist aus der Gruppe, bestehend aus Penicillinen, Penemen, Carbapenemen, Cephalosporinen und Monobactamen.
- 20 **10.** Pharmazeutische Zusammensetzung nach Anspruch 6, wobei das antibakterielle Mittel ein Cefalosporin-Antibiotikum ist, ausgewählt aus der Gruppe, bestehend aus Cephalothin, Cephalexin, Cefaclor, Cefadroxil, Cefamandol, Cefazolin, Cephalexin, Cephadrin, Ceftizoxim, Cefoxitin, Cephacetril, Cefotiam, Cefotaxim, Cefsulodin, Cefoperazon, Ceftizoxim, Cefmenoxim, Cefmetazol, Cephaloglycin, Cefonicid, Cefodizim, Cefpirom, Ceftazidim, Ceftriaxon, Cefpiramid, Cefbuperazon, Cefozopran, Cefepim, Cefoselis, Cefluprenam, Cefuzonam, Cefpimizol, Cefclidin, Cefixim, Ceftibuten, Cefdinir, Cefpodoximaxetil, Cefpodoximproxetil, Cefteterampivoxil, Cefetametpivoxil, Cefcapenpivoxil oder Cefditorenpivoxil, Cefuroxim, Cefuroximaxetil, Loracarbacef, Ceftarolin und Latamoxef.
- 25 **11.** Pharmazeutische Zusammensetzung nach Anspruch 6, wobei das antibakterielle Mittel ausgewählt ist aus der Gruppe, bestehend aus Ceftazidim, Cefepim, Cefpirom, Piperacillin, Doripenem, Meropenem, Imipenem, Ceftarolin und Ceftolozan.
- 30 **12.** Pharmazeutische Zusammensetzung nach Anspruch 6, wobei das antibakterielle Mittel aus der Gruppe ausgewählt ist, bestehend aus den antibakteriellen Mitteln Aminoglykosiden, Ansamycinen, Carbacephemem, Cephalosporinen, Cephamycinen, Lincosamiden, Lipopeptiden, Macroliden, Monobactamen, Nitrofuranen, Penicillinen, Polypeptiden, Chinolonen, Sulfonamiden, Tetracyclinen oder Oxazolidinon antibakterielle Mittel.
- 35 **13.** Pharmazeutische Zusammensetzung nach Anspruch 5, umfassend (a) trans-Schwefelsäure-mono-[2-(N'-[(R)-piperidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon und (b) Sulbactam oder ein pharmazeutisch annehmbares Salz, Hydrat, Solvat, Polymorph oder Enantiomer davon.
- 40 **14.** Pharmazeutische Zusammensetzung nach Anspruch 5, umfassend (a) trans-Schwefelsäure-mono-[2-(N'-[(R)-pyrrolidin-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon und (b) Sulbactam oder ein pharmazeutisch annehmbares Salz, Hydrat, Solvat, Polymorph oder Enantiomer davon.
- 45 **15.** Zusammensetzung nach irgendeinem der Ansprüche 1 bis 3 oder eine pharmazeutische Zusammensetzung zur Verwendung nach irgendeinem der Ansprüche 4 bis 14 zur Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt.
- 50 **16.** Verbindung nach irgendeinem der Ansprüche 1 bis 3 oder eine pharmazeutische Zusammensetzung nach irgendeinem der Ansprüche 4 bis 14 zur Verwendung bei der Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt nach Anspruch 15, wobei die Infektion durch Bakterien verursacht wird, die ein oder mehrere beta-
- 55

Lactamase-Enzyme erzeugen.

17. Verbindung der Formel (I) nach Anspruch 1 oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon zur Verwendung bei der Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt, wobei diese Verwendung die Verabreichung einer pharmazeutisch wirksamen Menge, (a) einer Verbindung der Formel (I) nach Anspruch 1 oder eines Stereoisomers oder eines pharmazeutisch annehmbaren Salzes davon und (b) mindestens eines antibakteriellen Mittels oder eines pharmazeutisch annehmbaren Salzes, Hydrats, Solvats, Polymorphs oder Enantiomers davon an das Subjekt umfasst.
18. Verbindung der Formel (I) nach Anspruch 1 oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon zur Verwendung bei der Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt, wobei diese Verwendung die Verabreichung einer pharmazeutisch wirksamen Menge, entsprechend Anspruch 17, (a) einer Verbindung der Formel (I) nach Anspruch 1 oder eines Stereoisomers oder eines pharmazeutisch annehmbaren Salzes davon und (b) eines antibakteriellen Mittels, das mindestens einen beta-Lactamase-Inhibitor darstellt, ausgewählt aus Sulbactam, Tazobactam, Clavulansäure, oder eines pharmazeutisch annehmbaren Salzes, Hydrats, Solvats, Polymorphs oder Enantiomers davon an das Subjekt umfasst.
19. Verbindung der Formel (I) nach Anspruch 1 oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon zur Verwendung bei der Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt, wobei diese Verwendung die Verabreichung einer pharmazeutisch wirksamen Menge an das Substrat umfasst, entsprechend Anspruch 18, (a) einer Verbindung der Formel (I) nach Anspruch 1 oder eines Stereoisomers oder eines pharmazeutisch annehmbaren Salzes davon, (b) eines antibakteriellen Mittels, das mindestens einen beta-Lactamase-Inhibitor darstellt, ausgewählt aus Sulbactam, Tazobactam, Clavulansäure, oder eines pharmazeutisch annehmbaren Salzes, Hydrats, Solvats, Polymorphs oder Enantiomers davon, und (c) mindestens eines antibakteriellen Mittels oder eines pharmazeutisch annehmbaren Salzes, Hydrats, Solvats, Polymorphs oder Enantiomers davon zur Verwendung bei der Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt.
20. Verbindung der Formel (I) nach Anspruch 1 oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon zur Verwendung nach Anspruch 17, Anspruch 18 oder Anspruch 19 zur Prävention oder Behandlung einer bakteriellen Infektion in einem Subjekt, wobei die Infektion durch Bakterien verursacht ist, die ein oder mehrere beta-Lactamase-Enzyme erzeugen.
21. Verbindung der Formel (I) nach Anspruch 1 oder ein Stereoisomer oder ein pharmazeutisch annehmbares Salz davon zur Verwendung bei der Steigerung der antibakteriellen Wirksamkeit eines antibakteriellen Mittels in einem Substrat.

Revendications

1. Composé ayant la formule (I)



ou un stéréo-isomère ou un sel pharmaceutiquement acceptable de celui-ci ; dans lequel :

R₁ est :

- (a) un hydrogène,
- (b) (CO)_n-R₃,
- (c) COOR₄ ou
- (d) COCH₂COR₃ n est égal à 0, 1 ou 2 ;

R₂ est :

- (a) SO_3M ,
- (b) SO_2NH_2 ,
- (c) PO_3M ,
- (d) CH_2COOM ,
- (e) CF_2COOM ,
- (f) CHFCOOM , ou
- (g) CF_3 ;

M est un hydrogène ou un cation ;

R_3 est :

- (a) un hydrogène,
- (b) un alkyle en $\text{C}_1\text{-C}_6$ substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un halogène, OR_5 , CN , COOR_5 , CONR_6R_7 , NR_6R_7 , NR_5COR_6 , $\text{NR}_5\text{CONR}_6\text{R}_7$, un hétérocyclyle, un hétéroaryle, un cycloalkyle ou un aryle,
- (c) CN ,
- (d) NR_6R_7 ,
- (e) CONR_6R_7 ,
- (f) $\text{NHCONR}_6\text{R}_7$,
- (g) un aryle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_6\text{R}_7$,
- (h) un hétérocyclyle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$, $\text{NHC}(\text{NH})\text{NR}_6\text{R}_7$, ou $\text{NHCONR}_6\text{R}_7$,
- (i) un hétéroaryle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_6\text{R}_7$,
- (j) un cycloalkyle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_6\text{R}_7$,
- (k) un cycloalkyle substitué par un alkyle en $\text{C}_1\text{-C}_6$, dans lequel l'alkyle en $\text{C}_1\text{-C}_6$ est substitué de plus par un ou plusieurs substituants indépendamment choisis parmi OR_5 , NR_6R_7 , halogène, CN , ou CONR_6R_7 , ou
- (l) OR_6 ;

R_4 est :

- (a) un hydrogène,
- (b) un alkyle en $\text{C}_1\text{-C}_6$ substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un halogène, OR_5 , CN , COOR_5 , CONR_6R_7 , NR_6R_7 , NR_5COR_8 , un hétérocyclyle, un hétéroaryle, un cycloalkyle ou un aryle,
- (c) un aryle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_6\text{R}_7$,
- (d) un hétérocyclyle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_6\text{R}_7$,
- (e) un hétéroaryle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_6\text{R}_7$, ou
- (f) un cycloalkyle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_6R_7 , halogène, CN , CONR_6R_7 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$, ou $\text{NHCONR}_6\text{R}_7$;

R_5 et R_8 sont chacun indépendamment :

- (a) un hydrogène, ou
- (b) un alkyle en $\text{C}_1\text{-C}_6$ substitué de manière facultative par un ou plusieurs substituants indépendamment

choisis parmi un halogène, CN, CONR_6R_7 , NR_6R_7 , un hétérocyclyle, un hétéroaryle, un cycloalkyle ou un aryle ;

R_6 et R_7 sont chacun indépendamment:

- (a) un hydrogène,
- (b) un alkyle en $\text{C}_1\text{-C}_6$ substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un halogène, OR_5 , CN, COOR_5 , CONR_5R_8 , NR_5R_8 , NR_5COR_8 , un hétérocyclyle, un hétéroaryle, un cycloalkyle ou un aryle,
- (c) un aryle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_5R_8 , halogène, CN, CONR_5R_8 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_5\text{R}_8$,
- (d) un hétérocyclyle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_5R_8 , halogène, CN, CONR_5R_8 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_5\text{R}_8$,
- (e) un hétéroaryle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_5R_8 , halogène, CN, CONR_5R_8 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_5\text{R}_8$,
- (f) un cycloalkyle substitué de manière facultative par un ou plusieurs substituants indépendamment choisis parmi un alkyle en $\text{C}_1\text{-C}_6$, OR_5 , NR_5R_8 , halogène, CN, CONR_5R_8 , $\text{SO}_2\text{-alkyle}$, $\text{SO}_2\text{-aryle}$, $\text{OSO}_2\text{-alkyle}$, $\text{OSO}_2\text{-aryle}$ ou $\text{NHCONR}_5\text{R}_8$, ou
- (g) R_6 et R_7 sont rattachés ensemble pour former un cycle de quatre à sept éléments.

2. Composé selon la revendication 1, choisi parmi:

- acide trans-sulfurique mono[2-(N'-[(S)-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-[(R)-pipéridine-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-[(R)-pyrrolidine-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- trans-{7-oxo-2-[N'-((R)-pipéridine-3-carbonyl)-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-acide acétique ;
- trans-difluoro-{7-oxo-2-[N'-((R)-pipéridine-3-carbonyl)-hydrazinocarbonyl]-1,6-diaza-bicyclo[3.2.1]oct-6-yloxy}-acide acétique ;
- acide trans-sulfurique mono[2-hydrazinocarbonyl]-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-(amino-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-(3-amino-propioyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-(4-amino-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-((2S)-2-amino-3-hydroxy-propioyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-[(2S,4S)-4-fluoro-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-[(2S,4R)-4-méthoxy-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-(pipéridine-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-((RS)-pipéridine-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-((S)-pipéridine-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-((RS)-pipéridine-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;
- acide trans-sulfurique mono[2-(N'-((S)-pipéridine-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-((R)-pipéridine-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-(pipérazine-4-yl-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

5 acide trans-sulfurique mono[2-(N'-((RS)-1-amino-1-phényl-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-((RS)-3-amino-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

10 acide trans-sulfurique mono[2-(N'-(3-amino-2,2-diméthyl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-(1-aminométhyl-cyclopropane-1-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-(2-amino-4-carboxamido-butanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

15 acide trans-sulfurique mono[2-(N'-(5-amino-pentanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-((2S)-2,6-diamino-hexanoyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-((2-aminoéthoxy)-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

20 acide trans-sulfurique mono[2-(N'-(azétidine-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-(pyrrolidine-1-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

25 acide trans-sulfurique mono[2-(N'-[(R)-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-[(S)-3-pyrrolidine-2-yl]-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-[(RS)-pyrrolidine-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

30 acide trans-sulfurique mono[2-(N'-[(S)-pyrrolidine-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-[(2S,4R)-4-hydroxy-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

35 acide trans-sulfurique mono[2-(N'-[(2S,4S)-4-amino-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-[(2S,4S)-4-guanidino-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-[(RS)-3-pipéridine-2-yl-propionyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

40 acide trans-sulfurique mono[2-(N'-((RS)-pipérazine-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-((S)-morpholine-3-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

45 acide trans-sulfurique mono[2-(N'-(3-oxo-3-pipérazine-1-yl-propionyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

acide trans-sulfurique mono[2-(N'-((RS)-1-amino-1-pyridine-2-yl-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

50 acide trans-sulfurique mono[2-(N'-(2-amino-thiazol-4-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ;

ou un stéréo-isomère ou un sel pharmaceutiquement acceptable de ceux-ci.

3. Composé selon la revendication 1, choisi parmi :

sel de sodium de l'acide trans-sulfurique mono[2-(N'-(cyano-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diazabicyclo[3.2.1]oct-6-yl]ester ;

sel de sodium de trans-N'-(7-oxo-6-sulfoxy-1,6-diaza-bicyclo[3.2.1]octane-2-acide carboxylique) -acide hydra-

zinecarboxylique tert-butyl ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-(morpholine-4-yl-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-(6-carboxamido-pyridine-2-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-(morpholine-4-oxo-carbonyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-[(S)-1-carbamoyl-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-[(2S,4S)-1-carbamoyl-4-fluoro-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-[(S)-1-méthanesulfonyl-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-(cyano-diméthyl-acétyl)-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;
 sel de sodium de l'acide trans-sulfurique mono[2-(N'-[(S)-5-oxo-pyrrolidine-2-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]-oct-6-yl]ester ;

ou un stéréo-isomère de ceux-ci.

4. Composition pharmaceutique comprenant un composé selon l'une quelconque des revendications 1 à 3.
5. Composition pharmaceutique selon la revendication 4, comprenant en outre au moins un inhibiteur de bêta-lactamase choisi parmi le sulbactam, le tazobactam, l'acide clavulanique, ou un sel pharmaceutiquement acceptable, un hydrate, un solvate, un polymorphe ou un énantiomère de ceux-ci.
6. Composition pharmaceutique selon la revendication 4 ou 5, comprenant en outre au moins un agent antibactérien ou un sel pharmaceutiquement acceptable, un hydrate, un solvate, un polymorphe ou un énantiomère de celui-ci.
7. Composition pharmaceutique selon la revendication 6, dans laquelle l'agent antibactérien est choisi parmi un groupe constitué d'aminoglycosides, ansamycines, carbacéphèmes, céphalosporines, céphamycines, lincosamides, lipopeptides, macrolides, monobactams, nitrofuranes, pénicillines, polypeptides, quinolones, sulfonamides, tétracyclines ou agents antibactériens à base d'oxazolidinone.
8. Composition pharmaceutique selon la revendication 6, dans laquelle l'agent antibactérien est un agent antibactérien à base de bêta-lactame.
9. Composition pharmaceutique selon la revendication 6, dans laquelle ledit agent antibactérien est choisi parmi un groupe constitué de pénicillines, penems, carbapenems, céphalosporines et monobactams.
10. Composition pharmaceutique selon la revendication 6, dans laquelle l'agent antibactérien est un antibiotique à base de céphalosporine choisi parmi un groupe constitué de céphalothine, céphaloridine, céfaclor, céfadroxil, céfamandole, céfazoline, céphalexine, céphradine, céftizoxime, céfoxitine, céphacétrile, céfotiam, céfotaxime, céfsulodine, céfopérazone, céftizoxime, céfménoxime, céfmétazole, céphaloglycine, céfonicide, céfodizime, céfpirome, céftazidime, céftriaxone, céfpiramide, céfbupérazone, céfopran, céfépime, céfoselis, céfluprenam, céfuzonam, céfpimizole, céfclidine, céfixime, céftibuten, céfdinir, céfpodoxime axétil, céfpodoxime proxétil, céfteram pivoxil, pivoxil céfétafet pivoxil, céfcapène pivoxil ou céfditoren pivoxil, céfuroxime, céfuroxime axétil, loracarbacef, céftaroline et latamoxef.
11. Composition pharmaceutique selon la revendication 6, dans laquelle l'agent antibactérien est choisi parmi un groupe constitué de céftazidime, céfépime, céfpirome, pipéracilline doripenem, méropenem, imipenem, céftaroline et céftolozane.
12. Composition pharmaceutique selon la revendication 6, dans laquelle l'agent antibactérien est choisi parmi un groupe constitué d'aminoglycosides, ansamycines, carbacéphèmes, céphalosporines, céphamycines, lincosamides, lipopeptides, macrolides, monobactams, nitrofuranes, pénicillines, polypeptides, quinolones, sulfonamides, tétracyclines ou agents antibactériens à base d'oxazolidinone.

13. Composition pharmaceutique selon la revendication 5, comprenant (a) l'acide trans-sulfurique mono-[2-(N'-[(R)-pi-péridine-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ou un stéréo-isomère ou un sel pharmaceutiquement acceptable de celui-ci, et (b) du sulbactam ou un sel pharmaceutiquement acceptable, un hydrate, un solvate, un polymorphe ou un énantiomère de celui-ci.
14. Composition pharmaceutical selon la revendication 5, comprenant (a) l'acide trans-sulfurique mono-[2-(N'-[(R)-pyrrolidine-3-carbonyl]-hydrazinocarbonyl)-7-oxo-1,6-diaza-bicyclo[3.2.1]oct-6-yl]ester ou un stéréo-isomère ou un sel pharmaceutiquement acceptable de celui-ci, et (b) du sulbactam ou un sel pharmaceutiquement acceptable, un hydrate, un solvate, un polymorphe ou un énantiomère de celui-ci.
15. Composé selon l'une quelconque des revendications 1 à 3 ou composition pharmaceutique pour une utilisation selon l'une quelconque des revendications 4 à 14 pour prévenir ou traiter une infection bactérienne chez un sujet.
16. Composé selon l'une quelconque des revendications 1 à 3 ou composition pharmaceutique selon l'une quelconque des revendications 4 à 14 pour une utilisation pour prévenir ou traiter une infection bactérienne chez un sujet selon la revendication 15, ladite infection étant provoquée par des bactéries produisant une ou plusieurs enzymes bêta-lactamases.
17. Composé ayant la formule (I) selon la revendication 1 ou stéréo-isomère ou sel pharmaceutiquement acceptable de celui-ci pour une utilisation pour prévenir ou traiter une infection bactérienne chez un sujet, ladite utilisation comprenant l'administration audit sujet d'une quantité pharmaceutiquement efficace : (a) d'un composé ayant la formule (I) selon la revendication 1 ou d'un stéréo-isomère ou d'un sel pharmaceutiquement acceptable de celui-ci, et (b) et moins d'un agent antibactérien ou d'un sel pharmaceutiquement acceptable, d'un hydrate, d'un solvate, d'un polymorphe ou d'un énantiomère de celui-ci.
18. Composé ayant la formule (I) selon la revendication 1 ou stéréoisomère ou sel pharmaceutiquement acceptable de celui-ci pour une utilisation pour prévenir ou traiter une infection bactérienne chez un sujet, ladite utilisation comprenant l'administration audit sujet d'une quantité pharmaceutiquement efficace selon la revendication 17 : (a) d'un composé ayant la formule (I) selon la revendication 1 ou d'un stéréo-isomère ou d'un sel pharmaceutiquement acceptable de celui-ci, et (b) d'un agent antibactérien, qui est au moins un inhibiteur de bêta-lactamase choisi parmi le sulbactam, le tazobactam, l'acide clavulanique, ou un sel pharmaceutiquement acceptable, un hydrate, un solvate, un polymorphe ou un énantiomère de ceux-ci.
19. Composé ayant la formule (I) selon la revendication 1 ou stéréo-isomère ou sel pharmaceutiquement acceptable de celui-ci pour une utilisation pour prévenir ou traiter une infection bactérienne chez un sujet, ladite utilisation comprenant l'administration audit sujet d'une quantité pharmaceutiquement efficace selon la revendication 18 : (a) d'un composé ayant la formule (I) selon la revendication 1 ou d'un stéréo-isomère ou d'un sel pharmaceutiquement acceptable de celui-ci, (b) d'un agent antibactérien, qui est au moins un inhibiteur de bêta-lactamase choisi parmi le sulbactam, le tazobactam, l'acide clavulanique, ou un sel pharmaceutiquement acceptable, un hydrate, un solvate, un polymorphe ou un énantiomère de ceux-ci, et (c) au moins d'un agent antibactérien ou d'un sel pharmaceutiquement acceptable, d'un hydrate, d'un solvate, d'un polymorphe ou d'un énantiomère de ceux-ci pour une utilisation pour prévenir ou traiter une infection bactérienne chez un sujet.
20. Composé ayant la formule (I) selon la revendication 1 ou stéréo-isomère ou sel pharmaceutiquement acceptable de celui-ci pour une utilisation selon la revendication 17, la revendication 18 ou la revendication 19 pour prévenir ou traiter une infection bactérienne chez un sujet, ladite infection étant provoquée par des bactéries produisant une ou plusieurs enzymes bêta-lactamases.
21. Composé ayant la formule (I) selon la revendication 1 ou stéréo-isomère ou sel pharmaceutiquement acceptable de celui-ci pour une utilisation afin d'augmenter l'efficacité antibactérienne d'un agent antibactérien chez un sujet.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- US 2010060923 W [0003]
- EP 2010067647 W [0003]
- US 2010052109 W [0003]
- US 2010048109 W [0003]
- GB 2009050609 W [0003]
- EP 2009056178 W [0003]
- US 2009041200 W [0003]
- WO 2009091856 A [0003] [0059]
- US 20100093784 A [0003]

Non-patent literature cited in the description

- **COATES et al.** *Br. J. Pharmacol.*, 2007, vol. 152 (8), 1147-1154 [0002]
- **GWYNN et al.** *Annals of the New York Academy of Sciences*, 2010, vol. 1213, 5-19 [0002]
- **S. M. BERGE et al.** *J. Pharmaceutical Sciences*, 1977, vol. 66, 1-19 [0038]
- Goodman and Gilman's: The Pharmacological Basis of Therapeutics. Pergamon Press, 1990 [0052]

Szabadalmi igénypontok

1. (I) képlettű vegyület



a képletben

R₁ jelentése

- (a) hidrogén,
- (b) (CO)_n-R₃,
- (c) COOR₄ vagy
- (d) COCH₂COR₃,



n értéke 0, 1 vagy 2;

R₂ jelentése

- (a) SO₃M,
- (b) SO₂NH₂,
- (c) PO₃M,
- (d) CH₂COOM,
- (e) CF₂COOM,
- (f) CHF₂COOM vagy
- (g) CF₃,

M jelentése hidrogén vagy egy kation;

R₃ jelentése

- (a) hidrogén,
- (b) C₁-C₆ alkil, amely adott esetben helyettesítve van halogén, OR₅, CN, COOR₅, CONR₆R₇, NR₆R₇, NR₅COR₈, NR₅CONR₆R₇, heterociklusos, heteroaril, cikloalkil vagy aril közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,
- (c) CN,
- (d) NR₆R₇,
- (e) CONR₆R₇,
- (f) NHCONR₆R₇,
- (g) aril, amely adott esetben helyettesítve van C₁-C₆ alkil, OR₅, NR₆R₇, halogén, CN, CONR₆R₇, SO₂-alkil, SO₂-aril, OSO₂-alkil, OSO₂-aril vagy NHCONR₆R₇ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(h) heterociklusos csoport, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril, $NHC(NH)NR_6R_7$ vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(i) heteroaril, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(j) cikloalkil, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(k) C_1 - C_8 alkil csoporttal helyettesített cikloalkil, ahol C_1 - C_6 alkil tovább van helyettesítve OR_5 , NR_6R_7 , halogén, CN vagy $CONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel

vagy

(l) OR_6 ;

R_4 jelentése

(a) hidrogén,

(b) C_1 - C_6 alkil, amely adott esetben helyettesítve van halogén, OR_5 , CN, $COOR_5$, $CONR_6R_7$, NR_6R_7 , NR_5COR_6 , heterociklusos, heteroaril, cikloalkil vagy aril közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(c) aril, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(d) heterociklusos csoport, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(e) heteroaril, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel, vagy

(f) cikloalkil, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_6R_7 , halogén, CN, $CONR_6R_7$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_6R_7$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

R_5 és R_8 jelentése egymástól függetlenül

(a) hidrogén, vagy

(b) C_1 - C_6 alkil, amely adott esetben helyettesítve van halogén, CN, $CONR_6R_7$, NR_6R_7 , heterociklusos, heteroaril, cikloalkil vagy aril közül egymástól függetlenül megválasztott egy vagy több helyettesítővel;

R_6 és R_7 jelentése egymástól függetlenül

(a) hidrogén,

(b) C_1 - C_6 alkil, amely adott esetben helyettesítve van halogén, OR_5 , CN, $COOR_5$, $CONR_5R_8$, NR_5R_8 , NR_5COR_8 , heterociklusos, heteroaril, cikloalkil vagy aril közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(c) aril, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_5R_8 , halogén, CN, $CONR_5R_8$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_5R_8$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(d) heterociklusos csoport, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_5R_8 , halogén, CN, $CONR_5R_8$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_5R_8$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(e) heteroaril, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_5R_8 , halogén, CN, $CONR_5R_8$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_5R_8$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel,

(f) cikloalkil, amely adott esetben helyettesítve van C_1 - C_6 alkil, OR_5 , NR_5R_8 , halogén, CN, $CONR_5R_8$, SO_2 -alkil, SO_2 -aril, OSO_2 -alkil, OSO_2 -aril vagy $NHCONR_5R_8$ közül egymástól függetlenül megválasztott egy vagy több helyettesítővel, vagy

(g) R_6 és R_7 együtt összekapcsolódva négy – hét tagú gyűrűt képeznek -

vagy egy sztereoizomerje vagy ezek egy gyógyszerileg elfogadható sója.

2. Az 1. igénypont szerinti vegyület az alábbiak közül kiválasztva:

transz-kénsav-mono-[2-(N'-{(S)-pirrolidin-2-karbonil}-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-{(R)-piperidin-3-karbonil}-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-{(R)-pirrolidin-3-karbonil}-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-[7-oxo-2-[N'-{(R)-piperidin-3-karbonil}-hidrazinokarbonil]-1,8-diaza-biciklo[3.2.1]-okt-6-iloxi]-ecetsav;

transz-difluor-[7-oxo-2-[N'-{(R)-piperidin-3-karbonil}-hidrazinokarbonil]-1,6-diaza-biciklo[3.2.1]-okt-6-iloxi]-ecetsav;

transz-kénsav-mono-[2-hidrazinokarbonil-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-(amino-acetil)hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-(3-amino-propanoil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-(4-amino-butanoil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-(2S)-2-amino-3-hidroxi-propionil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-{(2S,4S)-4-fluor-pirrolidin-2-karbonil}-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-{(2S,4R)-4-metoxi-pirrolidin-2-karbonil}-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-(piperidin-4-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]-észter;

transz-kénsav-mono-[2-(N'-((RS)-piperidin-3-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((S)-piperidin-3-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((RS)-piperidin-2-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((S)-piperidin-2-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((R)-piperidin-2-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-(piperazin-4-il-acetil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((RS)-1-amino-1-fenil-acetil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((RS)-3-amino-butanoil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-(3-amino-2,2-dimetil-propionil)-hidrazinokarbonil)-7-oxo-1,6-diazabicyclo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-(1-aminometil-ciklopropán-1-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-(2-amino-4-karboxamido-butanoil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-(5-amino-pentanoil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((2S)-2,6-diamino-hexanoil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((2-aminoetoxi)-acetil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-(azetidin-3-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]-észter;

transz-kénsav-mono-[2-(N'-(pirrolidin-1-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]-észter;

transz-kénsav-mono-[2-(N'-((R)-pirrolidin-2-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((S)-3-pirrolidin-2-il)-propionil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-((RS)-pirrolidin-3-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-ii]észter;

transz-kénsav-mono-[2-(N'-[(S)-pirrolidin-3-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter,

transz-kénsav-mono-[2-(N'-[(2S,4R)-4-hidroxi-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(2S,4S)-4-amino-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(2S,4S)-4-guanidino-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(RS)-3-piperidin-2-il-propionil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(RS)-piperazin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(S)-morfolin-3-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[3-oxo-3-piperazin-1-il-propionil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(RS)-1-amino-1-piridin-2-il-acetil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[3-amino-tiazol-4-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

vagy ezek egy sztereoisomerje vagy ezek egy gyógyászati lág elfogadható sója.

3. Az 1. igénypont szerinti vegyület az alábbiak közül kiválasztva:

transz-kénsav-mono-[2-(N'-(ciano-acetil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

transz-N'-(7-oxo-6-szulfonoxi-1,6-diaza-biciklo[3.2.1]oktán-2-karbonsav)-hidrazinkarbonsav-terc-butil észter-nátriumsó;

transz-kénsav-mono-[2-(N'-(morfolin-4-il-acetil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

transz-kénsav-mono-[2-(N'-(6-karboxamido-piridin-2-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

ransz-kénsav-mono-[2-(N'-(morfolin-4-oxo-karbonil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

transz-kénsav-mono-[2-(N'-[(S)-1-karbamoil-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

transz-kénsav-mono-[2-(N'-[(2S,4S)-1-karbamoil-4-fluor-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

transz-kénsav-mono-[2-(N'-[(S)-1-metánszulfonil-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;

transz-kénsav-mono-[2-(N'-(ciano-dimetil-acetil)-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter;

transz-kénsav-mono-[2-(N'-[(S)-5-oxo-pirrolidin-2-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észter-nátriumsó;
vagy ezek egy sztereoizomerje.

4. Gyógyászati készítmény, amely magában foglal egy, az 1 – 3. igénypontok bármelyike szerinti vegyületet.

5. A 4. igénypont szerinti gyógyászati készítmény, amely magában foglal továbbá szulbaktám, tazobaktám, klavulánsav vagy ezek egy gyógyászatilag elfogadható sója, hidrátja, szolvátja, polimorfja vagy enantiomerje közül megválasztott legalább egy béta-laktámáz inhibitor.

6. A 4. vagy 5. igénypont szerinti gyógyászati készítmény, amely magában foglal továbbá legalább egy antibakteriális ágens vagy ennek egy gyógyászatilag elfogadható sóját, hidrátját, szolvátját, polimorfját vagy enantiomerjét.

7. A 6. igénypont szerinti gyógyászati készítmény, ahol az antibakteriális ágens aminoglükozidok, anzamicinek, karbacefemek, cefalosporinok, cefamicinek, linkozamidok, lipopeptidek, makrolidok, monobaktámok, nitrofuránok, penicillinek, polipeptidek, kinolonok, szulfonamidok, tetraciklinek vagy oxazolidinon antibakteriális ágensek alkotta csoportból van megválasztva.

8. A 6. igénypont szerinti gyógyászati készítmény, ahol az antibakteriális ágens egy béta-laktám antibakteriális ágens.

9. A 6. igénypont szerinti gyógyászati készítmény, ahol az antibakteriális ágens a penicillinek, penemek, karbapenemek, cefalosporinok és monobaktámok alkotta csoportból van megválasztva.

10. A 6. igénypont szerinti gyógyászati készítmény, ahol az antibakteriális ágens cefalotin, cefaloridin, cefaclór, cefadroxil, cefamandol, cefazolin, cefalexin, cefradin, ceftizoxim, cefoxitin, cefacetil, cefotiam, cefotaxim, cefazulodin, cefoperazon, ceftizoxim, cefmenoxim, cefmetazol, cefaloglicin, cefonicid, cefodizim, cefpirom, ceftazidim, ceftriaxon, cefpiramid, cefbuperazon, cefozopran, cefepim, cefoszelis, cefuprenam, cefuzonam, cefpimizol, cefelidin, cefixim, cefibutén, cefdinir, cefpodoxim axetil, cefpodoxim proxetil, cefteram pivoxil, cefetamet pivoxil, cefcapen pivoxil vagy cefditoren pivoxil, cefuroxim, cefuroxim axetil, lorakarbacef, ceftarolin és latamoxef közül megválasztott cefalosporin antibiotikum.

11. A 6. igénypont szerinti gyógyászati készítmény, ahol az antibakteriális ágens ceftazidim, cefepim, cefpirom, piperacillin doripeném, meropeném, imipeném, kaftarolin és ceftolozan közül van megválasztva.

12. A 6. igénypont szerinti gyógyászati készítmény, ahol az antibakteriális ágens aminoglükozidok, anzamicinek, karbacefemek, cefalosporinok, cefamicinek, linkozamidok, lipopeptidek, makrolidok, monobaktámok, nitrofuránok, penicillinek, polipeptidek, kinolonok, szulfonamidok, tetraciklinek vagy oxazolidinon antibakteriális ágensek alkotta csoportból van megválasztva.

13. Az 5. igénypont szerinti gyógyászati készítmény, amely magában foglalja (a) transz-kénsav-mono-[2-(N'-[(R)-piperidin-3-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észtert vagy egy sztereoizomerjét vagy ezek egy gyógyászatilag elfogadható sóját és (b) szulbaktámot vagy egy gyógyászatilag elfogadható sóját, hidrátját, szolvátját, polimorfját vagy enantiomerjét.

14. Az 5. igénypont szerinti gyógyászati készítmény, amely magában foglalja (a) transz-kénsav-mono-[2-(N'-[(R)-pirrolidin-3-karbonil]-hidrazinokarbonil)-7-oxo-1,6-diaza-biciklo[3.2.1]okt-6-il]észtert vagy egy sztereoizomerjét vagy ezek egy gyógyászatilag elfogadható sóját és (b) szulbaktámot vagy egy gyógyászatilag elfogadható sóját, hidrátját, szolvátját, polimorfját vagy enantiomerjét.

15. Az 1 – 3. igénypontok bármelyike szerinti vegyület vagy a 4 – 14. igénypontok bármelyike szerinti gyógyászati készítmény egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére való alkalmazásra.

16. Az 1 – 3. igénypontok bármelyike szerinti vegyület vagy a 4 – 14. igénypontok bármelyike szerinti gyógyászati készítmény egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére való alkalmazásra a 15. igénypont szerint, ahol a fertőzést egy vagy több béta-laktamáz enzimet termelő baktériumok okozzák.

17. Az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére való alkalmazásra, ahol az alkalmazás során az említett egyednek beadásra kerül gyógyászatilag hatásos mennyiségben (a) egy, az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója és (b) legalább egy antibakteriális ágens vagy egy gyógyászatilag elfogadható sója, hidrátja, szolvátja, polimorfja vagy enantiomerje.

18. Az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére való alkalmazásra, ahol az alkalmazás során az említett egyednek beadásra kerül gyógyászatilag hatásos mennyiségben a 17. igénypont szerint (a) egy, az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója és (b) egy antibakteriális ágens, amely szulbaktám, tazobaktám, klavulánsav vagy ezek egy gyógyászatilag elfogadható sója, hidrátja, szolvátja, polimorfja vagy enantiomerje közül megválasztott legalább egy béta-laktamáz inhibitor.

19. Az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére való alkalmazásra, ahol az alkalmazás során az említett egyednek beadásra kerül gyógyászatilag hatásos mennyiségben a 18. igénypont szerint (a) egy, az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója, (b) egy antibakteriális ágens, amely szulbaktám, tazobaktám, klavulánsav vagy ezek egy gyógyászatilag elfogadható sója, hidrátja, szolvátja, polimorfja vagy enantiomerje közül megválasztott legalább egy béta-laktamáz inhibitor és (c) legalább egy antibakteriális ágens vagy ennek egy gyógyászatilag elfogadható sója, hidrátja, szolvátja, polimorfja vagy enantiomerje egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére való alkalmazásra.

20. Az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója a 17., 18. vagy 19. igénypont szerinti alkalmazásra egy egyedben bakteriális fertőzés megelőzésére vagy kezelésére, ahol a fertőzést egy vagy több béta-laktamáz enzimet termelő baktériumok okozzák.

21. Az 1. igénypont szerinti (I) képletű vegyület vagy egy sztereoizomerje vagy ezek egy gyógyászatilag elfogadható sója egy egyedben az antibakteriális ágens antibakteriális hatékonyságának növelésére való alkalmazásra.