

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12)

Oversættelse af europæisk patentskrift

(10) **DK/EP 3440102 T3**

-
- (51) Int.Cl.: **C 07 K 14/47 (2006.01)** **A 61 K 38/00 (2006.01)** **A 61 P 35/00 (2006.01)**
C 07 K 14/725 (2006.01) **C 12 N 15/62 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-07-22**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2024-06-05**
- (86) Europæisk ansøgning nr.: **17723015.8**
- (86) Europæisk indleveringsdag: **2017-04-10**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-02-13**
- (86) International ansøgning nr.: **EP2017058576**
- (87) Internationalt publikationsnr.: **WO2017174822**
- (30) Prioritet: **2016-04-08 GB 201606172**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Adaptimmune Limited, 60 Jubilee Avenue , Milton Park, Abingdon, Oxfordshire OX14 4RX, Storbritannien**
- (72) Opfinder: **TRIBBLE, Nicholas, c/o Adaptimmune Limited, 101 Park Drive, Milton Park, Abingdon, Oxfordshire OX14 4RY, Storbritannien**
LAWRANCE, William, c/o Adaptimmune Limited, 101 Park Drive, Milton Park, Abingdon, Oxfordshire OX14 4RY, Storbritannien
BAGG, Eleanor, c/o Adaptimmune Limited, 101 Park Drive, Milton Park, Abingdon, Oxfordshire OX14 4RY, Storbritannien
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **T-cellereceptorer**
- (56) Fremdragne publikationer:
WO-A2-02/094981
US-A1- 2004 077 045
HILLIG R C ET AL: "High-resolution structure of HLA-A*0201 in complex with a tumour-specific antigenic peptide encoded by the MAGE-A4 gene", JOURNAL OF MOLECULAR BIO, ACADEMIC PRESS, UNITED KINGDOM, vol. 310, no. 5, 27 July 2001 (2001-07-27) , pages 1167-1176, XP004466111, ISSN: 0022-2836, DOI: 10.1006/JMBI.2001.4816
Zheng-Cai Jia ET AL: "Identification of Two Novel HLA-A*0201-Restricted CTL Epitopes Derived from MAGE-A4", CLINICAL & DEVELOPMENTAL IMMUNOLOGY, vol. 2010, 1 January 2010 (2010-01-01), pages 1-7, XP055364879, US ISSN: 1740-2522, DOI: 10.1155/2010/567594
WAHL A ET AL: "HLA class I molecules reflect an altered host proteome after influenza virus infection", HUMAN IMMUNOLOGY, NEW YORK, NY, US, vol. 71, no. 1, 1 January 2010 (2010-01-01), pages 14-22, XP026794729, ISSN: 0198-8859 [retrieved on 2009-09-11]

Fortsættes ...

DESCRIPTION

Description

[0001] The present invention relates to T cell receptors (TCRs) which bind the HLA-A*0201 restricted decapeptide GVDGEEHSV derived from the melanoma-associated antigen (MAGE) B2 protein (amino acids 230-239). The TCRs of the invention demonstrate excellent specificity profiles for this MAGE epitope.

Background to the Invention

[0002] Cancer testis antigens (CTA) are a subclass of tumour-associated antigen (TAA) encoded by approximately 140 genes. Expression of these antigens is restricted in immune privileged sites such as the testes, placenta and fetal ovary; they are typically not expressed in other tissues. Expression of these genes has been observed in malignant tumors. The immunogenicity of CTA has led to the widespread development of cancer vaccines targeting these antigens in many solid tumors. Within this large class of TAA, melanoma-associated antigens (MAGE) have emerged as promising candidates for cancer immunotherapy.

[0003] More than 30 cancer testis (CT) genes have been reported as members of multi-gene families that are organized into gene clusters on chromosome X (CT-X antigens). The CT gene clusters are located between Xq24 and Xq28 and include gene families such as MAGE and NY-ESO-1. Type I MAGE gene clusters are the most extensively characterized and include the MAGE-A, MAGE-B and MAGE-C families. The MAGE-A proteins are encoded by 12 different MAGE-A gene family members (MAGE-A1 to MAGE-A12) and are defined by a conserved 165-171 amino acid base, called the MAGE homology domain (MHD). The MHD corresponds to the only region of shared amino acids by all of the MAGE-A family members.

[0004] T cells recognize and interact with complexes of cell surface molecules, referred to as human leukocyte antigens ("HLA"), or major histocompatibility complexes ("MHCs"), and peptides. The peptides are derived from larger molecules, which are processed by the cells which also present the HLA/MHC molecule. The interaction of T cells and HLA/peptide complexes is restricted, requiring a T cell specific for a particular combination of an HLA molecule and a peptide. If a specific T cell is not present, there is no T cell response even if its partner complex is present. Similarly, there is no response if the specific complex is absent, but the T cell is present. The mechanism is involved in the immune system's response to infection, in autoimmune disease, and in responses to abnormalities such as tumours.

[0005] Some MAGE gene family proteins are only expressed in germ cells and cancer (MAGE-A to MAGE-C families). Others are widely expressed in normal tissues (MAGE-D through to

MAGE-H). All these MAGE protein families have a homologous region that is closely matched to the sequence of the other MAGE proteins and comprises peptides displayed as HLA/peptide complexes in immune recognition. Hence, it is important to select TCR clinical candidates that are highly specific for the desired MAGE peptide/HLA-A2 antigen.

[0006] MAGE B2 is a CTA member of the MAGE B gene family. The function is unknown, though it is thought that it may play a role in embryonal development. In tumour pathogenesis, it appears to be involved in tumor transformation or aspects of tumor progression. MAGE B2 has been implicated in a large number of tumours. The peptide GVDGEEHSV (SEQ ID No: 1) corresponds to amino acid residue numbers 231-241 of the known MAGE-B2 protein.

[0007] MAGE A4 is a CTA of the MAGE A gene family. MAGE A4 is expressed in testis and placenta, and in a significant fraction of tumors of various histological types. The peptide GVDGREHTV (SEQ ID No. 2) shows cross-reactivity with MAGE B2, such that certain TCRs are able to bind to HLA molecules displaying both peptides.

[0008] US patent publication 2007/077045 describes MAGE peptides including MAGE A4 and MAGE B2, and TCRs which are specific for such antigens.

[0009] International patent application WO02/094981 describes the MAGE peptides GVDGEEHSV and GVDGREHTV.

[0010] Hillig et al., 2001 J. Mol. Biol. 310:1167-1176 describes a heterotrimeric complex of human HLA-A*0201, GVDGREHTV and beta-2 microglobulin.

[0011] Zheng-Cai et al., 2010 Clin. Dev. Immunol. 2010:107 identifies as a CTL epitope of Mage-A4. Wahl et al., 2010 Human Immunology, New York, 71:14-22 identified GVDGEEHSV bound to HLA-A*0201.

Summary of the Invention

[0012] We have developed a TCR which binds to HLA molecules displaying the MAGE B2 peptide GVDGEEHSV in preference to MAGE A4. In a first aspect, the present invention provides a T cell receptor (TCR) having the property of binding to GVDGEEHSV (SEQ ID No: 1) in complex with HLA-A*0201 with a dissociation constant of from about 0.05 μ M to about 20.0 μ M when measured with surface plasmon resonance at 25°C and at a pH between 7.1 and 7.5 using a soluble form of the TCR, and has at least a ten-fold selectivity of binding to SEQ ID No:1 in complex with HLA-A*0201 over binding to GVDGREHTV (SEQ ID No 2) in complex with HLA-A*0201, wherein the TCR comprises (A) the TCR alpha chain variable domain of a sequence in which: (i) amino acid residues 1-27 has (a) at least 90% identity to the sequence of amino acid residues 1-26 of SEQ ID No: 3 or (b) one, two or three amino acid insertions or deletions relative to the sequence of (a); (ii) amino acid residues 28-33 is VSPFSN; (iii) amino acid residues 34-47 has (a) at least 90% identity to the sequence of amino

acid residues 34-47 of SEQ ID NO: 3 or (b) one, two or three amino acid insertions or deletions relative to the sequence of (a); (iv) amino acid residues 48-53 is LTIMTF or LTRMTF; (v) amino acid residues 54-90 has (a) at least 90% identity to the sequence of amino acid residues 54-90 of SEQ ID No: 3 or (b) one, two or three amino acid insertions, deletions or substitutions relative to the sequence of (a); and (vi) amino acid residues 91-105 is CWSGGTDSWGKLQF; and (B) the TCR beta chain variable domain of a sequence in which: (i) amino acid residues 1-45 has (a) at least 90% identity to the amino acid sequence of residues 1-45 of SEQ ID No: 4 or (b) one, two or three amino acid insertions or deletions relative to the sequence of (a); (ii) amino acid residues 46-50 is KGHDR or KGRDR; (iii) amino acid residues 51-67 has (a) at least 90% identity to the sequence of amino acid residues 51-67 of SEQ ID NO: 4 or (b) one, two or three amino acid insertions or deletions relative to the sequence of (a); (iv) amino acid residues 68-72 is SFDVK; (v) amino acid residues 73-109 has (a) at least 90% identity to the sequence of amino acid residues 73-109 of SEQ ID NO: 4 or (b) one, two or three amino acid insertions or deletions relative to the sequence of (a); and (vi) amino acid residues 110-123 is CATSGQGAYREQFF or CATSGQGAYKEQFF, or CATSGQGAYNEQFF; and wherein the TCR variable domains form contacts with at least residues V2, Y3 and D4 of GVDGEEHSV (SEQ ID No: 1).

[0013] The invention also provides nucleic acid encoding a TCR of the invention. The invention further provides an isolated or non-naturally occurring cell, especially a T-cell, presenting a TCR of the invention. The invention also provides a cell harbouring (a) a TCR expression vector which comprises nucleic acid of the invention in a single open reading frame, or two distinct open reading frames encoding the alpha chain and the beta chain respectively; or (b) a first expression vector which comprises nucleic acid encoding the alpha chain of a TCR of the invention, and a second expression vector which comprises nucleic acid encoding the beta chain of a TCR of the invention. The invention also provides a pharmaceutical composition comprising a TCR of the invention, nucleic acid of the invention or a cell of the invention, together with one or more pharmaceutically acceptable carriers or excipients. The invention also provides a TCR of the invention for use in a method of treating cancer.

[0014] The following aspects and embodiments are limited to the invention as defined in the claims.

[0015] In embodiments, the TCR according to the invention has the property of binding to GVDGEEHSV (SEQ ID No: 1) in complex with HLA-A*0201 with a dissociation constant of from about 20 μM to about 50 μM when measured with surface plasmon resonance at 25°C and at a pH between 7.1 and 7.5 using a soluble form of the TCR, wherein the TCR comprises a TCR alpha chain variable domain and a TCR beta chain variable domain. In some embodiments, the dissociation constant is above 50 μM , such as 100 μM , 200 μM , 500 μM or more.

[0016] Accordingly, a TCR in accordance with the invention is capable of binding efficiently to HLA displaying GVDGEEHSV but not to HLA displaying GVDGREHTV.

[0017] The GVDGEEHSV HLA-A2 complex provides a cancer marker that the TCRs of the invention can target. The present invention provides such TCRs useful for the purpose of delivering cytotoxic or immune effector agents to the cancer cells and/or useful for use in adoptive therapy.

[0018] TCRs are described using the International Immunogenetics (IMGT) TCR nomenclature, and links to the IMGT public database of TCR sequences. Native alpha-beta heterodimeric TCRs have an alpha chain and a beta chain. Broadly, each chain comprises variable, joining and constant regions, and the beta chain also usually contains a short diversity region between the variable and joining regions, but this diversity region is often considered as part of the joining region. Each variable region comprises three CDRs (Complementarity Determining Regions) embedded in a framework sequence, one being the hypervariable region named CDR3. There are several types of alpha chain variable ($V\alpha$) regions and several types of beta chain variable ($V\beta$) regions distinguished by their framework, CDR1 and CDR2 sequences, and by a partly defined CDR3 sequence. The $V\alpha$ types are referred to in IMGT nomenclature by a unique TRAV number. Thus "TRAV21" defines a TCR $V\alpha$ region having unique framework and CDR1 and CDR2 sequences, and a CDR3 sequence which is partly defined by an amino acid sequence which is preserved from TCR to TCR but which also includes an amino acid sequence which varies from TCR to TCR. In the same way, "TRBV5-1" defines a TCR $V\beta$ region having unique framework and CDR1 and CDR2 sequences, but with only a partly defined CDR3 sequence.

[0019] The joining regions of the TCR are similarly defined by the unique IMGT TRAJ and TRBJ nomenclature, and the constant regions by the IMGT TRAC and TRBC nomenclature.

[0020] The beta chain diversity region is referred to in IMGT nomenclature by the abbreviation TRBD, and, as mentioned, the concatenated TRBD/TRBJ regions are often considered together as the joining region.

[0021] The α and β chains of $\alpha\beta$ TCRs are generally regarded as each having two "domains", namely variable and constant domains. The variable domain consists of a concatenation of variable region and joining region. In the present specification and claims, the term "TCR alpha variable domain" therefore refers to the concatenation of TRAV and TRAJ regions, and the term TCR alpha constant domain refers to the extracellular TRAC region, or to a C-terminal truncated TRAC sequence. Likewise the term "TCR beta variable domain" refers to the concatenation of TRBV and TRBD/TRBJ regions, and the term TCR beta constant domain refers to the extracellular TRBC region, or to a C-terminal truncated TRBC sequence.

[0022] The unique sequences defined by the IMGT nomenclature are widely known and accessible to those working in the TCR field. For example, they can be found in the IMGT public database. The "T cell Receptor Factsbook", (2001) LeFranc and LeFranc, Academic Press, ISBN 0-12-441352-8 also discloses sequences defined by the IMGT nomenclature, but because of its publication date and consequent time-lag, the information therein sometimes needs to be confirmed by reference to the IMGT database.

[0023] One TCR in accordance with the invention comprises an alpha chain extracellular domain as shown in SEQ ID No: 3 (TRAV10 + TRAC) and a beta chain extracellular domain as shown in SEQ ID No: 4 (TRBV24-1 + TRBC-2). The terms "parental TCR", "parental MAGE-A4 TCR", are used synonymously herein to refer to this TCR comprising the extracellular alpha and beta chain of SEQ ID Nos: 3 and 4 respectively. It is desirable to provide TCRs that are mutated or modified relative to the parental TCR that have a higher affinity and/or a slower off-rate for the peptide-HLA complex than the parental TCR.

[0024] For the purpose of providing a reference TCR against which the binding profile of such mutated or modified TCRs may be compared, it is convenient to use a soluble TCR in accordance with the invention having the extracellular sequence of the parental MAGE-A4 TCR alpha chain given in SEQ ID No. 3 and the extracellular sequence of the parental MAGE-A4 TCR beta chain given in SEQ ID No: 4. That TCR is referred to herein as the "the reference TCR" or "the reference MAGE-A4 TCR". Note that SEQ ID No: 5 comprises the parental alpha chain extracellular sequence of SEQ ID No: 3 and that C162 has been substituted for T162 (i.e. T48 of TRAC). Likewise SEQ ID No: 6 is the parental beta chain extracellular sequence of SEQ ID No: 4 and that C169 has been substituted for S169 (i.e. S57 of TRBC2), A187 has been substituted for C187 and D201 has been substituted for N201. These cysteine substitutions relative to the parental alpha and beta chain extracellular sequences enable the formation of an interchain disulfide bond which stabilises the refolded soluble TCR, i.e. the TCR formed by refolding extracellular alpha and beta chains. Use of the stable disulfide linked soluble TCR as the reference TCR enables more convenient assessment of binding affinity and binding half life. TCRs of the invention may comprise the mutations described above.

[0025] TCRs of the invention may be non-naturally occurring and/or purified and/or engineered. TCRs of the invention may have more than one mutation present in the alpha chain variable domain and/or the beta chain variable domain relative to the parental TCR. "Engineered TCR" and "mutant TCR" are used synonymously herein and generally mean a TCR which has one or more mutations introduced relative to the parental TCR, in particular in the alpha chain variable domain and/or the beta chain variable domain thereof. These mutation(s) may improve the binding affinity for GVDGEEHSV (SEQ ID No: 1) in complex with HLA-A*020101. In certain embodiments, there are 1, 2, 3, 4, 5, 6, 7 or 8 mutations in alpha chain variable domain, for example 4 or 8 mutations, and/or 1, 2, 3, 4 or 5 mutations in the beta chain variable domain, for example 5 mutations. In some embodiments, the α chain variable domain of the TCR of the invention may comprise an amino acid sequence that has at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98 % or at least 99% identity to the sequence of amino acid residues 1- 105 of SEQ ID No: 3. In some embodiments, the β chain variable domain of the TCR of the invention may comprise an amino acid sequence that has at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98 % or at least 99% identity to the sequence of amino acid residues 1-123 of SEQ ID No: 4.

[0026] One aspect disclosed herein for illustrative purposes provides a TCR in which the alpha

chain variable domain may have the following mutation:

CDR2 I3	R
---------	---

with reference to the numbering shown in SEQ ID No: 3, and/or

the beta chain variable domain may have at least one of the following mutations:

CDR1 H3	R
CDR3 N10	E
CDR3 N10	R

with reference to the numbering shown in SEQ ID No: 4.

[0027] The TCR of the invention may comprise an alpha chain variable domain of an amino acid sequence of amino acid residues 1-105 of SEQ ID No: 3 or 7

or an amino acid sequence in which amino acid residues 1-27, 34-47, and 54-90 thereof have at least 90% or 95% identity to the sequence of amino acid residues 1-27, 34-47, and 54-90 respectively of SEQ ID No: 3 or 5 or 7 and in which amino acid residues 28-33, 48-53 and 91-105 are amino acid residues 28-33, 48-53 and 91-105 respectively of SEQ ID No: 3 or 5 or 7.

[0028] The TCR comprises the alpha chain variable domain of a sequence in which:

1. (i) amino acid residues 1-27 thereof have (a) at least 90% identity to the sequence of amino acid residues 1-27 of SEQ ID No: 3 or (b) one, two or three amino acid insertions, deletions or substitutions relative to the sequence of (a);
2. (ii) amino acid residues 28-33 is VSPFSN
3. (iii) amino acid residues 34-47 thereof have (a) at least 90% identity to the sequence of amino acid residues 34-47 of SEQ ID NO: 3 or (b) one, two or three amino acid insertions, deletions or substitutions relative to the sequence of (a);
4. (iv) amino acid residues 48-53 may be LTIMTF or LTRMTF
5. (v) amino acid residues 54-90 thereof have (a) at least 90% identity to the sequence of amino acid residues 54-90 of SEQ ID No: 3 or (b) one, two or three insertions, deletions or substitutions relative to the sequence of (a); and
6. (vi) amino acid residues 91-105 may be CWSGGTDSWGKLQF

[0029] The TCR of the invention may comprise an beta chain variable domain of an amino acid sequence of SEQ ID No: 4 or 8-10

or an amino acid sequence in which amino acid residues 1-45, 51-67, 73-109 thereof have at least 90% or 95% identity to the sequence of amino acid residues 1-45, 51-67, 73-109 respectively of SEQ ID No: 4 or 8-10 and in which amino acid residues 46-50, 68-72 and 110-123 have at least 90% or 95% identity to the sequence of amino acid residues 46-50, 68-72 and 110-123 respectively of SEQ ID No: 4 or 8-10.

[0030] The TCR comprises the beta chain variable domain of a sequence in which:

1. (i) amino acid residues 1-45 thereof have (a) at least 90% identity to the amino acid

- sequence of residues 1-45 of SEQ ID No: 4 or (b) one, two or three amino acid insertions, deletions or substitutions relative to the sequence of (a);
2. (ii) amino acid residues 46-50 may be KGHDR or KGRDR
 3. (iii) amino acid residues 51-67 thereof have (a) at least 90% identity to the sequence of amino acid residues 51-67 of SEQ ID NO: 4 or (b) one, two or three amino acid insertions, deletions or substitutions relative to the sequence of (a);
 4. (iv) amino acid residues 68-72 may be SVFDK;
 5. (v) amino acid residues 73-109 thereof may have (a) at least 90% identity to the sequence of amino acid residues 73-109 of SEQ ID NO: 4 or (b) one, two or three amino acid insertions, deletions or substitutions relative to the sequence of (a); and
 6. (vi) amino acid residues 110-123 is CATSGQGAYREQFF or CATSGQGAYKEQFF, or CATSGQGAYNEQFF.

[0031] A TCR of the invention may have one of the following combinations of alpha and beta chain variable domains:

Alpha Chain SEQ ID No	Beta Chain SEQ ID No
3	4
3	6
3	8
3	9
3	10
5	4
5	6
5	8
5	9
5	10
7	4
7	6
7	8
7	9
7	10

[0032] Within the scope of the invention are phenotypically silent variants of any TCR of the invention disclosed herein. As used herein the term "phenotypically silent variants" is understood to refer to a TCR which incorporates one or more further amino acid changes in addition to those set out above which TCR has a similar phenotype to the corresponding TCR without said change(s). For the purposes of this application, TCR phenotype comprises

antigen binding specificity (K_D and/or binding half life) and antigen specificity. A phenotypically silent variant may have a K_D and/or binding half-life for the GVYDGEEHSV (SEQ ID No: 1) HLA-A*0201 complex within 10% of the measured K_D and/or binding half-life of the corresponding TCR without said change(s), when measured under identical conditions (for example at 25°C and on the same SPR chip). Suitable conditions are further defined in Example 3. Antigen specificity is further defined below. As is known to those skilled in the art, it may be possible to produce TCRs that incorporate changes in the constant and/or variable domains thereof compared to those detailed above without altering the affinity for the interaction with the GVYDGEEHSV (SEQ ID No: 1) HLA-A*0201 complex. In particular, such silent mutations may be incorporated within parts of the sequence that are known not to be directly involved in antigen binding (e.g. outside the CDRs). Such trivial variants are included in the scope of this invention. Those TCRs in which one or more conservative substitutions have been made also form part of this invention.

[0033] Mutations can be carried out using any appropriate method including, but not limited to, those based on polymerase chain reaction (PCR), restriction enzyme-based cloning, or ligation independent cloning (LIC) procedures. These methods are detailed in many of the standard molecular biology texts. For further details regarding polymerase chain reaction (PCR) and restriction enzyme-based cloning, see Sambrook & Russell, (2001) *Molecular Cloning - A Laboratory Manual* (3rd Ed.) CSHL Press. Further information on ligation independent cloning (LIC) procedures can be found in Rashtchian, (1995) *Curr Opin Biotechnol* 6(1): 30-6.

[0034] The TCRs of the invention have the property of binding the MAGE-B2 peptide, GVYDGEEHSV (SEQ ID No: 1) HLA-A2 complex. The TCRs of the invention have been found to be highly specific for those MAGE epitopes relative to other, irrelevant epitopes, and are thus particularly suitable as targeting vectors for delivery of therapeutic agents or detectable labels to cells and tissues displaying those epitopes. Specificity in the context of TCRs of the invention relates to their ability to recognise HLA-A*0201 target cells that are positive for the peptide GVYDGEEHSV, whilst having minimal ability to recognise HLA-A*0201 target cells that are negative for the peptide, or HLA cells that display the MAGE A4 peptide GVYDGREHTV. To test specificity, the TCRs may be in soluble form and/or may be expressed on the surface of T cells. Recognition may be determined by measuring the level of T cell activation in the presence of a TCR and target cells. In this case, minimal recognition of peptide negative or MAGE A4 target cells is defined as a level of T cell activation of less than 10%, preferably less than 5%, and more preferably less than 1%, of the level produced in the presence of peptide positive target cells, when measured under the same conditions. For soluble TCRs of the invention, specificity may be determined at a therapeutically relevant TCR concentration. A therapeutically relevant concentration may be defined as a TCR concentration of 10^{-9} M or below, and/or a concentration of up to 100, preferably up to 1000, fold greater than the corresponding EC₅₀ value. Peptide positive cells may be obtained by peptide-pulsing or, more preferably, they may naturally present said peptide. Preferably, both peptide positive and peptide negative cells are human cells.

[0035] Certain TCRs of the invention have been found to be highly suitable for use in adoptive therapy. Such TCRs may have a K_D for the complex of less than the 200 μM , for example from about 0.05 μM to about 20 μM or about 100 μM and/or have a binding half-life ($T_{1/2}$) for the complex in the range of from about 0.5 seconds to about 12 minutes. In some embodiments, TCRs of the invention may have a K_D for the complex of from about 0.05 μM to about 20 μM , about 0.1 μM to about 5 μM or about 0.1 μM to about 2 μM . Without wishing to be bound by theory, there seems to be an optimum window of affinity for TCRs with therapeutic use in adoptive cell therapy. Naturally occurring TCRs recognising epitopes from tumour antigens are generally of too low affinity (20 μM to 50 μM) and very high affinity TCRs (in the nanomolar range or higher) suffer from cross-reactivity issues (Robbins et al (2008) J. Immunol. 180 6116-6131; Zhao et al (2007) J. Immunol. 179 5845-5854; Schmid et al (2010) J. Immunol 184 4936-4946).

[0036] The TCRs of the invention may be $\alpha\beta$ heterodimers or may be in single chain format. Single chain formats include $\alpha\beta$ TCR polypeptides of the $V\alpha$ -L- $V\beta$, $V\beta$ -L- $V\alpha$, $V\alpha$ -C α -L- $V\beta$ or $V\alpha$ -L- $V\beta$ -C β types, wherein $V\alpha$ and $V\beta$ are TCR α and β variable regions respectively, C α and C β are TCR α and β constant regions respectively, and L is a linker sequence. For use as a targeting agent for delivering therapeutic agents to the antigen presenting cell the TCR may be in soluble form (i.e. having no transmembrane or cytoplasmic domains). For stability, soluble $\alpha\beta$ heterodimeric TCRs preferably have an introduced disulfide bond between residues of the respective constant domains, as described, for example, in WO 03/020763. One or both of the constant domains present in an $\alpha\beta$ heterodimer of the invention may be truncated at the C terminus or C termini, for example by up to 15, or up to 10 or up to 8 or fewer amino acids. For use in adoptive therapy, an $\alpha\beta$ heterodimeric TCR may, for example, be transfected as full length chains having both cytoplasmic and transmembrane domains. TCRs for use in adoptive therapy may contain a disulphide bond corresponding to that found in nature between the respective alpha and beta constant domains, additionally or alternatively a non-native disulphide bond may be present.

[0037] As will be obvious to those skilled in the art, it may be possible to truncate the sequences provided at the C-terminus and/or N-terminus thereof, by 1, 2, 3, 4, 5 or more residues, without substantially affecting the binding characteristics of the TCR. All such trivial variants are encompassed by the present invention.

[0038] Alpha-beta heterodimeric TCRs of the invention usually comprise an alpha chain TRAC constant domain sequence and a beta chain TRBC1 or TRBC2 constant domain sequence. The alpha and beta chain constant domain sequences may be modified by truncation or substitution to delete the native disulfide bond between Cys4 of exon 2 of TRAC and Cys2 of exon 2 of TRBC1 or TRBC2. The alpha and beta chain constant domain sequences may also be modified by substitution of cysteine residues for Thr 48 of TRAC and Ser 57 of TRBC1 or TRBC2, the said cysteines forming a disulfide bond between the alpha and beta constant domains of the TCR.

[0039] Some TCRs of the invention have a binding affinity for, and/or a binding half-life for, the

GVYDGEEHSV -HLA-A2 complex substantially higher than that of the reference MAGE-B2 TCR, Increasing the binding affinity of a native TCR often reduces the specificity of the TCR for its peptide-MHC ligand, and this is demonstrated in Zhao Yangbing et al., The Journal of Immunology, The American Association of Immunologists, US, vol. 179, No.9, 1 November 2007, 5845-5854. However, the TCRs of the invention which are derived from the parental TCR remain specific for the GVDGEEHSV -HLA-A2 complex, despite having substantially higher binding affinity than the parental TCR. Moreover, they are significantly more (e.g. at least ten-fold) selective for MAGE-B2 over MAGE-A4 than the parental TCR.

[0040] Binding affinity (inversely proportional to the equilibrium constant K_D) and binding half-life (expressed as $T_{1/2}$) can be determined using the Surface Plasmon Resonance (BIAcore) method of Example 3 herein. Measurements may be carried out at 25°C and at a pH between 7.1 and 7.5 using a soluble version of the TCR. It will be appreciated that doubling the affinity of a TCR results in halving the K_D . $T_{1/2}$ is calculated as $\ln 2$ divided by the off-rate (k_{off}). So doubling of $T_{1/2}$ results in a halving in k_{off} . K_D and k_{off} values for TCRs are usually measured for soluble forms of the TCR, i.e. those forms which are truncated to remove hydrophobic transmembrane domain residues. Therefore it is to be understood that a given TCR meets the requirement that it has a binding affinity for, and/or a binding half-life for, the GVDGREHTV -HLA-A2 complex if a soluble form of that TCR meets that requirement. Preferably the binding affinity or binding half-life of a given TCR is measured several times, for example 3 or more times, using the same assay protocol, and an average of the results is taken. The reference TCR has a K_D of approximately 17 μ M as measured by that method, and $T_{1/2}$ is approximately 1.6S

[0041] In a further aspect, the present invention provides nucleic acid encoding a TCR of the invention. In some embodiments, the nucleic acid is cDNA. In some embodiments, the invention provides nucleic acid comprising a sequence encoding an α chain variable domain of a TCR of the invention. In some embodiments, the invention provides nucleic acid comprising a sequence encoding a β chain variable domain of a TCR of the invention. The nucleic acid may be non-naturally occurring and/or purified and/or engineered.

[0042] In another aspect, the invention provides a vector which comprises nucleic acid of the invention. Preferably the vector is a TCR expression vector.

[0043] The invention also provides a cell harbouring a vector of the invention, preferably a TCR expression vector. The vector may comprise nucleic acid of the invention encoding in a single open reading frame, or two distinct open reading frames, the alpha chain and the beta chain respectively. Another aspect provides a cell harbouring a first expression vector which comprises nucleic acid encoding the alpha chain of a TCR of the invention, and a second expression vector which comprises nucleic acid encoding the beta chain of a TCR of the invention. Such cells are particularly useful in adoptive therapy. The cells of the invention may be isolated and/or recombinant and/or non-naturally occurring and/or engineered.

[0044] Since the TCRs of the invention have utility in adoptive therapy, the invention includes a non-naturally occurring and/or purified and/or or engineered cell, especially a T-cell, presenting a TCR of the invention. The invention also provides an expanded population of T cells presenting a TCR of the invention. There are a number of methods suitable for the transfection of T cells with nucleic acid (such as DNA, cDNA or RNA) encoding the TCRs of the invention (see for example Robbins et al., (2008) J Immunol. 180: 6116-6131). T cells expressing the TCRs of the invention will be suitable for use in adoptive therapy-based treatment of cancer. As will be known to those skilled in the art, there are a number of suitable methods by which adoptive therapy can be carried out (see for example Rosenberg et al., (2008) Nat Rev Cancer 8(4): 299-308).

[0045] Soluble TCRs of the invention are useful for delivering detectable labels or therapeutic agents to the antigen presenting cells and tissues containing the antigen presenting cells. They may therefore be associated (covalently or otherwise) with a detectable label (for diagnostic purposes wherein the TCR is used to detect the presence of cells presenting the GVDGEEHSV-HLA-A2 complex); a therapeutic agent; or a PK modifying moiety (for example by PEGylation).

[0046] Detectable labels for diagnostic purposes include for instance, fluorescent labels, radiolabels, enzymes, nucleic acid probes and contrast reagents.

[0047] Therapeutic agents which may be associated with the TCRs of the invention include immunomodulators, radioactive compounds, enzymes (perforin for example) or chemotherapeutic agents (cisplatin for example). To ensure that toxic effects are exercised in the desired location the toxin could be inside a liposome linked to TCR so that the compound is released slowly. This will prevent damaging effects during the transport in the body and ensure that the toxin has maximum effect after binding of the TCR to the relevant antigen presenting cells.

[0048] Other suitable therapeutic agents include:

- small molecule cytotoxic agents, i.e. compounds with the ability to kill mammalian cells having a molecular weight of less than 700 Daltons. Such compounds could also contain toxic metals capable of having a cytotoxic effect. Furthermore, it is to be understood that these small molecule cytotoxic agents also include pro-drugs, i.e. compounds that decay or are converted under physiological conditions to release cytotoxic agents. Examples of such agents include cis-platin, maytansine derivatives, rachelmycin, calicheamicin, docetaxel, etoposide, gemcitabine, ifosfamide, irinotecan, melphalan, mitoxantrone, sorfimer sodiumphotofrin II, temozolomide, topotecan, trimetreate glucuronate, auristatin E vincristine and doxorubicin;
- peptide cytotoxins, i.e. proteins or fragments thereof with the ability to kill mammalian cells. For example, ricin, diphtheria toxin, pseudomonas bacterial exotoxin A, DNase and RNase;
- radio-nuclides, i.e. unstable isotopes of elements which decay with the concurrent

emission of one or more of α or β particles, or γ rays. For example, iodine 131, rhenium 186, indium 111, yttrium 90, bismuth 210 and 213, actinium 225 and astatine 213; chelating agents may be used to facilitate the association of these radio-nuclides to the high affinity TCRs, or multimers thereof;

- immuno-stimulants, i.e. immune effector molecules which stimulate immune response. For example, cytokines such as IL-2 and IFN- γ ,
- Superantigens and mutants thereof;
- TCR-HLA fusions;
- chemokines such as IL-8, platelet factor 4, melanoma growth stimulatory protein, etc;
- antibodies or fragments thereof, including anti-T cell or NK cell determinant antibodies (e.g. anti-CD3, anti-CD28 or anti-CD16);
- alternative protein scaffolds with antibody like binding characteristics
- complement activators;
- xenogeneic protein domains, allogeneic protein domains, viral/bacterial protein domains, viral/bacterial peptides.

[0049] One preferred embodiment is provided by a TCR of the invention associated (usually by fusion to an N-or C-terminus of the alpha or beta chain) with an anti-CD3 antibody, or a functional fragment or variant of said anti-CD3 antibody. Antibody fragments and variants/analogues which are suitable for use in the compositions and methods described herein include minibodies, Fab fragments, F(ab')₂ fragments, dsFv and scFv fragments, Nanobodies[™] (these constructs, marketed by Ablynx (Belgium), comprise synthetic single immunoglobulin variable heavy domain derived from a camelid (e.g. camel or llama) antibody) and Domain Antibodies (Domantis (Belgium), comprising an affinity matured single immunoglobulin variable heavy domain or immunoglobulin variable light domain) or alternative protein scaffolds that exhibit antibody like binding characteristics such as Affibodies (Affibody (Sweden), comprising engineered protein A scaffold) or Anticalins (Pieris (German), comprising engineered anticalins) to name but a few.

[0050] For some purposes, the TCRs of the invention may be aggregated into a complex comprising several TCRs to form a multivalent TCR complex. There are a number of human proteins that contain a multimerisation domain that may be used in the production of multivalent TCR complexes. For example the tetramerisation domain of p53 which has been utilised to produce tetramers of scFv antibody fragments which exhibited increased serum persistence and significantly reduced off-rate compared to the monomeric scFv fragment. (Willuda et al. (2001) J. Biol. Chem. 276 (17) 14385-14392). Haemoglobin also has a tetramerisation domain that could potentially be used for this kind of application. A multivalent TCR complex of the invention may have enhanced binding capability for the GGYDGREHTV HLA-A2 complex compared to a non-multimeric wild-type or T cell receptor heterodimer of the invention. Thus, multivalent complexes of TCRs of the invention are also included within the invention. Such multivalent TCR complexes according to the invention are particularly useful for tracking or targeting cells presenting particular antigens *in vitro* or *in vivo*, and are also

useful as intermediates for the production of further multivalent TCR complexes having such uses.

[0051] As is well-known in the art, TCRs may be subject to post translational modifications. Glycosylation is one such modification, which comprises the covalent attachment of oligosaccharide moieties to defined amino acids in the TCR chain. For example, asparagine residues, or serine/threonine residues are well-known locations for oligosaccharide attachment. The glycosylation status of a particular protein depends on a number of factors, including protein sequence, protein conformation and the availability of certain enzymes. Furthermore, glycosylation status (i.e. oligosaccharide type, covalent linkage and total number of attachments) can influence protein function. Therefore, when producing recombinant proteins, controlling glycosylation is often desirable. Controlled glycosylation has been used to improve antibody-based therapeutics. (Jefferis R., Nat Rev Drug Discov. 2009 Mar;8(3):226-34.). For soluble TCRs of the invention glycosylation may be controlled in vivo, by using particular cell lines for example, or in vitro, by chemical modification. Such modifications are desirable, since glycosylation can improve pharmacokinetics, reduce immunogenicity and more closely mimic a native human protein (Sinclair AM and Elliott S., Pharm Sci. 2005 Aug; 94(8):1626-35).

[0052] For administration to patients, the TCRs, nucleic acids and/or cells of the invention (usually associated with a detectable label or therapeutic agent), may be provided in a pharmaceutical composition together with a pharmaceutically acceptable carrier or excipient. Therapeutic or imaging TCRs in accordance with the invention will usually be supplied as part of a sterile, pharmaceutical composition which will normally include a pharmaceutically acceptable carrier. This pharmaceutical composition may be in any suitable form, (depending upon the desired method of administering it to a patient). It may be provided in unit dosage form, will generally be provided in a sealed container and may be provided as part of a kit. Such a kit would normally (although not necessarily) include instructions for use. It may include a plurality of said unit dosage forms.

[0053] The pharmaceutical composition may be adapted for administration by any appropriate route, preferably a parenteral (including subcutaneous, intramuscular, or preferably intravenous) route. Such compositions may be prepared by any method known in the art of pharmacy, for example by mixing the active ingredient with the carrier(s) or excipient(s) under sterile conditions.

[0054] Dosages of the substances of the present invention can vary between wide limits, depending upon the disease or disorder to be treated, the age and condition of the individual to be treated, etc. and a physician will ultimately determine appropriate dosages to be used.

[0055] TCRs, pharmaceutical compositions, vectors, nucleic acids and cells of the invention may be provided in substantially pure form, for example at least 80%, at least 85 %, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% pure.

[0056] Also provided by the invention are:

- A TCR, nucleic acid or cell of the invention for use in medicine, preferably for use in a method of treating cancer, such as solid tumours (e.g., lung, liver and gastric metastases) and/or squamous cell carcinomas.
- the use of a TCR, nucleic acid or cell of the invention in the manufacture of a medicament for treating cancer.

[0057] Preferred features of each aspect of the invention are as for each of the other aspects *mutatis mutandis*.

[0058] The invention is further described in the following non-limiting examples.

[0059] Reference is made to the enclosed sequences, in which:

SEQ ID No. 1 is the MAGE B2 peptide

SEQ ID No. 2 is the MAGE A4 peptide

SEQ ID No: 3 is the amino acid sequence of the extracellular part of the alpha chain of a parental MAGE-A4-specific TCR, and SEQ ID No: 4 shows the amino acid sequence of the extracellular part of the beta chain of a parental MAGE-A4-specific TCR beta chain amino acid sequence.

[0060] SEQ ID No: 5 shows the amino acid sequence of the alpha chain of a native Lenti TCR (referred to herein as the "reference TCR"). The sequence is the same as that of The parental TCR except that a cysteine is substituted for T162 (i.e. T48 of the TRAC constant region). SEQ ID No: 6 is the beta chain of a native Lenti TCR (referred to herein as the "reference TCR"). The sequence is the same as that of the parental TCR except that a cysteine is substituted for S169 (i.e. S57 of the TRBC2 constant region) and A187 is substituted for C187 and D201 is substituted for N201.

[0061] SEQ ID No: 7 show the sequence of alpha chains which may be present in TCRs of the invention. The subsequences forming the CDR regions, or substantial parts of the CDR regions, are underlined.

[0062] SEQ ID Nos: 8, 9 and 10 show the sequence of the beta chain which may be present in TCRs of the invention. The subsequences forming the CDR regions, or substantial parts of the CDR regions are underlined.

[0063] SEQ ID Nos 11-16 show the sequences of TCRs which could not be improved by

selection and mutation in accordance with the present invention.

Examples

Example 1 - Cloning of the reference MAGE-A4 TCR alpha and beta chain variable region sequences into pGMT7-based expression plasmids

[0064] The parental MAGE-A4 TCR variable alpha and TCR variable beta domains of SEQ ID NOS: 3 and 4 respectively were cloned into pGMT7-based expression plasmids containing either C α or C β by standard methods described in (Molecular Cloning a Laboratory Manual Third edition by Sambrook and Russell). Plasmids were sequenced using an Applied Biosystems 3730xl DNA Analyzer. The reference MAGE-A4 TCR variable alpha and TCR variable beta domains of SEQ ID NOS: 3 and 4 respectively were cloned in the same way.

[0065] The DNA sequence encoding the TCR alpha chain variable region was ligated into pEX956, which was cut with restriction enzymes. The DNA sequence encoding the TCR beta chain variable region was ligated into pEXb21, which was also cut with restriction enzymes.

[0066] Ligated plasmids were transformed into competent *E.coli* strain XL1-blue cells and plated out on LB/agar plates containing 100 μ g/mL ampicillin. Following incubation overnight at 37°C, single colonies were picked and grown in 5 mL LB containing 100 μ g/mL ampicillin overnight at 37°C with shaking. Cloned plasmids were purified using a Miniprep kit (Qiagen) and the plasmids were sequenced using an Applied Biosystems 3730xl DNA Analyzer.

Example 2 - Expression, refolding and purification of soluble reference MAGE-A4 TCR

[0067] The expression plasmids containing the reference TCR α -chain and β -chain respectively, as prepared in Example 1, were transformed separately into *E.coli* strain BL21pLysS, and single ampicillin-resistant colonies were grown at 37°C in TYP (ampicillin 100 μ g/ml) medium to OD₆₀₀ of ~0.6-0.8 before inducing protein expression with 0.5 mM IPTG. Cells were harvested three hours post-induction by centrifugation for 30 minutes at 4000rpm in a Beckman J-6B. Cell pellets were lysed with 25 ml Bug Buster (NovaGen) in the presence of MgCl₂ and DNaseI. Inclusion body pellets were recovered by centrifugation for 30 minutes at 13000rpm in a Beckman J2-21 centrifuge. Three detergent washes were then carried out to remove cell debris and membrane components. Each time the inclusion body pellet was homogenised in a Triton buffer (50 mM Tris-HCl pH 8.0, 0.5% Triton-X100, 200 mM NaCl, 10 mM NaEDTA,) before being pelleted by centrifugation for 15 minutes at 13000rpm in a Beckman J2-21. Detergent and salt was then removed by a similar wash in the following buffer: 50 mM Tris-HCl pH 8.0, 1 mM NaEDTA. Finally, the inclusion bodies were divided into 30 mg aliquots and frozen at -70°C. Inclusion body protein yield was quantified by solubilising with 6

M guanidine-HCl and an OD measurement was taken on a Hitachi U-2001 Spectrophotometer. The protein concentration was then calculated using the extinction coefficient.

[0068] Approximately 15mg of TCR α chain and 15mg of TCR β chain solubilised inclusion bodies were thawed from frozen stocks and diluted into 10ml of a guanidine solution (6 M Guanidine-hydrochloride, 50 mM Tris HCl pH 8.1, 100 mM NaCl, 10 mM EDTA, 10 mM DTT), to ensure complete chain denaturation. The guanidine solution containing fully reduced and denatured TCR chains was then injected into 0.5 litre of the following refolding buffer: 100 mM Tris pH 8.1, 400 mM L-Arginine, 2 mM EDTA, 5 M Urea. The redox couple (cysteamine hydrochloride and cystamine dihydrochloride) to final concentrations of 6.6 mM and 3.7 mM respectively, were added approximately 5 minutes before addition of the denatured TCR chains. The solution was left for ~30 minutes. The refolded TCR was dialysed in Spectrapor 1 membrane (Spectrum; Product No. 132670) against 10 L H₂O for 18-20 hours. After this time, the dialysis buffer was changed twice to fresh 10 mM Tris pH 8.1 (10 L) and dialysis was continued at 5 °C $\pm$ 3 °C for another ~8 hours. Soluble TCR was separated from degradation products and impurities by loading the dialysed refold onto a POROS 50HQ anion exchange column and eluting bound protein with a gradient of 0-500mM NaCl in 10 mM Tris pH 8.1 over 50 column volumes using an Akta purifier (GE Healthcare). Peak fractions were pooled and a cocktail of protease inhibitors (Calbiochem) were added. The pooled fractions were then stored at 4 °C and analysed by Coomassie-stained SDS-PAGE before being pooled and concentrated. Finally, the soluble TCR was purified and characterised using a GE Healthcare Superdex 75HR gel filtration column pre-equilibrated in PBS buffer (Sigma). The peak eluting at a relative molecular weight of approximately 50 kDa was pooled and concentrated prior to characterisation by BIAcore surface plasmon resonance analysis.

Example 3 - Binding characterisation

BIAcore Analysis

[0069] A surface plasmon resonance biosensor (BIAcore 3000™) can be used to analyse the binding of a soluble TCR to its peptide-MHC ligand. This is facilitated by producing soluble biotinylated peptide-HLA ("pHLA") complexes which can be immobilised to a streptavidin-coated binding surface (sensor chip). The sensor chips comprise four individual flow cells which enable simultaneous measurement of T-cell receptor binding to four different pHLA complexes. Manual injection of pHLA complex allows the precise level of immobilised class I molecules to be manipulated easily.

[0070] Biotinylated class I HLA-A*0201 molecules were refolded *in vitro* from bacterially-expressed inclusion bodies containing the constituent subunit proteins and synthetic peptide, followed by purification and *in vitro* enzymatic biotinylation (O'Callaghan et al. (1999) Anal. Biochem. 266: 9-15). HLA-A*0201-heavy chain was expressed with a C-terminal biotinylation

tag which replaces the transmembrane and cytoplasmic domains of the protein in an appropriate construct. Inclusion body expression levels of ~75 mg/litre bacterial culture were obtained. The MHC light-chain or β 2-microglobulin was also expressed as inclusion bodies in *E.coli* from an appropriate construct, at a level of ~500 mg/litre bacterial culture.

[0071] *E. coli* cells were lysed and inclusion bodies are purified to approximately 80% purity. Protein from inclusion bodies was denatured in 6 M guanidine-HCl, 50 mM Tris pH 8.1, 100 mM NaCl, 10 mM DTT, 10 mM EDTA, and was refolded at a concentration of 30 mg/litre heavy chain, 30 mg/litre β 2m into 0.4 M L-Arginine, 100 mM Tris pH 8.1, 3.7 mM cystamine dihydrochloride, 6.6 mM cysteamine hydrochloride, 4 mg/L of the MAGE-A4 GGYDGREHTV or MAGE-B2 GGYDGEEHSV peptide required to be loaded by the HLA-A*02 molecule, by addition of a single pulse of denatured protein into refold buffer at $< 5^{\circ}\text{C}$. Refolding was allowed to reach completion at 4°C for at least 1 hour.

[0072] Buffer was exchanged by dialysis in 10 volumes of 10 mM Tris pH 8.1. Two changes of buffer were necessary to reduce the ionic strength of the solution sufficiently. The protein solution was then filtered through a $1.5\mu\text{m}$ cellulose acetate filter and loaded onto a POROS 50HQ anion exchange column (8 ml bed volume). Protein was eluted with a linear 0-500 mM NaCl gradient in 10 mM Tris pH 8.1 using an Akta purifier (GE Healthcare). HLA-A*0201-peptide complex eluted at approximately 250 mM NaCl, and peak fractions were collected, a cocktail of protease inhibitors (Calbiochem) was added and the fractions were chilled on ice.

[0073] Biotin-tagged pHLA molecules were buffer exchanged into 10 mM Tris pH 8.1, 5 mM NaCl using a GE Healthcare fast desalting column equilibrated in the same buffer. Immediately upon elution, the protein-containing fractions were chilled on ice and protease inhibitor cocktail (Calbiochem) was added. Biotinylation reagents were then added: 1 mM biotin, 5 mM ATP (buffered to pH 8), 7.5 mM MgCl_2 , and 5 $\mu\text{g/ml}$ BirA enzyme (purified according to O'Callaghan et al. (1999) Anal. Biochem. 266: 9-15). The mixture was then allowed to incubate at room temperature overnight.

[0074] The biotinylated pHLA-A*0201 molecules were purified using gel filtration chromatography. AGE Healthcare Superdex 75 HR 10/30 column was pre-equilibrated with filtered PBS and 1 ml of the biotinylation reaction mixture was loaded and the column was developed with PBS at 0.5 ml/min using an Akta purifier (GE Healthcare). Biotinylated pHLA-A*0201 molecules eluted as a single peak at approximately 15 ml. Fractions containing protein were pooled, chilled on ice, and protease inhibitor cocktail was added. Protein concentration was determined using a Coomassie-binding assay (PerBio) and aliquots of biotinylated pHLA-A*01 molecules were stored frozen at -20°C .

[0075] Such immobilised complexes are capable of binding both T-cell receptors and the coreceptor CD8 α , both of which may be injected in the soluble phase. The pHLA binding properties of soluble TCRs are observed to be qualitatively and quantitatively similar if the TCR is used either in the soluble or immobilised phase. This is an important control for partial activity of soluble species and also suggests that biotinylated pHLA complexes are biologically

as active as non-biotinylated complexes.

[0076] The BIAcore 3000™ surface plasmon resonance (SPR) biosensor measures changes in refractive index expressed in response units (RU) near a sensor surface within a small flow cell, a principle that can be used to detect receptor ligand interactions and to analyse their affinity and kinetic parameters. The BIAcore experiments were performed at a temperature of 25°C, using PBS buffer (Sigma, pH 7.1-7.5) as the running buffer and in preparing dilutions of protein samples. Streptavidin was immobilised to the flow cells by standard amine coupling methods. The pHLA complexes were immobilised via the biotin tag. The assay was then performed by passing soluble TCR over the surfaces of the different flow cells at a constant flow rate, measuring the SPR response in doing so.

Equilibrium binding constant

[0077] The above BIAcore analysis methods were used to determine equilibrium binding constants. Serial dilutions of the disulfide linked soluble heterodimeric form of the reference MAGE-A4 TCR were prepared and injected at constant flow rate of 5 µl min⁻¹ over two different flow cells; one coated with ~1000 RU of specific GVDGREHTV HLA-A*0201 complex, the second coated with ~1000 RU of non-specific complex. Response was normalised for each concentration using the measurement from the control cell. Normalised data response was plotted versus concentration of TCR sample and fitted to a non-linear curve fitting model in order to calculate the equilibrium binding constant, K_D . (Price & Dwek, Principles and Problems in Physical Chemistry for Biochemists (2nd Edition) 1979, Clarendon Press, Oxford). The disulfide linked soluble form of the reference MAGE-A4 TCR (Example 2) demonstrated a K_D of approximately 2.00 µM. From the same BIAcore data the $T_{1/2}$ was approximately 0.95 s.

Kinetic Parameters

[0078] The above BIAcore analysis methods were also used to determine equilibrium binding constants and off-rates.

[0079] For high affinity TCRs (see Example 4 below) K_D was determined by experimentally measuring the dissociation rate constant, k_{off} , and the association rate constant, k_{on} . The equilibrium constant K_D was calculated as k_{off}/k_{on} .

[0080] TCR was injected over two different cells one coated with ~1000 RU of specific GVDGREHTV

[0081] HLA-A*0201 complex, the second coated with ~1000 RU of non-specific complex. Flow rate was set at 50 µl/min. Typically 250 µl of TCR at ~ 1 µM concentration was injected. Buffer

was then flowed over until the response had returned to baseline or >2 hours had elapsed. Kinetic parameters were calculated using BIAevaluation software. The dissociation phase was fitted to a single exponential decay equation enabling calculation of half-life.

Example 4 - Preparation of high affinity TCRs of the invention

Expression plasmids containing the TCR α -chain and β -chain respectively were prepared as in Example 1:

[0082]

TCR ID	Alpha Chain SEQ ID No	Beta Chain SEQ ID No
TCR1 (parental)	3	4
TCR2 958	7 (I3R)	4
TCR3 962	3	8 (H3R)
TCR4 1028	3	9 (N10R)
TCR5 1030	7 (I3R)	9 (N10R)
TCR6 1034	3	10 (N10K)

[0083] The plasmids were transformed separately into *E.coli* strain BL21pLysS, and single ampicillin-resistant colonies grown at 37°C in TYP (ampicillin 100 μ g/ml) medium to OD₆₀₀ of ~0.6-0.8 before inducing protein expression with 0.5 mM IPTG. Cells were harvested three hours post-induction by centrifugation for 30 minutes at 4000rpm in a Beckman J-6B. Cell pellets were lysed with 25 ml Bug Buster (Novagen) in the presence of MgCl₂ and DNaseI. Inclusion body pellets were recovered by centrifugation for 30 minutes at 13000rpm in a Beckman J2-21 centrifuge. Three detergent washes were then carried out to remove cell debris and membrane components. Each time the inclusion body pellet was homogenised in a Triton buffer (50 mM Tris-HCl pH 8.0, 0.5% Triton-X100, 200 mM NaCl, 10 mM NaEDTA,) before being pelleted by centrifugation for 15 minutes at 13000rpm in a Beckman J2-21. Detergent and salt was then removed by a similar wash in the following buffer: 50 mM Tris-HCl pH 8.0, 1 mM NaEDTA. Finally, the inclusion bodies were divided into 30 mg aliquots and frozen at -70°C. Inclusion body protein yield was quantified by solubilising with 6 M guanidine-HCl and an OD measurement was taken on a Hitachi U-2001 Spectrophotometer. The protein concentration was then calculated using the extinction coefficient.

[0084] Approximately 10mg of TCR α chain and 10mg of TCR β chain solubilised inclusion bodies for each TCR of the invention were diluted into 10ml of a guanidine solution (6 M Guanidine-hydrochloride, 50 mM Tris HCl pH 8.1, 100 mM NaCl, 10 mM EDTA, 10 mM DTT), to ensure complete chain denaturation. The guanidine solution containing fully reduced and

denatured TCR chains was then injected into 0.5 litre of the following refolding buffer: 100 mM Tris pH 8.1, 400 mM L-Arginine, 2 mM EDTA, 5 M Urea. The redox couple (cysteamine hydrochloride and cystamine dihydrochloride) to final concentrations of 6.6 mM and 3.7 mM respectively, were added approximately 5 minutes before addition of the denatured TCR chains. The solution was left for ~30 minutes. The refolded TCR was dialysed in Spectrapor 1 membrane (Spectrum; Product No. 132670) against 10 L H₂O for 18-20 hours. After this time, the dialysis buffer was changed twice to fresh 10 mM Tris pH 8.1 (10 L) and dialysis was continued at 5 °C $\pm$ 3 °C for another ~8 hours.

[0085] Soluble TCR was separated from degradation products and impurities by loading the dialysed refold onto a POROS 50HQ anion exchange column and eluting bound protein with a gradient of 0-500mM NaCl in 10 mM Tris pH 8.1 over 15 column volumes using an Akta purifier (GE Healthcare). The pooled fractions were then stored at 4 °C and analysed by Coomassie-stained SDS-PAGE before being pooled and concentrated. Finally, the soluble TCRs were purified and characterised using a GE Healthcare Superdex 75HR gel filtration column pre-equilibrated in PBS buffer (Sigma). The peak eluting at a relative molecular weight of approximately 50 kDa was pooled and concentrated prior to characterisation by BIAcore surface plasmon resonance analysis.

[0086] The affinity profiles of the thus-prepared TCRs for the MAGE-A4 epitope or MAGE-B2 epitope were assessed using the method of Example 3, and compared with the reference TCR. The results are set forth in the following table:

	MAGE A4 K _D (μM)	MAGE-B2 K _D (μM)
Reference (TCR1)	65.1	17
TCR2	208.4	17.8
TCR3	195.5	16.61
TCR4	183.3	11.27
TCR5	484.0	18.77
TCR6	305.3	6.03

[0087] Attempts were also made to prepare high affinity TCRs based on combinations of SEQ ID Nos 11/12, 13/14, 15/16.

[0088] In the case of TCR A, which combines alpha chain of SEQ ID No 11 and the Beta chain of SEQ ID No. 12, cross-reactivity was noted between MAGE-A1, MAGE-A10 and PRAME. It was not possible to remove this cross-reactivity by mutation and selection.

[0089] TCR B combines the alpha chain of SEQ ID No 12 and the Beta chain of SEQ ID No. 14. TCR B could not be folded to form a soluble TCR, so no binding characterisation was possible.

[0090] TCR C combines the alpha chain of SEQ ID NO 15 and the Beta chain of SEQ ID No 16. This TCR was soluble when expressed and could bind to antigen. However, when expressed in T-cells, TCR C showed no activity.

Example 5 - Transfection of T-cells with parental and variant MAGE-A4 TCRs

(a) Lentiviral vector preparation by Express-In mediated transient transfection of 293T cells

[0091] A 3rd generation lentiviral packaging system was used to package lentiviral vectors containing the gene encoding the desired TCR. 293T cells were transfected with 4 plasmids (one lentiviral vector containing the TCR alpha chain-P2A-TCR beta chain single ORF gene described in Example 5c (below), and 3 plasmids containing the other components necessary to construct infective but non-replicative lentiviral particles) using Express-In mediated transfection (Open Biosystems).

[0092] For transfection one T150 flask of 293T cells in exponential growth phase was taken, with cells evenly distributed on the plate, and slightly more than 50% confluent. Express-In aliquots were brought to room temperature. 3 ml Serum-Free Medium (RPMI 1640 + 10mM HEPES) were placed in a sterile 15 ml conical tube. 174 µl of Express-In Reagent were added directly into the Serum-Free Medium (this provides for a 3.6:1 weight ratio of Reagent to DNA). This was mixed thoroughly by inverting tubes 3-4 times and incubated at room temperature for 5-20 minutes.

[0093] In a separate 1.5 ml microtube was added 15 µg plasmid DNA to premixed packaging mix aliquots (containing 18 µg pRSV.REV (Rev expression plasmid), 18 µg pMDLg/p.RRE (Gag/Pol expression plasmid), 7 µg pVSV-G (VSV glycoprotein expression plasmid), usually ~22 µl, and pipetted up and down to ensure homogeneity of the DNA mix. Approx 1 mL of Express-In/Serum-Free Medium was added to the DNA mix dropwise then pipetted up and down gently before transferring back to the remainder of the Express-In/Serum-Free Medium. The tube was inverted 3-4 times and incubated at room temperature for 15-30 minutes. Old culture medium was removed from the flask of cells. Express-In/medium/DNA (3mL) complex was added directly into the bottom of an upright flask of 293T cells. Slowly, the flask was placed flat to cover the cells and very gently rocked to ensure even distribution. After 1 minute 22 ml fresh culture medium (R10+HEPES: RPMI 1640, 10% heat-inactivated FBS, 1% Pen/Strep/L-glutamine, 10 mM HEPES) was added and the flask carefully returned to the incubator. This was incubated overnight at 37°C/5% CO₂. After 24 hours, the medium containing packaged lentiviral vectors was harvested.

[0094] To harvest the packaged lentiviral vectors, the cell culture supernatant was filtered through a 0.45micron nylon syringe filter, the culture medium centrifuged at 10,000 g for 18

hours (or 112,000 g for 2 hours), most of the supernatant removed (taking care not to disturb the pellet) and the pellet resuspended in the remaining few mL of supernatant (usually about 2 ml from a 31 ml starting volume per tube). This was snap frozen on dry ice in 1 ml aliquots and stored at -80°C.

(b) Transduction of T cells with packaged lentiviral vectors containing gene of interest

[0095] Prior to transduction with the packaged lentiviral vectors, human T cells (CD8 or CD4 or both depending on requirements) were isolated from the blood of healthy volunteers. These cells were counted and incubated overnight in R10 containing 50 U/mL IL-2 at 1×10^6 cells per ml (0.5 mL/well) in 48 well plates with pre-washed anti-CD3/CD28 antibody-coated microbeads (Dynabeads® T cell expander, Invitrogen) at a ratio of 3 beads per cell.

[0096] After overnight stimulation, 0.5 ml of neat packaged lentiviral vector was added to the desired cells. This was incubated at 37°C/5% CO₂ for 3 days. 3 days post-transduction the cells were counted and diluted to 0.5×10^6 cells/ml. Fresh medium containing IL-2 was added as required. Beads were removed 5-7 days post-transduction. Cells were counted and fresh medium containing IL-2 replaced or added at 2 day intervals. Cells were kept between 0.5×10^6 and 1×10^6 cells/mL. Cells were analysed by flow cytometry from day 3 and used for functional assays (e.g. ELISpot for IFN γ release, see Example 6) from day 5. From day 10, or when cells are slowing division and reduced in size, cells are frozen in aliquots of at least 4×10^6 cells/vial (at 1×10^7 cells/ml in 90% FBS/10% DMSO) for storage.

Example 6 - Activation of MAGE B2 TCR engineered T cells

[0097] The following assay was carried out to demonstrate the activation of TCR-transduced cytotoxic T lymphocytes (CTLs) in response to tumour cell lines. IFN- γ production, as measured using the ELISPOT assay, was used as a read-out for cytotoxic T lymphocyte (CTL) activation.

ELISPOTs

Reagents

[0098] Assay media: 10% FCS (Gibco, Cat# 2011-09), 88% RPMI 1640 (Gibco, Cat# 42401), 1% glutamine (Gibco Cat# 25030) and 1% penicillin/streptomycin (Gibco Cat# 15070-063).

Wash buffer: 0.01M PBS/0.05% Tween 20

PBS (Gibco Cat# 10010)

The Human IFN γ ELISPOT kit (BD Bioscience; Cat# 551849) containing capture and detection antibodies and Human IFN- γ PVDF ELISPOT 96 well plates, with associated AEC substrate set (BD Bioscience, Cat# 551951)

Methods

Target cell preparation

[0099] The target cells used in this method were natural epitope-presenting cells: A375 human melanoma cells which are both HLA-A2⁺ MAGE A10⁺. HCT116 human colon cancer, which are HLA-A2⁺ MAGE A10⁻, were used as a negative control. Sufficient target cells (50,000 cells/well) were washed by centrifugation three times at 1200 rpm, 10 min in a Megafuge[®] 1.0 (Heraeus). Cells were then re-suspended in assay media at 10⁶ cells/ml.

Effector Cell Preparation

[0100] The effector cells (T cells) used in this method were peripheral blood lymphocytes (PBL), obtained by negative selection using CD14 and CD25 microbead kits (Miltenyi Biotech Cat# 130-050-201 and 130-092-983 respectively) from freshly isolated peripheral blood mononuclear cells (PBMC) from the venous blood of healthy volunteers. Cells were stimulated with antiCD3/CD28 coated beads (Dynabeads[®] T cell expander, Invitrogen), transduced with lentivirus carrying the gene encoding the full $\alpha\beta$ TCR of interest (based on the construct described in Example 5) and expanded in assay media containing 50U/mL IL-2 until between 10 and 13 days post transduction. These cells were then placed in assay media prior to washing by centrifugation at 1200 rpm, 10 min in a Megafuge[®] 1.0 (Heraeus). Cells were then re-suspended in assay media at a 4X the final required concentration.

[0101] Plates were prepared as follows: 100 μ L anti-IFN- γ capture antibody was diluted in 10 ml sterile PBS per plate. 100 μ L of the diluted capture antibody was then dispensed into each well. The plates were then incubated overnight at 4°C. Following incubation the plates were washed (programme 1, plate type 2, Ultrawash Plus 96-well plate washer; Dynex) to remove the capture antibody. Plates were then blocked by adding 200 μ L of assay media to each well and incubated at room temperature for two hours. The assay media was then washed from the plates (programme 1, plate type 2, Ultrawash Plus 96-well plate washer, Dynex) and any

remaining media was removed by flicking and tapping the ELISPOT plates on a paper towel.

[0102] The constituents of the assay were then added to the ELISPOT plate in the following order:

50 µL of target cells 10^6 cells/ml (giving a total of 50,000 target cells/well)

50 µL media (assay media)

50 µL effector cells (20,000 TCR-transduced PBL cells/well)

[0103] The plates were then incubated overnight (37°C / 5%CO₂). The next day the plates were washed three times (programme 1, plate type 2, Ultrawash Plus 96-well plate washer, Dynex) with wash buffer and tapped dry on paper towel to remove excess wash buffer. 100 µl of primary detection antibody was then added to each well. The primary detection antibody was diluted into 10mL of dilution buffer (the volume required for a single plate) using the dilution specified in the manufacturer's instructions. Plates were then incubated at room temperature for at least 2 hours prior to being washed three times (programme 1, plate type 2, Ultrawash Plus 96-well plate washer, Dynex) with wash buffer; excess wash buffer was removed by tapping the plate on a paper towel.

[0104] Secondary detection was performed by adding 100 µL of diluted streptavidin-HRP to each well and incubating the plate at room temperature for 1 hour. The streptavidin-HRP was diluted into 10mL dilution buffer (the volume required for a single plate), using the dilution specified in the manufacturer's instructions. The plates were then washed three times (programme 1, plate type 2, Ultrawash Plus 96-well plate washer, Dynex) with wash buffer and tapped on paper towel to remove excess wash buffer. Plates were then washed twice with PBS by adding 200 µL to each well, flicking the buffer off and tapping on a paper towel to remove excess buffer. No more than 15 min prior to use, one drop (20 uL) of AEC chromogen was added to each 1 ml of AEC substrate and mixed. 10ml of this solution was prepared for each plate; 100 µL was added per well. The plate was then protected from light using foil, and spot development monitored regularly, usually occurring within 5-20 min. The plates were washed in tap water to terminate the development reaction, and shaken dry prior to their disassembly into three constituent parts. The plates were then allowed to dry at room temperature for at least 2 hours prior to counting the spots using an Immunospot[®] Plate reader (CTL; Cellular Technology Limited).

Example 7 - Identification of the binding motif by substitution with all alternative amino acids

[0105] Variants of the native MAGE-B2 peptide were obtained in which the amino acid residue

at each position was sequentially replaced with all 19 alternative naturally-occurring amino acid, such that 171 peptides were prepared in total. The native and amino-acid substituted peptides were pulsed on to antigen presenting cells, and interferon γ (IFN γ) production, as measured using the ELISpot assay, used as a read-out for the activation of T cells transduced with TCR1. Essential positions were defined by a greater than 50% reduction in T cell activity relative to the native peptide.

[0106] ELISpot assays were carried as described in Example 6.

[0107] The tolerated residues at each position of the peptide are shown below. Underlined amino acids represent the native residue at the corresponding position in the peptide.

Position	Tolerated residues
1	<u>G</u>
2	<u>V</u>
3	F <u>Y</u>
4	<u>D</u>
5	<u>G</u> N
6	DA <u>E</u> S
7	YSWTFQMHP <u>L</u> ANGDICE
8	FWWLMAYRKCTIQSH <u>G</u> PN
9	TVIAS <u>P</u> GKEQNH
10	FM <u>V</u> AI

[0108] It is therefore apparent that the MAGE B2 TCR4 makes contact with at least V2 Y3 and D4 of the peptide (SEQ ID no: 1) when in complex with HLA-A*0201 on the surface of antigen presenting cells.

SEQ ID No. 1 MAGE B2 Epitope
GVYDGEEHSV

SEQ ID No.2 MAGE A4 Epitope
GVYDGREHTV

SEQ ID No. 3 alpha variable chain
MKNQVEQSPQSLIILEGKNCTLQCNYTVSPFSNLRWYKQDTGRGPVSLTIMTFSENTKS
NGRYTATLDADTKQSSLHITASQLSDSASYICVVSGGTDSWGKLQF

SEQ ID No. 4 beta variable chain
MASLLFFCGAFYLLGTGSMADAVTQTPRNRITKTGKRIMLECSQTKGHDRMYWYRQDP
GLGLRLIYYSFDVKDINKGEISDGYSVSRQAQAKFSLSLSAIPNQATALYFCATSGQGAYN
EQFF

SEQ ID No. 5 alpha chain soluble form

MKKHLTTFLVILWLYFYRGNGKNQVEQSPQSLIILEGKNCTLQCNYTVSPFSNLRWYKQD
TGRGPVSLTIMTFSENTKSNGRYTATLDADTKQSSLHITASQLSDSASYICVVSGGTDSW
GKLQFGAGTQVVVTPDIQNPDPVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYI
TDKTVLDMRSMDFKSNSAVAWSNKSDFACANAFNNSIIPEDTFFPSPRESS

SEQ ID No 6 beta chain soluble form

MASLLFFCGAFYLLGTGSMADVTQTPRNRITKTGKRIMLECSQTKGHDRMYWYRQDP
GLGLRLIYYSFVDKINKGEISDGYSVSRQAQAKFSLSLESAIPNQTALYFCATSGQGAYN
EQFFGPGTRLTVLEDLKNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWW
NGKEVHSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRQVQFYGLSE
NDEWTQDRAKPVTQIVSAEAWGRAD

SEQ ID No. 7 mutant alpha variable chain

MKNQVEQSPQSLIILEGKNCTLQCNYTVSPFSNLRWYKQDTGRGPVSLTRMTFSENTKS
NGRYTATLDADTKQSSLHITASQLSDSASYICVVSGGTDSWGKLQF

SEQ ID No. 8 mutant beta variable chain

MASLLFFCGAFYLLGTGSMADVTQTPRNRITKTGKRIMLECSQTKGRDRMYWYRQDP
GLGLRLIYYSFVDKINKGEISDGYSVSRQAQAKFSLSLESAIPNQTALYFCATSGQGAYN
EQFF

SEQ ID No. 9 mutant beta variable chain

MASLLFFCGAFYLLGTGSMADVTQTPRNRITKTGKRIMLECSQTKGHDRMYWYRQDP
GLGLRLIYYSFVDKINKGEISDGYSVSRQAQAKFSLSLESAIPNQTALYFCATSGQGAYR
EQFF

SEQ ID No. 10 mutant beta variable chain

MASLLFFCGAFYLLGTGSMADVTQTPRNRITKTGKRIMLECSQTKGHDRMYWYRQDP
GLGLRLIYYSFVDKINKGEISDGYSVSRQAQAKFSLSLESAIPNQTALYFCATSGQGAYK
EQFF

SEQ ID No. 11 alpha chain soluble form

METLLGLLILWLQLQWVSSKQEVTPQPAALSVPEGENLVLNCSFTDSAIYNLQWFRQDPG
KGLTSLLLIQSSQREQTSGRLNASLDKSSGRSTLYIAASQPGDSATYLCVGGYSTLTFG
KGTVLLVSPDNIQNPDPVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKTVL
DMRSMDFKSNSAVAWSNKSDFACANAFNNSIIPEDTFFPSPRESSCDVKLVEKSFETDTN
LNFQNLSVIGFRILLKLVAGFNLLMTLRLWSS

SEQ ID No 12 beta chain soluble form

MSISLLCCAAFPLLWAGPVNAGVTQTPKFRILKIGQSMTLQCAQDMNHNMYWYRQDP
GMGLKLIYYSVGAGITDKGEVPNGYNVSRSTTEDFPLRLELAAPSQTSVYFCASSYSRW
SPLHFGNGTRLTVTEDLNKVFPEVAVFEPSEAEISHTQKATLVCLATGFFPDHVELSWWW
VNGKEVHSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPRNHFRQVQFYGLS
ENDEWTQDRAKPVTQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLGKATLYAVLVS
ALVLMAMVKRKDF

SEQ ID No. 13 alpha chain soluble form

MQKEVEQNSGPLSVPEGAIASLNCTYSDRGSQSFFWYRQYSGKSPELIMFIYSNGDKED
GRFTAQLNKASQYVSLIRDSQPSDSATYLCVVKMANQAGTALIFGKGTTLSVSSNIQNP
DPAVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYITDKCVLDMRSMDFKSNSAVA
WSNKSDFACANAFNNSIIPEDTFFPSPESS

SEQ ID No 14 beta chain soluble form

MQDGGITQSPKFQVLRGTGQSMTELLCAQDMNHEMYWYRQDPGMGLRLIHYSVGAGITD
QGEVPNGYNVSRNLNREFSLRLESAAPSQTSVYFCASLGGLADEQFFGPGTRLTVLEDL
KNVFPPEVAVFEPSEAEISHTQKATLVCLATGFYPDHVELSWWWNGKEVHSGVCTDPQP
LKEQPALNDSRYALSSRLRVSATFWQDPRNHFRQVQFYGLSENDEWTQDRAKPVTQI
VSAEAWGRAD

SEQ ID No. 15 alpha chain soluble form

MKTFAGFSFLFLWLQLDCMSRGEDVEQSLFLSVREGDSSVINCTYTDSSSTYLYWYKQE
PGAGLQLLTYIFSNMDMKQDQRLTVLLNKKDKHLSLRIADTQTGDSAIYFCAERNNGAGS
YQLTFGKGTKLSVIPNIQNPDPVYQLRDSKSSDKSVCLFTDFDSQTNVSQSKDSDVYIT
DKTVLDMRSMDFKSNSAVAWSNKSDFACANAFNNSIIPEDTFFPSPESSCDVKLVEKSF
ETDTNLFQNLQSVIGFRILLKLVAGFNLLMTLRLWSS

SEQ ID No 16 beta chain soluble form

MGTSLLCWMALCLLGADHADTGVSQNPRHKITKRGQNVTFRCDPSEHNRLYWYRQTL
GQGPEFLTYFQNEAQLEKSRLSDRFSAPKGSFSTLEIQRTEQGDSAMYLCASSLFS
GVNTEAFFGQGTSLTVVEDLNKVFPPEVAVFEPSEAEISHTQKATLVCLATGFFPDHVEL
SWWWNGKEVHSGVSTDPQPLKEQPALNDSRYCLSSRLRVSATFWQNPARNHFRQVQF
YGLSENDEWTQDRAKPVTQIVSAEAWGRADCGFTSVSYQQGVLSATILYEILLGKATLYA
VLVSALVLMAMVKRKDF

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US2007077045A](#) [0008]
- [WO02094981A](#) [0009]
- [WO03020763A](#) [0036]

Non-patent literature cited in the description

- HILLIG et al. J. Mol. Biol., 2001, vol. 310, 1167-1176 [0010]
- ZHENG-CAI et al. Clin. Dev. Immunol., 2010, vol. 107, [0011]
- WAHL et al. Human Immunology, 2010, vol. 71, 14-22 [0011]
- LEFRANCLEFRANCT cell Receptor Factsbook Academic Press 20010000 [0022]
- SAMBROOKRUSSELL Molecular Cloning - A Laboratory Manual CSHL Press 20010000 [0033]
- RASHTCHIAN Curr Opin Biotechnol, 1995, vol. 6, 130-6 [0033]
- ROBBINS et al. J. Immunol., 2008, vol. 180, 6116-6131 [0035]
- ZHAO et al. J. Immunol., 2007, vol. 179, 5845-5854 [0035]
- SCMID et al. J. Immunol., 2010, vol. 184, 4936-4946 [0035]
- ZHAO YANGBING et al. The American Association of Immunologists The Journal of Immunology, 2007, vol. 179, 5845-5854 [0039]
- ROBBINS et al. J Immunol., 2008, vol. 180, 6116-6131 [0044]
- ROSENBERG et al. Nat Rev Cancer, 2008, vol. 8, 4299-308 [0044]
- WILLUDA et al. J. Biol. Chem., 2001, vol. 276, 1714385-14392 [0050]
- JEFFERIS R. Nat Rev Drug Discov., 2009, vol. 8, 3226-34 [0051]
- SINCLAIR AMELLIOTT S. Pharm Sci., 2005, vol. 94, 81626-35 [0051]
- SAMBROOKRUSSELL Molecular Cloning a Laboratory Manual [0064]
- O'CALLAGHAN et al. Anal. Biochem., 1999, vol. 266, 9-15 [0070] [0073]
- PRICEDWEK Principles and Problems in Physical Chemistry for Biochemists Clarendon Press 19790000 [0077]

PATENTKRAV

1. T-cellereceptor (TCR), der kan binde til GGYDGEHHSV (SEQ ID NO: 1) i kompleks med HLA-A*0201 med en dissociationskonstant på fra 0,05 μ M til 20,0 μ M ved måling med overfladeplasmonresonans ved 25 °C og ved et pH mellem 7,1 og 7,5 ved hjælp af en opløselig form af TCR'en, og har mindst en ti-fold bindingsselektivitet til SEQ ID NO: 1 i kompleks med HLA-A*0201 over binding til GGYDGREHTV (SEQ ID No 2) i kompleks med HLA-A*0201, hvor TCR'en omfatter

(A) det variable domæne af en sekvens' TCR-alfa-kæde, hvori:

(i) aminosyreresterne 1-27 har (a) mindst 90 % identitet med aminosyreresterens sekvens 1-27 ifølge SEQ ID NO: 3 eller (b) én, to eller tre aminosyreindsættelser eller -deletioner i forhold til sekvensen fra (a);

(ii) aminosyreresterne 28-33 er VSPFSN;

(iii) aminosyreresterne 34-47 har (a) mindst 90 % identitet med aminosyreresterens sekvens 34-47 ifølge SEQ ID NO: 3 eller (b) én, to eller tre aminosyreindsættelser eller -deletioner i forhold til sekvensen fra (a);

(iv) aminosyreresterne 48-53 er LTIMTF eller LTRMTF;

(v) aminosyreresterne 54-90 har (a) mindst 90 % identitet med aminosyreresterens sekvens 54-90 ifølge SEQ ID NO: 3 eller (b) én, to eller tre aminosyreindsættelser, -deletioner eller -substitutioner i forhold til sekvensen fra (a);

og

(vi) aminosyreresterne 91-105 er CVVSGGTDSWGKLQF; og

(B) det variable domæne af en sekvens TCR-beta-kæde, hvori:

(i) aminosyreresterne 1-45 har (a) mindst 90 % identitet med aminosyresekvensen for resterne 1-45 ifølge SEQ ID NO: 4 eller (b) én, to eller tre aminosyreindsættelser eller -deletioner i forhold til sekvensen fra (a);

(ii) aminosyreresterne 46-50 er KGHDR eller KGRDR;

(iii) aminosyreresterne 51-67 har (a) mindst 90 % identitet med aminosyreresterens sekvens 51-67 ifølge SEQ ID NO: 4 eller (b) én, to eller tre aminosyreindsættelser eller -deletioner i forhold til sekvensen fra (a);

(iv) aminosyreresterne 68-72 er SFDVK;

(v) aminosyreresterne 73-109 har (a) mindst 90 % identitet med aminosyreresternes sekvens 73-109 ifølge SEQ ID NO: 4 eller (b) én, to eller tre aminosyreindsætninger eller -deletioner i forhold til sekvensen fra (a); og

5 (vi) aminosyreresterne 110-123 er CATSGQGAYREQFF eller CATSGQGAYKEQFF eller CATSGQGAYNEQFF;

og hvor de variable TCR-domæner kommer i kontakt med mindst resterne V2, Y3 og D4 af GYDGEHSV (SEQ ID NO: 1).

2. TCR ifølge krav 1, der er i enkeltkædeformet af typen $V\alpha$ -L- $V\beta$, $V\beta$ -L- $V\alpha$,
10 $V\alpha$ - $C\alpha$ -L- $V\beta$ eller $V\alpha$ -L- $V\beta$ - $C\beta$, hvor $V\alpha$ og $V\beta$ henholdsvis er TCR α og β variable områder, $C\alpha$ og $C\beta$ henholdsvis er TCR α og β konstante områder, og L er en linker-sekvens.

3. TCR ifølge krav 1 eller 2, der er forbundet med en detekterbar markør, et terapeutisk middel og en PK-modificerende del.

15 4. TCR ifølge et hvilket som helst foregående krav, hvor det variable domæne af TCR-alfa-kæden indgår i aminosyresekvensen af aminosyreresterne 1-105 ifølge SEQ ID NO: 3 eller 7.

5. TCR ifølge et hvilket som helst foregående krav, hvor det variable domæne af TCR-beta-kæden indgår i aminosyresekvensen ifølge SEQ ID NO: 4 eller 8-10.

20 6. Nukleinsyre, der koder for en TCR ifølge et hvilket som helst af de foregående krav.

7. Isoleret eller ikke-naturligt forekommende celle, navnlig en T-celle, der præsenterer en TCR ifølge et hvilket som helst af krav 1 til 5.

8. Celle, der rummer

25 (a) en TCR-ekspressionsvektor, som omfatter nukleinsyre ifølge krav 6 i en åben læseramme, eller to særskilte åbne læserammer, der henholdsvis koder for alpha-kæden og beta-kæden; eller

(b) en første ekspressionsvektor, der omfatter nukleinsyre, som koder for alfa-kæden af en TCR ifølge et hvilket som helst af krav 1 til 5, og en anden

ekspressionsvektor, der omfatter nukleinsyre, som koder for betakæden af en TCR ifølge et hvilket som helst af krav 1 til 5.

5 **9.** Farmaceutisk sammensætning, der omfatter en TCR ifølge et hvilket som helst af krav 1 til 5, nukleinsyre ifølge krav 6 eller celle ifølge krav 7 eller krav 8, sammen med ét eller flere farmaceutisk acceptable bærestoffer eller hjælpestoffer.

10. TCR ifølge et hvilket som helst af krav 1 til 5, nukleinsyre ifølge krav 6 eller celle ifølge krav 7 eller krav 8 til anvendelse i medicin.

11. TCR, nukleinsyre eller celle til anvendelse ifølge krav 10, til anvendelse i en fremgangsmåde til behandling af cancer.