



(51) International Patent Classification:

C07D 213/81 (2006.01) *A61P 31/18* (2006.01)
A61K 31/44 (2006.01) *C07D 405/12* (2006.01)
A61K 31/4427 (2006.01) *C07D 493/04* (2006.01)
A61P 31/12 (2006.01)

(21) International Application Number:

PCT/CA2009/000787

(22) International Filing Date:

4 June 2009 (04.06.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/130,874 6 June 2008 (06.06.2008) US

(71) Applicant (for all designated States except US): **AM-BRILIA BIOPHARMA INC.** [CA/CA]; 1000 Chemin du Golf, Verdun, Quebec, H3E 1 H4 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **STRANIX, Brent** [CA/CA]; 109 Maywood, Pointe-Claire, Quebec, H9R 3L7 (CA). **BEAULIEU, Francis** [CA/CA]; 370 Leotable Dubuc, Laprairie, Quebec, J5R 5M4 (CA). **BOUCHARD, Jean-Emmanuel** [CA/CA]; 4660 De la Roche App. 4, Montreal, Quebec, H2J 3J6 (CA). **MILOT, Guy** [CA/CA]; 1972 de l'Elan, Longueuil, Quebec, J4N 1N1 (CA). **ZHIGANG, Wang** [CA/CA]; 436 Emile-Pominville Ave, Lachine, Quebec, H8R 1 M3 (CA). **RUEL, Rejean** [CA/CA]; 205 Av Macaulay, St Lambert, Quebec, J4R 2H1 (CA).

(74) Agents: **CHATTERJEE, Alakananda, Ph.D.** et al.; Gowling Lafleur Henderson LLP, P.O. Box 30, Suite 2300, 550 Burrard Street, Vancouver, British Columbia, V6C 2B5 (CA).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with amended claims (Art. 19(1))

(54) Title: HIV INTEGRASE INHIBITORS FROM PYRIDOXINE



(57) Abstract: The present invention relates to pyridoxine (vitamin B₆) derived compounds of formula (I), pharmaceutically acceptable salts, or solvates thereof, wherein R₁, R₂, R₄, A, L B₁ and B₂ are as defined in the specification, and pharmaceutical compositions comprising the compounds. Compounds of formula (I) inhibit Human Immunodeficiency Virus (HIV)-integrase enzyme and are useful for preventing and treating HIV infection and AIDS.

HIV INTEGRASE INHIBITORS FROM PYRIDOXINE

FIELD OF THE INVENTION

5 The present invention is directed to pyridoxine (vitamin B₆) derived compounds of Formula I, pharmaceutically acceptable salts or solvates thereof, pharmaceutical formulations comprising one or more compounds of Formula I, their synthesis, and use as modulators or inhibitors of human immunodeficiency virus (HIV) integrase enzyme. Compounds of the present invention are useful for prophylaxis, treatment,
10 delay in the onset or delay in the progression of human immuno-deficiency virus (HIV) infection, acquired immune deficiency syndrome AIDS, AIDS-related complex (ARC), and other diseases and conditions caused or mediated by HIV infection.

BACKGROUND OF THE INVENTION

15 Retroviruses designated as human immunodeficiency virus (HIV), particularly strains known as HIV-1 and HIV-2, are the etiological agent of AIDS, ARC, and other diseases or conditions caused or mediated by HIV. HIV infection and AIDS are difficult to treat due to the ability of retroviruses to rapidly replicate, mutate and acquire drug resistance. To date, the treatment of AIDS and HIV infection and the development of new drugs for AIDS and HIV infection have focused primarily on the
20 inhibition of HIV replication by targeting key steps in retroviral replication, such as conversion of viral RNA to viral DNA (reverse transcription) and insertion (integration) of viral DNA into the host genome. These steps rely on the activity of HIV enzymes including reverse transcriptase, protease and integrase. Various synthetic antiviral agents that block various stages of the HIV replication cycle have
25 been developed and marketed including compounds that: interfere with viral binding to CD4 (-) T-lymphocytes (for example, soluble CD4), block viral reverse transcriptase (for example, didanosine and zidovudine (AZT)), block viral aspartyl protease (for example Ritonavir and Indinavir) and inhibit viron budding (for example interferon). Some of these agents have proved ineffective in clinical tests and others,
30 primarily those that target the early stages of viral replication, have no effect on the


production of infectious virions in chronically infected cells. Furthermore,
administration of therapeutic doses of these agents has commonly led to cell-toxicity,
unwanted side effects, such as anemia, neurotoxicity and bone marrow suppression,
and rapid emergence of drug resistance which limits safe and effective treatment of
AIDS, HIV infection and other HIV-caused diseases.

The use of combination therapy has suppressed the emergence of resistance
relative to monotherapy, however even with combination therapy there is a loss of
efficacy in 30-50% of patients due to the development of viral resistance.
Considering the shortcomings of reverse transcriptase and protease inhibitors, even
when used as part of a drug cocktail (combination therapy), there is a need for new
antiviral drugs and in particular drugs that do not lead to cross-resistance with the
current standard of care.

SUMMARY OF THE INVENTION

The compounds of the present invention are useful for inhibiting or modulating
HIV integrase enzyme activity and, in particular, for inhibiting HIV replication and
for treating HIV infection, AIDS, and HIV-mediated diseases and conditions. The
present invention relates to a series of integrase inhibitors derived from pyridoxine
and pharmaceutically acceptable derivatives thereof (e.g., salts and solvates).

In one aspect the present invention are compounds of formula I,



I

wherein:

A is a six membered carbocyclic or heterocyclic ring system;

R₁ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH,
O-(C₁₋₆ alkyl), (O-C₁₋₆ branched alkyl), CO(R₉), COO(R₉), CON(R₉)(R_{9a}), or
SO₂N(R₉)(R_{9a}), wherein said R₉ and R_{9a} are selected independently from the group

consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; R₂ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl), (O-C₁₋₆ branched alkyl), CO(R₁₀), COO(R₁₀), or CON(R₁₀)(R_{10a}), wherein said R₁₀ and R_{10a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; or R₁ and R₂ are ortho substituents that together form a carbocyclic or heterocyclic ring system;

L is -N(R')C(O)-; -C(O)N(R')-; -OC(O)-; -C(O)O-; -OC(O)N(R')-; -N(R')C(O)O-; -N(R')C(O)N(R')-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-;

10 -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-;

-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-;

-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})-;

-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-;

-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-;

15 -C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-;

-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-;

-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-;

-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-;

-C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-; -Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-;

20 -C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})-; -C(R_{a1})(R_{a2})-Z-; or -Z-C(R_{a1})(R_{a2})-; wherein each R_{a1},

R_{a2}, R_{b1}, R_{b2}, R_{c1}, and R_{c2} is, independently, selected from H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, hydroxy-alkyl, benzyl, phenyl and heterocycle, or, alternatively, one or more of R_{a1} and R_{a2}; R_{b1} and R_{b2}; and R_{c1} and R_{c2} combine to form a carbocyclic

ring, and wherein Z is selected from -N(R')C(O)-; -C(O)N(R')-; -OC(O)-; -C(O)O-; -OC(O)N(R')-; -N(R')C(O)O-; -N(R')C(O)N(R')-; -N(R')-; -SO₂-; and -O-; wherein

25 R' is selected from H, C₁₋₆ alkyl, benzyl, SO₂R'', and C(O)R'', C(O)OR'', and R'' is selected from C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, heteroalkyl, carbocyclic group, benzyl, phenyl and heterocycle;

B₁ is -R₃, CH₂OR₃, CH₂N(R₈)(R_{8a}), C(O)OR₃ or C(O)N(R₈)(R_{8a}), wherein each of R₈ and R_{8a} is, independently, selected from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle;

B₂ is H or OR₅;

R_3 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; and R_5 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; or, alternatively, R_3 and R_5 combine to form a heterocyclic ring system; and

R_4 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; and
 5 pharmaceutically acceptable salts and solvates thereof.


In certain embodiments of the compounds of formula I, A is a phenyl ring, a pyridine ring, or a cyclohexyl ring.

In still other embodiments of the compounds of formula I, B_1 is H, CH_3 , CH_2OH , or CH_2OCH_3 .

10 In particular embodiments of the compounds of formula I, L is $-CH_2OCH_2-$, $-CH_2CH_2OCH_2-$, $-OCH_2-$, $-CH_2NHCH_2-$, $-C(cyclo-C_2H_4)NHCH_2-$, $-NHCH_2-$, $-CH_2CH_2NHCH_2-$, $-CH_2NHC(O)-$, $-CH_2N(CH_3)C(O)-$, $-CH(CH_2OH)NHC(O)-$, $-C(cyclo-C_2H_4)NHC(O)-$, $-CH_2CH_2NHC(O)-$, $-C(O)NH-$, $-CH_2OC(O)NH-$, $-NHC(O)NH-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2-$, $-SO_2CH_2-$, or $-CH_2SO_2CH_2-$.

15 In certain embodiments of the compounds of formula I, R_1 is selected from a halogen, $-OH$ or $-OCH_3$, R_2 is selected from $-OH$, $-H$, and a halogen, or R_1 and R_2 combine to form a cyclic acetal or cyclic ketal, R_4 is $-H$ or benzyl, B_2 is OR_5 , and R_5 is $-H$ or benzyl.

20 In certain embodiments, the compounds of formula I are further described by formula Ia:



Ia

wherein:

Y-X is $-C(R_7)(R_{7a})N(R')C(O)-$; $-C(R_7)(R_{7a})OC(O)-$; $-C(R_7)(R_{7a})N(R')C(R_6)(R_{6a})-$; or $-C(R_7)(R_{7a})OC(R_6)(R_{6a})-$, wherein each of R_6 , R_{6a} , R_7 , and R_{7a} , is, independently, selected from H, C_{1-6} alkyl, C_{1-6} fluoro-alkyl, benzyl,

25

phenyl and heterocycle, R' is selected from H, C₁₋₆ alkyl, benzyl, SO₂R'', and C(O)R'', and R'' is selected from C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle;

Q is H, CH₂, CH₃, or CO;

5 P is H, O, N(R₈)(R_{8a}), or is absent, wherein said R₈ and R_{8a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle;

R₁ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl), (O-C₁₋₆ branched alkyl), CO(R₉), COO(R₉), CON(R₉)(R_{9a}), or
10 SO₂N(R₉)(R_{9a}), wherein said R₉ and R_{9a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; R₂ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl), (O-C₁₋₆ branched alkyl), CO(R₁₀), COO(R₁₀), or CON(R₁₀)(R_{10a}), wherein said
15 R₁₀ and R_{10a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; or R₁ and R₂ are ortho substituents that together form a carbocyclic or heterocyclic ring system;

R₃ is H, C₁₋₆ alkyl, C₁₋₆ fluoroalkyl, benzyl, phenyl, heterocycle, or is absent; and R₅ is H, C₁₋₆ alkyl, C₁₋₆ fluoroalkyl, benzyl, phenyl, or heterocycle; or R₃ and R₅ combine to form a heterocyclic ring system;

20 R₄ is H, C₁₋₆ alkyl, C₁₋₆ fluoroalkyl, benzyl, phenyl, or heterocycle; and m is 0 or 1; or

pharmaceutically acceptable salts or solvates thereof.

Further provided herein are compounds of formula Ia, wherein: Q is CH₂, P is O, R₃ is H, Y-X is CH₂NHCH₂ or CH₂NHCO, R₁ is selected from a halogen, -OH or
25 -OCH₃, R₂ is selected from -OH, or R₁ and R₂ combine to form a cyclic acetal or cyclic ketal, -H, and a halogen, R₄ is -H or benzyl, and R₅ is -H or benzyl, or pharmaceutically acceptable salts or solvates thereof.

Further provided herein are compounds of formula Ia, Q is CH₂, P is O, R₃ is CH₃, Y-X is CH₂OCH₂, R₁ is selected from a halogen, -O or -OCH₃, R₂ is selected
30 from -O, -H, and a halogen, or R₁ and R₂ combine to form a cyclic acetal or cyclic ketal, R₄ is -H or benzyl and R₅ is -H or benzyl or pharmaceutically acceptable salts or solvates thereof.

Attorney _____

Further provided herein are compounds of formula I selected from N⁵-(4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(3-chloro-4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(3,4-dichlorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, and 5-(benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide.

Further provided herein are compounds of formula I selected from N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N²,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N², 3-bis(benzyloxy)-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide, N², 3-dihydroxy-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide, N², 3-bis(benzyloxy)-N⁵-(3,5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(3,5-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, 5-((4-fluorobenzylamino)methyl)-N,3-dihydroxy-4-(methoxymethyl)picolinamide, 5-((3,5-difluorobenzylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide, 5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide, N⁵-(4-fluorobenzyl)-N²,3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2,5-dicarboxamide, N²,3-bis(benzyloxy)-N⁵-(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(3-chloro-4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N²,3-bis(benzyloxy)-N⁵-(3,4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N⁵-(3,4-dichlorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N,3-dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide, 5-(benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide, N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide, and N⁵-(3,4-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide.

Further provided herein are compounds of formula I selected from N, 9-bis(benzyloxy)-3,3-dimethyl-1,5-dihydro-[1,3]dioxepino[5,6-c]pyridine-8-carboxamide, N, 3-bis(benzyloxy)-4,5-bis(hydroxymethyl) picolinamide, N, 7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide, N², 3-

5 bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, N, 3-bis(benzyloxy)-5-((4-fluorobenzylamino)methyl)-4-(hydroxymethyl)picolinamide, 5-((3,5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl) picolinamide, 5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N, 3-bis(benzyloxy)-4-(hydroxymethyl)picolinamide, 5-(benzyloxy)-N-(4-

10 fluorobenzyl)-4-(hydroxymethyl)-6-methylnicotinamide, 5-(benzyloxy)-N-(4-fluorobenzyl)-4-(methoxymethyl)-N, 6-dimethylnicotinamide, 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl)carbamoyl)-4-(methoxymethyl)-2-methylpyridine 1-oxide, 5-(benzyloxy)-N-(4-fluorobenzyl)-6-(hydroxymethyl)-4-(methoxymethyl)-N-methylnicotinamide, 5-(benzyloxy)-N-(4-fluorobenzyl)-6-formyl-4-(methoxymethyl)-

15 N-methylnicotinamide, methyl 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl)carbamoyl)-4-(methoxymethyl)picolinate, N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide, N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide, 5-((4-methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1,3]dioxino[4,5-c]pyridine, 5-((4-

20 methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1,3]dioxino[4,5-c]pyridine 7-oxide, (5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-yl)methanol, 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carbaldehyde, ethyl 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate, ethyl 3-hydroxy-4-

25 (hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinate, N, 3-dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinamide, 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid, N-(benzyloxy)-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide, N-(benzyloxy)-3-hydroxy-4-

30 (hydroxymethyl)-5-((4-methoxybenzyloxy)methyl) picolinamide, N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide.

Further provided herein are pharmaceutical compositions, and pharmaceutically acceptable formulations, comprising a therapeutically effective amount of at least one compound of the present invention, and pharmaceutically acceptable salts or solvates thereof.

5 The compounds of the present invention inhibit HIV integrase including both HIV-1 and HIV-2 and may be used as antiviral agent against HIV, including HIV-1 and HIV-2 strains.

 The compounds of the present invention are useful for prophylaxis, treatment or delay in the onset or progression of HIV infection, or of a disease or condition caused
10 or mediated by HIV infection, including HIV-1 and HIV-2 infection.

 In one aspect, the present invention features a method of inhibiting HIV replication, in a mammal, that includes administering to the mammal a replication-inhibiting amount of at least one compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof.

15 Further provided are methods of inhibiting HIV replication in a cell, comprising contacting the cell with an inhibiting amount of at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

 Further provided are methods of inhibiting HIV integrase enzyme activity, that include contacting the integrase enzyme with an integrase-inhibiting amount of at
20 least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof. The method includes contacting a cell directly or administering the compound of the invention to a mammal suffering from an HIV infection.

 Another aspect of the present invention includes methods of treating HIV
25 infection in a mammal, comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

 Further provided are methods of treating AIDS in a mammal, comprising administering to the mammal at least one compound of the present invention, or a
30 pharmaceutically acceptable salt, solvate or formulation thereof.

 Further provided are methods of treating AIDS in a mammal, comprising administering to the mammal at least one compound of the present invention, or a

pharmaceutically acceptable salt, solvate or formulation thereof in combination with one or more additional HIV-inhibiting agent.

Further provided are methods of treating a disease or condition caused or mediated by HIV infection in a mammal, comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

Further provided are methods of prophylaxis or prevention of HIV infection in a mammal, comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

Further provided are methods of inhibiting HIV replication in a mammal comprising administering to the mammal at least one compound of the present invention or a pharmaceutically acceptable salt, solvate or formulation thereof.

Further provided are methods of inhibiting HIV replication in a mammal comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof and at least one other HIV-inhibiting agent.

Further provided are methods of inhibiting HIV replication in a mammal wherein the HIV is resistant to at least one HIV protease inhibitor, the method comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

Further provided are methods of inhibiting HIV replication in a mammal, having an HIV infection, wherein the HIV is resistant to at least one HIV reverse transcriptase inhibitor, the method comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

Further provided are methods of reducing HIV viral load in a mammal infected with HIV, comprising administering to the mammal at least one compound of the present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof.

Further provided are methods of reducing HIV viral load in a mammal infected with HIV, comprising administering to the mammal at least one compound of the

present invention, or a pharmaceutically acceptable salt, solvate or formulation thereof in combination with one or more additional HIV-inhibiting agents.

Further provided is the use of at least one compound of the present invention for the manufacture of a pharmaceutical composition for treatment of HIV infection.

5 Further provided is the use of at least one compound of the present invention for the manufacture of a pharmaceutical composition for treatment of AIDS or ARC.

Further provided is the use of at least one compound of the present invention for the manufacture of a pharmaceutical composition for prevention or prophylaxis of AIDS or ARC.

10 Further provided is the use of at least one compound of the present invention for the manufacture of a pharmaceutical composition for prevention or prophylaxis of HIV infection.

For any of the above aspects of the invention, the mammal (e.g., human) may have or be suspected of having an HIV infection or an AIDS or HIV mediated disease
15 or condition. The mammal (e.g., human) may or may not have been previously treated with anti-viral or other therapeutic compounds for the HIV infection or AIDS or HIV mediated disease or condition.

Definitions

20 The terms "human immunodeficiency virus," "HIV," "HIV-1," or "HIV-2" as used herein refer to a retrovirus that is the causative agent for acquired immunodeficiency Syndrome (AIDS) and diseases, conditions or opportunistic infections associated with AIDS. Previous names for HIV include human T-lymphotropic virus-III (HTLV-III), lymphadenopathy-associated virus (LAV), and
25 AIDS-associated retrovirus (ARV).

The terms "HIV reverse transcriptase," "reverse transcriptase," or "RT" as used herein refer to an enzyme, encoded by a retroviral genome, which catalyzes or mediates the conversion (reverse transcription) of viral RNA to DNA or generation of a provirus (Haseltine W. A. FASEB J. vol. 5, p. 2349 - 2360 (1991)).

30 The terms "reverse transcriptase inhibitor" or "HIV reverse transcriptase inhibitor," as used herein, refer to compounds or combinations of compounds that

interfere with the proper functioning of the HIV reverse transcriptase enzyme that is responsible for converting single-stranded HIV viral RNA into HIV viral DNA.

The terms "HIV integrase" or "integrase" as used herein refer to an enzyme, encoded by a retroviral genome, that catalyzes or mediates integration of provirus DNA (retroviral double stranded DNA) into the host genomic DNA. The integrase enzyme can serve as a template for viral gene expression by the host transcription system, leading to viral replication (Roth et al., Cell, 1989 Jul 14; 58(1):47-54.; Bukrinsky M. I., Proc. Natn. Acad. Sci. USA 1992, vol. 89 p.6580 - 6584; Galloway et al., Cell. 1995 Nov 17;83(4):569-76).

The terms "integrase inhibitor" or "HIV integrase inhibitor," as used herein, refer to a compound or combination of compounds that interfere with the proper functioning of the HIV integrase enzyme that is responsible for inserting the genes of HIV into the DNA of a host cell.

The term "integration" as used herein refers to insertion of viral DNA, retroviral DNA, provirus, or provirus DNA into the host genome mediated by integrase enzyme. Integration generally occurs following association of integrase and viral DNA with the pre-integration complex (PIC) at the host nucleus and transport of the viral DNA into the host nucleus as a component of the pre-integration complex (Goldgur Y et al Proc Natl Acad Sci U S A. 1999 Nov 9;96(23):13040-3; Sayasith K, Sauvé G and Yelle J. Expert Opin Ther Targets. 2001 Aug;5(4):443-464; Debyser Z et al Methods Mol Biol. 2001;160:139-55).

The terms "protease inhibitor" or "HIV protease inhibitor" as used herein mean compounds or combinations of compounds that interfere with the proper functioning of the HIV protease enzyme that is responsible for cleaving long strands of viral protein into the separate proteins making up the viral core.

The terms "fusion inhibitor" or "HIV fusion inhibitor," as used herein, refer to compounds or combinations of compounds that bind to the gp41 envelope protein on the surface of CD4 cells and block the structural changes necessary for the virus to fuse with the cell.

The terms "viral load" and "HIV viral load," as used herein, mean the amount of HIV in the circulating blood of a mammal, such as a human. The amount of HIV

virus in the blood of mammal can be determined by measuring the quantity of HIV RNA in the blood using methods known to those of ordinary skill in the art.

The term “retrovirus” as used herein refers to a virus belonging to the viral family Retroviridae, which includes viruses that possess an RNA genome, and that replicate via a DNA intermediate.

The term “Vitamin B₆” as used herein refers to one or more of three compounds that are commonly referred to as vitamin B₆ namely pyridoxal, pyridoxamine and pyridoxine. Pyridoxine differs from pyridoxamine by the substituent at the '4 position. Pyridoxine based on a pyridine ring, with hydroxyl, methyl, and hydroxymethyl substituents and is converted in vivo to pyridoxal 5-phosphate, the biologically active form of pyridoxine.

The terms “comprising” and “including” are used in their open, non-limiting sense.

The term “C₁₋₆ alkyl” as used herein means saturated monovalent hydrocarbon radicals having straight or branched moieties and containing from 1 to 6 carbon atoms. The C₁₋₆ alkyl group may be substituted or unsubstituted. Examples of such groups include, but are not limited to, methyl, ethyl, propyl, iso-propyl, n-butyl, iso-butyl, and tert-butyl. The term “C₁₋₆ fluoro-alkyl” refers to a C₁₋₆ alkyl substituted with one or more fluorine atoms. Exemplary C₁₋₆ fluoro-alkyl groups include, without limitation, fluoromethyl, trifluoromethyl, and pentafluoroethyl. The term “C₁₋₆ branched alkyl” refers to alkyl group that include one or more tertiary or quaternary carbon atoms.

By “C₂₋₆ alkenyl” is meant a branched or unbranched hydrocarbon group containing one or more double bonds and having from 2 to 6 carbon atoms. A C₂₋₆ alkenyl may optionally include monocyclic or polycyclic rings, in which each ring desirably has from three to six members. The C₂₋₄ alkenyl group may be substituted or unsubstituted.

By “carbocyclic group” or “carbocyclic ring” is meant a monocyclic or polycyclic ring system which is saturated, partially unsaturated, or unsaturated (aromatic), and which consists of 3 to 8 carbon atoms (unless otherwise specified). Carbocyclic groups include alkyl groups substituted with such a monocyclic or polycyclic ring system. Exemplary cyclic groups include phenyl, benzyl,

cyclopropyl, cyclobutyl, cyclopentyl, 2-phenylcyclopropane, and cyclohexyl. The carbocyclic group may be substituted or unsubstituted.

By "heteroalkyl" is meant a branched or unbranched alkyl or alkenyl group having from 1 to 10 carbon atoms in addition to 1, 2, 3 or 4 heteroatoms independently selected from the group consisting of N, O, S, and P. Heteroalkyls include, without limitation, tertiary amines, secondary amines, ethers, thioethers, amides, thioamides, carbamates, thiocarbamates, hydrazones, imines, urethanes, phosphodiester, phosphoramidates, sulfonamides, and disulfides. A heteroalkyl may optionally include monocyclic, bicyclic, or tricyclic rings (either aromatic or non-aromatic ring systems), in which each ring desirably has three to six members. The heteroalkyl group may be substituted or unsubstituted. Examples of C₁₋₇ heteroalkyls include, without limitation, methoxymethyl, benzyloxyethyl, and ethoxyethyl.

The term "heterocycle" and "heterocyclic ring" as used herein means aromatic or non-aromatic, monocyclic, bicyclic, tricyclic, tetracyclic, or spirocyclic group having a total of from 3 to 10 atoms in its ring system, and containing from 2 to 9 carbon atoms and from one to four heteroatoms each independently selected from O, S and N and with the proviso that the ring of said group does not contain two adjacent O atoms or two adjacent S atoms. Furthermore, such heterocycle groups may contain an oxo substituent at any available atom that will result in a stable compound. For example, such a group may contain an oxo atom at an available carbon or nitrogen atom. Such a group may contain more than one oxo substituent if chemically feasible. In addition, it is to be understood that when such a heterocycle group contains a sulfur atom, said sulfur atom may be oxidized with one or two oxygen atoms to afford either a sulfoxide or sulfone. An example of a 4 membered heterocyclic group is azetidiny (derived from azetidine). An example of a 5 membered heterocyclic group is thiazolyl and an example of a 10 membered heterocyclic group is quinolinyl. Further examples of such heterocycle groups include, but are not limited to, pyrrolidinyl, tetrahydrofuranyl, dihydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, dihydropyranyl, tetrahydrothiopyranyl, piperidino, morpholino, thiomorpholino, thioxanyl, piperazinyl, azetidiny, oxetanyl, thietanyl, homopiperidinyl, oxepanyl, thiepanyl, oxazepiny, diazepiny, thiazepiny, 1,2,3,6-tetrahydropyridiny, 2-pyrroliny, 3-pyrroliny, indoliny, 2H-pyranyl, 4H-pyranyl, dioxanyl, 1,3-dioxolanyl,

pyrazoliny, dithianyl, dithiolanyl, dihydropyranyl, dihydrothienyl, dihydrofuranyl, pyrazolidinyl, imidazoliny, imidazolidinyl, 3 azabicyclo[3.1.0] hexanyl, 3-azabicyclo[4.1.0]heptanyl, 3H-indolyl and quinoliziny. The heterocycle group may be substituted or unsubstituted.

5 As used herein, the terms “benzyl” and “phenyl” refer to both substituted and unsubstituted benzyl and phenyl groups, respectively.

As used herein, the term “substituted” refers to a group (e.g., a “C₁₋₆ alkyl”, “C₂₋₆ alkenyl”, “C₁₋₆ fluoroalkyl”, “benzyl”, “phenyl”, “heterocycle”, or “carbocyclic group”) in which one or more hydrogen atoms in the group are, independently,
 10 replaced with a substituent selected from methyl, ethyl, n-propyl, isopropyl, hydroxy, methoxy, ethoxy, fluorine, chlorine, bromine, iodine, cyano, nitro, amino, alkylamino, dialkylamino, carboxy, chloromethyl, trichloromethyl, trifluoromethyl, methoxyethyl, -CH₂C(O)NH₂, -C(O)CH₂N(CH₃)₂, -CH₂CH₂OH, -CH₂OC(O)NH₂, -CH₂CH₂NH₂, -CH₂CH₂CH₂NEt₂, -CH₂OCH₃, -C(O)NH₂, -C(=NH)NH₂, -C(=NH)OEt, -C(O)NH-cyclopropyl, -C(O)NHCH₂CH₂OCH₃, -C(O)CH₂CH₂NHCH₃, -CH₂CH₂F, or
 15 -CH₂C(O)NIICH₃.

The term “inhibiting HIV replication” means reducing or preventing (e.g., by at least 10%, 20% 30%, 40%, 50%, 60%, 70%, 80%, 90% or more) human immunodeficiency virus (HIV) replication in a cell. Such a cell may be present in
 20 vitro, or it may be present in vivo, such as in a mammal, such as a human. Such inhibition may be accomplished by administering a compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, directly to the cell, or to a mammal, in an amount sufficient to inhibit HIV replication. The inhibition of HIV replication in a cell, such as in a mammal, can be measured or monitored using
 25 methods known to those of ordinary skill in the art. For example, an amount of a compound of the invention may be administered to a mammal, either alone or as part of a pharmaceutically acceptable formulation. Blood samples may then be withdrawn from the mammal and the amount of HIV virus in the sample may be quantified using methods known to those of ordinary skill in the art. A reduction in the amount of HIV
 30 virus in the sample compared to the amount found in the blood before administration of a compound of the invention would represent inhibition of the replication of HIV virus in the mammal. In another example, a reduction in the amount of HIV virus in

the sample compared to the amount found in a positive reference sample (e.g., the blood from a subject having HIV but not treated with a compound of the invention) would represent inhibition of the replication of HIV virus in the mammal. The administration of a compound of the invention to the cell, such as in a mammal, may be in the form of single dose or a series of doses. In the case of more than one dose, the doses may be administered in one day or they may be administered over more than one day.

“HIV-inhibiting agent,” “HIV antiviral agent,” or “anti-HIV agent” as used herein means a compound, including but not limited to the compounds of the present invention, or a pharmaceutically acceptable salt thereof which is capable of inhibiting the replication of HIV in a cell, such as a cell in a mammal. Such compounds may inhibit HIV replication through any mechanism known to those of ordinary skill in the art. Non-limiting examples of HIV-inhibiting agents include an entry inhibitor, a protease inhibitor, a reverse transcriptase inhibitor, a fusion inhibitor, and an integrase inhibitor.

The terms “human immunodeficiency virus-inhibiting amount” or “HIV-inhibiting amount,” as used herein, refer to the amount of an HIV-inhibiting agent, or a pharmaceutically acceptable salt or solvate thereof, required to inhibit replication of the human immunodeficiency virus (HIV) in vivo, such as in a mammal, or in vitro. The amount of such compounds required to cause such inhibition can be determined without undue experimentation using methods described herein and those known to those of ordinary skill in the art.

The term “inhibiting HIV integrase enzyme activity,” as used herein, means decreasing (e.g., by at least 10%, 20% 30%, 40%, 50%, 60%, 70%, 80%, 90% or more) the activity or functioning of the HIV integrase enzyme either in vitro or in vivo, such as in a mammal, such as a human.

The term “HIV integrase enzyme-inhibiting amount,” as used herein, refers to the amount of an HIV-inhibiting agent or a pharmaceutically acceptable salt or solvate thereof, required to decrease the activity of the HIV integrase enzyme either in vivo, such as in a mammal, or in vitro, such as in a cultured cell line. In one example, such inhibition may take place by the compound of the present invention binding directly to the HIV integrase enzyme. In addition, the activity of the HIV integrase enzyme

may be decreased in the presence of a compound of the present invention when such direct binding between the enzyme and the compound does not take place.

Furthermore, such inhibition may be competitive, non-competitive, or uncompetitive. Inhibition of HIV integrase may be determined using in vitro or in vivo systems, or a combination of both, using methods known to those of ordinary skill in the art.

The term "solvate," as used herein, means a pharmaceutically acceptable solvate form of a compound of the present invention that retains the biological effectiveness of such compound. Examples of solvates include, but are not limited to, compounds of the invention in combination with water, isopropanol, ethanol, methanol, dimethylsulfoxide (DMSO), ethyl acetate, acetic acid, ethanolamine, or mixtures thereof. In one embodiment of the present invention, one solvent molecule is associated with one molecule of the compounds of the present invention, such as a hydrate. In another embodiment of the present invention, more than one solvent molecule may be associated with one molecule of the compounds of the present invention, such as a dihydrate. Additionally, it is specifically contemplated that in the present invention less than one solvent molecule may be associated with one molecule of the compounds of the present invention, such as a hemihydrate. Furthermore, solvates of the present invention include solvates of compounds of the present invention that retain the biological effectiveness of the non-hydrate form of the compounds.

A "pharmaceutically acceptable salt" as used herein means a salt that retains the biological effectiveness of the free acids and bases of the specified derivative, containing pharmacologically acceptable anions or cations, and is not biologically or otherwise undesirable. Examples of pharmaceutically acceptable salts include, but are not limited to, acetate, acrylate, benzenesulfonate, benzoate (such as chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, and methoxybenzoate), bicarbonate, bisulfate, bisulfite, bitartrate, borate, bromide, butyne-1, 4-dioate, calcium edetate, camsylate, carbonate, chloride, caproate, caprylate, clavulanate, citrate, decanoate, dihydrochloride, dihydrogenphosphate, edetate, edisylate, estolate, esylate, ethylsuccinate, formate, fumarate, gluceptate, gluconate, glutamate, glycollate, glycolylarsanilate, heptanoate, hexyne-1,6-dioate, hexylresorcinate, hydabamine, hydrobromide, hydrochloride, γ -hydroxybutyrate, iodide, isobutyrate,

isothionate, lactate, lactobionate, laurate, malate, maleate, malonate, mandelate, mesylate, metaphosphate, methane-sulfonate, methylsulfate, monohydrogenphosphate, mucate, napsylate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, nitrate, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phenylacetates, phenylbutyrate, phenylpropionate, phthalate, phosphate/diphosphate, polygalacturonate, propanesulfonate, propionate, propiolate, pyrophosphate, pyrosulfate, salicylate, stearate, subacetate, suberate, succinate, sulfate, sulfonate, sulfite, tannate, tartrate, teoclate, tosylate, triethiodode, valerate salts, and cations, such as sodium, potassium, calcium, magnesium, ammonium, and tetraalkylammonium, among others.

The term "pharmaceutically acceptable formulation," as used herein, means a combination of a compound of the invention, or a pharmaceutically acceptable salt or solvate thereof, and a carrier, diluent, and/or excipients that are compatible with a compound of the present invention, and is not deleterious to the recipient thereof.

Pharmaceutical formulations can be prepared by procedures known to those of ordinary skill in the art. For example, the compounds of the present invention can be formulated with common excipients, diluents, or carriers, and formed into tablets, capsules, and the like. Examples of excipients, diluents, and carriers that are suitable for such formulations include the following: fillers and extenders such as starch, sugars, mannitol, and silicic derivatives; binding agents such as carboxymethyl cellulose and other cellulose derivatives, alginates, gelatin, and polyvinyl pyrrolidone; moisturizing agents such as glycerol; disintegrating agents such as povidone, sodium starch glycolate, sodium carboxymethylcellulose, agar agar, calcium carbonate, and sodium bicarbonate; agents for retarding dissolution such as paraffin; resorption accelerators such as quaternary ammonium compounds; surfactant agents such as cetyl alcohol, glycerol monostearate; adsorptive carriers such as keolin and bentonite; and lubricants such as talc, calcium and magnesium stearate and solid polyethylene glycols. Final pharmaceutical forms may be pills, tablets, powders, lozenges, sachets, cachets, or sterile packaged powders, and the like, depending on the type of excipient used. Additionally, it is specifically contemplated that pharmaceutically acceptable formulations of the present invention can contain more than one active ingredient. For example, such formulations may contain more than one compound according to

the present invention. Alternatively, such formulations may contain one or more compounds of the present invention and one or more additional anti-HIV agents. A pharmaceutically acceptable formulation may also include but is not limited to compounds, other than the compounds of formula 1, having a structure such that, upon administration to a recipient or patient, a compound of this invention, active metabolite or residue thereof is directly or indirectly provided.

By “reduce” or “inhibit” is meant the ability to cause an overall decrease of 10%, 20% 30%, 40%, 50%, 60%, 70%, 80%, 90% or more.

The term “therapeutically effective amount,” as used herein, means an amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof, that, when administered to a mammal in need of such treatment, is sufficient to effect treatment, as defined herein. Thus, a therapeutically effective amount of a compound of the present invention, or a pharmaceutically acceptable salt thereof, is a quantity sufficient to modulate or inhibit the activity of the HIV integrase enzyme such that a disease condition that is mediated by activity of the HIV integrase enzyme is reduced or alleviated.

The terms “treat,” “treating,” or “treatment” refers to both therapeutic treatment and prophylactic or preventative measures for an HIV infection or an HIV or AIDS mediated disease or condition. To “treat disease” or use for “therapeutic treatment” refers to administering treatment to a subject already suffering from a disease to improve the subject’s condition. Treatment can include modulating or inhibiting the disease or condition, (e.g., arresting its development); relieving the disease or condition, (e.g., causing regression of the disease or condition); reduction in viral load; or relieving and/or alleviating the disease or condition or the symptoms resulting from the disease or condition with or without addressing the underlying disease or condition. To “prevent disease” refers to prophylactic treatment of a subject who is not yet ill, but who is susceptible to, or otherwise at risk of, developing a particular disease. Prophylactic treatment can also include the prevention of one or more symptoms associated with HIV or AIDS. Thus, in the claims and embodiments, treating includes the administration to a mammal either for therapeutic or prophylactic purposes.

By "subject" is meant a mammal, including, but not limited to, a human or non-human mammal, such as a simian, bovine, canine, equine or feline.

5 The terms "resistant," "resistance," and "resistant HIV," as used herein, refer to HIV virus demonstrating a reduction in sensitivity to a particular drug. A mammal infected with HIV that is resistant to a particular anti-HIV agent or combination of agents usually manifests an increase in HIV viral load despite continued administration of the agent or agents. Resistance may be either genotypic, meaning that a mutation in the HIV genetic make-up has occurred, or phenotypic, meaning that resistance is discovered by successfully growing laboratory cultures of HIV virus in the presence of an anti-HIV agent or a combination of such agents.

10 The terms "co-administration," "co-administering," "co-administer," "co-administered," or "combination therapy" as used herein, refer to the administration of a combination of at least a first agent and a second agent and can include two or more agents. Such co-administration can be performed such that two or multiple agents are part of the same composition or part of the same unitary dosage form, or in separate compositions or dosage forms. Co-administration also includes administering a first agent and a second agent, or more than two agents separately and as part of the same therapeutic regimen. The agents, if administered separately, need not necessarily be administered at essentially the same time, although they can be if so desired. Thus, co-administration includes, for example, administering a first agent and a second agent as separate dosages or dosage forms, but at the same time. Co-administration also includes separate administration at different times (e.g., sequentially or alternating one agent with the other) and in any order.

20 The term "compound of the present invention" refers to compounds of formulas I and Ia, as well as those in the Examples that follow, and includes pharmaceutically acceptable salts of these compounds.

The abbreviations used herein refer to the following:

Abbreviation	Meaning
AcOH	Acetic acid
Ar	Argon
AIDS	Acquired Immunodeficiency Syndrome
AZT	3-Azido-3-deoxythymine (Zidovudine)
BSA	Bovine serum albumin
DMF	Dimethylformamide
DNA	Deoxyribonucleic acid
EtOH	Ethyl alcohol
g	gram
HPLC	High pressure liquid chromatography
HIV-1, -2	Human immunodeficiency virus type 1, type 2
HTLV-I, -II	Human T-cell lymphotropic virus type I, type II
M	Molar
MeOH	Methyl alcohol
mg	Milligram
mp	Melting point
min	Minute
mL	Milliliter
mmol	Millimole
nM	Nanomolar
RNA	Ribonucleic acid
THF	Tetrahydrofuran

Detailed Description

Pharmaceutical compositions contemplated herein comprise at least one compound of the present invention, including pharmaceutically acceptable salts, solvate or formulations thereof, with a pharmaceutically acceptable carrier, adjuvant or vehicle. Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethyleneglycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol, liposomes and lanolin.

It is understood by those skilled in the art that the compounds of the present invention, salts, or solvates thereof may exist in different crystal or polymorphic forms that are within the scope of the present invention and specified formulas.

Compounds of the present invention that are basic may be prepared as a salt
5 using suitable methods known in the art, including treatment of the free base with an inorganic acid, such as hydrochloric acid; hydrobromic acid; sulfuric acid; nitric acid; phosphoric acid; and the like, or with an organic acid, such as acetic acid; maleic acid; succinic acid; mandelic acid; fumaric acid; malonic acid; pyruvic acid; oxalic acid; glycolic acid; salicylic acid; pyranosidyl acid, such as glucuronic acid or galacturonic
10 acid; alpha- hydroxy acid, such as citric acid or tartaric acid; amino acid, such as aspartic acid or glutamic acid; aromatic acid, such as benzoic acid or cinnamic acid; sulfonic acid, such as p-toluenesulfonic acid or ethanesulfonic acid; and the like.

Basic compounds of the present invention can form a variety of salts with various inorganic and organic acids. Although such salts must be pharmaceutically
15 acceptable for administration to animals, it is common practice to first isolate the compound of the present invention as a pharmaceutically unacceptable salt and then convert to a free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention can be
20 prepared by treating the base compound with a substantially equivalent amount of the selected mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol.

Compounds of the present invention that are acidic may be prepared as a salt using suitable methods known in the art, including treatment of the free acid with an
25 inorganic or organic base, such as an amine (primary, secondary, or tertiary); an alkali metal or alkaline earth metal hydroxide; or the like. Examples of suitable salts include organic salts derived from amino acids such as glycine and arginine; ammonia; primary, secondary, and tertiary amines; and cyclic amines, such as piperidine, morpholine, and piperazine; as well as inorganic salts derived from
30 sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum, and lithium.

Acidic compounds of the present invention can form base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline- earth metal salts and particularly, the sodium and potassium salts, which can be prepared using conventional techniques. The chemical bases suitable as reagents in preparing the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of the present invention. Such non- toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc. These salts can be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.

To treat or prevent diseases or conditions caused or mediated by HIV, a pharmaceutical composition, comprising at least one of the compounds of the present invention, is administered in a pharmaceutically acceptable formulation prepared by combining a therapeutically effective amount of the compound with one or more pharmaceutically suitable carriers including diluents, excipients and auxiliaries that facilitate processing of the active compounds into a pharmaceutically acceptable formulation. Carriers employed may be either solid or liquid. Exemplary solid carriers are lactose, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the inventive compositions may include time-delay or time-release material known in the art, such as glyceryl monostearate or glyceryl distearate alone or with a wax, ethylcellulose, hydroxypropylmethylcellulose, methylmethacrylate or the like. Further additives or excipients may be added to achieve the desired formulation properties. For example, a bioavailability enhancer, such as Labrasol®, Gelucire® or the like, or formulator, such as CHIC (carboxy-methylcellulose), PG (propyleneglycol), or PEG (polyethyleneglycol), may be added.

Gelucire®, a semi-solid vehicle that protects active ingredients from light, moisture and oxidation, may be added, e.g., when preparing a capsule formulation.

5 If a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form, or formed into a troche or lozenge. The amount of solid carrier may vary, but generally will be from about 25 mg to about 1g. If a liquid carrier is used, the preparation may be in the form of syrup, emulsion, soft gelatin capsule, sterile injectable solution or suspension in an ampoule or vial or non-aqueous liquid suspension. The inventive compositions are prepared in unit-dosage form appropriate for the mode of administration, e.g., parenteral or oral
10 administration.

To obtain a stable water-soluble dose form, a pharmaceutically acceptable salt of a compound of the present invention may be dissolved in an aqueous solution of an organic or inorganic acid, such as 0.3 M solution of succinic acid or citric acid. If a soluble salt form is not available, the agent may be dissolved in a suitable co-solvent or combinations of co-solvents. Examples of suitable co-solvents include alcohol,
15 propylene glycol, polyethylene glycol 300, polysorbate 80, glycerin and the like in concentrations ranging from 0-60% of the total volume. In an exemplary embodiment, a compound of the present invention is dissolved in DMSO and diluted with water. The composition may also be in the form of a solution of a salt form of the active ingredient in an appropriate aqueous vehicle such as water or isotonic saline or dextrose solution.
20

Pharmaceutical preparations for oral use can be obtained using a solid excipient in an admixture with the active ingredient (agent), optionally grinding the resulting mixture, and processing the mixture of granules after adding suitable auxiliaries, if
25 desired, to obtain tablets or dragee cores. Suitable excipients include: fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; and cellulose preparations, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as
30 crosslinked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

The pharmaceutical compositions, comprising the compounds of the present invention may also contain suitable solid- or gel- phase carriers or excipients. These carriers and excipients may provide marked improvement in the bioavailability of poorly soluble drugs. Examples of such carriers or excipients include calcium carbonate, calcium, phosphate, sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols. Furthermore, additives or excipients such as Gelucire®, Capryol®, Labrafil®, Labrasol®, Lauroglycol®, Plurol®, Peceol®, Transcutol® and the like may be used. Further, the pharmaceutical composition may be incorporated into a skin patch for delivery of the drug directly onto the skin.

Methods of prophylaxis and treatment, their dosage levels and requirements may be selected by those of ordinary skill in the art from available methods and techniques.

The compounds of this invention may be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological system (e.g., blood, lymphatic system, central nervous system), increase oral bioavailability, increase solubility to allow administration by injection, alter metabolism or alter rate of excretion (*Pharmacokinetic Optimization in Drug Research*, Testa, B. et al, 2001, Wiley-VCH, VCHA).

The pharmaceutical compositions of this invention may be administered orally, intravenously, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally, or via an implanted reservoir and are preferably administered orally or parenterally. The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically acceptable carriers, adjuvants or vehicles. The term "parenteral" or "parenterally" as used herein includes sub-cutaneous, intra-cutaneous, intra-venous, intra-muscular, intra-articular, intra-synovial, intra-sternal, intra-theal, intra-lesional and intracranial injection or infusion techniques.

For intravenous administration, pharmaceutical compositions of the invention may be in the form of a sterile injectable preparation, for example, as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable

preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solutions. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions.

Pharmaceutical compositions of the invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, and aqueous suspension and solutions. In the case of tablets for oral and carriers which are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are administered orally, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavouring and/or coloring agents may be added.

Pharmaceutical compositions of the invention may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a compound of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active components. Such materials include, but are not limited to, cocoa butter, beeswax, and polyethylene glycols.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

It will be appreciated that the actual dosages of the agents of this invention will vary according to the particular agent being used, the particular composition

formulated, the mode of administration, and the particular site, host, and disease being treated. Those skilled in the art using: conventional dosage-determination tests in view of the experimental data for a given compound may ascertain optimal dosages for a given set of conditions. For oral administration, an exemplary daily dose
5 generally employed will be from about 0.001 to about 1000 mg/kg of body weight, with courses of treatment repeated at appropriate intervals, preferably between 0.01 and about 25 mg/kg body weight per day, and more preferably between about 0.5 and about 25 mg/kg body weight per day of the active ingredient compound useful in the prevention and treatment of viral infection, including HIV infection.

10 Furthermore, the pharmaceutically acceptable formulations of the present invention may contain a compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, in an amount of about 10 mg to about 2000 mg, or from about 10 mg to about 1500 mg, or from about 10 mg to about 1000 mg, or from about 10 mg to about 750 mg, or from about 10 mg to about 500 mg, or from about 25
15 mg to about 500 mg, or from about 50 to about 500 mg, or from about 100 mg to about 500 mg. Additionally, the pharmaceutically acceptable formulations of the present invention may contain a compound of the present invention, or a pharmaceutically acceptable salt or solvate thereof, in an amount from about 0.5 w/w % to about 95 w/w %, or from about 1 w/w % to about 95 w/w %, or from about 1
20 w/w % to about 75 w/w %, or from about 5 w/w % to about 75 w/w %, or from about 10 w/w % to about 75 w/w %, or from about 10 w/w % to about 50 w/w %.

The pharmaceutical compositions of this invention may be administered as a continuous infusion, once per day, multiple times per day (e.g., from about 1 to about 5 times per day), once per week, twice per week, three times per week, every other
25 day, every other week or as determined by the practicing clinician. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the patient treated and the particular mode of administration. A typical preparation will contain from about 5% to about 75% active
30 compound (w/w). Preferably, such preparations contain from about 20% to about 50% active compound.

Upon improvement of a patient's condition, a maintenance dose of a compound, composition or combination of this invention may be administered if necessary or desired. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained or maintained. When the symptoms have been alleviated to the desired level, treatment should cease, at least in principle. Patients may, however, require intermittent treatment on a long-term basis, upon any recurrence of disease symptoms, especially for AIDS.

As the skilled artisan will appreciate, lower or higher doses than those recited above may be required. Specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of excretion, drug combination, the severity and course of the infection, the patient's disposition to the infection and the judgment of the treating physician.

With respect to the compounds of the present invention, the particular pharmaceutical formulation, the dosage, and the number of doses given per day to a mammal requiring such treatment, are all choices within the knowledge of one of ordinary skill in the art and can be determined without undue experimentation. For example, see "Guidelines for the Use of Antiretroviral Agents in HIV-1 Infected Adults and Adolescents," United States Department of Health and Human Services, available at <http://www.aidsinfo.nih.gov/guidelines>.

The compounds of this invention are also useful as commercial reagents which effectively bind to HIV integrase. As commercial reagent, the compounds of this invention, and their derivatives, may be used to block integration of a target DNA molecule by integrase, or may be derivatized to bind to a stable resin as a tethered substrate for affinity chromatography applications. These and other uses which characterize commercial integrase inhibitors will be evident to those of ordinary skill in the art.

The compounds of the present invention can be used alone (monotherapy) or administered in combination with one or more other HIV-inhibiting agents including but not limited to additional compounds of the invention or entry inhibitors, protease

inhibitors, reverse transcriptase inhibitors, fusion inhibitors, and integrase inhibitors, examples of which are described below and known to the skilled artisan.

In one example, the compounds of the invention can be used in combination with an additional HIV integrase inhibitor. Compounds that effectively inhibit HIV integrase may provide improved antiviral agents and compositions for treating HIV infection (Wai, J.S. et al., J. Med. Chem. 43:4923-4926 (2000); Grobler, J. et al., PNAS 99: 6661- 6666 (2002); Pals, G.C.G. et al., J. Med. Chem. 45: 3184-3194 (2002); Young, S. D., Curr. Opin. Drug Disc. & Devel. 4(4): 402-410 (2001); Godwin, C.G. et al., J. Med. Chem. 45: 3184-3194 (2002); Opar, A. Nature Reviews, Drug Discovery, vol. 6, p. 258-259, (2007)). Other integrase inhibitors known in the art include those disclosed in patent applications WO200510305, WO2004039803, WO2004067531, WO2008/048538, WO2003082881 WO2007000043, and .

The compounds of this invention may be administered in combination with antiviral agents which target other steps in the retroviral replication cycle. For example, the co-administered antiviral agent can be one that targets early events in the life cycle of the virus, such as cell entry, reverse transcription and viral DNA integration into cellular DNA. Antiviral agents targeting such early life cycle events include, didanosine (ddI), zalcitabine (ddC), stavudine (d4T), zidovudine (AZT), polysulfated polysaccharides, sT4 (soluble CD4) -- which blocks attachment or adsorption of the virus to host cells -- and other compounds which block binding of virus to CD4 receptors on CD4-bearing T-lymphocytes. Other retroviral reverse transcriptase inhibitors, such as derivatives of AZT, may also be co-administered with the compounds of this invention to provide therapeutic treatment for substantially reducing or eliminating viral infectivity and the symptoms associated therewith. Examples of other antiviral agents include ganciclovir, dideoxycytidine, trisodium phosphonoformate, efloornithine, ribavirin, acyclovir, alpha interferon and trimenotrexate. Additionally, non-ribonucleoside inhibitors of reverse transcriptase, such as TIBO, nevirapine or delavirdine, may be used to potentiate the effect of the compounds of this invention, as may viral uncoating inhibitors, inhibitors of trans-activating proteins such as tat or rev, or inhibitors of the viral protease. These compounds may also be co-administered with other inhibitors of HIV integrase.

Combination therapies according to this invention may exert an additive or combined inhibitory effect on HIV replication because each therapeutic agent of the combination acts on a different site of HIV replication or a synergistic effect. For example, the use of such combination therapies also advantageously enables a reduction in the dosage of each anti-retroviral agent, compared to administration of either agent alone as a monotherapy, while providing an equivalent or better therapeutic or prophylactic effect. Administration of lower doses of each therapeutic agent often reduces or even eliminates side effects or toxicity relative to monotherapy. Furthermore, combination therapies reduce the potential for the development of viral resistance to the agents administered compared to monotherapy.

Preferred combination therapies include the administration of a compound of this invention with AZT, 3TC, ddI, ddC, d4T, combivir, ziagen, sustiva, nevirapine and delavirdine. The compounds of this invention may also be co-administered with other HIV protease inhibitors such as saquinavir, indinavir, nelfinavir, ritonavir and amprenavir. Combination of the compounds of this invention with such protease inhibitors may increase the therapeutic or prophylactic against various HIV viral mutants, HIV quasi species or other closely related viruses.

The compounds of this invention may be administered in combination with nucleoside or non-nucleoside retroviral reverse transcriptase inhibitors (e.g. derivatives of AZT or HIV aspartyl protease inhibitors) HIV-entry inhibitors, HIV integrase inhibitors, immuno-modulators (e.g., broprimine, anti-human alpha interferon antibody, IL-2, GM-CSF, methionine enkephalin, interferon alpha, diethyldithiocarbante, tumor necrosis factor, naltrexone and rEPO); antibiotics (e.g., pentamidine isethionate), vaccines or a combination thereof.

Administration of the compounds of this invention in combination therapies with other agents to patients may be sequential or concurrent. Furthermore, pharmaceutical or prophylactic compositions of this invention may include a combination of an integrase inhibitor compound of the present invention and another therapeutic or prophylactic agent or HIV-inhibiting agent. Additional examples of agents useful for treating AIDS and HIV and suitable for combination therapies with the compounds of this invention are listed in Tables 1 and 2 below.

Table 1

Antiviral Drug	Manufacturer	Indication
097 (non-nucleoside reverse transcriptase inhibitor)	Hoechst/Bayer	HIV infection, AIDS, ARC
Amprenavir 141 W94 GW 141 (protease inhibitor)	Glaxo Wellcome	HIV infection, AIDS, ARC
GW 1592 (RT inhibitor)	Glaxo Wellcome	HIV infection, AIDS, ARC
Acemannan	Carrington Labs (Irving, TX)	ARC
Acyclovir	Burroughs Wellcome	HIV infection, AIDS, ARC, in combination with AZT
AD-439	Tanox Biosystems	HIV infection, AIDS, ARC
AD-519	Tanox Biosystems	HIV infection, AIDS, ARC
Adefovir dipivoxil AL-721	Gilead Sciences Ethigen (Los Angeles, CA)	HIV infection, ARC, PGL HIV positive, AIDS
Alpha Interferon HIV in combination w/Retrovir	Glaxo Wellcome	Kaposi's sarcoma
Ansamycin LM 427	Adria Laboratories (Dublin, OH) Erbamont (Stamford, CT)	ARC
Antibody which Neutralizes pH Labile alpha aberrant Interferon	Advanced Biotherapy Concepts (Rockville, MD)	AIDS, ARC
AR177	Aronex Pharm	HIV infection, AIDS, ARC
Beta-fluoro-ddA	Nat'l Cancer Institute	AIDS-associated diseases
BMS-232623 (CGP-73547) (protease inhibitor)	Bristol-Myers Squibb/Novartis	HIV infection, AIDS, ARC
BMS-234475 (CGP-61755) (protease inhibitor)	Bristol-Myers Squibb/Novartis	HIV infection, AIDS, ARC
CI-1012	Warner-Lambert	HIV-1 infection
Cidofovir	Gilead Science	CMV retinitis, herpes, papillomavirus
Curdan sulfate	AJI Pharma USA	HIV infection
Cytomegalovirus Immune globulin	MedImmune	CMV retinitis
Cytovene Ganciclovir	Syntex	Sight threatening CMV peripheral, CMV retinitis
Delaviridine (RT inhibitor)	Pharmacia-Upjohn	HIV infection, AIDS, ARC
Dextran Sulfate	Ueno Fine Chem. Ind. Ltd. (Osaka, Japan)	AIDS, ARC, HIV positive asymptomatic
ddCDideoxycytidine	Hoffman-La Roche	HIV infection, AIDS, ARC
ddI Dideoxyinosine	Bristol-Myers Squibb	HIV infection, AIDS, ARC; combination with AZT/d4T
DMP-450 (protease inhibitor)	AVID (Camden, NJ)	HIV infection, AIDS, ARC
Efavirenz (DMP 266)	DuPont Merck	HIV infection, AIDS, ARC
EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
Famciclovir	Smith Kline	herpes zoster, herpes simplex
FTC (reverse transcriptase inhibitor)	Emory University	HIV infection, AIDS, ARC
GS 840 (reverse transcriptase inhibitor)	Gilead	HIV infection, AIDS, ARC
HBV097 (non-nucleoside RT inhibitor)	Hoechst Marion Roussel	HIV infection, AIDS, ARC

Hypericin	VIMRx Pharm.	IHIV infection, AIDS, ARC
Recombinant Human Interferon Beta	Triton Biosciences (Alameda, CA)	AIDS, Kaposi's sarcoma, ARC
Interferon alfa-n3	Interferon Sciences	ARC, AIDS
Indinavir	Merck	HIV infection, AIDS, ARC, asymptomatic HIV positive, also in combination with AZT/ddI/ddC
ISIS 2922	ISIS Pharmaceuticals	CMV retinitis
KNI-272	Nat'l Cancer Institute	HIV-associated diseases
Lamivudine, 3TC (reverse transcriptase inhibitor)	Glaxo Wellcome	HIV infection, AIDS, ARC, also with AZT
Lobucavir	Bristol-Myers Squibb	CMV infection
Nelfinavir (protease inhibitor)	Agouron Pharmaceuticals	HIV infection, AIDS, ARC
Nevirapine (RT inhibitor)	Boehringer Ingelheim	HIV infection, AIDS, ARC
Novapren	Novaferon Labs, Inc. (Akron, OH)	HIV inhibitor
Peptide T Octapeptide Sequence	Peninsula Labs (Belmont, CA)	AIDS
Trisodium Phosphonoformate	Astra Pharm. Products, Inc.	CMV retinitis, HIV infection, other CMV infections
PNU-140690 (protease inhibitor)	Pharmacia Upjohn	HIV infection, AIDS, ARC
Probucol	Vyrex	HIV infection, AIDS
RBC-CD4	Sheffield Med. Tech (Houston, TX)	HIV infection, AIDS, ARC
Ritonavir (protease inhibitor)	Abbott	HIV infection, AIDS, ARC
Saquinavir (protease inhibitor)	Hoffmann-LaRoche	HIV infection, AIDS, ARC
Stavudine; d4T	Bristol-Myers Squibb	HIV infection, AIDS, ARC
Didehydrodeoxy-thymidine		
Valaciclovir	Glaxo Wellcome	Genital HSV & CMV infections
Virazole Ribavirin	Viratek/ICN (Costa Mesa, CA)	asymptomatic HIV-positive, LAS, ARC
VX-478	Vertex	HIV infection, AIDS, ARC
Zalcitabine	Hoffmann-LaRoche	HIV infection, AIDS, ARC, combination with AZT
Zidovudine; AZT	Glaxo Wellcome	HIV infection, AIDS, ARC, Kaposi's sarcoma, in combination with other therapies
Tenofovir disoproxil, fumarate salt (Viread®) (RT inhibitor)	Gilead	HIV infection, AIDS
Combivir® (RT inhibitor)	GSK	HIV infection, AIDS
abacavir succinate (or Ziagen®) (RT inhibitor)	GSK	HIV infection, AIDS
Reyataz® (atazanavir)	Bristol-Myers Squibb	HIV infection, AIDS
Fuzeon (Enfuvirtide, T-20)	Roche/Trimeris	HIV infection, AIDS, viral fusion inhibitor
Trizivir®	GSK	HIV infection, AIDS
Kaletra®	Abbott	HIV infection, AIDS, ARC

Table 2

Immuno-Modulator	Manufacturer	Indication
Acemannan	Carrington Labs Inc. (Irving, TX)	AIDS, ARC
Alpha-2 Interferon	Schering Plough	Kaposi's sarcoma w/AZT, AIDS
AS-101 Bropirimine	Wyeth-Ayerst/Pharmacia Upjohn	AIDS, advanced AIDS
CL246,738	American Cyanamid Lederle Labs	AIDS, Kaposi's sarcoma
EL10	Elan Corp PLC (Gainesville, GA)	HIV infection
FP-21399	Fuki ImmunoPharm	Blocks HIV fusion with CD4+ cells
Gamma Interferon	Genentech	ARC, in combination w/TNF (tumor necrosis factor)
Granulocyte Macrophage Colony Stimulating Factor	Genetics Institute/Sandoz	AIDS
Granulocyte Macrophage Colony Stimulating Factor	Hoechst-Roussel/Immunex	AIDS
Granulocyte Macrophage Colony Stimulating Factor	Schering-Plough	AIDS, combination w/AZT
HIV Core Particle Immunostimulant	Rorer	Seropositive HIV
IL-2 Interleukin-2	Cetus	AIDS, in combination w/AZT
IL-2 Interleukin-2	Hoffman-LaRoche/Immunex	AIDS, ARC, HIV, in combination w/AZT
IL-2 Interleukin-2 (aldeslukin)	Chiron	AIDS, increase in CD4 cell counts
IMERG-2	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
Immune Globulin Intravenous (human)	Cutter Biological (Berkeley, CA)	Pediatric AIDS, in combination w/AZT
IMREG-1	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
Imuthiol Diethyl Dithio Carbamate	Merieux Institute	AIDS, ARC
Interferon Alfa 2a	Hoffman-La Roche	Kaposi's sarcoma, AIDS, in combination w/AZT ARC
Methionine-Enkephalin	TNI Pharmaceutical (Chicago, IL)	AIDS, ARC
MTP-PE Muramyl-Tripeptide		
Granulocyte Colony Stimulating Factor	Ciba-Geigy Corp./Amgen	Kaposi's sarcoma AIDS in combination w/AZT
rCD4 Soluble Human CD4		AIDS, ARC Recombinant AIDS, ARC
rCD4-IgG	Genentech	hybrids
Recombinant Soluble Human CD4	Biogen	AIDS, ARC
Remune	Immune Response Corp.	Immunotherapeutic

Anti-infectives that may be used in combination with the compounds of the present invention include, but are not limited to, atovaquone, azithromycin, clarithromycin, trimethoprim, trovafloxacin, pyrimethamine, daunorubicin,

clindamycin with primaquine, fluconazole, pastill, ornidyl, eflornithine pentamidine, rifabutin, spiramycin, itraconazole-R51211, trimetrexate, daunorubicin, recombinant human erythropoietin, recombinant human growth hormone, megestrol acetate, testosterone, and total enteral nutrition.

5 Antifungals that may be used in combination with the compounds of the present invention include, but are not limited to, anidulafungin, C31G, caspofungin, DB-289, fluconazole, itraconazole, ketoconazole, micafungin, posaconazole, and voriconazole. [0454] Other compounds that may be used in combination with the compounds of the present invention include, but are not limited to, acmannan, ansamycin, LM 427,
10 AR177, BMS-232623, BMS-234475, CI-1012, curdlan sulfate, dextran sulfate, STOCRINE EL10, hypericin, lobucavir, novapren, peptide T octapeptide sequence, trisodium phosphonoformate, probucol, and RBC-CD4.

 In addition, the compounds of the present invention may be used in combination with anti-proliferative agents for the treatment of conditions such as Kaposi's
15 sarcoma. Such agents include, but are not limited to, inhibitors of metallo-matrix proteases, A-007, bevacizumab, BMS-275291, halofuginone, interleukin-12, rituximab, paclitaxel, porfimer sodium, rebimastat, and COL-3.


 Compounds of the present invention may be administered in combination with an additional agent or pharmaceutical composition that increases the bioavailability or slows the metabolism of the compounds. Agents or pharmaceutical compositions that
20 may increase the bioavailability or slow the metabolism of the compounds herein include inhibitors of at least one isoform of the cytochrome P450 (CYP450) enzymes, preferably CYP1A2, CYP2D6, CYP2C9, CYP2C19 and CYP3A4. Suitable agents that may be used to inhibit CYP 3A4 include, but are not limited to, delavirdine and
25 ritonavir. Such combinations may be administered such that a compound or compounds of the present invention are present in a single formulation or in the form of separate formulations that may be administered sequentially with an appropriate period of time in between or simultaneously. The choice of whether to include the compound or compounds of the present invention in the same formulation as the
30 additional agent or agents is within the knowledge of one of ordinary skill in the art.

Preparation of Intermediates and Compounds

Fifteen general approaches (synthetic schemes) were used to prepare the compounds of the present invention.

The first approach (Scheme 1) starts from pyridoxine which is modified to produce Intermediate I using methodologies described in Paul et al. J. Med. Chem., 1977, 20 p745,. Intermediate I is modified to produce a protected hydroxamic acid (II) by ester displacement and the isopropylidene protecting group of II is removed by mild hydrolysis with formic acid to give intermediate III. Selective oxidation of the 5-CH₂OH group of III is effected using Manganese dioxide and III spontaneously cyclizes to the corresponding lactone intermediate (IV). IV may be substituted, as illustrated in Scheme 1, with an amine producing the corresponding amide V. Protecting groups are removed using hydrogenolysis giving the desired product VI.


Scheme 1



The second approach (Scheme 2 below) starts with intermediate VIII. VIII is obtained from Intermediate I (Scheme 1) using steps 2 and 3 of Scheme 1; removal of the isopropylidene protecting group by hydrolysis (as step 2 of scheme 1) followed by selective oxidation of 5-CH₂OH with MnO₂ (as step 3 of scheme 1). VIII is then hydrolysed using potassium trimethylsilanoate to generate intermediate VII. VII is transformed to intermediate IV using O-protected hydroxylamine and aryl sulfonyl halide. VI is substituted with an amine producing the corresponding amide V and the


protecting groups are removed by hydrogenolysis (VI). Further hydrogenation yields the desired product IX.

Scheme 2



5 The third approach (Scheme 3) starts with the protected pyro as in scheme I and generates the lactol IV by controlled oxidation with MnO_2 . The lactol (IV) is readily converted to the amine X through reductive amination and further reduction by catalytic hydrogenation generating compound XI.

Scheme 3




10

15

The fourth approach (Scheme 4) starts from pyridoxine and generates intermediate XIII in a manner similar to that described in Adamczyk M. *et al*, Tetrahedron, 56, 2000 p 2379. XIII is then oxidized selectively at the 2 methyl group through an N-oxide intermediate XIV followed by rearrangement to the alcohol XV. Further

stepwise oxidation yields an aldehyde (XVI) followed by an ester (XVII). Hydrolysis of the isopropylidene of XVII and displacement of the ester XVIII with hydroxylamine yields the desired product compound XIX.


Scheme 4




5

10

The fifth approach (Scheme 5), for synthesis of integrase inhibitor compounds of invention, starts from the intermediate XVII of scheme 4. XVII is hydrolysed to the acid XX and XX is subsequently coupled to O-benzylhydroxylamine to give XXI. XXI may be subjected to hydrogenation yielding the product compound XXIII or to hydrolysis yielding the product compound XXII.


Scheme 5

The sixth approach (Scheme 6) begins from 7-(benzyloxy)-6-methylfuro[3,4-c]pyridin-3(1H)-one; intermediate XXIV (Paul, B., Korytnyk, W. J. *Het. Chem.*, 1976, v13, p701). XXIV is reacted with amine yielding intermediate XXV. XXV is then alkylated to XXVI and oxidized selectively at the 2 position through an N-oxide rearrangement, as described in Scheme 4, to yield intermediate XXX. XXX is then displaced and hydrogenated to yielding the desired product compound XXXII.

Scheme 6


The seventh approach begins from an analogue of compound XVII of scheme 5, XVIIe (17e below), which is acidified in anhydrous conditions to yield product XXXIII. This product is oxidized to give an intermediate aldehyde XXXIV and further oxidized, under controlled conditions, to give the carboxylic acid intermediate XXXV.

Scheme 7



The eighth approach consists of reductive amination of the intermediate XXXIV with either a primary or secondary amine to obtain XXXVI or XXXVII respectively. XXXVII analogues may be obtained by another reductive amination of aldehydes with XXXVI. Treatment of XXXVI and XXXVII with aqueous acids yield the intermediate XXXVIII which can be further reacted with hydroxylamine solutions to give product XXXIX.


Scheme 8



The ninth approach consists of controlled oxidation of intermediate XXXIV producing carboxylic acid XXXV. A Curtius rearrangement using diphenylphosphorylazide (DPPA) in the presence of benzyl alcohol affords the


intermediate XL. XL may be deprotected by hydrogenation to yield the amine XLI which may be reacted with activated carboxylic acids, acyl halides, isocyanates, chloroformated and other electrophiles to yield intermediate XLII. These may then be converted to product XLIII by exposure to formic acid, followed by reaction with hydroxylamine.

Scheme 9




The tenth approach begins with intermediate XXXIII (scheme 7) which is reacted in a Mitsunobu reaction with phenols to yield ether XLIV. XLIV is exposed to aqueous formic acid, and subsequently reacted with hydroxylamine to yield product XLV.


Scheme 10




The eleventh approach consists of acylation of compound XXXVI by activated carboxylic acids, acyl halides, chloroformates isocyanates and other electrophiles to obtain compound XLVI. XLVI is then exposed to aqueous formic acid and reacted with hydroxylamine to yield product XLVII.

Scheme 11


The twelfth approach begins with a Wittig reaction of intermediate XXXIV to yield alkenes of the form XLVII. XLVII is then hydrogenated producing saturated alkanes which are exposed to aqueous formic acid and subsequently reacted with hydroxylamine to yield product L.

Scheme 12


The thirteenth approach begins with intermediate XXXIII and reaction of XXXIII with methane sulfonyl chloride to yield the reactive intermediate alkyl chloride LI. LI is immediately and cautiously reacted with a mercaptan to yield the thioether LII. Reaction of LII with excess peroxide mCPBA yields a sulfone-N-Oxide intermediate LIII, which upon exposure to trifluoroacetic anhydride rearranges to products LIV. The intermediate alcohol may be oxidized in a stepwise manner, to LV esters, which are then exposed to aqueous formic acid and reacted with hydroxylamine to yield products LVI.

Scheme 13

The fourteenth approach consists of first deprotecting intermediate XXIV through catalytic hydrogenation to yield LVII. LVII is then alkylated with trimethylsilyl diazomethane to the methoxy derivative LVIII. Reaction of the lactone with a primary amine followed by protection of the liberated alcohol with tert-butyl diphenylsilyl chloride gives intermediates LIX. Reaction of LIX with peroxide, mCPBA, yields intermediates LX. LX can be rearranged with trifluoroacetic anhydride yielding LXI and oxidized in stepwise fashion to LXII. A protected hydroxylamide is formed from by first reacting O-benzylhydroxylamine hydrochloride with several equivalents of lithium hexamethyldisilazane and reacting this with intermediate LXII, quenched and exposed to fluoride ions, yielding LXIII. Catalytic hydrogenation of LXIII yields product LXIV.

Scheme 14

The fifteenth approach begins by exposing the compound XVIIe (of example 7) to aqueous formic acid to yield LXV. The liberated hydroxyl groups are then acylated with acetic anhydride to the acetyl esters and modified by catalytic hydrogenation to the methyl derivative LXVI in good yields. The PMB protecting group is removed by exposure to TFA and the liberated hydroxyl group oxidized in a stepwise manner to the aldehyde LXVII and carboxylic acid LXIX by reacting initially with manganese oxide then sodium chlorite respectively.

Scheme 15

It can be appreciated by those skilled in the art the above synthetic schemes are not intended to be a comprehensive list of all means by which the compounds above may be synthesized. Further methods will be evident to those of ordinary skill in the art.


5 **General Procedures**

Preparative chromatography was performed by flash chromatography, using silica gel 60 (EM Science) with the indicated solvent systems and positive air pressure, to allow for a proper rate of elution, or with a Biotage SP4™ automated chromatography system. Detection of the compounds was carried out by exposing
10 eluted plates (analytical or preparative) to iodine, UV light and/or treating analytical plates with a 2% solution of *p*-anisaldehyde in ethanol containing 3% sulfuric acid and 1% acetic acid, followed by heating. Alternatively, analytical plates can be treated with a 0.3% ninhydrin solution in ethanol containing 3% acetic acid and/or a CAM solution made of 20 g (NH₄)₆Mo₇O₂₄ and 8.3 g Ce(SO₄)₂ polyhydrate in water
15 (750 mL) containing concentrated sulfuric acid (90 mL).

Unless otherwise indicated: all starting materials were purchased from a commercial source such as Aldrich Co. or Sigma Co; melting points (mp) were determined on a Büchi 530 melting point apparatus in capillary tubes (uncorrected); mass spectra were recorded on a Hewlett Packard LC/MSD 1100 system APCI either
20 in negative mode or positive mode; nuclear magnetic resonance (NMR) spectra were recorded on a Bruker AMX 400 equipped with a reversed or QNP probe.

Samples were dissolved in deuterio-chloroform (CDCl₃), deuterium oxide (D₂O) or deuterio-dimethylsulfoxide (DMSO-*d*₆) for data acquisition and tetramethylsilane was used as internal standard. Chemical shifts (δ) are expressed in parts per million
25 (ppm), coupling constants (*J*) are expressed in hertz (Hz) and multiplicities are denoted as s for singlet, d for doublet, dd for doublet of doublets, t for triplet, q for quartet, quint for quintet, m for multiplet, and br s for broad singlet.

EXAMPLES

Example 1: Preparation of N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (Product 1)

5 **Step 1a: Preparation of N, 9-bis(benzyloxy)-3,3-dimethyl-1,5-dihydro-[1,3]dioxepino[5,6-c]pyridine-8-carboxamide (compound 1a)**

Lithium bis(trimethylsilyl)amide (30.0 mL, 30.0 mmol, 3.75 eq) was added to benzylhydroxylamine hydrochloride (1.41 g, 8.82 mmol, 1.1 eq) in tetrahydrofuran (40.0 mL) at -78 °C. The reaction mixture was stirred 10 min. and a solution of methyl 9-(benzyloxy)-3,3-dimethyl-1,5-dihydro-[1,3]dioxepino[5,6-c]pyridine-8-carboxylate, J. Med Chem, 1977, 20 ,p745) (2.75 g, 8.01 mmol, 1 eq) in tetrahydrofuran (20.0 mL) was added. This reaction mixture was stirred at -78 °C for 30 min. and a solution of saturated ammonium chloride was added. The reaction mixture was then extracted with ethyl acetate and the organic phase recovered was concentrated under reduced pressure yielding 3.48 g of crude **compound 1a** (100% yield) as a white solid; MS-ESI *m/z* 435 [MH]⁺.

15 **Step 1b: Preparation of N, 3-bis(benzyloxy)-4,5-bis(hydroxymethyl)picolinamide (compound 1b)**

Compound 1a was dissolved in water (50.0 mL), formic acid (5.0 mL) and tetrahydrofuran (25.0 mL) and stirred at 70 °C for 2 hours. Saturated aqueous sodium bicarbonate solution was added and the reaction mixture was extracted with ethyl acetate. The organic phase recovered was dried over magnesium sulfate and concentrated under reduced pressure yielding 3.44 g of crude **compound 1b** (100% yield) as a yellow oil; MS-ESI *m/z* 395 [MH]⁺.


25 **Step 1c: Preparation of N, 7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (compound 1c)**

Activated manganese dioxide (7 g, 80 mmol, 3 eq) and 3.4 g of **compound 1b** (80.0 mL) were stirred at room temperature in dichloromethane until the reaction was complete, as indicated by TLC. The reaction mixture was then filtered on Celite and concentrated under pressure. The crude product obtained was purified by chromatography on silica gel (30% ethyl acetate/hexane) yielding 0.4 g of **compound 1c** (13% yield) as a white solid and 1.8 g (58% yield) of the corresponding lactol, N,7-bis(benzyloxy)-3-hydroxy-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (**compound 1c-2**); ¹H-NMR (400MHz, MeOD) : δ = 8.74 (s, 1H), 7.46 (m, 10H), 5.52 (s, 2H), 5.28 (s, 2H), 4.98 (s, 2H); MS-ESI *m/z* 391 [MH]⁺.

Step 1d: Preparation of N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (compound 1d)


4-Fluorobenzylamine (0.160 g, 1.29 mmol, 2 eq) and **compound 1c** (0.250 g, 0.641 mmol, 1 eq) were heated neat at 70 °C for 30 min. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.275 g of N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide as a white solid; MS-ESI *m/z* 516 [MH]⁺.

Example 2: Preparation of N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (Product 2)




The product of example 1, N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide, (0.170 g, 0.330 mmol, 1 eq) and 10% Pd/C catalyst (5 mg) were stirred in 4.0 mL of methanol under an atmosphere of hydrogen for 1 hour. The catalyst was filtered and the reaction mixture was concentrated under vacuum yielding 0.100 g of N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (90% yield) as a white solid; ¹H-NMR (400MHz, MeOD) : δ = 8.17 (s, 1H), 7.44 (m, 2H), 7.09 (t, 2H), 4.78 (s, 2H), 4.58 (s, 2H); MS-ESI *m/z* 336 [MH]⁺.

Example 3: Preparation of N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N2,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (Product 3)




Piperonylamine (0.035 g, 0.288 mmol, 2 eq) and N,7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (0.056 g, 0.144 mmol, 1 eq), compound 1c of example 1, were heated neat at 70 °C for 30 min. The crude product was purified by silica gel (30% ethyl acetate) yielding 0.048 g of N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N2,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (62% yield) as a white solid; MS-ESI m/z 542 [MH]⁺.

Example 4: Preparation of N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N2,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (Product 4)




The final product of Example 3 above (0.170 g, 0.330 mmol, 1 eq) and 10% Pd/C (5 mg) catalyst were mixed in methanol (4.0 mL) under an atmosphere of hydrogen 1 hour. The catalyst was filtered and reaction mixture was concentrated under vacuum yielding 0.029 g of N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N2,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (90% yield) as a white solid; ¹H-NMR (400MHz, MeOD) : δ = 8.176 (s, 1H), 6.93 (m, 3H), 5.95 (s, 2H), 4.77 (s, 2H), 4.50 (s, 2H); MS-ESI m/z 362 [MH]⁺.

Example 5: Preparation of N², 3-bis(benzyloxy)-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide (Product 5)




4-Methoxyaniline (0.018 g, 0.128 mmol, 2.5 eq) and N,7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (0.020 g, 0.051 mmol, 1 eq) were heated neat at 70 °C for 30 min. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.011 g of N²,3-bis(benzyloxy)-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide (41% yield) as a white solid; MS-ESI *m/z* 528 [MH]⁺.

Example 6: Preparation of N², 3-dihydroxy-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide (Product 6)




The product of example 5 (0.010 g, 0.019 mmol, 1 eq) and 10% Pd/C catalyst (5 mg) were stirred in 4.0 mL of methanol under an atmosphere of hydrogen for 1 hour. The catalyst was filtered and reaction mixture was concentrated under vacuum yielding 0.006 g of N²,3-dihydroxy-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide (85% yield) as a white solid; MS-ESI *m/z* 348 [MH]⁺.

Example 7: Preparation of N², 3-bis(benzyloxy)-N⁵-(3,5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (Product 7)




3,5-difluorobenzylamine (0.02 g, 0.154 mmol, 2.5 eq) and N,7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (0.030 g, 0.077 mmol, 1 eq) were heated neat at 70 °C for 30 min. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.011 g of N²,3-bis(benzyloxy)-N⁵-(3,5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (15% yield) as a white solid; MS-ESI *m/z* 534 [MH]⁺.

Example 8: Preparation of N⁵-(3, 5-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (Product 8)



The product of example 7 (0.006 g, 0.011 mmol, 1 eq) and 10% Pd/C catalyst (5 mg) were mixed in 4.0 mL of methanol under an atmosphere of hydrogen 1 hour. The catalyst was filtered and the reaction mixture was concentrated under vacuum yielding 0.004 g of N⁵-(3, 5-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (67% yield) as a white solid; MS-ESI *m/z* 354 [MH]⁺.

Example 9: Preparation of 5-((4-fluorobenzylamino)methyl)-N, 3-dihydroxy-4-(methoxymethyl)picolinamide (Product 9)




Step 9a: preparation of N, 3-bis(benzyloxy)-5-((4-fluorobenzylamino)methyl)-4-(hydroxymethyl)picolinamide (compound 9a)

4-fluorobenzylamine (0.035 g, 0.281 mmol, 1.1 eq), followed by sodium cyanoborohydride (0.048 g, 0.765 mmol, 3 eq) were added to N,7-bis(benzyloxy)-3-hydroxy-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (0.100 g, 0.255 mmol, 1 eq), **compound 1c-2** of example 1, in 10.0 mL of ethanol. This reaction mixture was stirred at room temperature overnight. Saturated aqueous sodium bicarbonate solution was added and the reaction mixture was extracted with ethyl acetate. Organic phase recovered was dried over magnesium sulfate and concentrated under reduced pressure. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.050 g of **compound 9a** (39% yield) as a white solid; MS-ESI m/z 502 $[MH]^+$.

Step 9b: Preparation of 5-((4-fluorobenzylamino)methyl)-N, 3-dihydroxy-4-(methoxymethyl)picolinamide

Compound 9a (0.013 g, 0.019 mmol, 1 eq) and 10% Pd/C catalyst (5 mg) were stirred in 4 mL of methanol under an atmosphere of hydrogen for 12 hrs. The catalyst was filtered and reaction mixture was concentrated under vacuum yielding 0.006 g of 5-((4-fluorobenzylamino)methyl)-N,3-dihydroxy-4-(methoxymethyl)picolinamide (75% yield) as a white solid; MS-ESI m/z 322 $[MH]^+$.

Example 10: Preparation of 5-((3, 5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide




Step 10a: preparation of 5-((3,5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl) picolinamide (compound 10a)

N,7-bis(benzyloxy)-3-hydroxy-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (**compound 1c-2** of example 1) (0.200 g, 0.510 mmol, 1 eq) in 10.0 mL of ethanol was added to 3,5-difluorobenzylamine (0.080 g, 0.561 mmol, 1.1 eq), followed by sodium cyanoborohydride (0.096 g, 1.53 mmol, 3 eq). The reaction mixture was stirred at room temperature overnight. 1M potassium carbonate solution was added and the reaction mixture was extracted with ethyl acetate. Organic phase recovered was dried over magnesium sulfate and concentrated under reduced pressure. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.046 g of **compound 10a** (17% yield) as a white solid; MS-ESI m/z 520 [MH]⁺.

Step 10b: preparation of 5-((3, 5-difluorobenzylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl) picolinamide

Compound 10a (0.020 g, 0.039 mmol, 1 eq) and 10% Pd/C (5 mg) in 4.0 mL of methanol were stirred under an atmosphere of hydrogen overnight. The catalyst was filtered and reaction mixture was concentrated under vacuum to give 0.010 g of 5-((3,5-difluorobenzylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide (76% yield) as a white solid; MS-ESI m/z 340 [MH]⁺.

Example 11: Preparation of 5-((benzo[d][1, 3]dioxol-5-ylmethylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide (Product 11)



Step 11a: preparation of 5-((benzo[d][1, 3]dioxol-5-ylmethylamino)methyl)-N, 3-bis(benzyloxy)-4-(hydroxymethyl)picolinamide (compound 11a)


N, 7-bis(benzyloxy)-3-hydroxy-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (**compound 1c-2** of example 1) (0.150 g, 0.383 mmol, 1 eq) was added to 10.0 mL of ethanol followed by 0.072 g of sodium cyanoborohydride (1.15 mmol, 3 eq). The reaction mixture was stirred at room temperature overnight. A 1M potassium

carbonate solution was added to the reaction mixture followed by extraction with ethyl acetate. Organic phase recovered was dried over magnesium sulfate and concentrated under reduced pressure. The crude product was purified by silica gel chromatography (100% ethyl acetate) yielding 0.046 g of **compound 11a** (22% yield) as a white solid; MS-ESI m/z 528 $[MH]^+$.

Step 11b: preparation of 5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide

Compound 11a (0.024 g, 0.039 mmol, 1 eq) and 10% Pd/C catalyst (5 mg) were stirred in 4.0 mL of methanol overnight under an atmosphere of hydrogen. The catalyst was filtered and the reaction mixture was concentrated under vacuum yielding 0.011 g of 5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide (69% yield) as a white solid; MS-ESI m/z 348 $[MH]^+$.

Example 12: Preparation of N⁵-(4-fluorobenzyl)-N²,3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2,5-dicarboxamide (Product 12)



Step 12a: preparation of 5-(benzyloxy)-N-(4-fluorobenzyl)-4-(hydroxymethyl)-6-methylnicotinamide (compound 12a)

Benzyloxy)-6-methylfuro[3,4-c]pyridin-3(1H)-one (Paul, B., Korytnyk, W. J. Het. Chem., 1976, v13, p701) (1.00g, 3.92 mmol, 1 eq) and 4-fluorobenzylamine (0.491 g, 3.92 mmol, 1.1 eq) were heated neat at 70 °C for 1 hour. The crude product was purified by silica gel (100% ethyl acetate) yielding 1.42 g of **compound 12a** (95% yield) as a white solid; MS-ESI m/z 381 $[MH]^+$.

Step 12b: preparation of 5-(benzyloxy)-N-(4-fluorobenzyl)-4-(methoxymethyl)-N,6-dimethylnicotinamide (compound 12b)

Sodium hydride in tetrahydrofuran (10.0 mL) was added to **compound 12a** (0.690 g, 1.82 mmol, 1 eq) at room temperature. The reaction mixture was stirred at room temperature 20 min. and 0.541 g of iodomethane (3.81 mmol, 2.1 eq) was added. Water was added and reaction mixture was extracted with ethyl acetate.

Organic phase recovered was dried over magnesium sulfate and concentrated. The crude product was purified by silica gel (60% ethyl acetate/hexane) yielding 0.400 g of **compound 12b** (22% yield) as a white solid; MS-ESI m/z 409 $[MH]^+$.

Step 12c: preparation of 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl)carbamoyl)-4-(methoxymethyl)-2-methylpyridine 1-oxide (compound 12b)

Compound 12b (0.400 g, 0.980 mmol, 1 eq) in dichloromethane (20.0 mL) was added to 3-chloroperbenzoic acid (0.254 g, 1.47 mmol, 1.5 eq) at room temperature and the reaction mixture was stirred for 1 hour at room temperature. A 1M solution of potassium carbonate was added and the reaction mixture was extracted with dichloromethane. Organic phase recovered was dried over magnesium sulfate and concentrated yielding 0.400 g of **compound 12c** (95% yield) as a white solid; MS-ESI m/z 424 $[MH]^+$.

Step 12d: preparation of 5-(benzyloxy)-N-(4-fluorobenzyl)-6-(hydroxymethyl)-4-(methoxymethyl)-N-methylnicotinamide (compound 12d)

Compound 12c (0.400 g, 0.946 mmol, 1 eq) in dichloromethane (20.0 mL) was added to trifluoroacetic anhydride (1.30 g, 4.90 mmol, 5 eq) at room temperature. The reaction mixture was stirred overnight at room temperature. A 1M solution of potassium carbonate was added and the reaction mixture was extracted with ethyl acetate. Organic phase recovered was dried over magnesium sulfate and concentrated yielding 0.400 g of **compound 12d** (97% yield) as a white solid; MS-ESI m/z 424 $[MH]^+$.

Step 12e: preparation of 5-(benzyloxy)-N-(4-fluorobenzyl)-6-formyl-4-(methoxymethyl)-N-methylnicotinamide (compound 12d)

Compound 12d (0.400 g, 0.946 mmol, 1 eq) in dichloromethane (20.0 mL) was added to activated manganese oxide (0.852 g, 9.80 mmol, 10 eq) at room

temperature. The reaction mixture was stirred overnight at room temperature. The reaction mixture was filtered on Celite and concentrated under reduced pressure. A 1M solution of potassium carbonate was added and the reaction mixture was extracted with ethyl acetate. The crude product was purified by silica gel (40% ethyl acetate/hexane) yielding 0.122 g of **compound 12e** (30% yield) as a white solid; MS-ESI m/z 423 $[MH]^+$.

Step 12f: preparation methyl 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl) carbamoyl)-4-(methoxymethyl)picolinate (compound 12f)

Compound 12e (0.122 g, 0.289 mmol, 1 eq) in methanol (20.0 mL) was added to powdered potassium hydroxide (0.042 g, 0.751 mmol, 2.6 eq) at room temperature. The reaction mixture was stirred at room temperature 10 min. and iodine (0.095 g, 0.376 mmol, 1.3 eq) was added. The reaction mixture was then stirred for 4 hours at room temperature following which a solution of sodium bisulfite was added. The reaction mixture was extracted with ethyl acetate and organic phase recovered was dried over magnesium sulfate and concentrated to give 0.114 g of **compound 12f** (95%) as a colourless oil; MS-ESI m/z 453 $[MH]^+$.

Step 12g: preparation of N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide (compound 12g)

Benzylhydroxylamine hydrochloride (0.045 g, 0.282 mmol, 1.1 eq) in tetrahydrofuran (3.0 mL) was added lithium bis(trimethylsilyl)amide (1.30 mL, 1.27 mmol, 5 eq) at -78 °C. The reaction mixture was stirred for 10 min. following which a solution of **compound 12f** (0.115 g, 0.254 mmol, 1 eq) in tetrahydrofuran (2.0 mL) was added. The reaction mixture was then stirred at -78 °C for 30 min. and a solution of saturated ammonium chloride was added. The reaction mixture was extracted with ethyl acetate and organic phase recovered was concentrated under reduced pressure. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.100 g of **compound 12g** (72% yield) of a clear oil; MS-ESI m/z 544 $[MH]^+$.


Step 12h: preparation of N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide

Compound 12g (0.100 g, 0.184 mmol, 1 eq) and 10% Pd/C (5 mg) catalyst were stirred in methanol (4.0 mL) under a hydrogen atmosphere for 1 hour. The

catalyst was filtered and the reaction mixture was concentrated under vacuum yielding 0.040 g of **N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide** (60% yield) as a white solid; MS-ESI m/z 364 [MH]⁺.

5


Example 13: Preparation of N², 3-bis(benzyloxy)-N⁵-(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (Product 13)



4-Fluoro-3-Chloro benzylamine (0.060 g, 1.29 mmol, 2 eq) and N, 7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3, 4-c]pyridine-6-carboxamide (0.050 g, 0.141 mmol, 1 eq) (**compound 1c** of example 1) were heated in DMF at 90 °C for 120 min. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.03 g of **N², 3-bis(benzyloxy)-N⁵-(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide** as a white solid; MS-ESI m/z 573 [MH]⁻.

15


Example 14: Preparation of N⁵-(3-chloro-4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (Product 14)



The product of example 13 (0.03 g, 0.330 mmol,) and 10% Pd/C catalyst (25 mg) were stirred in methanol (4.0 mL) under an atmosphere of hydrogen for 1 hour. The catalyst was filtered and the reaction mixture was concentrated under vacuum yielding 0.015 g of **N⁵-(3-chloro-4-fluorobenzyl)-N², 3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide** as a white solid; ¹H-NMR (400 MHz, dmso) : δ = 13.01 (s, 1H), 12.0 (s, 1H), 9.51 (s 1H), 9.02 (t 1H), 8.11 (s, 1H), 7.60 (d 1H), 7.40 (m, 2H), , 4.68 (s, 2H), 4.48 (s, 2H); MS-ESI m/z 370 [MH]⁺.


25

Example 15: Preparation of N², 3-bis(benzyloxy)-N⁵-(3, 4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (Product 15)




3,4 diChloro benzylamine (0.060 g, 1.29 mmol, 2 eq) and N,7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide (0.050 g, 0.141 mmol, 1 eq) (**compound 1c** of example 1) were heated in DMF at 90 °C for 120 min. The crude product was purified by silica gel (100% ethyl acetate) yielding 0.03 g of N², 3-bis(benzyloxy)-N⁵-(3, 4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (29 %) as a white solid; MS-ESI *m/z* 589 [MH]⁺.

Example 16: Preparation of N⁵-(3, 4-dichlorobenzyl)-N²-(3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (Product 16)



Product 15 of example 15 (0.03 g, 0.330 mmol,) and 10% Pd/C (25 mg) catalyst were stirred in methanol (4.0 mL) under an atmosphere of hydrogen for 1 hour. The catalyst was filtered and the reaction mixture was concentrated under vacuum to give 0.006 g of N⁵-(3, 4-dichlorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide as a white solid; ¹H-NMR (400MHz, dmso) : δ = 13.01 (s,1H), 12.0 (s, 1H), 9.51 (s 1H), 9.02 (t 1H), 8.11 (s, 1H), 7.60 (d 1H), 7.40 (m, 2H), , 4.68 (s, 2H), 4.48 (s, 2H); MS-ESI *m/z* 386 [MH]⁺.

Example 17: Preparation of N, 3-Dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide (Product 17)



Step 17a: preparation of 5-((4-Methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine (compound 17a)

Anhydrous THF (200 ml) was added to NaH (60%, 24 g, 600 mmol) at 0°C under a nitrogen atmosphere. To this suspended mixture a solution of 2, 2, 8-trimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl)methanol (32.0 g, 150 mmol) in 400 ml of THF (J. Med Chem., 1977, 20, p745) was added. The resulting mixture was refluxed for 30 min; a significant amount of precipitate accumulated during the reflux. After cooling to room temperature, *p*-methoxybenzyl chloride (23.5 g, 150 mmol) was introduced drop-wise and the resulting mixture was refluxed for another 8 h. The reaction was quenched carefully by adding ice-cold water to the viscous mixture at 0°C and diluted with a saturated ammonium chloride solution followed by extraction with methylene chloride. The combined organic extracts were washed with brine, dried (Na₂SO₄), and concentrated yielding a brown oil. The crude product was purified by chromatography (10% ethyl acetate/petroleum ether) yielding 25.0 g of **compound 17a** (50% yield); LC-MS (M+H)⁺ m/z 331.

Step 17b: preparation of 5-((4-Methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine 7-oxide (compound 17b)

Compound 17a was dissolved in dry CH₂Cl₂ (500 mL) and the solution was cooled to 0°C and *m*-chloroperbenzoic acid (85% purity of the reagent, 37.0 g, 182 mmol, 1.2 equiv) was added. After stirring at 23°C for 12 h, the reaction mixture was extracted with Na₂SO₃ (10%, 2×200 mL), NaHCO₃ (5%, 2×200 mL), H₂O, dried (Na₂SO₄), filtered and the solvent was removed under reduced pressure. The crude product was purified by chromatography (10% methanol/ethyl acetate) yielding 35 g of **compound 17b** (68% yield) as pale-yellow solid; LC-MS (M+H)⁺ m/z 346).

Step 17c: preparation of 5-((4-Methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridin-8-yl)methanol (compound 17c)

Trifluoroacetic anhydride (4.5 mL, 32 mmol) was added to a solution of **compound 17b** (21.2 g, 61 mmol) in dry CH₂Cl₂ (200 ml) at 0°C and stirred for 5 min. An additional amount of trifluoroacetic anhydride (11.5 mL, 82.7 mmol) was added and the reaction mixture was stirred overnight at 23°C. The reaction mixture was cooled to 0 °C and MeOH (150 mL) was added while stirring was continued. The solvents were evaporated and the resulting residue was dissolved in CH₂Cl₂ and washed with Na₂CO₃ (20% aqueous) and H₂O until the pH was neutral. Organic phase recovered was dried (Na₂SO₄), filtered and concentrated under vacuum. The residue was crystallized from EtOH-CH₂Cl₂ yielding 17.5 g of **compound 17c** (83% yield); LC-MS (M+H)⁺ m/z 346.

Step 17d: preparation of 5-((4-Methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carbaldehyde (compound 17d)

IBX (35.5 g, 128 mmol) was added to a solution of 5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridin-8-yl)methanol (**compound 17c**) (14.60 g, 42.3 mmol) in ethyl acetate (500 mL) and the suspension was heated to reflux for 4 h. The precipitate was removed by filtration and the filtrate was concentrated under reduced pressure yielding 14.0 g of **compound 17d** (95% yield). Crude **compound 17d** was used for the next step without further purification; LC-MS (M+H)⁺ m/z 344.

Step 17e: preparation of ethyl 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxylate (compound 17e)

KOH (85%, 5.3 g, 78 mmol) and iodine (9.9 g, 39 mmol) were added to a solution of **compound 17d** (10.3 g, 30 mmol) in anhydrous MeOH (120 mL), at 0°C. The reaction mixture was kept at 23°C and stirred for 12 h until no starting material was detected by TLC. The solution was then treated with Na₂SO₃ (solid) and the pH was adjusted to 7. The solid was filtered and solvent was removed under reduced pressure. The residue was dissolved in EtOAc and washed with water. The combined organic extracts were dried (Na₂SO₄), filtered and concentrated in vacuum. The crude

residue was purified by chromatography (SiO₂) with petroleum ether:ethyl acetate (5:1) as eluent yielding 8.8 g of **compound 17e** (78% yield) as pale-yellow solid; LC-MS (M+H)⁺ m/z 374.


5 **Step 17f: preparation of ethyl 3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinate (compound 17f)**

A solution of **compound 17e** (8.8 g, 23.6 mmol) in 200 mL of HCl/MeOH was stirred at 23°C for 24 h. MeOH (500 mL) was added to dissolve the suspension and NaHCO₃ (solid) was added to neutralize the reaction mixture. Excess solid was filtered and the solvent was removed under reduced pressure. The residue was dissolved in EtOAc and washed with water. The combined organic extracts were dried (Na₂SO₄), filtered and concentrated in vacuum yielding 6.0 g of **compound 17f** (100% yield) as a light yellow solid; LC-MS (M+H)⁺ m/z 334.

10 **Step 17g: preparation of N, 3-Dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinamide**

Hydroxylamine hydrochloride (0.042 g, 0.60 mmol) and *N,N*-diisopropylethylamine (0.13 mL, 0.75 mmol) were added to a solution of **compound 17f** (0.050 g, 0.15 mmol) in MeOH (3 mL). The resulting mixture was placed in the microwave and heated to 80°C for 1.5 h. The crude mixture was diluted with EtOAc and washed with saturated aqueous ammonium chloride solution and brine. The organic extract was dried (Na₂SO₄), filtered and concentrated in vacuum to yielding 0.039 g of **N, 3-Dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide** (78% yield); LC-MS (M+H)⁺ m/z 335.


20 **Example 18: Preparation of 5-(Benzyloxymethyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide (Product 18)**



The procedure described in Example 17 was applied using benzyl bromide in step 17a yielding 0.054 g of **5-(Benzyloxymethyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide** as a white solid (45% yield); LC-MS (M+H)⁺ m/z 305.

5

Example 19: Preparation of N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide (Product 19)



Step 19a: preparation of 5-((4-Methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxylic acid (compound 19a)

Lithium hydroxide monohydrate (0.049 g, 1.18 mmol) was added to a solution of **compound 17e** (0.367 g, 0.98 mmol) in THF/H₂O (5/10 mL). The resulting mixture was stirred at 23°C for 18 h. The solution was then acidified with AcOH, extracted with EtOAc and washed with brine. The organic extract was dried (Na₂SO₄), filtered and concentrated in vacuum yielding 0.315 g of **compound 19a** (90% yield); LC-MS (M+H)⁺ m/z 360.

Step 19b: preparation of N-(Benzyloxy)-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide (compound 19a)

A solution of **compound 19a** (0.307 g, 0.86 mmol) and O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) (0.389 g, 1.03 mmol) in acetonitrile (20 mL) was stirred at 23°C for 15 minutes. Benzylhydroxylamine hydrochloride (0.137 g, 0.86 mmol) and N,N-diisopropylethylamine (0.45 mL, 2.58 mmol) were added to the solution and the reaction mixture was stirred at 23°C for 17 h. The solvent was then removed in vacuum, and saturated aqueous ammonium chloride solution was added, followed by extraction with EtOAc. The combined organic extracts were washed with brine, dried (MgSO₄), filtered and concentrated in

vacuum. The crude residue was purified by flash chromatography (SiO₂) with hexane:ethyl acetate as eluent yielding 0.380 g of **compound 19b** (95% yield); LC-MS (M+H)⁺ m/z 465.

5 **Step 19c: preparation of N-(Benzyloxy)-3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl) picolinamide (compound 19c)**

A solution of **compound 19b** (0.060 g, 0.13 mmol) in hydrogen chloride-ethanol solution (5 mL) was stirred at 23°C for 5 h. The solvent was then removed in vacuum and an aqueous solution of NaHCO₃ (1M) was added. The reaction mixture was extracted with CH₂Cl₂ and the combined organic extracts were washed with brine,
10 dried (MgSO₄), filtered and concentrated in vacuum yielding 0.030 g of **compound 19c** (54% yield); LC-MS (M+H)⁺ m/z 425.


Step 19d N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide

A solution of **compound 19c** (0.10 g, 0.22 mmol) in ethyl acetate (10 mL) was
15 hydrogenated under 1 atm of hydrogen at 23°C over 10% palladium on activated carbon for 1 h. The reaction mixture was filtered and the solution was concentrated in vacuum. The residue was purified on preparative TLC with CH₂Cl₂:MeOH (9:1) as eluent yielding 0.016 g of **N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide** (20% yield). LC-MS
20 (M+H)⁺ m/z 375.


Example 20: Preparation of N⁵-(3,4-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (Product 20)

N⁵-(3,4-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-
25 dicarboxamide was prepared using the procedure described in Example 2 and using 3,4 difluorobenzylamine as the starting material.

Example 21: Preparation of 5-[(4-Fluoro-phenylamino)-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 21)




Step 21a: Preparation of methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 21a)



Acetic acid (50 μ L, 0.87 mmol) was added to a mixture of 5-Formyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid methyl ester [*J. Org. Chem.* 1999, 64, 4537] (220 mg, 0.87 mmol) and 4-fluoroaniline (125 μ L, 1.3 mmol) in dry methanol (4 mL) and stirred at room temperature for 15 min. and sodium cyanoborohydride (72 mg, 1.3 mmol) was added. This mixture was stirred at room temperature for 3 h followed by evaporation of 90% of the methanol volume under reduced pressure. The residue was extracted with dichloromethane (3 x 25 mL) and the combined organic layers were dried (anh. Na_2SO_4), filtered and concentrated under reduced pressure. The crude product was purified by silica gel (methanol/dichloromethane, 0 to 10% methanol) to give 0.290 g of methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 21a) (96 %) as a white solid MS-ESI m/z 347 $[\text{MH}]^+$.

Step 21b: Preparation of 5-((4-fluorophenylamino)methyl)-3-hydroxy-4-(hydroxymethyl) picolinate (compound 21b)



Formic acid (2 mL) was added to methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (2) (180 mg, 0.5 mmol) at


0 °C and stirred at room temperature for 2 hours. Evaporation under reduced pressure afforded a residue 5-((4-fluorophenylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (**compound 21b**), which was triturated in acetonitrile. MS-ESI m/z 307 [MH]⁺

5 **Step 21c: Preparation of N⁵-(3,4-difluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (Product 21)**


Diisopropylethylamine (142 µL, 0.8 mmol) and hydroxylamine hydrochloride (45 mg, 0.64 mmol) were added to a solution of 5-((4-fluorophenylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (**compound 21b**) (50.0 mg, 0.16 mmol) in methanol (1.0 mL). The reaction mixture was heated to 55 °C for 16 h. The reaction mixture was allowed to cool to room temperature and a saturated solution of ammonium chloride was added. The reaction mixture was extracted with ethyl acetate (3 x 25 mL) and the combined organic layers were dried (anh. Na₂SO₄), filtered and concentrated under reduced pressure to give a solid residue which was recrystallised in acetonitrile to give compound 21.

¹H-NMR (500MHz, DMSO-d₆) δ ppm 8.24 (s, 1H), 7.45 (m, 4H), 4.94 (s, 2H), 3.77 (s, 2H), 3.33 (s, 2H); MS-ESI m/z 308 [MH]⁺

20 **Example 22: Preparation of 5-{[2-(4-Fluoro-phenyl)-ethylamino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 22)**




25 **Step 22a: Preparation of Methyl 5-((4-fluorophenethylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate**



Methyl 5-((4-fluorophenethylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound **22a**) was synthesized from methyl 5-formyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate and 4-fluorophenethylamine as described in example 21. MS-ESI m/z 375 $[MH]^+$.

5 **Step 22b: Preparation of Methyl 5-((4-fluorophenethylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate**




Methyl 5-((4-fluorophenethylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (**3**) was synthesized starting from methyl 5-((4-fluorophenethylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound **22a**) as described in example 21. MS-ESI m/z 335 $[MH]^+$


15 **Step 22c: Preparation of Product 22**

Product 22 was synthesized following a procedure similar to the one described above for the synthesis of 5-((4-fluorophenylamino)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide starting from methyl 5-((4-fluorophenethylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (compound **22b**). ^1H -NMR (500MHz, DMSO- d_6) δ ppm 8.08 (s, 1H), 7.24 (m, 2H), 7.09 (t, J = 7.2 Hz, 2H), 4.58 (s, 2H), 3.92 (s, 2H), 2.80 (m, 2H), 2.76 (m, 2H). MS-ESI m/z 336 $[MH]^+$

25 **Example 23: Preparation of 5-(4-Fluoro-benzoylamino)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 23)**




Step 23a: Preparation of 8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid (compound 23a)




8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid (23a) was synthesized from 5-Formyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid methyl ester [*J. Org. Chem.* 1999, 64, 4537] 1 mmol and 1.5 mmol NaClO₂ in t-butanol/water .

Step 23b: Preparation of Methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 23b)




Methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (23b) was made according to the following procedure: diphenyl phosphorylazide (275 μ L, 1.46 mmol) was added to a mixture of 8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid (compound 23a) (390 mg, 1.46 mmol) and triethylamine (205 μ L, 1.46 mmol) in DMF (7 mL). This mixture was stirred for 3h and water was added. The reaction mixture was then extracted with ethyl ether (3 x 25 mL) and the combined organic layers were washed with saturated sodium bicarbonate, dried (anh. Na₂SO₄), filtered and concentrated under reduced pressure to give a solid, the acylazide intermediate. This intermediate was dissolved in toluene (5 mL) and benzyl alcohol (1.1 mL) was added. The mixture was heated to reflux for 4 h. and allowed to cool at room temperature. Evaporation afforded a residue which was purified by silica gel with methanol / dichloromethane (5% to 35%).

Step 23c: Preparation of 5-Amino-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid methyl ester (compound 23c)




A mixture of methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (Compound 23b) (110 mg, 0.25 mmol) and palladium on charcoal 10% (20 mg) in methanol was vigorously stirred at room temperature for 12 h. under an atmosphere of hydrogen. Filtration through Celite and evaporation afforded the aniline, compound 23c. MS-ESI m/z 239 $[MH]^+$.

Step 23d: Preparation of Methyl 5-(4-fluorobenzamido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 23d)



A mixture of methyl 5-amino-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (23c) (70 mg, 0.25 mmol), 4-fluorobenzoyl chloride (35 μ L, 0.27 mmol) and 4-dimethylaminopyridine in pyridine (1 mL) was stirred at room temperature overnight and a saturated solution of ammonium chloride was added. The reaction mixture was extracted with dichloromethane (3 x 25 mL) and the combined organic layers were dried (anh. Na_2SO_4), filtered and concentrated under reduced pressure to give methyl 5-(4-fluorobenzamido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (23d) as a white solid MS-ESI m/z 361 $[MH]^+$.

Step 23e: Preparation of Methyl 5-(4-fluorobenzamido)-3-hydroxy-4-(hydroxymethyl)picolinate (compound 23e)




Methyl 5-(4-fluorobenzamido)-3-hydroxy-4-(hydroxymethyl)picolinate (**23e**) was synthesized from methyl 5-(4-fluorobenzamido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (Compound **23d**) and formic acid as described above. MS-ESI m/z 361 $[MH]^+$.

5 **Step 23f: Preparation of Product 23**

5-(4-fluorobenzamido)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide was synthesized from methyl 5-(4-fluorobenzamido)-3-hydroxy-4-(hydroxymethyl)picolinate (compound **23e**) using the procedure described in Example 22. MS-ESI m/z 322 $[MH]^+$.

10

Example 24: Preparation of (8-Hydroxycarbamoyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl)-carbamic acid benzyl ester (Product 24)




15

Benzyl 8-(hydroxycarbamoyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-ylcarbamate was prepared from methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (**23b**) according to the procedure described in Example 23. MS-ESI m/z 374 $[MH]^+$.


20

Example 25: Preparation of 5-[(Benzyl-(4-fluoro-phenyl)-amino)-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 25)



25

Step 25a: Preparation of Methyl 5-((benzyl(4-fluorophenyl)amino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 25a)




- 5 Methyl 5-((benzyl(4-fluorophenyl)amino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate **25a** was prepared from methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (**21a**) (120 mg, 0.3 mmol), benzaldehyde (100 μ L, 1.0 mmol) and sodium cyanoborohydride (28 mg, 0.45 mmol) according to the procedure described in
- 10 Example 24. MS-ESI m/z 437 $[MH]^+$.

Step 25b: Preparation of Product 25


Product 25 was obtained by analogous procedures to Example 21 Step 21b and 21c. MS-ESI m/z 398 $[MH]^+$.

- 15 **Example 26: Preparation of 5-((2-benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 26)**




20

Step 26a: Preparation of compound 26a



Methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (compound **22a**) was reacted with benzyloxycetaldehyde according to the procedure described above. MS-ESI m/z 509 $[MH]^+$.

Step 26b: Preparation of Methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (compound 26b)




Methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound **26a**) was reacted with formic acid according to the procedure described in example 21 step 21b. MS-ESI m/z 469 $[MH]^+$.


Step 26c: Preparation of 5-(((2-Benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino)-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 26)

Product 26 was made from methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (compound **23b**) according to the procedure described in example 21 step 21c. MS-ESI m/z 470 $[MH]^+$.

Example 27: Preparation of 5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 27)




Step 27a Preparation of Methyl 5-(3-(4-fluorophenyl)ureido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 27a)



Compound 27a was prepared according to the procedure described for the synthesis of methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (23b) above, except that in the second step, benzyl alcohol was substituted by 4-fluoroaniline. MS-ESI m/z 376 $[MH]^+$.

Step 27b: Preparation of Methyl 5-(3-(4-fluorophenyl)ureido)-3-hydroxy-4-(hydroxymethyl)picolinate (compound 27b)




Compound 27b was made from methyl 5-(3-(4-fluorophenyl)ureido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate 27a according to the procedure described in example 23, step 23e. MS-ESI m/z 337 $[MH]^+$.


Step 27c: Preparation of 5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 27)

Product 27 was made from methyl 5-(3-(4-fluorophenyl)ureido)-3-hydroxy-4-(hydroxymethyl)picolinate 27b according to the procedure described above. MS-ESI m/z 336 $[MH]^+$.

Example 28: Preparation of 5-(4-Fluoro-phenoxy-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 28)




Step 28a: Preparation of Methyl 5-((4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 28a)



Triphenylphosphine (0.105 g, 0.40 mmol), followed by diethyl azodicarboxylate (DEAD) (0.06 mL, 0.40 mmol) were added to a solution of methyl 5-(hydroxymethyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (0.102 g, 0.40 mmol) in 10 mL of THF, at room temperature under Ar. The resulting mixture was stirred at room temperature for 6 h and then concentrated. The crude residue was purified by chromatography (SiO₂) with hexanes: ethyl acetate (1:1) as eluent to afford the title compound with a contaminant. LC-MS (M+H)⁺ m/z 348.

Step 28b: Preparation of Methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (compound 28b)




A solution of methyl 5-((4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate **28a** (0.129 g, 0.37 mmol) in 3 mL of formic acid was stirred at 23°C for 2 h and then it was concentrated. The crude residue was purified by chromatography (SiO₂) with hexanes : ethyl acetate (3:7) as eluent to afford the title compound as a white solid: LC-MS (M+H)⁺ m/z 308; ¹H NMR (DMSO-*d*₆): 1.39 (s, 6H), 4.80 (s, 2H), 4.89 (s, 2H), 5.01 (s, 2H), 7.39-7.46 (m, 5H), 8.38 (s, 1H), 10.04 (s, 1H).

Step 28c: Preparation of Product 28


Hydroxylamine hydrochloride (0.045 g, 0.65 mmol) and *N,N*-diisopropylethylamine (0.14 mL, 0.81 mmol) were added to a solution of methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate **28b** (0.050 g, 0.16 mmol) in MeOH (3 mL) and heated to 70°C for 5 h. The crude mixture was diluted with EtOAc and washed with saturated aqueous ammonium chloride solution and

brine. The organic extract was dried (Na_2SO_4), filtered and concentrated in vacuo to afford the title compound (0.048 g, 96%): LC-MS ($\text{M}+\text{H}^+$) m/z 309.

Example 29: Preparation of 5-(3-Chloro-4-fluorophenoxy)methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 29)




Step 29a: Preparation of Methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (compound 29a)



Using similar procedure as described in the example of methyl 5-((4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (step 28a), we obtained compound 29a: ESI-MS ($\text{M}+\text{H}^+$) m/z 382.

Step 29b: Preparation of Methyl 5-((3-chloro-4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate (compound 29b)




Using similar procedure as described in the example 28, in the preparation of methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate 28b, we obtained compound 29b as a white solid: yield (63%); LC-MS ($\text{M}+\text{H}^+$) m/z 342.

Step 29c: Preparation of Product 29

Using similar procedure as described in the example 28, in the preparation of 5-((4-fluorophenoxy)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide 28c, we obtained Product 29 as a beige solid: yield (98%); LC-MS ($\text{M}+\text{H}^+$) m/z 342.


Example 30: Preparation of 5-(3-Chloro-4-fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 30)



Product 30 was prepared in a procedure analogous to that shown in example 20 with the exception of the use of 4-fluoro-3-chloro-benzyl chloride as alkylating agent.

LC-MS (M+H)⁺ m/z 357.7.


Example 31: Preparation of 5-[2-(4-Fluoro-phenyl)-ethoxymethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 31)



Product 31 was prepared in a procedure analogous to that shown in example 20 with the exception of the use of 4-fluoro-3-chloro-benzyl chloride as alkylating agent.

LC-MS (M+H)⁺ m/z 337.


Example 32: Preparation of 5-(2,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 32)



Product 32 was prepared in a procedure analogous to that shown in example 20 with the exception of the use of 2,4-difluoro-benzyl chloride as alkylating agent.

LC-MS (M+H)⁺ m/z 341.


Example 33: Preparation of 5-(3,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 33)



Product 33 was prepared in a procedure analogous to that shown in example 20 with the exception of the use of 3,4-difluoro -benzyl chloride as alkylating agent.

LC-MS (M+H)⁺ m/z 341.


Example 34: Preparation of 5-(4-Fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide




Product 34 was prepared in a procedure analogous to that shown in example 20 with the exception of the use of 4-fluoro-benzyl chloride as alkylating agent.

LC-MS (M+H)⁺ m/z 323.

Example 35: Preparation of 5-(4-Fluoro-phenoxy)methyl-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide (Product 35)




Step 35a: Preparation of Methyl 3-(benzyloxy)-5-((4-fluorophenoxy)methyl)-4-methylpicolinate (compound 35a)



Using a procedure similar to that described in the example 28 in the preparation of of methyl 5-((4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate **28a**, we obtained **35a** as a colorless oil: Yield (72%); LC-MS (M+H)⁺ m/z 382.

5 **Step 35b: Preparation of Methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate (compound 35b)**




A solution of methyl 3-(benzyloxy)-5-((4-fluorophenoxy)methyl)-4-methylpicolinate **35a** (0.245 g, 0.64 mmol) in ethyl acetate (10 mL) was hydrogenated under 1 atm of
10 hydrogen at 23°C over 10% palladium on activated carbon for 1 h. The reaction mixture was filtrated and the solution was concentrated in vacuo to afford **35b** as a white solid (0.171 g, 92%); LC-MS (M+H)⁺ m/z 292.


Step 35c: Preparation of Product 35

Using similar procedure as described in the example 28 in the preparation of 5-((4-fluorophenoxy)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide,(28c)
15 **Product 35** was obtained as an off-white solid: Yields (97%); LC-MS (M+H)⁺ m/z 293.

20 **Example 36: Preparation of 5-(3-Chloro-4-fluoro-phenoxy)methyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide (Product 36)**



Step 36a: Preparation of Methyl 3-(benzyloxy)-5-((3-chloro-4-fluorophenoxy)methyl)-4-methylpic (compound 36a)



Using similar procedure as described in the example of Methyl 3-(benzyloxy)-5-((4-fluorophenoxy)methyl)-4-methylpicolinate, (**example 35a**) we obtained the title compound as a colorless oil: Yields (68%); LC-MS (M+H)⁺ m/z 416.


Step 36b: Preparation of Methyl 5-((3-chloro-4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate (compound 36b)

A solution of methyl 3-(benzyloxy)-5-((3-chloro-4-fluorophenoxy)methyl)-4-methylpicolinate (0.167 g, 0.13 mmol) in trifluoroacetic acid (6 mL) was stirred at 23°C for 2 days. The solvent was then removed in vacuo and an aqueous solution of NaHCO₃ (1M) was added, followed by extraction with ethyl acetate. The combined organic extracts were washed with brine, dried (Na₂SO₄), filtered and concentrated. The crude residue was purified by flash chromatography (SiO₂) with hexane:ethyl acetate (7:3) as eluent to afford **36b** as a white solid (0.113 g, 86%): LC-MS (M+H)⁺ m/z 326.

Step 36c: Preparation of Product 36

Using a procedure similar to that described in the example 28 in the preparation of of 5-((4-fluorophenoxy)methyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide, (**28c**) **Product 36** was obtained as an off-white solid: Yield (100%); LC-MS (M+H)⁺ m/z 327.


Example 37: Preparation of 5-(2,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide (Product 37)



Product 37 was prepared using a procedure analogous to that shown in example 20 with the exception of the use of 4-fluoro-3-chloro-benzyl chloride as an alkylating agent, and 3-Benzyloxy-5-hydroxymethyl-4-methyl-pyridine-2-carboxylic acid methyl ester: LC-MS (M+H)⁺ m/z 325.

5

Example 38: Preparation of 5-(3,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide (Product 38)




10

Product 38 was prepared using a procedure analogous to that shown in example 20 with the exception of the use of 3,4-difluoro -benzyl chloride as the alkylating agent and 3-Benzyloxy-5-hydroxymethyl-4-methyl-pyridine-2-carboxylic acid methyl ester. LC-MS (M+H)⁺ m/z 325.

15


Example 39: Preparation of 5-(4-Fluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide (Product 39)



20

Product 39 was prepared using a procedure analogous to that shown in example 20 with the exception of the use of 4-fluorobenzyl chloride as the alkylating agent and 3-Benzyloxy-5-hydroxymethyl-4-methyl-pyridine-2-carboxylic acid methyl ester. LC-MS (M+H)⁺ m/z 307.


Example 40: Preparation of 5-Benzyloxymethyl-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide (Product 40)



Product 40 was prepared using a procedure analogous to that shown in example 20 with the exception of the use of benzyl chloride as the alkylating agent and 3-Benzyloxy-5-hydroxymethyl-4-methyl-pyridine-2-carboxylic acid methyl ester. LC-MS (M+H)⁺ m/z 389.


5

Example 41: Preparation of (S)- (-)-3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(1-phenyl-ethyl)-amide] (Product 41)



10

Step 41a: Preparation of compound 41a



15

(S)-(-)- α -Methylbenzylamine (0.027 g, 0.226 mmol, 2.2 eq) and **III** (scheme I) (0.040 g, 0.103 mmol, 1 eq) were heated neat at 70 °C for 20 min. The crude product was purified by silica gel (50% ethyl acetate/hexane) to give 0.043 g of **41a** (83%) as a white solid. MS-ESI *m/z* 512 [MH]⁺.


Step 41b: Preparation of Product 41

20


(0.040 g, 0.078 mmol, 1 eq) and 10% Pd/C (5 mg) in methanol (4.0 mL) were stirred under an atmosphere of hydrogen 1 hour. The catalyst was filtered and reaction mixture was concentrated under vacuum to give 0.015 g of **41b** (58%) as a white solid. ¹H-NMR (400MHz, MeOD) : δ = 8.16 (s, 1H), 7.42 (m, 5H), 4.75 (s, 2H), 4.59 (s, 2H), 1.58 (s, 3H); MS-ESI *m/z* 332 [MH]⁺.

25

Example 42: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(pyridin-2-ylmethyl)-amide] (Product 42)



Step 42a: Preparation of xxx (compound 42a)



5


2-(Aminomethyl)pyridine (0.024 g, 0.226 mmol, 2.2 eq) and **III** (Scheme I) (0.040 g, 0.103 mmol, 1 eq) were heated neat at 70 °C for 20 min to give compound 42a. MS-ESI m/z 499 $[MH]^+$.

Step 42b: Preparation of Product 42

10 Compound 42a (0.103 mmol, 1 eq) and 10% Pd/C (5 mg) in methanol (4.0 mL) were stirred under an atmosphere of hydrogen 1 hour. The catalyst was filtered and triturated with diethyl ether and acetonitrile and precipitated with MeOH/acetonitrile to give 0.003 g of Product **42** (1%) as a white solid. ^1H -NMR (400MHz, MeOD) : δ = 8.53 (d, J = 5.7 Hz, 1H), 8.12 (s, 1H), 7.84 (m, 1H), 7.55 (d, J = 7.7 Hz, 1H), 7.34


15 (m, 1H), 4.90 (s, 2H), 4.58 (s, 2H); MS-ESI m/z 319 $[MH]^+$.

Example 43: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-benzylamide 2-hydroxyamide (Product 43)



20

Step 43a: Preparation of compound 43a




Benzylamine (0.024 g, 0.235 mmol, 2.5 eq) and **III** (Scheme I) (0.040 g, 0.103 mmol, 1 eq) were heated neat at 70 °C for 20 min. The crude product was purified by silica gel (60% ethyl acetate/hexane) to give 0.012 g of compound **43a** (24%) as a white solid. MS-ESI m/z 498 $[MH]^+$.

5 **Step 43b: Preparation of Product 43**

Compound 43a (0.012 g, 0.024 mmol, 1 eq) and 10% Pd/C (5 mg) in methanol (4.0 mL) were stirred under an atmosphere of hydrogen 1 hour. The catalyst was filtered and reaction mixture was concentrated under vacuum to give 0.008 g of **43b** (75%) as a white solid. MS-ESI m/z 318 $[MH]^+$.


10

Example 44: Preparation of (S)-(-)-3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(2-hydroxy-1-phenyl-ethyl)-amide] (Product 44)



15

Step 44a: Preparation of compound 44a




20

N,N-diisopropylethylamine (0.275 g, 2.133 mmol, 3 eq) and *N,N,N',N'*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uranium hexafluorophosphate (0.404 g, 1.067 mmol, 1.5 eq) were added to 8-(methoxycarbonyl)-2,2-dimethyl-4*H*-[1,3]dioxino[4,5-*c*]pyridine-5-carboxylic acid (**23a**) (0.190 g, 0.711 mmol, 1eq) and (*S*)-phenylglycinol (0.107 g, 0.783 mmol, 1.1 eq) in *N,N*-dimethylformamide (5.0 mL). The reaction mixture was stirred at room temperature overnight. Water was added and reaction mixture was

25

extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated. The crude product was purified by silica gel (75% ethyl acetate/hexane) to give 0.254 g of **44a** (93%) as a white solid. MS-ESI m/z 387 $[MH]^+$.

5

Step 44b: Preparation of compound 44b

Hydroxylamine solution 50 wt. % in water (5.0 mL) was added to **44a** (0.178 g, 0.461 mmol, 1 eq) in tetrahydrofuran (5.0 mL). The reaction mixture was stirred at reflux 3hrs. The pH was adjusted to 6 and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated under vacuum to give 0.045 g of **44b** (25%) as a white solid. MS-ESI m/z 388 $[MH]^+$.



10

Step 44c: Preparation of Product 44

To **44b** (0.045 g, 0.116 mmol, 1 eq) was added formic acid (2 mL). The reaction mixture was stirred at room temperature 10 min. Formic acid was concentrated under vacuum and solid was triturated with diethyl ether to give 0.035 g of **44** (87%) as a white solid. MS-ESI m/z 348 $[MH]^+$.

15


20

Example 45: Preparation of Pyridine-2,5-dicarboxylic acid 5-(4-fluorobenzylamide) 2-hydroxyamide (Product 45)**Step 45a: Preparation of compound 45a**

25


4-fluorobenzylamine (1.00 g, 8.02 mmol, 1.1 eq), *N,N*-diisopropylethylamine (0.942 g, 21.87 mmol, 3 eq) and *N,N,N',N''*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uranium hexafluorophosphate (4.15 g, 10.93 mmol, 1.5 eq) were added to 6-methylnicotinic acid (1.00 g, 7.29 mmol, 1 eq) in *N,N*-dimethylformamide (10.0 mL). The reaction mixture was stirred at room temperature overnight. Water was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated. The crude product was purified by silica gel (75% ethyl acetate/hexane) to give 1.44 g of **45a** (80%) as a white solid. MS-ESI *m/z* 245 [MH]⁺.

Step 45b: Preparation of compound 45b




3-chloroperbenzoic acid (1.30 g, 7.56 mmol, 1.5 eq) at room temperature was added to **45a** (1.23 g, 5.04 mmol, 1 eq) in dichloromethane (30.0 mL). The reaction mixture was stirred 1 hour at room temperature. A 1M solution of potassium carbonate was added and reaction mixture was extracted with dichloromethane. Organic phases were combined, dried over magnesium sulfate and concentrated to give 0.908 g of **45b** (69%) as a white solid. MS-ESI *m/z* 260 [MH]⁺.

Step 45c: Preparation of compound 45a



Trifluoroacetic anhydride (3.18 g, 15.12 mmol, 4.3 eq) at room temperature was added to **45b** (0.908 g, 3.49 mmol, 1 eq) in dichloromethane (20.0 mL). The reaction mixture was stirred overnight at room temperature. A 1M solution of potassium carbonate was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated to give 0.900 g of **45c** (99%) as a white solid. MS-ESI *m/z* 260 [MH]⁺.


Step 45d: Preparation of compound 45a


Activated manganese oxide (0.083 g, 0.962 mmol, 5 eq) at room temperature was added to **45c** (0.050 g, 0.192 mmol, 1 eq) in chloroform (10.0 mL) and tetrahydrofuran (5.0 mL). The reaction mixture was stirred at reflux 1hr or until all starting material disappeared by LC-MS. Solvent was concentrated under vacuo and crude reaction mixture was dissolved in methanol (10.0 mL). A solution of sodium cyanide (0.011 g, 0.230 mmol, 1.2 eq) in methanol (3.0 mL) was added and the reaction mixture was stirred 20 min. at room temperature. The reaction mixture was filtered on Celite. Organic phase was washed with water and extracted with ethyl acetate. The crude product was purified by silica gel (100% ethyl acetate/hexane) to give 0.025 g of **45d** (46%) as a white solid. MS-ESI m/z 289 $[MH]^+$.

Step 45e: Preparation of Product 45

N,N-diisopropylethylamine (0.062 g, 0.479 mmol, 6 eq) and hydroxylamine hydrochloride (0.022 g, 0.319 mmol, 4 eq) were added to **45d** (0.023 g, 0.079 mmol, 1 eq) in methanol (5.0 mL). The reaction mixture was stirred at reflux 3hrs. Water was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated under vacuum to give 0.015 g of **45** (65%) as a white solid. MS-ESI m/z 290 $[MH]^+$.

Example 46: 2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-([1-(4-fluorophenyl)-cyclopropyl]-amide) 8-hydroxyamide (Product 46)



Step 46a: Preparation of compound 46a

5 *N,N*-diisopropylethylamine (0.145 g, 1.12 mmol, 3 eq) and *N,N,N',N''*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uranium hexafluorophosphate (0.213 g, 0.561 mmol, 1.5 eq) were added to 8-(methoxycarbonyl)-2,2-dimethyl-4*H*-[1,3]dioxino[4,5-*c*]pyridine-5-carboxylic acid (**23a**) (0.100 g, 0.374 mmol, 1 eq) and 1-(4-fluorophenyl)cyclopropylamine (0.062 g, 0.412 mmol, 1.1 eq) in *N,N*-dimethylformamide (5.0 mL). The reaction mixture was stirred at room temperature overnight. Water was added and reaction mixture was extracted with ethyl acetate.


10 Organic phases were combined, dried over magnesium sulfate and concentrated. The crude product was purified by silica gel (75% ethyl acetate/hexane) to give 0.254 g of **46a** (93%) as a white solid. MS-ESI *m/z* 401 [MH]⁺.

Step 46b: Preparation of Product 46

To **46b** (0.140 g, 0.350 mmol, 1 eq) in tetrahydrofuran (5.0 mL) was added

15 hydroxylamine solution 50 wt. % in water (5.0 mL). The reaction mixture was stirred at reflux 3hrs. Water was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated under vacuum to give 0.130 g of **46** (93%) as a white solid. MS-ESI *m/z* 402 [MH]⁺.


20 **Example 47: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-{{[1-(4-fluoro-phenyl)-cyclopropyl]-amide} 2-hydroxyamide (Product 47)**




To **46** (0.125 g, 0.312 mmol, 1 eq) was added formic acid (2 mL). The reaction

25 mixture was stirred at room temperature 10 min. Formic acid was concentrated under vacuum and solid was triturated with diethyl ether under to give 0.070 g of **47** (62%) as a white solid. MS-ESI *m/z* 362 [MH]⁺.

Example 48: Preparation of 2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-benzylamide) 8-(methoxy-amide) (Product 48)



Step 48a: Preparation of compound 48a




N,N,N',N'-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uranium hexafluorophosphate (0.287 g, 0.758 mmol, 1.5 eq) was added to 8-(methoxycarbonyl)-2,2-dimethyl-4*H*-[1,3]dioxino[4,5-*c*]pyridine-5-carboxylic acid (**23a**) (0.135 g, 0.505 mmol, 1 eq) and *N,N*-diisopropylethylamine (0.196 g, 1.515 mmol, 3 eq) in *N,N*-dimethylformamide (5.0 mL). Reaction was stirred 10 min. before 4-fluorobenzylamine (0.070 g, 0.556 mmol, 1.1 eq) was added. The reaction mixture was stirred at room temperature overnight. Water was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated. The crude product was purified by silica gel (50% ethyl acetate/hexane) to give 0.066 g of **48a** (34%) as a white solid. MS-ESI *m/z* 375 [MH]⁺.

Step 48b: Preparation of Product 48


Lithium bis(trimethylsilyl)amide (0.870 mL of 1M solution in tetrahydrofuran, 0.870 mmol, 5 eq) at -78 °C was added to methoxylamine hydrochloride (0.017 g, 0.209 mmol, 1.2 eq) in tetrahydrofuran (3.0 mL). The reaction mixture was stirred 10 min. then a solution of **48a** (0.065 g, 0.174 mmol, 1 eq) in tetrahydrofuran (3.0 mL) was added. The reaction mixture was stirred at -78 °C 30 min. before a solution of saturated ammonium chloride was added. This reaction mixture was extracted with ethyl acetate and organic phases were combined and concentrated under reduced pressure to give 0.50 g of crude product **48** (74%) as a white solid. MS-ESI *m/z* 390 [MH]⁺.

Example 49: 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(methoxy-amide)




Formic acid (2 mL) was added to **48** (0.048 g, 0.123 mmol, 1 eq). The reaction mixture was stirred at room temperature 10 min. Formic acid was concentrated under vacuum and solid was triturated with diethyl ether under to give 0.030 g of **49** (71%) as a white solid. MS-ESI m/z 350 $[MH]^+$.

Example 50: Preparation of 2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-cyclohexylmethyl-amide 8-hydroxyamide (Product 50)



Step 50a: Preparation of xxx (compound 50a)




To 8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid (**23a**) (0.100 g, 0.374 mmol, 1eq) and *N,N*-diisopropylethylamine (0.145 g, 1.122 mmol, 3 eq) in *N,N*-dimethylformamide (5.0 mL) was added *N,N,N',N'*-Tetramethyl-*O*-(1*H*-benzotriazol-1-yl)uranium hexafluorophosphate (0.213 g, 0.758 mmol, 1.5 eq). Reaction was stirred 10 min. before cyclohexanemethylamine (0.047 g, 0.412 mmol, 1.1 eq) was added. The reaction mixture was stirred at room temperature overnight. Water was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated. The crude

product was purified by silica gel (60% ethyl acetate/hexane) to give 0.040 g of **50a** (30%) as a white solid. MS-ESI m/z 363 $[MH]^+$.

Step 50b: Preparation of Product 50


5 To **50a** (0.140 g, 0.350 mmol, 1 eq) in tetrahydrofuran (5.0 mL) was added hydroxylamine solution 50 wt. % in water (5.0 mL). The reaction mixture was stirred at reflux 3hrs. The pH was adjusted to 6 and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated under vacuum to give 0.025 g of **50** (63%) as a white solid. MS-ESI m/z 364 $[MH]^+$.

Example 51: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide (Product 51)




15 To **50** (0.020 g, 0.055 mmol, 1 eq) was added formic acid (2.0 mL). The reaction mixture was stirred at room temperature 10 min. Formic acid was concentrated under vacuum and solid was triturated with diethyl ether under to give 0.015 g of **51** (88%) as a white solid. MS-ESI m/z 324 $[MH]^+$.

20 **Example 52: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide (Product 52)**




Step 52a: Preparation of compound 52a




The starting material was obtained by the catalytic hydrogenation of **XXIV** (2.0g) over Pd/C 5% in EtOAc. Filtration and evaporation gave 1.3g white powder quantitative conversion.

5

Step 52b: Preparation of compound 52b


Trimethylsilyl-diazomethane (8.40 mL, 16.8 mmol, 3 eq) at 0°C was added to **52a** (0.924 g, 0.516 mmol, 1 eq) in methanol (10.0 mL) and tetrahydrofuran (10.0 mL). The reaction mixture was stirred at room temperature 20 min. A saturated sodium bicarbonate solution was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated under reduced pressure. The crude product was purified by silica gel (50% ethyl acetate/hexane) to give 0.600 g of **52b** (60%) as a white solid. MS-ESI m/z 180 $[MH]^+$.

15

Step 52c: Preparation of compound 52c

4-Fluorobenzylamine (0.440 g, 3.519 mmol, 2 eq) and **52b** (0.315 g, 1.759 mmol, 1 eq) were heated neat at 80 °C for 30 min. The crude product was purified by silica gel (100% ethyl acetate) to give 0.350 g of **52c** (%) as a white solid. MS-ESI m/z 305 $[MH]^+$.

20


Step 52d: Preparation of compound 52d

tert-Butyl(chloro)diphenylsilane (0.380 g, 1.382 mmol, 1.2 eq) was added to imidazole (0.235 g, 3.45 mmol, 3 eq) in dichloromethane (30.0 mL) . The reaction mixture was stirred at room temperature 10 min. then a solution of **52c** (0.350 g,

25


1.151 mmol, 1 eq) in dichloromethane (5.0 mL) was added. The reaction was stirred at room temperature 2 hrs. Water was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated under reduced pressure. The crude product was purified by silica gel (70% ethyl acetate/hexane) to give 0.387 g of **52d** (62%) as a white solid. MS-ESI m/z 543 $[MH]^+$.

Step 52e: Preparation of compound 52e




3-chloroperbenzoic acid (0.222 g, 1.285 mmol, 1.8 eq) at room temperature was added to **52d** (0.387 g, 0.714 mmol, 1 eq) in dichloromethane (30.0 mL) was added. The reaction mixture was stirred 1 hour at room temperature. A 1M solution of potassium carbonate was added and reaction mixture was extracted with dichloromethane. Organic phases were combined, dried over magnesium sulfate and concentrated to give 0.398 g of **52e** (100%) as a white solid. MS-ESI m/z 559 $[MH]^+$.


Step 52f: Preparation of compound 52f



Trifluoroacetic anhydride (0.450 g, 2.142 mmol, 3 eq) at room temperature was added to **52e** (0.398 g, 0.714 mmol, 1 eq) in dichloromethane (30.0 mL). The reaction mixture was stirred overnight at room temperature. A 1M solution of potassium carbonate was added and reaction mixture was extracted with ethyl acetate. Organic phases were combined, dried over magnesium sulfate and concentrated. The crude product was purified by silica gel (60% ethyl acetate) to give 0.209 g of **52f** (53%) as a white solid. MS-ESI m/z 559 $[MH]^+$.

Step 52g: Preparation of compound 52g

Activated manganese oxide (0.164 g, 1.88 mmol, 5 eq) at room temperature was added to **52f** (0.210 g, 0.376 mmol, 1 eq) in chloroform (30.0 mL). The reaction mixture was stirred at reflux 5hr. Solvent was concentrated under vacuo and crude reaction mixture was dissolved in methanol (20.0 mL). A solution of sodium cyanide (0.022g, 0.451 mmol, 1.2 eq) in methanol (3.0 mL) was added and the reaction mixture was stirred 1hr at room temperature. The reaction mixture was filtered on Celite. Organic phase was washed with water and extracted with ethyl acetate. The crude product was purified by silica gel (30% ethyl acetate/hexane) to give 0.130 g of **52g** (59%) as a white solid. MS-ESI m/z 587 $[MH]^+$.

Step 52h: Preparation of compound 52h


Lithium bis(trimethylsilyl)amide (1.2 mL of a 1M solution in tetrahydrofuran, 1.135 mmol, 5 eq) at -78°C was added to benzylhydroxylamine hydrochloride (0.041 g, 0.249 mmol, 1.1 eq) in tetrahydrofuran (20.0 mL). The reaction mixture was stirred 10 min. then a solution of **52g** (0.133 g, 0.227 mmol, 1 eq) in tetrahydrofuran (3.0 mL) was added. The reaction mixture was stirred at -78°C 30 min. before a solution of saturated ammonium chloride was added. This reaction mixture was extracted with ethyl acetate and organic phases were combined and concentrated under reduced pressure to give 0.140 g of crude product **52h** (100%) as a white solid. MS-ESI m/z 678 $[MH]^+$.

Step 52i: Preparation of Product 52

A tetrabutylammonium fluoride solution (0.620 mL of a solution 1M in tetrahydrofuran, 0.620 mmol, 3 eq) was added to a solution of **52h** (0.140 g, 0.207 mmol, 1 eq) in tetrahydrofuran (10.0 mL) was added. The reaction mixture was stirred at room temperature 1 hr. and concentrated under vacuo. The crude product


was purified by silica gel (70% ethyl acetate/hexane) to give 0.062 g of **52** (62%) as a white solid. MS-ESI m/z 440 $[MH]^+$.

Example 53: Preparation of 4-Hydroxymethyl-3-methoxy-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide (Product 53)



Compound **52** (0.060 g, 0.137 mmol, 1 eq) and 10% Pd/C (5 mg) in methanol (4.0 mL) were stirred under an atmosphere of hydrogen 1 hour. The catalyst was filtered and reaction mixture was concentrated under vacuum to give 0.040 g of **53** (85%) as a white solid. MS-ESI m/z 350 $[MH]^+$.


Example 54: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(hydroxy-methyl-amide) (Product 54)



Product **54** was prepared using the procedure described in example 2 and using methoxyamine. MS-ESI m/z 350 $[MH]^+$.


MS-ESI m/z 350 $[MH]^+$.

Example 55: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-dibenzylamide 2-hydroxyamide (Product 55)




Compound **55** was synthesized in a manner similar to example **2** using dibenzylamine. MS-ESI m/z 408 $[MH]^+$.

Example 56: 5-{{1-(4-Fluoro-phenyl)-cyclopropylamino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 56)




Product 56 was synthesized in a manner analogous to that in example **9** using 1-(4-Fluoro-phenyl)-cyclopropylamine. MS-ESI m/z 354 $[MH]^+$.

Example 57: Preparation of 5-{{1-(4-Fluoro-phenyl)-cyclopropylamino]-methyl}-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide (Product 57)



Product 57 was synthesized in a manner analogous to that in example **48** using 1-(4-Fluoro-phenyl)-cyclopropylamine. MS-ESI m/z 354 $[MH]^+$.


Example 58: Preparation of 5-(4-Fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide



Product 58 was synthesized in a manner similar to that in example 20 using 4-fluorobenzyl chloride and methoxyamine.

MS-ESI m/z 337 $[MH]^+$.


5 **Example 59: Preparation of 2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-2-methylcarbamoyl-benzylamide) 8-hydroxyamide (Product 59)**



10 Product 59 was synthesized in a manner similar to example 48 using 4-fluoro-2-methylcarbamoyl-benzylamine.

MS-ESI m/z 433 $[MH]^+$.

Example 60: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-(4-methyl-benzylamide) (Product 60)




15 Product 60 was synthesized in a manner similar to example 2 using 4-methyl-benzylamine.

MS-ESI m/z 332 $[MH]^+$.

20


25

Example 61: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-{{2-(4-fluoro-phenyl)-ethyl}-amide} 2-hydroxyamide (Product 61)




5 **Product 61** was synthesized in a manner similar to example 2 using 2-(4-fluoro-phenyl)-ethylamine MS-ESI m/z 350 $[MH]^+$.

Example 62: Preparation of 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(2,4-difluoro-benzylamide) 2-hydroxyamide (Product 62)



10 **Product 62** was synthesized in a manner similar to example 2 using 2,4-difluoro-benzylamine MS-ESI m/z 354 $[MH]^+$.


Example 63: Preparation of (rac)- {2-(4-Chloro-phenyl)-1-[(4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamoyl]-ethyl}-carbamic acid methyl ester (Product 63)



Product 63 was obtained from the reaction of 3-(4-Chloro-phenyl)-2-methoxycarbonylamino-propionic acid chloride and compound obtained from example 9 step **9a** , followed by catalytic hydrogenation Over Pd/C 5% in ethyl acetate. MS-ESI m/z 561[MH]⁺.


5

Example 64: Preparation of (rac) 5-[[[4-Fluoro-benzyl)-(2-phenyl-cyclopropanecarbonyl)-amino]-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Compound 64)



Product 64 was obtained from the reaction of 2-phenyl-cyclopropanecarbonyl chloride and compound obtained from example 9, step **9a** , followed by catalytic hydrogenation over Pd/C 5% in ethyl acetate. MS-ESI m/z 466[MH]⁺.


Example 65: Preparation of (4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid methyl ester (Product 65)



Product 65 was obtained from the reaction of methyl chloroformate and compound obtained from example **9a** , followed by catalytic hydrogenation over Pd/C 5% in ethyl acetate. MS-ESI m/z 380[MH]⁺.


20

Example 66: Preparation of (3-Chloro-4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester (Product 66)




Product 66 was obtained in a similar manner as that of **example 65**, using 4-fluoro-3-Chlorobenzyl amine in the initial step and benzyl chloroformate as the acylating agent. MS-ESI m/z 490[MH]⁺.

Example 67: Preparation of 5-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide ((Product 67)




Product 67 was synthesized by reaction of 5-formyl (0.751 g, 3 mmol) and 1000 mg (3.5 mmol) (4-fluorobenzyl) triphenylphosphonium bromide in THF at -78°C then stirred overnight at R.T. The reaction was quenched with NH₄Cl (sat). The crude product was separated on silica gel to yield 417 mg (31%). This product was hydrogenated by catalytic hydrogenation over Pd/C 5% in EtOAc. Silica gel chromatography purification yielded 305 mg, which was reacted with hydroxylamine (excess) in ethanol (2h, 80°C). The compound was precipitated with 1N HCl, filtered and redissolved in MeOH with the addition of 6N HCl, stirred for 4hr, followed by evaporation; providing the desired product. MS-ESI m/z 307 [MH]⁺.

Example 68: Preparation of 3-Hydroxy-4-hydroxymethyl-5-(3-phenyl-propyl)-pyridine-2-carboxylic acid hydroxyamide ((Product 68)



Product 68 was synthesized in a manner similar to example 68 using
 5 (phenylacetylene lithium salt in the initial step. Catalytic hydrogenation (Pd/C 5% in EtOAc) was done in the presence of 3 eq of acetic anhydride, 1 atm 24h. The subsequent steps were as per example 68. MS-ESI m/z 303 [MH]⁺.


Example 69: Preparation of 5-Benzenesulfonylmethyl-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide ((Product 69)



550 mg of **Compound XI**, of scheme 3, was dissolved in 30 mL DCM was reacted with 2.5 eq of methane sulfonyl chloride in the presence of 5 eq triethyl amine. Extraction vs 5% citric acid, drying over Na₂SO₄, and evaporation yielded 500mg of
 15 desired mesylate. This mesylate was immediately reacted with 400 mg of benzene sulfinic acid in DMF 2mL. the product was isolated by precipitation in water and filtration. The crude was dissolved in CHCl₃ 30mL and 400 mg mCPBA (70%) was added. After 1 h stirring, the reaction was extracted using K₂CO₃ and the organic phase dried over CaCO₃ the evaporated. The residue was dissolved in 3 mL DCM
 20 and 3mL trifluoroacetic anhydride was added. Stirring at reflux 45°C for 20h affording the rearranged product, isolated through evaporation of solvent. The residue was then added to a solution of MnO₂ 2g in CHCl₃ (30ml) and stirred at reflux 1h. filtration and evaporation afforded the aldehyde (250mg) This was placed in 10mL MeOH with 1.2 eq I₂ and 3 eq KOTMS. Stirring at R.T. for 1h gave the ester
 25 in quantitative conversion. The product was purified on Silica gel. 100 mg of the ester was reacted with excess (hydroxylamine 50% aq) in pyridine to give the


hydroxamate. Dilution in EtOAc and extraction vs 5% citric acid gave the desired intermediate. The final product was obtained by adding 50 mg of the above acetonide to neat 70% formic acid. After 15 min the reaction is complete, the formic acid is evaporated off and the residue triturated with water to give **69** as a white powder. MS-ESI m/z 339 $[MH]^+$.

Example 70: Preparation of 5-(4-Fluoro-phenylmethanesulfonylmethyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxamide (Product 70)




550 mg of **Compound XI**, of scheme 3, was dissolved in 30 mL DCM was reacted with 2.5 eq of methane sulfonyl chloride in the presence of 5 eq triethyl amine. Extraction vs 5% citric acid, drying over Na_2SO_4 , and evaporation yielded 500mg of desired mesylate. This was immediately reacted with 400 mg of 4 fluoro benzyl mercaptan in DMF 2mL. the product was isolated by precipitation in water and filtration. The crude(0.5g) was dissolved in $CHCl_3$ 30mL and 1000 mg mCPBA (70%) was added. After 0.5 h stirring, the reaction was extracted using K_2CO_3 and the organic phase dried over $CaCO_3$ the evaporated. The residue was dissolved in 3 mL DCM and 3mL trifluoroacetic anhydride was added. Stirring at reflux 45°C for 20h affording the rearranged product, isolated through evaporation of solvent. The residue was then added to a solution of MnO_2 2g in $CHCl_3$ (30ml) and stirred at reflux 1h. filtration and evaporation afforded the aldehyde (300mg). This was placed in 10mL meOH with 1.2 eq NaCN and 0.5g MnO_2 . Stirring at R.T. for 1h gave the ester in quantitative conversion. The product was purified on Silica gel. 98 mg of the ester was reacted with excess (hydroxylamine 50% aq) in pyridine to give the hydroxamate. Dilution in EtOAc and extraction vs 5% citric acid gave the desired compound **70**. MS-ESI m/z 411 $[MH]^+$.

Example 71: Preparation of 5-(4-Fluorophenylmethanesulfonylmethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide (Product 71)




Product 71 was obtained by adding 50 mg of compound from example 70 to neat 70% formic acid. After 15 min the reaction is complete, the formic acid is evaporated off and the residue triturated with water to give 71 as a white powder. MS-ESI m/z 371 $[MH]^+$.

Example 72: Preparation of (4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester (Product 72)




Step 72a: Preparation of compound 72a



The product from example 20e of methyl 5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate (3.1 g, 8.3mmol) was dissolved in 70% formic acid and left at RT 1h. Evaporation of the acid gave a crude off-white powder (2.77g, 100%). 1.95g (5.9mmol) was reacted in 30ml DCM with 1.7ml Acetic anhydride, 2.5ml TEA and 36mg DMAP for 3h. Evaporation and Silica gel chromatography (EtOAc) of the crude gave 2.44g (94%). 2.3g was dissolved in 30mL THF containing 450mg 10% Pd/C and H_2 bubbled through for 3h. Filtration


and evaporation of the solvent yielded a residue used without further purification. This residue was dissolved in DCM (20ml) and 5mL of trifluoroacetic acid was added with a few drops of tripropylsilane. Evaporation yielded a crude product used without further purification. The crude product (1.31g 5.5mmol) was dissolved in 25 mL EtOAc and 1.85g IBX added. The mixture was refluxed 3h, cooled, filtered washed with EtOAc and the solvents removed. Purification by silica gel chromatography (EtOAc) gave 0.96g (72%) of the desired product.

Step 72b: Preparation of 5-[(4-Fluoro-benzylamino)-methyl]-3-hydroxy-4-methyl-pyridine-2-carboxylic acid methyl ester (compound 72b)




65mg of the aldehyde **72a** was dissolved in MeOH and 50mg 4-Fluorobenzyl amine was added followed by 20 mg NaCNBH₃. The mixture was stirred 2h and concentrated. The residue was purified by silica gel (EtOAc) to give 87 mg product. This was reacted with 1.25 eq of benzylchloroformate, 2 eq TEA in DCM to yield the desired carbamate. Extraction vs 5% citric acid and evaporation of the organic phase yielded the crude product. The crude (17mg) was dissolved in 2ml THF and 1mL (50% hydroxylamine aq) was added, heated for 1h at 60°C. The solvent was then evaporated and addition of 1 ml 10% citric acid gave a precipitate which was filtered and washed with water. Yield 11mg 70%. MS-ESI *m/z* 440 [MH]⁺.

Example 73: (4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid tert-butyl ester (Product 73)




Product **73** was synthesized using a similar procedure as **example 72** where di tert-butylpyrocarbonate replaces the benzylchloroformate. MS-ESI *m/z* 406 [MH]⁺.

Example 74: Preparation of 3-Hydroxy-4-methyl-pyridine-2,5-dicarboxylic acid 5-(3-chloro-4-fluoro-benzylamide) 2-hydroxyamide



Step 74a: Preparation of compound 74a



5

The compound from example 72 step a (608 mg, 2.6mmol) was dissolved in acetone (15ml). 2-methylbutene (2.0M in THF) 12mL was added followed by a solution of 1.85g Sodium Chlorite (NaClO_2) and 2.83g NaH_2PO_4 in 15 mL H_2O . After 20 min at RT the solution was concentrated to remove the organic solvent and the pH adjusted to ~4 with Citric acid. The resulting white precipitate was filtered and washed with water. Drying gave 424mg of desired product.

10


Step 74b: Preparation of compound 74b

60mg of the above acid was reacted with 30mg (3-chloro-4-fluorobenzyl amine) 75mg HBTU and 100 mg N-methylmorpholine in 2mL DMF. After 1h the solution was diluted in EtOAc and extracted vs 5% citric acid. The organic phase was evaporated to afford the crude product. This was dissolved in 1mL pyridine and 0.5 mL 50% hydroxylamine in water was added. After 1h at 45°C the pyridine/water azeotrope was removed under vacuum and the residue triturated with 10% citric acid. The precipitate was filtered and washed to give the desired product. MS-ESI m/z 354 $[\text{MH}]^+$.

15

20

Example 75: Preparation of 5-(4-Fluoro-benzylloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide (Product 75)



Product 75 was prepared in a procedure analogous to that shown in example 20 with the exception of the use of 4-fluoro -benzyl chloride as alkylating agent, and the use of methoxylamine instead of hydroxylamine.

LC-MS (M+H)⁺ m/z 337.

5

Table 3: Listing of Compounds in Examples

Example, step	Chemical Name
1	N ² , 3-bis(benzyloxy)-N ⁵ -(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
1, step 1a	N, 9-bis(benzyloxy)-3,3-dimethyl-1,5-dihydro-[1,3]dioxepino[5,6-c]pyridine-8-carboxamide
1, step 1b	N, 3-bis(benzyloxy)-4,5-bis(hydroxymethyl) picolinamide
1, step 1c	N, 7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide
1, step 1d	N ² , 3-bis(benzyloxy)-N ⁵ -(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
2	N ⁵ -(4-fluorobenzyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
3	N ⁵ -(benzo[d][1,3]dioxol-5-ylmethyl)-N ² ,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
4	N ⁵ -(benzo[d][1,3]dioxol-5-ylmethyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
5	N ² , 3-bis(benzyloxy)-4-(hydroxymethyl)-N ⁵ -(4-methoxybenzyl)pyridine-2,5-dicarboxamide
6	N ² , 3-dihydroxy-4-(hydroxymethyl)-N ⁵ -(4-methoxybenzyl)pyridine-2,5-dicarboxamide
7	N ² , 3-bis(benzyloxy)-N ⁵ -(3,5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
8	N ⁵ -(3, 5-difluorobenzyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide
9	5-((4-fluorobenzylamino)methyl)-N, 3-dihydroxy-4-(methoxymethyl)picolinamide
9, step 9a	N, 3-bis(benzyloxy)-5-((4-fluorobenzylamino)methyl)-4-(hydroxymethyl)picolinamide (compound 9a)
10	5-((3, 5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide
10, step 10a	5-((3,5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl) picolinamide
11	5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide
1, step 11a	5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N, 3-bis(benzyloxy)-4-(hydroxymethyl)picolinamide
12	N ⁵ -(4-fluorobenzyl)-N ² , 3-dihydroxy-4-(methoxymethyl)-N ⁵ -methylpyridine-2, 5-dicarboxamide
12, step 12a	5-(benzyloxy)-N-(4-fluorobenzyl)-4-(hydroxymethyl)-6-methylnicotinamide
12, step 12b	5-(benzyloxy)-N-(4-fluorobenzyl)-4-(methoxymethyl)-N, 6-dimethylnicotinamide
12, step 12c	3-(benzyloxy)-5-((4-fluorobenzyl)(methyl)carbamoyl)-4-(methoxymethyl)-2-methylpyridine 1-oxide
12, step 12d	5-(benzyloxy)-N-(4-fluorobenzyl)-6-(hydroxymethyl)-4-(methoxymethyl)-N-methylnicotinamide
12, step 12e	5-(benzyloxy)-N-(4-fluorobenzyl)-6-formyl-4-(methoxymethyl)-N-methylnicotinamide
12, step 12f	methyl 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl) carbamoyl)-4-(methoxymethyl)picolinate
12, step 12g	N ² , 3-bis(benzyloxy)-N ⁵ -(4-fluorobenzyl)-4-(methoxymethyl)-N ⁵ -methylpyridine-2, 5-dicarboxamide
12, step 12h	N ⁵ -(4-fluorobenzyl)-N ² , 3-dihydroxy-4-(methoxymethyl)-N ⁵ -methylpyridine-2, 5-dicarboxamide
13	N ² , 3-bis(benzyloxy)-N ⁵ -(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide
14	N ⁵ -(3-chloro-4-fluorobenzyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide
15	N ² , 3-bis(benzyloxy)-N ⁵ -(3, 4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide
16	N ⁵ -(3, 4-dichlorobenzyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide
17	N, 3-Dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide
17, step 17a	5-((4-Methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine
17, step 17b	5-((4-Methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine 7-oxide
17, step 17c	5-((4-Methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridin-8-yl)methanol
17, step 17d	5-((4-Methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carbaldehyde
17, step 17e	ethyl 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxylate
17, step 17f	ethyl 3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinate
17, step 17g	N, 3-Dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinamide
18	5-(Benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide
19	N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide
19, step 19a	5-((4-Methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridine-8-carboxylic acid
19, step 19b	N-(Benzyloxy)-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide
19, step 19c	N-(Benzyloxy)-3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl) picolinamide
19, step 19d	N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide
20	N ⁵ -(3,4-difluorobenzyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide
21	5-[(4-Fluoro-phenylamino)-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
21 step a	methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate

21 Step b	5-((4-fluorophenylamino)methyl)-3-hydroxy-4-(hydroxymethyl) picolinate
22	5-([2-(4-Fluoro-phenyl)-ethylamino]-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
22 Step 22b	Methyl 5-((4-fluorophenethylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate
23	5-(4-Fluoro-benzoylamino)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
23 Step 23a	8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid
23 Step 23b	Methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate
23 Step 23c	5-Amino-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid methyl ester
23 Step 23d	Methyl 5-(4-fluorobenzamido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate
23 Step 23e	Methyl 5-(4-fluorobenzamido)-3-hydroxy-4-(hydroxymethyl)picolinate
24	(8-Hydroxycarbamoyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl)-carbamic acid benzyl ester
25	5-([Benzyl-(4-fluoro-phenyl)-amino]-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
25 Step 25a	Methyl 5-((benzyl(4-fluorophenyl)amino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate
26	5-((2-Benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino)-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
26 Step 26b	Methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate
26 Step 26c	5-(((2-Benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino)-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
27	5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
27 Step 27b	Methyl 5-(3-(4-fluorophenyl)ureido)-3-hydroxy-4-(hydroxymethyl)picolinate
27 Step 27c	5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
28	5-(4-Fluoro-phenoxy)methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
28 Step 28 a	Methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate
28 Step 28b	Methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate
29	5-(3-Chloro-4-fluoro-phenoxy)methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
29 Step 29a	Methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate
29 Step 29b	Methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate
30	5-(3-Chloro-4-fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
31	5-[2-(4-Fluoro-phenyl)-ethoxymethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
32	5-(2,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
33	5-(3,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
34	5-(4-Fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
35	5-(4-Fluoro-phenoxy)methyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide
35, Step 35a	Methyl 3-(benzyloxy)-5-((4-fluorophenoxy)methyl)-4-methylpicolinate
35, Step 35b	Methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate
36	5-(3-Chloro-4-fluoro-phenoxy)methyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide
36, Step 36 a	Methyl 3-(benzyloxy)-5-((3-chloro-4-fluorophenoxy)methyl)-4-methylpicolinate
36, Step 36b	Methyl 5-((3-chloro-4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate
37	5-(2,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide
38	5-(3,4-Difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide
39	5-(4-Fluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide
40	5-Benzyloxymethyl-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide
41	(S)-(-)-3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(1-phenyl-ethyl)-amide]
42	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(pyridin-2-ylmethyl)-amide]
43	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-benzylamide 2-hydroxyamide
44	(S)-(-)-3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(2-hydroxy-1-phenyl-ethyl)-amide]
45	Pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide
46	2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-[[1-(4-fluoro-phenyl)-cyclopropyl]-amide] 8-hydroxyamide
47	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-[[1-(4-fluoro-phenyl)-cyclopropyl]-amide] 2-hydroxyamide
48	2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-benzylamide) 8-(methoxy-amide)
49	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(methoxy-amide)
50	2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-cyclohexylmethyl-amide 8-hydroxyamide
51	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide
52	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide
53	4-Hydroxymethyl-3-methoxy-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide

54-	- 3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(hydroxy-methyl-amide)
55	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-dibenzylamide 2-hydroxyamide
56	5-[[1-(4-Fluoro-phenyl)-cyclopropylamino]-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
57	5-[[1-(4-Fluoro-phenyl)-cyclopropylamino]-methyl]-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide
58	5-(4-Fluoro-benzoyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide
59	2,2-Dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-2-methylcarbamoyl-benzylamide) 8-hydroxyamide
60	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-(4-methyl-benzylamide)
61	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-[[2-(4-fluoro-phenyl)-ethyl]-amide] 2-hydroxyamide
62	3-Hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(2,4-difluoro-benzylamide) 2-hydroxyamide
63	(rac)- {2-(4-Chloro-phenyl)-1-[(4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamoyl]-ethyl}-carbamic acid methyl ester
64	(rac) 5-[[4-(4-Fluoro-benzyl)-(2-phenyl-cyclopropanecarbonyl)-amino]-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
65	(4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid methyl ester
66	(3-Chloro-4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester
67	5-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
68	3-Hydroxy-4-hydroxymethyl-5-(3-phenyl-propyl)-pyridine-2-carboxylic acid hydroxyamide
69	5-Benzenesulfonylmethyl-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
70	5-(4-Fluoro-phenylmethanesulfonylmethyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide
71	5-(4-Fluoro-phenylmethanesulfonylmethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide
72	(4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester
72 Step 72b	5-[[4-Fluoro-benzylamino]-methyl]-3-hydroxy-4-methyl-pyridine-2-carboxylic acid methyl ester
73	(4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid tert-butyl ester
74	3-Hydroxy-4-methyl-pyridine-2,5-dicarboxylic acid 5-(3-chloro-4-fluoro-benzylamide) 2-hydroxyamide
75	5-(4-Fluoro-benzoyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide

Example 76: Biological Evaluation, *In Vitro* Integrase Inhibition Assay

IC₅₀ was determined for the compounds of the inventions based on data generated in strand transfer assays. The IC₅₀ is a measure of the ability of the compounds tested to inhibit the integration of 3'-processed oligonucleotides by recombinant HIV-1 integrase.

Strand transfer assays were performed essentially as described in Hazuda, D. J.; Felock, P.; Hastings, J. C.; Pramanik, B.; Wolfe, A. J. *Viol.* 1997, 71, 7005-7011). Donor DNA (1.5 pmol/well), biotinylated on the 5' end of the strand processed by integrase, was immobilized onto streptavidin-coated microtiter plates. Recombinant integrase (250 ng/well) was assembled onto the immobilized donor oligonucleotide in reaction buffer (20 mM Hepes, pH 7.6, 5 mM B-mercaptoethanol, 50 ug/mL bovine serum albumin) containing 30 mM MnCl₂. Excess enzyme was removed, and the

complexes were washed extensively prior to the addition of the target DNA substrate. The target DNA (0.75 pmoles/well) substrate was labeled on each 3' end with FITC. After strand transfer, the FITC-labeled products were detected using an anti-FITC antibody conjugated with alkaline phosphatase (Roche) and a chemiluminescent substrate (Tropix CSPD with Sapphire II enhancer, Applied Biosystems). The assay was performed in a final concentration of 10% DMSO. To specifically evaluate inhibition of strand transfer, compounds were added after assembly, just prior to the addition of the target DNA.

The results of the integrase strand transfer assay are reported as IC₅₀ values. IC₅₀ values were determined using a sigmoidal dose-response equation. The formula used for calculating % inhibition was: % Inhibition = [1-(sample counts/average of positive control)]*100. The percent inhibition of HIV-1 integrase activity was graphed against the log of the compound concentration (M). Using GraphPad Prism or ActivityBase (IDBS) software IC₅₀ was determined using following sigmoidal dose-response equation:

$$Y = (A + ((B - A) / (1 + ((C / X)^D))))$$

Where A is the lower plateau (~0%), B is the higher plateau (~100%), C is the IC₅₀, D is the slope, X is the compound concentration (M), and Y is the % inhibition.

Inhibition of strand transfer, as determined by their IC₅₀, demonstrates that the compounds of the present invention inhibit HIV integrase and have IC₅₀s similar to that of Raltegravir, a marketed HIV integrase inhibitor, and L-708906, an integrase inhibitor currently in clinical development.

Compound	ST IC ₅₀
Raltegravir (MK-0518)	0.065
L-708906 (Merck)	0.045
Compound of Example 14	0.027
Compound of Example 2	0.303
Compound of Example 18	0.088

Example 77: Antiviral Efficacy

The antiviral efficacies of the integrase inhibitor compounds of the invention were evaluated based on EC₅₀ measures obtained from two different *in vitro* HIV infection assay using cultured MT-4 cells: (1) a multi-cycle infection where cells were infected with wild type NL-4.3) and (2) a single-cycle infection where the cells were infected with a luciferase-bearing, envelope defective (env-) NL-4.3 virus pseudotyped with HIV-1 env (HXBc2).

The incubation period for the multi-cycle infection assay was 6 days. Cell viability (cytoprotection) and EC₅₀ were determined using the colorimetric MTT assay (A. J. Japour et al, Antimicrobial Agents and Chemotherapy, 37, 1095-1101, 1993 and R. Pauwels et al. Journal of Virological Methods, 20, 309-321, 1988).

The incubation period for the single-cycle infection assay was 48 hours. EC₅₀ was determined, as described by Chen et al., Journal of Virology, Feb. 1994, Vol. 68, No. 2, p. 654-660, based on measures of luciferase signal over a range of drug concentrations.

The results of these assays are shown in Table 3 and integrase inhibitors of the invention were prepared using the synthetic methods described in Schemes 1-15; and the examples described herein. The reference numbers of the compounds listed in Table 4 (Ex. No.) correspond to the example numbers of examples 1 to 75 described above. These data demonstrate the antiviral efficacy of the compounds of the invention as integrase inhibitors and for treatment of HIV infection and AIDS. The compounds tested display potent antiviral activity. Furthermore, similar antiviral activity was observed when the HIV-1 envelope was replaced with VSV-G, validating that the compounds of the invention are post-entry inhibitors.

Table 4: Results of Cytoprotection-Cytotoxicity Assay

Ex. No.	Compound	EC ₅₀ (nM) multi-cycle
----	L-708906 (a integrase inhibitor currently in development at Merck)	5800
----	MK-0518 (Raltegravir, a marketed integrase inhibitor, brand name Isentress™)	20
1	N ² , 3-bis(benzyloxy)-N ⁵ -(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	>10000
2	N ² -(4-fluorobenzyl)-N ² ,3-dihydroxy-4-	358

	(hydroxymethyl)pyridine-2, 5-dicarboxamide	
<u>3</u>	N ⁵ -(benzo[d][1, 3]dioxol-5-ylmethyl)-N ² ,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	>10000
<u>4</u>	N ⁵ -(benzo[d][1, 3]dioxol-5-ylmethyl)-N ² ,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide	>10000
<u>5</u>	N ² , 3-bis(benzyloxy)-4-(hydroxymethyl)-N ⁵ -(4-methoxybenzyl)pyridine-2,5-dicarboxamide	900
<u>6</u>	N ² , 3-dihydroxy-4-(hydroxymethyl)-N ⁵ -(4-methoxybenzyl)pyridine-2, 5-dicarboxamide	1000
<u>7</u>	N ² , 3-bis(benzyloxy)-N ⁵ -(3, 5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	1500
<u>8</u>	N ⁵ -(3, 5-difluorobenzyl)-N ² , 3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	>10000
<u>9</u>	5-((4-fluorobenzylamino)methyl)-N, 3-dihydroxy-4-(methoxymethyl)picolinamide	1000
<u>10</u>	5-((3, 5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide	>10000
<u>11</u>	5-((benzo[d][1, 3]dioxol-5-ylmethylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide	300
<u>12</u>	N ⁵ -(4-fluorobenzyl)-N ² , 3-dihydroxy-4-(methoxymethyl)-N5-methylpyridine-2, 5-dicarboxamide	14000
<u>13</u>	N ² , 3-bis(benzyloxy)-N ⁵ -(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	>10000
<u>14</u>	N ⁵ -(3-chloro-4-fluorobenzyl)-N ² , 3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	123
<u>15</u>	N ² , 3-bis(benzyloxy)-N ⁵ -(3, 4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	>10000
<u>16</u>	N ⁵ -(3, 4-dichlorobenzyl)-N ² , 3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide	700
<u>17</u>	N, 3-Dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide	2000
<u>18</u>	5-(Benzyloxymethyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide	86
<u>19</u>	N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide	5058
<u>20</u>	N ² -(3, 4-difluorobenzyl)-N ² , 3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide	30
<u>21</u>	5-[(4-Fluoro-phenylamino)-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	4.3
<u>23</u>	5-(4-Fluoro-benzoylamino)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	6300
<u>25</u>	5-{{[Benzyl-(4-fluoro-phenyl)-amino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	175
<u>27</u>	5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	1096
<u>28</u>	5-(4-Fluoro-phenoxy)methyl-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	186
<u>65</u>	(4-Fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid methyl ester	>10000
<u>67</u>	5-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	235
<u>71</u>	5-(4-Fluoro-phenylmethanesulfonylmethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide	4.3
<u>74</u>	3-Hydroxy-4-methyl-pyridine-2,5-dicarboxylic acid 5-(3-chloro-4-fluoro-benzylamide) 2-hydroxyamide	267

Example 78: Inhibition of HIV-1 Clinical Isolates in PBMCs

Acute infection assays using fresh human phytohemagglutinin (PHA)-stimulated peripheral blood mononuclear cells (PBMCs) were carried out. PBMCs were stimulated with IL-2 and infected with one of two wild-type drug-sensitive clinical HIV-1 isolates. The strains HIV-1 used for the infections, 91US005 (primary R5 strain of HIV-1) and 94US33931N, are both Group M, Subtype B viruses. Antiviral activity was determined as a reduction in supernatant reverse transcriptase (RT) activity after a 6 day incubation.

Compounds of the invention N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide (product of example 2) and of N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N²,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide (product of example 3) and Retrovir (AZT), a reverse transcriptase inhibitor, were tested at 9 concentrations and the EC₅₀ of each compound was determined. These data demonstrate that the compounds of the invention inhibit infection of PBMCs by HIV-1 isolates.

The results of these assays are provided in Table 5 below.

Table 5: Results of assays in PMBCs

Compound	HIV-1 isolate	EC ₅₀ (uM)
Example 2	91US005	0.076
Example 2	94US33931N	0.123
Example 3	91US005	0.093
Example 3	94US33931N	0.03
Retrovir (AZT)	91US005	0.006
Retrovir (AZT)	94US33931N	0.003

Example 79: Effect of Protein Binding on Antiviral Activity

Protein binding was tested by Rapid Equilibrium Dialysis (RED) method (The RED (Rapid equilibrium Dialysis) Device inserts instruction manual. Pierce, Rockford IL). Compounds were spiked in human serum (HS) or 10% FBS-RPMI at a concentration of 1 uM. The red chamber was loaded with 0.3 mL of samples containing compounds. The white chamber contained 0.5 mL of D-PBS buffer only. The plate was incubated at 37°C while shaking at 150 rpm for 5 hours. Aliquots from both chambers were analyzed by LC/MS/MS. The multi-cycle antiviral activity in the

absence and presence of human serum was determined by p24 ELISA measurement after 6 days of NL4.3 virus infection. The results, showing a moderate effect of protein binding on antiviral activity, are provided in Table 6 below.

5 **Table 6: Effect of Protein Binding on Antiviral Activity**

Compound	% Protein Binding		EC ₅₀ (uM)		
Protein Spike	10% FBS	100% HS	10% FBS	10% FBS + 40% HS	Fold-change with HS
Raltegravir	34.7	86.7	0.047	0.064	1.4
Example 1	52.5	89	0.227	0.481	2.1
Example 2	31.9	84.5	na	na	na

Example 80: Human CYP Inhibition

For the human CYP inhibition experiments, the incubation medium contained 0.3 mg/mL human liver microsomes, 100 mM phosphate buffer pH 7.4, 5 mM MgCl₂, 1 mM EDTA. The respective substrates for CYP3A4, CYP2C9 and CYP2D6 were added at various concentrations in the presence or absence of the compounds tested. After pre-incubation, the reaction was initiated by the addition of NADPH at final concentration 1 mM. Incubation times were 10, 60 and 15 minutes for CYP3A4, CYP2C9 and CYP2D6, respectively (Walsky RL and Obach RS., Drug Metabolism and Disposition, 2004, 32, 647-660). GraphPad Prism software was used for data analysis. The results of these assays, showing the metabolic stability of the compounds of the invention are provided in Table 7 below.

Table 7: Results of CYP Inhibition Assays

Compound	CYP3A4		
	Vmax (pmol/min/mg)	Km (μM)	Ki (μM)
Example 1	3652	11.24	5.10
Example 2	1252	7.76	1.11E+16
Example 3	1247	8.30	586.80
	CYP2C9		
	Vmax (pmol/min/mg)	Km (μM)	Ki (μM)
Example 1	78.72	293.20	63.60
Example 2	NA	NA	NA
Example 3	NA	NA	NA
	CYP2D5		
	Vmax (pmol/min/mg)	Km (μM)	Ki (μM)

Example 1	NA	NA	NA
Example 2	32.16	5.24	1.84E+13
Example 3	36.02	5.79	4.87E+20

Example 81: Human Liver Microsome Stability

Human liver microsome stability was assayed by incubation of liver microsomes (0.6 mg/mL), in medium containing 100 mM phosphate buffer pH 7.4, 10 mM MgCl₂, 1 mM EDTA, 25 µg Alamethicin/mg protein, and 1 mM NADPH, 1 mM UDPGA, or 1 mM NADPH and 1 mM UDPGA. After pre-incubation, the assay was started by the addition of 1 µM of compound. Samples were taken after 0, 15, 30, 60, 90 and 120 minutes of incubation (Fisher MB, Drug Metabolism and Disposition, 2000, 28, 560-566). GraphPad Prism software was used for data analysis. The results, showing that the compounds of the invention are metabolically stable in Human liver microsomes, are provided in Table 7 below.

Table 7: Stability of Compounds in Human Liver Microsomes

Compound	Half-life in HLM (hrs)		
	<i>NADPH</i>	<i>UDPGA</i>	<i>NADPH/UDPGA</i>
Raltegravir	19.7	7.03	11
Compound 1	7.4	19.7	3.9
Compound 2	5	16	5.9
Compound 3	6.18	3.8	1.95

Example 82: Pharmacokinetic (PK) Profile in Rats

Female Sprague-Dawley rats were randomly selected and assigned to two groups. A group of 13 rats were administered 5 mg/kg of compound 2 intravenously. A second group of 8 rats were administered 50 mg/kg of compound 2 orally. Following the dosing, blood was collected at 7 different time points. Plasma samples obtained from the blood samples were analyzed by LC/MS/MS and the bioavailability of compound 2 was determined. Figure 1 shows the results of this experiment demonstrating the high adsorption and generally favourable profile of the compounds of the invention.

Other Embodiments


The examples, synthetic schemes and procedures provided herein are for the purpose of illustration only. They are not intended to be exhaustive or to limit the scope of the invention to the specific examples, synthetic schemes, and procedures described herein. Although the invention has been described with reference to several
5 embodiments, it will be understood by one of ordinary skill in the art that various modifications can be made without departing from the spirit and the scope of the invention, as set forth in the claims. Other embodiments are in the claims.

All patents, patent applications, and publications referenced herein, including
10 U.S. provisional application number 61/130,874, filed June 4, 2008, are hereby incorporated by reference.

What is claimed is:

Claims

1. A compound of formula I,



I

wherein:

A is a six membered carbocyclic or heterocyclic ring system;

R₁ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl), (O-C₁₋₆ branched alkyl), CO(R₉), COO(R₉), CON(R₉)(R_{9a}), or SO₂N(R₉)(R_{9a}), wherein said R₉ and R_{9a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; R₂ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl),

(O-C₁₋₆ branched alkyl), CO(R₁₀), COO(R₁₀), or CON(R₁₀)(R_{10a}), wherein said R₁₀ and R_{10a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; or R₁ and R₂ are ortho substituents that together form a carbocyclic or heterocyclic ring system;

L is -N(R')C(O)-; -C(O)N(R')-; -OC(O)-; -C(O)O-; -OC(O)N(R')-; -N(R')C(O)O-; -N(R')C(O)N(R')-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-;

$-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-$;
 $-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})-$; $-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-$;
 $-C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-$; $-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-$;
 $-C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})-$; $-C(R_{a1})(R_{a2})-Z-$; or $-Z-C(R_{a1})(R_{a2})-$; wherein each R_{a1} , R_{a2} , R_{b1} , R_{b2} , R_{c1} , and R_{c2} is, independently, selected from H, C_{1-6} alkyl, C_{1-6} fluoro-alkyl, hydroxy-alkyl, benzyl, phenyl and heterocycle, or, alternatively, one or more of R_{a1} and R_{a2} ; R_{b1} and R_{b2} ; and R_{c1} and R_{c2} combine to form a carbocyclic ring, and wherein Z is selected from $-N(R')C(O)-$; $-C(O)N(R')-$; $-OC(O)-$; $-C(O)O-$; $-OC(O)N(R')-$; $-N(R')C(O)O-$; $-N(R')C(O)N(R')-$; $-N(R')-$; $-SO_2-$; and $-O-$; wherein R' is selected from H, C_{1-6} alkyl, benzyl, SO_2R'' , and $C(O)R''$, $C(O)OR''$, and R'' is selected from C_{1-6} alkyl, C_{1-6} fluoro-alkyl, heteroalkyl, carbocyclic group, benzyl, phenyl and heterocycle;

B_1 is $-R_3$, CH_2OR_3 , $CH_2N(R_8)(R_{8a})$, $C(O)OR_3$ or $C(O)N(R_8)(R_{8a})$, wherein each of R_8 and R_{8a} is, independently, selected from the group consisting of H, C_{1-6} alkyl, C_{1-6} fluoro-alkyl, benzyl, phenyl and heterocycle;

B_2 is H or OR_5 ;

R_3 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; and R_5 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; or, alternatively, R_3 and R_5 combine to form a heterocyclic ring system; and

R_4 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; and pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein A is a phenyl ring, a pyridine ring, or a cyclohexyl ring.
3. A compound according to claim 1, wherein L is $-CH_2OCH_2-$, $-CH_2CH_2OCH_2-$, $-OCH_2-$, $-CH_2NHCH_2-$, $-C(cyclo-C_2H_4)NHCH_2-$, $-NHCH_2-$, $-CH_2CH_2NHCH_2-$, $-CH_2NHC(O)-$, $-CH_2N(CH_3)C(O)-$, $-CH(CH_2OH)NHC(O)-$, $-C(cyclo-C_2H_4)NHC(O)-$, $-CH_2CH_2NHC(O)-$, $-C(O)NH-$, $-CH_2OC(O)NH-$, $-NHC(O)NH-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2-$, $-SO_2CH_2-$, or $-CH_2SO_2CH_2-$.
4. A compound according to claim 1, wherein B_1 is H, CH_3 , CH_2OH , or CH_2OCH_3 .

5. A compound according to claim 1, wherein B₂ is H or -OR₅ and R₅ is H or benzyl.
6. A compound according to any of claims 2-5, wherein L is -CH₂NHCH₂- or -CH₂NHC(O)-.
7. A compound according to claim 6, wherein each of R₁ and R₂ is, independently, selected from halogen, -OCH₃, -OH, or R₁ and R₂ combine to form a cyclic acetal or cyclic ketal.
8. A compound according to claim 7 wherein, R₄ is -H or benzyl.
9. A compound according to any of claims 2-5, wherein L is -CH₂OCH₂-.
10. A compound according to claim 9, wherein each of R₁ and R₂ is, independently, selected from halogen, -OCH₃, -OH, or R₁ and R₂ combine to form a cyclic acetal or cyclic ketal.
11. A compound according to claim 9, wherein R₄ is -H or benzyl.
12. A compound according to any of claims 2-5, wherein L is -SO₂CH₂-, or -CH₂SO₂CH₂-.
13. A compound according to claim 12, wherein each of R₁ and R₂ is, independently, selected from halogen, -OCH₃, -OH, or R₁ and R₂ combine to form a cyclic acetal or cyclic ketal.
14. A compound according to claim 13, wherein R₄ is -H or benzyl.
15. A compound according to any of claims 3, 6, 10, or 12, wherein A is a phenyl ring.
16. A compound according to claim 1 selected from:
N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
N⁵-(4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N²,3-bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
N⁵-(benzo[d][1,3]dioxol-5-ylmethyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
N², 3-bis(benzyloxy)-4-(hydroxymethyl)-N⁵-(4-methoxybenzyl)pyridine-2,5-dicarboxamide;

N^2 , 3-dihydroxy-4-(hydroxymethyl)- N^5 -(4-methoxybenzyl)pyridine-2,5-dicarboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(3,5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(3, 5-difluorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
5-((4-fluorobenzylamino)methyl)-N, 3-dihydroxy-4-(methoxymethyl)picolinamide);
5-((3, 5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide;
5-((benzo[d][1, 3]dioxol-5-ylmethylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide;
 N^5 -(4-fluorobenzyl)- N^2 , 3-dihydroxy-4-(methoxymethyl)- N^5 -methylpyridine-2, 5-dicarboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N^5 -(3-chloro-4-fluorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(3, 4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N^5 -(3, 4-dichlorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
N, 3-dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide;
5-(benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide;
N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide;
 N^5 -(3,4-difluorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
5-[(4-fluoro-phenylamino)-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-{[2-(4-fluoro-phenyl)-ethylamino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(4-fluoro-benzoylamino)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

(8-hydroxycarbamoyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl)-carbamic acid benzyl ester;

5-{[benzyl-(4-fluoro-phenyl)-amino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-({(2-benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino}-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-[3-(4-fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(4-fluoro-phenoxyethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(3-chloro-4-fluoro-phenoxyethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;

5-(3-chloro-4-fluoro-benzyloxyethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-[2-(4-fluoro-phenyl)-ethoxymethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(2,4-difluoro-benzyloxyethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(3,4-difluoro-benzyloxyethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(4-fluoro-benzyloxyethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-(4-fluoro-phenoxyethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(3-chloro-4-fluoro-phenoxyethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(2,4-difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(3,4-difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(4-fluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-benzyloxymethyl-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

(S)-(-)-3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(1-phenyl-ethyl)-amide];

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(pyridin-2-ylmethyl)-amide];

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-benzylamide 2-hydroxyamide;

(S)-(-)-3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(2-hydroxy-1-phenyl-ethyl)-amide];

pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide;

2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-{[1-(4-fluoro-phenyl)-cyclopropyl]-amide} 8-hydroxyamide;

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-{[1-(4-fluoro-phenyl)-cyclopropyl]-amide} 2-hydroxyamide;

2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-benzylamide) 8-(methoxy-amide);

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(methoxy-amide);

2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-cyclohexylmethyl-amide 8-hydroxyamide;

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide;

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide;

4-hydroxymethyl-3-methoxy-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide;
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(hydroxy-methyl-amide);
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-dibenzylamide 2-hydroxyamide;
5-{{[1-(4-fluoro-phenyl)-cyclopropylamino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-{{[1-(4-fluoro-phenyl)-cyclopropylamino]-methyl}-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide;
5-(4-fluoro-benzylloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide;
2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-2-methylcarbamoyl-benzylamide) 8-hydroxyamide;
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-(4-methyl-benzylamide);
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-{{[2-(4-fluoro-phenyl)-ethyl]-amide} 2-hydroxyamide;
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(2,4-difluoro-benzylamide) 2-hydroxyamide;
(rac)- {2-(4-Chloro-phenyl)-1-[(4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamoyl]-ethyl}-carbamic acid methyl ester;
(rac) 5-{{[(4-Fluoro-benzyl)-(2-phenyl-cyclopropanecarbonyl)-amino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
(4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid methyl ester;
(3-chloro-4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester;
5-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

3-hydroxy-4-hydroxymethyl-5-(3-phenyl-propyl)-pyridine-2-carboxylic acid hydroxyamide;
 5-benzenesulfonylmethyl-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
 5-(4-fluoro-phenylmethanesulfonylmethyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide;
 5-(4-fluoro-phenylmethanesulfonylmethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
 (4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester;
 (4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid tert-butyl ester;
 3-hydroxy-4-methyl-pyridine-2,5-dicarboxylic acid 5-(3-chloro-4-fluoro-benzylamide) 2-hydroxyamide; and
 5-(4-fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide; and
 pharmaceutically acceptable salts thereof.

17. A compound according to claim 1 selected from:
 N, 9-bis(benzyloxy)-3,3-dimethyl-1,5-dihydro-[1,3]dioxepino[5,6-c]pyridine-8-carboxamide;
 N, 3-bis(benzyloxy)-4,5-bis(hydroxymethyl) picolinamide;
 N, 7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N, 3-bis(benzyloxy)-5-((4-fluorobenzylamino)methyl)-4-(hydroxymethyl)picolinamide;
 5-((3,5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide;
 5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N, 3-bis(benzyloxy)-4-(hydroxymethyl)picolinamide;
 5-(benzyloxy)-N-(4-fluorobenzyl)-4-(hydroxymethyl)-6-methylnicotinamide;

5-(benzyloxy)-N-(4-fluorobenzyl)-4-(methoxymethyl)-N, 6-dimethylnicotinamide;

3-(benzyloxy)-5-((4-fluorobenzyl)(methyl)carbamoyl)-4-(methoxymethyl)-2-methylpyridine 1-oxide;

5-(benzyloxy)-N-(4-fluorobenzyl)-6-(hydroxymethyl)-4-(methoxymethyl)-N-methylnicotinamide;

5-(benzyloxy)-N-(4-fluorobenzyl)-6-formyl-4-(methoxymethyl)-N-methylnicotinamide;

methyl 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl) carbamoyl)-4-(methoxymethyl)picolinate;

N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide;

N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide;

5-((4-methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine;

5-((4-methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine 7-oxide;

(5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridin-8-yl)methanol;

5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carbaldehyde;

ethyl 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxylate;

ethyl 3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinate;

N, 3-dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinamide;

5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridine-8-carboxylic acid;

N-(benzyloxy)-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide;

N-(benzyloxy)-3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl) picolinamide;
N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide;
5-((4-fluorophenylamino)methyl)-3-hydroxy-4-(hydroxymethyl) picolinate;
8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid;
methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
5-amino-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid methyl ester;
methyl 5-(4-fluorobenzamido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
methyl 5-(4-fluorobenzamido)-3-hydroxy-4-(hydroxymethyl)picolinate;
methyl 5-((benzyl(4-fluorophenyl)amino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate;
5-(((2-Benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino)-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
methyl 5-(3-(4-fluorophenyl)ureido)-3-hydroxy-4-(hydroxymethyl)picolinate;
5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate;
methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate;
methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
methyl 3-(benzyloxy)-5-((4-fluorophenoxy)methyl)-4-methylpicolinate;
methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate;
methyl 3-(benzyloxy)-5-((3-chloro-4-fluorophenoxy)methyl)-4-methylpicolinate;
methyl 5-((3-chloro-4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate;

5-[(4-Fluoro-benzylamino)-methyl]-3-hydroxy-4-methyl-pyridine-2-carboxylic acid methyl ester;
methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate; and
Methyl 5-((4-fluorophenethylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate; and
pharmaceutically acceptable salts thereof.

18. A compound selected from the group consisting of:
N⁵-(4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
N⁵-(3-chloro-4-fluorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
N⁵-(3,4-dichlorobenzyl)-N²,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide; and
5-(benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide; and
pharmaceutically acceptable salts thereof.
19. A pharmaceutical composition comprising a compound according to any one of claims 1 to 18 and a pharmaceutically acceptable carrier or diluent.
20. A pharmaceutical composition comprising a compound according to any one of claims 1 to 18, at least one additional HIV-inhibiting agent, and a pharmaceutically acceptable carrier or diluent.
21. Use of a compound according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof, in the manufacture of a drug for treatment of HIV infection.
22. Use of a compound according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof, in the manufacture of a drug for treatment of acquired immune deficiency syndrome (AIDS) or AIDS-related complex.
23. Use of a compound according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof, in the manufacture of a drug for treatment of hepatitis C infection.


24. A method of treating or preventing an HIV infection in a mammal, said method comprising administering to said mammal a compound according to any one of claims 1 to 18 in an amount effective for the treatment or prevention of said HIV infection.
25. A method of treating or preventing AIDS or AIDS-related complex in a mammal, said method comprising administering to said mammal a compound according to any one of claims 1 to 18 in an amount effective for the treatment or prevention of said AIDS or AIDS-related complex.
26. A method of inhibiting HIV replication in a mammal, said method comprising administering to said mammal a compound according to any one of claims 1 to 18 in an amount effective to inhibit HIV replication in a mammal.
27. A method of inhibiting HIV replication in a cell, said method comprising contacting said cell with a compound according to any one of claims 1 to 18 in an amount sufficient to inhibit HIV replication.
28. The method of any one of 24-27, said method further comprising administering to said mammal or contacting said cell with at least one additional HIV inhibiting agent.
29. The method of claim 28, wherein said HIV inhibiting agent is selected from the group consisting of an entry inhibitor, a protease inhibitor, a reverse transcriptase inhibitor, a fusion inhibitor, and an integrase inhibitor.
30. The method of claim 24, 26, or 27, wherein said HIV is resistant to at least one HIV inhibiting agent.

AMENDED CLAIMS

received by the International Bureau on 02 November 2009 (02.11.2009)

Claims

1. A compound of formula I,



I

wherein:

A is a six membered carbocyclic or heterocyclic ring system;

R₁ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl), (O-C₁₋₆ branched alkyl), CO(R₉), COO(R₉), CON(R₉)(R_{9a}), or SO₂N(R₉)(R_{9a}), wherein said R₉ and R_{9a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; R₂ is H, C₁₋₆ alkyl, C₁₋₆ branched alkyl, C₂₋₆ alkenyl, halogen (F, Cl, Br, I), OH, O-(C₁₋₆ alkyl),

(O-C₁₋₆ branched alkyl), CO(R₁₀), COO(R₁₀), or CON(R₁₀)(R_{10a}), wherein said R₁₀ and R_{10a} are selected independently from the group consisting of H, C₁₋₆ alkyl, C₁₋₆ fluoro-alkyl, benzyl, phenyl and heterocycle; or R₁ and R₂ are ortho substituents that together form a carbocyclic or heterocyclic ring system;

L is -N(R')C(O)-; -C(O)N(R')-; -OC(O)-; -C(O)O-; -OC(O)N(R')-; -N(R')C(O)O-; -N(R')C(O)N(R')-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-C(R_{a1})(R_{a2})-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-Z-; -C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-; -Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})C(R_{c1})(R_{c2})-;

$-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-$;
 $-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-C(R_{a1})(R_{a2})-$; $-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-Z-$;
 $-C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-$; $-Z-C(R_{a1})(R_{a2})C(R_{b1})(R_{b2})-$;
 $-C(R_{a1})(R_{a2})-Z-C(R_{a1})(R_{a2})-$; $-C(R_{a1})(R_{a2})-Z-$; or $-Z-C(R_{a1})(R_{a2})-$; wherein each R_{a1} , R_{a2} , R_{b1} , R_{b2} , R_{c1} , and R_{c2} is, independently, selected from selected from H, C_{1-6} alkyl, C_{1-6} fluoro-alkyl, hydroxy-alkyl, benzyl, phenyl and heterocycle, or, alternatively, one or more of R_{a1} and R_{a2} ; R_{b1} and R_{b2} ; and R_{c1} and R_{c2} combine to form a carbocyclic ring, and wherein Z is selected from $-N(R')C(O)-$; $-C(O)N(R')-$; $-OC(O)-$; $-C(O)O-$; $-OC(O)N(R')-$; $-N(R')C(O)O-$; $-N(R')C(O)N(R')-$; $-N(R')-$; $-SO_2-$; and $-O-$; wherein R' is selected from H, C_{1-6} alkyl, benzyl, SO_2R'' , and $C(O)R''$, $C(O)OR''$, and R'' is selected from C_{1-6} alkyl, C_{1-6} fluoro-alkyl, heteroalkyl, carbocyclic group, benzyl, phenyl and heterocycle;

B_1 is $-R_3$, CH_2OR_3 , $CH_2N(R_8)(R_{8a})$, $C(O)OR_3$ or $C(O)N(R_8)(R_{8a})$, wherein each of R_8 and R_{8a} is, independently, selected from the group consisting of H, C_{1-6} alkyl, C_{1-6} fluoro-alkyl, benzyl, phenyl and heterocycle;

B_2 is H or OR_5 ;

R_3 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; and R_5 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; or, alternatively, R_3 and R_5 combine to form a heterocyclic ring system; and

R_4 is H, C_{1-6} alkyl, C_{1-6} fluoroalkyl, benzyl, phenyl, or heterocycle; and
 pharmaceutically acceptable salts thereof.

2. A compound according to claim 1, wherein A is a phenyl ring, a pyridine ring, or a cyclohexyl ring.
3. A compound according to claim 1, wherein L is $-CH_2OCH_2-$, $-CH_2CH_2OCH_2-$, $-OCH_2-$, $-CH_2NHCH_2-$, $-C(cyclo-C_2H_4)NHCH_2-$, $-NHCH_2-$, $-CH_2CH_2NHCH_2-$, $-CH_2NHC(O)-$, $-CH_2N(CH_3)C(O)-$, $-CH(CH_2OH)NHC(O)-$, $-C(cyclo-C_2H_4)NHC(O)-$, $-CH_2CH_2NHC(O)-$, $-C(O)NH-$, $-CH_2OC(O)NH-$, $-NHC(O)NH-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2-$, $-SO_2CH_2-$, or $-CH_2SO_2CH_2-$.
4. A compound according to claim 1, wherein B_1 is H, CH_3 , CH_2OH , or CH_2OCH_3 .

5. A compound according to claim 1, wherein B_2 is H or $-OR_5$ and R_5 is H or benzyl.
6. A compound according to any of claims 2-5, wherein L is $-CH_2NHCH_2-$ or $-CH_2NHC(O)-$.
7. A compound according to claim 6, wherein each of R_1 and R_2 is, independently, selected from halogen, $-OCH_3$, $-OH$, or R_1 and R_2 combine to form a cyclic acetal or cyclic ketal.
8. A compound according to claim 7 wherein R_4 is $-H$ or benzyl.
9. A compound according to any of claims 2-5, wherein L is $-CH_2OCH_2-$.
10. A compound according to claim 9, wherein each of R_1 and R_2 is, independently, selected from halogen, $-OCH_3$, $-OH$, or R_1 and R_2 combine to form a cyclic acetal or cyclic ketal.
11. A compound according to claim 9, wherein R_4 is $-H$ or benzyl.
12. A compound according to any of claims 2-5, wherein L is $-SO_2CH_2-$, or $-CH_2SO_2CH_2-$.
13. A compound according to claim 12, wherein each of R_1 and R_2 is, independently, selected from halogen, $-OCH_3$, $-OH$, or R_1 and R_2 combine to form a cyclic acetal or cyclic ketal.
14. A compound according to claim 13, wherein R_4 is $-H$ or benzyl.
15. A compound according to any of claims 3, 6, 10, or 12, wherein A is a phenyl ring.
16. A compound according to claim 1 selected from:
 $N^2, 3$ -bis(benzyloxy)- N^5 -(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(4-fluorobenzyl)- $N^2, 3$ -dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(benzo[d][1,3]dioxol-5-ylmethyl)- $N^2, 3$ -bis(benzyloxy)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(benzo[d][1,3]dioxol-5-ylmethyl)- $N^2, 3$ -dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 $N^2, 3$ -bis(benzyloxy)-4-(hydroxymethyl)- N^5 -(4-methoxybenzyl)pyridine-2,5-dicarboxamide;

N^2 , 3-dihydroxy-4-(hydroxymethyl)- N^5 -(4-methoxybenzyl)pyridine-2,5-dicarboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(3,5-difluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(3, 5-difluorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 5-((4-fluorobenzylamino)methyl)-N, 3-dihydroxy-4-(methoxymethyl)picolinamide);
 5-((3, 5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide;
 5-((benzo[d][1, 3]dioxol-5-ylmethylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl)picolinamide;
 N^5 -(4-fluorobenzyl)- N^2 , 3-dihydroxy-4-(methoxymethyl)- N^5 -methylpyridine-2, 5-dicarboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(3-chloro-4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N^5 -(3-chloro-4-fluorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N^2 , 3-bis(benzyloxy)- N^5 -(3, 4-dichlorobenzyl)-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N^5 -(3, 4-dichlorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2, 5-dicarboxamide;
 N, 3-dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)-picolinamide;
 5-(benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide;
 N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxamide;
 N^5 -(3,4-difluorobenzyl)- N^2 ,3-dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 5-[(4-fluoro-phenylamino)-methyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

5-{[2-(4-fluoro-phenyl)-ethylamino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(4-fluoro-benzoylamino)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
(8-hydroxycarbamoyl-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridin-5-yl)-carbamic acid benzyl ester;
5-{[benzyl-(4-fluoro-phenyl)-amino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-({(2-benzyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino}-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-[3-(4-fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(4-fluoro-phenoxy-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(3-chloro-4-fluoro-phenoxy-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
5-(3-chloro-4-fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-[2-(4-fluoro-phenyl)-ethoxymethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(2,4-difluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(3,4-difluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(4-fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-(4-fluoro-phenoxy-methyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;
5-(3-chloro-4-fluoro-phenoxy-methyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(2,4-difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(3,4-difluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-(4-fluoro-benzyloxymethyl)-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

5-benzyloxymethyl-3-hydroxy-4-methyl-pyridine-2-carboxylic acid hydroxyamide;

(S)-(-)-3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(1-phenyl-ethyl)-amide];

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(pyridin-2-ylmethyl)-amide];

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-benzylamide 2-hydroxyamide;

(S)-(-)-3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-[(2-hydroxy-1-phenyl-ethyl)-amide];

pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide;

2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-[[1-(4-fluoro-phenyl)-cyclopropyl]-amide] 8-hydroxyamide;

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-[[1-(4-fluoro-phenyl)-cyclopropyl]-amide] 2-hydroxyamide;

2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-benzylamide) 8-(methoxy-amide);

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(methoxy-amide);

2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-cyclohexylmethyl-amide 8-hydroxyamide;

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide;

3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-cyclohexylmethyl-amide 2-hydroxyamide;

4-hydroxymethyl-3-methoxy-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-hydroxyamide;
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(4-fluoro-benzylamide) 2-(hydroxy-methyl-amide);
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-dibenzylamide 2-hydroxyamide;
5-{[1-(4-fluoro-phenyl)-cyclopropylamino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
5-{[1-(4-fluoro-phenyl)-cyclopropylamino]-methyl}-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide;
5-(4-fluoro-benzylloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide;
2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5,8-dicarboxylic acid 5-(4-fluoro-2-methylcarbamoyl-benzylamide) 8-hydroxyamide;
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 2-hydroxyamide 5-(4-methyl-benzylamide);
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-{[2-(4-fluoro-phenyl)-ethyl]-amide} 2-hydroxyamide;
3-hydroxy-4-hydroxymethyl-pyridine-2,5-dicarboxylic acid 5-(2,4-difluoro-benzylamide) 2-hydroxyamide;
(rac)- {2-(4-Chloro-phenyl)-1-[(4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamoyl]-ethyl}-carbamic acid methyl ester;
(rac) 5-{[(4-Fluoro-benzyl)-(2-phenyl-cyclopropanecarbonyl)-amino]-methyl}-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
(4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid methyl ester;
(3-chloro-4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-hydroxymethyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester;
5-[2-(4-Fluoro-phenyl)-ethyl]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;

3-hydroxy-4-hydroxymethyl-5-(3-phenyl-propyl)-pyridine-2-carboxylic acid hydroxyamide;
 5-benzenesulfonylmethyl-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
 5-(4-fluoro-phenylmethanesulfonylmethyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid hydroxyamide;
 5-(4-fluoro-phenylmethanesulfonylmethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
 (4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid benzyl ester;
 (4-fluoro-benzyl)-(5-hydroxy-6-hydroxycarbamoyl-4-methyl-pyridin-3-ylmethyl)-carbamic acid tert-butyl ester;
 3-hydroxy-4-methyl-pyridine-2,5-dicarboxylic acid 5-(3-chloro-4-fluoro-benzylamide) 2-hydroxyamide; and
 5-(4-fluoro-benzyloxymethyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid methoxy-amide; and
 pharmaceutically acceptable salts thereof.

17. A compound according to claim 1 selected from:
 N, 9-bis(benzyloxy)-3,3-dimethyl-1,5-dihydro-[1,3]dioxepino[5,6-c]pyridine-8-carboxamide;
 N, 3-bis(benzyloxy)-4,5-bis(hydroxymethyl) picolinamide;
 N, 7-bis(benzyloxy)-3-oxo-1,3-dihydrofuro[3,4-c]pyridine-6-carboxamide;
 N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N, 3-bis(benzyloxy)-5-((4-fluorobenzylamino)methyl)-4-(hydroxymethyl)picolinamide;
 5-((3,5-difluorobenzylamino)methyl)-N, 3-dihydroxy-4-(hydroxymethyl) picolinamide;
 5-((benzo[d][1,3]dioxol-5-ylmethylamino)methyl)-N, 3-bis(benzyloxy)-4-(hydroxymethyl)picolinamide;
 5-(benzyloxy)-N-(4-fluorobenzyl)-4-(hydroxymethyl)-6-methylnicotinamide;


5-(benzyloxy)-N-(4-fluorobenzyl)-4-(methoxymethyl)-N, 6-dimethylnicotinamide;
 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl)carbamoyl)-4-(methoxymethyl)-2-methylpyridine 1-oxide;
 5-(benzyloxy)-N-(4-fluorobenzyl)-6-(hydroxymethyl)-4-(methoxymethyl)-N-methylnicotinamide;
 5-(benzyloxy)-N-(4-fluorobenzyl)-6-formyl-4-(methoxymethyl)-N-methylnicotinamide;
 methyl 3-(benzyloxy)-5-((4-fluorobenzyl)(methyl) carbamoyl)-4-(methoxymethyl)picolinate;
 N², 3-bis(benzyloxy)-N⁵-(4-fluorobenzyl)-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide;
 N⁵-(4-fluorobenzyl)-N², 3-dihydroxy-4-(methoxymethyl)-N⁵-methylpyridine-2, 5-dicarboxamide;
 5-((4-methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine;
 5-((4-methoxybenzyloxy)methyl)-2, 2, 8-trimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine 7-oxide;
 (5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridin-8-yl)methanol;
 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carbaldehyde;
 ethyl 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1, 3]dioxino[4, 5-c]pyridine-8-carboxylate;
 ethyl 3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinate;
 N, 3-dihydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl)picolinamide;
 5-((4-methoxybenzyloxy)methyl)-2, 2-dimethyl-4H-[1,3]dioxino[4, 5-c]pyridine-8-carboxylic acid;
 N-(benzyloxy)-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide;

N-(benzyloxy)-3-hydroxy-4-(hydroxymethyl)-5-((4-methoxybenzyloxy)methyl) picolinamide;
 N-hydroxy-5-((4-methoxybenzyloxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxamide;
 5-((4-fluorophenylamino)methyl)-3-hydroxy-4-(hydroxymethyl) picolinate;
 8-(methoxycarbonyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-5-carboxylic acid;
 methyl 5-(benzyloxycarbonylamino)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
 5-amino-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylic acid methyl ester;
 methyl 5-(4-fluorobenzamido)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
 methyl 5-(4-fluorobenzamido)-3-hydroxy-4-(hydroxymethyl)picolinate;
 methyl 5-((benzyl(4-fluorophenyl)amino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
 methyl 5-(((2-(benzyloxy)ethyl)(4-fluorophenethyl)amino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate;
 5-({(2-Benzoyloxy-ethyl)-[2-(4-fluoro-phenyl)-ethyl]-amino}-methyl)-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
 methyl 5-(3-(4-fluorophenyl)ureido)-3-hydroxy-4-(hydroxymethyl)picolinate;
 5-[3-(4-Fluoro-phenyl)-ureido]-3-hydroxy-4-hydroxymethyl-pyridine-2-carboxylic acid hydroxyamide;
 methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate;
 methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate;
 methyl 5-((3-chloro-4-fluorophenoxy)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate;
 methyl 3-(benzyloxy)-5-((4-fluorophenoxy)methyl)-4-methylpicolinate;
 methyl 5-((4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate;
 methyl 3-(benzyloxy)-5-((3-chloro-4-fluorophenoxy)methyl)-4-methylpicolinate;
 methyl 5-((3-chloro-4-fluorophenoxy)methyl)-3-hydroxy-4-methylpicolinate;

5-[(4-Fluoro-benzylamino)-methyl]-3-hydroxy-4-methyl-pyridine-2-carboxylic acid methyl ester;
 methyl 5-((4-fluorophenylamino)methyl)-2,2-dimethyl-4H-[1,3]dioxino[4,5-c]pyridine-8-carboxylate; and
 Methyl 5-((4-fluorophenethylamino)methyl)-3-hydroxy-4-(hydroxymethyl)picolinate; and
 pharmaceutically acceptable salts thereof.

18. A compound selected from the group consisting of:
 N^5 -(4-fluorobenzyl)- $N^{2,3}$ -dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(3-chloro-4-fluorobenzyl)- $N^{2,3}$ -dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide;
 N^5 -(3,4-dichlorobenzyl)- $N^{2,3}$ -dihydroxy-4-(hydroxymethyl)pyridine-2,5-dicarboxamide; and
 5-(benzyloxymethyl)-N,3-dihydroxy-4-(hydroxymethyl)picolinamide; and
 pharmaceutically acceptable salts thereof.
19. A pharmaceutical composition comprising a compound according to any one of claims 1 to 18 and a pharmaceutically acceptable carrier or diluent.
20. A pharmaceutical composition comprising a compound according to any one of claims 1 to 18, at least one additional HIV-inhibiting agent, and a pharmaceutically acceptable carrier or diluent.
21. Use of a compound according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof, in the manufacture of a drug for treatment of HIV infection.
22. Use of a compound according to any one of claims 1 to 18 or a pharmaceutically acceptable salt thereof, in the manufacture of a drug for treatment of acquired immune deficiency syndrome (AIDS) or AIDS-related complex.

23. A method of treating or preventing an HIV infection in a mammal, said method comprising administering to said mammal a compound according to any one of claims 1 to 18 in an amount effective for the treatment or prevention of said HIV infection.
24. A method of treating or preventing AIDS or AIDS-related complex in a mammal, said method comprising administering to said mammal a compound according to any one of claims 1 to 18 in an amount effective for the treatment or prevention of said AIDS or AIDS-related complex.
25. A method of inhibiting HIV replication in a mammal, said method comprising administering to said mammal a compound according to any one of claims 1 to 18 in an amount effective to inhibit HIV replication in a mammal.
26. A method of inhibiting HIV replication in a cell, said method comprising contacting said cell with a compound according to any one of claims 1 to 18 in an amount sufficient to inhibit HIV replication.
27. The method of any one of 23-26 said method further comprising administering to said mammal or contacting said cell with at least one additional HIV inhibiting agent.
28. The method of claim 27, wherein said HIV inhibiting agent is selected from the group consisting of an entry inhibitor, a protease inhibitor, a reverse transcriptase inhibitor, a fusion inhibitor, and an integrase inhibitor.
29. The method of claim 23, 25 or 26 wherein said HIV is resistant to at least one HIV inhibiting agent.

Figure 1**PK profile in rats**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA2009/000787

A. CLASSIFICATION OF SUBJECT MATTER

IPC: *C07D 213/81* (2006.01) , *A61K 31/44* (2006.01) , *A61K 31/4427* (2006.01) , *A61P 31/12* (2006.01) ,
A61P 31/18 (2006.01) , *C07D 405/12* (2006.01), *C07D 493/04* (2006.01).

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC: *C07D 213/81* (2006.01) , *A61K 31/44* (2006.01) , *A61K 31/4427* (2006.01) , *A61P 31/12* (2006.01) ,
A61P 31/18 (2006.01) , *C07D 405/12* (2006.01), *C07D 493/04* (2006.01).

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic database(s) consulted during the international search (name of database(s) and, where practicable, search terms used)
STN (Registry file, CAPLUS) , Delphion, Canadian Patent Database, Scopus (search terms: "HIV integrase inhibitors", "integrase enzymes", "pyridoxine").

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 1867636, A1 (MIYAUCHI ET AL.) 19 December 2007 (19-12-2007) page 57, Ex. 78; claims 1-4, 7-9, 11, 16, 17, 29, 32	1, 2, 4, 5, 19

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier application or patent but published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"&" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

29 July 2009 (29-07-2009)

Date of mailing of the international search report

3 September 2009 (03-09-2009)

Name and mailing address of the ISA/CA
Canadian Intellectual Property Office
Place du Portage I, C114 - 1st Floor, Box PCT
50 Victoria Street
Gatineau, Quebec K1A 0C9
Facsimile No.: 001-819-953-2476

Authorized officer

Olusola Womiloju 819- 994-1689

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/CA2009/000787**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of the first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons :

1. ☒ Claim Nos. : 24-30
 because they relate to subject matter not required to be searched by this Authority, namely :

 Claims 24-30 are directed to a method for treatment of the human or animal body by surgery or therapy which the International Search Authority is not required to search. However, this Authority has carried out a search based on the alleged effects or purposes/uses of the product defined in claims 24-30.
2. ☐ Claim Nos. :
 because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically :
3. ☐ Claim Nos. :
 because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows :

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claim Nos. :
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim Nos. :

- Remark on Protest** ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.
PCT/CA2009/000787

Patent Document Cited in Search Report	Publication Date	Patent Family Member(s)	Publication Date
EP 1867636A1	19-12-2007	AU 2006234441A1	19-10-2006
		CA 2603320A1	19-10-2006
		CN 101163678A	16-04-2008
		KR 20070116865A	11-12-2007
		MX 2007012490A	09-11-2007
		NO 20075080A	05-11-2007
		US 2009149439A1	11-06-2009
		WO 2006109729A1	19-10-2006