

(19) **DANMARK**

(10) **DK/EP 3047269 T3**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **G 01 N 33/50 (2006.01)** **G 01 N 33/487 (2006.01)** **G 01 N 33/53 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-09-17**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-08-01**
- (86) Europæisk ansøgning nr.: **14772109.6**
- (86) Europæisk indleveringsdag: **2014-09-19**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-07-27**
- (86) International ansøgning nr.: **EP2014070053**
- (87) Internationalt publikationsnr.: **WO2015040190**
- (30) Prioritet: **2013-09-19 GB 201316668**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Salurate Limited, Unit 17 Petersfield Business Park , Bedford Road, Petersfield, Hampshire GU32 3QA, Storbritannien**
- (72) Opfinder: **MAGUIRE, Patrick Martin, Ballymore, Killinick, Co. Wexford, Irland**
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **DIAGNOSE AF FØTAL DISTRESS**
- (56) Fremdragne publikationer:
WO-A2-2004/113926
MANY A ET AL: "HYPERURICEMIA AND XANTHINE OXIDASE IN PREECLAMPSIA, REVISITED", AMERICAN JOURNAL OF OBSTETRICS & GYNECOLOGY, MOSBY, ST LOUIS, MO, US, vol. 174, no. 1, PART 01, 1 January 1996 (1996-01-01), pages 288-291, XP009042145, ISSN: 0002-9378, DOI: 10.1016/S0002-9378(96)70410-6
CHAPPELL L C ET AL: "A LONGITUDINAL STUDY OF BIOCHEMICAL VARIABLES IN WOMEN AT RISK OF PREECLAMPSIA", AMERICAN JOURNAL OF OBSTETRICS & GYNECOLOGY, MOSBY, ST LOUIS, MO, US, vol. 187, no. 1, 1 July 2002 (2002-07-01), pages 127-136, XP001205032, ISSN: 0002-9378, DOI: 10.1067/MOB.2002.122969
THANGARATINAM S ET AL: "Accuracy of serum uric acid in predicting complications of pre-eclampsia: a systematic review.", BJOG : AN INTERNATIONAL JOURNAL OF OBSTETRICS AND GYNAECOLOGY APR 2006, vol. 113, no. 4, April 2006 (2006-04), pages 369-378, ISSN: 1470-0328

DESCRIPTION

[0001] The present invention relates to diagnosis or monitoring of foetal distress, in particular distress caused by conditions in the mother other than pre-eclampsia.

[0002] Pregnancies are often associated with distress in the foetus. It is important to monitor foetal distress because this may indicate that the pregnancy or labour is compromised. A number of signs of foetal distress have been historically used such as decreased foetal movement felt by the mother, meconium in the amniotic fluid, abnormal indications in cardiotocography and biochemical signs typically assessed by collecting a small sample of baby's blood from a scalp prick through the open cervix during labour.

[0003] Foetal distress is thought to be one indicator of a multitude of possible issues in the foetus/pregnancy such as breathing problems, growth disorders, gestational diabetes, a breach position, umbilical cord problems, placental problems, uterine problems and pre-eclampsia.

[0004] Treatment of foetal distress depends on its severity, but can range from monitoring as part of a pre-natal care plan to emergency admission for induced labour or hitherto unplanned caesarean section birth.

[0005] Pregnancies at risk of developing problems that may cause foetal distress are currently identified by measurements of high gestational blood pressure and/or urinary protein levels. However, these measurements have been reported as being poor predictors of pregnancy problems. As a result a large proportion of apparently high-risk pregnancies, which have been allocated an intensive and expensive care, treatment and diagnostics plan, are trouble-free, and a large proportion of apparently low-risk pregnancies require unexpected medical intervention. Accordingly, current medical practice does not adequately identify pregnancies at risk of developing problems that may cause foetal distress.

[0006] Many et al (1996) Am J Obstet Gynecol 174(1 Pt 1):288-91 is a review article that discusses the association between preeclampsia and high serum uric acid concentration and hypothesizes about possible driving forces behind the hyperuricemia.

[0007] EP 1634082, originally published as WO 2004/113926, discloses a method for diagnosis of pre-eclampsia in the mother comprising measuring urate in a biological sample from maternal saliva. This enables identification of women at risk of or having pre-eclampsia by the presence of an elevated urate level in maternal saliva.

[0008] The method of EP 1634082 has been developed specifically to provide a warning signal for pre-eclampsia and the signal it monitors, salivary urate levels, is subject to regulation according to diurnal rhythms, the stage of pregnancy and the medical/emotional status of the mother.

[0009] JP 20062666721 relates to the use of salivary urate for monitoring the health of a subject.

[0010] Soukup et al (2012) Diabet. Metabol. Syndrome 4:14, which investigates correlations between salivary urate and cardiometabolic risk factors, suggests that salivary and serum uric acid levels have a linear relationship. Data to support this assertion is not presented. In any case, this study is unrelated to foetal health.

[0011] Gao et al (2008) Prenatal Diagnosis 28:810-814 studies whether urate in amniotic fluid is predictive of infant birth outcomes. Sampling of amniotic fluid as used in Gao et al is invasive and thus risks harm to the foetus.

[0012] Akahori et al (2012) Gynecol. Obs. Invest. 73:162-167 correlates maternal serum urate with foetal growth and maternal kidney function. Assays were performed at week 36, which is too late to provide any guidance for antenatal care.

[0013] Laughton et al (2009) Am. J. Obs. Gynecol. 201:582.e1-582.e6 correlates maternal plasma urate with foetal growth.

[0014] Some known studies are hence based on urate sampling of serum or plasma. The inventors have found, however, that salivary and serum urate levels are uncorrelated (Figures 1 and 2).

[0015] The inventors also note that salivary and serum urate are not correlated in conditions such as gout or during/following exercise (Owen-Smith et al (1998) The Lancet 351:1932). Without being bound by any theory, the decoupling of salivary and serum urate may be explained by the salivary urate being sensitive to urate levels in interstitial fluids, which may vary independently from blood urate levels.

[0016] There remains a need for an improved or alternative method for diagnosing foetal distress regardless of the cause.

[0017] Accordingly, the present invention provides a method of diagnosing and/or monitoring foetal distress comprising measuring uric acid levels in saliva in one or more samples obtained from the expectant mother wherein the distress is other than that caused by pre-eclampsia.

[0018] Foetal distress is defined according to the invention as stress in the foetus manifesting in metabolic stress. In making the invention, the inventors have found that metabolic stress in the foetus may be detected in the mother's saliva for example in elevated levels of uric acid, and that this metabolic stress is a warning sign for various specific disorders.

[0019] Collection of saliva is not invasive, unlike for example sampling of blood from the mother or the foetus and is not associated with fear for example of pain related to providing a

blood sample using a hypodermic needle or finger prick test. Collection of saliva is also more convenient than collection of urine. The level of patient compliance for saliva testing is therefore high, leading to fewer delays in testing and better clinical outcomes. Additionally, collection of saliva does not require the services of a skilled practitioner.

[0020] While the diagnosis or monitoring of foetal distress according to the invention does not provide a diagnosis of the specific cause of the distress, it does enable mothers to be allocated or transferred to a more appropriate care, treatment and diagnostics plan.

[0021] The method of the invention therefore provides an early warning of potential issues in a pregnancy and the more efficient care of at risk mothers and fetuses. Diagnosis or monitoring can be based upon a single sample, in which case it is preferable to monitor for causes of spikes in urate levels not associated with foetal distress, for example acute stress in the mother. In typical use of the invention, two or more samples, more preferably five or more samples, typically ten or more, or 20 or more samples, are obtained from the expectant mother.

[0022] The diagnosis or monitoring can be used to produce a wider ranging diagnosis of pregnancy/foetus difficulties than hitherto available. It is known in the art to provide a test indicative only of pre-eclampsia. The present invention provides diagnosis or monitoring of foetal distress wherein the distress is other than that caused by pre-eclampsia. Thus, for example, a mother known not to be suffering from pre-eclampsia can be tested for foetal distress other than caused by pre-eclampsia. The test of the invention can be carried out with an open mind as to the specific cause of the foetal distress.

[0023] Another embodiment of the invention provides a method wherein the mother is not known to have a condition that could cause foetal distress. Thus, the method enables mothers who are otherwise considered to be having a normal pregnancy, and so may not have particular testing done, to be transferred to a more appropriate care, treatment and diagnostics plan.

[0024] While the method of the invention may be useful for patients with known past or present issues with pregnancy, it is particularly suitable for mothers who are generally considered to be at low risk of experiencing complications with their pregnancy. Such mothers may be considered to be from around 12 or 16 years old, generally between around 16 and 50 years of age, preferably between around 18 and 40 years of age, and to have a clinically normal blood pressure, and/or a clinically normal level of protein in the urine.

[0025] A clinically normal blood pressure according to the invention is defined in the usual manner by clinicians according to local practices, and may comprise a systolic blood pressure of 140 mmHg or less and/or a diastolic blood pressure of 90 mmHg or less.

[0026] A clinically normal level of protein according to the invention is defined in the usual manner by clinicians according to local practices, and may comprise less than 300 mg,

preferably less than 150 mg of protein in a 24-hour urine sample.

[0027] Accordingly, the method of the invention works on the principle that a clinically high uric acid level in the sample indicates foetal distress.

[0028] A clinically high uric acid level in saliva according to the invention may be calibrated according to a baseline (normal) level at the beginning of pregnancy, or at another suitable milestone, which baseline may be stable or time/development varied. The normal level may be determined on an individual, group or population level. Typically, a clinically high uric acid level in saliva is above 275 $\mu\text{mol/l}$, preferably above 350 $\mu\text{mol/l}$, more preferably above 400 $\mu\text{mol/l}$. Generally, a pattern of high uric acid levels in the sample indicates foetal distress

[0029] The sensitivity of the test can be adjusted by setting the level at which the uric acid level is considered clinically high. There is a trade-off between sensitivity and specificity such that a lower threshold for considering the level as clinically high provides a highly sensitive test, which will have a low specificity (i.e. more false positives) due to detection of normal variations in the levels - e.g. diurnal variations. However, a higher threshold will conversely provide a lower sensitivity with higher specificity.

[0030] For diagnosis or monitoring of foetal stress, an embodiment of the invention requires that the uric acid level is clinically high in two or more samples, preferably three or more samples, optionally four or more samples, in which the samples are obtained on different days.

[0031] By incorporating a requirement for the detection of repeated high levels of uric acid to trigger a diagnosis or indication of foetal distress, an embodiment of the invention enables an increase in the specificity of the diagnosis by filtering out cases in which fewer than the required number of samples have a high level of uric acid.

[0032] In an embodiment of the invention, the sample is obtained once a week. This further enhances patient compliance by establishing a predictable routine.

[0033] In another embodiment of the invention, the sample is obtained more frequently than once a week, for example once a day for a period of e.g. 4 days or more, a week or more, two weeks or more, or even longer. The sampling frequency may be varied throughout the monitoring period according to clinical need or interest.

[0034] In another embodiment of the invention, the sample is obtained at approximately the same time in the day, which advantageously avoids false positive or false negative measurements e.g. due to natural diurnal rhythms.

[0035] Many complications may occur in the late second or third trimester of pregnancies. Accordingly, in an embodiment of the invention, the sample is obtained from week 30, preferably week 20, optionally from week 10 of the pregnancy until birth. Typically, the sample is obtained from week 20 of the pregnancy until birth, optionally at a higher frequency, for

example daily, approximately in the period prior to birth. Limiting sampling and testing to particular stages of pregnancy reduces costs, enabling large scale monitoring to be more financially feasible.

[0036] The collection of saliva samples from the mother may be carried out cheaply without technically sophisticated equipment, using for example a tube, in which the mother may spit, or an absorbent swab, which may be chewed or simply placed under the tongue. Preferably, the sample is a sublingual salivary sample.

[0037] The present invention may be used to detect warning signs for all pregnancy conditions that cause foetal distress, and may indicate one or more of HELLP syndrome, eclampsia, gestational diabetes, gestational hypertension, premature amniorrhexis, placenta accreta and intrauterine growth restriction. Additionally, the present invention may provide an early warning of the possibility of premature or other emergency birth.

[0038] Another aspect of the invention provides use of uric acid levels in maternal saliva as an indicator of foetal distress wherein the foetal distress is not caused by pre-eclampsia, in particular distress caused by one or more of HELLP syndrome, eclampsia, gestational diabetes, gestational hypertension, premature amniorrhexis, placenta accreta and intrauterine growth restriction. The indicator is suitable for use in monitoring or diagnostic purposes.

[0039] Other optional and preferred embodiments of the use are as described in relation generally to the method of diagnosis or monitoring of the invention.

[0040] Specific embodiments of the invention will now be described in the following Examples, with reference to the accompanying drawings, in which:

Fig. 1 shows levels of urate in saliva and blood measured every 2 hours over a 50 hour period; and

Fig. 2 plots the levels of salivary urate against blood urate using the data shown in Figure 1.

Example 1 - Sample Collection and Uric Acid Measurement

[0041] A sample of saliva was collected from a set of pregnant subjects attending antenatal clinic by placing an absorbent swab under the tongue. A drop of saliva was expressed from each swab onto a test strip for a standard hand-held uric acid meter intended for measuring uric acid in whole blood samples. The sample was then tested per the manufacturer's instructions and the measured uric acid level was indicated on the meter and noted.

[0042] The indicated uric acid level was calibrated to provide the uric acid concentration from a whole blood sample. Therefore, the salivary uric acid level was calculated from the indicated

uric acid level using a correction factor.

[0043] The salivary uric acid concentration was determined from samples collected once a week at around the same time of day from week six or week 20 of pregnancy to birth.

Example 2 - Data Analysis and Diagnosis

[0044] Uric acid concentrations were determined according to Example 1 from week six of pregnancy from a random sample of 149 women. These data show that mean salivary uric acid increased non-linearly with gestational age from approximately 100 $\mu\text{mol/l}$ at six weeks to around 250 $\mu\text{mol/l}$ at 40 weeks, the rate of increase rising from week 30 to week 40.

[0045] Therefore, the analysis for this Example assumes that salivary uric acid levels of less than 350 $\mu\text{mol/l}$ are normal, levels in the range 350 to 400 $\mu\text{mol/l}$ are marginal, and levels of more than 400 $\mu\text{mol/l}$ are abnormally high.

[0046] In 135 subjects, salivary uric acid concentrations were measured according to Example 1 weekly from week 20 of pregnancy until birth and the clinical outcome of each pregnancy noted. The results of this study are set out below.

[0047] Outcomes of the study were tabulated into four groups according to the clinical events in the pregnancy as follows:

Table 1.

Subjects diagnosed as having pre-eclampsia (not claimed)

Table 2.

Small babies ($\sim 2500\text{g}$ or less) and admissions to the neonatal unit

Table 3.

Subjects with other medical conditions including diabetes

Table 4.

Normal pregnancies with abnormal salivary uric acid results

Table 1 - Subjects diagnosed as having pre-eclampsia

No. of Subjects	135
Sensitivity of salivary uric acid	15 subjects were classified as having pre-eclampsia. All 15 showed repeated high salivary uric acid results
Diagnostic potential	Number of abnormal salivary uric acid readings per subject diagnosed with pre-eclampsia = 6.6 separate occasions on average of the 15 'positives'.
Predictive potential	Average week of first salivary uric acid flag = week 24

[0048] Referring to Table 1, 15 subjects were classified by the clinicians as having pre-eclampsia. Of these fifteen, ten exhibited gestational hypertension and urinary protein, whilst five showed hypertension only. All fifteen were marked by a pattern of elevated salivary urate levels appearing on six to seven occasions on average. In eleven of the fifteen pre-eclampsia subjects, the first abnormal salivary uric acid reading appeared on average 13 weeks before the first conventional marker (hypertension/proteinuria). The pre-eclampsia salivary uric acid pattern was also exhibited by two subjects who had had severe pre-eclampsia in an earlier pregnancy, but did not show hypertension or urinary protein on this occasion.

[0049] There were no false negatives. No subjects showing the 'normal' salivary uric acid pattern were found to have pre-eclampsia or other gestational abnormalities.

Table 2 - Small babies (~2500g or less) and admissions to the neonatal unit.

Subject No.	Baby weight	Flagged by salivary uric acid	Flagged by CMA	Detected clinical skills	Undiagnosed	Comments
1	2490g	✓	✓	✓		Previous HELLP; pre-eclampsia emergency caesarean section
49	1730g	✓	✓	✓		HELLP syndrome; admitted to neonatal unit
104	1610g	✓	✓	✓		Severe pre-eclampsia caesarean at 32 weeks; admitted to neonatal unit
37	2370g	✓		✓		Premature; ROM; delivered at 34 weeks
45	1940g	✓		✓		Fragmented placenta accreta; spontaneous labour at 32 weeks
64	2308g	✓		✓		IUGR diagnosed by ultrasound
20	2000g	✓			X	Planned home delivery; foetal distress; emergency admission to neonatal unit
129	2540g	✓			X	Precipitous home delivery at 38 weeks; breathing difficulty; admission to neonatal unit

CMA = Conventional Pre-eclampsia Marker Abnormality (gestational hypertension or proteinuria); HELLP = Haemolysis, Elevated Liver enzymes and Low Platelet

count; ROM = rupture of membranes (amniorrhexis); IUGR = intrauterine growth restriction.

[0050] Referring to Table 2, eight subjects delivered small babies. All eight were identifiable by their elevated salivary uric acid patterns. Only three of these exhibited conventional pre-eclampsia marker abnormalities. A further three did not show either hypertension or proteinuria, but were diagnosed by the professional expertise of the ante-natal care team.

[0051] Two of the eight subjects were undiagnosed. Both subjects had exhibited elevated salivary uric acid levels starting in week 24. Both were scheduled for home deliveries, and resulted in emergency admissions to the neonatal unit. One of these subjects who had no conventional markers of pre-eclampsia, was scheduled for home delivery but in week 38 required emergency admission to a neonatal unit.

Table 3 - Miscellaneous conditions showing elevated SalU results.

No. of Subjects	Condition
4	Diabetes (type 2 insulin dependent)
1	Multiple Sclerosis (in remission)
1	Pancreatitis
1	Campylobacter Infection/Diarrhoea

[0052] Referring to Table 3, seven subjects with medical disorders exhibited a pattern of elevated salivary uric acid results. These conditions were known to the obstetrics team prior to delivery.

Table 4 - Subjects with uneventful outcomes associated with abnormal salivary uric acid readings

No. of SUBJECTS	11
Frequency of abnormal salivary uric acid readings	Average of 1.5 abnormal readings per subject (compared to 6.6 abnormal readings per subject for pre-eclampsia subjects)
Timing of abnormal readings	Average timing of last abnormal reading seen 8.4 weeks before delivery (compared to week of delivery for pre-eclampsia subjects)

[0053] In eleven cases where deliveries were uneventful, the results showed either one or two instances of abnormal salivary uric acid levels prior to week 30, a pattern recognisably less marked than in the cases of gestational abnormality, where elevated salivary uric acid typically occurred on six / seven occasions from week 24 up to delivery.

[0054] Therefore, incorporating a requirement of at least two abnormal salivary uric acid

readings into the criteria for diagnosing foetal distress reduced the detection of false positives.

Example 3 - Alternative Data Analysis and Diagnosis

[0055] To develop a specific diagnosis method, the data set from Example 2 was reanalysed. Subjects were included for reanalysis if it was the subject's first pregnancy, there was a low clinical risk of pregnancy issues (defined as an absence of information to the contrary in subject's medical records) and the subject was aged 18-40.

[0056] The inclusion criteria were formulated to exclude subjects who would already have a known history of pregnancy issues and/or would be selected for a greater level of monitoring by more conventional methods.

[0057] Seventy-six subjects were selected according to the stated criteria. Of these, 26 subjects provided two or more saliva samples in which the uric acid level was 275 $\mu\text{mol/l}$ or greater. The clinical outcomes of the 17 of the 26 subjects that exhibited clinical indications are listed in Table 5.

Table 5 - Clinical outcomes

Confirmed Clinical Pre-eclampsia (not claimed)		Other Clinical Indications	
9 Subjects		8 Subjects	
Subject	Severity	Subject	Condition
A	2	J	Emergency admission at week 39 following planned homebirth. Admission to neonatal unit. Intrauterine growth restriction with a birth weight of 2000g.*
B	1		
C	3		
D	2	K	Gestational diabetes.
E	3	L	Emergency admission at week 34 following planned homebirth. Admission to neonatal unit. Birth weight of 2370g.*
F	1		
G	3		
H	1	M	Emergency C. section due to breech position at week 37 Pancreatitis.
I	1	N	
		O	Intrauterine growth restriction with a birth weight at week 38 was 2308g.
		P	Emergency C. section. Suboptimal cardiotocography.
		Q	Emergency admission at week 38. Admission to neonatal intensive care unit. Intrauterine growth restriction with a birth weight of 2540g.
			*Not detected by conventional diagnostic indicators

Confirmed Clinical Pre-eclampsia (not claimed)		Other Clinical Indications	
9 Subjects		8 Subjects	
Subject	Severity	Subject	Condition
Grading of pre-eclampsia severity was carried out by consultant obstetrician (retired), and codes indicate the following severity: 3 = severe: 2 = moderate; 1=mild.			

[0058] There were no false negative results, meaning that no subjects that provided one or no saliva samples in which the uric acid level was 275 $\mu\text{mol/l}$ or greater exhibited clinical indications detected by conventional diagnostic indicators. Thus, the diagnosis detected 100% of subjects in which there were the specified clinical issues with the pregnancies. The method therefore exhibits a higher sensitivity than known current clinical practice, which is reported to have a sensitivity of 46%. There were nine false positive results, meaning that seven subjects provided two or more saliva samples in which the uric acid level was 275 $\mu\text{mol/l}$ or greater, thus indicating apparent foetal distress, yet exhibited no clinical indications detected by conventional diagnostic indicators. Thus, the specificity of the method of the invention is such that a high proportion (85%) of subjects in which there was an indicator of foetal distress had clinical issues with the pregnancies. This level of specificity is higher than current clinical practice, which is reported to have a specificity of 83%.

[0059] The method of the invention accurately identified 46 out of 46 negative clinical outcomes.

Comparative Example 1 - Blood Urate Does Not Predict Salivary Urate

[0060] Salivary urate concentrations were determined according to Example 1 and blood urate concentrations were determined using standard methods. The levels of urate in saliva and blood were measured in one subject every 2 hours over a 50 hour period. Both salivary and blood urate varied over the monitoring period (Figure 1). However, an increase or decrease in the salivary urate level was not always associated with a corresponding increase or decrease in blood urate level (Figure 1). A plot of blood urate against salivary urate reveals no useful correlation between the two urate measurements with the data being widely scattered (Figure 2; linear $r^2 = 0.1028$). Thus, blood urate did not predict salivary urate.

[0061] Accordingly, the invention provides diagnosis of foetal distress other than pre-eclampsia and uses thereof.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [EP1634082A](#) [0007] [0008]
- [WO2004113926A](#) [0007]
- [JP20062666721B](#) [0009]

Non-patent literature cited in the description

- **MANY et al.** Am J Obstet Gynecol, 1996, vol. 174, 1288-91 [0006]
- **SOUKUP et al.** Diabet. Metabol. Syndrome, 2012, vol. 4, 14- [0010]
- **GAO et al.** Prenatal Diagnosis, 2008, vol. 28, 810-814 [0011]
- **AKAHORI et al.** Gynecol. Obs. Invest., 2012, vol. 73, 162-167 [0012]
- **LAUGHTON et al.** Am. J. Obs. Gynecol., 2009, vol. 201, 582.e1-582.e6 [0013]
- **OWEN-SMITH et al.** The Lancet, 1998, vol. 351, 1932- [0015]

Patentkrav

1. Fremgangsmåde til diagnosticering og/eller monitorering af føtal distress, hvilken fremgangsmåde omfatter måling af urinsyreniveauer i spyt i én eller flere prøver opnået fra den gravide mor, hvor den pågældende distress er andet end forårsaget af præeklampsi.
- 5 2. Fremgangsmåde ifølge krav 1, hvor moren ikke er kendt for at have en tilstand, der kunne forårsage føtal distress.
3. Fremgangsmåde ifølge krav 1 eller 2, hvor moren er mellem 18 og 40 år gammel.
4. Fremgangsmåde ifølge et hvilket som helst foregående krav, hvor moren har a) et systolisk blodtryk på 140 mmHg eller mindre og et diastolisk blodtryk på 90 mmHg eller
10 mindre og/eller b) mindre end 300 mg, fortrinsvis mindre end 150 mg, protein i en 24-timers urinprøve.
5. Fremgangsmåde ifølge et hvilket som helst foregående krav, hvor et klinisk højt urinsyreniveau i prøven indikerer føtal distress.
6. Fremgangsmåde ifølge krav 5, hvor urinsyreniveauet er over 275 $\mu\text{mol/l}$, fortrinsvis
15 over 350 $\mu\text{mol/l}$, mere fortrinsvis over 400 $\mu\text{mol/l}$.
7. Fremgangsmåde ifølge krav 5 eller krav 6, hvor urinsyreniveau, der er højt i to eller flere prøver, fortrinsvis tre eller flere prøver, eventuelt fire eller flere prøver, hvor prøverne er opnået på forskellige dage, indikerer føtal distress.
8. Fremgangsmåde ifølge et hvilket som helst foregående krav, hvor prøven opnås én
20 gang om ugen.
9. Fremgangsmåde ifølge et hvilket som helst foregående krav, hvor prøven opnås fra uge 20 af svangerskabet frem til fødslen.
10. Fremgangsmåde ifølge et hvilket som helst foregående krav, hvor prøven er en sublingual spytprøve.
- 25 11. Fremgangsmåde ifølge et hvilket som helst foregående krav, hvor føtal distress er forårsaget af én eller flere af HELLP-syndrom, eklampsi, gestational diabetes, gestational hypertension, præmatur amniorrhesis, placenta accreta og intrauterin vækstbegrænsning.
12. Anvendelse af urinsyreniveauer i en maternel spytprøve som en indikator for føtal distress, hvor føtal distress ikke er forårsaget af præeklampsi.
- 30 13. Anvendelse af urinsyreniveauer i maternelt spyt ifølge krav 12 som en indikator for én eller flere af HELLP-syndrom, eklampsi, gestational diabetes, gestational hypertension, præmatur amniorrhesis, placenta accreta og intrauterin vækstbegrænsning.

DRAWINGS




Fig. 1




Fig. 2