



(11)

EP 2 845 856 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
11.03.2015 Bulletin 2015/11

(51) Int Cl.:
C07D 473/34 ^(2006.01) **C07D 513/04** ^(2006.01)
A61K 31/52 ^(2006.01) **A61P 29/00** ^(2006.01)
A61P 37/00 ^(2006.01)

(21) Application number: **14169702.9**

(22) Date of filing: **28.06.2010**

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO SE SI SK SM TR**
Designated Extension States:
BA ME RS

(30) Priority: **29.06.2009 US 221160 P**
10.11.2009 US 259765 P

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
10729035.5 / 2 448 938

(71) Applicant: **Incyte Corporation**
Wilmington, DE 19803 (US)

(72) Inventors:
• **Li, Yun-Long**
Chadds Ford, PA Pennsylvania 19317 (US)
• **Metcalf, Brian, W.**
Moraga, CA California 94556 (US)
• **Combs, Andrew, P.**
Kennett Square, PA Pennsylvania 19348 (US)

(74) Representative: **Carpmaels & Ransford LLP**
One Southampton Row
London WC1B 5HA (GB)

Remarks:

This application was filed on 23-05-2014 as a
divisional application to the application mentioned
under INID code 62.

(54) **Pyrimidinones as PI3K inhibitors**

(57) The present invention provides pyrimidinones that modulate activity of phosphoinositide 3-kinases (PI3Ks) and are useful in the treatment of diseases related to the activity of PI3Ks including, for example, inflammatory disorders, immune-based disorders, cancer, and other diseases.

EP 2 845 856 A1

Description

[0001] This application claims the benefit of priority of U.S. Prov. Appl. No. 61/221,160, filed on June 29, 2009, and U.S. Prov. Appl. No. 61/259,765, filed on November 10, 2009, each of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] The present invention provides pyrimidinones that modulate the activity of phosphoinositide 3-kinases (PI3Ks) and are useful in the treatment of diseases related to the activity of PI3Ks including, for example, inflammatory disorders, immune-based disorders, cancer, and other diseases.

BACKGROUND OF THE INVENTION

[0003] The phosphoinositide 3-kinases (PI3Ks) belong to a large family of lipid signaling kinases that phosphorylate phosphoinositides at the D3 position of the inositol ring (Cantley, Science, 2002, 296(5573):1655-7). PI3Ks are divided into three classes (class I, II, and III) according to their structure, regulation and substrate specificity. Class I PI3Ks, which include PI3K α , PI3K β , PI3K γ , and PI3K δ , are a family of dual specificity lipid and protein kinases that catalyze the phosphorylation of phosphatidylinosito-4,5-bisphosphate (PIP₂) giving rise to phosphatidylinosito-3,4,5-trisphosphate (PIP₃). PIP₃ functions as a second messenger that controls a number of cellular processes, including growth, survival, adhesion and migration. All four class I PI3K isoforms exist as heterodimers composed of a catalytic subunit (p110) and a tightly associated regulatory subunit that controls their expression, activation, and subcellular localization. PI3K α , PI3K β , and PI3K δ associate with a regulatory subunit known as p85 and are activated by growth factors and cytokines through a tyrosine kinase-dependent mechanism (Jimenez, et al., J Biol Chem., 2002, 277(44):41556-62) whereas PI3K γ associates with two regulatory subunits (p101 and p84) and its activation is driven by the activation of G-protein-coupled receptors (Brock, et al., J Cell Biol., 2003, 160(1):89-99). PI3K α and PI3K β are ubiquitously expressed. In contrast, PI3K γ and PI3K δ are predominantly expressed in leukocytes (Vanhaesebroeck, et al., Trends Biochem Sci., 2005, 30(4):194-204).

[0004] The differential tissue distribution of the PI3K isoforms factors in their distinct biological functions. Genetic ablation of either PI3K α or PI3K β results in embryonic lethality, indicating that PI3K α and PI3K β have essential and non-redundant functions, at least during development (Vanhaesebroeck, et al., 2005). In contrast, mice which lack PI3K γ and PI3K δ are viable, fertile and have a normal life span although they show an altered immune system. PI3K γ deficiency leads to impaired recruitment of macrophages and neutrophils to sites of inflammation as well as impaired T cell activation (Sasaki, et al., Science, 2000, 287(5455):1040-6). PI3K δ -mutant mice have specific defects in B cell signaling that lead to impaired B cell development and reduced antibody responses after antigen stimulation (Clayton, et al., J. Exp. Med. 2002, 196(6):753-63; Jou, et al., Mol. Cell Biol. 2002, 22(24):8580-91; Okkenhaug, et al., Science, 2002, 297(5583):1031-4).

[0005] The phenotypes of the PI3K γ and PI3K δ -mutant mice suggest that these enzymes may play a role in inflammation and other immune-based diseases and this is borne out in preclinical models. PI3K γ -mutant mice are largely protected from disease in mouse models of rheumatoid arthritis (RA) and asthma (Camps, et al., Nat Med. 2005, 11(9):936-43; Thomas, et al., Eur. J. Immunol., 2005, 35(4):1283-91). In addition, treatment of wild-type mice with a selective inhibitor of PI3K γ was shown to reduce glomerulonephritis and prolong survival in the MRL-lpr model of systemic lupus nephritis (SLE) and to suppress joint inflammation and damage in models of RA (Barber, et al., Nat Med. 2005, 11(9):933-5; Camps, et al., 2005). Similarly, both PI3K δ -mutant mice and wild-type mice treated with a selective inhibitor of PI3K δ have been shown to have attenuated allergic airway inflammation and hyper-responsiveness in a mouse model of asthma (Ali, et al., Nature. 2004, 431(7011):1007-11; Lee, et al., FASEB J. 2006, 20(3):455-65) and to have attenuated disease in a model of RA (Randis, et al., Eur. J. Immunol., 2008,38(5):1215-24).


[0006] In addition to their potential role in inflammatory diseases, all four class I PI3K isoforms may play a role in cancer. The gene encoding p110 α is mutated frequently in common cancers, including breast, prostate, colon and endometrial (Samuels, et al., Science, 2004, 304(5670):554; Samuels, et al., Curr. Opin. Oncol. 2006, 18(1):77-82). Eighty percent of these mutations are represented by one of three amino acid substitutions in the helical or kinase domains of the enzyme and lead to a significant upregulation of kinase activity resulting in oncogenic transformation in cell culture and in animal models (Kang, et al., Proc. Natl. Acad. Sci. U.S.A. 2005, 102(3):802-7; Bader, et al., Proc. Natl. Acad. Sci. U.S.A. 2006, 103(5):1475-9). No such mutations have been identified in the other PI3K isoforms although there is evidence that they can contribute to the development and progression of malignancies. Consistent overexpression of PI3K δ is observed in acute myeloblastic leukemia (Sujobert, et al., Blood, 2005, 106(3):1063-6) and inhibitors of PI3K δ can prevent the growth of leukemic cells (Billottet, et al., Oncogene. 2006,25(50):6648-59). Elevated expression of PI3K γ is seen in chronic myeloid leukemia (Hickey, et al., J. Biol. Chem. 2006,281(5):2441-50). Alterations in expression of

PI3K β , PI3K γ and PI3K δ have also been observed in cancers of the brain, colon and bladder (Benistant, et al., Oncogene, 2000, 19(44):5083-90; Mizoguchi, et al., Brain Pathol. 2004, 14(4):372-7; Knobbe, et al., Neuropathol. Appl. Neurobiol. 2005, 31(5):486-90). Further, these isoforms have all been shown to be oncogenic in cell culture (Kang, et al., 2006).

[0007] Thus, new or improved agents which inhibit kinases such as PI3K are continually needed for developing new and more effective pharmaceuticals that are aimed at augmentation or suppression of the immune and inflammatory pathways (such as immunosuppressive agents for organ transplants), as well as agents for the prevention and treatment of autoimmune diseases (e.g., multiple sclerosis, rheumatoid arthritis, asthma, type I diabetes, inflammatory bowel disease, Crohn's disease, autoimmune thyroid disorders, Alzheimer's disease, nephritis), diseases involving a hyper-active inflammatory response (e.g., eczema), allergies, lung diseases, cancer (e.g., prostate, breast, leukemia, multiple myeloma), and some immune reactions (e.g., skin rash or contact dermatitis or diarrhea) caused by other therapeutics. The compounds, compositions, and methods described herein are directed toward these needs and other ends.

SUMMARY OF THE INVENTION

[0008] The present invention provides, *inter alia*, compounds of Formula I or II:



or pharmaceutically acceptable salts thereof, wherein constituent members are defined herein.

[0009] The present invention further provides pharmaceutical compositions comprising a compound of Formula I or II, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.

[0010] The present invention further provides methods of modulating an activity of one or more kinases (such as a PI3K) comprising contacting the kinase with a compound of Formula I or II, or a pharmaceutically acceptable salt thereof.

[0011] The present invention further provides methods of treating a disease or a disorder associated with abnormal kinase expression or activity in a patient by administering to a patient a therapeutically effective amount of a compound of Formula I or II, or a pharmaceutically acceptable salt of the same.

[0012] The present invention further provides methods of treating diseases such as immune-based diseases, cancer, and lung diseases in a patient by administering to the patient a therapeutically effective amount of a compound of Formula I or II, or a pharmaceutically acceptable salt thereof.

[0013] The present invention further provides a compound of Formula I or II, or a pharmaceutically acceptable salt thereof, for use in therapy.

[0014] The present invention further provides use of a compound of Formula I or II, or a pharmaceutically acceptable salt thereof, for the production of a medicament for use in therapy.

BRIEF DESCRIPTION OF THE DRAWINGS



[0015]

Figure 1. X-ray crystal structure of Example 15, step 5.

Figure 2. X-ray crystal structure lattice of Example 15, step 5.

DETAILED DESCRIPTION

[0016] The present invention provides, *inter alia*, compounds that modulate the activity of one or more PI3Ks and are useful, for example, in the treatment of various diseases such as those associated with expression or activity of one or more PI3Ks. The compounds of the invention include those of Formula I or II:



or pharmaceutically acceptable salts thereof, wherein:

A is C₁₋₁₀ alkyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

L is absent, (CR^{7a}R^{7b})_m, (CR^{7a}R^{7b})_pO(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(O)(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(O)₂(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(O)NR^{7c}(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}S(O)₂(CR^{7a}R^{7b})_q, or (CR^{7a}R^{7b})_pNR^{7c}C(=NR^{7d})NR^{7c}(CR^{7a}R^{7b})_q;

W is N or CR³;

X is N or CR⁴;

Y is N or CR⁵;

Z is N or CR⁶;

G is O, S, or NR^N;

R¹ is OR^a, SR^a, S(O)R^a, S(O)₂R^a, NR^aR^b, NR^cC(O)NR^aR^b, NR^cC(O)OR^a, NR^cC(=NR^e)NR^aR^b, NR^cS(O)₂R^a, NR^cS(O)₂NR^cR^a, heterocycloalkyl, or heteroaryl, wherein the heterocycloalkyl or heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

R^{2a} and R^{2b} are independently selected from H, halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

or R^{2a} and R^{2b} together with the carbon atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered cycloalkyl ring or a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl ring, each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)₂NR^{c3}R^{d3}, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3}; R^{7a} and R^{7b} are independently selected from H, halo, or C₁₋₄ alkyl;

R^{7c} is H or C₁₋₄ alkyl;

R^{7d} is H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5Rd5}, or C(O)NR^{c5Rd5};

R^A is heteroaryl, heterocycloalkyl, heteroarylalkyl, or heterocycloalkylalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1Rd1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1Rd1}, C(=NR^e)NR^{c1Rd1}, NR^{c1C(=NR^e)NR^{c1Rd1}, NR^{c1Rd1}, NR^{c1C(O)R^{b1}}, NR^{c1C(O)OR^{a1}}, NR^{c1C(O)NR^{c1Rd1}}, NR^{c1S(O)R^{b1}}, NR^{c1S(O)₂R^{b1}}, NR^{c1S(O)₂NR^{c1Rd1}}, S(O)R^{b1}, S(O)NR^{c1Rd1}, S(O)₂R^{b1}, and S(O)₂NR^{c1Rd1};}

R^B and R^C are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

R^E is H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5Rd5}, or C(O)NR^{c5Rd5};

R^N is H or C₁₋₄ alkyl;

Cy¹ is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1Rd1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1Rd1}, C(=NR^e)NR^{c1Rd1}, NR^{c1C(=NR^e)NR^{c1Rd1}, NR^{c1R^{a1}}, NR^{c1C(O)R^{b1}}, NR^{c1C(O)OR^{a1}}, NR^{c1C(O)NR^{c1Rd1}}, NR^{c1S(O)R^{b1}}, NR^{c1S(O)₂R^{b1}}, NR^{c1S(O)₂NR^{c1Rd1}}, S(O)R^{b1}, S(O)NR^{c1Rd1}, S(O)₂R^{b1}, and S(O)₂NR^{c1Rd1};}

R^a, R^b, R^c, and R^d are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

or R^c and R^d together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

R^{a1}, R^{b1}, R^{c1}, and R^{d1} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

R^{a2}, R^{b2}, R^{c2}, and R^{d2} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

or R^{c2} and R^{d2} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5Rd5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5Rd5}, NR^{c5Rd5}, NR^{c5C(O)R^{b5}}, NR^{c5C(O)NR^{c5Rd5}}, NR^{c5C(O)OR^{a5}}, C(=NR^f)NR^{c5Rd5}, NR^{c5C(=NR^f)NR^{c5Rd5}}, S(O)R^{b5}, S(O)NR^{c5Rd5}, S(O)₂R^{b5}, NR^{c5S(O)₂R^{b5}}, NR^{c5S(O)₂NR^{c5Rd5}}, and S(O)₂NR^{c5Rd5};

R^{a3}, R^{b3}, R^{c3}, and R^{d3} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl,

heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5}; or R^{c3} and R^{d3} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, halo, C₁₋₆ haloalkyl, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5}; R^e and R^f are independently selected from H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, and C(O)NR^{c5}R^{d5}; R^{a5}, R^{b5}, R^{c5}, and R^{d5} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents. independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy; or R^{c5} and R^{d5} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy; m is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10; p is 0, 1, 2, 3, or 4; q is 0, 1, 2, 3, or 4; and r is 0 or 1.

[0017] In some embodiments, the compounds of the invention have Formula I.

[0018] In some embodiments, the compounds of the invention have Formula II.

[0019] In some embodiments, A is C₁₋₁₀ alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

[0020] In some embodiments, A is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

[0021] In some embodiments, A is aryl or heteroaryl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b,

$\text{NR}^c\text{C}(\text{O})\text{OR}^a$, $\text{NR}^c\text{C}(\text{O})\text{NR}^c\text{R}^d$, $\text{NR}^c\text{S}(\text{O})\text{R}^b$, $\text{NR}^c\text{S}(\text{O})_2\text{R}^b$, $\text{NR}^c\text{S}(\text{O})_2\text{NR}^c\text{R}^d$, $\text{S}(\text{O})\text{R}^b$, $\text{S}(\text{O})\text{NR}^c\text{R}^d$, $\text{S}(\text{O})_2\text{R}^b$, and $\text{S}(\text{O})_2\text{NR}^c\text{R}^d$.

[0022] In some embodiments, L is absent.

[0023] In some embodiments, L is $(\text{CR}^{7a}\text{R}^{7b})_m$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{O}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{S}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{S}(\text{O})(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{S}(\text{O})_2(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{C}(\text{O})\text{NR}^{7c}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{C}(\text{O})\text{O}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{C}(=\text{NR}^{7d})\text{NR}^{7c}(\text{CR}^{7a}\text{R}^{7b})_q$, or $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{S}(\text{O})_2(\text{CR}^{7a}\text{R}^{7b})_q$.

[0024] In some embodiments, L is $(\text{CR}^{7a}\text{R}^{7b})_m$.

[0025] In some embodiments, W is N.

[0026] In some embodiments, W is CR^3 .

[0027] In some embodiments, X is N.

[0028] In some embodiments, X is CR^4 .

[0029] In some embodiments, Y is N.

[0030] In some embodiments, Y is CR^5 .

[0031] In some embodiments, Z is N.

[0032] In some embodiments, Z is CR^6 .

[0033] In some embodiments, not more than 2 of W, X, Y, or Z are N.

[0034] In some embodiments, not more than 3 of W, X, Y, or Z are N.

[0035] In some embodiments, one of W and X is N.

[0036] In some embodiments, G is O.

[0037] In some embodiments, G is S.

[0038] In some embodiments, G is NR^N .

[0039] In some embodiments, R^1 is heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from $-(\text{C}_{1-4}\text{ alkyl})_r\text{-Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $\text{C}(\text{O})\text{R}^{b1}$, $\text{C}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{C}(\text{O})\text{OR}^{a1}$, $\text{OC}(\text{O})\text{R}^{b1}$, $\text{OC}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{C}(=\text{NR}^e)\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}(\text{O})\text{R}^{b1}$, $\text{NR}^{c1}\text{C}(\text{O})\text{OR}^{a1}$, $\text{NR}^{c1}\text{C}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}\text{S}(\text{O})\text{R}^{b1}$, $\text{NR}^{c1}\text{S}(\text{O})_2\text{R}^{b1}$, $\text{NR}^{c1}\text{S}(\text{O})_2\text{NR}^{c1}\text{R}^{d1}$, $\text{S}(\text{O})\text{R}^{b1}$, $\text{S}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{S}(\text{O})_2\text{R}^{b1}$, and $\text{S}(\text{O})_2\text{NR}^{c1}\text{R}^{d1}$.

[0040] In some embodiments, R^1 is NR^AR^B .

[0041] In some embodiments, R^{2a} and R^{2b} are independently selected from H and C_{1-6} alkyl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a2} , SR^{a2} , $\text{C}(\text{O})\text{R}^{b2}$, $\text{C}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{C}(\text{O})\text{OR}^{a2}$, $\text{OC}(\text{O})\text{R}^{b2}$, $\text{OC}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{C}(=\text{NR}^e)\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{C}(=\text{NR}^e)\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{C}(\text{O})\text{R}^{b2}$, $\text{NR}^{c2}\text{C}(\text{O})\text{OR}^{a2}$, $\text{NR}^{c2}\text{C}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{S}(\text{O})\text{R}^{b2}$, $\text{NR}^{c2}\text{S}(\text{O})_2\text{R}^{b2}$, $\text{NR}^{c2}\text{S}(\text{O})_2\text{NR}^{c2}\text{R}^{d2}$, $\text{S}(\text{O})\text{R}^{b2}$, $\text{S}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{S}(\text{O})_2\text{R}^{b2}$, and $\text{S}(\text{O})_2\text{NR}^{c2}\text{R}^{d2}$.

[0042] In some embodiments, at least one of R^{2a} and R^{2b} is other than H.

[0043] In some embodiments, R^3 , R^4 , R^5 , and R^6 are independently selected from H or C_{1-6} alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3} , SR^{a3} , $\text{C}(\text{O})\text{R}^{b3}$, $\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{C}(\text{O})\text{OR}^{a3}$, $\text{OC}(\text{O})\text{R}^{b3}$, $\text{OC}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{R}^{b3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{OR}^{a3}$, $\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})\text{R}^{b3}$, $\text{S}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$, and $\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$.

[0044] In some embodiments, R^3 is C_{1-6} alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3} , SR^{a3} , $\text{C}(\text{O})\text{R}^{b3}$, $\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{C}(\text{O})\text{OR}^{a3}$, $\text{OC}(\text{O})\text{R}^{b3}$, $\text{OC}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{R}^{b3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{OR}^{a3}$, $\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})\text{R}^{b3}$, $\text{S}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$, and $\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$.

[0045] In some embodiments, R^4 , R^5 , and R^6 are each H.

[0046] In some embodiments, R^{7a} , R^{7b} , R^{7c} , and R^{7d} are each H.

[0047] In some embodiments, R^A is heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $\text{C}(\text{O})\text{R}^{b5}$, $\text{C}(\text{O})\text{NR}^{c5}\text{R}^{d5}$, $\text{C}(\text{O})\text{OR}^{a5}$, $\text{OC}(\text{O})\text{R}^{b5}$, $\text{OC}(\text{O})\text{NR}^{c5}\text{R}^{d5}$, $\text{NR}^{c5}\text{R}^{d5}$, $\text{NR}^{c5}\text{C}(\text{O})\text{R}^{b5}$, $\text{NR}^{c5}\text{C}(\text{O})\text{NR}^{c5}\text{R}^{d5}$, $\text{NR}^{c5}\text{C}(\text{O})\text{OR}^{a5}$, $\text{C}(=\text{NR}^f)\text{NR}^{c5}\text{R}^{d5}$, $\text{NR}^{c5}\text{C}(=\text{NR}^f)\text{NR}^{c5}\text{R}^{d5}$, $\text{S}(\text{O})\text{R}^{b5}$, $\text{S}(\text{O})\text{NR}^{c5}\text{R}^{d5}$, $\text{S}(\text{O})_2\text{R}^{b5}$, $\text{NR}^{c5}\text{S}(\text{O})_2\text{R}^{b5}$, $\text{NR}^{c5}\text{S}(\text{O})_2\text{NR}^{c5}\text{R}^{d5}$, and $\text{S}(\text{O})_2\text{NR}^{c5}\text{R}^{d5}$.




[0048] In some embodiments, R^B and R^C are independently selected from H and C_{1-6} alkyl.

[0049] In some embodiments, A is other than phenyl substituted at the 4-position by halogen.


[0050] In some embodiments, r is 0.

[0051] In some embodiments, r is 1.

[0052] In some embodiments, the compounds of the invention have Formula Ia, Ib, Ic, Id, Ie, IIa, IIb, or IIc:



[0053] In some embodiments, the compounds of the invention have Formula If or IId:



[0054] In some embodiments of compounds of Formulas I, Ia-If, II, and Ila-IId:

A is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

R¹ is OR^a, SR^a, S(O)R^a, S(O)₂R^a, NR^aR^b, NR^cC(O)NR^aR^b, NR^cC(O)OR^a, NR^cC(=NR^e)NR^aR^b, NR^cS(O)₂R^a, NR^cS(O)₂NR^cR^a, heterocycloalkyl, or heteroaryl, wherein the heterocycloalkyl or heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

R^{2a} is H, halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2},

$C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $C(=NR^e)NR^{c2}R^{d2}$, $NR^{c2}C(=NR^e)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, and $S(O)_2NR^{c2}R^{d2}$;

R^3 , R^4 , R^5 , and R^6 are independently selected from H, halo, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)_2NR^{c3}R^{d3}$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}C(O)OR^{a3}$, $C(=NR^e)NR^{c3}R^{d3}$, $NR^{c3}C(=NR^e)NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, and $S(O)_2NR^{c3}R^{d3}$; R^A is heteroaryl, heterocycloalkyl, heteroarylalkyl, or heterocycloalkylalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4} \text{ alkyl})_R-Cy^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^B and R^C are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

R^E is H, CN, NO_2 , OR^{a5} , SR^{b5} , $S(O)_2R^{b5}$, $C(O)R^{b5}$, $S(O)_2NR^{c5}R^{d5}$, or $C(O)NR^{c5}R^{d5}$;

Cy^1 is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^a , R^b , R^c , and R^d are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

or R^c and R^d together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

R^{a1} , R^{b1} , R^{c1} , and R^{d1} are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

R^{a2} , R^{b2} , R^{c2} , and R^{d2} are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$,

$$\text{NR}^{\text{c5}}\text{R}^{\text{d5}}, \text{NR}^{\text{c5}}\text{C}(\text{O})\text{R}^{\text{b5}}, \text{NR}^{\text{c5}}\text{C}(\text{O})\text{NR}^{\text{c5}}\text{R}^{\text{d5}}, \text{NR}^{\text{c5}}\text{C}(\text{O})\text{OR}^{\text{a5}}, \text{C}(\text{=NR}^{\text{f}})\text{NR}^{\text{c5}}\text{R}^{\text{d5}}, \text{NR}^{\text{c5}}\text{C}(\text{=NR}^{\text{f}})\text{NR}^{\text{c5}}\text{R}^{\text{d5}}, \text{S}(\text{O})\text{R}^{\text{b5}}, \text{S}(\text{O})\text{NR}^{\text{c5}}\text{R}^{\text{d5}}, \text{S}(\text{O})_2\text{R}^{\text{b5}}, \text{NR}^{\text{c5}}\text{S}(\text{O})_2\text{R}^{\text{b5}}, \text{NR}^{\text{c5}}\text{S}(\text{O})_2\text{NR}^{\text{c5}}\text{R}^{\text{d5}}, \text{and } \text{S}(\text{O})_2\text{NR}^{\text{c5}}\text{R}^{\text{d5}}.$$

or R^{c2} and R^{d2} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆-haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^{a3}, R^{b3}, R^{c3}, and R^{d3} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5}.

or R^{c3} and R^{d3} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, halo, C₁₋₆ haloalkyl, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^e and R^f are independently selected from H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, and C(O)NR^{c5}R^{d5}.

R^{a5}, R^{b5}, R^{c5}, and R^{d5} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy;

or R^{c5} and R^{d5} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy; and r is 0 or 1.

[0055] In some embodiments of compounds of Formulas I and Ia-If, when A is 3-fluorophenyl; R^{2a} is H; R³ is methyl; and R⁴, R⁵, and R⁶ are H; then R¹ is other than 4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl.

[0056] In some embodiments of compounds of Formulas I and Ia-If, when A is 1,3-dioxolan-2-yl; R^{2a}, R³, R⁴, and R⁶ are H; then R¹ is other than 1-(*tert*-butoxycarbonyl)piperidin-4-yl.

[0057] In some embodiments of compounds of Formulas I and Ia-If, R¹ is other than a substituted or unsubstituted pyrazolo[3,4-d]pyrimidin-1-yl group.

[0058] In some embodiments of compounds of Formulas I and Ia-If, R¹ is other than a substituted or unsubstituted piperidinyl group.

[0059] In some embodiments, the compound has Formula Ia.

[0060] In some embodiments, the compound has Formula Ib.

[0061] In some embodiments, the compound has Formula Ic.

[0062] In some embodiments, the compound has Formula Id.

[0063] In some embodiments, the compound has Formula 1e.

[0064] In some embodiments, the compound has Formula If.

[0065] In some embodiments, the compound has Formula IIa.

[0066] In some embodiments, the compound has Formula IIb.

[0067] In some embodiments, the compound has Formula IIc.

[0068] In some embodiments, the compound has Formula IIc.

[0069] In some embodiments of compounds of Formulas I, Ia-If, II, and IIa-IId, A is cycloalkyl or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d,

substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein the C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

[0079] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, A is pyrazolyl optionally substituted with 1 or 2 substituents independently selected from halo, C₁₋₆ alkyl, or C₁₋₆ haloalkyl.

[0080] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R¹ is heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

[0081] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R¹ is bicyclic heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

[0082] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R¹ is purinyl optionally substituted with -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, or S(O)₂NR^{c1}R^{d1}.

[0083] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R¹ is OR^A, SR^A, S(O)R^A, S(O)₂R^A, NR^AR^B, NR^CC(O)NR^AR^B, NR^CC(O)OR^A, NR^CC(=NR^E)NR^AR^B, NR^CS(O)₂R^A, or NR^CS(O)₂NR^CR^A.

[0084] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R¹ is NR^AR^B.


[0085] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^A is heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

[0086] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^A is bicyclic heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.


[0087] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^A is purinyl optionally substituted with 1 or 2 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

[0088] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^A is purinyl optionally substituted with 1 or 2 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5}.

[0089] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^A is:



[0090] In some embodiments, R^A is selected from:



[0091] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^A is bicyclic heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4} \text{ alkyl})_r$ -Cy¹, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

[0092] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^B and R^C are independently selected from H and C_{1-6} alkyl.

[0093] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^B and R^C are each H.

[0094] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^{2a} is H, halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl, wherein the C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2}.

[0095] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^{2a} is halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl, wherein the C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2}.

[0096] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^{2a} is H, halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl.

[0097] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^{2a} is C_{1-6} alkyl.

[0098] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^{2a} is methyl or ethyl.

[0099] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^3 is halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)₂NR^{c3}R^{d3}, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein the C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3}.

[0100] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^3 is H, halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, S(O)₂NR^{c3}R^{d3}, or C_{1-6} alkyl, wherein the C_{1-6} alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

[0101] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^3 is H or C_{1-6} alkyl.

[0102] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^3 is C_{1-6} alkyl.

[0103] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^3 is methyl.

[0104] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^4 , R^5 , and R^6 are independently selected from H, halo, C_{1-6} alkyl, or C_{1-6} haloalkyl.


[0105] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^4 , R^5 , and R^6 are each H.

[0106] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R^4 is H.

[0107] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R⁵ is H.


[0108] In some embodiments of compounds of Formulas I, Ia-Ic, II, and IIa-IIc, R⁶ is H.

[0109] In some embodiments, the compounds of the invention have Formula Ig or IIf:




[0110] In some embodiments of compounds of Formulas Ig and IIf, R¹ is according to any of the previously recited embodiments for R¹.

[0111] In some embodiments, the compounds of the invention have Formula Ih or IIIf:



[0112] In some embodiments, of compounds of Formulas Ih and IIIf, R^A is according to any of the previously recited embodiments for R^A.

[0113] In some embodiments, the compounds of the invention have Formula Ii or IIg:



wherein R⁸ and R⁹ are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5}.

[0114] In some embodiments, R⁸ and R⁹ are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, NR^{c5}S(O)₂R^{b5}, and NR^{c5}S(O)₂NR^{c5}R^{d5}.

[0115] In some embodiments, R⁸ and R⁹ are independently selected from H and C₁₋₆ alkyl.

[0116] In some embodiments of compounds of Formulas Ig-Ii and IIf-IIg, A is according to any of the previously recited embodiments for A.

[0117] In some embodiments of compounds of Formulas Ig-Ii and IIf-IIg, R³ is according to any of the previously recited embodiments for R³.

[0118] In some embodiments of compounds of Formulas Ig-Ii and IIf-IIg, R^{2a} is according to any of the previously recited embodiments for R^{2a}.

[0119] In some embodiments of compounds of Formulas Ig-Ii and IIf-IIg, R^B is according to any of the previously recited embodiments for R^B.

[0120] In some embodiments of compounds of Formula Ii or IIg:

A is aryl or heteroaryl, each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and OR^a;

R¹ is NR^AR^B or heteroaryl; wherein the heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and OR^{a1};

R^{2a} is C₁₋₆ alkyl;

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, and C₁₋₆ alkyl; wherein the C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo;

R^A is heteroaryl, which is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, CN, OR^{a1}, and NR^{c1}R^{d1};

R^B is H;

each R^a is independently selected from H and C₁₋₆ alkyl;


each R^{a1}, R^{c1}, and R^{d1} is independently selected from H and C₁₋₆ alkyl, wherein the C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo; and

each R^{a3} is independently selected from H and C₁₋₆ alkyl, wherein the C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo.

[0121] In some embodiments of compounds of Formula If or IId:

A is phenyl, 5-membered heteroaryl or 6-membered heteroaryl, each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and OR^a;



R¹ is NR^AR^B or



R^{2a} is C₁₋₆ alkyl;

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, and C₁₋₆ alkyl; wherein the C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo;

R^A is selected from:



R^B is H;

each R^a is independently selected from H and C₁₋₆ alkyl; and

each R^{a3} is independently selected from H and C₁₋₆ alkyl, wherein the C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo.

[0122] It is further appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

[0123] At various places in the present specification, linking substituents are described. It is specifically intended that each linking substituent include both the forward and backward forms of the linking substituent. For example, -NR(CR'R'')_n- includes both -NR(CR'R'')_n- and -(CR'R'')_nNR-. Where the structure clearly requires a linking group, the Markush variables listed for that group are understood to be linking groups. For example, if the structure requires a linking group and the

Markush group definition for that variable lists "alkyl" or "aryl" then it is understood that the "alkyl" or "aryl" represents a linking alkylene group or arylene group, respectively.

[0124] The term "n-membered" where n is an integer typically describes the number of ring-forming atoms in a moiety where the number of ring-forming atoms is n. For example, piperidinyl is an example of a 6-membered heterocycloalkyl ring, pyrazolyl is an example of a 5-membered heteroaryl ring, pyridyl is an example of a 6-membered heteroaryl ring, and 1,2,3,4-tetrahydro-naphthalene is an example of a 10-membered cycloalkyl group.

[0125] As used herein, the term "alkyl" is meant to refer to a saturated hydrocarbon group which is straight-chained or branched. Example alkyl groups include methyl (Me), ethyl (Et), propyl (e.g., n-propyl and isopropyl), butyl (e.g., n-butyl, isobutyl, sec-butyl, t-butyl), pentyl (e.g., n-pentyl, isopentyl, sec-pentyl, neopentyl), and the like. An alkyl group can contain from 1 to about 20, from 2 to about 20, from 1 to about 10, from 1 to about 8, from 1 to about 6, from 1 to about 4, or from 1 to about 3 carbon atoms.

[0126] As used herein, "alkenyl" refers to an alkyl group having one or more carbon-carbon double bonds. Example alkenyl groups include ethenyl, propenyl, cyclohexenyl, and the like.

[0127] As used herein, "alkynyl" refers to an alkyl group having one or more carbon-carbon triple bonds. Example alkynyl groups include ethynyl, propynyl, and the like.

[0128] As used herein, "haloalkyl" refers to an alkyl group having one or more halogen substituents. Example haloalkyl groups include CF₃, C₂F₅, CHF₂, CCl₃, CHCl₂, C₂Cl₅, and the like.

[0129] As used herein, "halosulfanyl" refers to a sulfur group having one or more halogen substituents. Example halosulfanyl groups include pentahalosulfanyl groups such as SF₅.

[0130] As used herein, "aryl" refers to monocyclic or polycyclic (e.g., having 2, 3 or 4 fused rings) aromatic hydrocarbons such as, for example, phenyl, naphthyl, anthracenyl, phenanthrenyl, indanyl, indenyl, and the like. In some embodiments, aryl groups have from 6 to about 20 carbon atoms.

[0131] As used herein, "cycloalkyl" refers to non-aromatic cyclic hydrocarbons including cyclized alkyl, alkenyl, and alkynyl groups. Cycloalkyl groups can include mono- or polycyclic (e.g., having 2, 3 or 4 fused rings) groups and spiro-cycles. Ring-forming carbon atoms of a cycloalkyl group can be optionally substituted by oxo or sulfido. Cycloalkyl groups also include cycloalkylidenes. Example cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclohexadienyl, cycloheptatrienyl, norbornyl, norpinyl, norcamyl, adamantyl, and the like. Also included in the definition of cycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the cycloalkyl ring, for example, benzo or thienyl derivatives of cyclopentane, cyclopentene, cyclohexane, and the like. A cycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom including a ring-forming atom of the fused aromatic ring.

[0132] As used herein, "heteroaryl" refers to an aromatic heterocycle having at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include monocyclic and polycyclic (e.g., having 2, 3 or 4 fused rings) systems. Examples of heteroaryl groups include without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrol, oxazolyl, benzofuryl, benzothienyl, benzothiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-thiadiazolyl, isothiazolyl, benzothienyl, purinyl, carbazolyl, benzimidazolyl, indolyl, and the like. Examples of bicyclic heteroaryl groups include without limitation, purinyl, indolyl, and the like. In some embodiments, any ring-forming N in a heteroaryl moiety can be substituted by oxo. In some embodiments, the heteroaryl group has from 1 to about 20 carbon atoms, and in further embodiments from about 3 to about 20 carbon atoms. In some embodiments, the heteroaryl group contains 3 to about 14, 4 to about 14, 9 to about 10, or 5 to 6 ring-forming atoms. In some embodiments, the heteroaryl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms.

[0133] As used herein, "heterocycloalkyl" refers to non-aromatic heterocycles having one or more ring-forming heteroatoms such as an O, N, or S atom. Heterocycloalkyl groups include monocyclic and polycyclic (e.g., having 2, 3 or 4 fused rings) systems as well as spirocycles. Example "heterocycloalkyl" groups include morpholino, thiomorpholino, piperazinyl, tetrahydrofuranyl, tetrahydrothienyl, 2,3-dihydrobenzofuryl, 1,3-benzodioxole, benzo-1,4-dioxane, piperidinyl, pyrrolidinyl, isoxazolidinyl, isothiazolidinyl, pyrazolidinyl, oxazolidinyl, thiazolidinyl, imidazolidinyl, and the like. Ring-forming carbon atoms and heteroatoms of a heterocycloalkyl group can be optionally substituted by oxo or sulfido. Also included in the definition of heterocycloalkyl are moieties that have one or more aromatic rings fused (i.e., having a bond in common with) to the nonaromatic heterocyclic ring, for example phthalimidyl, naphthalimidyl, and benzo derivatives of heterocycles. The heterocycloalkyl group can be attached through a ring-forming carbon atom or a ring-forming heteroatom. The heterocycloalkyl group containing a fused aromatic ring can be attached through any ring-forming atom including a ring-forming atom of the fused aromatic ring. In some embodiments, the heterocycloalkyl group has from 1 to about 20 carbon atoms, and in further embodiments from about 3 to about 20 carbon atoms. In some embodiments, the heterocycloalkyl group contains 3 to about 14, 4 to about 14, 3 to about 7, or 5 to 6 ring-forming atoms. In some embodiments, the heterocycloalkyl group has 1 to about 4, 1 to about 3, or 1 to 2 heteroatoms. In some embodiments, the heterocycloalkyl group contains 0 to 3 double or triple bonds. In some embodiments, the heterocycloalkyl group contains 0 to 2 double or triple bonds.

[0134] As used herein, "halo" or "halogen" includes fluoro, chloro, bromo, and iodo.

[0135] As used herein, "arylalkyl" refers to alkyl substituted by aryl and "cycloalkylalkyl" refers to alkyl substituted by cycloalkyl. An example arylalkyl group is benzyl.

[0136] As used herein, "heteroarylalkyl" refers to alkyl substituted by heteroaryl and "heterocycloalkylalkyl" refers to alkyl substituted by heterocycloalkyl.

[0137] As used herein, "amino" refers to NH_2 .

[0138] As used herein, "alkoxy" refers to an -O-alkyl group. Example alkoxy groups include methoxy, ethoxy, propoxy (e.g., n-propoxy and isopropoxy), t-butoxy, and the like.

[0139] As used herein, "haloalkoxy" refers to an -O-(haloalkyl) group.

[0140] As used herein, "alkylthio" refers to an -S-alkyl group. Example alkylthio groups include meththio, ethylthio, propylthio (e.g., n-propylthio and isopropylthio), and the like.

[0141] As used herein, "alkylamino" refers to an -NH-alkyl group. Example alkylamino groups include methylamino, ethylamino, propylamino (e.g., n-propylamino and isopropylamino), and the like.

[0142] As used herein, "di(alkyl)amino" refers to an -N(alkyl)₂ group. Example di(alkyl)amino groups include dimethylamino, diethylamino, dipropylamino (e.g., di(n-propyl)amino and di(isopropyl)amino), and the like.


[0143] It should be further appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, can also be provided in combination in a single embodiment. Conversely, various features of the invention which are, for brevity, described in the context of a single embodiment, can also be provided separately or in any suitable subcombination.

[0144] The compounds described herein can be asymmetric (e.g., having one or more stereocenters). All stereoisomers, such as enantiomers and diastereomers, are intended unless otherwise indicated. Compounds of the present invention that contain asymmetrically substituted carbon atoms can be isolated in optically active or racemic forms. Methods on how to prepare optically active forms from optically inactive starting materials are known in the art, such as by resolution of racemic mixtures or by stereoselective synthesis. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms.

[0145] Resolution of racemic mixtures of compounds can be carried out by any of numerous methods known in the art. An example method includes fractional recrystallization using a chiral resolving acid which is an optically active, salt-forming organic acid. Suitable resolving agents for fractional recrystallization methods are, for example, optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids such as β -camphorsulfonic acid. Other resolving agents suitable for fractional crystallization methods include stereoisomerically pure forms of α -methylbenzylamine (e.g., S and R forms, or diastereomerically pure forms), 2-phenylglycinol, norephedrine, ephedrine, N-methylephedrine, cyclohexylethylamine, 1,2-diaminocyclohexane, and the like.

[0146] Resolution of racemic mixtures can also be carried out by elution on a column packed with an optically active resolving agent (e.g., dinitrobenzoylphenylglycine). Suitable elution solvent composition can be determined by one skilled in the art.

[0147] Compounds of the invention also include tautomeric forms. Tautomeric forms result from the swapping of a single bond with an adjacent double bond together with the concomitant migration of a proton. Tautomeric forms include prototropic tautomers which are isomeric protonation states having the same empirical formula and total charge. Example prototropic tautomers include ketone - enol pairs, amide - imidic acid pairs, lactam - lactim pairs, enamine - imine pairs, and annular forms where a proton can occupy two or more positions of a heterocyclic system, for example, 1H- and 3H-imidazole, 1H-, 2H- and 4H- 1,2,4-triazole, 1H- and 2H- isoindole, and 1H- and 2H-pyrazole. Tautomeric forms can be in equilibrium or sterically locked into one form by appropriate substitution. For example, purine includes the 9H and a 7H tautomeric forms:



[0148] Compounds of the invention can include both the 9H and 7H tautomeric forms.

[0149] Compounds of the invention can also include all isotopes of atoms occurring in the intermediates or final compounds. Isotopes include those atoms having the same atomic number but different mass numbers. For example, isotopes of hydrogen include tritium and deuterium.

[0150] The term, "compound," as used herein is meant to include all stereoisomers, geometric isomers, tautomers,

and isotopes of the structures depicted. Compounds herein identified by name or structure as one particular tautomeric form are intended to include other tautomeric forms unless otherwise specified.

[0151] All compounds, and pharmaceutically acceptable salts thereof, can be found together with other substances such as water and solvents (e.g. hydrates and solvates) or can be isolated.

[0152] In some embodiments, the compounds of the invention, or salts thereof, are substantially isolated. By "substantially isolated" is meant that the compound is at least partially or substantially separated from the environment in which it was formed or detected. Partial separation can include, for example, a composition enriched in the compounds of the invention. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compounds of the invention, or salt thereof. Methods for isolating compounds and their salts are routine in the art.

[0153] The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0154] The expressions, "ambient temperature" and "room temperature," as used herein, are understood in the art, and refer generally to a temperature, e.g. a reaction temperature, that is about the temperature of the room in which the reaction is carried out, for example, a temperature from about 20 °C to about 30 °C.

[0155] The present invention also includes pharmaceutically acceptable salts of the compounds described herein. As used herein, "pharmaceutically acceptable salts" refers to derivatives of the disclosed compounds wherein the parent compound is modified by converting an existing acid or base moiety to its salt form. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts of the present invention include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, non-aqueous media like ether, ethyl acetate, alcohols (e.g., methanol, ethanol, iso-propanol, or butanol) or acetonitrile (ACN) are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and Journal of Pharmaceutical Science, 66, 2 (1977), each of which is incorporated herein by reference in its entirety.

Synthesis

[0156] Compounds of the invention, including salts thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes.

[0157] The reactions for preparing compounds of the invention can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates, or products at the temperatures at which the reactions are carried out, e.g., temperatures which can range from the solvent's freezing temperature to the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.


[0158] Preparation of compounds of the invention can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and the selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups can be found, for example, in T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Synthesis, 3rd Ed., Wiley & Sons, Inc., New York (1999), which is incorporated herein by reference in its entirety.

[0159] Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (e.g., ¹H or ¹³C), infrared spectroscopy, spectrophotometry (e.g., UV-visible), mass spectrometry, or by chromatographic methods such as high performance liquid chromatography (HPLC) or thin layer chromatography (TLC).

[0160] Example synthetic methods for preparing compounds of the invention are provided in the Schemes below. For instance, compounds of the invention can be prepared by the general synthetic procedure shown in Scheme 1. Heteroaryl compounds of formula 1 can react with 4-halo-3-oxo-pentanoates 2 in the presence of polyphosphoric acid (PPA) to provide the compounds of formula 3 via a cyclocondensation reaction. These can be subjected to halogenation reaction under suitable conditions, to provide halogenated compounds 4. Compounds of formula 4 can be transformed to the compounds of formula 5 through any variation of a sequence of steps. X¹ can be replaced with either an azide, an amine, or a heterocyclic group through an S_N2 displacement and eventually transformed to the R¹ group. X² can be converted


to a desired cyclic moiety (Cy) through any of the standard cross-coupling reactions, known to one skilled in the art, e.g., using boronic acid derivatives of the desired cyclic moiety.

Scheme 1




[0161] As shown in Scheme 2, the pyrido[1,2-a]pyrimidin-4-ones of the invention can be prepared by cyclocondensation of aminopyrimidines 6 with a β -keto ester 2. Halogenation of the resultant pyridopyrimidinones 7 under suitable conditions (such as NBS or bromine) provides compounds of formula 8. The latter can be transformed to the compounds of formula 9 through an S_N2 substitution of X^1 with a heterocycle R^1 followed by a coupling reaction of X^2 with L-A moiety (such as a Negishi coupling of organozinc reagent; a Suzuki or Stille coupling of an arylboronic acid or arylstannane, respectively). Alternatively, X^1 can be replaced with either an amine or an azide which can be reduced to amine. The amine can then be subjected to coupling reaction with a R^1 moiety to give compounds of formula 9.

Scheme 2




[0162] The thiazolo[3,2-a]pyrimidin-5-ones of the invention can be prepared according to Scheme 3. 2-Aminothiazole 10 condensed with a β -keto ester 2 provides thiazolopyrimidinone 11. Compounds of formula 11 can be converted to compounds of formula 13 through any variations of a sequence of steps as described above.

Scheme 3




[0163] Alternatively, compounds of the invention can be synthesized by reacting amino heterocycles 14 with an α -substituted β -keto ester 15 shown in Scheme 4. The cyclocondensation derivatives 16 can then be subjected to halogenation (such as NBS or bromine) or oxidation (such as SeO_2) to afford halogen compounds 17 (X^1 = halogen), or alcohol compounds 17 (X = OH), respectively. Compounds of formula 17 can then be transformed to compounds of formula 18 through any variations of a sequence of steps. X^1 can be coupled directly with a heterocycles under any of the cross coupling conditions know to one skilled in the art (such as Buchwald-Hartwig cross coupling conditions) or converted to a halogen then the latter can be coupled with a heterocycles through S_N2 substitutions.

Scheme 4



[0164] Compounds of Formula II of the invention can be prepared according to Scheme 5. Amino heterocycles **19** condensed with α -substituted β -ketone ester **15** affords compounds of formula **20**. The latter can be transformed to compounds of Formula II through any variation of steps described above.

Scheme 5



[0165] It should be noted that in all of the Schemes described herein, if there are functional groups present on a substituent group, further modification can be made if appropriate and desired. For example, a CN group can be hydrolyzed to afford an amide group; a carboxylic acid can be converted to an ester, which in turn can be reduced to an alcohol, which in turn can be further modified. In another example, an OH group can be converted into a better leaving group such as mesylate, which in turn is suitable for nucleophilic substitution, such as by CN. Furthermore, an OH group can be subjected to Mitsunobu reaction conditions with phenol, or heteroaryl alcohol, to afford aryl or heteroaryl ether compounds. One skilled in the art will recognize further modifications.

[0166] It should be further noted that the reaction sequences described above can be modified to suit different target molecules. For instance, Cy-boronic acid can be reacted with **4** to generate the Suzuki product first. The X^1 group of the Suzuki product can then be further functionalized with a nucleophilic reagent such as an azide or a heterocyclic amine.

Methods

[0167] The compounds of the invention can modulate activity of one or more of various kinases including, for example, phosphoinositide 3-kinases (PI3Ks). The term "modulate" is meant to refer to an ability to increase or decrease the activity of one or more members of the PI3K family. Accordingly, the compounds of the invention can be used in methods of modulating a PI3K by contacting the PI3K with any one or more of the compounds or compositions described herein. In some embodiments, compounds of the present invention can act as inhibitors of one or more PI3Ks. In further embodiments, the compounds of the invention can be used to modulate activity of a PI3K in an individual in need of modulation of the receptor by administering a modulating amount of a compound of the invention, or a pharmaceutically acceptable salt thereof. In some embodiments, modulating is inhibiting.

[0168] Given that cancer cell growth and survival is impacted by multiple signaling pathways, the present invention is useful for treating disease states characterized by drug resistant kinase mutants. In addition, different kinase inhibitors, exhibiting different preferences in the kinases which they modulate the activities of, may be used in combination. This approach could prove highly efficient in treating disease states by targeting multiple signaling pathways, reduce the likelihood of drug-resistance arising in a cell, and reduce the toxicity of treatments for disease.

[0169] Kinases to which the present compounds bind and/or modulate (e.g., inhibit) include any member of the PI3K family. In some embodiments, the PI3K is PI3K α , PI3K β , PI3K γ , or PI3K δ . In some embodiments, the PI3K is PI3K γ or PI3K δ . In some embodiments, the PI3K is PI3K γ . In some embodiments, the PI3K is PI3K δ . In some embodiments, the PI3K includes a mutation. A mutation can be a replacement of one amino acid for another, or a deletion of one or more amino acids. In such embodiments, the mutation can be present in the kinase domain of the PI3K.

[0170] In some embodiments, more than one compound of the invention is used to inhibit the activity of one kinase (e.g., PI3K γ or PI3K δ).

[0171] In some embodiments, more than one compound of the invention is used to inhibit more than one kinase, such as at least two kinases (e.g., PI3K γ and PI3K δ).

[0172] In some embodiments, one or more of the compounds is used in combination with another kinase inhibitor to inhibit the activity of one kinase (e.g., PI3K γ or PI3K δ).

[0173] In some embodiments, one or more of the compounds is used in combination with another kinase inhibitor to inhibit the activities of more than one kinase (e.g., PI3K γ or PI3K δ), such as at least two kinases.

[0174] The compounds of the invention can be selective. By "selective" is meant that the compound binds to or inhibits a kinase with greater affinity or potency, respectively, compared to at least one other kinase. In some embodiments, the compounds of the invention are selective inhibitors of PI3K γ or PI3K δ over PI3K α and/or PI3K β . In some embodiments, the compounds of the invention are selective inhibitors of PI3K δ (e.g., over PI3K α , PI3K β and PI3K γ). In some embodiments, the compounds of the invention are selective inhibitors of PI3K γ (e.g., over PI3K α , PI3K β and PI3K δ). In some embodiments, selectivity can be at least about 2-fold, 5-fold, 10-fold, at least about 20-fold, at least about 50-fold, at least about 100-fold, at least about 200-fold, at least about 500-fold or at least about 1000-fold. Selectivity can be measured by methods routine in the art. In some embodiments, selectivity can be tested at the K_m ATP concentration of each enzyme. In some embodiments, the selectivity of compounds of the invention can be determined by cellular assays associated with particular PI3K kinase activity.

[0175] Another aspect of the present invention pertains to methods of treating a kinase (such as PI3K)-associated disease or disorder in an individual (e.g., patient) by administering to the individual in need of such treatment a therapeutically effective amount or dose of one or more compounds of the present invention or a pharmaceutical composition thereof. A PI3K-associated disease can include any disease, disorder or condition that is directly or indirectly linked to expression or activity of the PI3K, including overexpression and/or abnormal activity levels. In some embodiments, the disease can be linked to Akt (protein kinase B), mammalian target of rapamycin (mTOR), or phosphoinositide-dependent kinase 1 (PDK1). In some embodiments, the mTOR-related disease can be inflammation, atherosclerosis, psoriasis, restenosis, benign prostatic hypertrophy, bone disorders, pancreatitis, angiogenesis, diabetic retinopathy, arthritis, immunological disorders, kidney disease, or cancer. A PI3K-associated disease can also include any disease, disorder or condition that can be prevented, ameliorated, or cured by modulating PI3K activity. In some embodiments, the disease is characterized by the abnormal activity of PI3K. In some embodiments, the disease is characterized by mutant PI3K. In such embodiments, the mutation can be present in the kinase domain of the PI3K.

[0176] Examples of PI3K-associated diseases include immune-based diseases involving the system including, for example, rheumatoid arthritis, allergy, asthma, glomerulonephritis, lupus, or inflammation related to any of the above.

[0177] Further examples of PI3K-associated diseases include cancers such as breast, prostate, colon, endometrial, brain, bladder, skin, uterus, ovary, lung, pancreatic, renal, gastric, or hematological cancer.

[0178] In some embodiments, the hematological cancer is acute myeloblastic leukemia (AML) or chronic myeloid leukemia (CML), or B cell lymphoma.

[0179] Further examples of PI3K-associated diseases include lung diseases such as acute lung injury (ALI) and adult respiratory distress syndrome (ARDS).

[0180] Further examples of PI3K-associated diseases include osteoarthritis, restenosis, atherosclerosis, bone disorders, arthritis, diabetic retinopathy, psoriasis, benign prostatic hypertrophy, inflammation, angiogenesis, pancreatitis, kidney disease, inflammatory bowel disease, myasthenia gravis, multiple sclerosis, or Sjogren's syndrome, and the like.

[0181] The present invention further provides a compound described herein, or a pharmaceutically acceptable salt thereof, for use in any of the methods described herein.

[0182] The present invention further provides use of a compound described herein, or a pharmaceutically acceptable salt thereof, for the production of a medicament for use in any of the methods described herein.

[0183] As used herein, the term "contacting" refers to the bringing together of indicated moieties in an *in vitro* system or an *in vivo* system. For example, "contacting" a PI3K with a compound of the invention includes the administration of a compound of the present invention to an individual or patient, such as a human, having a PI3K, as well as, for example, introducing a compound of the invention into a sample containing a cellular or purified preparation containing the PI3K.

[0184] As used herein, the term "individual" or "patient," used interchangeably, refers to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans.

[0185] As used herein, the phrase "therapeutically effective amount" refers to the amount of active compound or pharmaceutical agent that elicits the biological or medicinal response that is being sought in a tissue, system, animal, individual or human by a researcher, veterinarian, medical doctor or other clinician.

[0186] As used herein, the term "treating" or "treatment" refers to one or more of (1) preventing the disease; for example, preventing a disease, condition or disorder in an individual who may be predisposed to the disease, condition or disorder but does not yet experience or display the pathology or symptomatology of the disease; (2) inhibiting the disease; for example, inhibiting a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., arresting further development of the pathology and/or symptomatology); and (3) ameliorating the disease; for example, ameliorating a disease, condition or disorder in an individual who is experiencing or displaying the pathology or symptomatology of the disease, condition or disorder (i.e., reversing the pathology and/or symptomatology) such as decreasing the severity of disease.

Combination Therapies

[0187] One or more additional pharmaceutical agents such as, for example, chemotherapeutics, anti-inflammatory agents, steroids, immunosuppressants, as well as Bcr-Ab1, Flt-3, EGFR, HER2, JAK, c-MET, VEGFR, PDGFR, cKit, IGF-1R, RAF and FAK kinase inhibitors such as, for example, those described in WO 2006/056399, or other agents such as, therapeutic antibodies can be used in combination with the compounds of the present invention for treatment of PI3K-associated diseases, disorders or conditions. The one or more additional pharmaceutical agents can be administered to a patient simultaneously or sequentially.

[0188] Example antibodies for use in combination therapy include but are not limited to Trastuzumab (e.g. anti-HER2), Ranibizumab (e.g. anti-VEGF-A), Bevacizumab (trade name Avastin, e.g. anti-VEGF), Panitumumab (e.g. anti-EGFR), Cetuximab (e.g. anti-EGFR), Rituxan (anti-CD20) and antibodies directed to c-MET.

[0189] One or more of the following agents may be used in combination with the compounds of the present invention and are presented as a non limiting list: a cytostatic agent, cisplatin, doxorubicin, taxotere, taxol, etoposide, irinotecan, camptostar, topotecan, paclitaxel, docetaxel, epothilones, tamoxifen, 5-fluorouracil, methotrexate, temozolomide, cyclophosphamide, SCH 66336, R115777, L778,123, BMS 214662, Iressa, Tarceva, antibodies to EGFR, Gleevec™, intron, ara-C, adriamycin, cytoxan, gemcitabine, Uracil mustard, Chlormethine, Ifosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramine, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, oxaliplatin, leucovorin, ELOXATIN™, Pentostatine, Vinblastine, Vincristine, Vindesine, Bleomycin, Dactinomycin, Daunorubicin, Doxorubicin, Epirubicin, Idarubicin, Mithramycin, Deoxycoformycin, Mitomycin-C, L-Asparaginase, Teniposide 17.alpha.-Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrolacetate, Methylprednisolone, Methyltestosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, goserelin, Cisplatin, Carboplatin, Hydroxyurea, Amsacrine, Procarbazine, Mitotane, Mitoxantrone, Levamisole, Navelbene, Anastrozole, Letrozole, Capecitabine, Reloxafine, Droloxafine, Hexamethylmelamine, Avastin, herceptin, Bexxar, Velcade, Zevalin, Trisenox, Xeloda, Vinorelbine, Porfimer, Erbitux, Liposomal, Thiotepa, Altretamine, Melphalan, Trastuzumab, Lerozole, Fulvestrant, Exemestane, Fulvestrant, Ifosfomide, Rituximab, C225, Campath, Clofarabine, cladribine, aphidicolon, rituxan, sunitinib, dasatinib, tezacitabine, Sml1, fludarabine, pentostatin, triapine, didox, trimidox, amidox, 3-AP, and MDL-101,731.

[0190] Example chemotherapeutics include proteasome inhibitors (e.g., bortezomib), thalidomide, revlimid, and DNA-damaging agents such as melphalan, doxorubicin, cyclophosphamide, vincristine, etoposide, carmustine, and the like.

[0191] Example steroids include corticosteroids such as dexamethasone or prednisone.

[0192] Example Bcr-Abl inhibitors include the compounds, and pharmaceutically acceptable salts thereof, of the genera and species disclosed in U.S. Pat. No. 5,521,184, WO 04/005281, and U.S. Ser. No. 60/578,491.

[0193] Example suitable Flt-3 inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 03/037347, WO 03/099771, and WO 04/046120.

[0194] Example suitable RAF inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 00/09495 and WO 05/028444.

[0195] Example suitable FAK inhibitors include compounds, and their pharmaceutically acceptable salts, as disclosed in WO 04/080980, WO 04/056786, WO 03/024967, WO 01/064655, WO 00/053595, and WO 01/014402.

[0196] In some embodiments, the compounds of the invention can be used in combination with one or more other kinase inhibitors including imatinib, particularly for treating patients resistant to imatinib or other kinase inhibitors.

[0197] In some embodiments, the compounds of the invention can be used in combination with a chemotherapeutic in the treatment of cancer, such as multiple myeloma, and may improve the treatment response as compared to the response to the chemotherapeutic agent alone, without exacerbation of its toxic effects. Examples of additional pharmaceutical agents used in the treatment of multiple myeloma, for example, can include, without limitation, melphalan, melphalan plus prednisone [MP], doxorubicin, dexamethasone, and Velcade (bortezomib). Further additional agents used in the treatment of multiple myeloma include Bcr-Abl, Flt-3, RAF and FAK kinase inhibitors. Additive or synergistic effects are desirable outcomes of combining a PI3K inhibitor of the present invention with an additional agent. Furthermore, resistance of multiple myeloma cells to agents such as dexamethasone may be reversible upon treatment with the PI3K inhibitor of the present invention. The agents can be combined with the present compound in a single or continuous dosage form, or the agents can be administered simultaneously or sequentially as separate dosage forms.

[0198] In some embodiments, a corticosteroid such as dexamethasone is administered to a patient in combination with the compounds of the invention where the dexamethasone is administered intermittently as opposed to continuously.

[0199] In some further embodiments, combinations of the compounds of the invention with other therapeutic agents can be administered to a patient prior to, during, and/or after a bone marrow transplant or stem cell transplant.

[0200] When employed as pharmaceuticals, the compounds of the invention can be administered in the form of pharmaceutical compositions. These compositions can be prepared in a manner well known in the pharmaceutical art, and can be administered by a variety of routes, depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including transdermal, epidermal, ophthalmic and to mucous membranes including intranasal, vaginal and rectal delivery), pulmonary (e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal or intranasal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal intramuscular or injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Parenteral administration can be in the form of a single bolus dose, or may be, for example, by a continuous perfusion pump. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

[0201] This invention also includes pharmaceutical compositions which contain, as the active ingredient, the compound of the invention or a pharmaceutically acceptable salt thereof, in combination with one or more pharmaceutically acceptable carriers (excipients). In some embodiments, the composition is suitable for topical administration. In making the compositions of the invention, the active ingredient is typically mixed with an excipient, diluted by an excipient or enclosed within such a carrier in the form of, for example, a capsule, sachet, paper, or other container. When the excipient serves as a diluent, it can be a solid, semi-solid, or liquid material, which acts as a vehicle, carrier or medium for the active ingredient. Thus, the compositions can be in the form of tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), ointments containing, for example, up to 10% by weight of the active compound, soft and hard gelatin capsules, suppositories, sterile injectable solutions, and sterile packaged powders.

[0202] In preparing a formulation, the active compound can be milled to provide the appropriate particle size prior to combining with the other ingredients. If the active compound is substantially insoluble, it can be milled to a particle size of less than 200 mesh. If the active compound is substantially water soluble, the particle size can be adjusted by milling to provide a substantially uniform distribution in the formulation, e.g. about 40 mesh.

[0203] The compounds of the invention may be milled using known milling procedures such as wet milling to obtain a particle size appropriate for tablet formation and for other formulation types. Finely divided (nanoparticulate) preparations of the compounds of the invention can be prepared by processes known in the art, e.g., see International App. No. WO 2002/000196.

[0204] Some examples of suitable excipients include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, water, syrup, and methyl cellulose. The formulations can additionally include: lubricating agents such as talc, magnesium stearate, and mineral oil; wetting agents; emulsifying and suspending agents; preserving agents such as methyl- and propylhydroxy-benzoates; sweetening agents; and flavoring agents. The compositions of the invention can be formulated so as to provide quick, sustained or delayed release of the active ingredient after administration to the patient by employing procedures known in the art.

[0205] The compositions can be formulated in a unit dosage form, each dosage containing from about 5 to about 1000 mg (1 g), more usually about 100 to about 500 mg, of the active ingredient. The term "unit dosage forms" refers to physically discrete units suitable as unitary dosages for human subjects and other mammals, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical excipient.

[0206] In some embodiments, the compounds or compositions of the invention contain from about 5 to about 50 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 5 to about 10, about 10 to about 15, about 15 to about 20, about 20 to about 25, about 25 to about 30, about 30 to about 35, about 35 to about 40, about 40 to about 45, or about 45 to about 50 mg of the active ingredient.

[0207] In some embodiments, the compounds or compositions of the invention contain from about 50 to about 500 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 50 to about 100, about 100 to about 150, about 150 to about 200, about 200 to about 250, about 250 to about 300, about 350 to about 400, or about 450 to about 500 mg of the active ingredient.

[0208] In some embodiments, the compounds or compositions of the invention contain from about 500 to about 1000 mg of the active ingredient. One having ordinary skill in the art will appreciate that this embodies compounds or compositions containing about 500 to about 550, about 550 to about 600, about 600 to about 650, about 650 to about 700, about 700 to about 750, about 750 to about 800, about 800 to about 850, about 850 to about 900, about 900 to about 950, or about 950 to about 1000 mg of the active ingredient.

[0209] The active compound can be effective over a wide dosage range and is generally administered in a pharma-

ceutically effective amount. It will be understood, however, that the amount of the compound actually administered will usually be determined by a physician, according to the relevant circumstances, including the condition to be treated, the chosen route of administration, the actual compound administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the like.

[0210] For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical excipient to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention. When referring to these preformulation compositions as homogeneous, the active ingredient is typically dispersed evenly throughout the composition so that the composition can be readily subdivided into equally effective unit dosage forms such as tablets, pills and capsules. This solid preformulation is then subdivided into unit dosage forms of the type described above containing from, for example, about 0.1 to about 1000 mg of the active ingredient of the present invention.

[0211] The tablets or pills of the present invention can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which serves to resist disintegration in the stomach and permit the inner component to pass intact into the duodenum or to be delayed in release. A variety of materials can be used for such enteric layers or coatings, such materials including a number of polymeric acids and mixtures of polymeric acids with such materials as shellac, cetyl alcohol, and cellulose acetate.

[0212] The liquid forms in which the compounds and compositions of the present invention can be incorporated for administration orally or by injection include aqueous solutions, suitably flavored syrups, aqueous or oil suspensions, and flavored emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil, or peanut oil, as well as elixirs and similar pharmaceutical vehicles.

[0213] Compositions for inhalation or insufflation include solutions and suspensions in pharmaceutically acceptable, aqueous or organic solvents, or mixtures thereof, and powders. The liquid or solid compositions may contain suitable pharmaceutically acceptable excipients as described *supra*. In some embodiments, the compositions are administered by the oral or nasal respiratory route for local or systemic effect. Compositions can be nebulized by use of inert gases. Nebulized solutions may be breathed directly from the nebulizing device or the nebulizing device can be attached to a face mask, tent, or intermittent positive pressure breathing machine. Solution, suspension, or powder compositions can be administered orally or nasally from devices which deliver the formulation in an appropriate manner.

[0214] Topical formulations can contain one or more conventional carriers. In some embodiments, ointments can contain water and one or more hydrophobic carriers selected from, for example, liquid paraffin, polyoxyethylene alkyl ether, propylene glycol, white Vaseline, and the like. Carrier compositions of creams can be based on water in combination with glycerol and one or more other components, e.g. glycerinemonostearate, PEG-glycerinemonostearate and cetylstearyl alcohol. Gels can be formulated using isopropyl alcohol and water, suitably in combination with other components such as, for example, glycerol, hydroxyethyl cellulose, and the like. In some embodiments, topical formulations contain at least about 0.1, at least about 0.25, at least about 0.5, at least about 1, at least about 2, or at least about 5 wt % of the compound of the invention. The topical formulations can be suitably packaged in tubes of, for example, 100 g which are optionally associated with instructions for the treatment of the select indication, e.g., psoriasis or other skin condition.

[0215] The amount of compound or composition administered to a patient will vary depending upon what is being administered, the purpose of the administration, such as prophylaxis or therapy, the state of the patient, the manner of administration, and the like. In therapeutic applications, compositions can be administered to a patient already suffering from a disease in an amount sufficient to cure or at least partially arrest the symptoms of the disease and its complications. Effective doses will depend on the disease condition being treated as well as the judgment of the attending clinician depending upon factors such as the severity of the disease, the age, weight and general condition of the patient, and the like.

[0216] The compositions administered to a patient can be in the form of pharmaceutical compositions described above. These compositions can be sterilized by conventional sterilization techniques, or may be sterile filtered. Aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous carrier prior to administration. The pH of the compound preparations typically will be between 3 and 11, more preferably from 5 to 9 and most preferably from 7 to 8. It will be understood that use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of pharmaceutical salts.

[0217] The therapeutic dosage of a compound of the present invention can vary according to, for example, the particular use for which the treatment is made, the manner of administration of the compound, the health and condition of the patient, and the judgment of the prescribing physician. The proportion or concentration of a compound of the invention in a pharmaceutical composition can vary depending upon a number of factors including dosage, chemical characteristics (e.g., hydrophobicity), and the route of administration. For example, the compounds of the invention can be provided in an aqueous physiological buffer solution containing about 0.1 to about 10% w/v of the compound for parenteral administration. Some typical dose ranges are from about 1 µg/kg to about 1 g/kg of body weight per day. In some embodiments,

the dose range is from about 0.01 mg/kg to about 100 mg/kg of body weight per day. The dosage is likely to depend on such variables as the type and extent of progression of the disease or disorder, the overall health status of the particular patient, the relative biological efficacy of the compound selected, formulation of the excipient, and its route of administration. Effective doses can be extrapolated from dose-response curves derived from *in vitro* or animal model test systems.

[0218] The compositions of the invention can further include one or more additional pharmaceutical agents such as a chemotherapeutic, steroid, anti-inflammatory compound, or immunosuppressant, examples of which are listed herein.

Labeled Compounds and Assay Methods

[0219] Another aspect of the present invention relates to labeled compounds of the invention (radio-labeled, fluorescent-labeled, etc.) that would be useful not only in imaging techniques but also in assays, both *in vitro* and *in vivo*, for localizing and quantitating PI3K in tissue samples, including human, and for identifying PI3K ligands by inhibition binding of a labeled compound. Accordingly, the present invention includes PI3K assays that contain such labeled compounds.

[0220] The present invention further includes isotopically-labeled compounds of the invention. An "isotopically" or "radio-labeled" compound is a compound of the invention where one or more atoms are replaced or substituted by an atom having an atomic mass or mass number different from the atomic mass or mass number typically found in nature (*i.e.*, naturally occurring). Suitable radionuclides that may be incorporated in compounds of the present invention include but are not limited to ^2H (also written as D for deuterium), ^3H (also written as T for tritium), ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{18}F , ^{35}S , ^{36}Cl , ^{82}Br , ^{75}Br , ^{76}Br , ^{77}Br , ^{123}I , ^{124}I , ^{125}I and ^{131}I . The radionuclide that is incorporated in the instant radio-labeled compounds will depend on the specific application of that radio-labeled compound. For example, for *in vitro* PI3K labeling and competition assays, compounds that incorporate ^3H , ^{14}C , ^{82}Br , ^{125}I , ^{131}I , ^{35}S or will generally be most useful. For radio-imaging applications ^{11}C , ^{18}F , ^{125}I , ^{123}I , ^{124}I , ^{131}I , ^{75}Br , ^{76}Br or ^{77}Br will generally be most useful.

[0221] It is understood that a "radio-labeled" or "labeled compound" is a compound that has incorporated at least one radionuclide. In some embodiments the radionuclide is selected from the group consisting of ^3H , ^{14}C , ^{125}I , ^{35}S and ^{82}Br .

[0222] The present invention can further include synthetic methods for incorporating radio-isotopes into compounds of the invention. Synthetic methods for incorporating radio-isotopes into organic compounds are well known in the art, and an ordinary skill in the art will readily recognize the methods applicable for the compounds of invention.

[0223] A labeled compound of the invention can be used in a screening assay to identify/evaluate compounds. For example, a newly synthesized or identified compound (*i.e.*, test compound) which is labeled can be evaluated for its ability to bind a PI3K by monitoring its concentration variation when contacting with the PI3K, through tracking of the labeling. For example, a test compound (labeled) can be evaluated for its ability to reduce binding of another compound which is known to bind to a PI3K (*i.e.*, standard compound). Accordingly, the ability of a test compound to compete with the standard compound for binding to the PI3K directly correlates to its binding affinity. Conversely, in some other screening assays, the standard compound is labeled and test compounds are unlabeled. Accordingly, the concentration of the labeled standard compound is monitored in order to evaluate the competition between the standard compound and the test compound, and the relative binding affinity of the test compound is thus ascertained.


Kits

[0224] The present invention also includes pharmaceutical kits useful, for example, in the treatment or prevention of PI3K-associated diseases or disorders, such as cancer, which include one or more containers containing a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention. Such kits can further include, if desired, one or more of various conventional pharmaceutical kit components, such as, for example, containers with one or more pharmaceutically acceptable carriers, additional containers, etc., as will be readily apparent to those skilled in the art. Instructions, either as inserts or as labels, indicating quantities of the components to be administered, guidelines for administration, and/or guidelines for mixing the components, can also be included in the kit.

[0225] The invention will be described in greater detail by way of specific examples. The following examples are offered for illustrative purposes, and are not intended to limit the invention in any manner. Those of skill in the art will readily recognize a variety of non-critical parameters which can be changed or modified to yield essentially the same results. The compounds of the Examples have been found to be PI3K inhibitors according to at least one assay described herein.

EXAMPLES

[0226] The example compounds below containing one or more chiral centers were obtained in racemate form or as isomeric mixtures, unless otherwise specified.

Example 1. 2-[1-(6-amino-9H-purin-9-yl)ethyl]-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one**[0227]***Step 1. methyl 4-chloro-3-oxopentanoate*

[0228] To a solution of 3-oxopentanoic acid, methyl ester (Aldrich, 26.0 mL, 207.2 mmol) in methylene chloride (300 mL) was added in portions, N,N,N-trimethyl(phenyl) methanaminium dichloriodanuide (75.71 g, 217.5 mmol). The reaction mixture was stirred at room temperature (rt) for 2 h, then washed with saturated sodium thiosulfate, brine, dried over magnesium sulfate and concentrated. The crude product was used directly in next step (23 g, 67.4%).

Step 2. 2-(1-chloroethyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0229] To a manually stirred polyphosphoric acid (30 g, 200 mmol) in a 200 mL beaker was added 6-methyl-2-pyridinamine (Aldrich, 4.7 g, 43 mmol), followed by methyl 4-chloro-3-oxopentanoate (8.584 g, 52.15 mmol). The mixture was heated with stirring at 110 °C for 5 h. After being cooled, the dark slurry was transferred on to 100 g of ice. The pH of the mixture was adjusted to 6-7 with 10% sodium hydroxide. The mixture was extracted with methylene chloride. The combined organic layers were washed with water, brine, dried over magnesium sulfate and evaporated to dryness. The residue was purified on silica gel, eluting with 0-10% methanol in methylene chloride, to yield the desired product (3.16 g, 32.7%). LCMS calculated for $C_{11}H_{12}ClN_2O(M+H)^+$: $m/z = 223.1$; Found: 223.2.

Step 3. 3-bromo-2-(1-chloroethyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0230] To a stirred solution of 2-(1-chloroethyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (3.16 g, 14.2 mmol) in methylene chloride (30 mL) was added drop-wise bromine (0.804 mL, 15.6 mmol) in methylene chloride (7 mL). The reaction mixture was stirred at room temperature for 2 h. The product precipitated out and was collected by filtration (2.42 g, 56.6%). LCMS calculated for $C_{11}H_{11}BrClN_2O(M+H)^+$: $m/z = 301.0$; Found: 301.1.

Step 4. 2-[1-(6-amino-9H-purin-9-yl)ethyl]-3-bromo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0231] A mixture of 3-bromo-2-(1-chloroethyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (1.24 g, 4.11 mmol), adenine (Sigma, 1.08 g, 8.04 mmol), and potassium carbonate (1.11 g, 8.04 mmol) in *N,N*-dimethylformamide (20 mL) was stirred at room temperature overnight. The suspension was then poured into water and extracted with methylene chloride. The combined organic layers were washed with water, brine, and then the organic layers dried and evaporated to dryness. The residue was chromatographed on silica gel, eluting with 0 to 10% MeOH in methylene chloride, to provide the desired product (176 mg, 10.7%). LCMS calculated for $C_{16}H_{15}BrN_7O(M+H)^+$: $m/z = 400.1$; Found: 400.1. 1H NMR (DMSO- d_6 , 300 MHz) δ 8.44 (1H, s), 8.07 (1H, s), 7.67 (1H, dd, $J = 8.7$ and 7.2 Hz), 7.30 (1H, br d, $J = 8.7$ Hz), 7.20 (2H, s), 6.99 (1H, br d, $J = 7.2$ Hz), 6.15 (1H, q, $J = 7.2$ Hz), 2.91 (3H, s), 1.84 (3H, d, $J = 7.2$ Hz) ppm.


Step 5. 2-[1-(6-amino-9H-purin-9-yl)ethyl]-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0232] To a mixture of 2-[1-(6-amino-9H-purin-9-yl)ethyl]-3-bromo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (0.030 g, 0.075 mmol) and phenylboronic acid (11.0 mg, 0.0899 mmol) in 1,4-dioxane (0.6 mL) was added a 1 M solution of sodium carbonate (9.53 mg, 0.0899 mmol) in water (0.089 mL) and tetrakis(triphenylphosphine)palladium (0) (4.33 mg, 0.00375 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over $MgSO_4$, and concentrated. The residue was purified on RP-HPLC at pH 10 (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.15% NH_4OH) to provide the desired product. LCMS calculated for $C_{22}H_{20}N_7O(M+H)^+$: $m/z = 398.2$; Found: 398.3. 1H NMR (DMSO- d_6 , 300 MHz) δ 8.37 (1H, s), 7.97 (1H, s), 7.56 (1H, dd, $J = 8.7$ and 6.6 Hz), 7.41-7.36 (5H, m), 7.26 (1H, br d, $J = 9.0$ Hz), 7.09 (2H, br s), 6.85

(1H, br d, $J = 7.2$ Hz), 5.59 (1H, q, $J = 7.2$ Hz), 2.80 (3H, s), 1.64 (3H, d, $J = 7.2$ Hz) ppm.

Example 2. 6-methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one

[0233]



Step 1. methyl 4-bromo-3-oxohexanoate

[0234] Under a nitrogen atmosphere, a solution of bromine (8.61 mL, 167 mmol) in chloroform (20 mL) was added drop-wise over a period of 2 h to a solution of methyl 3-oxohexanoate (Fluka, 24.1 g, 167 mmol) in chloroform (147 mL), at 0 °C (ice bath). The reaction mixture was stirred for 30 min at 0 °C and then allowed to warm to room temperature overnight. While stirring, a stream of air was bubbled through the solution for 1 hour. The reaction mixture was dried over sodium sulfate and the solvent evaporated under reduced pressure to provide the desired compound.

Step 2. 2-(1-bromopropyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0235] To a manually stirred polyphosphoric acid (80 g, 800 mmol) in a 1000 mL beaker at room temperature was added 6-methyl-2-pyridinamine (15 g, 140 mmol), followed by methyl 4-bromo-3-oxohexanoate (37.3 g, 167 mmol). The mixture was heated with stirring at 110 °C for 5 h. After cooling, the dark slurry was transferred into 300 g of ice. The pH of the mixture was adjust to 6-7 with 10% sodium hydroxide. The precipitate was collected by filtration under reduced pressure, washed with water, and air dried to yield the desired product (25.4 g, 64.8%). LCMS calculated for $C_{12}H_{14}BrN_2O(M+H)^+$: $m/z = 281.0$; Found: 281.2. 1H NMR (DMSO- d_6 , 300 MHz) δ 7.66 (1H, d, $J = 9.0$ and 6.9 Hz), 7.39 (1H, d, $J = 9.0$ Hz), 6.90 (1H, d, $J = 6.9$ Hz), 6.33 (1H, s), 4.91 (1H, t, $J = 7.5$ Hz), 2.91 (3H, s), 2.15 (2H, qd, $J = 7.5$ and 7.5 Hz), 0.93 (3H, t, $J = 7.5$ Hz) ppm.

Step 3. 2-(1-bromopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0236] A mixture of 2-(1-bromopropyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (3.46 g, 12.3 mmol) and *N*-iodosuccinimide (4.15 g, 18.4 mmol) in acetonitrile (100 mL) was stirred at 80 °C, under nitrogen, overnight. After removal of acetonitrile in vacuum, the resulting solid was dissolved in methylene chloride, washed with water, saturated $Na_2S_2O_3$, saturated sodium bicarbonate, and brine; and then the organic layers dried over sodium sulfate and then filtered. The filtrate was concentrated under reduced pressure to provide the desired product (4.53 g, 90.4%). LCMS calculated for $C_{12}H_{13}BrIN_2O(M+H)^+$: $m/z = 406.9$; Found: 407.1.

Step 4. 2-(1-azidopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0237] A mixture of 2-(1-bromopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (4.50 g, 11.0 mmol) and sodium azide (3.59 g, 55.3 mmol) in DMF was stirred at room temperature for 2 h. After diluting with ethyl acetate, the mixture was washed with water, brine, dried over sodium sulfate, and evaporated under reduced pressure to provide the crude product, which was used directly in next step (3.35 g, 82.1 %). LCMS calculated for $C_{12}H_{13}IN_5O(M+H)^+$: $m/z = 370.0$; Found: 370.2.

Step 5. 2-(1-azidopropyl)-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0238] To a mixture of 2-(1-azidopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (0.11 g, 0.29 mmol) and phenylboronic acid (42.9 mg, 0.352 mmol) in 1,4-dioxane (2 mL) was added a 1 M solution of sodium carbonate (37.3 mg, 0.352 mmol) in water (0.35 mL) and tetrakis(triphenylphosphine)palladium (0) (16.9 mg, 0.0147 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over $MgSO_4$, and concentrated. The residue was purified on silica gel, eluting with 0 to 40% EtOAc in hexane,

to provide the desired product (50 mg, 53.4%). LCMS calculated for $C_{18}H_{18}N_5O(M+H)^+$: $m/z = 320.2$; Found: 320.3.

Step 6. 2-(1-aminopropyl)-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one


[0239] To a stirred solution of 2-(1-azidopropyl)-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one (0.030 g, 0.094 mmol) in tetrahydrofuran (0.24 mL) and water (0.06 mL) was added 1.0 M of trimethylphosphine in tetrahydrofuran (0.11 mL) at room temperature and the mixture was stirred at room temperature for 1 hour. To the mixture was added methylene chloride and the mixture was washed with brine, dried over magnesium sulfate, and evaporated to dryness under reduced pressure. The crude residue was used directly in next step. LCMS calculated for $C_{18}H_{20}N_3O(M+H)^+$: $m/z = 294.2$; Found: 294.3.

Step 7. 6-methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one

[0240] A mixture of 6-bromo-9H-purine (Aldrich, 0.0152 g, 0.07656 mmol), 2-(1-aminopropyl)-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one (0.019 g, 0.064 mmol), and *N,N*-diisopropylethylamine (0.0134 mL, 0.07666 mmol) in ethanol (0.5 mL) was refluxed under nitrogen overnight. The mixture was cooled and purified on RP-HPLC at pH 10 (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.15% NH_4OH) to provide the product as the free base. LCMS calculated for $C_{23}H_{22}N_7O(M+H)^+$: $m/z = 412.2$; Found: 412.4. 1H NMR ($DMSO-d_6$, 300 MHz) δ 8.07 (2H, m), 7.60 (1H, dd, $J = 9.0$ and 6.9 Hz), 7.39-7.32 (7H, m), 7.00 (1H, m), 6.85 (1H, br d, $J = 6.9$ Hz), 5.13 (1H, m), 2.81 (3H, s), 1.72 (2H, m), 0.65 (3H, t, $J = 7.2$ Hz) ppm.

Example 3. 3-(5-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one

[0241]



Step 1. 2-(1-aminopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0242] To a stirred solution of 2-(1-azidopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (3.10 g, 8.40 mmol) in tetrahydrofuran (20 mL) and water (6.06 mL) was added a 1.0 M solution of trimethylphosphine in tetrahydrofuran (0.1 mmol) at room temperature and stirred for 1 hour. To the mixture was added EtOAc and the mixture was extracted twice with 1 N HCl. The combined extracts were neutralized with solid sodium bicarbonate and extracted with methylene chloride. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step (2.58 g, 89.5%). LCMS calculated for $C_{12}H_{15}IN_3O(M+H)^+$: $m/z = 344.0$; Found: 344.2.

Step 2. 3-iodo-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one

[0243] A mixture of 6-bromo-9H-purine (1.65 g, 0.008270 mol), 2-(1-aminopropyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (2.58 g, 0.00752 mol), and *N,N*-diisopropylethylamine (1.571 mL, 0.009022 mol) in ethanol (60 mL) was refluxed under nitrogen overnight. The mixture was concentrated and the resulting residue was purified on silica gel, eluting with 0 to 10% methanol in methylene chloride, to provide the desired product (2.86 g, 82.5%). LCMS calculated for $C_{17}H_{17}IN_7O(M+H)^+$: $m/z = 462.1$; Found: 462.2. 1H NMR ($DMSO-d_6$, 300 MHz) δ 8.31 (2H, m), 8.19 (1H, s), 8.15 (1H, s), 7.69 (1H, dd, $J = 8.7$ and 6.9 Hz), 7.44 (1H, d, $J = 8.7$ Hz), 6.99 (1H, d, $J = 6.9$ Hz), 5.69 (1H, m), 2.89 (3H, s), 1.91 (2H, m), 0.95 (3H, t, $J = 7.2$ Hz) ppm.


Step 3. 3-(5-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one

[0244] To a mixture of 3-iodo-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one (0.030 g,

0.065 mmol) and (5-fluoropyridin-3-yl)boronic acid (Combi-Blocks, 11.0 mg, 0.0780 mmol) in 1,4-dioxane (0.5 mL) was added a 1 M solution of sodium carbonate (8.27 mg, 0.0780 mmol) in water (0.077 mL) and tetrakis(triphenylphosphine)palladium (0) (3.76 mg, 0.00325 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over MgSO₄, and concentrated. The residue was purified on RP-HPLC at pH 10 conditions (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to provide the desired product. LCMS calculated for C₂₂H₂₀FN₈O(M+H)⁺: m/z = 431.2; Found: 431.3.

Example 4. 3-(3-fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one


[0245]



[0246] To a mixture of 3-iodo-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one (from example 3, step 2; 0.030 g, 0.065 mmol) and (3-fluorophenyl)boronic acid (Aldrich, 10.9 mg, 0.0780 mmol) in 1,4-dioxane (0.5 mL) was added a 1 M solution of sodium carbonate (8.27 mg, 0.0780 mmol) in water (0.077 mL) and tetrakis(triphenylphosphine)palladium (0) (3.76 mg, 0.00325 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over MgSO₄, and concentrated. The residue was purified on RP-HPLC at pH 10 conditions (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to provide the desired product. LCMS calculated for C₂₃H₂₁FN₇O(M+H)⁺: m/z = 430.2; Found: 430.3. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 8.11 (2H, m), 7.65 (1H, m), 7.45 (2H, m), 7.24 (4H, m), 7.08 (1H, m), 6.91 (1H, m), 5.17 (1H, m), 2.87 (3H, s), 1.79 (2H, m), 0.72 ((3H, t, *J* = 7.2 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 282 MHz) δ -114 ppm.

Example 5. 3-(3,5-difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one


[0247]



[0248] To a mixture of 3-iodo-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one (from example 3, step 2; 0.030 g, 0.065 mmol) and (3,5-difluorophenyl)boronic acid (Aldrich, 12.3 mg, 0.0780 mmol) in 1,4-dioxane (0.5 mL) was added a 1 M solution of sodium carbonate (8.27 mg, 0.0780 mmol) in water (0.077 mL) and tetrakis(triphenylphosphine)palladium (0) (3.76 mg, 0.00325 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over MgSO₄, and concentrated. The residue was purified on RP-HPLC at pH 10 conditions (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to provide the desired product. LCMS calculated for C₂₃H₂₀F₂N₇O(M+H)⁺: m/z = 448.2; Found: 448.3. ¹H NMR (DMSO-*d*₆, 300 MHz) δ 8.06 (2H, m), 7.61 (1H, m), 7.38 (1H, m), 7.14 (5H, m), 6.88 (1H, m), 5.08 (1H, m), 2.83 (3H, s), 1.75 (2H, m), 0.70 ((3H, t, *J* = 7.8 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 282 MHz) δ -111 ppm.

Example 6. 3-(2-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one


[0249]



[0250] To a mixture of 3-iodo-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one (from example 3, step 2; 0.030 g, 0.065 mmol) and (2-fluoropyridin-3-yl)boronic acid (Alfa Aesar, 11.0 mg, 0.0780 mmol) in 1,4-dioxane (0.5 mL) was added a 1 M solution of sodium carbonate (8.27 mg, 0.0780 mmol) in water (0.077 mL) and tetrakis(triphenylphosphine)palladium (0) (3.76 mg, 0.00325 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over MgSO₄, and concentrated. The residue was purified on RP-HPLC at pH 2 conditions (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to provide the desired product as a TFA salt. LCMS calculated for free base C₂₂H₂₀FN₈O(M+H)⁺: m/z = 431.2; Found: 431.3.

Example 7. 6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-3-(1H-pyrazol-4-yl)-4H-pyrido[1,2-a]pyrimidin-4-one


[0251]



[0252] To a mixture of 3-iodo-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one (from example 3, step 2; 0.030 g, 0.065 mmol) and 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (Aldrich, 15.1 mg, 0.0780 mmol) in 1,4-dioxane (0.5 mL) was added a 1 M solution of sodium carbonate (8.27 mg, 0.0780 mmol) in water (0.077 mL) and tetrakis(triphenylphosphine)palladium (0) (3.76 mg, 0.00325 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over MgSO₄, and concentrated. The residue was purified on RP-HPLC at pH 2 conditions (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to provide the desired product as a TFA salt. LCMS calculated for free base C₂₀H₂₀N₉O(M+H)⁺: m/z = 402.2; Found: 402.1.

Example 8. 3-methyl-6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0253]



Step 1. methyl 4-bromo-3-oxopentanoate

[0254] Under a nitrogen atmosphere, a solution of bromine (8.61 mL, 167 mmol) in chloroform (20 mL, 200 mmol) was added dropwise over a period of 2 h to a solution of 3-oxopentanoic acid, methyl ester (Aldrich, 21.0 mL, 167 mmol) in chloroform (147 mL, 1840 mmol), at 0 °C (ice bath). The reaction mixture was stirred for 30 min at 0 °C and then

allowed to stand at room temperature overnight. While stirring, a stream of air was bubbled through the solution for 1 hour. After drying over sodium sulfate, the solvent was evaporated under reduced pressure leaving the desired compound.

Step 2. 7-(1-bromoethyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0255] To a manually stirred polyphosphoric acid (80 g, 800 mmol) in a 1000 mL beaker was added 4-methyl-1,3-thiazol-2-amine (Aldrich, 16 g, 140 mmol), followed by methyl 4-bromo-3-oxopentanoate (34.9 g, 167 mmol). The mixture was heated with stirring at 110 °C for 5 h. After cooling, the dark slurry was transferred into 300 g of ice. The pH of the mixture was adjust to 6-7 with 10% sodium hydroxide. The aqueous layer was discarded and the dark oil layer was diluted with methylene chloride and washed with 1 N NaOH, brine, dried over magnesium sulfate, and evaporated to dryness to yield the desired product (16.2 g, 42.6%). LCMS calculated for $C_9H_{10}BrN_2OS(M+H)^+$: $m/z = 273.0$; Found: 273.1.

Step 2. 6-bromo-7-(1-bromoethyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0256] A mixture of 7-(1-bromoethyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (16.2 g, 59.3 mmol) and *N*-bromosuccinimide (15.8 g, 89.0 mmol) in acetonitrile (500 mL) was stirred at 80 °C, under nitrogen, overnight. After removal of acetonitrile in vacuum, the resulting solid was dissolved in methylene chloride, washed with water, saturated $Na_2S_2O_3$, saturated sodium bicarbonate, and brine; and then the organic layers dried over sodium sulfate and filtered. The filtrate was concentrated under reduced pressure to provide the desired product (19.5 g, 93.4%). LCMS calculated for $C_9H_9Br_2N_2OS(M+H)^+$: $m/z = 350.9$; Found: 351.0.

Step 3. 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0257] A mixture of 6-bromo-7-(1-bromoethyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (11.1 g, 31.5 mmol) and sodium azide (6.15 g, 94.6 mmol) in dimethyl formamide (DMF) (100 mL) was stirred at room temperature for 2 h. After diluting with EtOAc, the mixture was washed with water, brine, dried over sodium sulfate, and evaporated under reduced pressure. The crude residue was purified on silica gel, eluting with 0 to 80% EtOAc in hexane, to provide the product (8.68 g, 87.6%). LCMS calculated for $C_9H_9BrN_5OS(M+H)^+$: $m/z = 314.0$; Found: 313.9. 1H NMR (DMSO- d_6 , 300 MHz) δ 7.15 (1H, s), 4.83 (1H, q, $J = 6.6$ Hz), 2.69 (3H, s), 1.48 (3H, d, $J = 6.6$ Hz) ppm.

Step 4. 7-(1-azidoethyl)-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0258] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.100 g, 0.318 mmol) and phenylboronic acid (46.6 mg, 0.382 mmol) in 1,4-dioxane (2 mL) was added a 1 M solution of sodium carbonate (40.5 mg, 0.382 mmol) in water (0.38 mL) and tetrakis(triphenylphosphine)palladium (0) (18.4 mg, 0.0159 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over $MgSO_4$, and concentrated. The residue was purified on silica gel, eluting with 0 to 50% EtOAc in hexane, to provide the desired product (44 mg, 44.4%). LCMS calculated for $C_{15}H_{14}N_5OS(M+H)^+$: $m/z = 312.1$; Found: 312.3.

Step 5. 7-(1-aminoethyl)-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0259] To a stirred solution of 7-(1-azidoethyl)-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.044 g, 0.14 mmol) in tetrahydrofuran (0.4 mL) and water (0.102 mL) was added 1.0 M of trimethylphosphine in tetrahydrofuran (0.17 mL) at room temperature and the mixture was stirred at room temperature for 1 hour. To the mixture was added EtOAc and the mixture was extracted twice with 1 N HCl. The combined extracts were neutralized with solid sodium bicarbonate and extracted with methylene chloride. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step (36 mg, 89.3%). LCMS calculated for $C_{15}H_{16}N_3OS(M+H)^+$: $m/z = 286.1$; Found: 286.0.


Step 6. 3-methyl-6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0260] A mixture of 6-bromo-9H-purine (0.01504 g, 0.0076 mmol), 7-(1-aminoethyl)-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.018 g, 0.063 mmol), and *N,N*-diisopropylethylamine (0.013 mL, 0.0076 mol) in ethanol (0.5 mL) was refluxed under nitrogen overnight. The mixture was concentrated under reduced pressure and the residue was purified on RP-HPLC at pH 2 to provide the product as a TFA salt. LCMS calculated for free base $C_{20}H_{18}N_7OS(M+H)^+$: $m/z = 404.1$; Found: 404.3. 1H NMR (DMSO- d_6 , 300 MHz) for a TFA salt: δ 8.54 (2H, m), 8.41 (2H, m), 7.44-7.36 (5H,

m), 7.08 (1H, d, J = 1.2 Hz), 5.21 (1H, m), 2.64 (3H, s), 1.38 (3H, d, J = 6.6 Hz) ppm.

Example 9. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0261]



[0262] A mixture of 2-amino-6-bromopurine (Aldrich, 0.01618 g, 0.007558 mmol), 7-(1-aminoethyl)-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from example 8, step 5; 0.018 g, 0.063 mmol), and *N,N*-diisopropylethylamine (0.01318 mL, 0.007569 mmol) in ethanol (0.5 mL) was refluxed under nitrogen overnight. The mixture was evaporated and the resulting residue was purified on RP-HPLC at pH 2 to provide the product as a TFA salt. LCMS calculated for $C_{20}H_{19}N_8OS(M+H)^+$: m/z = 419.1; Found: 419.3. 1H NMR (DMSO- d_6 , 300 MHz) for TFA salt: δ 8.74 (1H, m), 8.16 (1H, s), 7.46-7.33 (6H, m), 7.14-7.11 (3H, m), 5.20 (1H, m), 2.66 (3H, d, J = 1.5 Hz), 1.32 (3H, d, J = 6.6 Hz) ppm.

Example 10. 6-(3-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0263]



Step 1. 7-(1-azidoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0264] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from example 8, step 3; 0.100 g, 0.318 mmol) and (3-fluorophenyl)boronic acid (53.4 mg, 0.382 mmol) in 1,4-dioxane (2 mL) was added a 1 M solution of sodium carbonate (40.5 mg, 0.382 mmol) in water (0.38 mL) and tetrakis(triphenylphosphine)palladium (0) (18.4 mg, 0.0159 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over $MgSO_4$, and concentrated. The residue was purified on silica gel, eluting with 0 to 50% EtOAc in hexane, to provide the desired product (35 mg, 33.4%). LCMS calculated for $C_{15}H_{13}FN_5OS(M+H)^+$: m/z = 330.1; Found: 330.2.

Step 2. 7-(1-aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0265] To a stirred solution of 7-(1-azidoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.037 g, 0.11 mmol) in tetrahydrofuran (0.3 mL) and water (0.0811 mL) was added 1.0 M of trimethylphosphine in tetrahydrofuran (0.13 mmol) at room temperature and stirred for 1 hour. To the mixture was added EtOAc and was extracted twice with 1 N HCl. The combined extracts were neutralized with solid sodium bicarbonate and extracted with methylene chloride. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step (31 mg, 90.9%). LCMS calculated for $C_{15}H_{15}FN_3OS(M+H)^+$: m/z = 304.1; Found: 304.3.

Step 3. 6-(3-fluorophenyl)-3-methyl-7-(1-(9H-purin-6-ylamino)ethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0266] A mixture of 6-bromo-9H-purine (0.01258 g, 0.006320 mmol), 7-(1-aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.016 g, 0.053 mmol), and *N,N*-diisopropylethylamine (0.011 mL, 0.006329 mmol) in ethanol (0.4 mL) was refluxed under nitrogen overnight. The mixture was evaporated and the resulting residue was purified on RP-HPLC at pH 2 to provide the product as a TFA salt. LCMS calculated for $C_{20}H_{17}FN_7OS(M+H)^+$: m/z = 422.1; Found: 422.3. 1H NMR (DMSO- d_6 , 300 MHz) for TFA salt: δ 8.55 (2H, m), 8.40 (2H, m), 7.48 (1H, m), 7.25-7.22 (3H, m), 7.09 (1H, s), 5.19 (1H, m), 2.64 (3H, d, J = 0.9 Hz), 1.40 (3H, d, J = 6.6 Hz) ppm. ^{19}F NMR (DMSO- d_6 , 282 MHz) for TFA salt: δ -74.2, -114.0 ppm.

Example 11. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0267]



[0268] A mixture of 2-amino-6-bromopurine (0.01353 g, 0.006320 mmol), 7-(1-aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from example 10, step 2; 0.016 g, 0.053 mmol), and *N,N*-diisopropylethylamine (0.01102 mL, 0.006329 mmol) in ethanol (0.4 mL) was refluxed under nitrogen overnight. The mixture was evaporated and the resultant residue was purified on RP-HPLC at pH 2 to provide the product as a TFA salt. LCMS calculated for $C_{20}H_{18}FN_8OS(M+H)^+$: m/z = 437.1; Found: 437.3. 1H NMR (DMSO- d_6 , 300 MHz) for TFA salt: δ 8.74 (1H, m), 8.16 (1H, s), 7.46 (1H, m), 7.20-7.13 (7H, m), 5.18 (1H, m), 2.66 (3H, s), 1.33 (3H, d, J = 6.9 Hz) ppm. ^{19}F NMR (DMSO- d_6 , 282 MHz) for TFA salt: δ 74.0, -114.0 ppm.

Example 12. 6-(3,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0269]



Step 1. 7-(1-azidoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0270] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.10 g, 0.318 mmol) and (3,5-difluorophenyl)boronic acid (60.3 mg, 0.382 mmol) in 1,4-dioxane (2 mL) was added a 1 M solution of sodium carbonate (40.5 mg, 0.382 mmol) in water (0.38 mL) and tetrakis(triphenylphosphine)palladium (0) (18.4 mg, 0.0159 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to rt, the mixture was diluted with EtOAc, washed with water, brine, dried over $MgSO_4$, and concentrated. The residue was purified on silica gel, eluting with 0 to 40% EtOAc in hexane, to provide the desired product (42 mg, 38.0%). LCMS calculated for $C_{15}H_{12}F_2N_5OS(M+H)^+$: m/z = 348.1; Found: 348.2.

Step 2. 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0271] To a stirred solution of 7-(1-azidoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.042 g, 0.12 mmol) in tetrahydrofuran (0.3 mL) and water (0.0873 mL) was added 1.0 M of trimethylphosphine in tetrahydrofuran (0.14 mmol) at room temperature and stirred for 1 hour. To the mixture was added EtOAc and the mixture was extracted twice with 1 N HCl. The combined extracts were neutralized with solid sodium bicarbonate, and extracted with methylene chloride. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step (36 mg, 92.7%). LCMS calculated for $C_{15}H_{14}F_2N_3OS(M+H)^+$: $m/z = 322.1$; Found: 322.3.

Step 3. 6-(3,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0272] A mixture of 6-bromo-9H-purine (0.01258 g, 0.006320 mmol), 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.017 g, 0.053 mmol), and *N,N*-diisopropylethylamine (0.011 mL, 0.006329 mmol) in ethanol (0.4 mL) was refluxed under nitrogen overnight. The mixture was concentrated under reduced pressure and the resultant residue was purified on RP-HPLC at pH 2 to provide the product as a TFA salt. LCMS calculated for $C_{20}H_{16}F_2N_7OS(M+H)^+$: $m/z = 440.1$; Found: 440.3. 1H NMR (DMSO- d_6 , 300 MHz) for TFA salt: δ 8.55 (1H, m), 8.39 (2H, m), 7.29 (1H, m), 7.15-7.11 (3H, m), 5.17 (1H, m), 2.64 (3H, d, $J = 1.2$ Hz), 1.42 (3H, d, $J = 6.9$ Hz) ppm.

Example 13. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one**[0273]**

[0274] A mixture of 2-amino-6-bromopurine (0.01436 g, 0.006712 mmol), 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from example 12, step 2; 0.018 g, 0.056 mmol), and *N,N*-diisopropylethylamine (0.01171 mL, 0.006722 mmol) in ethanol (0.4 mL) was refluxed under nitrogen overnight. The mixture was evaporated and the resultant residue was purified on RP-HPLC at pH 2 to provide the product as a TFA salt. LCMS calculated for $C_{20}H_{17}F_2N_8OS(M+H)^+$: $m/z = 455.1$; Found: 455.3. 1H NMR (DMSO- d_6 , 300 MHz) for TFA salt: δ 8.74 (1H, m), 8.16 (1H, s), 7.26-7.06 (6H, m), 5.16 (1H, m), 2.66 (3H, d, $J = 1.2$ Hz), 1.36 (3H, d, $J = 6.9$ Hz) ppm.

Example 14. 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one**[0275]**

Step 1. 7-(1-azidoethyl)-3-methyl-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0276] A mixture of 2-(tributylstannyl)pyridine (Aldrich, 0.176 g, 0.382 mmol), 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (from example 8, step 3; 0.10 g, 0.318 mmol), and tetrakis(triphenylphosphine)pal-

ladium (0) (0.0184 g, 0.0159 mmol) in 1,4-dioxane (0.5 mL) was heated at 65 °C overnight. After being cooled and quenched with saturated ammonium chloride, the resulting mixture was extracted with EtOAc. The organic layers were combined, washed with brine, dried and evaporated to dryness. The residue was purified on silica gel, eluting with 0 to 100% EtOAc in hexane, to provide the desired product (13 mg, 13%). LCMS calculated for $C_{14}H_{13}N_6OS(M+H)^+$: m/z = 313.1; Found: 313.0.

Step 2. 7-(1-aminoethyl)-3-methyl-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0277] To a stirred solution of 7-(1-azidoethyl)-3-methyl-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.012 g, 0.039 mmol) in tetrahydrofuran (0.1 mL) and water (0.0285 mL) was added 1.0 M of trimethylphosphine in tetrahydrofuran (0.047 mmol) at room temperature and stirred for 1 hour. To the mixture was added EtOAc and the mixture was extracted twice with 1 N HCl. The combined extracts were neutralized with solid sodium bicarbonate and extracted with methylene chloride. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step. LCMS calculated for $C_{14}H_{15}N_4OS(M+H)^+$: m/z = 287.1; Found: 287.0.

Step 3. 3-methyl-7-(1-(9H-purin-6-ylamino)ethyl)-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0278] A mixture of 6-bromo-9H-purine (9.300 mg, 0.004673 mmol), 7-(1-aminoethyl)-3-methyl-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (11 mg, 0.039 mmol), and *N,N*-diisopropylethylamine (8.152 μ L, 0.004680 mmol) in ethanol (0.3 mL) was refluxed under nitrogen overnight. The mixture was concentrated under reduced pressure and the residue was purified on RP-HPLC (eluting with a gradient of methanol/water containing 1% TFA) to provide the product as a TFA salt. LCMS calculated for $C_{19}H_{17}N_8OS(M+H)^+$: m/z = 405.1; Found: 405.3.


Example 15. (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

[0279]



Step 1. 7-(1-Bromoethyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

[0280]



[0281] A solution of 3-oxopentanoic acid methyl ester (12.5 g, 96.0 mmol) in methylene chloride (50 mL) was cooled with an ice water bath. The outlet of the flask was attached to a NaOH trap. Bromine (5.19 mL, 101 mmol) in methylene chloride (10 mL, 200 mmol) was added dropwise over a 20-min period, and the reaction mixture was allowed to warm to room temperature and then stirred overnight. The reaction mixture was bubbled with nitrogen for 30 min and then concentrated to give an oil. This oil was used in the next step without further purification. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.89 (q, *J* = 6.9 Hz, 1H), 3.85 (s, 2H), 3.63 (s, 3H), 1.64 (d, *J* = 6.7 Hz, 3H).


[0282] Into a 3-neck flask fitted with a condenser, a thermometer, and a nitrogen inlet was added polyphosphoric acid (50.0 g, 458 mmol). The flask was heated to ~ 70 °C to give a liquid that was easy to stir. 4-Methyl-1,3-thiazol-2-amine (10.0 g, 87.6 mmol) was added in small portions with stirring. The internal temperature slowly increased to 78 °C upon mixing. The crude oil obtained above was then added to the flask *via* a pipette and the mixture was heated to 110 °C under nitrogen. After 6 h of heating, HPLC indicated that the reaction was complete.

[0283] The reaction mixture was cooled to ~ 35 °C. Water (70 mL) and EtOAc (200 mL) were added. The mixture was stirred until all solids dissolved. The organic layer was separated. The aqueous layer was extracted with EtOAc (200 mL x 2). The combined organic extracts were washed with 1 N aqueous HCl (40 mL x 2) with sat. NaHCO₃ (50 mL x 2) and brine (30 mL). The organic layer was dried and concentrated to give 7-(1-bromoethyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as a yellow solid (11.8 g, 49.3%). LCMS calculated for C₉H₁₀BrN₂OS (M+H)⁺: *m/z* 274.96, 272.96;

Found: 274.75, 272.75. ^1H NMR (400 MHz, DMSO- d_6) δ 7.05 (m, 1H), 6.27 (s, 1H), 5.17 (q, J = 6.9 Hz, 1H), 2.65 (s, 3H), 1.85 (d, J = 6.9 Hz, 3H).

Step 2. 6-Bromo-7-(1-bromoethyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one


[0284]



[0285] Under nitrogen, a suspension of 7-(1-bromoethyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (13.2 g, 48.3 mmol) in acetonitrile (100 mL, 2000 mmol) was stirred until a clear solution was obtained. *N*-Bromosuccinimide (9.891 g, 55.57 mmol) was then added and the reaction mixture was stirred at 50 °C. After 20 min, HPLC indicated that the reaction was complete. A solution of sodium sulfite (3.046 g, 24.16 mmol) in water (50 mL) was added and the mixture was stirred at room temperature for 20 min. Water (200 mL) was added slowly and the mixture stirred at room temperature for 30 min. and then filtered. The solid was washed with water (100 mL x 3) and dried to give 6-bromo-7-(1-bromoethyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (15.61 g, 91%). LCMS calculated for $\text{C}_9\text{H}_9\text{Br}_2\text{N}_2\text{OS}$ ($\text{M}+\text{H}^+$): m/z 352.87, 354.87; Found: 352.65, 354.60. ^1H NMR (400 MHz, DMSO- d_6) δ 7.15 (q, J = 1.3 Hz, 1H), 5.51 (q, J = 6.7 Hz, 1H), 2.66 (d, J = 1.2 Hz, 3H), 1.90 (d, J = 6.7 Hz, 3H).

Step 3. 7-(1-Azidoethyl)-6-bromo-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one


[0286]



[0287] To a suspension of 6-bromo-7-(1-bromoethyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (6.85 g, 19.4 mmol) in *N,N*-dimethylformamide (30.1 mL) was added sodium azide (1.45 g, 22.4 mmol). The mixture slowly turned clear after 5-10 min. After 50 min, a solution of sodium bicarbonate (4.7 g, 56 mmol) in water (90 mL) was added dropwise with stirring. The mixture was stirred at room temperature for 1 h and the solid precipitates were filtered off. The solid was then washed with water (30 mL x 3), and dried to give 7-(1-azidoethyl)-6-bromo-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one as an off-white solid (5.94 g, 97.2%). LCMS calculated for $\text{C}_9\text{H}_9\text{BrN}_5\text{OS}$ ($\text{M}+\text{H}^+$): m/z 313.96, 315.96; Found: 313.75, 315.75. ^1H NMR (400 MHz, DMSO- d_6) δ 7.15 (q, J = 1.3 Hz, 1H), 4.83 (q, J = 6.8 Hz, 1H), 2.67 (d, J = 1.4 Hz, 3H), 1.48 (d, J = 6.8 Hz, 3H).

Step 4. 7-(1-Aminoethyl)-6-bromo-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

[0288]




[0289] 7-(1-Azidoethyl)-6-bromo-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (22.6 g, 71.9 mmol) was mixed with methanol (200 mL). Sodium iodide (64.7 g, 432 mmol) was added and stirred at room temperature for 10 min. Chlorotrimethylsilane (54.8 mL, 432 mmol) was dissolved in methanol (29.1 mL) and added dropwise over 10 min. at 5-25 °C. The reaction mixture was stirred at room temperature for 10 min. HPLC and TLC showed that the reaction was complete. The reaction was quenched by addition of a solution of sodium thiosulfate (69.4 g, 439 mmol) in water (259 mL) while maintaining the batch temperature at 5- 25 °C. A large amount of solid was formed, and the pH of the mixture was 3.

The mixture was stirred at 0- 5 °C for 30 min. The pH was adjusted to 11 using 3 N aqueous sodium hydroxide (85 mL). In order to facilitate product purification and isolation, the *N*-Boc derivative of the product was prepared. To the mixture was added di-*t*-butyldicarbonate (28.3 g, 129 mmol) and the reaction mixture was stirred at room temperature for 2 h. HPLC indicated a small amount of amine remained unreacted. Additional di-*t*-butyldicarbonate (10.0 g, 45.8 mmol) was added followed by 3 N aqueous sodium hydroxide (15 mL) to adjust the pH to 11. The reaction mixture was stirred at room temperature for 30 min. The reaction mixture was extracted with ethyl acetate (150 mL x 3). The organic solution which contained the *N*-Boc derivative of the product was dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to give a residue. The residue was added to a 4 M solution of hydrogen chloride in 1,4-dioxane (206 mL, 824 mmol) and stirred at room temperature for 1.5 h. HPLC indicated the *N*-Boc-deprotection was complete. The hydrochloride salt of the product was isolated by filtration, the solid washed with MTBE, dried by suction filtration for 1 h to give 7-(1-aminoethyl)-6-bromo-3-methyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one hydrochloride salt (25.1 g) as a purple powder.

[0290] The hydrochloride salt was dissolved in water (50 mL) and a 50% solution of sodium hydroxide (about 5 mL) was added to adjust the pH to 11. The mixture was stirred at room temperature for 20 min. The product precipitated and was isolated by filtration. The wet solid was washed with water (10 mL) and dried on the filter under vacuum for 18 h to give 7-(1-aminoethyl)-6-bromo-3-methyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one (18.8 g, 65.2 mmol, 90.7 % yield) as a yellow powder. LCMS calculated for $C_9H_{11}BrN_3OS$ ($M+H$)⁺: m/z 287.97, 289.97; Found: 287.75, 289.75. ¹H NMR (400 MHz, DMSO-*d*₆) 7.08 (q, *J* = 1.3 Hz, 1H), 4.19 (q, *J* = 6.7 Hz, 1H), 2.65 (d, *J* = 1.3 Hz, 3H), 1.17 (d, *J* = 6.7 Hz, 3H).

Step 5. (S)-1-(6-bromo-3-methyl-5-oxo-5*H*-thiazolo[3,2-*a*]pyrimidin-7-yl)ethanaminium (S)-2-hydroxy-2-phenylacetate

[0291]



[0292] 7-(1-Aminoethyl)-6-bromo-3-methyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one (18.8 g, 65.2 mmol) was dissolved in isopropanol (375 mL) at reflux and then (S)-(+)-mandelic acid (4.84 g, 31.8 mmol) in isopropanol (375 mL) was added dropwise to the amine solution over 35 min. The reaction mixture was allowed to cool to about 72 °C and solid precipitation was observed. The slurry was cooled to room temperature and stirred for 1 hour. The solid product was collected by filtration. The wet cake was washed with isopropanol (100 mL) and dried on the filter under suction for 1 h to give the product (S)-1-(6-bromo-3-methyl-5-oxo-5*H*-thiazolo[3,2-*a*]pyrimidin-7-yl)ethanaminium (S)-2-hydroxy-2-phenylacetate (11.9 g) as a white solid. Chiral HPLC analysis was performed on a Lux Cellulose-2, 4.6 x 250 mm, 5 micron column using 60% ethanol/40% hexanes as the mobile phase at a flow rate of 1 mL/ min. The major enantiomer eluted at retention time 11.21 min (99.0 area %). The minor enantiomer eluted at retention time 14.31 min (0.96 area %). The e.e. of the desired product was 98.08%.


[0293] The product at 98.08% e.e. (11.9 g) was suspended in isopropanol (750 mL) and heated under reflux for 30 min. The slurry was cooled to room temperature with stirring. The solid was collected by filtration. The wet solid was washed with isopropanol (100 mL) and dried on the filter under suction for 18 h to give 10.9 g of white solid. Chiral HPLC by the method described above gave e.e. of 98.48%.

[0294] The product at 98.48% e.e. (10.9 g) was stirred in a solution of sodium carbonate (3.9 g, 37 mmol) in water (100 mL) at room temperature for 30 min. The solid free base was collected by filtration, washed with water (20 mL) and dried on the filter under suction for 2 h to give a slightly wet cake (13 g). The wet solid was dissolved in isopropanol (325 mL) at reflux and a solution of (S)-(+)-mandelic acid (3.613 g, 23.75 mmol) in isopropanol (325 mL) was added dropwise over 20 min to the free base solution. The solution was cooled to room temperature with stirring. The solid product was collected by filtration, washed with isopropanol (100 mL) and dried on the filter under suction for 48 h to give pure product (S)-1-(6-bromo-3-methyl-5-oxo-5*H*-thiazolo[3,2-*a*]pyrimidin-7-yl)ethanaminium (S)-2-hydroxy-2-phenylacetate (8.4 g, 19.08 mmol, 29.3 % yield) as a white solid. The e.e. of this sample was determined to be 100 % as no minor enantiomer (retention time = 14.31 min) was detected. LCMS calculated for $C_9H_{11}BrN_3OS$ ($M+H$)⁺ for the free base: m/z 289.97, 287.97; Found: 289.75, 287.75. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.33 (d, *J* = 7.5 Hz, 2H), 7.22 (dd, *J* = 7.1, 7.5 Hz, 2H), 7.16 (m, 2H), 4.61 (s, 1H), 4.47 (q, *J* = 6.9 Hz, 1H), 2.68 (d, *J* = 1.1 Hz, 3H), 1.31 (d, *J* = 6.8 Hz, 3H).

[0295] In order to determine the absolute stereochemistry of the product, a sample was sublimed at about 105 °C to provide colorless needles suitable for X-ray crystal structure analysis. The study determined the absolute configuration of the amine bearing carbon (C-8) is S (see Example 16 and Figure 1).

Step 6. (S)-tert-Butyl 1-(6-(3-fluorophenyl)-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethylcarbamate

[0296]




[0297] (S)-1-(6-Bromo-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethanaminium (S)-2-hydroxy-2-phenylacetate (4.93 g, 11.2 mmol) was dissolved in THF (100 mL) and water (33 mL). Di-*t*-butyldicarbonate (3.03 g, 13.9 mmol) was added, followed by sodium bicarbonate (1.88 g, 22.4 mmol). The reaction mixture was stirred at room temperature for 30 min. at which point the HPLC showed the reaction was near complete. Additional di-*t*-butyldicarbonate (0.49 g, 2.24 mmol) was then added and the reaction mixture was stirred at room temperature for 1 hour. The reaction was shown to be complete by HPLC. The reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (2 x 100 mL). The ethyl acetate solution was concentrated to give (S)-tert-butyl 1-(6-bromo-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethylcarbamate (5.46 g, 14.1 mmol, 126% yield) which was used in the subsequent Suzuki coupling reaction without further purification.

[0298] (S)-tert-Butyl 1-(6-bromo-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethylcarbamate (5.46 g, 14.1 mmol) and (3-fluorophenyl)boronic acid (2.95 g, 21.1 mmol) were suspended in 1,4-dioxane (110 mL). A solution of sodium carbonate (4.47g, 42.2 mmol) in water (27 mL) was added to the mixture followed by tetrakis(triphenylphosphine)palladium(0) catalyst (0.81 g, 0.70 mmol). The reaction mixture was degassed and heated under nitrogen at 100 °C for 16 h. HPLC indicated the starting material was consumed. The reaction mixture was cooled to room temperature and water (100 mL) was added. The resultant mixture was extracted with ethyl acetate (2 x 100 mL). The ethyl acetate solution was washed with saturated aqueous sodium bicarbonate (100 mL), dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to give a residue. The residue was purified by flash column chromatography on silica gel using 1- 50% ethyl acetate in hexane as eluent to give (S)-tert-butyl 1-(6-(3-fluorophenyl)-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethylcarbamate (4.34 g, 10.8 mmol, 76%) as an off-white solid. LCMS calculated for C₂₀H₂₃FN₃O₃S (M+H)⁺: m/z 404.1; Found 404.1. ¹H NMR (500 MHz, DMSO-*d*₆) δ 7.48 (ddd, *J* = 8.1, 7.8, 6.2 Hz, 1H), 7.18 (m, 3H), 7.05 (q, *J* = 1.3 Hz, 1H), 6.96 (d, *J* = 7.5 Hz, 1H), 4.41 (m, 1H), 2.66 (d, *J* = 1.3 Hz, 3H), 1.33 (s, 9H), 1.13 (d, *J* = 6.8 Hz, 3H).

Step 7. (S)-7-(1-Aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one


[0299]



[0300] (S)-tert-Butyl 1-(6-(3-fluorophenyl)-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethylcarbamate (4.15 g, 10.3 mmol) was dissolved in a 4.0 M solution of hydrogen chloride in 1,4-dioxane (25.7 mL, 102.8 mmol) and the solution was stirred at room temperature for 45 min. HPLC indicated that the reaction was complete. To the solution was added water (10 mL) followed by 3 N aqueous sodium hydroxide solution at 0-5 °C to adjust the pH to 10. The aqueous mixture was extracted with ethyl acetate (2 x 30 mL). The ethyl acetate solution was dried over anhydrous sodium sulfate, filtered, and concentrated under reduced pressure to give (S)-7-(1-Aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (3.30 g, 10.88 mmol, 103% yield). LCMS calculated for C₁₅H₁₅FN₃OS (M+H)⁺: m/z 304.08; Found 303.9. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.45 (ddd, *J* = 8.1, 7.9, 5.9 Hz, 1H), 7.19 (m, 1H), 7.12 (m, 2H) 7.04 (q, *J* = 1.1 Hz, 1H), 3.57 (q, *J* = 6.6 Hz, 1H), 2.64 (d, *J* = 1.3 Hz, 3H), 1.10 (d, *J* = 6.7 Hz, 3H)

Step 8. (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0301]



[0302] (S)-7-(1-Aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (2.30 g, 7.58 mmol), 6-bromo-9H-purine (2.716 g, 13.65 mmol), *N,N*-diisopropylethylamine (6.60 mL, 37.9 mmol) were dissolved in ethanol (15 mL) and the resultant mixture was heated at reflux under a nitrogen atmosphere for 17 h. HPLC indicated the reaction was complete. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by flash column chromatography on silica gel using gradient elution starting at 100% DCM with increasing polarity to 25% of a mixture of DCM/MeOH/ aq. NH_4OH (100:5:0.5, v/v/v) in DCM. After the silica chromatography, 2.1 g of crude product was obtained. This crude product was further purified by preparative reversed phase HPLC using 0.1% TFA in water and acetonitrile as mobile phases at a flow rate of 60 mL/min. on a SunFirc C 18, 5 μM , 30 x 100 mm column. Pure (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt (trifluoroacetic acid salt) (1.86 g, 3.47 mmol, 45.8 % yield) was obtained as a white solid after lyophilization. LCMS calculated for $\text{C}_{20}\text{H}_{17}\text{FN}_7\text{OS}$ ($\text{M}+\text{H}$)⁺ for the free base: *m/z* 422.1; Found: 422.0). ¹H NMR (500 MHz, $\text{DMSO}-d_6$) δ 9.03 (br s, 1H), 8.53 (s, 1H), 8.51 (s, 1H), 7.47 (m, 1H), 7.21 (m, 3H), 7.09 (s, 1H), 5.23 (m, 1H), 2.65 (d, *J* = 1.3 Hz, 3H), 1.43 (d, *J* = 7.0 Hz, 3H). ¹³C NMR (125 MHz, $\text{DMSO}-d_6$) δ 164.0, 162.1 (*J*_{CF} = 244.9 Hz), 160.5, 160.3, 150.9, 147.6, 147.5, 144.4, 135.9, 135.9, 130.2 (*J*_{CF} = 8.3 Hz), 126.9, 117.4 (*J*_{CF} = 22.6 Hz), 116.1, 114.8 (*J*_{CF} = 21.5 Hz), 111.1, 107.8, 48.5, 19.6, 18.0. Reversed phase analytical HPLC showed purity at 99.8 area %. Chiral HPLC analysis was performed on Chiralcel OJ-H, 4.6 x 250 mm, 5 micron column using 60 % ethanol/40 % hexanes as eluent at a flow rate of 0.5 mL/min. The peak for the desired enantiomer (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one was observed at a retention time of 21.171 min. (99.1 area %). The minor peak for the undesired enantiomer (R)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one was observed at a retention time of 13.358 min (0.9 area %). The enantiomeric excess of the desired enantiomer was 98.2%.

Example 15A. (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one

[0303]




[0304] A mixture of (S)-7-(1-aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (108.2 g, 357 mmol), 6-chloropurine (71.73 g, 464.1 mmol), and *N,N*-diisopropylethylamine (74.6 mL, 428.4 mmol) in 1-butanol (952 mL) was degassed with nitrogen bubbling for 5 minutes. The reaction mixture was heated at 105 °C under nitrogen for 15 hours, at which point HPLC indicated amine was consumed. The reaction mixture was cooled down to room temperature before being treated with water (200 mL) at room temperature. The resulting mixture was concentrated under reduced pressure to give an oily residue and the residue was treated with CH_2Cl_2 (1000 mL) to give a brownish clear solution. The resulting solution was washed with 2.5% aqueous sodium carbonate solution (Na_2CO_3 , 250 mL x 2) and the organic layer was concentrated under reduced pressure to afford the crude desired product as a brownish solid. The solution of the crude desired product in CH_2Cl_2 was absorbed onto silica gel (300 g) and the dried silica gel was loaded onto a flash column. The flash column was eluted with pure CH_2Cl_2 and a mixture of CH_2Cl_2 , MeOH and aqueous NH_4OH (2000:10:5) to afford pure desired product. The fractions containing pure desired product were combined and concentrated under reduced pressure. The resulting yellowish solid (90.3 g) was dissolved in a mixture of CH_2Cl_2 and methanol (500 : 50 mL). The resulting solution was treated with ethyl acetate (900 mL) and the resulting mixture was

distilled until the internal solution temperature reached 68 °C. The mixture was then cooled to room temperature and subsequently to 0 - 5 °C for 1 hour. The solids were collected by filtration, washed with cold ethyl acetate (100 mL), and dried overnight on the filter under vacuum to afford (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one (89.5 g, 59.4% yield, 99.4% ee) as a light yellowish solid. LCMS calculated for C₂₀H₁₇FN₇OS (M+H)⁺ for the free base: m/z 422.1; Found: 422.0; ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.40 (s, 1H), 7.99 (br s, 1H), 7.45 (m, 1H), 7.21-7.12 (m, 3H), 6.8 (m, 1H), 6.42 (s, 1H), 5.52 (br s, 1H), 2.79 (d, *J* = 1.3 Hz, 3H), 1.43 (d, *J* = 7.0 Hz, 3H).

Example 16. X-Ray Crystallography of (S)-1-(6-bromo-3-methyl-5-oxo-5H-thiazolo[3,2-a]pyrimidin-7-yl)ethan-aminium (S)-2-hydroxy-2-phenylacetate (From Example 15, Step 5)

[0305]



In order to determine the absolute stereochemistry of the product from Example 15, step 5, a sample was sublimed at about 105 °C to provide colorless needles suitable for X-ray crystal structure analysis. The study determined the absolute configuration of the amine bearing carbon (C-8) is S.

[0306] DATA COLLECTION: Bruker SMART APEX-II CCD system, MoK α radiation, standard focus tube, anode power = 50kV x 42 mA, crystal to plate distance = 5.0 cm, 512 x 512 pixels/frame, beam center = (256.13,253.14), total frames = 1081, oscillation/frame = 0.50°, exposure/frame = 300.1 sec/frame, SAINT integration, hkl min/max = (-4, 7, -14,14, -31, 35), data input to shelx = 11285, unique data = 3870, two-theta range = 3.82 to 53.64°, completeness to two-theta 53.64 = 99.70%, R(int- σ) = 0.0908, SADABS correction applied.

[0307] SOLUTION AND REFINEMENT: Structure solved using XS(Shelxtl), refined using shelxtl software package, refinement by full-matrix least squares on F², scattering factors from Int. Tab. Vol C Tables 4.2.6.8 and 6.1.1.4, number of data = 3870, number of restraints = 0, number of parameters = 309, data/parameter ratio = 12.52, goodness-of-fit on F² = 0.99, R indices[σ] R1 = 0.0455, wR2 = 0.0674, R indices(all data) R1 = 0.1059, wR2 = 0.0825, max difference peak and hole = 0.420 and -0.863 e/Å³, refined flack parameter = 0.025(11). All of the hydrogen atoms have been found from a difference map and fully refined.

[0308] CRYSTAL DATA: C₁₇H₁₈BrN₃O₄S, from sublimation @ 105 °C, colorless, needle, -0.160 x 0.020 x 0.020 mm, orthorhombic, P2₁2₁2₁, a = 5.5572(18) Å, b = 11.547(4) Å, c = 28.207(10) Å, Vol = 1810.1 (11) Å³, Z = 4, T = -100.°C, Formula weight = 440.31, Density = 1.616 g/cm³, μ (Mo) = 2.41 mm⁻¹.

[0309] RESULTS: This study determined the structure of C₁₇H₁₈N₃O₄S₁Br₁ for the product of Example 15, step 5. The asymmetric unit contains one of each molecule as shown in Figure 1 with thermal ellipsoids drawn to the 50% probability level. The predicted structure was confirmed. The molecules form an infinite hydrogen bonded chain via the NH₃'s along the a-axis which is the needle axis, as shown in Figure 2. The absolute configuration was determined to be S at both C8 and C16 based upon the refinement of the flack parameter = 0.02(5). The configuration of C16 was known to be S.

Table A1. Atomic coordinates (x 10⁴) and equivalent isotropic displacement parameters (Å² x 10³). U(eq) is defined as one third of the trace of the orthogonalized U_{ij} tensor.

	x	y	z	U(eq)
Br(1)	5816(1)	1172(1)	1875(1)	34(1)
S(1)	-2800(2)	-1679(1)	1030(1)	28(1)
O(1)	2992(7)	-875(4)	2278(1)	48(1)
O(2)	-1818(7)	-1946(3)	-193(1)	33(1)
O(3)	-1693(5)	1147(3)	-24(1)	27(1)
O(4)	-4576(7)	-176(3)	77(1)	32(1)
N(1)	285(6)	-1133(4)	1670(1)	22(1)
N(2)	444(8)	26(3)	965(2)	25(1)
N(3)	2679(10)	1576(4)	395(2)	23(1)
C(1)	2242(9)	648(4)	1173(2)	20(1)

(continued)

		x	y	z	U(eq)
5	C(2)	3198(8)	351(4)	1606(2)	22(1)
	C(3)	2267(10)	-571(4)	1891(2)	29(1)
	C(4)	-474(9)	-820(4)	1219(2)	20(1)
	C(5)	-1135(9)	-2023(4)	1878(2)	25(1)
	C(6)	-2805(11)	-2394(5)	1565(2)	26(1)
10	C(7)	-807(18)	-2456(6)	2365(2)	41(2)
	C(8)	2920(10)	1736(4)	917(2)	23(1)
	C(9)	1254(13)	2703(5)	1078(3)	32(2)
	C(10)	-755(11)	-530(4)	-802(2)	21(1)
	C(11)	-2631(10)	-858(4)	-1102(2)	26(1)
15	C(12)	-2597(11)	-528(5)	-1570(2)	28(1)
	C(13)	-732(12)	127(4)	-1755(2)	31(1)
	C(14)	1149(11)	434(5)	-1457(2)	31(1)
	C(15)	1156(10)	102(4)	-981(2)	26(1)
	C(16)	-926(12)	-817(4)	-274(2)	24(1)
20	C(17)	-2506(10)	130(4)	-50(2)	23(1)

Table A2. Bond lengths [Å] and angles (deg)

25	Br(1)-C(2)	1.896(5)
	S(1)-C(4)	1.714(5)
	S(1)-C(6)	1.719(6)
	O(1)-C(3)	1.215(6)
30	O(2)-C(16)	1.413(6)
	O(3)-C(17)	1.260(6)
	O(4)-C(17)	1.256(6)
	N(1)-C(4)	1.387(6)
	N(1)-C(3)	1.422(6)
35	N(1)-C(5)	1.423(6)
	N(2)-C(4)	1.314(6)
	N(2)-C(1)	1.362(6)
	N(3)-C(8)	1.489(6)
40	C(1)-C(2)	1.375(7)
	C(1)-C(8)	1.497(7)
	C(2)-C(3)	1.431(7)
	C(5)-C(6)	1.350(8)
	C(5)-C(7)	1.474(8)
45	C(8)-C(9)	1.521(8)
	C(10)-C(15)	1.385(8)
	C(10)-C(11)	1.395(7)
	C(10)-C(16)	1.529(7)
	C(11)-C(12)	1.375(7)
50	C(12)-C(13)	1.385(8)
	C(13)-C(14)	1.388(8)
	C(14)-C(15)	1.394(8)
	C(16)-C(17)	1.538(7)
55	C(4)-S(1)-C(6)	90.4(3)
	C(4)-N(1)-C(3)	121.2(4)
	C(4)-N(1)-C(5)	113.4(4)
	C(3)-N(1)-C(5)	125.4(4)

EP 2 845 856 A1

(continued)

5	C(4)-N(2)-C(1)	116.3(5)
	N(2)-C(1)-C(2)	122.2(5)
	N(2)-C(1)-C(8)	114.8(5)
	C(2)-C(1)-C(8)	122.7(5)
	C(1)-C(2)-C(3)	123.0(5)
10	C(1)-C(2)-Br(1)	121.9(4)
	C(3)-C(2)-Br(1)	115.1(4)
	O(1)-C(3)-N(1)	121.2(5)
	O(1)-C(3)-C(2)	126.8(5)
	N(1)-C(3)-C(2)	111.9(5)
15	N(2)-C(4)-N(1)	125.1(5)
	N(2)-C(4)-S(1)	123.6(4)
	N(1)-C(4)-S(1)	111.3(3)
	C(6)-C(5)-N(1)	110.0(5)
	C(6)-C(5)-C(7)	125.9(5)
20	N(1)-C(5)-C(7)	124.1(5)
	C(5)-C(6)-S(1)	114.8(5)
	N(3)-C(8)-C(1)	110.5(4)
	N(3)-C(8)-C(9)	109.4(5)
	C(1)-C(8)-C(9)	108.6(4)
25	C(15)-C(10)-C(11)	119.6(5)
	C(15)-C(10)-C(16)	121.2(5)
	C(11)-C(10)-C(16)	119.1(5)
	C(12)-C(11)-C(10)	119.8(5)
	C(11)-C(12)-C(13)	121.5(6)
30	C(12)-C(13)-C(14)	118.4(5)
	C(13)-C(14)-C(15)	121.0(5)
	C(10)-C(15)-C(14)	119.6(5)
	O(2)-C(16)-C(10)	112.3(4)
	O(2)-C(16)-C(17)	113.0(5)
35	C(10)-C(16)-C(17)	106.3(4)
	O(4)-C(17)-O(3)	125.1(5)
	O(4)-C(17)-C(16)	116.1(5)
	O(3)-C(17)-C(16)	118.8(5)
40		

Table A3. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) (symmetry transformations used to generate equivalent atoms. The anisotropic displacement factor exponent takes the form: $-2 \pi^2 [h^2 a^{*2} U_{11} + \dots + 2 h k a^* b^* U_{12}]$

45		U11	U22	U33	U23	U13	U12
	Bar(1)	27(1)	40(1)	36(1)	-4(1)	-6(1)	-9(1)
	S(1)	27(1)	24(1)	33(1)	2(1)	-5(1)	-7(1)
	O(1)	54(3)	56(3)	33(3)	9(2)	-21(2)	-11(2)
50	O(2)	41(3)	13(2)	46(3)	3(2)	6(2)	4(2)
	O(3)	33(2)	14(2)	35(2)	5(2)	-7(1)	0(2)
	O(4)	31(3)	21(2)	44(3)	-1(2)	8(2)	0(2)
	N(1)	22(2)	22(2)	21(2)	1(2)	-3(2)	-3(2)
55	N(2)	26(3)	21(2)	26(3)	5(2)	-2(2)	0(2)
	N(3)	27(3)	22(3)	19(3)	4(2)	3(2)	-8(2)
	C(1)	16(3)	19(3)	26(3)	-4(2)	0(2)	-3(2)
	C(2)	13(3)	23(3)	29(3)	1(3)	1(2)	-2(2)

(continued)

		U11	U22	U33	U23	U13	U12
5	C(3)	29(3)	31(3)	27(3)	-1(3)	-5(3)	2(3)
	C(4)	23(3)	17(3)	20(3)	4(2)	0(2)	1(2)
	C(5)	26(3)	22(3)	29(3)	11(3)	6(3)	-2(2)
	C(6)	22(3)	17(3)	40(4)	4(3)	0(3)	-4(3)
	C(7)	56(5)	36(4)	31(4)	11(3)	1(4)	-2(4)
10	C(8)	23(3)	21(3)	26(3)	-7(3)	-4(2)	-4(2)
	C(9)	34(5)	21(3)	41(5)	-3(3)	6(3)	-9(3)
	C(10)	23(3)	17(3)	23(3)	1(2)	6(3)	9(3)
	C(11)	26(3)	22(3)	31(4)	1(3)	3(3)	-6(3)
	C(12)	26(3)	19(3)	38(4)	-7(3)	-4(3)	11(3)
15	C(13)	37(3)	22(3)	32(4)	3(2)	5(3)	8(3)
	C(14)	22(4)	25(3)	45(4)	8(3)	10(3)	0(3)
	C(15)	27(3)	20(3)	33(3)	-1(3)	7(3)	6(3)
	C(16)	30(3)	10(3)	31(3)	-2(2)	1(3)	3(3)
	C(17)	29(3)	18(3)	22(3)	1(2)	-3(3)	6(3)
20							

Table A4. Hydrogen coordinates (x 10⁴) and isotropic displacement parameters (Å² x 10³)

		x	y	z	U(eq)
25	H(2)	-3600(200)	-1860(80)	-110(30)	160(40)
	H(3)	4250(110)	910(50)	254(19)	59(17)
	H(3A)	1260(150)	1240(60)	310(20)	80(20)
	H(3B)	2910(100)	2160(50)	266(19)	27(18)
	H(6)	-3610(100)	-2890(40)	1579(18)	25(18)
30	H(7)	-2240(160)	-3020(80)	2480(30)	120(30)
	H(7A)	-1360(90)	-1900(50)	2583(17)	30(17)
	H(7B)	640(120)	-2750(50)	2426(19)	41(19)
	H(8)	4800(80)	1970(40)	1003(17)	28(15)
	H(9)	2070(100)	3440(50)	950(20)	56(19)
35	H(9A)	1570(110)	2790(50)	1430(20)	60(20)
	H(9B)	-210(100)	2520(50)	1035(19)	34(19)
	H(11)	-3890(80)	-1350(40)	-963(14)	16(12)
	H(12)	-3720(100)	-740(50)	-1780(20)	60(20)
	H(13)	-670(100)	380(40)	-2129(18)	40(15)
40	H(14)	2390(110)	910(50)	-1573(18)	45(17)
	H(15)	2710(120)	320(50)	-760(20)	70(20)
	H(16)	780(100)	840(40)	-125(18)	50(16)
45					

Table A5. Torsion angles [deg].

50	C(4)-N(2)-C(1)-C(2)	5.5(7)
	C(4)-N(2)-C(1)-C(8)	-168.8(4)
	N(2)-C(1)-C(2)-C(3)	-5.2(8)
	C(8)-C(1)-C(2)-C(3)	168.7(5)
	N(2)-C(1)-C(2)-Br(1)	176.5(4)
55	C(8)-C(1)-C(2)-Br(1)	-9.6(7)
	C(4)-N(1)-C(3)-O(1)	-177.1(5)
	C(5)-N(1)-C(3)-O(1)	4.4(8)
	C(4)-N(1)-C(3)-C(2)	3.3(7)


EP 2 845 856 A1

(continued)

	C(5)-N(1)-C(3)-C(2)	-175.2(4)
	C(1)-C(2)-C(3)-O(1)	-179.0(5)
5	Br(1)-C(2)-C(3)-O(1)	-0.7(8)
	C(1)-C(2)-C(3)-N(1)	0.6(7)
	Br(1)-C(2)-C(3)-N(1)	179.0(3)
	C(1)-N(2)-C(4)-N(1)	-1.5(7)
10	C(1)-N(2)-C(4)-S(1)	178.2(4)
	C(3)-N(1)-C(4)-N(2)	-3.0(7)
	C(5)-N(1)-C(4)-N(2)	175.6(5)
	C(3)-N(1)-C(4)-S(1)	177.2(4)
	C(5)-N(1)-C(4)-S(1)	-4.1(5)
15	C(6)-S(1)-C(4)-N(2)	-177.2(5)
	C(6)-S(1)-C(4)-N(1)	2.5(4)
	C(4)-N(1)-C(5)-C(6)	3.8(6)
	C(3)-N(1)-C(5)-C(6)	-177.6(5)
20	C(4)-N(1)-C(5)-C(7)	-175.8(6)
	C(3)-N(1)-C(5)-C(7)	2.8(8)
	N(1)-C(5)-C(6)-S(1)	-1.8(6)
	C(7)-C(5)-C(6)-S(1)	177.7(5)
	C(4)-S(1)-C(6)-C(5)	-0.4(5)
25	N(2)-C(1)-C(8)-N(3)	-34.3(6)
	C(2)-C(1)-C(8)-N(3)	151.4(5)
	N(2)-C(1)-C(8)-C(9)	85.7(6)
	C(2)-C(1)-C(8)-C(9)	-88.6(6)
30	C(15)-C(10)-C(11)-C(12)	-1.5(8)
	C(16)-C(10)-C(11)-C(12)	174.9(5)
	C(10)-C(11)-C(12)-C(13)	0.0(8)
	C(11)-C(12)-C(13)-C(14)	1.0(8)
	C(12)-C(13)-C(14)-C(15)	-0.6(8)
35	C(11)-C(10)-C(15)-C(14)	1.9(8)
	C(16)-C(10)-C(15)-C(14)	-174.4(5)
	C(13)-C(14)-C(15)-C(10)	-0.9(8)
	C(15)-C(10)-C(16)-O(2)	-141.3(5)
40	C(11)-C(10)-C(16)-O(2)	42.3(7)
	C(15)-C(10)-C(16)-C(17)	94.7(6)
	C(11)-C(10)-C(16)-C(17)	-81.7(6)
	O(2)-C(16)-C(17)-O(4)	-15.1(7)
	C(10)-C(16)-C(17)-O(4)	108.4(5)
45	O(2)-C(16)-C(17)-O(3)	167.6(4)
	C(10)-C(16)-C(17)-O(3)	-68.9(6)

Example 17. 6-(3,5-difluorophenyl)-3-methyl-7-[1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0310]



[0311] A mixture of 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one hydrochloride (0.030 g, 0.084 mmol), 4-chloropyrrolo[2,3-d]pyrimidine (0.013 g, 0.084 mmol), and *N,N*-diisopropylethylamine (0.044 mL, 0.25 mmol) in isopropyl alcohol (0.2 mL) was heated at 100 °C, in a sealed tube, for three days. The resultant mixture was applied on RP-HPLC (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product. LCMS calculated for C₂₁H₁₇F₂N₆OS(M+H)⁺: m/z = 439.1; Found: 439.1.

Example 18. 6-(3,5-difluorophenyl)-7-{1-[(2-fluoro-9H-purin-6-yl)amino]ethyl}-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0312]



[0313] A mixture of 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one hydrochloride (0.030 g, 0.084 mmol), 2-fluoro-6-chloropurine (0.015 g, 0.084 mmol), and *N,N*-diisopropylethylamine (0.044 mL, 0.25 mmol) in isopropyl alcohol (0.2 mL) was heated at 100 °C for three days, in a sealed tube. The resultant mixture was purified on RP-HPLC (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product. LCMS calculated for C₂₀H₁₅F₃N₇OS(M+H)⁺: m/z = 458.1; Found: 458.0.


Example 19. 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0314]



Step 1. 7-(1-aminoethyl)-3-methyl-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0315]



[0316] To a stirred solution of 7-(1-azidoethyl)-3-methyl-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.050 g, 0.16 mmol) in tetrahydrofuran (0.5 mL) and water (0.12 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.19 mL, 0.19 mmol) at room temperature and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate (EtOAc) and the mixture was extracted with 1 N HCl two times. The combined extracts were neutralized with solid sodium bicarbonate, and extracted with dichloromethane. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue, shown two peaks with same desired mass, was used directly in next step. LCMS calculated for $C_{14}H_{15}N_4OS(M+H)^+$: $m/z = 287.1$; Found: 287.0.

Step 2. 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0317]



[0318] A mixture of 6-bromo-9H-purine (0.064 g, 0.32 mmol), 7-(1-aminoethyl)-3-methyl-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.046 g, 0.16 mmol), and *N,N*-diisopropylethylamine (0.056 mL, 0.32 mmol) in ethanol (0.5 mL) was heated at reflux under nitrogen overnight. The mixture was evaporated and the resultant residue was purified on RP-HPLC (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH_4OH) to give the product as the free base. LCMS calculated for $C_{19}H_{17}N_8OS(M+H)^+$: $m/z = 405.1$; Found: 405.1.


Example 20. 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0319]



Step 1. 7-(1-azidoethyl)-3-methyl-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0320]



[0321] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.10 g, 0.32 mmol), and 2-(tributylstannyl)-1,3-thiazole (143 mg, 0.382 mmol) in 1,4-dioxane (3 mL) was added tetrakis(triphenylphosphine)palladium(0) (18 mg, 0.016 mmol). The reaction mixture was heated at 120 °C overnight. After cooling to room temperature, the mixture concentrated under reduced pressure. The crude mixture was purified on silica gel, eluting with 0 to 60% ethyl acetate in hexane, to give the desired product (73 mg, 72%). LCMS calculated for $C_{12}H_{11}N_6OS_2(M+H)^+$: $m/z = 319.0$; Found: 319.0.

Step 2. 7-(1-aminoethyl)-3-methyl-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0322]



[0323] To a stirred solution of 7-(1-azidoethyl)-3-methyl-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.030 g, 0.094 mmol) in tetrahydrofuran (0.3 mL) and water (0.068 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.11 mL, 0.11 mmol) at room temperature and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate and the mixture was extracted with 1 N HCl two times. The combined extracts were neutralized with solid sodium bicarbonate, and extracted with dichloromethane. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step. LCMS calculated for $C_{12}H_{13}N_4OS_2(M+H)^+$: $m/z = 293.1$; Found: 293.0.

Step 3. 3-methyl-7-(1-(9H-purin-6-ylamino)ethyl)-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0324]



[0325] A mixture of 6-bromo-9H-purine (0.038 g, 0.19 mmol), 7-(1-aminoethyl)-3-methyl-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.028 g, 0.096 mmol), and *N,N*-diisopropylethylamine (0.033 mL, 0.19 mmol) in ethanol (0.3 mL) was heated at reflux under nitrogen overnight. The mixture was evaporated and the resultant residue was purified on RP-HPLC (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% trifluoroacetic acid (TFA)) to give the product as a TFA salt. LCMS calculated for $C_{17}H_{15}N_8OS_2(M+H)^+$: $m/z = 411.1$; Found: 411.0.


Example 21. 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0326]



Step 1. 7-(1-azidoethyl)-3-methyl-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0327]



[0328] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.10 g, 0.32 mmol) and 4-(tributylstannyl)-1,3-thiazole (143 mg, 0.382 mmol) in 1,4-dioxane (3 mL) was added tetrakis(triphenylphosphine)palladium(0) (18.4 mg, 0.0159 mmol). The reaction mixture was heated at 120 °C overnight. After cooling to room temperature, the mixture concentrated under reduced pressure. The crude mixture was purified on silica gel, eluting with 0 to 60% EtOAc in hexane, to give the desired product (82 mg, 81%). LCMS calculated for $C_{12}H_{11}N_6OS_2(M+H)^+$: $m/z = 319.0$; Found: 319.0.

Step 2. 7-(1-aminoethyl)-3-methyl-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0329]



[0330] To a stirred solution of 7-(1-azidoethyl)-3-methyl-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.030 g, 0.094 mmol) in tetrahydrofuran (0.3 mL) and water (0.068 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.113 mL, 0.113 mmol) at room temperature and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate and the mixture was extracted with 1 N HCl two times. The combined extracts were neutralized with solid sodium bicarbonate, and extracted with dichloromethane. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step. LCMS calculated for $C_{12}H_{13}N_4OS_2(M+H)^+$: $m/z = 293.1$; Found: 293.0.

Step 3. 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0331]



[0332] A mixture of 6-bromo-9H-purine (0.038 g, 0.19 mmol), 7-(1-aminoethyl)-3-methyl-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.028 g, 0.096 mmol), and *N,N*-diisopropylethylamine (0.033 mL, 0.19 mmol) in ethanol (0.3 mL) was heated at reflux under nitrogen overnight. The mixture was evaporated and the resultant residue was purified on RP-HPLC (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the product as a TFA salt. LCMS calculated for $C_{17}H_{15}N_8OS_2(M+H)^+$: $m/z = 411.1$; Found: 411.0.


Example 22. 6-(4-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0333]



Step 1. 7-(1-azidoethyl)-6-(4-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0334]



[0335] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.10 g, 0.32 mmol) and 4-fluorophenylboronic acid (53 mg, 0.38 mmol) in 1,4-dioxane (2 mL) was added a 1 M solution of sodium carbonate in water (0.38 mL, 0.38 mmol) and tetrakis(triphenylphosphine)palladium(0) (18 mg, 0.016 mmol). The reaction mixture was heated at 100 °C overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate, washed with water, brine, dried over $MgSO_4$, and concentrated. The crude mixture was purified on silica gel, eluting with 0 to 40% EtOAc in hexane, to give the desired product (69 mg, 66%). LCMS calculated for $C_{15}H_{13}FN_5OS(M+H)^+$: $m/z = 330.1$; Found: 330.0.

Step 2. 7-(1-aminoethyl)-6-(4-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0336]



[0337] To a stirred solution of 7-(1-azidoethyl)-6-(4-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.062 g, 0.19 mmol) in tetrahydrofuran (0.6 mL) and water (0.14 mL), was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.226 mL, 0.226 mmol) at room temperature, and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate, and then the mixture was extracted with 1 N HCl two times. The combined extracts were neutralized with solid sodium bicarbonate and extracted with dichloromethane. The combined organic layers were washed with brine, dried over magnesium sulfate, and concentrated under reduced pressure. The residue was used directly in next step. LCMS calculated for $C_{15}H_{15}FN_3OS(M+H)^+$: $m/z = 304.1$; Found: 304.1.

Step 3. 6-(4-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0338]



[0339] A mixture of 6-bromo-9H-purine (0.076 g, 0.38 mmol), 7-(1-aminoethyl)-6-(4-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.058 g, 0.19 mmol), and *N,N*-diisopropylethylamine (0.066 mL, 0.38 mmol) in ethanol (0.6 mL) was heated at reflux under nitrogen overnight. The mixture was evaporated, and the resultant residue was purified on RP-HPLC (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the product as a TFA salt. LCMS calculated for $C_{20}H_{17}FN_7OS(M+H)^+$: $m/z = 422.1$; Found: 422.1.


Example 23. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0340]



Step 1. 7-(1-azidoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0341]



[0342] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (1.24 g, 3.95 mmol) and (3,5-difluorophenyl)boronic acid (0.748 g, 4.74 mmol) in 1,4-dioxane (25 mL) was added a 1 N solution of sodium carbonate in water (5.92 mL, 5.92 mmol) and tetrakis(triphenylphosphine)palladium(O) (0.27 g, 0.24 mmol). The mixture was heated at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water and brine, dried over $MgSO_4$, and concentrated. The residue was purified on silica gel (0-40% EtOAc/Hex) to give the desired product (0.42 g, 31 %). LCMS calculated for $C_{15}H_{12}F_2N_5OS(M+H)^+$: $m/z = 348.1$; Found: 348.0. The product was subjected to chiral HPLC separation (ChiralPak IA Column: 20 x 250 mm, 5 μm ; Mobile Phase: 5% Ethanol - 95% Hexanes; Flow Rate: 15 mL/min) to give two enantiomers. On analytic HPLC (ChiralPak IA Column: 4.6 x 250 mm, 5 μm ; Mobile Phase: 5% Ethanol-95% Hexanes; Flow Rate: 1 mL/min), the first enantiomer has retention time of 7.78 min and the second peak has retention time of 8.61 minutes.

Step 2. 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0343]



[0344] To a stirred solutions of 7-(1-azidoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.15 g, 0.43 mmol) (1 st peak from chiral separation) in tetrahydrofuran (2 mL) and water (0.5 mL) were added 1.00 M of trimethylphosphine in tetrahydrofuran (0.52 mL, 0.52 mmol) at room temperature and the mixtures were stirred at room temperature for 1 hour. To the mixture was added EtOAc and the mixture was extracted with aqueous 1 N HCl solution (three times). The combined extracts were neutralized with solid Na₂CO₃ and extracted with dichloromethane (two times). The combined organic layers were washed with brine, dried over Na₂SO₄, and concentrated to give the desired product (134 mg, 96.6%). LCMS calculated for C₁₅H₁₄F₂N₃OS(M+H)⁺: m/z = 322.1; Found: 322.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3, 5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0345]



[0346] A mixture of optical pure 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.13 g, 0.40 mmol) made from above, 2-amino-6-bromopurine (0.10 g, 0.47 mmol), and *N,N*-diisopropylethylamine (0.085 mL, 0.49 mmol) in ethanol (1 mL) was heated at 110 °C overnight. LCMS showed incomplete conversion. An additional 0.5 equivalent of 2-amino-6-bromopurine and 1.0 equivalent of *N,N*-diisopropylethylamine was added, and the mixture was stirred at 110 °C for another day. The solid was shown to be 2-amino-6-bromopurine by LCMS. The mixture was filtered, and the filtrates were purified on preparative-LCMS ((XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product (0.095 g, 52%). LCMS calculated for C₂₀H₁₇F₂N₈OS(M+H)⁺: m/z = 455.1; Found: 455.1 ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.67 (1H, s), 7.27 (1H, m), 7.17 (3H, m), 7.07 (1H, s), 6.89 (1H, br s), 5.45 (2H, br s), 5.03 (1H, m), 2.63 (3H, s), 1.30 (3H, d, *J* = 6.8 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 376.3 MHz) δ - 111 ppm.


Example 24. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0347]



Step 1. 7-(1-bromoethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0348]



[0349] A mixture of polyphosphoric acid (73.8 g, 677 mmol), 1,3-thiazol-2-amine (12.3 g, 123 mmol), and methyl 4-bromo-3-oxopentanoate (34.8 g, 166 mmol) was stirred at 110 °C overnight. After cooling, an ice-cold 10% aq. NaOH solution was slowly added to adjust the pH to 7. The mixture was filtered, and the collected precipitate was air-dried to give crude product which was directly used in next step. LCMS calculated for $C_8H_8BrN_2OS(M+H)^+$: $m/z = 259.0$; Found: 259.0.

Step 2. 6-bromo-7-(1-bromoethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0350]



[0351] A mixture of 7-(1-bromoethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (17.5 g, 67.5 mmol) and *N*-bromosuccinimide (14.2 g, 79.8 mmol) in acetonitrile (400 mL) was stirred at 80 °C under N_2 overnight. After removal of the solvent under reduced pressure, the resulting solid was dissolved in dichloromethane, washed sequentially with water, saturated aqueous $Na_2S_2O_3$ and $NaHCO_3$ solution and brine, dried over Na_2SO_4 , and then concentrated to give crude product (3.7 g), which was used in the next step without further purification. LCMS calculated for $C_8H_7Br_2N_2OS(M+H)^+$: $m/z = 336.9$; Found: 336.9.

Step 3. 7-(1-azidoethyl)-6-bromo-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0352]



[0353] A mixture of 6-bromo-7-(1-bromoethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (3.7 g, 11 mmol) and sodium azide (1.4 g, 22 mmol) in *N,N*-dimethylformamide (30 mL) was stirred at room temperature for 1.5 hour. After diluted with ethyl acetate, the mixture was washed with water, dried over Na_2SO_4 , concentrated and purified on silica gel (0-60% ethyl acetate/hexanes) to give the desired product (2.16 g). LCMS calculated for $C_8H_7BrN_5OS(M+H)^+$: $m/z = 300.0$; Found: 300.0.

Step 4. 7-(1-azidoethyl)-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0354]



[0355] To a mixture of 7-(1-azidoethyl)-6-bromo-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.50 g, 1.7 mmol) and (3,5-difluorophenyl)boronic acid (0.31 g, 2.0 mmol) in 1,4-dioxane (10 mL) was added a 1 N solution of sodium carbonate in water (2.2 mL, 2.2 mmol) and tetrakis(triphenylphosphine)palladium(O) (0.096 g, 0.083 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water and brine, dried over MgSO_4 , concentrated, and purified on silica gel (0-45% ethyl acetate/hexanes) to give the desired product (0.30 g, 53%). LCMS calculated for $\text{C}_{14}\text{H}_{10}\text{F}_2\text{N}_5\text{OS}(\text{M}+\text{H})^+$: $m/z = 334.1$; Found: 334.0.

Step 5. 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0356]




[0357] To a stirred solution of 7-(1-azidoethyl)-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.295 g, 0.885 mmol) in tetrahydrofuran (5 mL) and water (1 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (1.06 mL, 1.06 mmol) at room temperature and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate, and the mixture was extracted with aqueous 1 N HCl solution (three times). The combined extract was neutralized with solid NaHCO_3 and extracted with dichloromethane (two times). The combined organic layers were washed with brine, dried over Na_2SO_4 , and concentrated to give the desired compound (0.241 g, 88.6%), which was used directly in next step. LCMS calculated for $\text{C}_{14}\text{H}_{12}\text{F}_2\text{N}_3\text{OS}(\text{M}+\text{H})^+$: $m/z = 308.1$; Found: 308.0.

Step 6. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0358]




[0359] A mixture of 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.040 g, 0.13 mmol), 2-amino-6-bromopurine (0.056 g, 0.26 mmol), and *N,N*-diisopropylethylamine (0.045 mL, 0.26 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH_4OH), to give the desired product. LCMS calculated for $\text{C}_{19}\text{H}_{15}\text{F}_2\text{N}_8\text{OS}(\text{M}+\text{H})^+$: $m/z = 441.1$; Found: 441.1.

Example 25. 6-(3,5-difluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one**[0360]**

[0361] A mixture of 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.037 g, 0.12 mmol), 6-bromo-9H-purine (0.048 g, 0.24 mmol), and *N,N*-diisopropylethylamine (0.042 mL, 0.24 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH_4OH) to give the desired product. LCMS calculated for $\text{C}_{19}\text{H}_{14}\text{F}_2\text{N}_7\text{OS}(\text{M}+\text{H})^+$: $m/z = 426.1$; Found: 426.0. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 7.99 (1H, d, $J = 4.8$ Hz), 7.63 (1H, s), 7.53 (1H, d, $J = 4.8$ Hz), 7.24 (1H, m), 7.16 (2H, m), 6.88 (1H, br s), 5.41 (2H, br s), 5.05 (1H, m), 1.27 (3H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR ($\text{DMSO}-d_6$, 376.3 MHz) δ -111 ppm.

Example 26. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one**[0362]**


Step 1. 7-(1-azidoethyl)-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0363]

[0364] To a mixture of 7-(1-azidoethyl)-6-bromo-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.48 g, 1.6 mmol) and (3-fluorophenyl)boronic acid (0.27 g, 2.0 mmol) in 1,4-dioxane (10 mL) was added a 1 N solution of sodium carbonate in water (2.1 mL, 2.1 mmol) and tetrakis(triphenylphosphine)palladium(O) (0.092 g, 0.080 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water and brine, dried over MgSO_4 , concentrated, and purified on silica gel (0-50% ethyl acetate/hexanes) to give the desired compound (0.32 g, 63%). LCMS calculated for $\text{C}_{14}\text{H}_{11}\text{FN}_5\text{OS}(\text{M}+\text{H})^+$: $m/z = 316.1$; Found: 316.0.

Step 2. 7-(1-aminoethyl)-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0365]



[0366] To a stirred solution of 7-(1-azidoethyl)-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.32 g, 1.0 mmol) in tetrahydrofuran (5 mL) and water (1 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (1.22 mL, 1.22 mmol) at room temperature, and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate, and the mixture was extracted with aqueous 1 N HCl solution (three times). The combined extract was neutralized with solid NaHCO₃ and extracted with dichloromethane (two times). The combined organic layer was washed with brine, dried over Na₂SO₄, and concentrated to give the desired product (0.17 g, 58%). LCMS calculated for C₁₄H₁₃FN₃OS(M+H)⁺: m/z = 290.1; Found: 290.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0367]



[0368] A mixture of 7-(1-aminoethyl)-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.025 g, 0.086 mmol), 2-amino-6-bromopurine (0.033 g, 0.16 mmol), and *N,N*-diisopropylethylamine (0.027 mL, 0.16 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product. LCMS calculated for C₁₉H₁₆FN₈OS(M+H)⁺: m/z = 423.1; Found: 423.0. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 7.98 (1H, d, *J* = 4.8 Hz), 7.63 (1H, s), 7.52 (1H, d, *J* = 4.8 Hz), 7.46 (1H, m), 7.20 (2H, m), 6.84 (1H, br s), 5.41 (1H, br s), 5.07 (1H, m), 1.26 (3H, d, *J* = 6.8 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 376.3 MHz) δ -114 ppm.


Example 27. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0369]



Step 1. 7-(1-azidoethyl)-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0370]



[0371] To a mixture of 7-(1-azidoethyl)-6-bromo-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.34 g, 1.1 mmol) and phenylboronic acid (0.16 g, 1.4 mmol) in 1,4-dioxane (10 mL) was added a 1 N solution of sodium carbonate in water (1.5 mL, 1.5 mmol) and tetrakis(triphenylphosphine)palladium(O) (0.065 g, 0.057 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water and brine, dried over MgSO₄, concentrated, and purified on silica gel (0-50% ethyl acetate/hexanes) to give the desired product (0.23 g, 68%). LCMS calculated for C₁₄H₁₂N₅OS(M+H)⁺: m/z = 298.1; Found: 298.0.

Step 2. 7-(1-aminoethyl)-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0372]



[0373] To a stirred solution of 7-(1-azidoethyl)-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.23 g, 0.77 mmol) in tetrahydrofuran (5 mL) and water (1 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.93 mL, 0.93 mmol) at room temperature, and the mixture was stirred at room temperature for 1 hour. To the mixture was added ethyl acetate, and the mixture was extracted with aqueous 1 N HCl solution (three times). The combined extract was neutralized with solid NaHCO₃ and extracted with dichloromethane (two times). The combined organic layer was washed with brine, dried over Na₂SO₄, and concentrated to give the desired compound (0.13 g, 62%). LCMS calculated for C₁₄H₁₄N₃OS(M+H)⁺: m/z = 272.1; Found: 272.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0374]



[0375] A mixture of 7-(1-aminoethyl)-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.025 g, 0.092 mmol), 2-amino-6-bromopurine (0.035 g, 0.16 mmol), and *N,N*-diisopropylethylamine (0.029 mL, 0.16 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product. LCMS calculated for C₁₉H₁₇N₈OS(M+H)⁺: m/z = 405.1; Found: 405.1. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 80.1 (1H, d, *J* = 4.8 Hz), 7.65 (1H, s), 7.55 (1H, d, *J* = 4.8 Hz), 7.47 (2H, m), 7.40 (3H, m), 6.79 (1H, br s), 5.48 (2H, br s), 5.13 (1H, m), 1.29 (3H, d, *J* = 6.8 Hz) ppm.

Example 28. 6-(3-fluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0376]



[0377] A mixture of 7-(1-aminoethyl)-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.025 g, 0.086 mmol), 6-bromo-9H-purine (0.031 g, 0.16 mmol), and *N,N*-diisopropylethylamine (0.027 mL, 0.16 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product. LCMS calculated for C₁₉H₁₅FN₇OS(M+H)⁺: m/z = 408.1; Found: 408.0.

Example 29. 6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0378]



[0379] A mixture of 7-(1-aminoethyl)-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.025 g, 0.092 mmol), 6-bromo-9H-purine (0.033 g, 0.16 mmol), and *N,N*-diisopropylethylamine (0.029 mL, 0.16 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product. LCMS calculated for C₁₉H₁₆N₇OS(M+H)⁺: m/z = 390.1; Found: 390.1. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.08 (1H, s), 8.06 (1H, s), 7.97 (1H, d, *J* = 4.8 Hz), 7.51 (1H, d, *J* = 4.8 Hz), 7.44~7.33 (6H, m), 5.15 (1H, m), 1.29 (3H, d, *J* = 7.2 Hz) ppm.


Example 30. 6-(3-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0380]



Step 1. 7-(1-azidoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0381]




[0382] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.50 g, 1.6 mmol) and (3-fluorophenyl)boronic acid (0.27 g, 1.9 mmol) in 1,4-dioxane (10 mL) was added 1 N solution of sodium carbonate in water (2.1 mL, 2.1 mmol) and tetrakis(triphenylphosphine)palladium(O) (0.092 g, 0.080 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na₂SO₄, concentrated and purified on silica gel (0-40% ethyl acetate/hexanes) to give the desired product (0.32 g, 61%). LCMS calculated for C₁₅H₁₃FN₅OS(M+H)⁺: m/z = 330.1; Found: 330.0. The product was subjected to chiral HPLC separation (ChiralPak IA Column: 20 x 250 mm, 5 μm; Mobile Phase: 10% Ethanol - 90% Hexanes; Flow Rate: 18 mL/min) to give

two enantiomers. On analytic HPLC (ChiralPak IA Column: 4.6 x 250 mm, 5 μ m; Mobile Phase: 10% Ethanol-90% Hexanex; Flow Rate: 1 mL/min), the first enantiomer has retention time of 6.38 minutes and the second peak has retention time of 6.99 minutes.

Step 2. 7-(1-aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0383]



[0384] To a stirred solution of 7-(1-azidoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.14 g, 0.42 mmol) (1st peak from chiral separation) in tetrahydrofuran (3 mL) and water (0.5 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.52 mL, 0.52 mmol), and the mixture was stirred at room temperature for 1 hour. To the mixtures were added ethyl acetate and the mixtures were extracted with aqueous 1 N HCl solution (three times). The combined extracts were neutralized with solid NaHCO₃, and extracted with dichloromethane (three times). The combined organic layers were washed with brine, dried over MgSO₄, and concentrated to give the crude product (0.125 g) used directly in next step. LCMS calculated for C₁₅H₁₅FN₃OS(M+H)⁺: m/z = 304.1; Found: 304.0.

Step 3. 6-(3-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0385]



[0386] A mixture of single enantiomer 7-(1-aminoethyl)-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.125 g, 0.412 mmol) made from above, 6-bromo-9H-purine (0.148 g, 0.742 mmol), and *N,N*-diisopropylethylamine (0.144 mL, 0.824 mmol) in ethanol (1.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product (0.076 g, 44%). LCMS calculated for C₂₀H₁₇FN₇OS(M+H)⁺: m/z = 422.1; Found: 422.0. ¹H NMR (DMSO-*d*₆, 500 MHz) δ 8.05 (2H, s), 7.43 (1H, m), 7.24~7.14 (5H, m), 6.99 (1H, s), 5.08 (1H, m), 2.59 (3H, s), 1.29 (3H, d, *J* = 6.5 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 376.3 MHz) δ -114 ppm.


Example 31. 3-methyl-6-(4-methylphenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0387]



Step 1. 7-(1-azidoethyl)-3-methyl-6-(4-methylphenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0388]



[0389] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-1H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.080 g, 0.25 mmol) and (4-methylphenyl)boronic acid (0.042 g, 0.31 mmol) in 1,4-dioxane (2 mL) was added 1 N solution of sodium carbonate in water (0.38 mL, 0.38 mmol) and dichloro(bis(di-tert-butyl[4-(dimethylamino)phenyl]phosphoranyl))palladium (0.011 g, 0.015 mmol). The mixture was heated at 100 °C overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate, washed with water, dried over MgSO₄, concentrated and then purified on silica gel (0-25% ethyl acetate/hexane) to give the desired product (50 mg). LCMS calculated for C₁₆H₁₆N₅OS(M+H)⁺: m/z = 326.1; Found: 326.0.

Step 2. 7-(1-aminoethyl)-3-methyl-6-(4-methylphenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0390]



[0391] To a solution of 7-(1-azidoethyl)-3-methyl-6-(4-methylphenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.050 g, 0.15 mmol) in tetrahydrofuran (2 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.23 mL, 0.23 mmol) and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (40 mg), which was used directly in next step. LCMS calculated for C₁₆H₁₈N₃OS(M+H)⁺: m/z = 300.1; Found: 300.1.

Step 3. 3-methyl-6-(4-methylphenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0392]



[0393] A mixture of 7-(1-aminoethyl)-3-methyl-6-(4-methylphenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.040 g, 0.13 mmol), 6-bromo-9H-purine (0.040 g, 0.20 mmol), and *N,N*-diisopropylethylamine (0.046 mL, 0.27 mmol) in ethanol (0.3 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA), to give the desired product as a TFA salt. LCMS calculated for C₂₁H₂₀N₇OS(M+H)⁺: m/z = 418.1; Found: 418.1. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.57 (1H, br s), 8.39 (1H, s), 8.38 (1H, s), 7.19 (4H, s), 7.02 (1H, d, *J* = 1.2 Hz), 5.17 (1H, m), 2.59 (3H, s), 2.30 (3H, s), 1.32 (3H, d, *J* = 6.8 Hz) ppm.


Example 32. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0394]



Step 1. 7-(1-azidoethyl)-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0395]



[0396] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.12 g, 0.38 mmol) and (3-chlorophenyl)boronic acid (0.072 g, 0.46 mmol) in 1,4-dioxane (3 mL) was added a 1 N solution of sodium carbonate in water (0.5 mL, 0.5 mmol) and tetrakis(triphenylphosphine)palladium(O) (0.022 g, 0.019 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na₂SO₄, concentrated, and purified on silica gel (0-30% ethyl acetate/hexanes) to give the desired product. LCMS calculated for C₁₅H₁₃ClN₅OS(M+H)⁺: m/z = 346.1; Found: 346.0.

Step 2. 7-(1-aminoethyl)-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0397]



[0398] To a stirred solution of 7-(1-azidoethyl)-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.10 g, 0.29 mmol) in tetrahydrofuran (3 mL, 40 mmol) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.35 mL, 0.35 mmol), and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (0.090 g), which was used directly in next step. LCMS calculated for C₁₅H₁₅ClN₃OS(M+H)⁺: m/z = 320.1; Found: 320.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0399]



[0400] A mixture of 7-(1-aminoethyl)-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.045 g, 0.14 mmol), 2-amino-6-bromopurine (0.060 g, 0.28 mmol), and *N,N*-diisopropylethylamine (0.049 mL, 0.28 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a TFA salt. LCMS calculated for $C_{20}H_{18}ClN_8OS(M+H)^+$: $m/z = 453.1$; Found: 453.1. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.13 (1H, s), 7.48~7.12 (8H, s), 6.55 (1H, br s), 5.14 (1H, m), 1.33 (3H, d, $J = 6.8$ Hz) ppm.


Example 33. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0401]



Step 1. 7-(1-azidoethyl)-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0402]



[0403] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.12 g, 0.38 mmol) and (2-fluorophenyl)boronic acid (0.064 g, 0.46 mmol) in 1,4-dioxane (4 mL) was added a 1 N solution of sodium carbonate in water (0.8 mL, 0.8 mmol) and dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium (0.014 g, 0.019 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na_2SO_4 , concentrated, and purified on silica gel (0-35% ethyl acetate/hexane) to give the desired product (87 mg). LCMS calculated for $C_{15}H_{13}FN_5OS(M+H)^+$: $m/z = 330.1$; Found: 330.0.

Step 2. 7-(1-aminoethyl)-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0404]



[0405] To a stirred solution of 7-(1-azidoethyl)-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.087 g, 0.26 mmol) in tetrahydrofuran (3 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.32 mL, 0.32 mmol), and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (0.080 g), which was used directly in next step. LCMS calculated for $C_{15}H_{15}FN_3OS(M+H)^+$: $m/z = 304.1$; Found: 304.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0406]



[0407] A mixture of 7-(1-aminoethyl)-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.040 g, 0.13 mmol), 2-amino-6-bromopurine (0.056 g, 0.26 mmol), and *N,N*-diisopropylethylamine (0.046 mL, 0.26 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a diastereoisomeric mixture (TFA salts). LCMS calculated for $C_{20}H_{18}FN_8OS(M+H)^+$: $m/z = 437.1$; Found: 437.1. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.77 (1H, br s), 8.14 (1H, m), 7.45 (2H, m), 7.28 (4H, m), 7.14 (1H, m), 5.13 (1H, m), 2.65 (3H, s), 1.42 (1.5H, d, $J = 6.8$ Hz), 1.28 (1.5H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR (DMSO- d_6 , 376.3 MHz) δ -113.8, -114 ppm.


Example 34. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0408]



Step 1. 7-(1-azidoethyl)-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0409]




[0410] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.12 g, 0.38 mmol) and (2,3-difluorophenyl)boronic acid (0.072 g, 0.46 mmol) in 1,4-dioxane (3 mL) was added a 1 N solution of sodium carbonate in water (0.57 mL, 0.57 mmol) and dichloro(bis(di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl)palladium

(0.014 g, 0.019 mmol). The mixture was stirred at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na₂SO₄, concentrated, and purified on silica gel (0-30% ethyl acetate/hexane) to give the desired product (83 mg). LCMS calculated for C₁₅H₁₂F₂N₅OS(M+H)⁺: m/z = 348.1; Found: 348.0.

Step 2. 7-(1-aminoethyl)-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0411]



[0412] To a solution of 7-(1-azidoethyl)-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.083 g, 0.24 mmol) in tetrahydrofuran (3 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.29 mL, 0.29 mmol), and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (0.076 g), which was used directly in next step. LCMS calculated for C₁₅H₁₄F₂N₃OS(M+H)⁺: m/z = 322.1; Found: 322.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0413]



[0414] A mixture of 7-(1-aminoethyl)-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.038 g, 0.12 mmol), 2-amino-6-bromopurine (0.051 g, 0.24 mmol), and *N,N*-diisopropylethylamine (0.041 mL, 0.24 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a mixture of two diastereomers (TFA salt). LCMS calculated for C₂₀H₁₇F₂N₈OS(M+H)⁺: m/z = 455.1; Found: 455.1. ¹H NMR (DMSO-d₆, 400 MHz) δ 8.12 (1H, d, *J* = 9.6 Hz), 7.45 (1H, m), 7.30~7.23 (3H, m), 7.18~7.11 (3H, m), 6.56 (1H, s), 5.16 (1H, m), 2.66 (3H, s), 1.44 (1.5H, d, *J* = 6.8 Hz), 1.30 (1.5H, d, *J* = 6.8 Hz) ppm.


Example 35. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0415]



Step 1. 7-(1-azidoethyl)-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0416]



[0417] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.12 g, 0.38 mmol) and (3-chloro-5-fluorophenyl)boronic acid (0.080 g, 0.46 mmol) in 1,4-dioxane (3 mL) was added a 1 N solution of sodium carbonate in water (0.5 mL, 0.5 mmol) and tetrakis(triphenylphosphine)palladium(0) (0.022 g, 0.019 mmol). The mixture was heated at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na₂SO₄, concentrated and purified on silica gel (0-25% ethyl acetate/hexanes) to give the desired product (0.077 g, 55%). LCMS calculated for C₁₅H₁₂ClFN₅OS(M+H)⁺: m/z = 364.0; Found: 364.0.

Step 2. 7-(1-aminoethyl)-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0418]




[0419] To a stirred solution of 7-(1-azidoethyl)-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.077 g, 0.21 mmol) in tetrahydrofuran (3 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.25 mL, 0.25 mmol) and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give crude product (0.070 g), which was used directly in the next step. LCMS calculated for C₁₅H₁₄ClFN₃OS(M+H)⁺: m/z = 338.1; Found: 338.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0420]




[0421] A mixture of 7-(1-aminoethyl)-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.035 g, 0.10 mmol), 2-amino-6-bromopurine (0.058 g, 0.27 mmol), and *N,N*-diisopropylethylamine (0.047 mL, 0.27 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a TFA salt. LCMS calculated for C₂₀H₁₇ClFN₈OS(M+H)⁺: m/z = 471.1; Found: 471.0. ¹H NMR (DMSO-*d*₆, 400 MHz) δ 8.79 (1H, d, *J* = 7.2 Hz), 8.16 (1H, s), 7.39 (3H, m), 7.21 (2H, s), 7.18 (1H, m), 7.14 (1H, d, *J* = 1.2 Hz), 5.13 (1H, m), 2.65 (3H, s), 1.37 (3H, d, *J* = 6.8 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 376.3 MHz) δ -112 ppm.

Example 36. 6-(3-chlorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt**[0422]**

[0423] A mixture of 7-(1-aminoethyl)-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.045 g, 0.14 mmol), 6-bromo-9H-purine (0.056 g, 0.28 mmol), and *N,N*-diisopropylethylamine (0.049 mL, 0.28 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA), to give the desired product as a TFA salt. LCMS calculated for $C_{20}H_{17}ClN_7OS(M+H)^+$: $m/z = 438.1$; Found: 438.0. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.29 (1H, s), 7.46 (3H, m), 7.37 (1H, m), 7.08 (1H, s), 5.14 (1H, m), 2.64 (3H, s), 1.37 (3H, d, $J = 6.8$ Hz) ppm.


Example 37. 6-(3-chloro-5-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt**[0424]**

[0425] A mixture of 7-(1-aminoethyl)-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.035 g, 0.10 mmol), 6-bromo-9H-purine (0.041 g, 0.21 mmol), and *N,N*-diisopropylethylamine (0.036 mL, 0.21 mmol) in ethanol (0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a TFA salt. LCMS calculated for $C_{20}H_{16}ClFN_7OS(M+H)^+$: $m/z = 456.1$; Found: 456.0. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.52 (1H, br s), 8.39 (1H, s), 8.36 (1H, s), 7.43 (1H, d, $J = 8.0$ Hz), 7.33~7.27 (3H, m), 7.10 (1H, s), 5.15 (1H, m), 2.64 (3H, s), 1.41 (3H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR (DMSO- d_6 , 376.3 MHz) δ -112 ppm.

Example 38. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt**[0426]**

Step 1. 7-(1-azidoethyl)-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0427]



[0428] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.12 g, 0.38 mmol) and 3-fluoro-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (0.10 g, 0.46 mmol) in 1,4-dioxane (3 mL) was added a 1 N solution of sodium carbonate in water (0.57 mL, 0.57 mmol) and dichloro(bis(di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl)palladium (0.014 g, 0.020 mmol). The mixture was heated at 100 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na₂SO₄, concentrated and purified on silica gel (0-45% ethyl acetate/hexanes) to give the desired product (0.020 g, 16%). LCMS calculated for C₁₄H₁₂FN₆OS(M+H)⁺: m/z = 331.1; Found: 331.0.

Step 2. 7-(1-aminoethyl)-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0429]



[0430] To a solution of 7-(1-azidoethyl)-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.020 g, 0.060 mmol) in tetrahydrofuran (3 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.079 mL, 0.079 mmol), and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (0.018 g), which was used directly in next step. LCMS calculated for C₁₄H₁₄FN₄OS(M+H)⁺: m/z = 305.1; Found: 305.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0431]




[0432] A mixture of 7-(1-aminoethyl)-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (9 mg, 0.03 mmol), 2-amino-6-bromopurine (9.5 mg, 0.044 mmol) and *N,N*-diisopropylethylamine (0.010 mL, 0.059 mmol) in ethanol (0.3 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on prep-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a TFA salt. LCMS calculated for C₁₉H₁₇FN₉OS(M+H)⁺: m/z = 438.1; Found: 438.0. ¹HNMR (DMSO-*d*₆, 400 MHz) δ 8.82 (1H, br s), 8.59 (1H, d, *J* = 2.8 Hz), 8.42 (1H, s), 8.16 (1H, s), 7.77 (1H, dt, *J* = 9.6 and 2.4 Hz), 7.22~7.16 (4H, m),

5.08 (1H, m), 2.66 (3H, s), 1.37 (3H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR (DMSO- d_6 , 376.3 MHz) δ -128 ppm.


Example 39.7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0433]



Step 1. 7-(1-azidoethyl)-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0434]



[0435] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.12 g, 0.38 mmol) and (2-chlorophenyl)boronic acid (0.072 g, 0.46 mmol) in 1,4-dioxane (3 mL) was added a 1 N solution of sodium carbonate in water (0.57 mL, 0.57 mmol) and dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium (0.014 g, 0.019 mmol). The mixture was stirred at 105 °C overnight. After cooling, the mixture was diluted with ethyl acetate, washed with water, dried over Na_2SO_4 , concentrated, and purified on silica gel (0-30% ethyl acetate/hexanes) to give the desired product (0.062 g). LCMS calculated for $\text{C}_{15}\text{H}_{13}\text{ClN}_5\text{OS}(\text{M}+\text{H})^+$: m/z =346.1; Found: 346.0.

Step 2. 7-(1-aminoethyl)-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0436]



[0437] To a stirred solution of 7-(1-azidoethyl)-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.062 g, 0.18 mmol) in tetrahydrofuran (3 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.22 mL, 0.22 mmol), and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (0.056 g), which was used directly in next step. $\text{C}_{15}\text{H}_{15}\text{ClN}_3\text{OS}(\text{M}+\text{H})^+$: m/z = 320.1; Found: 320.0.

Step 3. 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0438]



[0439] A mixture of 7-(1-aminoethyl)-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.028 g, 0.088 mmol), 2-amino-6-bromopurine (0.037 g, 0.18 mmol), and *N,N*-diisopropylethylamine (0.030 mL, 0.18 mmol) in ethanol (0.4 mL) was heated at 110 °C overnight. The mixture was filtered and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give two diastereomers as a TFA salts. On analytic HPLC (Waters SunFire C18, 2.1x50 mm, 5 μ M; injection volume 2 μ L; flow rate 3 mL/min; at gradient from 2% to 80% acetonitrile in water containing 0.15% NH_4OH in 3 min): First peak has retention time 1.296 min; LCMS calculated for $\text{C}_{20}\text{H}_{18}\text{ClN}_8\text{OS}(\text{M}+\text{H})^+$: $m/z = 453.1$; Found: 453.0. Second peak has retention time 1.431 min; LCMS calculated for $\text{C}_{20}\text{H}_{18}\text{ClN}_8\text{OS}(\text{M}+\text{H})^+$: $m/z = 453.1$; Found: 453.0.

Example 40. 6-(2-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0440]



[0441] A mixture of 7-(1-aminoethyl)-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.040 g, 0.13 mmol), 6-bromo-9H-purine (0.052 g, 0.26 mmol), and *N,N*-diisopropylethylamine (0.046 mL, 0.26 mmol) in ethanol (0.5 mL, 8 mmol) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA) to give the desired product as a diastereoisomeric mixture (TFA salt). LCMS calculated for $\text{C}_{20}\text{H}_{17}\text{FN}_7\text{OS}(\text{M}+\text{H})^+$: $m/z = 422.1$; Found: 422.1. ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ 8.50 (1H, br s), 8.40 (1H, s), 8.38 (1H, s), 7.50 (1H, m), 7.36~7.25 (3H, m), 7.10 (1H, s), 5.14 (1H, m), 2.64 (3H, s), 1.48 (1.5H, d, $J = 6.8$ Hz), 1.34 (1.5H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR ($\text{DMSO}-d_6$, 376.3 MHz) δ -112, -114 ppm.

Example 41. 6-(2,3-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0442]




[0443] A mixture of 7-(1-aminoethyl)-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.038 g, 0.12 mmol), 6-bromo-9H-purine (0.047 g, 0.24 mmol), and *N,N*-diisopropylethylamine (0.041 mL, 0.24 mmol) in ethanol

(0.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on prep-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA), to give the desired product as a mixture of two diastereomers (TFA salt). LCMS calculated for $C_{20}H_{16}F_2N_7OS(M+H)^+$: $m/z = 440.1$; Found: 440.0. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.38~8.34 (3H, m), 7.49~7.10 (4H, m), 5.12 (1H, m), 2.64 (3H, s), 1.50 (1.5H, d, $J = 6.8$ Hz), 1.36 (1.5H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR (DMSO- d_6 , 376.3 MHz) δ -137.8, -139.8, -140.0 ppm.

Example 42. 6-(5-fluoropyridin-3-yl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0444]



[0445] A mixture of 7-(1-aminoethyl)-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (9 mg, 0.03 mmol), 6-bromo-9H-purine (8.8 mg, 0.044 mmol), and *N,N*-diisopropylethylamine (0.010 mL, 0.059 mmol) in ethanol (0.3 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA), to give the desired product as a TFA salt. LCMS calculated for $C_{19}H_{16}FN_8OS(M+H)^+$: $m/z = 423.1$; Found: 423.1. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.60 (1H, d, $J = 2.8$ Hz), 8.47 (1H, s), 8.35 (1H, s), 8.33 (1H, s), 7.82 (1H, d, $J = 9.6$ Hz), 7.12 (1H, s), 5.09 (1H, m), 2.64 (3H, s), 1.43 (3H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR (DMSO- d_6 , 376.3 MHz) δ -128 ppm.

Example 43. 6-(2-chlorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt


[0446]



[0447] A mixture of 7-(1-aminoethyl)-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.028 g, 0.088 mmol), 6-bromo-9H-purine (0.035 g, 0.18 mmol), and *N,N*-diisopropylethylamine (0.030 mL, 0.18 mmol) in ethanol (0.4 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA), to give two diastereomers as a TFA salts. On an analytic HPLC (Waters SunFire C18, 2.1x50 mm, 5 μ M; injection volume 2 μ L; flow rate 3 mL/min; at gradient from 2% to 80% acetonitrile in water containing 0.15% NH_4OH in 3 min): First peak has retention time 1.421 min; LCMS calculated for $C_{20}H_{17}ClN_7OS(M+H)^+$: $m/z = 438.1$; Found: 438.0. Second peak has retention time 1.516 min; LCMS calculated for $C_{20}H_{17}ClN_7OS(M+H)^+$: $m/z = 438.1$; Found: 438.0. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.36 (1H, s), 8.32 (1H, s), 7.57 (2H, m), 7.44 (2H, m), 7.11 (1H, s), 5.04 (1H, m), 2.64 (1H, s), 1.34 (3H, d, $J = 6.8$ Hz) ppm.

Example 44. 6-(3,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0448]



[0449] A mixture of 7-(1-aminoethyl)-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.105 g, 0.327 mmol) (1st peak from Example 23, step 1 chiral separation), 6-bromo-9H-purine (0.117 g, 0.588 mmol), and *N,N*-diisopropylethylamine (0.114 mL, 0.654 mmol) in ethanol (1.5 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.15% NH₄OH) to give the desired product (0.073 g, 51%). LCMS calculated for C₂₀H₁₆F₂N₇OS(M+H)⁺: m/z = 440.1; Found: 440.0. ¹H NMR (DMSO-*d*₆, 500 MHz) δ 8.05 (2H, s), 7.34 (1H, br s), 7.18 (1H, m), 7.12 (2H, m), 6.84 (1H, s), 7.01 (1H, s), 5.07 (1H, m), 2.43 (3H, s), 1.31 (3H, d, *J* = 7.0 Hz) ppm. ¹⁹F NMR (DMSO-*d*₆, 376.3 MHz) δ -111 ppm.


Example 45. 6-(2,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0450]



Step 1. 7-(1-azidoethyl)-6-(2,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0451]



[0452] To a mixture of 7-(1-azidoethyl)-6-bromo-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.080 g, 0.25 mmol) and (2,5-difluorophenyl)boronic acid (0.048 g, 0.30 mmol) in 1,4-dioxane (2 mL) was added a 1 N solution of sodium carbonate in water (0.38 mL, 0.38 mmol) and dichloro(bis[di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl])palladium (0.011 g, 0.015 mmol). The mixture was stirred at 100 °C overnight. After cooled to room temperature, the mixture was diluted with ethyl acetate, washed with water, dried over MgSO₄, then concentrated and purified on silica gel (0-25% ethyl acetate/hexane) to give the desired product as a diastereoisomer mixture (54 mg). LCMS calculated for C₁₅H₁₂F₂N₅OS(M+H)⁺: m/z = 348.1; Found: 348.0.

Step 2. 7-(1-aminoethyl)-6-(2,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0453]



[0454] To a solution of 7-(1-aminoethyl)-6-(2,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.054 g, 0.16 mmol) in tetrahydrofuran (2 mL) was added 1.00 M of trimethylphosphine in tetrahydrofuran (0.23 mL, 0.23 mmol), and the mixture was stirred at room temperature for 1 hour. The mixture was concentrated to give the crude product (45 mg), which was used directly in the next step. LCMS calculated for $C_{15}H_{14}F_2N_3OS(M+H)^+$: $m/z = 322.1$; Found: 322.0.

Step 3. 6-(2,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0455]




[0456] A mixture of 7-(1-aminoethyl)-6-(2,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.045 g, 0.14 mmol), 6-bromo-9H-purine (0.042 g, 0.21 mmol), and *N,N*-diisopropylethylamine (0.049 mL, 0.28 mmol) in ethanol (0.3 mL) was heated at 110 °C overnight. The mixture was filtered, and the filtrate was purified on preparative-LCMS (XBridge C18 Column, eluting with a gradient of acetonitrile/water containing 0.05% TFA), to give the desired product as a mixture of two diastereomers (TFA salts). LCMS calculated for $C_{20}H_{16}F_2N_7OS(M+H)^+$: $m/z=440.1$; Found: 440.1. 1H NMR (DMSO- d_6 , 400 MHz) δ 8.64 (1H, br s), 8.38 (1H, s), 8.36 (1H, s), 7.34~7.19 (3H, m), 7.08 (1H, m), 5.06 (1H, m), 2.60 (3H, s), 1.46 (1.5H, d, $J = 6.8$ Hz), 1.33 (1.5H, d, $J = 6.8$ Hz) ppm. ^{19}F NMR (DMSO- d_6 , 376.3 MHz) δ - 117.8, -119.4, -119.8, -119.9 ppm.

Example 46. 6-(3-Fluorophenyl)-7-[(1S)-1-(3H-imidazo[4,5-b]pyridin-7-ylamino)ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0457]



[0458] A solution of 7-[(1S)-1-aminoethyl]-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (50 mg, 0.16 mmol), 7-chloro-3H-imidazo[4,5-b]pyridine (51 mg, 0.33 mmol), and *N,N*-diisopropylethylamine (57 μ L, 0.33 mmol) in 1-butanol (0.5 mL) in a sealable vial was degassed with nitrogen, sealed, and heated at 140 °C for 48 hours. The reaction mixture was diluted with methanol and purified by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.1 % ammonium hydroxide, at flow rate of 60 mL/min) to give the desired product (7 mg, 10%) as a white solid. LCMS for $C_{21}H_{18}FN_6OS(M+H)^+$: $m/z = 420.8$. 1H NMR (400 MHz, CD_3OD): δ 8.04 (s, 1 H), 7.83 (d, $J = 5.9$ Hz, 1 H), 7.55 - 7.49 (m, 1 H), 7.23 - 7.12 (m, 3 H), 6.86 (d, $J = 1.2$ Hz, 1 H), 5.96 (d, $J = 5.9$ Hz, 1 H), 4.69 - 4.67 (m, 1 H), 2.75 (s, 3 H), 1.57 (d, $J = 6.4$ Hz, 3 H).

Example 47. 6-(3-Fluorophenyl)-7-[(1S)-1-[(2-hydroxy-9H-purin-6-yl)amino]ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one**[0459]**

Step 1. 7-[(1S)-1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0460]


[0461] A solution of 7-[(1S)-1-aminoethyl]-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (0.10 g, 0.33 mmol) and 2-amino-6-bromopurine (0.11 g, 0.49 mmol) in 1-butanol (0.66 mL) was treated with *N,N*-diisopropylethylamine (86 μ L, 0.49 mmol), degassed with nitrogen for 5 min and heated at 100 °C for 18 hours. The reaction was not complete and was, therefore, heated at 115 °C for an additional 5 hours. The reaction mixture was diluted with methanol (10 mL), stirred, and filtered. The filtrate was purified by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.1% ammonium hydroxide, at flow rate of 60 mL/min) to give the desired product (27 mg, 19%) as a white solid. LCMS for $C_{20}H_{18}FN_8OS$ (M+H)⁺: m/z = 437.0.

Step 2. 6-(3-Fluorophenyl)-7-[(1S)-1-[(2-hydroxy-9H-purin-6-yl)amino]ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0462] A solution of 7-[(1S)-1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one (27 mg, 62 μ mol) in acetic acid (0.41 mL) and water (84 μ L) at 0 °C was treated with a solution of sodium nitrite (13 mg, 0.19 mmol) in water (0.15 mL) dropwise and stirred at 0 °C for 30 minutes and at 20 °C for 16 hours. The reaction mixture was concentrated and purified by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.1% ammonium hydroxide, at flow rate of 60 mL/min) to give the desired product (7 mg, 20%) as a white solid. LCMS for $C_{20}H_{17}FN_8O_2S$ (M+H)⁺: m/z = 437.8. ¹H NMR (300 MHz, DMSO-*d*₆): δ 7.79 (br s, 1 H), 7.68-7.61 (m, 1 H), 7.50 - 7.38 (m, 3 H), 7.23 - 7.14 (m, 2 H), 7.06 (br s, 1 H), 5.02 - 4.92 (m, 1 H), 2.64 (s, 3 H), 1.26 (d, *J* = 6.7 Hz, 3 H).


Example 48. 6-(3-Fluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-3-(trifluoromethyl)-5H [1,3]thiazolo[3,2-a]pyrimidin-5-one

[0463]



Step 1. 7-(1-Bromoethyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0464]



[0465] The desired compound was prepared according to the procedure of Example 8, step 2, using 4-(trifluoromethyl)-1,3-thiazol-2-amine as the starting material in 53% yield. LCMS for $C_9H_7BrF_3N_2OS$ (M+H)⁺: m/z = 326.8, 328.8.

Step 2. 6-Bromo-7-(1-bromoethyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0466]



[0467] The desired compound was prepared according to the procedure of Example 8, step 3, using 7-(1-bromoethyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as the starting material in quantitative yield. LCMS for $C_9H_6Br_2F_3N_2OS$ (M+H)⁺: m/z = 404.8, 406.7, 408.7.

Step 3. 7-(1-Azidoethyl)-6-bromo-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0468]



[0469] The desired compound was prepared according to the procedure of Example 8, step 4, using 6-bromo-7-(1-bromoethyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as the starting material in 84% yield. LCMS for $C_9H_6BrF_3N_5OS$ (M+H)⁺: m/z = 367.7, 369.8.

Step 4. 7-(1-Azidoethyl)-6-(3-fluorophenyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one


[0470]



[0471] The desired compound was prepared according to the procedure of Example 8, step 5, using 7-(1-azidoethyl)-6-bromo-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one and (3-fluorophenyl)boronic acid as the starting materials in 29% yield. LCMS for $C_{15}H_{10}F_4N_5OS$ ($M+H$)⁺: m/z = 383.9.

Step 5. 7-(1-Aminoethyl)-6-(3-fluorophenyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt

[0472]




[0473] The desired compound was prepared according to the procedure of Example 8, step 6, using 7-(1-azidoethyl)-6-(3-fluorophenyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one as the starting material in 79% yield after purification by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.05% TFA, at flow rate of 30 mL/min). LCMS for $C_{15}H_{12}F_4N_3OS$ ($M+H$)⁺: m/z = 357.9.

Step 6. 6-(3-Fluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one

[0474] The desired compound was prepared according to the procedure of Example 8, step 7, using 7-(1-aminoethyl)-6-(3-fluorophenyl)-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one trifluoroacetic acid salt as the starting material in 54% yield after purification by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.1% ammonium hydroxide, at flow rate of 30 mL/min). LCMS for $C_{20}H_{14}F_4N_7OS$ ($M+H$)⁺: m/z = 475.9. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.36 (s, 1 H), 8.14 - 8.08 (m, 2 H), 7.55 - 7.46 (m, 2 H), 7.32 - 7.21 (m, 3 H), 5.19 - 5.07 (m, 1 H), 1.37 (d, J = 7.0 Hz, 3 H).


Example 49. 6-Methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

[0475]



Step 1. 2-(1-Bromoethyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one


[0476]



[0477] The desired compound was prepared according to the procedure of Example 8, step 2, using 6-methyl-2-pyridinaminc as the starting material in 58% yield. LCMS for $C_{11}H_{12}BrN_2O$ ($M+H$)⁺: m/z = 267.0, 269.0.

Step 2. 2-(1-Bromoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one


[0478]



[0479] The desired compound was prepared according to the procedure of Example 8, step 3, using 2-(1-bromoethyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one and *N*-iodosuccinimide as the starting materials in 98% yield. LCMS for $C_{11}H_{11}BrIN_2O$ ($M+H$)⁺: m/z = 392.7, 394.7.

Step 3. 2-(1-Azidoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one


[0480]



[0481] The desired compound was prepared according to the procedure of Example 8, step 4, using 2-(1-bromoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one as the starting material in 99% yield. LCMS for $C_{11}H_{11}IN_5O$ ($M+H$)⁺: m/z = 356.0.

Step 4. 2-(1-Azidoethyl)-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one

[0482]




[0483] A solution of 2-(1-azidoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one (100 mg, 0.28 mmol) and phenylboronic acid (48 mg, 0.39 mmol) in 1,4-dioxane (2 mL) was treated with sodium carbonate (45 mg, 0.42 mmol), water (0.50 mL), and dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium (2.0 mg, 28 μ mol), degassed with nitrogen for 5 minutes, and heated at 110 °C for 18 hours. The reaction mixture was purified by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.05% TFA, at flow rate of 30 mL/min). LCMS for $C_{17}H_{16}N_5O$ ($M+H$)⁺: m/z = 306.1.

Step 5. 6-Methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

[0484] A solution of 2-(1-azidoethyl)-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one (31 mg, 0.10 mmol) in tetrahydrofuran (1 mL) and water (0.2 mL) was treated with 1 M of trimethylphosphine in tetrahydrofuran (0.20 mL, 0.20 mmol) and stirred at 20 °C for 1 hour. The reaction mixture was diluted with brine (2 mL) and extracted with dichloromethane (3 x 15 mL). The combined organic extracts were dried with sodium sulfate, filtered, and concentrated to a crude residue. This intermediate amine was used without further purification. A solution of the amine in ethanol (1 mL) was treated with 6-bromo-9H-purine (31 mg, 0.16 mmol) and *N,N*-diisopropylethylamine (24 mL, 0.14 mmol) and then heated at 90 °C for 18 hours. The reaction mixture was purified by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.05% TFA, at flow rate of 30 mL/min). LCMS for $C_{22}H_{20}N_7O$ ($M+H$)⁺: m/z = 398.1.

Example 50. 2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-6-methyl-3-phenyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0485]



[0486] The desired compound was prepared according to the procedure of Example 49 using 2-amino-6-bromopurine (instead of 6-bromo-9*H*-purine in step 5). LCMS for $C_{22}H_{21}N_8O$ ($M+H$)⁺: m/z = 413.0.

Example 51. 6-Methyl-3-(3-methylphenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0487]



[0488] The desired compound was prepared according to the procedure of Example 49 using (3-methylphenyl)boronic acid (instead of phenylboronic acid in step 4). LCMS for $C_{23}H_{22}N_7O$ ($M+H$)⁺: m/z = 411.9.

Example 52. 2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-6-methyl-3-(3-methylphenyl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0489]



[0490] The desired compound was prepared according to the procedure of Example 49 using (3-methylphenyl)boronic acid (instead of phenylboronic acid in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9*H*-purine in step 5). LCMS for $C_{23}H_{23}N_8O$ ($M+H$)⁺: m/z = 427.0.

Example 53. 3-(3-Chlorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0491]



10 **[0492]** The desired compound was prepared according to the procedure of Example 49 using (3-chlorophenyl)boronic acid (instead of phenylboronic acid in step 4). LCMS for $C_{22}H_{19}ClN_7O$ ($M+H$)⁺: m/z = 432.1.

15 **Example 54. 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3-chlorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


20 **[0493]**



30 **[0494]** The desired compound was prepared according to the procedure of Example 49 using (3-chlorophenyl)boronic acid (instead of phenylboronic acid in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{22}H_{20}ClN_8O$ ($M+H$)⁺: m/z = 447.1.

35 **Example 55. 3-(4-Chlorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


40 **[0495]**



50 **[0496]** The desired compound was prepared according to the procedure of Example 49 using (4-chlorophenyl)boronic acid (instead of phenylboronic acid in step 4). LCMS for $C_{22}H_{19}ClN_7O$ ($M+H$)⁺: m/z = 432.1.

55 **Example 56. 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(4-chlorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0497]



10 **[0498]** The desired compound was prepared according to the procedure of Example 49 using (4-chlorophenyl)boronic acid (instead of phenylboronic acid in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{22}H_{20}ClN_8O$ ($M+H$)⁺: m/z = 447.1.

15 **Example 57. 3-(2-Chlorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0499]



25 **[0500]** The desired compound was prepared according to the procedure of Example 49 using (2-chlorophenyl)boronic acid (instead of phenylboronic acid in step 4) as a mixture of atropisomers. LCMS for $C_{22}H_{19}ClN_7O$ ($M+H$)⁺: m/z = 432.1.

30 **Example 58. 2-[1-[(2-Amino-9H-purin-6-yl)amino]ethyl]-3-(2-chlorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0501]



40 **[0502]** The desired compound was prepared according to the procedure of Example 49 using (2-chlorophenyl)boronic acid (instead of phenylboronic acid in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5) as a mixture of atropisomers. LCMS for $C_{22}H_{20}ClN_8O$ ($M+H$)⁺: m/z = 447.1.

45 **Example 59. 3-(2-Fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0503]



10 **[0504]** The desired compound was prepared according to the procedure of Example 49 using (2-fluorophenyl)boronic acid (instead of phenylboronic acid in step 4). LCMS for $C_{22}H_{19}FN_7O$ ($M+H$)⁺: m/z = 416.1.

15 **Example 60. 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(2-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


20 **[0505]**



30 **[0506]** The desired compound was prepared according to the procedure of Example 49 using (2-fluorophenyl)boronic acid (instead of phenylboronic acid in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{22}H_{20}FN_8O$ ($M+H$)⁺: m/z = 431.1.

35 **Example 61. 4-{6-Methyl-4-oxo-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-3-yl}benzonitrile trifluoroacetic acid salt**


40 **[0507]**



50 **[0508]** The desired compound was prepared according to the procedure of Example 49 using (4-cyanophenyl)boronic acid (instead of phenylboronic acid in step 4). LCMS for $C_{23}H_{19}N_8O$ ($M+H$)⁺: m/z = 423.1.

55 **Example 62. 4-(2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-6-methyl-4-oxo-4H-pyrido[1,2-a]pyrimidin-3-yl)benzonitrile trifluoroacetic acid salt**


[0509]



10 **[0510]** The desired compound was prepared according to the procedure of Example 49 using (4-cyanophenyl)boronic acid (instead of phenylboronic acid in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{23}H_{20}N_9O$ ($M+H$)⁺: m/z = 438.2.

15 **Example 63. 6-Methyl-3-(2-methylphenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0511]



25 **[0512]** The desired compound was prepared according to the procedure of Example 49 using (2-methylphenyl)boronic acid (instead of phenylboronic acid in step 4), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (instead of dichloro(bis{di-*tert*-butyl}[4-(dimethylamino)phenyl]phosphoranyl)palladium in step 4), and potassium carbonate (instead of sodium carbonate in step 4). LCMS for $C_{23}H_{22}N_7O$ ($M+H$)⁺: m/z = 412.1.

30 **Example 64. 6-Methyl-3-(4-methylphenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


35 **[0513]**



45 **[0514]** The desired compound was prepared according to the procedure of Example 49 using (4-methylphenyl)boronic acid (instead of phenylboronic acid in step 4), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (instead of dichloro(bis{di-*tert*-butyl}[4-(dimethylamino)phenyl]phosphoranyl)palladium in step 4), and potassium carbonate (instead of sodium carbonate in step 4). LCMS for $C_{23}H_{22}N_7O$ ($M+H$)⁺: m/z = 412.1.

50 **Example 65. 3-(3-Methoxyphenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


55 **[0515]**



10 **[0516]** The desired compound was prepared according to the procedure of Example 49 using 3-methoxyphenylboronic acid (instead of phenylboronic acid in step 4), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4), and potassium carbonate (instead of sodium carbonate in step 4). LCMS for $C_{23}H_{22}N_7O_2$ ($M+H$)⁺: m/z = 428.1.

15 **Example 66. 3-(2,3-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0517]



30 **[0518]** The desired compound was prepared according to the procedure of Example 49 using (2,3-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4), and potassium carbonate (instead of sodium carbonate in step 4). LCMS for $C_{22}H_{18}F_2N_7O$ ($M+H$)⁺: m/z = 434.2.

35 **Example 67. 3-(2,5-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0519]



50 **[0520]** The desired compound was prepared according to the procedure of Example 49 using (2,5-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4), and potassium carbonate (instead of sodium carbonate in step 4). LCMS for $C_{22}H_{18}F_2N_7O$ ($M+H$)⁺: m/z = 434.1.

55 **Example 68. 3-(3,4-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**


[0521]



10 **[0522]** The desired compound was prepared according to the procedure of Example 49 using (3,4-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane (1:1) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4), and potassium carbonate (instead of sodium carbonate in step 4). LCMS for $C_{22}H_{18}F_2N_7O$ ($M+H$)⁺: m/z = 434.0.

15 **Example 69. 3-(3,5-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt**

[0523]




25

30 **[0524]** The desired compound was prepared according to the procedure of Example 49 using (3,5-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4) and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{22}H_{18}F_2N_7O$ ($M+H$)⁺: m/z = 434.0. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.80 (br s, 1 H), 8.48 (s, 2 H), 7.71 (dd, J =7.9, 7.6 Hz, 1 H), 7.45 (d, J = 8.2 Hz, 1 H), 7.24 (d, J = 9.7, 9.1 Hz, 1 H), 7.18 - 7.11 (m, 3 H), 6.97 (d, J =6.7 Hz, 1 H), 5.29 - 5.20 (m, 1 H), 2.89 (s, 3 H), 1.46 (d, J =6.7 Hz, 3 H).

35

Example 70. 3-(3-Fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

40 **[0525]**




50 **[0526]** The desired compound was prepared according to the procedure of Example 49 using (3-fluorophenyl)boronic acid (instead of phenylboronic acid in step 4) and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{22}H_{19}FN_7O$ ($M+H$)⁺: m/z = 416.1. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.85 (br s, 1 H), 8.50 (s, 2 H), 7.70 (dd, J = 7.9, 7.6 Hz, 1 H), 7.51 - 7.40 (m, 2 H), 7.27 - 7.16 (m, 3 H), 6.96 (d, J = 6.7 Hz, 1 H), 5.31 - 5.20 (m, 1 H), 2.88 (s, 3 H), 1.44 (d, J = 6.7 Hz, 3 H).

55


Example 71 and Example 72. Single enantiomers of 3-(3-Fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

[0527]



Step 1. Chiral separation of 2-(1-Azidoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one


[0528]



The racemic mixture of 2-(1-azidoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one was separated by HPLC (Chiracel OJ-H, eluting with 30% ethanol/70% hexanes, at flow rate of 20 mL/min) to give the two individual enantiomers (retention times = 21.6 min, 27.2 min). Both peaks were advanced to the next step.

Step 2. Single enantiomers of 2-(1-azidoethyl)-3-(3-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

[0530]




The desired compounds were prepared according to the procedure of Example 49, step 4, using peak 1 and peak 2 of 2-(1-azidoethyl)-3-iodo-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one and (3-fluorophenyl)boronic acid as the starting materials after purification by RP-HPLC (XBridge C18 column, eluting with a gradient of acetonitrile/water containing 0.05% TFA, at flow rate of 60 mL/min). From peak 1: LCMS for $C_{17}H_{15}FN_5O$ ($M+H$)⁺: m/z = 324.1. From peak 2: LCMS for $C_{17}H_{15}FN_5O$ ($M+H$)⁺: m/z = 323.9.

Step 3. Single enantiomers of 3-(3-Fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

The desired compounds were prepared according to the procedure of Example 49, step 5, using the single enantiomers of 2-(1-azidoethyl)-3-(3-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt and (3-fluorophenyl)boronic acid as the starting materials. Example 71 (from peak 1): LCMS for $C_{22}H_{19}FN_7O$ ($M+H$)⁺: m/z = 415.9; ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.78 (br s, 1 H), 8.48 (s, 2 H), 7.70 (dd, *J*=7.8, 7.7 Hz, 1 H), 7.50 - 7.41 (m, 2 H), 7.28 - 7.17 (m, 3 H), 6.96 (d, *J*=7.0 Hz, 1 H), 5.30 - 5.21 (m, 1 H), 2.88 (s, 3 H), 1.44 (d, *J*=6.7 Hz, 3 H). Example 72 (from peak 2): LCMS for $C_{22}H_{19}FN_7O$ ($M+H$)⁺: m/z = 416.1; ¹H NMR (400 MHz, DMSO-*d*₆): δ 8.78 (br s, 1 H), 8.48 (s, 2 H), 7.70 (dd, *J*=8.1, 7.5 Hz, 1 H), 7.50 - 7.42 (m, 2 H), 7.27 - 7.18 (m, 3 H), 6.96 (d, *J*=6.8 Hz, 1 H), 5.30 - 5.21 (m, 1 H), 2.88 (s, 3 H), 1.44 (d, *J*=6.7 Hz, 3 H).

Example 73. 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt


[0533]



[0534] The desired compound was prepared according to the procedure of Example 49 using (3-fluorophenyl)boronic acid (instead of phenylboronic acid in step 4), tetrakis(triphenylphosphine)palladium(0) (instead of dichloro[bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl}]palladium in step 4), and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{22}H_{20}FN_8O$ (N_4+H)⁺: $m/z = 431.2$. 1H NMR (300 MHz, DMSO- d_6): δ 8.72 (d, $J=7.3$ Hz, 1 H), 8.17 (s, 1 H), 7.74 - 7.68 (m, 1 H), 7.50 - 7.41 (m, 2 H), 7.29 - 7.14 (m, 5 H), 6.98 (d, $J=6.4$ Hz, 1 H), 5.26 - 5.17 (m, 1 H), 2.89 (s, 3 H), 1.37 (d, $J=6.7$ Hz, 3 H).

Example 74 and Example 75. Single enantiomers of 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt


[0535]



[0536] The desired compounds were prepared according to the procedure of Example 71 and 72. 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). Example 74 (from peak 1): $C_{22}H_{20}FN_8O$ ($M+H$)⁺: $m/z = 431.0$; 1H NMR (400 MHz, DMSO- d_6): δ 8.72 (d, $J=7.3$ Hz, 1 H), 8.17 (s, 1 H), 7.72 (dd, $J=8.8, 7.1$ Hz, 1 H), 7.48 - 7.40 (m, 2 H), 7.27 - 7.13 (m, 5 H), 6.98 (d, $J=6.8$ Hz, 1 H), 5.25 - 5.18 (m, 1 H), 2.89 (s, 3 H), 1.37 (d, $J=6.7$ Hz, 3 H). Example 75 (from peak 2): $C_{22}H_{20}FN_8O$ ($M+H$)⁺: $m/z = 431.1$; 1H NMR (400 MHz, DMSO- d_6): δ 8.72 (d, $J=7.1$ Hz, 1 H), 8.17 (s, 1 H), 7.71 (dd, $J=8.9, 7.1$ Hz, 1 H), 7.49 - 7.42 (m, 2 H), 7.28 - 7.15 (m, 5 H), 6.98 (d, $J=6.8$ Hz, 1 H), 5.25 - 5.18 (m, 1 H), 2.89 (s, 3 H), 1.37 (d, $J=6.8$ Hz, 3 H).

Example 76. 3-(3,5-Difluorophenyl)-6-ethyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt


[0537]



[0538] The desired compound was prepared according to the procedure of Example 49 using 6-ethylpyridin-2-amine (instead of 6-methyl-2-pyridinamine in step 1), *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (3,5-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{23}H_{20}F_2N_7O$ ($M+H$)⁺: m/z = 448.2. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.57 - 8.44 (m, 1 H), 8.40 (s, 2 H), 7.75 (dd, J =8.2, 7.9 Hz, 1 H), 7.48 (d, J =8.8 Hz, 1 H), 7.29 - 7.12 (m, 3 H), 7.04 (d, J =6.7 Hz, 1 H), 5.30 - 5.17 (m, 1 H), 3.33 (q, J =7.0 Hz, 2 H), 1.45 (d, J =6.7 Hz, 3 H), 1.14 (t, J =7.0 Hz, 3 H).

Example 77. 2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(3,5-difluorophenyl)-6-ethyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0539]



[0540] The desired compound was prepared according to the procedure of Example 49 using 6-ethylpyridin-2-amine (instead of 6-methyl-2-pyridinamine in step 1), *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (3,5-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9*H*-purine in step 5). LCMS for $C_{23}H_{21}F_2N_8O$ ($M+H$)⁺: m/z =463.2. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.74 - 8.69 (m, 1 H), 8.17 (s, 1 H), 7.78 (dd, J = 8.8, 7.0 Hz, 1 H), 7.50 (d, J =8.5 Hz, 1 H), 7.32 - 7.16 (m, 3 H), 7.14 - 7.04 (m, 2 H), 5.26 - 5.16 (m, 1 H), 3.35 (q, J = 7.3 Hz, 2 H), 1.41 (d, J = 6.7 Hz, 3 H), 1.15 (t, J =7.3 Hz, 3 H).

Example 78. 6-Ethyl-3-(4-fluorophenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0541]



[0542] The desired compound was prepared according to the procedure of Example 49 using 6-ethylpyridin-2-amine (instead of 6-methyl-2-pyridinamine in step 1), *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (4-fluorophenyl)boronic acid (instead of phenylboronic acid in step 4), and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{23}H_{21}FN_7O$ ($M+H$)⁺: m/z = 430.2.

Example 79. 3-(3,5-Difluorophenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one trifluoroacetic acid salt


[0543]



[0544] The desired compound was prepared according to the procedure of Example 49 using 2-pyridinamine (instead of 6-methyl-2-pyridinamine in step 1), *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (3,5-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{21}H_{16}F_2N_7O$ ($M+H$)⁺: m/z = 420.0. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.96 (d, J =7.3 Hz, 1 H), 8.36 (s, 2 H), 8.01 (dd, J =8.2, 7.3 Hz, 1 H), 7.74 (d, J =8.8 Hz, 1 H), 7.40 (dd, J =6.7, 6.4 Hz, 1 H), 7.32 - 7.16 (m, 3 H), 5.37 - 5.26 (m, 1 H), 1.46 (d, J = 6.7 Hz, 3 H).

Example 80. 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3,5-difluorophenyl)-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt


[0545]



[0546] The desired compound was prepared according to the procedure of Example 49 using 2-pyridinamine (instead of 6-methyl-2-pyridinamine in step 1), *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (3,5-difluorophenyl)boronic acid (instead of phenylboronic acid in step 4), tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4) and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{21}H_{17}F_2N_8O$ ($M+H$)⁺: m/z = 435.0. ¹H NMR (300 MHz, DMSO-*d*₆): δ 8.9 (d, J =7.0 Hz, 1 H), 8.82 - 8.72 (br s, 1 H), 8.17 (s, 1 H), 8.06 - 8.00 (m, 1 H), 7.76 (d, J = 9.1 Hz, 1 H), 7.43 (dd, J = 7.0, 5.6 Hz, 1 H), 7.30 - 7.07 (m, 5 H), 5.32 - 5.22 (m, 1 H), 1.42 (d, J = 6.7 Hz, 3 H).

Example 81. 3-(6-Chloro-5-methylpyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt

[0547]




[0548] The desired compound was prepared according to the procedure of Example 49 using *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (6-chloro-5-methylpyridin-3-yl)boronic acid (instead of phenylboronic acid in step 4), and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{22}H_{20}ClN_8O$ ($M+H$)⁺: m/z = 446.9. ¹H NMR (300 MHz, DMSO-*d*₆):

δ 8.37 (br s, 2 H), 8.26 (s, 1 H), 7.82 (s, 1 H), 7.71 (dd, J = 9.1, 7.0 Hz, 1 H), 7.45 (d, J = 8.8 Hz, 1 H), 6.97 (d, J = 6.2 Hz, 1 H), 5.21 - 5.10 (m, 1 H), 2.88 (s, 3 H), 2.33 (s, 3 H), 1.45 (d, J = 6.7 Hz, 3 H).

Example 82. 2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(6-chloro-5-methylpyridin-3-yl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one trifluoroacetic acid salt


[0549]



[0550] The desired compound was prepared according to the procedure of Example 49 using *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (6-chloro-5-methylpyridin-3-yl)boronic acid (instead of phenylboronic acid in step 4), tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4), and 2-amino-6-bromopurine (instead of 6-bromo-9H-purine in step 5). LCMS for $C_{22}H_{21}ClN_9O$ ($M+H$)⁺: m/z = 462.0. 1H NMR (500 MHz, DMSO- d_6): δ 8.74 (br s, 1 H), 8.24 (s, 1 H), 8.18 (s, 1 H), 7.79 - 7.71 (m, 2 H), 7.50 (d, J = 8.8 Hz, 1 H), 7.29 - 7.14 (m, 2 H), 7.03 (d, J = 6.9 Hz, 1 H), 5.23 - 5.16 (m, 1 H), 2.92 (s, 3 H), 2.31 (s, 3 H), 1.43 (d, J = 6.7 Hz, 3 H).

Example 83. 3-{6-Methyl-4-oxo-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-3-yl}benzonitrile trifluoroacetic acid salt


[0551]



[0552] The desired compound was prepared according to the procedure of Example 49 using *N*-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (3-cyanophenyl)boronic acid (instead of phenylboronic acid in step 4), and tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4). LCMS for $C_{23}H_{19}N_8O$ ($M+H$)⁺: m/z = 422.9. 1H NMR (500 MHz, DMSO- d_6): δ 8.46 (br s, 1 H), 8.40 (s, 2 H), 7.91 - 7.83 (m, 2 H), 7.81 - 7.76 (m, 1 H), 7.75 - 7.69 (m, 1 H), 7.69 - 7.63 (m, 1 H), 7.48 (d, J = 8.7 Hz, 1 H), 6.98 (d, J = 6.7 Hz, 1 H), 5.24 - 5.15 (br s, 1 H), 2.91 (s, 3 H), 1.46 (d, J = 6.6 Hz, 3 H).

Example 84. 3-(2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-6-methyl-4-oxo-4H-pyrido[1,2-a]pyrimidin-3-yl)benzonitrile trifluoroacetic acid salt

[0553]



[0554] The desired compound was prepared according to the procedure of Example 49 using N-bromosuccinimide (instead of *N*-iodosuccinimide in step 2), (3-cyanophenyl)boronic acid (instead of phenylboronic acid in step 4), tetrakis(triphenylphosphine)palladium(0) (instead of dichloro(bis{di-*tert*-butyl[4-(dimethylamino)phenyl]phosphoranyl})palladium in step 4), and 2-amino-6-bromopurine (instead of 6-bromo-9*H*-purine in step 5). LCMS for $C_{23}H_{20}N_9O$ ($M+H$)⁺: m/z = 438.0. ¹H NMR (500 MHz, DMSO-*d*₆): δ 8.73 (br s, 1 H), 8.18 (s, 1 H), 7.84-7.80 (m, 2 H), 7.78 - 7.71 (m, 2 H), 7.64 (dd, *J* = 8.2, 8.0 Hz, 1 H), 7.49 (d, *J* = 8.7 Hz, 1 H), 7.24 (br s, 2 H), 7.01 (d, *J* = 6.9 Hz, 1 H), 5.21 - 5.15 (m, 1 H), 2.92 (s, 3 H), 1.40 (d, *J* = 6.7 Hz, 3 H).

Example A1: PI3K Enzyme Assay

[0555] PI3-Kinase luminescent assay kit including lipid kinase substrate, D-myo-phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂)D (+)-sn-1,2-di-O-octanoylglycerol, 3-O-phospho linked (PIP₂), biotinylated I(1,3,4,5)P₄, PI(3,4,5)P₃ Detector Protein, was purchased from Echelon Biosciences (Salt Lake City, UT). AlphaScreen™ GST Detection Kit including donor and acceptor beads was purchased from PerkinElmer Life Sciences (Waltham, MA). PI3Kδ (p110δ /p85α) was purchased from Millipore (Bedford, MA). ATP, MgCl₂, DTT, EDTA, HEPES and CHAPS were purchased from Sigma-Aldrich (St. Louis, MO).

AlphaScreen™ Assay for PI3Kδ

[0556] The kinase reaction was conducted in 384-well REMP plate from Thermo Fisher Scientific in a final volume of 40 μL. Inhibitors were first diluted serially in DMSO and added to the plate wells before the addition of other reaction components. The final concentration of DMSO in the assay was 2%. The PI3K assays were carried out at room temperature in 50 mM HEPES, pH 7.4, 5mM MgCl₂, 50 mM NaCl, 5mM DTT and CHAPS 0.04%. Reactions were initiated by the addition of ATP, the final reaction mixture consisted of 20 μM PIP₂, 20 μM ATP, 1.2nM PI3Kδ were incubated for 20 min. 10 μL of reaction mixture was then transferred to 5 μL 50nM biotinylated I(1,3,4,5)P₄ in quench buffer: 50 mM HEPES pH 7.4, 150 mM NaCl, 10 mM EDTA, 5 mM DTT, 0.1% Tween-20, followed with the addition of 10 μL AlphaScreen™ donor and acceptor beads suspended in quench buffer containing 25nM PI(3,4,5)P₃ detector protein. The final concentration of both donor and acceptor beads is 20 mg/ml. After plate sealing, the plate was incubated in a dark location at room temperature for 2 hours. The activity of the product was determined on Fusion-alpha microplate reader (Perkin-Elmer). IC₅₀ determination was performed by fitting the curve of percent control activity versus the log of the inhibitor concentration using the GraphPad Prism 3.0 software.

Example A2: PI3K Enzyme Assay

[0557] Materials: Lipid kinase substrate, phosphoinositol-4,5-bisphosphate (PIP₂), was purchased from Echelon Biosciences (Salt Lake City, UT). PI3K isoforms α, β, δ and γ were purchased from Millipore (Bedford, MA). ATP, MgCl₂, DTT, EDTA, MOPS and CHAPS were purchased from Sigma-Aldrich (St. Louis, MO).

[0558] The kinase reaction was conducted in clear-bottom 96-well plate from Thermo Fisher Scientific in a final volume of 24 μL. Inhibitors were first diluted serially in DMSO and added to the plate wells before the addition of other reaction components. The final concentration of DMSO in the assay was 0.5%. The PI3K assays were carried out at room temperature in 20 mM MOPS, pH 6.7, 10 mM MgCl₂, 5 mM DTT and CHAPS 0.03%. The reaction mixture was prepared containing 50 μM PIP₂, kinase and varying concentration of inhibitors. Reactions were initiated by the addition of ATP containing 2.2 μCi [³³P]ATP to a final concentration of 1000 μM. The final concentration of PI3K isoforms α, β, δ and γ in the assay were 1.3, 9.4, 2.9 and 10.8 nM respectively. Reactions were incubated for 180 min and terminated by the addition of 100 μL of 1 M potassium phosphate pH 8.0, 30 mM EDTA quench buffer. A 100 μL aliquot of the reaction solution was then transferred to 96-well Millipore MultiScreen IP 0.45 μm PVDF filter plate (The filter plate was prewetted with 200 μL 100% ethanol, distilled water, and 1 M potassium phosphate pH 8.0, respectively). The filter plate was aspirated on a Millipore Manifold under vacuum and washed with 18 × 200 μL wash buffer containing 1 M potassium phosphate pH 8.0 and 1 mM ATP. After drying by aspiration and blotting, the plate was air dried in an incubator at 37

°C overnight. Packard TopCount adapter (Millipore) was then attached to the plate followed with addition of 120 μ L Microscint 20 scintillation cocktail (Perkin Elmer) in each well. After the plate sealing, the radioactivity of the product was determined by scintillation counting on Topcount (Perkin-Elmer). IC₅₀ determination was performed by fitting the curve of percent control activity versus the log of the inhibitor concentration using the GraphPad Prism 3.0 software. Compounds having and IC₅₀ value of 10 μ M or less are considered active. See Table 1 for data related to compounds of the invention.

Example A3: PI3K δ scintillation proximity assay

Materials

[0559] [γ -³³P]ATP (10mCi/mL) was purchased from Perkin-Elmer (Waltham, MA). Lipid kinase substrate, D-myo-Phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂)D (+)-sn-1,2-di-O-octanoylglycerol, 3-O-phospho linked (PIP₂), CAS 204858-53-7, was purchased from Echelon Biosciences (Salt Lake City, UT). PI3K δ (p110 δ /p85 α) was purchased from Millipore (Bedford, MA). ATP, MgCl₂, DTT, EDTA, MOPS and CHAPS were purchased from Sigma-Aldrich (St. Louis, MO). Wheat Germ Agglutinin (WGA) YSi SPA Scintillation Beads was purchased from GE healthcare life sciences (Piscataway, NJ).

[0560] The kinase reaction was conducted in polystyrene 384-well matrix white plate from Thermo Fisher Scientific in a final volume of 25 μ L. Inhibitors were first diluted serially in DMSO and added to the plate wells before the addition of other reaction components. The final concentration of DMSO in the assay was 0.5%. The PI3K assays were carried out at room temperature in 20 mM MOPS, pH 6.7, 10 mM MgCl₂, 5 mM DTT and CHAPS 0.03%. Reactions were initiated by the addition of ATP, the final reaction mixture consisted of 20 μ M PIP₂, 20 μ M ATP, 0.2 μ Ci [γ -³³P] ATP, 4 nM PI3K δ . Reactions were incubated for 210 min and terminated by the addition of 40 μ L SPA beads suspended in quench buffer: 150mM potassium phosphate pH 8.0, 20% glycerol, 25 mM EDTA, 400 μ M ATP. The final concentration of SPA beads was 1.0mg/mL. After the plate sealing, plates were shaken overnight at room temperature and centrifuged at 1800 rpm for 10 minutes, the radioactivity of the product was determined by scintillation counting on Topcount (Perkin-Elmer). IC₅₀ determination was performed by fitting the curve of percent control activity versus the log of the inhibitor concentration using the GraphPad Prism 3.0 software.

Table 1. IC₅₀ data for PI3K δ enzyme assays A1, A2, or A3*

Example	PI3K δ^a IC ₅₀ (nM)	PI3K δ^b IC ₅₀ (nM)	PI3K α^c IC ₅₀ (nM)	PI3K β^c IC ₅₀ (nM)	PI3K γ^c IC ₅₀ (nM)
1	+++	+++			
2	++	+			
3	+++	+++			
4	+++	++			
5	+++	++			
6	+++	+++			
7	++	++			
8	++	+			
9	+	+	++++	++++	++
10	+	+	++++	++++	++++
11	+	+	++++	++++	+++
12	+	+			
13	+	+	++++	++++	+++
14	++	++			
15	+	+	++++	++++	++++
17		+++			
18		+++			
19		+++			
20		+			

EP 2 845 856 A1

(continued)

	Example	PI3K δ^a IC ₅₀ (nM)	PI3K δ^b IC ₅₀ (nM)	PI3K α^c IC ₅₀ (nM)	PI3K β^c IC ₅₀ (nM)	PI3K γ^c IC ₅₀ (nM)
5	21		+			
	22		++			
	23		+			++++
	24		++++			
10	25		++++			
	26		+++			
	27		+++			
	28		++++			
15	29		++++			
	30	+	+	++++	++++	++++
	31		+++			
	32		++			
20	33		+			
	34		+			
	35		++			
	36		+++			
25	37		+++			
	38		++			
	39		1 st peak: +++ 2 nd peak: +			
	40		+			
30	41		++			
	42		+++			
	43		1 st peak: +++ 2 nd peak: +			
	44	+	+	++++	++++	++++
35	45		+			
	46		+			
	47		+++			
	48		++			
40	49		+			
	50		+			
	51		+++			
	52		+			
45	53		+++			
	54		+			
	55		++++			
	56		++++			

EP 2 845 856 A1

(continued)

Example	PI3K δ^a IC ₅₀ (nM)	PI3K δ^b IC ₅₀ (nM)	PI3K α^c IC ₅₀ (nM)	PI3K β^c IC ₅₀ (nM)	PI3K γ^c IC ₅₀ (nM)
57		+			
58		+			
59		+			
60		+			
61		++++			
62		+++			
63		++			
64		+			
65		+			
66		+			
67		+			
68		+++			
69		+++			
70		++			
71		+			++++
72		++++			
73		+			
74		+			
75		++++			
76		+++			
77		++			
78		+			
79		++++			
80		+++			
81		++++			
82		++++			
83		+++			
84		++			
* "+" = <50 nM; "++" = 50-100 nM; "+++" = 100-500 nM; "++++" =>500 nM. a. Results in this column were obtained by Assay A1, except for Examples 15, 30 and 44 which used Assay A2. b. Results in this column were obtained by Assay A3. c. Results in this column were obtained by Assay A2.					

Example B1: B cell proliferation assay

[0561] To acquire B cells, human PBMC were isolated from the peripheral blood of normal, drug free donors by standard density gradient centrifugation on Ficoll-Hypaque (GE Healthcare, Piscataway, NJ) and incubated with anti-CD19 microbeads (Miltenyi Biotech, Auburn, CA). The B cells were then purified by positive immunosorting using an autoMacs (Miltenyi Biotech) according to the manufacturer's instruction.

[0562] The purified B cells (2x 10⁵/well/200 μ L) were cultured in 96-well ultra-low binding plates (Corning, Corning, NY) in RPMI1640, 10% FBS and goat F(ab')₂ anti-human IgM (10 μ g/ml) (Invitrogen, Carlsbad, CA), in the presence of different amount of test compounds, for three days. [³H]-thymidine (1 μ Ci/well) (PerkinElmer, Boston, MA) in PBS

was then added to the B cell cultures for an additional 12 hrs before the incorporated radioactivity was separated by filtration with water through GF/B filters (Packard Bioscience, Meriden, CT) and measured by liquid scintillation counting with a TopCount (Packard Bioscience). Compounds having an IC_{50} value of 10 μ M or less are considered active. See Table 2 for data related to compounds of the invention.

Table 2. IC_{50} data for B cell proliferation assay*

Example	B cell IC_{50} (nM)
1	+
2	+++
3	+
4	+
5	+
6	++
7	++++
8	+
9	+
10	+
11	+
12	+
13	+
15	+
30	+
44	+
71	+
* "+" = <50 nM; "++" = 50-100 nM; "+++" = 100-500 nM; "++++" = >500 nM.	

Example B2: Pfeiffer cell proliferation assay

[0563] Pfeiffer cell line (diffuse large B cell lymphoma) was purchased from ATCC (Manassas, VA) and maintained in the culture medium recommended (RPMI and 10% FBS). To measure the anti-proliferation activity of the PI3K δ submittals, the Pfeiffer cells were plated with the culture medium (2×10^3 cells / well/ per 200 μ l) into 96-well ultra-low binding plates (Corning, Corning, NY), in the presence or absence of a concentration range of test compounds. After 3-4 days, [3 H]-thymidine (1 μ Ci/well) (PerkinElmer, Boston, MA) in PBS was then added to the cell culture for an additional 12 hrs before the incorporated radioactivity was separated by filtration with water through GF/B filters (Packard Bioscience, Meriden, CT) and measured by liquid scintillation counting with a TopCount (Packard Bioscience). See Table 3 for data related to compounds of the invention.

Table 3. IC_{50} data for Pfeiffer cell proliferation assay*

Example	IC_{50} (nM)
1	+++
2	+
3	+++
4	+++
5	++
6	+++

EP 2 845 856 A1

(continued)

	Example	IC ₅₀ (nM)
5	7	++++
	8	+
	9	+
	10	+
10	11	+
	12	+++
	13	+
	14	+++++
15	15	+
	17	+++
	18	++
	19	+++++
20	20	++
	21	++
	22	+++
	23	+
25	24	+++
	25	++++
	27	+++
	30	+
30	31	+++
	32	+
	33	+
	34	+
35	35	+
	36	++
	37	++
	38	++
40	39	First peak: +++ Second peak: +
	40	++
	41	++
	42	+++
45	43	First peak: +++ Second peak: +
	44	+
	45	++
	46	+

EP 2 845 856 A1

(continued)

	Example	IC ₅₀ (nM)
5	48	++
	49	++
	50	+
	51	+++
10	52	++
	53	++
	54	++
15	57	++
	58	+
	59	+++
	60	++
20	62	++++
	63	+++
	64	++
25	65	++
	66	+++
	67	++
	68	++++
30	69	++
	70	+
	71	+
35	72	++++
	73	+
	74	+
	75	++++
40	76	+++
	77	+
	78	+++
45	80	+++
	83	+++
	84	+++
50	* "+" = <50 nM; "++" = 50-100 nM; "+++" = 100-500 nM; "++++" = 500-1000 nM; "+++++" =>1000nM.	

Example C: Akt phosphorylation assay

[0564] Ramos cells (B lymphocyte from Burkitts lymphoma) were obtained from ATCC (Manassas, VA) and maintained in RPMI1640 and 10% FBS. The cells (3×10^7 cells /tube/3 mL in RPMI) were incubated with different amounts of test compounds for 2 hrs at 37 °C and then stimulated with goat F(ab')₂ anti-human IgM (5 µg/mL) (Invitrogen) for 17 min. in a 37 °C water bath. The stimulated cells were spun down at 4 °C with centrifugation and whole cell extracts prepared


using 300 μ L lysis buffer (Cell Signaling Technology, Danvers, MA). The resulting lysates were sonicated and supernatants were collected. The phosphorylation level of Akt in the supernatants were analyzed by using PathScan phospho-Akt (Ser473) sandwich ELISA kits (Cell Signaling Technology) according to the manufacturer's instruction.

[0565] Various modifications of the invention, in addition to those described herein, will be apparent to those skilled in the art from the foregoing description. Such modifications are also intended to fall within the scope of the appended claims. Each reference, including all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety.

Embodiments

[0566]

1. A compound of Formula I or II:



or a pharmaceutically acceptable salt thereof, wherein:

A is C_{1-10} alkyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO_2 , OR^a , SR^a , $C(O)R^b$, $C(O)NR^cR^d$, $C(O)OR^a$, $OC(O)R^b$, $OC(O)NR^cR^d$, NR^cR^d , $NR^cC(O)R^b$, $NR^cC(O)OR^3$, $NR^cC(O)NR^cR^d$, $C(=NR^e)R^b$, $C(=NR^e)NR^cR^d$, $NR^cC(=NR^e)NR^cR^d$, $NR^cS(O)R^b$, $NR^cS(O)_2R^b$, $NR^cS(O)_2NR^cR^d$, $S(O)R^b$, $S(O)NR^cR^d$, $S(O)_2R^b$, and $S(O)_2NR^cR^d$; wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^a , SR^3 , $C(O)R^b$, $C(O)NR^cR^d$, $C(O)OR^a$, $OC(O)R^b$, $OC(O)NR^cR^d$, $C(=NR^e)NR^cR^d$, $NR^cC(=NR^e)NR^cR^d$, NR^cR^d , $NR^cC(O)R^b$, $NR^cC(O)OR^a$, $NR^cC(O)NR^cR^d$, $NR^cS(O)R^b$, $NR^cS(O)_2R^b$, $NR^cS(O)_2NR^cR^d$, $S(O)R^b$, $S(O)NR^cR^d$, $S(O)_2R^b$, and $S(O)_2NR^cR^d$;

L is absent, $(CR^{7a}R^{7b})_m$, $(CR^{7a}R^{7b})_pO(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pS(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pS(O)(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pS(O)_2(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pNR^{7c}(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pNR^{7c}C(O)NR^{7c}(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pNR^{7c}C(O)O(CR^{7a}R^{7b})_q$, $(CR^{7a}R^{7b})_pNR^{7c}(=NR^{7d})NR^{7c}(CR^{7a}R^{7b})_q$, or $(CR^{7a}R^{7b})_pNR^{7c}S(O)_2(CR^{7a}R^{7b})_q$;

W is N or CR^3 ;

X is N or CR^4 ;

Y is N or CR^5 ;

Z is N or CR^6 ;

G is O, S, or NR^N ;

R^1 is OR^A , SR^A , $S(O)R^A$, $S(O)_2R^A$, NR^AR^B , $NR^cC(O)NR^AR^B$, $NR^cC(O)OR^A$, $NR^cC(=NR^E)NR^AR^B$, $NR^cS(O)_2R^A$, $NR^cS(O)_2NR^cR^A$, heterocycloalkyl, or heteroaryl, wherein said heterocycloalkyl or heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from $-(C_{1-4}alkyl)_r-Cy^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{a1}$;

R^{2a} and R^{2b} are independently selected from H, halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $C(=NR^e)NR^{c2}R^{d2}$,

$$\text{NR}^{\text{c}2}\text{C}(\text{=NR}^{\text{e}})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}, \text{NR}^{\text{c}2}\text{R}^{\text{d}2}, \text{NR}^{\text{c}2}\text{C}(\text{O})\text{R}^{\text{b}2}, \text{NR}^{\text{c}2}\text{C}(\text{O})\text{OR}^{\text{a}2}, \text{NR}^{\text{c}2}\text{C}(\text{O})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}, \text{NR}^{\text{c}2}\text{S}(\text{O})\text{R}^{\text{b}2}, \text{NR}^{\text{c}2}\text{S}(\text{O})_2\text{R}^{\text{b}2}, \text{NR}^{\text{c}2}\text{S}(\text{O})_2\text{NR}^{\text{c}2}\text{R}^{\text{d}2}, \text{S}(\text{O})\text{R}^{\text{b}2}, \text{S}(\text{O})\text{NR}^{\text{c}2}\text{R}^{\text{d}2}, \text{S}(\text{O})_2\text{R}^{\text{b}2}, \text{and } \text{S}(\text{O})_2\text{NR}^{\text{c}2}\text{R}^{\text{d}2};$$

or R^{2a} and R^{2b} together with the carbon atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered cycloalkyl ring or a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl ring, each optionally substituted by 1, 2, or 3

substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{a2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)₂NR^{c3}R^{d3}, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3};

R^{7a} and R^{7b} are independently selected from H, halo, or C₁₋₄ alkyl;

R^{7c} is H or C₁₋₄ alkyl;

R^{7d} is H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, or C(O)NR^{c5}R^{d5}.

RA is heteroaryl, heterocycloalkyl, heteroarylalkyl, or heterocycloalkylalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4} \text{ alkyl})_r\text{-Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , C(O)R^{b1} , $\text{C(O)NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{C(O)OR}^{\text{a1}}$, $\text{OC(O)R}^{\text{b1}}$, $\text{OC(O)NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{C(=NR}^{\text{e}})\text{NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{NR}^{\text{c1}}\text{C(=NR}^{\text{e}})\text{NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{NR}^{\text{c1}}\text{C(O)R}^{\text{b1}}$, $\text{NR}^{\text{c1}}\text{C(O)OR}^{\text{a1}}$, $\text{NR}^{\text{c1}}\text{C(O)NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{NR}^{\text{c1}}\text{S(O)R}^{\text{b1}}$, $\text{NR}^{\text{c1}}\text{S(O)}_2\text{R}^{\text{b1}}$, $\text{NR}^{\text{c1}}\text{S(O)}_2\text{NR}^{\text{c1}}\text{R}^{\text{d1}}$, S(O)R^{b1} , $\text{S(O)NR}^{\text{c1}}\text{R}^{\text{d1}}$, $\text{S(O)}_2\text{R}^{\text{b1}}$, and $\text{S(O)}_2\text{NR}^{\text{c1}}\text{R}^{\text{d1}}$.

R^B and R^C are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

RE is H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, or C(O)NR^{c5}R^{d5}.

R^N is H or C₁₋₄ alkyl;

Cy¹ is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a5}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

R^a, R^b, R^c, and R^d are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^c and R^d together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5} and S(O)₂NR^{c5}R^{d5};

R^{a1}, R^{b1}, R^{c1}, and R^{d1} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently

selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5} and S(O)₂NR^{c5}R^{d5};

R^{a2}, R^{b2}, R^{c2}, and R^{d2} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c2} and R^{d2} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5} and S(O)₂NR^{c5}R^{d5};

R^{a3}, R^{b3}, R^{c3}, and R^{d3} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c3} and R^{d3} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, halo, C₁₋₆ haloalkyl, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5} and S(O)₂NR^{c5}R^{d5};

R^e and R^f are independently selected from H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, and C(O)NR^{c1}R^{d5};

R^{a5}, R^{b5}, R^{c5}, and R^{d5} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆haloalkoxy;

or R^{c5} and R^{d5} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆alkyl)amino, C₁₋₆haloalkyl, and C₁₋₆haloalkoxy;

m is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

p is 0, 1, 2, 3, or 4;

q is 0, 1, 2, 3, or 4; and

r is 0 or 1.

2. The compound of Embodiment 1, or a pharmaceutically acceptable salt thereof, having Formula I.

3. The compound of Embodiment 1, or a pharmaceutically acceptable salt thereof, having Formula II.

4. The compound of any one of Embodiments 1-3, or a pharmaceutically acceptable salt thereof, wherein A is C₁₋₁₀ alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

5. The compound of any one of Embodiments 1-3, or a pharmaceutically acceptable salt thereof, wherein A is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

6. The compound of any one of Embodiments 1-3, or a pharmaceutically acceptable salt thereof, wherein A is aryl or heteroaryl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

7. The compound of any one of Embodiments 1-6, or a pharmaceutically acceptable salt thereof, wherein L is absent.

8. The compound of any one of Embodiments 1-6, or a pharmaceutically acceptable salt thereof, wherein L is (CR^{7a}R^{7b})_m, (CR^{7a}R^{7b})_pO(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(O)₂(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(O)NR^{7c}(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(O)O(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(=NR^{7d})NR^{7c}(CR^{7a}R^{7b})_q, or (CR^{7a}R^{7b})_pNR^{7c}S(O)₂(CR^{7a}R^{7b})_q.

9. The compound of any one of Embodiments 1-6, or a pharmaceutically acceptable salt thereof, wherein L is (CR^{7a}R^{7b})_m.

10. The compound of any one of Embodiments 1-9, or a pharmaceutically acceptable salt thereof, wherein W is N.

11. The compound of any one of Embodiments 1-9, or a pharmaceutically acceptable salt thereof, wherein W is CR³.

12. The compound of any one of Embodiments 1-11, or a pharmaceutically acceptable salt thereof, wherein X is N.

13. The compound of any one of Embodiments 1-11, or a pharmaceutically acceptable salt thereof, wherein X is CR⁴.

14. The compound of any one of Embodiments 1-13, or a pharmaceutically acceptable salt thereof, wherein Y is N.

15. The compound of any one of Embodiments 1-13, or a pharmaceutically acceptable salt thereof, wherein Y is CR⁵.

16. The compound of any one of Embodiments 1-15, or a pharmaceutically acceptable salt thereof, wherein Z is N.

17. The compound of any one of Embodiments 1-15, or a pharmaceutically acceptable salt thereof, wherein Z is CR⁶.

18. The compound of any one of Embodiments 1-17, or a pharmaceutically acceptable salt thereof, wherein G is O.

19. The compound of any one of Embodiments 1-17, or a pharmaceutically acceptable salt thereof, wherein G is S.

20. The compound of any one of Embodiments 1-17, or a pharmaceutically acceptable salt thereof, wherein G is NR^N.

21. The compound of any one of Embodiments 1-20, or a pharmaceutically acceptable salt thereof, wherein R¹ is heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a5}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

22. The compound of any one of Embodiments 1-20, or a pharmaceutically acceptable salt thereof, wherein R¹ is NR^AR^B.

23. The compound of any one of Embodiments 1-22, or a pharmaceutically acceptable salt thereof, wherein R^{2a} and R^{2b} are independently selected from H and C₁₋₆ alkyl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2}.

24. The compound of any one of Embodiments 1-23, or a pharmaceutically acceptable salt thereof, wherein R³, R⁴, R⁵, and R⁶ are independently selected from H or C₁₋₆ alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3}.

25. The compound of any one of Embodiments 1-23, or a pharmaceutically acceptable salt thereof, wherein R³ is C₁₋₆ alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3}.

26. The compound of any one of Embodiments 1-23, or a pharmaceutically acceptable salt thereof, wherein R⁴, R⁵, and R⁶ are each H.


27. The compound of any one of Embodiments 1-26, or a pharmaceutically acceptable salt thereof, wherein R^{7a}, R^{7b}, R^{7c}, and R^{7d} are each H.

28. The compound of any one of Embodiments 1-27, or a pharmaceutically acceptable salt thereof, wherein R^A is heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.


29. The compound of any one of Embodiments 1-28, or a pharmaceutically acceptable salt thereof, wherein R^B and R^C are independently selected from H and C₁₋₆ alkyl.

30. The compound of any one of Embodiments 1-29, or a pharmaceutically acceptable salt thereof, wherein A is other than phenyl substituted at the 4-position by halogen.


31. The compound of any one of Embodiments 1-30, or a pharmaceutically acceptable salt thereof, having Formula Ia, Ib, Ic, Id, Ie, IIa, IIb, or IIc:




Ia




Ib




Ic




Id




Ie



IIa




IIb




IIc.

32. The compound of Embodiment 1 having Formula If or IId:



If



IId

or a pharmaceutically acceptable salt thereof.

33. The compound of Embodiment 32, or pharmaceutically acceptable salt thereof, wherein:

A is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; R¹ is OR^a, SR^a, S(O)R^a, S(O)₂R^a, NR^aR^b, NR^cC(O)NR^aR^b, NR^cC(O)OR^a, NR^cC(=NR^e)NR^aR^b, NR^cS(O)₂R^a, NR^cS(O)₂NR^cR^a, heterocycloalkyl, or heteroaryl, wherein said heterocycloalkyl or heteroaryl is

optionally substituted with 1, 2, or 3 substituents independently selected from $-(C_{1-4}alkyl)_r-Cy^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a5}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^{2a} is H, halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $C(=NR^e)NR^{c2}R^{d2}$, $NR^{c2}C(=NR^e)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, and $S(O)_2NR^{c2}R^{d2}$;

R^3 , R^4 , R^5 , and R^6 are independently selected from H, halo, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, $S(O)_2NR^{c3}R^{d3}$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3} , SR^{a3} , $C(O)R^{b3}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}C(O)OR^{a3}$, $C(=NR^e)NR^{c3}R^{d3}$, $NR^{c3}C(=NR^e)NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, and $S(O)_2NR^{c3}R^{d3}$;

R^A is heteroaryl, heterocycloalkyl, heteroarylalkyl, or heterocycloalkylalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4}alkyl)_r-Cy^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^B and R^C are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

R^E is H, CN, NO_2 , OR^{a5} , SR^{b5} , $S(O)_2R^{b5}$, $C(O)R^{b5}$, $S(O)_2NR^{c5}R^{d5}$, or $C(O)NR^{c5}R^{d5}$;

Cy^1 is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^a , R^b , R^c , and R^d are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

or R^c and R^d together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

R^{a1} , R^{b1} , R^{c1} , and R^{d1} are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl,

cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^{a2}, R^{b2}, R^{c2}, and R^{d2} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c2} and R^{d2} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^{a3}, R^{b3}, R^{c3}, and R^{d3} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c3} and R^{d3} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, halo, C₁₋₆ haloalkyl, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^e and R^f are independently selected from H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, and C(O)NR^{c5}R^{d5};

R^{a5}, R^{b5}, R^{c5}, and R^{d5} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy;

or R^{c5} and R^{d5} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy; and

r is 0 or 1;

provided that:

a) in Formula If, when A is 3-fluorophenyl; R^{2a} is H; R³ is methyl; and R⁴, R⁵, and R⁶ are H; then, R¹ is other than 4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl; and

b) in Formula If, when A is 1,3-dioxolan-2-yl; R^{2a}, R³, R⁴, and R⁶ are H; then, R¹ is other than 1-(*tert*-butoxycarbonyl)piperidin-4-yl.

42. The compound of any one of Embodiments 33-35, or a pharmaceutically acceptable salt thereof, wherein A is heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

43. The compound of any one of Embodiments 33-35, or a pharmaceutically acceptable salt thereof, wherein A is 6-membered heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

44. The compound of any one of Embodiments 33-35, or a pharmaceutically acceptable salt thereof, wherein A is pyridyl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, or C₁₋₆ haloalkyl.

45. The compound of any one of Embodiments 33-35, or a pharmaceutically acceptable salt thereof, wherein A is 5-membered heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d.

46. The compound of any one of Embodiments 33-35, or a pharmaceutically acceptable salt thereof, wherein A is pyrazolyl optionally substituted with 1 or 2 substituents independently selected from halo, C₁₋₆ alkyl, or C₁₋₆ haloalkyl.

47. The compound of any one of Embodiments 33-46, or a pharmaceutically acceptable salt thereof, wherein R¹ is heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

48. The compound of any one of Embodiments 33-46, or a pharmaceutically acceptable salt thereof, wherein R¹ is bicyclic heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

49. The compound of any one of Embodiments 33-46, or a pharmaceutically acceptable salt thereof, wherein R¹ is purinyl optionally substituted with -(C₁₋₄alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, or S(O)₂NR^{c1}R^{d1}.

50. The compound of any one of Embodiments 33-46, or a pharmaceutically acceptable salt thereof, wherein R^1 is OR^A , SR^A , $S(O)R^A$, $S(O)_2R^A$, $NR^A R^B$, $NR^C C(O)NR^A R^B$, $NR^C C(O)OR^A$, $NR^C C(=NR^E)NR^A R^B$, $NR^C S(O)_2R^A$, or $NR^C S(O)_2NR^C R^A$.

51. The compound of any one of Embodiments 33-46, or a pharmaceutically acceptable salt thereof, wherein R^1 is $NR^A R^B$.


52. The compound of any one of Embodiments 33-51, or a pharmaceutically acceptable salt thereof, wherein R^A is heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4} \text{ alkyl})_r\text{-Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$.

53. The compound of any one of Embodiments 33-51, or a pharmaceutically acceptable salt thereof, wherein R^A is bicyclic heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4} \text{ alkyl})_r\text{-Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$.

54. The compound of any one of Embodiments 33-51, or a pharmaceutically acceptable salt thereof, wherein R^A is purinyl optionally substituted with 1 or 2 substituents independently selected from $-(C_{1-4} \text{ alkyl})_r\text{-Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$.

55. The compound of any one of Embodiments 33-51, or a pharmaceutically acceptable salt thereof, wherein R^A is purinyl optionally substituted with 1 or 2 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$.

56. The compound of any one of Embodiments 33-51, or a pharmaceutically acceptable salt thereof, wherein R^A is:



57. The compound of any one of Embodiments 33-51, or a pharmaceutically acceptable salt thereof, wherein R^A is bicyclic heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4} \text{ alkyl})_r\text{-Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$.

58. The compound of any one of Embodiments 33-57, or a pharmaceutically acceptable salt thereof, wherein R^B and R^C are independently selected from H and C_{1-6} alkyl.

59. The compound of any one of Embodiments 33-57, or a pharmaceutically acceptable salt thereof, wherein R^B and R^C are each H.

60. The compound of any one of Embodiments 33-59, or a pharmaceutically acceptable salt thereof, wherein R^{2a} is H, halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, or C_{2-6} alkynyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo,

C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2}.

61. The compound of any one of Embodiments 33-59, or a pharmaceutically acceptable salt thereof, wherein R^{2a} is halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2}.

62. The compound of any one of Embodiments 33-59, or a pharmaceutically acceptable salt thereof, wherein R^{2a} is H, halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl.

63. The compound of any one of Embodiments 33-59, or a pharmaceutically acceptable salt thereof, wherein R^{2a} is methyl or ethyl.

64. The compound of any one of Embodiments 33-63, or a pharmaceutically acceptable salt thereof, wherein R³ is halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)₂NR^{c3}R^{d3}, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3}.

65. The compound of any one of Embodiments 33-63, or a pharmaceutically acceptable salt thereof, wherein R³ is H, halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, S(O)₂NR^{c3}R^{d3}, or C₁₋₆ alkyl, wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, and S(O)₂NR^{c3}R^{d3}.

66. The compound of any one of Embodiments 33-63, or a pharmaceutically acceptable salt thereof, wherein R³ is H or C₁₋₆ alkyl.

67. The compound of any one of Embodiments 33-63, or a pharmaceutically acceptable salt thereof, wherein R³ is C₁₋₆ alkyl.

68. The compound of any one of Embodiments 33-63, or a pharmaceutically acceptable salt thereof, wherein R³ is methyl.

69. The compound of any one of Embodiments 33-68, or a pharmaceutically acceptable salt thereof, wherein R⁴, R⁵, and R⁶ are independently selected from H, halo, C₁₋₆ alkyl, or C₁₋₆ halo alkyl.

70. The compound of any one of Embodiments 33-68, or a pharmaceutically acceptable salt thereof, wherein R⁴, R⁵, and R⁶ are each H.


71. The compound of any one of Embodiments 33-68, or a pharmaceutically acceptable salt thereof, wherein R⁴ is H.

72. The compound of any one of Embodiments 33-68, or a pharmaceutically acceptable salt thereof, wherein R⁵ is H.


73. The compound of any one of Embodiments 33-68, or a pharmaceutically acceptable salt thereof, wherein R⁶ is H.

74. The compound of any one of Embodiments 33-73, or a pharmaceutically acceptable salt thereof, having Formula

Ig or Iie:




Ig




Iie.

75. The compound of any one of Embodiments 33-74, or a pharmaceutically acceptable salt thereof, having Formula Ih or IIf:




Ih




IIf.

76. The compound of any one of Embodiments 33-75, or a pharmaceutically acceptable salt thereof, having Formula Ii or IIg:




Ii




IIg

wherein R⁸ and R⁹ are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5}.

77. The compound of Embodiment 33, having having Formula If or IId:



If



IId

or a pharmaceutically acceptable salt thereof; wherein:

A is aryl or heteroaryl, each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and OR^a;

R¹ is NR^AR^B or heteroaryl; wherein said heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and OR^{a1};

R^{2a} is C₁₋₆ alkyl;

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, and C₁₋₆ alkyl; wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo;

R^A is heteroaryl, which is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, CN, OR^{a1}, and NR^{c1}R^{d1};

R^B is H;

each R^a is independently selected from H and C₁₋₆ alkyl;


each R^{a1}, R^{c1}, and R^{d1} is independently selected from H and C₁₋₆ alkyl, wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo; and

each R^{a3} is independently selected from H and C₁₋₆ alkyl, wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo.

78. The compound of Embodiment 77, or a pharmaceutically acceptable salt thereof, wherein:

A is phenyl, 5-membered heteroaryl or 6-membered heteroaryl, each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, CN, and OR^a;







R¹ is NR^AR^B or



R^{2a} is C₁₋₆ alkyl;

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, and C₁₋₆ alkyl; wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo;

R^A is selected from:



R^B is H;

each R^a is independently selected from H and C₁₋₆ alkyl; and

each R^{a3} is independently selected from H and C₁₋₆ alkyl, wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo.

79. The compound of Embodiment 1, selected from:

2-[1-(6-amino-9H-purin-9-yl)ethyl]-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 6-methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(5-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3-fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3,5-difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(2-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-3-(1H-pyrazol-4-yl)-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-methyl-6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one; and
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one, or a pharmaceutically acceptable salt of any of the aforementioned.

80. The compound of Embodiment 1, selected from:

6-(3,5-difluorophenyl)-3-methyl-7-[1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3,5-difluorophenyl)-7-[1-[(2-fluoro-9H-purin-6-yl)amino]ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(4-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3,5-difluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-fluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-6-(4-methylphenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-chlorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-chloro-5-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2,3-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(5-fluoropyridin-3-yl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2-chlorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-Fluorophenyl)-7-[(1S)-1-(3H-imidazo[4,5-b]pyridin-7-ylamino)ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyri-

midin-5-one;

6-(3-Fluorophenyl)-7-[(1*S*)-1-[(2-hydroxy-9*H*-purin-6-yl)amino]ethyl]-3-methyl-5*H*-[1,3]thiazolo[3,2-*a*]pyrimidin-5-one;

6-(3-Fluorophenyl)-7-[1-(9*H*-purin-6-ylamino)ethyl]-3-(trifluoromethyl)-5*H*-[1,3]thiazolo[3,2-*a*]pyrimidin-5-one;

6-Methyl-3-phenyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-6-methyl-3-phenyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

6-Methyl-3-(3-methylphenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-6-methyl-3-(3-methylphenyl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3-Chlorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(3-chlorophenyl)-6-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(4-Chlorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(4-chlorophenyl)-6-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(2-Chlorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(2-chlorophenyl)-6-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(2-Fluorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(2-fluorophenyl)-6-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

4-(6-Methyl-4-oxo-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-3-yl)benzonitrile;

4-(2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-6-methyl-4-oxo-4*H*-pyrido[1,2-*a*]pyrimidin-3-yl)benzonitrile;

6-Methyl-3-(2-methylphenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

6-Methyl-3-(4-methylphenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3-Methoxyphenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(2,3-Difluorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(2,5-Difluorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3,4-Difluorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3,5-Difluorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3-Fluorophenyl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(3-fluorophenyl)-6-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3,5-Difluorophenyl)-6-ethyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(3,5-difluorophenyl)-6-ethyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

6-Ethyl-3-(4-fluorophenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(3,5-Difluorophenyl)-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(3,5-difluorophenyl)-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(6-Chloro-5-methylpyridin-3-yl)-6-methyl-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-3-(6-chloro-5-methylpyridin-3-yl)-6-methyl-4*H*-pyrido[1,2-*a*]pyrimidin-4-one;

3-(6-Methyl-4-oxo-2-[1-(9*H*-purin-6-ylamino)ethyl]-4*H*-pyrido[1,2-*a*]pyrimidin-3-yl)benzonitrile; and

3-(2-{1-[(2-Amino-9*H*-purin-6-yl)amino]ethyl}-6-methyl-4-oxo-4*H*-pyrido[1,2-*a*]pyrimidin-3-yl)benzonitrile;

or a pharmaceutically acceptable salt of any of the aforementioned.

81. The compound of Embodiment 1, wherein the compound is (S)-7-(1-(9*H*-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5*H*-thiazolo[3,2-*a*]pyrimidin-5-one; or a pharmaceutically acceptable salt thereof.

82. A composition comprising a compound according to any one of Embodiments 1-81, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.

83. A method of modulating an activity of a PI3K kinase comprising contacting the kinase with a compound of any one of Embodiments 1-81, or a pharmaceutically acceptable salt thereof.

84. The method of Embodiment 83 wherein the PI3K is selected from the group consisting of PI3K α , PI3K β , PI3K γ , and PI3K δ .

85. The method of Embodiment 83 wherein said PI3K comprises a mutation.

86. The method of any one of Embodiments 83-85 wherein said modulating is inhibiting.

87. The method of any one of Embodiments 83-85 wherein said compound is a selective inhibitor for PI3K δ over one or more of PI3K α , PI3K β , and PI3K γ .

88. A method of treating a disease in a patient, wherein said disease is associated with abnormal expression or activity of a PI3K kinase, comprising administering to said patient a therapeutically effective amount of a compound of any one of Embodiments 1-81, or a pharmaceutically acceptable salt thereof.

89. The method of Embodiment 88 wherein said disease is osteoarthritis, restenosis, atherosclerosis, bone disorders, arthritis, diabetic retinopathy, psoriasis, benign prostatic hypertrophy, inflammation, angiogenesis, pancreatitis, kidney disease, inflammatory bowel disease, myasthenia gravis, multiple sclerosis, or Sjogren's syndrome.

90. The method of Embodiment 88 wherein the PI3K is selected from the group consisting of PI3K α , PI3K β , PI3K γ , and PI3K δ .

91. The method of Embodiment 88 wherein the PI3K is PI3K δ .

92. The method of any one of Embodiments 88-91 wherein more than one of said compounds is administered.

93. The method of Embodiment 92 wherein the compound is administered in combination with a kinase inhibitor that inhibits a kinase other than a PI3K kinase.

94. A method of treating an immune-based disease in a patient, comprising administering to said patient a therapeutically effective amount of a compound of any one of Embodiments 1-81, or a pharmaceutically acceptable salt thereof.

95. The method of Embodiment 94 wherein said immune-based disease is rheumatoid arthritis, allergy, asthma, glomerulonephritis, lupus, or inflammation related to any of the aforementioned.

96. A method of treating a cancer in a patient, comprising administering to said patient a therapeutically effective amount of a compound of any one of Embodiments 1-81, or a pharmaceutically acceptable salt thereof.

97. The method of Embodiment 96 wherein said cancer is breast, prostate, colon, endometrial, brain, bladder, skin, uterus, ovary, lung, pancreatic, renal, gastric, or a hematological cancer.


98. The method of Embodiment 97 wherein said hematological cancer is acute myeloblastic leukemia, chronic myeloid leukemia, or B cell lymphoma.

99. A method of treating a lung disease in a patient, comprising administering to said patient a therapeutically effective amount of a compound of any one of Embodiments 1-81, or a pharmaceutically acceptable salt thereof.

100. The method of Embodiment 99 wherein said lung disease is acute lung injury (ALI) or adult respiratory distress syndrome (ARDS).

Claims

1. A compound of Formula II:



II

or a pharmaceutically acceptable salt thereof, wherein:

A is C₁₋₁₀ alkyl, aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5

substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)²R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)²R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

L is absent, (CR^{7a}R^{7b})_m, (CR^{7a}R^{7b})_pO(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(O)(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pS(O)₂(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(O)NR^{7c}(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(O)O(CR^{7a}R^{7b})_q, (CR^{7a}R^{7b})_pNR^{7c}C(=NR^{7d})NR^{7c}(CR^{7a}R^{7b})_q, or (CR^{7a}R^{7b})_pNR^{7c}S(O)₂(CR^{7a}R^{7b})_q;

W is N or CR³;

X is N or CR⁴;

Y is N or CR⁵;

Z is N or CR⁶;

G is O, S, or NR^N;

R¹ is OR^a, SR^a, S(O)R^a, S(O)₂R^a, NR^aR^b, NR^cC(O)NR^aR^b, NR^cC(O)OR^a, NR^cC(=NR^e)NR^aR^b, NR^cS(O)₂R^a, NR^cS(O)₂NR^cR^a, heterocycloalkyl, or heteroaryl, wherein said heterocycloalkyl or heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

R^{2a} and R^{2b} are independently selected from H, halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

or R^{2a} and R^{2b} together with the carbon atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered cycloalkyl ring or a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl ring, each optionally substituted by 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)₂NR^{c3}R^{d3}, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3}, S(O)NR^{c3}R^{d3}, S(O)₂R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, and S(O)₂NR^{c3}R^{d3};

R^{7a} and R^{7b} are independently selected from H, halo, or C₁₋₄ alkyl;

R^{7c} is H or C₁₋₄ alkyl;

R^{7d} is H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, or C(O)NR^{c5}R^{d5};

R^a is heteroaryl, heterocycloalkyl, heteroarylalkyl, or heterocycloalkylalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

R^b and R^c are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl,

Ra³, kb³, Rc³, and Rd³ are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl,

cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c3} and R^{d3} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, halo, C₁₋₆ haloalkyl, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^e and R^f are independently selected from H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, and C(O)NR^{c5}R^{d1};

R^{a5}, R^{b5}, R^{c5}, and R^{d5} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy;

or R^{c5} and R^{d5} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆ alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy;

m is 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10;

p is 0, 1, 2, 3, or 4;

q is 0, 1, 2, 3, or 4; and

r is 0 or 1.

2. The compound of any one of claim 1, or a pharmaceutically acceptable salt thereof, wherein A is:

(a) C₁₋₁₀ alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(b) aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; or

(c) aryl or heteroaryl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d,

NR^cR^d , $\text{NR}^c\text{C}(\text{O})\text{R}^b$, $\text{NR}^c\text{C}(\text{O})\text{OR}^a$, $\text{NR}^c\text{C}(\text{O})\text{NR}^c\text{R}^d$, $\text{NR}^c\text{S}(\text{O})\text{R}^b$, $\text{NR}^c\text{S}(\text{O})_2\text{R}^b$, $\text{NR}^c\text{S}(\text{O})_2\text{NR}^c\text{R}^d$, $\text{S}(\text{O})\text{R}^b$, $\text{S}(\text{O})\text{NR}^c\text{R}^d$, $\text{S}(\text{O})_2\text{R}^b$, and $\text{S}(\text{O})_2\text{NR}^c\text{R}^d$.

3. The compound of any one of claims 1-2, or a pharmaceutically acceptable salt thereof, wherein L is:

(a) absent; or

(b) $(\text{CR}^{7a}\text{R}^{7b})_m$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{O}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{S}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{S}(\text{O})$ $(\text{CR}^{7a}\text{R}^{7b})_q$,
 $(\text{CR}^{7a}\text{R}^{7b})_p\text{S}(\text{O})_2(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{C}(\text{O})\text{NR}^{7c}(\text{CR}^{7a}\text{R}^{7b})_q$,
 $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{C}(\text{O})\text{O}(\text{CR}^{7a}\text{R}^{7b})_q$, $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{C}(=\text{NR}^{7d})\text{NR}^{7c}(\text{CR}^{7a}\text{R}^{7b})_q$, or
 $(\text{CR}^{7a}\text{R}^{7b})_p\text{NR}^{7c}\text{S}(\text{O})_2(\text{CR}^{7a}\text{R}^{7b})_q$;
 (c) $(\text{CR}^{7a}\text{R}^{7b})_m$,

4. The compound of any one of claims 1-3, or a pharmaceutically acceptable salt thereof, wherein W is:

(a) N; or

(b) CR^3 .

5. The compound of any one of claims 1-4, or a pharmaceutically acceptable salt thereof, wherein X is:

(a) N; or

(b) CR^4 .

6. The compound of any one of claims 1-5, or a pharmaceutically acceptable salt thereof, wherein G is:

(a) O;

(b) S; or

(c) NR^N .

7. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt thereof, wherein R^1 is:

(a) heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from from $-(\text{C}_{1-4}\text{alkyl})-\text{Cy}^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $\text{C}(\text{O})\text{R}^{b1}$, $\text{C}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{C}(\text{O})\text{OR}^{a1}$, $\text{OC}(\text{O})\text{R}^{b1}$, $\text{OC}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{C}(=\text{NR}^e)\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}\text{C}(=\text{NR}^e)\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}\text{C}(\text{O})\text{R}^{b1}$, $\text{NR}^{c1}\text{C}(\text{O})\text{OR}^{a1}$, $\text{NR}^{c1}\text{C}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{NR}^{c1}\text{S}(\text{O})\text{R}^{b1}$, $\text{NR}^{c1}\text{S}(\text{O})_2\text{R}^{b1}$, $\text{NR}^{c1}\text{S}(\text{O})_2\text{NR}^{c1}\text{R}^{d1}$, $\text{S}(\text{O})\text{R}^{b1}$, $\text{S}(\text{O})\text{NR}^{c1}\text{R}^{d1}$, $\text{S}(\text{O})_2\text{R}^{b1}$, and $\text{S}(\text{O})_2\text{NR}^{c1}\text{R}^{d1}$; or
 (b) NR^{ARB} .

8. The compound of any one of claims 1-7, or a pharmaceutically acceptable salt thereof, wherein R^{2a} and R^{2b} are independently selected from H and C_{1-6} alkyl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a2} , SR^{a2} , $\text{C}(\text{O})\text{R}^{b2}$, $\text{C}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{C}(\text{O})\text{OR}^{a2}$, $\text{OC}(\text{O})\text{R}^{b2}$, $\text{OC}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{C}(=\text{NR}^e)\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{C}(=\text{NR}^e)\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{C}(\text{O})\text{R}^{b2}$, $\text{NR}^{c2}\text{C}(\text{O})\text{OR}^{a2}$, $\text{NR}^{c2}\text{C}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{NR}^{c2}\text{S}(\text{O})\text{R}^{b2}$, $\text{NR}^{c2}\text{S}(\text{O})_2\text{R}^{b2}$, $\text{NR}^{c2}\text{S}(\text{O})_2\text{NR}^{c2}\text{R}^{d2}$, $\text{S}(\text{O})\text{R}^{b2}$, $\text{S}(\text{O})\text{NR}^{c2}\text{R}^{d2}$, $\text{S}(\text{O})_2\text{R}^{b2}$, and $\text{S}(\text{O})_2\text{NR}^{c2}\text{R}^{d2}$.

9. The compound of any one of claims 1-8, or a pharmaceutically acceptable salt thereof, wherein:


(a) R^3 , R^4 , R^5 , and R^6 are independently selected from H or C_{1-6} alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3} , SR^{a3} , $\text{C}(\text{O})\text{R}^{b3}$, $\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{C}(\text{O})\text{OR}^{a3}$, $\text{OC}(\text{O})\text{R}^{b3}$, $\text{OC}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{R}^{b3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{OR}^{a3}$, $\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})\text{R}^{b3}$, $\text{S}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$, and $\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$;

(b) R^3 is C_{1-6} alkyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a3} , SR^{a3} , $\text{C}(\text{O})\text{R}^{b3}$, $\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{C}(\text{O})\text{OR}^{a3}$, $\text{OC}(\text{O})\text{R}^{b3}$, $\text{OC}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{R}^{b3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(\text{O})\text{OR}^{a3}$, $\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{NR}^{c3}\text{C}(=\text{NR}^e)\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})\text{R}^{b3}$, $\text{S}(\text{O})\text{NR}^{c3}\text{R}^{d3}$, $\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{R}^{b3}$, $\text{NR}^{c3}\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$, and $\text{S}(\text{O})_2\text{NR}^{c3}\text{R}^{d3}$; or
 (c) R^4 , R^5 , and R^6 are each H.


10. The compound of any one of claims 1-9, or a pharmaceutically acceptable salt thereof, wherein R^{7a} , R^{7b} , R^{7c} , and

R^{7d} are each H.


11. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein R^A is heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.
12. The compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof, wherein R^B and R^C are independently selected from H and C₁₋₆ alkyl.
13. The compound of any one of claims 1-12, or a pharmaceutically acceptable salt thereof, wherein A is other than phenyl substituted at the 4-position by halogen.
14. The compound of any one of claims 1-13, or a pharmaceutically acceptable salt thereof, having Formula IIa, IIb, or IIc:



IIa




IIb



IIc.

15. The compound of claim 1 having Formula IIId:



IIId

or a pharmaceutically acceptable salt thereof.

16. The compound of claim 15, or pharmaceutically acceptable salt thereof, wherein:

A is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and

$S(O)_2NR^cR^d$; wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^a , SR^a , $C(O)R^b$, $C(O)NR^cR^d$, $C(O)OR^a$, $OC(O)R^b$, $OC(O)NR^cR^d$, $C(=NR^e)NR^cR^d$, $NR^cC(=NR^e)NR^cR^d$, NR^cR^d , $NR^cC(O)R^b$, $NR^cC(O)OR^a$, $NR^cC(O)NR^cR^d$, $NR^cS(O)R^b$, $NR^cS(O)_2R^b$, $NR^cS(O)_2NR^cR^d$, $S(O)R^b$, $S(O)NR^cR^d$, $S(O)_2R^b$, and $S(O)_2NR^cR^d$;

R^1 is OR^a , SR^a , $S(O)R^a$, $S(O)_2R^a$, NR^aR^b , $NR^cC(O)NR^aR^b$, $NR^cC(O)OR^a$, $NR^cC(=NR^e)NR^aR^b$, $NR^cS(O)_2R^a$, $NR^cS(O)_2NR^cR^a$, heterocycloalkyl, or heteroaryl, wherein said heterocycloalkyl or heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from $-(C_{1-4}alkyl)_r-Cy^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{a1}$;

R^{2a} is H, halo, OH, CN, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{1-6} haloalkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a2}$, $OC(O)R^{b2}$, $OC(O)NR^{c2}R^{d2}$, $C(=NR^e)NR^{c2}R^{d2}$, $NR^{c2}C(=NR^e)NR^{c2}R^{d2}$, $NR^{c2}R^{d2}$, $NR^{c2}C(O)R^{b2}$, $NR^{c2}C(O)OR^{a2}$, $NR^{c2}C(O)NR^{c2}R^{d2}$, $NR^{c2}S(O)R^{b2}$, $NR^{c2}S(O)_2R^{b2}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)R^{b2}$, $S(O)NR^{c2}R^{d2}$, $S(O)_2R^{b2}$, and $S(O)_2NR^{c2}R^{d2}$;

R^3 , R^4 , R^5 , and R^6 are independently selected from H, halo, CN, NO_2 , OR^{a3} , SR^{a3} , $C(O)R^{b2}$, $C(O)NR^{c2}R^{d2}$, $C(O)OR^{a3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c2}S(O)_2NR^{c2}R^{d2}$, $S(O)_2NR^{c3}R^{d3}$, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a2} , SR^{a2} , $C(O)R^{b2}$, $C(O)NR^{c3}R^{d3}$, $C(O)OR^{a3}$, $OC(O)R^{b3}$, $OC(O)NR^{c3}R^{d3}$, $NR^{c3}R^{d3}$, $NR^{c3}C(O)R^{b3}$, $NR^{c3}C(O)NR^{c3}R^{d3}$, $NR^{c3}C(O)OR^{a3}$, $C(=NR^e)NR^{c3}R^{d3}$, $NR^{c3}C(=NR^e)NR^{c3}R^{d3}$, $S(O)R^{b3}$, $S(O)NR^{c3}R^{d3}$, $S(O)_2R^{b3}$, $NR^{c3}S(O)_2R^{b3}$, $NR^{c3}S(O)_2NR^{c3}R^{d3}$, and $S(O)_2NR^{c3}R^{d3}$;

R^A is heteroaryl, heterocycloalkyl, heteroarylalkyl, or heterocycloalkylalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from $-(C_{1-4}alkyl)_r-Cy^1$, halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^B and R^C are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

R^E is H, CN, NO_2 , OR^{a5} , SR^{b5} , $S(O)_2R^{b5}$, $C(O)R^{b5}$, $S(O)_2NR^{c5}R^{d5}$, or $C(O)NR^{c5}R^{d5}$;

Cy^1 is aryl, heteroaryl, cycloalkyl, or heterocycloalkyl, each optionally substituted by 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, halosulfanyl, CN, NO_2 , OR^{a1} , SR^{a1} , $C(O)R^{b1}$, $C(O)NR^{c1}R^{d1}$, $C(O)OR^{a1}$, $OC(O)R^{b1}$, $OC(O)NR^{c1}R^{d1}$, $C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}C(=NR^e)NR^{c1}R^{d1}$, $NR^{c1}R^{d1}$, $NR^{c1}C(O)R^{b1}$, $NR^{c1}C(O)OR^{a1}$, $NR^{c1}C(O)NR^{c1}R^{d1}$, $NR^{c1}S(O)R^{b1}$, $NR^{c1}S(O)_2R^{b1}$, $NR^{c1}S(O)_2NR^{c1}R^{d1}$, $S(O)R^{b1}$, $S(O)NR^{c1}R^{d1}$, $S(O)_2R^{b1}$, and $S(O)_2NR^{c1}R^{d1}$;

R^a , R^b , R^c , and R^d are independently selected from H, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$;

or R^c and R^d together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently

selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)R^{b5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^{a1}, R^{b1}, R^{c1}, and R^{d1} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c1} and R^{d1} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^{a2}, R^{b2}, R^{c2}, and R^{d2} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c2} and R^{d2} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^{a3}, R^{b3}, R^{c3}, and R^{d3} are independently selected from H, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

or R^{c3} and R^{d3} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or a heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from C₁₋₆ alkyl, halo, C₁₋₆ haloalkyl, CN, OR^{a5}, SR^{a5}, C(O)R^{b5}, C(O)NR^{c5}R^{d5}, C(O)OR^{a5}, OC(O)R^{b5}, OC(O)NR^{c5}R^{d5}, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, C(=NR^f)NR^{c5}R^{d5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, S(O)R^{b5}, S(O)NR^{c5}R^{d5}, S(O)₂R^{b5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

R^e and R^f are independently selected from H, CN, NO₂, OR^{a5}, SR^{b5}, S(O)₂R^{b5}, C(O)R^{b5}, S(O)₂NR^{c5}R^{d5}, and C(O)NR^{c5}R^{d5};

R^{a5}, R^{b5}, R^{c5}, and R^{d5} are independently selected from H, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy;

or R^{c5} and R^{d5} together with the N atom to which they are attached form a 3-, 4-, 5-, 6-, or 7-membered heterocycloalkyl group or heteroaryl group, each optionally substituted with 1, 2, or 3 substituents independently selected from OH, CN, amino, halo, C₁₋₆ alkyl, C₁₋₆ alkoxy, C₁₋₆ alkylthio, C₁₋₆ alkylamino, di(C₁₋₆alkyl)amino, C₁₋₆ haloalkyl, and C₁₋₆ haloalkoxy; and

r is 0 or 1;
provided that:

- a) in Formula If, when A is 3-fluorophenyl; R^{2a} is H; R³ is methyl; and R⁴, R⁵, and R⁶ are H; then, R¹ is other than 4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl; and
b) in Formula If, when A is 1,3-dioxolan-2-yl; R^{2a}, R³, R⁴, and R⁶ are H; then, R¹ is other than 1-(*tert*-butoxycarbonyl)piperidin-4-yl.

17. The compound of claim 16, or a pharmaceutically acceptable salt thereof, wherein A is:

(a) cycloalkyl or heterocycloalkyl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(b) aryl or heteroaryl, each optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(c) aryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(d) phenyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(e) phenyl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, or C₁₋₆ haloalkyl;

(f) phenyl;

(g) heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(h) 6-membered heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d;

(i) pyridyl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, or C₁₋₆ haloalkyl;

(j) 5-membered heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, C(=NR^e)R^b, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, or heterocycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^a, SR^a, C(O)R^b, C(O)NR^cR^d, C(O)OR^a, OC(O)R^b, OC(O)NR^cR^d, C(=NR^e)NR^cR^d, NR^cC(=NR^e)NR^cR^d, NR^cR^d, NR^cC(O)R^b, NR^cC(O)OR^a, NR^cC(O)NR^cR^d, NR^cS(O)R^b, NR^cS(O)₂R^b, NR^cS(O)₂NR^cR^d, S(O)R^b, S(O)NR^cR^d, S(O)₂R^b, and S(O)₂NR^cR^d; or

(k) pyrazolyl optionally substituted with 1 or 2 substituents independently selected from halo, C₁₋₆ alkyl, or C₁₋₆ haloalkyl.

18. The compound of any one of claims 16-17, or a pharmaceutically acceptable salt thereof, wherein R¹ is:

(a) heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

(b) bicyclic heteroaryl optionally substituted with 1, 2, or 3 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

(c) purinyl optionally substituted with -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, or S(O)₂NR^{c1}R^{d1};

(d) OR^A, SR^A, S(O)R^A, S(O)₂R^A, NR^AR^B, NR^CC(O)NR^AR^B, NR^CC(O)OR^A, NR^CC(=NR^E)NR^AR^B, NR^CS(O)₂R^A, or NR^CS(O)₂NR^CR^A; or

(e) NR^AR^B.

19. The compound of any one of claims 16-18, or a pharmaceutically acceptable salt thereof, wherein R^A is:

(a) heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};


(b) bicyclic heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)_r-Cy¹,

alkyl)-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};


(c) purinyl optionally substituted with 1 or 2 substituents independently selected from -(C₁₋₄ alkyl)-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1};

(d) purinyl optionally substituted with 1 or 2 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, NR^{c5}R^{d5}, NR^{c5}C(O)R^{b5}, NR^{c5}C(O)NR^{c5}R^{d5}, NR^{c5}C(O)OR^{a5}, NR^{c5}C(=NR^f)NR^{c5}R^{d5}, NR^{c5}S(O)₂R^{b5}, NR^{c5}S(O)₂NR^{c5}R^{d5}, and S(O)₂NR^{c5}R^{d5};

(e) selected from the group consisting of:



or



; or

(f) bicyclic heteroaryl optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from -(C₁₋₄ alkyl)-Cy¹, halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a1}, SR^{a1}, C(O)R^{b1}, C(O)NR^{c1}R^{d1}, C(O)OR^{a1}, OC(O)R^{b1}, OC(O)NR^{c1}R^{d1}, C(=NR^e)NR^{c1}R^{d1}, NR^{c1}C(=NR^e)NR^{c1}R^{d1}, NR^{c1}R^{d1}, NR^{c1}C(O)R^{b1}, NR^{c1}C(O)OR^{a1}, NR^{c1}C(O)NR^{c1}R^{d1}, NR^{c1}S(O)R^{b1}, NR^{c1}S(O)₂R^{b1}, NR^{c1}S(O)₂NR^{c1}R^{d1}, S(O)R^{b1}, S(O)NR^{c1}R^{d1}, S(O)₂R^{b1}, and S(O)₂NR^{c1}R^{d1}.

20. The compound of any one of claims 16-19, or a pharmaceutically acceptable salt thereof, wherein R^B and R^C are:

- (a) independently selected from H and C₁₋₆ alkyl; or
(b) each H.

21. The compound of any one of claims 16-20, or a pharmaceutically acceptable salt thereof, wherein R^{2a} is:

(a) H, halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

(b) halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl, wherein said C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, halosulfanyl, CN, NO₂, OR^{a2}, SR^{a2}, C(O)R^{b2}, C(O)NR^{c2}R^{d2}, C(O)OR^{a2}, OC(O)R^{b2}, OC(O)NR^{c2}R^{d2}, C(=NR^e)NR^{c2}R^{d2}, NR^{c2}C(=NR^e)NR^{c2}R^{d2}, NR^{c2}R^{d2}, NR^{c2}C(O)R^{b2}, NR^{c2}C(O)OR^{a2}, NR^{c2}C(O)NR^{c2}R^{d2}, NR^{c2}S(O)R^{b2}, NR^{c2}S(O)₂R^{b2}, NR^{c2}S(O)₂NR^{c2}R^{d2}, S(O)R^{b2}, S(O)NR^{c2}R^{d2}, S(O)₂R^{b2}, and S(O)₂NR^{c2}R^{d2};

- (c) H, halo, OH, CN, C₁₋₆ alkyl, C₁₋₆ haloalkyl, C₂₋₆ alkenyl, or C₂₋₆ alkynyl; or
(d) methyl or ethyl.

22. The compound of any one of claims 16-21, or a pharmaceutically acceptable salt thereof, wherein R³ is:

(a) halo, CN, NO₂, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a1}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}S(O)₂R^{b3}, NR^{c3}S(O)₂NR^{c3}R^{d3}, S(O)₂NR^{c3}R^{d3}, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, and heterocycloalkylalkyl, wherein said C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, aryl, cycloalkyl, heteroaryl, heterocycloalkyl, arylalkyl, heteroarylalkyl, cycloalkylalkyl, or heterocycloalkylalkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from C₁₋₆ alkyl, C₁₋₆ haloalkyl, halo, CN, OR^{a3}, SR^{a3}, C(O)R^{b3}, C(O)NR^{c3}R^{d3}, C(O)OR^{a3}, OC(O)R^{b3}, OC(O)NR^{c3}R^{d3}, NR^{c3}R^{d3}, NR^{c3}C(O)R^{b3}, NR^{c3}C(O)NR^{c3}R^{d3}, NR^{c3}C(O)OR^{a3}, C(=NR^e)NR^{c3}R^{d3}, NR^{c3}C(=NR^e)NR^{c3}R^{d3}, S(O)R^{b3},

$S(O)NR^cR^d$, $S(O)_2R^b$, $NR^cS(O)_2R^b$, $NR^cS(O)_2NR^cR^d$, and $S(O)_2NR^cR^d$;

(b) H, halo, CN, NO_2 , OR^a , SR^a , $C(O)R^b$, $C(O)NR^cR^d$, $C(O)OR^a$, NR^cR^d , $NR^cC(O)R^b$, $NR^cS(O)_2R^b$, $S(O)_2NR^cR^d$, or C_{1-6} alkyl, wherein said C_{1-6} alkyl, is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, CN, OR^a , SR^a , $C(O)R^b$, $C(O)NR^cR^d$, $C(O)OR^a$, $OC(O)R^b$, $OC(O)NR^cR^d$, NR^cR^d , $NR^cC(O)R^b$, $NR^cC(O)NR^cR^d$, $NR^cC(O)OR^a$, $C(=NR^e)NR^cR^d$, $NR^cC(=NR^e)NR^cR^d$, $S(O)R^b$, $S(O)NR^cR^d$, $S(O)_2R^b$, $NR^cS(O)_2R^b$, and $S(O)_2NR^cR^d$;

(c) H or C_{1-6} alkyl;

(d) C_{1-6} alkyl; or

(e) methyl.

23. The compound of any one of claims 16-22, or a pharmaceutically acceptable salt thereof, wherein R^4 , R^5 , and R^6 are:

(a) independently selected from H, halo, C_{1-6} alkyl, or C_{1-6} haloalkyl; or

(b) are each H.


24. The compound of any one of claims 16-23, or a pharmaceutically acceptable salt thereof, wherein:

(a) R^4 is H; or

(b) R^6 is H.


25. The compound of any one of claims 16-24, or a pharmaceutically acceptable salt thereof, having:

(a) Formula IIe:




IIe;

(b) Formula II f:



II f; or


(c) Formula II g:



IIg

wherein R^8 and R^9 are independently selected from H, C_{1-6} alkyl, C_{1-6} haloalkyl, halo, CN, OR^{a5} , SR^{a5} , $C(O)R^{b5}$, $C(O)NR^{c5}R^{d5}$, $C(O)OR^{a5}$, $OC(O)R^{b5}$, $OC(O)NR^{c5}R^{d5}$, $NR^{c5}R^{d5}$, $NR^{c5}C(O)R^{b5}$, $NR^{c5}C(O)NR^{c5}R^{d5}$, $NR^{c5}C(O)OR^{a5}$, $C(=NR^f)NR^{c5}R^{d5}$, $NR^{c5}C(=NR^f)NR^{c5}R^{d5}$, $S(O)R^{b5}$, $S(O)NR^{c5}R^{d5}$, $S(O)_2R^{b5}$, $NR^{c5}S(O)_2R^{b5}$, $NR^{c5}S(O)_2NR^{c5}R^{d5}$, and $S(O)_2NR^{c5}R^{d5}$.

26. The compound of claim 16, having Formula IIId:




IIId

or a pharmaceutically acceptable salt thereof; wherein:

A is aryl or heteroaryl, each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, CN, and OR^a ;
 R^1 is NR^AR^B or heteroaryl; wherein said heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, CN, and OR^{a1} ;
 R^{2a} is C_{1-6} alkyl;
 R^3 , R^4 , R^5 , and R^6 are independently selected from H, halo, CN, NO_2 , OR^{a3} , and C_{1-6} alkyl; wherein said C_{1-6} alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo;
 R^A is heteroaryl, which is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo, C_{1-6} alkyl, CN, OR^{a1} , and $NR^{c1}R^{d1}$;
 R^B is H;
each R^a is independently selected from H and C_{1-6} alkyl;
each R^{a1} , R^{c1} , and R^{d1} is independently selected from H and C_{1-6} alkyl, wherein said C_{1-6} alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo; and
each R^{a3} is independently selected from H and C_{1-6} alkyl, wherein said C_{1-6} alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo.

27. The compound of claim 26, or a pharmaceutically acceptable salt thereof, wherein:


A is phenyl, 5-membered heteroaryl or 6-membered heteroaryl, each optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, CN, and OR^a ;
 R^1 is NR^AR^B or



R^{2a} is C₁₋₆ alkyl;

R³, R⁴, R⁵, and R⁶ are independently selected from H, halo, CN, NO₂, OR^{a3}, and C₁₋₆ alkyl; wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo;

R^A is selected from:



R^B is H;

each R^a is independently selected from H and C₁₋₆ alkyl; and

each R^{a3} is independently selected from H and C₁₋₆ alkyl, wherein said C₁₋₆ alkyl is optionally substituted with 1, 2, 3, 4, or 5 substituents independently selected from halo.


28. A compound, selected from:


- 2-[1-(6-amino-9H-purin-9-yl)ethyl]-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 6-methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(5-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3-fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3,5-difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(2-fluoropyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 6-methyl-2-[1-(9H-purin-6-ylamino)propyl]-3-(1H-pyrazol-4-yl)-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-methyl-6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-3-methyl-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3,5-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-2-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3,5-difluorophenyl)-3-methyl-7-[1-(7H-pyrrolo[2,3-d]pyrimidin-4-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3,5-difluorophenyl)-7-[1-[(2-fluoro-9H-purin-6-yl)amino]ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-pyridin-4-yl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-2-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-6-(1,3-thiazol-4-yl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(4-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-[1-[(2-amino-9H-purin-6-yl)amino]ethyl]-6-(3,5-difluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;

6-(3,5-difluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-fluorophenyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-phenyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-fluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 5 6-phenyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 3-methyl-6-(4-methylphenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2,3-difluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-
 10 one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(3-chloro-5-fluorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimi-
 din-5-one;
 6-(3-chlorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-chloro-5-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 15 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(5-fluoropyridin-3-yl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-
 one;
 7-{1-[(2-amino-9H-purin-6-yl)amino]ethyl}-6-(2-chlorophenyl)-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2-fluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2,3-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 20 6-(5-fluoropyridin-3-yl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2-chlorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(2,5-difluorophenyl)-3-methyl-7-[1-(9H-purin-6-ylamino)ethyl]-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-(3-Fluorophenyl)-7-[(1S)-1-(3H-imidazo[4,5-b]pyridin-7-ylamino)ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyri-
 midin-5-one;
 25 6-(3-Fluorophenyl)-7-[(1S)-1-[(2-hydroxy-9H-purin-6-yl)amino]ethyl]-3-methyl-5H-[1,3]thiazolo[3,2-a]pyrimi-
 din-5-one;
 6-(3-Fluorophenyl)-7-[1-(9H-purin-6-ylamino)ethyl]-3-(trifluoromethyl)-5H-[1,3]thiazolo[3,2-a]pyrimidin-5-one;
 6-Methyl-3-phenyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-6-methyl-3-phenyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 30 6-Methyl-3-(3-methylphenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-6-methyl-3-(3-methylphenyl)-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3-Chlorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3-chlorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(4-Chlorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 35 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(4-chlorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(2-Chlorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(2-chlorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(2-Fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(2-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 40 4-[6-Methyl-4-oxo-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-3-yl]benzonitrile;
 4-(2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-6-methyl-4-oxo-4H-pyrido[1,2-a]pyrimidin-3-yl)benzonitrile;
 6-Methyl-3-(2-methylphenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 6-Methyl-3-(4-methylphenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3-Methoxyphenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 45 3-(2,3-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(2,5-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3,4-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3,5-Difluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3-Fluorophenyl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 50 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3-fluorophenyl)-6-methyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3,5-Difluorophenyl)-6-ethyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3,5-difluorophenyl)-6-ethyl-4H-pyrido[1,2-a]pyrimidin-4-one;
 6-Ethyl-3-(4-fluorophenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 3-(3,5-Difluorophenyl)-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 55 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(3,5-difluorophenyl)-4H-pyrido 1,2-a]pyrimidin-4-one;
 3-(6-Chloro-5-methylpyridin-3-yl)-6-methyl-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-4-one;
 2- {1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-3-(6-chloro-5-methylpyridin-3-yl)-6-methyl-4H-pyrido[1,2-a]pyrimi-
 din-4-one;

3-{6-Methyl-4-oxo-2-[1-(9H-purin-6-ylamino)ethyl]-4H-pyrido[1,2-a]pyrimidin-3-yl}benzonitrile; and
3-(2-{1-[(2-Amino-9H-purin-6-yl)amino]ethyl}-6-methyl-4-oxo-4H-pyrido[1,2-a]pyrimidin-3-yl)benzonitrile;
or a pharmaceutically acceptable salt of any of the aforementioned.

- 5 **29.** The compound of claim 1, which is 7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one; or a pharmaceutically acceptable salt thereof.
- 10 **30.** The compound of claim 1, wherein the compound is (S)-7-(1-(9H-purin-6-ylamino)ethyl)-6-(3-fluorophenyl)-3-methyl-5H-thiazolo[3,2-a]pyrimidin-5-one; or a pharmaceutically acceptable salt thereof.
- 15 **31.** A composition comprising a compound according to any one of claims 1-30, or a pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier.
- 20 **32.** A compound of any one of claims 1 to 30, or a pharmaceutically acceptable salt thereof, for use in inhibiting an activity of a PI3K kinase.
- 25 **33.** The compound for use according to claim 32 wherein the PI3K is selected from the group consisting of PI3K α , PI3K β , PI3K γ , and PI3K δ .
- 30 **34.** The compound for use according to any one of claims 32-33 wherein said compound is a selective inhibitor for PI3K δ over one or more of PI3K α , PI3K β , and PI3K γ .
- 35 **35.** A compound of any one of claims 1 to 30, or a pharmaceutically acceptable salt thereof, for use a method of treating a disease in a patient, wherein said disease is associated with abnormal expression or activity of a PI3K kinase.
- 40 **36.** The compound for use according to claim 35 wherein:
 - (a) said disease is osteoarthritis, restenosis, atherosclerosis, bone disorders, arthritis, diabetic retinopathy, psoriasis, benign prostatic hypertrophy, inflammation, angiogenesis, pancreatitis, kidney disease, inflammatory bowel disease, myasthenia gravis, multiple sclerosis, or Sjogren's syndrome;
 - (b) the PI3K is: (i) selected from the group consisting of PI3K α , PI3K β , PI3K γ , and PI3K δ ; or (ii) is PI3K δ .
- 45 **37.** The compound for use according to claim 35 wherein the compound is administered in combination with a kinase inhibitor that inhibits a kinase other than a PI3K kinase.
- 50 **38.** A compound of any one of claims 1 to 30, or a pharmaceutically acceptable salt thereof, for use in method of treating (a) an immune-based disease in a patient; (b) cancer; or (c) a lung disease.
- 55 **39.** A compound of any one of claims 1 to 30, or a pharmaceutically acceptable salt thereof, for use in a method of treating:
 - (a) an immune-based disease selected from rheumatoid arthritis, allergy, asthma, glomerulonephritis, lupus, or inflammation related to any of the aforementioned;
 - (b) cancer wherein the cancer is breast, prostate, colon, endometrial, brain, bladder, skin, uterus, ovary, lung, pancreatic, renal, gastric, or a hematological cancer;
 - (c) hematological cancer wherein the hematological cancer is acute myeloblastic leukemia, chronic myeloid leukemia, or B cell lymphoma; or
 - (d) lung disease wherein the lung cancer is acute lung injury (ALI) or adult respiratory distress syndrome (ARDS).







EUROPEAN SEARCH REPORT

Application Number
EP 14 16 9702

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X	WO 2006/008523 A1 (ASTRAZENECA AB [SE]; ASTRAZENECA UK LTD [GB]; BLOCK MICHAEL HOWARD [US] 26 January 2006 (2006-01-26) * method 13; page 37 * * page 22, line 22 - page 23, line 3 * * claims 1-10, 13, 15, 16, 19 *	1-14, 31-36, 38,39	INV. C07D473/34 C07D513/04 A61K31/52 A61P29/00 A61P37/00
A	WO 2008/064018 A1 (LILLY CO ELI [US]; WHITE STEPHEN L [US] LILLY CO ELI [US]; WHITE STEPH) 29 May 2008 (2008-05-29) * paragraph [0001] * * examples 1-17 * * claims 1-12 *	1-39	
			TECHNICAL FIELDS SEARCHED (IPC)
			C07D
The present search report has been drawn up for all claims			
Place of search Munich		Date of completion of the search 30 January 2015	Examiner Koch, Kristian
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 14 16 9702

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

30-01-2015

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2006008523 A1	26-01-2006	AU 2005263969 A1	26-01-2006
		BR PI0513513 A	06-05-2008
		CA 2574204 A1	26-01-2006
		CN 101023082 A	22-08-2007
		EP 1773830 A1	18-04-2007
		JP 2008506759 A	06-03-2008
		KR 20070044458 A	27-04-2007
		RU 2007106552 A	27-08-2008
		UA 84954 C2	10-12-2008
		US 2007287703 A1	13-12-2007
		WO 2006008523 A1	26-01-2006
		ZA 200700227 A	28-05-2008
WO 2008064018 A1	29-05-2008	AU 2007323836 A1	29-05-2008
		CA 2669399 A1	29-05-2008
		CN 101605797 A	16-12-2009
		EP 2097422 A1	09-09-2009
		EP 2441768 A1	18-04-2012
		JP 5313909 B2	09-10-2013
		JP 2010509373 A	25-03-2010
		KR 20090087027 A	14-08-2009
		NZ 576997 A	24-02-2012
		SG 178786 A1	29-03-2012
		US 2010216820 A1	26-08-2010
		WO 2008064018 A1	29-05-2008

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61221160 A [0001]
- US 61259765 A [0001]
- WO 2006056399 A [0187]
- US 5521184 A [0192]
- WO 04005281 A [0192]
- US 60578491 B [0192]
- WO 03037347 A [0193]
- WO 03099771 A [0193]
- WO 04046120 A [0193]
- WO 0009495 A [0194]
- WO 05028444 A [0194]
- WO 04080980 A [0195]
- WO 04056786 A [0195]
- WO 03024967 A [0195]
- WO 01064655 A [0195]
- WO 00053595 A [0195]
- WO 01014402 A [0195]
- WO 2002000196 A [0203]

Non-patent literature cited in the description

- **CANTLEY**. *Science*, 2002, vol. 296 (5573), 1655-7 [0003]
- **JIMENEZ et al.** *J Biol Chem.*, 2002, vol. 277 (44), 41556-62 [0003]
- **BROCK et al.** *J Cell Biol.*, 2003, vol. 160 (1), 89-99 [0003]
- **VANHAESEBROECK et al.** *Trends Biochem Sci.*, 2005, vol. 30 (4), 194-204 [0003]
- **SASAKI et al.** *Science*, 2000, vol. 287 (5455), 1040-6 [0004]
- **CLAYTON et al.** *J. Exp. Med.*, 2002, vol. 196 (6), 753-63 [0004]
- **JOU et al.** *Mol. Cell Biol.*, 2002, vol. 22 (24), 8580-91 [0004]
- **OKKENHAUG et al.** *Science*, 2002, vol. 297 (5583), 1031-4 [0004]
- **CAMPS et al.** *Nat Med.*, 2005, vol. 11 (9), 936-43 [0005]
- **THOMAS et al.** *Eur. J. Immunol.*, 2005, vol. 35 (4), 1283-91 [0005]
- **BARBER et al.** *Nat Med.*, 2005, vol. 11 (9), 933-5 [0005]
- **ALI et al.** *Nature*, 2004, vol. 431 (7011), 1007-11 [0005]
- **LEE et al.** *FASEB J.*, 2006, vol. 20 (3), 455-65 [0005]
- **RANDIS et al.** *Eur. J. Immunol.*, 2008, vol. 38 (5), 1215-24 [0005]
- **SAMUELS et al.** *Science*, 2004, vol. 304 (5670), 554 [0006]
- **SAMUELS et al.** *Curr. Opin. Oncol.*, 2006, vol. 18 (1), 77-82 [0006]
- **KANG et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 2005, vol. 102 (3), 802-7 [0006]
- **BADER et al.** *Proc. Natl. Acad. Sci. U.S.A.*, 2006, vol. 103 (5), 1475-9 [0006]
- **SUJOBERT et al.** *Blood*, 2005, vol. 106 (3), 1063-6 [0006]
- **BILLOTTET et al.** *Oncogene*, 2006, vol. 25 (50), 6648-59 [0006]
- **HICKEY et al.** *J. Biol. Chem.*, 2006, vol. 281 (5), 2441-50 [0006]
- **BENISTANT et al.** *Oncogene*, 2000, vol. 19 (44), 5083-90 [0006]
- **MIZOGUCHI et al.** *Brain Pathol.*, 2004, vol. 14 (4), 372-7 [0006]
- **KNOBBE et al.** *Neuropathol. Appl. Neurobiol.*, 2005, vol. 31 (5), 486-90 [0006]
- Remington's Pharmaceutical Sciences. Mack Publishing Company, 1985, 1418 [0155]
- *Journal of Pharmaceutical Science*, 1977, vol. 66, 2 [0155]
- **T. W. GREENE ; P. G. M. WUTS.** Protective Groups in Organic Synthesis. Wiley & Sons, Inc, 1999 [0158]

摘要

本發明提供了調節磷酸肌醇 3-激酶(PI3Ks)活性並有用於治療與 PI3Ks 活性相關的疾病包括，例如，炎症性疾病，基於免疫的病症，癌症，和其他疾病的嘧啶酮。