



(11)

EP 1 448 205 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
23.03.2011 Bulletin 2011/12

(51) Int Cl.:
A61K 31/56 ^(2006.01) **A61K 31/519** ^(2006.01)
A61P 1/04 ^(2006.01) **A61P 29/00** ^(2006.01)

(21) Application number: **02800923.1**

(86) International application number:
PCT/US2002/031866

(22) Date of filing: **04.10.2002**

(87) International publication number:
WO 2003/030823 (17.04.2003 Gazette 2003/16)

(54) **COMBINATIONS FOR THE TREATMENT OF IMMUNOINFLAMMATORY DISORDERS**

KOMBINATIONEN FÜR DIE BEHANDLUNG VON IMMUN-ENTZÜNDLICHEN ERKRANKUNGEN

COMBINAISONS PERMETTANT LE TRAITEMENT DES TROUBLES IMMUNOINFLAMMATOIRES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
IE IT LI LU MC NL PT SE SK TR**
Designated Extension States:
AL LT LV MK RO SI

(74) Representative: **Bösl, Raphael Konrad et al**
Isenbruck Bösl Hörschler LLP
Patentanwälte
Prinzregentenstrasse 68
81675 München (DE)

(30) Priority: **05.10.2001 US 327674 P**

(56) References cited:
WO-A-01/47572 WO-A-01/68056
WO-A-92/16226 WO-A2-00/12497
GB-A- 2 292 079 US-A- 3 934 036
US-A- 4 554 271 US-A- 5 242 921
US-A- 5 314 688 US-A- 5 468 729
US-A- 6 071 514 US-A1- 2001 016 604

(43) Date of publication of application:
25.08.2004 Bulletin 2004/35

(60) Divisional application:
10013063.2 / 2 279 742

(73) Proprietor: **Zalicus Inc.**
Cambridge, MA 02142 (US)

- **NICOLAUS B.J.R.: "SYMBIOTIC APPROACH TO DRUG DESIGN" DECISION MAKING IN DRUG RESEARCH 1983,**
- **BERGER: "Long-term toxicology effects..." SEMINARS IN ONCOLOGY, 13,1,1,1988,, 2019, pages 8-13,**
- **BERGER: "comparative carcinogenic..." ARCH. GESCHWULSTFORSCH., 1985,**
- **RYEFELDT: "Liver tumors..." TOXICOLOGIC PATHOLOGY, 1992,**
- **CONWAY: "inhibition of cartilage..." J. EXP. MED., 1995,**
- **BADGER: "disease-modifying activity of..." ARTHRITIS AND RHEUMATISM, 2000,**
- **FELDMANN: "anti-TNFalpha therapy of trheumatoid arthritis..." ANNUA. REV. IMMUNOL., 2001,**
- **INTERNET CITATION, [Online] Retrieved from the Internet: <URL:en.wikipedia.org/wiki/prednisone>**
- **INTERNET CITATION, [Online] Retrieved from the Internet: <URL:en.wikipedia/wiki/prednisolone>**

(72) Inventors:

- **KEITH, Curtis**
Boston, MA 02116 (US)
- **BORISY, Alexis**
Boston, MA 02114 (US)
- **ZIMMERMANN, Grant**
Somerville, MA 02144 (US)
- **JOST-PRICE, Edward, Roydon**
West Roxbury, MA 02132 (US)
- **MANIVASAKAM, Palaniyandi**
Brighton, MA 02135 (US)
- **HURST, Nicole**
Boston, MA 02132 (US)
- **FOLEY, Michael, A.**
Chestnut Hill, MA 02467 (US)

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 1 448 205 B1

Remarks:

The file contains technical information submitted after the application was filed and not included in this specification

Description**Field of the Invention**

5 [0001] The invention relates to the treatment of immunoinflammatory disorders.

Background of the Invention

10 [0002] Immunoinflammatory disorders (e.g., rheumatoid arthritis, psoriasis, ulcerative colitis, Crohn's disease, stroke-induced brain cell death, ankylosing spondylitis, fibromyalgia, and inflammatory dermatoses, asthma, multiple sclerosis, type I diabetes, systemic lupus erythematosus, scleroderma, systemic sclerosis, and Sjögren's syndrome) are characterized by dysregulation of the immune system and inappropriate activation of body's defenses resulting in damage to healthy tissue.

15 [0003] One percent of humans world-wide are afflicted with rheumatoid arthritis (RA), a relentless, progressive disease causing severe swelling, pain, and eventual deformity and destruction of joints. According to the Arthritis Foundation, rheumatoid arthritis currently affects over two million Americans, of which women are three times more likely to be afflicted. Rheumatoid arthritis is characterized by inflammation of the lining of the joints and/or other internal organs, and the presence of elevated numbers of lymphocytes and high levels of proinflammatory cytokines.

20 [0004] Diagnosis of RA generally includes: (i) morning stiffness in joints lasting at least one hour before improvement, (ii) arthritis of three or more joint areas having simultaneously soft tissue swelling or fluid; (iii) arthritis of at least one hand joint; (iv) symmetric arthritis, i.e., simultaneous involvement of the same joint area on both sides of the body; (v) rheumatoid nodules; (vi) abnormal serum rheumatoid factor; and (vii) radiographic changes typical of rheumatoid arthritis on posteroanterior hand and wrist radiographs, which include erosions or unequivocal bony decalcification localized in or most marked adjacent to the involved joints. Patients are classified as having RA if at least four of these seven criteria, and (i) through (iv) must have been present for at least six weeks. (American College of Rheumatology, 1987 Criteria for the Classification of Acute Arthritis of Rheumatoid Arthritis, based on Arnett FC et al., Arthritis Rheum. 1988; 31: 315-324). Pain per se is not required for the diagnosis of RA.

25 [0005] Treatment of RA generally includes anti-inflammatory strategies directed at suppressing the clinical manifestations of joint inflammation, including synovial thickening, joint tenderness, and joint stiffness. Drugs used to address these signs and symptoms generally include (i) non-steroidal anti-inflammatory drugs (NSAIDs; e.g., detoprofen, diclofenac, diflunisal, etodolac, fenoprofen, flurbiprofen, ibuprofen, indomethacin, ketoprofen, meclofenamate, mefenamic acid, meloxicam, nabumetone, naproxen sodium, oxaprozin, piroxicam, sulindac, tolmetin, celecoxib, rofecoxib, aspirin, choline salicylate, salsalate, and sodium and magnesium salicylate)--these drugs may be adequate for mild RA, but do not appear to alter the longterm course of the disease; and (ii) steroids (e.g., cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone).

30 [0006] Treatment for RA may also include strategies directed at limiting the long term joint damage and deformity caused by the inflammation in the joints. Such treatments are generally described as DMARDs, i.e., disease modifying antirheumatic drugs (e.g., cyclosporine, azathioprine, methotrexate, leflunomide, cyclophosphamide, hydroxychloroquine, sulfasalazine, D-penicillamine, minocycline, gold, etanercept (soluble TNF receptor) and infliximab (a chimeric monoclonal anti-TNF antibody)).

35 [0007] There is a need to develop new compositions for the treatment of immunoinflammatory disorders.

Summary of the Invention

40 [0008] We have discovered that the combination of dipyradamole (also known as 2,6-bis(diethanolamino)-4,8-dipiperidinopyrimido(5,4-d)pyrimidine), and corticosteroids, such as fludrocortisone (as known as 9-alpha-fluoro-11-beta, 17-alpha, 21-trihydroxy-4-pregnene-3,20-dione acetate) or prednisolone (also known as 1-dehydrocortisol; 1-dehydrohydrocortisone; 1,4-pregnadiene-11 beta,17alpha,21-triol-3,20-dione; and 11beta,17alpha,21-trihydroxy-1,4-pregnadiene-3,20-dione), brings about substantial suppression of TNF α levels induced in peripheral blood mononuclear cells (PBMCs).

45 [0009] Accordingly, the invention features a composition for treating a patient having, or at risk for developing, an immunoinflammatory disorder. The method includes administering (i) a corticosteroid; and (ii) dipyradamole. The two compounds are each to be administered in an amount that, when combined is sufficient to treat or prevent the immunoinflammatory disorder.

50 [0010] Dipyradamole and the corticosteroid may be present in pharmaceutical compositions that contain a pharmaceutically acceptable carrier, diluent, or excipient, and are administered at dosages and frequencies sufficient to suppress TNF α levels enough to produce a therapeutic benefit to the patient. Dipyradamole and the corticosteroid can be administered within 14 days of each other (e.g., within 10 days, within five days, twenty-four hours, or one hour of each other,

or even simultaneously). Administration of each compound can occur, e.g. 1 to 5 times each day, or as necessary to alleviate symptoms.

[0011] Accordingly, this invention also features pharmaceutical compositions, pharmaceutical packs, and kits containing dipyridamole and one or more corticosteroid. Compositions (pharmaceutical compositions and pharmaceutical packs) of the invention may feature higher order combinations of dipyridamole and corticosteroids. Specifically, dipyridamole may be combined with one, two, three, or more corticosteroids. In preferred embodiments, the dipyridamole, the corticosteroid, or both are approved by the United States Food and Drug Administration (USFDA) for administration to a human.

[0012] The invention described herein has been exemplified using the corticosteroids fludrocortisone (as known as 9- α -fluoro-11- β , 17- α , 21-trihydroxy-4-pregnene-3,20-dione acetate) and prednisolone (also known as 1-dehydrocortisol; 1-dehydrohydrocortisone; 1,4-pregnadiene-11 β , 17 α , 21-triol-3,20-dione; and 11 β , 17 α , 21-trihydroxy-1,4-pregnadiene-3,20-dione); however, a skilled artisan will recognize that structural and functional analogs of these corticosteroids can also be used in combination with dipyridamole in the methods and compositions of the present invention. Other useful corticosteroids may be identified based on the shared structural features and apparent mechanism of action among the corticosteroid family.

[0013] Dipyridamole and the corticosteroid may be administered in the same or different pharmaceutical formulations. Pharmaceutical compositions or components of the pharmaceutical pack may be administered by the same or different routes and include oral, rectal, intravenous, intramuscular, subcutaneous, intra-articular, inhalation, topical or transdermal, vaginal, and ophthalmic administration.

[0014] Dosages of dipyridamole and the corticosteroid may be determined individually; however, dipyridamole is typically administered to human patients at 0.5-800 mg/day, 18-600 mg/day; or 50-400 mg/day. Corticosteroids are typically administered at 0.1-1500 mg/day, 0.5-30 mg/day, or 0.5-10 mg/day. Low doses of corticosteroids (e.g., 10 mg/day or less of prednisolone, or its equivalent) are preferred. The total daily dosage of dipyridamole and the corticosteroid may be administered in one, two, three, four, or more dosages. There is no need for dipyridamole and the corticosteroid to be administered in the same number of daily doses. Further, there is no need for dipyridamole and/or the corticosteroid to be administered every day or by the same route of administration. For example, dipyridamole may be administered by intravenous injection every second day and the corticosteroid administered by topical application twice every day. Accordingly, when administered in different compositions, pharmaceutical formulations, packs, and kits are prepared in form and dosage suitable for achieving the desired treatment regimen.

[0015] The diseases or disorders treated using the methods and compositions of this invention are immunoinflammatory disorders including, for example, rheumatoid arthritis, psoriasis, ulcerative colitis, Crohn's disease, stroke-induced brain cell death, ankylosing spondylitis, fibromyalgia, multiple sclerosis, type I diabetes, systemic lupus erythematosus, , or Sjögren's syndrome.

[0016] Compounds useful in the invention include those described herein in any of their pharmaceutically acceptable forms, including diastereomers and enantiomers, salts, solvates, and polymorphs thereof, as well as racemic mixtures of the compounds described herein.

[0017] By "treating" is meant administering or prescribing a pharmaceutical composition for the treatment or prevention of an inflammatory disease.

[0018] By "patient" is meant any animal (e.g., a human).

[0019] The term "immunoinflammatory disorder" encompasses a variety of conditions, including autoimmune diseases. Immunoinflammatory disorders result in the destruction of healthy tissue by an inflammatory process. The immunoinflammatory disorders are, rheumatoid arthritis, psoriasis ulcerative colitis, Crohn's disease, stroke-induced brain cell death, ankylosing spondylitis, fibromyalgia multiple sclerosis, type I diabetes, systemic lupus erythematosus, and Sjögren's syndrome.

[0020] By "corticosteroid" is meant any naturally occurring or synthetic steroid hormone which can be derived from cholesterol and is characterized by a hydrogenated cyclopentanoperhydrophenanthrene ring system. Naturally occurring corticosteroids are generally produced by the adrenal cortex. Synthetic corticosteroids may be halogenated. Functional groups required for activity include a double bond at Δ^4 , a C3 ketone, and a C20 ketone. Corticosteroids may have glucocorticoid and/or mineralocorticoid activity. In preferred embodiments, the corticosteroid is either fludrocortisone or prednisolone.

[0021] Exemplary corticosteroids include algestone, 6- α -fluoroprednisolone, 6- α -methylprednisolone, 6- α -methylprednisolone 21-acetate, 6- α -methylprednisolone 21-hemisuccinate sodium salt, 6- α ,9- α -difluoroprednisolone 21-acetate 17-butyrate, amcinafal, beclomethasone, beclomethasone dipropionate, beclomethasone dipropionate monohydrate, 6- β -hydroxycortisol, betamethasone, betamethasone-17-valerate, budesonide, clobetasol, clobetasol propionate, clobetasone, clocortolone, clocortolone pivalate, cortisone, cortisone acetate, cortodoxone, deflazacort, 21-deoxycortisol, deprodone, descinolone, desonide, desoximetasone, dexamethasone, dexamethasone-21-acetate, dichlorisone, diflorasone, diflorasone diacetate, diflucortolone, doxibetasol, fludrocortisone, flumethasone, flumethasone pivalate, flumoxonide, flunisolide, fluocinonide, fluocinolone acetonide, 9-fluorocortisone, fluorohydroxy-

androstenedione, fluorometholone, fluorometholone acetate, fluoxymesterone, flupredidene, fluprednisolone, flurandrenolide, formocortal, halcinonide, halometasone, halopredone, hyrcanoside, hydrocortisone, hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydrocortisone valerate, 6-hydroxydexamethasone, isoflupredone, isoflupredone acetate, isoprednidene, meclorisone, methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, paramethasone, paramethasone acetate, prednisolone, prednisolone acetate, prednisolone metasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hemisuccinate free acid, prednisolone-21-acetate, prednisolone-21(beta-D-glucuronide), prednisone, prednylidene, procinonide, tralonide, triamcinolone, triamcinolone acetonide, triamcinolone acetonide 21-palmitate, triamcinolone diacetate, triamcinolone hexacetonide, and wortmannin.

Desirably, the corticosteroid is fludrocortisone or prednisolone.

[0022] By "an effective amount" is meant the amount of a compound, in a combination of the invention, required to treat or prevent an immunoinflammatory disease. The effective amount of active compound(s) used to practice the present invention for therapeutic treatment of conditions caused by or contributing to an inflammatory disease varies depending upon the manner of administration, the age, body weight, and general health of the patient. Ultimately, the attending physician or veterinarian will decide the appropriate amount and dosage regimen. Such amount is referred to as an effective amount.

[0023] By the term "cytokine suppressing amount" is meant an amount of the combination which will cause a decrease in the *in vivo* presence or level of the proinflammatory cytokine, when given to a patient for the prophylaxis or therapeutic treatment of an immunoinflammatory disorder which is exacerbated or caused by excessive or unregulated proinflammatory cytokine production.

Detailed Description

[0024] We have discovered that the combination of dipyridamole with a corticosteroid substantially has substantial TNF α suppressing activity against stimulated white blood cells. The combinations of dipyridamole with fludrocortisone, and dipyridamole with prednisolone were particularly effective. Thus, the combination of dipyridamole with a corticosteroid is useful for the treatment of immunoinflammatory disorders.

Dipyridamole

[0025] Dipyridamole (2,6-bis(diethanolamino)-4,8-dipiperidinopyrimido(5,4-d)pyrimidine) is a tetra-substituted pyrimidopyrimidine that is used as a platelet inhibitor, e.g., to prevent blood clot formation following heart valve surgery and to reduced the moribundity associated with clotting disorders, including myocardial and cerebral infarction. Typically, anticoagulation therapy (prophylaxis or treatment) is effected by administering dipyridamole at about 75-200 mg b.i.d, t.i.d., or q.i.d. either alone or in combination with aspirin.

[0026] Combination therapy according to the invention may be performed alone or in conjunction with another therapy and may be provided at home, the doctor's office, a clinic, a hospital's outpatient department, or a hospital. Treatment generally begins at a hospital so that the doctor can observe the therapy's effects closely and make any adjustments that are needed. The duration of the combination therapy depends on the type of immunoinflammatory disorder being treated, the age and condition of the patient, the stage and type of the patient's disease, and how the patient responds to the treatment. Additionally, a person having a greater risk of developing an immunoinflammatory disorder (e.g., a person who is genetically predisposed or previously had an immunoinflammatory disorder) may receive prophylactic treatment to inhibit or delay an inflammatory response.

[0027] The dosage, frequency and mode of administration of each component of the combination can be controlled independently. For example, one compound may be administered orally three times per day, while the second compound may be administered intramuscularly once per day. Combination therapy may be given in on-and-off cycles that include rest periods so that the patient's body has a chance to recovery from any as yet unforeseen side-effects. The compounds may also be formulated together such that one administration delivers both compounds.

Formulation of pharmaceutical compositions

[0028] The administration of each compound of the combination may be by any suitable means that results in a concentration of the compound that, combined with the other component, is antiinflammatory upon reaching the target region. The compound may be contained in any appropriate amount in any suitable carrier substance, and is generally present in an amount of 1-95% by weight of the total weight of the composition. The composition may be provided in a dosage form that is suitable for the oral, parenteral (e.g., intravenously, intramuscularly), rectal, cutaneous, nasal, vaginal, inhalant, skin (patch), or ocular administration route. Thus, the composition may be in the form of, e.g., tablets, capsules, pills, powders, granulates, suspensions, emulsions, solutions, gels including hydrogels, pastes, ointments, creams,

plasters, drenches, osmotic delivery devices, suppositories, enemas, injectables, implants, sprays, or aerosols. The pharmaceutical compositions may be formulated according to conventional pharmaceutical practice (see, e.g., Remington: The Science and Practice of Pharmacy (20th ed.), ed. A.R. Gennaro, Lippincott Williams & Wilkins, 2000 and Encyclopedia of Pharmaceutical Technology, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York).

[0029] Pharmaceutical compositions according to the invention may be formulated to release the active compound substantially immediately upon administration or at any predetermined time or time period after administration. The latter types of compositions are generally known as controlled release formulations, which include (i) formulations that create a substantially constant concentration of the drug within the body over an extended period of time; (ii) formulations that after a predetermined lag time create a substantially constant concentration of the drug within the body over an extended period of time; (iii) formulations that sustain drug action during a predetermined time period by maintaining a relatively, constant, effective drug level in the body with concomitant minimization of undesirable side effects associated with fluctuations in the plasma level of the active drug substance (sawtooth kinetic pattern); (iv) formulations that localize drug action by, e.g., spatial placement of a controlled release composition adjacent to or in the diseased tissue or organ; and (v) formulations that target drug action by using carriers or chemical derivatives to deliver the drug to a particular target cell type.

[0030] Administration of compounds in the form of a controlled release formulation is especially preferred in cases in which the compound, either alone or in combination, has (i) a narrow therapeutic index (i.e., the difference between the plasma concentration leading to harmful side effects or toxic reactions and the plasma concentration leading to a therapeutic effect is small; in general, the therapeutic index, TI, is defined as the ratio of median lethal dose (LD50) to median effective dose (ED50)); (ii) a narrow absorption window in the gastrointestinal tract; or (iii) a very short biological half-life so that frequent dosing during a day is required in order to sustain the plasma level at a therapeutic level.

[0031] Any of a number of strategies can be pursued in order to obtain controlled release in which the rate of release outweighs the rate of metabolism of the compound in question. In one example, controlled release is obtained by appropriate selection of various formulation parameters and ingredients, including, e.g., various types of controlled release compositions and coatings. Thus, the drug is formulated with appropriate excipients into a pharmaceutical composition that, upon administration, releases the drug in a controlled manner. Examples include single or multiple unit tablet or capsule compositions, oil solutions, suspensions, emulsions, microcapsules, microspheres, nanoparticles, patches, and liposomes.

Solid dosage forms for oral use

[0032] Formulations for oral use include tablets containing the active ingredient(s) in a mixture with non-toxic pharmaceutically acceptable excipients. These excipients may be, for example, inert diluents or fillers (e.g., sucrose, sorbitol, sugar, mannitol, microcrystalline cellulose, starches including potato starch, calcium carbonate, sodium chloride, lactose, calcium phosphate, calcium sulfate, or sodium phosphate); granulating and disintegrating agents (e.g., cellulose derivatives including microcrystalline cellulose, starches including potato starch, croscarmellose sodium, alginates, or alginic acid); binding agents (e.g., sucrose, glucose, sorbitol, acacia, alginic acid, sodium alginate, gelatin, starch, pregelatinized starch, microcrystalline cellulose, magnesium aluminum silicate, carboxymethylcellulose sodium, methylcellulose, hydroxypropyl methylcellulose, ethylcellulose, polyvinylpyrrolidone, or polyethylene glycol); and lubricating agents, glidants, and antiadhesives (e.g., magnesium stearate, zinc stearate, stearic acid, silicas, hydrogenated vegetable oils, or talc). Other pharmaceutically acceptable excipients can be colorants, flavoring agents, plasticizers, humectants, buffering agents.

[0033] The tablets may be uncoated or they may be coated by known techniques, optionally to delay disintegration and absorption in the gastrointestinal tract and thereby providing a sustained action over a longer period. The coating may be adapted to release the active drug substance in a predetermined pattern (e.g., in order to achieve a controlled release formulation) or it may be adapted not to release the active drug substance until after passage of the stomach (enteric coating). The coating may be a sugar coating, a film coating (e.g., based on hydroxypropyl methylcellulose, methylcellulose, methyl hydroxyethylcellulose, hydroxypropylcellulose, carboxymethylcellulose, acrylate copolymers, polyethylene glycols and/or polyvinylpyrrolidone), or an enteric coating (e.g., based on methacrylic acid copolymer, cellulose acetate phthalate, hydroxypropyl methylcellulose phthalate, hydroxypropyl methylcellulose acetate succinate, polyvinyl acetate phthalate, shellac, and/or ethylcellulose). Furthermore, a time delay material such as, e.g., glyceryl monostearate or glyceryl distearate may be employed.

[0034] The solid tablet compositions may include a coating adapted to protect the composition from unwanted chemical changes, (e.g., chemical degradation prior to the release of the active drug substance). The coating may be applied on the solid dosage form in a similar manner as that described in Encyclopedia of Pharmaceutical Technology, *supra*.

[0035] The two drugs may be mixed together in the tablet, or may be partitioned. In one example, the first drug is contained on the inside of the tablet, and the second drug is on the outside, such that a substantial portion of the second drug is released prior to the release of the first drug.

[0036] Formulations for oral use may also be presented as chewable tablets, or as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent (e.g., potato starch, lactose, microcrystalline cellulose, calcium carbonate, calcium phosphate or kaolin), or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example, peanut oil, liquid paraffin, or olive oil. Powders and granulates may be prepared using the ingredients mentioned above under tablets and capsules in a conventional manner using, e.g., a mixer, a fluid bed apparatus or a spray drying equipment.

Controlled release oral dosage forms

[0037] Controlled release compositions for oral use may, e.g., be constructed to release the active drug by controlling the dissolution and/or the diffusion of the active drug substance.

[0038] Dissolution or diffusion controlled release can be achieved by appropriate coating of a tablet, capsule, pellet, or granulate formulation of compounds, or by incorporating the compound into an appropriate matrix. A controlled release coating may include one or more of the coating substances mentioned above and/or, e.g., shellac, beeswax, glycowax, castor wax, carnauba wax, stearyl alcohol, glyceryl monostearate, glyceryl distearate, glycerol palmitostearate, ethylcellulose, acrylic resins, dl-poly(lactic acid), cellulose acetate butyrate, polyvinyl chloride, polyvinyl acetate, vinyl pyrrolidone, polyethylene, polymethacrylate, methylmethacrylate, 2-hydroxymethacrylate, methacrylate hydrogels, 1,3 butylene glycol, ethylene glycol methacrylate, and/or polyethylene glycols. In a controlled release matrix formulation, the matrix material may also include, e.g., hydrated methylcellulose, carnauba wax and stearyl alcohol, carbopol 934, silicone, glyceryl tristearate, methyl acrylate-methyl methacrylate, polyvinyl chloride, polyethylene, and/or halogenated fluorocarbon.

[0039] A controlled release composition containing one or more of the compounds of the claimed combinations may also be in the form of a buoyant tablet or capsule (i.e., a tablet or capsule that, upon oral administration, floats on top of the gastric content for a certain period of time). A buoyant tablet formulation of the compound(s) can be prepared by granulating a mixture of the drug(s) with excipients and 20-75% w/w of hydrocolloids, such as hydroxyethylcellulose, hydroxypropylcellulose, or hydroxypropylmethylcellulose. The obtained granules can then be compressed into tablets. On contact with the gastric juice, the tablet forms a substantially water-impermeable gel barrier around its surface. This gel barrier takes part in maintaining a density of less than one, thereby allowing the tablet to remain buoyant in the gastric juice.

Liquids for oral administration

[0040] Powders, dispersible powders, or granules suitable for preparation of an aqueous suspension by addition of water are convenient dosage forms for oral administration. Formulation as a suspension provides the active ingredient in a mixture with a dispersing or wetting agent, suspending agent, and one or more preservatives. Suitable dispersing or wetting agents are, for example, naturally-occurring phosphatides (e.g., lecithin or condensation products of ethylene oxide with a fatty acid, a long chain aliphatic alcohol, or a partial ester derived from fatty acids) and a hexitol or a hexitol anhydride (e.g., polyoxyethylene stearate, polyoxyethylene sorbitol monooleate, polyoxyethylene sorbitan monooleate, and the like). Suitable suspending agents are, for example, sodium carboxymethylcellulose, methylcellulose, sodium alginate.

Parenteral compositions

[0041] The pharmaceutical composition for use in the methods of treatment as in the claims may also be administered parenterally by injection, infusion or implantation (intravenous, intramuscular, subcutaneous, or the like) in dosage forms, formulations, or via suitable delivery devices or implants containing conventional, non-toxic pharmaceutically acceptable carriers and adjuvants. The formulation and preparation of such compositions are well known to those skilled in the art of pharmaceutical formulation. Formulations can be found in Remington: The Science and Practice of Pharmacy, *supra*.

[0042] Compositions for parenteral use may be provided in unit dosage forms (e.g., in single-dose ampoules), or in vials containing several doses and in which a suitable preservative may be added (see below). The composition may be in form of a solution, a suspension, an emulsion, an infusion device, or a delivery device for implantation, or it may be presented as a dry powder to be reconstituted with water or another suitable vehicle before use. Apart from the active drug(s), the composition may include suitable parenterally acceptable carriers and/or excipients. The active drug(s) may be incorporated into microspheres, microcapsules, nanoparticles, liposomes, or the like for controlled release. Furthermore, the composition may include suspending, solubilizing, stabilizing, pH-adjusting agents, and/or dispersing agents.

[0043] As indicated above, the pharmaceutical compositions according to the invention may be in the form suitable for sterile injection. To prepare such a composition, the suitable active drug(s) are dissolved or suspended in a parenterally acceptable liquid vehicle. Among acceptable vehicles and solvents that may be employed are water, water adjusted to

a suitable pH by addition of an appropriate amount of hydrochloric acid, sodium hydroxide or a suitable buffer, 1,3-butanediol, Ringer's solution, and isotonic sodium chloride solution. The aqueous formulation may also contain, one or more preservatives (e.g., methyl, ethyl or n-propyl p-hydroxybenzoate). In cases where one of the compounds is only sparingly or slightly soluble in water, a dissolution enhancing or solubilizing agent can be added, or the solvent may include 10-60% w/w of propylene glycol.

Controlled release parenteral compositions

[0044] Controlled release parenteral compositions for use according to claims 1-11 may be in form of aqueous suspensions, microspheres, microcapsules, magnetic microspheres, oil solutions, oil suspensions, or emulsions. Alternatively, the active drug(s) may be incorporated in biocompatible carriers, liposomes, nanoparticles, implants, or infusion devices.

[0045] Materials for use in the preparation of microspheres and/or microcapsules are, e.g., biodegradable/bioerodible polymers such as polygalactin, poly-(isobutyl - cyanoacrylate), poly(2-hydroxyethyl-L-glutamine) and, poly(lactic acid). Biocompatible carriers that may be used when formulating a controlled release parenteral formulation are carbohydrates (e.g., dextrans), proteins (e.g., albumin), lipoproteins, or antibodies. Materials for use in implants can be non-biodegradable (e.g., polydimethyl siloxane) or biodegradable (e.g., poly(caprolactone), poly(lactic acid), poly(glycolic acid) or poly(ortho esters)).

Rectal compositions

[0046] For rectal application, suitable dosage forms for a composition for use according to claims 1-11 include suppositories (emulsion or suspension type), and rectal gelatin capsules (solutions or suspensions). In a typical suppository formulation, the active drug(s) are combined with an appropriate pharmaceutically acceptable suppository base such as cocoa butter, esterified fatty acids, glycerinated gelatin, and various water-soluble or dispersible bases like polyethylene glycols and polyoxyethylene sorbitan fatty acid esters. Various additives, enhancers, or surfactants may be incorporated.

Compositions for inhalation

[0047] For administration by inhalation, typical dosage forms for use according to claims 1-11 include nasal sprays and aerosols. In a typically nasal formulation, the active ingredient(s) are dissolved or dispersed in a suitable vehicle. The pharmaceutically acceptable vehicles and excipients (as well as other pharmaceutically acceptable materials present in the composition such as diluents, enhancers, flavoring agents, and preservatives) are selected in accordance with conventional pharmaceutical practice in a manner understood by the persons skilled in the art of formulating pharmaceuticals.

Percutaneous and topical compositions

[0048] The pharmaceutical compositions for use according to claims 1-11 may also be administered topically on the skin for percutaneous absorption in dosage forms or formulations containing conventionally non-toxic pharmaceutical acceptable carriers and excipients including microspheres and liposomes. The formulations for use according to claims 1-11 include creams, ointments, lotions, liniments, gels, hydrogels, solutions, suspensions, sticks, sprays, pastes, plasters, and other kinds of transdermal drug delivery systems. The pharmaceutically acceptable carriers or excipients may include emulsifying agents, antioxidants, buffering agents, preservatives, humectants, penetration enhancers, chelating agents, gel-forming agents, ointment bases, perfumes, and skin protective agents.

[0049] Examples of emulsifying agents are naturally occurring gums (e.g., gum acacia or gum tragacanth) and naturally occurring phosphatides (e.g., soybean lecithin and sorbitan monooleate derivatives). Examples of antioxidants are butylated hydroxy anisole (BHA), ascorbic acid and derivatives thereof, tocopherol and derivatives thereof, butylated hydroxy anisole, and cysteine. Examples of preservatives are parabens, such as methyl or propyl p-hydroxybenzoate, and benzalkonium chloride. Examples of humectants are glycerin, propylene glycol, sorbitol, and urea. Examples of penetration enhancers are propylene glycol, DMSO, triethanolamine, N,N-dimethylacetamide, N,N-dimethylformamide, 2-pyrrolidone and derivatives thereof, tetrahydrofurfuryl alcohol, and AZONE™. Examples of chelating agents are sodium EDTA, citric acid, and phosphoric acid. Examples of gel forming agents are CARBOPOL™, cellulose derivatives, bentonite, alginates, gelatin and polyvinylpyrrolidone. Examples of ointment bases are beeswax, paraffin, cetyl palmitate, vegetable oils, sorbitan esters of fatty acids (Span), polyethylene glycols, and condensation products between sorbitan esters of fatty acids and ethylene oxide (e.g., polyoxyethylene sorbitan monooleate (TWEEN™)).

[0050] The pharmaceutical composition described above for use according to claims 1-11 for topical administration

on the skin may also be used in connection with topical administration onto or close to the part of the body that is to be treated. The compositions for use according to claims 1-11 may be adapted for direct application or for introduction into relevant orifice(s) of the body (e.g., rectal, urethral, vaginal or oral orifices). The composition for use according to claims 1-11 may be applied by means of special drug delivery devices such as dressings or alternatively plasters, pads, sponges, strips, or other forms of suitable flexible material.

Controlled release percutaneous and topical compositions for use according to claims 1-11

[0051] There are several approaches for providing rate control over the release and transdermal permeation of a drug, for use according to claims 1-11 including: membrane-moderated systems, adhesive diffusion-controlled systems, matrix dispersion-type systems, and microreservoir systems. A controlled release percutaneous and/or topical composition may be obtained by using a suitable mixture of the above-mentioned approaches.

[0052] In a membrane-moderated system, the active drug is present in a reservoir which is totally encapsulated in a shallow compartment molded from a drug-impermeable laminate, such as a metallic plastic laminate, and a rate-controlling polymeric membrane such as a microporous or a non-porous polymeric membrane (e.g., ethylene-vinyl acetate copolymer). The active compound is only released through the rate-controlling polymeric membrane. In the drug reservoir, the active drug substance may either be dispersed in a solid polymer matrix or suspended in a viscous liquid medium such as silicone fluid. On the external surface of the polymeric membrane, a thin layer of an adhesive polymer is applied to achieve an intimate contact of the transdermal system with the skin surface. The adhesive polymer is preferably a hypoallergenic polymer that is compatible with the active drug.

[0053] In an adhesive diffusion-controlled system, a reservoir of the active drug is formed by directly dispersing the active drug in an adhesive polymer and then spreading the adhesive containing the active drug onto a flat sheet of substantially drug-impermeable metallic plastic backing to form a thin drug reservoir layer. A matrix dispersion-type system is characterized in that a reservoir of the active drug substance is formed by substantially homogeneously dispersing the active drug substance in a hydrophilic or lipophilic polymer matrix and then molding the drug-containing polymer into a disc with a substantially well-defined surface area and thickness. The adhesive polymer is spread along the circumference to form a strip of adhesive around the disc.

[0054] In a microreservoir system, the reservoir of the active substance is formed by first suspending the drug solids in an aqueous solution of water-soluble polymer, and then dispersing the drug suspension in a lipophilic polymer to form a plurality of microscopic spheres of drug reservoirs.

Dosages

[0055] The dosage of each compound of the claimed combinations depends on several factors, including: the administration method, the condition to be treated, the severity of the condition, whether the condition is to be treated or prevented, and the age, weight, and health of the person to be treated. Additionally, pharmacogenomic (the effect of genotype on the pharmacokinetic, pharmacodynamic or efficacy profile of a therapeutic) information about a particular patient may affect dosage used.

[0056] As described above, the compound in question may be administered orally in the form of tablets, capsules, elixirs or syrups, or rectally in the form of suppositories. Parenteral administration of a compound is suitably performed, for example, in the form of saline solutions or with the compound incorporated into liposomes. In cases where the compound in itself is not sufficiently soluble to be dissolved, a solubilizer such as ethanol can be applied. Below, for illustrative purposes, the dosages for dipyridamole and fludrocortisone are described.

Routes of administration

[0057] For oral, intramuscular, subcutaneous, topical, inhalation, rectal, vaginal and ophthalmic administration of the tetra-substituted pyrimidopyrimidine, the dosage is normally 0.5-800 mg/day, preferably 18-600 mg/day, and more preferably 50-400 mg/day. Administration can be one to four times daily for one day to one year, and may even be for the life of the patient. Chronic, long-term administration will be indicated in many cases. Dosages up to 1600 mg/day may be necessary. For intravenous administration of the tetra-substituted pyrimidopyrimidine, the dosage is normally 0.1-200 mg/day, preferably 0.5-150 mg/day, and more preferably 1-100 mg/day. Administration can be one to four times daily. Systemic dosing will result in steady-state plasma concentrations preferably of 0.1-7.0 μ M, more preferably, 0.5-5.0 μ M, and most preferably, 1.0-2.0 μ M.

[0058] The dosage of the corticosteroid for use in combination with the tetra-substituted pyrimidopyrimidine is normally 0.1-1500 mg/day, preferably 0.5-30 mg/day, and more preferably 0.5-10 mg/day. Administration can be one to four times daily for one day to one year, and may even be for the life of the patient. Chronic, long-term administration will be indicated in many cases. Dosages up to 3000 mg/day may be necessary.

[0059] The following examples are to illustrate the invention.

Example 1: Preparation of pairwise compound mixed combination serial dilution matrix

[0060] Stock solutions at 16 mg/ml of dipyridamole, and 1.6 mg/ml of fludrocortisone acetate (Sigma-Aldrich, St. Louis, MO; catalog numbers D9766 and F6127, respectively) were made in dimethylsulfoxide (DMSO). The dipyridamole master plates were made by adding 25 μ l of the concentrated stock solution to columns 3, 9, and 15 (rows C through N) of a polypropylene 384-well storage plate that had been pre-filled with 75 μ l of anhydrous DMSO. Using a TomTec Quadra Plus liquid handler, the 25 μ l of dipyridamole stock solution was serially diluted four times into the adjacent columns (columns 4-7, 10-13, 16-19). The sixth column (8, 14, and 20) did not receive any compound and served as a vehicle control. The fludrocortisone master plates were made by adding 25 μ l of the concentrated stock solution to the appropriate wells (row C, columns 3-8; row C, columns 9-14; row C, columns 15-20; row I, columns 3-8; row I, columns 9-14; row I, columns 15-20) of the appropriate master polypropylene 384-well storage plate. These master plates had been pre-filled with 75 μ l of anhydrous DMSO. Using the TomTec Quadra Plus liquid handler, the 25 μ l was serially diluted four times in the adjacent rows (rows D-G, and J-M). The sixth row (H and N) did not receive any compound to serve as a vehicle control. Master plates were sealed and stored at -20 C until ready for use.

[0061] The final dipyridamole/fludrocortisone combination plates were generated by transferring 1 μ l from each of the dipyridamole and fludrocortisone master plates to a dilution plate containing 100 μ l of media (RPMI; Gibco BRL, #11875-085), 10% Fetal Bovine Serum (Gibco BRL, #25140-097), 2% Penicillin/Streptomycin (Gibco BRL, #15140-122)) using the TomTec Quadra Plus liquid handler. This dilution plate was then mixed and a 10 μ l aliquot transferred to the final assay plate, which had been pre-filled with 40 μ l/well RPMI media containing the appropriate stimulant to activate TNF α secretion (see below).

Example 2: Assay for TNF α suppressing activity by the combination of dipyridamole and fludrocortisone

[0062] The compound dilution matrix was assayed using a TNF α ELISA method. Briefly, a 100 μ l suspension of diluted human white blood cells contained within each well of a polystyrene 384-well plate (NalgeNunc) was stimulated to secrete TNF α by treatment with a final concentration of 10 ng/ml phorbol 12-myristate 13-acetate (Sigma) and 750 ng/ μ l ionomycin (Sigma). Various concentrations of each test compound were added at the time of stimulation. After 16-18 hours of incubation at 37°C in a humidified incubator, the plate was centrifuged and the supernatant transferred to a white opaque polystyrene 384 well plate (NalgeNunc, Maxisorb) coated with an anti-TNF antibody (PharMingen, #18631D). After a two-hour incubation, the plate was washed (Tecan PowerWasher 384) with phosphate buffered saline (PBS) containing 0.1% Tween 20 (polyoxyethylene sorbitan monolaurate) and incubated for an additional one hour with another anti-TNF antibody that was biotin labeled (PharMingen, 18642D) and horseradish peroxidase (HRP) coupled to strepavidin (PharMingen, #13047E). After the plate was washed with 0.1% Tween 20/PBS, the HRP substrate (which contains luminol, hydrogen peroxide, and an enhancer such as para-iodophenol) was added to each well and light intensity measured using a LJL Analyst luminometer. Control wells contained a final concentration of 1 μ g/ml cyclosporin A (Sigma).

[0063] Together, dipyridamole and fludrocortisone were able to suppress TNF α secretion in blood stimulated with phorbol 12-myristate 13-acetate and ionomycin. As seen in Tables 1 and 2, dipyridamole was able to enhance the potency of fludrocortisone by 60-fold. At a concentration of 947 nM, fludrocortisone alone inhibited TNF α secretion by 39%. Addition of 124 nM dipyridamole to a concentration of only 15 nM fludrocortisone resulted in the inhibition of TNF α secretion by 39% (Table 1). Efficacy was maintained while reducing the total drug species by over 80%, from 947 nM to 163 nM. In the presence of 2 μ M dipyridamole, 50% TNF α inhibition is achieved by only 4 nM fludrocortisone. This level of inhibition is not possible with fludrocortisone alone at concentrations that would be expected to cause serious mineralocorticoid-induced side effects. Dipyridamole enhancement of fludrocortisone activity was observed in a secondary screen (Table 2). Again, a low dose of 495 nM dipyridamole enhanced the potency of fludrocortisone by over 135 fold. Specifically, 947 nM fludrocortisone alone was required to achieve a 52% reduction of TNF α secretion. A similar reduction (49%) was measured for the combination of 7 nM fludrocortisone and 495 nM dipyridamole. Further, the addition of 248 nM dipyridamole resulted in a supramaximal effect on the inhibition of TNF α secretion at fludrocortisone concentrations as low as 59 nM.

TABLE 1
Primary Screen Data of Fludrocortisone vs Dipyridamole Average Result of 2 Plates
 (%TNF α suppression from P/I-induced white blood cells)

Fludrocortisone [μ M]	Dipyridamole [μ M]					
	7.927	1.982	0.495	0.124	0.031	0.000
0.947	82.90	66.61	54.90	52.48	61.35	39.19
0.237	81.88	61.99	52.35	52.11	46.77	36.66
0.059	79.57	60.37	47.08	45.47	42.93	32.49
0.015	77.13	54.06	40.70	38.73	30.62	22.63
0.004	74.61	50.60	34.21	24.90	22.52	17.21
0.000	66.37	35.24	13.21	9.08	3.68	0.00

TABLE 2
Secondary Screen Data of Fludrocortisone vs Dipyridamole Average Result of 2 Plates
 (%TNF α Suppression from P/I-induced white blood cells)

Fludrocortisone [μ M]	Dipyridamole [μ M]									
	7.927	3.964	1.982	0.991	0.495	0.248	0.124	0.062	0.031	0.000
0.947	89.12	82.25	78.01	69.10	67.91	61.77	60.82	53.38	53.41	52.05
0.473	92.64	84.40	78.44	70.25	65.06	60.25	56.14	53.68	50.07	50.16
0.237	89.69	83.68	82.01	70.36	66.53	65.46	60.90	56.65	52.34	49.25
0.118	87.58	80.66	76.13	68.18	65.89	67.09	58.91	54.17	47.39	46.42
0.059	84.43	79.37	73.86	64.53	63.88	58.96	56.84	48.63	44.66	40.24
0.030	88.61	76.42	68.58	65.77	62.08	51.31	49.96	47.02	44.19	36.95
0.015	90.46	79.36	73.52	65.22	56.39	62.88	43.17	47.73	46.00	37.77
0.007	84.11	75.29	69.74	64.61	48.90	42.05	38.92	39.27	36.70	29.49
0.004	79.02	75.15	65.79	55.19	46.00	41.93	35.15	30.94	30.20	22.40
0.000	78.11	66.54	62.36	48.20	33.73	23.02	12.13	9.43	10.16	-3.50

Example 3: Preparation of pairwise compound mixed combination serial dilution matrix

[0064] A compound matrix of dipyridamole and prednisolone were prepared according to the method of Example 1. The initial stock solution of dipyridamole, was 16 mg/ml, and prednisolone was 1.6 mg/ml.

Example 4: Assay for TNF α suppressing activity by the combination of dipyridamole and prednisolone

[0065] The compound dilution matrix of Example 3, was assayed using the TNF α ELISA method of Example 2. The results are shown in Table 3. Together, dipyridamole and prednisolone were able to suppress TNF α secretion in blood stimulated with phorbol 12-myristate 13-acetate and ionomycin to a greater extent than either compound alone. Specifically, dipyridamole greatly increased the potency of prednisolone. Prednisolone alone, at a concentration of 250 nM, can suppress TNF α secretion by 38%. The same level of suppression (41%) can be achieved by only 1 nM prednisolone in combination with 2 μ M dipyridamole. This represents a shift in the potency of prednisolone of over 250-fold. Further, the addition of 2 μ M dipyridamole to 250 nM prednisolone resulted in a supramaximal effect (57%), compared to prednisolone alone (38%). The combination of low doses of prednisolone and dipyridamole, therefore, results in the inhibition TNF α to levels previously unattainable without a high risk of glucocorticoid-induced side effects.

TABLE 3
Primary Screen Data of Prednisolone vs Dipyridamole Average Result of 2 Plates
(%TNF α suppression from P/I-induced white blood cells)

	Prednisolone [μ M]	Dipyridamole [μ M]					
		7.93	1.98	0.50	0.12	0.031	0.00
	0.250	70.30	56.72	48.90	50.82	46.08	38.25
	0.063	68.53	57.68	51.61	47.24	37.57	33.00
	0.016	66.48	45.20	40.12	40.99	32.42	37.84
	0.004	61.06	47.25	34.66	33.48	32.42	19.99
	0.001	57.35	40.84	32.10	25.47	20.64	4.86
	0.000	47.51	27.21	18.30	12.63	11.24	0.00

Claims

1. A medicament comprising

- (a) a corticosteroid; and
- (b) dipyridamole

for use in a method of treating a patient who has an immunoinflammatory disorder or is at risk for developing an immunoinflammatory disorder, wherein said immunoinflammatory disorder is selected from the group consisting of rheumatoid arthritis, psoriasis, ulcerative colitis, Crohn's disease, ankylosing spondylitis, fibromyalgia, multiple sclerosis, type I diabetes, systemic lupus erythematosus, Sjögren's syndrome, and stroke induced brain cell death, wherein said medicament comprises said dipyridamole and said corticosteroid for simultaneous administration or for administration within 14 days of each other, in amounts and for a duration sufficient to treat said patient.

2. The medicament for use according to claim 1, wherein said corticosteroid is algestone, 6-alpha-fluoroprednisolone, 6-alpha-methylprednisolone, 6-alpha-methylprednisolone 21-acetate, 6-alpha-methylprednisolone 21-hemisuccinate sodium salt, 6-alpha,9-alpha-difluoroprednisolone 21-acetate 17-butyrate, amcinafal, beclomethasone, beclomethasone dipropionate, beclomethasone dipropionate monohydrate, 6-betahydroxycortisol, betamethasone, betamethasone-17-valerate, budesonide, clobetasol, clobetasol propionate, clobetasone, clocortolone, clocortolone pivalate, cortisone, cortisone acetate, cortodoxone, deflazacort, 21-deoxycortisol, deprodone, descinolone, desonide, desoximethasone, dexamethasone, dexamethasone-21-acetate, dichlorisone, diflorasone, diflorasone diacetate, diflucortolone, doxibetasol, fludrocortisone, flumethasone, flumethasone pivalate, flumoxonide, flunisolide, fluocinonide, fluocinolone acetonide, 9-fluorocortisone, fluorohydroxyandrostenedione, fluorometholone, fluorometholone acetate, fluoxymesterone, flupredidene, fluprednisolone, flurandrenolide, formocortal, halcinonide, halometasone, halopredone, hyrcanoside, hydrocortisone, hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydrocortisone valerate, 6-hydroxydexamethasone, isoflupredone, isoflupredone acetate, isoprednidene, meclorisonone, methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, paramethasone, paramethasone acetate, prednisolone, prednisolone acetate, prednisolone metasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hemisuccinate free acid, prednisolone-21-acetate, prednisolone-21(beta-D-glucuronide), prednisone, prednylidene, procinonide, tralonide, triamcinolone, triamcinolone acetonide, triamcinolone acetonide 21-palmitate, triamcinolone diacetate, triamcinolone hexacetone, or wortmannin.

3. The medicament for use according to claim 2, wherein said corticosteroid is fludrocortisone or prednisolone.

4. The medicament for use according to claim 3, wherein said corticosteroid is prednisolone or a pharmaceutically acceptable salt thereof.

5. The medicament for use according to claim 1 or 3, or 4, wherein said medicament comprises dipyridamole and said corticosteroid for administration within ten days of each other, for administration within five days of each other, for administration within twenty-four hours of each other, within one hour of each other, or for simultaneous administration of said dipyridamole and said corticosteroid.

6. The medicament for use according to claim 1, 3 or 4, wherein said medicament comprises said dipyridamole and

said corticosteroid for administration in the same pharmaceutical formulation.

7. The medicament for use according to claim 1, or 4, 5 or 6, wherein said immunoinflammatory disorder is rheumatoid arthritis, psoriasis, ulcerative colitis, Crohn's disease, ankylosing spondylitis, or multiple sclerosis.
8. The medicament for use according to claim 1, 4, 5, 6 or 7, wherein said medicament comprises said dipyridamole in an amount of 0.5-800 mg/day and comprises said corticosteroid in an amount of 0.1-1500 mg/day, wherein said medicament comprises said dipyridamole in an amount of 18-600 mg/day and comprises said corticosteroid in an amount of 0.5-30 mg/day, or wherein said medicament comprises said corticosteroid in an amount of 0.5-10 mg/day.
9. The medicament for use according to claim 1, 3, 4, or 7, wherein said medicament comprises a daily dosage of 18 to 600 mg dipyridamole or a daily dosage of 50 to 400 mg dipyridamole.
10. The medicament for use according to claim 1, 3, 4, or 7, wherein said medicament comprises a daily dosage of 0.5 to 10 mg of fludrocortisone or prednisolone.
11. The medicament for use according to claim 1, 3, or 4, wherein said medicament is formulated for oral or intravenous administration.
12. A medicament consisting of active ingredients and excipients, wherein said active ingredients consist of a corticosteroid and dipyridamole and wherein said medicament is formulated for oral administration.
13. The medicament of claim 12, wherein said corticosteroid is algestone, 6-alpha-fluoroprednisolone, 6-alpha-methylprednisolone, 6-alpha-methylprednisolone 21-acetate, 6-alpha-methylprednisolone 21-hemisuccinate sodium salt, 6-alpha,9-alpha-difluoroprednisolone 21-acetate 17-butyrate, amcinafal, beclomethasone, beclomethasone dipropionate, beclomethasone dipropionate monohydrate, 6-beta-hydroxycortisol, betamethasone, betamethasone-17-valerate, budesonide, clobetasol, clobetasol propionate, clobetasone, clocortolone, clocortolone pivalate, cortisone, cortisone acetate, cortodoxone, deflazacort, 21-deoxycortisol, deprodone, descinolone, desonide, desoximethasone, dexamethasone, dexamethasone-21-acetate, dichlorisone, diflorasone, diflorasone diacetate, diflucortolone, doxibetasol, fludrocortisone, flumethasone, flumethasone pivalate, flumoxonide, flurusolide, fluocinonide, fluocinolone acetonide, 9-fluorocortisone, fluorohydroxyandrostenedione, fluorometholone, fluorometholone acetate, fluoxymesterone, flupredidene, fluprednisolone, flurandrenolide, formocortal, halcinonide, halometasone, halopredone, hyrcanoside, hydrocortisone, hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydrocortisone valerate, 6-hydroxydexamethasone, isoflupredone, isoflupredone acetate, isoprednidene, meclorison, methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, paramethasone, paramethasone acetate, prednisolone, prednisolone acetate, prednisolone metasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hemisuccinate free acid, prednisolone-21-acetate, prednisolone-21 (beta-D- glucuronide), prednisone, prednylidene, procinonide, tralonide, triamcinolone, triamcinolone acetonide, triamcinolone acetonide 21-palmitate, triamcinolone diacetate, triamcinolone hexacetonide, or wortmannin.
14. The medicament of claim 13, wherein said corticosteroid is fludrocortisone or prednisolone.
15. The medicament of claim 14, wherein said corticosteroid is prednisolone or a pharmaceutically acceptable salt thereof.
16. The medicament of claim 14 or 15, wherein said medicament comprises a daily dosage of 0.5 to 10 mg of fludrocortisone or prednisolone.
17. The medicament of claim 12, 14, or 15, wherein said medicament comprises a daily dosage of 0.5 to 10 mg of corticosteroid.
18. The medicament of claim 12, 14, or 15, wherein said medicament comprises a daily dosage of 18 to 600 mg dipyridamole or a daily dosage of 50 to 400 mg dipyridamole.
19. A composition comprising two active ingredients, wherein the first of said active ingredients is a corticosteroid and the second of said active ingredients is dipyridamole, and wherein said composition is formulated for oral adminis-

tration.

20. The composition of claim 19, wherein said corticosteroid is algestone, 6- α -fluoroprednisolone, 6- α -methylprednisolone, 6- α -methylprednisolone 21-acetate, 6- α -methylprednisolone 21-hemisuccinate sodium salt, 6- α ,9- α -difluoroprednisolone 21-acetate 17-butyrate, amcinafal, beclomethasone, beclomethasone dipropionate, beclomethasone dipropionate monohydrate, 6- β -hydroxycortisol, betamethasone, betamethasone-17-valerate, budesonide, clobetasol, clobetasol propionate, clobetasone, clocortolone, clocortolone pivalate, cortisone, cortisone acetate, cortodoxone, deflazacort, 21-deoxycortisol, deprodone, descinolone, desonide, desoximethasone, dexamethasone, dexamethasone-21-acetate, dichlorisone, diflorasone, diflorasone diacetate, diflucortolone, doxibetasol, fludrocortisone, flumethasone, flumethasone pivalate, flumoxonide, flunisolide, fluocinonide, fluocinolone acetone, 9-fluorocortisone, fluorohydroxyandrostenedione, fluorometholone, fluorometholone acetate, fluoxymesterone, flupredidene, fluprednisolone, flurandrenolide, formocortol, halcinonide, halometasone, halopredone, hyrcanoside, hydrocortisone, hydrocortisone acetate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydrocortisone valerate, 6-hydroxydexamethasone, isoflupredone, isoflupredone acetate, isoprednidene, meclorisone, methylprednisolone, methylprednisolone acetate, methylprednisolone sodium succinate, paramethasone, paramethasone acetate, prednisolone, prednisolone acetate, prednisolone metasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hemisuccinate free acid, prednisolone-21-acetate, prednisolone-21 (β -D-glucuronide), prednisone, prednylidene, procinonide, tralonide, triamcinolone, triamcinolone acetonide, triamcinolone acetonide 21-palmitate, triamcinolone diacetate, triamcinolone hexacetonide, or wortmannin.

21. The composition of claim 19, wherein said composition comprises a daily dosage of 18 to 600 mg dipyridamole or a daily dosage of 50 to 400 mg dipyridamole.

22. The composition of claim 20 or 21, wherein said corticosteroid is fludrocortisone or prednisolone.

23. The composition of claim 22, wherein said corticosteroid is prednisolone or a pharmaceutically acceptable salt thereof.

24. The composition of claim 22 or 23, wherein said composition comprises a daily dosage of 0.5 to 10 mg of fludrocortisone or prednisolone.

Patentansprüche

1. Medikament enthaltend

- (a) ein Corticosteroid; und
- (b) Dipyridamol

zur Verwendung in einem Verfahren zur Behandlung eines Patienten, der eine immunoinflammatorische Krankheit hat oder gefährdet ist, eine immunoinflammatorische Krankheit zu entwickeln, worin die besagte immunoinflammatorische Krankheit ausgewählt wird aus der Gruppe bestehend aus rheumatoide Arthritis, Psoriasis, Colitis ulcerosa, Crohn-Krankheit, Spondylitis ankylosans, Fibromyalgia, Multiple Sklerose, Typ I Diabetes, systemischer Lupus Erythematodes, Sjögren-Syndrom und durch Gehirnschlag induzierter Gehirnzelltod, worin besagtes Medikament besagtes Dipyridamol und besagtes Corticosteroid zur simultanen Verabreichung oder zur Verabreichung innerhalb von 14 Tagen voneinander umfasst, in Mengen und für eine Dauer, die ausreichend ist, um besagten Patienten zu behandeln.

2. Medikament zur Verwendung gemäß Anspruch 1, worin besagtes Corticosteroid Algeston, 6- α -Fluorprednisolon, 6- α -Methylprednisolon, 6- α -Methylprednisolon 21-Azetat, 6- α -Methylprednisolon 21-Hemisuccinat Natriumsalz, 6- α , 9- α -Difluorprednisolon 21-Azetat 17-butyrat, Amcinafal, Beclomethason, Beclomethasondipropionat, Beclomethasondipropionat-Monohydrat, 6- β -Hydroxycortisol, Betamethason, Betamethason-17-Valerat, Budesonid, Clobetasol, Clobetasolpropionat, Clobetason, Clocortolon, Clocortolonpivalat, Cortison, Cortisonazetat, Cortodoxon, Deflazacort, 21-Deoxycortisol, Deprodon, Descinolone, Desonid, Desoximethason, Dexamethason, Dexamethason-21-Azetat, Dichlorison, Diflorason, Diflorasondiazetat, Diflucortolon, Doxibetasol, Fludrocortison, Flumethason, Flumethasonpivalat, Flumoxonid, Flunisolid, Fluocinonid, Fluocinoloneazetonid, 9-Fluorocortison, Fluorhydroxyandrostendion, Fluormetholon, Fluormetholonazetat, Fluoxymesteron, Fluprediden, Flupred-

nisolon, Flurandrenolid, Formocortal, Halcinonid, Halometason, Halopredon, Hyrcanosid, Hydrocortison, Hydrocortisonazetat, Hydrocortisonbutyrat, Hydrocortisoncypionat, Hydrocortisonnatriumphosphat, Hydrocortisonnatriumsuccinat, Hydrocortisonprobutat, Hydrocortisonvalerat, 6-Hydroxydexamethason, Isoflupredon, Isoflupredonazetat, Isopredniden, Meclorison, Methylprednisolon, Methylprednisolonazetat, Methylprednisolonnatriumsuccinat, Paramethason, Paramethasonazetat, Prednisolon, Prednisolonazetat, Prednisolonmetasulphobenzoat, Prednisolonnatriumphosphat, Prednisolontebutat, Prednisolon-21-Hemisuccinat freie Säure, Prednisolon-21-Azetat, Prednisolon-21(beta-D-Glucuronid), Prednison, Prednyliden, Procinonid, Tralonid, Triamcinolon, Triamcinolonacetamid, Triamcinolonacetamid 21-Palmitat, Triamcinolondiazetat, Triamcinolonhexacetamid, oder Wortmannin ist.

3. Medikament zur Verwendung gemäß Anspruch 2, worin besagtes Corticosteroid Fludrocortison oder Prednisolon ist.

4. Medikament zur Verwendung gemäß Anspruch 3, worin besagtes Corticosteroid Prednisolon oder ein pharmazeutisch verträgliches Salz davon ist.

5. Medikament zur Verwendung gemäß Anspruch 1 oder 3, oder 4, worin besagtes Medikament Dipyridamol und besagtes Corticosteroid zur Verabreichung innerhalb von zehn Tagen voneinander umfasst, zur Verabreichung innerhalb von fünf Tagen voneinander, zur Verabreichung innerhalb von vierundzwanzig Stunden voneinander, innerhalb einer Stunde voneinander, oder für die simultane Verabreichung von besagtem Dipyridamol und besagtem Corticosteroid.

6. Medikament zur Verwendung gemäß Anspruch 1, 3 oder 4, worin besagtes Medikament besagtes Dipyridamol und besagtes Corticosteroid zur Verabreichung in derselben pharmazeutischen Formulierung umfasst.

7. Medikament zur Verwendung gemäß Anspruch 1 oder 4, 5 oder 6, worin besagte immunoinflammatorische Krankheit rheumatoide Arthritis, Psoriasis, Colitis ulcerosa, Crohn-Krankheit, Spondylitis ankylosans, oder Multiple Sklerose ist.

8. Medikament zur Verwendung gemäß Anspruch 1, 4, 5, 6 oder 7, worin besagtes Medikament besagtes Dipyridamol in einer Menge von 0,5-800 mg/Tag umfasst und besagtes Corticosteroid in einer Menge von 0,1-1500 mg/Tag umfasst, worin besagtes Medikament besagtes Dipyridamol in einer Menge von 18-600 mg/Tag umfasst und besagtes Corticosteroid in einer Menge von 0,5-30 mg/Tag umfasst, oder worin besagtes Medikament besagtes Corticosteroid in einer Menge von 0,5-10 mg/Tag umfasst.

9. Medikament zur Verwendung gemäß Anspruch 1, 3, 4 oder 7, worin besagtes Medikament eine tägliche Dosis von 18 bis 600 mg Dipyridamol oder eine tägliche Dosis von 50 bis 400 mg Dipyridamol umfasst.

10. Medikament zur Verwendung gemäß Anspruch 1, 3, 4 oder 7, worin besagtes Medikament eine tägliche Dosis von 0,5 bis 10 mg Fludrocortison oder Prednisolon umfasst.

11. Medikament zur Verwendung gemäß Anspruch 1, 3 oder 4, worin besagtes Medikament für die orale oder intravenöse Verabreichung formuliert wird.

12. Medikament bestehend aus Wirkstoffen und Hilfsstoffen, worin besagte Wirkstoffe aus einem Corticosteroid und Dipyridamol bestehen und worin besagtes Medikament für die orale Verabreichung formuliert wird.

13. Medikament nach Anspruch 12, worin besagtes Corticosteroid Algeston, 6-alpha-Fluorprednisolon, 6-alpha-Methylprednisolon, 6-alpha-Methylprednisolon 21-Azetat, 6-alpha-Methylprednisolon 21-Hemisuccinat Natriumsalz, 6-alpha, 9-alpha-Difluorprednisolon 21-Azetat 17-butytrat, Amcinafal, Beclomethason, Beclomethasondipropionat, Beclomethasondipropionat-Monohydrat, 6-beta-Hydroxycortisol, Betamethason, Betamethason-17-Valerat, Budesonid, Clobetasol, Clobetasolpropionat, Clobetason, Clotololon, Clotololonpivalat, Cortison, Cortisonazetat, Cortodoxon, Deflazacort, 21-Deoxycortisol, Depredon, Descinololon, Desonid, Desoximethason, Dexamethason, Dexamethason-21-Azetat, Dichlorison, Diflorason, Diflorasondiazetat, Diflucortolon, Doxibetasol, Fludrocortison, Flumethason, Flumethasonpivalat, Flumoxonid, Flunisolid, Fluocinonid, Fluocinolonazetonid, 9-Fluorcortison, Fluorhydroxyandrostendion, Fluormetholon, Fluormetholonazetat, Fluoxymesteron, Fluprediden, Fluprednisolon, Flurandrenolid, Formocortal, Halcinonid, Halometason, Halopredon, Hyrcanosid, Hydrocortison, Hydrocortisonazetat, Hydrocortisonbutyrat, Hydrocortisoncypionat, Hydrocortisonnatriumphosphat, Hydrocortisonnatriumsuccinat, Hydrocortisonprobutat, Hydrocortisonvalerat, 6-Hydroxydexamethason, Isoflupredon, Isoflupredonazetat, Isopredniden, Meclorison, Methylprednisolon, Methylprednisolonazetat, Methylprednisolonnatriumsuccinat, Paramethason,

Paramethasonazetat, Prednisolon, Prednisolonazetat, Prednisolonmetasulphobenzoat, Prednisolonnatriumphosphat, Prednisolontebutat, Prednisolon-21-Hemisuccinat freie Säure, Prednisolon-21-Azetat, Prednisolon-21 (beta-D-Glucuronid), Prednison, Prednyliden, Procinonid, Tralonid, Triamcinolon, Triamcinolonacetamid, Triamcinolonacetamid 21-Palmitat, Triamcinolondiazetat, Triamcinolonhexacetamid, oder Wortmannin ist.

- 5 14. Medikament nach Anspruch 13, worin besagtes Corticosteroid Fludrocortison oder Prednisolon ist.
15. Medikament nach Anspruch 14, worin besagtes Corticosteroid Prednisolon oder ein pharmazeutisch verträgliches Salz davon ist.
- 10 16. Medikament nach Anspruch 14 oder 15, worin besagtes Medikament eine tägliche Dosis von 0,5 bis 10 mg Fludrocortison oder Prednisolon umfasst.
- 15 17. Medikament nach Anspruch 12, 14 oder 15, worin besagtes Medikament eine tägliche Dosis von 0,5 bis 10 mg des Corticosteroids umfasst.
18. Medikament des Anspruchs 12, 14 oder 15, worin besagtes Medikament eine tägliche Dosis von 18 bis 600 mg Dipyridamol oder eine tägliche Dosis von 50 bis 400 mg Dipyridamol umfasst.
- 20 19. Zusammensetzung enthaltend zwei Wirkstoffe, worin der erste der besagten Wirkstoffe ein Corticosteroid ist und der zweite der besagten Wirkstoffe Dipyridamol ist, und worin besagte Zusammensetzung für die orale Verabreichung formuliert ist.
- 25 20. Zusammensetzung nach Anspruch 19, worin besagtes Corticosteroid Algeston, 6-alpha-Fluorprednisolon, 6-alpha-Methylprednisolon, 6-alpha-Methylprednisolon 21-Azetat, 6-alpha-Methylprednisolon 21-Hemisuccinat Natriumsalz, 6-alpha, 9-alpha-Difluorprednisolon 21-Azetat 17-butytrat, Amcinafal, Beclomethason, Beclomethasondipropionat, Beclomethasondipropionat-Monohydrat, 6-beta-Hydroxycortisol, Betamethason, Betamethason-17-Valerat, Budesonid, Clobetasol, Clobetasolpropionat, Clobetason, Clodacortol, Clodacortolnivalat, Cortison, Cortisonazetat, Cortodoxon, Deflazacort, 21-Deoxycortisol, Depredon, Descinol, Desonid, Desoximethason, Dexamethason, Dexamethason-21-Azetat, Dichlorison, Diflorason, Diflorasondiazetat, Diflucortolon, Doxibetasol, Fludrocortison, Flumethason, Flumethasonpivalat, Flumoxonid, Flunisolid, Fluocinonid, Fluocinolonacetamid, 9-Fluorocortison, Fluorhydroxyandrostendion, Fluormetholon, Fluormetholonazetat, Fluoxymesteron, Flupredniden, Fluprednisolon, Flurandrenolid, Formocortal, Halcinonid, Halometason, Halopredon, Hyrcanosid, Hydrocortison, Hydrocortisonazetat, Hydrocortisonbutyrat, Hydrocortisoncypionat, Hydrocortisonnatriumphosphat, Hydrocortisonnatriumsuccinat, Hydrocortisonprobutat, Hydrocortisonvalerat, 6-Hydroxydexamethason, Isoflupredon, Isoflupredonazetat, Isopredniden, Meclorison, Methylprednisolon, Methylprednisolonazetat, Methylprednisolonnatriumsuccinat, Paramethason, Paramethasonazetat, Prednisolon, Prednisolonazetat, Prednisolonmetasulphobenzoat, Prednisolonnatriumphosphat, Prednisolontebutat, Prednisolon-21-Hemisuccinat freie Säure, Prednisolon-21-Azetat, Prednisolon-21 (beta-D-Glucuronid), Prednison, Prednyliden, Procinonid, Tralonid, Triamcinolon, Triamcinolonacetamid, Triamcinolonacetamid 21-Palmitat, Triamcinolondiazetat, Triamcinolonhexacetamid, oder Wortmannin ist.
- 30 21. Zusammensetzung nach Anspruch 19, worin besagte Zusammensetzung eine tägliche Dosis von 18 bis 600 mg Dipyridamol oder eine tägliche Dosis von 50 bis 400 mg Dipyridamol umfasst.
- 35 22. Zusammensetzung nach Anspruch 20 oder 21, worin besagtes Corticosteroid Fludrocortison oder Prednisolon ist.
- 40 23. Zusammensetzung nach Anspruch 22, worin besagtes Corticosteroid Prednisolon oder ein pharmazeutisch verträgliches Salz davon ist.
- 45 24. Zusammensetzung nach Anspruch 22 oder 23, worin besagte Zusammensetzung eine tägliche Dosis von 0,5 bis 10 mg Fludrocortison oder Prednisolon umfasst.

Revendications

1. Un médicament comprenant:

- (a) un corticostéroïde; et
- (b) du dipyridamole

pour son utilisation dans une méthode pour traiter un patient qui a un désordre immunoinflammatoire ou est à risque de développer un désordre immunoinflammatoire, dans lequel ledit désordre immunoinflammatoire est sélectionné dans le groupe consistant en l'arthrite rhumatoïde, le psoriasis, la colite ulcéreuse, la maladie de Crohn, la spondylite ankylosante, la fibromyalgie, la sclérose en plaque, les diabètes de type I, le lupus systémique érythémateux, le syndrome de Sjögren, et la mort des cellules du cerveau induite par un accident vasculaire cérébral, dans lequel ledit médicament comprend ledit dipyridamole et ledit corticostéroïde pour une administration simultanée ou pour une administration séquentielle avec un intervalle maximum de 14 jours entre l'un et l'autre, dans des quantités et pour une durée suffisante pour traiter ledit patient.

2. Le médicament pour utilisation selon la revendication 1, dans lequel ledit corticostéroïde est algestone, 6- α -fluoroprednisolone, 6- α -méthylprednisolone, 6- α -méthylprednisolone 21-acétate, 6- α -méthylprednisolone 21-hémisuccinate sel de sodium, 6- α ,9- α -difluoroprednisolone 21-acétate 17-butyrate, amcinafal, béclo-méthasone, béclo-méthasone dipropionate, béclo-méthasone dipropionate monohydrate, 6-bétahydroxycortisol, bétaméthasone, bétaméthasone-17-valérate, budésonide, clobétasol, clobétasol propionate, clobétasone, clocortolone, clocortolone pivalate, cortisone, cortisone acétate, cortodoxone, deflazacort, 21-déoxycortisol, déprodone, descinolone, désenide, désoxyméthasone, dexaméthasone, dexaméthasone-21-acétate, dichlorisone, diflorasone, diflorasone diacétate, difluocortolone, doxybétasol, fludrocortisone, fluméthasone, fluméthasone pivalate, flumoxonide, flunisolide, fluocinonide, fluocinolone acétonide, 9-fluorocortisone, fluorohydroxyandrostènedione, fluorométholone, fluorométholone acétate, fluoxymestérone, flupredidène, fluprednisolone, flurandrénolide, formocortal, halcinonide, halométhasone, haloprédone, hircanoside, hydrocortisone, hydrocortisone acétate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydrocortisone valérate, 6-hydroxydexaméthasone, isoflupredone, isoflupredone acétate, isoprednidène, méclorisone, méthylprednisolone, méthylprednisolone acétate, méthylprednisolone sodium succinate, paraméthasone, paraméthasone acétate, prednisolone, prednisolone acétate, prednisolone métasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hémisuccinate acide libre, prednisolone-21-acétate, prednisolone-21 (beta-D-glucuronide), prednisone, prednylidène, procinonide, tralonide, triamcinolone, triamcinolone acétonide, triamcinolone acétonide 21-palmitate, triamcinolone diacétate, triamcinolone hexacétonide ou wortmannine.
3. Le médicament pour utilisation selon la revendication 2, dans lequel ledit corticostéroïde est la fludrocortisone ou la prednisolone.
4. Le médicament pour utilisation selon la revendication 3, dans lequel ledit corticostéroïde est la prednisolone ou un sel de celle-ci pharmaceutiquement acceptable.
5. Le médicament pour utilisation selon la revendication 1 ou 3 ou 4, dans lequel ledit médicament comprend du dipyridamole et ledit corticostéroïde pour administration séquentielle avec un intervalle maximum de 10 jours entre l'un et l'autre, pour administration séquentielle avec un intervalle maximum de 5 jours entre l'un et l'autre, pour administration séquentielle avec un intervalle maximum de 24 heures entre l'un et l'autre, séquentielle avec un intervalle maximum de 1 heure entre l'un et l'autre, ou pour une administration simultanée dudit dipyridamole et dudit corticostéroïde.
6. Le médicament pour utilisation selon la revendication 1, 3 ou 4, dans lequel ledit médicament comprend ledit dipyridamole et ledit corticostéroïde pour administration dans la même formulation pharmaceutique.
7. Le médicament pour utilisation selon la revendication 1 ou 4, 5 ou 6, dans lequel ledit désordre immunoinflammatoire est l'arthrite rhumatoïde, le psoriasis, la colite ulcéreuse, la maladie de Crohn, la spondylite ankylosante ou la sclérose en plaque.
8. Le médicament pour utilisation selon la revendication 1, 4, 5, 6 ou 7, dans lequel ledit médicament comprend ledit dipyridamole dans une quantité de 0.5-800 mg/jour et comprend ledit corticostéroïde dans une quantité de 0.1-1500 mg/jour, dans lequel ledit médicament comprend ledit dipyridamole dans une quantité de 18-600 mg/jour et comprend ledit corticostéroïde dans une quantité de 0.5-30 mg/jour, ou dans lequel ledit médicament comprend ledit corticostéroïde dans une quantité de 0.5-10 mg/jour.

9. Le médicament pour utilisation selon la revendication 1, 3, 4 ou 7, dans lequel ledit médicament comprend un dosage quotidien de 18 à 600 mg de dipyridamole ou un dosage quotidien de 50 à 400 mg de dipyridamole.
- 5 10. Le médicament pour utilisation selon la revendication 1, 3, 4 ou 7, dans lequel ledit médicament comprend un dosage quotidien de 0.5 à 10 mg de fludrocortisone ou de prednisolone.
11. Le médicament pour utilisation selon la revendication 1, 3 ou 4, dans lequel ledit médicament est formulé pour administration orale ou intraveineuse.
- 10 12. Un médicament consistant en des ingrédients actifs et des excipients, dans lequel lesdits ingrédients actifs consistent en un corticostéroïde et du dipyridamole et dans lequel ledit médicament est formulé pour une administration orale.
13. Le médicament selon la revendication 12, dans lequel ledit corticostéroïde est algestone, 6-alpha-fluoroprednisolone, 6-alpha-méthylprednisolone, 6-alpha-méthylprednisolone 21-acétate, 6-alpha-méthylprednisolone 21-hémisuccinate sel de sodium, 6-alpha,9-alpha-difluoroprednisolone 21-acétate 17-butyrate, amcinafal, béclo-méthasone, béclo-méthasone dipropionate, béclo-méthasone dipropionate monohydrate, 6-bétahydroxycortisol, bétaméthasone, bétaméthasone-17-valérate, budésonide, clobétasol, clobétasol propionate, clobétasone, clocortolone, clocortolone pivalate, cortisone, cortisone acétate, cortodoxone, deflazacort, 21-déoxycortisol, déprodone, descinolone, désos-nide, désoxyméthasone, dexaméthasone, dexaméthasone-21-acétate, dichlorisone, diflorasone, diflorasone dia-cétate, diflucortolone, doxybétasol, fludrocortisone, fluméthasone, fluméthasone pivalate, flumoxonide, flunisolide, flucinonide, fluocinolone acétonide, 9-fluorocortisone, fluorohydroxyandrostènedione, fluorométholone, fluorométholone acétate, fluoxymestérone, flupredidène, fluprednisolone, flurandrénilide, formocortal, halcinonide, halométhasone, halopredone, hyrcanoside, hydrocortisone, hydrocortisone acétate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydro-cortisone valérate, 6-hydroxydexaméthasone, isoflupredone, isoflupredone acétate, isoprednidène, méclorisone, méthylprednisolone, méthylprednisolone acétate, méthylprédnisolone sodium succinate, paraméthasone, paraméthasone acétate, prednisolone, prednisolone acétate, prednisolone mé-tasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hémisuccinate acide libre, prednisolone-21-acétate, predniso-lone-21 (beta-D-glucuronide), prednisone, prednylidène, procinonide, tralonide, triamcinolone, triamcinolone acé-tonide, triamcinolone acétonide 21-palmitate, triamcinolone diacétate, triamcinolone hexacétonide ou wortmannine.
14. Le médicament selon la revendication 13, dans lequel ledit corticostéroïde est la fludrocortisone ou la prednisolone.
15. Le médicament selon la revendication 14, dans lequel ledit corticostéroïde est la prednisolone ou un sel de celle-ci pharmaceutiquement acceptable.
- 35 16. Le médicament selon la revendication 14 ou 15, dans lequel ledit médicament comprend un dosage quotidien de 0.5 à 10 mg de fludrocortisone ou de prednisolone.
17. Le médicament selon la revendication 12, 14 ou 15, dans lequel ledit médicament comprend un dosage quotidien de 0.5 à 10 mg de corticostéroïde.
- 40 18. Le médicament selon la revendication 12, 14 ou 15, dans lequel ledit médicament comprend un dosage quotidien de 18 à 600 mg de dipyridamole ou un dosage quotidien de 50 à 400 mg de dipyridamole.
- 45 19. Une composition comprenant deux ingrédients actifs, dans laquelle le premier desdits ingrédients actifs est un corticostéroïde et le second desdits ingrédients actifs est le dipyridamole, et dans laquelle ladite composition est formulée pour une administration orale.
- 50 20. La composition selon la revendication 19, dans laquelle ledit corticostéroïde est algestone, 6-alpha-fluoroprednisolone, 6-alpha-méthylprednisolone, 6-alpha-méthylprednisolone 21-acétate, 6-alpha-méthylprednisolone 21-hémisuccinate sel de sodium, 6-alpha,9-alpha-difluoroprednisolone 21-acétate 17-butyrate, amcinafal, béclo-méthasone, béclo-méthasone dipropionate, béclo-méthasone dipropionate monohydrate, 6-bétahydroxycortisol, bétaméthasone, bétaméthasone-17-valérate, budésonide, clobétasol, clobétasol propionate, clobétasone, clocortolone, clocortolone pivalate, cortisone, cortisone acétate, cortodoxone, deflazacort, 21-déoxycortisol, déprodone, descinolone, désos-nide, désoxyméthasone, dexaméthasone, dexaméthasone-21-acétate, dichlorisone, diflorasone, diflorasone dia-cétate, diflucortolone, doxybétasol, fludrocortisone, fluméthasone, fluméthasone pivalate, flumoxonide, flunisolide, flucinonide, fluocinolone acétonide, 9-fluorocortisone, fluorohydroxyandrostènedione, fluorométholone, fluorométholone acétate, fluoxymestérone, flupredidène, fluprednisolone, flurandrénilide, formocortal, halcinonide, halométhasone, halopredone, hyrcanoside, hydrocortisone, hydrocortisone acétate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydro-cortisone valérate, 6-hydroxydexaméthasone, isoflupredone, isoflupredone acétate, isoprednidène, méclorisone, méthylprednisolone, méthylprednisolone acétate, méthylprédnisolone sodium succinate, paraméthasone, paraméthasone acétate, prednisolone, prednisolone acétate, prednisolone mé-tasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hémisuccinate acide libre, prednisolone-21-acétate, predniso-lone-21 (beta-D-glucuronide), prednisone, prednylidène, procinonide, tralonide, triamcinolone, triamcinolone acé-tonide, triamcinolone acétonide 21-palmitate, triamcinolone diacétate, triamcinolone hexacétonide ou wortmannine.
- 55

tholone acétate, fluoxymestérone, flupredidène, fluprednisolone, flurandrénilide, formocortal, halcinonide, halométhasone, haloprédone, hyrcanoside, hydrocortisone, hydrocortisone acétate, hydrocortisone butyrate, hydrocortisone cypionate, hydrocortisone sodium phosphate, hydrocortisone sodium succinate, hydrocortisone probutate, hydrocortisone valérate, 6-hydroxydexaméthasone, isoflupredone, isoflupredone acétate, isoprednidène, méclorisone, méthylprednisolone, méthylprednisolone acétate, méthylprédnisolone sodium succinate, paraméthasone, paraméthasone acétate, prednisolone, prednisolone acétate, prednisolone métrasulphobenzoate, prednisolone sodium phosphate, prednisolone tebutate, prednisolone-21-hémisuccinate acide libre, prednisolone-21-acétate, prednisolone-21 (beta-D-glucuronide), prednisone, prednylidène, procinonide, tralonide, triamcinolone, triamcinolone acétonide, triamcinolone acétonide 21-palmitate, triamcinolone diacétate, triamcinolone hexacétonide ou wortmannine.

21. La composition selon la revendication 19, dans laquelle ladite composition comprend un dosage quotidien de 18 à 600 mg de dipyridamole ou un dosage quotidien de 50 à 400 mg de dipyridamole.
22. La composition selon la revendication 20 ou 21, dans laquelle ledit corticostéroïde est la fludrocortisone ou la prednisolone.
23. La composition selon la revendication 22, dans laquelle ledit corticostéroïde est la prednisolone ou un sel de celle-ci pharmaceutiquement acceptable.
24. La composition selon la revendication 22 ou 23, dans laquelle ladite composition comprend un dosage quotidien de 0.5 à 10 mg de fludrocortisone ou de prednisolone.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Non-patent literature cited in the description

- **Arnett FC et al.** *Arthritis Rheum.*, 1988, vol. 31, 315-324 **[0004]**
- Remington: The Science and Practice of Pharmacy. Lippincott Williams & Wilkins, 2000 **[0028]**
- Encyclopedia of Pharmaceutical Technology. Marcel Dekker, 1988 **[0028]**