



(86) Date de dépôt PCT/PCT Filing Date: 1998/06/15

(87) Date publication PCT/PCT Publication Date: 1998/12/23

(45) Date de délivrance/Issue Date: 2004/01/06

(85) Entrée phase nationale/National Entry: 1999/12/08

(86) N° demande PCT/PCT Application No.: US 1998/011502

(87) N° publication PCT/PCT Publication No.: 1998/057963

(30) Priorité/Priority: 1997/06/17 (08/877,399) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 401/14, A61K 31/505,
A61K 31/445


(72) Inventeurs/Inventors:
NJORGE, F. GEORGE, US;
VIBULBHAN, BANCHA, US;
PINTO, PATRICK A., US;
GIRIJAVALLABHAN, VIYYOOR M., US

(73) Propriétaire/Owner:
SCHERING CORPORATION, US

(74) Agent: OGILVY RENAULT

(54) Titre : UREES CYCLIQUES BENZO(5,6)CYCLOHEPTAPYRIDINE ET LACTAMES UTILES COMME INHIBITEURS
DE FARNESYL PROTEINE TRANSFERASE

(54) Title: BENZO(5,6)CYCLOHEPTAPYRIDINE CYCLIC UREAS AND LACTAMS USEFUL AS FARNESYL PROTEIN
TRANSFERASE INHIBITORS



(57) Abrégé/Abstract:

Compounds of formula (I) useful for inhibiting Ras function and therefore inhibiting or treating the abnormal growth of cells are disclosed, or a pharmaceutically acceptable salt or solvate thereof, wherein: R and R² are halo; R¹ and R³ are H and halo, provided that at least one is H; W is N, CH, or C when the double bond is present at the C-11 position; R⁴ is -(CH₂)_n-R⁵ or (II); R⁵ is (III); R⁶ is R⁵ or (IV); Z¹ and Z² are independently selected from the group consisting of =O and =S; n is 1-6; and n₁ is 0 or 1.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :

C07D 401/14, A61K 31/445, 31/505

A1

(11) International Publication Number:

WO 98/57963

(43) International Publication Date:

23 December 1998 (23.12.98)

(21) International Application Number: PCT/US98/11502

(22) International Filing Date: 15 June 1998 (15.06.98)

(30) Priority Data:

08/877,399

17 June 1997 (17.06.97)

US

(71) Applicant: SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: NJORGE, F., George; 2597 Juliat Place, Union, NJ 07083 (US). VIBULBHAN, Bancha; 201 North 24th Street, Kenilworth, NJ 07033 (US). PINTO, Patrick, A.; 34 Battle Ridge Road, Morris Plains, NJ 07950 (US). GIRIJAVALLABHAN, Viyyoor, M.; 10 Maplewood Drive, Parsippany, NJ 07054 (US).


(74) Agents: MAGATTI, Anita, W. et al.; Schering-Plough Corporation, Patent Dept., K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).

(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CZ, EE, GE, GW, HU, ID, IL, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).


Published

With international search report.




(54) Title: BENZO(5,6)CYCLOHEPTAPYRIDINE CYCLIC UREAS AND LACTAMS USEFUL AS FARNESYL PROTEIN TRANSFERASE INHIBITORS




(I)



(II)



(III)



(IV)

(57) Abstract


Compounds of formula (I) useful for inhibiting Ras function and therefore inhibiting or treating the abnormal growth of cells are disclosed, or a pharmaceutically acceptable salt or solvate thereof, wherein: R and R² are halo; R¹ and R³ are H and halo, provided that at least one is H; W is N, CH, or C when the double bond is present at the C-11 position; R⁴ is -(CH₂)_n-R⁵ or (II); R⁵ is (III); R⁶ is R⁵ or (IV); Z¹ and Z² are independently selected from the group consisting of =O and =S; n is 1-6; and n₁ is 0 or 1.

5

BENZO(5,6)CYCLOHEPTAPYRIDINE
CYCLIC UREAS AND LACTAMS USEFUL AS
FARNESYL PROTEIN TRANSFERASE INHIBITORS

10 **BACKGROUND**


International Publication Number WO95/10516, published April 20, 1995 discloses compounds of the formula:



15 wherein R can be heterocycloalkyl attached to the carbon of the -C(=Z)-group by a heteroatom, substituted piperidiny or substituted piperidinylmethyl. The compounds are said to be useful for inhibiting farnesyl protein transferase.

20 **SUMMARY OF THE INVENTION**

Compounds of the present invention are represented by Formula I:






or an N-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof, wherein:

- 2 -

R and R² are independently selected from halo;

R¹ and R³ are independently selected from the group consisting of H and halo, provided that at least one of R¹ and R³ is H;

W is N, CH or C when the double bond is present at the C-11 position;




Z¹ and Z² are independently selected from the group consisting of =O and =S;

n is 1-6; and

n₁ is 0 or 1.

In the compounds of the invention, preferably R is Br, R² is halo and R¹ is halo; or R is Br, R² is halo and R³ is halo; or R is Br, R² is halo and R¹ and R³ are each H. R² is preferably Br or Cl. When R¹ or R³ is halo, it is preferably Br or Cl. Z¹ is preferably =O. Z² is preferably =O. W is preferably CH. Preferred values for n are 1-3. R⁵ and R⁶ are



The compounds of this invention: (i) potently inhibit farnesyl protein transferase, but not geranylgeranyl protein transferase I, *in vitro*; (ii) block the phenotypic change induced by a form of transforming Ras which is a farnesyl acceptor but not by a form of transforming Ras engineered to be a geranylgeranyl acceptor; (iii) block intracellular processing of Ras which is a farnesyl acceptor but not of Ras engineered to be a geranylgeranyl acceptor; and (iv) block abnormal cell growth in culture induced by transforming Ras.

- 3 -

The compounds of this invention inhibit farnesyl protein transferase and the farnesylation of the oncogene protein Ras. This invention further provides a method of inhibiting ras farnesyl protein transferase, in mammals, especially humans, by the administration of an effective amount
5 of the tricyclic compounds described above. The administration of the compounds of this invention to patients, to inhibit farnesyl protein transferase, is useful in the treatment of the cancers described below.

This invention provides a method for inhibiting or treating the abnormal growth of cells, including transformed cells, by administering an
10 effective amount of a compound of this invention. Abnormal growth of cells refers to cell growth independent of normal regulatory mechanisms (e.g., loss of contact inhibition). This includes the abnormal growth of: (1) tumor cells (tumors) expressing an activated Ras oncogene; (2) tumor cells in which the Ras protein is activated as a result of oncogenic
15 mutation in another gene; and (3) benign and malignant cells of other proliferative diseases in which aberrant Ras activation occurs.

This invention also provides a method for inhibiting or treating tumor growth by administering an effective amount of the tricyclic compounds, described herein, to a mammal (e.g., a human) in need of
20 such treatment. In particular, this invention provides a method for inhibiting or treating the growth of tumors expressing an activated Ras oncogene by the administration of an effective amount of the above described compounds. Examples of tumors which may be inhibited or treated include, but are not limited to, breast cancer, prostate cancer, lung
25 cancer (e.g., lung adenocarcinoma), pancreatic cancers (e.g., pancreatic carcinoma such as, for example, exocrine pancreatic carcinoma), colon cancers (e.g., colorectal carcinomas, such as, for example, colon adenocarcinoma and colon adenoma), myeloid leukemias (for example, acute myelogenous leukemia (AML)), thyroid follicular cancer,
30 myelodysplastic syndrome (MDS), bladder carcinoma and epidermal carcinoma.

It is believed that this invention also provides a method for inhibiting or treating proliferative diseases, both benign and malignant, wherein Ras proteins are aberrantly activated as a result of oncogenic mutation in other
35 genes--i.e., the Ras gene itself is not activated by mutation to an oncogenic form--with said inhibition or treatment being accomplished by the administration of an effective amount of the tricyclic compounds described herein, to a mammal (e.g., a human) in need of such treatment.

- 4 -

For example, the benign proliferative disorder neurofibromatosis, or tumors in which Ras is activated due to mutation or overexpression of tyrosine kinase oncogenes (e.g., neu, src, abl, lck, and fyn), may be inhibited or treated by the tricyclic compounds described herein.

5 The tricyclic compounds useful in the methods of this invention inhibit or treat the abnormal growth of cells. Without wishing to be bound by theory, it is believed that these compounds may function through the inhibition of G-protein function, such as ras p21, by blocking G-protein isoprenylation, thus making them useful in the treatment of proliferative
10 diseases such as tumor growth and cancer. Without wishing to be bound by theory, it is believed that these compounds inhibit ras farnesyl protein transferase, and thus show antiproliferative activity against ras transformed cells.

15 DETAILED DESCRIPTION OF THE INVENTION

As used herein, the following terms are used as defined below unless otherwise indicated:

MH⁺ represents the molecular ion plus hydrogen of the molecule in the mass spectrum;


20 Bu represents butyl; Et represents ethyl; Me represents methyl; Ph represents phenyl; and

halo represents fluoro, chloro, bromo and iodo.

The following solvents and reagents may be referred to herein by the abbreviations indicated: tetrahydrofuran (THF); ethanol (EtOH);
25 methanol (MeOH); acetic acid (HOAc or AcOH); ethyl acetate (EtOAc); N,N-dimethylformamide (DMF); trifluoroacetic acid (TFA); trifluoroacetic anhydride (TFAA); 1-hydroxybenzotriazole (HOBT); m-chloroperbenzoic acid (MCPBA); triethylamine (Et₃N); diethyl ether (Et₂O); ethyl
30 chloroformate (ClCO₂Et); and 1-(3-dimethylaminopropyl)-3-ethyl carbodiimide hydrochloride (DEC).

Representative structures of Formula I with respect to W and the optional double bond are as follows:

- 5 -



Lines drawn into the ring systems indicate that the indicated bond may be attached to any of the substitutable ring carbon atoms.

Certain compounds of the invention may exist in different isomeric (e.g., enantiomers and diastereoisomers) forms. The invention contemplates all such isomers both in pure form and in admixture, including racemic mixtures. Enol forms are also included.

Certain tricyclic compounds will be acidic in nature, e.g. those compounds which possess a carboxyl or phenolic hydroxyl group. These compounds may form pharmaceutically acceptable salts. Examples of such salts may include sodium, potassium, calcium, aluminum, gold and silver salts. Also contemplated are salts formed with pharmaceutically acceptable amines such as ammonia, alkyl amines, hydroxyalkylamines, N-methylglucamine and the like.


Certain basic tricyclic compounds also form pharmaceutically acceptable salts, e.g., acid addition salts. For example, the pyrido-nitrogen atoms may form salts with strong acid, while compounds having basic substituents such as amino groups also form salts with weaker acids. Examples of suitable acids for salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, oxalic, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those in the art. The salts are prepared by contacting the free base form with a sufficient amount of the desired acid to produce a salt in the conventional manner. The free base forms may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia and sodium bicarbonate. The free base forms differ from their respective salt forms somewhat in certain physical properties, such as solubility in polar solvents, but the acid and base salts are otherwise equivalent to their respective free base forms for purposes of the invention.

- 6 -

All such acid and base salts are intended to be pharmaceutically acceptable salts within the scope of the invention and all acid and base salts are considered equivalent to the free forms of the corresponding compounds for purposes of the invention.


5 Compounds of the invention may be made by the methods described in the examples below, and by using the methods described in WO 95/10516 -- see, for example, the methods for preparing compounds of Formula 400.00.

10 Compounds of the invention wherein Z^1 and Z^2 are =O can be prepared by reacting a compound of the formula II or III



15 wherein all other substituents are as defined for Formula I, with an acid of the formula $\text{HOOC}-(\text{CH}_2)_n-\text{NHR}^7$ or $\text{HOOC}-(\text{CH}_2)_{n3}-\text{NHR}^7$, respectively, wherein n is as defined above and R^7 is an amino protecting group such as t-butoxycarbonyl (BOC). The reaction is carried out using standard amide coupling conditions, for example the reaction can be carried out at room temperature, in an inert solvent such as DMF, in the presence of a condensing agent such as 1-(3-dimethylaminopropyl)-3-ethyl-

20 activating agent such as 1-hydroxybenzotriazole. The R^7 protecting group is then removed, for example by treatment with trifluoroacetic acid, to obtain the corresponding amine of the formula IIA or IIIA:




- 7 -

To prepare compounds of formula I wherein R⁵ or R⁶ comprises a cyclic lactam, a compound of formula IIA or IIIA is reacted with 4-bromobutyryl chloride or 4-bromovaleryl chloride, followed by cyclization with a reagent such as NaH. Compounds of formula I wherein R⁵ or R⁶ comprises a cyclic urea are similarly prepared by reacting an amine of formula IIA or IIIA with 2-bromo ethyl isocyanate or 3-chloropropyl isocyanate, followed as before by cyclization with a reagent such as NaH.

Alternatively, an amine of formula IIA or IIIA can be reacted with lactam-substituted acetate under standard amide coupling conditions as described above.

When Z¹, or Z¹ and Z², represent sulfur, a compound of formula I wherein Z¹, or Z¹ and Z², is oxygen is reacted with P₂S₅, Lawesson's reagent, or another reagent capable of introducing sulfur in place of oxygen. The reaction may take place at elevated temperature in pyridine, toluene or other suitable solvents. For compounds wherein Z¹ and Z² are different, the conversion from oxygen to sulfur can be effected before the starting materials (i.e., compounds of formula IIIA and the alkanoyl chloride or isocyanate) are reacted.

Compounds of formula I comprising a pyridyl N-oxide in ring I of the tricyclic portion can be prepared by procedures well known in the art. For example, the compound of formula II wherein W is C or CH can be reacted with MCPBA in a suitable organic solvent, e.g., CH₂Cl₂ (usually anhydrous), at a suitable temperature, to obtain an N-oxide of formula IIa




Generally, the organic solvent solution of formula II is cooled to about 0°C before the MCPBA is added. The reaction is then allowed to warm to room temperature during the reaction period. The desired product can be recovered by standard separation means, for example, the reaction mixture can be washed with an aqueous solution of a suitable base, e.g., saturated NaHCO₃ or NaOH (e.g., 1 N NaOH), and then dried over anhydrous MgSO₄. The solution containing the product can be

- 8 -

concentrated in vacuo, and the product can be purified by standard means, e.g., by chromatography using silica gel (e.g., flash column chromatography).

Compounds of formula II are prepared by methods known in the art, for example by methods disclosed in WO 95/10516, in U.S. 5,151,423 and those described below. Compounds of formula II wherein the C-3 position of the pyridine ring in the tricyclic structure is substituted by bromo can also be prepared by a procedure comprising the following steps:


(a) reacting an amide of the formula



10


wherein R^{11a} is Br, R^{5a} is hydrogen and R^{6a} is C_1 - C_6 alkyl, aryl or heteroaryl; R^{5a} is C_1 - C_6 alkyl, aryl or heteroaryl and R^{6a} is hydrogen; R^{5a} and R^{6a} are independently selected from the group consisting of C_1 - C_6 alkyl and aryl; or R^{5a} and R^{6a} , together with the nitrogen to which they are attached, form a ring comprising 4 to 6 carbon atoms or comprising 3 to 5 carbon atoms and one hetero moiety selected from the group consisting of -O- and - NR^{9a} -, wherein R^{9a} is H, C_1 - C_6 alkyl or phenyl; with a compound of the formula

15



20

wherein R^{1a} , R^{2a} , R^{3a} and R^{4a} are independently selected from the group consisting of hydrogen and halo and R^{7a} is Cl or Br, in the presence of a strong base to obtain a compound of the formula




(b) reacting a compound of step (a) with


25

(i) $POCl_3$ to obtain a cyano compound of the formula


- 9 -




(ii) DIBALH to obtain an aldehyde of the formula



(c) reacting the cyano compound or the aldehyde with a piperidine
5 derivative of the formula




wherein L is a leaving group selected from the group consisting of Cl and Br, to obtain an aldehyde or an alcohol of the formula below, respectively:



10 (d)(i) cyclizing the aldehyde with $\text{CF}_3\text{SO}_3\text{H}$ to obtain a compound of formula II wherein the dotted line represents a double bond; or

(d)(ii) cyclizing the alcohol with polyphosphoric acid to obtain a compound of formula II wherein the dotted line represents a single bond.


Methods for preparing compounds of formula II disclosed in WO
15 95/10516, U.S. 5,151,423 and described below employ a tricyclic ketone intermediate. Such intermediates of the formula



- 10 -


wherein R^{11b} , R^{1a} , R^{2a} , R^{3a} and R^{4a} are independently selected from the group consisting of hydrogen and halo, can be prepared by the following process comprising :

(a) reacting a compound of the formula




5

(i) with an amine of the formula $NHR^{5a}R^{6a}$, wherein R^{5a} and R^{6a} are as defined in the process above; in the presence of a palladium catalyst and carbon monoxide to obtain an amide of the formula:



10


(ii) with an alcohol of the formula $R^{10a}OH$, wherein R^{10a} is C_1 - C_6 lower alkyl or C_3 - C_6 cycloalkyl, in the presence of a palladium catalyst and carbon monoxide to obtain the ester of the formula



followed by reacting the ester with an amine of formula $NHR^{5a}R^{6a}$ to obtain the amide;


15

(b) reacting the amide with an iodo-substituted benzyl compound of the formula



wherein R^{1a} , R^{2a} , R^{3a} , R^{4a} and R^{7a} are as defined above, in the presence of a strong base to obtain a compound of the formula

20



(c) cyclizing a compound of step (b) with a reagent of the formula $R^{8a}MgL$, wherein R^{8a} is C_1 - C_8 alkyl, aryl or heteroaryl and L is Br or Cl ,

- 11 -


provided that prior to cyclization, compounds wherein R^{5a} or R^{6a} is hydrogen are reacted with a suitable N-protecting group.

(+)-Isomers of compounds of formula II wherein X is CH can be prepared with high enantioselectivity by using a process comprising enzyme catalyzed transesterification. Preferably, a racemic compound of formula II, wherein X is C, the double bond is present and R³ is not H, is reacted with an enzyme such as Toyobo LIP-300 and an acylating agent such as trifluoroethyl isobutyrate; the resultant (+)-amide is then hydrolyzed, for example by refluxing with an acid such as H₂SO₄, to obtain the corresponding optically enriched (+)-isomer wherein X is CH and R³ is not H. Alternatively, a racemic compound of formula II, wherein X is C, the double bond is present and R³ is not H, is first reduced to the corresponding racemic compound of formula II wherein X is CH and then treated with the enzyme (Toyobo LIP-300) and acylating agent as described above to obtain the (+)-amide, which is hydrolyzed to obtain the optically enriched (+)-isomer.

Compounds of formula III can be prepared from compounds of formula II by procedures known in the art, for example by reacting 1-N-t-butoxy-carbonylpiperidiny-4-acetic acid with the compound of formula II under the standard amide coupling conditions described above.


Compounds useful in this invention are exemplified by the following preparative examples, which should not be construed to limit the scope of the disclosure. Alternative mechanistic pathways and analogous structures within the scope of the invention may be apparent to those skilled in the art.

PREPARATIVE EXAMPLE 1




Step A:

- 12 -



Combine 25.86 g (55.9 mmol) of 4-(8-chloro-3-bromo-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidine-1-carboxylic acid ethyl ester and 250 mL of concentrated H₂SO₄ at -5°C, then add 4.8 g (56.4 mmol) of NaNO₃ and stir for 2 hours. Pour the mixture into 600 g of ice and basify with concentrated NH₄OH (aqueous). Filter the mixture, wash with 300 mL of water, then extract with 500 mL of CH₂Cl₂. Wash the extract with 200 mL of water, dry over MgSO₄, then filter and concentrate *in vacuo* to a residue. Chromatograph the residue (silica gel, 10% EtOAc/ CH₂Cl₂) to give 24.4 g (86% yield) of the product. m.p. = 165-167°C, Mass Spec.: MH⁺ = 506, 508 (Cl).
 elemental analysis: calculated - C, 52.13; H, 4.17; N, 8.29
 found - C, 52.18; H, 4.51; N, 8.16

Step B:


15

Combine 20 g (40.5 mmol) of the product of Step A and 200 mL of concentrated H₂SO₄ at 20°C, then cool the mixture to 0°C. Add 7.12 g (24.89 mmol) of 1,3-dibromo-5,5-dimethyl-hydantoin to the mixture and stir for 3 hours at 20°C. Cool to 0°C, add an additional 1.0 g (3.5 mmol) of the dibromohydantoin and stir at 20°C for 2 hours. Pour the mixture into 400 g of ice, basify with concentrated NH₄OH (aqueous) at 0°C, and collect the resulting solid by filtration. Wash the solid with 300 mL of water, slurry in 200 mL of acetone and filter to provide 19.79 g (85.6% yield) of the product. m.p. = 236-237°C, Mass Spec.: MH⁺ = 586 (Cl).


20

- 13 -

elemental analysis: calculated - C, 45.11; H, 3.44; N, 7.17
 found - C, 44.95; H, 3.57; N, 7.16

Step C:

- 5 Combine 25 g (447 mmol) of Fe filings, 10 g (90 mmol) of CaCl_2 and a suspension of 20 g (34.19 mmol) of the product of Step B in 700 mL of 90:10 EtOH/water at 50°C . Heat the mixture at reflux overnight, filter through Celite® and wash the filter cake with 2 X 200 mL of hot EtOH. Combine the filtrate and washes, and concentrate *in vacuo* to a residue.
- 10 Extract the residue with 600 mL of CH_2Cl_2 , wash with 300 mL of water and dry over MgSO_4 . Filter and concentrate *in vacuo* to a residue, then chromatograph (silica gel, 30% EtOAc/ CH_2Cl_2) to give 11.4 g (60% yield) of the product. m.p. = $211-212^\circ\text{C}$, Mass Spec.: $\text{MH}^+ = 556$ (Cl).
 elemental analysis: calculated - C, 47.55; H, 3.99; N, 7.56
 found - C, 47.45; H, 4.31; N, 7.49
- 15

Step D:


- 20 Slowly add (in portions) 20 g (35.9 mmol) of the product of Step C to a solution of 8 g (116 mmol) of NaNO_2 in 120 mL of concentrated HCl (aqueous) at -10°C . Stir the resulting mixture at 0°C for 2 hours, then slowly add (dropwise) 150 mL (1.44 mole) of 50% H_3PO_2 at 0°C over a 1 hour period. Stir at 0°C for 3 hours, then pour into 600 g of ice and basify with concentrated NH_4OH (aqueous). Extract with 2 X 300 mL of CH_2Cl_2 .

- 14 -


dry the extracts over MgSO_4 , then filter and concentrate *in vacuo* to a residue. Chromatograph the residue (silica gel, 25% EtOAc/ hexanes) to give 13.67 g (70% yield) of the product. m.p. = 163-165°C, Mass Spec.: $\text{MH}^+ = 541$ (CI).

5 elemental analysis: calculated - C, 48.97; H, 4.05; N, 5.22
found - C, 48.86; H, 3.91; N, 5.18


10

Step E:

Combine 6.8 g (12.59 mmol) of the product of Step D and 100 mL
15 of concentrated HCl (aqueous) and stir at 85°C overnight. Cool the
mixture, pour it into 300 g of ice and basify with concentrated NH_4OH
(aqueous). Extract with 2 x 300 mL of CH_2Cl_2 , then dry the extracts over
 MgSO_4 . Filter, concentrate *in vacuo* to a residue, then chromatograph
(silica gel, 10% MeOH/EtOAc + 2% NH_4OH (aq.)) to give 5.4 g (92% yield)
20 of the title compound. m.p. = 172-174°C, Mass Spec.: $\text{MH}^+ = 469$ (FAB).
elemental analysis: calculated - C, 48.69; H, 3.65; N, 5.97
found - C, 48.83; H, 3.80; N, 5.97


PREPARATIVE EXAMPLE 225 Step A:

- 15 -




Hydrolyze 2.42 g of 4-(8-chloro-3-bromo-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidine-1-carboxylic acid ethyl ester by dissolving in concentrated HCl and heating to about 100°C for @ 16 hours. Cool the mixture, the neutralize with 1 M NaOH (aqueous). Extract with CH₂Cl₂, dry the extracts over MgSO₄, filter and concentrate *in vacuo* to give 1.39 g (69% yield) of the product.

Step B:




Combine 1 g (2.48 mmol) of the product of Step A and 25 mL of dry toluene, add 2.5 mL of 1 M DIBAL in toluene and heat the mixture at reflux. After 0.5 hours, add another 2.5 mL of 1 M DIBAL in toluene and heat at reflux for 1 hour. (The reaction is monitored by TLC using 50% MeOH/CH₂Cl₂ + NH₄OH (aqueous).) Cool the mixture to room temperature, add 50 mL of 1 N HCl (aqueous) and stir for 5 min. Add 100 mL of 1 N NaOH (aqueous), then extract with EtOAc (3 X 150 mL). Dry the extracts over MgSO₄, filter and concentrate *in vacuo* to give 1.1 g of the title compound.

PREPARATIVE EXAMPLE 3




- 16 -

[racemic as well as (+)- and (-)-isomers]

Step A:

- Combine 16.6 g (0.03 mole) of the product of Preparative Example 1, Step D, with a 3:1 solution of CH₃CN and water (212.65 mL CH₃CN and 70.8 mL of water) and stir the resulting slurry overnight at room temperature. Add 32.833 g (0.153 mole) of NaIO₄ and then 0.31 g (2.30 mmol) of RuO₂ and stir at room temperature give 1.39 g (69% yield) of the product. (The addition of RuO is accompanied by an exothermic reaction and the temperature climbs from 20° to 30°C.) Stir the mixture for 1.3 hrs. (temperature returned to 25°C after about 30 min.), then filter to remove the solids and wash the solids with CH₂Cl₂. Concentrate the filtrate *in vacuo* to a residue and dissolve the residue in CH₂Cl₂. Filter to remove insoluble solids and wash the solids with CH₂Cl₂. Wash the filtrate with water, concentrate to a volume of about 200 mL and wash with bleach, then with water. Extract with 6 N HCl (aqueous). Cool the aqueous extract to 0°C and slowly add 50% NaOH (aqueous) to adjust to pH = 4 while keeping the temperature <30°C. Extract twice with CH₂Cl₂, dry over MgSO₄ and concentrate *in vacuo* to a residue. Slurry the residue in 20 mL of EtOH and cool to 0°C. Collect the resulting solids by filtration and dry the solids *in vacuo* to give 7.95 g of the product. ¹H NMR (CDCl₃, 200 MHz): 8.7 (s, 1H); 7.85 (m, 6H); 7.5 (d, 2H); 3.45 (m, 2H); 3.15 (m, 2H).


Step B:

- Combine 21.58 g (53.75 mmol) of the product of Step A and 500 mL of an anhydrous 1:1 mixture of EtOH and toluene, add 1.43 g (37.8 mmol) of NaBH₄ and heat the mixture at reflux for 10 min. Cool the

- 17 -

5 mixture to 0°C, add 100 mL of water, then adjust to pH≈ 4-5 with 1 M HCl (aqueous) while keeping the temperature <10°C. Add 250 mL of EtOAc and separate the layers. Wash the organic layer with brine (3 X 50 mL) then dry over Na₂SO₄. Concentrate *in vacuo* to a residue (24.01 g) and chromatograph the residue (silica gel, 30 % hexane/CH₂Cl₂) to give the product. Impure fractions were purified by rechromatography. A total of 18.57 g of the product was obtained. ¹H NMR (DMSO-d₆, 400 MHz): 8.5 (s, 1H); 7.9 (s, 1H); 7.5 (d of d, 2H); 6.2 (s, 1H); 6.1 (s, 1H); 3.5 (m, 1H); 3.4 (m, 1H); 3.2 (m, 2H).

10


Step C:

15 Combine 18.57 g (46.02 mmol) of the product of Step B and 500 mL of CHCl₃, then add 6.70 mL (91.2 mmol) of SOCl₂, and stir the mixture at room temperature for 4 hrs. Add a solution of 35.6 g (0.413 mole) of piperazine in 800 mL of THF over a period of 5 min. and stir the mixture for 1 hr. at room temperature. Heat the mixture at reflux overnight, then cool to room temperature and dilute the mixture with 1 L of CH₂Cl₂. Wash with water (5 X 200 mL), and extract the aqueous wash with CHCl₃ (3 X 100 mL). Combine all of the organic solutions, wash with brine (3 X 200 mL) and dry over MgSO₄. Concentrate *in vacuo* to a residue and chromatograph (silica gel, gradient of 5%, 7.5%, 10% MeOH/CH₂Cl₂ + NH₄OH) to give 18.49 g of the title compound as a racemic mixture.

25

Step D - Separation of Enantiomers:

- 18 -




The racemic title compound of Step C is separated by preparative chiral chromatography (Chiralpak AD, 5 cm X 50 cm column, flow rate 100 mL/min., 20% iPrOH/hexane + 0.2% diethylamine), to give 9.14 g of the (+)-isomer and 9.30 g of the (-)-isomer. Chiralpak is a trade-mark.


Physical chemical data for (+)-isomer: m.p. = 74.5°-77.5°C; Mass Spec. MH^+ = 471.9; $[\alpha]_D^{25}$ = +97.4° (8.48 mg/ 2mL MeOH).

Physical chemical data for (-)-isomer: m.p. = 82.9°-84.5°C; Mass Spec. MH^+ = 471.8; $[\alpha]_D^{25}$ = -97.4° (8.32 mg/ 2mL MeOH).

PREPARATIVE EXAMPLE 4




Step A:



- 19 -


Combine 15 g (38.5 mmol) of 4-(8-chloro-3-bromo-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidine-1-carboxylic acid ethyl ester and 150 mL of conc. H_2SO_4 at -5°C , then add 3.89 g (38.5 mmol) of KNO_3 and stir for 4 h. Pour the mixture into 3 L of ice and basify with 50% NaOH (aqueous). Extract with CH_2Cl_2 , dry over MgSO_4 , then filter and concentrate *in vacuo* to a residue. Recrystallize the residue from acetone to give 6.69 g of the product. ^1H NMR (CDCl_3 , 200 MHz): 8.5 (s, 1H); 7.75 (s, 1H); 7.6 (s, 1H); 7.35 (s, 1H); 4.15 (q, 2H); 3.8 (m, 2H); 3.5-3.1 (m, 4H); 3.0-2.8 (m, 2H); 2.6-2.2 (m, 4H); 1.25 (t, 3H).

10 Step B:



Combine 6.69 g (13.1 mmol) of the product of Step A and 100 mL of 85% EtOH /water, add 0.66 g (5.9 mmol) of CaCl_2 and 6.56 g (117.9 mmol) of Fe and heat the mixture at reflux overnight. Filter the hot reaction mixture through celite® and rinse the filter cake with hot EtOH . Concentrate the filtrate *in vacuo* to give 7.72 g of the product. Mass Spec.: $\text{MH}^+ = 478.0$

Step C:



20 Combine 7.70 g of the product of Step B and 35 mL of HOAc , then add 45 mL of a solution of Br_2 in HOAc and stir the mixture at room temperature overnight. Add 300 mL of 1 N NaOH (aqueous), then 75 mL of 50% NaOH (aqueous) and extract with EtOAc . Dry the extract over MgSO_4 and concentrate *in vacuo* to a residue. Chromatograph the residue (silica gel, 20%-30% EtOAc /hexane) to give 3.47 g of the product


- 20 -

(along with another 1.28 g of partially purified product).

Mass Spec.: $MH^+ = 555.9$.


^1H NMR ($CDCl_3$, 300 MHz): 8.5 (s, 1H); 7.5 (s, 1H); 7.15 (s, 1H); 4.5 (s, 2H); 4.15 (m, 3H); 3.8 (br s, 2H); 3.4-3.1 (m, 4H); 9-2.75 (m, 1H); 2.7-2.5 (m, 2H); 2.4-2.2 (m, 2H); 1.25 (m, 3H).

Step D:



Combine 0.557 g (5.4 mmol) of t-butylnitrite and 3 mL of DMF, and heat the mixture at to 60°-70°C. Slowly add (dropwise) a mixture of 2.00 g (3.6 mmol) of the product of Step C and 4 mL of DMF, then cool the mixture to room temperature. Add another 0.64 mL of t-butylnitrite at 40°C and reheat the mixture to 60°-70°C for 0.5 hrs. Cool to room temperature and pour the mixture into 150 mL of water. Extract with CH_2Cl_2 , dry over $MgSO_4$ and concentrate *in vacuo* to a residue. Chromatograph the residue (silica gel, 10%-20% EtOAc/hexane) to give 0.74 g of the product. Mass Spec.: $MH^+ = 541.0$. ^1H NMR ($CDCl_3$, 200 MHz): 8.52 (s, 1H); 7.5 (d, 2H); 7.2 (s, 1H); 4.15 (q, 2H); 3.9-3.7 (m, 2H); 3.5-3.1 (m, 4H); 3.0-2.5 (m, 2H); 2.4-2.2 (m, 2H); 2.1-1.9 (m, 2H); 1.26 (t, 3H).

Step E:




Combine 0.70 g (1.4 mmol) of the product of Step D and 8 mL of concentrated HCl (aqueous) and heat the mixture at reflux overnight. Add 30 mL of 1 N NaOH (aqueous), then 5 mL of 50% NaOH (aqueous) and extract with CH_2Cl_2 . Dry the extract over $MgSO_4$ and concentrate *in*

- 21 -

vacuo to give 0.59 g of the title compound. Mass Spec.: $M^+ = 468.7$. m.p. = 123.9°-124.2°C.


PREPARATIVE EXAMPLE 5



5

[racemic as well as (+)- and (-)-isomers]


Step A:



Prepare a solution of 8.1 g of the title compound from Preparative Example 4 in toluene and add 17.3 mL of a 1M solution of DIBAL in toluene. Heat the mixture at reflux and slowly add (dropwise) another 21 mL of 1 M DIBAL/toluene solution over a period of 40 min. Cool the reaction mixture to about 0°C and add 700 mL of 1 M HCl (aqueous). Separate and discard the organic phase. Wash the aqueous phase with CH_2Cl_2 , discard the extract, then basify the aqueous phase by adding 50% NaOH (aqueous). Extract with CH_2Cl_2 , dry the extract over MgSO_4 and concentrate *in vacuo* to give 7.30 g of the title compound, which is a racemic mixture of enantiomers.

Step B - Separation of Enantiomers:

- 22 -




The racemic title compound of Step A is separated by preparative chiral chromatography (Chiralpack AD, 5 cm X 50 cm column, using 20% iPrOH/hexane + 0.2% diethylamine), to give the (+)-isomer and the (-)-isomer of the title compound.

Physical chemical data for (+)-isomer: m.p. = 148.8°C;
Mass Spec. MH^+ = 469; $[\alpha] = +65.6^\circ$ (mg/ 2mL MeOH).

Physical chemical data for (-)-isomer: m.p. = 112°C;
Mass Spec. MH^+ = 469; $[\alpha] = -65.2^\circ$ (mg/ 2mL MeOH).

10


PREPARATIVE EXAMPLE 6



[racemic as well as (+)- and (-)-isomers]


Step A:

- 23 -




Combine 40.0 g (0.124 mole) of the starting ketone and 200 mL of H_2SO_4 and cool to 0°C . Slowly add 13.78 g (0.136 mole) of KNO_3 over a period of 1.5 hrs., then warm to room temperature and stir overnight. Work up the reaction using substantially the same procedure as described for Preparative Example 1, Step A. Chromatograph (silica gel, 20%, 30%, 40%, 50% EtOAc/hexane, then 100% EtOAc) to give 28 g of the 9-nitro product, along with a smaller quantity of the 7-nitro product and 19 g of a mixture of the 7-nitro and 9-nitro compounds.

10 Step B:



React 28 g (76.2 mmol) of the 9-nitro product of Step A, 400 mL of 85% EtOH/water, 3.8 g (34.3 mmol) of CaCl_2 and 38.28 g (0.685 mole) of Fe using substantially the same procedure as described for Preparative Example 1, Step C, to give 24 g of the product


Step C:




Combine 13 g (38.5 mmol) of the product of Step B, 140 mL of HOAc and slowly add a solution of 2.95 mL (57.8 mmol) of Br_2 in 10 mL of HOAc over a period of 20 min. Stir the reaction mixture at room temperature, then concentrate *in vacuo* to a residue. Add CH_2Cl_2 and water, then adjust to pH = 8-9 with 50% NaOH (aqueous). Wash the

- 24 -


organic phase with water, then brine and dry over Na_2SO_4 . Concentrate *in vacuo* to give 11.3 g of the product.

Step D:

- 5 Cool 100 mL of concentrated HCl (aqueous) to 0°C , then add 5.61 g (81.4 mmol) of NaNO_2 and stir for 10 min. Slowly add (in portions) 11.3 g (27.1 mmol) of the product of Step C and stir the mixture at $0^\circ\text{--}3^\circ\text{C}$ for 2.25 hrs. Slowly add (dropwise) 180 mL of 50% H_3PO_2 (aqueous) and allow the mixture to stand at 0°C overnight. Slowly add (dropwise) 150 mL of 50% NaOH over 30 min., to adjust to pH = 9, then extract with CH_2Cl_2 . Wash the extract with water, then brine and dry over Na_2SO_4 . Concentrate *in vacuo* to a residue and chromatograph (silica gel, 2% EtOAc/ CH_2Cl_2) to give 8.6 g of the product.

Step E:

- 15 Combine 8.6 g (21.4 mmol) of the product of Step D and 300 mL of MeOH and cool to $0^\circ\text{--}2^\circ\text{C}$. Add 1.21 g (32.1 mmol) of NaBH_4 and stir at $\sim 0^\circ\text{C}$ for 1 hr. Add another 0.121 g (3.21 mmol) of NaBH_4 , stir for 2 hr. at 0°C , then let stand overnight at 0°C . Concentrate *in vacuo* to a residue then partition the residue between CH_2Cl_2 and water. Separate the organic phase and concentrate *in vacuo* (50°C) to give 8.2 g of the product.

Step F:

- 25 Combine 8.2 g (20.3 mmol) of the product of Step E and 160 mL of CH_2Cl_2 , cool to 0°C , then slowly add (dropwise) 14.8 mL (203 mmol) of SOCl_2 over a 30 min. period. Warm the mixture to room temperature and


- 25 -

stir for 4.5 hrs., then concentrate *in vacuo* to a residue, add CH_2Cl_2 and wash with 1 N NaOH (aqueous) then brine and dry over Na_2SO_4 .

Concentrate *in vacuo* to a residue, then add dry THF and 8.7 g (101 mmol) of piperazine and stir at room temperature overnight. Concentrate *in*

5 *vacuo* to a residue, add CH_2Cl_2 , and wash with 0.25 N NaOH (aqueous), water, then brine. Dry over Na_2SO_4 and concentrate *in vacuo* to give 9.46 g of the crude product. Chromatograph (silica gel, 5% MeOH/ CH_2Cl_2 + NH_3) to give 3.59 g of the title compound, as a racemate. ^1H NMR (CDCl₃, 200 MHz): 8.43 (d, 1H); 7.55 (d, 1H); 7.45 (d, 1H); 7.11 (d, 1H);
10 5.31 (s, 1H); 4.86-4.65 (m, 1H); 3.57-3.40 (m, 1H); 2.98-2.55 (m, 6H); 2.45-2.20 (m, 5H).

Step G - Separation of Enantiomers:




15 The racemic title compound from Step F (5.7 g) is chromatographed as described for Preparative Example 3, Step D, using 30% iPrOH/hexane + 0.2% diethylamine, to give 2.88 g of the R-(+)-isomer and 2.77 g of the S-(-)-isomer of the title compound.


Physical chemical data for the R-(+)-isomer: Mass Spec. $\text{MH}^+ = 470$; $[\alpha] = +12.1^\circ$ (10.9 mg/ 2mL MeOH).

20 Physical chemical data for the S-(-)-isomer: Mass Spec. $\text{MH}^+ = 470$; $[\alpha] = -13.2^\circ$ (11.51 mg/ 2mL MeOH).

- 26 -



[racemic as well as (+)- and (-)-isomers]


Step A:

- 5 Combine 13 g (33.3 mmol) of the title compound from Preparative Example 1, Step D, and 300 mL of toluene at 20°C, then add 32.5 mL (32.5 mmol) of a 1 M solution of DIBAL in toluene. Heat the mixture at reflux for 1 hr., cool to 20°C, add another 32.5 mL of 1 M DIBAL solution and heat at reflux for 1 hr. Cool the mixture to 20°C and pour it into a
- 10 mixture of 400 g of ice, 500 mL of EtOAc and 300 mL of 10% NaOH (aqueous). Extract the aqueous layer with CH₂Cl₂ (3 x 200 mL), dry the organic layers over MgSO₄, then concentrate *in vacuo* to a residue. Chromatograph (silica gel, 12% MeOH/CH₂Cl₂ + 4% NH₄OH) to give 10.4 g of the title compound as a racemate. Mass Spec.: MH⁺ = 469 (FAB).
- 15 partial ¹H NMR (CDCl₃, 400 MHz): 8.38 (s, 1H); 7.57 (s, 1H); 7.27 (d, 1H); 7.06-(d, 1H); 3.95 (d, 1H).

20

25 Step B - Separation of Enantiomers:

- 27 -



The racemic title compound of Step A is separated by preparative chiral chromatography (Chiralpack AD, 5 cm X 50 cm column, using 5% iPrOH/hexane + 0.2% diethylamine), to give the (+)-isomer and the (-)-isomer of the title compound.


Physical chemical data for (+)-isomer: Mass Spec.

MH⁺ = 470.9 (FAB); [α] = +43.5° (c=0.402, EtOH); partial ¹H NMR (CDCl₃, 400 MHz): 8.38 (s, 1H); 7.57 (s, 1H); 7.27 (d, 1H); 7.05 (d, 1H); 3.95 (d, 1H).

Physical chemical data for (-)-isomer: Mass Spec.

MH⁺ = 470.9 (FAB); [α] = -41.8° (c=0.328 EtOH); partial ¹H NMR (CDCl₃, 400 MHz): 8.38 (s, 1H); 7.57 (s, 1H); 7.27 (d, 1H); 7.05 (d, 1H); 3.95 (d, 1H).

PREPARATIVE EXAMPLE 8



[racemic as well as R-(+)- and S-(-)-isomers]

Treat 4-(8-chloro-3-bromo-5,6-dihydro-11H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-ylidene)-1-piperidine-1-carboxylic acid ethyl ester via substantially the same procedure as described in Preparative Example 3, Steps A-D, to give as the product of Step C, the racemic title compound,


- 28 -

and as the products of Step D the R-(+)-isomer and S-(-)-isomer of the title compound.


Physical chemical data for the R-(+)-isomer: ^{13}C NMR (CDCl_3):
 155.8 (C); 146.4 (CH); 140.5 (CH); 140.2 (C); 136.2 (C); 135.3 (C);
 5 133.4 (C); 132.0 (CH); 129.9 (CH); 125.6 (CH); 119.3 (C); 79.1 (CH);
 52.3 (CH_2); 52.3 (CH); 45.6 (CH_2); 45.6 (CH_2); 30.0 (CH_2); 29.8 (CH_2).
 $[\alpha] = +25.8^\circ$ (8.46 mg/2 mL MeOH).

Physical chemical data for the S-(-)-isomer: ^{13}C NMR (CDCl_3):
 155.9 (C); 146.4 (CH); 140.5 (CH); 140.2 (C); 136.2 (C); 135.3 (C);
 10 133.3 (C); 132.0 (CH); 129.9 (CH); 125.5 (CH); 119.2 (C); 79.1 (CH);
 52.5 (CH_2); 52.5 (CH); 45.7 (CH_2); 45.7 (CH_2); 30.0 (CH_2); 29.8 (CH_2).
 $[\alpha] = -27.9^\circ$ (8.90 mg/2 mL MeOH).

PREPARATIVE EXAMPLE 9




15 Step A:




Dissolve 9.90 g (18.9 mmol) of the product of Preparative Example 4, Step B, in 150 mL CH_2Cl_2 and 200 mL of CH_3CN and heat to 60°C . Add 2.77 g (20.8 mmol) N-chlorosuccinimide and heat to reflux for 3 h.,
 20 monitoring the reaction by TLC (30% EtOAc/ H_2O). Add an additional 2.35 g (10.4 mmol) of N-chlorosuccinimide and reflux an additional 45 min. Cool the reaction mixture to room temperature and extract with 1N NaOH and CH_2Cl_2 . Dry the CH_2Cl_2 layer over MgSO_4 , filter and purify by flash chromatography (1200 mL normal phase silica gel, eluting with 30%
 25 EtOAc/ H_2O) to obtain 6.24 g of the desired product. M.p. $193-195.4^\circ\text{C}$.
Step B:

- 29 -




To 160 mL of conc. HCl at -10°C add 2.07 g (30.1 mmol) NaNO_2 and stir for 10 min. Add 5.18 g (10.1 mmol) of the product of Step A and warm the reaction mixture from -10°C to 0°C for 2 h. Cool the reaction to -10°C , add 100 mL H_3PO_2 and let stand overnight. To extract the reaction mixture, pour over crushed ice and basify with 50% $\text{NaOH}/\text{CH}_2\text{Cl}_2$. Dry the organic layer over MgSO_4 , filter and concentrate to dryness. Purify by flash chromatography (600 mL normal phase silica gel, eluting with 20% EtOAc/hexane) to obtain 3.98 g of product. Mass spec.: $\text{MH}^+=497.2$.

10 Step C:



Dissolve 3.9 g of the product of Step B in 100 mL conc. HCl and reflux overnight. Cool the mixture, basify with 50 % w/w NaOH and extract the resultant mixture with CH_2Cl_2 . Dry the CH_2Cl_2 layer over MgSO_4 , evaporate the solvent and dry under vacuum to obtain 3.09 g of the desired product. Mass spec.: $\text{MH}^+=424.9$.


Step D:




- 30 -

Using a procedure similar to that described in Preparative Example 5, obtain 1.73 g of the desired product, m.p. 169.6-170.1°C; $[\alpha]_D^{25} = +48.2^\circ$ (c=1, MeOH).

PREPARATIVE EXAMPLE 10



5

Step A:

Combine 1.33 g of the (+)-enantiomer of the compound of Preparative Example 5, Step B, in anhydrous DMF with 1.37 g of 1-N-t-butoxy-carbonylpiperidiny-4-acetic acid, and with DEC, HOBT and N-methylmorpholine. Stir the mixture at room temperature overnight. Concentrate *in vacuo* to remove the DMF and add 50 mL of saturated NaHCO_3 (aqueous). Extract with CH_2Cl_2 (2 X 250 mL), wash the extracts with 50 mL of brine and dry over MgSO_4 . Concentrate *in vacuo* to a residue and chromatograph (silica gel, 2% $\text{CH}_3\text{OH}/\text{CH}_2\text{Cl}_2$ + 10% NH_4OH) to give 2.78 g of the product. Mass Spec.: $\text{MH}^+ = 694.0$ (FAB); $[\alpha]_D^{25} = +34.1^\circ$ (5.45 mg/2 mL, MeOH).

15

Step B:

Combine 2.78 g of the product of Step A and CH_2Cl_2 , then cool to 0°C and add TFA. Stir the mixture for 3 h at 0°C, then add 1 N NaOH (aqueous) followed by 50% NaOH (aqueous). Extract with CH_2Cl_2 , dry over MgSO_4 and concentrate *in vacuo* to give 1.72 g of the product. M.p. = 104.1°C; Mass Spec.: $\text{MH}^+ = 594$; $[\alpha]_D^{25} = +53.4^\circ$ (11.42 mg/2 mL, CH_3OH).


20

25


- 31 -

PREPARATIVE EXAMPLE 11

(+)-1-(Aminoacetyl)-4-(3-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) piperidine



- 5 Step 1: (+)-1,1-Dimethylethyl[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl]-carbamate



- 10 The product of Preparative Example 5, (+-isomer) (0.4 g, 0.85 mmol), was dissolved in DMF (10mL) and then cooled to ~ 4 °C. BOC-Glycine (0.19 g, 1.1 mmol) was then added, followed by DEC (0.2 g, 1.1 mmol), HOBT (0.15 g, 1.1 mmol) and 4-methylmorpholine (0.11 g, 0.12 μL, 1.1 mmol). The reaction was stirred at room temperature overnight, then was concentrated in vacuo to a residue and partitioned between CH₂Cl₂ and sat. NaHCO₃
- 15 (aqueous). The aqueous phase was extracted further with CH₂Cl₂, the combined CH₂Cl₂ fractions were dried over MgSO₄ and concentrated in vacuo to give a residue that was chromatographed on silica gel column using 5 % (NH₃ saturated CH₃OH)/ CH₂Cl₂ as eluent to give the title compound as a white solid: 0.52 g, 99% yield, m.p. = 95-96 °C, MH⁺ = 628.

20 Step 2:


- The product of Step 1 (2.65 g, 4.2 mmol) was dissolved in CH₂Cl₂ (20 mL) and cooled to 0 °C. Trifluoroacetic acid (10 mL) was then added. The reaction mixture was stirred at room temperature for 4h, then poured into ice and the pH was adjusted to 10 using 50%(w/v) aqueous NaOH.
- 25 The reaction mixture was extracted with CH₂Cl₂, the combined CH₂Cl₂ extracts were washed with H₂O and brine, and dried over Na₂SO₄. The

- 32 -


solvents were removed by rotary evaporation to give the title compound as a white solid: 2.18 g, 98% yield, m.p. = 150-152 °C, MH^+ = 528.

PREPARATIVE EXAMPLE 12

5 (+)-1-(3-Amino-1-oxopropyl)-4-(3-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) piperidine



Step 1: (+)-1,1-Dimethylethyl[3-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxopropyl]-carbamate



10


The title compound was prepared following essentially the same procedure as described in Preparative Example 11, Step 1, except that BOC-β-alanine was used instead of BOC-glycine to obtain a white solid. Yield = 99 %, MH^+ = 642.

15 Step 2:

The title compound was prepared following essentially the same procedure as described in Preparative Example 11, Step 2, to obtain a white solid. Yield = 100 %, m.p. = 136-137 °C, MH^+ = 642.


PREPARATIVE EXAMPLE 13

20 (+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-[4-amino]-1-oxobutyl] piperidine



- 33 -

Step 1: (+)-1,1-dimethylethyl[4-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxobutyl]-carboxamide




5 The title compound was prepared following essentially the same procedure as described in Preparative Example 11, Step 1, except that BOC- α -amino butyric acid was used instead of BOC-glycine to obtain a white solid. Yield = 79 %, m.p. = 102-103 °C, MH⁺ = 781.

Step 2:


10 The title compound was prepared following essentially the same procedure as described for Preparative Example 11, Step 2, to obtain a white solid. Yield = 94 %, m.p. = 114-115 °C, MH⁺ = 681.

PREPARATIVE EXAMPLE 14

15 (+)-1-(Aminoacetyl)-4-[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl] piperidine



20 Step 1: (+)-1,1-Dimethylethyl[2-[4-[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl]-1-piperidiny]-2-oxoethyl] carbamate



The title compound was prepared following essentially the same procedure as described for Preparative Example 11, Step 1, except that


- 34 -

the compound of Preparative Example 10 -(+)-isomer) was used instead of the compound from Preparative Example 5 to obtain a white solid. Yield = 82 %, m.p. = 98-99 °C, MH^+ = 753.


Step 2: The title compound was prepared following essentially the same procedure as described in Preparative Example 11, Step 2 to obtain a white solid. Yield = 89 %, m.p. = 130-131 °C, MH^+ = 653.

PREPARATIVE EXAMPLE 15

(+)-1-(3-Amino-1-oxopropyl)-4-[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl] piperidine



Step 1: (+)-1,1-Dimethylethyl[3-[4-[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl]-1-piperidiny]-3-oxopropyl] carbamate




The title compound was prepared following essentially the same procedure as described for Preparative Example 12, Step 1, except that the compound from Preparative Example 10 was used instead of the compound of Preparative Example 5 to obtain a white solid. Yield = 84 %, m.p. = 87-88 °C, MH^+ = 767.

Step 2: The title compound was prepared following essentially the same procedure as described for Preparative Example 11, Step 2, to obtain a white solid. Yield = 84 %, m.p. = 120-121 °C, MH^+ = 667.

- 35 -


PREPARATIVE EXAMPLE 16

(+)-1-(4-Amino-1-oxobutyl)-4-[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl] piperidine



5

Step 1: (+)-1,1-Dimethylethyl[4-[4-[2-[4-(3,10-dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclo-hepta[1,2-b]pyridin-11-yl) -1-piperidiny]-2-oxoethyl]-1-piperidiny]-4-oxobutyl] carbamate




10

The title compound was prepared following essentially the same procedure as described for Preparative Example 13, Step 1, except that the compound of Preparative Example 10 was used instead of the compound of Preparative Example 5 to obtain a white solid. Yield = 79 %, m.p. = 102-103 °C, MH^+ = 782.

15

Step2: The title compound was prepared following essentially the same procedure as described for Preparative Example 11, Step 2, to obtain a white solid. Yield = 94 %, m.p. = 114-115 °C, MH^+ = 681.

PREPARATIVE EXAMPLE 17

20


Dissolve 2 g (12.7 mmol) of methyl 2-oxo-1-pyrrolidine acetate in 20 mL of EtOH and then add 20 mL of 1M LiOH. Stir the reaction mixture at room temperature for 16 h. Strip off the solvents, dissolve the resulting material in water and adjust the pH to ~ 4. Concentrate the reaction mixture to give the product. Mass Spec.: MH^+ = 144.

25

- 36 -

EXAMPLE 1


(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[(2-oxo-1-pyrrolidiny]acetyl]piperidine



5 The compound of Preparative Example 15 (0.15 g, 0.32 mmol) was dissolved in DMF (5mL) and then cooled to ~ 4 °C. The compound of Preparative Example 17 (0.06 g, 0.4 mmol) was then added, followed by DEC (0.08 g, 0.4 mmol), HOBT (0.6 g, 0.4 mmol), and 4-methylmorpholine (0.04 g, 50 µL, 0.4 mmol), and the reaction was then stirred at room
10 temperature overnight. The reaction mixture was concentrated in vacuo to a residue that was partitioned between CH₂Cl₂ and sat. NaHCO₃ (aqueous). The aqueous phase was extracted further with CH₂Cl₂, the combined CH₂Cl₂ fractions were dried over MgSO₄ and concentrated in vacuo to give a residue that was chromatographed on silica gel column using 5 % (NH₃
15 saturated CH₃OH)/ CH₂Cl₂ as eluent to give the title compound as a white solid: 0.11 g, 61% yield, m.p. = 118-119 °C, MH⁺ = 596.

EXAMPLE 2

(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[1-oxo-3-(2-oxo-1-pyrrolidiny]propyl]piperidine



20


The title compound of Preparative Example 12 (0.4 g, 0.74 mmol) was dissolved in CH₂Cl₂ (10 mL), and 4-bromo butyryl chloride (0.2 g, 0.13 mL, 1.11 mmol) and Et₃N (0.164g, 0.23 ml, 1.62 mmol)-were then added. The reaction mixture was stirred at room temperature for 16h. The
25 reaction mixture was partitioned between sat. NaHCO₃ and CH₂Cl₂. The

- 37 -

aqueous phase was extracted with CH_2Cl_2 , the combined CH_2Cl_2 extracts were dried over MgSO_4 and the solvent was removed by rotary evaporation. The resulting product was dissolved in THF (10 mL), cooled to $-10\text{ }^\circ\text{C}$, NaH (0.09 g, 3.79 mmol) was added and the reaction mixture was stirred for 16h, allowing the temperature to come to room temperature. The reaction mixture was then partitioned between sat. NaHCO_3 and EtOAc. The organic phase was dried over MgSO_4 and purified by flash chromatography on silica gel, eluting with 3% CH_3OH (sat. with NH_3)/ CH_2Cl_2 to give the title compound as a white solid (0.07 g), m.p. = $128\text{-}129\text{ }^\circ\text{C}$, $\text{MH}^+ = 610$.

EXAMPLE 3


(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1-[1-oxo-4-(2-oxo-1-pyrrolidiny)butyl]piperidine



The title compound is prepared from the product of Preparative Example 13, following essentially the same procedure as described for Example 2 to obtain a white solid, m.p. = $127\text{-}128\text{ }^\circ\text{C}$, $\text{MH} = 624$.

EXAMPLE 4

(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1-[(2-oxo-1-piperidiny)acetyl]piperidine




The title compound is prepared from the product of Preparative Example 11, following essentially the same procedure as described for Example 2 except that 4-bromovaleryl chloride was used instead of 4bromo butyryl chloride to obtain a white solid. Yield = 50%, m.p. = $138\text{-}139\text{ }^\circ\text{C}$, $\text{MH} = 610$.

- 38 -

EXAMPLE 5

(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[1-oxo-3-(2-oxo-1-piperidiny)propyl]piperidine




5 The title compound was prepared from the product of Preparative Example 12, following essentially the same procedure as described for Example 2 except that 4-bromovaleryl chloride was used instead of 4-bromo butyryl chloride to obtain a white solid.
Yield = 50%, m.p. = 138-139 °C, MH= 610.

10

EXAMPLE 6


(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[1-oxo-4-(2-oxo-1-piperidiny)butyl]piperidine



15 The title compound is prepared from the product of Preparative Example 13, following essentially the same procedure as described for Example 2 except that 4-bromovaleryl chloride was used instead of 4-bromo butyryl chloride to obtain a white solid.
Yield = 81%, m.p. = 101-102 °C, MH= 638.

EXAMPLE 7

20 (+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[(2-oxo-1-imidazolidiny)acetyl]piperidine




- 39 -

The product of Preparative Example 11 (2.08 g, 3.9 mmol) was dissolved in CH₂Cl₂ (20 mL) and 2-bromo ethyl isocyanate (0.8 g, 0.5 mL, 7.9 mmol) was added. The reaction mixture was stirred at room temperature for 16h. The reaction mixture was partitioned between sat. NaHCO₃ and CH₂Cl₂. The aqueous phase was extracted with CH₂Cl₂, the combined CH₂Cl₂ extracts were dried over MgSO₄ and the solvent was removed by rotary evaporation. The resulting product was dissolved in THF (20 mL), cooled to -10 °C, NaH (0.46 g, 19.5 mmol) was added and the reaction mixture was stirred for 16 h, allowing the temperature to come to room temperature. The reaction mixture was then partitioned between sat. NaHCO₃ and EtOAc. The organic phase was dried over MgSO₄ and purified by flash chromatography on silica gel, eluting with 5% CH₃OH(saturated with NH₃)/ CH₂Cl₂ to give the title compound as a white solid: 1.95 g, yield = 80%, m.p. = 167-168 °C, MH⁺ = 597.

EXAMPLE 8

(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[3-(2-oxoimidazolidinyl)-1-oxopropyl]piperidine




The title compound is prepared from the product of Preparative Example 12, following essentially the same procedure as described in Example 7 to obtain a white solid.

Yield = 45%, m.p. = 202-203 °C, MH⁺ = 611.

EXAMPLE 9

((+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[1-oxo-4-(2-oxo-1-imidazolidinyl)butyl]piperidine



- 40 -


The title compound is prepared from the product of Preparative Example 13, following essentially the same procedure as described in Example 7 to obtain a white solid.

Yield = 52%, m.p. = 120-123 °C, MH^+ = 625.

5

EXAMPLE 10

(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[(hexahydro-2-oxo-1-pyrimidinyl)acetyl]piperidine




The title compound is prepared from the product of Preparative Example 11, following essentially the same procedure as described for Example 7, substituting 3-chloro propyl isocyanate for 2-bromo ethyl isocyanate to obtain a white solid.

Yield = 56%, m.p. = 155-156 °C, MH^+ = 611.

EXAMPLE 11

(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta-[1,2-b]pyridin-11-yl)-1-[1-oxo-3-(hexahydro-2-oxo-1-pyrimidinyl)-oxopropyl] piperidine




The title compound is prepared from the product of Preparative Example 12, following essentially the same procedure as described for Example 10 to obtain a white solid.

Yield = 40%, m.p. = 135-136°C, MH^+ = 625.

- 41 -

EXAMPLE 12


(+)-4-(3,10-Dibromo-8-chloro-6,11-dihydro-5H-benzo[5,6]cyclohepta[1,2-b]pyridin-11-yl)-1-[4-(hexahydro-2-oxo-1-pyrimidinyl)oxobutyl]piperidine



5 The title compound is prepared from the product of Preparative Example 13, following essentially the same procedure as described for Example 10 to obtain a white solid.






Yield = 63%, m.p. = 157-158°C, $MH^+ = 639$.

10 Using the starting materials described above and the appropriate procedure, compounds of the following structure are prepared:



Ex.	Starting Material	Procedure	$(CH_2)_n R^6$	Analytical Data
13	Prep. Ex. 10 and 17	Ex. 1		Mass spec: $MH^+ = 721$; m.p. = 140-141°C
14	Prep. Ex. 15	Ex. 2		Mass spec: $MH^+ = 735$; m.p. = 129-130°C
15	Prep. Ex. 16	Ex. 2		Mass spec: $MH^+ = 749$; m.p. = 141-142°C
16	Prep. Ex. 14	Ex. 5		Mass spec: $MH^+ = 735$; m.p. = 147-148°C
17	Prep. Ex. 15	Ex. 5		Mass spec: $MH^+ = 749$; m.p. = 130-131°C
18	Prep. Ex. 16	Ex. 5		Mass spec: $MH^+ = 763$; m.p. = 137-138°C

- 42 -

19	Prep. Ex. 15	Ex. 7		Mass spec: $MH^+=736$; m.p. = 184-185°C
20	Prep. Ex. 16	Ex. 7		Mass spec: $MH^+=750$
21	Prep. Ex. 14	Ex. 10		Mass spec: $MH^+=736$; m.p. = 168-169°C
22	Prep. Ex. 15	Ex. 10		Mass spec: $MH^+=750$; m.p. = 146-147°C
23	Prep. Ex. 16	Ex. 10		Mass spec: $MH^+=764$; m.p. = 148-149°C

FPT IC_{50} (inhibition of farnesyl protein transferase, in vitro enzyme assay), COS Cell IC_{50} (Cell-Based Assay), GGPT IC_{50} (inhibition of geranylgeranyl protein transferase, in vitro enzyme assay), Cell Mat

5 Assay, and anti-tumor activity (in vivo anti-tumor studies) are determined by the assay procedures described in WO 95/10516. The test protocol for the assay used to determine inhibition of human tumor cell growth, the Soft Agar Assay, is as follows:

Anchorage-independent growth is a characteristic of tumorigenic

10 cell lines. Human tumor cells are suspended in growth medium containing 0.3% agarose and an indicated concentration of a farnesyl transferase inhibitor. The solution is overlayed onto growth medium solidified with 0.6% agarose containing the same concentration of farnesyl transferase inhibitor as the top layer. After the top layer is solidified, plates

15 are incubated for 10-16 days at 37°C under 5% CO_2 to allow colony outgrowth. After incubation, the colonies are stained by overlaying the agar with a solution of MTT (3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyl-tetrazolium bromide, Thiazolyl blue) (1 mg/mL in PBS). Ras mutation status was determined by ELISA (Oncogene Science). Colonies are

20 counted and the IC_{50} 's determined.

FPT IC_{50} values determined for compounds of this invention ranged from 0.0014 to 0.085 μM .

IC_{50} values for inhibition of ras processing in COS cells ranged from <0.010 to 0.16 μM for compounds of this invention.

- 43 -

Results of the soft agar assay using the tumor cell line activated H ras NIH 3T3 ranged from 0.145 to >0.500 for compounds of the invention.

For preparing pharmaceutical compositions from the compounds
5 described by this invention, inert, pharmaceutically acceptable carriers
can be either solid or liquid. Solid form preparations include powders,
tablets, dispersible granules, capsules, cachets and suppositories. The
powders and tablets may be comprised of from about 5 to about 70
percent active ingredient. Suitable solid carriers are known in the art, e.g.
10 magnesium carbonate, magnesium stearate, talc, sugar, lactose. Tablets,
powders, cachets and capsules can be used as solid dosage forms
suitable for oral administration.

For preparing suppositories, a low melting wax such as a mixture of
fatty acid glycerides or cocoa butter is first melted, and the active
15 ingredient is dispersed homogeneously therein as by stirring. The molten
homogeneous mixture is then poured into convenient sized molds,
allowed to cool and thereby solidify.

Liquid form preparations include solutions, suspensions and
emulsions. As an example may be mentioned water or water-propylene
20 glycol solutions for parenteral injection.

Liquid form preparations may also include solutions for intranasal
administration.

Aerosol preparations suitable for inhalation may include solutions
and solids in powder form, which may be in combination with a
25 pharmaceutically acceptable carrier, such as an inert compressed gas.

Also included are solid form preparations which are intended to be
converted, shortly before use, to liquid form preparations for either oral or
parenteral administration. Such liquid forms include solutions,
suspensions and emulsions.

30 The compounds of the invention may also be deliverable
transdermally. The transdermal compositions can take the form of creams,
lotions, aerosols and/or emulsions and can be included in a transdermal
patch of the matrix or reservoir type as are conventional in the art for this
purpose.

35 Preferably the compound is administered orally.

Preferably, the pharmaceutical preparation is in unit dosage form.
In such form, the preparation is subdivided into unit doses containing

- 44 -

appropriate quantities of the active component, e.g., an effective amount to achieve the desired purpose.

The quantity of active compound in a unit dose of preparation may be varied or adjusted from about 0.1 mg to 1000 mg, more preferably from about 1 mg. to 300 mg, according to the particular application.

The actual dosage employed may be varied depending upon the requirements of the patient and the severity of the condition being treated. Determination of the proper dosage for a particular situation is within the skill of the art. Generally, treatment is initiated with smaller dosages which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired.

The amount and frequency of administration of the compounds of the invention and the pharmaceutically acceptable salts thereof will be regulated according to the judgment of the attending clinician considering such factors as age, condition and size of the patient as well as severity of the symptoms being treated. A typical recommended dosage regimen is oral administration of from 10 mg to 2000 mg/day preferably 10 to 1000 mg/day, in two to four divided doses to block tumor growth. The compounds are non-toxic when administered within this dosage range.

The following are examples of pharmaceutical dosage forms which contain a compound of the invention. The scope of the invention in its pharmaceutical composition aspect is not to be limited by the examples provided.

Pharmaceutical Dosage Form Examples

EXAMPLE A Tablets

No.	Ingredients	mg/tablet	mg/tablet
1.	Active compound	100	500
2.	Lactose USP	122	113
3.	Corn Starch, Food Grade, as a 10% paste in Purified Water	30	40
4.	Corn Starch, Food Grade	45	40
5.	Magnesium Stearate	<u>3</u>	<u>7</u>
Total		300	700

- 45 -

Method of Manufacture

- Mix Item Nos. 1 and 2 in a suitable mixer for 10–15 minutes. Granulate the mixture with Item No. 3. Mill the damp granules through a coarse screen (e.g., 1/4", 0.63 cm) if necessary. Dry the damp granules.
- 5 Screen the dried granules if necessary and mix with Item No. 4 and mix for 10–15 minutes. Add Item No. 5 and mix for 1–3 minutes. Compress the mixture to appropriate size and weigh on a suitable tablet machine.

EXAMPLE BCapsules

No.	Ingredient	mg/capsule	mg/capsule
1.	Active compound	100	500
2.	Lactose USP	106	123
3.	Corn Starch, Food Grade	40	70
4.	Magnesium Stearate NF	<u>7</u>	<u>7</u>
Total		253	700


10 Method of Manufacture

Mix Item Nos. 1, 2 and 3 in a suitable blender for 10-15 minutes. Add Item No. 4 and mix for 1-3 minutes. Fill the mixture into suitable two-piece hard gelatin capsules on a suitable encapsulating machine.

- 15 While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.

CLAIMS:

1. A compound represented by the structural formula



or an N-oxide thereof, or a pharmaceutically acceptable salt or solvate thereof,
wherein:

R and R^2 are independently selected from halo;

R^1 and R^3 are independently selected from the group consisting of H and halo, provided that at least one of R^1 and R^3 is H;

W is N, CH or C when the double bond is present at the C-11 position;

R^4 is $-(CH_2)_n-R^5$ or

$$-CH_2-\left(\text{cycloheptanone ring}\right)_{n_1}-C(=O)-\left(CH_2\right)_n-R^6$$

R⁵ is ,  or  ;

R⁶ is R⁵ or ;


Z^1 and Z^2 are independently selected from the group consisting of =O and =S;

n is 1-6; and


n_1 is 0 or 1.

2. A compound of claim 1 wherein Z¹ is =O.
3. A compound of any of claims 1 or 2 wherein R⁴ is







- 47 -




4. A compound of any of claims 1, 2 or 3 wherein R is bromo and R² is chloro or bromo.
5. A compound of any of claims 1, 2 or 3 wherein R is bromo, R² is chloro or bromo, R¹ is H, and R³ is chloro or bromo.
6. A compound of any of claims 1, 2 or 3 wherein R is bromo, R² is chloro or bromo, R³ is H, and R¹ is chloro or bromo.
7. A compound of claim 1 selected from the group consisting of



- 48 -



- 49 -



8. An N-oxide, pharmaceutically acceptable salt or solvate of a compound of the formula defined in any one of claims 1 to 7.

9. A pharmaceutical composition for inhibiting the abnormal growth of cells comprising an effective amount of compound of claim 1, 2, 3, 4, 5, 6 or 7 in combination with a pharmaceutically acceptable carrier.

10. A pharmaceutical composition for inhibiting the abnormal growth of cells comprising an effective amount of an N-oxide, pharmaceutically acceptable salt or solvate of a compound of the formula defined in claim 1, 2, 3, 4, 5, 6 or 7 in combination with a pharmaceutically acceptable carrier.

11. The use of a compound of claim 1, 2, 3, 4, 5, 6 or 7 for the preparation of a medicament for the treatment of tumor cells expressing an activated ras oncogene.

12. The use of an N-oxide, pharmaceutically acceptable salt or solvate of a compound of the formula defined in claim 1, 2, 3, 4, 5, 6 or 7 for the preparation of a medicament for the treatment of tumor cells expressing an activated ras oncogene.

- 50 -

13. The use of claim 11 or 12 wherein the cells are pancreatic tumor cells, breast cancer cells, prostate cancer cells, lung cancer cells, myeloid leukemia tumor cells, thyroid follicular tumor cells, myelodysplastic tumor cells, epidermal carcinoma tumor cells, bladder carcinoma tumor cells or colon tumors cells.

14. The use of a compound of claim 1, 2, 3, 4, 5, 6 or 7 for the preparation of a medicament for inhibiting farnesyl protein transferase.

15. The use of an N-oxide, pharmaceutically acceptable salt or solvate of a compound of the formula defined in claim 1, 2, 3, 4, 5, 6 or 7 for the preparation of a medicament for inhibiting farnesyl protein transferase.

