



(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2020/04/24
 (87) Date publication PCT/PCT Publication Date: 2020/10/29
 (85) Entrée phase nationale/National Entry: 2021/10/19
 (86) N° demande PCT/PCT Application No.: US 2020/029775
 (87) N° publication PCT/PCT Publication No.: 2020/219848
 (30) Priorités/Priorities: 2019/04/26 (US62/839,455);
 2020/04/03 (US63/005,041)

(51) Cl.Int./Int.Cl. *C07K 14/725* (2006.01)
 (71) Demandeur/Applicant:
 ALLOGENE THERAPEUTICS, INC., US
 (72) Inventeurs/Inventors:
 PERTEL, THOMAS CHARLES, US;
 SASU, BARBRA JOHNSON, US;
 LEONARD, MARK W., US
 (74) Agent: GOWLING WLG (CANADA) LLP

(54) Titre : RECEPTEURS D'ANTIGENES CHIMERIQUES RESISTANT AU RITUXIMAB ET LEURS UTILISATIONS
 (54) Title: RITUXIMAB-RESISTANT CHIMERIC ANTIGEN RECEPTORS AND USES THEREOF

(57) **Abrégé/Abstract:**

Provided herein are polynucleotides encoding chimeric antigen receptors (CARs) comprising a CD19 antigen binding domain that specifically binds to CD19 and is resistant to rituximab binding; and immune cells comprising these CD19-specific CARs, e.g., CAR-T cells. Also provided are methods of making and using these CD19-specific CARs, and immune cells comprising these CD19-specific CARs.

Date de soumission / Date Submitted: 2021/10/19

No de la demande can. / CA App. No.: 3134308

Abrégé:

L'invention concerne des polynucléotides codant pour des récepteurs d'antigènes chimériques (CAR) comprenant un domaine de liaison à l'antigène CD19 qui se lie spécifiquement à CD19 et qui est résistant à la liaison au rituximab; et des cellules immunitaires comprenant ces CAR spécifiques à CD19, par exemple des cellules CAR-T. L'invention concerne également des procédés de fabrication et d'utilisation de ces CAR spécifiques à CD19, et des cellules immunitaires comprenant ces CAR spécifiques à CD19.

Abstract:

Provided herein are polynucleotides encoding chimeric antigen receptors (CARs) comprising a CD19 antigen binding domain that specifically binds to CD19 and is resistant to rituximab binding; and immune cells comprising these CD19-specific CARs, e.g., CAR-T cells. Also provided are methods of making and using these CD19-specific CARs, and immune cells comprising these CD19-specific CARs.

RITUXIMAB-RESISTANT CHIMERIC ANTIGEN RECEPTORS AND USES THEREOF**CROSS REFERENCE TO RELATED APPLICATIONS**

[001] The present application claims the benefit of priority to U.S. Provisional Application No. 62/839,455, filed on April 26, 2019; and U.S. Provisional Application No. 63/005,041, filed on
5 April 3, 2020, the contents of both of which are hereby incorporated by reference in their entireties.

TECHNICAL FIELD

[002] This disclosure relates to chimeric antigen receptors (CARs) comprising an antigen binding molecule which binds to CD19, polynucleotides encoding the same, and methods of treating a cancer in a patient using the same.

10 **SEQUENCE LISTING**

[003] This application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on April 21, 2020, is named AT-028_03WO_SL.txt and is 81,460 bytes in size.

15 **BACKGROUND**

[004] Adoptive transfer of immune cells genetically modified to recognize malignancy-associated antigens is showing promise as a new approach to treating cancer (see, e.g., Brenner et al., *Current Opinion in Immunology*, 22(2): 251-257 (2010); Rosenberg et al., *Nature Reviews Cancer*, 8(4): 299-308 (2008)). Immune cells can be genetically modified to express chimeric
20 antigen receptors (CARs), fusion proteins comprised of a CD19 antigen recognition moiety and T cell activation domains (see, e.g., Eshhar et al., *Proc. Natl. Acad. Sci. USA*, 90(2): 720-724 (1993), and Sadelain et al., *Curr. Opin. Immunol*, 21(2): 215-223 (2009)). Immune cells that contain CARs, e.g., CAR-T cells (CAR-Ts), are engineered to endow them with antigen specificity while retaining or enhancing their ability to recognize and kill a target cell.

25 [005] There is a need for treatments for cancer and in particular malignancies involving aberrant expression of CD19. Provided herein are methods and compositions addressing this need.

SUMMARY

[006] Provided herein are chimeric antigen receptors (CARs) comprising a CD19 antigen binding domain that specifically binds to CD19; polynucleotides encoding these CARs; and immune
30 cells expressing these CD19-specific CARs, e.g., CAR-T cells. Also provided are methods of

making and using these CD19-specific CARs, and immune cells comprising these CD19-specific CARs.

5 [007] In one aspect, the disclosure provides an isolated polynucleotide encoding a polypeptide comprising an anti-CD19 chimeric antigen receptor (CAR) that is at least 70% identical to SEQ ID NO: 9, wherein the polypeptide does not comprise a rituximab binding site, and wherein the polynucleotide comprises a short EF1a promoter that is capable of expressing the anti-CD19 chimeric antigen receptor (CAR) in a mammalian T cell.

10 [008] In some embodiments, the short EF1a promoter does not comprise an intron sequence within the nucleic acid sequence of SEQ ID NO:15. In some embodiments, the intron comprises the nucleic acid sequence of SEQ ID NO:39.

[009] In some embodiments, the promoter comprises the nucleic acid sequence of SEQ ID NO:16,

[010] In some embodiments, the promoter is a full length EF1a promoter comprising the nucleic acid sequence of SEQ ID NO:15.

15 [011] In some embodiments, the promoter comprises the nucleic acid sequence of SEQ ID NO:15, and the polynucleotide encodes a polypeptide that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to any one of SEQ ID NO: 8-14.

[012] In some embodiments, the polypeptide further comprises a safety switch.

20 [013] In some embodiments, the safety switch is linked to the CD19 CAR using a linker peptide.

[014] In some embodiments, the safety switch is linked to the anti-CD19 CAR using a T2A linker.

[015] In some embodiments, the safety switch comprises an antibody binding site.

[016] In some embodiments, the safety switch comprises a mutated CD20 mimotope.

25 [017] In some embodiments, the polypeptide further comprises a CD8 hinge/transmembrane domain.

[018] In some embodiments, the polypeptide comprises a CD34 epitope.

[019] In some embodiments, the CD34 epitope is a QBEND-10 epitope.

30 [020] In some embodiments, the isolated polynucleotide comprises a nucleic acid sequence that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to any one of SEQ ID NO: 1-7.

[021] In some embodiments, the isolated polynucleotide encodes a polypeptide that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to any one of SEQ ID NO: 8-14.

35 [022] In another aspect, the disclosure provides a vector comprising the isolated polynucleotide described herein.

[023] In some embodiments, the vector is a retroviral vector, a DNA vector, a plasmid, an RNA vector, an adenoviral vector, an adenovirus associated vector, a lentiviral vector, or any combination thereof.

[024] In one aspect, the disclosure provides an engineered immune cell comprising the isolated polynucleotide described herein.

[025] In some embodiments, the disclosure provides an engineered immune cell comprising a polynucleotide that comprises a nucleic acid sequence that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to SEQ ID NO: 3.

[026] In some embodiments, the disclosure provides an engineered immune cell comprising a polynucleotide that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to the anti-CD19 CAR v1.2, v1.3, v1.4, v1.5 or v1.6 lentiviral construct as shown in Table 1.

[027] In some embodiments, the disclosure provides an engineered immune cell comprising a polynucleotide that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to the anti-CD19 CAR v1.2 lentiviral construct as shown in Table 1.

[028] In some embodiments, the disclosure provides an engineered immune cell comprising a polynucleotide that encodes a polypeptide that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to SEQ ID NO:9 or SEQ ID NO:10, with or without the signal sequence.

[029] In some embodiments, the disclosure provides an engineered immune cell comprising a polynucleotide that comprises the nucleic acid sequence of SEQ ID NO:3, driven by an EF1a short promoter comprising the nucleic acid of SEQ ID NO:16.

[030] In some embodiments, the disclosure provides an engineered immune cell comprising a polynucleotide that encodes a polypeptide sequence of SEQ ID NO:9 or 10, with or without the signal sequence, driven by an EF1a short promoter comprising the nucleic acid of SEQ ID NO:16. In some embodiments, the promoter does not comprise the first intron of the EF1a gene.

[031] In some embodiments, the engineered immune cell does not comprise a rituximab mimotope.

[032] In some embodiments, the engineered immune cell comprises a polynucleotide comprising the CD19 CAR v1.2 lentiviral construct as shown in FIG. 1 (may also be referred to as ALLO-501A).

[033] In some embodiments, the engineered immune cell comprises a vector described herein.

[034] In some embodiments, the immune cell is a T cell, tumor infiltrating lymphocyte (TIL), NK cell, TCR-expressing cell, dendritic cell, or NK-T cell.

[035] In some embodiments, the cell is an autologous T cell.

[036] In some embodiments, the cell is an allogeneic T cell.

[037] In one aspect, the disclosure provides an engineered immune cell described herein, wherein the cell is resistant to rituximab.

[038] In another aspect, the disclosure provides a pharmaceutical composition comprising the engineered immune cell described herein.

[039] In one aspect, the disclosure provides a method of treating a disease or disorder in a subject in need thereof comprising administering to the subject the engineered immune cell
5 described herein, or the pharmaceutical composition described herein.

[040] In some embodiments, the disease or disorder is Non-Hodgkin lymphoma (NHL).

[041] In some embodiments, the subject has been treated or is currently being treated with rituximab.

[042] In one aspect, the disclosure provides an article of manufacture comprising the
10 engineered immune cell or the pharmaceutical composition comprising the engineered immune cell expressing a chimeric antigen receptors described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[043] FIG. 1 shows schematic representations of rituximab-resistant CD19 chimeric antigen
15 receptors.

[044] FIG. 2A and 2B show flow cytometry plots demonstrating CAR expression on day 5
from Pan T cells transduced with the rituximab resistant CAR expression vectors. FIG. 2A shows
CAR expression on cells from donor 541 and 604. FIG. 2B shows CAR and CD34 expression on
cells from donor 410. A similar profile was observed with donor 2593 (data not shown).

[045] FIG. 3 shows normalized cell expansion and final CAR expression from all four donors
20 on day 13.

[046] FIG. 4A-4D show cell expansion and CAR expression over time of Pan T cells from
donor 541 (FIG. 4A), 604 (FIG. 4B), 410 (FIG. 4C), 2593 (FIG. 4D) transduced with rituximab
resistant CAR expression vectors.

[047] FIGs. 5A-5D show CD4/CD8 ratios on day 5, 9, and 13 of Pan T cells from donor 541
25 (FIG. 5A), 604 (FIG. 5B), 410 (FIG. 5C), 2593 (FIG. 5D) transduced with rituximab resistant CAR
expression vectors.

[048] FIGs. 6A-6D show phenotype and activation on day 9 of Pan T cells from donor 541
(FIG. 6A), 604 (FIG. 6B), 410 (FIG. 6C), 2593 (FIG. 6D) transduced with rituximab resistant CAR
expression vectors.

[049] FIG. 7 shows phenotype, activation %CD8+, and anergy measured using TIM3 and PD1
30 staining averaged from all four donors on day 9.

[050] FIGs. 8A-8D show phenotype and activation on day 13 of Pan T cells from donor 541
(FIG. 8A), 604 (FIG. 8B), 410 (FIG. 8C), 2593 (FIG. 8D) transduced with rituximab resistant CAR
expression vectors.

[051] FIG. 9 shows phenotype, activation %CD8+, and anergy measured using TIM3 and PD1 staining from all four donors on day 13.

[052] FIG. 10 shows average short-term (24hr) killing assays using Raji cells as target cells for each CAR construct.

5 [053] FIGs. 11A-11D shows average long-term killing assays using A549-CD19+ cells as target cells with an E:T of 8:1 (FIG. 11A), 4:1 (FIG. 11B), 2:1 (FIG. 11C), and 1:1 (FIG. 11D) for each CAR construct.

[054] FIG. 12 shows the level of % CAR+ T cells on Day 5 after Pan T cells were transduced with serial dilution of lentiviral preparations of rituximab-resistant lentiviral constructs (ALLO-501v1.2 and v1.3) or rituximab-sensitive lentiviral construct (ALLO-501v1.0).

10 [055] FIGs. 13A-13B show *in vivo* efficacy of pan T cells from donors 541 (FIG. 13A) and 604 (FIG. 13B) transduced with rituximab-resistant CAR expression vector (ALLO-501v1.2) or rituximab-sensitive CAR expression vector (ALLO-501v1.0) when tested in a Raji cells-bearing NSG mouse tumor model.

15

DETAILED DESCRIPTION

[056] Chimeric Antigen Receptor (CAR) therapy is a promising approach to cancer treatment. The CAR construct described herein as v1.0 is an exemplary anti-CD19 CAR that expresses a synthetic peptide, RQR8, which serves as a safety switch. RQR8 contains two rituximab-binding mimotopes. In the instance of an adverse event, patients may be treated with rituximab to deplete the levels of anti-CD19 v1.0 in circulation. Rituximab is also used as the standard of care in some Non-Hodgkin Lymphoma (NHL) indications and is administered at high doses. Due to the long half-life of rituximab, anti-CD19 v1.0 cannot be administered to patients until the level of circulating rituximab has reached a low concentration. Rituximab-resistant CD19 CAR therapy would allow patients who were previously treated with rituximab to receive CAR-T therapy immediately, without having to wait for rituximab levels to diminish and without patients being subjected to apheresis. Provided herein are anti-CD19 chimeric antigen receptors (CARs) that are resistant to the CD20 binding antibody rituximab. The novel CAR constructs are designed to eliminate rituximab binding while retaining CAR expression and activity.

30

I. Chimeric Antigen Receptors

[057] As used herein, chimeric antigen receptors (CARs) are proteins that specifically recognize target antigens (e.g., target antigens on cancer cells). When bound to the target antigen, the CAR may activate the immune cell to attack and destroy the cell bearing that antigen (e.g., the cancer cell). CARs may also incorporate costimulatory or signaling domains to increase their

35

potency. See Krause *et al.*, *J. Exp. Med.*, Volume 188, No. 4, 1998 (619–626); Finney *et al.*, *Journal of Immunology*, 1998, 161: 2791–2797, Song *et al.*, *Blood* 119:696-706 (2012); Kalos *et al.*, *Sci. Transl. Med.* 3:95 (2011); Porter *et al.*, *N. Engl. J. Med.* 365:725-33 (2011), and Gross *et al.*, *Annu. Rev. Pharmacol. Toxicol.* 56:59–83 (2016); U.S. Patent Nos. 7,741,465, and 6,319,494.

5 [058] Chimeric antigen receptors described herein comprise an extracellular domain, a transmembrane domain, and an intracellular domain, wherein the extracellular domain comprises a CD19 antigen binding domain that specifically binds to CD19. In some embodiments, the CD19 specific CAR comprises the following elements from 5' to 3': a signal sequence, a CD19 antigen binding domain (e.g., a scFv derived from 4G7), a hinge and transmembrane region, and one or
10 more successive signaling domains. In some embodiments, the antibody binding domain binds CD19 to treat a hematologic cancer associated with expression of CD19.

[059] The scFv portion of the chimeric antigen receptor (CAR) used in the allogeneic anti-CD19 CAR v1.0 is derived from the mouse anti-human CD19 antibody clone 4G7. 4G7 is a CD19 monoclonal antibody that recognizes CD19. Single chain variable fragments (scFv) formed from
15 4G7 comprise the targeting component of some chimeric antigen receptors (CARs) (See WO2014184143A1). In some embodiments, the scFv derived from the CD19 monoclonal antibody 4G7, comprises a part of the CD19 monoclonal antibody 4G7 immunoglobulin gamma 1 heavy chain (GenBank: CAD88275.1; SEQ ID NO: 17) and a part of CD19 monoclonal antibody 4G7 immunoglobulin kappa light chain (GenBank: CAD88204.1; SEQ ID NO: 35), linked together by a
20 flexible linker. (Peipp M., D. Saul, et al., 2004. Efficient eukaryotic expression of fluorescent scFv fusion proteins directed against CD antigens for FACS applications. *J. Immunol. Methods* 285: 265–280). In some embodiments, the scFv comprises the variable fragments of the CD19 monoclonal antibody 4G7 immunoglobulin gamma 1 heavy chain and the variable fragments of the CD19 monoclonal antibody 4G7 immunoglobulin kappa light chain linked together by a flexible linker.

25 [060] CD19 monoclonal antibody 4G7 immunoglobulin gamma 1 heavy chain (the signal sequence is underlined)

[061] MEWSWIFLFLLSGTAGVHSEVQLQQSGPELIKPGASVKMSCKASGYTFTSYVM
HWVKQKPGQGLEWIGYINPYNDGTYNEKFKGKATLTSDKSSSTAYMELSSLTSEDSAVY
YCARGTYYYGSRVFDYWGQGTTTLTVSSAKTTPPSVYPLAPGSAAQTNSMVTLGCLVKGYF
30 PEPVTVTWNSGSLSSGVHTFPAVLQSDLYTLSSSVTVPSSTWPSETVTCNVAHPASSTKVDK
KIVPRDCGCKPCICTVPEVSSVFIFPPKPKDVLITLTPKVTCTVVVDISKDDPEVQFSWFVDD
VEVHTAQTQPREEQFNSTFRSVSELPIMHQDWLNGKEFKCRVNSAAFPAPIEKTISKTKGRP
KAPQVYTIPPPKEQMAKDKVSLTCMITDFFPEDITVEWQWNGQPAENYKNTQPIMDTDGSY
FVYSKLVNPKSNWEAGNTFTCSVLHEGLHNHHTEKSLSHSPGK (SEQ ID NO: 17)

35 [062] CD19 monoclonal antibody 4G7 immunoglobulin kappa light chain (the signal sequence is underlined)

[063] MRCLAEFLGLLVLPWGAIGDIVMTQAAPSIPVTPGESVSISCRSSKSLNSNGNT
 YLYWFLQRPGQSPQLLIYRMSNLASGVDPDRFSGSGSGTAFTLRISRVEAEDVGVYYCMQHL
 EYPFTFGAGTKLELKRADAAPTVSIFPPSSEQLTSGGASVVCFLNMFYPKDINVKWKIDGSE
 RQNGVLNSWTDQDSKDYSTYSMSSTLTTLTKDEYERHNSYTCETHKSTSTSPIVKSFNREK
 5 (SEQ ID NO: 18)

[064] In some embodiments, the scFv comprises a part of amino acid sequences of SEQ ID NO: 17 and/or SEQ ID NO: 18. In some embodiments, the scFv comprises at least 70%, at least 80%, at least 90%, at least 95%, at least 97% or at least 99% sequence identity with the variable region of amino acid sequence of 34 and/or SEQ ID NO: 35. Disclosed herein are antigen binding
 10 molecules, including antibodies, that specifically bind to the anti-CD19 scFv derived from 4G7, as well as molecules comprising these sequences and cells presenting such molecules. Humanized forms of the antigen binding molecules also form as aspect of the disclosure. Applications and uses of these antigen binding molecules are also disclosed.

15 **a. Antigen Binding Domain**

[065] As discussed above, the CD19 CARs described herein comprise an antigen binding domain. An “antigen binding domain” as used herein means any polypeptide that binds a specified target antigen, for example the specified target antigen can be the CD19 protein or fragment thereof. In some embodiments, the antigen binding domain binds to a CD19 antigen on a tumor cell. In
 20 some embodiments, the antigen binding domain binds to a CD19 antigen on a cell involved in a hyperproliferative disease.

[066] In some embodiments, the antigen binding domain comprises a variable heavy chain, variable light chain, and/or one or more CDRs. In some embodiments, the antigen binding domain is a single chain variable fragment (scFv), comprising light chain CDRs CDR1, CDR2 and CDR3, and
 25 heavy chain CDRs CDR1, CDR2 and CDR3. Variants of the antigen binding domains (e.g., variants of the CDRs, VH and/or VL) are also within the scope of the disclosure, e.g., variable light and/or variable heavy chains that each have at least 70-80%, 80-85%, 85-90%, 90-95%, 95-97%, 97-99%, or above 99% identity to the amino acid sequences of the antigen binding domain sequences described herein. In some instances, such molecules include at least one heavy chain and one light
 30 chain, whereas in other instances the variant forms contain two variable light chains and two variable heavy chains (or subparts thereof). A skilled artisan will be able to determine suitable variants of the antigen binding domains as set forth herein using well-known techniques. In certain embodiments, one skilled in the art can identify suitable areas of the molecule that may be changed without destroying activity by targeting regions not believed to be important for activity.

35 [067] In certain embodiments, the polypeptide structure of the antigen binding domains is based on antibodies, including, but not limited to, monoclonal antibodies, bispecific antibodies,

minibodies, domain antibodies, synthetic antibodies (sometimes referred to herein as “antibody mimetics”), chimeric antibodies, humanized antibodies, human antibodies, antibody fusions (sometimes referred to herein as “antibody conjugates”), and fragments thereof, respectively. In some embodiments, the antigen binding domain comprises or consists of avimers.

5 [068] A CD19 antigen binding domain is said to be “selective” when it binds to one target more tightly than it binds to a second target. In some embodiments, the CD19 antigen binding domain is a scFv.

[069] In some embodiments, the disclosure relates to isolated polynucleotides encoding any one of the CD19 chimeric antigen receptors (CARs) described herein. In some embodiments, the
 10 disclosure relates to isolated polynucleotides encoding a CD19 CAR described in Table 1. Also provided herein are vectors comprising the polynucleotides, and methods of making the same.

Table 1. Polynucleotide Sequences of exemplary CD19 targeting CARs

SEQ ID NO	Description	Sequence
1	anti-CD19 CAR_v1.0	ATGCTGACCAGCCTGCTGTGCTGGATGGCCCTGTGCCTGCTG GGCGCCGACCACGCCGATGCCTGCCCTACAGCAACCCAG CCTGTGCAGCGGAGGCGGCGGAGCGAGCTGCCACCCAGG GCACCTTCTCCAACGTGTCCACCAACGTGAGCCCAGCCAAG CCCACCACCACCGCCTGTCCTTATTCCAATCCTTCCTGTGT AGCGGAGGGGGAGGCAGCCCAGCCCCAGACCTCCCACCCC AGCCCCACCATCGCCAGCCAGCCTCTGAGCCTGAGACCCG AGGCCTGCCGCCAGCCGCCGCGGCGCCGTGCACACCAGA GGCTGGATTTCGCCTGCGATATCTACATCTGGGCCCCACTG GCCGGCACCTGTGGCGTGCTGCTGCTGAGCCTGGTGATCACC CTGTACTGCAACCACCGCAACCGCAGGCGCGTGTGCAAGTG CCCCAGGCCCGTGGTGAGAGCCGAGGGCAGAGGCAGCCTGC TGACCTGCGGCGACGTGGAGGAGAACCAGGCCCCATGGAG ACCGACACCCTGCTGCTGTGGGTGCTGCTGCTGTGGGTGCCA GGCAGCACCGGCGAGGTGCAGCTGCAGCAGAGCGGACCCG AGCTGATCAAGCCAGGCGCCAGCGTGAAGATGAGCTGCAAG GCCAGCGGCTACACCTTACCAGCTACGTGATGCACTGGGT GAAGCAGAAGCCAGGCCAGGGCCTGGAGTGGATCGGCTAC ATCAACCCTACAACGACGGCACCAAGTACAACGAGAAGTT CAAGGGCAAGGCCACCCTGACCAGCGACAAGAGCAGCAGC ACCGCCTACATGGAGCTGAGCAGCCTGACCAGCGAGGACAG CGCCGTGTACTACTGCGCCAGAGGCACCTACTACTACGGCA GCCGGGTGTTGACTACTGGGGCCAGGGCACCCACCCTGACC

		<p>GTGAGCTCTGGCGGAGGCGGCTCTGGCGGAGGCGGCTCTGG CGGAGGCGGCAGCGACATCGTGATGACCCAGGCTGCCCCCA GCATCCCCGTGACCCCAGGCGAGAGCGTGAGCATCAGCTGC CGGAGCAGCAAGAGCCTGCTGAACAGCAACGGCAACACCTA CCTGTACTGGTTCCTGCAGCGGCCAGGCCAGAGCCCCCAGC TGCTGATCTACCGGATGAGCAACCTGGCCAGCGGCGTGCCC GACCGGTTACAGCGGCAGCGGCAGCGGCACCGCCTTCACCCT GCGGATCAGCCGGGTGGAGGCCGAGGACGTGGGCGTGTACT ACTGCATGCAGCACCTGGAGTACCCCTTCACCTTCGGAGCCC GCACCAAGCTGGAGCTGAAGCGGTCGGATCCCACCACCACC CCAGCCCCACGGCCACCTACCCCTGCCCAACCATCGCCAG CCAGCCCCTGAGCCTGCGGCCTGAAGCCTGCAGGCCTGCCG CCGGAGGAGCCGTGCACACAAGGGCCTGGACTTCGCCTGC GACATCTATATCTGGGCCCCCTGGCCGGGACATGCGGGGT GCTGCTGCTGTCCCTGGTGATTACACTGTATTGCAAACGGGG CCGGAAGAAGCTGCTGTACATCTTCAAGCAGCCCTTCATGC GGCCCGTGCAGACCACCCAGGAGGAGGACGGCTGCAGCTGC CGGTTCCCCGAGGAAGAGGAAGGCGGCTGCGAGCTGCGGGT GAAGTTCAGCCGGAGCGCCGACGCCCCAGCCTACCAGCAGG GCCAGAACCAGCTGTACAACGAGCTGAACCTGGGACGGCGG GAGGAGTACGACGTGCTGGACAAGCGGCGGGGACGGGACC CCGAGATGGGCGGCAAGCCTCGCCGGAAGAATCCCCAGGAG GGCCTGTACAACGAGCTGCAGAAGGACAAGATGGCCGAGG CCTACAGCGAGATCGGCATGAAGGGCGAGCGGCGCCGGGG CAAGGGCCACGACGGCCTGTACCAGGGCCTGAGCACCGCCA CCAAGGACACCTACGACGCCCTGCACATGCAGGCCCTGCCA CCCCGGTGA</p>
<p>2</p>	<p>anti-CD19 CAR _v1.1</p>	<p>ATGGAGACCGACACCCTGCTGCTGTGGGTGCTGCTGCTGTG GGTGCCAGGCAGCACCGGCGAGGTGCAGCTGCAGCAGAGC GGACCCGAGCTGATCAAGCCAGGCGCCAGCGTGAAGATGAG CTGCAAGGCCAGCGGCTACACCTTCACCAGCTACGTGATGC ACTGGGTGAAGCAGAAGCCAGGCCAGGGCCTGGAGTGGATC GGCTACATCAACCCCTACAACGACGGCACCAAGTACAACGA GAAGTTCAAGGGCAAGGCCACCCTGACCAGCGACAAGAGC AGCAGCACCGCCTACATGGAGCTGAGCAGCCTGACCAGCGA GGACAGCGCCGTGTACTACTGCGCCAGAGGCACCTACTACT ACGGCAGCCGGGTGTTCTGACTACTGGGGCCAGGGCACCA CTGACCGTGAGCTCTGGCGGAGGCGGCTCTGGCGGAGGCGG CTCTGGCGGAGGCGGCAGCGACATCGTGATGACCCAGGCTG</p>

		<p>CCCCAGCATCCCCGTGACCCCAGGCCGAGAGCGTGAGCATC AGCTGCCGGAGCAGCAAGAGCCTGCTGAACAGCAACGGCA ACACCTACCTGTACTGGTTCCTGCAGCGGCCAGGCCAGAGC CCCCAGCTGCTGATCTACCGGATGAGCAACCTGGCCAGCGG CGTGCCCCGACCGGTTTCAGCGGCAGCGGCAGCGGCACCGCCT TCACCTGCGGATCAGCCGGGTGGAGGCCGAGGACGTGGGC GTGTACTACTGCATGCAGCACCTGGAGTACCCCTTCACCTTC GGAGCCGGCACCAAGCTGGAGCTGAAGCGGTTCGGATCCCAC CACCACCCAGCCCCACGGCCACCTACCCCTGCCCAACCAT CGCCAGCCAGCCCCTGAGCCTGCGGCCTGAAGCCTGCAGGC CTGCCGCCGGAGGAGCCGTGCACACAAGGGGCCTGGACTTC GCCTGCGACATCTATATCTGGGCCCCCCTGGCCGGGACATGC GGGGTGCTGCTGCTGTCCCTGGTGATTACACTGTATTGCAA CGGGGCCGGAAGAAGCTGCTGTACATCTTCAAGCAGCCCTT CATGCGGCCCGTGCAGACCACCCAGGAGGAGGACGGCTGCA GCTGCCGGTTCGCCGAGGAAGAGGAAGGCGGCTGCGAGCTG CGGGTGAAGTTCAGCCGGAGCGCCGACGCCCCAGCCTACCA GCAGGGCCAGAACCAGCTGTACAACGAGCTGAACCTGGGAC GGCGGGAGGAGTACGACGTGCTGGACAAGCGGCCGGGACG GGACCCCGAGATGGGCGGCAAGCCTCGCCGGAAGAATCCCC AGGAGGGCCTGTACAACGAGCTGCAGAAGGACAAGATGGC CGAGGCCTACAGCGAGATCGGCATGAAGGGCGAGCGGGCGC CGGGGCAAGGGCCACGACGGCCTGTACCAGGGCCTGAGCAC CGCCACCAAGGACACCTACGACGCCCTGCACATGCAGGCC TGCCACCCCGGTGA</p>
<p>3</p>	<p>anti-CD19 CAR _v1.2</p>	<p>ATGGAGACAGATACCCCTGCTGCTGTGGGTGCTGCTGCTGTG GGTGCCTGGCTCCACAGGAGAGGTGCAGCTGCAGCAGTCTG GACCAGAGCTGATCAAGCCTGGAGCATCCGTGAAGATGTCT TGCAAGGCCAGCGGCTATACATTCACCAGCTACGTGATGCA CTGGGTGAAGCAGAAGCCTGGCCAGGGCCTGGAGTGGATCG GCTATATCAATCCATAACAACGACGGCACCAAGTATAATGAG AAGTTTAAGGGCAAGGCCACACTGACCTCTGATAAGAGCTC CTCTACAGCCTACATGGAGCTGAGCTCCCTGACCTCTGAGGA CAGCGCCGTGTACTATTGCGCCAGAGGCACATACTATTACG GCAGCAGGGTGTTTCGATTACTGGGGCCAGGGCACCACACTG ACCGTGTCTAGCGGAGGAGGAGGCTCCGGAGGAGGAGGCTC TGGCGGCCGGCGGCAGCGACATCGTGATGACACAGGCAGCAC CAAGCATCCCAGTGACCCCTGGCGAGAGCGTGTCCATCTCTT GTCGGTCTCTAAGTCCCTGCTGAACTCTAATGGCAACACCT</p>

		<p>ATCTGTA CTGGTTTCTGCAGCGGCCCGGACAGTCCCCACAGC TGCTGATCTATAGGATGAGCAACCTGGCATCCGGAGTGCCT GATCGCTTCAGCGGCTCCGGCTCTGGAACAGCCTTTACCCTG AGGATCTCTCGGGTGGAGGCAGAGGACGTGGGCGTGTATTA CTGCATGCAGCACCTGGAGTACCCCTTCACATTTGGCGCAGG AACCAAGCTGGAGCTGAAGCGGAGCGACCCACCACAACCC CTGCACCACGGCCCCCTACACCAGCACCTACCATCGCATCTC AGCCACTGAGCCTGCGGCCCGAGGCCTGTAGGCCTGCAGCA GGAGGAGCAGTGCACACCAGGGGCCTGGACTTCGCCTGCGA TATCTATATCTGGGCACCACTGGCAGGAACATGTGGCGTGTCT GCTGCTGAGCCTGGTCATCACCCCTGTATTGCAAGAGAGGCA GGAAGAAGCTGCTGTACATCTTCAAGCAGCCTTTTATGCGGC CAGTGCAGACAACCCAGGAGGAGGATGGCTGCTCCTGTAGA TTCCAGAGGAGGAGGAGGGAGGATGTGAGCTGCGCGTGA AGTTTAGCCGGTCCGCCGACGCACCAGCATATCAGCAGGGC CAGAATCAGCTGTACAATGAGCTGAACCTGGGCCGGAGAGA GGAGTACGACGTGCTGGATAAGAGGAGGGGAAGGGACCCC GAGATGGGAGGCAAGCCACGGAGAAAGAATCCCCAGGAGG GCCTGTATAACGAGCTGCAGAAGGATAAGATGGCCGAGGCC TACAGCGAGATCGGCATGAAGGGAGAGAGGCCGCCGGGGCA AGGGACACGACGGCCTGTATCAGGGCCTGTCCACAGCCACC AAGGACACCTACGATGCCCTGCACATGCAGGCCCTGCCACC AAGGTGA</p>
<p>4</p>	<p>anti-CD19 CAR _v1.3</p>	<p>ATGGGAACAAGCCTGCTGTGCTGGATGGCTCTGTGCCTGCTG GGGGCCGACCACGCTGACGCCTCCGGGGGGGGGGCTCTCC TGCCCTAGGCCCCCTACACCTGCACCAACCATCGCATCCCA GCCACTGTCTCTGCGCCCTGAGGCCTGCCGGCCAGCAGCAG GAGGAGCAGTGCACACCCGCGGCCTGGACTTCGCCTGCGAT ATCTATATCTGGGCACCACTGGCAGGCACATGTGGCGTGTCT GCTGCTGAGCCTGGTCATCACCCCTGTA CTGCAATCACAGGA ACCGGAGAAGGGTGTGCAAGTGTCCCCGGCCTGTGGTGAGA GCAGAGGGAAGGGGCAGCCTGCTGACATGTGGCGACGTGG AGGAGAATCCAGGCCCTATGGAGACAGATACCCTGCTGCTG TGGGTGCTGCTGCTGTGGGTGCCCGGCAGCACCGGAGAGGT GCAGCTGCAGCAGTCCGGACCAGAGCTGATCAAGCCTGGAG CCAGCGTGAAGATGTCTGTAAAGGCCTCTGGCTATACATTCA CCAGCTACGTGATGCACTGGGTGAAGCAGAAGCCTGGCCAG GGCCTGGAGTGGATCGGCTATATCAATCCATAACAACGACGG</p>

		<p>CACAAAGTATAACGAGAAGTTTAAGGGCAAGGCCACACTGA CCTCCGATAAGAGCTCCTCTACAGCCTACATGGAGCTGAGCT CCCTGACCTCTGAGGACAGCGCCGTGTACTATTGCGCCAGA GGCACATACTATTACGGCTCTAGGGTGTTCGATTACTGGGGC CAGGGCACCACACTGACCGTGTCTAGCGGAGGAGGAGGCAG CGGAGGAGGAGGCTCCGGCGGCGGGCTCTGACATCGTGA TGACACAGGCAGCACCATCCATCCCAGTGACCCCTGGCGAG AGCGTGTCCATCTCTTGTTCGGTCCTCTAAGAGCCTGCTGAAC TCCAATGGCAACACCTATCTGTACTGGTTTCTGCAGCGGCC GGACAGAGCCCACAGCTGCTGATCTATAGGATGTCTAATCT GGCAAGCGGCGTGCCCGATCGCTTCAGCGGCTCCGGCTCTG GCACAGCCTTTACCCTGAGGATCTCCCGCGTGGAGGCAGAG GACGTGGGCGTGTATTACTGCATGCAGCACCTGGAGTACCC CTTTACATTTGGCGCAGGCACCAAGCTGGAGCTGAAGCGGA GCGACCCACCACAACCCCTGCACCACGGCCACCCACACCA GCACCTACTATTGCATCCCAGCCACTGAGCCTGCGGCCGA GGCTGTAGGCCTGCCGCCGGCGGCAGTGCACACCCGGG GCCTGGACTTTGCCTGCGATATCTACATCTGGGCACCTCTGG CCGGCACATGCGGCGTGTGTTACTGAGCCTGGTTCATCACCC TGTATTGCAAGCGGGGCAGAAAGAAGCTGCTGTACATCTTC AAGCAGCCTTTTATGCGGCCAGTGCAGACAACCCAGGAGGA GGATGGCTGCTCCTGTAGATTCCCAGAGGAGGAGGAGGGAG GATGTGAGCTGCGCGTGAAGTTTAGCCGGTCCGCCGACGCA CCAGCATATCAGCAGGGCCAGAACCAGCTGTACAATGAGCT GAACCTGGGCCGGAGAGAGGAGTATGACGTGCTGGATAAG AGACGGGGCCGGGACCCCGAGATGGGAGGCAAGCCACGCC GGAAGAATCCCCAGGAGGGCCTGTATAACGAGCTGCAGAAG GATAAGATGGCCGAGGCCTACAGCGAGATCGGCATGAAGG GAGAGAGAAGGCGCGCAAGGGACACGACGGCCTGTACCA GGGCCTGAGCACAGCAACAAAAGACACCTACGACGCACTGC ACATGCAGGCTCTGCCCCCTCGGTAA</p>
<p>5</p>	<p>anti-CD19 CAR _v1.4</p>	<p>ATGGGAACCTCTCTGCTGTGCTGGATGGCTCTGTGCCTGCTG GGGGCCGATCACGCTGACGCAAGTGGCGGGGGGGGGTCCG AACTGCCACACAGGGCACCTTCTCCAACGTGAGCACCAAC GTGAGCTCCGGCGGAGGAGGCAGCCCTGCACCAAGGCCCC TACACCAGCACCTACCATCGCATCTCAGCCACTGAGCCTGCG CCCCAGGCCTGCCGGCCTGCAGCAGGCGGCGCCGTGCACA CCCGCGGCTGGACTTTGCCTGCGATATCTATATCTGGGCAC CTCTGGCAGGCACATGTGGCGTGTGCTGCTGAGCCTGGTCA</p>

		<p>TCACCCTGTA CTGCAATCACAGGAACCGGAGAAGGGTGTGC AAGTGTCCACGGCCCGTGGTGAGAGCAGAGGGAAGGGGCTC CCTGCTGACATGTGGCGACGTGGAGGAGAATCCTGGCCCAA TGGAGACAGATAACCCTGCTGCTGTGGGTGCTGCTGCTGTGG GTGCCCCGGCTCCACCGGAGAGGTGCAGCTGCAGCAGTCTGG ACCAGAGCTGATCAAGCCAGGAGCATCCGTGAAGATGTCTT GTAAGGCCAGCGGCTATACATTCACCAGCTACGTGATGCAC TGGGTGAAGCAGAAGCCAGGACAGGGCCTGGAGTGGATCG GCTATATCAATCCTTACAACGACGGCACCAAGTATAACGAG AAGTTTAAGGGCAAGGCCACACTGACCTCTGATAAGTCTAG CTCCACAGCCTACATGGAGCTGTCTAGCCTGACCAGCGAGG ACTCCGCCGTGTA CTATTGCGCCAGAGGCACATACTATTACG GCAGCAGGGTGTTCGATTACTGGGGCCAGGGCACCACTG ACCGTGTCTCTGGAGGAGGAGGCTCCGGAGGAGGAGGCTC TGGCGGCGGGCGGCAGCGACATCGTGATGACACAGGCAGCAC CTTCCATCCCAGTGACCCCAGGCGAGTCTGTGAGCATCTCCT GTCGGAGCTCCAAGTCCCTGCTGAACTCTAATGGCAACACCT ATCTGTA CTGGTTTCTGCAGCGGCCCGGACAGTCCCCACAGC TGCTGATCTATAGGATGAGCAATCTGGCCTCCGGCGTGCCA GATCGTTCTCTGGCAGCGGCTCCGGCACAGCCTTTACCCTG AGGATCTCTCGCGTGGAGGCAGAGGACGTGGGCGTGTATTA CTGCATGCAGCACCTGGAGTACCCATTACATTTGGCGCAG GCACCAAGCTGGAGCTGAAGCGGAGCGACCCCACCACAACC CCAGCACCTCGGCCACCCACACCAGCACCCACCATCGCATC TCAGCCTCTGAGCCTGCGGCCCGAGGCCTGTAGGCCCGCAG CAGGAGGAGCAGTGCACACCCGGGGCCTGGACTTCGCCTGC GATATCTACATCTGGGCACCACTGGCCGGCACATGCGGCGT GCTGTTACTGAGCCTGGTCATCACCCCTGTATTGCAAGCGGGG CAGAAAGAAGCTGCTGTACATCTTCAAGCAGCCCTTTATGC GGCCTGTGCAGACAACCCAGGAGGAGGATGGCTGCTCCTGT AGATTCCCTGAGGAGGAGGAGGGAGGATGTGAGCTGCGCGT GAAGTTTTCTCGGAGCGCCGACGCACCAGCATATCAGCAGG GACAGAACCAGCTGTACAATGAGCTGAACCTGGGCCGGAGA GAGGAGTATGACGTGCTGGATAAGAGACGGGGCCGGGACC CCGAGATGGGAGGCAAGCCTCGCCGGAAGAATCCACAGGA GGGCCTGTATAACGAGCTGCAGAAGGATAAGATGGCCGAGG CCTACAGCGAGATCGGCATGAAGGGAGAGAGAAGGCGCGG CAAGGGACACGACGGCCTGTACCAGGGCCTGAGCACAGCAA</p>
--	--	---

		<p>CAAAAGACACCTACGACGCACTGCACATGCAGGCTCTGCCA CCAAGATGA</p>
<p>6</p>	<p>anti-CD19 CAR _v1.5</p>	<p>ATGGGGACCTCACTGCTGTGCTGGATGGCTCTGTGCCTGCTG GGGGCCGACCACGCTGACGCCTGCTCTGGGGGGGGGGGGGG CTCATGCTCCGGAGGAGGAGGCTCTGAGCTGCCAACCAGG GCACATTCTCCAACGTGAGCACCAACGTGTCTCCTGCCAAGC CAACCACAACCGCATGCAGCGGCGGAGGAGGAGGCAGCTG TTCCGGCGGGCGGCGGCAGCCCTGCCCAAGGCCCCCTACCC CAGCACCTACAATCGCATCTCAGCCTCTGAGCCTGCGCCCAG AGGCCTGTGCGCCCGCAGCAGGAGGAGCAGTGCACACCCGC GGCCTGGACTTTGCCTGCGATATCTATATCTGGGCACCACTG GCAGGCACCTGTGGCGTGCTGCTGCTGAGCCTGGTCATCACC CTGTACTGCAATCACAGGAACCGGAGAAGGGTGTGCAAGTG TCCACGGCCCGTGGTGAGAGCAGAGGGAAGGGGCTCTCTGC TGACCTGTGGCGACGTGGAGGAGAATCCTGGCCCTATGGAG ACAGATACACTGCTGCTGTGGGTGCTGCTGCTGTGGGTGCC GGCAGCACAGGAGAGGTGCAGCTGCAGCAGTCCGGACCTGA GCTGATCAAGCCAGGCGCCTCCGTGAAGATGTCTTGCAAGG CCAGCGGCTATACCTTACAAGCTACGTGATGCACTGGGTG AAGCAGAAGCCAGGCCAGGGCCTGGAGTGGATCGGCTATAT CAATCCCTACAACGACGGCACCAAGTATAACGAGAAGTTTA AGGGCAAGGCCACCCTGACAAGCGATAAGAGCTCCTCTACC GCCTACATGGAGCTGAGCTCCCTGACAAGCGAGGACTCCGC CGTGTACTATTGCGCCAGAGGCACCTACTATTACGGCTCCAG GGTGTTCGATTACTGGGGCCAGGGCACCAACCCTGACAGTGT CTAGCGGAGGAGGAGGCAGCGGAGGAGGAGGCTCCGGCGG CGGCGGCTCTGACATCGTGATGACCCAGGCAGCACCATCCA TCCCTGTGACACCAGGCGAGTCTGTGAGCATCTCCTGTCCGT CCTCTAAGTCCCTGCTGAACTCTAATGGCAACACCTATCTGT ACTGGTTTCTGCAGCGGCCCGGACAGTCTCCTCAGCTGCTGA TCTATAGGATGAGCAATCTGGCCTCCGGCGTGCCTGATCGCT TCTCTGGCAGCGGCTCCGGCACCCGCTTTACACTGAGGATCA GCCGCGTGGAGGCAGAGGACGTGGGCGTGTATTACTGCATG CAGCACCTGGAGTACCCTTTCACCTTTGGCGCCGGCACAAA GCTGGAGCTGAAGCGGAGCGACCCCAACAACCACACCAGCAC CTCGGCCACCCACCCAGCACCAACAATCGCATCTCAGCCA CTGAGCCTGCGGCCCGAGGCCTGTAGGCCAGCCGCGGGCGG CGCAGTGCACACCCGGGGCCTGGACTTCGCCTGCGATATCT ACATCTGGGCCCTCTGGCCGGCACCTGCGGCGTGCTGTTAC</p>

		<p>TGAGCCTGGTCATCACCTGTATTGCAAGCGGGGCAGAAAG AAGCTGCTGTACATCTTCAAGCAGCCCTTCATGCGGCCCGTG CAGACCACACAGGAGGAGGATGGCTGCTCCTGTAGATTCCC AGAGGAGGAGGAGGGAGGATGTGAGCTGCGCGTGAAGTTTT CTCGGAGCGCCGACGCACCTGCATATCAGCAGGGACAGAAC CAGCTGTACAATGAGCTGAACCTGGGCCGGAGAGAGGAGTA TGACGTGCTGGATAAGAGACGGGGCCGGGACCCCGAGATGG GAGGCAAGCCCCGCCGGAAGAATCCTCAGGAGGGCCTGTAT AACGAGCTGCAGAAGGATAAGATGGCCGAGGCCTACAGCG AGATCGGCATGAAGGGAGAGAGAAGGCGCGGCAAGGGCCA CGACGGCCTGTACCAGGGCCTGTCCACAGCAACAAAGGATA CTTATGACGCTCTGCACATGCAGGCTCTGCCCCCTCGGTGA</p>
<p>7</p>	<p>anti-CD19 CAR _v1.6</p>	<p>ATGGGAACCAGCCTGCTGTGCTGGATGGCACTGTGCCTGCT GGGAGCAGACCACGCCGATGCCGAACGCCTACTCAGGGGA CATTCTCTAATGTGAGCACCAACGTGAGCTCTGGAGGAGGA GGCTCCGAGCTGCCAACCCAGGGCACATTCTCTAATGTGAG CACAAACGTGTCTCCCGCAAGCCTACCACAACCGCCGAAC TGCTACCCAGGGCACATTTTCCAACGTGTCTACCAACGTGT CTAGCGGAGGAGGAGGCTCCCCCGCACCTAGGCCCCCTACC CCAGCACCAACAATCGCAAGCCAGCCTCTGTCCCTGCGCCC AGAGGCATGCAGGCCAGCAGCAGGAGGAGCAGTGCACACC CGCGGCCTGGACTTTGCCTGCGATATCTATATCTGGGCACCA CTGGCAGGAACCTGTGGCGTGCTGCTGCTGTCTTGGTCATC ACCTGTACTGCAATCACAGAAACCGGAGAAGGGTGTGCAA GTGTCTCGGCCAGTGGTGAGAGCAGAGGGAAGGGGCAGCC TGCTGACCTGTGGCGACGTGGAGGAGAATCCCGGCCCTATG GAGACAGATACTGCTGCTGTGGGTGCTGCTGCTGTGGGT GCCAGGCTCTACAGGAGAGGTGCAGCTGCAGCAGAGCGGAC CTGAGCTGATCAAGCCAGGCGCCTCTGTGAAGATGAGCTGC AAGGCCTCCGGCTATACCTTCAAGCTACGTGATGCACTG GGTGAAGCAGAAGCCAGGCCAGGGCCTGGAGTGGATCGGCT ATATCAATCCCTACAACGACGGCACCAAGTATAACGAGAAG TTTAAGGGCAAGGCCACCCTGACATCCGATAAGAGCTCCTC TACCGCCTACATGGAGCTGAGCTCCCTGACATCCGAGGACT CTGCCGTGTAATAATTGCGCCAGAGGCACCTACTATTACGGCT CTAGGGTGTTCGATTACTGGGGCCAGGGCACAACCCTGACA GTGTCTAGCGGAGGAGGAGGCTCTGGAGGAGGAGGCAGCG GCGGCGGAGGCTCCGACATCGTGATGACCCAGGCAGCACCA TCCATCCAGTGACACCTGGCGAGAGCGTGTCCATCTCTTGT</p>

		<p>AGGTCCTCTAAGTCTCTGCTGAACAGCAATGGCAACACCTAT CTGTACTGGTTTCTGCAGCGGCCCGGACAGAGCCCTCAGCTG CTGATCTATAGGATGTCCAATCTGGCCTCTGGAGTGCCTGAT CGCTTCAGCGGCTCCGGCTCTGGAACCGCCTTTACACTGAGG ATCTCCC GCGTGGAGGCAGAGGACGTGGGCGTGTATTACTG CATGCAGCACCTGGAGTACCCTTTACCTTTGGCGCCGGCAC AAAGCTGGAGCTGAAGCGGAGCGACCCCAACACACCA GCACCCCGGCCACCAACCCCTGCCCTACAATCGCAAGCCA GCCACTGTCCCTGCGGCCCGAGGCCTGTAGACCTGCCGCCG GCGGCGCCGTCCATAACCGCGGCCTGGATTTGCCTGCGATA TCTACATTTGGGCCCTCTGGCCGGCACTTGGCGGTGCTGC TGCTGAGCCTGGTCATCACCTGTATTGCAAGCGGGGCAGA AAGAAGCTGCTGTACATCTTCAAGCAGCCCTTCATGCGGCC GTGCAGACCACACAGGAGGAGGATGGCTGCTCCTGTAGATT CCCAGAGGAGGAGGAGGGAGGATGTGAGCTGCGCGTGAAG TTTAGCCGGTCCGCCGACGCACCTGCATATCAGCAGGGCCA GAACCAGCTGTACAATGAGCTGAACCTGGGCCGGAGAGAGG AGTACGACGTGCTGGATAAGAGAAGGGGACGGGACCCCGA GATGGGAGGCAAGCCCGCCGGAAGAATCCTCAGGAGGGC CTGTATAACGAGCTGCAGAAGGATAAGATGGCCGAGGCCTA CAGCGAGATCGGCATGAAGGAGAGAGAAGGCGCGGCAAG GGACACGACGGCCTGTATCAGGGCCTGTCCACCGCCACAAA GGACACCTACGATGCCCTGCACATGCAGGCCCTGCCTCCAA GATGA</p>
--	--	---

b. Safety switches and monoclonal antibody specific-epitopes

Safety Switches

[070] It will be appreciated that adverse events may be minimized by transducing the immune cells (containing one or more CARs) with a suicide gene other than a rituximab-binding epitope. It may also be desired to incorporate an inducible “on” or “accelerator” switch into the immune cells. Suitable techniques include use of inducible caspase-9 (U.S. Appl. 2011/0286980) or a thymidine kinase, before, after or at the same time, as the cells are transduced with the CAR construct of the present disclosure. Additional methods for introducing suicide genes and/or “on” switches include TALENS, zinc fingers, RNAi, siRNA, shRNA, antisense technology, and other techniques known in the art.

[071] In accordance with the disclosure, additional on-off or other types of control switch techniques may be incorporated herein. These techniques may employ the use of dimerization domains and optional activators of such domain dimerization. These techniques include, e.g., those

described by Wu et al., Science 2014 350 (6258) utilizing FKBP/Rapalog dimerization systems in certain cells, the contents of which are incorporated by reference herein in their entirety. Additional dimerization technology is described in, e.g., Fegan et al. Chem. Rev. 2010, 110, 3315-3336 as well as U.S. Pat. Nos. 5,830,462; 5,834,266; 5,869,337; and 6,165,787, the contents of which are also
 5 incorporated by reference herein in their entirety. Additional dimerization pairs may include cyclosporine-A/cyclophilin, receptor, estrogen/estrogen receptor (optionally using tamoxifen), glucocorticoids/glucocorticoid receptor, tetracycline/tetracycline receptor, vitamin D/vitamin D receptor. Further examples of dimerization technology can be found in e.g., WO 2014/127261, WO 2015/090229, US 2014/0286987, US2015/0266973, US2016/0046700, U.S. Pat. No. 8,486,693, US
 10 2014/0171649, and US 2012/0130076, the contents of which are further incorporated by reference herein in their entirety.

[072] In some embodiments, the CAR-immune cell (e.g., CAR-T cell) of the disclosure comprises a polynucleotide encoding a suicide polypeptide that is deficient in rituximab binding. In some embodiments, the suicide peptide comprises a mutated RQR8 sequence. See, e.g.,
 15 WO2013153391A, which is hereby incorporated by reference in its entirety. In CAR-immune cell (e.g., CAR-T cell) cells comprising the polynucleotide, the suicide polypeptide is expressed at the surface of a CAR-immune cell (e.g., CAR-T cell). In some embodiments, the suicide polypeptide comprises the amino acid sequence shown in SEQ ID NO: 19.

CPYSNPSLCSGGGGSELPTQGTFNSVSTNVSPAKPTTTACPYSNPSLCSGGGGSP
 20 APRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYWAPLAGTCGVLLLS
 LVITLYCNHRNRRRVCKCPRPVV (SEQ ID NO: 19).

[073] The suicide polypeptide may also comprise a signal peptide at the amino terminus—for example, MGTSLLCWMALCLLGADHADA (SEQ ID NO: 20). In some embodiments, the suicide polypeptide comprises the amino acid sequence shown in SEQ ID NO: 21, which includes the signal
 25 sequence of SEQ ID NO: 20.

MGTSLLCWMALCLLGADHADACPYSNPSLCSGGGGSELPTQGTFNSVSTNVSPAKPTTTAC
 PYSNPSLCSGGGGSPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYWAPL
 AGTCGVLLLSLVITLYCNHRNRRRVCKCPRPVV (SEQ ID NO: 21).

[074] In certain embodiments, the suicide peptide comprises an amino acid sequence that
 30 comprises a one or more mutated residues, inserted residues, or deleted residues that reduce or eliminate rituximab binding.

[075] When the suicide polypeptide is expressed at the surface of a CAR-immune cell (e.g., CAR-T cell), antibody binding to the suicide gene epitopes of the polypeptide causes lysis of the cell. Deletion of CD19-specific CAR-immune cell (e.g., CAR-T cell) may occur *in vivo*, for
 35 example by administering a suicide agent to a patient. The decision to delete the transferred cells may arise from undesirable effects being detected in the patient which are attributable to the

transferred cells, such as for example, when unacceptable levels of toxicity are detected. As used herein, a “suicide agent” refers to a molecule that binds to the CAR immune cell and causes lysis of the CAR expressing immune cell.

5 [076] In some embodiments, a suicide polypeptide is expressed on the surface of the cell. In some embodiments, a suicide polypeptide is included in the CAR construct. In some embodiments, a suicide polypeptide is not part of the CD19 CAR construct.

[077] In some embodiments, the extracellular domain of any one of the CD19-specific CARs disclosed herein may comprise one or more epitopes specific for (*i.e.*, specifically recognized by) a monoclonal antibody. These epitopes are also referred to herein as mAb-specific epitopes.
10 Exemplary mAb-specific epitopes are disclosed in International Patent Publication No. WO 2016/120216, which is incorporated herein in its entirety. In these embodiments, the extracellular domain of the CARs comprise antigen binding domains that specifically bind to CD19 and one or more epitopes that bind to one or more monoclonal antibodies (mAbs). CARs comprising the mAb-specific epitopes can be single-chain or multi-chain.

15 [078] The inclusion of epitopes specific for monoclonal antibodies in the extracellular domain of the CARs described herein allows sorting and depletion of engineered immune cells expressing the CARs. In some embodiments, this feature also promotes recovery of endogenous CD19-expressing cells that were depleted by administration of engineered immune cells expressing the CARs. In some embodiments, allowing for depletion provides a safety switch in case of deleterious
20 effects, e.g., upon administration to a subject.

[079] Accordingly, in some embodiments, the present disclosure relates to a method for sorting and/or depleting the engineered immune cells endowed with the CARs comprising mAb-specific epitopes and a method for promoting recovery of endogenous CD19-expressing cells.

25 [080] Several epitope-monoclonal antibody couples can be used to generate CARs comprising monoclonal antibody specific epitopes; in particular, those already approved for medical use or for use in GMP manufacturing, such as CD34 epitope/QBEND-10 as a non-limiting example.

[081] The disclosure also encompasses methods for sorting the engineered immune cells endowed with the CD19-specific CARs expressing the mAb-specific epitope(s) and therapeutic
30 methods where the activation of the engineered immune cells endowed with these CARs is modulated by depleting the cells using an antibody that targets the external ligand binding domain of said CARs. Table 2 provides exemplary mimotope sequences that can be inserted into the extracellular domains of the CARs of the disclosure.

Table 2: Exemplary mimotope sequences

<u>Rituximab</u>		
Mimotope	SEQ ID NO: 22	CPYSNPSLC
<u>Palivizumab</u>		
Epitope	SEQ ID NO: 23	NSELLSLINDMPITNDQKKLMSNN
<u>Cetuximab</u>		
Mimotope 1	SEQ ID NO: 24	CQFDLSTRRLKC
Mimotope 2	SEQ ID NO: 25	CQYNLSSRALKC
Mimotope 3	SEQ ID NO: 26	CVWQRWQKSYVC
Mimotope 4	SEQ ID NO: 27	CMWDRFSRWYKC
<u>Nivolumab</u>		
Epitope 1	SEQ ID NO: 28	SFVLNWYRMSPSNQTDKLAAFPEDR
Epitope 2	SEQ ID NO: 29	SGTYLCGAISLAPKAQIKE
<u>OBEND-10</u>		
Epitope	SEQ ID NO: 30	ELPTQGTFSNVSTNVS
<u>Alemtuzumab</u>		
Epitope	SEQ ID NO: 31	GQNDTSQTSSPS

[082] In certain embodiments, the CAR comprises an epitope or mimotope amino acid sequence that is at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identical to the epitope or mimotope amino acid sequences set forth herein in Table 2. In certain embodiments, the CAR comprises an epitope or mimotope amino acid sequence that is not or does not comprise SEQ ID NO: 22. In certain embodiments, the CAR comprises an epitope or mimotope comprising the amino acid sequence of SEQ ID NO: 30.

10

c. Hinge Domain

[083] The extracellular domain of the CARs of the disclosure may comprise a “hinge” domain (or hinge region). The term generally to any polypeptide that functions to link the transmembrane domain in a CAR to the extracellular antigen binding domain in a CAR. In particular, hinge domains can be used to provide more flexibility and accessibility for the extracellular antigen binding domain.

15

[084] A hinge domain may comprise up to 300 amino acids—in some embodiments 10 to 100 amino acids or in some embodiments 25 to 50 amino acids. The hinge domain may be derived from all or part of naturally occurring molecules, such as from all or part of the extracellular region of CD8, CD4, CD28, 4-1BB, or IgG (in particular, the hinge region of an IgG; it will be appreciated that the hinge region may contain some or all of a member of the immunoglobulin family such as

20

IgG1, IgG2, IgG3, IgG4, IgA, IgD, IgE, IgM, or fragment thereof), or from all or part of an antibody heavy-chain constant region. Alternatively, the hinge domain may be a synthetic sequence that corresponds to a naturally occurring hinge sequence, or may be an entirely synthetic hinge sequence. In some embodiments said hinge domain is a part of human CD8 α chain (e.g., NP_001139345.1). In another particular embodiment, said hinge and transmembrane domains comprise a part of human CD8 α chain. In some embodiments, the hinge domain of CARs described herein comprises a subsequence of CD8 α , an IgG1, IgG4, PD-1 or an Fc γ RIII α , in particular the hinge region of any of an CD8 α , an IgG1, IgG4, PD-1 or an Fc γ RIII α . In some embodiments, the hinge domain comprises a human CD8 α hinge, a human IgG1 hinge, a human IgG4, a human PD-1 or a human Fc γ RIII α hinge. In some embodiments the CARs disclosed herein comprise a scFv, CD8 α human hinge and transmembrane domains, the CD3 ζ signaling domain, and 4-1BB signaling domain. Table 3 provides amino acid sequences for exemplary hinges provided herein.

Table 3 : Exemplary hinges

Domain	Amino Acid Sequence	SEQ ID NO:
Fc γ RIII α hinge	GLAVSTISSFFPPGYQ	32
CD8 α hinge	TTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACD	33
IgG1 hinge	EPKSPDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMIARTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGK	34

[085] In certain embodiments, the hinge region comprises an amino acid sequence that is at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or 100% identical to the hinge domain amino acid sequences set forth herein in Table 3.

d. Transmembrane Domain

[086] The CARs of the disclosure are designed with a transmembrane domain that is fused to the extracellular domain of the CAR. It can similarly be fused to the intracellular domain of the CAR. In some instances, the transmembrane domain can be selected or modified by amino acid substitution to avoid binding of such domains to the transmembrane domains of the same or

different surface membrane proteins to minimize interactions with other members of the receptor complex. In some embodiments, short linkers may form linkages between any or some of the extracellular, transmembrane, and intracellular domains of the CAR. In some embodiments, the linker comprises a glycine repeat sequence. In some embodiments, the linker comprises (GGGS)_n,
5 wherein n is 1, 2, 3, 4, or 5 (SEQ ID NO: 41).

[087] Suitable transmembrane domains for a CAR disclosed herein have the ability to (a) be expressed at the surface an immune cell such as, for example without limitation, a lymphocyte cell, such as a T helper (T_h) cell, cytotoxic T (T_c) cell, T regulatory (T_{reg}) cell, or Natural killer (NK) cells, and/or (b) interact with the extracellular antigen binding domain and intracellular signaling
10 domain for directing the cellular response of an immune cell against a target cell.

[088] The transmembrane domain may be derived either from a natural or from a synthetic source. Where the source is natural, the domain may be derived from any membrane-bound or transmembrane protein.

[089] Transmembrane regions of particular use in this disclosure may be derived from
15 (comprise, or correspond to) CD28, OX-40, 4-1BB/CD137, CD2, CD7, CD27, CD30, CD40, programmed death-1 (PD-1), inducible T cell costimulator (ICOS), lymphocyte function-associated antigen-1 (LFA-1, CD1-1a/CD18), CD3 gamma, CD3 delta, CD3 epsilon, CD247, CD276 (B7-H3), LIGHT, (TNFSF14), NKG2C, Ig alpha (CD79a), DAP-10, Fc gamma receptor, MHC class I molecule, TNF receptor proteins, an Immunoglobulin protein, cytokine receptor, integrins, Signaling
20 Lymphocytic Activation Molecules (SLAM proteins), activating NK cell receptors, BTLA, a Toll ligand receptors, ICAM-1, B7-H3, CDS, ICAM-1, GITR, BAFFR, LIGHT, HVEM (LIGHTR), KIRDS2, SLAMF7, NKp80 (KLRF1), NKp44, NKp30, NKp46, CD19, CD4, CD8alpha, CD8beta, IL-2R beta, IL-2R gamma, IL-7R alpha, ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD1 1d, ITGAE, CD103, ITGAL, CD1 1a, LFA-1, ITGAM, CD1 1b,
25 ITGAX, CD1 1c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, NKG2D, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRT AM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3), BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, PAG/Cbp, CD19a, a ligand that specifically binds with
30 CD83, or any combination thereof.

[090] As non-limiting examples, the transmembrane region can be a derived from, or be a portion of a T cell receptor such as α , β , γ or δ , polypeptide constituting CD3 complex, IL-2 receptor p55 (a chain), p75 (β chain) or γ chain, subunit chain of Fc receptors, in particular Fc γ receptor III or CD proteins. Alternatively, the transmembrane domain can be synthetic and can comprise
35 predominantly hydrophobic residues such as leucine and valine. In some embodiments said transmembrane domain is derived from the human CD8 α chain (*e.g.*, NP_001139345.1).

[091] In some embodiments, the transmembrane domain in the CAR of the disclosure is a CD8 α transmembrane domain. In some embodiments, the transmembrane domain in the CAR of the disclosure is a CD8 α transmembrane domain comprising the amino acid sequence

IYIWAPLAGTCGVLLLSLVIT (SEQ ID NO: 35). In some embodiments, the CD8 α

5 transmembrane domain comprises the nucleic acid sequence that encodes the transmembrane amino acid sequence of SEQ ID NO: 35. In some embodiments, the hinge and transmembrane domain in the CAR of the disclosure is a CD8 α hinge and transmembrane domain comprising the amino acid sequence

TTTTAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSL
10 VIT (SEQ ID NO: 36).

e. Intracellular Domain

[092] The intracellular (cytoplasmic) domain of the CARs of the disclosure can provide activation of at least one of the normal effector functions of the immune cell comprising the CAR.

15 Effector function of a T cell, for example, may refer to cytolytic activity or helper activity, including the secretion of cytokines.

In some embodiments, an activating intracellular signaling domain for use in a CAR can be the cytoplasmic sequences of, for example without limitation, the T cell receptor and co-receptors that act in concert to initiate signal transduction following antigen receptor engagement, as well as any
20 derivative or variant of these sequences and any synthetic sequence that has the same functional capability.

[093] It will be appreciated that suitable (e.g., activating) intracellular domains include, but are not limited to signaling domains derived from (or corresponding to) CD28, OX-40, 4-

1BB/CD137, CD2, CD7, CD27, CD30, CD40, programmed death-1 (PD-1), inducible T cell
25 costimulator (ICOS), lymphocyte function-associated antigen-1 (LFA-1, CD1-1a/CD18), CD3

gamma, CD3 delta, CD3 epsilon, CD247, CD276 (B7-H3), LIGHT, (TNFSF14), NKG2C, Ig alpha
(CD79a), DAP-10, Fc gamma receptor, MHC class I molecule, TNF receptor proteins, an

Immunoglobulin protein, cytokine receptor, integrins, Signaling Lymphocytic Activation Molecules
(SLAM proteins), activating NK cell receptors, BTLA, a Toll ligand receptors, ICAM-1, B7-H3,

30 CDS, ICAM-1, GITR, BAFFR, LIGHT, HVEM (LIGHTR), KIRDS2, SLAMF7, NKp80 (KLRF1),
NKp44, NKp30, NKp46, CD19, CD4, CD8alpha, CD8beta, IL-2R beta, IL-2R gamma, IL-7R alpha,

ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD1 1d, ITGAE,
CD103, ITGAL, CD1 1a, LFA-1, ITGAM, CD1 1b, ITGAX, CD1 1c, ITGB1, CD29, ITGB2,

CD18, LFA-1, ITGB7, NKG2D, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4

35 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRT AM, Ly9 (CD229), CD160 (BY55),

PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3),

BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, PAG/Cbp, CD19a, a ligand that specifically binds with CD83, or any combination thereof.

[094] The intracellular domains of the CARs of the disclosure may incorporate, in addition to the activating domains described above, co-stimulatory signaling domains (interchangeably referred to herein as costimulatory molecules) to increase their potency. Costimulatory domains can provide a signal in addition to the primary signal provided by an activating molecule as described herein.

[095] It will be appreciated that suitable costimulatory domains within the scope of the disclosure can be derived from (or correspond to) for example, CD28, OX40, 4-1BB/CD137, CD2, CD3 (alpha, beta, delta, epsilon, gamma, zeta), CD4, CD5, CD7, CD9, CD16, CD22, CD27, CD30, CD 33, CD37, CD40, CD 45, CD64, CD80, CD86, CD134, CD137, CD154, PD-1, ICOS, lymphocyte function-associated antigen-1 (LFA-1 (CD1 1a/CD18), CD247, CD276 (B7-H3), LIGHT (tumor necrosis factor superfamily member 14; TNFSF14), NKG2C, Ig alpha (CD79a), DAP-10, Fc gamma receptor, MHC class I molecule, TNFR, integrin, signaling lymphocytic activation molecule, BTLA, Toll ligand receptors, ICAM-1, B7-H3, CDS, ICAM-1, GITR, BAFFR, LIGHT, HVEM (LIGHTR), KIRDS2, SLAMF7, NKp80 (KLRF1), NKp44, NKp30, NKp46, CD19, CD4, CD8alpha, CD8beta, IL-2R beta, IL-2R gamma, IL-7R alpha, ITGA4, VLA1, CD49a, ITGA4, IA4, CD49D, ITGA6, VLA-6, CD49f, ITGAD, CD1-1d, ITGAE, CD103, ITGAL, CD1-1a, LFA-1, ITGAM, CD1-1b, ITGAX, CD1-1c, ITGB1, CD29, ITGB2, CD18, LFA-1, ITGB7, NKG2D, TNFR2, TRANCE/RANKL, DNAM1 (CD226), SLAMF4 (CD244, 2B4), CD84, CD96 (Tactile), CEACAM1, CRT AM, Ly9 (CD229), CD160 (BY55), PSGL1, CD100 (SEMA4D), CD69, SLAMF6 (NTB-A, Ly108), SLAM (SLAMF1, CD150, IPO-3), BLAME (SLAMF8), SELPLG (CD162), LTBR, LAT, GADS, SLP-76, PAG/Cbp, CD19a, CD83 ligand, or fragments or combinations thereof. It will be appreciated that additional costimulatory molecules, or fragments thereof, not listed above are within the scope of the disclosure.

[096] In some embodiments, the intracellular/cytoplasmic domain of the CAR can be designed to comprise the 41BB/CD137 domain by itself or combined with any other desired intracellular domain(s) useful in the context of the CAR of the disclosure. The complete native amino acid sequence of 41BB/CD137 is described in NCBI Reference Sequence: NP_001552.2. The complete native 41BB/CD137 nucleic acid sequence is described in NCBI Reference Sequence: NM_001561.5.

[097] In some embodiments, the intracellular/cytoplasmic domain of the CAR can be designed to comprise the CD28 domain by itself or combined with any other desired intracellular domain(s) useful in the context of the CAR of the disclosure. The complete native amino acid sequence of CD28 is described in NCBI Reference Sequence: NP_006130.1. The complete native CD28 nucleic acid sequence is described in NCBI Reference Sequence: NM_006139.1.

[098] In some embodiments, the intracellular/cytoplasmic domain of the CAR can be designed to comprise the CD3 zeta domain by itself or combined with any other desired intracellular domain(s) useful in the context of the CAR of the disclosure. In some embodiments, the intracellular signaling domain of the CAR can comprise the CD3 ζ signaling domain which has amino acid sequence with at least about 70%, at least 80%, at least 90%, 95%, 97%, or 99% sequence identity with an amino acid sequence shown in SEQ ID NO: 38. For example, the intracellular domain of the CAR can comprise a CD3 zeta chain portion and a portion of a costimulatory signaling molecule. The intracellular signaling sequences within the intracellular signaling portion of the CAR of the disclosure may be linked to each other in a random or specified order. In some embodiments, the intracellular domain is designed to comprise the activating domain of CD3 zeta and a signaling domain of CD28.

[099] In some embodiments, the intracellular domain is designed to comprise the activating domain of CD3 zeta and a signaling domain of 4-1BB. In some embodiments, the 4-1BB (intracellular domain) comprises the amino acid sequence

KRGRKKLLYIFKQPFMRPVQTTQEEDGCSRFPEEEEGGCEL (SEQ ID NO: 37).

[0100] The CD3 zeta amino acid sequence may comprise SEQ ID NO: 38.

LRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPENGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHGDLGQGLSTATKDTYDALHMQUALPPR (SEQ ID NO: 38).

[0101] In some embodiments the intracellular signaling domain of the CAR of the disclosure comprises a domain of a co-stimulatory molecule. In some embodiments, the intracellular signaling domain of a CAR of the disclosure comprises a part of co-stimulatory molecule selected from the group consisting of fragment of 41BB (GenBank: AAA53133.) and CD28 (NP_006130.1). In some embodiments, the intracellular signaling domain of the CAR of the disclosure comprises amino acid sequence which comprises at least 70%, at least 80%, at least 90%, 95%, 97%, or 99% sequence identity with an amino acid sequence shown in SEQ ID NO: 37 and SEQ ID NO: 38. In some embodiments, the intracellular signaling domain of the CAR of the disclosure comprises amino acid sequence which comprises at least 70%, at least 80%, at least 90%, 95%, 97%, or 99% sequence identity with an amino acid sequence shown in SEQ ID NO: 37 and/or at least 70%, at least 80%, at least 90%, 95%, 97%, or 99% sequence identity with an amino acid sequence shown in SEQ ID NO: 38.

[0102] In exemplary embodiments, a CAR of the disclosure comprises, from N-terminus to C-terminus: a CD8 α signal sequence, a CD19 scFv, a CD8 α hinge and transmembrane region, a 41BB cytoplasmic signaling domain, and a CD3 ζ cytoplasmic signaling domain.

III. Immune Cells Comprising CARs

a. Immune Cells

[0103] Provided herein are engineered immune cells expressing the CARs of the disclosure (e.g., CAR-T cells).

5 [0104] In some embodiments, an engineered immune cell comprises a population of CARs, each CAR comprising different extracellular antigen-binding domains. In some embodiments, an immune cell comprises a population of CARs, each CAR comprising the same extracellular antigen-binding domains.

[0105] The engineered immune cells can be allogeneic or autologous.

10 [0106] In some embodiments, the engineered immune cell is a T cell (e.g., inflammatory T-lymphocyte cytotoxic T-lymphocyte, regulatory T-lymphocyte, helper T-lymphocyte, tumor infiltrating lymphocyte (TIL)), NK cell, NK-T-cell, TCR-expressing cell, dendritic cell, killer dendritic cell, a mast cell, or a B-cell. In some embodiments, the cell can be derived from the group consisting of CD4+ T-lymphocytes and CD8+ T-lymphocytes. In some exemplary embodiments, the
15 engineered immune cell is a T cell. In some exemplary embodiments, the engineered immune cell is a gamma delta T cell. In some exemplary embodiments, the engineered immune cell is a macrophage.

[0107] In some embodiments, the engineered immune cell can be derived from, for example
20 without limitation, a stem cell. The stem cells can be adult stem cells, non-human embryonic stem cells, more particularly non-human stem cells, cord blood stem cells, progenitor cells, bone marrow stem cells, induced pluripotent stem cells, totipotent stem cells or hematopoietic stem cells.

[0108] In some embodiments, the cell is obtained or prepared from peripheral blood. In some
25 embodiments, the cell is obtained or prepared from peripheral blood mononuclear cells (PBMCs). In some embodiments, the cell is obtained or prepared from bone marrow. In some embodiments, the cell is obtained or prepared from umbilical cord blood. In some embodiments, the cell is a human cell. In some embodiments, the cell is transfected or transduced by the nucleic acid vector using a method selected from the group consisting of electroporation, sonoporation, biolistics (e.g., Gene Gun), lipid transfection, polymer transfection, nanoparticles, viral transfection (e.g., retrovirus, lentivirus, AAV) or polyplexes.

30 [0109] In some embodiments, the engineered immune cells expressing at their cell surface membrane a CD19-specific CAR of the disclosure comprise a percentage of stem cell memory and central memory cells greater than 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100%. In some embodiments, the engineered immune cells expressing at their cell surface membrane a CD19-specific CAR of the disclosure comprise a percentage of stem cell memory and central memory cells
35 of about 10% to about 100%, about 10% to about 90%, about 10% to about 80%, about 10% to about 70%, about 10% to about 60%, about 10% to about 50%, about 10% to about 40%, about 10%

to about 30%, about 10% to about 20%, about 15% to about 100%, about 15% to about 90%, about 15% to about 80%, about 15% to about 70%, about 15% to about 60%, about 15% to about 50%, about 15% to about 40%, about 15% to about 30%, about 20% to about 100%, about 20% to about 90%, about 20% to about 80%, about 20% to about 70%, about 20% to about 60%, about 20% to about 50%, about 20% to about 40%, about 20% to about 30%, about 30% to about 100%, about 30% to about 90%, about 30% to about 80%, about 30% to about 70%, about 30% to about 60%, about 30% to about 50%, about 30% to about 40%, about 40% to about 100%, about 40% to about 90%, about 40% to about 80%, about 40% to about 70%, about 40% to about 60%, about 40% to about 50%, about 50% to about 100%, about 50% to about 90%, about 50% to about 80%, about 50% to about 70%, about 50% to about 60%, about 60% to about 100%, about 60% to about 90%, about 60% to about 80%, about 60% to about 70%, about 70% to about 90%, about 70% to about 80%, about 80% to about 100%, about 80% to about 90%, about 90% to about 100%, about 25% to about 50%, about 75% to about 100%, or about 50% to about 75%.

[0110] In some embodiments, the immune cell is an inflammatory T-lymphocyte that expresses any one of the CARs described herein. In some embodiments, the immune cell is a cytotoxic T-lymphocyte that expresses any one of the CARs described herein. In some embodiments, the immune cell is a regulatory T-lymphocyte that expresses any one of the CARs described herein. In some embodiments, the immune cell is a helper T-lymphocyte that expresses any one of the CARs described herein.

[0111] Prior to expansion and genetic modification, a source of cells can be obtained from a subject through a variety of non-limiting methods. Cells can be obtained from a number of non-limiting sources, including peripheral blood mononuclear cells, bone marrow, lymph node tissue, cord blood, thymus tissue, stem cell- or iPSC-derived T cells or NK cells, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In some embodiments, any number of T cell lines available and known to those skilled in the art, may be used. In some embodiments, cells can be derived from a healthy donor, from a patient diagnosed with cancer or from a patient diagnosed with an infection. In some embodiments, cells can be part of a mixed population of cells which present different phenotypic characteristics.

[0112] Also provided herein are cell lines obtained from a transformed immune cell (e.g., T-cell) according to any of the above-described methods. Also provided herein are modified cells resistant to an immunosuppressive treatment. In some embodiments, an isolated cell according to the disclosure comprises a polynucleotide encoding a CAR.

[0113] The immune cells of the disclosure can be activated and expanded, either prior to or after genetic modification of the immune cells, using methods as generally known. Generally, the engineered immune cells of the disclosure can be expanded, for example, by contacting with an agent that stimulates a CD3 TCR complex and a co-stimulatory molecule on the surface of the T-

cells to create an activation signal for the T cell. For example, chemicals such as calcium ionophore A23187, phorbol 12-myristate 13-acetate (PMA), or mitogenic lectins like phytohemagglutinin (PHA) can be used to create an activation signal for the T cell.

[0114] In some embodiments, T cell populations may be stimulated in vitro by contact with, for example, an anti-CD3 antibody, or antigen-binding fragment thereof, or an anti-CD28 antibody immobilized on a surface, or by contact with a protein kinase C activator (*e.g.*, bryostatin) in conjunction with a calcium ionophore. For co-stimulation of an accessory molecule on the surface of the T cells, a ligand that binds the accessory molecule is used. For example, a population of T cells can be contacted with an anti-CD3 antibody and an anti-CD28 antibody, under conditions appropriate for stimulating proliferation of the T cells. The anti-CD3 antibody and an anti-CD28 antibody can be disposed on a bead or plate or other substrate. Conditions appropriate for T cell culture include an appropriate media (*e.g.*, Minimal Essential Media or RPMI Media 1640 or, X-vivo 15, (Lonza)) that may contain factors necessary for proliferation and viability, including serum (*e.g.*, fetal bovine or human serum), interleukin-2 (IL-2), insulin, IFN- γ , IL-4, IL-7, GM-CSF, IL-10, IL-2, IL-15, TGFbeta, and TNF, or any other additives for the growth of cells known to the skilled artisan. Other additives for the growth of cells include, but are not limited to, surfactant, plasmanate, and reducing agents such as N-acetyl- cysteine and 2-mercaptoethanoi. Media can include RPMI 1640, AIM-V, DMEM, MEM, a- MEM, F-12, X-Vivo 15, and X-Vivo 20, Optimizer, with added amino acids, sodium pyruvate, and vitamins, either serum-free or supplemented with an appropriate amount of serum (or plasma) or a defined set of hormones, and/or an amount of cytokine(s) sufficient for the growth and expansion of T cells (*e.g.*, IL-7 and/or IL-15). Antibiotics, *e.g.*, penicillin and streptomycin, are included only in experimental cultures, not in cultures of cells that are to be infused into a subject. The target cells are maintained under conditions necessary to support growth, for example, an appropriate temperature (*e.g.*, 37° C) and atmosphere (*e.g.*, air plus 5% CO₂). T cells that have been exposed to varied stimulation times may exhibit different characteristics

In some embodiments, the cells of the disclosure can be expanded by co-culturing with tissue or cells. The cells can also be expanded in vivo, for example in the subject's blood after administering the cell into the subject.

[0115] In some embodiments, an engineered immune cell according to the present disclosure may comprise one or more disrupted or inactivated genes. In some embodiments, an engineered immune cell according to the present disclosure comprises one disrupted or inactivated gene selected from the group consisting of CD52, CD19, GR, PD-1, CTLA-4, LAG3, TIM3, BTLA, BY55, TIGIT, B7H5, LAIR1, SIGLEC10, 2B4, HLA, TCR α and TCR β and/or expresses a CAR, a multi-chain CAR and/or a pT α transgene. In some embodiments, an isolated cell comprises polynucleotides encoding polypeptides comprising a multi-chain CAR. In some embodiments, the

isolated cell according to the present disclosure comprises two disrupted or inactivated genes selected from the group consisting of: CD52 and GR, CD52 and TCR α , CDR52 and TCR β , CD19 and CD52, CD19 and TCR α , CD19 and TCR β , GR and TCR α , GR and TCR β , TCR α and TCR β , PD-1 and TCR α , PD-1 and TCR β , CTLA-4 and TCR α , CTLA-4 and TCR β , LAG3 and TCR α , LAG3 and TCR β , TIM3 and TCR α , Tim3 and TCR β , BTLA and TCR α , BTLA and TCR β , BY55 and TCR α , BY55 and TCR β , TIGIT and TCR α , TIGIT and TCR β , B7H5 and TCR α , B7H5 and TCR β , LAIR1 and TCR α , LAIR1 and TCR β , SIGLEC10 and TCR α , SIGLEC10 and TCR β , 2B4 and TCR α , 2B4 and TCR β and/or expresses a CAR, a multi-chain CAR and a pT α transgene. In some embodiments the method comprises disrupting or inactivating one or more genes by introducing into the cells an endonuclease able to selectively inactivate a gene by selective DNA cleavage. In some embodiments the endonuclease can be, for example, a zinc finger nuclease (ZFN), megaTAL nuclease, meganuclease, transcription activator-like effector nuclease (TALE-nuclease), or CRIPR (e.g., Cas9) endonuclease.

[0116] In some embodiments, TCR is rendered not functional in the cells according to the disclosure by disrupting or inactivating TCR α gene and/or TCR β gene(s). In some embodiments, a method to obtain modified cells derived from an individual is provided, wherein the cells can proliferate independently of the major histocompatibility complex (MHC) signaling pathway. Modified cells, which can proliferate independently of the MHC signaling pathway, susceptible to be obtained by this method are encompassed in the scope of the present disclosure. Modified cells disclosed herein can be used in for treating patients in need thereof against Host versus Graft (HvG) rejection and Graft versus Host Disease (GvHD); therefore in the scope of the present disclosure is a method of treating patients in need thereof against Host versus Graft (HvG) rejection and Graft versus Host Disease (GvHD) comprising treating said patient by administering to said patient an effective amount of modified cells comprising disrupted or inactivated TCR α and/or TCR β genes.

[0117] In some embodiments, the immune cells are engineered to be resistant to one or more chemotherapy drugs. The chemotherapy drug can be, for example, a purine nucleotide analogue (PNA), thus making the immune cell suitable for cancer treatment combining adoptive immunotherapy and chemotherapy. Exemplary PNAs include, for example, clofarabine, fludarabine, cyclophosphamide, and cytarabine, alone or in combination. PNAs are metabolized by deoxycytidine kinase (dCK) into mono-, di-, and tri-phosphate PNA. Their tri-phosphate forms compete with ATP for DNA synthesis, act as pro-apoptotic agents, and are potent inhibitors of ribonucleotide reductase (RNR), which is involved in trinucleotide production. Provided herein are CD19-specific CAR-T cells comprising a disrupted or inactivated dCK gene. In some embodiments, the dCK knockout cells are made by transfection of T cells using polynucleotides encoding specific TAL-nulcease directed against dCK genes by, for example, electroporation of mRNA. The dCK knockout CD19-specific CAR-T cells are resistant to PNAs, including for

example clorofarabine and/or fludarabine, and maintain T cell cytotoxic activity toward CD19-expressing cells.

5 [0118] In some embodiments, isolated cells or cell lines of the disclosure can comprise a pTa or a functional variant thereof. In some embodiments, an isolated cell or cell line can be further genetically modified by disrupting or inactivating the TCR α gene.

[0119] The disclosure also provides engineered immune cells comprising any of the CAR polynucleotides described herein.

c. Methods of Making

10 [0120] Provided herein are methods of making the CARs and the CAR containing immune cells of the disclosure. A variety of known techniques can be utilized in making the polynucleotides, polypeptides, vectors, antigen binding domains, immune cells, compositions, and the like according to the disclosure.

15 Polynucleotides and Vectors

[0121] In some embodiments, a CAR can be introduced into an immune cell as a transgene via a plasmid vector. In some embodiments, the plasmid vector can also contain, for example, a selection marker which provides for identification and/or selection of cells which received the vector.

20 [0122] CAR polypeptides may be synthesized in situ in the cell after introduction of polynucleotides encoding the CAR polypeptides into the cell. Alternatively, CAR polypeptides may be produced outside of cells, and then introduced into cells. Methods for introducing a polynucleotide construct into cells are known in the art. In some embodiments, stable transformation methods (e.g., using a lentiviral vector) can be used to integrate the polynucleotide construct into the genome of the cell. In other embodiments, transient transformation methods can be used to transiently express the polynucleotide construct, and the polynucleotide construct not integrated into the genome of the cell. In other embodiments, virus-mediated methods can be used. The polynucleotides may be introduced into a cell by any suitable means such as for example, recombinant viral vectors (e.g., retroviruses, adenoviruses), liposomes, and the like. Transient transformation methods include, for example without limitation, microinjection, electroporation or particle bombardment. Polynucleotides may be included in vectors, such as for example plasmid vectors or viral vectors.

30 [0123] In some embodiments, isolated nucleic acids are provided comprising a promoter operably linked to a first polynucleotide encoding a CD19 antigen binding domain, at least one costimulatory molecule, and an activating domain. In some embodiments, the nucleic acid construct is contained within a viral vector. In some embodiments, the viral vector is selected from the group

consisting of retroviral vectors, murine leukemia virus vectors, SFG vectors, adenoviral vectors, lentiviral vectors, adeno-associated virus (AAV) vectors, Herpes virus vectors, and vaccinia virus vectors. In some embodiments, the nucleic acid is contained within a plasmid.

[0124] In one aspect, the present disclosure provides a polynucleotide sequence comprising a promoter that is capable of expressing a CAR transgene in a mammalian T cell. In some
5
embodiments, the promoter is the EF1a promoter. The native EF1a promoter drives expression of the alpha subunit of the elongation factor-1 complex, which is responsible for the enzymatic delivery of aminoacyl tRNAs to the ribosome. The EF1a promoter has been extensively used in mammalian expression plasmids and has been shown to be effective in driving CAR expression
10
from transgenes cloned into a lentiviral vector. See, e.g., Milone et al., Mol. Ther. 17(8): 1453-1464 (2009). In some embodiments, the EF1a promoter comprises the sequence provided as SEQ ID NO: 15.

CGGTGAGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCACATCGCCACAGTCCCCGAGA
AGTTGGGGGGAGGGGTCGGCAATTGAACCGGTGCCTAGAGAAGGTGGCGCGGGGTAA
15
ACTGGGAAAGTGATGTCGTGTAAGTGGCTCCGCCCTTTTCCCGAGGGTGGGGGAGAACCG
TATATAAGTGCAGTAGTCGCCGTGAACGTTCTTTTCGCAACGGGTTTGCCGCCAGA
ACACAGGTAAGTGCCGTGTGTGGTTCGCGGGCCTGGCCTCTTTACGGGTTATGGCCC
TTGCGTGCCTTGAATTACTTCCACGCCCTGGCTGCAGTACGTGATTCTTGATCCCCGAGC
TTCGGGTTGGAAGTGGGTGGGAGAGTTCGAGGCCTTGCCTTAAGGAGCCCCCTTCGCT
20
CGTGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGCCGCGCGTGCGAATCTGGTGGC
ACCTTCGCGCCTGTCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAAAATTTTTGATGAC
CTGCTGCGACGCTTTTTTTCTGGCAAGATAGTCTTGTAATGCGGGCCAAGATCTGCAC
ACTGGTATTTTCGGTTTTTGGGGCCGCGGGCGGCGACGGGGCCCGTGCCTCCAGCGCAC
ATGTTTCGGCGAGGCGGGCCTGCGAGCGCGGCCACCGAGAATCGGACGGGGGTAGTCT
25
CAAGCTGGCCGGCCTGCTCTGGTGCTGGCCTCGCGCCGCGTGTATCGCCCCGCCCTG
GGCGGCAAGGCTGGCCCGGTCCGCACCAGTTGCGTGAGCGGAAAGATGGCCGCTTCCC
GGCCCTGCTGCAGGGAGCTCAAAATGGAGGACGCGGCGCTCGGGAGAGCGGGCGGGT
GAGTCACCCACACAAAGGAAAAGGGCCTTTCCGTCTCAGCCGTCGCTTCATGTGACTC
CACGGAGTACCGGGCGCCGTCCAGGCACCTCGATTAGTTCTCGAGCTTTTGGAGTACGT
30
CGTCTTTAGGTTGGGGGGAGGGGTTTTATGCGATGGAGTTTCCCCACACTGAGTGGGTG
GAGACTGAAGTTAGGCCAGCTTGGCACTTGATGTAATTCCTTGGAAATTTGCCCTTTTT
GAGTTTGGATCTTGGTTCATTCTCAAGCCTCAGACAGTGGTTCAAAGTTTTTTCTTCCA
TTTCAGGTGTCGTGA (SEQ ID NO: 15)

[0125] The EF1a promoter sequence shown above comprises the first exon (bold) and the first
35
intron (underlined, SEQ ID NO: 39), followed by the N-terminal portion of the second exon, of the EF1a gene. In some embodiments, the polynucleotide provided herein comprises a short EF1a

promoter. In some embodiments, the polynucleotide provided herein comprises an EF1a promoter that is shorter than the nucleic acid sequence of SEQ ID NO:15. In some embodiments, the polynucleotide provided herein comprises an EF1a promoter that does not comprise the first intron of the EF1a gene. In some embodiments, the polynucleotide provided herein comprises an EF1a promoter that does not comprise the nucleic acid sequence of SEQ ID NO:39.

[0126] In some embodiments, the promoter comprises the sequence provided as SEQ ID NO:16.

GCGTGAGGCTCCGGTGCCCGTCAGTGGGCAGAGCGCACATCGCCCACAGTCCCCGAGA
 AGTTGGGGGGAGGGGTCGGCAATTGAACCGGTGCCTAGAGAAGGTGGCGCGGGGTAA
 10 ACTGGGAAAGTGATGTCGTGTACTGGCTCCGCCTTTTTCCCGAGGGTGGGGGAGAACCG
 TATATAAGTGCAGTAGTCGCCGTGAACGTTCTTTTTTCGCAACGGGTTTGCCGCCAGAAC
 ACAG (SEQ ID NO: 16)

[0127] Prior to the *in vitro* manipulation or genetic modification of the immune cells described herein, the cells may be obtained from a subject. The cells expressing a CD19 CAR may be derived from an allogenic or autologous process.

Source Material

[0128] In some embodiments, the immune cells comprise T cells. T cells can be obtained from a number of sources, including peripheral blood mononuclear cells (PBMCs), bone marrow, lymph nodes tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In certain embodiments, T cells can be obtained from a unit of blood collected from the subject using any number of techniques known to the skilled person, such as FICOLL™ separation.

[0129] Cells may be obtained from the circulating blood of an individual by apheresis. The apheresis product typically contains lymphocytes, including T cells, monocytes, granulocytes, B cells, other nucleated white blood cells, red blood cells, and platelets. In certain embodiments, the cells collected by apheresis may be washed to remove the plasma fraction, and placed in an appropriate buffer or media for subsequent processing.

[0130] In certain embodiments, T cells are isolated from PBMCs by lysing the red blood cells and depleting the monocytes, for example, using centrifugation through a PERCOLL™ gradient. A specific subpopulation of T cells, (e.g., CD28+, CD4+, CD5+, CD45RA-, and CD45RO+ T cells or CD28+, CD4+, CD5+, CD45RA-, CD45RO+, and CD62L+ T cells) can be further isolated by positive or negative selection techniques known in the art. For example, enrichment of a T cell population by negative selection can be accomplished with a combination of antibodies directed to surface markers unique to the negatively selected cells. One method for use herein is cell sorting and/or selection via negative magnetic immunoadherence or flow cytometry that uses a cocktail of

monoclonal antibodies directed to cell surface markers present on the cells negatively selected. For example, to enrich for CD4+ cells by negative selection, a monoclonal antibody cocktail typically includes antibodies to CD14, CD20, CD11b, CD16, HLA-DR, and CD8. Flow cytometry and cell sorting may also be used to isolate cell populations of interest for use in the present disclosure.

5 [0131] PBMCs may be used directly for genetic modification with the immune cells (such as CARs or TCRs) using methods as described herein. In certain embodiments, after isolating the PBMCs, T lymphocytes can be further isolated and both cytotoxic and helper T lymphocytes can be sorted into naive, memory, and effector T cell subpopulations either before or after genetic modification and/or expansion.

10 [0132] In some embodiments, CD8+ cells are further sorted into naive, stem cell memory, central memory, and effector cells by identifying cell surface antigens that are associated with each of these types of CD8+ cells. In some embodiments, the expression of phenotypic markers of central memory T cells include CD45RO, CD62L, CCR7, CD28, CD3, and CD127 and are negative for granzyme B. In some embodiments, stem cell memory T cells are CD45RO-, CD62L+, CD8+ T
15 cells. In some embodiments, central memory T cells are CD45RO+, CD62L+, CD8+ T cells. In some embodiments, effector T cells are negative for CD62L, CCR7, CD28, and CD127, and positive for granzyme B and perforin. In certain embodiments, CD4+ T cells are further sorted into subpopulations. For example, CD4+ T helper cells can be sorted into naive, central memory, and effector cells by identifying cell populations that have cell surface antigens.

20

Stem cell derived immune cells

[0133] In some embodiments, the immune cells may be derived from embryonic stem (ES) or induced pluripotent stem (iPS) cells. Suitable HSCs, ES cells, iPS cells and other stem cells may be cultivated immortal cell lines or isolated directly from a patient. Various methods for isolating,
25 developing, and/or cultivating stem cells are known in the art and may be used to practice the present invention.

[0134] In some embodiments, the immune cell is an induced pluripotent stem cell (iPSC) derived from a reprogrammed T-cell. In some embodiments, the source material may be an induced pluripotent stem cell (iPSC) derived from a T cell or non-T cell. The source material may be an
30 embryonic stem cell. The source material may be a B cell, or any other cell from peripheral blood mononuclear cell isolates, hematopoietic progenitor, hematopoietic stem cell, mesenchymal stem cell, adipose stem cell, or any other somatic cell type.

Genetic Modification of isolated cells

35 [0135] The immune cells, such as T cells, can be genetically modified following isolation using known methods, or the immune cells can be activated and expanded (or differentiated in the case of

progenitors) *in vitro* prior to being genetically modified. In some embodiments, the isolated immune cells are genetically modified to reduce or eliminate expression of endogenous TCR α and/or CD52. In some embodiments, the cells are genetically modified using gene editing technology (e.g., CRISPR/Cas9, a zinc finger nuclease (ZFN), a TALEN, a MegaTAL, a meganuclease) to reduce or eliminate expression of endogenous proteins (e.g., TCR α and/or CD52). In another embodiment, the immune cells, such as T cells, are genetically modified with the chimeric antigen receptors described herein (e.g., transduced with a viral vector comprising one or more nucleotide sequences encoding a CAR) and then are activated and/or expanded *in vitro*. Methods for activating and expanding T cells are known in the art and are described, for example, in U.S. Pat. No. 6,905,874; U.S. Pat. No. 6,867,041; U.S. Pat. No. 6,797,514; and PCT WO2012/079000, the contents of which are hereby incorporated by reference in their entirety. Generally, such methods include contacting PBMC or isolated T cells with a stimulatory molecule and a costimulatory molecule, such as anti-CD3 and anti-CD28 antibodies, generally attached to a bead or other surface, in a culture medium with appropriate cytokines, such as IL-2. Anti-CD3 and anti-CD28 antibodies attached to the same bead serve as a “surrogate” antigen presenting cell (APC). One example is The Dynabeads® system, a CD3/CD28 activator/stimulator system for physiological activation of human T cells. In other embodiments, the T cells may be activated and stimulated to proliferate with feeder cells and appropriate antibodies and cytokines using methods such as those described in U.S. Pat. No. 6,040,177; U.S. Pat. No. 5,827,642; and WO2012129514, the contents of which are hereby incorporated by reference in their entirety.

[0136] Certain methods for making the constructs and engineered immune cells of the disclosure are described in PCT application PCT/US15/14520, the contents of which are hereby incorporated by reference in their entirety.

[0137] It will be appreciated that PBMCs can further include other cytotoxic lymphocytes such as NK cells or NKT cells. An expression vector carrying the coding sequence of a chimeric receptor as disclosed herein can be introduced into a population of human donor T cells, NK cells or NKT cells. Successfully transduced T cells that carry the expression vector can be sorted using flow cytometry to isolate CD3 positive T cells and then further propagated to increase the number of these CAR expressing T cells in addition to cell activation using anti-CD3 antibodies and IL-2 or other methods known in the art as described elsewhere herein. Standard procedures are used for cryopreservation of T cells expressing the CAR for storage and/or preparation for use in a human subject. In one embodiment, the *in vitro* transduction, culture and/or expansion of T cells are performed in the absence of non-human animal derived products such as fetal calf serum and fetal bovine serum.

[0138] For cloning of polynucleotides, the vector may be introduced into a host cell (an isolated host cell) to allow replication of the vector itself and thereby amplify the copies of the

polynucleotide contained therein. The cloning vectors may contain sequence components generally include, without limitation, an origin of replication, promoter sequences, transcription initiation sequences, enhancer sequences, and selectable markers. These elements may be selected as appropriate by a person of ordinary skill in the art. For example, the origin of replication may be selected to promote autonomous replication of the vector in the host cell.

[0139] In certain embodiments, the present disclosure provides isolated host cells containing the vector provided herein. The host cells containing the vector may be useful in expression or cloning of the polynucleotide contained in the vector. Suitable host cells can include, without limitation, prokaryotic cells, fungal cells, yeast cells, or higher eukaryotic cells such as mammalian cells. Suitable prokaryotic cells for this purpose include, without limitation, eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacterales such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis*, *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*.

[0140] The vector can be introduced to the host cell using any suitable methods known in the art, including, without limitation, DEAE-dextran mediated delivery, calcium phosphate precipitate method, cationic lipids mediated delivery, liposome mediated transfection, electroporation, microprojectile bombardment, receptor-mediated gene delivery, delivery mediated by polylysine, histone, chitosan, and peptides. Standard methods for transfection and transformation of cells for expression of a vector of interest are well known in the art. In a further embodiment, a mixture of different expression vectors can be used in genetically modifying a donor population of immune effector cells wherein each vector encodes a different CAR as disclosed herein. The resulting transduced immune effector cells form a mixed population of engineered cells, with a proportion of the engineered cells expressing more than one different CARs.

[0141]

[0142] In some embodiments, the vector comprises a lentiviral vector. The lentiviral vector comprising a CAR coding sequence can be introduced into a lentiviral packaging cell line and the lentivirus produced by the packaging cell line can be used for transduction of T cells to generate CAR-T cells. To make lentivirus encoding CARs, HEK-293T cells can be plated at 0.4 million cells per mL in 2mL of DMEM (Gibco) supplemented with 10% FBS (Hyclone or JR Scientific) per well of a 6-well plate on Day 0. On Day 1, the lentivirus can be prepared by mixing together lentiviral packaging vectors 1.5ug psPAX2, 0.5ug pMD2G, and 0.5ug of the appropriate transfer CAR vector in 250uL Opti-MEM (Gibco) per well of the 6-well plate ("DNA mix"). 10uL Lipofectamine 2000 (Invitrogen) in 250uL Opti-MEM can be incubated at room temperature for 5 minutes and then added to the DNA mix. The mixture can be incubated at room temperature for 20 minutes and the total volume of 500uL was slowly added to the sides of the wells containing HEK-293T. The

general methods of CAR-containing lentivirus production and transduction are generally known in the art, for example, see Milone et al., *Leukemia*, 2018, 32:1529-1541; Sanber et al., Construction of stable packaging cell lines for clinical lentiviral vector production, *Nature* 2015, DOI: 10.1038; Roddie et al., *Cytotherapy* 2019, 21:327-340, all of which are incorporate herein by reference in their entireties. In one embodiment, the disclosure provides a method of storing genetically engineered cells expressing CARs or TCRs which target a CD19 protein. This involves cryopreserving the immune cells such that the cells remain viable upon thawing. A fraction of the immune cells expressing the CARs can be cryopreserved by methods known in the art to provide a permanent source of such cells for the future treatment of patients afflicted with a malignancy.

When needed, the cryopreserved transformed immune cells can be thawed, grown and expanded for more such cells.

[0143] In some embodiments, the cells are formulated by first harvesting them from their culture medium, and then washing and concentrating the cells in a medium and container system suitable for administration (a “pharmaceutically acceptable” carrier) in a treatment-effective amount. Suitable infusion media can be any isotonic medium formulation, typically normal saline, Normosol™ R (Abbott) or Plasma-Lyte™ A (Baxter), but also 5% dextrose in water or Ringer's lactate can be utilized. The infusion medium can be supplemented with human serum albumin.

Allogeneic CAR T cells

[0144] The process for manufacturing allogeneic CAR T therapy, or AlloCARs™, involves harvesting healthy, selected, screened and tested T cells from healthy donors. Next, the T cells are engineered to express CARs, which recognize certain cell surface proteins (e.g., CD19) that are expressed in hematologic or solid tumors. Allogeneic T cells are gene edited to reduce the risk of graft versus host disease (GvHD) and to prevent allogeneic rejection. A T cell receptor gene (e.g., TCR α , TCR β) is knocked out to avoid GvHD. The CD52 gene can be knocked out to render the CAR T product resistant to anti-CD52 antibody treatment. Anti-CD52 antibody treatment can therefore be used to suppress the host immune system and allow the CAR T to stay engrafted to achieve full therapeutic impact. The engineered T cells then undergo a purification step and are ultimately cryopreserved in vials for delivery to patients.

Autologous CAR T cells

[0145] Autologous chimeric antigen receptor (CAR) T cell therapy, involves collecting a patient's own cells (e.g., white blood cells, including T cells) and genetically engineering the T cells to express CARs that recognize target expressed on the cell surface of one or more specific cancer cells and kill cancer cells. The engineered cells are then cryopreserved and subsequently administered to the patient.

IV. Methods of Treatment

[0146] The disclosure comprises methods for treating or preventing a condition associated with undesired and/or elevated CD19 levels in a patient, comprising administering to a patient in need thereof an effective amount of at least one CAR, or immune-cell comprising a CAR disclosed
5 herein.

[0147] Methods are provided for treating diseases or disorders, including cancer. In some embodiments, the disclosure relates to creating a T cell-mediated immune response in a subject, comprising administering an effective amount of the engineered immune cells of the present application to the subject. In some embodiments, the T cell-mediated immune response is directed
10 against a target cell or cells. In some embodiments, the engineered immune cell comprises a chimeric antigen receptor (CAR). In some embodiments, the target cell is a tumor cell. In some aspects, the disclosure comprises a method for treating or preventing a malignancy, said method comprising administering to a subject in need thereof an effective amount of at least one isolated antigen binding domain described herein. In some aspects, the disclosure comprises a method for
15 treating or preventing a malignancy, said method comprising administering to a subject in need thereof an effective amount of at least one immune cell, wherein the immune cell comprises at least one chimeric antigen receptor, T cell receptor, and/or isolated antigen binding domain as described herein.

[0148] The CAR containing immune cells of the disclosure can be used to treat malignancies
20 involving aberrant expression of CD19. In some embodiments, CAR containing immune cells of the disclosure can be used to treat cancer. As used herein, the term "cancer" includes, but is not limited to, solid tumors and blood born tumors. The term "cancer" refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth,
25 neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor, malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal
30 adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi's sarcoma, karotype acute myeloblastic leukemia, Hodgkin lymphoma, non-Hodgkin lymphoma (NHL), cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, malignant melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal
35 carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans

progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation.

5 [0149] In exemplary embodiments, the CAR containing immune cells, e.g., CAR-T cells of the disclosure, are used to treat NHL.

10 [0150] Also provided are methods for reducing the size of a tumor in a subject, comprising administering to the subject an engineered cell of the present disclosure to the subject, wherein the cell comprises a chimeric antigen receptor comprising a CD19 antigen binding domain and binds to a CD19 antigen on the tumor.

[0151] In some embodiments, the subject has a solid tumor, or a blood malignancy such as lymphoma or leukemia. In some embodiments, the engineered cell is delivered to a tumor bed. In some embodiments, the cancer is present in the bone marrow of the subject. In some embodiments, the engineered cells are autologous immune cells, e.g., autologous T cells. In some embodiments, the engineered cells are allogeneic immune cells, e.g., allogeneic T cells. In some embodiments, the engineered cells are heterologous immune cells, e.g., heterologous T cells. In some embodiments, the engineered cells of the present application are transfected or transduced in vivo. In other 15 20 25 30 35
embodiments, the engineered cells are transfected or transduced ex vivo. As used herein, the term "in vitro cell" refers to any cell which is cultured ex vivo.

[0152] A "therapeutically effective amount," "effective dose," "effective amount," or "therapeutically effective dosage" of a therapeutic agent, e.g., engineered CART cells, is any amount that, when used alone or in combination with another therapeutic agent, protects a subject against the onset of a disease or promotes disease regression evidenced by a decrease in severity of disease symptoms, an increase in frequency and duration of disease symptom-free periods, or a prevention of impairment or disability due to the disease affliction. The ability of a therapeutic agent to promote disease regression can be evaluated using a variety of methods known to the skilled practitioner, such as in human subjects during clinical trials, in animal model systems predictive of efficacy in humans, or by assaying the activity of the agent in in vitro assays.

[0153] The terms "patient" and "subject" are used interchangeably and include human and non-human animal subjects as well as those with formally diagnosed disorders, those without formally recognized disorders, those receiving medical attention, those at risk of developing the disorders, etc.

[0154] The term “treat” and “treatment” includes therapeutic treatments, prophylactic treatments, and applications in which one reduces the risk that a subject will develop a disorder or other risk factor. Treatment does not require the complete curing of a disorder and encompasses embodiments in which one reduces symptoms or underlying risk factors. The term “prevent” does not require the 100% elimination of the possibility of an event. Rather, it denotes that the likelihood of the occurrence of the event has been reduced in the presence of the compound or method.

[0155] Desired treatment amounts of cells in the composition is generally at least 2 cells (for example, at least 1 CD8+ central memory T cell and at least 1 CD4+ helper T cell subset) or is more typically greater than 10^2 cells, and up to 10^6 , up to and including 10^8 or 10^9 cells and can be more than 10^{10} cells. The number of cells will depend upon the desired use for which the composition is intended, and the type of cells included therein. The density of the desired cells is typically greater than 10^6 cells/ml and generally is greater than 10^7 cells/ml, generally 10^8 cells/ml or greater. The clinically relevant number of immune cells can be apportioned into multiple infusions that cumulatively equal or exceed 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , or 10^{12} cells. In some aspects of the present disclosure, particularly since all the infused cells will be redirected to a particular target antigen (CD19), lower numbers of cells, in the range of 10^6 /kilogram (10^6 - 10^{11} per patient) may be administered. CAR treatments may be administered multiple times at dosages within these ranges. The cells may be autologous, allogeneic, or heterologous to the patient undergoing therapy.

[0156] In some embodiments, the therapeutically effective amount of the CAR T cells is about 1×10^5 cells/kg, about 2×10^5 cells/kg, about 3×10^5 cells/kg, about 4×10^5 cells/kg, about 5×10^5 cells/kg, about 6×10^5 cells/kg, about 7×10^5 cells/kg, about 8×10^5 cells/kg, about 9×10^5 cells/kg, 2×10^6 cells/kg, about 3×10^6 cells/kg, about 4×10^6 cells/kg, about 5×10^6 cells/kg, about 6×10^6 cells/kg, about 7×10^6 cells/kg, about 8×10^6 cells/kg, about 9×10^6 cells/kg, about 1×10^7 cells/kg, about 2×10^7 cells/kg, about 3×10^7 cells/kg, about 4×10^7 cells/kg, about 5×10^7 cells/kg, about 6×10^7 cells/kg, about 7×10^7 cells/kg, about 8×10^7 cells/kg, or about 9×10^7 cells/kg.

[0157] In some embodiments, target doses for CAR+/CAR-T+/TCR+ cells range from 1×10^6 - 2×10^8 cells/kg, for example 2×10^6 cells/kg. It will be appreciated that doses above and below this range may be appropriate for certain subjects, and appropriate dose levels can be determined by the healthcare provider as needed. Additionally, multiple doses of cells can be provided in accordance with the disclosure.

[0158] In some aspects, the disclosure comprises a pharmaceutical composition comprising at least one antigen binding domain as described herein and a pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition further comprises an additional active agent.

[0159] The CAR expressing cell populations of the present disclosure may be administered either alone, or as a pharmaceutical composition in combination with diluents and/or with other

components such as IL-2 or other cytokines or cell populations. Pharmaceutical compositions of the present disclosure may comprise a CAR or TCR expressing cell population, such as T cells, as described herein, in combination with one or more pharmaceutically or physiologically acceptable carriers, diluents or excipients. Such compositions may comprise buffers such as neutral buffered saline, phosphate buffered saline and the like; carbohydrates such as glucose, mannose, sucrose or dextrans, mannitol; proteins; polypeptides or amino acids such as glycine; antioxidants; chelating agents such as EDTA or glutathione; adjuvants (e.g., aluminum hydroxide); and preservatives. Compositions of the present disclosure are preferably formulated for intravenous administration.

[0160] The pharmaceutical compositions (solutions, suspensions or the like), may include one or more of the following: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono- or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. An injectable pharmaceutical composition is preferably sterile.

[0161] In some embodiments, upon administration to a patient, engineered immune cells expressing at their cell surface any one of the CD19-specific CARs described herein may reduce, kill or lyse endogenous CD19-expressing cells of the patient. In one embodiment, a percentage reduction or lysis of CD19-expressing endogenous cells or cells of a cell line expressing CD19 by engineered immune cells expressing any one of the CD19-specific CARs described herein is at least about or greater than 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%. In one embodiment, a percentage reduction or lysis of CD19-expressing endogenous cells or cells of a cell line expressing CD19 by engineered immune cells expressing any one of the CD19-specific CARs described herein is about 5% to about 95%, about 10% to about 95%, about 10% to about 90%, about 10% to about 80%, about 10% to about 70%, about 10% to about 60%, about 10% to about 50%, about 10% to about 40%, about 20% to about 90%, about 20% to about 80%, about 20% to about 70%, about 20% to about 60%, about 20% to about 50%, about 25% to about 75%, or about 25% to about 60%. In one embodiment, the endogenous CD19-expressing cells are endogenous CD19-expressing bone marrow cells.

[0162] In one embodiment, the percent reduction or lysis of target cells, e.g., a cell line expressing CD19, by engineered immune cells expressing at their cell surface membrane a CD19-specific CAR of the disclosure can be measured using the assay disclosed herein.

[0163] The methods can further comprise administering one or more chemotherapeutic agent. In certain embodiments, the chemotherapeutic agent is a lymphodepleting (preconditioning) chemotherapeutic. For example, methods of conditioning a patient in need of a T cell therapy comprising administering to the patient specified beneficial doses of cyclophosphamide (between 5 200 mg/m²/day and 2000 mg/m²/day, about 100 mg/m²/day and about 2000 mg/m²/day; *e.g.*, about 100 mg/m²/day, about 200 mg/m²/day, about 300 mg/m²/day, about 400 mg/m²/day, about 500 mg/m²/day, about 600 mg/m²/day, about 700 mg/m²/day, about 800 mg/m²/day, about 900 mg/m²/day, about 1000 mg/m²/day, about 1500 mg/m²/day or about 2000 mg/m²/day) and specified doses of fludarabine (between 20 mg/m²/day and 900 mg/m²/day, between about 10 mg/m²/day and 10 10 900 mg/m²/day; *e.g.*, about 10 mg/m²/day, about 20 mg/m²/day, about 30 mg/m²/day, about 40 mg/m²/day, about 40 mg/m²/day, about 50 mg/m²/day, about 60 mg/m²/day, about 70 mg/m²/day, about 80 mg/m²/day, about 90 mg/m²/day, about 100 mg/m²/day, about 500 mg/m²/day or about 900 mg/m²/day). A preferred dose regimen involves treating a patient comprising administering daily to 15 the patient about 300 mg/m²/day of cyclophosphamide and about 30 mg/m²/day of fludarabine for three days prior to administration of a therapeutically effective amount of engineered T cells to the patient.

[0164] In some embodiments, lymphodepletion further comprises administration of a CD52 antibody. In some embodiments, the CD52 antibody is administered at a dose of about 13 mg/day IV.

[0165] In other embodiments, the antigen binding domain, transduced (or otherwise 20 engineered) cells and the chemotherapeutic agent are administered each in an amount effective to treat the disease or condition in the subject.

[0166] In certain embodiments, compositions comprising CAR-expressing immune effector cells disclosed herein may be administered in conjunction with any number of chemotherapeutic 25 agents. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN™); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, trietylenephosphoramidate, triethylenethiophosphoramidate and trimethylolomelamine resins; nitrogen mustards such as 30 chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carabycin, carminomycin, carzinophilin, chromomycins, 35 dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins,

peplomycin, potfiromycin, puromycin, quelamycin, rodoxubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, 5
 enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitiostanol, mepitiothane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; 10
 diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea, lentinan; lonidamine; mitoguanzone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK®; razoxane; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2', 2''-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); 15
 cyclophosphamide; thiotepa; taxoids, e.g., paclitaxel (TAXOL™, Bristol-Myers Squibb) and doxetaxel (TAXOTERE®, Rhone-Poulenc Rorer); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; 20
 topoisomerase inhibitor RF S2000; difluoromethylomithine (DMFO); retinoic acid derivatives such as Targretin™ (bexarotene), Panretin™, (alitretinoin); ONTAK™ (denileukin diftitox); esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, 25
 aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Combinations of chemotherapeutic agents are also administered where appropriate, including, but not limited to CHOP, i.e., Cyclophosphamide (Cytoxan®), Doxorubicin 30
 (hydroxydoxorubicin), Vincristine (Oncovin®), and Prednisone.

[0167] In some embodiments, the chemotherapeutic agent is administered at the same time or within one week after the administration of the engineered cell, polypeptide, or nucleic acid. In other embodiments, the chemotherapeutic agent is administered from 1 to 4 weeks or from 1 week to 1 month, 1 week to 2 months, 1 week to 3 months, 1 week to 6 months, 1 week to 9 months, or 1 week 35
 to 12 months after the administration of the engineered cell, polypeptide, or nucleic acid. In other embodiments, the chemotherapeutic agent is administered at least 1 month before administering the

cell, polypeptide, or nucleic acid. In some embodiments, the methods further comprise administering two or more chemotherapeutic agents.

[0168] A variety of additional therapeutic agents may be used in conjunction with the compositions described herein. For example, potentially useful additional therapeutic agents include PD-1 inhibitors such as nivolumab (Opdivo®), pembrolizumab (Keytruda®), pembrolizumab, pidilizumab, and atezolizumab.

[0169] Additional therapeutic agents suitable for use in combination with the disclosure include, but are not limited to, ibrutinib (Imbruvica®), ofatumumab (Arzerra®), rituximab (Rituxan®), bevacizumab (Avastin®), trastuzumab (Herceptin®), trastuzumab emtansine (KADCYLA®), imatinib (Gleevec®), cetuximab (Erbix®), panitumumab (Vectibix®), catumaxomab, ibritumomab, ofatumumab, tositumomab, brentuximab, alemtuzumab, gemtuzumab, erlotinib, gefitinib, vandetanib, afatinib, lapatinib, neratinib, axitinib, masitinib, pazopanib, sunitinib, sorafenib, toceranib, lestaurtinib, axitinib, cediranib, lenvatinib, nintedanib, pazopanib, regorafenib, semaxanib, sorafenib, sunitinib, tivozanib, toceranib, vandetanib, entrectinib, cabozantinib, imatinib, dasatinib, nilotinib, ponatinib, radotinib, bosutinib, lestaurtinib, ruxolitinib, pacritinib, cobimetinib, selumetinib, trametinib, binimetinib, alectinib, ceritinib, crizotinib, aflibercept, adipotide, denileukin diftitox, mTOR inhibitors such as Everolimus and Temsirolimus, hedgehog inhibitors such as sonidegib and vismodegib, CDK inhibitors such as CDK inhibitor (palbociclib).

[0170] In some embodiments, the composition comprising CAR-containing immune cells may be administered with a therapeutic regimen to prevent cytokine release syndrome (CRS) or neurotoxicity. The therapeutic regimen to prevent cytokine release syndrome (CRS) or neurotoxicity may include lenzilumab, tocilizumab, atrial natriuretic peptide (ANP), anakinra, iNOS inhibitors (e.g., L-NIL or 1400W). In additional embodiments, the composition comprising CAR-containing immune cells can be administered with an anti-inflammatory agent. Anti-inflammatory agents or drugs include, but are not limited to, steroids and glucocorticoids (including betamethasone, budesonide, dexamethasone, hydrocortisone acetate, hydrocortisone, hydrocortisone, methylprednisolone, prednisolone, prednisone, triamcinolone), nonsteroidal anti-inflammatory drugs (NSAIDs) including aspirin, ibuprofen, naproxen, methotrexate, sulfasalazine, leflunomide, anti-TNF medications, cyclophosphamide and mycophenolate. Exemplary NSAIDs include ibuprofen, naproxen, naproxen sodium, Cox-2 inhibitors, and sialylates. Exemplary analgesics include acetaminophen, oxycodone, tramadol or propoxyphene hydrochloride. Exemplary glucocorticoids include cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisolone, or prednisone. Exemplary biological response modifiers include molecules directed against cell surface markers (e.g., CD4, CD5, etc.), cytokine inhibitors, such as the TNF antagonists, (e.g., etanercept (ENBREL®), adalimumab (HUMIRA®) and infliximab (REMICADE®), chemokine inhibitors and

adhesion molecule inhibitors. The biological response modifiers include monoclonal antibodies as well as recombinant forms of molecules. Exemplary DMARDs include azathioprine, cyclophosphamide, cyclosporine, methotrexate, penicillamine, leflunomide, sulfasalazine, hydroxychloroquine, Gold (oral (auranofin) and intramuscular) and minocycline.

5 [0171] In certain embodiments, the compositions described herein are administered in conjunction with a cytokine. Examples of cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormones such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor (HGF); fibroblast growth factor (FGF); prolactin; placental lactogen; mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors (NGFs) such as NGF-beta; platelet-growth factor; transforming growth factors (TGFs) such as TGF-alpha and TGF-beta; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon-alpha, beta, and -gamma; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-1alpha, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12; IL-15, IL-21 a tumor necrosis factor such as TNF-alpha or TNF-beta; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture, and biologically active equivalents of the native sequence cytokines.

V. Methods of Sorting and Depletion

25 [0172] In some embodiments, provided are methods for *in vitro* sorting of a population of immune cells, wherein a subset of the population of immune cells comprises engineered immune cells expressing any one of the CD19-specific CARs comprising epitopes specific for monoclonal antibodies (e.g., exemplary mimotope sequences). The method comprises contacting the population of immune cells with a monoclonal antibody specific for the epitopes and selecting the immune cells that bind to the monoclonal antibody to obtain a population of cells enriched in engineered immune cells expressing the CD19-specific CAR.

30 [0173] In some embodiments, said monoclonal antibody specific for said epitope is optionally conjugated to a fluorophore. In this embodiment, the step of selecting the cells that bind to the monoclonal antibody can be done by Fluorescence Activated Cell Sorting (FACS).

35 [0174] In some embodiments, said monoclonal antibody specific for said epitope is optionally conjugated to a magnetic particle. In this embodiment, the step of selecting the cells that bind to the monoclonal antibody can be done by Magnetic Activated Cell Sorting (MACS).

[0175] In some embodiments, the mAb used in the method for sorting immune cells expressing the CAR is chosen from alemtuzumab, ibritumomab tiuxetan, muromonab-CD3, tositumomab, abciximab, basiliximab, brentuximab vedotin, cetuximab, infliximab, rituximab, bevacizumab, certolizumab pegol, daclizumab, eculizumab, efalizumab, gemtuzumab, natalizumab, omalizumab, palivizumab, ranibizumab, tocilizumab, trastuzumab, vedolizumab, adalimumab, belimumab, canakinumab, denosumab, golimumab, ipilimumab, ofatumumab, panitumumab, QBEND-10 and/or ustekinumab. In some embodiments, said mAb is rituximab. In another embodiment, said mAb is QBEND-10.

[0176] In some embodiments, the population CAR-expressing immune cells obtained when using the method for *in vitro* sorting CAR-expressing immune cells described above, comprises at least 70%, 75%, 80%, 85%, 90%, 95% of CAR-expressing immune cells. In some embodiments, the population of CAR-expressing immune cells obtained when using the method for *in vitro* sorting CAR-expressing immune cells, comprises at least 85% CAR-expressing immune cells.

[0177] In some embodiments, the population of CAR-expressing immune cells obtained when using the method for *in vitro* sorting CAR-expressing immune cells described above shows increased cytotoxic activity *in vitro* compared with the initial (non-sorted) cell population. In some embodiments, said cytotoxic activity *in vitro* is increased by 10%, 20%, 30% or 50%. In some embodiments, the immune cells are T-cells.

[0178] In some embodiments, the mAbs are previously bound onto a support or surface. Non-limiting examples of solid support may include a bead, agarose bead, a magnetic bead, a plastic well plate, a glass well plate, a ceramic well plate, a column, or a cell culture bag.

[0179] The CAR-expressing immune cells to be administered to the recipient may be enriched *in vitro* from the source population. Methods of expanding source populations may include selecting cells that express an antigen such as CD34 antigen, using combinations of density centrifugation, immuno-magnetic bead purification, affinity chromatography, and fluorescent activated cell sorting.

[0180] Flow cytometry may be used to quantify specific cell types within a population of cells. In general, flow cytometry is a method for quantitating components or structural features of cells primarily by optical means. Since different cell types can be distinguished by quantitating structural features, flow cytometry and cell sorting can be used to count and sort cells of different phenotypes in a mixture.

[0181] A flow cytometry analysis involves two primary steps: 1) labeling selected cell types with one or more labeled markers, and 2) determining the number of labeled cells relative to the total number of cells in the population. In some embodiments, the method of labeling cell types includes binding labeled antibodies to markers expressed by the specific cell type. The antibodies may be either directly labeled with a fluorescent compound or indirectly labeled using, for example, a fluorescent-labeled second antibody which recognizes the first antibody.

[0182] In some embodiments, the method used for sorting T cells expressing CAR is the Magnetic- Activated Cell Sorting (MACS). Magnetic-activated cell sorting (MACS) is a method for separation of various cell populations depending on their surface antigens (CD molecules) by using superparamagnetic nanoparticles and columns. MACS may be used to obtain a pure cell population.

5 Cells in a single-cell suspension may be magnetically labeled with microbeads. The sample is applied to a column composed of ferromagnetic spheres, which are covered with a cell-friendly coating allowing fast and gentle separation of cells. The unlabeled cells pass through while the magnetically labeled cells are retained within the column. The flow-through can be collected as the unlabeled cell fraction. After a washing step, the column is removed from the separator, and the
10 magnetically labeled cells are eluted from the column.

[0183] Detailed protocol for the purification of specific cell population such as T-cell can be found in Basu S et al. (2010). (Basu S, Campbell HM, Dittel BN, Ray A. Purification of specific cell population by fluorescence activated cell sorting (FACS). J Vis Exp. (41): 1546).

[0184] In some aspects, the present disclosure provides a method for depleting CD19 specific
15 CAR-expressing immune cells by *in vivo* depletion. *in vivo* depletion may include the administration of a treatment (e.g., a molecule that binds an epitope on the CAR) to a mammalian organism aiming to stop the proliferation of the CAR-expressing immune cells by inhibition or elimination.

[0185] One aspect of the invention is related to a method for *in vivo* depleting an engineered immune cell expressing a CD19 CAR comprising a mAb specific epitope, comprising contacting
20 said engineered immune cell or said CAR-expressing immune cell with at least one epitope-specific mAb. Another aspect of the invention relates to a method for *in vivo* depleting CAR-expressing immune cell which comprises a chimeric scFv (e.g., formed by insertion of a mAb-specific epitope) by contacting said engineered immune cell with epitope- specific antibodies. In some embodiments, the immune cells are T-cells and/or the antibodies are monoclonal.

25 [0186] According to one embodiment, the *in vivo* depletion of the immune engineered cells is performed on engineered immune cells which has been previously sorted using the *in vitro* method of the present invention. In this case, the same infused mAb may be used. In some embodiments, the mAb-specific antigen is CD20 antigen and the epitope- specific mAb is rituximab. In some
30 embodiments, the invention relates to a method for *in vivo* depleting an engineered immune cell expressing a CAR comprising an mAb-specific epitope (CAR-expressing immune cell) in a patient comprising contacting said CAR-expressing immune cell with at least one epitope-specific mAb

[0187] In some embodiments, the step of contacting said engineered immune cell or said CAR-
35 expressing immune cell with at least one epitope-specific mAb comprises infusing the patient with epitope- specific mAb (e.g., rituximab). In some embodiments, the amount of epitope-specific mAb administered to the patient is sufficient to eliminate at least 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90% of the CAR-expressing immune cell in the patient.

[0188] In some embodiments, the step of contacting said engineered immune cell or said CAR-expressing immune cell with at least one epitope-specific mAb comprises infusing the patient with 375mg/m² of rituximab, once or several times. In some embodiments, the mAb (e.g., rituximab) is administered once weekly.

5 [0189] In some embodiments, when immune cells expressing a CAR comprising an mAb-specific epitope (CAR-expressing immune cells) are depleted in a complement dependent cytotoxicity (CDC) assay using epitope-specific mAb, the amount of viable CAR-expressing immune cells decreases. In some embodiments, the amount of viable CAR-expressing immune cells decreases by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80% or 90%. In some embodiments, 10 said mAb-specific epitope is a CD20 epitope or mimotope and/or the epitope-specific mAb is rituximab.

[0190] In certain embodiments, the *in vivo* depletion of CAR-engineered immune cells is performed by infusing bi-specific antibodies. By definition, a bispecific monoclonal antibody (BsAb) is an artificial protein that is composed of fragments of two different monoclonal antibodies 15 and consequently binds to two different types of antigen. These BsAbs and their use in immunotherapy have been reviewed in Muller D and Kontermann R.E. (2010) Bispecific Antibodies for Cancer Immunotherapy, *BioDrugs* 24 (2): 89-98.

[0191] According to another particular embodiment, the infused bi-specific mAb is able to bind 20 both the mAb-specific epitope borne on engineered immune cells expressing the chimeric scFv and to a surface antigen on an effector and cytotoxic cell (e.g., immune cells such as lymphocytes, macrophages, dendritic cells, natural killer cells (NK Cell), cytotoxic T lymphocytes (CTL)). By doing so, the depletion of engineered immune cells triggered by the BsAb may occur through antibody- dependent cellular cytotoxicity (ADCC). (Deo Y M, Sundarapandiyan K, Keler T, Wallace PK, and Graziano RF, (2000), *Journal of Immunology*, 165 (10):5954-5961).

25 [0192] In some embodiments, a cytotoxic drug is coupled to the epitope-specific mAbs which may be used to deplete CAR-expressing immune cells. By combining targeting capabilities of monoclonal antibodies with the cancer-killing ability of cytotoxic drugs, antibody-drug conjugate (ADC) allows a sensitive discrimination between healthy and diseased tissue when compared to the use of the drug alone. Market approvals were received for several ADCs; the technology for making 30 them -particularly on linkers- are described in (Payne, G. (2003) *Cancer Cell* 3:207-212; Trail et al (2003) *Cancer Immunol. Immunother.* 52:328-337; Syrigos and Epenetos (1999) *Anticancer Research* 19:605-614; Niculescu-Duvaz and Springer (1997) *Adv. Drug Del. Rev.* 26:151-172; U.S. Pat. No. 4,975,278).

[0193] In some embodiments, the epitope-specific mAb to be infused is conjugated beforehand 35 with a molecule able to promote complement dependent cytotoxicity (CDC). Therefore, the complement system helps or complements the ability of antibodies to clear pathogens from the

organism. When stimulated an activation cascade is triggered as a massive amplification of the response and activation of the cell-killing membrane attack complex. Different molecule may be used to conjugate the mAb, such as glycans [Courtois, A, Gac-Breton, S., Berthou, C, Guezennec, J., Bordron, A. and Boisset, C. (2012), Complement dependent cytotoxicity activity of therapeutic antibody fragments may be acquired by immunogenic glycan coupling, Electronic Journal of Biotechnology ISSN: 0717-3458; <http://www.ejbiotechnology.info> DOI: 10.2225/voll5-issue5).

VI. Kits and Articles of Manufacture

[0194] The present application provides kits comprising any one of the CD19 containing CARs or CD19 CAR containing immune cells described herein, and pharmaceutical compositions of the same. In some exemplary embodiments, a kit of the disclosure comprises allogeneic CD19 CAR-containing T-cells and a CD52 antibody for administering to the subject a lymphodepletion regimen and a CAR-T regimen.

[0195] The present application also provides articles of manufacture comprising any one of the therapeutic compositions or kits described herein. Examples of an article of manufacture include vials (e.g., sealed vials).

EXAMPLES

Example 1: Generation of rituximab-resistant CD19 CAR immune cells

[0196] Rituximab-resistant anti-CD19 chimeric antigen receptor constructs that do not express a rituximab binding site as shown in FIG. 1 and Table 4 were generated. The lentiviral vectors constructs were introduced into viral packaging cell line and anti-CD19 CAR containing lentiviruses were produced at Allogene.

[0197] Pan T cells from four human donors (541, 604, 410 and 2593) were thawed and activated at 1.5×10^6 cells/ml with Large-scale T Cell TransAct™ (1:15 ratio) in the presence of IL-2 (100IU/ml). After 2 days 1.5×10^6 cells (in 3ml) were transduced with 2ml of fresh lentivirus comprising the vectors described in Table 4. Schematics of the modified vectors are shown in Figure 1. IL-2 (100 IU/ml) was added on days 0, 2, 5, 7, 9 and 12. 6×10^6 total cells were transferred to a 6-well G-Rex plate on day 5 and media exchange performed on day 9 and 12. Cells were frozen on day 13.

Table 4: Rituximab-resistant CD19 CAR vectors

Vector:	Name:
anti-CD19 CAR v1.0	pCLS-m4G7 CAR (with the RQR8 safety switch)
anti-CD19 CAR v1.1	pCLS-4G7_CAR (same EF1a promoter sequence as in v1.0)
anti-CD19 CAR v1.2	pCLS-EF1a(short)-4G7(co)

anti-CD19 CAR v1.3	pCLS-(deltaRQR)8-4G7(co) (same EF1a promoter sequence as in v1.0)
anti-CD19 CAR v1.4	pCLS-1Q8-4G7(co) (same EF1a promoter sequence as in v1.0)
anti-CD19 CAR v1.5	pCLS-LQL8-4G7(co) (same EF1a promoter sequence as in v1.0)
anti-CD19 CAR v1.6	pCLS-Q38-4G7(co) (same EF1a promoter sequence as in v1.0)

co = codon optimized

[0198] Flow cytometry experiments were performed on transduced cells gated on lymphocytes, live CD3+, CAR+, CD4/CD8 and downstream markers. A human transduction check and CD34 panel was performed on Day 5 and Day 13 using a panel of CD3, CD4, CD8, viability, CD34 and an anti-idiotype to antibody to the anti-CD19 CAR (4G7 anti-Id). FIGs. 2A and 2B show flow cytometry plots demonstrating CAR expression on day 5 from Pan T cells transduced with the CAR expression vectors shown in Table 4 using the anti-CD19 CAR anti-Id antibody.

[0199] Flow cytometry with a human phenotype and activation panel was performed on day 9 and 13. The panel included CD3, CD4, CD8, viability, CD45RO, CD62L, CD25, 4-1BB, PD-1, an anti-idiotype to antibody to the anti-CD19 CAR, and TIM3. Cells were normalized for cell expansion and final CAR expression from all four donors on day 13 (FIG. 3). The data in FIG. 3 show that although v1.2 exhibited higher transduction rate (%CAR+), v1.2 transduced cells exhibited lower levels of CAR expression (CAR MFI), as compared to, e.g., v1.0 and v1.1. Cell expansion and CAR expression was monitored over time of the Pan T cells from donor 541 (FIG. 4A), 604 (FIG. 4B), 410 (FIG. 4C), 2593 (FIG. 4D) transduced with rituximab resistant CAR expression vectors.

[0200] CD4/CD8 ratios were measured on day 5, 9, and 13 of Pan T cells from donor 541 (FIG. 5A), 604 (FIG. 5B), 410 (FIG. 5C), 2593 (FIG. 5D) transduced with rituximab resistant CAR expression vectors. FIGs. 6A-6D show phenotype and activation on day 9 of Pan T cells from donor 541 (FIG. 6A), 604 (FIG. 6B), 410 (FIG. 6C), 2593 (FIG. 6D) transduced with rituximab resistant CAR expression vectors. FIG. 7 shows phenotype, activation %CD8+, and T cell anergy measured using TIM3 and PD1 staining averaged from all four donors on day 9. Phenotype and activation were measured on day 13 of Pan T cells from donor 541 (FIG. 8A), 604 (FIG. 8B), 410 (FIG. 8C), 2593 (FIG. 8D) transduced with rituximab resistant CAR expression vectors. FIG. 9 shows phenotype, activation %CD8+, and T cell anergy measured using TIM3 and PD1 staining from all four donors on day 13.

Example 2: Short- and long-term in vitro killing assays

[0201] Transduced CAR cells from Example 1 were tested for short- and long-term killing capabilities. CAR T cell co-cultures with Raji cells (2:1 E-to-T) were prepared for posterior a Luminex assay. Average short-term (24hr) killing assays using Raji cells as target cells was determined for each CAR construct (FIG. 10). FIGs. 11A-11D shows average long-term killing

assays using A549-CD19+ cells as target cells with an E:T of 8:1 (FIG. 11A), 4:1 (FIG. 11B), 2:1 (FIG. 11C), and 1:1 (FIG. 11D) for each CAR construct.

[0202] Killing assay results were analyzed for correlation with phenotypic characteristics. Day 7 percent killing at 1:1 E-to-T negatively correlates with CAR+CD4+41BB+, CAR+CD4+Tim3+ (p=0.0352), CAR+CD4+T_{EM}+ (p=0.0328), CAR+CD8+PD-1+ (p=0.0269) and % CAR expression on day 13 (p=0.0245). Day 7 percent killing at 1:1 E-to-T positively correlates with CAR+CD8+T_{SCM}+. Day 9 percent killing at 1:1 E-to-T negatively correlates with CAR+CD4+Tim3+, CAR+CD4+T_{EM}+ (p=0.0031), CAR+CD8+T_{cm}+ (p=0.0182) and %CAR expression day 13 (p=0.0469). Day 9 percent killing at 1:1 E-to-T positively correlates with CAR+CD8+T_{SCM}+, CAR+CD8+Tim3-PD-1- (p=0.0318), and CAR+CD4+T_{SCM}+ (p=0.0289).

Example 3: Analysis of titers of lentivirus containing different lentiviral constructs

[0203] In this experiment, the lentiviral vectors constructs were introduced into viral packaging cell line and anti-CD19 CAR containing lentiviruses were produced and titers determined at Lentigen (Gaithersburg, MD) under a similar protocol as in Example 1.

[0204] To lentiviral titers were assessed either by measuring the physical titers of the levels of viral protein p24 or by measuring the transducing titers. It was unexpectedly found that when the safety switch RQR8 was removed from the lentiviral construct v1.0, viral titer significantly decreased (compare v1.0 with v1.1 in Table 5). The titer was improved when the EF1a promoter in v1.1 was replaced with a short or truncated EF1a promoter as in v1.2 (“EF1a(short) promoter”).

Table 5 Viral Titers of lentivirus bearing rituximab-sensitive and -resistant CD19 CAR constructs

Construct	Physical titer p24	Transducing titer
anti-CD19 CAR v1.0	9293 ng/mL	1.9 x 10 ⁹ TU/mL ¹
anti-CD19 CAR v1.1	2464 ng/mL	5.4 x 10 ⁸ TU/mL
anti-CD19 CAR v1.2	8208 ng/mL	7 x 10 ⁹ TU/mL
anti-CD19 CAR v1.3	4494 ng/mL	7 x 10 ⁸ TU/mL
anti-CD19 CAR v1.4	9686 ng/mL	2.4 x 10 ⁹ TU/mL

¹TU = Transducing Unit

25

[0205] To analyze the robustness of the lentiviral preparations of anti-CD19 CAR v1.0, v1.2 and v1.3, a viral titration assay was performed. Serial volumetric dilution of lentiviral preparations of v1.0, v1.2 and v1.3 was analyzed for % CAR+ T cells on Day 5 after transduction of pan T cells. The results show that at low dilution (e.g., 10% v/v), all three constructs exhibited similar acceptable transduction efficiency. At increasing dilutions (e.g., 3.3%, 1.1% v/v), however, the transduction

30

efficiency of rituximab-resistant anti-CD19 CAR construct v1.3 dropped more significantly as compared to the other rituximab-resistant anti-CD19 CAR construct v1.2. See FIG. 12. Construct v1.2 was selected for *in vivo* analysis.

5 **Example 4: *In vivo* potency assay**

[0206] In this experiment, *in vivo* anti-tumor potency of ALLO-501v1.2 was analyzed as compared to ALLO- 501v1.0 in a mice tumor model. CD19 positive Raji cells that carry a luciferase reporter gene were injected into NSG mice. Lentivirus containing the v1.0 or v1.2 lentiviral construct was transduced to pan T cells of two donors 541 and 604. NSG mice were inoculated via
10 tail vein injection with 100,000 luciferase Raji cells. On day 4 post-inoculation Raji-bearing NSG mice were administered the CAR construct at the indicated doses. Raji engraftment and progression was evaluated by i.p. injection of luciferase substrate, followed by measurement of cumulative luciferase signals. Results are shown in FIG. 13A (donor 541) and FIG. 13B (donor 604).

[0207] Although the disclosed teachings have been described with reference to various
15 applications, methods, kits, and compositions, it will be appreciated that various changes and modifications can be made without departing from the teachings herein and the claimed invention below. The foregoing examples are provided to better illustrate the disclosed teachings and are not intended to limit the scope of the teachings presented herein. While the present teachings have been described in terms of these exemplary embodiments, the skilled artisan will readily understand that
20 numerous variations and modifications of these exemplary embodiments are possible without undue experimentation. All such variations and modifications are within the scope of the current teachings.

25

SEQ ID NO. CHART

SEQ ID NO	Description	Sequence
1	ALLO-501_v1.0	atgctgaccagcctgctgtgctggatggcctgtgctgctggcgccgaccag ccgatgcctgccctacagcaaccccagcctgtgcagcggaggcggcggcagc gagtgcgccaccagggcaccttctccaacgtgtccaccaacgtgagcccagcc aagcccaccaccagcctgtccttattccaatcctcctgtgtagcggaggggg aggcagcccagccccagacctcccaccccagccccaccatcgccagccagc ctctgagcctgagaccgaggcctgcccccagccgcccggcgccgctgac accagaggcctggatttcgctgcgatactacatctgggccccactggccgca cctgtggcgtgctgctgctgagcctggtgatcacctgtactgcaaccaccgcaac cgcaggcgctgtgcaagtgcccaggcccgtggtgagagccgagggcagag gcagcctgctgacctggcgacgtggaggagaaccaggccccatggagacc gacacctgctgctgtgggtgctgctgctgtgggtgccaggcagcaccggcgag gtgcagctgcagcagcggaccgagctgatcaagccaggcgccagcgtgaa gatgagctgcaaggccagcggctacacctaccagctacgtgatgactgggtg aagcagaagccaggccaggcctggagtggatcggtacatcaaccctacaac gacggcaccaagtacaacgagaagttcaaggcaaggccaccctgaccagcga caagagcagcagcaccgctacatggagctgagcagcctgaccagcgaggaca gcgccgtgactactgcgccagaggcacctactactacggcagccgggtgtcga ctactggggccaggggcaccacctgaccgtgagctctggcggaggcggctctgg cggaggcggctctggcggaggcggcagcgcacatcgtgatgaccaggctgccc ccagcatccccgtgaccaggcgagagcgtgagcatcagctgccggagcagc aagagcctgctgaacagcaacggcaacacctactgtactgttctgcagcggc caggccagagccccagctgctgatctaccggatgagcaacctggccagcggc gtgcccgaccggtcagcggcagcggcagcggcaccgcttcacctgcccgatc agccgggtggaggccgaggacgtggcgtgtactactgcatgcagcacctgga gtaccttcacctcggagccggcaccgaagctggagctgaagcggctggatccc accaccaccccagccccagggccacctaccctgcccccaaccatgccagcca gccctgagcctgcggcctgaagcctgcaggcctgccgcccggaggagccgtgc acacaaggggctggaactgcctgcgacatctatctgggccccctggccgg gacatgcggggtgctgctgtcctggtgattacactgattgcaaacggggcc ggaagaagctgctgtacatctcaagcagccctcatgcggcccgtgcagaccac ccaggaggaggacggctgcagctgccggtcccagaggaagagggaaggcggc tgcgagctgcgggtgaagttcagccggagcggcagccccagcctaccagca gggccagaaccagctgtacaacgagctgaacctgggacggcgggaggagtac gacgtgctggacaagcggcggggacgggaccccagatgggcccgaagcctc gccggaagaatcccaggagggcctgtacaacgagctgcagaaggacaagatg gccgagcctacagcgatcggcatgaagggcgagcggcggcggggcaag ggccacgacggcctgtaccaggcctgagcaccgccaccaaggacacctacga cgccctgcacatgcaggcctgccaccccggtga
2	ALLO-501_v1.1	atggagaccgacacctgctgctgtgggtgctgctgctggtgccaggcagca ccggcgagggtgcagctgcagcagagcggaccggagctgatcaagccaggcgc cagcgtgaagatgagctgcaaggccagcggctacacctaccagctacgtgatg cactgggtgaagcagaagccaggccaggcctggagtggatcggtacatcaa cccctacaacgacggcaccgaagttcaagggcaaggccacct

		<p>gaccagcgacaagagcagcagcaccgcctacatggagctgagcagcctgacca gcgaggacagcgccgtgtaactactgcgccagagggcacctactactacggcagcc ggglgttcgactactggggccagggcaccacccctgaccgtgagctctggcggag gcggctctggcggagggcggctctggcggagggcggcagcgcacatcgtgatgacc caggctgccccagcatccccgtgaccccagggcagagcgtgagcatcagctg ccggagcagcaagagcctgctgaacagcaacggcaacacctacgtactggtt cctgcagcggccagggcagagccccagctgctgatctacgggatgagcaacct ggccagcggcgtgcccaccgggtcagcggcagcggcagcggcaccgccttca ccctgggatcagccgggtggagggcaggagcgtggcgctgtaactactgcatgc agcacctggagtacccttcaccttcggagccggcaccaagctggagctgaagc ggtcggatcccaccaccaccccagccccacggccacctaccccctgccccacca tcgccagccagccccctgagcctgaggcctgaagcctgaggcctgccggcga ggagccgtgcacacaaggggctggacttcgctgcgacatctatactggggccc ccctggccgggacatgcgggggtgctgctgctgctcctgggtgattacactgtattgc aaacggggccggaagaagctgctgtacatctcaagcagccctcatgcggccccg tgcagaccaccagggagggagcggctgcagctgccgggtccccgaggaagag gaaggcggctgcgagctgcgggtgaagttcagccggagcggcagcccag cctaccagcagggccagaaccagctgtacaacgagctgaacctgggacggcgg gaggagtacgacgtgctggacaagcggcgggggacgggacccccagatgggcg gcaagcctgccggaagaatcccagggggcctgtacaacgagctgcagaag gacaagatggccgagcctacagcgagatcggcatgaagggcagcggcgcc ggggcaagggccacgcagcggcctgtaccagggcctgagcaccgccaccaagga cacctacgacgcctgcacatgcaggccctgccaccgggtga</p>
3	ALLO-501_v1.2	<p>atggagacagataccctgctgctgtgggtgctgctgctgtgggtgcctggtccac aggagagggtgcagctgcagcagctggaccagagctgatcaagcctggagcatc cgtgaagatgtcttgcaaggccagcggctatacattcaccagctacgtgatgact gggtgaagcagaagcctggccagggcctggagtggatcggctatatcaatcata caacgacggcaccaagtataatgagaagtttaagggcaaggccacactgacctct gataagagctcctctacagcctacatggagctgagctccctgacctctgaggacag cgccgtgtaactattgcccagagggcacatactattacggcagcagggtgttcgatt actggggccagggcaccacactgaccgtgtctagcggaggaggaggctccgga ggaggaggctctggcggcggcggcagcgcacatcgtgatgacacaggcagcac caagcatcccagtgaccctggcagagcgtgtccatctctgtcggctccttaagt ccctgctgaactctaattgcaaacctatctgtactggttctgcagcggccccggac agtccccacagctgctgatctataggatgagcaacctggcatccggagtgcctgat cgctcagcggctccggctctggaacagcctttacctgaggatctctcggtgga ggcagaggacgtgggcgtgtattactgcatgcagcacctggagtacccctcacat ttggcgcaggaaccaagctggagctgaagcggagcgaacccaccacaaccct gcaccacggccccctacaccagcacctaccatcgcactcagccactgagcctgc ggcccagggcctgtaggcctgcagcaggaggagcagtgacaccaggggct ggacttcgctgcatatctatactgggcaccactggcaggaacatgtggcgtgc tgctgctgagcctggtcatcaccctgtattgcaagagaggcaggaagaagctgct glacatctcaagcagcctttatgcggccagtgacagacaaccaggaggaggat ggctgctcctgtagattccagaggaggaggaggatgtgagctgcgctg aagtttagccggtccgcccagcaccagcatabcagcagggccagaatcagctgt acaatgagctgaacctggccggagagaggagtacgacgtgctggataagagg aggggaagggaccccagatgggagggcaagccacggagaaagaatcccag gagggcctgtataacgagctgcagaaggataagatggccgaggcctacagcga gatcggcatgaagggagagaggcggcggggcaagggacacgacggcctgtat</p>

		cagggcctgtccacagccaccaaggacacctacgatgccttgacatgcaggcc ctgccaccaaggtga
4	ALLO-501_v1.3	atgggaacaagcctgtgtgctggatggctctgtgctgctggggccgaccacg ctgacgcctccggggggggggctctcctgcccctaggccccctacactgcac caaccatcgcacccagccactgtctctgcgccttgaggcctgcccggcagcagc aggaggagcagtgcacacccgcggcctggacttcgctgcgatctatctatctgg gcaccactggcaggcacatgtggcgtgctgctgctgagcctggatcacacctgta ctgcaatcacaggaaccggagaagggtgtgcaagtgtccccggcctgtggtgag agcagaggggaaggggcagcctgtgacatgtggcgacgtggaggagaatccag gccctatggagacagataccctgctgctgtgggtgctgctgctgtgggtgcccgg cagcaccggagaggtgcagctgcagcagtccggaccagagctgatcaagcctg gagccagcgtgaagatgtcctgtaaggcctctggctatactaccagctacgtg atgcactgggtgaagcagaagcctggccagggcctggagtggatcggctatact aatccatacaacgacggcacaagataacgagaagttaagggaaggccaca ctgacctccgataagagctcctctacagcctacatggagctgagctccctgacctt gaggacagcgcctgtactattgcgcagaggcacatactattacgctctagggf glitcattactggggccagggcaccacactgaccgtctagcggaggaggagg cagcggaggaggaggctccggcggcggcggcctgacatcgtgatgacacagg cagcaccatccatcccagtgacccctggcgagagcgtgtccatctctgtcgtcc tctaagagcctgctgaactcaatggcaacacctatctgactggfctctgcagcgg cccggacagagcccacagctgctgatctataggatgtctaactggcaagcggcg tgcccgatcgttcagcggctccggctctggcacagccttaccctgaggatctcc cgcgtggaggcagaggacgtggcgtgtattactgcatgcagcactggagtac cccttcacatttggcgcagggaccaagctggagctgaagcggagcgcaccccacc acaaccctgcaccacggccacccacaccagcactactattgcatcccagccac tgagcctgcggcccaggcctgtaggcctgcccggcggcggcagtgacacacc cggggcctggactttgctgcgatatctacatctgggcacctctggccggcacatg cggcgtgctgtactgagcctggatcacacctgtattgcaagcggggcagaaag aagctgctgtacatctcaagcagcctttatgcccggcagtgacagacaaccagga ggaggatggctgctcctgtagattcccagaggaggaggaggaggatgtgagct gcgcgtgaagtttagcggctccggcgcagcaccagcatatcagcagggccagaa ccagctgtacaatgagctgaacctgggcccggagagaggatgactgtctgga taagagacggggccgggaccccagatgggaggcaagccacgcccggaaagaat ccccaggaggcctgtataacgagctgcagaaggataagatggccgaggcctac agcagatcggcatgaagggagagagaaggcggcaagggacacgacggc ctgtaccagggcctgagcacagcaacaaaagacacctacgacgcactgcacatg caggctctgccccctcggtaa
5	>ALLO-501_v1.4	atgggaacctctctgtgtgctggatggctctgtgctgctggggccgatcacgc tgacgcaagtgccgggggggggtccgaactgcccacacagggcaccttctcca acgtgagcaccacagtgagctccggcggaggaggcagccctgcaccaaggcc ccctacaccagcacctaccatcgcacatcagccactgagcctgcccggaggcc tgccggcctgcagcaggcggcgcctgacacaccgcggcctggactttgctgc gatctatctatctgggcacctctggcaggcacatgtggcgtgctgctgagcctg gtcatcacctgtactgcaatcacaggaaccggagaagggtgtgcaagtgtccac ggcccgtggtgagagcagaggggaaggggtccctgctgacatgtggcgacgtg gaggagaatcctggcccaatggagacagataccctgctgctgtgggtgctgctgc tgtgggtgcccggctccaccggagaggtgcagctgcagcagctgtggaccagagc tgatcaagccaggagcatccgtgaagatgtcttgaaggccagcggctatacttc

		<p>accagctacgtgatgactgggtgaagcagaagccaggacagggcctggagtg gatcggctatatcaatccttacaacgacggcaccaagtataacgagaagtttaagg gcaaggccaactgacctctgataagctlagctccacagcctacatggagctgct agcctgaccagcgaggactccgccgtgtactattgcccagagggcacatactatta cggcagcaggggtgttcgattactggggccagggcaccacactgacctgtctct ggaggaggaggctccggaggaggaggctctggcggcggcggcagcgacatc gtgatgacacaggcagcaccttccatcccagtgaccccaggcgagtctgtgagca tctctgtcggagctccaagtccctgctgaactctaattggcaacacctatctgtactg gtttctgcagcggcccggacagtccccacagtctgtgatctataggatgagcaatc tggcctccggcgtgccagatcgcttctctggcagcggctccggcacagcctttacc ctgaggatctctcgcgtggaggcagaggacgtgggctgtattactgcatgcagc acctggagtaccattcacatttggcgcaggcaccaagctggagctgaagcggga gacccccaccacaaccccagcacctcggccaccacaccagcaccaccatc gcatctcagcctctgagcctcggcccggagcctgtaggcccagcaggagg agcagtgacaccccggggcctggacttcgctgcgatatctacatctgggcacca ctggcccggcacatcgggcgtgctgttactgagcctggtcatcacctgtattgcaa gcggggcagaaagaagctgctgtacatctcaagcagcccttatgcggcctgtgc agacaaccaggaggaggatggctgctcctgtagattccctgaggaggaggagg gaggatgtgagctgcgctgaagtttctcggagcggccagcaccagcatatca gcagggacagaaccagctgtacaatgagctgaacctgggcccggagagaggagt atgacgtgctggataagagacggggccgggaccccagatgggaggcgaagcct cgccggaagaatccacaggaggcctgtataacgagctgcagaaggataagatg gcccaggcctacagcgagatcggcatgaaggagagagaaggcggcggcaag ggacacgacggcctgtaccaggcctgagcacagcaaaaagacacctacga cgactgcacatgcaggctctgccaccaagatga</p>
6	ALLO-501_v1.5	<p>atggggacctcactgctgtgctggatggctctgtgcctgctgggggcccaccacg ctgacgcctgctctggggggggggggggctcatgctccggaggaggaggctct gagctgccaaccaggggcacattctccaacgtgagcaccaacgtgtctcctgcca agccaaccacaaccgcatgcagcggcggaggaggagcagctgtccggcgg cggcggcagccctgccccaggccccctaccccagcacctacaatcgatctca gcctctgagcctgcgcccagaggcctgtcggcccgcagcaggaggagcagctgc acaccgcggcctggactttgcctgcgatatctatatctgggaccactggcaggc acctgtggcgtgctgctgctgagcctggatcacctgtactgcaatcacaggaa ccggagaagggtgtcaagtgtccacggcccgtggtgagagcagagggaagg ggctctctgctgacctgtggcagctggaggagaatcctggccctatggagacag atacactgctgctggtgctgctgctgtgggtgcccggcagcacaggagaggf gcagctgcagcagtcggacctgagctgatcaagccaggcgcctccgtgaagat gtcttgcaaggccagcggctataccttcacaagctacgtgatgactgggtgaagc agaagccaggccaggcctggagtggatcggctatatcaatcctacaacgacg gcaccaagtataacgagaagtttaagggaaggccaccctgacaagcgataaga gtcctctaccgctacatggagctgagctccctgacaagcgaggactccgccgt gtactattgcgccagaggcacctactattacggctccagggtgttcgattactgggg ccagggcacaacctgacagtgcttagcggaggaggaggcagcggaggaggga ggctccggcggcggcggctctgacatcgtgatgaccaggcagcaccatccatc cctgtgacaccaggcgagtctgtgagcatctcctgtcggctccttaagtcctgtg aacttaatggcaacacctatctgtactggttctgcagcggcccggacagctcct cagctgctgatctataggatgagcaatctggcctccggcgtgctgatcgcttct ggcagcggctccggcaccgctttactgaggatcagccgctggaggcaga ggacgtgggctgtattactgcatgcagcacctggagtaccctttaccctttggcgc</p>

		<p>cggcacaagctggagctgaagcggagcagccccacaaccacaccagcacctc ggccaccaccccagcaccaacaatcgcatctcagccactgagcctgcgccccg aggcctgtagccagccgcccggcggcagtgacacccggggcctggacttc gcctgcatatctacatctggcccctctggccggcactgcgccgtgctgtact gagcctggatcacctgtattgcaagcggggcagaaagaagctgctgtacatct tcaagcagccctcatgcccgtgcagaccacacaggaggaggatggctgct cctgtagattcccagaggaggaggaggatgtgagctgcgctgaaagtctc tcggagcggcagcgcactgcatatcagcagggaagaaccagctgtacaatga gctgaacctggccggagagaggatgacgtgctggataagagacggggcc gggaccccagatgggaggcaagccccgggaagaatcctcaggaggccct gtataacgagctgcagaaggataagatggccgaggcctacagcgagatcgccat gaaggagagagaaggcggcaaggccacgacggcctgtaccaggccct gtccacagcaacaaggatactatgacgctctgcacatgcaggctctgccccctc gggga</p>
7	ALLO-501_v1.6	<p>atgggaaccagcctgctgtgctggatggcactgtgcctgctgggagcagaccac gccgatgccgaactgcctactcaggggacattcttaatgtgagcaccaacgtga gctctggaggaggaggctccgagctgccaaccaggccacattcttaatgtgag cacaacgltctcccgaagcctaccacaaccgccgaactgcctaccagggg cacattttcaacgtgtctaccaacgtgtctagcggaggaggaggctccccgcac ctaggccccctaccccagcaccaacaatgcaagccagcctctgtccctgcccc agaggcatgcaggccagcagcaggaggagcagtgacacccgcggcctggac ttgctgcatatctatctgggaccactggcaggaacctgtggcgtgctgctg ctgtctctgctcatccctgtactgcaatcacagaaccggagaagggtgtgcaa gtgtcctcggccagtggtagagcagaggggaaggggcagcctgctgacctgtgg cgactggaggagaatcccggccctatggagacagatacactgctgctgtgggt gctgctgctgtgggtgccaggctctacaggagagggtgcagctgcagcagagcgg acctgagctgatcaagccaggcgcctctgtgaagatgagctgcaaggcctccgg ctataccttcacaagctacgtgatgcactgggtgaagcagaagccaggccagggc ctggagtgatcggctataatccctacaacgacggcaccaagtataacgaga agtttaagggaaggccaccctgacatccgataagagctcctctaccgcctacatg gagctgagctccctgacatccgaggactctgcccgtgactattgcgccagaggca cctactattacggctctagggtgttcgattactggggccaggggcacaacctgaca gtgttagcggaggaggaggctctggaggaggaggcagcggcggcggaggct ccgacatcgtgatgaccaggcagcaccatccatcccagtgacacctggcgaga gcgtgtccatctctgtaggtccttaagtctctgctgaacagcaatggcaacaccta tctgtactggttctgcagcggcccggacagagccctcagctgctgatctataggat gtccaatctggcctctggagtgcctgatcgttcagcggctccggctctggaaccg ccttlacactgaggatctcccgtggaggcagaggacgtggcgctgtattactgc atgcagcacctggagtaccctttcacctttggcggccggcacaagctggagctga agcggagcagccccacaaccaaccagcaccggccaccaacctgccccct acaatcgcaagccagccactgtccctgcccggaggcctgtagacctgcccgc ggcggcggcctccatacccggcctggatttcgctgcatatctacattgggc ccctctggccggcacttgcggcgtgctgctgctgagcctggctacacctgtattg caagcggggcagaaagaagctgctgtacatctcaagcagccctcatgccc gtgcagaccacacaggaggaggatggctgctcctgtagattcccagaggaggag gaggaggatgtgagctgcccgtgaagtttagccggctccggcagcaccctgca tatcagcaggccagaaccagctgtacaatgagctgaacctggccggagagag gagtagcagctgctggataagagaaggggacgggaccccagatgggaggca agccccggcgaagaatcctcaggaggcctgtataacgagctgcagaaggata</p>

		<p>agatggccgaggcctacagcgagatcggcatgaagggagagagaaggcgcgg caagggacacgacggcctgtatcagggcctgtccaccgccacaaaggcacaccta cgalgccctgcacatgcaggccctgcctccaagatga</p>
8	<p>ALLO-501_v1.0 (or ALLO-501) Exemplary signal sequence is underlined</p>	<p><u>MLTSLLCWMALCLLGADHADAC</u>PYSNPSLCSGGGG SELPTQGTFSNVSTNVSPAKPTTTACPYSNPSLCSGG GGSPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHT RGLDFACDIYWAPLAGTCGVLLLSL VITLYCNHRN RRRVCKCPRPVVRA [EGRGSLTTCGDVEENPGP] <u>METDLLLLWVLLLWVPGSTGEVQLQQSGPELIKPG</u> ASVKMSCKASGYTFTSYVMHWWKQKPGQGLEWIG YINPYNDGTKYNEKFKGKATLTSDKSSSTAYMELSS LTSEDSAVYYCARGTYYYGSRVFDYWGQGTTLTVS SGGGGSGGGGSGGGGSDIVMTQAAPSIPVTPGESVSI SCRSSKLLNSNGNTYL YWFLQRPGQSPQLLIYRMS NLASGVDPDRFSGSGSGTAFTLRISRVEAEDVGVYYC MQHLEYPFTEGAGTKLELKRSDPTTTPAPRPPTPAPT IASQPLSLRPEACRPAAGGAVHTRGLDFACDIYWAP LAGTCGVLLLSL VITLYCKRGRKLLYIFKQPFMRP VQTTQEEDGCSCRFP EEEEGGCELRVKFSRSADAPA YQQGQNQLYNELNLGRREEYDVLDKRRGRDP EMG GKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERR RGKGHDGLYQGLSTATKDTYDALHMQUALPPR</p>
9	<p>ALLO-501_v1.1</p>	<p><u>METDLLLLWVLLLWVPGSTGEVQLQQSGPELIKPG</u> ASVKMSCKASGYTFTSYVMHWWKQKPGQGLEWIG YINPYNDGTKYNEKFKGKATLTSDKSSSTAYMELSS LTSEDSAVYYCARGTYYYGSRVFDYWGQGTTLTVS SGGGGSGGGGSGGGGSDIVMTQAAPSIPVTPGESVSI SCRSSKLLNSNGNTYL YWFLQRPGQSPQLLIYRMS NLASGVDPDRFSGSGSGTAFTLRISRVEAEDVGVYYC MQHLEYPFTEGAGTKLELKRSDPTTTPAPRPPTPAPT IASQPLSLRPEACRPAAGGAVHTRGLDFACDIYWAP LAGTCGVLLLSL VITLYCKRGRKLLYIFKQPFMRP VQTTQEEDGCSCRFP EEEEGGCELRVKFSRSADAPA YQQGQNQLYNELNLGRREEYDVLDKRRGRDP EMG GKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERR RGKGHDGLYQGLSTATKDTYDALHMQUALPPR</p>
10	<p>ALLO-501_v1.2</p>	<p><u>METDLLLLWVLLLWVPGSTGEVQLQQSGPELIKPG</u> ASVKMSCKASGYTFTSYVMHWWKQKPGQGLEWIG YINPYNDGTKYNEKFKGKATLTSDKSSSTAYMELSS LTSEDSAVYYCARGTYYYGSRVFDYWGQGTTLTVS SGGGGSGGGGSGGGGSDIVMTQAAPSIPVTPGESVSI SCRSSKLLNSNGNTYL YWFLQRPGQSPQLLIYRMS NLASGVDPDRFSGSGSGTAFTLRISRVEAEDVGVYYC MQHLEYPFTEGAGTKLELKRSDPTTTPAPRPPTPAPT IASQPLSLRPEACRPAAGGAVHTRGLDFACDIYWAP</p>

		LAGTCGVLLLSLVITLYCKRGRKLLYIFKQPFMRP VQTTQEEDGCSCRFP EEEEGGCEL RVKFSRSADAPA YQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMG GKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERR RGKGHDGLYQGLSTATKDTYDALHMQUALPPR
11	ALLO-501_v1.3	<u>MGTSLLCWMALCLLGADHADASGGGGSPAPRPPTP</u> APTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYI WAPLAGTCGVLLLSLVITLYCNHRNRRRVCKCPRPV VRAEGRGSLTTCGDVEENPGP <u>METDLLLLWVLLLW</u> <u>VPGSTGEVQLQQSGPELIKPGASVKMSCKASGYTFT</u> SYVMHWVKQKPGQGLEWIGYINPYNDGTKYNEKF KGKATLTSDKSSSTAYMELSSLTSEDSAVYYCARGT YYYGSRVFDYWGQGTTLTVSSGGGGSGGGGSGGG GSDIVMTQAAPSIPVTPGESVSISCRSSKSLNNSGNT YLYWFLQRPQGQSPQLLIYRMSNLASGVPDRFSGSGS GTAFTLRISRVEAEDVGVYYCMQHLEYPFTFGAGTK LELKRSDPTTTPAPRPPTPAPTIASQPLSLRPEACRPA AGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSLVIT LYCKRGRKLLYIFKQPFMRPVQTTQEEDGCSCRFP EEEEGGCEL RVKFSRSADAPAYQQGQNQLYNELNL GRREEYDVLDKRRGRDPEMGGKPRRKNPQEGLYNE LQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLST ATKDTYDALHMQUALPPR
12	ALLO-501_v1.4	<u>MGTSLLCWMALCLLGADHADASGGGGSELPTQGT</u> SNVSTNVSSGGGGSPAPRPPTPAPTIASQPLSLRPEAC RPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSL VITLYCNHRNRRRVCKCPRPVVRAEGRGSLTTCGD VEENPGP <u>METDLLLLWVLLLWVPGSTGEVQLQQSG</u> <u>PELIKPGASVKMSCKASGYTFTSYVMHWVKQKPGQ</u> GLEWIGYINPYNDGTKYNEKFKGKATLTSDKSSSTA YMELSSLTSEDSAVYYCARGTYYYGSRVFDYWGQ GTTLTVSSGGGGSGGGGSGGGGSDIVMTQAAPSIPV TPGESVSISCRSSKSLNNSGNTYLYWFLQRPQGQSPQ LLIYRMSNLASGVPDRFSGSGSGTAFTLRISRVEAED VGVYYCMQHLEYPFTFGAGTKLELKRSDPTTTPAPR PPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFAC DIYIWAPLAGTCGVLLLSLVITLYCKRGRKLLYIFK QPFMRPVQTTQEEDGCSCRFP EEEEGGCEL RVKFSR SADAPAYQQGQNQLYNELNLGRREEYDVLDKRRG RDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIG MKGERRRGKGHDGLYQGLSTATKDTYDALHMQUAL PPR
13	ALLO-501_v1.5	<u>MGTSLLCWMALCLLGADHADACSGGGGGSCSGGG</u> GSELPTQGTFSNVSTNVSPAKPTTACSGGGGGSCS GGGGSPAPRPPTPAPTIASQPLSLRPEACRPAAGGAV HTRGLDFACDIYIWAPLAGTCGVLLLSLVITLYCNH

		<p>RNRRRVCKCPRPVVRAEGRGSLTTCGDVEENPGPM <u>ETD</u>TLLWVLLWVPGSTGEVQLQSGPELIKPGAS VKMSCKASGYTFTSYVMHWVKQKPGQGLEWIGYI NPYNDGTKYNEKFKGKATLTSDKSSSTAYMELSSLT SEDSAVYYCARGTYYYGSRVFDYWQGTTLTVSSG GGGSGGGSGGGSDIVMTQAAPSIPVTPGESVSISC RSSKSLNSNGNTYLYWFLQRPQSQPQLLIYRMSNL ASGVPDRFSGSGSGTAFTLRISRVEAEDGVVYYCMQ HLEYPFTFGAGTKLELKRSDPTTTPAPRPPTPAPTAS QPLSLRPEACRPAAGGAVHTRGLDFACDIYWAPLA GTCGVLLLSLVITLYCKRGRKLLYIFKQPFMRPVQ TTQEEDGCSCRFPEEEEGGCELRVKFSRSADAPAYQ QGQNQLYNELNLGRREEYDVLDRRRGRDPEMGGK PRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRG KGHGGLYQGLSTATKDTYDALHMQALPPR</p>
14	ALLO-501_v1.6	<p><u>MGT</u>SLLCWMALCLLGADHADAELPTQGTFSNVSTN VSSGGGGSELPTQGTFSNVSTNVSPAKPTTAAELPTQ GTFSNVSTNVSSGGGGSPAPRPPTPAPTASQPLSLRP EACRPAAGGAVHTRGLDFACDIYWAPLAGTCGVL LLSLVITLYCNHRNRRRVCKCPRPVVRAEGRGSLT CGDVEENPGPM<u>ETD</u>TLLWVLLWVPGSTGEVQLQ QSGPELIKPGASVKMSCKASGYTFTSYVMHWVKQK PGQGLEWIGYINPYNDGTKYNEKFKGKATLTSDKSS STAYMELSSLTSEDSAVYYCARGTYYYGSRVFDYW GQGTTLTVSSGGGGSGGGSGGGSDIVMTQAAPSI PVTGPESVSISSKSLNSNGNTYLYWFLQRPQSQ PQLLIYRMSNLASGVPDRFSGSGSGTAFTLRISRVEA EDVGVVYYCMQHLEYPFTFGAGTKLELKRSDPTTTP APRPPTPAPTASQPLSLRPEACRPAAGGAVHTRGLD FACDIYWAPLAGTCGVLLLSLVITLYCKRGRKLL YIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCELRV KFSRSADAPAYQGGQNQLYNELNLGRREEYDVLDR RRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAY SEIGMKGERRRGKGHGGLYQGLSTATKDTYDALH MQALPPR</p>
15	EF1alpha (long_1189)_promoter_GRCh38.p12 (v1.0_v1.1_v1.3_v1.4_v1.5_v1.6)	<p>gcgtgaggctccggtgcccgtcagtgggcagagcgcacatgccccacagtcccc gagaagtggggggagggtcggcaattgaaccggtgcctagagaagggtggcg cggggtaaactgggaaagtgatgctgtgactggctcccttttcccgagggtg ggggagaaccgtatataagtgcagtagtcgcccgtgaacgttCTTTTCGC AACGGGTTTGCCGCCAGAACACAGgtaagtgcctgtgtgg tccccgggcctggcctctttacgggttatggcccttgcgtgccctgaattactcc acgcccctggctgcagtagtgattcttgatccccgagcttcgggttgaagtgggt gggagagtgcaggccttgcgcttaaggagcccccttcgctcgtgcttgagttgag gcctggcctggcgctggggccgccgctgcgaatctggtggcaccttcgcgcc tgtctcgtgcttgcataagtctctagccatttaaaattttgatgacctgctgcgacg cttttttctggcaagatagctttaaagtgggccaagatctgcacactggtatttc ggltttggggccgggcgggcgacggggcccgtgcgtcccagcgcacatgttc ggcgaggcggggcctgcgagcggccaccgagaatcgacgggggtagtct caagctggccggcctgctctggtgctggcctcgccgcccgtgtatgccccgc</p>

		cctggggcggcaaggctggcccggcggcaccagttgctgagcggaaagatgg ccgcttcccggccctgctgcagggagctcaaatggaggacgcggcgctcggg agagcggcgggtgagtcaccacacaaaggaaaagggccttccgctcagc cgtcgcttcatgtgactccacggagtaccggcgccgctccaggcacctcgattag tctcagcctttggagtacgctgcttttaggtggggggaggggtttatgcgatgga gttccccacactgagtgggtggagactgaagttaggccagctggcactgatgta attctccttgaattgcccttttgagtttgatcttggttcattctcaagcctcagaca gtggtcaaagtttttcttccattcagGTGTCGTGA
16	>EF1alpha_(short _237)_promoter_ GRCh38.p12_(v1 .2)	gcgtgaggctccgggtgcccgtcagtgggcagagcgcacatcgcccacagtccc gagaagttggggggaggggtcggcaattgaaccgggtgcctagagaaggtggcg cggggtaaactgggaaagtgatgctgtgactggctccgcttttcccagggtg ggggagaaccgtatataagtgcagtagtcgccgtgaacgtCTTTTTCGC AACGGGTTTGCCGCCAGAACACAG
17	4G7 immunoglobulin gamma 1 heavy chain	MEWSWIFLFLLSGTAGVHSEVQLQQSGPELIKPGAS VKMSCKASGYTFTSYVMHWVKQKPGQGLEWIGYI NPYNDGTKYNEKFKGKATLTSKSSSTAYMELSSLT SEDSAVYYCARGTYYYGSRVFDYWGQGTTLTVSSA KTPPSVYPLAPGSAAQTNMVTLGCLVKGYFPEPV TVTWNSGSLSSGVHTFPAVLQSDLYTLSSSVTPVPSST WPSETVTCNVAHPASSTKVDKKIVPRDCGCKPCICT VPEVSSVFIFPPKPKDVLITLTPKVTCVVVDISKDDP EVQFSWFVDDVEVHTAQTQPREEQFNSTFRSVSELPI MHQDWLNGKEFKCRVNSAAFPAPIEKTISKTKGRPK APQVYTIPPPKEQMAKDKVSLTCMITDFFPEDITVE WQWNGQPAENYKNTQPIMDTDGSYFVYSKLNQVK SNWEAGNTFTCSVLHEGLHNHHTKSLSHSPGK
18	4G7 immunoglobulin kappa light chain	MRCLAEFLGLLVLWIPGAIGDIVMTQAAPSIPVTPGE SVSISCRSSKLLNSNGNTLYWFLQRPQSPQLLIY RMSNLAGVPPDRFSGSGSGTAFTLRISRVEAEDVGV YYCMQHLEYPFTFGAGTKLELKRADAAPTVSIFPPS SEQLTSGGASVVCFLNRFYPKDINVKWKIDGSERQN GVLNSWTDQDSKDYSTYSMSSTLTLTKDEYERHNSY TCEATHKTSTSPIVKSFNREK
19	RQR8	CPYSNPSLCSGGGGSELPTQGTFSNVSTNVSPAKPTT TACPYSNPSLCSGGGGSP APRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLD FACDIYWAPLAGTCGVLLLS LVITLYCNHRNRRRVCKCPRPVV
20	Suicide peptide signal sequence	MGTSLLCWMALCLLGADHADA
21	Signal Sequence and RQR8	MGTSLLCWMALCLLGADHADACPYSNPSLCSGGG GSELPTQGTFSNVSTNVSPAKPTTTACPYSNPSLCSG GGGSPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVH TRGLDFACDIYWAPLAGTCGVLLLSLVITLYCNHR NRRRVCKCPRPVV

22	Rituximab mimotope	CPYSNPSLC
23	Palivizumab epitope	NSELLSLINDMPITNDQKKLMSNN
24	Cetuximab mimotope 1	CQFDLSTRRLKC
25	Cetuximab mimotope 2	CQYNLSSRALKC
26	Cetuximab mimotope 3	CVWQRWQKSYVC
27	Cetuximab mimotope 4	CMWDRFSRWYKC
28	Nivolumab Epitope 1	SFVLNWYRMSPSNQTDKLAAPEDR
29	Nivolumab Epitope 2	SGTYLCGAISLAPKAQIKE
30	QBEND-10 Epitope	ELPTQGTFSNVSTNVS
31	Alemtuzumab epitope	GQNDTSQTSSPS
32	FcγRIIIα hinge	GLAVSTISSFFPPGYQ
33	CD8α hinge	TTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHT RGLDFACD
34	IgG1 hinge	EPKSPDKTHTCPPCPAPPVAGPSVFLFPPKPKDTLMI ARTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKC KVSNAKALPAPIEKTISKAKGQPREPQVYTLPPSRDEL TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTT PPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMH EALHNHYTQKSLSLSPGK
35	CD8α transmembrane	IYIWAPLAGTCGVLLLSLVIT
36	CD8α hinge/ transmembrane	TTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHT RGLDFACDIYIWAPLAGTCGVLLLSLVIT
37	41BB (intracellular domain)	KRGRKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEE EGGCEL

38	CD3ζ cytoplasmic signaling domain	LRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDV LDKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMA EAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDA LHMQUALPPR
39	EF1a first intron	GTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCT CTTTACGGGTTATGGCCCTTGCGTGCCTTGAATTA CTTCCACGCCCTGGCTGCAGTACGTGATTCTTGA TCCCGAGCTTCGGGTTGGAAGTGGGTGGGAGAGT TCGAGGCCTTGCGCTTAAGGAGCCCCCTTCGCCTCG TGCTTGAGTTGAGGCCTGGCCTGGGCGCTGGGGC CGCCGCGTGCGAATCTGGTGGCACCTTCGCGCCTG TCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAAA ATTTTTGATGACCTGCTGCGACGCTTTTTTTCTGGC AAGATAGTCTTGTAATGCGGGCCAAGATCTGCA CACTGGTATTTTCGGTTTTTGGGGCCGCGGGC ACGGGGCCCGTGCCTCCAGCGCACATGTTCCGGC GAGGCGGGGCTGCGAGCGCGGCCACCGAGAATC GGACGGGGGTAGTCTCAAGCTGGCCGGCCTGCTC TGGTGCCTGGCCTCGCGCCGCGGTGTATCGCCCCG CCCTGGGCGGCAAGGCTGGCCCGGTCGGCACCAG TTGCGTGAGCGGAAAGATGGCCGCTTCCCGGCC TGCTGCAGGGAGCTCAAATGGAGGACGCGGGCGC TCGGGAGAGCGGGCGGGTGAGTCACCCACACAAA GGAAAAGGGCCTTTCGTCCTCAGCCGTCGCTTCA TGTGACTCCACGGAGTACCGGGCGCCGTCCAGGC ACCTCGATTAGTTCTCGAGCTTTTGGAGTACGTCG TCTTTAGGTTGGGGGGAGGGGTTTTATGCGATGG AGTTTCCCACACTGAGTGGGTGGAGACTGAAGT TAGGCCAGCTTGGCACTTGATGTAATCTCCTTGG AATTTGCCCTTTTGGAGTTGGATCTTGGTTCATTC TCAAGCCTCAGACAGTGGTTCAAAGTTTTTTCTT CCATTTAG
40	LQL8	CSGGGGGSCSGGGGSELPTQGTFSNVSTNVSPAKPT TTACSGGGGGSCSGGGGSPAPRPPTPAPTIASQPLSL RPEACRPAAGGAVHTRGLDFACDIYTWAPLAGTCG VLLLSLVITLYCNHRNRRRVCKCPRPVV

WHAT IS CLAIMED IS:

1. An isolated polynucleotide encoding a polypeptide comprising an anti-CD19 chimeric antigen receptor (CAR) that is at least 70% identical to SEQ ID NO: 9, wherein the polypeptide does not comprise a rituximab binding site, and wherein the polynucleotide comprises a short EF1a promoter that is capable of expressing the anti-CD19 chimeric antigen receptor (CAR) in a mammalian T cell.
2. The isolated polynucleotide of claim 1, wherein the short EF1a promoter does not comprise an intron comprised by the nucleic acid sequence of SEQ ID NO:15.
3. The isolated polynucleotide of claim 2, wherein the intron comprises the nucleic acid sequence of SEQ ID NO:39.
4. The isolated polynucleotide of any one of the preceding claims, wherein the short EF1a promoter comprises the nucleic acid sequence of SEQ ID NO:16,
5. The isolated polynucleotide of any one of the preceding claims, wherein the polypeptide further comprises a safety switch.
6. The isolated polynucleotide of claim 5, wherein the safety switch is linked to the CD19 CAR using a linker peptide.
7. The isolated polynucleotide of claim 5, wherein the safety switch is linked to the anti-CD19 CAR using a T2A linker.
8. The isolated polynucleotide of any one of claims 5-7, wherein the safety switch comprises an antibody binding site.
9. The isolated polynucleotide of any one of claims 5-7, wherein the safety switch comprises a mutated CD20 mimotope.

10. The isolated polynucleotide of any one of claims 5-9, wherein the polypeptide comprises a CD34 epitope.
11. The isolated polynucleotide of claim 10, wherein the CD34 epitope is a QBEND-10 epitope.
12. The isolated polynucleotide of any one of claims 5-11, wherein the polypeptide further comprises a CD8 hinge/transmembrane domain.
13. The isolated polynucleotide of any one of the preceding claims, comprising a nucleic acid sequence that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to any one of the nucleic acid sequences of SEQ ID NOs: 1-7.
14. The isolated polynucleotide of any one of the preceding claims, wherein the polynucleotide encodes a polypeptide that is at least about 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to any one of the amino acid sequences of SEQ ID NOs: 8-14.
15. A vector comprising the isolated polynucleotide of any one of the preceding claims.
16. The vector of claim 15, wherein the vector is a retroviral vector, a DNA vector, a plasmid, an RNA vector, an adenoviral vector, an adenovirus associated vector, a lentiviral vector, or any combination thereof.
17. An engineered immune cell comprising the isolated polynucleotide of any one of claims 1-14, wherein the engineered immune cell does not express a rituximab binding site.
18. An engineered immune cell comprising the vector of claim 15 or 16, wherein the engineered immune cell does not express a rituximab binding site.
19. The engineered immune cell of claim 17 or 18, wherein the immune cell is a T cell, tumor infiltrating lymphocyte (TIL), NK cell, TCR-expressing cell, dendritic cell, or NK-T cell.
20. The engineered immune cell of claim 19, wherein the cell is an autologous T cell.

21. The engineered immune cell of claim 19, wherein the cell is an allogeneic T cell.
22. The engineered immune cell of any one of claims 17-21, wherein the cell comprises the polynucleotide that is at least 80%, 85%, 90%, 95%, 96%, 98%, 99% or 100% identical to the nucleic acid sequence of SEQ ID NO: 3.
23. The engineered immune cell of any one of claims 17-22, wherein the cell is resistant to rituximab.
24. A pharmaceutical composition comprising the engineered immune cell of any one of claims 17-23.
25. A method of treating a disease or disorder in a subject in need thereof, comprising administering to the subject the engineered immune cell of any one of claims 17-23.
26. A method of treating a disease or disorder in a subject in need thereof, comprising administering to the subject the pharmaceutical composition of claim 24.
27. The method of claim 25 or 26, wherein the disease or disorder is Non-Hodgkin lymphoma (NHL).
28. The method of any one of claims 25-27, wherein the subject has been treated or is currently being treated with rituximab.
29. An article of manufacture comprising the engineered immune cell of any one of claims 17-23, or a pharmaceutical composition thereof.
30. Use of the engineered immune cell of any one of claims 17-23, for treating a disease or disorder in a subject in need thereof.
31. Use of the engineered immune cell of any one of claims 17-23, for the preparation of a medicament for treating a disease or disorder in a subject in need thereof.

32. Use of the pharmaceutical composition of claim 24, for treating a disease or disorder in a subject in need thereof.
33. Use of the pharmaceutical composition of claim 24, for the preparation of a medicament for treating a disease or disorder in a subject in need thereof.




Fig. 1

SUBSTITUTE SHEET (RULE 26)

Day 5 CAR expression (541 and 604)




Fig. 2A

Day 5 CAR and CD34 expression (Donor 410)




Fig. 2B

SUBSTITUTE SHEET (RULE 26)

Normalized cell expansion and final CAR expression (4 donors)




Fig. 3

Cell expansion and CAR expression (Donor 541)




Fig. 4A




Fig. 4C




Fig. 4D




Fig. 5A




Fig. 5B

11/27






Fig. 5C




Fig. 5D

Day 9 Phenotype and Activation




Fig. 6A




Fig. 6B

SUBSTITUTE SHEET (RULE 26)

Day 9 Phenotype and Activation




Fig. 6C




Fig. 6D

SUBSTITUTE SHEET (RULE 26)




Fig. 8A




Fig. 8B




Fig. 8C




Fig. 8D




Fig. 9

Short-term (24hr) killing assays (Targets = Raji cells)

All donors




Fig. 10




Fig. 11B




Fig. 11A




Fig. 11D




Fig. 11C




Fig. 12




Fig. 13A




Fig. 13B