



(51) International Patent Classification:

A61K 31/7088 (2006.01) A61P 37/06 (2006.01)
A61K 35/17 (2015.01) A61P 37/08 (2006.01)
A61K 48/00 (2006.01)

(21) International Application Number:

PCT/US2016/053893

(22) International Filing Date:

27 September 2016 (27.09.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/233,517 28 September 2015 (28.09.2015) US

(71) Applicant: TRUSTEES OF DARTMOUTH COLLEGE
[US/US]; 11 Rope Ferry Road, Hanover, NH 03755 (US).(72) Inventors: SENTMAN, Charles, L.; 25 Wedgewood
Drive, Grantham, NH 03753 (US). ARELLANO, Ben-
jamin, H.; 40 Wolf Road, Unit #13, Lebanon, NH 03766
(US).(74) Agents: LICATA, Jane, Massey et al.; Licata & Tyrrell
P.C., 66 E. Main Street, Marlton, NJ 08053 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: CHIMERIC ANTIGEN RECEPTOR ANTI-INFLAMMATORY CELLS AND METHODS OF USE




FIG. 1

(57) Abstract: A nucleic acid construct and an immune cell, which harbor nucleic acids encoding a CAR and nucleic acids encoding at least one anti-inflammatory or immunosuppressant protein and methods of using the same in treatment or amelioration of inflammation or immune-mediated autoimmunity are described.

Chimeric Antigen Receptor Anti-Inflammatory Cells and Methods of Use

Introduction

[0001] This application claims the benefit of priority of U.S. Provisional Application No. 62/233,517, filed September 28, 2015, the content of which is incorporated herein by reference in its entirety.

Background

[0002] Although genetic modifications to stimulate the immune system are beneficial for battling infectious organisms and cancer, there also exists a set of devastating diseases that are caused by an over-zealous and unchecked immune response. The targeting of self-antigens under normal physiologic conditions can cause a range of serious ailments including type 1 diabetes, multiple sclerosis, rheumatoid arthritis, systemic lupus erythematosus, and autoimmune encephalomyelitis. A relatively new set of autoimmune diseases, categorized as autoinflammatory diseases, have also been characterized including familial Mediterranean fever (FMF), neonatal onset multisystem inflammatory disease (NOMID), tumor necrosis factor (TNF) receptor-associated periodic syndrome (TRAPS), deficiency of the Interleukin-1 receptor antagonist (DIRA) and Behcet's disease. Additional inflammatory diseases that cause morbidity and mortality in a large number of patients include inflammatory bowel disease (Crohn's disease and ulcerative colitis), chronic granulomatous disease (CGD), and the various forms of vasculitis.

[0003] In general, dampening the immune response is the ideal treatment for autoimmune and inflammatory diseases

and current therapies revolve around the use of steroids, cytokine antagonists, or directly down-regulating the immune system using various modalities. Gene therapies may provide a viable biological alternative to directly blunt an overactive immune response. Over-expressing anti-inflammatory cytokines such as IL-10, TGF- β , IL-30, IL-35 or IL-37 in T cells or monocyte/dendritic cell populations *ex vivo* with a reinfusion of the modified cells has been suggested to aid in decreasing inflammatory driven symptoms (Kerkar (2013) *Frontiers Immunol.* 4:1-7; Mathisen et al. (1997) *J. Exp. Med.* 186:159-164). Another suggested alternative is to construct a decoy cytokine receptor that contains the correct receptor sequence to enable for binding of pro-inflammatory cytokines such as IL-12 combined with a non-functioning cytoplasmic signal transducing sequence. Over-expressing these "dominant-negative" receptors would enable reinfused immune cells to function as sinks for the inflammatory cytokines responsible for the pathophysiology of the disease.

[0004] US 2015/0038684 discloses a bispecific antigen receptor containing at least two antigen-specific targeting regions that target antigens specific for cancer, inflammatory disease, neuronal disorder, diabetes, cardiovascular disease, infectious disease, autoimmune disease, and combinations thereof.

[0005] Similarly, US 2010/0135974 describes a redirected regulatory T lymphocyte endowed with specificity toward a selected target antigen or ligand by expressing a chimeric receptor polypeptide. This reference indicates that the redirected regulatory T cells at sites of inflammation results in suppression of inflammatory conditions, commonly part of organ-specific autoimmune disease. In addition, this reference suggests that redirected regulatory T cells

of the invention can be triggered or activated to release suppressive cytokines that will result in suppression of any "bystander" effector T-cells, and by this mechanism, quell an ongoing inflammatory/autoimmune response.

Summary of the Invention

[0006] The invention provides an immune cell and nucleic acid constructs harboring nucleic acids encoding a chimeric antigen receptor and nucleic acids encoding at least one exogenous anti-inflammatory or immunosuppressant protein. In some embodiments, the cell is a T lymphocyte. In one embodiment, the chimeric antigen receptor is a single chain variable fragment. In another embodiment, the chimeric antigen receptor includes an antigen targeting domain or recognition domain that binds an antigen or ligand at a site of inflammation or autoimmunity and an immune signaling receptor containing an immunoreceptor tyrosine-based activation motif. In a further embodiment, the chimeric antigen receptor further includes a costimulatory endodomain. An anti-inflammatory or immunosuppressant protein includes a cytokine, a virokin (e.g., viral IL-10 protein) or a C-type lectin protein (e.g., a helminth C-type lectin protein). Further, in some embodiments, expression of the anti-inflammatory or immunosuppressant protein is regulated by a promoter that is enhanced by signaling from the chimeric antigen receptor. A method for treating inflammation or immune-mediated autoimmunity using an immune cell of the invention and kits containing one or more immune cells or the nucleic acid construct of the invention are also provided.

Brief Description of the Drawings

[0007] Figure 1 shows that CD4⁺ CD25⁺ CD127⁻ Foxp3⁺ Treg cells (Treg) incubated in the presence of stimulus (Stim) and carboxyfluorescein succinimidyl ester (CFSE)-labeled CD4⁺ T cells (Teff) are capable of suppressing CD4⁺ T cell proliferation in a dose dependent manner, as measured by CFSE dilution assays.

[0008] Figure 2 shows that CD4⁺ CD25⁺ CD127⁻ Foxp3⁺ Treg cells can be transduced with a CAR construct without losing Foxp3 expression.

Detailed Description of the Invention

[0009] The present invention is directed to chimeric immune cells that reduce or inhibit inflammation and subsequent tissue damage. In particular, this invention provides chimeric immune cells that secrete at least one anti-inflammatory or immunosuppressant protein that inhibits or reduces a proliferative and/or proinflammatory immune response preferentially at the site of inflammation or immune-mediated autoimmunity. In addition to an anti-inflammatory or immunosuppressant protein, the immune cells express a chimeric antigen receptor (CAR) that binds a specific ligand(s) expressed at the site of inflammation and/or autoimmunity. Binding of the CAR to the ligand(s) allows the chimeric immune cells to localize and become activated at the site of tissue damage. Secretion of the anti-inflammatory or immunosuppressant protein by the activated CAR cells suppresses the activation and function of local immune cells thereby reducing or inhibiting an immune response. The chimeric immune regulatory cells of this invention are useful in the treatment or amelioration of diseases where chronic inflammation or immune-mediated

autoimmunity leads to tissue destruction and/or disease progression.

[0010] This invention provides nucleic acid constructs and immune cells, which harbor nucleic acids encoding a CAR and nucleic acids encoding an anti-inflammatory or immunosuppressant protein. As used herein, "anti-inflammatory," and "immunosuppressant" or "immunosuppressive" protein refers to a protein that inhibits local immune responses. Such immune responses include vasodilatation; leukocyte infiltration; redness; heat; activation of local cells and immune cells; expression of major histocompatibility complex (MHC) class II molecules, costimulatory molecules (e.g., B7-1 and B7-2), and adhesion proteins (e.g., ICAM-1); IL-1 induced proliferation; and the release of pro-inflammatory cytokines (e.g., IL-1, IL-2, IL-6, IL-8, IL-12, GM-CSF and TNF- α) and/or inflammatory mediators (e.g., NO, free radicals, prostaglandins and metalloproteases) from these cells. Further, "anti-inflammatory" and "immunosuppressant" or "immunosuppressive" protein refers to a protein that affects the diseased area by suppressing or countering immunologic and/or pro-inflammatory activities. In certain embodiments, the anti-inflammatory or immunosuppressant protein is selected for having minimal or no immune activation or immunostimulatory activity. By way of illustration, whereas human IL-10 can costimulate thymocyte and mast cell proliferation and B cell MHC class II expression, viral IL-10 cannot (Vieira, et al. (1991) *Proc. Natl. Acad. Sci. USA*. 88:1172-1176; MacNeil, et al. (1990) *J. Immunol.* 145:4167-4173; Fei, et al. (1990) *J. Exp. Med.* 172:1625-1631). Accordingly, while some embodiments embrace a human anti-inflammatory or immunosuppressant protein, in other embodiments, the anti-inflammatory or

immunosuppressant protein is a non-human or non-mammalian protein.

[0011] The anti-inflammatory or immunosuppressant protein is exogenous in the sense that the recipient host cell (*i.e.*, immune cell) either does not naturally express the anti-inflammatory or immunosuppressant protein or does not express the anti-inflammatory or immunosuppressant protein in concert with a CAR, *e.g.*, on the same chimeric nucleic acid construct, from the same promoter, and/or from a promoter that is dependent upon CAR signaling.

[0012] Anti-inflammatory or immunosuppressant proteins of use in this invention include cytokines, virokines and C-type lectin proteins. Examples of anti-inflammatory or immunosuppressant cytokines include, but are not limited to, human Interleukin (IL)-10, IL-4, IL-35, IL-37, IL-27, IL-13, IL-1 receptor antagonist (IL-1ra), Interferon β (IFN- β), and Transforming Growth Factor β (TGF- β). An exemplary virokine is viral IL-10. Examples of anti-inflammatory or immunosuppressant C-type lectin proteins include helminth C-type lectin proteins such as *Heligmosomoides polygyrus* C-type lectin-1 (*Hp*-CTL-1) and *Nippostrongylus brasiliensis* C-type lectin-2 (*Nb*-CTL-2).

[0013] Human IL-10, also referred to as cellular IL-10 (cIL-10), exerts its function both directly and indirectly through upregulation of molecules including IL-1 receptor antagonists (IL-1ra) and soluble tumor necrosis factor α (TNF- α) receptor (sTNFR) and down-regulation of IL-8 and MCP-1 levels (Makhija & Kingsnorth (2002) *J. Hepatobiliary. Pancreat. Surg.* 4:401-410; Bhatia, et al. (2005) *Pancreatology* 2-3:132-144). Ultimately, the net effect is down-regulation of macrophages, neutrophils, and CD4+ T cells, decreased antigen presentation, augmented CD8+ T cell cytolytic activity, and enhanced

proliferation/differentiation of B cells. IL-10 has been shown to provide beneficial effects in TH1-mediated models diseases including experimental allergic encephalomyelitis ((EAE) Dai, et al. (2012) *J. Neuroimmunol.* 245(1-2):1-7; Rott, et al. (1994) *Eur. J. Immunol.* 24:1434-1440); pancreatitis (Van Laethem, et al. (1995) *Gastroenterology* 108:1917-1922), diabetes mellitus (Pennline, et al. (1994) *Clin. Immunol. Immunopathol.* 71:169-175), and experimental endotoxemia (Gerard, et al. (1993) *J. Exp. Med.* 177:547-550). IL-10 has also been effective in various animal models of arthritis, in reducing inflammation, in cellular infiltrates, and in joint destruction (Persson, et al. (1996) *Scand. J. Immunol.* 44:607-614; Tanaka, et al. (1996) *Inflamm. Res.* 45:283-288). These models have provided the basis for human trials in diseases such as multiple sclerosis, rheumatoid arthritis, psoriasis and Crohn's disease (Van Deventer, et al. (1997) *Gastroenterology* 113:383-389; Keystone, et al. (1998) *Rheum. Dis. Clin. N. Am.* 24:629-39; Asadullah, et al. (1998) *J. Clin. Investig.* 101:783-794). Exemplary IL-10 proteins of use in this invention include, but are not limited to, human IL-10 (GENBANK Accession No. NP_000563).

[0014] Like IL-10, human IL-4 is an important member of the TH2 family of anti-inflammatory cytokines, which has been shown to inhibit the ability of human monocytes to product TNF- α , IL-1 and prostaglandin E₂ (Hart, et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3803-3807; Essner, et al. (1989) *J. Immunol.* 142(11):3857-61). Because IL-4 antagonizes the effects of IL-12, a macrophage-derived cytokine promoting TH1 development, IL-4 represents an important regulator of the TH1/TH2 balance. The activity of IL-4 has been studied in several animal models of TH1-mediated autoimmunity, including CIA and streptococcal cell wall-induced arthritis

(Horsfall, et al. (1997) *J. Immunol.* 159(11):5687-96; Allen, et al. (1993) *J. Immunol.* 151:4344-4351). Further, local expression of IL-4 has been shown to protect against metalloproteinase-induced cartilage destruction during immune complex-mediated arthritis (van Lent, et al. (2002) *Osteoarthritis Cartilage* 10(3):234-43). In addition, IL-4 has been shown to synergize with IL-10 to limit mononuclear cell infiltration of synovial tissue and prevent cartilage degradation (van Meegeren, et al. (2012) *Osteoarthritis Cartilage* 20(7):764-772; Joosten, et al. (1997) *Arthritis Rheum.* 40:249-260). Therefore, the combined use of IL-4 and IL-10 is expected to be of particular use in the treatment of human autoimmune disease. Exemplary IL-4 proteins of use in this invention include, e.g., human IL-4 available under GENBANK Accession Nos. NP_000580 and NP_758858.

[0015] Human IL-35 is a heterodimeric protein with two subunits, IL-12 α (also known as IL-12p35) and Epstein-Barr virus induced 3 (EBI3; also known as IL-27 β) (Collison & Vignali (2008) *Immunol. Rev.* 226:248-262; Collison, et al. (2007) *Nature* 450:566-569; Devergne, et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:12041-6). IL-35 down-regulates Th17 cell development and inhibits autoimmune inflammation and has been shown to inhibit inflammation in various autoimmunity models such as experimental colitis (Collison, et al. (2007) *Nature* 450:566-569; Wirtz, et al. (2011) *Gastroenterology* 141:1875-86), collagen-induced autoimmune arthritis (Niedbala, et al. (2007) *Eur. J. Immunol.* 37:3021-3029; Kochetkova, et al. (2010) *J. Immunol.* 184:7144-53), autoimmune demyelination in central nervous system (Zandian, et al. (2011) *Gene Ther.* 18:734-42), and type 2 T helper cell (Th2)-mediated allergic asthma (Huang, et al. (2011) *J. Immunol.* 187:462-71). An exemplary IL-35 protein of use in this invention can be prepared using,

e.g., human IL-12 α available under GENBANK Accession No. NP_000873, and EBI3 available under GENBANK Accession No. NP_005755. The IL-12 α and EBI3 proteins of IL-35 can be expressed as separate proteins or as a single chain protein (Collison, et al. (2007) *Nature* 450:566-9). When expressed as a single chain protein, the IL-12 α and EBI3 proteins can be directly fused to one another as a contiguous protein sequence or fused together via a linker, e.g., (GGGS)₄ (SEQ ID NO:1).

[0016] IL-37 is encoded by the *IL1F7* gene that undergoes alternative splicing, which results in the expression of five different isoforms of IL-37. *IL1F7a* (isoform 5, IL-37a) uses a unique start codon in exon 3 (prodomain), which is then spliced in exon 4 to 6 (forming the putative 12 β -strand-containing protein structure). *IL1F7b* (isoform 1, IL-37b) encodes the longest transcript variant, composed of exons 1 and 2 (prodomain) and exons 4 to 6. *IL1F7c* (isoform 4, IL-37c) is a transcript variant encompassing exons 1 and 2 (prodomain) followed by exons 5 and 6. *IL1F7d* (isoform 2, IL-37d) has a prodomain limited to exon 1 (exon 2 is missing) followed by the complete IL-1-like sequence encoded by exons 4 to 6. *IL1F7e* (isoform 3, IL-37e) is only composed of exon 1, 5, and 6. IL-37b is the best characterized IL-37 isoform and has been shown to suppress the production of soluble TNF- α , IL-6, MIP-2, and of cell-associated IL-1 α in IL-37b-overexpressing cells in response to LPS (Sharma, et al. (2008) *J. Immunol.* 180:5477). Further, the *in vivo* effect of IL-37b has been examined in IL-37b transgenic (tg) mice, wherein sublethal endotoxic shock was significantly decreased in IL-37b-tg mice as compared to normal littermates, in terms of hypothermia, metabolic acidosis, dehydration, rise in potassium concentration, and liver damage (Nold, et al. (2010) *Nat.*

Immunol. 11:1014). In addition, levels of inflammatory cytokines (IL-6, IL-1 β , IL-17, IFN- γ , etc.) are dampened in IL-37b-tg mice in response to LPS challenge (Nold, et al. (2010) *Nat. Immunol.* 11:1014). In an experimental model of intestinal bowel disease (dextran sodium sulfate-induced colitis), the severity of the intestinal inflammation is significantly lower in IL-37b-tg mice as compared to wild-type controls (McNamee, et al. (2011) *Proc. Natl. Acad. Sci. USA* 108:16711-6). Exemplary IL-37 proteins of use in this invention include, e.g., human IL-37a available under GENBANK Accession No. NP_775297; human IL-37b available under GENBANK Accession No. NP_055254; human IL-37c available under GENBANK Accession No. NP_775296; human IL-37d available under GENBANK Accession No. NP_775294 and human IL-37d available under GENBANK Accession No. NP_775295.

[0017] IL-27 is a heterodimeric protein composed of EBI3 (also known as IL-27 β) and IL-27 α (also known as IL-30 or IL-27p28) (Pflanz, et al. (2002) *Immunity* 16:779-790), which signals through the IL27 receptor complex formed by WSX-1 and gp130 subunits. IL-27 has been shown to inhibit the Th17 response during inflammation (Stumhofer, et al. (2006) *Nat. Immunol.* 7:937-945; Batten, et al. (2006) *Nat. Immunol.* 7:929-936; Colgan & Rothman (2006) *Nat. Immunol.* 7:899-901), which contributes to the development of autoimmune inflammation in multiple sclerosis and rheumatoid arthritis. Consistent with the ability of IL-27 to decrease Th17 responses, continual delivery of IL-27 significantly suppresses the establishment of clinical disease in EAE, associated with a decreased proportion of Th17 cells in the CNS (Fitzgerald, et al. (2007) *J. Immunol.* 179:3268-3275). In addition, it has been found that IL-27 can also promote IL-10 synthesis by CD4⁺ and CD8⁺

T cells (Stumhofer, et al. (2007) *Nat. Immunol.* 8:1363-71; Fitzgerald, et al. (2007) *Nat. Immunol.* 8(12):1372-9; Awasthi, et al. (2007) *Nat. Immunol.* 8(12):1380-9). An exemplary IL-27 protein of use in this invention can be prepared using, e.g., human IL-27 α available under GENBANK Accession No. NP_663634, and EBI3 available under GENBANK Accession No. NP_005755. The IL-27 α and EBI3 proteins of IL-27 can be expressed as separate proteins or as a single chain protein. When expressed as a single chain protein, the IL-27 α and EBI3 proteins can be directly fused to one another as a contiguous protein sequence or fused together via a linker, e.g., (GGGS)₄ (SEQ ID NO:1).

[0018] IL-13 is implicated as a key player in allergic disease. IL-13 induces immunoglobulin (Ig) class switching to IgE (Punnonen & de Vries (1994) *J. Immunol.* 152:1094-1102) and upregulate vascular cell adhesion molecule (VCAM)-1 expression (Bochner, et al. (1995) *J. Immunol.* 154:799-803). However, actions of IL-13 consistent with an anti-inflammatory role have been indicated by its activity in several systems. In particular, IL-13 inhibits IL-1, TNF- α , and nitric oxide synthesis but increases IL-1 receptor antagonist production (Hart, et al. (1989) *Proc. Nat. Acad. Sci. USA* 86(10):3803-7; Minty, et al. (1993) *Nature* 361:248-250; De Wall, et al. (1993) *J. Immunol.* 151:6370-6381; Yanagawa, et al. (1995) *Am. J. Respir. Cell Mol. Biol.* 12:71-76; Berkman, et al. (1996) *Immunology* 89:363-367). IL-13 inhibits chemokine synthesis by vascular endothelial and airway smooth-muscle cells (Marfaing-Koka, et al. (1995) *J. Immunol.* 154:1870-1878; John, et al. (1997) *J. Immunol.* 158:1841-1847; John, et al. (1998) *Am. J. Respir. Cell Mol. Biol.* 18:84-90). *In vivo*, IL-13 has been shown to inhibit TNF- α release, neutrophil accumulation, and TNF- α production in rat lung immune

complex injury (Mulligan, et al. (1997) *J. Immunol.* 159:3483-3489), and endotoxin-induced lethality and increased serum TNF- α in the mouse (Di Santo, et al. (1997) *J. Immunol.* 159:379-382; Nicoletti, et al. (1997) *Eur. J. Immunol.* 27:1580-1583; Baumhofer, et al. (1998) *Eur. J. Immunol.* 28:610-615). Furthermore, in osteoarthritis synovial membrane treated with LPS, IL-13 inhibits the synthesis of IL-1 β , TNF- α and stromelysin-1, reduces the level of IL-1 β mRNA and stimulates the level of IL-1Ra mRNA (Jovanovic, et al. (1998) *Osteoarthritis Cartilage* 6(1):40-9). Therefore, IL-13 reduces the production of proinflammatory cytokines and metalloproteases, and favors the production of IL-1Ra. Exemplary IL-13 proteins of use in this invention include, e.g., human IL-13 available under GENBANK Accession Nos. NP_002179.

[0019] IL-1ra is a naturally occurring inhibitor of IL-1 (Arend (1993) *Adv. Immunol.* 54:167-227). IL-1ra is produced by a variety of cells including macrophages, fibroblasts, and keratinocytes (Arend, et al. (1985) *J. Immunol.* 134:3668-3875). IL-1ra binds to IL-1 receptor I on the cell surface but fails to trigger signal transduction, thereby acting as a competitive inhibitor of IL-1 binding to target cells (Dripps, et al. (1991) *J. Biol. Chem.* 266:10331-6). In inflammatory bowel disease, severe cases are associated with a decrease in IL-1ra leading to an imbalance of IL-1ra and IL-1 (Casini-Raggi, et al. (1995) *J. Immunol.* 154:2434-2440). The decrease of the IL-1ra:IL-1 ratio has been shown to account for acute cases of Crohn's disease and ulcerative colitis (Andus, et al. (1997) *Gut* 41:651-7; Heresbach, et al. (1997) *Am. J. Gastroenterol.* 92:1164-9). In the rabbit ligated-loop model of *Shigella* infection, administration of IL-1ra reduces intestinal inflammation (Perdomo, et al. (1994) *J. Exp. Med.* 180:1307-19),

indicating that the imbalance between IL-1 and IL-1ra accounts for disease severity (Arondel, et al. (1999) *Infect. Immun.* 67:6056-6066). Further, in animal models of osteoarthritis (OA), IL-1 blockade by IL-1Ra has been shown to slow the progression of disease by blocking PGE₂ synthesis, collagenase and nitric oxide (NO) production by chondrocytes, and cartilage matrix degradation (Martel-Pelletier, et al. (1999) *Front. Biosci.* 4:D694-703; Caron, et al. (1996) *Arthritis Rheum.* 39:1535-44). In addition, in patients with rheumatoid arthritis, IL-1ra injected subcutaneously daily demonstrates a disease-modifying antirheumatic effect (Bresnihan (2002) *Clin. Exp. Rheumatol.* 20:S32-S34). Exemplary IL-1ra proteins of use in this invention include, e.g., human IL-1ra available under GENBANK Accession Nos. NP_000568, NP_776213 or NP_776214, as well as the IL-1ra protein used in Anakinra, which is a 153 amino acid residue, recombinant human IL-1ra that differs from native human IL-1ra by the addition of a single methionine residue on its amino terminus.

[0020] IFN- β down-regulates the expression of IL-1 β and TNF- α and enhances IL-10 and IL-1 receptor antagonist production by lymphocytes *in vitro* (Tak, et al. (1999) *Rheumatology (Oxford)* 38:362-369; Rep, et al. (1996) *J. Neuroimmunol.* 67:111-118; Rep, et al. (1999) *J. Neuroimmunol.* 96:92-100), increases IL-1 receptor antagonist production by fibroblast-like synoviocytes (Palmer, et al. (2004) *Ann. Rheum. Dis.* 63:43-49), inhibits T-cell proliferation and migration, and prevents contact-dependent T-cell activation of monocytes (Jungo, et al. (2001) *Cytokine* 14:272-282). IFN- β also suppresses IFN- γ production and MHC class II expression by activated peripheral blood mononuclear cells (Yong, et al. (1998) *Neurology* 51:682-689). Further, studies have also found

that IFN- β enhances expression of TGF- β 1 and TGF- β 1 receptor type II by peripheral blood mononuclear cells (Ossege, et al. (2001) *Int. Immunopharmacol.* 1:1085-1100). IFN- β has shown therapeutic promise as an anti-inflammatory cytokine in multiple sclerosis (Li, et al. (1999) *Ann. Neurol.* 46:197-206) and in reducing inflammation and slowing cartilage destruction in CIA (van Holten, et al. (2004) *Arthritis Res. Ther.* 6(3):R239-49). In addition, *in vivo* data from Lewis rats with EAE have shown IFN- β -induced down-regulation of critical adhesion molecules such as ICAM-1 and monocyte VLA-4 that clearly play a role in the pathogenesis of multiple sclerosis by controlling cellular trafficking within the CNS (Ruuls & Sedgwick (1998) *Immunol. Cell Biol.* 76:65-73). Exemplary IFN- β proteins of use in this invention include, e.g., human IFN- β available under GENBANK Accession No. NP_002167, as well as the IFN β -1b protein used in Betaferon®/Betaseron® or Extavia®, and IFN β -1a protein used in Avonex® or Rebif®.

[0021] TGF- β antagonizes the activation of important target genes of pro-inflammatory stimuli of NF- κ B in macrophages and lymphocytes, such as inducible nitric oxide synthetase (iNOS) and MHC class I and class II antigens (Geiser, et al. (1993) *Proc. Natl. Acad. Sci.* 90:9944-9948; Vodovotz, et al. (1996) *J. Exp. Med.* 183:2337-2342). In addition, TGF- β decreases the expression of Type I and II IL-1 receptors and upregulates IL-1ra (Turner, et al. (1991) *Euro. J. Immunol.* 21:1635-39; Dubois, et al. (1990) *J. Exp. Med.* 172:737-744). Applied in the animal model EAE, TGF- β delays disease development and decreases the rate of recurrence in established disease (Kuruvilla, et al. (19910) *Proc. Natl. Acad. Sci. USA* 88:18-21). Further, in CIA, TGF- β decreases antibody production and alters histopathology with decreased lining membrane hyperplasia

and limited mononuclear cell infiltration (Kuruville, et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:18-21). Exemplary TGF- β proteins of use in this invention include, e.g., human TGF- β 1 available under GENBANK Accession No. NP_000651.

[0022] Viral IL-10 proteins are functional orthologues of cIL-10, which have been found thus far in the *Herpesviridae*, *Alloherpesviridae* and *Poxviridae* families of viruses. Viral IL-10 proteins have been found to inhibit PBMC proliferation and production of proinflammatory cytokines IL-1 α , IL-6, granulocyte-macrophage colony-stimulating factor, and TNF- α in lipopolysaccharide (LPS)-treated PMBC and monocytes (Spencer, et al. (2008) *Virology* 374:164-9; de Waal, et al. (1991) *J. Exp. Med.* 174:1209-1220; Hsu, et al. (1990) *Science* 250:830-2; Hsu, et al. (1992) *Int. Immunol.* 4:563-9; Vieira, et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:1172-6) and reduce the levels of transcription of components of the MHC class II biosynthesis pathway (Spencer, et al. (2008) *Virology* 374:164-9; Jenkins, et al. (2008) *J. Virol.* 82:3736-50; de Waal, et al. (1991) *J. Exp. Med.* 174:915-24). However, the viral cytokine had not retained all properties of cIL-10. While human and murine IL-10 costimulate mouse thymocyte proliferation and mast cell proliferation and promote upregulation of MHC class II surface expression on B cells, Epstein-Barr virus (ebv) IL-10 lacks these functions (Go, et al. (1990) *J. Exp. Med.* 172:1625-31; MacNeil, et al. (1990) *J. Immunol.* 145:4167-73; Vieira, et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:1172-6). Further examination has also revealed that the ability to inhibit IL-2 production by a CD4 T-cell clone is greatly reduced in comparison with that of the cIL-10s and that ebvIL-10 has an approximately 1,000-fold-lower affinity for the cellular IL-10R than hIL-

10 (Liu, et al. (1997) *J. Immunol.* 152:1821-9). In particular, ebvIL-10 has been shown to be effective in inhibiting CIA (Keravala, et al. (2006) *Arthritis Res. Ther.* 8:R91; Kim, et al. (2000) *J. Immunol.* 164:1576-81; Ma, et al. (1998) *J. Immunol.* 161:1516-24), suppressing autoimmune diabetes (Yang, et al. (2002) *J. Immunol.* 168:6479-85), ameliorating symptoms of necrotizing pancreatitis (Minter, et al. (2001) *J. Immunol.* 167:1053-9), and improving survival of sepsis (Oberholzer, et al. (2001) *Proc. Natl. Acad. Sci. USA* 98:11503-8). Exemplary viral IL-10 proteins of use in this invention include, e.g., Human herpes 5 IL-10/UL111A (GENBANK Accession No. AAR31656, AAF36285, or YP_081552), Human herpesvirus 5 latency associated cmvIL-10 (GENBANK Accession No. ACR49217), Rhesus cytomegalovirus UL111A/vIL-10 (GENBANK Accession No. AF200417), African green monkey cytomegalovirus vIL-10 (GENBANK Accession No. AAF63435, Baboon cytomegalovirus IL-10-like protein (GENBANK Accession No. AAF63436), Epstein-Barr virus protein BCRF1 (GENBANK Accession No. CAD53385 or P03180), Rhesus lymphocryptovirus BCFR1 (GENBANK Accession No. AAK95412), Baboon lymphocryptovirus IL-10 (GENBANK Accession No. AAF23949), Equine herpesvirus 2 IL-10 (GENBANK Accession No. AAB26148), Ovine herpesvirus 2 IL-10 (GENBANK Accession No. ABB22222), Koi herpesvirus IL-10 (GENBANK Accession No. YP_001096169), Orf virus IL-10 (GENBANK Accession No. AAC57332), *Pan paniscus* viral IL-10 homolog (GENBANK Accession No. XP_003804206), Sheeppox virus IL-10 (GENBANK Accession No. NP_659579), Goatpox virus IL-10 (GENBANK Accession No. YP_001293197) and Lumpy skin disease virus IL-10 (GENBANK Accession No. AAN02729). See also, Slobedman, et al. (2009) *J. Virol.* 83:9618-29.

[0023] Helminth C-type lectins, sharing sequence and structural similarity with mammalian immune cell lectins, have been identified from nematode parasites and shown to exert profound anti-inflammatory effects on the host immune system (Pritchard, et al. (1994) *Int. J. Parasitol.* 24:495-500; Holland, et al. (2000) *Eur. J. Immunol.* 30:1977-87; Balic, et al. (2004) *Eur. J. Immunol.* 34:3407-59). Exemplary helminth C-type lectins of use in this invention include, e.g., *Heligmosomoides polygyrus* C-type lectin-1 (GENBANK Accession No. FJ456978) and *Nippostrongylus brasiliensis* C-type lectin-2 (GENBANK Accession No. ACS37723).

[0024] Other anti-inflammatory and immunosuppressant proteins of use in the invention include, e.g., Glucocorticoid-induced leucine zipper (GILZ), a protein which interacts with and inhibits the function of NF κ B and AP-1 (Ayroldi & Riccardi (2009) *FASEB J.* 23:3649-3658); annexin AI (AnxA1) and its mimetic peptides (e.g., Ac2-26), which reduce the production of inflammatory mediators such as TNF- α , IL-1 β , IL-6, and NO (Perretti & D'Acquisto (2009) *Nat. Rev. Immunol.* 9:62-70; Girol, et al. (2013) *J. Immunol.* 190:5689-5701); IL-18 binding protein from vaccinia virus, cowpox virus or *Molluscum contagiosum* virus, which blocks the effects of IL-18 (Smith, et al. (2000) *J. Gen. Virol.* 81:1223-30); the vaccinia virus IFN- α binding protein (Colamonici, et al. (1995) *J. Biol. Chem.* 270:15974-8); and viral serpins (Chen, et al. (2011) *Methods Enzymol.* 499:301-29).

[0025] It is also to be understood that by any particular gene/protein, the invention encompasses the gene/protein and any obvious variants thereby, which may be allelic variants or other modifications, which maintain the immunosuppressive and/or anti-inflammatory activities. The

present invention also includes families of the gene/protein, which are related by sequence similarity or function.

[0026] A chimeric antigen receptor, also known as a CAR, artificial T cell receptor, chimeric T cell receptor, or chimeric immunoreceptor, of this invention is a fusion protein composed of at least one antigen targeting domain or recognition domain, a transmembrane region that anchors the antigen targeting domain to the cell surface, at least one signaling endodomain and an optional extracellular spacer/hinge domain between the antigen targeting domain or recognition domain and transmembrane region.

[0027] Antigen targeting or antigen recognition by CAR molecules can involve the use of a single chain variable fragment (scFv) that has been assembled from a monoclonal antibody. Alternatively, targeting moieties can include receptors such as NKG2D, NKp30 or dectin1 (Zhang & Sentman (2013) *J. Immunol.* 190:2455-63; Zhang, et al. (2012) *J. Immunol.* 189:2290-9); ligands (Altenschmidt, et al. (1996) *Clin. Cancer Res.* 2:1001-8; Muniappan, et al. (2000) *Cancer Gene Ther.* 7:128-134), peptides (Pameijer, et al. (2007) *Cancer Gene Ther.* 14:91-97), chimeric ligands (Davies, et al. (2012) *Mol. Med.* 18:565-576), receptor derivatives (Scholler, et al. (2012) *Sci. Translation. Med.* 4:Article IDS 132ra53; Zhang, et al. (2012) *J. Immunol.* 189:2290-9), and single domain antibodies (Sharifzadeh, et al. (2012) *Cancer Res.* 72:1844-52). When two or more antigen-specific targeting domains target at least two different antigens, the domains may be arranged in tandem and separated by linker sequences. In other embodiments, the extracellular spacer domain is optional.

[0028] Antigens or ligands, which can be used as targets at a site of inflammation or autoimmunity, include molecules

specific for inflammatory diseases or autoimmune diseases, as well as tissue- or cell-specific molecules. Examples of antigens or ligands specific for inflammatory diseases or autoimmune diseases, which may be targeted by the CARs of the invention include, but are not limited to, one or more of the antigens listed in Table 1.

TABLE 1

Disease or Condition	Target
Inflammatory bowel disease (IBD)	Antigen or ligand expressed in diseased colon or ileum
Rheumatoid arthritis	Antigen or ligand is an epitope of collagen or an antigen present in joints, e.g., Rheumatoid factor IgG complexes
Type I diabetes mellitus or autoimmune insulinitis	Pancreatic β cell antigen and/or insulin
Multiple sclerosis	A myelin basic protein (MBP) antigen or MOG-I or MOG2-2, proteolipid protein, myelin oligodendrocyte glycoprotein and/or a neuronal antigen
Autoimmune uveitis or uveoretinitis	S-antigen or another uveal or retinal antigen
Autoimmune orchitis	Testicular antigen
Autoimmune oophoritis	An ovarian antigen
Psoriasis	Keratinocyte antigen or another antigen present in dermis or epidermis
Vitiligo	Melanocyte antigen such as melanin or tyrosinase
Autoimmune prostatitis	Prostate antigen
Autoimmune hemolytic anemia	Rh blood group antigen
Autoimmune thrombocytopenic purpura	Platelet integrin GpIIb:IIIa
Goodpasture's syndrome	Noncollagenous domain of basement membrane collagen type IV
Pemphigus vulgaris	Epidermal cadherin
Graves' Disease	Thyroid Peroxidase and/or thyroid-stimulating hormone receptor
Hashimoto Thyroiditis	Thyroglobulin
Systemic Vasculitides	Myeloperoxidase
Crohn's Disease	Glycoprotein 2

Primary Biliary Cirrhosis	M2 or components thereof (e.g., BCOADC-E2, OGDC-E2, PDC-E2), Sp100, Gp210 and/or Nup62
Autoimmune Hepatitis II	Formiminotransferase Cyclodeaminase and/or Cytochrome P450 2D6
Celiac Disease	Tissue transglutaminase and/or gliadin
Thromboembolic Syndrome	β 2 Glycoprotein I
Systemic Vasculitides / Wegener's Granulomatosis	Proteinase 3

[0029] When targeting an inflammatory response, the target of the CAR of the invention can be an antigen expressed on or by a dendritic cell, macrophage/monocyte, granulocyte or eosinophil present at the inflammation site. Such antigens include, but are not limited to, AOC3 (VAP-1), CCL11 (eotaxin-1), CD20, CD3, CD4, CD5, IFN- γ , IgE, IgE Fc region, IL-1, IL-12, IL-23, IL-17, IL-17A, IL-22, IL-6, integrin α 4 β 7, LFA-1 (CD11a), OX-40L, and TNF- α .

[0030] When targeting particular tissues or cells, the target of the CAR of the invention can be a biomarker or cell ligand including, but not limited to, those listed in Table 2.

TABLE 2

Tissue	Biomarker/Cell Ligand
Lung ^a	FGFR2 Sprouty Homolog 2 (SPRY2) Stimulated by retinoic acid 6 (STRA6) CD68 Mucin 1 Bone morphogenetic Protein Receptor Type 1A
Neuron ^a	S-100 protein
Cardiovascular System	Caveolin 1
Liver ^a	SLCO1B1 CYP1A2 CYP3A4
Kidney ^a	Glomeruli (NPHS2)
Neurons (CNS) ^a	Myelin Basic Protein

	Myelin Oligodendrocyte Glycoprotein Proteolipid Protein
Glial cells (CNS) ^a	Aquaporin-4 (AQP4)
Heart ^a	β -adrenergic receptor
Pancreas/Islet Cells ^b	Retinoic acid early inducible (RAE)-1
minor salivary glands ^c	B7H6
Liver (bile duct epithelial cells) ^a	Carbonic anhydrase IX (CIAX)
CNS ^a	Mutant Superoxide dismutase (SOD)-1
Inflammatory tissues ^a	Autoimmune TCRs
Inflammatory tissues ^d	MHCI/II + Autoimmune peptide

^aScFv as antigen-specific targeting domain; ^bNKG2D as antigen-specific targeting domain; ^cNKp30 as antigen-specific targeting domain; ^dAutoimmune TCR as antigen-specific targeting domain.

[0031] Exemplary antigen-specific targeting domains of the instant CAR include, but are not limited to, Cam-3001, CD125, CD154, CD2, CD23 (IgE receptor), CD25 (a chain of IL-2 receptor), IL-6 receptor, rhuMAb β 7 and OX-40.

[0032] In addition to antigen-specific approaches, two "universal" CAR systems have been described. These generic CARs containing avidin (Urbanska, et al. (2012) *Cancer Res.* 72:1844-52) or antiluorescein isothiocyanate (FITC) scFv (Ang, et al. (2011) *Mol. Ther.* 19:abstract 353; Chmielewski, et al. (2004) *J. Immunol.* 173:7647-7653), enabling their use in conjunction with separate targeting moieties that have been biotinylated or conjugated to FITC, respectively.

[0033] In embodiments wherein the antigen targeting domain is a scFv, the scFv can be derived from the variable heavy chain (VH) and variable light chain (VL) regions of an antigen-specific mAb linked by a flexible linker. The scFv

retains the same specificity as the full antibody from which it was derived (Muniappan, et al. (2000) *Cancer Gene Ther.* 7:128-134). Various methods for preparing an scFv can be used including methods described in US 4,694,778; Bird, et al. (1988) *Science* 242:423-442; Ward, et al. (1989) *Nature* 334:544-54; and Skerra, et al. (1988) *Science* 242:1038-1041. In certain embodiments, the scFv is humanized or is a fully human scFv.

[0034] As indicated, the CAR of the invention may also have an extracellular spacer/hinge domain and a transmembrane region or domain. The transmembrane domain may be derived from a natural polypeptide, or may be artificially designed. The transmembrane domain derived from a natural polypeptide can be obtained from any membrane-binding or transmembrane protein. For example, a transmembrane domain of a T cell receptor α or β chain, a CD3 ζ chain, CD28, CD3 ϵ , CD45, CD4, CD5, CD7, CD8, CD9, CD16, CD22, CD33, CD37, CD64, CD80, CD86, CD134, CD137, ICOS, CD154, H2-Kb, Fc ϵ RI γ or a GITR can be used. See, e.g., Kahlon, et al. (2004) *Cancer Res.* 64:9160-9166; Schambach, et al. (2009) *Methods Mol. Biol.* 506:191-205; Jensen, et al. (1998) *Biol. Blood Marrow Transplant* 4:75-83; Patel, et al. (1999) *Gene Ther.* 6:412; Song, et al. (2012) *Blood* 119:696-706; Carpenito, et al. (2009) *Proc. Natl. Acad. Sci. USA* 106:3360-5; Hombach, et al. (2012) *Oncoimmunology* 1:458-66) and Geiger, et al. (2001) *Blood* 98:2364-71. The artificially designed transmembrane domain is a polypeptide mainly composed hydrophobic residues such as leucine and valine. It is preferable that a triplet of phenylalanine, tryptophan and valine is found at each end of the synthetic transmembrane domain. See, U.S. 7,052,906. In one embodiment, the transmembrane domain is composed of residues 153 to 180 of CD28 (GENBANK Accession No.

NP_006130). As another embodiment, the transmembrane domain is composed of residues 162 to 183 of a GITR (GENBANK Accession No. NP_004186).

[0035] In the CAR of the invention, a spacer or hinge domain can be arranged between the extracellular antigen targeting domain and the transmembrane domain, and/or between the intracellular signaling endodomain and the transmembrane domain. A spacer domain refers to any oligopeptide or polypeptide that serves to link the transmembrane domain with the antigen targeting domain and/or the transmembrane domain with the intracellular signaling endodomain. The spacer domain can be up to 300 amino acids, preferably 10 to 100 amino acids, 25 to 50 amino acids or 2 to 10 amino acids in length.

[0036] The spacer domain preferably has a sequence that promotes binding of a CAR with an antigen and enhances signaling in a cell. Examples of an amino acid that is expected to promote the binding include cysteine, a charged amino acid, and serine and threonine in a potential glycosylation site, and these amino acids can be used as an amino acid constituting the spacer domain.

[0037] As the spacer domain, all or a part of residues 118 to 178 of CD8 α (GENBANK Accession No. NP_001759.3), residues 135 to 195 of CD8 β (GENBANK Accession No. AAA35664), residues 315 to 396 of CD4 (GENBANK Accession No. NP_000607.1), or residues 137 to 152 of CD28 (GENBANK Accession No. NP_006130.1) can be used. Also, as the spacer domain, a part of a constant region of an antibody H chain or L chain (CH1 region or CL region) can be used. Further, the spacer domain may be an artificially synthesized sequence.

[0038] The intracellular signaling endodomain used in this invention is a molecule that can transmit a signal into a

cell when the extracellular antigen targeting domain present within the same molecule binds to (interacts with) an antigen. Natural T cell-activation is transmitted by two different kinds of cytoplasmic signaling endodomain, that is, a sequence for initiating antigen-dependent primary activation via a TCR complex (primary cytoplasmic signaling endodomain) and a sequence for acting antigen-independently to provide a secondary or costimulating signal (secondary cytoplasmic signaling endodomain or costimulatory endodomain). Therefore, while some embodiments embrace a CAR with only a primary cytoplasmic signaling endodomain, in other embodiments, a CAR of the invention includes a primary signaling endodomain and a secondary cytoplasmic signaling endodomain.

[0039] The primary cytoplasmic signaling endodomain regulates primary activation of a TCR complex. The primary cytoplasmic signaling sequence that stimulates the activation may include a signal transduction motif known as an immunoreceptor tyrosine-based activation motif (ITAM) (Asp/Glu)-Xaa-Xaa-Tyr*-Xaa-Xaa-(Ile/Leu)-Xaa₆₋₈-Tyr*-Xaa-Xaa-(Ile/Leu) (SEQ ID NO:2) (Reth, et al. (1989) *Nature* 338:383-384).

[0040] Examples of proteins having an ITAM that can be used in the present invention include CD3 ζ , FcR γ , FcR β , CD3 γ , CD3 δ , CD3 ϵ , CD5, CD22, CD79a, CD79b, and CD66d. Specifically, examples of the ITAM include residues 51 to 164 of CD3 ζ (GENBANK Accession No. NP_932170), residues 45 to 86 of Fc ϵ RI γ (GENBANK Accession No. NP_004097), residues 201 to 244 of Fc ϵ RI β (GENBANK Accession No. NP_000130), residues 139 to 182 of CD3 γ (GENBANK Accession No. NP_000064), residues 128 to 171 of CD3 δ (GENBANK Accession No. NP_000723), residues 153 to 207 of CD3 ϵ (GENBANK Accession No. NP_000724), residues 402 to 495 of CD5

(GENBANK Accession No. NP_055022), residues 707 to 847 of CD22 (GENBANK Accession No. NP_001762), residues 166 to 226 of CD79a (GENBANK Accession No. NP_001774), residues 182 to 229 of CD79b (GENBANK Accession No. NP_000611), and residues 177 to 252 of CD66d (GENBANK Accession No. NP_001806), and their variants having the same function as these peptides have. The referenced residues are based on amino acid sequence information from GENBANK and is based on the full length of the precursor (including a signal peptide sequence etc.) of each protein.

[0041] Examples of secondary cytoplasmic signaling endodomains or costimulatory endodomains that can be used in the present invention include sequences derived from CD2, CD4, CD5, CD8 α , CD8 β , CD28, CD134, CD137, ICOS, CD122, CD132, Dap10, Dap12, CD40 and CD154. Specific examples thereof include residues 236 to 351 of CD2 (GENBANK Accession No. NP_001758), residues 421 to 458 of CD4 (GENBANK Accession No. NP_000607), residues 402 to 495 of CD5 (GENBANK Accession No. NP_055022), residues 207 to 235 of CD8 α (GENBANK Accession No. NP_001759), residues 196 to 210 of CD8 β (GENBANK Accession No. AAA35664), residues 181 to 220 of CD28 (GENBANK Accession No. NP_006130), residues 214 to 255 of CD137 (4-1BB, GENBANK Accession No. NP_001552), residues 241 to 277 of CD134 (OX40, GENBANK Accession No. NP_003318), and residues 166 to 199 of ICOS (GENBANK Accession No. NP_036224), and their variants having the same function as these peptides have. In some embodiments, the costimulatory endodomain is from CD28, 4-1BB, OX40, ICOS, or a combination thereof.

[0042] While any suitable endodomain can be used in the CAR of the invention, in certain embodiments, the invention embraces the use of all or a part of the endodomains of CD28 and CD3 ζ . In specific embodiments, intracellular

signaling endodomains are those of the T cell antigen receptor complex, e.g., CD28, DAP10, CD137, CD2, which are used either alone or in a series with CD3 ζ . One or multiple endodomains may be employed, as so-called third generation CARs have at least 2 or 3 signaling domains fused together for additive or synergistic effect, for example.

[0043] In a CAR containing more than one intracellular endodomain, an oligopeptide linker or a polypeptide linker can be inserted between the intracellular endodomains to link the domains. Preferably, a linker having a length of 2 to 10 amino acids can be used. Particularly, a linker having a glycine-serine continuous sequence can be used.

[0044] In addition to the antigen targeting domain, optional extracellular spacer/hinge domain, transmembrane domain, and signaling endodomain, the CAR of the invention can also include a signal peptide sequence linked to the N-terminus of the CAR. Signal peptide sequences exist at the N-terminus of many secretory proteins and membrane proteins, and have a length of 15 to 30 amino acids. Since many of the protein molecules mentioned above have signal peptide sequences, these signal peptides can be used as a signal peptide for the CAR of this invention.

[0045] As will be appreciated by one of skill in the art, in some instances, a few amino acids at the ends of the antigen targeting domain can be deleted, usually not more than 10, more usually not more than 5 residues. Also, it may be desirable to introduce a small number of amino acids at the borders, usually not more than 10, more usually not more than 5 residues. The deletion or insertion of amino acids will usually be as a result of the needs of the construction, providing for convenient restriction sites, ease of manipulation, improvement in levels of expression, or the like. In addition, the substitute of one or more

amino acids with a different amino acid can occur for similar reasons, usually not substituting more than about five amino acids in any one domain.

[0046] For the purposes of this invention, "nucleic acids" refer to single or double stranded nucleic acid molecules, which are isolated and provided in the form of RNA, a complementary polynucleotide (cDNA), a genomic polynucleotide and/or a composite polynucleotide (e.g., a combination of the above). As used herein, the term "nucleic acid construct" refers to a nucleic acid molecule, which includes nucleic acids encoding a CAR and nucleic acids encoding an anti-inflammatory or immunosuppressant protein. In some embodiments, the nucleic acid construct is a linear naked molecule or a vector, e.g., a plasmid, a bacmid, a phagemid, a cosmid, a phage, a virus or an artificial chromosome.

[0047] In accordance with the present invention, the nucleic acid construct is transformed or introduced into an immune cell and is transcribed and translated to produce a product (e.g., a chimeric receptor, anti-inflammatory or immunosuppressant protein, and optionally a suicide protein). Thus, the nucleic acid construct further includes at least one promoter for directing transcription of the CAR and anti-inflammatory or immunosuppressant protein. According to some embodiments, nucleic acids encoding the CAR and anti-inflammatory or immunosuppressant protein are operably linked to at least one promoter sequence. A coding nucleic acid is "operably linked" to a regulatory sequence (e.g., promoter) if the regulatory sequence is capable of exerting a regulatory effect on the coding sequence linked thereto.

[0048] In the nucleic acid construct of the invention, at least one promoter directs transcription of the CAR and

anti-inflammatory or immunosuppressant protein. According to some embodiments, nucleic acids encoding the CAR and anti-inflammatory or immunosuppressant protein are independently expressed via different promoters, *i.e.*, nucleic acids encoding the CAR are operably linked to a first promoter and nucleic acids encoding the anti-inflammatory or immunosuppressant protein are operably linked to a second promoter, which may be the same or different than the first promoter. In accordance with particular embodiments, expression of the anti-inflammatory or immunosuppressant protein by the second promoter is enhanced by CAR signaling. In this respect, CAR binding to a ligand expressed preferentially at the site of tissue damage will activate expression of the anti-inflammatory or immunosuppressant protein in a location specific manner and dampen inflammation at that site. Examples of promoter elements that are responsive to CAR signaling include, but are not limited to, the nuclear factor of activated T-cells (NFAT)-responsive promoter (Zhang, et al. (2011) *Mol. Ther.* 19:751-759), viral LTR, EF1-alpha promoter, or a doxycycline-responsive promoter. While it is contemplated that a CAR and anti-inflammatory or immunosuppressant protein expressed via different promoters can be achieved using two independent nucleic acid constructs, in accordance with the present invention, it is preferable that the nucleic acids encoding the CAR and anti-inflammatory or immunosuppressant protein reside on a single nucleic acid construct.

[0049] According to certain embodiments of the invention, nucleic acids encoding the CAR and anti-inflammatory or immunosuppressant protein are co-expressed via a single promoter, *i.e.*, nucleic acids encoding the CAR and nucleic acids encoding the anti-inflammatory or immunosuppressant

protein are in tandem and operably linked to a single promoter. A coding nucleic acid is "operably linked" to a regulatory sequence (e.g., promoter) if the regulatory sequence is capable of exerting a regulatory effect on the coding sequence linked thereto. In other words, the promoter(s) of the invention is positioned so as to promote transcription of the messenger RNA from the DNA encoding the CAR and anti-inflammatory or immunosuppressant protein.

[0050] The promoter(s) of the invention can be of genomic origin or synthetically generated. A variety of promoters for use in T cells have been described in the art. For example, the CD4 promoter is disclosed by Marodon, et al. ((2003) *Blood* 101(9):3416-23). The promoter can be constitutive or inducible, where induction is associated with the specific cell type, a specific level of maturation, or drug (e.g., tetracycline or doxorubicin). Alternatively, a number of viral promoters are also suitable. Promoters of interest include the β -actin promoter, SV40 early and late promoters, immunoglobulin promoter, human cytomegalovirus promoter, retrovirus promoter, and the Friend spleen focus-forming virus promoter. The promoters may or may not be associated with enhancers, wherein the enhancers may be naturally associated with the particular promoter or associated with a different promoter.

[0051] The simultaneous or co-expression of a CAR and anti-inflammatory or immunosuppressant protein via a single promoter may be achieved by the use of an internal ribosomal entry site (IRES) or cis-acting hydrolase element. The term "internal ribosome entry site" or "IRES" defines a sequence motif that promotes attachment of ribosomes to that motif on internal mRNA sequences. Consequently, an mRNA containing an IRES sequence motif

results in two translational products, one initiating from the 5'-end of the mRNA and the other by an internal translation mechanism mediated by the IRES. A number of IRES have been described and can be used in the nucleic acid construct of this invention. See, e.g., US 8,192,984; WO 2010/119257; and US 2005/0112095.

[0052] A "cis-acting hydrolase element" or "CHYSEL" refers to a peptide sequence that causes a ribosome to release the growing polypeptide chain that it is being synthesizes without dissociation from the mRNA. In this respect, the ribosome continues translating and therefore produces a second polypeptide. Peptides such as the foot and mouth disease virus (FMDV) 2A sequence (GSGSRVTELLYRMKRAETYC PRPLLAIHPTEARHKQKIVAPVKQLLNFDLLKLAGDVESNPGP, SEQ ID NO:3), sea urchin (*Strongylocentrotus purpuratus*) 2A sequence (DGFCILYLLLILLMRSGDVETNPGP, SEQ ID NO:4); Sponge (*Amphimedon queenslandica*) 2A sequence (LLCFMLLLLLSGDVELNPGP, SEQ ID NO:5; or HHFMFLLLLL AGDIELNPGP, SEQ ID NO:6); acorn worm (*Saccoglossus kowalevskii*) (WFLVLLSFILSGDIEVNPGP, SEQ ID NO:7) 2A sequence; amphioxus (*Branchiostoma floridae*) (KNCAMYMLLSGDVETNPGP, SEQ ID NO:8; or MVISQLMLKLAGDVEENPGP, SEQ ID NO:9) 2A sequence; porcine teschovirus-1 (GSGATNFSLLKQAGDVEENPGP, SEQ ID NO:10) 2A sequence (P2A); *Thosea asigna* virus (GSGEGRGSLT TCGDVEENPGP, SEQ ID NO:11) 2A sequence (T2A); and equine rhinitis A virus (GSGQCTNYALLKLAGDVESNPGP, SEQ ID NO:12) 2A sequence are CHYSELS of use in this invention. In some embodiments, the 2A sequence is a naturally occurring or synthetic sequence that includes the 2A consensus sequence D-X-E-X-NPGP (SEQ ID NO:13), in which X is any amino acid residue. In preferred embodiments, a furin sequence is

included upstream of the 2A sequence to allow the proteins to be separated.

[0053] The sequence of the open reading frames encoding the CAR and anti-inflammatory or immunosuppressant protein can be obtained from a genomic DNA source, a cDNA source, or can be synthesized (e.g., via PCR), or combinations thereof. Depending upon the size of the genomic DNA and the number of introns, it may be desirable to use cDNA or a combination thereof as it is found that introns stabilize the mRNA or provide T cell-specific expression (Barthel & Goldfeld (2003) *J. Immunol.* 171(7):3612-9). Also, it may be further advantageous to use endogenous or exogenous non-coding regions to stabilize the mRNA.

[0054] For expression of a CAR or anti-inflammatory or immunosuppressant protein, the naturally occurring or endogenous transcriptional initiation region of the nucleic acid sequence encoding N-terminal component of the CAR or anti-inflammatory or immunosuppressant protein can be used to generate the CAR or anti-inflammatory or immunosuppressant protein in the target host. Alternatively, an exogenous transcriptional initiation region can be used which allows for constitutive or inducible expression, wherein expression can be controlled depending upon the target host, the level of expression desired, the nature of the target host, and the like.

[0055] The termination region(s) of the construct may be provided by the naturally occurring or endogenous transcriptional termination regions of the nucleic acids encoding the C-terminal component of the last gene. Alternatively, the termination region may be derived from a different source. For the most part, the source of the termination region is generally not considered to be critical to the expression of a recombinant protein and a

wide variety of termination regions can be employed without adversely affecting expression.

[0056] In some embodiments of the invention, a nucleic acid construct or cell harboring the nucleic acid construct includes a nucleic acid encoding a protein that is capable of triggering cell death or elimination. Examples of such proteins include suicide proteins such as thymidine kinase (TK) of the HSV virus (herpesvirus) type I (Bonini, et al. (1997) *Science* 276:1719-1724), a Fas-based "artificial suicide gene" (Thomis, et al. (2001) *Blood* 97:1249-1257), *E. coli* cytosine deaminase gene or caspase-9, which are activated by gancyclovir, AP1903, 5-fluorocytosine or a specific chemical inducer of dimerization (CID), respectively.

[0057] The nucleic acid encoding the protein for cell death or elimination is advantageously provided in the nucleic acid construct of the invention to allow for the opportunity to ablate the transduced immune cells in case of toxicity and to destroy the chimeric construct once the signs or symptoms of disease have been reduced or ameliorated. The use of suicide genes for eliminating transformed or transduced cells is described in the art. For example, Bonini, et al. ((1997) *Science* 276:1719-1724) teach that donor lymphocytes transduced with the HSV-TK suicide gene provide antitumor activity in patients for up to one year and elimination of the transduced cells is achieved using ganciclovir. Further, Gonzalez, et al. ((2004) *J. Gene Med.* 6:704-711) describe the targeting of neuroblastoma with cytotoxic T lymphocyte clones genetically modified to express a chimeric scFvFc:ζ immunoreceptor specific for an epitope on L1-CAM, wherein the construct further expresses the hygromycin thymidine

kinase (HyTK) suicide gene to eliminate the transgenic clones.

[0058] It is contemplated that the nucleic acid encoding the protein for cell death or elimination can be expressed from the same promoter as the CAR and/or anti-inflammatory or immunosuppressant protein or from a different promoter. Generally, however, nucleic acid encoding the protein for cell death or elimination, CAR and anti-inflammatory or immunosuppressant protein reside on the same construct or vector. Expression of the protein for cell death or elimination from the same promoter as the CAR and/or anti-inflammatory or immunosuppressant protein can be accomplished using the IRES or CHYSEL sequences described herein.

[0059] In certain embodiments of the invention, a nucleic acid construct or cell harboring the nucleic acid construct uses a detectable marker so that the cell that harbors the nucleic acid construct is identifiable, for example for qualitative and/or quantitative purposes. The detectable marker may be detectable by any suitable means in the art, including by flow cytometry, fluorescence, spectrophotometry, and so forth. An example of a detectable marker is one that encodes a nonfunctional gene product but that is still detectable by flow cytometry means, for example, or can be used to select transgenic cells by flow cytometry or magnetic selection. In addition to detection, the marker protein can be used as a means to eliminate the transduced cells *in vivo* via an antibody that recognizes the marker protein. Examples of marker proteins of use in cell elimination include, e.g., truncated CD19 (Tey, et al. (2007) *Biol. Blood Marrow Transplant* 13:913-24), the extracellular region of CD20 (Introna, et al. (2000) *Hum. Gene Ther.* 11:611-20; Griffioen, et al. (2009)

Haematologica 94:1316-20), and the extracellular region of EGFR (Terakura, et al. (2012) *Blood* 119:72-82). See also, Lang, et al. (2004) *Blood* 103:3982-5. Incorporation of these proteins into gene-modified T cells renders the cells susceptible to elimination by clinically used anti-CD19 antibodies, anti-CD20 antibodies, and anti-EGFR antibodies (e.g., cetuximab).

[0060] A nucleic acid construct according to the present invention can be produced by any means known in the art, though preferably it is produced using recombinant DNA techniques. Nucleic acids encoding the CAR can be prepared and assembled into a complete coding sequence by standard techniques of molecular cloning (genomic library screening, PCR, primer-assisted ligation, site-directed mutagenesis, etc.). Nucleic acids encoding the other moieties (e.g., anti-inflammatory or immunosuppressant protein, IRES or CHYSEL) may be similarly prepared. The resulting nucleic acids are preferably inserted into an expression vector and used to transform suitable mammalian host cells, preferably immune cells such as T lymphocyte cells.

[0061] The constructs and mammalian cells of this invention find application in subjects having or suspected of having an inflammatory condition, in particular a chronic inflammatory condition, or immune-mediated autoimmunity. Chronic inflammatory conditions and autoimmune diseases that can be treated using the immune cells and nucleic acid constructs of this invention include, for example, rheumatoid arthritis, reactive arthritis, multiple sclerosis, Type I diabetes mellitus or autoimmune insulinitis, systemic lupus erythematosus, autoimmune uveoretinitis, autoimmune vasculitis, bullous pemphigus, myasthenia gravis, autoimmune thyroiditis or Hashimoto's disease, Sjogren's syndrome, granulomatous orchitis,

autoimmune oophoritis, Inflammatory bowel disease, Crohn's disease, sarcoidosis, rheumatic carditis, ankylosing spondylitis, Grave's disease, Scleroderma, Amyotrophic Lateral Sclerosis, autoimmune hemolytic anemia, psoriasis, vitiligo, eczema, primary biliary cirrhosis, autoimmune prostatitis, Goodpasture's syndrome, autoimmune hepatitis II, celiac disease, ulcerative colitis, thromboembolic syndrome, systemic vasculitides/Wegener's granulomatosis, autoimmune thrombocytopenic purpura, arthritis deformans, Lyme disease arthritis, osteoarthritis, psoriatic arthritis, gout, fibromyalgia, Still's disease, chronic uveitis, chronic back or neck pain and sciatica, Addison's disease, Gaucher's disease, Huntington's disease, muscular dystrophy, cystic fibrosis and idiopathic pulmonary fibrosis. See, e.g., Paul, W.E. (1993) *Fundamental Immunology*, Third Edition, Raven Press, New York, Chapter 30, pp. 1033-1097; and Cohen, et al . (1994) *Autoimmune Disease Models, A Guidebook*, Academic Press, 1994.

[0062] Accordingly, the invention further relates to a method for treating inflammation, in particular chronic inflammation, or immune-mediated autoimmunity by delivering to a subject in need of treatment an immune cell of this invention. The step of delivering the immune cell to the subject generally involves introducing, e.g., via transduction, transposons or electroporation, a nucleic acid construct of the invention into an isolated immune cell (e.g., an autologous or third party-derived immune cell) and introducing into the subject the transduced/transformed immune cell, thereby effecting inflammatory or immunosuppressive responses in the subject to treat or ameliorate chronic inflammation or immune-mediated autoimmunity.

[0063] "Immune cell" as used herein refers to the cells of the mammalian immune system including but not limited to antigen presenting cells, B-cells, basophils, cytotoxic T-cells, dendritic cells, eosinophils, granulocytes, helper T-cells, leukocytes, lymphocytes, macrophages, mast cells, memory cells, monocytes, natural killer cells, neutrophils, phagocytes, plasma cells and T-cells (e.g., naive T cells, central memory T cells, effector memory T cells). In particular embodiments, the immune cell of the invention is a T cell. In certain embodiments, the T cell is a CD4+ T cell.

[0064] Suitable T cells that can be used include autologous T lymphocyte cells, third party-derived T cells, transformed tumor or xenogenic immunologic effector cells, tumor infiltrating lymphocytes, cytotoxic lymphocytes or other cells that are capable of producing effector proteins when activated. As is known to one of skill in the art, various methods are readily available for isolating these cells from a subject. For example, using cell surface marker expression or using commercially available kits (e.g., ISOCELL from Pierce, Rockford, IL).

[0065] It is contemplated that the nucleic acid construct can be introduced into the immune cells as naked DNA or in a suitable vector. Methods of stably transfecting immune cells by electroporation using naked DNA or capped mRNA are known in the art. See, e.g., US 6,410,319. Naked DNA generally refers to the DNA encoding the nucleic acid construct of the invention contained in an expression vector in proper orientation for expression.

[0066] Alternatively, a viral vector (e.g., a retroviral vector, adenoviral vector, adeno-associated viral vector, or lentiviral vector) can be used to introduce the nucleic acid construct of the invention into immune cells. Suitable

vectors for use in accordance with the method of the invention are non-replicating in the immune cells. A large number of vectors are known that are based on viruses, where the copy number of the virus maintained in the cell is low enough to maintain the viability of the cell. Illustrative vectors include the pFB-neo vectors (STRATAGENE), as well as vectors based on HIV, SV40, EBV, HSV or BPV.

[0067] Both lentiviruses and retroviruses have been widely used as gene transfer vectors, and they compose the vector system that is currently used in the majority of clinical gene therapy trials (Sinn, et al. (2005) *Gene Ther.* 12:1089-1098). However, the lentiviral vectors have some advantages because they mediate the efficient transduction of cells, can be used with both dividing and nondividing cells, result in long-term, stable transgene expression and appear to be less prone to gene silencing (Sinn, et al. (2005) *Gene Ther.* 12:1089-1098).

[0068] Nonviral gene transfer technologies have also been explored for gene therapy. One approach includes the electrotransfer of DNA plasmids using the Sleeping Beauty (SB) transposon/transposase system into primary human immune cells, which has been shown to provide efficient and stable CD19-specific CAR gene expression (Singh, et al. (2008) *Cancer Res.* 68:2961-71; Maiti, et al. (2013) *J. Immunother.* 36:112-123). An alternative non-viral approach that does not rely on transgene integration, which uses RNA electroporation, results in transient CAR expression, precluding effective T-cell persistence beyond a week (Zhao, et al. (2006) *Mol. Ther.* 13:151-159). The use of transient CAR immune cells, which require multiple injections to provide meaningful tumor responses, may reduce the destruction of normal tissues or prevent T cell

accumulations to levels that increase the risk of cytokine storms (Zhao, et al. (2010) *Cancer Res.* 70:9053-61). Moreover, mRNA CAR T cells have been shown to mediate antitumor activity in patients with advanced solid tumors (Beatty, et al. (2014) *Cancer Immunology Res.* 2:112-20).

[0069] Once it is established that the transfected or transduced immune cell is capable of expressing proteins of the nucleic acid construct with the desired regulation and at a desired level, the activity of the CAR and/or anti-inflammatory or immunosuppressant protein can be determined. Subsequently, the transduced immune cells are reintroduced or administered to the subject to activate anti-inflammatory or immunosuppressive responses in the subject.

[0070] To facilitate administration, the transduced immune cells according to the invention can be made into a pharmaceutical composition or made implant-appropriate for administration *in vivo*, with appropriate carriers or diluents, which further can be pharmaceutically acceptable. The means of making such a composition or an implant have been described in the art (see, for instance, Remington: *The Science and Practice of Pharmacy*, Lippincott Williams & Wilkins, 21st edition (2005). Where appropriate, the transduced immune cells can be formulated into a preparation in semisolid or liquid form, such as a capsule, solution, injection, inhalant, or aerosol, in the usual ways for their respective route of administration. Means known in the art can be used to prevent or minimize release and absorption of the composition until it reaches the target tissue or organ, or to ensure timed-release of the composition. Desirably, however, a pharmaceutically acceptable form is employed which does not ineffectuate the cells of the invention. Thus, desirably the transduced

immune cells can be made into a pharmaceutical composition containing a balanced salt solution, preferably Hanks' balanced salt solution, or normal saline. Additional examples of carriers or diluents include fats, oils, water, saline solutions, lipids, liposomes, resins, binders, fillers and the like, or combinations thereof. The composition may also include various antioxidants to retard oxidation of one or more component.

[0071] A pharmaceutical composition of the invention can be used alone or in combination with other well-established agents useful for treating inflammation or an autoimmune disease. Whether delivered alone or in combination with other agents, the pharmaceutical composition of the invention can be delivered via various routes and to various sites in a mammalian, particularly human, body to achieve a particular effect. One skilled in the art will recognize that, although more than one route can be used for administration, a particular route can provide a more immediate and more effective reaction than another route. For example, intradermal delivery may be advantageously used over inhalation for the treatment of psoriasis, vitiligo or eczema. Local or systemic delivery can be accomplished by administration comprising application or instillation of the formulation into body cavities, inhalation or insufflation of an aerosol, or by parenteral introduction, comprising intramuscular, intravenous, intraportal, intrahepatic, peritoneal, subcutaneous, or intradermal administration.

[0072] Although systemic (intravenous, IV) injection is favored in clinical applications because of its ease of administration several preclinical studies (Carpenito, et al. (2009) *Proc. Natl. Acad. Sci. USA* 106:3360-3365; Song, et al. (2011) *Cancer Res.* 71:4617-4627; Parente-Pereira, et

al. (2011) *J. Clin. Immunol.* 31:710-718) suggest that the regional (intratumoral, IT or intraperitoneal, IP) administration of T cells may provide optimal therapeutic effects, which may be in part due to increased T-cell trafficking. For example, it has been shown that CAR T cells remain at the site of inoculation with minimal systemic absorption when delivered via IP or IT routes (Parente-Pereira, et al. (2011) *J. Clin. Immunol.* 31:710-718). In contrast, after IV administration, CAR immune cells initially reach the lungs and then are redistributed to the spleen, liver, and lymph nodes. In addition, RNA CAR-electroporated immune cells may be particularly suitable for regional administration, due to the transient nature of the CAR expression on the immune cells (Zhao, et al. (2010) *Cancer Res.* 70:9053-9061). Furthermore, clinical studies have shown the feasibility and safety of both the intratumoral and intraperitoneal injection of immune cells (Canevari, et al. (1995) *J. Natl. Cancer Inst.* 87:1463-1469; Duval, et al. (2006) *Clin. Cancer Res.* 12:1229-123680). Overall, a local route of administration of the chimeric immune cells may provide the optimal therapeutic effect and decrease the potential for the "on-target, off-organ" toxicity discussed below.

[0073] A composition of the invention can be provided in unit dosage form wherein each dosage unit, e.g., an injection, contains a predetermined amount of the composition, alone or in appropriate combination with other active agents including conventional immunosuppressants and anti-inflammatory agents. The term unit dosage form, as used herein, refers to physically discrete units suitable as unitary dosages for human and animal subjects, each unit containing a predetermined quantity of the composition of the invention, alone or in combination with other active

agents, calculated in an amount sufficient to produce the desired effect, in association with a pharmaceutically acceptable diluent, carrier, or vehicle, where appropriate. The specifications for the novel unit dosage forms of the invention depend on the particular pharmacodynamics associated with the pharmaceutical composition in the particular subject.

[0074] Desirably an effective amount or sufficient number of the isolated transduced immune cells is present in the composition and introduced into the subject such that long-term, specific, anti-inflammatory or immunosuppressive responses are established to reduce or ameliorate one or more signs or symptoms associated with inflammation or autoimmunity than would otherwise result in the absence of such treatment.

[0075] Accordingly, the amount of transduced immune cells administered should take into account the route of administration and should be such that a sufficient number of the transduced immune cells will be introduced so as to achieve the desired therapeutic response. Furthermore, the amounts of each active agent included in the compositions described herein (e.g., the amount per each cell to be contacted or the amount per certain body weight) can vary in different applications. In general, the concentration of transduced immune cells desirably should be sufficient to provide in the subject being treated at least from about 1×10^6 to about 1×10^9 transduced immune cells, even more desirably, from about 1×10^7 to about 5×10^8 transduced immune cells, although any suitable amount can be utilized either above, e.g., greater than 5×10^8 cells, or below, e.g., less than 1×10^7 cells. The dosing schedule can be based on well-established cell-based therapies (see, e.g., Topalian & Rosenberg (1987) *Acta Haematol.* 78 Suppl 1:75-6; US

4,690,915) or an alternate continuous infusion strategy can be employed.

[0076] Any of the compositions described herein may be included in a kit. The kits will thus include, in suitable container means, cells or nucleic acid constructs or related reagents of the present invention. In some embodiments, the kit further includes an additional immunosuppressive or anti-inflammatory agent. Examples of currently available immunosuppressants and anti-inflammatory agents include, but are not limited to, cyclosporine A, cyclophosphamide, prednisone, tacrolimus (FK506), nonsteroidal anti-inflammatory agents such as aspirin, ibuprofen, diclofenac, etodolac, fenoprofen, flurbiprofen, naproxen, and oxaprozin. In certain embodiments, the additional agent may be combined with the nucleic acid construct(s) or cells of the invention or may be provided separately in the kit. In some embodiments, means of taking a sample from an individual and/or of assaying the sample may be provided in the kit. In certain embodiments the kit includes cells, buffers, cell media, vectors, primers, restriction enzymes, salts, and so forth, for example.

[0077] The components of the kits may be packaged either in aqueous media or in lyophilized form. The container means of the kits will generally include at least one vial, test tube, flask, bottle, syringe or other container means, into which a component may be placed, and preferably, suitably aliquoted. Where there is more than one component in the kit, the kit also will generally contain a second, third or other additional container into which the additional components may be separately placed. However, various combinations of components may be comprised in a vial. Such

containers may include injection or blow-molded plastic containers into which the desired vials are retained.

[0078] When the components of the kit are provided in one and/or more liquid solutions, the liquid solution is an aqueous solution, with a sterile aqueous solution being particularly preferred. The compositions may also be formulated into a syringeable composition. In which case, the container means may itself be a syringe, pipette, and/or other such like apparatus, from which the formulation may be applied to an infected area of the body, injected into an animal, and/or even applied to and/or mixed with the other components of the kit. However, the components of the kit may be provided as dried powder(s). When reagents and/or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container means.

[0079] The following non-limiting examples are provided to further illustrate the present invention.

Example 1: Isolation and Transduction of Treg Cells with a CAR Construct

[0080] A method to isolate and expand natural T-regs was developed using commonly available magnet sort systems (Liu, et al. (2006) *J. Exp. Med.* 203:1701-11). The isolated T cells were CD4⁺ CD25⁺ CD127⁻ Foxp3⁺ and had a suppressive phenotype characterized by their ability to suppress CD4⁺ T cell proliferation (Figure 1). These cells were isolated, expanded, and transduced with a CAR construct composed of nucleic acids encoding a CAR protein fused to a truncated CD19 via T2A. Notably, Foxp3 expression was not lost after transduction (Figure 2).

Example 2: Coexpression of a CAR and BCRF1

[0081] Nucleic acids encoding BCRF1 were inserted downstream of nucleic acids encoding a CAR and Foxp3 protein. Coding sequences for the CAR and Foxp3 were linked via CHYSEL peptide T2A and coding sequences for Foxp3 and BCRF1 were linked via P2A. The construct was transduced into an E86 packing cell line and BCRF1 was found to be actively secreted. ELISA analysis showed that high levels of human IL-10 were present in the E86 supernatants (~49000 pg/mL) compared to untransduced control cells (~30 pg/mL). It is expected that this construct will be viable in T cells and that BCRF1 will exert a suppressive affect on various lymphocytes.

What is claimed is:

1. An immune cell comprising nucleic acids encoding a chimeric antigen receptor and nucleic acids encoding at least one exogenous anti-inflammatory or immunosuppressant protein.

2. The immune cell of claim 1, wherein the cell is a T lymphocyte.

3. The immune cell of claim 1, wherein the chimeric antigen receptor comprises a single chain variable fragment.

4. The immune cell of claim 1, wherein the chimeric antigen receptor comprises an antigen targeting domain or recognition domain that binds an antigen or ligand at a site of inflammation or autoimmunity and an immune signaling receptor containing an immunoreceptor tyrosine-based activation motif.

5. The immune cell of claim 3 or 4, wherein the chimeric antigen receptor further comprises a costimulatory endodomain.

6. The immune cell of claim 1, wherein the anti-inflammatory or immunosuppressant protein comprises a cytokine, virokinin or a C-type lectin protein.

7. The immune cell of claim 6, wherein the virokinin is a viral IL-10 protein.

8. The immune cell of claim 6, wherein the C-type lectin protein is a helminth C-type lectin protein.

9. The immune cell of claim 1, wherein expression of the anti-inflammatory or immunosuppressant protein is regulated by a promoter that is dependent upon signaling from the chimeric antigen receptor.

10. A nucleic acid construct comprising nucleic acids encoding a chimeric antigen receptor and nucleic acids encoding at least one anti-inflammatory or immunosuppressant protein.

11. The nucleic acid construct of claim 10, wherein the construct comprises a vector.

12. The nucleic acid construct of claim 10, wherein the chimeric antigen receptor comprises a single chain variable fragment.

13. The nucleic acid construct of claim 10, wherein the chimeric antigen receptor comprises an antigen targeting domain or recognition domain that binds a ligand at a site of inflammation or autoimmunity and an immune signaling receptor containing an immunoreceptor tyrosine-based activation motif.

14. The nucleic acid construct of claim 12 or 13, wherein the chimeric antigen receptor further comprises a costimulatory endodomain.

15. The nucleic acid construct of claim 10, wherein the anti-inflammatory or immunosuppressant protein comprises a cytokine, virokinin or a C-type lectin protein.

16. The nucleic acid construct of claim 15, wherein the virokinin is a viral IL-10 protein.

17. The nucleic acid construct of claim 15, wherein the C-type lectin protein is a helminth C-type lectin protein.

18. The nucleic acid construct of claim 10, wherein expression of the anti-inflammatory or immunosuppressant protein is regulated by a promoter that is enhanced upon signaling from the chimeric antigen receptor.

19. A method for treating inflammation or immune-mediated autoimmunity comprising delivering to a subject in need of treatment an immune cell of claim 1 thereby treating the subject's inflammation or immune-mediated autoimmunity.

20. A kit comprising one or more immune cells of claim 1.

21. A kit comprising the nucleic acid construct of claim 10.

1/1




FIG. 1




FIG. 2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 16/53893

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/7088, A61K 35/17, A61K 48/00, A61P 37/06, A61P 37/08 (2016.01)

CPC - C07K 14/705, A61K 35/17, A61K 48/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8) - A61K 31/7088, A61K 35/17, A61K 48/00, A61P 37/06, A61P 37/08 (2016.01)

CPC - C07K 14/705, A61K 35/17, A61K 48/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched (keyword limited; terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Patbase, Google Scholar, Google Patents

Search terms: chimeric antigen receptor, chimeric, receptor, antigen receptor, cytokine, virokin, C-type lectin, lectin, calcium-dependent, anti-inflammatory, immunosuppressant, inflammation, IL10, IL-10, interleukin 10, viral, virus, helminth, encode, express, promoter, single

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---	US 2012/0029063 A1 (ZHANG et al.) 02 February 2012 (02.02.2012) para [0008], [0015], [0032], [0035], [0039], [0041], [0046], [0048], [0052], [0056]	1-4, 6, 9-13, 15, 18-21
Y		5, 7-8, 14, 16-17
Y	US 2014/0370017 A1 (THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA) 18 December 2014 (18.12.2014) abstract; para [0005], [0124]	5, 14
Y	LUCAS et al., Secreted immunomodulatory viral proteins as novel biotherapeutics. J Immunol. 15 October 2004, Vol 173, No 8, pp 4765-74. Especially p 4766, col 1, para 2; p 4766, col 2, para 1; p 4766, Figure 1	7, 16
Y	DITGEN et al., Harnessing the Helminth Secretome for Therapeutic Immunomodulators. Rinmed Res Int. 2014, Vol 2014, page 964350 (pp 1-15). Especially p 7, col 1, para 4	8, 17

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 November 2016

Date of mailing of the international search report

09 DEC 2016

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents

P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774