

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2014243796 B2**

(54) Title  
**Non-replicative transduction particles and transduction particle-based reporter systems**

(51) International Patent Classification(s)  
**A61K 35/76** (2006.01)      **C12N 15/74** (2006.01)  
**C12N 7/01** (2006.01)

(21) Application No: **2014243796**      (22) Date of Filing: **2014.03.13**

(87) WIPO No: **WO14/160418**

(30) Priority Data

(31) Number <b>61/897,040</b>	(32) Date <b>2013.10.29</b>	(33) Country <b>US</b>
<b>61/779,177</b>	<b>2013.03.13</b>	<b>US</b>
<b>61/939,126</b>	<b>2014.02.12</b>	<b>US</b>

(43) Publication Date: **2014.10.02**  
(44) Accepted Journal Date: **2018.07.19**

(71) Applicant(s)  
**GeneWEAVE Biosciences, Inc.**

(72) Inventor(s)  
**Rey, Diego Ariel;Deforest, Nikol;Cox, Heather;Shukla, Soni**

(74) Agent / Attorney  
**FB Rice Pty Ltd, L 23 44 Market St, Sydney, NSW, 2000, AU**

(56) Related Art  
**WO 2008131230 A1**  
**WO 9621007 A2**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2014/160418 A3

(43) International Publication Date  
2 October 2014 (02.10.2014)

WIPO | PCT

(51) International Patent Classification:  
*A61K 35/76 (2015.01) C12N 7/01 (2006.01)*  
*C12N 15/74 (2006.01)*

sciences, Inc., 983 University Avenue, Bldg. B200, Los Gatos, CA 95032 (US).

(21) International Application Number:  
PCT/US2014/026536

(74) Agents: YEE, Gene et al.; Fenwick & West LLP, 801 California Street, Mountai View, CA 94041 (US).

(22) International Filing Date:  
13 March 2014 (13.03.2014)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(25) Filing Language:  
English

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(26) Publication Language:  
English


(30) Priority Data:  
61/779,177 13 March 2013 (13.03.2013) US  
61/897,040 29 October 2013 (29.10.2013) US  
61/939,126 12 February 2014 (12.02.2014) US

(71) Applicant: GENEWEEVE BIOSCIENCES, INC.  
[US/US]; 983 University Avenue, Bldg. B200, Los Gatos, CA 95032 (US).

(72) Inventors: REY, Diego, Ariel; Geneweave Biosciences, Inc., 983 University Avenue, Bldg. B200, Los Gatos, CA 95032 (US). DEFOREST, Nikol; Geneweave Biosciences, Inc., 983 University Avenue, Bldg. B200, Los Gatos, CA 95032 (US). COX, Heather; Geneweave Biosciences, Inc., 983 University Avenue, Bldg. B200, Los Gatos, CA 95032 (US). SHUKLA, Soni; Geneweave Bios-

[Continued on next page]

(54) Title: NON-REPLICATIVE TRANSDUCTION PARTICLES AND TRANSDUCTION PARTICLE-BASED REPORTER SYSTEMS



Schematic of a P1-based packaging system (100)

FIGURE 1

(57) Abstract: Methods and systems are provided for packaging reporter nucleic acid molecules into non-replicative transduction particles for use as reporter molecules. The non-replicative transduction particles can be constructed from viruses and use viral transduction and replication systems. The reporter nucleic acid molecules include a reporter gene, such as a reporter molecule or selectable marker, for detecting target genes or cells. Methods and systems are provided for detection of cells and target nucleic acid molecules using the non-replicative transduction particles as reporter molecules.

**Declarations under Rule 4.17:**

— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

**Published:**

— *with international search report (Art. 21(3))*

— *with sequence listing part of description (Rule 5.2(a))*

**(88) Date of publication of the international search report:**

19 March 2015

## **TITLE**

**[0001]** Non-Replicative Transduction Particles and Transduction Particle-Based Reporter Systems

## **CROSS REFERENCE TO RELATED APPLICATIONS**

**[0002]** This application claims the benefit of U.S. Provisional Application No. 61/779,177, filed on March 13, 2013, U.S. Provisional Application No. 61/897,040, filed on Oct. 29, 2013, and U.S. Provisional Application No. 61/939,126, filed on Feb. 12, 2014, each of which is hereby incorporated in its entirety by reference.

## **SEQUENCE LISTING**

**[0002.1]** The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on April 10, 2014, is named 25938PCT\_CRF\_SequenceListing.txt and is 65,305 bytes in size.

## **BACKGROUND OF THE INVENTION**

### **Field of the invention**

**[0003]** The invention relates to methods and compositions for packaging and delivery of non-replicative transduction reporter molecules into cells for detecting target genes in cells.

### **Description of the Related Art**

**[0004]** A transduction particle refers to a virus capable of delivering a non-viral nucleic acid into a cell. Viral-based reporter systems have been used to detect the presence of cells and rely on the lysogenic phase of the virus to allow expression of a reporter molecule from the cell. These viral-based reporter systems use replication-competent transduction particles that express reporter molecules and cause a target cell to emit a detectable signal.

**[0005]** However, the lytic cycle of the virus has been shown to be deleterious to viral-based reporter assays. Carriere, C. *et al.*, *Conditionally replicating luciferase reporter phages: Improved sensitivity for rapid detection and assessment of drug susceptibility of Mycobacterium tuberculosis*. Journal of Clinical Microbiology, 1997. **35**(12): p. 3232-3239. Carrière *et al.* developed *M. tuberculosis*/bacillus Calmette-Guérin (BCG) luciferase reporter phages that have their lytic cycles suppressed at 30°C, but active at 37°C. Using this system, Carrière *et al.* have demonstrated the detection of BCG using phage reporters with a suppressed lytic cycle.

**[0006]** There are disadvantages, however, associated with suppressing but not eliminating the replication functions of the bacteriophage in bacteriophage-based reporter assays. First,

controlling replication functions of the bacteriophage imposes limiting assay conditions. For example, the lytic cycle of the reporter phage phAE40 used by Carrière *et al.* was repressed when the phage was used to infect cells at the non-permissive temperature of 30°C. This temperature requirement imposed limiting conditions on the reporter assay in that the optimum temperature for the target bacteria was 37°C. These limiting conditions hinder optimum assay performance.

[0007] Moreover, the replication functions of the virus are difficult to control. The replication of the virus should be suppressed during the use of the transduction particles as a reporter system. For example, the lytic activity of the reporter phage phAE40 reported by Carrière *et al.* was reduced but was not eliminated, resulting in a drop in luciferase signal in the assay. Carrière *et al.* highlighted possible causes for the resulting drop in reporter signal, such as intact phage-expressed genes and temperature limitations of the assay, all stemming from the fact that the lytic cycle of the phage reporter was not eliminated.

[0008] Reporter assays relying on the natural lysogenic cycle of phages can be expected to exhibit lytic activity sporadically. In addition, assays that rely on the lysogenic cycle of the phage can be prone to superinfection immunity from target cells already lysogenized with a similar phage, as well as naturally occurring host restriction systems that target incoming virus nucleic acid, thus limiting the host range of these reporter phages.

[0009] In other examples, transduction particle production systems are designed to package exogenous nucleic acid molecules, but the transduction particle often contains a combination of exogenous nucleic acid molecules and native progeny virus nucleic acid molecules. The native virus can exhibit lytic activity that is a hindrance to assay performance, and the lytic activity of the virus must be eliminated in order to purify transduction particles. However, this purification is generally not possible. In U.S. 2009/0155768 A, entitled *Reporter Plasmid Packaging System for Detection of Bacteria*, Scholl *et al.* describes the development of such a transduction particle system. The product of the system is a combination of reporter transduction particles and native bacteriophage (Figure 8 in the reference). Although the authors indicate that the transduction particle and native bacteriophage can be separated by ultracentrifugation, this separation is only possible in a system where the transduction particle and the native virus exhibit different densities that would allow separation by ultracentrifugation. While this characteristic is exhibited by the bacteriophage T7-based packaging system described in the reference, this is not a characteristic that is generally applicable for other virus systems. It is common for viral packaging machinery to exhibit headful packaging that would result in native virus and transduction particles to exhibit indistinguishable densities that cannot be separated by ultracentrifugation. Virus packaging systems also rely on a minimum amount of packaging as a requirement for proper virus structural assembly that results in native virus and transduction particles with indistinguishable densities.

[0010] Thus, there is a need for non-replicative transduction particles that do not suffer from the deleterious effects from lytic functions of the virus and the possibility of being limited by

superinfection immunity and host restriction mechanisms that target virus nucleic acid molecules and viral functions, all of which can limit the performance of the reporter assay by increasing limits of detection and resulting in false negative results.

**[0011]** Even where transduction particles have been engineered, methods for using the transduction particles to detect and report the presence of target nucleic acid molecules in cells have limitations. Some methods require disruption of the cell and cumbersome techniques to isolate and detect transcripts in the lysate. Detection methods include using labeled probes such as antibodies, aptamers, or nucleic acid probes. Labeled probes directed to a target gene can result in non-specific binding to unintended targets or generate signals that have a high signal-to-noise ratio. Therefore, there is a need for specific, effective and accurate methods for detection and reporting of endogenous nucleic acid molecules in cells.

**[0012]** Accordingly, methods and systems are needed for generating non-replicative transduction particles that allow packaging and expression of reporter molecules in cells, while eliminating replication-competent progeny virus. Effective and accurate methods for detecting molecules in cells using the expressed reporter molecules are also needed.

**[0012A]** Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed before the priority date of each of the appended claims.

**[0012B]** Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

## **SUMMARY OF THE INVENTION**

**[0013]** Disclosed herein is a bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising: a lysogenized bacteriophage genome comprising a first bacteriophage gene comprising a deletion of a packaging initiation site sequence of said first bacteriophage gene that prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage gene comprising a second packaging initiation site sequence that facilitates the packaging a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said

second bacteriophage gene encodes a protein, wherein said replica of said reporter nucleic acid molecule plasmid forms a replicon for packaging into said non-replicative transduction particle.

**[0013A]** Also disclosed herein is a bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising: a lysogenized bacteriophage genome lacking a bacteriophage gene encoding a packaging initiation site sequence, wherein deletion of said bacteriophage gene prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage gene, wherein said second bacteriophage gene encodes a packaging initiation site sequence and facilitates the packaging a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said second bacteriophage gene is capable of expressing a protein that is encoded by said gene, wherein said replica of said reporter nucleic acid molecule plasmid forms a replicon amenable to packaging into said non-replicative transduction particle.

**[0014]** In some embodiments, the reporter nucleic acid molecule plasmid is operatively linked to a promoter. In another embodiment, the promoter is selected for contributing to reactivity of a reporter molecule expressed from said reporter nucleic acid molecule plasmid in said bacterial cell. In one embodiment, the reporter nucleic acid molecule plasmid comprises an origin of replication. In yet another embodiment, the replicon comprises a concatamer amenable to packaging into said non-replicative transduction particle.

**[0015]** In an embodiment, the first and said second bacteriophage genes each comprises a pacA gene of the Enterobacteriaceae bacteriophage P1 and comprises said packaging initiation site sequence. In one embodiment, the second bacteriophage gene comprises the sequence of SEQ ID NO:9. In another embodiment, the replicon is the Enterobacteriaceae bacteriophage P1 lytic replicon. In certain embodiments, the replicon comprises a C1 repressor-controlled P53 promoter, a promoter P53 antisense, a repL gene, and an in-frame deletion of a kilA gene. In one embodiment, the replicon comprises of the sequence of SEQ ID NO:3.

**[0016]** In yet another embodiment, the first and said second bacteriophage genes each comprises a small terminase (terS) gene comprising said packaging initiation site sequence. In one embodiment, the terS gene is a *S. aureus* bacteriophage φ11 or φ80α terS gene.

**[0017]** In another embodiment, the replicon is derived from a *S. aureus* pT181 plasmid origin of replication. In yet another embodiment, the replicon comprises the sequence of SEQ ID NO:5. In some embodiments, the packaging initiation site sequence of said second bacteriophage gene comprises a pac-site. In other embodiments, the pac-site of said second bacteriophage gene comprises the sequence of SEQ ID NO:7. In one aspect, the packaging initiation site sequence of said second bacteriophage gene comprises a cos-site. In another aspect, the packaging initiation site sequence of said second bacteriophage gene comprises a concatamer junction.

**[0018]** In another aspect, a plasmid comprises said reporter nucleic acid molecule plasmid. In one aspect, the second bacteriophage gene is operatively linked to a promoter. In another embodiment, the promoter is an inducible promoter or a constitutive promoter. In one embodiment, the bacteriophage comprises the Enterobacteriaceae bacteriophage P1. In yet another embodiment, the bacteriophage comprises a *S. aureus* bacteriophage φ80α or a bacteriophage φ11. In one aspect, the bacterial cell comprises an *E. coli* cell. In another aspect, the bacterial cell comprises an *S. aureus* cell. In yet another embodiment, the bacterial cell comprises a Gram-negative cell. In other embodiments, the bacterial cell comprises a Gram-positive cell.

**[0019]** In another aspect, the reporter nucleic acid molecule plasmid comprises a reporter gene. In one aspect, the reporter gene encodes a detectable and/or a selectable marker. In certain aspects, the reporter gene is selected from the group consisting of enzymes mediating luminescence reactions (luxA, luxB, luxAB, luc, ruc, nluc), enzymes mediating colorimetric reactions (lacZ, HRP), fluorescent proteins (GFP, eGFP, YFP, RFP, CFP, BFP, mCherry, near-infrared fluorescent proteins), affinity peptides (His-tag, 3X-FLAG), and selectable markers (ampC, tet(M), CAT, erm). In another aspect, the reporter nucleic acid molecule plasmid comprises an aptamer. In yet another aspect, the reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said reporter nucleic acid molecule plasmid.

**[0020]** In one embodiment, the nucleic acid transcript sequence is complementary to a cellular transcript. In another embodiment, the nucleic acid transcript sequence comprises a cis-repressing sequence. In yet another embodiment, the replica of said reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said replica of said reporter nucleic acid molecule plasmid, wherein the nucleic acid transcript sequence is complementary to a cellular transcript and wherein said nucleic acid transcript sequence comprises a cis-repressing sequence.

**[0021]** In some embodiments, the method for packaging a reporter nucleic acid molecule plasmid into a non-replicative transduction particle, comprising providing conditions to said bacterial cell packaging system described herein that induce a lytic phase of said bacteriophage to produce non-replicative transduction particles packaged with said reporter nucleic acid molecule plasmid; and isolating said non-replicative transduction particle comprising said reporter nucleic acid molecule plasmid. In one embodiment, the non-replicative transduction particle does not contain a replicated bacteriophage genome. In another embodiment, induction of said lytic phase triggers excision of said genomic island nucleic acid molecule from said genome of said bacterial cell.

**[0022]** In another embodiment, the composition comprising said non-replicative transduction particle comprising a replica of said reporter nucleic acid molecule plasmid produced from the method described herein.

**[0023]** The invention also provides a bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising: a lysogenized bacteriophage genome comprising a first bacteriophage packaging initiation site sequence, wherein said first bacteriophage packaging initiation site sequence comprises a mutation that prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage packaging initiation site sequence, wherein said second bacteriophage packaging initiation site sequence lacks said mutation and facilitates the packaging of a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said replica of said reporter nucleic acid molecule plasmid forms a replicon for packaging into said non-replicative transduction particle.

**[0024]** In one embodiment, the reporter nucleic acid molecule is operatively linked to a promoter. In another embodiment, the promoter is selected for contributing to reactivity of a reporter molecule expressed from said reporter nucleic acid molecule in said bacterial cell. In yet another embodiment, the reporter nucleic acid molecule comprises an origin of replication. In one embodiment, the replicon comprises a concatamer amenable to packaging into said non-replicative transduction particle. In another aspect, the first and said second bacteriophage packaging initiation site sequences each comprise a packaging initiation site sequence from a small terminase gene. In one aspect, the first and said second bacteriophage packaging initiation site sequences each comprise a pac-site sequence from a pacA gene of the Enterobacteriaceae bacteriophage P1. In another aspect, the first bacteriophage packaging

initiation site sequence comprises SEQ ID NO:2. In yet another aspect, the second bacteriophage packaging initiation site sequence comprises SEQ ID NO:1. In one embodiment, the replicon comprises an Enterobacteriaceae bacteriophage P1 lytic replicon. In another embodiment, the replicon comprises a C1 repressor-controlled P53 promoter, a promoter P53 antisense, a repL gene, and an in-frame deletion of a kilA gene. In another aspect, the replicon comprises the sequence of SEQ ID NO:3. In certain aspects, the first and said second bacteriophage packaging initiation site sequences each comprise a pac-site sequence from a small terminase (terS) gene of an *S. aureus* bacteriophage φ11 or φ80α. In another aspect, the replicon is derived from a *S. aureus* pT181 plasmid origin of replication. In yet another aspect, the replicon comprises the sequence of SEQ ID NO:5. In one aspect, the first bacteriophage packaging initiation site sequence comprises the sequence of SEQ ID NO:2. In some embodiments, the second bacteriophage packaging initiation site sequence comprises the sequence of SEQ ID NO:1. In other embodiments, the packaging initiation site sequence comprises a pac-site. In another embodiment, the packaging initiation site sequence comprises a cos-site. In yet another embodiment, the packaging initiation site sequence comprises a concatamer junction. In some embodiments, the mutation in said first bacteriophage packaging initiation site sequence comprises a silent mutation. In another embodiment, the mutation in said first bacteriophage packaging initiation site sequence prevents cleavage of said packaging initiation sequence. In another embodiment, a plasmid comprises said reporter nucleic acid molecule plasmid. In one embodiment, the bacteriophage comprises Enterobacteriaceae bacteriophage P1.

**[0025]** In another embodiment, the bacteriophage comprises the *S. aureus* bacteriophage φ11 or φ80α. In one embodiment, the bacterial cell comprises an *E. coli* cell. In another embodiment, the bacterial cell comprises an *S. aureus* cell. In some embodiments, the bacterial cell comprises a Gram-negative bacterial cell. In one aspect, the bacterial cell comprises a Gram-positive bacterial cell. In another aspect, the reporter nucleic acid molecule plasmid comprises a reporter gene. In yet another aspect, the reporter gene encodes a detectable marker and/or a selectable marker.

**[0026]** In other aspects, the reporter gene is selected from the group consisting of: genes encoding enzymes mediating luminescence reactions (luxA, luxB, luxAB, luc, ruc, nluc), genes encoding enzymes mediating colorimetric reactions (lacZ, HRP), genes encoding fluorescent proteins (GFP, eGFP, YFP, RFP, CFP, BFP, mCherry, near-infrared fluorescent proteins), nucleic acid molecules encoding affinity peptides (His-tag, 3X-FLAG), and genes encoding selectable markers (ampC, tet(M), CAT, erm). In another aspect, the reporter

nucleic acid molecule plasmid comprises an aptamer. In other aspects, the replicon is packaged into said non-replicative transduction particle by bacteriophage packaging machinery. In some embodiments, the reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said reporter nucleic acid molecule plasmid. In another embodiment, the nucleic acid transcript sequence is complementary to a cellular transcript.

**[0027]** In one aspect, the nucleic acid transcript sequence comprises a cis-repressing sequence. In another aspect, the replica of said reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said replica of said reporter nucleic acid molecule plasmid, wherein said nucleic acid transcript sequence is complementary to a cellular transcript, and wherein said nucleic acid transcript sequence comprises a cis-repressing sequence.

**[0028]** In certain aspects, the method for packaging a reporter nucleic acid molecule plasmid into a non-replicative transduction particle, comprising: providing conditions to said bacterial cell packaging system described herein that induce a lytic phase of said bacteriophage to produce non-replicative transduction particles packaged with said reporter nucleic acid molecule plasmid; and isolating said non-replicative transduction particle comprising said reporter nucleic acid molecule plasmid.

**[0029]** In other aspects, the non-replicative transduction particle does not contain a replicated bacteriophage genome. In one aspect, the induction of said lytic phase triggers excision of said genomic island nucleic acid molecule from said genome of said bacterial cell.

**[0030]** In another aspect, the invention comprises a composition comprising said non-replicative transduction particle comprising a replica of said reporter nucleic acid molecule plasmid produced from said method described herein.

**[0031]** In one aspect, the invention includes a bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising: a lysogenized bacteriophage genome comprising a first bacteriophage gene comprising a deletion of a packaging initiation site sequence of said first bacteriophage gene that prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage gene comprising a second packaging initiation site sequence that facilitates the packaging a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said second bacteriophage gene encodes a protein, wherein said

replica of said reporter nucleic acid molecule plasmid forms a replicon for packaging into said non-replicative transduction particle.

**[0032]** In another aspect, the reporter nucleic acid molecule plasmid is operatively linked to a promoter. In one aspect, the promoter is selected for contributing to reactivity of a reporter molecule expressed from said reporter nucleic acid molecule plasmid in said bacterial cell. In certain aspects, the reporter nucleic acid comprises an origin of replication. In another aspect, the replicon comprises a concatamer amenable to packaging into said non-replicative transduction particle. In one aspect, the first and said second bacteriophage genes each comprises a pacA gene of the Enterobacteriaceae bacteriophage P1 and comprise said packaging initiation site sequence. In another aspect, the first bacteriophage gene comprises the sequence of SEQ ID NO:6. In certain aspects, the second bacteriophage gene comprises the sequence SEQ ID NO:7. In one aspect, the replicon comprises an Enterobacteriaceae bacteriophage P1 lytic replicon. In yet another aspect, the replicon comprises a C1 repressor-controlled P53 promoter, a promoter P53 antisense, a repL gene, and an in-frame deletion of a kilA gene. In another aspect, the replicon comprises the sequence of SEQ ID NO:3. In other aspects, the first and said second bacteriophage genes each comprises a small terminase (terS) gene comprising said packaging initiation site sequence. In one aspect, the terS gene is a *S. aureus* bacteriophage φ11 or φ80α terS gene. In another aspect, the first bacteriophage gene comprises the sequence of SEQ ID NO:8. In yet another aspect, the second bacteriophage gene comprises the sequence of SEQ ID NO:9. In one aspect, the replicon is derived from a *S. aureus* pT181 plasmid origin of replication. In one embodiment, the replicon comprises the sequence of SEQ ID NO:5. In another embodiment, the packaging initiation site sequence of said second bacteriophage gene comprises a pac-site. In yet another embodiment, the packaging initiation site sequence of said second bacteriophage gene comprises a cos-site.

**[0033]** In certain embodiments, the packaging initiation site sequence of said second bacteriophage gene comprises a concatamer junction. In one embodiment, a plasmid comprises said reporter nucleic acid molecule plasmid. In another embodiment, the second bacteriophage gene is operatively linked to a promoter. In yet another embodiment, the promoter is an inducible promoter or a constitutive promoter. In certain embodiments, the bacteriophage comprises the Enterobacteriaceae bacteriophage P1. In one embodiment, the bacteriophage comprises the *S. aureus* bacteriophage φ80α or bacteriophage φ11. In other embodiments, the bacterial cell comprises an *E. coli* cell. In another embodiment, the bacterial cell comprises an *S. aureus* cell. In one embodiment, the bacterial cell comprises a Gram-negative cell. In another embodiment, the bacterial cell comprises a Gram-positive cell.

**[0034]** In another aspect, the reporter nucleic acid molecule plasmid comprises a reporter gene. In one aspect, the reporter gene encodes a detectable and/or a selectable marker. In another aspect, the reporter gene is selected from the group consisting of genes encoding enzymes mediating luminescence reactions (luxA, luxB, luxAB, luc, ruc, nluc), genes encoding enzymes mediating colorimetric reactions (lacZ, HRP), genes encoding fluorescent proteins (GFP, eGFP, YFP, RFP, CFP, BFP, mCherry, near-infrared fluorescent proteins), nucleic acid molecules encoding affinity peptides (His-tag, 3X-FLAG), and genes encoding selectable markers (ampC, tet(M), CAT, erm). In one embodiment, the reporter nucleic acid molecule plasmid comprises an aptamer. In another embodiment, the replicon is packaged into said non-replicative transduction particle by bacteriophage packaging machinery. In yet another embodiment, the reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said reporter nucleic acid molecule plasmid. In one embodiment, the nucleic acid transcript sequence is complementary to a cellular transcript. In another embodiment, the nucleic acid transcript sequence comprises a cis-repressing sequence. In certain embodiments, the replica of said reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said replica of said reporter nucleic acid molecule plasmid, wherein said nucleic acid transcript sequence is complementary to a cellular transcript and wherein said nucleic acid transcript sequence comprises a cis-repressing sequence.

**[0035]** The invention includes a method for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, comprising: providing conditions to said bacterial cell packaging system as described herein that induce a lytic phase of said bacteriophage to produce non-replicative transduction particles packaged with said reporter nucleic acid molecule plasmid; and isolating said non-replicative transduction particle comprising said reporter nucleic acid molecule. In one embodiment, the non-replicative transduction particle does not contain a replicated bacteriophage genome. In another embodiment, the induction of said lytic phase triggers excision of said genomic island nucleic acid molecule from said genome of said bacterial cell.

**[0036]** In some aspects, the invention includes a composition comprising said non-replicative transduction particle comprising a replica of said reporter nucleic acid molecule plasmid produced from said method described herein.

**[0037]** In another aspect, the invention includes a bacterial cell packaging system for packaging a reporter nucleic acid molecule plasmid into a non-replicative transduction

particle, said bacterial cell comprising: a lysogenized bacteriophage genome lacking a packaging gene and comprising genes that encode proteins that form said non-replicative transduction particle; and a genomic island nucleic acid molecule comprising a reporter nucleic acid molecule plasmid and a packaging gene. In one aspect, the packaging gene comprises a small terminase (terS) gene. terS gene comprises a *S. aureus* bacteriophage φ80α terS gene or a bacteriophage φ11 terS gene.

**[0038]** In one aspect, the terS gene comprises the sequence of SEQ ID NO:9. In another aspect, the genomic island nucleic acid molecule comprises a SaPIbov2 genomic island nucleic acid molecule. In yet another aspect, the genomic island nucleic acid molecule is selected from the group consisting of a SaPI, a SaPII, a SaPI2, a SaPIbov1 and a SaPIbov2 genomic island nucleic acid molecule. In another embodiment, the reporter nucleic acid molecule plasmid is operatively linked to a promoter. In yet another embodiment, the reporter nucleic acid molecule plasmid comprises an origin of replication. In some embodiments, the bacteriophage comprises a *S. aureus* bacteriophage φ80α or bacteriophage φ11. In other embodiments, the bacterial cell comprises an *S. aureus* cell. In one embodiment, the genomic island nucleic acid molecule comprises an integrase gene and wherein said integrase gene encodes an integrase protein for excising and integrating said genomic island nucleic acid molecule out of and into a bacterial genome of said bacterial cell. In another embodiment, the integrase gene comprises the sequence of SEQ ID NO:10. In yet another embodiment, the genomic island nucleic acid molecule is integrated into a bacterial genome of said bacterial cell.

**[0039]** In certain aspects, the genomic island nucleic acid molecule can be replicated and forms molecule replicon that is amenable to packaging by the bacteriophage packaging machinery in said bacterial cell. In another aspect, the nucleic acid molecule forms a concatamer. In yet another aspect, the replicated genomic island nucleic acid molecule is capable of being packaged into said non-replicative transduction particle. In certain aspects, the packaging gene comprises a pac site sequence. In another aspect, the packaging gene comprises a cos-site sequence. In yet another embodiment, the packaging gene comprises a concatamer junction.

**[0040]** In other embodiments, the reporter nucleic acid molecule plasmid comprises a reporter gene. In some embodiments, the reporter gene encodes a selectable marker and/or a selectable marker. In another embodiment, the reporter gene is selected from the group consisting of enzymes mediating luminescence reactions (luxA, luxB, luxAB, luc, ruc, nluc), enzymes mediating colorimetric reactions (lacZ, HRP), fluorescent proteins (GFP, eGFP,

YFP, RFP, CFP, BFP, mCherry, near-infrared fluorescent proteins), affinity peptides (His-tag, 3X-FLAG), and selectable markers (ampC, tet(M), CAT, erm). In certain embodiments, the reporter nucleic acid molecule plasmid comprises an aptamer. In other embodiments, the genomic island nucleic acid molecule lacks an integrase gene. In another embodiment, the invention includes a bacterial gene comprising an integrase gene operatively linked to a promoter and wherein said integrase gene encodes an integrase protein for excising and integrating said genomic island nucleic acid molecule out of and into a bacterial genome of said bacterial cell. In one embodiment, the reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said reporter nucleic acid molecule plasmid. In other embodiments, the nucleic acid transcript sequence is complementary to a cellular transcript. In yet other embodiments, the nucleic acid transcript sequence comprises a cis-repressing sequence. In another embodiment, the replica of said reporter nucleic acid molecule plasmid comprises a nucleic acid transcript sequence that is complementary to a second sequence in said replica of said reporter nucleic acid molecule plasmid. In other embodiments, the nucleic acid transcript sequence is complementary to a cellular transcript. In other embodiments, the nucleic acid transcript sequence comprises a cis-repressing sequence.

**[0041]** The invention includes a method for packaging a reporter nucleic acid molecule plasmid into a non-replicative transduction particle, comprising: providing conditions to said bacterial cell packaging system as described herein that induce a lytic phase of said bacteriophage to produce non-replicative transduction particles packaged with said reporter nucleic acid molecule plasmid; and isolating said non-replicative transduction particle comprising said reporter nucleic acid molecule plasmid. In some embodiments, the non-replicative transduction particle does not contain a replicated bacteriophage genome. In one embodiment, the induction of said lytic phase triggers excision of said genomic island nucleic acid molecule from said genome of said bacterial cell.

**[0042]** In another embodiment, the invention includes a composition comprising said non-replicative transduction particle comprising a replica of said reporter nucleic acid molecule plasmid produced from said method described herein.

**[0043]** The invention also includes a method for detecting a presence or an absence of a bacterial cell in a sample, comprising: introducing into a sample a non-replicative transduction particle comprising a reporter gene encoding a reporter molecule and lacking a bacteriophage genome under conditions such that said non-replicative transduction particle can transduce said bacterial cell and wherein said reporter gene can be expressed in said

bacterial cell; providing conditions for activation of said reporter molecule; and detecting for a presence or an absence of a reporter signal transmitted from said expressed reporter molecule, wherein a presence of said reporter signal correctly indicates said presence of said bacterial cell.

**[0044]** In one embodiment, the method achieves at least 80% specificity of detection with reference to a standard, at least 90% specificity of detection with reference to a standard, or at least 95% specificity of detection with reference to a standard. In another embodiment, the method achieves at least 80% sensitivity of detection with reference to a standard, at least 85% sensitivity of detection with reference to a standard, or at least 90% sensitivity of detection with reference to a standard, or at least 95% sensitivity of detection with reference to a standard. In yet another embodiment, the method achieves at least 95% specificity of detection and at least 90% sensitivity of detection with reference to a standard. In another embodiment, the standard is a Gold standard. In yet another embodiment, the bacterial cell comprises a Methicillin Resistant *Staphylococcus aureus* (MRSA) cell. In other embodiments, the bacterial cell comprises a Methicillin Sensitive *Staphylococcus aureus* (MSSA) cell.

**[0045]** In another embodiment, the reporter gene encodes a detectable or selectable marker. In one embodiment, the reporter gene is selected from the group consisting of genes encoding enzymes mediating luminescence reactions (luxA, luxB, luxAB, luc, ruc, nluc), genes encoding enzymes mediating colorimetric reactions (lacZ, HRP), genes encoding fluorescent proteins (GFP, eGFP, YFP, RFP, CFP, BFP, mCherry, near-infrared fluorescent proteins), nucleic acid molecules encoding affinity peptides (His-tag, 3X-FLAG), and genes encoding selectable markers (ampC, tet(M), CAT, erm). In one embodiment, the reporter gene is operatively linked to a constitutive promoter.

**[0046]** In another aspect, the reporter signal can be detected from a sample at a limit of detection (LoD) of less than 1,000 colony forming units (CFU). In other aspects, the reporter signal can be detected from a sample at a limit of detection (LoD) of less than 100 colony forming units (CFU). In one aspect, the reporter signal can be detected from a sample at a limit of detection (LoD) of less than 10 colony forming units (CFU). In other aspects, the reporter signal can be detected from a sample at a LoD less than five CFU. In another aspect, the reporter signal can be detected from a sample at a LoD of three or less CFU.

**[0047]** In one embodiment, the method includes providing an antibiotic to said sample at a pre-determined concentration and detecting a presence or absence of said reporter signal to determine whether said bacterial cell is resistant or sensitive to said antibiotic. In another

embodiment, the method includes providing varying pre-determined concentrations antibiotic to said sample and detecting the amount of said reporter signal to determine the minimum inhibitory concentration of said bacterial cell to said antibiotic.

**[0048]** In one aspect, the invention includes a composition comprising a nucleic acid construct that encodes a nucleic acid reporter transcript that is capable of forming at least two conformations comprising a first conformation that prevents reporter expression comprising an intramolecular double stranded region comprising a first subsequence and a second subsequence, and a second conformation that lacks said intramolecular double-stranded region and allows reporter gene expression, wherein conversion between said first and second conformations is mediated by competitive binding of a cellular transcript to said first and/or said second subsequence.

**[0049]** In another aspect, the invention includes a non-replicative transduction particle comprising said nucleic acid construct. In yet another aspect, the competitive binding of said cellular transcript to said first and/or said second subsequence results in said second conformation of said nucleic acid reporter construct. In one aspect, the first subsequence or said second subsequence comprises a cis-repressing sequence. In another aspect, the cis-repressing sequence comprises a sequence that is complementary or substantially complementary to a portion of said cellular transcript. In other aspects, the first subsequence or said second subsequence comprises a reporter gene sequence. In yet another aspect, the reporter gene sequence comprises a ribosome binding site. In other aspects, the reporter gene sequence encodes a detectable molecule. In another aspect, the detectable marker comprises a fluorescent molecule or an enzyme capable of mediating a luminescence or colorimetric reaction. In one embodiment, the reporter gene sequence encodes a selectable marker. In another embodiment, the selectable marker comprises an antibiotic resistance gene.

**[0050]** In other embodiments, the first subsequence and said second subsequence are located *cis* to each other on said nucleic acid construct to form said intramolecular double stranded region. In certain embodiments, the first subsequence and said second subsequence are complementary or substantially complementary to each other to form said intramolecular double stranded region. In one embodiment, the first subsequence or said second subsequence of said first conformation comprises a transcriptional enhancer sequence, and wherein said transcriptional enhancer sequence is upstream from a coding region of said reporter gene sequence. In another embodiment, the first conformation of said nucleic acid reporter transcript is capable of binding to a cleaving enzyme. In other embodiments, the first conformation of said nucleic acid reporter transcript is a target for degradation by a cellular

enzyme. In other aspects, the first conformation comprises a non-binding intramolecular region. In another aspect, the non-binding intramolecular region is located 3' of said first subsequence and 5' of said second subsequence. In other aspects, the non-binding intramolecular region comprises a sequence YUNR, wherein Y is a pyrimidine, U is a Uracil, N is any nucleotide, and R is a purine.

**[0051]** In one embodiment, the first subsequence or said second subsequence comprises a modified sequence of said cellular transcript. In another embodiment, the modified sequence comprises a nucleotide substitution. In yet another embodiment, the modified sequence comprises a sequence insertion, a deletion or an inversion of said cellular transcript.

**[0052]** The method includes a composition comprising a nucleic acid construct that encodes a nucleic acid reporter transcript comprising a gene reporter sequence and that is capable of forming at least two conformations of said nucleic acid reporter transcript, a first unstable conformation that prevents translation of said reporter gene sequence in said nucleic acid reporter transcript, and a second stable conformation resulting from binding of said first unstable conformation with a cellular transcript, said second stable secondary conformation allowing translation of said reporter gene sequence of said nucleic acid reporter transcript.

**[0053]** In one embodiment, the composition comprises a non-replicative transduction particle comprising said nucleic acid construct. In another embodiment, the cellular transcript binds at a 3'UTR sequence of said nucleic acid reporter transcript. In one embodiment, the second stable secondary conformation is formed by cleavage of a portion of a sequence of said first unstable secondary conformation. In another embodiment, the reporter gene sequence encodes a detectable molecule. In some embodiments, the detectable marker comprises a fluorescent molecule or an enzyme capable of mediating a luminescence or colorimetric reaction. In other embodiments, the reporter gene sequence encodes a selectable marker. In another embodiment, the selectable marker comprises an antibiotic resistance gene.

**[0054]** The invention also includes a composition comprising a nucleic acid construct that encodes a nucleic acid reporter transcript comprising a reporter gene sequence and that is capable of forming at least two conformations of said nucleic acid reporter transcript, comprising a first conformation that prevents further transcription of said nucleic acid construct, and a second conformation formed upon binding of said first conformation with a cellular transcript, wherein said second conformation allows transcription of said nucleic acid construct. In some embodiments, the composition comprises a non-replicative transduction

particle comprising said nucleic acid construct. In another embodiment, the nucleic acid reporter transcript comprises a cis-repressing sequence.

**[0055]** In one embodiment, the nucleic acid reporter transcript comprises a reporter gene sequence. In another embodiment, the first conformation forms from a binding of said cis-repressing sequence to said reporter gene sequence. In some embodiments, the first conformation is a substrate for a cleaving enzyme. In one embodiment, the first conformation of said nucleic acid reporter transcript comprises a sequence that forms a transcription termination structure. In other embodiments, the binding of said cellular transcript to said sequence that forms a transcription termination structure results in cleavage of a portion of said nucleic acid reporter transcript and formation of said second conformation.

**[0056]** The invention comprises a vector comprising a regulatory sequence operably linked to a nucleic acid sequence that encodes said nucleic acid reporter transcript described herein.

**[0057]** The invention includes a method for detecting a target transcript in a cell, comprising: introducing into said cell said nucleic acid reporter construct described herein; and detecting the presence or absence of an output signal from said cell, wherein said presence of said output signal indicates the presence of the target transcript in said cell. The method includes detecting a presence of a bacterial cell based on detecting said presence of said target transcript.

**[0058]** In one embodiment, the method for detecting a presence of a bacterial cell in a sample comprising introducing into said sample said nucleic acid reporter construct described herein; and detecting the presence or absence of an output signal from said sample, wherein said presence of said output signal indicates the presence of the bacterial cell in said sample.

**[0059]** The invention comprises a kit, comprising a compartment for holding a sample comprising a cell and said nucleic acid reporter construct described herein; and instructions for detecting the presence or absence of an output signal from said sample, wherein the presence of the output signal indicates the presence of a target transcript in said cell

**[0060]** The invention comprises a composition, comprising a non-replicative transduction particle comprising a nucleic acid reporter construct, the nucleic acid reporter construct comprising a first promoter operatively linked a reporter gene, wherein said first promoter is capable of being induced by an inducer protein endogenous in a bacterial cell.

**[0061]** The invention includes a method for detecting a presence of a bacterial cell in a sample comprising contacting said sample with a non-replicative transduction particle comprising nucleic acid reporter construct comprising a first promoter operatively linked to a

reporter gene, wherein said first promoter is capable of being induced by an inducer protein endogenous to said bacterial cell; and detecting the presence or absence of an output signal from said reporter gene, wherein said presence of said output signal indicates the presence of said bacterial cell in said sample.

**[0062]** In one embodiment, the first promoter is the same as an inducible promoter operatively linked to a target nucleic acid molecule in said bacterial cell.

**[0063]** The invention comprises a composition, comprising a non-replicative transduction particle comprising a nucleic acid reporter construct, the nucleic acid reporter construct comprising a reporter gene that encodes a reporter molecule, the non-replicative transduction particle capable of entering a bacterial cell; and a caged substrate that exogenous to said bacterial cell that once un-caged is capable of reacting to said reporter molecule in said cell.

**[0064]** The invention comprises a method for detecting a presence of a bacterial cell in a sample comprising contacting said sample with a caged substrate and a non-replicative transduction particle comprising a nucleic acid reporter construct, the nucleic acid reporter construct comprising a reporter gene that encodes a reporter molecule, the caged substrate exogenous to said cell that once un-caged is capable of binding to said reporter molecule in said bacterial cell; and detecting the presence or absence of an output signal from said reporter molecule, wherein said presence of said output signal indicates the presence of said bacterial cell in said sample.

**[0065]** In one embodiment, a target enzyme in said cell binds said caged substrate to produce an un-caged substrate. In some embodiments, the un-caged substrate reacts with said reporter molecule to produce said output signal.

**[0066]** The invention also includes a composition, comprising a non-replicative transduction particle comprising a nucleic acid reporter construct, the nucleic acid reporter construct encoding a switchable molecule capable of binding to a target molecule in a bacterial cell to form a complex; and a substrate capable of penetrating said cell and binding said complex to produce a detectable signal from said cell.

**[0067]** The invention includes a method for detecting a presence of a bacterial cell in a sample comprising contacting said sample with a substrate and a non-replicative transduction particle comprising a nucleic acid reporter construct encoding a switchable molecule, the switchable molecule capable of binding a target molecule in said cell to form a complex, the substrate capable of binding said complex to form a substrate-bound complex; and detecting the presence or absence of an output signal from said substrate-bound complex, wherein said presence of said output signal indicates the presence of said bacterial cell in said sample. In

one embodiment, the binding of said switchable molecule to said target molecule produces a conformational change in said switchable molecule. In another embodiment, the conformational change in said switchable molecule allows said substrate to bind to said complex.

#### **BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS**

**[0068]** These and other features, aspects, and advantages of the present invention will become better understood with regard to the following description, and accompanying drawings, where:

**[0069]** Figure 1 illustrates an example of the design and function of the silent mutation/complementation-based P1 plasmid packaging system, according to an embodiment of the invention.

**[0070]** Figure 2 illustrates a schematic of the pGWP10001 vector, according to an embodiment of the invention.

**[0071]** Figure 3 illustrates an example of the design and function of a pac-site deletion/complementation plasmid packaging system, according to an embodiment of the invention.

**[0072]** Figure 4 illustrates a schematic of the pGW80A0001 vector, according to an embodiment of the invention.

**[0073]** Figure 5 depicts the process for genomic island (GI) packaging by a bacteriophage, according to an embodiment of the invention.

**[0074]** Figure 6 depicts an example of the design and function of a GI-based packaging system, according to an embodiment of the invention.

**[0075]** Figure 7 depicts the design and function of a GI-based packaging system that lacks the integrase gene, according to an embodiment of the invention.

**[0076]** Figure 8 depicts the design and function of a SaPIbov2-based packaging system that lacks the integrase gene, according to an embodiment of the invention.

[0077] Figure 9 depicts a system for the use of NRTPs for the detection of inducers to target gene promoters within viable cells, according to an embodiment of the invention.

[0078] Figure 10 depicts a reporter system that includes a reporter nucleic acid molecule (e.g., plasmid) that is constructed for detecting VanR, the inducer of the promoter of the vancomycin resistance (vanA) gene in *Enterococcus faecium* (or *E. faecalis*), according to an embodiment of the invention. The reporter plasmid carries a reporter gene that is operatively linked to the vanA gene promoter.

[0079] Figure 11 depicts a reporter system that includes a reporter nucleic acid molecule constructed for detecting TcdD, the inducer of the promoters of the toxins A and B genes (tcdA and tcdB, respectively) of *C. difficile*, according to an embodiment of the invention. The reporter nucleic acid molecule includes a reporter gene that is operatively linked to the tcdA gene promoter.

[0080] Figure 12 depicts a reporter system that includes a reporter nucleic acid molecule is constructed for detecting SarS, the inducer of the promoter of the Protein A gene (spa) in *S. aureus*, according to an embodiment of the invention. The reporter nucleic acid molecule includes the bacterial luciferase genes luxA and luxB operatively linked to the spa gene promoter ( $P_{spa}$ ).

[0081] Figure 13 shows a reporter system that comprises a system for the detection of intracellular enzymes within viable cells that employs caged substrate molecules that can be un-caged by a target intracellular enzyme, according to an embodiment of the invention.

[0082] Figure 14 depicts the design and function of a  $\beta$ -lactamase enzyme detection system, according to an embodiment of the invention.

[0083] Figure 15 shows a reporter system for the detection of intracellular molecules within viable cells that employs switchable molecules capable of generating a detectable signal upon their binding to a target molecule, according to an embodiment of the invention.

[0084] Figure 16 depicts the design and function of a bacteriophage/switchable-aptamer (SA)-based intracellular molecule reporter system, according to an embodiment of the invention.

[0085] Figure 17 depicts an example of a system that uses a cis-repression mechanism that can target the 5' UTR (untranslated region) of a reporter sequence on a reporter transcript, according to an embodiment of the invention.

[0086] Figure 18 shows an example of a system for detecting the presence of a target transcript in a cell that is based on a cis-repression mechanism targeting the ribosome binding

site (RBS) of a reporter sequence in a reporter transcript, according to an embodiment of the invention.

**[0087]** Figure 19 illustrates an exemplary system for detecting the presence of a target transcript in a cell that is based on a *cis*-repression mechanism targeting the coding region (“AUG”) of a reporter sequence in a reporter transcript, according to an embodiment of the invention.

**[0088]** Figure 20 illustrates an example system for detecting the presence of a target transcript in a cell that is based on a repression mechanism using an unstable reporter transcript, according to an embodiment of the invention.

**[0089]** Figure 21 shows the results of the transduction assay in which 36 tetracycline-sensitive MRSA were exposed to transduction particles carrying pGW80A0001 and then were spotted onto media plates containing 5 ug/mL of tetracycline, according to an embodiment of the invention.

**[0090]** Figure 22 illustrates the luminescence measured from 80 clinical isolates of MRSA and 28 clinical isolates of methicillin sensitive *S. aureus* (MSSA) transduced with the transduction particle, according to an embodiment of the invention.

**[0091]** Figure 23 shows the results of *S. aureus* growth at 4, 8, 16, 32, 64, and 128 ug/mL of cefoxitin.

**[0092]** Figure 24 shows the RLU values obtained by the NRTP assay in the presence of 4, 8, 16, 32, 64, and 128 ug/mL cefoxitin. The x-axis in Figure 24 is set at the MSSA RLU cutoff value.

**[0093]** Figure 25 shows a secondary structure of the *mecA* transcript (SEQ ID NO: 16) generated based on the lowest energy conformation calculated by MFold and visualized with VARNA.

**[0094]** Figure 26 shows the terminal loop 23 (T23) of the *mecA* transcript (nucleotides 1,464-1,519 of SEQ ID NO: 16) that contains a YUNR consensus sequence.

**[0095]** Figure 27 depicts a *cis*-repressing sequence added to the 5’ terminus of the *luxAB* genes and designed to form a stem-loop structure that blocks the RBS sequence (“AAGGAA”) of the *luxA* gene (nucleotides 1-61 of SEQ ID NO: 19).

**[0096]** Figure 28 shows a diagram of base pairing between the target transcript (nucleotides 1,464-1,519 of SEQ ID NO: 16) and the *cis*-repressing sequence of the reporter transcript (nucleotides 1-61 of SEQ ID NO: 19).

[0097] Figure 29 shows an example of a target *mecA* gene sequence (SEQ ID NO: 15), according to an embodiment of the invention.

[0098] Figure 30 shows an exemplary *mecA* transcript sequence that can be used for designing a reporter transcript (SEQ ID NO:16), according to an embodiment of the invention.

[0099] Figure 31 is an example of a *luxAB* gene loci DNA sequence (SEQ ID NO: 17) that can be used for designing a reporter transcript, according to an embodiment of the invention.

[00100] Figure 32 is an example of a *luxAB* transcript sequence that can be used for designing a reporter transcript (SEQ ID NO:18), according to an embodiment of the invention.

[00101] Figure 33 is an example of a *luxAB* cis-repressed transcript sequence that can be used in a reporter transcript (SEQ ID NO:19), according to an embodiment of the invention.

[00102] Figure 34 shows an example of a cell comprising a vector that encodes a reporter transcript, where there is no endogenous *mecA* transcript in the cell, according to an embodiment of the invention.

[00103] Figure 35 shows a vector introduced into a cell, where the vector encodes the reporter transcript, which includes a cis-repressing sequence and a reporter sequence (*luxA* and *luxB* genes). When the *mecA* transcript present in the cell binds to the cis-repressing sequence, the inhibitory hairpin loop opens up and the RBS for the *luxA* gene is exposed. Translation of the reporter sequences (*luxA* and *luxB*) can occur, resulting in the formation of a *luxAB* enzyme. The *luxAB* enzyme produces a detectable luminescent signal. In this manner, the transcript reporter vector reports the presence of endogenous *mecA* transcripts within a cell.

## DETAILED DESCRIPTION OF THE INVENTION

### I. Definitions

[00104] Terms used in the claims and specification are defined as set forth below unless otherwise specified.

[00105] As used herein, “reporter nucleic acid molecule” refers to a nucleotide sequence comprising a DNA or RNA molecule. The reporter nucleic acid molecule can be naturally occurring or an artificial or synthetic molecule. In some embodiments, the reporter nucleic acid molecule is exogenous to a host cell and can be introduced into a host cell as part of an exogenous nucleic acid molecule, such as a plasmid or vector. In certain embodiments, the reporter nucleic acid molecule can be complementary to a target gene in a cell. In other embodiments, the reporter nucleic acid molecule comprises a reporter gene encoding a reporter molecule (e.g., reporter enzyme, protein). In some embodiments, the reporter

nucleic acid molecule is referred to as a “reporter construct” or “nucleic acid reporter construct.”

**[00106]** A “reporter molecule” or “reporter” refers to a molecule (*e.g.*, nucleic acid or protein) that confers onto an organism a detectable or selectable phenotype. The detectable phenotype can be colorimetric, fluorescent or luminescent, for example. Reporter molecules can be expressed from reporter genes encoding enzymes mediating luminescence reactions (*luxA*, *luxB*, *luxAB*, *luc*, *ruc*, *nluc*), genes encoding enzymes mediating colorimetric reactions (*lacZ*, HRP), genes encoding fluorescent proteins (GFP, eGFP, YFP, RFP, CFP, BFP, mCherry, near-infrared fluorescent proteins), nucleic acid molecules encoding affinity peptides (His-tag, 3X-FLAG), and genes encoding selectable markers (*ampC*, *tet(M)*, CAT, erm). The reporter molecule can be used as a marker for successful uptake of a nucleic acid molecule or exogenous sequence (plasmid) into a cell. The reporter molecule can also be used to indicate the presence of a target gene, target nucleic acid molecule, target intracellular molecule, or a cell, as described herein. Alternatively, the reporter molecule can be a nucleic acid, such as an aptamer or ribozyme.

**[00107]** In some aspects of the invention, the reporter nucleic acid molecule is operatively linked to a promoter. In other aspects of the invention, the promoter can be chosen or designed to contribute to the reactivity and cross-reactivity of the reporter system based on the activity of the promoter in specific cells (*e.g.*, specific species) and not in others. In certain aspects, the reporter nucleic acid molecule comprises an origin of replication. In other aspects, the choice of origin of replication can similarly contribute to reactivity and cross-reactivity of the reporter system, when replication of the reporter nucleic acid molecule within the target cell contributes to or is required for reporter signal production based on the activity of the origin of replication in specific cells (*e.g.*, specific species) and not in others. In some embodiments, the reporter nucleic acid molecule forms a replicon capable of being packaged as concatameric DNA into a progeny virus during virus replication.

**[00108]** As used herein, a “target transcript” refers to a portion of a nucleotide sequence of a DNA sequence or an mRNA molecule that is naturally formed by a target cell including that formed during the transcription of a target gene and mRNA that is a product of RNA processing of a primary transcription product. The target transcript can also be referred to as a cellular transcript or naturally occurring transcript.

**[00109]** As used herein, the term “transcript” refers to a length of nucleotide sequence (DNA or RNA) transcribed from a DNA or RNA template sequence or gene. The transcript can be a cDNA sequence transcribed from an RNA template or an mRNA sequence

transcribed from a DNA template. The transcript can be protein coding or non-coding. The transcript can also be transcribed from an engineered nucleic acid construct.

**[00110]** A transcript derived from a reporter nucleic acid molecule can be referred to as a “reporter transcript.” The reporter transcript can include a reporter sequence and a cis-repressing sequence. The reporter transcript can have sequences that form regions of complementarity, such that the transcript includes two regions that form a duplex (e.g., an intermolecular duplex region). One region can be referred to as a “cis-repressing sequence” and has complementarity to a portion or all of a target transcript and/or a reporter sequence. A second region of the transcript is called a “reporter sequence” and can have complementarity to the cis-repressing sequence. Complementarity can be full complementarity or substantial complementarity. The presence and/or binding of the cis-repressing sequence with the reporter sequence can form a conformation in the reporter transcript, which can block further expression of the reporter molecule. The reporter transcript can form secondary structures, such as a hairpin structure, such that regions within the reporter transcript that are complementary to each other can hybridize to each other.

**[00111]** “Introducing into a cell,” when referring to a nucleic acid molecule or exogenous sequence (e.g., plasmid, vector, construct), means facilitating uptake or absorption into the cell, as is understood by those skilled in the art. Absorption or uptake of nucleic acid constructs or transcripts can occur through unaided diffusive or active cellular processes, or by auxiliary agents or devices including via the use of bacteriophage, virus, and transduction particles. The meaning of this term is not limited to cells *in vitro*; a nucleic acid molecule may also be “introduced into a cell,” wherein the cell is part of a living organism. In such instance, introduction into the cell will include the delivery to the organism. For example, for *in vivo* delivery, nucleic acid molecules, constructs or vectors of the invention can be injected into a tissue site or administered systemically. *In vitro* introduction into a cell includes methods known in the art, such as electroporation and lipofection. Further approaches are described herein or known in the art.

**[00112]** A “transduction particle” refers to a virus capable of delivering a non-viral nucleic acid molecule into a cell. The virus can be a bacteriophage, adenovirus, etc.

**[00113]** A “non-replicative transduction particle” refers to a virus capable of delivering a non-viral nucleic acid molecule into a cell, but does not package its own replicated viral genome into the transduction particle. The virus can be a bacteriophage, adenovirus, etc.

**[00114]** A “plasmid” is a small DNA molecule that is physically separate from, and can replicate independently of, chromosomal DNA within a cell. Most commonly found as small

circular, double-stranded DNA molecules in bacteria, plasmids are sometimes present in archaea and eukaryotic organisms. Plasmids are considered replicons, capable of replicating autonomously within a suitable host.

[00115] A “vector” is a nucleic acid molecule used as a vehicle to artificially carry foreign genetic material into another cell, where it can be replicated and/or expressed.

[00116] A “virus” is a small infectious agent that replicates only inside the living cells of other organisms. Virus particles (known as virions) include two or three parts: i) the genetic material made from either DNA or RNA molecules that carry genetic information; ii) a protein coat that protects these genes; and in some cases, iii) an envelope of lipids that surrounds the protein coat.

[00117] “MRSA” refers to Methicillin-resistant *Staphylococcus aureus*.

[00118] “MSSA” refers to Methicillin-sensitive *Staphylococcus aureus*.

[00119] The term “ameliorating” refers to any therapeutically beneficial result in the treatment of a disease state, *e.g.*, a disease state, including prophylaxis, lessening in the severity or progression, remission, or cure thereof.

[00120] The term “*in situ*” refers to processes that occur in a living cell growing separate from a living organism, *e.g.*, growing in tissue culture.

[00121] The term “*in vivo*” refers to processes that occur in a living organism.

[00122] The term “mammal” as used herein includes both humans and non-humans and include but is not limited to humans, non-human primates, canines, felines, murines, bovines, equines, and porcines.

[00123] “G,” “C,” “A” and “U” each generally stand for a nucleotide that contains guanine, cytosine, adenine, and uracil as a base, respectively. “T” and “dT” are used interchangeably herein and refer to a deoxyribonucleotide wherein the nucleobase is thymine, *e.g.*, deoxyribothymine. However, it will be understood that the term “ribonucleotide” or “nucleotide” or “deoxyribonucleotide” can also refer to a modified nucleotide, as further detailed below, or a surrogate replacement moiety. The skilled person is well aware that guanine, cytosine, adenine, and uracil may be replaced by other moieties without substantially altering the base pairing properties of an oligonucleotide comprising a nucleotide bearing such replacement moiety. For example, without limitation, a nucleotide comprising inosine as its base may base pair with nucleotides containing adenine, cytosine, or uracil. Hence, nucleotides containing uracil, guanine, or adenine may be replaced in the nucleotide sequences of the invention by a nucleotide containing, for example, inosine. Sequences comprising such replacement moieties are embodiments of the invention.

**[00124]** As used herein, the term “complementary,” when used to describe a first nucleotide sequence in relation to a second nucleotide sequence, refers to the ability of an oligonucleotide or polynucleotide comprising the first nucleotide sequence to hybridize and form a duplex structure under certain conditions with an oligonucleotide or polynucleotide comprising the second nucleotide sequence, as will be understood by the skilled person. Complementary sequences are also described as binding to each other and characterized by binding affinities.

**[00125]** For example, a first nucleotide sequence can be described as complementary to a second nucleotide sequence when the two sequences hybridize (*e.g.*, anneal) under stringent hybridization conditions. Hybridization conditions include temperature, ionic strength, pH, and organic solvent concentration for the annealing and/or washing steps. The term stringent hybridization conditions refers to conditions under which a first nucleotide sequence will hybridize preferentially to its target sequence, *e.g.*, a second nucleotide sequence, and to a lesser extent to, or not at all to, other sequences. Stringent hybridization conditions are sequence dependent, and are different under different environmental parameters. Generally, stringent hybridization conditions are selected to be about 5°C lower than the thermal melting point ( $T_m$ ) for the nucleotide sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the first nucleotide sequences hybridize to a perfectly matched target sequence. An extensive guide to the hybridization of nucleic acids is found in, *e.g.*, Tijssen (1993) Laboratory Techniques in Biochemistry and Molecular Biology--Hybridization with Nucleic Acid Probes part I, chap. 2, “Overview of principles of hybridization and the strategy of nucleic acid probe assays,” Elsevier, N.Y. (“Tijssen”). Other conditions, such as physiologically relevant conditions as may be encountered inside an organism, can apply. The skilled person will be able to determine the set of conditions most appropriate for a test of complementarity of two sequences in accordance with the ultimate application of the hybridized nucleotides.

**[00126]** This includes base-pairing of the oligonucleotide or polynucleotide comprising the first nucleotide sequence to the oligonucleotide or polynucleotide comprising the second nucleotide sequence over the entire length of the first and second nucleotide sequence. Such sequences can be referred to as “fully complementary” with respect to each other herein. However, where a first sequence is referred to as “substantially complementary” with respect to a second sequence herein, the two sequences can be fully complementary, or they may form one or more, but generally not more than 4, 3 or 2 mismatched base pairs upon

hybridization, while retaining the ability to hybridize under the conditions most relevant to their ultimate application. However, where two oligonucleotides are designed to form, upon hybridization, one or more single stranded overhangs, such overhangs shall not be regarded as mismatches with regard to the determination of complementarity. For example, a dsRNA comprising one oligonucleotide 21 nucleotides in length and another oligonucleotide 23 nucleotides in length, wherein the longer oligonucleotide comprises a sequence of 21 nucleotides that is fully complementary to the shorter oligonucleotide, may yet be referred to as “fully complementary” for the purposes described herein.

**[00127]** “Complementary” sequences, as used herein, may also include, or be formed entirely from, non-Watson-Crick base pairs and/or base pairs formed from non-natural and modified nucleotides, in as far as the above requirements with respect to their ability to hybridize are fulfilled. Such non-Watson-Crick base pairs includes, but not limited to, G:U Wobble or Hoogstein base pairing.

**[00128]** The terms “complementary,” “fully complementary” and “substantially complementary” herein may be used with respect to the base matching between two strands of a dsRNA, or between the antisense strand of a dsRNA and a target sequence, between complementary strands of a single stranded RNA sequence or a single stranded DNA sequence, as will be understood from the context of their use.

**[00129]** As used herein, a “duplex structure” comprises two anti-parallel and substantially complementary nucleic acid sequences. Complementary sequences in a nucleic acid construct, between two transcripts, between two regions within a transcript, or between a transcript and a target sequence can form a “duplex structure.” In general, the majority of nucleotides of each strand are ribonucleotides, but as described in detail herein, each or both strands can also include at least one non-ribonucleotide, *e.g.*, a deoxyribonucleotide and/or a modified nucleotide. The two strands forming the duplex structure may be different portions of one larger RNA molecule, or they may be separate RNA molecules. Where the two strands are part of one larger molecule, and therefore are connected by an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting RNA chain is referred to as a “hairpin loop.” Where the two strands are connected covalently by means other than an uninterrupted chain of nucleotides between the 3'-end of one strand and the 5'-end of the respective other strand forming the duplex structure, the connecting structure is referred to as a “linker.” The RNA strands may have the same or a different number of nucleotides. The maximum number of base pairs is the number of nucleotides in the shortest strand of the duplex minus any

overhangs that are present in the duplex. Generally, the duplex structure is between 15 and 30 or between 25 and 30, or between 18 and 25, or between 19 and 24, or between 19 and 21, or 19, 20, or 21 base pairs in length. In one embodiment the duplex is 19 base pairs in length. In another embodiment the duplex is 21 base pairs in length. When two different siRNAs are used in combination, the duplex lengths can be identical or can differ.

**[00130]** As used herein, the term “region of complementarity” refers to the region on the antisense strand that is substantially complementary to a sequence, for example a target sequence, as defined herein. Where the region of complementarity is not fully complementary to the target sequence, the mismatches are most tolerated in the terminal regions and, if present, are generally in a terminal region or regions, *e.g.*, within 6, 5, 4, 3, or 2 nucleotides of the 5’ and/or 3’ terminus.

**[00131]** The term percent “identity,” in the context of two or more nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences that have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms described below (*e.g.*, BLASTP and BLASTN or other algorithms available to persons of skill) or by visual inspection. Depending on the application, the percent “identity” can exist over a region of the sequence being compared, *e.g.*, over a functional domain, or, alternatively, exist over the full length of the two sequences to be compared.

**[00132]** For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

**[00133]** Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Nat'l. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally Ausubel *et al.*, *infra*).

**[00134]** One example of an algorithm that is suitable for determining percent sequence identity and sequence similarity is the BLAST algorithm, which is described in Altschul *et*

al., J. Mol. Biol. 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information ([www.ncbi.nlm.nih.gov/](http://www.ncbi.nlm.nih.gov/)).

[00135] The term “sufficient amount” means an amount sufficient to produce a desired effect, e.g., an amount sufficient to produce a detectable signal from a cell.

[00136] The term “therapeutically effective amount” is an amount that is effective to ameliorate a symptom of a disease. A therapeutically effective amount can be a “prophylactically effective amount” as prophylaxis can be considered therapy.

[00137] It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise.

## II. Lysogenic and Lytic Cycle of Viruses

[00138] Viruses undergo lysogenic and lytic cycles in a host cell. If the lysogenic cycle is adopted, the phage chromosome can be integrated into the bacterial chromosome, or it can establish itself as a stable plasmid in the host, where it can remain dormant for long periods of time. If the lysogen is induced, the phage genome is excised from the bacterial chromosome and initiates the lytic cycle, which culminates in lysis of the cell and the release of phage particles. The lytic cycle leads to the production of new phage particles which are released by lysis of the host.

[00139] Certain temperate phage can exhibit lytic activity, and the propensity for this may vary with varying host bacteria. To illustrate this phenomenon, the lytic activity of two temperate *S. aureus* phages on ten MRSA clinical isolates was examined via plaque assay (Table 1). The phage  $\phi$ 11 exhibited lytic activity on 10 out of 10 clinical MRSA isolates and  $\phi$ 80 $\alpha$  exhibited lytic activity on six of the 10 clinical MRSA isolates. Thus, reporter assays relying on the natural lysogenic cycle of phages can be expected to exhibit lytic activity sporadically.

Table 1: Lytic activity (denoted by the letter “x”) of the *S. aureus* temperate phages  $\phi$ 11 and  $\phi$ 80 $\alpha$  on ten clinical MRSA isolates

MRSA isolate	$\phi$ 11	$\phi$ 80 $\alpha$
1	x	
2	x	
3	x	x
4	x	x

<b>5</b>	x	x
<b>6</b>	x	
<b>7</b>	x	x
<b>8</b>	x	
<b>9</b>	x	x
<b>10</b>	x	x

**[00140]** In addition, virus-based reporter assays, such as phage-based reporters, can suffer from limited reactivity (*i.e.*, analytical inclusivity) due to limits in the phage host range caused by host-based and prophage-derived phage resistance mechanisms. These resistance mechanisms target native phage nucleic acid that can result in the degradation or otherwise inhibition of the phage DNA and functions. Such resistance mechanisms include restriction systems that cleave phage DNA and CRISPR systems that inhibit phage-derived transcripts.

**[00141]** Both lytic activity and phage resistance can be inhibitory to assays based on reporter phages. Lytic activity can inhibit signal by destroying or otherwise inhibiting the cell in its ability to generate a detectable signal and thus affecting limits of detection by reducing the amount of detectable signal or preventing the generation of a detectable signal. Phage resistance mechanisms can limit the host range of the phage and limit the inclusivity of the phage-based reporter, similarly affecting limits of detection by reducing the amount of detectable signal or preventing the generation of a detectable signal. Both lytic activity and phage resistance caused by the incorporation of phage DNA in a reporter phage can lead to false-negative results in assays that incorporate these phage reporters.

### **III. Methods for Producing Non-Replicative Transduction Particles (N RTP)**

#### **A. Disruption/Complementation-Based Methods For Producing Non-Replicative Transduction Particles**

##### **1) Silent Mutation/Complementation Packaging System**

**[00142]** The invention includes methods for producing N RTPs using a silent mutation/complementation-based method.

**[00143]** This non-replicative transduction particle packaging system is based on introducing a silent mutation into a component of the genome of a virus that is recognized by the viral packaging machinery as the element from which genomic packaging is initiated during viral production. Examples of such an element include the pac-site sequence of pac-type bacteriophages and the cos-site sequence of cos-type bacteriophages.

**[00144]** Because these packaging initiation sites are often found within coding regions of genes that are essential to virus production, the silent mutation is introduced such that the pac-site is no longer recognized as a site of packaging initiation by the viral packaging machinery. At the same time, the mutation does not disrupt the gene in which the site is encoded. By disrupting the packaging site sequence, the mutated virus is able to undergo a lytic cycle, but is unable to package its genomic DNA into its packaging unit.

**[00145]** An exogenous reporter nucleic acid molecule, such as plasmid DNA, can be introduced into a host cell that has been lysogenized with a viral genome with a mutated packaging initiation site sequence. The exogenous reporter nucleic acid molecule can include a native packaging initiation site sequence. The exogenous reporter nucleic acid molecule can be introduced into the cell and replicated in the cell. When the mutated virus is undergoing a lytic cycle, the expressed viral packaging machinery packages the exogenous reporter nucleic acid molecule with the native packaging initiation site sequence into the viral packaging unit. The viral genome is not packaged into the packaging unit because its packaging initiation site sequence has been mutated. In certain embodiments, the mutation in the packaging initiation site sequence comprises a silent mutation that prevents cleavage of the packaging initiation sequence, but does not disrupt the expression of the gene product that encompasses the packaging initiation site sequence. This produces non-replicative transduction particles, *e.g.*, viral structural components carrying the replicated exogenous nucleic acid molecule.

**[00146]** An example of such a system is based on the bacteriophage P1, a pac-type phage. In an embodiment, a plasmid including a native P1 pac site is transformed into a cell. The cell is lysogenized with a P1 prophage genome. The P1 prophage genome includes a silent mutation in the pac-site sequence encoded within the pacA gene of P1. When the lytic cycle of the prophage is induced, the system results in the production of P1-based transduction particles carrying the plasmid DNA. An example of a silent mutation that is suitable for this system is described in U.S. Pub. No. 2005/0118719, filed on November 7, 2002, which is incorporated by reference in its entirety. An example is also found in SEQ ID NO: 2, listed below (P1 pac-site with silent mutations, lower case letters signify mutated bases).

**[00147]** Figure 1 illustrates an example of the design and function of the silent mutation/complementation-based P1 plasmid packaging system 100, according to an embodiment of the invention. In this system, an *E. coli* cell 101 is lysogenized with a P1 prophage 102 that includes a silent mutation in its packaging initiation site sequence (*e.g.*, pac-site). The cell is transformed with a plasmid containing the native pac-site 103, and the

plasmid is replicated in the cell to form plasmid concatamers 104. The plasmid can also include a reporter gene that encodes a reporter molecule. When the lytic cycle of the P1 prophage is induced, the P1 prophage is excised from the bacterial genome and the P1 structural components, such as capsid proteins, 105 are expressed. The P1 structural components only package DNA that contains a native pac-site (e.g., plasmid DNA), thus producing non-replicative transduction particles carrying plasmid DNA 106 (e.g., a reporter gene).

**[00148]** An example vector for use in the silent mutation/complementation-based P1 plasmid packaging system is shown in Figure 2. Details about how to construct the strains and vectors of the silent mutation/complementation-based P1 plasmid packaging system are described in detail in Example 1 below.

## 2) Deletion/Complementation-Based Packaging System

**[00149]** The invention includes methods for producing NRTPs using a deletion/complementation-based method.

**[00150]** This non-replicative transduction particle packaging system is based on deletion of a component of the genome of a virus that is recognized by the viral packaging machinery as the element from which genomic packaging is initiated during viral production. Examples of such an element include the pac-site sequence of pac-type bacteriophages and the cos-site sequence of cos-type bacteriophages. These packaging initiation sites are often found within coding regions of genes that are essential to virus production. In some embodiments, the packaging initiation site alone is deleted, which allows the mutated virus to undergo a lytic cycle but does not allow the virus to package its genomic DNA. For example, SEQ ID NO: 6 is an example of a P1 pacA gene with a deleted pac-site sequence (lower case letters indicate the deleted pac-site sequence). In other embodiments, the entire gene comprising the packaging initiation site is deleted. For example, SEQ ID NO: 8 shows the deletion of the terS gene (lower case characters show the deleted sequence).

**[00151]** In one example, a cell's genome is lysogenized with a viral genome where the packaging initiation site has been deleted. A complementing plasmid is introduced into the cell, and the plasmid DNA includes a gene with a packaging initiation site sequence that complements the deleted packaging initiation site sequence in the viral genome. When the mutated virus is undergoing a lytic cycle, the viral packaging proteins package a replicon of the plasmid DNA into the packaging unit because of its packaging initiation site, and non-replicative transduction particles are produced carrying the replicated plasmid DNA.

**[00152]** In some embodiments, it is preferable that the deletion/complementation is designed such that there is no homology between the mutated virus DNA and the complementing exogenous DNA. This is because lack of homology between the mutated virus DNA and the complementing exogenous DNA avoids the possibility of homologous recombination between the two DNA molecules that can result in re-introduction of a packaging sequence into the virus genome. To accomplish a lack of homology, one strategy is to delete the entire gene that contains the packaging initiation site sequence from the virus genome and then complement this gene with an exogenous DNA molecule that contains no more than exactly the DNA sequence that was deleted from virus. In this strategy, the complementing DNA molecule is designed to express the gene that was deleted from the virus.

**[00153]** Another example of such a system is provided using the bacteriophage  $\varphi 80\alpha$ , a pac-type phage. The phage genome is lysogenized in a host bacterial cell, and the phage genome includes a small terminase gene where the pac-site of a pac-type prophage  $\varphi 80\alpha$  has been deleted. A plasmid including a complementary small terminase gene with a native pac-site is transformed into the cell. When the lytic cycle of the lysogenized prophage is induced, the bacteriophage packaging system packages plasmid DNA into progeny bacteriophage structural components, rather than packaging the native bacteriophage DNA. The packaging system thus produces non-replicative transduction particles carrying plasmid DNA.

**[00154]** Figure 3 illustrates an example of the design and function of a pac-site deletion/complementation plasmid packaging system 300, according to an embodiment of the invention. A bacterial cell 301 is lysogenized with a pac-type phage 302 that has its small terminase (terS) gene deleted. The cell is transformed with a rolling circle replication plasmid 303 that includes a small terminase gene that complements the terS gene deletion in the phage. The small terminase gene contains the packaging initiation site sequence, *e.g.*, a pac-site. The plasmid 303 can also include a reporter gene that encodes a reporter molecule.

**[00155]** A protein complex comprising the small terminase and large terminase proteins is able to recognize and cleave a double-stranded DNA molecule at or near the pac-site, and this allows the plasmid DNA molecule to be packaged into a phage capsid. When the prophage in the cell is induced, the lytic cycle of the phage produces the phage's structural proteins 304 and the phage's large terminase protein 305. The complementing plasmid is replicated, and the small terminase protein 306 is expressed. The replicated plasmid DNA 307 containing the terS gene (and the reporter gene) are packaged into phage capsids, resulting in non-replicative transduction particles carrying only plasmid DNA 308. Figure 4 shows an example

of a resulting vector used in the pac-site deletion/complementation plasmid packaging system. Further details about the components and construction of pac-site deletion/complementation plasmid packaging system are in Example 2 below.

[00156]

**B. Pathogenicity Island-Based Packaging System**

[00157] Pathogenicity islands (PTIs) are a subset of horizontally transferred genetic elements known as genomic islands. There exists a particular family of highly mobile PTIs in *Staphylococcus aureus* that are induced to excise and replicate by certain resident prophages. These PTIs are packaged into small headed phage-like particles and are transferred at frequencies commensurate with the plaque-forming titer of the phage. This process is referred to as the SaPI excision replication- packaging (ERP) cycle, and the high-frequency SaPI transfer is referred to as SaPI-specific transfer (SPST) to distinguish it from classical generalized transduction (CGT). The SaPIs have a highly conserved genetic organization that parallels that of bacteriophages and clearly distinguishes them from all other horizontally acquired genomic islands. The SaPI1-encoded and SaPIbov2-encoded integrases are required for both excision and integration of the corresponding elements, and it is assumed that the same is true for the other SaPIs. Phage 80 $\alpha$  can induce several different SaPIs, including SaPI1, SaPI2, and SaPIbov1, whereas  $\varphi$ 11 can induce SaPIbov1 but neither of the other two SaPIs.

[00158] Figure 5 depicts the natural process for genomic island (GI) packaging 500 by a bacteriophage. In nature, a bacterial cell 501 lysogenized with a suitable prophage 503 and carrying a GI 504 can produce phage particles carrying GI concatamers 512. In this process, when the phage is induced into its lytic cycle, the phage genome is excised (not shown) from the bacterial genome 502, which then expresses bacteriophage proteins including capsid constituents 505 and the large terminase protein (TerL) 506. Prophage induction also triggers GI excision via the expression of the GI integrase protein (*int*) 507. In a similar manner to the excised phage genome (not shown), the GI circularizes 508, expresses its own small terminase protein (TerS) 509, and begins to replicate forming a GI concatamer 510. The phage TerL gene and GI TerS gene can then combine bind and cleave the GI concatamer via a pac-site sequence in the GI genome, and the GI concatamer can then be packaged into phage capsids 511 resulting in phage particles carrying GI concatamers 512.

[00159] In natural systems, as depicted in Figure 5, the resulting lysate produced from phage production includes both native phage particles, as well as GI-containing phage

particles. The native phage particles are a result of packaging of the native phage genome due to recognition of the pac-site within phage genome concatamers.

## 1) Genomic Island (GI) Packaging System Design and Function

[00160] Methods of the invention for producing NRTPs include a GI based-packaging system.

[00161] Compared to a plasmid packaging system, the natural GI-packaging system benefits from the fact that the DNA that is packaged is derived from a genomic region within the bacterial genome and thus does not require the maintenance of a plasmid by the bacterial host.

[00162] In some embodiments, the invention includes a bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, wherein the bacterial cell comprises a lysogenized bacteriophage genome lacking a packaging gene, and a genomic island, cryptic phage, or other nucleic acid molecule requiring a bacteriophage (e.g., a heper phage) for mobilization of the nucleic acid molecule and comprising a reporter nucleic acid molecule and a packaging gene. Genomic island-based systems can be based on *S. aureus* Pathogenicity Islands (SaPIs), the *E. coli* criptic phage P4 and helper phage P2, and the *Enterococci* criptic phage P7 and helper phage P1, for example.

[00163] GI-packaging systems can be exploited such that exogenous nucleic acid sequences are packaged by the bacteriophage. This can be accomplished by incorporating such exogenous nucleic acids sequences into the GI.

[00164] In order to eliminate the native phage from this process, the small terminase gene of the prophage can be deleted. The small terminase gene sequence contains the pac-site sequence of the native phage, and this deletion has the effect of preventing the packaging of native phage DNA. In other embodiments, only the pac site of the small terminase gene can be deleted. The GI that will be packaged includes its own pac-site and a small terminase gene that expresses a suitable small terminase protein, and only GI DNA will be amenable for packaging in this system.

[00165] Figure 6 depicts an example of the design and function of a GI-based packaging system 600, according to an embodiment of the invention. In this system, a bacterial cell 601 has its genome lysogenized with a suitable prophage 603 that has its small terminase gene deleted, and the cell's genome 602 carries a GI 604. When the phage is induced into its lytic cycle, the phage genome is excised (not shown) from the bacterial genome 602. The phage

genome expresses bacteriophage proteins, including capsid constituents 605 and the large terminase protein (TerL) 606. Prophage induction also triggers GI excision via the expression of the GI integrase protein (*int*) 607. In a similar manner to the excised phage genome (not shown), the GI circularizes 608 and expresses its own small terminase protein (TerS) 609 and is replicated forming a GI concatamer 610. The phage TerL gene and GI TerS gene can then combine, bind and cleave the GI concatamer via a pac-site sequence in the GI DNA. The GI concatamer can then be packaged into phage capsids 611 resulting in phage particles carrying GI concatamers 612. In this system, phage DNA will not be packaged into phage particles, since it lacks the *terS* gene that contains the phage's pac-site sequence, and thus cannot be recognized by the expressed GI TerS and phage TerL proteins.

**[00166]** When phage particles containing packaged GI DNA are administered to a recipient cell, the phage will bind to the recipient cell's surface and then introduce the packaged GI DNA concatamer into the cell. Once inside the cell, the GI can again express its integrase protein, and the GI can then integrate into its specific site in the recipient cell's genome. If exogenous DNA sequences are included in the GI prior to packaging, the packaging system thus allows for delivering exogenous DNA sequences to a recipient cell and integrating these exogenous DNA sequences into the recipient cell's genome.

## 2) GI-Based Packaging System Lacking Integrase

**[00167]** In another embodiment, the packaging system described above is designed such that packaged GI DNA cannot integrate into a recipient cell's genome. This can be accomplished by deleting the integrase gene in the GI and complementing the deletion by causing the expression of the integrase gene in trans from the GI. In this manner, the integrase protein is available for excision of the GI in the packaging host cell, and the GI DNA that has been packaged in a bacteriophage does not contain the integrase gene and cannot express the integrase protein, thus preventing integration of the delivered GI.

**[00168]** Figure 7 depicts the design and function of a GI-based packaging system that lacks the *int* gene 700, according to an embodiment of the invention. In this system, a bacterial cell 701 is lysogenized with a suitable prophage that has had its small terminase gene deleted 703. The cell's genome 702 carries a GI that has its integrase (*int*) gene deleted 704 and also carries the deleted *int* gene operatively linked to a suitable promoter 705. The *int* gene can thus express the integrase protein (Int) in trans from the GI 706. When the phage is induced into its lytic cycle, the phage genome is excised (not shown) from the bacterial genome 702, which then expresses bacteriophage proteins including capsid constituents 707 and the large terminase protein (TerL) 708. Prophage induction also triggers

GI excision via the expression of the integrase protein 707. In a similar manner to the excised phage genome (not shown), the excised GI circularizes 709, expresses its own small terminase protein (TerS) 710, and begins to replicate forming a GI concatamer 711. The phage TerL gene and GI TerS gene can then combine, bind and cleave the GI concatamer via a pac-site sequence in the GI DNA, and the GI concatamer can then be packaged into phage capsids 712 resulting in phage particles carrying GI concatamers 713. In this system, phage DNA will not be packaged since it lacks the terS gene that contains the phage's pac-site sequence and thus cannot be recognized by the expressed GI TerS and phage TerL proteins.

**[00169]** When phage particles containing packaged GI DNA lacking the *int* gene are administered to a recipient cell, the phage will bind to the recipient cell's surface and then introduce the packaged GI DNA concatamer into the cell. Once inside the cell, the GI cannot express its integrase protein due to the lack of the integrase gene and the GI cannot then integrate into its specific site in the recipient cell's genome. If exogenous DNA sequences are included in the GI prior to packaging, the packaging system thus allows for delivering exogenous DNA sequences to a recipient cell and the delivered DNA sequences do not integrate into the recipient cell's genome at the specific site for GI integration.

### **3) Design and Function of SaPIbov2-Based Packaging Lacking Integrase**

**[00170]** In some embodiments, the method of producing NRTPs employ a GI SaPIbov2 and a bacteriophage  $\phi$ 11 in a GI-based packaging system. Alternative embodiments can employ other SaPI GI's and other suitable bacteriophages, including the SaPI's SaPI1, SaPI2, SaPIbov1, and SaPIbov2 along with the bacteriophage 80 $\alpha$ , and the SaPI's SaPIbov1 and SaPIbov2 along with the bacteriophage  $\phi$ 11. Based on the description below, one of skill in the art would know how to develop a GI-based packaging system that does not lack the *int* gene, as described in Section II A.

**[00171]** Figure 8 depicts the design and function of a SaPIbov2-based packaging system 800 that lacks the *int* gene, according to an embodiment of the invention. In this system, a *S. aureus* cell 801 is lysogenized with  $\phi$ 11 that has its small terminase gene deleted 803. The cell's genome 802 carries SaPIbov2 that has its integrase (*int*) gene deleted 804 and also carries the deleted *int* gene operatively linked the constitutively expressed *PclpB* gene promoter 805. The *int* gene can express the integrase protein (Int) in trans from SaPIbov2 806. When the phage is induced into its lytic cycle, the phage genome is excised (not shown) from the bacterial genome 802, which then expresses bacteriophage proteins including capsid constituents 807 and the large terminase protein (TerL) 808. Prophage induction also triggers

SaPIbov2 excision via the expression of the integrase protein 806. In a similar manner to the excised phage genome (not shown), the excised SaPIbov2 circularizes 809, expresses its own small terminase protein (TerS) 810 and begins to replicate forming a SaPIbov2 concatamer 811. The phage TerL gene and SaPIbov2 TerS gene can then combine bind and cleave the SaPIbov2 concatamer via a pac-site sequence in the SaPIbov2 DNA and the SaPIbov2 concatamer can then be packaged into phage capsids 812 resulting in phage particles carrying SaPIbov2 concatamers 813. In this system, phage DNA will not be packaged since it lacks the terS gene that contains the phage's pac-site sequence and thus cannot be recognized by the expressed SaPIbov2 TerS and phage TerL proteins.

#### IV. Reporters

**[00172]** In some embodiments, the NRTPs and constructs of the invention comprise a reporter nucleic acid molecule including a reporter gene. The reporter gene can encode a reporter molecule, and the reporter molecule can be a detectable or selectable marker. In certain embodiments, the reporter gene encodes a reporter molecule that produces a detectable signal when expressed in a cell.

**[00173]** In certain embodiments, the reporter molecule can be a fluorescent reporter molecule, such as, but not limited to, a green fluorescent protein (GFP), enhanced GFP, yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), blue fluorescent protein (BFP), red fluorescent protein (RFP) or mCherry, as well as near-infrared fluorescent proteins.

**[00174]** In other embodiments, the reporter molecule can be an enzyme mediating luminescence reactions (luxA, luxB, luxAB, luc, ruc, nluc, *etc.*). Reporter molecules can include a bacterial luciferase, a eukaryotic luciferase, an enzyme suitable for colorimetric detection (lacZ, HRP), a protein suitable for immunodetection, such as affinity peptides (His-tag, 3X-FLAG), a nucleic acid that function as an aptamer or that exhibits enzymatic activity (ribozyme), or a selectable marker, such as an antibiotic resistance gene (ampC, tet(M), CAT, erm). Other reporter molecules known in the art can be used for producing signals to detect target nucleic acids or cells.

**[00175]** In other aspects, the reporter molecule comprises a nucleic acid molecule. In some aspects, the reporter molecule is an aptamer with specific binding activity or that exhibits enzymatic activity (*e.g.*, aptazyme, DNAzyme, ribozyme).

**[00176]** Reporters and reporter assays are described further in Section V herein.

## V. NRTPs and Reporter Assays

### A. Inducer Reporter Assay

[00177] The invention comprises methods for the use of NRTPs as reporter molecules for use with endogenous or native inducers that target gene promoters within viable cells. The NRTPs of the invention can be engineered using the methods described in Section III and below in Examples 1-6.

[00178] In some embodiments, the method comprises employing a NRTP as a reporter, wherein the NRTP comprises a reporter gene that is operably linked to an inducible promoter that controls the expression of a target gene within a target cell. When the NRTP that includes the reporter gene is introduced into the target cell, expression of the reporter gene is possible via induction of the target gene promoter in the reporter nucleic acid molecule.

[00179] Figure 9 depicts a genomic locus of a target cell 900 with two genes, a gene encoding an inducer 902 and a target gene 903. Also depicted is a reporter nucleic acid molecule 904 that includes a reporter gene 905 that is operatively linked to the promoter 906 of the target gene of the target cell. The reporter nucleic acid molecule 904 can be introduced into the cell via a NRTP. In the native cell, when the inducer gene 902 is expressed and produces the inducer protein 907, the inducer protein 907 is able to induce the target gene promoter 906 that is operatively linked to the target gene, thus causing the expression of the target gene and the production of the target gene product 908.

[00180] When the reporter nucleic acid molecule 904 is present within the target organism, the inducer 907 is also able to induce the target gene promoter 906 present within the reporter nucleic acid molecule 904, thus causing the expression of the reporter gene 905 resulting in the production of a reporter molecule 909 capable of generating a detectable signal.

[00181] Thus, the production of a detectable signal from the reporter molecule 909 is indicative of the presence of the cell, based on the presence of the inducer protein 907 within a target cell.

#### 1) VanR Reporter System

[00182] In one embodiment, the reporter system includes NRTP comprising a reporter nucleic acid molecule (e.g., plasmid). The reporter nucleic acid molecule can be constructed for detecting VanR, the inducer of the promoter of the vancomycin resistance (vanA) gene in *Enterococcus faecium* (or *E. faecalis*). The reporter plasmid carries a reporter gene that is operatively linked to the vanA gene promoter.

[00183] Figure 10 outlines the design and function of a VanR reporter system. Figure 10 depicts a region of the transposon Tn1546 1001 that may be present in *E. faecium*. The

Tn1546 transposon can include the vanR inducer gene 1002 and the vanA target gene 1003. Also depicted in the figure is a reporter nucleic acid molecule 1004 that can be packaged in a NRTP and introduced into the cell. The reporter nucleic acid molecule 1004 includes a reporter gene 1005 that is operatively linked to a promoter  $P_H$  1006 that controls the expression of the vanHAX operon that includes the vanA gene. In the native cell, when the vanR gene 1002 is expressed and produces the VanR protein 1007, VanR is able to induce  $P_H$  1006 in the Tn1546 transposon, thus causing the expression of the vanA gene and thus producing the VanA protein 1008. When the reporter nucleic acid molecule 1003 (vector) is present within the target organism, VanR is also able to induce  $P_H$  1006 within the reporter nucleic acid molecule 1003, thus causing the expression of a reporter molecule 1009. Thus, the production of a reporter molecule is indicative of the presence of VanR within a target cell.

**[00184]** Examples of promoters that are suitable for the development of a VRE assay include: the vanA gene promoter and a vanB gene promoter. Arthur, M., et al., *The VanS sensor negatively controls VanR-mediated transcriptional activation of glycopeptide resistance genes of Tn1546 and related elements in the absence of induction*. J. Bacteriol., 1997. **179**(1): p. 97-106.

## 2) TcdD reporter system

**[00185]** In another embodiment of this system, a reporter nucleic acid molecule is introduced into a cell using a NRTP. The reporter nucleic acid molecule can be constructed for detecting TcdD, the inducer of the promoters of the toxins A and B genes (tcdA and tcdB, respectively) of *C. difficile*. The reporter nucleic acid molecule includes a reporter gene that is operatively linked to the tcdA gene promoter.

**[00186]** Figure 11 outlines the design and function of a TcdD reporter system, according to an embodiment of the invention. Figure 11 depicts a region of the transposon PaLoc 1101 that may be present in *C. difficile*. The PaLoc transposon may contain the tcdD gene 1102 and the tcdA target gene 1103. Also depicted in the figure is a reporter nucleic acid molecule 1104 (e.g., vector) that is introduced into the cell using a NRTP. The reporter nucleic acid molecule 1104 includes the reporter gene 1105 that operatively linked to the tcdA gene promoter ( $P_{tcdA}$ ) 1106.

**[00187]** In the native cell, when the tcdD gene is expressed and produces the TcdD protein 1107, TcdD is able to induce  $P_{tcdA}$  1106 in the PaLoc transposon 1101, thus causing the expression of the tcdA gene 1103 and thus producing the toxin A protein 1108.

**[00188]** When the reporter nucleic acid molecule 1104 is present within the target organism, TcdD is also able to induce  $P_{tcdA}$  1106 within the reporter vector, thus causing the expression of a reporter molecule 1109. Thus, the production of a reporter molecule 1109 is indicative of the presence of TcdD within a target cell.

**[00189]** Examples of promoters suitable for the development of a *C. difficile* assay include: the *tcdA* gene promoter and the *tcdB* gene promoter. Karlsson, S., et al., *Expression of Clostridium difficile Toxins A and B and Their Sigma Factor TcdD Is Controlled by Temperature*. Infect. Immun., 2003. 71(4): p. 1784-1793.

**[00190]** Target cells and inducers: Target cells can include eukaryotic and prokaryotic cell targets and associated inducers.

**[00191]** Vector delivery systems: The delivery of the vector containing the recombinant DNA can be performed by abiologic or biologic systems. Including but not limited to liposomes, virus-like particles, transduction particles derived from phage or viruses, and conjugation.

### 3) Bacteriophage-based SarS reporter system

**[00192]** In another embodiment of the invention, a reporter nucleic acid molecule is constructed for detecting SarS, the inducer of the promoter of the Protein A gene (*spa*) in *S. aureus*. The reporter nucleic acid molecule can be introduced into the cell in a NRTP and includes the bacterial luciferase genes *luxA* and *luxB* operatively linked to the *spa* gene promoter ( $P_{spa}$ ). The reporter nucleic acid molecule is delivered to *S. aureus* via a NRTP, for example. If SarS is present in the cell, it will induce the expression of the *luxAB* genes, thus producing luciferase enzyme that is capable of generating a luminescent signal.

**[00193]** Figure 12 outlines the design and function of a SarS reporter system, according to one embodiment of the invention. Figure 12 depicts a region of the *S. aureus* genome 1201 that contain the *sarS* gene 1202 and *spa* gene 1203. Also depicted in the figure is a reporter nucleic acid molecule (e.g., vector) 1204 delivered by NRTP to the cell and that includes the *luxAB* reporter genes 1205 that operatively linked to the promoter  $P_{spa}$  1206 that controls the expression of the *spa* gene 1203.

**[00194]** In the native cell, when the *sarS* gene 1202 is expressed, producing SarS protein 1207, the protein is able to induce  $P_{spa}$  1206 in the *S. aureus* genome transposon, thus causing the expression of the *spa* gene 1203 and producing the Protein A 1208.

**[00195]** When the reporter nucleic acid molecule 1204 is present within the target organism, SarS 1207 is also able to induce  $P_{spa}$  1206 within the reporter nucleic acid molecule 1204, thus causing the expression of *luxAB* resulting in the production of the luciferase

enzyme 1209 that can generate a luminescent signal. Thus, the production of luciferase is indicative of the presence of SarS within a target cell.

### ***B. Enzyme Reporter Assay***

**[00196]** The invention comprises a system for the detection of intracellular enzymes within viable cells that employs caged substrate molecules that can be un-caged by a target intracellular enzyme, according to an embodiment of the invention.

**[00197]** Figure 13 depicts the design and function of an intracellular enzyme detection system. A reporter molecule-expressing vector 1301 is delivered to a target cell 1302 with a NRTP (not shown). The reporter molecule-expressing vector 1301 is able to penetrate the target cell 1302 via the NRTP and deliver a reporter molecule gene 1303 into the target cell 1302, and a reporter molecule 1304 can then be expressed from the reporter molecule gene 1303. A caged substrate 1305 is also added to the target cell 1302 and is able to penetrate into the target cell 1302. If a target intracellular enzyme 1307 is present in the target cell 1306, the enzyme 1307 is able to remove the caging component of the caged substrate 1305, thus producing an un-caged substrate 1308. The un-caged substrate 1308 can then react with the reporter molecule 1304 inside of the cell 1302, and the product of this reaction results in a detectable signal 1309.

**[00198] Target cells and enzymes:** Target cells can include eukaryotic and prokaryotic cell targets and associated enzymes, including, for example,  $\beta$ -lactamases in *S. aureus*.

**[00199] Vector delivery systems:** The delivery of the vector containing the recombinant DNA can be performed by abiologic or biologic systems. Including but not limited to liposomes, virus-like particles, transduction particles derived from phage or viruses, and conjugation.

**[00200] Reporter molecules and caged substrates:** Various reporter molecules and caged substrates can be employed as those described in Daniel Sobek, J.R., Enzyme detection system with caged substrates, 2007, Zymera, Inc.

#### **1) Bacteriophage-based $\beta$ -lactamase Reporter**

**[00201]** In one embodiment, a reporter molecule-expressing vector can be carried by a NRTP, such that the vector can be delivered into a bacterial cell. The reporter molecule to be expressed can be Renilla luciferase, and the caged substrate can be Renilla luciferin that is caged, such that a  $\beta$ -lactamase enzyme that is endogenous to the target cell is able to cleave the caging compound from the caged luciferin and release un-caged luciferin.

**[00202]** Figure 14 depicts the design and function of a  $\beta$ -lactamase enzyme detection system, according to an embodiment of the invention. A Renilla luciferase-expressing vector carried by a bacteriophage-based NRTP 1401 is added to a target *S. aureus* cell 1402. The Renilla luciferase-expressing vector is able to penetrate the target cell 1402 using a NRTP comprising the vector. The NRTP delivers the Renilla luciferase gene 1403 into the target cell 1402, and Renilla luciferase 1404 can then be expressed from its gene. Caged Renilla luciferin 1405 is also added to the target cell 1402 and is able to penetrate into the target cell 1402. If an intracellular  $\beta$ -lactamase 1407 is present in the target cell 1402, the enzyme is able to remove the caging component of the caged luciferin 1406, thus producing an uncaged luciferin 1408. The un-caged luciferin 1408 can then react with the Renilla luciferase 1404 inside of the cell 1402, and the product of this reaction results in luminescence 1409.

**[00203]** In this manner, when a target cell that contains the  $\beta$ -lactamase is exposed to the NRTP and caged luciferin, the cell will exhibit a luminescent signal that is indicative of the presence of the  $\beta$ -lactamase present in the cell.

### ***C. Intracellular Molecule Reporter***

**[00204]** The invention includes a system for the detection of intracellular molecules within viable cells that employs switchable molecules capable of generating a detectable signal upon their binding to a target molecule.

**[00205]** Figure 15 depicts the design and function of a switchable molecule (SM)-based intracellular molecule detection system. A SM-expressing vector 1501 is delivered to a target cell 1502 in a NRTP. The SM-expressing vector 1501 is able to penetrate the target cell 1502 and deliver a SM gene 1503 into the target cell 1502. A SM protein 1504 can then be expressed from the SM gene 1503. The SM protein 1504 can then bind to a target molecule 1505 inside of the cell and thus forms an SM-target molecule complex 1506. The binding of the SM 1504 to the target molecule 1505 results in a conformational change in the SM 1504 that makes the bound SM amenable to binding of a substrate. A substrate 1508 is added to the cell 1507 and is able to penetrate into the cell 1502. Bound SM inside of the cell 1502 is able to also bind the substrate, thus forming a SM-target molecule-substrate complex 1509. Finally, the binding of the substrate 1508 by the target molecule-bound SM has the effect of producing a detectable signal 1510. Thus a detectable signal generated by the system is indicative of the presence of a target molecule inside of a cell.

**[00206] Target cells and molecules:** Various eukaryotic and prokaryotic cell targets can be employed and switchable aptamer-based SM's can be designed to target various nucleic

acid and amino acid-based intracellular molecular targets as described in Samie Jaffrey, J.P., *Coupled recognition/detection system for in vivo and in vitro use*, 2010, Cornell University.

**[00207] Vector delivery systems:** The delivery of the vector containing the recombinant DNA can be performed by abiologic or biologic systems. Including but not limited to liposomes, virus-like particles, transduction particles derived from phage or viruses, and conjugation.

**1) Non-Replicative Transduction Particle/Switchable Aptamer-Based Intracellular Molecule Reporter System**

**[00208]** In one example of this method, a switchable molecule-expressing vector can be carried by a bacteriophage-based transduction particle such that the vector can be delivered into a bacterial cell. The switchable molecule to be expressed can be a switchable aptamer that is designed to undergo a conformational change upon its binding to an intracellular target molecule. The conformational change allows the aptamer to then bind a fluorophore that exhibits enhanced fluorescence when bound by the aptamer.

**[00209]** Figure 16 depicts the design and function of a bacteriophage/switchable-aptamer (SA)-based intracellular molecule reporter system. A SA-expressing vector carried by a NRTP 1601 is added to a target cell 1602. The NRTP 1601 is able to deliver the SA-expressing vector and the SA-expressing gene 1603 into the target cell 1602. An SA protein 1604 can then be expressed from the SA gene 1603. The SA protein 1604 can then bind to a target molecule 1605 inside of the cell and thus form an SA-target molecule complex 1606. The binding of the SA 1604 to the target molecule 1605 results in a conformational change in the SA that makes the bound SA amenable to binding of a fluorophore 1608. A fluorophore 1607 is added to the cell and is able to penetrate into the cell 1608. Bound SA inside of the cell is able to also bind the fluorophore thus forming an SA-target molecule-fluorophore complex 1609. Finally, the binding of the fluorophore by the target molecule-bound SA has the effect of enhancing the fluorescence of the fluorophore 1610. Thus, a detectable fluorescent signal generated by the system is indicative of the presence of a target molecule inside of a cell.

***D. Transcript Reporter Assay***

**[00210]** The invention comprises a reporter assay comprising an antisense RNA-based method for detecting target transcripts within viable cells by causing the expression of a reporter molecule if a target transcript is present within a cell.

**[00211]** Certain intracellular methods in the art for inhibiting gene expression employ small interfering RNA, such as double-stranded RNA (dsRNA), to target transcribed genes in

cells. The dsRNA comprise antisense and sense strands that are delivered into or expressed in cells, and the strands of the dsRNA act via a trans-acting inhibition mechanism, where one strand (typically the antisense strand) binds to a target gene sequence (RNA transcript) and prevents expression of the target gene sequence. Double-stranded RNA molecules have been shown to block (knock down) gene expression in a highly conserved regulatory mechanism known as RNA interference (RNAi). WO 99/32619 (Fire *et al.*) discloses the use of a dsRNA of at least 25 nucleotides in length to inhibit the expression of genes in *C. elegans*. dsRNA has also been shown to degrade target RNA in other organisms, including plants (see, e.g., WO 99/53050, Waterhouse *et al.*; and WO 99/61631, Heifetz *et al.*), Drosophila (see, e.g., Yang, D., *et al.*, Curr. Biol. (2000) 10:1191-1200), and mammals (see WO 00/44895, Limmer; and DE 101 00 586.5, Kreutzer *et al.*). However, binding of a strand of the dsRNA to the target gene can be non-specific. If a similar mechanism were to be applied to a detection system, this non-specific binding can result in high false positive rates, which make it unsuitable for the development of clinically useful detection systems.

**[00212]** Previous trans-acting inhibition mechanisms have been shown to be unsuitable for development of clinically useful detection systems. For example, some methods result in high levels of non-specific signals and up to 90% false positive rate, when achieving a 90% sensitivity of the assay. See U.S. Patent No. 8,329,889. Certain methods for post-transcriptional regulation of gene expression have been developed that use a *cis*-repressed marker transcript, such as a green fluorescent protein marker, where the ribosomal binding site of the marker is blocked by the *cis*-repressing sequence, along with a trans-activating RNA transcript. When the trans-activating RNA transcript binds to the *cis*-repressed marker transcript, the hairpin structure of the *cis*-repressed marker transcript is altered, and the upstream ribosome binding site of the marker gene is exposed, allowing transcription and expression of the marker gene. However, these methods have not previously been used for the detection of endogenous transcripts, nor successful beyond a basic switching mechanism for controlling expression of genes in cells.

### **1) Nucleic Acid Molecule Interactions and Mechanisms**

**[00213]** The methods of the invention take advantage of the transcript-level regulation mechanisms, including antisense RNA (asRNA) mechanism in cells, to deliver nucleic acid molecules into cells. The antisense mechanism includes all forms of sequence-specific mRNA recognition leading to reduced, eliminated, increased, activated, or otherwise altered expression of a target transcript. See Good, L., *Translation Repression By Antisense Sequences*. Cellular and Molecular Life Sciences, 2003. 60(5): p. 854-861, and Lioliou, E.,

*RNA-mediated regulation in bacteria: from natural to artificial systems*, New Biotechnology. 2010. 27(3): p. 222-235. Naturally occurring asRNAs are found in all three kingdoms of life, and they affect messenger RNA (mRNA) destruction, repression and activation, as well as RNA processing and transcription. See Sabine, B., *Antisense-RNA regulation and RNA Interference*. Biochimica et Biophysica Acta (BBA) – Gene Structure and Expression, 2001. 1575(1-3): p. 15-25. This mechanism has been exploited in inhibiting protein synthesis for therapeutic applications.

**[00214]** Antisense RNA is a single-stranded RNA that is complementary to a messenger RNA (mRNA) strand transcribed within a cell. asRNA may be introduced into a cell to inhibit translation of a complementary mRNA by base pairing to it and physically obstructing the translation machinery. Antisense RNA anneal to a complementary mRNA target sequence, and translation of the mRNA target sequence is disrupted as a result of steric hindrance of either ribosome access or ribosomal read through.

**[00215]** The antisense RNA mechanism is different from RNA interference (RNAi), a related process in which double-stranded RNA fragments (dsRNA, also called small interfering RNAs (siRNAs)) trigger catalytically mediated gene silencing, most typically by targeting the RNA-induced silencing complex (RISC) to bind to and degrade the mRNA. Annealing of a strand of the dsRNA molecule to mRNA or DNA can result in fast degradation of duplex RNA, hybrid RNA/DNA duplex, or duplex RNA resembling precursor tRNA by ribonucleases in the cell, or by cleavage of the target RNA by the antisense compound itself.

**[00216]** The RNAi pathway is found in many eukaryotes and is initiated by the enzyme Dicer, which cleaves long double-stranded RNA (dsRNA) molecules into short double stranded fragments of ~20 nucleotides that are called siRNAs. Each siRNA is unwound into two single-stranded RNAs (ssRNA), namely the passenger strand and the guide strand. The passenger strand is degraded, and the guide strand is incorporated into the RNA-induced silencing complex (RISC). In post-transcriptional gene silencing, the guide strand base pairs with a complementary sequence in a messenger RNA molecule, and cleavage is induced by a protein called Argonaute, the catalytic component of the RISC complex.

**[00217]** In regards to the nucleic acid interactions of the mechanisms of the invention, interactions between a reporter transcript and a target transcript can rely on base pairings between loops present in both transcripts (e.g., “kissing complexes”), or between a loop and a single-stranded (ss) region. In some cases, the kissing complex formation suffices for

mediating the desired effect of the interaction, and in other cases, propagation of the primary contacts will lead to an interaction resulting in the desired effect.

## 2) Mechanisms for Cis-Repression and Trans-Activation of Translation of a Reporter Construct via Transcript-Level Regulation

**[00218]** The following description illustrates transcript reporter systems based on various repression/activation mechanisms that can be used, according to embodiments of this invention. In each of Figures 17-20, a vector includes a reporter construct comprising a reporter sequence, and the regions on the reporter construct are shown in each of the figures, including regions that can be targeted for repression by a cis-repressing sequence. The description below provides non-limiting examples of various inhibition mechanisms, including transcription attenuation, translation attenuation, and destabilization of the transcript, and various activation mechanisms including conformational changes and cleavage.

**[00219]** Figure 17 depicts an example of a system 1700 that uses a cis-repression mechanism that can target the 5' UTR (untranslated region) 1701 of the reporter sequence 1702 on a reporter transcript 1703. The regions within the reporter sequence 1702 (5'UTR (1701), RBS, Coding Region and 3'UTR) are also shown. The cis-repressing sequence 1705 is upstream of the reporter sequence and up to the 5' UTR 1701 of the reporter sequence. An RNA polymerase 1704 transcribes the sequence of the reporter construct 1703 from the vector 1706.

**[00220]** At some point during transcription, the transcription process is stopped by the formation of a transcription termination (TT) stem-loop structure 1707 in the reporter transcript 1703, due to an interaction within the transcribed cis-repressing sequence 1705. The transcription termination 1707 structure stops 1708 the RNA polymerase 1704 from transcribing the vector 1706. In some embodiments, a transcription termination protein (*e.g.*, NusA in *E. coli*) binds to RNA polymerase and/or to the transcription termination 1707 structure to cease transcription of the reporter construct.

**[00221]** When a target transcript 1709 is present in the cell, the target transcript 1709 binds to the reporter transcript 1703. In some embodiments, the binding between the target transcript and the reporter transcript is by base pairing of the nucleotides in each sequence. The interaction between the target transcript 1709 and the reporter transcript 103 causes the transcription termination (TT) stem-loop structure 1707 to be cleaved 1710. Cleavage of the reporter transcript 1703 can occur by a cellular enzyme, such as RNase III, for example. In

this case, the secondary structure of a target transcript is analyzed for the presence of an RNase III consensus sequence among the ssRNA regions of the secondary structure, for example 5'-nnWAWGNNUUN-3' (SEQ ID NO: 20) or 5'-NAGNNNNCWUWnn-3' (SEQ ID NO: 21) where "N" and "n" are any nucleotide and "W" is A or U and "N" indicates a relatively strict requirement for Watson-Crick base pairing, while "n" indicates a minimal requirement for base pairing. When such a consensus sequence is found on a target transcript, the loop of the transcription termination structure 1707 can be designed to be complementary to said RNase III consensus sequence such that when the ssRNA in each RNA molecule hybridize, the RNase III cleavage site is formed allowing for cleavage of the transcription termination structure 1707. In the *mecA* transcript, loop T23, starting at nucleotide 1,404, has the sequence CAGAUAAACAUUUU (SEQ ID NO: 22) that is suitable for such an approach.

**[00222]** In some embodiments, a cleavage site is engineered in the reporter construct, such that the reporter transcript is cleaved after transcription. The cleavage, in the example provided, can occur immediately adjacent to the location of the loop in the transcription terminator structure. Transcription is re-initiated 1711 by the RNA polymerase 104. Cleavage of the transcription termination (TT) stem-loop structure 1707 allows the remainder of the reporter sequence 1702 to be transcribed and subsequently translated. This results in the production of a detectable or selectable marker from the translated reporter molecule.

**[00223]** In prokaryotes, the transcription termination structure 1707 involves a Rho-independent mechanism with a stem-loop structure that is 7-20 base pairs in length, rich in cytosine-guanine base pairs and is followed by a chain of uracil residues. NusA binds to the transcription termination stem-loop structure 1707 causing RNA polymerase to stall during transcription of the poly-uracil sequence. Weak Adenine-Uracil bonds lower the energy of destabilization for the RNA-DNA duplex, allowing it to unwind and dissociate from the RNA polymerase. In eukaryotes, the transcription termination structure 1707 is recognized by protein factors and involves cleavage of the new transcript followed by polyadenylation.

**[00224]** Figure 18 shows an example of a system 1800 for detecting the presence of a target transcript in a cell that is based on a cis-repression mechanism targeting the ribosome binding site (RBS) 1801 of the reporter sequence 1702 in the reporter transcript 1703. The RBS 1801 is a sequence of mRNA that is bound by the ribosome 1802 when initiating protein translation. The cis-repressing sequence 1705 is designed to bind to the RBS 1801 (*e.g.*, the cis-repressing sequence 1705 is complementary to the RBS sequence 1801). The RBS 1801

binds to the cis-repressing sequence 1705 and becomes sequestered (inaccessible by a ribosome 1802), preventing the translation of the reporter transcript 1703. When a target

transcript 109 from the cell binds to the reporter transcript 1703, the target transcript 1709 has a higher binding affinity for the RBS sequence 1801, and a conformational change occurs in the reporter transcript 1703 in a manner that releases the binding between the cis-repressing 1705 sequence and the RBS sequence 1801. This allows the ribosome 1802 to bind to the RBS 1801, thereby allowing for translation of the reporter transcript 1703.

**[00225]** Figure 19 illustrates an exemplary system 1900 for detecting the presence of a target transcript in a cell that is based on a cis-repression mechanism targeting the coding region (“AUG”) 1901 of the reporter sequence 1702 in the reporter transcript 1703. The cis-repressing sequence 1705 is constructed such that it binds with (e.g., complementary to) the coding region 1901 of the reporter sequence 1702. The “AUG” start codon is shown as part of the coding region 1901. The binding of the cis-repressing sequence 1705 and the coding region 1901 results in a conformation that leads to cleavage 1902 of the reporter construct 1703. Cleavage of the reporter transcript 1703 prevents translation.

**[00226]** When a target transcript 1709 is present in the cell, the target transcript 1709 binds to the cis-repressing sequence 1705 in a manner that causes a conformational change in the reporter transcript 1703. This conformational change prevents or removes the interaction between the cis-repressing sequence 1705 and the coding region 1901 of the reporter sequence 1702, thereby allowing for translation of the reporter sequence 1702.

**[00227]** Figure 20 illustrates an example system 2000 for detecting the presence of a target transcript in a cell that is based on a repression mechanism using an unstable reporter transcript 2001. The reporter transcript 2001 is designed to be unstable such that it forms an unstable conformation that prevents the translation of the reporter transcript 2001. A reporter transcript 2001 is defined to be unstable if it is prone to rapid degradation due to a variety of factors, such as activity of exosome complexes or a degradosome. A target transcript 1709 in the cell binds to a portion of the unstable reporter transcript 2001. In this example, the portion responsible for destabilizing the transcript is located in the 3' UTR 2005 of the reporter sequence, and the 3' UTR 2005 acts like the cis-repressing sequence of the reporter construct 1703. The binding of the target transcript 1709 with the 3' UTR 2005 of the reporter sequence results in a cleaving event 2003 that stabilizes the reporter transcript 2001 and allows for translation 2004 of the reporter transcript 2001. Cleavage occurs upon binding of the target transcript 1709 and serves to remove the portion of sequence that is responsible for destabilizing the transcript. In this example, the target transcript 1709 binds to the 3' UTR 405 of the reporter sequence, but the system 400 can also be designed such that binding and cleavage occurs in the 5' UTR, upstream of the 5' UTR, or downstream of the 3' UTR.

Binding and cleavage can occur anywhere outside of regions necessary for translation of the reporter sequence 1702.

**[00228]** In some embodiments, the cis-repressing sequence itself comprises two sequences that can bind to each other (e.g., complementary to each other), and the conformation of the reporter transcript that results from the binding of the two sequences of the cis-repressing sequence prevents translation of the reporter sequence in the reporter transcript.

### **3) Naturally Occurring and Synthetic Systems for Repression/Activation Mechanisms**

**[00229]** Several naturally occurring and synthetically produced transcript-level mechanisms have been described that demonstrate the individual mechanisms (i.e., conformational change and cleavage) employed in each of the examples illustrated in Figures 17-20.

**[00230]** Transcription termination has been observed in antisense RNA (asRNA)-mediated transcriptional attenuation. In one example, two loop-loop interactions between RNAIII/repR mRNA are subsequently followed by the formation of a stable duplex. This complex stabilizes a Rho-independent terminator structure to arrest elongation by RNA polymerase (RNAP).

**[00231]** The RBS sequestration mechanism has been described via the development of a synthetic riboswitch system. In this system, a sequence complementary to a RBS is placed upstream of the RBS, allowing the presence of a linker sequence between the two regions. After transcription of the mRNA, the two complementary regions hybridize, creating a hairpin that prevents docking of the ribosome. To activate translation, a synthetic trans-activating RNA carrying the RBS sequence binds to the hybridized RNA, allowing the RBS to be exposed and available for translation.

**[00232]** The prevention of translation due to the cleaving of RNA has also been described in a natural system where the asRNA MicC targets a sequence inside the coding region of ompD mRNA. The interaction, which is promoted by Hfq, causes the cleaving of the mRNA by RNase E.

**[00233]** Yet another natural mechanism demonstrates a cleaving event to activate translation rather than inhibiting it. The *E. coli* GadY asRNA targets the intergenic region between two genes of the gadXW operon. Following the formation of a stable helix between GadY and the 3'UTR of gadX, an RNase cleavage occurs in the transcript and stabilizes gadX transcript allowing for its translation.

#### 4) Mechanism of Conformational Change by Cis-Repression of the Reporter Sequence and by Binding of a Target Transcript

**[00234]** The general mechanisms employed in the invention are intermolecular nucleic acid molecule interactions that may result in two subsequent mechanisms: (1) a conformational change in the secondary structure of the nucleic acid molecules, and (2) a cleaving event. Described herein are methods for designing reporter transcripts that can undergo a conformational change between a cis-repressed conformation and a de-repressed conformation, such that the conformational change is induced by binding of a target transcript to the reporter transcript.

**[00235]** As described above, a reporter transcript can comprise a reporter sequence and be designed such that translation of the reporter gene sequence is blocked by cis-repression of the ribosome binding site (RBS) of the reporter gene.

**[00236]** In some embodiments, the following tools can be used for designing the reporter transcripts of the invention.

**[00237]** 1) RNA secondary structure is calculated using secondary structure program, such as Mfold available at a server maintained by The RNA Institute College of Arts and Sciences, University at Albany, State University of New York (Mfold web server for nucleic acid folding and hybridization prediction. Nucleic Acids Res. 31 (13), 3406-15, (2003)) (<http://mfold.rna.albany.edu/?q=mfold/RNA-Folding-Form>).

**[00238]** 2) Intermolecular RNA interactions are calculated using a software program such as RNA-RNA InterACTion prediction using Integer Programming (RactIP) available at a server maintained by the Graduate School of Information Science, Nara Institute of Science and Technology (NAIST), Department of Biosciences and Informatics, Keio University Japan (<http://rna.naist.jp/ractip/>).

**[00239]** 3) RNA secondary structure is visualized using Visualization Applet for RNA (VARNA) (<http://varna.lri.fr/>), which is a Java lightweight Applet dedicated to drawing the secondary structure of RNA.

**[00240]** A secondary structure of the target transcript can be generated based on the lowest energy conformation calculated by MFold and visualized with VARNA.

**[00241]** ssRNA regions or target regions can be identified within the target transcript that can be ideal for binding to a reporter transcript. In some instances, the secondary structure of the target transcript includes a consensus sequence or loop sequence that can bind to a portion of the reporter sequence. For example, in the *mecA* transcript of methicillin-resistant *S. aureus*, there is a terminal loop that includes a consensus YUNR sequence (“UUGG”) that

can be used to bind to a cis-repressing sequence of a reporter transcript. Analysis of the secondary structure of the target transcript can reveal these one or more ssRNA regions that can be suitable for binding to a cis-repressing sequence. The cis-repressing sequence of the reporter transcript can then be designed to bind to these one or more ssRNA regions.

**[00242]** In some embodiments, the cis-repressing sequence can be designed to bind to the RBS of the reporter sequence in the reporter transcript and form a stem-loop structure within the reporter transcript, such that the cis-repressing sequence blocks binding of an RNA polymerase to the RBS of the reporter sequence. Upon binding of the cis-repressing sequence to the ssRNA region of the target transcript, the RBS of the reporter sequence can be exposed and translation of the reporter sequence can be initiated.

**[00243]** In some embodiments, the cis-repressing sequence of the reporter transcript can be designed to be positioned at the 5' terminus of the reporter sequence and designed to generate a stem-loop structure in the reporter sequence, such that the RBS sequence of the reporter sequence is blocked. The cis-repressing stem-loop structure can be designed to block the RBS sequence based on the lowest energy conformation of the reporter transcript, as calculated by MFold and visualized with VARNA. The predicted inter-molecular interactions between the target transcript and the cis-repressing sequence of the reporter transcript can be calculated by RactIP and visualized by VARNA. A diagram can be drawn to visualize the base pairing between the target transcript and the cis-repressing sequence of the reporter transcript, as shown in Figure 28 below.

**[00244]** The interaction can include base pairing between 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 or more nucleotides in the target sequence and cis-repressing sequence. The complementary binding between the two sequences can be fully complementary, substantially complementary or partially complementary. The base pairing can be across contiguous nucleotide sequences or regions within the target and cis-repressing sequences, for example, as shown in Figure 28.

## 5) Cleavage Mechanisms for Cis-repressed Transcripts or Reporter Transcripts

**[00245]** The general mechanisms employed in the invention are intermolecular nucleic acid molecule interactions that may result in two subsequent mechanisms: (1) a conformational change in the secondary structure of the nucleic acid molecules, and (2) a cleaving event. Described herein are methods and systems for designing reporter transcripts that employ a cleaving event.

**[00246]** In some embodiments, a cleaving mechanism can be employed in the system and methods of the invention for cis-repression or for trans-activation. For example, as described above in Figures 17, 19 and 20, a system can be designed to take advantage of a cleaving mechanism by exposing a nucleic acid sequence of the reporter transcript to a cleaving enzyme (RNase) or sequestering a single-stranded sequence that is recognized by a sequence specific RNAase.

**[00247]** In one example, an ribonuclease E (RNase E) site can be designed in the reporter transcript (“\*” indicates the cleaving site):

(G,A)N(C,A)N(G)(G,U,A)\*(A,U)(C,U)N(C,A)(C,A). *See Kaberdin et al., Probing the substrate specificity of E. coli RNase E using a novel oligonucleotide-based assay. Nucleic Acids Research, 2003, Vol. 31, No. 16 (doi: 10.1093/nar/gkg690).*

**[00248]** In a cis-repression system, a cis-repressing sequence can be incorporated in the design of a reporter transcript, such that when transcribed, the conformation of the reporter transcript exposes a single stranded region containing a sequence RNase E recognition motif at the desired site to be cleaved. In some embodiments, the cleavage site can be involved in repression of the transcription of the reporter transcript, for example, if the cleavage site is within the coding region of the reporter gene.

**[00249]** For a trans-derepression system, the cis-repressed transcript can be engineered to bind to a target transcript, such that the interaction causes a conformational change in the reporter transcript that sequesters the single-stranded region containing the RNase E site.

**[00250]** The system can be designed such that the cis-repressing mechanism is due to a specific secondary structure generated by a conformation of the cis-repressing sequence, such as the transcription termination structure described above. In this example, a cleaving event serves to de-repress the reporter sequence. This can be accomplished by designing the cis-repressing sequence to interact with (bind to) a naturally-occurring plasmid or other cellular transcript, such that the interaction results in the generation of a single-stranded region containing the RNase E site that can be cleaved and thus removes the cis-repressing sequence from the reporter transcript.

**[00251]** In some embodiments, when a cleavage event is employed for expression of the reporter, the RNase E site is designed to be outside of the coding region of a reporter sequence with enough sequence length in the 5' and 3' UTR in order to allow for a viable reporter transcript. In this case, the RNase E site is designed to be at least 0, 1, 2, 3, 4, 5, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, or more base pairs upstream of the start codon in prokaryotic systems and at least 18, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200,

300, 400, 500, or more base pairs upstream of the start codon in eukaryotic systems or at least 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, or more base pairs downstream of the stop codon. In other embodiments, when a cleavage event is employed for repression of the reporter, the RNase E site is designed to be within the coding region of the reporter sequence or otherwise placed in order to inhibit expression of the reporter.

### 6) Transcripts

**[00252]** As described above, a transcript is a length of nucleotide sequence (DNA or RNA) transcribed from a DNA or RNA template sequence or gene. The transcript can be a cDNA sequence transcribed from an RNA template or an mRNA sequence transcribed from a DNA template. The transcript can be transcribed from an engineered nucleic acid construct. The transcript can have regions of complementarity within itself, such that the transcript includes two regions that can form an intra-molecular duplex. One region can be referred to as a “cis-repressing sequence” that binds to and blocks translation of a reporter sequence. A second region of the transcript is called a “reporter sequence” that encodes a reporter molecule, such as a detectable or selectable marker.

**[00253]** The transcripts of the invention can be a transcript sequence that can be 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotides in length. In other embodiments, the transcript can be at least 25, 30, 40, 50, 60, 70, 80, 90, 100, 500, 1000, 1500, 2000, 3000, 4000, 5000 or more nucleotides in length. The cis-repressing sequence and the reporter sequence can be the same length or of different lengths.

**[00254]** In some embodiments, the cis-repressing sequence is separated from the reporter sequence by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, or more spacer nucleotides.

### 7) Vectors

**[00255]** In another aspect, the transcripts (including antisense and sense sequences) of the invention are expressed from transcription units inserted into DNA or RNA vectors (see, e.g., Couture, A, *et al.*, *TIG*. (1996), 12:5-10; Skillern, A., *et al.*, International PCT Publication No. WO 00/22113, Conrad, International PCT Publication No. WO 00/22114, and Conrad, U.S. Pat. No. 6,054,299). These sequences can be introduced as a linear construct, a circular plasmid, or a viral vector, including bacteriophage-based vectors, which can be incorporated and inherited as a transgene integrated into the host genome. The transcript can also be constructed to permit it to be inherited as an extrachromosomal plasmid (Gassmann, *et al.*, *Proc. Natl. Acad. Sci. USA* (1995) 92:1292).

**[00256]** The transcript sequences can be transcribed by a promoter located on the expression plasmid. In one embodiment, the cis-repressing and reporter sequences are expressed as an inverted repeat joined by a linker polynucleotide sequence such that the transcript has a stem and loop structure.

**[00257]** Recombinant expression vectors can be used to express the transcripts of the invention. Recombinant expression vectors are generally DNA plasmids or viral vectors. Viral vectors expressing the transcripts can be constructed based on, but not limited to, adeno-associated virus (for a review, see Muzyczka, *et al.*, *Curr. Topics Micro. Immunol.* (1992) 158:97-129)); adenovirus (see, for example, Berkner, *et al.*, *BioTechniques* (1998) 6:616), Rosenfeld *et al.* (1991, *Science* 252:431-434), and Rosenfeld *et al.* (1992, *Cell* 68:143-155)); or alphavirus as well as others known in the art. Retroviruses have been used to introduce a variety of genes into many different cell types, including epithelial cells, *in vitro* and/or *in vivo* (see, *e.g.*, Eglitis, *et al.*, *Science* (1985) 230:1395-1398; Danos and Mulligan, *Proc. Natl. Acad. Sci. USA* (1998) 85:6460-6464; Wilson *et al.*, 1988, *Proc. Natl. Acad. Sci. USA* 85:3014-3018; Armentano *et al.*, 1990, *Proc. Natl. Acad. Sci. USA* 87:61416145; Huber *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88:8039-8043; Ferry *et al.*, 1991, *Proc. Natl. Acad. Sci. USA* 88:8377-8381; Chowdhury *et al.*, 1991, *Science* 254:1802-1805; van Beusechem. *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:7640-19 ; Kay *et al.*, 1992, *Human Gene Therapy* 3:641-647; Dai *et al.*, 1992, *Proc. Natl. Acad. Sci. USA* 89:10892-10895; Hwu *et al.*, 1993, *J. Immunol.* 150:4104-4115; U.S. Patent No. 4,868,116; U.S. Patent No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573). Recombinant retroviral vectors capable of transducing and expressing genes inserted into the genome of a cell can be produced by transfecting the recombinant retroviral genome into suitable packaging cell lines such as PA317 and Psi-CRIP (Comette *et al.*, 1991, *Human Gene Therapy* 2:5-10; Cone *et al.*, 1984, *Proc. Natl. Acad. Sci. USA* 81:6349). Recombinant adenoviral vectors can be used to infect a wide variety of cells and tissues in susceptible hosts (*e.g.*, rat, hamster, dog, and chimpanzee) (Hsu *et al.*, 1992, *J. Infectious Disease*, 166:769), and also have the advantage of not requiring mitotically active cells for infection.

**[00258]** Any viral vector capable of accepting the coding sequences for the transcript(s) to be expressed can be used, for example, vectors derived from adenovirus (AV); adeno-associated virus (AAV); retroviruses (*e.g.*, lentiviruses (LV), Rhabdoviruses, murine leukemia virus); herpes virus, and the like. The tropism of viral vectors can be modified by

pseudotyping the vectors with envelope proteins or other surface antigens from other viruses, or by substituting different viral capsid proteins, as appropriate.

**[00259]** For example, lentiviral vectors featured in the invention can be pseudotyped with surface proteins from vesicular stomatitis virus (VSV), rabies, Ebola, Mokola, and the like. AAV vectors featured in the invention can be made to target different cells by engineering the vectors to express different capsid protein serotypes. Techniques for constructing AAV vectors which express different capsid protein serotypes are within the skill in the art; see, *e.g.*, Rabinowitz J E *et al.* (2002), *J Virol* 76:791-801, the entire disclosure of which is herein incorporated by reference.

**[00260]** Selection of recombinant viral vectors suitable for use in the invention, methods for inserting nucleic acid sequences for expressing the transcripts into the vector, and methods of delivering the viral vector to the cells of interest are within the skill in the art. See, for example, Dornburg R (1995), *Gene Therap.* 2: 301-310; Eglitis M A (1988), *Biotechniques* 6: 608-614; Miller A D (1990), *Hum Gene Therap.* 1: 5-14; Anderson W F (1998), *Nature* 392: 25-30; and Rubinson D A *et al.*, *Nat. Genet.* 33: 401-406, the entire disclosures of which are herein incorporated by reference.

**[00261]** Viral vectors can be derived from AV and AAV. A suitable AV vector for expressing the transcripts featured in the invention, a method for constructing the recombinant AV vector, and a method for delivering the vector into target cells, are described in Xia H *et al.* (2002), *Nat. Biotech.* 20: 1006-1010. Suitable AAV vectors for expressing the transcripts featured in the invention, methods for constructing the recombinant AV vector, and methods for delivering the vectors into target cells are described in Samulski R *et al.* (1987), *J. Virol.* 61: 3096-3101; Fisher K J *et al.* (1996), *J. Virol.* 70: 520-532; Samulski R *et al.* (1989), *J. Virol.* 63: 3822-3826; U.S. Pat. No. 5,252,479; U.S. Pat. No. 5,139,941; International Patent Application No. WO 94/13788; and International Patent Application No. WO 93/24641, the entire disclosures of which are herein incorporated by reference.

**[00262]** The promoter driving transcript expression in either a DNA plasmid or viral vector featured in the invention may be a eukaryotic RNA polymerase I (*e.g.*, ribosomal RNA promoter), RNA polymerase II (*e.g.*, CMV early promoter or actin promoter or U1 snRNA promoter) or generally RNA polymerase III promoter (*e.g.*, U6 snRNA or 7SK RNA promoter) or a prokaryotic promoter, for example the T7 promoter, provided the expression plasmid also encodes T7 RNA polymerase required for transcription from a T7 promoter. The promoter can also direct transgene expression to the pancreas (*see, e.g.*, the insulin

regulatory sequence for pancreas (Bucchini *et al.*, 1986, Proc. Natl. Acad. Sci. USA 83:2511-2515)).

**[00263]** In addition, expression of the transcript can be precisely regulated, for example, by using an inducible regulatory sequence and expression systems such as a regulatory sequence that is sensitive to certain physiological regulators, *e.g.*, circulating glucose levels, or hormones (Docherty *et al.*, 1994, FASEB J. 8:20-24). Such inducible expression systems, suitable for the control of transgene expression in cells or in mammals include regulation by ecdysone, by estrogen, progesterone, tetracycline, chemical inducers of dimerization, and isopropyl-beta-D-1-thiogalactopyranoside (IPTG). A person skilled in the art would be able to choose the appropriate regulatory/promoter sequence based on the intended use of the dsRNA transgene.

**[00264]** Generally, recombinant vectors capable of expressing transcript molecules are delivered as described below, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of transcript molecules. Such vectors can be repeatedly administered as necessary. Once expressed, the transcript binds to target RNA and modulates its function or expression. Delivery of transcript expressing vectors can be systemic, such as by intravenous or intramuscular administration, by administration to target cells ex-planted from the patient followed by reintroduction into the patient, or by any other means that allows for introduction into a desired target cell.

**[00265]** Transcript expression DNA plasmids are typically transfected into target cells as a complex with cationic lipid carriers (*e.g.*, Oligofectamine) or non-cationic lipid-based carriers (*e.g.*, Transit-TKO<sup>TM</sup>). Multiple lipid transfections for dsRNA-mediated knockdowns targeting different regions of a single PROC gene or multiple PROC genes over a period of a week or more are also contemplated by the invention. Successful introduction of vectors into host cells can be monitored using various known methods. For example, transient transfection can be signaled with a reporter, such as a fluorescent marker, such as Green Fluorescent Protein (GFP). Stable transfection of cells *ex vivo* can be ensured using markers that provide the transfected cell with resistance to specific environmental factors (*e.g.*, antibiotics and drugs), such as hygromycin B resistance.

**[00266]** The delivery of the vector containing the recombinant DNA can be performed by abiotic or biologic systems. Including but not limited to liposomes, virus-like particles, transduction particles derived from phage or viruses, and conjugation.

### 8) Reporters for Transcript Assay

**[00267]** In some embodiments, the nucleic acid construct comprises a reporter sequence (e.g., a reporter gene sequence). The reporter gene encodes a reporter molecule that produces a signal when expressed in a cell. In some embodiments, the reporter molecule can be a detectable or selectable marker. In certain embodiments, the reporter molecule can be a fluorescent reporter molecule, such as a green fluorescent protein (GFP), yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), blue fluorescent protein (BFP), or red fluorescent protein (RFP). In other embodiments, the reporter molecule can be a chemiluminescent protein.

**[00268]** Reporter molecules can be a bacterial luciferase, an eukaryotic luciferase, a fluorescent protein, an enzyme suitable for colorimetric detection, a protein suitable for immunodetection, a peptide suitable for immunodetection or a nucleic acid that function as an aptamer or that exhibits enzymatic activity.

**[00269]** Selectable markers can also be used as a reporter. The selectable marker can be an antibiotic resistance gene, for example.

### 9) Cells and Target Genes for Transcript Reporter Assay

**[00270]** Examples of cells that can be used for detection include Gram-positive and Gram-negative bacteria, such as *S. aureus*, *E. coli*, *K. pneumoniae*, etc., fungi such as *Streptomyces coelicolor*, and other eukaryotic cells, including cells from humans, other mammals, insects, invertebrates, or plants.

**[00271]** Target transcripts can include any endogenous transcript, whether coding or non-coding. Target transcripts can be derived from eukaryotic and prokaryotic cells, including, for example, *mecA* transcript in *S. aureus* cells (indicative of MRSA), the *tcdB* transcript in *C. difficile* (indicative of toxigenic C. diff), and HPV E6/E7 transcripts in cervical epithelial cells (indicative of cervical cancer). Genes associated with infectious agents, such as viruses, can be targets as well, including HIV, HPV, etc. Other examples of target genes include non-coding RNA such as transfer RNA (tRNA) and ribosomal RNA (rRNA), as well as RNAs such as snoRNAs, microRNAs, siRNAs, snRNAs, exRNAs, and piRNAs and ncRNAs.

## EXAMPLES

**[00272]** Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

**[00273]** The practice of the present invention will employ, unless otherwise indicated, conventional methods of protein chemistry, biochemistry, recombinant DNA techniques and pharmacology, within the skill of the art. Such techniques are explained fully in the literature. See, e.g., T.E. Creighton, *Proteins: Structures and Molecular Properties* (W.H. Freeman and Company, 1993); A.L. Lehninger, *Biochemistry* (Worth Publishers, Inc., current addition); Sambrook, et al., *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *Methods In Enzymology* (S. Colowick and N. Kaplan eds., Academic Press, Inc.); *Remington's Pharmaceutical Sciences*, 18th Edition (Easton, Pennsylvania: Mack Publishing Company, 1990); Carey and Sundberg *Advanced Organic Chemistry 3<sup>rd</sup> Ed.* (Plenum Press) Vols A and B(1992).

**Example 1: Silent Mutation/Complementation Packaging System**

**[00274]** The following is an example of the design and construction of a silent mutation/complementation-based packaging system for producing non-replicative transduction particles.

**[00275]** The materials used for developing the packaging system are listed below:

**[00276]** Bacterial Strains:

- N1706, an E. coli K-12 P1 c1-100 Tn9 lysogen

**[00277]** Vectors:

- Y14439 (pBHR1 backbone)

**[00278]** The following GenBank accession numbers (N.B., the sequences referred to by accession number are those listed in the database as of the priority date of this application) or SEQ ID NOs. can be used for the vector backbone and cassette sequences:

- X06758 (bacterial luciferase genes luxAB)
- SEQ ID NO:1 (Native P1 pac-site)
- SEQ ID NO:3 (P1 lytic replicon containing the C1 repressor-controlled P53 promoter, the promoter P53 antisense, the repL genes, and an in-frame deletion of the kilA gene)
- SEQ ID NO:4 (Pblast promoter driving luxAB expression)

**[00279]** Construction of N1706(pac): pacA mutated strain: An exemplary sequence of a pacA mutated sequence is shown in SEQ ID NO: 2, shown in the informal sequence listing below. The mutation can be accomplished by constructing the mutated sequence via gene synthesis and then replacing the native sequence in N1706 with the mutated sequence via an allelic exchange approach.

**[00280]** Construction of the GWP10001 reporter vector: The GWP10001 vector contains the pBHR1 origin of replication exhibiting broad Gram-negative activity, two selectable markers for kanamycin and chloramphenicol, the native bacteriophage P1 pac-site sequence, the luxA and luxB genes are from *Vibrio harveyi* operatively linked to the constitutive blasticillin promoter (Pblast), and the P1 lytic replicon containing the C1 repressor-controlled P53 promoter, the promoter P53 antisense, the repL genes, and an in-frame deletion of the kilA gene.

**[00281]** Figure 2 shows the resulting vector (GWP10001, SEQ ID NO:11), which can be constructed in a variety of manners that are known to one of skill in the art including obtaining the cassettes via PCR from their native sources or via gene synthesis and assembly of the vector via traditional restriction enzyme-based cloning or alternative techniques such as Gibson assembly.

**[00282]** Silent/Complementation Packaging System: The packaging system includes the pacA mutant strain N1706(pac) complemented with the vector pGWP10001. As known to one of skill in the art, the manner of constructing this system can be accomplished by transformation N1706(pac) with vector pGWP10001. The vector pGWP10001 can be maintained in cultures of the transformed N1706(pac) by growing the transformant in the presence of 50 ug/mL of kanamycin.

**[00283]** Production of Transduction Particles Carrying Plasmid DNA: Non-replicative transduction particles carrying vector pGWP10001 can be produced from N1706(pac) transformants via thermal induction at 42°C. Incubation at 42°C results in induction of the P1 lytic cycle in which the prophage excises from the N1706 genome, produces phage structural elements, and packages pGWP10001 concatameric DNA formed by the lytic replicon in progeny phage particles, as depicted in Figure 1. The resulting cell lysate is then collected and contains non-replicative transduction particles, each consisting of bacteriophage P1 particles carrying a linear concatamer of pGWP10001 DNA.

#### **Example 2: Deletion/Complementation Packaging System**

**[00284]** The following is an example of the design and construction of a deletion/complementation-based packaging system for producing non-replicative transduction particles.

**[00285]** The materials used for developing the packaging system are listed below:

**[00286]** Bacterial Strains:

**[00287]** RN4220 is a restriction defective *S. aureus* strain that is a non-lysogenic derivative of NCTC 8325 and is an efficient recipient for *E. coli* DNA. It was first described

in Kreiswirth, B.N. *et al.*, *The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage*. Nature, 1983. 305(5936): p. 709-712.

[00288] RN10616 is derived by lysogenizing RN4220 with bacteriophage  $\varphi$ 80 $\alpha$ . Ubeda, C. *et al.*, *Specificity of staphylococcal phage and SaPI DNA packaging as revealed by integrase and terminase mutations*. Molecular Microbiology, 2009. 72(1): p. 98-108.

[00289] ST24 is derived from deleting the small terminase gene terS from the lysogenized bacteriophage  $\varphi$ 80 $\alpha$  in RN10616. Ubeda, C. *et al.*, *Specificity of staphylococcal phage and SaPI DNA packaging as revealed by integrase and terminase mutations*. Molecular Microbiology, 2009. 72(1): p. 98-108.

[00290] Vectors:

[00291] Examples of plasmids that can be used as source plasmids for cassettes, in some embodiments of the invention are described in Charpentier, E., *et al.*, *Novel Cassette-Based Shuttle Vector System for Gram-Positive Bacteria*. Appl. Environ. Microbiol., 2004. 70(10): p. 6076-6085.

[00292] The following GenBank accession numbers can be used for cassette sequences:

- SEQ ID NO:5 (*S. aureus* pT181 plasmid origin or replication copy number variant pT181cop-623 repC)
- M21136 (tetA(M))
- SEQ ID NO:12 (P<sub>clpB</sub> promoter sequence)
- SEQ ID NO:9 ( $\varphi$ 11 small terminase (terS) gene sequence)
- L09137 (amp ColE1 ori)
- X06758 (luxAB)
- M62650 (Transcription Termination)

[00293] terS Deletion: The construction of the terS knockout strain ST24 can be accomplished via an allelic-exchange-based strategy resulting in an in-frame deletion removing most of the coding sequence of the  $\varphi$ 80 $\alpha$  small terminase gene. The details of this strategy are described in Ubeda, C. *et al.*, *Specificity of staphylococcal phage and SaPI DNA packaging as revealed by integrase and terminase mutations*. Molecular Microbiology, 2009. 72(1): p. 98-108.

[00294] An exemplary sequence of a terS knockout strain is shown in SEQ ID NO:13, (shown in the sequence listing below). SEQ ID NO:13 is a RN10616 genomic sequence loci showing the  $\varphi$ 80 $\alpha$  terS deletion and complementation.

**[00295] Vector Construction:** The GW80A0001 vector is an *E. coli/S. aureus* shuttle vector. The vector contains *S. aureus* (pT181cop-623 repC) and *E. coli* (ColE1ori) origins of replication, the selectable markers for ampicillin (amp) and tetracycline (tet(M)) resistance for selection in *E. coli* and *S. aureus*, respectively, the  $\varphi$ 11 small terminase (terS) gene sequence that includes its own promoter, the luxA and luxB genes are from *Vibrio harveyi* operatively linked to the constitutive *S. aureus*  $P_{clpB}$  promoter, and a transcription termination sequence (TT).

**[00296]** Figure 4 shows the resulting vector (pGW80A0001, SEQ ID NO:14), which can be constructed in a variety of manners that are known to one of skill in the art. In one example, the tet(M) cassette and luxAB genes can be obtained via PCR amplification from the publically available pCN36 and pCN58 vectors (Charpentier, E., *et al.*).  $P_{clpB}$  can be obtained from PCR amplification from *S. aureus* RN4220 and terS can be obtained via PCR amplification from RN10616. A vector backbone can be obtained by removing the ermC gene from the publically available vector pCN48 (Charpentier, E., *et al.*), and the various components of the final vector pGW80A0001 can be assembled onto this vector backbone via appropriately designed restriction enzyme-based cloning.

**[00297] Deletion/Complementation Packaging System:** The packaging system can include the terS knockout strain ST24 complemented with the vector pGW80A0001 to generate strain GW24. As known to one of skill in the art, the manner of constructing this system can be accomplished by transformation ST24 with vector pGW80A0001. The vector pGW80A0001 can be maintained in cultures of the transformed ST24 by growing the transformant in the presence of 5 ug/mL of tetracycline.

**[00298] Production of Transduction Particles Carrying Plasmid DNA:** Non-replicative transduction particles carrying vector pGW80A0001 can be produced from GW24 via a Mitomycin C-induction method that was first demonstrated in *E. coli* and is now a standard technique for obtaining prophages from lysogenized bacteria. Otsuji, N. *et al.*, *Induction of Phage Formation in the Lysogenic Escherichia coli K-12 by Mitomycin C*. *Nature*, 1959. 184(4692): p. 1079-1080. This prophage induction method results in induction of the  $\varphi$ 80 $\alpha$  lytic cycle in which the prophage excises from the GW24 genome, produces phage structural elements, and packages pGW80A0001 concatameric DNA in progeny phage particles, as depicted in Figure 2. The resulting cell lysate is then collected and contains non-replicative transduction particles, each consisting of bacteriophage  $\varphi$ 80 $\alpha$  particles carrying a linear concatamer of pGW80A0001 DNA.

**Example 3: SaPIbov2-Based Packaging System Lacking Integrase**

[00299] The following is an example of the design and construction of a SaPIbov2-based packaging system for producing non-replicative transduction particles.

[00300] The materials used for developing the packaging system are listed below:

[00301] The following materials can be used to develop a SaPIbov2-based packaging system lacking integrase.

[00302] Bacterial Strains:

[00303] RN451 is a *S. aureus* strain lysogenized with bacteriophage  $\phi$ 11.

[00304] JP2131 is RN451 that has been lysogenized with SaPIbov2. See Maiques, E. et al., Role of Staphylococcal Phage and SaPI Integrase in Intra- and Interspecies SaPI Transfer. *J. Bacteriol.*, 2007. 189(15): p. 5608-5616.

[00305] JP2488 is strain JP2131 in which the *int* gene has been deleted from SaPIbov2 (SaPIbov2 $\Delta$ int). Maiques, E. et al., Role of Staphylococcal Phage and SaPI Integrase in Intra- and Interspecies SaPI Transfer. *J. Bacteriol.*, 2007. 189(15): p. 5608-5616.

[00306] Bacteriophage:

[00307] Bacteriophage  $\phi$ 11 can be obtained from *S. aureus* strain RN0451 via a Mitomycin C-induction method that was first described in *E. coli* and is now a standard technique for obtaining prophages from lysogenized bacteria. Otsuji, N. et al., *Induction of Phage Formation in the Lysogenic Escherichia coliK-12 by Mitomycin C*. *Nature*, 1959. 184(4692): p. 1079-1080.

[00308] Promoters:

[00309]  $P_{clpB}$  can be used as a promoter in this example. The *clpB* gene promoter is a constitutive promoter used for controlling the expression of the *int* gene. The *S. aureus* *clpB* ( $P_{clpB}$ ) gene promoter sequence was first described in 2004. Frees, D., et al., Clp ATPases are required for stress tolerance, intracellular replication and biofilm formation in *Staphylococcus aureus*. *Molecular Microbiology*, 2004. 54(5): p. 1445-1462. It was also first employed for controlling the gene expression in a plasmid in 2004. Arnaud, M., A. Chastanet, and M. Debarbouille, *New Vector for Efficient Allelic Replacement in Naturally Nontransformable, Low-GC-Content, Gram-Positive Bacteria*. *Appl. Environ. Microbiol.*, 2004. 70(11): p. 6887-6891. The promoter can be obtained from *S. aureus* RN4220 using primers described in 2004. *Id.*

[00310] Production of  $\phi$ 11/ SaPIbov2 $\Delta$ int co-lysogen (RN451( $\phi$ 11 SaPIbov2 $\Delta$ int)): The strain JP2488( $\phi$ 11 SaPIbov2 $\Delta$ int) can be produced by lysogenizing JP2488 with  $\phi$ 11.

**[00311]** Deletion of  $\varphi 11$  terS (RN451( $\varphi 11\Delta$ terS SaPIbov2 $\Delta$ int)): The strain RN451( $\varphi 11\Delta$ terS SaPIbov2 $\Delta$ int) can be produced by deleting the  $\varphi 11$  terS gene from RN451( $\varphi 11$ SaPIbov2 $\Delta$ int), as described in Tormo, M.A. *et al.*, *Staphylococcus aureus Pathogenicity Island DNA Is Packaged in Particles Composed of Phage Proteins*. J. Bacteriol., 2008. 190(7): p. 2434-2440.

**[00312]** Incorporation of  $P_{clpB}$ -int into *S. aureus* genome (RN451( $\varphi 11\Delta$ terS SaPIbov2 $\Delta$ int  $P_{clpB}$ -int)): RN451( $\varphi 11\Delta$ terS SaPIbov2 $\Delta$ int  $P_{clpB}$ -int) can be produced by first fusing  $P_{clpB}$  and *int* via standard molecular biology techniques then inserting the  $P_{clpB}$ -int fusion into the genome of RN451( $\varphi 11\Delta$ terS SaPIbov2 $\Delta$ int) and then selecting clones that have  $P_{clpB}$ -int inserted outside of the  $\varphi 11$  and SaPIbov2 regions.

**[00313]** Production of  $\varphi 11$  particles carrying only SaPIbov2 $\Delta$ int  $P_{clpB}$ -int concatamers:  $\varphi 11$  particles carrying only SaPIbov2 $\Delta$ int  $P_{clpB}$ -int concatamers can be produced via mitomycin-C induction of RN451( $\varphi 11\Delta$ terS SaPIbov2 $\Delta$ int  $P_{clpB}$ -int), as described by Otsuji, N. *et al.*, *Induction of Phage Formation in the Lysogenic Escherichia coliK-12 by Mitomycin C*. Nature, 1959. 184(4692): p. 1079-1080. The cell lysate contains non-replicative transduction particles, each consisting of bacteriophage  $\varphi 11$  structural proteins carrying a linear concatamer of GI-derived DNA.

**[00314]** One of skill in the art will understand how to construct the NRTPs of the invention using the above-referenced materials and well-known molecular biology and genetic techniques in the art.

#### **Example 4: terS Deletion/Complementation-based SarS Reporter Transduction Particles**

**[00315]** The following is an example of an inducer reporter-based SarS reporter system that employs a terS deletion/complementation-based non-replicative transduction particle.

**[00316]** *Reporter gene:* Bacterial luciferase (luxAB). The luxA and luxB genes are from *Vibrio harveyi*. They lack a transcriptional promoter and each contains their own ribosomal binding site.

**[00317]** Spa gene promoter ( $P_{spa}$ ): The spa gene promoter will be used for controlling the expression of the luxAB genes.

**[00318]** *Construction of  $P_{spa}$ -luxAB fusion:* The luxAB genes can be fused to the  $P_{spa}$  promoter sequence such that the luxAB genes are operatively linked to the  $P_{spa}$  promoter.

**[00319]** *Construction of the luxAB-expressing reporter vector*

[00320] The luxAB-expressing reporter vector can be constructed via standard molecular biological techniques by incorporating the  $P_{spa}$ -luxAB fusion product into the MCS of the shuttle vector depicted below.



[00321] *E. coli/S. aureus* shuttle vector that carries a *S. aureus* (pT181cop-623 repC) and *E. coli* (ColE1ori) origins of replication, genes for ampicillin (amp) and tetracycline (tet(M)) resistance, the  $\phi$ 11 small terminase (terS) gene under the control of a constitutive promoter ( $P_{clpB}$ ), a multiple cloning site (MCS), and a transcription termination sequence (TT).

[00322] GenBank accession numbers for cassette sequences:

[00323] J01764 (pT181 replicons)

[00324] M21136 (tetA(M))

[00325] Accession number not yet available ( $P_{clpB}$ )

[00326] AF424781 REGION: 16526..16966 (terS)

[00327] L09137 (amp ColE1 ori)

[00328] M62650 (TT)

[00329] Propagation of the vector for conducting *in vitro* manipulations and for verification of manipulations can be accomplished via the *E. coli* Top 10 and the final modified vector can then be introduced into *S. aureus* RN0451 $\Delta$ terS. Transduction particles carrying shuttle vector can be produced from the RN0451 $\Delta$ terS transformants via a Mitomycin C-induction method that was first described in *E. coli* 1959 and is now a standard technique for obtaining prophages from lysogenized bacteria. Otsuji, N., et al., Induction of Phage Formation in the Lysogenic Escherichia coliK-12 by Mitomycin C. *Nature*, 1959, 184(4692): p. 1079-1080. The cell lysate is then collected and contains non-replicative transduction particles each consisting of bacteriophage  $\phi$ 11 structural proteins carrying a linear concatamer of plasmid DNA capable of reporting on the presence of SarS in *target S. aureus* cells.

**Example 5: terS Deletion/Complementation-based  $\beta$ -lactamase Reporter Transduction Particles**

[00330] The following is an example of an intracellular enzyme reporter-based  $\beta$ -lactamase reporter system that employs a terS deletion/complementation-based non-replicative transduction particle.

[00331] *Reporter gene:* Renilla luciferase (ruc)

[00332] *Promoter:* The promoter can be  $P_{blaZ}$ . The constitutive beta-lactamase promoter can be used for driving the expression of the ruc gene.

**[00333]** *Caged substrate:* Caged coelenterazine-phosphate as described in Daniel Sobek, J.R., *Enzyme detection system with caged substrates*, 2007, Zymera, Inc.

**[00334]** *Construction of  $P_{blaZ}$ -ruc fusion:* The *ruc* genes can be fused to the  $P_{blaZ}$  promoter sequence such that the *ruc* genes are operatively linked to the  $P_{blaZ}$  promoter.

**[00335]** *Construction of the ruc-expressing reporter vector:* The ruc-expressing reporter vector can be constructed via standard molecular biological techniques by incorporating the  $P_{blaZ}$ -ruc fusion product into the MCS of the shuttle vector depicted in Section V, A, 3), i) above.

**[00336]** Propagation of the vector for conducting in vitro manipulations and for verification of manipulations can be accomplished via the *E. coli* Top 10 and the final modified vector can then be introduced into *S. aureus* RN0451 $\Delta$ terS. Transduction particles carrying shuttle vector can be produced from the RN0451 $\Delta$ terS transformants via a Mitomycin C-induction method that was first described in *E. coli* 1959 and is now a standard technique for obtaining prophages from lysogenized bacteria. Otsuji, N., et al., Induction of Phage Formation in the Lysogenic Escherichia coliK-12 by Mitomycin C. *Nature*, 1959.

184(4692): p. 1079-1080. The cell lysate is then collected and contains NRTPs each consisting of bacteriophage  $\phi$ 11 structural proteins carrying a linear concatamer of plasmid DNA capable of expressing Renilla luciferase within viable *S. aureus* cells within the  $\phi$ 11 host range.

**Example 6: terS Deletion/Complementation-based Intracellular Molecule Reporter Transduction Particles**

**[00337]** The following is an example of an intracellular molecule reporter-based\_reporter system that employs a terS deletion/complementation-based non-replicative transduction particle.

**[00338]** *Promoter:* The promoter can be  $P_{blaZ}$ . The constitutive beta-lactamase promoter can be used for driving the expression of the *ruc* gene.

**[00339]** *Switchable aptamer:* Switchable aptamers can be designed and constructed as described in Samie Jaffrey, J.P., *Coupled recognition/detection system for in vivo and in vitro use*, 2010, Cornell University.

**[00340]** *Fluorophore substrate:* Corresponding fluorophore substrates in conjunction with the above switchable aptamers can be designed and constructed as described in Samie Jaffrey, J.P., *Coupled recognition/detection system for in vivo and in vitro use*, 2010, Cornell University.

**[00341]** *Construction of P<sub>blaZ</sub>-SA fusion:* The SA gene can be fused to the P<sub>blaZ</sub> promoter sequence such that the SA gene is operatively linked to the P<sub>blaZ</sub> promoter.

**[00342]** *Construction of the SA-expressing reporter vector:* The SA-expressing reporter vector can be constructed via standard molecular biological techniques by incorporating the P<sub>blaZ</sub>-SA fusion product into the MCS of the shuttle vector depicted in Example 4 above. Propagation of the vector for conducting in vitro manipulations and for verification of manipulations can be accomplished via the *E. coli* Top 10 and the final modified vector can then be introduced into *S. aureus* RN0451ΔterS. Transduction particles carrying shuttle vector can be produced from the RN0451ΔterS transformants via a Mitomycin C-induction method that was first described in *E. coli* 1959 and is now a standard technique for obtaining prophages from lysogenized bacteria. Otsuji, N. *et al.*, *Induction of Phage Formation in the Lysogenic Escherichia coli K-12 by Mitomycin C*. *Nature*, 1959. 184(4692): p. 1079-1080. The cell lysate is then collected and contains non-replicative transduction particles each consisting of bacteriophage φ11 structural proteins carrying a linear concatamer of plasmid DNA capable of expressing the SA within viable *S. aureus* cells within the φ11 host range.

#### **Example 7: Non-Replicative Transduction Particle-Based Reporter System**

**[00343]** The non-replicative transduction particles described above can be used in a reporter system for detecting the presence of viable bacteria via the expression of a reporter molecule (e.g. luxAB). When this transduction particle introduces a reporter vector (e.g. pGW80A0001) into a cell within the host range of the transduction particle, cells in which the promoter (e.g. P<sub>clpB</sub>) is recognized by the cells transcription machinery are able to drive the expression of the reporter molecule within that cell.

**[00344]** To test the functionality of non-replicative transduction particles as reporters for detecting the presence of *S. aureus* cells, various MSSA/MRSA reporter assays were developed. In an embodiment, a non-replicative transduction particle was developed from a *S. aureus*-specific bacteriophage, and the bacterial luciferase genes luxAB under the control of a constitutive promoter were incorporated. When the non-replicative transduction particle delivered the reporter nucleic acid into *S. aureus*, the constitutive promoter expressed luxAB suitable for reporting on the presence of a viable *S. aureus*.

**[00345]** In addition, the antibiotic cefoxitin was added prior to, simultaneously with, or after the addition of the transduction particles to a sample containing *S. aureus* cells. If the cells were not phenotypically resistant to cefoxitin (i.e., were not MRSA), luminescence was decreased or eliminated, indicating that the cells were MSSA. If, however, the cells were

phenotypically resistant to cefoxitin (*i.e.*, were MRSA), increased or detectable luminescence was observed, indicating that the cells were MRSA.

**Non-Replicative Transduction Particle-Based Viable Cell Reporter Assay Function**

**[00346]** The function of the non-replicative transduction particle as a reporter was assayed. The transduction host range of the bacteriophage  $\varphi 80\alpha$ -based non-replicative transduction particle was examined in 101 clinical MRSA isolates. The transduction assay was conducted by exposing cultures of each bacterial isolate grown in modified TSB to GW24 cell lysate containing the non-replicative transduction particles and culturing the mixture on solid media containing tetracycline.

**[00347]** In this example, the non-replicative transduction particle carried a tetracycline selectable marker. Cells transduced with the non-replicative transduction particles were expected to be resistant to tetracycline. In addition, transduction was examined via luminescence assay by exposing each bacterial isolate in liquid culture to cell lysate containing the non-replicative transduction particles and evaluating the mixture for bacterial luciferase luminescence activity after an incubation period.

**[00348]** The transduction assay showed that the  $\varphi 80\alpha$ -based non-replicative transduction particle was able to transduce all of the 101 clinical isolates of MRSA and none of the non-*S. aureus* Staphylococci.

**[00349]** Figure 21 shows the results of the transduction assay in which 36 tetracycline-sensitive MRSA were exposed to transduction particles carrying pGW80A0001 and then were spotted onto media plates containing 5 ug/mL of tetracycline. The results show that all 36 MRSA strains grew on the media containing tetracycline due to transduction with pGW80A0001. Control experiments in which MRSA isolates were spotted onto tetracycline containing media without exposure to transduction particles showed no growth (not shown). Furthermore, plasmid isolation from transduced MRSA strains demonstrated recovery of the pGW80A0001 plasmid as confirmed via sequencing of the isolated plasmid. The transduction results thus demonstrated that the origin of replication of the reporter plasmid exhibits activity on all of the MRSA isolates tested.

**[00350]** Figure 22 illustrates the luminescence measured from 80 clinical isolates of MRSA and 28 clinical isolates of methicillin sensitive *S. aureus* (MSSA) transduced with the transduction particle. In the experiment, cultures of MRSA and MSSA were grown to an optical density at 600nm of 0.1 and then 100 uL of the cultures grown in Modified TSB were mixed with 10 uL of GW24 cell lysates containing transduction particles and further

incubated at 37°C for a period of 4 hours prior to assaying for luminescence. Luminescence measurements were conducted by adding 10 uL of a 1 mM solution of Decanal, an aldehyde that triggers a luminescent reaction within cells expressing bacterial luciferase. As expected, luminescence was observed from both MRSA and MSSA transduced with the *S. aureus*-specific non-replicative transduction particle. Furthermore, when cefoxitin was added to the cell cultures at the same time as the addition of transduction particles, luminescence was observed from MRSA but not from MSSA, thus demonstrating the ability for the transduction particles to report on both the presence of MSSA and of MRSA. The luminescence results thus demonstrate that the promoter driving luxAB expression exhibits activity on all of the *S. aureus* isolates tested.

**Optimization of Non-Replicative Transduction Particle-Based Viable Cell**

**Reporter MRSA Assay – Transduction Particle Reagent Formulation**

**[00351]** The production and formulation of the non-replicative transduction particle reagent was optimized to a final formulation. In summary, a 15 L scale fermentation was performed using TSB media including peroxide induction of GW24. The 15 L fermenter batch was inoculated from a 200 mL overnight seed culture (an inoculum ratio of 1.3% (v/v)). The culture was induced at an O.D. of 0.8 with hydrogen peroxide and cooled to 25°C post induction without pH or DO control. Culture supernatant was harvested by tangential flow filtration (TFF) the following morning for the purpose of clarifying the phage transduction particles from the cell debris. The material was then further concentrated and diafiltered into SM Buffer without gelatin and stored at 2-8°C prior to final sterile filtration and storage.

**[00352]** A detailed summary of the process is outlined below:

**Seed flask growth**

- (1) Inoculate 200 mL of TSB containing 5 ug/mL tetracycline with GW24
- (2) Incubate at 37°C, 200 RPM for 10-18 hours.

**Fermentation Innoculation (15L TSB with 5 ug/mL tetracycline)**

- (1) Prepare the fermenter skid with the following fermentation conditions: 37°C, agitation at 250 RPM, airflow at 15 LPM, and backpressure at 3 psig.
- (2) Inoculate the fermentor using the 200 mL overnight seed culture.

**Induce Culture**

- (1) Once the OD600nm reaches 0.8 (0.6-0.9), induce the culture with 0.5 mM H2O2
- (2) Increase temperature setpoint of fermenter to 42°C

**Post-Induction Conditions and Monitoring**

- (1) Once 30 minute induction is complete, reset temperature target for fermenter to 25 °C
- (2) One hour post cooling, turn off air feed to fermenter and set agitation to zero
- (3) Monitor the fermentation culture at hour intervals, or more frequently as necessary, until the OD600nm has decreased to or below 0.40.

**Harvest/Clarification**

- (1) After the fermentation culture OD600 has reached a minimum less than or equal to 0.40, take a 20mL aseptic sample and add 30µL of Benzonase to the fermentor.
- (2) Reset agitation to 250 rpm. Allow 60 minutes with agitation for Benzonase incubation.
- (3) Clarify the EOF sample with a 15 minute centrifugation at 3000g.
- (4) Pass the clarified material through a 0.45uM membrane filter

**Concentration and Buffer Exchange**

- (1) Concentrate the clarified culture by TFF using 500 kDa flat sheet membrane 10-fold.
- (2) Diafilter the concentrated culture at a constant volume against SM Buffer without gelatin using the 500kDa TFF membrane used for concentration

**Final Filtration**

- (1) Filter the concentrated buffer exchanged material through a 0.2µm filter.
- (2) Store the final filtered phage material at 2-8 °C.

[00353] Various other reagents and formulations can be used as known to those of skill in the art to derive the formulation.

### **Optimization of Non-Replicative Transduction Particle-Based Viable Cell**

#### **Reporter MRSA Assay – Growth Media Formulation**

[00354] A growth media formulation was optimized for the NRTP-based viable cell reporter MRSA assay. In order to produce luminescence in the NRTP-based MRSA Assay, the media needs to be balanced for *Staphylococcus aureus* growth and have adequate concentration of cations and additives to favor NRTP transduction. The TSBmod media used in assays prior to this development study was known to have precipitation issues that would affect the stability of the media. Growth media formulation required stability in the final formulation with a goal of 1 year at room temperature.

[00355] Methods/Procedures: Cells preparation for MRSA Assay

(1) Ten unique strains of MRSA for the Subset Assay and one unique strains of MSSA were tested in the MRSA Assay.
(2) Overnight cultures were started in a deep 96 well plate at a 1:50 dilution in TSB from a frozen one-time use stock and incubated at 37°C on an orbital shaker for >15 hours. MRSA/MSSA (8µl) in TSB (392µl)
(3) The next day, a day culture at a 1:50 dilution from the overnight culture was started in TSB in 96 well deep well plate (392µl TSB + 8ul cells) and incubated at 37°C on an orbital shaker for 4 hours.
(4) Cells were spun in a centrifuge for 5 minutes at 1800g force and 10°C, spent media was aspirated without disturbing the pellet.
(5) Spun cells were washed in 50mM Tris-HCl pH 7.2, centrifuged, buffer aspirated without disturbing pellet and re-suspended in 400µl RPMI. RPMI is used in order to reduce variability in the metabolic state of cells and to mimic low metabolism as found in clinical samples.
(6) Plate was covered with airpore seal and incubated on bench for 48hrs.
(7) OD was read by transferring 200µl of RPMI culture in a shallow well OD plate and blank well with RPMI media alone was used to subtract blank OD.
(8) Cells were normalized to OD 0.1 in 100µl
(9) Another dilution was made 1:10 in RPMI to yield an OD of 0.005

**[00356]** Assay base media was prepared to be tested as shown in Table 2 and a representative set of media modifications in preparation for MRSA assay are shown in Table 3.

**Table 2: Base Media for Growth Media Formulation Development**

Components	TSB	B2	BSS-2	Notes
Enzymatic Digest of Soybean Meal (g)	3	0	3	
Enzymatic Digest of Casein (g)	17	10	10	
Yeast Extract (g)	N/A	25	25	
Sodium Chloride (g)	5	25	25	
Dipotassium Phosphate (g)	2.5	1	0	
alpha-D Glucose (g)	2.5	5	5	
Volume (litre)	1	1	1	Adjust pH to 7.2 with 10N NaOH. Autoclave or filter sterilize

**Table 3: Base Media Modifications for Growth Media Formulation Development**

**Development**

Concentration of salt/ additives for modification							
Base media (30ml)	Mod number	CaCl <sub>2</sub> (mM)	MgCl <sub>2</sub> (mM)	BGP (mM)	Tris-HCl pH7.0 (mM)	EDTA (mM)	HEPES (mM)
B2	M53	5.0	2.0	0.0	50.0	10.0	0.0
BSS-2	M50	10.0	2.0	60.0	50.0	10.0	0.0
BSS-2	M54	6.7	3.3	60.0	50.0	0.0	0.0
BSS-2	M55	5.0	5.0	60.0	50.0	0.0	0.0
BSS-2	M56	6.7	3.3	60.0	0.0	0.0	10.0
BSS-2	M57	5.0	5.0	60.0	0.0	0.0	10.0
TSB	M1 (original)	5.0	10.0	60.0	0.0	0.0	0.0
TSB	M58	5.0	10.0	60.0	0.0	11.1	0.0

**[00357]** To each media preparation, NRTP and Cefoxitin was added according to Table 4 below to make the NRTP media reagent:

**Table 4: MRSA Assay Growth Media/ Transduction particle reagent combination**

30ml media	Final Concentration
<b>Cefoxitin</b>	5ug/ml
<b>GW24 Lysate</b>	30X

[00358] The MRSA Assay was run with the following steps:

(1) Assay Plate Setup: Add 198 $\mu$ l of Phage Media Reagent and 2.0 $\mu$ l of each dilution of bacteria 0.05 OD and 0.005 OD in RPMI (roughly equivalent to 20,000 and 2,000 CFU/mL, respectively) or 2.0 $\mu$ L of RPMI as a blank.
(2) Incubate Assay Plate: Incubate Assay plate on orbital shaker at ~100 rpm for 4 hours at 37°C.
(3) Prepare Luminometer (Molecular Devices SpectraMax L): Wash reagent line with 70% ethanol followed by DI water then prime with the substrate reagent. Set up software as Fast Kinetic with injection of 50 $\mu$ L of substrate reagent at 250 $\mu$ l/sec after 10 baseline points and read at 40 points every 0.25 seconds.
(4) Run Assay: Test each bacterial dilution plate, after letting plate equilibrate to room temperature for 5 minutes.

[00359] Analysis

(1) Determine cutoff by averaging blank RLU across all replicates and time points and adding three standard deviations.
(2) Determine maximum RLU for each sample using SoftMaxPro.
(3) Determine if the maximum RLU was greater than the cutoff RLU, and if so, then the sample data was used for comparisons of media performance.
(4) Normalize all max RLU values to the Max RLU in TSB M1 (media in use until development started) for the strain being analyzed at the specific dilution.
(5) Average the normalized RLU values across all MRSA strains for a particular media and its modification

(6) Average the averages for the two dilution plates, ultimately leading to a single numerical value representing the fold increase in performance based on RLU of a particular media across 10 different MRSA strains in 2 cell dilutions tested.

**Results of NRTP-Based Viable Cell Reporter MRSA Assay**

**[00360]** Determination of Cutoff RLU: The average and standard deviation of the RLU was calculated across all time points (25) for each blank replicate (4). The cutoff was calculated for each plate as the average blank RLU plus three standard deviations.

**[00361]** Determination of Relative Improvements: The maximum RLU was exported for each sample (blanks, MSSA and MRSA at all dilutions) from SoftMaxPro and compared to the cutoff RLU. If the sample had 2 data points greater than the cutoff for phage concentration, then the max RLU value was utilized for analysis.

**[00362]** The values were normalized by dividing a particular max RLU by the max RLU of its control condition (that strain in TSB M1-origianl media, at the dilution being analyzed). The ratios obtained were averaged across 10 MRSA for each media condition and each dilution, as shown in Table 5. The average across the two dilutions is also shown in the table.

***Table 5: MRSA Assay Results from Various Growth Media Formulations***

<b>Media</b>		<b>Plate 1</b>	<b>Plate 2</b>	<b>Average for both dilutions</b>
B2	M53	1.89	1.88	1.89
BSS-2	M50	1.37	1.47	1.42
BSS-2	M54	1.50	1.76	1.63
BSS-2	M55	1.82	2.90	2.36
BSS-2	M56	2.38	6.00	4.19
BSS-2	M57	2.00	3.92	2.96
TSB	M1	1.00	1.00	1.00
TSB	M58	1.18	0.96	1.07

**Conclusions**

**[00363]** BSS2-M56 exhibited the best performance on average across the various media tested. HEPES buffer based media performed better than Tris-HCl buffered media. HEPES

is known to be a biologically favorable buffering system as opposed to Tris-HCl. B2 based base/broth had better performance than TSB based broth.

**[00364]** Various other reagents and formulations can be used as known to those of skill in the art to derive the formulation. Other suitable formulations were developed via similar experiments as described above. Examples of other suitable formulations are included below in Tables 6, 7, and 8.

***Table 6: BSC Media Formulation***

<b>BSC Components</b>	<b>Amount</b>
Enzymatic Digest of Casein	14.5g
Yeast Extract	35.5g
Sodium Chloride	35.5g
alpha-D Glucose	7g
Total Vol	1 L

***Table 7: BSC Media Modification***

<b>BSC-M64</b>	
<b>Chemical Name</b>	<b>Final (Assay) Conc</b>
BGP (mM)	60.0
HEPES (mM)	10.0
LiCl(mM)	84.0
BSC	To 1 L

**Table 8: Transduction Particle Media Modification**

<b><u>Transduction Particle Formulation (PM4)</u></b>	
<b><u>Chemicals</u></b>	<b><u>Final (Assay ) conc</u></b>
CaCl <sub>2</sub> (M)	0.00667
MgCl <sub>2</sub> (M)	0.00335
HEPES (M)	0.01000
GW24 lysate stock	0.01250
Sodium Azide (%)	0.0006
Water	To 1 mL

**Optimization of Non-Replicative Transduction Particle-Based Viable Cell****Reporter MRSA Assay – Substrate Reagent Formulation**

**[00365]** In order to produce luminescence in the MRSA Assay, the Substrate Reagent must include an aldehyde as a substrate for luciferase. An initially developed aliphatic aldehyde formulation (4.2 mM Tridecanal in TSB) was not stable and formed a heterogeneous emulsion rather than a solution. This example outlines the development of a Substrate Reagent formulation that addresses these issues with a goal of 6 months at room temperature or 2-8°C stability.

**[00366]** This example describes the steps that were taken to develop the Substrate Reagent to a final formulation.

**Methods/Procedures**

**[00367]** All screening and stability experiments were tested using a “Model System” that consists of *S. aureus* strain RN4220 harboring a LuxAB-expressing plasmid. The typical preparation and testing method was as follows.

- (1) Overnight Culture: 2 mL TSB+1 uL of 10 mg/mL Tetracycline+1 colony of Model System Bacteria from TSA plate, shaking at 225 rpm overnight at 37°C
- (2) Day Culture: Diluted overnight culture 1:50 or 1:100 into TSB+5 ug/mL Tetracycline, shaking at 225 rpm for 1.5-2 hours at 37°C.
- (3) Normalize Day Culture: Measured 1 mL of day culture on Nanodrop with cuvette at 600 nm, blanking with TSB+5 ug/mL Tetracycline. Diluted to 0.1 OD with TSB+ 5ug/mL Tetracycline.
- (4) Dilute Culture for Testing: Diluted 0.1 OD Culture with TSB+ 5ug/mL Tetracycline to a 1:200, 1:2000 and 1:20000 dilution which was roughly equivalent to 100000, 10000 and 1000 CFU/mL.
- (5) Plate Bacteria: Added 200 uL of each dilution and a blank (TSB+5 ug/mL Tetracycline with no bacteria) in three replicates to a Greiner Bio-one white assay plate for each substrate to be tested.
- (6) Prepare Luminometer (SpectraMax L): Wash reagent line with 70% ethanol followed by DI water then prime with the substrate. Set up software as Fast Kinetic with injection of 50 uL substrate at 250 ul/sec after 10 baseline points and read at 40 points every 0.25 seconds.
- (7) Run Assay: Test each formulation of Substrate Reagents with washing and priming SpectraMax L between each substrate. Bring all Substrate Reagents to room temperature before testing.

**[00368]** All confirmation experiments were tested using the MRSA Assay in order to ensure similar results on the actual assay as the Model System used to screen new formulations.

- (1) Prepare Culture: Ten MRSA low performing strains and one MSSA strain were grown to log-phase in TSB in a 2 mL deep well block. Cells were spun down, washed with 1x PBS then resuspended in RPMI media.
- (2) Normalize Bacteria: Measure 200 uL of RPMI culture and RPMI blank in Greiner Bio-one clear plate on VersaMax at 600 nm. Subtract blank OD from each strain. Normalize each strain to 0.05 OD in RPMI media.
- (3) Dilute Bacteria: Dilute 0.05 OD culture 1:10 in RPMI media to 0.005 OD.
- (4) Prepare Phage Media Reagent: Add Phage, Cefoxitin and Sodium Pyruvate to BSS-M56 including:
  - a. Cefoxitin (5 ug/mL)
  - b. GW24 lysate stock (0.03X)
  - c. Sodium Pyruvate (0.025M)
- (5) Set up Assay Plate = Add 198 uL of Phage Media Reagent and 2 uL of each dilution of bacteria (0.05 OD and 0.005 OD in RPMI, roughly equivalent to 20000 or 2000 CFU/mL) or 2 uL of RPMI as a blank in two replicates.
- (6) Incubate Assay Plate = Incubate Assay plate on orbital shaker at ~100 rpm (speed 3) for 4 hours at 37°C.
- (7) Prepare Luminometer (SpectraMax L) = Wash reagent line with 70% ethanol followed by DI water then prime with the substrate. Set up software as Fast Kinetic with injection of 50 uL substrate at 250 ul/sec after 10 baseline points and read at 40 points every 0.25 seconds.
- (8) Run Assay = Test each formulation of Substrate Reagents with washing and priming SpectraMax L between each substrate.

[00369] Experiments for the development of Substrate Reagent formulation were designed to improve the following:

- (1) Improve solubility via adding surfactants (Tween 20, Triton X-100, NP-40, Brij-35, SNS, etc.), adding solvents (Ethanol, Methanol, DMSO, etc.), adding non-volatile oils (Castor Oil)
- (2) Improve stability via adding stabilizers (Triethanolamine, Cyclodextrin etc.), adding antioxidants (Vitamin E, Vitamin E Acetate, Vitamin E PEG 1000, Oxyrase, etc.), adjust method of Tridecanal addition (with surfactant, with solvent, into final solution, with antioxidant, etc.), storing Tridecanal and Substrate Reagent under nitrogen to reduce oxidation of aldehyde, and reducing possibility of microbial contamination by adding preservatives such as ProClin and by sterile filtration of the Substrate Reagent.
- (3) Improve Assay Performance via adjustment of the pH of the formulation and the pH buffer system
- (4) Improve overall performance via determining the aldehyde with highest RLU output (tested aldehydes from 6-14 carbons in multiple formulations to determine if an improvement in solubility, stability and assay performance was observed).
- (5) Improve overall performance via adding antifoam in order to reduce foaming during preparation of reagent and addition of reagent to sample during the assay.

### Analysis and Results

[00370] The kinetic reaction was plotted for each sample and a line fit to the average at each read point of three replicates. Typically results showed at 1:2000 dilution of 0.1 OD model system bacteria, roughly equivalent to 10,000 CFU/mL or 2,000 CFU/assay.

[00371] The normalized maximum RLU to that of the reference substrate reagent was analyzed for stability experiments. At each stability time point, maximum RLU for each sample was normalized to the reference substrate maximum RLU. Normalized Maximum RLU was plotted over time points and linear regression with 95% CI was plotted.

### Conclusions

[00372] The key parameters adjusted from the reference formulation for producing a final Substrate Reagent formulation are summarized in Table 9.

***Table 9: Summary of Reagent Formulation Development Results***

Modification to Substrate Reagent	Reason
4.2 mM Tridecanal+TSB	Original Substrate Reagent
Remove TSB	Reduce possibility of contamination
Add 1% Tween 20	Improve Solubility
Adjust to pH3 with 79.45% 0.1 M Citric Acid-19.55% 0.2 M Sodium Phosphate Dibasic Buffer	Improve Assay Performance
Add Tridecanal directly to concentrated surfactant	Improve Stability
Add Filtering of Substrate Reagent through 0.2 um PES membrane	Improve Stability
Add 0.05% ProClin 300	Improve Stability
Add Triethanolamine	Improve Stability
Change 1% Tween 20 to 0.5% Triton X-100	Improve Stability, Improve Solubility
Change from 79.45% 0.1 M Citric Acid-19.55% 0.2 M Sodium Phosphate Dibasic Buffer to 82% 0.1 M Citric Acid-18% 0.1 M Sodium Citrate Buffer, remain at pH3	Improve Assay Performance, reduce possibility of precipitation with removal of phosphate buffer
Add 100 ppm Antifoam Y30	Improve Assay Performance
Add 0.5% Vitamin E Acetate	Improve Stability, reduce precipitation
Change Primary Tridecanal Manufacturer from Alfa Aesar to Sigma/OmegaChem	Improve Assay Performance
Change 0.5% Vitamin E Acetate to 1-2% Vitamin E PEG 1000	Improve Assay Performance, Improve Solubility, Improve Stability

**[00373]** Two Substrate Reagent Formulations were prepared for two different storage temperatures, one for storage at 2-8°C and one at 18-24°C.

**[00374]** Final Substrate Reagent Formulations stored at 2-8°C. Formulation: 0.5% Triton X-100+ 4.2mM Tridecanal+ 0.5% Vitamin E Acetate+ 100 ppm Antifoam Y30+ 0.5%

Triethanolamine+ 82% 0.1 M Citric Acid + 18% 0.1 M Sodium Citrate @pH3+ 0.05% ProClin 300. The formulation did not precipitate after 1 month at 2-8°C and was able to detect MRSA strains the same as on Day 0.

[00375] Final Substrate Reagent Formulations stored at 18-24°C. Formulation: 0.5% Triton X-100+ 6.3mM Tridecanal+ 100 ppm Antifoam Y30+ 0.5% Triethanolamine+ 82% 0.1 M Citric Acid + 18% 0.1 M Sodium Citrate @pH3+ 2%  $\alpha$ -Tocopherol-PEG 1000 Succinate+ 0.05% ProClin 300. The formulation did not precipitate after 1 month at 18-24°C and was able to detect MRSA strains the same as on Day 0.

[00376] Various other reagents and formulations can be used as known to those of skill in the art to derive the formulation.

**Analytical Performance of Non-Replicative Transduction Particle-Based Viable Cell Reporter MRSA Assay**

[00377] The analytical performance of the optimized NRTP MRSA assay was examined, including an analysis of the assay's limit of detection and an analysis of the cross-reactivity and microbial interference of the assay when challenged with non-target organisms.

**A) Limit of Detection Assay**

[00378] The Limit of Detection of the NRTP assay was assessed via determining the lowest amount of MRSA cells representing various strains that could produce a relative light unit (RLU) signal above that of a threshold determined from blank samples. MRSA strains included the SCCmec Types I, II, and IV as well as a MRSA strain carrying the *mecA* gene variant *mecC* – a strain of MRSA that conventional FDA-cleared MRSA PCR assays have failed to detect.

[00379] The following key materials were used in the clinical performance study:

[00380] Growth Media Reagent: BSS-M56

[00381] Substrate Reagent: Final Substrate Reagent Formulations to be stored at 18-24°C as described above.

[00382] Transduction Particle Reagent: BSS-M56 base with 10ug/mL (*i.e.* 2X concentration) cefoxitin and transduction particle reagent as described above at 2X concentration.

[00383] LoD Study Protocol:

[00384] Overnight Culture: For each MRSA strain and a MSSA negative control strain, 2 mL of TSB were inoculated with a colony of the strain previously grown on TSA plates.

Overnight MRSA cultures included 5ug/mL cefoxitin. All samples were incubated overnight at 37°C in a shaking incubator.

[00385] Day Culture: 20 uL of each of the overnight cultures were transferred into a new culture tube containing 2 mL of Growth Media Reagent. The inoculums were then incubated at 37C with shaking for approximately 1hr 45min, until the OD(600nm) reached 0.1.

[00386] Serial dilutions:

- a) 1000uL of each of the samples were dispensed into row A of 2mL deep well 96-well plate.
- b) The remaining rows (B-H) were then filled with 900 uL of Growth Media Reagent.
- c) 10-fold serial dilutions were then prepared taking 100uL from row A and mixing in row B, etc., such that row H contained samples of row A material at  $10^{-7}$  dilution.

[00387] Enumeration of bacterial load: 5 uL of each well of row E was spotted onto a TSA plate which was then tilted to allow the spot of liquid to spread onto the plate (in order to later facilitate colony counting). (Row E is a  $10^{-4}$  dilution of row A). Plates were then incubated overnight at 37°C.

[00388] Assay preparation:

[00389] a) Wells of a white 96 well assay plate were filled with 100 uL of 2x Transduction Particle Reagent.

[00390] b) Rows F and G (*i.e.*,  $10^{-5}$  and  $10^{-6}$ -fold dilutions of row A, respectively) were then used to fill wells of the 96 well assay plate containing Transduction Particle Reagent such that each sample was added to the plate in four-replicates.

[00391] c) The plate was then sealed with a breathable seal and incubated for 4 hours at 37°C with moderate shaking, 50rpm.

[00392] At the end of 4 hours, the plate was remove from incubator and immediately measured for luminescence on a SpectraMax L that injected 50  $\mu$ l of the Substrate Reagent and measured luminescence for a period of 1 minute.

[00393] Analysis:

[00394] The luminescence data from each sample was plotted as RLU vs. time. Blank samples were used to determine a Cutoff calculated from all time points of the blank samples using the following formula: (Mean Blank RLU + 3\* SD Blank RLU)

[00395] The average peak RLU post-substrate injection was then obtained for each sample in order to determine the sample of highest dilution for which an RLU value was generated

that was above the blank samples Cutoff. The colony forming unit (CFU) counts at the highest dilution for which an RLU value was generated that was above the blank samples Cutoff was determined from the enumeration study, and this CFU count was reported as the LoD in the study.

**[00396] Results:**

**[00397]** The LoD for all MRSA samples tested was determined to be below 10 CFU.

Table 11 summarizes the results of the lowest LoDs obtained in the study.

**[00398] Table 11: Results of the lowest LoDs obtained in the LoD study.**

SCCmec Type	LoD (CFU)
I	3
II	2
IV	3
mecC	1

**[00399]** All MRSA strains tested resulted in fewer than 10 CFU detected with the NRTP assay above a Cutoff calculated from blank samples. MSSA did not generate RLU values above the blank samples Cutoff.

**[00400]** RLU values are shown at the highest dilution for which an RLU value was generated that was above the blank samples Cutoff were plotted as the average RLU value and standard deviation for the four replicates tested for each sample. The horizontal axis is set at the blank samples Cutoff and the CFU counts for the sample that generated each RLU data point is superimposed with the data. All MRSA samples generated RLU values above the Cutoff while MSSA did not.

#### **Cross-Reactivity and Microbial Interference Study**

**[00401]** A cross-reactivity and microbial interference study was performed. The purpose of the study was to test a set of bacterial strains commonly encountered in clinical samples and known to potentially be in the host range of the bacteriophage  $\phi 80\alpha$  in the MRSA Assay to see if there was cross reactivity or interference of these strains with phage or substrate used in the test.

**[00402]** Previous experiments with clinical samples had resulted in false positive results with a presence of *Enterococci faecalis* and *Staphylococcus epidermidis* as indicated from the presence of blue and white colonies when plating on BBL™ CHROMagar™ Staph aureus plates. In addition, *Listeria monocytogenes* and *Listeria innocua* may be within the infective

or penetrative host range of the phage  $\varphi 80\alpha$  which may also contribute to cross-reactivity in the MRSA assay. The study tested *Enterococci faecalis*, *Staphylococcus epidermidis*, *Listeria monocytogenes* and *Listeria innocua* for cross reactivity/ interference with Viability MRSA assay. Each strain was tested at high cell numbers in the order of  $10^6$ ,  $10^7$  or  $10^8$  cells in the assay volume. Tests were done without the addition of GW24 lysate to address potential autoluminescence of strains.

**[00403]** Experiment 1 tested various strains (MSSA-S121, NRS# 9-*Staphylococcus haemolyticus*, NRS # 6- *Staphylococcus epidermidis*, ATCC 12228-*Staphylococcus epidermidis*, ATCC 15305- *Staphylococcus saprophyticus*, ATCC 29212-*Enterococcus faecalis*, ATCC 60193-*Candida albicans*, ATCC 12453-*Proteus mirabilis*) for luminescence at high cell numbers under normal assay conditions.

**[00404]** Experiment 2: A subset of strains that were luminescent from Experiment 1 were re-assayed in the presence of various antibiotics at various concentrations to quench background luminescence.

**[00405]** Experiment 3: *E. faecalis* and S32 (MRSA) were tested with various substrate formulations developed as described above without GW24 lysate and without incubation.

**[00406]** Experiment 4: ATCC 33090-*Listeria innocua* and ATCC 19111-*Listeria monocytogenes* were tested for background signal and non-specific luminescence and retested with various substrate formulations developed as described above along with *E. faecalis* and *S. epidermidis*.

**[00407]** Experiment 5: *E. faecalis* was retested with a final substrate formulation developed as described above.

**[00408]** Substrate Reagent formulations tested in this study are summarized in Table 9.

**Table 10: Substrate Reagent Formulations**

Experiment	Substrate	Description
1	Original Substrate	1% Tween20 + 4.2 mM Tridecanal, pH3.0
2	Original Substrate	
3	Substrate 1	6.3 mM Tridecanal + 0.5% Vitamin E Acetate, pH 3.0
	Substrate 2	20 mM Nonanal + 0.5% Vitamin E Acetate, pH 3.0
	Substrate 3	8.4 mM Tridecanal + 0.5% Vitamin E Acetate, pH 3.0
	Substrate 4	6.3 mM Tridecanal + 1% $\alpha$ -Tocopherol-PEG 1000 Succinate, pH 3.0
4	Original substrate	1% Tween20 + 4.2 mM Tridecanal, pH 3.0

	Substrate 5	0.5% Triton + 4.2 mM Tridecanal (Sigma) + 0.5% Vitamin E Acetate, pH 3.0
5	Substrate 6	6.3mM Tridecanal + 2% VitE PEG, pH 3.0

**Methods/Procedures:**

[00409] The following were steps performed for the MRSA Assay.

[00410] A) Strains Grown for Experiments 1-5

[00411] On the day before the assay, an overnight culture was started in a deep 96 well plate at a 1:50 dilution in TSB from a frozen one-time use stock and incubated at 37°C on an orbital shaker for >15 hours. Bacteria (8 uL) in TSB (392 uL).

[00412] The absorbance of culture was measured on Versamax. The TSB was set as blank in template on SoftmaxPro. Optical density (OD) was measured at 600nm.

[00413] On the day of the assay, cells were re-suspended to an OD 0.5 to set up the assays. Prepared BSS-M56 for Experiments 1-5.

[00414] B) Transduction particle media reagent was prepared for all Experiments 1, 2, 4 and 5 (no transduction particle reagent used in Experiment 3): 15 ug/mL cefoxitin + GW24 lysate stock from as described above at 30X.

[00415] C) Sample Preparation: Various dilutions were made from overnight cultures of strains. All strains were diluted BSS M56.

[00416] D) MRSA Assay was run for Experiments 1-5

[00417] Media was loaded with or without phage and cefoxitin at 5 $\mu$ g/ml to assay plate. 2.5ul cells were added. The assay plate was incubated with a plate lid at 37°C on an orbital shaker with the speed set to approximately 100 rpm for 4 hours.

[00418] Next, the assay plates were measured on the SpectraMax L with the following standard assay parameters:

[00419] Fast Kinetic Luminescence

[00420] Read for 20 time points at 0.5 second intervals. Substrate was injected with M injector with 50 uL/well at 250 ul/sec including 5 baseline reads. No incubation temperature was set and was read at room temperature.

[00421] The SpectraMax L was primed with Substrate Reagent before running the assay.

[00422] The results were analyzed with the following:

[00423] A) Determined cutoff by averaging blank RLU across all replicates and time points and adding three standard deviations.

[00424] B) Determined maximum RLU for each sample using SoftMaxPro.

[00425] C) Determined if the maximum RLU was greater than the cutoff RLU, and if so, then the sample data was used for analysis.

### **Results Summary**

[00426] Experiment 1: Various strains were tested for cross reactivity and interference using the Original Substrate formulation, out of those tested, NRS# 9- *S. haemolyticus*, NRS # 6- *S. epidermidis* and *E. faecalis* tested false positive in MRSA assay.

[00427] Experiment 2: Out of the three strains tested, NRS #9 and *E. faecalis* tested MRSA positive with all Cefoxitin conditions tested. All three strains (NRS #9, *E. faecalis*, NRS #6) tested positive when no transduction particle reagent was used in the assay, indicating that non-specific luminescence was not transduction particle reagent-dependent but rather strain and substrate reagent dependent. Carb (Carbencillin) at all concentrations tested was effective in removing the false positive signal.

[00428] Experiment 3: *E. faecalis* gave a positive signal without transduction particle reagent. MRSA strain S32 also gave a positive signal without transduction particle reagent. This result was indicative of the substrate reagent causing background luminescence. Substrate 4 was effective in eliminating background signal in the assay.

[00429] Experiment 4: Strains ATCC 33090 -*Listeria innocua*, ATCC 19111-*Listeria monocytogenes*, were tested for luminescence with transduction particle reagent and substrate reagent as *Listeria sp.* can be within the host range of the bacteriophage used in the MRSA assay. Luminescence was observed from *L. innocua* with and without transduction particle reagent using Original Substrate formulation indicating that the luminescence was due to non-specific reaction potentially with the substrate. Substrate 5 was effective in eliminating luminescence from *Listeria* but not *E. faecalis*.

[00430] Experiment 5: Retested *E. faecalis* with Substrate 6. In two independent runs on two different days with high load of cells at 0.5 OD, the assay yielded negative results.

### **Conclusions**

[00431] The cross-reactivity study demonstrated background luminescence from several bacterial species at high loads. The light output did not require transduction particle reagent and certain substrate formulations utilizing phosphate ions contributed to non-specific signal. Because no light output from cross-reactive species was observed from the use of transduction particle reagent, in the case that  $\phi$ 80 $\alpha$  penetrates cross-reactive species, light output is prevented from the lack of activity of the *S. aureus* *PclpB* promoter that is operatively linked to the bacterial luciferase genes and/or the lack of activity of the *S. aureus* *pT181* origin of replication within these species.

[00432] Replacing the buffer from sodium phosphate dibasic in the formulation with sodium citrate and citric acid eliminated background luminescence from all cross-reactive species tested except for *E. faecalis*. Substrate 6 with the added ingredient of  $\alpha$  tocopherol-PEG 1000 Succinate eliminated the remaining non-specific signal from *E. faecalis*.

**Clinical Performance of Non-Replicative Transduction Particle-Based Viable Cell Reporter MRSA Assay - Results with Reference to Direct Plating onto CHROMAgar MRSA II**

[00433] A MRSA screening assay was developed employing  $\varphi 80\alpha$ -based luxAB expressing non-replicative transduction particles (NRTP). The assay consisted of adding NRTP to a clinical sample suspected of containing MRSA, incubating the sample for a period of 4 hours at 37°C, and then assaying the incubated sample by injecting an aldehyde into the sample while measuring for luminescence with a photomultiplier tube. The results of the assay were compared to that of commercially available chromogenic media designed for the detection of MRSA as a reference in order to determine the sensitivity and specificity of the assay. The NRTP-based assay was expected to correlate well with the culture-based reference since both require the presence of viable MRSA cells and both rely on the expression of the MRSA phenotype. The results showed excellent correlation with the reference.

[00434] The purpose of the study was to determine the performance of the NRTP-based MRSA Assay with reference to CHROMAgar MRSA II from testing remnant nasal swab samples collected for the purpose of MRSA screening.

[00435] Scope:

[00436] De-identified nasal swab samples collected from patients for the purpose of MRSA surveillance by a clinical institution were tested for the presence of MRSA using the NRTP-based MRSA Assay, CHROMAgar MRSA II, CHROMAgar SA and Blood Agar TSA via direct plating and via enriched culture followed by plating. The results of the NRTP-based MRSA Assay were compared with the results of the CHROMAgar MRSA II assay in order to calculate the sensitivity and specificity of the NRTP-based MRSA Assay with reference to CHROMAgar MRSA II.

[00437] The following key materials were used in the clinical performance study:

[00438] Growth Media Reagent: BSS-M56

[00439] Substrate Reagent: Final Substrate Reagent Formulations to be stored at 18-24°C as described as described above

[00440] Transduction Particle Reagent: BSS-M56 base with 10ug/mL (*i.e.* 2X concentration) cefoxitin and transduction particle reagent as described above at 2X concentration.

**Methods/Procedures**

[00441] Clinical Sample Description: Sample transport tubes containing liquid Amies (220093 - BD BBL™ CultureSwab™ Liquid Amies) were provided to a clinical institution for collecting de-identified remnant nasal swabs collected by the clinical institution. Prior to placing the nasal swabs into the provided sample transport tube, the clinical institution used the swab for performing their own direct culture MRSA screening by streaking the swab onto a culture plate. More specifically, anterior nares specimens were collected at the clinical institution internal standard procedures and using the clinical institution's standard collection swab. The clinical institution then performed direct culture screening with the swab. The remnant swab was then added to the sample transport tube in which the swab tip was submerged in the Amies buffer in the sample transport tube. Samples were then kept at room temperature for 2-24 hours prior to further processing.

[00442] Sample Handling: Upon receipt, samples were stored overnight at room temperature in a biosafety cabinet upright to ensure swab immersion in the sample transport tube Amies buffer. After overnight storage, samples were further processed as follows.

[00443] Clinical Sample Preparation

[00444] Using a 1 mL Pipette, 300  $\mu$ l of Growth Media Reagent was added to 15 mL falcon tubes.

[00445] The swabs from remnant nasal swabs were removed from the original transport tube and immersed into the Growth Media Reagent in a corresponding falcon tube. The swab contents were then eluted into the Growth Media Reagent in the falcon tube by rolling it back and forth in the Growth Media Reagent 4-6 times. The swab was then placed back into the original transport tube and stored at 2-8 °C until the end of the study while the eluted clinical samples in the falcon tube were transferred to 1.5 mL tubes and kept at room temperature until further processing.

[00446] Running the NRTP MRSA Assay: The following samples were loaded directly into a white 96 well assay plate.

[00447] Clinical Samples: 100  $\mu$ l of the eluted material of each clinical sample in singlet.

[00448] MRSA positive control: 2  $\mu$ l of a thoroughly mixed 0.1 OD culture of a known MRSA isolate into 98 uL of Growth Media Reagent in triplicate.

[00449] MSSA negative control: 2  $\mu$ L of a thoroughly mixed 0.1 OD culture of a known MSSA isolate into 98  $\mu$ L of Growth Media Reagent in triplicate.

[00450] Blanks: 100  $\mu$ L of Growth Media Reagent in triplicate.

[00451] To each sample, 100  $\mu$ L of Transduction Particle Reagent was added. The assay plate was then placed in an incubator set at 37°C, shaking on orbital shaker for 4 hours. At the end of 4 hours, the plate was removed from incubator and immediately measured for luminescence on a SpectraMax L that injected 50  $\mu$ L of the Substrate Reagent and measured luminescence for a period of 1 minute.

[00452] Bacteria Plating for clinical sample CFU enumeration: Each eluted clinical sample was plated in order to determine bacterial colony counts on CHROMAgar MRSA II, CHROMAgar SA and Blood Agar (TSA II) via direct and enriched culture as follows. Organism CFU counts were determined by direct plating. MRSA CFU counts were determined by plating on CHROMAgar MRSA II. *S. aureus* CFU counts were determined by plating on CHROMAgar SA plate. CFU counts of any organism whose growth is supported by Blood Agar TSA were determined by plating on Blood Agar TSA. In the case that direct plating did not produce colonies due to the load of organisms being below the limit of detection of the plates used, sample enrichment was also performed by incubating a portion of the eluted clinical sample in TSB overnight at 37°C with shaking and then again plating the enriched culture on CHROMAgar MRSA II. All plates were incubated for 20-24 hours at 37°C. After incubation, the CFU counts of any colonies appearing on each plate were recorded.

[00453] Analysis: The presence and CFU load of MRSA, *S. aureus*, and total organisms per eluted clinical sample were calculated based on the CFU counts obtained on CHROMAgar MRSA II, CHROMAgar SA, and Blood Agar TSA, respectively.

[00454] NRTP Assay analysis: Data from each sample were plotted as RLU vs. time.

[00455] Cutoff Determination: The Assay Cutoff was calculated from all time points of the blank samples using the following formula: (Mean Blank RLU + 3\* SD Blank RLU).

[00456] MRSA Positive Determination: The RLU of each time point after substrate injection was determined to be above or below the Assay Cutoff. If two or more data points after injection were above the Assay Cutoff then the sample was designated as “MRSA Positive.”

[00457] Results: The MRSA positive results of the NRTP Assay were compared to those of the direct and enriched culture plating onto CHROMAgar MRSA II. The following

calculations were conducted in order to determine the NTRP Assay Sensitivity and Specificity with reference to CHROMAgar MRSA II.

• True Positive (TP)
○ Sample that produced a MRSA positive result on both the NTRP Assay and CHROMAgar MRSA II
• True Negative (TN)
○ Sample that produced a MRSA negative result on both the NTRP Assay and CHROMAgar MRSA II
• False Positive (FP)
○ Sample that produced a MRSA positive result on the NTRP Assay and a MRSA negative result on CHROMAgar MRSA II
• False Negative (FN)
○ Sample that produced a MRSA negative result on the NTRP Assay and a MRSA positive result on CHROMAgar MRSA II
• Sensitivity = TP/(TP+FN)
• Specificity = TN/(TN+FP)

**Results with Reference to Direct Plating onto CHROMAgar MRSA II**

[00458] Table 11 shows following results were obtained comparing the NTRP Assay with reference to direct plating on CHROMAgar MRSA II.

**Table 11. NTRP Assay Results vs. Direct Plating on CHROMAgar MRSA II Results**

Total Samples	CHROMAgar MRSA II Positive	CHROMAgar MRSA II Negative	NTRP ASSAY Positive	NTRP ASSAY Negative	True Positive	True Negative	False Positive	False Negative
69	7	62	12	57	7	57	5	0

[00459] Based on the above data, the sensitivity and specificity of the assay with reference to direct plating onto CHROMAgar MRSA II were calculated to be:

[00460] Sensitivity = 100%

[00461] Specificity = 92%

**Clinical Performance of Non-Replicative Transduction Particle-Based Viable Cell Reporter MRSA Assay - Results with Reference to Enriched Culture, followed by Plating onto CHROMAgar MRSA II**

**[00462]** Based on the results with reference to direct plating on CHROMAgar MRSA II, all clinical samples were re-tested with reference to enriched culture, followed by plating on CHROMAgar MRSA II. The rationale for the follow-on testing was based on the possibility that false positive results when compared to direct plating may indeed be true positives that were detected by the NRTP assay but may have been missed by direct plating. A portion of the remaining eluted swab samples were re-tested via the NRTP assay as described above. Another portion of the remaining eluted swab samples were also tested via enriched culture, followed by plating onto CHROMAgar MRSA II. Enriched culture testing consisted of adding 100 uL of the remaining eluted swab material to 2 mL of TSB and incubating at 37C with shaking for a period of 18-24 hours. The resulting culture was then streaked onto CHROMAgar MRSA II in order to determine the presence of MRSA in the culture. Table 12 summarizes the data from both the direct plating and enrichment followed by plating assays – only the samples that produced a MRSA positive result on either NRTP Assay or CHROMAgar MRSA II are shown.

**Table 12: NRTP Assay Results vs. Direct Plating and Enriched Culture followed by Plating on CHROMAgar MRSA II**

**[00463]** Only the samples that produced a MRSA positive result on either NRTP Assay or CHROMAgar MRSA II are shown.

Sample #	NRTP Assay	Direct CHROMagar MRSA II	Enrichment + NRTP Asssay	Enrichment + CHROMagar MRSA II
1	+	+	+	+
2	+	+	+	+
3	+	+	+	+
4	+	+	+	+
5	+	+	+	+
6	+	+	+	+
7	+	+	+	+
8	+	-	+	+

9	+	-	+	+
10	+	-	+	+
11	+	-	+	+
12	+	-	+	-

[00464] Table 13 shows following results were obtained comparing the NRTP Assay with reference to enriched culture of clinical samples, followed by plating on CHROMAgar MRSA II.

**Table 13. NRTP Assay Results vs. Enriched Culture Followed By Plating on CHROMAgar MRSA II Results**

Total Samples	CHROMA gar MRSA II Positive	CHROMA gar MRSA II Negative	NRTP ASSA Y Positive	NRTP ASSA Y Negative	True Positive	True Negative	False Positive	False Negative
69	11	58	12	57	11	57	1	0

[00465] Based on the above data, the sensitivity and specificity of the assay with reference to enriched culture followed by plating onto CHROMAgar MRSA II was calculated to be:

- Sensitivity = 100%
- Specificity = 98.3%

**Example 8: NRTP-Based Assay For Antimicrobial Susceptibility Testing - Correlation Of Minimum Inhibitory Concentration To Luminescence Output**

[00466] In another example, a *S. aureus* cefoxitin susceptibility assay was developed to determine the minimum inhibitory concentration of cefoxitin required to inhibit the growth of cefoxitin resistant *S. aureus*. Unlike a MRSA cefoxitin resistance assay as described above, which differentiates cefoxitin sensitive from cefoxitin resistant *S. aureus*, the MRSA cefoxitin susceptibility assay in this example describes the development of an assay to determine the minimum amount of cefoxitin needed to inhibit the growth of *S. aureus* in the presence of cefoxitin.

[00467] The following key materials were used in the clinical performance study:

[00468] Growth Media Reagent: BSS-M56

[00469] Substrate Reagent: Final Substrate Reagent Formulations to be stored at 18-24°C as described in Example 7.

[00470] Transduction Particle Reagent: BSS-M56 base with 10ug/mL (*i.e.* 2X concentration) cefoxitin and transduction particle reagent as described in Example 7 at 2X concentration MIC Study Protocol.

[00471] Overnight Culture: For each MRSA strain (NRS35 and S7) and a MSSA negative control strain (MSSA121), 2 mL of TSB were inoculated with a colony of the strain previously grown on TSA plates. Overnight MRSA cultures included 5ug/mL cefoxitin. All samples were incubated overnight at 37°C in a shaking incubator.

[00472] Day Culture: 20 uL of each of the overnight cultures were transferred into a new culture tube containing 2 mL of Growth Media Reagent. The inoculums were then incubated at 37C with shaking for approximately 1hr 45min, until the OD(600nm) reached 0.1.

[00473] MIC determination via plating:

[00474] a) Each of the day cultures was streaked onto TSA plates containing cefoxitin at 4, 8, 16, 32, 64, and 128 ug/mL.

[00475] b) Plates were incubated for 18 hours at 37C to determine growth.

[00476] NRTP Assay preparation:

[00477] a) Wells of a white 96 well assay plate were filled with 100 uL of 2x Transduction Particle Reagent.

[00478] b) For each of the day cultures, five wells were then filled with 100 uL of day culture.

[00479] c) For each of the day cultures, cefoxitin was added to one well each such that the cefoxitin concentration in the well was at 4, 8, 16, 32, 64, and 128 ug/mL.

[00480] d) The plate was then sealed with a breathable seal and incubated for 4 hours at 37°C with moderate shaking, 50rpm.

[00481] At the end of 4 hours, the plate was remove from incubator and immediately measured for luminescence on a SpectraMax L that injected 50  $\mu$ l of the Substrate Reagent and measured luminescence for a period of 1 minute.

Analysis:

[00482] The maximum luminescence value after Substrate Reagent addition from each sample was plotted. MSSA sample RLU values were used to determine a Cutoff calculated using the following formula: (Mean MSSA RLU + 3\* SD MSSA RLU).

[00483] Results:

**[00484]** Figure 23 shows the results of *S. aureus* growth at 4, 8, 16, 32, 64, and 128 ug/mL of cefoxitin. Figure 24 shows the RLU values obtained by the NRTP assay in the presence of 4, 8, 16, 32, 64, and 128 ug/mL cefoxitin. The x-axis in Figure 24 is set at the MSSA RLU cutoff value.

**[00485]** As can be seen in Figure 23, MRSA NRS25 exhibited a MIC of 128 ug/mL cefoxitin while MRSA S7 exhibited a MIC of 64 ug/mL cefoxitin. Correspondingly, MRSA NRS25 exhibited appreciable luminescence above the MSSA RLU cutoff to a cefoxitin concentration up to 64 ug/mL cefoxitin while MRSA S7 exhibited luminescence above the MSSA RLU cutoff to a cefoxitin concentration up to 32 ug/mL.

**[00486]** Based on the above data, the NRTP assay demonstrates that RLU values obtained from the assay correlate with MIC results and thus the NRTP assay may be used to develop antibiotic susceptibility assays.

**Example 9: Transcript Reporter Assay: Mechanism of Conformational Change by RBS-Blocking Cis-Repression Of *Luxab* Translation Activated By The *mecA* Gene Transcript Of *Mrsa***

**[00487]** As described above, a reporter transcript can be designed such that translation of the reporter gene sequence is blocked by cis-repression of the ribosome-binding site (RBS) of the reporter gene.

**[00488]** The following tools were used for designing the reporter transcripts of the invention.

**[00489]** 1) RNA secondary structure was calculated using secondary structure program, such as Mfold (<http://mfold.rna.albany.edu/?q=mfold/RNA-Folding-Form>).

**[00490]** 2) Intermolecular RNA interactions were calculated using a software program such as RNA-RNA InterACTion prediction using Integer Programming (RactIP) (<http://rna.naist.jp/ractip/>).

**[00491]** 3) RNA secondary structure was visualized using Visualization Applet for RNA (VARNA) (<http://varna.lri.fr/>).

**[00492]** Figure 25 shows a secondary structure of the *mecA* transcript generated based on the lowest energy conformation calculated by MFold and visualized with VARNA. The terminal loop 23 (T23) contains a YUNR sequence UUGG consisting of bases 1,487-1,490 of the *mecA* transcript sequence. Analysis of the secondary structure of the *mecA* gene transcript revealed several ssRNA regions that were suitable for designing a cis-repressed *luxAB* reporter that can be de-repressed via interactions between the reporter and an ssRNA region.

**[00493]** As shown in detail in Figure 26, the terminal loop 23 (T23) of the *mecA* transcript contains a YUNR consensus sequence. A YUNR (pYrimidine-Uracil-Nucleotide-puRine) consensus sequence has been shown to be a critical target for intermolecular RNA complexes in natural systems. A cis-repressing sequence was designed to form a stem-loop structure with the RBS of the reporter sequence, such that the cis-repressing sequence blocks binding of an RNA polymerase to the RBS of the reporter sequence. The reporter sequence was exposed upon binding of the loop of the cis-repressing stem-loop structure with T23 of the *mecA* transcript.

**[00494]** As shown in Figure 27, a cis-repressing sequence 2701 was added to the 5' terminus of the *luxAB* genes and designed to form a stem-loop structure that blocks the RBS sequence (“AAGGAA”) 2702 of the *luxA* gene. The cis-repressing stem-loop structure was predicted to block the *luxA* RBS (“AAGGAA”) sequence, based on the lowest energy conformation of the *luxAB* transcript including the cis-repressing sequence at the 5' terminus of the *luxAB* transcript, as calculated by MFold and visualized with VARNA.

**[00495]** The first 61 nucleotides of the cis-repressed *luxAB* genes are shown in FIG. 7, up to the start codon AUG of the *luxA* gene. The RBS sequence “AAGGAA” includes bases 47-52. This terminal loop of the reporter transcript was designed to interact with (bind to) the terminal loop 23 (T23) of the *mecA* transcript, which contains a YUNR sequence.

**[00496]** The terminal loop of the cis-repressing sequence was designed to interact with T23 of the *mecA* transcript, such that hybridization of the cis-repressed *luxAB* transcript and the *mecA* transcript via the interaction of the loop from the cis-repressing stem-loop structure and T23 of the *mecA* transcript results in exposure of the RBS of the *luxA* gene. Figure 28 shows the predicted inter-molecular interactions between the *mecA* T23 sequence and the cis-repressing sequence on the *luxAB* transcript calculated by RactIP and visualized by VARNA. Lines indicate base pairing between the *mecA* transcript and the cis-repressed *luxAB* transcript. The interaction between the two sequences results in exposure of the *luxA* RBS sequence AAGGAA and thus de-repression of the *luxAB* reporter.

**Example 10: Transcript Reporter Assay: Methods of Detecting Target**

**Transcripts or Genes Using a *mecA* – *luxAB* Reporter System**

**[00497]** In another example, a method for detecting a target *mecA* gene is provided using a *mecA*-*luxAB* reporter system. Here, *mecA* is the target transcript, and *luxAB* is the reporter molecule.

### 1. Construction of the Reporter Construct

**[00498]** A vector comprising a reporter construct encoding *luxAB* can be constructed via standard molecular biological techniques by incorporating the reporter construct into a shuttle vector capable of propagating in both *E. coli* and *S. aureus*. The vector can contain an origin of replication that is functional in *E. coli* and a selectable marker that is expressed in *E. coli* and suitable for allowing the growth of *E. coli* cells transformed with the vector and grown under selective conditions. The vector can also contain an origin of replication that is functional in *S. aureus* and a selectable marker that is expressed in *S. aureus* and suitable for allowing the growth of *E. coli* cells transformed with the vector and grown under selective conditions. Propagation of the vector for conducting *in vitro* manipulations and for verification of manipulations can be accomplished via a suitable laboratory cloning strain of *E. coli* and the final modified vector can then be introduced into *S. aureus* strains.

**[00499]** The reporter construct can be first introduced into a *S. aureus* cell for transcribing the construct and producing the reporter transcript.

### 2. Construction of a cis-Repressed Reporter Transcript

**[00500]** Methods are provided for constructing a cis-repressed reporter transcript that can bind to a *mecA*-target transcript. The reporter transcript can be constructed via standard molecular biological techniques. The *luxA* and *luxB* genes serve as reporter genes and can be derived from *Vibrio harveyi*. The genes lack a transcriptional promoter, and each contains its own ribosomal binding site (RBS). When both the *luxA* and *luxB* genes are translated in a cell, the *luxA* and *luxB* proteins complex to form the active luciferase enzyme (LuxAB). See Farinha, M.A. and A.M. Kropinski, Construction of broad-host-range plasmid vectors for easy visible selection and analysis of promoters. *J. Bacteriol.*, 1990. 172(6): p. 3496-3499.

**[00501]** The cis-repressing sequence can be situated upstream of the *luxAB* genes and downstream of a promoter and includes a sequence that is complementary to the *luxA* RBS. A linker sequence can separate the complementary regions of the cis-repressing sequence and the *luxA* sequence. After transcription of the vector, the complementary regions of the cis-repressing sequence and the *luxA* RBS sequence complex, creating a stem loop that prevents docking of a ribosome and hence translation.

**[00502]** The stem loop of the reporter transcript is designed to destabilize and form an open complex when it interacts with a naturally-occurring *mecA* transcript sequence (endogenous to the cell). To activate translation of the *luxA* gene sequence, the natural *mecA* transcript serves as a trans-activating RNA that binds to the cis-repressed reporter transcript and opens the inhibitory stem loop that sequesters the RBS of the *luxA* gene. Once the RBS

is not sequestered by the cis-repressing sequence, translation of *luxA* can occur.

Transcription of the reporter construct is accomplished via operatively linking the reporter sequence to a constitutive promoter, upstream of the cis-repressing sequence.

**[00503]** An example of a target *mecA* gene sequence is shown in Figure 29. The sequence is a *mecA* gene loci DNA sequence (from *Staphylococcus aureus* subsp. *aureus* SA40, complete genome GenBank: CP003604.1; SEQ ID NO:15) and can be used for generating a reporter construct comprising a reporter sequence and a cis-repressing sequence. The -10 position 2901, the transcription start position 2902, the RBS 2903, the coding region (in grey 904) and the transcription termination sequence 2905 are shown.

**[00504]** Figure 30 shows an exemplary *mecA* transcript sequence that can be used for designing a reporter transcript (SEQ ID NO:16), according to an embodiment of the invention. The RBS 3001 and the coding sequence 3002 are shown for *mecA*.

**[00505]** Figure 31 is an example of a *luxAB* gene loci DNA sequence that can be used for designing a reporter transcript, according to an embodiment of the invention. The *luxAB* gene loci DNA sequence was obtained from *Vibrio fischeri* genes *luxA* and *luxB* for luciferase alpha and beta subunits (GenBank: X06758.1) (SEQ ID NO: 17). The -10 position 3101, the transcription start position 3102, the RBS for lux A 3103, the *luxA* coding sequence 3104 (gray shading), the RBS for *luxB* 3105, and the *luxB* coding sequence (gray shading) 3106 are shown.

**[00506]** Figure 32 is an example of a *luxAB* transcript sequence that can be used for designing a reporter transcript (SEQ ID NO:18). The RBS for lux A 3201, the *luxA* coding sequence 3202 (gray shading), the RBS for *luxB* 3203, and the *luxB* coding sequence (gray shading) 3204 are shown.

**[00507]** Figure 33 is an example of a *luxAB* cis-repressed transcript sequence that can be used in a reporter transcript (SEQ ID NO:19). The cis-repressing sequence (dotted line box) 3301, the RBS for lux A 3302, the *luxA* coding sequence 3303 (gray shading), the RBS for *luxB* 3304, and the *luxB* coding sequence (gray shading) 3305 are shown.

### 3. *Methods for Detecting the Presence or Absence of a *mecA* Target Transcript Using the Reporter Transcript*

**[00508]** Examples are provided for detecting the presence or absence of a *mecA* target transcript in a cell using the reporter transcripts of the invention. Figure 34 shows an example of a cell comprising a vector 3400 that encodes a reporter transcript 1410, where there is no endogenous *mecA* transcript in the cell 3401 (e.g., the cell's genome does not contain the *mecA* gene). In this case, the cis-repressing sequence 3420 binds to the RBS 3430

of the *luxAB* genes. In some embodiments, the cis-repressing sequence 3420 can bind to a portion of or all of the RBS of the *luxA* gene, the RBS of the *luxB* gene, or both. This binding event blocks and prevents the translation of the *luxAB* genes, and the reporter molecule (e.g., luciferase) is not produced in the cell. Thus, no signal is detected, indicating the absence of the *mecA* gene in the cell.

**[00509]** In another example, the cell includes an endogenous *mecA* transcript (e.g., the cell's genome contains the *mecA* gene). Figure 35 shows a vector 3400 introduced into a cell 3401. The vector 3400 encodes the reporter transcript 3410, which includes a cis-repressing sequence 3420 and a reporter sequence (*luxA* and *luxB* genes). When the *mecA* transcript 3510 present in the cell binds to the cis-repressing sequence 1420, the inhibitory hairpin loop opens up and the RBS 3430 for the *luxA* gene is exposed. Translation of the reporter sequences (*luxA* and *luxB*) can occur, resulting in the formation of a *luxAB* enzyme 3520. The *luxAB* enzyme 3520 produces a detectable luminescent signal 3530. In this manner, the transcript reporter vector 3400 reports the presence of endogenous *mecA* transcripts 3510 within a cell 3401.

**[00510]** While the invention has been particularly shown and described with reference to a preferred embodiment and various alternate embodiments, it will be understood by persons skilled in the relevant art that various changes in form and details can be made therein without departing from the spirit and scope of the invention.

**[00511]** All references, issued patents and patent applications cited within the body of the instant specification are hereby incorporated by reference in their entirety, for all purposes.

**REFERENCES CITED**

1. Michael G. Schmidt, D.A.S., Caroline Westwater, Joseph W. Dolan, Brian D. Hoel, Philip A. Werner, James S. Norris, Laura M. Kasman, Nucleic Acid Delivery and Expression, 2005.
2. Kreiswirth, B.N. *et al.*, The toxic shock syndrome exotoxin structural gene is not detectably transmitted by a prophage. *Nature*, 1983. 305(5936): p. 709-712.
3. Ubeda, C. *et al.*, Specificity of staphylococcal phage and SaPI DNA packaging as revealed by integrase and terminase mutations. *Molecular Microbiology*, 2009. 72(1): p. 98-108.
4. Otsuji, N. *et al.*, Induction of Phage Formation in the Lysogenic *Escherichia coli*K-12 by Mitomycin C. *Nature*, 1959. 184(4692): p. 1079-1080.
5. Brantl, S. (2007) Regulatory mechanisms employed by cis-encoded antisense RNAs. *Curr. Opin. Microbiol.* 10, 102–109.
6. Isaacs, F.J. *et al.* (2004) Engineered riboregulators enable post-transcriptional control of gene expression. *Nat. Biotechnol.* 22, 841–847.
7. Pfeiffer, V. *et al.* (2009) Coding sequence targeting by MicC RNA reveals bacterial mRNA silencing downstream of translational initiation. *Nat. Struct. Mol. Biol.* 16, 840–846.
8. Opdyke, J.A. *et al.* (2004) GadY, a small-RNA regulator of acid response genes in *Escherichia coli*. *J. Bacteriol.* 186, 6698–6705.
9. Carriere, C., et al., *Conditionally replicating luciferase reporter phages: Improved sensitivity for rapid detection and assessment of drug susceptibility of Mycobacterium tuberculosis*. *Journal of Clinical Microbiology*, 1997. 35(12): p. 3232-3239.
10. Merten, O.-W. and M. Al-Rubeai, *Viral Vectors for Gene Therapy : Methods and Protocols*. Methods in Molecular Biology. Vol. 737. 2011.
11. Lofdahl, S., J.E. Sjostrom, and L. Philipson, *CLONING OF RESTRICTION FRAGMENTS OF DNA FROM STAPHYLOCOCCAL BACTERIOPHAGE-PHI-11*. *Journal of Virology*, 1981. 37(2): p. 795-801.
12. Charpentier, E., et al., *Novel Cassette-Based Shuttle Vector System for Gram-Positive Bacteria*. *Appl. Environ. Microbiol.*, 2004. 70(10): p. 6076-6085.
13. Novick, R.P., I. Edelman, and S. Lofdahl, *Small staphylococcus-auerus plasmids are transduced as linear multimers that are formed and resolved by replicative processes*. *Journal of Molecular Biology*, 1986. 192(2): p. 209-220.

14. Westwater, C., et al., Development of a P1 phagemid system for the delivery of DNA into Gram-negative bacteria. *Microbiology*, 2002. 148(4): p. 943-950.
15. Norris, J.U., et al., *Tissue-Specific and Pathogen-Specific Toxic Agents and Ribozymes*. 1999.
16. Maiques, E., et al., Role of Staphylococcal Phage and SaPI Integrase in Intra- and Interspecies SaPI Transfer. *J. Bacteriol.*, 2007. 189(15): p. 5608-5616.
17. Frees, D., et al., Clp ATPases are required for stress tolerance, intracellular replication and biofilm formation in *Staphylococcus aureus*. *Molecular Microbiology*, 2004. 54(5): p. 1445-1462.
18. Arnaud, M., A. Chastanet, and M. Debarbouille, New Vector for Efficient Allelic Replacement in Naturally Nontransformable, Low-GC-Content, Gram-Positive Bacteria. *Appl. Environ. Microbiol.*, 2004. 70(11): p. 6887-6891.
19. Tormo, M.A., et al., *Staphylococcus aureus* Pathogenicity Island DNA Is Packaged in Particles Composed of Phage Proteins. *J. Bacteriol.*, 2008. 190(7): p. 2434-2440.
20. Arthur, M., et al., The VanS sensor negatively controls VanR-mediated transcriptional activation of glycopeptide resistance genes of Tn1546 and related elements in the absence of induction. *J. Bacteriol.*, 1997. 179(1): p. 97-106.
21. Karlsson, S., et al., Expression of *Clostridium difficile* Toxins A and B and Their Sigma Factor TcdD Is Controlled by Temperature. *Infect. Immun.*, 2003. 71(4): p. 1784-1793.
22. Daniel Sobek, J.R., Enzyme detection system with caged substrates, 2007, Zymera, Inc.
23. Samie Jaffrey, J.P., Coupled recognition/detection system for in vivo and in vitro use, 2010, Cornell University.
24. Good, L., Translation repression by antisense sequences. *Cellular and Molecular Life Sciences*, 2003. 60(5): p. 854-861.
25. Sabine, B., Antisense-RNA regulation and RNA interference. *Biochimica et Biophysica Acta (BBA) - Gene Structure and Expression*, 2002. 1575(1-3): p. 15-25.

## INFORMAL SEQUENCE LISTING

SEQ ID NO: 1

Native P1 pac-site

CCACTAAAAAGCATGATCATTGATCACTCTAATGATCAACATGCAGGTGATCACATTGCGGC  
 TGAAATAGCGGAAAAACAAAGAGTTAATGCCGTTGTCAGTGCCGAGTCGAGAAATGCGAAGC  
 GCCAAAATAAGCGCATAAAATGATCGTTCAGATGATCATGACGTGATCACCCGC

SEQ ID NO:2

P1 pac-site with silent mutations, lower case letters signify mutated bases

CCACTAAAAAGCATGATaATaGAcCACTCTAACGAcCAACATGCAGGgGAgCACATTGCGGC  
 TGAAATAGCGGAAAAGCAGAGGTgAATGCCGTTGTCAGTGCCGAGTCGAGAAATGCGAAGC  
 GCCAAAATAAGCGCATAAAACGAcCGTTCAGACGAcCATGACGTtATTACCCGC

SEQ ID NO: 3

P1 lytic replicon containing the C1 repressor-controlled P53 promoter, the promoter P53 antisense, the repL genes, and an in-frame deletion of the kilA gene

CACTATAGGGCGAATTGGCGGAAGGCCGTCAAGGCCGCATTGGGCCGGCGGCCGGATCC  
 GCTAGCTCTAGACTGGCAGGTTCTGAGCAGATCGTCCAACCCGATCTGGATCGGGTCAGAA  
 AAATTTGCTCTAATAAATTCTGTTCTAAGTGCAGAAAGAATCACCATTGAGCTGGTGTATT  
 GAAGGTTGATGCAAATTGGAGAAAAATGCAACAAACATTCAATGCCGATATGAATATATC  
 AACACCTTCATCAAAATGTCGATCCTCAACCACACTGCCGTATTGTTGTTGAAATT  
 CGACCGACCGCGCTGGCGTTACACCTTAATGCTCTACACAGAGCGAGCGGACTCGGTGCC  
 CATAAAGCGCCAGCTCAATGGCTAAGAACGCTGTCAGCTAACAGCTCATCGAAAGAGCTTGA  
 AAAAGAAACTATGCAAAATTGCATAGTTCTGTTACAAGCAATGGAAGCAGGATTCTTCA  
 CGACTCGTATAACCGCAAAGGTCAAGCAGTGGCTGATGAAGCGATTGCTGATGCTGGTGTG  
 CTGGTACCTGTCGCGCAACCGCTAACAGACGTAGTAAGAACCCACAGCATTGTAATGCTG  
 GCTAAAGTCACTTCTGAGCTGTATAACGATGAGCGATTCTGCTGATGCTGGTGTG  
 GGCCTGCTTGTAAACACACTCCGGTCTATCCCGTAGCGCCGGCATATCCTGTCGCAATGTC  
 CAAATCTCGCGCAACAAACCGTGAATACTCATTACAAGCCTCACCGCCTGATCGCGCA  
 GAAACTGGTTAGCCAATCAACCGTCGTTGTCGATTCCGTGAAGCTGTAAACAAAGGAAT  
 TCTGTCGTTAGAGATTGTTATCGCGATCACCGTGAACGTCGCGCTAACCTGTACCGGTTA  
 CACCATCCTTTGGCCTTCGCACAACAAGCCAAAATGCGCTGATAGAAAGCAAATTAAAG  
 ATCTCTTCAGCGGCAACCAAGGTTAAAGCTGTTCTGCTAAGACATTGGCTTATTAAATT  
 TTTATCCACACCCCCATGTCAAAATGATAACCCCTCCCCGTCAAGGATGACGTGGCAATAA

AGAATAAGAAGTCACAAGTTAAAAAACAAAAAGATCAGTTCCGGCGGTGCCGGAACAACC  
AGCCTCAAAAATTGACTTCATGGATCGCTAAGGCAAAGCAAAGGCTGACAATCTGCGGTT  
ATCCAAAAACGCACTCAAAACATGAGTTCAAGCAGAAAGTAGAGGCGGCTGCGCGGAAAT  
ATGCTTACCTGAAGAACAGCGTTCGCCTGATATTGGCGGGATATCAAACCTCGATAACCTA  
CCGCATTGCATGACGGTAAACGAAGCTCTTAATGCGGTTTAGCCAAAATAAGATAACGA  
ACAATGGGTATAACCGCAGGATTCAAGAGGGTAATGAATTGCTCTAATTATAACCATGCATA  
CTTTCAACACCTCTAGTTGCCATGAGGCAAACACTCATAGGTGTCCTGGTAAGAGGACACTGT  
TGCCAAAACGGACGCCATTATTGCAATTAAACAACTAACGGACAATTCTACCTAAC  
AATAAGTGGCTTAAAAAACCGCCCCGGCGGGTTTTTATCTAGAGCTAGCGGATCCGGC  
GCGCCGGGCCCTCTGGGCCTCATGGCCTCCGCTCACTGCCGCTTCCAG

SEQ ID NO: 4

Pblast promoter sequence

CGTCAGGTGGCACTTTCGGAAATGTGCGCGGAACCCCTATTGTTATTTCTAAATACA  
TTCAAATATGTATCCGCTCATGAGACAATAACCTGATAATGCTCAATAATATTGAAAAG  
GAAGAGT

SEQ ID NO: 5

*S. aureus* pT181 plasmid origin or replication copy number variant pT181cop-623 repC

TTTGCAGAAAGAGTTAGTAAGTTAACAGAAGACGAGCCAAACCTAAATGGTTAGCAGGAAA  
CTTAGATAAAAAAAATGAATCCAGAATTATATTAGAACAGGAACAGCAACAAGAGCAACAAA  
AGAATCAAAACGAGATAGAGGTATGCACTTATAGAACATGCATTATGCCAGAAAAACTTA  
TTGGTTGGAATGGGCTATGTGTTAGCTAACTGTTAGCGAGTGGTGGACTTGAATTGGGA  
TTAATCCAAGAAAGTACCGGCTCAACAACCCATAAAGCCCTGTAGGTTCCGNCCAATAAGG  
AAATTGGAATAAAGCAATAAAAGGAGTTGAAGAAATGAAATTCAAGAGAACGCTTGAGAATT  
TTATAACAAGTAAGTATGTACTGGTGTAGTAGTCTTAACGTGTTACCAAGATAATACAA  
ATGCTTAAATAAAAAGACTTGATCTGATTAGACCAAATCTTTGATAGTGTATATTAAAT  
AACAAAATAAAAGGAGTCGCTCACGCCCTACCAAGTTGTGAACGACATCATTCAAAGAA  
AAAAACACTGAGTTGTTTATAATCTGTATATTAGATATTAAACGATATTAAATATAC  
ATCAAGATATATTGGGTGAGCGATTACTAAACGAAATTGAGATTAAGGAGTCGATT  
TTATGTATAAAACAATCATGCAAATCATTCAAATCATTGGAAATCACGATTAGACAAT  
TTTCTAAACCGGCTACTCTAATAGCCGGTGGACGCACATACTGTGTGCATATCTGATCC  
AAAATTAAGTTGATGCAATGACGATCGTGGAAATCTCAACCGAGACACGCTCAGGCC  
TTTCTAAATTATGAGTGTAGAGCCCCAAATAAGACTTGGATATTCTCAAACAAAGTT  
AAAGCTAAAGCACCTCAAGAAAAAGTTATATTGAATATGACAAAGTGAAGCAGATAGTTG  
GGATAGACGTAATATCGTATTGAATTAAATCCAAACAAACTACACGAGATGAAATGATT  
GGTTAAACAAATATAATAAGCTACATGGAAGATGACGGTTTACAAGATTAGATTAGCC  
TTGATTTGAAGATGATTGAGTGACTIONTGCATGTCTGATAAAGCAGTTAAGAAAAC  
TATTTTTATGGTCGAATGGTAAGCCAGAAACAAATTTGGCGTGAGAGATAGTAATA  
GATTATTAGAATTATAATAAAAGCAAGAACGTAAGATAATGCAGATGCTGAAGTTATG  
TCTGAACATTATGGCGTGTAGAAATCGAACTTAAAGAGATATGGTGGATTACTGGAATGA  
TTGCTTTAGTGTATTACATATCTTGCACCAACAGATTGGAAAATATCCAACGCAGTCGGATA  
GAGCAATAGTTTATGTTATTGAGTGTAGAAGAAGAATGGGAAAGCTTCACAGAAATTCT  
AGAACAAATATAAGAATTGATAAAAGAAATTGCGCAGTCGATTAAACGGACTTAATGAA  
ATCGACTTTAAAGCGAACGAAAAACAATTGCAAAACAAATCGATTTGGCAACATGAAT  
TTAAATTGGAAATAGTGTACATATTAATATTACTGAACAAAATGATATATTAAACTAT

TCTAATTAGGAGGATTTTTATGAAGTGTCTATTAAAAATTGGGAATTATGAGG  
TGAAAGAATAATTACCCCTATAAACTTAGCCACCTCAAGTAAAGAGGTAAAATTGTTAG  
TTTATATAAAAAATTAAAGGTTGTTTATAGCGTTTATTGGCTTGTATTCTTCAT  
TTTTAGTGTATTAAATGAAATGGTTAAATGTTCTTACCTGATATTGCAAATCATT  
AATACTACTCCTGGAATTACAAACTGGTAAACACTGCATATATGTTAATTTCGATAGG  
AACAGCAGTATATGAAAATTATCTGATTATATAAATATAAAAAATTGTTAATTGGTA  
TTAGTTGAGCTGCTTGGTTATTGATTGCTTTATTGGGCCACCTAGGCAAATATGCTC  
TTACGTGCTATTATTAAGTGACTATTAAAAGGAGTTAATAATATGCGGCAAGGTATTCT  
TAAATAACTGTCAATTGATAGCGGGAAACAAATAATTAGATGCTTTTAGGAGGGCTT  
AGTTTTGTACCCAGTTAAGAATACCTTATCATGTGATTCTAAAGTATCCAGAGAATAT  
CTGTATGCTTGTATACCTATGGTTATGCATAAAAATCCCAGTGATAAAAGTATTACT  
GGGATTTTATGCCCTTTGGTTTGAAATGGAGGAAATCACATGAAAATTATTAATATT  
GGAGTTTAGCTCATGTTGATGCAGGAAAAACTACCTAACAGAAAGCTTATTATATAACAG  
TGGAGCGATTACAGAATTAGGAAGCGTGGACAAAGGTACAACGGAGGACGGATAATACGCTT  
TAGAACGTCAGAGAGGAATTACAATTACAGACAGGAATAACCTTTAGTGGAAAATACG  
AAGGTGAACATCATAGACACGCCAGGACATATGGATTCTAGCAGAAGTATATCGTTCATT  
ATCAGTTTAGATGGGCAATTCTACTGATTCTGCAAAGATGGCGTACAAGCACAAACTC  
GTATATTATTCATGCACTTAGGAAATGGGATTCCCACAATCTTTTATCAATAAGATT  
GACCAAAATGGAATTGATTCAACGGTTATCAGGATATTAAAGAGAAACTTCTGCCGA  
AATTGTAATCAAACAGAAGGTAGAACTGTATCCTAATATGTGTGACGAACCTACCGAAT  
CTGAACAATGGGATACGGTAATAGAGGGAAACGATAACCTTTAGAGAAATATGTCCGGT  
AAATCATTAGAACGATTGGAACTCGAACAAGAGGAAGCATAAGATTCAAATTGTTCT  
GTTCCCTTTATCATGGAAGTGCAAAAGTAATATAGGGATTGATAACCTTATAGAAGTTA  
TTACTAATAAATTATTATTCATCAACACATCGAGGTCCGTCTGAACCTTGCAGAAATGTTTC  
AAAATTGAATACAAAAAGACAACGTCTGCATATACGCCTTATAGTGGAGTACT  
ACATTTACGAGATTGGTTAGAGTATCAGAAAAAGAAAAATAAAAGTTACAGAAATGTATA  
CTTCAATAATGGGAATTATGTAAGATTGATAGAGCTTATTCTGGAGAAATTGTTATTTG  
CAAAATGAGTTTGAAAGTTAAATAGTGTCTGGAGATAACAAACTATTGCCACAGAGAAA  
AAAGATTGAAAATCCGCACCCCTACTACAAACAACTGTTGAACCGAGTAAACCTGAACAGA  
GAGAAATGTTGCTGATGCCCTTGGAAATCTCAGATAGTGTACCGCTTACGATATTAC  
GTGGATTCTACGACACATGAAATTATACTTCTTCTAGGAAAGTACAATGGAAGTGAT  
TAGTGCACGTGCAAGAAAAGTATCATGTGGAGATAGAACTAAAAGAGCCTACAGTCATT  
ATATGGAGAGACCGTTAAAAATGCAGAATATACCATTACATCGAAGTGCCTACAGTATGA  
TTCTGGCTTCCATTGGTTATCTGTATCACCGCTCCGGTGGAAAGTGGAAATGCAGTATGA  
GAGCTCGGTTCTCTGGATACTTAAATCAATCATTCAAAATGCAGTTATGGAAGGGTAC  
GCTATGGTTGCGAACAGGATTATATGGTTGGAATGTGACGGATTGAAAATCTGTTAAG  
TACGGTTATACTATAGCCCTGTTAGTACTCCAGCAGATTTCGGATGCTTACTCCTATTGT  
ACTGGAGCAAGCCTTAAAGCTGGAACAGAATTGTTAGAGCCATATCTTAGTTAAAG  
TTTATGCACCACAGGAATATCTTCNCGGCATATAACGATGCTCCAAATATTGTGCAAAT

ATCGTAAATACTCAACTGAAAAATAATGAGGT CATTATTATTGGAGAAATTCTGCTCGATG  
 TATTCAAGATTATCGCAATGATTTAACTTTTTACAAATGGGCTTAGTGTGTTAGCAG  
 AGCTAAAAGGATATCAGGTTACCACTGGCGAACCTGTTGCCAGACCCGTCGTCTAAATAGT  
 CGGATAGATAAAGTAAGATATATGTTCAATAAAACTTAGTGCCTTATGTTGTTATAT  
 AAATATGGTTCTTATTAAATAAGATGAAATATTCTTAATATAGATTGAATTAAAGTGG  
 AAGGAGGAGATTGTTATTATAAACTACAAGTGGATATTGTGCCTATTGTGAAATAAAC  
 AAGACTACGAATACGAGTGGACTACTACTTAAACCTACAGCCCCAAATGTA  
 AGAACGAAACTTAATTATGTTCAAAAAATGAATATAACAATCAAAGAGCCAGACGCC  
 AAGACGAGGCCGATAATTGAGAAATGAAACTCTCATCTTATCGGCTTTTGTATC  
 TGAATTTACTGACTAGCCTCAATATTCC

SEQ ID NO: 6

P1 pacA gene, lower case letters signify deleted pac-site sequence

GTGACCTGGGACGATCACAAGAAGAATTTGCTCGCTGGCGCGAGATGGTGGTTACACCAT  
 CGCACAGTATGCCGCCAGTTAATCTAACCTAATACCGCACGTCGTTATCTCCGTGCCT  
 TCAAAGAAGACACCAAGGACTACGGACAGCCGAAGCCAAATAAGCCAGTCAGGAAGGccacta  
 aaaagcatgatcattgatcactctaattatgatcaacatgcaggtgatcacattgcggctgaaat  
 agcggaaaaacaaagagttaatgccgtgtcagtgccgcagtcgagaatgcgaagcgccaaa  
 ataagcgcataaatgatcggtcagatgatcatgacgtgatcacccgcGCCCACCGGACCTTA  
 CGTGTGATCGCCTGGAACCGACACCCCTGGATGATGATGGTGAACGCTTGAATTGAAGTTGG  
 CGATTACCTGATAGATAACGTTGAAGCGCGGAAGGCCGCGCGCTATGTTGCGTCGGTCCG  
 GGGCGATGTTCTGGAAACCACTCTCTGGAAAAGTCTCTTCTATCTCCTTATGCTGGAG  
 AACGCCAGGGATACGTGTATTGCCCTGGTGCAGGAAATGCGCGATCAGCAAAAGACGATGA  
 TGAAGGTACTCCGCTGAATACCGTATCGCGAGCATGCTAACAGCTGTTCCGCGCAGATAA  
 GCAGCCTGATCAACACCATTACAGCATCCGAATAACTATCGAAAAGAAAGCCGGAGGCG  
 GAAAAGCACGCTTATCTATGGGCAAGCTGGCATTGTTAAGCTGGCATACGAACGAAAGCG  
 TGAAAATAACTGGTCAGTGCTGGAAGCGGCTGAATTCATCGAGGCGCATGGAGGAAAAGTGC  
 CGCCCCCTGATGCTGGAGCAAATCAAAGCCGATCTGCGTGCCTAAGACCAATACCGATGAT  
 GAGGAAAACCAACAGCATCTGGCGCTCCATCACTTGAAGATCTGGATAAAATCGCGCAGA  
 ACGGGCCGCCAGCCGCGCTGATGCCGATTGTGGATTGAGCATCGTAGAGAAGAAATTG  
 CCGATATCGTCGATACAGGTGGTTATGGTGTGCGATGCCGAAGGCATATCAAACGAAGCA  
 TGGCTTGAACAGGATCTGGACGAAGACGAGGAGGAAGACGAAGAAGTTACCCGCAAACGTGTA  
 CGGGGATGATGATTAA

SEQ ID NO: 7

Native P1 pacA gene

GTGACCTGGACGATCACAGAAGAATTTGCTGCCTGGCGAGATGGTGGTTACACCAT  
CGCACAGTATGCCGCCGAGTTAACCTAACCCTAACCGCACGTCGTTATCTCCGTGCCT  
TCAAAGAACGACACCAGGACTACGGACAGCCGAAGCCAAATAAGCCAGTCAGGAAGCCACTA  
AAAAGCATGATCATTGATCACTCTAACATGATCACATGCAGGTGATCACATTGCGGCTGAAAT  
AGCGGAAAAACAAAGAGTTAACGCCGTTGTCAGTGCCGCAGTCGAGAATGCGAACGCCAAA  
ATAAGCGCATAATGATCGTTAGATGATCATGACGTGATCACCCGCGCCACCGGACCTTA  
CGTGATCGCCTGGAACGCGACACCCCTGGATGATGATGGTGAACGCTTGAATTGAAAGTGG  
CGATTACCTGATAAGATAACGTTGAAGCGCGGAAGGCCGCGCGCTATGTTGCGTCGGTCCG  
GGGCCGATGTTCTGGAAACCACTCTCTGGAAAAGTCTCTTCTCATCTCCTTATGCTGGAG  
AACGCCAGGGATACGTGTATTGCCTGGTGCAGGAAATGCGCGATCAGCAAAAGACGATGA  
TGAAGGTACTCCGCCTGAATACCGTATCGCGAGCATGCTAACAGCTGTTCCGCGCAGATAA  
GCAGCCTGATCAACACCATTACAGCATCCGAATAACTATCGAAAAGAAAGCCGGAGGCG  
GAAAAGCAGCTTATCTATGGGGCAAGCTGGCATTGTTAAGCTGGCATACGAACGAAAGCG  
TGAAAATAACTGGTCAGTGCTGGAAGCGGCTGAATTCATCGAGGCGCATGGAGGAAAAGTGC  
CGCCCTGATGCTGGAGCAAATCAAAGCCGATCTGCGTGCTCTAACGACCAATACCGATGAT  
GAGGAAAACAAACAGCATCTGGCGCTCCATCACTTGAAGATCTGGATAAAATCGCGCAGA  
ACGGGCCGCCAGCCGCCGCTGATGCCGCATTGTGGATTGAGCATCGTAGAGAAGAAATTG  
CCGATATCGTCGATACAGGTGGTTATGGTGAATGTCGATGCGGAAGGCATATCAAACGAAGCA  
TGGCTTGAACAGGATCTGGACGAAGACGAGGAGGAAGACGAAGAAGTTACCCGCAAACGTAA  
CGGGGATGATGATTAA

## SEQ ID NO:8

terS gene, lower case characters signify deleted sequence

ATGAACGAAAAACAAAAGAGATT CGCAGATGAATATATAATGAATGGATGTAATGGTAAAAA  
 AGCAGCAATTCAGCAggttatagtaagaaaacagcagagtcttagcaagtcgattgttaa  
 gaaatgttaatgttcggaatatattaaagaacgattagaacacagatacaagaagagcgtta  
 atgagcattacagaagcttagcgttatctgctctattgcttagaggagaacctcaagaggc  
 ttacagtaagaaatatgaccattaaacgatgaagtggaaaaagaggttacttacacaatca  
 caccactttgaagagcgtcagagatctattgaccacataactaaaagttcatggtgcgtat  
 atcgacaaaaaagaattactcagaagaatattgagattaatattAGATCTATTGACCACAT  
 ACTAAAAGTTCATGGTGCCTATATCGACAAAAAGAAATTACTCAGAAGAATATTGAGATTA  
 ATATTGGTGAGTACGATGACGAAAGTTAA

## SEQ ID NO:9

Sequence containing native terS gene

AATTGGCAGTAAAGTGGCAGTTTGATACCTAAAATGAGATATTATGATAGTAGGATAT  
 TGACTATCTTACTCGCTTCCCTTATCGCAATTAGGAATAAAGGATCTATGTGGGTTGGCTG  
 ATTATAGCCAATCCTTTTAATTAAAAGCGTATAGCGCGAGAGTTGGTGGTAAATGAA  
 ATGAACGAAAAACAAAAGAGATT CGCAGATGAATATATAATGAATGGATGTAATGGTAAAAA  
 AGCAGCAATTCAGCAGGTTATAGTAAGAAAACAGCAGAGTCTTAGCAAGTCGATTGTTAA  
 GAAATGTTAATGTTCGGAATATATTAAAGAACGATTAGAACAGATAACAAGAAGAGCGTTA  
 ATGAGCATTACAGAAGCTTAGCGTTATCTGCTTCTATTGCTAGAGGAGAACCTCAAGAGGC  
 TTACAGTAAGAAATATGACCATTAAACGATGAAGTGGAAAAAGAGGTTACTTACACAATCA  
 CACCAACTTTGAAGAGCGTCAGAGATCTATTGACCACATACTAAAAGTTCATGGTGCCTAT  
 ATCGACAAAAAAGAAATTACTCAGAAGAATATTGAGATTAATATTGGTGAGTACGATGACGA  
 AAGTTAAATTAAACTTAAACAAACCCTAATGTTCAACAG

## SEQ ID NO:10

SaPibov2 integrase gene

TCATAAATATTAACTATTCTTCTGTGACTAGGGTACAAATGACCGTATCGTTATATA  
 CTTCATTACTATCAGCATGGCCTAAACGCTGTGCTATTACCATGATACTTGCCTGATTG  
 ACTAGCATAGACGCATGGCTATGTCTTAACTCATGAATTACAATTCTAGGAAATGTCTGACC  
 GTCTGGTAGTTGTCATCTAATACTTTAATGCAGCGGTAAACCAACGATCTATAGTTGATT  
 CACTATAAGCTTGAAGAATGTACCGAATAACATAATCATCTTATATACATTGTTCT

TTGTACCATTTAAATATTCTTGATATCATTGATCATGTGAAACAGGTAAGTATATACAG  
TATTGCTGCTTTGTTAGGGCTGTCACTCACCGTGATAGTCTGTTGTTAATATGTA  
TGAAATCATCATCATAGTTAATATCACGCCATGTGAGGGCTCTAATTCGCCCTACGTGCA  
CCAGAGTAAAACAGTAGCTTAAAGAATAACTTTGTTGTTAGCTAAAGCCTCATAGAA  
TTGATTGAATTGTTCTAATGTCCAATAGTTCAAACGCTTATTGATTCTATTCAAAGTTAC  
CTACTAGAGAGGCTACATTTGCTTAGATCATGAAACCTCATAGCATGGTTAAGTAACGAT  
ACTAAGAACACGTGCATTTCTTAGGTACTCTCCAGAGTGTCCCTCTTTAACTTCGTATT  
CTGAAACTTCATAATATCTTGTGTAGTCATATTAAACACGTCCATAGACTAAATAGGGTA  
GCAAATGGTTGTTGTATGTCCTTAATGCTTCACACTAGATGACTTACGACGTGCAGAA  
TACCACTCTATATACTCATCTACGAGCTTACAAAGGGCAGTTGTTATCTGCTCCTACACC  
CTCTAACTCGTCCATAATTCATTACATTCTCAATGCCTCTTACGCTGTTAAAGCCAC  
TCTTTTTATTCCTTACGTTGATTAAATTATCATAGTATTATACGAAAATAGTATGTA  
CCACGTTAGCGTCTTATATATGTTGGGATAGGTTAAGTTGTTCTATGGGAATCAC

SEQ ID NO:11

pGWP10001 full sequence

CTCGGGCCGTCTTGGGCTTGTACGGCCTTGCATCTCACGCGCTCCTGCGGCGGCC  
TGTAGGGCAGGCTCATACCCCTGCCGAACCGCTTGTCAAGCCGGTGGCCACGGCTCCGG  
CGTCTCAACGCGCTTGTAGATTCCCAGCTTTCGGCCAATCCCTGCGGTGCATAGGCGCGTG  
GCTCGACCGCTTGCAGGGCTGATGGTGACGTGGCCACTGGTGGCCGCTCCAGGGCTCGTAG  
AACGCCTGAATGCGCGTGTGACGTGCCTGCTGCCCTCGATGCCCGTTGCAGCCCTAGATC  
GGCCACAGCGGCCGAAACGTGGTCTGGTCGCGGTATCTGCGCTTGTGCCGATGAAC  
CCTTGGCCGACAGCCTGCCGTGCCTGCGTCAGCGGACCCACGAACGCGGTATGTGCGGGCTG  
GTTTCGTACGGTGGATGCTGGCCGTACGATGCGATCCGCCCGTACTTGTCCGCCAGCCA  
CTTGTGCGCCTCTCGAAGAACGCCCTGCTGTTCTGGCTGGCCGACTTCCACCATTCCG  
GGCTGGCCGTATGACGTACTCGACGCCAACACAGCGTCCTGCGCCGTTCTGGCAG  
AACTCGCGCAGTCGGCCCATCGCTTCATCGGTGCTGCTGGCCGCCAGTGCTCGTTCTGG  
CGTCCTGCTGGCGTCAGCGTTGGCGTCTCGCGTCGCGTAGGCGTGCTTGAGACTGGCCG  
CCACGTTGCCCATTTGCCAGCTTGCATCGATGATCGGTATGCCGATGCCCTGCC  
CCTCCCTTTGGTGTCCAACCGGCTCGACGGGGCAGCGCAAGGCGGTGCCTCCGGCGGCC  
ACTCAATGCTTGAGTATACTCACTAGACTTGCCTCGCAAAGTCGTGACCGCCTACGGCGGC  
TGCAGGCCCTACGGCTTGCCTCCGGCTTCGCCCTGCGCGTCGCTGCCCTTGCC  
AGCCCGTGGATATGTGGACGATGGCGCGAGCGGCCACCGGCTGGCTCGCTCGCTGGCC  
GTGGACAACCGTGGCTACCACTATAGGGCGAATTGGCGGAAGGCCGTCAAGGCCGATTGG  
GCCCGCGCGCCGGATCCGCTAGCTCTAGACCTCTAGACCGAGGACAGAAATGCCCTGA  
CTTCGCTGCTGCCAAGGTTGCCGGTGACGCACACCGTGGAAACGGATGAAGGCACGAACC  
CAGTGGACATAAGCCTGTTCGTTCTGAAGCTGTAATGCAAGTAGCGTGCCTCCGGGCC  
TTTGACATGTGACTTCGTTACCCCTCGCTCAAAAGAGTTTACGAAAGGAAGCATAAG  
TGACCTGGGACGATCACAAGAAGAATTTGCTCGCTGGCGAGATGGTGGTTACACCAC  
GCACAGTATGCCGCCGAGTTAATCTAACCTAATACCGCACGTCGTTATCTCCGTGCC  
CAAAGAACGACACCAGGACTACGGACAGCCGAAGCCAAATAAGCCAGTCAGGAAGCCACTAA  
AAAGCATGATCATTGATCACTCTAAATGATCAACATGCAGGTGATCACATTGCCGCTGAAATA  
GCGGAAAAACAAAGAGTTAATGCCGTTGTCAGTGCAGTCAGGAAATGCGAAGGCCAAAA  
TAAGCGCATAAATGATCGTTAGATGATCATGACGTGATCACCCCGGCCACCGGACCTAC  
GTGATCGCCTGGAACCGACACCCCTGGATGATGATGGTAACGCTTGAATTGAAAGTTGGC  
GATTACCTGATAGATAACGTTGAAGCGCGGAAGGCCGCGCGCTATGTTGCGTCGGTCCGG  
GGCCGATGTTCTGGAAACCACTCTTCTGGAAAAGTCTCTTCTATCTCCTATGCTGGAGA  
ACGCCAGGGATACGTGTATTGCCCTGGTGCAGGAAATGCGCGATCAGAAAAAGACGATGAT  
GAAGGTACTCCGCCTGAATACCGTATCGCGAGCATGCTAACACAGCTGTTCCGCGCAGATAAG  
CAGCCTGATCAACACCATTACAGCATCCGGAATAACTATCGAAAAGAAAGCCGGGAGGCGG

AAAAGCACGCTTATCTATGGGGCAAGCTGGCATTGTAAGCTGGCATACGAACGAAAGCGT  
GAAAATAACTGGTCAGTGCTGGAAGCGGCTGAATTCATCGAGGCGATGGAGGAAAAGTGC  
GCCCTGATGCTGGAGCAAATCAAAGCCGATCTCGTGCTCCTAACGACCAATACCGATGATG  
AGGAAAACCAAACAGCATCTGGCGCTCCATCACTGAAAGATCTGGATAAAATCGCGAGAA  
CGGGCCGCCAGCCGCCGCTGATGCCGATTGTGGATTGAGCATCGTAGAGAAGAAATTGC  
CGATATCGTCGATACAGGTGGTTATGGTATGTCATGCCGAAGGCATATCAAACGAAGCAT  
GGCTGAAACAGGATCTGGACGAAGACGAGGGAGGAAGACGAAGAAGTTACCCGCAAACGTAC  
GGGGATGATGATTAATTAAAAACCGCCCCGGCGGGTTTTTATCTAGAGCTAGCGGAT  
CCGGCGGCCGGGCCCTCTGGCCTCATGGCCTTCGCTACTGCCGTTCCAGCACT  
ATAGGGCGAATTGGCGGAAGGCCGTCAAGGCCGATTGGGCCGGCGCCGGATCCGCTA  
GCTCTAGACTGGCAGGTTCTGAGCAGATCGTCAAACCGATCTGGATCGGTAGAAAAAT  
TTGCTCTAATAAAATTCTGTTCTAAGTCAAAGAACATTCAATGCCGATATGAATATATCAAAC  
CTTCATCAAATGTCGATCCTCAACCACTCTGCCGTTATTGTGGTGTGAAATTACGAC  
CGACCGCGCTGGCGTTACAACCTTAATGCTCTACACAGAGCGAGCGGACTCGGTGCCATA  
AAGGCCAGCTCAATGGCTAAGAACGCTGTCAGCTAACAGCTCATCGAAGAGCTTGAAAAA  
GAAACTATGCAGAATTGCATAGTTCTACAAGCAATGGAAGCAGGATTCTTCACGAC  
TCGTATAACCGCAAAGGTCAAGCAGTCAGCTGCTGATGAAGCGATTGCTGATGGTGTGCTGG  
TACCTGTCGGCAACGCGCTAACAGACGTAGTAAGAACCAACCAGCATTGTAATGCTGGCTA  
AAAGTCACTTCTGAGCTGTATAACGATGAGCGATTACTTTCTGGCTATGAATTGCC  
TGCTTGTAAACACACTCCGGCTATCCGTAGCGCCGGCATATCCTGCGCAATGTGAAA  
TCTCGGGCAACAACCAGTGAATACTCATTACAAGCCTCACCGCCTGATCGGGCAGAAA  
CTGGTTAGCCAATCAACCGCTCGCATCCGTGAAGCTGTAACAAAGGAATTCTG  
TCTGTAGAGATTGTTATCGCGATCACCCTGAGCTGCTGCAACGCTAACCTGTACCGGTTACACC  
ATCCTTTGGCCTCGACAACAAGCCAAAATGCGCTGATAGAAAGCAAATTAAAGATCT  
CTTCAGCGGCAACCAAGGTTAAAGCTGTTCTGCTAACGACATTGGCTTATTAAATTGTTA  
TCCACACCCCCATGTCAAAATGATAACCCCTCCCCCTGTCAGGATGACGTGGCAATAAGAA  
TAAGAAGTCACAAGTTAAAAAAACAAAAAGATCAGTTCCGGCGGTGCCGAACAACCAGCC  
TCAAAAAATTGACTCATGGATCGCTAACGGAAAAGCAAAGGCTGACAATCTCGGTTATCC  
AAAAAACGCACTAAAAACATGAGTTCAAGCAGAAAGTAGAGGGCGCTCGCGGAAATATGC  
TTACCTGAAGAACAAAGCGTTCGCCTGATATTGGCGGGATATCAAACCTCGATAACCTACCGC  
ATTGCATGACGGTAAACGAAGCTCTTAATGCGGTTTAGCCAAAATAAGATAACGAACAA  
TGGGGTACCGGCAGGATTCAAGAGGTAATGAATTGCTCTAACCATGCATACTTT  
CAACACCTCTAGTTGCCATGAGGCAAACCTCATAGGTGTCCTGGTAAGAGGACACTGTTGCC  
AAAACGGACGCCCAATTATGCAATTAAACAACTAACGGACAATTCTACCTAACATA  
AGTGGCTAAAAAAACCGCCCCGGCGGGTTTTTATCTAGAGCTAGCGGATCCGGCGCG  
CGGGCCCTCTGGCCTCATGGCCTCCGCTCACTGCCGCTTCCAGCCAGCCTCGACC  
ACATACCCACCGGCTCCAACGCGCCCTGCCCTGCCCCATCAATTTTTAATTTC  
TCTGGGGAAAAGCCTCCGGCTGCGCCTGCGCCTGCGCTTGGACACCAAGTGG

AAGGCGGGTCAAGGCTCGCGCAGCGACCGCGCAGCGCTTGGCCTTGACGCGCTGGAACGA  
CCCAAGCCTATGCGAGTGGGGCAGTCGAAGGCGAAGCCGCCCTGCCCCCGAGACCT  
GCAGGGGGGGGGGGCGCTGAGGTCTGCCTCGTAAGAAGGTGTTGCTGACTCATACCAGGC  
CTGAATGCCCATCATCCAGCCAGAAAGTGAGGGAGCCACGGTGTGAGAGCTTGTG  
AGGTGGACCAGTTGGTGAACCTTGCTTGCCACGGAACGGTCTGCGTTGTCGGGA  
AGATGCGTGATCTGATCCTCAACTCAGCAAAAGTCGATTATTCAACAAAGCCCGTCC  
CGTCAAGTCAGCGTAATGCTCTGCCAGTGTACAACCAATTAAACCAATTGATTAGAAAAA  
CTCATCGAGCATCAAATGAAACTGCAATTATTATCATATCAGGATTATCAATACCATATTTT  
GAAAAAGCCGTTCTGTAATGAAGGAGAAAACTCACCGAGGCAGTCCATAGGATGGCAAGA  
TCCTGGTATCGGTCTGCGATTCCGACTCGTCCAACATCAATACAACCTATTAATTCCCCTC  
GTCAAAAATAAGTTATCAAGTGAGAAATCACCATGAGTGACCGACTGAATCCGGTGAGAATG  
GCAAAAGCTTATGCATTCTTCCAGACTTGTCAACAGGCCAGCCATTACGCTCGTCATCA  
AAATCACTCGCATCAACCAAACCGTTATTCAATTGTGATTGCCCTGAGCGAGACGAAATAC  
GCGATCGCTGTTAAAGGACAATTACAAACAGGAATCGAATGCAACCGGCGCAGGAACACTG  
CCAGCGCATCAACAATATTTCACCTGAATCAGGATATTCTCTAATACCTGGAATGCTGTT  
TTCCCGGGATCGCAGTGGTGAACCATGCATCATCAGGAGTACGGATAAAATGCTTGAT  
GGTCGGAAGAGGCATAAATTCCGTAGCCAGTTAGTCTGACCATCTCATCTGTAACATCAT  
TGGCAACGCTACCTTGCCATGTTCAAGAACAACTCTGGCGATCGGGCTTCCCATACAAT  
CGATAGATTGTCGCACCTGATTGCCGACATTATCGCGAGGCCATTATACCCATATAAAC  
AGCATCCATGTTGGAATTAAATCGGGCCTCGAGCAAGACGTTCCGTTGAATATGGCTCA  
TAACACCCCTGTATTACTGTTATGTAAGCAGACAGTTATTGTTCATGATGATATATTT  
TTATCTTGCAATGTAACATCAGAGATTGAGACACAACGTGGCTTCCCCCCCCCCT  
GCAGGTCCGAGCCTCACGGCGCGAGTGCAGGGGTTCCAAGGGGAGCGCCACCTGGC  
AAGGCCGAAGGCCGCGCAGTCATCAACAAGCCCCGGAGGGCCACTTTGCCGGAGGGGG  
AGCCGCGCGAAGCGTGGGGAACCCCGCAGGGTCCCTTGGCACCAAAGAACTA  
GATATAGGGCGAAATGCGAAAGACTTAAACAAACTTAAAGGGGGTACGCAACA  
GCTCATTGCGGCACCCCCCGCAATAGCTCATTGCGTAGGTTAAAGAAAATCTGTAATTGACT  
GCCACTTTACGCAACGATAATTGTTGCGCCTGCCGAAAGTTGCGAGCTGATTGCGCAT  
GGTGCCGCAACCGTGGCACCCCTACCGCATGGAGATAAGCATGCCACGCAGCAGAGA  
AATCGGCATTCAAGCCAAGAACAAAGCCGGTCACTGGTGCAAACGGAACGCAAAGCGCATG  
AGGCGTGGGCCGGCTTATTGCGAGGAAACCCACGGCGCAATGCTGCTGCATCACCTCG  
GCGCAGATGGGCCACCAGAACGCCGTGGTGGTCAGCCAGAAAGACACTTCCAAGCTCATCGG  
ACGTTCTTGCGGACGGTCCAATACGCACTGAGACTTGGTGGCCAGCGCTGGATCTCCG  
TCGTGAAGCTCAACGGCCCCGGCACCGTGTGGCCTACGTGGTCAATGACCGCGTGGCGTGG  
GGCCAGCCCCGCGACCAGTTGCGCCTGCGGTGTTCAAGTGCAGGGTGGTGGTGTGATCACGA  
CGACCAGGACGAATCGCTGTTGGGCATGGCGACCTGCGCCGATCCGACCCCTGTATCCGG  
GCGAGCAGCAACTACCGACCGCCCCGGCGAGGAGCCGCCAGCCAGCCGGCATTCCGG  
ATGGAACCAGACCTGCCAGCCTGACCGAAACGGAGGAATGGGAACGGCGCGGGAGCAGCG  
CCTGCCGATGCCGATGAGCCGTGTTCTGGACGATGGCGAGCCGTTGGAGCCGCCGACAC

GGGTACGCTGCCCGCCGGTAGCACTGGGTTGCGCAGCAACCGTAAGTGCCTGTTCCA  
GAATCGGCTGTAGCCGCCTGCCGCCCTACCTGTCTGCCCTCCCCGCGTGCCTCGCG  
GTGCATGGAGCCGGGCACCTGACCTGAATGGAAGCCGGCAGCCTCGCTAACGGATTCA  
CCGTTTATCAGGCTCTGGGAGGCAGAATAATGATCATCGTCAATTATTACCTCCACG  
GGGAGAGCCTGAGCAAACCTGGCCTCAGGCATTGAGAAGCACACGGTACACTGCTCCGGT  
AGTCAATAAACCGTAAACCAGCAATAGACATAAGCGGCTATTAAACGACCCTGCCCTGAAC  
CGACGACCGGGTCGAATTGCTTCGAATTCTGCCATTCATCCGTTATTATACTTATTCA  
GGCGTAGCACCAGGCCTTAAGGGCACCAATAACTGCCTTAAAAAAATTACGCCCGCCCTG  
CCACTCATCGCACTCGTCAGGTGGCACTTTCGGGAAATGTGCGCGAACCCCTATTGTT  
TATTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCTGATAATGCTT  
CAATAATATTGAAAAGGAAGAGTATGAAGTTGGAAATATTGTTTCGTATCAACCACC  
AGGTGAAACTCATAAGCTAAGTAATGGATCGCTTGGCTGGTATCGCCTCAGAAGAG  
TAGGGTTGATACATATTGGACCTAGAACATCATTACAGAGTTGGCTTACGGGAAAT  
TTATTGTTGCTGCCGCTAACCTGTTAGGAAGAACTAAAACATTAAATGTTGGCACTATGGG  
GGTTGTTATTCCGACAGCACACCCAGTCGACAGTTAGAACAGCTTTATTAGATCAA  
TGTGAAAGGTCGTTAATTGGAACCGTTGAGGGCTATACCATAAGATTTCGAGTA  
TTGGTGTGATATGGAAGAGTCTCGAGCAATTACTCAAATTCCTAACGGTTGATG  
TATATCCAAAGTGTACTCAAATGTACCAACCTGTATGACTGCTGAGTCCGCAAGTACG  
ACAGAACGGCTAGCAATAAGGGCTACCAATGGTTCTAGTGGATTATTGGTACTAATGA  
AAAAAAAGCACAGATGGAACCTATAATGAAATTGCGACAGAATATGGTCATGATATATCTA  
AAATAGATCATTGTATGACTTATATTGTTCTGTTGATGATGATGCACAAAGGCGCAAGAT  
GTTTGTGGAGTTCTGAAAATTGGTATGACTCATATGAAATGCGACCAATATCTTAA  
TGATAGCAATCAAACCTCGTGGTTATGATTATCATAAAGGTCAATGGCGTATTGTTTAC  
AAGGACATACAAACACCAATCGACGTGGATTATAGCAATGGTATTAAACCTGTAGGCACT  
CCTGAGCAGTGTATTGAAATCATTCAACGTGATATTGATGCAACGGTATTACAAACATTAC  
ATGCGGATTGAAAGCTAATGGAACGTGAAAGATGAAATAATTGCTTCCATGCGACGCTTATGA  
CACAAGTCGCTCCTTCTAAAAGAACCTAAATAATTACTTATTGATACTAGAGATAATA  
AGGAACAAGTTATGAAATTGGATTATTTCTAAACTTCAGAAAGATGGAATAACATCT  
GANGAAACGTTGGATAATATGGTAAAGACTGTCACGTTAATTGATTCAACTAAATATCATT  
TAATACTGCCTTGTATGAAACATCACTTTCAAAATGGTATTGTTGGAGCACCTATTA  
CCGCAGCTGGTTTATTAGGGTTAACAAATAATTACATATTGTTCTAACATCAAGTA  
ATTACCACCCATCACCTGTACGTGAGCAGAAGAACCCAGTTATTAGATCAAATGTCAGA  
GGGACGCTTCATTCTGGTTAGTGACTGCGAAAGTGATTGAAATGGAATTAGAC  
GTCATATCTCATCAAGGCAACAACAATTGAAAGCATGCTATGAAATAATTGACGCATTA  
ACTACAGGTTATTGTATCCCCAAAACGACTTTATGATTTCAGAAAGGTTCAATTAAATCC  
ACACTGTTACAGTGAGAATGGACCTAACGAAATATGTATCCGCTACATCAAAGAAGTCGTCA  
TGTGGCAGCGAAAAGGCAGCTGCCTTAACATTAAAGTGGAGGATAATTAGAAACCAA  
GAACGCTATGCAATTCTATATAATAACAGCACAACAATGAGTATTGATATTGGATGT

TGATCATCAATTAACGTAAATTGCGAACTTAAATGCTGATAGAAGTACGGCTCAAGAAGAAG  
 TGAGAGAATACCTAAAAGACTATATCACTGAAACTTACCCCAAATGGACAGAGATGAAAAAA  
 ATTAACACTGCATTATTGAAGAGAATGCAGTTGGGTCTCATGATGACTATTATGAATCGACAAA  
 ATTAGCAGTGGAAAAAACAGGGTCTAAAATATTATTATCCTTGAATCAATGTCCGATA  
 TTAAAGATGTAAAAGATATTATTGATATGTTGAACCAAAAATCGAAATGAATTACCATAA  
 AGTAGTACTGTTGTAATTCAATTAAAGCATTCTGCCGACATGGAAGCCATCACAGACGGCATGA  
 TGAACCTGAATGCCAGCGGCATCAGCACCTGTCGCCCTGCGTATAATATTGCCATGGT  
 GAAAACGGGGCGAAGAAGTTGCCATTGGCACGTTAAATCAAACACTGGTGAAACTCA  
 CCCAGGGATTGGCTGAGACGAAAAACATATTCTCAATAAACCTTAGGGAAATAGGCCAGG  
 TTTTCAACCGTAACACGCCACATCTTGCATGAAATATGTGTAGAAACTGCCGAAATCGTCGTG  
 GTATTCACTCCAGAGCGATGAAAACGTTAGTTGCTCATGGAAAACGGTGTAAACAGGGT  
 GAACACTATCCCATATCACCAGCTACCGTCTTCATTGCCATACGGAATTCCGGATGAGCA  
 TTCATCAGGCGGGCAAGAATGTGAATAAAGGCCGGATAAAACTTGTGCTTATTTCATTAC  
 GGTCTTAAAAAGGCCGTAAATATCCAGCTGAACGGTCTGGTTAGGTACATTGAGCAACTG  
 ACTGAAATGCCTCAAAATGTTCTTACGATGCCATTGGATATATCACCGTGGTATATCCA  
 GTGATTTTTCTCCATTAGCTTCCTAGCTCCTGAAAATCTGATAACTCAAAAAATAC  
 GCCCGGTAGTGTATCTTACGGTAAAGTTGGAACCTCTTACGTGCCGATCAACGT  
 CTCATTTGCCAAAAGTTGGCCAGGGCTTCCCGGTATCACAGGGACACCAGGATTATT  
 TATTCTGCGAAGTGTATCTCCGTACAGGTATTATTGAAGACGAAAGGGCCTCGTGTAC  
 GCCTATTTTATAGGTTAATGTATGATAATAATGGTTCTTAGACGTCAGGTGGCACTTT  
 CGGGAAATGTGCGCGCCCGTCTGCTGGCGCTGGCCTGTTCTGGCGCTGGACTTCC  
 CGCTGTTCCGTACAGCAGCTTCCGCCACGGCCTGATGATCGCGCGCCTGGCCTGCAT  
 ATCCCGATTCAACGGCCCCAGGGCGTCCAGAACGGCTTCAGGCGCTCCGAAGGT

SEQ ID NO:12

S. aureus P<sub>clpB</sub> Promoter Sequence

GTCTAGTTAATGTGTAACGTAACATTAGCTAGATTTTTATTCAAAAAAATATTACAAAT  
 ATTAGGAAATTAAAGTGTAAAAGAGTTGATAATGATTATATTGGACTATAATATAATTAA  
 GGTC

SEQ ID NO:13

RN10616 genomic sequence loci showing the  $\varphi 80\alpha$  terS deletion and complementation.

terS=Bracketed Text, Deletion=Underlined, Complement=Bold

ATTAGACAACAAACAAGTCATTGAAAATTCCGACTTATTATTCAAAAAGAAATTGATAGCG  
 CAGATATAACAAGCTAGGTTAAAAGTAGGCGATAAGGTAGAAGTTAAAACAATCGGTTATAGA

ATACACTTTAAATTATACCGGTCTTACGAAGTAAAGAAGGTAGATAAACAAATGATT  
 AAACAAATACTAAGACTATTATTCTACTAGCAATGTATGAGTTAGGTAAGTATGTAACTGA  
 GCAAGTATATATTATGATGACGGCTAATGATGTAGAGGTGCCGAGTGACTTCGCGAAGT  
 TGAGCGATCAGTCAGATTGATGAGGGCGGAGGTGACGGAGTAGATGATGTGGTTAGTCATA  
 GCAATTATATTACTAGTCATCTTATTGTTGGTGTGATGTTGCAAGCTGAACAGTTAAAGG  
 CGATGTGAAAGTTAAAGAGCGGGAGATAGAGATATTAAGAAGTAGATTGAGACATTTGAAG  
 ATTAAAAATATTGTATGGAGGGTATTGACTAAAAAGAAATATGGATTAAATTATCAA  
 CAGTTCGAAAGTTAGAAGATGAGTTGTGATTATCCTAATTATCATAAGCAACTCGAAGAT  
 TTAAGAAGTGAATAATGACACCAGGATTCCAACAGATAACAAATATAGCGGGAGTTGT  
 ACCGTCTAATACATCGAAAACAGAAATGGCAGTAACTAATTATCTTGATGACAGAG  
 GTAAAATCCTTGAGTTAAGAGCGCTATTGAAACGTATAATCAACACATCAAGTAGGAAAGAA  
 CGCGAATTCAAGAGTATTATTAAATAAAAGGAATTAGTGAAAGTTGTGATGACAT  
 ACACATTCTGATAGAACTGCTCATAGAATCAAAAGAAAATCATATCTAGATTGGCGGAAG  
 AGTTAGGGGAAGAGTGAAATTGGCAGTAAAGTGGCAGTTTGATACCTAAAATGAGATATT  
**ATGATAGTGTAGGATATTGACTATCTTACTGCGTTCCCTATCGCAATTAGGAATAAAGGA**  
**TCTATGTGGTTGGCTGATTATGCCAACCTTTTAATTAAAAGCGTATAGCGCGAG**  
**AGTTGGTGGTAAATGAA**

[ ATGAACGAAAACAAAAGAGATTGCGAGATGAATATATAATGAATGGATGTAATGGTAA  
AAAGCAGCAATTTCAGCAGGTATAGTAAGAAAACAGCAGAGTCTTAGCAAGTCGATTGTT  
AAGAAATGTTAATGTTCGGAATATATTAAAGAACGATTAGAACAGATAACAAGAAGAGCGTT  
TAATGAGCATTACAGAAGCTTAGCGTTATCGCTCTATTGCTAGAGGAGAACCTCAAGAG  
GCTTACAGTAAGAAATATGACCATTAAACGATGAAGTGGAAAAGAGGTTACTTACACAAT  
CACACCAACTTTGAAGAGCGTCAGAGATCTATTGACCACATACTAAAAGTTCATGGTGCCT  
ATATCGACAAAAAGAAATTACTCAGAAGAATATTGAGATTAATATTGGTAGTACGATGAC  
GAAAGTTAA ]

**ATTAAACTTAACAAACCACATCTAATGTTCAACAGAAACATATTGAAATACTAACCAATT**  
 ACGATAACTCACTGAAGTACATTACGGTGGAGGTTCGAGTGTAACTCAGGGCGTTATA  
 CAAAAAGTTGTACTTAAAGCATTGCAAGACTGGAAATATCCTAGGCGTATACTATGGCTTAG  
 AAAAGTCCAATCAACAATTAAAGATAGTTATTGAAAGATGTCAAAGATTGTTGATAAAACT  
 TCGGTATTGGACATGTGCCTTGGATAAGACTGATAACAAAGTTGAATTGCCAACGGC  
 GCAGTTTTGTTAAAGGATTAGATAACCCAGAGAAAATAAGTCGATAAAAGGCATATC  
 AGACATAGTCATGGAAGAAGCGTCTGAATTCACACTAAATGATTACACGCAATTACGTTGC  
 GTTGAGGGAGCGTAAACACGTGAATAAGCAAATATTGATGTTAACCGAGTATCTAAA  
 CTGAATTGGTTATAAGTATTCTTGAAACATGGTAACCAATGGAAAATGTATGATTAG  
 ACAATCTAGTTATCGAGATAATAAGTTCTTGATGAAATGACACGACAAAATGAGTTGT  
 TAGCAAATCGTAATCCAGCATATTACAAAATTATGCGTTAGGTGAATTCTACACTAGAC  
 AAATTGGTTCCCTAAGTATGAAAACGTTAATAAAAGATGAGTTAAGACACATTAC  
 TTCTTATTGAGTGGACTTGGCTACGTTAATGATCCTAGTGCTTTATACATTCTAAA  
 TAGATGTAAGAAAAAGAAGTTATACATCATTGAAGAGTATGTTAAACAAAGGTATGCTGAAT

GATGAAATAGCTAATGTCATAAAGCAACTGGTTATGCTAAAGAAGAAATTACAGCAGATAG  
TGCAGAACAAAAAGTATACTGAATTAAGGAATCTAGGGCTAAAAGGATTTACCAACCA  
AAAAAGGGAAAGGGCTCGGTTACAAGGGTTACAATTCTTAATGCAATTGAAATCATTGTT  
GATGAACGTTGTTCAAGACTATTGAAGAGTTGACAACATACACATGGCAAAAGGACAAAGA  
TACAGGTGAATATACCAATGAACCAGTAGATACATACAATCATTGTATCGATTGTTGCGTT  
ATTCACTGGAACGATT

SEQ ID NO:14

pGW80A0001 Full Sequence

GGCGCCATGGTTAAGGGCCCTTGCAGAAAGAGTTAGTAAGTTAACAGAACGAAACCAAAA  
CTAAATGGTTAGCAGGAAACTTAGATAAAAAAATGAATCCAGAATTATATTCAAGAACAGGA  
ACAGCAACAAGAACAAACAAAAGAATCAAAACGAGATAGAGGTATGCACCTATAGAACATGC  
ATTTATGCCGAGAAAACCTATTGGTTGGAATGGCTATGTGTTAGCTAACTGTTAGCGAGT  
TGGTTGGACTTGAATTGGGATTAATCCAAGAAAAGTACCAACTCAACAAACACATAAAGCCCT  
GTAGGTTCCGACCAATAAGGAAATTGGAATAAGCAATAAAGGAGTTGAAGAAATGAAATT  
CAGAGAACGCTTGAGAATTTTATAACAAGTAAGTATGTTAGCTTGTGTTAGTCTAA  
CTGTTACAGATAATACAATGCTAAATAAAAAAGACTTGATCTGATTAGACCAAATCT  
TTTGATAGTGTATATTAAATAACAAATAAAAGGAGTCGCTCACGCCCTACCAAAGTTGT  
GAACGACATCATTCAAAGAAAAACACTGAGTTGTTTATAATCTTGTATATTAGATAT  
TAAACGATATTAAATATACATCAAGATATATATTGGGTGAGCGATTACTAAACGAAATT  
GAGATTAAGGAGTCGATTTTATGTATAAAACAATCATGCAAATCATTCAAATCATTGG  
AAAATCACGATTTAGACAATTTCTAAACCGGCTACTCTAAATAGCCGGTGGACGCACAT  
ACTGTGTGCATATCTGATCCAAAATTAAGTTGATGCAATGACGATCGTTGGAAATCTCAA  
CCGAGACACGCTCAGGCCCTTCTAAATTATGAGTGTAGAGCCCCAAATAAGACTTGGG  
ATATTCTCAAACAAAGTTAAAGCTAAAGCAGCTTCAAGAAAAAGTTATATTGAATATGAC  
AAAGTGAAGCAGATAGTTGGGATAGACGTAATATGCGTATTGAATTAAATCCAAACAAACT  
TACACGAGATGAAATGATTGGTAAAACAAAATATAAGCTACATGGAAGATGACGGTT  
TTACAAGATTAGATTAGCCTTGATTTGAAGATGATTGAGTGACTACTATGCAATGTCT  
GATAAAGCAGTTAAGAAAATTTTATGGCGTAATGGTAAGCCAGAAACAAAATATT  
TGGCGTGAGAGATAGTAATAGATTATTAGAATTATAATAAAAGCAAGAACGTAAGATA  
ATGCAGATGCTGAAGTTATGTCTGAACATTATGGCGTGTAGAAATCGAACTAAAAGAGAT  
ATGGTGGATTACTGGAATGATTGCTTGTGATTACATATCTGCAACCAGATTGGAAAAC  
TATCCAACGCACTGCGGATAGAGCAATAGTTTATGTTATTGAGTGATGAAGAAGAATGGG  
GAAAGCTTCACAGAAATTCTAGAACAAAATATAAGAATTGATAAAAGAAATTCGCCAGTC  
GATTTAACGGACTTAATGAAATCGACTTAAAAGCGAACGAAAAACAATTGCAAAACAAAT  
CGATTGGCAACATGAATTAAATTTGGAAATAGTGTACATATTAATATTACTGAACAA  
AAATGATATATTAAACTATTCTAATTAGGAGGATTTTATGAAGTGTCTATTAAAAA

TTTGGGGAATTATGAGGTGAAAGAATAATTACCCATAAACTTAGCCACCTCAAGT  
AAAGAGGTAAAATTGTTAGTTATATAAAAATTAAAGGTTGTTTATAGCGTTTATT  
TTGGCTTGTATTCTTCATTTTAGTGTATTAAATGAAATGGTTAAATGTTCTTAC  
CTGATATTGCAAATCATTAAACTACTCCTGAAATTACAAACTGGTAAACACTGCATAT  
ATGTTAACTTTTCGATAGGAACAGCAGTATATGGAAAATTATCTGATTATATAAATATAA  
AAAATTGTTAATTATTGGTATTAGTTGAGCTGTCTGGTCATTGATTGCTTTATTGGC  
CCACCTAGGCAAATATGCTCTTACGTGCTATTATTAAGTGACTATTAAGGAGTTAATA  
AATATGCGGCAAGGTATTCTTAAATAACTGTCAATTGATAGCGGGAACAAATAATTAGAT  
GTCCTTTTAGGAGGGCTTAGTTTGTACCCAGTTAAGAATACCTTATCATGTGATT  
CTAAAGTATCCAGAGAATATCTGTATGCTTGTACCTATGGTTATGCATAAAAATCCCAG  
TGATAAAAGTATTATCACTGGGATTTATGCCCTTGGGTTTGAATGGAGGAAATC  
ACATGAAAATTATTAATATTGGAGTTTAGCTCATGTTGATGCAGGAAAAACTACCTAAC  
GAAAGCTTATTATATAACAGTGGAGCGATTACAGAATTAGGAAGCGTGGACAAAGGTACAAC  
GAGGACGGATAATACGCTTTAGAACGTAGAGAGGAATTACAATTAGACAGGAATAACCT  
CTTTCACTGGAAAATACGAAGGTGAACATCATAGACACGCCAGGACATATGGATTCTTA  
GCAGAAGTATATCGTCATTATCAGTTAGATGGCAATTCTACTGATTCTGCAAAAGA  
TGGCGTACAAGCACAAACTCGTATATTATTCATGCACTTAGAAAATGGGATTCCCACAA  
TCTTTTATCAATAAGATTGACAAAATGGAATTGATTCAACGGTTATCAGGATATT  
AAAGAGAAACTTCTGCCGAAATTGTAATCAAACAGAAGGTAGAACTGTATCCTAATATGTG  
TGTGACGAACCTTACGAATCTGAACAAATGGGATACGTAATAGAGGAAACGATAACCTT  
TAGAGAAATATATGTCCGGAAATCATTAGAACGTAGGAAACTCGAACAGAGGAAAGCATA  
AGATTCAGAATTGTTCTGTCCCTTTATCATGGAAGTGCAAAAGTAATATAGGGAT  
TGATAACCTTATAGAAGTTATTACTAATAAATTATTATCATCAACACATCGAGGTCCGTCTG  
AACTTTCGGAAATGTTCAAAATTGAATATACAAAAAGACAAACGTCTGCATATATA  
CGCCTTATAGTGGAGTACTACATTACGAGATTGGTTAGAGTATCAGAAAAGAAAAAT  
AAAAGTTACAGAAATGTATACTCAATAATGGTAATTATGTAAGATTGATAGAGCTTATT  
CTGGAGAAATTGTTATTGCAAAATGAGTTTGAAGTTAAATAGTGTCTTGAGATA  
AAACTATTGCCACAGAGAAAAAGATTGAAAATCCGCACCCCTCTACTACAAACAATGTTGA  
ACCGAGTAAACCTGAACAGAGAGAAATGTTGCTGATGCCCTTGGAAATCTCAGATAGTG  
ATCCGCTCTACGATATTACGTGGATTCTACGACACATGAAATTATACTTCTTCTAGGG  
AAAGTACAAATGGAAGTGATTAGTGCAGTGTGCAAGAAAAGTATCATGTGGAGATAGAACT  
AAAAGAGCCTACAGTCATTATATGGAGAGACCGTTAAAATGCAGAATATACCATTACA  
TCGAAGTGCCGCAAATCCTTCTGGCTTCCATTGGTTATCTGTATGCCGCTCCGTG  
GGAAAGTGGAAATGCACTGAGAGCTGGTTCTGGATACTTAAATCAATCATTCAAAA  
TGCAGTTATGGAAGGGTACGCTATGGTGCAGAACAGGATTATATGGTGGAAATGTGACGG  
ATTGTAATCTGTTAAGTACGGTTATACTATAGCCCTGTTAGTACTCCAGCAGATT  
CGGATGCTTACTCCTATTGTAAGTACGGTTATACTATAGCCCTGTTAGTACTCCAGCAGATT  
GCCATATCTTAGTTAAAGTTATGCACCAACAGGAATATCTTCACGGCATATAACGATG  
CTCCCAAATATTGTGCAAATATCGTAAACTCAACTGAAAATAATGAGGTCAATTATT

GGAGAAATTCTGCTCGATGTATTCAAGATTATCGCAATGATTAACCTTTTACAAATGG  
GCTTAGTGTGTTAGCAGAGCTAAAAGGATATCAGGTTACCACTGGCGAACCTGTTGCC  
AGACCCGTCGTCTAAATAGTCGGATAGATAAGTAAGATATGTTCAATAAAACTTAG  
TGCCTTTATGTTATATAATGGTTCTTATTAAATAAGATGAAATATTCTTTAATA  
TAGATTGAATTAAAGTGGAAAGGAGGAGATTGTTATTATAAACTACAAGTGGATATTGTGT  
CCTAGTTGTGGAAATAAAACAAGACTACGAATACGAGTGGATACTATACTAAAAATTCCC  
TTTATACAGCCCCAAATGTAAGAACGAACTTTAATTAATGTCAAAAATGAATATAATAA  
CAATCAAAGAGCCAGACGCCAAGACGCAGAGCCGATAATTGAGAAATGAAACTCTCATCTT  
ATCGGCTCTTTGTTATCTGAATTTACTGACTAGCCTCAATATTCCGGCCAGCTT  
ACTATGCCATTATTAAGCTTGTAAATATCGGAGGGTTATTAAATTGGCAGTAAAGTGGCAGTT  
TTGATACCTTAAATGAGATATTATGATAGTGTAGGATATTGACTATCGTACTGCGTTCCC  
TACCGCAAATTAGGAATAAAGGATCTATGTGGGTTGGCTGATTATAGCCAATCCTTTAA  
TTTAAAAAGCGTATGCCGAGAGTTGGTAAATGAAATGAACGAAAAACAAAAGAGAT  
TCGCAGATGAATATATAATGAATGGATGTAATGGTAAAAAGCAGCAATTACAGTAGGTTAT  
AGTAAGAAAACAGCAGAGTCTTAGCAAGTCGATTGTTAAGAAATGTTATGTTCGGAATA  
TATTAAAGAACGATTAGAACAGGTACAAGAACAGCGTTAATGAGTATTACAGAACGTTAG  
CGTTATCTGCTTCTATTGCTAGAGGAGAACCTCAAGAGGCTACAGTAAGAAATATGACCAT  
TTAACGATGAAGTGGAAAAGAGGTTACTTACACAATCACACCAACTTTGAAGAGCGTCA  
GAGATCTATTGACCACACATAACTAAAGTACATGGTGCATATCGATAAAAAAGAAATTACTC  
AGAAGAATATTGAGATTAATATTGGTGAGTACGATGACGAAAGTTAAATTAAACTTAAACAA  
ACCGTCTAATGTTCAATAGCCGGGGGCCAACACACCAACTTTGAAGAGCGTCAGAG  
ATCTATTGACCACACATAACTAAAGTACATGGTGCATATCGATAAAAAAGAAATTACTCAGA  
AGAATATTGAGATTAATATTGGTGAGTACGATGACGAAAGTTAAATTAAACTTAAACAAACC  
GTCTAATGTTCAATAGCCGGGGGCCAACGAGCGGCCATAGTTAAGCCAGCCCCGA  
CACCGCCAACACCCGCTGACGCCCTGACGGCTGTCTGCCGCATCCGCTTACAG  
ACAAGCTGTGACCGTCTCCGGAGCTGCATGTGTCAGAGGTTTACCGTCATACCGAAAC  
GGCGAGACGAAAGGGCTCGTACCGCTATTGTTAGGTTAATGTCATGATAATAATG  
GTTCTTAGACGTCAGGTGGCACTTTGGGAAATGTGCGCGAACCCCTATTGTTATT  
TTCTAAATACATTCAAATATGTATCCGCTCATGAGACAATAACCTGATAATGCTTCAAT  
AATATTGAAAAAGGAAGAGTATGAGTATTCAACATTCCGTGCGCCCTATTCCCTTTT  
GCGGCATTGCTTGCCTTGTGTCACCCAGAAACGCTGGTAAAGTAAAAGATGCTGA  
AGATCAGTTGGGTGCACGAGTGGTTACATCGAACTGGATCTCAACAGCGGTAAGATCCTG  
AGAGTTTCGCCCGAAGAACGTTCCAATGATGAGCAGTAAAGTTCTGCTATGTC  
GCGGTATTATCCCGTATTGACGCCGGCAAGAGCAACTCGGTGCCGCATACACTATTCTCA  
GAATGACTTGGTTGAGTACTCACCAGTCAGAGCAACTCGGTGCCGCATGACAGTAA  
GAGAATTATGCAGTGCTGCCATAACCATGAGTGATAACACTGCGGCCACTTACTCTGACA  
ACGATCGGAGGACCGAAGGAGCTAACCGCTTTTGACAAACATGGGGATCATGTAACCTG  
CCTTGATCGTGGGAACCGGAGCTGAATGAAGCCATACCAACGACGAGCGTACACCACGA  
TGCCTGTAGCAATGGCAACACGTTGCGCAAACATTAAACTGGCGAACTACTTACTCTAGCT

TCCCGGCAACAATTAAATAGACTGGATGGAGGCAGATAAAGTTGCAGGACCACTCTGCGCTC  
GGCCCTCCGGCTGGCTGGTTATTGCTGATAAATCTGGAGCCGGTAGACGTGGGTCTCGCG  
GTATCATTGCAGCACTGGGCCAGATGGTAAGCCCTCCGTATCGTAGTTATCTACACGACG  
GGGAGTCAGGCAACTATGGATGAACGAAATAGACAGATCGCTGAGATAGGTGCCTCACTGAT  
TAAGCATTGGTAACTGTCAGACCAAGTTACTCATATATACTTTAGATTGATTAAAACCTC  
ATTTTAATTAAAAGGATCTAGGTGAAGATCCTTTGATAATCTCATGACCAAAATCCCT  
TAACGTGAGTTTCGTTCCACTGAGCGTAGACCCCGTAGAAAAGATCAAAGGATCTTCTTG  
AGATCCTTTCTGCGCGTAATCTGCTGCTTGCAAACAAAAAACCGACCGCTACCAGCGG  
TGGTTTTTGCCGGATCAAGAGCTACCAACTCTTTCCGAAGGTAACGGCTTCAGCAGA  
GCGCAGATACCAAATACTGTTCTTAGTGTAGCCGTAGTTAGGCCACCACTCAAGAACTC  
TGTAGCACCGCCTACATACCTCGCTCTGCTAACCTGTTACCGAGTGGCTGCTGCCAGTGGCG  
ATAAGTCGTGTCTTACCGGGTTGGACTCAAGACGATAGTTACCGATAAGGCGCAGCGGTG  
GGCTGAACGGGGGTTCGTCACACAGCCCAGCTGGAGCGAACGACCTACACCGAACCTGA  
GATACCTACAGCGTGAGCTATGAGAAAGGCCACGCTCCGAAGGGAGAAAGGGGACAGG  
TATCCGGAAGCGGCAGGGTCGGAACAGGAGAGCGCACGAGGGAGCTCCAGGGGAAACGC  
CTGGTATCTTATAGTCCTGTCGGGTTGCCACCTCTGACTTGAGCGTCGATTTGTGAT  
GCTCGTCAGGGGGCGGAGCCTATGGAAAACGCCAGCAACGCCCTTTACGGTTCTG  
GCCTTTGCTGGCCTTGCTCACATGTTCTCGTTATCCCCTGATTCTGTGGATAA  
CCGTATTACCGCCTTGAGTGAGCTGGCGGTCTAGTTAATGTGTAACGTAACATTAGCTAG  
ATTTTTTATTCAAAAAATTTACAAATATTAGGAAATTAAAGTGTAAAAGAGTTGATAA  
ATGATTATATTGGGACTATAATATAATTAAAGGTCATTGAAATTGTTAACTAATTATCACC  
AAAAAGGAATAGAGTATGAAGTTGGAAATTGTTCTGATCAACCACCAAGGTGAAAC  
TCATAAGCAAGTAATGGATCGCTTGGCTGGTATGCCCTCAGAAGAGGTAGGGTTG  
ATACATATTGGACCTAGAACATCATTACAGAGTTGGCTTACGGAAATTATTGTT  
GCTCGGCTAACCTGTTAGGAAGAACTAAACATTAAATGTTGGCACTATGGGGTTGTTAT  
TCCGACAGCACACCCAGTCGACAGTTAGAAGACGTTATTAGATCAAATGTCGAAAG  
GTCGTTTAATTGGAACCGTTGAGGGCTATACCAAAAGATTTCGACTATTGGT  
GATATGGAAGAGTCTCGAGCAATTACTCAAATTCACCAGATGATAATGGAAAGCTTACA  
GACAGGAACCATTAGCTCTGATAGTGTGATTACATTCAATTCTAAGGTTGATGTATATCCA  
AAAGTGTACTCAAAATGTACCAACCTGTATGACTGCTGAGTCCGCAAGTACGACAGAATGG  
CTAGCAATACAAGGGCTACCAATGGTTCTAGTTGGATTATTGGTACTAATGAAAAAAAGC  
ACAGATGGAACCTATAATGAAATTGCGACAGAAATGGTCAATGATGATATCTAAAATAGATC  
ATTGTATGACTTATATTGTTCTGTTGATGATGATGACAAAAGGCGCAAGATGTTGCGG  
GAGTTCTGAAAAATTGGTATGACTCATATGAAATGCGACCAATATCTTAATGATAGCAA  
TCAAAATCGTGGTTATGATTATCATAAAGGTCATGGCGTGTGTTTACAAGGACATA  
CAAACACCAATCGACGTGTTGATTATAGCAATGGTATTAACCCGTAGGCACCTGAGCAG  
TGTATTGAAATCATTCAACGTGATATTGATGCAACGGGTATTACAAACATTACATGCGGATT  
TGAAGCTAATGGAACGTGAAAGATGAAATAATTGCTTCCATGCGACGCTTATGACACAAGTCG  
CTCCTTCTAAAAGAACCTAAATAATTACTTATTGATACTAGAGATAATAAGGAACAAG

TTATGAAATGGATTATTTCTAAACTTCAGAAAGATGGAATAACATCTGAAGAACG  
TTGGATAATATGGTAAAGACTGTCACGTTAATTGATTCAACTAAATATCATTAAACTGC  
CTTGTTAACATCACCTTCAAAAAATGGTATTGGAGCACCTATTACCGCAGCTG  
GTTTTTATTAGGGTTAACAAATAAATTACATATTGGTCATTAATCAAGTAATTACCA  
CATCACCCGTACGTGTAGCAGAAGGCCAGTTATTAGATCAAATGTCAGAGGGACGCTT  
CATTCTGGTTTAGTGAUTGCGAAAGTGATTGAAATGGAATTAGACGTCATATCT  
CATCAAGGCAACAACAATTGAAGCATGCTATGAAATAATTAATGACGCATTAACACTACAGGT  
TATTGCCATCCCCAAAACGACTTTATGATTGAAAGGTTCAATTAATCCACACTGTTA  
CAGTGAGAATGGACCTAACGAAATATGTATCCGCTACATCAAAGAAGTCGTATGGCAG  
CGAAAAAGGCAGTGCCTTAACGTTAAGTGGAGGATAATTAGAAACCAAAGCCTAT  
GCAATTCTATATAATAAAACAGCACAAACATATGGTATTGATATTGGATGTTGATCATCA  
ATTAACGTAAATTGCGAACTTAAATGCTGATAGAAGTACGGCTCAAGAAGAAGTGAGAGAAT  
ACTTAAAAGACTATACACTGAAACTTACCCCTCAAATGGACAGAGATGAAAAAAATTAACTGC  
ATTATTGAAGAGAATGCAGTTGGGCTCATGACTATTATGAATCGACAAAATTAGCAGT  
GGAAAAAAACAGGGCTAAAAATATTATTATCCTTGAATCAATGTCGATATTAAAGATG  
TAAAAGATATTATTGATATGTTGAACCAAAAAATCGAAATGAATTACCATATAAAATTAA  
AGGCAATTCTATATTAGATTGCCTTTGGCGCCTATTCTAATGCATAATAACTGA  
TAACATCTTATATTGTATTATTTGTATTATCGTTGACATGTATAATTGATATCAA  
AAACTGATTTCCCTCTATTATTCGAGATTATTTCTTAATTCTCTTAAACAAACTAGA  
AATATTGTATATAACAAAAATTATAAATAATAGATGAATAGTTAATTATAGGTGTTCATCA  
ATCGAAAAAGCAACGTATCTTATTAAAGTGCCTGCTTTCTCATTTATAAGGTTAAAT  
AATTCTCATATATCAAGCAAAGTGACA

SEQ ID NO:15 (*mecA* gene loci DNA Sequence (from *Staphylococcus aureus* subsp. *aureus* SA40, complete genome GenBank: CP003604.1))

TATACTACAAATGTAGTCTTATATAAGGAGGATATTGATGAAAAAGATAAAATTGTTCCAC  
TTATTTAATAGTTGAGTGTGGTTGGTATATATTTTATGCTCAAAAGATAAAGAA  
ATTAATAACTATTGATGCAATTGAAGATAAAAATTCAAACAAAGTTATAAAGATAGCAG  
TTATATTCTAAAAGCGATAATGGTGAAGTAGAAATGACTGAACGTCCGATAAAATATA  
ATAGTTAGCGTAAAGATATAAACATTCAAGGATCGTAAAATAAAAGTATCTAAAAT  
AAAAACGAGTAGATGCTCAATATAAAATTAAAACAAACTACGGTAACATTGATCGCAACGT  
TCAATTAAATTGTTAAAGAAGATGGTATGTGGAAAGTTAGATTGGATCATCGTCATTA  
TTCCAGGAATGCAGAAAGACCAAAGCATAACATATTGAAAATTAAAATCAGAACGTGGTAA  
ATTTAGACGAAACAATGTGGATTGGCCAATACAGGAACAGCATATGAGATAGGCATCGT  
TCCAAAGAATGTATCTAAAAAGATTATAAGCAATCGCTAAAGAACTAAAGTATTCTGAAG  
ACTATATCAAACAAACAAATGGATCAAAATTGGGTACAAGATGATACTTCGTTCCACTAAA  
ACCGTTAAAAAAATGGATGAATATTAAGTGGTACATTTGGCAAAAAATTTCATCTTACAAC  
TGAAACAGAAAGTCGTAACTATCCTCTAGAAAAGCGACTTCACATCTATTAGGTTATGTT  
GTCCCATTAACTCTGAAGAATTAAAACAAAAGAATATAAAGGCTATAAGATGATGCAGTT  
ATTGGTAAAAGGGACTCGAAAAACTTACGATAAAAAGCTCCAACATGAAGATGGCTATCG  
TGTCACAATCGTTGACGATAATAGCAATACAATCGCACATACATTAATAGAGAAAAGAAA  
AAGATGGCAAAGATATTCAACTAACTATTGATGCTAAAGTTCAAAGAGTATTATAACAAC  
ATGAAAATGATTATGGCTCAGGTACTGCTATCCACCCCTCAAACAGGTGAATTATTAGCACT  
TGTAAGCACACCTCATATGACGTCTATCCATTATGTATGGCATGAGTAACGAAGAATATA  
ATAAATTAAACCGAAGATAAAAAGAACCTCTGCTCAACAAGTCCAGATTACAACCTCACCA  
GGTTCAACTCAAAAATATTAACAGCAATGATTGGTTAAATAACAAACATTAGACGATAA  
AACAAAGTTATAAAATCGATGGTAAAGGTTGGCAAAAGATAAATCTGGGGTGGTTACAACG  
TTACAAGATATGAAGTGGTAAATGGTAATATCGACTAAAACAAGCAATAGAATCATCAGAT  
AACATTCTTGCTAGAGTAGCACTCGAATTAGGCAGTAAGAAATTGAAAAGGCATGAA  
AAAATAGGTGTTGGTGAAGATATACCAAGTGATTCCATTATGCTCAAATTCAA  
ACAAAATTAGATAATGAAATATTATTAGCTGATTCAAGGTACGGACAAGGTGAAATACTG  
ATTAACCCAGTACAGATCCTTCAATCTATAGCGCATTAGAAAATAATGGCAATATTAACGC  
ACCTCACTTATTAAAAGACACGAAAAACAAAGTTGGAAGAAAATATTATTCAGAAAGAAA  
ATATCAATCTATTAACTGATGGTATGCAACAAGTCGTAATAAAACACATAAAGAAGATATT  
TATAGATCTTATGCAAACCTTAATTGGCAAATCCGGTACTGCGAACTCAAATGAAACAAGG  
AGAAAATGGCAGACAAATTGGGTGGTTATATCATATGATAAAGATAATCCAAACATGATGA  
TGGCTATTAAATGTTAAAGATGTACAAGATAAAGGAATGGCTAGCTACAATGCCAAATCTCA  
GGTAAAGTGTATGATGAGCTATATGAGAACGGTAATAAAAGTACGATATAGATGAATAACA  
AACAGTGAAGCAATCCGTAACGATGGTTGCTTCACTGTTT

SEQ ID NO:16 (*mecA* transcript sequence)

UAGUCUUUAUAAGGAGGAUUGAUGAAAAAGAUAAAAUUGUUCCACUUUUUUAAUAGU  
UGUAGUUGUCGGGUUUGGUUAUAUUUUUAUGCUUCAAAAGAUAAAGAAUUAUAAUACUA  
UUGAUGCAAUUGAAGAUAAAUAUUCAAACAAGUUUAAGAUAGCAGUUUAUUCUAAA  
ACGGAUAAUGGUGAAGUAGAAAUGACUGAACGUCCGAUAAAUAUAAUAGUUUAGGCGU  
UAAAGAUUAACAUUCAGGAUCGUAAAUAUAAAAGUAUCUAAAUAUAAAACGAGUAG  
AUGCUCAAUUAUAAAUAUAAAACAUACGGUACAUUGAUCGCAACGUUCAUUAUUCUAAA  
GUUAAAGAAGAUGGUAGUGGAAGGUAGAUUGGGAUCAUAGCGUCAUUUCCAGGAAUGCA  
GAAAGACCAAAGCAUACAUUUGAAAUAUAAAUCAGAACGUGGUAAAUAUAGACCGAA  
ACAAUUGGGAUUGGCCAAUACAGGAACAGCAUAUGAGAUAGGCAUCGUUCCAAAGAAUGUA  
UCUAAAAAGAUUAUAAAGCAAUCGUAAAAGAACUAAGUAUUCUGAAGACUUAUCAAAACA  
ACAAAUGGAUAAAUAUUGGUACAAGAUGAUACCUUCGUUCCACUUAAAACGUUAAAAAAA  
UGGAUGAAUUAUUAAGUGAUUUCGCAAAAAAAUUCAUUACAACUAAUGAAACAGAAAGU  
CGUAACUAUCCUCUAGAAAAGCGACUUCACAUUAGGUUAUGUUGGUCCCAUUAACUC  
UGAAGAAUAAAACAAAAGAAUUAUAAAGGCUAAAGAUGAUGCAGUUAUUGGUAAAAGG  
GACUCGAAAACUUUACGAUAAAAGCUCCAACAUGAAGAUGGCUAUCGUGUCACAAUCGUU  
GACGAUAAUAGCAAUACAUUCGCACAUACAUUAAUAGAGAAAAGAAAAAGAUGGCAAAGA  
UAUUCACUAACUAUUGAUGCUAAGUUCAAAAGAGUAUUAACACAAUCAUGAAAAGAUU  
AUGGCUCAGGUACUGCUAUCCACCCUCAAACAGGUGAAUUAUAGCACUUGUAAGCACACC  
UCAUAUGACGUCUAUCCAUUUAUGUAUGGCAUGAGUAACGAAGAAUUAUAAAACCGA  
AGAUAAAAAGAACCUUCUGCUACAAGUUCCAGAUUACAACUUCACCAGGUUCAACUAAA  
AAAUAUUAACAGCAAUGAUUGGUAAAACAAAACAUUAGACGAUAAAACAAGUUUAAAA  
AUCGAUGGUAAAGGUUGGCAAAAAGAUAAAUCUUGGGUGGUACACGUUACAAGAUUAUGA  
AGUGGUAAAUGGUAAUACGACUAAAACAAGCAAUAGAAUCAUCAGAUUACAUUUUCUUUG  
CUAGAGUAGCACUCGAAUUAGGCAGUAAGAAAUAUUGAAAAGGCAUGAAAAACUAGGUU  
GGUGAAGAUUAACCAAGUGAUUAUCCAUUUUAUAGCUAAAUAUCAACAAAUAUUGA  
UAAUGAAAUAUUAUAGCUGAUUACGGACAAGGUGAAAACUGAUUAACCCAGUAC  
AGAUCCUUCAAUCUAAUAGCGCAUJAGAAAAUAUAGGCAAUUAACGCACCUCUUAUUA  
AAAGACACGAAAACAAAGUUUGGAAGAAAAUAUUAUUCAAAGAAAUAUCAACUUAU  
AACUGAUGGUAAUGCAACAAGUCGUAAAACACAUAAAGAAGAUUUUAUAGAUUCUUAUG  
CAAACUUUAUUGGCAAAUCGGUACUGCAGAACUCAAAUAGAAACAAGGAGAACUGGCAGA  
CAAUUGGGUGGUUUAUCAUAUGUAAAAGAUAAAUCCAAACAUGAUGAUGGUCAUUUAUGU  
UAAAGAUGUACAAGAUAAAGGAUUGGUAGCUACAUUGCCAAAUCUCAGGUAAAGUGUAUG  
AUGAGCUUAUAGAGAACGGUAAAUAUACGAUUAUAGAUGAUUAACAAAACAGUGAAGCA  
AUCCGUACGAUGGUUGCUUCACUGUUUU

SEQ ID NO:17 (*luxAB* gene loci DNA sequence (from *Vibrio fischeri* genes *luxA* and *luxB* for luciferase alpha and beta subunits - GenBank: X06758.1))

GGCTTAAATAAACAGAACATCACCAAAAGGAATAGAGTATGAAGTTGAAATATTTGTTTT  
CGTATCAACCACCAGGTGAAACTCATAAGCTAAGTAATGGATCGCTTGGCTGGTAT  
CGCCTCAGAAGAGTAGGGTTGATACATATTGGACCTAGAACATCATTTACAGAGTTGG  
TCTTACGGAAATTATTTGTTGCTCGGCTAACCTGTTAGGAAGAACTAAAACATTAAATG  
TTGGCACTATGGGGTTGTTATTCCGACAGCACACCCAGTCGACAGTTAGAAGACGTTTA  
TTATTAGATCAAATGTCGAAAGGTGTTGATATGGAAGAGTCTCGAGCAATTACTCAAATTCACC  
AGATTTCGAGTATTGGTGTGATATGGAAGAGTCTCGAGCAATTACTCAAATTCACC  
AGATGATAATGGAAGCTTACAGACAGGAACCATTAGCTCTGATAGTGATTACATTCAATT  
CCTAAGGTTGATGTATATCCAAAGTGTACTCAAAAAATGTACCAACCTGTATGACTGCTGA  
GTCCGCAAGTACGACAGAATGGCTAGCAATACAAGGGCTACCAATGGTCTTAGTTGGATTA  
TTGGTACTAATGAAAAAAAGCACAGATGGAACTCTATAATGAAATTGCGACAGAATATGGT  
CATGATATATCTAAAATAGATCATTGTATGACTTATATTGTTCTGTTGATGATGATGCACA  
AAAGGCGCAAGATGTTGTCGGAGTTCTGAAAATTGGTATGACTCATATGAAATGCGA  
CCAATATCTTAATGATAGCAATCAAACCTCGTGGTTATGATTATCATAAAGGTCAATGGCGT  
GATTTGTTTACAAGGACATACAAACACCAATCGACGTGTTGATTAGCAATGGTATTAA  
CCCTGTAGGCACCTCTGAGCAGTGTATTGAAATCATTCAACGTGATATTGATGCAACGGTA  
TTACAAACATTACATCGGATTGAAAGCTAATGGAACTGAAGATGAAATAATTGCTCCATG  
CGACGCTTATGACACAAGTCGCTCCTTCTAAAAGAACCTAAATAATTACTTATTGAT  
ACTAGAGATAATAAGGAACAAGTTATGAAATTGGATTATTTCTAAACTTCAGAAAGA  
TGGAAATAACATCTGANGAACGTTGGATAATATGGTAAAGACTGTCACGTTAATTGATTCAA  
CTAAATATCATTAAACTGCCTTGTAAATGAACATCACTTTCAAAAATGGTATTGTT  
GGAGCACCTATTACCGCAGCTGGTTTTATTAGGTTAACAAATAATTACATATTGGTTC  
ATTAAATCAAGTAATTACCAACCCATCACCTGTACGTGTTAGCAGAAGAACCGAGTTATTAG  
ATCAAATGTCAGAGGGACGCTTCATTCTGGTTTAGTGAATGCGAAAGTGAATTGAAATG  
GAATTGGTAGACGTATCTCATCAAGGCAACAACAATTGAAGCATGCTATGAAATAAT  
TAATGACGCATTAACTACAGGTTATTGTATCCCAAAACGACTTTATGATTTCCAAAGG  
TTTCAATTAACTCCACACTGTTACAGTGAGAATGGACCTAACAGCAATATGATCCGCTACATCA  
AAAGAAGTCGTATGTGGCAGCGAAAAGGCAGTCACGTTAACATTAAAGTGGGAGGATAA  
TTTAGAAACCAAAAGAACGCTATGCAATTCTATATAATAAAACAGCACAACAATTGGTATTG  
ATATTTCGGATGTTGATCATCAATTAACTGTAATTGCGAACTAAATGCTGATAGAAGTACG  
GCTCAAGAAGAAGTGAGAGAATACTTAAAGACTATATCACTGAAACTTACCCCTCAAATGGA  
CAGAGATGAAAAAAATTAACTGCATTATTGAAGAGAATGCAGTTGGTCTCATGATGACTATT  
ATGAATCGACAAATTAGCAGTGGAAAAACAGGGTCTAAAATATTTATTATCCTTGAA  
TCAATGTCCGATATTAAAGATGTAAAAGATATTATTGATATGTTGAACCAAAATCGAAAT  
GAATTTACCATATAAAATTAAAGGCAATTCTATATTAGATTGCCTTTAAATTTC

SEQ ID NO: 18 (*luxAB* Transcript Sequence)

AAUCACCAAAAGGAAUAGAGUAUGAAGGUUUGGAAUUAUUGUUUUUCGUAUCAACCACCAAG  
GUGAAACUCAUAAGCUAAGUAAUGGAUCGCUUJGUUCGGCUUGGUACGCCUCAGAAGAGUA  
GGGUUJUGAUACAUAUUGGACCUCUAGAACAUCAUUUACAGAGUUUGGUCUACGGGAAAUU  
AUJUGUUGCUGCGCUAACCUUGGUAGGAAGAACUAAAACAUAAAUGUUGGCACUAUGGGGG  
UUGUUAUUCGACAGCACCCCAGUUCGACAGUAGAACGCUUUUAAUJAGAUCAAAUG  
UCGAAAGGUCGUUUUAAUUGGAACCGUUCGAGGGCUAUACCAUAAAGAUUUUCGAGUAUU  
UGGUGUUGAUUAUGGAAGAGUCUCGAGCAAUACUCAAAAUUCUACCAGAUGAUAAUGGAAA  
GUUACAGACAGGAACCAUAGCUCUGAUAGUGAUUACAUCAUUUCCUAAAGGUUGAUGUA  
UAUCCCAAAGUGUACUCAAAAAGUACCAACCUGUAUGACUGCUGAGUCCGCAAGUACGAC  
AGAAUGGCUAGCAAUACAAGGGCUACCAAUGGUUCUAGUUGGAUUAUGGUACUAAUGAAA  
AAAAGCACAGAUGGAACUCUAAUUGAAAUGCGACAGAAUAAUGGUCAUGAUUAUCUAAA  
AUAGAUCAUUGUAUGACUUUAUUGUUCGUUGAUGAUGAUGGCACAAAGGCGCAAGAUGU  
UUGUCGGAGUUUCUGAAAAAUUGGU AUGACUCAUAUGUAAAUGCGACCAAUACUUUUAUG  
AUAGCAAUCAAACUCGUGGUUAUGAUUAUCAUAAAGGUCAAUGGCUGAUUUGUUUUACAA  
GGACAUACAAACACCAAUCGACGUGUUGAUUAUAGCAAUGGUAAAACCCUGUAGGCACUCC  
UGAGCAGUGUAUUGAAUCAUCAGUGAUUUGAUGCAACGGGUUUACAAACAUUACAU  
GCGGUUUGAAGCUAAUGGAACUGAAGAUGAAAUAUGCUUCCAUGCGACGCCUUUAUGACA  
CAAGUCGCUCCUUUCUUAAAAGAACCUAAAUAUACUUAUUGAUACUAGAGAUUAAG  
GAACAAGUUUAUGAAAUAUGGUUAUUGGUUACUUAUUCAGAAAGAUGGAUUAACUUGA  
AGAACGUUGGAUUAUUGGUAAAGACUGUCACGUAAAUGAUUCAACUAAAUAUCAUUUA  
AUACUGCCUUUGUUAUGAACAUACACUUUCAAAAAAUGGUUUGUUGGAGCACCACUAC  
GCAGCUGGUUUUUUUUAGGGUUAACAAUAAAUAUACAUUUGGUCAUAAAUAUCAAGUAAU  
UACCACCCAUCAACCCUGUACGUGUAGCAGAAGAAGCCAGUUUAUAGAUCAAAUGUCAGAGG  
GACGCUUCAUUCUUGGUUUAGUGACUGCGAAAGUGAUUUCGAAAUGGAUUUUUAGACGU  
CAUAUCUCAAGCAACAACAUUUGAAGCAUGCUAUGAAAUAUUAUGACGCAUUAAC  
UACAGGUUAUUGUCAUCCCCAAACGACUUUAUGAUUUUCAAAGGUUCAUAAAUAUCCAC  
ACUGUUACAGUGAGAAUGGACCUAAGCAAAUUGUAUCCGUACAUCAAAAGAACGUCGUAC  
UGGGCAGCGAAAAGGCACUGCCUUUACAUUUAUGGGAGGAUAAAAGAACCAAAGA  
ACGCUAUGCAAUUCUAAUAAAACAGCACAACAUUACUGGUUUGAUUUCGGAUGUUG  
AUCAUCAUUAACUGUAUUGCGAACAUAAAUGCUGAUAGAACGUACGGCUCAAGAACGAG  
AGAGAAUACUAAAAGACUUAUCACUGAAACUUACCCUAAAUGGACAGAGAUAAAAG  
UAACUGCAUUAUUGAAGAGAAUGCAGUUGGGUCUCAUGAUGACUUAUUAUGAACGAC  
UAGCAGUGGAAAAACAGGGUCUAAAUAUUAUCCUUUGAAUCAUUGUCCGUAUUAU

AAAGAUGUAAAAGAUUUUUGAUUUGAACCACCAAAUCGAAUUGAAUUAACAUAA  
AAAUUAAGGCAAUUCUAUUAUAGAUUGCCUUU

SEQ ID NO: 19 (cis-repressed *luxAB* transcript sequence)

AAAGGCAUGAAAAACUJGGUAUCUUCACCAACACCUAGCUUUJUGAAGGAAUJUGAGUAUGA  
AGUUJGGAAAUUJUGGUUGUUCGUAUCAACCACCAGGUGAACUCAUAAGCUAAAGGCAUGA  
AAAAACUAGGUGAUCUUCACCAACACCUAGUUUUUCAAGGAAUJUGAGUAUGAAGUJUGGAA  
AUAUUJGUUUUCGUAUCAACCACCAGGUGAACUCAUAAGCUAAGUAAUGGAUCGCUUJGU  
UCGGCUJGGUAUCGCCUCAGAAGAGUAGGGUUUGAUACAUUJUGGACCUUAGAACAUCAUU  
UACAGAGUJUGGUUCGUACGGAAUJGUUGCUGCGCUACCUGUUAGGAAGAACAUCAUU  
AAACAUAAAUGUJUGGCACUAUGGGGUUGUUAUCCGACAGCACACCCAGUUCGACAGUUA  
GAAGACGUUUUAUUAUAGAUCAAAUGUCGAAAGGUJGUUUUAAUJUGGAAACCGUUCGAGG  
GCUAUACCAUAAAGAUUUUCGAGUAUUJUGGUUGAUUAUGGAAGAGUCUCGAGCAAUUACUC  
AAAUUUCUACCAGAUGAUAUAGGAAAGCUJACAGACAGGAACCAUJAGCUCUGAUAGUGAU  
UACAUUCAAUUCCUAAGGUUGAUGUUAUCCCAAAGGUACUCAAAAAAUGUACCAACCUG  
UAUGACUGCUGAGUCCGCAAGUACGACAGAAUGGUAGCAAUACAAGGGCUACCAAUGGUUC  
UUAGUJGGAUUAUJUGGUACUAAUGGAAAAAGCACAGAUGGAACUCUUAUUAUGGAAUJUGCG  
ACAGAAUUAUGGUCAUGAUUAUCUAAAUAUGAUCAUUGUAUGACUUUAUUAUGGUUCUGUJUGA  
UGAUGAUGCACAAAAGGCGCAAGAUGUUJUGUCCGGAGUUUCUGAAAAUJUGGUAGACUCAU  
AUGUAAAUGCGACCAAUAUCUUUAUGAUAGCAAUACUCGUGGUUAUGAUUAUCAUAAA  
GGUCAAUGGCGUGAUUUJGUUUUACAAGGACAUACAAACACCAUCGACGUGUJGUUAUAG  
CAAUGGUAUUAACCCUGUAGGCACUCCUGAGCAGGUUAUGAAAUCUCAACGUGUUAUUG  
AUGCAACGGGUUAUACAAACAUUACAUGCGGAAUJUGAAGCUAAUGGAACUGAAGAUGGAAU  
AUUGCUUCCAUGCGACGCUUUAUGACACAAGUCGCUCUUCUAAAAGAACCUAAAUAU  
UACUUUAUJUGAUACAGAGAUUAAGGAACAAGGUUAUGGAAUJUGGUUAUUAUUCUAAA  
CUUUCAGAAAGAUGGAAUACAUCUGAAGAACGUJUGGUUAUUAUGGUAAAGACUGUCACGU  
UAAUJUGAUCAACAAUAUCUUUAUACUGCCUUJGUUAUGAACAUCAUCACUUUCAAAA  
AAUGGUAUJUGGUAGCACCUCUACCGCAGCUGGUUUUUUUAUGGUUAACAAUAAAUAU  
ACAUAUJGGUCAUAAAUCAGUAAUJACCCACCAUCACCCUGUACGUGUAGCAGAAGAAC  
CCAGUUUAUAGAUCAAUGUCAGAGGGACGGCUUCAUUCUUGGUUUUAGUGACUGCGAAAGU  
GAUUUCGAAAUGGAAUJGUUGACGUCAUACUCAAGGCAACAACAUJUGAAGCAUG  
CUAUGAAUUAUUAUGACGCAUUAACUACAGGUUAUJGUCAUCCCAAAACGACUUUAUG  
AUUUUCCAAAGGUUCAUAAAUCACACUGUUACAGUGAGAAUGGACCUAGCAUAUJUGU  
UCCGUACAUCAAAAGAACGUCAUGUGGGCAGCGAAAAGGCACUGGUUUACAUUAAA  
GUGGGAGGAUAAAAGAACCAAAGAACGCUAUGCAAUUCUUAUAAAACAGCACAAC  
AAUAUGGUAUJUGAUUUJCGAUGUUGAUCAUAAAUCAGGUUAUJUGGUACUJGUAAAUGCU  
GAUAGAAGUACGGCUCAAGAAGAAGUGAGAGAAUACUAAAAGACUUAUACACUGAAACUUA  
CCCUCAAAUGGACAGAGAUGGAAAAACUGCAUUAUGGUAGAGAAUGCAGUUGGGCUC  
AUGAUGACUUAUUAUGAAUCGACAAAUAUGCAGUGGAAAAACAGGGGUCAAAAUAU  
UUAUCCUUJUGAAUCAAUGUCCGAUAAAAGAUGUAAAAGAUUAUJUGUUAUGUUGAACCA  
AAAAACGAAAUGGUUACCAUAAAUAAGGCAUUCUUAUAGAUJUGGUUU

**THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:**

1. A bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising:
  - a lysogenized bacteriophage genome comprising a first bacteriophage gene comprising a deletion of a packaging initiation site sequence of said first bacteriophage gene that prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and
  - a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage gene comprising a second packaging initiation site sequence that facilitates the packaging a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said second bacteriophage gene encodes a protein, wherein said replica of said reporter nucleic acid molecule plasmid forms a replicon for packaging into said non-replicative transduction particle.
2. A bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising:
  - a lysogenized bacteriophage genome lacking a bacteriophage gene encoding a packaging initiation site sequence, wherein deletion of said bacteriophage gene prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and
  - a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage gene, wherein said second bacteriophage gene encodes a packaging initiation site sequence and facilitates the packaging a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said second bacteriophage gene is capable of expressing a protein that is encoded by said gene, wherein said replica of said reporter nucleic acid molecule plasmid forms a replicon amenable to packaging into said non-replicative transduction particle.

3. A bacterial cell packaging system for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, said bacterial cell packaging system comprising a host bacteria cell comprising:
  - a lysogenized bacteriophage genome comprising a first bacteriophage packaging initiation site sequence, wherein said first bacteriophage packaging initiation site sequence comprises a mutation that prevents packaging of a bacteriophage nucleic acid molecule into said non-replicative transduction particle; and
  - a reporter nucleic acid molecule plasmid comprising a reporter gene and comprising a second bacteriophage packaging initiation site sequence, wherein said second bacteriophage packaging initiation site sequence lacks said mutation and facilitates the packaging of a replica of said reporter nucleic acid molecule plasmid into said non-replicative transduction particle, wherein said replica of said reporter nucleic acid molecule plasmid forms a replicon for packaging into said non-replicative transduction particle.
4. The bacterial cell packaging system of any one of claims 1 to 3, wherein said reporter nucleic acid molecule plasmid is operatively linked to a promoter.
5. The bacterial cell packaging system of any one of claims 1 to 4, wherein said reporter nucleic acid molecule plasmid comprises an origin of replication.
6. The bacterial cell packaging system of any one of claims 1 to 5, wherein said replicon comprises a concatamer amenable to packaging into said non-replicative transduction particle.
7. The bacterial cell packaging system of claim 1 or 2, wherein said second bacteriophage gene comprises the sequence set forth in SEQ ID NO:9.
8. The bacterial cell packaging system of claim 1, wherein said first bacteriophage gene comprises the sequence of SEQ ID NO:8.
9. The bacterial cell packaging system of claim 1 or 2, wherein said first and said second bacteriophage genes each comprises a small terminase (terS) gene comprising said packaging initiation site sequence, optionally wherein said terS gene is a *Staphylococcus aureus* (*S. aureus*) bacteriophage φ11 or φ80α terS gene.

10. The bacterial cell packaging system of claim 3, wherein said first and said second bacteriophage packaging initiation site sequences each comprise a packaging initiation site sequence from a small terminase gene, or said first and said second bacteriophage packaging initiation site sequences each comprise a pac-site sequence from a small terminase (terS) gene of an *S. aureus* bacteriophage φ11 or φ80α..
11. The bacterial cell packaging system of any one of claims 1 to 3, wherein said replicon is derived from a *S. aureus* pT181 plasmid origin of replication.
12. The bacterial cell packaging system of any one of claims 1 to 3, wherein said replicon comprises the sequence set forth in SEQ ID NO:5.
13. The bacterial cell packaging system of any one of claims 1 to 3, wherein said packaging initiation site sequence of said second bacteriophage gene or said packaging gene comprises a pac-site.
14. The bacterial cell packaging system of claim 1 or 2, wherein said second bacteriophage gene is operatively linked to a promoter, optionally wherein said promoter is an inducible promoter or a constitutive promoter.
15. The bacterial cell packaging system of any one of claims 1 to 3 wherein said bacteriophage comprises a *S. aureus* bacteriophage φ80α or a bacteriophage φ11.
16. The bacterial cell packaging system of any one of claims 1 to 3, wherein said bacterial cell comprises:
  - an *Escherichia coli* (*E. coli*) cell;
  - a Gram-negative cell; or
  - a Gram-positive cell.
17. The bacterial cell packaging system of any one of claims 1 to 34, wherein said bacterial cell comprises an *S. aureus* cell.

18. The bacterial cell packaging system of any one of claims 1 to 3, wherein said reporter gene encodes a detectable and/or a selectable marker,
  - or optionally wherein said reporter gene is selected from the group consisting of (i) genes encoding enzymes mediating luminescence reactions optionally selected from luxA, luxB, luxAB, luc, ruc, or nluc, (ii) genes encoding enzymes mediating colorimetric reactions optionally selected from lacZ or HRP, (iii) genes encoding fluorescent proteins optionally selected from GFP, eGFP, YFP, RFP, CFP, BFP, mCherry or near-infrared fluorescent proteins, (iv) nucleic acid molecules encoding affinity peptides optionally selected from His-tag or 3X-FLAG, and genes encoding selectable markers optionally selected from ampC, tet(M), CAT or erm.
19. A method for packaging a reporter nucleic acid molecule into a non-replicative transduction particle, comprising:
  - providing conditions to said bacterial cell packaging system of any one of claims 1 to 18 that induce a lytic phase of said bacteriophage to produce non-replicative transduction particles packaged with said reporter nucleic acid molecule; and
  - isolating said non-replicative transduction particle comprising a replica of said reporter nucleic acid molecule plasmid, optionally wherein said non-replicative transduction particle does not contain a replicated bacteriophage genome, or optionally wherein said induction of said lytic phase triggers excision of said genomic island nucleic acid molecule from said genome of said bacterial cell.
20. A composition comprising a non-replicative transduction particle comprising a replica of said reporter nucleic acid molecule plasmid produced by the method of claim 19.
21. A method for detecting a presence or an absence of a bacterial cell in a sample, comprising:
  - introducing into a sample a non-replicative transduction particle produced by the method of claim 19 comprising a reporter gene encoding a reporter molecule and lacking a bacteriophage genome under conditions such that said non-replicative transduction particle can transduce said bacterial cell and wherein said reporter gene can be expressed in said bacterial cell;
  - providing conditions for activation of said reporter molecule; and

detecting for a presence or an absence of a reporter signal transmitted from said expressed reporter molecule, wherein a presence of said reporter signal correctly indicates said presence of said bacterial cell.

22. The method of claim 21, wherein said bacterial cell is a Methicillin Resistant *Staphylococcus aureus* (MRSA) cell or a Methicillin Sensitive *Staphylococcus aureus* (MSSA) cell.

23 The method of claim 21 or 22, further comprising:  
providing an antibiotic to said sample at a pre-determined concentration and detecting a presence or absence of said reporter signal to determine whether said bacterial cell is resistant or sensitive to said antibiotic; or  
providing varying pre-determined concentrations antibiotic to said sample and detecting the amount of said reporter signal to determine the minimum inhibitory concentration of said bacterial cell to said antibiotic.






FIGURE 1




Schematic of *pGWP10001*

FIGURE 2



Schematic of a pac-site deletion/complementation packaging system (300)

FIGURE 3



Schematic of *pGW80A0001*

FIGURE 4






FIGURE 5

*Schematic of a natural genomic-island packaging system (500)*




Schematic of a genomic island (GI)-based packaging system.

FIGURE 6



Schematic of a genomic-island packaging system that does not integrate into host genome (700)

FIGURE 7



*Schematic of a SaP1bov2 packaging system that does not integrate into host genome (800)*

FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15





FIGURE 16



GURE 17





FIGURE 18








FIGURE 20



*Photograph of transduced MRSA clinical isolates*

FIGURE 21



*Luminescence in *S. aureus* and distinguishing MRSA vs. MSSA*

FIGURE 22

23/35





FIGURE 23

Category	Subcategory	Value
System	Processor	Intel Core i7-9700K
System	RAM	16GB DDR4 3200MHz
System	SSD	1TB NVMe
System	HDD	2TB
System	Power Supply	650W
System	Case	MSI Aegis 3
System	GPU	NVIDIA RTX 2080 Super
System	Monitor	24" 144Hz 1080p
System	Keyboard	Razer BlackWidow
System	Mouse	Razer DeathAdder
System	Mousepad	Razer Goliathus
System	Headphones	SteelSeries Arctis 7
System	Microphone	SteelSeries Arctis 1
System	Mousepad	SteelSeries QcK



*Luminescence obtained from MRSA and MSSA at various concentrations of cefoxitin*

FIGURE 24




FIGURE 25




FIGURE 26




FIGURE 27




FIGURE 28




FIGURE 29

Transcription Terminator Sequence  
2905

FIGURE 30




FIGURE 31




FIGURE 32




FIGURE 33




FIGURE 34




FIGURE 35

MRSA: *mecA* (+) *S. aureus*

25938PCT\_CRF\_SequenceListing.txt  
SEQUENCE LISTING

<110> GENEWAVE BIOSCIENCES, INC.

<120> NON-REPLICATIVE TRANSDUCTION PARTICLES AND TRANSDUCTION PARTICLE-BASED REPORTER SYSTEMS

<130> 28421-25938/PCT

<140> PCT/US2014/026536

<141> 2014-03-13

<150> 61/939, 126

<151> 2014-02-12

<150> 61/897, 040

<151> 2013-10-29

<150> 61/779, 177

<151> 2013-03-13

<160> 22

<170> PatentIn version 3.5

<210> 1

<211> 177

<212> DNA

<213> Enterobacteria phage P1

<400> 1

ccactaaaaa gcatgatcat tgcactctt aatgatcaac atgcaggta tcacattgc 60

gctgaaatag cgaaaaaca aagatataat gccgttgtca gtgccgcagt cgagaatgc 120

aagcgccaaa ataagcgcat aaatgatcgt tcagatgatc atgacgtgat caccgc 177

<210> 2

<211> 177

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 2

ccactaaaaa gcatgataat agaccactt aacgaccaac atgcaggaga gcacattgc 60

gctgaaatag cgaaaaagca gagggtaat gccgttgtca gtgccgcagt cgagaatgc 120

aagcgccaaa ataagcgcat aaacgaccgt tcagacgacc atgacgttat taccgc 177

<210> 3

<211> 1727

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 3

cactataggg cgaattggcg gaaggccgtc aaggccgtt ttggggccgg cgccgcggat 60

ccgctagctc tagactggca ggttctgag cagatgtcc aacccgatct ggatcgggtc 120

25938PCT\_CRF\_SequenceListing.txt

agaaaaat	tttcgagctg	180
gtgattgaag	tttgcataa	240
aatatataa	acccatca	300
gttgaat	cgaccgaccg	360
ggactcggt	cccataaagc	420
atcgaagagc	ttgaaaaaga	480
agcaggattt	cttacacgac	540
ttgcttgc	tcgtataacc	600
ccaccagcat	ggcaaagg	660
ttacttttc	ttggctatgaa	720
ccgggcata	ttggcctgct	780
acaaggcct	ttgtcaaaatc	840
cattccgt	tcgcggcaac	900
gtgaacgt	aaccagt	960
ccaaaaat	ttggttat	1020
ctgttctcg	ttttttat	1080
atacccc	ccctgtc	1140
aaacaaaa	tttttatt	1200
atcagtttcc	ttttttat	1260
ggatcgct	ttttttat	1320
aacatgag	ttttttat	1380
taaacgaag	ttttttat	1440
cgccaggatt	ttttttat	1500
ctagtttgcc	ttttttat	1560
ggacgccc	ttttttat	1620
gcttaaaaa	ttttttat	1680
ggcccttct	ttttttat	1727

<210> 4  
 <211> 131  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Description of Artificial Sequence: Synthetic  
 polynucleotide

<400> 4  
 cgtcagggtgg cactttcgg gaaatgtcg cgaaaccctt atttgtttat tttctaaata

60

25938PCT\_CRF\_SequenceListing.txt

cattcaaata	tgtatccgct	catgagacaa	taaccctgat	aatgcttca	ataatattga	120
aaaggaagag	t					131
<210> 5						
<211> 4681						
<212> DNA						
<213> Artificial Sequence						
<220>						
<223> Description of Artificial Sequence: Synthetic polynucleotide						
<220>						
<221> modified_base						
<222> (301)..(301)						
<223> a, c, t, g, unknown or other						
<220>						
<221> modified_base						
<222> (4056)..(4056)						
<223> a, c, t, g, unknown or other						
<400> 5						
tttgcggaaa	gagttagtaa	gttaacagaa	gacgaggccaa	acctaaatgg	tttagcagga	60
aacttagata	aaaaatgaa	tccagaatta	tattcagaac	aggaacagca	acaagagcaa	120
caaaagaatc	aaaacgaga	tagaggtatg	cacttataga	acatgcattt	atgccgagaa	180
aacttattgg	ttgaaatggg	ctatgtgtta	gctaacttgt	tagcgagttg	gttggacttg	240
aattgggatt	aatcccaaga	aagtaccggc	tcaacaaccc	ataaaggccct	gtaggttccg	300
nccaaaataagg	aaatttggaaat	aaagcaataa	aaggagttga	agaaaatgaaa	ttcagagaag	360
cctttgagaa	ttttataaca	agtaagtatg	tacttggtgt	tttagtagtc	ttaactgttt	420
accagataat	acaaatgctt	aaataaaaaaa	agacttgatc	tgatttagacc	aaatcttttg	480
atagtgttat	attaataaca	aaataaaaag	gagtcgtca	cggccatcca	aagtttgtga	540
acgacatcat	tcaaagaaaa	aaacactgag	ttgtttttat	aatcttgtat	atttagatat	600
taaacgatat	ttaaatatac	atcaagatat	atattgggt	gagcgattac	ttaaacgaaa	660
ttgagattaa	ggagtcgatt	tttatgtat	aaaaacaatc	atgcaaatca	ttcaaatacat	720
ttggaaaatc	acgatttaga	caattttct	aaaaccggct	actctaatacg	ccgggttggac	780
gcacatactg	tgtgcataatc	tgtccaaaa	ttaagtttg	atgcaatgac	gatcggttgg	840
aatctcaacc	gagacaacgc	tcaggccctt	tctaaattta	ttagtgtaga	gccccaataa	900
agactttggg	atattcttca	aacaaagttt	aaagctaaag	cacttcaaga	aaaagtttat	960
attgaatatg	acaaagtcaa	agcagatagt	tggatagac	gtaatatgcg	tattgaattt	1020
aatccaaaca	aacttacacg	agatgaaatg	atttggtaa	aacaaaatat	aataagctac	1080
atggaagatg	acggttttac	aagatttagat	ttagcctttg	atttgaaga	tgatttgagt	1140
gactactatg	caatgtctga	taaagcagtt	aagaaaacta	tttttatgg	tcgtaatgg	1200
aagccagaaaa	caaaatattt	tggcgtgaga	gatagtaata	gatttattag	aatttataat	1260

25938PCT_CRF_SequenceListing.txt	
aaaaagcaag aacgtaaaga taatgcagat gctgaagtttgtctgaaca tttatggcgt	1320
gtagaaatcg aactaaaag agatatggg gattactgga atgattgctt tagtattta	1380
catacttgc aaccagattg gaaaactatc caacgcactg cgatagagc aatagtttt	1440
atgttattga gtatgaaga agaatggg aagtttcaca gaaattctag aacaaaatat	1500
aagaatttga taaaagaaat ttgcgcagtc gatttaacgg acttaatgaa atcgacttta	1560
aaagcgaacg aaaaacaatt gcaaaaacaa atcgattttt ggcaacatga atttaaattt	1620
tggaaatagt gtacatatta atattactga acaaaaatga tatatttaaa ctattctaat	1680
ttaggaggat tttttatga agtgtctatt taaaaatttgg gggattttat atgaggtgaa	1740
agaataaattt acccctataa actttagcca cctcaagtaa agaggtaaaa ttgttttagtt	1800
tatataaaaa atttaaaggt ttgtttata gcgtttattt ttggctttgt attcttcatt	1860
tttttagtgt attaaatgaa atggtttaa atgtttctt acctgatatt gcaaatcatt	1920
ttaatactac tcctggatt acaaactggg taaacactgc atatatgttta acttttcga	1980
taggaacagc agtataatgga aaattatctg attatataaa tataaaaaaaa ttgttaattt	2040
ttggatttag tttgagctgt ctgggttcat tgattgtttt tattgggccc acctaggcaa	2100
atatgctctt acgtgttattt attaaatgta ctataaaaa ggagttataa aatatgcggc	2160
aaggattttt taaataaact gtcaatttga tagcgggaac aaataattttag atgtcctttt	2220
ttaggaggc ttgtttttt gtacccagtt taagaatacc tttatcatgt gattctaaag	2280
tatccagaga atatctgtat gcattgtata cctatggta tgcataaaaa tcccaatgtat	2340
aaaagtattt atcaactggg tttttatgcc cttttgggtt ttgaatgga ggaaaatcac	2400
atgaaaatta ttaatattgg agtttttagt catgttgatg cagaaaaaac tacctaaca	2460
gaaagcttat tatataacag tggagcgatt acagaattttag gaagcgtgga caaaggta	2520
acgaggacgg ataatacgct ttttagaacgt cagagaggaa ttacaattca gacaggaata	2580
acctcttttca agtggaaaaa tacgaaggtg aacatcatag acacgccagg acatatggat	2640
ttcttagcag aagtataatcg ttcatatca gtttttagatg gggcaattct actgatttct	2700
gcaaaagatg gcgtacaagc acaaactcgt atatttttc atgcacttag gaaaatgggg	2760
attcccacaa tctttttat caataagatt gaccaaaatg gaattgattt atcaacggtt	2820
tatcaggata ttaaagagaa actttctgcc gaaattgtaa tcaaacagaa ggtagaactg	2880
tatcctaata tgtgtgtac gaactttacc gaatctgaac aatgggatac ggtatagag	2940
ggaaacgata acctttttaga gaaatataatg tccggtaaat cattagaagc attggaactc	3000
gaacaagagg aaagcataag atttcagaat tggatctgt tcccttttca tcatgaaat	3060
gcaaaaaatgtat atataggat tgataaccctt atagaatgtt ttactaataa attttattca	3120
tcaacacatc gaggtccgtc tgaactttgc ggaaatgttt tcaaaattga atataaaaaa	3180
aaaagacaac gtcttgata tatacgctt tatagtggag tactacattt acgagattcg	3240
gttagagtat cagaaaaaga aaaaataaaa gttacagaaa tgtatacttc aataaatggt	3300

25938PCT\_CRF\_SequenceListing.txt

gaattatgt	agattgatag	agcttattct	ggagaaattg	ttat	ttgc	aat	atgagtt	3360
ttgaagttaa	atagtgttct	tggagataca	aaactattgc	cacagagaaa	aaagattgaa	3420		
aatccgcacc	ctctactaca	aacaactgtt	gaaccgagta	aacctgaaca	gagagaaatg	3480		
ttgcttgc	ccctttgga	aatctcgat	agtgatccgc	ttctacgata	ttacgtggat	3540		
tctacgacac	atgaaattat	acttcttgc	ttagggaaag	tacaaatgga	agtgattagt	3600		
gcactgtgc	aagaaaagta	tcatgtggag	atagaactaa	aagagcctac	agtcatttat	3660		
atggagagac	cgttaaaaaaa	tgcagaatat	accattcaca	tgcagaatg	gccaaatcct	3720		
ttctggc	ccattgg	tttcttatca	ccgcttccgt	tggaagtgg	aatgcagtat	3780		
gagagctcg	tttctcttgc	atacttaat	caatcatttc	aaaatgcagt	tatggaaagg	3840		
gtacgctatg	gttgcgaaca	aggattat	ggttggaaatg	tgacggattt	taaaatctgt	3900		
tttaagtacg	gtttatacta	tagccctgtt	agtactccag	cagat	ttcgat	3960		
cctattgtac	tggagcaagc	ctttagaaaaa	gctggaaacag	aattgtttaga	gccatatctt	4020		
agttttaag	tttatgcacc	acaggaatat	cttcncgg	catataacga	tgctccaaa	4080		
tatttgcaa	atatcgaaa	tactcaactg	aaaaataatg	agtcattat	tattggagaa	4140		
attcctgctc	gatgtattca	agattatcgc	aatgatttaa	cttttttac	aatggc	4200		
agtgtttgtt	tagcagagct	aaaaggat	caggttacca	ctggcgaacc	tggttgc	4260		
acccgtcg	taaatagtcg	gata	gata	tgttcaataa	aataacttag	4320		
tgcgtttat	gttgttat	aaatatgg	tcttattaa	taagatgaa	tattctttaa	4380		
tatagatttgc	aattaaagt	gaaaggag	gattgttatt	ataaactaca	agtggatatt	4440		
gtgtcctatt	tgtggaaata	aaacaagact	acgaatacga	gtggata	tactaaaaaa	4500		
tttccctta	tacagcccc	aatgtaagaa	cgaaacttta	attaatgtt	aaaaatgaa	4560		
tataataaca	atcaaagagc	cagacgcca	gacgcagagc	cgataattt	agaaatgaaa	4620		
ctctcatctt	atcggtctt	tttgttatac	tgaattttac	tgactagc	cataatattc	4680		
c						4681		

<210> 6  
<211> 1194  
<212> DNA  
<213> Enterobacteri a phage P1

<400> 6	gtgacctgg	acgatcacaa	gaagaattt	gctcgctgg	cgcgagatgg	tgg	ttacacc	60
atcgacagt	atgccc	ccga	gtttaatctt	aaccctaata	ccgcacgtcg	ttat	ctccgt	120
gccttcaaag	aagacaccag	gactacggac	agccgcaagc	caaataagcc	agtca	ggaaag		180
ccactaaaaa	gcatgatcat	tgtactct	aatgatcaac	atgcagg	tg	acatt	gc	240
gctgaaatag	cgaaaaaca	aagagttaat	gccgttgc	gtgccc	cagt	cgagaat	gcg	300
aagcgc	aaaataagc	cat	aatgatcg	tc	agatgatc	atgacgt	gat	360
cacccggac	ct	acgtgatcg	cctgg	gacacc	tgg	atgatgatgg	tgaacg	420

25938PCT\_CRF\_SequenceListing.txt

gaattcgaag ttggcgatta cctgatagat aacgttgaag cgcggaaaggc cgcgccgcgt	480
atgttgcgtc ggtccggggc cgatgttctg gaaaccactc ttctggaaaa gtctcttct	540
catctcctta tgctggagaa cgccagggat acgtgtattc gcctggtgca ggaaatgcgc	600
gatcagcaaa aagacgtatga tgaaggtact ccgcctgaat accgtatcgc gagcatgcta	660
aacagctgtt ccgcgcagat aagcagcctg atcaacacca tttacagcat ccggaataac	720
tatcgaaaag aaagccggga ggcggaaaag cacgctttat ctatggggca agctggcatt	780
gttaagctgg catacgaacg aaagcgtgaa aataactggt cagtgttggaa agcggctgaa	840
ttcatcgagg cgcatggagg aaaagtgcgc cccctgtatgc tggagcaaat caaagccat	900
ctgcgtgctc ctaagaccaa taccgtatgat gaggaaaacc aaacagcatc tggcgttcca	960
tcacttgaag atctggataa aatcgcgca gaacggccg ccagccgcgc cgctgtatgcc	1020
gcattgttggaa ttgagcatcg tagagaagaa attggcata tcgtcgatatac aggtggttat	1080
ggtgatgtcg atgcggaaagg catatcaaac gaagcatggc ttgaacagga tctggacgaa	1140
gacgaggagg aagacgaaga agttacccgc aaactgtacg gggatgtatga ttaa	1194

<210> 7  
 <211> 1194  
 <212> DNA  
 <213> Enterobacteria phage P1

<400> 7	
gtgacctggg acgatcacaa gaagaatttt gctgcctgg cgcgagatgg tggttacacc	60
atcgacagt atgcccggaa gttaatctt aaccctaata ccgcacgtcg ttatctccgt	120
gccttcaaag aagacaccag gactacggac agccgcaagc caaataagcc agtcaggaag	180
ccactaaaaa gcatgtatcat tgcgtactt aatgtatcaac atgcagggtga tcacattgcg	240
gctgaaatag cgaaaaaaca aagagttaat gccgttgtca gtgcccgcagt cgagaatgcg	300
aagcgccaaa ataagcgcat aaatgtatcg tcaagatgatc atgacgtatgc caccgcgc	360
cacggaccc tacgtgtatcg cctggAACGC gacaccctgg atgatgtatgg tgaacgcgtt	420
gaattcgaag ttggcgatta cctgatagat aacgttgaag cgcggaaaggc cgcgccgcgt	480
atgttgcgtc ggtccggggc cgatgttctg gaaaccactc ttctggaaaa gtctcttct	540
catctcctta tgctggagaa cgccagggat acgtgtattc gcctggtgca ggaaatgcgc	600
gatcagcaaa aagacgtatga tgaaggtact ccgcctgaat accgtatcgc gagcatgcta	660
aacagctgtt ccgcgcagat aagcagcctg atcaacacca tttacagcat ccggaataac	720
tatcgaaaag aaagccggga ggcggaaaag cacgctttat ctatggggca agctggcatt	780
gttaagctgg catacgaacg aaagcgtgaa aataactggt cagtgttggaa agcggctgaa	840
ttcatcgagg cgcatggagg aaaagtgcgc cccctgtatgc tggagcaaat caaagccat	900
ctgcgtgctc ctaagaccaa taccgtatgat gaggaaaacc aaacagcatc tggcgttcca	960
tcacttgaag atctggataa aatcgcgca gaacggccg ccagccgcgc cgctgtatgcc	1020

25938PCT\_CRF\_SequenceListing.txt

gcattgtgga	ttgagcatcg	tagagaagaa	attgccata	tcgtcgatac	aggtggttat	1080
ggtgatgtcg	atgcggaagg	catacaaac	gaagcatggc	ttgaacagga	tctggacgaa	1140
gacgaggagg	aagacgaaga	agttacccgc	aaactgtacg	gggatgatga	ttaa	1194
<210> 8						
<211> 525						
<212> DNA						
<213> Artificial Sequence						
<220>						
<223> Description of Artificial Sequence: Synthetic						
polynucleotide						
<400> 8						
atgaacgaaa	aacaaaagag	attcgagat	gaatataaa	tgaatggatg	taatggtaaa	60
aaagcagcaa	tttcagcagg	ttatagtaag	aaaacagcag	agtctttagc	aagtgcattg	120
ttaagaatg	ttaatgttc	ggaatataatt	aaagaacgat	tagaacagat	acaagaagag	180
cgttaatga	gcattacaga	agcttagcg	ttatctgctt	ctattgctag	aggagaacct	240
caagaggctt	acagtaagaa	atatgaccat	ttaaacgatg	aagtggaaaa	agaggttact	300
tacacaatca	caccaactt	tgaagagcgt	cagagatcta	ttgaccacat	actaaaagtt	360
catggtgcgt	atatcgacaa	aaaagaaatt	actcagaaga	atattgagat	taatattaga	420
tctattgacc	acatactaaa	agttcatggt	gcgtatatcg	acaaaaaaga	aattactcag	480
aagaatattg	agattaatat	tggtgagtac	gatgacgaaa	gttaa		525
<210> 9						
<211> 663						
<212> DNA						
<213> Staphylococcus phage 80al pha						
<400> 9						
aattggcagt	aaagtggcag	ttttgatac	ctaaaatgag	atattatgat	agtgtaggat	60
attgactatc	ttactgcgtt	tcccttatcg	caattaggaa	taaaggatct	atgtggggtt	120
gctgattata	gccaatcctt	tttaatttt	aaaaagcgt	tagcgcgaga	gttgggtgta	180
aatgaaatga	acgaaaaaca	aaagagattc	gcagatgaat	atataatgaa	tggatgtaat	240
ggtaaaaaaag	cagcaattt	agcaggttat	agtaagaaaa	cagcagagtc	tttagcaagt	300
cgattgttaa	gaaatgttaa	tgttgcgaa	tatattaaag	aacgattaga	acagatacaa	360
gaagagcgtt	taatgagcat	tacagaagct	ttagcgttat	ctgcttctat	tgctagagga	420
gaacctcaag	aggcttacag	taagaaatat	gaccattaa	acgatgaagt	ggaaaaaagag	480
gttacttaca	caatcacacc	aactttgaa	gagcgtcaga	gatctattga	ccacatacta	540
aaagttcatg	gtgcgtatat	cgacaaaaaa	gaaattactc	agaagaatat	tgagattaat	600
attggtgagt	acgatgacga	aagttaaatt	aaacttaac	aaaccatcta	atgtttcaa	660
cag						663

<210> 10

25938PCT\_CRF\_SequenceListing.txt

<211> 1116

<212> DNA

<213> *Staphylococcus aureus*

<400> 10

tcataaaat	ttaactat	tttctgtgt	actagggtac	aatgaccgt	atcggttata	60
tacttcatta	ctatcagcat	ggctaaacg	ctgtgttatt	accatgatac	ttgcgccatg	120
attgactagc	atagacgcat	ggctatgtct	taactcatga	attacaattc	tagggatgt	180
ctgaccgtct	ggtagttgtt	catctaatac	tttaatgca	gcggtaaacc	aacgatctat	240
agttgattca	ctataagctt	tgaagaatgt	accgaataat	acataatcat	ctttatatac	300
attgtttct	ttgtaccatt	ttaaatattc	tttgatatca	ttcatcatgt	gaacaggtaa	360
gtatatatca	cgtattgctg	ctttgtttt	aggggctgtc	acttcaccgt	gatagtctgt	420
tttgttaata	tgtatgaaat	catcatcata	gttaatatac	cgccatgtga	gggctctaata	480
ttcgcctta	cgtgcaccag	agtaaaacag	tagctaaag	aataactttt	gttgtgtgt	540
agctaaagcc	tcatagaatt	gatgaattt	ttctaatgtc	caatagttca	aacgcttatt	600
tgattctatt	tcaaagttac	ctactagaga	ggctacattt	tgcttttagat	catgaaactt	660
catagcatgg	ttaagtaacg	atactaagaa	cacgtgcatt	ttcttttagt	actctccaga	720
gtgtccctct	tttaacttcg	tattctgaaa	tttcataata	tcttgtgtag	tcatattaaa	780
cacgtccata	gacttaaaat	aggtagcaa	atggttgttt	gtatgtgtct	ttaatgcttt	840
cacactagat	gacttacgac	gtgcagaata	ccactctata	tactcatcta	cgagcttatac	900
aaagggcagt	ttgttatct	gtcctacacc	ctctaactcg	tccataattt	cattacattt	960
cttcaatgcc	tcttacgct	gtttaaagcc	actctttttt	atttctttac	gttgattaaa	1020
tttatcatag	tatttatac	gaaaatagta	tgtaccacgt	ttagcgtctt	tatatatgtt	1080
gtggatagg	tttaagttgt	gttctatggg	aatcac			1116

<210> 11

<211> 10844

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<220>

<221> modified\_base

<222> (8807)...(8807)

<223> a, c, t, g, unknown or other

<400> 11

ctcggccgt	ctttggcgt	tgtcggcct	tcttgcgtat	ctcacgcgt	cctgcggcgg	60
cctgtaggc	aggctcatac	ccctgccc	ccgttttgt	cagccggcgt	gccacggctt	120
ccggcgtctc	aacgcgttt	gagattccca	gctttcggc	caatccctgc	ggtgcataagg	180
cgcgtggctc	gaccgcgttgc	gggctgatgg	tgacgtggcc	cactggcgttgc	cgctccaggg	240

25938PCT\_CRF\_SequenceListing.txt

cctcgtagaa	cgcctgaatg	cgctgtgac	gtgccttgct	gccctcgatg	ccccgttgc	300
gccctagatc	ggccacagcg	gccgcaaacg	tggctggtc	gcgggtcatc	tgcgcttgc	360
tgccgatgaa	ctccttggcc	gacagcctgc	cgtcctgcgt	cagcggcacc	acgaacgcgg	420
tcatgtcg	gctggtttcg	tcacggtgga	tgctggccgt	cacgatgcga	tccgccccgt	480
acttgtccgc	cagccacttgc	tgcccttct	cgaagaacgc	cgcctgctgt	tcttgctgg	540
ccgacttcca	ccattccggg	ctggccgtca	tgacgtactc	gaccgccaac	acagcgtcct	600
tgcgccgctt	ctctggcagc	aactcgcgc	gtcggccat	cgcttcatcg	gtgctgctgg	660
ccgcccagtgc	ctcggtctct	ggcgtcctgc	tggcgtcagc	gttggcgctc	tgcgcgtcgc	720
gttaggcgtg	ctttagactg	gccgccacgt	tgcccatttt	cgcagcttc	ttgcatcgca	780
tgatcgcgt	tgccgcatg	cctgcccctc	cctttggtg	tccaaccggc	tcgacggggg	840
cagcgcaagg	cggtgccctcc	ggcggccac	tcaatgcttgc	agtataactca	ctagactttg	900
cttcgcaaaag	tcgtgaccgc	ctacggcggc	tgcggccccc	tacgggcttgc	ctctccgggc	960
ttcgcctgc	cggtcgctgc	cgctcccttg	ccagccgtg	gatatgtgga	cgatggccgc	1020
gagcggccac	cggctggctc	gctcgctcg	gcccggtggac	aaccctgctg	gacaagctga	1080
tggacaggct	gcccctgccc	acgagcttgc	ccacaggat	tgcccaccgg	ctaccactat	1140
aggcgaatt	ggcggaaaggc	cgtcaaggcc	gcattttggc	ccggcgcgc	ggatccgcta	1200
gctctagacc	tctagaccag	ccaggacaga	aatgcctgc	cttcgctgct	gcccaaggtt	1260
gccgggtgac	gcacaccgtg	gaaacggatg	aaggcacgaa	cccagtggac	ataaggctgt	1320
tcggttcgta	agctgtatgc	caagtagcgt	gcccgttccg	ggcctttga	catgtgactt	1380
tcgttaccct	cgcgtcaaaa	agagtttttgc	cgaaaggaag	cataagtgc	ctgggacgat	1440
cacaagaaga	atttgctcg	cctggcgcga	gatgggtgg	acaccatgc	acagtatgcc	1500
gccgagtttgc	atcttaaccc	taataccgca	cgtcgatatac	tccgtgcctt	caaagaagac	1560
accaggacta	cggacagccg	caagccaaat	aagccagtca	ggaagccact	aaaaagcatg	1620
atcattgtatc	actctaata	tcaacatgc	ggtgatcaca	ttggcgtcgt	aatagcggaa	1680
aaacaaagag	ttaatgccgt	tgtcagtgc	gcagtcgaga	atgcgaagcg	ccaaaataag	1740
cgcataaaatg	atcggtcaga	tgtatcatgc	gtgatcaccc	gcgcaccgc	gaccttacgt	1800
gatgcctgg	aacgcgacac	cctggatgt	gatggtaac	gcttgaatt	cgaagttggc	1860
gattacctga	tagataacgt	tgaagcgcgg	aaggccgcgc	gcgctatgtt	gcgtcggtcc	1920
ggggccgatg	ttctggaaac	cactttctg	gaaaagtctc	tttctcatct	ccttatgctg	1980
gagaacgcca	gggatacgt	tattgcctg	gtgcaggaaa	tgcgcgtatca	gcaaaaagac	2040
gatgatgaag	gtactccgc	tgaataccgt	atcgcgagca	tgctaaacag	ctgttccgc	2100
cagataagca	gcctgatcaa	caccattac	agcatccgga	ataactatcg	aaaagaaagc	2160
cgggaggcgg	aaaagcacgc	tttatctatg	gggcaagctg	gcattgttaa	gctggcatac	2220
gaacgaaagc	gtgaaaataa	ctggtcagtg	ctggaaacgcgg	ctgaattcat	cgaggcgcac	2280

25938PCT\_CRF\_SequenceListi ng. txt

ggaggaaaag	tgccgcccct	gatgctggag	caaataaaag	ccgatctgcg	tgctccctaa	2340
accaataaccg	atgatgagga	aaaccaaaca	gcatctggcg	ctccatcaact	tgaagatctg	2400
gataaaaatcg	cgcgagaacg	ggccgcccagc	cgccgcgctg	atgcccgcatt	gtggatttag	2460
catcgtagag	aagaaattgc	cgcatatcgctc	gatacaggtg	gttatggtga	tgtcgatgcg	2520
gaaggcatat	caaacaaggc	atggcttgaa	caggatctgg	acgaagacga	ggaggaagac	2580
gaagaagtta	cccgcaaaact	gtacggggat	gatgattaat	taaaaaaaacc	cgccccggcg	2640
gttttttta	tctagagcta	gcccgcggc	cgcgcggc	ccttctggc	ctcatggcc	2700
ttccgctcac	tgcccgttt	ccagcactat	agggcgaatt	ggcggaaaggc	cgtcaaggcc	2760
gcatttggc	ccggcgcgcc	ggatccgcta	gctctagact	ggcagggttc	tgagcagatc	2820
gtccaacccg	atctggatcg	ggtcagaaaaa	atttgctcta	ataaatttcg	ttttctaagt	2880
gcaaagaatc	accatttcga	gctggtgatt	gaaggttgat	gcaaatttg	agaaaaaaatg	2940
caacaaacat	tcaatgcgga	tatgaatata	tcaaacccttc	atcaaataatgt	cgatcctca	3000
accactctgc	ccgttatttg	tggtggtaaa	attacgaccg	accgcgctgg	ccgttacaac	3060
cttaatgctc	tacacagagc	gaggggactc	ggtgcccata	aagcgccagc	tcaatggcta	3120
agaacgctgt	cagctaaaca	gctcatcgaa	gagttgaaa	aagaaactat	gcagaattgc	3180
atagtttgcgt	tcacaagcaa	tggaaagcagg	atttcttca	cgactcgtat	aaccggcaaa	3240
ggtcagcagt	ggctgatgaa	gcgattgctt	gatgctggtg	tgctggtacc	tgtcgccgca	3300
acgcgctaac	agacgttagta	agaaccacca	gcattgtaat	gctggctaaa	gtcactttcc	3360
tgagctgtat	aacgatgagc	gattttactt	tttctggcta	tgaattggcc	tgctttgtaa	3420
cacactccgg	tctatccgt	agcgccggc	atatccgtc	gcaatgtgca	aatctcgccg	3480
caacaaccag	tgaataacttc	attcacaagc	ctcaccgcct	gatcgccgca	gaaactggtt	3540
atagccaatc	aaccgtcgtt	cgtgcattcc	gtgaagctgt	aaacaaagga	attctgtctg	3600
tagagattgt	tatcggcgat	caccgtgaac	gtcgcgctaa	cctgtaccgg	tttacaccat	3660
cctttttggc	cttcgcacaa	caagccaaaa	atgcgctgat	agaaagcaaa	ttaaagatct	3720
tttcagcggc	aaccaaggtt	aaagctgttc	tcgctaagac	attggcttta	tttaattttt	3780
tatccacacc	cccatgtcaa	aatgatacc	cctccccc	tcaggatgac	gtggcaataa	3840
agaataagaa	gtcacaagtt	aaaaaaacaa	aaagatcagt	ttccggcggt	gccggaaacaa	3900
ccagcctcaa	aaaattgact	tcatggatcg	ctaaggcaaa	agcaaaggct	gacaatctgc	3960
ggttatccaa	aaaacgcact	caaaaacatg	agttcaagca	gaaagttagag	gcggctgcgc	4020
ggaaatatgc	ttacctgaag	aacaagcg	cgcctgat	tggcgggata	tcaaacttcg	4080
ataacctacc	gcattgcatt	acgtaaacg	aagctttaa	tgccgtttta	gccaaaata	4140
aagataacga	acaatgggg	ataccggcag	gattcagagg	gtaatgaatt	gctctaatta	4200
taaccatgca	tacttcaac	acctcttagt	tgccatgagg	caaactcata	ggtgtcctgg	4260
taagaggaca	ctgttgccaa	aactggacgc	cccattattt	caattaataa	acaactaact	4320

## 25938PCT\_CRF\_SequenceList.sting.txt

gacaattcta	cctaacaata	agtggctaa	aaaaaccgc	ccggcgggt	tttttatct	4380
agagctagcg	gatccggcgc	gccggccct	tctggcctc	atggccttc	cgctcaactgc	4440
ccgctttcca	gccagccttc	gaccacatac	ccaccggctc	caactgcgcg	gcctgcggcc	4500
ttgccccatc	aatttttta	atttctctg	gggaaaagcc	tccggcctgc	ggcctgcgcg	4560
cttcgcttgc	cgttggaca	ccaagtggaa	ggcgggtcaa	ggctcgcgca	gchgaccgcgc	4620
agcggcttgg	ccttgacgcg	ccttggaaacga	cccaaggcta	tgcgagtggg	ggcagtcgaa	4680
ggcgaagccc	gcccgcctgc	cccccgagac	ctgcaggggg	ggggggggcgc	tgaggtctgc	4740
ctcgtgaaga	agggtttgt	gactcatacc	aggcctgaat	cgccccatca	tccagccaga	4800
aagtgaggga	gccacggttg	atgagagctt	tgtttaggt	ggaccagttg	gtgattttga	4860
acttttgctt	tgccacggaa	cggctgcgt	tgtcggaaag	atgcgtgatc	tgatccttca	4920
actcagcaaa	agttcgattt	attcaacaaa	gccgcgtcc	cgtcaagtca	gcgtaatgct	4980
ctgccagtgt	tacaaccaat	taaccaattc	tgattagaaa	aactcatcga	gcatcaaatg	5040
aaactgcaat	ttattcatat	caggattatc	aataccatat	tttggaaaaaa	gccgtttctg	5100
taatgaagga	gaaaactcac	cgaggcagtt	ccataggatg	gcaagatcct	ggtatcggtc	5160
tgcgattccg	actcgccaa	catcaataca	acctattaaat	ttcccctcg	caaaaataag	5220
gttatcaagt	gagaaatcac	catgagtgc	gactgaatcc	ggtgagaatg	gcaaaagctt	5280
atgcatttct	ttccagactt	gttcaacagg	ccagccatta	cgctcgcat	caaaaatcact	5340
cgcataacc	aaaccgttat	tcattcgtga	ttgcgcctga	gcgagacgaa	atacgcgatc	5400
gctgttaaaa	ggacaattac	aaacaggaat	cgaatgcaac	cggcgcagga	acactgccag	5460
cgcataaca	atattttcac	ctgaatcagg	atattttct	aataccttga	atgcttttt	5520
cccggggatc	gcagtggta	gtaaccatgc	atcatcagga	gtacggataa	aatgcttgat	5580
ggtcggaaga	ggcataaatt	ccgtcagcca	gttttagtctg	accatctcat	ctgtaacatc	5640
attggcaacg	ctaccttgc	catgtttcag	aaacaactct	ggcgcatcgg	gcttcccata	5700
caatcgatag	attgtcgac	ctgattgccc	gacattatcg	cgagccatt	tataccata	5760
taaatcagca	tccatgttgg	aatttaatcg	cgccctcgag	caagacgttt	cccggttaat	5820
atggctcata	acacccttg	tattactgtt	tatgttaagca	gacagtttta	ttgttcatga	5880
tgtatatattt	ttatcttgc	caatgttaaca	tcagagattt	tgagacacaa	cgtggtttc	5940
ccccccccc	ctgcagggtcc	cgagcctcac	ggcggcaggt	gccccgggttc	caagggggca	6000
gcccaccc	ggcaaggcc	gaaggccgcg	cagtcgatca	acaagcccg	gagggggcac	6060
tttttgcgg	agggggagcc	gcccgaagg	cgtggggaa	ccccgcaggg	gtgcccattct	6120
ttgggcacca	aagaactaga	tataggcga	aatgcgaaag	actaaaaat	caacaactta	6180
aaaaaggggg	gtacgcaaca	gctcattgc	gcacccccc	caatagctca	ttgcgttaggt	6240
taaagaaaat	ctgtattga	ctgccactt	tacgcaacgc	ataattgtt	tcgcgtcgcc	6300
gaaaagttgc	agctgattgc	gcatggtgc	gcaaccgtgc	ggcaccccta	ccgcatggag	6360

## 25938PCT\_CRF\_SequenceList.txt

ataagcatgg ccacgcagtc cagagaaatc ggcattcaag ccaagaacaa gcccgtcac	6420
tgggtgcaaa cggAACGCAA agcgcattgag gcgtggccg ggcttattgc gagaaaccc	6480
acggcggcaa tgctgctgca tcacctcgta gcgcagatgg gccaccagaa cgccgtggtg	6540
gtcagccaga agacacttgc caagctcatc ggacgttctt tgccggacggt ccaatacgca	6600
gtcaaggact tggtggccga gcgcgtggatc tccgtcgta agctcaacgg ccccgccacc	6660
gtgtcggcct acgtggtcaa tgaccgcgtg gcgtggggcc agcccgccga ccagttgcgc	6720
ctgtcggtgt tcagtgcgc cgtggtggtt gatcacgacg accaggacga atcgctgttg	6780
gggcatggcg acctgcgccc catccgacc ctgtatccgg gcgagcagca actaccgacc	6840
ggcccccggcg aggagccgcc cagccagccc ggcattccgg gcatggaaacc agacctgcca	6900
gccttgaccg aaacggagga atggaaacgg cgccggcagc agcgcctgccc gatgcccgt	6960
gagccgtgtt ttctggacga tggcgagccg ttggagccgc cgacacgggt cacgctgccc	7020
cgccggtagc acttgggttg cgacgcaacc cgtaagtgcg ctgttccaga ctatcgctg	7080
tagccgcctc gccgccttat accttgtctg cctcccgcc ttgcgtcgcc gtgcattggag	7140
ccgggccacc tcgacctgaa tggaaagccgg cgccaccccg ctaacggatt caccgttttt	7200
atcaggctct gggaggcaga ataaatgtatc atatcgtaa ttattacctc cacggggaga	7260
gcctgagcaa actggcctca ggcatttgag aagcacacgg tcacactgct tccggtagtc	7320
aataaaccgg taaaccagca atagacataa gcggcttattt aacgaccctg ccctgaaccg	7380
acgaccgggt cgaatttgct ttcaatttc tgccattcat ccgcttatta tacttattca	7440
ggcgtagcac caggcggtta agggcaccaa taactgcctt aaaaaaattt cgcccccggcc	7500
tgccactcat cgcaactcgctc aggtggact ttccggggaa atgtgcgcgg aaccctatt	7560
tgtttatttt tctaaataca ttcaaataatg tatccgctca tgagacaata accctgataa	7620
atgcttcaat aatattgaaa aaggaagagt atgaagttt gaaatattt ttttcgtat	7680
caaccaccag gtgaaactca taagctaagt aatggatcgc ttgttgcggc ttggatcgcc	7740
ctcagaagag tagggttga tacatattgg accttagaac atcattttac agagttgggt	7800
cttacggaa atttatttgt tgctgcggct aacctgttag gaagaactaa aacattaaat	7860
gttggcacta tgggggttgtt tattccgaca gcacacccag ttgcacagtt agaagacgtt	7920
ttattatttag atcaaataatgtc gaaaggctgt tttaattttt gaaaccgttcg agggctatac	7980
cataaagatt ttgcagtatt tgggtttgat atggaagagt ctcgagcaat tactcaaaat	8040
ttctaccaga tgataatgga aagcttacag acaggaacca ttagctctga tagtattac	8100
attcaatttc ctaagggtga tgtatatccc aaagtgtact caaaaaatgt accaacctgt	8160
atgactgctg agtccgcaag tacgacagaa tggctagcaa tacaagggtt accaatgggtt	8220
cttagttgga ttattggatc taatgaaaaaa aaagcacaga tggaaactcta taatgaaatt	8280
gacacagaat atggtcatga tataatctaaa atagatcatt gtatgactta tattttttct	8340
gttgcgtatg atgcacaaaaa ggcgcacat gtttgcggg agtttctgaa aaattggat	8400

25938PCT\_CRF\_SequenceList.txt

gactcatatg taaatgcgac caatatctt aatgatagca atcaaactcg tggttatgtat 8460  
tatcataaag gtcaatggcg tgattttgtt ttacaaggac atacaaacac caatcgacgt 8520  
gttGattata gcaatggtat taaccctgta ggcactcctg agcagtgtat tgaatcatt 8580  
caacgtgata ttgatgcaac gggatttaca aacattacat gcggatttga agctaattgaa 8640  
actgaagatg aaataattgc ttccatgcga cgctttatga cacaagtcgc tccttctt 8700  
aaagaaccta aataaattac ttatttgata ctagagataa taaggaacaa gttatgaaat 8760  
ttggattatt tttctaaac tttcagaaag atgaaataac atctgangaa acgttggata 8820  
atatggtaaa gactgtcacf ttaattgatt caactaaata tcattttat actgccttg 8880  
ttaatgaaca tcactttca aaaaatggta ttgttggagc acctattacc gcagctggtt 8940  
ttttatttagg gtaacaaat aaattacata ttggttcatt aaatcaagta attaccaccc 9000  
atcaccctgt acgtgttagca gaagaagcca gtttatttga tcaaattgtca gagggacgct 9060  
tcattcttgg ttttagtgac tgcgaaagtg atttcgaaat ggaattttt agacgtcata 9120  
tctcatcaag gcaacaacaa tttgaagcat gctatgaaat aattaatgac gcattaacta 9180  
caggttatttgc tcatccccaa aacgactttt atgattttcc aaaggttca attaatccac 9240  
actgttacag tgagaatgga cctaagcaat atgtatccgc tacatcaaaa gaagtcgtca 9300  
tgtggcagc gaaaaaggca ctgcctttaa catttaagtgg ggaggataat ttagaaacca 9360  
aagaacgcta tgcaattcta tataataaaa cagcacaaca atatggattt gatatttcgg 9420  
atgttgcatt tcaattaact gtaattgcga acttaaatgc tgatagaagt acggctcaag 9480  
aagaagttag agaataactta aaagactata tcactgaaac ttaccctcaa atggacagag 9540  
atgaaaaaat taactgcattt attgaagaga atgcagttgg gtctcatgtat gactattatg 9600  
aatcgacaaa attagcagt gaaaaaacag ggtctaaaaa tattttttaa tccttgaat 9660  
caatgtccga tattaaagat gtaaaagata ttattgat gttgaaccaa aaaatcgaaa 9720  
tgaatttacc ataaagtatgactgttgcattt ttcattaaactg attctgcga catgaaagcc 9780  
atcacagacg gcatgatgaa cctgaatgcg cagcggcatc agcacctgt cgccttgcgt 9840  
ataatatttgc cccatggtaa aaacgggggc gaagaagtttgc tccatattgg ccacgtttaa 9900  
atcaaaactg gtgaaactca cccagggattt ggctgagacg aaaaacatat tctcaataaa 9960  
ccctttaggg aaataggcca ggttttcacc gtaacacgccc acatcttgcg aatataatgtg 10020  
tagaaactgc cgaaatcgt cgtggatttgc actccagagc gatgaaaacg tttcagtttgc 10080  
ctcatggaaa acggtgtac aagggtgaac actatcccat atcaccagct caccgtctt 10140  
cattgccata cgaaattccg gatgagcattt catcaggccg gcaagaatgtt gataaaaggc 10200  
cgataaaaac ttgtgcttac ttttcttac ggtctttaaa aaggccgtaa tatccagctg 10260  
aacggctgg ttataggtac attgagcaac tgactgaaat gcctcaaaaat gttcttacg 10320  
atgcattgg gatatatcaa cggtggata tccagtgattt ttttcttca ttttagcttc 10380  
cttagctcctt gaaaatctcg ataaactcaaa aaatacgccc ggttagtgc tttatcatt 10440

25938PCT\_CRF\_SequenceListing.txt

atggtaaaag ttggAACCTC ttacgtgccc atcaacgtct catttcgccc	aaaaggttggc	10500
ccagggtttc ccggtatcaa cagggacacc aggattatt tattctgcga	agtgtatccc	10560
cgtcacaggt atttattcga agacgaaagg gcctcgat acgcctattt	ttataggtta	10620
atgtcatgat aataatggtt tcttagacgt caggtggcac ttttcgggaa	aatgtgcgcg	10680
cccgcttcc tgctggcgt gggctgtt ctggcgctgg acttcccgct	gttccgtcag	10740
cagctttcg cccacggcct tcatgtatgcg ggcggccttgc	gcctgcataat cccgattcaa	10800
cggccccagg gcgtccagaa cgggcttcag	gcgtcccga aggt	10844

<210> 12

<211> 128

<212> DNA

<213> Staphylococcus aureus

<400> 12

gtctagttaa tgtgtaacgt aacattagct agatTTTTT attcaaaaaa	atatttacaa	60
atatttagaa atttaagtgt aaaagagttt ataaatgatt atattggac	tataatataa	120
ttaaggtc		128

<210> 13

<211> 2769

<212> DNA

<213> Staphylococcus phage 80al pha

<400> 13

attagacaac aaacaagtca ttgaaaattc cgacttatta ttcaaaaaga	aatttgatag	60
cgcagatata caagctaggtaaaaatgg cgataaggta gaagttaaaa	caatcggtta	120
tagaatacac ttttaaatt tataatccgt cttatacgaa gtaaagaagg	tagataaaca	180
atgattaaac aaatactaag actattttc ttactagcaa tgtatgagtt	aggtaagtat	240
gtaactgagc aagtataat tatgatgacg gctaatgatg atgttagaggt	gccgagtgac	300
ttcgcgaagt tgagcgatca gtcagattt atgagggcgg aggtgacgga	gtagatgatg	360
tggtagtca tagcaattat attactagtc atcttattgt ttgggtgtat	gttgcaagct	420
gaacagttaa aaggcgatgt gaaagttaaa gagcgggaga tagagatatt	aagaagtaga	480
ttgagacatt ttgaagatta aaaatattt tatggagggt attcatgact	aaaaagaaat	540
atggattaaa attatcaaca gttcgaaagt tagaagatga gttgtgtat	tatcctaatt	600
atcataagca actcgaagat ttaagaagtg aaataatgac accatggatt	ccaacagata	660
caaataatagg cggggagttt gtaccgtcta atacatcgaa aacagaaatg	gcagtaacta	720
attatcttg tagtatacga agagttaaa tccttgagtt taagagcgct	attgaacgta	780
taatcaacac atcaagttagg aaagaacgca aattcattca agagtattat	tttaataaaa	840
aggaattagt gaaagtttgt gatgacatac acatttctga tagaactgct	catagaatca	900
aaaggaaaat catatctaga ttggcggaaag agtttagggaa agagtgaaat	tggcagtaaa	960
gtggcagttt ttgataccat aatgagata ttatgatagt gtaggatatt	gactatctta	1020

25938PCT\_CRF\_SequenceListing.txt

ctgcgtttcc	cttatcgcaa	ttagaaataa	aggatctatg	tgggttggct	gattatagcc	1080
aatcctttt	taatttaaa	aagcgtatag	cgcgagagtt	ggtggtaaat	gaaatgaacg	1140
aaaaacaaaa	gagattcgca	gatgaatata	taatgaatgg	atgtaatggt	aaaaaagcag	1200
caatttcagc	aggttatagt	aagaaaacag	cagagtctt	agcaagtcga	ttgttaagaa	1260
atgttaatgt	ttcggatat	attaaagaac	gattagaaca	gatacaagaa	gagcgtttaa	1320
tgagcattac	agaagcttta	gcgttatctg	cttctattgc	tagaggagaa	cctcaagagg	1380
cttacagtaa	gaaatatgac	cattaaacg	atgaagtgg	aaaagaggtt	acttacacaa	1440
tcacaccaac	tttgaagag	cgtcagagat	ctattgacca	catactaaaa	gttcatggtg	1500
cgtatatcga	caaaaaagaa	attactcaga	agaatattga	gattaatatt	ggtgagtacg	1560
atgacgaaag	ttaaattaaa	ctttaacaaa	ccatctaattg	tttcaacag	aaacatattc	1620
gaaatactaa	ccaattacga	taacttcact	gaagtacatt	acggtgagg	ttcgagtgg	1680
aagtctcacg	gcgttataca	aaaagttgt	cttaaagcat	tgcaagactg	gaaatatcct	1740
aggcgtatac	tatggcttag	aaaagtccaa	tcaacaatta	aagatagttt	attcgaagat	1800
gtcaaagatt	gtttgataaa	cttcggatt	tggacatgt	gccttggaa	taagactgat	1860
aacaaagtt	aattgccaaa	cggcgcagtt	ttttgttta	aaggattaga	taacccagag	1920
aaaataaaagt	cgataaaagg	catatcagac	atagtcattg	aagaagcgtc	tgaattcaca	1980
ctaaatgatt	acacgcaatt	aacgttgcgt	ttgagggagc	gtaaacacgt	gaataagcaa	2040
atattttga	tgttaaccc	agtatctaa	ctgaattggg	tttataagta	tttcttgaa	2100
catggtaac	caatggaaaa	tgtcatgatt	agacaatcta	gttatcgaga	taataagttt	2160
cttgcgtaaa	tgacacgaca	aaacttagag	ttgttagcaa	atcgtaatcc	agcatattac	2220
aaaatttatg	cgttaggtga	atttctaca	ctagacaaat	tggtttccc	taagtatgaa	2280
aaacgtttaa	taaataaaga	tgagttaga	catttacctt	cttattttgg	attggacttt	2340
ggctacgtt	atgatcctag	tgctttata	cattctaaaa	tagatgtaaa	aaaaaagaag	2400
ttatacatca	ttgaagagta	tgttaaccaa	ggtatgctga	atgatgaaat	agctaatgtc	2460
ataaagcaac	ttggttatgc	taaagaagaa	attacagcag	atagtgcaga	acaaaaaaagt	2520
atagctgaat	taaggaatct	aggcgttaaa	aggattttac	caaccaaaaa	aggaaaggc	2580
tcgggtgtac	aagggttaca	attcttaatg	caatttggaa	tcattgtga	tgaacgttgt	2640
ttcaagacta	ttgaagagtt	tgacaactac	acatggcaaa	aggacaaaga	tacaggtgaa	2700
tataccaaatg	aaccagtaga	tacatacaat	cattgtatcg	attcgttgcg	ttattcagtg	2760
gaacgattc						2769

<210> 14

<211> 10319

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic

## 25938PCT\_CRF\_SequenceListing.txt

polynucleotide

<400> 14	
ggcgccatgg ttaaggccc tttcgaaaa gagttagtaa gttAACAGAA gacgaaccaa	60
aactaaatgg tttAGCAGGA aacttagata aaaaaatgaa tccagaatta tattcagaac	120
aggaacagca acaagaacaa caaaaAGAATC aaaaACGAGA tagAGGTATG cacttataga	180
acatgcattt atGCCGAGAA aacttattgg ttGGAATGGG ctatgttta gctaacttgt	240
tagcgAGTTG gttGGACTTG aattGGGATT aatCCAAAGA aagtACCAAC tcaacaacac	300
ataaAGCCCT gtagGTTCCG accaATAAGG aaattGGAAT aaAGCAATAA aaggAGTTGA	360
agaaatgaaa ttCAAGAGAAAG CCTTGGAGAA ttttATAACA agtaAGTATG tactGGTGT	420
tttagtagTC ttaactgttt accagataat acaaATGCTT aaataaaaaa agacttgatc	480
tgatttagacc aaatCTTTG atagtGTTat attaATAACA aaataaaaaa gagTCGCTCA	540
cGCCCTACCA aagTTGTGA acgACATCAT tcaaAGAAAA aaACACTGAG ttGTTTTAT	600
aatCTTGTat atttagatAT taaACGATAT ttaaATATAc ATCAAGATAT atattGGGT	660
gagCGATTAC ttaaACGAAA ttGAGATTAA ggAGTCGATT ttttATGTAT aaaaACAATC	720
atGCAAATCA ttCAAATCAT ttGAAAATC acgATTAGA caATTTTCTT aaaACCGCT	780
actCTAATAG CCGGTTGGAC gcACATACTG tGTGCAATC tGATCCAAAA ttaAGTTTG	840
atGCAATGAC gatCGTTGGA aatCTCAACC gagACAAACG tcAGGCCCTT tCTAAATTAA	900
tgAGTGTAGA gccccAAATA agactttGGG atattCTCA aacaAGTTT aaAGCTAAAG	960
cacttCAAGA aaaAGTTTAT attGAATATG acaaAGTGAAG AGCAGATAGT tGGGATAGAC	1020
gtaATATGCG tattGAATTtT aatCCAAACA aacttACACG agatGAAATG atttGGTTAA	1080
aacAAATAT aataAGCTAC atGGAAGATG acggTTTAC aagATTAGAT ttagCCTTTG	1140
atTTTGAAGA tgatttgAGT gACTACTATG caatGTCGA taaAGCAGTT aagAAAactA	1200
tttttATGG tcGTAATGGT aagCCAGAAA caaaATATTt tggcGTGAGA gatAGTAATA	1260
gatttATTAG aatttATAAT aaaaAGCAAG AACGTAAGA TAATGCAAGT gCTGAAGTTA	1320
tGTCTGAACA tttatGGCGT gtagAAATCG aacttAAAG AGATATGGT gattACTGGA	1380
atGATTGCTT tagtGATTa catatCTTC aaccAGATTG gaaaACTATC caACGCACTG	1440
cggATAGAGC aatAGTTTT atGTTATTGA gtGATGAAGA AGAATGGGGa aagCTTCACA	1500
gaaATTCTAG aacAAATAT aagaATTGA taaaAGAAAT ttCGCCAGTC gatTTAACGG	1560
acttAAATGAA atCGACTTTA aaAGCGAACG aaaaACAAATT GCAAAACAA atCGATTTT	1620
ggcaACATGA atttAAATTtT tggAAATAGT gtACATATTa atattACTGA acaAAATGA	1680
tatATTTAA ctattCTAAT ttaggAGGAT tttttATGA agtGTCtatt taaaATTTG	1740
ggaATTtAT atGAGGTGAA agaATAATTt ACCCCTATAA actttAGCCA CCTCAAGTAA	1800
agAGGTAAAGG ttGTTAGTT tatATAAAATtTtTAAAGGT ttGTTTATAA gCGTTTATT	1860
ttGGCTTGTtTtCATT ttTTAGTGT attAAATGAA atGGTTTAA atGTTCTTT	1920
acCTGATATT GCAAAATCATT ttaataCTAC tcCTGGAATT ACAAACTGGG taaACACTGC	1980

25938PCT\_CRF\_SequenceListing.txt

atatatgtta	acttttcga	taggaacagc	agtatatgga	aaattatctg	attatataaa	2040
tataaaaaaa	ttgttaatta	ttggtattag	tttgagctgt	cttggttcat	tgattgcttt	2100
tattggcc	acctaggcaa	atatgctctt	acgtgctatt	atthaagtga	ctatttaaaa	2160
ggagttaaa	aatatgcggc	aaggattct	taaataaact	gtcaatttga	tagcggaaac	2220
aaataattag	atgcctttt	ttaggaggc	ttagttttt	gtacccagtt	taagaatacc	2280
tttatcatgt	gattctaaag	tatccagaga	atatctgtat	gctttgtata	cctatggta	2340
tgcataaaaa	tcccagtat	aaaagtattt	atcactggga	tttttatgcc	cttttggggtt	2400
tttgaatgga	ggaaaatcac	atgaaaattt	ttaatattgg	agtttttagct	catgttgatg	2460
cagaaaaaac	tacctaaca	gaaagcttat	tatataacag	tggagcgatt	acagaattag	2520
gaagcgtgga	caaaggtaca	acgaggacgg	ataatacgct	tttagaacgt	cagagaggaa	2580
ttacaattca	gacaggaata	acctctttc	agtggggaaa	tacgaaggtg	aacatcatag	2640
acacgccagg	acatatggat	ttcttagcag	aagtataatcg	ttcattatca	gttttagatg	2700
gggcaattct	actgatttct	gcaaaagatg	gcgtacaagc	acaaactcg	atattatttc	2760
atgcacttag	gaaaatgggg	attccacaa	tctttttat	caataagatt	gaccaaaatg	2820
gaattgattt	atcaacggtt	tatcaggata	ttaaagagaa	actttctgcc	gaaattgtaa	2880
tcaaacagaa	ggtagaactg	tatcctaata	tgtgtgtgac	gaactttacc	aatctgaac	2940
aatgggatac	ggtaatagag	ggaaacgata	accttttaga	gaaatataatg	tccggtaaat	3000
cattagaagc	attggaactc	gaacaagagg	aaagcataag	atttcagaat	tgttctctgt	3060
tccctctta	tcatggaagt	gcaaaaagta	atataggat	tgataacctt	atagaagtt	3120
ttactaataa	attttattca	tcaacacatc	gaggtccgtc	tgaactttgc	gaaatgttt	3180
tcaaaattga	atatacaaaa	aaaagacaac	gtcttgata	tatacgctt	tatagtggag	3240
tactacattt	acgagattcg	gttagagtat	cagaaaaaga	aaaaataaaa	gttacagaaa	3300
tgtatacttc	aataaatggt	gaattatgta	agattgatag	agcttattct	ggagaaattg	3360
ttatttgca	aatgagttt	ttgaagttaa	atagtgtct	tggagataca	aaactattgc	3420
cacagagaaa	aaagattgaa	aatccgcacc	ctctactaca	aacaactgtt	gaaccgagta	3480
aacctgaaca	gagagaaatg	ttgcttgatg	ccctttgga	aatctcagat	agtgatccgc	3540
ttctacgata	ttacgtggat	tctacgacac	atgaaattat	actttctttc	ttaggaaag	3600
tacaaatgga	agtgattagt	gcactgtgc	aagaaaagta	tcatgtggag	atagaactaa	3660
aagagcctac	agtcatat	atggagagac	cgttaaaaaa	tgcagaatat	accattcaca	3720
tcgaagtgcc	gccaaatcct	ttctggcctt	ccattggttt	atctgtatcg	ccgcttccgt	3780
tgggaagtgg	aatgcagtat	gagagctcg	tttctttgg	atacttaat	caatcatttc	3840
aaaatgcagt	tatggaaggg	gtacgctatg	gttgcgaaca	aggattatat	ggttggaaatg	3900
tgacggattg	taaaatctgt	tttaagtacg	gtttatacta	tagccctgtt	agtactccag	3960
cagattttcg	gatgcttact	cctattgtac	tggagcaagc	cttttagaaaa	gctggaacag	4020

25938PCT\_CRF\_SequenceListing.txt

aattgttaga	gccatatctt	agtttaaag	tttatgcacc	acaggaatat	cttcacggg	4080
catataacga	tgctccaaa	tattgtcaa	atatcgaaa	tactcaactg	aaaaataatg	4140
aggcattat	tattggagaa	attcctgctc	gatgtattca	agattatcgc	aatgatttaa	4200
ctttttac	aatggcctt	agtgttggtt	tagcagagct	aaaaggatat	caggttacca	4260
ctggcgaacc	tgttgccag	acccgtcgtc	taaatagtcg	gatagataaa	gtaagatata	4320
tgttcaataa	aataacttag	tgcgtttat	gttgttat	aaatatggtt	tcttattaa	4380
taagatgaaa	tattcttaa	tatagatttgc	aattaaagtgc	gaaaggagga	gattgttatt	4440
ataaaactaca	agtggatatt	gtgtcctagt	tgtggaaata	aaacaagact	acgaatacga	4500
gtggatacta	tacttaaaaa	tttccctta	taacagcccc	aatgtaaagaa	cgaaacttta	4560
attaatgttc	aaaaaatgaa	tataataaca	atcaaagagc	cagacgcca	gacgcagagc	4620
cgataatttgc	agaaatgaaa	ctctcatctt	atcggctctt	tttggttatc	tgaattttac	4680
tgactagcct	tcaatatttc	cgcggccagc	ttactatgcc	attattaagc	ttgtatatac	4740
ggagggttta	ttaattggca	gtaaagtggc	agttttgtat	accttaaatg	agatattatg	4800
atagtgttag	atattgacta	tcgtactgcg	tttccctacc	gcaaatttagg	aataaaggat	4860
ctatgtgggt	tggctgatta	tagccaatcc	tttttaattt	ttaaaaagcg	tatagcgcga	4920
gagttggtgg	taaatgaaat	gaacgaaaaaa	caaaagagat	tcgcagatga	atataatgc	4980
aatggatgt	atggtaaaaa	agcagcaattt	acagtaggtt	atagtaagaa	aacagcagag	5040
tctttagcaa	gtcgattgtt	aagaaatgtt	aatgtttcg	aatatattaa	agaacgatta	5100
gaacaggtac	aagaagagcg	ttaatgagt	attacagaag	ctttagcgtt	atctgcttct	5160
attgctagag	gagaacctca	agaggcttac	agtaagaaat	atgaccattt	aaacgatgaa	5220
gtggaaaaag	aggttactta	cacaatcaca	ccaaacttttgc	aagagcgtca	gagatctatt	5280
gaccacatac	taaaagtaca	tggtgctat	atcgataaaaa	aagaaattac	tcagaagaat	5340
attgagatta	atattggta	gtacgtgac	gaaagttaaa	ttgaacttta	acaaaccgtc	5400
taatgtttc	aatagccgcg	ggggcccaac	acaccaactt	ttgaagagcg	tcagagatct	5460
attgaccaca	tactaaaagt	acatggtgcg	tatatcgata	aaaaagaaat	tactcagaag	5520
aatattgaga	ttaatattgg	tgagtacgt	gacgaaagtt	aaattaaact	ttaacaaacc	5580
gtctaattgtt	ttcaatagcc	gcggggccccc	aacgagccgc	cgcatacgta	agccagcccc	5640
gacacccgccc	aacacccgct	gacgcgcct	gacggcttg	tctgctcccg	gcatccgctt	5700
acagacaagc	tgtgaccgtc	tccggagct	gcatgtgtca	gaggtttca	ccgtcatcac	5760
cgaaacgcgc	gagacgaaag	ggcctcgtga	tacgcctatt	tttatacggtt	aatgtcatga	5820
taataatgg	ttcttagacg	tcaggtggca	cttttcgggg	aaatgtgcgc	ggaaccccta	5880
tttggttatt	tttctaaata	cattcaaata	tgtatccgct	catgagacaa	taaccctgat	5940
aatgcttca	ataatattga	aaaaggaaga	gtatgagtat	tcaacatttc	cgtgtcgccc	6000
ttattccctt	tttgcggca	tttgccttc	ctgttttgc	tcacccagaa	acgctggtga	6060

25938PCT\_CRF\_SequenceListing.txt

aagtaaaaga	tgctgaagat	cagttgggtg	cacgagtggg	ttacatcgaa	ctggatctca	6120
acagcgtaa	gatcctttag	agtttcgcc	ccgaagaacg	tttccaatg	atgagcactt	6180
ttaaagttct	gctatgtggc	gcggatttat	cccgatttga	cgccggcaa	gagcaactcg	6240
gtcggccat	acactattct	cagaatgact	tggttgagta	ctcaccggc	acagaaaagc	6300
atcttacgga	tggcatgaca	gtaagagaat	tatgcagtgc	tgcataacc	atgaggata	6360
acactgcggc	caacttactt	ctgacaacga	tcggaggacc	gaaggagcta	accgctttt	6420
tgcacaacat	gggggatcat	gtaactcgcc	ttgatcgttg	ggaaccggag	ctgaatgaag	6480
ccataccaaa	cgacgaggct	gacaccacga	tgcctgttagc	aatggcaaca	acgttgcgca	6540
aactattaac	tggcgaacta	cttactctag	cttccggca	acaattaata	gactggatgg	6600
aggcggataa	agttgcagga	ccacttctgc	gctcgccct	tccggctggc	tggtttattg	6660
ctgataaattc	tggagccggt	gagcgtgggt	ctcgcggtat	cattgcagca	ctggggccag	6720
atggtaagcc	ctcccgatc	gtagttatct	acacgacggg	gagtcaggca	actatggatg	6780
aacgaaatag	acagatcgct	gagataggtg	cctcaactgat	taagcattgg	taactgtcag	6840
accaagtttta	ctcatatata	ctttagattt	atttaaaact	tcattttaa	tttaaaagga	6900
tctaggtgaa	gatcctttt	gataatctca	tgacaaaat	cccttaacgt	gagtttcgt	6960
tccactgagc	gtcagacccc	gtagaaaaga	tcaaaggatc	ttcttgagat	ctttttttc	7020
tgcgcgtaat	ctgctgcttgc	caaacaaaaa	aaccaccgct	accagcggtg	gttttttgc	7080
cggatcaaga	gctaccaact	cttttccga	aggttaactgg	cttcagcaga	gcfgagatac	7140
caaatactgt	tcttctagtg	tagccgtat	taggccacca	cttcaagaac	tctgttagcac	7200
cgcctacata	cctcgctctg	ctaattctgt	taccagtggc	tgctgccagt	ggcgataagt	7260
cgtgtttac	cgggttggac	tcaagacgt	agttaccgga	taaggcgcag	cggtcggcgt	7320
gaacgggggg	ttcgtgcaca	cagccagct	tggagcgaac	gacctacacc	gaacctgaga	7380
tacctacagc	gtgagctatg	agaaagcgcc	acgcttcccg	aaggagaaaa	ggcggacagg	7440
tatccgtaa	gcggcagggt	cggAACAGGA	gagcgcacga	gggagcttcc	agggggaaac	7500
gcctggatc	ttttagtcc	tgtcggttt	cgcacctct	gacttgagcg	tcgatttttgc	7560
tgatgctcg	cagggggcgc	gagcctatgg	aaaaacgcca	gcaacgcggc	cttttacgg	7620
ttcctggcct	tttgctggcc	tttgctcac	atgttcttc	ctgcgttatac	ccctgattct	7680
gtggataacc	gtattaccgc	ctttagtga	gctggcggt	ctagttatg	tgtAACGTA	7740
cattagctag	atttttttat	tcaaaaaat	atttacaaat	attaggaaat	ttaagtgtaa	7800
aagagttgat	aatgattat	atgggacta	taatataatt	aaggtcgatt	gaattcgta	7860
actaattat	caccaaaaag	gaatagagta	tgaagttgg	aaatatttg	tttgcgtatc	7920
aaccaccagg	tgaaactcat	aagcaagtaa	tggatcgctt	tgttcggc	ggtatcgcc	7980
cagaagaggt	agggtttgat	acatattgga	ccttagaaca	tcattttaca	gagtttggtc	8040
ttacggaaaa	tttatttgtt	gctgcggcta	acctgttagg	aagaactaaa	acattaaatg	8100

25938PCT\_CRF\_SequenceListing.txt

ttggcactat	gggggttgtt	attccgacag	cacacccagt	tcgacagttt	gaagacgttt	8160
tattattaga	tcaaatgtcg	aaaggtcggtt	ttaatttgg	aaccgttcga	gggctataacc	8220
ataaaagattt	tcgagtattt	ggtgttgata	tggaagagtc	tcgagcaatt	actcaaattt	8280
tctaccagat	gataatggaa	agcttacaga	caggaaccat	tagctctgat	agtgattaca	8340
ttcaatttcc	taaggttgat	gtatatccca	aagtgtactc	aaaaaaatgta	ccaacctgta	8400
tgactgctga	gtccgcaagt	acgacagaat	ggctagcaat	acaagggcta	ccaatggttc	8460
tttagttggat	tattggtact	aatgaaaaaa	aagcacagat	ggaactctat	aatgaaattt	8520
cgacagaata	tggtcatgat	atatctaaaa	tagatcattt	tatgacttat	atttggttctg	8580
tttagatgatga	tgcacaaaaag	gcgcaagatg	tttgcggga	gtttctgaaa	aattggtata	8640
actcatatgt	aatgcgacc	aatatctta	atgatagcaa	tcaaactcg	ggttatgatt	8700
atcataaaagg	tcaatggcgt	gatttgttt	tacaaggaca	tacaaacacc	aatcgacgt	8760
ttgattatag	caatggtatt	aacccctgt	gcactcctga	gcagtgtatt	gaaatcattt	8820
aacgtgatat	tgtgcaacg	ggtattacaa	acattacatg	cggatttgaa	gctaattggaa	8880
ctgaagatga	aataattgct	tccatgcgac	gctttatgac	acaagtcgct	cctttcttaa	8940
aagaacctaa	ataaattact	tattgtatac	tagagataat	aaggaacaag	ttatgaaatt	9000
tggattat	tttctaaact	ttcagaaaga	tggaataaca	tctgaagaaa	cgttggataa	9060
taatggtaaag	actgtcacgt	taattgattt	aactaaatat	cattttataa	ctgcctttgt	9120
taatgaacat	cactttcaa	aaaatggat	tgttggagca	cctattaccg	cagctggttt	9180
tttatttaggg	ttaacaataa	aattacatat	tggttcatta	aatcaagtaa	ttaccaccca	9240
tcaccctgta	cgtgtacgag	aagaagccag	tttatttagat	caaattgtcag	agggacgctt	9300
cattcttgg	tttagtgact	gcgaaagtga	tttcgaaatg	gaatttttta	gacgtcatat	9360
ctcatcaagg	caacaacaat	ttgaagcatg	ctatgaaata	attaatgacg	cattaactac	9420
agtttattgc	catccccaaa	acgacttttta	tgattttcca	aaggttcaa	ttaatccaca	9480
ctgttacagt	gagaatggac	ctaagcaata	tgtatccgct	acatcaaag	aagtgcgtat	9540
gtgggcagcg	aaaaaggcac	tgccttaac	gtttaagtgg	gaggataatt	tagaaaccaa	9600
agaacgctat	gcaattctat	ataataaaac	agcacaacaa	tatggtattt	atatttcgga	9660
tgttgcattat	caattaactg	taattgcgaa	cttaaatgct	gatagaagta	cggtcaaga	9720
agaagtgaga	gaataactaa	aagactata	cactgaaact	taccctcaaa	tggacagaga	9780
tgaaaaattt	aactgcatta	ttgaagagaa	tgcagttggg	tctcatgat	actattatga	9840
atcgacaaaa	ttagcagtgg	aaaaaacagg	gtctaaaaat	attttattat	cctttgaatc	9900
aatgtccgat	attaaagatg	taaaagatat	tattgtatg	ttgaaccaaa	aaatcgaaat	9960
gaatttacca	taataaaattt	aaaggcaatt	tctatattt	attgcctttt	tggcgcgcct	10020
attctaattgc	ataataaata	ctgataacat	cttataatttt	gtattatattt	ttgttattatc	10080
gttgacatgt	ataattttga	tatcaaaaac	tgattttccc	tctattattt	tcgagattt	10140

25938PCT\_CRF\_SequenceListing.txt

ttttcttaat tctcttaac aaactagaaa tattgtatat acaaaaaatt ataaataata	10200
gatgaatagt ttaattatag gtgtcatca atcgaaaaag caacgtatct tattaaagt	10260
gcgttgcttt tttctcattt ataaggtaa ataattctca tataatcaagc aaagtgaca	10319

<210> 15  
 <211> 2088  
 <212> DNA  
 <213> Staphylococcus aureus

<400> 15	
tatactacaa atgtagtctt atataaggag gatattgatg aaaaagataa aaattgttcc	60
acttatttta atagttgtag ttgtcgggtt tggtatatat ttttatgctt caaaagataa	120
agaaattaat aatactattg atgcaattga agataaaaat ttcaaaacaag tttataaaga	180
tagcagttat atttctaaaa gcgataatgg tgaagtagaa atgactgaac gtccgataaa	240
aatatataat agtttaggcg ttaaagatata aacattcag gatcgtaaaa taaaaaaagt	300
atctaaaaat aaaaacgag tagatgctca atataaaattt aaaaacaaact acggtacat	360
tgatcgcaac gttcaattta attttgttaa agaagatggt atgtggaagt tagattggaa	420
tcatagcgtc attattccag gaatgcagaa agaccaaagc atacatattt aaaaattttaa	480
atcagaacgt ggtttttttt tagaccgaaa caatgtggaa ttggccaata caggaacagc	540
atatgagata ggcatcggtt caaagaatgt atctaaaaaa gattataaag caatcgctaa	600
agaactaagt atttctgaag actatataa acaacaaatg gatcaaaattt gggtaacaaga	660
tgtatccctt gttccactt aaaccgttaa aaaaatggat gaatatttaa gtgatttcgc	720
aaaaaaattt catcttacaa ctaatgaaac agaaagtctgt aactatcctc tagaaaaagc	780
gacttcacat ctatttagttt atgttggtcc cattaactct gaagaattaa aacaaaaaga	840
atataaaggc tataaagatg atgcagttat tggtaaaaag ggactcgaaa aactttacga	900
taaaaaagctc caacatgaag atggctatcg tgcacaatc gttgacgata atagcaatac	960
aatcgcacat acattaatag agaaaaagaa aaaagatggc aaagatattt aactaactat	1020
tgtatcgaaa gttcaaaaga gtatttataa caacatgaaa aatgattatg gctcaggat	1080
tgctatccac cctcaaacag gtgaattattt agcacttgta agcacacctt catatgacgt	1140
ctatccattt atgtatggca tgagtaacga agaatataat aaattaaccg aagataaaaaa	1200
agaacctctg ctcaacaagt tccagattac aacttcacca ggttcaactc aaaaatattt	1260
aacagcaatg attgggttaa ataacaaaac attagacgtt aaaaacaagtt ataaaatcga	1320
tggtaaaggat tggcaaaaag ataaatcttg ggggtggttac aacgttacaa gatatgaagt	1380
ggtaaatggat aatatcgact taaaacaagc aatagaatca tcagataaca ttttcttgc	1440
tagatgtca ctcgaattttt gcagtaagaa atttggaaaaa ggcgttacaa aacttaggtgt	1500
tggtaaaggat ataccaagtg attatccatt ttataatgct caaatttcaa acaaaaaattt	1560
agataatgaa atatttattt ctgattcagg ttacggacaa ggttacaa aacttaggtgt	1620

25938PCT\_CRF\_SequenceListing.txt

agtacagatc	ctttcaatct	atagcgcatt	agaaaataat	ggcaatatta	acgcaccta	1680
cttattaaaa	gacacgaaaa	acaaagttt	gaagaaaaat	attatttcca	aagaaaatat	1740
caatctatta	actgatggta	tgcaacaagt	cgtaaataaa	acacataaag	aagatattta	1800
tagatcttat	gcaaacttaa	ttggcaaatc	cggtaactgca	gaactcaaaa	tgaaacaagg	1860
agaaaactggc	agacaaattt	ggtggtttat	atcatatgtat	aaagataatc	caaacatgtat	1920
gatggctatt	aatgttaaag	atgtacaaga	taaaggaatg	gctagctaca	atgc当地at	1980
ctcaggtaaa	gtgtatgtat	agctatatga	gaacggtaat	aaaaaatacg	atatacatgtat	2040
ataacaaaac	agtgaagcaa	tccgtaacga	tggttgcttc	actgtttt		2088

<210> 16

<211> 2075

<212> RNA

<213> Staphylococcus aureus

<400> 16

uagucuuuaa	uaaggaggau	auugaugaaa	aagauaaaaa	uuguuccacu	uauuuuuuaaua	60
guuguaguug	ucggguuugg	uauauuuuuu	uaugcuucaa	aagauaaaga	auuuuauuaau	120
acuauugaug	caauugaaga	aaaaaaauuuc	aaacaaguuu	auaaagauag	caguuaauuu	180
ucuaaaagcg	auaaugguga	aguagaaaug	acugaacguc	cgauaaaaau	auauuaauagu	240
uuaggcguua	aagauuaaaa	cauucaggau	cguaaaaauaa	aaaaaguauuc	uaaaaauuaaa	300
aaacgaguag	augcucaaaua	aaaaauuuaaa	acaaacuacg	guaacauuga	ucgcaacguu	360
caauuuuaauu	uuguuaaaga	agaugguaug	uggaaguuaug	auugggauca	uagcgucauu	420
auuccaggaa	ugcagaaaga	ccaaagcaua	cauauugaaa	auuuaaaauc	agaacguggu	480
aaaauuuuag	accgaaacaa	uguggaaauug	gccaaauacag	gaacagcaua	ugagauaggc	540
aucguuccaa	agaauguauc	aaaaaaagau	uauaaagcaa	ucgcuaaaga	acuaaguauu	600
ucugaagacu	auaucaaaca	acaaauggau	caaaaauuggg	uacaagauga	uaccuuucguu	660
ccacuuaaaa	ccguuuaaaa	aauggaugaa	uauuuuaagug	auuucgcaaa	aaaauuuucau	720
cuuacaacua	augaaacaga	aagucguaac	uauccucuag	aaaaagcgac	uucacacuua	780
uuagguuaug	uuggucccau	uaacucugaa	gaauuaaaac	aaaaagaaua	uaaaggcua	840
aaagaugaug	caguauuugg	aaaaaaggga	cucgaaaaac	uuuacgauaa	aaagcuccaa	900
caugaagaug	gcuaucgugu	cacaaucggu	gacgauuaau	gcaauacaau	cgcacauaca	960
uuauuagaga	aaaagaaaaaa	agauggcaaa	gauauucaac	uaacuauuga	ugcuaaaguu	1020
caaaagagua	uuuauaacaa	caugaaaaau	gauuauggcu	cagguacugc	uauccacccu	1080
caaacaggug	aaauuuuagc	acuuguaagc	acaccuucau	augacgucua	uccauuuuug	1140
uauggcauga	guaacgaaga	auauuaauaa	uuuacccgaag	auaaaaaaaga	accucugcuc	1200
aacaaguucc	agauuacaac	uucaccaggu	ucaacucaa	aaaauuuuaac	agcaaugauu	1260
ggguuaauua	acaaaacauu	agacgauaaa	acaaguuaua	aaaucgaugg	aaaaggguugg	1320
caaaaaagaua	aaucuugggg	ugguuacaac	guuacaagau	augaaguggu	aaaugguaau	1380

25938PCT\_CRF\_SequenceListing.txt

aucgacuuua aacaagcaau agaaucauca gauaacuuuu ucuuugcuag aguagcacuc	1440
gaauuaggca guaagaaauu ugaaaaaggc augaaaaaac uagguguugg ugaagauaua	1500
ccaagugauu auccauuuua uaaugcucaa auuucaaaca aaaaauuuaga uaaugaaaua	1560
uuauuagcug auucagguua cggacaaggua gaaauacuga uuaacccagu acagauccuu	1620
ucaaucuua ggcacauua gaaauauggc aauauuaacg caccucacuu auuaaaagac	1680
acgaaaaaca aaguuuggaa gaaaaauuuu auuuuccaaag aaaaauucaa ucuauuaacu	1740
gaugguaugc aacaagucgu aaaaaca cauaagaag auuuuauag aucuuaugca	1800
aacuuauug gcaaaucgg uacugcagaa cuaaaauga aacaaggaga aacuggcaga	1860
caaaauuggu gguuuauauc auugauaaa gauaauccaa acaugaugau ggcuauuaau	1920
guuaaagaug uacaagauaa aggauggcu agcuacaug ccaaaaucuc agguaaagug	1980
uaugaugagc uuaugagaa cgguaauaaa aaauacgaua uagaugaaau acaaaacagu	2040
gaagcaaucc guaacgaugg uugcuucacu guuuu	2075

<210> 17

<211> 2166

<212> DNA

<213> Virofischi

<220>

<221> modified\_base

<222> (1194)..(1194)

<223> a, c, t, g, unknown or other

<400> 17

ggcttaata aacagaatca ccaaaaagga atagagtatg aagtttggaa atatttgg	60
ttcgtatcaa ccaccaggtaa aaactcataa gctaagtaat ggatcgctt gttcggctt	120
gtatcgccctc agaagagtag ggttgatac atattggacc ttagaacatc atttacaga	180
gtttggctt acggaaatt tatttggc tgcggctaac ctgttaggaa gaactaaaac	240
attaaatgtt ggcactatgg gggttgttat tccgacagca cacccagttc gacagttga	300
agacgttttta ttatttagatc aaatgtcgaa aggtcgaaaaatttggaa ccgttcgagg	360
gctataccat aaagattttc gagtatttgg tggatgtatg gaagagtctc gagcaattac	420
tcaaaaatttc taccagatga taatggaaag cttacagaca ggaaccatta gctctgatag	480
tgattacatt caatttccta aggttgatgt atatccaaa gtgtactcaa aaaatgtacc	540
aacctgtatg actgctgagt ccgcaagtac gacagaatgg ctagcaatac aagggttacc	600
aatggttctt agttggatta ttggatctaa tggaaaaaaa gcacagatgg aactctataa	660
tggaaatttgc acagaatatg gtcgtatgtat atctaaataa gatcattgtt tgacttataat	720
ttgttctgtt gatgtatgtt cacaaggc gcaagatgtt tgcggagtttctgaaaa	780
ttggatgtac tcatatgtaa atggcaccat tatcttaat gatagcaatc aaactcgtgg	840
ttatgattat cataaaggc aatggcgttga ttttggatcaaggacata caaacaccaa	900

25938PCT\_CRF\_SequenceListing.txt

tcgacgtgtt	gattatagca	atggtattaa	ccctgttaggc	actcctgagc	agtgtattga	960
aatcattcaa	cgtgatattg	atgcaacggg	tattacaaac	attacatgcg	gatttgaagc	1020
taatggaact	gaagatgaaa	taattgcttc	catgcgacgc	tttatgacac	aagtgcgtcc	1080
tttcttaaaa	gaacctaaat	aaattactta	tttgatacta	gagataataa	ggaacaagtt	1140
atgaaatttg	gattattttt	tctaaacttt	cagaaagatg	gaataacatc	tgangaaacg	1200
ttggataata	tggtaaagac	tgtcacgtt	attgattcaa	ctaaatatca	ttttaatact	1260
gcctttgtta	atgaacatca	ctttcaaaa	aatggtattg	ttggagcacc	tattaccgca	1320
gctggttttt	tattagggtt	aacaaataaa	ttacatattg	gttcattaaa	tcaagtaatt	1380
accacccatc	accctgtacg	tgtacgagaa	gaagccagtt	tattagatca	aatgtcagag	1440
ggacgcttca	ttcttggttt	tagtgactgc	gaaagtgatt	tcgaaatgga	attttttaga	1500
cgtcatatct	catcaaggca	acaacaattt	gaagcatgct	atgaaataat	taatgacgca	1560
ttaactacag	gttattgtca	tccccaaac	gactttatg	atttccaaa	ggtttcaatt	1620
aatccacact	gttacagtga	aatggacct	aagcaatatg	tatccgctac	atcaaaagaa	1680
gtcgtcatgt	gggcagcgaa	aaaggcactg	cctttaacat	ttaagtggga	ggataattta	1740
gaaaccaaag	aacgctatgc	aattctatat	aataaaacag	cacaacaata	tgttattgtat	1800
atttcgatg	ttgatcatca	attaactgta	attgcact	taaatgctga	tagaagtacg	1860
gctcaagaag	aagtgagaga	atacttaaaa	gactatatca	ctgaaactta	ccctcaaatg	1920
gacagagatg	aaaaaaattaa	ctgcattatt	gaagagaatg	cagttgggtc	tcatgatgac	1980
tattatgaat	cgacaaaatt	agcagtggaa	aaaacagggt	ctaaaaatat	tttattatcc	2040
tttgaatcaa	tgtccgatat	taaagatgta	aaagatatta	ttgatatgtt	gaaccaaaaa	2100
atcgaaatga	atttaccata	ataaaattaa	aggcaatttc	tatatttagat	tgcctttta	2160
aatttc						2166

<210> 18

<211> 2143

<212> RNA

<213> *Vi brio fischeri*

<400> 18

aaucacaaa	aaggaauaga	guaugaaguu	uggaaauuu	uguuuuuucgu	aucaaccacc	60
aggugaaacu	cauaagcuua	guauuggauc	gcuuuguucg	gcuugguauc	gccucagaag	120
aguaggguuu	gauacauuu	ggacccuaga	acaucuuuu	acagaguuug	gcuuuacggg	180
aaauuuuuu	guugcugcgg	cuaaccuguu	aggaagaacu	aaaacauuaa	auguuggcac	240
uaugggguuu	guuauuccga	cagcacaccc	aguucgacag	uuagaagacg	uuuuauuuuu	300
agaucaaaug	ucgaaagguc	guuuuuauuu	uggaaccguu	cgagggcuau	accauaaaga	360
uuuucgagua	uuugguguug	auauggaaga	gucucgagca	auuacucaa	auuucuacca	420
gaugauaaug	gaaagcuuac	agacaggaac	cauuagcucu	gauagugauu	acauucaauu	480
uccuaagguu	gauguauauc	ccaaagugua	cuaaaaaaau	guaccaaccu	guaugacugc	540

25938PCT\_CRF\_SequenceList.txt

ugaguccgca	aguacgacag	aauggcuagc	aauacaaggg	cuaccaaugg	uucuuaguug	600
gauuauuggu	acuaaugaaa	aaaaagcaca	gauggaacuc	uauaaugaaa	uugcgacaga	660
auauggucau	gauauaucua	aaauagauca	uuguaugacu	uauauuuguu	cuguugauga	720
ugaugcaca	aaggcgcaag	auguuugucg	ggaguuucug	aaaaauuggu	augacuaua	780
uguaaaugcg	accaauaucu	uuaaugauag	caaucaa	acuacuauaa	cgugguuau	840
agguaugg	cgugauuuug	uuuuacaagg	acauaca	accaaucgac	guguugauua	900
uagcauugg	auuaacccug	uaggcacucc	ugagcagugu	auugaaauca	uucaacguga	960
uauugaugca	acggguauua	caaacauuac	augcgauuu	gaagcuaau	gaacugaaga	1020
ugaaaauuuu	gcuucca	gacgcuuau	gacacaaguc	gcuccuuucu	aaaaagaacc	1080
uaaaauuuu	acuuauuuga	uacuagagau	aauaaggaac	aaguuaugaa	auuuggauua	1140
uuuuuucuaa	acuuuucagaa	agauggaa	acaucugaag	aaacguugga	uaauauggua	1200
aagacuguca	cguaauuuga	uucaacuaaa	uaucuuuuua	auacugccuu	uguuuaugaa	1260
caucacuuu	caaaaaaugg	uauguuugga	gcaccuuua	ccgcagcugg	uuuuuuuuua	1320
ggguuaacaa	auaaauuaca	uuuggu	uuaaaucaag	uaauuaccac	ccaucacccu	1380
guacguguag	cagaagaagc	caguuua	gaucaa	cagagggacg	cuucauucuu	1440
gguuuuagug	acugcgaaag	ugauuucgaa	auggaauuuu	uuagacgu	uaucucau	1500
aggcaacaac	aaauugaagc	augcuau	auaauua	acgcauu	uacagguuau	1560
ugucauccc	aaaacgacuu	uuugauuuu	ccaaagg	cauuua	acacuguuac	1620
agugagaaug	gaccuaagca	auaugua	gcuacau	aagaagucgu	cauguggc	1680
gcgaaaaagg	cacugccuu	aacuuuaag	ugggagg	aa	caaagaacgc	1740
uaugcaauuc	uaauua	aacagcacaa	cauaug	uugauuu	ggaugugau	1800
caucauuua	cuguaauugc	gaacuu	gcugau	guacgg	agaagaagug	1860
agagaauacu	aaaaagacua	uaucacugaa	acuuaccc	aaaugg	agaugaaaaa	1920
auuaacugca	uuauugaaga	gaaugcagu	gggucu	augacua	ugaaucgaca	1980
aaauuagcag	uggaaaaa	aggucua	aaauuuu	uauccuu	aucaaugucc	2040
gauauuaag	auguaaaaga	uaauuaugau	auguuga	aaaaaaucg	aaugauuuua	2100
ccauua	aa	aaaggca	uuuucua	uagauug	ccu	2143

<210> 19

<211> 2294

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic polynucleotide

<400> 19

aaaggcauga aaaaacuugg uaucuucacc aacaccuagc uuuuugaagg aauugaguau

60

## 25938PCT\_CRF\_SequenceList.txt

gaaguuugga	aauuuugguu	guucguauca	accaccaggu	gaaacucua	agcuaaaggc	120
augaaaaaac	uaggugau	ucaccaacac	cuaguuuuuu	caaggaauug	aguaugaagu	180
uuggaaauau	uuguuuuuucg	uaucuaccac	caggugaaac	ucauaagcua	aguaauggau	240
cgcuuuguuc	ggcuugguau	cgcucagaa	gaguaggguu	ugauacauau	uggaccuuag	300
aacaucuuuu	uacagaguuu	ggucuuacgg	gaaauuuuu	uguugcugcg	gcuaaccugu	360
uaggaagaac	aaaaacauua	aauguuggca	cuauggggu	uguuauuucg	acagcacacc	420
caguucgaca	guuagaagac	guuuuauua	uagaucaaau	gucgaaaggu	cguuuuaauu	480
uuggaaccgu	ucgagggcua	uaccauaaag	auuuucgagu	auuugguguu	gauauggaag	540
agucucgagc	aaucacuca	aaucuacc	agaugauaa	gaaagcua	cagacaggaa	600
ccauuagcuc	ugauagugau	uacauucaau	uuccuaaggu	ugauguaau	cccaaagugu	660
acucaaaaaa	uguaccaacc	uguaugacug	cugaguccgc	aaguacgaca	gaauggcuag	720
caauacaagg	gcuaccaaug	guucuuaguu	ggauuauugg	uacuaaugaa	aaaaaagcac	780
agauggaacu	cuauaaugaa	auugcgacag	aaauaugguca	ugauauaucu	aaaauagauc	840
auuguaugac	uuauuuuugu	ucuguugaug	augaugcaca	aaaggcgcaa	gauguuuguc	900
gggaguuucu	gaaaaauugg	uaugacucau	auguaaaugc	gaccaauauc	uuuaaugaua	960
gcaucaaac	ucgugguuau	gauuaucua	aaggucaaug	gcgugauuuu	guuuuacaag	1020
gacauacaaa	caccaaucga	cguguugauu	auagcaugg	uauuaacccu	guaggcacuc	1080
cugagcagug	uuugaaaauc	auucaacgug	auauugaugc	aacggguauu	acaaacauua	1140
caugcggauu	ugaagcuaau	ggaacugaag	augaaaauau	ugcuucca	cgacgcuuua	1200
ugacacaagu	cgcuccuuuc	uuaaaagaac	cuaaaaaaau	uacuuuuuug	auacuagaga	1260
uaauaaggaa	caaguuauga	aauiuggauu	auuuuuuucua	aacuuucaga	aagauggaa	1320
aacaucugaa	gaaacguugg	auaaauuggu	aaagacuguc	acguuaauug	auucaacuaa	1380
auaucuuuu	aaucugccu	uuguuuauga	acaucacuuu	ucaaaaaaaug	guauuguugg	1440
agcaccuuuu	accgcagcug	guuuuuuauu	aggguuaca	aaaaaauuac	auauugguuc	1500
auuaaauc	guaauuacca	cccaucaccc	uguacgugua	gcagaagaag	ccaguuuauu	1560
agauaaaaug	ucagagggac	gcuucuuucu	ugguuuuagu	gacugcgaaa	gugauuuucga	1620
aauggaauuu	uuuagacguc	auaucucauc	aaggcaacaa	cauuugaag	caugcuauga	1680
aauaauuaau	gacgcacuu	cuacagguua	uugucaucc	caaaacgacu	uuuaugauuu	1740
uccaaagguu	ucaauuaauc	cacacuguua	cagugagaau	ggaccuaagc	aauauguauc	1800
cgcuacauca	aaagaagucg	ucaugugggc	agcgaaaaag	gcacugccuu	uaacauuuua	1860
gugggaggau	aaaaaagaaaa	ccaaagaacg	cuaugcauuu	cuauuaauua	aaacagcaca	1920
acaauauggu	auugauauuu	cggauguuga	ucaucaauua	acuguaauug	cgaacuuaaa	1980
ugcugauaga	aguacggcuc	aagaagaagu	gagagaauac	uuaaaagacu	auaucacuga	2040
aacuuacccu	caauggaca	gagaugaaaa	aauuaacugc	auuauugaag	agaaugcagu	2100

25938PCT_CRF_SequenceList.txt		
ugggucuau	gaugacuau	2160
aaaauuuua	uuauccuuug	2220
uauguugaac	caaaaaaucg	2280
uuagauugcc	aaaaaaucg	2294
	aaaugaauuu	
	accauaauaa	
	aauuaaaggc	
	aaauucuaua	
	uuuu	
<210>	20	
<211>	12	
<212>	RNA	
<213>	Artificial Sequence	
<220>		
<223>	Description of Artificial Sequence: Synthetic oligonucleotide	
<220>		
<221>	modified_base	
<222>	(1)..(2)	
<223>	a, c, u, g, unknown or other	
<220>		
<221>	modified_base	
<222>	(7)..(9)	
<223>	a, c, u, g, unknown or other	
<220>		
<221>	modified_base	
<222>	(12)..(12)	
<223>	a, c, u, g, unknown or other	
<220>		
<223>	See specification as filed for detailed description of substitutions and preferred embodiments	
<400>	20	
nnwawgnnnu	un	12
<210>	21	
<211>	13	
<212>	RNA	
<213>	Artificial Sequence	
<220>		
<223>	Description of Artificial Sequence: Synthetic oligonucleotide	
<220>		
<221>	modified_base	
<222>	(1)..(1)	
<223>	a, c, u, g, unknown or other	
<220>		
<221>	modified_base	
<222>	(4)..(7)	
<223>	a, c, u, g, unknown or other	
<220>		
<221>	modified_base	
<222>	(12)..(13)	
<223>	a, c, u, g, unknown or other	
<220>		

25938PCT\_CRF\_SequenceListing.txt

<223> See specification as filed for detailed description of substitutions and preferred embodiments

<400> 21

nagnnnncwu wnn

13

<210> 22

<211> 13

<212> RNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic oligonucleotide

<400> 22

cagauaacau uuu

13