

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
4 January 2007 (04.01.2007)

PCT

(10) International Publication Number
WO 2007/001358 A2

(51) International Patent Classification: Not classified

(21) International Application Number:
PCT/US2005/032328(22) International Filing Date:
9 September 2005 (09.09.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/948,097 22 September 2004 (22.09.2004) US(71) Applicant (for all designated States except US): **ALLERGAN, INC.** [US/US]; 2525 Dupont Drive, Irvine, CA 92612 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **WILLIAMS, Dudley, J.** [US/US]; 11 Dunn Street, Laguna Niguel, CA 92677 (US). **GILMORE, Marcella, A.** [US/US]; 13151 St. Thomas Drive, Santa Ana, CA 92705 (US). **STEWART, Lance, E.** [US/US]; 9 Woodfern, Irvine, CA 92614 (US). **VERHAGEN, Marc** [NL/US]; 14 Thunder Trail, Irvine, CA 92614 (US). **AOKI, Kei, Roger** [US/US]; 2 Ginger Lily Court, Coto de Caza, CA 92679 (US).(74) Agents: **STATHAKIS, Dean, G.** et al.; c/o Allergan, Inc., 2525 Dupont Drive, Irvine, CA 92612 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: FLUORESCENCE POLARIZATION ASSAYS FOR DETERMINING CLOSTRIDIAL TOXIN ACTIVITY



(57) Abstract: The present invention provides a method of determining the presence or activity of a clostridial toxin by (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate which includes a fluorophore; a bulking group; and a clostridial toxin recognition sequence containing a cleavage site that intervenes between the fluorophore and the bulking group; (b) exciting the fluorophore with plane polarized light; and (c) determining fluorescence polarization of the treated substrate relative to a control substrate, where a change in fluorescence polarization of the treated substrate as compared to fluorescence polarization of the control substrate is indicative of the presence or activity of the clostridial toxin.

Fluorescence Polarization Assays for Determining Clostridial Toxin Activity

[001] The present invention relates generally to protease assays, and more specifically, to methods for determining the presence or activity of clostridial toxins such as botulinum toxins and tetanus toxins using fluorescence polarization.

[002] The neuromuscular syndrome of tetanus and the rare but potentially fatal disease, botulism, are caused by neurotoxins produced by bacteria of the genus *Clostridium*. These clostridial neurotoxins are highly potent and specific poisons of neural cells, with the human lethal dose of the botulinum toxins on the order of nanograms. Thus, the presence of even minute levels of botulinum toxins in foodstuffs represents a public health hazard that must be avoided through rigorous testing.

[003] However, in spite of their potentially deleterious effects, low controlled doses of botulinum neurotoxins have been successfully used as therapeutics and for some cosmetic applications. In particular, botulinum toxins have been used in the therapeutic management of a variety of focal and segmental dystonias, strabismus, and other conditions in which a reversible depression of cholinergic nerve terminal activity is desired. Established therapeutic uses of botulinum neurotoxins in humans include, without limitation, treatment of blepharospasm, hemifacial spasm, laryngeal dysphonia, focal hyperhidrosis, hypersalivation, oromandibular dystonia, cervical dystonia, torticollis, strabismus, limb dystonia, occupational cramps and myokymia (Rossetto et al., *Toxicon* 39:27-41 (2001)). As an example, intramuscular injection of spastic tissue with small quantities of botulinum neurotoxin A has been used effectively to treat spasticity due to brain injury, spinal cord injury, stroke, multiple sclerosis and cerebral palsy. Additional possible clinical uses of clostridial neurotoxins are currently being investigated.

[004] Given the potential danger associated with small quantities of botulinum toxins in foodstuffs and the need to prepare accurate pharmaceutical formulations, assays for botulinum neurotoxins presently are employed in the food and pharmaceutical industries. The food industry requires assays for the botulinum neurotoxins to validate new food packaging methods and to ensure food safety. The growing clinical use of the botulinum toxins necessitates accurate assays for botulinum neurotoxin activity for product formulation as well as quality control. In both industries, a mouse lethality test currently is the only acceptable assay for botulinum neurotoxin potency.

[005] Unfortunately, the mouse lethality assay suffers from several drawbacks: cost due to the large numbers of laboratory animals required; lack of specificity; potential for inaccuracy unless large animal groups are used; and sacrifice of animal life. Thus, there is a need for new methods based on convenient synthetic substrates that can complement and reduce the need for the mouse lethality assay. The present invention satisfies this need by providing novel assays for determining the presence or activity of a clostridial toxin and provides related advantages as well.

[006] The present invention provides a method of determining the presence or activity of a clostridial toxin by (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate which includes a fluorophore; a bulking group; and a clostridial toxin recognition sequence containing a cleavage site that intervenes between the fluorophore and the bulking group; (b) exciting the fluorophore with plane polarized light; and (c) determining fluorescence polarization of the treated substrate relative to a control substrate, where a change in fluorescence polarization of the treated substrate as compared to fluorescence polarization of the control substrate is indicative of the presence or activity of the clostridial toxin.

[007] Further provided herein is a method of determining the presence or activity of a clostridial toxin by (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate containing (i) a donor fluorophore; (ii) an acceptor having an absorbance spectrum overlapping the emission spectrum of the donor fluorophore; and (iii) a clostridial toxin recognition sequence containing a cleavage site, where the cleavage site intervenes between the donor fluorophore and the acceptor and where, under the appropriate conditions, resonance energy transfer is exhibited between the donor fluorophore and the acceptor; (b) exciting the donor fluorophore with plane polarized light; and (c) determining fluorescence polarization of the treated substrate relative to a control substrate, where a change in fluorescence polarization of the treated substrate as compared to fluorescence polarization of the control substrate is indicative of the presence or activity of the clostridial toxin.

BRIEF DESCRIPTION OF THE DRAWINGS

[008] Figure 1 shows a schematic of the four steps required for tetanus and botulinum toxin activity in central and peripheral neurons.

[009] Figure 2 shows the subcellular localization and sites of cleavage of SNAP-25, VAMP and syntaxin. VAMP is bound to synaptic vesicle membrane, whereas SNAP-25 and syntaxin are bound to the target plasma membrane. BoNT/A and /E cleave SNAP-25 close to the carboxy-terminus, releasing nine or 26 residues, respectively. BoNT/B, /D, /F, /G and TeNT act on the conserved central portion of VAMP (dotted) and release the amino-terminal portion of VAMP into the cytosol. BoNT/C1 cleaves SNAP-25 close to the carboxy-terminus as well as cleaving syntaxin at a single site near the cytosolic membrane surface. The action of BoNT/B, /C1, /D, /F, /G and TeNT results in release of a large portion of the cytosolic domain of VAMP or syntaxin, while only a small portion of SNAP-25 is released by selective proteolysis of BoNT/A, /C1 or /E.

[0010] Figure 3 shows an alignment of various SNAP-25 proteins. Human SNAP-25 (SEQ ID NO: 1; GenBank accession g4507099; see, also, related human SNAP-25 sequence g2135800); mouse SNAP-25 (SEQ ID NO: 2; GenBank accession G6755588); Drosophila SNAP-25 (SEQ ID NO: 3; GenBank accession g548941); goldfish SNAP-25 (SEQ ID NO: 4; GenBank accession g2133923); sea urchin

SNAP-25 (SEQ ID NO: 5; GenBank accession g2707818) and chicken SNAP-25 (SEQ ID NO: 6; GenBank accession g481202) are depicted.

[0011] Figure 4 shows an alignment of various VAMP proteins. Human VAMP-1 (SEQ ID NO: 7; GenBank accession g135093); human VAMP-2 (SEQ ID NO: 8; GenBank accession g135094); mouse VAMP-2 (SEQ ID NO: 9; GenBank accession g2501081); bovine VAMP (SEQ ID NO: 10; GenBank accession g89782); frog VAMP (SEQ ID NO: 11; GenBank accession g6094391); and sea urchin VAMP (SEQ ID NO: 12; GenBank accession g5031415) are depicted.

[0012] Figure 5 shows an alignment of various syntaxin proteins. Human syntaxin 1A (SEQ ID NO: 13; GenBank accession g15079184), human syntaxin 1B2 (SEQ ID NO: 14; GenBank accession g15072437), mouse syntaxin 1A (SEQ ID NO: 15; GenBank accession g15011853), Drosophila syntaxin 1A (SEQ ID NO: 16; GenBank accession g2501095); C. elegans syntaxin A (SEQ ID NO: 17; GenBank accession g7511662) and sea urchin syntaxin (SEQ ID NO: 18; GenBank accession g13310402) are depicted.

[0013] Figure 6 shows (A) a schematic of plasmid pQBI GFP-SNAP25₍₁₃₄₋₂₀₆₎-6XHIS-C and (B) the nucleic acid and amino acid sequences (SEQ ID NOS: 19 and 20) of pQBI GFP-SNAP25₍₁₃₄₋₂₀₆₎-6XHIS-C.

[0014] Figure 7 shows (A) the absorption spectrum and (B) the excitation (dotted) and emission (bold) spectra of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6C.

[0015] Figure 8 shows (A) the UV-VIS absorption spectrum and (B) the excitation (bold) and emission (dotted) spectra of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6C-Alexa Fluor® 594.

[0016] Figure 9 shows turnover of the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6C-Alexa Fluor® 594 substrate using reduced BoNT/A at various concentrations. The arrow indicates when the reduced toxin complex was added.

[0017] Figure 10 shows turnover of the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6C-Alexa Fluor® 546 substrate using recombinant BoNT/A light chain. The arrow indicates addition of the BoNT/A light chain.

DETAILED DESCRIPTION

[0018] The invention provides novel methods for determining the presence or activity of clostridial toxins including botulinum toxins of all serotypes as well as tetanus toxins. The novel methods of the invention, which rely on a clostridial toxin substrate useful for fluorescence polarization analysis, reduce the need for animal toxicity studies and can be used to analyze crude and bulk samples as well as highly purified dichain or single chain toxins or formulated toxin products. The fluorescence polarization-based methods

of the invention are advantageous in that they are sensitive assays which are robust in terms of interference from background fluorescence present in samples. Furthermore, the novel methods of the invention can be performed as homogeneous solution-phase assays and are amenable to automated high-throughput formats.

[0019] As disclosed herein in Example I, a clostridial toxin substrate was prepared with Alexa Fluor[®] 594 as a fluorophore, green fluorescent protein (GFP) as a bulking group, and a portion of SNAP-25 (residues 134-206) as a clostridial toxin recognition sequence for BoNT/A. The absorption spectrum of the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-Cys protein labeled with Alexa Fluor[®] 594 is shown herein in Figure 8A, and the excitation and emission spectra of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor[®] 594 are shown herein in Figure 8B. As further disclosed herein in Example II, the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor[®] 594 substrate was tested for its utility as a suitable substrate by assaying for the activity of BoNT/A reduced bulk toxin by recording the change in fluorescence polarization over time. As shown in Figure 9, there was a reduction in fluorescence polarization at or shortly after the time the diluted bulk BoNT/A toxin was added, and toxin activity was detected at a concentration of as little as about 50 ng/ml (see panel 9D). These results demonstrate that the presence or activity of clostridial toxins can be determined using synthetic substrates assayed by fluorescence polarization.

[0020] As further disclosed herein, fluorescence polarization can be combined with fluorescence resonance energy transfer to sensitively assay for the presence or activity of a clostridial toxin. As disclosed in Example I, a GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein was site-specifically labeled at the carboxy-terminal cysteine residue with Alexa Fluor[®] 546; the photoselection properties of GFP and Alexa Fluor[®] 546 provide for fluorescence resonance energy transfer (FRET) between the donor fluorophore GFP and the acceptor Alexa Fluor[®] 546. As disclosed in Example III and shown in Figure 10, fluorescence polarization increased upon addition of recombinant BoNT/A light chain. Without wishing to be bound by the following, FRET in the intact substrate leads to an apparent depolarization of Alexa Fluor[®] 546 emission due to the significant angle between the initially selected dipole (GFP) and the dipole which would be selected by direct excitation of Alexa Fluor[®] 546. Upon proteolysis, the FRET effect is abolished, and polarization consequently increases even though rotation of the Alexa Fluor[®] dye is increased. The combination of fluorescence resonance energy transfer with fluorescence polarization enhanced the polarization change upon turnover, increasing the sensitivity of the assay (see Figure 10). These results indicate that fluorescence polarization can be combined with fluorescence resonance energy transfer for enhanced sensitivity in assaying for the presence or activity of a clostridial toxin.

[0021] Based on these findings, the present invention provides a method of determining the presence or activity of a clostridial toxin by (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate which includes a fluorophore; a bulking group; and a clostridial toxin recognition sequence containing a cleavage site that intervenes between the fluorophore and the bulking group; (b) exciting the fluorophore with plane polarized light; and (c) determining

fluorescence polarization of the treated substrate relative to a control substrate, where a change in fluorescence polarization of the treated substrate as compared to fluorescence polarization of the control substrate is indicative of the presence or activity of the clostridial toxin. In one embodiment, the change in fluorescence polarization is a decrease in fluorescence polarization. In another embodiment, step (c) includes determining the change in fluorescence polarization of the treated substrate over time.

[0022] In a method of the invention, a fluorophore can have, without limitation, a fluorescence lifetime of at least 0.5 nanoseconds, or at least 5 nanoseconds, or at least 10 nanoseconds. Any of a variety of fluorophores can be useful in the methods of the invention including, but not limited to, Alexa Fluor® dyes; fluorescein and fluorescein derivatives such as diaminotriazinylamino-fluorescein (DTAF); biarsenic derivatives of fluorescein such as fluorescein arsenical hairpin binding dye (FIAsH™) and red biarsenical dye (ReAsH™); carboxyfluorescein (FAM); Texas Red™; tetramethylcarboxyrhodamine (TMR); carboxy-x-rhodamine (ROX); rhodamine green; Oregon Green 488; BODIPY-TR®; BODIPY-TMR; BODIPY®-FL; Cy3; Cy™3B and Dansyl. In one embodiment, the fluorophore is an Alexa Fluor® dye such as, without limitation, Alexa Fluor® 594. In other embodiments, the fluorophore is FIAsH™ or ReAsH™.

[0023] A variety of bulking groups are useful in the methods of the invention, including, without limitation, fluorescent proteins such as green fluorescent protein. In one embodiment, a method of the invention is practiced such that the change in molecular mass upon cleavage of the clostridial toxin substrate is at least 1000 Da. In a further embodiment, a method of the invention is practiced such that the decrease in fluorescence polarization is at least 5 millipolarization units (mP). In still a further embodiment, a method of the invention is practiced such that the decrease in fluorescence polarization is at least 15 mP.

[0024] A variety of recognition sequences can be included in a clostridial toxin substrate useful in a method of the invention. In one embodiment, the recognition sequence is a BoNT/A recognition sequence such as, without limitation, a BoNT/A recognition sequence containing at least six consecutive residues of SNAP-25, where the six consecutive residues include Gln-Arg, or a peptidomimetic thereof. Such a BoNT/A recognition sequence can include, for example, residues 134 to 206 of SEQ ID NO: 2. A recognition sequence included in a clostridial toxin substrate useful in a method of the invention also can be, without limitation, a BoNT/B recognition sequence. Such a BoNT/B recognition sequence can contain, for example, at least six consecutive residues of VAMP, where the six consecutive residues include Gln-Phe, or a peptidomimetic thereof. In a further embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a BoNT/C1 recognition sequence. Such a BoNT/C1 recognition sequence can contain, without limitation, at least six consecutive residues of syntaxin, where the six consecutive residues include Lys-Ala, or a peptidomimetic thereof. A BoNT/C1 recognition sequence useful in the invention also can contain at least six consecutive residues of SNAP-25, where the six consecutive residues include Arg-Ala, or a peptidomimetic thereof.

[0025] In a further embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a BoNT/D recognition sequence. Such a BoNT/D recognition sequence can contain, for example, at least six consecutive residues of VAMP, where the six consecutive residues include Lys-Leu, or a peptidomimetic thereof. A recognition sequence useful in the invention also can be, for example, a BoNT/E recognition sequence. Such a BoNT/E recognition sequence can include, without limitation, residues 134 to 206 of SEQ ID NO: 2, or can contain at least six consecutive residues of SNAP-25, where the six consecutive residues include Arg-Ile, or a peptidomimetic thereof. In yet another embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a BoNT/F recognition sequence. BoNT/F recognition sequences useful in the invention encompass, without limitation, those having at least six consecutive residues of VAMP, where the six consecutive residues include Gln-Lys, or a peptidomimetic thereof. A recognition sequence included in a clostridial toxin substrate useful in a method of the invention also can be a BoNT/G recognition sequence. Such BoNT/G recognition sequences encompass, without limitation, those having at least six consecutive residues of VAMP, where the six consecutive residues include Ala-Ala, or a peptidomimetic thereof. In still a further embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a TeNT recognition sequence. Such a TeNT recognition sequence can be, without limitation, a sequence containing at least six consecutive residues of VAMP, where the six consecutive residues include Gln-Phe, or a peptidomimetic thereof.

[0026] Any of a variety of clostridial toxin substrates are useful for determining the presence or activity of a clostridial toxin according to a method of the invention. In one embodiment, a clostridial toxin substrate is a peptide or peptidomimetic having at least 100 residues. In another embodiment, a clostridial toxin substrate is a peptide or peptidomimetic having at least 200 residues. Furthermore, any of a variety of samples can be assayed according to a method of the invention including, but not limited to, crude cell lysates, isolated clostridial toxins including isolated clostridial toxin light chains; and formulated clostridial toxin products such as, without limitation, formulated BoNT/A, BoNT/B or BoNT/E toxin products.

[0027] The tetanus and botulinum neurotoxins which can be assayed according to a method of the invention are produced by *Clostridia*. These toxins cause the neuromuscular syndromes of tetanus and botulism, with tetanus toxin acting mainly within the central nervous system and botulinum toxin acting on the peripheral nervous system. Clostridial neurotoxins share a similar mechanism of cell intoxication in which the release of neurotransmitters is blocked. In these toxins, which are composed of two disulfide-linked polypeptide chains, the larger subunit is responsible for neurospecific binding and translocation of the smaller subunit into the cytoplasm. Upon translocation and reduction in neurons, the smaller chain displays peptidase activity specific for protein components involved in neuroexocytosis. The "SNARE" protein targets of clostridial toxins are common to exocytosis in a variety of non-neuronal types; in these cells, as in neurons, light chain peptidase activity inhibits exocytosis.

[0028] Tetanus neurotoxin and botulinum neurotoxins B, D, F, and G specifically recognize VAMP (also known as synaptobrevin), an integral protein of the synaptic vesicle membrane. VAMP is cleaved at distinct bonds depending on the neurotoxin. Botulinum A and E neurotoxins recognize and cleave specifically SNAP-25, a protein of the presynaptic membrane, at two different sites in the carboxy-terminal portion of the protein. Botulinum neurotoxin C cleaves syntaxin, a protein of the nerve plasmalemma, in addition to SNAP-25. The three protein targets of the Clostridial neurotoxins are conserved from yeast to humans although cleavage sites and toxin susceptibility are not necessarily conserved (see below; see, also, Humeau et al., Biochimie 82:427-446 (2000); Niemann et al., Trends in Cell Biol. 4:179-185 (1994); and Pellizzari et al., Phil. Trans. R. Soc. London 354:259-268 (1999)).

[0029] Naturally occurring tetanus and botulinum neurotoxins are produced as polypeptide chains of 150 kDa without a leader sequence. These toxins may be cleaved by bacterial or tissue proteinases at an exposed protease-sensitive loop, generating active dichain toxin. Selective proteolytic cleavage activates the toxins by generating two disulfide-linked chains: an L chain of 50 kDa and an H chain of 100 kDa, which is composed of two domains denoted H_N and H_C. This dichain toxin is more active than unnicked toxin. Naturally occurring clostridial toxins contain a single interchain disulfide bond bridging the heavy chain and light chain; such a bridge is important for neurotoxicity of toxin added extracellularly (Montecucco and Schiavo, Quarterly Rev. Biophysics 28:423-472 (1995)).

[0030] The clostridial toxins appear to be folded into three distinct domains of about 50 kDa which are connected by loops, with each domain having a distinct functional role. As illustrated in Figure 1, the cell intoxication mechanism of the clostridial toxins consists of four distinct steps: (1) binding; (2) internalization; (3) membrane translocation; and (4) enzymatic target modification. The carboxy-terminal domain of the heavy chain (H_C) functions in neurospecific binding, while the amino-terminal domain of the H chain (H_N) functions in membrane translocation from endosome to cell cytoplasm. Following reduction of the disulfide linkage inside the cell, the zinc-endopeptidase activity of the L chain is liberated (Montecucco and Schiavo, *supra*, 1995).

[0031] The amino acid sequences of eight human clostridial neurotoxin serotypes have been derived from the corresponding genes (Niemann, "Molecular Biology of Clostridial Neurotoxins" in Sourcebook of Bacterial Protein Toxins Alouf and Freer (Eds.) pp. 303-348 London: Academic Press 1991). The L chain and H chain are composed of roughly 439 and 843 residues, respectively. Homologous segments are separated by regions of little or no similarity. The most well conserved regions of the L chain are the amino-terminal portion (100 residues) and central region (corresponding to residues 216 to 244 of TeNT), as well as the two cysteines forming the interchain disulfide bond. The 216 to 244 region contains a His-Glu-X-X-His binding motif characteristic of zinc-endopeptidases.

[0032] The clostridial toxin heavy chains are less well conserved than the light chains, with the carboxy-terminal portion of H_C corresponding to residues 1140 to 1315 of TeNT the most variable. This is

consistent with the involvement of the H_C domain in binding to nerve terminals and the fact that different neurotoxins appear to bind different receptors.

[0033] Comparison of the nucleotide and amino acid sequences of the clostridial toxins indicates that they derive from a common ancestral gene. Spreading of these genes may have been facilitated by the fact that the clostridial neurotoxin genes are located on mobile genetic elements. As discussed further below, sequence variants of the seven botulinum toxins are known in the art. See, for example, Humeau et al., *supra*, 2000.

[0034] As discussed above, natural targets of the clostridial neurotoxins include VAMP, SNAP-25, and syntaxin. VAMP is associated with the synaptic vesicle membrane, whereas SNAP-25 and syntaxin are associated with the target membrane (see Figure 2). BoNT/A and BoNT/E cleave SNAP-25 in the carboxy-terminal region, releasing nine or twenty-six amino acid residues, respectively, and BoNT/C1 also cleaves SNAP-25 near the carboxy-terminus. The botulinum serotypes BoNT/B, BoNT/D, BoNT/F and BoNT/G, and tetanus toxin, act on the conserved central portion of VAMP, and release the amino-terminal portion of VAMP into the cytosol. BoNT/C1 cleaves syntaxin at a single site near the cytosolic membrane surface. Thus, BoNT/B, BoNT/C1, BoNT/D, BoNT/F, BoNT/G or TeNT proteolysis results in release of a large portion of the cytosolic domain of VAMP or syntaxin, while only a small portion of SNAP-25 is released by BoNT/A, BoNT/C1 or BoNT/E cleavage (Montecucco and Schiavo, *supra*, 1995).

[0035] Naturally occurring SNAP-25, a protein of about 206 residues lacking a transmembrane segment, is associated with the cytosolic surface of the nerve plasmalemma (Figure 2; see, also, Hodel et al., Int. J. Biochemistry and Cell Biology 30:1069-1073 (1998)). In addition to homologs highly conserved from *Drosophila* to mammals, SNAP-25-related proteins also have been cloned from yeast. SNAP-25 is required for axonal growth during development and may be required for nerve terminal plasticity in the mature nervous system. In humans, two isoforms are differentially expressed during development; isoform a is constitutively expressed during fetal development, while isoform b appears at birth and predominates in adult life. SNAP-25 analogues such as SNAP-23 also are expressed outside the nervous system, for example, in pancreatic cells.

[0036] Naturally occurring VAMP is a protein of about 120 residues, with the exact length depending on the species and isotype. As shown in Figure 2, VAMP contains a short carboxy-terminal segment inside the vesicle lumen while most of the molecule is exposed to the cytosol. The proline-rich amino-terminal thirty residues are divergent among species and isoforms while the central portion of VAMP (residues 30 to 96), which is rich in charged and hydrophilic residues and includes known cleavage sites, is highly conserved. VAMP colocalizes with synaptophysin on synaptic vesicle membranes.

[0037] A variety of species homologs of VAMP are known in the art including human, rat, bovine, *Torpedo*, *Drosophila*, yeast, squid and *Aplysia* homologs. In addition, multiple isoforms of VAMP have been identified including VAMP-1, VAMP-2 and cellubrevin, and forms insensitive to toxin cleavage have been identified in non-neuronal cells. VAMP appears to be present in all vertebrate tissues although the distribution of VAMP-1 and VAMP-2 varies in different cell types. Chicken and rat VAMP-1 are not cleaved by TeNT or BoNT/B. These VAMP-1 homologs have a valine in place of the glutamine present in human and mouse VAMP-1 at the TeNT or BoNT/B cleavage site. The substitution does not affect BoNT/D, /F or /G, which cleave both VAMP-1 and VAMP-2 with similar rates.

[0038] Syntaxin is located on the cytosolic surface of the nerve plasmalemma and is membrane-anchored via a carboxy-terminal segment, with most of the protein exposed to the cytosol. Syntaxin colocalizes with calcium channels at the active zones of the presynaptic membrane, where neurotransmitter release takes place. In addition, syntaxin interacts with synaptotagmin, a protein of the SSV membrane, that forms a functional bridge between the plasmalemma and the vesicles. A variety of syntaxin isoforms have been identified. Two isoforms of slightly different length (285 and 288 residues) have been identified in nerve cells (isoforms 1A and 1B), with isoforms 2, 3, 4 and 5 expressed in other tissues. The different isoforms have varying sensitivities to BoNT/C1, with the 1A, 1B, 2 and 3 syntaxin isoforms cleaved by this toxin.

[0039] The methods of the invention rely, in part, on the use of fluorescence polarization. According to the theory of fluorescence polarization, when a fluorescently labeled molecule is excited with plane polarized light, it emits lights that has a degree of polarization which is inversely proportional to its molecular rotation. As a consequence, for large fluorescently labeled molecules, which remain relatively stationary during their excited state (about 4 ns for fluorescein), polarization remains relatively constant between excitation and emission. In contrast, small fluorescently labeled molecules rotate rapidly during the excited state, such that polarization of the light changes significantly between excitation and emission. Therefore, as a generalization, small molecules have low polarization values, and large molecules have high polarization values. See, for example, Weber, "Polarization of the Fluorescence of Solutions" in Fluorescence & Phosphorescence Analysis pages 217-241 Wiley Interscience (1996), and Jameson and Seifried, Methods Enzym. 19:222-233 (1999).

[0040] Fluorescence polarization assays are homogeneous in that they do not require a separation step and do not require attachment of substrate to an immobilized phase. Furthermore, polarization values can be measured repeatedly. In addition, fluorescence polarization is a sensitive technique which can be used to measure polarization values of fluorophores from low picomolar and micromolar levels. Polarization is also independent of fluorescence intensity.

[0041] Fluorescence anisotropy (commonly denoted as "r" or sometimes "A") is an alternative definition of how a plane of polarized light changes between excitation and emission with a rotating fluorophore.

Fluorescence polarization and anisotropy are well known in the art as described in Lundblad et al., Mol. Endocrin. 10:607-612 (1996); Nasir et al., Comb. Chem. High Throughput Screen. 2:177-190 (1999); Sittampalam et al., Curr. Opin. Chem. Biol. 1:384-391 (1997); Thompson et al., Biotechniques 32:34-40 (1997); Lakowicz et al., J. Biomol. Screen. 5:123-132 (2000); and Fernandes, Curr. Opin. Chem. Biol. 2:597-603 (1998).

[0042] In particular, fluorescence polarization (P) and anisotropy (r) are defined as follows:

$$\text{Polarization} = P = \frac{I_{\text{Vertical}} - I_{\text{Horizontal}}}{I_{\text{Vertical}} + I_{\text{Horizontal}}}$$

and

$$\text{Anisotropy} = r = \frac{I_{\text{Vertical}} - I_{\text{Horizontal}}}{I_{\text{Vertical}} + 2I_{\text{Horizontal}}}$$

[0043] where I_{Vertical} is the intensity of the emission light parallel to the excitation light plane and $I_{\text{Horizontal}}$ is the intensity of the emission light perpendicular to the excitation light plane. P and r , being ratios of light intensities, are dimensionless. Experimental data can be expressed in millipolarization units, where 1 polarization unit = 1000 mP units, or in millianisotropy units, where 1 anisotropy unit = 1000 mA units.

[0044] The formulae to interconvert polarization and anisotropy are as follows:

$$P = \frac{3r}{(2+r)} \quad \text{and} \quad r = \frac{2P}{(3-P)}$$

[0045] Fundamentally, polarization is a relationship of fluorescence lifetime and how fast a fluorophore rotates in the time between excitation and emission. The principal factors controlling rotation are molar volume (V), absolute temperature (T), and viscosity (η). The rotational correlation time (Θ) and the rotational relaxation time (ρ_o) are taken from the work of Perrin and Weber. In particular, the rotational correlation time (Θ) is taken from the Perrin equation as follows:

$$\left(\frac{1}{P} - \frac{1}{3} \right) = \left(\frac{1}{P_o} - \frac{1}{3} \right) * \left(1 + T/\Theta \right)$$

and is defined as:

Rotational Correlation Time

$$(\Theta) = \frac{\eta V}{RT}$$

[0046] Furthermore, the rotational relaxation time (ρ_o) is taken from the Perrin/Weber equation (Perrin, J. Phys. Rad. 7:390-401 (1926)), as follows:

$$\left(\frac{1}{P} - \frac{1}{3} \right) = \left(\frac{1}{P_o} - \frac{1}{3} \right) * \left(1 + 3\tau/P \right)$$

and is defined as:

Rotational Relaxation Time

$$(\rho_o) = \frac{3\eta V}{RT}$$

where R is the gas constant, τ is the fluorescence lifetime, P is the polarization, and P_o is the limiting polarization.

[0047] From the above, it can be seen that, where lifetime, viscosity, and temperature are held constant, the molecular volume (and thus the polarization or anisotropy) determines the rotation. The larger the molecular volume, the slower the molecule rotates and the higher the polarization and anisotropy values. Furthermore, as is evident from the equations above, the rotational relaxation time will be exactly three times longer than the rotational correlation time.

[0048] A method of the invention relies on a clostridial toxin substrate which includes, in part, a fluorophore. As used herein, the term "fluorophore" means a molecule that, when irradiated with light of a certain wavelength, emits light, also denoted fluorescence, of a different wavelength. The term fluorophore is synonymous in the art with the term "fluorochrome."

[0049] Fluorophores useful in the invention, as well as donor fluorophores which are discussed further below, include those having fluorescence lifetimes suitable for fluorescence polarization analysis. Useful fluorophores include, without limitation, Alexa Fluor[®] dyes; fluorescein and fluorescein derivatives such as diaminotriazinylamino-fluorescein (DTAF); biarsenic derivatives of fluorescein such as fluorescein arsenical hairpin binding dye (FIAsH[™]) and red biarsenical dye (ReAsH[™]); carboxyfluorescein (FAM); Texas Red[™]; tetramethylcarboxyrhodamine (TMR); carboxy-x-rhodamine (ROX); rhodamine green; Oregon Green 488; BODIPY-TR[®]; BODIPY-TMR; BODIPY[®]-FL; Cy3, Cy[™]3B and Dansyl. Additional fluorophores suitable for fluorescence polarization are known in the art, including, but not limited to, long-wavelength fluorophores such as BODIPY-TMR and BODIPY-TR[®] (Molecular Probes), which tend to minimize assay interference, and pH insensitive fluorophores such as BODIPY-FL. See, for example, Owicki, J. Biomol. Screening 5:297-306 (2000); Burke et al., Comb. Chem. & High Throughput Screen. 6:183-194 (2003); and Jameson and Croney, Comb. Chem. & High Throughput Screen. 6:167-176 (2003). A variety of fluorophores and donor fluorophores useful for fluorescence polarization are commercially available from various sources such as Molecular Probes (Eugene, Oregon) and Amersham Pharmacia Biotech (Piscataway, New Jersey). One skilled in the art understands that these as well as

other fluorophores suitable for fluorescence polarization are known in the art and can be useful in the methods of the invention.

[0050] As used herein, the term "bulking group" means a moiety having sufficient hydrodynamic volume such that, upon cleavage of a clostridial toxin substrate into which the bulking group is incorporated, there is a change in polarization of at least 3 millipolarization units (mP).

[0051] Any of a variety of moieties can be useful as a bulking group in a method of the invention including physical, chemical and biological moieties which can be covalently or non-covalently incorporated into a clostridial toxin substrate. In one embodiment, the bulking group is expressed as a fusion protein with another component of the clostridial toxin substrate. Bulking groups useful in the invention encompass natural and man-made moieties and further encompass, without limitation, inert moieties as well as those with biological or other activity. A bulking group useful in the invention can be, without limitation, a moiety having a size of greater than 1000 Da. A bulking group useful in the invention also can be, without limitation, a moiety having a size of greater than 2 kDa, 3 kDa, 4 kDa, 5 kDa, 10 kDa, 15 kDa, 20 kDa, 25 kDa, 30 kDa, 35 kDa or 40 kDa. See, also, Mattison et al., Application Note for Protein Solutions Inc. February 2001. One skilled in the art understands that a fluorophore with a suitable lifetime will be selected depending, in part, on the size of the bulking group.

[0052] A variety of bulking groups can be useful in the invention. As non-limiting examples, a bulking group useful in the invention can be an inert or active protein, peptide or peptidomimetic; an antibody; organic chemical; latex or other bead; or moiety such as streptavidin. Additional bulking groups useful in the invention encompass, without limitation, phage and other viruses; cells; liposomes; polymeric and non-polymeric matrices; gold and other particles; and microdevices and nanodevices. As non-limiting examples, a bulking group useful in the invention can be a fluorescent protein such as GFP or BFP, or a fragment thereof; a protein useful for affinity purification such as glutathione-S-transferase (GST) or maltose-binding protein (MBP); or an antibody such as, without limitation, an anti-FLAG, anti-hemagglutinin (HA) or anti-myc antibody. Streptavidin also can be a bulking group useful in the invention. As a non-limiting example, a biotinylation sequence can be covalently included in a clostridial toxin substrate, providing for association with streptavidin; enzymatic cleavage can be detected by following the fluorescence polarization change upon addition of streptavidin as described in Levine et al., "Measurement of specific protease activity utilizing fluorescence polarization, " Anal. Biochem. 247:83-88 (1997).

[0053] A clostridial toxin substrate useful in the invention contains a cleavage site that "intervenes" between a fluorophore and a bulking group. Thus, the cleavage site is positioned in between the fluorophore and the bulking group such that proteolysis at the cleavage site results in a first cleavage product containing the fluorophore and a second cleavage product containing the bulking group. It is

understood that all or only a portion of the clostridial toxin recognition sequence may intervene between the fluorophore and the bulking group.

[0054] A clostridial toxin substrate useful in the invention contains, in part, a clostridial toxin recognition sequence which includes a cleavage site. By definition, a clostridial toxin substrate is susceptible to cleavage by at least one clostridial toxin under conditions suitable for clostridial toxin protease activity.

[0055] As used herein, the term "clostridial toxin recognition sequence" means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a clostridial toxin under conditions suitable for clostridial toxin protease activity. A variety of clostridial toxin recognition sequences are discussed herein below.

[0056] In particular embodiments, a clostridial toxin substrate useful in the invention is a peptide or peptidomimetic having a defined length. A clostridial toxin substrate can be, for example, a peptide or peptidomimetic having at least 50, at least 100, at least 150, at least 200, at least 250, at least 300, at least 350, at least 500, at least 600, at least 700, at least 800 or at least 900 residues. In other embodiments, a clostridial toxin substrate has at most 20 residues, at most 30 residues, at most 40 residues, at most 50 residues, at most 60 residues, at most 70 residues, at most 80 residues, at most 90 residues, at most 100 residues, at most 150 residues, at most 200 residues, at most 250 residues, at most 300 residues, at most 350 residues or at most 400 residues.

[0057] It is understood that a clostridial toxin substrate useful in the invention optionally can include one or more additional components. As a non-limiting example, a flexible spacer sequence such as GGGGS (SEQ ID NO: 21) can be included in a clostridial toxin substrate useful in the invention. A useful clostridial toxin substrate further can include, without limitation, one or more of the following: an affinity tag such as HIS6; biotin or a biotinylation sequence; or an epitope such as FLAG, hemagglutinin (HA), c-myc, or AU1; an immunoglobulin hinge region; an N-hydroxysuccinimide linker; a peptide or peptidomimetic hairpin turn; or a hydrophilic sequence or another component or sequence that, for example, facilitates purification or promotes the solubility or stability of the clostridial toxin substrate.

[0058] As discussed further below, it is understood that the methods of the invention are applicable to crude samples as well as highly purified dichain and single chain toxins. As non-limiting examples, a method of the invention can be useful to determine the presence or activity of a clostridial toxin in a food or beverage sample; to assay a sample from a human or animal, for example, exposed to a clostridial toxin or having one or more symptoms of a clostridial toxin; to follow activity during production and purification of clostridial toxin; or to assay formulated clostridial toxin products such as pharmaceuticals or cosmetics.

[0059] A variety of samples are useful in the methods of the invention. As used herein, the term "sample" means any biological matter that contains or potentially contains an active clostridial toxin. Thus, the term sample encompasses, but is not limited to, purified or partially purified clostridial toxin; recombinant single chain or dichain toxin with a naturally or non-naturally occurring sequence; recombinant clostridial toxin with a modified protease specificity; recombinant clostridial toxin with an altered cell specificity; chimeric toxin containing structural elements from multiple clostridial toxin species or subtypes; bulk toxin; formulated toxin product; cells or crude, fractionated or partially purified cell lysates, for example, engineered to include a recombinant nucleic acid encoding a clostridial toxin; bacterial, baculoviral and yeast lysates; raw, cooked, partially cooked or processed foods; beverages; animal feed; soil samples; water samples; pond sediments; lotions; cosmetics; and clinical formulations. It further is understood that the term sample encompasses tissue samples, including, without limitation, mammalian tissue samples, livestock tissue samples such as sheep, cow and pig tissue samples; primate tissue samples; and human tissue samples. Such samples encompass, without limitation, intestinal samples such as infant intestinal samples, and tissue samples obtained from a wound.

[0060] As discussed further below, a variety of conditions suitable for clostridial toxin protease activity are useful in the methods of the invention. For example, conditions suitable for clostridial toxin protease activity can be provided such that at least 10% of the substrate is cleaved. Similarly, conditions suitable for clostridial toxin protease activity can be provided such that at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 95% of the clostridial toxin substrate is cleaved, or such that 100% of the clostridial toxin substrate is cleaved. In one embodiment, the conditions suitable for clostridial toxin protease activity are selected such that the assay is linear. In another embodiment, conditions suitable for clostridial toxin protease activity are provided such that at least 90% of the clostridial toxin substrate is cleaved. In a further embodiment, conditions suitable for clostridial toxin protease activity are provided such that at most 25% of the clostridial toxin substrate is cleaved. In yet further embodiments, conditions suitable for clostridial toxin protease activity are provided such that at most 5%, 10%, 15% or 20% of the clostridial toxin substrate is cleaved.

[0061] In the methods of the invention, the clostridial toxin substrate can be treated with a sample in solution phase. As used herein in reference to a clostridial toxin substrate, the term "in solution phase" means that the substrate is soluble and, during proteolysis, is not constrained or immobilized on a solid support such as a column or dish.

[0062] In the methods of the invention, a sample is treated with a clostridial toxin substrate under conditions suitable for clostridial toxin protease activity. Exemplary conditions suitable for clostridial toxin protease activity are well known in the art, and further can be determined by routine methods. See, for example, Hallis et al., *J. Clin. Microbiol.* 34:1934-1938 (1996); Ekong et al., *Microbiol.* 143:3337-3347 (1997); Shone et al., WO 95/33850; Schmidt and Bostian, *supra*, 1995; Schmidt and Bostian, *supra*, 1997; Schmidt et al., *supra*, 1998; and Schmidt and Bostian, U.S. Patent No. 5,965,699. It is understood

that conditions suitable for clostridial toxin protease activity can depend, in part, on the specific clostridial toxin type or subtype being assayed and the purity of the toxin preparation. Conditions suitable for clostridial toxin protease activity generally include a buffer, such as HEPES, Tris or sodium phosphate, typically in the range of pH 5.5 to 9.5, for example, in the range of pH 6.0 to 9.0, pH 6.5 to 8.5 or pH 7.0 to 8.0. Conditions suitable for clostridial toxin protease activity also can include, if desired, dithiothreitol, β -mercaptoethanol or another reducing agent, for example, where a dichain toxin is being assayed (Ekong et al., *supra*, 1997). In one embodiment, the conditions include DTT in the range of 0.01 mM to 50 mM; in other embodiments, the conditions include DTT in the range of 0.1 mM to 20 mM, 1 to 20 mM, or 5 to 10 mM. If desired, an isolated clostridial toxin or sample can be pre-incubated with a reducing agent, for example, with 10 mM dithiothreitol (DTT) for about 30 minutes prior to addition of clostridial toxin substrate.

[0063] Clostridial toxins are zinc metalloproteases, and a source of zinc, such as zinc chloride or zinc acetate, typically in the range of 1 to 500 μ M, for example, 5 to 10 μ M can be included, if desired, as part of the conditions suitable for clostridial toxin protease activity. One skilled in the art understands that zinc chelators such as EDTA generally are excluded from a buffer for determining the activity of a clostridial toxin.

[0064] Conditions suitable for clostridial toxin protease activity can optionally include a detergent such as TWEEN-20, which can be used, for example, in place of bovine serum albumin. TWEEN-20 can be provided, for example, in the range of 0.001% to 10% (v/v), or in the range of 0.01% to 1.0% (v/v). As a non-limiting example, TWEEN-20 can be included at a concentration of 0.1 % (v/v).

[0065] Conditions suitable for clostridial toxin protease activity also can include, if desired, bovine serum albumin (BSA) or another agent which acts as a protein stabilizer, solubilizing agent or blocker of surface loss. As an example, when included, BSA typically is provided in the range of 0.1 mg/ml to 10 mg/ml. In one embodiment, BSA is included at a concentration of 1 mg/ml. See, for example, Schmidt and Bostian, *supra*, 1997. In another embodiment, BSA is included at a concentration of 0.1% (w/v).

[0066] The amount of clostridial toxin substrate can be varied in a method of the invention. A clostridial toxin substrate can be supplied, for example, at a concentration of 1 μ M to 500 μ M, 1 μ M to 50 μ M, 1 μ M to 30 μ M, 5 μ M to 20 μ M, 50 μ M to 3.0 mM, 0.5 mM to 3.0 mM, 0.5 mM to 2.0 mM, or 0.5 mM to 1.0 mM. The skilled artisan understands that the concentration of clostridial toxin substrate or the amount of sample can be limited, if desired, such that the assay is linear. In one embodiment, a method of the invention relies on a clostridial toxin substrate concentration of less than 100 μ M. In further embodiments, a method of the invention relies on a clostridial toxin substrate concentration of less than 50 μ M or less than 25 μ M. In a further embodiment, a method of the invention relies on a clostridial toxin substrate concentration of 10 μ M to 20 μ M. If desired, a linear assay also can be performed by mixing clostridial toxin substrate with corresponding, "unlabeled" substrate which lacks a fluorophore. The

appropriate dilution can be determined, for example, by preparing serial dilutions of clostridial toxin substrate in the corresponding unlabeled substrate.

[0067] The concentration of purified or partially purified clostridial toxin to be assayed in a method of the invention generally is in the range of about 0.0001 ng/ml to 500 µg/ml toxin, for example, about 0.0001 ng/ml to 50 µg/ml toxin, 0.001 ng/ml to 500 µg/ml toxin, 0.001 ng/ml to 50 µg/ml toxin, 0.0001 to 5000 ng/ml toxin, for example, about 0.001 ng/ml to 5000 ng/ml, 0.01 ng/ml to 5000 ng/ml, 0.1 ng/ml to 5000 ng/ml, 1 ng/ml to 5000 ng/ml, 10 ng/ml to 5000 ng/ml, 50 ng/ml to 5000 ng/ml, 50 ng/ml to 500 ng/ml or 100 ng/ml to 5000 ng/ml toxin, which can be, for example, purified recombinant dichain toxin or formulated clostridial toxin product containing human serum albumin and excipients. Generally, the amount of purified toxin assayed in a method of the invention is in the range of 0.1 pg to 100 µg, for example, 0.1 pg to 50 µg or 0.1 pg to 10 µg.

[0068] The concentration of purified or partially purified clostridial toxin assayed in a method of the invention can be, for example, in the range of about 0.1 pM to 100 µM, 0.1 pM to 10 µM, 0.1 pM to 1 µM, 0.1 pM to 500 nM, 0.1 pM to 100 nM, for example, 1 pM to 2000 pM, 1 pM to 200 pM, 1 pM to 50 pM, 1 nM to 1 µM, 1 nM to 500 nM, 1 nM to 200 nM, 1 nM to 100 nM or 3 nM to 100 nM toxin, which can be, for example, purified native or recombinant light chain or dichain toxin or formulated clostridial toxin product containing human serum albumin and excipients. In particular embodiments, the concentration of purified or partially purified recombinant BoNT/A or BoNT/E light chain or dichain or formulated toxin product is in the range of 1 pM to 2000 pM, 10 pM to 2000 pM, 20 pM to 2000 pM, 40 pM to 2000 pM, or 1 pM to 200 pM. In further embodiments, the concentration of purified or partially purified recombinant BoNT/C light chain or dichain or formulated toxin product is in the range of 1 to 200 nM, 4 to 100 nM, 10 to 100 nM or 4 to 60 nM. One skilled in the art understands that the concentration of purified or partially purified clostridial toxin will depend on the serotype of the toxin assayed, as well as the purity or recombinant sequence of the toxin, the presence of inhibitory components, and the assay conditions. It is additionally understood that purified, partially purified or crude samples can be diluted to within a convenient range for assaying for clostridial toxin protease activity against a standard curve. Similarly, it is understood that a sample can be diluted, if desired, such that the assay is linear.

[0069] Conditions suitable for clostridial toxin protease activity also generally include, for example, temperatures in the range of about 20°C to about 45°C, for example, in the range of 25°C to 40°C, or the range of 35°C to 39°C. Assay volumes often are in the range of about 5 to about 200 µl, for example, in the range of about 10 µl to 100 µl or about 0.5 µl to 100 µl, although nanoliter reaction volumes also can be used with the methods of the invention. Assay volumes also can be, for example, in the range of 100 µl to 2.0 ml or in the range of 0.5 ml to 1.0 ml.

[0070] One skilled in the art understands that fluorescence polarization reactions may or may not be terminated and that assay times can be varied as appropriate by the skilled artisan. Assay times

generally depend, in part, on the concentration, purity and activity of the clostridial toxin and generally vary, without limitation, in the range of about 15 minutes to about 5 hours. As non-limiting examples, exemplary assay times include incubation, for example, at 37°C for 30 minutes, 45 minutes, 60 minutes, 75 minutes or 90 minutes. In particular embodiments, at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 100% of the clostridial toxin substrate is cleaved. In further embodiments, the protease reaction is stopped before more than 5%, 10%, 15%, 20%, 25% or 50% of the clostridial toxin substrate is cleaved. It is understood that protease reactions can be terminated by the appropriate reagent, which generally depends on the fluorophore and other components of the substrate. As a non-limiting example, a protease reaction based on a substrate containing GFP as the donor fluorophore can be terminated by the addition of guanidinium chloride, for example, to a final concentration of 1 to 2 M. Protease reactions also can be terminated by addition of H₂SO₄; addition of about 0.5 to 1.0 sodium borate, pH 9.0 to 9.5; or addition of zinc chelators. One skilled in the art understands that protease reactions can be terminated, if desired, prior to exciting the fluorophore or donor fluorophore with plane polarized light.

[0071] As a non-limiting example, conditions suitable for clostridial toxin protease activity such as BoNT/A protease activity can be incubation at 37°C for 90 minutes in a buffer containing 50 mM HEPES (pH 7.2), 10 μ M ZnCl₂, 10 mM DTT, and 0.1% (v/v) TWEEN-20 with 10-16 μ M substrate. If desired, samples containing BoNT/A, particularly dichain BoNT/A, can be preincubated with dithiothreitol, for example, for 20 or 30 minutes before addition of substrate. As a further non-limiting example, conditions suitable for BoNT/A protease activity can be incubation at 37°C in a buffer such as 30 mM HEPES (pH 7.3) containing a reducing agent such as 5 mM dithiothreitol; and a source of zinc such as 25 μ M zinc chloride (approximately 7 nM; Schmidt and Bostian, *supra*, 1997). BSA in the range of 0.1 mg/ml to 10 mg/ml, for example, 1 mg/ml BSA, also can be included when a sample is treated with a clostridial toxin substrate (Schmidt and Bostian, *supra*, 1997). As another non-limiting example, conditions suitable for clostridial toxin protease activity, for example BoNT/B activity, can be incubation in 50 mM HEPES, pH 7.4, with 10 μ M zinc chloride, 1% fetal bovine serum and 10 mM dithiothreitol, with incubation for 90 minutes at 37°C (Shone and Roberts, *Eur. J. Biochem.* 225:263-270 (1994); Hallis et al., *supra*, 1996); or can be, for example, incubation in 40 mM sodium phosphate, pH 7.4, with 10 mM dithiothreitol, optionally including 0.2% (v/v) Triton X-100, with incubation for 2 hours at 37°C (Shone et al., *supra*, 1993). Conditions suitable for tetanus toxin protease activity or other clostridial toxin protease activity can be, for example, incubation in 20 mM HEPES, pH 7.2, and 100 mM NaCl for 2 hours at 37°C with 25 μ M peptide substrate (Cornille et al., *supra*, 1994).

[0072] In one embodiment, conditions suitable for clostridial toxin protease activity include cationic polyamino acids such as polyarginine in a buffer of suitable ionic strength. Where there is a charge difference in the clostridial toxin substrate as compared to the cleavage product, fluorescence polarization can be observed in the presence of polyarginine or another cationic polyamino acid (Simeonov et al., *Analytical Biochemistry* 304:193-199 (2002)). As a non-limiting example, if the net ionic charge of a

fluorescently labeled cleavage product becomes negative following treatment of clostridial toxin substrate with a toxin sample, polyarginine will selectively bind to the fluorescently labeled cleavage product, thereby generating a measurable increase in polarization.

[0073] It is understood that any of a variety of control substrates are useful in the methods of the invention. A control substrate can be, for example, a clostridial toxin substrate which is not treated with active, toxin-containing sample; a polarization value determined before addition of the sample; or a similar, but different substrate which does not contain a toxin cleavage site or functional recognition sequence.

[0074] It is understood that the methods of the invention can be automated and can be configured in a high-throughput or ultra high-throughput format using, without limitation, 96-well, 384-well or 1536-well plates. Any of a variety of spectrofluorometers equipped with an appropriate polarizer can be used to assay the change in fluorescence polarization over time including, without limitation a Cary Eclipse spectrofluorometer; the Beckmann Affinity™ Multi-Mode plate reader; TECAN GeniusPro; and other systems from, for example, Perkin Elmer.

[0075] Further provided herein are methods of determining the presence or activity of a clostridial toxin by (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate containing (i) a donor fluorophore; (ii) an acceptor having an absorbance spectrum overlapping the emission spectrum of the donor fluorophore; and (iii) a clostridial toxin recognition sequence containing a cleavage site, where the cleavage site intervenes between the donor fluorophore and the acceptor and where, under the appropriate conditions, resonance energy transfer is exhibited between the donor fluorophore and the acceptor; (b) exciting the donor fluorophore with plane polarized light; and (c) determining fluorescence polarization of the treated substrate relative to a control substrate, where a change in fluorescence polarization of the treated substrate as compared to fluorescence polarization of the control substrate is indicative of the presence or activity of the clostridial toxin. In one embodiment, step (c) includes determining the change in fluorescence polarization of the treated substrate over time.

[0076] In the methods of the invention based on FRET-assisted fluorescence polarization, the change in fluorescence polarization can be an increase or decrease in fluorescence polarization. In one embodiment, the donor fluorophore has a fluorescence lifetime of at least 0.5 nanoseconds. In another embodiment, the donor fluorophore has a fluorescence lifetime of at least 5 nanoseconds. A donor fluorophore useful in the invention can be, without limitation, a green fluorescent protein (GFP); blue fluorescent protein (BFP); cyan fluorescent protein (CFP); yellow fluorescent protein (YFP); red fluorescent protein (RFP); Alexa Fluor® dye; fluorescein; a fluorescein derivative; diaminotriazinylamino-fluorescein (DTAF); a biarsenic derivative of fluorescein; fluorescein arsenical hairpin binding dye (FIAsH™); red biarsenical dye (ReAsH™); carboxyfluorescein (FAM); Texas Red™; tetramethylcarboxy-

rhodamine (TMR); carboxy-x-rhodamine (ROX); rhodamine green; Oregon Green 488; BODIPY®-TR; BODIPY®-TMR; BODIPY®-FL; Cy3, Cy™3B or Dansyl. In particular embodiments, the donor fluorophore is a green fluorescent protein; blue fluorescent protein; cyan fluorescent protein; yellow fluorescent protein or red fluorescent protein. In one embodiment, the donor fluorophore is a green fluorescent protein (GFP). In another embodiment, the acceptor fluorophore is Alexa Fluor® 546.

[0077] Any of a variety of recognition sequences can be included in a clostridial toxin substrate useful in a method of the invention. In one embodiment, the recognition sequence is a BoNT/A recognition sequence such as, without limitation, a BoNT/A recognition sequence containing at least six consecutive residues of SNAP-25, where the six consecutive residues include Gln-Arg, or a peptidomimetic thereof. Such a BoNT/A recognition sequence can include, for example, residues 134 to 206 of SEQ ID NO: 2. A recognition sequence included in a clostridial toxin substrate useful in a method of the invention also can be, without limitation, a BoNT/B recognition sequence. Such a BoNT/B recognition sequence can contain, for example, at least six consecutive residues of VAMP, where the six consecutive residues include Gln-Phe, or a peptidomimetic thereof. In a further embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a BoNT/C1 recognition sequence. Such a BoNT/C1 recognition sequence can contain, without limitation, at least six consecutive residues of syntaxin, where the six consecutive residues include Lys-Ala, or a peptidomimetic thereof. A BoNT/C1 recognition sequence useful in the invention also can contain at least six consecutive residues of SNAP-25, where the six consecutive residues include Arg-Ala, or a peptidomimetic thereof.

[0078] In a further embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a BoNT/D recognition sequence. Such a BoNT/D recognition sequence can contain, for example, at least six consecutive residues of VAMP, where the six consecutive residues include Lys-Leu, or a peptidomimetic thereof. A recognition sequence useful in the invention also can be, for example, a BoNT/E recognition sequence. Such a BoNT/E recognition sequence can contain, without limitation, at least six consecutive residues of SNAP-25, where the six consecutive residues include Arg-Ile, or a peptidomimetic thereof. In yet another embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a BoNT/F recognition sequence. BoNT/F recognition sequences useful in the invention encompass, without limitation, those having at least six consecutive residues of VAMP, where the six consecutive residues include Gln-Lys, or a peptidomimetic thereof. A recognition sequence included in a clostridial toxin substrate useful in a method of the invention also can be a BoNT/G recognition sequence. Such BoNT/G recognition sequences encompass, without limitation, those having at least six consecutive residues of VAMP, where the six consecutive residues include Ala-Ala, or a peptidomimetic thereof. In still a further embodiment, a recognition sequence included in a clostridial toxin substrate useful in a method of the invention is a TeNT recognition sequence. Such a TeNT recognition sequence can be, without limitation, a sequence containing at least six consecutive residues of VAMP, where the six consecutive residues include Gln-Phe, or a peptidomimetic thereof.

[0079] Any of a variety of clostridial toxin substrates can be useful in the methods of the invention, including peptides and peptidomimetics having at least 100 residues, or having at least 200 residues. Furthermore, any of a variety of samples can be assayed according to a method of the invention including, without limitation, crude cell lysates, isolated clostridial toxins including isolated clostridial toxin light chains; and formulated clostridial toxin products such as formulated BoNT/A, BoNT/B or BoNT/E toxin products.

[0080] Where a method of the invention involves fluorescence resonance energy transfer, the method relies on a clostridial toxin substrate which includes, in part, a donor fluorophore. Like a "fluorophore," a "donor fluorophore" is a molecule that, when irradiated with light of a certain wavelength, emits light, also denoted fluorescence, of a different wavelength. A donor fluorophore is a fluorophore which, when paired with a suitable acceptor, transfers energy to the acceptor.

[0081] As used herein, the term "acceptor" means a molecule that can absorb energy from, and upon excitation of, a donor fluorophore. An acceptor useful in a clostridial toxin substrate has an absorbance spectrum which overlaps the emission spectrum of a donor fluorophore included in the substrate. An acceptor useful in the invention generally has rather low absorption at a wavelength suitable for excitation of the donor fluorophore.

[0082] As set forth above, an acceptor has an absorbance spectrum that overlaps the emission spectrum of the donor fluorophore. The term "overlapping," as used herein in reference to the absorbance spectrum of an acceptor and the emission spectrum of a donor fluorophore, means an absorbance spectrum and emission spectrum that are partly or entirely shared. Thus, in such overlapping spectra, the high end of the range of the donor fluorophore's emission spectrum is higher than the low end of the range of the acceptor's absorbance spectrum.

[0083] As set forth above, any of a variety of donor fluorophores can be useful in the invention, including, without limitation, green fluorescent protein; blue fluorescent protein; cyan fluorescent protein; yellow fluorescent protein; red fluorescent protein; an Alexa Fluor[®] dye; fluorescein; a fluorescein derivative; diaminotriazinylamino-fluorescein (DTAF); a biarsenic derivative of fluorescein; fluorescein arsenical hairpin binding dye (FIAsH[™]); red biarsenical dye (ReAsH[™]); carboxyfluorescein (FAM); Texas Red[™]; tetramethylcarboxy-rhodamine (TMR); carboxy-x-rhodamine (ROX); rhodamine green; Oregon Green 488; BODIPY[®]-TR; BODIPY[®]-TMR; BODIPY[®]-FL; Cy3, Cy[™]3B or Dansyl. A variety of acceptors also can be useful in the invention including, but not limited to, Alexa Fluor[®] dyes such as Alexa Fluor[®] 546, Alexa Fluor[®] 568, Alexa Fluor[®] 610, Alexa Fluor[®] 660 and Alexa Fluor[®] 750; QSY[®] 7; tetramethylrhodamine; octadecylrhodamine; flavodoxin, cytochrome c peroxidase; and rubredoxin.

[0084] Exemplary donor fluorophore-acceptor pairs which exhibit FRET and are useful in the methods of the invention encompass, without limitation, GFP and Alexa Fluor[®] 546; fluorescein and QSY[®] 7; fluorescein and tetramethylrhodamine; and dansyl and octadecylrhodamine. Further exemplary donor fluorophore-acceptor pairs which are useful in the methods of the invention encompass, without limitation, Alexa Fluor[®] 633 and Alexa Fluor[®] 660; Alexa Fluor[®] 594 and Alexa Fluor[®] 610; Alexa Fluor[®] 700 and Alexa Fluor[®] 750; and Alexa Fluor[®] 555 and Alexa Fluor[®] 568. Additional acceptors useful in the invention include those in which the acceptor is a protein with a visible chromophore such as, without limitation, flavodoxin, cytochrome c peroxidase or rubredoxin; such a protein can have, for example, a molecular weight in the range of 6 to 34 kDa and a chromophore which absorbs strongly in the region between 400-500 nm. Exemplary donor fluorophore-acceptor pairs based on such proteins include, but are not limited to, 5 -(((2-iodoacetyl)amino)ethyl)amino)naphthalene-1-sulfonic acid (1,5 IAEDANS) and flavodoxin; 4-acetamido-4' maleimidylstilbene 2,2' disulfonic acid and cytochrom c peroxidase; and Alexa Fluor[®] 488 and rubredoxin. These and other donor fluorophores suitable for fluorescence polarization can be paired with any of a variety of acceptors having an absorbance spectrum which overlaps the emission spectrum of the donor fluorophore.

[0085] One skilled in the art understands that the methods of the invention based on FRET-assisted fluorescence polarization optionally utilize a substrate which includes a bulking group in addition to the donor fluorophore and acceptor. One skilled in the art understands that the optional inclusion of a bulking group depends on the molecular weight and bulking characteristics of the selected donor fluorophore and acceptor. A variety of bulking groups are optionally useful in the invention, including those described herein above.

[0086] Substrates useful in the invention can be prepared by recombinant methods or using synthetic chemical methods, or a combination thereof. As described herein in Example I, a fusion protein containing a bulking group fused to a BoNT/A clostridial toxin recognition sequence and a carboxy-terminal cysteine was prepared by recombinant methods. The carboxy-terminal cysteine was used for attachment of a fluorophore to produce the complete clostridial toxin substrate. Recombinant methods for preparation of clostridial toxin substrates which are fusion proteins are well known in the art as described, for example, in Ausubel, Current Protocols in Molecular Biology John Wiley & Sons, Inc., New York 2000.

[0087] Chemical methods for modifying a protein, peptide or peptidomimetic to contain a fluorophore and bulking group, or a donor fluorophore and acceptor, are well known in the art (Fairclough and Cantor, Methods Enzymol. 48:347-379 (1978); Glaser et al., Chemical Modification of Proteins Elsevier Biochemical Press, Amsterdam (1975); Haugland, Excited States of Biopolymers (Steiner Ed.) pp. 29-58, Plenum Press, New York (1983); Means and Feeney, Bioconjugate Chem. 1:2-12 (1990); Matthews et al., Methods Enzymol. 208:468-496 (1991); Lundblad, Chemical Reagents for Protein Modification 2nd Ed., CRC Press, Boca Ratan, Florida (1991); Haugland, *supra*, 1996). A variety of groups can be used to

couple a fluorophore, bulking group, donor fluorophore or acceptor, for example, to a peptide or peptidomimetic containing a clostridial toxin recognition sequence. A thiol group, for example, can be used to couple a fluorophore, bulking group, donor fluorophore or acceptor to the desired position in a peptide or peptidomimetic to produce a clostridial toxin substrate useful in the invention (see Example I). Haloacetyl and maleimide labeling reagents also can be used to couple a fluorophore, bulking group, donor fluorophore or acceptor in preparing a clostridial toxin substrate useful in the invention. See, for example, Wu and Brand, *supra*, 1994.

[0088] Cross-linker moieties also can be useful for preparing a clostridial toxin substrate. Cross-linkers are well known in the art and include homo- and hetero-bifunctional cross-linkers such as BMH and SPDP. Where a fluorophore, bulking group, donor fluorophore or acceptor is a protein, well known chemical methods for specifically linking molecules to the amino- or carboxy-terminus of a protein can be employed. See, for example, "Chemical Approaches to Protein Engineering" in Protein Engineering: A Practical Approach Rees et al. (Eds) Oxford University Press, 1992.

[0089] Where a clostridial toxin substrate contains a fluorophore and bulking group, the clostridial toxin cleavage site is positioned between the fluorophore and bulking group. In one embodiment, the fluorophore is positioned carboxy-terminal of the cleavage site while the bulking group is positioned amino-terminal of the cleavage site. In another embodiment, the fluorophore is positioned amino-terminal of the cleavage site while the bulking group is positioned carboxy-terminal of the cleavage site.

[0090] Where a clostridial toxin substrate contains a donor fluorophore and an acceptor, the clostridial toxin cleavage site is positioned between the donor fluorophore and the acceptor. In one embodiment, the donor fluorophore is positioned amino-terminal of the cleavage site while the acceptor is positioned carboxy-terminal of the cleavage site. In another embodiment, the donor fluorophore is positioned carboxy-terminal of the cleavage site while the acceptor is positioned amino-terminal of the cleavage site.

[0091] One skilled in the art understands that there are several considerations in selecting and positioning a fluorophore and a bulking group, or a donor fluorophore and an acceptor, in a clostridial toxin substrate useful in the invention. The fluorophore and bulking group, or donor fluorophore and acceptor, generally are positioned to minimize interference with substrate binding to, or proteolysis by, the clostridial toxin. Thus, a fluorophore and bulking group, or donor fluorophore and acceptor, can be selected and positioned, for example, so as to minimize the disruption of bonded and non-bonded interactions that are important for binding, and to minimize steric hindrance. In addition, as discussed further below, the spatial distance between an acceptor and donor fluorophore generally is limited to achieve efficient energy transfer from the donor fluorophore to the acceptor.

[0092] As discussed above, efficiency of energy transfer from a donor fluorophore to an acceptor is dependent, in part, on the spatial separation of the donor fluorophore and acceptor molecules. As the distance between the donor fluorophore and acceptor increases, there is less energy transfer to the

acceptor, and the donor fluorescence signal therefore increases. The overall energy transfer between the donor fluorophore and acceptor is dependent upon many factors, including the separation distance between the donor fluorophore and acceptor in the substrate, the spectral overlap between donor fluorophore and acceptor, and the substrate concentration. One skilled in the art understands that, as the concentration of substrate increases, intermolecular quenching of the donor, even after proteolytic cleavage, can become a factor. This phenomenon is denoted the "inner filter effect." One skilled in the art further understands that the concentration of substrate can be controlled as described above.

[0093] The Förster distance, which is the separation between a donor fluorophore and an acceptor for 50% energy transfer, represents a spatial separation between donor fluorophore and acceptor that provides a good sensitivity. For peptide substrates, adjacent residues are separated by a distance of approximately 3.6Å in the most extended conformation. For example, the calculated Förster distance for a fluorescein/tetramethylrhodamine pair is 55Å, which would represent a spatial separation between fluorescein and tetramethylrhodamine of about 15 residues in the most extended conformation. Because peptides and peptidomimetics in solution rarely have a fully extended conformation, donor fluorophores and acceptors can be more widely separated than expected based on a calculation performed using 3.6 Å per residue and still remain within the Förster distance as shown, for example, by the occurrence of FRET between donor-acceptor pairs separated by about 50 amino acids (Graham et al., Analyt. Biochem. 296: 208-217 (2001)).

[0094] Förster theory is based on very weak interactions between a donor fluorophore and an acceptor; spectroscopic properties such as absorption of one fluorophore should not be altered in the presence of the other, defining the shortest distance range over which the theory is valid. It is understood that, for many donor fluorophore-acceptor pairs, Förster theory is valid when donor fluorophores and acceptors are separated by about 10Å to 100Å. However, for particular donor fluorophore-acceptor pairs, Förster theory is valid below 10Å as determined by subpicosecond techniques (Kaschke and Ernstring, *Ultrafast Phenomenon in Spectroscopy* (Klose and Wilhelmi (Eds.)) Springer-Verlag, Berlin 1990).

[0095] In particular embodiments, the invention provides a method that relies on a clostridial toxin substrate in which the donor fluorophore is spatially separated from the acceptor by a distance of at most 100Å. In other embodiments, the invention provides a method that relies on a clostridial toxin substrate in which the donor fluorophore is spatially separated from the acceptor by a distance of at most 90Å, 80Å, 70Å, 60Å, 50Å, 40Å, 30Å or 20Å. In further embodiments, the invention provides a method that relies on a clostridial toxin substrate in which the donor fluorophore is spatially separated from the acceptor by a distance of 10Å to 100Å, 10Å to 80Å, 10Å to 60Å, 10Å to 40Å, 10Å to 20Å, 20Å to 100Å, 20Å to 80Å, 20Å to 60Å, 20Å to 40Å, 40Å to 100Å, 40Å to 80Å or 40Å to 60Å. In still further embodiments, the invention provides a method that relies on a clostridial toxin substrate in which the donor fluorophore and the acceptor are separated in the primary amino acid sequence by at most six residues, at most eight residues, at most ten residues, at most twelve residues, at most fifteen residues, at most twenty residues,

at most twenty-five residues, at most thirty residues, at most thirty-five residues, at most forty residues, at most forty-five residues, at most fifty residues, at most sixty residues, at most seventy residues, at most eighty residues, at most ninety residues, at most 100 residues, at most 150 residues, at most 200 residues or up to the full-length of a naturally occurring clostridial toxin target protein.

[0096] One skilled in the art understands that a clostridial toxin substrate useful in the invention can be designed, if desired, to optimize the efficiency of FRET. One skilled in the art understands that a donor fluorophore can be selected, if desired, with a high quantum yield, and acceptor can be selected, if desired, with a high extinction coefficient to maximize the Förster distance. One skilled in the art further understands that fluorescence arising from direct excitation of an acceptor can be difficult to distinguish from fluorescence resulting from resonance energy transfer. Thus, it is recognized that a donor fluorophore and acceptor can be selected which have relatively little overlap of their excitation spectra such that the donor can be excited at a wavelength that does not result in direct excitation of the acceptor. It further is recognized that a clostridial toxin substrate useful in the invention can be designed so that the emission spectra of the donor fluorophore and acceptor overlap relatively little such that the two emissions can be readily distinguished.

[0097] Specific and distinct cleavage sites for different clostridial toxins are well known in the art. BoNT/A cleaves a Gln-Arg bond; BoNT/B and TeNT cleave a Gln-Phe bond; BoNT/C1 cleaves a Lys-Ala or Arg-Ala bond; BoNT/D cleaves a Lys-Leu bond; BoNT/E cleaves an Arg-Ile bond; BoNT/F cleaves a Gln-Lys bond; and BoNT/G cleaves an Ala-Ala bond (see Table A). In standard nomenclature, the sequence surrounding a clostridial toxin cleavage site is denoted P_5 - P_4 - P_3 - P_2 - P_1 - P_1' - P_2' - P_3' - P_4' - P_5' , with P_1 - P_1' representing the scissile bond. It is understood that a P_1 or P_1' site, or both, can be substituted with another amino acid or amino acid mimetic in place of the naturally occurring residue. As an example, BoNT/A substrates have been prepared in which the P_1 position (Gln) is modified to be an alanine, 2-aminobutyric acid or asparagine residue; these substrates were hydrolyzed by BoNT/A at the P_1 -Arg bond (Schmidt and Bostian, *J. Protein Chem.* 16:19-26 (1997)). While it is recognized that substitutions can be introduced at the P_1 position of the scissile bond, for example, a BoNT/A scissile bond, it is further recognized that conservation of the P_1' residue can be advantageous (Vaidyanathan et al., *J. Neurochem.* 72:327-337 (1999)). Thus, in particular embodiments, the invention provides a method which relies on a clostridial toxin substrate having a clostridial toxin recognition sequence in which the P_1' residue is not modified or substituted relative to the naturally occurring residue in a target protein cleaved by the clostridial toxin. In other embodiments, the invention provides a method which relies on a clostridial toxin substrate having a recognition sequence in which the P_1 residue is modified or substituted relative to the naturally occurring residue in a target protein cleaved by the clostridial toxin; such a clostridial toxin substrate retains susceptibility to peptide bond cleavage between the P_1 and P_1' residues.

TABLE A
BONDS CLEAVED IN HUMAN VAMP-2, SNAP-25 OR SYNTAXIN

Toxin	Target	P ₄ -P ₃ -P ₂ -P ₁ --P ₁ '-P ₂ '-P ₃ '-P ₄ '	SEQ ID NO:
BoNT/A	SNAP-25	Glu-Ala-Asn- Gln—Arg* -Ala-Thr-Lys	22
BoNT/B	VAMP-2	Gly-Ala-Ser- Gln—Phe* -Glu-Thr-Ser	23
BoNT/C1	syntaxin	Asp-Thr-Lys- Lys—Ala* -Val-Lys-Tyr	24
BoNT/D	VAMP-2	Arg-Asp-Gln- Lys—Leu* -Ser-Glu-Leu	25
BoNT/E	SNAP-25	Gln-Ile-Asp- Arg—Ile* -Met-Glu-Lys	26
BoNT/F	VAMP-2	Glu-Arg-Asp- Gln—Lys* -Leu-Ser-Glu	27
BoNT/G	VAMP-2	Glu-Thr-Ser- Ala—Ala* -Lys-Leu-Lys	28
TeNT	VAMP-2	Gly-Ala-Ser- Gln—Phe* -Glu-Thr-Ser	29

* Scissile bond shown in bold

[0098] SNAP-25, VAMP and syntaxin share a short motif located within regions predicted to adopt an α -helical conformation. This motif is present in SNAP-25, VAMP and syntaxin isoforms expressed in animals sensitive to the neurotoxins. In contrast, *Drosophila* and yeast homologs that are resistant to these neurotoxins and syntaxin isoforms not involved in exocytosis contain sequence variations in the α -helical motif regions of these VAMP and syntaxin proteins.

[0099] Multiple repetitions of the α -helical motif are present in proteins sensitive to cleavage by clostridial toxins: Four copies are naturally present in SNAP-25; two copies are naturally present in VAMP; and two copies are naturally present in syntaxin. Furthermore, peptides corresponding to the specific sequence of the α -helical motifs can inhibit neurotoxin activity *in vitro* and *in vivo*, and such peptides can cross-inhibit different neurotoxins. In addition, antibodies raised against such peptides can cross-react among the three target proteins, indicating that this α -helical motif is exposed on the protein surface and adopts a similar configuration in each of the three target proteins. Consistent with these findings, SNAP-25-specific, VAMP-specific and syntaxin-specific neurotoxins cross-inhibit each other by competing for the same binding site, although they do not cleave targets non-specifically. These results indicate that a clostridial toxin recognition sequence can include, if desired, at least one α -helical motif. It is recognized that an α -helical motif is not required for cleavage by a clostridial toxin, as evidenced by 16-mer and 17-mer substrates for BoNT/A known in the art.

[00100] Although multiple α -helical motifs are found in the naturally occurring SNAP-25, VAMP and syntaxin target proteins, a clostridial toxin recognition sequence useful in a clostridial toxin substrate can have a single α -helical motif. In particular embodiments, a method of the invention relies on a clostridial toxin recognition sequence including two or more α -helical motifs. A BoNT/A or BoNT/E recognition sequence can include, for example, the S4 α -helical motif, alone or combined with one or more additional α -helical motifs; a BoNT/B, BoNT/G or TeNT recognition sequence can include, for example, the V2 α -helical motif, alone or combined with one or more additional α -helical motifs; a BoNT/C1 recognition

sequence can include, for example, the S4 α -helical motif, alone or combined with one or more additional α -helical motifs, or the X2 α -helical motif, alone or combined with one or more additional α -helical motifs; and a BoNT/D or BoNT/F recognition sequence can include, for example, the V1 α -helical motif, alone or combined with one or more additional α -helical motifs.

[00101] As used herein, the term "botulinum toxin serotype A recognition sequence" is synonymous with "BoNT/A recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/A under conditions suitable for clostridial toxin protease activity. A scissile bond cleaved by BoNT/A can be, for example, Gln-Arg.

[00102] A variety of BoNT/A recognition sequences are well known in the art and are useful in the invention. A BoNT/A recognition sequence can have, for example, residues 134 to 206 or residues 137 to 206 of human SNAP-25 (Ekong et al., *supra*, 1997; U.S. Patent No. 5,962,637). A BoNT/A recognition sequence also can include, without limitation, the sequence Thr-Arg-Ile-Asp-Glu-Ala-Asn-Gln-Arg-Ala-Thr-Lys-Met (SEQ ID NO: 30) or a peptidomimetic thereof, which corresponds to residues 190 to 202 of human SNAP-25; Ser-Asn-Lys-Thr-Arg-Ile-Asp-Glu-Ala-Asn-Gln-Arg-Ala-Thr-Lys (SEQ ID NO: 31) or a peptidomimetic thereof, which corresponds to residues 187 to 201 of human SNAP-25; Ser-Asn-Lys-Thr-Arg-Ile-Asp-Glu-Ala-Asn-Gln-Arg-Ala-Thr-Lys-Met (SEQ ID NO: 32) or a peptidomimetic thereof, which corresponds to residues 187 to 202 of human SNAP-25; Ser-Asn-Lys-Thr-Arg-Ile-Asp-Glu-Ala-Asn-Gln-Arg-Ala-Thr-Lys-Met-Leu (SEQ ID NO: 33) or a peptidomimetic thereof, which corresponds to residues 187 to 203 of human SNAP-25; Asp-Ser-Asn-Lys-Thr-Arg-Ile-Asp-Glu-Ala-Asn-Gln-Arg-Ala-Thr-Lys-Met (SEQ ID NO: 34) or a peptidomimetic thereof, which corresponds to residues 186 to 202 of human SNAP-25; or Asp-Ser-Asn-Lys-Thr-Arg-Ile-Asp-Glu-Ala-Asn-Gln-Arg-Ala-Thr-Lys-Met-Leu (SEQ ID NO: 35) or a peptidomimetic thereof, which corresponds to residues 186 to 203 of human SNAP-25. See, for example, Schmidt and Bostian, *J. Protein Chem.* 14:703-708 (1995); Schmidt and Bostian, *supra*, 1997; Schmidt et al., *FEBS Letters* 435:61-64 (1998); and Schmidt and Bostian, U.S. Patent No. 5,965,699. If desired, a similar BoNT/A recognition sequence can be prepared from a corresponding (homologous) segment of another BoNT/A-sensitive SNAP-25 isoform or homolog such as, for example, murine, rat, goldfish or zebrafish SNAP-25 or can be any of the peptides described herein or known in the art, for example, in U.S. Patent No. 5,965,699.

[00103] A BoNT/A recognition sequence useful in the invention can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype A, or can be substantially similar to a segment of a BoNT/A-sensitive protein. As illustrated in Table B, a variety of naturally occurring proteins sensitive to cleavage by BoNT/A are known in the art and include, for example, human, mouse and rat SNAP-25; and goldfish SNAP-25A and SNAP-25B. Thus, a BoNT/A recognition sequence useful in the invention can correspond, for example, to a segment of human SNAP-25, mouse SNAP-25, rat SNAP-25, goldfish

SNAP-25A or 25B, or another naturally occurring protein sensitive to cleavage by BoNT/A. Furthermore, comparison of native SNAP-25 amino acid sequences cleaved by BoNT/A reveals that such sequences are not absolutely conserved (see Table B and Figure 3), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/A-sensitive SNAP-25 sequence can be tolerated in a BoNT/A recognition sequence useful in the invention.

TABLE B

TABLE B
Cleavage of SNAP-25 and related proteins^{a,b,c,d}

Species		Isoform		Cleavage Sites					SEQ ID NO:	Resistance to Cleavage by	
				BoNT/E		BoNT/A		BoNT/C			
			▼				▼ ▼				
Human			174								
Mouse	-----	SNAP-25		qnrqid	ri	mekadsnkrtridean	qra	tkmlgsg	206	36	none ^a
Human	-----	SNAP-23	180	qnrqid	ri	tdkadtnrdridian	qra	kkldls	end	37	all ^b
Mouse	-----	SNAP-23	179	qnqqiq	ri	tekadttnkridian	qra	kkldls	end	38	BoNT/A&C
Chicken	-----	SNAP-25	174	qnrqid	ri	meklipikpglmkpt	qra	qresavvk	end	39	BoNT/A&C
Goldfish	-----	SNAP-25 A	171	qnrqid	ri	mdmdadsnkrtridean	qra	tkmlgsg	end	40	none
	-----	SNAP-25 B	172	qnrqid	ri	mekadsnkrtridean	qra	tkmlgsg	end	41	none
Torpedo	-----	SNAP-25	180	qnaqvd	ri	v ¹⁸⁹ kgdmnkaridean	qra	tkml	end	42	BoNT/E ^c & A ^d
Sea urchin	-----	SNAP-25	180	qnsqvg	ri	tskaesnegrinsad	qra	knilmk	end	43	(?) ^a
C. elegans	-----	SNAP-25	203	qnrqid	ri	hdkqsnevrvesank	qra	nlitk	end	44	BoNT/A & C
Drosophila	-----	SNAP-25	182	qnrqid	ri	nrgesneariavan	qra	hqllk	end	45	BoNT/E & A ^c
Fish	-----	SNAP-25	181	qnrqvd	ri	nnkmtnsqhrisdan	qra	skllke	end	46	BoNT/A ^c

^a = In vitro cleavage of SNAP-25 requires 1000-fold higher BoNT/C concentration than BoNT/A or /E.

^b = Substitution of p182r, or k185dd (boxes) induces susceptibility toward BoNT/E.

^c = Resistance to BoNT/A possibly due to d189 or e189 substitution by v189, see box.

^d = Note that Torpedo is susceptible to BoNT/A.

^e = Note the presence of several non-conservative mutations around putative cleavage sites.

[00104] A clostridial toxin substrate, such as a substrate containing a BoNT/A recognition sequence, can have one or multiple modifications as compared to a naturally occurring sequence that is cleaved by the corresponding clostridial toxin. As an example, as compared to a 17-mer corresponding to residues 187 to 203 of human SNAP-25, substitution of Asp193 with Asn in the BoNT/A substrate resulted in a relative rate of proteolysis of 0.23; substitution of Glu194 with Gln resulted in a relative rate of 2.08; substitution of Ala195 with 2-aminobutyric acid resulted in a relative rate of 0.38; and substitution of Gln197 with Asn, 2-aminobutyric acid or Ala resulted in a relative rate of 0.66, 0.25, or 0.19, respectively (see Table C). Furthermore, substitution of Ala199 with 2-aminobutyric acid resulted in a relative rate of 0.79; substitution of Thr200 with Ser or 2-aminobutyric acid resulted in a relative rate of 0.26 or 1.20, respectively; substitution of Lys201 with Ala resulted in a relative rate of 0.12; and substitution of Met202 with Ala or norleucine resulted in a relative rate of 0.38 or 1.20, respectively. See Schmidt and Bostian, *supra*, 1997. These results indicate that a variety of residues can be substituted in a clostridial toxin substrate as compared to a naturally occurring toxin-sensitive sequence. In the case of BoNT/A, these results indicate that residues including but not limited to Glu194, Ala195, Gln197, Ala199, Thr200 and Met202, Leu203, Gly204, Ser205, and Gly206, as well as residues more distal from the Gln-Arg scissile bond, can be substituted or conjugated to a fluorophore, bulking group, donor fluorophore or acceptor in a BoNT/A substrate useful in the invention. Such a BoNT/A substrate is detectably proteolyzed at the scissile bond by BoNT/A under conditions suitable for clostridial toxin protease activity. Thus, a BoNT/A substrate can include, if desired, one or several amino acid substitutions, additions or deletions relative to a naturally occurring SNAP-25 sequence.

TABLE C
KINETIC PARAMETERS OF BONT/A SYNTHETIC PEPTIDE SUBSTRATES

Peptide	Sequence ^a	SEQ ID NO:	Relative Rate ^b
[1-15]	SNKTRIDEANQRATK	31	0.03
[1-16]	SNKTRIDEANQRATKM	32	1.17
[1-17]	SNKTRIDEANQRATKML	33	1.00
M16A	SNKTRIDEANQRATK <u>A</u> L	50	0.38
M16X	SNKTRIDEANQRATK <u>X</u> L	51	1.20
K15A	SNKTRIDEANQRAT <u>A</u> ML	52	0.12
T14S	SNKTRIDEANQRAS <u>S</u> KML	53	0.26
T14B	SNKTRIDEANQRAB <u>B</u> KML	54	1.20
A13B	SNKTRIDEANQRB <u>B</u> TKML	55	0.79
Q11A	SNKTRIDEAN <u>A</u> RATKML	56	0.19
Q11B	SNKTRIDEAN <u>B</u> RATKML	57	0.25
Q11N	SNKTRIDEAN <u>N</u> RATKML	58	0.66
N10A	SNKTRIDEA <u>A</u> QRATKML	59	0.06
A9B	SNKTRIDE <u>B</u> NQRATKML	60	0.38
E8Q	SNKTRID <u>Q</u> ANQRATKML	61	2.08
D7N	SNKTRID <u>N</u> ANQRATKML	62	0.23

^a Nonstandard amino acid abbreviations are: B, 2-aminobutyric acid; X, 2-aminohexanoic acid (norleucine)

^b Initial hydrolysis rates relative to peptide [1-17]. Peptide concentrations were 1.0 mM.

[00105] As used herein, the term "botulinum toxin serotype B recognition sequence" is synonymous with "BoNT/B recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/B under appropriate conditions. A scissile bond cleaved by BoNT/B can be, for example, Gln-Phe.

[00106] A variety of BoNT/B recognition sequences are well known in the art or can be defined by routine methods. Such BoNT/B recognition sequences can include, for example, a sequence corresponding to some or all of the hydrophilic core of a VAMP protein such as human VAMP-1 or human VAMP-2. A BoNT/B recognition sequence can include, without limitation, residues 33 to 94, residues 45 to 94, residues 55 to 94, residues 60 to 94, residues 65 to 94, residues 60 to 88 or residues 65 to 88 of human VAMP-2 (SEQ ID NO: 8), or residues 60 to 94 of human VAMP-1 (SEQ ID NO: 7). See, for example, Shone et al., Eur. J. Biochem. 217: 965-971 (1993). and U.S. Patent No. 5,962,637. If desired, a similar BoNT/B recognition sequence can be prepared from a corresponding (homologous) segment of another BoNT/B-sensitive VAMP isoform or homolog such as human VAMP-1 or rat or chicken VAMP-2.

[00107] Thus, it is understood that a BoNT/B recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype B, or can be substantially similar to such a segment of a BoNT/B-sensitive protein. As shown in Table D, a variety of naturally occurring proteins sensitive to cleavage by BoNT/B are known in the art and include, for example, human, mouse and bovine VAMP-1 and VAMP-2; rat VAMP-2; rat cellubrevin; chicken VAMP-2; *Torpedo* VAMP-1; sea urchin VAMP; *Aplysia* VAMP; squid VAMP; *C. elegans* VAMP; *Drosophila* n-syb; and leech VAMP. Thus, a BoNT/B recognition sequence included in a BoNT/B substrate can correspond, for example, to a segment of human VAMP-1 or VAMP-2, mouse VAMP-1 or VAMP-2, bovine VAMP-1 or VAMP-2, rat VAMP-2, rat cellubrevin, chicken VAMP-2, *Torpedo* VAMP-1, sea urchin VAMP, *Aplysia* VAMP, squid VAMP, *C. elegans* VAMP, *Drosophila* n-syb, leech VAMP, or another naturally occurring protein sensitive to cleavage by BoNT/B. Furthermore, as shown in Table D, comparison of native VAMP amino acid sequences cleaved by BoNT/B reveals that such sequences are not absolutely conserved (see, also, Figure 4), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring VAMP sequence can be tolerated in a BoNT/B substrate of the invention.

TABLE D

TABLE D
Cleavage of VAMP^{a,b}

Species	Isoform	Cleavage Sites	SEQ ID NO:	Resistance to Cleavage by
		BoNT/B TeNT		
		BoNT/F ⌋ ⌋ BoNT/D ⌋ ⌋ BoNT/G		
human		VAMP-1 ⁵³ dkvlerd qkl selddradalqagas qf ess aa klkrkyww ⁹²	63	none
mouse		VAMP-2 ⁵¹ dkvlerd qkl selddradalqagas qf ets aa klkrkyww ⁹⁰	64	none
ovine		VAMP-1 ⁵³ dkvlerd qkl selddradalqagas qf ess aa klkrkyww ⁹²	65	TeNT & BoNT/B
		VAMP-2 ⁵¹ dkvlerd qkl selddradalqagas qf ets aa klkrkyww ⁹⁰	66	none
rat		Cellubrevin ³⁸ dkvlerd qkl selddradalqagas qf ets aa klkrkyww ⁷⁷	67	none
		TI-VAMP ¹⁴⁶ dlvaqrg qkl ellidktenlvdsav qf ktt aa nlaramcm ¹⁷⁵	68	all
		VAMP-1 ⁵³ ----erd qkl selddradalqagas qf ess aa klkr----	69	TeNT & BoNT/B
chicken		VAMP-2 ⁵¹ ----erd qkl selddradalqagas qf ets aa klkr----	70	none
torpedo		VAMP-1 ⁵⁵ dkvlerd qkl selddradalqagas qf ess aa klkrkyww ⁹⁴	71	none
sea urchin		VAMP ³⁵ dkvlrd qkl svlddradalqagas qf etn aa klkrkyww ⁷⁴	72	BoNT/F, D & G

TABLE D

TABLE D
Cleavage of VAMP^{a,b}

Species	Isoform	Cleavage Sites								SEQ ID NO:	Resistance to Cleavage by
			BoNT/F		BoNT/D		BoNT/B		TeNT		
			↓	↓	↓	↓	↓	↓	↓		
plysia	VAMP ⁴¹	ekvldrd	qk ^a	sqlddraealqagas	qf	eas	a ^a	klkrkyww	⁸⁰	73	BoNT/G
quid	VAMP ⁶⁰	dkvlerd	sk ^a	selddradalqagas	qf	eas	a ^a	klkrkfw	⁹⁹	74	BoNT/F & G
C. elegans	VAMP ⁸⁶	nkvmnerd	sk ^a	nsldhraevlqngas	qf	qqs	a ^a	tlrqkyww	¹¹⁵	75	BoNT/F, D & G
	syb ^a ⁶⁷	ekvlerd	qkl	selgeradqleqgas	q ^a	eqq	a ^a	klkrkqww	¹⁰⁶	76	TeNT & BoNT/B & G
Drosophila	n-syb ^b ⁶¹	ekvlerd	sk ^a	selddradalqqgas	qf	eqq	a ^a	klkrkfwl	¹⁰⁰	77	BoNT/F & G
Xech	VAMP ⁴⁹	dkvlekd	qkl	aeldgradalqagas	qf	eas	a ^a	klkrkfw	⁸⁸	78	BoNT/G

^a = Sequence corrected in position 93 (f>s).^b = Sequence corrected in position 68 (t>s).

[00108] As used herein, the term "botulinum toxin serotype C1 recognition sequence" is synonymous with "BoNT/C1 recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/C1 under appropriate conditions. A scissile bond cleaved by BoNT/C1 can be, for example, Lys-Ala or Arg-Ala.

[00109] It is understood that a BoNT/C1 recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype C1, or can be substantially similar to a segment of a BoNT/C1-sensitive protein. As shown in Table E, a variety of naturally occurring proteins sensitive to cleavage by BoNT/C1 are known in the art and include, for example, human, rat, mouse and bovine syntaxin 1A and 1B; rat syntaxins 2 and 3; sea urchin syntaxin; *Aplysia* syntaxin 1; squid syntaxin; *Drosophila* Dsynt1; and leech syntaxin 1. Thus, a BoNT/C1 recognition sequence useful in a BoNT/C1 substrate can correspond, for example, to a segment of human, rat, mouse or bovine syntaxin 1A or 1B, rat syntaxin 2, rat syntaxin 3, sea urchin syntaxin, *Aplysia* syntaxin 1, squid syntaxin, *Drosophila* Dsynt1, leech syntaxin 1, or another naturally occurring protein sensitive to cleavage by BoNT/C1. Furthermore, comparison of native syntaxin amino acid sequences cleaved by BoNT/C1 reveals that such sequences are not absolutely conserved (see Table E and Figure 5), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/C1-sensitive syntaxin sequence can be tolerated in a BoNT/C1 substrate useful in the invention.

TABLE E

TABLE E
Cleavage of syntaxin

Species	Isoform	Cleavage Sites	SEQ ID NO:	Resistance to Cleavage by
		BoNT/C		
human, rat	syntaxin 1A	²⁴⁵ eravsdtk ka vkyqskar ²⁶²	79	<i>no</i>
mouse				
bovine	syntaxin 1B	²⁴⁴ eravsdtk ka vkyqskar ²⁶¹	80	<i>no</i>
	syntaxin 2	²⁴⁵ ehakeetk ka ikyqskar ²⁶²	81	<i>no</i>
rat	syntaxin 3	²⁴⁴ ekardetr ka mkyqgqar ²⁶¹	82	<i>no</i>
	syntaxin 4	²⁴⁴ ergqehvk ka lenqlkar ²⁶¹	83	<i>yes</i>
chicken	syntaxin 1B	²³⁹ vpevfvtk ka vmyqcksr ²⁵⁹	84	<i>expected</i>
sea urchin	syntaxin	²⁴³ vrrqndtk ka vkyqskar ²⁶⁰	85	<i>no</i>
<i>Aplysia</i>	syntaxin 1	²⁴⁷ etakmdtk ka vkyqskar ²⁶⁴	86	<i>no</i>
squid	syntaxin	²⁴⁸ etakvdtk ka vkyqskar ²⁶⁵	87	<i>no</i>
<i>Drosophila</i>	Dsynt 1	²⁴⁸ qtatqdtk ka lkyqskar ²⁶⁵	88	<i>no</i>
leech	syntaxin 1	²⁵¹ etaaadtk ka mkyqsaar ²⁶⁸	89	<i>no</i>

[00110] A variety of naturally occurring SNAP-25 proteins also are sensitive to cleavage by BoNT/C1, including human, mouse and rat SNAP-25; goldfish SNAP-25A and 25B; and *Drosophila* and leech SNAP-25. Thus, a BoNT/C1 recognition sequence useful in a BoNT/C1 substrate can correspond, for example, to a segment of human, mouse or rat SNAP-25, goldfish SNAP-25A or 25B, *Torpedo* SNAP-25, zebrafish SNAP-25, *Drosophila* SNAP-25, leech SNAP-25, or another naturally occurring protein sensitive to cleavage by BoNT/C1. As discussed above in regard to variants of naturally occurring syntaxin sequences, comparison of native SNAP-25 amino acid sequences cleaved by BoNT/C1 reveals significant sequence variability (see Figure 3 and Table B above), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/C1-sensitive SNAP-25 sequence can be tolerated in a BoNT/C1 substrate useful in the invention.

[00111] The term "botulinum toxin serotype D recognition sequence" is synonymous with "BoNT/D recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/D under appropriate conditions. A scissile bond cleaved by BoNT/D can be, for example, Lys-Leu.

[00112] A variety of BoNT/D recognition sequences are well known in the art or can be defined by routine methods. A BoNT/D recognition sequence can include, for example, residues 27 to 116; residues 37 to 116; residues 1 to 86; residues 1 to 76; or residues 1 to 69 of rat VAMP-2 (SEQ ID NO: 90; Yamasaki et al., *J. Biol. Chem.* 269:12764-12772 (1994)). Thus, a BoNT/D recognition sequence can include, for example, residues 27 to 69 or residues 37 to 69 of rat VAMP-2 (SEQ ID NO: 90). If desired, a similar BoNT/D recognition sequence can be prepared from a corresponding (homologous) segment of another BoNT/D-sensitive VAMP isoform or homolog such as human VAMP-1 or human VAMP-2.

[00113] A BoNT/D recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype D, or can be substantially similar to a segment of a BoNT/D-sensitive protein. As shown in Table D, a variety of naturally occurring proteins sensitive to cleavage by BoNT/D are known in the art and include, for example, human, mouse and bovine VAMP-1 and VAMP-2; rat VAMP-1 and VAMP-2; rat cellubrevin; chicken VAMP-1 and VAMP-2; *Torpedo* VAMP-1; *Aplysia* VAMP; squid VAMP; *Drosophila* syb and n-syb; and leech VAMP. Thus, a BoNT/D recognition sequence can correspond, for example, to a segment of human VAMP-1 or VAMP-2, mouse VAMP-1 or VAMP-2, bovine VAMP-1 or VAMP-2, rat VAMP-1 or VAMP-2, rat cellubrevin, chicken VAMP-1 or VAMP-2, *Torpedo* VAMP-1, *Aplysia* VAMP, squid VAMP, *Drosophila* syb or n-syb, leech VAMP, or another naturally occurring protein sensitive to cleavage by BoNT/D. Furthermore, as shown in Table D above, comparison of native VAMP amino acid sequences cleaved by BoNT/D reveals significant sequence variability (see, also, Figure 4), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/D-sensitive VAMP sequence can be tolerated in a BoNT/D substrate useful in the invention.

[00114] As used herein, the term "botulinum toxin serotype E recognition sequence" is synonymous with "BoNT/E recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/E under appropriate conditions. A scissile bond cleaved by BoNT/E can be, for example, Arg-Ile.

[00115] One skilled in the art appreciates that a BoNT/E recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype E, or can be substantially similar to a segment of a BoNT/E-sensitive protein. A variety of naturally occurring proteins sensitive to cleavage by BoNT/E are known in the art and include, for example, human, mouse and rat SNAP-25; mouse SNAP-23; chicken SNAP-25; goldfish SNAP-25A and SNAP-25B; zebrafish SNAP-25; *C. elegans* SNAP-25; and leech SNAP-25 (see Table B). Thus, a BoNT/E recognition sequence can correspond, for example, to a segment of human SNAP-25, mouse SNAP-25, rat SNAP-25, mouse SNAP-23, chicken SNAP-25, goldfish SNAP-25A or 25B, *C. elegans* SNAP-25, leech SNAP-25, or another naturally occurring protein sensitive to cleavage by BoNT/E. Furthermore, as shown in Table B and Figure 3 above, comparison of native SNAP-23 and SNAP-25 amino acid sequences cleaved by BoNT/E reveals that such sequences are not absolutely conserved, indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/E-sensitive SNAP-23 or SNAP-25 sequence can be tolerated in a BoNT/E substrate useful in the invention.

[00116] The term "botulinum toxin serotype F recognition sequence," as used herein, is synonymous with "BoNT/F recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/F under appropriate conditions. A scissile bond cleaved by BoNT/F can be, for example, Gln-Lys.

[00117] A variety of BoNT/F recognition sequences are well known in the art or can be defined by routine methods. A BoNT/F recognition sequence can include, for example, residues 27 to 116; residues 37 to 116; residues 1 to 86; residues 1 to 76; or residues 1 to 69 of rat VAMP-2 (SEQ ID NO: 90; Yamasaki et al., *supra*, 1994). A BoNT/F recognition sequence also can include, for example, residues 27 to 69 or residues 37 to 69 of rat VAMP-2 (SEQ ID NO: 90). It is understood that a similar BoNT/F recognition sequence can be prepared, if desired, from a corresponding (homologous) segment of another BoNT/F-sensitive VAMP isoform or homolog such as human VAMP-1 or human VAMP-2.

[00118] A BoNT/F recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype F, or can be substantially similar to a segment of a BoNT/F-sensitive protein. A variety of naturally occurring proteins sensitive to cleavage by BoNT/F are known in the art and include, for example, human, mouse and bovine VAMP-1 and VAMP-2; rat VAMP-1 and VAMP-2; rat cellubrevin; chicken VAMP-1 and VAMP-2; *Torpedo* VAMP-1; *Aplysia* VAMP; *Drosophila* syb; and leech VAMP (see Table D). Thus, a BoNT/F recognition sequence can correspond, for example, to a segment of human VAMP-1 or VAMP-2, mouse VAMP-1 or VAMP-2, bovine VAMP-1 or

VAMP-2, rat VAMP-1 or VAMP-2, rat cellubrevin, chicken VAMP-1 or VAMP-2, *Torpedo* VAMP-1, *Aplysia* VAMP, *Drosophila* syb, leech VAMP, or another naturally occurring protein sensitive to cleavage by BoNT/F. Furthermore, as shown in Table D above, comparison of native VAMP amino acid sequences cleaved by BoNT/F reveals that such sequences are not absolutely conserved (see, also, Figure 4), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/F-sensitive VAMP sequence can be tolerated in a BoNT/F substrate useful in the invention.

[00119] As used herein, the term "botulinum toxin serotype G recognition sequence" is synonymous with "BoNT/G recognition sequence" and means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a BoNT/G under appropriate conditions. A scissile bond cleaved by BoNT/G can be, for example, Ala-Ala.

[00120] A BoNT/G recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by botulinum toxin serotype G, or can be substantially similar to such a BoNT/G-sensitive segment. As illustrated in Table D above, a variety of naturally occurring proteins sensitive to cleavage by BoNT/G are known in the art and include, for example, human, mouse and bovine VAMP-1 and VAMP-2; rat VAMP-1 and VAMP-2; rat cellubrevin; chicken VAMP-1 and VAMP-2; and *Torpedo* VAMP-1. Thus, a BoNT/G recognition sequence can correspond, for example, to a segment of human VAMP-1 or VAMP-2, mouse VAMP-1 or VAMP-2, bovine VAMP-1 or VAMP-2, rat VAMP-1 or VAMP-2, rat cellubrevin, chicken VAMP-1 or VAMP-2, *Torpedo* VAMP-1, or another naturally occurring protein sensitive to cleavage by BoNT/G. Furthermore, as shown in Table D above, comparison of native VAMP amino acid sequences cleaved by BoNT/G reveals that such sequences are not absolutely conserved (see, also, Figure 4), indicating that a variety of amino acid substitutions and modifications relative to a naturally occurring BoNT/G-sensitive VAMP sequence can be tolerated in a BoNT/G substrate useful in the invention.

[00121] As used herein, the term "tetanus toxin recognition sequence" means a scissile bond together with adjacent or non-adjacent recognition elements, or both, sufficient for detectable proteolysis at the scissile bond by a tetanus toxin under appropriate conditions. A scissile bond cleaved by TeNT can be, for example, Gln-Phe.

[00122] A variety of TeNT recognition sequences are well known in the art or can be defined by routine methods and include sequences corresponding to some or all of the hydrophilic core of a VAMP protein such as human VAMP-1 or human VAMP-2. A TeNT recognition sequence can include, for example, residues 25 to 93 or residues 33 to 94 of human VAMP-2 (SEQ ID NO: 8; Cornille et al., Eur. J. Biochem. 222:173-181 (1994); Foran et al., Biochem. 33: 15365-15374 (1994)); residues 51 to 93 or residues 1 to 86 of rat VAMP-2 (SEQ ID NO: 90; Yamasaki et al., *supra*, 1994); or residues 33 to 94 of human VAMP-1 (SEQ ID NO: 7). A TeNT recognition sequence also can include, for example, residues 25 to 86, residues 33 to 86 or residues 51 to 86 of human VAMP-2 (SEQ ID NO: 8) or rat VAMP-2 (SEQ ID NO:

90). It is understood that a similar TeNT recognition sequence can be prepared, if desired, from a corresponding (homologous) segment of another TeNT-sensitive VAMP isoform or species homolog such as human VAMP-1 or sea urchin or *Aplysia* VAMP.

[00123] Thus, a TeNT recognition sequence can correspond to a segment of a protein that is sensitive to cleavage by tetanus toxin, or can be substantially similar to a segment of a TeNT-sensitive protein. As shown in Table D above, a variety of naturally occurring proteins sensitive to cleavage by TeNT are known in the art and include, for example, human, mouse and bovine VAMP-1 and VAMP-2; rat VAMP-2; rat cellubrevin; chicken VAMP-2; *Torpedo* VAMP-1; sea urchin VAMP; *Aplysia* VAMP; squid VAMP; *C. elegans* VAMP; *Drosophila* n-syb; and leech VAMP. Thus, a TeNT recognition sequence can correspond, for example, to a segment of human VAMP-1 or VAMP-2, mouse VAMP-1 or VAMP-2, bovine VAMP-1 or VAMP-2, rat VAMP-2, rat cellubrevin, chicken VAMP-2, *Torpedo* VAMP-1, sea urchin VAMP, *Aplysia* VAMP, squid VAMP, *C. elegans* VAMP, *Drosophila* n-syb, leech VAMP, or another naturally occurring protein sensitive to cleavage by TeNT. Furthermore, comparison of native VAMP amino acid sequences cleaved by TeNT reveals that such sequences are not absolutely conserved (Table D and Figure 4). This finding indicates that a variety of amino acid substitutions and modifications relative to a naturally occurring TeNT-sensitive VAMP sequence can be tolerated in a TeNT substrate useful in the invention.

[00124] As used herein, the term "peptidomimetic" is used broadly to mean a peptide-like molecule that is cleaved by the same clostridial toxin as the peptide substrate upon which it is structurally based. Such peptidomimetics include chemically modified peptides, peptide-like molecules containing non-naturally occurring amino acids, and peptoids, which are peptide-like molecules resulting from oligomeric assembly of N-substituted glycines, and are cleaved by the same clostridial toxin as the peptide substrate upon which the peptidomimetic is derived (see, for example, Goodman and Ro, Peptidomimetics for Drug Design, in "Burger's Medicinal Chemistry and Drug Discovery" Vol. 1 (ed. M.E. Wolff; John Wiley & Sons 1995), pages 803-861).

[00125] A variety of peptidomimetics are known in the art including, for example, peptide-like molecules which contain a constrained amino acid, a non-peptide component that mimics peptide secondary structure, or an amide bond isostere. A peptidomimetic that contains a constrained, non-naturally occurring amino acid can include, for example, an α -methylated amino acid; an α,α -dialkyl-glycine or α -aminocycloalkane carboxylic acid; an N^α -C $^\alpha$ cyclized amino acid; an N^α -methylated amino acid; a β - or γ -amino cycloalkane carboxylic acid; an α,β -unsaturated amino acid; a β , β -dimethyl or β -methyl amino acid; a β -substituted-2,3-methano amino acid; an NC $^\delta$ or C $^\alpha$ -C $^\delta$ cyclized amino acid; or a substituted proline or another amino acid mimetic. In addition, a peptidomimetic which mimics peptide secondary structure can contain, for example, a nonpeptidic β -turn mimic; γ -turn mimic; mimic of β -sheet structure; or mimic of helical structure, each of which is well known in the art. A peptidomimetic also can be a peptide-like molecule which contains, for example, an amide bond isostere such as a retro-inverso

modification; reduced amide bond; methylenethioether or methylenesulfoxide bond; methylene ether bond; ethylene bond; thioamide bond; trans-olefin or fluoroolefin bond; 1,5-disubstituted tetrazole ring; ketomethylene or fluoroketomethylene bond or another amide isostere. One skilled in the art understands that these and other peptidomimetics are encompassed within the meaning of the term "peptidomimetic" as used herein.

[00126] In any of the methods of the invention, a clostridial toxin substrate can include one or multiple clostridial toxin cleavage sites for the same or different clostridial toxins. In particular embodiments, the invention provides methods that rely on a clostridial toxin substrate which contains a single clostridial toxin cleavage site. In other embodiments, the invention provides methods which rely on a clostridial toxin substrate which contains multiple cleavage sites for the same clostridial toxin. These cleavage sites can be incorporated within the same or different clostridial toxin recognition sequences. As non-limiting examples, a clostridial toxin substrate can have multiple cleavage sites for the same clostridial toxin intervening between the same fluorophore and bulking group or the same donor fluorophore and acceptor. A clostridial toxin substrate useful in the invention can contain, for example, two or more, three or more, five or more, or ten or more cleavage sites for the same clostridial toxin. A clostridial toxin substrate useful in the invention also can have, for example, two, three, four, five, six, seven, eight, nine or ten cleavage sites for the same clostridial toxin; the multiple cleavage sites can intervene between the same or different fluorophores and bulking groups, or between the same or different donor fluorophores and acceptors.

[00127] A clostridial toxin substrate useful in the invention also can include cleavage sites for different clostridial toxins. In particular embodiments, the invention provides a method that relies on a clostridial toxin substrate which includes multiple cleavage sites for different clostridial toxins all intervening between the same fluorophore and bulking group, or between the same donor fluorophore and acceptor. A clostridial toxin substrate can include, for example, cleavage sites for two or more, three or more, or five or more different clostridial toxins all intervening between the same fluorophore and bulking group. A clostridial toxin substrate also can include, for example, cleavage sites for two or more, three or more, or five or more different clostridial toxins all intervening between the same donor fluorophore and acceptor. A clostridial toxin substrate also can incorporate, for example, cleavage sites for two or more, three or more, or five or more different clostridial toxins intervening between at least two fluorophore-bulking group pairs or between at least two donor fluorophore-acceptor pairs. In particular embodiments, the invention provides a clostridial toxin substrate having cleavage sites for two, three, four, five, six, seven or eight different clostridial toxins, where the cleavage sites intervene between the same or different fluorophores and bulking groups, or between the same or different donor fluorophores and acceptors. In further embodiments, the invention provides a clostridial toxin substrate which has any combination of two, three, four, five, six, seven or eight cleavage sites for any combination of the following clostridial toxins: BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G and TeNT.

[00128] A method of the invention optionally can be performed with multiple substrates. In a method of the invention which relies on a clostridial toxin substrate containing a donor fluorophore-acceptor pair, a clostridial toxin substrate is treated with a sample, the substrate including a first donor fluorophore, a first acceptor having an absorbance spectrum which overlaps the emission spectrum of the first donor fluorophore, and a first clostridial toxin recognition sequence containing a cleavage site, where the cleavage site intervenes between the donor fluorophore and the acceptor and where, under the appropriate conditions, resonance energy transfer is exhibited between the first donor fluorophore and the first acceptor. If desired, a second clostridial toxin substrate can be included in the same assay; this second substrate contains a second donor fluorophore and second acceptor having an absorbance spectrum which overlaps the emission spectrum of the second donor fluorophore, and a second clostridial toxin recognition sequence that is cleaved by a different clostridial toxin than the toxin that cleaves the first clostridial toxin recognition sequence. The donor fluorophore-acceptor pair in the second substrate can be the same or different from the donor fluorophore-acceptor pair in the first substrate. In this way, a single sample can be simultaneously assayed for the presence of more than one clostridial toxin.

[00129] In a method of the invention which relies on a clostridial toxin substrate containing a donor fluorophore-acceptor pair, it is understood that one can assay for any combination of clostridial toxins, for example, two, three, four, five, six, seven, eight, or more clostridial toxins. One can assay, for example, any combination of two, three, four, five, six, seven or eight of BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G and TeNT. As an example, an assay can be performed with seven substrates, each of which includes fluorescein and tetramethylrhodamine flanking a BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F or BoNT/G recognition sequence and cleavage site. These substrates can be treated with a sample under conditions suitable for botulinum toxin activity before exciting the donor fluorescein with plane polarized light at an absorption wavelength of about 488 nm and determining fluorescence polarization. A change in the fluorescence polarization is indicative of the presence or activity of at least one clostridial toxin. Such an assay can be useful, for example, for assaying food samples or tissue samples for the presence of any botulinum or other clostridial toxin and can be combined, if desired, with one or more subsequent assays for individual clostridial toxins or specific combinations of clostridial toxins.

[00130] In another embodiment, a single sample is assayed for two or more different clostridial toxins using two or more different clostridial toxin substrates, with each substrate containing a different donor fluorophore-acceptor pair. The use of multiple substrates can be useful for extending the dynamic range of an assay, as described, for example, in U.S. Patent No. 6,180,340. As an example of the use of multiple clostridial toxin substrates, a single sample can be assayed for the presence or activity of BoNT/A and BoNT/B using first and second clostridial toxin substrates: the first clostridial toxin substrate contains the donor fluorophore Alexa Fluor® 555 and the acceptor Alexa Fluor® 568 with an intervening BoNT/A recognition sequence, and a second clostridial toxin substrate contains the donor fluorophore Alexa Fluor® 700 and the acceptor Alexa Fluor® 750 with an intervening BoNT/B recognition sequence.

Those skilled in the art understand that the first donor fluorophore can be excited before or after excitation of the second donor fluorophore, and that the change in fluorescence polarization of the first substrate can be determined before, at the same time, or after determining energy transfer of the second substrate.

[00131] In a further embodiment, a method of the invention is useful for assaying two or more different purified or isolated clostridial toxins using two or more different clostridial toxin substrates, with each substrate containing the same donor fluorophore-acceptor pair. In the endpoint format, the presence or activity of different serotypes is assayed by adding the serotypes sequentially and waiting between additions for the response to stabilize.

EXAMPLES

[00132] The following examples are intended to illustrate but not limit the present invention.

EXAMPLE I: Preparation of GFP-Snap25-His6-C- Alexa Fluor® 594 And GFP-Snap25-His6-C- Alexa Fluor® 546 Substrates

[00133] This example describes construction of substrates suitable for assaying for the presence or activity of a clostridial toxin using fluorescence polarization.

A. Construction of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C

[00134] A substrate was prepared as a fusion protein containing green fluorescent protein (GFP), murine SNAP-25 residues 134-206, a polyhistidine affinity tag (6xHis), and a carboxy-terminal cysteine, with several components separated by peptide linkers. As described further below, the substrate was designed such that the GFP and terminal cysteine were present at opposite ends of SNAP-25₍₁₃₄₋₂₀₆₎.

[00135] The SNAP-25 sequence was obtained from pT25FL, a plasmid which contains the full-length mouse SNAP-25 gene inserted in frame with the 3' terminus of the glutathione-S-transferase (GST) gene (GST-SNAP25₍₁₋₂₀₆₎), provided by Professor Dolly (O'Sullivan et al., *J. Biol. Chem.* 274:36897-36904 (1999)). The SNAP-25 sequence from pT25FL was incorporated into a second expression vector, which was designed to have a BirAsp signal sequence for biotinylation and a polyhistidine affinity tag fused to the amino-terminus of residues 134 to 206 of SNAP-25 (BirAsp-polyHis-SNAP25₍₁₃₄₋₂₀₆₎, denoted "BA-SNAP"). The DNA sequence encoding SNAP25₍₁₃₄₋₂₀₆₎ was generated by PCR amplification of the appropriate region of the pT25FL plasmid with PCR primers 5'-GCT AGA TCT CGA GTT AAC CAC TTC CCA GCA TCT TTG-3' (SEQ ID NO: 91; antisense) and 5'-ATC CGG AGG GTA ACA AAC GAT GCC-3' (SEQ ID NO: 92; sense) to produce a SNAP25₍₁₃₄₋₂₀₆₎ PCR product containing a *Bgl* II restriction site (PCR product A).

[00136] The BirAsp sequence, a natural substrate for biotinylation, as well as a polyhistidine affinity tag, were engineered for fusion upstream and in frame with the SNAP25₍₁₃₄₋₂₀₆₎ sequence using synthetic oligonucleotides SEQ ID NOS: 93 and 94, which contained a 20 bp complementary region. These oligonucleotides, 5'-CGA ATT CCG CGG GCC ACC ATG GGA GGA GGA CTG AAC GAC ATC TTC GAG GCT CAA AAG ATC-3' (SEQ ID NO: 93; sense; *Sac* II site underlined) and 5'-TCG TTT GTT ACC CTC CGG ATA TGA TGA TGA TGA TGA TGA TGA TGG GAT CCA TGC CAC TCG ATC TTT TGA GCC TCG AAG A-3' (SEQ ID NO: 94; antisense), were annealed, and the single strand overhangs filled by PCR amplification to yield PCR product B.

[00137] The two double stranded PCR products containing the coding sequences for SNAP25₍₁₃₄₋₂₀₆₎, denoted PCR product A, and BirAsp and polyhistidine, denoted PCR product B, were denatured and annealed. The 20 bp complementary sequence in the two gene fragments is shown in italics in PCR primers SEQ ID NO: 92 and SEQ ID NO: 94). After filling in the overhangs by PCR, the product was amplified with primers SEQ ID NO: 93 and SEQ ID NO: 91. The resulting PCR product, which encoded BirAsp-polyHis-SNAP25₍₁₃₄₋₂₀₆₎ (designated "BA-SNAP"), was digested with *Sac*II and *Bgl*II, and the isolated gene insert ligated into pQBI25fA2 vector digested with *Sac*II and *Bam*HI, to yield plasmid pNTP12 (pQBI25fA2 containing BA-SNAP).

[00138] For expression and purification from *E. coli*, the BA-SNAP gene was transferred into a pTrc99A plasmid (Amersham Pharmacia Biotech). The BA-SNAP gene was isolated from pNTP12 by digestion with *Nco*I and *Xho*I followed by gel purification. Separately, the pTrc99A plasmid was digested with *Nco*I and *Sal*I, and the isolated vector ligated to the BA-SNAP gene to yield plasmid pNTP14 (pTrc99A containing BA-SNAP).

[00139] For cloning of the BA-SNAP gene into plasmid pQE-50, the BA-SNAP fragment was PCR amplified from pNTP14 with primer SEQ ID NO: 91 and primer SEQ ID NO: 95 (5'-CGA AGA TCT GGA GGA CTG AAC GAC ATC TTC-3' (sense; *Bgl* II site underlined)). After digestion with *Bgl*II and *Xho*I, the amplified PCR product was ligated into vector pQE-50, which had been digested with *Bam*HI and *Sal*I. The resulting plasmid, which represents pQE50 containing BA-SNAP, was designated pNTP26.

[00140] A plasmid encoding the green fluorescent protein (GFP) fusion protein substrate was prepared by modifying vector pQBI T7-GFP (Quantum Biotechnologies; Carlsbad, CA) in three phases as described below. First, vector pQBI T7-GFP was PCR-modified to remove the stop codon at the 3' terminus of the GFP-coding sequence and to insert the coding sequence for a portion of the peptide linker separating GFP from the SNAP-25 fragment. Second, a DNA fragment coding for SNAP-25₍₁₃₄₋₂₀₆₎ was PCR amplified from pNTP26 using PCR primers designed to incorporate the coding sequence for the remainder of the peptide linker fused 5' to the SNAP-25₍₁₃₄₋₂₀₆₎ gene and a 6xHis affinity tag fused 3' of the gene. The resultant PCR product was cloned into the modified pQBI vector described above to yield pQBI GFP-SNAP25₍₁₃₄₋₂₀₆₎.

[00141] Plasmid pQBI GFP-SNAP25₍₁₃₄₋₂₀₆₎ was then modified by site-directed mutagenesis to add a cysteine codon at the carboxy-terminus using primer SEQ ID NO: 96 (5'-GATGGTGATGGTGATGACAGCCGCCACC GCCACC-3' (antisense primer, with the added nucleotides underlined) and its reverse complement (sense primer). The resulting plasmid, designated pQBI GFP-SNAP25 (Cys-Stop), is shown in Figure 6A and was used for expression of GFP-SNAP25₍₁₃₄₋₂₀₆₎-6xHis-Cys. The nucleic acid and predicted amino acid sequence for the GFP-SNAP25₍₁₃₄₋₂₀₆₎-6xHis-cysteine construct is shown herein in Figure 6B.

B. Expression and characterization of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C

[00142] The pQBI GFP-SNAP25 (Cys-Stop) expression vector was transformed into *E. coli* BL21(DE3) cells (Novagen; Madison, WI; or Invitrogen; Carlsbad, CA) or into *E. coli* BL21-CodonPlus[®](DE3)-RIL cells (Stratagene) containing the T7 RNA polymerase gene. Transformed cells were selected on LB-ampicillin plates overnight at 37 °C. Single colonies were used to inoculate 1-3 mL starter cultures, which were in turn used to inoculate 0.5 to 1.0 L cultures. The large cultures were grown at 37 °C with shaking until A₅₉₅ reached 0.5-0.6, at which time they were removed from the incubator and were allowed to cool briefly. After induction of protein expression with 1 mM IPTG, GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C substrate was expressed from the pQBI GFP-SNAP25 (Cys-Stop) plasmid overnight with shaking at 16 °C in order to facilitate formation of the GFP fluorophore. Cells from 250 mL aliquots of the expression cultures were collected by centrifugation (30 minutes, 6,000 x g, 4 °C) and stored at -80 °C until needed.

[00143] Substrates were purified at 4 °C by a two-step procedure involving IMAC purification, followed by a de-salting step to remove NaCl and imidazole, typically yielding greater than 150 mg/L of purified substrate as follows. Cell pellets from 250 mL cultures were each resuspended in 7-12 mL Column Binding Buffer (25 mM HEPES, pH 8.0; 500 mM NaCl; 1 mM β-mercaptoethanol; 10 mM imidazole), lysed by sonication (1 minute 40 seconds in 10-second pulses at 38% amplitude), and clarified by centrifugation (16000 rpm, 4 °C, 1 hour). Affinity resin (3-5 mL Talon SuperFlow Co²⁺ per cell pellet) was equilibrated in a glass or disposable column support (Bio-Rad) by rinsing with 4 column volumes of sterile ddH₂O and 4 column volumes of Column Binding Buffer. Clarified lysate was applied to the column in one of two ways: (1) Lysate was added to the resin and batch bound by horizontal incubation for 1 hour with gentle rocking or (2) Lysate was applied to the vertical column and allowed to enter the column slowly by gravity flow. Following batch binding only, the column was righted and the solution drained, collected, and passed over the resin again. In both cases, after the lysate had been applied, the column was washed with 4-5 column volumes of Column Binding Buffer. In some cases, the column was further washed with 1-2 column volumes of Column Wash Buffer (25 mM HEPES, pH 8.0; 500 mM NaCl; 1 mM β-mercaptoethanol; 20 mM imidazole). Protein was eluted with 1.5 to 2.0 column volumes of Column Elution Buffer (25 mM HEPES, pH 8.0; 500 mM NaCl; 1 mM β-mercaptoethanol; 250 mM imidazole), which was collected in fractions of ~1.4 mL. The green fractions were combined, concentrated with a

centrifugal filter (10,000 or 30,000 molecular weight cut-off) and desalted by FPLC (BioRad Biologic DuoLogic, QuadTec UV-Vis detector) with a HiPrep 26/10 size exclusion column (Pharmacia) and an isocratic mobile phase of chilled Fusion Protein Desalting Buffer (50 mM HEPES, pH 7.4, 4 °C) at a flow rate of 10 mL/minute. Desalted protein was collected as a single fraction, and the concentration determined using a BioRad Protein Assay. The GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C substrate was analyzed by reducing SDS-PAGE. The protein solution was subsequently divided into 500 μ L aliquots, flash-frozen with liquid nitrogen and stored at -80 °C. Once defrosted, a working aliquot was stored at 4 °C, protected from light.

C. Labeling with Alexa Fluor[®] 594 and Alexa Fluor[®] 546

[00144] The GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C construct contains a single cysteine residue which is solvent exposed although there are three buried cysteine residues within GFP which are not available for chemical modification (Selvin, *supra*, 2000; Heyduk, *Curr. Opin. Biotech.* 13:292-296 (2002)). The carboxy-terminal cysteine residue can therefore be selectively labeled using a fluorophore-maleimide at neutral pH. Shown in Figures 7A and 7B, respectively, are the absorption and emission/excitation spectra of purified GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein. The concentration of the protein solution was determined to be 2.74 mg/ml based on the theoretical molar extinction coefficient of 20250 M⁻¹cm⁻¹ as calculated from the primary sequence of the construct. The molecular weight of the purified GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein was confirmed to be about 37,000 using Matrix Assisted Laser Desorption Time of Flight mass spectrometry (MALDI-TOF).

[00145] Labeling with Alexa Fluor[®] 594 was performed essentially as follows. The C-terminal cysteine residue of the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein was labeled by adding a concentrated solution of Alexa Fluor[®] 594 (Molecular Probes, Inc.) in dry dimethyl formamide (DMF) to a final concentration of 20:1 molar excess of fluorophore to protein. The protein/fluorophore solution was kept at 4°C in the refrigerator overnight and subsequently dialyzed against 20mM HEPES pH 6.9.

[00146] The absorption spectrum of the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein labeled with Alexa Fluor[®] 594 is shown in Figure 8A following dialysis against 20 mM HEPES pH 6.9, which is the pH used for assaying enzymatic activity of reduced bulk toxin or purified BoNT-A light chain. The labeling ratio, as calculated from the absorption spectrum using the theoretical extinction coefficient of the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C construct, was approximately 3:1 (protein: Alexa probe). Shown in Figure 8B are the excitation and emission spectra of labeled GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor[®] 594 after extensive dialysis for 20 hours with three changes of buffer to remove free probe.

[00147] Labeling with Alexa Fluor[®] 546 was performed essentially as follows with all procedures carried out on ice or at 4 °C. Four microliters of a 10 mM aqueous solution of Alexa Fluor[®] 546 C₅ maleimide (MW 1,034.37; Molecular Probes) were added to 200 μ L of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C (135 μ M in 25

mM HEPES buffer, pH 7.2), mixed well, and incubated at 4 °C overnight. The reactions were transferred to Biomax Ultrafree centrifugal filters (30 KDa NMWL; Millipore), concentrated, and then reconcentrated two times from 25 mM HEPES, pH 7.2, to remove most of the excess Alexa Fluor® 546. To remove the remaining unreacted Alexa Fluor® 546, the concentrated solutions were transferred to Spin Microdialyzers (Harvard Apparatus) and each was dialyzed against 500 mL 20 mM HEPES, pH 6.9, for 1 hour, and against 3 x 250 mL of that buffer for about 1.5 hours each. A small aliquot was removed for fluorescence measurements, and the balance of the reaction was flash-frozen in liquid nitrogen and stored at -80 °C.

EXAMPLE II: Clostridial Toxin Complex Activity Assayed Using Fluorescence Polarization

[00148] This example demonstrates that a fluorescence polarization assay can be used to determine the presence or activity of a clostridial toxin.

[00149] The GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein labeled with Alexa Fluor® 594 was tested for its utility as a suitable substrate for BoNT/A reduced bulk toxin by recording the change in polarization over time. A Cary Eclipse spectrofluorometer (Varian, Inc.; Palo Alto, California) equipped with motorized thin-film polarizers was used to monitor the reaction. The excitation wavelength was set at 590 nm, and polarized emission was recorded at 620 nm.

[00150] Several dilutions of BoNT/A bulk toxin or BoNT/A light chain were prepared, and fluorescence polarization monitored over time. For both bulk toxin and light chain, fluorescence polarization was reduced at or shortly after the time the diluted toxin was added. Figure 9 shows the data for bulk BoNT/A toxin proteolysis of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor® 594. As shown in panel 9D, toxin was detected at a concentration of as little as about 50 ng/ml.

[00151] These results demonstrate that the presence or activity of a clostridial toxin can be sensitively determined using synthetic substrates assayed by fluorescence polarization.

EXAMPLE III: Clostridial Toxin Complex Activity Assayed Using Fluorescence Polarization in Combination with Fluorescence Resonance Energy Transfer

[00152] This example demonstrates that fluorescence polarization can be assayed to determine the presence or activity of a clostridial toxin using a substrate which exhibits fluorescence resonance energy transfer.

[00153] The GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C protein labeled with Alexa Fluor® 546 as described above was utilized as a substrate for BoNT/A. As indicated above, the photoselection properties of GFP and Alexa Fluor® 546 provide for fluorescence resonance energy transfer (FRET) between the donor fluorophore GFP and the acceptor Alexa Fluor® 546. Steady-state polarization measurements were carried out in a

Cary Eclipse spectrophotometer (Varian). Excitation was at 474 nm, the excitation maximum of the GFP component. Emission was measured at the Alexa Fluor® 546 fluorescence maximum of 570 nm. In all cases, a dual path length cuvette (10 mm by 2 mm) was utilized, and the emission viewed through the 2 mm path. A solution of 390 μ L Toxin Reaction Buffer (50 mM HEPES, pH 7.2; 0.1 % v/v TWEEN-20; 10 μ M Zn Cl₂, 10 mM DTT) and 10 μ L of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor® 546 was placed in the cuvette and allowed to equilibrate to 30 °C. When the polarization measurements, which were taken at 30 second intervals, were stabilized, 10 μ L of recombinant BoNT/A light chain (rLC/A) at a concentration of 1.0 μ g/ μ L, 0.5 μ g/ μ L, 0.25 μ g/ μ L, or 0.1 μ g/ μ L was added to the cuvette. Measurements continued to be taken until the polarization again stabilized.

[00154] As shown in Figure 10, fluorescence polarization increased upon addition of recombinant BoNT/A light chain which results in substrate cleavage. As compared to the substrate having GFP and Alexa Fluor® 594, the fluorescence resonance energy transfer enhanced the polarization change upon turnover, thereby increasing the sensitivity of the assay. The overall change in polarization using the GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor® 546 substrate was about 40 mP, twice the magnitude of the depolarization of approximately 20 mP observed during proteolysis of GFP-SNAP25₍₁₃₄₋₂₀₆₎-His6-C-Alexa Fluor® 594.

[00155] These results indicate that fluorescence polarization can be combined with fluorescence resonance energy transfer for enhanced sensitivity in assaying for the presence or activity of a clostridial toxin.

[00156] All journal article, reference and patent citations provided above, in parentheses or otherwise, whether previously stated or not, are incorporated herein by reference in their entirety.

[00157] Although the invention has been described with reference to the examples provided above, it should be understood that various modifications can be made without departing from the spirit of the invention. Accordingly, the invention is limited only by the following claims.

We claim:

1. A method of determining the presence or activity of a clostridial toxin, comprising the steps of:
 - (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate comprising
 - (i) a fluorophore;
 - (ii) a bulking group; and
 - (iii) a clostridial toxin recognition sequence comprising a cleavage site, wherein said cleavage site intervenes between said fluorophore and said bulking group;
 - (b) exciting said fluorophore with plane polarized light; and
 - (c) determining fluorescence polarization of said treated substrate relative to a control substrate, wherein a change in fluorescence polarization of said treated substrate as compared to fluorescence polarization of said control substrate is indicative of the presence or activity of said clostridial toxin.
2. The method of claim 1, wherein said change in fluorescence polarization is a decrease in fluorescence polarization.
3. The method of claim 1, wherein said fluorophore has a fluorescence lifetime of at least 0.5 nanoseconds.
4. The method of claim 1, wherein said fluorophore has a fluorescence lifetime of at least 5 nanoseconds.
5. The method of claim 1, wherein said fluorophore has a fluorescence lifetime of at least 10 nanoseconds.
6. The method of claim 1, wherein said fluorophore is selected from the group of an Alexa Fluor[®] dye; fluorescein; a fluorescein derivative; diaminotriazinylamino-fluorescein (DTAF); a biarsenic derivative of fluorescein; fluorescein arsenical hairpin binding dye (FIAsH[™]); red biarsenical dye (ReAsH[™]); carboxyfluorescein (FAM); Texas Red[™]; tetramethylcarboxyrhodamine (TMR); carboxy-x-rhodamine (ROX); rhodamine green; Oregon Green 488; BODIPY-TR[®]; BODIPY-TMR; BODIPY[®]-FL; Cy3, Cy[™]3B and Dansyl.
7. The method of claim 6, wherein said fluorophore is an Alexa Fluor[®] dye.
8. The method of claim 7, wherein said Alexa Fluor[®] dye is Alexa Fluor[®] 594.
9. The method of claim 6, wherein said fluorophore is FIAsH[™] or ReAsH[™].
10. The method of claim 1, wherein said bulking group is a fluorescent protein.
11. The method of claim 10, wherein said bulking group is green fluorescent protein.
12. The method of claim 1, wherein the change in molecular mass upon cleavage of said clostridial toxin substrate is at least 1000 Da.
13. The method of claim 1, wherein said decrease in fluorescence polarization is at least 5 millipolarization units (mP).
14. The method of claim 13, wherein said decrease in fluorescence polarization is at least 15 mP.
15. The method of claim 1, wherein said recognition sequence is a BoNT/A recognition sequence.

16. The method of claim 15, wherein said BoNT/A recognition sequences comprises at least six consecutive residues of SNAP-25, said six consecutive residues comprising Gln-Arg, or a peptidomimetic thereof.
17. The method of claim 16, wherein said BoNT/A recognition sequence comprises residues 134 to 206 of SEQ ID NO: 2.
18. The method of claim 1, wherein said recognition sequence is a BoNT/B recognition sequence.
19. The method of claim 18, wherein said BoNT/B recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Gln-Phe, or a peptidomimetic thereof.
20. The method of claim 1, wherein said recognition sequence is a BoNT/C1 recognition sequence.
21. The method of claim 20, wherein said BoNT/C1 recognition sequence comprises at least six consecutive residues of syntaxin, said six consecutive residues comprising Lys-Ala, or a peptidomimetic thereof.
22. The method of claim 21, wherein said BoNT/C1 recognition sequence comprises at least six consecutive residues of SNAP-25, said six consecutive residues comprising Arg-Ala, or a peptidomimetic thereof.
23. The method of claim 1, wherein said recognition sequence is a BoNT/D recognition sequence.
24. The method of claim 23, wherein said BoNT/D recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Lys-Leu, or a peptidomimetic thereof.
25. The method of claim 1, wherein said recognition sequence is a BoNT/E recognition sequence.
26. The method of claim 25, wherein said BoNT/E recognition sequence comprises at least six consecutive residues of SNAP-25, said six consecutive residues comprising Arg-Ile, or a peptidomimetic thereof.
27. The method of claim 1, wherein said recognition sequence is a BoNT/F recognition sequence.
28. The method of claim 27, wherein said BoNT/F recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Gln-Lys, or a peptidomimetic thereof.
29. The method of claim 1, wherein said recognition sequence is a BoNT/G recognition sequence.
30. The method of claim 29, wherein said BoNT/G recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Ala-Ala, or a peptidomimetic thereof.
31. The method of claim 1, wherein said recognition sequence is a TeNT recognition sequence.
32. The method of claim 31, wherein said TeNT recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Gln-Phe, or a peptidomimetic thereof.
33. The method of claim 1, wherein said clostridial toxin substrate is a peptide or peptidomimetic having at least 100 residues.
34. The method of claim 1, wherein said clostridial toxin substrate is a peptide or peptidomimetic having at least 200 residues.

35. The method of claim 1, wherein said sample is a crude cell lysate.
36. The method of claim 1, wherein said sample is isolated clostridial toxin.
37. The method of claim 1, wherein said sample is isolated clostridial toxin light chain.
38. The method of claim 1, wherein said sample is a formulated clostridial toxin product.
39. The method of claim 38, wherein said formulated product is formulated BoNT/A, BoNT/B or BoNT/E toxin product.
40. The method of claim 39, wherein said formulated product is formulated BoNT/A toxin product.
41. The method of claim 1, wherein step (c) comprises determining the change in fluorescence polarization of said treated substrate over time.
42. A method of determining the presence or activity of a clostridial toxin, comprising the steps of:
 - (a) treating with a sample, under conditions suitable for clostridial toxin protease activity, a clostridial toxin substrate comprising
 - (i) a donor fluorophore;
 - (ii) an acceptor having an absorbance spectrum overlapping the emission spectrum of said donor fluorophore; and
 - (iii) a clostridial toxin recognition sequence comprising a cleavage site, wherein said cleavage site intervenes between said donor fluorophore and said acceptor and wherein, under the appropriate conditions, resonance energy transfer is exhibited between said donor fluorophore and said acceptor;
 - (b) exciting said donor fluorophore with plane polarized light; and
 - (c) determining fluorescence polarization of said treated substrate relative to a control substrate, wherein a change in fluorescence polarization of said treated substrate as compared to fluorescence polarization of said control substrate is indicative of the presence or activity of said clostridial toxin.
43. The method of claim 42, wherein said change in fluorescence polarization is an increase in fluorescence polarization.
44. The method of claim 42, wherein said donor fluorophore has a fluorescence lifetime of at least 0.5 nanoseconds.
45. The method of claim 42, wherein said donor fluorophore has a fluorescence lifetime of at least 5 nanoseconds.
46. The method of claim 42, wherein said donor fluorophore is selected from the group of green fluorescent protein (GFP); blue fluorescent protein (BFP); cyan fluorescent protein (CFP); yellow fluorescent protein (YFP) and red fluorescent protein (RFP); an Alexa Fluor[®] dye; fluorescein; a fluorescein derivative; diaminotriazinylamino-fluorescein (DTAF); a biarsenic derivative of fluorescein; fluorescein arsenical hairpin binding dye (FIAsh[™]); red biarsenical dye (ReAsH[™]); carboxyfluorescein (FAM); Texas Red[™]; tetramethylcarboxyrhodamine (TMR); carboxy-x-rhodamine (ROX); rhodamine green; Oregon Green 488; BODIPY-TR[®]; BODIPY-TMR; BODIPY[®]-FL; Cy3; Cy[™]3B and Dansyl.
47. The method of claim 46, wherein said donor fluorophore is selected from the group green fluorescent protein; blue fluorescent protein; cyan fluorescent protein; yellow fluorescent protein; and red fluorescent protein (RFP).

48. The method of claim 47, wherein said donor fluorophore is GFP.
49. The method of claim 42 or claim 48, wherein said acceptor is Alexa Fluor[®] 546.
50. The method of claim 42, wherein said acceptor is selected from the group Alexa Fluor[®] 546; Alexa Fluor[®] 568; Alexa Fluor[®] 610; Alexa Fluor[®] 660; Alexa Fluor[®] 750; QSY[®] 7; tetramethylrhodamine; octadecylrhodamine; flavodoxin, cytochrome c peroxidase; and rubredoxin.
51. The method of claim 42, wherein said recognition sequence is a BoNT/A recognition sequence.
52. The method of claim 51, wherein said BoNT/A recognition sequences comprises at least six consecutive residues of SNAP-25, said six consecutive residues comprising Gln-Arg, or a peptidomimetic thereof.
53. The method of claim 52, wherein said BoNT/A recognition sequence comprises residues 134 to 206 of SEQ ID NO: 2.
54. The method of claim 42, wherein said recognition sequence is a BoNT/B recognition sequence.
55. The method of claim 54, wherein said BoNT/B recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Gln-Phe, or a peptidomimetic thereof.
56. The method of claim 42, wherein said recognition sequence is a BoNT/C1 recognition sequence.
57. The method of claim 56, wherein said BoNT/C1 recognition sequence comprises at least six consecutive residues of syntaxin, said six consecutive residues comprising Lys-Ala, or a peptidomimetic thereof.
58. The method of claim 56, wherein said BoNT/C1 recognition sequence comprises at least six consecutive residues of SNAP-25, said six consecutive residues comprising Arg-Ala, or a peptidomimetic thereof.
59. The method of claim 42, wherein said recognition sequence is a BoNT/D recognition sequence.
60. The method of claim 59, wherein said BoNT/D recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Lys-Leu, or a peptidomimetic thereof.
61. The method of claim 60, wherein said recognition sequence is a BoNT/E recognition sequence.
62. The method of claim 61, wherein said BoNT/E recognition sequence comprises at least six consecutive residues of SNAP-25, said six consecutive residues comprising Arg-Ile, or a peptidomimetic thereof.
63. The method of claim 42, wherein said recognition sequence is a BoNT/F recognition sequence.
64. The method of claim 63, wherein said BoNT/F recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Gln-Lys, or a peptidomimetic thereof.
65. The method of claim 42, wherein said recognition sequence is a BoNT/G recognition sequence.
66. The method of claim 65, wherein said BoNT/G recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Ala-Ala, or a peptidomimetic thereof.
67. The method of claim 42, wherein said recognition sequence is a TeNT recognition sequence.

68. The method of claim 67, wherein said TeNT recognition sequence comprises at least six consecutive residues of VAMP, said six consecutive residues comprising Gln-Phe, or a peptidomimetic thereof.
69. The method of claim 42, wherein said clostridial toxin substrate is a peptide or peptidomimetic having at least 100 residues.
70. The method of claim 42, wherein said clostridial toxin substrate is a peptide or peptidomimetic having at least 200 residues.
71. The method of claim 42, wherein said sample is a crude cell lysate.
72. The method of claim 42, wherein said sample is isolated clostridial toxin.
73. The method of claim 42, wherein said sample is isolated clostridial toxin light chain.
74. The method of claim 42, wherein said sample is formulated clostridial toxin product.
75. The method of claim 74, wherein said formulated clostridial toxin product is BoNT/A, BoNT/B or BoNT/E toxin product.
76. The method of claim 75, wherein said formulated product is formulated BoNT/A toxin product.
77. The method of claim 42, wherein step (c) comprises determining the change in fluorescence polarization of said treated substrate over time.

FIG. 1




FIG. 2




FIG. 3

SNAP-25 Human	(1)	1	MAEDADMRNELIEMQ	PRADQ	LADESL	STRMLQ	LVESKD	AGIRTL	VLMD	EQGEQ	LERIE	GMDQ	IN	75		
SNAP-25 Mouse	(1)	1	MAEDADMRNELIEMQ	PRADQ	LADESL	STRMLQ	LVESKD	AGIRTL	VLMD	EQGEQ	LERIE	GMDQ	IN			
SNAP-25 Drosophila	(1)	1	MPADPEEVAVQVPEKLEIEE	QINAQ	GVAD	ESL	STRMLQ	VEESKD	AGIRTL	VLMD	EQGEQ	LERIE	GMDQ	IN		
SNAP-25 Goldfish	(1)	1	MAEDADMRNELIEMQ	PRADQ	LADESL	STRMLQ	LVESKD	AGIRTL	VLMD	EQGEQ	LERIE	GMDQ	IN			
SNAP-25 Sea Urchin	(1)	1	MEDQNDKMRSELIEE	QWOSNMQ	DES	LESTRMSQ	VAEESQ	MGIRK	ITLVLMD	EQGEQ	LERIE	GMDQ	IN			
SNAP-25 Chicken	(1)	1	MAEDADMRNELIEMQ	PRADQ	LADESL	STRMLQ	LVESKD	AGIRTL	VLMD	EQGEQ	LERIE	GMDQ	IN			
SNAP-25 Human	(69)	76	KDMKEAEKNI	TDIGKFCGL	CVCPCN	---	KLKSSD	PAKKK	AMGNNQ	DGVVAS	---	QPARVV	DEREQMAISGGF	IRRVTN	150	
SNAP-25 Mouse	(69)		KDMKEAEKNI	TDIGKFCGL	CVCPCN	---	KLKSSD	PAKKK	AMGNNQ	DGVVAS	---	QPARVV	DEREQMAISGGF	IRRVTN		
SNAP-25 Drosophila	(76)		ADMPEAEKNI	SGMEKCCGFCV	PCNKSQ	SPKEDD	---	TKGND	DGKVVNNQ	PEQRYV	TD	PKNG	WVAQA	YIGRI	IN	
SNAP-25 Goldfish	(69)		KDMKEAEKNI	TDIGKFCGL	CVCPCN	---	KLKSSD	PAKKK	AMGNNQ	DGVVAS	---	QPARVV	DEREQMAISGGF	IRRVTN		
SNAP-25 Sea Urchin	(71)		TDMPPEAEKNI	TDIGKFCGL	CVCPCN	---	KLKSSD	PAKKK	AMGNNQ	DGVVAS	---	QPARVV	DEREQMAISGGF	IRRVTN		
SNAP-25 Chicken	(69)		QDMKEAEKNI	KDIGKCCGFC	PCN	---	KLKSSD	PAKKK	AMGNNQ	DGVVAS	---	QPARVV	DEREQMAISGGF	IRRVTN		
SNAP-25 Human	(140)	151	DARENEMDEN	LEQVSGIT	IGNLRH	WALDMGNE	IDTONRQ	IDRIMEK	ADSNKTR	IDEAN	---	QRA	PKML	GSG	225	
SNAP-25 Mouse	(140)		DARENEMDEN	LEQVSGIT	IGNLRH	WALDMGNE	IDTONRQ	IDRIMEK	ADSNKTR	IDEAN	---	QRA	PKML	GSG		
SNAP-25 Drosophila	(148)		DAREME	ENMGQVNT	MTIGNLRH	WALDMGSE	LENONRQ	IDRIMR	KGESNEAR	IAVAN	---	QRA	HOIL	K		
SNAP-25 Goldfish	(137)		DARENEMDEN	LEQVSGIT	IGNLRH	WALDMGNE	IDTONRQ	IDRIMEK	ADSNKTR	IDEAN	---	QRA	PKML	GSG		
SNAP-25 Sea Urchin	(146)		DARE	EMDEN	LEQVSGIT	IGNLRH	WALDMGSE	LENONRQ	IDRIMEK	ADSNKTR	IDEAN	---	QRA	PKML	GSG	
SNAP-25 Chicken	(140)		DARENEMDEN	LEQVSGIT	IGNLRH	WALDMGNE	IDTONRQ	IDRIMEK	ADSNKTR	IDEAN	---	QRA	PKML	GSG		
SNAP-25 Human	(207)	226	---	---	---	---	---	---	---	---	---	---	---	---		
SNAP-25 Mouse	(207)		---	---	---	---	---	---	---	---	---	---	---	---		
SNAP-25 Drosophila	(213)		---	---	---	---	---	---	---	---	---	---	---	---		
SNAP-25 Goldfish	(204)		---	---	---	---	---	---	---	---	---	---	---	---		
SNAP-25 Urchin	(213)		---	---	---	---	---	---	---	---	---	---	---	---		
SNAP-25 Chicken	(215)		IMLSQ	RAVP	SCFYHGI	YLLGLHTCT	YQPHCKCCPV	---	---	---	---	---	---	---		

FIG. 4

VAMP-1 HUMAN	(1)	1	MSAPAQPEAEGTEGTAHCGGGPRGPPPNMTSNRRLQQTQAQVEVVDILRVNVDKVLERDQKLSLDDRADALQA	75
VAMP-2 HUMAN	(1)		---MSATAATAPPAAPAGEGPPAPPPNLTNSNRLQQTQAQVDEVVDIMRVNVDKVLERDQKLSLDDRADALQA	
VAMP-2 MOUSE	(1)		---MSATAATVPPAAPAGEGPPAPPPNLTNSNRLQQTQAQVDEVVDIMRVNVDKVLERDQKLSLDDRADALQA	
VAMP Bovine	(1)		---MSATAATAPPAAPAGEGPPAPPPNLTNSNRLQQTQAQVDEVVDIMRVNVDKVLERDQKLSLDDRADALQA	
VAMP-2 Frog	(1)		---MSAPFAAGPPAALAFGSAFHQGPFLNLTNSNRLQQTQAQVDEVVDIMRVNVDKVLERDQKLSLDDRADALQA	
VAMP Sea Urchin	(1)		-----MAAPPPFQPAFSSNKRRLQQTQAQVDEVVDIMRVNVDKVLERDQALSLVLLDDRADALQA	
VAMP-1 HUMAN	(75)	76	GASQFESGSAKKLKRKYWWKNCKMMIMLGAICAITVVVIVIFETI	123
VAMP-2 HUMAN	(73)		GASQFETSAKKLKRKYWWKNLKMIIILGVICAIILIIIVYFS---	
VAMP-2 MOUSE	(73)		GASQFETSAKKLKRKYWWKNLKMIIILGVICAIILIIIVYFS---	
VAMP Bovine	(73)		GASQFETSAKKLKRKYWWKNLKMIIILGVICAIILIIIVYFS---	
VAMP-2 Frog	(71)		GASQFETSAKKLKRKYWWKNCKMMIIMGVICAIILIIIVYFS---	
VAMP Sea Urchin	(57)		GASQFETINAEKLRKYWWKNCKMMIILAIILIVLIIIVVAVIVQSOKK	

FIG. 5

Syntaxin 1A human	(1)	1	-MKDRIDQLRIQAK-DSDDDDVAVIVD-RDRFMDFFFEQVEEIRGFTDKIAENVEEVRKHSAILIASPNPDEKTK	75
Syntaxin 1B2 human	(1)		-MKDRIDQLRIQAK-DSDDDEEV-NHND-RDRFMDFFFEQVEEIRGFTDKIAENVEEVRKHSAILIASPNPDEKTK	
Syntaxin 1A mouse	(1)		-MKDRIDQLRIQAK-DSDDDDVAVIVD-RDRFMDFFFEQVEEIRGFTDKIAENVEEVRKHSAILIASPNPDEKTK	
Syntaxin 1a drosophila	(1)		MKDRIDQLRIQAK-DSDDDEEV-NHND-RDRFMDFFFEQVEEIRGFTDKIAENVEEVRKHSAILIASPNPDEKTK	
Syntaxin A C. elegans	(1)		MKDRIDQLRIQAK-DSDDDDVAVIVD-RDRFMDFFFEQVEEIRGFTDKIAENVEEVRKHSAILIASPNPDEKTK	
Syntaxin Sea urchin	(1)		-MKDRIDQLRIQAK-EDDVGPVAVIVD-SEKEMFFFEQVEEIRGFTDKIAENVEEVRKHSAILIASPNPDEKTK	
Syntaxin 1A human	(73)	76	EELEELMSDIIKKTANKVRSKLKSIEQSIIEQEGINRSSADLRIRKTOHSTLSRKFEVWVSEYNATQSDYRERCKG	150
Syntaxin 1B2 human	(72)		QELDELADIKKTANKVRSKLKALIEQSIIEQEGINRSSADLRIRKTOHSTLSRKFEVWVTEYNATQSKYRDRCKD	
Syntaxin 1A mouse	(73)		EELEELMSDIIKKTANKVRSKLKSIEQSIIEQEGINRSSADLRIRKTOHSTLSRKFEVWVSEYNATQSDYRERCKG	
Syntaxin 1a drosophila	(76)		QELDELADIKKTANKVRSKLKALIEQSIIEQEGINRSSADLRIRKTOHSTLSRKFEVWVTEYNATQSDYRERCKG	
Syntaxin A C. elegans	(75)		EELEELMSDIIKKTANKVRSKLKSIEQSIIEQEGINRSSADLRIRKTOHSTLSRKFEVWVTEYNATQSDYRERCKG	
Syntaxin Sea urchin	(73)		DELEELMSDIIKKTANKVRSKLKALIEQSIIEQEGINRSSADLRIRKTOHSTLSRKFEVWVTEYNATQSDYRERCKG	
Syntaxin 1A human	(148)	151	RIQRQLETTGRTTISEELEMDLSEGNPAITAGIIMDSISKQALISEIIRHSEIKLENSIRELHDMFMDMAMTL	225
Syntaxin 1B2 human	(147)		RIQRQLETTGRTTISEELEMDLSEGNPAITAGIIMDSISKQALISEIIRHSEIKLENSIRELHDMFMDMAMTL	
Syntaxin 1A mouse	(148)		RIQRQLETTGRTTISEELEMDLSEGNPAITAGIIMDSISKQALISEIIRHSEIKLENSIRELHDMFMDMAMTL	
Syntaxin 1a drosophila	(151)		RIQRQLETTGRTTISEELEMDLSEGNPAITAGIIMDSISKQALISEIIRHSEIKLENSIRELHDMFMDMAMTL	
Syntaxin A C. elegans	(149)		RIQRQLETTGRTTISEELEMDLSEGNPAITAGIIMDSISKQALISEIIRHSEIKLENSIRELHDMFMDMAMTL	
Syntaxin Sea urchin	(148)		RIQRQLETTGRTTISEELEMDLSEGNPAITAGIIMDSISKQALISEIIRHSEIKLENSIRELHDMFMDMAMTL	
Syntaxin 1A human	(223)	226	VESQGENMDRIEYVNEHADVVERAVSDTKKAVKYQSKARRKKIMIIICCVILGLVJASIVGGLFA--	293
Syntaxin 1B2 human	(222)		VESQGENMDRIEYVNEHADVVERAVSDTKKAVKYQSKARRKKIMIIICCVILGLVJASIVGGLFA--	
Syntaxin 1A mouse	(223)		VESQGENMDRIEYVNEHADVVERAVSDTKKAVKYQSKARRKKIMIIICCVILGLVJASIVGGLFA--	
Syntaxin 1a drosophila	(226)		VESQGENMDRIEYVNEHADVVERAVSDTKKAVKYQSKARRKKIMIIICCVILGLVJASIVGGLFA--	
Syntaxin A C. elegans	(224)		VESQGENMDRIEYVNEHADVVERAVSDTKKAVKYQSKARRKKIMIIICCVILGLVJASIVGGLFA--	
Syntaxin Sea urchin	(223)		VESQGENMDRIEYVNEHADVVERAVSDTKKAVKYQSKARRKKIMIIICCVILGLVJASIVGGLFA--	


FIG. 6A

FIG. 6B

```

                                     GFP
~~~~~
M A S K G E E L F T G V V P I L V E L D G
ATGGCTAGC AAAGGAGAA GAACTCTTC ACTGGAGTT GTCCCAATT CTGTTGAA TTAGATGGT
                                     GFP
~~~~~
D V N G H K F S V S G E G E G D A T Y G K
GATGTTAAC GGCCACAAG TTCTCTGTC AGTGGAGAG GGTGAAGGT GATGCAACA TACGGAAAA
                                     GFP
~~~~~
L T L K F I C T T G K L P V P W P T L V T
CTTACCCTG AAGTTCATC TGCACTACT GGCAAACTG CCTGTTCCA TGGCCAACA CTAGTCACT
                                     GFP
~~~~~
T L C Y G V Q C F S R Y P D H M K R H D F
ACTCTGTGC TATGGTGT TCAATGCTTT TCAAGATAC CCGGATCAT ATGAAACGG CATGACTTT
                                     GFP
~~~~~
F K S A M P E G Y V Q E R T I F F K D D G
TTCAAGAGT GCCATGCCC GAAGGTTAT GTACAGGAA AGGACCATC TTCTTCAA GATGACGGC
                                     GFP
~~~~~
N Y K T R A E V K F E G D T L V N R I E L
AACTACAAG ACACGTGCT GAAGTCAAG TTTGAAGGT GATACCCTT GTTAATAGA ATCGAGTTA
                                     GFP
~~~~~
K G I D F K E D G N I L G H K L E Y N Y N
AAAGGTATT GACTTCAAG GAAGATGGC AACATTCTG GGACACAAA TTGGAATAC AACTATAAC
                                     GFP
~~~~~
S H N V Y I M A D K Q K N G I K V N F K T
TCACACAAT GTATACATC ATGGCAGAC AAACAAAAG AATGGAATC AAAGTGAAC TTCAAGACC
                                     GFP
~~~~~
R H N I E D G S V Q L A D H Y Q Q N T P I
CGCCACAAC ATTGAAGAT GGAAGCGTT CAAC TAGCA GACCATTAT CAACAAAAT ACTCCAATT
                                     GFP
~~~~~
G D G P V L L P D N H Y L S T Q S A L S K
GGCGATGGC CCTGTCTT TTACCAGAC AACCATTAC CTGTCCACA CAATCTGCC CTTTCGAAA
                                     GFP
~~~~~
D P N E K R D H M V L L E F V T A A G I T
GATCCCAAC GAAAAGAGA GACCACATG GTCCTTCTT GAGTTTGTA ACAGCTGCT GGGATTACA
                                     Linker
                                     GFP
~~~~~
H G M D E L Y N G G A G S G A G G G G I R
CATGGCATG GATGAACTG TACAACGGC GGTGCAGGA TCCGGTGCG GGTGGCGGT GGCATCCGG
                                     SNAP25 (134-206)
~~~~~
R V T N D A R E N E M D E N L E Q V S G I
AGGGTAACA AACGATGCC CGGGAAAAT GAGATGGAT GAGAACCTG GAGCAGGTG AGCGGCATC
                                     SNAP25 (134-206)
~~~~~
I G N L R H M A L D M G N E I D T Q N R Q
ATCGGAAAC CTCCGCCAT ATGGCTCTA GACATGGGC AATGAGATT GACACCCAG AATCGCCAG
                                     SNAP25 (134-206)
~~~~~
I D R I M E K A D S N K T R I D E A N Q R
ATCGACAGG ATCATGGAG AAGGCTGAT TCCAACAAA ACCAGAATT GATGAAGCC AACCAACGT
                                     Linker
                                     Cys
                                     SNAP25 (134-206)
~~~~~
A T K M L G S G G G G G H H H H H H C *
GCAACAAAG ATGCTGGGA AGTGGTGGC GGTGGCGGC CATCACCAT CACCATCAC TGCTAA

```

FIG. 7A




FIG. 7B




FIG. 8A

Rxtn Conditions: GFP-SNAP-His6-C 100ul at 2.74mg/ml. Overnight
4C with 3.6ul 22mM Alexa594 in DMF





FIG. 8B

FIG. 9A

**Regression Cubic Hill Plot 4 Parameters:
20ul of BCK2034 Reduced BoNT-A, 37 °C
1750 ng/ml**




FIG. 9B

**Regression Cubic Hill Plot 4 Parameters:
150ul 100:1 dilution of BCK 2034, 37 °C
131.25 ng/ml**




FIG. 9C

**Regression Cubic Hill Plot 4 Parameters:
100ul of 100:1 dilution of BCK2034, 37 °C
87.5 ng/ml**




FIG. 9D

Regression Cubic Hill Plot 4 Parameters:
60ul of 100-1 dilution of BCK2034, 37 °C
52.5 ng/ml




FIG. 9E

Regression Cubic Hill Plot 4 Parameters:
20ul of 100:1 dilution of BCK2034, 37 °C
17.5 ng/ml






FIG. 10A

FIG. 10B

SEQUENCE LISTING

<110> Verhagen, Marc
 Williams, Dudley J.
 Gilmore, Marcella
 Steward, Lance
 Aoki, Kei Roger

<120> Fluorescence Polarization Assays For
 Determining Clostridial Toxin Activity

<130> 66872-040

<160> 96

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 206

<212> PRT

<213> Homo sapiens

<400> 1

Met	Ala	Glu	Asp	Ala	Asp	Met	Arg	Asn	Glu	Leu	Glu	Glu	Met	Gln	Arg	1	5	10	15
Arg	Ala	Asp	Gln	Leu	Ala	Asp	Glu	Ser	Leu	Glu	Ser	Thr	Arg	Arg	Met	20	25	30	
Leu	Gln	Leu	Val	Glu	Glu	Ser	Lys	Asp	Ala	Gly	Ile	Arg	Thr	Leu	Val	35	40	45	
Met	Leu	Asp	Glu	Gln	Gly	Glu	Gln	Leu	Glu	Arg	Ile	Glu	Glu	Gly	Met	50	55	60	
Asp	Gln	Ile	Asn	Lys	Asp	Met	Lys	Glu	Ala	Glu	Lys	Asn	Leu	Thr	Asp	65	70	75	80
Leu	Gly	Lys	Phe	Cys	Gly	Leu	Cys	Val	Cys	Pro	Cys	Asn	Lys	Leu	Lys	85	90	95	
Ser	Ser	Asp	Ala	Tyr	Lys	Lys	Ala	Trp	Gly	Asn	Asn	Gln	Asp	Gly	Val	100	105	110	
Val	Ala	Ser	Gln	Pro	Ala	Arg	Val	Val	Asp	Glu	Arg	Glu	Gln	Met	Ala	115	120	125	
Ile	Ser	Gly	Gly	Phe	Ile	Arg	Arg	Val	Thr	Asn	Asp	Ala	Arg	Glu	Asn	130	135	140	
Glu	Met	Asp	Glu	Asn	Leu	Glu	Gln	Val	Ser	Gly	Ile	Ile	Gly	Asn	Leu	145	150	155	160
Arg	His	Met	Ala	Leu	Asp	Met	Gly	Asn	Glu	Ile	Asp	Thr	Gln	Asn	Arg	165	170	175	
Gln	Ile	Asp	Arg	Ile	Met	Glu	Lys	Ala	Asp	Ser	Asn	Lys	Thr	Arg	Ile	180	185	190	
Asp	Glu	Ala	Asn	Gln	Arg	Ala	Thr	Lys	Met	Leu	Gly	Ser	Gly			195	200	205	

<210> 2

<211> 206

<212> PRT

<213> Mus musculus

<400> 2

Met	Ala	Glu	Asp	Ala	Asp	Met	Arg	Asn	Glu	Leu	Glu	Glu	Met	Gln	Arg	1	5	10	15
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	---	---	----	----

```

Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
      20      25      30
Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
      35      40      45
Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met
      50      55      60
Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp
      65      70      75      80
Leu Gly Lys Phe Cys Gly Leu Cys Val Cys Pro Cys Asn Lys Leu Lys
      85      90      95
Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
      100      105      110
Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
      115      120      125
Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn
      130      135      140
Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile Gly Asn Leu
      145      150      155      160
Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg
      165      170      175
Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys Thr Arg Ile
      180      185      190
Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly
      195      200      205

```

<210> 3

<211> 212

<212> PRT

<213> Drosophila sp.

<400> 3

```

Met Pro Ala Asp Pro Ser Glu Glu Val Ala Pro Gln Val Pro Lys Thr
  1      5      10      15
Glu Leu Glu Glu Leu Gln Ile Asn Ala Gln Gly Val Ala Asp Glu Ser
      20      25      30
Leu Glu Ser Thr Arg Arg Met Leu Ala Leu Cys Glu Glu Ser Lys Glu
      35      40      45
Ala Gly Ile Arg Thr Leu Val Ala Leu Asp Asp Gln Gly Glu Gln Leu
      50      55      60
Asp Arg Ile Glu Glu Gly Met Asp Gln Ile Asn Ala Asp Met Arg Glu
      65      70      75      80
Ala Glu Lys Asn Leu Ser Gly Met Glu Lys Cys Cys Gly Ile Cys Val
      85      90      95
Leu Pro Cys Asn Lys Ser Gln Ser Phe Lys Glu Asp Asp Gly Thr Trp
      100      105      110
Lys Gly Asn Asp Asp Gly Lys Val Val Asn Asn Gln Pro Gln Arg Val
      115      120      125
Met Asp Asp Arg Asn Gly Met Met Ala Gln Ala Gly Tyr Ile Gly Arg
      130      135      140
Ile Thr Asn Asp Ala Arg Glu Asp Glu Met Glu Glu Asn Met Gly Gln
      145      150      155      160
Val Asn Thr Met Ile Gly Asn Leu Arg Asn Met Ala Leu Asp Met Gly
      165      170      175
Ser Glu Leu Glu Asn Gln Asn Arg Gln Ile Asp Arg Ile Asn Arg Lys
      180      185      190
Gly Glu Ser Asn Glu Ala Arg Ile Ala Val Ala Asn Gln Arg Ala His
      195      200      205
Gln Leu Leu Lys

```

210

<210> 4
 <211> 203
 <212> PRT
 <213> Carassius auratus

<400> 4

Met Ala Asp Glu Ala Asp Met Arg Asn Glu Leu Thr Asp Met Gln Ala

1	5	10	15
Arg Ala Asp Gln Leu Gly Asp Glu Ser Leu Glu Ser Thr Arg Arg Met			
	20	25	30
Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val			
	35	40	45
Met Leu Asp Glu Gln Gly Glu Gln Leu Glu Arg Ile Glu Glu Gly Met			
	50	55	60
Asp Gln Ile Asn Lys Asp Met Lys Glu Ala Glu Lys Asn Leu Thr Asp			
65	70	75	80
Leu Gly Asn Leu Cys Gly Leu Cys Pro Cys Pro Cys Asn Lys Leu Lys			
	85	90	95
Gly Gly Gly Gln Ser Trp Gly Asn Asn Gln Asp Gly Val Val Ser Ser			
	100	105	110
Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala Ile Ser Gly			
	115	120	125
Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn Glu Met Asp			
	130	135	140
Glu Asn Leu Glu Gln Val Gly Ser Ile Ile Gly Asn Leu Arg His Met			
145	150	155	160
Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg Gln Ile Asp			
	165	170	175
Arg Ile Met Asp Met Ala Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala			
	180	185	190
Asn Gln Arg Ala Thr Lys Met Leu Gly Ser Gly			
	195	200	

<210> 5
 <211> 212
 <212> PRT
 <213> Strongylcentrotas purpuratus

<400> 5

Met Glu Asp Gln Asn Asp Met Asn Met Arg Ser Glu Leu Glu Glu Ile	
1	5
Gln Met Gln Ser Asn Met Gln Thr Asp Glu Ser Leu Glu Ser Thr Arg	
	20
Arg Met Leu Gln Met Ala Glu Glu Ser Gln Asp Met Gly Ile Lys Thr	
	35
Leu Val Met Leu Asp Glu Gln Gly Glu Gln Leu Asp Arg Ile Glu Glu	
	50
Gly Met Asp Gln Ile Asn Thr Asp Met Arg Glu Ala Glu Lys Asn Leu	
65	70
Thr Gly Leu Glu Lys Cys Cys Gly Ile Cys Val Cys Pro Trp Lys Lys	
	85
Leu Gly Asn Phe Glu Lys Gly Asp Asp Tyr Lys Lys Thr Trp Lys Gly	
	100
Asn Asp Asp Gly Lys Val Asn Ser His Gln Pro Met Arg Met Glu Asp	
	115
	120
	125

Asp Arg Asp Gly Cys Gly Gly Asn Ala Ser Met Ile Thr Arg Ile Thr
 130 135 140
 Asn Asp Ala Arg Glu Asp Glu Met Asp Glu Asn Leu Thr Gln Val Ser
 145 150 155 160
 Ser Ile Val Gly Asn Leu Arg His Met Ala Ile Asp Met Gln Ser Glu
 165 170 175
 Ile Gly Ala Gln Asn Ser Gln Val Gly Arg Ile Thr Ser Lys Ala Glu
 180 185 190
 Ser Asn Glu Gly Arg Ile Asn Ser Ala Asp Lys Arg Ala Lys Asn Ile
 195 200 205
 Leu Arg Asn Lys
 210

<210> 6
 <211> 249
 <212> PRT
 <213> Gallus gallus

<400> 6
 Met Ala Glu Asp Ala Asp Met Arg Asn Glu Leu Glu Glu Met Gln Arg
 1 5 10 15
 Arg Ala Asp Gln Leu Ala Asp Glu Ser Leu Glu Ser Thr Arg Arg Met
 20 25 30
 Leu Gln Leu Val Glu Glu Ser Lys Asp Ala Gly Ile Arg Thr Leu Val
 35 40 45
 Met Leu Asp Glu Gln Gly Glu Gln Leu Asp Arg Val Glu Glu Gly Met
 50 55 60
 Asn His Ile Asn Gln Asp Met Lys Glu Ala Glu Lys Asn Leu Lys Asp
 65 70 75 80
 Leu Gly Lys Cys Cys Gly Leu Phe Ile Cys Pro Cys Asn Lys Leu Lys
 85 90 95
 Ser Ser Asp Ala Tyr Lys Lys Ala Trp Gly Asn Asn Gln Asp Gly Val
 100 105 110
 Val Ala Ser Gln Pro Ala Arg Val Val Asp Glu Arg Glu Gln Met Ala
 115 120 125
 Ile Ser Gly Gly Phe Ile Arg Arg Val Thr Asn Asp Ala Arg Glu Asn
 130 135 140
 Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly Ile Ile Gly Asn Leu
 145 150 155 160
 Arg His Met Ala Leu Asp Met Gly Asn Glu Ile Asp Thr Gln Asn Arg
 165 170 175
 Gln Ile Asp Arg Ile Met Glu Lys Leu Ile Pro Ile Lys Pro Gly Leu
 180 185 190
 Met Lys Pro Thr Ser Val Gln Gln Arg Cys Ser Ala Val Val Lys Cys
 195 200 205
 Ser Lys Val His Phe Leu Leu Met Leu Ser Gln Arg Ala Val Pro Ser
 210 215 220
 Cys Phe Tyr His Gly Ile Tyr Leu Leu Gly Leu His Thr Cys Thr Tyr
 225 230 235 240
 Gln Pro His Cys Lys Cys Cys Pro Val
 245

<210> 7
 <211> 118
 <212> PRT
 <213> Homo sapiens

<400> 7

```

Met Ser Ala Pro Ala Gln Pro Pro Ala Glu Gly Thr Glu Gly Thr Ala
 1      5      10      15
Pro Gly Gly Gly Pro Pro Gly Pro Pro Asn Met Thr Ser Asn Arg
      20      25      30
Arg Leu Gln Gln Thr Gln Ala Gln Val Glu Glu Val Val Asp Ile Ile
      35      40      45
Arg Val Asn Val Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu
      50      55      60
Leu Asp Asp Arg Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu
65      70      75      80
Ser Ser Ala Ala Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Cys Lys
      85      90      95
Met Met Ile Met Leu Gly Ala Ile Cys Ala Ile Ile Val Val Val Ile
      100      105      110
Val Ile Tyr Phe Phe Thr
      115

```

<210> 8

<211> 116

<212> PRT

<213> Homo sapiens

<400> 8

```

Met Ser Ala Thr Ala Ala Thr Ala Pro Pro Ala Ala Pro Ala Gly Glu
 1      5      10      15
Gly Gly Pro Pro Ala Pro Pro Pro Asn Leu Thr Ser Asn Arg Arg Leu
      20      25      30
Gln Gln Thr Gln Ala Gln Val Asp Glu Val Val Asp Ile Met Arg Val
      35      40      45
Asn Val Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp
      50      55      60
Asp Arg Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser
65      70      75      80
Ala Ala Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Leu Lys Met Met
      85      90      95
Ile Ile Leu Gly Val Ile Cys Ala Ile Ile Leu Ile Ile Ile Val
      100      105      110
Tyr Phe Ser Ser
      115

```

<210> 9

<211> 116

<212> PRT

<213> Mus musculus

<400> 9

```

Met Ser Ala Thr Ala Ala Thr Val Pro Pro Ala Ala Pro Ala Gly Glu
 1      5      10      15
Gly Gly Pro Pro Ala Pro Pro Pro Asn Leu Thr Ser Asn Arg Arg Leu
      20      25      30
Gln Gln Thr Gln Ala Gln Val Asp Glu Val Val Asp Ile Met Arg Val
      35      40      45
Asn Val Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp
      50      55      60
Asp Arg Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser
65      70      75      80
Ala Ala Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Leu Lys Met Met
      85      90      95

```

Ile Ile Leu Gly Val Ile Cys Ala Ile Ile Leu Ile Ile Ile Ile Val
 100 105 110
 Tyr Phe Ser Thr
 115

<210> 10
 <211> 116
 <212> PRT
 <213> Bos taurus

<400> 10
 Met Ser Ala Thr Ala Ala Thr Ala Pro Pro Ala Ala Pro Ala Gly Glu
 1 5 10 15
 Gly Gly Pro Pro Ala Pro Pro Pro Asn Leu Thr Ser Asn Arg Arg Leu
 20 25 30
 Gln Gln Thr Gln Ala Gln Val Asp Glu Val Val Asp Ile Met Arg Val
 35 40 45
 Asn Val Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp
 50 55 60
 Asp Arg Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser
 65 70 75 80
 Ala Ala Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Leu Lys Met Met
 85 90 95
 Ile Ile Leu Gly Val Ile Cys Ala Ile Ile Leu Ile Ile Ile Ile Val
 100 105 110
 Tyr Phe Ser Ser
 115

<210> 11
 <211> 114
 <212> PRT
 <213> Xenopus laevis

<400> 11
 Met Ser Ala Pro Ala Ala Gly Pro Pro Ala Ala Ala Pro Gly Asp Gly
 1 5 10 15
 Ala Pro Gln Gly Pro Pro Asn Leu Thr Ser Asn Arg Arg Leu Gln Gln
 20 25 30
 Thr Gln Ala Gln Val Asp Glu Val Val Asp Ile Met Arg Val Asn Val
 35 40 45
 Asp Lys Val Leu Glu Arg Asp Thr Lys Leu Ser Glu Leu Asp Asp Arg
 50 55 60
 Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser Ala Ala
 65 70 75 80
 Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Met Lys Met Met Ile Ile
 85 90 95
 Met Gly Val Ile Cys Ala Ile Ile Leu Ile Ile Ile Ile Val Tyr Phe
 100 105 110
 Ser Thr

<210> 12
 <211> 104
 <212> PRT
 <213> Strongylocentrotus purpuratus

<400> 12
 Met Ala Ala Pro Pro Pro Pro Gln Pro Ala Pro Ser Asn Lys Arg Leu

```

      1           5           10           15
Gln Gln Thr Gln Ala Gln Val Asp Glu Val Val Asp Ile Met Arg Val
      20           25           30
Asn Val Asp Lys Val Leu Glu Arg Asp Gln Ala Leu Ser Val Leu Asp
      35           40           45
Asp Arg Ala Asp Ala Leu Gln Gln Gly Ala Ser Gln Phe Glu Thr Asn
      50           55           60
Ala Gly Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Cys Lys Met Met
      65           70           75           80
Ile Ile Leu Ala Ile Ile Ile Ile Val Ile Leu Ile Ile Ile Ile Val
      85           90           95
Ala Ile Val Gln Ser Gln Lys Lys
      100

```

<210> 13
 <211> 288
 <212> PRT
 <213> Homo sapiens

```

<400> 13
Met Lys Asp Arg Thr Gln Glu Leu Arg Thr Ala Lys Asp Ser Asp Asp
      1           5           10           15
Asp Asp Asp Val Ala Val Thr Val Asp Arg Asp Arg Phe Met Asp Glu
      20           25           30
Phe Phe Glu Gln Val Glu Glu Ile Arg Gly Phe Ile Asp Lys Ile Ala
      35           40           45
Glu Asn Val Glu Glu Val Lys Arg Lys His Ser Ala Ile Leu Ala Ser
      50           55           60
Pro Asn Pro Asp Glu Lys Thr Lys Glu Glu Leu Glu Glu Leu Met Ser
      65           70           75           80
Asp Ile Lys Lys Thr Ala Asn Lys Val Arg Ser Lys Leu Lys Ser Ile
      85           90           95
Glu Gln Ser Ile Glu Gln Glu Glu Gly Leu Asn Arg Ser Ser Ala Asp
      100           105           110
Leu Arg Ile Arg Lys Thr Gln His Ser Thr Leu Ser Arg Lys Phe Val
      115           120           125
Glu Val Met Ser Glu Tyr Asn Ala Thr Gln Ser Asp Tyr Arg Glu Arg
      130           135           140
Cys Lys Gly Arg Ile Gln Arg Gln Leu Glu Ile Thr Gly Arg Thr Thr
      145           150           155           160
Thr Ser Glu Glu Leu Glu Asp Met Leu Glu Ser Gly Asn Pro Ala Ile
      165           170           175
Phe Ala Ser Gly Ile Ile Met Asp Ser Ser Ile Ser Lys Gln Ala Leu
      180           185           190
Ser Glu Ile Glu Thr Arg His Ser Glu Ile Ile Lys Leu Glu Asn Ser
      195           200           205
Ile Arg Glu Leu His Asp Met Phe Met Asp Met Ala Met Leu Val Glu
      210           215           220
Ser Gln Gly Glu Met Ile Asp Arg Ile Glu Tyr Asn Val Glu His Ala
      225           230           235           240
Val Asp Tyr Val Glu Arg Ala Val Ser Asp Thr Lys Lys Ala Val Lys
      245           250           255
Tyr Gln Ser Lys Ala Arg Arg Lys Lys Ile Met Ile Ile Ile Cys Cys
      260           265           270
Val Ile Leu Gly Ile Val Ile Ala Ser Thr Val Gly Gly Ile Phe Ala
      275           280           285

```

<210> 14

<211> 288

<212> PRT

<213> Homo sapiens

<400> 14

```

Met Lys Asp Arg Thr Gln Glu Leu Arg Ser Ala Lys Asp Ser Asp Asp
 1           5           10           15
Glu Glu Glu Val Val His Val Asp Arg Asp His Phe Met Asp Glu Phe
           20           25           30
Phe Glu Gln Val Glu Glu Ile Arg Gly Cys Ile Glu Lys Leu Ser Glu
           35           40           45
Asp Val Glu Gln Val Lys Lys Gln His Ser Ala Ile Leu Ala Ala Pro
           50           55           60
Asn Pro Asp Glu Lys Thr Lys Gln Glu Leu Glu Asp Leu Thr Ala Asp
65           70           75           80
Ile Lys Lys Thr Ala Asn Lys Val Arg Ser Lys Leu Lys Ala Ile Glu
           85           90           95
Gln Ser Ile Glu Gln Glu Gly Leu Asn Arg Ser Ser Ala Asp Leu
           100          105          110
Arg Ile Arg Lys Thr Gln His Ser Thr Leu Ser Arg Lys Phe Val Glu
           115          120          125
Val Met Thr Glu Tyr Asn Ala Thr Gln Ser Lys Tyr Arg Asp Arg Cys
           130          135          140
Lys Asp Arg Ile Gln Arg Gln Leu Glu Ile Thr Gly Arg Thr Thr Thr
145           150           155           160
Asn Glu Glu Leu Glu Asp Met Leu Glu Ser Gly Lys Leu Ala Ile Phe
           165          170          175
Thr Asp Asp Ile Lys Met Asp Ser Gln Met Thr Lys Gln Ala Leu Asn
           180          185          190
Glu Ile Glu Thr Arg His Asn Glu Ile Ile Lys Leu Glu Thr Ser Ile
           195          200          205
Arg Glu Leu His Asp Met Phe Val Asp Met Ala Met Leu Val Glu Ser
           210          215          220
Gln Gly Glu Met Ile Asp Arg Ile Glu Tyr Asn Val Glu His Ser Val
225           230           235           240
Asp Tyr Val Glu Arg Ala Val Ser Asp Thr Lys Lys Ala Val Lys Tyr
           245          250          255
Gln Ser Lys Ala Arg Arg Lys Lys Ile Met Ile Ile Ile Cys Cys Val
           260          265          270
Val Leu Gly Val Val Leu Ala Ser Ile Gly Gly Thr Leu Gly Leu
           275          280          285

```

<210> 15

<211> 288

<212> PRT

<213> Mus musculus

<400> 15

```

Met Lys Asp Arg Thr Gln Glu Leu Arg Thr Ala Lys Asp Ser Asp Asp
 1           5           10           15
Asp Asp Asp Val Thr Val Thr Val Asp Arg Asp Arg Phe Met Asp Glu
           20           25           30
Phe Phe Glu Gln Val Glu Glu Ile Arg Gly Phe Ile Asp Lys Ile Ala
           35           40           45
Glu Asn Val Glu Glu Val Lys Arg Lys His Ser Ala Ile Leu Ala Ser
           50           55           60
Pro Asn Pro Asp Glu Lys Thr Lys Glu Glu Leu Glu Glu Leu Met Ser
65           70           75           80

```

Asp Ile Lys Lys Thr Ala Asn Lys Val Arg Ser Lys Leu Lys Ser Ile
 85 90 95
 Glu Gln Ser Ile Glu Gln Glu Glu Gly Leu Asn Arg Ser Ser Ala Asp
 100 105 110
 Leu Arg Ile Arg Lys Thr Gln His Ser Thr Leu Ser Arg Lys Phe Val
 115 120 125
 Glu Val Met Ser Glu Tyr Asn Ala Thr Gln Ser Asp Tyr Arg Glu Arg
 130 135 140
 Cys Lys Gly Arg Ile Gln Arg Gln Leu Glu Ile Thr Gly Arg Thr Thr
 145 150 155 160
 Thr Ser Glu Glu Leu Glu Asp Met Leu Glu Ser Gly Asn Pro Ala Ile
 165 170 175
 Phe Ala Ser Gly Ile Ile Met Asp Ser Ser Ile Ser Lys Gln Ala Leu
 180 185 190
 Ser Glu Ile Glu Thr Arg His Ser Glu Ile Ile Lys Leu Glu Thr Ser
 195 200 205
 Ile Arg Glu Leu His Asp Met Phe Met Asp Met Ala Met Leu Val Glu
 210 215 220
 Ser Gln Gly Glu Met Ile Asp Arg Ile Glu Tyr Asn Val Glu His Ala
 225 230 235 240
 Val Asp Tyr Val Glu Arg Ala Val Ser Asp Thr Lys Lys Ala Val Lys
 245 250 255
 Tyr Gln Ser Lys Ala Arg Arg Lys Lys Ile Met Ile Ile Ile Cys Cys
 260 265 270
 Val Ile Leu Gly Ile Ile Ile Ala Ser Thr Ile Gly Gly Ile Phe Gly
 275 280 285

<210> 16

<211> 291

<212> PRT

<213> Drosophila sp.

<400> 16

Met Thr Lys Asp Arg Leu Ala Ala Leu His Ala Ala Gln Ser Asp Asp
 1 5 10 15
 Glu Glu Glu Thr Glu Val Ala Val Asn Val Asp Gly His Asp Ser Tyr
 20 25 30
 Met Asp Asp Phe Phe Ala Gln Val Glu Glu Ile Arg Gly Met Ile Asp
 35 40 45
 Lys Val Gln Asp Asn Val Glu Glu Val Lys Lys Lys His Ser Ala Ile
 50 55 60
 Leu Ser Ala Pro Gln Thr Asp Glu Lys Thr Lys Gln Glu Leu Glu Asp
 65 70 75 80
 Leu Met Ala Asp Ile Lys Lys Asn Ala Asn Arg Val Arg Gly Lys Leu
 85 90 95
 Lys Gly Ile Glu Gln Asn Ile Glu Gln Glu Glu Gln Gln Asn Lys Ser
 100 105 110
 Ser Ala Asp Leu Arg Ile Arg Lys Thr Gln His Ser Thr Leu Ser Arg
 115 120 125
 Lys Phe Val Glu Val Met Thr Glu Tyr Asn Arg Thr Gln Thr Asp Tyr
 130 135 140
 Arg Glu Arg Cys Lys Gly Arg Ile Gln Arg Gln Leu Glu Ile Thr Gly
 145 150 155 160
 Arg Pro Thr Asn Asp Asp Glu Leu Glu Lys Met Leu Glu Glu Gly Asn
 165 170 175
 Ser Ser Val Phe Thr Gln Gly Ile Ile Met Glu Thr Gln Gln Ala Lys
 180 185 190
 Gln Thr Leu Ala Asp Ile Glu Ala Arg His Gln Asp Ile Met Lys Leu

195 200 205
 Glu Thr Ser Ile Lys Glu Leu His Asp Met Phe Met Asp Met Ala Met
 210 215 220
 Leu Val Glu Ser Gln Gly Glu Met Ile Asp Arg Ile Glu Tyr His Val
 225 230 235 240
 Glu His Ala Met Asp Tyr Val Gln Thr Ala Thr Gln Asp Thr Lys Lys
 245 250 255
 Ala Leu Lys Tyr Gln Ser Lys Ala Arg Arg Lys Lys Ile Met Ile Leu
 260 265 270
 Ile Cys Leu Thr Val Leu Gly Ile Leu Ala Ala Ser Tyr Val Ser Ser
 275 280 285
 Tyr Phe Met
 290

<210> 17
 <211> 291
 <212> PRT
 <213> C. elegans

<400> 17
 Met Thr Lys Asp Arg Leu Ser Ala Leu Lys Ala Ala Gln Ser Glu Asp
 1 5 10 15
 Glu Gln Asp Asp Asp Met His Met Asp Thr Gly Asn Ala Gln Tyr Met
 20 25 30
 Glu Glu Phe Phe Glu Gln Val Glu Glu Ile Arg Gly Ser Val Asp Ile
 35 40 45
 Ile Ala Asn Asn Val Glu Glu Val Lys Lys Lys His Ser Ala Ile Leu
 50 55 60
 Ser Asn Pro Val Asn Asp Gln Lys Thr Lys Glu Glu Leu Asp Glu Leu
 65 70 75 80
 Met Ala Val Ile Lys Arg Ala Ala Asn Lys Val Arg Gly Lys Leu Lys
 85 90 95
 Leu Ile Glu Asn Ala Ile Asp His Asp Glu Gln Gly Ala Gly Asn Ala
 100 105 110
 Asp Leu Arg Ile Arg Lys Thr Gln His Ser Thr Leu Ser Arg Arg Phe
 115 120 125
 Val Glu Val Met Thr Asp Tyr Asn Lys Thr Gln Thr Asp Tyr Arg Glu
 130 135 140
 Arg Cys Lys Gly Arg Ile Gln Arg Gln Leu Asp Ile Ala Gly Lys Gln
 145 150 155 160
 Val Gly Asp Glu Asp Leu Glu Glu Met Ile Glu Ser Gly Asn Pro Gly
 165 170 175
 Val Phe Thr Gln Gly Ile Ile Thr Asp Thr Gln Gln Ala Lys Gln Thr
 180 185 190
 Leu Ala Asp Ile Glu Ala Arg His Asn Asp Ile Met Lys Leu Glu Ser
 195 200 205
 Ser Ile Arg Glu Leu His Asp Met Phe Met Asp Met Ala Met Leu Val
 210 215 220
 Glu Ser Gln Gly Glu Met Val Asp Arg Ile Glu Tyr Asn Val Glu His
 225 230 235 240
 Ala Lys Glu Phe Val Asp Arg Ala Val Ala Asp Thr Lys Lys Ala Val
 245 250 255
 Gln Tyr Gln Ser Lys Ala Arg Arg Lys Lys Ile Cys Ile Leu Val Thr
 260 265 270
 Gly Val Ile Leu Ile Thr Gly Leu Ile Ile Phe Ile Leu Phe Tyr Ala
 275 280 285
 Lys Val Leu
 290

<210> 18
 <211> 288
 <212> PRT
 <213> Strongylocentrotus purpuratus

<400> 18
 Met Arg Asp Arg Leu Gly Ser Leu Lys Arg Asn Glu Glu Asp Asp Val
 1 5 10 15
 Gly Pro Glu Val Ala Val Asn Val Glu Ser Glu Lys Phe Met Glu Glu
 20 25 30
 Phe Phe Glu Gln Val Glu Glu Val Arg Asn Asn Ile Asp Lys Ile Ser
 35 40 45
 Lys Asn Val Asp Glu Val Lys Lys Lys His Ser Asp Ile Leu Ser Ala
 50 55 60
 Pro Gln Ala Asp Glu Lys Val Lys Asp Glu Leu Glu Glu Leu Met Ser
 65 70 75 80
 Asp Ile Lys Lys Thr Ala Asn Lys Val Arg Ala Lys Leu Lys Met Met
 85 90 95
 Glu Gln Ser Ile Glu Gln Glu Glu Ser Ala Lys Met Asn Ser Ala Asp
 100 105 110
 Val Arg Ile Arg Lys Thr Gln His Ser Thr Leu Ser Arg Lys Phe Val
 115 120 125
 Glu Val Met Thr Asp Tyr Asn Ser Thr Gln Thr Asp Tyr Arg Glu Arg
 130 135 140
 Cys Lys Gly Arg Ile Gln Arg Gln Leu Glu Ile Thr Gly Lys Ser Thr
 145 150 155 160
 Thr Asp Ala Glu Leu Glu Asp Met Leu Glu Ser Gly Asn Pro Ala Ile
 165 170 175
 Phe Thr Ser Gly Ile Ile Met Asp Thr Gln Gln Ala Lys Gln Thr Leu
 180 185 190
 Arg Asp Ile Glu Ala Arg His Asn Asp Ile Ile Lys Leu Glu Ser Ser
 195 200 205
 Ile Arg Glu Leu His Asp Met Phe Met Asp Met Ala Met Leu Val Glu
 210 215 220
 Ser Gln Gly Glu Met Ile Asp Arg Ile Glu Tyr Asn Val Glu Gln Ser
 225 230 235 240
 Val Asp Tyr Val Glu Thr Ala Lys Met Asp Thr Lys Lys Ala Val Lys
 245 250 255
 Tyr Gln Ser Lys Ala Arg Arg Lys Lys Phe Tyr Ile Ala Ile Cys Cys
 260 265 270
 Gly Val Ala Leu Gly Ile Leu Val Leu Val Leu Ile Ile Val Leu Ala
 275 280 285

<210> 19
 <211> 1005
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Plasmid pQPIGFP-SNAP25

<221> CDS
 <222> (1)...(1005)

<400> 19
 atg gct agc aaa gga gaa gaa ctc ttc act gga gtt gtc cca att ctt
 Met Ala Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu

48

1	5	10	15	
ggt gaa tta gat ggt gat gtt aac ggc cac aag ttc tct gtc agt gga				96
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly				
20	25	30		
gag ggt gaa ggt gat gca aca tac gga aaa ctt acc ctg aag ttc atc				144
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile				
35	40	45		
tgc act act ggc aaa ctg cct gtt cca tgg cca aca cta gtc act act				192
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr				
50	55	60		
ctg tgc tat ggt gtt caa tgc ttt tca aga tac ccg gat cat atg aaa				240
Leu Cys Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys				
65	70	75	80	
cgg cat gac ttt ttc aag agt gcc atg ccc gaa ggt tat gta cag gaa				288
Arg His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu				
85	90	95		
agg acc atc ttc ttc aaa gat gac ggc aac tac aag aca cgt gct gaa				336
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu				
100	105	110		
gtc aag ttt gaa ggt gat acc ctt gtt aat aga atc gag tta aaa ggt				384
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly				
115	120	125		
att gac ttc aag gaa gat ggc aac att ctg gga cac aaa ttg gaa tac				432
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr				
130	135	140		
aac tat aac tca cac aat gta tac atc atg gca gac aaa caa aag aat				480
Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn				
145	150	155	160	
gga atc aaa gtg aac ttc aag acc cgc cac aac att gaa gat gga agc				528
Gly Ile Lys Val Asn Phe Lys Thr Arg His Asn Ile Glu Asp Gly Ser				
165	170	175		
gtt caa cta gca gac cat tat caa caa aat act cca att ggc gat ggc				576
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly				
180	185	190		
cct gtc ctt tta cca gac aac cat tac ctg tcc aca caa tct gcc ctt				624
Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu				
195	200	205		
tcg aaa gat ccc aac gaa aag aga gac cac atg gtc ctt ctt gag ttt				672
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe				
210	215	220		
gta aca gct gct ggg att aca cat ggc atg gat gaa ctg tac aac ggc				720
Val Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Asn Gly				
225	230	235	240	

```

ggt gca gga tcc ggt gcg ggt ggc ggt ggc atc cgg agg gta aca aac      768
Gly Ala Gly Ser Gly Ala Gly Gly Gly Gly Ile Arg Arg Val Thr Asn
                245                        250                        255

gat gcc cgg gaa aat gag atg gat gag aac ctg gag cag gtg agc ggc      816
Asp Ala Arg Glu Asn Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly
                260                        265                        270

atc atc gga aac ctc cgc cat atg gct cta gac atg ggc aat gag att      864
Ile Ile Gly Asn Leu Arg His Met Ala Leu Asp Met Gly Asn Glu Ile
                275                        280                        285

gac acc cag aat cgc cag atc gac agg atc atg gag aag gct gat tcc      912
Asp Thr Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser
                290                        295                        300

aac aaa acc aga att gat gaa gcc aac caa cgt gca aca aag atg ctg      960
Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu
305                        310                        315                        320

gga agt ggt ggc ggt ggc ggc cat cac cat cac cat cac tgc taa      1005
Gly Ser Gly Gly Gly Gly Gly His His His His His His Cys *
                325                        330

```

<210> 20 ,

<211> .334

<212> PRT

<213> Artificial Sequence

<220>

<223> Plasmid pQBI GFP-SNAP25

<400> 20

```

Met Ala Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 1          5          10          15
Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
 20          25          30
Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
 35          40          45
Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
 50          55          60
Leu Cys Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
 65          70          75          80
Arg His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
 85          90          95
Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100          105          110
Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115          120          125
Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130          135          140
Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145          150          155          160
Gly Ile Lys Val Asn Phe Lys Thr Arg His Asn Ile Glu Asp Gly Ser
165          170          175
Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180          185          190

```

```

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
    195                                200                205
Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
    210                                215                220
Val Thr Ala Ala Gly Ile Thr His Gly Met Asp Glu Leu Tyr Asn Gly
    225                                230                235                240
Gly Ala Gly Ser Gly Ala Gly Gly Gly Gly Ile Arg Arg Val Thr Asn
    245                                250                255
Asp Ala Arg Glu Asn Glu Met Asp Glu Asn Leu Glu Gln Val Ser Gly
    260                                265                270
Ile Ile Gly Asn Leu Arg His Met Ala Leu Asp Met Gly Asn Glu Ile
    275                                280                285
Asp Thr Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser
    290                                295                300
Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu
    305                                310                315                320
Gly Ser Gly Gly Gly Gly Gly His His His His His His Cys
    325                                330

```

<210> 21
 <211> 5
 <212> PRT
 <213> Clostridium sp.

<400> 21
 Gly Gly Gly Gly Ser
 1 5

<210> 22
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> synthetic construct

<400> 22
 Glu Ala Asn Gln Arg Ala Thr Lys
 1 5

<210> 23
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> synthetic construct

<400> 23
 Gly Ala Ser Gln Phe Glu Thr Ser
 1 5

<210> 24
 <211> 8
 <212> PRT
 <213> Artificial Sequence

<220>

- <223> synthetic construct

<400> 24
Asp Thr Lys Lys Ala Val Lys Trp
1 5

<210> 25
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<400> 25
Arg Asp Gln Lys Leu Ser Glu Leu
1 5

<210> 26
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<400> 26
Gln Ile Asp Arg Ile Met Glu Lys
1 5

<210> 27
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<400> 27
Glu Arg Asp Gln Lys Leu Ser Glu
1 5

<210> 28
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<400> 28
Glu Thr Ser Ala Ala Lys Leu Lys
1 5

<210> 29
<211> 8
<212> PRT
<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 29

Gly Ala Ser Gln Phe Glu Thr Ser
1 5

<210> 30

<211> 13

<212> PRT

<213> Homo sapiens

<400> 30

Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met
1 5 10

<210> 31

<211> 15

<212> PRT

<213> Homo sapiens

<400> 31

Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys
1 5 10 15

<210> 32

<211> 16

<212> PRT

<213> Homo sapiens

<400> 32

Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met
1 5 10 15

<210> 33

<211> 17

<212> PRT

<213> Homo sapiens

<400> 33

Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met
1 5 10 15
Leu

<210> 34

<211> 17

<212> PRT

<213> Homo sapiens

<400> 34

Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys
1 5 10 15
Met

<210> 35
 <211> 18
 <212> PRT
 <213> Homo sapiens

<400> 35
 Asp Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys
 1 5 10 15
 Met Leu

<210> 36
 <211> 33
 <212> PRT
 <213> Mus musculus

<400> 36
 Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys
 1 5 10 15
 Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser
 20 25 30
 Gly

<210> 37
 <211> 32
 <212> PRT
 <213> Homo sapiens

<400> 37
 Gln Asn Pro Gln Ile Lys Arg Ile Thr Asp Lys Ala Asp Thr Asn Arg
 1 5 10 15
 Asp Arg Ile Asp Ile Ala Asn Ala Arg Ala Lys Lys Leu Ile Asp Ser
 20 25 30

<210> 38
 <211> 32
 <212> PRT
 <213> Mus musculus

<400> 38
 Gln Asn Gln Gln Ile Gln Lys Ile Thr Glu Lys Ala Asp Thr Asn Lys
 1 5 10 15
 Asn Arg Ile Asp Ile Ala Asn Thr Arg Ala Lys Lys Leu Ile Asp Ser
 20 25 30

<210> 39
 <211> 34
 <212> PRT
 <213> Gallus gallus

<400> 39
 Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Leu Ile Pro Ile Lys
 1 5 10 15
 Pro Gly Leu Met Lys Pro Thr Ser Val Gln Gln Arg Cys Ser Ala Val
 20 25 30
 Val Lys

<210> 40
 <211> 33
 <212> PRT
 <213> *Carassius auratus*

<400> 40
 Gln Asn Arg Gln Ile Asp Arg Ile Met Asp Met Ala Asp Ser Asn Lys
 1 5 10 15
 Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser
 20 25 30
 Gly

<210> 41
 <211> 33
 <212> PRT
 <213> *Carassius auratus*

<400> 41
 Gln Asn Arg Gln Ile Asp Arg Ile Met Glu Lys Ala Asp Ser Asn Lys
 1 5 10 15
 Thr Arg Ile Asp Glu Ala Asn Gln Arg Ala Thr Lys Met Leu Gly Ser
 20 25 30
 Gly

<210> 42
 <211> 30
 <212> PRT
 <213> *Torpedo sp.*

<400> 42
 Gln Asn Ala Gln Val Asp Arg Ile Val Val Lys Gly Asp Met Asn Lys
 1 5 10 15
 Ala Arg Ile Asp Glu Ala Asn Lys His Ala Thr Lys Met Leu
 20 25 30

<210> 43
 <211> 33
 <212> PRT
 <213> *Strongylocentrotus purpuratus*

<400> 43
 Gln Asn Ser Gln Val Gly Arg Ile Thr Ser Lys Ala Glu Ser Asn Glu
 1 5 10 15
 Gly Arg Ile Asn Ser Ala Asp Lys Arg Ala Lys Asn Ile Leu Arg Asn
 20 25 30
 Lys

<210> 44
 <211> 31
 <212> PRT
 <213> *C. elegans*

<400> 44
 Gln Asn Arg Gln Leu Asp Arg Ile His Asp Lys Gln Ser Asn Glu Val

1		5		10		15
Arg	Val	Glu	Ser	Ala	Asn	Lys
				Arg	Ala	Lys
		20		25		30

<210> 45

<211> 31

<212> PRT

<213> Drosophila sp.

<400> 45

Gln	Asn	Arg	Gln	Ile	Asp	Arg	Ile	Asn	Arg	Lys	Gly	Glu	Ser	Asn	Glu
1			5					10						15	
Ala	Arg	Ile	Ala	Val	Ala	Asn	Gln	Arg	Ala	His	Gln	Leu	Leu	Lys	
			20					25					30		

<210> 46

<211> 32

<212> PRT

<213> Hirudinida sp.

<400> 46

Gln	Asn	Arg	Gln	Val	Asp	Arg	Ile	Asn	Asn	Lys	Met	Thr	Ser	Asn	Gln
1			5					10						15	
Leu	Arg	Ile	Ser	Asp	Ala	Asn	Lys	Arg	Ala	Ser	Lys	Leu	Leu	Lys	Glu
			20					25					30		

<210> 47

<400> 47

000

<210> 48

<400> 48

000

<210> 49

<400> 49

000

<210> 50

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 50

Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu	Ala	Asn	Gln	Arg	Ala	Thr	Lys	Ala
1			5					10					15		
Leu															

<210> 51

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<221> MOD_RES

<222> 16

<223> Xaa=Nle

<400> 51

Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu	Ala	Asn	Gln	Arg	Ala	Thr	Lys	Xaa
1				5					10					15	
Leu															

<210> 52

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 52

Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu	Ala	Asn	Gln	Arg	Ala	Thr	Ala	Met
1				5					10					15	
Leu															

<210> 53

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 53

Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu	Ala	Asn	Gln	Arg	Ala	Ser	Lys	Met
1				5					10					15	
Leu															

<210> 54

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<221> MOD_RES

<222> 14

<223> Xaa=Abu

<400> 54

Ser	Asn	Lys	Thr	Arg	Ile	Asp	Glu	Ala	Asn	Gln	Arg	Ala	Xaa	Lys	Met
1				5					10					15	

Leu

<210> 55
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<221> MOD_RES
<222> 13
<223> Xaa=Abu

<400> 55
Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Gln Arg Xaa Thr Lys Met
1 5 10 15
Leu

<210> 56
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<400> 56
Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Ala Arg Ala Thr Lys Met
1 5 10 15
Leu

<210> 57
<211> 17
<212> PRT
<213> Artificial Sequence

<220>
<223> synthetic construct

<221> MOD_RES
<222> 11
<223> Xaa=Abu

<400> 57
Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Xaa Arg Ala Thr Lys Met
1 5 10 15
Leu

<210> 58
<211> 17
<212> PRT
<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 58

Ser Asn Lys Thr Arg Ile Asp Glu Ala Asn Asn Arg Ala Thr Lys Met
 1 5 10 15
 Leu

<210> 59

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 59

Ser Asn Lys Thr Arg Ile Asp Glu Ala Ala Gln Arg Ala Thr Lys Met
 1 5 10 15
 Leu

<210> 60

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<221> MOD_RES

<222> 9

<223> Xaa=Abu

<400> 60

Ser Asn Lys Thr Arg Ile Asp Glu Xaa Asn Gln Arg Ala Thr Lys Met
 1 5 10 15
 Leu

<210> 61

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 61

Ser Asn Lys Thr Arg Ile Asp Gln Ala Asn Gln Arg Ala Thr Lys Met
 1 5 10 15
 Leu

<210> 62

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> synthetic construct

<400> 62

Ser Asn Lys Thr Arg Ile Asn Glu Ala Asn Gln Arg Ala Thr Lys Met
 1 5 10 15
 Leu

<210> 63

<211> 40

<212> PRT

<213> Homo sapiens

<400> 63

Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg
 1 5 10 15
 Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Ser Ser Ala Ala
 20 25 30
 Lys Leu Lys Arg Lys Tyr Trp Trp
 35 40

<210> 64

<211> 40

<212> PRT

<213> Bos taurus

<400> 64

Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg
 1 5 10 15
 Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser Ala Ala
 20 25 30
 Lys Leu Lys Arg Lys Tyr Trp Trp
 35 40

<210> 65

<211> 40

<212> PRT

<213> Rattus sp.

<400> 65

Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg
 1 5 10 15
 Ala Asp Ala Leu Gln Ala Gly Ala Ser Val Phe Glu Ser Ser Ala Ala
 20 25 30
 Lys Leu Lys Arg Lys Tyr Trp Trp
 35 40

<210> 66

<211> 40

<212> PRT

<213> Rattus sp.

<400> 66

Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg
 1 5 10 15

Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser Ala Ala
 20 25 30
 Lys Leu Lys Arg Lys Tyr Trp Trp
 35 40

<210> 67
 <211> 40
 <212> PRT
 <213> Rattus sp.

<400> 67
 Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg
 1 5 10 15
 Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser Ala Ala
 20 25 30
 Lys Leu Lys Arg Lys Tyr Trp Trp
 35 40

<210> 68
 <211> 40
 <212> PRT
 <213> Rattus sp.

<400> 68
 Asp Leu Val Ala Gln Arg Gly Glu Arg Leu Glu Leu Leu Ile Asp Lys
 1 5 10 15
 Thr Glu Asn Leu Val Asp Ser Ser Val Thr Phe Lys Thr Thr Ser Arg
 20 25 30
 Asn Leu Ala Arg Ala Met Cys Met
 35 40

<210> 69
 <211> 32
 <212> PRT
 <213> Gallus gallus

<400> 69
 Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg Ala Asp Ala Leu
 1 5 10 15
 Gln Ala Gly Ala Ser Val Phe Glu Ser Ser Ala Ala Lys Leu Lys Arg
 20 25 30

<210> 70
 <211> 32
 <212> PRT
 <213> Gallus gallus

<400> 70
 Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg Ala Asp Ala Leu
 1 5 10 15
 Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser Ala Ala Lys Leu Lys Arg
 20 25 30

<210> 71
 <211> 40
 <212> PRT
 <213> Torpedo sp.

<400> 71

```

Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp Asp Arg
 1           5           10           15
Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Ser Ser Ala Ala
          20           25           30
Lys Leu Lys Arg Lys Tyr Trp Trp
      35           40

```

<210> 72

<211> 40

<212> PRT

<213> Strongylocentrotus purpuratus

<400> 72

```

Asp Lys Val Leu Asp Arg Asp Gly Ala Leu Ser Val Leu Asp Asp Arg
 1           5           10           15
Ala Asp Ala Leu Gln Gln Gly Ala Ser Gln Phe Glu Thr Asn Ala Gly
          20           25           30
Lys Leu Lys Arg Lys Tyr Trp Trp
      35           40

```

<210> 73

<211> 40

<212> PRT

<213> Aplysia sp.

<400> 73

```

Glu Lys Val Leu Asp Arg Asp Gln Lys Ile Ser Gln Leu Asp Asp Arg
 1           5           10           15
Ala Glu Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Ala Ser Ala Gly
          20           25           30
Lys Leu Lys Arg Lys Tyr Trp Trp
      35           40

```

<210> 74

<211> 40

<212> PRT

<213> Teuthoida sp.

<400> 74

```

Asp Lys Val Leu Glu Arg Asp Ser Lys Ile Ser Glu Leu Asp Asp Arg
 1           5           10           15
Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Ala Ser Ala Gly
          20           25           30
Lys Leu Lys Arg Lys Phe Trp Trp
      35           40

```

<210> 75

<211> 40

<212> PRT

<213> C. elegans

<400> 75

```

Asn Lys Val Met Glu Arg Asp Val Gln Leu Asn Ser Leu Asp His Arg
 1           5           10           15
Ala Glu Val Leu Gln Asn Gly Ala Ser Gln Phe Gln Gln Ser Ser Arg
          20           25           30
Glu Leu Lys Arg Gln Tyr Trp Trp

```

35

40

<210> 76
 <211> 40
 <212> PRT
 <213> *Drosophila melanogaster*

<400> 76
 Glu Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Gly Glu Arg
 1 5 10 15
 Ala Asp Gln Leu Glu Gly Gly Ala Ser Gln Ser Glu Gln Gln Ala Gly
 20 25 30
 Lys Leu Lys Arg Lys Gln Trp Trp
 35 40

<210> 77
 <211> 40
 <212> PRT
 <213> *Drosophila melanogaster*

<400> 77
 Glu Lys Val Leu Glu Arg Asp Ser Lys Leu Ser Glu Leu Asp Asp Arg
 1 5 10 15
 Ala Asp Ala Leu Gln Gln Gly Ala Ser Gln Phe Glu Gln Gln Ala Gly
 20 25 30
 Lys Leu Lys Arg Lys Phe Trp Leu
 35 40

<210> 78
 <211> 39
 <212> PRT
 <213> *Hirudinida sp.*

<400> 78
 Asp Lys Val Leu Glu Lys Asp Gln Lys Leu Ala Glu Leu Asp Arg Ala
 1 5 10 15
 Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Ala Ser Ala Gly Lys
 20 25 30
 Leu Lys Arg Lys Phe Trp Trp
 35

<210> 79
 <211> 18
 <212> PRT
 <213> *Homo sapiens*

<400> 79
 Glu Arg Ala Val Ser Asp Thr Lys Lys Ala Val Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 80
 <211> 18
 <212> PRT
 <213> *Bos taurus*

<400> 80

Glu Arg Ala Val Ser Asp Thr Lys Lys Ala Val Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 81
 <211> 18
 <212> PRT
 <213> Rattus rattus

<400> 81
 Glu His Ala Lys Glu Glu Thr Lys Lys Ala Ile Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 82
 <211> 18
 <212> PRT
 <213> Rattus rattus

<400> 82
 Glu Lys Ala Arg Asp Glu Thr Arg Lys Ala Met Lys Tyr Gln Gly Gly
 1 5 10 15
 Ala Arg

<210> 83
 <211> 18
 <212> PRT
 <213> Rattus rattus

<400> 83
 Glu Arg Gly Gln Glu His Val Lys Ile Ala Leu Glu Asn Gln Lys Lys
 1 5 10 15
 Ala Arg

<210> 84
 <211> 18
 <212> PRT
 <213> Gallus gallus

<400> 84
 Val Pro Glu Val Phe Val Thr Lys Ser Ala Val Met Tyr Gln Cys Lys
 1 5 10 15
 Ser Arg

<210> 85
 <211> 18
 <212> PRT
 <213> Strongylocentrotus purpuratus

<400> 85
 Val Arg Arg Gln Asn Asp Thr Lys Lys Ala Val Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 86
 <211> 18
 <212> PRT
 <213> *Aplysia* sp.

<400> 86
 Glu Thr Ala Lys Met Asp Thr Lys Lys Ala Val Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 87
 <211> 18
 <212> PRT
 <213> *Teuthoida* sp.

<400> 87
 Glu Thr Ala Lys Val Asp Thr Lys Lys Ala Val Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 88
 <211> 18
 <212> PRT
 <213> *Drosophila melanogaster*

<400> 88
 Gln Thr Ala Thr Gln Asp Thr Lys Lys Ala Leu Lys Tyr Gln Ser Lys
 1 5 10 15
 Ala Arg

<210> 89
 <211> 18
 <212> PRT
 <213> *Hirudinida* sp.

<400> 89
 Glu Thr Ala Ala Ala Asp Thr Lys Lys Ala Met Lys Tyr Gln Ser Ala
 1 5 10 15
 Ala Arg

<210> 90
 <211> 116
 <212> PRT
 <213> *Rattus* sp.

<400> 90
 Met Ser Ala Thr Ala Ala Thr Val Pro Pro Ala Ala Pro Ala Gly Glu
 1 5 10 15
 Gly Gly Pro Pro Ala Pro Pro Pro Asn Leu Thr Ser Asn Arg Arg Leu
 20 25 30
 Gln Gln Thr Gln Ala Gln Val Asp Glu Val Val Asp Ile Met Arg Val
 35 40 45

Asn Val Asp Lys Val Leu Glu Arg Asp Gln Lys Leu Ser Glu Leu Asp
 50 55 60
 Asp Arg Ala Asp Ala Leu Gln Ala Gly Ala Ser Gln Phe Glu Thr Ser
 65 70 75 80
 Ala Ala Lys Leu Lys Arg Lys Tyr Trp Trp Lys Asn Leu Lys Met Met
 85 90 95
 Ile Ile Leu Gly Val Ile Cys Ala Ile Ile Leu Ile Ile Ile Ile Val
 100 105 110
 Tyr Phe Ser Thr
 115

<210> 91
 <211> 36
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 91
 gctagatctc gagttaacca cttcccagca tctttg 36

<210> 92
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 92
 atccggaggg taacaaacga tgcc 24

<210> 93
 <211> 60
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 93
 cgaattccgc gggccaccat gggaggagga ctgaacgaca tcttcgaggc tcaaaagatc 60

<210> 94
 <211> 79
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> primer

<400> 94
 tcgtttgtta ccctccggat atgatgatga tgatgatgat gatgggatcc atgccactcg 60
 atcttttgag cctcgaaga 79

<210> 95
 <211> 30

<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 95
cgaagatctg gaggactgaa cgacatcttc 30

<210> 96
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> primer

<400> 96
gatggtgatg gtgatgacag ccgccaccgc cacc 34