



US 20090069310A1

(19) **United States**
(12) **Patent Application Publication**
Flynn et al.

(10) **Pub. No.: US 2009/0069310 A1**
(43) **Pub. Date: Mar. 12, 2009**

(54) **ANTI-INFLAMMATORY MEDICAMENTS**

ation No. 60/437,304, filed on Dec. 31, 2002, provi-
sional application No. 60/437,804, filed on Jan. 3,
2003.

(76) Inventors: **Daniel L. Flynn**, Lawrence, KS
(US); **Peter A. Petillo**, Arlington,
MA (US)

Correspondence Address:
HOVEY WILLIAMS LLP
10801 Mastin Boulevard, Suite 1000
Overland Park, KS 66210 (US)

(21) Appl. No.: **11/450,852**

(22) Filed: **Jun. 9, 2006**

Related U.S. Application Data

(63) Continuation of application No. 10/746,460, filed on
Dec. 24, 2003, now Pat. No. 7,144,911.


(60) Provisional application No. 60/437,487, filed on Dec.
31, 2002, provisional application No. 60/437,403,
filed on Dec. 31, 2002, provisional application No.
60/437,415, filed on Dec. 31, 2002, provisional appli-

Publication Classification

- (51) **Int. Cl.**
A61K 31/4155 (2006.01)
A61K 31/506 (2006.01)
A61K 31/4709 (2006.01)
A61K 31/5377 (2006.01)
A61K 31/541 (2006.01)
A61P 29/00 (2006.01)
A61K 31/4439 (2006.01)
A61K 31/427 (2006.01)
A61K 31/4184 (2006.01)
- (52) **U.S. Cl.** **514/227.8**; 514/256; 514/313;
514/370; 514/395; 514/407; 514/236.5; 514/341

(57) **ABSTRACT**

Novel compounds and methods of using those compounds for
the treatment of inflammatory conditions are provided. In a
preferred embodiment, modulation of the activation state of
p38 kinase protein comprises the step of contacting the kinase
protein with the novel compounds.






Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.




Fig. 4a.




Fig. 5.




Fig. 6.




Fig. 7.

ANTI-INFLAMMATORY MEDICAMENTS

RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. patent application Ser. No. 10/746,460 filed Dec. 24, 2003, provisional applications entitled Process For MODULATING PROTEIN FUNCTION, Ser. No. 60/437,487 filed Dec. 31, 2002, ANTI-CANCER MEDICAMENTS, Ser. No. 60/437,403 filed Dec. 31, 2002, ANTI-INFLAMMATORY MEDICAMENTS, Ser. No. 60/437,415 filed Dec. 31, 2002, ANTI-INFLAMMATORY MEDICAMENTS, Ser. No. 60/437,304 filed Dec. 31, 2002, and MEDICAMENTS FOR THE TREATMENT OF NEURODEGENERATIVE DISORDERS OR DIABETES, Ser. No. 60/463,804 filed Apr. 18, 2003. Each of these applications is incorporated by reference herein.

SEQUENCE LISTING

[0002] The following application contains a sequence listing in computer readable format (CRF). The content of the enclosed CRF is hereby incorporated by reference.

BACKGROUND OF THE INVENTION

[0003] 1. Field of the Invention

[0004] The present invention relates to novel compounds and methods of using those compounds to treat anti-inflammatory diseases.

[0005] 2. Description of the Prior Art

[0006] Basic research has recently provided the life sciences community with an unprecedented volume of information on the human genetic code and the proteins that are produced by it. In 2001, the complete sequence of the human genome was reported (Lander, E. S. et al. Initial sequencing and analysis of the human genome. *Nature* (2001) 409:860; Venter, J. C. et al. The sequence of the human genome. *Science* (2001) 291:1304). Increasingly, the global research community is now classifying the 50,000+ proteins that are encoded by this genetic sequence, and more importantly, it is attempting to identify those proteins that are causative of major, under-treated human diseases.

[0007] Despite the wealth of information that the human genome and its proteins are providing, particularly in the area of conformational control of protein function, the methodology and strategy by which the pharmaceutical industry sets about to develop small molecule therapeutics has not significantly advanced beyond using native protein active sites for binding to small molecule therapeutic agents. These native active sites are normally used by proteins to perform essential cellular functions by binding to and processing natural substrates or transducing signals from natural ligands. Because these native pockets are used broadly by many other proteins within protein families, drugs which interact with them are often plagued by lack of selectivity and, as a consequence, insufficient therapeutic windows to achieve maximum efficacy. Side effects and toxicities are revealed in such small molecules, either during preclinical discovery, clinical trials, or later in the marketplace. Side effects and toxicities continue to be a major reason for the high attrition rate seen within the drug development process. For the kinase protein family of proteins, interactions at these native active sites have been recently reviewed: see J. Dumas, Protein Kinase Inhibitors: Emerging Pharmacophores 1997-2001, *Expert Opinion on Therapeutic Patents* (2001) 11: 405-429; J.


Dumas, Editor, New challenges in Protein Kinase Inhibition, in *Current Topics in Medicinal Chemistry* (2002) 2: issue 9.

[0008] It is known that proteins are flexible, and this flexibility has been reported and utilized with the discovery of the small molecules which bind to alternative, flexible active sites with proteins. For review of this topic, see Teague, *Nature Reviews/Drug Discovery*, Vol. 2, pp. 527-541 (2003). See also, Wu et al., *Structure*, Vol. 11, pp. 399-410 (2003). However these reports focus on small molecules which bind only to proteins at the protein natural active sites. Peng et al., *Bio. Organic and Medicinal Chemistry Ltrs.*, Vol. 13, pp. 3693-3699 (2003), and Schindler, et al., *Science*, Vol. 289, p. 1938 (2000) describe inhibitors of abl kinase. These inhibitors are identified in WO Publication No. 2002/034727. This class of inhibitors binds to the ATP active site while also binding in a mode that induces movement of the kinase catalytic loop. Pargellis et al., *Nature Structural Biology*, Vol. 9, p. 268 (2002) reported inhibitors p38 alpha-kinase also disclosed in WO Publication No. 00/43384 and Regan et al., *J. Medicinal Chemistry*, Vol. 45, pp. 2994-3008 (2002). This class of inhibitors also interacts with the kinase at the ATP active site involving a concomitant movement of the kinase activation loop.

[0009] More recently, it has been disclosed that kinases utilize activation loops and kinase domain regulatory pockets to control their state of catalytic activity. This has been recently reviewed (see, e.g., M. Huse and J. Kuriyan, *Cell* (2002) 109:275).

SUMMARY OF THE INVENTION

[0010] The present invention is broadly concerned with new compounds for use in treating anti-inflammatory conditions and methods of treating such conditions. In more detail, the inventive compounds have the formula



wherein:

[0011] R^1 is selected from the group consisting of aryls (preferably C_6-C_{18} , and more preferably C_6-C_{12}) and heteroaryls;

[0012] each X and Y is individually selected from the group consisting of —O—, —S—, —NR₆—, —NR₆SO₂—, —NR₆CO—, alkynyls (preferably C_1-C_{18} , and more preferably C_1-C_{12}), alkenyls (preferably C_1-C_{18} , and more preferably C_1-C_{12}), alkylenes (preferably C_1-C_{18} , and more preferably C_1-C_{12}), —O(CH₂)_h—, and —NR₆(CH₂)_h—, where each h is individually selected from the group consisting of 1, 2, 3, or 4, and where for each of alkylenes (preferably C_1-C_{18} , and more preferably C_1-C_{12}), —O(CH₂)_h—, and —NR₆(CH₂)_h—, one of the methylene groups present therein may be optionally double-bonded to a side-chain oxo group except that where —O(CH₂)_h— the introduction of the side-chain oxo group does not form an ester moiety;

[0013] A is selected from the group consisting of aromatic (preferably C_6-C_{18} , and more preferably C_6-C_{12}), monocycloheterocyclic, and bicycloheterocyclic rings;

[0014] D is phenyl or a five- or six-membered heterocyclic ring selected from the group consisting of pyrazolyl, pyrrolyl, imidazolyl, oxazolyl, thiazolyl, furyl, pyridyl, and pyrimidyl;

[0015] E is selected from the group consisting of phenyl, pyridinyl, and pyrimidinyl;

[0016] L is selected from the group consisting of $-C(O)-$ and $-S(O)_2-$;

[0017] j is 0 or 1;

[0018] m is 0 or 1;

[0019] n is 0 or 1;


[0020] p is 0 or 1;

[0021] q is 0 or 1;


[0022] t is 0 or 1;

[0023] Q is selected from the group consisting of


-continued




Q-7




Q-8




Q-1




Q-9




Q-2




Q-10




Q-3




Q-11




Q-4




Q-12




Q-5



Q-13




Q-6




Q-14


-continued




Q-15




Q-16




Q-17




Q-18




Q-19



Q-20




Q-21




Q-22


-continued




Q-23




Q-24




Q-25



Q-26




Q-27




Q-28


-continued




Q-29




Q-30




Q-31



Q-32




Q-33



Q-34

-continued



Q-35

[0024] each R_4 group is individually selected from the group consisting of —H, alkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aminoalkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), alkoxyalkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aryls (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), aralkyls (preferably C_6 - C_{18} , and more preferably C_6 - C_2 and preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), heterocyclyls, and heterocyclylalkyls except when the R_4 substituent places a heteroatom on an alpha-carbon directly attached to a ring nitrogen on Q;

[0025] when two R_4 groups are bonded with the same atom, the two R_4 groups optionally form an alicyclic or heterocyclic 4-7 membered ring;

[0026] each R_5 is individually selected from the group consisting of —H, alkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aryls (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), heterocyclyls, alkylaminos (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), arylaminos (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), cycloalkylaminos (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), heterocyclylaminos, hydroxys, alkoxys (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aryloxys (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), alkylthios (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), arylthios (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), cyanos, halogens, perfluoroalkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), alkylcarbonyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), and nitros;

[0027] each R_6 is individually selected from the group consisting of —H, alkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), allyls, and β -trimethylsilylethyl;

[0028] each R_8 is individually selected from the group consisting of alkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aralkyls (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}) preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), heterocyclyls, and heterocyclylalkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12});

[0029] each R_9 group is individually selected from the group consisting of —H, —F, and alkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), wherein when two R groups are geminal alkyl groups, said geminal alkyl groups may be cyclized to form a 3-6 membered ring;




[0030] each Z is individually selected from the group consisting of —O— and —N(R_4)—; and

[0031] each ring of formula (I) optionally includes one or more of R_7 , where R_7 is a noninterfering substituent individually selected from the group consisting of —H,

alkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aryls (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), heterocyclyls, alkylaminos (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), arylaminos (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), cycloalkylaminos (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), heterocyclylaminos, hydroxys, alkoxy (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aryloxy (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), alkylthio (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), arylthio (preferably C_6 - C_{18} , and more preferably C_6 - C_{12}), arylthio, cyano, halogens, nitrilo, nitro, alkylsulfinyl (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), alkylsulfonyl (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), aminosulfonyl, and perfluoroalkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}).

[0032] In one preferred embodiment, the compound has the structure of formula (I) except that:

[0033] when Q is Q-3 or Q-4, then the compound of formula (I) is not











[0034] when Q is Q-7, q is 0, and R_5 and D are phenyl, then A is not phenyl, oxazolyl, pyridyl, pyrimidyl, pyrazolyl, or imidazolyl;

[0035] when Q is Q-7, R_5 is —OH, Y is —O—, —S—, or —CO—, m is 0, n is 0, p is 0, and A is phenyl, pyridyl, or thiazolyl, then D is not thienyl, thiazolyl, or phenyl;




[0036] when Q is Q-7, R_5 is —OH, m is 0, n is 0, p is 0, t is 0, and A is phenyl, pyridyl, or thiazolyl, then D is not thienyl, thiazolyl, or phenyl;

[0037] when Q is Q-7, then the compound of formula (I) is not








[0038] when Q is Q-8, then Y is not $-\text{CH}_2\text{O}-$;
 [0039] when Q is Q-8, the compound of formula (I) is not



[0040] when Q is Q-9, then the compound of formula (I) is not




[0041] when Q is Q-10, t is 0, and E is phenyl, then any R₇ on E is not an o-alkoxy;
 [0042] when Q is Q-10, then the compound of formula (I) is not







[0043] when Q is Q-11, t is 0, and E is phenyl, then any R₇ on E is not an o-alkoxy;

[0044] when Q is Q-11, then the compound of formula (I) is not




[0045] when Q is Q-15, then the compound of formula (I) is not




R₂₀ = substituted phenyl, R₂₁ = H, alkyl

[0046] when Q is Q-16 and Y is —NH—, then






of formula (I) is not biphenyl;

[0047] when Q is Q-16 and Y is —S—, then




of formula (I) is not phenylsulfonylaminophenyl or phenylcarbonylaminophenyl;


[0048] when Q is Q-16 and Y is —SO₂NH—, then the compound of formula (I) is not




R₂₂ = Me, OH



-continued




when Q is Q-16 and Y is ---CONH--- , then




of formula (I) is not imidazophenyl;

[0049] when Q is Q-16 and Y is ---CONH--- , then the compound of formula (I) is not




[0050] when Q is Q-16 and t is 0, then




of formula (I) is not phenylcarbonylphenyl, pyrimidophenyl, phenylpyrimidyl, pyrimidyl, or N-pyrolyl;


[0051] when Q is Q-17, then the compound of formula (I) is not





[0052] when Q is Q-21, then the compound of formula (I) is not







[0053] when Q is Q-22, then the compound of formula (I) is selected from the group consisting of







-continued




[0054] when Q is Q-22 and q is 0, then the compound of formula (I) is selected from the group consisting of



-continued




[0055] but excluding




meta or para-

R34 = Me, Cl
 R35 = $\text{N}(\text{Me})_2$, morpholino
 R36 = H, F




meta or para-


R37 = $\text{N}(\text{Me})_2$, morpholino, OMe, OH, H
 R38 = H, CN, OMe, OH, benzyloxy, phenyl, nitro
 R39 = H, OH
 R40 = H, F
 R41 = H, Cl




meta or para-



-continued





meta or para-





meta or para-

[0056] when Q is Q-23, then the compound of formula (I) is not











R42 = H, Me







[0057] when Q is Q-24, Q-25, Q-26, or Q-31, then the compound of formula (I) is selected from the group consisting of



-continued




-continued



[0061] wherein each Z is individually selected from the group consisting of $-O-$ and $-N(R_4)-$;


[0062] when Q is Q-31, then the compound of formula (I) is not




[0058] wherein each W is individually selected from the group consisting of $-CH-$ and $-N-$;

[0059] each G_1 is individually selected from the group consisting of $-O-$, $-S-$, and $-N(R_4)-$; and

[0060] $*$ denotes the point of attachment to Q-24, Q-25, Q-26, or Q-31 as follows:




[0063] when Q is Q-28 or Q-29 and t is 0, then the compound of formula (I) is not




R_{46} = hydrogen, hydroxyalkyl, alkoxyalkyloxy, hydroxy


[0064] when Q is Q-28 or Q-29 and Y is an ether linkage, then the compound of formula (I) is not



[0065] when Q is Q-28 or Q-29 and Y is —CONH—, then the compound of formula (I) is not




[0066] when Q is Q-32, then




is not biphenyl, benzoxazolylphenyl, pyridylphenyl or bipyridyl;

[0067] when Q is Q-32, Y is —CONH—, q is 0, m is 0, and





of formula (I) is —CONH—, then A is not phenyl;

[0068] when Q is Q-32, q is 0, m is 0, and



is —CONH—, then the compound of formula (I) is not




R₄₇ = alkyl, substituted phenyl, thienyl, phenacetyl naphthyl
 R₄₈ = H, alkyl, Br, substituted phenyl, benzoyl, phenylsulfonyl
 R₄₉ = H, alkyl, phenyl
 R₅₀ = substituted phenyl
 R₅₁ = benzoyl, phenylalkylaminocarbonyl, substituted phenylaminocarbonyl H, Br
 R₅₂ = Cl, Br, SPh, benzoyl, phenylsulfonyl
 R₅₃ = H, phenylsulfonyl, phenyl, benzyl
 R₅₄ = Et, i-Pr
 R₅₅ = substituted phenyl, substituted benzyl
 X₁ = O, N-Ph, N-alkyl, N-carbamoyl
 Z₁ = N(R₅₀), O

[0069] when Q is Q-32, D is thiazolyl, q is 0, t is 0, p is 0, n is 0, and m is 0, then A is not phenyl or 2-pyridone;

[0070] when Q is Q-32, D is oxazolyl or isoxazolyl, q is 0, t is 0, p is 0, n is 0, and m is 0, then A is not phenyl;


[0071] when Q is Q-32, D is pyrimidyl q is 0, t is 0, p is 0, n is 0, and m is 0, then A is not phenyl;

[0072] when Q is Q-32 and Y is an ether linkage, then




of formula (I) is not biphenyl or phenyloxazolyl;

[0073] when Q is Q-32 and Y is —CH=CH—, then



of formula (I) is not phenylaminophenyl;

[0074] when Q is Q-32, then the compound of formula (I) is not




b = 0-1


X₁ = O, S

R₅₆ = H, CF₃, Cl, imidazolyl, amino, morpholino, phenylthio, cycloalkyl, benzyl, phenyl, phenoxy, thienyl, substituted alkyl, pyridylthio, pyrimidyl, benzylamino, N-benimidazolyl, pyridylcarbonylamino, ureido, N-thiourea, substituted alkanoylamino, phenylsulfonyl, substituted benzoyl, phenylalkenoyl, furanoyl, thienoyl, pyridinoyl, R₅₇ = substituted phenyl, substituted biphenyl


-continued




[0075] when Q is Q-35 as shown












[0076] wherein G is selected from the group consisting of —O—, —S—, —NR₄—, and —CH₂—, k is 0 or 1, and u is 1, 2, 3, or 4, then









[0077] is selected from the group consisting of




-continued




-continued



[0078] except that the compound of formula (I) is not




R71 = H, Me
W4 = N, CH




R72 = thiazolyl, isoxazolyl
imidazolyl, furyl

-continued




28.1 R73 = —OCH2CO2H
 R74 = oxazolyl, imidazolyl
 28.2 R73 = CO2Me
 R74 = chlorophenyl




meta, para

R75 = H, Et
 R76 = H, NH2, NO2
 n = 0-1




R77 = H, alkyl
 X3 = O or CH2





meta, para



R78 = H, alkyl




-continued



meta, para




R67 = OH, NH2
 R68 = CF3, Me
 R70 = 2-MeSO2-phen-1-yl,
 2-NH2SO2-phen-1-yl,
 morpholino, imidazolyl, N(Et)2
 W2 = CR69 or N




K65 = H, Et

-continued



[0079] Even more preferably, R₁ as discussed above is selected from the group consisting of 6-5 fused heteroaryls, 6-5 fused heterocyclyls, 5-6 fused heteroaryls, and 5-6 fused heterocyclyls, and even more preferably, R₁ is selected from the group consisting of




[0080] each R₂ is individually selected from the group consisting of —H, alkyls (preferably C₁-C₁₈, and more preferably C₁-C₁₂), aminos, alkylaminos (preferably C₁-C₁₈, and more preferably C₁-C₁₂), arylaminos (preferably C₆-C₁₈ and more preferably C₆-C₁₂), cycloalkylaminos (preferably C₁-C₁₈, and more preferably C₁-C₁₂), heterocyclaminos, halogens, alkoxy (preferably C₁-C₁₈, and more preferably C₁-C₁₂), and hydroxys;

[0081] each R₃ is individually selected from the group consisting of —H, alkyls (preferably C₁-C₁₈, and more preferably C₁-C₁₂), alkylaminos (preferably C₁-C₁₈, and more preferably C₁-C₁₂), arylaminos (preferably C₆-C₁₈, and more preferably C₆-C₁₂), cycloalkylaminos (preferably C₁-C₁₈, and more preferably C₁-C₁₂), heterocyclaminos, alkoxy (preferably C₁-C₁₈, and more preferably C₁-C₁₂), hydroxys, cyanos, halogens, per-

fluoroalkyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), alkylsulfinyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), alkylsulfonyls (preferably C_1 - C_{18} , and more preferably C_1 - C_{12}), R_4NHSO_2 —, and — $NHSO_2R_4$; and

[0082] V is selected from the group consisting of O and H_2 .

[0083] Finally, in another preferred embodiment, A as described above is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidyl, thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, indolyl, indazolyl, benzimidazolyl, benzotriazolyl, isoquinolyl, quinolyl, benzothiazolyl, benzofuranyl, benzothienyl, pyrazolylpyrimidinyl, imidazopyrimidinyl, purinyl, and



[0084] where each W_1 is individually selected from the group consisting of —CH— and —N—.

[0085] With respect to the method of using the novel compounds, the activation state of a kinase is determined by the interaction of switch control ligands and complementary switch control pockets. One conformation of the kinase may result from the switch control ligand's interaction with a particular switch control pocket while another conformation may result from the ligand's interaction with a different switch control pocket. Generally interaction of the ligand with one pocket, such as the "on" pocket, results in the kinase assuming an active conformation wherein the kinase is biologically active. Similarly, an inactive conformation (wherein the kinase is not biologically active) is assumed when the ligand interacts with another of the switch control pockets, such as the "off" pocket. The switch control pocket can be selected from the group consisting of simple, composite and combined switch control pockets. Interaction between the switch control ligand and the switch control pockets is dynamic and therefore, the ligand is not always interacting with a switch control pocket. In some instances, the ligand is not in a switch control pocket (such as occurs when the protein is changing from an active conformation to an inactive conformation). In other instances, such as when the ligand is interacting with the environment surrounding the protein in order to determine with which switch control pocket to interact, the ligand is not in a switch control pocket. Interaction of the ligand with particular switch control pockets is controlled in part by the charge status of the amino acid residues of the switch control ligand. When the ligand is in a neutral charge state, it interacts with one of the switch control pockets and when it is in a charged state, it interacts with the other of the switch control pockets. For example, the switch control ligand may have a plurality of OH groups and be in a neutral charge state. This neutral charge state results in a ligand that is more likely to interact with one of the switch control pockets through hydrogen bonding between the OH groups and selected residues of the pocket, thereby resulting in whichever protein conformation results from that interaction. However, if the OH groups of the switch control ligand become charged through phosphorylation or some other means, the

propensity of the ligand to interact with the other of the switch control pockets will increase and the ligand will interact with this other switch control pocket through complementary covalent binding between the negatively or positively charged residues of the pocket and ligand. This will result in the protein assuming the opposite conformation assumed when the ligand was in a neutral charge state and interacting with the other switch control pocket.

[0086] Of course, the conformation of the protein determines the activation state of the protein and can therefore play a role in protein-related diseases, processes, and conditions. For example, if a metabolic process requires a biologically active protein but the protein's switch control ligand remains in the switch control pocket (i.e. the "off" pocket) that results in a biologically inactive protein, that metabolic process cannot occur at a normal rate. Similarly, if a disease is exacerbated by a biologically active protein and the protein's switch control ligand remains in the switch control pocket (i.e. the "on" pocket) that results in the biologically active protein conformation, the disease condition will be worsened. Accordingly, as demonstrated by the present invention, selective modulation of the switch control pocket and switch control ligand by the selective administration of a molecule will play an important role in the treatment and control of protein-related diseases, processes, and conditions.

[0087] One aspect of the invention provides a method of modulating the activation state of a kinase, preferably p38 α -kinase and including both the consensus wild type sequence and disease polymorphs thereof. The activation state is generally selected from an upregulated or downregulated state. The method generally comprises the step of contacting the kinase with a molecule having the general formula (I). When such contact occurs, the molecule will bind to a particular switch control pocket and the switch control ligand will have a greater propensity to interact with the other of the switch control pockets (i.e., the unoccupied one) and a lesser propensity to interact with the occupied switch control pocket. As a result, the protein will have a greater propensity to assume either an active or inactive conformation (and consequently be upregulated or downregulated), depending upon which of the switch control pockets is occupied by the molecule. Thus, contacting the kinase with a molecule modulates that protein's activation state. The molecule can act as an antagonist or an agonist of either switch control pocket. The contact between the molecule and the kinase preferably occurs at a region of a switch control pocket of the kinase and more preferably in an interlobe oxyanion pocket of the kinase. In some instances, the contact between the molecule and the pocket also results in the alteration of the conformation of other adjacent sites and pockets, such as an ATP active site. Such an alteration can also effect regulation and modulation of the active state of the protein. Preferably, the region of the switch control pocket of the kinase comprises an amino acid residue sequence operable for binding to the Formula I molecule. Such binding can occur between the molecule and a specific region of the switch control pocket with preferred regions including the α -C helix, the α -C helix, the catalytic loop, the activation loop, and the C-terminal residues or C-lobe residues (all residues located downstream (toward the C-end) from the Activation loop), and combinations thereof. When the binding region is the α -C helix, one preferred binding sequence in this helix is the sequence IXXKRXRR-EXXLLXXM, (SEQ ID NO. 2). When the binding region is the catalytic loop, one preferred binding sequence in this loop

is DIIHRD (SEQ ID NO. 3). When the binding region is the activation loop, one preferred binding sequence in this loop is a sequence selected from the group consisting of DFGLARHTDD (SEQ ID NO.4), EMTGYVATR WYR (SEQ ID NO. 5), and combinations thereof. When the binding region is in the C-lobe residues, one preferred binding sequence is WMHY (SEQ ID NO. 6). When a biologically inactive protein conformation is desired, molecules which interact with the switch control pocket that normally results in a biologically active protein conformation (when interacting with the switch control ligand) will be selected. Similarly, when a biologically active protein conformation is desired, molecules which interact with the switch control pocket that normally results in a biologically inactive protein conformation (when interacting with the switch control ligand) will be selected. Thus, the propensity of the protein to assume a desired conformation will be modulated by administration of the molecule. In preferred forms, the molecule will be administered to an individual undergoing treatment for a condition selected from the group consisting of human inflammation, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, asthma, gouty arthritis, sepsis, septic shock, endotoxemic shock, Gram-negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, stroke, reperfusion injury, neural trauma, neural ischemia, psoriasis, restenosis, chronic pulmonary inflammatory disease, bone resorptive diseases, graft-versus-host reaction, Chron's disease, ulcerative colitis, inflammatory bowel disease, pyresis, and combinations thereof. In such forms, it will be desired to select molecules that interact with the switch control pocket that generally leads to a biologically active protein conformation so that the protein will have the propensity to assume the biologically inactive form and thereby alleviate the condition. It is contemplated that the molecules of the present invention will be administerable in any conventional form including oral, parenteral, inhalation, and subcutaneous. It is preferred for the administration to be in the oral form. Preferred molecules include the preferred compounds of formula (I), as discussed above.

[0088] Another aspect of the present invention provides a method of treating an inflammatory condition of an individual comprising the step of administering a molecule having the general formula (I) to the individual. Such conditions are often the result of an overproduction of the biologically active form of a protein, including kinases. The administering step generally includes the step of causing said molecule to contact a kinase involved with the inflammatory process, preferably p38 α -kinase. When the contact is between the molecule and a kinase, the contact preferably occurs in an interlobe oxyanion pocket of the kinase that includes an amino acid residue sequence operable for binding to the Formula I molecule. Preferred binding regions of the interlobe oxyanion pocket include the α -C helix region, the α -C helix region, the catalytic loop, the activation loop, the C-terminal residues, and combinations thereof. When the binding region is the α -C helix, one preferred binding sequence in this helix is the sequence IIXXRXXREXXLLXXM, (SEQ ID NO. 2). When the binding region is the catalytic loop, one preferred binding sequence in this loop is DIIHRD (SEQ ID NO. 3). When the binding region is the activation loop, one preferred binding sequence in this loop is a sequence selected from the group consisting of DFGLARHTDD (SEQ ID NO.4), EMTGYVATR WYR (SEQ ID NO. 5), and combinations thereof. Such a method permits treatment of the condition by virtue of

the modulation of the activation state of a kinase by contacting the kinase with a molecule that associates with the switch control pocket that normally leads to a biologically active form of the kinase when interacting with the switch control ligand. Because the ligand cannot easily interact with the switch control pocket associated with or occupied by the molecule, the ligand tends to interact with the switch control pocket leading to the biologically inactive form of the protein, with the attendant result of a decrease in the amount of biologically active protein. Preferably, the inflammatory condition is selected from the group consisting of human inflammation, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis, asthma, gouty arthritis, sepsis, septic shock, endotoxemic shock, Gram-negative sepsis, toxic shock syndrome, adult respiratory distress syndrome, stroke, reperfusion injury, neural trauma, neural ischemia, psoriasis, restenosis, chronic pulmonary inflammatory disease, bone resorptive diseases, graft-versus-host reaction, Chron's disease, ulcerative colitis, inflammatory bowel disease, pyresis, and combinations thereof. As with the other methods of the invention, the molecules may be administered in any conventional form, with any convention excipients or ingredients. However, it is preferred to administer the molecule in an oral dosage form. Preferred molecules are again selected from the group consisting of the preferred formula (I) compounds discussed above.

BRIEF DESCRIPTION OF THE DRAWINGS

[0089] FIG. 1 is a schematic representation of a naturally occurring mammalian protein in accordance with the invention including "on" and "off" switch control pockets, a transiently modifiable switch control ligand, and an active ATP site;

[0090] FIG. 2 is a schematic representation of the protein of FIG. 1, wherein the switch control ligand is illustrated in a binding relationship with the off switch control pocket, thereby causing the protein to assume a first biologically downregulated conformation;

[0091] FIG. 3 is a view similar to that of FIG. 1, but illustrating the switch control ligand in its charged-modified condition wherein the OH groups of certain amino acid residues have been phosphorylated;

[0092] FIG. 4 is a view similar to that of FIG. 2, but depicting the protein wherein the switch control ligand is in a binding relationship with the on switch control pocket, thereby causing the protein to assume a second biologically-active conformation different than the first conformation of FIG. 2;

[0093] FIG. 4a is an enlarged schematic view illustrating a representative binding between the phosphorylated residues of the switch control ligand, and complementary residues from the on switch control pocket;

[0094] FIG. 5 is a view similar to that of FIG. 1, but illustrating in schematic form possible small molecule compounds in a binding relationship with the on and off switch control pockets;

[0095] FIG. 6 is a schematic view of the protein in a situation where a composite switch control pocket is formed with portions of the switch control ligand and the on switch control pocket, and with a small molecule in binding relationship with the composite pocket; and

[0096] FIG. 7 is a schematic view of the protein in a situation where a combined switch control pocket is formed with portions of the on switch control pocket, the switch control

ligand sequence, and the active ATP site, and with a small molecule in binding relationship with the combined switch control pocket.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

[0097] The present invention provides a way of rationally developing new small molecule modulators which interact with naturally occurring proteins (e.g., mammalian, and especially human proteins) in order to modulate the activity of the proteins. Novel protein-small molecule adducts are also provided. The invention preferably makes use of naturally occurring proteins having a conformational property whereby the proteins change their conformations *in vivo* with a corresponding change in protein activity. For example, a given enzyme protein in one conformation may be biologically upregulated, while in another conformation, the same protein may be biologically downregulated. The invention preferably makes use of one mechanism of conformation change utilized by naturally occurring proteins, through the interaction of what are termed “switch control ligands” and “switch control pockets” within the protein.

[0098] As used herein, “switch control ligand” means a region or domain within a naturally occurring protein and having one or more amino acid residues therein which are transiently modified *in vivo* between individual states by biochemical modification, typically phosphorylation, sulfation, acylation or oxidation. Similarly, “switch control pocket” means a plurality of contiguous or non-contiguous amino acid residues within a naturally occurring protein and comprising residues capable of binding *in vivo* with transiently modified residues of a switch control ligand in one of the individual states thereof in order to induce or restrict the conformation of the protein and thereby modulate the biological activity of the protein, and/or which is capable of binding with a non-naturally occurring switch control modulator molecule to induce or restrict a protein conformation and thereby modulate the biological activity of the protein.

[0099] A protein-modulator adduct in accordance with the invention comprises a naturally occurring protein having a switch control pocket with a non-naturally occurring molecule bound to the protein at the region of said switch control pocket, said molecule serving to at least partially regulate the biological activity of said protein by inducing or restricting the conformation of the protein. Preferably, the protein also has a corresponding switch control ligand, the ligand interacting *in vivo* with the pocket to regulate the conformation and biological activity of the protein such that the protein will assume a first conformation and a first biological activity upon the ligand-pocket interaction, and will assume a second, different conformation and biological activity in the absence of the ligand-pocket interaction.

[0100] The nature of the switch control ligand/switch control pocket interaction may be understood from a consideration of schematic FIGS. 1-4. Specifically, in FIG. 1, a protein 100 is illustrated in schematic form to include an “on” switch control pocket 102, and “off” switch control pocket 104, and a switch control ligand 106. In addition, the schematically depicted protein also includes an ATP active site 108. In the exemplary protein of FIG. 1, the ligand 106 has three amino acid residues with side chain OH groups 110. The off pocket 104 contains corresponding X residues 112 and the on pocket 102 has Z residues 114. In the exemplary instance, the protein 100 will change its conformation depending upon the charge

status of the OH groups 110 on ligand 106, i.e., when the OH groups are unmodified, a neutral charge is presented, but when these groups are phosphorylated a negative charge is presented.

[0101] The functionality of the pockets 102, 104 and ligand 106 can be understood from a consideration of FIGS. 2-4. In FIG. 2, the ligand 106 is shown operatively interacted with the off pocket 104 such that the OH groups 110 interact with the X residues 112 forming a part of the pocket 104. Such interaction is primarily by virtue of hydrogen bonding between the OH groups 110 and the residues 112. As seen, this ligand/pocket interaction causes the protein 100 to assume a conformation different from that seen in FIG. 1 and corresponding to the off or biologically downregulated conformation of the protein.

[0102] FIG. 3 illustrates the situation where the ligand 106 has shifted from the off pocket interaction conformation of FIG. 2 and the OH groups 110 have been phosphorylated, giving a negative charge to the ligand. In this condition, the ligand has a strong propensity to interact with on pocket 102, to thereby change the protein conformation to the on or biologically upregulated state (FIG. 4). FIG. 4a illustrates that the phosphorylated groups on the ligand 106 are attracted to positively charged residues 114 to achieve an ionic-like stabilizing bond. Note that in the on conformation of FIG. 4, the protein conformation is different than the off conformation of FIG. 2, and that the ATP active site is available and the protein is functional as a kinase enzyme.

[0103] FIGS. 1-4 illustrate a simple situation where the protein exhibits discrete pockets 102 and 104 and ligand 106. However, in many cases a more complex switch control pocket pattern is observed. FIG. 6 illustrates a situation where an appropriate pocket for small molecule interaction is formed from amino acid residues taken both from ligand 106 and, for example, from pocket 102. This is termed a “composite switch control pocket” made up of residues from both the ligand 106 and a pocket, and is referred to by the numeral 120. A small molecule 122 is illustrated which interacts with the pocket 120 for protein modulation purposes.

[0104] Another more complex switch pocket is depicted in FIG. 7 wherein the pocket includes residues from on pocket 102, and ATP site 108 to create what is termed a “combined switch control pocket.” Such a combined pocket is referred to as numeral 124 and may also include residues from ligand 106. An appropriate small molecule 126 is illustrated with pocket 124 for protein modulation purposes.

[0105] It will thus be appreciated that while in the simple pocket situation of FIGS. 1-4, the small molecule will interact with the simple pocket 102 or 104, in the more complex situations of FIGS. 6 and 7 the interactive pockets are in the regions of the pockets 120 or 124. Thus, broadly the small molecules interact “at the region” of the respective switch control pocket.

Materials and Methods


General Synthesis of Compounds

[0106] In the synthetic schemes of this section, q is 0 or 1. When q=0, the substituent is replaced by a synthetically non-interfering group R₇.



[0107] Compounds of Formula I wherein Q is taken from Q-1 or Q-2 and Y is alkylene are prepared according to the synthetic route shown in Scheme 1.1. Reaction of isothiocyanate 1 with chlorine, followed by addition of isocyanate 2

affords 3-oxo-thiadiazolium salt 3. Quenching of the reaction with air affords compounds of Formula 1-4. Alternatively, reaction of isothiocyanate 1 with isothiocyanate 5 under the reaction conditions gives rise to compounds of Formula I-7. See A. Martinez et al, *Journal of Medicinal Chemistry* (2002) 45: 1292.


[0108] Intermediates 1, 2 and 5 are commercially available or prepared according to Scheme 1.2. Reaction of amine 8 with phosgene or a phosgene equivalent affords isocyanate 2. Similarly, reaction of amine 8 with thiophosgene affords isothiocyanate 5. Amine 8 is prepared by palladium(0)-catalyzed amination of 2, wherein M is a group capable of oxidative insertion into palladium(0), according to methodology reported by S. Buchwald. See M. Wolter et al, *Organic Letters* (2002) 4:973; B. H. Yang and S. Buchwald, *Journal of Organometallic Chemistry* (1999) 576(I-2):125. In this reaction sequence, P is a suitable amine protecting group. Use of and removal of amine protecting groups is accomplished by methodology reported in the literature (Protective Groups in Organic Synthesis, Peter G. M. Wutts, Theodora Greene (Editors) 3rd edition (April 1999) Wiley, John & Sons, Incorporated; ISBN: 0471160199). Starting compounds 9 are commercially available or readily prepared by one of ordinary skill in the art: See March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, Michael B. Smith & Jerry March (Editors) 5th edition (January 2001) Wiley John & Sons; ISBN: 0471585890.




-continued





[0109] Compounds of Formula I wherein Q is taken from Q1 or Q-2 and Y is alkylene are also available via the synthetic route shown in Scheme 1.3. Reaction of amine 8 with isocyanate or isothiocyanate 2a yields the urea/thiourea 8a which can be cyclized by the addition of chlorocarbonyl sulfonyl chloride. See GB 1115350 and U.S. Pat. No. 3,818,024, Revankar et. al U.S. Pat. No. 4,093,624, and Klayman et. al *JOC* 1972, 37(10), 1532 for further details. Where R₄ is a readily removable protecting group (e.g. R=3,4-d-methoxybenzyl amine), the action of mild, acidic deprotection conditions such as CAN or TFA will reveal the parent ring system of I-4 (X=O) and I-7 (X=S).








[0110] I-7 is also available as shown in Scheme 1.4. Condensation of isocyanate or isothiocyanate 2a with amine R_5NH_2 yields urea/thiourea 2b, which, when reacted with chlorocarbonyl sulfonyl chloride according to GB1115350 and U.S. Pat. No. 3,818,024 yields 2c. Where R_4 is a readily removable protecting group (e.g. R=3,4-d-methoxybenzyl amine), the action of mild, acidic deprotection conditions such as CAN or TFA will reveal the parent ring system of 2d. Reaction of 2d with NaH in DMF, and displacement wherein M is a suitable leaving group such as chloride, bromide or iodide yields I-4' (X=O) and I-7' (X=S).




[0111] Compounds of Formula I wherein Q is taken from Q-1' or Q-2' and Y is alkylene are available via the synthetic route shown in Scheme 1.3. Condensation of isocyanate or isothiocyanate 2a with ammonia yields urea/thiourea 2e, which, when reacted with chlorocarbonyl sulfonyl chloride according to GB1115350 and U.S. Pat. No. 3,818,024 yields 2f. Reaction of 2f with NaH in DMF, and displacement wherein M is a suitable leaving group such as chloride, bromide or iodide yields I-4' (X=O) and I-7' (X=S).




[0112] Compounds of Formula I wherein Q is taken from Q-3 or Q-4 and Y is alkylene, are prepared according to the synthetic route shown in Schemes 2.1 and 2.2, respectively. Reaction of 12, wherein M is a suitable leaving group, with the carbamate-protected hydrazine 13 affords intermediate 14. Reaction of 14 with an isocyanate gives rise to intermediate 15. Thermal cyclization of 15 affords 1,2,4-triazolidinedione of Formula I-16. By analogy, scheme 2.2 illustrates the preparation of 3-thio-5-oxo-1,2,4-triazolidines of Formula I-18 by reaction of intermediate 14 with an isothiocyanate and subsequent thermal cyclization.




[0113] Intermediates 12 wherein p is 1 are readily available or are prepared by reaction of 19 with carbamates 10 under palladium(0)-catalyzed conditions. M₁ is a group which oxidatively inserts palladium(0), preferably iodo or bromo, and is of greater reactivity than M. Compounds 19 are either commercially available or prepared by one of ordinary skill in the art.



Compounds of Formula I wherein D is taken from Q-3 or Q-4 and Y is alkylene, are also prepared according to the synthetic route shown in Scheme 2.4. Oxidation of amine R₄NH₂ to the corresponding hydrazine, condensation with ethyl chloroformate subsequent heating yields 1,2,4-triazolidinedione 15a. After the action of NaH in DMF, displacement wherein M is a suitable leaving group such as chloride, bromide or iodide yields I-16 (X=O) and I-18 (X=S).







[0114] Compounds of Formula I wherein D is taken from D-3' or D-4' and Y is alkylene, are also prepared according to the synthetic route shown in Scheme 2.4. When R₅ is a readily removable protecting group (e.g. R=3,4-d-methoxybenzyl amine), the action of mild, acidic deprotection conditions such as CAN or TFA on 15a will reveal 1,2,4-triazolidinone 15b. After deprotonation of 15b by NaH in DMF, displacement wherein M is a suitable leaving group such as chloride, bromide or iodide yields I-16' (X=O) and I-18' (X=S).

[0115] Compounds of Formula I wherein Q is taken from Q-5 or Q-6 and Y is alkylene are prepared according to the


synthetic route shown in Scheme 3. Reaction of hydrazine 20 with chlorosulfonylisocyanate and base, such as triethylamine, gives rise to a mixture of intermediates 21A and 21B which are not isolated but undergo cyclization in situ to afford compounds of Formulae I-22A and I-22B. Compounds I-22A and I-22B are separated by chromatography or fractional crystallization. Optionally, compounds I-22A and I-22B can undergo Mitsunobu reaction with alcohols R₄OH to give compounds of Formulae I-23A and I-23B. Compounds 20 are prepared by acid-catalyzed deprotection of t-butyl carbamates of structure 14, wherein R₁₀ is t-butyl.




[0116] Compounds of Formula I wherein Q is Q-7 and Y is alkylene are prepared as shown in Scheme 4. Reaction of amine 8 with maleimide 24, wherein M is a suitable leaving group, affords compounds of Formula I-25. Reaction of compound 26, wherein M is a group which can oxidatively insert Pd(0), can participate in a Heck reaction with maleimide 27, affording compounds of Formula I-28. Maleimides 24 and 27 are commercially available or prepared by one of ordinary skill in the art.




[0117] Compounds of Formula I wherein Q is Q-8 and Y is alkylene are prepared as shown in Scheme 5, according to methods reported by M. Tremblay et al, *Journal of Combinatorial Chemistry* (2002) 4:429. Reaction of polymer-bound activated ester 29 (polymer linkage is oxime activated-ester) with chlorosulfonyl isocyanate first with benzyl alcohol then 31a yields 31b, which after heating yields I-32. Optionally, intermediate 30 is treated with acid, preferably trifluoroacetic acid, to afford the N-unsubstituted sulfahydantoin I-33.





[0118] Compounds of Formula I wherein Q is Q-8 and Y is alkylene are also prepared as shown in Scheme 5a. Amine 8 is condensed with the glyoxal hemiacetal to yield 31a. Reaction of chlorosulphonyl isocyanate first with benzyl alcohol then 31a yields 31b, which after heating yields I-32.







[0119] Compounds of Formula I wherein Q is taken from Q-8', are prepared according to the synthetic route shown in Scheme 5.2. Formation of 31c by the method of Muller and DuBois *JOC* 1989, 54, 4471 and its deprotonation with NaH/DMF or NaH/DMF and subsequently alkylation wherein M is a suitable leaving group such as chloride, bromide or iodide yields I-32'. Alternatively, I-32' is also available as shown in Scheme 5.3. Mitsunobu reaction of boc-sulfamide amino ethyl ester with alcohol 8b (made by methods analogous to that for amine 8) yields 31c, which after Boc removal with 2N HCl in dioxane is cyclized by the action of NaH on 31d results in I-32'.




[0120] Compounds of Formula I wherein Q is Q-9 and Y is alkylene are prepared as shown in Scheme 6. Reaction of polymer-bound amino acid ester 34 with an isocyanate affords intermediate urea 35. Treatment of 35 with base, preferably pyridine or triethylamine, with optional heating, gives rise to compounds of Formula I-36.




[0121] Compounds of Formula I wherein Q is Q-9 and Y is alkylene are also prepared as shown in Scheme 6.1. Reaction of aldehyde 8c under reductive amination conditions with the t-butyl ester of glycine yields 35a. Isocyanate 2a is condensed with p-nitrophenol (or the corresponding R₄NH₂ amine is condensed with p-nitrophenyl chloroformate) to yield the carbamic acid p-nitrophenyl ester, which when reacted with deprotonated 35a and yields the urea that when deprotected with acid yields 35b. Formula I-36 is directly available from 35b by the action of NaH and heat.




[0122] Compounds of Formula I wherein Q is taken from Q-9', are prepared according to the synthetic route shown in Scheme 6.2. Formation of 35c by the method described in JP10007804A2 and Zvilichovsky and Zucker, Israel Journal of Chemistry, 1969, 7(4), 547-54 and its deprotonation with NaH/DMF or NaH/DMF and its subsequent displacement of M, wherein M is a suitable leaving group such as chloride, bromide or iodide, yields I-36'.




[0123] Compounds of Formula I wherein Q is Q-0 or Q-11, and Y is alkylene are prepared as shown in Schemes 7.1 and 7.2, respectively. Treatment of alcohol 37 (Z=O) or amine 37 (Z=NH) with chlorosulfonylisocyanate affords intermediate carbamate or urea of structure 38. Treatment of 38 with an amine of structure HN(R₄)₂ and base, preferably triethylamine or pyridine, gives sulfonylureas of Formula I-39. Reaction of chlorosulfonylisocyanate with an alcohol (Z=O) or amine (Z=NR₄) 40 affords intermediate 41. Treatment of 41 with an amine 8 and base, preferably triethylamine or pyridine, gives sulfonylureas of Formula I-42.



[0124] Compounds of Formula I wherein Q is taken from Q-12 are prepared according to the synthetic route shown in Scheme 8. Alkylation of pyridine 43, wherein TIPS is triisopropylsilyl, under standard conditions (K_2CO_3 , DMF, R_4-I or Mitsunobu conditions employing R_4-OH) yields pyridine derivative 44 which is reacted with compound 12, wherein M is a suitable leaving group, to afford pyridones of formula I-45.




-continued



[0125] Compounds of Formula I wherein Q is taken from Q-13 are prepared according to the synthetic route shown in Scheme 9. Starting from readily available pyridine 46, alkylation under standard conditions (K_2CO_3 , DMF, R_4-I or Mitsunobu conditions employing R_4-OH) yields pyridine derivative 47. N-alkylation with K_2CO_3 , DMF, R_4-I affords pyridones of formula 48. Intermediate 48 is partitioned to undergo a Heck reaction, giving I-49; a Buchwald amination reaction, giving I-51; or a Buchwald Cu(I) catalyzed O-arylation reaction, to give I-52. The Heck reaction product I-49 may be optionally hydrogenated to afford the saturated compound I-50. Wherein the phenyl ether R_4 group is methyl, compounds of formula I-49, I-50, I-51, or I-52 are treated with boron tribromide or lithium chloride to afford compounds of Formula I-53, wherein R_4 is hydrogen.


Scheme 9



[0126] Compounds of Formula I wherein Q is taken from Q-14 are prepared according to the synthetic route shown in Scheme 10. Starting from readily available pyridine 54, alkylation under standard conditions (K_2CO_3 , DMF, R_4-I or Mitsunobu conditions employing R_4-OH) yields pyridine derivative 55. N-alkylation with K_2CO_3 , DMF, R_4-I affords pyridones of formula 56. Intermediate 56, wherein M is a suitable leaving group, preferably bromine or chlorine, is



partitioned to undergo a Heck reaction, giving I-57; a Buchwald amination reaction, giving I-59; or a Buchwald Cu(I) catalyzed O-arylation reaction, to give I-60. The Heck reaction product I-57 may be optionally hydrogenated to afford the saturated compound I-58. Wherein R_4 is methyl, compounds of formula I-57, I-58, I-59, or I-60 are treated with boron tribromide or lithium chloride to afford compounds of Formula I-61, wherein R_4 is hydrogen.

Scheme 10




[0127] Compounds of Formula I wherein Q is taken from Q-15 are prepared according to the synthetic routes shown in Schemes 11 and 12. Starting esters 62 are available from the corresponding secoacids via TBS-ether and ester formation under standard conditions. Reaction of protected secoester 62 with Meerwin's salt produces the vinyl ether 63 as a pair of regioisomers. Alternatively, reaction of 62 with dimethylamine affords the vinylogous carbamate 64. Formation of the dihydropyrimidinedione 66 proceeds by condensation with urea 65 with azeotropic removal of dimethylamine or methanol. Dihydropyrimidinedione 66 may optionally be further substituted by Mitsunobu reaction with alcohols R_4OH to give rise to compounds 67.

[0128] Scheme 12 illustrates the further synthetic elaboration of intermediates 67. Removal of the silyl protecting group (TBS) is accomplished by treatment of 67 with fluoride (tetra-n-butylammonium fluoride or cesium fluoride) to give primary alcohols 68. Reaction of 68 with isocyanates 2 gives rise to compounds of Formula I-69. Alternatively, reaction of 68 with $[R_6O_2 C(NH)p]q-D-E-M$, wherein M is a suitable leaving group, affords compounds of Formula I-70. Oxidation of 68 using the Dess-Martin periodinane (D. Dess, J. Martin, *J. Am. Chem. Soc.* (1991) 113:7277) or tetra-n-alkyl peruthenate (W. Griffith, S. Ley, *Aldrichimica Acta* (1990) 23:13) gives the aldehydes 71. Reductive amination of 71 with amines 8 gives rise to compounds of Formula I-72. Alternatively, aldehydes 71 may be reacted with ammonium acetate under reductive alkylation conditions to give rise to the primary amine 73. Reaction of 73 with isocyanates 2 affords compounds of Formula I-74.







Scheme 12




[0129] Compounds of Formula I wherein Q is taken from Q-16 are prepared according to the synthetic routes shown in Schemes 13 and 14. Starting esters 75 are available from the corresponding secoacids via TBS-ether and ester formation under standard conditions. Reaction of protected secoester 75 with Meerwin's salt produces the vinyl ether 76 as a pair of regioisomers. Alternatively, reaction of 75 with dimethylamine affords the vinylogous carbamate 77. Formation of the dihydropyrimidinedione 78 proceeds by condensation with urea 65 with azeotropic removal of dimethylamine or methanol. Dihydropyrimidinedione 78 may optionally be further substituted by Mitsunobu reaction with alcohols R_4OH to give rise to compounds 79. Compounds of Formulae I-81, I-82, I-84, and I-86 are prepared as shown in Scheme 14 by analogy to the sequence previously described in Scheme 12.

-continued







[0130] Alkyl acetoacetates 87 are commercially available and are directly converted into the esters 88 as shown in Scheme 15. Treatment of 87 with NaHMDS in THF, followed by quench with formaldehyde and TBSCl (n=1) or Q-(CH₂)_n-OTBS (n=2-4), gives rise to compounds 88.






[0131] Compounds of Formula I wherein Q is taken from Q-17 are prepared according to the synthetic routes shown in Schemes 16.1 and 16.2, and starts with the BOC-protected hydrazine 13, which is converted to the 1,2-disubstituted hydrazine 89 by a reductive alkylation with a glyoxal derivative mediated by sodium cyanoborohydride and acidic workup. Condensation of 89 with diethyl malonate in benzene under reflux yields the heterocycle 90. Oxidation with N₂O₄ in benzene (see Cardillo, Merlini and Boeri *Gazz. Chim. Ital.*, (1966) 9:8) to the nitromalonohydrazide 91 and further treatment with P₂O₅ in benzene (see: Cardillo, G. et al, *Gazz. Chim. Ital.* (1966) 9:973-985) yields the tricarbonyl 92. Alternatively, treatment of 90 with Brederick's reagent (t-BuOCH(N(Me)₂)₂), gives rise to 93, which is subjected to ozonolysis, with a DMS and methanol workup, to afford the protected tricarbonyl 92. Compound 92 is readily deprotected by the action of CsF in THF to yield the primary alcohol 94. Alcohol 94 is optionally converted into the primary amine 95 by a sequence involving tosylate formation, azide displacement, and hydrogenation.






[0132] Reaction of 94 with (hetero)aryl halide 26, wherein M is iodo, bromo, or chloro, under copper(I) catalysis affords compounds I-96. Optional deprotection of the di-methyl ketal with aqueous acid gives rise to compounds of Formula I-98. By analogy, reaction of amine 95 with 26 under palladium(0) catalysis affords compounds of Formula I-97. Optional deprotection of the di-methyl ketal with aqueous acid gives rise to compounds of Formula I-99.







[0133] Compounds of Formula I wherein Q is taken from Q-17 are also prepared according to the synthetic route shown in Scheme 16.3. Deprotonation of 4,4-dimethyl-3,5-dioxopyrazolidine (95a, prepared according to the method described in Zinner and Boese, *D. Pharmazie* 1970, 25(5-6), 309-12 and Bausch, M. J. et. al *J. Org. Chem.* 1991, 56(19), 5643) with NaH/DMF or NaH/DMF and its subsequent displacement of M, wherein M is a suitable leaving group such as chloride, bromide or iodide yields I-99a.





[0134] Compounds of Formula I wherein Q is taken from Q-18 are prepared as shown in Schemes 17.1 and 17.2. Amine esters 100 are subjected to reductive alkylation conditions to give rise to intermediates 101. Condensation of amines 101 with carboxylic acids using an acid activating reagent such as dicyclohexylcarbodiimide (DCC)/hydroxybenzotriazole (HOBt) affords intermediate amides 102. Cyclization of amides 102 to tetramic acids 104 is mediated by Amberlyst A-26 hydroxide resin after trapping of the in situ generated alkoxide 103 and submitting 103 to an acetic acid-mediated resin-release.




M' is t-BuOCH₂—, BOCNH(CH₂)₃—, BOCNH(CH₂)₄—, HC≡C—CH₂—
 Y is HOCH₂—; H₂N—(CH₂)₃—;
 H₂N—(CH₂)₄—; HC≡C—CH₂—

[0135] Scheme 17.2 illustrates the synthetic sequences for converting intermediates 104 to compounds of Formula I. Reaction of alcohol 104.1 with aryl or heteroaryl halide (Q=halogen) under copper(I) catalysis gives rise to compounds of Formula I-105.1. Reaction of amines 104.2 and 104.3 with 26 under Buchwald palladium(0) catalyzed amination conditions affords compounds of Formulae I-105.2 and I-105.3. Reaction of acetylene 104.4 with 26 under Sonogashira coupling conditions affords compounds of Formula I-105.4. Compounds I-105.4 may optionally be reduced to the corresponding saturated analogs I-105.5 by standard hydrogenation.







anhydride 107 using a dehydrating reagent, preferably diisopropylcarbodiimide (DIC) or 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDC). Reaction of 107 with amines R₄NH₂ affords the intermediate amides which are cyclized to the imides 108 by reaction with DIC or EDC. Alternatively, 107 is reacted with amines 8 to afford amides of Formula I-110. Amides I-110 may optionally be further reacted with DIC or EDC to give rise to compounds of Formula I-111. Acid 108 is further reacted with amines 8 to give compounds of Formula I-109.




[0136] Compounds of Formula I wherein Q is taken from Q-1 g, Q-20, or Q-21 are prepared as illustrated in Scheme 18. Commercially available Kemp's acid 106 is converted to its


[0137] Compounds of Formula I wherein Q is taken from Q-22 or Q-23 are prepared as shown in Schemes 19.1 through 19.3. Preparation of intermediates 113 and 114 are prepared as shown in Scheme 19.1 from di-halo(hetero)aryls 112, wherein M_2 is a more robust leaving group than M_1 . Reaction of 112 with amines 37 ($Z=NH$) either thermally in the presence of base or by palladium(0) catalysis in the presence of base and phosphine ligand affords compounds 113. Alternatively, reaction of 112 with alcohols 37 ($X=O$) either thermally in the presence of base or by copper(I) catalysis in the presence of base affords compounds 114.



[0138] Scheme 19.2 illustrates the conversion of intermediates 113 into compounds of Formula I-115, I-118, or 117. Treatment of 113 with aqueous copper oxide or an alkaline hydroxide affords compounds of Formula I-115. Alternatively, treatment of 113 with t-butylmercaptan under copper (I) catalysis in the presence of ethylene glycol and potassium carbonate gives rise to 116 (see F. Y. Kwong and S. L. Buchwald, *Organic Letters* (2002) 4:3517. Treatment of the t-butyl sulfide 116 with acid affords the desired thiols of Formula I-118. Alternatively, 113 may be treated with excess ammonia under pressurized conditions to afford compound 117.




[0139] Scheme 19.3 illustrates the conversion of intermediate 114 into compounds of Formula I-119, I-122, and 121, by analogy to the sequence described in Scheme 19.2.





[0140] Compounds of Formula I wherein q is taken from Q-24, Q-25, or Q-26 are prepared as shown in Scheme 20. Reaction of compounds I-115 or I-119 with chlorosulfonylisocyanate, followed by in situ reaction with amines $\text{HN}(\text{R}_4)_2$ gives rise to compounds of Formulae I-123 or I-124. Reaction of compounds I-118 or I-122 with a peracid, preferably per-

acetic acid or trifluoroperacetic acid, affords compounds of Formula I-125 or I-126. Reaction of compounds 117 or 121 with chlorosulfonylisocyanate, followed by in situ reaction with amines $\text{HN}(\text{R}_4)_2$ or alcohols R_4OH , affords compounds of Formulae I-127, I-128, I-129, or I-130.


Scheme 20




[0141] Compounds of Formula I wherein Q is taken from Q-27 are prepared as illustrated in Scheme 21. Reductive alkylation of thiomorpholine with aldehydes 131 affords benzylic amines 132, which are then subjected to peracid oxidation to give rise to the thiomorpholine sulfones 133 (see C. R. Johnson et al, *Tetrahedron* (1969) 25: 5649). Intermediates 133 are reacted with amines 8 (Z=NH₂) under Buchwald palladium-catalyzed amination conditions to give rise to compounds of Formula I-134. Alternatively, compounds 133 are reacted with alcohols 8 (Z=OH) under Buchwald copper (I) catalyzed conditions to afford compounds of Formula I-135. Alternatively, intermediates 133 are reacted with alkenes under palladium(0)-catalyzed Heck reaction conditions to give compounds of Formula I-136. Compounds I-136 are optionally reduced to the corresponding saturated analogs I-137 by standard hydrogenation conditions or by the action of diimide.



[0142] Compounds of Formula I wherein Q is taken from Q-27 are also prepared as illustrated in Scheme 21.1. Aldehyde 8c is reductively aminated with ammonia, and the resultant amine condensed with divinyl sulphone to yield I-134. Intermediate 134a is also available by reduction of amide 8d under a variety of standard conditions.




[0143] More generally, amines 134c are available via the reduction of amides 134b as shown in Scheme 21.2. The morpholine amide analogues 134d and morpholine analogues 134e are also available as shown in Scheme 21.2.




[0144] Compounds of Formula I wherein Q is taken from Q-28 or Q-29 are prepared according to the sequences illustrated in Scheme 22. Readily available amides 138 are reacted with chlorosulfonylisocyanate to give intermediates 140,


which are reacted in situ with amines $HN(R_4)_2$ or alcohols R_4OH to afford compounds of Formulae I-141 or I-142, respectively. Alternatively, amides 138 are reacted with sulfonylchlorides to give compounds of Formula I-139.




-continued



[0145] Compounds of Formula I wherein Q is taken from Q-30 are prepared as shown in Scheme 23. Readily available N—BOC anhydride 143 (see S. Chen et al, *J. Am. Chem. Soc.* (1996) 118:2567) is reacted with amines $\text{HN}(\text{R}_4)_2$ or alcohols R_6OH to afford acids 144 or 145, respectively. Intermediates 144 or 145 are further reacted with amines $\text{HN}(\text{R}_4)_2$ in the presence of an acid-activating reagent, preferably PyBOP and di-isopropylethylamine, to give diamides 146 or ester-amides 147. Intermediate 145 is converted to the diesters 148 by reaction with an alkyl iodide in the presence of base, preferably potassium carbonate. Intermediates 146-148 are treated with HCl/dioxane to give the secondary amines 149-151, which are then condensed with acids 152 in the presence of PyBOP and di-isopropylethylamine to give compounds of Formula I-153.




-continued




[0146] Compounds of Formula I wherein Q is taken from Q-31 or Q-32 are prepared according to the sequences illustrated in Scheme 24. Treatment of readily available sulfenamides 154 with amines 37 ($Z=NH$), alcohols 37 ($Z=O$), or alkenes 37 ($Z=CH=CH_2$), gives rise to compounds of Formula I-155. Treatment of sulfenamides I-155 with iodosobenzene in the presence of alcohols R_6OH gives rise to the sulfonimidates of Formula I-157 (see D. Leca et al,

Organic Letters (2002) 4:4093). Alternatively, compounds I-155 ($Z=CH=CH$) may be optionally reduced to the saturated analogs I-156 ($Z=CH_2-CH_2-$), which are converted to the corresponding sulfonimidates I-157.


[0147] Treatment of readily available sulfonylchlorides 154.1 with amines $HN(R_4)_2$ and base gives rise to compounds of Formula I-154.2.






[0148] Compounds of Formula I wherein Q is taken from Q-33 are prepared as shown in Scheme 25. Readily available nitrites 158 are reacted with amines 37 (Z=NH), alcohols 37 (Z=O), or alkenes 37 (Z=—CH=CH₂) to afford compounds of Formula I-159. Compounds I-159 (wherein

Z=CH=CH—) are optionally reduced to their saturated analogs I-160 by standard catalytic hydrogenation conditions. Treatment of compounds I-159 or I-160 with a metal azide (preferably sodium azide or zinc azide) gives rise to tetrazoles of Formula I-161.







[0149] Compounds of Formula I wherein Q is taken from Q-34 are prepared as shown in Scheme 26. Readily available esters 162 are reacted with amines 37 ($Z=\text{NH}$), alcohols 37 ($Z=\text{O}$), or alkenes 37 ($Z=\text{—CH=CH}_2$) to afford compounds of Formula I-163. Compounds I-163 (wherein Z is —CH=CH—) are optionally converted to the saturated analogs I-164 by standard hydrogenation conditions. Com-

pounds I-163 or I-164 are converted to the desired phosphonates I-165 by an Arbuzov reaction sequence involving reduction of the esters to benzylic alcohols, conversion of the alcohols to the benzylic bromides, and treatment of the bromides with a tri-alkylphosphite. Optionally, phosphonates I-165 are converted to the fluorinated analogs I-166 by treatment with diethylaminosulfur trifluoride (DAST).


Scheme 26




[0150] Compounds of Formula I wherein Q is taken from Q-35 are prepared according to Scheme 27. Readily available acid chlorides 167 are reacted with oxazolidinones in the presence of base to afford the N-acyl oxazolidinones 168. Intermediate 168 are reacted with amines 37 (Z=NH), alcohols 37 (Z=O), or alkenes 37 (Z=—CH=CH₂) to afford the N-acyl oxazolidinones of Formula I-169. Compounds I-169 (wherein Z is —CH=CH—) are optionally converted to the saturated analogs I-170 under standard hydrogenation conditions.





[0151] Compounds of Formula I wherein Q is taken from Q-35 are also prepared as illustrated in Scheme 27.1. Intermediate 8a, wherein M is a suitable leaving group such as chloride, bromide or iodide, is refluxed with triethyl phosphite and the resulting phosphoryl intermediate saponified under mild conditions to yield I-165.




[0152] Compounds of Formula I wherein Q is taken from Q-36 are prepared as illustrated in Schemes 28.1 and 28.2. Reductive alkylation of the t-butylsulfide substituted piperazines with the readily available aldehydes 131 gives rise to the benzylic piperazines 171. Intermediates 171 are reacted with amines 37 (Z=NH), alcohols 37 (Z=O), or alkenes 37 (Z=—CH=CH₂) to give compounds 172, 173, or 174, respectively. Optionally, intermediates 174 are converted to the saturated analogs 175 under standard hydrogenation conditions.






[0153] Scheme 28.2 illustrates the conversion of intermediate t-butylsulfides 172-175 to the sulfonic acids, employing a two step process involving acid-catalyzed deprotection of the t-butyl sulfide to the corresponding mercaptans, and subsequent peracid oxidation (preferably with peracetic acid or trifluoroperacetic acid) of the mercaptans to the desired sulfonic acids of Formula I-176.




Z = NH, O, CH=CH, CH₂-CH₂


[0154] In some instances a hybrid p38-alpha kinase inhibitor is prepared which also contains an ATP-pocket binding moiety or an allosteric pocket binding moiety R_1-X-A . The synthesis of functionalized intermediates of formula R_1-X-A are accomplished as shown in Scheme 29. Readily available intermediates 177, which contain a group M capable of oxidative addition to palladium(0), are reacted with amines 178 ($X=NH$) under Buchwald Pd(0) amination conditions to afford 179. Alternatively amines or alcohols 178 ($X=NH$ or O) are reacted thermally with 177 in the presence of base under nuclear aromatic substitution reaction conditions to afford 179. Alternatively, alcohols 178 ($X=O$) are reacted with 177 under Buchwald copper(I)-catalyzed conditions to afford 179. In cases where $p=1$, the carbamate of 179 is removed, preferably under acidic conditions when R_6 is t-butyl, to afford amines 180. In cases where $p=0$, the esters 179 are converted to the acids 181 preferably under acidic conditions when R_6 is t-butyl.



[0155] Another sequence for preparing amines 180 is illustrated in Scheme 30. Reaction of amines or alcohols 178 with nitro(hetero)arenes 182 wherein M is a leaving group, preferably M is fluoride, or M is a group capable of oxidative insertion into palladium(0), preferably M is bromo, chloro, or iodo, gives intermediates 183. Reduction of the nitro group under standard hydrogenation conditions or treatment with a reducing metal, such as stannous chloride, gives amines 180.




[0156] In instances when hybrid p38-alpha kinase inhibitors are prepared, compounds of Formula I-184 wherein q is 1 may be converted to amines I-185 ($p=1$) or acids I-186 ($p=0$) by analogy to the conditions described in Scheme 29. Compounds of Formula I-184 are prepared as illustrated in previous schemes 1.1, 2.1, 2.2, 3, 4, 5, 6, 7.1, 7.2, 8, 9, 10, 12, 14, 16.2, 17.2, 18, 19.1, 19.2, 19.3, 20, 21, 22, 23, 24, 25, 26, 27, 28.2.




Compounds I-184 are taken from schemes 1.1, 2.1, 2.2, 3, 4, 5, 6, 7.1, 7.2, 8, 9, 10, 12, 14, 16.2, 17.2, 18, 19.1, 19.2, 19.3, 20, 21, 22, 23, 24, 25, 26, 27, 28.2


[0157] The preparation of inhibitors of Formula I which contain an amide linkage $-CO-NH-$ connecting the oxyanion pocket binding moieties and R_1-X-A moieties are shown in Scheme 32. Treatment of acids 181 with an activating agent, preferably PyBOP in the presence of di-iso-propylethylamine, and amines I-185 gives compounds of Formula I. Alternatively, retroamides of Formula I are formed by treatment of acids I-186 with PyBOP in the presence of di-iso-propylethylamine and amines 180.



[0158] The preparation of inhibitors of Formula I which contain an urea linkage $NH-CO-NH-$ connecting the oxyanion pocket binding moieties and the R_1-X-A moieties are shown in Scheme 33. Treatment of amines I-185 with p-nitrophenyl chloroformate and base affords carbamates 187. Reaction of 187 with amines 180 gives ureas of Formula I.



[0159] Alternatively, inhibitors of Formula I which contain an urea linkage NH—CO—NH— connecting the oxyanion pocket binding moieties and the R₁-X-A moieties are prepared as shown in Scheme 33. Treatment of amines 180 with p-nitrophenyl chloroformate and base affords carbamates 188. Reaction of 188 with amines I-185 gives ureas of Formula I.



Affinity and Biological Assessment Of P38-Alpha Kinase Inhibitors

[0160] A fluorescence binding assay is used to detect binding of inhibitors of Formula I with unphosphorylated p38-alpha kinase as previously described: see J. Regan et al, *Journal of Medicinal Chemistry* (2002) 45:2994.

1. P38 MAP Kinase Binding Assay

[0161] The binding affinities of small molecule modulators for p38 MAP kinase were determined using a competition assay with SKF 86002 as a fluorescent probe, modified based on published methods (C. Pargellis, et al *Nature Structural Biology* (2002) 9, 268-272. J. Regan, et al *J. Med. Chem.* (2002) 45, 2994-3008). Briefly, SKF 86002, a potent inhibitor of p38 kinase (K_d=180 nM), displays an emission fluorescence around 420 nm when excited at 340 nm upon its

binding to the kinase. Thus, the binding affinity of an inhibitor for p38 kinase can be measured by its ability to decrease the fluorescence from SKF 86002. The assay was performed in a 384 plate (Greiner uclear 384 plate) on a Polarstar Optima plate reader (BMG). Typically, the reaction mixture contained 1 μM SKF 86002, 80 nM p38 kinase and various concentrations of an inhibitor in 20 mM Bis-Tris Propane buffer, pH 7, containing 0.15% (w/v) n-octylglucoside and 2 mM EDTA in a final volume of 65 μl. The reaction was initiated by addition of the enzyme. The plate was incubated at room temperature (~25° C.) for 2 hours before reading at emission of 420 nm and excitation at 340 nm. By comparison of rfu (relative fluorescence unit) values with that of a control (in the absence of an inhibitor), the percentage of inhibition at each concentration of the inhibitor was calculated. IC₅₀ value for the inhibitor was calculated from the % inhibition values obtained at a range of concentrations of the inhibitor using Prism. When time-dependent inhibition was assessed, the plate was read at multiple reaction times such as 0.5, 1, 2, 3, 4 and 6 hours. The IC₅₀ values were calculated at the each time point. An inhibition was assigned as time-dependent if the IC₅₀ values decrease with the reaction time (more than two-fold in four hours).

Example #	IC50, nM	Time-dependent
1	292	Yes
2	997	No
2	317	No
3	231	Yes
4	57	Yes
5	1107	No
6	238	Yes
7	80	Yes
8	66	Yes
9	859	No
10	2800	No
11	2153	No
12	~10000	No
13	384	Yes
15	949	No
19	~10000	No
21	48	Yes
22	666	No
25	151	Yes
26	68	Yes
29	45	Yes
30	87	Yes
31	50	Yes
32	113	Yes
37	497	No
38	508	No
41	75	Yes
42	373	No
43	642	No
45	1855	No
46	1741	No
47	2458	No
48	3300	No
57	239	Yes

IC50 values obtained at 2 hours reaction time

[0162] Biological assessment of p38-alpha kinase inhibitors of Formula I is performed in a THP-1 cell assay, measuring inhibition of LPS-stimulated TNF-alpha production. See J. Regan et al, *Journal of Medicinal Chemistry* (2002) 45:2994.

EXAMPLES


[0163] The following examples set forth preferred methods in accordance with the invention. It is to be understood, how-

ever, that these examples are provided by way of illustration and nothing therein should be taken as a limitation upon the overall scope of the invention.

[0164] [Boc-sulfamide] aminoester (Reagent AA), 1,5,7-trimethyl-2,4-dioxo-3-aza-bicyclo[3.3.1]nonane-7-carboxylic acid (Reagent BB), and Kemp acid anhydride (Reagent CC) was prepared according to literature procedures. See Askew et. al *J. Am. Chem. Soc.* 1989, 111, 1082 for further details.

EXAMPLE A

[0165]




[0166] To a solution (200 mL) of m-amino benzoic acid (200 g, 1.46 mol) in concentrated HCl was added an aqueous solution (250 mL) of NaNO₂ (102 g, 1.46 mol) at 0° C. The reaction mixture was stirred for 1 h and a solution of SnCl₂·2H₂O (662 g, 2.92 mol) in concentrated HCl (2 L) was then added at 0° C., and the reaction stirred for an additional 2 h at RT. The precipitate was filtered and washed with ethanol and ether to yield 3-hydrazino-benzoic acid hydrochloride as a white solid.

[0167] The crude material from the previous reaction (200 g, 1.06 mol) and 4,4-dimethyl-3-oxo-pentanenitrile (146 g, 1.167 mol) in ethanol (2 L) were heated to reflux overnight. The reaction solution was evaporated in vacuo and the residue purified by column chromatography to yield ethyl 3-(3-tert-butyl-5-amino-1H-pyrazol-1-yl)benzoate (Example A, 116 g, 40%) as a white solid together with 3-(5-amino-3-tert-butyl-1H-pyrazol-1-yl)benzoic acid (93 g, 36%). ¹H NMR (DMSO-d₆): 8.09 (s, 1H), 8.05 (brd, J=8.0 Hz, 1H), 7.87 (brd, J=8.0 Hz, 1H), 7.71 (t, J=8.0 Hz, 1H), 5.64 (s, 1H), 4.35 (q, J=7.2 Hz, 2H), 1.34 (t, J=7.2 Hz, 3H), 1.28 (s, 9H).

EXAMPLE B


[0168]



[0169] To a solution of 1-naphthyl isocyanate (9.42 g, 55.7 mmol) and pyridine (44 mL) in THF (100 mL) was added a solution of Example A (8.0 g, 27.9 mmol) in THF (200 mL) at 0° C. The mixture was stirred at RT for 1 h, heated until all solids were dissolved, stirred at RT for an additional 3 h and quenched with H₂O (200 mL). The precipitate was filtered, washed with dilute HCl and H₂O, and dried in vacuo to yield ethyl 3-[3-tert-butyl-5-(3-naphthalen-1-yl)ureido]-1H-pyrazol-1-yl]benzoate (12.0 g, 95%) as a white powder. ¹H NMR (DMSO-d₆): 9.00 (s, 1H), 8.83 (s, 1H), 8.25 7.42 (m, 11H), 6.42 (s, 1H), 4.30 (q, J=7.2 Hz, 2H), 1.26 (s, 9H), 1.06 (t, J=7.2 Hz, 3H); MS (ESI) m/z: 457.10 (M+H⁺).

EXAMPLE C


[0170]



[0171] To a solution of Example A (10.7 g, 70.0 mmol) in a mixture of pyridine (56 mL) and THF (30 μL) was added a solution of 4-nitrophenyl 4-chlorophenylcarbamate (10 g, 34.8 mmol) in THF (150 mL) at 0° C. The mixture was stirred at RT for 1 h and heated until all solids were dissolved, and stirred at RT for an additional 3 h. H₂O (200 mL) and CH₂Cl₂ (200 mL) were added, the aqueous phase separated and extracted with CH₂Cl₂ (2×100 mL). The combined organic layers were washed with 1N NaOH, and 0.1N HCl, saturated brine and dried over anhydrous Na₂SO₄. The solvent was removed in vacuo to yield ethyl 3-{3-tert-butyl-5-[3-(4-chlorophenyl)ureido]-1H-pyrazol-1-yl}benzoate (8.0 g, 52%). ¹H NMR (DMSO-d₆): δ 9.11 (s, 1H), 8.47 (s, 1H), 8.06 (m, 1H), 7.93 (d, J=7.6 Hz, 1H), 7.81 (d, J=8.0 Hz, 1H), 7.65 (dd, J=8.0, 7.6 Hz, 1H), 7.43 (d, J=8.8 Hz, 2H), 7.30 (d, J=8.8 Hz, 2H), 6.34 (s, 1H), 4.30 (q, J=6.8 Hz, 2H), 1.27 (s, 9H), 1.25 (t, J=6.8 Hz, 3H); MS (ESI) m/z: 441 (M⁺+H).

EXAMPLE D


[0172]



[0173] To a stirred solution of Example B (8.20 g, 18.0 mmol) in THF (500 mL) was added LiAlH_4 powder (2.66 g, 70.0 mmol) at -10°C . under N_2 . The mixture was stirred for 2 h at RT and excess LiAlH_4 destroyed by slow addition of ice. The reaction mixture was acidified to pH=7 with dilute HCl, concentrated in vacuo and the residue extracted with EtOAc. The combined organic layers were concentrated in vacuo to yield 1-{3-tert-butyl-1-[3-(hydroxymethyl)phenyl]-1H-pyrazol-5-yl}-3-(naphthalen-1-yl)urea (7.40 g, 99%) as a white powder. $^1\text{H NMR}$ (DMSO-d_6): 9.19 (s, 1H), 9.04 (s, 1H), 8.80 (s, 1H), 8.26-7.35 (m, 11H), 6.41 (s, 1H), 4.60 (s, 2H), 1.28 (s, 9H); MS (ESI) m/z: 415 ($\text{M}+\text{H}^+$).

EXAMPLE E


[0174]



[0175] A solution of Example C (1.66 g, 4.0 mmol) and SOCl_2 (0.60 mL, 8.0 mmol) in CH_2Cl_2 (100 mL) was refluxed for 3 h and concentrated in vacuo to yield 1-{3-tert-butyl-1-[3-(chloromethyl)phenyl]-1H-pyrazol-5-yl}-3-(naphthalen-1-yl)urea (1.68 g, 97%) was obtained as white powder. $^1\text{H NMR}$ (DMSO-d_6): δ 9.26 (s, 1H), 9.15 (s, 1H), 8.42-7.41 (m, 11H), 6.40 (s, 1H), 4.85 (s, 2H), 1.28 (s, 9H). MS (ESI) m/z: 433 ($\text{M}+\text{H}^+$).

EXAMPLE F

[0176]




[0177] To a stirred solution of Example C (1.60 g, 3.63 mmol) in THF (200 mL) was added LiAlH_4 powder (413 mg, 10.9 mmol) at -10°C . under N_2 . The mixture was stirred for 2 h and excess LiAlH_4 was quenched by adding ice. The solution was acidified to pH=7 with dilute HCl. Solvents were slowly removed and the solid was filtered and washed with EtOAc (200+100 mL). The filtrate was concentrated to yield 1-{3-tert-butyl-1-[3-(hydroxymethyl)phenyl]-1H-pyrazol-5-yl}-3-(4-chlorophenyl)urea (1.40 g, 97%). $^1\text{H NMR}$ (DMSO-d_6): δ 9.11 (s, 1H), 8.47 (s, 1H), 7.47-7.27 (m, 8H), 6.35 (s,

1H), 5.30 (t, J=5.6 Hz, 1H), 4.55 (d, J=5.6 Hz, 2H), 1.26 (s, 9H); MS (ESI) m/z: 399 ($\text{M}+\text{H}^+$).

EXAMPLE G


[0178]



[0179] A solution of Example F (800 mg, 2.0 mmol) and SOCl_2 (0.30 mL, 4 mmol) in CHCl_3 (30 mL) was refluxed gently for 3 h. The solvent was evaporated in vacuo and the residue was taken up to in CH_2Cl_2 (2x20 mL). After removal of the solvent, 1-{3-tert-butyl-1-[3-(chloromethyl)phenyl]-1H-pyrazol-5-yl}-3-(4-chlorophenyl)urea (812 mg, 97%) was obtained as white powder. $^1\text{H NMR}$ (DMSO-d_6): δ 9.57 (s, 1H), 8.75 (s, 1H), 7.63 (s, 1H), 7.50-7.26 (m, 7H), 6.35 (s, 1H), 4.83 (s, 2H), 1.27 (s, 9H); MS (ESI) m/z: 417 ($\text{M}+\text{H}^+$).

EXAMPLE H

[0180]




[0181] To a suspension of LiAlH_4 (5.28 g, 139.2 mmol) in THF (1000 mL) was added Example A (20.0 g, 69.6 mmol) in portions at 0°C . under N_2 . The reaction mixture was stirred for 5 h, quenched with 1 N HCl at 0°C . and the precipitate was filtered, washed by EtOAc and the filtrate evaporated to yield [3-(5-amino-3-tert-butyl-1H-pyrazol-1-yl)phenyl]methanol (15.2 g, 89%). $^1\text{H NMR}$ (DMSO-d_6): 7.49 (s, 1H), 7.37 (m, 2H), 7.19 (d, J=7.2 Hz, 1H), 5.35 (s, 1H), 5.25 (t, J=5.6 Hz, 1H), 5.14 (s, 2H), 4.53 (d, J=5.6 Hz, 2H), 1.19 (s, 9H); MS (ESI) m/z: 246.19 ($\text{M}+\text{H}^+$).

[0182] The crude material from the previous reaction (5.0 g, 20.4 mmol) was dissolved in dry THF (50 mL) and SOCl_2 (4.85 g, 40.8 mmol), stirred for 2 h at RT, concentrated in vacuo to yield 3-tert-butyl-1-(3-chloromethylphenyl)-1H-pyrazol-5-amine (5.4 g), which was added to N_3 (3.93 g, 60.5 mmol) in DMF (50 mL). The reaction mixture was heated at 30°C . for 2 h, poured into H_2O (50 mL), and extracted with CH_2Cl_2 . The organic layers were combined, dried over

MgSO₄, and concentrated in vacuo to yield crude 3-tert-butyl-1-[3-(azidomethyl)phenyl]-1H-pyrazol-5-amine (1.50 g, 5.55 mmol).

EXAMPLE I

[0183]




[0184] Example H was dissolved in dry THF (10 mL) and added a THF solution (10 mL) of 1-isocyano naphthalene (1.13 g, 6.66 mmol) and pyridine (5.27 g, 66.6 mmol) at RT. The reaction mixture was stirred for 3 h, quenched with H₂O (30 mL), the resulting precipitate filtered and washed with 1 N HCl and ether to yield 1-[2-(3-azidomethyl-phenyl)-5-tert-butyl-2H-pyrazol-3-yl]-3-naphthalen-1-yl-urea (2.4 g, 98%) as a white solid.

[0185] The crude material from the previous reaction and Pd/C (0.4 g) in THF (30 mL) was hydrogenated under 1 atm at RT for 2 h. The catalyst was removed by filtration and the filtrate concentrated in vacuo to yield 1-{3-tert-butyl-1-[3-(amonomethyl)phenyl]-1H-pyrazol-5-yl}-3-(naphthalene-1-yl)urea (2.2 g, 96%) as a yellow solid. ¹H NMR (DMSO-d₆): 9.02 (s, 1H), 7.91 (d, J=7.2 Hz, 1H), 7.89 (d, J=7.6 Hz, 2H), 7.67-7.33 (m, 9H), 6.40 (s, 1H), 3.81 (s, 2H), 1.27 (s, 9H); MS (ESI) m/z: 414 (M+H⁺).

EXAMPLE J

[0186]




[0187] To a solution of Example H (1.50 g, 5.55 mmol) in dry THF (10 mL) was added a THF solution (10 mL) of 4-chlorophenyl isocyanate (1.02 g, 6.66 mmol) and pyridine (5.27 g, 66.6 mmol) at RT. The reaction mixture was stirred for 3 h and then H₂O (30 mL) was added. The precipitate was filtered and washed with 1N HCl and ether to give 1-{3-tert-butyl-1-[3-(amonomethyl)phenyl]-1H-pyrazol-5-yl}-3-(4-

chlorophenyl)urea (2.28 g, 97%) as a white solid, which was used for next step without further purification. MS (ESI) m/z: 424 (M+H⁺).

EXAMPLE K

[0188]




[0189] To a solution of benzyl amine (16.5 g, 154 mmol) and ethyl bromoacetate (51.5 g, 308 mmol) in ethanol (500 mL) was added K₂CO₃ (127.5 g, 924 mmol). The mixture was stirred at RT for 3 h, was filtered, washed with EtOH, concentrated in vacuo and chromatographed to yield N-(2-ethoxy-2-oxoethyl)-N-(phenylmethyl)-glycine ethyl ester (29 g, 67%). ¹H NMR (CDCl₃): δ 7.39-7.23 (m, 5H), 4.16 (q, J=7.2 Hz, 4H), 3.91 (s, 2H), 3.54 (s, 4H), 1.26 (t, J=7.2 Hz, 6H); MS (ESI): m/e: 280 (M⁺+H).

[0190] A solution of N-(2-ethoxy-2-oxoethyl)-N-(phenylmethyl)-glycine ethyl ester (7.70 g, 27.6 mmol) in methanol solution (25-30%, 50 mL) was heated to 50° C. in a sealed tube for 3 h, cooled to RT and concentrated in vacuo to yield N-(2-methylamino-2-oxoethyl)-N-(phenylmethyl)-glycine methylamide in quantitative yield (7.63 g). ¹H NMR (CDCl₃): δ 7.35-7.28 (m, 5H), 6.75 (br s, 2H), 3.71 (s, 2H), 3.20 (s, 4H), 2.81 (d, J=5.6 Hz, 6H); MS (ESI) m/e 250 (M+H⁺).

[0191] The mixture of N-(2-methylamino-2-oxoethyl)-N-(phenylmethyl)-glycine methylamide (3.09 g, 11.2 mmol) in MeOH (30 mL) was added 10% Pd/C (0.15 g). The mixture was stirred and heated to 40° C. under 40 psi H₂ for 10 h, filtered and concentrated in vacuo to yield N-(2-methylamino-2-oxoethyl)-glycinemethylamide in quantitative yield (1.76 g). ¹H NMR (CDCl₃): δ 6.95 (br s, 2H), 3.23 (s, 4H), 2.79 (d, J=6.0, 4.8 Hz), 2.25 (br s 1H); MS (ESI) m/e 160 (M+H⁺).

EXAMPLE 1

[0192]




[0193] To a solution of 1-methyl-[1,2,4]triazolidine-3,5-dione (188 mg, 16.4 mmol) and sodium hydride (20 mg, 0.52 mmol) in DMSO (1 mL) was added Example E (86 mg, 0.2 mmol). The reaction was stirred at RT overnight, quenched with H₂O (10 mL), extracted with CH₂Cl₂, and the organic

layer was separated, washed with brine, dried over Na_2SO_4 and concentrated in vacuo. The residue was purified by preparative HPLC to yield 1-(3-tert-butyl-1-{3-[(1-methyl-3,5-dioxo-1,2,4-triazolidin-4-yl)methyl]phenyl}-1H-pyrazol-5-yl)-3-(naphthalene-1-yl)urea (Example 1, 14 mg). $^1\text{H NMR}$ (CD_3OD): δ 7.88-7.86 (m, 2H), 7.71-7.68 (m, 2H), 7.58 (m, 2H), 7.60-7.42 (m, 5H), 6.49 (s, 1H), 4.85 (s, 1H), 1.34 (s, 9H), 1.27 (s, 6H); MS (ESI) m/z : 525 ($\text{M}+\text{H}^+$).

EXAMPLE 2


[0194]



[0195] The title compound was synthesized in a manner analogous to Example 1, utilizing Example G to yield 1-(3-tert-butyl-1-{3-[(1-methyl-3,5-dioxo-1,2,4-triazolidin-4-yl)methyl]phenyl}-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea. $^1\text{H NMR}$ (CD_3OD): δ 7.2-7.5 (m, 7H), 6.40 (s, 1H), 4.70 (s, 2H), 2.60 (d, $J=14$ Hz, 2H), 1.90 (m, 1H), 1.50 (m, 1H), 1.45 (s, 9H), 1.30 (m, 2H), 1.21 (s, 3H), 1.18 (s, 6H); MS (ESI) m/z : 620 ($\text{M}+\text{H}^+$).

EXAMPLE 3

[0196]




[0197] A mixture of compound 1,1-Dioxo-[1,2,5]thiadiazolidin-3-one (94 mg, 0.69 mmol) and NaH (5.5 mg, 0.23 mmol) in THF (2 mL) was stirred at -10°C . under N_2 for 1 h until all NaH was dissolved. Example E (100 mg, 0.23 mmol) was added and the reaction was allowed to stir at RT overnight, quenched with H_2O , and extracted with CH_2Cl_2 . The combined organic layers were concentrated in vacuo and the residue was purified by preparative HPLC to yield 1-(3-tert-butyl-1-{3-[(1,1,3-trioxo-[1,2,5]thiadiazolidin-2-yl)methyl]phenyl}-1H-pyrazol-5-yl)-3-(naphthalen-1-yl)urea (18 mg)

as a white powder. $^1\text{H NMR}$ (CD_3OD): δ 7.71-7.44 (m, 11H), 6.45 (s, 1H), 4.83 (s, 2H), 4.00 (s, 2H), 1.30 (s, 9H). MS (ESI) m/z : 533.40 ($\text{M}+\text{H}^+$).

EXAMPLE 4


[0198]



[0199] The title compound was obtained in a manner analogous to Example 3 utilizing Example G. to yield 1-(3-tert-butyl-1-{3-[(1,1,3-trioxo-[1,2,5]thiadiazolidin-2-yl)methyl]phenyl}-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea. $^1\text{H NMR}$ (CD_3OD): δ 7.38-7.24 (m, 8H), 6.42 (s, 1H), 4.83 (s, 2H), 4.02 (s, 2H), 1.34 (s, 9H); MS (ESI) m/z : 517 ($\text{M}+\text{H}^+$).

EXAMPLE 5

[0200]




[0201] To a stirred solution of chlorosulfonyl isocyanate (19.8 μL , 0.227 mmol) in CH_2Cl_2 (0.5 mL) at 0°C . was added pyrrolidine (18.8 μL , 0.227 mmol) at such a rate that the reaction solution temperature did not rise above 5°C . After stirring for 1.5 h, a solution of Example J (97.3 mg, 0.25 mmol) and Et_3N (95 μL , 0.678 mmol) in CH_2Cl_2 (1.5 mL) was added at such a rate that the reaction temperature didn't rise above 5°C . When the addition was completed, the reaction solution was warmed to RT and stirred overnight. The reaction mixture was poured into 10% HCl, extracted with CH_2Cl_2 , the organic layer washed with saturated NaCl, dried over MgSO_4 , and filtered. After removal of the solvents, the crude product was purified by preparative HPLC to yield 1-(3-tert-butyl-1-{3-[(1-pyrrolidinylcarbonyl)amino]sulphonylaminomethyl]phenyl}-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea. $^1\text{H NMR}$ (CD_3OD): δ 7.61 (s, 1H), 7.43-7.47 (m, 3H), 7.23-7.25 (dd, $J=6.8$ Hz, 2H), 7.44 (dd, $J=6.8$

Hz, 2H), 6.52 (s, 1H), 4.05 (s, 2H), 3.02 (m, 4H), 1.75 (m, 4H), 1.34 (s, 9H); MS (ESI) m/z: 574.00 (M+H⁺).

EXAMPLE 6


[0202]



[0203] The title compound was made in a manner analogous to Example 5 utilizing Example I to yield 1-(3-tert-butyl-1-[[3-N-[[1-(pyrrolidinylsulfonyl)amino]sulfonyl]aminomethyl]-phenyl]-1H-pyrazol-5-yl)-3-(naphthalen-1-yl)urea. ¹H NMR (CDCl₃): δ 7.88 (m, 2H), 7.02-7.39 (m, 2H), 7.43-7.50 (m, 7H), 6.48 (s, 1H), 4.45 (s, 1H), 3.32-3.36 (m, 4H), 1.77-1.81 (m, 4H), 1.34 (s, 9H); MS (ESI) m/z: 590.03 (M+H⁺).

EXAMPLE 7

[0204]




[0205] To a stirred solution of chlorosulfonyl isocyanate (19.8 μL, 0.227 μmol) in CH₂Cl₂ (0.5 mL) at 0° C., was added Example J (97.3 mg, 0.25 mmol) at such a rate that the reaction solution temperature did not rise above 5° C. After being stirred for 1.5 h, a solution of pyrrolidine (18.8 μL, 0.227 mmol) and Et₃N (95 μL, 0.678 mmol) in CH₂Cl₂ (1.5 mL) was added at such a rate that the reaction temperature didn't rise above 5° C. When addition was completed, the reaction solution was warmed to RT and stirred overnight. The reaction mixture was poured into 10% HCl, extracted with CH₂Cl₂, the organic layer was washed with saturated NaCl, dried over Mg₂SO₄, and filtered. After removal of the solvents, the crude product was purified by preparative HPLC to yield 1-(3-tert-butyl-1-[[3-N-[[1-(pyrrolidinylsulfonyl)amino]sulfonyl]aminomethyl]phenyl]-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea. ¹H NMR (CDCl₃): δ 7.38 (m, 1H), 7.36-7.42 (m, 3H), 7.23 (d, J=8.8 Hz, 2H), 7.40 (d, J=8.8 Hz,

2H), 6.43 (s, 1H), 4.59 (s, 1H), 4.43 (s, 2H), 1.81 (s, 2H), 1.33 (s, 9H); MS (ESI) m/z: 574.10 (M+H⁺).

EXAMPLE 8


[0206]



[0207] The title compound was made in a manner analogous to Example 7 utilizing Example I to yield 1-(3-tert-butyl-1-[[3-N-[[1-(pyrrolidinylsulfonyl)amino]sulfonyl]aminomethyl]-phenyl]-1H-pyrazol-5-yl)-3-(naphthalen-1-yl)urea. ¹H NMR (CDCl₃): δ 7.88 (m, 2H), 7.02-7.39 (m, 2H), 7.43-7.50 (m, 7H), 6.48 (s, 1H), 4.45 (s, 1H), 3.32-3.36 (m, 4H), 1.77-1.81 (m, 4H), 1.34 (s, 9H); MS (ESI) m/z: 590.03 (M+H⁺).

EXAMPLE 9

[0208]




[0209] To a solution of Reagent BB (36 mg, 0.15 mmol), Example I (62 mg, 0.15 mmol), HOBt (40 mg, 0.4 mmol) and NMM (0.1 mL, 0.9 mmol) in DMF (10 mL) was added EDCI (58 mg, 0.3 mmol). After being stirred overnight, the mixture was poured into water (15 mL) and extracted with EtOAc (3 x 5 mL). The organic layers were combined, washed with brine, dried with Na₂SO₄, and concentrated in vacuo. The residue was purified by preparative TLC to yield 1,5,7-trimethyl-2,4-dioxo-3-azabicyclo[3.3.1]nonane-7-carboxylic acid 3-[3-tert-butyl-5-(3-naphthalen-1-yl-ureido)-pyrazol-1-yl]benzylamide (22 mg). ¹H NMR (CDCl₃): δ 8.40 (s, 1H), 8.14 (d, J=8.0 Hz, 2H), 7.91 (s, 1H), 7.87 (s, 1H), 7.86 (d, J=7.2 Hz,

1H), 7.78 (d, J=7.6 Hz, 1H), 7.73 (d, J=8.4 Hz, 1H), 7.69 (d, J=8.4 Hz, 1H), 7.57-7.40 (m, 4H), 7.34 (d, J=7.6 Hz, 1H), 6.69 (s, 1H), 6.32 (t, J=5.6 Hz, 1H), 5.92 (brs, 1H), 4.31 (d, J=5.6 Hz, 2H), 2.37 (d, J=14.8 Hz, 2H), 1.80 (d, J=13.2 Hz, 1H), 1.35 (s, 9H), 1.21 (d, J=13.2 Hz, 1H), 1.15 (s, 3H), 1.12 (d, J=12.8 Hz, 2H), 1.04 (s, 6H); MS (ESI) m/z: 635 (M+H⁺).

EXAMPLE 10


[0210]



[0211] The title compound, was synthesized in a manner analogous to Example 9 utilizing Example J to yield 1,5,7-trimethyl-2,4-dioxo-3-aza-bicyclo[3.3.1]nonane-7-carboxylic acid 3-{3-t-butyl-5-[3-(4-chloro-phenyl)-ureido]-pyrazol-1-yl}benzylamide. ¹H NMR (CDCl₃): δ 8.48 (s, 1H), 7.78 (s, 1H), 7.75 (d, J=8.0 Hz, 1H), 7.69 (s, 1H), 7.53 (t, J=8.0 Hz, 1H), 7.48 (d, J=8.8 Hz, 2H), 7.26 (m, 3H), 6.62 (s, 1H), 6.35 (t, J=6.0 Hz, 1H), 5.69 (brs, 1H), 4.26 (d, J=6.0 Hz, 2H), 2.48 (d, J=14.0 Hz, 2H), 1.87 (d, J=13.6 Hz, 1H), 1.35 (s, 9H), 1.25 (m, 6H), 1.15 (s, 6H); MS (ESI) m/z: 619 (M+H⁺).

EXAMPLE 11


[0212]



[0213] A mixture of Example I (41 mg, 0.1 mmol), Kemp acid anhydride (24 mg, 0.1 mmol) and Et₃N (100 mg, 1 mmol) in anhydrous CH₂Cl₂ (2 mL) were stirred overnight at RT, and concentrated in vacuo. Anhydrous benzene (20 mL) was added to the residue, the mixture was refluxed for 3 h, concentrated in vacuo and purified by preparative HPLC to yield 3-{3-[3-t-butyl-5-(3-naphthalen-1-yl-ureido)-pyrazol-1-yl]-benzyl}-1,5-di-methyl-2,4-dioxo-3-aza-bicyclo[3.3.1]nonane-7-carboxylic acid (8.8 mg, 14%). ¹H NMR (CD₃OD): δ 7.3-7.4 (m, 2H), 7.20 (m, 2H), 7.4-7.6 (m, 7H), 6.50 (m, 1H), 4.80 (s, 2H), 2.60 (d, J=14 Hz, 2H), 1.90 (m, 1H), 1.40 (m, 1H), 1.30 (m, 2H), 1.20 (s, 3H), 1.15 (s, 6H); MS (ESI) m/z: 636 (M+H⁺).

EXAMPLE 12


[0214]



[0215] The title compound, was synthesized in a manner analogous to Example 11 utilizing Example J to yield 3-{3-[3-t-butyl-5-(3-naphthalen-1-yl-ureido)-pyrazol-1-yl]-benzyl}-1,5-dimethyl-2,4-dioxo-3-aza-bicyclo[3.3.1]nonane-7-carboxylic acid. ¹H NMR (CD₃OD): δ 7.2-7.5 (m, 7H), 6.40 (s, 1H), 4.70 (s, 2H), 2.60 (d, J=14 Hz, 2H), 1.90 (m, 1H), 1.50 (m, 1H), 1.45 (s, 9H), 1.30 (m, 2H), 1.21 (s, 3H), 1.18 (s, 6H); MS (ESI) m/z: 620 (M+H⁺).

EXAMPLE 13


[0216]



[0217] The title compound was synthesized in a manner analogous to Example 1 utilizing Example E and 4,4-dimethyl-3,5-dioxo-pyrazolidine to yield 1-(3-tert-butyl-1-{3-[(4,4-dimethyl-3,5-dioxopyrazolidin-1-yl)methyl]phenyl}-1H-pyrazol-5-yl)-3-(naphthalen-1-yl)urea. ¹H NMR (CD₃OD): δ 7.88-7.86 (m, 2H), 7.71-7.68 (m, 2H), 7.58 (m, 2H), 7.60-7.42 (m, 5H), 6.49 (s, 1H), 4.85 (s, 1H), 1.34 (s, 9H), 1.27 (s, 6H); MS (ESI) m/z: 525 (M+H⁺).

EXAMPLE 14


[0218]



[0219] The title compound was synthesized in a manner analogous to Example 1 utilizing Example G and 4,4-dimethyl-3,5-dioxo-pyrazolidine to yield 1-(3-tert-butyl-1-{3-[(4,4-dimethyl-3,5-dioxopyrazolidin-1-yl)methyl]phenyl}-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea. ¹H NMR (CD₃OD): δ 7.60-7.20 (m, 8H), 6.43 (s, 1H), 4.70 (s, 1H), 1.34 (s, 9H), 1.26 (s, 6H); MS (ESI) m/z: 509, 511 (M+H⁺).

EXAMPLE 15

[0220]




[0221] Example B was saponified with 2N LiOH in MeOH, and to the resulting acid (64.2 mg, 0.15 mmol) were added HOBt (30 mg, 0.225 mmol), Example K (24 mg, 0.15 mmol) and 4-methylmorpholine (60 mg, 0.60 mmol 4.0 equiv), DMF (3 mL) and EDCI (43 mg, 0.225 mmol). The reaction mixture was stirred at RT overnight and poured into H₂O (3 mL), and

a white precipitate collected and further purified by preparative HPLC to yield 1-[1-(3-{bis[(methylcarbamoyl)methyl]phenyl}-3-tert-butyl-1H-pyrazol-5-yl)-3-(naphthalen-1-yl)urea (40 mg). ¹H NMR (CDCl₃): δ 8.45 (brs, 1H), 8.10 (d, J=7.6 Hz, 1H), 7.86-7.80 (m, 2H), 7.63-7.56 (m, 2H), 7.52 (s, 1H), 7.47-7.38 (m, 3H), 7.36-7.34 (m, 1H), 7.26 (s, 1H), 7.19-7.17 (m, 2H), 6.60 (s, 1H), 3.98 (s, 2H), 3.81 (s, 3H), 2.87 (s, 3H), 2.63 (s, 3H), 1.34 (s, 9H); MS (ESI) m/z: 570 (M+H⁺).

EXAMPLE 16


[0222]



[0223] The title compound was synthesized in a manner analogous to Example 15 utilizing Example C (37 mg) and Example K to yield 1-[1-(3-{bis[(methylcarbamoyl)methyl]phenyl}-3-tert-butyl-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea. ¹H NMR (CD₃OD): δ 8.58 (brs, 1H), 8.39 (brs, 1H), 7.64-7.62 (m, 3H), 7.53-7.51 (m, 1H), 7.38 (d, J=9.2 Hz, 2H), 7.25 (d, J=8.8 Hz, 2H), 6.44 (s, 1H), 4.17 (s, 2H), 4.11 (s, 2H), 2.79 (s, 3H), 2.69 (s, 3H), 1.34-1.28 (m, 12H); MS (ESI) m/z: 554 (M+H⁺).

EXAMPLE 17

[0224]




[0225] Example B was saponified with 2N LiOH in MeOH, and to the resulting acid (0.642 g, 1.5 mmol) in dry THF (25 mL) at -78° C. were added freshly distilled triethylamine (0.202 g, 2.0 mmol) and pivaloyl chloride (0.216 g, 1.80 mmol) with vigorous stirring. After stirring at -78° C. for 15 min and at 0° C. for 45 min, the mixture was again cooled to

-78° C. and then transferred into the THF solution of lithium salt of D-4-phenyl-oxazolidin-2-one [*]: The lithium salt of the oxazolidinone reagent was previously prepared by the slow addition of n-BuLi (2.50M in hexane, 1.20 mL, 3.0 mmol) into THF solution of D-4-phenyl-oxazolidin-2-one at -78° C.]. The reaction solution was stirred at -78° C. for 2 h and RT overnight, and then quenched with aq. ammonium chloride and extracted with dichloromethane (100 mL). The combined organic layers were dried (Na₂SO₄) and concentrated in vacuo. The residue was purified by preparative HPLC to yield D-1-{5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl-3-carbonyl)phenyl]-2H-pyrazol-3-yl}-3-(naphthalen-1-yl)urea (207 mg, 24%). ¹H NMR (CDCl₃): δ 8.14-8.09 (m, 2H), 8.06 (s, 1H), 7.86-7.81 (m, 4H), 7.79 (s, 1H), 7.68-7.61 (m, 2H), 7.51-7.40 (m, 9H), 6.75 (s, 1H), 5.80 (t, J=9.2, 7.6 Hz, 1H), 4.89 (t, J=9.2 Hz, 1H), 4.42 (dd, J=9.2, 7.6 Hz, 1H), 1.37 (s, 9H); MS (ESI) m/z: 574 (M+H⁺).

EXAMPLE 18


[0226]



[0227] The title compound was synthesized in a manner analogous to Example 17 utilizing Example B and L-4-phenyl-oxazolidin-2-one to yield L-1-{5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl-3-carbonyl)phenyl]-2H-pyrazol-3-yl}-3-(naphthalen-1-yl)urea. ¹H NMR (CDCl₃): δ 8.14-8.09 (m, 2H), 8.06 (s, 1H), 7.86-7.81 (m, 4H), 7.79 (s, 1H), 7.68-7.61 (m, 2H), 7.51-7.40 (m, 9H), 6.75 (s, 1H), 5.80 (t, J=9.2, 7.6 Hz, 1H), 4.89 (t, J=9.2 Hz, 1H), 4.42 (dd, J=9.2, 7.6 Hz, 1H), 1.37 (s, 9H); MS (ESI) m/z: 574 (M+H⁺).

EXAMPLE 19


[0228]



[0229] The title compound was synthesized in a manner analogous to Example 17 utilizing Example C and D-4-phenyl-oxazolidin-2-one to yield D-1-{5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl-3-carbonyl)phenyl]-2H-pyrazol-3-yl}-3-(4-chlorophenyl)urea. ¹H NMR (CDCl₃): δ 7.91 (s, 1H), 7.85 (d, J=8.0 Hz, 1H), 7.79 (d, J=7.6 Hz, 1H), 7.71 (m, 1H), 7.65 (m, 1H), 7.49-7.40 (m, 8H), 7.26-7.24 (m, 2H), 6.68 (s, 1H), 5.77 (dd, J=8.8, 8.0 Hz, 1H), 4.96 (t, 8.8 Hz, 1H), 4.44 (dd, J=8.8, 8.0 Hz, 1H), 1.36 (s, 9H); MS (ESI) m/z: 558 (M+H⁺).

EXAMPLE 20


[0230]



[0231] The title compound was synthesized in a manner analogous to Example 17 utilizing Example C and L-4-phenyl-oxazolidin-2-one to yield L-1-{5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl-3-carbonyl)phenyl]-2H-pyrazol-3-yl}-3-(4-chlorophenyl)urea. ¹H NMR (CDCl₃): δ 7.91 (s, 1H), 7.85 (d, J=8.0 Hz, 1H), 7.79 (d, J=7.6 Hz, 1H), 7.71 (m, 1H), 7.65 (m, 1H), 7.49-7.40 (m, 8H), 7.26-7.24 (m, 2H), 6.68 (s, 1H), 5.77 (dd, J=8.8, 8.0 Hz, 1H), 4.96 (t, 8.8 Hz, 1H), 4.44 (dd, J=8.8, 8.0 Hz, 1H), 1.36 (s, 9H); MS (ESI) m/z: 558 (M+H⁺).


EXAMPLE L

[0232]




[0233] To a stirred suspension of (3-nitro-phenyl)-acetic acid (2 g) in CH_2Cl_2 (40 ml, with a catalytic amount of DMF) at 0°C . under N_2 was added oxalyl chloride (1.1 ml) drop wise. The reaction mixture was stirred for 40 min morpholine (2.5 g) was added. After stirring for 20 min, the reaction mixture was filtered. The filtrate was concentrated in vacuo to yield 1-morpholin-4-yl-2-(3-nitro-phenyl)-ethanone as a solid (2 g). A mixture of 1-morpholin-4-yl-2-(3-nitro-phenyl)-ethanone (2 g) and 10% Pd on activated carbon (0.2 g) in ethanol (30 ml) was hydrogenated at 30 psi for 3 h and filtered over Celite. Removal of the volatiles in vacuo provided 2-(3-amino-phenyl)-1-morpholin-4-yl-ethanone (1.7 g). A solution of 2-(3-amino-phenyl)-1-morpholin-4-yl-ethanone (1.7 g, 7.7 mmol) was dissolved in 6 N HCl (15 ml), cooled to 0°C ., and vigorously stirred. Sodium nitrite (0.54 g) in water (8 ml) was added. After 30 min, tin (II) chloride dihydrate (10 g) in 6 N HCl (30 ml) was added. The reaction mixture was stirred at 0°C . for 3 h. The pH was adjusted to pH 14 with solid potassium hydroxide and extracted with EtOAc. The combined organic extracts were concentrated in vacuo provided 2-(3-hydrazin-phenyl)-1-morpholin-4-yl-ethanone (1.5 g). 2-(3-Hydrazinophenyl)-1-morpholin-4-yl-ethanone (3 g) and 4,4-dimethyl-3-oxopentanenitrile (1.9 g, 15 mmol) in ethanol (60 ml) and 6 N HCl (1 ml) were refluxed for 1 h and cooled to RT. The reaction mixture was neutralized by adding solid sodium hydrogen carbonate. The slurry was filtered and removal of the volatiles in vacuo provided a residue that was extracted with ethyl acetate. The volatiles were removed in vacuo to provide 2-[3-(3-tert-butyl-5-amino-1H-pyrazol-1-yl)phenyl]-1-morpholinoethanone (4 g), which was used without further purification.

EXAMPLE 21

[0234]


[0235] A mixture of Example L (0.2 g, 0.58 mmol) and 1-naphthylisocyanate (0.10 g, 0.6 mmol) in dry CH_2Cl_2 (4 ml) was stirred at RT under N_2 for 18 h. The solvent was removed in vacuo and the crude product was purified by column chromatography using ethyl acetate/hexane/ CH_2Cl_2 (3/1/0.7) as the eluent (0.11 g, off-white solid) to yield 1-[3-tert-butyl-1-[3-(2-morpholino-2-oxoethyl)phenyl]-1H-pyrazol-5-yl]-3-(naphthalene-1-yl)urea. mp: 194-196; ^1H NMR (200 MHz, DMSO-d_6): δ 9.07 (1H, s), 8.45 (s, 1H), 8.06-7.93 (m, 3H), 7.69-7.44 (m, 7H), 7.33-7.29 (d, 6.9 Hz, 1H), 6.44 (s, 1H), 3.85 (m, 2H), 3.54-3.45 (m, 8H), 1.31 (s, 9H); MS:

EXAMPLE 22

[0236]


[0237] The title compound was synthesized in a manner analogous to Example 21 utilizing Example L (0.2 g, 0.58 mmol) and 4-chlorophenylisocyanate (0.09 g, 0.6 mmol) to yield 1-[3-tert-butyl-1-[3-(2-morpholino-2-oxoethyl)phenyl]-1H-pyrazol-5-yl]-3-(4-chlorophenyl)urea. mp: 100-104; ^1H NMR (200 MHz, DMSO-d_6): δ 9.16 (s, 1H), 8.45 (s, 1H), 7.52-7.30 (m, 8H), 6.38 (s, 1H), 3.83 (m, 1H), 3.53-3.46 (m, 8H), 1.30 (s, 9H); MS:

EXAMPLE 23

[0238]

[0239] The title compound is synthesized in a manner analogous to Example 21 utilizing Example L (0.2 g, 0.58 mmol) and phenylisocyanate (0.09 g, 0.6 mmol) to yield 1-[3-tert-butyl-1-[3-(2-morpholino-2-oxoethyl)phenyl]-1H-pyrazol-5-yl]-3-phenylurea.


EXAMPLE 24

[0240]

[0241] The title compound is synthesized in a manner analogous to Example 21 utilizing Example L (0.2 g, 0.58 mmol) and 1-isocyanato-4-methoxy-naphthalene to yield 1-{3-tert-butyl-1-[3-(2-morpholino-2-oxoethyl)phenyl]-1H-pyrazol-5-yl}-3-(1-methoxynaphthalen-4-yl)urea.

EXAMPLE M


[0242]



[0243] The title compound is synthesized in a manner analogous to Example C utilizing Example A and phenylisocyanate to yield ethyl 3-(3-tert-butyl-5-(3-phenylureido)-1H-pyrazol-1-yl)benzoate.

EXAMPLE N

[0244]



[0245] A solution of (3-nitrophenyl)acetic acid (23 g, 127 mmol) in methanol (250 ml) and a catalytic amount of concentrated in vacuo H_2SO_4 was heated to reflux for 18 h. The reaction mixture was concentrated in vacuo to a yellow oil. This was dissolved in methanol (250 ml) and stirred for 18 h in an ice bath, whereupon a slow flow of ammonia was charged into the solution. The volatiles were removed in vacuo. The residue was washed with diethyl ether and dried to afford 2-(3-nitrophenyl)acetamide (14 g, off-white solid). 1H NMR ($CDCl_3$): δ 8.1 (s, 1H), 8.0 (d, 1H), 7.7 (d, 1H), 7.5 (m, 1H), 7.1 (bd s, 1H), 6.2 (brs, 1H), 3.6 (s, 2H).


[0246] The crude material from the previous reaction (8 g) and 10% Pd on activated carbon (1 g) in ethanol (100 ml) was hydrogenated at 30 psi for 18 h and filtered over Celite. Removal of the volatiles in vacuo provided 2-(3-aminophenyl)acetamide (5.7 g). A solution of this material (7 g, 46.7 mmol) was dissolved in 6 N HCl (100 ml), cooled to 0° C.,

and vigorously stirred. Sodium nitrite (3.22 g, 46.7 mmol) in water (50 ml) was added. After 30 min, tin (II) chloride dihydrate (26 g) in 6 N HCl (100 ml) was added. The reaction mixture was stirred at 0° C. for 3 h. The pH was adjusted to pH 14 with 50% aqueous NaOH solution and extracted with ethyl acetate. The combined organic extracts were concentrated in vacuo provided 2-(3-hydrazinophenyl)acetamide.

[0247] The crude material from the previous reaction (ca. 15 mmol) and 4,4-dimethyl-3-oxopentenenitrile (1.85 g, 15 mmol) in ethanol (60 ml) and 6 N HCl (1.5 ml) was refluxed for 1 h and cooled to RT. The reaction mixture was neutralized by adding solid sodium hydrogen carbonate. The slurry was filtered and removal of the volatiles in vacuo provided a residue, which was extracted with ethyl acetate. The solvent was removed in vacuo to provide 2-[3-(3-tert-butyl-5-amino-1H-pyrazol-1-yl)phenyl]acetamide as a white solid (3.2 g), which was used without further purification.

EXAMPLE 25


[0248]



[0249] A mixture of Example N (2 g, 0.73 mmol) and 1-naphthylisocyanate (0.124 g, 0.73 mmol) in dry CH_2Cl_2 (4 ml) was stirred at RT under N_2 for 18 h. The solvent was removed in vacuo and the crude product was washed with ethyl acetate (8 ml) and dried in vacuo to yield 1-{3-tert-butyl-1-[3-(3-tert-butyl-5-(1-naphthylureido)-1H-pyrazol-5-yl)-3-(naphthalene-1-yl)urea]phenyl}acetamide as a white solid (0.22 g). mp: 230 (dec.); 1H NMR (200 MHz, $DMSO-d_6$): δ 9.12 (s, 1H), 8.92 (s, 1H), 8.32-8.08 (m, 3H), 7.94-7.44 (m, 8H), 6.44 (s, 1H), 3.51 (s, 2H), 1.31 (s, 9H); MS:

EXAMPLE 26


[0250]



[0251] The title compound was synthesized in a manner analogous to Example 23 utilizing Example N (0.2 g, 0.73 mmol) and 4-chlorophenylisocyanate (0.112 g, 0.73 mmol) to yield 1-{3-tert-butyl-1-[3-(carbamoylmethyl)phenyl]-1H-pyrazol-5-yl}-3-(4-chlorophenyl)urea as a white solid (0.28 g). mp: 222-224. (dec.); ¹H NMR (200 MHz, DMSO-d₆); δ 9.15 (s, 1H), 8.46 (s, 1H), 7.55-7.31 (m, 8H), 6.39 (s, 1H), 3.48 (s, 2H), 1.30 (s, 9H); MS:

EXAMPLE O


[0252]



[0253] The title compound is synthesized in a manner analogous to Example C utilizing Example A and 1-isocyanato-4-methoxy-naphthalene to yield ethyl 3-(3-tert-butyl-5-(3-(1-methoxynaphthalen-4-yl)ureido)-1H-pyrazol-1-yl)benzoate.

EXAMPLE 27


[0254]



[0255] The title compound is synthesized in a manner analogous to Example 17 utilizing Example M and D-4-phenyl-oxazolidin-2-one to yield D-1-{5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl-3-carbonyl)phenyl]-2H-pyrazol-3-yl}-3-phenylurea.

EXAMPLE 28


[0256]



[0257] The title compound is synthesized in a manner analogous to Example 17 utilizing Example M and L-4-phenyl-oxazolidin-2-one to yield L-1-{5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl-3-carbonyl)phenyl]-2H-pyrazol-3-yl}-3-phenylurea.

EXAMPLE P

[0258]



[0259] A mixture of 3-(3-amino-phenyl)-acrylic acid methyl ester (6 g) and 10% Pd on activated carbon (1 g) in ethanol (50 ml) was hydrogenated at 30 psi for 18 h and filtered over Celite. Removal of the volatiles in vacuo provided 3-(3-amino-phenyl)propionic acid methyl ester (6 g).


[0260] A vigorously stirred solution of the crude material from the previous reaction (5.7 g, 31.8 mmol) dissolved in 6 N HCl (35 ml) was cooled to 0° C., and sodium nitrite (2.2 g) in water (20 ml) was added. After 1 h, tin (II) chloride dihydrate (18 g) in 6 N HCl (35 ml) was added. And the mixture was stirred at 0° C. for 3 h. The pH was adjusted to pH 14 with solid KOH and extracted with EtOAc. The combined organic extracts were concentrated in vacuo provided methyl 3-(3-hydrazino-phenyl)propionate (1.7 g).

[0261] A stirred solution of the crude material from the previous reaction (1.7 g, 8.8 mmol) and 4,4-dimethyl-3-oxopentanenitrile (1.2 g, 9.7 mmol) in ethanol (30 ml) and 6 N HCl (2 ml) was refluxed for 18 h and cooled to RT. The volatiles were removed in vacuo and the residue dissolved in EtOAc and washed with 1 N aqueous NaOH. The organic layer was dried (Na₂SO₄) and concentrated in vacuo and the residue was purified by column chromatography using 30% ethyl acetate in hexane as the eluent to provide methyl 3-[3-

(3-tert-butyl-5-amino-1H-pyrazol-1-yl)phenyl]propionate (3.2 g), which was used without further purification

EXAMPLE 29


[0262]



[0263] A mixture of Example P (0.35 g, 1.1 mmol) and 1-naphthylisocyanate (0.19 g, 1.05 mmol) in dry CH_2Cl_2 (5 ml) was stirred at RT under N_2 for 20 h. The solvent was removed in vacuo and the residue was stirred in a solution of THF (3 ml)/MeOH (2 ml)/water (1.5 ml) containing lithium hydroxide (0.1 g) for 3 h at RT, and subsequently diluted with EtOAc and dilute citric acid solution. The organic layer was dried (Na_2SO_4), and the volatiles removed in vacuo. The residue was purified by column chromatography using 3% methanol in CH_2Cl_2 as the eluent to yield 3-(3-{3-tert-butyl-5-(naphthalen-1-yl)ureido}-1H-pyrazol-1-yl)phenylpropionic acid (0.22 g, brownish solid), mp: 105-107; $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 7.87-7.36 (m, 10H), 7.18-7.16 (m, 1H), 6.52 (s, 1H), 2.93 (t, J=6.9 Hz, 2H), 2.65 (t, J=7.1 Hz, 2H), 1.37 (s, 9H); MS

EXAMPLE 30


[0264]



[0265] The title compound was synthesized in a manner analogous to Example 29 utilizing Example P (0.30 g, 0.95 mmol) and 4-chlorophenylisocyanate (0.146 g, 0.95 mmol) to yield 3-(3-{3-tert-butyl-5-(4-chlorophenyl)ureido}-1H-pyrazol-1-yl)phenylpropionic acid (0.05 g, white solid), mp: δ 87; $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 8.21 (s, 1H), 7.44-7.14 (m, 7H), 6.98 (s, 1H), 6.55 (s, 1H), 2.98 (t, J=5.2 Hz, 2H), 2.66 (t, J=5.6 Hz, 2H), 1.40 (s, 9H); MS

EXAMPLE Q

[0266]




A mixture of ethyl 3-(4-aminophenyl)acrylate (1.5 g) and 10% Pd on activated carbon (0.3 g) in ethanol (20 ml) was hydrogenated at 30 psi for 18 h and filtered over Celite. Removal of the volatiles in vacuo provided ethyl 3-(4-aminophenyl)propionate (1.5 g).

[0267] A solution of the crude material from the previous reaction (1.5 g, 8.4 mmol) was dissolved in 6 N HCl (9 ml), cooled to 0°C ., and vigorously stirred. Sodium nitrite (0.58 g) in water (7 ml) was added. After 1 h, tin (II) chloride dihydrate (5 g) in 6 N HCl (10 ml) was added. The reaction mixture was stirred at 0°C . for 3 h. The pH was adjusted to pH 14 with solid KOH and extracted with EtOAc. The combined organic extracts were concentrated in vacuo provided ethyl 3-(4-hydrazino-phenyl)propionate (1 g).

[0268] The crude material from the previous reaction (1 g, 8.8 mmol) and 4,4-dimethyl-3-oxopentenenitrile (0.7 g) in ethanol (8 ml) and 6 N HCl (1 ml) was refluxed for 18 h and cooled to RT. The volatiles were removed in vacuo. The residue was dissolved in ethyl acetate and washed with 1 N aqueous sodium hydroxide solution. The organic layer was dried (Na_2SO_4) and concentrated in vacuo. The residue was purified by column chromatography using 0.7% methanol in CH_2Cl_2 as the eluent to provide ethyl 3-{4-[3-tert-butyl-5-(3-(naphthalen-1-yl)ureido)-1H-pyrazol-1-yl]phenyl}propionate (0.57 g).

EXAMPLE 31

[0269]




[0270] A mixture of Example Q (0.25 g, 0.8 mmol) and 1-naphthylisocyanate (0.13 g, 0.8 mmol) in dry CH_2Cl_2 (5 ml) was stirred at RT under N_2 for 20 h. The solvent was removed in vacuo and the residue was stirred in a solution of THF (3 ml)/MeOH (2 ml)/water (1.5 ml) containing lithium hydroxide (0.1 g) for 3 h at RT and diluted with EtOAc and

diluted citric acid solution. The organic layer was dried (Na_2SO_4), and the volatiles removed in vacuo. The residue was purified by column chromatography using 4% methanol in CH_2Cl_2 as the eluent to yield 3-{4-[3-tert-butyl-5-(3-(naphthalene-1-yl)ureido)-1H-pyrazol-1-yl]phenyl}propanoic acid (0.18 g, off-white solid). mp: 120-122; $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 7.89-7.06 (m, 11H), 6.5 (s, 1H), 2.89 (m, 2H), 2.61 (m, 2H), 1.37 (s, 9H); MS

EXAMPLE 32


[0271]



[0272] The title compound was synthesized in a manner analogous to Example 31 utilizing Example Q (0.16 g, 0.5 mmol) and 4-chlorophenylisocyanate (0.077 g, 0.5 mmol) to yield 3-{4-[3-tert-butyl-5-(3-(4-chlorophenyl)ureido)-1H-pyrazol-1-yl]phenyl}propanoic acid (0.16 g, off-white solid). mp: 112-114; $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 8.16 (s, 1H), 7.56 (s, 1H), 7.21 (s, 2H), 7.09 (s, 2H), 6.42 (s, 1H), 2.80 (m, 2H), 2.56 (m, 2H), 1.32 (s, 9H); MS

EXAMPLE R

[0273]




[0274] A 250 mL pressure vessel (ACE Glass Teflon screw cap) was charged with 3-nitrobiphenyl (20 g, 0.10 mol) dissolved in THF (~100 mL) and 10% Pd/C (3 g). The reaction vessel was charged with H_2 (g) and purged three times. The reaction was charged with 40 psi H_2 (g) and placed on a Parr shaker hydrogenation apparatus and allowed to shake overnight at RT. HPLC showed that the reaction was complete thus the reaction mixture was filtered through a bed of Celite and evaporated to yield the amine: 16.7 g (98% yield)

[0275] In a 250 mL Erlenmeyer flask with a magnetic stir bar, the crude material from the previous reaction (4.40 g, 0.026 mol) was added to 6 N HCl (40 mL) and cooled with an ice bath to 0° C. A solution of NaNO_2 (2.11 g, 0.0306 mol, 1.18 eq.) in water (5 mL) was added drop wise. After 30 min, $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (52.0 g, 0.23 mol, 8.86 eq.) in 6N HCl (100 mL) was added and the reaction mixture was allowed to stir for 3 h, then subsequently transferred to a 500 mL round bottom flask. To this, 4,4-dimethyl-3-oxopentanenitrile (3.25 g, 0.026 mol) and EtOH (100 mL) were added and the mixture refluxed for 4 h, concentrated in vacuo and the residue extracted with EtOAc (2x100 mL). The residue was purified by column chromatograph using hexane/EtOAc/ Et_3N (8:2:0.2) to yield 0.53 g of Example R. $^1\text{H NMR}$ (CDCl_3): δ 7.5 (m, 18H), 5.8 (s, 1H), 1.3 (s, 9H).

EXAMPLE 33


[0276]



[0277] In a dry vial with a magnetic stir bar, Example R (0.145 g; 0.50 mmol) was dissolved in 2 mL CH_2Cl_2 (anhydrous) followed by the addition of phenylisocyanate (0.0544 mL; 0.50 mmol; 1 eq.). The reaction was kept under argon and stirred for 17 h. Evaporation of solvent gave a crystalline mass that was triturated with hexane/EtOAc (4:1) and filtered to yield 1-(3-tert-butyl-1-(3-phenylphenyl)-1H-pyrazol-5-yl)-3-phenylurea (0.185 g, 90%). HPLC purity: 96%; mp: 80-84; $^1\text{H NMR}$ (CDCl_3): δ 7.3 (m, 16H), 6.3 (s, 1H), 1.4 (s, 9H).

EXAMPLE 34


[0278]



[0279] The title compound was synthesized in a manner analogous to Example 33 utilizing Example R (0.145 g; 0.50 mmol) and p-chlorophenylisocyanate (0.0768 g, 0.50 mmol, 1 eq.) to yield 1-(3-tert-butyl-1-(3-phenylphenyl)-1H-pyrazol-5-yl)-3-(4-chlorophenyl)urea (0.205 g, 92%). HPLC purity: 96.5%; mp: 134-136; $^1\text{H NMR}$ (CDCl_3): δ 7.5 (m, 14H), 7.0 (s, 1H), 6.6 (s, 1H), 6.4 (s, 1H), 1.4 (s, 9H).

EXAMPLE S


[0280]



[0281] The title compound is synthesized in a manner analogous to Example C utilizing Example A and 4-fluorophenyl isocyanate yield ethyl 3-(3-tert-butyl-5-(3-(4-fluorophenyl)ureido)-1H-pyrazol-1-yl)benzoate.

EXAMPLE 35


[0282]



[0283] The title compound is synthesized in a manner analogous to Example 17 utilizing Example M and D-4-phenyl-oxazolidin-2-one to yield D-1-({5-tert-butyl-2-[3-(2-oxo-4-phenyl-oxazolidinyl)-3-carbonyl]phenyl}-2H-pyrazol-3-yl)-3-(naphthalen-1-yl)urea.

EXAMPLE 36


[0284]



[0285] The title compound is synthesized in a manner analogous to Example 29 utilizing Example P (0.30 g, 0.95 mmol) and 4-fluorophenylisocyanate (0.146 g, 0.95 mmol) to yield 3-(3-(3-tert-butyl-5-(3-(4-fluorophenyl)ureido)-1H-pyrazol-1-yl)phenyl)propanoic acid.

EXAMPLE T

[0286]




[0287] To a stirred solution of Example N (2 g, 7.35 mmol) in THF (6 ml) was added borane-methylsulfide (18 mmol). The mixture was heated to reflux for 90 min and cooled to RT, after which 6 N HCl was added and heated to reflux for 10 min. The mixture was basified with NaOH and extracted with EtOAc. The organic layer was dried (Na_2SO_4) filtered and concentrated in vacuo to yield 3-tert-butyl-1-[3-(2-aminoethyl)phenyl]-1H-pyrazol-5-amine (0.9 g).

[0288] A mixture of the crude material from the previous reaction (0.8 g, 3.1 mmol) and di-tert-butylcarbonate (0.7 g, 3.5 mmol) and catalytic amount of DMAP in dry CH_2Cl_2 (5 ml) was stirred at RT under N_2 for 18 h. The reaction mixture was concentrated in vacuo and the residue was purified by column chromatography using 1% methanol in CH_2Cl_2 as the eluent to yield tert-butyl 3-(3-tert-butyl-5-amino-1H-pyrazol-1-yl)phenylcarbamate (0.5 g).

EXAMPLE 37


[0289]



[0290] A mixture of Example T (0.26 g, 0.73 mmol) and 1-naphthylisocyanate (0.123 g, 0.73 mmol) in dry CH_2Cl_2 (5 mL) was stirred at RT under N_2 for 48 h. The solvent was removed in vacuo and the residue was purified by column chromatography using 1% methanol in CH_2Cl_2 as the eluent (0.15 g, off-white solid). The solid was then treated with TFA (0.2 mL) for 5 min and diluted with EtOAc. The organic layer was washed with saturated NaHCO_3 solution and brine, dried (Na_2SO_4), filtered and concentrated in vacuo to yield 1-[3-tert-butyl-1-[3-(2-aminoethyl)phenyl]-1H-pyrazol-5-yl]-3-(naphthalen-1-yl)urea as a solid (80 mg). mp: 110-112; ^1H NMR (200 MHz, DMSO-d_6): δ 9.09 (s, 1H), 8.90 (s, 1H), 8.01-7.34 (m, 11H), 6.43 (s, 1H), 3.11 (m, 2H), 2.96 (m, 2H), 1.29 (s, 9H); MS

EXAMPLE 38


[0291]



[0292] The title compound was synthesized in a manner analogous to Example 37 utilizing Example T (0.15 g, 0.42 mmol) and 4-chlorophenylisocyanate (0.065 g, 0.42 mmol) to yield 1-[3-tert-butyl-1-[3-(2-aminoethyl)phenyl]-1H-pyrazol-5-yl]-3-(4-chlorophenyl)urea as an off-white solid (20 mg). mp: 125-127; ^1H NMR (200 MHz, CDCl_3): δ 8.81 (s, 1H), 8.66 (s, 1H), 7.36-7.13 (m, 8H), 6.54 (s, 1H), 3.15 (brs, 2H), 2.97 (brs, 2H), 1.32 (s, 9H); MS

EXAMPLE U

[0293]




[0294] In a 250 mL Erlenmeyer flask with a magnetic stir bar, m-anisidine (9.84 g, 0.052 mol) was added to 6 N HCl (80 mL) and cooled with an ice bath to 0°C . A solution of NaNO_2 (4.22 g, 0.0612 mol, 1.18 eq.) in water (10 mL) was added drop wise. After 30 min, $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (104.0 g, 0.46 mol, 8.86 eq.) in 6 N HCl (200 mL) was added and the reaction mixture was allowed to stir for 3 h., and then subsequently transferred to a 1000 mL round bottom flask. To this, 4,4-dimethyl-3-oxopentenenitrile (8.00 g, 0.064 mol) and EtOH (200 mL) were added and the mixture refluxed for 4 h, concentrated in vacuo and the residue recrystallized from CH_2Cl_2 to yield 3-tert-butyl-1-(3-methoxyphenyl)-1H-pyrazol-5-amine as the HCl salt (13.9 g).

[0295] The crude material from the previous reaction (4.65 g, 0.165 mol) was dissolved in 30 mL of CH_2Cl_2 with Et_3N (2.30 mL, 0.0165 mol, 1 eq.) and stirred for 30 min. Extraction with water followed by drying of the organic phase with Na_2SO_4 and concentration in vacuo yielded a brown syrup that was the free base, 3-tert-butyl-1-(3-methoxyphenyl)-1H-pyrazol-5-amine (3.82 g, 94.5%), which was used without further purification.

EXAMPLE 39


[0296]



[0297] In a dry vial with a magnetic stir bar, Example U (2.62 g, 0.0107 mol) was dissolved in CH_2Cl_2 (5 mL, anhydrous) followed by the addition of 1-naphthylisocyanate (1.53 mL, 0.0107 mol, 1 eq.). The reaction was kept under Ar and stirred for 18 h. Evaporation of solvent followed by column chromatography with EtOAc/hexane/ Et_3N (7:2:0.5) as the eluent yielded 1-[3-tert-butyl-1-(3-methoxyphenyl)-1H-pyrazol-5-yl]-3-(naphthalen-1-yl)urea (3.4 g, 77%). HPLC: 97%; mp: 78-80; ^1H NMR (CDCl_3): δ 7.9-6.8 (m, 15H), 6.4 (s, 1H), 3.7 (s, 3H), 1.4 (s, 9H).

EXAMPLE 40


[0298]



[0299] The title compound was synthesized in a manner analogous to Example 39 utilizing Example U (3.82 g; 0.0156 mol) and p-chlorophenylisocyanate (2.39 g, 0.0156 mol, 1 eq.), purified by trituration with hexane/EtOAc (4:1) and filtered to yield 1-[3-tert-butyl-1-(3-methoxyphenyl)-1H-pyrazol-5-yl]-3-(4-chlorophenyl)urea (6.1 g, 98%). HPLC purity: 95%; mp: 158-160; $^1\text{H NMR}$ (CDCl_3): δ 7.7 (s, 1H); δ 7.2-6.8 (m, 8H), 6.4 (s, 1H), 3.7 (s, 3H), 1.3 (s, 9H).

EXAMPLE 41


[0300]



[0301] In a 100 ml round bottom flask equipped with a magnetic stir bar, Example 39 (2.07 g) was dissolved in CH_2Cl_2 (20 mL) and cooled to 0°C . with an ice bath. BBr_3 (1 M in CH_2Cl_2 ; 7.5 mL) was added slowly. The reaction mixture was allowed to warm to RT overnight. Additional BBr_3 (1 M in CH_2Cl_2 , 2x1 mL, 9.5 mmol total added) was added and the reaction was quenched by the addition of MeOH. Evaporation of solvent led to a crystalline material that was chromatographed on silica gel (30 g) using $\text{CH}_2\text{Cl}_2/\text{MeOH}$ (9.6:0.4) as the eluent to yield 1-[3-tert-butyl-1-(3-hydroxyphenyl)-1H-pyrazol-5-yl]-3-(naphthalene-1-yl)urea (0.40 g, 20%). $^1\text{H NMR}$ ($\text{DMSO}-d_6$): δ 9.0 (s, 1H), 8.8 (s, 1H), 8.1-6.8 (m, 11H), 6.4 (s, 1H), 1.3 (s, 9H). MS (ESI) m/z: 401 ($\text{M}+\text{H}^+$).

EXAMPLE 42


[0302]



[0303] The title compound was synthesized in a manner analogous to Example 41 utilizing Example 40 (2.00 g, 5 mmol) that resulted in a crystalline material that was filtered and washed with MeOH to yield 1-[3-tert-butyl-1-(3-hydroxyphenyl)-1H-pyrazol-5-yl]-3-(4-chlorophenyl)urea (1.14 g, 60%). HPLC purity: 96%; mp: 214-216; $^1\text{H NMR}$ (CDCl_3): δ 8.4 (s, 1H), 7.7 (s, 1H), 7.4-6.6 (m, 9H), 1.3 (s, 9H).

EXAMPLE V

[0304]



[0305] The starting material, 1-[4-(aminomethyl)phenyl]-3-tert-butyl-N-nitroso-1H-pyrazol-5-amine, was synthesized in a manner analogous to Example A utilizing 4-aminobenzamide and 4,4-dimethyl-3-oxopentenenitrile.


[0306] A 1 L four-necked round bottom flask was equipped with a stir bar, a source of dry Ar, a heating mantle, and a reflux condenser. The flask was flushed with Ar and charged with the crude material from the previous reaction (12 g, 46.5 mmol; 258.1 g/mol) and anhydrous THF (500 ml). This solution was treated cautiously with LiAlH_4 (2.65 g, 69.8 mmol) and the reaction was stirred overnight. The reaction was heated to reflux and additional LiAlH_4 was added complete (a total of 8.35 g added). The reaction was cooled to 0 and H_2O (8.4 ml), 15% NaOH (8.4 ml) and H_2O (24 ml) were added sequentially; The mixture was stirred for 2 h, the solids filtered through Celite, and washed extensively with THF, the solution was concentrated in vacuo to yield 1-(4-(aminomethyl)-3-methoxyphenyl)-3-tert-butyl-1H-pyrazol-5-amine (6.8 g) as an oil.

[0307] A 40 mL vial was equipped with a stir bar, a septum, and a source of Ar. The vial was charged with the crude material from the previous reaction (2 g, 8.2 mmol, 244.17

g/mol) and CHCl_3 (15 mL) were cooled to 0 under Ar and di-tert-butylcarbonate (1.9 g, 9.0 mmol) dissolved in CHCl_3 (5 mL) was added drop wise over a 2 min period. The mixture was treated with 1N KOH (2 mL), added over a 2 h period. The resulting emulsion was broken with the addition of saturated NaCl solution, the layers were separated and the aqueous phase extracted with CH_2Cl_2 (2x1.5 ml). The combined organic phases were dried over Na_2SO_4 , filtered, concentrated in vacuo to yield tert-butyl [4-(3-tert-butyl-5-amino-1H-pyrazol-1-yl)-2-methoxybenzyl]carbamate (2.23 g, 79%) as a light yellow solid. $^1\text{H NMR}$ (CDCl_3): δ 7.4 (m, 5H), 5.6 (s, 1H), 4.4 (d, 2H), 1.5 (s, 9H), 1.3 (s, 9H).

EXAMPLE 43


[0308]



[0309] A 40 mL vial was equipped with a septum, a stir bar and a source of Ar, and charged with Example V (2 g, 5.81 mmol), flushed with Ar and dissolved in CHCl_3 (20 mL). The solution was treated with 2-naphthylisocyanate (984 mg, 5.81 mmol) in CHCl_3 (5 mL) and added over 1 min. The reaction was stirred for 8 h, and additional 1-naphthylisocyanate (81 mg) was added and the reaction stirred overnight. The solid was filtered and washed with CH_2Cl_2 to yield tert-butyl 4-[3-tert-butyl-5-(3-naphthalen-1-yl)ureido]-1H-pyrazol-1-yl]benzyl]carbamate (1.2 g). HPLC purity: 94.4%; $^1\text{H NMR}$ (DMSO-d_6): δ 9.1 (s, 1H), 8.8 (s, 1H), 8.0 (m, 3H), 7.6 (m, 9H), 6.4 (s, 1H), 4.2 (d, 2H), 1.4 (s, 9H), 1.3 (s, 9H).

EXAMPLE 44


[0310]



[0311] The title compound was synthesized in a manner analogous to Example 43 utilizing Example V (2.0 g, 5.81 mmol) and p-chlorophenylisocyanate (892 mg) to yield tert-butyl 4-[3-tert-butyl-5-(3-(4-chlorophenyl)ureido)-1H-pyrazol-1-yl]benzyl]carbamate (1.5 g). HPLC purity: 97%; $^1\text{H NMR}$ (DMSO-d_6): δ 9.2 (s, 1H), 8.4 (s, 1H), 7.4 (m, 8H), 6.4 (s, 1H), 4.2 (d, 2H), 1.4 (s, 9H), 1.3 (s, 9H).

EXAMPLE 45


[0312]



[0313] A 10 mL flask equipped with a stir bar was flushed with Ar and charged with Example 43 (770 mg, 1.5 mmol) and CH_2Cl_2 (1 ml) and 1:1 CH_2Cl_2 :TFA (2.5 mL). After 1.5 h, reaction mixture was concentrated in vacuo, the residue was dissolved in EtOAc (15 mL), washed with saturated NaHCO_3 (10 mL) and saturated NaCl (10 mL). The organic layers were dried, filtered and concentrated in vacuo to yield 1-[3-tert-butyl-1-[4-(aminomethyl)phenyl]-1H-pyrazol-5-yl]-3-(naphthalen-1-yl)urea (710 mg). $^1\text{H NMR}$ (DMSO-d_6): δ 7.4 (m, 1H), 6.4 (s, 1H), 3.7 (s, 2H), 1.3 (s, 9H).

EXAMPLE 46


[0314]



[0315] The title compound was synthesized in a manner analogous to Example 45 utilizing Example 44 (1.5 g, 1.5 mmol) to yield 1-{3-tert-butyl-1-[4-(aminomethyl)phenyl]-1H-pyrazol-5-yl}-3-(4-chlorophenyl)urea (1.0 g). HPLC purity: 93.6%; mp: 100-102; ¹H NMR (CDCl₃): δ 8.6 (s, 1H), 7.3 (m, 8H), 6.3 (s, 1H), 3.7 (brs, 2H), 1.3 (s, 9H).

EXAMPLE 47


[0316]



[0317] A 10 ml vial was charged with Example 45 (260 mg, 63 mmol) and absolute EtOH (3 mL) under Ar. Divinylsulfone (63 uL, 74 mg, 0.63 mmol) was added drop wise over 3 min and the reaction was stirred at RT for 1.5 h. and concentrated in vacuo to yield a yellow solid, which was purified via preparative TLC, developed in 5% MeOH:CH₂Cl₂. The predominant band was cut and eluted off the silica with 1:1 EtOAc:MeOH, filtered and concentrated in vacuo to yield 1-{3-tert-butyl-1-[4-(1,1-dioxothiomorpholin-4-yl)methylphenyl]-1H-pyrazol-5-yl}-3-(naphthalen-1-yl)urea (150 mg). HPLC purity: 96%; ¹H NMR (DMSO-d₆): δ 9.1 (s, 1H), 9.0 (s, 1H), 7.9 (m, 3H), 7.5 (m, 8H), 6.4 (s, 1H), 3.1 (brs, 4H), 2.9 (brs, 4H), 1.3 (s, 9H).

EXAMPLE 48


[0318]



[0319] The title compound was synthesized in a manner analogous to Example 47 utilizing Example 46 (260 mg, 0.66 mmol) to yield 1-{3-tert-butyl-1-[4-(1,1-dioxothiomorpholin-4-yl)methylphenyl]-1H-pyrazol-5-yl}-3-(4-chlorophenyl)urea (180 mg). HPLC purity: 93%; mp: 136-138; ¹H NMR (DMSO-d₆): δ 9.2 (s, 1H), 8.5 (s, 1H), 7.4 (m, 9H), 6.4 (s, 1H), 3.1 (brs, 4H), 3.0 (brs, 4H), 1.3 (s, 9H).

EXAMPLE 49


[0320]



[0321] To a stirring solution of chlorosulfonyl isocyanate (0.35 g, 5 mmol) in CH₂Cl₂ (20 mL) at 0° C. was added pyrrolidine (0.18 g, 5 mmol) at such a rate that the reaction temperature did not rise above 5° C. After stirring for 2 h, a solution of Example 41 (1.10 g, 6.5 mmol) and triethylamine (0.46 g, 9 mmol) in CH₂Cl₂ (20 mL) was added. When the addition was complete, the mixture was allowed to warm to RT and stirred overnight. The reaction mixture was poured into 10% HCl (10 mL) saturated with NaCl, the organic layer was separated and the aqueous layer extracted with ether (20 mL). The combined organic layers were dried (Na₂SO₄) and concentrated in vacuo, purified by preparative HPLC to yield (pyrrolidine-1-carbonyl)sulfamic acid 3-[3-tert-butyl-5-(3-naphthalen-1-yl-ureido)-pyrazol-1-yl]phenyl ester (40 mg). ¹H NMR (CDCl₃): δ 9.12 (brs, 1H), 8.61 (brs, 1H), 7.85-7.80 (m, 3H), 7.65 (d, J=8.0 Hz, 2H), 7.53-7.51 (m, 1H), 7.45-7.25 (m, 5H), 6.89 (s, 4H), 3.36-3.34 (brs, 1H), 3.14-3.13 (brs, 2H), 1.69 (brs, 2H), 1.62 (brs, 2H), 1.39 (s, 9H); MS (ESI) m/z: 577 (M+H⁺).

EXAMPLE 50

[0322]




[0323] The title compound was synthesized in a manner analogous to Example 49 utilizing Example 42 to yield (pyr-

olidine-1-carbonyl)sulfamic acid 3-[3-tert-butyl-5-(4-chlorophenyl-1-yl-ureido)pyrazol-1-yl]phenyl ester. MS (ESI) m/z: 561 (M+H⁺).

EXAMPLE W


[0324]



[0325] Solid 4-methoxyphenylhydrazine hydrochloride (25.3 g) was suspended in toluene (100 mL) and treated with triethylamine (20.2 g). The mixture was stirred at RT for 30 min and treated with pivaloylacetone (18 g). The reaction was heated to reflux and stirred overnight. The hot mixture was filtered, the solids washed with hexane and dried in vacuo to afford 3-tert-butyl-1-(4-methoxyphenyl)-1H-pyrazol-5-amine (25 g, 70%). ¹H NMR (DMSO-d₆): δ 7.5 (d, 2H), 7.0 (d, 1H), 6.4 (s, 1H), 6.1 (s, 2H), 3.9 (s, 3H), 1.3 (s, 9H).

EXAMPLE 51


[0326]



[0327] To a solution of 1-isocyanato-4-methoxy-naphthalene (996 mg) in anhydrous CH₂Cl₂ (20 mL) of was added Example W (1.23 g). The reaction solution was stirred for 3 h, the resulting white precipitate filtered, treated with 10% HCl and recrystallized from MeOH, and dried in vacuo to yield 1-[3-tert-butyl-1-(4-methoxyphenyl)-1H-pyrazol-5-yl]-3-(1-methoxynaphthalen-4-yl)-urea as white crystals (900 mg, 40%). HPLC purity: 96%; mp: 143-144; ¹H NMR (DMSO-d₆): δ 8.8 (s, 1H), 8.5 (s, 1H), 8.2 (d, 1H), 8.0 (d, 1H), 7.6 (m, 5H), 7.1 (d, 2H), 7.0 (d, 1H), 6.3 (s, 1H), 4.0 (s, 3H), 3.9 (s, 3H); 1.3 (s, 9H).

EXAMPLE 52


[0328]



[0329] The title compound was synthesized in a manner analogous to Example 51 utilizing Example W and p-bromophenylisocyanate (990 mg) to yield 1-[3-tert-butyl-1-(4-methoxyphenyl)-1H-pyrazol-5-yl]-3-(4-bromophenyl)urea as off-white crystals (1.5 g, 68%). HPLC purity: 98%; mp: 200-201; ¹H NMR (DMSO-d₆): δ 9.3 (s, 1H), 8.3 (s, 1H), 7.4 (m, 6H), 7.0 (d, 2H), 6.3 (s, 1H), 3.8 (s, 3H), 1.3 (s, 9H).

EXAMPLE 53


[0330]



[0331] The title compound was synthesized in a manner analogous to Example 51 utilizing Example W and p-chlorophenylisocyanate (768 mg) into yield 1-[3-tert-butyl-1-(4-methoxyphenyl)-1H-pyrazol-5-yl]-3-(4-chlorophenyl)urea as white crystals (1.3 g, 65%). HPLC purity: 98%; mp: 209-210; ¹H NMR (DMSO-d₆): δ 9.1 (s, 1H), 8.3 (s, 1H), 7.4 (m, 4H), 7.3 (d, 2H), 7.1 (d, 2H), 6.3 (s, 1H), 3.8 (s, 3H), 1.3 (s, 9H).

EXAMPLE 54


[0332]



[0333] The title compound was synthesized in a manner analogous to Example 41 utilizing Example 53 (500 mg) to yield 1-{3-tert-butyl-1-(4-hydroxyphenyl)-1H-pyrazol-5-yl}-3-(4-chlorophenyl)urea as white crystals (300 mg, 62%). HPLC purity: 94%; mp: 144-145; ¹H NMR (DMSO-d₆): δ 9.7 (s, 1H), 9.1 (s, 1H), 8.3 (s, 1H), 7.4 (d, 2H), 7.3 (m, 4H); 6.9 (d, 2H), 6.3 (s, 1H), 1.3 (s, 9H)

EXAMPLE 55


[0334]



[0335] The title compound was synthesized in a manner analogous to Example 41 utilizing Example 52 (550 mg) to yield 1-{3-tert-butyl-1-(4-hydroxyphenyl)-1H-pyrazol-5-yl}-3-(4-bromophenyl)urea as a white crystalline solid (400 mg, 70%). HPLC purity: 93%; mp: 198-200; ¹H NMR (DMSO-d₆): δ 9.7 (s, 1H), 9.2 (s, 1H), 8.3 (s, 1H), 7.4 (d, 4H), 7.2 (m, 2H), 6.9 (d, 2H), 6.3 (s, 1H), 1.3 (s, 9H).

EXAMPLE X


[0336]



[0337] Methyl 4-(3-tert-butyl-5-amino-1H-pyrazol-1-yl)benzoate (3.67 mmol) was prepared from methyl 4-hydrazinobenzoate and pivaloylacetonitrile by the procedure of Regan, et al., *J. Med. Chem.*, 45, 2994 (2002).

EXAMPLE 56


[0338]



[0339] A 500 mL round bottom flask was equipped with a magnetic stir bar and an ice bath. The flask was charged with Example X (1 g) and this was dissolved in CH₂Cl₂ (100 mL). Saturated sodium bicarbonate (100 mL) was added and the mixture rapidly stirred, cooled in an ice bath and treated with diphosgene (1.45 g) and the heterogeneous mixture stirred for 1 h. The layers were separated and the CH₂Cl₂ layer treated with tert-butanol (1.07 g) and the solution stirred overnight at RT. The solution was washed with H₂O (2×150 mL), dried (Na₂SO₄), filtered, concentrated in vacuo, and purified by flash chromatography using 1:2 ethyl acetate:hexane as the eluent to yield tert-butyl-1-(4-(methoxycarbonyl)phenyl)-3-tert-butyl-1H-pyrazol-5-ylcarbamate (100 mg) as an off-white solid. ¹H NMR (DMSO-d₆): δ 9.2 (s, 1H), 8.1 (d, 2H), 7.7 (d, 2H), 6.3 (s, 1H), 3.3 (s, 3H), 1.3 (s, 18H).

EXAMPLE 57


[0340]



[0341] The title compound was synthesized in a manner analogous to Example 41 utilizing Example X (1.37 g) and p-chlorophenylisocyanate (768 mg) to yield methyl 4-{3-tert-butyl-5-[3-(4-chlorophenyl)ureido]-1H-pyrazol-1-yl}benzoate as white crystals (1.4 g 66%). HPLC purity: 98%; mp: 160-161; ¹H NMR (DMSO-d₆): δ 9.2 (s, 1H), 8.6 (s, 1H), 8.1 (d, 2H), 7.8 (d, 2H), 7.5 (d, 2H), 7.3 (d, 2H), 6.4 (s, 1H), 3.9 (s, 3H), 1.3 (s, 9H).

EXAMPLE 58


[0342]



[0343] The title compound was synthesized in a manner analogous to Example 41 utilizing Example X (1.27 g) and 1-isocyanato-4-methoxy-naphthalene (996 mg) to yield methyl 4-{3-tert-butyl-5-[3-(1-methoxynaphthalen-4-yl)ureido]-1H-pyrazol-1-yl}benzoate as white crystals (845 mg, 36%). HPLC purity: 98%; mp: 278-280; $^1\text{H NMR}$ (DMSO- d_6): δ 8.76 (s, 1H), 8.73 (s, 1H), 8.1 (m, 3H), 7.9 (d, 1H), 7.7 (d, 2H), 7.6 (m, 3H), 7.0 (d, 1H), 7.0 (d, 1H), 6.3 (s, 1H), 4.0 (s, 3H), 3.9 (s, 3H), 1.3 (s, 9H).

EXAMPLE 59


[0344]



[0345] The title compound was synthesized in a manner analogous to Example 41 utilizing Example X (1.37 g) and p-bromophenyliisocyanate (990 mg) to yield methyl 4-{3-tert-butyl-5-[3-(4-bromophenyl)ureido]-1H-pyrazol-1-yl}benzoate as white crystals (1.4 g, 59%). HPLC purity: 94%; mp: 270-272; $^1\text{H NMR}$ (DMSO- d_6): δ 9.2 (s, 1H), 8.6 (s, 1H), 8.1 (d, 2H), 7.7 (d, 2H), 7.4 (d, 4H), 6.4 (s, 1H), 3.9 (s, 3H), 1.3 (s, 9H).


EXAMPLE 60

[0346]




[0347] To a solution of Example 59 (700 mg) in 30 mL of toluene at -78°C ., was added dropwise a solution of diisobutylaluminum hydride in toluene (1M in toluene, 7.5 mL) over 10 min. The reaction mixture was stirred for 30 min at -78°C ., and then 30 min at 0°C . The reaction mixture was concentrated in vacuo to dryness and treated with H_2O . The solid was filtered and treated with acetonitrile. The solution was evaporated to dryness and the residue was dissolved in ethyl acetate, and precipitated by hexanes to afford yellow solid which was dried under vacuum to give 1-[3-tert-butyl-1-(4-hydroxymethylphenyl)pyrazol-5-yl]urea (400 mg, 61%). HPLC purity: 95%; $^1\text{H NMR}$ (DMSO- d_6): δ 9.2 (s, 1H), 8.4 (s, 1H), 7.5 (m, 8H), 6.4 (s, 1H), 5.3 (t, 1H), 4.6 (d, 2H), 1.3 (s, 9H).


Example 1



Example 2



-continued



Example 3















Chart 1








Example 4



Example 5















Example 6












Wherein Y is O, S, NR₆, —NR₆SO₂—, NR₆CO—, alkylene, O—(CH₂)_n—, NR₆—(CH₂)_n—, wherein one of the methylene units may be substituted with an oxo group, or Y is a direct bond; D is taken from the groups identified in Chart 1:


-continued




-continued




-continued




Example 7




Example 10




Example 8



Example 11




Example 9




Example 12

-continued


-continued




Example 13




Example 16




Example 14



Example 17




Example 15




Example 18

-continued


-continued




Example 19




Example 22




Example 20



Example 23




Example 21




Example 24

-continued


-continued




Example 25




Example 28




Example 26



Example 29




Example 27




Example 30

-continued


-continued




Example 31




Example 34




Example 32



Example 35




Example 33




Example 36

-continued




Example 37


-continued




Example 40




Example 38



Example 41




Example 39




Example 42

-continued




Example 43


-continued




Example 46




Example 44



Example 47




Example 45




Example 48


-continued




Example 49



Example 50




Example 51




Example 52


-continued




Example 53



Example 54




Example 55




Example 56


-continued



Example 57




Example 58




Example 59


-continued




Example 60



Example 61




Example 62




Example 63

-continued




Example 64


-continued




Example 68




Example 65




Example 69




Example 66



Example 70




Example 67



Example 71


-continued

Example 72




-continued


Example 76




Example 73




Example 77




Example 74



Example 78



Example 75



wherein X or Y is O, S, NR₆, —NR₆SO₂—, NR₆CO—, alkylene, O—(CH₂)_n—, NR₆—(CH₂)_n—, wherein one of the methylene units may be substituted with an oxo group, or X or Y is a direct bond; D is taken from the groups identified in Chart I:







Chart I


-continued




D-3




D-4




D-5




D-6




D-7




D-8




D-9




D-10




D-11




D-12




D-13



D-14




D-15




D-16


-continued




D-17




D-18




D-19




D-20




D-21




D-22




D-23




D-24




D-25



D-26




D-27




D-28


-continued




D-29




D-30




D-31




D-32




D-33



D-34




D-35




D-36


Specific examples of the present invention are illustrated by their structural formulae below:



Example 79




Example 80



Example 81


-continued

Example 82




-continued


Example 85




Example 83




Example 86



Example 84




Example 87




-continued

-continued


Example 88




Example 91




Example 89




Example 92



Example 90




Example 93




-continued

-continued


Example 94




Example 97




Example 95




Example 98



Example 96




Example 99




-continued

Example 100




-continued


Example 103




Example 101




Example 104



Example 102




Example 105




-continued

-continued


Example 106




Example 109




Example 107




Example 110



Example 108




Example 111




-continued

-continued


Example 112




Example 115




Example 113




Example 116



Example 114




Example 117




-continued

-continued


Example 118




Example 121




Example 119




Example 122



Example 120




Example 123




-continued

-continued


Example 124




Example 127




Example 125




Example 128



Example 126




Example 129




-continued


Example 130



Example 131




Example 132




-continued


Example 133




Example 134



Example 135




Example 136




-continued


Example 137




Example 138



Example 139




Example 140




-continued


Example 141




Example 142



Example 143



Example 144



[0348] All of the references above identified are incorporated by reference herein. In addition, two simultaneously applications are also incorporated by reference, namely

Modulation of Protein Functionalities, Ser. No. _____, filed December _____, 2003, and Anti-Cancer Medicaments, Ser. No. _____ filed December _____, 2003.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 6

<210> SEQ ID NO 1

<211> LENGTH: 360

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Ser Gln Glu Arg Pro Thr Phe Tyr Arg Gln Glu Leu Asn Lys Thr
 1          5          10          15

Ile Trp Glu Val Pro Glu Arg Tyr Gln Asn Leu Ser Pro Val Gly Ser
 20          25          30

Gly Ala Tyr Gly Ser Val Cys Ala Ala Phe Asp Thr Lys Thr Gly Leu
 35          40          45

Arg Val Ala Val Lys Lys Leu Ser Arg Pro Phe Gln Ser Ile Ile His
 50          55          60

Ala Lys Arg Thr Tyr Arg Glu Leu Arg Leu Leu Lys His Met Lys His
 65          70          75          80

Glu Asn Val Ile Gly Leu Leu Asp Val Phe Thr Pro Ala Arg Ser Leu
 85          90          95

Glu Glu Phe Asn Asp Val Tyr Leu Val Thr His Leu Met Gly Ala Asp
100          105          110

Leu Asn Asn Ile Val Lys Cys Gln Lys Leu Thr Asp Asp His Val Gln
115          120          125

Phe Leu Ile Tyr Gln Ile Leu Arg Gly Leu Lys Tyr Ile His Ser Ala
130          135          140

Asp Ile Ile His Arg Asp Leu Lys Pro Ser Asn Leu Ala Val Asn Glu
145          150          155          160

Asp Cys Glu Leu Lys Ile Leu Asp Phe Gly Leu Ala Arg His Thr Asp
165          170          175

Asp Glu Met Thr Gly Tyr Val Ala Thr Arg Trp Tyr Arg Ala Pro Glu
180          185          190

Ile Met Leu Asn Trp Met His Tyr Asn Gln Thr Val Asp Ile Trp Ser
195          200          205

Val Gly Cys Ile Met Ala Glu Leu Leu Thr Gly Arg Thr Leu Phe Pro
210          215          220

Gly Thr Asp His Ile Asp Gln Leu Lys Leu Ile Leu Arg Leu Val Gly
225          230          235          240

Thr Pro Gly Ala Glu Leu Leu Lys Lys Ile Ser Ser Glu Ser Ala Arg
245          250          255

Asn Tyr Ile Gln Ser Leu Thr Gln Met Pro Lys Met Asn Phe Ala Asn
260          265          270

Val Phe Ile Gly Ala Asn Pro Leu Ala Val Asp Leu Leu Glu Lys Met
275          280          285

Leu Val Leu Asp Ser Asp Lys Arg Ile Thr Ala Ala Gln Ala Leu Ala
290          295          300

His Ala Tyr Phe Ala Gln Tyr His Asp Pro Asp Asp Glu Pro Val Ala
305          310          315          320

```

-continued

Asp Pro Tyr Asp Gln Ser Phe Glu Ser Arg Asp Leu Leu Ile Asp Glu
 325 330 335

Trp Lys Ser Leu Thr Tyr Asp Glu Val Ile Ser Phe Val Pro Pro Pro
 340 345 350

Leu Asp Gln Glu Glu Met Glu Ser
 355 360

<210> SEQ ID NO 2
 <211> LENGTH: 17
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (2)..(16)
 <223> OTHER INFORMATION: X is any amino acid

<400> SEQUENCE: 2

Ile Ile Xaa Xaa Lys Arg Xaa Xaa Arg Glu Xaa Xaa Leu Leu Xaa Xaa
 1 5 10 15

Met

<210> SEQ ID NO 3
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Asp Ile Ile His Arg Asp
 1 5

<210> SEQ ID NO 4
 <211> LENGTH: 10
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Asp Phe Gly Leu Ala Arg His Thr Asp Asp
 1 5 10

<210> SEQ ID NO 5
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Glu Met Thr Gly Tyr Val Ala Thr Arg Trp Tyr Arg
 1 5 10


<210> SEQ ID NO 6
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Trp Met His Tyr
 1


1-31. (canceled)

32. A method of modulating the activation state of a p38 α -kinase comprising the step of contacting said kinase with a molecule having the formula



wherein:

R_1 is selected from the group consisting of



each R_2 is individually selected from the group consisting of —H, alkyls, aminos, alkylaminos, arylaminos, cycloalkylaminos, heterocyclylaminos, halogens, alkoxy, and hydroxys;


each R_3 is individually selected from the group consisting of —H, alkyls, alkylaminos, arylaminos,

cycloalkylaminos, heterocyclylaminos, alkoxy, hydroxys, cyanos, halogens, perfluoroalkyls, alkylsulfinyls, alkylsulfonyls, R_4NHSO_2 —, and $-NHSO_2R_4$;

each R_5 is individually selected from the group consisting of —H, alkyls, aryls, heterocyclyls, alkylaminos, arylaminos, cycloalkylaminos, heterocyclylaminos, hydroxys, alkoxy, aryloxy, alkylthios, arylthios, cyanos, halogens, perfluoroalkyls, alkylcarbonyls, and nitros; and each W is individually selected from the group consisting of CH and N;

each X is individually selected from the group consisting of —O—, —S—, —NR₆—, —NR₆SO₂—, —NR₆CO—, alkynyls, alkenyls, alkylenes, —O(CH₂)_h—, and —NR₆(CH₂)_h—, where each h is individually selected from the group consisting of 1, 2, 3, or 4, and where for each of alkylenes, —O(CH₂)_h—, and —NR₆(CH₂)_h—, one of the methylene groups present therein may be optionally double-bonded to a side-chain oxo group except that where —O(CH₂)_h— the introduction of the side-chain oxo group does not form an ester moiety;

A is selected from the group consisting of phenyl, naphthyl, pyridyl, pyrimidyl, thienyl, furyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, indolyl, indazolyl, benzimidazolyl, benzotriazolyl, isoquinolyl, quinolyl, benzothiazolyl, benzofuranyl, benzothienyl, pyrazolylpyrimidinyl, imidazopyrimidinyl, purinyl, and



where each W_1 is individually selected from the group consisting of —CH— and —N—;


cycloalkylaminos, heterocyclylaminos, hydroxys, alkoxy, aryloxy, alkylthios, arylthios, cyanos, halogens, nitrilos, nitros, alkylsulfinyls, alkylsulfonyls, aminosulfonyls, and perfluoroalkyls.

33. The method of claim 32, wherein m is 0; wherein D is substituted by a first R_7 substituent.

34. The method of claim 33, wherein A is phenyl substituted by one or more second R_7 substituents, said second R_7 substituents being halogens.


35. The method of claim 33, wherein A is an R_7 -substituted naphthyl.

36. The method of claim 32, wherein m is 1; R_1 is taken from the group consisting of



37. The method of claim 36, wherein A is taken from phenyl or naphthyl.


38. The method of claim 32 having the formula IB



39. A method of claim 32 having the formula IC

IB

IC



* * * * *