

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2016377681 B2

(54) Title
Compounds and compositions for intracellular delivery of agents

(51) International Patent Classification(s)

C07D 211/14 (2006.01)	A61K 31/495 (2006.01)
A61K 9/14 (2006.01)	C07D 211/16 (2006.01)
A61K 31/445 (2006.01)	C07D 295/13 (2006.01)
A61K 31/45 (2006.01)	C07D 295/185 (2006.01)

(21) Application No: **2016377681** **(22) Date of Filing:** **2016.12.22**

(87) WIPO No: **WO17/112865**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/271,160	2015.12.22	US
62/271,146	2015.12.22	US
62/271,179	2015.12.22	US
62/271,137	2015.12.22	US
62/271,200	2015.12.22	US
62/338,474	2016.05.18	US
62/413,345	2016.10.26	US

(43) Publication Date: **2017.06.29**

(44) Accepted Journal Date: **2021.05.13**

(71) Applicant(s)
ModernaTX, Inc.

(72) Inventor(s)
Benenato, Kerry E.;Butcher, William

(74) Agent / Attorney
Griffith Hack, Level 10 161 Collins St, MELBOURNE, VIC, 3000, AU

(56) Related Art

WO 2016/118724 A1
WO 2009/129395 A1
WO 2015/199952 A1
WO 2009/086558 A1

Advances in Genetics, 2014, 88, 71-110

European Journal of Pharmaceutics and Biopharmaceutics, 2000, 50(1), 161-77

US 2012/0295832 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

WO 2017/112865 A1

(43) International Publication Date

29 June 2017 (29.06.2017)

WIPO | PCT

(51) International Patent Classification:

C07D 211/14 (2006.01) *A61K 31/495* (2006.01)
C07D 211/16 (2006.01) *A61K 31/445* (2006.01)
C07D 295/13 (2006.01) *A61K 31/45* (2006.01)
C07D 295/185 (2006.01) *A61K 9/14* (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/US2016/068300

(22) International Filing Date:

22 December 2016 (22.12.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/271,160	22 December 2015 (22.12.2015)	US
62/271,146	22 December 2015 (22.12.2015)	US
62/271,179	22 December 2015 (22.12.2015)	US
62/271,137	22 December 2015 (22.12.2015)	US
62/271,200	22 December 2015 (22.12.2015)	US
62/338,474	18 May 2016 (18.05.2016)	US
62/413,345	26 October 2016 (26.10.2016)	US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- Published:
- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))



WO 2017/112865 A1

(54) Title: COMPOUNDS AND COMPOSITIONS FOR INTRACELLULAR DELIVERY OF AGENTS

(57) Abstract: The disclosure features amino lipids and compositions involving the same. Nanoparticle compositions include an amino lipid as well as additional lipids such as phospholipids, structural lipids, PEG lipids, or a combination thereof. Nanoparticle compositions further including therapeutic and/or prophylactic agents such as RNA are useful in the delivery of therapeutic and/or prophylactic agents to mammalian cells or organs to, for example, regulate polypeptide, protein, or gene expression.

COMPOUNDS AND COMPOSITIONS FOR INTRACELLULAR DELIVERY OF AGENTS

RELATED APPLICATIONS

5 [001] This application claims priority to, and the benefit of, U.S. Provisional Application Nos. 62/271,160, filed December 22, 2015, 62/271,179, filed December 22, 2015, 62/271,137, filed December 22, 2015, 62/271,200, filed December 22, 2015, 62/271,146, filed December 22, 2015; 62/338,474, filed May 18, 2016; 62/413,345, filed October 26, 2016; the entire contents of each of which are incorporated herein by reference in their entireties.

10

TECHNICAL FIELD

15 [002] The present disclosure provides compounds, compositions comprising such compounds, and methods involving lipid nanoparticle compositions to deliver one or more therapeutic and/or prophylactic agents to and/or produce polypeptides in mammalian cells or organs. In addition to an amino lipid, lipid nanoparticle compositions of the disclosure may include one or more cationic and/or ionizable amino lipids, phospholipids including polyunsaturated lipids, PEG lipids, structural lipids, and/or therapeutic and/or prophylactic agents in specific fractions.

BACKGROUND

20 [003] The effective targeted delivery of biologically active substances such as small molecule drugs, proteins, and nucleic acids represents a continuing medical challenge. In particular, the delivery of nucleic acids to cells is made difficult by the relative instability and low cell permeability of such species. Thus, there exists a need to develop methods and compositions to facilitate the delivery of therapeutic and/or prophylactic agents such as nucleic acids to cells.


25 [004] Lipid-containing nanoparticle compositions, liposomes, and lipoplexes have proven effective as transport vehicles into cells and/or intracellular compartments for biologically active substances such as small molecule drugs, proteins, and nucleic acids. Such compositions generally include one or more “cationic” and/or amino (ionizable) lipids, phospholipids including polyunsaturated lipids, structural lipids (e.g., sterols), and/or lipids containing

30 polyethylene glycol (PEG lipids). Cationic and/or ionizable lipids include, for example, amine-containing lipids that can be readily protonated. Though a variety of such lipid-containing nanoparticle compositions have been demonstrated, improvements in safety, efficacy, and specificity are still lacking.


SUMMARY

[005] The present disclosure provides compounds and compositions and methods involving the same.

5 [006] In one aspect, the disclosure provides a compound having the formula (I)



or a salt or isomer thereof, wherein



t is 1 or 2;

10 A₁ and A₂ are each independently selected from CH or N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R"MR', -R*YR", -YR", and -R*OR";

15 each M is independently selected from the group consisting

of -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)O₂-, an aryl group, and a heteroaryl group;

X¹, X², and X³ are independently selected from the group consisting of a bond, -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-,

20 -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

each Y is independently a C₃₋₆ carbocycle;


each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆

25 carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl,









5 wherein when ring A is , then

- at least one of X¹, X², and X³ is not -CH₂-; and/or
- ii) at least one of R₁, R₂, R₃, R₄, and R₅ is -R''MR'.

[007] The compounds of formula (I) may include one or more of the following features when applicable.

10 [008] In some embodiments, the compound is of any of formulae (Ia1)-(Ia6):






[0009] In some embodiments, at least one of X^1 , X^2 , and X^3 is not $-\text{CH}_2-$. For example, in certain embodiments, X^1 is not $-\text{CH}_2-$. In some embodiments, at least one of X^1 , X^2 , and X^3 is $-\text{C}(\text{O})-$. In some embodiments, X^3 is a bond while each of X^1 and X^2 is not a bond. In some embodiments, none of X^1 , X^2 , and X^3 is a bond.

[0010] In some embodiments, R_1 and R_2 are the same. In certain embodiments, R_1 , R_2 , and R_3 are the same. In some embodiments, R_4 and R_5 are the same. In certain embodiments, R_1 , R_2 , R_3 , R_4 , and R_5 are the same.

[0011] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-\text{R}''\text{MR}'$. In some embodiments, at most one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-\text{R}''\text{MR}'$. For example, at least one of R_1 , R_2 , and R_3 may be $-\text{R}''\text{MR}'$, and/or at least one of R_4 and R_5 is $-\text{R}''\text{MR}'$. In certain embodiments, at least one M is $-\text{C}(\text{O})\text{O}-$. In some embodiments, each M is $-\text{C}(\text{O})\text{O}-$. In some embodiments, at least one M is $-\text{OC}(\text{O})-$. In some embodiments, each M is $-\text{OC}(\text{O})-$. In some embodiments, at least one R'' is C_3 alkyl. In certain embodiments, each R'' is C_3 alkyl. In some embodiments, at least one R'' is C_5 alkyl. In certain embodiments, each R'' is C_5 alkyl. In some embodiments, at least one R'' is C_6 alkyl. In certain embodiments, each R'' is C_6 alkyl. In some embodiments, at least one R'' is C_7 alkyl. In certain embodiments, each R'' is C_7 alkyl. In some embodiments, at least one R' is C_5 alkyl. In certain embodiments, each R' is C_5 alkyl. In other embodiments, at least one R' is C_1 alkyl. In certain embodiments, each R' is C_1 alkyl. In some embodiments, at least one R' is C_2 alkyl. In certain embodiments, each R' is C_2 alkyl.

[0012] In some embodiments, at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is C_{12} alkyl. In certain embodiments, each of R_1 , R_2 , R_3 , R_4 , and R_5 are C_{12} alkyl.

[0013] In another aspect, the disclosure provides a compound having formula (II):




or a salt or isomer thereof, wherein

5 A_1 and A_2 are each independently selected from CH or N and at least one of A_1 and A_2 is N;

Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , R_3 , R_4 , and R_5 are independently selected from the group consisting of C_{6-20} alkyl and C_{6-20} alkenyl;




10 wherein when ring A is , then

- i) R_1 , R_2 , R_3 , R_4 , and R_5 are the same, wherein R_1 is not C_{12} alkyl, C_{18} alkyl, or C_{18} alkenyl;
- ii) only one of R_1 , R_2 , R_3 , R_4 , and R_5 is selected from C_{6-20} alkenyl;
- iii) at least one of R_1 , R_2 , R_3 , R_4 , and R_5 have a different number of carbon atoms than at least one other of R_1 , R_2 , R_3 , R_4 , and R_5 ;
- iv) R_1 , R_2 , and R_3 are selected from C_{6-20} alkenyl, and R_4 and R_5 are selected from C_{6-20} alkyl; or
- v) R_1 , R_2 , and R_3 are selected from C_{6-20} alkyl, and R_4 and R_5 are selected from C_{6-20} alkenyl.

20 [0014] The compounds of formula (II) may include one or more of the following features when applicable.

[0015] In some embodiments, the compound is of formula (IIa):



[0016] In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same, and are not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl. In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same and are C₉ alkyl or C₁₄ alkyl.


[0017] In some embodiments, only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl. In certain such embodiments, R₁, R₂, R₃, R₄, and R₅ have the same number of carbon atoms. In some embodiments, R₄ is selected from C₅₋₂₀ alkenyl. For example, R₄ may be C₁₂ alkenyl or C₁₈ alkenyl.

10 [0018] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅.

[0019] In certain embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl. In other embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ have the same number of carbon atoms, and/or R₄ and R₅ have the same number of carbon atoms. For example, R₁, R₂, and R₃, or R₄ and R₅, may have 6, 8, 9, 12, 14, or 18 carbon atoms. In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are C₁₈ alkenyl (e.g., linoleyl). In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are alkyl groups including 6, 8, 9, 12, or 14 carbon atoms.

15 [0020] In some embodiments, R₁ has a different number of carbon atoms than R₂, R₃, R₄, and R₅. In other embodiments, R₃ has a different number of carbon atoms than R₁, R₂, R₄, and R₅. In further embodiments, R₄ has a different number of carbon atoms than R₁, R₂, R₃, and R₅.

[0021] In another aspect, the disclosure provides a compound according to formula (III):



or a salt or isomer thereof, in which

A_3 is CH or N;

A_4 is CH_2 or NH; and at least one of A_3 and A_4 is N or NH;

5 Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , and R_3 are independently selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R''MR'$, $-R^*YR''$, $-YR''$, and $-R^*OR''$;

each M is independently selected

10 from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$,
 $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

X^1 and X^2 are independently selected from the group consisting of $-CH_2-$,
 $-(CH_2)_2-$, $-CHR-$, $-CHY-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-C(O)-CH_2-$, $-CH_2-C(O)-$,
 $-C(O)O-CH_2-$, $-OC(O)-CH_2-$, $-CH_2-C(O)O-$, $-CH_2-OC(O)-$, $-CH(OH)-$, $-C(S)-$, and $-CH(SH)-$;

15 each Y is independently a C_{3-6} carbocycle;

each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;


each R is independently selected from the group consisting of C_{1-3} alkyl and a C_{3-6} carbocycle;

20 each R' is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each R'' is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl.

25 **[0022]** The compounds of formula (III) may include one or more of the following features when applicable.

[0023] In some embodiments, the compound is of formula (IIIa):




[0024] In some embodiments, R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ are the same.

5 In certain embodiments, R₁, R₂, and R₃ are C₆, C₉, C₁₂, or C₁₄ alkyl. In other embodiments, R₁, R₂, and R₃ are C₁₈ alkenyl. For example, R₁, R₂, and R₃ may be linoleyl.

[0025] In some embodiments, at least one of X¹ and X² is not -CH₂-¹. For example, in certain embodiments, X¹ is not -CH₂-¹. In some embodiments, at least one of X¹ and X² is -C(O)-¹.

[0026] In another aspect, the disclosure provides a compound according to formula (Ib):



or a salt or isomer thereof, in which

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

15 Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-¹, -(CH₂)₂-¹, -CHR-¹, -CHY-¹, -C(O)-¹, -C(O)O-¹, -OC(O)-¹, -C(O)-CH₂-¹, -CH₂-C(O)-¹, -C(O)O-CH₂-¹, -OC(O)-CH₂-¹, -CH₂-C(O)O-¹, -CH₂-OC(O)-¹, -CH(OH)-¹, -C(S)-¹, and -CH(SH)-¹;

20 R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl, -R"MR', -R*YR", -YR", and -R*OR";

each M is independently selected from the group consisting of -C(O)O-¹, -OC(O)-¹, -C(O)N(R')-¹, -N(R')C(O)-¹, -C(O)-¹, -C(S)-¹, -C(S)S-¹, -SC(S)-¹, -CH(OH)-¹, -P(O)(OR')O-¹, -S(O)O₂-¹, an aryl group, and a heteroaryl group;

each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

5 each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R" is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

[0027] The compounds of formula (Ib) may include one or more of the following features when

10 applicable.


[0028] In some embodiments, R₁ and R₂ are the same. In certain embodiments, R₁, R₂, and R₃ are the same. In some embodiments, R₄ and R₅ are the same. In certain embodiments, R₁, R₂, R₃, R₄, and R₅ are the same.

15 [0029] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is C₉₋₁₂ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ independently is C₉, C₁₂ or C₁₄ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ is C₉ alkyl.


[0030] In some embodiments, A₆ is N and A₇ is N. In some embodiments, A₆ is CH and A₇ is N.

20 [0031] In some embodiments, X⁴ is -CH₂- and X⁵ is -C(O)-. In some embodiments, X⁴ and X⁵ are -C(O)-.

[0032] In an embodiment, the compound has the formula (IV)



[0033] In another aspect, the disclosure provides a compound having the formula (17-I):



25 or a salt or isomer thereof, wherein R_{1a} is -(CH₂)_{n^a}Q^a, where Q^a is selected from a heterocycle, -OR^a, -O(CH₂)_{n^a}N(R^a)₂, -C(O)OR^a, -OC(O)R^a, -CX^a₃, -CX^a₂H, -CX^aH₂, -CN, -N(R^a

)₂, -C(O)N(R^a)₂, -N(R^a)C(O)R^a, and -N(R^a)S(O)₂R^a and each n^a is independently selected from 1, 2, 3, 4, and 5;

R_{2a} and R_{3a} are each independently selected from the group consisting of C₃₋₂₄ alkyl, C₃₋₂₄ alkenyl, -R^{a*}Y^aR^{a*}, -Y^aR^{a*}, and -R^{a*}OR^{a*};

5 each Y^a is independently a C₃₋₆ carbocycle;

each R^{a*} is independently selected from the group consisting of C₁₋₁₂ alkyl and C₁₋₁₂ alkenyl;

each X^a is independently selected from the group consisting of F, Cl, Br, and I;

10 each R^a is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H; and

each R^{a*} is selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl;


wherein R_{2a} includes 7 or fewer carbon atoms.

[0034] In some embodiments, Q^a is -OR^a. In certain embodiments, R^a is H. In other embodiments, R^a is -CH₃.

15 [0035] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[0036] In some embodiments, R_{3a} includes 7 or fewer carbon atoms.

[0037] In another aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(CH_2)_n^aQ^a$, where Q^a is selected from a heterocycle, $-OR^a$, $-O(CH_2)_n^aN(R^a)_2$, $-C(O)OR^a$, $-OC(O)R^a$, $-CX_3^a$, $-CX_2^aH$, $-CX^aH_2$, $-CN$, $-N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)R^a$, and $-N(R^a)S(O)_2R^a$ and each n^a is independently selected from 1, 2, 3, 4, and 5;

5 each X^a is independently selected from the group consisting of F, Cl, Br, and I;

R_{2a} is selected from the group consisting of C_{8-24} alkenyl;

R_{3a} is selected from the group consisting of C_{8-24} alkyl; and

each R^a is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.


10 [0038] In some embodiments, Q^a is $-OR^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-CH_3$.

[0039] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[0040] In some embodiments, R_{3a} is an alkyl including 9, 12, 14, or 18 carbon atoms.

15 [0041] In some embodiments, R_{2a} is C_{18} alkenyl (e.g., linoleyl).

[0042] In a further aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(CH_2)_n^aQ^a$, where Q^a is selected from a heterocycle, $-OR^a$, $-O(CH_2)_n^aN(R^a)_2$, $-C(O)OR^a$, $-OC(O)R^a$, $-CX_3^a$, $-CX_2^aH$, $-CX^aH_2$, $-CN$, $-N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)R^a$, and $-N(R^a)S(O)_2R^a$ and each n^a is independently selected from 1, 2, 3, 4, and 5;

each X^a is independently selected from the group consisting of F, Cl, Br, and I;

R_{2a} is selected from the group consisting of C_{13-20} alkyl;

R_{3a} is selected from the group consisting of C_{8-20} alkyl; and

25 each R^a is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.


[0043] In some embodiments, Q^a is $-OR^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-CH_3$.

[0044] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[0045] In some embodiments, R_{2a} and R_{3a} are the same.

[0046] In some embodiments, R_{2a} and/or R_{3a} is C_{14} alkyl.

5 [0047] In a further aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(CH_2)_n^a Q^a$, where Q^a is $-OR^a$, R^a is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H, and n^a is selected from 1, 2, 3, 4, and 5; and

10 R_{2a} and R_{3a} are each independently selected from the group consisting of C_{8-20} alkenyl, wherein

- i) R^a is selected from the group consisting of C_{1-3} alkyl and C_{2-3} alkenyl; or
- ii) R_{1a} is $-(CH_2)_2OH$, and R_{2a} and R_{3a} each include one or fewer double bonds.

[0048] In some embodiments, R^a is H. In other embodiments, R^a is $-CH_3$.


15 [0049] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[0050] In certain embodiments, R_{1a} is $-(CH_2)_2OCH_3$. In other embodiments, R_{1a} is $-(CH_2)_2OH$.

[0051] In some embodiments, R_{2a} is C_{18} alkenyl (e.g., linoleyl). In certain embodiments, R_{3a} is C_{18} alkenyl (e.g., linoleyl).

20 [0052] In some embodiments, R_{2a} and R_{3a} are the same.

[0053] In another aspect, the disclosure provides a compound of formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(CH_2)_n^a Q^a$, where Q^a is selected from a heterocycle, $-OR^a$, $-O(CH_2)_n^a N(R^a)_2$, $-C(O)OR^a$, $-OC(O)R^a$, $-CX^a_3$, $-CX^a_2H$, $-CX^aH_2$, $-CN$, $-N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)R^a$, and $-N(R^a)S(O)_2R^a$ and each n^a is independently selected from 1, 2, 3, 4, and 5;

5 each X^a is independently selected from the group consisting of F, Cl, Br, and I;

R_{2a} is selected from the group consisting of C_{8-12} alkyl;

R_{3a} is selected from the group consisting of C_{8-20} alkyl; and

each R^a is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.

10 [0054] In some embodiments, Q^a is $-OR^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-CH_3$.


[0055] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[0056] In certain embodiments, Q^a is $-OR^a$ and n^a is selected from 2, 3, and 4.

15 [0057] In some embodiments, R_{2a} is C_9 alkyl. In other embodiments, R_{2a} is C_{12} alkyl.

[0058] In some embodiments, R_{2a} and R_{3a} are the same.

[0059] In another aspect, the disclosure provides a compound having the formula (19-I),



(19-I),

or a salt or isomer thereof, wherein

20 R_{1b} is selected from the group consisting of H, C_{1-5} alkyl, C_{2-5} alkenyl, $-R^{b*}M^bR^b$, a C_{3-6} carbocycle, $-(CH_2)_nQ^b$, and $-(CH_2)_nCHQ^bR^b$, where Q^b is selected from a heterocycle, $-OR^b$, $-O(CH_2)_nN(R^b)_2$, $-C(O)OR^b$, $-OC(O)R^b$, $-CX^b_3$, $-CX^b_2H$, $-CX^bH_2$, $-CN$, $-N(R^b)_2$, $-C(O)N(R^b)_2$, $-N(R^b)C(O)R^b$, and $-N(R^b)S(O)_2R^b$ and each n is independently selected from 1, 2, 3, 4, and 5;

25 R_{2b} and R_{3b} are independently selected from the group consisting of C_{1-20} alkyl, C_{2-20} alkenyl, $-R^{b*}MR^b$, $-R^{b*}YR^b$, $-YR^b$, and $-R^{b*}OR^b$,

each M^b is independently selected from the group consisting of $-C(O)O-$, $-OC(O)-$, $-C(O)N(R^{b'})-$, $-N(R^{b'})C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR^{b'})O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

W is selected from the group consisting of $-CH_2-$, $-CHR^{b'}-$, $-C(O)-$, $-CH(OH)-$, $-C(S)-$,
5 and $-CH(SH)-$;

each X^b is independently selected from the group consisting of F, Cl, Br, and I;

each Y^b is independently a C_{3-6} carbocycle;

each R^{b*} is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

10 each R^b is independently selected from the group consisting of C_{1-3} alkyl, a C_{3-6} carbocycle, C_{2-3} alkenyl, and H;

each $R^{b''}$ is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

15 each $R^{b'''}$ is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl.

[0060] In some embodiments, W is not $-CH_2-$. In particular such embodiments, W is $-C(O)-$.


[0061] In some embodiments, at least one of R_{2b} and R_{3b} is $-R^{b''}M^bR^{b'}$. In certain embodiments, at least one M^b is $-C(O)O-$. In some embodiments, at least one $R^{b''}$ is C_5 alkyl. In certain embodiments, at least one $R^{b'}$ is C_5 alkyl.

20 **[0062]** In some embodiments, R_{2b} and/or R_{3b} are selected from the group consisting of C_{1-20} alkyl. For example, R_{2b} and/or R_{3b} may be alkyl groups including 9 or 12 carbon atoms. In other embodiments, R_{2b} and/or R_{3b} are selected from the group consisting of C_{2-20} alkenyl. For example, R_{2b} and/or R_{3b} may be alkenyl groups including 18 carbon atoms (e.g., linoleyl groups). In certain embodiments, R_{2b} and R_{3b} are the same.

25 **[0063]** In some embodiments, R_{1b} is H, while in other embodiments, R_{1b} is selected from C_{1-5} alkyl. For example, R_{1b} may be C_1 alkyl.

[0064] In certain embodiments, R_{1b} is $-(CH_2)_nQ^b$. In such embodiments, Q^b is a heterocycle such as a phenyl group. For example, Q^b may be a phenyl group with one or more substituents, as described herein.

30 **[0065]** Also disclosed herein are compounds of formula (19-II):



or salts or isomers thereof, wherein

R_{1b} is selected from the group consisting of C_{6-20} alkyl; and


R_{2b} and R_{3b} are independently selected from the group consisting of C_{6-20} alkenyl.

5 [0066] In particular embodiments, R_{1b} is C_{12} alkyl.

[0067] In some embodiments, R_{2b} and/or R_{3b} are C_{18} alkenyl (e.g., linoleyl).

[0068] In certain embodiments, R_{2b} and R_{3b} are both linoleyl.

[0069] In another aspect, the disclosure provides a compounds of formula (20-I):



10 or a salt or isomer thereof, wherein

R_{1c} is selected from the group consisting of a C_{3-6}

carbocycle, $-(CH_2)_n^cQ^c$, $-(CH_2)_n^cCHQ^cR^c$,

$-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from a heterocycle, $-OR^c$, $-O(CH_2)_n^cN(R^c)_2$,

$-C(O)OR^c$, $-OC(O)R^c$, $-CX^c_3$, $-CX^c_2H$, $-CX^cH_2$, $-CN$, $-N(R^c)_2$, $-C(O)N(R^c)_2$, $-N(R^c)C(O)R^c$,

15 and $-N(R^c)S(O)_2R^c$ and each n^c is independently selected from 1, 2, 3, 4, and 5;

R_{2c} , R_{3c} , and R_{4c} are independently selected from the group consisting of C_{1-20} alkyl, C_{2-20} alkenyl, $-R^{c*}M^cR^c$, $-R^{c*}Y^cR^c$, $-Y^cR^c$, and $-R^{c*}OR^c$;

each M^c is independently selected from the group consisting of $-C(O)O-$, $-OC(O)-$, $-C(O)N(R^c)-$, $-N(R^c)C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$,

20 $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

each X^c is independently selected from the group consisting of F, Cl, Br, and I;

each Y^c is independently a C_{3-6} carbocycle;

each R^{c*} is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R^{c*} is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

5 each R^{c**} is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl,

wherein

i) R_{1c} is selected from the group consisting of a C_{3-6}

carbocycle, $-(CH_2)_n^cQ^c$, $-(CH_2)_n^cCHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from a heterocycle, $-O(CH_2)_n^cN(R^c)_2$, $-C(O)OR^c$, $-OC(O)R^c$, $-CX^c_3$, $-CX^c_2H$, $-CX^cH_2$, $-C(O)N(R^c)_2$, $-N(R^c)C(O)R^c$, and $-N(R^c)S(O)_2R^c$ and each n^c is independently selected from 1, 2, 3, 4, and 5; and/or

ii) at least one of R_{2c} , R_{3c} , and R_{4c} is $-R^{c**}M^cR^c$.

[0070] In some embodiments, R_1 is selected from the group consisting

15 of $-(CH_2)_n^cQ^c$, $-(CH_2)_n^cCHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from a heterocycle, $-O(CH_2)_n^cN(R^c)_2$, $-C(O)OR^c$, $-OC(O)R^c$, $-CX^c_3$, $-CX^c_2H$, $-CX^cH_2$, $-CN$, $-C(O)N(R^c)_2$, $-N(R^c)C(O)R^c$, and $-N(R^c)S(O)_2R^c$ and each n is independently selected from 1, 2, 3, 4, and 5. In certain embodiments, R_{1c} is $-(CH_2)_n^cQ^c$. In some embodiments, n^c is 2. In some embodiments, Q^c is $-C(O)OR^c$, where R^c is, for example, H.

20 **[0071]** In some embodiments, at least one of R_{2c} , R_{3c} , and R_{4c} is $-R^{c**}M^cR^c$. For example, R_{2c} , R_{3c} , and/or R_{4c} may be $-R^{c**}M^cR^c$. In some embodiments, at least one M^c is $-C(O)O-$. In certain embodiments, each M^c is $-C(O)O-$. In some embodiments, at least one R^{c**} is C_5 or C_7 alkyl. In certain embodiments, each R^{c**} is C_5 alkyl. In other embodiments, each R^{c**} is C_7 alkyl. In some embodiments, at least one R^{c*} is C_5 , C_7 , or C_9 alkyl. In certain embodiments, each R^{c*} is C_5 alkyl. In other embodiments, each R^{c*} is C_7 alkyl. In other embodiments, each R^{c*} is C_9 alkyl. In some embodiments, R^{c*} is branched.

[0072] In some embodiments, R_{2c} , R_{3c} , and R_{4c} are selected from the group consisting of C_{5-20} alkyl. In certain embodiments, R_{2c} , R_{3c} , and R_{4c} are C_{12} alkyl.


30 **[0073]** In some embodiments, R_{2c} is selected from the group consisting of C_{5-20} alkyl and C_{5-20} alkenyl. For example, R_{2c} may be C_{12} alkyl.

[0074] In some embodiments, R_{3c} is selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. For example, R_{3c} may be C₆, C₉, or C₁₂ alkyl.

[0075] In some embodiments, R_{4c} is selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. For example, R_{4c} may be C₆, C₉, or C₁₂ alkyl.

5 [0076] In some embodiments, R_{3c} and R_{4c} are the same.

[0077] In yet another aspect, the disclosure provides a compound according to formula (20-I):



or a salt or isomer thereof, wherein

R₁ is selected from the group consisting of -(CH₂)_{n^c}Q^c, -(CH₂)_{n^c}CHQ^cR^c, -CHQ^cR^c,
10 and -CQ^c(R^c)₂, where Q^c is selected from -OR^c, -CN, and -N(R^c)₂, and n^c is selected from 1, 2, 3, 4, and 5;

R_{2c} and R_{3c} are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl;

15 R_{4c} is selected from the group consisting of C₁₃₋₂₀ alkyl and C₅₋₂₀ alkenyl; and each R^c is independently selected from the group consisting of C₁₋₃ alkyl, C₂₋₃ alkenyl, and H.

[0078] In some embodiments, R_{3c} is C₁₄ alkyl.

[0079] In some embodiments, R_{4c} is C₁₄ alkyl.

[0080] In some embodiments, R_{3c} is C₁₈ alkenyl. For example, R_{3c} may be linoleyl.


20 [0081] In some embodiments, R_{4c} is C₁₈ alkenyl. For example, R_{4c} may be linoleyl.

[0082] In some embodiments, R_{2c} is C₁₂ alkyl. In other embodiments, R_{2c} is C₁₄ alkyl. In some embodiments, R_{2c} is C₁₈ alkenyl. For example, R_{2c} may be linoleyl.

[0083] In some embodiments, R_{3c} and R_{4c} are the same.

25 [0084] In some embodiments, R_{1c} is -(CH₂)_{n^c}Q^c. In some embodiments, Q^c is -OR^c. For example, Q^c may be -OH. In some embodiments, n^c is 2 or 3.

[0085] The disclosure also provides a compound having formula (20-I):



or a salt or isomer thereof, wherein

R_1 is selected from the group consisting of $-(\text{CH}_2)_{n^c}\text{Q}^c$, $-(\text{CH}_2)_{n^c}\text{CHQ}^c\text{R}^c$, $-\text{CHQ}^c\text{R}^c$, and $-\text{CQ}^c(\text{R}^c)_2$, where Q^c is selected from $-\text{OR}^c$, $-\text{CN}$, and $-\text{N}(\text{R}^c)_2$, and n^c is selected from 1, 2, 5 3, 4, and 5;

R_{2c} , R_{3c} , and R_{4c} are independently selected from the group consisting of C_{6-20} alkyl and C_{6-20} alkenyl; and

each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H ;

10 wherein

- i) R_{2c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl, and/or
- ii) R_{3c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl.

[0086] In some embodiments, R_{2c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl. For example, R_{2c} may be C_6 or C_9 alkyl.

15 **[0087]** In some embodiments, R_{3c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl. For example, R_{3c} may be C_6 or C_9 alkyl.


[0088] In some embodiments, R_{3c} is C_{12} alkyl.

[0089] In some embodiments, R_{2c} is C_{12} alkyl.

[0090] In some embodiments, R_{4c} is C_6 , C_9 , or C_{12} alkyl.

20 **[0091]** In some embodiments, R_{1c} is $-(\text{CH}_2)_{n^c}\text{Q}^c$. In certain embodiments, Q^c is $-\text{OR}^c$. In some embodiments, R^c is H . In some embodiments, n^c is 2 or 3.

[0092] In still another aspect, the disclosure provides a compound according to formula (20-I):



or a salt or isomer thereof, wherein

R_{1c} is selected from the group consisting of $-(CH_2)_{n^c}Q^c$, $-(CH_2)_{n^c}CHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from $-OR^c$, $-CN$, and $-N(R^c)_2$, and n^c is selected from 1, 2, 3, 4, and 5;

R_{2c} is selected from the group consisting of H, C_{12-20} alkyl, and C_{6-20} alkenyl;


5 R_{3c} and R_{4c} are C_{12} alkyl; and

each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.

[0093] In some embodiments, R_{2c} is H. In other embodiments, R_{2c} is C_{12} alkyl or alkenyl. In some embodiments, R_{2c} is C_{14} alkyl. In other embodiments, R_{2c} is C_{18} alkenyl. For example, 10 R_{2c} may be linoleyl.

[0094] In some embodiments, R_{1c} is $-(CH_2)_{n^c}Q^c$. In certain embodiments, Q^c is $-OR^c$. For example, Q^c may be OH. In some embodiments, n^c is 2, 3, or 4.

[0095] In another aspect, the disclosure provides compounds of formula (21-I):



(21-I),

15 or salts or isomers thereof, wherein

R_{1d} and R_{2d} are independently selected from the group consisting of H, C_{1-5} alkyl, and C_{2-5} alkenyl;

n^d is selected from 1, 2, 3, 4, and 5;

k is selected from 0, 1, 2, and 3;

20 R_{3d} and R_{4d} are independently selected from the group consisting of C_{1-20} alkyl, C_{2-20} alkenyl,

$-R^{d*}M^dR^{d*}$, $-R^{d*}Y^dR^{d*}$, $-Y^dR^{d*}$, and $-R^{d*}OR^{d*}$;

each M^d is independently selected from the group consisting of $-C(O)O-$, $-OC(O)-$, $-C(O)N(R^d)-$, $-N(R^d)C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR^d)O-$,

25 and $-S(O)_2-$, or is absent;

each R^{d*} is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; each Y^d is independently a C_{3-6} carbocycle;

each R^{d*} is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

each R^{d*} is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

5 each R^{d*} is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl,

wherein R_{3d} and R_{4d} are bound to either i) the same carbon atom or ii) adjacent carbon atoms.

[0096] In some embodiments, R_{3d} and R_{4d} are bound to the same carbon atom. For example, R_{3d} and R_{4d} may be bound to a carbon atom adjacent to C^* . In certain embodiments, R_{3d} and R_{4d} are not bound to a carbon atom adjacent to C^* .

[0097] In other embodiments, R_{3d} and R_{4d} are bound to adjacent carbon atoms. In certain embodiments, one or both of R_{3d} and R_{4d} are bound to carbon atoms adjacent to C^* .

[0098] In some embodiments, k is 0. In other embodiments, k is 1, 2, or 3.

15 **[0099]** In certain embodiments, M^d is absent. In other embodiments, M^d is selected from the group consisting


of $-C(O)O-$, $-OC(O)-$, $-C(O)N(R^{d*})-$, $-N(R^{d*})C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR^{d*})O-$, and $-S(O)_2-$. In particular such embodiments, M^d is $-C(O)O-$.

[0100] In some embodiments, n^d is 1, 2, or 3.

20 **[00101]** In some embodiments, R_{1d} and/or R_{2d} are selected from C_{1-5} alkyl. In certain embodiments, R_{1d} and/or R_{2d} are C_1 alkyl.

[00102] In certain embodiments, R_{3d} and/or R_{4d} are selected from C_{2-20} alkenyl. In certain embodiments, R_{3d} and/or R_{4d} are alkenyl groups including 17, 18, or 19 carbon atoms. For example, R_{3d} and/or R_{4d} may be C_{18} alkenyl groups (e.g., linoleyl).

25 **[00103]** In a further aspect, the disclosure features a nanoparticle composition including a lipid component comprising a compound as described above (e.g., a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), and (20-I)). In some embodiments, the lipid component of the nanoparticle composition includes a phospholipid. In certain embodiments, a phospholipid of a nanoparticle composition includes a phospholipid moiety and one or more fatty acid moieties, one or more of which may be unsaturated. For example, a nanoparticle composition may include a lipid according to formula 30 (V)



in which R_p represents a phospholipid moiety and R₁ and R₂ represent unsaturated fatty acid moieties that may be the same or different.

[00104] A phospholipid moiety may be selected from the non-limiting group consisting of

5 phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl glycerol, phosphatidyl serine, phosphatidic acid, 2-lysophosphatidyl choline, and a sphingomyelin. A fatty acid moiety may be selected from the non-limiting group consisting of lauric acid, myristic acid, myristoleic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid, linoleic acid, alpha-linolenic acid, erucic acid, arachidic acid, arachidonic acid, phytanic acid, eicosapentaenoic acid, behenic acid, 10 docosapentaenoic acid, and docosahexaenoic acid. For example, in certain embodiments, a phospholipid is selected from the group consisting of 1,2-dilinoleoyl-sn-glycero-3-phosphocholine (DLPC), 1,2-dimyristoyl-sn-glycero-phosphocholine (DMPC), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), 15 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC), 1,2-diundecanoyl-sn-glycero-phosphocholine (DUPC), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), 1,2-di-O-octadecenyl-sn-glycero-3-phosphocholine (18:0 Diether PC), 1-oleoyl-2-cholesterylhemisuccinoyl-sn-glycero-3-phosphocholine (OChemSPC), 20 1-hexadecyl-sn-glycero-3-phosphocholine (C16 Lyso PC), 1,2-dilinolenoyl-sn-glycero-3-phosphocholine, 1,2-diarachidonoyl-sn-glycero-3-phosphocholine, 1,2-didocosahexaenoyl-sn-glycero-3-phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphoethanol 25 amine (DOPE), 1,2-diphytanoyl-sn-glycero-3-phosphoethanolamine (ME 16.0 PE), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine, 1,2-dilinoleoyl-sn-glycero-3-phosphoethanolamine, 1,2-dilinolenoyl-sn-glycero-3-phosphoethanolamine, 1,2-diarachidonoyl-sn-glycero-3-phosphoethanolamine,

1,2-didocosahexaenoyl-sn-glycero-3-phosphoethanolamine, 1,2-dioleoyl-sn-glycero-3-phospho-rac-(1-glycerol) sodium salt (DOPG), and sphingomyelin. In certain embodiments, the phospholipid is DOPE. In other embodiments, the phospholipid is DSPC. Non-natural species including natural species with modifications and substitutions including branching, oxidation, cyclization, and alkynes are also contemplated.

5 [00105] In some embodiments, the lipid component of the nanoparticle composition includes a structural lipid. In certain embodiments, a structural lipid is selected from the group consisting of cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, ursolic acid, and alpha-tocopherol. In certain embodiments, the 10 structural lipid is cholesterol.

10 [00106] In some embodiments, the lipid component of the nanoparticle composition includes a PEG lipid. In certain embodiments, the PEG lipid is selected from the group consisting of a PEG-modified phosphatidylethanolamine, a PEG-modified phosphatidic acid, a PEG-modified ceramide, a PEG-modified dialkylamine, a PEG-modified diacylglycerol, and a 15 PEG-modified dialkylglycerol.

15 [00107] In some embodiments, the nanoparticle composition includes a lipid component comprising a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), a phospholipid (which may or may not be unsaturated), a PEG lipid, and a structural lipid. In certain embodiments, the lipid component of 20 the nanoparticle composition includes about 30 mol % to about 60 mol % compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), about 0 mol % to about 30 mol % phospholipid, about 18.5 mol % to about 48.5 mol % structural lipid, and about 0 mol % to about 10 mol % of PEG lipid. In some embodiments, the lipid component of the nanoparticle composition includes about 30 mol % to about 45 mol % compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), about 5 mol % to about 25 mol % phospholipid, about 30 mol % to 25 about 40 mol % structural lipid, and about 0 mol % to about 10 mol % of PEG lipid. In some 30 embodiments, the lipid component of the nanoparticle composition includes about 35 mol % to about 55 mol % compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), about 5 mol % to about 25 mol % phospholipid, about 30 mol % to about 40 mol % structural lipid, and about 0 mol % to about 10 mol % of PEG lipid. In certain embodiments, the lipid component includes about 50 mol % said compound, about 10 mol % phospholipid, about 38.5 mol % structural lipid, and about 1.5 mol % of PEG

lipid. In other embodiments, the lipid component includes about 40 mol % said compound, about 20 mol % phospholipid, about 38.5 mol % structural lipid, and about 1.5 mol % of PEG lipid. In some of these embodiments, the phospholipid is DOPE, while in other embodiments the phospholipid is DSPC. In certain embodiments, the structural lipid is cholesterol. In certain 5 embodiments, the PEG lipid is PEG-DMG. In any of the above, the total content of the lipid component may not exceed 100%.

[00108] In some embodiments, the nanoparticle composition includes more than one phospholipid, PEG lipid, structural lipid, or other lipid. In certain embodiments, the nanoparticle composition further includes a cationic and/or ionizable lipid such as an amino-lipid. In certain embodiments, a cationic and/or ionizable lipid is selected from the group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridodecyl-15,18,21,24-tetraaza-octatriacontane (KL25), 1,2-dilinoleyl-1,2-dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA), 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA), 1,2-dioleyloxy-N,N-dimethylaminopropane (DODMA), 2-((8-[(3 β)-cholest-5-en-3-yloxy]octyl)oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA), (2R)-2-((8-[(3 β)-cholest-5-en-3-yloxy]octyl)oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)), (2S)-2-((8-[(3 β)-cholest-5-en-3-yloxy]octyl)oxy)-N,N-dimethyl-3-[(9Z,12Z)-octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)), (12Z, 15Z)-N,N-dimethyl-2-nonylhenicosa-12,15-dien-1-amine, and N,N-dimethyl-1-((1S,2R)-2-octylcyclopropyl)heptadecan-8-amine.

[00109] In some embodiments, the nanoparticle composition includes a therapeutic and/or prophylactic agent. In certain embodiments, the therapeutic and/or prophylactic agent may be selected from the group consisting of a protein, a small molecule drug, a cytotoxic agent, a radioactive ion, a chemotherapeutic agent, a vaccine, a compound that elicits an immune response, and/or a nucleic acid (such as a deoxyribonucleic acid or a ribonucleic acid). In certain embodiments, the therapeutic and/or prophylactic agent is a ribonucleic acid (RNA). An RNA may be selected from the group consisting of a small interfering RNA (siRNA), an

asymmetrical interfering RNA (aiRNA), a microRNA (miRNA), a Dicer-substrate RNA (dsRNA), a small hairpin RNA (shRNA), a messenger RNA (mRNA), and mixtures thereof. In certain embodiments, the therapeutic and/or prophylactic agent is a messenger RNA (mRNA).

5 An RNA of a nanoparticle composition may be naturally or non-naturally occurring and may include one or more of a stem loop, a chain terminating nucleoside, a polyA sequence, a polyadenylation signal, and/or a 5' cap structure.

[00110] In some embodiments, the nanoparticle composition includes more than one therapeutic and/or prophylactic agent, such as one or more RNAs. The therapeutic and/or prophylactic agents may be of the same or different types (e.g., two mRNAs, two siRNAs, one mRNA and one siRNA, one mRNA and one small molecule drug, etc.).

[00111] In some embodiments, the encapsulation efficiency of a therapeutic and/or prophylactic agent of a nanoparticle composition is at least 50%. In certain embodiments, the encapsulation efficiency is at least 80%. In certain embodiments, the encapsulation efficiency is greater than 90%.

15 [00112] In some embodiments, the wt/wt ratio of the lipid component to a therapeutic and/or prophylactic agent in the nanoparticle composition is from about 10:1 to about 60:1. In certain embodiments, the wt/wt ratio is about 20:1.

[00113] In some embodiments, the N:P ratio of the nanoparticle composition is from about 2:1 to about 30:1. In certain embodiments, the N:P ratio is from about 2:1 to about 8:1. In 20 certain embodiments, the N:P ratio is from about 5:1 to about 8:1. For example, the N:P ratio may be about 5.0:1, about 5.5:1, about 5.67:1, about 6.0:1, about 6.5:1, or about 7.0:1. In some embodiments, the mean size of a nanoparticle composition is from about 40 nm to about 150 nm. In certain embodiments, the mean size is from about 70 nm to about 100 nm. In one embodiment, the mean size may be about 80 to about 100 nm. In certain embodiments, the 25 mean size may be about 80 nm. In other embodiments, the mean size may be about 100 nm.

[00114] The polydispersity index of the nanoparticle composition is from about 0 to about 0.25 in certain embodiments. In certain embodiments, the polydispersity index is from about 0.10 to about 0.20.

30 [00115] In some embodiments, the nanoparticle composition has a zeta potential of about -10 mV to about +20 mV.

[00116] In some embodiments, upon contacting the compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) (e.g., any of Compounds 1-109) or a nanoparticle composition thereof with a mammalian cell, the cell uptake of the compound or

nanoparticle composition is LDLR-independent. In some embodiments, the cell uptake of the compound or nanoparticle composition is LDLR-dependent. In some embodiments, the cell uptake of the compound or nanoparticle composition is apoE-independent. In some embodiments, the cell uptake of the compound or nanoparticle composition is apoE-dependent.

5 In some embodiments, the cell uptake of the compound or nanoparticle composition is LDLR-apoE-interaction independent. In some embodiments, the cell uptake of the compound or nanoparticle composition is LDLR-apoE-interaction dependent.

[00117] In some embodiments, upon contacting the compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) (e.g., any of Compounds 1-109) or 10 the nanoparticle composition thereof with a mammalian cell to produce a polypeptide, the production of the polypeptide is higher in mammalian hepatocytes than cells from a different tissue (e.g., spleen or kidney).

[00118] In some embodiments, upon contacting the compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) (e.g., any of Compounds 1-109) or 15 the nanoparticle composition thereof with a mammalian cell to produce a polypeptide, the production of the polypeptide occurs substantively in mammalian hepatocytes (e.g., little or no production of the polypeptide in other cells, e.g., spleen cells or renal cells).

[00119] In some embodiments, the nanoparticle composition includes one or more other components including, but not limited to, one or more pharmaceutically acceptable excipients, 20 small hydrophobic molecules, therapeutic and/or prophylactic agents, carbohydrates, polymers, permeability enhancing molecules, buffers, and surface altering agents.

[00120] In yet another aspect, the disclosure features a pharmaceutical composition comprising a nanoparticle composition according to the preceding aspects and a pharmaceutically acceptable carrier. For example, the pharmaceutical composition is 25 refrigerated or frozen for storage and/or shipment (e.g., being stored at a temperature of 4 °C or lower, such as a temperature between about -150 °C and about 0 °C or between about -80 °C and about -20 °C (e.g., about -5 °C, -10 °C, -15 °C, -20 °C, -25 °C, -30 °C, -40 °C, -50 °C, -60 °C, -70 °C, -80 °C, -90 °C, -130 °C or -150 °C). For example, the pharmaceutical composition is a solution that is refrigerated for storage and/or shipment at, for example, about -20° C, -30 °C, - 30 40 °C, -50 °C, -60 °C, -70 °C, or -80 °C.

[00121] In a further aspect, the disclosure provides a method of delivering a therapeutic and/or prophylactic agent (e.g., an mRNA) to a cell (e.g., a mammalian cell). This method

includes the step of administering to a subject (e.g., a mammal, such as a human) a nanoparticle composition including (i) a lipid component including a phospholipid (such as a polyunsaturated lipid), a PEG lipid, a structural lipid, and a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) and (ii) a therapeutic and/or prophylactic agent, in which administering involves contacting the cell with the nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the cell.

5 [00122] In another aspect, the disclosure provides a method of producing a polypeptide of interest in a cell (e.g., a mammalian cell). The method includes the step of contacting the cell with a nanoparticle composition including (i) a lipid component including a phospholipid (such as a polyunsaturated lipid), a PEG lipid, a structural lipid, and a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) and (ii) an mRNA encoding the polypeptide of interest, whereby the mRNA is capable of being translated in the cell to produce the polypeptide.

10 15 [00123] In yet another aspect, the disclosure provides a method of treating a disease or disorder in a mammal (e.g., a human) in need thereof. The method includes the step of administering to the mammal a therapeutically effective amount of a nanoparticle composition including (i) a lipid component including a phospholipid (such as a polyunsaturated lipid), a PEG lipid, a structural lipid, and a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) and (ii) a therapeutic and/or prophylactic agent (e.g., an mRNA). In some embodiments, the disease or disorder is characterized by dysfunctional or aberrant protein or polypeptide activity. For example, the disease or disorder is selected from the group consisting of rare diseases, infectious diseases, cancer and proliferative diseases, genetic diseases (e.g., cystic fibrosis), autoimmune diseases, 20 25 diabetes, neurodegenerative diseases, cardio- and reno-vascular diseases, and metabolic diseases.

30 [00124] In a further aspect, the disclosure provides a method of delivering (e.g., specifically delivering) a therapeutic and/or prophylactic agent to a mammalian organ (e.g., a liver, spleen, lung, or femur). This method includes the step of administering to a subject (e.g., a mammal) a nanoparticle composition including (i) a lipid component including a phospholipid, a PEG lipid, a structural lipid, and a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) and (ii) a therapeutic and/or prophylactic agent (e.g., an mRNA), in which administering involves contacting the cell with the

nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the organ such as liver.

[00125] In a further aspect, the disclosure features a method for the enhanced delivery of a therapeutic and/or prophylactic agent (e.g., an mRNA) to a target tissue (e.g., a liver, spleen, 5 lung, or femur). This method includes administering to a subject (e.g., a mammal) a nanoparticle composition, the composition including (i) a lipid component including a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), a phospholipid, a structural lipid, and a PEG lipid; and (ii) a therapeutic and/or prophylactic agent, the administering including contacting the target tissue 10 with the nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the target tissue. In some embodiments, the delivery is enhanced as compared to a reference composition which comprises a reference lipid instead of a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I).

15 **[00126]** In yet another aspect, the disclosure features a method of lowering immunogenicity comprising introducing the nanoparticle composition of the disclosure into cells, wherein the nanoparticle composition reduces the induction of the cellular immune response of the cells to the nanoparticle composition, as compared to the induction of the cellular immune response in cells induced by a reference composition which comprises a reference lipid instead of a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I). For example, the cellular immune response is an innate immune response, an adaptive immune response, or both.

20 **[00127]** In certain embodiments of the above aspects, a cell contacted in a method is in a mammal.

25 **[00128]** In any of the preceding aspects, a mammal may be, for example, a rodent, non-human primate, or a human. In certain embodiments, the mammal is a human. In certain embodiments, the mammal is LDLR-deficient, or apoE-deficient, or both. In certain embodiments, the mammal is not LDLR-deficient. In certain embodiments, the mammal is not apoE-deficient. In certain embodiments, the mammal is neither LDLR-deficient nor apoE-deficient. In certain embodiments, the mammal has an abnormal LDLR-apoE interaction. In certain embodiments, the mammal has a normal LDLR-apoE interaction.

30 **[00129]** In any of the preceding aspects, a therapeutic and/or prophylactic agent may be an mRNA.

[00130] In some embodiments of the above methods, the therapeutic and/or prophylactic agent may be specifically delivered to a target tissue of interest (e.g., a mammalian liver, spleen, lung, or femur).

[00131] In some embodiments of the above methods, a polypeptide of interest may be 5 specifically produced in a target cell or tissue of interest (e.g., a hepatocyte, a mammalian liver, spleen, lung, or femur), e.g., the production of polypeptide is substantively higher in the target cell or tissue than in a non-target cell/tissue.

[00132] In some embodiments, the nanoparticle composition is administered 10 intravenously, intramuscularly, intradermally, subcutaneously, intra-arterially, intra-tumor, or by inhalation. A dose of about 0.001 mg/kg to about 10 mg/kg of therapeutic and/or prophylactic agent (e.g., mRNA) is administered to a mammal in certain embodiments.

[00133] In any of the preceding aspects, in some embodiments, the delivery (e.g., delivery efficiency) of the therapeutic and/or prophylactic agent to the mammalian cell is LDLR-independent. In some embodiments, the delivery of the therapeutic and/or prophylactic agent to 15 the mammalian cell is LDLR-dependent. In some embodiments, the delivery of the therapeutic and/or prophylactic agent to the mammalian cell is apoE-independent. In some embodiments, the delivery of the therapeutic and/or prophylactic agent to the mammalian cell is apoE-dependent. In some embodiments, the delivery of the therapeutic and/or prophylactic agent to the mammalian cell is LDLR-apoE-interaction independent. In some embodiments, the delivery 20 of the therapeutic and/or prophylactic agent to the mammalian cell is LDLR-apoE-interaction dependent.

[00134] In any of the preceding aspects, in some embodiments, the production (e.g., yield) of the polypeptide of interest in the mammalian cell is LDLR-independent. In some 25 embodiments, the production of the polypeptide of interest in the mammalian cell is LDLR-dependent. In some embodiments, the production of the polypeptide of interest in the mammalian cell is apoE-independent. In some embodiments, the production of the polypeptide of interest in the mammalian cell is apoE-dependent. In some embodiments, the production of the polypeptide of interest in the mammalian cell is LDLR-apoE-interaction independent. In some 30 embodiments, the production of the polypeptide of interest in the mammalian cell is LDLR-apoE-interaction dependent.


[00135] In the preceding aspects, one or more nanoparticle compositions each including one or more therapeutic and/or prophylactic agents may be used in combination. In some 28

and/or prophylactic agents may be simultaneously contacted with a cell or delivered to a mammalian cell or organ. In other embodiments, the one or more nanoparticle compositions are contacted with a cell or delivered to a mammalian cell or organ at different times.


[00136] In the preceding aspects, one or more additional therapeutic and/or prophylactic agents or compounds may be used in combination with a nanoparticle composition including a therapeutic and/or prophylactic agent. In some embodiments, an additional therapeutic and/or prophylactic agent or compound may be administered at or near the same time as a nanoparticle composition (e.g., within one hour). In other embodiments, an additional therapeutic and/or prophylactic agent or compound may be administered before or after (e.g., one or more hours before or after) a nanoparticle composition as a pretreatment or post-treatment therapy. In some embodiments, an additional therapeutic and/or prophylactic agent or compound is selected from the group consisting of an anti-inflammatory compound, a steroid (e.g., a corticosteroid), a statin, an estradiol, a BTK inhibitor, an S1P1 agonist, a glucocorticoid receptor modulator (GRM), or an anti-histamine. In certain embodiments, an additional therapeutic and/or prophylactic agent or compound is selected from the group consisting of dexamethasone, methotrexate, acetaminophen, an H1 receptor blocker, or an H2 receptor blocker.

The present invention as claimed herein is described in the following items 1 to 21:

1. A compound having formula (I) or formula (Ib):



or a salt or isomer thereof, wherein



t is 1 or 2;

A₁ and A₂ are each independently selected from CH or N;

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

Z is CH₂ or absent, wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R"MR', -R*YR", -YR", and -R*OR";

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group; wherein when the compound is of formula (Ib), then M is not -OC(O)O-;

X¹, X², and X³ are independently selected from the group consisting of a bond, -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;


each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and


each R" is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl;



wherein when the compound is of formula (I), and when ring A is then

- i) at least one of X¹, X², and X³ is not -CH₂-; and/or
- ii) at least one of R₁, R₂, R₃, R₄, and R₅ is -R"MR';


or



when the compound is of Formula (I) and ring A is , and X¹, X², and X³ are each -CH₂, then

at least one of A¹ and A² is N,


R₁, R₂, R₃, R₄ and R₅ are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl, and




when ring A is , then

- i) R₁, R₂, R₃, R₄, and R₅ are the same, wherein R₁ is not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl;
- ii) only one of R₁, R₂, R₃, R₄ and R₅ is selected from C₆₋₂₀ alkenyl;
- iii) at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅;
- iv) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl; or
- v) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl.

2. The compound of item 1, wherein ring A is



or






, in which the N atom is connected with X^2 .

3. The compound of any one of the preceding items, wherein in formula (I):

- (i) at least one of A_1 and A_2 is N;
- (ii) A_1 is N and A_2 is CH;
- (iii) A_1 is CH and A_2 is N;
- (iv) A_1 is N and A_2 is N; or
- (v) each of A_1 and A_2 is CH.

4. The compound of item 1, being of any of formulae (Ia1)-(Ia6):





5. The compound of any one of the preceding items, wherein R₁, R₂, and R₃ are:

- (i) independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl;
- (ii) the same;
- (iii) C₆ alkyl;
- (iv) C₉ alkyl;
- (v) C₁₂ alkyl;
- (vi) C₁₄ alkyl;
- (vii) C₁₈ alkenyl; or
- (viii) linoleyl.

6. The compound of any one of the preceding items, wherein

- (i) at least one of X¹ and X² is not -CH₂-;
- (ii) X¹ is not -CH₂-;


(iii) X^1 is $-C(O)-$;

(iv) X^2 is not $-CH_2-$; or

(v) X^2 is $-C(O)-$.

7. The compound of any one of the preceding items, wherein R_2 , R_3 , R_4 , and R_5 are the same.

8. The compound of item 1, wherein the compound is Compound 4:



(Compound 4)

or a salt or isomer thereof.

9. A nanoparticle composition comprising a lipid component comprising a compound of any one of the preceding items, wherein the lipid component further comprises (i) a phospholipid; (ii) a structural lipid, and/or (iii) a PEG lipid.

10. The nanoparticle composition of item 9, wherein the phospholipid is selected from the group consisting of 1,2-dilinoleoyl-sn-glycero-3-phosphocholine (DLPC), 1,2-dimyristoyl-sn-glycero-phosphocholine (DMPC), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), 1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC), 1,2-diundecanoyl-sn-glycero-phosphocholine (DUPC), 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), 1,2-di-O-octadecenyl-sn-glycero-3-phosphocholine (18:0 Diether PC), 1-oleoyl-2-cholesterylhemisuccinoyl-sn-glycero-3-phosphocholine (OChemSPC), 1-hexadecyl-sn-glycero-3-phosphocholine (C16 Lyso PC), 1,2-dilinolenoyl-sn-glycero-3-phosphocholine, 1,2-diarachidonoyl-sn-glycero-3-phosphocholine, 1,2-didocosahexaenoyl-sn-glycero-3-phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), 1,2-diphytanoyl-sn-glycero-3-phosphoethanolamine (ME 16.0 PE), 1,2-distearoyl-sn-glycero-3-phosphoethanolamine,

1,2-dilinoleoyl-sn-glycero-3-phosphoethanolamine,
1,2-dilinolenoyl-sn-glycero-3-phosphoethanolamine,
1,2-diarachidonoyl-sn-glycero-3-phosphoethanolamine,
1,2-didocosahexaenoyl-sn-glycero-3-phosphoethanolamine,
1,2-dioleoyl-sn-glycero-3-phospho-rac-(1-glycerol) sodium salt (DOPG), sphingomyelin, and mixtures thereof.

11. The nanoparticle composition of item 9 or 10, wherein the structural lipid is selected from the group consisting of cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, ursolic acid, alpha-tocopherol, and mixtures thereof.
12. The nanoparticle composition of any one of items 9-11, wherein the PEG lipid is selected from the group consisting of a PEG-modified phosphatidylethanolamine, a PEG-modified phosphatidic acid, a PEG-modified ceramide, a PEG-modified dialkylamine, a PEG-modified diacylglycerol, a PEG-modified dialkylglycerol, and mixtures thereof.
13. The nanoparticle composition of any one of items 9-12, further comprising a therapeutic and/or prophylactic agent.
14. The nanoparticle composition of item 13, wherein the therapeutic and/or prophylactic agent is a ribonucleic acid (RNA).
15. The nanoparticle composition of item 14, wherein the RNA is selected from the group consisting of a small interfering RNA (siRNA), an asymmetrical interfering RNA (aiRNA), a microRNA (miRNA), a Dicer-substrate RNA (dsRNA), a small hairpin RNA (shRNA), a messenger RNA (mRNA), and mixtures thereof.
16. A method of delivering a therapeutic and/or prophylactic agent to a mammalian cell, the method comprising administering to a subject the nanoparticle composition of any one of items 13-15, said administering comprising contacting the cell with the nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the cell.
17. A method of producing a polypeptide of interest in a mammalian cell, the method comprising contacting the cell with the nanoparticle composition of any one of items 13-15,

wherein the therapeutic and/or prophylactic agent is an mRNA, and wherein the mRNA encodes the polypeptide of interest, whereby the mRNA is capable of being translated in the cell to produce the polypeptide of interest.

18. A method of treating a disease or disorder in a mammal in need thereof, the method comprising administering to the mammal a therapeutically effective amount of the nanoparticle composition of any one of items 13-15, wherein the disease or disorder is characterized by dysfunctional or aberrant protein or polypeptide activity.

19. The method of item 18, wherein the mammal (a) is LDLR-deficient; (b) is apoE-deficient; and/or (c) mammal has an abnormal LDLR-apoE interaction.

20. The method of item 18, wherein the mammal (a) is not LDLR-deficient; (b) is not apoE-deficient; and/or (c) has a normal LDLR-apoE interaction.

21. Use of the compound of any one of items 1 to 28, or the nanoparticle composition of any one of items 13-15, in the manufacture of a medicament for delivering a therapeutic and/or prophylactic agent to a mammalian cell and/or treating a disease or disorder in a mammal in need thereof, wherein the disease or disorder is characterized by dysfunctional or aberrant protein or polypeptide activity.

BRIEF DESCRIPTION OF THE DRAWINGS

[00137] The skilled artisan will understand that the drawings primarily are for illustrative purposes and are not intended to limit the scope of the inventive subject matter described herein. The drawings are not necessarily to scale; in some instances, various aspects of the inventive subject matter disclosed herein may be shown exaggerated or enlarged in the drawings to facilitate an understanding of different features. In the drawings, like reference characters generally refer to like features (e.g., functionally similar and/or structurally similar elements).

[00138] The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

[00139] The above and further features will be more clearly appreciated from the following detailed description when taken in conjunction with the accompanying drawings.

[00140] Figure 1 is a pair of graphs comparing luciferase expression levels in mice (whole body) after administration of nanoparticle compositions containing compounds of the disclosure over time.

[00141] Figure 2 is a graph summarizing luciferase expression levels at 3 h after administration of nanoparticle compositions containing compounds of the disclosure. Total light flux values were acquired via body luminescent imaging (BLI) 3 h after administration. In this Figure, the numbers 1-12 refer to the compositions containing Compounds 42-52 and MC3 respectively.

[00142] Figure 3 is a graph summarizing luciferase expression levels at 6 hr after administration of nanoparticle compositions containing compounds of the disclosure. Total light flux values were acquired via BLI 6 h after administration. In this Figure, the numbers 1-12 refer to the compositions containing Compounds 42-52 and MC3 respectively.

[00143] Figure 4 is graph summarizing luciferase expression levels at 24 h after administration of nanoparticle compositions containing compounds of the disclosure. Total light flux values were acquired via BLI 24 h after administration. In this Figure, the numbers 1-12 refer to the compositions containing Compounds 42-52 and MC3 respectively.

[00144] Figure 5 is graph summarizing expression levels of luciferase in mouse liver 6 hours after administration of nanoparticle compositions including compounds of the disclosure. In this Figure, the numbers 1-12 refer to the compositions containing Compounds 42-52 and MC3 respectively.

[00145] Figure 6 is a graph summarizing expression levels of luciferase in mouse lungs 6 hours after administration of nanoparticle compositions including compounds of the disclosure. In this Figure, the numbers 1-12 refer to the compositions containing Compounds 42-52 and MC3 respectively.

[00146] Figure 7 is a graph summarizing expression levels of luciferase in mouse spleen 6 hours after administration of nanoparticle compositions including compounds of the disclosure. In this Figure, the numbers 1-12 refer to the compositions containing Compounds 42-52 and MC3 respectively.

[00147] Figure 8 is a pair of graphs illustrating hEPO expression levels in rats dosed with compounds of the disclosure as compared to with KL22, showing that KL22 and its chain length derivatives (previously showing improved protein expression in mice), do not express hEPO in rats. PBS (phosphate buffered saline) is used as control. Graph A compares the hEPO concentration after administration of nanoparticle compositions containing Compound 23, Compound 11, KL22, and MC3 at 2 mpk, i.v. administration. Compound 11 LNP showed hEPO expression comparable to MC3 and improved tolerability as compared to KL22. Graph B illustrates the results of a dose response study using Compound 20, KL22, and MC3 at dose of

0.2, 0.5 and 1 mpk. KL22 and Compound 23 were shown to be toxic at the 2 mg/kg dose. In Graph A, the numbers 1-5 refer to compositions containing the following: 1: Compound 23, 2: Compound 11, 3: KL22; 4: MC3; 5: PBS. In Graph B, the numbers 1-9 refer to compositions containing the following: 1: Compound 20, 0.2 mpk; 2: Compound 20, 0.5 mpk; 3: Compound 20, 1 mpk; 4: KL22, 0.2 mpk; 5: KL22, 0.5 mpk; 6: KL22, 1 mpk; 7: MC3, 0.2 mpk; 8: MC3, 0.5 mpk; 9: MC3, 1 mpk; 10: PBS.

5 [00148] Figure 9 is a pair of graphs illustrating performance of Compound 11 as an LDLr independent lipid. Graph A is a bar graph showing expression of luciferase induced by administration of nanoparticle compositions including Compound 11 at dosages of 0.05 mpk, 0.25 mpk, and 0.5 mpk to LDLR -/- knockout and wild-type mice. Graph B shows LDL-c levels in LDLR knockout mice after administration of a control mRNA, i.e., non-translating Factor IX (“NT-FIX”) and various compositions comprising mRNAs encoding the LDL receptor in mice, with KL22 at 0.5 mpk, or with Compound 11 at 0.5 and 1 mpk. LDL-c levels in mice were found to drop with nanoparticle composition containing Compound 11.

10 [00149] Figure 10 is a graph showing hEPO levels in non-human primates up to ~50 h after administration of a nanoparticle composition containing Compound 4, compared to a composition containing MC3. The Compound 4 LNP demonstrated 3-fold expression of hEPO compared to MC3, establishing Compound 4 as an LDLr independent lipid that translates to higher species.

15 [00150] Figure 11 is a pair of images comparing the results of a mouse liver immunohistochemistry (IHC) using mRNA expressing green fluorescent protein (GFP) after administration of nanoparticle compositions containing Compound 4 and MC3. Graph A shows CD-1 mouse liver cells after administration of GFP mRNA in a MC3 LNP, 6h after intravenous administration at a dose of 0.5 mpk. GFP mRNA Protein expression from the MC3 LNP composition was observed in both hepatocytes and Kupffer cells. Graph B shows LDLR knockout mouse liver cells after administration of GFP mRNA in a Compound 4 LNP, 8h after intravenous administration at a dose of 0.5 mpk. In contrast to MC3, the LNP containing Compound 4 appears to show less protein expression in Kupffer cells.

20 [00151] Figure 12 is a set of graphs illustrating hEPO expression levels in CD-1 mice dosed with compounds of the disclosure, compared to MC3. PBS is used as control. Graph A shows the hEPO concentration 3 h after administration of the nanoparticle compositions. Graph B shows the hEPO concentration 6 h after administration of the nanoparticle compositions. Graph C shows the hEPO concentration 24 h after administration of the nanoparticle

compositions. In graphs A-C numbers 1-14 refer to compositions containing the following: 1: Compound 73, 2: Compound 80, 3: Compound 70; 4: Compound 81; 5: Compound 69; 6: Compound 82; 7: Compound 83; 8: Compound 62; 9: Compound 84; 10: Compound 85; 11: Compound 86; 12: Compound 87; 13: MC3; 14: PBS.

5 [00152] Figure 13 is a graph showing hEPO levels (pg/mL) in CD-1 mice up to ~25 h after administration of a nanoparticle composition containing compounds of the disclosure, compared to a composition containing MC3. Numbers 1-13 refer to compositions containing the following: 1: Compound 73, 2: Compound 80, 3: Compound 70; 4: Compound 81; 5: Compound 69; 6: Compound 82; 7: Compound 83; 8: Compound 62; 9: Compound 84; 10: Compound 85; 11: Compound 86; 12: Compound 87; 13: MC3.

10

[00153] Figure 14 is a pair of graphs showing percentages of activated B-cells in the spleens of CD-1 mice dosed with compounds of the disclosure, compared to MC3, and compared to mice not having received any treatment (naïve test subject). PBS is used as control. Graph A shows the percentage of CD19+ cells. Graph B shows the percentage of CD19+ CD69+ CD86+ cells. Numbers 1-13 refer to compositions containing the following: 1: Compound 73, 2: Compound 80, 3: Compound 70; 4: Compound 81; 5: Compound 69; 6: Compound 82; 7: Compound 83; 8: Compound 62; 9: Compound 84; 10: Compound 85; 11: Compound 86; 12: Compound 87; 13: MC3; 14: PBS; 15: treatment naïve subject.

15 [00154] Figure 15 is a graph summarizing luciferase expression levels at 6 h after administration of nanoparticle compositions containing compounds of the disclosure to CD-1 mice at a dose of 0.5 mpk. Total light flux values were acquired via body luminescent imaging (BLI) 6 h after administration. In this Figure, the numbers 1-7 and 10 refer to the compositions containing Compounds 4-10 and MC3 respectively.

20 [00155] Figure 16 is a graph showing GFP levels in the livers of LDLR knockout mice at 30 min to 24 h after intravenous administration of an eGFP RNA in a lipid composition containing Compound 4. The liver GFP levels were determined via IHC. The square markers represent the number of GFP positive cells following administration of a dose of 0.1 mpk of the composition. The circular markers represent the number of GFP positive cells following administration of a dose of 0.5 mpk of the composition.

25 [00156] Figure 17 is a pair of graphs showing the ApoE dependence of luciferase (“Luc”) expression following administration of a composition containing Luc mRNA and Compound 4 to mice at a dose 0.5 mpk. The expression following administration of a composition containing Luc mRNA and MC3 is presented for comparison. Graph A shows the total flux in the liver 6h

after administration. The % change in Luc expression in livers of ApoE knockout vs. wild-type mice (i.e., (WT mean expression-KO mean expression)/WT mean expression]*100%) was 91.9 % for Compound 4 and 97.5 % for MC3. Graph B shows the total flux in the spleen 6h after administration. The % change in expression in spleens of ApoE knockout vs. wild-type mice

5 was 4.34 % for Compound 4 and 72.2 % for MC3. Numbers 1-4 refer to the following: 1: Composition containing Compound 4, administered to ApoE knockout mice; 2: Composition containing Compound 4, administered to wild-type mice; 3: Composition containing MC3, administered to ApoE knockout mice; 4: Composition containing MC3, administered to wild-type mice.

10 [00157] Figure 18 is a pair of graphs showing the effect of a composition containing Compound 4 on liver enzymes. The composition was administered to rats at 0.1 mpk and 1 mpk. The effects of MC3 are shown for comparison. PBS is used as a control. Graph A shows the effect on aspartate aminotransferase (AST). Graph B shows the effect on alanine aminotransferase (ALT).

15 [00158] Figure 19 is a set of graphs showing immune cell activation by a composition containing Compound 4. The effects of MC3 are shown for comparison. PBS (phosphate buffered saline) is used as control. Compositions were administered to rats at 0.1 mpk or 1 mpk. Graph A shows the effect on activation of neutrophil. Graph B shows the effect on activation of lymphocytes. Graph C shows the effect on activation of monocytes. Numbers 1-5 in Graphs A-20 C refer to the following: 1: Compound 4, 0.1 mpk; 2: Compound 4; 1 mpk; 3: MC3, 0.1 mpk 4: MC3, 1 mpk; 5: PBS.

[00159] Figure 20 is a graph showing the expression of Stefin A Quadruple Mutant-Tracy (SQT) protein in mouse liver determined via FLAG IHC at different time points following intravenous administration of various nanoparticle compositions comprising SQT mRNA and 25 lipids disclosed herein. Numbers 1-11 in the figure refer to the following: 1: 0 h, PBS; 2: 0 h, Compound 4; 3: 0.5 h, Compound 4; 4: 4 h, Compound 4; 5: 8 h, Compound 4; 6: 24 h, Compound 4; 7: 0 h, MC3; 8: 0.5 h, MC3; 9: 4 h, MC3; 10: 8 h, MC3; 11: 24 h, MC3.

DETAILED DESCRIPTION


30 [00160] The disclosure relates to novel lipids and lipid nanoparticle compositions including a novel lipid. The disclosure also provides methods of delivering a therapeutic and/or prophylactic agent to a mammalian cell, specifically delivering a therapeutic and/or prophylactic agent to a mammalian organ, producing a polypeptide of interest in a mammalian cell, and

treating a disease or disorder in a mammal in need thereof. For example, a method of producing a polypeptide of interest in a cell involves contacting a nanoparticle composition comprising an mRNA with a mammalian cell, whereby the mRNA may be translated to produce the polypeptide of interest. A method of delivering a therapeutic and/or prophylactic agent to a mammalian cell or organ may involve administration of a nanoparticle composition including the therapeutic and/or prophylactic agent to a subject, in which the administration involves contacting the cell or organ with the composition, whereby the therapeutic and/or prophylactic agent is delivered to the cell or organ.


10 Lipids

[00161] The present disclosure provides lipids including a central piperazine moiety. The lipids described herein may be advantageously used in lipid nanoparticle compositions for the delivery of therapeutic and/or prophylactic agents to mammalian cells or organs. For example, the lipids described herein have little or no immunogenicity. For example, the lipid compound of any of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) has a lower immunogenicity as compared to a reference lipid (*e.g.*, MC3, KC2, or DLinDMA). For example, a formulation comprising a lipid disclosed herein and a therapeutic or prophylactic agent has an increased therapeutic index as compared to a corresponding formulation which comprises a reference lipid (*e.g.*, MC3, KC2, or DLinDMA) and the same therapeutic or prophylactic agent.

[00162] Lipids may be compounds of formula (I),



or salts or isomers thereof, wherein



25 t is 1 or 2;

A_1 and A_2 are each independently selected from CH or N;

Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , R_3 , R_4 , and R_5 are independently selected from the group consisting of C_{5-20}

5 alkyl, C_{5-20} alkenyl, $-R''MR'$, $-R^*YR''$, $-YR''$, and $-R^*OR''$;

each M is independently selected from the group consisting

of $-C(O)O-$, $-OC(O)-$, $-OC(O)O-$, $-C(O)N(R')$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

X^1 , X^2 , and X^3 are independently selected from the group consisting of a bond, $-CH_2-$,

10 $-(CH_2)_2-$, $-CHR-$, $-CHY-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-C(O)-CH_2-$, $-CH_2-C(O)-$,

$-C(O)O-CH_2-$, $-OC(O)-CH_2-$, $-CH_2-C(O)O-$, $-CH_2-OC(O)-$, $-CH(OH)-$, $-C(S)-$, and $-CH(SH)-$;

each Y is independently a C_{3-6} carbocycle;


each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;


15 each R is independently selected from the group consisting of C_{1-3} alkyl and a C_{3-6} carbocycle;

each R' is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each R'' is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12}

20 alkenyl,









wherein when ring A is , then

i) at least one of X^1 , X^2 , and X^3 is not $-CH_2-$; and/or

ii) at least one of R_1 , R_2 , R_3 , R_4 , and R_5 is $-R''MR'$.

[00163] In some embodiments, the compound is of any of formulae (Ia1)-(Ia6):








5

[00164] The compounds of Formula (I) or any of (Ia1)-(Ia6) include one or more of the following features when applicable.


[00165] In some embodiments, ring A is




[00166] In some embodiments, ring A is





[00167] In some embodiments, ring A is



[00168] In some embodiments, ring A is



[00169] In some embodiments, ring A is



, or

[00170] In some embodiments, ring A is

wherein ring, in which the N atom is connected with X^2 .

[00171] In some embodiments, Z is CH_2 .

[00172] In some embodiments, Z is absent.

10 [00173] In some embodiments, at least one of A_1 and A_2 is N.

[00174] In some embodiments, each of A_1 and A_2 is N.

[00175] In some embodiments, each of A_1 and A_2 is CH.

[00176] In some embodiments, A_1 is N and A_2 is CH.

[00177] In some embodiments, A_1 is CH and A_2 is N.

[00178] In some embodiments, at least one of X¹, X², and X³ is not -CH₂- . For example, in certain embodiments, X¹ is not -CH₂- . In some embodiments, at least one of X¹, X², and X³ is -C(O)-.

[00179] In some embodiments, X² is -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, 5 -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, or -CH₂-OC(O)-.

[00180] In some embodiments, X³ is -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, or -CH₂-OC(O)-. In other embodiments, X³ is -CH₂-.


[00181] In some embodiments, X³ is a bond or -(CH₂)₂-.

10 **[00182]** In some embodiments, R₁ and R₂ are the same. In certain embodiments, R₁, R₂, and R₃ are the same. In some embodiments, R₄ and R₅ are the same. In certain embodiments, R₁, R₂, R₃, R₄, and R₅ are the same.


15 **[00183]** In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is -R"MR'. In some embodiments, at most one of R₁, R₂, R₃, R₄, and R₅ is -R"MR'. For example, at least one of R₁, R₂, and R₃ may be -R"MR', and/or at least one of R₄ and R₅ is -R"MR'. In certain 20 embodiments, at least one M is -C(O)O-. In some embodiments, each M is -C(O)O-. In some embodiments, at least one M is -OC(O)-. In some embodiments, each M is -OC(O)-. In some embodiments, at least one M is -OC(O)O-. In some embodiments, each M is -OC(O)O-. In some embodiments, at least one R" is C₃ alkyl. In certain embodiments, each R" is C₃ alkyl. In some embodiments, at least one R" is C₅ alkyl. In certain embodiments, each R" is C₅ alkyl. In some embodiments, at least one R" is C₆ alkyl. In certain embodiments, each R" is C₆ alkyl. In some embodiments, at least one R" is C₇ alkyl. In certain embodiments, each R" is C₇ alkyl. In some embodiments, at least one R' is C₅ alkyl. In certain embodiments, each R' is C₅ alkyl. In other 25 embodiments, at least one R' is C₁ alkyl. In certain embodiments, each R' is C₁ alkyl. In some embodiments, at least one R' is C₂ alkyl. In certain embodiments, each R' is C₂ alkyl.


[00184] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is C₁₂ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ are C₁₂ alkyl.

[00185] In certain embodiments, the compound is selected from the group consisting of:




(Compound 1),







(Compound 11),




(Compound 12),




(Compound 13),




(Compound 14),




(Compound 15),




(Compound 16),




(Compound 42),




(Compound 43),




(Compound 44),




(Compound 45),




(Compound 46),




(Compound 47),




(Compound 48),




(Compound 49),




(Compound 50),




(Compound 51),




(Compound 52),




(Compound 53),




(Compound 54),




(Compound 55),




(Compound 56),




(Compound 57),




(Compound 58),




(Compound 59),




(Compound 60),




(Compound 61),




(Compound 62),



(Compound 63),




(Compound 64),




5


(Compound 65),




(Compound 66),




(Compound 68),



(Compound 69),




(Compound 70),




(Compound 71),


5




(Compound 72),




(Compound 73),




(Compound 74),




(Compound 75),




(Compound 76),




(Compound 78),




(Compound 79)




(Compound 80),




(Compound 81),




(Compound 82),




(Compound 83),




(Compound 84),




(Compound 85),




(Compound 86),




(Compound 87),




(Compound 88),




(Compound 89),




(Compound 90),




5 (Compound 91),




(Compound 92),




(Compound 93),




(Compound 94),



(Compound 95),





(Compound 96),




(Compound 97),

10





[00186] In other embodiments, a lipid has the formula (II)



5

(II),


or a salt or isomer thereof, wherein


A₁ and A₂ are each independently selected from CH or N and at least one of A₁ and A₂ is

N;

10 Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl;




wherein when ring A is , then

- i) R₁, R₂, R₃, R₄, and R₅ are the same, wherein R₁ is not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl;
- ii) only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl;
- iii) at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅;

iv) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl; or

v) R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl.

5 [00187] In some embodiments, the compound is of formula (IIa):



(IIa).

[00188] The compounds of Formula (II) or (IIa) include one or more of the following features when applicable.

[00189] In some embodiments, Z is CH₂.

10 [00190] In some embodiments, Z is absent.

[00191] In some embodiments, at least one of A₁ and A₂ is N.

[00192] In some embodiments, each of A₁ and A₂ is N.

[00193] In some embodiments, each of A₁ and A₂ is CH.

[00194] In some embodiments, A₁ is N and A₂ is CH.

15 [00195] In some embodiments, A₁ is CH and A₂ is N.

[00196] In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same, and are not C₁₂ alkyl, C₁₈ alkyl, or C₁₈ alkenyl. In some embodiments, R₁, R₂, R₃, R₄, and R₅ are the same and are C₉ alkyl or C₁₄ alkyl.

[00197] In some embodiments, only one of R₁, R₂, R₃, R₄, and R₅ is selected from C₆₋₂₀ alkenyl. In certain such embodiments, R₁, R₂, R₃, R₄, and R₅ have the same number of carbon atoms. In some embodiments, R₄ is selected from C₅₋₂₀ alkenyl. For example, R₄ may be C₁₂ alkenyl or C₁₈ alkenyl.

[00198] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ have a different number of carbon atoms than at least one other of R₁, R₂, R₃, R₄, and R₅.


25 [00199] In certain embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkenyl, and R₄ and R₅ are selected from C₆₋₂₀ alkyl. In other embodiments, R₁, R₂, and R₃ are selected from C₆₋₂₀ alkyl, and R₄ and R₅ are selected from C₆₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃

have the same number of carbon atoms, and/or R₄ and R₅ have the same number of carbon atoms. For example, R₁, R₂, and R₃, or R₄ and R₅, may have 6, 8, 9, 12, 14, or 18 carbon atoms. In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are C₁₈ alkenyl (e.g., linoleyl). In some embodiments, R₁, R₂, and R₃, or R₄ and R₅, are alkyl groups including 6, 8, 9, 12, or 14 carbon atoms.

5 atoms.


[00200] In some embodiments, R₁ has a different number of carbon atoms than R₂, R₃, R₄, and R₅. In other embodiments, R₃ has a different number of carbon atoms than R₁, R₂, R₄, and R₅. In further embodiments, R₄ has a different number of carbon atoms than R₁, R₂, R₃, and R₅.

[00201] In some embodiments, the compound is selected from the group consisting of:




10

(Compound 17),




(Compound 18),




15


(Compound 19),




(Compound 20),




(Compound 21),



(Compound 22),




(Compound 23),




5


(Compound 24),




(Compound 25),




10 (Compound 26),




(Compound 27),




(Compound 28),



(Compound 29),




(Compound 30),




5


(Compound 31),



(Compound 32),




10 (Compound 33), and



(Compound 34).

[00202] In other embodiments, the compound has the formula (III)



or a salt or isomer thereof, in which

A_3 is CH or N;

A_4 is CH_2 or NH; and at least one of A_3 and A_4 is N or NH;

5 Z is CH_2 or absent wherein when Z is CH_2 , the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R_1 , R_2 , and R_3 are independently selected from the group consisting of C_{5-20} alkyl, C_{5-20} alkenyl, $-R''MR'$, $-R^*YR''$, $-YR''$, and $-R^*OR''$;

each M is independently selected

10 from $-C(O)O-$, $-OC(O)-$, $-C(O)N(R')-$, $-N(R')C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR')O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

X^1 and X^2 are independently selected from the group consisting of $-CH_2-$, $-(CH_2)_2-$, $-CHR-$, $-CHY-$, $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-C(O)-CH_2-$, $-CH_2-C(O)-$, $-C(O)O-CH_2-$, $-OC(O)-CH_2-$, $-CH_2-C(O)O-$, $-CH_2-OC(O)-$, $-CH(OH)-$, $-C(S)-$, and $-CH(SH)-$;

15 each Y is independently a C_{3-6} carbocycle;


each R^* is independently selected from the group consisting of C_{1-12} alkyl and C_{2-12} alkenyl;

each R is independently selected from the group consisting of C_{1-3} alkyl and a C_{3-6} carbocycle;

20 each R' is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each R'' is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl.

[00203] In some embodiments, the compound is of formula (IIIa):



25

[00204] The compounds of Formula (III) or (IIIa) include one or more of the following features when applicable.

[00205] In some embodiments, Z is CH₂.

[00206] In some embodiments, Z is absent.

5 [00207] In some embodiments, at least one of A₃ and A₄ is N or NH.

[00208] In some embodiments, A₃ is N and A₄ is NH.

[00209] In some embodiments, A₃ is N and A₄ is CH₂.






[00210] In some embodiments, A₃ is CH and A₄ is NH.


10 [00211] In some embodiments, at least one of X¹ and X² is not -CH₂- . For example, in certain embodiments, X¹ is not -CH₂- . In some embodiments, at least one of X¹ and X² is -C(O)-.

[00212] In some embodiments, X² is -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂- , -CH₂-C(O)-, -C(O)O-CH₂- , -OC(O)-CH₂- , -CH₂-C(O)O-, or -CH₂-OC(O)-.


15 [00213] In some embodiments, R₁, R₂, and R₃ are independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. In some embodiments, R₁, R₂, and R₃ are the same. In certain embodiments, R₁, R₂, and R₃ are C₆, C₉, C₁₂, or C₁₄ alkyl. In other embodiments, R₁, R₂, and R₃ are C₁₈ alkenyl. For example, R₁, R₂, and R₃ may be linoleyl.

[00214] In some embodiments, the compound is selected from the group consisting of:





[00215] In another aspect, the disclosure provides a compound according to formula (Ib):



or a salt or isomer thereof, in which

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

Z is CH₂ or absent wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R"MR', -R*YR", -YR", and -R*OR";

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group;

each Y is independently a C₃₋₆ carbocycle;

each R* is independently selected from the group consisting of C₁₋₁₂ alkyl and C₂₋₁₂ alkenyl;

each R is independently selected from the group consisting of C₁₋₃ alkyl and a C₃₋₆ carbocycle;

each R' is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R'' is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl.

5 [00216] In some embodiments, R₁, R₂, R₃, R₄, and R₅ each are independently selected from the group consisting of C₆₋₂₀ alkyl and C₆₋₂₀ alkenyl.

[00217] In some embodiments, R₁ and R₂ are the same. In certain embodiments, R₁, R₂, and R₃ are the same. In some embodiments, R₄ and R₅ are the same. In certain embodiments, R₁, R₂, R₃, R₄, and R₅ are the same.

10 [00218] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is C₉₋₁₂ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ independently is C₉, C₁₂ or C₁₄ alkyl. In certain embodiments, each of R₁, R₂, R₃, R₄, and R₅ is C₉ alkyl.


[00219] In some embodiments, A₆ is N and A₇ is N. In some embodiments, A₆ is CH and A₇ is N.

15 [00220] In some embodiments, X⁴ is -CH₂- and X⁵ is -C(O)-. In some embodiments, X⁴ and X⁵ are -C(O)-.

[00221] In some embodiments, when A₆ is N and A₇ is N, at least one of X⁴ and X⁵ is not -CH₂-, e.g., at least one of X⁴ and X⁵ is -C(O)-. In some embodiments, when A₆ is N and A₇ is N, at least one of R₁, R₂, R₃, R₄, and R₅ is -R''MR'.


20 [00222] In some embodiments, at least one of R₁, R₂, R₃, R₄, and R₅ is not -R''MR'.

[00223] In some embodiments, the compound is




(Compound 67).

[00224] In an embodiment, the compound has the formula (IV)



25 (IV).

[00225] In another aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein R_{1a} is $-(\text{CH}_2)_n^a \text{Q}^a$, where Q^a is selected from a heterocycle, $-\text{OR}^a$, $-\text{O}(\text{CH}_2)_n^a \text{N}(\text{R}^a)_2$, $-\text{C}(\text{O})\text{OR}^a$, $-\text{OC}(\text{O})\text{R}^a$, $-\text{CX}^a_3$, $-\text{CX}^a_2\text{H}$, $-\text{CX}^a\text{H}_2$, $-\text{CN}$, $-\text{N}(\text{R}^a)_2$, $-\text{C}(\text{O})\text{N}(\text{R}^a)_2$, $-\text{N}(\text{R}^a)\text{C}(\text{O})\text{R}^a$, and $-\text{N}(\text{R}^a)\text{S}(\text{O})_2\text{R}^a$ and each n^a is independently selected from 5 1, 2, 3, 4, and 5;

R_{2a} and R_{3a} are each independently selected from the group consisting of C_{3-24} alkyl, C_{3-24} alkenyl, $-\text{R}^{a*}\text{Y}^a\text{R}^{a*}$, $-\text{Y}^a\text{R}^{a*}$, and $-\text{R}^{a*}\text{OR}^{a*}$,

each Y^a is independently a C_{3-6} carbocycle;

each R^{a*} is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

each X^a is independently selected from the group consisting of F, Cl, Br, and I;

each R^a is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H; and

each R^{a*} is selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl;



wherein R_{2a} includes 7 or fewer carbon atoms.

[00226] In some embodiments, Q^a is $-\text{OR}^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-\text{CH}_3$.


[00227] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[00228] In some embodiments, R_{3a} includes 7 or fewer carbon atoms.

[00229] In some embodiments, the compound is selected from the group consisting of:



[00230] In another aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(\text{CH}_2)_{n^a}\text{Q}^a$, where Q^a is selected from a heterocycle, $-\text{OR}^a$, $-\text{O}(\text{CH}_2)_{n^a}\text{N}(\text{R}^a)_2$, $\text{C}(\text{O})\text{OR}^a$, $-\text{OC}(\text{O})\text{R}^a$, $-\text{CX}^a_3$, $-\text{CX}^a_2\text{H}$, $-\text{CX}^a\text{H}_2$, $-\text{CN}$, $-\text{N}(\text{R}^a)_2$, $-\text{C}(\text{O})\text{N}(\text{R}^a)_2$, $-\text{N}(\text{R}^a)\text{C}(\text{O})\text{R}^a$, and $-\text{N}(\text{R}^a)\text{S}(\text{O})_2\text{R}^a$ and each n^a is independently selected from 1, 2, 3, 4, and 5;

5 each X^a is independently selected from the group consisting of F, Cl, Br, and I;

R_{2a} is selected from the group consisting of C_{8-24} alkenyl;

R_{3a} is selected from the group consisting of C_{8-24} alkyl; and

each R is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl,

10 and H.



[00231] In some embodiments, Q^a is $-\text{OR}^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-\text{CH}_3$.

[00232] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[00233] In some embodiments, R_{3a} is an alkyl including 9, 12, 14, or 18 carbon atoms.


[00234] In some embodiments, R_{2a} is C_{18} alkenyl (e.g., linoleyl).

[00235] In some embodiments, the compound is selected from the group consisting of:




20

(Compound 17-5), and



[00236] In a further aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(\text{CH}_2)_{n^a}\text{Q}^a$, where Q^a is selected from a heterocycle, $-\text{OR}^a$, $-\text{O}(\text{CH}_2)_{n^a}\text{N}(\text{R}^a)_2$, $\text{C}(\text{O})\text{OR}^a$, $-\text{OC}(\text{O})\text{R}^a$, $-\text{CX}^a_3$, $-\text{CX}^a_2\text{H}$, $-\text{CX}^a\text{H}_2$, $-\text{CN}$, $-\text{N}(\text{R}^a)_2$, $-\text{C}(\text{O})\text{N}(\text{R}^a)_2$, $-\text{N}(\text{R}^a)\text{C}(\text{O})\text{R}^a$, and $-\text{N}(\text{R}^a)\text{S}(\text{O})_2\text{R}^a$ and each n^a is independently selected from 1, 2, 3, 4, and 5;

5 each X^a is independently selected from the group consisting of F, Cl, Br, and I;

R_{2a} is selected from the group consisting of C_{13-20} alkyl;

R_{3a} is selected from the group consisting of C_{8-20} alkyl; and

each R^a is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl,

10 and H.


[00237] In some embodiments, Q^a is $-\text{OR}^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-\text{CH}_3$.

[00238] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.


15 **[00239]** In some embodiments, R_{2a} and R_{3a} are the same.

[00240] In some embodiments, R_{2a} and/or R_{3a} is C_{14} alkyl.

[00241] In some embodiments, the compound is



[00242] In a further aspect, the disclosure provides a compound having the formula (17-I)



or a salt or isomer thereof, wherein

R_{1a} is $-(\text{CH}_2)_{n^a}\text{Q}^a$, where Q^a is $-\text{OR}^a$, R^a is selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H, and n^a is selected from 1, 2, 3, 4, and 5; and

5 R_{2a} and R_{3a} are each independently selected from the group consisting of C_{8-20} alkenyl, wherein

- iii) R^a is selected from the group consisting of C_{1-3} alkyl and C_{2-3} alkenyl; or
- iv) R_{1a} is $-(\text{CH}_2)_2\text{OH}$, and R_{2a} and R_{3a} each include one or fewer double bonds.

[00243] In some embodiments, R^a is H. In other embodiments, R^a is $-\text{CH}_3$.



10 **[00244]** In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n^a is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[00245] In certain embodiments, R_{1a} is $-(\text{CH}_2)_2\text{OCH}_3$. In other embodiments, R_{1a} is $-(\text{CH}_2)_2\text{OH}$.

15 **[00246]** In some embodiments, R_{2a} is C_{18} alkenyl (e.g., linoleyl). In certain embodiments, R_{3a} is C_{18} alkenyl (e.g., linoleyl).


[00247] In some embodiments, R_{2a} and R_{3a} are the same.

[00248] In some embodiments, the compound is selected from the group consisting of



20

[00249] In another aspect, the disclosure provides a compound of formula (17-I)



or a salt or isomer thereof, wherein

R_1 is $-(CH_2)_n^a Q^a$, where Q^a is selected from a heterocycle, $-OR^a$, $-O(CH_2)_n^a N(R^a)_2$, $-C(O)OR^a$, $-OC(O)R^a$, $-CX^a_3$, $-CX^a_2H$, $-CX^aH_2$, $-CN$, $-N(R^a)_2$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)R^a$, and $-N(R^a)S(O)_2R^a$ and each n^a is independently selected from 1, 2, 3, 4, and 5;

5 each X^a is independently selected from the group consisting of F, Cl, Br, and I;
 R_{2a} is selected from the group consisting of C_{8-12} alkyl;
 R_{3a} is selected from the group consisting of C_{8-20} alkyl; and
each R^a is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.

10 [00250] In some embodiments, Q^a is $-OR^a$. In certain embodiments, R^a is H. In other embodiments, R^a is $-CH_3$.





[00251] In some embodiments, n^a is 1. In other embodiments, n^a is 2. In other embodiments, n is 3. In other embodiments, n^a is 4. In some embodiments, n^a is 5.

[00252] In certain embodiments, Q^a is $-OR^a$ and n^a is selected from 2, 3, and 4.


15 [00253] In some embodiments, R_{2a} is C_9 alkyl. In other embodiments, R_{2a} is C_{12} alkyl.

[00254] In some embodiments, R_{2a} and R_{3a} are the same.

[00255] In some embodiments, the compound is selected from the group consisting of:



[00256] In another aspect, the disclosure provides a compound having the formula (19-I),



(19-I),

or a salt or isomer thereof, wherein

R_{1b} is selected from the group consisting of H, C_{1-5} alkyl, C_{2-5} alkenyl, $-R^{b,*}M^bR^{b,*}$, a C_{3-6} carbocycle, $-(CH_2)_nQ^b$, and $-(CH_2)_nCHQ^bR^b$, where Q^b is selected from a heterocycle, $-OR^b$,

5 $-O(CH_2)_nN(R^b)_2$, $-C(O)OR^b$, $-OC(O)R^b$, $-CX^b_3$, $-CX^b_2H$, $-CX^bH_2$, $-CN$,
 $-N(R^b)_2$, $-C(O)N(R^b)_2$, $-N(R^b)C(O)R^b$, and $-N(R^b)S(O)_2R^b$ and each n is independently selected from 1, 2, 3, 4, and 5;

R_{2b} and R_{3b} are independently selected from the group consisting of C_{1-20} alkyl, C_{2-20} alkenyl, $-R^{b,*}MR^{b,*}$, $-R^{b,*}YR^{b,*}$, $-YR^{b,*}$, and $-R^{b,*}OR^{b,*}$;

10 each M^b is independently selected from the group consisting of $-C(O)O-$, $-OC(O)-$, $-C(O)N(R^b)-$, $-N(R^b)C(O)-$, $-C(O)-$, $-C(S)-$, $-C(S)S-$, $-SC(S)-$, $-CH(OH)-$, $-P(O)(OR^b)O-$, $-S(O)_2-$, an aryl group, and a heteroaryl group;

W is selected from the group consisting of $-CH_2-$, $-CHR^b-$, $-C(O)-$, $-CH(OH)-$, $-C(S)-$, and $-CH(SH)-$;

15 each X^b is independently selected from the group consisting of F, Cl, Br, and I;

each Y^b is independently a C_{3-6} carbocycle;

each $R^{b,*}$ is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

20 each R^b is independently selected from the group consisting of C_{1-3} alkyl, a C_{3-6} carbocycle, C_{2-3} alkenyl, and H;

each $R^{b,*}$ is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each $R^{b,*}$ is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl.

25 [00257] In some embodiments, W is not $-CH_2-$, e.g., W is $-C(O)-$.


[00258] In some embodiments, at least one of R_{2b} and R_{3b} is $-R^{b,*}M^bR^{b,*}$. In certain embodiments, at least one M^b is $-C(O)O-$. In some embodiments, at least one $R^{b,*}$ is C_5 alkyl. In certain embodiments, at least one $R^{b,*}$ is C_5 alkenyl.

[00259] In some embodiments, R_{2b} and/or R_{3b} are selected from the group consisting of C_{1-20} alkyl. For example, R_{2b} and/or R_{3b} may be alkyl groups including 9 or 12 carbon atoms. In other embodiments, R_{2b} and/or R_{3b} are selected from the group consisting of C_{2-20} alkenyl. For example, R_{2b} and/or R_{3b} may be alkenyl groups including 18 carbon atoms (e.g., linoleyl groups). In certain embodiments, R_{2b} and R_{3b} are the same.


5 [00260] In some embodiments, R_{1b} is H, while in other embodiments, R_{1b} is selected from C_{1-5} alkyl. For example, R_{1b} may be C_1 alkyl.

10 [00261] In certain embodiments, R_{1b} is $-(CH_2)_nQ^b$. In such embodiments, Q^b is a heterocycle such as a phenyl group. For example, Q^b may be a phenyl group with one or more substituents, as described herein.


[00262] In certain embodiments, the compound is selected from the group consisting of:




(Compound 19-1),




(Compound 19-2),



(Compound 19-3),




(Compound 19-4), and



(Compound 19-5).

[00263] In other embodiments, lipids are compounds of formula (19-II)



or a salt or isomer thereof, wherein

R_{1b} is selected from the group consisting of C₆₋₂₀ alkyl; and


R_{2b} and R_{3b} are independently selected from the group consisting of C₆₋₂₀ alkenyl.

5 [00264] In particular embodiments, R_{1b} is C₁₂ alkyl.


[00265] In some embodiments, R_{2b} and/or R_{3b} are C₁₈ alkenyl (e.g., linoleyl).

[00266] In certain embodiments, R_{2b} and R_{3b} are both linoleyl.

[00267] In one embodiment, the compound is



10 [00268] In another aspect, lipids may be compounds of formula (20-I),



or a salt or isomer thereof, wherein

R_{1c} is selected from the group consisting of a C₃₋₆

carbocycle, -(CH₂)_{n^c}Q^c, -(CH₂)_{n^c}CHQ^cR^c, -CHQ^cR^c, and -CQ^c(R^c)₂, where Q^c is selected from a

15 heterocycle, -OR^c, -O(CH₂)_{n^c}N(R^c)₂, -C(O)OR^c, -

OC(O)R^c, -CX^c₃, -CX^c₂H, -CX^cH₂, -CN, -N(R^c)₂,

-C(O)N(R^c)₂, -N(R^c)C(O)R^c, and -N(R^c)S(O)₂R^c and each n^c is independently selected from 1, 2, 3, 4, and 5;

R_{2c}, R_{3c}, and R_{4c} are independently selected from the group consisting of C₁₋₂₀ alkyl, C₂₋

20 alkenyl, -R^c*M^cR^c, -R^c*Y^cR^c, -Y^cR^c, and -R^c*OR^c;

each M^c is independently selected from the group consisting

of -C(O)O-, -OC(O)-, -C(O)N(R^c)-, -N(R^c)C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR^c)O-, -S(O)₂-, an aryl group, and a heteroaryl group;

each X^c is independently selected from the group consisting of F, Cl, Br, and I;

each Y^c is independently a C_{3-6} carbocycle;

each R^{c*} is independently selected from the group consisting of C_{1-12} alkyl and C_{1-12} alkenyl;

5 each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

each R^{c*} is independently selected from the group consisting of C_{1-12} alkyl, C_{2-12} alkenyl, and H; and

each R^{c*} is independently selected from the group consisting of C_{3-12} alkyl and C_{3-12} alkenyl,

10 wherein

i) R_{1c} is selected from the group consisting of a C_{3-6} carbocycle, $-(CH_2)_n^cQ^c$, $-(CH_2)_n^cCHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from a heterocycle, $-O(CH_2)_n^cN(R^c)_2$, $-C(O)OR^c$, $-OC(O)R^c$, $-CX^c_3$, $-CX^c_2H$, $-CX^cH_2$, $-C(O)N(R^c)_2$, $-N(R^c)C(O)R^c$, and $-N(R^c)S(O)_2R^c$ and each n^c is independently selected from 1, 2, 3, 4, and 5; and/or

ii) at least one of R_{2c} , R_{3c} , and R_{4c} is $-R^{c*}M^cR^{c*}$.

[00269] In some embodiments, R_{1c} is selected from the group consisting of $-(CH_2)_n^cQ^c$, $-(CH_2)_n^cCHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from a heterocycle, $-O(CH_2)_n^cN(R^c)_2$, $-C(O)OR^c$, $-OC(O)R^c$, $-CX^c_3$, $-CX^c_2H$, $-CX^cH_2$, $-CN$, $-C(O)N(R^c)_2$, $-N(R^c)C(O)R^c$, and $-N(R^c)S(O)_2R^c$ and each n is independently selected from 1, 2, 3, 4, and 5. In certain embodiments, R_{1c} is $-(CH_2)_n^cQ^c$. In some embodiments, n^c is 2. In some embodiments, Q^c is $-C(O)OR^c$, where R^c is, for example, H.

[00270] In some embodiments, at least one of R_{2c} , R_{3c} , and R_{4c} is $-R^{c*}M^cR^{c*}$. For example, R_{2c} , R_{3c} , and/or R_{4c} may be $-R^{c*}M^cR^{c*}$. In some embodiments, at least one M^c is $-C(O)O-$. In certain embodiments, each M^c is $-C(O)O-$. In some embodiments, at least one R^{c*} is C_5 or C_7 alkyl. In certain embodiments, each R^{c*} is C_5 alkyl. In other embodiments, each R^{c*} is C_7 alkyl. In some embodiments, at least one R^c is C_5 , C_7 , or C_9 alkyl. In certain embodiments, each R^c is C_5 alkyl. In other embodiments, each R^c is C_7 alkyl. In other embodiments, each R^c is C_9 alkyl. In some embodiments, R^c is branched.

[00271] In some embodiments, R_{2c} , R_{3c} , and R_{4c} are selected from the group consisting of C_{5-20} alkyl. In certain embodiments, R_{2c} , R_{3c} , and R_{4c} are C_{12} alkyl.


[00272] In some embodiments, R_{2c} is selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. For example, R_{2c} may be C₁₂ alkyl.

[00273] In some embodiments, R_{3c} is selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. For example, R_{3c} may be C₆, C₉, or C₁₂ alkyl.


5 **[00274]** In some embodiments, R_{4c} is selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl. For example, R_{4c} may be C₆, C₉, or C₁₂ alkyl.

[00275] In some embodiments, R_{3c} and R_{4c} are the same.


[00276] In some embodiments, the compound is selected from the group consisting of:




(Compound 20-1),




(Compound 20-2),




(Compound 20-3),




(Compound 20-4),



(Compound 20-5),




(Compound 20-6),




(Compound 20-7),


15



(Compound 20-8),




(Compound 20-9), and



(Compound 20-10).

[00277] In other embodiments, the lipid is a compound according to formula (20-I)



(20-I),

or a salt or isomer thereof, wherein

R_{1c} is selected from the group consisting of $-(CH_2)_nQ^c$, $-(CH_2)_nCHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from $-OR^c$, $-CN$, and $-N(R^c)_2$, and n^c is selected from 1, 2, 3, 4, and 5;

10 R_{2c} and R_{3c} are independently selected from the group consisting of C_{6-20} alkyl and C_{6-20} alkenyl;

R_{4c} is selected from the group consisting of C_{13-20} alkyl and C_{5-20} alkenyl; and each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.

15 [00278] In some embodiments, R_{3c} is C_{14} alkyl.

[00279] In some embodiments, R_{4c} is C_{14} alkyl.

[00280] In some embodiments, R_{3c} is C_{18} alkenyl. For example, R_{3c} may be linoleyl.

[00281] In some embodiments, R_{4c} is C_{18} alkenyl. For example, R_{4c} may be linoleyl.


[00282] In some embodiments, R_{2c} is C_{12} alkyl. In other embodiments, R_{2c} is C_{14} alkyl.

20 In some embodiments, R_{2c} is C_{18} alkenyl. For example, R_{2c} may be linoleyl.


[00283] In some embodiments, R_{3c} and R_{4c} are the same.

[00284] In some embodiments, R_{1c} is $-(CH_2)_{n^c}Q^c$. In some embodiments, Q^c is $-OR^c$. For example, Q^c may be $-OH$. In some embodiments, n^c is 2 or 3.


[00285] In some embodiments, the compound is selected from the group consisting of:




(Compound 20-11),



(Compound 20-12),




(Compound 20-13), and



(Compound 20-14).

[00286] In other embodiments, the lipid is a compound having formula (20-I)



10 or a salt or isomer thereof, wherein

R_{1c} is selected from the group consisting of $-(CH_2)_{n^c}Q^c$, $-(CH_2)_{n^c}CHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from $-OR^c$, $-CN$, and $-N(R^c)_2$, and n^c is selected from 1, 2, 3, 4, and 5;

15 R_{2c} , R_{3c} , and R_{4c} are independently selected from the group consisting of C_{6-20} alkyl and C_{6-20} alkenyl; and

each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H;

wherein

i) R_{2c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl, and/or

20 ii) R_{3c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl.

[00287] In some embodiments, R_{2c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl. For example, R_{2c} may be C_6 or C_9 alkyl.

[00288] In some embodiments, R_{3c} is selected from the group consisting of C_{1-11} alkyl and C_{2-5} alkenyl. For example, R_{3c} may be C_6 or C_9 alkyl.


[00289] In some embodiments, R_{3c} is C_{12} alkyl.

[00290] In some embodiments, R_{2c} is C_{12} alkyl.


5 [00291] In some embodiments, R_{4c} is C_6 , C_9 , or C_{12} alkyl.

[00292] In some embodiments, R_{1c} is $-(CH_2)_n^cQ^c$. In certain embodiments, Q^c is $-OR^c$. In some embodiments, R^c is H. In some embodiments, n^c is 2 or 3.


[00293] In some embodiments, the compound is selected from the group consisting of:




(Compound 20-15),



(Compound 20-16),




(Compound 20-17), and



(Compound 20-18).

[00294] In other embodiments, the lipid is a compound according to formula (20-I)



(20-I),

15 or a salt or isomer thereof, wherein

R_{1c} is selected from the group consisting of $-(CH_2)_n^cQ^c$, $-(CH_2)_n^cCHQ^cR^c$, $-CHQ^cR^c$, and $-CQ^c(R^c)_2$, where Q^c is selected from $-OR^c$, $-CN$, and $-N(R^c)_2$, and n^c is selected from 1, 2, 3, 4, and 5;

R_{2c} is selected from the group consisting of H, C_{12-20} alkyl, and C_{6-20} alkenyl;

20 R_{3c} and R_{4c} are C_{12} alkyl; and


each R^c is independently selected from the group consisting of C_{1-3} alkyl, C_{2-3} alkenyl, and H.

[00295] In some embodiments, R_{2c} is H. In other embodiments, R_{2c} is C_{12} alkyl or alkenyl. In some embodiments, R_{2c} is C_{14} alkyl. In other embodiments, R_{2c} is C_{18} alkenyl. For example, R_{2c} may be linoleyl.


[00296] In some embodiments, R_{1c} is $-(CH_2)_{n^c}Q^c$. In certain embodiments, Q^c is $-OR^c$.

5 For example, Q^c may be OH. In some embodiments, n^c is 2, 3, or 4.


[00297] In some embodiments, the compound is selected from the group consisting of:




(Compound 20-19),




(Compound 20-20),




(Compound 20-21),




(Compound 20-22),



(Compound 20-23),




(Compound 20-24), and



(Compound 20-25).

[0001] In another aspect, lipids may be compounds of formula (21-I),



(21-I),

or a salt or isomer thereof, wherein

R_{1d} and R_{2d} are independently selected from the group consisting of H, C₁₋₅ alkyl, and C₂₋₅ alkenyl;

n^d is selected from 1, 2, 3, 4, and 5;

k is selected from 0, 1, 2, and 3;

5 R_{3d} and R_{4d} are independently selected from the group consisting of C₁₋₂₀ alkyl, C₂₋₂₀ alkenyl, - $R^{d*}M^dR^d$, - $R^{d*}Y^dR^{d*}$, - Y^dR^{d*} , and - $R^{d*}OR^{d*}$;

each M^d is independently selected from the group consisting of -C(O)O-, -OC(O)-, -C(O)N(R^{d*})-, -N(R^{d*})C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR^{d*})O-, and -S(O)₂-, or is absent;

10 each R^{d*} is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; each Y^d is independently a C₃₋₆ carbocycle;

each R^{d*} is independently selected from the group consisting of C₁₋₁₂ alkyl and C₁₋₁₂ alkenyl;

15 each R^{d*} is independently selected from the group consisting of C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, and H; and

each R^{d*} is independently selected from the group consisting of C₃₋₁₂ alkyl and C₃₋₁₂ alkenyl,

wherein R_{3d} and R_{4d} are bound to either i) the same carbon atom or ii) adjacent carbon atoms.

20 [0002] In some embodiments, R_{3d} and R_{4d} are bound to the same carbon atom. For example, R_{3d} and R_{4d} may be bound to a carbon atom adjacent to C*. In certain embodiments, R_{3d} and R_{4d} are not bound to a carbon atom adjacent to C*.

[0003] In other embodiments, R_{3d} and R_{4d} are bound to adjacent carbon atoms. In certain embodiments, one or both of R_{3d} and R_{4d} are bound to carbon atoms adjacent to C*.

25 [0004] In some embodiments, k is 0. In other embodiments, k is 1, 2, or 3.







[0005] In certain embodiments, M^d is absent. In other embodiments, M^d is selected from the group consisting of -C(O)O-, -OC(O)-, -C(O)N(R^{d*})-, -N(R^{d*})C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR^{d*})O-, and -S(O)₂-. In particular such embodiments, M^d is -C(O)O-.

30 [0006] In some embodiments, n^d is 1, 2, or 3.

[0007] In some embodiments, R_{1d} and/or R_{2d} are selected from C₁₋₅ alkyl. In certain embodiments, R_{1d} and/or R_{2d} are C₁ alkyl.

[0008] In certain embodiments, R_{3d} and/or R_{4d} are selected from C₂₋₂₀ alkenyl. In certain embodiments, R_{3d} and/or R_{4d} are alkenyl groups including 17, 18, or 19 carbon atoms. For example, R_{3d} and/or R_{4d} may be C₁₈ alkenyl groups (e.g., linoleyl).

[0009] In certain embodiments, the compound is selected from the group consisting of:



[00298] As used herein, the term “alkyl” or “alkyl group” means a linear or branched, saturated hydrocarbon including one or more carbon atoms (e.g., one, two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, or more carbon atoms), which is optionally substituted. For example, the notation “C₁₋₂₄ alkyl” means an optionally substituted linear or branched, saturated hydrocarbon including 1-24 carbon atoms. An alkyl group described herein refers to both unsubstituted and substituted alkyl group unless otherwise specified.

[00299] As used herein, the term “alkenyl” or “alkenyl group” means a linear or branched hydrocarbon including two or more carbon atoms (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, sixteen, seventeen, eighteen, nineteen, twenty, or more carbon atoms) and at least one double bond, which is optionally substituted. The notation “C₂₋₂₄ alkenyl” means an optionally substituted linear or branched hydrocarbon including 2 to 24 carbon atoms and at least one carbon-carbon double bond. An alkenyl group may include one, two, three, four, or more carbon-carbon double bonds. For example, C₁₈

alkenyl may include one or more double bonds. A C₁₈ alkenyl group including two double bonds may be a linoleyl group. An alkenyl group described herein refers to both unsubstituted and substituted unless otherwise specified.

[00300] As used herein, the term “carbocycle” or “carbocyclic group” means an 5 optionally substituted mono- or multi-cyclic system including one or more rings of carbon atoms. Rings may be three, four, five, six, seven, eight, nine, ten, eleven, or twelve membered rings. The notation “C₃₋₆ carbocycle” means a carbocycle including a single ring having 3-6 carbon atoms. Carbocycles may include one or more carbon-carbon double or triple bonds and may be non-aromatic or aromatic (e.g., cycloalkyl or aryl groups). Examples of carbocycles 10 include cyclopropyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, and 1,2-dihydronaphthyl groups. Carbocycles described herein refers to both unsubstituted and substituted carbocycles unless otherwise specified. The term “cycloalkyl” as used herein means a non-aromatic carbocycle and may or may not include any double or triple bond. Unless otherwise specified, carbocycles described herein refers to both unsubstituted and substituted carbocycle groups, i.e., 15 optionally substituted carbocycles.

[00301] As used herein, the term “heterocycle” or “heterocyclic group” means an 20 optionally substituted mono- or multi-cyclic system including one or more rings, where at least one ring includes at least one heteroatom. Heteroatoms may be, for example, nitrogen, oxygen, or sulfur atoms. Rings may be three, four, five, six, seven, eight, nine, ten, eleven, or twelve membered rings. Heterocycles may include one or more double or triple bonds and may be non-aromatic or aromatic. Examples of heterocycles include imidazolyl, imidazolidinyl, oxazolyl, 25 oxazolidinyl, thiazolyl, thiazolidinyl, pyrazolidinyl, pyrazolyl, isoxazolidinyl, isoxazolyl, isothiazolidinyl, isothiazolyl, morpholinyl, pyrrolyl, pyrrolidinyl, furyl, tetrahydrofuryl, thiophenyl, pyridinyl, piperidinyl, quinolyl, and isoquinolyl groups. Heterocycles may be optionally substituted.

[00302] As used herein, a “biodegradable group” is a group that may facilitate faster 30 metabolism of a lipid in a mammalian entity. A biodegradable group may be selected from the group consisting of, but is not limited to, -C(O)O-, -OC(O)-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-, -CH(OH)-, -P(O)(OR')O-, -S(O)₂-, an aryl group, and a heteroaryl group. As used herein, an “aryl group” is a carbocyclic group including one or more aromatic rings. Examples of aryl groups include phenyl and naphthyl groups. As used herein, a “heteroaryl group” is a heterocyclic group including one or more aromatic rings. Examples of heteroaryl groups include pyrrolyl, furyl, thiophenyl, imidazolyl, oxazolyl, and thiazolyl. Aryl

and heteroaryl groups may be optionally substituted. For example, each M, M^b, M^c, or M^d can be independently selected from the non-limiting group consisting of phenyl, oxazole, and thiazole. In the formulae above, each M, M^b, M^c, or M^d can be independently selected from the list of biodegradable groups above.

5 [00303] Alkyl, alkenyl, and cyclyl (e.g., carbocyclyl and heterocyclyl) groups may be optionally substituted unless otherwise specified. Optional substituents may be selected from the group consisting of, but are not limited to, a halogen atom (e.g., a chloride, bromide, fluoride, or iodide group), a carboxylic acid (e.g., -C(O)OH), an alcohol (e.g., a hydroxyl, -OH), an ester (e.g., -C(O)OR or -OC(O)R), an aldehyde (e.g., -C(O)H), a carbonyl (e.g., -C(O)R, alternatively represented by C=O), an acyl halide (e.g., -C(O)X, in which X is a halide selected from bromide, fluoride, chloride, and iodide), a carbonate (e.g., -OC(O)OR), an alkoxy (e.g., -OR), an acetal (e.g., -C(OR)₂R[”]“, in which each OR are alkoxy groups that can be the same or different and R[”]“ is an alkyl or alkenyl group), a phosphate (e.g., P(O)₄³⁻), a thiol (e.g., -SH), a sulfoxide (e.g., -S(O)R), a sulfinic acid (e.g., -S(O)OH), a sulfonic acid (e.g., -S(O)₂OH), a thial (e.g., -C(S)H), a sulfate (e.g., S(O)₄²⁻), a sulfonyl (e.g., -S(O)₂-), an amide (e.g., -C(O)NR₂, or -N(R)C(O)R), an azido (e.g., -N₃), a nitro (e.g., -NO₂), a cyano (e.g., -CN), an isocyano (e.g., -NC), an acyloxy (e.g., -OC(O)R), an amino (e.g., -NR₂, -NRH, or -NH₂), a carbamoyl (e.g., -OC(O)NR₂, -OC(O)NRH, or -OC(O)NH₂), a sulfonamide (e.g., -S(O)₂NR₂, -S(O)₂NRH, -S(O)₂NH₂, -N(R)S(O)₂R, -N(H)S(O)₂R, -N(R)S(O)₂H, or -N(H)S(O)₂H), a cyclyl (e.g., carbocyclyl or heterocyclyl) group, an alkyl group, and an alkenyl group. In any of the preceding, R is an alkyl or alkenyl group, as defined herein. In some embodiments, the substituent groups themselves may be further substituted with, for example, one, two, three, four, five, or six substituents as defined herein. For example, a C₅₋₂₀ alkyl group may be further substituted with one, two, three, four, five, six, or more substituents as described herein.

10

15

20

25

[00304] An amine moiety of a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) may be protonated at a physiological pH. Thus, a lipid may have a positive or partial positive charge at physiological pH. Such lipids may be referred to as cationic or ionizable (amino)lipids. Lipids may be zwitterionic, i.e., neutral molecules having both a positive and a negative charge.

30

Nanoparticle compositions

[00305] The disclosure also features nanoparticle compositions comprising a lipid component comprising a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) as described herein. In some embodiments, the largest dimension of a nanoparticle composition is 1 μ m or shorter (e.g., 1

5 μ m, 900 nm, 800 nm, 700 nm, 600 nm, 500 nm, 400 nm, 300 nm, 200 nm, 175 nm, 150 nm, 125 nm, 100 nm, 75 nm, 50 nm, or shorter), e.g., when measured by dynamic light scattering (DLS), transmission electron microscopy, scanning electron microscopy, or another method.


Nanoparticle compositions include, for example, lipid nanoparticles (LNPs), liposomes, lipid vesicles, and lipoplexes. In some embodiments, nanoparticle compositions are vesicles 10 including one or more lipid bilayers. In certain embodiments, a nanoparticle composition includes two or more concentric bilayers separated by aqueous compartments. Lipid bilayers may be functionalized and/or crosslinked to one another. Lipid bilayers may include one or more ligands, proteins, or channels.

[00306] Nanoparticle compositions comprise a lipid component including at least one 15 lipid, such as a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), as described herein. For example, in some embodiments, a nanoparticle composition may include a lipid component including one of Compounds 1 through 88, Compounds 17-1 through 17-13, Compounds 19-1 through 19-6, Compounds 20-1 through 20-25 and Compounds 21-1 through 21-6. Nanoparticle compositions 20 may also include a variety of other components. For example, the lipid component of a nanoparticle composition may include one or more other lipids in addition to a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I).

25 *Cationic/ionizable lipids*


[00110] A nanoparticle composition may include one or more cationic and/or ionizable lipids 30 (e.g., lipids that may have a positive or partial positive charge at physiological pH) in addition to a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I). Cationic and/or ionizable lipids may be selected from the non-limiting group consisting of 3-(didodecylamino)-N1,N1,4-tridodecyl-1-piperazineethanamine (KL10), N1-[2-(didodecylamino)ethyl]-N1,N4,N4-tridodecyl-1,4-piperazinediethanamine (KL22), 14,25-ditridodecyl-15,18,21,24-tetraaza-octatriacontane (KL25),

1,2-dilinoleyloxy-N,N-dimethylaminopropane (DLinDMA),
 2,2-dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane (DLin-K-DMA),
 heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate (DLin-MC3-DMA),
 2,2-dilinoleyl-4-(2-dimethylaminoethyl)-[1,3]-dioxolane (DLin-KC2-DMA),
 5 1,2-dioleyloxy-N,N-dimethylaminopropane (DODMA),
 2-({8-[(3β) -cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[$(9Z,12Z)$ -octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA),
 (2R)-2-({8-[(3β) -cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[$(9Z,12Z)$ -octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2R)),
 10 (2S)-2-({8-[(3β) -cholest-5-en-3-yloxy]octyl}oxy)-N,N-dimethyl-3-[$(9Z,12Z)$ -octadeca-9,12-dien-1-yloxy]propan-1-amine (Octyl-CLinDMA (2S)),



(i.e., (12Z, 15Z)-N,N-dimethyl-2-

nonylnheicos-12,15-dien-1-amine), and



(i.e., N,N-dimethyl-1-[(1S,2R)-2-

15 octylcyclopropyl]heptadecan-8-amine).

[00307] In addition to these, a cationic lipid may also be a lipid including a cyclic amine group. Additional cationic and/or ionizable lipids that are suitable for the formulations and methods disclosed herein include those described in WO2015199952, WO2016176330, and WO2015011633, the entire contents of each of which are hereby incorporated by reference in 20 their entireties.

PEG lipids

[00308] The lipid component of a nanoparticle composition may include one or more PEG or PEG-modified lipids. Such species may be alternately referred to as PEGylated lipids.
 25 A PEG lipid is a lipid modified with polyethylene glycol. A PEG lipid may be selected from the non-limiting group consisting of PEG-modified phosphatidylethanolamines, PEG-modified phosphatidic acids, PEG-modified ceramides, PEG-modified dialkylamines, PEG-modified diacylglycerols, PEG-modified dialkylglycerols, and mixtures thereof. For example, a PEG


lipid may be PEG-c-DOMG, PEG-DMG, PEG-DLPE, PEG-DMPE, PEG-DPPC, or a PEG-DSPE lipid.

Structural lipids

5 [00309] The lipid component of a nanoparticle composition may include one or more structural lipids. Structural lipids can be selected from the group consisting of, but are not limited to, cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, tomatine, ursolic acid, alpha-tocopherol, and mixtures thereof. In certain embodiments, the structural lipid is cholesterol. In some embodiments, the structural lipid 10 includes cholesterol and a corticosteroid (such as prednisolone, dexamethasone, prednisone, and hydrocortisone), or a combination thereof.

Phospholipids

15 [00310] The lipid component of a nanoparticle composition may include one or more phospholipids, such as one or more (poly)unsaturated lipids. Phospholipids may assemble into one or more lipid bilayers. In general, phospholipids may include a phospholipid moiety and one or more fatty acid moieties. For example, a phospholipid may be a lipid according to formula (V)



20 in which R_p represents a phospholipid moiety and R₁ and R₂ represent fatty acid moieties with or without unsaturation that may be the same or different. A phospholipid moiety may be selected from the non-limiting group consisting of phosphatidyl choline, phosphatidyl ethanolamine, phosphatidyl glycerol, phosphatidyl serine, phosphatidic acid, 2-lysophosphatidyl choline, and a sphingomyelin. A fatty acid moiety may be selected from the non-limiting group consisting of 25 lauric acid, myristic acid, myristoleic acid, palmitic acid, palmitoleic acid, stearic acid, oleic acid, linoleic acid, alpha-linolenic acid, erucic acid, phytanic acid, arachidic acid, arachidonic acid, eicosapentaenoic acid, behenic acid, docosapentaenoic acid, and docosahexaenoic acid. Non-natural species including natural species with modifications and substitutions including

branching, oxidation, cyclization, and alkynes are also contemplated. For example, a phospholipid may be functionalized with or cross-linked to one or more alkynes (e.g., an alkenyl group in which one or more double bonds is replaced with a triple bond). Under appropriate reaction conditions, an alkyne group may undergo a copper-catalyzed cycloaddition upon 5 exposure to an azide. Such reactions may be useful in functionalizing a lipid bilayer of a nanoparticle composition to facilitate membrane permeation or cellular recognition or in conjugating a nanoparticle composition to a useful component such as a targeting or imaging moiety (e.g., a dye).

[00311] Phospholipids useful in the compositions and methods described herein may be 10 selected from the non-limiting group consisting of 1,2-distearoyl-*sn*-glycero-3-phosphocholine (DSPC), 1,2-dioleoyl-*sn*-glycero-3-phosphoethanolamine (DOPE), 1,2-dilinoleoyl-*sn*-glycero-3-phosphocholine (DLPC), 1,2-dimyristoyl-*sn*-glycero-phosphocholine (DMPC), 1,2-dioleoyl-*sn*-glycero-3-phosphocholine (DOPC), 1,2-dipalmitoyl-*sn*-glycero-3-phosphocholine (DPPC), 15 1,2-diundecanoyl-*sn*-glycero-phosphocholine (DUPC), 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC), 1,2-di-O-octadecenyl-*sn*-glycero-3-phosphocholine (18:0 Diether PC), 1-oleoyl-2-cholesterylhemisuccinoyl-*sn*-glycero-3-phosphocholine (OChemSPC), 1-hexadecyl-*sn*-glycero-3-phosphocholine (C16 Lyso PC), 20 1,2-dilinolenoyl-*sn*-glycero-3-phosphocholine, 1,2-diarachidonoyl-*sn*-glycero-3-phosphocholine, 1,2-didocosahexaenoyl-*sn*-glycero-3-phosphocholine, 1,2-diphytanoyl-*sn*-glycero-3-phosphoethanolamine (ME 16.0 PE), 1,2-distearoyl-*sn*-glycero-3-phosphoethanolamine, 25 1,2-dilinoleoyl-*sn*-glycero-3-phosphoethanolamine, 1,2-dilinolenoyl-*sn*-glycero-3-phosphoethanolamine, 1,2-diarachidonoyl-*sn*-glycero-3-phosphoethanolamine, 1,2-didocosahexaenoyl-*sn*-glycero-3-phosphoethanolamine, 1,2-dioleoyl-*sn*-glycero-3-phospho-rac-(1-glycerol) sodium salt (DOPG), and sphingomyelin. 30 In certain embodiments, a nanoparticle composition includes DSPC. In certain embodiments, a nanoparticle composition includes DOPE. In some embodiments, a nanoparticle composition includes both DSPC and DOPE.

Adjuvants

[00312] In some embodiments, a nanoparticle composition that includes one or more lipids described herein may further include one or more adjuvants, e.g., Glucopyranosyl Lipid Adjuvant (GLA), CpG oligodeoxynucleotides (e.g., Class A or B), poly(I:C), aluminum hydroxide, and Pam3CSK4.

Therapeutic agents

[00313] Nanoparticle compositions may include one or more therapeutic and/or prophylactic agents. The disclosure features methods of delivering a therapeutic and/or prophylactic agent to a mammalian cell or organ, producing a polypeptide of interest in a mammalian cell, and treating a disease or disorder in a mammal in need thereof comprising administering to a mammal and/or contacting a mammalian cell with a nanoparticle composition including a therapeutic and/or prophylactic agent.

[00314] Therapeutic and/or prophylactic agents include biologically active substances and are alternately referred to as “active agents.” A therapeutic and/or prophylactic agent may be a substance that, once delivered to a cell or organ, brings about a desirable change in the cell, organ, or other bodily tissue or system. Such species may be useful in the treatment of one or more diseases, disorders, or conditions. In some embodiments, a therapeutic and/or prophylactic agent is a small molecule drug useful in the treatment of a particular disease, disorder, or condition. Examples of drugs useful in the nanoparticle compositions described herein include, but are not limited to, antineoplastic agents (e.g., vincristine, doxorubicin, mitoxantrone, camptothecin, cisplatin, bleomycin, cyclophosphamide, methotrexate, and streptozotocin), antitumor agents (e.g., actinomycin D, vincristine, vinblastine, cytosine arabinoside, anthracyclines, alkylating agents, platinum compounds, antimetabolites, and nucleoside analogs, such as methotrexate and purine and pyrimidine analogs), anti-infective agents, local anesthetics (e.g., dibucaine and chlorpromazine), beta-adrenergic blockers (e.g., propranolol, timolol, and labetalol), antihypertensive agents (e.g., clonidine and hydralazine), anti-depressants (e.g., imipramine, amitriptyline, and doxepin), anti-convulsants (e.g., phenytoin), antihistamines (e.g., diphenhydramine, chlorpheniramine, and promethazine), antibiotic/antibacterial agents (e.g., gentamycin, ciprofloxacin, and cefoxitin), antifungal agents (e.g., miconazole, terconazole, econazole, isoconazole, butaconazole, clotrimazole, itraconazole, nystatin, naftifine, and amphotericin B), antiparasitic agents, hormones, hormone antagonists, immunomodulators, neurotransmitter antagonists, antiglaucoma agents, vitamins, narcotics, and imaging agents.

[00315] In some embodiments, a therapeutic and/or prophylactic agent is a cytotoxin, a radioactive ion, a chemotherapeutic, a vaccine, a compound that elicits an immune response, and/or another therapeutic and/or prophylactic agent. A cytotoxin or cytotoxic agent includes any agent that may be detrimental to cells. Examples include, but are not limited to, taxol, 5 cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicine, doxorubicin, daunorubicin, dihydroxyanthracinedione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, maytansinoids, e.g., maytansinol, rachelmycin (CC-1065), and analogs or homologs thereof. Radioactive ions include, but are not limited to 10 iodine (e.g., iodine 125 or iodine 131), strontium 89, phosphorous, palladium, cesium, iridium, phosphate, cobalt, yttrium 90, samarium 153, and praseodymium. Vaccines include compounds and preparations that are capable of providing immunity against one or more conditions related to infectious diseases such as influenza, measles, human papillomavirus (HPV), rabies, meningitis, whooping cough, tetanus, plague, hepatitis, and tuberculosis and can include 15 mRNAs encoding infectious disease derived antigens and/or epitopes. Vaccines also include compounds and preparations that direct an immune response against cancer cells and can include mRNAs encoding tumor cell derived antigens, epitopes, and/or neoepitopes. Compounds eliciting immune responses may include vaccines, corticosteroids (e.g., dexamethasone), and other species. In some embodiments, a vaccine and/or a compound capable of eliciting an 20 immune response is administered intramuscularly via a composition including a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I). Other therapeutic and/or prophylactic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil, dacarbazine), alkylating agents (e.g., mechlorethamine, thiotepa chlorambucil, 25 rachelmycin (CC-1065), melphalan, carmustine (BSNU), lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine, 30 vinblastine, taxol and maytansinoids).

[00316] In other embodiments, a therapeutic and/or prophylactic agent is a protein.

Therapeutic proteins useful in the nanoparticles of the disclosure include, but are not limited to, gentamycin, amikacin, insulin, erythropoietin (EPO), granulocyte-colony stimulating factor (G-

CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), Factor VIR, luteinizing hormone-releasing hormone (LHRH) analogs, interferons, heparin, Hepatitis B surface antigen, typhoid vaccine, and cholera vaccine.

5 *Polynucleotides and Nucleic Acids*

[00317] In some embodiments, a therapeutic and/or prophylactic agent is a polynucleotide or nucleic acid (e.g., ribonucleic acid or deoxyribonucleic acid). The term “polynucleotide,” in its broadest sense, includes any compound and/or substance that is or can be incorporated into an oligonucleotide chain. Exemplary polynucleotides for use in accordance with the present disclosure include, but are not limited to, one or more of deoxyribonucleic acid (DNA), ribonucleic acid (RNA) including messenger mRNA (mRNA), hybrids thereof, RNAi-inducing agents, RNAi agents, siRNAs, shRNAs, miRNAs, antisense RNAs, ribozymes, catalytic DNA, RNAs that induce triple helix formation, aptamers, vectors, etc. In certain embodiments, a therapeutic and/or prophylactic agent is an RNA. RNAs useful in the compositions and methods described herein can be selected from the group consisting of, but are not limited to, shortmers, antagonists, antisense, ribozymes, small interfering RNA (siRNA), asymmetrical interfering RNA (aiRNA), microRNA (miRNA), Dicer-substrate RNA (dsRNA), small hairpin RNA (shRNA), transfer RNA (tRNA), messenger RNA (mRNA), and mixtures thereof. In certain embodiments, the RNA is an mRNA.

10 [00318] In certain embodiments, a therapeutic and/or prophylactic agent is an mRNA. An mRNA may encode any polypeptide of interest, including any naturally or non-naturally occurring or otherwise modified polypeptide. A polypeptide encoded by an mRNA may be of any size and may have any secondary structure or activity. In some embodiments, a polypeptide encoded by an mRNA may have a therapeutic effect when expressed in a cell.

15 [00319] In other embodiments, a therapeutic and/or prophylactic agent is an siRNA. An siRNA may be capable of selectively knocking down or down regulating expression of a gene of interest. For example, an siRNA could be selected to silence a gene associated with a particular disease, disorder, or condition upon administration to a subject in need thereof of a nanoparticle composition including the siRNA. An siRNA may comprise a sequence that is complementary to an mRNA sequence that encodes a gene or protein of interest. In some embodiments, the siRNA may be an immunomodulatory siRNA.

20 [00320] In some embodiments, a therapeutic and/or prophylactic agent is an shRNA or a vector or plasmid encoding the same. An shRNA may be produced inside a target cell upon

delivery of an appropriate construct to the nucleus. Constructs and mechanisms relating to shRNA are well known in the relevant arts.

[00321] Nucleic acids and polynucleotides useful in or suitable for the compounds and methods of the disclosure typically include a first region of linked nucleosides encoding a polypeptide of interest (e.g., a coding region), a first flanking region located at the 5'-terminus of the first region (e.g., a 5'-UTR), a second flanking region located at the 3'-terminus of the first region (e.g., a 3'-UTR), at least one 5'-cap region, and a 3'-stabilizing region. In some embodiments, a nucleic acid or polynucleotide further includes a poly-A region or a Kozak sequence (e.g., in the 5'-UTR). In some cases, polynucleotides may contain one or more 10 intronic nucleotide sequences capable of being excised from the polynucleotide. In some embodiments, a polynucleotide or nucleic acid (e.g., an mRNA) may include a 5' cap structure, a chain terminating nucleotide, a stem loop, a polyA sequence, and/or a polyadenylation signal. Any one of the regions of a nucleic acid may include one or more alternative components (e.g., an alternative nucleoside). For example, the 3'-stabilizing region may contain an alternative 15 nucleoside such as an L-nucleoside, an inverted thymidine, or a 2'-O-methyl nucleoside and/or the coding region, 5'-UTR, 3'-UTR, or cap region may include an alternative nucleoside such as a 5-substituted uridine (e.g., 5-methoxyuridine), a 1-substituted pseudouridine (e.g., 1-methyl-pseudouridine or 1-ethyl-pseudouridine), and/or a 5-substituted cytidine (e.g., 5-methyl-cytidine).

[00322] Generally, the shortest length of a polynucleotide can be the length of the polynucleotide sequence that is sufficient to encode for a dipeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for a tripeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for a tetrapeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for a pentapeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for a hexapeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for a heptapeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for an octapeptide. In another embodiment, the length of the polynucleotide sequence is sufficient to encode for a nonapeptide. In another embodiment, the 25 length of the polynucleotide sequence is sufficient to encode for a decapeptide.

[00323] Examples of dipeptides that the alternative polynucleotide sequences can encode for include, but are not limited to, carnosine and anserine.

[00324] In some cases, a polynucleotide is greater than 30 nucleotides in length. In another embodiment, the polynucleotide molecule is greater than 35 nucleotides in length. In another embodiment, the length is at least 40 nucleotides. In another embodiment, the length is at least 45 nucleotides. In another embodiment, the length is at least 55 nucleotides. In another embodiment, the length is at least 50 nucleotides. In another embodiment, the length is at least 60 nucleotides. In another embodiment, the length is at least 80 nucleotides. In another embodiment, the length is at least 90 nucleotides. In another embodiment, the length is at least 100 nucleotides. In another embodiment, the length is at least 120 nucleotides. In another embodiment, the length is at least 140 nucleotides. In another embodiment, the length is at least 160 nucleotides. In another embodiment, the length is at least 180 nucleotides. In another embodiment, the length is at least 200 nucleotides. In another embodiment, the length is at least 250 nucleotides. In another embodiment, the length is at least 300 nucleotides. In another embodiment, the length is at least 350 nucleotides. In another embodiment, the length is at least 400 nucleotides. In another embodiment, the length is at least 450 nucleotides. In another embodiment, the length is at least 500 nucleotides. In another embodiment, the length is at least 600 nucleotides. In another embodiment, the length is at least 700 nucleotides. In another embodiment, the length is at least 800 nucleotides. In another embodiment, the length is at least 900 nucleotides. In another embodiment, the length is at least 1000 nucleotides. In another embodiment, the length is at least 1100 nucleotides. In another embodiment, the length is at least 1200 nucleotides. In another embodiment, the length is at least 1300 nucleotides. In another embodiment, the length is at least 1400 nucleotides. In another embodiment, the length is at least 1500 nucleotides. In another embodiment, the length is at least 1600 nucleotides. In another embodiment, the length is at least 1800 nucleotides. In another embodiment, the length is at least 2000 nucleotides. In another embodiment, the length is at least 2500 nucleotides. In another embodiment, the length is at least 3000 nucleotides. In another embodiment, the length is at least 4000 nucleotides. In another embodiment, the length is at least 5000 nucleotides, or greater than 5000 nucleotides.

[00325] Nucleic acids and polynucleotides may include one or more naturally occurring components, including any of the canonical nucleotides A (adenosine), G (guanosine), C (cytosine), U (uridine), or T (thymidine). In one embodiment, all or substantially all of the nucleotides comprising (a) the 5'-UTR, (b) the open reading frame (ORF), (c) the 3'-UTR, (d) the poly A tail, and any combination of (a, b, c, or d above) comprise naturally occurring canonical nucleotides A (adenosine), G (guanosine), C (cytosine), U (uridine), or T (thymidine).

[00326] Nucleic acids and polynucleotides may include one or more alternative components, as described herein, which impart useful properties including increased stability and/or the lack of a substantial induction of the innate immune response of a cell into which the polynucleotide is introduced. For example, an alternative polynucleotide or nucleic acid

5 exhibits reduced degradation in a cell into which the polynucleotide or nucleic acid is introduced, relative to a corresponding unaltered polynucleotide or nucleic acid. These alternative species may enhance the efficiency of protein production, intracellular retention of the polynucleotides, and/or viability of contacted cells, as well as possess reduced immunogenicity.

10 **[00327]** Polynucleotides and nucleic acids may be naturally or non-naturally occurring. Polynucleotides and nucleic acids may include one or more modified (e.g., altered or alternative) nucleobases, nucleosides, nucleotides, or combinations thereof. The nucleic acids and polynucleotides useful in the nanoparticle compositions described herein can include any useful modification or alteration, such as to the nucleobase, the sugar, or the internucleoside linkage

15 (e.g., to a linking phosphate / to a phosphodiester linkage / to the phosphodiester backbone). In certain embodiments, alterations (e.g., one or more alterations) are present in each of the nucleobase, the sugar, and the internucleoside linkage. Alterations according to the present disclosure may be alterations of ribonucleic acids (RNAs) to deoxyribonucleic acids (DNAs), e.g., the substitution of the 2'-OH of the ribofuranosyl ring to 2'-H, threose nucleic acids

20 (TNAs), glycol nucleic acids (GNAs), peptide nucleic acids (PNAs), locked nucleic acids (LNAs), or hybrids thereof. Additional alterations are described herein.

25 **[00328]** Polynucleotides and nucleic acids may or may not be uniformly altered along the entire length of the molecule. For example, one or more or all types of nucleotide (e.g., purine or pyrimidine, or any one or more or all of A, G, U, C) may or may not be uniformly altered in a polynucleotide or nucleic acid, or in a given predetermined sequence region thereof. In some instances, all nucleotides X in a polynucleotide (or in a given sequence region thereof) are altered, wherein X may any one of nucleotides A, G, U, C, or any one of the combinations A+G, A+U, A+C, G+U, G+C, U+C, A+G+U, A+G+C, G+U+C or A+G+C.

30 **[00329]** Different sugar alterations and/or internucleoside linkages (e.g., backbone structures) may exist at various positions in a polynucleotide. One of ordinary skill in the art will appreciate that the nucleotide analogs or other alteration(s) may be located at any position(s) of a polynucleotide such that the function of the polynucleotide is not substantially decreased. An alteration may also be a 5'- or 3'-terminal alteration. In some embodiments, the

polynucleotide includes an alteration at the 3'-terminus. The polynucleotide may contain from about 1% to about 100% alternative nucleotides (either in relation to overall nucleotide content, or in relation to one or more types of nucleotide, i.e., any one or more of A, G, U or C) or any intervening percentage (e.g., from 1% to 20%, from 1% to 25%, from 1% to 50%, from 1% to 5 60%, from 1% to 70%, from 1% to 80%, from 1% to 90%, from 1% to 95%, from 10% to 20%, from 10% to 25%, from 10% to 50%, from 10% to 60%, from 10% to 70%, from 10% to 80%, from 10% to 90%, from 10% to 95%, from 10% to 100%, from 20% to 25%, from 20% to 50%, from 20% to 60%, from 20% to 70%, from 20% to 80%, from 20% to 90%, from 20% to 95%, 10 from 20% to 100%, from 50% to 60%, from 50% to 70%, from 50% to 80%, from 50% to 90%, from 50% to 95%, from 50% to 100%, from 70% to 80%, from 70% to 90%, from 70% to 95%, from 70% to 100%, from 80% to 90%, from 80% to 95%, from 80% to 100%, from 90% to 95%, from 90% to 100%, and from 95% to 100%). It will be understood that any remaining percentage is accounted for by the presence of a canonical nucleotide (e.g., A, G, U, or C).

[00330] Polynucleotides may contain at a minimum zero and at maximum 100% 15 alternative nucleotides, or any intervening percentage, such as at least 5% alternative nucleotides, at least 10% alternative nucleotides, at least 25% alternative nucleotides, at least 50% alternative nucleotides, at least 80% alternative nucleotides, or at least 90% alternative nucleotides. For example, polynucleotides may contain an alternative pyrimidine such as an alternative uracil or cytosine. In some embodiments, at least 5%, at least 10%, at least 25%, at 20 least 50%, at least 80%, at least 90% or 100% of the uracil in a polynucleotide is replaced with an alternative uracil (e.g., a 5-substituted uracil). The alternative uracil can be replaced by a compound having a single unique structure, or can be replaced by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures). In some instances, at least 5%, at least 10%, at least 25%, at least 50%, at least 80%, at least 90% or 100% of the cytosine 25 in the polynucleotide is replaced with an alternative cytosine (e.g., a 5-substituted cytosine). The alternative cytosine can be replaced by a compound having a single unique structure, or can be replaced by a plurality of compounds having different structures (e.g., 2, 3, 4 or more unique structures).

[00331] In some instances, nucleic acids do not substantially induce an innate immune 30 response of a cell into which the polynucleotide (e.g., mRNA) is introduced. Features of an induced innate immune response include 1) increased expression of pro-inflammatory cytokines, 2) activation of intracellular PRRs (RIG-I, MDA5, etc., and/or 3) termination or reduction in protein translation.

[00332] The nucleic acids can optionally include other agents (e.g., RNAi-inducing agents, RNAi agents, siRNAs, shRNAs, miRNAs, antisense RNAs, ribozymes, catalytic DNA, tRNA, RNAs that induce triple helix formation, aptamers, and vectors). In some embodiments, the nucleic acids may include one or more messenger RNAs (mRNAs) having one or more 5 alternative nucleoside or nucleotides (i.e., alternative mRNA molecules).

[00333] In some embodiments, a nucleic acid (e.g. mRNA) molecule, formula, composition or method associated therewith comprises one or more polynucleotides comprising features as described in WO2002/098443, WO2003/051401, WO2008/052770, WO2009127230, WO2006122828, WO2008/083949, WO2010088927, WO2010/037539, 10 WO2004/004743, WO2005/016376, WO2006/024518, WO2007/095976, WO2008/014979, WO2008/077592, WO2009/030481, WO2009/095226, WO2011069586, WO2011026641, WO2011/144358, WO2012019780, WO2012013326, WO2012089338, WO2012113513, WO2012116811, WO2012116810, WO2013113502, WO2013113501, WO2013113736, WO2013143698, WO2013143699, WO2013143700, WO2013/120626, WO2013120627, 15 WO2013120628, WO2013120629, WO2013174409, WO2014127917, WO2015/024669, WO2015/024668, WO2015/024667, WO2015/024665, WO2015/024666, WO2015/024664, WO2015101415, WO2015101414, WO2015024667, WO2015062738, WO2015101416, all of which are incorporated by reference herein.

20 *Nucleobase Alternatives*

[00334] The alternative nucleosides and nucleotides can include an alternative nucleobase. A nucleobase of a nucleic acid is an organic base such as a purine or pyrimidine or a derivative thereof. A nucleobase may be a canonical base (e.g., adenine, guanine, uracil, thymine, and cytosine). These nucleobases can be altered or wholly replaced to provide 25 polynucleotide molecules having enhanced properties, e.g., increased stability such as resistance to nucleases. Non-canonical or modified bases may include, for example, one or more substitutions or modifications including but not limited to alkyl, aryl, halo, oxo, hydroxyl, alkyloxy, and/or thio substitutions; one or more fused or open rings; oxidation; and/or reduction.

[00335] Alternative nucleotide base pairing encompasses not only the standard adenine-thymine, adenine-uracil, or guanine-cytosine base pairs, but also base pairs formed between 30 nucleotides and/or alternative nucleotides including non-standard or alternative bases, wherein the arrangement of hydrogen bond donors and hydrogen bond acceptors permits hydrogen bonding between a non-standard base and a standard base or between two complementary non-

standard base structures. One example of such non-standard base pairing is the base pairing between the alternative nucleotide inosine and adenine, cytosine, or uracil.

[00336] In some embodiments, the nucleobase is an alternative uracil. Exemplary nucleobases and nucleosides having an alternative uracil include pseudouridine (ψ), pyridin-4-one ribonucleoside, 5-aza-uracil, 6-aza-uracil, 2-thio-5-aza-uracil, 2-thio-uracil (s^2U), 4-thio-uracil (s^4U), 4-thio-pseudouridine, 2-thio-pseudouridine, 5-hydroxy-uracil (ho^5U), 5-aminoallyl-uracil, 5-halo-uracil (e.g., 5-iodo-uracil or 5-bromo-uracil), 3-methyl-uracil (m^3U), 5-methoxy-uracil (mo^5U), uracil 5-oxyacetic acid (cmo^5U), uracil 5-oxyacetic acid methyl ester ($mcmo^5U$), 5-carboxymethyl-uracil (cm^5U), 1-carboxymethyl-pseudouridine, 5-carboxyhydroxymethyl-uracil (chm^5U), 5-carboxyhydroxymethyl-uracil methyl ester ($mchm^5U$), 5-methoxycarbonylmethyl-uracil (mcm^5U), 5-methoxycarbonylmethyl-2-thio-uracil (mcm^5s^2U), 5-aminomethyl-2-thio-uracil (nm^5s^2U), 5-methylaminomethyl-uracil (mnm^5U), 5-methylaminomethyl-2-thio-uracil (mnm^5s^2U), 5-methylaminomethyl-2-seleno-uracil (mnm^5se^2U), 5-carbamoylmethyl-uracil (ncm^5U), 5-carboxymethylaminomethyl-uracil ($cmnm^5U$), 5-carboxymethylaminomethyl-2-thio-uracil ($cmnm^5s^2U$), 5-propynyl-uracil, 1-propynyl-pseudouracil, 5-taurinomethyl-uracil (tm^5U), 1-taurinomethyl-pseudouridine, 5-taurinomethyl-2-thio-uracil (tm^5s^2U), 1-taurinomethyl-4-thio-pseudouridine, 5-methyl-uracil (m^5U , i.e., having the nucleobase deoxythymine), 1-methyl-pseudouridine ($m^1\psi$), 1-ethyl-pseudouridine ($Et^1\psi$), 5-methyl-2-thio-uracil (m^5s^2U), 1-methyl-4-thio-pseudouridine ($m^1s^4\psi$), 4-thio-1-methyl-pseudouridine, 3-methyl-pseudouridine ($m^3\psi$), 2-thio-1-methyl-pseudouridine, 1-methyl-1-deaza-pseudouridine, 2-thio-1-methyl-1-deaza-pseudouridine, dihydrouracil (D), dihydropseudouridine, 5,6-dihydrouracil, 5-methyl-dihydrouracil (m^5D), 2-thio-dihydrouracil, 2-thio-dihydropseudouridine, 2-methoxy-uracil, 2-methoxy-4-thio-uracil, 4-methoxy-pseudouridine, 4-methoxy-2-thio-pseudouridine, N1-methyl-pseudouridine, 3-(3-amino-3-carboxypropyl)uracil (acp^3U), 1-methyl-3-(3-amino-3-carboxypropyl)pseudouridine ($acp^3\psi$), 5-(isopentenylaminomethyl)uracil (inm^5U), 5-(isopentenylaminomethyl)-2-thio-uracil (inm^5s^2U), 5,2'-O-dimethyl-uridine (m^5Um), 2-thio-2'-O_methyl-uridine (s^2Um), 5-methoxycarbonylmethyl-2'-O_methyl-uridine (mcm^5Um), 5-carbamoylmethyl-2'-O_methyl-uridine (ncm^5Um), 5-carboxymethylaminomethyl-2'-O_methyl-uridine ($cmnm^5Um$), 3,2'-O-dimethyl-uridine (m^3Um), and 5-(isopentenylaminomethyl)-2'-O_methyl-uridine (inm^5Um), 1-thio-uracil, deoxythymidine, 5-(2-carbomethoxyvinyl)-uracil, 5-(carbamoylhydroxymethyl)-uracil, 5-carbamoylmethyl-2-thio-uracil, 5-carboxymethyl-2-thio-uracil, 5-cyanomethyl-uracil, 5-methoxy-2-thio-uracil, and 5-[3-(1-E-propenylamino)]uracil.

[00337] In some embodiments, the nucleobase is an alternative cytosine. Exemplary nucleobases and nucleosides having an alternative cytosine include 5-aza-cytosine, 6-aza-cytosine, pseudoisocytidine, 3-methyl-cytosine (m3C), N4-acetyl-cytosine (ac4C), 5-formyl-cytosine (f5C), N4-methyl-cytosine (m4C), 5-methyl-cytosine (m5C), 5-halo-cytosine (e.g., 5-iodo-cytosine), 5-hydroxymethyl-cytosine (hm5C), 1-methyl-pseudoisocytidine, pyrrolo-cytosine, pyrrolo-pseudoisocytidine, 2-thio-cytosine (s2C), 2-thio-5-methyl-cytosine, 4-thio-pseudoisocytidine, 4-thio-1-methyl-pseudoisocytidine, 4-thio-1-methyl-1-deaza-pseudoisocytidine, 1-methyl-1-deaza-pseudoisocytidine, zebularine, 5-aza-zebularine, 5-methyl-zebularine, 5-aza-2-thio-zebularine, 2-thio-zebularine, 2-methoxy-cytosine, 2-methoxy-5-methyl-cytosine, 4-methoxy-pseudoisocytidine, 4-methoxy-1-methyl-pseudoisocytidine, lysidine (k2C), 5,2'-O-dimethyl-cytidine (m5Cm), N4-acetyl-2'-O-methyl-cytidine (ac4Cm), N4,2'-O-dimethyl-cytidine (m4Cm), 5-formyl-2'-O-methyl-cytidine (f5Cm), N4,N4,2'-O-trimethyl-cytidine (m42Cm), 1-thio-cytosine, 5-hydroxy-cytosine, 5-(3-azidopropyl)-cytosine, and 5-(2-azidoethyl)-cytosine.

[00338] In some embodiments, the nucleobase is an alternative adenine. Exemplary nucleobases and nucleosides having an alternative adenine include 2-amino-purine, 2,6-diaminopurine, 2-amino-6-halo-purine (e.g., 2-amino-6-chloro-purine), 6-halo-purine (e.g., 6-chloro-purine), 2-amino-6-methyl-purine, 8-azido-adenine, 7-deaza-adenine, 7-deaza-8-aza-adenine, 7-deaza-2-amino-purine, 7-deaza-8-aza-2-amino-purine, 7-deaza-2,6-diaminopurine, 7-deaza-8-aza-2,6-diaminopurine, 1-methyl-adenine (m1A), 2-methyl-adenine (m2A), N6-methyl-adenine (m6A), 2-methylthio-N6-methyl-adenine (ms2m6A), N6-isopentenyl-adenine (i6A), 2-methylthio-N6-isopentenyl-adenine (ms2i6A), N6-(cis-hydroxyisopentenyl)adenine (i6A), 2-methylthio-N6-(cis-hydroxyisopentenyl)adenine (ms2i6A), N6-glycylcarbamoyl-adenine (g6A), N6-threonylcarbamoyl-adenine (t6A), N6-methyl-N6-threonylcarbamoyl-adenine (m6t6A), 2-methylthio-N6-threonylcarbamoyl-adenine (ms2g6A), N6,N6-dimethyl-adenine (m62A), N6-hydroxynorvalylcarbamoyl-adenine (hn6A), 2-methylthio-N6-hydroxynorvalylcarbamoyl-adenine (ms2hn6A), N6-acetyl-adenine (ac6A), 7-methyl-adenine, 2-methylthio-adenine, 2-methoxy-adenine, N6,2'-O-dimethyl-adenosine (m6Am), N6,N6,2'-O-trimethyl-adenosine (m62Am), 1,2'-O-dimethyl-adenosine (m1Am), 2-amino-N6-methyl-purine, 1-thio-adenine, 8-azido-adenine, N6-(19-amino-pentaoxonanadecyl)-adenine, 2,8-dimethyl-adenine, N6-formyl-adenine, and N6-hydroxymethyl-adenine.

[00339] In some embodiments, the nucleobase is an alternative guanine. Exemplary nucleobases and nucleosides having an alternative guanine include inosine (I), 1-methyl-inosine


(m1I), wyosine (imG), methylwyosine (mimG), 4-demethyl-wyosine (imG-14), isowyosine (imG2), wybutosine (yW), peroxywybutosine (o2yW), hydroxywybutosine (OHyW), undermodified hydroxywybutosine (OHyW*), 7-deaza-guanine, queuosine (Q), epoxyqueuosine (oQ), galactosyl-queuosine (galQ), mannosyl-queuosine (manQ), 7-cyano-7-deaza-guanine

5 (preQ0), 7-aminomethyl-7-deaza-guanine (preQ1), archaeosine (G+), 7-deaza-8-aza-guanine, 6-thio-guanine, 6-thio-7-deaza-guanine, 6-thio-7-deaza-8-aza-guanine, 7-methyl-guanine (m7G), 6-thio-7-methyl-guanine, 7-methyl-inosine, 6-methoxy-guanine, 1-methyl-guanine (m1G), N2-methyl-guanine (m2G), N2,N2-dimethyl-guanine (m22G), N2,7-dimethyl-guanine (m2,7G), N2,10 N2,7-dimethyl-guanine (m2,2,7G), 8-oxo-guanine, 7-methyl-8-oxo-guanine, 1-methyl-6-thio-guanine, N2-methyl-6-thio-guanine, N2,N2-dimethyl-6-thio-guanine, N2-methyl-2'-O-methyl-guanosine (m2Gm), N2,N2-dimethyl-2'-O-methyl-guanosine (m22Gm), 1-methyl-2'-O-methyl-guanosine (m1Gm), N2,7-dimethyl-2'-O-methyl-guanosine (m2,7Gm), 2'-O-methyl-inosine (Im), 1,2'-O-dimethyl-inosine (m1Im), 1-thio-guanine, and O-6-methyl-guanine.

[00340] The alternative nucleobase of a nucleotide can be independently a purine, a pyrimidine, a purine or pyrimidine analog. For example, the nucleobase can be an alternative to adenine, cytosine, guanine, uracil, or hypoxanthine. In another embodiment, the nucleobase can also include, for example, naturally-occurring and synthetic derivatives of a base, including pyrazolo[3,4-d]pyrimidines, 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo (e.g., 8-bromo), 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxy and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, deazaguanine, 7-deazaguanine, 3-deazaguanine, deazaadenine, 7-deazaadenine, 3-deazaadenine, pyrazolo[3,4-d]pyrimidine, imidazo[1,5-a]1,3,5 triazinones, 9-deazapurines, imidazo[4,5-d]pyrazines, thiazolo[4,5-d]pyrimidines, pyrazin-2-ones, 1,2,4-triazine, pyridazine; or 1,3,5 triazine. When the nucleotides are depicted using the shorthand A, G, C, T or U, each letter refers to the representative base and/or derivatives thereof, e.g., A includes adenine or adenine analogs, e.g., 7-deaza adenine).

Alterations on the Sugar


[00341] Nucleosides include a sugar molecule (e.g., a 5-carbon or 6-carbon sugar, such as pentose, ribose, arabinose, xylose, glucose, galactose, or a deoxy derivative thereof) in combination with a nucleobase, while nucleotides are nucleosides containing a nucleoside and a phosphate group or alternative group (e.g., boranophosphate, thiophosphate, selenophosphate, phosphonate, alkyl group, amide, and glycerol). A nucleoside or nucleotide may be a canonical species, e.g., a nucleoside or nucleotide including a canonical nucleobase, sugar, and, in the case of nucleotides, a phosphate group, or may be an alternative nucleoside or nucleotide including one or more alternative components. For example, alternative nucleosides and nucleotides can be altered on the sugar of the nucleoside or nucleotide. In some embodiments, 5 the alternative nucleosides or nucleotides include the structure: 10



Formula VI

Formula VII

Formula VIII



Formula IX

15 In each of the Formulae VI, VII, VIII, and IX, each of m and n is independently, an integer from 0 to 5, each of U and U' independently, is O, S, N(R^U)_{nu}, or C(R^U)_{nu}, wherein nu is an integer from 0 to 2 and each R^U is, independently, H, halo, or optionally substituted alkyl; each of R¹, R², R^{1'}, R^{2'}, R¹, R², R³, R⁴, and R⁵ is, independently, if present, H, halo, 20 hydroxy, thiol, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, optionally substituted aminoalkoxy, optionally substituted alkoxyalkoxy, optionally substituted hydroxyalkoxy, optionally substituted amino, azido, optionally substituted aryl, optionally substituted aminoalkyl, optionally substituted aminoalkenyl, optionally substituted aminoalkynyl, or absent; wherein the combination of R³ 25 with one or more of R¹, R^{1'}, R², R^{2'}, or R⁵ (e.g., the combination of R¹ and R³, the combination

of $R^{1''}$ and R^3 , the combination of $R^{2'}$ and R^3 , the combination of $R^{2''}$ and R^3 , or the combination of R^5 and R^3) can join together to form optionally substituted alkylene or optionally substituted heteroalkylene and, taken together with the carbons to which they are attached, provide an optionally substituted heterocycl (e.g., a bicyclic, tricyclic, or tetracyclic heterocycl);

5 wherein the combination of R^5 with one or more of R^1' , $R^{1''}$, $R^{2'}$, or $R^{2''}$ (e.g., the combination of R^1' and R^5 , the combination of $R^{1''}$ and R^5 , the combination of $R^{2'}$ and R^5 , or the combination of $R^{2''}$ and R^5) can join together to form optionally substituted alkylene or optionally substituted heteroalkylene and, taken together with the carbons to which they are attached, provide an optionally substituted heterocycl (e.g., a bicyclic, tricyclic, or tetracyclic heterocycl); and

10 wherein the combination of R^4 and one or more of R^1' , $R^{1''}$, $R^{2'}$, $R^{2''}$, R^3 , or R^5 can join together to form optionally substituted alkylene or optionally substituted heteroalkylene and, taken together with the carbons to which they are attached, provide an optionally substituted heterocycl (e.g., a bicyclic, tricyclic, or tetracyclic heterocycl); each of m' and m'' is, independently, an integer from 0 to 3 (e.g., from 0 to 2, from 0 to 1, from 1 to 3, or from 1 to 2);

15 each of Y^1 , Y^2 , and Y^3 , is, independently, O, S, Se, $—NR^{N1}$, optionally substituted alkylene, or optionally substituted heteroalkylene, wherein R^{N1} is H, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted aryl, or absent;

20 each Y^4 is, independently, H, hydroxy, thiol, boranyl, optionally substituted alkyl, optionally substituted alkenyl, optionally substituted alkynyl, optionally substituted alkoxy, optionally substituted alkenyloxy, optionally substituted alkynyloxy, optionally substituted thioalkoxy, optionally substituted alkoxyalkoxy, or optionally substituted amino;

each Y^5 is, independently, O, S, Se, optionally substituted alkylene (e.g., methylene), or optionally substituted heteroalkylene; and

25 B is a nucleobase, either modified or unmodified. In some embodiments, the 2'-hydroxy group (OH) can be modified or replaced with a number of different substituents. Exemplary substitutions at the 2'-position include, but are not limited to, H, azido, halo (e.g., fluoro), optionally substituted C_{1-6} alkyl (e.g., methyl); optionally substituted C_{1-6} alkoxy (e.g., methoxy or ethoxy); optionally substituted C_{6-10} aryloxy; optionally substituted C_{3-8} cycloalkyl; optionally substituted C_{6-10} aryl- C_{1-6} alkoxy, optionally substituted C_{1-12} (heterocycl)oxy; a sugar (e.g., ribose, pentose, or any described herein); a polyethyleneglycol (PEG), -
30 $O(CH_2CH_2O)_nCH_2CH_2OR$, where R is H or optionally substituted alkyl, and n is an integer from 0 to 20 (e.g., from 0 to 4, from 0 to 8, from 0 to 10, from 0 to 16, from 1 to 4, from 1 to 8, from

1 to 10, from 1 to 16, from 1 to 20, from 2 to 4, from 2 to 8, from 2 to 10, from 2 to 16, from 2 to 20, from 4 to 8, from 4 to 10, from 4 to 16, and from 4 to 20); “locked” nucleic acids (LNA) in which the 2'-hydroxy is connected by a C₁₋₆ alkylene or C₁₋₆ heteroalkylene bridge to the 4'-carbon of the same ribose sugar, where exemplary bridges included methylene, propylene, ether, 5 or amino bridges; aminoalkyl, as defined herein; aminoalkoxy, as defined herein; amino as defined herein; and amino acid, as defined herein.

[00343] Generally, RNA includes the sugar group ribose, which is a 5-membered ring having an oxygen. Exemplary, non-limiting alternative nucleotides include replacement of the oxygen in ribose (e.g., with S, Se, or alkylene, such as methylene or ethylene); addition of a 10 double bond (e.g., to replace ribose with cyclopentenyl or cyclohexenyl); ring contraction of ribose (e.g., to form a 4-membered ring of cyclobutane or oxetane); ring expansion of ribose (e.g., to form a 6- or 7-membered ring having an additional carbon or heteroatom, such as for anhydrohexitol, altritol, mannitol, cyclohexanyl, cyclohexenyl, and morpholino (that also has a phosphoramidate backbone)); multicyclic forms (e.g., tricyclo and “unlocked” forms, such as 15 glycol nucleic acid (GNA) (e.g., R-GNA or S-GNA, where ribose is replaced by glycol units attached to phosphodiester bonds), threose nucleic acid (TNA, where ribose is replaced with α-L-threofuranosyl-(3'→2')), and peptide nucleic acid (PNA, where 2-amino-ethyl-glycine linkages replace the ribose and phosphodiester backbone).

[00344] In some embodiments, the sugar group contains one or more carbons that possess 20 the opposite stereochemical configuration of the corresponding carbon in ribose. Thus, a polynucleotide molecule can include nucleotides containing, e.g., arabinose or L-ribose, as the sugar.

[00345] In some embodiments, the polynucleotide includes at least one nucleoside 25 wherein the sugar is L-ribose, 2'-O-methyl-ribose, 2'-fluoro-ribose, arabinose, hexitol, an LNA, or a PNA.

Alterations on the Internucleoside Linkage

[00346] Alternative nucleotides can be altered on the internucleoside linkage (e.g., phosphate backbone). Herein, in the context of the polynucleotide backbone, the phrases “phosphate” and “phosphodiester” are used interchangeably. Backbone phosphate groups can 30 be altered by replacing one or more of the oxygen atoms with a different substituent.

[00347] The alternative nucleotides can include the wholesale replacement of an unaltered phosphate moiety with another internucleoside linkage as described herein. Examples of alternative phosphate groups include, but are not limited to, phosphorothioate,

phosphoroselenates, boranophosphates, boranophosphate esters, hydrogen phosphonates, phosphoramidates, phosphorodiamidates, alkyl or aryl phosphonates, and phosphotriesters. Phosphorodithioates have both non-linking oxygens replaced by sulfur. The phosphate linker can also be altered by the replacement of a linking oxygen with nitrogen (bridged phosphoramidates), sulfur (bridged phosphorothioates), and carbon (bridged methylene-phosphonates).

5 [00348] The alternative nucleosides and nucleotides can include the replacement of one or more of the non-bridging oxygens with a borane moiety (BH₃), sulfur (thio), methyl, ethyl, and/or methoxy. As a non-limiting example, two non-bridging oxygens at the same position 10 (e.g., the alpha (α), beta (β) or gamma (γ) position) can be replaced with a sulfur (thio) and a methoxy.

15 [00349] The replacement of one or more of the oxygen atoms at the α position of the phosphate moiety (e.g., α-thio phosphate) is provided to confer stability (such as against exonucleases and endonucleases) to RNA and DNA through the unnatural phosphorothioate backbone linkages. Phosphorothioate DNA and RNA have increased nuclease resistance and subsequently a longer half-life in a cellular environment.

20 [00350] Other internucleoside linkages that may be employed according to the present disclosure, including internucleoside linkages which do not contain a phosphorous atom, are described herein.

20

Internal ribosome entry sites

[00351] Polynucleotides may contain an internal ribosome entry site (IRES). An IRES may act as the sole ribosome binding site, or may serve as one of multiple ribosome binding sites of an mRNA. A polynucleotide containing more than one functional ribosome binding site 25 may encode several peptides or polypeptides that are translated independently by the ribosomes (e.g., multicistronic mRNA). When polynucleotides are provided with an IRES, further optionally provided is a second translatable region. Examples of IRES sequences that can be used according to the present disclosure include without limitation, those from picornaviruses (e.g., FMDV), pest viruses (CFFV), polio viruses (PV), encephalomyocarditis viruses (ECMV), 30 foot-and-mouth disease viruses (FMDV), hepatitis C viruses (HCV), classical swine fever viruses (CSFV), murine leukemia virus (MLV), simian immune deficiency viruses (SIV) or cricket paralysis viruses (CrPV).

5'-cap structure

[00352] A polynucleotide (e.g., an mRNA) may include a 5'-cap structure. The 5'-cap structure of a polynucleotide is involved in nuclear export and increasing polynucleotide stability and binds the mRNA Cap Binding Protein (CBP), which is responsible for polynucleotide stability in the cell and translation competency through the association of CBP with poly-A binding protein to form the mature cyclic mRNA species. The cap further assists the removal of 5'-proximal introns removal during mRNA splicing.

[00353] Endogenous polynucleotide molecules may be 5'-end capped generating a 5'-PPP-5'-triphosphate linkage between a terminal guanosine cap residue and the 5'-terminal transcribed sense nucleotide of the polynucleotide. This 5'-guanylate cap may then be methylated to generate an N7-methyl-guanylate residue. The ribose sugars of the terminal and/or anteterminal transcribed nucleotides of the 5' end of the polynucleotide may optionally also be 2'-O-methylated. 5'-decapping through hydrolysis and cleavage of the guanylate cap structure may target a polynucleotide molecule, such as an mRNA molecule, for degradation.

[00354] Alterations to polynucleotides may generate a non-hydrolyzable cap structure preventing decapping and thus increasing polynucleotide half-life. Because cap structure hydrolysis requires cleavage of 5'-PPP-5' phosphorodiester linkages, alternative nucleotides may be used during the capping reaction. For example, a Vaccinia Capping Enzyme from New England Biolabs (Ipswich, MA) may be used with α -thio-guanosine nucleotides according to the manufacturer's instructions to create a phosphorothioate linkage in the 5'-PPP-5' cap.

[00355] Additional alternative guanosine nucleotides may be used such as α -methyl-phosphonate and seleno-phosphate nucleotides.

[00356] Additional alterations include, but are not limited to, 2'-O-methylation of the ribose sugars of 5'-terminal and/or 5'-anteterminal nucleotides of the polynucleotide (as mentioned above) on the 2'-hydroxy group of the sugar. Multiple distinct 5'-cap structures can be used to generate the 5'-cap of a polynucleotide, such as an mRNA molecule.

[00357] 5'-Cap structures include those described in International Patent Publication Nos. WO2008127688, WO 2008016473, and WO 2011015347, the cap structures of each of which are incorporated herein by reference.

[00358] Cap analogs, which herein are also referred to as synthetic cap analogs, chemical caps, chemical cap analogs, or structural or functional cap analogs, differ from natural (i.e., endogenous, wild-type, or physiological) 5'-caps in their chemical structure, while retaining cap function. Cap analogs may be chemically (i.e., non-enzymatically) or enzymatically synthesized and/linked to a polynucleotide.

[00358] For example, the Anti-Reverse Cap Analog (ARCA) cap contains two guanosines linked by a 5'-5'-triphosphate group, wherein one guanosine contains an N7-methyl group as well as a 3'-O-methyl group (i.e., N7,3'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine, m⁷G-3'mppp-G, which may equivalently be designated 3' O-Me-m7G(5')ppp(5')G). The 3'-O atom of the other, unaltered, guanosine becomes linked to the 5'-terminal nucleotide of the capped polynucleotide (e.g., an mRNA). The N7- and 3'-O-methylated guanosine provides the terminal moiety of the capped polynucleotide (e.g., mRNA).

5 **[00359]** Another exemplary cap is mCAP, which is similar to ARCA but has a 2'-O-methyl group on guanosine (i.e., N7,2'-O-dimethyl-guanosine-5'-triphosphate-5'-guanosine, 10 m⁷Gm-ppp-G).

15 **[00360]** A cap may be a dinucleotide cap analog. As a non-limiting example, the dinucleotide cap analog may be modified at different phosphate positions with a boranophosphate group or a phosphoroselenoate group such as the dinucleotide cap analogs described in US Patent No. 8,519,110, the cap structures of which are herein incorporated by reference.

20 **[00361]** Alternatively, a cap analog may be a N7-(4-chlorophenoxyethyl) substituted dinucleotide cap analog known in the art and/or described herein. Non-limiting examples of N7-(4-chlorophenoxyethyl) substituted dinucleotide cap analogs include a N7-(4-chlorophenoxyethyl)-G(5')ppp(5')G and a N7-(4-chlorophenoxyethyl)-m3'-OG(5')ppp(5')G cap analog (see, e.g., the various cap analogs and the methods of synthesizing cap analogs described in Kore et al. *Bioorganic & Medicinal Chemistry* 2013 21:4570-4574; the cap structures of which are herein incorporated by reference). In other instances, a cap analog useful in the polynucleotides of the present disclosure is a 4-chloro/bromophenoxyethyl analog.

25 **[00362]** While cap analogs allow for the concomitant capping of a polynucleotide in an *in vitro* transcription reaction, up to 20% of transcripts remain uncapped. This, as well as the structural differences of a cap analog from endogenous 5'-cap structures of polynucleotides produced by the endogenous, cellular transcription machinery, may lead to reduced translational competency and reduced cellular stability.

30 **[00363]** Alternative polynucleotides may also be capped post-transcriptionally, using enzymes, in order to generate more authentic 5'-cap structures. As used herein, the phrase “more authentic” refers to a feature that closely mirrors or mimics, either structurally or functionally, an endogenous or wild type feature. That is, a “more authentic” feature is better representative of an endogenous, wild-type, natural or physiological cellular function, and/or

structure as compared to synthetic features or analogs of the prior art, or which outperforms the corresponding endogenous, wild-type, natural, or physiological feature in one or more respects. Non-limiting examples of more authentic 5'-cap structures useful in the polynucleotides of the present disclosure are those which, among other things, have enhanced binding of cap binding 5 proteins, increased half-life, reduced susceptibility to 5'-endonucleases, and/or reduced 5'- decapping, as compared to synthetic 5'-cap structures known in the art (or to a wild-type, natural or physiological 5'-cap structure). For example, recombinant Vaccinia Virus Capping Enzyme and recombinant 2'-O-methyltransferase enzyme can create a canonical 5'-5'-triphosphate linkage between the 5'-terminal nucleotide of a polynucleotide and a guanosine cap nucleotide 10 wherein the cap guanosine contains an N7-methylation and the 5'-terminal nucleotide of the polynucleotide contains a 2'-O-methyl. Such a structure is termed the Cap1 structure. This cap results in a higher translational-competency, cellular stability, and a reduced activation of cellular pro-inflammatory cytokines, as compared, e.g., to other 5' cap analog structures known in the art. Other exemplary cap structures include 7mG(5')ppp(5')N,pN2p (Cap 0), 15 7mG(5')ppp(5')NlmpNp (Cap 1), 7mG(5')-ppp(5')NlmpN2mp (Cap 2), and m(7)Gpppm(3)(6,6,2')Apm(2')Apm(2')Cpm(2)(3,2')Up (Cap 4).

20 [00364] Because the alternative polynucleotides may be capped post-transcriptionally, and because this process is more efficient, nearly 100% of the alternative polynucleotides may be capped. This is in contrast to ~80% when a cap analog is linked to an polynucleotide in the course of an in vitro transcription reaction.

25 [00365] 5'-terminal caps may include endogenous caps or cap analogs. A 5'-terminal cap may include a guanosine analog. Useful guanosine analogs include inosine, N1-methyl-guanosine, 2'-fluoro-guanosine, 7-deaza-guanosine, 8-oxo-guanosine, 2-amino-guanosine, LNA-guanosine, and 2-azido-guanosine.

30 [00366] In some cases, a polynucleotide contains a modified 5'-cap. A modification on the 5'-cap may increase the stability of polynucleotide, increase the half-life of the polynucleotide, and could increase the polynucleotide translational efficiency. The modified 5'-cap may include, but is not limited to, one or more of the following modifications: modification at the 2'- and/or 3'-position of a capped guanosine triphosphate (GTP), a replacement of the sugar ring oxygen (that produced the carbocyclic ring) with a methylene moiety (CH₂), a modification at the triphosphate bridge moiety of the cap structure, or a modification at the nucleobase (G) moiety.

5'-UTRs

[00367] A 5'-UTR may be provided as a flanking region to polynucleotides (e.g., mRNAs). A 5'-UTR may be homologous or heterologous to the coding region found in a polynucleotide. Multiple 5'-UTRs may be included in the flanking region and may be the same or of different sequences. Any portion of the flanking regions, including none, may be codon 5 optimized and any may independently contain one or more different structural or chemical alterations, before and/or after codon optimization.

[00368] Shown in Table 21 in US Provisional Application No 61/775,509, and in Table 21 and in Table 22 in US Provisional Application No. 61/829,372, of which are incorporated herein by reference, is a listing of the start and stop site of alternative polynucleotides (e.g., mRNA). In Table 21 each 5'-UTR (5'-UTR-005 to 5'-UTR 68511) is identified by its start and stop site relative to its native or wild type (homologous) transcript (ENST; the identifier used in the ENSEMBL database).

[00369] To alter one or more properties of a polynucleotide (e.g., mRNA), 5'-UTRs which are heterologous to the coding region of an alternative polynucleotide (e.g., mRNA) may 15 be engineered. The polynucleotides (e.g., mRNA) may then be administered to cells, tissue or organisms and outcomes such as protein level, localization, and/or half-life may be measured to evaluate the beneficial effects the heterologous 5'-UTR may have on the alternative polynucleotides (mRNA). Variants of the 5'-UTRs may be utilized wherein one or more nucleotides are added or removed to the termini, including A, T, C or G. 5'-UTRs may also be 20 codon-optimized, or altered in any manner described herein.

5'-UTRs, 3'-UTRs, and Translation Enhancer Elements (TEEs)

[00370] The 5'-UTR of a polynucleotides (e.g., mRNA) may include at least one translation enhancer element. The term “translational enhancer element” refers to sequences that increase the amount of polypeptide or protein produced from a polynucleotide. As a non-25 limiting example, the TEE may be located between the transcription promoter and the start codon. The polynucleotides (e.g., mRNA) with at least one TEE in the 5'-UTR may include a cap at the 5'-UTR. Further, at least one TEE may be located in the 5'-UTR of polynucleotides (e.g., mRNA) undergoing cap-dependent or cap-independent translation.

[00371] In one aspect, TEEs are conserved elements in the UTR which can promote 30 translational activity of a polynucleotide such as, but not limited to, cap-dependent or cap-independent translation. The conservation of these sequences has been previously shown by Panek et al. (Nucleic Acids Research, 2013, 1-10) across 14 species including humans.

[00372] In one non-limiting example, the TEEs known may be in the 5'-leader of the Gtx homeodomain protein (Chappell et al., Proc. Natl. Acad. Sci. USA 101:9590-9594, 2004, the TEEs of which are incorporated herein by reference).

[00373] In another non-limiting example, TEEs are disclosed in US Patent Publication Nos. 2009/0226470 and 2013/0177581, International Patent Publication Nos. WO2009/075886, WO2012/009644, and WO1999/024595, US Patent Nos. 6,310,197 and 6,849,405, the TEE sequences disclosed in each of which are incorporated herein by reference.

[00374] In yet another non-limiting example, the TEE may be an internal ribosome entry site (IRES), HCV-IRES or an IRES element such as, but not limited to, those described in US Patent No. 7,468,275, US Patent Publication Nos. 2007/0048776 and 2011/0124100 and International Patent Publication Nos. WO2007/025008 and WO2001/055369, the IRES sequences of each of which are incorporated herein by reference. The IRES elements may include, but are not limited to, the Gtx sequences (e.g., Gtx9-nt, Gtx8-nt, Gtx7-nt) described by Chappell et al. (Proc. Natl. Acad. Sci. USA 101:9590-9594, 2004) and Zhou et al. (PNAS 102:6273-6278, 2005) and in US Patent Publication Nos. 2007/0048776 and 2011/0124100 and International Patent Publication No. WO2007/025008, the IRES sequences of each of which are incorporated herein by reference.

[00375] “Translational enhancer polynucleotides” are polynucleotides which include one or more of the specific TEE exemplified herein and/or disclosed in the art (see e.g., U.S. Patent Nos. 6,310,197, 6,849,405, 7,456,273, 7,183,395, U.S. Patent Publication Nos. 20090/226470, 2007/0048776, 2011/0124100, 2009/0093049, 2013/0177581, International Patent Publication Nos. WO2009/075886, WO2007/025008, WO2012/009644, WO2001/055371, WO1999/024595, and European Patent Nos. 2610341 and 2610340; the TEE sequences of each of which are incorporated herein by reference) or their variants, homologs or functional derivatives. One or multiple copies of a specific TEE can be present in a polynucleotide (e.g., mRNA). The TEEs in the translational enhancer polynucleotides can be organized in one or more sequence segments. A sequence segment can harbor one or more of the specific TEEs exemplified herein, with each TEE being present in one or more copies. When multiple sequence segments are present in a translational enhancer polynucleotide, they can be homogenous or heterogeneous. Thus, the multiple sequence segments in a translational enhancer polynucleotide can harbor identical or different types of the specific TEEs exemplified herein, identical or different number of copies of each of the specific TEEs, and/or identical or different organization of the TEEs within each sequence segment.

[00376] A polynucleotide (e.g., mRNA) may include at least one TEE that is described in International Patent Publication Nos. WO1999/024595, WO2012/009644, WO2009/075886, WO2007/025008, WO1999/024595, European Patent Publication Nos. 2610341 and 2610340, US Patent Nos. 6,310,197, 6,849,405, 7,456,273, 7,183,395, and US Patent Publication Nos.

5 2009/0226470, 2011/0124100, 2007/0048776, 2009/0093049, and 2013/0177581 the TEE sequences of each of which are incorporated herein by reference. The TEE may be located in the 5'-UTR of the polynucleotides (e.g., mRNA).

[00377] A polynucleotide (e.g., mRNA) may include at least one TEE that has at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95% or at least 99% identity with the TEEs described in US Patent Publication Nos. 2009/0226470, 2007/0048776, 2013/0177581 and 2011/0124100, International Patent Publication Nos. WO1999/024595, WO2012/009644, WO2009/075886 and WO2007/025008, European Patent Publication Nos. 2610341 and 2610340, US Patent Nos. 6,310,197, 6,849,405, 7,456,273, 7,183,395, the TEE sequences of each of which are incorporated herein by reference.

[00378] The 5'-UTR of a polynucleotide (e.g., mRNA) may include at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18 at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55 or more than 60 TEE sequences. The TEE sequences in the 5'-UTR of a polynucleotide (e.g., mRNA) may be the same or different TEE sequences. The TEE sequences may be in a pattern such as ABABAB, AABBAABBAABB, or ABCABCABC, or variants thereof, repeated once, twice, or more than three times. In these patterns, each letter, A, B, or C represent a different TEE sequence at the nucleotide level.

25 **[00379]** In some cases, the 5'-UTR may include a spacer to separate two TEE sequences. As a non-limiting example, the spacer may be a 15 nucleotide spacer and/or other spacers known in the art. As another non-limiting example, the 5'-UTR may include a TEE sequence-spacer module repeated at least once, at least twice, at least 3 times, at least 4 times, at least 5 times, at least 6 times, at least 7 times, at least 8 times, at least 9 times, or more than 9 times in the 5'-UTR.

[00380] In other instances, the spacer separating two TEE sequences may include other sequences known in the art which may regulate the translation of the polynucleotides (e.g., mRNA) of the present disclosure such as, but not limited to, miR sequences (e.g., miR binding

sites and miR seeds). As a non-limiting example, each spacer used to separate two TEE sequences may include a different miR sequence or component of a miR sequence (e.g., miR seed sequence).

[00381] In some instances, the TEE in the 5'-UTR of a polynucleotide (e.g., mRNA) may include at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 99% or more than 99% of the TEE sequences disclosed in US Patent Publication Nos. 2009/0226470, 2007/0048776, 2013/0177581 and 2011/0124100, International Patent Publication Nos.

10 WO1999/024595, WO2012/009644, WO2009/075886 and WO2007/025008, European Patent Publication Nos. 2610341 and 2610340, and US Patent Nos. 6,310,197, 6,849,405, 7,456,273, and 7,183,395 the TEE sequences of each of which are incorporated herein by reference. In another embodiment, the TEE in the 5'-UTR of the polynucleotides (e.g., mRNA) of the present disclosure may include a 5-30 nucleotide fragment, a 5-25 nucleotide fragment, a 5-20 nucleotide fragment, a 5-15 nucleotide fragment, a 5-10 nucleotide fragment of the TEE sequences disclosed in US Patent Publication Nos. 2009/0226470, 2007/0048776, 2013/0177581 and 2011/0124100, International Patent Publication Nos. WO1999/024595, WO2012/009644, WO2009/075886 and WO2007/025008, European Patent Publication Nos. 2610341 and 2610340, and US Patent Nos. 6,310,197, 6,849,405, 7,456,273, and 7,183,395; the 15 TEE sequences of each of which are incorporated herein by reference.

[00382] In certain cases, the TEE in the 5'-UTR of the polynucleotides (e.g., mRNA) of the present disclosure may include at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 99% or more than 99% of the TEE sequences disclosed in Chappell et al. (Proc. Natl. Acad. Sci. USA 101:9590-9594, 2004) and Zhou et al. (PNAS 102:6273-6278, 2005), in Supplemental Table 1 and in Supplemental Table 2 disclosed by Wellensiek et al (Genome-wide profiling of human cap-independent translation-enhancing elements, Nature Methods, 2013; DOI:10.1038/NMETH.2522); the TEE sequences of each of which are herein incorporated by reference. In another embodiment, the TEE in the 5'-UTR of the polynucleotides (e.g., mRNA) of the present disclosure may include a 5-30 nucleotide fragment, a 5-25 nucleotide fragment, a 5-20 nucleotide fragment, a 5-15 nucleotide fragment, a 5-10 nucleotide fragment of the TEE sequences disclosed in Chappell et al. (Proc. Natl. Acad. Sci. USA 101:9590-9594, 2004) and

Zhou et al. (PNAS 102:6273-6278, 2005), in Supplemental Table 1 and in Supplemental Table 2 disclosed by Wellensiek et al (Genome-wide profiling of human cap-independent translation-enhancing elements, Nature Methods, 2013; DOI:10.1038/NMETH.2522); the TEE sequences of each of which is incorporated herein by reference.

5 [00383] In some cases, the TEE used in the 5'-UTR of a polynucleotide (e.g., mRNA) is an IRES sequence such as, but not limited to, those described in US Patent No. 7,468,275 and International Patent Publication No. WO2001/055369, the TEE sequences of each of which are incorporated herein by reference.

10 [00384] In some instances, the TEEs used in the 5'-UTR of a polynucleotide (e.g., mRNA) may be identified by the methods described in US Patent Publication Nos. 2007/0048776 and 2011/0124100 and International Patent Publication Nos. WO2007/025008 and WO2012/009644, the methods of each of which are incorporated herein by reference.

15 [00385] In some cases, the TEEs used in the 5'-UTR of a polynucleotide (e.g., mRNA) of the present disclosure may be a transcription regulatory element described in US Patent Nos. 7,456,273 and 7,183,395, US Patent Publication No. 2009/0093049, and International Publication No. WO2001/055371, the TEE sequences of each of which is incorporated herein by reference. The transcription regulatory elements may be identified by methods known in the art, such as, but not limited to, the methods described in US Patent Nos. 7,456,273 and 7,183,395, US Patent Publication No. 2009/0093049, and International Publication No. WO2001/055371, the methods of each of which is incorporated herein by reference.

20 [00386] In yet other instances, the TEE used in the 5'-UTR of a polynucleotide (e.g., mRNA) is a polynucleotide or portion thereof as described in US Patent Nos. 7,456,273 and 7,183,395, US Patent Publication No. 2009/0093049, and International Publication No. WO2001/055371, the TEE sequences of each of which are incorporated herein by reference.

25 [00387] The 5'-UTR including at least one TEE described herein may be incorporated in a monocistronic sequence such as, but not limited to, a vector system or a polynucleotide vector. As a non-limiting example, the vector systems and polynucleotide vectors may include those described in US Patent Nos. 7,456,273 and 7,183,395, US Patent Publication Nos. 2007/0048776, 2009/0093049 and 2011/0124100, and International Patent Publication Nos. 30 WO2007/025008 and WO2001/055371, the TEE sequences of each of which are incorporated herein by reference.

[00388] The TEEs described herein may be located in the 5'-UTR and/or the 3'-UTR of the polynucleotides (e.g., mRNA). The TEEs located in the 3'-UTR may be the same and/or different than the TEEs located in and/or described for incorporation in the 5'-UTR.

[00389] In some cases, the 3'-UTR of a polynucleotide (e.g., mRNA) may include at least 5 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18 at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55 or more than 60 TEE sequences. The TEE sequences in the 3'-UTR of the polynucleotides (e.g., mRNA) of the present disclosure may be the same or 10 different TEE sequences. The TEE sequences may be in a pattern such as ABABAB, AABBAABBAABB, or ABCABCABC, or variants thereof, repeated once, twice, or more than three times. In these patterns, each letter, A, B, or C represent a different TEE sequence at the nucleotide level.

[00390] In one instance, the 3'-UTR may include a spacer to separate two TEE sequences. 15 As a non-limiting example, the spacer may be a 15 nucleotide spacer and/or other spacers known in the art. As another non-limiting example, the 3'-UTR may include a TEE sequence-spacer module repeated at least once, at least twice, at least 3 times, at least 4 times, at least 5 times, at least 6 times, at least 7 times, at least 8 times, at least 9 times, or more than 9 times in the 3'-UTR.

[00391] In other cases, the spacer separating two TEE sequences may include other 20 sequences known in the art which may regulate the translation of the polynucleotides (e.g., mRNA) of the present disclosure such as, but not limited to, miR sequences described herein (e.g., miR binding sites and miR seeds). As a non-limiting example, each spacer used to separate two TEE sequences may include a different miR sequence or component of a miR 25 sequence (e.g., miR seed sequence).

[00392] In yet other cases, the incorporation of a miR sequence and/or a TEE sequence changes the shape of the stem loop region which may increase and/or decrease translation. (see e.g., Kedde et al. A Pumilio-induced RNA structure switch in p27-3'UTR controls miR-221 and miR-22 accessibility. *Nature Cell Biology*. 2010).

30 Stem Loops

[00393] Polynucleotides (e.g., mRNAs) may include a stem loop such as, but not limited to, a histone stem loop. The stem loop may be a nucleotide sequence that is about 25 or about 26 nucleotides in length such as, but not limited to, those as described in International Patent

Publication No. WO2013/103659, which is incorporated herein by reference. The histone stem loop may be located 3'-relative to the coding region (e.g., at the 3'-terminus of the coding region). As a non-limiting example, the stem loop may be located at the 3'-end of a polynucleotide described herein. In some cases, a polynucleotide (e.g., an mRNA) includes 5 more than one stem loop (e.g., two stem loops). Examples of stem loop sequences are described in International Patent Publication Nos. WO2012/019780 and WO201502667, the stem loop sequences of which are herein incorporated by reference. In some instances, a polynucleotide includes the stem loop sequence CAAAGGCTCTTCAGAGCCACCA (SEQ ID NO: 1). In others, a polynucleotide includes the stem loop sequence 10 CAAAGGCUCUUUCAGAGCCACCA (SEQ ID NO: 2).

[00394] A stem loop may be located in a second terminal region of a polynucleotide. As a non-limiting example, the stem loop may be located within an untranslated region (e.g., 3'-UTR) in a second terminal region.

[00395] In some cases, a polynucleotide such as, but not limited to mRNA, which 15 includes the histone stem loop may be stabilized by the addition of a 3'-stabilizing region (e.g., a 3'-stabilizing region including at least one chain terminating nucleoside). Not wishing to be bound by theory, the addition of at least one chain terminating nucleoside may slow the degradation of a polynucleotide and thus can increase the half-life of the polynucleotide.

[00396] In other cases, a polynucleotide such as, but not limited to mRNA, which 20 includes the histone stem loop may be stabilized by an alteration to the 3'-region of the polynucleotide that can prevent and/or inhibit the addition of oligo(U) (see e.g., International Patent Publication No. WO2013/103659).

[00397] In yet other cases, a polynucleotide such as, but not limited to mRNA, which 25 includes the histone stem loop may be stabilized by the addition of an oligonucleotide that terminates in a 3'-deoxynucleoside, 2',3'-dideoxynucleoside 3'-O-methyl nucleosides, 3'-O-ethyl nucleosides, 3'-arabinosides, and other alternative nucleosides known in the art and/or described herein.

[00398] In some instances, the polynucleotides of the present disclosure may include a histone stem loop, a poly-A region, and/or a 5'-cap structure. The histone stem loop may be 30 before and/or after the poly-A region. The polynucleotides including the histone stem loop and a poly-A region sequence may include a chain terminating nucleoside described herein.

[00399] In other instances, the polynucleotides of the present disclosure may include a histone stem loop and a 5'-cap structure. The 5'-cap structure may include, but is not limited to, those described herein and/or known in the art.

5 **[00400]** In some cases, the conserved stem loop region may include a miR sequence described herein. As a non-limiting example, the stem loop region may include the seed sequence of a miR sequence described herein. In another non-limiting example, the stem loop region may include a miR-122 seed sequence.

[00401] In certain instances, the conserved stem loop region may include a miR sequence described herein and may also include a TEE sequence.

10 **[00402]** In some cases, the incorporation of a miR sequence and/or a TEE sequence changes the shape of the stem loop region which may increase and/or decrease translation. (see e.g., Kedde et al. A Pumilio-induced RNA structure switch in p27-3'UTR controls miR-221 and miR-22 accessibility. *Nature Cell Biology*. 2010, herein incorporated by reference in its entirety).

15 **[00403]** Polynucleotides may include at least one histone stem-loop and a poly-A region or polyadenylation signal. Non-limiting examples of polynucleotide sequences encoding for at least one histone stem-loop and a poly-A region or a polyadenylation signal are described in International Patent Publication No. WO2013/120497, WO2013/120629, WO2013/120500, WO2013/120627, WO2013/120498, WO2013/120626, WO2013/120499 and WO2013/120628, the sequences of each of which are incorporated herein by reference. In certain cases, the polynucleotide encoding for a histone stem loop and a poly-A region or a polyadenylation signal may code for a pathogen antigen or fragment thereof such as the polynucleotide sequences described in International Patent Publication No WO2013/120499 and WO2013/120628, the sequences of both of which are incorporated herein by reference. In other cases, the 20 polynucleotide encoding for a histone stem loop and a poly-A region or a polyadenylation signal may code for a therapeutic protein such as the polynucleotide sequences described in International Patent Publication No WO2013/120497 and WO2013/120629, the sequences of both of which are incorporated herein by reference. In some cases, the polynucleotide encoding for a histone stem loop and a poly-A region or a polyadenylation signal may code for a tumor 25 antigen or fragment thereof such as the polynucleotide sequences described in International Patent Publication No WO2013/120500 and WO2013/120627, the sequences of both of which are incorporated herein by reference. In other cases, the polynucleotide encoding for a histone stem loop and a poly-A region or a polyadenylation signal may code for a allergenic antigen or 30

an autoimmune self-antigen such as the polynucleotide sequences described in International Patent Publication No WO2013/120498 and WO2013/120626, the sequences of both of which are incorporated herein by reference.

5 *Poly-A Regions*

[00404] A polynucleotide or nucleic acid (e.g., an mRNA) may include a polyA sequence and/or polyadenylation signal. A polyA sequence may be comprised entirely or mostly of adenine nucleotides or analogs or derivatives thereof. A polyA sequence may be a tail located adjacent to a 3' untranslated region of a nucleic acid.

10 **[00405]** During RNA processing, a long chain of adenosine nucleotides (poly-A region) is normally added to messenger RNA (mRNA) molecules to increase the stability of the molecule. Immediately after transcription, the 3'-end of the transcript is cleaved to free a 3'-hydroxy. Then poly-A polymerase adds a chain of adenosine nucleotides to the RNA. The process, called polyadenylation, adds a poly-A region that is between 100 and 250 residues long.

15 **[00406]** Unique poly-A region lengths may provide certain advantages to the alternative polynucleotides of the present disclosure.

[00407] Generally, the length of a poly-A region of polynucleotides of the present disclosure is at least 30 nucleotides in length. In another embodiment, the poly-A region is at least 35 nucleotides in length. In another embodiment, the length is at least 40 nucleotides. In 20 another embodiment, the length is at least 45 nucleotides. In another embodiment, the length is at least 55 nucleotides. In another embodiment, the length is at least 60 nucleotides. In another embodiment, the length is at least 70 nucleotides. In another embodiment, the length is at least 80 nucleotides. In another embodiment, the length is at least 90 nucleotides. In another embodiment, the length is at least 100 nucleotides. In another embodiment, the length is at least 120 nucleotides. In another embodiment, the length is at least 140 nucleotides. In another embodiment, the length is at least 160 nucleotides. In another embodiment, the length is at least 180 nucleotides. In another embodiment, the length is at least 200 nucleotides. In another embodiment, the length is at least 250 nucleotides. In another embodiment, the length is at least 300 nucleotides. In another embodiment, the length is at least 350 nucleotides. In another 30 embodiment, the length is at least 400 nucleotides. In another embodiment, the length is at least 450 nucleotides. In another embodiment, the length is at least 500 nucleotides. In another embodiment, the length is at least 600 nucleotides. In another embodiment, the length is at least 700 nucleotides. In another embodiment, the length is at least 800 nucleotides. In another

embodiment, the length is at least 900 nucleotides. In another embodiment, the length is at least 1000 nucleotides. In another embodiment, the length is at least 1100 nucleotides. In another embodiment, the length is at least 1200 nucleotides. In another embodiment, the length is at least 1300 nucleotides. In another embodiment, the length is at least 1400 nucleotides. In 5 another embodiment, the length is at least 1500 nucleotides. In another embodiment, the length is at least 1600 nucleotides. In another embodiment, the length is at least 1700 nucleotides. In another embodiment, the length is at least 1800 nucleotides. In another embodiment, the length is at least 1900 nucleotides. In another embodiment, the length is at least 2000 nucleotides. In another embodiment, the length is at least 2500 nucleotides. In another embodiment, the length 10 is at least 3000 nucleotides.

[00408] In some instances, the poly-A region may be 80 nucleotides, 120 nucleotides, 160 nucleotides in length on an alternative polynucleotide molecule described herein.

[00409] In other instances, the poly-A region may be 20, 40, 80, 100, 120, 140 or 160 nucleotides in length on an alternative polynucleotide molecule described herein.

[00410] In some cases, the poly-A region is designed relative to the length of the overall alternative polynucleotide. This design may be based on the length of the coding region of the alternative polynucleotide, the length of a particular feature or region of the alternative polynucleotide (such as mRNA), or based on the length of the ultimate product expressed from the alternative polynucleotide. When relative to any feature of the alternative polynucleotide (e.g., other than the mRNA portion which includes the poly-A region) the poly-A region may be 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100% greater in length than the additional feature. The poly-A region may also be designed as a fraction of the alternative polynucleotide to which it belongs. In this context, the poly-A region may be 10, 20, 30, 40, 50, 60, 70, 80, or 90% or more of the total length of the construct or the total length of the construct minus the poly-A region.

[00411] In certain cases, engineered binding sites and/or the conjugation of polynucleotides (e.g., mRNA) for poly-A binding protein may be used to enhance expression. The engineered binding sites may be sensor sequences which can operate as binding sites for ligands of the local microenvironment of the polynucleotides (e.g., mRNA). As a non-limiting example, the polynucleotides (e.g., mRNA) may include at least one engineered binding site to alter the binding affinity of poly-A binding protein (PABP) and analogs thereof. The incorporation of at least one engineered binding site may increase the binding affinity of the PABP and analogs thereof.

[00412] Additionally, multiple distinct polynucleotides (e.g., mRNA) may be linked together to the PABP (poly-A binding protein) through the 3'-end using alternative nucleotides at the 3'-terminus of the poly-A region. Transfection experiments can be conducted in relevant cell lines at and protein production can be assayed by ELISA at 12 hours, 24 hours, 48 hours, 72 hours, and day 7 post-transfection. As a non-limiting example, the transfection experiments may be used to evaluate the effect on PABP or analogs thereof binding affinity as a result of the addition of at least one engineered binding site.

[00413] In certain cases, a poly-A region may be used to modulate translation initiation. While not wishing to be bound by theory, the poly-A region recruits PABP which in turn can interact with translation initiation complex and thus may be essential for protein synthesis.

[00414] In some cases, a poly-A region may also be used in the present disclosure to protect against 3'-5'-exonuclease digestion.

[00415] In some instances, a polynucleotide (e.g., mRNA) may include a polyA-G Quartet. The G-quartet is a cyclic hydrogen bonded array of four guanosine nucleotides that can be formed by G-rich sequences in both DNA and RNA. In this embodiment, the G-quartet is incorporated at the end of the poly-A region. The resultant polynucleotides (e.g., mRNA) may be assayed for stability, protein production and other parameters including half-life at various time points. It has been discovered that the polyA-G quartet results in protein production equivalent to at least 75% of that seen using a poly-A region of 120 nucleotides alone.

[00416] In some cases, a polynucleotide (e.g., mRNA) may include a poly-A region and may be stabilized by the addition of a 3'-stabilizing region. The polynucleotides (e.g., mRNA) with a poly-A region may further include a 5'-cap structure.

[00417] In other cases, a polynucleotide (e.g., mRNA) may include a poly-A-G Quartet. The polynucleotides (e.g., mRNA) with a poly-A-G Quartet may further include a 5'-cap structure.

[00418] In some cases, the 3'-stabilizing region which may be used to stabilize a polynucleotide (e.g., mRNA) including a poly-A region or poly-A-G Quartet may be, but is not limited to, those described in International Patent Publication No. WO2013/103659, the poly-A regions and poly-A-G Quartets of which are incorporated herein by reference. In other cases, the 3'-stabilizing region which may be used with the polynucleotides of the present disclosure include a chain termination nucleoside such as 3'-deoxyadenosine (cordycepin), 3'-deoxyuridine, 3'-deoxycytosine, 3'-deoxyguanosine, 3'-deoxythymine, 2',3'-dideoxynucleosides, such as 2',3'-dideoxyadenosine, 2',3'-dideoxyuridine, 2',3'-

dideoxycytosine, 2',3'- dideoxyguanosine, 2',3'-dideoxythymine, a 2'-deoxynucleoside, or an O-methylnucleoside.

[00419] In other cases, a polynucleotide such as, but not limited to mRNA, which includes a polyA region or a poly-A-G Quartet may be stabilized by an alteration to the 3'-region of the polynucleotide that can prevent and/or inhibit the addition of oligo(U) (see e.g., International Patent Publication No. WO2013/103659).

[00420] In yet other instances, a polynucleotide such as, but not limited to mRNA, which includes a poly-A region or a poly-A-G Quartet may be stabilized by the addition of an oligonucleotide that terminates in a 3'-deoxynucleoside, 2',3'-dideoxynucleoside 3'-O-methylnucleosides, 3'-O-ethylnucleosides, 3'-arabinosides, and other alternative nucleosides known in the art and/or described herein.

Chain terminating nucleosides

[00421] A nucleic acid may include a chain terminating nucleoside. For example, a chain terminating nucleoside may include those nucleosides deoxygenated at the 2' and/or 3' positions of their sugar group. Such species may include 3'-deoxyadenosine (cordycepin), 3'-deoxyuridine, 3'-deoxycytosine, 3'-deoxyguanosine, 3'-deoxythymine, and 2',3'-dideoxynucleosides, such as 2',3'-dideoxyadenosine, 2',3'-dideoxyuridine, 2',3'-dideoxycytosine, 2',3'-dideoxyguanosine, and 2',3'-dideoxythymine.

20 Other components

[00422] A nanoparticle composition may include one or more components in addition to those described in the preceding sections. For example, a nanoparticle composition may include one or more small hydrophobic molecules such as a vitamin (e.g., vitamin A or vitamin E) or a sterol.

[00423] Nanoparticle compositions may also include one or more permeability enhancer molecules, carbohydrates, polymers, surface altering agents, or other components. A permeability enhancer molecule may be a molecule described by U.S. patent application publication No. 2005/0222064, for example. Carbohydrates may include simple sugars (e.g., glucose) and polysaccharides (e.g., glycogen and derivatives and analogs thereof).

[00424] A polymer may be included in and/or used to encapsulate or partially encapsulate a nanoparticle composition. A polymer may be biodegradable and/or biocompatible. A polymer may be selected from, but is not limited to, polyamines, polyethers, polyamides, polyesters, polycarbamates, polyureas, polycarbonates, polystyrenes, polyimides, polysulfones,

polyurethanes, polyacetylenes, polyethylenes, polyethyleneimines, polyisocyanates, polyacrylates, polymethacrylates, polyacrylonitriles, and polyarylates. For example, a polymer may include poly(caprolactone) (PCL), ethylene vinyl acetate polymer (EVA), poly(lactic acid) (PLA), poly(L-lactic acid) (PLLA), poly(glycolic acid) (PGA), poly(lactic acid-co-glycolic acid) (PLGA), poly(L-lactic acid-co-glycolic acid) (PLLGA), poly(D,L-lactide) (PDLA), poly(L-lactide) (PLLA), poly(D,L-lactide-co-caprolactone), poly(D,L-lactide-co-caprolactone-co-glycolide), poly(D,L-lactide-co-PEO-co-D,L-lactide), poly(D,L-lactide-co-PPO-co-D,L-lactide), polyalkyl cyanoacralate, polyurethane, poly-L-lysine (PLL), hydroxypropyl methacrylate (HPMA), polyethyleneglycol, poly-L-glutamic acid, poly(hydroxy acids), polyanhydrides, 10 polyorthoesters, poly(ester amides), polyamides, poly(ester ethers), polycarbonates, polyalkylenes such as polyethylene and polypropylene, polyalkylene glycols such as poly(ethylene glycol) (PEG), polyalkylene oxides (PEO), polyalkylene terephthalates such as poly(ethylene terephthalate), polyvinyl alcohols (PVA), polyvinyl ethers, polyvinyl esters such as poly(vinyl acetate), polyvinyl halides such as poly(vinyl chloride) (PVC), 15 polyvinylpyrrolidone (PVP), polysiloxanes, polystyrene (PS), polyurethanes, derivatized celluloses such as alkyl celluloses, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, hydroxypropylcellulose, carboxymethylcellulose, polymers of acrylic acids, such as poly(methyl(meth)acrylate) (PMMA), poly(ethyl(meth)acrylate), poly(butyl(meth)acrylate), poly(isobutyl(meth)acrylate), poly(hexyl(meth)acrylate), 20 poly(isodecyl(meth)acrylate), poly(lauryl(meth)acrylate), poly(phenyl(meth)acrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), poly(octadecyl acrylate) and copolymers and mixtures thereof, polydioxanone and its copolymers, polyhydroxyalkanoates, polypropylene fumarate, polyoxymethylene, poloxamers, polyoxamines, poly(ortho)esters, poly(butyric acid), poly(valeric acid), poly(lactide-co-caprolactone), trimethylene carbonate, poly(N-acryloylmorpholine) (PAcM), poly(2-methyl-2-oxazoline) (PMOX), poly(2-ethyl-2-oxazoline) (PEOZ), and polyglycerol.

[00425] Surface altering agents may include, but are not limited to, anionic proteins (e.g., bovine serum albumin), surfactants (e.g., cationic surfactants such as dimethyldioctadecylammonium bromide), sugars or sugar derivatives (e.g., cyclodextrin), nucleic acids, polymers (e.g., heparin, polyethylene glycol, and poloxamer), mucolytic agents (e.g., acetylcysteine, mugwort, bromelain, papain, clerodendrum, bromhexine, carbocisteine, eprazinone, mesna, ambroxol, sofrerol, domiodol, letosteine, stepronin, tiopronin, gelsolin, thymosin β 4, dornase alfa, neltenexine, and erdosteine), and DNases (e.g., rhDNase). A surface altering agent may be

disposed within a nanoparticle and/or on the surface of a nanoparticle composition (e.g., by coating, adsorption, covalent linkage, or other process).

[00426] A nanoparticle composition may also comprise one or more functionalized lipids. For example, a lipid may be functionalized with an alkyne group that, when exposed to an azide under appropriate reaction conditions, may undergo a cycloaddition reaction. In particular, a lipid bilayer may be functionalized in this fashion with one or more groups useful in facilitating membrane permeation, cellular recognition, or imaging. The surface of a nanoparticle composition may also be conjugated with one or more useful antibodies. Functional groups and conjugates useful in targeted cell delivery, imaging, and membrane permeation are well known in the art.

[00427] In addition to these components, nanoparticle compositions may include any substance useful in pharmaceutical compositions. For example, the nanoparticle composition may include one or more pharmaceutically acceptable excipients or accessory ingredients such as, but not limited to, one or more solvents, dispersion media, diluents, dispersion aids, suspension aids, granulating aids, disintegrants, fillers, glidants, liquid vehicles, binders, surface active agents, isotonic agents, thickening or emulsifying agents, buffering agents, lubricating agents, oils, preservatives, and other species. Excipients such as waxes, butters, coloring agents, coating agents, flavorings, and perfuming agents may also be included. Pharmaceutically acceptable excipients are well known in the art (see for example Remington's *The Science and Practice of Pharmacy*, 21st Edition, A. R. Gennaro; Lippincott, Williams & Wilkins, Baltimore, MD, 2006).

[00428] Examples of diluents may include, but are not limited to, calcium carbonate, sodium carbonate, calcium phosphate, dicalcium phosphate, calcium sulfate, calcium hydrogen phosphate, sodium phosphate lactose, sucrose, cellulose, microcrystalline cellulose, kaolin, mannitol, sorbitol, inositol, sodium chloride, dry starch, cornstarch, powdered sugar, and/or combinations thereof. Granulating and dispersing agents may be selected from the non-limiting list consisting of potato starch, corn starch, tapioca starch, sodium starch glycolate, clays, alginic acid, guar gum, citrus pulp, agar, bentonite, cellulose and wood products, natural sponge, cation-exchange resins, calcium carbonate, silicates, sodium carbonate, cross-linked poly(vinyl-pyrrolidone) (crospovidone), sodium carboxymethyl starch (sodium starch glycolate), carboxymethyl cellulose, cross-linked sodium carboxymethyl cellulose (croscarmellose), methylcellulose, pregelatinized starch (starch 1500), microcrystalline starch, water insoluble

starch, calcium carboxymethyl cellulose, magnesium aluminum silicate (VEEGUM®), sodium lauryl sulfate, quaternary ammonium compounds, and/or combinations thereof.

[00429] Surface active agents and/or emulsifiers may include, but are not limited to,

natural emulsifiers (e.g. acacia, agar, alginic acid, sodium alginate, tragacanth, chondrus,

5 cholesterol, xanthan, pectin, gelatin, egg yolk, casein, wool fat, cholesterol, wax, and lecithin),

colloidal clays (e.g. bentonite [aluminum silicate] and VEEGUM® [magnesium aluminum silicate]), long chain amino acid derivatives, high molecular weight alcohols (e.g. stearyl

alcohol, cetyl alcohol, oleyl alcohol, triacetin monostearate, ethylene glycol distearate, glyceryl

monostearate, and propylene glycol monostearate, polyvinyl alcohol), carbomers (e.g. carboxy

10 polymethylene, polyacrylic acid, acrylic acid polymer, and carboxyvinyl polymer), carrageenan,

cellulosic derivatives (e.g. carboxymethylcellulose sodium, powdered cellulose, hydroxymethyl

cellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose, methylcellulose), sorbitan

fatty acid esters (e.g. polyoxyethylene sorbitan monolaurate [TWEEN®20], polyoxyethylene

sorbitan [TWEEN® 60], polyoxyethylene sorbitan monooleate [TWEEN®80], sorbitan

15 monopalmitate [SPAN®40], sorbitan monostearate [SPAN®60], sorbitan tristearate

[SPAN®65], glyceryl monooleate, sorbitan monooleate [SPAN®80]), polyoxyethylene esters

(e.g. polyoxyethylene monostearate [MYRJ® 45], polyoxyethylene hydrogenated castor oil,

polyethoxylated castor oil, polyoxymethylene stearate, and SOLUTOL®), sucrose fatty acid

esters, polyethylene glycol fatty acid esters (e.g. CREMOPHOR®), polyoxyethylene ethers,

20 (e.g. polyoxyethylene lauryl ether [BRIJ® 30]), poly(vinyl-pyrrolidone), diethylene glycol

monolaurate, triethanolamine oleate, sodium oleate, potassium oleate, ethyl oleate, oleic acid,

ethyl laurate, sodium lauryl sulfate, PLURONIC®F 68, POLOXAMER® 188, cetrimonium

bromide, cetylpyridinium chloride, benzalkonium chloride, docusate sodium, and/or

combinations thereof.

25 **[00430]** A binding agent may be starch (e.g. cornstarch and starch paste); gelatin; sugars

(e.g. sucrose, glucose, dextrose, dextrin, molasses, lactose, lactitol, mannitol); natural and

synthetic gums (e.g. acacia, sodium alginate, extract of Irish moss, panwar gum, ghatti gum,

mucilage of isapol husks, carboxymethylcellulose, methylcellulose, ethylcellulose,

hydroxyethylcellulose, hydroxypropyl cellulose, hydroxypropyl methylcellulose,

30 microcrystalline cellulose, cellulose acetate, poly(vinyl-pyrrolidone), magnesium aluminum

silicate (VEEGUM®), and larch arabogalactan); alginates; polyethylene oxide; polyethylene

glycol; inorganic calcium salts; silicic acid; polymethacrylates; waxes; water; alcohol; and

combinations thereof, or any other suitable binding agent.

[00431] Examples of preservatives may include, but are not limited to, antioxidants, chelating agents, antimicrobial preservatives, antifungal preservatives, alcohol preservatives, acidic preservatives, and/or other preservatives. Examples of antioxidants include, but are not limited to, alpha tocopherol, ascorbic acid, acorbyl palmitate, butylated hydroxyanisole,

5 butylated hydroxytoluene, monothioglycerol, potassium metabisulfite, propionic acid, propyl gallate, sodium ascorbate, sodium bisulfite, sodium metabisulfite, and/or sodium sulfite.

Examples of chelating agents include ethylenediaminetetraacetic acid (EDTA), citric acid monohydrate, disodium edetate, dipotassium edetate, edetic acid, fumaric acid, malic acid, phosphoric acid, sodium edetate, tartaric acid, and/or trisodium edetate. Examples of

10 antimicrobial preservatives include, but are not limited to, benzalkonium chloride, benzethonium chloride, benzyl alcohol, bronopol, cetrimide, cetylpyridinium chloride, chlorhexidine, chlorobutanol, chlorocresol, chloroxylenol, cresol, ethyl alcohol, glycerin, hexetidine, imidurea, phenol, phenoxyethanol, phenylethyl alcohol, phenylmercuric nitrate, propylene glycol, and/or thimerosal. Examples of antifungal preservatives include, but are not limited to, butyl paraben,

15 methyl paraben, ethyl paraben, propyl paraben, benzoic acid, hydroxybenzoic acid, potassium benzoate, potassium sorbate, sodium benzoate, sodium propionate, and/or sorbic acid.

Examples of alcohol preservatives include, but are not limited to, ethanol, polyethylene glycol, benzyl alcohol, phenol, phenolic compounds, bisphenol, chlorobutanol, hydroxybenzoate, and/or phenylethyl alcohol. Examples of acidic preservatives include, but are not limited to, vitamin A,

20 vitamin C, vitamin E, beta-carotene, citric acid, acetic acid, dehydroascorbic acid, ascorbic acid, sorbic acid, and/or phytic acid. Other preservatives include, but are not limited to, tocopherol, tocopherol acetate, detersoxime mesylate, cetrimide, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), ethylenediamine, sodium lauryl sulfate (SLS), sodium lauryl ether sulfate (SLES), sodium bisulfite, sodium metabisulfite, potassium sulfite, potassium

25 metabisulfite, GLYDANT PLUS®, PHENONIP®, methylparaben, GERMALL® 115, GERMABEN®II, NEOLONE™, KATHON™, and/or EUXYL®.

[00432] Examples of buffering agents include, but are not limited to, citrate buffer solutions, acetate buffer solutions, phosphate buffer solutions, ammonium chloride, calcium carbonate, calcium chloride, calcium citrate, calcium glubionate, calcium gluceptate, calcium gluconate, d-gluconic acid, calcium glycerophosphate, calcium lactate, calcium lactobionate, propanoic acid, calcium levulinate, pentanoic acid, dibasic calcium phosphate, phosphoric acid, tribasic calcium phosphate, calcium hydroxide phosphate, potassium acetate, potassium chloride, potassium gluconate, potassium mixtures, dibasic potassium phosphate, monobasic

potassium phosphate, potassium phosphate mixtures, sodium acetate, sodium bicarbonate, sodium chloride, sodium citrate, sodium lactate, dibasic sodium phosphate, monobasic sodium phosphate, sodium phosphate mixtures, tromethamine, amino-sulfonate buffers (e.g. HEPES), magnesium hydroxide, aluminum hydroxide, alginic acid, pyrogen-free water, isotonic saline, 5 Ringer's solution, ethyl alcohol, and/or combinations thereof. Lubricating agents may be selected from the non-limiting group consisting of magnesium stearate, calcium stearate, stearic acid, silica, talc, malt, glyceryl behenate, hydrogenated vegetable oils, polyethylene glycol, sodium benzoate, sodium acetate, sodium chloride, leucine, magnesium lauryl sulfate, sodium lauryl sulfate, and combinations thereof.

10 [00433] Examples of oils include, but are not limited to, almond, apricot kernel, avocado, babassu, bergamot, black current seed, borage, cade, camomile, canola, caraway, carnauba, castor, cinnamon, cocoa butter, coconut, cod liver, coffee, corn, cotton seed, emu, eucalyptus, evening primrose, fish, flaxseed, geraniol, gourd, grape seed, hazel nut, hyssop, isopropyl myristate, jojoba, kukui nut, lavandin, lavender, lemon, litsea cubeba, macadamia nut, mallow, 15 mango seed, meadowfoam seed, mink, nutmeg, olive, orange, orange roughy, palm, palm kernel, peach kernel, peanut, poppy seed, pumpkin seed, rapeseed, rice bran, rosemary, safflower, sandalwood, sasquana, savoury, sea buckthorn, sesame, shea butter, silicone, soy bean, sunflower, tea tree, thistle, tsubaki, vetiver, walnut, and wheat germ oils as well as butyl stearate, caprylic triglyceride, capric triglyceride, cyclomethicone, diethyl sebacate, dimethicone 20 360, simethicone, isopropyl myristate, mineral oil, octyldodecanol, oleyl alcohol, silicone oil, and/or combinations thereof.

Formulations

[00434] Nanoparticle compositions may include a lipid component and one or more 25 additional components, such as a therapeutic and/or prophylactic agent. A nanoparticle composition may be designed for one or more specific applications or targets. The elements of a nanoparticle composition may be selected based on a particular application or target, and/or based on the efficacy, toxicity, expense, ease of use, availability, or other feature of one or more elements. Similarly, the particular formulation of a nanoparticle composition may be selected 30 for a particular application or target according to, for example, the efficacy and toxicity of particular combinations of elements.

[00435] The lipid component of a nanoparticle composition may include, for example, a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I),

(19-II), (20-I) and (21-I), a phospholipid (such as an unsaturated lipid, e.g., DOPE or DSPC), a PEG lipid, and a structural lipid. The elements of the lipid component may be provided in specific fractions.

[00436] In some embodiments, the lipid component of a nanoparticle composition 5 includes a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), a phospholipid, a PEG lipid, and a structural lipid. In certain embodiments, the lipid component of the nanoparticle composition includes about 30 mol % to about 60 mol % compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), about 0 mol % to about 30 mol % 10 phospholipid, about 18.5 mol % to about 48.5 mol % structural lipid, and about 0 mol % to about 10 mol % of PEG lipid, provided that the total mol % does not exceed 100%. In some embodiments, the lipid component of the nanoparticle composition includes about 35 mol % to about 55 mol % compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), about 5 mol % to about 25 mol % 15 phospholipid, about 30 mol % to about 40 mol % structural lipid, and about 0 mol % to about 10 mol % of PEG lipid. In certain embodiments, the lipid component includes about 50 mol % said compound, about 10 mol % phospholipid, about 38.5 mol % structural lipid, and about 1.5 mol % of PEG lipid. In other embodiments, the lipid component includes about 40 mol % said compound, about 20 mol % phospholipid, about 38.5 mol % structural lipid, and about 1.5 mol 20 % of PEG lipid. In some embodiments, the phospholipid may be DOPE or DSPC. In other embodiments, the PEG lipid may be PEG-DMG and/or the structural lipid may be cholesterol.

[00437] Nanoparticle compositions may be designed for one or more specific applications or targets. For example, a nanoparticle composition may be designed to deliver a therapeutic and/or prophylactic agent such as an RNA to a particular cell, tissue, organ, or system or group 25 thereof in a mammal's body. Physiochemical properties of nanoparticle compositions may be altered in order to increase selectivity for particular bodily targets. For instance, particle sizes may be adjusted based on the fenestration sizes of different organs. The therapeutic and/or prophylactic agent included in a nanoparticle composition may also be selected based on the desired delivery target or targets. For example, a therapeutic and/or prophylactic agent may be 30 selected for a particular indication, condition, disease, or disorder and/or for delivery to a particular cell, tissue, organ, or system or group thereof (e.g., localized or specific delivery). In certain embodiments, a nanoparticle composition may include an mRNA encoding a polypeptide of interest capable of being translated within a cell to produce the polypeptide of interest. Such

a composition may be designed to be specifically delivered to a particular organ. In certain embodiments, a composition may be designed to be specifically delivered to a mammalian liver.

[00438] The amount of a therapeutic and/or prophylactic agent in a nanoparticle composition may depend on the size, composition, desired target and/or application, or other properties of the nanoparticle composition as well as on the properties of the therapeutic and/or prophylactic agent. For example, the amount of an RNA useful in a nanoparticle composition may depend on the size, sequence, and other characteristics of the RNA. The relative amounts of a therapeutic and/or prophylactic agent and other elements (e.g., lipids) in a nanoparticle composition may also vary. In some embodiments, the wt/wt ratio of the lipid component to a therapeutic and/or prophylactic agent in a nanoparticle composition may be from about 5:1 to about 60:1, such as 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 25:1, 30:1, 35:1, 40:1, 45:1, 50:1, and 60:1. For example, the wt/wt ratio of the lipid component to a therapeutic and/or prophylactic agent may be from about 10:1 to about 40:1. In certain embodiments, the wt/wt ratio is about 20:1. The amount of a therapeutic and/or prophylactic agent in a nanoparticle composition may, for example, be measured using absorption spectroscopy (e.g., ultraviolet-visible spectroscopy).

[00439] In some embodiments, a nanoparticle composition includes one or more RNAs, and the one or more RNAs, lipids, and amounts thereof may be selected to provide a specific N:P ratio. The N:P ratio of the composition refers to the molar ratio of nitrogen atoms in one or more lipids to the number of phosphate groups in an RNA. In general, a lower N:P ratio is preferred. The one or more RNA, lipids, and amounts thereof may be selected to provide an N:P ratio from about 2:1 to about 30:1, such as 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 12:1, 14:1, 16:1, 18:1, 20:1, 22:1, 24:1, 26:1, 28:1, or 30:1. In certain embodiments, the N:P ratio may be from about 2:1 to about 8:1. In other embodiments, the N:P ratio is from about 5:1 to about 8:1. For example, the N:P ratio may be about 5.0:1, about 5.5:1, about 5.67:1, about 6.0:1, about 6.5:1, or about 7.0:1. For example, the N:P ratio may be about 5.67:1.

Physical properties

[00440] The characteristics of a nanoparticle composition may depend on the components thereof. For example, a nanoparticle composition including cholesterol as a structural lipid may have different characteristics than a nanoparticle composition that includes a different structural lipid. Similarly, the characteristics of a nanoparticle composition may depend on the absolute or relative amounts of its components. For instance, a nanoparticle composition including a higher

molar fraction of a phospholipid may have different characteristics than a nanoparticle composition including a lower molar fraction of a phospholipid. Characteristics may also vary depending on the method and conditions of preparation of the nanoparticle composition.

[00441] Nanoparticle compositions may be characterized by a variety of methods. For example, microscopy (e.g., transmission electron microscopy or scanning electron microscopy) may be used to examine the morphology and size distribution of a nanoparticle composition. Dynamic light scattering or potentiometry (e.g., potentiometric titrations) may be used to measure zeta potentials. Dynamic light scattering may also be utilized to determine particle sizes. Instruments such as the Zetasizer Nano ZS (Malvern Instruments Ltd, Malvern, 10 Worcestershire, UK) may also be used to measure multiple characteristics of a nanoparticle composition, such as particle size, polydispersity index, and zeta potential.

[00442] The mean size of a nanoparticle composition may be between 10s of nm and 100s of nm. For example, the mean size may be from about 40 nm to about 150 nm, such as about 40 nm, 45 nm, 50 nm, 55 nm, 60 nm, 65 nm, 70 nm, 75 nm, 80 nm, 85 nm, 90 nm, 95 nm, 100 nm, 15 105 nm, 110 nm, 115 nm, 120 nm, 125 nm, 130 nm, 135 nm, 140 nm, 145 nm, or 150 nm. In some embodiments, the mean size of a nanoparticle composition may be from about 50 nm to about 100 nm, from about 50 nm to about 90 nm, from about 50 nm to about 80 nm, from about 50 nm to about 70 nm, from about 50 nm to about 60 nm, from about 60 nm to about 100 nm, from about 60 nm to about 90 nm, from about 60 nm to about 80 nm, from about 60 nm to about 20 70 nm, from about 70 nm to about 100 nm, from about 70 nm to about 90 nm, from about 70 nm to about 80 nm, from about 80 nm to about 100 nm, from about 80 nm to about 90 nm, or from about 90 nm to about 100 nm. In certain embodiments, the mean size of a nanoparticle composition may be from about 70 nm to about 100 nm. In some embodiments, the mean size may be about 80 nm. In other embodiments, the mean size may be about 100 nm.

[00443] A nanoparticle composition may be relatively homogenous. A polydispersity index may be used to indicate the homogeneity of a nanoparticle composition, e.g., the particle size distribution of the nanoparticle compositions. A small (e.g., less than 0.3) polydispersity index generally indicates a narrow particle size distribution. A nanoparticle composition may have a polydispersity index from about 0 to about 0.25, such as 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 30 0.07, 0.08, 0.09, 0.10, 0.11, 0.12, 0.13, 0.14, 0.15, 0.16, 0.17, 0.18, 0.19, 0.20, 0.21, 0.22, 0.23, 0.24, or 0.25. In some embodiments, the polydispersity index of a nanoparticle composition may be from about 0.10 to about 0.20.

[00444] The zeta potential of a nanoparticle composition may be used to indicate the electrokinetic potential of the composition. For example, the zeta potential may describe the surface charge of a nanoparticle composition. Nanoparticle compositions with relatively low charges, positive or negative, are generally desirable, as more highly charged species may interact undesirably with cells, tissues, and other elements in the body. In some embodiments, the zeta potential of a nanoparticle composition may be from about -10 mV to about +20 mV, from about -10 mV to about +15 mV, from about -10 mV to about +10 mV, from about -10 mV to about +5 mV, from about -10 mV to about 0 mV, from about -10 mV to about -5 mV, from about -5 mV to about +20 mV, from about -5 mV to about +15 mV, from about -5 mV to about +10 mV, from about -5 mV to about +5 mV, from about -5 mV to about 0 mV, from about 0 mV to about +20 mV, from about 0 mV to about +15 mV, from about 0 mV to about +10 mV, from about 0 mV to about +5 mV, from about +5 mV to about +20 mV, from about +5 mV to about +15 mV, or from about +5 mV to about +10 mV.

[00445] The efficiency of encapsulation of a therapeutic and/or prophylactic agent describes the amount of therapeutic and/or prophylactic agent that is encapsulated or otherwise associated with a nanoparticle composition after preparation, relative to the initial amount provided. The encapsulation efficiency is desirably high (e.g., close to 100%). The encapsulation efficiency may be measured, for example, by comparing the amount of therapeutic and/or prophylactic agent in a solution containing the nanoparticle composition before and after breaking up the nanoparticle composition with one or more organic solvents or detergents. Fluorescence may be used to measure the amount of free therapeutic and/or prophylactic agent (e.g., RNA) in a solution. For the nanoparticle compositions described herein, the encapsulation efficiency of a therapeutic and/or prophylactic agent may be at least 50%, for example 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%. In some embodiments, the encapsulation efficiency may be at least 80%. In certain embodiments, the encapsulation efficiency may be at least 90%.

[00446] A nanoparticle composition may optionally comprise one or more coatings. For example, a nanoparticle composition may be formulated in a capsule, film, or tablet having a coating. A capsule, film, or tablet including a composition described herein may have any useful size, tensile strength, hardness, or density.

Pharmaceutical compositions

[00447] Nanoparticle compositions may be formulated in whole or in part as pharmaceutical compositions. Pharmaceutical compositions may include one or more nanoparticle compositions. For example, a pharmaceutical composition may include one or more nanoparticle compositions including one or more different therapeutic and/or prophylactic agents. Pharmaceutical compositions may further include one or more pharmaceutically acceptable excipients or accessory ingredients such as those described herein. General guidelines for the formulation and manufacture of pharmaceutical compositions and agents are available, for example, in Remington's *The Science and Practice of Pharmacy*, 21st Edition, A. R. Gennaro; Lippincott, Williams & Wilkins, Baltimore, MD, 2006. Conventional excipients and accessory ingredients may be used in any pharmaceutical composition, except insofar as any conventional excipient or accessory ingredient may be incompatible with one or more components of a nanoparticle composition. An excipient or accessory ingredient may be incompatible with a component of a nanoparticle composition if its combination with the component may result in any undesirable biological effect or otherwise deleterious effect.

[00448] In some embodiments, one or more excipients or accessory ingredients may make up greater than 50% of the total mass or volume of a pharmaceutical composition including a nanoparticle composition. For example, the one or more excipients or accessory ingredients may make up 50%, 60%, 70%, 80%, 90%, or more of a pharmaceutical composition. In some embodiments, a pharmaceutically acceptable excipient is at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% pure. In some embodiments, an excipient is approved for use in humans and for veterinary use. In some embodiments, an excipient is approved by United States Food and Drug Administration. In some embodiments, an excipient is pharmaceutical grade. In some embodiments, an excipient meets the standards of the United States Pharmacopoeia (USP), the European Pharmacopoeia (EP), the British Pharmacopoeia, and/or the International Pharmacopoeia.

[00449] Relative amounts of the one or more nanoparticle compositions, the one or more pharmaceutically acceptable excipients, and/or any additional ingredients in a pharmaceutical composition in accordance with the present disclosure will vary, depending upon the identity, size, and/or condition of the subject treated and further depending upon the route by which the composition is to be administered. By way of example, a pharmaceutical composition may comprise between 0.1% and 100% (wt/wt) of one or more nanoparticle compositions.

[00450] In certain embodiments, the nanoparticle compositions and/or pharmaceutical compositions of the disclosure are refrigerated or frozen for storage and/or shipment (e.g., being

stored at a temperature of 4 °C or lower, such as a temperature between about -150 °C and about 0 °C or between about -80 °C and about -20 °C (e.g., about -5 °C, -10 °C, -15 °C, -20 °C, -25 °C, -30 °C, -40 °C, -50 °C, -60 °C, -70 °C, -80 °C, -90 °C, -130 °C or -150 °C). For example, the pharmaceutical composition comprising a compound of any of Formulae (I)-(IV) is a

5 solution that is refrigerated for storage and/or shipment at, for example, about -20 °C, -30 °C, -40 °C, -50 °C, -60 °C, -70 °C, or -80 °C. In certain embodiments, the disclosure also relates to a method of increasing stability of the nanoparticle compositions and/or pharmaceutical compositions comprising a compound of any of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) by storing the nanoparticle compositions and/or pharmaceutical compositions at a temperature of 4 °C or lower, such as a temperature between about -150 °C and about 0 °C or between about -80 °C and about -20 °C, e.g., about -5 °C, -10 °C, -15 °C, -20 °C, -25 °C, -30 °C, -40 °C, -50 °C, -60 °C, -70 °C, -80 °C, -90 °C, -130 °C or -150 °C). For example, the nanoparticle compositions and/or pharmaceutical compositions disclosed herein are stable for about at least 1 week, at least 2 weeks, at least 3 weeks, at least 4 weeks, at least 5 weeks, at least 6 weeks, at least 1 month, at least 2 months, at least 4 months, at least 6 months, at least 8 months, at least 10 months, at least 12 months, at least 14 months, at least 16 months, at least 18 months, at least 20 months, at least 22 months, or at least 24 months, e.g., at a temperature of 4 °C or lower (e.g., between about 4 °C and -20 °C). In one embodiment, the formulation is stabilized for at least 4 weeks at about 4 °C. In certain embodiments, the pharmaceutical composition of the disclosure comprises a nanoparticle composition disclosed herein and a pharmaceutically acceptable carrier selected from one or more of Tris, an acetate (e.g., sodium acetate), an citrate (e.g., sodium citrate), saline, PBS, and sucrose. In certain embodiments, the pharmaceutical composition of the disclosure has a pH value between about 7 and 8 (e.g., 6.8, 6.9, 7.0, 7.1, 7.2, 7.3, 7.4, 7.5, 7.6, 7.7, 7.8, 7.9 or 8.0, or between 7.5 and 8 or between 7 and 7.8). For example, a pharmaceutical composition of the disclosure comprises a nanoparticle composition disclosed herein, Tris, saline and sucrose, and has a pH of about 7.5-8, which is suitable for storage and/or shipment at, for example, about -20 °C. For example, a pharmaceutical composition of the disclosure comprises a nanoparticle composition disclosed herein and PBS and has a pH of about 7-7.8, suitable for storage and/or shipment at, for example, about 4 °C or lower. “Stability,” “stabilized,” and “stable” in the context of the present disclosure refers to the resistance of nanoparticle compositions and/or pharmaceutical compositions disclosed herein to chemical or physical changes (e.g.,

10

15

20

25

30

degradation, particle size change, aggregation, change in encapsulation, etc.) under given manufacturing, preparation, transportation, storage and/or in-use conditions, e.g., when stress is applied such as shear force, freeze/thaw stress, etc.

[00451] Nanoparticle compositions and/or pharmaceutical compositions including one or more nanoparticle compositions may be administered to any patient or subject, including those patients or subjects that may benefit from a therapeutic effect provided by the delivery of a therapeutic and/or prophylactic agent to one or more particular cells, tissues, organs, or systems or groups thereof, such as the renal system. Although the descriptions provided herein of nanoparticle compositions and pharmaceutical compositions including nanoparticle

compositions are principally directed to compositions which are suitable for administration to humans, it will be understood by the skilled artisan that such compositions are generally suitable for administration to any other mammal. Modification of compositions suitable for administration to humans in order to render the compositions suitable for administration to various animals is well understood, and the ordinarily skilled veterinary pharmacologist can design and/or perform such modification with merely ordinary, if any, experimentation.

Subjects to which administration of the compositions is contemplated include, but are not limited to, humans, other primates, and other mammals, including commercially relevant mammals such as cattle, pigs, horses, sheep, cats, dogs, mice, and/or rats.

[00452] A pharmaceutical composition including one or more nanoparticle compositions may be prepared by any method known or hereafter developed in the art of pharmacology. In general, such preparatory methods include bringing the active ingredient into association with an excipient and/or one or more other accessory ingredients, and then, if desirable or necessary, dividing, shaping, and/or packaging the product into a desired single- or multi-dose unit.

[00453] A pharmaceutical composition in accordance with the present disclosure may be prepared, packaged, and/or sold in bulk, as a single unit dose, and/or as a plurality of single unit doses. As used herein, a “unit dose” is discrete amount of the pharmaceutical composition comprising a predetermined amount of the active ingredient (e.g., nanoparticle composition). The amount of the active ingredient is generally equal to the dosage of the active ingredient which would be administered to a subject and/or a convenient fraction of such a dosage such as, for example, one-half or one-third of such a dosage.

[00454] Pharmaceutical compositions may be prepared in a variety of forms suitable for a variety of routes and methods of administration. For example, pharmaceutical compositions may be prepared in liquid dosage forms (e.g., emulsions, microemulsions, nanoemulsions,

solutions, suspensions, syrups, and elixirs), injectable forms, solid dosage forms (e.g., capsules, tablets, pills, powders, and granules), dosage forms for topical and/or transdermal administration (e.g., ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants, and patches), suspensions, powders, and other forms.

5 [00455] Liquid dosage forms for oral and parenteral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, nanoemulsions, solutions, suspensions, syrups, and/or elixirs. In addition to active ingredients, liquid dosage forms may comprise inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, 10 ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, oral compositions can include additional therapeutic and/or prophylactic agents, additional agents such as wetting agents, emulsifying and 15 suspending agents, sweetening, flavoring, and/or perfuming agents. In certain embodiments for parenteral administration, compositions are mixed with solubilizing agents such as Cremophor®, alcohols, oils, modified oils, glycols, polysorbates, cyclodextrins, polymers, and/or combinations thereof.

20 [00456] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing agents, wetting agents, and/or suspending agents. Sterile injectable preparations may be sterile injectable solutions, suspensions, and/or emulsions in nontoxic parenterally acceptable diluents and/or solvents, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles 25 and solvents that may be employed are water, Ringer's solution, U.S.P., and isotonic sodium chloride solution. Sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. Fatty acids such as oleic acid can be used in the preparation of injectables.

30 [00457] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, and/or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00458] In order to prolong the effect of an active ingredient, it is often desirable to slow the absorption of the active ingredient from subcutaneous or intramuscular injection. This may

be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or 5 suspending the drug in an oil vehicle. Injectable depot forms are made by forming microencapsulated matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are 10 prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

[00459] Compositions for rectal or vaginal administration are typically suppositories which can be prepared by mixing compositions with suitable non-irritating excipients such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature 15 but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active ingredient.

[00460] Solid dosage forms for oral administration include capsules, tablets, pills, films, powders, and granules. In such solid dosage forms, an active ingredient is mixed with at least one inert, pharmaceutically acceptable excipient such as sodium citrate or dicalcium phosphate 20 and/or fillers or extenders (e.g. starches, lactose, sucrose, glucose, mannitol, and silicic acid), binders (e.g. carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia), humectants (e.g. glycerol), disintegrating agents (e.g. agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate), solution retarding agents 25 (e.g. paraffin), absorption accelerators (e.g. quaternary ammonium compounds), wetting agents (e.g. cetyl alcohol and glycerol monostearate), absorbents (e.g. kaolin and bentonite clay, silicates), and lubricants (e.g. talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate), and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may comprise buffering agents.

[00461] Solid compositions of a similar type may be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. Solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings 30 well known in the pharmaceutical formulating art. They may optionally comprise opacifying

agents and can be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. Solid compositions of a similar type may be employed as fillers in soft and hard-filled gelatin capsules 5 using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00462] Dosage forms for topical and/or transdermal administration of a composition may include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants, and/or patches. Generally, an active ingredient is admixed under sterile conditions with a 10 pharmaceutically acceptable excipient and/or any needed preservatives and/or buffers as may be required. Additionally, the present disclosure contemplates the use of transdermal patches, which often have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms may be prepared, for example, by dissolving and/or dispensing the compound in the proper medium. Alternatively or additionally, rate may be controlled by either 15 providing a rate controlling membrane and/or by dispersing the compound in a polymer matrix and/or gel.

[00463] Suitable devices for use in delivering intradermal pharmaceutical compositions described herein include short needle devices such as those described in U.S. Patents 4,886,499; 5,190,521; 5,328,483; 5,527,288; 4,270,537; 5,015,235; 5,141,496; and 5,417,662. Intradermal 20 compositions may be administered by devices which limit the effective penetration length of a needle into the skin, such as those described in PCT publication WO 99/34850 and functional equivalents thereof. Jet injection devices which deliver liquid compositions to the dermis via a liquid jet injector and/or via a needle which pierces the stratum corneum and produces a jet which reaches the dermis are suitable. Jet injection devices are described, for example, in U.S. 25 Patents 5,480,381; 5,599,302; 5,334,144; 5,993,412; 5,649,912; 5,569,189; 5,704,911; 5,383,851; 5,893,397; 5,466,220; 5,339,163; 5,312,335; 5,503,627; 5,064,413; 5,520,639; 4,596,556; 4,790,824; 4,941,880; 4,940,460; and PCT publications WO 97/37705 and WO 97/13537. Ballistic powder/particle delivery devices which use compressed gas to accelerate 30 vaccine in powder form through the outer layers of the skin to the dermis are suitable. Alternatively or additionally, conventional syringes may be used in the classical mantoux method of intradermal administration.

[00464] Formulations suitable for topical administration include, but are not limited to, liquid and/or semi liquid preparations such as liniments, lotions, oil in water and/or water in oil

emulsions such as creams, ointments and/or pastes, and/or solutions and/or suspensions.

Topically-administrable formulations may, for example, comprise from about 1% to about 10% (wt/wt) active ingredient, although the concentration of active ingredient may be as high as the solubility limit of the active ingredient in the solvent. Formulations for topical administration

5 may further comprise one or more of the additional ingredients described herein.

[00465] A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for pulmonary administration via the buccal cavity. Such a formulation may comprise dry particles which comprise the active ingredient. Such compositions are conveniently in the form of dry powders for administration using a device comprising a dry

10 powder reservoir to which a stream of propellant may be directed to disperse the powder and/or using a self propelling solvent/powder dispensing container such as a device comprising the active ingredient dissolved and/or suspended in a low-boiling propellant in a sealed container.

Dry powder compositions may include a solid fine powder diluent such as sugar and are conveniently provided in a unit dose form.

[00466] Low boiling propellants generally include liquid propellants having a boiling point of below 65 °F at atmospheric pressure. Generally the propellant may constitute 50% to 99.9% (wt/wt) of the composition, and active ingredient may constitute 0.1% to 20% (wt/wt) of the composition. A propellant may further comprise additional ingredients such as a liquid non-ionic and/or solid anionic surfactant and/or a solid diluent (which may have a particle size of the same order as particles comprising the active ingredient).

[00467] Pharmaceutical compositions formulated for pulmonary delivery may provide an active ingredient in the form of droplets of a solution and/or suspension. Such formulations may be prepared, packaged, and/or sold as aqueous and/or dilute alcoholic solutions and/or suspensions, optionally sterile, comprising active ingredient, and may conveniently be

25 administered using any nebulization and/or atomization device. Such formulations may further comprise one or more additional ingredients including, but not limited to, a flavoring agent such as saccharin sodium, a volatile oil, a buffering agent, a surface active agent, and/or a preservative such as methylhydroxybenzoate. Droplets provided by this route of administration may have an average diameter in the range from about 1 nm to about 200 nm.

[00468] Formulations described herein as being useful for pulmonary delivery are useful for intranasal delivery of a pharmaceutical composition. Another formulation suitable for intranasal administration is a coarse powder comprising the active ingredient and having an average particle from about 0.2 µm to 500 µm. Such a formulation is administered in the

manner in which snuff is taken, i.e. by rapid inhalation through the nasal passage from a container of the powder held close to the nose.

[00469] Formulations suitable for nasal administration may, for example, comprise from about as little as 0.1% (wt/wt) and as much as 100% (wt/wt) of active ingredient, and may 5 comprise one or more of the additional ingredients described herein. A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for buccal administration. Such formulations may, for example, be in the form of tablets and/or lozenges made using conventional methods, and may, for example, 0.1% to 20% (wt/wt) active ingredient, the balance comprising an orally dissolvable and/or degradable composition and, 10 optionally, one or more of the additional ingredients described herein. Alternately, formulations suitable for buccal administration may comprise a powder and/or an aerosolized and/or atomized solution and/or suspension comprising active ingredient. Such powdered, aerosolized, and/or aerosolized formulations, when dispersed, may have an average particle and/or droplet size in the range from about 0.1 nm to about 200 nm, and may further comprise one or more of any 15 additional ingredients described herein.

[00470] A pharmaceutical composition may be prepared, packaged, and/or sold in a formulation suitable for ophthalmic administration. Such formulations may, for example, be in the form of eye drops including, for example, a 0.1/1.0% (wt/wt) solution and/or suspension of the active ingredient in an aqueous or oily liquid excipient. Such drops may further comprise 20 buffering agents, salts, and/or one or more other of any additional ingredients described herein. Other ophthalmically-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form and/or in a liposomal preparation. Ear drops and/or eye drops are contemplated as being within the scope of this present disclosure.

25 *Methods of producing polypeptides in cells*

[00471] The present disclosure provides methods of producing a polypeptide of interest in a mammalian cell. Methods of producing polypeptides involve contacting a cell with a nanoparticle composition including an mRNA encoding the polypeptide of interest. Upon 30 contacting the cell with the nanoparticle composition, the mRNA may be taken up and translated in the cell to produce the polypeptide of interest.

[00472] In general, the step of contacting a mammalian cell with a nanoparticle composition including an mRNA encoding a polypeptide of interest may be performed *in vivo*, *ex vivo*, in culture, or *in vitro*. The amount of nanoparticle composition contacted with a cell,

and/or the amount of mRNA therein, may depend on the type of cell or tissue being contacted, the means of administration, the physiochemical characteristics of the nanoparticle composition and the mRNA (e.g., size, charge, and chemical composition) therein, and other factors. In 5 general, an effective amount of the nanoparticle composition will allow for efficient polypeptide production in the cell. Metrics for efficiency may include polypeptide translation (indicated by polypeptide expression), level of mRNA degradation, and immune response indicators.

[00473] The step of contacting a nanoparticle composition including an mRNA with a cell may involve or cause transfection. A phospholipid including in the lipid component of a nanoparticle composition may facilitate transfection and/or increase transfection efficiency, for 10 example, by interacting and/or fusing with a cellular or intracellular membrane. Transfection may allow for the translation of the mRNA within the cell.

[00474] In some embodiments, the nanoparticle compositions described herein may be used therapeutically. For example, an mRNA included in a nanoparticle composition may encode a therapeutic polypeptide (e.g., in a translatable region) and produce the therapeutic 15 polypeptide upon contacting and/or entry (e.g., transfection) into a cell. In other embodiments, an mRNA included in a nanoparticle composition may encode a polypeptide that may improve or increase the immunity of a subject. For example, an mRNA may encode a granulocyte-colony stimulating factor or trastuzumab.

[00475] In certain embodiments, an mRNA included in a nanoparticle composition may 20 encode a recombinant polypeptide that may replace one or more polypeptides that may be substantially absent in a cell contacted with the nanoparticle composition. The one or more substantially absent polypeptides may be lacking due to a genetic mutation of the encoding gene or a regulatory pathway thereof. Alternatively, a recombinant polypeptide produced by 25 translation of the mRNA may antagonize the activity of an endogenous protein present in, on the surface of, or secreted from the cell. An antagonistic recombinant polypeptide may be desirable to combat deleterious effects caused by activities of the endogenous protein, such as altered activities or localization caused by mutation. In another alternative, a recombinant polypeptide produced by translation of the mRNA may indirectly or directly antagonize the activity of a 30 biological moiety present in, on the surface of, or secreted from the cell. Antagonized biological moieties may include, but are not limited to, lipids (e.g., cholesterol), lipoproteins (e.g., low density lipoprotein), nucleic acids, carbohydrates, and small molecule toxins. Recombinant polypeptides produced by translation of the mRNA may be engineered for localization within

the cell, such as within a specific compartment such as the nucleus, or may be engineered for secretion from the cell or for translocation to the plasma membrane of the cell.

[00476] In some embodiments, contacting a cell with a nanoparticle composition including an mRNA may reduce the innate immune response of a cell to an exogenous nucleic acid. A cell may be contacted with a first nanoparticle composition including a first amount of a first exogenous mRNA including a translatable region and the level of the innate immune response of the cell to the first exogenous mRNA may be determined. Subsequently, the cell may be contacted with a second composition including a second amount of the first exogenous mRNA, the second amount being a lesser amount of the first exogenous mRNA compared to the first amount. Alternatively, the second composition may include a first amount of a second exogenous mRNA that is different from the first exogenous mRNA. The steps of contacting the cell with the first and second compositions may be repeated one or more times. Additionally, efficiency of polypeptide production (e.g., translation) in the cell may be optionally determined, and the cell may be re-contacted with the first and/or second composition repeatedly until a target protein production efficiency is achieved.

[00477] In some embodiments, a method of producing a polypeptide of interest in a mammalian cell involves contacting the cell with a nanoparticle composition including (i) a lipid component including a phospholipid, a structural lipid, a PEG lipid, and a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), as described herein; and (ii) an mRNA encoding the polypeptide of interest, whereby the mRNA is capable of being translated in the cell to produce the polypeptide of interest.

Methods of delivering therapeutic agents to cells and organs

[00478] The present disclosure provides methods of delivering a therapeutic and/or prophylactic agent to a mammalian cell or organ. Delivery of a therapeutic and/or prophylactic agent to a cell involves administering a nanoparticle composition including the therapeutic and/or prophylactic agent to a subject, where administration of the composition involves contacting the cell with the composition. For example, a protein, cytotoxic agent, radioactive ion, chemotherapeutic agent, or nucleic acid (such as an RNA, e.g., mRNA) may be delivered to a cell or organ. In the instance that a therapeutic and/or prophylactic agent is an mRNA, upon contacting a cell with the nanoparticle composition, a translatable mRNA may be translated in the cell to produce a polypeptide of interest. However, mRNAs that are substantially not translatable may also be delivered to cells. Substantially non-translatable mRNAs may be useful

as vaccines and/or may sequester translational components of a cell to reduce expression of other species in the cell.

[00479] In some embodiments, a nanoparticle composition may target a particular type or class of cells (e.g., cells of a particular organ or system thereof). For example, a nanoparticle composition including a therapeutic and/or prophylactic agent of interest may be specifically delivered to a mammalian liver, kidney, spleen, femur, or lung. Specific delivery to a particular class of cells, an organ, or a system or group thereof implies that a higher proportion of nanoparticle compositions including a therapeutic and/or prophylactic agent are delivered to the destination (e.g., tissue) of interest relative to other destinations, e.g., upon administration of a nanoparticle composition to a mammal. In some embodiments, specific delivery may result in a greater than 2 fold, 5 fold, 10 fold, 15 fold, or 20 fold increase in the amount of therapeutic and/or prophylactic agent per 1 g of tissue of the targeted destination (e.g., tissue of interest, such as a liver) as compared to another destination (e.g., the spleen). In certain embodiments, the tissue of interest is selected from the group consisting of a liver, kidney, a lung, a spleen, a femur, vascular endothelium in vessels (e.g., intra-coronary or intra-femoral) or kidney, and tumor tissue (e.g., via intratumoral injection).

[00480] As another example of targeted or specific delivery, an mRNA that encodes a protein-binding partner (e.g., an antibody or functional fragment thereof, a scaffold protein, or a peptide) or a receptor on a cell surface may be included in a nanoparticle composition. An mRNA may additionally or instead be used to direct the synthesis and extracellular localization of lipids, carbohydrates, or other biological moieties. Alternatively, other therapeutic and/or prophylactic agents or elements (e.g., lipids or ligands) of a nanoparticle composition may be selected based on their affinity for particular receptors (e.g., low density lipoprotein receptors) such that a nanoparticle composition may more readily interact with a target cell population including the receptors. For example, ligands may include, but are not limited to, members of a specific binding pair, antibodies, monoclonal antibodies, Fv fragments, single chain Fv (scFv) fragments, Fab' fragments, F(ab')2 fragments, single domain antibodies, camelized antibodies and fragments thereof, humanized antibodies and fragments thereof, and multivalent versions thereof; multivalent binding reagents including mono- or bi-specific antibodies such as disulfide stabilized Fv fragments, scFv tandems, diabodies, tridobodies, or tetrabodies; and aptamers, receptors, and fusion proteins.

[00481] In some embodiments, a ligand may be a surface-bound antibody, which can permit tuning of cell targeting specificity. This is especially useful since highly specific

antibodies can be raised against an epitope of interest for the desired targeting site. In one embodiment, multiple antibodies are expressed on the surface of a cell, and each antibody can have a different specificity for a desired target. Such approaches can increase the avidity and specificity of targeting interactions.

5 [00482] A ligand can be selected, e.g., by a person skilled in the biological arts, based on the desired localization or function of the cell. For example an estrogen receptor ligand, such as tamoxifen, can target cells to estrogen-dependent breast cancer cells that have an increased number of estrogen receptors on the cell surface. Other non-limiting examples of ligand/receptor interactions include CCR1 (e.g., for treatment of inflamed joint tissues or brain 10 in rheumatoid arthritis, and/or multiple sclerosis), CCR7, CCR8 (e.g., targeting to lymph node tissue), CCR6, CCR9, CCR10 (e.g., to target to intestinal tissue), CCR4, CCR10 (e.g., for targeting to skin), CXCR4 (e.g., for general enhanced transmigration), HCELL (e.g., for treatment of inflammation and inflammatory disorders, bone marrow), Alpha4beta7 (e.g., for intestinal mucosa targeting), and VLA-4NCAM-1 (e.g., targeting to endothelium). In general, 15 any receptor involved in targeting (e.g., cancer metastasis) can be harnessed for use in the methods and compositions described herein.

[00483] Targeted cells may include, but are not limited to, hepatocytes, epithelial cells, hematopoietic cells, epithelial cells, endothelial cells, lung cells, bone cells, stem cells, mesenchymal cells, neural cells, cardiac cells, adipocytes, vascular smooth muscle cells, 20 cardiomyocytes, skeletal muscle cells, beta cells, pituitary cells, synovial lining cells, ovarian cells, testicular cells, fibroblasts, B cells, T cells, reticulocytes, leukocytes, granulocytes, and tumor cells.

[00484] In certain embodiments, a nanoparticle composition may target hepatocytes. Apolipoproteins such as apolipoprotein E (apoE) have been shown to associate with neutral or 25 near neutral lipid-containing nanoparticle compositions in the body, and are known to associate with receptors such as low-density lipoprotein receptors (LDLRs) found on the surface of hepatocytes. See, e.g., Akinc, A. *et al.*, *Mol. Ther.* 2010, 18, 1357–1364 and Dong, Y. *et al.*, *PNAS* 2014, 111, 3955-3960, the contents of each of which are incorporated herein by reference in their entireties. Thus, a nanoparticle composition including a lipid component with a neutral 30 or near neutral charge that is administered to a subject may acquire apoE in a subject's body and may subsequently deliver a therapeutic and/or prophylactic agent (e.g., an RNA) to hepatocytes including LDLRs in a targeted manner.

[00485] In certain embodiments, cell uptake of a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) as described herein or a nanoparticle composition comprising the compound may be dependent on levels and/or activities of LDLRs, or cell uptake of the nanoparticle composition is LDLR-dependent. For example, if the cell is LDLR-deficient

5 (e.g., having an aberrant LDLR activity and/or an abnormally low level of LDLRs), the cell uptake of the compound or nanoparticle composition may decrease as compared to the uptake by a normal cell.

[00486] In certain embodiments, cell uptake of a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) as described herein or a nanoparticle composition

10 comprising the compound may be independent on levels and/or activities of LDLRs, or cell uptake of the nanoparticle composition is LDLR-independent. For example, if the cell is LDLR-deficient (e.g., having an aberrant LDLR activity and/or an abnormally low level of LDLRs), the cell uptake of the compound or nanoparticle composition is substantively the same as the uptake by a normal cell.

15 **[00487]** In certain embodiments, cell uptake of a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) as described herein or a nanoparticle composition comprising the compound may be dependent on levels and/or activities of apoE, or cell uptake of the nanoparticle composition is apoE-dependent. For example, if the cell is apoE-deficient (e.g., having an aberrant apoE activity and/or an abnormally low level of apoE), the cell uptake

20 of the compound or nanoparticle composition may decrease as compared to the uptake by a normal cell.

25 **[00488]** In certain embodiments, cell uptake of a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), and (IV) as described herein or a nanoparticle composition comprising the compound may be independent on levels and/or activities of apoE, or cell uptake of the nanoparticle composition is apoE-independent. For example, if the cell is apoE-deficient (e.g., having an aberrant apoE activity and/or an abnormally low level of apoE), the cell uptake

30 of the compound or nanoparticle composition is substantively the same as the uptake by a normal cell.

[00489] In certain embodiments, cell uptake of the compound or nanoparticle composition disclosed herein may be both LDLR-dependent and apoE-dependent.

[00490] In certain embodiments, cell uptake of the compound or nanoparticle composition disclosed herein may be dependent on the interaction of LDLR and apoE. For example, if the interaction of LDLR and apoE is abnormal (e.g., leading to an abnormally low level of

downstream signaling), the cell uptake of the compound or nanoparticle composition may decrease as compared to the uptake by a normal cell.

[00491] In certain embodiments, cell uptake of the compound or nanoparticle composition disclosed herein may be both LDLR-independent and apoE-independent.

5 **[00492]** In certain embodiments, cell uptake of the compound or nanoparticle composition disclosed herein may be independent on the interaction of LDLR and apoE. For example, if the interaction of LDLR and apoE is abnormal (e.g., leading to an abnormally low level of downstream signaling), the cell uptake of the compound or nanoparticle composition is substantively the same as the uptake by a normal cell.

10 **[00493]** In certain embodiments, the apoE is apoE3.

[00494] In some embodiments, a method of delivering a therapeutic and/or prophylactic agent to a mammalian cell involves administering to a subject a nanoparticle composition including (i) a lipid component including a phospholipid, a structural lipid, a PEG lipid, and a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), as described herein; and (ii) a therapeutic and/or prophylactic agent (e.g., an mRNA), where administering involves contacting the cell with the nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the cell.

15 **[00495]** In further embodiments, a method of specifically delivering a therapeutic and/or prophylactic agent to a mammalian organ involves administering to a mammal a nanoparticle composition including (i) a lipid component including a phospholipid, a structural lipid, a PEG lipid, and a compound of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), as described herein; and (ii) a therapeutic and/or prophylactic agent (e.g., an mRNA), where administering involves contacting the mammalian organ with the nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the organ.

20 **[00496]** In certain embodiments, the delivery efficiency of the therapeutic and/or prophylactic agent is LDLR-independent or apoE-independent, or both. In certain embodiments, the delivery efficiency of the therapeutic and/or prophylactic agent is LDLR-dependent or apoE-dependent, or both. In certain embodiments, the delivery efficiency of the therapeutic and/or prophylactic agent is independent of LDLR-apoE interaction. In certain embodiments, the delivery efficiency of the therapeutic and/or prophylactic agent is dependent 25 on LDLR-apoE interaction.

Methods of treating diseases and disorders

[00497] Nanoparticle compositions may be useful for treating a disease, disorder, or condition. In particular, such compositions may be useful in treating a disease, disorder, or condition characterized by missing or aberrant protein or polypeptide activity. For example, a nanoparticle composition comprising an mRNA encoding a missing or aberrant polypeptide may be administered or delivered to a cell. Subsequent translation of the mRNA may produce the polypeptide, thereby reducing or eliminating an issue caused by the absence of or aberrant activity caused by the polypeptide. Because translation may occur rapidly, the methods and compositions may be useful in the treatment of acute diseases, disorders, or conditions such as sepsis, stroke, and myocardial infarction. A therapeutic and/or prophylactic agent included in a nanoparticle composition may also be capable of altering the rate of transcription of a given species, thereby affecting gene expression.

[00498] Diseases, disorders, and/or conditions characterized by dysfunctional or aberrant protein or polypeptide activity for which a composition may be administered include, but are not limited to, rare diseases, infectious diseases (as both vaccines and therapeutics), cancer and proliferative diseases, genetic diseases (e.g., cystic fibrosis), autoimmune diseases, diabetes, neurodegenerative diseases, cardio- and reno-vascular diseases, and metabolic diseases.

Multiple diseases, disorders, and/or conditions may be characterized by missing (or substantially diminished such that proper protein function does not occur) protein activity. Such proteins may not be present, or they may be essentially non-functional. A specific example of a dysfunctional protein is the missense mutation variants of the cystic fibrosis transmembrane conductance regulator (CFTR) gene, which produce a dysfunctional protein variant of CFTR protein, which causes cystic fibrosis. The present disclosure provides a method for treating such diseases, disorders, and/or conditions in a subject by administering a nanoparticle composition including an RNA and a lipid component including a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), a phospholipid (optionally unsaturated), a PEG lipid, and a structural lipid, wherein the RNA may be an mRNA encoding a polypeptide that antagonizes or otherwise overcomes an aberrant protein activity present in the cell of the subject.

[00499] In some embodiments, a method of treating a disease or disorder in a mammal in need involves administering to the mammal a therapeutically effective amount of a nanoparticle composition including (i) a lipid component including a phospholipid, a structural lipid, a PEG lipid, and a compound of formula (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I),

(19-II), or (20-I), as described herein; and (ii) a therapeutic and/or prophylactic agent (e.g., an mRNA).

[00500] The disclosure provides methods involving administering nanoparticle compositions including one or more therapeutic and/or prophylactic agents and pharmaceutical compositions including the same. The terms therapeutic and prophylactic can be used interchangeably herein with respect to features and embodiments of the present disclosure. Therapeutic compositions, or imaging, diagnostic, or prophylactic compositions thereof, may be administered to a subject using any reasonable amount and any route of administration effective for preventing, treating, diagnosing, or imaging a disease, disorder, and/or condition and/or any other purpose. The specific amount administered to a given subject may vary depending on the species, age, and general condition of the subject; the purpose of the administration; the particular composition; the mode of administration; and the like. Compositions in accordance with the present disclosure may be formulated in dosage unit form for ease of administration and uniformity of dosage. It will be understood, however, that the total daily usage of a composition of the present disclosure will be decided by an attending physician within the scope of sound medical judgment. The specific therapeutically effective, prophylactically effective, or otherwise appropriate dose level (e.g., for imaging) for any particular patient will depend upon a variety of factors including the severity and identify of a disorder being treated, if any; the one or more therapeutic and/or prophylactic agents employed; the specific composition employed; the age, body weight, general health, sex, and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific pharmaceutical composition employed; the duration of the treatment; drugs used in combination or coincidental with the specific pharmaceutical composition employed; and like factors well known in the medical arts.

[00501] A nanoparticle composition including one or more therapeutic and/or prophylactic agents may be administered by any route. In some embodiments, compositions, including prophylactic, diagnostic, or imaging compositions including one or more nanoparticle compositions described herein, are administered by one or more of a variety of routes, including oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, subcutaneous, intraventricular, trans- or intra-dermal, interdermal, rectal, intravaginal, intraperitoneal, intraocular, subretinal, intravitreal, topical (e.g. by powders, ointments, creams, gels, lotions, and/or drops), mucosal, nasal, buccal, enteral, vitreal, intratumoral, sublingual, intranasal; by intratracheal instillation, bronchial instillation, and/or inhalation; as an oral spray and/or powder, nasal spray, and/or aerosol, and/or through a portal vein catheter. In some embodiments, a

composition may be administered intravenously, intramuscularly, intradermally, intra-arterially, intratumorally, subcutaneously, intraocular, subretinal, intravitreal, or by inhalation. However, the present disclosure encompasses the delivery or administration of compositions by any appropriate route taking into consideration likely advances in the sciences of drug delivery. In 5 general, the most appropriate route of administration will depend upon a variety of factors including the nature of the nanoparticle composition including one or more therapeutic and/or prophylactic agents (e.g., its stability in various bodily environments such as the bloodstream and gastrointestinal tract), the condition of the patient (e.g., whether the patient is able to tolerate particular routes of administration), etc.

10 [00502] In certain embodiments, compositions in accordance with the present disclosure may be administered at dosage levels sufficient to deliver from about 0.0001 mg/kg to about 10 mg/kg, from about 0.001 mg/kg to about 10 mg/kg, from about 0.005 mg/kg to about 10 mg/kg, from about 0.01 mg/kg to about 10 mg/kg, from about 0.05 mg/kg to about 10 mg/kg, from about 0.1 mg/kg to about 10 mg/kg, from about 1 mg/kg to about 10 mg/kg, from about 2 mg/kg to about 10 mg/kg, from about 5 mg/kg to about 10 mg/kg, from about 0.0001 mg/kg to about 5 mg/kg, from about 0.001 mg/kg to about 5 mg/kg, from about 0.005 mg/kg to about 5 mg/kg, from about 0.01 mg/kg to about 5 mg/kg, from about 0.05 mg/kg to about 5 mg/kg, from about 0.1 mg/kg to about 5 mg/kg, from about 1 mg/kg to about 5 mg/kg, from about 2 mg/kg to about 5 mg/kg, from about 0.0001 mg/kg to about 2.5 mg/kg, from about 0.001 mg/kg to about 2.5 mg/kg, from about 0.005 mg/kg to about 2.5 mg/kg, from about 0.01 mg/kg to about 2.5 mg/kg, from about 0.05 mg/kg to about 2.5 mg/kg, from about 0.1 mg/kg to about 2.5 mg/kg, from about 0.1 mg/kg to about 2.5 mg/kg, from about 0.0001 mg/kg to about 0.25 mg/kg, from about 0.001 mg/kg to about 0.25 mg/kg, from about 0.005 mg/kg to about 0.25 mg/kg, from about 0.01 mg/kg to about 0.25 mg/kg, from about 0.05 mg/kg to about 0.25 mg/kg, or from about 0.1 mg/kg to about 0.25 mg/kg of a therapeutic and/or prophylactic agent (e.g., an mRNA) in a given dose, where a dose of 1 mg/kg (mpk) provides 1 mg of a therapeutic and/or prophylactic agent per 1 kg of subject body weight. In certain embodiments, a dose of about 0.001 mg/kg to about 10 mg/kg of a therapeutic and/or prophylactic agent (e.g., mRNA) of a nanoparticle composition may be administered. In other embodiments, a dose of about 0.005 mg/kg to about 2.5 mg/kg of a therapeutic and/or prophylactic agent may be administered. In 15 20 25 30

certain embodiments, a dose of about 0.1 mg/kg to about 1 mg/kg may be administered. In other embodiments, a dose of about 0.05 mg/kg to about 0.25 mg/kg may be administered. A dose may be administered one or more times per day, in the same or a different amount, to obtain a desired level of mRNA expression and/or therapeutic, diagnostic, prophylactic, or imaging effect. The desired dosage may be delivered, for example, three times a day, two times a day, once a day, every other day, every third day, every week, every two weeks, every three weeks, or every four weeks. In certain embodiments, the desired dosage may be delivered using multiple administrations (e.g., two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, or more administrations). In some embodiments, a single dose may be administered, for example, prior to or after a surgical procedure or in the instance of an acute disease, disorder, or condition.

[00503] Nanoparticle compositions including one or more therapeutic and/or prophylactic agents may be used in combination with one or more other therapeutic, prophylactic, diagnostic, or imaging agents. By “in combination with,” it is not intended to imply that the agents must be administered at the same time and/or formulated for delivery together, although these methods of delivery are within the scope of the present disclosure. For example, one or more nanoparticle compositions including one or more different therapeutic and/or prophylactic agents may be administered in combination. Compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. In general, each agent will be administered at a dose and/or on a time schedule determined for that agent. In some embodiments, the present disclosure encompasses the delivery of compositions, or imaging, diagnostic, or prophylactic compositions thereof in combination with agents that improve their bioavailability, reduce and/or modify their metabolism, inhibit their excretion, and/or modify their distribution within the body.

[00504] It will further be appreciated that therapeutically, prophylactically, diagnostically, or imaging active agents utilized in combination may be administered together in a single composition or administered separately in different compositions. In general, it is expected that agents utilized in combination will be utilized at levels that do not exceed the levels at which they are utilized individually. In some embodiments, the levels utilized in combination may be lower than those utilized individually.

[00505] The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the

therapies employed may achieve a desired effect for the same disorder (for example, a composition useful for treating cancer may be administered concurrently with a chemotherapeutic agent), or they may achieve different effects (e.g., control of any adverse effects, such as infusion related reactions).

5 [00506] A nanoparticle composition may be used in combination with an agent to increase the effectiveness and/or therapeutic window of the composition. Such an agent may be, for example, an anti-inflammatory compound, a steroid (e.g., a corticosteroid), a statin, an estradiol, a BTK inhibitor, an S1P1 agonist, a glucocorticoid receptor modulator (GRM), or an anti-histamine. In some embodiments, a nanoparticle composition may be used in combination 10 with dexamethasone, methotrexate, acetaminophen, an H1 receptor blocker, or an H2 receptor blocker. In certain embodiments, a method of treating a subject in need thereof or of delivering a therapeutic and/or prophylactic agent to a subject (e.g., a mammal) may involve pre-treating the subject with one or more agents prior to administering a nanoparticle composition. For example, a subject may be pre-treated with a useful amount (e.g., 10 mg, 20 mg, 30 mg, 40 mg, 15 50 mg, 60 mg, 70 mg, 80 mg, 90 mg, 100 mg, or any other useful amount) of dexamethasone, methotrexate, acetaminophen, an H1 receptor blocker, or an H2 receptor blocker. Pre-treatment may occur 24 or fewer hours (e.g., 24 hours, 20 hours, 16 hours, 12 hours, 8 hours, 4 hours, 2 hours, 1 hour, 50 minutes, 40 minutes, 30 minutes, 20 minutes, or 10 minutes) before 20 administration of the nanoparticle composition and may occur one, two, or more times in, for example, increasing dosage amounts.

[00507] In any method or use described herein, in certain embodiments, the subject in need thereof is LDLR-deficient or apoE-deficient or both. In certain embodiments, the subject in need thereof is not LDLR-deficient or has normal LDLR levels and/or activities. In certain embodiments, the subject in need thereof is not apoE-deficient or has normal apoE levels and/or 25 activities. In certain embodiments, the subject in need thereof has an abnormal interaction of LDLR and apoE. In certain embodiments, the subject in need thereof has a normal interaction of LDLR and apoE.

[00508] About, Approximately: As used herein, the terms “approximately” and “about,” 30 as applied to one or more values of interest, refer to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated

reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value). For example, when used in the context of an amount of a given compound in a lipid component of a nanoparticle composition, “about” may mean +/- 10% of the recited value. For instance, a nanoparticle composition including a

5 lipid component having about 40% of a given compound may include 30-50% of the compound.

10 [00509] Compound: As used herein, the term “compound,” is meant to include all isomers and isotopes of the structure depicted. “Isotopes” refers to atoms having the same atomic number but different mass numbers resulting from a different number of neutrons in the nuclei. For example, isotopes of hydrogen include tritium and deuterium. Further, a compound, salt, or complex of the present disclosure can be prepared in combination with solvent or water molecules to form solvates and hydrates by routine methods.

15 [00510] Contacting: As used herein, the term “contacting” means establishing a physical connection between two or more entities. For example, contacting a mammalian cell with a nanoparticle composition means that the mammalian cell and a nanoparticle are made to share a physical connection. Methods of contacting cells with external entities both in vivo and ex vivo are well known in the biological arts. For example, contacting a nanoparticle composition and a mammalian cell disposed within a mammal may be performed by varied routes of administration (e.g., intravenous, intramuscular, intradermal, and subcutaneous) and may involve varied amounts of nanoparticle compositions. Moreover, more than one mammalian cell may be

20 contacted by a nanoparticle composition.

25 [00511] Delivering: As used herein, the term “delivering” means providing an entity to a destination. For example, delivering a therapeutic and/or prophylactic agent to a subject may involve administering a nanoparticle composition including the therapeutic and/or prophylactic agent to the subject (e.g., by an intravenous, intramuscular, intradermal, or subcutaneous route). Administration of a nanoparticle composition to a mammal or mammalian cell may involve

contacting one or more cells with the nanoparticle composition.

30 [00512] Enhanced delivery: As used herein, the term “enhanced delivery” means delivery of more (e.g., at least 1.5 fold more, at least 2-fold more, at least 3-fold more, at least 4-fold more, at least 5-fold more, at least 6-fold more, at least 7-fold more, at least 8-fold more, at least 9-fold more, at least 10-fold more) of a therapeutic and/or prophylactic agent by a nanoparticle to a target tissue of interest (e.g., mammalian liver) compared to the level of delivery of a therapeutic and/or prophylactic agent by a control nanoparticle to a target tissue of interest (e.g., MC3, KC2, or DLinDMA). The level of delivery of a nanoparticle to a particular tissue may be

measured by comparing the amount of protein produced in a tissue to the weight of said tissue, comparing the amount of therapeutic and/or prophylactic agent in a tissue to the weight of said tissue, comparing the amount of protein produced in a tissue to the amount of total protein in said tissue, or comparing the amount of therapeutic and/or prophylactic agent in a tissue to the amount of total therapeutic and/or prophylactic agent in said tissue. It will be understood that the enhanced delivery of a nanoparticle to a target tissue need not be determined in a subject being treated, it may be determined in a surrogate such as an animal model (e.g., a rat model).

5 [00513] Specific delivery: As used herein, the term “specific delivery,” “specifically deliver,” or “specifically delivering” means delivery of more (e.g., at least 1.5 fold more, at least 10 2-fold more, at least 3-fold more, at least 4-fold more, at least 5-fold more, at least 6-fold more, at least 7-fold more, at least 8-fold more, at least 9-fold more, at least 10-fold more) of a therapeutic and/or prophylactic agent by a nanoparticle to a target tissue of interest (e.g., mammalian liver) compared to an off-target tissue (e.g., mammalian spleen). The level of delivery of a nanoparticle to a particular tissue may be measured by comparing the amount of 15 protein produced in a tissue to the weight of said tissue, comparing the amount of therapeutic and/or prophylactic agent in a tissue to the weight of said tissue, comparing the amount of protein produced in a tissue to the amount of total protein in said tissue, or comparing the amount of therapeutic and/or prophylactic agent in a tissue to the amount of total therapeutic and/or prophylactic agent in said tissue. For example, for renovascular targeting, a therapeutic 20 and/or prophylactic agent is specifically provided to a mammalian kidney as compared to the liver and spleen if 1.5, 2-fold, 3-fold, 5-fold, 10-fold, 15 fold, or 20 fold more therapeutic and/or prophylactic agent per 1 g of tissue is delivered to a kidney compared to that delivered to the liver or spleen following systemic administration of the therapeutic and/or prophylactic agent. It will be understood that the ability of a nanoparticle to specifically deliver to a target tissue need 25 not be determined in a subject being treated, it may be determined in a surrogate such as an animal model (e.g., a rat model).

30 [00514] Encapsulation efficiency: As used herein, “encapsulation efficiency” refers to the amount of a therapeutic and/or prophylactic agent that becomes part of a nanoparticle composition, relative to the initial total amount of therapeutic and/or prophylactic agent used in the preparation of a nanoparticle composition. For example, if 97 mg of therapeutic and/or prophylactic agent are encapsulated in a nanoparticle composition out of a total 100 mg of therapeutic and/or prophylactic agent initially provided to the composition, the encapsulation

efficiency may be given as 97%. As used herein, “encapsulation” may refer to complete, substantial, or partial enclosure, confinement, surrounding, or encasement.

5 [00515] Expression: As used herein, “expression” of a nucleic acid sequence refers to translation of an mRNA into a polypeptide or protein and/or post-translational modification of a polypeptide or protein.

[00516] In vitro: As used herein, the term “in vitro” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, in a Petri dish, etc., rather than within an organism (e.g., animal, plant, or microbe).

10 [00517] In vivo: As used herein, the term “in vivo” refers to events that occur within an organism (e.g., animal, plant, or microbe or cell or tissue thereof).

[00518] Ex vivo: As used herein, the term “ex vivo” refers to events that occur outside of an organism (e.g., animal, plant, or microbe or cell or tissue thereof). Ex vivo events may take place in an environment minimally altered from a natural (e.g., in vivo) environment.

15 [00519] Isomer: As used herein, the term “isomer” means any geometric isomer, tautomer, zwitterion, stereoisomer, enantiomer, or diastereomer of a compound. Compounds may include one or more chiral centers and/or double bonds and may thus exist as stereoisomers, such as double-bond isomers (i.e., geometric E/Z isomers) or diastereomers (e.g., enantiomers (i.e., (+) or (-)) or *cis/trans* isomers). The present disclosure encompasses any and all isomers of the compounds described herein, including stereomerically pure forms (e.g., 20 geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric and stereoisomeric mixtures, e.g., racemates. Enantiomeric and stereomeric mixtures of compounds and means of resolving them into their component enantiomers or stereoisomers are well-known.

25 [00520] Lipid component: As used herein, a “lipid component” is that component of a nanoparticle composition that includes one or more lipids. For example, the lipid component may include one or more cationic/ionizable, PEGylated, structural, or other lipids, such as phospholipids.

30 [00521] Linker: As used herein, a “linker” is a moiety connecting two moieties, for example, the connection between two nucleosides of a cap species. A linker may include one or more groups including but not limited to phosphate groups (e.g., phosphates, boranophosphates, thiophosphates, selenophosphates, and phosphonates), alkyl groups, amidates, or glycerols. For example, two nucleosides of a cap analog may be linked at their 5' positions by a triphosphate group or by a chain including two phosphate moieties and a boranophosphate moiety.

[00522] Methods of administration: As used herein, “methods of administration” may include intravenous, intramuscular, intradermal, subcutaneous, or other methods of delivering a composition to a subject. A method of administration may be selected to target delivery (e.g., to specifically deliver) to a specific region or system of a body.

5 **[00523]** Modified: As used herein, “modified” means non-natural. For example, an RNA may be a modified RNA. That is, an RNA may include one or more nucleobases, nucleosides, nucleotides, or linkers that are non-naturally occurring. A “modified” species may also be referred to herein as an “altered” species. Species may be modified or altered chemically, structurally, or functionally. For example, a modified nucleobase species may include one or
10 more substitutions that are not naturally occurring.

[00524] N:P ratio: As used herein, the “N:P ratio” is the molar ratio of ionizable (in the physiological pH range) nitrogen atoms in a lipid to phosphate groups in an RNA, e.g., in a nanoparticle composition including a lipid component and an RNA.

15 **[00525]** Nanoparticle composition: As used herein, a “nanoparticle composition” is a composition comprising one or more lipids. Nanoparticle compositions are typically sized on the order of micrometers or smaller and may include a lipid bilayer. Nanoparticle compositions encompass lipid nanoparticles (LNPs), liposomes (e.g., lipid vesicles), and lipoplexes. For example, a nanoparticle composition may be a liposome having a lipid bilayer with a diameter of 500 nm or less.

20 **[00526]** Naturally occurring: As used herein, “naturally occurring” means existing in nature without artificial aid.

[00527] Patient: As used herein, “patient” refers to a subject who may seek or be in need of treatment, requires treatment, is receiving treatment, will receive treatment, or a subject who is under care by a trained professional for a particular disease or condition.

25 **[00528]** PEG lipid: As used herein, a “PEG lipid” or “PEGylated lipid” refers to a lipid comprising a polyethylene glycol component.

30 **[00529]** Pharmaceutically acceptable: The phrase “pharmaceutically acceptable” is used herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[00530] Pharmaceutically acceptable excipient: The phrase “pharmaceutically acceptable excipient,” as used herein, refers to any ingredient other than the compounds described herein

(for example, a vehicle capable of suspending, complexing, or dissolving the active compound) and having the properties of being substantially nontoxic and non-inflammatory in a patient. Excipients may include, for example: anti-adherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluents), film 5 formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, and waters of hydration. Exemplary excipients include, but are not limited to: butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose, gelatin, hydroxypropyl cellulose, 10 hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E (alpha-tocopherol), vitamin C, xylitol, and other species disclosed herein.

15 **[00531]** In the present specification, the structural formula of the compound represents a certain isomer for convenience in some cases, but the present disclosure includes all isomers, such as geometrical isomers, optical isomers based on an asymmetrical carbon, stereoisomers, tautomers, and the like, it being understood that not all isomers may have the same level of 20 activity. In addition, a crystal polymorphism may be present for the compounds represented by the formula. It is noted that any crystal form, crystal form mixture, or anhydride or hydrate thereof is included in the scope of the present disclosure.

25 **[00532]** The term “crystal polymorphs”, “polymorphs” or “crystal forms” means crystal structures in which a compound (or a salt or solvate thereof) can crystallize in different crystal packing arrangements, all of which have the same elemental composition. Different crystal forms usually have different X-ray diffraction patterns, infrared spectral, melting points, density hardness, crystal shape, optical and electrical properties, stability and solubility. Recrystallization solvent, rate of crystallization, storage temperature, and other factors may 30 cause one crystal form to dominate. Crystal polymorphs of the compounds can be prepared by crystallization under different conditions.

[00533] Pharmaceutically acceptable salts: Compositions may also include salts of one or more compounds. Salts may be pharmaceutically acceptable salts. As used herein, “pharmaceutically acceptable salts” refers to derivatives of the disclosed compounds wherein the

parent compound is altered by converting an existing acid or base moiety to its salt form (e.g., by reacting a free base group with a suitable organic acid). Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

5 Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, 10 maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, valerate salts, and the like. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like, as well as nontoxic ammonium, 15 quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, ethylamine, and the like. The pharmaceutically acceptable salts of the present disclosure include the conventional non-toxic salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. The pharmaceutically acceptable salts of the present disclosure can be synthesized from the parent compound which contains a basic or acidic moiety 20 by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts 25 are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418, Pharmaceutical Salts: Properties, Selection, and Use, P.H. Stahl and C.G. Wermuth (eds.), Wiley-VCH, 2008, and Berge et al., Journal of Pharmaceutical Science, 66, 1-19 (1977), each of which is incorporated herein by reference in its entirety.

[00534] Phospholipid: As used herein, a "phospholipid" is a lipid that includes a 30 phosphate moiety and one or more carbon chains, such as unsaturated fatty acid chains. A phospholipid may include one or more multiple (e.g., double or triple) bonds (e.g., one or more unsaturations). Particular phospholipids may facilitate fusion to a membrane. For example, a cationic phospholipid may interact with one or more negatively charged phospholipids of a

membrane (e.g., a cellular or intracellular membrane). Fusion of a phospholipid to a membrane may allow one or more elements of a lipid-containing composition to pass through the membrane permitting, e.g., delivery of the one or more elements to a cell.

5 [00535] Polydispersity index: As used herein, the “polydispersity index” is a ratio that describes the homogeneity of the particle size distribution of a system. A small value, e.g., less than 0.3, indicates a narrow particle size distribution.

[00536] Polypeptide: As used herein, the term “polypeptide” or “polypeptide of interest” refers to a polymer of amino acid residues typically joined by peptide bonds that can be produced naturally (e.g., isolated or purified) or synthetically.

10 [00537] RNA: As used herein, an “RNA” refers to a ribonucleic acid that may be naturally or non-naturally occurring. For example, an RNA may include modified and/or non-naturally occurring components such as one or more nucleobases, nucleosides, nucleotides, or linkers. An RNA may include a cap structure, a chain terminating nucleoside, a stem loop, a polyA sequence, and/or a polyadenylation signal. An RNA may have a nucleotide sequence 15 encoding a polypeptide of interest. For example, an RNA may be a messenger RNA (mRNA). Translation of an mRNA encoding a particular polypeptide, for example, *in vivo* translation of an mRNA inside a mammalian cell, may produce the encoded polypeptide. RNAs may be selected from the non-liming group consisting of small interfering RNA (siRNA), asymmetrical interfering RNA (aiRNA), microRNA (miRNA), Dicer-substrate RNA (dsRNA), small hairpin 20 RNA (shRNA), mRNA, and mixtures thereof.

[00538] Single unit dose: As used herein, a “single unit dose” is a dose of any therapeutic administered in one dose/at one time/single route/single point of contact, i.e., single administration event.

25 [00539] Split dose: As used herein, a “split dose” is the division of single unit dose or total daily dose into two or more doses.

[00540] Total daily dose: As used herein, a “total daily dose” is an amount given or prescribed in 24 hour period. It may be administered as a single unit dose.

[00541] Size: As used herein, “size” or “mean size” in the context of nanoparticle compositions refers to the mean diameter of a nanoparticle composition.

30 [00542] Subject: As used herein, the term “subject” or “patient” refers to any organism to which a composition in accordance with the disclosure may be administered, e.g., for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include

animals (e.g., mammals such as mice, rats, rabbits, non-human primates, and humans) and/or plants.

5 [00543] Targeted cells: As used herein, “targeted cells” refers to any one or more cells of interest. The cells may be found in vitro, in vivo, in situ, or in the tissue or organ of an organism. The organism may be an animal, preferably a mammal, more preferably a human and most preferably a patient.

10 [00544] Target tissue: As used herein “target tissue” refers to any one or more tissue types of interest in which the delivery of a therapeutic and/or prophylactic agent would result in a desired biological and/or pharmacological effect. Examples of target tissues of interest include specific tissues, organs, and systems or groups thereof. In particular applications, a target tissue may be a kidney, a lung, a spleen, vascular endothelium in vessels (e.g., intra-coronary or intra-femoral), or tumor tissue (e.g., via intratumoral injection). An “off-target tissue” refers to any one or more tissue types in which the expression of the encoded protein does not result in a desired biological and/or pharmacological effect. In particular applications, off-target tissues 15 may include the liver and the spleen.

20 [00545] Therapeutic and/or prophylactic agent: The term “therapeutic agent” refers to any agent that, when administered to a subject, has a therapeutic and/or diagnostic effect and/or elicits a desired biological and/or pharmacological effect. The term “prophylactic agent” refers to any agent that, when administered to a subject, has a prophylactic effect. Therapeutic and/or prophylactic agents are also referred to as “actives” or “active agents.” Such agents include, but are not limited to, cytotoxins, radioactive ions, chemotherapeutic agents, small molecule drugs, 25 proteins, and nucleic acids.

25 [00546] Therapeutically effective amount: As used herein, the term “therapeutically effective amount” means an amount of an agent to be delivered (e.g., nucleic acid, drug, composition, therapeutic agent, diagnostic agent, prophylactic agent, etc.) that is sufficient, when administered to a subject suffering from or susceptible to an infection, disease, disorder, and/or condition, to treat, improve symptoms of, diagnose, prevent, and/or delay the onset of the infection, disease, disorder, and/or condition.

30 [00547] Transfection: As used herein, “transfection” refers to the introduction of a species (e.g., an RNA) into a cell. Transfection may occur, for example, in vitro, ex vivo, or in vivo.

[00548] Treating: As used herein, the term “treating” refers to partially or completely alleviating, ameliorating, improving, relieving, delaying onset of, inhibiting progression of, reducing severity of, and/or reducing incidence of one or more symptoms or features of a

particular infection, disease, disorder, and/or condition. For example, “treating” cancer may refer to inhibiting survival, growth, and/or spread of a tumor. Treatment may be administered to a subject who does not exhibit signs of a disease, disorder, and/or condition and/or to a subject who exhibits only early signs of a disease, disorder, and/or condition for the purpose of
5 decreasing the risk of developing pathology associated with the disease, disorder, and/or condition.

[00549] Zeta potential: As used herein, the “zeta potential” is the electrokinetic potential of a lipid e.g., in a particle composition.

[00550] Those skilled in the art will recognize, or be able to ascertain using no more than
10 routine experimentation, many equivalents to the specific embodiments in accordance with the present disclosure. The scope of the present disclosure is not intended to be limited to the above Description, but rather is as set forth in the appended claims.

[00551] In the claims, articles such as “a,” “an,” and “the” may mean one or more than
15 one unless indicated to the contrary or otherwise evident from the context. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The disclosure includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The disclosure
20 includes embodiments in which more than one, or all, of the group members are present in, employed in, or otherwise relevant to a given product or process. As used herein, the expressions “one or more of A, B, or C,” “one or more A, B, or C,” “one or more of A, B, and C,” “one or more A, B, and C”, “selected from A, B, and C,” “selected from the group
25 consisting of A, B, and C,” and the like are used interchangeably and all refer to a selection from a group consisting of A, B, and /or C, i.e., one or more As, one or more Bs, one or more Cs, or any combination thereof, unless otherwise specified.

[00552] It is also noted that the term “comprising” is intended to be open and permits but does not require the inclusion of additional elements or steps. When the term “comprising” is used herein, the terms “consisting essentially of” and “consisting of” are thus also encompassed and disclosed. Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where methods or processes are described as having, including, or comprising specific process steps, the processes also consist
30

essentially of, or consist of, the recited processing steps. Further, it should be understood that the order of steps or order for performing certain actions is immaterial so long as the invention remains operable. Moreover, two or more steps or actions can be conducted simultaneously.

[00553] Where ranges are given, endpoints are included. Furthermore, it is to be

5 understood that unless otherwise indicated or otherwise evident from the context and understanding of one of ordinary skill in the art, values that are expressed as ranges can assume any specific value or sub-range within the stated ranges in different embodiments of the disclosure, to the tenth of the unit of the lower limit of the range, unless the context clearly dictates otherwise.

10 [00554] The synthetic processes of the disclosure can tolerate a wide variety of functional groups, therefore various substituted starting materials can be used. The processes generally provide the desired final compound at or near the end of the overall process, although it may be desirable in certain instances to further convert the compound to a pharmaceutically acceptable salt thereof.

15 [00555] Compounds of the present disclosure can be prepared in a variety of ways using commercially available starting materials, compounds known in the literature, or from readily prepared intermediates, by employing standard synthetic methods and procedures either known to those skilled in the art, or which will be apparent to the skilled artisan in light of the teachings herein. Standard synthetic methods and procedures for the preparation of organic molecules and 20 functional group transformations and manipulations can be obtained from the relevant scientific literature or from standard textbooks in the field. Although not limited to any one or several sources, classic texts such as Smith, M. B., March, J., *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 5th edition, John Wiley & Sons: New York, 2001; Greene, T.W., Wuts, P.G. M., *Protective Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons: New York, 1999; R. Larock, *Comprehensive Organic Transformations*, VCH Publishers (1989); L. Fieser and M. Fieser, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995), incorporated by reference herein, are useful and 25 recognized reference textbooks of organic synthesis known to those in the art. The following descriptions of synthetic methods are designed to illustrate, but not to limit, general procedures for the preparation of compounds of the present disclosure.

[00556] The compounds of this disclosure having any of the formulae described herein may be prepared according to the procedures illustrated in Schemes 1-3 below, from

commercially available starting materials or starting materials which can be prepared using literature procedures. The variables in the Schemes (e.g., R₁, R₂, R₃, R₄, and R₅) are as defined herein, e.g., R₁, R₂, R₃, R₄, and R₅ are each independently alkyl. One of ordinary skill in the art will note that, during the reaction sequences and synthetic schemes described herein, the order 5 of certain steps may be changed, such as the introduction and removal of protecting groups.

[00557] One of ordinary skill in the art will recognize that certain groups may require protection from the reaction conditions via the use of protecting groups. Protecting groups may also be used to differentiate similar functional groups in molecules. A list of protecting groups and how to introduce and remove these groups can be found in Greene, T.W., Wuts, P.G. M., 10 *Protective Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons: New York, 1999.

Preferred protecting groups include, but are not limited to:

For a hydroxyl moiety: TBS, benzyl, THP, Ac;

For carboxylic acids: benzyl ester, methyl ester, ethyl ester, allyl ester;

For amines: Fmoc, Cbz, BOC, DMB, Ac, Bn, Tr, Ts, trifluoroacetyl, phthalimide, 15 benzylideneamine;

For diols: Ac (x2) TBS (x2), or when taken together acetonides;


For thiols: Ac;

For benzimidazoles: SEM, benzyl, PMB, DMB;

For aldehydes: di-alkyl acetals such as dimethoxy acetal or diethyl acetyl.

[00558] In the reaction schemes described herein, multiple stereoisomers may be produced. When no particular stereoisomer is indicated, it is understood to mean all possible stereoisomers that could be produced from the reaction. A person of ordinary skill in the art will recognize that the reactions can be optimized to give one isomer preferentially, or new schemes may be devised to produce a single isomer. If mixtures are produced, techniques such as 20 preparative thin layer chromatography, preparative HPLC, preparative chiral HPLC, or preparative SFC may be used to separate the isomers.


Scheme 1



[00559] Scheme 1 above illustrates an 11-step procedure for the synthesis of lipids of the disclosure. (a) Aldehyde (**I-1**) is reacted with glycine methyl ester hydrochloride in the presence of a reducing agent, e.g. $NaBH(OAc)_3$, and a base, e.g. NEt_3 , in an appropriate solvent, e.g. $AcOH$, to yield methyl dialkylglycinate (**I-2**). This reaction can take place in an organic solvent, e.g. dichloroethane. Methyl dialkylglycinate (**I-2**) is hydrolyzed using lithium hydroxide, e.g. in THF , to produce the respective lithium alkyl glycinate (**I-3**). A solution of (**I-3**), e.g. in THF , is reacted with 1-*tert*-butyl-piperazine in the presence of a base, e.g. diisopropylethylamine (DIPEA), and a coupling agent, e.g. propylphosphonic acid anhydride, to form *tert*-butyl 4-(alkyl)piperazine-1-carboxylate (**I-4**), which is then deprotected, using e.g. trifluoroacetic acid (TFA), to yield 2-(alkyl)-1-(piperazine-1-yl)ethan-1-one (**I-5**). This reaction can take place in an

organic solvent, e.g. dichloromethane (DCM). (b) Bromoalkane (**I-6**) is reacted with *tert*-butyl-methyl alkyl glycinate in the presence of a strong base, e.g. NaH, and in an appropriate solvent, e.g. dimethylformamide (DMF), to form methyl N-(*tert*-butyl)-N-alkylglycinate (**I-7**), which is then deprotected, using e.g. TFA, to yield methyl alkylglycinate (**I-8**). The deprotection reaction 5 can take place in an organic solvent, e.g. dichloromethane. (c) Bromoalkane (**I-6**) is reacted with ethanol-1-amine under alkaline conditions (e.g. K₂CO₃) and in the presence of a catalyst, e.g. KI in an appropriate solvent (e.g. acetonitrile), to form 2-(dialkylamino)ethanol (**I-9**), which is converted to *N*-(2-chloroethyl)-*N*-alkylalkan-1-amine (**I-10**) using a suitable reagent, e.g. mesylchloride in the presence of a base, e.g. triethylamine, and an appropriate solvent, e.g. 10 DCM. (d) methyl alkylglycinate (**I-8**), obtained according to (b) and *N*-(2-chloroethyl)-*N*-alkylalkan-1-amine (**I-10**) obtained according to (c) are coupled in the presence of a base, e.g. K₂CO₃ and a nucleophilic catalyst, e.g. KI in an appropriate solvent (e.g. acetonitrile), to form the methyl glycinate intermediate (**I-11**), which is then hydrolyzed using lithium hydroxide in an appropriate solvent, e.g. tetrahydrofuran (THF), to yield the lithium glycinate compound 15 (**I-12**). (e) 2-(alkyl)-1-(piperazine-1-yl)ethan-1-one (**I-5**) obtained according to (a) and compound (**I-12**) obtained according to (d) are reacted in the presence of a base, e.g. diisopropylethylamine (DIPEA), and a coupling agent, e.g. propylphosphonic acid anhydride, to yield the product (**P-1**). This reaction can take place in an organic solvent, e.g. THF.


20 **Scheme 2**



[00560] Scheme 2 above illustrates a 6-step procedure for the synthesis of lipids of the disclosure. (a) Commercially available *tert*-butyl 4-glycylpiperazine-1-carboxylate hydrochloride (**II-1**) is reacted with bromoalkane in the presence of a base, e.g. K_2CO_3 and a

nucleophilic catalyst, e.g. KI in an appropriate solvent, e.g. cyclopentyl methyl ether/acetonitrile, and deprotected using, e.g. trifluoroacetic (TFA) to yield 2-(dialkylamino)-1-(piperazin-1-yl)ethan-1-one. The deprotection step can take place in an organic solvent, e.g. dichloromethane (DCM) (b) tert-Butyl 2-bromoacetate (**II-4**) is reacted with 1,2-diamino ethane, 5 in an appropriate solvent, e.g. DCM to yield tert-butyl (2-aminoethyl)glycinate (**II-5**) which is coupled with bromoalkane in the presence of a base, e.g. K_2CO_3 and a nucleophilic catalyst, e.g. KI in an appropriate solvent, e.g. acetonitrile, to yield tert-butyl N-(2-(dialkylamino)ethyl)-N-alkylglycinate (**II-6**). Deprotection of **II-6**, using e.g. TFA, yields the corresponding glycine compound (**II-7**). (c) The reaction of (**II-3**), obtained according to (a), and (**II-7**), obtained 10 according to (b), in the presence of a base, e.g. diisopropylethylamine (DIPEA), and a coupling agent, e.g. propylphosphonic acid anhydride, yields the product (**P-2**). This reaction can take place in an organic solvent, e.g. 2-methyltetrahydrofuran.


Scheme 3



15

[00561] As illustrated in Scheme 3 above, intermediates for the synthesis of certain compounds of the disclosure may be obtained by alkylating the amino group of tert-butyl 4-(2-aminoethyl)piperidine-1-carboxylate. Similar reactions can be performed with a different

starting material such as



20

[00562] In addition, it is to be understood that any particular embodiment of the present disclosure that falls within the prior art may be explicitly excluded from any one or more of the claims. Since such embodiments are deemed to be known to one of ordinary skill in the art, they may be excluded even if the exclusion is not set forth explicitly herein.

25

[00563] All cited sources, for example, references, publications, databases, database entries, and art cited herein, are incorporated into this application by reference, even if not expressly stated in the citation. In case of conflicting statements of a cited source and the instant application, the statement in the instant application shall control. Citation of publications and

patent documents is not intended as an admission that any is pertinent prior art, nor does it constitute any admission as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description and examples 5 below are for purposes of illustration and not limitation of the claims that follow.

Examples

Example 1: Synthesis of compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I).

10 A. General Considerations

[00564] All solvents and reagents used were obtained commercially and used as such unless noted otherwise. ^1H NMR spectra were recorded in CDCl_3 , at 300 K using a Bruker Ultrashield 300 MHz instrument or a Varian Unity Inova 400MHz Instrument. Chemical shifts are reported as parts per million (ppm) relative to TMS (0.00) for ^1H . Silica gel

15 chromatographies were performed on ISCO CombiFlash Rf+ Lumen Instruments using ISCO RediSep Rf Gold Flash Cartridges (particle size: 20-40 microns). Reverse phase chromatographies were performed on ISCO CombiFlash Rf+ Lumen Instruments using RediSep Rf Gold C18 High Performance columns. All final compounds were determined to be greater than 85% pure *via* analysis by reverse phase UPLC-MS (retention times, RT, in minutes) using 20 Waters Acquity UPLC instrument with DAD and ELSD and a ZORBAX Rapid Resolution High Definition (RRHD) SB-C18 LC column, 2.1 mm, 50 mm, 1.8 μm , and a gradient of 65 to 100% acetonitrile in water with 0.1% TFA over 5 minutes at 1.2 mL/min. Injection volume was 5 μL and the column temperature was 80 °C. Detection was based on electrospray ionization (ESI) in positive mode using Waters SQD mass spectrometer (Milford, MA, USA) and evaporative light 25 scattering detector.

[00565] The procedures described for the synthesis of Compounds 12 and 19 are applicable to the synthesis of compounds according to formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) generally.

30 The following abbreviations are employed herein:

rt: Room Temperature

MeOH: Methanol

DCM: Dichloromethane

DCE: Dichloroethane

DMAP: 4-Dimethylaminopyridine

DMF: *N,N*-Dimethylformamide

EtOAc: Ethylacetate

5 MeCN: Acetonitrile


THF: Tetrahydrofuran

EDC·HCl: N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide hydrochloride

B. Compound 1: 2-(Didodecylamino)-*N*-(2-(4-(didodecylamino)ethyl)piperazin-1-yl)ethyl)-

10 *N*-dodecylacetamide

Step 1: *tert*-Butyl 4-(2-(dodecylamino)ethyl)piperazine-1-carboxylate



Chemical Formula: C₂₃H₄₇N₃O₂


Molecular Weight: 397.65

15 [00566] In the same manner as Step 3 for Compound 18, *tert*-butyl 4-(2-(dodecylamino)ethyl)piperazine-1-carboxylate was synthesized from 1-bromododecane (3.3 g, 13.1 mmol), 4-(2-aminoethyl)-1-boc-piperazine (3.0 g, 13.1 mmol), K₂CO₃ (3.62 g, 26.2 mmol), and KI (217 mg, 1.31 mmol) in MeCN (60 mL). Yield (1.42 g, 27%).

UPLC/ELSD: RT = 1.18 min. MS (ES): *m/z* (MH⁺) 398.56 for C₂₃H₄₇N₃O₂

20 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.45 (br. m, 4H); 2.75 (br. m, 2H); 2.65 (br. m, 2H); 2.55 (br. m, 2H); 2.42 (br. m, 4H); 1.60-1.22 (br. m, 29H); 0.91 (br. m, 3H).

Step 2: *tert*-Butyl 4-(2-(2-(didecylamino)-*N*-dodecylacetamido)ethyl)piperazine-1-carboxylate



Chemical Formula: C₄₉H₉₈N₄O₃

Molecular Weight: 791.35


[00567] In the same manner as Step 3 for Compound 11, *tert*-butyl 4-(2-(2-(didecylamino)-*N*-dodecylacetamido)ethyl)piperazine-1-carboxylate was synthesized from *tert*-butyl 4-(2-(dodecylamino)ethyl)piperazine-1-carboxylate (100 mg, 0.25 mmol), lithium

didodecylglycine (0.10 g, 0.25 mmol), propylphosphonic acid anhydride (50% EtOAc solution, 0.45 mL, 0.75 mmol), and *i*-Pr₂EtN (0.044 mL, 0.25 mmol) in THF (2 mL). Yield (0.12 g, 63%).

UPLC/ELSD: RT = 3.36 min. MS (ES): *m/z* (MH⁺) 792.082 for C₄₉H₉₈N₄O₃

5

Step 3: 2-(Didodecylamino)-*N*-dodecyl-*N*-(2-(piperazin-1-yl)ethyl)acetamide



Chemical Formula: C₄₄H₉₀N₄O


Molecular Weight: 691.23

10 [00568] In the same manner as Step 4 for Compound 11, 2-(didodecylamino)-*N*-dodecyl-*N*-(2-(piperazin-1-yl)ethyl)acetamide was synthesized from *tert*-butyl 4-(2-(didodecylamino)-*N*-dodecylacetamido)ethyl)piperazine-1-carboxylate (0.12 g, 0.16 mmol) and TFA (0.25 mL, 3.2 mmol) in 0.25 mL DCM.

UPLC/ELSD: RT = 3.06 min. MS (ES): *m/z* (MH⁺) 692.984 for C₄₄H₉₀N₄O

15

Step 4: Compound 1: 2-(Didodecylamino)-*N*-(2-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*-dodecylacetamide



Chemical Formula: C₇₀H₁₄₃N₅O

Molecular Weight: 1070.95

20


[00569] In the same manner as Step 6 for Compound 18, 2-(didodecylamino)-*N*-(2-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*-dodecylacetamide was synthesized from 2-(didodecylamino)-*N*-dodecyl-*N*-(2-(piperazin-1-yl)ethyl)acetamide (65 mg, 0.094 mmol), *N*-(2-chloroethyl)-*N*-dodecyldecane-1-amine (42 mg, 0.10 mmol), K₂CO₃ (13 mg, 0.094 mmol) and KI (2 mg, 0.0094 mmol) in 0.5 mL MeCN to afford 58.5 mg for 58% yield.

UPLC/ELSD: RT = 3.75 min. MS (ES): *m/z* (MH⁺) 1072.585 for C₇₀H₁₄₃N₅O

¹H NMR (300 MHz, CDCl₃) δ: ppm 3.82-3.23 (br. m. 8H); 3.04-2.90 (br. m., 2H); 2.47 (m, 18H); 1.24 (m, 100H); 0.96 (m, 15H).

C. Compound 2: 2-((2-(Didodecylamino)ethyl)(dodecyl)amino)-1-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethan-1-one

Step 1: *tert*-Butyl 4-(2-(didodecylamino)ethyl)piperazine-1-carboxylate



5

Chemical Formula: C₃₅H₇₁N₃O₂


Molecular Weight: 565.97

[00570] A mixture of 1-bromododecane (1.1 mL, 4.6 mmol), 4-(2-aminoethyl)-1-boc-piperazine (1.0 g, 4.4 mmol), K₂CO₃ (0.61 g, 4.4 mmol), in 10 mL MeCN was allowed to stir at 10 rt for 12 h. After this time the reaction was filtered and concentrated. The crude material was purified by silica gel chromatography (0-20% MeOH in DCM with 1% NH₄OH to afford *tert*-butyl 4-(2-(didodecylamino)ethyl)piperazine-1-carboxylate (450 mg, 0.80 mmol, 18%).

UPLC/ELSD: RT = 2.87 min. MS (ES): *m/z* (MH⁺) 566.655 for C₃₅H₇₁N₃O₂

[00571] ¹H NMR (400 MHz, CDCl₃) δ: ppm 3.40 (m, 4H); 2.56 (m, 2H); 2.40 (m, 10H); 1.44 (s, 9H); 1.40-1.24 (m, 40H); 0.86 (t, 6H).

Step 2: *N*-Dodecyl-*N*-(2-(piperazin-1-yl)ethyl)dodecan-1-amine



Chemical Formula: C₃₀H₆₃N₃


Molecular Weight: 465.86

[00571] In the same manner as Step 5 for Compound 18, *N*-dodecyl-*N*-(2-(piperazin-1-yl)ethyl)dodecan-1-amine was synthesized from *tert*-butyl 4-(2-(didodecylamino)ethyl)piperazine-1-carboxylate (0.92 g, 1.63 mmol), TFA (6.2 mL, 82 mmol) in 6 mL DCM to afford 490 mg for 65% yield.

UPLC/ELSD: RT = 2.10 min. MS (ES): *m/z* (MH⁺) 466.379 for C₃₀H₆₃N₃

[00571] ¹H NMR (400 MHz, CDCl₃) δ: ppm 2.88 (t, 4H); 2.61 (m, 2H); 2.45 (m, 10H); 1.43-1.24 (m, 40H); 0.86 (t, 6H).

Step 3: Compound 2: 2-((2-(Didodecylamino)ethyl)(dodecyl)amino)-1-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethan-1-one



Chemical Formula: C₇₀H₁₄₃N₅O

Molecular Weight: 1070.95


In the same manner as Step 11 for Compound 11, 2-((2-

5 (didodecylamino)ethyl)(dodecyl)amino)-1-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethan-1-one was synthesized from *N*-dodecyl-*N*-(2-(piperazin-1-yl)ethyl)dodecan-1-amine (32 mg, 0.069 mmol), *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycine (43 mg, 0.069 mmol), propylphosphonic acid anhydride (50% EtOAc solution, 0.12 mL, 0.21 mmol) and *i*-Pr₂EtN (0.024 mL, 0.14 mmol) in 0.5 mL THF to provide 17.7 mg (17%).

10 UPLC: RT = 3.90 min. MS (ES): *m/z* (MH⁺) 1071.475 for C₇₀H₁₄₃N₅O
¹H NMR (400 MHz, CDCl₃) δ: ppm 3.65 (m, 2H); 3.57 (m, 2H); 3.26 (s, 2H); 2.33-2.57 (m, 22H); 1.24-1.39 (m, 100H); 0.88 (t, 15H).

D. Compound 3: 2-(Didodecylamino)-1-(4-(2-((2-

15 (didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethan-1-one
Step 1: *tert*-Butyl 4-((2-(didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazine-1-carboxylate




Chemical Formula: C₄₉H₁₀₀N₄O₂

Molecular Weight: 777.37

20 [00572] In the same manner as Step 4 for Compound 18, *tert*-butyl 4-((2-(didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazine-1-carboxylate was synthesized from *tert*-butyl 4-(2-(dodecylamino)ethyl)piperazine-1-carboxylate (700 mg, 1.76 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (806 mg, 1.93 mmol), K₂CO₃ (486 mg, 3.52 mmol), and KI (29 mg, 0.176 mmol) in THF (15 mL). Yield (683 mg, 50%).

UPLC/ELSD: RT = 3.35 min. MS (ES): *m/z* (MH⁺) 778.16 for C₄₉H₁₀₀N₄O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.44 (t, 4H); 3.11-2.86 (br. m, 4H); 2.78-2.32 (br. m, 14H); 1.80-1.05 (br. m, 69H); 0.91 (t, 9H).

Step 2: *N*¹,*N*¹,*N*²-Tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamineChemical Formula: C₄₄H₉₂N₄

Molecular Weight: 677.25


5

[00573] In the same manner as Step 5 for Compound 18, *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine was synthesized from *tert*-butyl 4-((2-(didodecylamino)ethyl)(dodecylamino)ethyl)piperazine-1-carboxylate (683 mg, 0.879 mmol), and TFA (3.4 mL, 43.9 mmol) in DCM (3.4 mL). Yield (595 mg, 99%).

10 UPLC/ELSD: RT = 2.94 min. MS (ES): *m/z* (MH⁺) 678.16 for C₄₄H₉₂N₄

Step 3: Compound 3:

2-(Didodecylamino)-1-(4-((2-(didodecylamino)ethyl)(dodecylamino)ethyl)piperazin-1-yl)ethan-1-one



15

Chemical Formula: C₇₀H₁₄₃N₅O

Molecular Weight: 1070.95


[00574] In the same manner as Step 11 for Compound 11, 2-(didodecylamino)-1-(4-((2-(didodecylamino)ethyl)(dodecylamino)ethyl)piperazin-1-yl)ethan-1-one was from *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (50 mg, 0.074 mmol), lithium didodecylglycine (33 mg, 0.078 mmol), propylphosphonic acid anhydride (50% in EtOAc, 0.13 mL, 0.22 mmol) and *i*-Pr₂EtN (0.026 mL) in 0.5 mL THF to afford 33.9 mg (43%).

UPLC: RT = 3.90 min. MS (ES): *m/z* (MH⁺) 1071.475 for C₇₀H₁₄₃N₅O

¹H NMR (400 MHz, CDCl₃) δ : ppm 3.65 (m, 2H); 3.57 (m, 2H); 3.26 (s, 2H); 2.33-2.57 (m, 22H); 1.24-1.39 (m, 100H); 0.88 (t, 15H).

E. Compound 4: 2-(Dinonylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one

Step 1: Methyl dinonylglycinate

Chemical Formula: C₂₁H₄₃NO₂


Molecular Weight: 341.58

5 [00575] In the same manner as Step 1 for Compound 11, methyl dinonylglycinate was synthesized from glycine methyl ester hydrochloride (5.0 g, 39.8 mmol), triethylamine (8.3 mL, 59.7 mmol), 95% nonanal (15.0 g, 99.6 mmol), sodium triacetoxyborohydride (21.1 g, 99.6 mmol), and acetic acid (5.7 mL, 99.6 mmol) in DCE (50 mL). Yield (3.5 g, 26%).

UPLC/ELSD: RT = 1.82 min. MS (ES): *m/z* (MH⁺) 343.62 for C₂₁H₄₃NO₂

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.72 (s, 3H); 3.35 (s, 2H); 2.57 (t, 4H); 1.46 (br. m, 4H); 1.29 (br. m, 24H); 0.90 (t, 6H).

Step 2: Lithium dinonylglycinate


Chemical Formula: C₂₀H₄₀LiNO₂

Molecular Weight: 333.49

[00576] In the same manner as Step 2 for Compound 11, lithium dinonylglycinate was synthesized from methyl dinonylglycinate (3.5 g, 10.2 mmol) and 1M LiOH (50 mL, 50 mmol) in THF (50 mL). Yield (3.0 g, 88%).

UPLC/ELSD: RT = 1.71 min. MS (ES): *m/z* (MH⁺) 328.37 for C₂₀H₄₁NO₂

¹H-NMR (300 MHz, CD₃OD) δ: ppm 3.13 (s, 2H); 2.59 (t, 4H); 1.51 (br. m, 4H); 1.32 (br. m, 24H); 0.92 (t, 6H).

Step 3: *tert*-Butyl 4-(dinonylglycyl)piperazine-1-carboxylateChemical Formula: C₂₉H₅₇N₃O₃


Molecular Weight: 495.79

25

[00577] In the same manner as Step 3 for Compound 11, *tert*-butyl 4-(dinonylglycyl)piperazine-1-carboxylate was synthesized from lithium dinonylglycinate (2.0 g, 6.00 mmol), 1-boc-piperazine (1.23 g, 6.58 mmol), *i*-Pr₂EtN (2.3 mL, 13.2 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 10.7 mL, 17.9 mmol). Yield (824 mg, 5 28%).

UPLC/ELSD: RT = 2.19 min. MS (ES): *m/z* (MH⁺) 496.72 for C₂₉H₅₇N₃O₃

Step 4: 2-(Dinonylamino)-1-(piperazin-1-yl)ethan-1-one



Chemical Formula: C₂₄H₄₉N₃O


10 Molecular Weight: 395.68

[00578] In the same manner as Step 4 for Compound 11, 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one was synthesized from *tert*-butyl 4-(dinonylglycyl)piperazine-1-carboxylate (824 mg, 1.66 mmol) and TFA (6.4 mL, 83.1 mmol) in DCM (6.4 mL). Yield (246 mg, 37%).

UPLC/ELSD: RT = 1.25 min. MS (ES): *m/z* (MH⁺) 396.68 for C₂₄H₄₉N₃O

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.63 (br. m, 4H); 3.28 (s, 2H); 2.89 (br. m, 4H); 2.48 (t, 4H); 1.45 (br. m, 4H); 1.28 (br. m, 24H); 0.90 (t, 6H).

Step 5: Methyl N-(*tert*-butoxy carbonyl)-N-nonylglycinate




20 Chemical Formula: C₁₇H₃₃NO₄

Molecular Weight: 315.45

[00579] In the same manner as Step 5 for Compound 11, methyl N-(*tert*-butoxy carbonyl)-N-nonylglycinate was synthesized from N-(*tert*-butoxy carbonyl)glycine methyl ester (7.7 g, 40.7 mmol) and NaH (60%, 1.71 g, 42.7 mmol) in DMF (100 mL). Yield (3.32 g, 26%).

25 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.02-3.84 (br. m, 2H); 3.75 (s, 3H); 3.26 (br. m, 2H); 1.65-1.39 (br. m, 11H); 1.28 (br. m, 12H); 0.90 (t, 3H).

Step 6: Methyl nonylglycinate




Chemical Formula: C₁₂H₂₅NO₂

Molecular Weight: 215.34

[00580] In the same manner as Step 6 for Compound 11, methyl nonylglycinate was synthesized from methyl *N*-(*tert*-butoxycarbonyl)-*N*-nonylglycinate (3.32 g, 10.5 mmol) and TFA (16 mL, 210 mmol) in DCM (16 mL). Yield (2.23 g, 98%).
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.75 (s, 3H); 3.44 (s, 2H); 2.61 (t, 2H); 1.69 (br. m, 1H); 1.51 (br. m, 2H); 1.28 (br. m, 12H); 0.90 (t, 3H).

10 Step 7: Methyl *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycinate




Chemical Formula: C₃₂H₆₆N₂O₂

Molecular Weight: 510.89

[00581] In the same manner as Step 9 for Compound 11, methyl *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycinate was synthesized from methyl nonylglycinate (449 mg, 2.08 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (830 mg, 2.50 mmol), K₂CO₃ (576 mg, 4.16 mmol), and KI (35 mg, 0.208 mmol) in MeCN (13 mL). Yield (958 mg, 90%).
UPLC/ELSD: RT = 3.11 min. MS (ES): *m/z* (MH⁺) 511.97 for C₃₂H₆₆N₂O₂
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.72 (s, 3H); 3.42 (s, 2H); 2.95-2.15 (br. m, 10H); 1.85-1.00 (br. m, 42H); 0.90 (t, 9H).

Step 8: *N*-(2-(Dinonylamino)ethyl)-*N*-nonylglycine




Chemical Formula: C₃₁H₆₄N₂O₂

Molecular Weight: 496.87

25

[00582] In the same manner as Step 10 for Compound 11, *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine was synthesized from methyl *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycinate (958 mg, 1.88 mmol), and 1M LiOH (10 mL, 10 mmol) in THF (10 mL). Yield (514 mg, 55%).
 UPLC/ELSD: RT = 2.75 min. MS (ES): *m/z* (MH⁺) 497.95 for C₃₁H₆₄N₂O₂
 5 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.92 (br. m, 6H); 3.14 (br. m, 6H); 1.77 (br. m, 6H); 1.45-1.13 (br. m, 36H); 0.90 (t, 9H).

Step 9: Compound 4: 2-(Dinonylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: C₅₅H₁₁₁N₅O₂


Molecular Weight: 874.53

[00583] In the same manner as Step 11 for Compound 11, 2-(dinonylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (61.5 mg, 0.155 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (85 mg, 0.171 mmol), *i*-Pr₂EtN (60 μL, 0.342 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.278 mL, 0.466 mmol). Yield (38 mg, 28%).

UPLC/ELSD: RT = 3.13 min. MS (ES): *m/z* (MH⁺) 875.76 for C₅₅H₁₁₁N₅O₂
 20 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.82-3.49 (br. m, 8H); 3.33 (s, 2H); 3.27 (s, 2H); 2.68-2.18 (br. m, 14H); 1.82-1.02 (br. m, 70H); 0.90 (t, 15H).

F. Compound 5: 2-(Dinonylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one

25 Step 1: Methyl *N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycinate




Chemical Formula: C₃₅H₇₂N₂O₂

Molecular Weight: 552.97

[00584] In the same manner as Step 9 for Compound 11, methyl *N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycinate was synthesized from methyl dodecylglycinate (535 mg, 2.08 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (830 mg, 2.50 mmol), K_2CO_3 (576 mg, 4.16 mmol), and KI (35 mg, 0.208 mmol) in MeCN (13 mL). Yield (385 mg, 34%).

5 UPLC/ELSD: RT = 3.34 min. MS (ES): m/z (MH^+) 553.96 for $C_{35}H_{72}N_2O_2$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.72 (s, 3H); 3.41 (s, 2H); 2.90-2.20 (br. m, 10H); 1.85-1.05 (br. m, 48H); 0.90 (t, 9H).

Step 2: *N*-(2-(Dinonylamino)ethyl)-*N*-dodecylglycine



10

Chemical Formula: $C_{34}H_{70}N_2O_2$

Molecular Weight: 538.95


[00585] In the same manner as Step 10 for Compound 11, *N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycine was synthesized from methyl *N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycinate (385 mg, 0.696 mmol), and 1M LiOH (3.5 mL, 3.5 mmol) in THF (3.5 mL). Yield (225 mg, 60%).

UPLC/ELSD: RT = 3.13 min. MS (ES): m/z (MH^+) 539.93 for $C_{34}H_{70}N_2O_2$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.73 (s, 2H); 3.62-3.39 (br. m, 4H); 3.09 (br. m, 6H); 1.76 (br. m, 6H); 1.28 (br, 42H); 0.90 (t, 9H).

20

Step 3: Compound 5:

2-(Dinonylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: $C_{58}H_{117}N_5O_2$

Molecular Weight: 916.61

25

[00586] In the same manner as Step 11 for Compound 11, 2-(dinonylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (62 mg, 0.155 mmol), *N*-(2-

(dinonylamino)ethyl)-*N*-dodecylglycine (92 mg, 0.171 mmol), *i*-Pr₂EtN (60 μ L, 0.342 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.278 mL, 0.466 mmol). Yield (38 mg, 26%).


UPLC/ELSD: RT = 3.32 min. MS (ES): *m/z* (MH⁺) 917.67 for C₅₈H₁₁₇N₅O₂

5 ¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.86-3.45 (br. m, 8H); 3.33 (s, 2H); 3.28 (s, 2H); 2.73-2.27 (br. m, 14H); 1.86-1.00 (76H); 0.91 (t, 15H).

G. Compound 6:

2-((2-(Didodecylamino)ethyl)(nonyl)amino)-1-(4-(dinonylglycyl)piperazin-1-yl)ethan-1-one

10 Step 1: Methyl *N*-(2-(didodecylamino)ethyl)-*N*-nonylglycinate



Chemical Formula: C₃₈H₇₈N₂O₂

Molecular Weight: 595.05


[00587] In the same manner as Step 9 for Compound 11, methyl *N*-(2-

15 (didodecylamino)ethyl)-*N*-nonylglycinate was synthesized from methyl nonylglycinate (355 mg, 1.65 mmol), *N*-(2-chloroethyl)-*N*-dodecyldecylamine (825 mg, 1.98 mmol), K₂CO₃ (457 mg, 3.30 mmol), and KI (27 mg, 0.165 mmol) in MeCN (10 mL). Yield (460 mg, 47%).

UPLC/ELSD: RT = 3.62 min. MS (ES): *m/z* (MH⁺) 596.03 for C₃₈H₇₈N₂O₂

19 ¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.72 (s, 3H); 3.42 (s, 2H); 2.80-2.24 (br. m, 10H); 1.56-2.00 (br. m, 54H); 0.90 (t, 9H).

Step 2: *N*-(2-(Didodecylamino)ethyl)-*N*-nonylglycine



Chemical Formula: C₃₇H₇₆N₂O₂

Molecular Weight: 581.03


[00588] In the same manner as Step 10 for Compound 11, *N*-(2-(didodecylamino)ethyl)-*N*-nonylglycine was synthesized from methyl *N*-(2-(didodecylamino)ethyl)-*N*-nonylglycinate (460 mg, 0.773 mmol), and 1M LiOH (3.9 mL, 3.9 mmol) in THF (3.9 mL). Yield (323 mg, 72%).

UPLC/ELSD: RT = 3.37 min. MS (ES): m/z (MH^+) 582.00 for $C_{37}H_{76}N_2O_2$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.17 (s, 2H); 4.00 (br. m, 2H); 3.84 (br. m, 2H); 3.34 (br. m, 2H); 3.18 (br. m, 4H); 1.82 (br. m, 6H); 1.27 (br. m, 48H); 0.91 (t, 9H).

5 Step 3: Compound 6:

2-((2-(Didodecylamino)ethyl)(nonyl)amino)-1-(4-(dinonylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: $C_{61}H_{123}N_5O_2$

Molecular Weight: 958.69


10 [00589] In the same manner as Step 11 for Compound 11, 2-((2-(didodecylamino)ethyl)(nonyl)amino)-1-(4-(dinonylglycyl)piperazin-1-yl)ethan-1-one was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (62 mg, 0.155 mmol), *N*-(2-(didodecylamino)ethyl)-*N*-nonylglycine (99 mg, 0.171 mmol), *i*-Pr₂EtN (60 μ L, 0.342 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.278 mL, 0.466 mmol). Yield (45 mg, 30%).

15 UPLC/ELSD: RT = 3.46 min. MS (ES): m/z (MH^+) 959.98 for $C_{61}H_{123}N_5O_2$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.81-3.49 (br. m, 8H); 3.33 (s, 2H); 3.27 (s, 2H); 2.70-2.25 (br. m, 14H); 1.90-1.00 (br. m, 82H); 0.90 (t, 15H).

20 H. Compound 7:

2-((2-(Didodecylamino)ethyl)(dodecyl)amino)-1-(4-(dinonylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: $C_{64}H_{129}N_5O_2$

Molecular Weight: 1000.77


25 [00590] In the same manner as Step 11 for Compound 11, 2-((2-(didodecylamino)ethyl)(dodecyl)amino)-1-(4-(dinonylglycyl)piperazin-1-yl)ethan-1-one was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (62 mg, 0.155 mmol), *N*-(2-

(didodecylamino)ethyl)-*N*-dodecylglycine (107 mg, 0.171 mmol), *i*-Pr₂EtN (60 μ L, 0.342 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.278 mL, 0.466 mmol). Yield (34 mg, 20%).

5 UPLC/ELSD: RT = 3.60 min. MS (ES): *m/z* (MH⁺) 1001.97 for C₆₄H₁₂₉N₅O₂
¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.85-2.18 (br. m, 26H); 1.91-1.00 (br. m, 88H); 0.90 (t, 15H).

I. Compound 8:

2-(Didodecylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one



10

Chemical Formula: C₆₁H₁₂₃N₅O₂


Molecular Weight: 958.69

[00591] In the same manner as Step 11 for Compound 11, 2-(didodecylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one was synthesized from 2-(didodecylamino)-1-(piperazin-1-yl)ethan-1-one (202 mg, 0.421 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (230 mg, 0.463 mmol), *i*-Pr₂EtN (0.162 mL, 0.926 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.752 mL, 1.26 mmol). Yield (148 mg, 37%).

15 UPLC/ELSD: RT = 3.41 min. MS (ES): *m/z* (MH⁺) 959.74 for C₆₁H₁₂₃N₅O₂
¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.82-3.49 (br. m, 8H); 3.33 (s, 2H); 3.27 (s, 2H); 2.66-2.30 (br. m, 14H); 1.85-1.02 (br. m, 82H), 0.90 (t, 15H).

J. Compound 9:

2-(Didodecylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: C₆₄H₁₂₉N₅O₂


Molecular Weight: 1000.77

[00592] In the same manner as Step 11 for Compound 11, 2-(didodecylamino)-1-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one was synthesized from (76 mg, 0.157 mmol), (93 mg, 0.173 mmol), *i*-Pr₂EtN (60 μ L, 0.342 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.278 mL, 0.466 mmol). Yield (59 mg, 37%).

5 UPLC/ELSD: RT = 3.57 min. MS (ES): *m/z* (MH⁺) 1001.65 for C₆₄H₁₂₉N₅O₂
¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.95-2.23 (br. m, 26H); 2.05-1.00 (br. m, 88H); 0.90 (t, 15H).

10 K. Compound 10:

2-(Didodecylamino)-1-(4-(*N*-(2-(didodecylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: C₆₇H₁₃₅N₅O₂

15 Molecular Weight: 1042.85

[00593] In the same manner as Step 11 for Compound 11, 2-(didodecylamino)-1-(4-(*N*-(2-(didodecylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)ethan-1-one was synthesized from 2-(didodecylamino)-1-(piperazin-1-yl)ethan-1-one (76 mg, 0.157 mmol), *N*-(2-(didodecylamino)ethyl)-*N*-nonylglycine (101 mg, 0.173 mmol), *i*-Pr₂EtN (60 μ L, 0.342 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 0.278 mL, 0.466 mmol). Yield (56 mg, 34%).

20 UPLC/ELSD: RT = 3.72 min. MS (ES): *m/z* (MH⁺) 1043.88 for C₆₇H₁₃₅N₅O₂


¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.95-2.15 (br. m, 26H); 1.90-1.05 (br. m, 94H); 0.90 (t, 15H).

25

L. Compound 11:

2-(Didodecylamino)-1-(4-(*N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one

Step 1: Methyl didodecylglycinate

Chemical Formula: C₂₇H₅₅NO₂

Molecular Weight: 425.74


[00594] A solution of glycine methyl ester hydrochloride (5.0 g, 39.8 mmol) and

5 triethylamine (8.3 mL, 59.7 mmol) in DCE (50 mL) was allowed to stir at room temperature for 15 minutes. A solution of 92% dodecanol (20.0 g, 99.6 mmol) in DCE (50 mL) was added and the mixture was cooled to 0 °C. Sodium triacetoxyborohydride (21.1 g, 99.6 mmol) and acetic acid (5.7 mL, 99.6 mmol) were added and the reaction was allowed to return to room temperature and stir for 16 hours. The reaction was quenched by slow addition of saturated 10 sodium bicarbonate and extracted with DCM. The combined extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-30% EtOAc/hexanes) provided methyl didodecylglycinate (7.7 g, 45%).

UPLC/ELSD: RT = 2.82 min. MS (ES): *m/z* (MH⁺) 426.69 for C₂₇H₅₅NO₂

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.72 (s, 3H); 3.35 (s, 2H); 2.57 (t, 4H); 1.46 (m, 4H); 1.28 (br. m, 36H); 0.91 (t, 6H).

Step 2: Lithium didodecylglycinate

Chemical Formula: C₂₆H₅₂LiNO₂


Molecular Weight: 417.65

[00595] A solution of methyl didodecylglycinate (7.7 g, 18.1 mmol) in THF (100 mL)

and 1M LiOH (90.4 mL, 90.4 mmol) was allowed to stir at 65 °C for 16 hours. The reaction 25 was cooled to room temperature and concentrated to a white powder. The powder was suspended in water, filtered, washed with water and diethyl ether, and dried under vacuum to provide lithium didodecylglycinate (7.0 g, 93%).

UPLC/ELSD: RT = 2.74 min. MS (ES): *m/z* (MH⁺) 412.83 for C₂₆H₅₃NO₂

¹H-NMR (300 MHz, CD₃OD) δ: ppm 3.14 (s, 2H); 2.60 (t, 4H); 1.51 (m, 4H); 1.31 (br. m, 36H); 0.92 (t, 6H).


Step 3: *tert*-Butyl 4-(didodecylglycyl)piperazine-1-carboxylateChemical Formula: C₃₅H₆₉N₃O₃

Molecular Weight: 579.96

5 [00596] A solution of lithium didodecylglycinate (2.0 g, 4.79 mmol), 1-boc-piperazine (978 mg, 5.25 mmol), *i*-Pr₂EtN (1.84 mL, 10.5 mmol), and propylphosphonic acid anhydride (50% EtOAc solution, 8.53 mL, 14.3 mmol) in THF (24 mL) was allowed to stir at room temperature for 48 hours. The reaction was diluted with water and extracted with EtOAc. The organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided *tert*-butyl 4-(didodecylglycyl)piperazine-1-carboxylate (983 mg, 35%).

10 UPLC/ELSD: RT = 3.06 min. MS (ES): *m/z* (MH⁺) 581.02 for C₃₅H₆₉N₃O₃

Step 4: 2-(Didodecylamino)-1-(piperazin-1-yl)ethan-1-one

Chemical Formula: C₃₀H₆₁N₃O


Molecular Weight: 479.84

15 [00597] To a 0 °C solution of *tert*-butyl 4-(didodecylglycyl)piperazine-1-carboxylate (983 mg, 1.69 mmol) in DCM (6.5 mL) was added dropwise TFA (6.5 mL, 84.7 mmol). The reaction was allowed to return to room temperature and stir for 16 hours. The reaction mixture was concentrated *in vacuo* and the crude material was dissolved in CHCl₃. The solution was washed with 5% Na₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo* to provide 2-(didodecylamino)-1-(piperazin-1-yl)ethan-1-one (163 mg, 20%).

UPLC/ELSD: RT = 2.07 min. MS (ES): *m/z* (MH⁺) 480.89 for C₃₀H₆₁N₃O

20 [00597] ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.67 (br. m, 4H); 3.32 (s, 2H); 2.92 (br. m, 4H); 2.53 (br. m, 4H); 1.48 (br. m, 4H); 1.28 (br. m, 36H); 0.91 (t, 6H).

Step 5: Methyl *N*-(*tert*-butoxycarbonyl)-*N*-dodecylglycinate




Chemical Formula: C₂₀H₃₉NO₄

Molecular Weight: 357.54

[00598] A 0 °C solution of *N*-(*tert*-butoxy carbonyl)glycine methyl ester (7.7 g, 40.7 mmol) in DMF (100 mL) was treated with NaH (60%, 1.71 g, 42.7 mmol) and the mixture was allowed to stir for 30 minutes. The solution was allowed to return to room temperature before 1-bromododecane (15.2 g, 61.0 mmol) was added. The reaction was quenched with water and extracted with EtOAc. The organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% EtOAc/hexanes) provided methyl *N*-(*tert*-butoxy carbonyl)-*N*-dodecylglycinate (4.03 g, 28%).
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.01-3.84 (br. m, 2H); 3.75 (s, 3H); 3.27 (br. m, 2H); 1.67-1.39 (br. m, 11H); 1.28 (br. m, 18H); 0.90 (t, 3H).

Step 6: Methyl dodecylglycinate




Chemical Formula: C₁₅H₃₁NO₂

Molecular Weight: 257.42

[00599] To a 0 °C solution of methyl N-(tert-butoxycarbonyl)-N-dodecylglycinate (4.03 g, 11.3 mmol) in DCM (17 mL) was added dropwise TFA (17 mL, 226 mmol). The reaction was allowed to return to room temperature and stir for 6 hours. The reaction mixture was concentrated in vacuo and the crude material was dissolved in DCM. The solution was washed with 10% NaOH, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated in vacuo to provide methyl dodecylglycinate (2.84 g, 98%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.75 (s, 3H); 3.44 (s, 2H); 2.62 (t, 2H); 1.70 (br, 1H); 1.51 (m, 2H); 1.29 (br, m, 18H); 0.90 (t, 3H).

Step 7: 2-(Didodecylamino)ethan-1-ol



Chemical Formula: C₂₆H₅₅NO


Molecular Weight: 397.73

[00600] In the same manner as Step 1 for Compound 18, 2-(didodecylamino)ethan-1-ol was synthesized from 1-bromododecane (10 g, 40.1 mmol), ethanolamine (1.10 mL, 18.2 mmol), K_2CO_3 (11.1 g, 80.1 mmol), and KI (302 mg, 1.82 mmol) in MeCN (84 mL). Yield 5 (3.87 g, 53%).

UPLC/ELSD: RT = 2.69 min. MS (ES): m/z (MH^+) 398.56 for $C_{26}H_{55}NO$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.57 (t, 2H); 2.63 (t, 2H); 2.49 (br. m, 4H); 1.48 (br. m, 4H); 1.29 (br. m, 36H); 0.91 (t, 6H).

10 Step 8: *N*-(2-Chloroethyl)-*N*-dodecylidodecan-1-amine



Chemical Formula: $C_{26}H_{54}ClN$


Molecular Weight: 416.18

[00601] In the same manner as Step 2 for Compound 18, *N*-(2-chloroethyl)-*N*-dodecylidodecan-1-amine was synthesized from 2-(didodecylamino)ethan-1-ol (3.87 g, 9.73 mmol), triethylamine (1.76 mL, 12.6 mmol), and methanesulfonyl chloride (0.941 mL, 12.2 mmol) in DCM (50 mL). Yield (1.92 g, 47%).

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.51 (t, 2H); 2.78 (t, 2H); 2.47 (br. m, 4H); 1.44 (br. m, 4H); 1.28 (br. m, 36H); 0.90 (t, 6H).

20

Step 9: Methyl *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycinate



Chemical Formula: $C_{41}H_{84}N_2O_2$

Molecular Weight: 637.14


[00602] To a solution of methyl dodecylglycinate (425 mg, 1.65 mmol) in MeCN (10 mL) was added *N*-(2-chloroethyl)-*N*-dodecylidodecan-1-amine (825 mg, 1.98 mmol), K_2CO_3 (457 mg, 3.30 mmol), and KI (27 mg, 0.165 mmol). The reaction was allowed to stir at 82 °C for 72 hours. The reaction mixture was filtered and the solids were washed with hexanes. The filtrate was concentrated *in vacuo* to provide the crude product. Purification by ISCO silica flash

chromatography (0-20% MeOH/DCM) provided methyl *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycinate (652 mg, 62%).

UPLC/ELSD: RT = 3.77 min. MS (ES): *m/z* (MH⁺) 638.18 for C₄₁H₈₄N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.72 (s, 3H); 3.41 (s, 2H); 2.90-2.20 (br. m, 10H); 1.60-5 1.00 (br. m, 60H); 0.90 (t, 9H).

Step 10: *N*-(2-(Didodecylamino)ethyl)-*N*-dodecylglycine



Chemical Formula: C₄₀H₈₂N₂O₂


Molecular Weight: 623.11

[00603] A solution of methyl *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycinate (652 mg, 1.02 mmol) in THF (6 mL) and 1M LiOH (5 mL, 5 mmol) was allowed to stir at 65 °C for 16 hours. The reaction was cooled to room temperature and acidified with 10% HCl. The mixture was extracted with chloroform, and the organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycine (153 mg, 24%).

UPLC/ELSD: RT = 3.60 min. MS (ES): *m/z* (MH⁺) 624.07 for C₄₀H₈₂N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.02-3.40 (br. m, 6H); 3.16 (br. m, 6H); 1.78 (br. m, 6H); 1.46-1.01 (br. m, 54H); 0.90 (t, 9H).

Step 11: Compound 11: 2-(Didodecylamino)-1-(4-(*N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycyl)piperazin-1-yl)ethan-1-one



Chemical Formula: C₇₀H₁₄₁N₅O₂

Molecular Weight: 1084.93


[00604] To a solution of *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycine (212 mg, 0.340 mmol) and 2-(didodecylamino)-1-(piperazin-1-yl)ethan-1-one (163 mg, 0.340 mmol) in THF (4 mL) was added *i*-Pr₂EtN (0.119 mL, 0.680 mmol), and propylphosphonic acid anhydride (50% 25

EtOAc solution, 0.606 mL, 1.02 mmol). The reaction was allowed to stir at room temperature overnight. The reaction mixture was diluted with water and extracted with EtOAc. The organics were washed with brine, dried over anhydrous Na₂SO₄, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% [DCM, 20% MeOH, 1% NH₄OH]/MeOH) provided 2-(didecylamino)-1-(4-(N-(2-(didecylamino)ethyl)-N-dodecylglycyl)piperazin-1-yl)ethan-1-one (148 mg, 37%).

5 UPLC/ELSD: RT = 3.81 min. MS (ES): *m/z* (MH⁺) 1086.94 for C₇₀H₁₄₁N₅O₂
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.00-2.20 (br. m, 26H); 1.77 (br. m, 6H); 1.54-1.02 (br. m, 94H); 0.90 (t, 15H).

10

M. Compound 12: Petyl 6-((2-(4-(2-((2-(didecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)hexanoate
Step 1: Petyl 6-bromohexanoate



15


Chemical Formula: C₁₁H₂₁BrO₂

Molecular Weight: 265.19

[00605] To a solution of 6-bromohexanoic acid (2 g, 10.3 mmol) and pentan-1-ol (2.2 mL, 20.5 mmol) in 26 mL DCM, EDC·HCl (1.97 g, 10.3 mmol) and DMAP (0.26 g, 2.1 mmol) were added. The solution was allowed to stir at rt overnight. After this time the reaction was 20 quenched by the addition of water. The mixture was extracted three times with DCM. The organics were pooled and washed with saturated NaHCO₃, 10% citric acid and brine. The organics were then dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was purified via silica gel chromatography (0-30% EtOAc in hexanes) to afford the desired product (2.3 g, 8.67 mmol).

25 ¹H NMR (400 MHz, CDCl₃) δ: ppm 4.06 (t, 2H); 3.39 (t, 2H); 2.30 (t, 2H); 1.84 (m, 2H); 1.62 (m, 4H); 1.46 (m, 2H); 1.31 (m, 4H); 0.88 (t, 3H).

Step 2: 2-(Dodecylamino)ethan-1-ol



30


Chemical Formula: C₁₄H₃₁NO

Molecular Weight: 229.41

[00606] Methyl dodecylglycinate (3.4 g, 13.2 mmol) was dissolved in 2 mL THF under N₂ atmosphere and the reaction flask was allowed to cool in an ice bath. To the solution LiAlH₄ (0.55 g, 14.5 mmol) was slowly added. The reaction was allowed to stir at the same temperature for 1 h. After this time the reaction was quenched by the subsequent addition of 0.55 mL H₂O, 5 0.55 mL 10% NaOH and then 1.65 mL of H₂O. The reaction was then filtered and the filtrate was concentrated *in vacuo*. The crude material was purified *via* silica gel chromatography (0-20% MeOH in DCM, with 1% NaOH) to afford the desired alcohol (1.9 g, 8.28 mmol, 63% yield).

¹H NMR (400 MHz, CDCl₃) δ: ppm 3.63 (t, 2H); 2.78 (t, 2H); 2.63 (t, 2H); 1.48 (m, 2H); 2.14 10 (m, 18H); 0.88 (t, 3H).

Step 3: Pentyl 6-(dodecyl(2-hydroxyethyl)amino)hexanoate




Chemical Formula: C₂₅H₅₁NO₃

Molecular Weight: 413.69

[00607] In the same manner as Step 1 for Compound 18, pentyl 6-(dodecyl(2-hydroxyethyl)amino)hexanoate was synthesized from pentyl 6-bromohexanoate (0.87 g, 3.27 mmol), 2-(dodecylamino)ethan-1-ol (0.50 g, 2.18 mmol), K₂CO₃ (0.60 g, 4.36 mmol) and KI (36 mg, 0.22 mmol) in 10 mL THF to afford 0.30 g of the desired product (33%).

¹H NMR (400 MHz, CDCl₃) δ: ppm 4.04 (t, 2H); 3.51 (m, 2H); 2.56 (m, 2H); 2.42 (m, 4H); 2.28 (t, 2H); 1.60 (m, 4H); 1.42 (m, 4H); 1.30-1.24 (m, 24); 0.87 (m, 6H).

Step 4: Pentyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate



Chemical Formula: C₂₅H₅₀ClNO₂

Molecular Weight: 432.13


[00608] In the same manner as Step 2 for Compound 18, pentyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate was synthesized from pentyl 6-(dodecyl(2-hydroxyethyl)amino)hexanoate (300 mg, 0.73 mmol), methanesulfonyl chloride (0.062 mL, 0.80

mmol) and triethylamine (0.13 mL, 1.3 mmol) in 2 mL DCM to afford 285 mg of the desired product (66%).

¹H NMR (400 MHz, CDCl₃) δ: ppm 4.04 (t, 2H); 3.45 (t, 2H); 2.74 (t, 2H); 2.43 (m, 4H); 2.28 (t, 2H); 1.65-1.59 (m, 4H); 1.31-1.24 (m, 32H); 0.88 (m, 6H).

5

Step 5: Compound 12: Penty 6-((2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)hexanoate



Chemical Formula: C₆₉H₁₄₁N₅O₂

10

Molecular Weight: 1072.92

In the same manner as Step 6 for Compound 18, penty 6-((2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)hexanoate was synthesized from penty 6-((2-chloroethyl)(dodecyl)amino)hexanoate (75 mg, 0.17 mmol), N¹,N¹,N²-tridodecyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (107 mg, 0.16 mmol), K₂CO₃ (23 mg, 0.17 mmol) and KI (2.7 mg, 0.1 mmol) in 1 mL MeCN to afford 99 mg of the desired product (58%).


UPLC: RT = 3.53 min. MS (ES): *m/z* (MH⁺) 1073.325 for C₆₉H₁₄₁N₅O₂

¹H NMR (400 MHz, CDCl₃) δ: ppm 4.03 (t, 2H); 2.56-2.37 (br. m., 30H); 2.27 (t, 2H); 1.61 (m, 4H); 1.40-1.23 (br. m.; 90H); 0.87 (m, 15H).

20

N. Compound 13: Penty 6-((2-(4-(2-((2-ditetradecylamino)ethyl)(tetradecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)hexanoate

Step 1: 2-(Ditetradecylamino)ethan-1-ol



Chemical Formula: C₃₀H₆₃NO


25

Molecular Weight: 453.84

[00609] In the same manner as Step 1 for Compound 18, 2-(ditetradecylamino)ethan-1-ol was synthesized from 1-bromotetradecane (21.6 mL, 72.8 mmol), ethanolamine (2 mL, 33.1 mmol), K₂CO₃ (20 g, 145.5 mmol), and KI (549 mg, 3.31 mmol) in MeCN (165 mL). Yield (12 g, 81%).

5 UPLC/ELSD: RT = 3.30 min. MS (ES): *m/z* (MH⁺) 454.46 for C₃₀H₆₃NO
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.54 (br. m, 2H); 2.59 (br. m, 2H); 2.46 (br. m, 4H); 1.56-1.17(br. m, 48H); 0.90 (br. m, 6H).

Step 2: *N*-(2-Chloroethyl)-*N*-tetradecyltetradecan-1-amine




Chemical Formula: C₃₀H₆₂ClN

Molecular Weight: 472.28

[00610] In the same manner as Step 2 for Compound 18, *N*-(2-chloroethyl)-*N*-tetradecyltetradecan-1-amine was synthesized from 2-(ditetradecylamino)ethan-1-ol (10 g, 22.0 mmol), triethylamine (4.0 mL, 28.6 mmol), and methanesulfonyl chloride (2.75 mL, 27.5 mmol) in DCM (110 mL). Crude material was carried onto next step without purification. Yield (10.2 g, 98%).

UPLC/ELSD: RT = 3.37 min. MS (ES): *m/z* (MH⁺) 472.45 for C₃₀H₆₂ClN
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.27-2.20 (br. m, 8H); 1.96-1.17 (br. m, 48H); 0.90 (br. m, 6H).

Step 3: *tert*-Butyl 4-(2-(tetradecylamino)ethyl)piperazine-1-carboxylate



Chemical Formula: C₂₅H₅₁N₃O₂


Molecular Weight: 425.70

[00611] In the same manner as Step 3 for Compound 18, *tert*-butyl 4-(2-(tetradecylamino)ethyl)piperazine-1-carboxylate was synthesized from 1-bromotetradecane (3.63 g, 13.1 mmol), 4-(2-aminoethyl)-1-boc-piperazine (3.0 g, 13.1 mmol), K₂CO₃ (3.62 g, 26.2 mmol), and KI (217 mg, 1.31 mmol) in MeCN (60 mL). Yield (1.42 g, 27%).

UPLC/ELSD: RT = 1.58 min. MS (ES): m/z (MH^+) 426.61 for $C_{25}H_{51}N_3O_2$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.45 (t, 4H); 2.75 (t, 2H) 2.65 (t, 2H); 2.54 (t, 2H); 2.42 (t, 4H); 1.61-1.41 (br. m, 11H); 1.40-1.20 (br. m, 22H); 0.90 (t, 3H).

5 Step 4: *tert*-Butyl 4-((2-(ditetradecylamino)ethyl)(tetradecyl)amino)ethyl)piperazine-1-carboxylate




Chemical Formula: $C_{55}H_{112}N_4O_2$

Molecular Weight: 861.53

10 [00612] In the same manner as Step 4 for Compound 18, *tert*-butyl 4-((2-(ditetradecylamino)ethyl)(tetradecyl)amino)ethyl)piperazine-1-carboxylate was synthesized from *tert*-butyl 4-(tetradecylamino)ethyl)piperazine-1-carboxylate (700 mg, 1.64 mmol), *N*-(2-chloroethyl)-*N*-tetradecyltetradecan-1-amine (1.01 g, 2.14 mmol), K_2CO_3 (455 mg, 3.29 mmol), and KI (27 mg, 0.164 mmol) in THF (15 mL). Yield (740 mg, 52%).

15 UPLC/ELSD: RT = 3.81 min. MS (ES): m/z (MH^+) 862.47 for $C_{55}H_{112}N_4O_2$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.45 (br. m, 4H); 3.10-2.83 (br. m, 4H); 2.74-2.34 (br. m, 14H); 1.75-1.20 (br. m, 81H); 0.91 (t, 9H).

Step 5: N^1 -(2-(Piperazin-1-yl)ethyl)- N^1,N^2 -tritetradecylethane-1,2-diamine




Chemical Formula: $C_{50}H_{104}N_4$

Molecular Weight: 761.41

20 [00613] In the same manner as Step 5 for Compound 18, N^1 -(2-(piperazin-1-yl)ethyl)- N^1,N^2 -tritetradecylethane-1,2-diamine was synthesized from *tert*-butyl 4-((2-(ditetradecylamino)ethyl)(tetradecyl)amino)ethyl)piperazine-1-carboxylate (740 mg, 0.859 mmol), and TFA (3.3 mL, 42.9 mmol) in DCM (3.3 mL). Yield (661 mg, 99%).

UPLC/ELSD: RT = 3.38 min. MS (ES): m/z (MH^+) 762.42 for $C_{50}H_{104}N_4$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 2.92 (t, 4H); 2.70-2.30 (br. m, 18H); 1.46 (br. m, 6H); 1.37-1.20 (br. m, 66H); 0.90 (t, 9H).

Step 6: Compound 13: Pentyl 6-((2-(4-(2-((2-ditetradeclamino)ethyl)(tetradecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)hexanoate




5

Chemical Formula: C₇₅H₁₅₃N₅O₂

Molecular Weight: 1157.08

[00614] In the same manner as Step 6 for Compound 18, pentyl 6-((2-(4-(2-((2-ditetradeclamino)ethyl)(tetradecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)hexanoate was synthesized from N¹-(2-(piperazin-1-yl)ethyl)-N¹,N²,N²-tritetradeclylethane-1,2-diamine (66 mg, 0.087 mmol), pentyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate (42 mg, 0.095 mmol) K₂CO₃ (24 mg, 0.17 mmol), and KI (2 mg, 0.012 mmol) in THF (2 mL). Yield (38 mg, 38%).
 UPLC/ELSD: RT = 3.81 min. MS (ES): *m/z* (MH⁺) 1157.70 for C₇₅H₁₅₃N₅O₂
 15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (m, 2H); 3.16-2.15 (br. m, 32H); 1.65 (br. m, 4H); 1.54-1.00 (br. m, 100H); 0.91 (br. m, 15H).

O. Compound 14: Dipentyl 6,6'-((2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)azanediyl)dihexanoate
 20 Step 1: Dipentyl 6,6'-((2-hydroxyethyl)azanediyl)dihexanoate



Chemical Formula: C₂₄H₄₇NO₅

Molecular Weight: 429.64


[00615] In the same manner as Step 1 for Compound 18, dipentyl 6,6'-((2-hydroxyethyl)azanediyl)dihexanoate was synthesized from pentyl 6-bromohexanoate (0.50 g,

1.89 mmol), ethanolamine (0.052 mL, 0.86 mmol), K_2CO_3 (0.52 g, 3.77 mmol) and KI (14 mg, 0.098 mmol) in 4 mL MeCN to provide 234 mg (55%).

1H NMR (400 MHz, $CDCl_3$) δ : ppm 4.08 (t, 4H); 3.62 (m, 2H); 2.68-2.56 (br. m., 6H); 2.33 (t, 4H); 1.64-1.54 (m, 13H); 1.35 (m, 12H); 0.93 (t, 6H).

5

Step 2: Dipentyl 6,6'-(2-chloroethyl)azanediyldihexanoate




Chemical Formula: $C_{24}H_{46}ClNO_4$

Molecular Weight: 448.09

10 [00616] In the same manner as Step 2 for Compound 18, dipentyl 6,6'-(2-chloroethyl)azanediyldihexanoate was synthesized from dipentyl 6,6'-(2-hydroxyethyl)azanediyldihexanoate (124 mg, 0.29 mmol), methanesulfonyl chloride (0.025 mL, 0.32 mmol) and triethylamine (0.060 mL, 0.44 mmol) in 1.5 mL DCM to provide 84 mg (65%).

15 1H NMR (400 MHz, $CDCl_3$) δ : ppm 4.04 (t, 4H); 3.46 (t, 2H); 2.73 (t, 2H); 2.43 (t, 4H); 2.28 (t, 4H); 1.60 (m, 8H); 1.40 (m, 4H); 1.29 (m, 12H); 0.89 (t, 6H).

Step 3: Compound 14: Dipentyl 6,6'-(2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)azanediyldihexanoate



20 Chemical Formula: $C_{68}H_{137}N_5O_4$


Molecular Weight: 1088.88

[00617] In the same manner as Step 6 for Compound 18, dipentyl 6,6'-(2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)azanediyldihexanoate was synthesized from N^1,N^1,N^2 -tridodecyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (105 mg,

0.16 mmol), dipentyl 6,6'-(2-chloroethyl)azanediyl)dihexanoate (84 mg, 0.19 mmol) and K₂CO₃ (22 mg, 0.16 mmol) in 1 mL MeCN. Yield (53 mg, 0.049 mmol, 30%).
 UPLC: RT = 3.47 min. MS (ES): *m/z* (MH⁺) 1089.53 for C₆₈H₁₃₇N₅O₄
¹H NMR (400 MHz, CDCl₃) δ: ppm 4.04 (t, 4H); 2.89-2.98 (m, 4H); 2.39-2.68 (m, 26H); 2.27
 5 (t, 4H); 1.57-1.71 (m, 10H); 1.35 (m, 4H); 1.28-1.35 (m, 74H); 0.90 (m, 15H).

P. Compound 15: Methyl 12-((2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)dodecanoate
 Step 1: Methyl 12-bromododecanoate

10




Chemical Formula: C₁₃H₂₅BrO₂

Molecular Weight: 293.25

[00618] To a solution of 12-bromododecanoic acid (2.5 g, 8.95 mmol) in THF (7 mL) was added methanol (7.2 mL, 179 mmol). Sulfuric acid (0.50 mL, 8.95 mmol) was added dropwise and the reaction was allowed to stir at 65 °C for two hours. The reaction mixture was washed with 5% NaHCO₃ and brine. The organic layer was dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% EtOAc/hexanes) provided methyl 12-bromododecanoate (2.40 g, 92%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.44 (t, 2H); 2.33 (t, 2H); 1.88 (br. m, 2H); 1.64 (br. m, 2H); 1.45 (br. m, 2H); 1.31 (br. m, 12H).

Step 2: Methyl 12-(dodecyl(2-hydroxyethyl)amino)dodecanoate



Chemical Formula: C₂₇H₅₅NO₃

25


Molecular Weight: 441.74

[00619] To a solution of methyl 12-((2-hydroxyethyl)amino)dodecanoate (413 mg, 1.51 mmol) in MeCN (5 mL) was added 1-bromododecane (452 mg, 1.81 mmol), K₂CO₃ (418 mg, 3.02 mmol), and KI (25 mg, 0.151 mmol). The reaction was allowed to stir at 82 °C for 16 hours. The reaction mixture was cooled to room temperature, diluted with H₂O, and extracted

with EtOAc. The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% [DCM, 20% MeOH, 1% NH₄OH]/MeOH) provided methyl 12-(dodecyl(2-hydroxyethyl)amino)dodecanoate (409 mg, 61%).

5 UPLC/ELSD: RT = 2.39 min. MS (ES): *m/z* (MH⁺) 442.60 for C₂₇H₅₅NO₃
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.61 (t, 2H); 2.68 (t, 2H); 2.54 (t, 4H); 2.32 (t, 2H); 1.64 (m, 2H); 1.50 (br. m, 4H); 1.28 (br. m, 32H); 0.90 (t, 3H).

Step 3: Methyl 12-((2-chloroethyl)(dodecyl)amino)dodecanoate



10


Chemical Formula: C₂₇H₅₄ClNO₂

Molecular Weight: 460.18

[00620] In the same manner as Step 2 for Compound 18, methyl 12-((2-chloroethyl)(dodecyl)amino)dodecanoate was synthesized from methyl 12-((2-hydroxyethyl)amino)dodecanoate (409 mg, 0.926 mmol), triethylamine (0.168 mL, 1.20 mmol), and methanesulfonyl chloride (0.090 mL, 1.16 mmol) in DCM (5 mL). Yield (307 mg, 72%).
 15 UPLC/ELSD: RT = 4.30 min. MS (ES): *m/z* (MH⁺) 460.80 for C₂₇H₅₄ClNO₂
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.59 (s, 3H); 3.42 (br. m, 2H); 2.70 (br. m, 2H); 2.38 (br. m, 4H); 2.30 (t, 2H); 1.55 (m, 2H); 1.36 (br. m, 4H); 1.27-0.96 (br. m, 32H); 0.81 (t, 3H).

20

Step 4: Compound 15: Methyl 12-((2-(4-(2-((2-didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)dodecanoate



Chemical Formula: C₇₁H₁₄₅N₅O₂

Molecular Weight: 1100.97

25


[00621] In the same manner as Step 6 for Compound 18, methyl 12-((2-(4-((2-(didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)(dodecyl)amino)dodecanoate was synthesized from *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (150 mg, 0.221 mmol), methyl 12-((2-chloroethyl)(dodecyl)amino)dodecanoate (134 mg, 0.266 mmol) K₂CO₃ (61 mg, 0.443 mmol), and KI (4 mg, 0.024 mmol) in THF (5 mL). Yield (32 mg, 15%).

5 UPLC/ELSD: RT = 4.83 min. MS (ES): *m/z* (MH⁺) 1102.11 for C₇₁H₁₄₅N₅O₂
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 2.75-2.24 (br. m, 32H); 1.64 (m, 2H); 1.52-1.00 (br. m, 96H); 0.90 (t, 12H).

10

Q. Compound 16: Dimethyl 12,12'-((2-(4-((2-(didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)azanediyil)didecanoate

Step 1: Dimethyl 12,12'-((2-hydroxyethyl)azanediyil)didecanoate



15

Chemical Formula: C₂₈H₅₅NO₅


Molecular Weight: 485.75

[00622] In the same manner as Step 1 for Compound 18, dimethyl 12,12'-((2-hydroxyethyl)azanediyil)didecanoate was synthesized from methyl 12-bromododecanoate (1.5 g, 5.12 mmol), ethanolamine (0.310 mL, 5.12 mmol), K₂CO₃ (1.42 g, 10.2 mmol), and KI (85 mg, 0.512 mmol) in MeCN (11 mL). Yield (563 mg, 45%).

20 UPLC/ELSD: RT = 1.81 min. MS (ES): *m/z* (MH⁺) 486.63 for C₂₈H₅₅NO₅
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 6H); 3.59 (br. m, 2H); 2.75-2.40 (br. m, 6H); 2.32 (t, 4H); 1.64 (m, 4H); 1.48 (br. m, 4H); 1.29 (br. m, 28H).

25

Step 2: Dimethyl 12,12'-((2-chloroethyl)azanediyil)didecanoate



Chemical Formula: C₂₈H₅₄ClNO₄


Molecular Weight: 504.19

[00623] In the same manner as Step 2 for Compound 18, dimethyl 12,12'-(2-chloroethyl)azanediyl)didodecanoate was synthesized from dimethyl 12,12'-(2-hydroxyethyl)azanediyl)didodecanoate (518 mg, 1.07 mmol), triethylamine (0.193 mL, 1.39 mmol), and methanesulfonyl chloride (0.103 mL, 1.33 mmol) in DCM (5.5mL). Yield (376 mg, 70%).

UPLC/ELSD: RT = 2.17 min. MS (ES): m/z (MH^+) 504.75 for $\text{C}_{28}\text{H}_{54}\text{ClNO}_4$

10

Step 3: Compound 16: Dimethyl 12,12'-(2-(4-(2-((2-(didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)azanediyil)didodecanoate



Chemical Formula: C₇₂H₁₄₅N₅O₄

Molecular Weight: 1144.98

15

[00624] In the same manner as Step 6 for Compound 18, dimethyl 12,12'-(2-(4-(2-(2-(didodecylamino)ethyl)(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)azanediyl)didodecanoate was synthesized from *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (150 mg, 0.221 mmol), dimethyl 12,12'-(2-chloroethyl)azanediyl)didodecanoate (134 mg, 0.266 mmol) 20 K_2CO_3 (61 mg, 0.443 mmol), and KI (4 mg, 0.024 mmol) in THF (5 mL). Yield (32 mg, 15%)


UPLC/ELSD: RT = 3.46 min, MS (ES): m/z (MH $^+$) 1146.07 for C₇₀H₁₄₅N₅O₄

¹H NMR (300 MHz, CDCl₃) δ: ppm 3.60 (s, 6H); 2.75, 2.24 (br, m, 3H); 1.64

¹H NMR (300 MHz, CDCl₃, δ, ppm) 5.89 (s, 9H), 2.75–2.24 (br, m, 9H), 1.84 (m, 1H), 1.52–1.00 (br, m, 92H); 0.90 (t, 9H).

R. Compound 17: *N*¹-(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-trihexylethane-1,2-diamine

Step 1: 2-(Dihexylamino)ethan-1-ol



5

Chemical Formula: C₁₄H₃₁NO


Molecular Weight: 229.41

[00625] In the same manner as Step 1 for Compound 18, 2-(dihexylamino)ethan-1-ol was synthesized from 1-bromohexane (5 g, 82 mmol), ethanolamine (11.5 mL, 82 mmol), K₂CO₃ (22.7 g, 164 mmol), and KI (1.36 g, 8.2 mmol) in MeCN (380mL). Yield (2.58g, 14%).

10 UPLC/ELSD: RT = 0.41 min. MS (ES): *m/z* (MH⁺) 229.95 for C₁₄H₃₁NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.62 (t, 2H); 2.70 (t, 2H), 2.57 (t, 4H); 1.50 (br. m, 4H); 1.30 (br, 12H); 0.91 (t, 6H).

Step 2: *N*-(2-Chloroethyl)-*N*-hexylhexan-1-amine



15

Chemical Formula: C₁₄H₃₀ClN

Molecular Weight: 247.85


[00626] In the same manner as Step 2 for Compound 18, *N*-(2-chloroethyl)-*N*-hexylhexan-1-amine was synthesized from 2-(dihexylamino)ethan-1-ol (2.50 g, 10.9 mmol), triethylamine (2.0 mL, 14.2 mmol), and methanesulfonyl chloride (1.0 mL, 13.6 mmol) in DCM (56 mL). Yield (1.93 g, 71%).

UPLC/ELSD: RT = 0.42 min. MS (ES): *m/z* (MH⁺) 247.86 for C₁₄H₃₀ClN

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.50 (t, 2H); 2.77 (t, 2H); 2.51 (t, 4H); 1.42 (br. m, 4H); 1.27 (br, 12H); 0.89 (t, 6H).

25

Step 3: *tert*-Butyl 4-(2-(hexylamino)ethyl)piperazine-1-carboxylate




Chemical Formula: C₁₇H₃₅N₃O₂

Molecular Weight: 313.49

[00627] In the same manner as Step 3 for Compound 18, *tert*-butyl 4-(2-(hexylamino)ethyl)piperazine-1-carboxylate was synthesized from 1-bromohexane (1.44 g, 8.72 mmol), 4-(2-aminoethyl)-1-boc-piperazine (2.0 g, 8.72 mmol), K₂CO₃ (2.4 g, 17.4 mmol), and KI (145 mg, 0.872 mmol). Yield (446 mg, 16%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.44 (br. m, 4H); 2.75 (br. m, 2H); 2.65 (br. m, 2H); 2.54 (br. m, 2H); 2.42 (br. m, 4H); 1.60-1.43 (br. m, 11H); 1.40-1.05 (br. m, 6H); 0.91 (br. m, 3H).

Step 4: *tert*-Butyl 4-((2-(dihexylamino)ethyl)(hexylamino)ethyl)piperazine-1-carboxylate



10

Chemical Formula: C₃₁H₆₄N₄O₂


Molecular Weight: 524.88

[00628] In the same manner as Step 4 for Compound 18, *tert*-butyl 4-((2-(dihexylamino)ethyl)(hexylamino)ethyl)piperazine-1-carboxylate was synthesized from *tert*-butyl 4-(2-(hexylamino)ethyl)piperazine-1-carboxylate (250 mg, 0.797 mmol), N-(2-chloroethyl)-N-hexylhexan-1-amine (217 mg, 0.877 mmol), K₂CO₃ (220 mg, 1.59 mmol), and KI (13 mg, 0.0797 mmol) in THF (5 mL). Yield 308 mg, 74%).

UPLC/ELSD: RT = 1.40 min. MS (ES): *m/z* (MH⁺) 525.83 for C₃₁H₆₄N₄O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.45 (br. m, 4H); 3.15-2.15 (br. m, 18H); 1.85-1.00 (br. m, 33H); 0.91 (9H).

Step 5: N¹,N¹,N²-Trihexyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine



Chemical Formula: C₂₆H₅₆N₄


25

Molecular Weight: 424.76

[00629] In the same manner as Step 5 for Compound 18, N¹,N¹,N²-trihexyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine was synthesized from *tert*-butyl 4-((2-(dihexylamino)ethyl)(hexylamino)ethyl)piperazine-1-carboxylate (308 mg, 0.587 mmol), and TFA (2.25 ml, 29.3 mmol) in DCM (2.5 mL). Yield (220 mg, 88%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.92 (br. m, 4H); 2.70-2.20 (br. m, 18H), 1.54-1.22 (br. m, 24H); 0.91 (br. m, 9H).

Step 6: Compound 17: *N*¹-(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²-5 trihexylethane-1,2-diamine



Chemical Formula: C₅₂H₁₀₉N₅

Molecular Weight: 804.48

[00630] In the same manner as Step 6 for Compound 18, *N*¹-(2-(4-(2-


10 (didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²-trihexylethane-1,2-diamine was synthesized from *N*¹,*N*¹,*N*²-trihexyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (110 mg, 0.259 mmol), *N*-(2-chloroethyl)-*N*-dodecyldecane-1-amine (162 mg, 0.388 mmol), K₂CO₃ (72 mg, 0.518 mmol), and KI (5 mg, 0.0259 mmol) in THF (6 mL). Yield (81 mg, 39%).

UPLC/ELSD: RT = 2.79 min. MS (ES): *m/z* (MH⁺) 806.30 for C₅₂H₁₀₉N₅

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.05-2.10 (br. m, 30H); 1.80-1.05 (br. m, 64H); 0.91 (br. m, 15H).

S: Compound 18: *N*¹-(2-(4-(2-(Dinonylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²-trinonylethane-1,2-diamine

20 Step 1: 2-(Dinonylamino)ethan-1-ol



Chemical Formula: C₂₀H₄₃NO

Molecular Weight: 313.57

[00631] To a solution of 1-bromononane (8.31 g, 40.1 mmol) in MeCN (84 mL) was


25 added ethanolamine (1.10 mL, 18.2 mmol), K₂CO₃ (11.1 g, 80.1 mmol), and KI (302 mg, 1.82 mmol). The reaction was allowed to stir at 82 °C for 48 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by

ISCO silica flash chromatography (0-20% MeOH/DCM) provided 2-(dinonylamino)ethan-1-ol (4.06 g, 71%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.57 (t, 2H); 2.63 (t, 2H); 2.49 (br. m, 4H); 1.48 (br. m, 4H); 1.29 (br. m, 24H); 0.91 (t, 6H).

5

Step 2: *N*-(2-Chloroethyl)-*N*-nonylnonan-1-amine



Chemical Formula: C₂₀H₄₂ClN


Molecular Weight: 332.01

10 [00632] To a 0 °C solution of 2-(dinonylamino)ethan-1-ol (4.06 g, 12.9 mmol) and triethylamine (2.35 ml, 16.8 mmol) in DCM (65 mL) was added dropwise a solution of methanesulfonyl chloride (1.25 mL, 16.18 mmol) in DCM (5 mL). The reaction was allowed to return to room temperature and stir for 16 hours. The mixture was quenched by the addition of water and extracted with DCM. The organic layer was washed with saturated NaHCO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-10% EtOAc/hexanes) provided *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (2.58 g, 60%).

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.51 (t, 2H); 2.78 (t, 2H); 2.47 (br. m, 4H); 1.44 (br. m, 4H); 1.28 (br. m, 24H); 0.90 (t, 6H).

20

Step 3: *tert*-Butyl 4-(2-(nonylamino)ethyl)piperazine-1-carboxylate



Chemical Formula: C₂₀H₄₁N₃O₂

Molecular Weight: 355.57

25 [00633] To a solution of 1-bromononane (1.81 g, 8.72 mmol) in MeCN (44 mL) was added 4-(2-aminoethyl)-1-boc-piperazine (2.0 g, 8.72 mmol), K₂CO₃ (2.4 g, 17.4 mmol), and KI (145 mg, 0.872 mmol). The reaction was allowed to stir at 65 °C for 16 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*.


Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided *tert*-butyl 4-(2-(nonylamino)ethyl)piperazine-1-carboxylate (775 mg, 25%).

UPLC/ELSD: RT = 0.47 min. MS (ES): *m/z* (MH⁺) 356.41 for C₂₀H₄₁N₃O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.44 (br. m, 4H); 2.74 (t, 2H); 2.63 (t, 2H); 2.53 (t, 2H);

5 2.41 (br. m, 4H); 1.48 (br. m, 9H); 1.30 (br. m, 14H); 0.90 (t, 3H).

Step 4: *tert*-Butyl 4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperazine-1-carboxylate



Chemical Formula: C₄₀H₈₂N₄O₂


10 Molecular Weight: 651.12

[00634] To a solution of *tert*-butyl 4-(2-(nonylamino)ethyl)piperazine-1-carboxylate (500 mg, 1.41 mmol) in THF (9 mL) was added *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (514 mg, 1.55 mmol), K₂CO₃ (390 mg, 2.82 mmol), and KI (23 mg, 0.141 mmol). The reaction was allowed to stir at 65 °C for 72 hours. The reaction mixture was cooled to room temperature, diluted with water, and extracted with EtOAc. The combined extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-15% MeOH/DCM) provided *tert*-butyl 4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperazine-1-carboxylate (763 mg, 83%).

UPLC/ELSD: RT = 2.61 min. MS (ES): *m/z* (MH⁺) 651.91 for C₄₀H₈₂N₄O₂

20 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.45 (br. m, 4H); 2.75-2.30 (br. m, 18H); 1.55-1.20 (br. m, 51H); 0.91 (br. m, 9H).

Step 5: N¹,N¹,N²-Trinonyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine




25 Chemical Formula: C₃₅H₇₄N₄

Molecular Weight: 551.01

[00635] To a 0 °C solution of *tert*-Butyl 4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperazine-1-carboxylate (763 mg, 1.17 mmol) in DCM (4.5 mL) was added dropwise TFA (4.5 mL, 58.5 mmol). The reaction was allowed to

return to room temperature and stir for 16 hours. The reaction mixture was concentrated *in vacuo* and the crude material was dissolved in CHCl₃. The solution was washed with 5% Na₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% [DCM, 20% MeOH, 1% NH₄OH]/MeOH) 5 provided N¹,N¹,N²-trinonyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (218 mg, 34%). UPLC/ELSD: RT = 1.81 min. MS (ES): *m/z* (MH⁺) 551.78 for C₃₅H₇₄N₄ ¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.91 (br. m, 4H); 2.70-2.35 (br. m, 18H); 1.46 (br. m, 6H); 1.29 (br. m, 36H); 0.91 (br. m, 9H)

10 Step 6: Compound 18: N¹-(2-(4-(2-(Dinonylamino)ethyl)piperazin-1-yl)ethyl)-N¹,N²,N²-trinonylethane-1,2-diamine



Chemical Formula: C₅₅H₁₁₅N₅


Molecular Weight: 846.56

15 [00636] To a solution of N¹,N¹,N²-trinonyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (74 mg, 0.134 mmol) and N-(2-chloroethyl)-N-nonylnonan-1-amine (58 mg, 0.175 mmol) in THF (4 mL) was added K₂CO₃ (37 mg, 0.269 mmol), and KI (3 mg, 0.0134 mmol). The reaction allowed to stir at 65 °C for 48 hours. The reaction mixture was cooled to room temperature, diluted with water, and extracted with EtOAc. The combined extracts were washed 20 with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO C18 flash chromatography (50-100% [MeCN 0.1% TFA]/[H₂O 0.1% TFA]) afforded the desired product as a TFA salt. The salt was dissolved in CHCl₃ and the solution was washed with 5% Na₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo* to provide N¹-(2-(4-(2-(dinonylamino)ethyl)piperazin-1-yl)ethyl)-N¹,N²,N²-trinonylethane-1,2-diamine (66 mg, 58%).

UPLC/ELSD: RT = 2.91 min. MS (ES): *m/z* (MH⁺) 847.30 for C₅₅H₁₁₅N₅

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.10-2.25 (br. m, 30H); 1.90-1.35 (br. m, 10H); 1.29 (br. m, 60H); 0.91 (br. m, 15H).

T: Compound 19: N^1 -(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -trinonylthane-1,2-diamine




Chemical Formula: $C_{61}H_{127}N_5$

Molecular Weight: 930.72

5 [00637] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -trinonylthane-1,2-diamine was synthesized from N^1,N^1,N^2 -trinonyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (70 mg, 0.127 mmol), N -(2-chloroethyl)- N -dodecyldecane-1-amine (79 mg, 0.191 mmol), K_2CO_3 (35 mg, 0.254 mmol), and KI (2 mg, 0.0127 mmol) in THF (3 mL). Yield (52 mg, 44%).
10 UPLC/ELSD: RT = 3.35 min. MS (ES): m/z (MH^+) 931.61 for $C_{61}H_{127}N_5$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 2.70 (br. m, 30H); 1.56-1.02 (br. m, 82H); 0.90 (t, 15H).

15 U: Compound 20: N^1 -(2-(4-(2-(Ditetradecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -trinonylthane-1,2-diamine




Chemical Formula: $C_{65}H_{135}N_5$

Molecular Weight: 986.83

20 [00638] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(ditetradecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -trinonylthane-1,2-diamine was synthesized from N^1,N^1,N^2 -trinonyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (74 mg, 0.134 mmol), N -(2-chloroethyl)- N -tetradecyltetradecan-1-amine (95 mg, 0.201 mmol), K_2CO_3 (37 mg, 0.269 mmol), and KI (3 mg, 0.0134 mmol) in THF (2 mL). Yield (50 mg, 38%).
UPLC/ELSD: RT = 3.55 min. MS (ES): m/z (MH^+) 987.87 for $C_{65}H_{135}N_5$
25 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.20-2.25 (br. m, 30H); 1.85-1.00 (br. m, 90H); 0.91 (t, 15H).

V: Compound 21: N^1 -(2-(4-(2-(Dihexylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tridodecylethane-1,2-diamine



Chemical Formula: $C_{58}H_{121}N_5$

5

Molecular Weight: 888.64


[00639] In the same manner Step 6 for Compound 18, N^1 -(2-(4-(2-(dihexylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tridodecylethane-1,2-diamine was synthesized from N^1,N^1,N^2 -tridodecyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (67 mg, 0.099 mmol), *N*-(2-chloroethyl)-*N*-hexylhexan-1-amine (32 mg, 0.129 mmol), K_2CO_3 (28 mg, 0.198 mmol), and KI (2 mg, 0.0099 mmol) in THF (2 mL). Yield (30 mg, 34%).

UPLC/ELSD: RT = 3.24 min. MS (ES): m/z (MH^+) 890.58 for $C_{58}H_{121}N_5$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.15-2.20 (br. m, 30H); 1.85-1.00 (br. m, 76H); 0.91 (br. m, 15H).

15 W: Compound 22: N^1 -(2-(4-(2-(Dioctylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tridodecylethane-1,2-diamine

Step 1: 2-(Dioctylamino)ethan-1-ol



Chemical Formula: $C_{18}H_{39}NO$

20


Molecular Weight: 285.52

[00640] In the same manner as Step 1 for Compound 18, compound was synthesized from ethanolamine (5 g, 82 mmol), 1-bromoocetane (14 mL, 82 mmol), and K_2CO_3 (11 g, 82 mmol) in 200 mL MeCN. Yield (3.13 g, 11 mmol, 13%).

UPLC/ELSD: RT = 1.78 min. MS (ES): m/z (MH^+) 286.22 for $C_{18}H_{39}NO$

25 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.54 (m, 2H); 2.60-2.47 (m, 6H); 1.44 (m, 4H); 1.26 (m, 21H); 0.86 (t, 6H).

Step 2: *N*-(2-Chloroethyl)-*N*-octyloctan-1-amine



Chemical Formula: C₁₈H₃₈ClN


Molecular Weight: 303.96

[00641] In the same manner as Step 2 for Compound 18, *N*-(2-chloroethyl)-*N*-octyloctan-1-amine was synthesized from 2-(diethylamino)ethan-1-ol (3.13 g, 11 mmol), methanesulfonyl chloride (0.85 mL, 11 mmol) and Et₃N (1.5 mL, 11 mmol) in 30 mL DCM. Yield (1.55 g, 5.1 mmol, 46%).

UPLC/ELSD: RT = 3.86 min. MS (ES): *m/z* (MH⁺) 304.43 for C₁₈H₃₈ClN

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.48 (2, 2H); 2.75 (t, 2H); 2.43 (t, 4H); 1.40-1.25 (m, 24H); 10 0.86 (t, 6H).

Step 3: Compound 22: *N*¹-(2-(4-(2-(Diethylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine



15 Chemical Formula: C₆₂H₁₂₉N₅

Molecular Weight: 944.75


[00642] In the same manner as Step 6 for Compound 18, *N*¹-(2-(4-(2-(diethylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine was synthesized from *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (150 mg, 0.221 mmol), *N*-(2-chloroethyl)-*N*-octyloctan-1-amine 54 (81 mg, 0.266 mmol), K₂CO₃ (61 mg, 0.443 mmol), and KI (4 mg, 0.024 mmol) in THF (5 mL). Yield (200 mg, 96%).

UPLC/ELSD: RT = 3.41 min. MS (ES): *m/z* (MH⁺) 945.96 for C₆₂H₁₂₉N₅

¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.76-2.10 (br. m, 30H); 1.56-1.00 (br. m, 84H); 0.90 (t, 15H).

25

X: Compound 23: *N*¹-(2-(4-(2-(Dinonylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine



Chemical Formula: C₆₄H₁₃₃N₅

Molecular Weight: 972.80

[00643] In the same manner as Step 6 for Compound 18, N¹-(2-(4-(2-


5 (dinonylamino)ethyl)piperazin-1-yl)ethyl)-N¹,N²,N²-tridodecylethane-1,2-diamine was synthesized from N¹,N¹,N²-tridodecyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (595 mg, 0.879 mg), N-(2-chloroethyl)-N-nonylnonan-1-amine (350 mg, 1.05 mmol), K₂CO₃ (243 mg, 1.76 mmol), and KI (15 mg, 0.0879 mmol) in THF (13 mL). Yield (534 mg, 62%).

UPLC/ELSD: RT = 3.50 min. MS (ES): *m/z* (MH⁺) 973.60 for C₆₄H₁₃₃N₅

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.70-2.30 (br. m, 30H); 1.56-1.37 (br. m, 10H); 1.28 (br. m, 78H); 0.90 (t, 15H).

Y: Compound 24: (Z)-N¹-(2-(4-(2-(Dodec-6-en-1-yl)dodecyl)amino)ethyl)piperazin-1-yl)ethyl)-N¹,N²,N²-tridodecylethane-1,2-diamine

15 Step 1: (6-Hydroxyhexyl)triphenylphosphonium bromide




Chemical Formula: C₂₄H₂₈BrOP

Molecular Weight: 443.36

20 [00644] 6-Bromo-1-hexanol (4.89 g, 27 mmol) and triphenylphosphine (7.87 g, 30 mmol) and 50 mL MeCN were combined in a round bottomed flask. The flask was fitted with a condenser and placed in a heating mantel and the reaction was allowed to stir at 82 °C for 48 h. After this time the reaction was allowed to cool to rt and the solution was cannulated into 200 mL Et₂O, producing a white precipitate. The solids were allowed to settle and the solvent was 25 decanted off. 20 mL DCM was added to dissolve the solids and then 100 mL Et₂O was slowly added to afford a white precipitate. The solvent was then removed *in vacuo* to afford clean (6-hydroxyhexyl)triphenylphosphonium bromide (9.4 g, 21.2 mmol, for 78% yield).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 7.80 (m, 15H); 3.80 (m, 2H); 3.65 (m, 2H); 2.23 (m, 2H); 1.68 (m, 4H); 1.52 (m, 4H).

Step 2: (Z)-Dodec-6-en-1-ol

Chemical Formula: C₁₂H₂₄O

Molecular Weight: 184.32

5

[00645] A solution of (6-hydroxyhexyl)triphenylphosphonium bromide (3.0 g, 6.77 mmol) in 25 mL THF was allowed to cool in a -78 °C dry ice/acetone bath. Once cool *n*-BuLi (2.5 M in hexanes) (5.7 mL, 14.2 mmol) was added dropwise. After 1 h, an additional 10 mL THF and *n*-BuLi (1.35 mL) were added and stirring was continued at the same temperature for 10 h. After this time 1-hexanal (1.6 mL, 13.5 mmol) was added and the reaction was allowed to warm to rt and stir for 3 h. After this time the reaction was quenched by addition of excess saturated NH₄Cl. The solution was extracted three times with EtOAc. The pooled organics were washed with brine, dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was purified by silica gel chromatography (0-50% EtOAc in hexanes) to afford the 15 desired product as a clear oil (0.76 g, 4.1 mmol, 61%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 2H); 3.62 (t, 2H); 2.01 (m, 4H); 1.56 (m, 2H); 1.35-1.27 (m, 11H); 0.87 (t, 3H).

Step 3: (Z)-Dodec-6-en-1-yl methanesulfonate

20


Chemical Formula: C₁₃H₂₆O₃S

Molecular Weight: 262.41

[00646] To a 0 °C solution of (Z)-dodec-6-en-1-ol (1.81 g, 9.3 mmol) in 20 mL DCM, was added Et₃N (1.7 mL, 12.1 mmol) and methanesulfonyl chloride (0.80 mL, 10.2 mmol). The 25 reaction was allowed to slowly warm to rt and stir overnight. The reaction was quenched by the addition of water and the mixture was extracted two times with DCM. The organics were pooled, washed with brine, dried over MgSO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford clean desired product (2.2 g, 8.4 mmol, 90%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 2H); 4.20 (t, 2H); 2.98 (s, 3H); 2.01 (m, 4H); 1.74 (m, 2H); 1.38-1.27 (m, 10H); 0.87 (t, 3H).

Step 4: (Z)-1-Bromododec-6-ene


Chemical Formula: C₁₂H₂₃Br

Molecular Weight: 247.22

5 [00647] In a round bottomed flask, under N₂, (Z)-dodec-6-en-1-yl methanesulfonate (2.2 g, 8.3 mmol) was dissolved in 40 mL Et₂O. MgBr₂·Et₂O (6.5 g, 25 mmol) was added and the reaction was allowed to stir for 48 h. After this time the reaction was quenched by the addition of ice. The mixture was then extracted with Et₂O three times. The pooled organics were washed with brine, dried over MgSO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford the desired product (1.8 g, 7.28 mmol, 88%).

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 2H); 3.39 (t, 2H); 2.01-1.84 (m, 6H); 1.28 (m, 10H); 0.87 (t, 3H).

15 Step 5: (Z)-2-(Dodec-6-en-1-yl(dodecyl)amino)ethan-1-ol


Chemical Formula: C₂₆H₅₃NO

Molecular Weight: 395.72

20 [00648] In the same manner as Step 1 for Compound 18, (Z)-2-(Dodec-6-en-1-yl(dodecyl)amino)ethan-1-ol was synthesized from (Z)-1-bromododec-6-ene (0.25 g, 1.0 mmol), 2-(dodecylamino)ethan-1-ol (0.23 g, 1.0 mmol), K₂CO₃ (0.14 g, 1.0 mmol) and KI (2 mg, 0.01 mmol) in 5 mL MeCN.

25 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 2H); 3.65 (br. m., 2H); 2.64 (br. m, 6H); 2.00 (m, 4H); 1.55 (m, 6H); 1.24 (m, 26H); 0.86 (t, 6H).

Step 6: (Z)-N-(2-Chloroethyl)-N-dodecylidodec-6-en-1-amine


Chemical Formula: C₂₆H₅₂ClN

Molecular Weight: 414.16

[00649] In the same manner as Step 2 for Compound 18, (Z)-N-(2-chloroethyl)-N-dodecyl dodec-6-en-1-amine was synthesized from (Z)-2-(dodec-6-en-1-yl(dodecyl)amino)ethan-1-ol (35 mg, 0.088 mmol), methanesulfonyl chloride (0.008 mL, 0.097 mmol) and triethylamine (0.018 mL, 0.13 mmol) in 0.5 mL DCM. Yield (17.3 mg, 47%).

5 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.34 (m, 2H); 3.47 (t, 2H); 2.74 (t, 2H); 2.43 (t, 4H); 2.0 (m, 4H); 1.24 (m, 32H); 0.86 (t, 6H).

Step 7: Compound 24: (Z)- N^1 -(2-(4-(2-(Dodec-6-en-1-yl(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tridodecylethane-1,2-diamine



10

Chemical Formula: $\text{C}_{70}\text{H}_{143}\text{N}_5$

Molecular Weight: 1054.95


[00650] In the same manner as Step 6 for Compound 18, (Z)- N^1 -(2-(4-(2-(dodec-6-en-1-yl(dodecyl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tridodecylethane-1,2-diamine was synthesized from N^1,N^1,N^2 -tridodecyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (27 mg, 0.040 mmol) and (Z)-N-(2-chloroethyl)-N-dodecyl dodec-6-en-1-amine (17.3 mg, 0.042 mmol), K_2CO_3 (6 mg, 0.040 mmol) in 0.5 mL DCM. Yield (24 mg, 0.023 mmol, 57%).

UPLC: RT = 3.78 min. MS (ES): m/z (MH^+) 1056.376 for $\text{C}_{70}\text{H}_{143}\text{N}_5$

15 $^1\text{H NMR}$ (400 MHz, CDCl_3) δ : ppm 5.36 (m, 2H); 2.53-2.35 (m, 30H); 2.00 (m, 4H); 1.39-1.24 (m, 92H); 0.86 (m, 15H).

Z: Compound 25: N^1 -(2-(4-(2-(Di((Z)-dodec-6-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tridodecylethane-1,2-diamine

25 Step 1: 2-(Di((Z)-dodec-6-en-1-yl)amino)ethan-1-ol




Chemical Formula: $\text{C}_{26}\text{H}_{51}\text{NO}$

Molecular Weight: 393.70

[00651] In the same manner as Step 1 for Compound 18, 2-(di((Z)-dodec-6-en-1-yl)amino)ethan-1-ol was synthesized from ethanolamine (60 mg, 1.0 mmol), (Z)-1-bromododec-6-ene (0.51 g, 2.1 mmol) and K_2CO_3 (0.14 g, 1.0 mmol) in 5 mL DCM. Yield (0.22 g, 0.56 mmol, 56%).

5 1H NMR (400 MHz, $CDCl_3$) δ : ppm 5.34 (m, 4H); 3.59 (m, 2H); 2.65-2.53 (m, 6H); 2.00 (m, 9H); 1.49 (m, 4H); 1.23 (m, 20H); 0.86 (t, 6H).

Step 2: (Z)-N-(2-Chloroethyl)-N-((Z)-dodec-6-en-1-yl)dodec-6-en-1-amine




10 Chemical Formula: $C_{26}H_{50}ClN$

Molecular Weight: 412.14

[00652] In the same manner as Step 2 for Compound 18, (Z)-N-(2-chloroethyl)-N-((Z)-dodec-6-en-1-yl)dodec-6-en-1-amine was synthesized from 2-(di((Z)-dodec-6-en-1-yl)amino)ethan-1-ol (0.22 g, 0.56 mmol), methanesulfonyl chloride (0.047 mL, 0.61 mmol) and triethylamine (0.12 mL, 0.84 mmol) in 3 mL DCM. (Yield (150 mg, 0.36 mmol, 65%).

15 1H NMR (400 MHz, $CDCl_3$) δ : ppm 5.34 (m, 4H); 3.47 (t, 2H); 2.74 (t, 2H); 2.43 (t, 4H); 2.00 (m, 8H); 1.41-1.27 (m, 24H); 0.87 (m, 6H).

Step 3: Compound 25: N^1 -(2-(4-(2-(Di((Z)-dodec-6-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)-
20 N^1,N^2,N^2 -tridodecylethane-1,2-diamine



Chemical Formula: $C_{70}H_{141}N_5$

Molecular Weight: 1052.93


[00653] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(di((Z)-dodec-6-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tridodecylethane-1,2-diamine was synthesized from (Z)-N-(2-chloroethyl)-N-((Z)-dodec-6-en-1-yl)dodec-6-en-1-amine (56 mg, 0.14 mmol), N^1,N^1,N^2 -tridodecyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (84 mg, 0.12 mmol) and K_2CO_3 (17 mg, 0.12 mmol) in 1 mL MeCN. Yield (41.9 mg, 0.040 mmol, 33%).

UPLC: RT = 3.74 min. MS (ES): m/z (MH^+) 1053.564 for $\text{C}_{70}\text{H}_{141}\text{N}_5$

^1H NMR (400 MHz, CDCl_3) δ : ppm 5.33 (m, 4H); 2.55-2.35 (br. m, 30H); 1.98 (m, 8H); 1.32-1.24 (m, 84H); 0.86 (m, 15H).

[00654]

5 AA: Compound 26: N^1 -(2-(4-(2-(Ditetradecylamino)ethyl)piperazin-1-yl)ethyl)- $N^1,\text{N}^2,\text{N}^2$ -tridodecylethane-1,2-diamine



Chemical Formula: $\text{C}_{74}\text{H}_{153}\text{N}_5$

10 Molecular Weight: 1113.07


[00655] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(ditetradecylamino)ethyl)piperazin-1-yl)ethyl)- $N^1,\text{N}^2,\text{N}^2$ -tridodecylethane-1,2-diamine was synthesized from $N^1,\text{N}^1,\text{N}^2$ -tridodecyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (75 mg, 0.111 mmol), N -(2-chloroethyl)- N -tetradecyltetradecan-1-amine (58 mg, 0.122 mmol), K_2CO_3 (31 mg, 0.221 mmol), and KI (3 mg, 0.0181 mmol) in THF (4 mL). Yield (17 mg, 7%).

UPLC/ELSD: RT = 3.88 min. MS (ES): m/z (MH^+) 1113.59 for $\text{C}_{74}\text{H}_{153}\text{N}_5$

^1H -NMR (300 MHz, CDCl_3) δ : ppm 3.15-2.00 (br. m, 30H); 1.75-0.90 (br. m, 108H); 0.81 (t, 15H).

20 AB: Compound 27: $N^1,\text{N}^1,\text{N}^2$ -Tridodecyl- N^2 -(2-(4-(2-(dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazin-1-yl)ethyl)ethane-1,2-diamine

Step 1: (6Z,9Z)-18-(Methylsulfonyl)octadeca-6,9-diene



Chemical Formula: $\text{C}_{19}\text{H}_{36}\text{O}_3\text{S}$


25 Molecular Weight: 344.55

[00656] To a 0 °C solution of linoleyl alcohol (10 mL, 31.2 mmol) and triethylamine (5.68 mL, 40.5 mmol) in DCM (50 mL) was added dropwise a solution of methanesulfonyl chloride (2.66 mL, 34.3 mmol) in DCM (20 mL). The reaction was allowed to return to room temperature and let stir for 4 hours. The mixture was quenched by the addition of water and

extracted with DCM. The organic layer was washed with saturated NaHCO_3 , brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-40% EtOAc/hexanes) provided (*6Z,9Z*)-18-(methylsulfonyl)octadeca-6,9-diene (10.0 g, 93%).

5 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.35 (m, 4H); 4.22 (t, 2H); 2.99 (s, 3H); 2.77 (t, 2H); 2.04 (q, 4H); 1.74 (m, 2H); 1.30 (br. m, 16H); 0.89 (t, 3H).

Step 2: (*6Z,9Z*)-18-Bromoctadeca-6,9-diene




10 Chemical Formula: $\text{C}_{18}\text{H}_{33}\text{Br}$
Molecular Weight: 329.37

[00657] To a solution of (*6Z,9Z*)-18-(methylsulfonyl)octadeca-6,9-diene (10.0 g, 29.0 mmol) in diethyl ether (372 mL) was added magnesium bromide ethyl etherate (22.5 g, 87.1 mmol). The reaction was let stir at room temperature for 16 hours. The mixture was quenched 15 by the addition of water and extracted with diethyl ether. The combined organic layers were washed with 1% K_2CO_3 , brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography provided (*6Z,9Z*)-18-bromoctadeca-6,9-diene (8.9 g, 93%).

15 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.36 (m, 4H); 3.41 (t, 2H); 2.77 (t, 2H); 2.05 (q, 4H); 1.86 (m, 2H); 1.48-1.22 (br. m, 16H); 0.89 (t, 3H).

Step 3: Methyl *N*-dodecyl-*N*-((*9Z,12Z*)-octadeca-9,12-dien-1-yl)glycinate



Chemical Formula: $\text{C}_{33}\text{H}_{63}\text{NO}_2$

25 Molecular Weight: 505.87


[00658] To a solution of methyl dodecylglycinate·HCl (1 g, 3.4 mmol) in 8.5 mL DMF, (*Z*)-1-bromoctadec-9-ene (1.68 g, 5.1 mmol) and K_2CO_3 (1.4 g, 10.2 mmol) were added. The reaction was then allowed to stir at 85 °C for 12 h. After this time the reaction was allowed to cool to rt and was quenched by the addition of excess H_2O . The mixture was extracted 3 times 30 with EtOAc. The organics were pooled and washed with brine, dried over MgSO_4 , filtered and

concentrated. The crude material was purified by silica gel chromatography (0-10% EtOAc in hexanes) to afford the desired product (0.87 g, 1.72 mmol, 50%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.33 (m, 4H); 3.68 (s, 3H); 3.30 (s, 2H); 2.74 (t, 2H); 2.52 (m, 4H); 2.02 (m, 4H); 1.23 (m, 38H); 0.86 (m, 6H).

5

Step 4: 2-(Dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol



Chemical Formula: C₃₂H₆₃NO

Molecular Weight: 477.86


10 [00659] To a 0 °C solution of methyl N-dodecyl-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)glycinate (980 mg, 1.94 mmol) in THF (10 mL) was added dropwise lithium aluminum hydride (184 mg, 4.85 mmol). The reaction was allowed to return to room temperature and let stir for 3 hours. The mixture was slowly quenched by the stepwise addition of water (0.184 mL), 10% NaOH (0.552 mL) and water (0.184 mL). The reaction mixture was filtered, washed 15 with THF and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% [DCM, 20% MeOH, 1% NH₄OH]/MeOH) provided 2-(dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol (660 mg, 71%).

UPLC/ELSD: RT = 3.13 min. MS (ES): *m/z* (MH⁺) 478.52 for C₃₂H₆₃NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.39 (br. m, 4H); 3.56 (br. m, 2H); 2.80 (br. m, 2H); 2.61

20 (br. m, 2H); 2.48 (br. m, 4H); 2.09 (br. m, 4H); 1.57-1.17 (br. m, 38H); 0.91 (br. m, 6H).

Step 5: (9Z,12Z)-N-(2-Chloroethyl)-N-dodecyloctadeca-9,12-dien-1-amine



Chemical Formula: C₃₂H₆₂ClN


Molecular Weight: 496.31

25 [00660] In a same manner as Step 2 for Compound 18, (9Z,12Z)-N-(2-chloroethyl)-N-dodecyloctadeca-9,12-dien-1-amine was synthesized from 2-(dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol (660 mg, 1.38 mmol), triethylamine (0.249 mL, 1.80 mmol), and methanesulfonyl chloride (0.172 mL, 1.73 mmol) in DCM (7 mL). Yield (123 mg, 18%).

30 UPLC/ELSD: RT = 3.23 min. MS (ES): *m/z* (MH⁺) 496.72 for C₃₂H₆₂ClN

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.29 (br. m, 4H); 3.43 (br. m, 2H); 2.71 (br. m, 4H); 2.38 (br. m, 4H); 1.98 (br. m, 4H); 1.45-1.07 (br. m, 38H); 0.82 (br. m, 6H).

Step 6: Compound 27: *N*¹,*N*¹,*N*²-Tridodecyl-*N*²-(2-(4-(2-(dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazin-1-yl)ethyl)ethane-1,2-diamine



Molecular Weight: 1137.10


[00661] In the same manner as Step 6 for Compound 18, *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(4-(2-(dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazin-1-yl)ethyl)ethane-1,2-diamine was synthesized from *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (67 mg, 0.099 mmol), (9Z,12Z)-*N*-(2-chloroethyl)-*N*-dodecyl octadeca-9,12-dien-1-amine (64 mg, 0.129 mmol) K₂CO₃ (28 mg, 0.198 mmol), and KI (2 mg, 0.012 mmol) in THF (2 mL). Yield (48 mg, 43%).

UPLC/ELSD: RT = 3.90 min. MS (ES): *m/z* (MH⁺) 1137.95 for C₇₆H₁₅₃N₅

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.48-5.29 (m, 4H); 3.15-2.15 (br. m, 32H); 2.07 (br. m, 4H); 1.83-1.00 (br. m, 98H); 0.91 (br. m, 15H).

AC: Compound 28: *N*¹-(2-(4-(2-(Di((Z)-octadec-9-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine

Step 1: (Z)-1-(Methylsulfonyl)octadec-9-ene



Chemical Formula: C₁₉H₃₈O₃S


Molecular Weight: 346.57

[00662] In the same manner as Step 1 for Compound 27, (Z)-1-(methylsulfonyl)octadec-9-ene was synthesized from oleyl alcohol (10 mL, 31.7 mmol), triethylamine (5.74 mL, 41.2 mmol), and methanesulfonyl chloride (2.70 mL, 34.9 mmol) in DCM (50 mL). Yield (8.55 g, 78%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.33 (m, 2H); 4.20 (t, 2H); 2.98 (s, 3H); 2.00 (m, 4H); 1.73 (m, 2H); 1.44-1.16 (br. m, 22H); 0.87 (t, 3H).

Step 2: (Z)-1-Bromoocatadec-9-ene

5



Chemical Formula: C₁₈H₃₅Br


Molecular Weight: 331.38

[00663] In the same manner as Step 2 for Compound 27, (Z)-1-bromoocatadec-9-ene was synthesized from (Z)-1-(methylsulfonyl)octadec-9-ene (8.55 g, 24.7 mmol), and magnesium bromide ethyl etherate (19.1 g, 74.1 mmol) in diethyl ether (317 mL). Yield (7.42 g, 91%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.35 (m, 2H); 3.41 (t, 2H); 2.01 (m, 4H); 1.85 (m, 2H); 1.48-1.14 (br. m, 22H); 0.88 (t, 3H).

Step 3: 2-(Di((Z)-octadec-9-en-1-yl)amino)ethan-1-ol

15



Chemical Formula: C₃₈H₇₅NO

Molecular Weight: 562.02


[00664] In the same manner as Step 1 for Compound 18, 2-(di((Z)-octadec-9-en-1-yl)amino)ethan-1-ol was synthesized from (Z)-1-bromoocatadec-9-ene (5 g, 15.1 mmol), ethanolamine, (0.414 mL, 6.86 mmol), K₂CO₃ (4.17 g, 30.2 mmol), and KI (114 mg, 0.686 mmol) in MeCN (32 mL). Yield (3.2 g, 83%).

UPLC/ELSD: RT = 7.325 min. MS (ES): *m/z* (MH⁺) 562.60 for C₃₈H₇₅NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 4H); 3.53 (t, 2H); 2.58 (t, 2H); 2.45 (t, 4H); 2.01 (m, 8H); 1.44 (m, 4H); 1.38-1.18 (br. m, 44H); 0.88 (t, 6H).

25

Step 4: (Z)-N-(2-Chloroethyl)-N-((Z)-octadec-9-en-1-yl)octadec-9-en-1-amine




Chemical Formula: C₃₈H₇₄ClN

Molecular Weight: 580.47

[00665] In the same manner as Step 2 for Compound 18, *(Z*)-*N*-(2-chloroethyl)-*N*-((*Z*)-octadec-9-en-1-yl)octadec-9-en-1-amine was synthesized from 2-(di(*(Z*)-octadec-9-en-1-yl)amino)ethan-1-ol (1.64 g, 2.92 mmol) triethylamine (0.529 mL, 3.79 mmol), and methanesulfonyl chloride (0.282 mL, 3.65 mmol) in DCM (15 mL). Yield (1.47 g, 87%).

5 UPLC/ELSD: RT = 3.75 min. MS (ES): *m/z* (MH⁺) 580.64 for C₃₈H₇₄ClN
¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.35 (m, 4H); 3.48 (br. m, 2H); 2.77 (br. m, 2H); 2.45 (br. m, 4H); 2.02 (br. m, 8H); 1.62-1.05 (br. m, 48H); 0.89 (t, 6H).

Step 5: Compound 28: *N*¹-(2-(4-(2-(Di(*(Z*)-octadec-9-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)-
10 *N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine



Chemical Formula: C₈₂H₁₆₅N₅


Molecular Weight: 1221.26

[00666] In the same manner as Step 6 for Compound 18, *N*¹-(2-(4-(2-(di(*(Z*)-octadec-9-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine was synthesized from *N*¹,*N*¹,*N*²-tridodecyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (75 mg, 0.111 mmol), (*Z*)-*N*-(2-chloroethyl)-*N*-((*Z*)-octadec-9-en-1-yl)octadec-9-en-1-amine (71 mg, 0.122 mmol) K₂CO₃ (31 mg, 0.222 mmol), and KI (3 mg, 0.018 mmol) in THF (1.5 mL). Yield (20 mg, 15%).

20 UPLC/ELSD: RT = 4.05 min. MS (ES): *m/z* (MH⁺) 1221.72 for C₈₂H₁₆₅N₅
¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.29-5.14 (br. m, 4H); 2.95-2.00 (br. m, 30H); 1.96-1.77 (br. m, 8H); 1.60-0.85 (br. m, 108H); 0.76 (br. m, 15H).

AD: Compound 29: *N*¹-(2-(4-(2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²,*N*²-tridodecylethane-1,2-diamine
25

Step 1: 2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol




Chemical Formula: C₃₈H₇₁NO

Molecular Weight: 557.99

[00667] In the same manner as Step 1 for Compound 18, 2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol was synthesized from (6Z,9Z)-18-bromooctadeca-6,9-diene (4 g, 12.1 mmol), ethanolamine, (0.334 mL, 5.52 mmol), K_2CO_3 (3.36 g, 24.3 mmol), and KI (92 mg, 0.552 mmol) in MeCN (26 mL). Yield (1.9 g, 62%).

5 UPLC/ELSD: RT = 6.80 min. MS (ES): m/z (MH^+) 557.94 for $C_{38}H_{71}NO$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 5.35 (m, 8H); 3.52 (t, 2H); 2.77 (t, 4H); 2.57 (t, 2H); 2.43 (t, 4H); 2.04 (q, 8H); 1.48-1.18 (br. m, 36H); 0.89 (t, 6H).

Step 2: (9Z,12Z)-*N*-(2-Chloroethyl)-*N*-(9Z,12Z)-octadeca-9,12-dien-1-amine




Chemical Formula: $C_{38}H_{70}ClN$

Molecular Weight: 576.44

[00668] In a same manner as Step 2 for Compound 18, (9Z,12Z)-*N*-(2-chloroethyl)-*N*-(9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine was synthesized from 2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol (250 mg, 0.45 mmol), triethylamine (81 μ L, 0.58 mmol), and methanesulfonyl chloride (38 μ L, 0.49 mmol) in DCM (2 mL). Yield (134 mg, 52%).

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 5.36 (m, 8H); 3.49 (t, 2H); 2.78 (m, 6H); 2.45 (t, 4H); 2.05 (q, 8H); 1.48-1.18 (br. m, 36H); 0.89 (t, 6H).

Step 3: Compound 29: N^1 -(2-(4-(2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tridodecylethane-1,2-diamine



Chemical Formula: $C_{82}H_{161}N_5$

Molecular Weight: 1217.23


[00669] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tridodecylethane-1,2-diamine was synthesized from N^1,N^1,N^2 -tridodecyl- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-

diamine (75 mg, 0.111 mmol), (9Z,12Z)-*N*-(2-chloroethyl)-*N*-(9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine (71 mg, 0.122) K_2CO_3 (31 mg, 0.222 mmol), and KI (3 mg, 0.018 mmol) in THF (3 mL). Yield (20 mg, 15%)

UPLC/ELSD: RT = 3.97 min. MS (ES): m/z (MH^+) 1217.95 for $C_{82}H_{161}N_5$

5 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 5.48-5.28 (m, 12H); 3.30-2.20 (br. m, 36H); 2.17-1.92 (br. m, 12H); 1.90-1.00 (br. m, 94H); 0.87 (br. m, 15H).

AE: Compound 30: N^1 -(2-(4-(2-(Dinonylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine



10

Chemical Formula: $C_{70}H_{145}N_5$


Molecular Weight: 1056.97

[00670] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(dinonylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine was synthesized from N^1 -(2-(piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine (150 mg, 0.197 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (79 mg, 0.236 mmol), K_2CO_3 (54 mg, 0.394 mmol), and KI (3 mg, 0.0134 mmol) in THF (4 mL). Yield (50 mg, 24%).

UPLC/ELSD: RT = 3.79 min. MS (ES): m/z (MH^+) 1057.74 for $C_{70}H_{145}N_5$

15 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.15-2.20 (br. m, 30H); 1.90-1.00 (br. m, 100H); 0.90 (t, 20 15H).

AF: Compound 31: N^1 -(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine



25


Chemical Formula: $C_{76}H_{157}N_5$

Molecular Weight: 1141.13

[00671] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine was synthesized from N^1 -(2-(piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine (150 mg, 0.197 mmol), *N*-(2-chloroethyl)-*N*-dodecyldecan-1-amine (98 mg, 0.236 mmol), K_2CO_3 (54 mg, 0.394 mmol), and KI (3 mg, 0.0134 mmol) in THF (4 mL). Yield (42 mg, 19%).
UPLC/ELSD: RT = 3.98 min. MS (ES): *m/z* (MH^+) 1142.14 for $C_{76}H_{157}N_5$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.20-2.20 (br. m, 30H); 1.90-1.00 (br. m, 112H); 0.90 (t, 15H).

10

AG: Compound 32: N^1 -(2-(4-(2-(Ditetradeceylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine



Chemical Formula: $C_{80}H_{165}N_5$

15


Molecular Weight: 1197.24

[00672] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(ditetradeceylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine was synthesized from N^1 -(2-(piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tritetradeceylethane-1,2-diamine (150 mg, 0.197 mmol), *N*-(2-chloroethyl)-*N*-tetradecyltetradecan-1-amine (130 mg, 0.276 mmol), K_2CO_3 (54 mg, 0.394 mmol), and KI (3 mg, 0.0134 mmol) in THF (4 mL). Yield (17 mg, 7%).
UPLC/ELSD: RT = 4.11 min. MS (ES): *m/z* (MH^+) 1198.32 for $C_{80}H_{165}N_5$
 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.20-2.15 (br. m, 30H); 1.90-1.00 (br. m, 120H); 0.90 (t, 15H).

25

AH: Compound 33: N^1 -(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2,N^2 -tri((*Z*)-octadec-9-en-1-yl)ethane-1,2-diamine

Step 1: *tert*-Butyl (*Z*)-4-(2-(octadec-9-en-1-ylamino)ethyl)piperazine-1-carboxylate




Chemical Formula: C₂₉H₅₇N₃O₂

Molecular Weight: 479.79

[00673] In the same manner as Step 3 for Compound 18, *tert*-butyl (Z)-4-(2-(octadec-9-en-1-ylamino)ethyl)piperazine-1-carboxylate was synthesized from (Z)-1-bromoocatadec-9-ene (1.95 g, 11.8 mmol), 4-(2-aminoethyl)-1-boc-piperazine (1.35 g, 5.89 mmol), K₂CO₃ (1.60 g, 11.8 mmol), and KI (98 mg, 0.689 mmol) in MeCN (30 mL). Yield (790 mg, 28%).

5 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.32 (m, 2H); 3.45 (br, 4H); 2.75 (t, 2H); 2.65 (t, 2H); 2.57 (t, 2H); 2.40 (br, 4H); 2.09 (br, 4H); 1.48 (br, m, 11H); 1.41-1.10 (br, 22H); 0.89 (t, 3H).

10 Step 2: *tert*-Butyl 4-(2-((2-(di((Z)-octadec-9-en-1-yl)amino)ethyl)((Z)-octadec-9-en-1-yl)amino)ethyl)piperazine-1-carboxylate

Chemical Formula: C₆₇H₁₃₀N₄O₂


Molecular Weight: 1023.80

15 [00674] In the same manner as Step 4 for Compound 18, *tert*-butyl 4-(2-((2-(di((Z)-octadec-9-en-1-yl)amino)ethyl)((Z)-octadec-9-en-1-yl)amino)ethyl)piperazine-1-carboxylate was synthesized from *tert*-butyl (Z)-4-(2-(octadec-9-en-1-ylamino)ethyl)piperazine-1-carboxylate (573 mg, 1.19 mmol), (Z)-N-(2-chloroethyl)-N-((Z)-octadec-9-en-1-yl)octadec-9-en-1-amine (693 mg, 1.19 mmol), K₂CO₃ (329 mg, 2.38 mmol), and KI (20 mg, 0.119 mmol) in 20 THF (6 mL). Yield (918 mg, 75%).

1H-NMR (300 MHz, CDCl₃) δ: ppm 5.37 (m, 6H); 3.45 (br, m, 4H); 2.72-2.18 (br, m, 18H); 2.04 (br, m, 12H); 1.65-1.05 (br, m, 81H) 0.91 (br, m, 9H).

Step 3: N¹,N¹,N²-Tri((Z)-octadec-9-en-1-yl)-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine

25


Chemical Formula: C₆₂H₁₂₂N₄

Molecular Weight: 923.69

[00675] In the same manner as Step 5 for Compound 18, N^1,N^1,N^2 -tri((Z)-octadec-9-en-1-yl)- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine was synthesized from, *tert*-butyl 4-(2-((2-(di((Z)-octadec-9-en-1-yl)amino)ethyl)((Z)-octadec-9-en-1-yl)amino)ethyl)piperazine-1-carboxylate (740 mg, 0.859 mmol), and TFA (3.3 mL, 42.9 mmol) in DCM (3.3 mL). Yield 5 (115 mg, 14%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.36 (m, 6H); 2.91 (br, 4H); 2.74-2.34 (br. m, 18H); 2.03 (br. m, 12H); 1.54-1.04 (br. m, 72H); 0.90 (br. m, 9H).

Step 4: Compound 33: N^1 -(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tri((Z)-octadec-9-en-1-yl)ethane-1,2-diamine 10



Chemical Formula: $\text{C}_{88}\text{H}_{175}\text{N}_5$

Molecular Weight: 1303.40


[00676] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tri((Z)-octadec-9-en-1-yl)ethane-1,2-diamine was synthesized from N^1,N^1,N^2 -tri((Z)-octadec-9-en-1-yl)- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (58 mg, 0.063 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (31 mg, 0.075 mmol), K_2CO_3 (17 mg, 0.13 mmol), and KI (1 mg, 0.006 mmol) in THF 15 (1.5 mL). Yield (30 mg, 37%).

UPLC/ELSD: RT = 4.16 min. MS (ES): m/z (MH^+) 1304.03 for $\text{C}_{88}\text{H}_{175}\text{N}_5$

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.36-5.24 (br. m, 6H); 2.66-2.08 (br. m, 30H); 2.04-1.82 (br. m, 12H); 1.57-0.87 (br. m, 112H); 0.81 (br. m, 15H).

AI: Compound 34: N^1 -(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tri((9Z,12Z)-octadeca-9,12-dien-1-yl)ethane-1,2-diamine

Step 1: *tert*-Butyl 4-(2-((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate 25

Chemical Formula: C₂₉H₅₅N₃O₂

Molecular Weight: 477.78


[00677] In the same manner as Step 3 for Compound 18, *tert*-butyl 4-((9Z,12Z)-

5 octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate was synthesized from (6Z,9Z)-18-bromo-octadeca-6,9-diene (3.0 g, 9.11 mmol), 4-(2-aminoethyl)-1-boc-piperazine (2.09 g, 9.11 mmol), K₂CO₃ (2.52 g, 18.22 mmol), and KI (151 mg, 0.911 mmol) in MeCN (44 mL). Yield (1.20 g, 27%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.35 (m, 4H); 3.42 (t, 4H); 2.77 (t, 2H); 2.73 (t, 2H); 2.62

10 (t, 2H); 2.51 (t, 2H); 2.38 (t, 4H); 2.04 (q, 4H); 1.60-1.20 (br. m, 27H); 0.89 (t, 3H).

Step 2: *tert*-Butyl 4-(2-((2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate

Chemical Formula: C₆₇H₁₂₄N₄O₂

Molecular Weight: 1017.76


[00678] In the same manner as Step 4 for Compound 18, *tert*-butyl 4-((2-(di((9Z,12Z)-

15 octadeca-9,12-dien-1-yl)amino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-

20 yl)amino)ethyl)piperazine-1-carboxylate was synthesized from *tert*-butyl 4-((2-((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate (600 mg, 1.26 mmol), (9Z,12Z)-N-(2-chloroethyl)-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine (796 mg, 1.38 mmol), K₂CO₃ (347 mg, 2.51 mmol), and KI (21 mg, 0.126 mmol) in THF (8 mL). Yield (793 mg, 62%).

25 UPLC/ELSD: RT = 3.95 min. MS (ES): *m/z* (MH⁺) 1018.19 for C₆₇H₁₂₄N₄O₂

Step 3: N¹,N¹,N²-Tri((9Z,12Z)-octadeca-9,12-dien-1-yl)-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine

Chemical Formula: C₆₂H₁₁₆N₄

Molecular Weight: 917.64


[00679] In the same manner as Step 5 for Compound 18, *N*¹,*N*¹,*N*²-tri((9Z,12Z)-octadeca-

5 9,12-dien-1-yl)-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine was synthesized from *tert*-butyl 4-((2-((2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate (793 mg, 0.779 mmol), and TFA (3.0 mL, 39.0 mmol) in DCM (3.0 mL). Yield (374 mg, 52%).

UPLC/ELSD: RT = 3.68 min. MS (ES): *m/z* (MH⁺) 918.84 for C₆₂H₁₁₆N₄

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.36 (m, 12H); 4.12 (m, 6H); 3.30-2.55 (22H); 2.04 (q, 12H); 1.80-1.00 (br. m, 54H); 0.89 (t, 9H).

Step 4: *N*¹-(2-(4-(2-(Didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²-tri((9Z,12Z)-octadeca-9,12-dien-1-yl)ethane-1,2-diamine

Chemical Formula: C₈₈H₁₆₉N₅

Molecular Weight: 1297.36

[00680] In the same manner as Step 6 for Compound 18, *N*¹-(2-(4-(2-


(didodecylamino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²-tri((9Z,12Z)-octadeca-9,12-dien-1-yl)ethane-1,2-diamine was synthesized from *N*¹,*N*¹,*N*²-tri((9Z,12Z)-octadeca-9,12-dien-1-yl)-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (75 mg, 0.082 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (37 mg, 0.090) K₂CO₃ (23 mg, 0.163 mmol), and KI (2 mg, 0.012 mmol) in THF (3 mL). Yield (20 mg, 15%).

UPLC/ELSD: RT = 4.00 min. MS (ES): *m/z* (MH⁺) 1297.88 for C₈₈H₁₆₉N₅

25 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.48-5.28 (Br, 12H); 3.30-2.20 (br. m, 36H); 2.17-1.92 (br. m, 12H); 1.90-1.00 (br. m, 94H); 0.87 (br. m., 15H).

AJ: Compound 35: *N*¹,*N*¹,*N*²-Trihexyl-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine

Compound 35 was synthesized according to Steps 1-5 for Compound 17.




Chemical Formula: C₂₆H₅₆N₄

Molecular Weight: 424.76

5

AK: Compound 36: N¹,N¹,N²-Trinonyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine

Compound 36 was synthesized according to Steps 1-5 outlined for Compound 18.




Chemical Formula: C₃₅H₇₄N₄

Molecular Weight: 551.01

10

AL: Compound 37: N¹,N¹,N²-Tridodecyl-N²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine

Compound 37 was synthesized according to Steps 1 and 2 for Compound 3.



Chemical Formula: C₄₄H₉₂N₄


Molecular Weight: 677.25

15

AM: Compound 38: N¹-(2-(Piperazin-1-yl)ethyl)-N¹,N²,N²-tritetradeceylethane-1,2-diamine

20

[00681] Compound 38 was synthesized according to Steps 1-5 for Compound 13.




Chemical Formula: C₅₀H₁₀₄N₄

Molecular Weight: 761.41

25

AN: Compound 39: 2-(Didodecylamino)-N-dodecyl-N-(2-(piperazin-1-yl)ethyl)acetamide


[00682] Compound 39 was synthesized according to Steps 1-3 for Compound 1.

Chemical Formula: C₄₄H₉₀N₄O

Molecular Weight: 691.23

5 AO: Compound 40: *N*¹,*N*¹,*N*²-Tri((*Z*)-octadec-9-en-1-yl)-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine


[00683] Compound 40 was synthesized according to Steps 1-3 for Compound 33.

Chemical Formula: C₆₂H₁₂₂N₄

10 Molecular Weight: 923.69

AP: Compound 41: *N*¹,*N*¹,*N*²-Tri((9*Z*,12*Z*)-octadeca-9,12-dien-1-yl)-*N*²-(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine

[00684] Compound 41 was synthesized according to Steps 1-3 for Compound 34.




Chemical Formula:

Molecular Weight:

15

AQ: Compound according to formula (IV): *N*¹-(2-(4-(2-(Di((*Z*)-octadec-9-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)-*N*¹,*N*²-tri((*Z*)-octadec-9-en-1-yl)ethane-1,2-diamine

Chemical Formula: C₁₀₀H₁₉₅N₅

Molecular Weight: 1467.70

[00685] In the same manner as Step 6 for Compound 18, N^1 -(2-(4-(2-(di((Z)-octadec-9-en-1-yl)amino)ethyl)piperazin-1-yl)ethyl)- N^1,N^2 -tri((Z)-octadec-9-en-1-yl)ethane-1,2-diamine was synthesized from N^1,N^1,N^2 -tri((Z)-octadec-9-en-1-yl)- N^2 -(2-(piperazin-1-yl)ethyl)ethane-1,2-diamine (75 mg, 0.0812 mmol), (Z)- N -(2-chloroethyl)- N -((Z)-octadec-9-en-1-yl)octadec-9-en-1-amine (57 mg, 0.0974 mmol) K_2CO_3 (22 mg, 0.162 mmol), and KI (2 mg, 0.012 mmol) in THF (1.5 mL). Yield (30 mg, 25%)


5 UPLC/ELSD: RT = 4.41 min. MS (ES): m/z (MH^+) 1469.08 for $C_{100}H_{195}N_5$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 5.40-5.27 (br. m, 10H); 3.18-2.22 (br. m, 30H); 2.06-1.89 (m, 20H); 1.80-0.97 (br. m, 120H); 0.88 (t, 15H).

10

AR:Compound 42: 2-(Dinonylamino)-1-(5-(N -(2-(dinonylamino)ethyl)- N -nonylglycyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethan-1-one

Step 1: *tert*-Butyl 5-(dinonylglycyl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate



15

Chemical Formula: $C_{30}H_{57}N_3O_3$


Molecular Weight: 507.80

[00686] In the same manner as Step 3 for Compound 11, *tert*-butyl 5-(dinonylglycyl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate was synthesized from lithium dinonylglycinate (500 mg, 1.50 mmol), *tert*-butyl 2,5-diazabicyclo[2.2.1]heptane-2-carboxylate (357 mg, 1.80 mmol), iPr_2EtN (628 μ L, 3.60 mmol), and T3P (50% EtOAc solution, 2.68 mL, 4.50 mmol) in THF (15 mL). Yield (710 mg, 78%).

UPLC/ELSD: RT = 0.87 min. MS (ES): m/z (MH^+) 508.44 for $C_{30}H_{57}N_3O_3$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.98-4.46 (br. m, 2H); 4.30-3.15 (br. m, 10H); 2.14-1.60 (br. m, 6H); 1.49 (s, 9H); 1.40-1.00 (br. m, 24H); 0.89 (t, 6H).

Step 2: 1-(2,5-Diazabicyclo[2.2.1]heptan-2-yl)-2-(dinonylamino)ethan-1-one




Chemical Formula: C₂₅H₄₉N₃O

Molecular Weight: 407.69

[00687] In the same manner as Step 4 for Compound 11, 1-(2,5-diazabicyclo[2.2.1]heptan-2-yl)-2-(dinonylamino)ethan-1-one was synthesized from *tert*-butyl 5-(dinonylglycyl)-2,5-diazabicyclo[2.2.1]heptane-2-carboxylate (710 mg, 1.40 mmol) and TFA (5.4 mL, 70 mmol) in DCM (5 mL). Yield (446 mg, 78%)
 UPLC/ELSD: RT = 0.67 min. MS (ES): *m/z* (MH⁺) 408.64 for C₂₅H₄₉N₃O
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.90-3.00 (br. m, 8H); 2.49 (br. m, 4H); 1.79 (br. m, 2H); 1.58-1.08 (br. m, 28H); 0.90 (t, 6H).

10

Step 3: 2-(Dinonylamino)-1-(5-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethan-1-one



Chemical Formula: C₅₆H₁₁₁N₅O₂

Molecular Weight: 886.54

15


[00688] In the same manner as Step 11 for Compound 11, 2-(dinonylamino)-1-(5-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethan-1-one was synthesized from 1-(2,5-diazabicyclo[2.2.1]heptan-2-yl)-2-(dinonylamino)ethan-1-one (100 mg, 0.25 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (134 mg, 0.27 mmol), iPr₂EtN (94 μL, 0.54 mmol), and T3P (50% EtOAc solution, 438 μL, 0.74 mmol) in THF (20 mL). Yield (50 mg, 23%).

UPLC/ELSD: RT = 3.60 min. MS (ES): *m/z* (MH⁺) 887.12 for C₅₆H₁₁₁N₅O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.15-2.28 (br. m, 26H); 2.16-1.00 (br. m, 70H); 0.90 (t, 15H).

25

AS: Compound 43: 2-((2-(Didodecylamino)ethyl)(dodecyl)amino)-1-(5-(dinonylglycyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethan-1-one 3



Chemical Formula: $C_{65}H_{129}N_5O_2$

Molecular Weight: 1012.78

[00689] In the same manner as Step 11 for Compound 11, 2-((2-

5 (didodecylamino)ethyl)(dodecyl)amino)-1-(5-(dinonylglycyl)-2,5-diazabicyclo[2.2.1]heptan-2-yl)ethan-1-one was synthesized from 1-(2,5-diazabicyclo[2.2.1]heptan-2-yl)-2-(dinonylamino)ethan-1-one (100 mg, 0.25 mmol), N-(2-(didodecylamino)ethyl)-N-dodecylglycine (168 mg, 0.27 mmol), iPr₂EtN (94 μ L, 0.54 mmol), and T3P (50% EtOAc solution, 438 μ L, 0.74 mmol) in THF (20 mL). Yield (150 mg, 60%).


10 UPLC/ELSD: RT = 3.60 min. MS (ES): *m/z* (MH⁺) 1013.24 for $C_{65}H_{129}N_5O_2$

¹H-NMR (300 MHz, CDCl₃) δ : ppm 5.15-2.26 (br. m, 24H); 2.13-1.09 (br. m, 90H); 0.90 (t, 15H).

AT: Compound 44: Methyl 8-((2-(4-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-

15 2-oxoethyl)(nonyl)amino)octanoate

Step 1: *tert*-Butyl nonylglycinate



Chemical Formula: $C_{15}H_{31}NO_2$

Molecular Weight: 257.42


[00690] To a mixture of *tert*-butyl glycine (3.0 g, 23 mmol) and 1-bromononane (2.4 g,

11.5 mmol) in MeCN (100 mL) was added K₂CO₃ (3.2 g, 23 mmol) and KI (190 mg, 1.1 mmol) and the mixture was allowed to stir at 82 °C for 24 hours. The suspension was cooled to RT and filtered through a celite plug, rinsing with hexanes. The MeCN was extracted 3x with hexanes,

25 and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-15% MeOH/DCM) provided *tert*-butyl nonylglycinate as a clear colorless oil (848 mg, 29%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.31 (s, 2H); 2.60 (t, 2H); 1.82-1.63 (br, 1H); 1.56-1.20 (br. m, 23H); 0.90 (t, 3H).

Step 2: Methyl 8-bromooctanoate




Chemical Formula: C₉H₁₇BrO₂

Molecular Weight: 237.14

[00691] In the same manner as Step 1 for Compound 15, methyl 8-bromooctanoate was synthesized from 8-bromooctanoic acid (5.0 g, 22 mmol), methanol (20 mL, 450 mmol), and 10 H₂SO₄ (1.2 mL, 22 mmol) in THF (20mL). Yield (5.0 g, 95%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.42 (t, 2H); 2.33 (t, 2H); 1.88 (quint, 2H); 1.65 (quint, 2H); 1.54-1.27 (br. m, 6H).

Step 3: Methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)octanoate



Chemical Formula: C₂₄H₄₇NO₄


Molecular Weight: 413.64

[00692] To a mixture of *tert*-butyl nonylglycinate (300 mg, 1.17 mmol) and methyl 8-bromooctanoate (290 mg, 1.22 mmol) in MeCN (12mL) was added K₂CO₃ (341 mg, 2.45 mmol) and KI (19 mg, 0.12 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT and filtered through a celite plug, rinsing with hexanes. The MeCN was extracted 3x with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-10% EtOAc/Hexanes) provided methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)octanoate (353 mg, 73%).

25 UPLC/ELSD: RT = 1.60 min. MS (ES): *m/z* (MH⁺) 414.51 for C₂₄H₄₇NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.23 (s, 2H); 2.57 (t, 4H); 2.32 (t, 2H); 1.70-1.18 (br. m, 33H); 0.90 (t, 3H).

Step 4: *N*-(8-Methoxy-8-oxooctyl)-*N*-nonyl glycine

Chemical Formula: C₂₀H₃₉NO₄

Molecular Weight: 357.54


[00693] To a solution of methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)octanoate

5 (353 mg, 0.85 mmol) in DCM (4mL) was added TFA (3.3 mL, 43 mmol) and the solution was allowed to stir at RT for 4 hours. The solution was concentrated *in vacuo*, taken up in DCM, and washed with 5% Na₂CO₃ and brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide *N*-(8-methoxy-8-oxoethyl)-*N*-nonylglycine (305 mg, 99%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.49 (s, 2H); 3.06 (t, 4H); 2.32 (t, 2H); 1.79-

10 1.14 (br. m, 24H); 0.90 (t, 3H).

Step 5: *tert*-Butyl 4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazine-1-carboxylate

Chemical Formula: C₄₀H₈₀N₄O₃

Molecular Weight: 665.11

[00694] To a solution of *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (1.0 g, 2.0 mmol)

and 1-boc-piperazine (412 mg, 2.2 mmol) in THF (20 mL) was added iPr₂EtN (773 μL, 4.4 mmol) and T3P (50% EtOAc solution, 3.6 mL, 6.0 mmol) and the solution was allowed to stir at RT for 12 hours. The reaction was quenched with water and extracted 3x with EtOAc. The


20 combined organics were washed with brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided *tert*-butyl 4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazine-1-carboxylate (961 mg, 72%).

UPLC/ELSD: RT = 3.27 min. MS (ES): *m/z* (MH⁺) 665.79 for C₄₀H₈₀N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.43-2.90 (br. m, 20H); 2.04-0.99 (br. m, 51H); 0.90 (t,

25 9H).

Step 6: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one

Chemical Formula: C₃₅H₇₂N₄O

Molecular Weight: 564.99

[00695] To a solution of *tert*-butyl 4-(2-(dinonylamino)ethyl)-*N*-


5 nonylglycyl)piperazine-1-carboxylate (961 mg, 1.44 mmol) in DCM (6 mL) was added TFA (5.5 mL, 72 mmol) and the solution was allowed to stir for 4 hours. The solution was concentrated *in vacuo*, taken up in DCM, and washed with 5% Na₂CO₃, brine, dried over Na₂SO₄, filtered, and concentrated *in vacuo* to provide 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (743 mg, 91%).

10 UPLC/ELSD: RT = 2.14 min. MS (ES): *m/z* (MH⁺) 565.82 for C₃₅H₇₂N₄O

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.59 (br, 4H); 3.36 (br, 2H); 2.99-2.03 (br. m, 14H); 1.74-1.01 (br. m, 42H); 0.90 (t, 9H).

Step 7: Methyl 8-((2-(4-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-

15 oxoethyl)(nonyl)amino)octanoate

Chemical Formula: C₅₅H₁₀₉N₅O₄

Molecular Weight: 904.51

20 [00696] To a solution of *N*-(8-methoxy-8-oxooctyl)-*N*-nonylglycine (100 mg, 0.28 mmol) and 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (174 mg, 0.31

mmol) in THF (25 mL) was added, iPr₂EtN (107 μL, 0.62 mmol), and T3P (50% EtOAc

solution, 0.50 mL, 0.84 mmol) and the reaction was allowed to stir at room temperature

overnight. The reaction mixture was diluted with water and extracted with EtOAc. The

25 organics were dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by

ISCO silica flash chromatography (0-100% DCM/[DCM 20% MeOH 1% NH₄OH]) provided


methyl 8-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)octanoate (121 mg, 48%).

UPLC/ELSD: RT = 2.88 min. MS (ES): *m/z* (MH⁺) 905 for C₅₅H₁₀₉N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.64-3.36 (br. m, 11H); 3.18 (s, 2H); 3.13 (s, 2H); 2.54-5 2.09 (br. m, 16H); 1.55-0.88 (br. m, 66H); 0.76 (t, 12H).

AU: Compound 45: Pentyl 4-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)butanoate

Step 1: Pentyl 4-bromobutanoate



Chemical Formula: C₉H₁₇BrO₂


Molecular Weight: 237.14

[00697] In the same manner as Step 1 for Compound 15, pentyl 4-bromobutanoate was synthesized from 4-bromobutanoic acid (2.0 g, 12 mmol), pentanol (1.7 mL, 15.6 mmol), and

15 H₂SO₄ (0.65 mL, 12 mmol) in THF (20mL). Yield (1.26 g, 44%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.10 (t, 2H); 3.49 (t, 2H); 2.52 (t, 2H); 2.20 (quint, 2H); 1.65 (quint, 2H); 1.35 (m, 4H); 0.93 (t, 3H).

Step 2: Pentyl 4-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)butanoate




Chemical Formula: C₂₄H₄₇NO₄

Molecular Weight: 413.64

[00698] In the same manner as Step 3 for Compound 44, pentyl 4-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)butanoate was synthesized from *tert*-butyl nonylglycinate (300 mg, 1.17 mmol) and pentyl 4-bromobutanoate (290 mg, 1.22 mmol) in MeCN (12 mL) was added K₂CO₃ (341 mg, 2.45 mmol) and KI (19 mg, 0.12 mmol). Yield (343 mg, 71%).

UPLC/ELSD: RT = 1.81 min. MS (ES): *m/z* (MH⁺) 415 for C₂₄H₄₇NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.23 (s, 2H); 2.60 (br. m, 4H); 2.37 (t, 2H); 1.78 (m, 2H); 1.64 (m, 2H); 1.52-1.20 (br. m, 27H); 0.90 (m, 6H).

Step 3: *N*-Nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycineChemical Formula: C₂₀H₃₉NO₄


Molecular Weight: 357.54

5

[00699] In the same manner as Step 4 for Compound 44, *N*-nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycine was synthesized from pentyl 4-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)butanoate (343 mg, 0.83 mmol) and TFA (3.17 mL, 41.5 mmol) in DCM (4 mL). Yield (296 mg, 99%).

10 UPLC/ELSD: RT = 1.29 min. MS (ES): *m/z* (MH⁺) 358 for C₂₀H₃₉NO₄
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.43 (br, 2H); 2.94 (br, 4H); 2.41 (t, 2H); 1.98 (br, m, 2H); 1.74-1.54 (br, m, 4H); 1.40-1.16 (br, m, 16H); 0.91 (m, 6H).

15 Step 4: Pentyl 4-((2-(4-(*N*-(2-(dinonylamo)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)butanoate

Chemical Formula: C₅₅H₁₀₉N₅O₄


Molecular Weight: 904.51

20 **[00700]** In the same manner as Step 7 for Compound 44, pentyl 4-((2-(4-(*N*-(2-(dinonylamo)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)butanoate was synthesized from *N*-nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycine (100 mg, 0.28 mmol), 2-((2-(dinonylamo)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (174 mg, 0.31 mmol), iPr₂EtN (107 μL, 0.62 mmol), and T3P (50% EtOAc solution, 0.50 mL, 0.84 mmol) in THF (25 mL). Yield (121 mg, 48%).

25 UPLC/ELSD: RT = 3.01 min. MS (ES): *m/z* (MH⁺) 905 for C₅₅H₁₀₉N₅O₄
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.93 (t, 2H); 3.61-3.31 (br, m, 8H); 3.17 (m, 4H); 2.55-2.08 (br, m, 16H); 1.71-0.90 (br, m, 64H); 0.75 (m, 15H).

AV: Compound 46: Methyl 8-((2-((2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)octanoate

5 Step 1: 2-(Nonylamino)ethan-1-ol




Chemical Formula: C₁₁H₂₅NO

Molecular Weight: 187.33

[00701] To a mixture of ethanolamine (4.4 mL, 72 mmol) and 1-bromononane (3.0g, 14.5 mmol) in MeCN (150 mL) was added K₂CO₃ (4.0 g, 29 mmol) and KI (240 mg, 1.5 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT and filtered over a pad of celite, rinsing with hexanes. The MeCN was extracted with hexanes 3x, and the combined hexanes were concentrated. Purification by ISCO silica flash chromatography (0-100% DCM/[DCM 20% MeOH 1% NH₄OH]) provided 2-(nonylamino)ethan-1-ol (1.0 g, 15 38%)

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.66 (t, 2H); 2.80 (t, 2H); 2.62 (t, 2H); 1.96 (br. m, 2H); 1.50 (br. m, 2H); 1.28 (br. m, 12H); 0.90 (t, 3H).

Step 2: Methyl 8-((2-hydroxyethyl)(nonyl)amino)octanoate



Chemical Formula: C₂₀H₄₁NO₃


Molecular Weight: 343.55

[00702] In the same manner as Step 1 for Compound 18, methyl 8-((2-hydroxyethyl)(nonyl)amino)octanoate was synthesized from 2-(nonylamino)ethan-1-ol (500 mg, 2.67 mmol), methyl 8-bromooctanoate (665 mg, 2.8 mmol), K₂CO₃ (780 mg, 5.6 mmol), and KI (44 mg, 0.27 mmol) in MeCN (30 mL). Yield (578 mg, 63%).

UPLC/ELSD: RT = 1.01 min. MS (ES): *m/z* (MH⁺) 344.31 for C₂₀H₄₁NO₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.59 (t, 2H), 2.65 (br, 2H); 2.51 (t, 4H); 2.32 (t, 2H); 1.65 (br. m, 2H); 1.49 (br. m, 4H); 1.30 (br. m, 18H); 0.90 (t, 3H).


Step 3: Methyl 8-((2-chloroethyl)(nonyl)amino)octanoate

Chemical Formula: C₂₀H₄₀ClNO₂

Molecular Weight: 362.00

5 [00703] In the same manner as Step 2 for Compound 18, methyl 8-((2-chloroethyl)(nonyl)amino)octanoate was synthesized from methyl 8-((2-hydroxyethyl)(nonyl)amino)octanoate (578 mg, 1.68 mmol), methanesulfonyl chloride (163 μ L, 2.10 mmol) and trimethylamine (305 μ L, 2.20 mmol) in DCM (10 mL). Yield (418 mg, 69%). UPLC/ELSD: RT = 1.21 min. MS (ES): *m/z* (MH⁺) 363 for C₂₀H₄₀ClNO₂


10 ¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.69 (br, 3H); 3.51 (br, 2H), 2.78 (br. m, 2H); 2.47 (br. m, 4H); 2.33 (t, 2H); 1.72-1.20 (br. m, 24H); 0.91 (t, 3H).

Step 4: Methyl 8-((2-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)octanoateChemical Formula: C₃₅H₇₀N₂O₄

Molecular Weight: 582.96

15 [00704] To a mixture of *tert*-butyl nonylglycinate (218 mg, 0.85 mmol) and methyl 8-((2-chloroethyl)(nonyl)amino)octanoate (337 mg, 0.93 mmol) in MeCN (10 mL) was added K₂CO₃ (236 mg, 1.69 mmol) and KI (14 mg, 0.08 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT and filtered through a pad of celite, rinsing with hexanes. The mixture was extracted with hexanes 3x and the combined hexanes were concentrated. Purification by ISCO silica flash chromatography (0-10% MeOH/DCM) provided methyl 8-((2-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)octanoate (283 mg, 57%).

25 UPLC/ELSD: RT = 2.92 min. MS (ES): *m/z* (MH⁺) 584 for C₃₅H₇₀N₂O₄ ¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.69 (s, 3H); 3.28 (s, 2H); 2.80-2.20 (br. m, 12H); 1.85-1.10 (br. m, 47H); 0.91 (t, 6H).


Step 5: *N*-(2-((8-Methoxy-8-oxooctyl)(nonyl)amino)ethyl)-*N*-nonylglycineChemical Formula: C₃₁H₆₂N₂O₄

Molecular Weight: 526.85

5 [00705] In the same manner as Step 4 for Compound 44, *N*-(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)-*N*-nonylglycine was synthesized from methyl 8-((2-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)octanoate (283 mg, 0.49 mmol) and TFA (1.86 mL, 24.3 mmol) in DCM (2 mL). Yield (255 mg, 99%).
 UPLC/ELSD: RT = 2.18 min. MS (ES): *m/z* (MH⁺) 528 for C₃₁H₆₂N₂O₄

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H), 3.26 (s, 2H), 2.79 (br. m, 8H), 2.59 (t, 2H), 2.33 (t, 2H), 1.76-1.08 (br. m, 38H); 0.90 (t, 6H).

Step 6: Methyl 8-((2-((4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)octanoate



15


Chemical Formula: C₅₅H₁₀₉N₅O₄

Molecular Weight: 904.51

[00706] In the same manner as Step 11 for Compound 11, methyl 8-((2-((4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)octanoate was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (75 mg, 0.20 mmol), *N*-(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)-*N*-nonylglycine (150 mg, 0.28 mmol), iPr₂EtN (88 μL, 0.50 mmol) and T3P (50% EtOAc solution, 409 μL, 0.69 mmol). Yield (57 mg, 32%).
 UPLC/ELSD: RT = 2.88 min. MS (ES): *m/z* (MH⁺) 905 for C₅₅H₁₀₉N₅O₄
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.82-2.87 (br. m, 27H); 2.64 (m, 2H); 2.33 (t, 2H); 1.80-1.15 (br. m, 66H); 0.90 (t, 12H).

AW: Compound 47: Methyl 8-((2-(dinonylamino)ethyl)(2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)amino)octanoate

Step 1: Methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)amino)octanoate




Chemical Formula: C₁₅H₂₉NO₄

Molecular Weight: 287.40

[00707] In the same manner as Step 1 for Compound 44, methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)amino)octanoate was synthesized from *tert*-butyl glycine (2.0 g, 12 mmol), methyl 8-bromooctanoate (2.8 g, 12 mmol), K₂CO₃ (3.3 g, 24 mmol), and KI (198 mg, 1.2 mmol) in MeCN (100 mL). Yield (1.16 g, 34%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.33 (s, 2H); 2.62 (t, 2H); 2.32 (s, 2H); 2.16-1.80 (br, 1H); 1.72-1.42 (br. m, 13H); 1.34 (br. m, 6H).

15 Step 2: Methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(2-(dinonylamino)ethyl)amino)octanoate



Chemical Formula: C₃₅H₇₀N₂O₄


Molecular Weight: 582.96

[00708] In the same manner as Step 4 for Compound 46, methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(2-(dinonylamino)ethyl)amino)octanoate was synthesized from methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)amino)octanoate (300 mg, 1.0 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (381 mg, 1.15 mmol), K₂CO₃ (320 mg, 2.3 mmol), and KI (17 mg, 0.10 mmol) in MeCN (10 mL). Yield (285 mg, 47%).

UPLC/ELSD: RT = 2.89 min. MS (ES): *m/z* (MH⁺) 584 for C₃₅H₇₀N₂O₄

25 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.29 (s, 2H); 3.10 (br. m, 8H); 2.59 (t, 2H); 2.32 (t, 2H); 1.82 (br. m, 4H); 1.74-1.16 (br. m, 43H); 0.91 (t, 6H).

Step 3: *N*-(2-(dinonylamino)ethyl)-*N*-(8-methoxy-8-oxooctyl)glycine



Chemical Formula: C₃₁H₆₂N₂O₄

Molecular Weight: 526.85

[00709] In the same manner as Step 5 for Compound 46, *N*-(2-(dinonylamino)ethyl)-*N*-(8-


5 methoxy-8-oxooctyl)glycine was synthesized from methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(2-(dinonylamino)ethyl)amino)octanoate (285 mg, 0.50 mmol), and TFA (1.87 mL, 24.4 mmol) in DCM (2 mL). Yield (254 mg, 98%).

UPLC/ELSD: RT = 2.16 min. MS (ES): *m/z* (MH⁺) 528 for C₃₁H₆₂N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.25 (s, 2H); 2.90-2.72 (br. m, 8H); 2.59 (t,

10 2H); 2.32 (t, 2H); 1.66 (br. m, 6H); 1.48 (br. m, 2H); 1.40-1.20 (br. m, 30H); 0.91 (t, 6H).

Step 4: Methyl 8-((2-(dinonylamino)ethyl)(2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)amino)octanoate



Chemical Formula: C₅₅H₁₀₉N₅O₄

Molecular Weight: 904.51

[00710] In the same manner as Step 11 for Compound 11, methyl 8-((2-

(dinonylamino)ethyl)(2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)amino)octanoate was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (75 mg, 0.20 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-(8-methoxy-8-oxooctyl)glycine (150 mg, 0.28 mmol), iPr₂EtN (88 μL,


20 0.50 mmol) and T3P (50% EtOAc solution, 409 μL, 0.69 mmol). Yield (80 mg, 39%).

UPLC/ELSD: RT = 2.87 min. MS (ES): *m/z* (MH⁺) 905 for C₅₅H₁₀₉N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.81-2.21 (br. m, 31H); 1.89-1.05 (br. m, 66H); 0.90 (t, 12H).

25 AX: Compound 48: Methyl 8-((2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)amino)octanoate

Step 1: Methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)amino)octanoate



5

Chemical Formula: C₃₅H₆₈N₂O₆


Molecular Weight: 612.94

[00711] In the same manner as Step 4 for Compound 46, methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)amino)octanoate was synthesized from methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)amino)octanoate (65 mg, 0.23 mmol), methyl 8-((2-chloroethyl)(nonyl)amino)octanoate (86 mg, 0.24 mmol), K₂CO₃ (69 mg, 0.50 mmol), and KI (4 mg, 0.02 mmol) in MeCN (4 mL). Yield (60 mg, 43%).

UPLC/ELSD: RT = 2.42 min. MS (ES): *m/z* (MH⁺) 614 for C₃₅H₆₈N₂O₆
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.66 (s, 6H); 3.25 (s, 2H); 2.66 (m, 2H); 2.54 (m, 4H); 2.38 (m, 4H); 2.28 (t, 4H); 1.61 (m, 4H); 1.54-1.10 (br. m, 39H); 0.87 (t, 3H).

15

Step 2: *N*-(8-methoxy-8-oxooctyl)-*N*-(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)glycine




Chemical Formula: C₃₁H₆₀N₂O₆

Molecular Weight: 556.83

[00712] In the same manner as Step 5 for Compound 46, *N*-(8-methoxy-8-oxooctyl)-*N*-(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)glycine was synthesized from methyl 8-((2-(*tert*-butoxy)-2-oxoethyl)(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)amino)octanoate (60 mg, 0.10 mmol), and TFA (0.37 mL, 4.9 mmol) in DCM (1 mL). Yield (54 mg, 99%).
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.60 (s, 6H); 3.15 (s, 2H); 2.72 (br. m, 8H); 2.50 (t, 2H); 2.22 (t, 4H); 1.70-1.05 (br. m, 34H); 0.81 (t, 3H).

Step 3: Methyl 8-((2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)amino)octanoate



5

Chemical Formula: C₅₅H₁₀₇N₅O₆


Molecular Weight: 934.49

[00713] In the same manner as Step 11 for Compound 11, methyl 8-((2-(4-(dinonylglycyl)piperazin-1-yl)-2-oxoethyl)(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)amino)octanoate was synthesized from 2-(dinonylamino)-1-(piperazin-1-yl)ethan-1-one (26 mg, 0.07 mmol), N-(8-methoxy-8-oxooctyl)-N-(2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)glycine (54 mg, 0.10 mmol), iPr₂EtN (30 μL, 0.17 mmol) and T3P (50% EtOAc solution, 140 μL, 0.24 mmol). Yield (20 mg, 27%).
 UPLC/ELSD: RT = 2.56 min. MS (ES): *m/z* (MH⁺) 935 for C₅₅H₁₀₇N₅O₆
 15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.70-3.00 (br. m, 18H); 2.55-2.05 (br. m, 18H); 1.70-0.95 (br. m, 62H); 0.76 (t, 9H).

AY: Compound 49: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(4-(2-(dinonylamino)ethyl)piperidin-1-yl)ethan-1-one

20

Step 1: *tert*-Butyl 4-(2-(dinonylamino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₆₀N₂O₂


Molecular Weight: 480.82

25 [00714] To a mixture of *tert*-butyl 4-(2-aminoethyl)piperidine-1-carboxylate (1.50 g, 6.6 mmol) and 1-bromomononane (1.36 g, 6.57 mmol) in MeCN (100 mL) was added K₂CO₃ (1.83 g, 13.1 mmol) and KI (109 mg, 0.66 mmol) and the mixture was allowed to stir at 82 °C for 12

hours. The suspension was cooled to RT, filtered over a pad of celite rinsing with hexanes, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% DCM/[DCM 20% MeOH 1% NH₄OH]) provided *tert*-butyl 4-(2-(dinonylamino)ethyl)piperidine-1-carboxylate (602 mg, 19%).

5 UPLC/ELSD: RT = 2.41 min. MS (ES): *m/z* (MH⁺) 482 for C₃₀H₆₀N₂O₂
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.20-2.26 (br. m, 10H); 1.77-1.10 (br. m, 44H); 0.91 (t, 6H).

Step 2: *N*-Nonyl-*N*-(2-(piperidin-4-yl)ethyl)nonan-1-amine



10

Chemical Formula: C₂₅H₅₂N₂


Molecular Weight: 380.71

[00715] In the same manner as Step 4 for Compound 11, *N*-nonyl-*N*-(2-(piperidin-4-yl)ethyl)nonan-1-amine was synthesized from *tert*-butyl 4-(2-(dinonylamino)ethyl)piperidine-1-carboxylate (602 mg, 1.25 mmol) and TFA (4.8 mL, 63 mmol) in DCM (5 mL). Yield (406 mg, 85%).

UPLC/ELSD: RT = 1.27 min. MS (ES): *m/z* (MH⁺) 382 for C₂₅H₅₂N₂
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.15 (br. m, 2H); 2.65 (br. m, 2H); 2.42 (br. m, 6H); 1.83-1.04 (br. m, 35H); 0.90 (t, 6H).

20

Step 3: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(4-(2-(dinonylamino)ethyl)piperidin-1-yl)ethan-1-one



25

Chemical Formula: C₅₆H₁₁₄N₄O

Molecular Weight: 859.56

[00716] In the same manner as Step 11 for Compound 11, 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(4-(2-(dinonylamino)ethyl)piperidin-1-yl)ethan-1-one


was synthesized from *N*-nonyl-*N*-(2-(piperidin-4-yl)ethyl)nonan-1-amine (200 mg, 0.53 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (287 mg, 0.58 mmol), iPr₂EtN (201 μ L, 1.2 mmol), and T3P (50% EtOAc solution, 938 μ L, 1.6 mmol) in THF (10 mL). Yield (90 mg, 20%).

UPLC/ELSD: RT = 3.26 min. MS (ES): m/z (MH $^+$) 860 for C₅₆H₁₁₄N₄O

5 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 4.62-4.09 (br. m, 2H); 3.55-2.21 (br. m, 20H); 1.94-1.00 (br. m, 77H); 0.91 (t, 15H).

AZ: Compound 50: 2-(Dinonylamino)-1-(4-(2-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperidin-1-yl)ethan-1-one

10 Step 1: *tert*-Butyl 4-(2-(nonylamino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₂₁H₄₂N₂O₂


Molecular Weight: 354.58

[00717] In the same manner as Step 1 for Compound 49, *tert*-butyl 4-(2-(nonylamino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(2-aminoethyl)piperidine-1-carboxylate (1.50 g, 6.6 mmol), 1-bromononane (1.36 g, 6.57 mmol), K₂CO₃ (1.83 g, 13.1 mmol), and KI (109 mg, 0.66 mmol) in MeCN (100 mL). Yield (288 mg, 13%).

UPLC/ELSD: RT = 1.23 min. MS (ES): m/z (MH $^+$) 356 for C₂₁H₄₂N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (br. m, 2H); 2.67 (br. m, 6H); 1.80-0.98 (br. m, 30H); 0.90 (t, 3H).

Step 2: *tert*-Butyl 4-((2-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperidine-1-carboxylate




Chemical Formula: C₄₁H₈₃N₃O₂

Molecular Weight: 650.13

[00718] To a mixture of *tert*-butyl 4-(2-(nonylamino)ethyl)piperidine-1-carboxylate (288 mg, 0.81 mmol) and *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (297 mg, 0.89 mmol) in MeCN (20 mL) was added K₂CO₃ (249 mg, 1.79 mmol), and KI (13 mg, 0.08 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT and filtered over a

pad of celite rinsing with hexanes, and concentrated. Purification by ISCO silica flash chromatography (0-100% DCM/[DCM 20% MeOH 1% NH₄OH]) provided *tert*-butyl 4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (216 mg, 41%)
 UPLC/ELSD: RT = 2.72 min. MS (ES): *m/z* (MH⁺) 651 for C₄₁H₈₃N₃O₂
 5 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (br, 2H); 2.83-2.29 (br. m, 14H); 1.75-1.00 (br. m, 58H); 0.90 (t, 9H).

Step 3: *N*¹,*N*¹,*N*²-Trinonyl-*N*²-(2-(piperidin-4-yl)ethyl)ethane-1,2-diamine




10

Chemical Formula: C₃₆H₇₅N₃

Molecular Weight: 550.02

[00719] In the same manner as Step 6 for Compound 44, *N*¹,*N*¹,*N*²-trinonyl-*N*²-(2-(piperidin-4-yl)ethyl)ethane-1,2-diamine was synthesized from *tert*-butyl 4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (216 mg, 0.33 mmol) and TFA (1.27 mL, 16.6 mmol) in DCM (2 mL). Yield (178 mg, 97%).
 UPLC/ELSD: RT = 1.84 min. MS (ES): *m/z* (MH⁺) 551 for C₃₆H₇₅N₃
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.08 (br. m, 2H); 2.70-2.25 (br. m, 14H); 2.0 (br, 1H); 1.80-1.02 (br. m, 49H); 0.90 (t, 9H).

20 Step 4: 2-(Dinonylamino)-1-(4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperidin-1-yl)ethan-1-one



Chemical Formula: C₅₆H₁₁₄N₄O

Molecular Weight: 859.56

25 [00720] In the same manner as Step 7 for Compound 44, 2-(dinonylamino)-1-(4-((2-(dinonylamino)ethyl)(nonyl)amino)ethyl)piperidin-1-yl)ethan-1-one was synthesized from dinonylglycine (96 mg, 0.29 mmol), *N*¹,*N*¹,*N*²-trinonyl-*N*²-(2-(piperidin-4-yl)ethyl)ethane-1,2-

diamine (178 mg, 0.32 mmol), iPr₂EtN (112 μ L, 0.65 mmol), and T3P (50% EtOAc solution, 525 μ L, 0.88 mmol) in THF (6 mL). Yield (121 mg, 48%).


UPLC/ELSD: RT = 2.96 min. MS (ES): *m/z* (MH⁺) 860 for C₅₆H₁₁₄N₄O

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.63-4.08 (br. m, 4H); 3.34-2.25 (br. m, 18H); 1.90-1.01

5 (br. m, 77H); 0.91 (t, 15H).

BA: Compound 51: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(4-ditetradecylglycyl)piperazin-1-yl)ethan-1-one

10 Step 1: Methyl ditetradecylglycinate



Chemical Formula: C₃₁H₆₃NO₂

Molecular Weight: 481.85


[00721] In the same manner as Step 1 from Compound 11, methyl 3-

15 (ditetradecylamino)propanoate was synthesized from glycine methyl ester hydrochloride (564 mg, 4.49 mmol), tetradecanal (2.1 g, 9.89 mmol), sodium triacetoxyborohydride (2.1 g, 9.89 mmol), acetic acid (0.6 mL, 9.89 mmol), trimethylamine (0.93 mL, 6.74 mmol), in DCE (22 mL). Yield (1.93 g, 89%).

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.72 (s, 3H); 3.34 (s, 2H); 1.56 (t, 4H); 1.60-1.03 (br. m,

20 48H); 0.91 (t, 6H).

Step 2: Lithium ditetradecylglycinate



Chemical Formula: C₃₀H₆₀LiNO₂


Molecular Weight: 473.76

[00722] In the same manner as Step 2 from Compound 11, lithium ditetradecylglycinate was synthesized from methyl ditetradecylglycinate (1.93 g, 4.0 mmol) and 1M LiOH (20 mL, 20 mmol) in THF (20 mL). Yield (1.81 g, 97%).

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.17 (s, 2H); 2.64 (t, 4H); 1.52 (br. m, 4H); 1.31 (br. m,

30 44H); 0.93 (t, 6H).

Step 3: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(4-(ditetradecylglycyl)piperazin-1-yl)ethan-1-one



5

Chemical Formula: C₆₅H₁₃₁N₅O₂

Molecular Weight: 1014.80

[00723] In the same manner as Step 7 for Compound 44, 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(4-(ditetradecylglycyl)piperazin-1-yl)ethan-1-one was synthesized from lithium ditetradecylglycinate (126 mg, 0.26 mmol), 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (134 mg, 0.24 mmol), iPr₂EtN (91 μ L, 0.52 mmol), and T3P (50% EtOAc solution, 424 μ L, 0.71 mmol) in THF (4 mL)


UPLC/ELSD: RT = 3.64 min. MS (ES): *m/z* (MH⁺) 1016 for C₆₅H₁₃₁N₅O₂

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.84-2.34 (br. m, 26H); 1.88-0.99 (br. m, 90H); 0.90 (t, 15H).

15

BB: Compound 52: 3-(Dinonylamino)-1-(4-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)propan-1-one

20 Step 1: Methyl 3-(dinonylamino)propanoate



Chemical Formula: C₂₂H₄₅NO₂

Molecular Weight: 355.61


[00724] To a mixture of methyl 3-aminopropanoate hydrochloride (2.0 g, 14 mmol) and 1-bromononane (2.7 mL, 14 mmol) in MeCN (100 mL) was added K₂CO₃ (4.0 g, 29 mmol) and KI (238 mg, 1.4 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT and filtered over a pad of celite washing with hexanes, and

concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-10% MeOH/DCM) provided methyl 3-(dinonylamino)propanoate (663 mg, 13%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 2.80 (t, 2H); 2.41 (br. m, 6H); 1.70-1.10 (br. m, 28H); 0.90 (t, 6H).

5


Step 2: Lithium 3-(dinonylamino)propanoate



Chemical Formula: C₂₁H₄₂LiNO₂

Molecular Weight: 347.51

10 [00725] In the same manner as Step 2 from Compound 11, lithium 3-(dinonylamino)propanoate was synthesized from methyl 3-(dinonylamino)propanoate (663 mg, 1.86 mmol) and 1M LiOH (9.32 mL, 9.32 mmol) in THF (10 mL). Yield (636 mg, 99%).
¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.94-1.65 (br. m, 8H); 1.65-1.04 (br. m, 28H); 0.90 (t, 6H).
tert-Butyl 4-(3-(dinonylamino)propanoyl)piperazine-1-carboxylate




Chemical Formula: C₃₀H₅₉N₃O₃

Molecular Weight: 509.82

15 [00726] In the same manner as Step 3 for Compound 11, *tert*-butyl 4-(3-(dinonylamino)propanoyl)piperazine-1-carboxylate was synthesized from lithium 3-(dinonylamino)propanoate (636 mg, 1.83 mmol), 1-boc-piperazine (388 mg, 2.08 mmol), iPr₂EtN (726 μL, 4.17 mmol), and T3P (50% EtOAc solution, 3.4 mL, 5.68 mmol) in THF (20 mL). Yield (839 mg, 87%).
UPLC/ELSD: RT = 2.26 min. MS (ES): *m/z* (MH⁺) 511 for C₃₀H₅₉N₃O₃
¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.71-2.21 (br. m, 16H); 1.92-0.98 (br. m, 37H); 0.90 (t, 6H).

Step 3: 3-(Dinonylamino)-1-(piperazin-1-yl)propan-1-one

Chemical Formula: C₂₅H₅₁N₃O

Molecular Weight: 409.70


[00727] In the same manner as Step 4 for Compound 11, 3-(dinonylamino)-1-(piperazin-1-yl)propan-1-one was synthesized from *tert*-butyl 4-(dinonylamino)propanoyl)piperazine-1-carboxylate (839 mg, 1.65 mmol), and TFA (6.3 mL, 83 mmol) in DCM (7 mL). Yield (501 mg, 74%).

UPLC/ELSD: RT = 1.19 min. MS (ES): *m/z* (MH⁺) 411 for C₂₅H₅₁N₃O

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.61 (t, 2H); 3.47 (t, 2H); 2.86 (br. m, 6H); 2.46 (br. m,

10 6H); 1.80 (br, 1H); 1.56-1.08 (br. m, 28H); 0.90 (t, 6H).

Step 4: Methyl 3-(nonylamino)propanoate


Chemical Formula: C₁₃H₂₇NO₂

Molecular Weight: 229.36

[00728] In the same manner as Step 1 for Compound 52, methyl 3-(nonylamino)propanoate was synthesized from methyl 3-aminopropanoate hydrochloride (2.0 g, 14 mmol), 1-bromononane (2.7 mL, 14 mmol), K₂CO₃ (4.0 g, 29 mmol) and KI (238 mg, 1.4 mmol) in MeCN (100 mL). Yield (300 mg, 9%)

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.70 (s, 3H); 2.91 (t, 2H); 2.60 (br. m, 4H); 1.90 (br, 1H); 1.58-1.02 (br. m, 14H); 0.90 (t, 3H).

Step 5: Methyl 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoate

Chemical Formula: C₃₃H₆₈N₂O₂

Molecular Weight: 524.92


25

[00729] In the same manner as Step 9 from Compound 10, methyl 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoate was synthesized from methyl 3-(nonylamino)propanoate (300 mg, 1.31 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (478 mg, 1.44 mmol), K_2CO_3 (400 mg, 2.88 mmol), and KI (22 mg, 0.13 mmol) in MeCN (20 mL).

5 Yield (348 mg, 51%).

UPLC/ELSD: RT = 2.66 min. MS (ES): m/z (MH $^+$) 526 for C₃₃H₆₈N₂O₂
¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.69 (s, 3H); 2.82 (t, 2H); 2.44 (br. m, 12H); 1.85-1.05 (br. m, 42H); 0.90 (t, 9H).

10 Step 6: 3-((2-(Dinonylamino)ethyl)(nonyl)amino)propanoic acid




Chemical Formula: C₃₂H₆₆N₂O₂

Molecular Weight: 510.89

[00730] In the same manner as Step 10 from Compound 10, 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid was synthesized from methyl 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoate (348 mg, 0.66 mmol) and 1M LiOH (3.3 mL, 3.3 mmol) in THF (3.3 mL). Yield (338 mg, 99%).

UPLC/ELSD: RT = 2.29 min. MS (ES): m/z (MH^+) 512 for $\text{C}_{32}\text{H}_{66}\text{N}_2\text{O}_2$

20 Step 7: 3-(Dinonylamino)-1-(4-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)propan-1-one



Chemical Formula: C₅₇H₁₁₅N₅O₂

Molecular Weight: 902.58

[00731] In the same manner as Step 11 for Compound 11, 3-(dinonylamino)-1-(4-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)propan-1-one was synthesized from 3-(dinonylamino)-1-(piperazin-1-yl)propan-1-one (298 mg, 0.73 mmol), 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid (338 mg, 0.68 mmol), iPr₂EtN (254 μ L, 1.46


mmol), and T3P (50% EtOAc solution, 1.18 mL, 1.98 mmol) in THF (10 mL). Yield (218 mg, 37%).

UPLC/ELSD: RT = 2.89 min. MS (ES): m/z (MH⁺) 903 for C₅₇H₁₁₅N₅O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.65 (br. m, 4H); 3.50 (br. m, 4H); 2.82 (br. m, 4H); 2.66-5 2.30 (br. m, 18H); 1.61-1.02 (br. m, 70H); 0.90 (t, 15H).

BC: Compound 53: Ethyl 7-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)heptanoate

10 Step 1: Ethyl 7-bromoheptanoate



Chemical Formula: C₉H₁₇BrO₂


Molecular Weight: 237.14

[00732] In the same manner as Step 1 for Compound 15, ethyl 7-bromoheptanoate was

15 synthesized from 7-bromoheptanoic acid (1.0 g, 4.8 mmol), ethanol (5.6 mL, 96 mmol), and H₂SO₄ (0.25 mL, 4.8 mmol) in THF (6 mL). Yield (911 mg, 80%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.15 (q, 2H); 3.42 (t, 2H); 2.33 (t, 2H); 1.87 (m, 2H); 1.66 (m, 2H); 1.57-1.14 (br. m, 7H).

20 Step 2: Ethyl 7-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)heptanoate



Chemical Formula: C₂₄H₄₇NO₄


Molecular Weight: 413.64

[00733] In the same manner as Step 3 for Compound 44, ethyl 7-((2-(*tert*-butoxy)-2-

25 oxoethyl)(nonyl)amino)heptanoate was synthesized from *tert*-butyl nonylglycinate (250 mg, 0.97 mmol), ethyl 7-bromoheptanoate (253 mg, 1.07 mmol), K₂CO₃ (270 mg, 1.94 mmol), and KI (16 mg, 0.10 mmol) in MeCN (10 mL). Yield (298 mg, 74%).

UPLC/ELSD: RT = 1.60 min. MS (ES): m/z (MH⁺) 414.68 for C₂₄H₄₇NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.14 (q, 2H); 3.23 (s, 2H); 2.57 (t, 4H); 2.31 (t, 2H); 1.74-30 1.12 (br. m, 34H); 0.90 (t, 3H).

Step 3: *N*-(7-Ethoxy-7-oxoheptyl)-*N*-nonylglycineChemical Formula: C₂₀H₃₉NO₄

5

Molecular Weight: 357.54


[00734] In the same manner as Step 4 for Compound 44, *N*-(7-ethoxy-7-oxoheptyl)-*N*-nonylglycine was synthesized from ethyl 7-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)heptanoate (298 mg, 0.72 mmol), and TFA (2.8 mL, 36 mmol) in DCM (3 mL). Yield (244 mg, 95%).

10 UPLC/ELSD: RT = 1.07 min. MS (ES): *m/z* (MH⁺) 358.50 for C₂₀H₃₉NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.15 (q, 2H); 3.46 (br, 2H); 3.01 (br, 4H); 2.31 (t, 2H); 1.86-1.10 (br. m, 25H); 0.91 (t, 3H).

Step 4: Ethyl 7-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-

15 oxoethyl)(nonyl)amino)heptanoate

Chemical Formula: C₅₅H₁₀₉N₅O₄

Molecular Weight: 904.51

[00735] In the same manner as Step 7 for Compound 44, ethyl 7-((2-(4-(*N*-(2-


20 (dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)heptanoate was synthesized from *N*-(7-ethoxy-7-oxoheptyl)-*N*-nonylglycine (111 mg, 0.31 mmol), 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (160 mg, 0.28 mmol), iPr₂EtN (109 μL, 0.62 mmol), and T3P (50% EtOAc solution, 0.51 mL, 0.81 mmol) in THF (10 mL). Yield (70 mg, 27%).

25 UPLC/ELSD: RT = 2.88 min. MS (ES): *m/z* (MH⁺) 905.33 for C₅₅H₁₀₉N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.00 (q, 2H); 3.65-3.34 (br. m, 8H); 3.19 (s, 2H); 3.13 (s, 2H); 2.50-2.10 (br. m, 16H); 1.65-0.90 (br. m, 67H); 0.75 (t, 12H).

BD: Compound 54: Propyl 6-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)hexanoate

Step 1: Propyl 6-bromohexanoate




Chemical Formula: C₉H₁₇BrO₂

Molecular Weight: 237.14

[00736] In the same manner as Step 1 for Compound 15, propyl 6-bromohexanoate was synthesized from 6-bromohexanoic acid (1.0 g, 5.1 mmol), 1-propanol (1.5 g, 26 mmol), and 10 H₂SO₄ (0.27 mL, 5.1 mmol) in THF (5 mL). Yield (1.14 g, 94%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.05 (t, 2H); 3.43 (t, 2H); 2.34 (t, 2H); 1.90 (m, 2H); 1.68 (m, 4H); 1.50 (m, 2H); 0.96 (t, 3H).

Step 2: Propyl 6-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)hexanoate



Chemical Formula: C₂₄H₄₇NO₄


Molecular Weight: 413.64

[00737] In the same manner as Step 3 for Compound 44, propyl 6-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)hexanoate was synthesized from *tert*-butyl nonylglycinate (250 mg, 0.97 mmol), propyl 6-bromohexanoate (253 mg, 1.07 mmol), K₂CO₃ (270 mg, 1.94 mmol), and KI (16 mg, 0.10 mmol) in MeCN (10 mL). Yield (258 mg, 64%).

UPLC/ELSD: RT = 1.62 min. MS (ES): *m/z* (MH⁺) 414.59 for C₂₄H₄₇NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.05 (t, 2H); 3.23 (s, 2H); 2.58 (br. m, 4H); 2.33 (t, 2H); 1.75-1.15 (br. m, 31H); 0.91 (m, 6H).

25 Step 3: *N*-Nonyl-*N*-(6-oxo-6-propoxyhexyl)glycine




Chemical Formula: C₂₀H₃₉NO₄

Molecular Weight: 357.54

5 [00738] In the same manner as Step 4 for Compound 44, *N*-nonyl-*N*-(6-oxo-6-propoxyhexyl)glycine was synthesized from propyl 6-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)hexanoate (258 mg, 0.62 mmol), and TFA (2.4 mL, 31 mmol) in DCM (3 mL). Yield (223 mg, 99%).

UPLC/ELSD: RT = 1.13 min. MS (ES): *m/z* (MH⁺) 358.50 for C₂₀H₃₉NO₄
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (t, 2H); 3.34 (s, 2H); 2.87 (br. m, 4H); 2.36 (t, 2H); 1.77-1.10 (br. m, 22H); 0.92 (m, 6H).

10 10 Step 4: Propyl 6-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)hexanoate



Chemical Formula: C₅₅H₁₀₉N₅O₄


Molecular Weight: 904.51

15 [00739] In the same manner as Step 7 for Compound 44, propyl 6-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)hexanoate was synthesized from *N*-nonyl-*N*-(6-oxo-6-propoxyhexyl)glycine (111 mg, 0.31 mmol), 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (160 mg, 0.28 mmol), iPr₂EtN (109 μL, 0.62 mmol), and T3P (50% EtOAc solution, 0.51 mL, 0.81 mmol) in THF (10 mL). Yield (72 mg, 28%).

UPLC/ELSD: RT = 2.91 min. MS (ES): *m/z* (MH⁺) 905.33 for C₅₅H₁₀₉N₅O₄
¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (t, 2H); 3.78-3.46 (br. m, 8H); 3.34 (s, 2H); 3.28 (s, 2H); 2.68-2.24 (br. m, 16H); 1.85-1.10 (br. m, 64H); 0.92 (m, 15H).

25 25 BE: Compound 55: Butyl 5-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)pentanoate

Step 1: Butyl 5-bromopentanoate




Chemical Formula: C₉H₁₇BrO₂

Molecular Weight: 237.14

[00740] In the same manner as Step 1 for Compound 15, butyl 5-bromopentanoate was synthesized from 5-bromopentanoic acid (1.47 g, 8.1 mmol), 1-butanol (0.50 g, 6.8 mmol), and H₂SO₄ (0.36 mL, 6.8 mmol) in THF (7 mL). Yield (1.42 g, 0.89%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.10 (t, 2H); 4.34 (t, 2H); 2.36 (t, 2H); 1.93 (m, 2H); 1.80 (m, 2H); 1.62 (m, 2H); 1.40 (m, 2H); 0.96 (t, 3H).

10 Step 2: Butyl 5-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)pentanoate



Chemical Formula: C₂₄H₄₇NO₄


Molecular Weight: 413.64

[00741] In the same manner as Step 3 for Compound 44, butyl 5-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)pentanoate was synthesized from *tert*-butyl nonylglycinate (250 mg, 0.97 mmol), butyl 5-bromopentanoate (253 mg, 1.07 mmol), K₂CO₃ (270 mg, 1.94 mmol), and KI (16 mg, 0.10 mmol) in MeCN (10 mL). Yield (284 mg, 71%).

UPLC/ELSD: RT = 1.67 min. MS (ES): *m/z* (MH⁺) 414.59 for C₂₄H₄₇NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (t, 2H); 3.23 (s, 2H); 2.58 (m, 4H); 2.34 (t, 2H); 1.74-2.0 (br. m, 31H); 0.93 (m, 6H).

Step 3: *N*-(5-butoxy-5-oxopentyl)-*N*-nonyl glycine



Chemical Formula: C₂₀H₃₉NO₄

Molecular Weight: 357.54

[00742] In the same manner as Step 4 for Compound 44, *N*-(5-butoxy-5-oxopentyl)-*N*-nonyl glycine was synthesized from butyl 5-((2-(*tert*-butoxy)-2-


oxoethyl)(nonyl)amino)pentanoate (284 mg, 0.69 mmol), and TFA (2.6 mL, 34 mmol) in DCM (3 mL). Yield (245 mg, 99%).

UPLC/ELSD: RT = 1.09 min. MS (ES): m/z (MH⁺) 358.50 for C₂₀H₃₉NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.05 (t, 2H); 3.48 (s, 2H); 3.03 (br. m, 4H); 2.34 (t, 2H);

5 1.85-1.15 (br. m, 22H); 0.93 (m, 6H).

Step 4: Butyl 5-((2-(4-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)pentanoate



10

Chemical Formula: C₅₅H₁₀₉N₅O₄

Molecular Weight: 904.51

[00743] In the same manner as Step 7 for Compound 44, butyl 5-((2-(4-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)pentanoate was synthesized from *N*-(5-butoxy-5-oxopentyl)-*N*-nonylglycine (111 mg, 0.31 mmol), 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (160 mg, 0.28 mmol), iPr₂EtN (109 μL, 0.62 mmol), and T3P (50% EtOAc solution, 0.51 mL, 0.81 mmol) in THF (10 mL). Yield (92 mg, 36%).


UPLC/ELSD: RT = 2.88 min. MS (ES): m/z (MH⁺) 905.33 for C₅₅H₁₀₉N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.90 (t, 2H); 3.65-3.35 (br. m, 8H); 3.19 (s, 2H); 3.13 (t,

20 2H); 2.52-2.06 (br. m, 16H); 1.65-0.95 (br. m, 64H); 0.77 (m, 15H).

BF: Compound 56: 3-((2-(4-(N-(2-(Dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)propyl hexanoate

Step 1: 3-Bromopropyl hexanoate



25


Chemical Formula: C₉H₁₇BrO₂

Molecular Weight: 237.14

[00744] In the same manner as Step 1 for Compound 15, 3-bromopropyl hexanoate was synthesized from 3-bromopropan-1-ol (0.87 mL, 9.6 mmol), hexanoic acid (1.0 mL, 8.0 mmol), and H₂SO₄ (1.0 mL, 8.0 mmol) in THF (10 mL). Yield (823 mg, 44%).

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.23 (t, 2H); 3.49 (t, 2H), 2.33 (t, 2H); 2.20 (m, 2H); 1.65 (m, 2H); 1.34 (m, 4H), 0.91 (t, 3H).

Step 2: 3-((2-(*tert*-Butoxy)-2-oxoethyl)(nonyl)amino)propyl hexanoate



Chemical Formula: C₂₄H₄₇NO₄


10 Molecular Weight: 413.64

[00745] In the same manner as Step 3 for Compound 44, 3-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)propyl hexanoate was synthesized from *tert*-butyl nonylglycinate (250 mg, 0.97 mmol), 3-bromopropyl hexanoate (253 mg, 1.07 mmol), K₂CO₃ (270 mg, 1.94 mmol), and KI (16 mg, 0.10 mmol) in MeCN (10 mL). Yield (335 mg, 83%).

15 UPLC/ELSD: RT = 1.78 min. MS (ES): *m/z* (MH⁺) 414.59 for C₂₄H₄₇NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.14 (t, 2H); 3.23 (s, 2H); 2.68 (t, 2H); 2.58 (t, 2H); 2.30 (t, 2H); 1.79 (m, 2H); 1.64 (m, 2H); 1.55-1.20 (br. m, 27H); 0.91 (m, 6H).

Step 3: *N*-(3-(Hexanoyloxy)propyl)-*N*-nonylglycine



20 Chemical Formula: C₂₀H₃₉NO₄


Molecular Weight: 357.54

[00746] In the same manner as Step 4 for Compound 44, *N*-(3-(hexanoyloxy)propyl)-*N*-nonylglycine was synthesized from 3-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)propyl

25 hexanoate (335 mg, 0.81 mmol), and TFA (3.1 mL, 40 mmol) in DCM (4 mL). Yield (284 mg, 98%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.11 (t, 2H); 3.17 (s, 2H); 2.68 (br. m, 4H); 2.30 (t, 2H); 1.87 (m, 2H); 1.63 (m, 2H); 1.48 (m, 2H); 1.28 (br. m, 16H); 0.91 (m, 6H).

Step 4: 3-((2-(4-(*N*-(2-(Dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)propyl hexanoate



5

Chemical Formula: C₅₅H₁₀₉N₅O₄

Molecular Weight: 904.51


[00747] In the same manner as Step 7 for Compound 44, 3-((2-(4-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperazin-1-yl)-2-oxoethyl)(nonyl)amino)propyl hexanoate was synthesized from *N*-(3-(hexanoyloxy)propyl)-*N*-nonylglycine (111 mg, 0.31 mmol), 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (160 mg, 0.28 mmol), iPr₂EtN (109 μ L, 0.62 mmol), and T3P (50% EtOAc solution, 0.51 mL, 0.81 mmol) in THF (10 mL). Yield (55 mg, 21%).

UPLC/ELSD: RT = 2.94 min. MS (ES): *m/z* (MH⁺) 905.25 for C₅₅H₁₀₉N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.01 (t, 2H); 3.77-3.43 (br. m, 8H), 3.23 (m, 4H); 2.64-2.15 (m, 16H); 1.70 (m, 2H); 1.54 (m, 2H); 1.50-0.96 (br. m, 60H); 0.81 (m, 15H).

BG: Compound 57: Methyl 8-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)octanoate

20 Step 1: *tert*-Butyl 4-((2-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₅₈N₂O₄

Molecular Weight: 510.80


[00748] To a mixture of *tert*-butyl 4-(nonylamino)ethyl)piperidine-1-carboxylate (239 mg, 0.67 mmol) and methyl 8-bromo-8-oxooctanoate (192 mg, 0.81 mmol) in MeCN (10 mL) was added K₂CO₃ (188 mg, 1.35 mmol) and KI (11 mg, 0.07 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT and filtered over a pad of celite

rinsing with EtOAc, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-80% EtOAc/hexanes) provided *tert*-butyl 4-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (241 mg, 70%).

UPLC/ELSD: RT = 1.87 min. MS (ES): *m/z* (MH⁺) 512.76 for C₃₀H₅₈N₂O₄

5 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.07 (br. m, 2H); 3.69 (s, 3H); 2.70 (m, 2H); 2.37 (br. m, 8H); 1.75-1.00 (br. m, 40H); 0.90 (t, 3H).

Step 2: Methyl 8-(nonyl(2-(piperidin-4-yl)ethyl)amino)octanoate




10 Chemical Formula: C₂₅H₅₀N₂O₂

Molecular Weight: 410.69

[00749] In the same manner as Step 4 for Compound 11, methyl 8-(nonyl(2-(piperidin-4-yl)ethyl)amino)octanoate was synthesized from *tert*-butyl 4-((8-methoxy-8-oxooctyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (241 mg, 0.47 mmol), and TFA (1.8 mL, 15 24 mmol) in DCM (2 mL). Yield (193 mg, 99%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69 (s, 3H); 3.08 (m, 2H); 2.60 (m, 2H); 2.49-2.24 (br. m, 8H); 2.06 (br, 1H); 1.78-1.02 (br. m, 31H); 0.90 (t, 3H).

Step 3: Methyl 8-((2-(1-(*N*-(2-(dinonylamoxy)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)octanoate



Chemical Formula: C₅₀H₁₁₂N₄O₃

Molecular Weight: 889.54

[00750] In the same manner as Step 11 for Compound 11, methyl 8-((2-(1-(*N*-(2-(dinonylamoxy)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)octanoate was synthesized from methyl 8-(nonyl(2-(piperidin-4-yl)ethyl)amino)octanoate (141 mg, 0.34

mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (188 mg, 0.38 mmol), iPr₂EtN (132 μ L, 0.76 mmol), and T3P (50% EtOAc solution, 614 μ L, 1.03 mmol) in THF (10 mL). Yield (70 mg, 23%).


UPLC/ELSD: RT = 2.97 min. MS (ES): *m/z* (MH⁺) 890.24 for C₅₆H₁₁₂N₄O₃

5 ¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.64-4.09 (br. m, 2H); 3.69 (s, 3H); 3.42-2.83 (br. m, 3H); 2.69-2.24 (br. m, 19H); 1.81-0.99 (br. m, 73H); 0.90 (t, 12H).

BH: Compound 58: Ethyl 7-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)heptanoate

10

Step 1: *tert*-Butyl 4-(2-((7-ethoxy-7-oxoheptyl)(nonyl)amino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₅₈N₂O₄


Molecular Weight: 510.80

15 [00751] In the same manner as Step 1 for Compound 57, *tert*-butyl 4-(2-((7-ethoxy-7-oxoheptyl)(nonyl)amino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(2-(nonylamino)ethyl)piperidine-1-carboxylate (239 mg, 0.67 mmol), ethyl 7-bromoheptanoate (192 mg, 0.81 mmol), K₂CO₃ (188 mg, 1.35 mmol) and KI (11 mg, 0.07 mmol) in MeCN (10 mL). Yield (247 mg, 72%).

20 UPLC/ELSD: RT = 1.91 min. MS (ES): *m/z* (MH⁺) 511.62 for C₃₀H₅₈N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.12 (br. m, 4H); 2.80-2.15 (br. m, 10H); 1.75-1.00 (br. m, 41H); 0.90 (t, 3H).

Step 2: Ethyl 7-(nonyl(2-(piperidin-4-yl)ethyl)amino)heptanoate



Chemical Formula: C₂₅H₅₀N₂O₂

Molecular Weight: 410.69


25

[00752] In the same manner as Step 4 for Compound 11, ethyl 7-(nonyl(2-(piperidin-4-yl)ethyl)amino)heptanoate was synthesized from *tert*-butyl 4-(2-((7-ethoxy-7-

oxoheptyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (247 mg, 0.48 mmol), and TFA (1.9 mL, 24 mmol) in DCM (2 mL). Yield (194 mg, 98%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.14 (t, 2H); 3.08 (m, 2H); 2.60 (m, 2H); 2.52-2.24 (br. m, 8H); 2.12 (br, 1H); 1.77-1.05 (br. m, 32H); 0.90 (t, 3H).

5 Ethyl 7-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)heptanoate



Chemical Formula: C₅₆H₁₁₂N₄O₃

Molecular Weight: 889.54


10 [00753] In the same manner as Step 11 for Compound 11, ethyl 7-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)heptanoate was synthesized from ethyl 7-(nonyl(2-(piperidin-4-yl)ethyl)amino)heptanoate (141 mg, 0.34 mmol), N-(2-(dinonylamino)ethyl)-N-nonylglycine (188 mg, 0.38 mmol), iPr₂EtN (132 μL, 0.76 mmol), and T3P (50% EtOAc solution, 614 μL, 1.03 mmol) in THF (10 mL). Yield (42 mg, 14%).

15 UPLC/ELSD: RT = 3.00 min. MS (ES): *m/z* (MH⁺) 890.32 for C₅₆H₁₁₂N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.55-3.95 (br. m, 4H); 3.38-2.72 (br. m, 4H); 2.66-2.10 (br. m, 18H); 1.72-0.91 (br. m, 74H); 0.81 (t, 12H).

20 BI: Compound 59: Propyl 6-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)hexanoate

Step 1: *tert*-Butyl 4-(2-(nonyl(6-oxo-6-propoxyhexyl)amino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₅₈N₂O₄

25 Molecular Weight: 510.80


[00754] In the same manner as Step 1 for Compound 57, *tert*-butyl 4-(2-(nonyl(6-oxo-6-propoxyhexyl)amino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(2-

(nonylamino)ethyl)piperidine-1-carboxylate (239 mg, 0.67 mmol), propyl 6-bromohexanoate (192 mg, 0.81 mmol), K_2CO_3 (188 mg, 1.35 mmol) and KI (11 mg, 0.07 mmol) in MeCN (10 mL). Yield (240 mg, 70%).

UPLC/ELSD: RT = 1.93 min. MS (ES): m/z (MH^+) 511.78 for $C_{30}H_{58}N_2O_4$

5 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.05 (br. m, 4H); 2.80-2.20 (br. m, 10H); 1.85-1.04 (br. m, 38H); 0.92 (t, 6H).

Step 2: Propyl 6-(nonyl(2-(piperidin-4-yl)ethyl)amino)hexanoate



10 Chemical Formula: $C_{25}H_{50}N_2O_2$


Molecular Weight: 410.69

[00755] In the same manner as Step 4 for Compound 11, propyl 6-(nonyl(2-(piperidin-4-yl)ethyl)amino)hexanoate was synthesized from *tert*-butyl 4-(2-(nonyl(6-oxo-6-propoxyhexyl)amino)ethyl)piperidine-1-carboxylate (240 mg, 0.47 mmol), and TFA (1.8 mL, 23 mmol) in DCM (2 mL). Yield (183 mg, 95%).

15 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.04 (t, 2H); 3.08 (m, 2H); 2.60 (m, 2H); 2.35 (br. m, 8H); 1.95 (br, 1H); 1.75-1.00 (br. m, 29H); 0.92 (m, 6H).

Step 3: Propyl 6-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-

20 yl)ethyl)(nonyl)amino)hexanoate



Chemical Formula: $C_{56}H_{112}N_4O_3$

Molecular Weight: 889.54

[00756] In the same manner as Step 11 for Compound 11, propyl 6-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)hexanoate was synthesized from propyl 6-(nonyl(2-(piperidin-4-yl)ethyl)amino)hexanoate (141 mg, 0.34

mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (188 mg, 0.38 mmol), iPr₂EtN (132 μ L, 0.76 mmol), and T3P (50% EtOAc solution, 614 μ L, 1.03 mmol) in THF (10 mL). Yield (67 mg, 22%).


UPLC/ELSD: RT = 3.02 min. MS (ES): *m/z* (MH⁺) 890.32 for C₅₆H₁₁₂N₄O₃

5 ¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.65-4.13 (br. m, 2H); 4.05 (t, 2H); 3.50-2.81 (br. m, 4H); 2.69-2.18 (br. m, 18H); 1.98-1.02 (br. m, 71H); 0.92 (br. m, 15H).

BJ: Compound 60: Butyl 5-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)pentanoate

10

Step 1: *tert*-Butyl 4-(2-((5-butoxy-5-oxopentyl)(nonyl)amino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₅₈N₂O₄


Molecular Weight: 510.80

15 [00757] In the same manner as Step 1 for Compound 57, *tert*-butyl 4-(2-((5-butoxy-5-oxopentyl)(nonyl)amino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(2-(nonylamino)ethyl)piperidine-1-carboxylate (239 mg, 0.67 mmol), butyl 5-bromopentanoate (192 mg, 0.81 mmol), K₂CO₃ (188 mg, 1.35 mmol) and KI (11 mg, 0.07 mmol) in MeCN (10 mL). Yield (211 mg, 61%).

20 UPLC/ELSD: RT = 1.95 min. MS (ES): *m/z* (MH⁺) 511.78 for C₃₀H₅₈N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.09 (br. m, 4H); 2.70 (br. m, 2H); 2.38 (br. m, 8H); 1.73-1.02 (br. m, 38H); 0.93 (br. m, 6H).

Step 2: Butyl 5-(nonyl(2-(piperidin-4-yl)ethyl)amino)pentanoate



Chemical Formula: C₂₅H₅₀N₂O₂

Molecular Weight: 410.69


25

[00758] In the same manner as Step 4 for Compound 11, butyl 5-(nonyl(2-(piperidin-4-yl)ethyl)amino)pentanoate was synthesized from *tert*-butyl 4-(2-((5-butoxy-5-

oxopentyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (211 mg, 0.41 mmol), and TFA (1.6 mL, 21 mmol) in DCM (2 mL). Yield (169 mg, 99%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (t, 2H); 3.07 (m, 2H); 2.60 (m, 2H); 2.37 (br. m, 8H); 1.83 (br, 1H); 1.76-1.04 (br. m, 29H); 0.93 (br. m, 6H).

5 Step 3: Butyl 5-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)pentanoate



Chemical Formula: C₅₆H₁₁₂N₄O₃

Molecular Weight: 889.54

10 [00759] In the same manner as Step 11 for Compound 11, butyl 5-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)pentanoate was synthesized from butyl 5-(nonyl(2-(piperidin-4-yl)ethyl)amino)pentanoate (141 mg, 0.34 mmol), N-(2-(dinonylamino)ethyl)-N-nonylglycine (188 mg, 0.38 mmol), iPr₂EtN (132 μL, 0.76 mmol), and T3P (50% EtOAc solution, 614 μL, 1.03 mmol) in THF (10 mL). Yield (46 mg, 15%).


15 UPLC/ELSD: RT = 3.03 min. MS (ES): *m/z* (MH⁺) 890.32 for C₅₆H₁₁₂N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.62-4.13 (br. m, 2H); 4.09 (t, 2H); 3.41-2.84 (br. m, 4H); 2.72-2.25 (br. m, 18H); 1.82-1.02 (br. m, 71H); 0.91 (br. m, 15H).

BK: Compound 61: 3-((2-(1-(N-(2-(Dinonylamino)ethyl)-N-nonylglycyl)piperidin-4-

20 yl)ethyl)(nonyl)amino)propyl hexanoate

Step 1: *tert*-Butyl 4-((2-((3-(hexanoyloxy)propyl)(nonyl)amino)ethyl)piperidine-1-carboxylate



25 Chemical Formula: C₃₀H₅₈N₂O₄

Molecular Weight: 510.80


[00760] In the same manner as Step 1 for Compound 57, *tert*-butyl 4-((3-(hexanoyloxy)propyl)(nonyl)amino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-

butyl 4-(2-(nonylamino)ethyl)piperidine-1-carboxylate (239 mg, 0.67 mmol), 3-bromopropyl hexanoate (192 mg, 0.81 mmol), K_2CO_3 (188 mg, 1.35 mmol) and KI (11 mg, 0.07 mmol) in MeCN (10 mL). Yield (195 mg, 57%).

UPLC/ELSD: RT = 1.97 min. MS (ES): m/z (MH^+) 511.86 for $C_{30}H_{58}N_2O_4$

5 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.11 (br. m, 4H); 2.69 (m, 2H); 2.56-2.22 (br. m, 8H); 1.76 (m, 2H); 1.65 (m, 4H); 1.55-1.05 (br. m, 32H); 0.91 (m, 6H).

Step 2: 3-(Nonyl(2-(piperidin-4-yl)ethyl)amino)propyl hexanoate



10

Chemical Formula: $C_{25}H_{50}N_2O_2$


Molecular Weight: 410.69

[00761] In the same manner as Step 4 for Compound 11, 3-(nonyl(2-(piperidin-4-yl)ethyl)amino)propyl hexanoate was synthesized from *tert*-butyl 4-(2-((3-

15 (hexanoyloxy)propyl)(nonyl)amino)ethyl)piperidine-1-carboxylate (195 mg, 0.38 mmol), and TFA (1.5 mL, 19 mmol) in DCM (2 mL). Yield (149 mg, 95%).

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.12 (t, 2H); 3.08 (m, 2H); 2.70-2.22 (br. m, 10H); 2.07 (br, 1H); 1.70 (br. m, 6H); 1.48-1.00 (br. m, 23H); 0.91 (m, 6H).

20 Step 3: 3-((2-(1-(*N*-(2-(Dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)propyl hexanoate



Chemical Formula: $C_{56}H_{112}N_4O_3$

Molecular Weight: 889.54

25 [00762] In the same manner as Step 11 for Compound 11, 3-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)propyl hexanoate was synthesized from 3-(nonyl(2-(piperidin-4-yl)ethyl)amino)propyl hexanoate (141 mg, 0.34 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (188 mg, 0.38 mmol), iPr_2EtN (132 μ L, 0.76


mmol), and T3P (50% EtOAc solution, 614 μ L, 1.03 mmol) in THF (10 mL). Yield (64 mg, 21%).

UPLC/ELSD: RT = 3.02 min. MS (ES): m/z (MH $^+$) 890.41 for C₅₆H₁₁₂N₄O₃
¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.67 (br. m, 4H); 3.42-2.81 (br. m, 3H); 2.73-2.23 (br. m, 19H); 1.87-1.00 (br. m, 71H); 0.90 (t, 15H).

BL: Compound 62: Pentyl 4-((3-(4-(3-((2-dinylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)-3-oxopropyl)(nonyl)amino)butanoate

10

Step 1: *tert*-Butyl 3-(nonylamino)propanoate




Chemical Formula: C₁₆H₃₃NO₂

Molecular Weight: 271.45

15 [00763] In the same manner as Step 1 for Compound 44, *tert*-butyl 3-(nonylamino)propanoate was synthesized from *tert*-butyl 3-aminopropanoate hydrochloride (2.8 g, 15 mmol), 1-bromononane (3.2 g, 15 mmol), K₂CO₃ (4.3 g, 31 mmol), and KI (256 mg, 1.54 mmol) in MeCN (200 mL). Yield (1.74 g, 42%).

20 ¹H-NMR (300 MHz, CDCl₃) δ : ppm 2.86 (t, 2H); 2.63 (t, 2H); 2.46 (t, 2H); 1.65 (br, 1H); 1.47 (br. m, 11H); 1.29 (br. m, 12H); 0.90 (t, 3H).

Step 2: Pentyl 4-((3-(*tert*-butoxy)-3-oxopropyl)(nonyl)amino)butanoate



Chemical Formula: C₂₅H₄₉NO₄

Molecular Weight: 427.67


25 [00764] In the same manner as Step 3 for Compound 44, pentyl 4-((3-(*tert*-butoxy)-3-oxopropyl)(nonyl)amino)butanoate was synthesized from *tert*-butyl 3-(nonylamino)propanoate

(750 mg, 2.76 mmol), pentyl 4-bromobutanoate (786 mg, 3.31 mmol), K_2CO_3 (764 mg, 5.52 mmol), and KI (46 mg, 0.28 mmol) in MeCN (30 mL). Yield (934 mg, 79%).

UPLC/ELSD: RT = 1.82 min. MS (ES): m/z (MH^+) 428.62 for $C_{25}H_{49}NO_4$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.08 (t, 2H); 2.74 (t, 2H); 2.50-2.28 (br. m, 8H); 1.76 (m, 2H); 1.64 (m, 2H); 1.50-1.14 (br. m, 27H); 0.91 (m, 6H).

Step 3: 3-(Nonyl(4-oxo-4-(pentyloxy)butyl)amino)propanoic acid



Chemical Formula: $C_{21}H_{41}NO_4$

10 Molecular Weight: 371.56


[00765] In the same manner as Step 4 for Compound 44, 3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propanoic acid was synthesized from pentyl 4-((3-(*tert*-butoxy)-3-oxopropyl)(nonyl)amino)butanoate (934 mg, 2.18 mmol), and TFA (8.4 mL, 109 mmol) in DCM (10 mL). Yield (793 mg, 98%).

15 UPLC/ELSD: RT = 1.23 min. MS (ES): m/z (MH^+) 372.52 for $C_{21}H_{41}NO_4$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.10 (t, 2H); 2.88 (t, 2H); 2.70 (br. m, 4H); 2.52 (t, 2H); 2.38 (t, 2H); 1.90 (m, 2H); 1.73-1.49 (br. m, 4H); 1.47-1.17 (br. m, 16H); 0.92 (m, 6H).

Step 4: *tert*-Butyl 4-(3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propanoyl)piperazine-1-

20 carboxylate



Chemical Formula: $C_{30}H_{57}N_3O_5$


Molecular Weight: 539.80

[00766] In the same manner as Step 3 for Compound 11, *tert*-butyl 4-(3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propanoyl)piperazine-1-carboxylate was synthesized from 3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propanoic acid (793 mg, 2.13 mmol), 1-boc-piperazine (477 mg,

2.56 mmol), iPr₂EtN (0.82 mL, 4.7 mmol), and T3P (50% EtOAc solution, 3.8 mL, 6.4 mmol). Yield (1.15 g, 99%).

UPLC/ELSD: RT = 1.86 min. MS (ES): *m/z* (MH⁺) 540.65 for C₃₀H₅₇N₃O₅

5 Step 5: Pentyl 4-(nonyl(3-oxo-3-(piperazin-1-yl)propyl)amino)butanoate



Chemical Formula: C₂₅H₄₉N₃O₃

Molecular Weight: 439.69

[00767] In the same manner as Step 4 for Compound 44, pentyl 4-(nonyl(3-oxo-3-


10 (piperazin-1-yl)propyl)amino)butanoate was synthesized from *tert*-butyl 4-(3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propanoyl)piperazine-1-carboxylate (1.15 g, 2.13 mmol), and TFA (8.2 mL, 106 mmol) in DCM (10 mL). Yield (901 mg, 96%).

UPLC/ELSD: RT = 0.75 min. MS (ES): *m/z* (MH⁺) 440.47 for C₂₅H₄₉N₃O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.70-3.40 (br. m, 4H); 2.88 (br. m, 6H); 2.57

15 (br. m, 6H); 2.36 (t, 2H); 1.83 (m, 2H); 1.64 (m, 2H); 1.49 (m, 2H); 1.41-1.18 (br. m, 17H); 0.91 (m, 6H).

Step 6: *tert*-Butyl 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoate



20 Chemical Formula: C₃₆H₇₄N₂O₂

Molecular Weight: 567.00

[00768] In the same manner as Step 4 for Compound 46, *tert*-butyl 3-((2-

(dinonylamino)ethyl)(nonyl)amino)propanoate was synthesized from *tert*-butyl 3-

(nonylamino)propanoate (1.13 mg, 4.14 mol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (1.65 g,


25 4.97 mmol), K₂CO₃ (1.15 g, 8.33 mmol), and KI (138 mg, 0.83 mmol) in MeCN (100 mL).

Yield (1.41 g, 60%).

UPLC/ELSD: RT = 2.90 min. MS (ES): *m/z* (MH⁺) 567.79 for C₃₆H₇₄N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.78 (t, 2H); 2.69-2.29 (br. m, 12H); 1.55-1.15 (br. m, 51H); 0.90 (t, 9H).

Step 7: 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid



5

Chemical Formula: C₃₂H₆₆N₂O₂

Molecular Weight: 510.89


[00769] In the same manner as Step 4 for Compound 44, 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid was synthesized from *tert*-butyl 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoate (1.41 g, 2.49 mmol), and TFA (9.6 mL, 124 mmol) in DCM (10 mL). Yield (924 mg, 73%).

UPLC/ELSD: RT = 2.26 min. MS (ES): m/z (MH $^+$) 511.78 for C₃₂H₆₆N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 2.76 (br. m, 6H); 2.61 (br. m, 6H); 2.47 (t, 2H); 1.52 (br. m, 6H); 1.40-1.10 (br. m, 36H); 0.90 (t, 9H).

15

Step 8: Penty 4-((3-(4-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)-3-oxopropyl)(nonyl)amino)butanoate



Chemical Formula: C₅₇H₁₁₃N₅O₄

20

Molecular Weight: 932.56


[00770] In the same manner as Step 11 for Compound 11, pentyl 4-((3-(4-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)-3-oxopropyl)(nonyl)amino)butanoate was synthesized from pentyl 4-(nonyl(3-oxo-3-(piperazin-1-yl)propyl)amino)butanoate (268 mg, 0.61 mmol), 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid (343 mg, 0.67 mmol), iPr₂EtN (234 μ L, 1.34 mmol), and T3P (50% EtOAc solution, 1.09 mL, 1.83 mmol) in THF (20 mL). Yield (243 mg, 43%).

UPLC/ELSD: RT = 2.26 min. MS (ES): m/z (MH $^+$) 933.10 for C₅₇H₁₁₃N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.07 (t, 2H); 3.72-3.40 (br. m, 8H); 2.81 (m, 4H); 2.66-2.28 (br. m, 20H); 1.77 (m, 2H); 1.64 (m, 2H); 1.54-1.08 (br. m, 60H); 0.90 (t, 15H).

5 BM: Compound 69: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(3-(2-(dinonylamino)ethyl)pyrrolidin-1-yl)ethan-1-one

Step 1: *tert*-Butyl 3-(2-(dinonylamino)ethyl)pyrrolidine-1-carboxylate



Chemical Formula: C₂₉H₅₈N₂O₂


10 Molecular Weight: 466.80

[00771] In the same manner as Step 1 for Compound 49, *tert*-butyl 3-(2-(dinonylamino)ethyl)pyrrolidine-1-carboxylate was synthesized from *tert*-butyl 3-(2-aminoethyl)pyrrolidine-1-carboxylate (1.25 g, 5.47 mmol), 1-bromononane (1.13 g, 5.47 mmol), K₂CO₃ (757 mg, 5.47 mmol), and KI (91 mg, 0.55 mmol) in MeCN (100 mL). Yield (710 mg, 28%).

UPLC/ELSD: RT = 2.23 min. MS (ES): *m/z* (MH⁺) 467.74 for C₂₉H₅₈N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.67-3.34 (br. m, 2H); 3.34-2.75 (br. m 2H); 2.52-1.89 (br. m, 8H); 1.70-1.03 (br. m, 40H); 0.90 (t, 6H).

20 Step 2: *N*-Nonyl-*N*-(2-(pyrrolidin-3-yl)ethyl)nonan-1-amine



Chemical Formula: C₂₄H₅₀N₂


Molecular Weight: 366.68

[00772] In the same manner as Step 4 for Compound 11, *N*-nonyl-*N*-(2-(pyrrolidin-3-yl)ethyl)nonan-1-amine was synthesized from *tert*-butyl 3-(2-(dinonylamino)ethyl)pyrrolidine-1-carboxylate (710 mg, 1.52 mmol), and TFA (5.8 mL, 76 mmol) in DCM (6 mL). Yield (541 mg, 97%).

UPLC/ELSD: RT = 1.23 min. MS (ES): *m/z* (MH⁺) 367.70 for C₂₄H₅₀N₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.32-1.90 (br. m, 11H); 1.66-1.14 (br. m, 33H); 0.90 (t, 6H).

Step 3: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(3-(2-(dinonylamino)ethyl)pyrrolidin-1-yl)ethan-1-one



Chemical Formula: C₅₅H₁₁₂N₄O

Molecular Weight: 845.53


[00773] In the same manner as Step 11 for Compound 11, 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(3-(2-(dinonylamino)ethyl)pyrrolidin-1-yl)ethan-1-one was synthesized from *N*-nonyl-*N*-(2-(pyrrolidin-3-yl)ethyl)nonan-1-amine (250 mg, 0.68 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (308 mg, 0.62 mmol), iPr₂EtN (0.24 mL, 1.4 mmol), and T3P (50% EtOAc solution, 1.1 mL, 1.9 mmol) in THF (10 mL). Yield (100 mg, 19%).

UPLC/ELSD: RT = 3.17 min. MS (ES): *m/z* (MH⁺) 846.20 for C₅₅H₁₁₂N₄O

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.82-2.90 (br. m, 6H); 2.74-1.94 (br. m, 16H); 1.83-1.00 (br. m, 75H); 0.90 (t, 15H).

BN: Compound 70: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(3-((dinonylamino)methyl)pyrrolidin-1-yl)ethan-1-one

Step 1: *tert*-Butyl 3-((dinonylamino)methyl)pyrrolidine-1-carboxylate



Chemical Formula: C₂₈H₅₆N₂O₂


Molecular Weight: 452.77

[00774] In the same manner as Step 1 for Compound 49, *tert*-butyl 3-((dinonylamino)methyl)pyrrolidine-1-carboxylate was synthesized from *tert*-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate (2.0 g, 10.0 mmol), 1-bromononane (2.07 g, 10.0 mmol), K₂CO₃ (1.39 g, 10.0 mmol), and KI (166 mg, 1.00 mmol) in MeCN (100 mL). Yield (1.16 g, 26%).

UPLC/ELSD: RT = 2.17 min. MS (ES): m/z (MH $^+$) 453.72 for C₂₉H₅₈N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.72-2.90 (br. m, 4H); 2.36 (br. m, 6H); 2.04-1.04 (br. m, 40H); 0.90 (t, 6H).

5 Step 2: *N*-Nonyl-*N*-(pyrrolidin-3-ylmethyl)nonan-1-amine



Chemical Formula: C₂₃H₄₈N₂


Molecular Weight: 352.65

[00775] In the same manner as Step 4 for Compound 11, *N*-nonyl-*N*-(pyrrolidin-3-ylmethyl)nonan-1-amine was synthesized from *tert*-butyl 3-((dinonylamino)methyl)pyrrolidine-1-carboxylate (1.16 g, 2.56 mmol), and TFA (9.8 mL, 128 mmol) in DCM (10 mL). Yield (900 mg, 99%).

UPLC/ELSD: RT = 1.17 min. MS (ES): m/z (MH $^+$) 353.66 for C₂₃H₄₈N₂

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.33-2.23 (br. m, 10H); 1.99 (br, 1H); 1.65-1.00 (br. m, 31H); 0.90 (t, 6H).

Step 3: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(3-((dinonylamino)methyl)pyrrolidin-1-yl)ethan-1-one



20 Chemical Formula: C₅₄H₁₁₀N₄O

Molecular Weight: 831.50


[00776] In the same manner as Step 11 for Compound 11, 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(3-((dinonylamino)methyl)pyrrolidin-1-yl)ethan-1-one was synthesized from *N*-nonyl-*N*-(pyrrolidin-3-ylmethyl)nonan-1-amine (200 mg, 0.57 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (256 mg, 0.52 mmol), iPr₂EtN (0.198 mL, 1.14 mmol), and T3P (50% EtOAc solution, 0.92 mL, 1.56 mmol). Yield (114 mg, 27%).

UPLC/ELSD: RT = 3.22 min. MS (ES): m/z (MH $^+$) 832.26 for C₅₄H₁₁₀N₄O

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.79-2.96 (br. m, 6H); 2.75-2.18 (br. m, 16H); 2.12-1.01 (br. m, 73H); 0.90 (t, 15H).

BO: Compound 72: Dipentyl 4,4'-(2-(4-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-2-oxoethyl)azanediyl)dibutyrate

5 Step 1: Dipentyl 4,4'-(2-(*tert*-butoxy)-2-oxoethyl)azanediyl)dibutyrate



Chemical Formula: C₂₄H₄₅NO₆

Molecular Weight: 443.63


[00777] To a mixture of *tert*-butyl glycine (200 mg, 1.52 mmol) and pentyl 4-

10 bromobutanoate (759 mg, 3.2 mmol) in MeCN (30 mL) was added K₂CO₃ (637 mg, 4.6 mmol) and KI (51 mg, 0.30 mmol) and the mixture was allowed to stir at 82 °C for 12 hours. The suspension was cooled to RT, filtered over a pad of celite rinsing with EtOAc, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% EtOAc/hexanes) provided dipentyl 4,4'-(2-(*tert*-butoxy)-2-oxoethyl)azanediyl)dibutyrate (230 mg, 34%).

15 UPLC/ELSD: RT = 1.54 min. MS (ES): *m/z* (MH⁺) 444.61 for C₂₄H₄₅NO₆

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.07 (t, 4H); 3.22 (s, 2H); 2.63 (t, 4H); 2.36 (t, 4H); 1.77 (m, 4H); 1.64 (m, 4H); 1.47 (s, 9H); 1.35 (br. m, 8H); 0.93 (t, 6H).

Step 2: Bis(4-oxo-4-(pentyloxy)butyl)glycine



20

Chemical Formula: C₂₀H₃₇NO₆

Molecular Weight: 387.52


[00778] In the same manner as Step 4 for Compound 44, bis(4-oxo-4-

(pentyloxy)butyl)glycine was synthesized from dipentyl 4,4'-(2-(*tert*-butoxy)-2-oxoethyl)azanediyl)dibutyrate (230 mg, 0.52 mmol), and TFA (2 mL, 26 mmol) in DCM (2 mL). Yield (200 mg, 99%).

UPLC/ELSD: RT = 0.80 min. MS (ES): *m/z* (MH⁺) 388.51 for C₂₀H₃₇NO₆

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.05 (t, 4H); 3.10 (s, 2H); 2.58 (m, 4H); 2.32 (t, 4H); 1.80 (br. m, 4H); 1.63 (br. m, 4H); 1.32 (br. m, 8H); 0.92 (t, 6H).

Step 3: Dipentyl 4,4'-(2-(4-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-2-oxoethyl)azanediyl)dibutyrate



Chemical Formula: C₅₅H₁₀₇N₅O₆

Molecular Weight: 934.49


[00779] In the same manner as Step 7 for Compound 44, dipentyl 4,4'-(2-(4-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperazin-1-yl)-2-oxoethyl)azanediyl)dibutyrate was synthesized from bis(4-oxo-4-(pentyloxy)butyl)glycine (200 mg, 0.52 mmol), 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(piperazin-1-yl)ethan-1-one (265 mg, 0.47 mmol), iPr₂EtN (180 μL, 1.03 mmol) and T3P (50% EtOAc solution, 838 μL, 1.41 mmol) in THF (10 mL). Yield (250 mg, 57%).

UPLC/ELSD: RT = 2.85 min. MS (ES): *m/z* (MH⁺) 935.26 for C₅₅H₁₀₇N₅O₆

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 4H); 3.78-3.46 (br. m, 8H); 3.34 (br. m, 4H); 2.72-2.24 (br. m, 18H); 1.78 (m, 4H); 1.64 (m, 4H); 1.50-1.16 (br. m, 50H); 0.91 (m, 15H).

BP: Compound 73: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(3-(2-(dinonylamino)ethyl)piperidin-1-yl)ethan-1-one

Step 1: *tert*-Butyl 3-(2-(dinonylamino)ethyl)piperidine-1-carboxylate




Chemical Formula: C₃₀H₆₀N₂O₂

Molecular Weight: 480.82

[00780] In the same manner as Step 1 for Compound 49, *tert*-butyl 3-(2-(dinonylamino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 3-(2-aminoethyl)piperidine-1-carboxylate (1.00 g, 4.38 mmol), 1-bromononane (907 mg, 4.38

mmol), K_2CO_3 (610 mg, 4.38 mmol), and KI (73 mg, 0.44 mmol) in $MeCN$ (50 mL). Yield (514 mg, 24%). 1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.12-2.24 (br. m, 10H); 1.92-1.00 (br. m, 44H); 0.90 (t, 6H).

N-Nonyl-*N*-(2-(piperidin-3-yl)ethyl)nonan-1-amine



5

Chemical Formula: $C_{25}H_{52}N_2$


Molecular Weight: 380.71

[00781] In the same manner as Step 4 for Compound 11, *N*-nonyl-*N*-(2-(piperidin-3-yl)ethyl)nonan-1-amine was synthesized from *tert*-butyl 3-(2-(dinonylamino)ethyl)piperidine-1-carboxylate (514 mg, 1.07 mmol), and TFA (4.1 mL, 53 mmol) in DCM (4 mL). Yield (378 mg, 93%).

UPLC/ELSD: RT = 1.27 min. MS (ES): m/z (MH^+) 381.62 for $C_{25}H_{52}N_2$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 3.12-1.95 (br. m, 11H); 1.93-0.98 (br. m, 35H); 0.90 (t, 6H).

15 2-((2-(Dinonylamino)ethyl)(nonyl)amino)-1-(3-(2-(dinonylamino)ethyl)piperidin-1-yl)ethan-1-one



Chemical Formula: $C_{56}H_{114}N_4O$

Molecular Weight: 859.56


20 [00782] In the same manner as Step 11 for Compound 11, 2-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(3-(2-(dinonylamino)ethyl)piperidin-1-yl)ethan-1-one was synthesized from *N*-nonyl-*N*-(2-(piperidin-3-yl)ethyl)nonan-1-amine (250 mg, 0.66 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (297 mg, 0.60 mmol), iPr_2EtN (0.23 mL, 1.3 mmol), and T3P (50% $EtOAc$ solution, 1.06 mL, 1.8 mmol) in THF (10 mL). Yield (136 mg, 27%).

25 UPLC/ELSD: RT = 3.22 min. MS (ES): m/z (MH^+) 860.39 for $C_{56}H_{114}N_4O$

1H -NMR (300 MHz, $CDCl_3$) δ : ppm 4.56-4.01 (br. m, 2H); 3.48-2.20 (br. m, 20H), 1.99-1.00 (br. m, 77H); 0.90 (t, 15H).

BQ: Compound 71: Pentyl 4-(nonyl(2-(4-(*N*-nonyl-*N*-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)glycyl)piperazin-1-yl)-2-oxoethyl)amino)butanoate

Step 1: *tert*-Butyl 4-(*N*-nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycyl)piperazine-1-carboxylate



5

Chemical Formula: C₂₉H₅₅N₃O₅

Molecular Weight: 525.78


[00783] In the same manner as Step 3 for Compound 11, *tert*-butyl 4-(*N*-nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycyl)piperazine-1-carboxylate was synthesized from *N*-nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycine (480 mg, 1.34 mmol), 1-boc-piperazine (275 mg, 1.48 mmol), iPr₂EtN (5.14 μ L, 2.95 mmol) and T3P (50% EtOAc solution, 2.40 mL, 4.03 mmol) in THF (15 mL). Yield (700 mg, 99%).

UPLC/ELSD: RT = 1.90 min. MS (ES): *m/z* (MH⁺) 526.79 for C₂₉H₅₅N₃O₅

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.08 (t, 2H); 3.70-3.10 (br. m, 14H); 2.45 (t, 2H); 2.13 (br. m, 2H); 2.00-1.00 (br. m, 29H); 0.91 (br. m, 6H).

15

Step 2: Pentyl 4-(nonyl(2-oxo-2-(piperazin-1-yl)ethyl)amino)butanoate



Chemical Formula: C₂₄H₄₇N₃O₃


Molecular Weight: 425.66

[00784] In the same manner as Step 4 for Compound 11, pentyl 4-(nonyl(2-oxo-2-(piperazin-1-yl)ethyl)amino)butanoate was synthesized from *tert*-butyl 4-(*N*-nonyl-*N*-(4-oxo-4-(pentyloxy)butyl)glycyl)piperazine-1-carboxylate (700 mg, 1.33 mmol), and TFA (5.1 mL, 66.6 mmol) in DCM (5 mL). Yield (560 mg, 99%).

UPLC/ELSD: RT = 0.77 min. MS (ES): *m/z* (MH⁺) 426.65 for C₂₄H₄₇N₃O₃

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.07 (t, 2H); 3.59 (br. m, 4H); 3.28 (s, 2H); 2.86 (br. m, 4H); 2.50 (br. m, 4H); 2.33 (t, 2H); 2.05 (br. 1H); 1.77 (m, 2H); 1.63 (m, 2H); 1.30 (br. m, 18H); 0.91 (m, 6H).

Step 3: Pentyl 4-((2-hydroxyethyl)(nonyl)amino)butanoate

Chemical Formula: C₂₀H₄₁NO₃

Molecular Weight: 343.55


5

[00785] In the same manner as Step 1 for Compound 18, pentyl 4-((2-hydroxyethyl)(nonyl)amino)butanoate was synthesized from 2-(nonylamino)ethan-1-ol (350 mg, 1.87 mmol), pentyl 4-bromobutanoate (487 mg, 2.06 mmol), K₂CO₃ (572 mg, 4.11 mmol), and KI (31 mg, 0.19 mmol) in MeCN (40 mL). Yield (427 mg, 66%).

10 UPLC/ELSD: RT = 1.25 min. MS (ES): *m/z* (MH⁺) 344.55 for C₂₀H₄₁NO₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (t, 2H); 3.61 (t, 2H); 2.67 (t, 2H); 2.56 (m, 4H); 2.36 (t, 2H); 1.85 (m, 2H); 1.65 (m, 2H); 1.49 (m, 2H); 1.42-1.18 (br. m, 16H); 0.91 (m, 6H).

Step 4: Pentyl 4-((2-chloroethyl)(nonyl)amino)butanoate

Chemical Formula: C₂₀H₄₀ClNO₂

Molecular Weight: 362.00

15


[00786] In the same manner as Step 2 for Compound 18, pentyl 4-((2-chloroethyl)(nonyl)amino)butanoate was synthesized from pentyl 4-((2-hydroxyethyl)(nonyl)amino)butanoate (427 mg, 1.27 mmol), methanesulfonyl chloride (120 μL, 1.55 mmol), and triethylamine (225 μL, 1.62 mmol) in DCM (8 mL). Yield (448 mg, 99%).

20 UPLC/ELSD: RT = 1.52 min. MS (ES): *m/z* (MH⁺) 362.51 for C₂₀H₄₀ClNO₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.07-3.71 (br. m, 4H); 3.45-2.76 (br. m, 6H); 2.30 (br. m, 2H); 2.24-1.05 (br. m, 22H); 0.82 (br. m, 6H).

25

Step 5: Pentyl 4-((2-((2-(*tert*-butoxy)-2-oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)butanoate



Chemical Formula: C₃₅H₇₀N₂O₄

Molecular Weight: 582.96


[00787] In the same manner as Step 4 for Compound 46, pentyl 4-((2-((2-(*tert*-butoxy)-2-

5 oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)butanoate was synthesized from *tert*-butyl nonylglycinate (338 mg, 1.31 mmol), pentyl 4-((2-chloroethyl)(nonyl)amino)butanoate (527 mg, 1.46 mmol), K₂CO₃ (402 mg, 2.89 mmol), and KI (22 mg, 0.13 mmol) in MeCN (30 mL). Yield (200 mg, 26%).

UPLC/ELSD: RT = 3.03 min. MS (ES): *m/z* (MH⁺) 583.95 for C₃₅H₇₀N₂O₄

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.07 (t, 2H); 3.27 (s, 2H); 2.76-2.24 (br. m, 12H); 1.85-1.10 (br. m, 45H); 0.90 (m, 9H).

Step 6: *N*-Nonyl-*N*-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)glycine



15 Chemical Formula: C₃₁H₆₂N₂O₄

Molecular Weight: 526.85


[00788] In the same manner as Step 5 for Compound 46, *N*-nonyl-*N*-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)glycine was synthesized from pentyl 4-((2-((2-(*tert*-butoxy)-2-

20 oxoethyl)(nonyl)amino)ethyl)(nonyl)amino)butanoate (200 mg, 0.34 mmol), and TFA (1.31 mL, 17.2 mmol), in DCM (2 mL). Yield (160 mg, 89%).

UPLC/ELSD: RT = 2.39 min. MS (ES): *m/z* (MH⁺) 527.77 for C₃₁H₆₂N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (t, 2H); 3.27 (s, 2H); 2.94-2.74 (br. m, 6H); 2.61 (t, 2H); 2.37 (m, 2H); 2.15-1.90 (br. m, 2H); 1.80-1.05 (br. m, 36H); 0.90 (m, 9H).

25 Step 7: Pentyl 4-(nonyl(2-(4-(*N*-nonyl-*N*-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)glycyl)piperazin-1-yl)-2-oxoethyl)amino)butanoate

Chemical Formula: C₅₅H₁₀₇N₅O₆


Molecular Weight: 934.49

5 [00789] In the same manner as Step 11 for Compound 11, pentyl 4-(nonyl(2-(4-(N-nonyl-N-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)glycyl)piperazin-1-yl)-2-oxoethyl)amino)butanoate was synthesized from pentyl 4-(nonyl(2-oxo-2-(piperazin-1-yl)ethyl)amino)butanoate (142 mg, 0.33 mmol), N-nonyl-N-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)glycine (160 mg, 0.30 mmol), iPr₂EtN (116 μ L, 0.67 mmol), and 10 T3P (50% EtOAc solution, 542 μ L, 0.91 mmol). Yield (53 mg, 19%).

UPLC/ELSD: RT = 2.79 min. MS (ES): *m/z* (MH⁺) 935.34 for C₅₅H₁₀₇N₅O₆

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.08 (t, 4H); 3.78-3.53 (br. m, 8H); 3.32 (br. m, 4H); 2.76-2.24 (br. m, 18H); 1.87-1.10 (br. m, 58H); 0.91 (br. m, 15H).

15 BR: Compound 80: Pentyl 4-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-3-yl)ethyl)(nonyl)amino)butanoate

Step 1: *tert*-butyl 3-(2-(nonylamino)ethyl)piperidine-1-carboxylateChemical Formula: C₂₁H₄₂N₂O₂


Molecular Weight: 354.58

20 [00790] In the same manner as Step 1 for Compound 49, *tert*-butyl 3-(2-(nonylamino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 3-(2-aminoethyl)piperidine-1-carboxylate (1.00 g, 4.38 mmol), 1-bromononane (907 mg, 4.38 mmol), K₂CO₃ (610 mg, 4.38 mmol), and KI (73 mg, 0.44 mmol) in MeCN (50 mL). Yield (474 mg, 31%).

UPLC/ELSD: RT = 1.23 min. MS (ES): m/z (MH⁺) 355.58 for C₂₁H₄₂N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.88 (br, 2H); 3.00-2.43 (br. m, 6H); 1.92-0.97 (br. m, 30H), 0.90 (t, 3H).

5 Step 2: *tert*-Butyl 3-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₅₈N₂O₄


Molecular Weight: 510.80

[00791] In the same manner as Step 1 for Compound 57, *tert*-butyl 3-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 3-(2-(nonylamino)ethyl)piperidine-1-carboxylate (474 mg, 1.34 mmol), pentyl 4-bromobutanoate (380 mg, 1.6 mmol), K₂CO₃ (223 mg, 1.60 mmol) and KI (44 mg, 0.27 mmol) in MeCN (15 mL). Yield (492 mg, 72%).

UPLC/ELSD: RT = 2.09 min. MS (ES): m/z (MH⁺) 511.70 for C₃₀H₅₈N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.90 (br. m, 2H); 2.87-2.22 (br. m, 10H); 1.91-1.00 (br. m, 38H); 0.91 (m, 6H).

Step 3: Pentyl 4-(nonyl(2-(piperidin-3-yl)ethyl)amino)butanoate



Chemical Formula: C₂₅H₅₀N₂O₂


Molecular Weight: 410.69

[00792] In the same manner as Step 4 for Compound 11, pentyl 4-(nonyl(2-(piperidin-3-yl)ethyl)amino)butanoate was synthesized from *tert*-butyl 3-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)piperidine-1-carboxylate (492 mg, 0.96 mmol), and TFA (3.7 mL, 48 mmol) in DCM (4 mL). Yield (390 mg, 99%).

UPLC/ELSD: RT = 0.85 min. MS (ES): m/z (MH⁺) 411.72 for C₂₅H₅₀N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.03 (br. m, 2H); 2.66-2.18 (br. m, 10H); 2.18-0.98 (br. m, 30H); 0.91 (m, 6H).

Step 4: Penty 4-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-3-yl)ethyl)(nonyl)amino)butanoate



Chemical Formula: C₅₆H₁₁₂N₄O₃

Molecular Weight: 889.54

[00793] In the same manner as Step 11 for Compound 11, penty 4-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)piperidin-3-yl)ethyl)(nonyl)amino)butanoate was synthesized from penty 4-(nonyl(2-(piperidin-3-yl)ethyl)amino)butanoate (250 mg, 0.61 mmol), N-(2-(dinonylamino)ethyl)-N-nonylglycine (275 mg, 0.55 mmol), iPr₂EtN (0.21 mL, 1.2 mmol), and T3P (50% EtOAc solution, 0.98 mL, 1.7 mmol). Yield (96 mg, 20%).


UPLC/ELSD: RT = 3.08 min. MS (ES): *m/z* (MH⁺) 890.32 for C₅₆H₁₁₂N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.55-4.01 (br. m, 4H); 3.48-2.21 (br. m, 22H); 1.95-1.00 (br. m, 71H); 0.90 (m, 15H).

BS: Compound 81: Penty 4-((2-(1-(N-(2-(dinonylamino)ethyl)-N-nonylglycyl)pyrrolidin-3-yl)ethyl)(nonyl)amino)butanoate

20

Step 1: *tert*-Butyl 3-(2-(nonylamino)ethyl)pyrrolidine-1-carboxylate



Chemical Formula: C₂₀H₄₀N₂O₂


Molecular Weight: 340.55

[00794] In the same manner as Step 1 for Compound 49, *tert*-butyl 3-(2-(nonylamino)ethyl)pyrrolidine-1-carboxylate was synthesized from *tert*-butyl 3-(2-aminoethyl)pyrrolidine-1-carboxylate (1.25 g, 5.47 mmol), 1-bromononane (1.13 g, 5.47 mmol), K₂CO₃ (757 mg, 5.47 mmol), and KI (91 mg, 0.55 mmol) in MeCN (100 mL). Yield (420 mg, 23%).

UPLC/ELSD: RT = 1.08 min. MS (ES): m/z (MH⁺) 341.52 for C₂₀H₄₀N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ : ppm 3.68-1.91 (br. m, 9H); 1.71-1.12 (br. m, 28H); 0.90 (t, 3H).

Step 2: *tert*-Butyl 3-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)pyrrolidine-1-carboxylate



Chemical Formula: C₂₉H₅₆N₂O₄


Molecular Weight: 496.78

[00795] In the same manner as Step 1 for Compound 57, *tert*-butyl 3-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)pyrrolidine-1-carboxylate was synthesized from *tert*-butyl 3-(2-(nonylamino)ethyl)pyrrolidine-1-carboxylate (420 mg, 1.23 mmol), penty 4-bromobutanoate (321 mg, 1.36 mmol), K₂CO₃ (187 mg, 1.36 mmol), and KI (41 mg, 0.25 mmol). Yield (390 mg, 64%).

UPLC/ELSD: RT = 1.98 min. MS (ES): m/z (MH⁺) 497.67 for C₂₉H₅₆N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.08 (t, 2H); 3.50 (br. m, 2H); 3.34-2.76 (br. m, 2H); 2.52-1.87 (br. m, 10H); 1.87-1.02 (br. m, 34H); 0.91 (t, 6H).

Step 3: Penty 4-(nonyl(2-(pyrrolidin-3-yl)ethyl)amino)butanoate



Chemical Formula: C₂₄H₄₈N₂O₂


Molecular Weight: 396.66

[00796] In the same manner as Step 4 for Compound 11, penty 4-(nonyl(2-(pyrrolidin-3-yl)ethyl)amino)butanoate was synthesized from *tert*-butyl 3-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)pyrrolidine-1-carboxylate (390 mg, 0.79 mmol), and TFA (3.0 mL, 40 mmol) in DCM (3 mL). Yield (298 mg, 96%).

25 UPLC/ELSD: RT = 0.81 min. MS (ES): m/z (MH⁺) 397.62 for C₂₄H₄₈N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.08 (t, 2H); 3.20-2.82 (br. m, 4H); 2.58-2.24 (br. m, 8H); 2.11-1.11 (br. m, 28H); 0.91 (m, 6H).

Step 4: Pentyl 4-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)pyrrolidin-3-yl)ethyl)(nonyl)amino)butanoate



Chemical Formula: C₅₅H₁₁₀N₄O₃

Molecular Weight: 875.51

5

[00797] In the same manner as Step 11 for Compound 11, pentyl 4-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)pyrrolidin-3-yl)ethyl)(nonyl)amino)butanoate was synthesized from pentyl 4-(nonyl(2-(pyrrolidin-3-yl)ethyl)amino)butanoate (202 mg, 0.51 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (230 mg, 0.46 mmol), iPr₂EtN (0.177 mL, 1.0 mmol), and T3P (50% EtOAc solution, 0.82 mL, 1.4 mmol) in THF (10 mL). Yield (109 mg, 27%).


UPLC/ELSD: RT = 3.06 min. MS (ES): *m/z* (MH⁺) 876.30 for C₅₅H₁₁₀N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.83-2.85 (br. m, 7H); 2.78-1.88 (br. m, 19H); 1.83-1.14 (br. m, 67H); 0.90 (m, 15H).

15

BT: Compound 82: Pentyl 4-(((1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)pyrrolidin-3-yl)methyl)(nonyl)amino)butanoate

Step 1: *tert*-Butyl 3-((nonylamino)methyl)pyrrolidine-1-carboxylate



Chemical Formula: C₁₉H₃₈N₂O₂


Molecular Weight: 326.53

20

[00798] In the same manner as Step 1 for Compound 49, *tert*-butyl 3-((nonylamino)methyl)pyrrolidine-1-carboxylate was synthesized from *tert*-butyl 3-(aminomethyl)pyrrolidine-1-carboxylate (2.0 g, 10.0 mmol), 1-bromononane (2.07 g, 10.0 mmol), K₂CO₃ (1.39 g, 10.0 mmol), and KI (166 mg, 1.00 mmol) in MeCN (100 mL). Yield (1.53 g, 47%).

UPLC/ELSD: RT = 0.92 min. MS (ES): *m/z* (MH⁺) 327.54 for C₁₉H₃₈N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.69-1.79 (br. m, 9H); 1.74-1.13 (br. m, 26H); 0.89 (t, 3H).

Step 2: *tert*-Butyl 3-((nonyl(4-oxo-4-(pentyloxy)butyl)amino)methyl)pyrrolidine-1-carboxylateChemical Formula: C₂₈H₅₄N₂O₄


Molecular Weight: 482.75

5 [00799] In the same manner as Step 1 for Compound 57, *tert*-butyl 3-((nonyl(4-oxo-4-(pentyloxy)butyl)amino)methyl)pyrrolidine-1-carboxylate was synthesized from *tert*-butyl 3-((nonylamino)methyl)pyrrolidine-1-carboxylate (500 mg, 1.53 mmol), pentyl 4-bromobutanoate (400 mg, 1.68 mmol), K₂CO₃ (423 mg, 3.07 mmol), and KI (51 mg, 0.31 mmol) in MeCN (100 mL). Yield (233 mg, 32%).

10 UPLC/ELSD: RT = 1.85 min. MS (ES): *m/z* (MH⁺) 483.65 for C₂₈H₅₄N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.59-2.91 (br. m, 4H); 2.49-1.83 (br. m, 10H); 1.83-1.13 (br. m, 32H); 0.91 (m, 6H).

15 Step 3: Pentyl 4-(nonyl(pyrrolidin-3-ylmethyl)amino)butanoate

Chemical Formula: C₂₃H₄₆N₂O₂

Molecular Weight: 382.63


20 [00800] In the same manner as Step 4 for Compound 11, pentyl 4-(nonyl(pyrrolidin-3-ylmethyl)amino)butanoate was synthesized from *tert*-butyl 3-((nonyl(4-oxo-4-(pentyloxy)butyl)amino)methyl)pyrrolidine-1-carboxylate (233 mg, 0.48 mmol), and TFA (1.84 mL, 24 mmol) in DCM (2 mL). Yield (179 mg, 97%).

25 UPLC/ELSD: RT = 0.70 min. MS (ES): *m/z* (MH⁺) 383.51 for C₂₃H₄₆N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.11-2.81 (br. m, 3H); 2.67-1.51 (br. m, 16H);

1.51-1.03 (br. m, 19H); 0.91 (m, 6H).

Step 4: Pentyl 4-(((1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)pyrrolidin-3-yl)methyl)(nonyl)amino)butanoate



Chemical Formula: C₅₄H₁₀₈N₄O₃

Molecular Weight: 861.48

[00801] In the same manner as Step 11 for Compound 11, pentyl 4-((1-(N-(2-

5 (dinoylamino)ethyl)-N-nonylglycyl)pyrrolidin-3-yl)methyl)(nonyl)amino)butanoate was synthesized from pentyl 4-(nonyl(pyrrolidin-3-ylmethyl)amino)butanoate (179 mg, 0.47 mmol), N-(2-(dinoylamino)ethyl)-N-nonylglycine (211 mg, 0.43 mmol), iPr₂EtN (163 µL, 0.95 mmol), and T3P (50% EtOAc solution, 0.76 mL, 1.1 mmol). Yield (88 mg, 24%).


UPLC/ELSD: RT = 3.05 min. MS (ES): *m/z* (MH⁺) 862.28 for C₅₄H₁₀₈N₄O₃

10 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.76-2.17 (br. m, 24H); 2.12-1.05 (br. m, 67H); 0.90 (m, 15H).

BU: Compound 83: Pentyl 4-((2-(1-(N-(2-(dinoylamino)ethyl)-N-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)butanoate

15

Step 1: *tert*-Butyl 4-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)piperidine-1-carboxylate



Chemical Formula: C₃₀H₅₈N₂O₄


Molecular Weight: 510.80

20 **[00802]** In the same manner as Step 1 for Compound 57, *tert*-butyl 4-(2-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)ethyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(2-(nonyl)amino)ethyl)piperidine-1-carboxylate (500 mg, 1.41 mmol), pentyl 4-bromobutanoate (368 mg, 1.55 mmol), K₂CO₃ (390 mg, 2.82 mmol), and KI (23 mg, 0.14 mmol) in MeCN (100 mL). Yield (487 mg, 68%).

25 UPLC/ELSD: RT = 2.03 min. MS (ES): *m/z* (MH⁺) 511.57 for C₃₀H₅₈N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (m, 4H); 2.69 (m, 2H); 2.51-2.25 (br. m, 8H); 1.83-1.55 (br. m, 6H); 1.53-1.02 (br. m, 32H); 0.91 (m, 6H).

Step 2: Pentyl 4-(nonyl(2-(piperidin-4-yl)ethyl)amino)butanoate

Chemical Formula: C₂₅H₅₀N₂O₂


Molecular Weight: 410.69

5

[00803] In the same manner as Step 4 for Compound 11, pentyl 4-(nonyl(2-(piperidin-4-yl)ethyl)amino)butanoate was synthesized from *tert*-butyl 4-(2-(nonyl(4-oxo-4-pentyloxy)butyl)amino)ethyl)piperidine-1-carboxylate (487 mg, 0.953 mmol), and TFA (3.6 mL, 48 mmol) in DCM (4 mL). Yield (386 mg, 98%).

10 UPLC/ELSD: RT = 0.87 min. MS (ES): *m/z* (MH⁺) 411.43 for C₂₅H₅₀N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.07 (m, 2H); 2.60 (m, 2H); 2.50-2.28 (br. m, 8H); 2.03 (br, 1H); 1.86-1.55 (br. m, 6H); 1.52-1.02 (br. m, 23H); 0.91 (m, 6H).

Step 3: Pentyl 4-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)butanoateChemical Formula: C₅₆H₁₁₂N₄O₃

Molecular Weight: 889.54


[00804] In the same manner as Step 11 for Compound 11, pentyl 4-((2-(1-(*N*-(2-(dinonylamino)ethyl)-*N*-nonylglycyl)piperidin-4-yl)ethyl)(nonyl)amino)butanoate was synthesized from pentyl 4-(nonyl(2-(piperidin-4-yl)ethyl)amino)butanoate (351 mg, 0.855 mmol), *N*-(2-(dinonylamino)ethyl)-*N*-nonylglycine (467 mg, 0.941 mmol), iPr₂EtN (328 μL, 1.88 mmol), and T3P (50% EtOAc solution, 1.53 mL, 2.56 mmol) in THF (15 mL). Yield (192 mg, 25%).

25 UPLC/ELSD: RT = 3.00 min. MS (ES): *m/z* (MH⁺) 890.13 for C₅₆H₁₁₂N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.61-4.14 (br. m, 2H); 4.08 (t, 2H); 3.40-2.24 (br. m, 22H); 1.86-0.99 (br. m, 71H); 0.90 (m, 15H).

5 BV: Compound 84: Petyl 4-((3-(1-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperidin-4-yl)propyl)(nonyl)amino)butanoate

Step 1: *tert*-Butyl 4-(3-(nonylamino)propyl)piperidine-1-carboxylate




Chemical Formula: C₂₂H₄₄N₂O₂

10 Molecular Weight: 368.61

[00805] In the same manner as Step 1 for Compound 49, *tert*-butyl 4-(3-(nonylamino)propyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(3-aminopropyl)piperidine-1-carboxylate (2.50 g, 10.3 mmol), 1-bromononane (2.14 g, 10.3 mmol), K₂CO₃ (2.85 g, 20.6 mmol), and KI (171 mg, 0.10 mmol) in MeCN (200 mL). Yield 15 (1.27 g, 33%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (m, 2H), 2.69 (br. m, 6H); 1.79-0.98 (br. m, 32H); 0.89 (t, 3H).

20 Step 2: *tert*-Butyl 4-(3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propyl)piperidine-1-carboxylate



Chemical Formula: C₃₁H₆₀N₂O₄


Molecular Weight: 524.83

[00806] In the same manner as Step 1 for Compound 57, *tert*-butyl 4-(3-(nonyl(4-oxo-4-(pentyloxy)butyl)amino)propyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(3-(nonylamino)propyl)piperidine-1-carboxylate (500 mg, 1.36 mmol), petyl 4-bromobutanoate (354 mg, 1.49 mmol), K₂CO₃ (375 mg, 2.71 mmol), and KI (23 mg, 0.14 mmol) in MeCN (20 mL). Yield (624 mg, 88%).

UPLC/ELSD: RT = 2.12 min. MS (ES): *m/z* (MH⁺) 525.60 for C₃₁H₆₀N₂O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (br. m, 4H); 2.69 (m, 2H); 2.38 (br. m, 8H); 1.85-1.55 (br. m, 6H); 1.54-1.00 (br. m, 34H); 0.91 (m, 6H).

Step 3: Pentyl 4-(nonyl(3-(piperidin-4-yl)propyl)amino)butanoate



5

Chemical Formula: C₂₆H₅₂N₂O₂

Molecular Weight: 424.71

[00807] In the same manner as Step 4 for Compound 44, pentyl 4-(nonyl(3-(piperidin-4-yl)propyl)amino)butanoate was synthesized from *tert*-butyl 4-(3-(nonyl(4-oxo-4-


10 (pentyloxy)butyl)amino)propyl)piperidine-1-carboxylate (624 mg, 1.19 mmol), and TFA (4.5 mL, 60 mmol) in DCM (5 mL). Yield (467 mg, 92%).

UPLC/ELSD: RT = 0.94 min. MS (ES): *m/z* (MH⁺) 424.62 for C₂₆H₅₂N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.16 (m, 2H); 2.65 (m, 2H); 2.39 (br. m, 8H); 1.84-1.57 (br. m, 6H); 1.52-1.04 (br. m, 26H); 0.91 (m, 6H).

15

Step 4: Pentyl 4-((3-(1-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperidin-4-yl)propyl)(nonyl)amino)butanoate



Chemical Formula: C₅₈H₁₁₆N₄O₃

Molecular Weight: 917.59


[00808] In the same manner as Step 11 for Compound 11, pentyl 4-((3-(1-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperidin-4-yl)propyl)(nonyl)amino)butanoate was synthesized from pentyl 4-(nonyl(3-(piperidin-4-yl)propyl)amino)butanoate (259 mg, 0.61 mmol), 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid (343 mg, 0.67 mmol), iPr₂EtN (234 μL, 1.34 mmol), and T3P (50% EtOAc solution, 1.09 mL, 1.83 mmol) in THF (20 mL). Yield (270 mg, 48%).

UPLC/ELSD: RT = 2.85 min. MS (ES): *m/z* (MH⁺) 918.18 for C₅₈H₁₁₆N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.61 (m, 1H); 4.08 (t, 2H); 4.08 (m, 1H); 3.08-2.72 (br. m, 4H); 2.63-2.26 (br. m, 20H); 1.87-1.57 (br. m, 6H); 1.54-1.00 (br. m, 67H); 0.90 (m, 15H).

BW: Compound 85: 3-((2-(Dinylamino)ethyl)(nonyl)amino)-1-(4-(3-(dinylamino)propyl)piperidin-1-yl)propan-1-one

5 Step 1: *tert*-Butyl 4-(3-(dinylamino)propyl)piperidine-1-carboxylate



Chemical Formula: C₃₁H₆₂N₂O₂

Molecular Weight: 494.85


[00809] In the same manner as Step 1 for Compound 49 *tert*-butyl 4-(3-

10 (dinylamino)propyl)piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(3-aminopropyl)piperidine-1-carboxylate (2.50 g, 10.3 mmol), 1-bromononane (2.14 g, 10.3 mmol), K₂CO₃ (2.85 g, 20.6 mmol), and KI (171 mg, 0.10 mmol) in MeCN (200 mL). Yield (1.03 g, 20%).

UPLC/ELSD: RT = 2.46 min. MS (ES): *m/z* (MH⁺) 495.66 for C₃₁H₆₂N₂O₂

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.09 (br. m, 2H); 2.69 (br. m, 2H); 2.39 (br. m, 6H); 1.75-1.00 (br. m, 46H); 0.90 (t, 6H).

Step 2: *N*-Nonyl-*N*-(3-(piperidin-4-yl)propyl)nonan-1-amine



20 Chemical Formula: C₂₆H₅₄N₂

Molecular Weight: 394.73

[00810] In the same manner as Step 4 for Compound 11, *N*-nonyl-*N*-(3-(piperidin-4-


yl)propyl)nonan-1-amine was synthesized from *tert*-butyl 4-(3-(dinylamino)propyl)piperidine-1-carboxylate (1.03 g, 2.08 mmol), and TFA (8.0 mL, 104

25 mmol) in DCM (10 mL). Yield (778 mg, 95%).

UPLC/ELSD: RT = 1.31 min. MS (ES): *m/z* (MH⁺) 395.61 for C₂₆H₅₄N₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.18-2.50 (br. m, 4H); 2.40 (br. m, 6H); 1.70 (m, 2H); 1.58-1.03 (br. m, 36H); 0.90 (t, 6H).

30 Step 3: 3-((2-(Dinylamino)ethyl)(nonyl)amino)-1-(4-(3-(dinylamino)propyl)piperidin-1-yl)propan-1-one



Chemical Formula: C₅₈H₁₁₈N₄O

Molecular Weight: 887.61


5 [00811] In the same manner as Step 11 for Compound 11, 3-((2-(dinonylamino)ethyl)(nonyl)amino)-1-(4-(3-(dinonylamino)propyl)piperidin-1-yl)propan-1-one was synthesized from *N*-nonyl-*N*-(3-(piperidin-4-yl)propyl)nonan-1-amine (247 mg, 0.63 mmol), 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid (352 mg, 0.69 mmol), iPr₂EtN (240 μ L, 1.4 mmol), and T3P (50% EtOAc solution, 1.1 mL, 1.9 mmol) in THF (20 mL). Yield 10 (293 mg, 53%).

UPLC/ELSD: RT = 3.01 min. MS (ES): *m/z* (MH⁺) 888.08 for C₅₈H₁₁₈N₄O

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.61 (m, 1H); 3.86 (m, 1H); 2.99 (m, 1H); 2.82 (m, 2H); 2.61-2.29 (br. m, 19H); 1.75 (m, 2H); 1.60-1.00 (br. m, 77H); 0.89 (t, 15H).

15 BX: Compound 86: 3-((3-(4-(3-((2-(Dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)-3-oxopropyl)(nonyl)amino)propyl hexanoate

Step 1: 3-((3-(*tert*-Butoxy)-3-oxopropyl)(nonyl)amino)propyl hexanoate



20

Chemical Formula: C₂₅H₄₉NO₄


Molecular Weight: 427.67

25 [00812] In the same manner as Step 3 for Compound 44, 2-((3-(*tert*-butoxy)-3-oxopropyl)(nonyl)amino)ethyl heptanoate was synthesized from *tert*-butyl 3-(nonylamino)propanoate (750 mg, 2.76 mmol), 3-bromopropyl hexanoate (786 mg, 3.32 mmol), K₂CO₃ (764 mg, 5.53 mmol), and KI (46 mg, 0.28 mmol) in MeCN (100 mL). Yield (661 mg, 56%).

UPLC/ELSD: RT = 1.80 min. MS (ES): *m/z* (MH⁺) 428.49 for C₂₅H₄₉NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.12 (t, 2H); 2.73 (t, 2H); 2.56-2.24 (br. m, 8H); 1.77 (m, 2H); 1.64 (m, 2H); 1.55-1.10 (br. m, 27H); 0.91 (m, 6H).

Step 2: 3-((3-(Hexanoyloxy)propyl)(nonyl)amino)propanoic acid



5

Chemical Formula: C₂₁H₄₁NO₄

Molecular Weight: 371.56


[00813] In the same manner as Step 4 for Compound 44, 3-((3-(hexanoyloxy)propyl)(nonyl)amino)propanoic acid was synthesized from 3-((3-(*tert*-butoxy)-3-oxopropyl)(nonyl)amino)propyl hexanoate (661 mg, 1.55 mmol), and TFA (5.9 mL, 77 mmol) in DCM (6 mL). Yield (556 mg, 97%).

UPLC/ELSD: RT = 1.14 min. MS (ES): *m/z* (MH⁺) 372.31 for C₂₁H₄₁NO₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.13 (t, 2H); 2.84 (t, 2H); 2.72 (t, 2H); 2.62 (t, 2H); 2.46 (t, 2H); 2.31 (t, 2H); 1.90 (m, 2H); 1.72-1.10 (br. m, 20H); 0.90 (M, 6H).

15

Step 3: *tert*-Butyl 4-((3-(hexanoyloxy)propyl)(nonyl)amino)propanoyl)piperazine-1-carboxylate



Chemical Formula: C₃₀H₅₇N₃O₅


Molecular Weight: 539.80

[00814] In the same manner as Step 3 for Compound 11, *tert*-butyl 4-((3-(hexanoyloxy)propyl)(nonyl)amino)propanoyl)piperazine-1-carboxylate was synthesized from 3-((3-(hexanoyloxy)propyl)(nonyl)amino)propanoic acid (570 mg, 1.49 mmol), 1-boc-piperazine (334 mg, 1.80 mmol), iPr₂EtN (573 μL, 3.29 mmol), and T3P (50% EtOAc solution, 2.67 mL, 4.49 mmol) in THF (20 mL). Yield (635 mg, 79%).

UPLC/ELSD: RT = 1.85 min. MS (ES): *m/z* (MH⁺) 540.52 for C₃₀H₅₇N₃O₅

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.12 (t, 2H); 3.60 (m, 2H); 3.46 (br. m, 6H); 2.80 (m, 2H); 2.58-2.37 (br. m, 6H); 2.30 (t, 2H); 1.78 (m, 2H); 1.63 (m, 2H); 1.54-1.10 (br. m, 27H); 0.90 (m, 6H).

Step 4: 3-(Nonyl(3-oxo-3-(piperazin-1-yl)propyl)amino)propyl hexanoate

Chemical Formula: C₂₅H₄₉N₃O₃

Molecular Weight: 439.69


[00815] In the same manner as Step 4 for Compound 11, 3-(nonyl(3-oxo-3-(piperazin-1-yl)propyl)amino)propyl hexanoate was synthesized from *tert*-butyl 4-(3-((3-(hexanoyloxy)propyl)(nonyl)amino)propanoyl)piperazine-1-carboxylate (635 mg, 1.18 mmol), and TFA (4.5 mL, 59 mmol) in DCM (5 mL). Yield (510 mg, 99%).

10 UPLC/ELSD: RT = 0.72 min. MS (ES): *m/z* (MH⁺) 440.47 for C₂₅H₄₉N₃O₃

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.12 (t, 2H); 3.60 (m, 2H) 3.46 (m, 2H); 2.85 (br. m, 6H); 2.49 (br. m, 6H); 2.30 (t, 2H); 1.80 (m, 2H); 1.64 (m, 2H); 1.52-1.10 (br. m, 19H); 0.91 (m, 6H).

Step 5: 3-((3-(4-(3-((2-(Dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)-3-

15 oxopropyl)(nonyl)amino)propyl hexanoate

Chemical Formula: C₅₇H₁₁₃N₅O₄

Molecular Weight: 932.56

[00816] In the same manner as Step 11 for Compound 11, 3-((3-(4-(3-((2-


20 (dinonylamino)ethyl)(nonyl)amino)propanoyl)piperazin-1-yl)-3-oxopropyl)(nonyl)amino)propyl hexanoate was synthesized from 3-(nonyl(3-oxo-3-(piperazin-1-yl)propyl)amino)propyl hexanoate (154 mg, 0.351 mmol), 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid (197 mg, 0.386 mmol), iPr₂EtN (134 μL, 0.77 mmol), and T3P (50% EtOAc solution, 616 μL, 1.05 mmol) in THF (10 mL). Yield (69 mg, 21%).

25 UPLC/ELSD: RT = 2.70 min. MS (ES): *m/z* (MH⁺) 933.10 for C₅₇H₁₁₃N₅O₄

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.12 (t, 2H); 3.72-3.40 (br. m, 8H); 2.81 (br. m, 4H); 2.61-2.36 (br. m, 18H); 2.30 (t, 2H); 1.78 (m, 2H); 1.64 (m, 2H); 1.54-1.06 (br. m, 60H); 0.90 (m, 15H).

BY: Compound 87: 3-((3-(1-(3-((2-(Dinonylamino)ethyl)(nonyl)amino)propanoyl)piperidin-4-yl)propyl)(nonyl)amino)propyl hexanoate

5 Step 1: *tert*-Butyl 4-((3-(hexanoyloxy)propyl)(nonyl)amino)propyl piperidine-1-carboxylate



Chemical Formula: C₃₁H₆₀N₂O₄

Molecular Weight: 524.83


[00817] In the same manner as Step 1 for Compound 57, *tert*-butyl 4-((3-

10 (hexanoyloxy)propyl)(nonyl)amino)propyl piperidine-1-carboxylate was synthesized from *tert*-butyl 4-(3-(nonylamino)propyl)piperidine-1-carboxylate (500 mg, 1.36 mmol), 3-bromopropyl hexanoate (386 mg, 1.63 mmol), K₂CO₃ (375 mg, 2.71 mmol), and KI (45 mg, 0.27 mmol) in MeCN (100 mL). Yield (322 mg, 45%).

UPLC/ELSD: RT = 2.09 min. MS (ES): *m/z* (MH⁺) 525.60 for C₃₁H₆₀N₂O₄

15 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.12 (br. m, 4H); 2.67 (m, 2H); 2.56-2.24 (br. m, 8H); 1.90-1.00 (br. m, 40H); 0.91 (m, 6H).

Step 2: 3-(nonyl(3-(piperidin-4-yl)propyl)amino)propyl hexanoate



Chemical Formula: C₂₆H₅₂N₂O₂

Molecular Weight: 424.71

[00818] In the same manner as Step 4 for Compound 11, 3-(nonyl(3-(piperidin-4-


yl)propyl)amino)propyl hexanoate was synthesized from *tert*-butyl 4-((3-(hexanoyloxy)propyl)(nonyl)amino)propyl piperidine-1-carboxylate (322 mg, 0.614 mmol), and

25 TFA (2.3 mL, 31 mmol) in DCM (2.5 mL). Yield (260 mg, 99%).

UPLC/ELSD: RT = 0.89 min. MS (ES): *m/z* (MH⁺) 425.54 for C₂₆H₅₂N₂O₂

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.12 (t, 2H); 3.12 (m, 2H); 2.75-2.24 (br. m, 10H); 1.84-1.54 (br. m, 6H); 1.54-1.02 (br. m, 26H); 0.90 (m, 6H).

Step 3: 3-((3-(1-(3-((2-(Dinonylamino)ethyl)(nonyl)amino)propanoyl)piperidin-4-yl)propyl)(nonyl)amino)propyl hexanoate



Chemical Formula: C₅₈H₁₁₆N₄O₃

Molecular Weight: 917.59

5


[00819] In the same manner as Step 11 for Compound 11, 3-((3-(1-(3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoyl)piperidin-4-yl)propyl)(nonyl)amino)propyl hexanoate was synthesized from 3-(nonyl(3-(piperidin-4-yl)propyl)amino)propyl hexanoate (149 mg, 0.351 mmol), 3-((2-(dinonylamino)ethyl)(nonyl)amino)propanoic acid (197 mg, 0.386 mmol), iPr₂EtN (134 μ L, 0.77 mmol), and T3P (50% EtOAc solution, 616 μ L, 1.05 mmol) in THF (10 mL). Yield (39 mg, 12%).

10 UPLC/ELSD: RT = 2.83 min. MS (ES): *m/z* (MH⁺) 918.01 for C₅₈H₁₁₆N₄O₃

¹H-NMR (300 MHz, CDCl₃) δ : ppm 4.61 (m, 1H); 4.12 (t, 2H); 3.87 (m, 1H); 3.08-2.74 (br. m, 4H); 2.70-2.23 (br. m, 20H); 1.82-1.56 (br. m, 6H); 1.56-1.00 (br. m, 67H); 0.90 (m, 15H).

15

BZ. Compound 17-1: 2-(Dihexylamino)ethan-1-ol



Chemical Formula: C₁₄H₃₁NO

Molecular Weight: 229.41

20 [00820] To a solution of 1-bromohexane (5 g, 82 mmol) in MeCN (380 mL) was added ethanolamine (11.5 mL, 82 mmol), K₂CO₃ (22.7 g, 164 mmol), and KI (1.36 g, 8.2 mmol). The reaction was allowed to stir at 82 °C for 48 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica 25 flash chromatography (0-20% MeOH/DCM) provided 2-(dihexylamino)ethan-1-ol (2.58 g, 14%).

UPLC/ELSD: RT = 0.41 min. MS (ES): *m/z* (MH⁺) 229.95 for C₁₄H₃₁NO


¹H NMR (300 MHz, CDCl₃) δ: ppm 3.62 (t, 2H); 2.70 (t, 2H), 2.57 (t, 4H); 1.50 (br. m, 4H); 1.30 (br, 12H); 0.91 (t, 6H).

CA. Compound 17-2: 2-(Hexyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol

5

Step 1: 2-(Hexylamino)ethan-1-ol

[00821] 2-(Hexylamino)ethan-1-ol was isolated from the same reaction that produced with Compound 1, 2-(dihexylamino)ethan-1-ol.




Chemical Formula: C₈H₁₉NO

Molecular Weight: 145.25

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.62 (t, 2H); 2.78 (t, 2H); 2.62 (t, 2H); 2.10-1.80 (br. m, 2H); 1.49 (m, 2H); 1.30 (br. m, 6H); 0.89 (t, 3H).

15 Step 2: 2-(Hexyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol



Chemical Formula: C₂₆H₅₁NO

Molecular Weight: 393.70


[00822] To a solution of (6Z,9Z)-18-bromoocatadeca-6,9-diene (0.2 g, 0.61 mmol) in MeCN (3.5 mL) was added 2-(hexylamino)ethan-1-ol (80 mg, 0.55 mmol), K₂CO₃ (76 mg, 0.55 mmol), and KI (9 mg, 0.06 mmol). The reaction was allowed to stir at room temperature for 18 hours. The reaction mixture was cooled to room temperature, added ethyl acetate and extracted with water. The combined extracts were dried with Na₂SO₄, filtered and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-10% MeOH/DCM) provided 2-(Hexyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol (23 mg, 11%).

LC/ELSD: RT = 2.47 min. MS (ES): m/z (MH $^+$) 394.60 for C₂₆H₅₁NO

¹H NMR (300 MHz, CDCl₃) δ: ppm 5.36 (m, 4H); 4.07 (m, 2H); 3.23-3.13 (m, 6H); 2.77 (m, 2H); 2.04 (m, 4H); 1.86 (m, 4H); 1.34 (m, 23H); 0.89 (m, 6H)

30 CB. Compound 17-3: 2-(Nonyl(9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol

[00823] Compound 17-3 was synthesized according to the same procedure as Compound 17-2.



Chemical Formula: C₂₉H₅₇NO


5 Molecular Weight: 435.78

LC/ELSD: RT = 2.72 min. MS (ES): *m/z* (MH⁺) 436.63 for C₂₉H₅₇NO

¹H NMR (300 MHz, CDCl₃) δ: ppm 5.36 (m, 4H); 3.55 (t, 2H); 2.77 (t, 2H); 2.60 (t, 2H); 2.47 (m, 4H); 2.04 (m, 4H); 1.55-1.18 (br. m, 33H); 0.87 (m, 6H).

10 CD. Compound 17-4: 2-(Dodecyl((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol

[00824] Compound 17-4 was synthesized according to the same procedure as Compound 17-2.



Chemical Formula: C₃₂H₆₃NO


15 Molecular Weight: 477.86

UPLC: RT = 3.18 min. MS (ES): *m/z* (MH⁺) 478.516 for C₃₂H₆₃NO

¹H NMR (400 MHz, CDCl₃) δ: ppm 5.33 (m, 4H); 3.53 (s, 2H); 2.75 (t, 2H); 2.58 (m, 2H); 2.45 (m, 4H); 2.03 (dt, 4H); 1.43 (m, 4H); 1.24 (m, 34H); 0.86 (m, 6H).

20 CE. Compound 17-5: 2-((9Z,12Z)-Octadeca-9,12-dien-1-yl)(tetradecyl)aminoethan-1-ol

[00825] Compound 17-5 was synthesized according to the same procedure as Compound 17-2.



Chemical Formula: C₃₄H₆₇NO

25 Molecular Weight: 505.92


LC/ELSD: RT = 3.39 min. MS (ES): *m/z* (MH⁺) 506.56 for C₃₄H₆₇NO

¹H NMR (300 MHz, CDCl₃) δ: ppm 5.37 (m, 4H); 3.58 (m, 2H); 2.80 (m, 2H); 2.69 -2.42 (br. m, 5H); 2.07 (m, 4H); 1.56-1.18 (br. m, 44H); 0.91 (m, 6H).

CF. Compound 17-6: 2-(((9Z,12Z)-Octadeca-9,12-dien-1-yl)(octadecyl)amino)ethan-1-ol

[00826] Compound 17-6 was synthesized according to the same procedure as Compound 17-2.

5



Chemical Formula: C₃₈H₇₅NO

Molecular Weight: 562.02

LC/ELSD: RT = 3.68 min. MS (ES): *m/z* (MH⁺) 562.58 for C₃₈H₇₅NO


¹H NMR (300 MHz, CDCl₃) δ: ppm 5.39 (m, 4H); 3.58 (m, 2H); 2.80 (m, 2H); 2.68 -2.44 (br.

10 m, 5H); 2.07 (m, 4H); 1.57-1.20 (br. m, 52H); 0.91 (m, 6H).

CG. Compound 17-7: 2-(Ditetradecylamino)ethan-1-ol

[00827] Compound 17-7 was synthesized according to the same procedure as Compound 17-1.

15



Chemical Formula: C₃₀H₆₃NO

Molecular Weight: 453.84

UPLC/ELSD: RT = 3.30 min. MS (ES): *m/z* (MH⁺) 454.46 for C₃₀H₆₃NO


¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.54 (br. m, 2H); 2.59 (br. m, 2H); 2.46 (br. m, 4H); 1.56-

20 1.17(br. m, 48H); 0.90 (br. m, 6H).

CG. Compound 17-8: 2-(Di((Z)-octadec-9-en-1-yl)amino)ethan-1-ol

[00828] Compound 17-8 was synthesized according to the same procedure as Compound 17-1.

25



Chemical Formula: C₃₈H₇₅NO

Molecular Weight: 562.02


UPLC/ELSD: RT = 7.325 min. MS (ES): *m/z* (MH⁺) 562.60 for C₃₈H₇₅NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 4H); 3.53 (t, 2H); 2.58 (t, 2H); 2.45 (t, 4H); 2.01 (m, 8H); 1.44 (m, 4H); 1.38-1.18 (br. m, 44H); 0.88 (t, 6H).

CH. Compound 17-9:

5 (9Z,12Z)-N-(2-Methoxyethyl)-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine

[00829] Compound 17-9 was synthesized according to the same procedure as Compound 17-1.



Chemical Formula: C₃₉H₇₃NO


10 Molecular Weight: 572.02

LC/ELSD: RT = 3.53 min. MS (ES): *m/z* (MH⁺) 572.72 for C₃₉H₇₃NO

¹H NMR (300 MHz, CDCl₃) δ: ppm 5.39 (m, 8H); 3.47 (m, 2H); 3.37 (s, 3H); 2.80 (m, 4H); 2.5 (m, 2H); 2.46 (m, 4H); 2.09 (m, 8H); 1.50-1.22 (m, 36H); 0.92 (m, 6H).

15 CI. Compound 17-10: 2-(Dinonylamino)ethan-1-ol

Compound 17-10 was synthesized according to the same procedure as Compound 17-1.




Chemical Formula: C₂₀H₄₃NO

Molecular Weight: 313.57

20 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.57 (t, 2H); 2.63 (t, 2H); 2.49 (br. m, 4H); 1.48 (br. m, 4H); 1.29 (br. m, 24H); 0.91 (t, 6H).

CJ: Compound 17-11: 2-(Didodecylamino)ethan-1-ol

Compound 17-11 was synthesized according to the same procedure as Compound 17-1.



Chemical Formula: C₂₆H₅₄ClN


Molecular Weight: 416.18

UPLC/ELSD: RT = 2.69 min. MS (ES): *m/z* (MH⁺) 398.56 for C₂₆H₅₅NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.57 (t, 2H); 2.63 (t, 2H); 2.49 (br. m, 4H); 1.48 (br. m, 4H); 1.29 (br. m, 36H); 0.91 (t, 6H).

CK. Compound 17-12: 3-(Didodecylamino)propan-1-ol

5 [00830] Compound 17-12 was synthesized according to the same procedure as Compound 1.



Chemical Formula: C₂₇H₅₇NO


Molecular Weight: 411.76

10 UPLC/ELSD: RT = 2.75 min. MS (ES): *m/z* (MH⁺) 412.36 for C₂₇H₅₇NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.79 (t, 2H); 2.66 (t, 2H); 2.43 (br. m, 4H); 1.69 (br. m, 2H); 1.47 (br. m, 4H) 1.25 (br. m, 36H); 0.87 (t, 6H).

CL. Compound 17-13: 4-(Didodecylamino)butan-1-ol

15 Compound 17-13 was synthesized according to the same procedure as Compound 17-1.



Chemical Formula: C₂₈H₅₉NO


Molecular Weight: 425.79

UPLC/ELSD: RT = 2.80 min. MS (ES): *m/z* (MH⁺) 426.42 for C₂₈H₅₉NO

20 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.56 (br. m, 2H); 2.46 (br. m, 6H); 1.66 (br. m, 4H); 1.48 (br. m, 4H); 1.26 (br. m, 36H); 0.88 (t, 6H).

CM. Compound 19-1: *N*-Nonyl-*N*-(2-(piperazin-1-yl)ethyl)nonan-1-amine

25 Step 1: *tert*-Butyl 4-(2-(dinonylamino)ethyl)piperazine-1-carboxylate




Chemical Formula: C₂₉H₅₉N₃O₂

Molecular Weight: 481.81

[00831] A mixture of 1-bromononane (1.81 g, 8.72 mmol), 4-(2-aminoethyl)-1-boc-piperazine (2.0 g, 8.72 mmol), K_2CO_3 (2.4 g, 17.4 mmol), KI (145 mg, 0.872 mmol) in 44 mL MeCN was allowed to stir at 65 °C for 16 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided *tert*-butyl 4-(2-(dinonylamino)ethyl)piperazine-1-carboxylate (924 mg, 1.92 mmol, 44%).
 UPLC/ELSD: RT = 1.99 min. MS (ES): m/z (MH^+) 482.36 for $C_{29}H_{59}N_3O_2$
 1H NMR (400 MHz, $CDCl_3$) δ : ppm 3.45 (br. m, 4H); 3.10 (br. m, 2H); 2.59 (br. m, 2H); 2.44 (br. m, 8H); 1.60-1.00 (br. m, 37H); 0.91 (t, 6H).

Step 2: Compound 19-1: *N*-Nonyl-*N*-(2-(piperazin-1-yl)ethyl)nonan-1-amine




Chemical Formula: $C_{24}H_{51}N_3$

15 Molecular Weight: 381.69

[00832] A solution of *tert*-butyl 4-(2-(dinonylamino)ethyl)piperazine-1-carboxylate (924 mg, 1.92 mmol) in 8 mL DCM was treated with TFA (7.4 mL, 96 mmol). The reaction was allowed to stir at room temperature for 16 hours. The reaction was concentrated, and the crude residue was taken up in chloroform and washed with 5% Na_2CO_3 and brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% DCM/[DCM, 20% MeOH, 1% NH_4OH]) provided *N*-nonyl-*N*-(2-(piperazin-1-yl)ethyl)nonan-1-amine (563 mg, 1.48 mmol, 77%).
 UPLC/ELSD: RT = 1.27 min. MS (ES): m/z (MH^+) 382.54 for $C_{24}H_{51}N_3$
 1H NMR (400 MHz, $CDCl_3$) δ : ppm 2.92 (br. m, 4H); 2.62 (br. m, 2H); 2.48 (br. m, 10H); 2.40-1.88 (br. m, 1H); 1.46 (br. m, 4H); 1.29 (br. m, 24H), 0.91 (t, 6H).

CN. Compound 19-2: *N*-Dodecyl-*N*-(2-(piperazin-1-yl)ethyl)dodecan-1-amine

Step 1: *tert*-Butyl 4-(2-(didodecylamino)ethyl)piperazine-1-carboxylate



Chemical Formula: C₃₅H₇₁N₃O₂

Molecular Weight: 565.97


[00833] A mixture of 1-bromododecane (1.1 mL, 4.6 mmol), 4-(2-aminoethyl)-1-boc-

5 piperazine (1.0 g, 4.4 mmol), K₂CO₃ (0.61 g, 4.4 mmol), in 10 mL MeCN was allowed to stir at room temperature for 12 h. After this time the reaction was filtered and concentrated. The crude material was purified by silica gel chromatography (0-20% MeOH in DCM with 1% NH₄OH to afford *tert*-butyl 4-(2-(dodecylamino)ethyl)piperazine-1-carboxylate (450 mg, 0.80 mmol, 18%).

10 UPLC/ELSD: RT = 2.87 min. MS (ES): *m/z* (MH⁺) 566.655 for C₃₅H₇₁N₃O₂

¹H NMR (400 MHz, CDCl₃) δ: ppm 3.40 (m, 4H); 2.56 (m, 2H); 2.40 (m, 10H); 1.44 (s, 9H); 1.40-1.24 (m, 40H); 0.86 (t, 6H).

Step 2: Compound 19-2: *N*-Dodecyl-*N*-(2-(piperazin-1-yl)ethyl)dodecan-1-amine



15

Chemical Formula: C₃₀H₆₃N₃

Molecular Weight: 465.86

[00834] A solution of *tert*-butyl 4-(2-(dodecylamino)ethyl)piperazine-1-carboxylate

(154 mg, 0.27 mmol) in 1 mL DCM was treated with TFA (0.21 mL, 2.7 mmol). The reaction 20 was allowed to stir overnight. After this time addition TFA (0.1 mL, 1.3 mmol) was added.

After an additional 3 h the reaction was concentrated. The crude residue was taken up in DCM and washed with 5% K₂CO₃ and brine, dried over Na₂SO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (0-20% MeOH in DCM with 1% NH₄OH) to provide *N*-dodecyl-*N*-(2-(piperazin-1-yl)ethyl)dodecan-1-amine (109 mg, 87%).


25 UPLC/ELSD: RT = 2.10 min. MS (ES): *m/z* (MH⁺) 466.379 for C₃₀H₆₃N₃

¹H NMR (400 MHz, CDCl₃) δ: ppm 2.88 (t, 4H); 2.61 (m, 2H); 2.45 (m, 10H); 1.43-1.24 (m, 40H); 0.86 (t, 6H).

CO. Compound 19-3: (9Z,12Z)-N-((9Z,12Z)-Octadeca-9,12-dien-1-yl)-N-(2-(piperazin-1-yl)ethyl)octadeca-9,12-dien-1-amine

Step 1: (9Z,12Z)-Octadeca-9,12-dien-1-yl methanesulfonate

5



Chemical Formula: C₁₉H₃₆O₃S


Molecular Weight: 344.55

[00835] To a 0 °C solution of linoleyl alcohol (10 mL, 31.2 mmol) and trimethylamine (5.68 mL, 40.5 mmol) in DCM (50 mL) was added dropwise a solution of methanesulfonyl chloride (2.66 mL, 34.3 mmol) in DCM (20 mL). The reaction was allowed to return to room temperature and let stir for 4 hours. The mixture was quenched by the addition of water and extracted with DCM. The organic layer was washed with saturated NaHCO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-40% EtOAc/hexanes) provided (9Z,12Z)-octadeca-9,12-dien-1-yl methanesulfonate (10.0 g, 93%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.35 (m, 4H); 4.22 (t, 2H); 2.99 (s, 3H); 2.77 (t, 2H); 2.04 (q, 4H); 1.74 (m, 2H); 1.30 (br. m, 16H); 0.89 (t, 3H).

Step 2: (6Z,9Z)-18-Bromoocatadeca-6,9-diene

20




Chemical Formula: C₁₈H₃₃Br

Molecular Weight: 329.37

[00836] To a solution of (9Z,12Z)-octadeca-9,12-dien-1-yl methanesulfonate (10.0 g, 29.0 mmol) in diethyl ether (372 mL) was added magnesium bromide ethyl etherate (22.5 g, 87.1 mmol). The reaction was let stir at room temperature for 16 hours. The mixture was quenched by the addition of water and extracted with diethyl ether. The combined organic layers were washed with 1% K₂CO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography provided (6Z,9Z)-18-bromoocatadeca-6,9-diene (8.9 g, 93%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.36 (m, 4H); 3.41 (t, 2H); 2.77 (t, 2H); 2.05 (q, 4H); 1.86 (m, 2H); 1.48-1.22 (br. m, 16H); 0.89 (t, 3H).

Step 3: *tert*-Butyl 4-(2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate




Chemical Formula: C₄₇H₈₇N₃O₂

Molecular Weight: 726.23

[00837] A mixture of (6Z,9Z)-18-bromoocatadeca-6,9-diene (1.5 g, 4.55 mmol), 4-(2-aminoethyl)-1-boc-piperazine (1.04 g, 4.54 mmol), K₂CO₃ (1.27 g, 9.10 mmol), KI (75 mg, 0.452 mmol), in 22 mL MeCN was allowed to stir at room temperature for 48 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-50% DCM/[DCM, 20% MeOH, 1% NH₄OH]) provided *tert*-butyl 4-(2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate (1.08 g, 1.49 mmol, 65%).

¹H NMR (400 MHz, CDCl₃) δ: ppm 5.43-5.26 (br. m, 8H); 3.42 (t, 4H); 2.77 (m, 4H); 2.57 (m, 2H); 2.41 (br. m, 10H); 2.04 (br. m, 8H); 1.60-1.00 (br. m, 45H); 0.89 (t, 6H).

Step 4: Compound 19-3: (9Z,12Z)-N-((9Z,12Z)-Octadeca-9,12-dien-1-yl)-N-(2-(piperazin-1-yl)ethyl)octadeca-9,12-dien-1-amine



Chemical Formula: C₄₂H₇₉N₃

Molecular Weight: 626.12

[00838] A solution of *tert*-butyl 4-(2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)piperazine-1-carboxylate (1.06 g, 1.46 mmol) in 6 mL DCM was treated with TFA (5.6 mL, 73 mmol). After 4 hours the mixture was concentrated. The crude residue was taken up in chloroform, washed with 5% Na₂CO₃, brine, dried over anhydrous Na₂SO₄ and


concentrated *in vacuo*. The residue was purification by ISCO silica flash chromatography (0-100% DCM/[DCM, 20% MeOH, 1% NH₄OH]) and ISCO C18 flash chromatography (50-100% [MeCN 1% TFA]/[H₂O 1% TFA]). The desired fractions were washed with 5% Na₂CO₃ and extracted with hexanes. The hexanes were washed with brine, dried over anhydrous Na₂SO₄, and concentrated *in vacuo* to provide (9Z,12Z)-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)-N-(2-(piperazin-1-yl)ethyl)octadeca-9,12-dien-1-amine (108 mg, 12%).

5 UPLC/ELSD: RT = 2.98 min. MS (ES): *m/z* (MH⁺) 626.75 for C₄₂H₇₉N₃
¹H NMR (400 MHz, CDCl₃) δ: ppm 5.47-5.25 (br. m, 8H); 2.92 (m, 4H); 2.76 (m, 4H); 2.66 (br. m, 2H); 2.50 (br. m, 10H); 2.05 (m, 8H); 1.60-1.10 (br. m, 36H), 0.89 (t, 6H).

10

CP. Compound 19-4: *N*-Dodecyl-*N*-(2-(4-methylpiperazin-1-yl)ethyl)dodecan-1-amine

Intermediate 1: 2-(Didodecylamino)ethan-1-ol



Chemical Formula: C₂₆H₅₅NO

15

Molecular Weight: 397.73

20


[00839] To a solution of 1-bromododecane (10 g, 40.1 mmol) in MeCN (84 mL) was added ethanolamine (1.10 mL, 18.2 mmol), K₂CO₃ (11.1 g, 80.1 mmol), and KI (302 mg, 1.82 mmol). The reaction was allowed to stir at 82 °C for 48 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided 2-(didodecylamino)ethan-1-ol (3.87 g, 53%).

UPLC/ELSD: RT = 2.69 min. MS (ES): *m/z* (MH⁺) 398.56 for C₂₆H₅₅NO

25

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.57 (t, 2H); 2.63 (t, 2H); 2.49 (br. m, 4H); 1.48 (br. m, 4H); 1.29 (br. m, 36H); 0.91 (t, 6H).

Step 2: *N*-(2-Chloroethyl)-*N*-dodecylidodecan-1-amine



Chemical Formula: C₂₆H₅₄ClN


30

Molecular Weight: 416.18

[00840] To a 0 °C solution of 2-(didodecylamino)ethan-1-ol (3.87 g, 9.73 mmol) triethylamine (1.76 mL, 12.6 mmol) in DCM (50 mL) was added dropwise a solution of methanesulfonyl chloride (0.941 mL, 12.2 mmol) in DCM (5 mL). The reaction was allowed to return to room temperature and stir for 16 hours. The mixture was quenched by the addition of 5 water and extracted with DCM. The organic layer was washed with saturated NaHCO_3 , brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-10% EtOAc/hexanes) provided *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (1.92 g, 47%).

10 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 3.51 (t, 2H); 2.78 (t, 2H); 2.47 (br. m, 4H); 1.44 (br. m, 4H); 1.28 (br. m, 36H); 0.90 (t, 6H).

Step 3: Compound 19-4: *N*-Dodecyl-*N*-(2-(4-methylpiperazin-1-yl)ethyl)dodecan-1-amine



Chemical Formula: $\text{C}_{31}\text{H}_{65}\text{N}_3$

15 Molecular Weight: 479.88


[00841] A mixture of *N*-methylpiperazine (40 μL , 0.36 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (166 mg, 0.4 mmol), and K_2CO_3 (50 mg, 0.36 mmol) in 2 mL MeCN was allowed to stir at 82 °C for 12 h. The reaction was allowed to cool room temperature, was filtered and concentrated. The crude material was purified by silica gel chromatography (0-20% 20 MeOH in DCM with 1% NH₄OH) to provide *N*-dodecyl-*N*-(2-(4-methylpiperazin-1-yl)ethyl)dodecan-1-amine (87.9 mg, 51%).

UPLC: RT = 2.24 min. MS (ES): m/z (MH^+) 480.662 for $\text{C}_{31}\text{H}_{65}\text{N}_3$

$^1\text{H NMR}$ (400 MHz, CDCl_3) δ : ppm 2.49 (m, 16H); 2.36 (s, 3H); 1.50 (m, 4H); 1.34 (m, 36H); 0.96 (t, 6H).

25

CQ. Compound 19-5: *N*-Dodecyl-*N*-(2-(4-(4-methoxybenzyl)piperazin-1-yl)ethyl)dodecan-1-amine



Chemical Formula: C₃₈H₇₁N₃O


Molecular Weight: 586.01

[00842] A mixture of 1-(4-methoxybenzyl)piperazine (206 mg, 1.0 mmol), *N*-(2-chloroethyl)-*N*-dodecyldecane-1-amine (289 mg, 0.69 mmol), K₂CO₃ (286 mg, 2.07 mmol) and KI (11 mg, 0.069 mmol) in 3.5 mL MeCN was allowed to stir at 80 °C for 2 h. After this time the reaction was allowed to cool to room temperature and was quenched with water. The mixture was extracted with EtOAc three times. The pooled organics were washed with brine, dried over MgSO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (0-20% MeOH in DCM) for provide *N*-dodecyl-*N*-(2-(4-(4-methoxybenzyl)piperazin-1-yl)ethyl)dodecan-1-amine (0.24 g, 59%).

UPLC: RT = 2.30 min. MS (ES): *m/z* (MH⁺) 586.92 for C₃₈H₇₁N₃O

¹H NMR (400 MHz, CDCl₃) δ: ppm 7.19 (d, 2H); 6.83 (d, 2H); 3.78 (s, 3H); 3.42 (s, 2H); 2.99-2.45 (br. m, 16H); 1.71-1.24 (br. m, 40H); 0.86 (t, 6H).

15 CR. Compound 19-6: (9Z,12Z)-*N*-(2-(4-Dodecylpiperazin-1-yl)ethyl)-*N*-(9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine

Chemical Formula: C₅₄H₁₀₃N₃

20 Molecular Weight: 794.44


[00843] A mixture of (9Z,12Z)-*N*-(9Z,12Z)-Octadeca-9,12-dien-1-yl)-*N*-(2-(piperazin-1-yl)ethyl)octadeca-9,12-dien-1-amine (54 mg, 0.086 mmol), 1-bromododecane (24 mg, 0.095 mmol), K₂CO₃ (24 mg, 0.172 mmol), KI (2 mg, 0.012 mmol), in 1.5 mL THF was allowed to stir at 65 °C for 16 hours. The reaction was cooled to room temperature, diluted with H₂O, and extracted with EtOAc. The organics were washed with brine, dried over anhydrous MgSO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% DCM/[DCM 20% MeOH 1% Et₃N]) provided (9Z,12Z)-*N*-(2-(4-dodecylpiperazin-1-yl)ethyl)-*N*-(9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine (51 mg, 0.064 mmol, 74%).

UPLC: RT = 3.40 min. MS (ES): *m/z* (MH⁺) 795.12 for C₅₄H₁₀₃N₃.

30

CS. Compound 20-1: *N*-(2-(Didodecylamino)ethyl)-*N*-dodecylglycine

Step 1: Methyl *N*-(*tert*-butoxy carbonyl)-*N*-dodecylglycinate



5


Chemical Formula: C₂₀H₃₉NO₄

Molecular Weight: 357.54

[00844] A 0 °C solution of *N*-(*tert*-butoxycarbonyl)glycine methyl ester (7.7 g, 40.7 mmol) in DMF (100 mL) was treated with NaH (60%, 1.71 g, 42.7 mmol) and the mixture was allowed to stir for 30 minutes. The solution was allowed to return to room temperature before 1-bromododecane (15.2 g, 61.0 mmol) was added and the reaction was allowed to stir overnight. The reaction was quenched with water and extracted with EtOAc. The organics were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% EtOAc/hexanes) provided methyl *N*-(*tert*-butoxycarbonyl)-*N*-dodecylglycinate (4.03 g, 28%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 4.01-3.84 (br. m, 2H); 3.75 (s, 3H); 3.27 (br. m, 2H); 1.67-1.39 (br. m, 11H); 1.28 (br, 18H); 0.90 (t, 3H).

Step 2: Methyl dodecylglycinate



20

Chemical Formula: C₁₅H₃₁NO₂


Molecular Weight: 257.42

[00845] To a 0 °C solution of methyl N-(tert-butoxy carbonyl)-N-dodecylglycinate (4.03 g, 11.3 mmol) in DCM (17 mL) was added dropwise TFA (17 mL, 226 mmol). The reaction was allowed to return to room temperature and stir for 6 hours. The reaction mixture was concentrated in vacuo and the crude material was dissolved in DCM. The solution was washed with 10% NaOH, brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo to provide methyl dodecylglycinate (2.84 g, 98%).

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.75 (s, 3H); 3.44 (s, 2H); 2.62 (t, 2H); 1.70 (br, 1H); 1.51 (m, 2H); 1.29 (br, 18H); 0.90 (t, 3H).

Step 3: 2-(Didodecylamino)ethan-1-ol

5



Chemical Formula: C₂₆H₅₅NO


Molecular Weight: 397.73

[00846] To a solution of 1-bromododecane (10 g, 40.1 mmol) in MeCN (84 mL) was 10 added ethanolamine (1.10 mL, 18.2 mmol), K₂CO₃ (11.1 g, 80.1 mmol), and KI (302 mg, 1.82 mmol). The reaction was allowed to stir at 82 °C for 48 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes. The filtrate was extracted with hexanes, and the combined extracts were concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided 2-(didodecylamino)ethan-1- 15 ol (3.87 g, 53%).

UPLC/ELSD: RT = 2.69 min. MS (ES): *m/z* (MH⁺) 398.56 for C₂₆H₅₅NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.57 (t, 2H); 2.63 (t, 2H); 2.49 (br. m, 4H); 1.48 (br. m, 4H); 1.29 (br, 36H); 0.91 (t, 6H).

20 Step 4: *N*-(2-Chloroethyl)-*N*-dodecylidodecan-1-amine



Chemical Formula: C₂₆H₅₄ClN


Molecular Weight: 416.18

[00847] To a 0 °C solution of 2-(didodecylamino)ethan-1-ol (3.87 g, 9.73 mmol) and 25 triethylamine (1.76 mL, 12.6 mmol) in DCM (50 mL) was added dropwise a solution of methanesulfonyl chloride (0.941 mL, 12.2 mmol) in DCM (5 mL). The reaction was allowed to return to room temperature and stir for 16 hours. The mixture was quenched by the addition of water and extracted with DCM. The organic layer was washed with saturated NaHCO₃, brine,

dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-10% EtOAc/hexanes) provided *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (1.92 g, 47%).

¹H-NMR (300 MHz, CDCl_3) δ : ppm 3.51 (t, 2H); 2.78 (t, 2H); 2.47 (br. m, 4H); 1.44 (br. m, 4H); 1.28 (br, 36H); 0.90 (t, 6H).

Step 5: Methyl *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycinate



Chemical Formula: $\text{C}_{41}\text{H}_{84}\text{N}_2\text{O}_2$


10 Molecular Weight: 637.14

[00848] To a solution of methyl dodecylglycinate (425 mg, 1.65 mmol) in MeCN (10 mL) was added *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (825 mg, 1.98 mmol), K_2CO_3 (457 mg, 3.30 mmol), and KI (27 mg, 0.165 mmol). The reaction was allowed to stir at 82 °C for 72 hours. The reaction mixture was filtered and the solids were washed with hexanes. The filtrate 15 was concentrated *in vacuo* to provide the crude product. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided methyl *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycinate (652 mg, 62%).

UPLC/ELSD: RT = 3.77 min. MS (ES): m/z (MH^+) 638.18 for $\text{C}_{41}\text{H}_{84}\text{N}_2\text{O}_2$

¹H-NMR (300 MHz, CDCl_3) δ : ppm 3.72 (s, 3H); 3.41 (s, 2H); 2.90-2.20 (br. m, 10H); 1.60-20 1.00 (br. m, 60H); 0.90 (t, 9H).

Step 6: *N*-(2-(Didodecylamino)ethyl)-*N*-dodecylglycine



Chemical Formula: $\text{C}_{40}\text{H}_{82}\text{N}_2\text{O}_2$

25 Molecular Weight: 623.11


[00849] A solution of methyl *N*-(2-(didodecylamino)ethyl)-*N*-dodecylglycinate (652 mg, 1.02 mmol) in THF (6 mL) and 1M LiOH (5 mL, 5 mmol) was allowed to stir at 65 °C for 16

hours. The reaction was cooled to room temperature and acidified with 10% HCl. The mixture was extracted with chloroform, and the organics were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-20% MeOH/DCM) provided *N*-(2-(dodecylamino)ethyl)-*N*-dodecylglycine (153 mg, 24%).

5 UPLC/ELSD: RT = 3.60 min. MS (ES): m/z (MH^+) 624.07 for $\text{C}_{40}\text{H}_{82}\text{N}_2\text{O}_2$
¹H-NMR (300 MHz, CDCl_3) δ : ppm 4.02-3.40 (br. m, 6H); 3.16 (br, 6H); 1.78 (br, 6H); 1.46-1.01 (br. m, 54H); 0.90 (t, 9H).

10 CT. Compound 20-2: Petyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate

Step 1: 2-(Dodecylamino)ethan-1-ol




Chemical Formula: $\text{C}_{14}\text{H}_{31}\text{NO}$

15 Molecular Weight: 229.41

[00850] Methyl dodecylglycinate (3.4 g, 13.2 mmol) was dissolved in 2 mL THF under N_2 atmosphere and the reaction flask was allowed to cool in an ice bath. To the solution LiAlH_4 (0.55 g, 14.5 mmol) was slowly added. The reaction was allowed to stir at the same temperature for 1 h. After this time the reaction was quenched by the subsequent addition of 0.55 mL H_2O , 20 0.55 mL 10% NaOH and then 1.65 mL of H_2O . The reaction was then filtered and the filtrate was concentrated *in vacuo*. The crude material was purified *via* silica gel chromatography (0-20% MeOH in DCM, with 1% NH_4OH) to afford 2-(dodecylamino)ethan-1-ol (1.9 g, 8.28 mmol, 63% yield).

25 ¹H NMR (400 MHz, CDCl_3) δ : ppm 3.63 (t, 2H); 2.78 (t, 2H); 2.63 (t, 2H); 1.48 (m, 2H); 2.14 (m, 18H); 0.88 (t, 3H).

Step 2: Petyl 6-bromohexanoate




Chemical Formula: $\text{C}_{11}\text{H}_{21}\text{BrO}_2$

30 Molecular Weight: 265.19

[00851] To a solution of 6-bromohexanoic acid (2 g, 10.3 mmol) and pentan-1-ol (2.2 mL, 20.5 mmol) in 26 mL DCM, EDC·HCl (1.97 g, 10.3 mmol) and DMAP (0.26 g, 2.1 mmol) were added. The solution was allowed to stir at room temperature overnight. After this time the reaction was quenched by the addition of water. The mixture was extracted three times with 5 DCM. The organics were pooled and washed with saturated NaHCO₃, 10% citric acid and brine. The organics were then dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was purified *via* silica gel chromatography (0-30% EtOAc in hexanes) to afford the desired product (2.3 g, 8.67 mmol).

10 ¹H NMR (400 MHz, CDCl₃) δ: ppm 4.06 (t, 2H); 3.39 (t, 2H); 2.30 (t, 2H); 1.84 (m, 2H); 1.62 (m, 4H); 1.46 (m, 2H); 1.31 (m, 4H); 0.88 (t, 3H).

Step 3: Pentyl 6-(dodecyl(2-hydroxyethyl)amino)hexanoate




Chemical Formula: C₂₅H₅₁NO₃

15 Molecular Weight: 413.69

[00852] To a solution of 2-(dodecylamino)ethan-1-ol (0.50 g, 2.18 mmol) in 10 mL THF, pentyl 6-bromohexanoate (0.87 g, 3.27 mmol) was added followed by K₂CO₃ (0.60 g, 4.36 mmol) and KI (36 mg, 0.22 mmol). The reaction was allowed to stir under N₂ at 65 °C for 24 h. After this time the reaction was allowed to cool to room temperature and the reaction was 20 diluted with water. The mixture was extracted three times with EtOAc. The pooled organics were washed with brine, dried over MgSO₄, filtered and concentrated. The crude material was purified by silica gel chromatography (0-20% EtOAc in hexanes) to afford pentyl 6-(dodecyl(2-hydroxyethyl)amino)hexanoate (300 mg, 33%).

25 ¹H NMR (400 MHz, CDCl₃) δ: ppm 4.04 (t, 2H); 3.51 (m, 2H); 2.56 (m, 2H); 2.42 (m, 4H); 2.28 (t, 2H); 1.60 (m, 4H); 1.42 (m, 4H); 1.30-1.24 (m, 24); 0.87 (m, 6H).

Step 4: Pentyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate


Chemical Formula: C₂₅H₅₀ClNO₂

Molecular Weight: 432.13

5 [00853] To a 0 °C solution of pentyl 6-(dodecyl(2-hydroxyethyl)amino)hexanoate (300 mg, 0.73 mmol) in 2 mL DCM, methanesulfonyl chloride (0.062 mL, 0.80 mmol) was added, followed by triethylamine (0.13 mL, 1.3 mmol). The reaction was allowed to slowly warm to room temperature and stir for 12 h under N₂. The reaction was quenched by the addition of water and was extracted with DCM. The pooled organics were dried over MgSO₄, filtered and concentrated. The aqueous layer was re-extracted with EtOAc three times. The organics were pooled and washed with brine, dried over MgSO₄, filtered and concentrated. The crude material was combined and purification by silica gel chromatography (0-30% EtOAc in hexanes) afforded pentyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate (285 mg, 66%).

10 15 ¹H NMR (400 MHz, CDCl₃) δ: ppm 4.04 (t, 2H); 3.45 (t, 2H); 2.74 (t, 2H); 2.43 (m, 4H); 2.28 (t, 2H); 1.65-1.59 (m, 4H); 1.31-1.24 (m, 32H); 0.88 (m, 6H).

Step 5: Pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate

Chemical Formula: C₃₉H₈₀N₂O₃

Molecular Weight: 625.08


20 [00854] To a solution of pentyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate (94 mg, 0.22 mmol) in MeCN (2 mL) and THF (2 mL) was added 2-(dodecylamino)ethan-1-ol (50 mg, 0.22 mmol), K₂CO₃ (60 mg, 0.44 mmol), and KI (4 mg, 0.022 mmol). The reaction was allowed to stir at 65 °C for 18 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes and EtOAc. The filtrate was extracted with EtOAc three times. The pooled organics were washed with water and brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% DCM, [20%

MeOH, 1% NH₄OH]/DCM) provided pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate (21 mg, 15%).

UPLC/ELSD: RT = 2.86 min. MS (ES): *m/z* (MH⁺) 625.86 for C₃₉H₈₀N₂O₃
¹H NMR (300 MHz, CDCl₃) δ: ppm 4.07-4.05 (m, 2H); 3.53 (m, 2H), 2.60-2.43 (br. m, 12H);
5 2.33-2.29 (m, 2H); 1.65-1.64 (m, 4H); 1.46 (m, 6H); 1.34-1.28 (br. m, 42H); 0.92-0.90 (m, 9H).

CU. Compound 20-3: Pentyl 6-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)hexanoate

Step 1: Pentyl 6-((2-hydroxyethyl)amino)hexanoate



Chemical Formula: C₁₃H₂₇NO₃


Molecular Weight: 245.36

[00855] To a solution of pentyl 6-bromohexanoate (4.65 g, 17.5 mmol) in MeCN (88 mL) was added ethanolamine (1.10 mL, 17.5 mmol), K₂CO₃ (4.85 g, 35.1 mmol), and KI (291 mg, 1.75 mmol). The reaction was allowed to stir at 82 °C for 48 hours. The reaction mixture was cooled to room temperature, filtered, and the solids were washed with hexanes and EtOAc. The filtrate was extracted with EtOAc three times. The pooled organics were washed with water and brine, dried over Na₂SO₄, filtered and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-100% DCM, [20% MeOH, 1% NH₄OH]/DCM) provided pentyl 6-((2-hydroxyethyl)amino)hexanoate (1.74 g, 41%).

UPLC/ELSD: RT = 0.30 min. MS (ES): *m/z* (MH⁺) 246.21 for C₁₃H₂₇NO₃

¹H NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 2H); 3.69 (t, 2H), 2.82 (t, 2H); 2.68 (t, 2H); 2.35-2.31 (m, 4H); 1.72-1.52 (br. m, 6H); 1.39-1.32 (br. m, 6H); 0.93 (t, 3H).

25 Step 2: Pentyl 6-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)hexanoate




Chemical Formula: C₃₉H₈₀N₂O₃

Molecular Weight: 625.08

[00856] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, pentyl 6-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)hexanoate was synthesized from pentyl 6-((2-hydroxyethyl)amino)hexanoate (108 mg, 0.44 mmol), *N*-(2-chloroethyl)-*N*-dodecylhexan-1-amine (183 mg, 0.44 mmol), K_2CO_3 (122 mg, 0.88 mmol), and KI (7.3 mg, 0.044 mmol) in MeCN (1 mL) and THF (1 mL). Yield (88 mg, 32%).

5 UPLC/ELSD: RT = 2.92 min. MS (ES): m/z (MH^+) 626.0 for $C_{39}H_{80}N_2O_3$,
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 4.07 (t, 2H); 3.53 (t, 2H), 2.62-2.40 (br. m, 12H); 2.31 (t, 10 2H); 1.70-1.60 (m, 4H); 1.53-1.43 (m, 6H); 1.27 (br. m, 42H); 0.91 (m, 9H).

CV. Compound 20-4: Dipentyl 6,6'-(2-(dodecyl(2-hydroxyethyl)amino)ethyl)azanediyldihexanoate



15 Chemical Formula: $C_{38}H_{76}N_2O_5$


Molecular Weight: 641.04

[00857] In the same manner as pentyl-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, dipentyl 6,6'-(2-(dodecyl(2-hydroxyethyl)amino)ethyl)azanediyldihexanoate was synthesized from 20 2-(dodecylamino)ethan-1-ol (60 mg, 0.26 mmol), dipentyl 6,6'-(2-chloroethyl)azanediyldihexanoate (118 mg, 0.26 mmol), K_2CO_3 (73 mg, 0.53 mmol), and KI (5 mg, 0.026 mmol) in MeCN (1 mL) and THF (1 mL). Yield (60 mg, 36%).

UPLC/ELSD: RT = 2.37 min. MS (ES): m/z (MH^+) 641.95 for $C_{38}H_{76}N_2O_5$,
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 4.08 (t, 4H); 3.57 (m, 2H), 2.64-2.53 (br. m, 12H); 2.32 (t, 25 4H); 1.72-1.60 (m, 8H); 1.50 (m, 6H); 1.38-1.28 (br. m, 30H); 0.95-0.88 (m, 9H).

CW. Compound 20-5:

Diheptyl 6,6'-(2-((6-(heptyloxy)-6-oxohexyl)(2-hydroxyethyl)amino)ethyl)azanediyil)dihexanoate



(Compound 20-5)

Chemical Formula: C₄₃H₈₄N₂O₇

5

Molecular Weight: 741.15

[00858] In the same manner as pentyl

6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, diheptyl

6,6'-(2-((6-(heptyloxy)-6-oxohexyl)(2-hydroxyethyl)amino)ethyl)azanediyil)dihexanoate was

synthesized from heptyl 6-((2-hydroxyethyl)amino)hexanoate (100 mg, 0.37 mmol), diheptyl


10 6,6'-(2-chloroethyl)azanediyil)dihexanoate (184 mg, 0.37 mmol), K₂CO₃ (101 mg, 0.73 mmol), and KI (6 mg, 0.037 mmol) in MeCN (2 mL) and THF (2 mL). Yield (91 mg, 34%).

UPLC/ELSD: RT = 3.33 min. MS (ES): *m/z* (MH⁺) 742.08 for C₄₃H₈₄N₂O₇

¹H NMR (300 MHz, CDCl₃) δ: ppm 4.08 (t, 6H); 3.66 (m, 2H), 3.23-2.53 (br. m, 12H); 2.37-2.30 (m, 6H); 1.74-1.31 (br. m, 48H); 0.93-0.89 (m, 9H).

15

CX. Compound 20-6: Pentyl 6-((2-(dinonylamo)ethyl)(2-hydroxyethyl)amino)hexanoate



(Compound 20-6)

Chemical Formula: C₃₃H₆₈N₂O₃

Molecular Weight: 540.92

20 [00859] In the same manner as pentyl 6-(dodecyl(2-

hydroxyethyl)amino)ethyl)amino)hexanoate, pentyl 6-((2-(dinonylamo)ethyl)(2-


hydroxyethyl)amino)hexanoate was synthesized from pentyl 6-((2-

hydroxyethyl)amino)hexanoate (100 mg, 0.41 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine

(101 mg, 0.41 mmol), K_2CO_3 (108 mg, 0.82 mmol), and KI (7 mg, 0.041 mmol) in MeCN (1 mL) and THF (1 mL). Yield (25 mg, 13%).

UPLC/ELSD: RT = 3.37 min. MS (ES): m/z (MH^+) 541.90 for $C_{33}H_{68}N_2O_3$
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 4.08 (t, 2H); 3.54 (t, 2H), 2.63-2.42 (br. m, 12H); 2.32 (t, 5H); 1.71-1.61 (m, 4H); 1.51-1.46 (m, 6H); 1.35-1.29 (br. m, 30H); 0.95-0.88 (m, 9H).

CX. Compound 20-7: Heptyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate




10 Chemical Formula: $C_{41}H_{84}N_2O_3$

Molecular Weight: 653.13

[00860] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, heptyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate was synthesized from 2-(dodecylamino)ethan-1-ol (100 mg, 0.37 mmol), heptyl 6-((2-chloroethyl)(dodecyl)amino)hexanoate (152 mg, 0.37 mmol), K_2CO_3 (101 mg, 0.73 mmol), and KI (6 mg, 0.037 mmol) in MeCN (2 mL) and THF (2 mL). Yield (41 mg, 17%).

UPLC/ELSD: RT = 3.14 min. MS (ES): m/z (MH^+) 654.0 for $C_{41}H_{84}N_2O_3$
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 4.08 (t, 2H); 3.55 (t, 2H), 2.63-2.45 (br. m, 12H); 2.32 (t, 2H); 1.71-1.59 (m, 4H); 1.54-1.28 (br. m, 52H); 0.92-0.88 (m, 9H).

CY. Compound 20-8: Nonyl 8-((2-(didecylamino)ethyl)(2-hydroxyethyl)amino)octanoate



Chemical Formula: $C_{45}H_{92}N_2O_3$

Molecular Weight: 709.24

[00861] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, nonyl 8-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)octanoate was synthesized from nonyl 8-((2-hydroxyethyl)amino)octanoate (240 mg, 0.73 mmol), *N*-(2-chloroethyl)-*N*-dodecyldecane-1-amine (335 mg, 0.80 mmol), 5 K_2CO_3 (121 mg, 0.88 mmol), and KI (12 mg, 0.072 mmol) in MeCN (1.5 mL) and THF (1.5 mL). Yield (122 mg, 24%).


UPLC/ELSD: RT = 3.41 min. MS (ES): m/z (MH^+) 709.93 for $\text{C}_{45}\text{H}_{92}\text{N}_2\text{O}_3$
 ^1H NMR (300 MHz, CDCl_3) δ : ppm 4.08 (t, 2H); 3.56 (m, 2H), 2.91-2.37 (br. m, 12H); 2.31 (m, 2H); 1.64(br. m, 4H); 1.55-1.20(br. m, 60H); 0.91 (m, 9H).

10

CZ: Compound 20-9: Heptadecan-9-yl 8-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)octanoate

Step 1: Heptadecan-9-yl 8-bromooctanoate

15




[00862] To a solution of 8-bromooctanoic acid (1.04 g, 4.6 mmol) and heptadecan-9-ol (1.5 g, 5.8 mmol) in dichloromethane (20 mL) was added *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (1.1 g, 5.8 mmol), *N,N*-diisopropylethylamine (3.3 mL, 18.7 mmol) and DMAP (114 mg, 0.9 mmol). The reaction was allowed to stir at room temperature 20 for 18 h. The reaction was diluted with dichloromethane and extracted with saturated sodium bicarbonate. The organic layer was separated, washed with brine and dried over MgSO_4 . The organic layer was filtered and the filtrate was evaporated *in vacuo*. The residue was purified by silica gel chromatography (0-10% ethyl acetate in hexanes) to obtain heptadecan-9-yl 8-bromooctanoate (875 mg, 1.9 mmol, 41%).

25

^1H NMR (300 MHz, CDCl_3) δ : ppm 4.89 (m, 1H); 3.42 (m, 2H); 2.31 (m, 2H); 1.89 (m, 2H); 1.73-1.18 (br. m, 36H); 0.88 (m, 6H).

Step 2: Heptadecan-9-yl 8-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)octanoate



Chemical Formula: C₅₃H₁₀₈N₂O₃

Molecular Weight: 821.46

5


[00863] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, heptadecan-9-yl 8-((2-(didodecylamino)ethyl)(2-hydroxyethyl)amino)octanoate was synthesized from heptadecan-9-yl 8-((2-hydroxyethyl)amino)octanoate (100 mg, 0.23 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (94 mg, 0.23 mmol), K₂CO₃ (63 mg, 0.45 mmol), and KI (4 mg, 0.023 mmol) in MeCN (1 mL) and THF (1 mL). Yield (107 mg, 57%).

10 UPLC/ELSD: RT = 3.91 min. MS (ES): *m/z* (MH⁺) 822.3 for C₅₃H₁₀₈N₂O₃

¹H NMR (300 MHz, CDCl₃) δ: ppm 4.89 (p, 1H); 3.56 (m, 2H), 2.62-2.45 (br. m, 12H); 2.30 (t, 2H); 1.88-1.11 (br. m, 78H); 0.92-0.88 (m, 12H).

15

DA. Compound 20-10: Dinonyl 8,8'-(2-(dodecyl(2-hydroxyethyl)amino)ethyl)azanediyl)dioctanoate



Chemical Formula: C₅₀H₁₀₀N₂O₅

Molecular Weight: 809.36


20

[00864] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, dinonyl 8,8'-(2-(dodecyl(2-hydroxyethyl)amino)ethyl)azanediyl)dioctanoate was synthesized from 2-(dodecylamino)ethan-1-ol (100 mg, 0.44 mmol), dinonyl 8,8'-(2-chloroethyl)azanediyl)dioctanoate (269 mg, 0.44

mmol), K_2CO_3 (121 mg, 0.87 mmol), and KI (72 mg, 0.044 mmol) in $MeCN$ (2.5 mL) and THF (2.5 mL). Yield (172 mg, 49%).

UPLC/ELSD: $RT = 4.09$ min. MS (ES): $m/z (MH^+)$ 810.31 for $C_{50}H_{100}N_2O_5$
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 4.07 (t, 4H); 3.55 (m, 2H), 2.64-2.47 (br. m, 12H); 2.31 (t, 5H); 1.66-1.59 (br. m, 8H); 1.46-1.28 (br. m, 60H); 0.92-0.88 (m, 9H).

DB: Compound 20-11: 3-((2-(Ditetradecylamino)ethyl)(dodecyl)amino)propan-1-ol




Chemical Formula: $C_{45}H_{94}N_2O$

10 Molecular Weight: 679.26

[00865] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 3-((2-(ditetradecylamino)ethyl)(dodecyl)amino)propan-1-ol was synthesized from 3-(dodecylamino)propan-1-oloctanoate (50 mg, 0.21 mmol), N -(2-chloroethyl)- N -tetradecyltetradecan-1-amine (109 mg, 0.23 mmol), K_2CO_3 (57 mg, 0.41 mmol), and KI (3.4 mg, 0.021 mmol) in $MeCN$ (1 mL) and THF (1 mL). Yield (65 mg, 46%).

15 UPLC/ELSD: $RT = 3.65$ min. MS (ES): $m/z (MH^+)$ 679.81 for $C_{45}H_{94}N_2O$
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 3.76 (t, 2H); 2.63-2.42 (br. m, 12H), 1.66-1.26 (br. m, 70H); 0.90-0.86 (m, 9H).

20 DC: Compound 20-12: 2-((2-(Ditetradecylamino)ethyl)(tetradecyl)amino)ethan-1-ol



Chemical Formula: $C_{46}H_{96}N_2O$

Molecular Weight: 693.29

25 [00866] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(ditetradecylamino)ethyl)(tetradecyl)amino)ethan-1-ol was synthesized from 2-

(tetradecylamino)ethan-1-ol (100 mg, 0.39 mmol), *N*-(2-chloroethyl)-*N*-tetradecyltetradecan-1-amine (184 mg, 0.39 mmol), K_2CO_3 (107 mg, 0.78 mmol), and KI (6.5 mg, 0.039 mmol) in MeCN (2 mL) and THF (2 mL). Yield (87 mg, 32%).


UPLC/ELSD: RT = 3.81 min. MS (ES): m/z (MH^+) 694.02 for $C_{46}H_{96}N_2O$

5 1H NMR (300 MHz, $CDCl_3$) δ : ppm 3.56 (m, 2H); 2.61-2.45 (br. m, 12H), 1.47-1.29 (br. m, 72H); 0.91 (m, 9H).

DD: Compound 20-13: 2-((2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)(dodecyl)amino)ethan-1-ol

10

Step 1: (6Z,9Z)-18-(Methylsulfonyl)octadeca-6,9-diene



Chemical Formula: $C_{19}H_{36}O_3S$

15

Molecular Weight: 344.55


[00867] To a 0 °C solution of linoleyl alcohol (10 mL, 31.2 mmol) and triethylamine (5.68 mL, 40.5 mmol) in DCM (50 mL) was added dropwise a solution of methanesulfonyl chloride (2.66 mL, 34.3 mmol) in DCM (20 mL). The reaction was allowed to return to room temperature and let stir for 4 hours. The mixture was quenched by the addition of water and extracted with DCM. The organic layer was washed with saturated $NaHCO_3$, brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography (0-40% EtOAc/hexanes) provided (6Z,9Z)-18-(methylsulfonyl)octadeca-6,9-diene (10.0 g, 93%).

1H-NMR (300 MHz, $CDCl_3$) δ : ppm 5.35 (m, 4H); 4.22 (t, 2H); 2.99 (s, 3H); 2.77 (t, 2H); 2.04

25

(q, 4H); 1.74 (m, 2H); 1.30 (br. m, 16H); 0.89 (t, 3H).

Step 2: (6Z,9Z)-18-Bromoocatadeca-6,9-diene



Chemical Formula: $C_{18}H_{33}Br$

30


Molecular Weight: 329.37

[00868] To a solution of (6Z,9Z)-18-(methylsulfonyl)octadeca-6,9-diene (10.0 g, 29.0 mmol) in diethyl ether (372 mL) was added magnesium bromide ethyl etherate (22.5 g, 87.1 mmol). The reaction was let stir at room temperature for 16 hours. The mixture was quenched by the addition of water and extracted with diethyl ether. The combined organic layers were 5 washed with 1% K_2CO_3 , brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography provided (6Z,9Z)-18-bromo-octadeca-6,9-diene (8.9 g, 93%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.36 (m, 4H); 3.41 (t, 2H); 2.77 (t, 2H); 2.05 (q, 4H); 1.86 (m, 2H); 1.48-1.22 (br. m, 16H); 0.89 (t, 3H).

10

Step 3: 2-((2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)(dodecyl)amino)ethan-1-ol



Chemical Formula: $\text{C}_{52}\text{H}_{100}\text{N}_2\text{O}$

Molecular Weight: 769.39


[00869] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)(dodecyl)amino)ethan-1-ol was synthesized from 2-(dodecylamino)ethan-1-ol (50 mg, 0.22 mmol), (9Z,12Z)-*N*-(2-chloroethyl)-*N*-(9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine (126 mg, 0.22 mmol), K_2CO_3 (60 mg, 0.44 mmol), and KI (3.6 mg, 0.022 mmol) in MeCN (2 mL). Yield (33 mg, 20%).

UPLC/ELSD: RT = 3.74 min. MS (ES): m/z (MH^+) 770.20 for $\text{C}_{52}\text{H}_{100}\text{N}_2\text{O}$

$^1\text{H NMR}$ (300 MHz, CDCl_3) δ : ppm 5.39 (m, 8H); 3.55 (m, 2H), 2.80 (m, 4H); 2.61-2.44 (br. m, 12H); 2.07 (m, 8H); 1.46-1.29 (br. m, 56H); 0.92 (m, 9H).

25 DE: Compound 20-14: 2-((2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol

Step 1: 2-((9Z,12Z)-octadeca-9,12-dien-1-yl)aminoethan-1-ol



Chemical Formula: C₂₀H₃₉NO

Molecular Weight: 309.54


[00870] In the same manner as 2-(dodecylamino)ethan-1-ol, 2-((9Z,12Z)-octadeca-9,12-dien-1-yl)aminoethan-1-ol was synthesized from ethanolamine (0.37 mL, 6.1 mmol), (6Z,9Z)-18-bromoocatadeca-6,9-diene (2.0 g, 6.1 mmol), K₂CO₃ (1.67 g, 12.1 mmol), and KI (101 mg, 0.607 mmol) in MeCN (28 mL). Yield (453 mg, 24%).

UPLC/ELSD: RT = 5.457 min. MS (ES): *m/z* (MH⁺) 311.38 for C₂₀H₃₉NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.36 (m, 4H); 3.62 (t, 2H); 2.78 (m, 4H); 2.61 (t, 2H); 2.05 (m, 4H); 1.49 (m, 2H); 1.30 (br. m, 16H); 0.89 (t, 3H).

10

Step 2: 2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol



Chemical Formula: C₃₈H₇₁NO

Molecular Weight: 557.99

15


[00871] In the same manner as 2-(didodecylamino)ethan-1-ol, 2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol was synthesized from (6Z,9Z)-18-bromoocatadeca-6,9-diene (4 g, 12.1 mmol), ethanolamine, (0.334 mL, 5.52 mmol), K₂CO₃ (3.36 g, 24.3 mmol), and KI (92 mg, 0.552 mmol) in MeCN (26 mL). Yield (1.9 g, 62%).

UPLC/ELSD: RT = 6.80 min. MS (ES): *m/z* (MH⁺) 557.94 for C₃₈H₇₁NO

20

¹H-NMR (300 MHz, CDCl₃) δ: ppm 5.35 (m, 8H); 3.52 (t, 2H); 2.77 (t, 4H); 2.57 (t, 2H); 2.43 (t, 4H); 2.04 (q, 8H); 1.48-1.18 (br. m, 36H); 0.89 (t, 6H).

Step 3: (9Z,12Z)-N-(2-chloroethyl)-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine



Chemical Formula: C₃₈H₇₀ClN

Molecular Weight: 576.44


25

[00872] In a same manner as compound N-(2-chloroethyl)-N-dodecyldecane-1-amine, (9Z,12Z)-N-(2-chloroethyl)-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine

was synthesized from 2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol (250 mg, 0.45 mmol), triethylamine (81 μ L, 0.58 mmol), and methanesulfonyl chloride (38 μ L, 0.49 mmol) in DCM (2 mL). Yield (134 mg, 52%).

5 1 H-NMR (300 MHz, CDCl₃) δ : ppm 5.36 (m, 8H); 3.49 (t, 2H); 2.78 (m, 6H); 2.45 (t, 4H); 2.05 (q, 8H); 1.48-1.18 (br. m, 36H); 0.89 (t, 6H).

Step 4; 2-((2-(Di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol



10 Chemical Formula: C₅₈H₁₀₈N₂O


Molecular Weight: 849.52

[00873] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(di((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol was synthesized from 2-((9Z,12Z)-octadeca-9,12-dien-1-yl)aminoethan-1-ol (75 mg, 0.24 mmol), (9Z,12Z)-N-(2-chloroethyl)-N-((9Z,12Z)-octadeca-9,12-dien-1-yl)octadeca-9,12-dien-1-amine (154 mg, 0.27 mmol), K₂CO₃ (67 mg, 0.49 mmol), and KI (4 mg, 0.024 mmol) in MeCN (2 mL). Yield (35 mg, 17%).

UPLC/ELSD: RT = 3.94 min. MS (ES): *m/z* (MH⁺) 850.03 for C₅₈H₁₀₈N₂O

20 1 H-NMR (300 MHz, CDCl₃) δ : ppm 5.35 (br. m, 12H); 2.77 (t, 6H) 2.70-2.38 (br. m, 14H); 2.05 (m, 12H); 1.50-1.00 (br. m, 54H); 0.88 (t, 9H).

DF: Compound 20-15: 2-((2-(Didodecylamino)ethyl)(hexyl)amino)ethan-1-ol




25 Chemical Formula: C₃₄H₇₂N₂O

Molecular Weight: 524.96

[00874] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(didodecylamino)ethyl)(hexyl)amino)ethan-1-ol was synthesized from 2-(hexylamino)ethan-1-ol (50 mg, 0.34 mmol), *N*-(2-chloroethyl)-*N*-dodecylidodecan-1-amine (143 mg, 0.34 mmol), K_2CO_3 (95 mg, 0.69 mmol), and KI (5.7 mg, 0.034 mmol) in MeCN (2 mL). Yield (145 mg, 80%).

5 UPLC/ELSD: RT = 2.73min. MS (ES): m/z (MH^+) 525.66 for $C_{34}H_{72}N_2O$
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 3.54 (m, 2H); 2.61-2.44 (br. m, 12H), 1.46-1.28 (br. m, 48H); 0.90 (m, 9H).

10 DG: Compound 20-16: 2-((2-(Dinonylamino)ethyl)(nonyl)amino)ethan-1-ol




Chemical Formula: $C_{31}H_{66}N_2O$

Molecular Weight: 482.88

[00875] To a solution of 2-((2-aminoethyl)amino)ethan-1-ol (2.0 g, 18.6 mmol) and DCE (50 mL) at 0 °C was added nonanal (12.8 mL, 74.6 mmol), followed by AcOH (3.2 mL, 55.9 mmol). The reaction was allowed to stir at 0 °C for 20 min. $Na(OAc)_3BH$ (15.8 g, 74.6 mmol) was added and the reaction was allowed to warm to room temperature and stir for 18 hours at room temperature. The mixture was quenched by the slow addition of aqueous saturated $NaHCO_3$ and extracted with DCM three times. The combined organic layers were dried over anhydrous Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification by ISCO silica flash chromatography provided 2-((2-(dinonylamino)ethyl)(nonyl)amino)ethan-1-ol. Yield (75 mg, 0.8%).

15 UPLC/ELSD: RT = 2.28 min. MS (ES): m/z (MH^+) 483.47 for $C_{21}H_{66}N_2O$
 1H NMR (300 MHz, $CDCl_3$) δ : ppm 3.53 (m, 2H); 2.61-2.41 (br. m, 12H), 1.43-1.25 (br. m, 42H); 0.86 (m, 9H).

DH: Compound 20-17: 2-((2-(Didodecylamino)ethyl)(nonyl)amino)ethan-1-ol




Chemical Formula: C₃₇H₇₈N₂O

Molecular Weight: 567.04

[00876] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(didodecylamino)ethyl)(nonyl)amino)ethan-1-ol was synthesized from 2-(nonylamino)ethan-1-ol (50 mg, 0.27 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (111 mg, 0.27 mmol), K_2CO_3 (74 mg, 0.53 mmol), and KI (4.4 mg, 0.027 mmol) in 1,4-dioxane (1.5 mL). Yield (29 mg, 19%).

10 UPLC/ELSD: RT = 3.05 min. MS (ES): m/z (MH $^+$) 567.91 for C₃₇H₇₈N₂O
¹H NMR (300 MHz, CDCl₃) δ : ppm 3.71 (m, 2H); 3.14-2.97 (br. m, 8H), 2.80 (m, 2H); 2.66 (m, 2H); 1.70 (m, 4H); 1.53 (m, 2H); 1.34-1.28 (br. m, 48H); 0.90 (m, 9H).

DI: Compound 20-18: 2-((2-(Dinonylamo)ethyl)(dodecyl)amino)ethan-1-ol




Chemical Formula: C₃₄H₇₂N₂O

Molecular Weight: 524.96

[00877] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(dinonylaminoo)ethyl)(dodecyl)amino)ethan-1-ol was synthesized from 2-(dodecylamino)ethan-1-ol (100 mg, 0.44 mmol), *N*-(2-chloroethyl)-*N*-nonylnonan-1-amine (145 mg, 0.44 mmol), K_2CO_3 (120 mg, 0.87 mmol), and KI (7.2 mg, 0.044 mmol) in MeCN (1 mL) and THF (1 mL). Yield (155 mg, 67%).

UPLC/ELSD: RT = 2.78 min. MS (ES): m/z (MH⁺) 525.99 for C₃₄H₇₂N₂O

¹H NMR (300 MHz, CDCl₃) δ: ppm 3.55 (m, 2H); 2.63-2.47 (br. m, 12H), 1.47-1.28 (br. m, 48H); 0.90 (m, 9H).



Chemical Formula: $C_{28}H_{60}N_2O$

Molecular Weight: 440.80


[00878] Ethanolamine (50 mg, 0.82 mmol), *N*-(2-chloroethyl)-*N*-dodecylidodecan-1-

5 amine (0.75 g, 1.8 mmol), K_2CO_3 (0.25 g, 1.8 mmol) and KI (14 mg, 0.082) and 4 mL THF were combined in a round bottomed flask. The reaction was placed in a 65 °C heating mantle and was allowed to stir under N_2 for 12 h. After this time the reaction was allowed to cool to room temperature and was filtered. The filtrate was washed with H_2O and brine, dried over $MgSO_4$, filtered and concentrated. The crude material was purified by C18 reverse phase chromatography (5-100% MeCN in H_2O with 0.1% TFA). The fractions were pooled and concentrated. The isolated material was taken up in $CHCl_3$, washed with 10% NaOH and brine, dried over $MgSO_4$, filtered and concentrated. The product was then re-purified *via* silica gel chromatography (0-20% MeOH in DCM with 1% NH_4OH) to afford 2-((2-(didecylamino)ethyl)amino)ethan-1-ol (0.15 g, 41%).

10 UPLC/ELSD: RT = 2.15 min. MS (ES): m/z (MH^+) 441.37 for $C_{28}H_{60}N_2O$

1H -NMR (400 MHz, $CDCl_3$) δ : ppm 3.59 (t, 2H); 2.75 (t, 2H); 2.62 (t, 2H); 2.50 (t, 2H); 2.37 (t, 4H); 1.39 (m, 4H); 1.24 (m 38H); 0.86 (t, 6H).

DK: Compound 20-20: 2-((2-(Didodecylamino)ethyl)(dodecyl)amino)ethan-1-ol



Chemical Formula: $C_{40}H_{84}N_2O$

Molecular Weight: 609.13

[00879] A solution of 2-((2-aminoethyl)amino)ethan-1-ol (2 g, 19.2 mmol) in 50 mL DCE

was allowed to cool under N_2 in an ice bath. Dodecanal (26 mL, 76.8 mmol) was added


25 followed by acetic acid (3.3 mL, 57.6 mmol). After 20 min., $Na(OAc)_3BH$ (16.3 g, 76.8 mmol) was added. The reaction was allowed to slowly warm to room temperature and stir for 48 h. After this time the reaction was quenched *via* portion wise addition of saturated $NaHCO_3$. The mixture was extracted three times with DCM. The pooled organics were washed with brine,

dried over MgSO_4 , filtered and concentrated. The crude material was purified by silica gel chromatography (0-100% EtOAc in hexanes) twice to afford clean 2-((2-(didodecylamino)ethyl)(dodecyl)amino)ethan-1-ol (7.4 g, 63%).

UPLC/ELSD: RT = 3.20 min. MS (ES): m/z (MH^+) 609.97 for $\text{C}_{40}\text{H}_{84}\text{N}_2\text{O}$

5 $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : ppm 3.51 (t, 2H); 2.57-2.40 (br. m, 12H); 1.41-1.23 (br. m, 60H); 0.86 (t, 9H).

DL: Compound 20-21: 3-((2-(Didodecylamino)ethyl)(dodecyl)amino)propan-1-ol



10 Chemical Formula: $\text{C}_{41}\text{H}_{86}\text{N}_2\text{O}$


Molecular Weight: 623.15

[00880] In the same manner as pentyl 6-(dodecyl(2-(hydroxyethyl)amino)ethyl)amino)hexanoate, 3-((2-(didodecylamino)ethyl)(dodecyl)amino)propan-1-ol was synthesized from 3-(dodecylamino)propan-1-ol (39 mg, 0.16 mmol), *N*-(2-chloroethyl)-*N*-dodecyl dodecan-1-amine (75 mg, 0.18 mmol), K_2CO_3 (44 mg, 0.32 mmol), and KI (2.7 mg, 0.016 mmol) in THF (1 mL). Yield (170 mg, >98%).

15 UPLC/ELSD: RT = 3.29 min. MS (ES): m/z (MH^+) 623.71 for $\text{C}_{41}\text{H}_{86}\text{N}_2\text{O}$

$^1\text{H NMR}$ (300 MHz, CDCl_3) δ : ppm 3.76 (m, 2H); 2.64-2.39 (br. m, 12H), 1.66 (m, 2H); 1.44-1.26 (br. m, 60H); 0.88 (m, 9H).

20 DM: Compound 20-22: 4-((2-(Didodecylamino)ethyl)(dodecyl)amino)butan-1-ol
4-(Dodecylamino)butan-1-ol



25 Chemical Formula: $\text{C}_{16}\text{H}_{35}\text{NO}$


Molecular Weight: 257.46

[00881] In the same manner as 2-(dodecylamino)ethan-1-ol, 4-(dodecylamino)butan-1-ol was synthesized from 4-aminobutan-1-ol (2.5 mL, 27 mmol), 1-bromododecane (6.75 g, 27 mmol), K_2CO_3 (7.5 g, 54 mmol), and KI (450 mg, 2.7 mmol) in MeCN (125 mL). Yield (303 mg, 4%).

UPLC/ELSD: RT = 1.09 min. MS (ES): m/z (MH⁺) 258.22 for C₁₆H₃₅NO

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.60 (t, 2H); 2.76-2.62 (br. m, 4H); 1.72-1.58 (br. m, 6H); 1.29 (br. m, 18H); 0.89 (t, 3H).

5



Chemical Formula: C₄₂H₈₈N₂O

Molecular Weight: 637.18

[00882] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-

10 hydroxyethyl)amino)ethyl)amino)hexanoate,
4-((2-(didodecylamino)ethyl)(dodecyl)amino)butan-1-ol was synthesized from 4-(dodecylamino)butan-1-ol (75 mg, 0.29 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (133 mg, 0.32 mmol), K₂CO₃ (80 mg, 0.58 mmol), and KI (5 mg, 0.029 mmol) in MeCN (2 mL). Yield (104 mg, 56%).


15 UPLC/ELSD: RT = 3.27 min. MS (ES): m/z (MH⁺) 637.85 for C₄₂H₈₈N₂O

¹H-NMR (300 MHz, CDCl₃) δ: ppm 3.56 (br. m, 2H); 2.58 (br. m, 4H); 2.45 (br. m, 8H); 1.65 (br. m, 4H); 1.45 (br. m, 6H); 1.25 (br. m, 54H); 0.88 (t, 9H).

DN: Compound 20-23: (Z)-2-((2-(Didodecylamino)ethyl)(dodec-6-en-1-yl)amino)ethan-1-ol

20

Step 1: (6-Hydroxyhexyl)triphenylphosphonium bromide



Chemical Formula: C₂₄H₂₈BrOP


Molecular Weight: 443.36

25 [00883] 6-Bromo-1-hexanol (4.89 g, 27 mmol) and triphenylphosphine (7.87 g, 30 mmol) and 50 mL MeCN were combined in a round bottomed flask. The flask was fitted with a condenser and placed in a heating mantel and the reaction was allowed to stir at 82 °C for 48 h. After this time the reaction was allowed to cool to room temperature and the solution was cannulated into 200 mL Et₂O, producing a white precipitate. The solids were allowed to settle

and the solvent was decanted off. 20 mL DCM was added to dissolve the solids and then 100 mL Et₂O was slowly added to afford a white precipitate. The solvent was then removed *in vacuo* to afford clean (6-hydroxyhexyl)triphenylphosphonium bromide (9.4 g, 21.2 mmol, for 78% yield).

5 ¹H-NMR (300 MHz, CDCl₃) δ: ppm 7.80 (m, 15H); 3.80 (m, 2H); 3.65 (m, 2H); 2.23 (m, 2H); 1.68 (m, 4H); 1.52 (m, 4H).

Step 2: (Z)-Dodec-6-en-1-ol



10 Chemical Formula: C₁₂H₂₄O
Molecular Weight: 184.32

[00884] A solution of (6-hydroxyhexyl)triphenylphosphonium bromide (3.0 g, 6.77 mmol) in 25 mL THF was allowed to cool in a -78 °C dry ice/acetone bath. Once cool *n*-BuLi (2.5 M in hexanes) (5.7 mL, 14.2 mmol) was added dropwise. After 1 h, an additional 10 mL 15 THF and *n*-BuLi (1.35 mL) were added and stirring was continued at the same temperature for 1 h. After this time 1-hexanal (1.6 mL, 13.5 mmol) was added and the reaction was allowed to warm to rt and stir for 3 h. After this time the reaction was quenched by addition of excess sat'd NH₄Cl. The solution was extracted three times with EtOAc. The pooled organics were washed with brine, dried over MgSO₄, filtered and concentrated *in vacuo*. The crude material was 20 purified by silica gel chromatography (0-50% EtOAc in hexanes) to afford the desired product as a clear oil (0.76 g, 4.1 mmol, 61%).

1H-NMR (300 MHz, CDCl₃) δ: ppm 5.34 (m, 2H); 3.62 (t, 2H); 2.01 (m, 4H); 1.56 (m, 2H); 1.35-1.27 (m, 11H); 0.87 (t, 3H).

25 Step 3: (Z)-Dodec-6-en-1-yl methanesulfonate




Chemical Formula: C₁₃H₂₆O₃S
Molecular Weight: 262.41

[00885] To a 0 °C solution of (Z)-dodec-6-en-1-ol (1.81 g, 9.3 mmol) in 20 mL DCM, 30 was added Et₃N (1.7 mL, 12.1 mmol) and methanesulfonyl chloride (0.80 mL, 10.2 mmol). The reaction was allowed to slowly warm to room temperature and stir overnight. The reaction was

quenched by the addition of water and the mixture was extracted two times with DCM. The organics were pooled, washed with brine, dried over MgSO_4 , filtered and concentrated. The crude material was purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford clean desired product (2.2 g, 8.4 mmol, 90%).

5 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.34 (m, 2H); 4.20 (t, 2H); 2.98 (s, 3H); 2.01 (m, 4H); 1.74 (m, 2H); 1.38-1.27 (m, 10H); 0.87 (t, 3H).

Step 4: (Z)-1-Bromododec-6-ene




10 Chemical Formula: $\text{C}_{12}\text{H}_{23}\text{Br}$
Molecular Weight: 247.22

[00886] In a round bottomed flask, under N_2 , (Z)-dodec-6-en-1-yl methanesulfonate (2.2 g, 8.3 mmol) was dissolved in 40 mL Et_2O . $\text{MgBr}_2 \cdot \text{Et}_2\text{O}$ (6.5 g, 25 mmol) was added and the reaction was allowed to stir for 48 h. After this time the reaction was quenched by the addition 15 of ice. The mixture was then extracted with Et_2O three times. The pooled organics were washed with brine, dried over MgSO_4 , filtered and concentrated. The crude material was purified by silica gel chromatography (0-30% EtOAc in hexanes) to afford the desired product (1.8 g, 7.28 mmol, 88%).

20 $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ : ppm 5.34 (m, 2H); 3.39 (t, 2H); 2.01-1.84 (m, 6H); 1.28 (m, 10H); 0.87 (t, 3H).

Step 5: (Z)-2-((2-(Didodecylamino)ethyl)(dodec-6-en-1-yl)amino)ethan-1-ol



Chemical Formula: $\text{C}_{40}\text{H}_{82}\text{N}_2\text{O}$


25 Molecular Weight: 607.11

[00887] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, (Z)-2-((2-(didodecylamino)ethyl)(dodec-6-en-1-yl)amino)ethan-1-ol was synthesized from (Z)-2-(dodec-6-en-1-ylamino)ethan-1-ol (100 mg, 0.44 mmol), *N*-(2-chloroethyl)-*N*-dodecyl dodecan-1-amine (183 mg, 0.44 mmol), K_2CO_3 (122

mg, 0.88 mmol), and KI (7.3 mg, 0.044 mmol) in MeCN (1 mL) and THF (1 mL). Yield (90 mg, 34%).

UPLC/ELSD: RT = 3.24 min. MS (ES): m/z (MH⁺) 608.08 for C₄₀H₈₂N₂O
¹H NMR (300 MHz, CDCl₃) δ: ppm 5.42-5.35 (m, 2H); 3.55 (m, 2H), 2.62-2.45 (br. m, 12H);
5 2.06-2.00 (m, 4H); 1.48-1.28 (br. m, 52H); 0.91 (m, 9H).

DO: Compound 20-24: 2-((2-(Didodecylamino)ethyl)(tetradecyl)amino)ethan-1-ol




Chemical Formula: C₄₂H₈₈N₂O

Molecular Weight: 637.18

[00888] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(didodecylamino)ethyl)(tetradecyl)amino)ethan-1-ol was synthesized from 2-(tetradecylamino)ethan-1-ol (100 mg, 0.39 mmol), N-(2-chloroethyl)-N-dodecyldecane-1-amine (162 mg, 0.39 mmol), K₂CO₃ (107 mg, 0.78 mmol), and KI (6.5 mg, 0.039 mmol) in MeCN (3 mL). Yield (128 mg, 52%).

UPLC/ELSD: RT = 3.47 min. MS (ES): m/z (MH⁺) 637.92 for C₄₂H₈₈N₂O
¹H NMR (300 MHz, CDCl₃) δ: ppm 3.54 (m, 2H); 2.61-2.44 (br. m, 12H); 1.46-1.28 (br. m, 64H); 0.91 (m, 9H).

DP: Compound 20-25: 2-((2-(Didodecylamino)ethyl)((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol



Chemical Formula: C₄₆H₉₂N₂O

Molecular Weight: 689.26

[00889] In the same manner as pentyl 6-(dodecyl(2-(dodecyl(2-hydroxyethyl)amino)ethyl)amino)hexanoate, 2-((2-(didodecylamino)ethyl)((9Z,12Z)-octadeca-

9,12-dien-1-yl)amino)ethan-1-ol was synthesized from 2-(((9Z,12Z)-octadeca-9,12-dien-1-yl)amino)ethan-1-ol (50 mg, 0.16 mmol), *N*-(2-chloroethyl)-*N*-dodecyldodecan-1-amine (67 mg, 0.16 mmol), K_2CO_3 (45 mg, 0.32 mmol), and *KI* (3 mg, 0.016 mmol) in MeCN (2 mL). Yield (45 mg, 41%).

5 UPLC/ELSD: RT = 3.64 min. MS (ES): *m/z* (MH^+) 689.95 for $C_{46}H_{92}N_2O$
¹H NMR (300 MHz, $CDCl_3$) δ : ppm 5.39-5.32 (m, 4H); 3.56 (m, 2H), 2.80 (m, 2H); 2.62-2.52 (br. m, 12H); 2.08 (m, 4H); 1.48-1.28 (br. m, 58H); 0.91 (m, 9H).


DQ. Compound 21-1: 1-(2,2-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopropyl)-*N,N*-

10 dimethylmethanamine




[0011] At 0 °C, a solution of nonane-1,9-diol (96.16 g, 0.60 mol) in 100 mL DMF was slowly added into a suspension of NaH (24.0 g, 0.60 mol) in 800 mL DMF. After stirring for 1 h, a

15 solution of benzyl bromide (71.4 mL, 0.60 mole) in 200 mL DMF was slowly added. After addition, the reaction mixture was warmed up to room temperature and stirred overnight. TLC showed starting material was almost consumed. The reaction mixture was poured onto ice, and then extracted with EtOAc (3 X). The combined organic layers were washed with water and brine, and then dried over sodium sulfate. After filtration and concentration, the crude was 20 purified by flash column chromatography (SiO_2 : 0 to 100% EtOAc/hexanes then 0 to 5% MeOH/dichloromethane) to provide the product as a colorless oil (74.4 g, 50%).




25 [0012] At 0 °C, to a solution of 9-(benzyloxy)nonan-1-ol (14.88 g, 61.5 mmol) in 150 mL dichloromethane, CBr_4 (30.6 mmol, 92.2 mmol) was added. And then triphenylphosphine (27.4 g, 0.104 mole) was added portionwise. After stirring at room temperature overnight, TLC showed completed reaction. The reaction mixture was poured onto ice, and then extracted with dichloromethane (2 X). The combined organic layers were washed with water and brine, and 30 then dried over magnesium sulfate. After filtration and concentration, the residue was purified

by ISCO (SiO₂: 0 to 10% EtOAc/hexanes) to provide the product as colorless oil (21.0 g, quant.).




5 [0013] To a suspension of magnesium (3.23 g, 0.134 mole) in 80 mL THF, catalytic amount of iodine was added, and then stirred until the color disappeared. A solution of (((9-bromononyl)oxy)methyl)benzene (21.0 g, 67.2 mmol) in 40 mL THF was slowly added in 15 min at room temperature, and then the mixture was heated to reflux for 1 h. After cooling to room temperature, a solution of methyl formate (4.2 mL, 67.2 mmol) in 10 mL THF was added 10 dropwise, and the mixture was stirred overnight. The reaction was quenched by addition of 5 N HCl and water, and the mixture was extracted with EtOAc (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was dissolved in EtOH, and then KOH and water were added. After stirring overnight, the reaction mixture was concentrated to dryness. Water was added, and then adjusted pH ~7 15 with 1 N HCl. The mixture was extracted with EtOAc (2 X), and the combined organic layer was dried over magnesium sulfate. After filtration and concentration, the crude was purified by ISCO (SiO₂: 0 to 10% EtOAc/hexanes) to provide the product as colorless oil (6.64 g, 40%).



20


[0014] At 0 °C, a solution of 1,19-bis(benzyloxy)nonadecan-10-ol (6.64 g, 13.4 mmol) in 30 mL dichloromethane was slowly added into a solution of Dess-Martin periodinane (7.94 g, 18.7 mmol) in 70 mL dichloromethane, and then the reaction mixture was stirred at this temperature for 3 h. TLC showed starting material was consumed. The reaction mixture was diluted with dichloromethane, and then 10% Na₂S₂O₃ solution and saturated sodium bicarbonate solution were added. After extraction with dichloromethane (2 X), the combined organic layers were washed with brine and concentrated. The residue was dissolved in ether and washed with saturated sodium bicarbonate and brine. After drying over sodium sulfate, the solution was

filtered and concentrated. The crude was purified by ISCO (SiO_2 : 0 to 20% EtOAc/hexanes) to provide the product as colorless oil (6.23 g, 94%).




5 [0015] At 0 °C, potassium *tert*-butoxide (1.70 g, 15.1 mmol) was added into a solution of methylphosphonium bromide (5.40 g, 15.1 mmol) in 80 mL THF which was purged with nitrogen 3 times. After 1 h, a solution of 1,19-bis(benzyloxy)nonadecan-10-one (6.23 g, 12.6 mmol) in 20 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight.

10 TLC showed completed reaction, and the reaction mixture was filtered through Celite. After concentration, the crude was purified by ISCO (SiO_2 : 0 to 10% EtOAc/hexanes) to provide the product as colorless oil (6.0 g, 96%).




15 [0016] To a refluxing solution of (((10-methylenenonadecane-1,19-diy1)bis(oxy))bis(methylene))dibenzene (2.99 g, 6.08 mmol) in 160 mL dichloromethane, a solution of $\text{Cu}(\text{acac})_2$ (180 mg, 0.69 mmol) in 40 mL dichloromethane was added. And then ethyl diazoacetate (contains 13% dichloromethane, 9 X 1.1 mL) was added every 30 min. MS showed the formation of product. The reaction was quenched with MeOH and stirred for 1 h at room temperature. After concentration, the crude was purified by flash column chromatography (SiO_2 : 0 to 10% EtOAc/hexanes) to provide the product as colorless oil (3.65 g, contains 1 equivalent of acetate by-product).




20 [0017] A mixture of ethyl 2,2-bis(9-(benzyloxy)nonyl)cyclopropane-1-carboxylate (2.8 g, 4.8 mmol) and Palladium on carbon (10wt%, 500 mg) in 500 mL EtOAc was stirred at room temperature under hydrogen balloon for 4.5 h. MS and TLC showed completed reaction. The reaction mixture was filtered through Celite and washed with EtOAc. The filtrate was

concentrated to provide the product mixed with by-product, diethyl succinate (2.53 g, contains 0.94 equivalent of by-product, 94%).




5 [0018] At 0 °C, a solution of ethyl 2,2-bis(9-hydroxynonyl)cyclopropane-1-carboxylate (2.4 g, 6 mmol) in 100 mL dichloromethane was slowly added into a suspension of Dess-Martin periodinane (7.67 g, 18 mmol) in 100 mL dichloromethane, and then the reaction mixture was stirred at room temperature for 4 h. After quenching with 10% aqueous $\text{Na}_2\text{S}_2\text{O}_3$ and saturated sodium bicarbonate, the mixture was extracted with dichloromethane (2 X). The combined 10 organic layers were dried over sodium sulfate and concentrated to give the product as colorless oil (2.3 g, contains about 0.7 g by-product).

Preparation of (Z)-non-3-en-1-yltriphenylphosphonium iodide:




15 At 0 °C, a solution of triphenylphosphine (110 g, 0.419 mole) in 200 mL dichloromethane was slowly added into a solution of (Z)-non-3-en-1-ol (49.6 g, 0.349 mole), imidazole (50.0 g, 0.732 mole) and iodine (124 g, 0.488 mole) in 800 mL dichloromethane, and then the reaction mixture was allowed to room temperature overnight. TLC showed small amount of (Z)-non-3-en-1-ol 20 left. The reaction mixture was concentrated, and the residue was triturated with hexanes. The solution was filtered through a plug of silica gel and eluted with hexanes to provide the iodide as a colorless liquid (81 g, 92%).


25 [0019] A solution of (Z)-1-iodonon-3-ene (81 g, 0.321 mole) and triphenylphosphine (169 g, 0.643 mole) in acetonitrile (1.1 L) was refluxed overnight. After concentrated to dryness, the residue was triturated with hexanes. The white gum was dissolved in dichloromethane and purified by flash column chromatography (SiO_2 : 0 to 5% $\text{MeOH}/\text{CH}_2\text{Cl}_2$) to provide the product as colorless oil which then turned into white solid (114 g, 69%).




[0020] At 0 °C, potassium *tert*-butoxide (1.98 g, 17.7 mmol) was added into a solution of (Z)-non-3-en-1-yltriphenylphosphonium iodide (9.2 g, 17.9 mmol) in 300 mL THF which was purged with nitrogen 3 times. After 1 h, a solution of ethyl 2,2-bis(9-oxononyl)cyclopropane-1-carboxylate (2.05 g, 5.2 mmol) in 100 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed completed reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with hexanes (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by ISCO (SiO₂: 0 to 5% EtOAc/hexanes) to provide the product as a colorless oil (1.54 g, 48%).



[0021] A solution of lithium aluminum hydride (2.0 M in THF, 1.9 mL, 3.8 mmol) was slowly added into a solution of ethyl 2,2-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopropane-1-carboxylate (1.54 g, 2.52 mmol) in 150 mL THF, and then the reaction mixture was stirred at room temperature for 30 min. TLC showed completed reaction. The reaction was quenched by slow addition of Na₂SO₄ 10H₂O, then the mixture was filtered and washed with THF. The filtrate was concentrated and purified by flash column chromatography (SiO₂: 0 to 15% EtOAc/Hexanes) to give the product as a colorless oil (1.2 g, 84%).



[0022] At 0 °C, a solution of (2,2-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopropyl)methanol (1.04 g, 1.83 mmol) in 100 mL dichloromethane was slowly added into a suspension of Dess-Martin periodinane (1.18 g, 2.77 mmol) in 200 mL dichloromethane, and then the reaction mixture was stirred at room temperature for 3 h. After quenched with saturated sodium bicarbonate, the mixture was extracted with dichloromethane (2 X). The combined organic layer was dried over sodium sulfate and concentrated to give the product as a colorless oil (0.95 g, 91%).




[0023] To a solution of 2,2-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopropane-1-carbaldehyde (0.95 g, 1.68 mmol) in 300 mL THF, dimethylamine (2.0 M in THF, 2 mL, 4 mmol), sodium triacetoxyborohydride (840 mg, 4 mmol) and acetic acid (0.23 mL, 4 mmol) were added sequentially, and the reaction mixture was stirred at room temperature overnight. MS showed completed reaction, and saturated sodium bicarbonate was added to quench the reaction. The mixture was extracted with EtOAc (2 X), and the combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by flash column chromatography (SiO₂: 0 to 10% MeOH/dichloromethane) to give the product 1-(2,2-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopropyl)-N,N-dimethylmethanamine as a colorless oil (0.56 g, 56%).

¹H NMR (300 MHz, CDCl₃) δ 5.27-5.42 (m, 8 H), 2.76 (t, 4 H, *J* = 6.2 Hz), 2.38 (bs, 8 H), 2.04 (q, 8 H, *J* = 6.6 Hz), 1.18-1.41 (m, 38 H), 0.96-1.17 (m, 2 H), 0.88 (t, 6 H, *J* = 6.6 Hz), 0.66-0.76 (m, 1 H), 0.48-0.56 (m, 1 H), 0.05-0.13 (m, 1 H).

APCI: m/z = 596.6 [M + H]⁺.


5

DR. Compound 21-2: 3,3-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclobutyl 4-(dimethylamino)butanoate




10 [0024] Preparation of Zn-Cu couple: A suspension of zinc dust (10 g) in 10 mL 4 M HCl was stirred for 10 min, and then the aqueous phase was decanted. After the solid was washed with water (2 X 20 mL), 20 mL water and copper sulfate (0.75 g) were added subsequently. After stirring overnight, water was decanted, and then the residue was washed with THF (2 X 10 mL). The black solid was dried under vacuum and store under nitrogen.


15 [0025] To a suspension of Zn-Cu couple (1.295 g, 19.8 mmol) in 30 mL ether purged with nitrogen 3 times, a solution of (((10-methylenonadecane-1,19-diy)bis(oxy))bis(methylene))dibenzene (2.96 g, 6.0 mmol) in 10 mL ether purged with nitrogen was added, and then a solution of POCl₃ (1.85 mL, 19.8 mmol) and 2,2,2-trichloroacetyl chloride (2.23 mL, 19.8 mmol) in 15 mL ether purged with nitrogen was added dropwise. After 20 the addition, the mixture was heated to reflux for 22 h. TLC showed trace amounts of starting material. The reaction mixture was cooled in ice bath, and then 8.0 g potassium carbonate was added. 30 mL MeOH was added dropwise and stirred until no gas evolution. EtOAc was added and the mixture was filtered through Celite. The filtrate was concentrated and the residue was purified by ISCO (SiO₂: 0 to 10% EtOAc/hexanes) to provide the product as a colorless oil (3.02 g, 83%).




[0026] To a solution of 3,3-bis(9-(benzyloxy)nonyl)-2,2-dichlorocyclobutan-1-one (3.02 g, 5.0 mmol) in 80 mL MeOH, Zn dust (1.96 g, 30 mmol) was added. After stirring for 15 min, ammonium chloride (1.6 g, 30 mmol) was added, and the reaction mixture was stirred at room temperature for 3 h. TLC showed completed reaction, and the mixture was concentrated to dryness. 100 mL water and 100 mL EtOAc were added, and the mixture was filtered through Celite. The filtrate was washed with brine and dried over magnesium sulfate. After filtration and concentration, the product was obtained (2.58 g, 97%), which was used for the next step without purification.




[0027] At 0 °C, sodium borohydride (0.51 g, 13.51 mmol) was added into a solution of 3,3-bis(9-(benzyloxy)nonyl)cyclobutan-1-one (2.58 g, 4.82 mmol) in 48 mL MeOH/THF (5:1), and then the reaction was stirred at this temperature for 1 h. TLC showed completed reaction. The reaction was quenched with saturated sodium bicarbonate, and then extracted with EtOAc (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the product was obtained as a colorless oil (2.68 g, quant.), which was used for the next step without purification.



20 [0028] To a solution of 3,3-bis(9-(benzyloxy)nonyl)cyclobutan-1-ol (3.31 g, 6.17 mmol) and imidazole (0.92 g, 13.57 mmol) in 50 mL dichloromethane, *tert*-butyldimethylsilyl chloride (1.15 g, 7.28 mmol) was added and the reaction mixture was stirred at room temperature for 4 h. TLC showed completed reaction. Water was added to quench the reaction, and the mixture was extracted with dichloromethane (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by ISCO (SiO₂: 0 to 20% EtOAc/hexanes) to provide the product as a colorless oil (3.53 g, 90%).




[0029] A mixture of (3,3-bis(9-(benzyloxy)nonyl)cyclobutoxy)(tert-butyl)dimethylsilane (3.53 g, 5.42 mmol) and palladium on carbon (10wt%, 0.71g) in 350 mL EtOAc was purged with nitrogen and hydrogen, respectively. After stirring under hydrogen balloon overnight, TLC 5 showed completed reaction, and then the reaction mixture was filtered through Celite. After concentration, the residue was purified by ISCO (SiO₂: 0 to 70% EtOAc/hexanes) to provide the product as a colorless oil (2.35 g, 92%).




10

[0030] At 0 °C, a solution of 9,9'-(3-((tert-butyldimethylsilyl)oxy)cyclobutane-1,1-diyl)bis(nanon-1-ol) (1.49 g, 3.16 mmol) in 20 mL dichloromethane was slowly added into a solution of Dess-Martin periodinane (2.68 g, 6.33 mmol) in 70 mL dichloromethane, and then the reaction mixture was stirred at this temperature for 3 h. After stirring at room temperature for 15 1 h, the reaction mixture was diluted with dichloromethane, and then 10% Na₂S₂O₃ solution and saturated sodium bicarbonate solution were added. After extraction with dichloromethane (2 X), the combined organic layers were washed with saturated sodium bicarbonate and brine. The organic layer was dried over sodium sulfate and concentrated. The crude was purified by ISCO (SiO₂: 0 to 10% EtOAc/hexanes) to provide the product as a colorless oil (0.88 g, 60%).


20



[0031] At 0 °C, potassium *tert*-butoxide (0.51 g, 4.53 mmol) was added into a solution of (Z)-non-3-en-1-yltriphenylphosphonium iodide (2.33 g, 4.53 mmol) in 30 mL THF which was purged with nitrogen 3 times. After 1 h, a solution of 9,9'-(3-((*tert*-butyldimethylsilyl)oxy)cyclobutane-1,1-diyl)dinonanal (0.88 g, 1.89 mmol) in 25 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed complete reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with EtOAc (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by ISCO (SiO₂: 0 to 20% EtOAc/hexanes) to provide the product as a colorless oil (543 mg, 42%).



[0032] To a solution of *tert*-butyl(3,3-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclobutoxy)dimethylsilane (0.67 g, 0.98 mmol) in 60 mL THF, a solution of TBAF (1.0 M in THF, 9.8 mL, 9.8 mmol) was added and the reaction mixture was stirred at room temperature for 3 h. TLC showed completed reaction. The solvent was removed under vacuum and the residue was purified by ISCO (SiO₂: 0 to 20% EtOAc/hexanes) to provide the product as a colorless oil (0.64 g, Quant.).



Chemical Formula: C₄₆H₈₃NO₂

Molecular Weight: 682.18

[0033] At 0 °C, pyridine (2.4 mL) and propylphosphonic anhydride solution (50wt% in DMF, 2.4 mL, 4.16 mmol) were added into a solution of 4-(dimethylamino)butanoic acid


5 hydrochloride (564 mg, 3.37 mmol) in 6 mL DMF. After stirring for 10 min, a solution of 3,3-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclobutan-1-ol (0.64 g, 1.12 mmol) in 4 mL DMF was added and the reaction mixture was stirred at room temperature overnight. MS and TLC showed the formation of product. Saturated sodium bicarbonate solution was added to quench the reaction, and then extracted with EtOAc (2 X). The combined organic layers were washed with 10 water and brine. After dried over sodium sulfate and concentration, the residue was purified by ISCO (SiO₂: 0 to 100% EtOAc/hexanes) to provide the product 3,3-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclobutyl 4-(dimethylamino)butanoate as a slight yellow oil (479 mg, 63%).

15 ¹H NMR (300 MHz, CDCl₃) δ 5.28-5.42 (m, 8 H), 4.89-4.99 (m, 1 H), 2.76 (t, 4 H, *J* = 6.1 Hz), 2.30 (t, 4 H, *J* = 7.4 Hz), 2.23 (s, 6 H), 2.15-2.21 (m, 2H), 2.04 (q, 8 H, *J* = 6.6 Hz), 1.68-1.84 (m, 4 H), 1.08-1.40 (m, 40 H), 0.88 (t, 6 H, *J* = 6.6 Hz).


APCI: m/z = 682.6 [M + H]⁺.

DS. Compound 21-3: 3,3-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentyl 3-(dimethylamino)propanoate

20




[0034] At 0 °C, to a solution of 1,4-dibromobutan-2-ol (75.0 g, 0.328 mole) and imidazole (49.0 g, 0.72 mole) in 500 mL dichloromethane, a solution of *tert*-butyldimethylsilyl chloride (57.0 g, 0.36 mole) in 300 mL dichloromethane was added dropwise. After the addition, the reaction mixture was warmed up to room temperature and kept stirring overnight. TLC showed clean conversion. The reaction mixture was filtered and washed with dichloromethane. After concentration, the residue was taken up with dichloromethane and washed with water and brine. The organic layer was dried over Na₂SO₄. After filtration and concentration, the crude was purified by flash column chromatography (SiO₂: 0 to 10% ether/hexanes) to provide pure product as a colorless liquid (83.05 g, 71%).




[0035] A solution of *tert*-butyl((1,4-dibromobutan-2-yl)oxy)dimethylsilane (53.7 g, 0.152 mole), diethyl malonate (10.0 g, 0.138 mole), potassium carbonate (47.6 g, 0.345 mole) and tetrabutylammonium bromide (4.45 g, 13.8 mmol) in 700 mL DMF was stirred at room temperature for 3 days. TLC showed almost no starting material. The reaction mixture was diluted with water and extracted by EtOAc (3 X), and the combined organic layers were washed with saturated ammonium chloride and brine. After drying over sodium sulfate, the solution was filtered and concentrated. The residue was purified by flash column chromatography (SiO₂: 0 to 10% EtOAc/hexanes) to give the desired product as a colorless oil (36.92 g, 77%).

10



[0036] At 0 °C, a solution of lithium aluminium hydride (2.0 M in THF, 43.2 mL, 86.4 mmol) was added into a solution of diethyl 3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-dicarboxylate (14.89 g, 43.2 mmol) in 60 mL THF, and then the reaction mixture was stirred at room temperature overnight. TLC showed clean conversion. The reaction was quenched by slowly adding of water (6 mL) and 1 N NaOH (20 mL), and then stirred for 30 min. The suspension was filtered through Celite and washed with EtOAc. After concentration, the residue was purified by flash column chromatography (SiO₂: 0 to 90% EtOAc/hexanes) to provide the product as a colorless oil (9.86 g, 88%).


20



[0037] At -78 °C, a solution of DMSO (2.15 mL, 30.3 mmol) in 10 mL dichloromethane was added dropwise into a solution of oxalyl chloride (1.35 mL, 15.2 mmol) in 15 mL dichloromethane, and then a solution of (3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-diol)dimethanol (1.88 g, 7.2 mmol) in 15 mL dichloromethane was added immediately. After

stirring 30 min, triethylamine (7.25 mL, 52.0 mmol) was added and the reaction mixture was warmed up to room temperature. TLC showed clean conversion. The reaction was quenched with water and extracted with ether (2 X). The combined organic layers were washed with saturated ammonium chloride and brine. After drying over sodium sulfate, the solution was 5 filtered and concentrated to give the product as a yellow oil (2.00 g, quant.), which was used for the next step without further purification.


Preparation of (8-(benzyloxy)octyl)triphenylphosphonium iodide:



10 [0038] At 0 °C, a solution of octane-1,8-diol (100 g, 0.684 mol) in 100 mL DMF was slowly added into a suspension of NaH (27.35 g, 0.684 mol) in 700 mL DMF. After stirring for 30 min, a solution of benzyl chloride (78.7 mL, 0.684 moles) in 200 mL DMF was slowly added. After addition, the reaction mixture was warmed up to room temperature and stirred overnight. TLC showed starting material was almost consumed. The reaction mixture was poured onto ice, and 15 then extracted by EtOAc (2 X). The combined organic layers were washed with water and brine, and then dried over sodium sulfate. After filtration and concentration, the crude was purified by flash column chromatography (SiO₂: 0 to 60% EtOAc/hexanes) to provide the product as a colorless oil (85.83 g, 53%).

20 [0039] At 0 °C, a solution of triphenylphosphine (114.4 g, 0.436 mole) in 300 mL dichloromethane was slowly added into a solution of 8-(benzyloxy)octan-1-ol (85.83 g, 0.363 mole), imidazole (52 g, 0.76 mole) and iodine (129.1 g, 0.51 mole) in 1200 mL dichloromethane, and then the reaction mixture was allowed to equilibrate to room temperature over 3 days. After filtration, the filtrate was concentrated and the residue was triturated with hexanes. The solution was filtered through a plug of silica gel and eluted with 10% ether in hexanes to provide the product as a cloudy liquid (81.09 g). The gummy solid was dissolved in dichloromethane and passed through silica gel and eluted with 10% ether in hexanes to provide a 25 cloudy liquid (20.0 g). Total yield: 101.1 g (80%).

[0040] A solution of (((8-iodooctyl)oxy)methyl)benzene (101.1 g, 0.293 mole) and triphenylphosphine (154.1 g, 0.586 mole) in acetonitrile (1 L) was refluxed overnight. After concentrated to dryness, the residue was dissolved in dichloromethane and purified by flash column chromatography (SiO₂: 0 to 10% MeOH/CH₂Cl₂) to provide the product as a yellow oil 5 (144.1 g, 81%).




[0041] At 0 °C, potassium tert-butoxide (2.42 g, 21.6 mmol) was added into a solution of (8-(benzyloxy)octyl)triphenylphosphonium iodide (14.2 g, 23.3 mmol) in 80 mL THF which 10 was purged with nitrogen 3 times. After 1 h, a solution of 3-((tert-butyldimethylsilyl)oxy)cyclopentane-1,1-dicarbaldehyde (2.00 g, 7.2 mmol) in 20 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed completed 15 reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with ether (2 X). The combined organic layer was washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by flash column chromatography (SiO₂: 0 to 6% ether/hexanes) to provide the product as a colorless oil (3.77 g, 79%).



20


[0042] A mixture of ((3,3-bis((Z)-9-(benzyloxy)non-1-en-1-yl)cyclopentyl)oxy)(tert-butyl)dimethylsilane (3.04 g, 4.6 mmol) and palladium on carbon (10%, 600 mg) in 200 mL EtOAc was purged with nitrogen then hydrogen, and then stirred under hydrogen overnight. TLC and MS showed complete reaction. The reaction mixture was filtered through Celite and 25 washed by EtOAc. The filtrate was concentrated to give the product as a colorless oil (2.32 g, Quant.).



[0043] At 0 °C, a solution of 9,9'-(3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-diyl)bis(nonan-1-ol) (2.32 g, 4.6 mmol) in 30 mL dichloromethane was slowly added into a solution of Dess-Martin periodinate (5.46 g, 12.88 mmol) in 70 mL dichloromethane, and then


5 the reaction mixture was stirred at this temperature for 4 h. TLC showed starting material was consumed. The reaction mixture was diluted with dichloromethane, and then 10% Na₂S₂O₃ solution and saturated sodium bicarbonate solution were added. After extraction with dichloromethane (2 X), the combined organic layers were washed with brine and concentrated. The residue was dissolved in ether and washed with saturated sodium bicarbonate and brine.

10 After drying over sodium sulfate, the solution was filtered and concentrated. The crude was purified by ISCO (SiO₂: 0 to 50% EtOAc/hexanes) to provide the product as a colorless oil (0.73 g, 16%).




[0044] At 0 °C, potassium *tert*-butoxide (363 mg, 3.23 mmol) was added into a solution of (Z)-

15 non-3-en-1-yltriphenylphosphonium iodide (1.66 g, 3.23 mmol) in 30 mL THF which was purged with nitrogen 3 times. After 1 h, a solution of 9,9'-(3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-diyl)dinonanal (0.52 g, 1.08 mmol) in 10 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed complete reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with ether (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by ISCO (SiO₂: 0 to 5% EtOAc/hexanes) to provide the product as a colorless oil (170 mg, 22%).



To a solution of *tert*-butyl((3,3-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentyl)oxy)dimethylsilane (170 mg, 0.24 mmol) in 20 mL THF, a solution of TBAF (1.0 M in THF, 2.4 mL, 2.4 mmol) was added and the reaction mixture was stirred at room temperature for 2 h. TLC showed complete reaction. The solvent was removed under vacuum and the residue was purified by ISCO (SiO₂: 0 to 20% EtOAc/hexanes) to provide the product as a colorless oil (80 mg, 57%).

5



(Compound 21-3)

Chemical Formula: C₄₆H₈₃NO₂

Molecular Weight: 682.18


10

[0045] At 0 °C, pyridine (0.1 mL) and propylphosphonic anhydride solution (50wt% in EtOAc, 0.51 mmol) were added into a solution of 3-(dimethylamino)propanoic acid hydrochloride (63 mg, 0.41 mmol) in 3 mL DMF. After stirring for 10 min, a solution of 3,3-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentan-1-ol (80 mg, 0.14 mmol) in 2 mL DMF was added and the reaction mixture was stirred at room temperature overnight. MS and TLC showed the formation of product. Saturated sodium bicarbonate solution was added to quench the reaction, and then extracted by EtOAc (2 X). The combined organic layer was washed with water and brine. After drying over sodium sulfate and concentration, the residue was purified by ISCO (SiO₂: 0 to 100% EtOAc/hexanes) to provide the product 3,3-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentyl 3-(dimethylamino)propanoate as a colorless oil (59 mg, 62%).

¹H NMR (300 MHz, CDCl₃) δ 5.28-5.42 (m, 8 H), 5.12-5.17 (m, 1 H), 2.76 (t, 4 H, *J* = 6.0 Hz), 2.59 (t, 2 H, *J* = 6.9 Hz), 2.42 (t, 2 H, *J* = 6.9 Hz), 2.23 (s, 6 H), 2.04 (q, 8 H, *J* = 6.9 Hz), 1.75-2.00 (m, 2 H), 1.38-1.72 (m, 5 H), 1.14-1.39 (m, 39 H), 0.88 (t, 6 H, *J* = 6.9 Hz).
APCI: m/z = 682.6 [M + H]⁺

5

DT. Compound 21-4: 3,3-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentyl 4-(dimethylamino)butanoate



10 4)

Chemical Formula: C₄₇H₈₅NO₂


Molecular Weight: 696.20

[0046] At 0 °C, pyridine (0.6 mL) and propylphosphonic anhydride solution (50wt% in DMF, 15 2.4 mL, 4.16 mmol) were added into a solution of 4-(dimethylamino)butanoic acid hydrochloride (565 mg, 3.37 mmol) in 3 mL DMF. After stirring for 10 min, a solution of 3,3-di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentan-1-ol (0.65 g, 1.115 mmol) in 1 mL DMF was added and the reaction mixture was stirred at room temperature overnight. MS and TLC showed the formation of product. Saturated sodium bicarbonate solution was added to quench the 20 reaction, and then extracted with EtOAc (2 X). The combined organic layer was washed with water and brine. After dried over sodium sulfate and concentration, the residue was purified by ISCO (SiO₂: 0 to 100% EtOAc/hexanes) to provide the product 3,3-Di((9Z,12Z)-octadeca-9,12-dien-1-yl)cyclopentyl 4-(dimethylamino)butanoate as colorless oil (556 mg, 72%).


¹H NMR (300 MHz, CDCl₃) δ 5.28-5.42 (m, 8 H), 5.06-5.15 (m, 1 H), 2.76 (t, 4 H, *J* = 6.0 Hz), 2.29 (t, 4 H, *J* = 7.4 Hz), 2.23 (s, 6 H), 2.04 (q, 8 H, *J* = 6.6 Hz), 1.14-1.99 (m, 48 H), 0.88 (t, 6 H, *J* = 6.9 Hz).

APCI: m/z = 696.6 [M + H]⁺.


DU. Compound 21-5: 3,3-Di((10 Z ,13 Z)-nonadeca-10,13-dien-1-yl)cyclopentyl 3-(dimethylamino)propanoate



[0047] At 0 °C, potassium *tert*-butoxide (10.03 g, 89.4 mmol) was added into a solution of (9-(benzyloxy)nonyl)triphenylphosphonium iodide (43.0 g, 69.0 mmol) in 160 mL THF which was purged with nitrogen 3 times. After 1 h, a solution of 3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-dicarbaldehyde (7.70 g, 29.8 mmol) in 60 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed complete reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with ether (2 X). The combined organic layer was washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by flash column chromatography (SiO₂: 0 to 6% ether/hexanes) to provide the product as a colorless oil (6.52 g, 32%).




[0048] A mixture of ((3,3-bis((Z)-10-(benzyloxy)dec-1-en-1-yl)cyclopentyl)oxy)(*tert*-butyl)dimethylsilane (6.52 g, 9.86 mmol) and palladium on carbon (10%, 1.30 g) in 400 mL EtOAc was purged with nitrogen then hydrogen, and then stirred under hydrogen overnight. TLC and MS showed complete reaction. The reaction mixture was filtered through Celite and washed with EtOAc. The filtrate was concentrated, and the residue was purified by flash column chromatography (SiO₂: 0 to 80% EtOAc/hexanes) to give the product as a semi-solid (4.60 g, 96%).



[0049] At 0 °C, a solution of 10,10'-(3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-diy)bis(decan-1-ol) (4.60 g, 9.49 mmol) in 100 mL dichloromethane was slowly added into a solution of Dess-Martin periodinane (11.276 g, 26.6 mmol) in 50 mL dichloromethane, and then the reaction mixture was stirred at this temperature for 2 h. The reaction mixture was diluted with dichloromethane, and then 10% Na₂S₂O₃ solution and saturated sodium bicarbonate solution were added. After extracted with dichloromethane (2 X), the combined organic layer was washed with saturated sodium bicarbonate and brine. The organic layer was dried over sodium sulfate and concentrated. The crude was purified by ISCO (SiO₂: 0 to 80% EtOAc/hexanes) to provide the product as a colorless oil (1.20 g, 26%).

Preparation of (9-(benzyloxy)nonyl)triphenylphosphonium iodide:




[0050] At 0 °C, a solution of nonane-1,9-diol (96.16 g, 0.60 mol) in 100 mL DMF was slowly added into a suspension of NaH (24.0 g, 0.60 mol) in 800 mL DMF. After stirring for 1 h, a solution of benzyl bromide (71.4 mL, 0.60 mole) in 200 mL DMF was slowly added. After addition, the reaction mixture was warmed up to room temperature and stirred overnight. TLC showed was almost consumed. The reaction mixture was poured onto ice, and then extracted with EtOAc (3 X). The combined organic layer was washed with water and brine, and then dried over sodium sulfate. After filtration and concentration, the crude was purified by flash column chromatography (SiO₂: 0 to 100% EtOAc/hexanes then 0 to 5% MeOH/dichloromethane) to provide the product as a colorless oil (74.4 g, 50%).

[0051] At 0 °C, a solution of triphenylphosphine (30.65 g, 0.117 mole) in 100 mL dichloromethane was slowly added into a solution of 9-(benzyloxy)nonan-1-ol (23.0 g, 0.097 mole), imidazole (13.93 g, 0.204 mole) and iodine (34.60 g, 0.136 mole) in 200 mL dichloromethane, and then the reaction mixture was allowed to room temperature overnight.

5 After filtration, the filtrate was concentrated and the residue was triturated with hexanes. The solution was filtered through a plug of silica gel and eluted with 10% ether in hexanes to provide the product as a cloudy liquid (32.2 g, 95%).


[0052] A solution of (((9-iodononyl)oxy)methyl)benzene (32.2 g, 0.093 mole) and 10 triphenylphosphine (48.93 g, 0.186 mole) in acetonitrile (500 mL) was refluxed overnight. After concentrated to dryness, the residue was dissolved in dichloromethane and purified by flash column chromatography (SiO₂: 0 to 10% MeOH/CH₂Cl₂) to provide the product as a yellow oil (44.4 g, 78%).




[0053] At 0 °C, potassium *tert*-butoxide (587 mg, 5.23 mmol) was added into a solution of (Z)-non-3-en-1-yltriphenylphosphonium iodide (3.69 g, 7.17 mmol) in 70 mL THF which was purged with nitrogen 3 times. After 1 h, a solution of 10,10'-(3-((*tert*-butyldimethylsilyl)oxy)cyclopentane-1,1-diyl)bis(decanal) **9'** (838 mg, 1.65 mmol) in 30 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed complete reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with ether (2 X). The combined organic layer was washed with brine and dried over sodium sulfate.

20 After filtration and concentration, the residue was purified by ISCO (SiO₂: 0 to 5% EtOAc/hexanes) to provide the product as a colorless oil (926 mg, 77%).

25



[0054] To a solution of *tert*-butyl((3,3-di((10Z,13Z)-nonadeca-10,13-dien-1-yl)cyclopentyl)oxy)dimethylsilane (926 mg, 1.27 mmol) in 100 mL THF, a solution of TBAF (1.0 M in THF, 13.3 mL, 13.3 mmol) was added and the reaction mixture was stirred at room temperature for 2 h. TLC showed complete reaction. The solvent was removed under vacuum and the residue was purified by ISCO (SiO₂: 0 to 20% EtOAc/hexanes) to provide the product as a colorless oil (830 mg, Quant.).



Chemical Formula: C₄₈H₈₇NO₂

Molecular Weight: 710.23

15 [0055] At 0 °C, pyridine (0.35 mL) and propylphosphonic anhydride solution (50wt% in EtOAc, 0.7 mL, 1.21 mmol) were added into a solution of 3-(dimethylamino)propanoic acid hydrochloride (151 mg, 0.98 mmol) in 6 mL DMF. After stirring for 10 min, a solution of 3,3-di((10Z,13Z)-nonadeca-10,13-dien-1-yl)cyclopentan-1-ol (200 mg, 0.33 mmol) in 4 mL DMF was added and the reaction mixture was stirred at room temperature overnight. MS and TLC showed the formation of product. Saturated sodium bicarbonate solution was added to quench the reaction, and then extracted with EtOAc (2 X). The combined organic layers were washed with water and brine. After drying over sodium sulfate and concentration, the residue was

20


purified by ISCO (SiO₂: 0 to 100% EtOAc/hexanes) to provide the product 3,3-Di((10Z,13Z)-nonadeca-10,13-dien-1-yl)cyclopentyl 3-(dimethylamino)propanoate as a colorless oil (135 mg, 58%).

¹H NMR (300 MHz, CDCl₃) δ 5.28-5.42 (m, 8 H), 5.10-5.17 (m, 1 H), 2.76 (t, 4 H, *J* = 6.1 Hz), 5.25 (t, 2 H, *J* = 6.9 Hz), 2.42 (t, 2 H, *J* = 6.9 Hz), 2.23 (s, 6 H), 2.04 (q, 8 H, *J* = 6.9 Hz), 1.75-2.00 (m, 2 H), 1.38-1.72 (m, 5 H), 1.14-1.39 (m, 43 H), 0.88 (t, 6 H, *J* = 6.9 Hz).
APCI: m/z = 710.7 [M + H]⁺.


DV. Compound 21-6: 1-(2,3-Di((8Z,11Z)-heptadeca-8,11-dien-1-yl)cyclopropyl)-N,N-dimethylmethanamine
10




[0056] At 0 °C, a solution of nonane-1,9-diol (96.16 g, 0.60 mol) in 100 mL DMF was slowly added into a suspension of NaH (24.0 g, 0.60 mol) in 800 mL DMF. After stirring for 1 h, a solution of benzyl bromide (71.4 mL, 0.60 mole) in 200 mL DMF was slowly added. After addition, the reaction mixture was warmed up to room temperature and stirred overnight. TLC showed starting material was almost consumed. The reaction mixture was poured onto ice, and then extracted with EtOAc (3 X). The combined organic layers were washed with water and brine, and then dried over sodium sulfate. After filtration and concentration, the crude was purified with flash column chromatography (SiO₂: 0 to 100% EtOAc/hexanes then 0 to 5% MeOH/dichloromethane) to provide the product as a colorless oil (74.4 g, 50%).




[0057] At 0 °C, a solution of triphenylphosphine (6.29 g, 24 mmol) in 100 mL dichloromethane was slowly added into a solution of 9-(benzyloxy)nonan-1-ol (5.0 g, 20 mmol), imidazole (2.9 g, 42 mmol) and iodine (8.5 g, 33.6 mmol) in 100 mL dichloromethane, and then the reaction mixture was allowed to room temperature overnight. After concentration, the residue was triturated with hexanes to provide the product as a cloudy liquid (5.38 g, 75%).



[0058] A solution of (((9-iodononyl)oxy)methyl)benzene (5.38 g, 14.9 mmol) and triphenylphosphine (7.8 g, 29.8 mmol) in acetonitrile (100 mL) was refluxed overnight. After concentrated to dryness, the residue was dissolved in dichloromethane and purified with flash column chromatography (SiO₂: 0 to 10% MeOH/CH₂Cl₂) to provide the product as a yellow oil 5 (8.5 g, 92%).




[0059] At 0 °C, a solution of 9-(benzyloxy)nonan-1-ol (5.0 g, 20 mmol) in 50 mL dichloromethane was slowly added into a suspension of Dess-Martin periodinane (12.5 g, 29.5 10 mmol) in 100 mL dichloromethane, and then the reaction mixture was stirred at room temperature overnight. After quenching with saturated sodium bicarbonate, the mixture was extracted with dichloromethane (2 X). The combined organic layer was dried over sodium sulfate and concentrated to give the product as a colorless oil (4.0 g, 80%).




[0060] At 0 °C, potassium *tert*-butoxide (1.54 g, 13.7 mmol) was added into a suspension of (9-(benzyloxy)nonyl)triphenylphosphonium iodide (8.5 g, 13.6 mmol) in 150 mL THF. After 1 h, a solution of 9-(benzyloxy)nonanal (3.0 g, 12 mmol) in 50 mL THF was added dropwise into the reaction mixture, and then the reaction was allowed to warm up to room temperature for 4 h. 20 TLC showed completed reaction. The reaction was quenched with saturated ammonium chloride, and then extracted with hexanes (2 X). The combined organic layers were dried over sodium sulfate. After filtration and concentration, the residue was purified with flash column chromatography (SiO₂: 0 to 100% dichloromethane/hexanes) to provide the product as colorless oil (3.5 g, 63%).


25




[0061] To a refluxing solution of (Z)-1,18-bis(benzyloxy)octadec-9-ene (3.5 g, 7.5 mmol) in 200 mL dichloromethane, a solution of Cu(acac)₂ (200 mg, 0.76 mmol) in 40 mL dichloromethane was added. And then ethyl diazoacetate (contains 13% dichloromethane, 9 X 1.3 mL) was added every 30 min. MS showed the formation of product. The reaction was quenched with MeOH and stirred for 1 h at room temperature. After concentration, the crude was purified by flash column chromatography (SiO₂: 0 to 10% EtOAc/hexanes) to provide the product as colorless oil (2.23 g, 82% pure mixed with acetate by-product).



[0062] A mixture of ethyl 2,3-bis(8-(benzyloxy)octyl)cyclopropane-1-carboxylate (2.23 g, 4.05 mmol) and Palladium on carbon (10wt%, 200 mg) in 200 mL EtOAc was stirred at room temperature under hydrogen balloon for 4.5 h. MS and TLC showed completed reaction. The reaction mixture was filtered through Celite and washed with EtOAc. The filtrate was concentrated to provide the product mixed with by-product, diethyl succinate (1.62 g, contains 0.55 equivalent of by-product, 84%).




[0063] At 0 °C, a solution of ethyl 2,3-bis(8-hydroxyoctyl)cyclopropane-1-carboxylate (1.43 g, 4.03 mmol) in 100 mL dichloromethane was slowly added into a suspension of Dess-Martin periodinane (3.63 g, 8.5 mmol) in 150 mL dichloromethane, and then the reaction mixture was stirred at room temperature for 2 h. After quenching with saturated sodium bicarbonate, the mixture was extracted with dichloromethane (2 X). The combined organic layer was dried over sodium sulfate and concentrated to give the product as colorless oil (1.2 g, 81%).




[0064] At 0 °C, potassium *tert*-butoxide (1.7 g, 15.1 mmol) was added into a solution of (Z)-non-3-en-1-yltrifphenylphosphonium iodide (5.8 g, 15.1 mmol) in 150 mL THF which was

purged with nitrogen 3 times. After 1 h, a solution of ethyl 2,3-bis(8-oxooctyl)cyclopropane-1-carboxylate (1.2 g, 3.28 mmol) in 50 mL THF (purged with nitrogen 3 times) was transferred via cannula into the reaction mixture, and then the reaction was allowed to warm up to room temperature overnight. TLC showed completed reaction. The reaction was quenched with 5 saturated ammonium chloride, and then extracted with hexanes (2 X). The combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by ISCO (SiO₂: 0 to 3% EtOAc/hexanes) to provide the product as colorless oil (1.3 g, 68%).




[0065] A solution of lithium aluminum hydride (2.0 M in THF, 1.7 mL, 3.35 mmol) was slowly added into a solution of ethyl 2,3-di((8Z,11Z)-heptadeca-8,11-dien-1-yl)cyclopropane-1-carboxylate (1.3 g, 2.23 mmol) in 150 mL THF, and then the reaction mixture was stirred at 15 room temperature for 30 min. TLC showed completed reaction. The reaction was quenched by slow addition of Na₂SO₄·10H₂O, then the mixture was filtered and washed with THF. The filtrate was concentrated and purified by flash column chromatography (SiO₂: 0 to 10% EtOAc/Hexanes) to give the product as colorless oil (1.1 g, 91%).



[0066] At 0 °C, a solution of (2,3-di((8Z,11Z)-heptadeca-8,11-dien-1-yl)cyclopropyl)methanol (1.1 g, 2.03 mmol) in 50 mL dichloromethane was slowly added into a suspension of Dess-Martin periodinane (1.30 g, 3.05 mmol) in 150 mL dichloromethane, and then the reaction

mixture was stirred at room temperature for 2 h. After quenching with saturated sodium bicarbonate, the mixture was extracted with dichloromethane (2 X). The combined organic layers were dried over sodium sulfate and concentrated to give the product as a colorless oil (0.87 g, 79%).

5



Chemical Formula: C₄₀H₇₃N

Molecular Weight: 568.03

10 [0067] To a solution of 2,3-di((8Z,11Z)-heptadeca-8,11-dien-1-yl)cyclopropane-1-carbaldehyde (0.87 g, 1.61 mmol) in 200 mL THF, dimethylamine (2.0 M in THF, 1.61 mL, 3.22 mmol), sodium triacetoxyborohydride (682 mg, 3.22 mmol) and acetic acid (0.19 mL, 3.22 mmol) were added subsequently, and the reaction mixture was stirred at room temperature overnight. MS showed completed reaction, and saturated sodium bicarbonate was added to quench the reaction.

15 The mixture was extracted with EtOAc (2 X), and the combined organic layers were washed with brine and dried over sodium sulfate. After filtration and concentration, the residue was purified by flash column chromatography (SiO₂: 0 to 10% MeOH/dichloromethane) to give the product 1-(2,3-di((8Z,11Z)-heptadeca-8,11-dien-1-yl)cyclopropyl)-N,N-dimethylmethanamine as a colorless oil (620 mg, 65%).

20 ¹H NMR (300 MHz, CDCl₃) δ 5.27-5.42 (m, 8 H), 2.76 (t, 4 H, *J* = 6.0 Hz), 2.38 (bs, 8 H), 2.04 (q, 8 H, *J* = 6.6 Hz), 1.18-1.59 (m, 36 H), 0.88 (t, 6 H, *J* = 6.6 Hz), 0.52-0.58 (m, 2 H), 0.28-0.38 (m, 1 H).

APCI: m/z = 568.6 [M + H]⁺.

25 Example 2: Production of nanoparticle compositions
Production of nanoparticle compositions

[00890] In order to investigate safe and efficacious nanoparticle compositions for use in the delivery of therapeutic and/or prophylactic agents to cells, a range of formulations are prepared and tested. Specifically, the particular elements and ratios thereof in the lipid component of nanoparticle compositions are optimized.

5 **[00891]** Nanoparticles can be made with mixing processes such as microfluidics and T-junction mixing of two fluid streams, one of which contains the therapeutic and/or prophylactic agent and the other has the lipid components.

[00892] Lipid compositions are prepared by combining a lipid according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) 10, a phospholipid (such as DOPE or DSPC, obtainable from Avanti Polar Lipids, Alabaster, AL), a PEG lipid (such as 1,2-dimyristoyl-*sn*-glycerol methoxypolyethylene glycol, also known as PEG-DMG, obtainable from Avanti Polar Lipids, Alabaster, AL), and a structural lipid (such as cholesterol, obtainable from Sigma-Aldrich, Taufkirchen, Germany) at concentrations of about 50 mM in ethanol. Solutions should be refrigeration for storage at, for example, -20° C. Lipids 15 are combined to yield desired molar ratios (see, for example, Table 1) and diluted with water and ethanol to a final lipid concentration of between about 5.5 mM and about 25 mM.

[00893] Nanoparticle compositions including a therapeutic and/or prophylactic agent and a lipid component are prepared by combining the lipid solution with a solution including the therapeutic and/or prophylactic agent at lipid component to therapeutic and/or prophylactic agent wt:wt ratios between about 5:1 and about 50:1. The lipid solution is rapidly injected using 20 a NanoAssemblr microfluidic based system at flow rates between about 10 ml/min and about 18 ml/min into the therapeutic and/or prophylactic agent solution to produce a suspension with a water to ethanol ratio between about 1:1 and about 4:1.

[00894] For nanoparticle compositions including an RNA, solutions of the RNA at 25 concentrations of 0.1 mg/ml in deionized water are diluted in 50 mM sodium citrate buffer at a pH between 3 and 4 to form a stock solution.

[00895] Nanoparticle compositions can be processed by dialysis to remove ethanol and achieve buffer exchange. Formulations are dialyzed twice against phosphate buffered saline (PBS), pH 7.4, at volumes 200 times that of the primary product using Slide-A-Lyzer cassettes 30 (Thermo Fisher Scientific Inc., Rockford, IL) with a molecular weight cutoff of 10 kD. The first dialysis is carried out at room temperature for 3 hours. The formulations are then dialyzed overnight at 4° C. The resulting nanoparticle suspension is filtered through 0.2 µm sterile filters

(Sarstedt, Nümbrecht, Germany) into glass vials and sealed with crimp closures. Nanoparticle composition solutions of 0.01 mg/ml to 0.10 mg/ml are generally obtained.

[00896] The method described above induces nano-precipitation and particle formation. Alternative processes including, but not limited to, T-junction and direct injection, may be used 5 to achieve the same nano-precipitation.

Characterization of nanoparticle compositions

[00897] A Zetasizer Nano ZS (Malvern Instruments Ltd, Malvern, Worcestershire, UK) can be used to determine the particle size, the polydispersity index (PDI) and the zeta potential 10 of the nanoparticle compositions in 1×PBS in determining particle size and 15 mM PBS in determining zeta potential.

[00898] Ultraviolet-visible spectroscopy can be used to determine the concentration of a therapeutic and/or prophylactic agent (e.g., RNA) in nanoparticle compositions. 100 µL of the diluted formulation in 1×PBS is added to 900 µL of a 4:1 (v/v) mixture of methanol and 15 chloroform. After mixing, the absorbance spectrum of the solution is recorded, for example, between 230 nm and 330 nm on a DU 800 spectrophotometer (Beckman Coulter, Beckman Coulter, Inc., Brea, CA). The concentration of therapeutic and/or prophylactic agent in the nanoparticle composition can be calculated based on the extinction coefficient of the therapeutic and/or prophylactic agent used in the composition and on the difference between the absorbance 20 at a wavelength of, for example, 260 nm and the baseline value at a wavelength of, for example, 330 nm.

[00899] For nanoparticle compositions including an RNA, a QUANT-IT™ RIBOGREEN® RNA assay (Invitrogen Corporation Carlsbad, CA) can be used to evaluate the encapsulation of an RNA by the nanoparticle composition. The samples are diluted to a 25 concentration of approximately 5 µg/mL in a TE buffer solution (10 mM Tris-HCl, 1 mM EDTA, pH 7.5). 50 µL of the diluted samples are transferred to a polystyrene 96 well plate and either 50 µL of TE buffer or 50 µL of a 2% Triton X-100 solution is added to the wells. The plate is incubated at a temperature of 37° C for 15 minutes. The RIBOGREEN® reagent is diluted 1:100 in TE buffer, and 100 µL of this solution is added to each well. The fluorescence 30 intensity can be measured using a fluorescence plate reader (Wallac Victor 1420 Multilabel Counter; Perkin Elmer, Waltham, MA) at an excitation wavelength of, for example, about 480 nm and an emission wavelength of, for example, about 520 nm. The fluorescence values of the reagent blank are subtracted from that of each of the samples and the percentage of free RNA is

determined by dividing the fluorescence intensity of the intact sample (without addition of Triton X-100) by the fluorescence value of the disrupted sample (caused by the addition of Triton X-100).

In vivo formulation studies

5 [00900] In order to monitor how effectively various nanoparticle compositions deliver therapeutic and/or prophylactic agents to targeted cells, different nanoparticle compositions including a particular therapeutic and/or prophylactic agent (for example, a modified or naturally occurring RNA such as an mRNA) are prepared and administered to rodent populations. Mice are intravenously, intramuscularly, intraarterially, or intratumorally administered a single dose 10 including a nanoparticle composition with a formulation such as those provided in Table 1. In some instances, mice may be made to inhale doses. Dose sizes may range from 0.001 mg/kg to 10 mg/kg, where 10 mg/kg describes a dose including 10 mg of a therapeutic and/or prophylactic agent in a nanoparticle composition for each 1 kg of body mass of the mouse. A control composition including PBS may also be employed.

15 [00901] Upon administration of nanoparticle compositions to mice, dose delivery profiles, dose responses, and toxicity of particular formulations and doses thereof can be measured by enzyme-linked immunosorbent assays (ELISA), bioluminescent imaging, or other methods. For nanoparticle compositions including mRNA, time courses of protein expression can also be evaluated. Samples collected from the rodents for evaluation may include blood, sera, and tissue 20 (for example, muscle tissue from the site of an intramuscular injection and internal tissue); sample collection may involve sacrifice of the animals.

[00902] Nanoparticle compositions including mRNA are useful in the evaluation of the efficacy and usefulness of various formulations for the delivery of therapeutic and/or prophylactic agents. Higher levels of protein expression induced by administration of a 25 composition including an mRNA will be indicative of higher mRNA translation and/or nanoparticle composition mRNA delivery efficiencies. As the non-RNA components are not thought to affect translational machineries themselves, a higher level of protein expression is likely indicative of a higher efficiency of delivery of the therapeutic and/or prophylactic agent by a given nanoparticle composition relative to other nanoparticle compositions or the absence 30 thereof.

Example 3: Sample formulations

5 [00903] Nanoparticle compositions including a therapeutic and/or prophylactic agent can be optimized according to the selection of a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) , the selection of additional lipids, the amount of each lipid in the lipid component, and the wt:wt ratio of the lipid component to the therapeutic and/or prophylactic agent, as described herein.

10 [00904] Initial studies were performed to compare the delivery efficiency of nanoparticle compositions including various compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) . The cationic lipid MC3 is a current standard in the art. Accordingly, the standard MC3 formulation including about 50 mol % MC3, about 10 mol % DSPC, about 38.5 mol % cholesterol, and about 1.5 mol % PEG-DMG was used as a basis for this study. Nanoparticle compositions including DOPE or DSPC as a phospholipid, cholesterol as a structural lipid, PEG-DMG as a PEG lipid, an RNA, and a compound according to one of formulae disclosed herein, e.g., selected from compounds of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), were prepared according to Examples 1 and 2. The ratios of the lipids were 40:20:38.5:1.5 mol% for the lipid described herein:DOPE:cholesterol:PEG-DMG or 50:10:38.5:1.5 mol% for the lipid described herein:DSPC:cholesterol:PEG-DMG. The RNA used was an mRNA encoding G5 luciferase (Luc) or G5 hEPO. Tables 1, 1b, 17-1, 19-1, 20-1 and 21-1 summarize the content and characteristics of the formulations.

20 [00905] As shown in Tables 1 and 1a, nanoparticle compositions including Compound 1 produced the largest particles amongst those of Tables 1 and 1a, while those including Compounds 34 and 50 produced the smallest particles amongst those of Tables 1 and 1da. Encapsulation efficiencies amongst those of Tables 1 and 1a were highest for compositions including Compounds 36, 37, 40, and 41 and lowest for those including Compounds 1 and 24.

25 Table 1. Characteristics of nanoparticle compositions including compounds of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	Size (nm)	PDI	EE (%)	pKa
1	203.2		46.2	5.95
2*	94.7	0.108	96	6.39
3	73.5	0.044	92.5	n.d.
4	89.7	0.120	96.78	7.12
5	85.1	0.140	98.1	7.01

Compound No.	Size (nm)	PDI	EE (%)	pKa
6	81.4	0.160	98.9	6.62
7	85.1	0.130	99.1	6.94
8	83.3	0.110	98.5	6.87
9	78.0	0.170	99.5	6.76
10	81.3	0.130	99.2	6.58
11	87.4	0.099	99.5	6.54
12	87.8	0.096	96.9	5.44
13	97.7	0.080	64.2	6.30
14	88.7	0.008	96.6	6.31
15 [#]	100.3	0.120	90.2	6.32
16 [#]	77.4	0.140	98.2	6.28
17	82.3	0.180	96.6	6.67
18	76.7	0.120	98.4	6.17
19	76.1	0.100	97.2	6.29
20	106.4	0.150	84.2	6.12
21	98.3	0.239	98.6	6.29
22 [#]	75.4	0.130	98.3	6.15
23	85.4	0.058	82.9	6.07
24*	110.4	0.131	36.4	6.01
25	90.0	0.186	97.0	6.20
26*	74.2	0.112	84.9	6.19
27	86.4	0.211	97.9	6.14
28	87.4	0.099	80.2	6.04
29	105.3	0.060	48.8	5.97
30	95.0	0.110	74.3	6.09
31	87.9	0.130	77.5	6.31
32	79.3	0.160	83.6	6.28
33	79.7	0.138	98.1	6.06
34*	66.0	0.077	98.1	5.74
36*	100.8	0.110	100.2	7.81
37*	86.6	0.107	99.9	6.45

Compound No.	Size (nm)	PDI	EE (%)	pKa
40*	78.9	0.210	100.0	6.78
41*	69.0	0.239	99.9	7.02
42	116.4	0.190	97.1	6.77
43	99.0	0.220	99.1	6.72
44	94.9	0.190	89.5	6.82
45	100.2	0.200	94.9	6.77
46	81.8	0.160	97.5	6.77
47	89.8	0.180	53.1	6.82
48	111.4	0.099	79.3	6.99
49	95.8	0.200	98.8	6.4
50	65.6	0.190	98.7	5.55
51	76.6	0.190	98.4	6.44
52	94.4	0.100	97.5	6.77
Formula IV	94.2		97.6	6.25
MC3	86.2	0.117	97.70	n.d.

n.d.=not determined

*=Formulated with lipid:DSPC:Chol:PEG-DMG 50:10:38.5:1.5

#=Formulated with hEPO mRNA

5 Table 1b. Characteristics of nanoparticle compositions including compounds of one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	Size (nm)	PDI	EE (%)
53	103.0	0.23	82.8
54	93.0	0.23	96.9
55	119.5	0.23	95.2
56	117.4	0.24	99.0
57	101.9	0.23	98.8
58	112.8	0.23	98.9
59	104.7	0.23	98.6
60	105.7	0.23	98.8
61	86.8	0.23	99.1
62	61.7	0.18	97.73

69	74.4	0.24	99.31
70	79.8	0.24	99.14
71	99.9	0.18	91.0
72	102.5	0.22	92.7
73	84.3	0.25	98.85
80	65.8	0.2	98.93
81	65.3	0.17	99.27
82	76.1	0.24	99.23
83	73.2	0.22	99.12
84	68.6	0.19	98.48
85	69.9	0.24	99.18
86	53.6	0.14	97.42
87	80.9	0.21	98.67
MC3	74.7	0.17	97.3

[00906] As shown in Table 17-1, Compounds 17-7 and 17-12 produced the smallest particles amongst those of Table 17-1, while Compounds 17-2 and 17-10 produced the largest particles amongst those of Table 17-1. The encapsulation efficiencies for Compounds 17-6 and 5 17-8 were comparable to that for MC3. Compounds 17-2 and 17-10 did not encapsulate RNA with high efficiency.

Table 17-1. Characteristics of nanoparticle compositions including compounds according to formula (17-I).

Compound No.	Size (nm)	PDI	EE (%)	pKa
17-2	136.9	0.104	57.2	6.92
17-3	117.9	0.095	82.1	6.69
17-4	95.6	0.154	94.5	6.34
17-5	88.4	0.137	94.2	6.92
17-6	80.2	0.117	97.3	6.73
17-7	68.5	0.110	95.6	5.68
17-8	86.4	0.20	96.9	6.16
17-9	87.1	0.138	95.0	4.93
17-10	165.0	0.239	30.3	2.85

Compound No.	Size (nm)	PDI	EE (%)	pKa
17-11	87.3	0.24	88.7	6.76
17-12	76.1	0.148	95.04	6.83
17-13	90.4	0.176	89.22	6.89
MC3	83.8	0.138	98.0	n.d.

n.d.=not determined

[001] As shown in Table 19-1, compositions including Compound 19-6 produced the largest particles amongst those of Table 19-1 with the lowest encapsulation efficiency, while those including Compound 19-3 produced the smallest particles amongst those of Table 19-1 with the highest encapsulation efficiency.

5

Table 19-1. Characteristics of nanoparticle compositions including compounds according to formula (19-I) or (19-II).

Compound No.	Size (nm)	PDI	EE (%)	pKa
19-1	98.0	0.071	83.4	6.76
19-2	72.0	0.239	98.8	7.32
19-3	47.9	0.076	99.7	7.24
19-4	111.4	0.071	95.2	7.09
19-5	106.9	0.204	93.6	6.32
19-6	137.7	0.088	16.9	5.89
MC3	83.3	0.122	97.6	n.d.

n.d.=not determined

[00907] As shown in Table 20-1, compositions including Compound 20-12 produced the largest particles amongst those of Table 20-1, while compositions including Compounds 20-8, 10 20-9, and 20-15 produced the smallest particles amongst those of Table 20-1. Encapsulation efficiencies for the compounds of Table 20-1 were highest for compositions including 20-19.

Table 20-1. Characteristics of nanoparticle compositions including compounds according to

15 formula (20-I).

Compound No.	Size (nm)	PDI	EE (%)	pKa
20-1*	87.8	0.078	90.7	4.48

Compound No.	Size (nm)	PDI	EE (%)	pKa
20-2	71.8	0.130	91.6	6.60
20-3	76.4	0.110	95.8	6.82
20-4	82.1	0.160	91.3	6.86
20-5	119.3	0.210	78.1	6.79
20-6	82.5	0.220	94.7	6.73
20-7	73.6	0.170	92.4	6.66
20-8	67.9	0.141	96.6	6.43
20-9	64.5	0.130	95.9	6.22
20-10	85.7	0.130	91.4	6.42
20-11	96.1	0.118	97.5	5.38
20-12	147.1	0.217	82.3	6.58
20-13	74.3	0.057	97.7	5.83
20-14	124.0	0.215	94.5	5.89
20-15	67.4	0.118	97.4	6.48
20-16	73.5	0.225	97.2	6.62
20-17	71.2	0.092	98.2	6.25
20-18	70.1	0.150	91.1	6.31
20-19	74.9	0.145	99.5	4.73
20-20	86.8	0.159	95.2	6.41
20-21	78.6	0.238	84.7	5.78
20-22	73.8	0.146	95.3	5.90
20-23	88.1	0.080	95.3	6.56
20-24	90.6	0.038	96.9	6.06
20-25	71.7	0.171	98.4	6.23
MC3	84.0	0.117	97.4	n.d.

n.d.=not determined

*=Formulated with DOPE

[0068] As shown in Table 21-1, compositions including Compound 21-1 produced the largest particles, amongst those of Table 21-1 while those including Compound 21-2 produced the

smallest particles amongst those of Table 21-1. The encapsulation efficiency for all compositions of Table 21-1 was greater than 98%.

Table 21-1. Characteristics of nanoparticle compositions including compounds according to formula (21-I).

Compound No.	Size (nm)	PDI	EE (%)	pKa
21-1	119.9	0.237	98.4	6.07
21-2	83.0	0.200	99.4	6.98
21-4	94.9	0.193	98.8	7.01
21-6	86.8	0.110	98.9	6.99
MC3	91.4	0.1191	98.1	n.d.

5

n.d.=not determined

Example 4: Expression of Luc induced by sample formulations

[00908] The efficacy of the nanoparticle compositions presented in Tables 1, 1a, 17-1, 19-1, 20-1 and 21-1 was evaluated with a bioluminescence study. Formulations were administered 10 intravenously to mice (n = 6) at a dosage of 0.5 mg/kg (mpk) and bioluminescence measured at 3, 6, and 24 hour time points. The standard MC3 formulation and a PBS control were evaluated for comparison.

[00909] As is evident in Table 2, the total flux for the compositions presented therein was generally comparable at 3 and 6 hours. The total flux after 24 hours was generally lower than 15 that at earlier time points. Amongst the compositions of Table 2, compositions including Compounds 18, 23, and 30 displayed the highest flux after 3 hours. Of the compositions of Table 2, compositions including Compounds 36 and 37 displayed the lowest flux after 24 hours. In general, these results suggest that the compounds described herein may be useful in transfection applications.

20

Table 2. Expression of luciferase induced by administration of nanoparticle compositions including compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
2	6.01E+09	3.23E+09	3.23E+09
3	3.75E+08	1.12E+09	n.d.
11	1.23E+10	3.81E+09	8.20E+08
12	1.06E+10	1.38E+10	6.03E+08
13	3.59E+09	3.80E+09	3.11E+08
14	9.86E+08	1.56E+09	1.02E+08
17	7.55E+09	2.49E+09	7.61E+08
18	2.13E+10	1.76E+10	7.00E+08
19	1.06E+10	6.52E+09	2.65E+09
20	1.00E+11	1.11E+11	4.60E+09
21	1.13E+10	1.08E+10	1.18E+08
23	2.33E+10	3.40E+10	1.06E+09
25	1.06E+10	1.08E+10	2.72E+08
26	1.65E+07	1.04E+07	2.75E+06
27	4.56E+09	4.70E+09	1.36E+08
28	6.18E+09	7.28E+09	4.02E+08
29	1.22E+08	2.51E+08	2.69E+07
30	2.87E+10	1.59E+10	1.95E+09
31	1.43E+10	1.42E+10	3.07E+08
32	6.85E+08	5.88E+08	4.37E+07
33	1.64E+09	4.71E+09	1.54E+08
34	7.77E+06	1.88E+07	2.19E+06
36	6.90E+05	3.93E+05	1.68E+05
37	1.19E+07	6.66E+06	9.38E+05
40	1.24E+08	1.07E+07	5.62E+06
41	4.06E+07	2.04E+07	6.05E+07
42	n.d	4.99E+10	n.d

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
43	n.d	4.54E+09	n.d
44	n.d	1.07E+10	n.d
45	n.d	7.86E+10	n.d
46	n.d	5.26E+09	n.d
47	n.d	2.64E+09	n.d
48	n.d	1.05E+08	n.d
49	n.d	5.67E+10	n.d
50	n.d	1.48E+08	n.d
51	n.d	6.70E+10	n.d
52	n.d	9.85E+10	n.d
MC3	1.63E+10	1.73E+10	1.16E+09

n.d.=not determined

[001] As is evident in Table 17-2, nanoparticle compositions including MC3 displayed the highest total flux of Table 17-2, while those including Compounds 17-4 and 17-8 displayed substantially higher flux than compositions including Compounds 17-2, 17-3, and 17-7. The total flux at 6 hours was higher than that at 3 hours for some compositions of Table 17-2. Generally, for the compositions of Table 17-2, the total flux at 24 hours was lower than the total flux measured at 3 or 6 hours.

Table 17-2. Expression of luciferase induced by administration of nanoparticle compositions including compounds according to formula (17-I).

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
17-2	2.83E+05	3.76E+05	2.64E+05
17-3	5.58E+05	1.38E+06	9.90E+05
17-4	1.22E+09	3.60E+08	6.05E+07
17-5	6.64E+08	8.54E+08	5.97E+07
17-6	6.27E+07	1.19E+08	1.27E+07
17-7	4.68E+05	6.82E+05	5.46E+05
17-8	1.02E+09	5.94E+08	3.79E+07

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
17-9	6.87E+06	9.97E+06	3.43E+05
17-10	6.61E+05	1.16E+06	4.06E+05
17-11	1.86E+06	2.82E+06	2.70E+05
17-12	2.94E+06	2.54E+06	4.66E+05
17-13	1.10E+06	1.80E+06	7.95E+05
MC3	1.63E+10	2.13E+10	1.01E+09

[00910] As is evident in Table 19-2, compositions including MC3 induced the highest expression of the compositions of Table 19-2 at each time point. Compositions including Compounds 19-5 and 19-6 produced the next highest flux of Table 19-2 at each time point, 5 while those including Compound 19-2 produced the lowest total flux of Table 19-2 at each time point.

Table 19-2. Expression of luciferase induced by administration of nanoparticle compositions including compounds according to formula (19-I) or (19-II).

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
19-1	6.77E+05	1.85E+06	2.75E+05
19-2	1.96E+05	3.51E+05	2.29E+05
19-3	1.69E+06	1.21E+06	3.39E+05
19-4	6.61E+05	4.75E+05	3.44E+05
19-5	2.37E+07	1.44E+07	2.76E+06
19-6	1.07E+07	1.27E+07	1.29E+06
MC3	1.53E+10	1.73E+10	1.13E+09

10 [00911] As is evident in Table 20-2, flux for the compositions of Table 20-2 was generally highest 3 hours after administration. Total flux for the compositions of Table 20-2 was highest after 3 hours for compositions including MC3, Compound 20-6, or Compound 20-7. Amongst the compositions of Table 20-2, expression 24 hours after administration was lowest for compositions including Compounds 20-1, 20-6, and 20-16.

Table 20-2. Expression of luciferase induced by administration of nanoparticle compositions including compounds according to formula (20-I).

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
20-1	1.44E+06	8.30E+05	4.53E+05
20-2	1.35E+09	2.18E+09	6.49E+07
20-3	5.35E+09	3.00E+09	6.49E+07
20-4	3.01E+06	1.75E+06	6.82E+08
20-5	6.07E+07	1.99E+07	2.07E+06
20-6	3.39E+10	5.44E+06	5.48E+05
20-7	1.22E+10	8.94E+09	2.65E+08
20-8	2.76E+09	4.37E+09	1.30E+08
20-9	3.65E+08	6.91E+08	4.81E+07
20-10	5.05E+09	2.16E+09	2.54E+08
20-11	1.44E+09	8.83E+08	2.40E+07
20-12	1.57E+09	1.84E+09	1.24E+08
20-13	7.01E+08	1.82E+09	7.39E+07
20-14	1.76E+09	4.07E+08	8.10E+07
20-15	3.36E+08	2.25E+08	2.08E+07
20-16	7.88E+05	6.25E+05	2.16E+05
20-17	1.97E+07	1.44E+07	1.54E+06
20-18	5.80E+09	6.48E+09	1.54E+09
20-19	5.37E+05	7.60E+05	6.08E+05
20-20	4.56E+09	3.27E+09	3.56E+08
20-21	1.43E+09	1.02E+09	1.13E+08
20-22	4.00E+09	3.01E+09	2.20E+08
20-23	1.91E+09	1.42E+09	1.13E+08
20-24	7.47E+08	1.64E+08	9.72E+06
20-25	1.16E+09	1.78E+09	3.82E+07
MC3	1.73E+10	1.94E+10	8.48E+08

[0069] As is evident in Table 21-2, compositions including MC3 produced the highest luciferase expression of Table 21-2 at each time point, while, amongst the compositions of Table 21-2, those including Compound 21-2 showed the least expression at 3 and 6 hours. Each composition of Table 21-2 showed substantial expression. Flux generally decreased over time.

5 These results suggest that the compounds described herein may be useful in transfection applications.

Table 21-2. Expression of luciferase induced by administration of nanoparticle compositions including compounds according to formula (21-I).

Compound No.	Total Flux		
	3 hours	6 hours	24 hours
21-1	3.66E+09	4.19E+09	4.19E+09
21-2	5.99E+08	8.09E+08	8.09E+08
21-4	1.15E+09	1.26E+09	1.26E+09
21-6	2.92E+09	4.54E+09	1.40E+08
MC3	1.69E+10	2.87E+10	2.23E+10

10 Example 5: Expression of Luc induced by sample formulations in different organs

[00912] The efficacy of the nanoparticle compositions presented in Tables 1, 1a, 17-1, 19-1, 20-1 and 21-1 was further evaluated by measuring the expression of modified luciferase in the liver, lung, spleen, and femur upon administration of a given composition. Formulations were administered intravenously to mice (n = 3) at a dosage of 0.5 mpk and bioluminescence measured after 6 hours. The standard MC3 formulation and a PBS control were also tested.

[00913] As is evident in Table 3, expression was highest in the liver for all formulations of Table 3. Of the compositions of Table 3, the highest total flux was measured for compositions including Compound 20. Lung and spleen expression were generally comparable for compounds of Table 3, while expression in the femur, where measured, was somewhat lower.

20 Table 3. Expression of luciferase in various organs 6 hours after administration of nanoparticle compositions including compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	Total Flux			
	Liver	Lung	Spleen	Femur
2	3.33E+08	4.76E+05	1.14E+07	n.d.

Compound No.	Total Flux			
	Liver	Lung	Spleen	Femur
3	1.36E+08	8.47E+05	9.51E+05	n.d.
11	1.67E+09	2.21E+06	7.21E+06	1.47E+06
12	1.05E+09	6.76E+06	1.11E+07	n.d.
13	6.10E+08	2.89E+06	3.63E+07	1.99E+06
14	2.62E+08	3.56E+06	1.46E+07	n.d.
17	4.26E+08	7.26E+05	3.20E+06	6.71E+05
18	3.91E+09	1.87E+07	1.60E+07	3.31E+06
19	1.89E+09	2.27E+06	8.28E+06	2.75E+06
20	1.42E+10	1.46E+08	6.11E+07	8.91E+06
21	1.24E+09	2.51E+06	1.17E+07	n.d.
23	4.94E+09	1.51E+07	2.95E+07	3.17E+06
25	2.68E+09	5.88E+06	6.00E+06	n.d.
26	2.35E+06	3.49E+04	2.30E+05	n.d.
27	7.84E+08	3.56E+06	3.34E+06	n.d.
28	8.10E+08	5.73E+06	5.67E+06	1.16E+06
29	2.27E+07	4.70E+05	2.97E+06	n.d.
30	2.42E+09	1.61E+07	7.18E+06	2.22E+06
31	1.54E+09	9.81E+06	1.28E+07	9.89E+05
32	8.36E+07	6.75E+05	9.38E+05	1.02E+05
33	6.15E+08	2.84E+06	4.82E+06	1.18E+06
34	2.79E+06	5.63E+04	1.22E+06	n.d.
36	5.85E+04	2.74E+04	1.24E+05	n.d.
37	1.92E+06	6.90E+05	9.75E+05	n.d.
40	1.33E+06	1.42E+05	5.68E+05	n.d.
41	3.00E+06	1.34E+05	2.13E+06	n.d.
42	5.53E+09	n.d.	2.29E+08	n.d.
43	2.60E+08	n.d.	4.52E+07	n.d.
44	1.11E+09	n.d.	1.19E+08	n.d.
45	7.87E+09	n.d.	1.70E+08	n.d.
46	3.84E+08	n.d.	4.35E+07	n.d.

Compound No.	Total Flux			
	Liver	Lung	Spleen	Femur
47	4.95E+08	n.d	1.42E+08	n.d
48	1.04E+07	n.d	1.50E+07	n.d
49	1.21E+10	n.d	6.65E+07	n.d
50	2.14E+07	n.d	1.94E+05	n.d
51	3.55E+09	n.d	2.24E+07	n.d
52	1.18E+10	n.d	8.74E+08	n.d
IV	9.15E+08	n.d	6.15E+08	n.d
MC3	2.31E+09	8.61E+06	1.95E+07	3.08E+06

n.d.=not determined

[00914] As is evident in Table 17-3, the total flux for compositions of Table 17-3 was generally higher in the liver than in other organs. The total flux in the liver for nanoparticle compositions of Table 17-3 including Compounds 17-5 and 17-8 was somewhat comparable to those including MC3, which displayed the highest total flux of Table 17-3 in each organ. For compositions of Table 17-3, the total flux in the spleen was generally higher than that in the lung and was highest for compositions including Compounds 17-4, 17-5, and 17-8.

Table 17-3. Expression of luciferase in various organs 6 hours after administration of nanoparticle compositions including compounds according to formula (17-I).

Compound No.	Total Flux			
	Liver	Lung	Spleen	Femur
17-2	6.42E+04	1.96E+04	4.16E+04	n.d.
17-3	2.51E+05	1.56E+04	1.02E+05	n.d.
17-4	8.00E+07	1.95E+05	2.30E+06	n.d.
17-5	1.17E+08	8.13E+05	2.13E+06	n.d.
17-6	2.16E+07	1.54E+05	9.06E+05	n.d.
17-7	7.44E+04	1.35E+04	3.46E+04	n.d.
17-8	2.27E+08	6.11E+05	4.25E+06	n.d.
17-9	1.39E+06	1.40E+04	3.72E+05	2.29E+04
17-10	2.00E+05	2.52E+04	1.74E+04	n.d.
17-11	5.06E+05	1.20E+04	3.37E+05	n.d.

Compound No.	Total Flux			
	Liver	Lung	Spleen	Femur
17-12	4.94E+05	4.89E+04	7.81E+04	n.d.
17-13	3.56E+05	4.66E+04	1.90E+05	n.d.
MC3	2.75E+09	7.25E+06	2.62E+07	6.03E+06

n.d.=not determined

[00915] As is evident in Table 19-3, total flux for the compositions of Table 19-3 was highest for compositions including MC3. For compositions of Table 19-3, expression in the liver was higher than expression in the lung and spleen for all compounds of Table 19-3 tested. Of the compositions of Table 19-3, compositions including Compound 19-2 yielded the lowest total flux in the liver.

Table 19-3. Expression of luciferase in various organs 6 hours after administration of nanoparticle compositions including compounds according to formula (19-I) or (19-II).

Compound No.	Total Flux		
	Liver	Lung	Spleen
19-1	8.53E+05	1.06E+04	1.58E+04
19-2	5.73E+04	2.33E+04	3.56E+04
19-3	2.65E+05	3.00E+04	2.55E+05
19-4	1.46E+05	4.49E+04	3.69E+04
19-5	5.99E+06	3.46E+04	2.10E+05
19-6	3.27E+06	1.81E+05	3.47E+06
MC3	2.39E+09	5.83E+06	2.45E+07

10

[00916] As is evident in Table 20-3, expression for the compounds therin was generally highest in the liver and lowest in the lung and femur. Total flux for the compounds of Table 20-3 in the liver was highest for compositions including MC3 or Compound 20-7 and lowest for those including Compound 20-16.

15 Table 20-3. Expression of luciferase in various organs 6 hours after administration of nanoparticle compositions including compounds according to formula (20-I).

Compound No.	Total Flux			
	Liver	Lung	Spleen	Femur
20-1	3.19E+05	3.79E+04	1.70E+05	n.d.
20-2	6.01E+08	4.52E+06	3.12E+07	1.07E+06
20-3	4.33E+08	6.08E+05	6.35E+06	1.73E+06
20-4	2.93E+05	1.11E+04	4.51E+04	2.09E+04
20-5	2.67E+06	5.80E+04	3.40E+06	9.53E+04
20-6	7.75E+05	2.32E+04	2.35E+05	n.d.
20-7	1.00E+09	1.38E+07	4.58E+07	n.d.
20-8	8.09E+08	3.45E+06	2.70E+07	1.62E+06
20-9	1.57E+08	1.30E+06	5.11E+06	2.89E+05
20-10	4.40E+08	5.01E+06	9.67E+07	3.09E+06
20-11	3.60E+08	5.86E+05	9.24E+06	n.d.
20-12	3.08E+08	3.58E+06	6.15E+07	1.37E+06
20-13	1.65E+08	9.09E+05	1.04E+07	3.11E+05
20-14	5.22E+07	1.08E+05	7.03E+05	n.d.
20-15	6.74E+07	5.56E+05	1.86E+06	n.d.
20-16	9.55E+04	1.44E+04	6.57E+04	n.d.
20-17	2.99E+06	2.79E+04	9.56E+04	n.d.
20-18	4.83E+08	1.84E+06	2.36E+06	3.78E+05
20-19	3.80E+05	1.96E+04	2.74E+04	n.d.
20-20	8.92E+08	1.12E+06	1.39E+07	n.d.
20-21	1.27E+08	1.98E+05	2.20E+06	n.d.
20-22	2.88E+08	5.04E+05	1.65E+06	n.d.
20-23	3.06E+08	1.28E+06	4.19E+06	5.10E+05
20-24	4.05E+07	1.08E+05	1.11E+06	n.d.
20-25	1.88E+08	7.85E+05	4.16E+06	3.13E+05
MC3	2.48E+09	1.28E+07	2.85E+07	2.60E+06

n.d.=not determined

[0070] As is evident in Table 21-3, expression was highest in the liver and lowest in the lung for all compositions of Table 21-3. Of the compositions of Table 21-3, compositions including

MC3 produced the highest expression in each organ, while compositions including Compounds 21-2 and 21-4 produced the lowest expression in each organ.

Table 21-3. Expression of luciferase in various organs 6 hours after administration of nanoparticle compositions including compounds according to formula (21-I).

Compound No.	Total Flux		
	Liver	Lung	Spleen
21-1	3.62E+08	5.44E+05	3.27E+07
21-2	5.50E+07	2.53E+05	1.56E+07
21-4	9.89E+07	2.13E+05	1.27E+07
21-6	4.79E+08	2.09E+06	1.94E+07
MC3	3.05E+09	6.60E+06	4.73E+07

5

Example 6: Cytokine production induced by sample formulations

[00917] The introduction of foreign material into a mammalian body induces an innate immune response that promotes cytokine production. Such immune responses to, for example, nanoparticle compositions including therapeutic and/or prophylactic agents, are undesirable.

10 The induction of certain cytokines is thus measured to evaluate the efficacy of nanoparticle compositions. The concentrations of various cytokines in mice upon intravenous administration of nanoparticle compositions presented in Tables 1, 1a, 17-1, 19-1, 20-1 and 21-1 at a dosage of 0.5 mpk was measured at 6 hours. The standard MC3 formulation and a PBS control were also tested.

15 **[00918]** As is evident in Table 4, IP-10 expression was lower than IL-6 expression for compositions of Table 4. Of the compositions of Table 4, compositions including Compound 13 induced the highest expression of both IL-6 and IP-10, while compositions including Compound 3 induced the lowest IL-6 expression and those including Compound 36 induced the lowest IP-10 expression.

20 Table 4. Cytokine induction 6 hours after administration of nanoparticle compositions including compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	IL-6	IP-10
2	n.d.	3365
3	7	341
11	250	3305

Compound No.	IL-6	IP-10
12	31	2382
13	301	7266
14	17	209
17	164	545
18	263	655
19	70	2326
20	127	2080
21	48	652
23	259	3702
25	131	1823
26	17	175
27	42	2564
28	73	5364
29	108	3454
30	300	4235
31	188	2513
32	174	727
33	37	1577
34	28	159
36	41	118
37	n.d.	198
40	134	919
41	116	350
MC3	92	438

n.d.=not determined

[002] As is evident in Table 17-4, nanoparticle compositions of Table 17-4 induced higher IP-10 levels than IL-6 levels. Of the nanoparticle compositions of Table 17-4, compositions including MC3 and Compound 17-4 induced the highest IL-6 and IP-10 levels while those including Compounds 17-2, 17-3, and 17-10 induced the lowest IL-6 levels and compositions including Compounds 17-2 and 17-3 induced the lowest IP-10 levels.

5

Table 17-4. Cytokine induction 6 hours after administration of nanoparticle compositions including compounds according to formula (17-I).

Compound No.	IL-6	IP-10
17-2	9.88	72.8
17-3	6.97	66.8
17-4	80.7	560
17-5	49.2	300
17-6	25.0	134
17-7	49.9	100
17-8	n.d.	135
17-9	35.2	112
17-10	7.33	243
17-11	n.d.	n.d.
17-12	83.1	148
17-13	84.3	222
MC3	107	500

[002] As is evident in Table 19-4, of the nanoparticle compositions of Table 19-4,

5 compositions including Compound 19-6 induced the highest IL-6 expression, while those including Compound 19-3 induced the lowest IL-6 expression. Of the nanoparticle compositions of Table 19-4, compositions including Compounds 19-4 and 19-6 yielded the highest IP-10 expression, while those including Compound 19-1 induced the lowest IP-10 expression.

Table 19-4. Cytokine induction 6 hours after administration of nanoparticle compositions

10 including compounds according to formula (19-I) or (19-II).

Compound No.	IL-6	IP-10
19-1	144.4	51.1
19-2	n.d.	n.d.
19-3	5.4	177.5
19-4	129.2	430.9
19-5	n.d.	n.d.
19-6	7571	433
MC3	65.7	323.9

n.d.=not determined

[00919] As is evident in Table 20-4, for nanoparticle compositions of Table 20-4, IP-10 induction was generally higher than IL-6 induction. For nanoparticle compositions of Table 20-4, IP-10 induction was highest for compositions including Compound 20-14 and lowest for compositions including Compound 20-6. IL-6 induction was highest for compositions including Compound 20-10 and lowest for compositions including Compound 20-6 for the nanoparticle compositions of Table 20-4.

Table 20-4. Cytokine induction 6 hours after administration of nanoparticle compositions including compounds according to formula (20-I).

Compound No.	IL-6	IP-10
20-1	62.4	2065.8
20-2	118.5	522.9
20-3	105.8	671.5
20-4	169.3	270.5
20-5	140.8	2012.3
20-6	0	25.3
20-7	24.5	696.3
20-8	18.4	134.0
20-9	152.1	271.1
20-10	739.0	2356.8
20-11	46.9	1700.5
20-12	445.7	3864.5
20-13	87.8	70.0

Compound No.	IL-6	IP-10
20-14	311.1	8436.6
20-15	49.4	136.3
20-16	n.d.	395.7
20-17	6.3	1207.0
20-18	102	572.7
20-19	n.d.	127.2
20-21	68.4	900.4
20-22	118.6	810.6
20-23	236.4	457.8
20-24	67.1	1678.6
20-25	20.6	188.0
MC3	119.5	499.1

[0071] As is evident in Table 21-4, IP-10 induction by compositions therein was substantially higher than IL-6 induction.

Table 20-4. Cytokine induction 6 hours after administration of nanoparticle compositions including compounds according to formula (21-I).

Compound No.	IL-6	IP-10
21-1	n.d.	1265.1
21-2	n.d.	477.9
21-4	n.d.	577.2
21-6	10.2	641.3
MC3	31.1	304.5

5

Example 7: Expression of hEPO induced by sample formulations

[00920] Formulations were prepared according to Table 5 and included mRNA encoding hEPO.

Table 5. Characteristics of nanoparticle compositions including compounds according to one of 10 formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	Composition (mol %)	Components	Size (nm)	PDI	EE (%)	pKa	Conc. (ug/ml)
12	40:20:38.5:1.5	Lipid:DOPE:Chol:	87.5	0.13	93.79	5.444	320.14

Compound No.	Composition (mol %)	Components	Size (nm)	PDI	EE (%)	pKa	Conc. (ug/ml)
		PEG-DMG					
14	40:20:38.5:1.5	Lipid:DOPE:Chol:PEG-DMG	76.8	0.14	98.91	6.308	603.76
15	40:20:38.5:1.5	Lipid:DOPE:Chol:PEG-DMG	100.3	0.12	90.15	6.323	713.00
16	40:20:38.5:1.5	Lipid:DOPE:Chol:PEG-DMG	77.4	0.14	98.22	6.282	665.11
20	40:20:38.5:1.5	Lipid:DOPE:Chol:PEG-DMG	114.5	0.14	94.39	n.d.	1264.28
22	40:20:38.5:1.5	Lipid:DOPE:Chol:PEG-DMG	75.4	0.13	98.29	6.153	564.97
23	40:20:38.5:1.5	Lipid:DOPE:Chol:PEG-DMG	98.5	0.16	77.19	6.070	438.20
23	50:10:38.5:1.5	Lipid:DSPC:Chol:PEG-DMG	95.2	0.11	51.46	6.164	454.58
MC3	50:10:38.5:1.5	Lipid:DSPC:Chol:PEG-DMG	76.5	0.11	97.37	n.d.	470.45

n.d.=not determined

[00921] Formulations were administered intravenously to rats (n=3 or 6) at a dosage of 0.2 mg/kg or 0.5 mg/kg (mpk) and hEPO levels measured at 3, 6, and 24 hour time points.

5 After the 48 hour time point, livers and spleens were harvested and frozen. As is evident in Table 6, compositions including MC3 yielded the highest hEPO expression at each time point, while compositions including Compound 16 yielded the lowest hEPO expression at each time point.

Table 6. Expression of hEPO induced by administration of nanoparticle compositions including 10 compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III) and (IIIa).

Compound No.	hEPO (pg/mL)		
	3 hours	6 hours	24 hours
12	592260	424740	165404
14	280973	158520	58805

Compound No.	hEPO (pg/mL)		
	3 hours	6 hours	24 hours
15	103773	125320	67965
16	35387	41720	17184
20	n.d.	227420	n.d.
22	181627	267680	75571
23 (DOPE)	249213	275440	120104
23 (DSPC)	86627	71360	29008
MC3	1407947	1036013	436243

n.d.=not determined

[00922] As shown in Table 7a, hEPO expression in mice was substantially higher for compositions including Compound 12 than those including MC3. In contrast, hEPO expression in rats induced by administration of a nanoparticle composition including a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), and (IIIa) was substantially lower than that measured for MC3. Luc expression in mice was several fold higher for Compound 23 than for MC3 but significantly lower for Compounds 12 and 14. Table 7b shows the Luc expressions in CD-1 mice vs. LDLr^{-/-} mice. Table 7c and 7d show additional protein expressions and clearance data from compositions with various compounds disclosed herein as compared to with MC3. Table 7e shows hEPO expression data in CD-1 mice at a dose of 0.5 mpk. Similar results were achieved with different strains of mice, e.g. chimeric mice with humanized livers (PXB) or immunodeficient mice (SCID).

Table 7a. Comparison of expression induced by administration of nanoparticle compositions including MC3 or compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), and (IIIa).

Compound No.	Lipid/MC3 Luc CD-1 (mice)	Lipid/MC3 hEPO CD-1 (mice)	Lipid/MC3 hEPO S.D. (rats)	%Dose remaining 6 h CD-1 mice	%Dose remaining 24 h CD-1 mice
11	1.03	n.d.	n.d.		
12	0.21	2.3	0.32		
14	0.070	n.d.	0.12		
15	n.d.	n.d.	0.095		
16	n.d.	n.d.	0.031		

Compound No.	Lipid/MC3 Luc CD-1 (mice)	Lipid/MC3 hEPO CD-1 (mice)	Lipid/MC3 hEPO S.D. (rats)	%Dose remaining 6 h CD-1 mice	%Dose remaining 24 h CD-1 mice
20	5.0	n.d.	n.d.		
22	n.d.	n.d.	0.20		
23 (DOPE)	5.2	2.4	0.21		
42	3.99	n.d.	n.d.	58	53
43	0.34			69	64
44	0.82	n.d.	n.d.	<1	<1
45	6.50	n.d.	n.d.	33	24
46	0.46			1	<1
47	0.22			1	<1
48	0.01			<1	<1
49	5.23	n.d.	n.d.	49	40
50	0.01			56	47
51	5.22	n.d.	n.d.	54	41
52	7.46	n.d.	n.d.	46	40
MC3	1	1	1	88	55

n.d.=not determined

Table 7b. Comparison of Luc expression induced by administration of nanoparticle compositions including MC3 or compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), and (IIIa) in CD-1 mice and LDLr^{-/-} mice.

5

Compound No.	Lipid/MC3 AUC CD-1 mice 0.5 mpk	Lipid/KL22 AUC LDLr ^{-/-} mice 0.5 mpk, LDLr	LDL levels lipid/untreated LDLr ^{-/-} mice 0.5 mpk, LDLr
4	3.70	2.03	0.54
5	2.62	1.86	0.47
6	1.72	0.37	0.91
7	1.51	2.46	0.63
8	2.33	3.74	0.66
9	0.73	0.58	0.87

10	1.14	0.71	0.98
MC3	1	0.15	0.55
KL22		1	0.51

Table 7c. Comparison of expression induced by administration of nanoparticle compositions including MC3 or compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), and (IIIa).

Compound No.	AUC (p/s*h)	Lipid/MC3 AUC	Liver/Spleen Ratio	% Dose Remaining 6hr	% Dose Remaining 24hr
53	9.22 E+09	0.07	17	1.35	0.77
54	1.89 E+10	0.13	32	5.18	4.77
55	2.00 E+10	0.14	9	2.58	1.60
56	1.77 E+11	1.25	29	1.21	0.16
57	1.21 E+11	0.85	15	0.88	0.24
58	1.38 E+11	0.97	11	1.37	0.61
59	1.19 E+11	0.84	5	6.99	5.03
60	2.84 E+11	2.00	15	21.18	15.98
61	4.65 E+11	3.27	30	1.31	0.13
71	1.77 E+11	1.25	25	12.39	9.25
72	6.53 E+10	0.46	6	7.06	6.40
MC3	1.42 E+11	-	55	55.70	55.43

5

Table 7d. Comparison of hEPO expression in S.D. rats induced by administration of nanoparticle compositions including MC3 or compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), and (IIIa).

Compound No	AUC (0.1 mpk)	AUC/MC3 0.1 mpk	AUC (1 mpk)	AUC/MC3 1 mpk	% dose remaining Rat liver (48 h)	% dose remaining Mouse liver (24 h)
-------------	---------------	-----------------	-------------	---------------	-----------------------------------	-------------------------------------

4	2.12 E+06	1.3	6.79 E+07	0.58	37.82	N.A.
45	1.45 E+06	0.90	2.00 E+08	1.7	8.57	23.7
49	5.98 E +06	3.7	1.44 E+08	1.2	38.45	40.3
MC3	1.62 E +06	-	1.17 E+08	-	43.11	55

Table 7e. Comparison of hEPO expression in CD-1 mice induced by administration of nanoparticle compositions including MC3 or compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), and (IIIa) at a dose of 0.5 mpk.

Compound No.	AUC (p/s*h) CD-1 mice 0.5 mpk	Lipid/MC3 AUC
62	3.46 E+7	0.97
69	2.33 E+8	6.52
70	6.34 E+7	1.77
73	1.41 E+8	3.93
80	6.24 E+7	1.74
81	1.08 E+8	3.01
82	1.29 E+8	3.62
83	5.21 E+7	1.46
84	5.10 E+7	1.43
85	1.27 E+8	3.54
86	1.75 E+7	0.49
87	2.86 E+7	0.80

5

[00923] The amount of lipid in the liver and spleen 48 hours after administration of a nanoparticle composition was also measured. As shown in Table 8, less than 6% of doses including Compounds 14, 15, and 16 remained in the liver after 48 hours. In contrast,

approximately 60% of doses including MC3 or Compound 22 remained in the liver after 48 hours. Less than 3% of the dose remained in the spleen for each composition tested.

Table 8. Lipids levels in the liver and spleen following administration of nanoparticle compositions including MC3 or compounds according to one of formulae (I), (Ia1)-(Ia6), (Ib), 5 (II), (IIa), (III), and (IIIa).

Compound No.	Lipid in liver (ng/g)	% remaining dose in liver	Lipid in spleen (ng/g)	% remaining dose in spleen
4	21950	38	5345	0.61
12	16850	23.8	2325	0.22
14	3990	5.54	1620	0.15
15	3070	4.22	971	0.089
16	597	0.79	293	0.026
22	36800	58.7	3887	0.41
23 (DOPE)	32900	51.4	26100	2.72
MC3	21750	51	2785	0.44

Example 8: Optimization of lipid:therapeutic agent ratios

[00924] The relative amounts of lipid component and therapeutic and/or prophylactic agent in a nanoparticle composition can be optimized according to considerations of efficacy and tolerability. For compositions including an RNA as a therapeutic and/or prophylactic agent, the N:P ratio can serve as a useful metric.

[00925] As the N:P ratio of a nanoparticle composition controls both expression and tolerability, nanoparticle compositions with low N:P ratios and strong expression are desirable. N:P ratios vary according to the ratio of lipids to RNA in a nanoparticle composition. Thus, the 15 wt/wt ratio of total lipid to RNA is varied between 10:1, 15:1, 20:1, 32:1, 40:1, 50:1, and 60:1 for a lipid formulation including about 50 mol % of a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I), about 10 mol % phospholipid (e.g., DOPE or DSPC), about 38.5 mol % structural lipid (e.g., cholesterol), and about 1.5 mol % PEG lipid (e.g., PEG-DMG). N:P ratios are calculated for each 20 nanoparticle composition assuming a single protonated nitrogen atom. The encapsulation efficiency (EE), size, and polydispersity index of each composition are also measured.

[00926] Generally, compositions with higher total lipid:RNA ratios yield smaller particles with higher encapsulation efficiencies, both of which are desirable. However, the N:P ratio for

such formulations generally exceeds 4. Current standards in the art such as the MC3 formulation described above have N:P ratios of 5.67. Thus, a balance between the N:P ratio, size, and encapsulation efficiency should be struck.

[00927] In order to explore the efficacy of nanoparticle compositions with different N:P ratios, the expression of luciferase (Luc) or human erythropoietin (hEPO) in mice after low (0.05 mg/kg) or high (0.5 mg/kg) doses of intravenously administered nanoparticle compositions is examined. The concentration of Luc or hEPO expressed is measured 3, 6, and/or 24 hours after administration.

10 Example 9: Optimization of content of a composition comprising a compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I)

15 **[00928]** As smaller particles with higher encapsulation efficiencies are generally desirable, the relative amounts of various elements in lipid components of nanoparticle compositions are optimized according to these parameters.

[00929] A compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) is selected for optimization. The relative amount of the compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) is varied between 30 mol % and 60 mol % in compositions including DOPE or DSPC as phospholipids to determine the optimal amount of the compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) in the formulations. Formulations are prepared using a standardized process with a water to ethanol ratio in the lipid-mRNA solution of 3:1 and a rate of injection of the lipid solution into the mRNA solution of 12 mL/min on a NanoAssemblr 25 microfluidic based system. This method induces nano-precipitation and particle formation. Alternative processes including, but not limited to, T-junction or direct injection, may also be used to achieve the same nano-precipitation.

[00930] Formulations producing the smallest particles with the highest encapsulation efficiencies are generally preferred, however larger or smaller particle sizes may be desirable based on a given application (e.g., based on the fenestration size of a target organ). Compositions are also evaluated for their Luc or hEPO expression levels and cytokine profiles.

Example 10: Optimization of phospholipid

[00931] The relative amount of phospholipid in a lipid component of a nanoparticle composition is varied to further optimize the formulation. A compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) is selected for use in the nanoparticle composition and DOPE and DSPC are selected as

5 phospholipids. Additional phospholipids can also be evaluated. Nanoparticle compositions are prepared with the relative phospholipid content varying between 0 mol % and 30 mol %. Compositions are evaluated for their size, encapsulation efficiency, Luc or hEPO expression levels, and cytokine profiles.

10 Example 11: Optimization of structural lipid

[00932] The relative amount of structural lipid in a lipid component of a nanoparticle composition is varied to further optimize the formulation. A compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) is selected for use in the nanoparticle composition and cholesterol is selected as a structural 15 lipid. Additional structural lipids can also be evaluated. Nanoparticle compositions are prepared with the relative structural lipid content varying between 18.5 mol % and 48.5 mol %. Compositions are evaluated for their size, encapsulation efficiency, Luc or hEPO expression levels, and cytokine profiles.

Example 12: Optimization of PEG lipid

20 **[00933]** The relative amount of PEG lipid in a lipid component of a nanoparticle composition is varied to further optimize the formulation. A compound according to one of formulae (I), (Ia1)-(Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I) is selected for use in the nanoparticle composition and PEG-DMG is selected as a PEG lipid. Additional PEG lipids can also be evaluated. Nanoparticle compositions are prepared with the 25 relative PEG lipid content varying between 0 mol % and 10 mol %. Compositions are evaluated for their size, encapsulation efficiency, Luc or hEPO expression levels, and cytokine profiles.

[00934] Exemplary formulations useful in the optimization of nanoparticle composition formulations are presented in Table 9.

Table 9. Exemplary formulations including compounds according to one of formulae (I), (Ia1)-

30 (Ia6), (Ib), (II), (IIa), (III), (IIIa), (IV), (17-I), (19-I), (19-II), (20-I) and (21-I).

Composition (mol %)	Components
40:20:38.5:1.5	Compound:Phospholipid:Chol:PEG-DMG

Composition (mol %)	Components
45:15:38.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
50:10:38.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
55:5:38.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
60:5:33.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
45:20:33.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
50:20:28.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
55:20:23.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
60:20:18.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
40:15:43.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
50:15:33.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
55:15:28.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
60:15:23.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
40:10:48.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
45:10:43.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
55:10:33.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
60:10:28.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
40:5:53.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
45:5:48.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
50:5:43.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
40:20:40:0	Compound:Phospholipid:Chol:PEG-DMG
45:20:35:0	Compound:Phospholipid:Chol:PEG-DMG
50:20:30:0	Compound:Phospholipid:Chol:PEG-DMG
55:20:25:0	Compound:Phospholipid:Chol:PEG-DMG
60:20:20:0	Compound:Phospholipid:Chol:PEG-DMG
40:15:45:0	Compound:Phospholipid:Chol:PEG-DMG
45:15:40:0	Compound:Phospholipid:Chol:PEG-DMG
50:15:35:0	Compound:Phospholipid:Chol:PEG-DMG
55:15:30:0	Compound:Phospholipid:Chol:PEG-DMG
60:15:25:0	Compound:Phospholipid:Chol:PEG-DMG
40:10:50:0	Compound:Phospholipid:Chol:PEG-DMG

Composition (mol %)	Components
45:10:45:0	Compound:Phospholipid:Chol:PEG-DMG
50:0:48.5:1.5	Compound:Phospholipid:Chol:PEG-DMG
50:10:40:0	Compound:Phospholipid:Chol:PEG-DMG
55:10:35:0	Compound:Phospholipid:Chol:PEG-DMG
60:10:30:0	Compound:Phospholipid:Chol:PEG-DMG

Example 13: Optimization of particle sizes

[00935] The fenestration sizes for different bodily organs often vary; for example, the kidney is known to have a smaller fenestration size than the liver. Thus, targeting delivery of a therapeutic and/or prophylactic agent (e.g., specifically delivering) to a particular organ or group of organs may require the administration of nanoparticle compositions with different particle sizes. In order to investigate this effect, nanoparticle compositions with formulations such as those included in Table 9 are prepared with a variety of particle sizes using a Nanoassemblr instrument. Nanoparticle compositions include an RNA encoding Luc. Each differently sized nanoparticle composition is subsequently administered to mice to evaluate the effect of particle size on delivery selectivity. Luc expression in two or more organs or groups of organs can be measured using bioluminescence to evaluate the relative expression in each organ.

Example 14: Administration following pretreatment

[00936] Administration of nanoparticle compositions to subjects can result in inflammation, infusion related reactions, and other undesirable effects indicative of low tolerability. These effects can be attributed to undesirable immunoactivity.

[00937] In order to combat negative effects, nanoparticle compositions are co-administered with one or more substances (e.g., co-medications or additional therapeutic and/or prophylactic agents) to subjects. Potentially useful additional therapeutic and/or prophylactic agents include steroids (e.g., corticosteroids), anti-histamines, H1 receptor blockers, H2 receptor blockers, anti-inflammatory compounds, statins, BTK inhibitors, S1P1 agonists, glucocorticoid receptor modulators (GRMs), and estradiols. Non-human primates are pretreated with one or more additional therapeutic agents selected from dexamethasone and acetaminophen. The additional therapeutic agent is administered either 24 hours, 1 hour, or both 24 hours and 1 hour before administration of a nanoparticle composition. Sample protocol are summarized in Table

10. Cytokine profiles, inflammation, and other parameters are measured and compared to evaluate the effectiveness of pretreatment.

Table 10. Sample protocol for pretreatment study.

Group	Pretreatment Time	Additional Therapeutic Agent(s) Administered
1	None	None
2	24 hours	Dexamethasone
3	24 hours	Acetaminophen
4	24 hours	Dexamethasone and Acetaminophen
5	1 hour	Dexamethasone
6	1 hour	Acetaminophen
7	1 hour	Dexamethasone and Acetaminophen
8	24 hours and 1 hour	Dexamethasone
9	24 hours and 1 hour	Acetaminophen
10	24 hours and 1 hour	Dexamethasone and Acetaminophen

5 [00938] For example, a useful therapeutic treatment course may involve administering an additional therapeutic and/or prophylactic agent both the day before and the day of (one hour prior) to administration of a nanoparticle composition at a dose level of 1.3 mpk. Additional therapeutic and/or prophylactic agents can be formulated for delivery by a variety of different routes. For example, dexamethasone may be delivered orally. In general, additional therapeutic 10 and/or prophylactic agents are administered at clinically approved or typical dosage levels.

Example 15: Administration to non-human primates

[00939] The tolerability and efficacy of nanoparticle compositions to non-human primates is evaluated in Cynomolgus monkeys. Monkeys are administered an optimized nanoparticle 15 composition including an mRNA encoding hEPO once weekly for four weeks. The levels of hEPO protein, mRNA, and cytokine profiles are measured using ELISA-based techniques before and 2, 6, 12, 24, 48, 72, and 120 hours after each administration.

[00940] The effects of pretreatment to non-human primates are evaluated using a standard MC3 formulation including an mRNA encoding hEPO. The study design is summarized in

Table 11. Male monkeys are administered the nanoparticle composition once weekly for four weeks at a dose rate of 5 ml/kg/h and are pretreated with either methotrexate or dexamethasone.

Table 11. Protocol for pretreatment study in Cynomolgus monkeys.

Group	Test Material	Dose level (mg/kg)	Additional Therapeutic Agent Administered	Dose concentration (mg/ml)	Number of monkeys
1	MC3	0	None	0	3
2	hEPO mRNA in MC3	0.3	None	0.06	3
3	hEPO mRNA in MC3	0.3	Methotrexate	0.06	3
4	hEPO mRNA in MC3	0.3	Dexamethasone	0.06	3

5 Example 16: Methods of treating diseases and disorders

[00941] A nanoparticle composition formulation having high tolerability (e.g., provoking a low immune response) and efficacy (e.g., facilitating efficient and effective encapsulation of a therapeutic and/or prophylactic agent and delivery of the agent to a desired target) is selected for use. A therapeutic and/or prophylactic agent for formulation with the nanoparticle composition

10 is selected for use based on the condition of a subject. For example, an mRNA encoding a vascular endothelial growth factor A (VEGF-A) may be selected to promote angiogenesis to treat atherosclerotic renovascular disease, while an siRNA capable of knocking down apolipoprotein B (apoB) may be selected to treat a metabolic disease or disorder such as dyslipidemia.

15 [00942] A subject in need of treatment is pretreated with a small dose of dexamethasone one or more hours prior to treatment with the nanoparticle composition. The nanoparticle composition is preferably administered to the subject intravenously, however intramuscular, intradermal, subcutaneous, intranasal, or inhalation administration routes are also acceptable. Treatment is provided in a dose of about 0.001 mg/kg to about 10 mg/kg of therapeutic and/or 20 prophylactic agent and is repeated daily, weekly, biweekly, or monthly according to needs of the subject.

Example 17: Expression induced by sample formulations upon intramuscular administration

[0072] Sample formulations including both modified luciferase (Luc) mRNA and H10 mRNA were prepared and administered intramuscularly at day 1 and day 21 and the resulting

5 expression and immunogenicity were evaluated simultaneously. Formulations including compounds according to formula (20-I) were prepared and administered at doses of 0.001 and 0.01 mpk (e.g., doses of 0.0005 mpk of a formulation including Luc mRNA and a formulation including H10 mRNA or doses of 0.005 mpk of a formulation including Luc mRNA and a formulation including H10 mRNA). As shown in Table 20-4, the total flux was higher upon 10 administration of the second dose for each composition and at each dose level. Total flux was higher for the higher dose level.

Table 12. Total flux (p/s) measured 6 hours after intramuscular administration of nanoparticle compositions including compounds according to formula (20-I).

Compound	0.001 mpk Dose 1	0.001 mpk Dose 2	0.01 mpk Dose 1	0.01 mpk Dose 2
20-1	3.50E+05	1.99E+06	3.16E+06	1.04E+07
20-4	1.04E+06	n.d.	9.46E+06	n.d.
20-6	8.57E+05	9.79E+05	3.41E+06	6.04E+06
MC3	1.23E+06	1.09E+06	1.79E+07	3.13E+07


15 [00943] It is to be understood that while the compounds and methods of the present disclosure have been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the present disclosure, which is defined by the scope of the appended claims. Other aspects, advantages, and alterations are within the scope of the following claims.

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.


It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

What is claimed is

1. A compound having formula (I) or formula (Ib):



or a salt or isomer thereof, wherein



t is 1 or 2;

A₁ and A₂ are each independently selected from CH or N;

A₆ and A₇ are each independently selected from CH or N, wherein at least one of A₆ and A₇ is N;

Z is CH₂ or absent, wherein when Z is CH₂, the dashed lines (1) and (2) each represent a single bond; and when Z is absent, the dashed lines (1) and (2) are both absent;

R₁, R₂, R₃, R₄, and R₅ are independently selected from the group consisting of C₅₋₂₀ alkyl, C₅₋₂₀ alkenyl, -R"MR', -R*YR", -YR", and -R*OR";

each M is independently selected from the group consisting of -C(O)O-, -OC(O)-, -OC(O)O-, -C(O)N(R')-, -N(R')C(O)-, -C(O)-, -C(S)-, -C(S)S-, -SC(S)-,

-CH(OH)-, -P(O)(OR')O-, -S(O)2-, an aryl group, and a heteroaryl group; wherein when the compound is of formula (Ib), then M is not -OC(O)O-;

X¹, X², and X³ are independently selected from the group consisting of a bond, -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;

X⁴ and X⁵ are independently selected from the group consisting of -CH₂-, -(CH₂)₂-, -CHR-, -CHY-, -C(O)-, -C(O)O-, -OC(O)-, -C(O)-CH₂-, -CH₂-C(O)-, -C(O)O-CH₂-, -OC(O)-CH₂-, -CH₂-C(O)O-, -CH₂-OC(O)-, -CH(OH)-, -C(S)-, and -CH(SH)-;


each Y is independently a C₃-6 carbocycle;

each R* is independently selected from the group consisting of C₁-12 alkyl and C₂-12 alkenyl;

each R is independently selected from the group consisting of C₁-3 alkyl and a C₃-6 carbocycle;

each R' is independently selected from the group consisting of C₁-12 alkyl, C₂-12 alkenyl, and H; and


each R'' is independently selected from the group consisting of C₃-12 alkyl and C₃-12 alkenyl;



wherein when the compound is of formula (I), and when ring A is , then

- i) at least one of X¹, X², and X³ is not -CH₂-; and/or
- ii) at least one of R₁, R₂, R₃, R₄, and R₅ is -R''MR';


or




when the compound is of Formula (I) and ring A is , and X¹, X², and X³ are each -CH₂, then


at least one of A¹ and A² is N,

R_1, R_2, R_3, R_4 and R_5 are independently selected from the group consisting of C_{6-20} alkyl and C_{6-20} alkenyl, and





when ring A is , then

- i) R_1, R_2, R_3, R_4 , and R_5 are the same, wherein R_1 is not C_{12} alkyl, C_{18} alkyl, or C_{18} alkenyl;
- ii) only one of R_1, R_2, R_3, R_4 and R_5 is selected from C_{6-20} alkenyl;
- iii) at least one of R_1, R_2, R_3, R_4 , and R_5 have a different number of carbon atoms than at least one other of R_1, R_2, R_3, R_4 , and R_5 ;
- iv) R_1, R_2 , and R_3 are selected from C_{6-20} alkenyl, and R_4 and R_5 are selected from C_{6-20} alkyl; or
- v) R_1, R_2 , and R_3 are selected from C_{6-20} alkyl, and R_4 and R_5 are selected from C_{6-20} alkenyl.



2. The compound of claim 1, wherein ring A is




or , in which the N atom is connected with X^2 .


3. The compound of any one of the preceding claims, wherein in formula (I):

- (i) at least one of A_1 and A_2 is N;
- (ii) A_1 is N and A_2 is CH;
- (iii) A_1 is CH and A_2 is N;
- (iv) A_1 is N and A_2 is N; or
- (v) each of A_1 and A_2 is CH.


4.. The compound of claim 1, being of any of formulae (Ia1)-(Ia6):




(Ia1),




(Ia2),




(Ia3),



(Ia4),



(Ia5), or



(Ia6).

5. The compound of any one of the preceding claims, wherein R₁, R₂, and R₃ are:


- (i) independently selected from the group consisting of C₅₋₂₀ alkyl and C₅₋₂₀ alkenyl;
- (ii) the same;
- (iii) C₆ alkyl;
- (iv) C₉ alkyl;
- (v) C₁₂ alkyl;
- (vi) C₁₄ alkyl;
- (vii) C₁₈ alkenyl; or
- (viii) linoleyl.

6. The compound of any one of the preceding claims, wherein

- (i) at least one of X¹ and X² is not -CH₂-;
- (ii) X¹ is not -CH₂-;
- (iii) X¹ is -C(O)-;
- (iv) X² is not -CH₂-; or
- (v) X² is -C(O)-.

7. The compound of any one of the preceding claims, wherein R₂, R₃, R₄, and R₅ are the same.

8. The compound of claim 1, wherein the compound is Compound 4:



or a salt or isomer thereof.

9. A nanoparticle composition comprising a lipid component comprising a compound of any one of the preceding claims, wherein the lipid component further comprises (i) a phospholipid; (ii) a structural lipid, and/or (iii) a PEG lipid.

10. The nanoparticle composition of claim 9, wherein the phospholipid is selected from the group consisting of 1,2-dilinoleoyl-sn-glycero-3-phosphocholine (DLPC),

1,2-dimyristoyl-sn-glycero-phosphocholine (DMPC), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC),
1,2-distearoyl-sn-glycero-3-phosphocholine (DSPC),
1,2-diundecanoyl-sn-glycero-phosphocholine (DUPC),
1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC),
1,2-di-O-octadecenyl-*sn*-glycero-3-phosphocholine (18:0 Diether PC),
1-oleoyl-2-cholesterylhemisuccinoyl-*sn*-glycero-3-phosphocholine (OChemSPC),
1-hexadecyl-sn-glycero-3-phosphocholine (C16 Lyso PC),
1,2-dilinolenoyl-sn-glycero-3-phosphocholine, 1,2-diarachidonoyl-sn-glycero-3-phosphocholine,
1,2-didocosahexaenoyl-sn-glycero-3-phosphocholine, 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (DOPE), 1,2-diphytanoyl-sn-glycero-3-phosphoethanolamine (ME 16.0 PE),
1,2-distearoyl-sn-glycero-3-phosphoethanolamine,
1,2-dilinoleoyl-sn-glycero-3-phosphoethanolamine,
1,2-dilinolenoyl-sn-glycero-3-phosphoethanolamine,
1,2-diarachidonoyl-sn-glycero-3-phosphoethanolamine,
1,2-didocosahexaenoyl-sn-glycero-3-phosphoethanolamine,
1,2-dioleoyl-sn-glycero-3-phospho-rac-(1-glycerol) sodium salt (DOPG), sphingomyelin, and mixtures thereof.

11. The nanoparticle composition of claim 9 or 10, wherein the structural lipid is selected from the group consisting of cholesterol, fecosterol, sitosterol, ergosterol, campesterol, stigmasterol, brassicasterol, tomatidine, ursolic acid, alpha-tocopherol, and mixtures thereof.

12. The nanoparticle composition of any one of claims 9-11, wherein the PEG lipid is selected from the group consisting of a PEG-modified phosphatidylethanolamine, a PEG-modified phosphatidic acid, a PEG-modified ceramide, a PEG-modified dialkylamine, a PEG-modified diacylglycerol, a PEG-modified dialkylglycerol, and mixtures thereof.

13. The nanoparticle composition of any one of claims 9-12, further comprising a therapeutic and/or prophylactic agent.

14. The nanoparticle composition of claim 13, wherein the therapeutic and/or prophylactic agent is a ribonucleic acid (RNA).

15. The nanoparticle composition of claim 14, wherein the RNA is selected from the group consisting of a small interfering RNA (siRNA), an asymmetrical interfering RNA (aiRNA), a microRNA (miRNA), a Dicer-substrate RNA (dsRNA), a small hairpin RNA (shRNA), a messenger RNA (mRNA), and mixtures thereof.
16. A method of delivering a therapeutic and/or prophylactic agent to a mammalian cell, the method comprising administering to a subject the nanoparticle composition of any one of claims 13-15, said administering comprising contacting the cell with the nanoparticle composition, whereby the therapeutic and/or prophylactic agent is delivered to the cell.
17. A method of producing a polypeptide of interest in a mammalian cell, the method comprising contacting the cell with the nanoparticle composition of any one of claims 13-15, wherein the therapeutic and/or prophylactic agent is an mRNA, and wherein the mRNA encodes the polypeptide of interest, whereby the mRNA is capable of being translated in the cell to produce the polypeptide of interest.
18. A method of treating a disease or disorder in a mammal in need thereof, the method comprising administering to the mammal a therapeutically effective amount of the nanoparticle composition of any one of claims 13-15, wherein the disease or disorder is characterized by dysfunctional or aberrant protein or polypeptide activity.
19. The method of claim 18, wherein the mammal (a) is LDLR-deficient; (b) is apoE-deficient; and/or (c) mammal has an abnormal LDLR-apoE interaction.
20. The method of claim 18, wherein the mammal (a) is not LDLR-deficient; (b) is not apoE-deficient; and/or (c) has a normal LDLR-apoE interaction.
21. Use of the compound of any one of claims 1 to 28, or the nanoparticle composition of any one of claims 13-15, in the manufacture of a medicament for delivering a therapeutic and/or prophylactic agent to a mammalian cell and/or treating a disease or disorder in a mammal in need thereof, wherein the disease or disorder is characterized by dysfunctional or aberrant protein or polypeptide activity.

FIGURE 1
Whole Body Luc Expression





**FIGURE 2****FIGURE 3**






FIGURE 4**FIGURE 5****FIGURE 6****FIGURE 7****FIGURE 8**




FIGURE 9**FIGURE 10****FIGURE 11**



FIGURE 12**FIGURE 13**




FIGURE 14**FIGURE 15**





FIGURE 16**FIGURE 17****FIGURE 18**

FIGURE 19**FIGURE 20**