

(19) **DANMARK**

(10) **DK/EP 3075855 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 15/113 (2010.01)** **A 61 K 31/713 (2006.01)** **C 12 N 15/11 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-10-08**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2018-08-15**
- (86) Europæisk ansøgning nr.: **16169874.1**
- (86) Europæisk indleveringsdag: **2012-06-08**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-10-05**
- (30) Prioritet: **2011-06-08 US 201161494832 P** **2011-06-15 US 201161497447 P**
- (62) Stamansøgningsnr: **12730301.4**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **NITTO DENKO CORPORATION, 1-1-2, Shimohozumi, Ibaraki-shi, Osaka 567-8680, Japan**
- (72) Opfinder: **AKOPIAN, Violetta, 12610 Torrey Bluff Dr., Apt. 370, San Diego, CA 92130, USA**
MINOMI, Kenjiro, 1-1-2 Shimohozumi, Ibaraki, Osaka, 567-8680, Japan
NIITSU, Yoshiro, 2-1-901, Kita 3-jo, Nishi 30-chome, Chuo-ku, Sapporo-shi, Hokkaido, 064-0823, Japan
PAYNE, Joseph E., 4810 Milano Way, Oceanside, CA 92057, USA
KNOPOV, Victor, 2104 SeaCliff Way, Oceanside, CA 92056, USA
WITTE, Richard P., 3462 Spring Tide Terrace, San Diego, CA 92110, USA
AHMADIAN, Mohammad, 3230 Rancho Quartillo, Carlsbad, CA 92009, USA
PERELMAN, Loren A., 458 Alcatraz Ave, Oakland, CA 94609, USA
TANAKA, Yasunobu, 1-1-2 Shimohozumi Ibaraki, Osaka, 567-8680, Japan
FEINSTEIN, Elena, 13/55 HaCarmel Street, Rehovot, Israel
AVKIN-NACHUM, Sharon, 4 Hakramim St., Nes Ziona, Israel
KALINSKI, Hagar, 1 Haim Landau Street, Richon Le-Zion, Israel
METT, Igor, 9/1 Paldi Street, Rehovot, Israel
YING, Wenbin, 7478 Park Village Road, San Diego, CA 92129, USA
LUI, Yun, 8428 Florissant Court, San Diego, CA 92129, USA
- (74) Fuldmægtig i Danmark: **Plougmann Vingtoft A/S, Rued Langgaards Vej 8, 2300 København S, Danmark**
- (54) Benævnelse: **RETINOID-LIPOSOMER TIL FORBEDRING AF MODULERING AF HSP47-EKSPRESSION**
- (56) Fremdragne publikationer:
EP-A1- 1 842 557
YASUSHI SATO ET AL.: "Resolution of liver cirrhosis using vitamin A-coupled liposomes to deliver siRNA against a collagen-specific chaperone", NATURE BIOTECHNOLOGY, vol. 26, no. 4, 1 April 2008 (2008-04-01), pages 431-442, XP055000721, ISSN: 1087-0156, DOI: 10.1038/nbt1396 -& YASUSHI SATO ET AL.: "Supplementary Text and Figures", NATURE BIOTECHNOLOGY, vol. 26, no. 04 April 2008 (2008-04), XP002689068, Retrieved from the Internet: URL:http://www.nature.com/nbt/journal/v26/n4/supinfo/nbt1396_S1.html [retrieved on 2012-12-04]

Fortsættes ...

XIA Z. ET AL.: "Suppression of renal tubulointerstitial fibrosis by small interfering RNA targeting heat shock protein 47", AMERICAN JOURNAL OF NEPHROLOGY, S. KARGER AG, CH, vol. 28, no. 1, 21 September 2007 (2007-09-21), pages 34-46, XP008161356, ISSN: 0250-8095

WANG Z. ET AL.: "Adenovirus-mediated RNA interference against collagen-specific molecular chaperone 47-KDa heat shock protein suppresses scar formation on mouse wounds", CELL BIOLOGY INTERNATIONAL, ACADEMIC PRESS, GB, vol. 32, no. 5, 1 May 2008 (2008-05-01), pages 484-493, XP022681968, ISSN: 1065-6995, DOI: 10.1016/J.CELLBI.2007.10.009 [retrieved on 2007-11-05]

DESCRIPTION

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of United States Provisional Application No. 61/497,447 filed June 15, 2011 and United States Provisional Application No. 61/494,832 filed June 8, 2011.

TECHNICAL FIELD

[0002] Provided herein are pharmaceutical compositions comprising retinoid-liposomes for enhancing the modulation of hsp47 expression by siRNA.

BACKGROUND

[0003] Fibrosis of the liver can be caused by activated hepatic stellate cells (HSC), resulting in a plurality of types of collagen molecules and fibronectin being deposited on interstitial tissue. This can lead to hepatic cirrhosis, hepatic failure, and/or hepatocellular carcinoma. Further, chronic pancreatitis develops as a result of pancreatic fibrosis by the same mechanism as that for hepatic fibrosis (Madro, et al., 2004; Med Sci Monit. 10:RA166-70; Jaster, 2004, Mol Cancer. 6:26). Furthermore, stellate cells are present in disorders of the vocal cord and larynx such as vocal cord scarring, vocal cord mucosal fibrosis, and laryngeal fibrosis. To prevent or treat fibrosis in these organs and elsewhere in the body, there is a desire for the development of a drug carrier and drug carrier kit.

[0004] Stellate cells are one of the important target candidates for treating fibrosis (Fallowfield et al., 2004, Expert Opin Ther Targets. 8:423-35; Pinzani, et al., 2004, Dig Liver Dis.36:231-42). During fibrosis, stellate cells are activated by cytokines from nearby cells to produce many factors that cause hepatic fibrosis. Stellate cells store vitamin A, and belong to the myofibroblast family.

[0005] Therapeutic methods to prevent or treat fibrosis attempt to control collagen metabolism, promotion of the collagen degradation system, and inhibition of activation of stellate cells. However, in all these cases, the low specificity of action and/or the low organ specificity, limited efficacy and adverse side effects create problems.

[0006] Inhibition of collagen protein synthesis has not been established as a therapeutic method. The potency of molecules targeting collagen production is limited due to the possibility of causing side effects. Inhibiting collagen production directly provides another therapeutic method to prevent or treat fibrosis. Such a method requires controlling one or more of the various types of collagen Types I to IV. A method for accomplishing this may be through heat shock protein47 (HSP47), a collagen-specific molecular chaperone that is essential for intracellular transport and molecular maturation necessary for various types of collagen. Therefore, if the function of HSP47 can be specifically controlled in stellate cells, there is a possibility of inhibiting hepatic fibrosis.


[0007] EP1842557 concerns drug carrier and drug carrier kit for inhibiting fibrosis.

[0008] Sato et al. (Nature Biotechnology; Volume 26, Number 4; April 2008; pages 431 - 442) concerns resolution of liver cirrhosis using vitamin A-coupled liposomes to deliver siRNA against a collagen-specific chaperone.

SUMMARY

[0009] The current invention is defined, *inter alia*, by the following items:

1. 1. A pharmaceutical composition comprising:
 1. (i) a double-stranded nucleic acid molecule and drug carrier comprising a mixture of a retinoid and a lipid, wherein the double-stranded nucleic acid molecule comprises the structure (A1):



wherein each of N and N' is a nucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein each of Z and Z' is independently present or absent, but if present independently includes 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present;


wherein Z'' may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of (N')_y;

wherein each of x and y is independently an integer between 18 and 40;


wherein the sequence of (N')_y has complementary to the sequence of (N)_x;

wherein (N)_x includes an antisense sequence to the mRNA coding sequence for human hsp47 exemplified by SEQ ID NO:1; and

wherein the lipid comprises a lipid selected from the group consisting of




and




or

2. (ii) a double-stranded nucleic acid molecule comprising a sense strand and an antisense strand wherein the sense and antisense strands are selected from the oligonucleotides described as SERPINH1_2 (SEQ ID NOS: 60 and 127), SERPINH1_45a (SEQ ID NOS: 98 and 165), and SERPINH1_51 (SEQ ID NOS: 101 and 168), and a drug carrier comprising a mixture of a lipid vesicle and a retinoid or a retinoid conjugate, wherein a lipid of the lipid vesicle comprises a lipid selected from the group consisting of



and



2. The pharmaceutical composition of item 1, wherein the retinoid conjugate is a PEG-conjugate.
3. The pharmaceutical composition according to any one of items 1-2, wherein the antisense strand comprises SEQ ID NO:127, and the sense strand comprises SEQ ID NO:60; wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 1, 5, 6, or 7; and a non-nucleotide moiety covalently attached to the 3'-terminus; and wherein the sense strand further comprises at least one 2'-5'-ribonucleotide or 2'-O-methyl modified ribonucleotide; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus.
4. The pharmaceutical composition according to any one of items 1-2, wherein the antisense strand comprises SEQ ID NO:127, and the sense strand comprises SEQ ID NO:60; wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides at positions 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
5. The pharmaceutical composition according to any one of items 1-2, wherein the antisense strand comprises SEQ ID NO:127, and the sense strand comprises SEQ ID NO:60; wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides at positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; and a C3C3 non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises a C3Pi non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
6. The pharmaceutical composition according to any one of items 1-2, wherein the antisense strand comprises SEQ ID NO:127, and the sense strand comprises SEQ ID NO:60; wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides at positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a C3Pi non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
7. The pharmaceutical composition according to any one of items 1-2, wherein the antisense strand comprises SEQ ID NO:127, and the sense strand comprises SEQ ID NO:60; wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide at position 9; a C3 non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
8. The pharmaceutical composition according to any one of items 1-2, wherein the sense strand comprises SEQ ID NO:98 and the antisense strand comprises SEQ ID NO:165; wherein the sense strand further comprises 2'-5'-ribonucleotides in positions at or near the 3'-terminus; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises a 2'-O-methyl modified ribonucleotide; a 2'-5'-ribonucleotide in at least one of positions 5, 6 or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus.
9. The pharmaceutical composition according to any one of items 1-2, wherein the sense strand comprises SEQ ID NO:98 and the antisense strand comprises SEQ ID NO:165; wherein the sense strand further comprises 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3-OH 3' moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

10. 10. The pharmaceutical composition according to any one of items 1-2, wherein the sense strand comprises SEQ ID NO:101 and the antisense strand comprises SEQ ID NO:168; wherein the sense strand further comprises 2'-O-methyl modified pyrimidine ribonucleotides; an optional 2'-5'-ribonucleotide in one of position 9 or 10; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises a 2'-O-methyl modified ribonucleotide; a 2'-5'-ribonucleotide in at least one of positions 5, 6, or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus.
11. 11. The pharmaceutical composition according to any one of items 1-2, wherein the sense strand comprises SEQ ID NO:101 and the antisense strand comprises SEQ ID NO:168; wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3PI-C3OH moiety covalently attached at the 3'-terminus.
12. 12. The pharmaceutical composition according to any one of items 1-2, wherein the sense strand comprises SEQ ID NO:101 and the antisense strand comprises SEQ ID NO:168; wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 4, 11, 13, and 17; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 4, 8, 13 and 15; a 2-5-ribonucleotide in position 6; and a C3PI-C3OH moiety covalently attached at the 3'-terminus.
13. 13. The pharmaceutical composition according to any one of items 1-2, wherein the sense strand comprises SEQ ID NO:101 and the antisense strand comprises SEQ ID NO:168; wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 2, 4, 11, 13, and 17; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3PI-C3OH non-nucleotide moiety covalently attached at the 3'-terminus.

[0010] The present description relates to a drug carrier and a drug carrier kit that enable a diagnostic and/or therapeutic drug to be specifically transported to stellate cells. The drug carrier in the present description may be selected from polymer micelles, liposomes, emulsions, microspheres, and nanosphere forms, and by bonding thereto or including therein a retinoid or retinoid conjugate and a therapeutic drug that can be transported specifically to HSC. Retinoids include vitamin A, retinal, retinoic acid, saturated Vitamin A, tretinoin, adapalene, or retinol palmitate, and fenretinide (4-HPR). Furthermore, by preparing the drug carrier to include one molecule or a plurality of molecules selected from TGF β activity inhibitors such as a truncated TGF β type II receptor and a soluble TGF β type II receptor, growth factor preparations such as HGF, MMP production promoters such as an MMP gene-containing adenovirus vector, a cell activation inhibitors and/or growth inhibitors including a PPAR γ -ligand, an angiotensin-II type I receptor antagonist, a PDGF tyrosine kinase inhibitor, and a sodium channel inhibitor such as amiloride, and apoptosis inducers such as compound 861 and gliotoxin; and by administering it for example, orally, parenterally, intravenously or intraperitoneally to a patient having a risk of fibrosis or fibrosis symptoms, or patients having various fibrosis-related disorders such as, for example, hepatic cirrhosis, hepatic failure, liver cancer, or chronic pancreatitis, the activation of stellate cells can be suppressed, and thereby preventing, inhibiting or improving the fibrosis and/or fibrosis-related disease conditions in said patient. Alternatively, or in addition thereto, by using the drug carrier which encloses therein a ribozyme, an antisense RNA, or an siRNA that specifically inhibits HSP47 or TIMP, which is an MMP inhibitor, secretion of type I to IV collagens can be simultaneously inhibited, and as a result fibrogenesis can be inhibited effectively.

[0011] An embodiment of the description is a pharmaceutical composition comprising a double-stranded nucleic acid molecule comprising a sense strand and an antisense strand wherein the sense and antisense strands are selected from the oligonucleotides described as SERPINH1_2 (SEQ ID NOS: 60 and 127), SERPINH1_45a (SEQ ID NOS: 98 and 165), and SERPINH1_51 (SEQ ID NOS: 101 and 168) in Table 4, *infra*, and a drug carrier comprising a mixture of a lipid vesicle and a retinoid or a retinoid conjugate. The retinoid can be one or more of the following: vitamin A, retinoic acid, saturated Vitamin A, retinal, tretinoin, adapalene, retinol palmitate, or fenretinide. Preferably, the retinoid comprises a conjugate of retinoic acid, most preferably a retinoid-PEG conjugate. The lipid vesicle can be comprised of a bilayer of lipid molecules, and can further be comprised of the retinoid. The retinoid is preferably at a concentration of 0.2 to 20 wt % in the drug carrier. The lipid vesicle can be comprised of an interior surface that encapsulates the interior of the lipid vesicle, and an exterior surface that is accessible to an aqueous medium outside of the lipid vesicle. The retinoid may be associated with the lipid bilayer. The double-stranded nucleic acid can be exposed on the exterior surface of the lipid vesicle.


[0012] In some embodiments, the double-stranded nucleic acid molecule includes an antisense strand having SEQ ID NO:127 and comprising 2'-O-methyl sugar (2'OMe)-modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 1, 5, 6, or 7; and a 3'-terminal non-nucleotide moiety covalently attached to the 3'-terminus; and a sense strand having SEQ ID NO:60 and comprising at least one 2'-5'-ribonucleotide or 2'OMe modified ribonucleotide; a non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus. In preferred embodiments the antisense strand is SEQ ID NO:127 and comprises 2'OMe modified ribonucleotides at positions 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a non-nucleotide moiety covalently attached at the 3'-terminus; and the sense strand is SEQ ID NO:60 and comprises five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus. In some embodiments the double-stranded nucleic acid molecule further includes a 2'OMe modified ribonucleotide or a 2'-5'-ribonucleotide at position 1 of the antisense strand. (All reference herein to nucleotide positions are expressed based on the 5'→3' direction of the oligonucleotide for both sense and antisense strands of the double-stranded nucleic acid molecule.)

[0013] In various embodiments, the sense strand is SEQ ID NO:98 and comprises 2'-5'-ribonucleotides in positions at the 3'-terminus; a non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:165 and comprises 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 5, 6 or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus. In preferred embodiments the sense strand is SEQ ID NO:98 and comprises 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3-OH 3' moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:165 and comprises 2'OMe modified

ribonucleotides in positions 4, 6, 8, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3PI-C3OH moiety covalently attached at the 3'-terminus. In some embodiments the double-stranded nucleic acid further comprises a 2'OMe modified ribonucleotide in position 2.

[0014] In various embodiments, the sense strand is SEQ ID NO:101 and comprises 2'OMe modified ribonucleotides; an optional 2'-5'-ribonucleotide in one of position 9 or 10; a non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:168 and comprises 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 5, 6, or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus. In preferred embodiments the sense strand is SEQ ID NO:101 and comprises 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:168 and comprises 2'OMe modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3PI-C3OH moiety covalently attached at the 3'-terminus. In certain embodiments the double-stranded nucleic acid molecule further comprises a 2'OMe modified ribonucleotide in position 13 in the antisense strand and or in position 2 in the sense strand.

[0015] Another aspect is a pharmaceutical composition comprising a double-stranded nucleic acid molecule and drug carrier comprising a mixture of a retinoid and a lipid, wherein the double-stranded oligonucleotide compound comprising the structure (A1):



wherein each of N and N' is a nucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of $(N)_x$ and $(N')_y$ is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein each of Z and Z' is independently present or absent, but if present independently includes 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present;

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of (N')_n;

wherein each of x and y is independently an integer between 18 and 40; wherein the sequence of (N')_y has complementary to the sequence of (N)_x; and wherein (N)_x includes an antisense sequence to the mRNA coding sequence for human hsp47 exemplified by SEQ ID NO:1, which is shown as follows.

[illegible]

[0016] Compositions, methods and kits for modulating expression of target genes are provided herein. In various aspects and embodiments, compositions, methods and kits provided herein modulate expression of heat shock protein 47 (hsp47), also known as SERPINH1 (SEQ ID NO:1). The compositions, methods and kits may involve use of nucleic acid molecules (for example, short interfering nucleic acid (siNA), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), or short hairpin RNA (shRNA)) that bind a nucleotide sequence (such as an mRNA sequence) encoding hsp47, SEQ ID NO:1. In certain preferred embodiments, the compositions, methods and kits disclosed herein inhibit expression of hsp47. For example, siNA molecules (e.g., RNA-induced silencing complex (RISC) length dsNA molecules or Dicer length dsNA molecules) are provided that reduce or inhibit hsp47 expression. Also provided are compositions, methods and kits for treating and/or preventing diseases, conditions or disorders associated with hsp47, such as liver fibrosis, cirrhosis, pulmonary fibrosis including lung fibrosis (including interstitial lung fibrosis (ILF)), kidney fibrosis resulting from any condition (e.g., chronic kidney disease (CKD) including End-Stage Renal Disease (ESRD)), peritoneal fibrosis, chronic hepatic damage, fibrillogenesis, fibrotic diseases in other organs, abnormal scarring (keloids) associated with all possible types of skin injury accidental and iatrogenic (operations); scleroderma; cardiofibrosis, failure of glaucoma filtering operation; and intestinal adhesions.

[0017] In one aspect, provided are the pharmaceutical compositions, above, comprising nucleic acid molecules (e.g., siRNA molecules) as a component of a pharmaceutical formulation in which the nucleic acid molecule includes a sense strand and an antisense strand; each strand of the nucleic acid molecule is independently 15 to 49 nucleotides in length; a 15 to 49 nucleotide sequence of the antisense strand is complementary to a sequence of an mRNA encoding human hsp47 (e.g., SEQ ID NO: 1); and a 15 to 49 nucleotide sequence of the sense strand is complementary

to the a sequence of the antisense strand and includes a 15 to 49 nucleotide sequence of an mRNA encoding human hsp47 (e.g., SEQ ID NO: 1).

[0018] In certain embodiments, the sequence of the antisense strand that is complementary to a sequence of an mRNA encoding human hsp47 includes a sequence complimentary to a sequence between nucleotides 600-800; or 801-899; or 900-1000; or 1001-1300 of SEQ ID NO: 1; or between nucleotides 650-730; or 900-975 of SEQ ID NO: 1. In some embodiments, the antisense strand includes a sequence that is complementary to a sequence of an mRNA encoding human hsp47 corresponding to nucleotides 674-693 of SEQ ID NO: 1 or a portion thereof; or nucleotides 698-716 of SEQ ID NO: 1 or a portion thereof; or nucleotides 698-722 of SEQ ID NO: 1 or a portion thereof; or nucleotides 701-720 of SEQ ID NO: 1 or a portion thereof; or nucleotides 920-939 of SEQ ID NO: 1 or a portion thereof; or nucleotides 963-982 of SEQ ID NO: 1 or a portion thereof; or nucleotides 947-972 of SEQ ID NO: 1 or a portion thereof; or nucleotides 948-966 of SEQ ID NO: 1 or a portion thereof; or nucleotides 945-969 of SEQ ID NO: 1 or a portion thereof; or nucleotides 945-963 of SEQ ID NO: 1 or a portion thereof.

[0019] In certain embodiments, the antisense strand of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein as a component of a pharmaceutical formulation includes a sequence corresponding to SEQ ID NO: 4 or a portion thereof; or SEQ ID NO: 6 or a portion thereof; or SEQ ID NO: 8 or a portion thereof; or SEQ ID NO: 10 or a portion thereof; or SEQ ID NO: 12 or a portion thereof; or SEQ ID NO: 14 or a portion thereof; or SEQ ID NO: 16 or a portion thereof; or SEQ ID NO: 18 or a portion thereof; or SEQ ID NO: 20 or a portion thereof; or SEQ ID NO: 22 or a portion thereof; or SEQ ID NO: 24 or a portion thereof; or SEQ ID NO: 26 or a portion thereof; or SEQ ID NO: 28 or a portion thereof; or SEQ ID NO: 30 or a portion thereof; or SEQ ID NO: 32 or a portion thereof; or SEQ ID NO: 34 or a portion thereof; or SEQ ID NO: 36 or a portion thereof; or SEQ ID NO: 38 or a portion thereof; or SEQ ID NO: 40 or a portion thereof; or SEQ ID NO: 42 or a portion thereof; or SEQ ID NO: 44 or a portion thereof; or SEQ ID NO: 46 or a portion thereof; or SEQ ID NO: 48 or a portion thereof; or SEQ ID NO: 50 or a portion thereof; or SEQ ID NO: 52 or a portion thereof; or SEQ ID NO: 54 or a portion thereof; or SEQ ID NO: 56 or a portion thereof; or SEQ ID NO: 58 or a portion thereof. In certain embodiments, the sense strand of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein includes a sequence corresponding to SEQ ID NO: 3 or a portion thereof; or SEQ ID NO: 5 or a portion thereof; or SEQ ID NO: 7 or a portion thereof; or SEQ ID NO: 9 or a portion thereof; or SEQ ID NO: 11 or a portion thereof; or SEQ ID NO: 13 or a portion thereof; or SEQ ID NO: 15 or a portion thereof; or SEQ ID NO: 17 or a portion thereof; or SEQ ID NO: 19 or a portion thereof; or SEQ ID NO: 21 or a portion thereof; or SEQ ID NO: 23 or a portion thereof; or SEQ ID NO: 25 or a portion thereof; or SEQ ID NO: 27 or a portion thereof; or SEQ ID NO: 29 or a portion thereof; or SEQ ID NO: 31 or a portion thereof; or SEQ ID NO: 33 or a portion thereof; or SEQ ID NO: 35 or a portion thereof; or SEQ ID NO: 37 or a portion thereof; or SEQ ID NO: 39 or a portion thereof; or SEQ ID NO: 41 or a portion thereof; or SEQ ID NO: 43 or a portion thereof; or SEQ ID NO: 45 or a portion thereof; or SEQ ID NO: 47 or a portion thereof; or SEQ ID NO: 49 or a portion thereof; or SEQ ID NO: 51 or a portion thereof; or SEQ ID NO: 53 or a portion thereof; or SEQ ID NO: 55 or a portion thereof; or SEQ ID NO: 57 or a portion thereof.

[0020] In certain preferred embodiments, the antisense strand of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein as a component of a pharmaceutical formulation includes a sequence corresponding to any one of the antisense sequences shown on Table 4. In certain preferred embodiments the antisense strand and the strand are selected from the sequence pairs shown in Table 4. In some embodiments the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 and SERPINH1_88. In some embodiments the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_4 (SEQ ID NOS:195 and 220), SERPINH1_12 (SEQ ID NOS:196 and 221), SERPINH1_30 (SEQ ID NOS:199 and 224), and SERPINH1_58 (SEQ ID NOS:208 and 233).

[0021] In some embodiments, the antisense and sense strands of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein as a component of a pharmaceutical formulation includes the sequence pairs set forth in SERPINH1_4 (SEQ ID NOS:195 and 220). In some embodiments of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein includes the antisense and sense strands of the sequence pairs set forth in SERPINH1_12 (SEQ ID NOS:196 and 221). In some embodiments the antisense and sense strands of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein includes the sequence pairs set forth in SERPINH1_30 (SEQ ID NOS:199 and 224). In some embodiments of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein includes the antisense and sense strands of the sequence pairs set forth in SERPINH1_58 (SEQ ID NOS:208 and 233).

[0022] In certain embodiments, the antisense strand of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein as a component of a pharmaceutical formulation includes a sequence corresponding to any one of the antisense sequences shown on any one of Tables B or C.

[0023] In certain preferred embodiments, the antisense strand of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein as a component of a pharmaceutical formulation includes a sequence corresponding to any one of the antisense sequences shown on Table 5. In certain preferred embodiments the antisense strand and the strand are selected from the sequence pairs shown in Table 5. In some embodiments of a nucleic acid molecule (e.g., a siNA molecule) as disclosed herein includes the antisense and sense strands selected from the sequence pairs set forth in

- SERPINH1_2 (SEQ ID NOS: 60 and 127),
- SERPINH1_6 (SEQ ID NOS: 63 and 130),
- SERPINH1_11 (SEQ ID NOS: 68 and 135),
- SERPINH1_13 (SEQ ID NOS: 69 and 136),
- SERPINH1_45 (SEQ ID NOS: 97 and 164),
- SERPINH1_45a (SEQ ID NOS: 98 and 165),
- SERPINH1_51 (SEQ ID NOS: 101 and 168),
- SERPINH1_52 (SEQ ID NOS: 102 and 169) or
- SERPINH1_86 (SEQ ID NOS: 123 and 190).

[0024] In some preferred embodiments the antisense and sense strands are selected from the sequence pairs set forth in

- SERPINH1_2 (SEQ ID NOS: 60 and 127),
- SERPINH1_6 (SEQ ID NOS: 63 and 130),
- SERPINH1_45a (SEQ ID NOS: 98 and 165), and
- SERPINH1_51 (SEQ ID NOS: 101 and 168).

[0025] In some preferred embodiments of a nucleic acid molecule (*e.g.*, a siNA molecule) as disclosed herein as a component of a pharmaceutical formulation includes the antisense and sense strands selected from the sequence pairs set forth in SERPINH1_2 (SEQ ID NOS: 60 and 127). In some embodiments the antisense and sense strands include the sequence pairs set forth in SERPINH1_6 (SEQ ID NOS: 63 and 130). In some embodiments of a nucleic acid molecule (*e.g.*, a siNA molecule) as disclosed herein includes the antisense and sense strands of the sequence pairs set forth in SERPINH1_11 (SEQ ID NOS: 68 and 135). In some embodiments the antisense and sense strands are the sequence pairs set forth in SERPINH1_13 (SEQ ID NOS: 69 and 136). In some embodiments the antisense and sense strands are the sequence pairs set forth in SERPINH1_45 (SEQ ID NOS: 97 and 164). In some embodiments the antisense and sense strands are the sequence pairs set forth in SERPINH1_45a (SEQ ID NOS: 98 and 165). In some embodiments the antisense and sense strands are the sequence pairs set forth in SERPINH1_51 (SEQ ID NOS: 101 and 168).

[0026] In certain embodiments, the antisense strand of a nucleic acid molecule (*e.g.*, a siNA molecule) as disclosed as a component of a pharmaceutical formulation herein includes a sequence corresponding to any one of the antisense sequences shown on any one of Tables D or E.

[0027] In various embodiments of nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein as a component of a pharmaceutical formulation, the antisense strand may be 15 to 49 nucleotides in length (*e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 or 49 nucleotides in length); or 17-35 nucleotides in length; or 17-30 nucleotides in length; or 15-25 nucleotides in length; or 18-25 nucleotides in length; or 18-23 nucleotides in length; or 19-21 nucleotides in length; or 25-30 nucleotides in length; or 26-28 nucleotides in length. In some embodiments of nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein, the antisense strand may be 19 nucleotides in length. Similarly the sense strand of nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein may be 15 to 49 nucleotides in length (*e.g.*, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 or 49 nucleotides in length); or 17-35 nucleotides in length; or 17-30 nucleotides in length; or 15-25 nucleotides in length; or 18-25 nucleotides in length; or 18-23 nucleotides in length; or 19-21 nucleotides in length; or 25-30 nucleotides in length; or 26-28 nucleotides in length. In some embodiments of nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein, the sense strand may be 19 nucleotides in length. In some embodiments of nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein, the antisense strand and the sense strand may be 19 nucleotides in length. The duplex region of the nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein may be 15-49 nucleotides in length (*e.g.*, about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 or 49 nucleotides in length), 15-35 nucleotides in length; or 15-30 nucleotides in length; or about 15-25 nucleotides in length; or 17-25 nucleotides in length; or 17-23 nucleotides in length; or 17-21 nucleotides in length; or 25-30 nucleotides in length; or 25-28 nucleotides in length. In various embodiments of nucleic acid molecules (*e.g.*, siNA molecules) as disclosed herein, the duplex region may be 19 nucleotides in length.

[0028] In certain embodiments, the sense and antisense strands of a nucleic acid (*e.g.*, an siNA nucleic acid molecule) as provided herein as a component of a pharmaceutical formulation are separate polynucleotide strands. In some embodiments, the separate antisense and sense strands form a double-stranded structure via hydrogen bonding, for example, Watson-Crick base pairing. In some embodiments the sense and antisense strands are two separate strands that are covalently linked to each other. In other embodiments, the sense and antisense strands are part of a single polynucleotide strand having both a sense and antisense region; in some preferred embodiments the polynucleotide strand has a hairpin structure.

[0029] In certain embodiments, the nucleic acid molecule (*e.g.*, siNA molecule) is a double-stranded nucleic acid (dsNA) molecule that is symmetrical with regard to overhangs, and has a blunt end on both ends. In other embodiments, the nucleic acid molecule (*e.g.*, siNA molecule) is a dsNA molecule that is symmetrical with regard to overhangs, and has an overhang on both ends of the dsNA molecule; preferably the molecule has overhangs of 1, 2, 3, 4, 5, 6, 7, or 8 nucleotides; preferably the molecule has 2 nucleotide overhangs. In some embodiments, the overhangs are 5' overhangs; in alternative embodiments the overhangs are 3' overhangs. In certain embodiments, the overhang nucleotides are modified with modifications as disclosed herein. In some embodiments, the overhang nucleotides are 2'-deoxynucleotides.

[0030] In certain preferred embodiments, the nucleic acid molecule (*e.g.*, siNA molecule) as a component of a pharmaceutical formulation is a dsNA molecule that is asymmetrical with regard to overhangs, and has a blunt end on one end of the molecule and an overhang on the other end of the molecule. In certain embodiments, the overhang is 1, 2, 3, 4, 5, 6, 7, or 8 nucleotides; preferably the overhang is 2 nucleotides. In some preferred embodiments, an asymmetrical dsNA molecule has a 3'-overhang (for example a two nucleotide 3'-overhang) on one side of a duplex occurring on the sense strand; and a blunt end on the other side of the molecule. In some preferred embodiments, an asymmetrical dsNA molecule has a 5'-overhang (for example a two nucleotide 5'-overhang) on one side of a duplex occurring on the sense strand; and a blunt end on the other side of the molecule. In other preferred embodiments, an asymmetrical dsNA molecule has a 3'-overhang (for example a two nucleotide 3'-overhang) on one side of a duplex occurring on the antisense strand; and a blunt end on the other side of the molecule. In some preferred embodiments, an asymmetrical dsNA molecule has a 5'-overhang (for example a two nucleotide 5'-overhang) on one side of a duplex occurring on wherein the antisense strand; and a blunt end on the other side of the molecule. In certain preferred embodiments, the overhangs are 2'-deoxynucleotides.

[0031] In some embodiments, the nucleic acid molecule (*e.g.*, siNA molecule) as a component of a pharmaceutical formulation has a hairpin structure (having the sense strand and antisense strand on one polynucleotide), with a loop structure on one end and a blunt end on the other end. In some embodiments, the nucleic acid molecule has a hairpin structure, with a loop structure on one end and an overhang end on the other end (for example a 1, 2, 3, 4, 5, 6, 7, or 8 nucleotide overhang); in certain embodiments, the overhang is a 3'-overhang; in certain embodiments

the overhang is a 5'-overhang; in certain embodiments the overhang is on the sense strand; in certain embodiments the overhang is on the antisense strand.

[0032] In some preferred embodiments, the nucleic acid molecule is selected from the nucleic acid molecules shown on Table 3.

[0033] The nucleic acid molecules (e.g., siNA molecule) disclosed herein as a component of a pharmaceutical formulation may include one or more modifications or modified nucleotides such as described herein. For example, a nucleic acid molecule (e.g., siNA molecule) as provided herein may include a modified nucleotide having a modified sugar; a modified nucleotide having a modified nucleobase; or a modified nucleotide having a modified phosphate group. Similarly, a nucleic acid molecule (e.g., siNA molecule) as provided herein may include a modified phosphodiester backbone and/or may include a modified terminal phosphate group.

[0034] Nucleic acid molecules (e.g., siNA molecules) as provided as a component of a pharmaceutical formulation may have one or more nucleotides that include a modified sugar moiety, for example as described herein. In some preferred embodiments, the modified sugar moiety is selected from the group consisting of 2'OMe, 2'-methoxyethoxy, 2'-deoxy, 2'-fluoro, 2'-allyl, 2'-O-(2-(methylamino)-2-oxoethyl), 4'-thio, 4'-(CH₂)₂-O-2'-bridge, 2'-LNA (the ribose moiety of an LNA nucleotide is modified with an extra bridge connecting the 2' oxygen and 4' carbon), and 2'-O-(N-methylcarbamate).

[0035] Nucleic acid molecules (e.g., siNA molecules) as provided as a component of a pharmaceutical formulation may have one or more modified nucleobase(s) for example as described herein, which preferably may be one selected from the group consisting of xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 5-halo uracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, amino, thiol, thioalkyl, hydroxyl and other 8-substituted adenines and guanines, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine, and acyclonucleotides.

[0036] Nucleic acid molecules (e.g., siNA molecules) as provided as a component of a pharmaceutical formulation may have one or more modifications to the phosphodiester backbone, for example as described herein. In some preferred embodiments, the phosphodiester bond is modified by substituting the phosphodiester bond with a phosphorothioate, 3'-(or 5'-)deoxy-3'-(or 5'-)thio-phosphorothioate, phosphorodithioate, phosphoroselenates, 3'-(or 5'-)deoxy phosphinates, borano phosphates, 3'-(or 5'-)deoxy-3'-(or 5'-)amino phosphoramidates, hydrogen phosphonates, borano phosphate esters, phosphoramidates, alkyl or aryl phosphonates and phosphotriester or phosphorus linkages.

[0037] In various embodiments, the provided nucleic acid molecules (e.g., siNA molecules) as a component of a pharmaceutical formulation may include one or modifications in the sense strand but not the antisense strand. In some embodiments the provided nucleic acid molecules (e.g., siNA molecules) include one or more modifications in the antisense strand but not the sense strand. In some embodiments the provided nucleic acid molecules (e.g., siNA molecules) include one or more modifications in the both the sense strand and the antisense strand.

[0038] In some embodiments in which the provided nucleic acid molecules (e.g., siNA molecules) as a component of a pharmaceutical formulation have modifications, the sense strand includes a pattern of alternating modified and unmodified nucleotides, and/or the antisense strand includes a pattern of alternating modified and unmodified nucleotides; in some preferred versions of such embodiments, the modification is a 2'OMe moiety. The pattern of alternating modified and unmodified nucleotides may start with a modified nucleotide at the 5'-end or 3'-end of one of the strands; for example the pattern of alternating modified and unmodified nucleotides may start with a modified nucleotide at the 5'-end or 3'-end of the sense strand and/or the pattern of alternating modified and unmodified nucleotides may start with a modified nucleotide at the 5'-end or 3'-end of the antisense strand. When both the antisense and sense strand include a pattern of alternating modified nucleotides, the pattern of modified nucleotides may be configured such that modified nucleotides in the sense strand are opposite modified nucleotides in the antisense strand; or there may be a phase shift in the pattern such that modified nucleotides of the sense strand are opposite unmodified nucleotides in the antisense strand and vice-versa.

[0039] The nucleic acid molecules (e.g., siNA molecules) as provided herein as a component of a pharmaceutical formulation may include one to three (i.e., 1, 2 or 3) deoxynucleotides at the 3'-end of the sense and/or antisense strand.

[0040] The nucleic acid molecules (e.g., siNA molecules) as provided herein as a component of a pharmaceutical formulation may include a phosphate group at the 5'-end of the sense and/or antisense strand.

[0041] In one aspect, provided as a component of a pharmaceutical formulation are double-stranded nucleic acid molecules having the structure (A1):

(A1)	5'	(N) _x -Z	3' (antisense strand)
	3'	Z'-(N') _y -z"	5' (sense strand)

wherein each of N and N' is a nucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein each of Z and Z' is independently present or absent, but if present independently includes 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present;

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of (N')_y;

wherein each of x and y is independently an integer between 18 and 40; wherein the sequence of (N')_y has complementary to the sequence of (N)_x; and wherein (N)_x includes an antisense sequence to SEQ ID NO:1.

[0042] In some embodiments, $(N)_x$ includes an antisense oligonucleotide present in Table 4. In other embodiments, $(N)_x$ is selected from an antisense oligonucleotide present in Tables B or C.

[0043] In some embodiments, the covalent bond joining each consecutive N or N' is a phosphodiester bond.

[0044] In some embodiments, $x = y$ and each of x and y is 19, 20, 21, 22 or 23. In various embodiments, $x = y = 19$.


[0045] In some embodiments of nucleic acid molecules (e.g., siRNA molecules) as disclosed herein, the double-stranded nucleic acid molecule is a siRNA, siNA or a miRNA.

[0046] In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in

- SERPINH1_4 (SEQ ID NOS:195 and 220),
- SERPINH1_12 (SEQ ID NOS:196 and 221),
- SERPINH1_30 (SEQ ID NOS:199 and 224), and
- SERPINH1_58 (SEQ ID NOS:208 and 233).

[0047] In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_4 (SEQ ID NOS:195 and 220). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_12 (SEQ ID NOS:196 and 221). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_30 (SEQ ID NOS:199 and 224). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_58 (SEQ ID NOS:208 and 233).

[0048] In some embodiments, the double-stranded nucleic acid molecules as a component of a pharmaceutical formulation comprise a DNA moiety or a mismatch to the target at position 1 of the antisense strand (5'-terminus). Such a structure is described herein. According to one embodiment provided are modified nucleic acid molecules having a structure (A2) set forth below:



wherein each of N^2 , N and N' is an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of $(N)_x$ and $(N')_y$ is an oligonucleotide in which each consecutive N or N' is joined to the adjacent N or N' by a covalent bond;

wherein each of x and y is independently an integer between 17 and 39;

wherein the sequence of $(N')_y$ has complementarity to the sequence of $(N)_x$ and $(N)_x$ has complementarity to a consecutive sequence in a target RNA;

wherein N^1 is covalently bound to $(N)_x$ and is mismatched to the target RNA or is a complementary DNA moiety to the target RNA;

wherein N^1 is a moiety selected from the group consisting of natural or modified uridine, deoxyribouridine, ribothymidine, deoxyribothymidine, adenosine or deoxyadenosine;

wherein z'' may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of $N^2-(N')_y$; and

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides, consecutive non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present.

[0049] In some embodiments, the sequence of $(N')_y$ is fully complementary to the sequence of $(N)_x$. In various embodiments, sequence of $N^2-(N')_y$ is complementary to the sequence of $N^1-(N)_x$. In some embodiments, $(N)_x$ comprises an antisense that is fully complementary to about 17 to about 39 consecutive nucleotides in a target RNA. In other embodiments, $(N)_x$ comprises an antisense that is substantially complementary to about 17 to about 39 consecutive nucleotides in a target RNA.

[0050] In some embodiments, N^1 and N^2 form a Watson-Crick base pair. In some embodiments, N^1 and N^2 form a non-Watson-Crick base pair. In some embodiments, a base pair is formed between a ribonucleotide and a deoxyribonucleotide.

[0051] In some embodiments, $x = y = 18$, $x = y = 19$ or $x = y = 20$. In preferred embodiments, $x = y = 18$. When $x = 18$ in $N^1-(N)_x$, N^1 refers to position 1 and positions 2-19 are included in $(N)_{18}$. When $y = 18$ in $N^2-(N')_y$, N^2 refers to position 19 and positions 1-18 are included in $(N')_{18}$.

[0052] In some embodiments, N^1 is covalently bound to $(N)_x$ and is mismatched to the target RNA. In various embodiments, N^1 is covalently bound to $(N)_x$ and is a DNA moiety complementary to the target RNA.

[0053] In some embodiments, a uridine in position 1 of the antisense strand is substituted with an N^1 selected from adenosine, deoxyadenosine, deoxyuridine (dU), ribothymidine or deoxythymidine. In various embodiments, N^1 selected from adenosine, deoxyadenosine or deoxyuridine.

[0054] In some embodiments, guanosine in position 1 of the antisense strand is substituted with an N^1 selected from adenosine, deoxyadenosine, uridine, deoxyuridine, ribothymidine or deoxythymidine. In various embodiments, N^1 is selected from adenosine, deoxyadenosine, uridine or deoxyuridine.

[0055] In some embodiments, cytidine in position 1 of the antisense strand is substituted with an N^1 selected from adenosine, deoxyadenosine, uridine, deoxyuridine, ribothymidine or deoxythymidine. In various embodiments, N^1 is selected from adenosine, deoxyadenosine, uridine or deoxyuridine.

[0056] In some embodiments, adenosine in position 1 of the antisense strand is substituted with an N^1 selected from deoxyadenosine, deoxyuridine, ribothymidine or deoxythymidine. In various embodiments, N^1 selected from deoxyadenosine or deoxyuridine.

[0057] In some embodiments, N^1 and N^2 form a base pair between uridine or deoxyuridine, and adenosine or deoxyadenosine. In other embodiments, N^1 and N^2 form a base pair between deoxyuridine and adenosine.

[0058] In some embodiments, the double-stranded nucleic acid molecule as a component of a pharmaceutical formulation is a siRNA, siNA or a miRNA. The double-stranded nucleic acid molecules as provided herein are also referred to as "duplexes".

[0059] In some embodiments $(N)_x$ includes an antisense oligonucleotide present in Table 5. In some embodiments, $x=y=18$ and $N'-(N)_x$ includes an antisense oligonucleotide present in Table 4. In some embodiments $x=y=19$ or $x=y=20$. In certain preferred embodiments, $x=y=18$. In some embodiments $x=y=18$ and the sequences of $N^1-(N)_x$ and $N^2-(N')_y$ are selected from the pair of oligonucleotides set forth in Table 4. In some embodiments, $x=y=18$ and the sequences of $N^1-(N)_x$ and $N^2-(N')_y$ are selected from the pair of oligonucleotides set forth in Tables D and E. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in

- SERPINH1_2 (SEQ ID NOS: 60 and 127),
- SERPINH1_6 (SEQ ID NOS: 63 and 130),
- SERPINH1_11 (SEQ ID NOS: 68 and 135),
- SERPINH1_13 (SEQ ID NOS: 69 and 136),
- SERPINH1_45 (SEQ ID NOS: 97 and 164),
- SERPINH1_45a (SEQ ID NOS: 98 and 165),
- SERPINH1_51 (SEQ ID NOS: 101 and 168),
- SERPINH1_51a (SEQ ID NOS: 105 and 172),
- SERPINH1_52 (SEQ ID NOS: 102 and 169), and
- SERPINH1_86 (SEQ ID NOS: 123 and 190).

[0060] In some preferred embodiments, the antisense and sense strands are selected from the sequence pairs set forth in

- SERPINH1_2 (SEQ ID NOS: 60 and 127),
- SERPINH1_6 (SEQ ID NOS: 63 and 130),
- SERPINH1_45a (SEQ ID NOS: 98 and 165),
- SERPINH1_51 (SEQ ID NOS: 101 and 168), and
- SERPINH1_51a (SEQ ID NOS: 105 and 172).

[0061] In some preferred embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2 (SEQ ID NOS: 60 and 127). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_6 (SEQ ID NOS: 63 and 130). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_11 (SEQ ID NOS: 68 and 135). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_13 (SEQ ID NOS: 69 and 136). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_45 (SEQ ID NOS: 97 and 164). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_45a (SEQ ID NOS: 98 and 165). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_51 (SEQ ID NOS: 101 and 168). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_51a (SEQ ID NOS: 105 and 172). In some embodiments, the antisense and sense strands are the sequence pairs set forth in SERPINH1_52 (SEQ ID NOS: 102 and 169). In some embodiments the antisense and sense strands are the sequence pairs set forth in (SEQ ID NOS: 123 and 190). In some preferred embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2 (SEQ ID NOS: 60 and 127), SERPINH1_6 (SEQ ID NOS: 63 and 130), SERPINH1_45a (SEQ ID NOS: 98 and 165), SERPINH1_51 (SEQ ID NOS: 101 and 168), and SERPINH1_51a (SEQ ID NOS: 105 and 172).

[0062] In some embodiments, N^1 and N^2 form a Watson-Crick base pair. In other embodiments, N^1 and N^2 form a non-Watson-Crick base pair. In some embodiments N^1 is a modified riboadenosine or a modified ribouridine.

[0063] In some embodiments, N¹ and N² form a Watson-Crick base pair. In other embodiments, N¹ and N² form a non-Watson-Crick base pair. In certain embodiments, N¹ is selected from the group consisting of riboadenosine, modified riboadenosine, deoxyriboadenosine, modified deoxyriboadenosine. In other embodiments, N¹ is selected from the group consisting of ribouridine, deoxyribouridine, modified ribouridine, and modified deoxyribouridine.

[0064] In certain embodiments, position 1 in the antisense strand (5'-terminus) includes deoxyribouridine (dU) or adenosine. In some embodiments, N¹ is selected from the group consisting of riboadenosine, modified riboadenosine, deoxyriboadenosine, modified deoxyriboadenosine and N² is selected from the group consisting of ribouridine, deoxyribouridine, modified ribouridine, and modified deoxyribouridine. In certain embodiments, N¹ is selected from the group consisting of riboadenosine and modified riboadenosine and N² is selected from the group consisting of ribouridine and modified ribouridine.

[0065] In certain embodiments, N¹ is selected from the group consisting of ribouridine, deoxyribouridine, modified ribouridine, and modified deoxyribouridine and N² is selected from the group consisting of riboadenosine, modified riboadenosine, deoxyriboadenosine, and modified deoxyriboadenosine. In certain embodiments, N¹ is selected from the group consisting of ribouridine and deoxyribouridine and N² is selected from the group consisting of riboadenosine and modified riboadenosine. In certain embodiments, N¹ is ribouridine and N² is riboadenosine. In certain embodiments, N¹ is deoxyribouridine and N² is riboadenosine.

[0066] In some embodiments of Structure (A2), N¹ includes 2'OMe modified ribouracil or 2'OMe modified riboadenosine. In certain embodiments of structure (A2), N² includes a 2'OMe modified ribonucleotide or deoxyribonucleotide.

[0067] In some embodiments of Structure (A2), N¹ includes 2'OMe modified ribouracil or 2'OMe modified ribocytosine. In certain embodiments of structure (A2), N² includes a 2'OMe modified ribonucleotide.

[0068] In some embodiments each of N and N' is an unmodified nucleotide. In some embodiments, at least one of N or N' includes a chemically modified nucleotide or an unconventional moiety. In some embodiments, the unconventional moiety is selected from a mirror nucleotide, an abasic ribose moiety and an abasic deoxyribose moiety. In some embodiments, the unconventional moiety is a mirror nucleotide, preferably an L-DNA moiety. In some embodiments, at least one of N or N' includes a 2'OMe modified ribonucleotide.

[0069] In some embodiments, the sequence of (N')_y is fully complementary to the sequence of (N)_x. In other embodiments, the sequence of (N')_y is substantially complementary to the sequence of (N)_x.

[0070] In some embodiments, (N)_x includes an antisense sequence that is fully complementary to about 17 to about 39 consecutive nucleotides in a target mRNA. In other embodiments, (N)_x includes an antisense that is substantially complementary to about 17 to about 39 consecutive nucleotides in a target mRNA.

[0071] In some embodiments of Structure A1 and Structure A2 the compound is blunt ended, for example wherein both Z and Z' are absent. In an alternative embodiment, at least one of Z or Z' is present. Z and Z' independently include one or more covalently linked modified and or unmodified nucleotides, including deoxyribonucleotides and ribonucleotides, or an unconventional moiety for example inverted abasic deoxyribose moiety or abasic ribose moiety; a non-nucleotide C3, C4 or C5 moiety, an amino-6 moiety, a mirror nucleotide, and the like. In some embodiments each of Z and Z' independently includes a C3 moiety or an amino-C6 moiety. In some embodiments Z' is absent and Z is present and includes a non-nucleotide C3 moiety. In some embodiments Z is absent and Z' is present and includes a non-nucleotide C3 moiety.

[0072] In some embodiments of Structure A1 and Structure A2, each N consists of an unmodified ribonucleotide. In some embodiments of Structure A1 and Structure A2, each N' consists of an unmodified nucleotide. In preferred embodiments, at least one of N and N' is a modified ribonucleotide or an unconventional moiety.

[0073] In other embodiments, the compound of Structure A1 or Structure A2 includes at least one ribonucleotide modified in the sugar residue. In some embodiments, the compound includes a modification at the 2' position of the sugar residue. In some embodiments, the modification in the 2' position includes the presence of an amino, a fluoro, an alkoxy or an alkyl moiety. In certain embodiments, the 2' modification includes an alkoxy moiety. In preferred embodiments, the alkoxy moiety is a methoxy moiety (2'OMe). In some embodiments, the nucleic acid compound includes 2'OMe modified alternating ribonucleotides in one or both of the antisense and the sense strands. In other embodiments, the compound includes 2'OMe modified ribonucleotides in the antisense strand, (N)_x or N¹-(N)_x, only. In certain embodiments, the middle ribonucleotide of the antisense strand; e.g. ribonucleotide in position 10 in a 19-mer strand is unmodified. In various embodiments, the nucleic acid compound includes at least 5 alternating 2'OMe modified and unmodified ribonucleotides. In additional embodiments, the compound of Structure A1 or Structure A2 includes modified ribonucleotides in alternating positions wherein each ribonucleotide at the 5' and 3'-termini of (N)_x or N¹-(N)_x are modified in their sugar residues, and each ribonucleotide at the 5' and 3'-termini of (N')_y or N²-(N')_y are unmodified in their sugar residues.

[0074] In some embodiments, the double-stranded molecule as a component of a pharmaceutical formulation includes one or more of the following modifications

- N in at least one of positions 5, 6, 7, 8, or 9 of the antisense strand is selected from a 2'-5'-nucleotide or a mirror nucleotide;
- N' in at least one of positions 9 or 10 of the sense strand is selected from a 2'-5'-nucleotide and a pseudo-uridine; and
- N' in 4, 5, or 6 consecutive positions at the 3'-terminus positions of (N')_y comprises a 2'-5'-nucleotide.

[0075] In some embodiments, the double-stranded molecule includes a combination of the following modifications

- the antisense strand includes a 2'-5'-nucleotide or a mirror nucleotide in at least one of positions 5, 6, 7, 8, or 9; and
- the sense strand includes at least one of a 2'-5'-nucleotide and a pseudo-uridine in positions 9 or 10.

[0076] In some embodiments, the double-stranded molecule includes a combination of the following modifications

- the antisense strand includes a 2'-5'-nucleotide or a mirror nucleotide in at least one of positions 5, 6, 7, 8, or 9; and
- the sense strand includes 4, 5, or 6 consecutive 2'-5'-nucleotides at the 3'-penultimate or 3'-terminal positions.

[0077] In some embodiments, the sense strand $((N)_x$ or $N^1-(N)_x$) includes 1, 2, 3, 4, 5, 6, 7, 8, or 9 2'OMe modified ribonucleotides. In some embodiments, the antisense strand includes 2'OMe modified ribonucleotides at positions 2, 4, 6, 8, 11, 13, 15, 17 and 19. In other embodiments, antisense strand includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 7, 9, 11, 13, 15, 17 and 19. In other embodiments, the antisense strand includes 2'OMe modified ribonucleotides at positions 3, 5, 7, 9, 11, 13, 15, 17 and 19. In some embodiments the antisense strand includes one or more 2'OMe modified pyrimidines. In some embodiments, all the pyrimidine nucleotides in the antisense strand are 2'OMe modified. In some embodiments, the sense strand includes 2'OMe modified pyrimidines.

[0078] In some embodiments of Structure A1 and Structure A2, neither the sense strand nor the antisense strand is phosphorylated at the 3'- and 5'-termini. In other embodiments, one or both of the sense strand or the antisense strand are phosphorylated at the 3'-termini.

[0079] In some embodiments of Structure A1 and Structure A2, $(N)_y$ includes at least one unconventional moiety selected from a mirror nucleotide, a 2'-5'-nucleotide and a TNA. In some embodiments, the unconventional moiety is a mirror nucleotide. In various embodiments, the mirror nucleotide is selected from an L-ribonucleotide (L-RNA) and an L-deoxyribonucleotide (L-DNA). In preferred embodiments, the mirror nucleotide is L-DNA. In certain embodiments, the sense strand comprises an unconventional moiety in position 9 or 10 (from the 5'-terminus). In preferred embodiments, the sense strand includes an unconventional moiety in position 9 (from the 5'-terminus). In some embodiments, the sense strand is 19 nucleotides in length and comprises 4, 5, or 6 consecutive unconventional moieties in positions 15, (from the 5'-terminus). In some embodiments, the sense strand includes 4 consecutive 2'-5'-ribonucleotides in positions 15, 16, 17, and 18. In some embodiments, the sense strand includes 5 consecutive 2'-5'-ribonucleotides in positions 15, 16, 17, 18 and 19. In various embodiments, the sense strand further comprises Z'. In some embodiments, Z' includes a C3OH moiety or a C3Pi moiety.

[0080] In some embodiments, of Structure A1 $(N')_y$ includes at least one L-DNA moiety. In some embodiments, $x=y=19$ and $(N')_y$ consists of unmodified ribonucleotides at positions 1-17 and 19 and one L-DNA at the 3' penultimate position (position 18). In other embodiments, $x=y=19$ and $(N')_y$ consists of unmodified ribonucleotides at positions 1-16 and 19 and two consecutive L-DNA at the 3' penultimate position (positions 17 and 18). In various embodiments, the unconventional moiety is a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate linkage. According to various embodiments, $(N')_y$ includes 2, 3, 4, 5, or 6 consecutive ribonucleotides at the 3'-terminus linked by 2'-5' internucleotide linkages. In one embodiment, four consecutive nucleotides at the 3'-terminus of $(N')_y$ are joined by three 2'-5' phosphodiester bonds, wherein one or more of the 2'-5' nucleotides which form the 2'-5' phosphodiester bonds further includes a 3'-O-methyl (3'OMe) sugar modification. Preferably the 3'-terminal nucleotide of $(N')_y$ includes a 2'OMe modification. In certain embodiments, $x=y=19$ and $(N')_y$ includes two or more consecutive nucleotides at positions 15, 16, 17, 18 and 19 include a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide bond (2'-5' nucleotide). In various embodiments the nucleotide forming the 2'-5' internucleotide bond includes a 3' deoxyribose nucleotide or a 3' methoxy nucleotide (3' H or 3'OMe in place of a 3' OH). In some embodiments $x=y=19$ and $(N')_y$ includes 2'-5' nucleotides at positions 15, 16 and 17 such that adjacent nucleotides are linked by a 2'-5' internucleotide bond between positions 15-16, 16-17 and 17-18; or at positions, 15, 16, 17, 18, and 19 such that adjacent nucleotides are linked by a 2'-5' internucleotide bond between positions 15-16, 16-17, 17-18 and 18-19 and a 3'OH is available at the 3'-terminal nucleotide or at positions 16, 17 and 18 such that adjacent nucleotides are linked by a 2'-5' internucleotide bond between positions 16-17, 17-18 and 18-19. In some embodiments $x=y=19$ and $(N')_y$ includes 2'-5'-nucleotides at positions 16 and 17 or at positions 17 and 18 or at positions 15 and 17 such that adjacent nucleotides are linked by a 2'-5' internucleotide bond between positions 16-17 and 17-18 or between positions 17-18 and 18-19 or between positions 15-16 and 17-18, respectively. In other embodiments, the pyrimidine ribonucleotides (rU, rC) in $(N')_y$ are substituted with nucleotides joined to the adjacent nucleotide by a 2'-5' internucleotide bond. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 or SERPINH1_88, and $x=y=19$ and $(N')_y$ comprises five consecutive nucleotides at the 3'-terminus joined by four 2'-5' linkages, specifically the linkages between the nucleotides position 15-16, 16-17, 17-18 and 18-19.

[0081] In some embodiments, the linkages include phosphodiester bonds. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 or SERPINH1_88 and $x=y=19$ and $(N')_y$ comprises five consecutive nucleotides at the 3'-terminus joined by four 2'-5' linkages and optionally further includes Z' and Z' independently selected from an inverted abasic moiety and a C3 alkyl [C3; 1,3-propanediol mono(dihydrogen phosphate)] cap. The C3 alkyl cap is covalently linked to the 3' or 5' terminal nucleotide. In some embodiments, the 3' C3 terminal cap further comprises a 3' phosphate. In some embodiments, the 3' C3 terminal cap further comprises a 3'-terminal hydroxy group.

[0082] In some embodiments, the antisense and sense strands as components of a pharmaceutical formulation are selected from the sequence pairs set forth in SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 or SERPINH1_88 and $x=y=19$ and $(N')_y$ includes an

L-DNA position 18; and (N')_y optionally further includes Z' and z' independently selected from an inverted abasic moiety and a C3 alkyl (C3; 1,3-propanediol mono(dihydrogen phosphate)) cap.

[0083] In some embodiments, (N')_y includes a 3'-terminal phosphate. In some embodiments, (N')_y includes a 3'-terminal hydroxyl.

[0084] In some embodiments, the antisense and sense strands as components of a pharmaceutical formulation are selected from the sequence pairs set forth in SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 or SERPINH1_88 and x=y=19 and (N)_x includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 7, 9, 11, 13, 15, 17, 19 or at positions 2, 4, 6, 8, 11, 13, 15, 17, 19. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 and SERPINH1_88 and x=y=19 and (N)_x includes 2'OMe modified pyrimidines. In some embodiments, all pyrimidines in (N)_x include the 2'OMe modification.

[0085] In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18 and N² is a riboadenosine moiety.

[0086] In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18, and N²-(N')_y includes five consecutive nucleotides at the 3'-terminus joined by four 2'-5' linkages, specifically the linkages between the nucleotides position 15-16, 16-17, 17-18 and 18-19. In some embodiments the linkages include phosphodiester bonds.

[0087] In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18 and N²-(N')_y includes five consecutive nucleotides at the 3'-terminus joined by four 2'-5' linkages and optionally further includes Z' and z' independently selected from an inverted abasic moiety and a C3 alkyl [C3; 1,3-propanediol mono(dihydrogen phosphate)] cap.

[0088] In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18 and N²-(N')_y includes an L-DNA position 18; and (N')_y optionally further includes Z' and z' independently selected from an inverted abasic moiety and a C3 alkyl [C3; 1,3-propanediol mono(dihydrogen phosphate)] cap.

[0089] In some embodiments, N²-(N')_y comprises a 3'-terminal phosphate. In some embodiments, N²-(N')_y comprises a 3'-terminal hydroxyl.

[0090] In some embodiments, the antisense and sense strands as components of a pharmaceutical formulation are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18 and N¹-(N)_x includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 11, 13, 15, 17, 19 or in positions 1, 3, 5, 9, 11, 13, 15, 17, 19, or in positions 3, 5, 9, 11, 13, 15, 17, or in positions 2, 4, 6, 8, 11, 13, 15, 17, 19. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_52 or SERPINH1_86 and x=y=18 and N¹-(N)_x includes 2'OMe modified ribonucleotides at positions 11, 13, 15, 17 and 19. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18 and N¹-(N)_x includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 11, 13, 15, 17, 19 or in positions 3, 5, 7, 9, 11, 13, 15, 17, 19. In some embodiments, the antisense and sense strands are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_52 or SERPINH1_86 and x=y=18 and N¹-(N)_x includes 2'OMe modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17, 19.

[0091] In some embodiments, the antisense and sense strands as components of a pharmaceutical formulation are selected from the sequence pairs set forth in SERPINH1_2, SERPINH1_6, SERPINH1_11, SERPINH1_13, SERPINH1_45, SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 or SERPINH1_86 and x=y=18 and N¹-(N)_x includes 2'OMe modified pyrimidines. In some embodiments, all pyrimidines in (N)_x include the 2'OMe modification. In some embodiments, the antisense strand further includes an L-DNA or a 2'-5' nucleotide in position 5, 6 or 7. In other embodiments, the antisense strand further includes a ribonucleotide which generates a 2'-5' internucleotide linkage in between the ribonucleotides in positions 5-6 or 6-7.


[0092] In additional embodiments, N¹-(N)_x further includes Z wherein Z includes a non-nucleotide overhang. In some embodiments the non-nucleotide overhang is C3 -C3 [1,3-propanediol mono(dihydrogen phosphate)]₂.

[0093] In some embodiments, of Structure A2, (N)_y includes at least one L-DNA moiety. In some embodiments x=y=18 and (N')_y consists of unmodified ribonucleotides at positions 1-16 and 18 and one L-DNA at the 3' penultimate position (position 17). In other embodiments, x=y=18 and (N')_y consists of unmodified ribonucleotides at position 1-15 and 18 and two consecutive L-DNA at the 3' penultimate position (positions 16 and 17). In various embodiments, the unconventional moiety is a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate linkage. According to various embodiments, (N')_y includes 2, 3, 4, 5, or 6 consecutive ribonucleotides at the 3'-terminus linked by 2'-5' internucleotide linkages. In one embodiment, four consecutive nucleotides at the 3'-terminus of (N')_y are joined by three 2'-5' phosphodiester bonds, wherein one or more of the 2'-5' nucleotides which form the 2'-5' phosphodiester bonds further includes a 3'-O-methyl (3'OMe) sugar

modification. Preferably, the 3'-terminal nucleotide of (N')_y includes a 2'OMe modification. In certain embodiments, x=y=18 and in (N')_y two or more consecutive nucleotides at positions 14, 15, 16, 17, and 18 include a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide bond. In various embodiments, wherein the nucleotide forming the 2'-5' internucleotide bond includes a 3' deoxyribose nucleotide or a 3' methoxy nucleotide. In some embodiments, x=y=18 and (N')_y includes nucleotides joined to the adjacent nucleotide by a 2'-5' internucleotide bond between positions 15-16, 16-17 and 17-18 or between positions 16-17 and 17-18. In some embodiments, x=y=18 and (N')_y includes nucleotides joined to the adjacent nucleotide by a 2'-5' internucleotide bond between positions 14-15, 15-16, 16-17, and 17-18 or between positions 15-16, 16-17, and 17-18 or between positions 16-17 and 17-18 or between positions 17-18 or between positions 15-16 and 17-18. In other embodiments, the pyrimidine ribonucleotides (rU, rC) in (N')_y are substituted with nucleotides joined to the adjacent nucleotide by a 2'-5' internucleotide bond.

[0094] In some embodiments, the antisense and sense strands are selected from the oligonucleotide pairs set forth in Table 5 and identified herein as SERPINH1_2 (SEQ ID NOS: 60 and 127), SERPINH1_6 (SEQ ID NOS: 63 and 130), SERPINH1_45a (SEQ ID NOS: 98 and 165), SERPINH1_51 (SEQ ID NOS: 101 and 168), and SERPINH1_51a (SEQ ID NOS: 105 and 172).

[0095] In some embodiments, the double-stranded nucleic acid molecule as a component of a pharmaceutical formulation includes the antisense strand set forth in SEQ ID NO:127 and sense strand set forth in SEQ ID NO:60; identified herein as SERPINH1_2. In some embodiments the double-stranded nucleic acid molecule has the structure



wherein each "|" represents base pairing between the ribonucleotides; wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of N²-(N')_y.

[0096] In some embodiments, provided is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes one or more 2'OMe modified pyrimidines and/or purines, a 2'-5'-ribonucleotide in position 5, 6, 7 or 8, and a 3'-terminal nucleotide or non-nucleotide overhang. In some embodiments, the sense strand (SEQ ID NO:60) includes 4 or 5 consecutive 2'-5'-nucleotides at the 3'-terminal or penultimate positions, a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus and a cap moiety covalently attached at the 5'-terminus. In other embodiments, the sense strand (SEQ ID NO:60) includes one or more 2'OMe pyrimidine, a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus and a cap moiety covalently attached at the 5'-terminus.

[0097] In some embodiments, provided is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 15, 17 and 19; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:60) is selected from a sense strand which includes

- 2'-5'-ribonucleotides at positions 15, 16, 17, 18 and 19; a C3OH 3'-terminal non-nucleotide overhang; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
- 2'-5'-ribonucleotides at positions 15, 16, 17, 18 and 19; a 3'-terminal phosphate; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
- 2'OMe modified ribonucleotides at positions 5, 7, 13, and 16; a 2'-5'-ribonucleotide at position 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
- 2'OMe modified ribonucleotides at positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide at position 9; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
- 2'-5'-ribonucleotides at positions 15, 16, 17, 18, and 19; a C3-Pi moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0098] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:60) includes 2'-5'-ribonucleotides at positions 15, 16, 17, 18, and 19; a C3 3'-terminal overhang; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0099] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH 3'-terminal overhang; and the sense strand (SEQ ID NO:60) includes 2'-5'-ribonucleotides at positions 15, 16, 17, 18, and 19; a 3'-terminal phosphate; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0100] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide at position 7 and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:60) includes 2'OMe modified ribonucleotides at positions 5, 7, 13, and 16; a 2'-5'-ribonucleotide at position 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0101] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide at position 7 and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:60) includes 2'OMe modified ribonucleotides at positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide at position 9; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0102] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:60) includes 2'-5'-ribonucleotides at positions 15, 16, 17, 18, and 19; a C3-Pi moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0103] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:127) includes 2'OMe modified ribonucleotides at positions 1, 3, 5, 9, 11, 13, 15, 17, 19; and a C3-C3 3'-terminal overhang; and the sense strand (SEQ ID NO:60) includes 2'OMe modified ribonucleotides at positions 7, 9, 13, 16 and 18; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0104] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:60) includes 2'-5'-ribonucleotides at positions 15, 16, 17, 18, and 19; a 3'-terminal phosphate; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:127) includes an antisense strand selected from one of

- 2'OMe modified ribonucleotides at positions 1, 3, 5, 7, 9, 11, 13, 15, 17, 19; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
- 2'OMe modified ribonucleotides at positions 1, 3, 6, 8, 10, 12, 14, 17, 18; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0105] In some embodiments provided herein is a double-stranded nucleic acid molecule as a component of a pharmaceutical formulation which includes the antisense strand set forth in SEQ ID NO:130 and the sense strand set forth in SEQ ID NO:63; identified herein as SERPINH1_6. In some embodiments the duplex comprises the structure

```

5'      UACUCGUCUCGCAUCUUGU  ~Z  3'      (antisense SEQ ID NO:130)
|||||
3'  Z' ~AUGAGCAGAGCGUAGAACA  ~z"  5'      (sense SEQ ID NO:63)

```

wherein each "|" represents base pairing between the ribonucleotides; wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of N²-(N')_y.

[0106] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:63) includes one or more 2'OMe modified pyrimidines; a 3'-terminal nucleotide or non-nucleotide overhang; and cap moiety covalently attached at the 5'-terminus. In some embodiments, the antisense strand (SEQ ID NO:130) includes one or more 2'OMe modified pyrimidine; a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus.

[0107] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:63) includes 2'OMe modified ribonucleotides at positions 2, 14 and 18; a C3OH or C3Pi moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:130) is selected from an antisense strand that includes

- 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
- 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 12, 13 and 17; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
- 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
- 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 13, 15 and 17; a dU in position 1; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0108] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:63) includes 2'OMe modified ribonucleotides in positions 2, 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:130) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0109] In some embodiments provided herein is a duplex oligonucleotide molecule wherein the sense strand (SEQ ID NO:63) includes 2'OMe

modified ribonucleotides in positions 14 and 18 and optionally in position 2; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:130) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 12, 13, and 17; a 2'-5'-ribonucleotide at position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0110] In some embodiments provided herein is a duplex oligonucleotide molecule wherein the sense strand (SEQ ID NO:63) includes 2'OMe modified ribonucleotides in positions 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:130) is selected from an antisense strand which includes

- 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3Pi or C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
- 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 12, 13, and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3Pi or C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0111] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:63) includes 2'OMe modified ribonucleotides in positions 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:130) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0112] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:63) includes 2'OMe modified ribonucleotides in positions 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:130) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 12, 13, and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH 3'-terminal overhang.

[0113] In some embodiments, the duplex is a component of a pharmaceutical formulation includes the antisense strand set forth in SEQ ID NO:165 and sense strand set forth in SEQ ID NO:98; identified herein as SERPINH1_45a. In some embodiments, the duplex comprises the structure

```

5'   AGGAAGUUGAUUCCUGGAGU   -Z 3'   (antisense SEQ ID NO:165)
      |||||
3'   Z'-UCCUUCACUAGAACCUCU   -Z" 5'   (sense SEQ ID NO:98)

```

wherein each "|" represents base pairing between the ribonucleotides;

wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of N²-(N)_y

[0114] In some embodiments, the sense strand (SEQ ID NO:98) includes 2'-5'-ribonucleotides in positions 15, 16, 17, and 18 or 15, 16, 17, 18, and 19; a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus. In some embodiments the antisense strand (SEQ ID NO:165) includes 2'OMe modified pyrimidine and/or purines; a 2'-5' nucleotide in position 5, 6, 7, or 8; and a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus.

[0115] In some embodiments, the sense strand (SEQ ID NO:98) includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3Pi or C3-OH 3'-terminal non-nucleotide moiety; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:165) includes an antisense strand selected from one of

- 2'OMe modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3Pi or C3Pi-C3OH 3'-terminal overhang; or
- 2'OMe modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17 and 19; and a C3Pi-C3Pi or C3Pi-C3OH 3'-terminal overhang;
- 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3Pi or C3Pi-C3OH 3'-terminal overhang; or
- 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 11, 13, 15, 17 and 19; and a C3Pi-C3Pi or C3Pi-C3OH 3'-terminal overhang.

[0116] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:98) includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3-OH 3'-terminal moiety; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:165) includes 2'OMe modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-COH 3'-terminal overhang.

[0117] In some embodiments, the double-stranded nucleic acid molecule is a component of a pharmaceutical formulation includes the antisense strand set forth in SEQ ID NO:168 and sense strand set forth in SEQ ID NO:101; identified herein as SERPINH1_51. In some embodiments, the duplex comprises the structure

```

5'   UCACCCAUGUGUCACGGA   -Z 3'   (antisense SEQ ID NO:168)
      |||||

```

|||||
3' Z' -AGUGGGUACACAGAGUCCU -z" 5' (sense SEQ ID NO:101)

wherein each "|" represents base pairing between the ribonucleotides;

wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of N²-(N')_y.

[0118] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:101) includes 2'OMe modified pyrimidines, optionally a 2'-5'-ribonucleotide in position 9 or 10; a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus; and optionally a cap moiety covalently attached at the 5'-terminus. In some embodiments the antisense strand (SEQ ID NO:168) includes 2'OMe modified pyrimidine and or purines; a 2'-5' nucleotide in position 5, 6, 7, or 8; and a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus.

[0119] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:101) includes 2'OMe modified pyrimidines in positions 4, 11, 13, and 17; optionally a 2'-5'-ribonucleotide in position 9 or 10; a C3Pi or C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:168) is selected from an antisense strand which includes

1. a) 2'OMe modified ribonucleotides in positions 1, 8, and 15; a 2'-5'-ribonucleotide in position 6 or 7; and a C3Pi-C3OH overhang covalently attached at the 3'-terminus; or
2. b) 2'OMe modified ribonucleotides in positions 1, 4, 8, 13 and 15; a 2'-5'-ribonucleotide in position 6 or 7; and a C3Pi-C3OH overhang covalently attached at the 3'-terminus; or
3. c) 2'OMe modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH overhang covalently attached at the 3'-terminus; or
4. d) 2'OMe modified ribonucleotides in positions 1, 3, 8, 12, 13, and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0120] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:101) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; optionally a 2'-5'-ribonucleotide in position 9; a C3-OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:168) includes 2'OMe modified ribonucleotides in positions 1, 8, and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0121] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:101) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; optionally a 2'-5'-ribonucleotide in position 9; a C3-OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:168) includes 2'OMe modified ribonucleotides in positions 1, 4, 8, 13 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0122] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:101) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:168) includes 2'OMe modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0123] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:101) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:168) includes 2'OMe modified ribonucleotides in positions 1, 3, 8, 12, 13, and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0124] In some embodiments, the double-stranded nucleic acid molecule is a component of a pharmaceutical formulation includes the antisense strand set forth in SEQ ID NO:168 and sense strand set forth in SEQ ID NO:101; identified herein as SERPINH1_51a. In some embodiments the duplex comprises the structure

5' ACACCAUGUGUCACGA -Z 3' (antisense SEQ ID NO:172)
|||||
3' Z' -UGUGGGUACACAGAGUCCU -z" 5' (sense SEQ ID NO:105)

wherein each "|" represents base pairing between the ribonucleotides;

wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of N²-(N')_y.

[0125] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:105) includes 2'OMe modified pyrimidines; optionally a 2'-5'-ribonucleotide in position 9 or 10; a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus; and optionally a cap moiety covalently attached at the 5'-terminus. In some embodiments the antisense strand (SEQ ID NO:172) includes 2'OMe modified pyrimidine and or purines; a 2'-5' nucleotide in position 5, 6, 7, or 8; and a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus.

[0126] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:105) includes 2'OMe modified pyrimidines in positions 4, 11, 13, and 17; optionally a 2'-5'-ribonucleotide in position 9 or 10; a C3Pi or C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:172) is selected from an antisense strand which includes

1. a) 2'OMe modified ribonucleotides in positions 8, and 15; a 2'-5'-ribonucleotide in position 6 or 7; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus; or
2. b) 2'OMe modified ribonucleotides in positions 4, 8, 13 and 15; a 2'-5'-ribonucleotide in position 6 or 7; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus; or
3. c) 2'OMe modified ribonucleotides in positions 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus; or
4. d) 2'OMe modified ribonucleotides in positions 3, 8, 12, 13, and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0127] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:105) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; optionally a 2'-5'-ribonucleotide in position 9; a C3-OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:172) includes 2'OMe modified ribonucleotides in positions 8 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0128] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:105) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; optionally a 2'-5'-ribonucleotide in position 9; a C3-OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:172) includes 2'OMe modified ribonucleotides in positions 4, 8, 13 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0129] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:105) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3-OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:172) includes 2'OMe modified ribonucleotides in positions 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0130] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:105) includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:172) includes 2'OMe modified ribonucleotides in positions 3, 8, 12, 13, and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0131] In some embodiments the antisense and sense strands are selected from the oligonucleotide pairs set forth in Table 5 and identified herein as SERPINH1_4 (SEQ ID NOS: 195 and 220) and SERPINH1_12 (SEQ ID NOS: 196 and 221).

[0132] In some embodiments the double-stranded nucleic acid molecule is a component of a pharmaceutical formulation includes the antisense strand set forth in SEQ ID NO:220 and sense strand set forth in SEQ ID NO:195; identified herein as SERPINH1_4. In some embodiments the double-stranded nucleic acid molecule has the structure

```

5'   AAUAGCACCCAUGUGUCUC   -Z 3'   (antisense SEQ ID NO:220)
      |||||
3' z'-UUAUCGUGGGUACACAGAG - z" 5'   (sense SEQ ID NO:195)

```

wherein each "|" represents base pairing between the ribonucleotides;

wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of N²-(N')_y.

[0133] In some embodiments provided is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 15, 17 and 19, a 2'-5'-ribonucleotide in position 7, and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) is selected from a sense strand which includes

1. a) 2'-5'-ribonucleotides in positions 15, 16, 17, 18 and 19, a C3OH moiety covalently attached to the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
2. b) 2'-5'-ribonucleotides in positions 15, 16, 17, 18 and 19, a 3'-terminal phosphate; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
3. c) 2'OMe modified ribonucleotides in positions 5, 7, 13, and 16; a 2'-5'-ribonucleotide in position 18; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
4. d) 2'OMe modified ribonucleotides in positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide in position 9; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; or
5. e) 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3Pi moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0134] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3 moiety covalently attached to the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0135] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a 3'-terminal phosphate; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0136] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) includes 2'OMe modified ribonucleotides in positions 5, 7, 13, and 16; a 2'-5'-ribonucleotide in position 18; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0137] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) includes 2'OMe modified ribonucleotides in positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide in position 9; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0138] Provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 15, 17, 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3Pi moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0139] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15, 17, 19; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; and the sense strand (SEQ ID NO:195) includes 2'OMe modified ribonucleotides in positions 7, 9, 13, 16 and 18; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus.

[0140] In some embodiments provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:195) includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a 3'-terminal phosphate and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:220) includes an antisense strand selected from one of

1. a) 2'OMe modified ribonucleotides in positions 3, 5, 7, 9, 11, 13, 15, 17, 19; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
2. b) 2'OMe modified ribonucleotides in positions 1, 3, 6, 8, 10, 12, 14, 17, 18; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0141] In some embodiments provided herein is a double-stranded nucleic acid molecule as a component of a pharmaceutical formulation which includes the antisense strand set forth in SEQ ID NO:130 and the sense strand set forth in SEQ ID NO:63; identified herein as SERPINH1_12. In some embodiments the duplex comprises the structure

```

5' AACUCGUCUCGACUCUUGU -Z 3' (antisense SEQ ID NO:221)
|||||
3' Z'-UUGAGCAGAGCGUAGAACA -Z" 5' (sense SEQ ID NO:196)

```

wherein each "|" represents base pairing between the ribonucleotides; wherein each of A, C, G, U is independently an unmodified or modified ribonucleotide, or an unconventional moiety;

wherein each of Z and Z' is independently present or absent, but if present is independently 1-5 consecutive nucleotides or non-nucleotide

moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present; and

wherein z'' may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of $N^2-(N')_y$.

[0142] In some embodiments provided is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:196) includes one or more 2'OMe modified pyrimidines; a 3'-terminal nucleotide or non-nucleotide overhang; and a cap moiety covalently attached at the 5'-terminus. In some embodiments the antisense strand (SEQ ID NO:221) includes one or more 2'OMe modified pyrimidines; a nucleotide or non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus.

[0143] In some embodiments provided is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:196) includes 2'OMe modified ribonucleotides in positions 2, 14 and 18; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:221) is selected from an antisense strand which includes

1. a) 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
2. b) 2'OMe modified ribonucleotides in positions 3, 5, 7, 9, 12, 13 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0144] In some embodiments provided is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:196) includes 2'OMe modified ribonucleotides in positions 2, 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:221) includes 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0145] In some embodiments provided is a duplex oligonucleotide molecule wherein the sense strand (SEQ ID NO:196) includes 2'OMe modified ribonucleotides in positions 14 and 18 and optionally in position 2; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:221) includes 2'OMe modified ribonucleotides in positions 3, 5, 7, 9, 12, 13, and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0146] In some embodiments provided is a duplex oligonucleotide molecule wherein the sense strand (SEQ ID NO:196) includes 2'OMe modified ribonucleotides in positions 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:221) is selected from an antisense strand which includes

1. a) 2'OMe modified ribonucleotides in positions 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus; or
2. b) 2'OMe modified ribonucleotides in positions 3, 5, 7, 9, 12, 13 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0147] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:196) includes 2'OMe modified ribonucleotides in positions 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15 and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0148] Provided herein is a double-stranded nucleic acid molecule wherein the sense strand (SEQ ID NO:196) includes 2'OMe modified ribonucleotides in positions 14 and 18; a C3-OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand (SEQ ID NO:220) includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 7, 9, 12, 13, and 17; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached to the 3'-terminus.

[0149] In further embodiments of Structures A1 and A2, $(N')_y$ includes 1-8 modified ribonucleotides wherein the modified ribonucleotide is a DNA nucleotide. In certain embodiments $(N')_y$ includes 1, 2, 3, 4, 5, 6, 7, or up to 8 DNA moieties.

[0150] In some embodiments, either Z or Z' is present and independently includes two non-nucleotide moieties.

[0151] In additional embodiments, Z and Z' are present and each independently includes two non-nucleotide moieties.

[0152] In some embodiments, each of Z and Z' includes an abasic moiety, for example a deoxyribo-abasic moiety (referred to herein as "dAb") or ribo-abasic moiety (referred to herein as "rAb"). In some embodiments each of Z and/or Z' includes two covalently linked abasic moieties and is for example dAb-dAb or rAb-rAb or dAb-rAb or rAb-dAb, wherein each moiety is covalently attached to an adjacent moiety, preferably via a phospho-based bond. In some embodiments the phospho-based bond includes a phosphorothioate, a phosphonoacetate or a phosphodiester bond. In preferred embodiments the phospho-based bond includes a phosphodiester bond.

[0153] In some embodiments, each of Z and/or Z' independently includes an alkyl moiety, optionally propane $((CH_2)_3)$ moiety ("C3") or a


derivative thereof including propanediol (C3OH) and phospho derivative of propanediol ("C3Pi"). In some embodiments each of Z and/or Z' includes two alkyl moieties covalently linked to the 3'-terminus of the antisense strand or sense strand via a phosphodiester or phosphorothioate linkage and covalently linked to one another via a phosphodiester or phosphorothioate linkage and in some examples is C3Pi-C3Pi or C3Pi-C3OH. The 3'-terminus of the antisense strand and/or the 3'-terminus of the sense strand is covalently attached to a C3 moiety via a phospho-based bond and the C3 moiety is covalently conjugated a C3OH moiety via a phospho-based bond. In some embodiments the phospho-based bonds include a phosphorothioate, a phosphonoacetate or a phosphodiester bond. In preferred embodiments the phospho-based bond includes a phosphodiester bond.

[0154] In various embodiments of Structure A1 or Structure A2, Z and Z' are absent. In other embodiments Z or Z' is present. In some embodiments, each of Z and/or Z' independently includes a C2, C3, C4, C5 or C6 alkyl moiety, optionally a C3 moiety or a derivative thereof including propanol (C3OH/C3OH), propanediol, and phosphodiester derivative of propanediol (C3Pi). In preferred embodiments, each of Z and/or Z' includes two hydrocarbon moieties and in some examples is C3Pi-C3OH or C3Pi-C3Pi. Each C3 is covalently conjugated to an adjacent C3 via a covalent bond, preferably a phospho-based bond. In some embodiments, the phospho-based bond is a phosphorothioate, a phosphonoacetate or a phosphodiester bond.

[0155] In specific embodiments, $x=y=19$ and Z comprises at least one C3 alkyl overhang. In some embodiments the C3-C3 overhang is covalently attached to the 3'-terminus of $(N)_x$ or $(N')_y$ via a covalent linkage, preferably a phosphodiester linkage. In some embodiments, the linkage between a first C3 and a second C3 is a phosphodiester linkage. In some embodiments, the 3' non-nucleotide overhang is C3Pi-C3Pi. In some embodiments the 3' non-nucleotide overhang is C3Pi-C3OH. In some embodiments the 3' non-nucleotide overhang is C3Pi-C3OH.

[0156] In various embodiments, the alkyl moiety comprises an alkyl derivative including a C3 alkyl, C4 alkyl, C5 alkyl or C6 alkyl moiety comprising a terminal hydroxyl, a terminal amino, or terminal phosphate group. In some embodiments, the alkyl moiety is a C3 alkyl or C3 alkyl derivative moiety. In some embodiments, the C3 alkyl moiety comprises propanol, propylphosphate, propylphosphorothioate or a combination thereof. The C3 alkyl moiety is covalently linked to the 3'-terminus of $(N')_y$ and/or the 3'-terminus of $(N)_x$ via a phosphodiester bond. In some embodiments, the alkyl moiety comprises propanol, propyl phosphate or propyl phosphorothioate. In some embodiments, each of Z and Z' is independently selected from propanol, propyl phosphate propyl phosphorothioate, combinations thereof or multiples thereof in particular 2 or 3 covalently linked propanol, propyl phosphate, propyl phosphorothioate or combinations thereof. In some embodiments, each of Z and Z' is independently selected from propyl phosphate, propyl phosphorothioate, propyl phospho-propanol; propyl phospho-propyl phosphorothioate; propylphospho-propyl phosphate; (propyl phosphate)₃, (propyl phosphate)₂-propanol, (propyl phosphate)₂-propyl phosphorothioate. Any propane or propanol conjugated moiety can be included in Z or Z'.

[0157] The structures of exemplary 3'-terminal non-nucleotide moieties are as follows:



[0158] In some embodiments, each of Z and Z' is independently selected from propanol, propyl phosphate, propyl phosphorothioate, combinations thereof or multiples thereof.

[0159] In some embodiments, each of Z and Z' is independently selected from propyl phosphate, propyl phosphorothioate, propyl phospho-propanol; propyl phospho-propyl phosphorothioate; propylphospho-propyl phosphate; (propyl phosphate)₃, (propyl phosphate)₂-propanol, (propyl phosphate)₂-propyl phosphorothioate. Any propane or propanol conjugated moiety can be included in Z or Z'.

[0160] In additional embodiments, each of Z and/or Z' includes a combination of an abasic moiety and an unmodified deoxyribonucleotide or ribonucleotide or a combination of a hydrocarbon moiety and an unmodified deoxyribonucleotide or ribonucleotide or a combination of an abasic moiety (deoxyribo or ribo) and a hydrocarbon moiety. In such embodiments, each of Z and/or Z' includes C3-rAb or C3-dAb wherein each moiety is covalently bond to the adjacent moiety via a phospho-based bond, preferably a phosphodiester, phosphorothioate or phosphonoacetate bond.

[0161] In certain embodiments, nucleic acid molecules as disclosed herein include a sense oligonucleotide sequence selected from any one of oligonucleotide shown *infra* in Tables 4 and 5 (SEQ ID NOS:60-126 and 194-218).

[0162] In certain preferred embodiments, compounds provided include Compound_1, Compound_2, Compound_3, Compound_4, Compound_5, Compound_6, Compound_7, Compound_8, and Compound_9, described *infra*.

[0163] In some embodiments, (such as, for example, Compound_1, Compound_5, and Compound_6) provided are 19-mer double-stranded nucleic acid molecules wherein the antisense strand is SEQ ID NO:127 and the sense strand is SEQ ID NO:60. In certain embodiments, provided are 19-mer double-stranded nucleic acid molecules wherein the antisense strand is SEQ ID NO:127 and includes 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 1, 5, 6, or 7; and a non-nucleotide moiety covalently attached to the 3'-terminus; and the sense strand is SEQ ID NO:60 and includes at least one 2'-5'-ribonucleotide or 2'OMe modified ribonucleotide; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus. In some embodiments, provided are 19-mer double-stranded nucleic acid molecule wherein the antisense strand is SEQ ID NO:127; and includes 2'OMe modified ribonucleotides at positions 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and the sense strand is SEQ ID NO:60 and includes five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a C3PI non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.

[0164] In one embodiment, provided is Compound_1 that is a 19-mer double-stranded nucleic acid molecule wherein the antisense strand is SEQ ID NO:127 and includes 2'OMe modified ribonucleotides at positions 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3PI-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and the sense strand is SEQ ID NO:60 and includes five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a C3PI non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus; and that further includes a 2'OMe modified ribonucleotide at position 1 of the antisense strand.

[0165] In one embodiment, provided is Compound_6 that is a 19-mer double-stranded nucleic acid molecule wherein the antisense strand is SEQ ID NO:127 and includes 2'OMe modified ribonucleotides at positions 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3PI-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and the sense strand is SEQ ID NO:60 and includes five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a C3PI non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus; and that further includes a 2'-5'-ribonucleotide at position 1 of the antisense strand.

[0166] In one embodiment, provided is Compound_5 that is a 19-mer double-stranded nucleic acid molecule wherein the antisense strand is SEQ ID NO:127 and includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3PI-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and the sense strand is SEQ ID NO:60 and includes 2'OMe modified ribonucleotides in positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide at position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.

[0167] In some embodiments, (such as, for example, Compound_2, and Compound_7, described *infra*) provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:63 and the antisense strand is SEQ ID NO:130. In some embodiments provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:63 and includes 2'OMe modified pyrimidine ribonucleotides; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:130 and includes 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide at position 7; and a non-nucleotide moiety covalently attached at the 3'-terminus. In some embodiments provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:63 and includes 2'OMe modified ribonucleotides; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:130 and includes 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 5, 6 or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus.

[0168] In one embodiment, provided is Compound_2 that is a 19-mer double-stranded nucleic acid molecule wherein the sense strand is SEQ ID NO:63 and includes 2'OMe modified ribonucleotides in positions 2, 14 and 18; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:130 and includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 12, 13, and 17; a 2'-5'-ribonucleotide in at least one of positions 5, 6 or 7; and C3PI-C3OH non-nucleotide moiety covalently attached at the 3'-terminus.

[0169] In one embodiment, provided is Compound_7 that is a 19-mer double-stranded nucleic acid molecule wherein the sense strand is SEQ ID NO:63 and includes 2'OMe modified ribonucleotides in positions 2, 14 and 18; a C3OH moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:130 and includes 2'OMe modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, and 17; a 2'-5'-ribonucleotide at position 7; and a C3PI-C3OH non-nucleotide moiety covalently attached at the 3'-terminus.

[0170] In some embodiments, (such as, for example, Compound_3, described *infra*) provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:98 and the antisense strand is SEQ ID NO:165. In some embodiments, provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:98 and includes 2'-5'-ribonucleotides in positions at the 3'-terminus; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:165 and includes 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 5, 6 or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus. In one embodiment, provided is Compound_3 that is a 19-mer double-stranded nucleic acid molecule wherein the sense strand is SEQ ID NO:98 and includes 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3-OH 3' moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:165 and includes 2'OMe modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3PI-C3OH covalently attached at the 3'-terminus.

[0171] In some embodiments, (such as, for example, Compound_4, Compound_8 and Compound_9, described *infra*) provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:101 and the antisense strand is SEQ ID NO:168. In some embodiments provided are 19-mer double-stranded nucleic acid molecules wherein the sense strand is SEQ ID NO:101 and includes 2'OMe modified pyrimidine ribonucleotides; an optional 2'-5'-ribonucleotide in one of position 9 or 10; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:168 and includes 2'OMe modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 5, 6, or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus.

[0172] In one embodiment, provided is Compound_4 that is a 19-mer double-stranded nucleic acid molecule wherein sense strand is SEQ ID NO:101 and includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:168 and includes 2'OMe modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; a C3Pi-C3OH overhang covalently attached at the 3'-terminus.

[0173] In one embodiment, provided is Compound_8 that is a 19-mer double-stranded nucleic acid molecule wherein sense strand is SEQ ID NO:101 and includes 2'OMe modified ribonucleotides in positions 4, 11, 13, and 17; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand is SEQ ID NO:168 and includes 2'OMe modified ribonucleotides in positions 1, 4, 8, 13 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH overhang covalently attached at the 3'-terminus.

[0174] In one embodiment, provided is Compound_9 that is a 19-mer double-stranded nucleic acid molecule wherein the sense strand is SEQ ID NO:101 and includes 2'OMe modified ribonucleotides in positions 2, 4, 11, 13, and 17; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and the antisense strand is SEQ ID NO:168 and includes 2'OMe modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.

[0175] In another aspect, provided are methods for reducing the expression of hsp47 in a cell by introducing into a cell a nucleic acid molecule as provided herein in an amount sufficient to reduce expression of hsp47. In one embodiment, the cell is hepatocellular stellate cell. In another embodiment, the cell is a stellate cell in renal or pulmonary tissue. In certain embodiment, the method is performed *in vitro*. In another embodiment, the method is performed *in vivo*.

[0176] In another aspect, provided are methods for treating an individual suffering from a disease associated with hsp47. The methods include administering to the individual a nucleic acid molecule such as provided herein in an amount sufficient to reduce expression of hsp47. In certain embodiments, the disease associated with hsp47 is a disease selected from the group consisting of liver fibrosis, cirrhosis, pulmonary fibrosis including lung fibrosis (including ILF), any condition causing kidney fibrosis (e.g., CKD including ESRD), peritoneal fibrosis, chronic hepatic damage, fibrillogenesis; fibrotic diseases in other organs; abnormal scarring (keloids) associated with all possible types of skin injury, accidental and iatrogenic (operations), scleroderma, cardiofibrosis, failure of glaucoma filtering operation, and intestinal adhesions. In some embodiments, the compounds may be useful in treating organ-specific indications, for example, indications including those shown in Table 1 below, listing organs and respective indications:

TABLE 1

Skin
• Pathologic scarring as keloid and hypertrophic scar
• Surgical scarring
• Injury scarring
• keloid, or nephrogenic fibrosing dermatopathy
Peritoneum
• Peritoneal fibrosis
• Adhesions
• Peritoneal Sclerosis associated with continual ambulatory peritoneal dialysis (CAPD)
Liver
• Cirrhosis including post-hepatitis C cirrhosis, primary biliary cirrhosis
• Liver fibrosis, e.g. Prevention of Liver Fibrosis in Hepatitis C carriers
• schistomasomiasis
• cholangitis
• Liver cirrhosis due to Hepatitis C post liver transplant or Non-Alcoholic Steatohepatitis (NASH)
Pancreas
• inter(peri)lobular fibrosis (e.g., alcoholic chronic pancreatitis), periductal fibrosis (e.g., hereditary pancreatitis), periductal and interlobular fibrosis (e.g., autoimmune pancreatitis), diffuse inter- and intralobular fibrosis (e.g., obstructive chronic pancreatitis)
Kidney
• Chronic Kidney Disease (CKD) of any etiology. Treatment of early stage CKD (elevated SCr) in diabetic patients (prevent further deterioration in renal function)
• kidney fibrosis associated with lupus glomerulosclerosis
• Diabetic Nephropathy

Heart
• Congestive heart failure,
• Endomyocardial fibrosis,
• cardiofibrosis
• fibrosis associated with myocardial infarction
Lung
• Asthma, Idiopathic pulmonary fibrosis (IPF);
• Interstitial lung fibrosis (ILF)
• Radiation Pneumonitis leading to Pulmonary Fibrosis (e.g. due to cancer treating radiation)
Bone marrow
• Myeloproliferative disorders: Myelofibrosis (MF), Polycythemia vera (PV), Essential thrombocythemia (ET)
• idiopathic myelofibrosis
• drug induced myelofibrosis.
Eye
• <u>Anterior segment</u> : Corneal opacification e.g. following inherited dystrophies, herpetic keratitis or pterygia; Glaucoma
• <u>Posterior segment</u> fibrosis and traction retinal detachment, a complication of advanced diabetic retinopathy (DR); Fibrovascular scarring and gliosis in the retina;
• <u>Under the retina</u> fibrosis for example subsequent to subretinal hemorrhage associated with neovascular AMD
• Retro-orbital fibrosis, postcataract surgery, proliferative vitreoretinopathy.
• Ocular cicatricial pemphigoid
Intestine
• Intestinal fibrosis,
• Crohn's disease
Vocal cord
• Vocal cord scarring,
• vocal cord mucosal fibrosis,
• laryngeal fibrosis
Vasculature
• Atherosclerosis,
• postangioplasty arterial restenosis
Multisystemic
• Scleroderma systemic sclerosis;
• multifocal fibrosclerosis;
• sclerodermatous graft-versus-host disease in bone marrow transplant recipients, and
• nephrogenic systemic fibrosis (exposure to gadolinium-based contrast agents (GBCAs), 30% of MRIs)
Malignancies of various origin
• Metastatic and invasive cancer by inhibiting function of activated tumor associated myofibroblasts

[0177] Another embodiment of the description is a method for treating a stellate cell-related disorder, the method comprising administering an effective amount of the pharmaceutical composition described *infra* to a subject in need thereof. The disorder includes hepatitis, hepatic fibrosis, hepatic cirrhosis, liver cancer, pancreatitis, pancreatic fibrosis, pancreatic cancer, vocal cord scarring, vocal cord mucosal fibrosis, and laryngeal fibrosis. In some embodiments the preferred indications include, liver cirrhosis due to Hepatitis C post liver transplant; liver cirrhosis due to Non-Alcoholic Steatohepatitis (NASH); idiopathic pulmonary fibrosis; radiation pneumonitis leading to pulmonary fibrosis; diabetic nephropathy; peritoneal sclerosis associated with continual ambulatory peritoneal dialysis (CAPD) and ocular cicatricial pemphigoid.

[0178] Fibrotic liver indications include Alcoholic Cirrhosis, Hepatitis B cirrhosis, Hepatitis C cirrhosis, Hepatitis C (Hep C) cirrhosis post-orthotopic liver transplant (OLT), non-alcoholic steatohepatitis/nonalcoholic fatty liver disease (NASH/NAFLD), primary biliary cirrhosis (PBC), primary sclerosing cholangitis (PSC), biliary atresia, alpha-1 antitrypsin deficiency (A1AD), copper storage diseases (Wilson's disease), fructosemia, galactosemia, glycogen storage diseases (especially types III, IV, VI, IX, and X), iron-overload syndromes (hemochromatosis), lipid abnormalities (e.g., Gaucher's disease), peroxisomal disorders (e.g., Zellweger syndrome), tyrosinemia, congenital hepatic fibrosis, bacterial infections (e.g., brucellosis), parasitic (e.g., echinococcosis), Budd-Chiari syndrome (hepatic veno-occlusive disease).

[0179] Pulmonary indications include Idiopathic Pulmonary Fibrosis, Silicosis, Pneumoconiosis, bronchopulmonary dysplasia in newborn following neonatal respiratory distress syndrome, bleomycin/chemolung injury, bronchiolitis obliterans (BOS) Post-lung transplant, Chronic obstructive pulmonary disorder (COPD), cystic fibrosis, and asthma.

[0180] Cardiac indications include cardiomyopathy, atherosclerosis (Bergers disease, etc), endomyocardial fibrosis, atrial fibrillation, scarring post-myocardial infarction (MI)

[0181] Other Thoracic indications include Radiation-induced capsule tissue reactions around textured breast implants, and oral submucosal fibrosis.

[0182] Renal indications include autosomal dominant polycystic kidney disease (ADPKD), Diabetic nephropathy (diabetic glomerulosclerosis), focal segmental glomerulosclerosis (FSGS) (collapsing vs. other histologic variants), IgA Nephropathy (Berger Disease), Lupus Nephritis, Wegner's, Scleroderma, Goodpasture Syndrome, tubulointerstitial fibrosis: drug induced (protective) penicillins, cephalosporins, analgesic nephropathy, membrano-proliferative glomerulonephritis (MPGN), Henoch-Schonlein purpura, Congenital nephropathies: Medullary Cystic Disease, Nail-Patella Syndrome and Alport Syndrome.

[0183] Bone Marrow indications include lymphangiolymphomatosis (LAM), chronic graft vs. host disease, polycythemia vera, essential thrombocythemia, and myelofibrosis.

[0184] Ocular indications include retinopathy of prematurity (RoP), ocular cicatricial pemphigoid, Lacrimal gland fibrosis, Retinal attachment surgery, corneal opacity, herpetic keratitis, pterygia, Glaucoma, age-related macular degeneration (AMD/ARMD), retinal fibrosis associated diabetes mellitus (DM) retinopathy

[0185] Gynecological indications include Endometriosis add on to hormonal therapy for prevention of scarring, post-STD fibrosis/salpingitis,

[0186] Systemic indications include Dupuytren's disease, Palmar fibromatosis, Peyronie's disease, Ledderhose disease, keloids, multifocal fibrosclerosis, nephrogenic systemic fibrosis and myelofibrosis (anemia).

[0187] Injury-associated fibrotic diseases include burn-induced (chemical included) skin and soft tissue scarring and contraction, radiation-induced skin and organ scarring, post cancer therapeutic radiation treatment, keloid (skin).

[0188] Surgical indications include peritoneal fibrosis post-peritoneal dialysis catheter, corneal implant, cochlear implant, other implants , silicone implants in breasts, chronic sinusitis; adhesions, pseudointimal hyperplasia of dialysis grafts.

[0189] Other indications include chronic pancreatitis.

[0190] In some embodiments, provided is a method for treatment of a subject suffering from liver fibrosis comprising administering to the subject an effective amount of a nucleic acid molecule disclosed herein, thereby treating liver fibrosis. In some embodiments the subject is suffering from cirrhosis of the liver due to hepatitis. In some embodiments the subject is suffering from cirrhosis of the liver due to NASH.

[0191] In some embodiments, provided is the use of a nucleic acid molecule disclosed herein for the manufacture of a medicament to treat liver fibrosis. In some embodiments, the liver fibrosis is due to hepatitis. In some embodiments, the liver fibrosis is due to NASH.

[0192] In some embodiments, provided is a method for remodeling of scar tissue comprising administering to a subject in need thereof an effective amount of a nucleic acid molecule disclosed herein, thereby effecting scar tissue remodeling. In some embodiments, the scar tissue is in the liver. In some embodiments, the subject is suffering from cirrhosis of the liver due to hepatitis. In some embodiments, the subject is suffering from cirrhosis of the liver due to NASH.

[0193] In some embodiments, a method for modulating fibrosis regression is provided comprising administering to a subject in need thereof an effective amount of a nucleic acid molecule disclosed herein, thereby effecting fibrosis regression.

[0194] In some embodiments, provided is a method for reduction of scar tissue in a subject comprising the step of administering to the subject an effective amount of a nucleic acid molecule disclosed herein to reduce the scar tissue. In some embodiments, provided is a method for reducing scar tissue in a subject comprising the step of topically applying to scar tissue an effective amount of a nucleic acid molecule disclosed herein to reduce scar tissue.

[0195] In some embodiments, provided is a method for improving the appearance of scar tissue comprising the step of topically applying to scar tissue an effective amount of a nucleic acid molecule disclosed herein to improve the appearance of the scar tissue.

[0196] In some embodiments, provided is a method for treatment of a subject suffering from lung fibrosis comprising administering to the subject an effective amount of a nucleic acid molecule disclosed herein, thereby treating the lung fibrosis. In some embodiments, the subject is suffering from interstitial lung fibrosis (ILF). In some embodiments, the subject is suffering from Radiation Pneumonitis leading to Pulmonary Fibrosis. In some embodiments, the subject is suffering from drug induced lung fibrosis.

[0197] In some embodiments, provided is the use of a nucleic acid molecule disclosed herein for the manufacture of a medicament to treat lung fibrosis. In some embodiments, the lung fibrosis is ILF. In some embodiments, the lung fibrosis drug- or radiation- induced lung fibrosis.

[0198] In one aspect, provided are pharmaceutical compositions that include a nucleic acid molecule (e.g., a siRNA molecule) as described herein in a pharmaceutically acceptable carrier. In certain embodiments, the pharmaceutical formulation includes, or involves, a delivery system suitable for delivering nucleic acid molecules (e.g., siRNA molecules) to an individual such as a patient; for example delivery systems described in more detail below.

[0199] In a related aspect, provided are compositions or kits that include a nucleic acid molecule (e.g., an siRNA molecule) packaged for use by a patient. The package may be labeled or include a package label or insert that indicates the content of the package and provides certain

information regarding how the nucleic acid molecule (e.g., an siNA molecule) should be or can be used by a patient, for example the label may include dosing information and/or indications for use. In certain embodiments, the contents of the label will bear a notice in a form prescribed by a government agency, for example the United States Food and Drug administration (FDA). In certain embodiments, the label may indicate that the nucleic acid molecule (e.g., an siNA molecule) is suitable for use in treating a patient suffering from a disease associated with hsp47; for example, the label may indicate that the nucleic acid molecule (e.g., an siNA molecule) is suitable for use in treating fibroids; or for example the label may indicate that the nucleic acid molecule (e.g., an siNA molecule) is suitable for use in treating a disease selected from the group consisting of fibrosis, liver fibrosis, cirrhosis, pulmonary fibrosis, kidney fibrosis, peritoneal fibrosis, chronic hepatic damage, and fibrillogenesis.

[0200] As used herein, the term "heat shock protein 47" or "hsp47" or "HSP47" are used interchangeably and refer to any heat shock protein 47, peptide, or polypeptide having any hsp47 protein activity. Heat shock protein 47 is a serine proteinase inhibitor (serpin) also known, for example, as serpin peptidase inhibitor, clade H, member 1 (SERPINH1), SERPINH2, collagen binding protein 1 (CBP1), CBP2, gp46; arsenic-transactivated protein 3 (AsTP3); HSP47; proliferation-inducing gene 14 (PIG14); PPR0M; rheumatoid arthritis antigen A-47 (RA-A47); colligin-1; and colligin-2. In certain preferred embodiments, "hsp47" refers to human hsp47. Heat shock protein 47 (or more particularly human hsp47) may have an amino acid sequence that is the same, or substantially the same, as SEQ ID NO. 2.

[0201] As used herein the term "nucleotide sequence encoding hsp47" means a nucleotide sequence that codes for an hsp47 protein, or portion thereof. The term "nucleotide sequence encoding hsp47" is also meant to include hsp47 coding sequences such as hsp47 isoforms, mutant hsp47 genes, splice variants of hsp47 genes, and hsp47 gene polymorphisms. A nucleic acid sequence encoding hsp47 includes mRNA sequences encoding hsp47, which can also be referred to as "hsp47 mRNA." An exemplary sequence of human hsp47 mRNA is SEQ ID. NO. 1.

[0202] As used herein, the term "nucleic acid molecule" or "nucleic acid" are used interchangeably and refer to an oligonucleotide, nucleotide or polynucleotide. Variations of "nucleic acid molecule" are described in more detail herein. A nucleic acid molecule encompasses both modified nucleic acid molecules and unmodified nucleic acid molecules as described herein. A nucleic acid molecule may include deoxyribonucleotides, ribonucleotides, modified nucleotides or nucleotide analogs in any combination.

[0203] As used herein, the term "nucleotide" refers to a chemical moiety having a sugar (or an analog thereof, or a modified sugar), a nucleotide base (or an analog thereof, or a modified base), and a phosphate group (or analog thereof, or a modified phosphate group). A nucleotide encompasses modified nucleotides or unmodified nucleotides as described herein. As used herein, nucleotides may include deoxyribonucleotides (e.g., unmodified deoxyribonucleotides), ribonucleotides (e.g., unmodified ribonucleotides), and modified nucleotide analogs including, inter alia, LNA and UNA, peptide nucleic acids, L-nucleotides (also referred to as mirror nucleotides), ethylene-bridged nucleic acid (ENA), arabinoside, PACE, nucleotides with a 6 carbon sugar, as well as nucleotide analogs (including abasic nucleotides) often considered to be non-nucleotides. In some embodiments, nucleotides may be modified in the sugar, nucleotide base and/or in the phosphate group with any modification known in the art, such as modifications described herein. A "polynucleotide" or "oligonucleotide" as used herein, refers to a chain of linked nucleotides; polynucleotides and oligonucleotides may likewise have modifications in the nucleotide sugar, nucleotide bases and phosphate backbones as are well known in the art and/or are disclosed herein.

[0204] As used herein, the term "short interfering nucleic acid", "siNA", or "short interfering nucleic acid molecule" refers to any nucleic acid molecule capable of modulating gene expression or viral replication. Preferably siNA inhibits or down regulates gene expression or viral replication. siNA includes without limitation nucleic acid molecules that are capable of mediating sequence specific RNAi, for example short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), short hairpin RNA (shRNA), short interfering oligonucleotide, short interfering nucleic acid, short interfering modified oligonucleotide, chemically-modified siRNA, post-transcriptional gene silencing RNA (ptgsRNA), and others. As used herein, "short interfering nucleic acid", "siNA", or "short interfering nucleic acid molecule" has the meaning described in more detail elsewhere herein.

[0205] As used herein, the term "complementary" means that a nucleic acid can form hydrogen bond(s) with another nucleic acid sequence by either traditional Watson-Crick or other non-traditional types. In reference to the nucleic molecules disclosed herein, the binding free energy for a nucleic acid molecule with its complementary sequence is sufficient to allow the relevant function of the nucleic acid to proceed, e.g., RNAi activity. Determination of binding free energies for nucleic acid molecules is well known in the art (see, e.g., Turner et al., 1987, CSH Symp. Quant. Biol. LII pp. 123-133; Frier et al., 1986, Proc. Nat. Acad. Sci. USA 83:9373-9377; Turner et al., 1987, J. Am. Chem. Soc. 109:3783-3785). A percent complementarity indicates the percentage of contiguous residues in a nucleic acid molecule that can form hydrogen bonds (e.g., Watson-Crick base pairing) with a second nucleic acid sequence (e.g., 5, 6, 7, 8, 9, or 10 nucleotides out of a total of 10 nucleotides in the first oligonucleotide being base paired to a second nucleic acid sequence having 10 nucleotides represents 50%, 60%, 70%, 80%, 90%, and 100% complementary respectively). "Fully complementary" means that all the contiguous residues of a nucleic acid sequence will hydrogen bond with the same number of contiguous residues in a second nucleic acid sequence. In one embodiment, a nucleic acid molecule disclosed herein includes about 15 to about 35 or more (e.g., about 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 or 35 or more) nucleotides that are complementary to one or more target nucleic acid molecules or a portion thereof.

[0206] As used herein, the term "sense region" refers to a nucleotide sequence of a siNA molecule complementary (partially or fully) to an antisense region of the siNA molecule. The sense strand of a siNA molecule can include a nucleic acid sequence having homology with a target nucleic acid sequence. As used herein, "sense strand" refers to nucleic acid molecule that includes a sense region and may also include additional nucleotides. Nucleotide positions of the sense strand are herein numbered 5'>3'.

[0207] As used herein, the term "antisense region" refers to a nucleotide sequence of a siNA molecule complementary (partially or fully) to a target nucleic acid sequence. The antisense strand of a siNA molecule can optionally include a nucleic acid sequence complementary to a sense region of the siNA molecule. As used herein, "antisense strand" refers to nucleic acid molecule that includes an antisense region and may also include additional nucleotides. Nucleotide positions of the antisense strand are herein numbered 5'>3'.

[0208] As used herein, the term "RNA" refers to a molecule that includes at least one ribonucleotide residue.

[0209] As used herein, the term "duplex region" refers to the region in two complementary or substantially complementary oligonucleotides that form base pairs with one another, either by Watson-Crick base pairing or any other manner that allows for a duplex between oligonucleotide strands that are complementary or substantially complementary. For example, an oligonucleotide strand having 21 nucleotide units can base pair with another oligonucleotide of 21 nucleotide units, yet only 19 bases on each strand are complementary or substantially complementary, such that the "duplex region" consists of 19 base pairs. The remaining base pairs may, for example, exist as 5' and 3' overhangs. Further, within the duplex region, 100% complementarity is not required; substantial complementarity is allowable within a duplex region. Substantial complementarity refers to complementarity between the strands such that they are capable of annealing under biological conditions. Techniques to empirically determine if two strands are capable of annealing under biological conditions are well known in the art. Alternatively, two strands can be synthesized and added together under biological conditions to determine if they anneal to one another.

[0210] As used herein, the terms "non-pairing nucleotide analog" means a nucleotide analog which includes a non-base pairing moiety including but not limited to: 6-des-amino adenosine (nebularine), 4-Me-indole, 3-nitroprole, 5-nitroindole, Ds, Pa, N3-Me ribo U, N3-Me riboT, N3-Me dC, N3-Me-dT, N1-Me-dG, N1-Me-dA, N3-ethyl-dC, N3-Me dC. In some embodiments the non-base pairing nucleotide analog is a ribonucleotide. In other embodiments it is a deoxyribonucleotide.

[0211] As used herein, the term, "terminal functional group" includes without limitation a halogen, alcohol, amine, carboxylic, ester, amide, aldehyde, ketone, ether groups.

[0212] An "abasic nucleotide" or "abasic nucleotide analog" is as used herein may also be often referred to herein and in the art as a pseudo-nucleotide or an unconventional moiety. While a nucleotide is a monomeric unit of nucleic acid, generally consisting of a ribose or deoxyribose sugar, a phosphate, and a base (adenine, guanine, thymine, or cytosine in DNA; adenine, guanine, uracil, or cytosine in RNA), an abasic or pseudo-nucleotide lacks a base, and thus is not strictly a nucleotide as the term is generally used in the art. Abasic deoxyribose moieties include for example, abasic deoxyribose-3'-phosphate; 1,2-dideoxy-D-ribofuranose-3-phosphate; 1,4-anhydro-2-deoxy-D-ribitol-3-phosphate. Inverted abasic deoxyribose moieties include inverted deoxyriboabasic; 3',5' inverted deoxyabasic 5'-phosphate.

[0213] The term "capping moiety" (z") as used herein includes a moiety which can be covalently linked to the 5'-terminus of (N')_y and includes abasic ribose moiety, abasic deoxyribose moiety, modifications abasic ribose and abasic deoxyribose moieties including 2' O alkyl modifications; inverted abasic ribose and abasic deoxyribose moieties and modifications thereof; C₆-imino-Pi; a mirror nucleotide including L-DNA and L-RNA; 5'OMe nucleotide; and nucleotide analogs including 4',5'-methylene nucleotide; 1-(β-D-erythrofuranosyl)nucleotide; 4'-thio nucleotide, carbocyclic nucleotide; 5'-amino-alkyl phosphate; 1,3-diamino-2-propyl phosphate, 3-aminopropyl phosphate; 6-aminoethyl phosphate; 12-aminododecyl phosphate; hydroxypropyl phosphate; 1,5-anhydrohexitol nucleotide; alpha-nucleotide; threo-pentofuranosyl nucleotide; acyclic 3',4'-seco nucleotide; 3,4-dihydroxybutyl nucleotide; 3,5-dihydroxypentyl nucleotide, 5'-5'-inverted abasic moiety; 1,4-butanediol phosphate; 5'-amino; and bridging or non-bridging methylphosphonate and 5'-mercapto moieties.

[0214] Certain capping moieties may be abasic ribose or abasic deoxyribose moieties; inverted abasic ribose or abasic deoxyribose moieties; C₆-amino-Pi; a mirror nucleotide including L-DNA and L-RNA. The nucleic acid molecules as disclosed herein may be synthesized using one or more inverted nucleotides, for example inverted thymidine or inverted adenine (for example see Takei, et al., 2002. JBC 277(26):23800-06).

[0215] The term "unconventional moiety" as used herein refers to non-nucleotide moieties including an abasic moiety, an inverted abasic moiety, a hydrocarbon (alkyl) moiety and derivatives thereof, and further includes a deoxyribonucleotide, a modified deoxyribonucleotide, a mirror nucleotide (L-DNA or L-RNA), a non-base pairing nucleotide analog and a nucleotide joined to an adjacent nucleotide by a 2'-5' internucleotide phosphate bond; bridged nucleic acids including LNA and ethylene bridged nucleic acids, linkage modified (e.g. PACE) and base modified nucleotides as well as additional moieties explicitly disclosed herein as unconventional moieties.

[0216] As used herein, the term "inhibit", "down-regulate", or "reduce" with respect to gene expression means the expression of the gene, or level of RNA molecules or equivalent RNA molecules encoding one or more proteins or protein subunits (e.g., mRNA), or activity of one or more proteins or protein subunits, is reduced below that observed in the absence of an inhibitory factor (such as a nucleic acid molecule, e.g., an siRNA, for example having structural features as described herein); for example the expression may be reduced to 90%, 80%, 70%, 60%, 50%, 40%, 30%, 20%, 10%, 5% or less than that observed in the absence of an inhibitor.

BRIEF DESCRIPTION OF THE DRAWINGS

[0217]

FIG. 1 is a diagram showing a protocol with respect to assessment of the effect of gp46-siRNA *in vitro* using NRK cells, and determination of optimal sequence, timing, and concentration.

FIG. 2 is a photographic diagram showing the result of western blotting of gp46 and actin (24 hour culturing, examination of optimal sequence).

FIG. 3 is a photographic diagram showing the result of western blotting of gp46 and actin (24 hour culturing, examination of optimal concentration).

FIG. 4 is a photographic diagram showing the result of western blotting of gp46 and actin (concentration 50 nM, examination of optimal culturing time).

FIG. 5 is a diagram showing a protocol for evaluating inhibition of expression of collagen by gp46-siRNA in NRK cells.

FIG. 6 is a graph showing inhibition of collagen synthesis by siRNA.

FIG. 7 is a photographic diagram showing HSC-specific siRNA transfection.

FIG. 8 is a photographic diagram for evaluating HSC-specific siRNA transfection percentage.

FIG. 9 is a photographic diagram for evaluating inhibition of expression of gp46 by siRNA.

FIG. 10 is a photographic diagram showing azan staining of rat liver to which DMN had been administered.

FIG. 11 is a diagram showing an LC rat treatment protocol.

FIG. 12 is a photographic diagram showing azan staining of LC rat liver to which VA-Lip-gp46siRNA had been administered.

FIG. 13. is a diagram showing a method for extracting a stained portion by means of NIH Image (6 positions being randomly taken from an azan-stained image).

FIG. 14 is a graph showing the ratio by area occupied by fibrotic portions in liver histology (Collagen ratio by area, %).

FIG. 15 is a graph showing the amount of hydroxyproline in hepatic tissue.

FIG. 16 is a graph showing a survival curve for hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intraportally administered.

FIG. 17 is a photographic diagram showing azan staining of hepatic tissue of hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intraportally administered.

FIG. 18 is a graph showing a survival curve for hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intraportally administered.

FIG. 19 is a photographic diagram showing azan staining of hepatic tissue of hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intraportally administered.

FIG. 20 is a graph showing a survival curve for hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intravenously administered.

FIG. 21 is a graph showing a survival curve for hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intravenously administered.

FIG. 22 is a photographic diagram showing azan staining of hepatic tissue of hepatic cirrhosis rat to which VA-Lip-gp46siRNA had been intravenously administered.

FIG. 23 is a diagram showing improvement of VA-Lip-gp46siRNA transfection efficiency by RBP.

FIG. 24 is a diagram showing inhibition of VA-Lip-gp46siRNA transfection by anti-RBP antibody.

Fig. 25 is a bar graph showing the effect of GFP siNA on various reporter cell lines. Cell lines were established by lenti-viral induction of human HSP47 cDNA-GFP or rat GP46 cDNA-GFP construct into HEK293, human fibrosarcoma cell line HT1080, human HSC line hTERT or NRK cell line. Negative control siNA or siNA against GFP was introduced into the cells and GFP fluorescence was measured. The results showed that siNA against GFP knocks down the fluorescence to different extent in different cell lines. 293_HSP47-GFP and 293_GP46-GFP cell lines were selected for siHsp47 screening due to their easiness of being transfected and sensitivity to fluorescence knockdown.

Fig. 26A, 26B, 26C, and 26D are a series of bar graphs showing the cytotoxicity and knockdown efficiency of various siHsp47s in 293_HSP47-GFP and 293_GP46-GFP cell lines. The result showed that siHsp47-C, siHsp47-2 and siHsp47-2d efficiently knockdown both human HSP47 and rat GP46 (the human hsp47 homolog) without substantial cytotoxicity. siGp46A against GP46 does not knock down human HSP47. Additionally, the newly designed siHsp47s outperformed siGp46A in knocking down rat GP46.

Fig. 27 is a bar graph showing the knock down effect of various siHsp47s on hsp47 mRNA, measured by TaqMan® qPCR using the human HSC cell line hTERT. The Y axis represents the remaining mRNA level of hsp47. HSP47-C was most effective among all the hsp47 siNAs tested.

Fig. 28 is a bar graph showing the effect of different hsp47 siNAs on collagen I expression in hTERT cells. The level of collagen I mRNA levels were measured by real-time quantitative PCR using TaqMan® probe. The Y axis represents the remaining mRNA expression level of collagen I. The result showed that collagen I mRNA level is significantly reduced in the cells treated with some of the candidates (siHsp47-2, siHsp47-2d, and their combination with siHsp47-1).

Fig. 29 is a graph showing a decrease in fibrotic areas of the liver in animals treated with siHSP47.

Fig. 30 is a schematic of the treatment schedule and evaluation method used in Example 22.

Fig. 31 shows the results of histological staining with Azan in the pulmonary field. The pictures display representative lung field of Azan-stained sections of each group at 80X magnification. (a) Pretreatment (BLM IT -2W); (b) Disease rat (BLM IT-5W+PBS i.v.); and (c) Treatment (BLM IT +siRNA i.v.), (d) Sham (Saline-IT+PBS i.v.).

Fig. 32 shows fibrosis scoring, showing the results of evaluating twenty randomly selected lung fields under 80X magnification for each rat. The bar graph summarizes the fibrosis scoring of Azan-stained section for each group. Statistical analysis were used One-way-ANOVA Bonferroni multi comparison test using Prism5 software.

DETAILED DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

RNA Interference and siNA Nucleic Acid Molecules

[0218] RNA interference refers to the process of sequence-specific post-transcriptional gene silencing in animals mediated by short interfering RNAs (siRNAs) (Zamore et al., 2000, Cell, 101, 25-33; Fire et al., 1998, Nature, 391, 806; Hamilton et al., 1999, Science, 286, 950-951; Lin et al., 1999, Nature, 402, 128-129; Sharp, 1999, Genes & Dev., 13:139-141; and Strauss, 1999, Science, 286, 886). The presence of dsRNA in cells triggers the RNAi response through a mechanism that has yet to be fully characterized.

[0219] Dicer is involved in the processing of the dsRNA into short pieces of dsRNA known as short interfering RNAs (siRNAs) (Zamore et al., 2000, Cell, 101, 25-33; Bass, 2000, Cell, 101, 235; Bernstein et al., 2001, Nature, 409, 363). siRNAs derived from dicer activity can be about 21 to about 23 nucleotides in length and include about 19 base pair duplexes (Zamore et al., 2000, Cell, 101, 25-33; Elbashir et al., 2001, Genes Dev., 15, 188). Dicer has also been implicated in the excision of 21- and 22-nucleotide small temporal RNAs (stRNAs) from precursor RNA of conserved structure that are implicated in translational control (Hutvagner et al., 2001, Science, 293, 834). The RNAi response also features an endonuclease complex, commonly referred to as an RNA-induced silencing complex (RISC), which mediates cleavage of single-stranded RNA having sequence complementary to the antisense strand of the siRNA duplex. Cleavage of the target RNA takes place in the middle of the region complementary to the antisense strand of the siRNA duplex (Elbashir et al., 2001, Genes Dev., 15, 188).

[0220] RNAi has been studied in a variety of systems. Fire et al., 1998, Nature, 391, 806, were the first to observe RNAi in *C. elegans*. Bahramian and Zarbl, 1999, Molecular and Cellular Biology, 19, 274-283 and Wianny and Goetz, 1999, Nature Cell Biol., 2, 70, describe RNAi mediated by dsRNA in mammalian systems. Elbashir et al., 2001, Nature, 411, 494 and Tuschl et al., WO0175164, describe RNAi induced by introduction of duplexes of synthetic 21-nucleotide RNAs in cultured mammalian cells including human embryonic kidney and HeLa cells. Recent work (Elbashir et al., 2001, EMBO J., 20, 6877 and Tuschl et al., WO0175164) has revealed certain requirements for siRNA length, structure, chemical composition, and sequence that are essential to mediate efficient RNAi activity.

[0221] Nucleic acid molecules (for example comprising structural features as disclosed herein) may inhibit or down regulate gene expression or viral replication by mediating RNA interference "RNAi" or gene silencing in a sequence-specific manner. (See, e.g., Zamore et al., 2000, Cell, 101, 25-33; Bass, 2001, Nature, 411, 428-429; Elbashir et al., 2001, Nature, 411, 494-498; and Kreutzer et al., WO0044895; Zemicka-Goetz et al., WO0136646; Fire, WO9932619; Plaetinck et al., WO0001846; Mello and Fire, WO0129058; Deschamps-Depaillette, WO9907409; and Li et al., WO0044914; Allshire, 2002, Science, 297, 1818-1819; Volpe et al., 2002, Science, 297, 1833-1837; Jenuwein, 2002, Science, 297, 2215-2218; and Hall et al., 2002, Science, 297, 2232-2237; Hutvagner and Zamore, 2002, Science, 297, 2056-60; McManus et al., 2002, RNA, 8, 842-850; Reinhart et al., 2002, Gene & Dev., 16, 1616-1626; and Reinhart & Bartel, 2002, Science, 297, 1831)

[0222] An siNA nucleic acid molecule can be assembled from two separate polynucleotide strands, where one strand is the sense strand and the other is the antisense strand in which the antisense and sense strands are self-complementary (i.e. each strand includes nucleotide sequence that is complementary to nucleotide sequence in the other strand); such as where the antisense strand and sense strand form a duplex or double-stranded structure having any length and structure as described herein for nucleic acid molecules as provided, for example wherein the double-stranded region (duplex region) is about 15 to about 49 (e.g., about 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, or 49 base pairs); the antisense strand includes nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule (i.e., hsp47 mRNA) or a portion thereof and the sense strand includes nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof (e.g., about 17 to about 49 or more nucleotides of the nucleic acid molecules herein are complementary to the target nucleic acid or a portion thereof).

[0223] In certain aspects and embodiments, a nucleic acid molecule (e.g., a siNA molecule) provided herein may be a "RISC length" molecule or may be a Dicer substrate as described in more detail below.

[0224] An siNA nucleic acid molecule may include separate sense and antisense sequences or regions, where the sense and antisense regions are covalently linked by nucleotide or non-nucleotide linkers molecules as is known in the art, or are alternately non-covalently linked by ionic interactions, hydrogen bonding, van der Waals interactions, hydrophobic interactions, and/or stacking interactions. Nucleic acid molecules may include a nucleotide sequence that is complementary to nucleotide sequence of a target gene. Nucleic acid molecules may interact with nucleotide sequence of a target gene in a manner that causes inhibition of expression of the target gene.

[0225] Alternatively, an siNA nucleic acid molecule is assembled from a single polynucleotide, where the self-complementary sense and antisense regions of the nucleic acid molecules are linked by means of a nucleic acid based or non-nucleic acid-based linker(s), i.e., the antisense strand and the sense strand are part of one single polynucleotide that having an antisense region and sense region that fold to form a duplex region (for example to form a "hairpin" structure as is well known in the art). Such siNA nucleic acid molecules can be a polynucleotide with a duplex, asymmetric duplex, hairpin or asymmetric hairpin secondary structure, having self-complementary sense and antisense regions, wherein the antisense region includes nucleotide sequence that is complementary to nucleotide sequence in a separate target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence (e.g., a sequence of hsp47 mRNA). Such siNA nucleic acid molecules can be a circular single-stranded polynucleotide having two or more loop structures and a stem comprising self-complementary sense and antisense regions, wherein the antisense region includes nucleotide sequence that is complementary to nucleotide sequence in a target nucleic acid molecule or a portion thereof and the sense region having nucleotide sequence corresponding to the target nucleic acid sequence or a portion thereof, and wherein the circular polynucleotide can be processed either *in vivo* or *in vitro* to generate an active nucleic acid molecule capable of mediating RNAi.

[0226] The following nomenclature is often used in the art to describe lengths and overhangs of siNA molecules and may be used throughout the specification and Examples. In all descriptions of oligonucleotides herein, the identification of nucleotides in a sequence is given in the 5' to 3' direction for both sense and antisense strands. Names given to duplexes indicate the length of the oligomers and the presence or absence of overhangs. For example, a "21+2" duplex contains two nucleic acid strands both of which are 21 nucleotides in length, also termed a 21-mer siRNA duplex or a 21-mer nucleic acid and having a 2 nucleotides 3'-overhang. A "21-2" design refers to a 21-mer nucleic acid duplex with a 2

nucleotides 5'-overhang. A 21-0 design is a 21-mer nucleic acid duplex with no overhangs (blunt). A "21+2UU" is a 21-mer duplex with 2-nucleotides 3'-overhang and the terminal 2 nucleotides at the 3'-ends are both U residues (which may result in mismatch with target sequence). The aforementioned nomenclature can be applied to siNA molecules of various lengths of strands, duplexes and overhangs (such as 19-0, 21+2, 27+2, and the like). In an alternative but similar nomenclature, a "25/27" is an asymmetric duplex having a 25 base sense strand and a 27 base antisense strand with a 2-nucleotides 3'-overhang. A "27/25" is an asymmetric duplex having a 27 base sense strand and a 25 base antisense strand.

Chemical Modifications

[0227] In certain aspects and embodiments, nucleic acid molecules (e.g., siNA molecules) as provided herein include one or more modifications (or chemical modifications). In certain embodiments, such modifications include any changes to a nucleic acid molecule or polynucleotide that would make the molecule different than a standard ribonucleotide or RNA molecule (i.e., that includes standard adenosine, cytosine, uracil, or guanosine moieties); which may be referred to as an "unmodified" ribonucleotide or unmodified ribonucleic acid. Traditional DNA bases and polynucleotides having a 2'-deoxy sugar represented by adenosine, cytosine, thymine, or guanosine moieties may be referred to as an "unmodified deoxyribonucleotide" or "unmodified deoxyribonucleic acid"; accordingly, the term "unmodified nucleotide" or "unmodified nucleic acid" as used herein refers to an "unmodified ribonucleotide" or "unmodified ribonucleic acid" unless there is a clear indication to the contrary. Such modifications can be in the nucleotide sugar, nucleotide base, nucleotide phosphate group and/or the phosphate backbone of a polynucleotide.

[0228] In certain embodiments, modifications as disclosed herein may be used to increase RNAi activity of a molecule and/or to increase the *in vivo* stability of the molecules, particularly the stability in serum, and/or to increase bioavailability of the molecules. Non-limiting examples of modifications include internucleotide or internucleoside linkages; deoxynucleotides or dideoxynucleotides at any position and strand of the nucleic acid molecule; nucleic acid (e.g., ribonucleic acid) with a modification at the 2'-position preferably selected from an amino, fluoro, methoxy, alkoxy and alkyl; 2'-deoxyribonucleotides, 2'OMe ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl nucleotides, biotin group, and terminal glyceryl and/or inverted deoxy abasic residue incorporation, sterically hindered molecules, such as fluorescent molecules and the like. Other nucleotide modifiers could include 3'-deoxyadenosine (cordycepin), 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxyinosine (ddI), 2',3'-dideoxy-3'-thiacytidine (3TC), 2',3'-didehydro-2',3'-dideoxythymidine (d4T) and the monophosphate nucleotides of 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxy-3'-thiacytidine (3TC) and 2',3'-didehydro-2',3'-dideoxythymidine (d4T). Further details on various modifications are described in more detail below.

[0229] Modified nucleotides include those having a Northern conformation (e.g., Northern pseudorotation cycle, see for example Sanger, Principles of Nucleic Acid Structure, Springer-Verlag ed., 1984). Non-limiting examples of nucleotides having a northern configuration include LNA nucleotides (e.g., 2'-O, 4'-C-methylene-(D-ribofuranosyl) nucleotides); 2'-methoxyethoxy (MOE) nucleotides; 2'-methyl-thio-ethyl, 2'-deoxy-2'-fluoro nucleotides, 2'-deoxy-2'-chloro nucleotides, 2'-azido nucleotides, and 2'OMe nucleotides. LNAs are described, for example, in Elman *et al.*, 2005; Kurreck *et al.*, 2002; Crinelli *et al.*, 2002; Braasch and Corey, 2001; Bondensgaard *et al.*, 2000; Wahlestedt *et al.*, 2000; and WO0047599, WO9914226, WO9839352, and WO04083430. In one embodiment, an LNA is incorporated at the 5'-terminus of the sense strand.

[0230] Chemical modifications also include UNAs, which are non-nucleotide, acyclic analogues, in which the C2'-C3' bond is not present (although UNAs are not truly nucleotides, they are expressly included in the scope of "modified" nucleotides or modified nucleic acids as contemplated herein). In particular embodiments, nucleic acid molecules with an overhang may be modified to have UNAs at the overhang positions (i.e., 2 nucleotide overhang). In other embodiments, UNAs are included at the 3'- or 5'- ends. A UNA may be located anywhere along a nucleic acid strand, i.e. in position 7. Nucleic acid molecules may contain one or more than UNA. Exemplary UNAs are disclosed in Nucleic Acids Symposium Series No. 52 p. 133-134 (2008). In certain embodiments, nucleic acid molecules (e.g., siNA molecules) as described herein, include one or more UNAs; or one UNA. In some embodiments, a nucleic acid molecule (e.g., a siNA molecule) as described herein that has a 3'-overhang include one or two UNAs in the 3' overhang. In some embodiments, a nucleic acid molecule (e.g., a siNA molecule) as described herein includes a UNA (for example one UNA) in the antisense strand; for example in position 6 or position 7 of the antisense strand. Chemical modifications also include non-pairing nucleotide analogs, for example as disclosed herein. Chemical modifications further include unconventional moieties as disclosed herein.

[0231] Chemical modifications also include terminal modifications on the 5' and/or 3' part of the oligonucleotides and are also known as capping moieties. Such terminal modifications are selected from a nucleotide, a modified nucleotide, a lipid, a peptide, and a sugar.

[0232] Chemical modifications also include six membered "six membered ring nucleotide analogs." Examples of six-membered ring nucleotide analogs are disclosed in Allart, et al (Nucleosides & Nucleotides, 1998, 17:1523-1526.; and Perez-Perez, et al., 1996, Bioorg. and Medicinal Chem Letters 6:1457-1460) Oligonucleotides including 6-membered ring nucleotide analogs including hexitol and altritol nucleotide monomers are disclosed in WO2006047842.

[0233] Chemical modifications also include "mirror" nucleotides which have a reversed chirality as compared to normal naturally occurring nucleotide; that is, a mirror nucleotide may be an "L-nucleotide" analogue of naturally occurring D-nucleotide (see US6602858). Mirror nucleotides may further include at least one sugar or base modification and/or a backbone modification, for example, as described herein, such as a phosphorothioate or phosphonate moiety. US6602858 discloses nucleic acid catalysts including at least one L-nucleotide substitution. Mirror nucleotides include for example L-DNA (L-deoxyriboadenosine-3'-phosphate (mirror dA); L-deoxyribocytidine-3'-phosphate (mirror dC); L-deoxyriboguanosine-3'-phosphate (mirror dG); L-deoxyribothymidine-3'-phosphate (mirror image dT)) and L-RNA (L-riboadenosine-3'-phosphate (mirror rA); L-ribocytidine-3'-phosphate (mirror rC); L-riboguanosine-3'-phosphate (mirror rG); L-ribouracil-3'-phosphate (mirror dU).

[0234] In some embodiments, modified ribonucleotides include modified deoxyribonucleotides, for example 5'OMe DNA (5-methyl-deoxyriboguanosine-3'-phosphate) which may be useful as a nucleotide in the 5' terminal position (position number 1); PACE (deoxyriboadenosine 3' phosphonoacetate, deoxyribocytidine 3' phosphonoacetate, deoxyriboguanosine 3' phosphonoacetate, deoxyribothymidine 3' phosphonoacetate).

[0235] Modifications may be present in one or more strands of a nucleic acid molecule disclosed herein, *e.g.*, in the sense strand, the antisense strand, or both strands. In certain embodiments, the antisense strand may include modifications and the sense strand may only include unmodified RNA.

Nucleobases

[0236] Nucleobases of the nucleic acid disclosed herein may include unmodified ribonucleotides (purines and pyrimidines) such as adenine, guanine, cytosine, uracil. The nucleobases in one or both strands can be modified with natural and synthetic nucleobases such as, thymine, xanthine, hypoxanthine, inosine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, any "universal base" nucleotides; 2-propyl and other alkyl derivatives of adenine and guanine, 5-halo uracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudo-uracil), 4-thiouracil, 8-halo, amino, thiol, thioalkyl, hydroxyl and other 8-substituted adenines and guanines, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine, deazapurines, heterocyclic substituted analogs of purines and pyrimidines, *e.g.*, aminoethoxy phenoxazine, derivatives of purines and pyrimidines (*e.g.*, 1-alkyl-, 1-alkenyl-, heteroaromatic- and 1-alkynyl derivatives) and tautomers thereof, 8-oxo-N6-methyladenine, 7-diazaxanthine, 5-methylcytosine, 5-methyluracil, 5-(1-propynyl)uracil, 5-(1-propynyl) cytosine and 4,4-ethanocytosine). Other examples of suitable bases include non-puriny and non-pyrimidiny bases such as 2-aminopuridine and triazines.

Sugar moieties

[0237] Sugar moieties in a nucleic acid disclosed herein may include 2'-hydroxyl-pentofuranosyl sugar moiety without any modification. Alternatively, sugar moieties can be modified such as, 2'-deoxy-pentofuranosyl sugar moiety, D-ribose, hexose, modification at the 2' position of the pentofuranosyl sugar moiety such as 2'-O-alkyl (including 2'OMe and 2'-O-ethyl), i.e., 2'-alkoxy, 2'-amino, 2'-O-allyl, 2'-S-alkyl, 2'-halogen (including 2'-fluoro, chloro, and bromo), 2'-methoxyethoxy, 2'-O-methoxyethyl, 2'-O-2-methoxyethyl, 2'-allyloxy ($-\text{OCH}_2\text{CH}=\text{CH}_2$), 2'-propargyl, 2'-propyl, ethynyl, ethenyl, propenyl, CF, cyano, imidazole, carboxylate, thioate, C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl, OCF₃, OCN, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃; SO₂CH₃; ONO₂; NO₂, N₃; heterozycloalkyl; heterozycloalkaryl; aminoalkylamino; polyalkylamino or substituted silyl, as, among others, for example as described in European patents EP0586520 or EP0618925.

[0238] Alkyl group includes saturated aliphatic groups, including straight-chain alkyl groups (*e.g.*, methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, etc.), branched-chain alkyl groups (isopropyl, tert-butyl, isobutyl, etc.), cycloalkyl (alicyclic) groups (cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl), alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. In certain embodiments, a straight chain or branched chain alkyl has six or fewer carbon atoms in its backbone (*e.g.*, C₁-C₆ for straight chain, C₃-C₆ for branched chain), and more preferably four or fewer. Likewise, preferred cycloalkyls may have from three to eight carbon atoms in their ring structure, and more preferably have five or six carbons in the ring structure. The term C₁-C₆ includes alkyl groups containing one to six carbon atoms. The alkyl group can be substituted alkyl group such as alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinate, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamide, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0239] Alkoxy group includes substituted and unsubstituted alkyl, alkenyl, and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups include methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. Examples of substituted alkoxy groups include halogenated alkoxy groups. The alkoxy groups can be substituted with groups such as alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinate, cyano, amino (including alkyl amino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarbonylate, sulfates, alkylsulfanyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkaryl, or an aromatic or heteroaromatic moieties. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy, trichloromethoxy, etc.

[0240] In some embodiments, the pentafuronosyl ring may be replaced with acyclic derivatives lacking the C₂'-C₃'-bond of the pentafuronosyl ring. For example, acyclonucleotides may substitute a 2-hydroxyethoxymethyl group for the 2'-deoxyribofuranosyl sugar normally present in dNMPs.

[0241] Halogens include fluorine, bromine, chlorine, iodine.

Backbone

[0242] The nucleoside subunits of the nucleic acid disclosed herein may be linked to each other by a phosphodiester bond. The phosphodiester bond may be optionally substituted with other linkages. For example, phosphorothioate, thiophosphate-D-ribose entities, triester, thioate, 2'-5' bridged backbone (may also be referred to as 5'-2' or 2'-5'-nucleotide or 2'-5'-ribonucleotide), PACE, 3'(or -5')-deoxy-3'(or -5')-thiophosphorothioate, phosphorodithioate, phosphoroselenates, 3'(or -5')-deoxy phosphinates, borano phosphates, 3'(or -5')-deoxy-3'(or 5')-amino

or other polymeric compounds (e.g. polyethylene glycols such as those having between 2 and 100 ethylene glycol units). Specific examples include those described by Seela and Kaiser, *Nucleic Acids Res.* 1990, 18:6353 and *Nucleic Acids Res.* 1987, 15:3113; Cload and Schepartz, *J. Am. Chem. Soc.* 1991, 113:6324; Richardson and Schepartz, *J. Am. Chem. Soc.* 1991, 113:5109; Ma et al., *Nucleic Acids Res.* 1993, 21:2585 and *Biochemistry* 1993, 32:1751; Durand et al., *Nucleic Acids Res.* 1990, 18:6353; McCurdy et al., *Nucleosides & Nucleotides* 1991, 10:287; Jschke et al., *Tetrahedron Lett.* 1993, 34:301; Ono et al., *Biochemistry* 1991, 30:9914; Arnold et al., WO8902439; Usman et al., WO9506731; Dudycz et al., WO9511910 and Ferentz and Verdine, *J. Am. Chem. Soc.* 1991, 113:4000.

5'-ends, 3'-ends, and Overhangs

[0249] Nucleic acid molecules disclosed herein (e.g., siNA molecules) may be blunt-ended on both sides, have overhangs on both sides, or a combination of blunt and overhang ends. Overhangs may occur on either the 5'- or 3'-end of the sense or antisense strand.

[0250] 5'- and/or 3'- ends of double-stranded nucleic acid molecules (e.g., siNA) may be blunt ended or have an overhang. The 5'-end may be blunt ended and the 3'-end has an overhang in either the sense strand or the antisense strand. In other embodiments, the 3'-end may be blunt ended and the 5'-end has an overhang in either the sense strand or the antisense strand. In yet other embodiments, both the 5'- and 3'-end are blunt ended or both the 5'- and 3'-ends have overhangs.

[0251] The 5'- and/or 3'-end of one or both strands of the nucleic acid may include a free hydroxyl group. The 5'- and/or 3'-end of any nucleic acid molecule strand may be modified to include a chemical modification. Such modification may stabilize nucleic acid molecules, e.g., the 3'-end may have increased stability due to the presence of the nucleic acid molecule modification. Examples of end modifications (e.g., terminal caps) include, but are not limited to, abasic, deoxy abasic, inverted (deoxy) abasic, glyceryl, dinucleotide, acyclic nucleotide, amino, fluoro, chloro, bromo, CN, CF, methoxy, imidazole, carboxylate, thioate, C₁ to C₁₀ lower alkyl, substituted lower alkyl, alkaryl or aralkyl, OCF₃, OCN, O-, S-, or N-alkyl; O-, S-, or N-alkenyl; SOCH₃; SO₂CH₃; ONO₂; NO₂; N₃; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino or substituted silyl, as, among others, described in European patents EP586520 and EP618925 and other modifications disclosed herein.

[0252] Nucleic acid molecules include those with blunt ends, i.e., ends that do not include any overhanging nucleotides. A nucleic acid molecule can include one or more blunt ends. The blunt ended nucleic acid molecule has a number of base pairs equal to the number of nucleotides present in each strand of the nucleic acid molecule. The nucleic acid molecule can include one blunt end, for example where the 5'-end of the antisense strand and the 3'-end of the sense strand do not have any overhanging nucleotides. Nucleic acid molecule may include one blunt end, for example where the 3'-end of the antisense strand and the 5'-end of the sense strand do not have any overhanging nucleotides. A nucleic acid molecule may include two blunt ends, for example where the 3'-end of the antisense strand and the 5'-end of the sense strand as well as the 5'-end of the antisense strand and 3'-end of the sense strand do not have any overhanging nucleotides. Other nucleotides present in a blunt ended nucleic acid molecule can include, for example, mismatches, bulges, loops, or wobble base pairs to modulate the activity of the nucleic acid molecule to mediate RNA interference.

[0253] In certain embodiments of the nucleic acid molecules (e.g., siNA molecules) provided herein, at least one end of the molecule has an overhang of at least one nucleotide (for example one to eight overhang nucleotides). For example, one or both strands of a double-stranded nucleic acid molecule disclosed herein may have an overhang at the 5'-end or at the 3'-end or both. An overhang may be present at either or both the sense strand and antisense strand of the nucleic acid molecule. The length of the overhang may be as little as one nucleotide and as long as one to eight or more nucleotides (e.g., 1, 2, 3, 4, 5, 6, 7 or 8 nucleotides; in some preferred embodiments an overhang is 2, 3, 4, 5, 6, 7 or 8 nucleotides; for example an overhang may be 2 nucleotides. The nucleotide(s) forming the overhang may include deoxyribonucleotide(s), ribonucleotide(s), natural and non-natural nucleobases or any nucleotide modified in the sugar, base or phosphate group such as disclosed herein. A double-stranded nucleic acid molecule may have both 5'- and 3'-overhangs. The overhangs at the 5'- and 3'-end may be of different lengths. An overhang may include at least one nucleic acid modification which may be deoxyribonucleotide. One or more deoxyribonucleotides may be at the 5'-terminal. The 3'-end of the respective counter-strand of the nucleic acid molecule may not have an overhang, more preferably not a deoxyribonucleotide overhang. The one or more deoxyribonucleotide may be at the 3'-terminal. The 5'-end of the respective counter-strand of the dsRNA may not have an overhang, more preferably not a deoxyribonucleotide overhang. The overhang in either the 5'- or the 3'-end of a strand may be one to eight (e.g., about 1, 2, 3, 4, 5, 6, 7 or 8) unpaired nucleotides, preferably, the overhang is two to three unpaired nucleotides; more preferably two unpaired nucleotides. Nucleic acid molecules may include duplex nucleic acid molecules with overhanging ends of about 1 to about 20 (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20); preferably one to eight (e.g., about 1, 2, 3, 4, 5, 6, 7 or 8) nucleotides, for example, about 21-nucleotide duplexes with about 19 base pairs and 3'-terminal mononucleotide, dinucleotide, or trinucleotide overhangs. Nucleic acid molecules herein may include duplex nucleic acid molecules with blunt ends, where both ends are blunt, or alternatively, where one of the ends is blunt. Nucleic acid molecules disclosed herein can include one or more blunt ends, i.e. where a blunt end does not have any overhanging nucleotides. In one embodiment, the blunt ended nucleic acid molecule has a number of base pairs equal to the number of nucleotides present in each strand of the nucleic acid molecule. The nucleic acid molecule may include one blunt end, for example where the 5'-end of the antisense strand and the 3'-end of the sense strand do not have any overhanging nucleotides. The nucleic acid molecule may include one blunt end, for example where the 3'-end of the antisense strand and the 5'-end of the sense strand do not have any overhanging nucleotides. A nucleic acid molecule may include two blunt ends, for example where the 3'-end of the antisense strand and the 5'-end of the sense strand as well as the 5'-end of the antisense strand and 3'-end of the sense strand do not have any overhanging nucleotides. In certain preferred embodiments the nucleic acid compounds are blunt ended. Other nucleotides present in a blunt ended siNA molecule can include, for example, mismatches, bulges, loops, or wobble base pairs to modulate the activity of the nucleic acid molecule to mediate RNA interference.

[0254] In many embodiments one or more, or all, of the overhang nucleotides of a nucleic acid molecule (e.g., a siNA molecule) as described herein includes are modified such as described herein; for example one or more, or all, of the nucleotides may be 2'-deoxynucleotides.

Amount, Location and Patterns of Modifications

[0255] Nucleic acid molecules (e.g., siNA molecules) disclosed herein may include modified nucleotides as a percentage of the total number of nucleotides present in the nucleic acid molecule. As such, a nucleic acid molecule may include about 5% to about 100% modified nucleotides (e.g., about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% modified nucleotides). The actual percentage of modified nucleotides present in a given nucleic acid molecule will depend on the total number of nucleotides present in the nucleic acid. If the nucleic acid molecule is single-stranded, the percent modification can be based upon the total number of nucleotides present in the single-stranded nucleic acid molecule. Likewise, if the nucleic acid molecule is double-stranded, the percent modification can be based upon the total number of nucleotides present in the sense strand, antisense strand, or both the sense and antisense strands.

[0256] Nucleic acid molecules disclosed herein may include unmodified RNA as a percentage of the total nucleotides in the nucleic acid molecule. As such, a nucleic acid molecule may include about 5% to about 100% modified nucleotides (e.g., about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or 100% of total nucleotides present in a nucleic acid molecule).

[0257] A nucleic acid molecule (e.g., an siNA molecule) may include a sense strand that includes about one to about five, specifically about 1, 2, 3, 4, or 5 phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, or more) 2'-deoxy, 2'OMe, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the sense strand; and wherein the antisense strand includes about one to about five or more, specifically about 1, 2, 3, 4, 5, or more phosphorothioate internucleotide linkages, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) 2'-deoxy, 2'OMe, 2'-deoxy-2'-fluoro, and/or one or more (e.g., about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more) universal base modified nucleotides, and optionally a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of the antisense strand. A nucleic acid molecule may include about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more, pyrimidine nucleotides of the sense and/or antisense nucleic acid strand are chemically-modified with 2'-deoxy, 2'OMe and/or 2'-deoxy-2'-fluoro nucleotides, with or without about one to about five or more, for example about 1, 2, 3, 4, 5, or more phosphorothioate internucleotide linkages and/or a terminal cap molecule at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends, being present in the same or different strand.

[0258] A nucleic acid molecule may include about one to about five or more (specifically about 1, 2, 3, 4, 5, or more) phosphorothioate internucleotide linkages in each strand of the nucleic acid molecule.

[0259] A nucleic acid molecule may include 2'-5' internucleotide linkages, for example at the 3'-end, the 5'-end, or both of the 3'- and 5'-ends of one or both nucleic acid sequence strands. In addition, the 2'-5' internucleotide linkage(s) can be present at various other positions within one or both nucleic acid sequence strands, for example, about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a pyrimidine nucleotide in one or both strands of the siNA molecule can include a 2'-5' internucleotide linkage, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more including every internucleotide linkage of a purine nucleotide in one or both strands of the siNA molecule can include a 2'-5' internucleotide linkage.

[0260] A chemically-modified short interfering nucleic acid (siNA) molecule may include an antisense region, wherein any (e.g., one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (e.g., one or more or all) purine nucleotides present in the antisense region are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides).

[0261] A chemically-modified short interfering nucleic acid (siNA) molecule may include an antisense region, wherein any (e.g., one or more or all) pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and wherein any (e.g., one or more or all) purine nucleotides present in the antisense region are 2'OMe purine nucleotides (e.g., wherein all purine nucleotides are 2'OMe purine nucleotides or alternately a plurality of purine nucleotides are 2'OMe purine nucleotides).

[0262] A chemically-modified siNA molecule capable of mediating RNA interference (RNAi) against hsp47 inside a cell or reconstituted *in vitro* system may include a sense region, wherein one or more pyrimidine nucleotides present in the sense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more purine nucleotides present in the sense region are 2'-deoxy purine nucleotides (e.g., wherein all purine nucleotides are 2'-deoxy purine nucleotides or alternately a plurality of purine nucleotides are 2'-deoxy purine nucleotides), and an antisense region, wherein one or more pyrimidine nucleotides present in the antisense region are 2'-deoxy-2'-fluoro pyrimidine nucleotides (e.g., wherein all pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides or alternately a plurality of pyrimidine nucleotides are 2'-deoxy-2'-fluoro pyrimidine nucleotides), and one or more purine nucleotides present in the antisense region are 2'OMe purine nucleotides (e.g., wherein all purine nucleotides are 2'OMe purine nucleotides or alternately a plurality of purine nucleotides are 2'OMc purine nucleotides). The sense region and/or the antisense region can have a terminal cap modification, such as any modification that is optionally present at the 3'-end, the 5'-end, or both of the 3' and 5'-ends of the sense and/or antisense sequence. The sense and/or antisense region can optionally further include a 3'-terminal nucleotide overhang having about 1 to about 4 (e.g., about 1, 2, 3, or 4) 2'-deoxynucleotides. The overhang nucleotides can further include one or more (e.g., about 1, 2, 3, 4 or more) phosphorothioate, phosphonoacetate, and/or thiophosphonoacetate internucleotide linkages. The purine nucleotides in the sense region may alternatively be 2'OMe purine nucleotides (e.g., wherein all purine nucleotides are 2'OMe purine nucleotides or alternately a plurality of purine nucleotides are 2'OMe purine nucleotides) and one or more purine nucleotides present in the antisense region are 2'OMe purine nucleotides (e.g., wherein all purine nucleotides are 2'OMe purine nucleotides or alternately a plurality of purine nucleotides are 2'OMe purine nucleotides). One or more purine nucleotides in the sense region may alternatively be purine ribonucleotides (e.g., wherein all purine nucleotides are purine ribonucleotides or alternately a plurality of purine nucleotides are purine ribonucleotides) and any purine nucleotides present in the antisense region are 2'OMe purine nucleotides (e.g., wherein all purine nucleotides are 2'OMe purine nucleotides or alternately a plurality of purine nucleotides are 2'OMe purine nucleotides). One or more purine nucleotides in the sense region and/or present in the antisense region may alternatively selected from the group consisting of 2'-deoxy nucleotides, LNA nucleotides, 2'-methoxyethyl nucleotides,

4'-thionucleotides, and 2'OMe nucleotides (e.g., wherein all purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, LNA nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'OMe nucleotides or alternately a plurality of purine nucleotides are selected from the group consisting of 2'-deoxy nucleotides, LNA nucleotides, 2'-methoxyethyl nucleotides, 4'-thionucleotides, and 2'OMc nucleotides).

[0263] In some embodiments, a nucleic acid molecule (e.g., a siNA molecule) as described herein includes a modified nucleotide (for example one modified nucleotide) in the antisense strand; for example in position 6 or position 7 of the antisense strand.

Modification Patterns and Alternating Modifications

[0264] Nucleic acid molecules (e.g., siNA molecules) provided herein may have patterns of modified and unmodified nucleic acids. A pattern of modification of the nucleotides in a contiguous stretch of nucleotides may be a modification contained within a single nucleotide or group of nucleotides that are covalently linked to each other via standard phosphodiester bonds or, at least partially, through phosphorothioate bonds. Accordingly, a "pattern" as contemplated herein, does not necessarily need to involve repeating units, although it may. Examples of modification patterns that may be used in conjunction with the nucleic acid molecules (e.g., siNA molecules) provided herein include those disclosed in Giese, US7452987. For example, nucleic acid molecules (e.g., siNA molecules) provided herein include those with modification patterns such as, similar to, or the same as, the patterns shown diagrammatically in Figure 2 of US7452987.

[0265] A modified nucleotide or group of modified nucleotides may be at the 5'-end or 3'-end of the sense or antisense strand, a flanking nucleotide or group of nucleotides is arrayed on both sides of the modified nucleotide or group, where the flanking nucleotide or group either is unmodified or does not have the same modification of the preceding nucleotide or group of nucleotides. The flanking nucleotide or group of nucleotides may, however, have a different modification. This sequence of modified nucleotide or group of modified nucleotides, respectively, and unmodified or differently modified nucleotide or group of unmodified or differently modified nucleotides may be repeated one or more times.

[0266] In some patterns, the 5'-terminal nucleotide of a strand is a modified nucleotide while in other patterns the 5'-terminal nucleotide of a strand is an unmodified nucleotide. In some patterns, the 5'-end of a strand starts with a group of modified nucleotides while in other patterns, the 5'-terminal end is an unmodified group of nucleotides. This pattern may be either on the first stretch or the second stretch of the nucleic acid molecule or on both.

[0267] Modified nucleotides of one strand of the nucleic acid molecule may be complementary in position to the modified or unmodified nucleotides or groups of nucleotides of the other strand.

[0268] There may be a phase shift between modifications or patterns of modifications on one strand relative to the pattern of modification of the other strand such that the modification groups do not overlap. In one instance, the shift is such that the modified group of nucleotides of the sense strand corresponds to the unmodified group of nucleotides of the antisense strand and vice versa.

[0269] There may be a partial shift of the pattern of modification such that the modified groups overlap. The groups of modified nucleotides in any given strand may optionally be the same length, but may be of different lengths. Similarly, groups of unmodified nucleotides in any given strand may optionally be the same length, or of different lengths.

[0270] In some patterns, the second (penultimate) nucleotide at the terminus of the strand, is an unmodified nucleotide or the beginning of group of unmodified nucleotides. Preferably, this unmodified nucleotide or unmodified group of nucleotides is located at the 5'-end of the either or both the sense and antisense strands and even more preferably at the terminus of the sense strand. An unmodified nucleotide or unmodified group of nucleotide may be located at the 5'-end of the sense strand. In a preferred embodiment the pattern consists of alternating single modified and unmodified nucleotides.

[0271] In some double-stranded nucleic acid molecules include a 2'OMe modified nucleotide and a non-modified nucleotide, preferably a nucleotide which is not 2'OMe modified, are incorporated on both strands in an alternating fashion, resulting in a pattern of alternating 2'OMe modified nucleotides and nucleotides that are either unmodified or at least do not include a 2'OMe modification. In certain embodiments, the same sequence of 2'OMe modification and non-modification exists on the second strand; in other embodiments the alternating 2'OMe modified nucleotides are only present in the sense strand and are not present in the antisense strand; and in yet other embodiments the alternating 2'OMe modified nucleotides are only present in the sense strand and are not present in the antisense strand. In certain embodiments, there is a phase shift between the two strands such that the 2'OMe modified nucleotide on the first strand base pairs with a non-modified nucleotide(s) on the second strand and vice versa. This particular arrangement, i.e. base pairing of 2'OMe modified and non-modified nucleotide(s) on both strands is particularly preferred in certain embodiments. In certain embodiments, the pattern of alternating 2'OMe modified nucleotides exists throughout the entire nucleic acid molecule; or the entire duplex region. In other embodiments the pattern of alternating 2'OMe modified nucleotides exists only in a portion of the nucleic acid; or the entire duplex region.

[0272] In "phase shift" patterns, it may be preferred if the antisense strand starts with a 2'OMe modified nucleotide at the 5'-end whereby consequently the second nucleotide is non-modified, the third, fifth, seventh and so on nucleotides are thus again 2'OMe modified whereas the second, fourth, sixth, eighth and the like nucleotides are non-modified nucleotides.

Modification Locations and Patterns

[0273] While exemplary patterns are provided in more detail below, all permutations of patterns with of all possible characteristics of the nucleic acid molecules disclosed herein and those known in the art are contemplated (e.g., characteristics include, but are not limited to, length of sense

strand, length of antisense strand, length of duplex region, length of hangover, whether one or both ends of a double-stranded nucleic acid molecule is blunt or has an overhang, location of modified nucleic acid, number of modified nucleic acids, types of modifications, whether a double overhang nucleic acid molecule has the same or different number of nucleotides on the overhang of each side, whether a one or more than one type of modification is used in a nucleic acid molecule, and number of contiguous modified/unmodified nucleotides). With respect to all detailed examples provided below, while the duplex region is shown to be 19 nucleotides, the nucleic acid molecules provided herein can have a duplex region ranging from 1 to 49 nucleotides in length as each strand of a duplex region can independently be 17-49 nucleotides in length Exemplary patterns are provided herein.

[0274] Nucleic acid molecules may have a blunt end (when n is 0) on both ends that include a single or contiguous set of modified nucleic acids. The modified nucleic acid may be located at any position along either the sense or antisense strand. Nucleic acid molecules may include a group of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48 or 49 contiguous modified nucleotides. Modified nucleic acids may make up 1%, 2%, 3%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 100% of a nucleic acid strand. Modified nucleic acids of the examples immediately below may be in the sense strand only, the antisense strand only, or in both the sense and antisense strand.

[0275] General nucleic acid patters are shown below where X = sense strand nucleotide in the duplex region; X_a = 5'-overhang nucleotide in the sense strand; X_b = 3'-overhang nucleotide in the sense strand; Y = antisense strand nucleotide in the duplex region; Y_a = 3'-overhang nucleotide in the antisense strand; Y_b = 5'-overhang nucleotide in the antisense strand; and M = a modified nucleotide in the duplex region. Each a and b are independently 0 to 8 (e.g., 0, 1, 2, 3, 4, 5, 6, 7 or 8). Each X, Y, a and b are independently modified or unmodified. The sense and antisense strands can are each independently 17-49 nucleotides in length. The examples provided below have a duplex region of 19 nucleotides; however, nucleic acid molecules disclosed herein can have a duplex region anywhere between 17 and 49 nucleotides and where each strand is independently between 17 and 49 nucleotides in length.

```
5'  XaXXXXXXXXXXXXXXXXXXXXXb
3'  YbYYYYYYYYYYYYYYYYYYYa
```

[0276] Further exemplary nucleic acid molecule patterns are shown below where X = unmodified sense strand nucleotides; x = an unmodified overhang nucleotide in the sense strand; Y = unmodified antisense strand nucleotides; y = an unmodified overhang nucleotide in the antisense strand; and M = a modified nucleotide. The sense and antisense strands can are each independently 17-49 nucleotides in length. The examples provided below have a duplex region of 19 nucleotides; however, nucleic acid molecules disclosed herein can have a duplex region anywhere between 17 and 49 nucleotides and where each strand is independently between 17 and 49 nucleotides in length.

```
5'  MaXXXXXXXXXXMXXXXXXXXXb
3'  MaYYYYYYYYYYYYYYYYYYb
```

```
5'  XXXXXXXXXXXXXXXXXXXX
3'  YYYYYYYYMYYYYYYYY
```

```
5'  XXXXXXXXMMXXXXXXXXX
3'  YYYYYYYYYYYYYYYYYY
```

```
5'  XXXXXXXXXXXXXXXXXXXX
3'  YYYYYYYYMMYYYYYYY
```

```
5'  XXXXXXXXMXXXXXXXXXX
3'  YYYYYYYYMYYYYYYYY
```

```
5'  XXXXMXXXXXXXXXXXXXX
3'  YYYYYYYYMYYYYYYYY
```

```
5'  MXXXXXXXXXXXXXXXXXX
3'  YYYYYYYYMYYYYYY
```

```
5'  XXXXXXXXXXXXXXXXXXXM
3'  YYYMYYYYYYYYYYYY
```

```
5'  XXXXXXXXMXXXXXXX
3'  MYYYYYYYYYYYYYY
```

```
5'  XXXXXXXMXXXXXXX
3'  YYYYYYYYYYYYYMY
```

```
5'  XXXXXXXXXXMXXXX
3'  MYYYYYYYYYYYYYY
```

```
5'  MMMMMMMMMMMMMMMM
3'  MMMMMMMMMMMMMMMM
```

[0277] Nucleic acid molecules may have blunt ends on both ends with alternating modified nucleic acids. The modified nucleic acids may be located at any position along either the sense or antisense strand.

```
5'  MXXMXMXMXMXMXMXMXM
3'  MYMYMYMYMYMYMYMY
```

```
5'  XMXMXMXMXMXMXMXMX
3'  MYMYMYMYMYMYMYMY
```

```
5'  MXXMXXMXMXMXMXMXM
3'  YMYMYMYMYMYMYMY
```

```
5'  XMXMXMXMXMXMXMXMX
3'  YMYMYMYMYMYMYMY
```

```

3'      MMYMMYMMYMMYMMYMMYMM
5'      MMYMMYMMYMMYMMYMMYMM
3'      YMMYMMYMMYMMYMMYMMYMM

5'      XMMYMMYMMYMMYMMYMMYMM
3'      MMYMMYMMYMMYMMYMMYMM

```

[0278] Nucleic acid molecules with a blunt 5'-end and 3'-end overhang end with a single modified nucleic acid.

[0279] Nucleic acid molecules with a 5'-end overhang and a blunt 3'-end with a single modified nucleic acid.

[0280] Nucleic acid molecules with overhangs on both ends and all overhangs are modified nucleic acids. In the pattern immediately below, M is n number of modified nucleic acids, where n is an integer from 0 to 8 (i.e., 0, 1, 2, 3, 4, 5, 6, 7 and 8).

```

5'      XXXXXXXXXXXXXXXXXXXX
3'      MYYYYYYYYYYYYYYYYY

```

[0281] Nucleic acid molecules with overhangs on both ends and some overhang nucleotides are modified nucleotides. In the patterns immediately below, M is n number of modified nucleotides, x is n number of unmodified overhang nucleotides in the sense strand, y is n number of unmodified overhang nucleotides in the antisense strand, where each n is independently an integer from 0 to 8 (i.e., 0, 1, 2, 3, 4, 5, 6, 7 and 8), and where each overhang is maximum of 20 nucleotides; preferably a maximum of 8 nucleotides (modified and/or unmodified).

```

5'      XXXXXXXXXXXXXXXXXXXX
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMx
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMxM
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMxMx
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMxMxM
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMxMxMx
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMxMxMxM
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      XXXXXXXXXXXXXXXXXXXMxMxMxMx
3'      yYYYYYYYYYYYYYYYYY

```

```

5'      MXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      xMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      MxMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      xMxMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      MxMxMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      xMxMxMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      MxMxMxMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      xMxMxMxMXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYY

```

```

5'      xXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYM

```

```

5'      xXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYMy

```

```

5'      xXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYMyM

```

```

5'      xXXXXXXXXXXXXXXXXX
3'      YYYYYYYYYYYYYYYYYMyMy

```

```

5'      xXXXXXXXXXXXXXXXXX

```

```

3'      YYYYYYYYYYYYYYYYYYMyMyM
5'      xxxxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYYMyMyMy

5'      xxxxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYYMyMyMyM

5'      xxxxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYYMyMyMyMy

5'      xxxxxxxxxxxxxxxxxxxx
3'      MyYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxx
3'      yMYYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxx
3'      MyMYYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxx
3'      yMyMYYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxx
3'      MyMyMYYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxx
3'      yMyMyMYYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxx
3'      MyMyMyMYYYYYYYYYYYYYYYYYY

```

Modified nucleotides at the 3'-end of the sense region.

```

5'      xxxxxxxxxxxxxxxxxxxxM
3'      YYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxxMM
3'      YYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxxMMM
3'      YYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxxMMMM
3'      YYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxxMMMMM
3'      YYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxxMMMMMM
3'      YYYYYYYYYYYYYYYYYY

5'      xxxxxxxxxxxxxxxxxxxxMMMMMMM
3'      YYYYYYYYYYYYYYYYYY

```

Overhang at the 5'-end of the sense region.

```

5'      Mxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

5'      MMMxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

5'      MMMMxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

5'      MMMMMxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

5'      MMMMMxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

5'      MMMMMxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

5'      MMMMMxxxxxxxxxxxxxxxxx
3'      YYYYYYYYYYYYYYYYYY

```

[0282] Overhang at the 3'-end of the antisense region.

```

5'      xxxxxxxxxxxxxxxxxxxx
3'      MYYYYYYYYYYYYYYYYYY

```

5' YYYYYYYYYYYYYYYYYY

[illegible]

[0283] Exemplary nucleic acid molecules are provided below along with the equivalent general structure in line with the symbols used above:

[0284] siHSP47-C siRNA to human and rat hsp47 having a 19 nucleotide (i.e., 19mer) duplex region and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

5' GGACAGGCCUCUACAACUAdTdT
3' dTdTCCUGUCCGAGAUUGUAU 5'

5' XXXXXXXXXXXXXXXXXXXXMM
3' MMYYYYYYYYYYYYYYYYYYYYYY

[0285] siHSP47-Cd siRNA to human and rat hsp47 having a 25-mer duplex region, a 2 nucleotide overhang at the 3'-end of the antisense strand and 2 modified nucleotides at the 5'-terminal and penultimate positions of the sense strand.

[illegible]

[0286] siHSP47-1 siRNA to human and rat hsp47 cDNA 719-737 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

5' CAGGCCUCUACAACUACUAdTdT 3'
3' dTdTGUCCGGAGAUGUUGAUGAU 5'

5' XXXXXXXXXXXXXXXXXXXXMM 3'
3' MMYYYYYYYYYYYYYYYYYYYYY 5'

[0287] siHSP47-ld siRNA to human hsp47 cDNA 719-743 having a 25-mer with a blunt end at the 3'-end of the sense strand and a 2 nucleotide overhang at the 3'-end of the antisense strand, and 2 modified nucleotides at the 5'- terminal and penultimate positions of the sense strand.

overhang at the 3' end of the antisense strand

5' CAGGCCUCUACAACUACAGCAdGa 3'

3' UUGUCCGGAGAUGUUGAUGCUGCU 5'

5' XXXXXXXXXXXXXXXXXXMM 3'

3' yYYYYYYYYYYYYYYYYYYY 5'

[0288] siHSP47-2 siRNA to human hsp47 cDNA 469-487 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

5' GAGCACUCCAAGAUCAACUdTdT 3'
3' dTdTCCUGGAGGUUCUAGUUGA 5'

5' XXXXXXXXXXXXXXXXXXXXMM 3'
3' MMYYYYYYYYYYYYYYYYYYYY 5'

[0289] siHSP47-2d siRNA to human hsp47 cDNA 469-493 having a 25-mer duplex region with a blunt end at the 3'-end of the sense strand and a 2 nucleotide overhang at the 3'-end of the antisense strand, and 2 modified nucleotides at the 5'- terminal and penultimate positions of the sense strand.

```

5'      GAGCACUCCAAGAUCAACUCCGdCdG  3'
3'      UUCUCUGAGGUUCUAGUUGAAGGC GC  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      yyYYYYYYYYYYYYYYYYYYYYYY  5'

```

[0290] siHSP47-2d rat siRNA to rat Gp46 cDNA 466-490 having a 25-mer duplex region with a blunt end at the 3'-end of the sense strand and a 2 nucleotide overhang at the 3'-end of the antisense strand, and 2 modified nucleotides at the 5'- terminal and penultimate positions of the sense strand.

```

5'      GAACACUCCAAGAUCAACUCCGdaadg  3'
3'      UUCUUGUGAGGUUCUAGUUGAAGGCUC  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      yyYYYYYYYYYYYYYYYYYYYYYY  5'

```

[0291] siHSP47-3 siRNA to human hsp47 cDNA 980-998 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

```

5'      CTGAGGCCATTGACAAGAAdTdT  3'
3'      dTdTGACUCCGGUAAACUGUUCUU  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      MMYYYYYYYYYYYYYYYYYYYYYY  5'

```

[0292] siHSP47-3d siRNA to human hsp47 cDNA 980-1004 having a 25-mer duplex region with a blunt end at the 3'-end of the sense strand and a 2 nucleotide overhang at the 3'-end of the antisense strand, and 2 modified nucleotides at the 5'- terminal and penultimate positions of the sense strand.

```

5'      CTGAGGCCATTGACAAGAAGdgdC  3'
3'      UUGACUCCGGUAAACUGUUCUUGCCG  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      yyYYYYYYYYYYYYYYYYYYYYYY  5'

```

[0293] siHSP47-4 siRNA to human hsp47 cDNA 735-753 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

```

5'      CUACGACGACGAGAGGAAdTdT  3'
3'      dTdTGAUGCUGCUGCUCUUCUU  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      MMYYYYYYYYYYYYYYYYYYYYYY  5'

```

[0294] siHSP47-4d siRNA to human hsp47 cDNA 735-759 having a 25-mer duplex region with a blunt end at the 3'-end of the sense strand and a 2 nucleotide overhang at the 3'-end of the antisense strand, and 2 modified nucleotides at the 5'- terminal and penultimate positions of the sense strand.

```

5'      CUACGACGACGAGAGGAAAAGCdTdg  3'
3'      UUGAUGCUGCUGCUCUCCUUCGAC  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      yyYYYYYYYYYYYYYYYYYYYYYY  5'

```

siHSP47-5 siRNA to human hsp47 cDNA 621-639 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

```

5'      GCCACACUGGGAUGAGAAAdTdT  3'
3'      dTdTCCGUGUGACCCUACUCUU  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      MMYYYYYYYYYYYYYYYYYYYYYY  5'

```

siHSP47-6 siRNA to human hsp47 cDNA 446-464 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

```

5'      GCAGCAAGCAGCACUACAAdTdT  3'
3'      dTdTCGUCGUCGUCGUGAUGUU  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      MMYYYYYYYYYYYYYYYYYYYYYY  5'

```

siHSP47-7 siRNA to human hsp47 cDNA 692-710 having a 19-mer duplex region, and modified 2 nucleotide (i.e., deoxynucleotide) overhangs at the 3'-ends of the sense and antisense strands.

```

5'      CCGUGGUGUCAUGAUGAdTdT  3'
3'      dTdTGGCACCACAGUACUACUA  5'

5'      XXXXXXXXXXXXXXXXXXXXXXXMM  3'
3'      MMYYYYYYYYYYYYYYYYYYYYYY  5'

```

Nicks and Gaps in Nucleic Acid Strands

[0295] Nucleic acid molecules (e.g., siRNA molecules) provided herein may have a strand, preferably, the sense strand, that is nicked or gapped.

As such, nucleic acid molecules may have three or more strand, for example, such as a meroduplex RNA (mdRNA) disclosed in PCT/US07/081836. Nucleic acid molecules with a nicked or gapped strand may be between about 1-49 nucleotides, or may be RISC length (e.g., about 15 to 25 nucleotides) or Dicer substrate length (e.g., about 25 to 30 nucleotides) such as disclosed herein.

[0296] Nucleic acid molecules with three or more strands include, for example, an 'A' (antisense) strand, 'S1' (second) strand, and 'S2' (third) strand in which the 'S1' and 'S2' strands are complementary to and form base pairs with non-overlapping regions of the 'A' strand (e.g., an mdRNA can have the form of A:S1S2). The S1, S2, or more strands together form what is substantially similar to a sense strand to the 'A' antisense strand. The double-stranded region formed by the annealing of the 'S1' and 'A' strands is distinct from and non-overlapping with the double-stranded region formed by the annealing of the 'S2' and 'A' strands. A nucleic acid molecule (e.g., an siRNA molecule) may be a "gapped" molecule, meaning a "gap" ranging from 0 nucleotides up to about 10 nucleotides (e.g., 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 nucleotides). Preferably, the sense strand is gapped. In some embodiments, the A:S1 duplex is separated from the A:S2 duplex by a gap resulting from at least one unpaired nucleotide (up to about 10 unpaired nucleotides) in the 'A' strand that is positioned between the A:S1 duplex and the A:S2 duplex and that is distinct from any one or more unpaired nucleotide at the 3'-end of one or more of the 'A', 'S1', or 'S2' strands. The A:S1 duplex may be separated from the A:S2 duplex by a gap of zero nucleotides (i.e., a nick in which only a phosphodiester bond between two nucleotides is broken or missing in the polynucleotide molecule) between the A:S1 duplex and the A:S2 duplex-which can also be referred to as nicked dsRNA (ndsRNA). For example, A:S1S2 may be include a dsRNA having at least two double-stranded regions that combined total about 14 base pairs to about 40 base pairs and the double-stranded regions are separated by a gap of about 0 to about 10 nucleotides, optionally having blunt ends, or A:S1S2 may include a dsRNA having at least two double-stranded regions separated by a gap of up to ten nucleotides wherein at least one of the double-stranded regions includes between about five base pairs and thirteen base pairs.

Dicer Substrates

[0297] In certain embodiments, the nucleic acid molecules (e.g., siRNA molecules) provided herein may be a precursor "Dicer substrate" molecule, e.g., double-stranded nucleic acid, processed *in vivo* to produce an active nucleic acid molecules, for example as described in Rossi, US20050244858. In certain conditions and situations, it has been found that these relatively longer dsRNA siRNA species, e.g., of from about 25 to about 30 nucleotides, can give unexpectedly effective results in terms of potency and duration of action. Without wishing to be bound by any particular theory, it is thought that the longer dsRNA species serve as a substrate for the enzyme Dicer in the cytoplasm of a cell. In addition to cleaving double-stranded nucleic acid into shorter segments, Dicer may facilitate the incorporation of a single-stranded cleavage product derived from the cleaved dsRNA into the RNA-induced silencing complex (RISC complex) that is responsible for the destruction of the cytoplasmic RNA derived from the target gene.

[0298] Dicer substrates may have certain properties which enhance its processing by Dicer. Dicer substrates are of a length sufficient such that it is processed by Dicer to produce an active nucleic acid molecule and may further include one or more of the following properties: (i) the dsRNA is asymmetric, e.g., has a 3'-overhang on the first strand (antisense strand) and (ii) the dsRNA has a modified 3'-end on the antisense strand (sense strand) to direct orientation of Dicer binding and processing of the dsRNA to an active siRNA. In certain embodiments, the longest strand in the Dicer substrate may be 24-30 nucleotides.

[0299] Dicer substrates may be symmetric or asymmetric. The Dicer substrate may have a sense strand includes 22-28 nucleotides and the antisense strand may include 24-30 nucleotides; thus, in some embodiments the resulting Dicer substrate may have an overhang on the 3' end of the antisense strand. Dicer substrate may have a sense strand 25 nucleotides in length, and the antisense strand having 27 nucleotides in length with a two base 3'-overhang. The overhang may be 1-3 nucleotides, for example 2 nucleotides. The sense strand may also have a 5'-phosphate.

[0300] An asymmetric Dicer substrate may further contain two deoxynucleotides at the 3'-end of the sense strand in place of two of the ribonucleotides. Some exemplary Dicer substrates lengths and structures are 21+0, 21+2, 21-2, 22+0, 22+1, 22-1, 23+0, 23+2, 23-2, 24+0, 24+2, 24-2, 25+0, 25+2, 25-2, 26+0, 26+2, 26-2, 27+0, 27+2, and 27-2.

[0301] The sense strand of a Dicer substrate may be between about 22 to about 30 (e.g., about 22, 23, 24, 25, 26, 27, 28, 29 or 30); about 22 to about 28; about 24 to about 30; about 25 to about 30; about 26 to about 30; about 26 and 29; or about 27 to about 28 nucleotides in length. In certain preferred embodiments Dicer substrates contain sense and antisense strands, that are at least about 25 nucleotides in length and no longer than about 30 nucleotides; between about 26 and 29 nucleotides; or 27 nucleotides in length. The sense and antisense strands may be the same length (blunt ended), different lengths (have overhangs), or a combination. The sense and antisense strands may exist on the same polynucleotide or on different polynucleotides. A Dicer substrate may have a duplex region of about 19, 20, 21, 22, 23, 24, 25 or 27 nucleotides.

[0302] Like other siRNA molecules provided herein, the antisense strand of a Dicer substrate may have any sequence that anneals to the antisense strand under biological conditions, such as within the cytoplasm of a eukaryotic cell.

[0303] Dicer substrates may have any modifications to the nucleotide base, sugar or phosphate backbone as known in the art and/or as described herein for other nucleic acid molecules (such as siRNA molecules). In certain embodiments, Dicer substrates may have a sense strand is modified for Dicer processing by suitable modifiers located at the 3'-end of the sense strand, i.e., the dsRNA is designed to direct orientation of Dicer binding and processing. Suitable modifiers include nucleotides such as deoxyribonucleotides, dideoxyribonucleotides, acyclo-nucleotides and the like and sterically hindered molecules, such as fluorescent molecules and the like. Acyclo-nucleotides substitute a 2-hydroxyethoxymethyl group for the 2'-deoxyribofuranosyl sugar normally present in dNMPs. Other nucleotides modifiers that could be used in Dicer substrate siRNA molecules include 3'-deoxyadenosine (cordycepin), 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxyinosine (ddI), 2',3'-dideoxy-3'-thiacytidine (3TC), 2',3'-didehydro-2',3'-dideoxythymidine (d4T) and the monophosphate nucleotides of 3'-azido-3'-deoxythymidine (AZT), 2',3'-dideoxy-3'-thiacytidine (3TC) and 2',3'-didehydro-2',3'-dideoxythymidine (d4T). In one embodiment, deoxynucleotides are used as the modifiers. When nucleotide modifiers are utilized, they may replace ribonucleotides (e.g., 1-3 nucleotide modifiers, or 2 nucleotide modifiers are substituted for the ribonucleotides on the 3'-end of the sense strand) such that the length of the Dicer substrate does not change. When sterically hindered molecules

are utilized, they may be attached to the ribonucleotide at the 3'-end of the antisense strand. Thus, in certain embodiments the length of the strand does not change with the incorporation of the modifiers. In certain embodiments, two DNA bases in the dsRNA are substituted to direct the orientation of Dicer processing of the antisense strand. In a further embodiment of, two terminal DNA bases are substituted for two ribonucleotides on the 3'-end of the sense strand forming a blunt end of the duplex on the 3'-end of the sense strand and the 5'-end of the antisense strand, and a two-nucleotide RNA overhang is located on the 3'-end of the antisense strand. This is an asymmetric composition with DNA on the blunt end and RNA bases on the overhanging end.

[0304] In certain embodiments modifications are included in the Dicer substrate such that the modification does not prevent the nucleic acid molecule from serving as a substrate for Dicer. In one embodiment, one or more modifications are made that enhance Dicer processing of the Dicer substrate. One or more modifications may be made that result in more effective RNAi generation. One or more modifications may be made that support a greater RNAi effect. One or more modifications are made that result in greater potency per each Dicer substrate to be delivered to the cell. Modifications may be incorporated in the 3'-terminal region, the 5'-terminal region, in both the 3'-terminal and 5'-terminal region or at various positions within the sequence. Any number and combination of modifications can be incorporated into the Dicer substrate so long as the modification does not prevent the nucleic acid molecule from serving as a substrate for Dicer. Where multiple modifications are present, they may be the same or different. Modifications to bases, sugar moieties, the phosphate backbone, and their combinations are contemplated. Either 5'-terminus can be phosphorylated.

[0305] Examples of Dicer substrate phosphate backbone modifications include phosphonates, including methylphosphonate, phosphorothioate, and phosphotriester modifications such as alkylphosphotriesters, and the like. Examples of Dicer substrate sugar moiety modifications include 2'-alkyl pyrimidine, such as 2'OMe, 2'-fluoro, amino, and deoxy modifications and the like (see, e.g., Amarzguioui *et al.*, 2003). Examples of Dicer substrate base group modifications include abasic sugars, 2-O-alkyl modified pyrimidines, 4-thiouracil, 5-bromouracil, 5-iodouracil, and 5-(3-aminoallyl)-uracil and the like. LNAs could also be incorporated.

[0306] The sense strand may be modified for Dicer processing by suitable modifiers located at the 3'-end of the sense strand, i.e., the Dicer substrate is designed to direct orientation of Dicer binding and processing. Suitable modifiers include nucleotides such as deoxyribonucleotides, dideoxyribonucleotides, acyclo-nucleotides and the like and sterically hindered molecules, such as fluorescent molecules and the like. Acyclo-nucleotides substitute a 2-hydroxyethoxymethyl group for the 2'-deoxyribofuranosyl sugar normally present in dNMPs. Other nucleotides modifiers could include cordycepin, AZT, ddl, 3TC, d4T and the monophosphate nucleotides of AZT, 3TC and d4T. In one embodiment, deoxynucleotides are used as the modifiers. When nucleotide modifiers are utilized, 1-3 nucleotide modifiers, or 2 nucleotide modifiers are substituted for the ribonucleotides on the 3'-end of the sense strand. When sterically hindered molecules are utilized, they are attached to the ribonucleotide at the 3'-end of the antisense strand. Thus, the length of the strand does not change with the incorporation of the modifiers. In another embodiment, the description contemplates substituting two DNA bases in the Dicer substrate to direct the orientation of Dicer processing of the antisense strand. In a further embodiment of the present description, two terminal DNA bases are substituted for two ribonucleotides on the 3'-end of the sense strand forming a blunt end of the duplex on the 3'-end of the sense strand and the 5'-end of the antisense strand, and a two-nucleotide RNA overhang is located on the 3'-end of the antisense strand. This is an asymmetric composition with DNA on the blunt end and RNA bases on the overhanging end.

[0307] The antisense strand may be modified for Dicer processing by suitable modifiers located at the 3'-end of the antisense strand, i.e., the dsRNA is designed to direct orientation of Dicer binding and processing. Suitable modifiers include nucleotides such as deoxyribonucleotides, dideoxyribonucleotides, acyclo-nucleotides and the like and sterically hindered molecules, such as fluorescent molecules and the like. Acyclo-nucleotides substitute a 2-hydroxyethoxymethyl group for the 2'-deoxyribofuranosyl sugar normally present in dNMPs. Other nucleotides modifiers could include cordycepin, AZT, ddl, 3TC, d4T and the monophosphate nucleotides of AZT, 3TC and d4T. In one embodiment, deoxynucleotides are used as the modifiers. When nucleotide modifiers are utilized, 1-3 nucleotide modifiers, or 2 nucleotide modifiers are substituted for the ribonucleotides on the 3'-end of the antisense strand. When sterically hindered molecules are utilized, they are attached to the ribonucleotide at the 3'-end of the antisense strand. Thus, the length of the strand does not change with the incorporation of the modifiers. In another embodiment, the description contemplates substituting two DNA bases in the dsRNA to direct the orientation of Dicer processing. In a further description, two terminal DNA bases are located on the 3'-end of the antisense strand in place of two ribonucleotides forming a blunt end of the duplex on the 5'-end of the sense strand and the 3'-end of the antisense strand, and a two-nucleotide RNA overhang is located on the 3'-end of the sense strand. This is an asymmetric composition with DNA on the blunt end and RNA bases on the overhanging end.

[0308] Dicer substrates with a sense and an antisense strand can be linked by a third structure. The third structure will not block Dicer activity on the Dicer substrate and will not interfere with the directed destruction of the RNA transcribed from the target gene. The third structure may be a chemical linking group. Suitable chemical linking groups are known in the art and can be used. Alternatively, the third structure may be an oligonucleotide that links the two oligonucleotides of the dsRNA in a manner such that a hairpin structure is produced upon annealing of the two oligonucleotides making up the Dicer substrate. The hairpin structure preferably does not block Dicer activity on the Dicer substrate or interfere with the directed destruction of the RNA transcribed from the target gene.

[0309] The sense and antisense strands of the Dicer substrate are not required to be completely complementary. They only need to be substantially complementary to anneal under biological conditions and to provide a substrate for Dicer that produces a siRNA sufficiently complementary to the target sequence.

[0310] Dicer substrate can have certain properties that enhance its processing by Dicer. The Dicer substrate can have a length sufficient such that it is processed by Dicer to produce an active nucleic acid molecules (e.g., siRNA) and may have one or more of the following properties: the Dicer substrate is asymmetric, e.g., has a 3'-overhang on the first strand (antisense strand) and/or the Dicer substrate has a modified 3' end on the second strand (sense strand) to direct orientation of Dicer binding and processing of the Dicer substrate to an active siRNA. The Dicer substrate can be asymmetric such that the sense strand includes 22-28 nucleotides and the antisense strand includes 24-30 nucleotides. Thus, the resulting Dicer substrate has an overhang on the 3' end of the antisense strand. The overhang is 1-3 nucleotides, for example two nucleotides. The sense strand may also have a 5' phosphate.

[0311] A Dicer substrate may have an overhang on the 3'-end of the antisense strand and the sense strand is modified for Dicer processing. The 5'-end of the sense strand may have a phosphate. The sense and antisense strands may anneal under biological conditions, such as the conditions found in the cytoplasm of a cell. A region of one of the strands, particularly the antisense strand, of the Dicer substrate may have a sequence length of at least 19 nucleotides, wherein these nucleotides are in the 21-nucleotide region adjacent to the 3'-end of the antisense strand and are sufficiently complementary to a nucleotide sequence of the RNA produced from the target gene. A Dicer substrate may also have one or more of the following additional properties: the antisense strand has a right shift from a corresponding 21-mer (i.e., the antisense strand includes nucleotides on the right side of the molecule when compared to the corresponding 21-mer), the strands may not be completely complementary, i.e., the strands may contain simple mismatch pairings and base modifications such as LNA may be included in the 5'-end of the sense strand.

[0312] An antisense strand of a Dicer substrate nucleic acid molecule may be modified to include 1-9 ribonucleotides on the 5' -end to give a length of 22-28 nucleotides. When the antisense strand has a length of 21 nucleotides, then 1-7 ribonucleotides, or 2-5 ribonucleotides and or 4 ribonucleotides may be added on the 3'-end. The added ribonucleotides may have any sequence. Although the added ribonucleotides may be complementary to the target gene sequence, full complementarity between the target sequence and the antisense strands is not required. That is, the resultant antisense strand is sufficiently complementary with the target sequence. A sense strand may then have 24-30 nucleotides. The sense strand may be substantially complementary with the antisense strand to anneal to the antisense strand under biological conditions. In one embodiment, the antisense strand may be synthesized to contain a modified 3' -end to direct Dicer processing. The sense strand may have a 3' overhang. The antisense strand may be synthesized to contain a modified 3' -end for Dicer binding and processing and the sense strand has a 3' overhang.

Heat Shock Protein 47

[0313] Heat shock protein 47 (HSP47) is a collagen-specific molecular chaperone and resides in the endoplasmic reticulum. It interacts with procollagen during the process of folding, assembling and transporting from the endoplasmic reticulum (Nagata Trends Biochem Sci 1996; 21:22-6; Razzaque et al. 2005; Contrib Nephrol 2005; 148: 57-69; Koide et al. 2006 J. Biol. Chem.; 281: 3432-38; Leivo et al. Dev. Biol. 1980; 76:100-114; Masuda et al. J. Clin. Invest. 1994; 94:2481-2488; Masuda et al. Cell Stress Chaperones 1998; 3:256-264). HSP47 has been reported to have an upregulated expression in various tissue fibrosis (Koide et al. J Biol Chem 1999; 274: 34523-26), such as liver cirrhosis (Masuda et al. J Clin Invest 1994; 94:2481-8), pulmonary fibrosis (Razzaque et al. Virchows Arch 1998; 432:455-60; Kakugawa et al. Eur Respir J 2004; 24: 57-65), and glomerulosclerosis (Moriyama et al. Kidney Int 1998; 54: 110-19). Exemplary nucleic acid sequence of target human hsp47 cDNA is disclosed in GenBank accession number: NM_001235 and the corresponding mRNA sequence, for example as listed as SEQ ID NO:1. One of ordinary skill in the art would understand that a given sequence may change over time and to incorporate any changes needed in the nucleic acid molecules herein accordingly.

[0314] The specific association of HSP47 with a diverse range of collagen types makes HSP47 a potential target for the treatment of fibrosis. Inhibition of hsp47 expression may prevent extracellular collagen I secretion. Sato et al. (Nat Biotechnol 2008; 26:431-442) explored this possibility by using siRNA for the inhibition hsp47 expression and preventing the progression of hepatic fibrosis in rats. Similarly, Chen et al. (Br J Dermatol 2007; 156: 1188-1195) and Wang et al. (Plast. Reconstr Surg 2003; 111: 1980-7) investigated the inhibition hsp47 expression by RNA interference technology.

Methods and Compositions for Inhibiting hsp47

[0315] Provided are compositions and methods for inhibition of hsp47 expression by using small nucleic acid molecules, such as short interfering nucleic acid (siNA), interfering RNA (RNAi), short interfering RNA (siRNA), double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating or that mediate RNA interference against hsp47 gene expression. The composition and methods disclosed herein are also useful in treating various fibrosis such as liver fibrosis, lung fibrosis, and kidney fibrosis.

[0316] Nucleic acid molecule(s) and/or methods of the description are used to down regulate the expression of gene(s) that encode RNA referred to, by example, Genbank Accession NM_001235.

[0317] Compositions, methods and kits provided herein may include one or more nucleic acid molecules (e.g., siNA) and methods that independently or in combination modulate (e.g., downregulate) the expression of hsp47 protein and/or genes encoding hsp47 proteins, proteins and/or genes encoding hsp47 (e.g., genes encoding sequences comprising those sequences referred to by GenBank Accession Nos. NM_001235), or an hsp47 gene family member where the genes or gene family sequences share sequence homology associated with the maintenance and/or development of diseases, conditions or disorders associated with hsp47, such as liver fibrosis, cirrhosis, pulmonary fibrosis, kidney fibrosis, peritoneal fibrosis, chronic hepatic damage, and fibrillogenesis. The description of the various aspects and embodiments is provided with reference to exemplary gene hsp47. However, the various aspects and embodiments are also directed to other related hsp47 genes, such as homolog genes and transcript variants, and polymorphisms (e.g., single nucleotide polymorphism, (SNPs)) associated with certain hsp47 genes. As such, the various aspects and embodiments are also directed to other genes that are involved in hsp47 mediated pathways of signal transduction or gene expression that are involved, for example, in the maintenance or development of diseases, traits, or conditions described herein. These additional genes can be analyzed for target sites using the methods described for the hsp47 gene herein. Thus, the modulation of other genes and the effects of such modulation of the other genes can be performed, determined, and measured as described herein.

[0318] In one embodiment, compositions and methods provided herein include a double-stranded short interfering nucleic acid (siNA) molecule that down-regulates expression of a hsp47 gene (e.g., human hsp47 exemplified by SEQ ID NO:1), where the nucleic acid molecule includes about 15 to about 49 base pairs.

[0319] In one embodiment, a nucleic acid disclosed may be used to inhibit the expression of the hsp47 gene or an hsp47 gene family where the genes or gene family sequences share sequence homology. Such homologous sequences can be identified as is known in the art, for example using sequence alignments. Nucleic acid molecules can be designed to target such homologous sequences, for example using perfectly complementary sequences or by incorporating non-canonical base pairs, for example mismatches and/or wobble base pairs that can provide additional target sequences. In instances where mismatches are identified, non-canonical base pairs (for example, mismatches and/or wobble bases) can be used to generate nucleic acid molecules that target more than one gene sequence. In a non-limiting example, non-canonical base pairs such as UU and CC base pairs are used to generate nucleic acid molecules that are capable of targeting sequences for differing hsp47 targets that share sequence homology. As such, one advantage of using siNAs disclosed herein is that a single nucleic acid can be designed to include nucleic acid sequence that is complementary to the nucleotide sequence that is conserved between the homologous genes. In this approach, a single nucleic acid can be used to inhibit expression of more than one gene instead of using more than one nucleic acid molecule to target the different genes.

[0320] Nucleic acid molecules may be used to target conserved sequences corresponding to a gene family or gene families such as hsp47 family genes. As such, nucleic acid molecules targeting multiple hsp47 targets can provide increased therapeutic effect. In addition, nucleic acid can be used to characterize pathways of gene function in a variety of applications. For example, nucleic acid molecules can be used to inhibit the activity of target gene(s) in a pathway to determine the function of uncharacterized gene(s) in gene function analysis, mRNA function analysis, or translational analysis. The nucleic acid molecules can be used to determine potential target gene pathways involved in various diseases and conditions toward pharmaceutical development. The nucleic acid molecules can be used to understand pathways of gene expression involved in, for example fibroses such as liver, kidney or pulmonary fibrosis, and/or inflammatory and proliferative traits, diseases, disorders, and/or conditions.

[0321] In one embodiment, the compositions and methods provided herein include a nucleic acid molecule having RNAi activity against hsp47 RNA, where the nucleic acid molecule includes a sequence complementary to any RNA having hsp47 encoding sequence, such as those sequences having sequences as shown in Table 3. In another embodiment, a nucleic acid molecule may have RNAi activity against hsp47 RNA, where the nucleic acid molecule includes a sequence complementary to an RNA having variant hsp47 encoding sequence, for example other mutant hsp47 genes not shown in Table 3 but known in the art to be associated with the maintenance and/or development of fibrosis. Chemical modifications as shown in Table 3 or otherwise described herein can be applied to any nucleic acid construct disclosed herein. In another embodiment, a nucleic acid molecule disclosed herein includes a nucleotide sequence that can interact with nucleotide sequence of a hsp47 gene and thereby mediate silencing of hsp47 gene expression, for example, wherein the nucleic acid molecule mediates regulation of hsp47 gene expression by cellular processes that modulate the chromatin structure or methylation patterns of the hsp47 gene and prevent transcription of the hsp47 gene.

[0322] Nucleic acid molecules disclosed herein may have RNAi activity against hsp47 RNA, where the nucleic acid molecule includes a sequence complementary to any RNA having hsp47 encoding sequence, such as those sequences having GenBank Accession Nos. NM_001235. Nucleic acid molecules may have RNAi activity against hsp47 RNA, where the nucleic acid molecule includes a sequence complementary to an RNA having variant hsp47 encoding sequence, for example other mutant hsp47 genes known in the art to be associated with the maintenance and/or development of fibrosis.

[0323] Nucleic acid molecules disclosed herein include a nucleotide sequence that can interact with nucleotide sequence of a hsp47 gene and thereby mediate silencing of hsp47 gene expression, *e.g.*, where the nucleic acid molecule mediates regulation of hsp47 gene expression by cellular processes that modulate the chromatin structure or methylation patterns of the hsp47 gene and prevent transcription of the hsp47 gene.

Methods of Treatment

[0324] The specific association of HSP47 with a diverse range of collagen types makes hsp47 a target for the treatment of fibrosis. Inhibition of hsp47 expression may prevent extracellular collagen I secretion. Sato et al. (Nat Biotechnol 2008; 26:431-442) explored this possibility by using siRNA for the inhibition hsp47 expression and preventing the progression of hepatic fibrosis in rats. Similarly, Chen et al. (Br J Dermatol 2007; 156: 1188-1195) and Wang et al. (Plast Reconstr Surg 2003; 111: 1980-7) investigated the inhibition hsp47 expression by RNA interference technology.

[0325] In one embodiment, nucleic acid molecules may be used to down regulate or inhibit the expression of hsp47 and/or hsp47 proteins arising from hsp47 and/or hsp47 haplotype polymorphisms that are associated with a disease or condition, (*e.g.*, fibrosis). Analysis of hsp47 and/or hsp47 genes, or hsp47 and/or hsp47 protein or RNA levels can be used to identify subjects with such polymorphisms or those subjects who are at risk of developing traits, conditions, or diseases described herein. These subjects are amenable to treatment, for example, treatment with nucleic acid molecules disclosed herein and any other composition useful in treating diseases related to hsp47 and/or hsp47 gene expression. As such, analysis of hsp47 and/or hsp47 protein or RNA levels can be used to determine treatment type and the course of therapy in treating a subject. Monitoring of hsp47 and/or hsp47 protein or RNA levels can be used to predict treatment outcome and to determine the efficacy of compounds and compositions that modulate the level and/or activity of certain hsp47 and/or hsp47 proteins associated with a trait, condition, or disease.

[0326] Provided are compositions and methods for inhibition of hsp47 expression by using small nucleic acid molecules as provided herein, such as siNA, RNAi, siRNA, double-stranded RNA (dsRNA), micro-RNA (miRNA), and short hairpin RNA (shRNA) molecules capable of mediating or that mediate RNA interference against hsp47 gene expression. The composition and methods disclosed herein are also useful in treating various fibrosis such as liver fibrosis, lung fibrosis, and kidney fibrosis.

[0327] The nucleic acid molecules disclosed herein individually, or in combination or in conjunction with other drugs, can be used for preventing or treating diseases, traits, conditions and/or disorders associated with hsp47, such as liver fibrosis, cirrhosis, pulmonary fibrosis, kidney fibrosis, peritoneal fibrosis, chronic hepatic damage, and fibrillogenesis.

[0328] The nucleic acid molecules disclosed herein are able to inhibit the expression of hsp47 in a sequence specific manner. The nucleic acid molecules may include a sense strand and an antisense strand which include contiguous nucleotides that are at least partially complementary (antisense) to an hsp47 mRNA.

[0329] In some embodiments, dsRNA specific for hsp47 can be used in conjunction with other dsRNA specific for other molecular chaperones that assist in the folding of newly synthesized proteins such as, calnexin, calreticulin, and/or BiP (Bergeron et al. Trends Biochem. Sci. 1994; 19:124-128; Herbert et al. 1995; Cold Spring Harb. Symp. Quant. Biol. 60:405-415)

[0330] Fibrosis can be treated by RNA interference using nucleic acid molecules as disclosed herein. Exemplary fibroses include liver fibrosis, peritoneal fibrosis, lung fibrosis, kidney fibrosis. The nucleic acid molecules disclosed herein may inhibit the expression of hsp47 in a sequence specific manner.

[0331] Treatment of fibrosis can be monitored by determining the level of extracellular collagen using suitable techniques known in the art such as, using anti-collagen I antibodies. Treatment can also be monitored by determining the level of hsp47 mRNA or the level of HSP47 protein in the cells of the affected tissue. Treatment can also be monitored by non-invasive scanning of the affected organ or tissue such as by computer assisted tomography scan, magnetic resonance elastography scans.

[0332] A method for treating or preventing hsp47 associated disease or condition in a subject or organism may include contacting the subject or organism with a nucleic acid molecule as provided herein under conditions suitable to modulate the expression of the hsp47 gene in the subject or organism.

[0333] A method for treating or preventing fibrosis in a subject or organism may include contacting the subject or organism with a nucleic acid molecule under conditions suitable to modulate the expression of the hsp47 gene in the subject or organism.

[0334] A method for treating or preventing one or more fibroses selected from the group consisting of liver fibrosis, kidney fibrosis, and pulmonary fibrosis in a subject or organism may include contacting the subject or organism with a nucleic acid molecule under conditions suitable to modulate the expression of the hsp47 gene in the subject or organism.

Fibrotic Diseases

[0335] Fibrotic diseases are generally characterized by the excess deposition of a fibrous material within the extracellular matrix, which contributes to abnormal changes in tissue architecture and interferes with normal organ function.

[0336] All tissues damaged by trauma respond by the initiation of a wound-healing program. Fibrosis, a type of disorder characterized by excessive scarring, occurs when the normal self-limiting process of wound healing response is disturbed, and causes excessive production and deposition of collagen. As a result, normal organ tissue is replaced with scar tissue, which eventually leads to the functional failure of the organ.

[0337] Fibrosis may be initiated by diverse causes and in various organs. Liver cirrhosis, pulmonary fibrosis, sarcoidosis, keloids and kidney fibrosis are all chronic conditions associated with progressive fibrosis, thereby causing a continuous loss of normal tissue function.

[0338] Acute fibrosis (usually with a sudden and severe onset and of short duration) occurs as a common response to various forms of trauma including accidental injuries (particularly injuries to the spine and central nervous system), infections, surgery, ischemic illness (e.g. cardiac scarring following heart attack), burns, environmental pollutants, alcohol and other types of toxins, acute respiratory distress syndrome, radiation and chemotherapy treatments).

[0339] Fibrosis, a fibrosis related pathology or a pathology related to aberrant crosslinking of cellular proteins may all be treated by the siRNAs disclosed herein. Fibrotic diseases or diseases in which fibrosis is evident (fibrosis related pathology) include both acute and chronic forms of fibrosis of organs, including all etiological variants of the following: pulmonary fibrosis, including interstitial lung disease and fibrotic lung disease, liver fibrosis, cardiac fibrosis including myocardial fibrosis, kidney fibrosis including chronic renal failure, skin fibrosis including scleroderma, keloids and hypertrophic scars; myelofibrosis (bone marrow fibrosis); all types of ocular scarring including proliferative vitreoretinopathy (PVR) and scarring resulting from surgery to treat cataract or glaucoma; inflammatory bowel disease of variable etiology, macular degeneration, Grave's ophthalmopathy, drug induced ergotism, keloid scars, scleroderma, psoriasis, glioblastoma in Li-Fraumeni syndrome, sporadic glioblastoma, myeloid leukemia, acute myelogenous leukemia, myelodysplastic syndrome, myeloproliferative syndrome, gynecological cancer, Kaposi's sarcoma, Hansen's disease, and collagenous colitis .

[0340] In various embodiments, the compounds (nucleic acid molecules) as disclosed herein may be used to treat fibrotic diseases, for example as disclosed herein, as well as many other diseases and conditions apart from fibrotic diseases, for example such as disclosed herein. Other conditions to be treated include fibrotic diseases in other organs - kidney fibrosis for any reason (CKD including ESRD); lung fibrosis (including ILF); myelofibrosis, abnormal scarring (keloids) associated with all possible types of skin injury accidental and iatrogenic (operations); scleroderma; cardiofibrosis, failure of glaucoma filtering operation; intestinal adhesions.

Ocular surgery and fibrotic complications

[0341] Contracture of scar tissue resulting from eye surgery may often occur. Glaucoma surgery to create new drainage channels often fails due to scarring and contraction of tissues and the generated drainage system may be blocked requiring additional surgical intervention. Current anti-

scarring regimens (Mitomycin C or 5FU) are limited due to the complications involved (*e.g.* blindness) *e.g.* see Cordeiro MF, et al., Human anti-transforming growth factor-beta2 antibody: a new glaucoma anti-scarring agent Invest Ophthalmol Vis Sci. 1999 Sep;40(10):2225-34. There may also be contraction of scar tissue formed after corneal trauma or corneal surgery, for example laser or surgical treatment for myopia or refractive error in which contraction of tissues may lead to inaccurate results. Scar tissue may be formed on/in the vitreous humor or the retina, for example, and may eventually causes blindness in some diabetics, and may be formed after detachment surgery, called proliferative vitreoretinopathy (PVR). PVR is the most common complication following retinal detachment and is associated with a retinal hole or break. PVR refers to the growth of cellular membranes within the vitreous cavity and on the front and back surfaces of the retina containing retinal pigment epithelial (RPE) cells. These membranes, which are essentially scar tissues, exert traction on the retina and may result in recurrences of retinal detachment, even after an initially successful retinal detachment procedure.

[0342] Scar tissue may be formed in the orbit or on eye and eyelid muscles after squint, orbital or eyelid surgery, or thyroid eye disease, and where scarring of the conjunctiva occurs as may happen after glaucoma surgery or in cicatricial disease, inflammatory disease, for example, pemphigoid, or infective disease, for example, trachoma. A further eye problem associated with the contraction of collagen-including tissues is the opacification and contracture of the lens capsule after cataract extraction. Important role for MMPs has been recognized in ocular diseases including wound healing, dry eye, sterile corneal ulceration, recurrent epithelial erosion, corneal neovascularization, pterygium, conjunctivochalasis, glaucoma, PVR, and ocular fibrosis.

Liver fibrosis

[0343] Liver fibrosis (LF) is a generally irreversible consequence of hepatic damage of several etiologies. In the Western world, the main etiologic categories are: alcoholic liver disease (30-50%), viral hepatitis (30%), biliary disease (5-10%), primary hemochromatosis (5%), and drug-related and cryptogenic cirrhosis of, unknown etiology, (10-15%). Wilson's disease, α_1 -antitrypsin deficiency and other rare diseases also have liver fibrosis as one of the symptoms. Liver cirrhosis, the end stage of liver fibrosis, frequently requires liver transplantation and is among the top ten causes of death in the Western world.

Kidney fibrosis and related conditions

Chronic Renal Failure (CRF)

[0344] Chronic renal failure is a gradual and progressive loss of the ability of the kidneys to excrete wastes, concentrate urine, and conserve electrolytes. CRF is slowly progressive. It most often results from any disease that causes gradual loss of kidney function, and fibrosis is the main pathology that produces CRF.

Diabetic nephropathy

[0345] Diabetic nephropathy, hallmarks of which are glomerulosclerosis and tubulointerstitial fibrosis, is the single most prevalent cause of end-stage renal disease in the modern world, and diabetic patients constitute the largest population on dialysis. Such therapy is costly and far from optimal. Transplantation offers a better outcome but suffers from a severe shortage of donors.

Chronic Kidney Disease

[0346] Chronic kidney disease (CKD) is a worldwide public health problem and is recognized as a common condition that is associated with an increased risk of cardiovascular disease and chronic renal failure (CRF).

[0347] The Kidney Disease Outcomes Quality Initiative (K/DOQI) of the National Kidney Foundation (NKF) defines chronic kidney disease as either kidney damage or a decreased kidney glomerular filtration rate (GFR) for three or more months. Other markers of CKD are also known and used for diagnosis. In general, the destruction of renal mass with irreversible sclerosis and loss of nephrons leads to a progressive decline in GFR. Recently, the K/DOQI published a classification of the stages of CKD, as follows:

Stage 1: Kidney damage with normal or increased GFR (>90 mL/min/1.73 m²)

Stage 2: Mild reduction in GFR (60-89 mL/min/1.73 m²)

Stage 3: Moderate reduction in GFR (30-59 mL/min/1.73 m²)

Stage 4: Severe reduction in GFR (15-29 mL/min/1.73 m²)

Stage 5: Kidney failure (GFR <15 mL/min/1.73 m² or dialysis)

[0348] In stages 1 and 2 CKD, GFR alone does not confirm the diagnosis. Other markers of kidney damage, including abnormalities in the composition of blood or urine or abnormalities in imaging tests, may be relied upon.

Pathophysiology of CKD

[0349] Approximately 1 million nephrons are present in each kidney, each contributing to the total GFR. Irrespective of the etiology of renal injury, with progressive destruction of nephrons, the kidney is able to maintain GFR by hyperfiltration and compensatory hypertrophy of the remaining healthy nephrons. This nephron adaptability allows for continued normal clearance of plasma solutes so that substances such as urea and creatinine start to show significant increases in plasma levels only after total GFR has decreased to 50%, when the renal reserve has been exhausted. The plasma creatinine value will approximately double with a 50% reduction in GFR. Therefore, a doubling in plasma creatinine from a baseline value of 0.6 mg/dL to 1.2 mg/dL in a patient actually represents a loss of 50% of functioning nephron mass.

[0350] The residual nephron hyperfiltration and hypertrophy, although beneficial for the reasons noted, is thought to represent a major cause of progressive renal dysfunction. This is believed to occur because of increased glomerular capillary pressure, which damages the capillaries and leads initially to focal and segmental glomerulosclerosis and eventually to global glomerulosclerosis. This hypothesis has been based on studies of five-sixths nephrectomized rats, which develop lesions that are identical to those observed in humans with CKD.

[0351] The two most common causes of chronic kidney disease are diabetes and hypertension. Other factors include acute insults from nephrotoxins, including contrasting agents, or decreased perfusion; proteinuria; increased renal ammoniogenesis with interstitial injury; hyperlipidemia; Hyperphosphatemia with calcium phosphate deposition; decreased levels of nitrous oxide and smoking.

[0352] In the United States, the incidence and prevalence of CKD is rising, with poor outcomes and high cost to the health system. Kidney disease is the ninth leading cause of death in the US. The high rate of mortality has led the US Surgeon General's mandate for America's citizenry, Healthy People 2010, to contain a chapter focused on CKD. The objectives of this chapter are to articulate goals and to provide strategies to reduce the incidence, morbidity, mortality, and health costs of chronic kidney disease in the United States.

[0353] The incidence rates of end-stage renal disease (ESRD) have also increased steadily internationally since 1989. The United States has the highest incident rate of ESRD, followed by Japan. Japan has the highest prevalence per million population followed by the United States.

[0354] The mortality rates associated with hemodialysis are striking and indicate that the life expectancy of patients entering into hemodialysis is markedly shortened. At every age, patients with ESRD on dialysis have significantly increased mortality when compared with nondialysis patients and individuals without kidney disease. At age 60 years, a healthy person can expect to live for more than 20 years, whereas the life expectancy of a 60-year-old patient starting hemodialysis is closer to 4 years.

Pulmonary fibrosis

[0355] Interstitial pulmonary fibrosis (IPF) is scarring of the lung caused by a variety of inhaled agents including mineral particles, organic dusts, and oxidant gases, or by unknown reasons (idiopathic lung fibrosis). The disease afflicts millions of individuals worldwide, and there are no effective therapeutic approaches. A major reason for the lack of useful treatments is that few of the molecular mechanisms of disease have been defined sufficiently to design appropriate targets for therapy (Lasky, et al., 2000, Environ Health Perspect;108:751-62).

Cardiac fibrosis

[0356] Heart failure is unique among the major cardiovascular disorders in that it alone is increasing in prevalence while there has been a striking decrease in other conditions. Some of this can be attributed to the aging of the populations of the United States and Europe. The ability to salvage patients with myocardial damage is also a major factor, as these patients may develop progression of left ventricular dysfunction due to deleterious remodelling of the heart.

[0357] The normal myocardium is composed of a variety of cells, cardiac myocytes and noncardiomyocytes, which include endothelial and vascular smooth muscle cells and fibroblasts.

[0358] Structural remodeling of the ventricular wall is a key determinant of clinical outcome in heart disease. Such remodeling involves the production and destruction of extracellular matrix proteins, cell proliferation and migration, and apoptotic and necrotic cell death. Cardiac fibroblasts are crucially involved in these processes, producing growth factors and cytokines that act as autocrine and paracrine factors, as well as extracellular matrix proteins and proteinases. Recent studies have shown that the interactions between cardiac fibroblasts and cardiomyocytes are essential for the progression of cardiac remodeling of which the net effect is deterioration in cardiac function and the onset of heart failure (Manabe, et al., 2002, Circ Res. 13:1103-13).

Burns and scars

[0359] A particular problem which may arise, particularly in fibrotic disease, is contraction of tissues, for example contraction of scars. Contraction of tissues including extracellular matrix components, especially of collagen-including tissues, may occur in connection with many different pathological conditions and with surgical or cosmetic procedures. Contracture, for example, of scars, may cause physical problems, which may lead to the need for medical treatment, or it may cause problems of a purely cosmetic nature. Collagen is the major component of scar and other contracted tissue and as such is the most important structural component to consider. Nevertheless, scar and other contracted tissue also include

other structural components, especially other extracellular matrix components, for example, elastin, which may also contribute to contraction of the tissue.

[0360] Contraction of collagen-including tissue, which may also include other extracellular matrix components, frequently occurs in the healing of burns. The burns may be chemical, thermal or radiation burns and may be of the eye, the surface of the skin or the skin and the underlying tissues. It may also be the case that there are burns on internal tissues, for example, caused by radiation treatment. Contraction of burnt tissues is often a problem and may lead to physical and/or cosmetic problems, for example, loss of movement and/or disfigurement.

[0361] Skin grafts may be applied for a variety of reasons and may often undergo contraction after application. As with the healing of burnt tissues the contraction may lead to both physical and cosmetic problems. It is a particularly serious problem where many skin grafts are needed as, for example, in a serious burns case.

[0362] Contraction is also a problem in production of artificial skin. To make a true artificial skin it is necessary to have an epidermis made of epithelial cells (keratinocytes) and a dermis made of collagen populated with fibroblasts. It is important to have both types of cells because they signal and stimulate each other using growth factors. The collagen component of the artificial skin often contracts to less than one tenth of its original area when populated by fibroblasts.


[0363] Cicatricial contraction, contraction due to shrinkage of the fibrous tissue of a scar, is common. In some cases the scar may become a vicious cicatrix, a scar in which the contraction causes serious deformity. A patient's stomach may be effectively separated into two separate chambers in an hour-glass contracture by the contraction of scar tissue formed when a stomach ulcer heals. Obstruction of passages and ducts, cicatricial stenosis, may occur due to the contraction of scar tissue. Contraction of blood vessels may be due to primary obstruction or surgical trauma, for example, after surgery or angioplasty. Stenosis of other hollow visci, for examples, ureters, may also occur. Problems may occur where any form of scarring takes place, whether resulting from accidental wounds or from surgery. Conditions of the skin and tendons which involve contraction of collagen-including tissues include post-trauma conditions resulting from surgery or accidents, for example, hand or foot tendon injuries, post-graft conditions and pathological conditions, such as scleroderma, Dupuytren's contracture and epidermolysis bullosa. Scarring and contraction of tissues in the eye may occur in various conditions, for example, the sequelae of retinal detachment or diabetic eye disease (as mentioned above). Contraction of the sockets found in the skull for the eyeballs and associated structures, including extra-ocular muscles and eyelids, may occur if there is trauma or inflammatory damage. The tissues contract within the sockets causing a variety of problems including double vision and an unsightly appearance.

[0364] For further information on different types of fibrosis see: Molina V, et al., 2002, Harefuah, 141: 973-8, 1009; Yu, et al., 2002, , Curr Opin Pharmacol. 2(2):177-81; Keane, et al., 2003, Am J Kidney Dis. 41: S22-5; Bohle, et al., 1989, Pathol Res Pract. 185:421-40; Kikkawa, et al., 1997, Kidney Int Suppl. 62:S39-40; Bataller et al., 2001, Semin Liver Dis. 21:437-51; Gross, et al., 2001 N Engl J Med. 345:517-25; Frohlich, 2001, Am J Hypertens;14:194S-199S; Friedman, 2003, J Hepatol. 38:S38-53; Albanis, et al., 2003, Curr Gastroenterol Rep. 5:48-56; Weber, 2000, Curr Opin Cardiol. 15:264-72).

Delivery of Nucleic Acid Molecules and Pharmaceutical Formulations

[0365] The retinoid or retinoid conjugate useful for delivery of nucleic acid is in a state in which it is dissolved in or mixed with a medium that can dissolve or retain it.

[0366] Any retinoid or retinoid conjugate may be used in the present description as long as it is actively accumulated by stellate cells; examples of retinoid include, but are not limited to, tretinoin, adapalene, retinol palmitate, and in particular vitamin A, saturated vitamin A, retinoic acid, and retinal. Examples of the retinoid-conjugate include PEG-retinoid conjugates. The present description utilizes the property of stellate cells to positively incorporate a retinoid and/or a retinoid conjugate, and by using the retinoid and/or retinoid conjugate as a drug carrier or by bonding to or being included in another drug carrier component, a desired material or body is transported specifically to stellate cells. A retinoid is a member of the class of compounds having a skeleton in which four isoprenoid units are bonded in a head-to-tail manner. See G. P. Moss, "Biochemical Nomenclature and Related Documents," 2nd Ed. Portland Press, pp. 247-251 (1992). Vitamin A is a generic descriptor for a retinoid qualitatively showing the biological activity of retinol. The retinoid in the present description promotes specific substance delivery to a cancer cell and a CAF (that is, the substance is targeted at these cells). Such a retinoid is not particularly limited, and examples thereof include retinol, Vitamin A, saturated Vitamin A, retinal, retinoic acid, an ester of retinol and a fatty acid, an ester of an aliphatic alcohol and retinoic acid, etretinate, tretinoin, isotretinoin, adapalene, acitretine, tazarotene, and retinol palmitate, and vitamin A analogues such as fenretinide, and bexarotene. Retinoid-conjugates include PEG-conjugates, e.g., diVA-PEG-diVA, shown in the following structure.



[0367] The drug carrier of the present description therefore may contain a drug carrier component other than a retinoid and/or retinoid-conjugate. Such a component is not particularly limited, and any component known in the fields of medicine and pharmacy may be used, but it is preferable for it to be capable of including a retinoid and/or retinoid conjugate. Examples of such a component include a lipid, for example, a phospholipid such as glycerophospholipid, a sphingolipid such as sphingomyelin, a sterol such as cholesterol, a vegetable oil such as soybean oil or poppy seed oil, mineral oil, and a lecithin such as egg-yolk lecithin, but the examples are not limited thereto. Among them, those that can form a liposome are preferable, for example, natural phospholipids such as lecithin, semisynthetic phospholipids such as dimyristoylphosphatidylcholine (DMPC),


dipalmitoylphosphatidylcholine (DPPC), and distearoylphosphatidylcholine (DSPC), and cholesterol.

[0368] Furthermore, the drug carrier of the present description may contain a substance that improves incorporation into stellate cells, for example, retinol-binding protein (RBP).



[0369] The bonding or inclusion of the retinoid and/or retinoid conjugate with the drug carrier of the present description may also be carried out by bonding or including the retinoid and/or retinoid conjugate with another component of the drug carrier by chemical and/or physical methods. Alternatively, bonding or inclusion of the retinoid and/or retinoid conjugate with the drug carrier of the present description may also be carried out by mixing the retinoid and/or retinoid conjugate having formation-affinity and basic components of the drug carrier, into the drug carrier components during preparation of the drug carrier. The amount of retinoid and/or retinoid conjugate bonded to or included in the drug carrier of the present description may be 0.01% to 100% as a ratio by weight relative to the drug carrier components, preferably 0.2% to 20%, and more preferably 1% to 5%.

[0370] Nucleic acid delivery systems may include, for example, aqueous and nonaqueous gels, creams, multiple emulsions, microemulsions, liposomes, ointments, aqueous and nonaqueous solutions, lotions, aerosols, hydrocarbon bases and powders, and can contain excipients such as solubilizers, permeation enhancers (e.g., fatty acids, fatty acid esters, fatty alcohols and amino acids), and hydrophilic polymers (e.g., polycarbophil and polyvinylpyrrolidone). In one embodiment, the pharmaceutically acceptable carrier is a liposome or a transdermal enhancer. Examples of liposomes which can be used in this description include the following:

- CellFectin, a 1:1.5 (M/M) liposome formulation of the cationic lipid N,N^I,N^{II},N^{III}-tetramethyl-N,N^I,N^{II},N^{III}-tetrapalmityl-spermine and dioleoyl phosphatidylethanolamine (DOPE) (GIBCO BRL);
- Cytofectin GSV, a 2:1 (M/M) liposome formulation of a cationic lipid and DOPE (Glen Research);
- DOTAP (N-[1-(2,3-dioleoyloxy)-N,N,N-tri-methyl-ammoniummethylsulfate] (Boehringer Mannheim);
- Lipofectamine, a 3:1 (M/M) liposome formulation of the polycationic lipid DOSPA, the neutral lipid DOPE (GIBCO BRL) and Di-Alkylated Amino Acid (DiLA2);
- Lipotrust, a 4:3:3 (M/M) liposome formulation of O,O'-ditetradecanoyl-N-(α -trimethylammonioacetyl) diethanolamine chloride (DC-6-14, cholesterol and dioleoylphosphatidylethanolamine (Hokkaido System Science). DC-6-14 consists of the following structure.



[0371] Other lipids may be useful: permanent cationic lipids and ionizable cationic lipids, including



and PEG-lipids, including

- 1,2-dimyristeoyl-sn-glycero-3-phosphoethanolamine-N-PEG (PEG-DMPE)
- 1,2-dipalmitoyl-sn-glycero-3-phosphoethanolamine-N-PEG (PEG-DPPE),
- 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-N-PEG (PEG-DSPE), or
- 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine-N-PEG (PEG-DOPE) and/or
- PEG-ceramide.

[0372] Delivery systems may include patches, tablets, suppositories, pessaries, gels and creams, and can contain excipients such as solubilizers and enhancers (e.g., propylene glycol, bile salts and amino acids), and other vehicles (e.g., polyethylene glycol, fatty acid esters and derivatives, and hydrophilic polymers such as hydroxypropylmethylcellulose and hyaluronic acid).

[0373] The drug carrier of the present description may be in any form as long as a desired material or body can be transported to target stellate cells, and examples of the form include, but are not limited to, polymer micelle, liposome, emulsion, microsphere, and nanosphere. Furthermore, the drug carrier of the present description may include in its interior the substance that is to be transported, be attached to the exterior of the substance that is to be transported, or be mixed with the substance that is to be transported as long as the retinoid and/or retinoid conjugate included therein is at least partially exposed on the exterior of the preparation before it reaches the stellate cells at the latest.

[0374] The drug carrier of the present description specifically targets stellate cells and enables a desired effect such as, for example, inhibition or prevention of fibrosis to be exhibited with the maximum effect and minimum side effects by efficiently transporting to stellate cells a desired material or body such as, for example, a drug for controlling the activity or growth of stellate cells. The material or body that the present drug

carrier delivers is not particularly limited, but it preferably has a size that enables physical movement in a living body from an administration site to the liver, pancreas, etc., where stellate cells are present. The drug carrier of the present description therefore can transport not only a material such as an atom, a molecule, a compound, a protein, or a nucleic acid but also a body such as a vector, a virus particle, a cell, a drug release system constituted from one or more elements, or a micromachine. The material or body preferably has the property of exerting some effect on stellate cells, and examples thereof include one that labels stellate cells and one that controls the activity or growth of stellate cells.

[0375] Therefore, in one embodiment of the present description, it is a drug for controlling the activity or growth of stellate cells that the drug carrier delivers. This may be any drug that directly or indirectly inhibits the physicochemical actions of stellate cells involved in the promotion of fibrosis, and examples thereof include, but are not limited to, TGF β activity inhibitors such as a truncated TGF β type II receptor and a soluble TGF β type II receptor, growth factor preparations such as HGF and expression vectors therefor, MMP production promoters such as an MMP gene-containing adenovirus vector, TIMP production inhibitors such as an antisense TIMP nucleic acid, a PPAR γ ligand, cell activation inhibitors and/or cell growth inhibitors such as an angiotensin activity inhibitor, a PDGF activity inhibitor, and a sodium channel inhibitor, and also apoptosis inducers such as compound 861 and gliotoxin, adiponectin, and a compound having Rho kinase inhibitory activity such as (+)-trans-4-(1-aminoethyl)-1-(4-pyridylcarbamoyl)cyclohexane. Furthermore, the 'drug for controlling the activity or growth of stellate cells' in the present description may be any drug that directly or indirectly promotes the physicochemical actions of stellate cells directly or indirectly involved in the inhibition of fibrosis, and examples thereof include, but are not limited to, a drug for promoting a collagen degradation system, e.g., MMP production promoters such as an MMP expression vector, HGF, and drugs having HGF-like activity such as HGF analogues and expression vectors therefor.

[0376] Other examples of the drug for controlling the activity or growth of stellate cells in the present description include a drug for controlling the metabolism of an extracellular matrix such as collagen, for example, a substance having an effect in inhibiting the expression of a target molecule, such as siRNA, ribozyme, and antisense nucleic acid (including RNA, DNA, PNA, and a composite thereof), a substance having a dominant negative effect, and vectors expressing same, that target, for example, an extracellular matrix constituent molecule produced by stellate cells or target one or more molecules that have the function of producing or secreting the extracellular matrix constituent molecule.

[0377] The present description also relates to a medicine for treating a stellate cell-related disorder, the medicine containing the drug carrier and the drug for controlling the activity or growth of stellate cells, and relates to the use of the drug carrier in the production of a pharmaceutical composition for treating a stellate cell-related disorder. The stellate cell-related disorder referred to here means a disorder in which stellate cells are directly or indirectly involved in the process of the disorder, that is, the onset, exacerbation, improvement, remission, cure, etc. of the disorder, and examples thereof include hepatic disorders such as hepatitis, in particular chronic hepatitis, hepatic fibrosis, hepatic cirrhosis, and liver cancer, and pancreatic disorders such as pancreatitis, in particular chronic pancreatitis, pancreatic fibrosis, and pancreatic cancer.

[0378] In the medicine of the present description, the drug carrier may include a drug in its interior, be attached to the exterior of a drug-containing substance, or be mixed with a drug as long as the retinoid and/or retinoid-conjugate included in the drug carrier is at least partially exposed on the exterior of the preparation before it reaches the stellate cells at the latest. Therefore, depending on the route of administration or manner in which the drug is released, the medicine may be covered with an appropriate material, such as, for example, an enteric coating or a material that disintegrates over time, or may be incorporated into an appropriate drug release system.

[0379] The present description therefore includes a drug carrier or medicine preparation kit containing one or more containers containing one or more of a drug carrier constituent, a retinoid and/or a retinoid conjugate, and/or a drug, and also includes an essential component for the drug carrier or the medicine provided in the form of such a kit. The kit of the present description may contain, in addition to those described above, a description, etc. in which a preparation method or an administration method for the drug carrier and the medicine of the present description is described. Furthermore, the kit of the present description may contain all components for completing the drug carrier or the medicine of the present description but need not necessarily contain all of the components. The kit of the present description therefore need not contain a reagent or a solvent that is normally available at a place of medical treatment, an experimental facility, etc. such as, for example, sterile water, saline, or a glucose solution.

[0380] The present description further relates to a method for treating a stellate cell-related disorder, the method including administering an effective amount of the medicine to a subject in need thereof. The effective amount referred to here is an amount that suppresses onset of the target disorder, reduces symptoms thereof, or prevents progression thereof, and is preferably an amount that prevents onset of the target disorder or cures the target disorder. It is also preferably an amount that does not cause an adverse effect that exceeds the benefit from administration. Such an amount may be determined as appropriate by an *in vitro* test using cultured cells, etc. or by a test in a model animal such as a mouse, a rat, a dog, or a pig, and such test methods are well known to a person skilled in the art.

[0381] In the method of the present description, the term 'subject' means any living individual, preferably an animal, more preferably a mammal, and yet more preferably a human individual. In the present description, the subject may be healthy or affected with some disorder, and in the case of treatment of a disorder being intended, the subject typically means a subject affected with the disorder or having a risk of being affected.

[0382] Furthermore, the term 'treatment' includes all types of medically acceptable prophylactic and/or therapeutic intervention for the purpose of the cure, temporary remission, prevention, etc. of a disorder. For example, when the disorder is hepatic fibrosis, the term 'treatment' includes medically acceptable intervention for various purposes including delaying or halting the progression of fibrosis, regression or disappearance of lesions, prevention of the onset of fibrosis, or prevention of recurrence.

[0383] The present description also relates to a method for delivering a drug to stellate cells using the drug carrier. This method includes, but is not limited to, a step of supporting a substance to be delivered on the drug carrier, and a step of administering or adding the drug carrier carrying the substance to be delivered to a stellate cell-containing living body or medium, such as, for example, a culture medium. These steps may be achieved as appropriate in accordance with any known method, the method described in the present specification, etc. This delivery method may be combined with another delivery method, for example, another delivery method in which an organ where stellate cells are present is the target,

etc.

[0384] Nucleic acid molecules may be adapted for use to prevent or treat fibroses (e.g., liver, kidney, peritoneal, and pulmonary) diseases, traits, conditions and/or disorders, and/or any other trait, disease, disorder or condition that is related to or will respond to the levels of hsp47 in a cell or tissue, alone or in combination with other therapies. A nucleic acid molecule may include a delivery vehicle, including liposomes, for administration to a subject, carriers and diluents and their salts, and/or can be present in pharmaceutically acceptable formulations.

[0385] The nucleic acid molecules of the description may include sequences shown in Table 3. Examples of such nucleic acid molecules consist essentially of sequences provided in Table 3.

[0386] The nucleic acid molecules may be administered via pulmonary delivery, such as by inhalation of an aerosol or spray dried formulation administered by an inhalation device or nebulizer, providing rapid local uptake of the nucleic acid molecules into relevant pulmonary tissues. Solid particulate compositions containing respirable dry particles of micronized nucleic acid compositions can be prepared by grinding dried or lyophilized nucleic acid compositions, and then passing the micronized composition through, for example, a 400 mesh screen to break up or separate out large agglomerates. A solid particulate composition comprising the nucleic acid compositions of the description can optionally contain a dispersant which serves to facilitate the formation of an aerosol as well as other therapeutic compounds. A suitable dispersant is lactose, which can be blended with the nucleic acid compound in any suitable ratio, such as a 1 to 1 ratio by weight.

[0387] Aerosols of liquid particles may include a nucleic acid molecules disclosed herein and can be produced by any suitable means, such as with a nebulizer (see e.g., U.S. Pat. No. 4,501,729). Nebulizers are commercially available devices which transform solutions or suspensions of an active ingredient into a therapeutic aerosol mist either by means of acceleration of a compressed gas, typically air or oxygen, through a narrow venturi orifice or by means of ultrasonic agitation. Suitable formulations for use in nebulizers include the active ingredient in a liquid carrier in an amount of up to 40% w/w preferably less than 20% w/w of the formulation. The carrier is typically water or a dilute aqueous alcoholic solution, preferably made isotonic with body fluids by the addition of, e.g., sodium chloride or other suitable salts. Optional additives include preservatives if the formulation is not prepared sterile, e.g., methyl hydroxybenzoate, anti-oxidants, flavorings, volatile oils, buffering agents and emulsifiers and other formulation surfactants. The aerosols of solid particles including the active composition and surfactant can likewise be produced with any solid particulate aerosol generator. Aerosol generators for administering solid particulate therapeutics to a subject produce particles which are respirable, as explained above, and generate a volume of aerosol containing a predetermined metered dose of a therapeutic composition at a rate suitable for human administration. One illustrative type of solid particulate aerosol generator is an insufflator. Suitable formulations for administration by insufflation include finely comminuted powders which can be delivered by means of an insufflator. In the insufflator, the powder, e.g., a metered dose thereof effective to carry out the treatments described herein, is contained in capsules or cartridges, typically made of gelatin or plastic, which are either pierced or opened in situ and the powder delivered by air drawn through the device upon inhalation or by means of a manually-operated pump. The powder employed in the insufflator consists either solely of the active ingredient or of a powder blend comprising the active ingredient, a suitable powder diluent, such as lactose, and an optional surfactant. The active ingredient typically includes from 0.1 to 100 w/w of the formulation. A second type of illustrative aerosol generator includes a metered dose inhaler. Metered dose inhalers are pressurized aerosol dispensers, typically containing a suspension or solution formulation of the active ingredient in a liquefied propellant. During use these devices discharge the formulation through a valve adapted to deliver a metered volume to produce a fine particle spray containing the active ingredient. Suitable propellants include certain chlorofluorocarbon compounds, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane and mixtures thereof. The formulation can additionally contain one or more co-solvents, for example, ethanol, emulsifiers and other formulation surfactants, such as oleic acid or sorbitan trioleate, anti-oxidants and suitable flavoring agents. Other methods for pulmonary delivery are described in, e.g., US20040037780, US6592904, US6582728, and US6565885. WO08132723 relates to aerosol delivery of oligonucleotides in general, and of siRNA in particular, to the respiratory system.

[0388] Nucleic acid molecules may be administered to the central nervous system (CNS) or peripheral nervous system (PNS). Experiments have demonstrated the efficient *in vivo* uptake of nucleic acids by neurons. See e.g., Sommer et al., 1998, *Antisense Nuc. Acid Drug Dev.*, 8:75; Epa et al., 2000, *Antisense Nuc. Acid Drug Dev.*, 10:469; Broaddus et al., 1998, *J. Neurosurg.*, 88:734; Karle et al., 1997, *Eur. J. Pharmacol.*, 340:153; Bannai et al., 1998, *Brain Research*, 784:304; Rajakumar et al., 1997, *Synapse*, 26:199; Wu-pong et al., 1999, *BioPharm*, 12:32; Bannai et al., 1998, *Brain Res. Protoc.*, 3:83; and Simantov et al., 1996, *Neuroscience*, 74:39. Nucleic acid molecules are therefore amenable to delivery to and uptake by cells in the CNS and/or PNS.

[0389] Delivery of nucleic acid molecules to the CNS is provided by a variety of different strategies. Traditional approaches to CNS delivery that can be used include, but are not limited to, intrathecal and intracerebroventricular administration, implantation of catheters and pumps, direct injection or perfusion at the site of injury or lesion, injection into the brain arterial system, or by chemical or osmotic opening of the blood-brain barrier. Other approaches can include the use of various transport and carrier systems, for example though the use of conjugates and biodegradable polymers. Furthermore, gene therapy approaches, e.g., as described in Kaplitt et al., U.S. Pat. No. 6,180,613 and Davidson, WO 04/013280, can be used to express nucleic acid molecules in the CNS.

[0390] Nucleic acid molecules may be formulated or complexed with polyethylenimine (e.g., linear or branched PEI) and/or polyethylenimine derivatives, including for example grafted PEIs such as galactose PEI, cholesterol PEI, antibody derivatized PEI, and polyethylene glycol PEI (PEG-PEI) derivatives thereof (see for example Ogris et al., 2001, *AAPA PharmSci*, 3, 1-11; Furgeson et al., 2003, *Bioconjugate Chem.*, 14, 840-847; Kunath et al., 2002, *Pharm Res* 19:810-17; Choi et al., 2001, *Bull. Korean Chem. Soc.*, 22:46-52; Bettinger et al., 1999, *Bioconjugate Chem.*, 10:558-561; Peterson et al., 2002, *Bioconjugate Chem.* 13:845-54; Erbacher et al., 1999, *J Gene Med* 1:1-18; Godbey et al., 1999., *PNAS*, 96:5177-81; Godbey et al., 1999, *J Controlled Release*, 60:149-60; Diebold et al., 1999, *J Biol Chem*, 274:19087-94; Thomas et al., 2002, *PNAS*, 99, 14640-45; and Sagara, U.S. Pat. No. 6,586,524).

[0391] Nucleic acid molecules may include a bioconjugate, for example a nucleic acid conjugate as described in Vargeese et al., U.S. Ser. No. 10/427,160; U.S. Pat. No. 6,528,631; U.S. Pat. No. 6,335,434; U.S. Pat. No. 6,235,886; U.S. Pat. No. 6,153,737; U.S. Pat. No. 5,214,136; U.S. Pat. No. 5,138,045.

[0392] Compositions, methods and kits disclosed herein may include an expression vector that includes a nucleic acid sequence encoding at least one nucleic acid molecule of the description in a manner that allows expression of the nucleic acid molecule. Methods of introducing nucleic acid molecules or one or more vectors capable of expressing the strands of dsRNA into the environment of the cell will depend on the type of cell and the make up of its environment. The nucleic acid molecule or the vector construct may be directly introduced into the cell (i.e., intracellularly); or introduced extracellularly into a cavity, interstitial space, into the circulation of an organism, introduced orally, or may be introduced by bathing an organism or a cell in a solution containing dsRNA. The cell is preferably a mammalian cell; more preferably a human cell. The nucleic acid molecule of the expression vector can include a sense region and an antisense region. The antisense region can include a sequence complementary to a RNA or DNA sequence encoding hsp47 and the sense region can include a sequence complementary to the antisense region. The nucleic acid molecule can include two distinct strands having complementary sense and antisense regions. The nucleic acid molecule can include a single-strand having complementary sense and antisense regions.

[0393] Nucleic acid molecules that interact with target RNA molecules and down-regulate gene encoding target RNA molecules (e.g., target RNA molecules referred to by Genbank Accession numbers herein) may be expressed from transcription units inserted into DNA or RNA vectors. Recombinant vectors can be DNA plasmids or viral vectors. Nucleic acid molecule expressing viral vectors can be constructed based on, but not limited to, adeno-associated virus, retrovirus, adenovirus, or alphavirus. The recombinant vectors capable of expressing the nucleic acid molecules can be delivered as described herein, and persist in target cells. Alternatively, viral vectors can be used that provide for transient expression of nucleic acid molecules. Such vectors can be repeatedly administered as necessary. Once expressed, the nucleic acid molecules bind and down-regulate gene function or expression via RNA interference (RNAi). Delivery of nucleic acid molecule expressing vectors can be systemic, such as by intravenous or intramuscular administration, by administration to target cells ex-planted from a subject followed by reintroduction into the subject, or by any other means that would allow for introduction into the desired target cell.

[0394] Expression vectors may include a nucleic acid sequence encoding at least one nucleic acid molecule disclosed herein, in a manner which allows expression of the nucleic acid molecule. For example, the vector may contain sequence(s) encoding both strands of a nucleic acid molecule that include a duplex. The vector can also contain sequence(s) encoding a single nucleic acid molecule that is self-complementary and thus forms a nucleic acid molecule. Non-limiting examples of such expression vectors are described in Paul et al., 2002, Nature Biotech 19, 505; Miyagishi et al., 2002, Nature Biotech 19, 497; Lee et al., 2002, Nature Biotech 19, 500; and Novina et al., 2002, Nature Med:10.1038/nm725. Expression vectors may also be included in a mammalian (e.g., human) cell.

[0395] An expression vector may include a nucleic acid sequence encoding two or more nucleic acid molecules, which can be the same or different. Expression vectors may include a sequence for a nucleic acid molecule complementary to a nucleic acid molecule referred to by a Genbank Accession number NM_001235, for example those shown in Table 2.

[0396] An expression vector may encode one or both strands of a nucleic acid duplex, or a single self-complementary strand that self hybridizes into a nucleic acid duplex. The nucleic acid sequences encoding nucleic acid molecules can be operably linked in a manner that allows expression of the nucleic acid molecule (see for example Paul et al., 2002, Nature Biotech, 19:505; Miyagishi and Taira, 2002, Nature Biotech 19:497; Lee et al., 2002, Nature Biotech 19:500; and Novina et al., 2002, Nature Med, 10.1038/nm725).

[0397] An expression vector may include one or more of the following: a) a transcription initiation region (e.g., eukaryotic pol I, II or III initiation region); b) a transcription termination region (e.g., eukaryotic pol I, II or III termination region); c) an intron and d) a nucleic acid sequence encoding at least one of the nucleic acid molecules, wherein said sequence is operably linked to the initiation region and the termination region in a manner that allows expression and/or delivery of the nucleic acid molecule. The vector can optionally include an open reading frame (ORF) for a protein operably linked on the 5' side or the 3'-side of the sequence encoding the nucleic acid molecule; and/or an intron (intervening sequences).

[0398] Transcription of the nucleic acid molecule sequences can be driven from a promoter for eukaryotic RNA polymerase I (pol I), RNA polymerase II (pol II), or RNA polymerase III (pol III). Transcripts from pol II or pol III promoters are expressed at high levels in all cells; the levels of a given pol II promoter in a given cell type depends on the nature of the gene regulatory sequences (enhancers, silencers, etc.) present nearby. Prokaryotic RNA polymerase promoters are also used, providing that the prokaryotic RNA polymerase enzyme is expressed in the appropriate cells (Elroy et al., 1990, PNAS, 87:6743-47; Gao et al., 1993, Nucleic Acids Res 21:2867-72; Lieber et al., 1993, Methods Enzymol., 217:47-66; Zhou et al., 1990, Mol. Cell. Biol. 10:4529-37). Several investigators have demonstrated that nucleic acid molecules expressed from such promoters can function in mammalian cells (e.g. Kashani-Sabet et al., 1992, Antisense Res. Dev., 2:3-15; Ojwang et al., 1992, PNAS 89:10802-06; Chen et al., 1992, Nucleic Acids Res., 20:4581-89; Yu et al., 1993, PNAS, 90:6340-44; L'Huillier et al., 1992, EMBO J., 11:4411-18; Lisiewicz et al., 1993, PNAS 90: 8000-04; Thompson et al., 1995, Nucleic Acids Res., 23:2259; Sullenger et al., 1993, Science, 262:1566). More specifically, transcription units such as the ones derived from genes encoding U6 small nuclear (snRNA), transfer RNA (tRNA) and adenovirus VA RNA are useful in generating high concentrations of desired RNA molecules such as siRNA in cells (Thompson *et al.*, supra; Couture and Stinchcomb, 1996, supra; Noonberg et al., 1994, Nucleic Acid Res. 22:2830; Noonberg et al., U.S. Pat. No. 5,624,803; Good et al., 1997, Gene Ther., 4:45; Beigelman et al., Int'l PCT Publication No. WO 96/18736. The above nucleic acid transcription units can be incorporated into a variety of vectors for introduction into mammalian cells, including but not restricted to, plasmid DNA vectors, viral DNA vectors (such as adenovirus or adeno-associated virus vectors), or viral RNA vectors (such as retroviral or alphavirus vectors) (see Couture and Stinchcomb, 1996 supra).

[0399] Nucleic acid molecule may be expressed within cells from eukaryotic promoters (e.g., Izant and Weintraub, 1985, Science, 229, 345; McGarry and Lindquist, 1986, PNAS 83, 399; Scanlon et al., 1991, PNAS 88:10591-95; Kashani-Sabet et al., 1992, Antisense Res. Dev., 2:3-15; Dropulic et al., 1992, J. Virol., 66:1432-41; Weerasinghe et al., 1991, J. Virol., 65:5531-34; Ojwang et al., 1992, PNAS, 89:10802-06; Chen et al., 1992, Nucleic Acids Res., 20:4581-89; Sarver et al., 1990 Science, 247:1222-25; Thompson et al., 1995, Nucleic Acids Res., 23:2259; Good et al., 1997, Gene Therapy, 4:45. Those skilled in the art realize that any nucleic acid can be expressed in eukaryotic cells from the appropriate DNA/RNA vector. The activity of such nucleic acids can be augmented by their release from the primary transcript by a enzymatic nucleic acid (Draper et al., PCT WO 93/23569, and Sullivan et al., PCT WO 94/02595; Ohkawa et al., 1992, Nucleic Acids Symp. Ser., 27:15-6; Taira et al., 1991, Nucleic Acids Res., 19:5125-30; Ventura et al., 1993, Nucleic Acids Res., 21:3249-55; Chowrira et al., 1994, J. Biol. Chem., 269:25856.

[0400] A viral construct packaged into a viral particle would accomplish both efficient introduction of an expression construct into the cell and transcription of dsRNA construct encoded by the expression construct.

[0401] Methods for oral introduction include direct mixing of RNA with food of the organism, as well as engineered approaches in which a species that is used as food is engineered to express an RNA, then fed to the organism to be affected. Physical methods may be employed to introduce a nucleic acid molecule solution into the cell. Physical methods of introducing nucleic acids include injection of a solution containing the nucleic acid molecule, bombardment by particles covered by the nucleic acid molecule, soaking the cell or organism in a solution of the RNA, or electroporation of cell membranes in the presence of the nucleic acid molecule.

[0402] Other methods known in the art for introducing nucleic acids to cells may be used, such as lipid-mediated carrier transport, chemical mediated transport, such as calcium phosphate, and the like. Thus the nucleic acid molecules may be introduced along with components that perform one or more of the following activities: enhance RNA uptake by the cell, promote annealing of the duplex strands, stabilize the annealed strands, or other-wise increase inhibition of the target gene.

Dosages

[0403] The useful dosage to be administered and the particular mode of administration will vary depending upon such factors as the cell type, or for *in vivo* use, the age, weight and the particular animal and region thereof to be treated, the particular nucleic acid and delivery method used, the therapeutic or diagnostic use contemplated, and the form of the formulation, for example, suspension, emulsion, micelle or liposome, as will be readily apparent to those skilled in the art. Typically, dosage is administered at lower levels and increased until the desired effect is achieved.

[0404] When lipids are used to deliver the nucleic acid, the amount of lipid compound that is administered can vary and generally depends upon the amount of nucleic acid being administered. For example, the weight ratio of lipid compound to nucleic acid is preferably from about 1:1 to about 30:1, with a weight ratio of about 5:1 to about 15:1 being more preferred.

[0405] A suitable dosage unit of nucleic acid molecules may be in the range of 0.001 to 0.25 milligrams per kilogram body weight of the recipient per day, or in the range of 0.01 to 20 micrograms per kilogram body weight per day, or in the range of 0.01 to 10 micrograms per kilogram body weight per day, or in the range of 0.10 to 5 micrograms per kilogram body weight per day, or in the range of 0.1 to 2.5 micrograms per kilogram body weight per day.

[0406] Suitable amounts of nucleic acid molecules may be introduced and these amounts can be empirically determined using standard methods. Effective concentrations of individual nucleic acid molecule species in the environment of a cell may be about 1 femtomolar, about 50 femtomolar, 100 femtomolar, 1 picomolar, 1.5 picomolar, 2.5 picomolar, 5 picomolar, 10 picomolar, 25 picomolar, 50 picomolar, 100 picomolar, 500 picomolar, 1 nanomolar, 2.5 nanomolar, 5 nanomolar, 10 nanomolar, 25 nanomolar, 50 nanomolar, 100 nanomolar, 500 nanomolar, 1 micromolar, 2.5 micromolar, 5 micromolar, 10 micromolar, 100 micromolar or more.

[0407] Dosage levels of the order of from about 0.1 mg to about 140 mg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5 mg to about 7 g per subject per day). The amount of active ingredient that can be combined with the carrier materials to produce a single dosage form varies depending upon the host treated and the particular mode of administration. Dosage unit forms generally contain between from about 1 mg to about 500 mg of an active ingredient.

[0408] It is understood that the specific dose level for any particular subject depends upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

[0409] Pharmaceutical compositions that include the nucleic acid molecule disclosed herein may be administered once daily, qid, tid, bid, QD, or at any interval and for any duration that is medically appropriate. However, the therapeutic agent may also be dosed in dosage units containing two, three, four, five, six or more sub-doses administered at appropriate intervals throughout the day. In that case, the nucleic acid molecules contained in each sub-dose may be correspondingly smaller in order to achieve the total daily dosage unit. The dosage unit can also be compounded for a single dose over several days, *e.g.*, using a conventional sustained release formulation which provides sustained and consistent release of the dsRNA over a several day period. Sustained release formulations are well known in the art. The dosage unit may contain a corresponding multiple of the daily dose. The composition can be compounded in such a way that the sum of the multiple units of a nucleic acid together contain a sufficient dose.

Pharmaceutical compositions, kits, and containers

[0410] Also provided are compositions, kits, containers and formulations that include a nucleic acid molecule (*e.g.*, an siRNA molecule) as provided herein for reducing expression of hsp47 for administering or distributing the nucleic acid molecule to a patient. A kit may include at least one container and at least one label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers can be formed from a variety of materials such as glass, metal or plastic. The container can hold amino acid sequence(s), small molecule(s), nucleic acid sequence(s), cell population(s) and/or antibody(s). In one embodiment, the container holds a polynucleotide for use in examining the mRNA expression profile of a cell, together with reagents used for this purpose. In another embodiment a container includes an antibody, binding fragment thereof or specific binding protein for use in evaluating hsp47 protein expression cells and tissues, or for relevant laboratory, prognostic, diagnostic, prophylactic and therapeutic purposes; indications and/or directions for such uses can be included on or with such container, as can reagents and other compositions or tools used for these purposes. Kits may further include associated indications and/or directions; reagents and

other compositions or tools used for such purpose can also be included.

[0411] The container can alternatively hold a composition that is effective for treating, diagnosis, prognosing or prophylaxing a condition and can have a sterile access port (for example the container can be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agents in the composition can be a nucleic acid molecule capable of specifically binding hsp47 and/or modulating the function of hsp47.

[0412] A kit may further include a second container that includes a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and/or dextrose solution. It can further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, stirrers, needles, syringes, and/or package inserts with indications and/or instructions for use.

[0413] The units dosage ampoules or multidose containers, in which the nucleic acid molecules are packaged prior to use, may include an hermetically sealed container enclosing an amount of polynucleotide or solution containing a polynucleotide suitable for a pharmaceutically effective dose thereof, or multiples of an effective dose. The polynucleotide is packaged as a sterile formulation, and the hermetically sealed container is designed to preserve sterility of the formulation until use.

[0414] The container in which the polynucleotide including a sequence encoding a cellular immune response element or fragment thereof may include a package that is labeled, and the label may bear a notice in the form prescribed by a governmental agency, for example the Food and Drug Administration, which notice is reflective of approval by the agency under Federal law, of the manufacture, use, or sale of the polynucleotide material therein for human administration.

[0415] Federal law requires that the use of pharmaceutical compositions in the therapy of humans be approved by an agency of the Federal government. In the United States, enforcement is the responsibility of the Food and Drug Administration, which issues appropriate regulations for securing such approval, detailed in 21 U.S.C. § 301-392. Regulation for biologic material, including products made from the tissues of animals is provided under 42 U.S.C. § 262. Similar approval is required by most foreign countries. Regulations vary from country to country, but individual procedures are well known to those in the art and the compositions and methods provided herein preferably comply accordingly.

[0416] The dosage to be administered depends to a large extent on the condition and size of the subject being treated as well as the frequency of treatment and the route of administration. Regimens for continuing therapy, including dose and frequency may be guided by the initial response and clinical judgment. The parenteral route of injection into the interstitial space of tissues is preferred, although other parenteral routes, such as inhalation of an aerosol formulation, may be required in specific administration, as for example to the mucous membranes of the nose, throat, bronchial tissues or lungs.

[0417] As such, provided herein is a pharmaceutical product which may include a polynucleotide including a sequence encoding a cellular immune response element or fragment thereof in solution in a pharmaceutically acceptable injectable carrier and suitable for introduction interstitially into a tissue to cause cells of the tissue to express a cellular immune response element or fragment thereof, a container enclosing the solution, and a notice associated with the container in form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of manufacture, use, or sale of the solution of polynucleotide for human administration.

Indications

[0418] The nucleic acid molecules disclosed herein can be used to treat diseases, conditions or disorders associated with hsp47, such as liver fibrosis, cirrhosis, pulmonary fibrosis, kidney fibrosis, peritoneal fibrosis, chronic hepatic damage, and fibrillogenesis and any other disease or conditions that are related to or will respond to the levels of hsp47 in a cell or tissue, alone or in combination with other therapies. As such, compositions, kits and methods disclosed herein may include packaging a nucleic acid molecule disclosed herein that includes a label or package insert. The label may include indications for use of the nucleic acid molecules such as use for treatment or prevention of liver fibrosis, peritoneal fibrosis, kidney fibrosis and pulmonary fibrosis, and any other disease or conditions that are related to or will respond to the levels of hsp47 in a cell or tissue, alone or in combination with other therapies. A label may include an indication for use in reducing expression of hsp47. A "package insert" is used to refer to instructions customarily included in commercial packages of therapeutic products, that contain information about the indications, usage, dosage, administration, contraindications, other therapeutic products to be combined with the packaged product, and/or warnings concerning the use of such therapeutic products, etc.

[0419] Those skilled in the art will recognize that other anti-fibrosis treatments, drugs and therapies known in the art can be readily combined with the nucleic acid molecules herein (e.g. siRNA molecules) and are hence contemplated herein.

[0420] The methods and compositions provided herein will now be described in greater detail by reference to the following non-limiting examples.

EXAMPLE 1 Preparation of siRNA for gp46

[0421] Among optimal sequences for siRNA recognition in targeting a base sequence of HSP47, which is a common molecular chaperone for collagens (types I to IV), Sequences A and B were prepared in accordance with an siRNA oligo design program by iGENE Therapeutics, Inc. Sequence C was prepared by searching on the Internet using the siRNA Target Finder from Ambion, Inc. and selecting 19 base sequences that would become a target for rat gp46 (human HSP47 homolog, GenBank Accession No. M69246). When carrying out the design, care was taken in starting at 75 to 100 bases downstream from the initiation codon, positioning the first AA dimer, and making sure that the GC content was 30%

to 70%. In this example, siRNAs having the sequences below were prepared.

1. A: GUUCCACCAUAAGAUGGUAGACAAAC
2. B: CCACAAGUUUUUAUCCAUCUAGC
3. C: GAAACCUGUAGAGGCCGCA

EXAMPLE 2 Inhibition of gp46 Expression by Prepared siRNA

[0422] Normal rat kidney cells (NRK cells), which had rat gp46 and were fibroblasts producing collagen, were transfected with 0.1 nM to 50 nM siRNA and cultured for 12 to 48 hours (FIG. 1). The amount of expression of gp46 was checked by the western blot method (FIGS. 2 to 4, upper band corresponding to gp46, lower band corresponding to actin control). All of the siRNAs inhibited the expression of gp46 protein remarkably compared with a vehicle (FIG. 2). In the experiment below, siRNA Sequence A, which showed the strongest effect, was used. Inhibition by siRNA was concentration dependent (FIG. 3); protein expression by gp46 was about 90% inhibited by 50 nM siRNA at 48 hours (FIG. 4).

EXAMPLE 3 Inhibition of Collagen Synthesis by Prepared siRNA

[0423] In order to examine the amount of collagen synthesized, ^3H -proline was added to the culture supernatant of rat fibroblasts (NRK cells) under the above-mentioned conditions (siRNA concentration 50 nM, time 48 hours), and after transfection the amount of ^3H in secreted protein was examined (FIG. 5). The amount of collagen synthesized was calculated from the ratio of protein secreted in the supernatant to protein degraded by collagenase when culturing gp46siRNA-transfected fibroblasts in the presence of ^3H -proline in accordance with Peterkofsky et al., 1971 Biochemistry 10:988-94.

$$\text{Collagen synthesis ratio} = \frac{\text{collagenase - sensitive fraction}}{\text{collagenase - insensitive fraction} + \text{collagenase - sensitive fraction}} \times 100 \quad (5.4 \times)$$

[0424] The collagen synthesis ratio in rat fibroblasts decreased by about 40% compared with a control group (FIG. 6).

EXAMPLE 4 Specific Transfection of Nucleic Acid into HSC

[0425] An emulsion (VA-Lip-GFP) was prepared by mixing GFP expression plasmid and liposome-encapsulated VA formed by mixing 10% Vitamin A (VA) and liposome. Cationic liposomes containing O,O'-ditetradecanoyl-N-(α -trimethylammonioacetyl) diethanolamine chloride (DC-6-14) as a cationic lipid, cholesterol, and dioleoylphosphatidylethanolamine at a molar ratio of 4:3:3 were dissolved in a chloroform-methanol mixture (4:1, v:v), and the solvent was removed by evaporation under vacuum. The lipids were mixed with an aqueous 9% sucrose solution and hydrated at 60° C. and homogenized to uniformity. The dispersion was extruded twice through a polyvinylidenedifluoride membrane filter with a 0.22 μm pore size. The dispersion was aliquotted into glass vials and frozen and then lyophilized. The dried liposomes were reconstituted with distilled water at a concentration of 1 mM DC-6-14 under vortexing before use. Specifically, 25 mg of Vitamin A was first dissolved in 87 μL of DMSO thus to give a 100 mM stock solution. To prepare VA-coupled liposomes, 200 nmol of VA dissolved in DMSO was mixed with 100 nmole of DC-6-14 by vortexing at room temperature. The VA-siRNA-liposomes were intraportally administered to a rat, hepatic tissue was collected and fixed. The emulsion was prepared by supposing that the amount of plasma for a 200 g rat was about 10 mL, and setting the concentrations of VA and GFP in portal blood at 10 μM . 1 μL of this VA stock solution was mixed with 10 μL of the VA-liposomes and 179 μL of PBS, 10 μg of GFP expression plasmid was further added thereto to give a total of 200 μL , and the mixture was vortexed for three minutes to give VA-Lip-GFP. The abdomen of an SD rat was opened, and the VA-Lip-GFP was slowly injected into a peripheral portal vein. Forty-eight hours after the injection, hepatic tissue was harvested. Since compared with other hepatic cells intermediate filament desmin is specifically expressed in HSC, when fixed hepatic tissue was stained with Alexa Fluor 568-labeled anti-desmin antibody, and a fluorescence double image with GFP was examined, it was confirmed that GFP was expressed within the HSC (FIG. 7). For untreated controls and a group to which the GFP expression plasmid vector alone was administered, expression in rat HSC was not observed, but in a group to which VA-Lip-GFP was administered, expression of GFP was observed specifically in stellate cells.

EXAMPLE 5 Quantitative Analysis of Nucleic Acid Transfection Rate

[0426] In the same manner as in Example 4, except that FITC-labeled gp46siRNA was used instead of the GFP expression plasmid, an emulsion (VA-Lip-gp46siRNA (FITC)) containing VA-encapsulated liposome and FITC-labeled gp46siRNA was prepared. A solution of siRNA_{gp46} (580 pmole/ μL in distilled water) was added to the VA-coupled liposome solution of Example 4 with stirring at room temperature. The ratio of siRNA to DC-6-14 was 1:11.5 (mol/mol) and the siRNA to liposome ratio (wt/wt) was 1:1. Free VA or siRNA was removed by micropartition using VIVASPIN concentrator, 30K MWCO, by three passes. Material trapped on the membrane was reconstituted with PBS and intraportally administered to an SD rat (10 μg as the amount of siRNA/200 μL). Forty-eight hours after administration hepatic tissue was harvested, αSMA (smooth muscle actin), which compared with other hepatic cells is expressed specifically in HSC, was stained with Alexa Fluor 568-labeled anti- αSMA antibody, cell nuclei were stained with DAPI, and a fluorescence image was examined by a confocal laser scanning microscope (LSM). As shown on the left-hand side of FIG. 8, in a group to which VA-Lip-gp46siRNA (FITC) was administered, a large number of cells emitting both green fluorescence due to FITC and red fluorescence due to Alexa Fluor 568 were observed, and when a quantitative analysis was carried out by NIH Image (the number of cells was counted by selecting any 10 fields from a X1000 fluorescence microscope photograph), the transfection efficiency was 77.6% (average of 10

fields). On the other hand, in a group to which Lip-gp46siRNA (FITC) containing no VA was administered, the transfection efficiency was a low value of 14.0% and, moreover, transfection into cells other than stellate cells was observed at 3.0% (right-hand side of FIG. 8). It has been found from the results above that the transfection efficiency into stellate cells is increased remarkably by including VA.

EXAMPLE 6 Inhibition of Expression of gp46 by VA-Lip-gp46siRNA

[0427] With regard to another section of the tissue harvested in Example 5, gp46 was stained with Alexa Fluor 568-labeled anti-HSP47 antibody and cell nuclei were stained with DAPI, and a fluorescence image was examined by a confocal laser scanning microscope. As shown in FIG. 9, it was observed that in a group to which VA-Lip-gp46siRNA was administered, expression of gp46, which can be observed as a red fluorescence (right-hand side in the figure), was markedly reduced compared with a control group to which was administered VA-Lip-random siRNA containing random siRNA, which was not specific to gp46 (left-hand side in the figure). The expression inhibition rate relative to an average of six fields of the control group was 75%, which was extremely high, when the number of gp46-negative cells was examined by selecting any ten fields from a X1000 fluorescence microscope photograph using NIH Image in the same manner as in Example 7.

EXAMPLE 7 Treatment of LC Rat (Intraportal Administration 1)

[0428] In accordance with a report by Jezequel *et al.* (Jezequel *et al.*, 1987, J Hepatol. 5:174-81), an LC model rat was prepared using dimethylnitrosamine (DMN) (FIG. 10). Specifically, a 1 mL/kg dose of 1% DMN (intraperitoneal administration) was administered to a five week-old SD rat (male) three straight days per week. As already reported, an increase in fiber was observed from the 2nd week, and in the 4th week this was accompanied by the findings of marked fibrosis, destruction of hepatic lobule structure, and formation of regenerative nodules being observed (FIG. 11). Then, by the same method as in Examples 4 and 5, an emulsion (VA-Lip-gp46siRNA) was prepared by formulating gp46siRNA as a liposome and mixing with 10% VA, and was administered. Administration of VA-Lip-gp46siRNA was started in the 3rd week, by which time sufficient fibrosis was observed, and evaluation was carried out in the 4th and 5th weeks. Since it was confirmed by Example 2 that the effects were observed for up to 48 hours *in vitro*, administration was carried out twice a week (FIG. 11). The amount administered was determined in accordance with a report in which siRNA was directly injected (McCaffery *et al.*, 2002, Nature 418: 38-39), and was 40 µg as the total amount of siRNA. From azan staining of the liver after administration of siRNA, in the 4th week there was no apparent difference between a group to which saline had been administered, a group to which siRNA (random) had been administered, and a group to which siRNA (gp46) had been administered, but in the 5th week a decrease in the amount of fiber was observed for the group to which gp46siRNA had been administered (FIG. 12). In order to quantitatively analyze the amount of fiber, an unstained portion was extracted using NIH Image, its area was measured (FIG. 13), and a significant decrease in the area of collagen was observed for the group to which gp46siRNA had been administered (FIG. 14). Furthermore, in order to evaluate the degree of fibrosis using another measure, the amount of hydroxyproline, which is an indicator for fibrosis, was quantitatively measured by a standard method. Specifically, after 20 mg of freeze-dried hepatic tissue was hydrolyzed with HCl for 24 hours, the reaction liquid was centrifuged, and the supernatant was treated with a reagent such as Ehrlich's solution and centrifuged. The supernatant was recovered, and the amount of hydroxyproline in the hepatic tissue was measured by measuring the absorbance at 560 nm (Hepatology 1998, 28:1247-52). As shown in FIG. 15, in the group to which gp46siRNA had been administered, the amount of hydroxyproline became very small.

EXAMPLE 8 Treatment of LC Rat (Intraportal Administration 2)

[0429] Furthermore, in order to examine a change in the survival rate by administration of the medicine of the present description, in accordance with a method by Qi Z *et al.* (PNAS 1999; 96:2345-49), an LC model rat was prepared using DMN in an amount that was increased by 20% over the normal amount. In this model, a total of four intraportal administrations were carried out in the first and second weeks. Administration details were: PBS, Lip-gp46siRNA, VA-Lip-random siRNA, and VA-Lip-gp46siRNA (n=7 for each group). After the third week, all of the controls (the group to which PBS had been administered, the group to which VA-Lip-random siRNA had been administered, and the group to which Lip-gp46siRNA had been administered) were dead, but 6 out of 7 survived for the group to which VA-Lip-gp46siRNA had been administered (FIG. 16). Furthermore, in azan staining of the liver on the 21st day, an apparent decrease in the amount of fiber was observed for the group to which gp46siRNA had been administered (FIG. 17).

EXAMPLE 9 Treatment of LC Rat (Intraportal Administration 3)

[0430] In another experiment, intraportal administration was carried out from the 3rd week for LC model rats (1% DMN 1 mg/kg intraperitoneally administered 3 times a week) prepared in accordance with the method by Qi *et al.* and a method by Ueki *et al.*, 1999, Nat Med. 5:226-30, as shown in Table 2 below (n=6 for each group). PBS was added to each substance to be administered so as to make a total volume of 200 µL, and the frequency of administration was once a week.

TABLE 2

Treatment group	Content of Administration	Dosage	Frequency of Administration
10-1	VA	VA 200 nmol	Twice a week 10-2
10-2	Lip-gp46siRNA	liposome 100 nmol; gp46siRNA 100 µg	
10-3	VA-Lip-random siRNA	VA 200 nmol; siRNA liposome 100 nmol; random-100 µg	
10-4	VA-Lip-gp46siRNA	VA 200 nmol; liposome 100 nmol; gp46siRNA 100 µg	
10-5	PBS	200 µL	Three times a week

Treatment group	Content of Administration	Dosage	Frequency of Administration
10-6	VA	VA 200 nmol	
10-7	VA-Lip VA	200 nmol; liposome 100 nmol	
10-8	Lip-gp46siRNA	liposome 100 nmol; gp46siRNA 150 µg	
10-9	VA-Lip-random siRNA	VA 200 nmol; liposome 100 nmol; random-siRNA 150 µg	
10-10	VA-Lip-gp46siRNA	VA 200 nmol; liposome 100 nmol; gp46siRNA 150 µg	

[0431] From the results, in the groups other than the group to which the medicine of the present description had been administered (treatment group 9-4), all 6 rats were dead by the 45th day after starting administration of DMN, but in the group to which the medicine of the present description had been administered, all of the individuals apart from one case, which was dead on the 36th day, survived for more than 70 days after starting administration of DMN (FIG. 18). For the dead individuals, the amount of hepatic fiber was quantitatively analyzed based on the area of collagen in the same manner as in Example 7, and the increase in the amount of hepatic fiber was remarkably inhibited by administration of VA-Lip-gp46siRNA (FIG. 19).

EXAMPLE 10 Treatment of LC Rat (Intravenous Administration)

[0432] Intravenous administration was carried out from the 3rd week for LC model rats (1% DMN 1 µg/BW (g) intraperitoneally administered 3 times a week) prepared in the same manner as in Example 9, as shown in the table below (n=6 for each group). PBS was added to each substance to be administered so as to make a total volume of 200 µL. The administration period was up to death except that it was up to the 7th week for Group 10-4 and the 6th week for Group 10-10.

[0433] From the results, in the groups other than the groups to which the medicine of the present description had been administered (treatment groups 10-4 and 10-10), all 6 rats were dead by the 45th day after starting administration of DMN, but in the groups to which the medicine of the present description had been administered, all of the individuals, apart from a case in which two rats were dead on the 45th day in treatment group 10-4, survived for more than 70 days after starting administration of DMN (FIGS. 20 and 21). For the dead individuals, the amount of hepatic fiber was quantitatively analyzed in the same manner as in Example 7, and the increase in the amount of hepatic fiber was remarkably inhibited by administration of VA-Lip-gp46siRNA (FIG. 22).

[0434] The above-mentioned results show that the medicine of the present description is extremely effective for the prevention and treatment of fibrosis, in which stellate cells are involved.

EXAMPLE 11 Improvement of Results by RBP (Retinol-Binding Protein)

[0435] The influence of RBP on VA-Lip-gp46siRNA transfection efficiency was examined using LI90, a cell line derived from human HSC. 100 nM of VA-Lip-gp46siRNA (FITC) prepared in Example 5, together with various concentrations (i.e. 0, 0.1, 0.5, 1, 2, 4, or 10%) of FBS (fetal bovine serum), were added to LI90 during culturing and incubated for 48 hours, a fluorescence image was observed by LSM, and the amount of siRNA incorporated into individual cells was quantitatively analyzed by FACS. FBS contained about 0.7 mg/dL of RBP. As shown in FIG. 23, FBS (RBP) gave a concentration-dependent increase in the amount of siRNA transfection. Subsequently, 100 nM of VA-Lip-gp46siRNA (FITC) and 4% FBS, together with 10 µg (21.476 nmol) of anti-RBP antibody, were added to LI90 during culturing, and the siRNA transfection efficiency was evaluated in the same manner. As shown in FIG. 24, the increase in the amount of transfection by RBP was markedly decreased by the addition of anti-RBP antibody. The above-mentioned results show that RBP is effective in further enhancing transfection of the medicine of the present description.

EXAMPLE 12 Selecting hsp47 Nucleic Acid Molecule Sequences

[0436] Nucleic acid molecules (e.g., siRNA ≤ 25 nucleotides) against Hsp47 were designed using several computer programs including siRNA at Whitehead Institute for Biomedical Research, siRNA Design (Integrated DNA Technologies), BLOCK-IT RNAi Designer (Invitrogen), siDESIGN Center (Dharmacon), and BIOPREDSi (Friedrich Miescher Institute for Biomedical Research). The sequences of top scored siRNAs from these programs were compared and selected (see Table 3) based on the algorithms as well as the sequence homology between human and rat. Candidate sequences were validated by *in vitro* knocking down assays.

[0437] Several parameters were considered for selecting a nucleic acid molecule (e.g., a 21-mer siRNA) sequence. Exemplary parameters include:

- thermodynamic stability (RISC favors the strand with less stable 5'-end)
- 30-52% GC content
- positional nucleotide preference:
(C/G)₁NNNNNNNN(A/U)₁₀NNNNNNNN(A/U)₁₉
where N is any nucleotide
- devoid of putative immunostimulatory motifs
- 2-nucleotide 3' overhang

- position of siRNA within the transcript (preferably within cDNA region)
- sequence specificity (checked by using BLAST)
- variations in single nucleotide by checking SNP database

[0438] iRNA sequences having ≤ 25 nucleotides were designed based on the foregoing methods. Corresponding Dicer substrate siRNA (e.g., ≥ 26 nucleotides) were designed based on the smaller sequences and extend the target site of the siRNA ≤ 25 nucleotide by adding four bases to the 3'-end of the sense strand and 6 bases to the 5'-end of the antisense strand. The Dicer substrates that were made generally have a 25 base sense strand a 27 base antisense strand with an asymmetric blunt ended and 3'-overhang molecule. The sequences of the sense and the antisense strand without base modification (base sequence) and with modifications (experimental sequence) are provided in Table 3.

TABLE 3

siRNA	Target region		Base sequence (corresponding nucleotides of SEQ ID NO: 1)	Experimental sequence (corresponding nucleotides of SEQ ID NO: 1)
siHSP47-C	human/rat hsp47	sense	5'GGACAGGCCUCUACAACUAAU (SEQ ID NO: 3) [945-963]	5'GGACAGGCCUCUACAACUAdTdT (SEQ ID NO: 5) [945-963]
		antisense	5' UAGUUGUAGAGGCCUGUCCUU (SEQ ID NO: 4) [945-963]	5' UAGUUGUAGAGGCCUGUCCdTdT (SEQ ID NO: 6) [945-963]
siHSP47-Cd	human/rat hsp47	sense	5'GGACAGGCCUCUACAACUACUACGA (SEQ ID NO: 7) [945-969]	5'GGACAGGCCUCUACAACUACUACdGdA (SEQ ID NO: 9) [945-969]
		antisense	5'UCGUAGUAGUUGUAGAGGCCUGUCCUU (SEQ ID NO: 8) [945-969]	5'UCGUAGUAGUUGUAGAGGCCUGUCCUU (SEQ ID NO: 10) [945-969]
siHSP47-1	human/rat hsp47	sense	5'CAGGCCUCUACAACUACUAAU (SEQ ID NO: 11) [948-966]	5'CAGGCCUCUACAACUACUAdTdT (SEQ ID NO: 13) [948-966]
		antisense	5'UAGUAGUUGUAGAGGCCUGUU (SEQ ID NO: 12) [948-966]	5'UAGUAGUUGUAGAGGCCUGdTdT (SEQ ID NO: 14) [948-966]
siHSP47-1d	human hsp47	sense	5'CAGGCCUCUACAACUACUACGACGA (SEQ ID NO: 15) [948-972]	5'CAGGCCUCUACAACUACUACGdGdA (SEQ ID NO: 17) [948-972]
		antisense	5'CGUCGUAGUAGUUGUAGAGGCCUGUU (SEQ ID NO: 16) [948-972]	5'CGUCGUAGUAGUUGUAGAGGCCUGUU (SEQ ID NO: 18) [948-972]
siHsp47-2	human hsp47	sense	5'GAGCACUCCAAGAUCAACUUU (SEQ ID NO: 19) [698-717]	5'GAGCACUCCAAGAUCAACUdTdT (SEQ ID NO: 21) [698-717]
		antisense	5'AGUUGAUCUUGGAGUGCUCUU (SEQ ID NO: 20) [698-716]	5' AGUUGAUCUUGGAGUGCUCdTdT (SEQ ID NO: 22) [698-716]
siHsp47-2d	human hsp47	sense	5'GAGCACUCCAAGAUCAACUCCGCG (SEQ ID NO: 23) [698-722]	5'GAGCACUCCAAGAUCAACUCCGdCdG (SEQ ID NO: 25) [698-722]
		antisense	5'CGCGGAAGUUGAUCUUGGAGUGCUCUU (SEQ ID NO: 24) [698-722]	5'CGCGGAAGUUGAUCUUGGAGUGCUCUU (SEQ ID NO: 26) [698-722]
siHsp47-2d rat	rat Gp46	sense	5'GAACACUCCAAGAUCAACUCCGAG (SEQ ID NO: 27) [587-611]	5'GAACACUCCAAGAUCAACUCCGdAdG (SEQ ID NO: 29) [587-611]
		antisense	5'UCGGAAGUUGAUCUUGGAGUGUUCUU (SEQ ID NO: 28) [587-611]	5'UCGGAAGUUGAUCUUGGAGUGUUCUU (SEQ ID NO: 30) [587-611]
siHsp47-3	human hsp47	sense	5'CUGAGGCCAUUGACAAGAAUU (SEQ ID NO: 31) [1209-1227]	5'CUGAGGCCAUUGACAAGAAAdTdT (SEQ ID NO: 33) [1209-1227]
		antisense	5'UUCUUGUCAAUUGGCCUCAGUU (SEQ ID NO: 32) [1209-1227]	5' UUCUUGUCAAUUGGCCUCAGdTdT (SEQ ID NO: 34) [1209-1227]
siHsp47-3d	human hsp47	sense	5'CUGAGGCCAUUGACAAGAAAGGC (SEQ ID NO: 35) [1209-1233]	5' CUGAGGCCAUUGACAAGAAAGdGdC (SEQ ID NO: 37) [1209-1233]
		antisense	5'CCUUGUUCUUGUCAUUGGCCUCAGUU (SEQ ID NO: 36) [1209-1233]	5'GCCUUGUUCUUGUCAUUGGCCUCAGUU (SEQ ID NO: 38) [1209-1233]
siHsp47-4	human hsp47	sense	5'CUACGACGACGAGAAGGAAUU (SEQ ID NO: 39) [964-982]	5' CUACGACGACGAGAAGGAAAdTdT (SEQ ID NO: 41) [964-982]
		antisense	5'UCCUUCUCGUCGUCGUAGUU (SEQ ID NO: 40) [964-982]	5' UCCUUCUCGUCGUCGUAGdTdT (SEQ ID NO: 42) [964-982]
siHsp47-4d	human hsp47	sense	5'CUACGACGACGAGAAGGAAAAGCUG (SEQ ID NO: 43) [964-988]	5' CUACGACGACGAGAAGGAAAAGCdTdG (SEQ ID NO: 45) [964-988]
		antisense	5'AGCUUUUCCUUCUCGUCGUAGUU (SEQ ID NO: 44) [964-988]	5' AGCUUUUCCUUCUCGUCGUAGUU (SEQ ID NO: 46) [964-988]
siHsp47-5	human hsp47	sense	5'GCCACACUGGGAUGAGAAUU (SEQ ID NO: 47) [850-870]	5' GCCACACUGGGAUGAGAAAdTdT (SEQ ID NO: 49) [850-870]
		antisense	5' UUCUCAUCCAGUGUGGCUU (SEQ ID NO: 48) [850-868]	5' UUCUCAUCCAGUGUGGCUdTdT (SEQ ID NO: 50) [850-868]
siHsp47-6	human hsp47	sense	5'GCAGCAAGCAGCACUACAAUU (SEQ ID NO: 51) [675-693]	5' GCAGCAAGCAGCACUACAAAdTdT (SEQ ID NO: 53) [675-693]
		antisense	5' UUGUAGUGCUGCUUGCUGCUU (SEQ ID NO: 52) [675-693]	5' UUGUAGUGCUGCUUGCUGCUdTdT (SEQ ID NO: 54) [675-693]

siRNA	Target region		Base sequence (corresponding nucleotides of SEQ ID NO: 1)	Experimental sequence (corresponding nucleotides of SEQ ID NO: 1)
siHsp47-7	human hsp47	sense	5'CCGUGGGUGUCAUGAUGAUUU (SEQ ID NO: 55) [921-939]	5' CCGUGGGUGUCAUGAUGAUdTdT (SEQ ID NO: 57) [921-939]
		antisense	5' AUCAUCAUGACACCCACGGUU (SEQ ID NO: 56) [921-939]	5' AUCAUCAUGACACCCACGGdTdT (SEQ ID NO: 58) [921-939]

EXAMPLE 13

[0439] In order to screen for the potent of various siRNA molecules against both the human and rat hsp47 genes, various reporter cell lines were established by lenti-viral induction of human HSP47 cDNA-green fluorescent protein (GFP) or rat GP46 cDNA-GFP construct into 293, HT1080, human HSC line hTERT, or NRK cell lines. These cell lines were further evaluated by siRNA against GFP. The remaining fluorescence signal was measured and normalized to scrambled siRNA (Ambion) and subsequently normalized to cell viability. The results showed that siRNA against GFP knocks down the fluorescence to different extent in different cell lines (FIG. 25). 293_HSP47-GFP and 293_GP46-GFP cell lines were selected for siHsp47 screening due to their ease of transfection and sensitivity to fluorescence knockdown.

[0440] Cells were transfected with 1.5 pmol per well of siRNA against GFP in 96-well tissue culture plates using Lipofectamine RNAiMAX (Invitrogen) in a reverse transfection manner. Cells were seeded at 6,000 cells per well and mixed with the siRNA complexes. Fluorescence readings were taken after 72 hours incubation on a Synergy 2 Multi-Mode Microplate Reader (BioTek).

[0441] Cells treated with or without siRNA were measured for viability after 72 hours incubation using CellTiter-Glo Luminescent Cell Viability Assay Kit according to the manual (Promega). The readings were normalized to samples treated with scrambled siRNA molecules.

EXAMPLE 14 Evaluation of inhibitory efficiency of siHsp47 on hsp47 expression in reporter cell lines

[0442] siRNAs against hsp47 were evaluated for their inhibitory efficiency in 293_HSP47-GFP and 293_GP46-GFP cell lines by evaluating the change in fluorescent signal from the reporter GFP. The experiments were carried out as described in Example 2. The fluorescent signals were normalized to fluorescent signals from cells treated with scrambled siRNA (Ambion) which served as a control. The results indicate that the tested hsp47 siRNA molecules were effective in inhibiting hsp47 mRNA in both cell lines. However, siRNA against GP46 mRNA (as published in the 2008 Sato et al paper) was effective only in the 293_GP46-GFP cell line. The results are shown in FIGS. 26A and 26B.

[0443] The 293_HSP47-GFP and 293_GP46-GFP cell lines treated with siRNA against hsp47 and gp46 were evaluated for viability using the methods described in Example 2. The cell viability was normalized to cells treated with scrambled siRNA (Ambion). The results indicate that the cell viability was not affected significantly by the treatment with siRNA molecules. However, the cell viability of 293_HSP47-GFP cell lines treated with different hsp47 siRNA molecules varied depending on the siRNA molecules used, while the viability of 293_GP46-GFP cell lines were similar. Viability for 293_HSP47-GFP cells was lower for siHsp47-6 and Hsp47-7 treated cells than the rest. The results are shown in FIGS. 26C and 26D.

EXAMPLE 15 Evaluation of siHsp47 inhibitory effect on hsp47 mRNA by TaqMan® qPCR

[0444] In Example 14, the knock down efficiency of siHsp47s in reporter cell lines was evaluated by change in fluorescent signal. To validate the results at the mRNA level, siRNAs targeting endogenous hsp47 were transfected into cells of the human HSC cell line hTERT using Lipofectamine RNAiMAX (Invitrogen) in a reverse transfection manner as described in Example 13

[0445] The hsp47 mRNA level was evaluated for knock down efficiency of the various tested siHsp47 siRNA molecules. Briefly, mRNA were isolated from hTERT cells after 72 hours after transfection using an RNeasy mini kit (Qiagen). The level of hsp47 mRNA was determined by reverse transcription coupled with quantitative PCR using TaqMan® probes. Briefly, cDNA synthesis was carried out using High-Capacity cDNA Reverse Transcription Kit (ABI) according to the manufacturer's instruction, and subjected to TaqMan Gene Expression Assay (ABI, hsp47). The level of hsp47 mRNA was normalized to the level of GAPDH mRNA according to the manufacturer's instruction (ABI). The results indicate that siHsp47-C was the most effective among all the hsp47 siRNAs, siHsp47-2 and siHsp47-2d were the next most effective. The combinations of siHsp47-1 with siHsp47-2 or siHsp47-1 with siHsp47-2d were more effective than siHsp47-1 alone. The results are shown in FIG. 27.

EXAMPLE 16 Validation of siHsp47 knock down effect at the protein level

[0446] The inhibitory effect of different Hsp47 siRNA molecules (siHsp47) on hsp47 mRNA expression were validated at the protein level by measuring the HSP47 in hTERT cells transfected with different siHsp47. Transfection of hTERT cells with different siHsp47 were performed as described in Example 13. Transfected hTERT cells were lysed and the cell lysate were clarified by centrifugation. Proteins in the clarified cell lysate were resolved by SDS polyacrylamide gel electrophoresis. The level of HSP47 protein in the cell lysate were determined using an anti-HSP47 antibody (Assay Designs) as the primary antibody, Goat anti-mouse IgG conjugated with HRP (Millipore) as the secondary antibody, and subsequently detected by Supersignal West Pico Chemiluminescence kit (Pierce). Anti-actin antibody (Abcam) was used as a protein loading control. The result showed significant decrease in the level of Hsp47 protein in cells treated with siHsp47-C, siHsp47-2d, alone or combination of siHsp47-1 with siHsp47-2d.

EXAMPLE 17 Downregulation of Collagen I expression by hsp47 siRNA

[0447] To determine the effect of siHsp47 on collagen I expression level, collagen I mRNA level in hTERT cells treated with different siRNA against hsp47 was measured. Briefly, hTERT cells were transfected with different siHsp47 as described in Example 13. The cells were lysed after 72 hours and mRNA were isolated using RNeasy mini kit according to the manual (Qiagen). The level of collagen 1 mRNA was determined by reverse transcription coupled with quantitative PCR using TaqMan® probes. Briefly, cDNA synthesis was carried out using High-Capacity cDNA Reverse Transcription Kit (ABI) according to the manual, and subjected to TaqMan Gene Expression Assay (ABI, COL1A1 assay). The level of collagen I mRNA was normalized to the level of GAPDH mRNA according to the manufacturer's instruction (ABI). The signals were normalized to the signal obtained from cells transfected with scrambled siRNA. The result indicated that collagen 1 mRNA level is significantly reduced in the cells treated with some of the candidates siHsp47-2, siHsp47-2d, and their combination with siHsp47-1 and shown in FIG. 28.

EXAMPLE 18 Immunofluorescence staining of hsp47 siRNA treated hTERT cells

[0448] To visualize the expression of two fibrosis markers, collagen I and alpha-smooth muscle actin (SMA), in hTERT cells transfected with or without siHsp47, the cells were stained with rabbit anti-collagen I antibody (Abcam) and mouse anti-alpha-SMA antibody (Sigma). Alexa Fluor 594 goat anti-mouse IgG and Alexa Fluor 488 goat anti-rabbit IgG (Invitrogen (Molecular Probes)) were used as secondary antibodies to visualize collagen I (green) and alpha-SMA (red). Hoechst was used to visualize nucleus (blue). The results indicate correlation between siRNA knocking down of some of the target genes and collagen/SMA expression.

EXAMPLE 19 *In vivo* testing of siHSP47 in animal models of liver fibrosis

[0449] The siRNA duplex sequence for HSP47 (siHSP47C) is as listed below.

Sense (5'->3')	ggacagggccucuaacaacuaTT
Antisense (5'->3')	uaguuguagaggccuguccTT

[0450] 10 mg/ml siRNA stock solution was prepared by dissolving in nuclease free water (Ambion). For treatment of cirrhotic rats, siRNA was formulated with vitamin A-coupled liposome as described by Sato et al (Sato Y. et al. 2008 Nature Biotech. 26:431) in order to target activated HSC that produce collagen. The vitamin A (VA)-liposome-siRNA formulation consists of 0.33 µmol/ml of VA, 0.33 µmol/ml of liposome (Coatsome EL, NOF Corporation) and 0.5 µg/µl of siRNA in 5% glucose solution.

[0451] Four week-old male SD rats were induced with liver cirrhosis with 0.5% dimethylnitrosamine (DMN) (Wako Chemicals, Japan) in phosphate-buffered saline (PBS). A dose of 2 ml/kg per body weight was administered intraperitoneally for 3 consecutive days per week on days 0, 2, 4, 7, 9, 11, 14, 16, 18, 21, 23, 25, 28, 30, 32, 34, 36, 38 and 40

[0452] siRNA treatment: siRNA treatment was carried out from day 32 and for 5 intravenous injections. In detail, rats were treated with siRNA on day-32, 34, 36, 38 and 40. Then rats were sacrificed on day-42 or 43. 3 different siRNA doses (1.5 mg siRNA per kg body weight, 2.25 mg siRNA per kg body weight, 3.0 mg siRNA per kg body weight) were tested.

[0453] Detail of tested groups and number of animals in each group are as follows:

1. 1) Cirrhosis was induced by DMN injection, then 5% glucose was injected instead of siRNA) (n=10)
2. 2) VA-Lip-siHSP47C 1.5 mg/kg (n=10)
3. 3) VA-Lip-siHSP47C 2.25 mg/kg (n=10)
4. 4) VA-Lip-siHSP47C 3.0 mg/Kg (n=10)
5. 5) Sham (PBS was injected instead of DMN; 5%Glucose was injected instead of siRNA) (n=6)
6. 6) No treatment control (Intact) (n=6)

VA-Lip refers to vitamin A - liposome complex.

[0454] Evaluation of therapeutic efficacy: On day 43, 2 out of 10 animals in the "diseased rat" group and 1 out of 10 animals in "VA-Lip-siHSP47C siRNA 1.5 mg/kg" died due to development of liver cirrhosis. The remainder of the animals survived. After rats were sacrificed, liver tissues were fixed in 10% formalin. Then, the left lobule of each liver was embedded in paraffin for histology. Tissue slides were stained with Sirius red, and hematoxylin and eosin (HE). Sirius red staining was employed to visualize collagen-deposits and to determine the level of cirrhosis. HE staining was for nuclei and cytoplasm as counter-staining. Each slide was observed under microscope (BZ-8000, Keyence Corp. Japan) and percentage of Sirius red-stained area per slide was determined by image analysis software attached to the microscope. At least 4 slides per each liver were prepared for image analysis, and whole area of each slide (slice of liver) was captured by camera and analyzed. Statistical analysis was carried out by Student's t-test.

[0455] Results: Figure 29 shows the fibrotic areas. The area of fibrosis in "diseased rats" was higher than in the "sham" or "no treatment control" groups. Therefore, DMN treatment induced collagen deposition in the liver, which was a typical observation of liver fibrosis. However, the area of fibrosis was significantly reduced by the treatment of siRNA targeting HSP47 gene, compared with "disease rat" group (FIG. 29). This result indicates that siRNA as disclosed herein has a therapeutic efficacy in actual disease.

[0456] Additional siRNA compounds are tested in the liver fibrosis animal model, and were shown to reduce the fibrotic area in the liver.

EXAMPLE 20 Generation of sequences for active double-stranded RNA compounds to HSP47/SERPINH1 and production of the siRNAs shown in Tables 4, 5, B, C, D and E.

[0457] Duplexes are generated by annealing complementary single-stranded oligonucleotides. In a laminar flow hood, a 500 μ M stock solution of single-stranded oligonucleotide is prepared by diluting in WFI (water for injection). Actual ssRNA concentrations are determined by diluting each 500 μ M ssRNA 1:200 using WFI, and measuring the OD using Nano Drop. The procedure is repeated 3 times and the average concentration is calculated. The stock solution was then diluted to a final concentration of 250 μ M. Complementary single-strands were annealed by heating to 85°C and allowing to cool to room temperature over at least 45 minutes. Duplexes were tested for complete annealing by testing 5 μ l on a 20% polyacrylamide gel and staining. Samples were stored at -80°C.

[0458] Tables 4, 5, B, C, D and E provide siRNAs for HSP47/SERPINH1. For each gene there is a separate list of 19-mer siRNA sequences, which are prioritized based on their score in the proprietary algorithm as the best sequences for targeting the human gene expression.

[0459] The following abbreviations are used in the Tables 4, 5, B, C, D and E herein: "other spec or Sp." refers to cross species identity with other animals: D-dog, Rt-rat, Rb-Rabbit, Rh-rhesus monkey, P- Pig, M-Mouse; ORF: open reading frame. 19-mers (Tables 5, B, C), and 18+1-(Tables 4, D, E) mers refer to oligomers of 19 and 18+1 (U in position 1 of Antisense, A in position 19 of sense strand) ribonucleic acids in length, respectively.

[0460] siRNA oligonucleotides useful in generating double-stranded RNA molecules are disclosed in Tables 4, 5, B, C, D and E below.

TABLE 4

Name	SEQ ID NO SEN	Sense (5' > 3')	SEQ ID NO AS	Antisense (5' > 3')	Cross Species	Ident Human gi"32454740
SERPINH1_2	60	GAGACACAUGGGUGCUAUA	127	UAUAGCACCCAUGUGUCUC	H, Rt, Rh, M, D	[1533-1551] (18/19)
SERPINH1_3	61	GGGAAGAUGCAGAAGAAGA	128	UCUUCUUCUGCAUCUCCCC	H, Rt, Rh, Rb	[1112-1130] (18/19)
SERPINH1_5	62	GAAGAAGGCUGUUGCCAU	129	UAUGGCAACAGCCUUCUUC	H, Rt	[1123-1141] (18/19)
SERPINH1_6	63	ACAAGAUGCAGACGAGUA	130	UACUCGUCUCGCAUCUUGU	H, Rt, Rh,	[1464-1482] (18/19)
SERPINH1_7	64	GGACAACCGUGGCUUCAUA	131	UAUGAAGCCACGGUUGUCC	H, Rh, M	[886-904] (18/19)
SERPINH1_8	65	UGCAGUCCAUAACGAGUA	132	UACUCGUUGAUGGACUGCA	H, Rt, Rh, M	[738-756] (18/19)
SERPINH1_9	66	GCCUCAUCAUCCUCAUGCA	133	UGCAUGAGGAUGAUGAGGC	H, Rt, Rh,	[1026-1044] (18/19)
SERPINH1_10	67	CGCGCUGCAGUCCAUA	134	UUUGAUGGACUGCAGCGCG	H, Rt, Rh	[733-751] (18/19)
SERPINH1_11	68	CGGACAGGCCUCUACAACA	135	UGUUGUAGAGGCCUGUCCG	H, Rt, Rh, P	[944-962] (18/19)
SERPINH1_13	69	UGACAAGAUGCAGACGAA	136	UUCGUCUCGCAUCUUGUCA	H, Rh	[1462-1480] (18/19)
SERPINH1_14	70	CCAGCCUCAUCAUCCUCAA	137	UUGAGGAUGAUGAGGCUGG	H, M, Rt, Rh, D-	[1023-1041] (18/19)
SERPINH1_15	71	GCUGCAGUCCAUAACGAA	138	UUCGUUGAUGGACUGCAGC	H, Rt, Rh	[736-754] (18/19)
SERPINH1_16	72	GCAGCGCGCUGCAGUCCAA	139	UUGGACUGCAGCGCGCUGC	H, Rt, Rh	[729-747] (18/19)
SERPINH1_17	73	UGAGACACAUGGGUGCUAA	140	UUAGCACCCAUGUGUCUCA	H, Rt, Rh, M, D	[1532-1550] (18/19)
SERPINH1_19	74	GGUGGAGGUGACCCAUGAA	141	UUCAUGGGUCACCUCCACC	H, Rt, Rh, M	[1159-1177] (18/19)
SERPINH1_20	75	CUUUGACCAGGACAUCUAA	142	UUAGAUGUCCUGGUCAAAG	H, Rt, Rh	[1324-1342] (18/19)
SERPINH1_21	76	GGAGGUGACCCAUGACCUA	143	UAGGUCAUGGGUCACCUCC	H, Rt, Rh, D	[1162-1180] (18/19)
SERPINH1_22	77	CUCCUGAGACACAUGGGUA	144	UACCCAUGUGUCUCAGGAG	H, D	[1528-1546] (18/19)
SERPINH1_23	78	AGAAGAAGGCUGUUGCCAA	145	UUGGCAACAGCCUUCUUCU	H, Rt	[1122-1140] (18/19)
SERPINH1_24	79	AGCUCUCCAGCCUCAUCAA	146	UUGAUGAGGCUGGAGAGCU	H, Rt, D, M, P, Rh	[1017-1035] (18/19)

Name	SEQ ID NO SEN	Sense (5' > 3')	SEQ ID NO AS	Antisense (5' > 3')	Cross Species	Ident Human gi"32454740
SERPINH1_25	80	CUGCAGUCCAUCAACGAGA	147	UCUCGUUGAUGGACUGCAG	H, Rt, Rh M	[737-755] (18/19)
SERPINH1_26	81	CCGGACAGGCCUCUACAAA	148	UUUGUAGAGGCCUGUCCGG	H, Rt, Rh, Rb, P	[943-961] (18/19)
SERPINH1_27	82	GCACCGGACAGGCCUCUAA	149	UUAGAGGCCUGUCCGGUGC	H, Rt, Rh, Rb, P	[940-958] (18/19)
SERPINH1_28	83	GCAGAAGAAGGCUGUUGCA	150	UGCAACAGCCUUCUUCUGC	H, Rt	[1120-1138] (18/19)
SERPINH1_31	84	AGAAGGCUGUUGCCAUCUA	151	UAGAUGGCAACAGCCUUCU	H, Rt	[1125-1143] (18/19)
SERPINH1_32	85	AGCGCAGCGCGCUGCAGUA	152	UACUGCAGCGCGCUGCGCU	H, Rt, Rh,	[726-744] (18/19)
SERPINH1_33	86	GACACAUGGGUGCUAUUGA	153	UCAAUAGCACCCAUGUGUC	H, Rt, Rh, M	[1535-1553] (18/19)
SERPINH1_34	87	GGGCCUGACUGAGGCCAUA	154	UAUGGCCUCAGUCAGGCC	H, Rt	[1201-1219] (18/19)
SERPINH1_35	88	AGACACAUGGGUGCUAUUA	155	UAAUAGCACCCAUGUGUCU	H, Rt, Rh, M	[1534-1552] (18/19)
SERPINH1_36	89	CCAUGACCUGCAGAAACAA	156	UUGUUUCUGCAGGUCAUGG	H, Rt, Rh, M	[1171-1189] (18/19)
SERPINH1_37	90	AGAUGCAGAAGAAGGCUGA	157	UCAGCCUUCUUCUGCAUCU	H, Rt, Rh, M	[1116-1134] (18/19)
SERPINH1_38	91	CAAGCUCUCCAGCCUCAUA	158	UAUGAGGCUGGAGAGCUUG	H, Rt, Rh, P, D	[1015-1033] (18/19)
SERPINH1_39	92	UGCAGAAGAAGGCUGUUGA	159	UCAACAGCCUUCUUCUGCA	H, Rt	[1119-1137] (18/19)
SERPINH1_41	93	CAGCCUCAUCAUCCUCAUA	160	UAUGAGGAUGAUGAGGCUG	H, Rt, Rh, M, D	[1024-1042] (18/19)
SERPINH1_42	94	GACAGGCCUCUACAACUAA	161	UUAGUUGUAGAGGCCUGUC	H, Rt, Rh, Rb, P	[946-964] (18/19)
SERPINH1_43	95	GAUGCAGAAGAAGGCUGUA	162	UACAGCCUUCUUCUGCAUC	H, Rt, Rh, M	[1117-1135] (18/19)
SERPINH1_44	96	ACCCAUGACCUGCAGAAAA	163	UUUUCUGCAGGUCAUGGGU	H, Rt, Rh, M	[1169-1187] (18/19)
SERPINH1_45	97	ACUCCAAGAUCAACUCCA	164	UGGAAGUUGAUCUUGGAGU	H, Rt, Rh, M, D	[702-720] (18/19)
SERPINH1_45 a	98	ACUCCAAGAUCAACUCCU	165	AGGAAGUUGAUCUUGGAGU	H, Rt, Rh, M, D	[702-720] (18/19)
SERPINH1_48	99	AGGCCUCUACAACUACUAA	166	UUAGUAGUUGUAGAGGCCU	H, Rt, Rh, Rb, P, D	[949-967] (18/19)
SERPINH1_49	100	CACUCCAAGAUCAACUUCA	167	UGAAGUUGAUCUUGGAGUG	H, Rt, Rh, M, D	[701-719] (18/19)
SERPINH1_51	101	UCCUGAGACACAUGGGUGA	168	UCACCCAUGUGUCUCAGGA	H, Rt, D, M	[1529-1547] (18/19)
SERPINH1_52	102	GACAAGAUGCGAGACGAGA	169	UCUCGUCUCGCAUCUUGUC	H, Rt, Rh,	[1463-1481] (18/19)
SERPINH1_53	103	GGUGACCCAUGACCUGCAA	170	UUGCAGGUCAUGGGUCACC	H, Rt, Rh, M	[1165-1183] (18/19)
SERPINH1_59	104	CCGAGGUGAAGAAACCUGA	171	UCAGGUUUCUUCACCUCGG	H, Rt, Rh,	[285-303] (18/19)
SERPINH1_51 a	105	UCCUGAGACACAUGGGUGU	172	ACACCCAUGUGUCUCAGGA	H, Rt, D, M	[1529-1547] (18/19)
SERPINH1_61	106	GCACUCCAAGAUCAACUUA	173	UAAGUUGAUCUUGGAGUGC	H, Rh, D	[700-718] (18/19)
SERPINH1_62	107	GUGGUGGAGGUGACCCAUA	174	UAUGGGUACCUCCACCAC	H, Rt, Rh, M, Rb	[1157-1175] (18/19)
SERPINH1_64	108	GCCGAGGUGAAGAAACCUA	175	UAGGUUUCUUCACCUCGGC	H, Rt, Rh,	[284-302] (18/19)
SERPINH1_65	109	GCUCUCCAGCCUCAUCAUA	176	UAUGAUGAGGCUGGAGAGC	H, Rt, D, M, P, Rh	[1018-1036] (18/19)
SERPINH1_66	110	GAUGCACCGGACAGGCCUA	177	UAGGCCUGUCCGGUGCAUC	H, Rt, Rh, M, Rb, P	[937-955] (18/19)

Name	SEQ ID NO SEN	Sense (5' > 3')	SEQ ID NO AS	Antisense (5' > 3')	Cross Species	Ident Human gi"32454740
SERPINH1_68	111	CUCUCCAGCCUCAUCAUCA	178	UGAUGAUGAGGCUGGAGAG	H, Rt, D, M, P, Rh	[1019-1037] (18/19)
SERPINH1_69	112	GCAGACCACCGACGGCAAA	179	UUUGCCGUCGGUGGUCUGC	H, Rt, D	[763-781] (18/19)
SERPINH1_70	113	AGUCCAUCAACGAGUGGGA	180	UCCCACUCGUUGAUGGACU	H, Rt, Rh, M	[741-759] (18/19)
SERPINH1_71	114	ACCGUGGCUUCAUGGUGAA	181	UUCACCAUGAAGCCACGGU	H, Rt, Rh, M	[891-909] (18/19)
SERPINH1_74	115	GAAGGCUGUUGCCAUCUCA	182	UGAGAUGGCAACAGCCUUC	H, Rt,	[1126-1144] (18/19)
SERPINH1_75	116	GAAGAUGCAGAAGAAGGCA	183	UGCCUUCUUCUGCAUCUUC	H, Rt, Rh, Rb	[1114-1132] (18/19)
SERPINH1_77	117	UGAUGAUGCACCGGACAGA	184	UCUGUCCGGUGCAUCAUCA	H, Rh,	[933-951] (18/19)
SERPINH1_78	118	CCCUUUGACCAGGACAUCA	185	UGAUGUCCUGGUCAAAGGG	H, Rt, Rh,	[1322-1340] (18/19)
SERPINH1_80	119	CAGUCCAUCAACGAGUGGA	186	UCCACUCGUUGAUGGACUG	H, Rt, Rh, M	[740-758] (18/19)
SERPINH1_82	120	CAACCGUGGCUUCAUGGUA	187	UACCAUGAAGCCACGGUUG	M Rt, Rh,	[889-907] (18/19)
SERPINH1_83	121	CGACAAGCGCAGCGCGCUA	188	UAGCGCGCUGCGCUUGUCG	H	[721-739] (18/19)
SERPINH1_84	122	GCAGUCCAUCAACGAGUGA	189	UCACUCGUUGAUGGACUGC	H, Rt, Rh, M	[739-757] (18/19)
SERPINH1_86	123	ACAGGCCUCUACAACUACA	190	UGUAGUUGUAGAGGCCUGU	H, Rt, Rb, P, D	[947-965] (18/19)
SERPINH1_87	124	AAGAUGCAGAAGAAGGCUA	191	UAGCCUUCUUCUGCAUCUU	H, Rt, Rh, M	[1115-1133] (18/19)
SERPINH1_89	125	CAGCGCGCUGCAGUCCAUA	192	UAUGGACUGCAGCGCGCUG	H, Rt, Rh,	[730-748] (18/19)
SERPINH1_90	126	GCGCAGCGCGCUGCAGUCA	193	UGACUGCAGCGCGCUGCGC	H, Rt, Rh,	[727-745] (18/19)

TABLE 5:

Name	SEQ ID NO SEN	Sense (5' > 3')	SEQ ID NO AS	Antisense (5' > 3')	Species	Lg	Ident Human gi 32454740
SERPINH1_1	194	GGACAGGCCUCUACAACUA	219	UAGUUGUAGAGGCCUGUCC	H, Rt, Rh, Rb, P	19	[945-963] (19/19)
SERPINH1_4	195	GAGACACAUGGGUGCUAUU	220	AAUAGCACCCAUGUGUCUC	H, Rt, Rh, M, D	19	[1533-1551] (19/19)
SERPINH1_12	196	ACAAGAUGCGAGACGAGUU	221	AACUCGUCUCGCAUCUUGU	H, Rt, Rh,	19	[1464-1482] (19/19)
SERPINH1_18	197	CCUUUGACCAGGACAUCUA	222	UAGAUGUCCUGGUCAAAGG	H, Rt, Rh,	19	[1323-1341] (19/19)
SERPINH1_29	198	GACCCAUGACCUGCAGAAA	223	UUUCUGCAGGUCAUGGGUC	H, Rt, Rh, M	19	[1168-1186] (19/19)
SERPINH1_30	199	CGGACAGGCCUCUACAACU	224	AGUUGUAGAGGCCUGUCCG	H, Rt, Rh, Rb, P	19	[944-962] (19/19)
SERPINH1_40	200	ACCGGACAGGCCUCUACAA	225	UUGUAGAGGCCUGUCCGGU	H, Rt, Rh, Rb, P,	19	[942-960] (19/19)
SERPINH1_46	201	GCAGCGCGCUGCAGUCCA	226	AUGGACUGCAGCGCGCUGC	H, Rt, Rh,	19	[729-747] (19/19)
SERPINH1_47	202	GCGCGCUGCAGUCCAUCAA	227	UUGAUGGACUGCAGCGCGC	H, Rt, Rh,	19	[732-750] (19/19)
SERPINH1_50	203	CUGAGACACAUGGGUGCUA	228	UAGCACCCAUGUGUCUCAG	H, Rt, Rh, M, D	19	[1531-1549] (19/19)
SERPINH1_54	204	AGAAGAAGGCUGUUGCCA	229	AUGGCAACAGCCUUCUUCU	H, Rt	19	[1122-1140] (19/19)
SERPINH1_55	205	AGCUCUCCAGCCUCAUCAU	230	AUGAUGAGGCUGGAGAGCU	H, Rt, D, P, Rh	19	[1017-1035] (19/19)

Name	SEQ ID NO SEN	Sense (5' > 3')	SEQ ID NO AS	Antisense (5' > 3')	Species	Lg	Ident Human gi 32454740
SERPINH1_56	206	CUGCAGUCCAUAACGAGU	231	ACUCGUUGAUGGACUGCAG	H, Rt, Rh, M	19	[737-755] (19/19)
SERPINH1_57	207	CGCUGCAGUCCAUAACGA	232	UCGUUGAUGGACUGCAGCG	H, Rt, Rh,	19	[735-753] (19/19)
SERPINH1_58	208	GACAAGAUGCGAGACGAGU	233	ACUCGUCUCGCAUCUUGUC	H, Rt, Rh,	19	[1463-1481] (19/19)
SERPINH1_63	209	GGGCCUGACUGAGGCCAUU	234	AAUGGCCUCAGUCAGGCC	H, Rt	19	[1201-1219] (19/19)
SERPINH1_67	210	GAUGCAGAAGAAGGCUGUU	235	AACAGCCUUCUUCUGCAUC	H, Rt, Rh, M	19	[1117-1135] (19/19)
SERPINH1_72	211	CACCGGACAGGCCUCUACA	236	UGUAGAGGCCUGUCCGGUG	H, Rt, Rh, Rb, P	19	[941-959] (19/19)
SERPINH1_73	212	AGAUGCAGAAGAAGGCUGU	237	ACAGCCUUCUUCUGCAUCU	H, Rt, Rh, M	19	[1116-1134] (19/19)
SERPINH1_76	213	AGCGCGCUGCAGUCCAUA	238	UGAUGGACUGCAGCGCGCU	H, Rt, Rh	19	[731-749] (19/19)
SERPINH1_79	214	GGAAGAUGCAGAAGAAGGC	239	GCCUUCUUCUGCAUCUUC	H, Rt, Rh, Rb	19	[1113-1131] (19/19)
SERPINH1_81	215	GAAGAAGGCUGUUGCCAUC	240	GAUGGCAACAGCCUUCUUC	H, Rt	19	[1123-1141] (19/19)
SERPINH1_83	216	UGCAGUCCAUAACGAGUG	241	CACUCGUUGAUGGACUGCA	H, Rt, Rh, M	19	[738-756] (19/19)
SERPINH1_88	217	CCUGAGACACAUGGGUGCU	242	AGCACCCAUGUGUCUCAGG	H, Rt, D,	19	[1530-1548] (19/19)
SERPINH1_91	218	CGCAGCGCGCUGCAGUCCA	243	UGGACUGCAGCGCGCUGCG	H, Rt, Rh,	19	[728-746] (19/19)

TABLE B: Additional Active 19-mer SERPINH1 siRNAs

No	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
1	244	GGCAGACUCUGGUCAAGAA	460	UUCUUGACCAGAGUCUGCC	Rh	[2009-2027] 3'UTR
2	245	CAGUGAGGCGGAUUGAGAA	461	UUCUCAUCCGCCUCACUG		[1967-1985] 3'UTR
3	246	AGCCUUUGUUGCUAUCAAU	462	AUUGAUAGCAACAAAGGCU	Rh	[2117-2135] 3'UTR
4	247	CCAUGUUCUUAAGCCACA	463	UGUGGCUUGAAGAACAUGG	Rh,Rb,D	[837-855] ORF
5	248	CCCUCUUCUGACACUAAAA	464	UUUUAGUGUCAGAAGAGGG		[1850-1868] 3'UTR
6	249	CCUCAAUCAUAUUAUUAU	465	AUAUGAAUACUGAUUGAGG		[1774-1792] 3'UTR
7	250	GAGACACAUGGGUGCUAUU	466	AAUAGCACCCAUGUGUCUC	Rh,D,Rt,M	[1533-1551] 3'UTR
8	251	GUGACAAGAUGCGAGACGA	467	UCGUCUCGCAUCUUGUCAC	Rh	[1461-1479] ORF
9	252	GCCACACUGGGAUGAGAAA	468	UUUCUCAUCCAGUGUGGC	Rh,Rb,M	[850-868] ORF
10	253	AGAUGCGAGACGAGUUUAU	469	UAUAACUCGUCUCGCAUCU	Rh	[1467-1485] ORF
11	254	ACGACGACGAGAAGGAAAA	470	UUUUCCUUCUCGUCGUCGU		[966-984] ORF
12	255	GCCUCUACAACUACUACGA	471	UCGUAGUAGUUGUAGAGGC	Rb,D	[951-969] ORF
13	256	AGAUAACUCCGCGACAA	472	UUGUCGCGGAAGUUGAUCU	D	[708-726] ORF
14	257	ACUACUACGACGACGAGAA	473	UUCUCGUCGUCGUAGUAGU	Rb	[960-978] ORF
15	258	AGCCCUCUUCUGACACUAA	474	UUAGUGUCAGAAGAGGGCU		[1848-1866] 3'UTR
16	259	ACAAGAUGCGAGACGAGUU	475	AACUCGUCUCGCAUCUUGU	Rh,Rt	[1464-1482] ORF
17	260	AGCCACACUGGGAUGAGAA	476	UUCUCAUCCAGUGUGGCU	Rh,Rb,M	[849-867] ORF
18	261	AGGACCAGGCAGUGGAGAA	477	UUCUCCACUGCCUGGUCCU	Rh	[408-426] ORF
19	262	CAGGCAAGAAGGACCUGUA	478	UACAGGUCCUUCUUGCCUG	Rh,D	[1251-1269] ORF
20	263	ACCUGUGAGACCAAUUGA	479	UCAUUUUGGUCUCACAGGU	Rh	[1813-1831] 3'UTR
21	264	CUUUGUUGCUAUCAAUCCA	480	UGGAUUGAUAGCAACAAAG	Rh	[2120-2138] 3'UTR
22	265	GUGAGACCAAUUGAGCUA	481	UAGCUCAAUUUGGUCUCAC	Rh	[1817-1835] 3'UTR
23	266	CCUGAAAGUCCAGAUCA	482	UGAUCUGGGACUUUCAGGG		[1749-1767] 3'UTR
24	267	CCUUUGACCAGGACAUCUA	483	UAGAUGUCCUGGUCAAAGG	Rh,Rt	[1323-1341] ORF
25	268	GACCAGGCAGUGGAGAACA	484	UGUUCUCCACUGCCUGGUC	Rh	[410-428] ORF
26	269	CGCGCAACGUGACCUGGAA	485	UUCCAGGUCACGUUGCGCG	M	[597-615] ORF
27	270	AUGAGAAAUCCACCACAA	486	UUGUGGUGGAUUUCUCAU	Rh	[861-879] ORF

No	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
28	271	GAAGAAACCUGCAGCCGCA	487	UGCGGCUGCAGGUUUUCUUC		[292-310] ORF
29	272	CUCUCGAGCGCCUUGAAAA	488	UUUUCAAGGCGCUCGAGAG		[1059-1077] ORF
30	273	GGAACAUGAGCCUUUGUUG	489	CAACAAAGGCUCAUGUUC	Rh	[2109-2127] 3'UTR
31	274	CUCACCUGUGAGACCAAAU	490	AUUUGGUCUCACAGGUGAG	Rh	[1810-1828] 3'UTR
32	275	CUACGACGACGAGAAGGAA	491	UUCUUCUCGUCGUCGUAG	Rb	[964-982] ORF
33	276	ACCACAAGAUGGUGGACAA	492	UUGUCCACCAUCUUGUGGU	Rh,Rb,M,P	[873-891] ORF
34	277	CUGGCACUGCGGAGAAGUU	493	AACUUCUCCGCAGUGCCAG		[318-336] ORF
35	278	GGUCCUAUACCGUGGGUGU	494	ACACCCACGGUAUAGGACC	Rh	[912-930] ORF
36	279	CCAACCUCUCCCAACUAUA	495	UAUAGUUGGGAGAGGUUGG	Rh	[1896-1914] 3'UTR
37	280	GAGAAGGAAAAGCUGCAAA	496	UUUGCAGCUUUUCCUUCUC	Rh	[974-992] ORF
38	281	GCCUCUCGAGCGCCUUGAA	497	UUCAAGGCGCUCGAGAGGC		[1057-1075] ORF
39	282	AGGCCAUUGACAAGAACAA	498	UUGUUCUUGUCAUUGCCU	Rh,D	[1212-1230] ORF
40	283	3ACCCAUGACCUGCAGAAA	499	UUUCUGCAGGUCAUGGGUC	Rh,Rt,M	[1168-1186] ORF
41	284	CUCCUGGCACUGCGGAGAA	500	UUCUCCGCAGUGCCAGGAG		[315-333] ORF
42	285	CGGACAGGCCUCUACAACU	501	AGUUGUAGAGGCCUGUCCG	Rh,Rb,Rt,P	[944-962] ORF
43	286	3AUGAGAAAUCCACCACA	502	UGUGGUGGAAUUUCUCAUC	Rh	[860-878] ORF
44	287	CACGCAUGUCAGGCAAGAA	503	UUCUUGCCUGACAUGCGUG	Rh,D	[1242-1260] ORF
45	288	ACCUCUCCCAACUAUAAAA	504	UUUUUAAGUUGGGAGAGGU	Rh	[1899-1917] 3'UTR
46	289	ACCAGGCAGUGGAGAACAU	505	AUGUUCUCCACUGCCUGGU	Rh	[411-429] ORF
47	290	GGGAACAUGAGCCUUUGUU	506	AACAAAGGCUCAUGUCCCC	Rh	[2108-2126] 3'UTR
48	291	AGAAUUCACUCCACUUGGA	507	UCCAAGUGGAGUGAAUUCU	Rh	[1653-1671] 3'UTR
49	292	GGGCAGACUCUGGUCAAGA	508	UCUUGACCAGAGUCUGCCC	Rh	[2008-2026] 3'UTR
50	293	AGAAGGAAAAGCUGCAAAU	509	AUUUGCAGCUUUUCCUUCU	Rh	[975-993] ORF
51	294	GGCAGUGGAGAACAUCCUG	510	CAGGAUGUUCUCCACUGCC	Rh	[415-433] ORF
52	295	GGGAUGAGAAAUCCACCA	511	UGGUGGAAUUUCUCAUCCC	Rh	[858-876] ORF
53	296	CCAAGCUGUUCUACGCCGA	512	UCGGCGUAGAACAGCUUGG	Rh	[1365-1383] ORF
54	297	ACCGGACAGGCCUCUACAA	513	UUGUAGAGGCCUGUCCGGU	Rh,Rb,Rt,P	[942-960] ORF
55	298	CUGCCUCAAUACAGUAUUA	514	UGAAUACUGAUUGAGGCAG		[1771-1789] 3'UTR
56	299	CAGCCUCUUCUGACACUA	515	UAGUGUCAGAAGAGGGCUG		[1847-1865] 3'UTR
57	300	CCAGCCUCAUCAUCCUCAU	516	AUGAGGAUGAUAGAGGCUGG	Rh,D,Rt,M	[1023-1041] ORF
58	301	AGGGUGACAAGAUGCGAGA	517	UCUCGCAUCUUGUCACCCU	Rh,D	[1458-1476] ORF
59	302	GGACCAGGCAGUGGAGAAC	518	GUUCUCCACUGCCUGGUCC	Rh	[409-427] ORF
60	303	GCAGCGCGCUGCAGUCCAU	519	AUGGACUGCAGCGCGCUGC	Rh,Rt	[729-747] ORF
51	304	GCGCGCUGCAGUCCAUA	520	UUGAUGGACUGCAGCGCGC	Rh,Rt	[732-750] ORF
62	305	CCAGAUACCAUGAUGCUGA	521	UCAGCAUCAUGGUAUCUGG	Rh	[1680-1698] 3'UTR
63	306	CUAGUGCGGGACACCCAAA	522	UUUGGGUGUCCCGCACUAG		[1400-1418] ORF
64	307	AGGCAGUGGAGAACAUCU	523	AGGAUGUUCUCCACUGCCU	Rh	[414-432] ORF
55	308	CUGAGACACAUGGGUGCUA	524	UAGCACCCAUGUGUCUCAG	Rh,D,Rt,M	[1531-1549] 3'UTR
66	309	GAUUGAGAAGGAGCUCCCA	525	UGGGAGCUCCUUCUCAAUC		[1977-1995] 3'UTR
67	310	CGCAGACCACCGACGGCAA	526	UUGCCGUCGGUGGUCUGCG	D,Rt	[762-780] ORF
58	311	CCACACUGGGAUGAGAAAU	527	AUUUCUCAUCCAGUGUGG	Rh	[851-869] ORF
59	312	GCUCAGUGAGCUUCGCUGA	528	UCAGCGAAGCUCACUGAGC		[642-660] ORF
70	313	CGCCUUUGAGUUGGACACA	529	UGUGUCCAACUCAAGGCG	Rh	[1294-1312] ORF
71	314	GGUCAGCCAGCCCUCUUC	530	GAAGAGGGCUGGCUGACCC	Rh	[1839-1857] 3'UTR
72	315	GGGCUUCUGGGCAGACUCU	531	AGAGUCUGCCCAGAAGCCC	Rh	[2000-2018] 3'UTR
73	316	GGUACCUUCUCACCUGUGA	532	UCACAGGUGAGAAGGUACC	Rh	[1802-1820] 3'UTR
74	317	GCCUGCCUCAAUACAGUAU	533	AAUACUGAUUGAGGCAGGC		[1769-1787] 3'UTR
75	318	UCUACAACUACUACGACGA	534	UCGUCGUAGUAGUUGUAGA	Rb	[954-972] ORF
76	319	GGGAAGAUGCAGAAGAAGG	535	CCUUCUUCUGCAUCUCCCC	Rh,Rb,Rt	[1112-1130] ORF
77	320	CGAGAAGGAAAAGCUGCAA	536	UUGCAGCUUUUCCUUCUCG	Rh	[973-991] ORF
78	321	AGAAGAAGGCUGUUGCCAU	537	AUGGCAACAGCCUUCUUCU	Rt	[1122-1140] ORF
79	322	CACAAGCUCUCCAGCCUCA	538	UGAGGCUGGAGAGCUUGUG	Rh,D,M,P	[1013-1031] ORF
80	323	GGGUGACAAGAUGCGAGAC	539	GUCUCGCAUCUUGUCACCC	Rh,D	[1459-1477] ORF

No	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
81	324	UGUUGGAGCGUGGAAAAAA	540	UUUUUCCACGCUCCAACA		[2190-2208] 3'UTR
82	325	CUUUGAGUUGGACACAGAU	541	AUCUGUGUCCAACUCAAAAG	Rh	[1297-1315] ORF
83	326	AGCUCUCCAGCCUCAUCAU	542	AUGAUGAGGCUGGAGAGCU	Rh,D,Rt,M, P	[1017-1035] ORF
84	327	AGCUGUUCUACGCCGACCA	543	UGGUCGGCGUAGAACAGCU	Rh	[1368-1386] ORF
85	328	CUGCAGUCCAUCAACGAGU	544	ACUCGUUGAUGGACUGCAG	Rh,Rt,M	[737-755] ORF
86	329	UACGACGACGAGAAGGAAA	545	UUUCCUUCUCGUCGUCGUA		[965-983] ORF
87	330	CCUAGUGCGGGACACCCAA	546	UUGGGUGUCCCGCACUAGG		[1399-1417] ORF
88	331	CUUCUCACCUGUGAGACCA	547	UGGUCUCACAGGUGAGAAG	Rh	[1807-1825] 3'UTR
89	332	AGUUGGACACAGAUGGCAA	548	UUGCCAUCUGUGUCCAACU		[1302-1320] ORF
90	333	CAGUGGAGAACAUCCUGGU	549	ACCAGGAUGUUCUCCACUG	Rh	[417-435] ORF
91	334	CCAGCUAGAAUUCACUCCA	550	UGGAGUGAAUUCUAGCUGG	Rh	[1647-1665] 3'UTR
92	335	CGCUGCAGUCCAUCAACGA	551	UCGUUGAUGGACUGCAGCG	Rh,Rt	[735-753] ORF
93	336	CCAAGGACCAGGCAGUGGA	552	UCCACUGCCUGGUCCUUGG	Rh	[405-423] ORF
94	337	AGUUCUCAAAGAUAGGGA	553	UCCCUAUCUUUGAAGAACU		[2082-2100] 3'UTR
95	338	CGGACCUUCCCAGCUAGAA	554	UUCUAGCUGGGAAGGUCCG	Rh	[1638-1656] 3'UTR
96	339	GACAAGAUGCGAGACGAGU	555	ACUCGUCUCGCAUCUUGUC	Rh,Rt	[1463-1481] ORF
97	340	CCAAGAUCAACUCCGCGA	556	UCGCGGAAGUUGAUCUUGG	D	[705-723] ORF
98	341	CCCAUCACGUGGAGCCUCU	557	AGAGGCUCCACGUGAUGGG	Rh	[1044-1062] ORF
99	342	CCAUGAUGCUGAGCCCGGA	558	UCCGGGCUCAGCAUCAUGG		[1687-1705] 3'UTR
100	343	AGCCUGCCUCAAUCAGUUA	559	AUACUGAUUAGGCGAGGCU		[1768-1786] 3'UTR
101	344	CGGCCUAAGGGUGACAAGA	560	UCUUGUCACCCUAGGCCG	Rh	[1451-1469] ORF
102	345	GGGCCUGACUGAGGCCAUU	561	AAUGGCCUCAGUCAGGCC	Rt	[1201-1219] ORF
103	346	UACCUGUGAGACCAAAUU	562	AAUUUGGUCUCACAGGUGA	Rh	[1811-1829] 3'UTR
104	347	GAGGCCAUUGACAAGAACA	563	UGUUCUUGUCAUUGGCCUC	Rh,D	[1211-1229] ORF
105	348	GCUCCUGGCACUGCGGAGA	564	UCUCCGCAGUGCCAGGAGC	ORF	[314-332] ORF
106	349	GGCGCCUGGUCCGGCCUAA	565	UUAGGCCGGACCAGGCGCC	Rh	[1440-1458] ORF
107	350	CCAGCCCUCUUCUGACACU	566	AGUGUCAGAAGAGGGCUGG		[1846-1864] 3'UTR
108	351	ACUACGACGACGAGAAGGA	567	UCCUUCUCGUCGUCGUAGU	Rb	[963-981] ORF
109	352	CCUAUACCGUGGGUGUCAU	568	AUGACACCCACGGUAUAGG	Rh,D,P	[915-933] ORF
110	353	GACCCAGCUCAGUGAGCUU	569	AAGCUCACUGAGCUGGGUC		[636-654] ORF
111	354	UGGGUGUCAUGAUGAUGCA	570	UGCAUCAUCAUGACACCCA	Rh	[924-942] ORF
112	355	CCAAGGGUGUGGUGGAGGU	571	ACCUCCACCACCCUUGG	Rh,D	[1149-1167] ORF
113	356	AGGUCACCAAGGACGUGGA	572	UCCACGUCCUUGGUGACCU	Rh,D	[789-807] ORF
114	357	CCUGGCCGCCGAGGUGAA	573	UUCACCUCGGCGGCCAGGG		[276-294] ORF
115	358	AGCACUCCAAGAUCAACUU	574	AAGUUGAUCUUGGAGUGCU	Rh,D	[699-717] ORF
116	359	CCUGGCACUGCGGAGAAGU	575	ACUUCUCCGCAGUGCCAGG		[317-335] ORF
117	360	GAUGCAGAAGAAGGCUGUU	576	AACAGCCUUCUUCUGCAUC	Rh,Rt,M	[1117-1135] ORF
118	361	CCCACAAGCUCUCCAGCCU	577	AGGCUGGAGAGCUUGUGGG	Rh,D,P	[1011-1029] ORF
119	362	CUCUUCUGACACUAAAACA	578	UGUUUUAGUGUCAGAAGAG		[1852-1870] 3'UTR
120	363	ACGAGAAGGAAAAGCUGCA	579	UGCAGCUUUUCCUUCUCGU	Rh	[972-990] ORF
121	364	UGAAAAGCUGCUAACCAAA	580	UUUGGUUAGCAGCUUUUCA		[1072-1090] ORF
122	365	UCUCACCUGUGAGACCAAA	581	UUUGGUCUCACAGGUGAGA	Rh	[1809-1827] 3'UTR
123	366	CAUGAUGAUGCACCGGACA	582	UGUCCGGUGCAUCAUCAUG	Rh	[931-949] ORF
124	367	GGAUUGAGAAGGAGCUCCC	583	GGGAGCUCCUUCUCAAUCC		[1976-1994] 3'UTR
125	368	CCUUAUCUUCUAGUGCG	584	CGCACUAGGAAGAUGAAGG		[1389-1407] ORF
126	369	GGCCUGGCCUUCAGCUUGU	585	ACAAGCUGAAGGCCAGGCC		[374-392] ORF
127	370	GGUCAGCCAGCCUCUUCU	586	AGAAGAGGGCUGGCUGACC	Rh	[1840-1858] 3'UTR
128	371	UUCUCACCUGUGAGACCAA	587	UUGGUCUCACAGGUGAGAA	Rh	[1808-1826] 3'UTR
129	372	CGCAGCAGCUCCUGGCACU	588	AGUGCCAGGAGCUGCUGCG		[307-325] ORF
130	373	GCCAUGUUCUUAAGCCAC	589	GUGGCUUGAAGAACAUGGC	Rh,Rb,D	[836-854] ORF
131	374	AGGCAGUGCUGAGCGCCGA	590	UCGGCGCUCAGCACUGCCU		[510-528] ORF
132	375	CACCUGUGAGACCAAAUUG	591	CAAUUUGGUCUCACAGGUG	Rh	[1812-1830] 3'UTR
133	376	CACCGGACAGGCCUCUACA	592	UGUAGAGGCCUGUCCGGUG	Rh,Rb,Rt,P	[941-959] ORF

No	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
134	377	AGCUAGAAUUCACUCCACU	593	AGUGGAGUGAAUUCUAGCU	Rh	[1649-1667] 3'UTR
135	378	AGAUGCAGAAGAAGGCUGU	594	ACAGCCUUCUUCUGCAUCU	Rh,Rt,M	[1116-1134] ORF
136	379	CCCUGCUAGUCAACGCCAU	595	AUGGCGUUGACUAGCAGGG	Rh	[822-840] ORF
137	380	ACAACUACUACGACGACGA	596	UCGUCGUCGUAGUAGUUGU	Rb	[957-975] ORF
138	381	GCUCCUGAGACACAUGGGU	597	ACCCAUGUGUCUCAGGAGC	D	[1527-1545] 3'UTR
139	382	UGGAGAAUUCUGGUGUC	598	GACACCAGGAUGUUCUCCA	Rh	[420-438] ORF
140	383	AGCGCGCUGCAGUCCAUC	599	UGAUGGACUGCAGCGCGCU	Rh,Rt	[731-749] ORF
141	384	CGCCUUGAAAAGCUGCUAA	600	UUAGCAGCUUUUCAAAGGCG		[1067-1085] ORF
142	385	GCCUUUGUUGCUAUAUAAUC	601	GAUUGAUAGCAACAAAGGC	Rh	[2118-2136] 3'UTR
143	386	CUCUACAACUACUACGACG	602	CGUCGUAGUAGUUGUAGAG	Rb	[953-971] ORF
144	387	CGCUCACUCAGCAACUCCA	603	UGGAGUUGCUGAGUGAGCG	Rh	[575-593] ORF
145	388	GGUACCAGCCUUGGAUACU	604	AGUAUCCAAGGCUGGUACC	Rh	[1571-1589] 3'UTR
146	389	GCCUGACUGAGGCCAUUGA	505	UCAAUGGCCUCAGUCAGGC	Rh	[1203-1221] ORF
147	390	UGAGCUUUCGUGAUGACUU	606	AAGUCAUCAGCGAAGCUCA	Rh	[648-666] ORF
148	391	CCAGCCUUGGAUACUCCA	607	AUGGAGUAUCCAAGGCUGG	Rh	[1575-1593] 3'UTR
149	392	AAAGGCUCUGAGACACAU	608	AUGUGUCUCAGGAGCCUUU		[1523-1541] 3'UTR
150	393	UGACCCAUGACCUGCAGAA	609	UUCUGCAGGUCAUGGGUCA	Rh,Rt,M	[1167-1185] ORF
151	394	CCUGUGAGACCAAUUGAG	610	CUCAAUUUGGUCUCACAGG	Rh	[1814-1832] 3'UTR
152	395	GCGGACCUUCCAGCUAGA	611	UCUAGCUGGGAAGGUCCGC	Rh	[1637-1655] 3'UTR
153	396	GGAAGAUGCAGAAGAAGGC	612	3CCUUCUUCUGCAUCUCC	Rh,Rb,Rt	[1113-1131] ORF
154	397	UGCCCAAGGGUGUGGUGGA	613	UCCACCACACCCUUGGGCA	Rh,D	[1146-1164] ORF
155	398	GGAGCCUCUCGAGCGCCUU	614	AAGGCGCUCGAGAGGCUCC		[1054-1072] ORF
156	399	GACUCUGGUCAAGAAGCAU	615	AUGCUUCUUGACCAGAGUC	Rh	[2013-2031] 3'UTR
157	400	CAGGCAGUGGAGAACAUC	616	3GAUGUUCUCCACUGCCUG	Rh	[413-431] ORF
158	401	CAAGCCUGCCCUAAUCAGU	617	ACUGAUUGAGGCAGGCUUG	Rh	[1766-1784] 3'UTR
159	402	CUGGAAGCUGGGCAGCCGA	618	UCGGCUGCCCAGCUUCCAG		[610-628] ORF
160	403	GAAGAAGGCUGUUGCCAUC	619	GAUGGCAACAGCCUUCUUC	Rt	[1123-1141] ORF
161	404	GGGCGAGCUGCUGCGCUCA	620	UGAGCGCAGCAGCUCGCC	Rh	[562-580] ORF
162	405	AAGCCACACUGGGAUGAGA	621	UCUCAUCCAGUGUGGCUU	Rh,Rb,M	[848-866] ORF
163	406	GUGUGGUGGAGGUGACCCA	622	UGGGUACCCUCCACCACAC	Rh,D	[1155-1173] ORF
164	407	CCGCCUUUGAGUUGGACAC	623	GUGUCCAACUCAAGGCGG	Rh	[1293-1311] ORF
165	408	3GCCAUUGACAAGAACAAG	624	CUUGUUCUUGUCAAUAGGCC	Rh,D	[1213-1231] ORF
166	409	UGCCUCAAUACGUAUUCAU	625	AUGAAUACUGAUUGAGGCA		[1772-1790] 3'UTR
167	410	CCUUCCCAGCUAGAAUUCA	626	UGAAUUCUAGCUGGGAAGG	Rh	[1642-1660] 3'UTR
168	411	GGGACCUGGGCCAUAGUCA	627	UGACUAUGGCCAGGUCCC		[1721-1739] 3'UTR
169	412	CGAGGUGAAGAAACCUGCA	628	UGCAGGUUUCUUCACCUCG	Rh	[286-304] ORF
170	413	GCCUUUGAGUUGGACACAG	629	CUGUGUCCAACUCAAGGC	Rh	[1295-1313] ORF
171	414	AGCGGACCUUCCAGCUAG	630	CUAGCUGGGAAGGUCCGCU	Rh	[1636-1654] 3'UTR
172	415	CGCAUGUCAGGCAAGAAGG	631	CCUUCUUGCCUGACAUGCG	Rh,D	[1244-1262] ORF
173	416	ACAACUGCGAGCACUCCAA	632	UUGGAGUGCUCGCAGUUGU	Rh,D	[690-708] ORF
174	417	3AGGCGGAUUGAGAAGGAG	633	CUCCUUCUCAAUCCGCCUC		[1971-1989] 3'UTR
175	418	GGCCGCCGAGGUGAAGAAA	634	UUUCUUCACCUCGGCGGCC		[280-298] ORF
176	419	CAGCUCUAUCCCAACCUCU	635	AGAGGUUGGGAUAGAGCUG		[1886-1904] 3'UTR
177	420	AGCUGGGCAGCCGACUGUA	636	UACAGUCGGCUGCCCAGCU		[615-633] ORF
178	421	GCCAUUGACAAGAACAAGG	637	CCUUGUUCUUGUCAAUGGC	Rh,D	[1214-1232] ORF
179	422	CGCCAUGUUCUUAAGCCA	638	UGGCUUGAAGAACAUGGCG	Rh,Rb,P	[835-853] ORF
180	423	CCGAGGUCACCAAGGACGU	639	ACGUCCUUGGUGACCUCGG	Rh,D	[786-804] ORF
181	424	GGACCCAGCUCAGUGAGCU	640	AGCUCACUGAGCUGGGUCC		[635-653] ORF
182	425	CAAUAGACAUUUUGUUGGA	641	UCCAACAAAUGUCAUUGG		[2178-2196] 3'UTR
183	426	AGUGAGGCGGAUUGAGAAG	642	CUUCUCAAUCCGCCUCACU		[1968-1986] 3'UTR
184	427	UGCAGUCCAUAACGAGUG	643	CACUCGUUGAUGGACUGCA	Rh,Rt,M	[738-756] ORF
185	428	UGUCACGCAUGUCAGGCAA	644	UUGCCUGACAUGCGUGACA	Rh,D	[1239-1257] ORF
186	429	CGACGACGAGAAGGAAAAG	645	CUUUUCCUUCUCGUCGUCG		[967-985] ORF

No	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
187	430	ACAAGAACAAGGCCGACUU	646	AAGUCGGCCUUGUUCUUGU	Rh	[1221-1239] ORF
188	431	CUUCAAGCCACACUGGGAU	647	AUCCAGUGUGGCUUGAAG	Rh,Rb,D	[844-862] ORF
189	432	CCUGGGCCAUAGUCAUUCU	648	AGAAUGACUAUGGCCAGG		[1725-1743] 3'UTR
190	433	UUUGUUGGAGCGUGGAAAA	649	UUUUCACGCUCCAACAAA		[2188-2206] 3'UTR
191	434	AGAACAUCCUGGUGUCACC	650	GGUGACACCAGGAUGUUCU		[423-441] ORF
192	435	ACGCCACCGCCUUUGAGUU	651	AACUCAAGGCGUGGCGU	Rh	[1287-1305] ORF
193	436	GUGAGGUACCAGCCUUGGA	652	UCCAAGGCUUGUACCUAC	Rh	[1567-1585] 3'UTR
194	437	GCGCCUUCUGCCUCCUGGA	653	UCCAGGAGGCAGAAGGCGC		[252-270] ORF
195	438	GCCUGGCCUUCAGCUUGUA	654	UACAAGCUGAAGGCCAGGC		[375-393] ORF
196	439	CCCGGAAACUCCACAUCU	655	AGGAUGUGGAGUUUCCGGG		[1700-1718] 3'UTR
197	440	UCUUAAGCCACACUGGGA	656	UCCCAGUGUGGCUUGAAGA	Rh,Rb,D	[843-861] ORF
198	441	UGUUGCUAUCAUCCAAGA	657	UCUUGGAUUGAUAGCAACA	Rh	[2123-2141] 3'UTR
199	442	GAGUGGGCCGCCGACACCA	658	UGGUCUGCGCGGCCACUC		[752-770] ORF
200	443	CCUGAGACACAUGGGUGCU	659	AGCACCCAUGUGUCUCAGG	D,Rt,M	[1530-1548] 3'UTR
201	444	AGCCGACUGUACGGACCCA	660	UGGGUCCGUACAGUCGGCU		[623-641] ORF
202	445	GGGCCUCAGGGUGCACACA	661	UGUGUGCACCCUGAGGCC		[1486-1504] 3'UTR
203	446	ACUGGGAUGAGAAAUUCCA	662	UGGAAUUUCUCAUCCAGU	Rh	[855-873] ORF
204	447	AGAAUGACCUGGCCGAGU	663	ACUGCGCCAGGUCAUUCU		[1952-1970] 3'UTR
205	448	CAUAUUUAUAGCCAGGUAC	664	GUACCUGGCUAUAUAUUG	Rh	[1788-1806] 3'UTR
206	449	AGGUGACCCAUGACCUGCA	665	UGCAGGUCAUGGGUCACCU	Rh,Rt,M	[1164-1182] ORF
207	450	GCGCUGCAGUCCAUAACG	666	CGUUGAUGGACUGCAGCGC	Rh,Rt	[734-752] ORF
208	451	GGUGACAAGAUGCGAGACG	667	CGUCUCGCAUCUUGUCACC	Rh	[1460-1478] ORF
209	452	CUUCAAGAUAGGGAGGGA	668	UCCCUCCUAUCUUUGAAG		[2086-2104] 3'UTR
210	453	AGCUGCAAAUCGUGGAGAU	669	AUCUCCACGAUUUGCAGCU	Rh	[984-1002] ORF
211	454	GUGGAGAACAUCCUGGUGU	670	ACACCAGGAUGUUCUCCAC	Rh	[419-437] ORF
212	455	GCAACAAGGCCGACUUGUCA	671	UGACAAGUCGGCCUUGUUC	Rh	[1225-1243] ORF
213	456	CAUGAUGCUGAGCCCGGAA	672	UUCGGGCUCAGCAUCAUG		[1688-1706] 3'UTR
214	457	GCGCCUUGAAAAGCUGCUA	673	UAGCAGCUUUUCAAGGCGC	Rh	[1066-1084] ORF
215	458	GCAGACUCUGGUCAAGAAG	674	CUUCUUGACCAGAGUCUGC	Rh	[2010-2028] 3'UTR
216	459	CCAGGCAGUGGAGAACAUC	675	GAUGUUCUCCACUGCCUGG	Rh	[412-430] ORF

TABLE C: Cross-Species 19-mer SERPINH1 siRNAs

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
1	676	CACUACAACUGCGAGCACU	973	AGUGCUCGCAGUUGUAGUG	Rh,D	[686-704] ORF
2	677	AACCGUGGCUUCAUGGUGA	974	UCACCAUGAAGCCACGGUU	Rh,Rt,M	[890-908] ORF
3	678	GGCAAGAAGGACCUGUACC	975	GGUACAGGUCCUUCUUGCC	Rh,D,M	[1253-1271] ORF
4	679	GGUGGACAACCGUGGCUUC	976	GAAGCCACGGUUGUCCACC	Rh,M	[883-901] ORF
5	680	AGGCCAUGGCCAAGGACCA	977	UGGUCCUUGGCCAUGGCCU	Rh,D	[396-414] ORF
6	681	CGCAGCGCGCUGCAGUCCA	978	UGGACUGCAGCGCGCUGCG	Rh,Rt	[728-746] ORF
7	682	AGCAGCAAGCAGCACUACA	979	UGUAGUGCUGCUUGCUGCU	Rh,D	[674-692] ORF
8	683	GGCCUCUACAACUACUACG	980	CGUAGUAGUUGUAGAGGCC	Rb,D	[950-968] ORF
9	684	GAAGAUGCAGAAGAAGGCU	981	AGCCUUCUUCUGCAUCUUC	Rh,Rb,Rt	[1114-1132] ORF
10	685	GGCUCCUGAGACACAUGGG	982	CCCAUGUGUCUCAGGAGCC	D	[1526-1544] 3'UTR
11	686	AGCAAGCAGCACUACAACU	983	AGUUGUAGUGCUGCUUGCU	Rh,D	[677-695] ORF
12	687	GGAGGUGACCCAUGACCUG	984	CAGGUCAUGGGUACCUCC	Rh,Rt,M	[1162-1180] ORF
13	688	CCUUUGACCAGGACAUCU	985	AGAUGUCCUGGUCAAAGGG	Rh,Rt	[1322-1340] ORF
14	689	CUCCUGAGACACAUGGGUG	986	CACCAUGUGUCUCAGGAG	D	[1528-1546] 3'UTR
15	690	AAGGCUCCUGAGACACAUG	987	CAUGUGUCUCAGGAGCCUU	D	[1524-1542] 3'UTR
16	691	CGCGCUGCAGUCCAUAAC	988	GUUGAUGGACUGCAGCGCG	Rh,Rt	[733-751] ORF
17	692	AGGGUGUGGUGGAGGUGAC	989	GUCACCUCCACCACCCU	Rh,D	[1152-1170] ORF
18	693	AGCACUACAACUGCGAGCA	990	UGCUCGCAGUUGUAGUGCU	Rh,D	[684-702] ORF
19	694	GGCUCCUGCUAUUCAUUG	991	CAAUGAAUAGCAGGGAGCC	D	[1421-1439] ORF
20	695	GCGCGCAACGUGACCUGGA	992	UCCAGGUCACGUUGCGCGC	M	[596-614] ORF
21	696	GCUGCAGUCCAUAACGAG	993	CUCGUUGAUGGACUGCAGC	Rh,Rt	[736-754] ORF

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
22	697	ACCAAAGAGCAGCUGAAGA	994	UCUUCAGCUGCUCUUUGGU	Rh,Rb,P	[1085-1103] ORF
23	698	CCAAGGACGUGGAGCGCAC	995	GUGCGCUCCACGUCCUUGG	Rh,D	[795-813] ORF
24	699	UGUUCUUAAGCCACACUG	996	CAGUGUGGCUUGAAGAACA	Rh,Rb,D	[840-858] ORF
25	700	GCCCAAGGGUGUGGUGGAG	997	CUCCACCACACCCUUGGGC	Rh,D	[1147-1165] ORF
26	701	ACAGGCCUCUACAACUACU	998	AGUAGUUGUAGAGGCCUGU	Rh,Rb,D,Rt,P	[947-965] ORF
27	702	UGCAGCAGCAGCAAGCAGCA	999	UGCUGCUUGCUGCUGCGCA	Rh,D	[689-687] ORF
28	703	GGUGGAGGUGACCCAUGAC	1000	GUCAUGGGUCACCUCACC	Rh,Rt,M	[1159-1177] ORF
29	704	CUUUGACCAGGACAUCUAC	1001	GUAGAUGUCCUGGUCAAAG	Rh,Rt	[1324-1342] ORF
30	705	AAGGGUGUGGUGGAGGUGA	1002	UCACCUCACCACACCCUU	Rh,D	[1151-1169] ORF
31	706	UCCUAUACCGUGGGUGUCA	1003	UGACACCCACGGUAUAGGA	Rh,D,P	[914-932] ORF
32	707	GCGCAGACCACCGACGGCA	1004	UGCCGUCGGUGGUCUGCGC	D	[761-779] ORF
33	708	CGCAGCAGCAAGCAGCACU	1005	AGUGCUGCUUGCUGCUGCG	Rh,D	[671-689] ORF
34	709	GCCUCAUCAUCCUCAUGCC	1006	GGCAUGAGGAUGAUGAGGC	Rh,D,Rt,M	[1026-1044] ORF
35	710	UCUCCAGCCUCAUCAUCCU	1007	AGGAUGAUGAGGCUGGAGA	Rh,D,Rt,M	[1020-1038] ORF
36	711	CCAUUGACAAGAACAAGGC	1008	GCCUUGUUCUUGUCAUUGG	Rh,D	[1215-1233] ORF
37	712	AGCAGCACUACAACUGCGA	1009	UCGCAGUUGUAGUOCUGCU	Rh,D	[681-699] ORF
38	713	UGCACCGGACAGGCCUCUA	1010	UAGAGGCCUGUCCGGUGCA	Rh,Rb,Rt,P	[939-957] ORF
39	714	ACUCCAAGAUCAACUCCG	1011	CGGAAGUUGAUUUUGGAGU	Rh,D,Rt,M	[702-720] ORF
40	715	UGGACAACCGUGGCUUCAU	1012	AUGAAGCCACGGUUGUCCA	Rh,M	[885-903] ORF
41	716	GAGCAGCUGAAGAUCUGGA	1013	UCCAGAUUCUAGCUGCUC	Rh,D	[1091-1109] ORF
42	717	CAGAAGAAGGCUUUGCCA	1014	UGGCAACAGCCUUCUUCUG	Rt	[1121-1139] ORF
43	718	AGGCAAGAAGGACCUGUAC	1015	GUACAGGUCCUUCUUGCCU	Rh,D	[1252-1270] ORF
44	719	CCUCUACAACUACUACGAC	1016	GUCGUAGUAGUUGUAGAGG	Rb,D	[952-970] ORF
45	720	AGCAGCUGAAGAUUGGAU	1017	AUCCAGAUCUUCAGCUGCU	Rh,D	[1092-1110] ORF
46	721	AACUACUACGACGACGAGA	1018	UCUCGUCGUCGUAGUAGUU	Rb	[959-977] ORF
47	722	GGCAAGCUGCCGAGGUCA	1019	UGACCUCGGGCAGCUUGCC	Rh,D	[776-794] ORF
48	723	CCGGACAGGCCUCUACAAC	1020	GUUGUAGAGGCCUGUCCGG	Rh,Rb,Rt,P	[943-961] ORF
49	724	GCUCCUGCUAUUCAUUGG	1021	CCAAUGAAUAGCAGGGAGC	D	[1422-1440] ORF
50	725	AACUGCGAGCACUCCAAGA	1022	UCUUGGAGUGCUCGCAGUU	Rh,D	[692-710] ORF
51	726	GACACAUUGGUGCUAUUGG	1023	CCAAUAGCACCCAUGUGUC	Rh,Rt,M	[1535-1553] 3'UTR
52	727	GCACCGGACAGGCCUCUAC	1024	GUAGAGGCCUGUCCGGUGC	Rh,Rb,Rt,P	[940-958] ORF
53	728	AGCGCAGCGCGCUGCAGUC	1025	GACUCGAGCGCGCUGCGCU	Rh,Rt	[726-744] ORF
54	729	GGACGUGGAGCGCACGGAC	1026	GUCCGUGCGCUCCACGUCC	Rh,D	[799-817] ORF
55	730	CAGCCUCAUCAUCCUCAUG	1027	CAUGAGGAUGAUGAGGCUG	Rh,D,Rt,M	[1024-1042] ORF
56	731	AAGAUCAACUCCGCGACA	1028	UGUCGCGGAAGUUGAUCUU	D	[707-725] RF
57	732	GCGCAACGUGACCUGGAAG	1029	CUUCCAGGUCACGUUGCGC	M	[598-616] ORF
58	733	ACUGCGAGCACUCCAAGAU	1030	AUCUUGGAGUGCUCGCAGU	Rh,D	[693-711] ORF
59	734	3UGGACAACCGUGGCUUCA	1031	UGAAGCCACGGUUGUCCAC	Rh,M	[884-902] ORF
60	735	CCACAAGCUCUCCAGCCUC	1032	GAGGCUGGAGAGCUUGUGG	Rh,D,P	[1012-1030] ORF
61	736	CAAGAUGGUGGACAACCGU	1033	ACGGUUGUCCACCAUCUUG	Rh,Rb,M,P	[877-895] ORF
62	737	CGAGCACUCCAAGAUCAAC	1034	GUUGAUCUUGGAGUGCUCG	Rh,D	[697-715] ORF
63	738	AGCUGCCCCGAGGUCACCAA	1035	UUGGUGACCUCGGGCAGCU	Rh,D	[780-798] ORF
64	739	GGACAUCUACGGGCGCGAG	1036	CUCGCGCCCGUAGAUGUCC	D	[1333-1351] ORF
65	740	AGGACAUCUACGGGCGCGA	1037	UCGCGCCCGUAGAUGUCCU	D	[1332-1350] ORF
66	741	UGUCAGGCAAGAAGGACCU	1038	AGGUCCUUCUUGCCUGACA	Rh,D	[1248-1266] ORF
67	742	GGGUGUGGUGGAGGUGACC	1039	GGUACCUCCACCACACC	Rh,D	[1153-1171] ORF
68	743	CAAGCUCUCCAGCCUCAUC	1040	GAUGAGGCUGGAGAGCUUG	Rh,D,M,P	[1015-1033] ORF
69	744	GUGACCCAUGACCUGCAGA	1041	UCUGCAGGUCAUGGGUCAC	Rh,Rt,M	[1166-1184] ORF
70	745	GUUCUUAAGCCACACUGG	1042	CCAGUGUGGCUUGAAGAAC	Rh,Rb,D	[841-859] ORF
71	746	ACAUCUACGGGCGCGAGGA	1043	UCCUCGCGCCCGUAGAUGU	D,M	[1335-1353] ORF
72	747	UGGAGGUGACCCAUGACCU	1044	AGGUCAUGGGUACCUCCA	Rh,Rt,M	[1161-1179] ORF
73	748	UGCAGAGAAGGCUUUGC	1045	GCAACAGCCUUCUUCUGCA	Rt	[1119-1137] ORF
74	749	UGUACCAGGCCAUGGCCAA	1046	UUGGCCAUGGCCUGGUACA	Rh,D	[390-408] ORF

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
75	750	UGUGGUGGAGGUGACCCAU	1047	AUGGGUACACCUCCACCACA	Rh,D	[1156-1174] ORF
76	751	AGAAGGACCUUGUACCUGGC	1048	GCCAGGUACAGGUCCUUCU	Rh,D	[1257-1275] ORF
77	752	AGCAGCUGCGCGACGAGGA	1049	UCCUCGUCGCGCAGCUGCU	Rh,D	[528-546] ORF
78	753	ACGCCAUGUUCUUAAGCC	1050	GGCUUGAAGAACAUGGCGU	Rh,Rb,P	[834-852] ORF
79	754	ACAAGAUGGUGGACAACCG	1051	CGGUUGUCCACCAUCUUGU	Rh,Rb,M,P	[876-894] ORF
80	755	CUGCGAGCACUCCAAGAUC	1052	GAUCUUGGAGUGCUCGCAG	Rh,D	[694-712] ORF
81	756	GUCACGCAUGUCAGGCAAG	1053	CUUGCCUGACAUGCGUGAC	Rh,D	[1240-1258] ORF
82	757	ACGCAUGUCAGGCAAGAAG	1054	CUUCUUGCCUGACAUGCGU	Rh,D	[1243-1261] ORF
83	758	UGCUAUUAUUGGGCGCCU	1055	AGGCGCCCAUGAAUAGCA	D	[1428-1446] ORF
84	759	UGC GCGACGAGGAGGUGCA	1056	UGCACCUCCUCGUCGCGCA	Rh,D	[534-552] ORF
85	760	GCAGCUGAAGAUCUGGAUG	1057	CAUCCAGAUUCUACGCGC	Rh,D	[1093-1111] ORF
86	761	CCAUGACCUGCAGAAACAC	1058	GUGUUUCUGCAGGUCAUGG	Rh,Rt,M	[1171-1189] ORF
87	762	AAGCUCUCCAGCCUCAUCA	1059	UGAUGAGGCUUGGAGAGCUU	Rh,D,Rt,M,P	[1016-1034] ORF
88	763	CAGCAAGCAGCACUACAAC	1060	GUUGUAGUGCUGCUUGCUG	Rh,D	[676-694] ORF
89	764	AUGUUCUUAAGCCACACU	1061	AGUGUGGCUUGAAGAACA	Rh,Rb,D	[839-857] ORF
90	765	UCCUGAGACACAUGGGUGC	1062	GCACCAUGUGUCUCAGGA	D,Rt,M	[1529-1547] 3'UTR
91	766	CACUCCAAGAUCAACUUC	1063	GGAAGUUGAUUCUUGGAGUG	Rh,D,Rt,M	[701-719] ORF
92	767	AAGGGUGACAAGAUCCGAG	1064	CUCGCAUCUUGUCACCCUU	Rh,D	[1457-1475] ORF
93	768	GACAGGCCUCUACAACUAC	1065	GUAGUUGUAGAGGCCUGUC	Rh,Rb,Rt,P	[946-964] ORF
94	769	ACCCAUGACCUGCAGAAAC	1066	GUUUCUGCAGGUCAUGGGU	Rh,Rt,M	[1169-1187] ORF
95	770	CACCAACAAGAUUGGUGACA	1067	UGUCCACCAUCUUGUGGUG	Rh,Rb,M,P	[872-890] ORF
96	771	GCAGAAGAAGGCUGUUGCC	1068	GGCAACAGCCUUCUUCUGC	Rt	[1120-1138] ORF
97	772	GUGGUGGAGGUGACCCAUG	1069	CAUGGGUACACCUCCACCAC	Rh,Rb,Rt,M	[1157-1175] ORF
98	773	AGGCCUCUACAACUACUAC	1070	GUAGUAGUUGUAGAGGCCU	Rh,Rb,D,Rt,P	[949-967] ORF
99	774	GGUGACCCAUGACCUGCAG	1071	CUGCAGGUCAUGGGUACCC	Rh,Rt,M	[1165-1183] ORF
100	775	GCCGAGGUGAAGAAACCUG	1072	CAGGUUUCUUCACCUCGGC	Rh,Rt	[284-302] ORF
101	776	CAACUACUACGACGACGAG	1073	CUCGUCGUCGUAGUAGUUG	Rb	[958-976] ORF
102	777	CAAGAAGGACCUUGUACCUG	1074	CAGGUACAGGUCCUUCUUG	Rh,D,M	[1255-1273] ORF
103	778	UGUUCCACGCCACCGCCUU	1075	AAGGCGGUGGCGUGGAACA	D	[1281-1299] ORF
104	779	CCCUGCUAUUCAUUGGGCG	1076	CGCCCAUGAAUAGCAGGG	D	[1425-1443] ORF
105	780	CCGUGGCUUCAUGGUGACU	1077	AGUCACCAUGAAGCCACGG	Rh,Rt,M	[892-910] ORF
106	781	CUACAACUACUACGACGAC	1078	GUCGUCGUAGUAGUUGUAG	Rb	[955-973] ORF
	107 782	GCAGCACUACAACUGCGAG	1079	CUCGCGAGUUGUAGUGCUGC	Rh,D	[682-700] ORF
108	783	UGGUGGACAACCGUGGCUU	1080	AAGCCACGGUUGUCCACCA	Rh,M	[882-900] ORF
109	784	AGACCACCGACGGAAGCU	1081	AGCUUGCCGUCGGUGGUCU	D,Rt	[765-783] ORF
110	785	AGAAACACCUGGCUGGGCU	1082	AGCCAGCCAGGUGUUUCU	D	[1182-1200] ORF
111	786	ACCAAGGACGUGGAGCGCA	1083	UGCUCUCCAGUCCUUGGU	Rh,D	[794-812] ORF
112	787	CCGAGGUGAAGAAACCUGC	1084	GCAGGUUUCUUCACCUCGG	Rh,Rt	[285-303] ORF
113	788	ACUACAACUGCGAGCACUC	1085	GAGUGCUCGCAGUUGUAGU	Rh,D	[687-705] ORF
114	789	ACAAGCUCUCCAGCCUCAU	1086	AUGAGGCUUGGAGAGCUUGU	Rh,D,M,P	[1014-1032] ORF
115	790	AGGACGUGGAGCGCACGGA	1087	UCCGUGCGCUCCACGUCCU	Rh,D	[798-816] ORF
116	791	GCUAUUCAUUGGGCGCCUG	1088	CAGGCGCCCAUGAAUAGC	D	[1429-1447] ORF
117	792	AACUUCGCGACAAGCGCA	1089	UGCUCUUGUCGCGGAAGUU	D	[713-731] ORF
118	793	GCUCUCCAGCCUCAUCAUC	1090	GAUGAUGAGGCUUGGAGAGC	Rh,D,Rt,M,P	[1018-1036] ORF
119	794	AGAAGGCUUGUCCAUCUC	1091	GAGAUGGCAACAGCCUUCU	Rt	[1125-1143] ORF
120	795	GGUACCAAGGACGUGGAG	1092	CUCCACGUCCUUGGUGACC	Rh,D	[790-808] ORF
121	796	AGCUGCGCGACGAGGAGGU	1093	ACCUCUCGUCGCGCAGCU	Rh,D	[531-549] ORF
122	797	CCCGAGGUCACCAAGGACG	1094	CGUCCUUGGUGACCUCGGG	Rh,D	[785-803] ORF
123	798	AUGUCAGGCAAGAAGGACC	1095	GGUCCUUCUUGCCUGACAU	Rh,D	[1247-1265] ORF
124	799	CGAGGUCACCAAGGACGUG	1096	CACGUCCUUGGUGACCUCG	Rh,D	[787-805] ORF
125	800	GAUGCACCGGACAGGCCUC	1097	GAGGCCUGUCCGGUGCAUC	Rh,Rb,Rt,M,P	[937-955] ORF
126	801	3CACUACAACUGCGAGCAC	1098	GUGCUCGCAGUUGUAGUGC	Rh,D	[685-703] ORF
127	802	CCACAAGAUGGUGGACAAC	1099	GUUGUCCACCAUCUUGUGG	Rh,Rb,M,P	[874-892] ORF

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
128	803	CAAGGGUGUGGUGGAGGUG	1100	CACCUCCACCACACCCUUG	Rh,D	[1150-1168] ORF
129	804	AGCUGAAGAUUCUGGAUGGG	1101	CCCAUCCAGAUCUUCAGCU	Rh,D	[1095-1113] ORF
130	805	ACCAGGCCAUGGCCAAGGA	1102	UCCUUGGCCAUGGCCUGGU	Rh,D	[393-411] ORF
131	806	CAUGUUCUUC AAGCCACAC	1103	GUGUGGCUUGAAGAACAUG	Rh,Rb,D	[838-856] ORF
132	807	CAAGAUCAACUCCGCGAC	1104	GUCGCGGAAGUUGAUCUUG	D	[706-724] ORF
133	808	UCCAGCCUCAUCAUCCUCA	1105	UGAGGAUGAUGAGGCUGGA	Rh,D,Rt,M	[1022-1040] ORF
134	809	GCCCCAGGUCACCAAGGAC	1106	GUCCUUGGUGACCUCGGGC	Rh,D	[784-802] ORF
135	810	UCAAGCCACACUGGGAUGA	1107	UCAUCCCAGUGUGGCUUGA	Rh,Rb	[846-864] ORF
136	811	AGUCCAUAACGAGUGGGC	1108	GCCCACUCGUUGAUGGACU	Rh,Rt,M	[741-759] ORF
137	1812	GACUUCGUGCGCAGCAGCA	1109	UGCUGCUGCGCACGAAGUC	Rh,D,M	[662-680] ORF
138	813	CUCUCCAGCCUCAUCAUCC	1110	GGAUGAUGAGGCUGGAGAG	Rh,D,Rt,M,P	[1019-1037] ORF
139	814	GCAGACCACCGACGGCAAG	1111	CUUGCCGUCGGUGGUCUGC	D,Rt	[763-781] ORF
140	815	AUGCAGAAGAAGGCUGUUG	1112	CAACAGCCUUCUUCUGCAU	Rt	[1118-1136] ORF
141	816	CAACCGUGGCUUCAUGGUG	1113	CACCAUGAAGCCACGGUUG	Rh,Rt,M	[889-907] ORF
142	817	UACUACGACGACGAGAAGG	1114	CCUUCUCGUCGUCGUAGUA	Rb	[962-980] ORF
143	818	GAAGGCUGUUGCCAUCUCC	1115	GGAGAUGGCAACAGCCUUC	Rt	[1126-1144] ORF
144	819	UCACCAAGGACGUGGAGCG	1116	CGCUCCACGUCCUUGGUGA	Rh,D	[792-810] ORF
145	820	CAGCUGAAGAUUCUGGAUGG	1117	CCAUCCAGAUCUUCAGCUG	Rh,D	[1094-1112] ORF
146	821	UGGGCCUGACUGAGGCCAU	1118	AUGGCCUCAGUCAGGCCCA	Rt	[1200-1218] ORF
147	822	ACCGUGGCUUCAUGGUGAC	1119	GUCACCAUGAAGCCACGGU	Rh,Rt,M	[891-909] ORF
148	823	CAGUCCAUAACGAGUGGG	1120	CCCACUCGUUGAUGGACUG	Rh,Rt,M	[740-758] ORF
149	824	CCGACGGCAAGCUGCCCCGA	1121	UCGGGCAGCUUGCCGUCGG	D	[771-789] ORF
150	825	ACAAGCGCAGCGCGCUGCA	1122	UGCAGCGCGCUGCGCUUGU	Rh,Rt	[723-741] ORF
151	826	GAAACACCUGGCUGGGCUG	1123	CAGCCCAGCCAGGUGUUUC	D	[1183-1201] ORF
152	827	AGGCUCUGAGACACAUGG	1124	CCAUGUGUCUCAGGAGCCU	D	[1525-1543] 3'UTR
153	828	CAAGGACGUGGAGCGCACG	1125	CGUGCGCUCCACGUCCUUG	Rh,D	[796-814] ORF
154	829	GCAGUCCAUAACGAGUGG	1126	CCACUCGUUGAUGGACUGC	Rh,Rt,M	[739-757] ORF
155	830	AGAUGGUGGACAACCGUGG	1127	CCACGGUUGUCCACCAUCU	Rh,M	[879-897] ORF
156	831	AAGCGCAGCGCGCUGCAGU	1128	ACUGCAGCGCGCUGCGCUU	Rh,Rt	[725-743] ORF
157	832	CAUGUCAGGCAAGAAGGAC	1129	GUCCUUCUUGCCUGACAUG	Rh,D	[1246-1264] ORF
158	833	CAAGCCACACUGGGAUGAG	1130	CUCAUCCAGUGUGGCUUG	Rh,Rb	[847-865] ORF
159	834	AAGAUGCAGAAGAAGGCUG	1131	CAGCCUUCUUCUGCAUCUU	Rh,Rt,M	[1115-1133] ORF
160	835	GGCCAUGGCCAAGGACCAG	1132	CUGGUCCUUGGCCAUGGCC	Rh,D	[397-415] ORF
161	836	GUGCGCAGCAGCAAGCAGC	1133	GCUGCUUGCUGCUGCGCAC	Rh,D	[668-686] ORF
162	837	CAACUGCGAGCACUCCAAG	1134	CUUGGAGUGCUCGCAGUUG	Rb,D	[691-709] ORF
163	838	UACAACUGCGAGCACUCCA	1135	UGGAGUGCUCGCAGUUGUA	Rh,D	[689-707] ORF
164	839	CAUUGACAAGAACAAGGCC	1136	GGCCUUGUUCUUGUCAUUG	Rh,D	[1216-1234] ORF
165	840	CAAGCAGCACUACAACUGC	1137	GCAGUUGUAGUGCUGCUUG	Rh,D	[679-697] ORF
166	841	GUGUUCCACGCCACCGCCU	1138	AGGCGGUGGCGUGGAACAC	D	[1280-1298] ORF
167	842	CCUGCUAUUCAUUGGGCGC	1139	GCGCCCAAUGAAUAGCAGG	D	[1426-1444] ORF
168	843	GCCCCAAGCUCUCCAGCC	1140	GGCUGGAGAGCUUGUGGGC	Rh,D,P	[1010-1028] ORF
169	844	CAGCAGCAAGCAGCACUAC	1141	3UAGUGCUGCUUGCUGCUG	Rh,D	[673-691] ORF
170	845	UGAUGCACCGGACAGGCCU	1142	AGGCCUGUCCGGUGCAUCA	Rb,Rb,Rt,M,P	[936-954] ORF
171	846	UCAACUUCGCGACAAGCG	1143	CGCUUGUCGCGGAAGUUGA	D	[711-729] ORF
172	847	UCAGGCAAGAAGGACCUGU	1144	ACAGGUCCUUCUUGCCUGA	Rh,D	[1250-1268] ORF
173	848	ACUUCGUGCGCAGCAGCAA	1145	UUGCUGCUGCGCACGAAGU	Rh,D,M	[663-681] ORF
174	849	ACAACGUGGCUUCAUGGU	1146	ACCAUGAAGCCACGGUUGU	Rh,Rt,M	[888-906] ORF
175	850	AAGGCUGUUGCCAUCUCCU	1147	AGGAGAUGGCAACAGCCUU	D,Rt	[1127-1145] ORF
176	851	GCAGCUGCGCGACGAGGAG	1148	CUCCUCGUCGCGCAGCUGC	Rh,D	[529-547] ORF
177	852	UAUUCAUUGGGCGCCUGGU	1149	ACCAGGCGCCCAUGAAUA	D	[1431-1449] ORF
178	853	UCCACCACAAGAUGGUGGA	1150	UCCACCAUCUUGUGGUGGA	Rh,Rb,D,P	[870-888] ORF
179	854	CCCUUGGCCACAAGCUCUC	1151	GAGAGCUUGUGGGCCAGGG	Rh,D,P	[1005-1023] ORF
180	855	ACCAGGACAUCUACGGGCG	1152	CGCCCGUAGAUGUCCUGGU	D,Rt	[1329-1347] ORF

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
181	856	GAUGAUGCACCGGACAGGC	1153	GCCUGUCCGGUGCAUCAUC	Rh,Rb,Rt,M	[934-952] ORF
182	857	CAACGCCAUGUUCUUAAG	1154	CUUGAAGAACAUGGCGUUG	Rh,Rb,P	[832-850] ORF
183	858	ACGGCAAGCUGCCCGAGGU	1155	ACCUCGGGCAGCUUGCCGU	Rh,D	[774-792] ORF
184	859	CAGCGCGCUGCAGUCCAUC	1156	GAUGGACUGCAGCGCGCUG	Rh,Rt	[730-748] ORF
185	860	CCCAAGGGUGUGGUGGAGG	1157	CCUCCACCACACCCUUGGG	Rh,D	[1148-1166] ORF
186	861	CAUGGCCAAGGACCAGGCA	1158	UGCCUGGUCCUUGGCCAUG	Rh,D	[400-418] ORF
187	862	CUCCAGCCUCAUCAUCCUC	1159	GAGGAUGAUGAGGCUGGAG	Rh,D,Rt,M	[1021-1039] ORF
188	863	UCUACGGGCGCGAGGAGCU	1160	AGCUCCUCGCGCCCGUAGA	D,M	[1338-1356] ORF
189	864	3GCCCACAAGCUCUCCAGC	1161	GCUGGAGAGCUUGUGGGCC	Rh,D,P	[1009-1027] ORF
190	865	GUCAGGCAAGAAGGACCUG	1162	CAGGUCCUUCUUGCCUGAC	Rh,D	[1249-1267] ORF
191	866	CAUCUACGGGCGCGAGGAG	1163	CUCCUCGCGCCCGUAGAUG	D,M	[1336-1354] ORF
192	867	CGUGCGCAGCAGCAAGCAG	1164	CUGCUUGCUGCUGCGCACG	Rh,D,M	[667-685] ORF
193	868	AGCCUCAUCAUCCUCAUGC	1165	GCAUGAGGAUGAUGAGGCU	Rh,D,Rt,M	[1025-1043] ORF
194	869	UUCAAGCCACACUGGGAUG	1166	CAUCCAGUGUGGCUUGAA	Rh,Rb	[845-863] ORF
195	870	AAGAAGGCGUGUUGCCAUCU	1167	AGAUGGCAACAGCCUUCUU	Rt	[1124-1142] ORF
196	871	GGUGUGGUGGAGGUGACCC	1168	GGGUCACCUCCACCACACC	Rh,D	[1154-1172] ORF
197	872	3AGGUGACCCAUGACCUGC	1169	GCAGGUCAUGGGUCACCUC	Rh,Rt,M	[1163-1181] ORF
198	873	GUGGAGGUGACCCAUGACC	1170	GGUCAUGGGUCACCUCAC	Rh,Rt,M	[1160-1178] ORF
199	874	CACAAGAUGGUGGACAACC	1171	GGUUGUCCACCAUCUUGUG	Rh,Rb,M,P	[875-893] ORF
200	875	CUGGCCCAAGCUCUCCA	1172	UGGAGAGCUUGUGGGCCAG	Rh,D,P	[1007-1025] ORF
201	876	GAUGACUUCGUGCGCAGCA	1173	UGCUGCGCACGAAGUCAUC	Rh,Rt,M	[659-677] ORF
202	877	ACUUCGCGACAAGCGCAG	1174	CUGCGCUUGUCGCGGAAGU	D	[714-732] ORF
203	878	AACGCCAUGUUCUUAAGC	1175	GCUUGAAGAACAUGGCGUU	Rh,Rb,P	[833-851] ORF
204	879	GGACCUGUACCUGGCCAGC	1176	GCUGGCCAGGUACAGGUCC	Rh,D	[1261-1279] ORF
205	880	GCGACGAGGAGGUGCACGC	1177	GCGUGCACCUCCUCGUCGC	D	[537-555] ORF
206	881	GCAAGCUGCCCGAGGUCAC	1178	GUGACCUCGGGCAGCUUGC	Rh,D	[777-795] ORF
207	882	AUUCAUUGGGCGCCUGGUC	1179	GACCAGGCGCCCAUGAAU	D	[1432-1450] ORF
208	883	GAGGUCACCAAGGACGUGG	1180	CCACGUCCUUGGUGACCUC	Rh,D	[788-806] ORF
209	884	AAGAAGGACCUGUACCUGG	1181	CCAGGUACAGGUCCUUCUU	Rh,D	[1256-1274] ORF
210	885	GACAACCGUGGCUUCAUGG	1182	CCAUGAAGCCACGGUUGUC	Rh,Rt,M	[887-905] ORF
211	886	CUGGGCCUGACUGAGGCCA	1183	UGGCCUCAGUCAGGCCAG	Rt	[1199-1217] ORF
212	887	CUCCAAGAUCAACUCCGC	1184	GCGGAAGUUGAUCUUGGAG	Rh,D,Rt,M	[703-721] ORF
213	888	CAACUUCGCGACAAGCGC	1185	GCGCUUGUCGCGGAAGUUG	D	[712-730] ORF
214	889	CUCCUGGCUAUUCAUUGGG	1186	CCCAAUGAAUAGCAGGGAG	D	[1423-1441] ORF
215	890	AAGCAGCACUACAACUGCG	1187	CGCAGUUGUAGUGCUGCUU	Rh,D	[680-698] ORF
216	891	GCGCAGCAGCAAGCAGCAC	1188	GUGCUGCUUGCUGCUGCGC	Rh,D	[670-688] ORF
217	892	CAGGCCAUGGCCAAGGACC	1189	GUCCUUGGCCAUGGCCUG	Rh,D	[395-413] ORF
218	893	GUACCAGGCCAUGGCCAAG	1190	CUUGGCCAUGGCCUGGUAC	Rh,D	[391-409] ORF
219	894	CUUCGUGCGCAGCAGCAAG	1191	CUUGCUGCUGCGCACGAAG	Rh,D,M	[664-682] ORF
220	895	CAGCACUACAACUGCGAGC	1192	GCUCGCAGUUGUAGUGCUG	Rh,D	[683-701] ORF
221	896	UACAACUACUACGACGACG	1193	CGUCGUCGUAGUAGUUGUA	Rb	[956-974] ORF
222	897	GAUGGUGGACAACCGUGGC	1194	GCCACGGUUGUCCACCAUC	Rh,M	[880-898] ORF
223	898	CUACAACUGCGAGCACUCC	1195	GGAGUGCUCGCGAGUUGUAG	Rh,D	[688-706] ORF
224	899	AAGGACCUGUACCUGGCCA	1196	UGGCCAGGUACAGGUCCUU	Rh,D	[1259-1277] ORF
225	900	GCUGCCCGAGGUCACCAAG	1197	CUUGGUGACCUCGGGCAGC	Rh,D	[781-799] ORF
226	901	GACAUCUACGGGCGCGAGG	1198	CCUCGCGCCCGUAGAUGUC	D,M	[1334-1352] ORF
227	902	CCACCACAAGAUGGUGGAC	1199	GUCCACCAUCUUGUGGUGG	Rh,Rb,D,P	[871-889] ORF
228	903	GCGCGACGAGGAGGUGCAC	1200	GUGCACCUCUCGUCGCGC	Rh,D	[535-553] ORF
229	904	CUAUUCAUUGGGCGCCUGG	1201	CCAGGCGCCCAUGAAUAG	D	[1430-1448] ORF
230	905	CCAGGACAUCUACGGGCGC	1202	GCGCCCGUAGAUGUCCUGG	D,Rt	[1330-1348] ORF
231	906	AAGAUGGUGGACAACCGUG	1203	CACGGUUGUCCACCAUCUU	Rh,M	[878-896] ORF
232	907	CAGGACAUCUACGGGCGCG	1204	CGCGCCCGUAGAUGUCCUG	D	[1331-1349] ORF
233	908	UCCAAGAUCAACUCCGCG	1205	CGCGGAAGUUGAUCUUGGA	D	[704-722] ORF

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
234	909	GUCACCAAGGACGUGGAGC	1206	GCUCCACGUCCUUGGUGAC	Rh,D	[791-809] ORF
235	910	CUGCCCCGAGGUACCAAGG	1207	CCUUGGUGACCUCGGGCAG	Rh,D	[782-800] ORF
236	911	GACCAGGACAUCUACGGGC	1208	3CCCGUAGAUGUCCUGGUC	D,Rt	[1328-1346] ORF
237	912	CCAUGGCCAAGGACCAGGC	1209	GCCUGGUCCUUGGCCAUGG	Rh,D	[399-417] ORF
238	913	CACCAAGGACGUGGAGCGC	1210	GCGCUCCACGUCCUUGGUG	Rh,D	[793-811] ORF
239	914	3ACAAGCGCAGCGCGCUGC	1211	GCAGCGCGCUGCGCUUGUC	Rh,Rt	[722-740] ORF
240	915	CAAGCGCAGCGCGCUGCAG	1212	CUGCAGCGCGCUGCGCUUG	Rh,Rt	[724-742] ORF
241	916	CAGACCACGACGGCAAGC	1213	GCUUGCCGUCGGUGGUCUG	D,Rt	[764-782] ORF
242	917	GACCACGACGGCAAGCUG	1214	CAGCUUGCCGUCGGUGGUC	D,Rt	[766-784] ORF
243	918	AGGACCUGUACCUGGCCAG	1215	CUGGCCAGGUACAGGUCCU	Rh,D	[1260-1278] ORF
244	919	CUGCUAUUCAUUGGGCGCC	1216	GGCGCCCAAUGAAUAGCAG	D	[1427-1445] ORF
245	920	UCAUUGGGCGCCUGGUCCG	1217	GGACCAGGCGCCCAAUGA	Rh,D	[1434-1452] ORF
246	921	GCUGCGCGACGAGGAGGUG	1218	CACCUCCUCGUCGCGCAGC	Rh,D	[532-550] ORF
247	922	CGGCAAGCUGCCGAGGUC	1219	GACCUCGGGCAGCUUGCCG	Rh,D	[775-793] ORF
248	923	CCUCAUCAUCCUCAUGCCC	1220	GGGCAUGAGGAUGAUGAGG	Rh,D,Rt,M	[1027-1045] ORF
249	924	CCAGGCCAUGGCCAAGGAC	1221	GUCCUUGGCCAUGGCCUGG	Rh,D	[394-412] ORF
250	925	GCCAUGGCCAAGGACCAGG	1222	CCUGGUCCUUGGCCAUGGC	Rh,D	[398-416] ORF
251	926	CCACCGACGGCAAGCUGCC	1223	GGCAGCUUGCCGUCGGUGG	D,Rt	[768-786] ORF
252	927	AUGGUGGACAACCGUGGCU	1224	AGCCACGGUUGUCCACCAU	Rh,M	[881-899] ORF
253	928	CUUCCGCGACAAGCGCAGC	1225	GCUGCGCUUGUCGCGGAAG	D	[715-733] ORF
254	929	CGCGACGAGGAGGUGCAGC	1226	CGUGCACCUCUCGUCGCG	Rh,D	[536-554] ORF
255	930	UGGCCACACAAGCUCUCCAG	1227	CUGGAGAGCUUGUGGGCCA	Rh,D,P	[1008-1026] ORF
256	931	GAGCAGCUGCGCGACGAGG	1228	CCUCGUCGCGCAGCUGCUC	Rh,D	[527-545] ORF
257	932	UGACCAGGACAUCUACGGG	1229	CCCGUAGAUGUCCUGGUCA	Rt	[1327-1345] ORF
258	933	ACCACGACGGCAAGCUGC	1230	GCAGCUUGCCGUCGGUGGU	D,Rt	[767-785] ORF
259	934	GAAGGACCUGUACCUGGCC	1231	GGCCAGGUACAGGUCCUUC	Rh,D	[1258-1276] ORF
260	935	CAUUGGGCGCCUGGUCCGG	1232	CCGGACCAGGCGCCCAAUG	Rh,D	[1435-1453] ORF
261	936	AUGCACCGACAGGCCUCU	1233	AGAGGCCUGUCCGGUGCAU	Rh,Rb,Rt,P	[938-956] ORF
262	937	AUACAUCUCCGCGACAAGC	1234	GCUUGUCGCGGAAGUUGAU	D	[710-728] ORF
263	938	CAGCUGCGCGACGAGGAGG	1235	CCUCCUCGUCGCGCAGCUG	Rh,D	[530-548] ORF
264	939	CAGAAACACCUGGCUGGGC	1236	GCCAGCCAGGUGUUUCUG	D	[1181-1199] ORF
265	940	CUACGGGCGCGAGGAGCUG	1237	CAGCUCCUCGCGCCCGUAG	D,M	[1339-1357] ORF
266	941	CGACGAGGAGGUGCACGCC	1238	GGCGUGCACCUCUCGUCG	D	[538-556] ORF
267	942	UUUGACCAGGACAUCUACG	1239	CGUAGAUGUCCUGGUCAAA	Rt	[1325-1343] ORF
268	943	GUCCAUCAACGAGUGGGCC	1240	GGCCACUCGUUGAUGGAC	Rh,Rt,M	[742-760] ORF
269	944	AUGACUUCGUGCGCAGCAG	1241	CUGCUGCGCACGAAGUCAU	Rh,Rt,M	[660-678] ORF
270	945	UCCUGCUAUUCAUUGGGC	1242	GCCCAAUGAAUAGCAGGGA	D	[1424-1442] ORF
271	946	CUGCGCGACGAGGAGGUGC	1243	GCACCUCCUCGUCGCGCAG	Rh,D	[533-551] ORF
272	947	CAAGCUGCCCGAGGUCACC	1244	GGUGACCUCGGGCAGCUUG	Rh,D	[778-796] ORF
273	948	AAGCUGCCCGAGGUCACCA	1245	UGGUGACCUCGGGCAGCUU	Rh,D	[779-797] ORF
274	949	UUCUUCAAGCCACACUGGG	1246	CCCAGUGUGGCUUGAAGAA	Rh,Rb,D	[842-860] ORF
275	950	ACACCUGGCUGGGCUCGGGC	1247	GCCAGCCCAGCCAGGUGU	D	[1186-1204] ORF
276	951	UCCAUCAACGAGUGGGCCG	1248	CGGCCACUCGUUGAUGGA	Rt,M	[743-761] ORF
277	952	AUCUACGGGCGCGAGGAGC	1249	GCUCCUCGCGCCCGUAGAU	D,M	[1337-1355] ORF
278	953	UCGUGCGCAGCAGCAAGCA	1250	UGCUGCUGCUGCGCACGA	Rh,D,M	[666-684] ORF
279	954	CGACGGCAAGCUGCCCGAG	1251	CUCGGGCAGCUUGCCGUCG	D	[772-790] ORF
280	955	UUCAUUGGGCGCCUGGUCC	1252	GGACCAGGCGCCCAAUGAA	Rh,D	[1433-1451] ORF
281	956	UUGACCAGGACAUCUACGG	1253	CCGUAGAUGUCCUGGUCAA	Rt	[1326-1344] ORF
282	957	CCUGGCCACACAAGCUCUCC	1254	GGAGAGCUUGUGGGCCAGG	Rh,D,P	[1006-1024] ORF
283	958	UGACUUCGUGCGCAGCAGC	1255	GCUGCUGCGCACGAAGUCA	Rh,Rt,M	[661-679] ORF
284	959	AUGAUGCACCGGACAGGCC	1256	GGCCUGUCCGGUGCAUCAU	Rh,Rb,Rt,M,P	[935-953] ORF
285	960	CACCGACGGCAAGCUGCCC	1257	GGGCAGCUUGCCGUCGGUG	D,Rt	[769-787] ORF
286	961	GACGGCAAGCUGCCCGAGG	1258	CCUCGGGCAGCUUGCCGUC	Rh,D	[773-791] ORF

No.	SEQ ID SEN	Sense siRNA	SEQ ID AS	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
287	962	UACCAGGCCAUGGCCAAGG	1259	CCUUGGCCAUGGCCUGGUA	Rh,D	[392-410] ORF
288	963	UCCGCGACAAGCGCAGCGC	1260	GCGCUGCGCUUGUCGCGGA	D	[717-735] ORF
289	964	UUCGCGACAAGCGCAGCG	1261	CGCUGCGCUUGUCGCGGAA	D	[716-734] ORF
290	965	AAGGACGUGGAGCGCACGG	1262	CCGUGCGCUCCACGUCCUU	Rh,D	[797-815] ORF
291	166	UUCCACCACAAGAUGGUGG	1263	CCACCAUCUUGUGGUGGAA	Rh,Rb,D,P	[869-887] ORF
292	967	UACGGGCGCGAGGAGCUGC	1264	GCAGCUCCUCGCGCCCGUA	D,M	[1340-1358] ORF
293	968	AAACACCUGGCUGGGCUGG	1265	CCAGCCCAGCCAGGUGUUU	D	[1184-1202] ORF
294	969	AACACCUGGCUGGGCUGGG	1266	CCCAGCCCAGCCAGGUGUU	D	[1185-1203] ORF
295	970	AUUGGGCGCCUGGUCCGGC	1267	GCCGGACCAGGCGCCCAU	Rh,D	[1436-1454] ORF
296	971	ACCGACGGCAAGCUGCCCG	1268	CGGGCAGCUUGCCGUCGGU	D	[770-788] ORF
297	972	UUCGUGCGCAGCAGCAAGC	1269	GCUUGCUGCUGCGCACGAA	Rh,D,M	[665-683] ORF

Table D: SERPINH1 Active 18+1-mer siRNAs

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human- 32454740 ORF:230- 1486
1	1270	AGCCUUUGUUGCUAUCAAA	1849	UUUGAUAGCAACAAAGGCU	Rh	[2117-2135] 3'UTR
2	1271	GCCUAAGGGUGACAAGAU	1850	UAUCUUGUCACCCUAGGC	Rh	[1453-1471] ORF
3	1272	GGCCUAAGGGUGACAAGAA	1851	UUCUUGUCACCCUAGGCC	Rh	[1452-1470] ORF
4	1273	CCUCAAUCAAGUAUUAUAA	1852	UUUUGAAUACUGAUUGAG G		[1774-1792] 3'UTR
5	1274	GGCGGAUUGAGAAGGAGCA	1853	UGCUCUUCUCAAUCCGCC		[1973-1991] 3'UTR
5	1275	GGCAGUGGAGAACAUCUA	1854	UAGGAUGUUCUCCACUGCC	Rh	[415-433] ORF
7	1276	GGGUCAGCCAGCCUCUUA	1855	UAAGAGGGCUGGCUGACCC	Rh	[1839-1857] 3'UTR
8	1277	GGGUGACAAGAUGCGAGAA	1856	UUCUCGCAUCUUGUCACCC	Rh,D	[1459-1477] ORF
9	1278	GGACCAGGCAGUGGAGAAA	1857	UUUCUCCACUGCCUGGUCC	Rh	[409-427] ORF
10	1279	GAGACACAUGGGUGCUAUA	1858	UAUAGCACCCAUGUGUCUC	Rh,D,Rt, M	[1533-1551] 3'UTR
11	1280	GUUGGAGCGUGGAAAAAAA	1859	UUUUUUUCCACGCUCCAAC		[2191-2208] 3'UTR
12	1281	GGAACAUGAGCCUUUGUUA	1860	UAACAAAGGCUCAUGUCC	Rh	[2109-2127] 3'UTR
13	1282	GCCAUGUUCUUAAGCCAA	1861	UUGGCUUGAAGAACAUGGC	Rh,Rb,D	[836-854] ORF
14	1283	GGAUUGAGAAGGAGCUCCA	1862	UGGAGCUCCUUCUCAUCC		[1976-1994] 3'UTR
15	1284	GGGAUGAACUUUUUGUUUA	1863	UAAACAAAAGUUCAUCCC	Rh	[2048-2066] 3'UTR
16	1285	GCCGAGUGAGGCGGAUUA	1864	UAAUCCGCCUCACUGCGGC		[1963-1981] 3'UTR
17	1286	GGACCUUCCAGCUAGAAA	1865	UUUCUAGCUGGGAAGGUCC	Rh	[1639-1657] 3'UTR
18	1287	GACCUUCCAGCUAGAAUA	1866	UAUUCUAGCUGGGAAGGUC	Rh	[1640-1658] 3'UTR
19	1288	CCUGUGAGACCAAAUUGAA	1867	UUCAAUUUGGUCUCACAGG	Rh	[1814-1832] 3'UTR
20	1289	UGGAGAACAUCUGGUGUA	1868	UACACCAGGAUGUUCUCCA	Rh	[420-438] ORF
21	1290	GCCUUUGUUGCUAUCAAUA	1869	UAUUGAUAGCAACAAAGGC	Rh	[2118-2136] 3'UTR
22	1291	CCGCCUUUGAGUUGGACAA	1870	UUGUCCAACUCAAGGCGG	Rh	[1293-1311] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
23	1292	CAGGCAGUGGAGAACAUCA	1871	UGAUGUUCUCCACUGCCUG	Rh	[413-431] ORF
24	1293	CACCUGUGAGACCAAAUUA	1872	UAAUUUGGUCUCACAGGUG	Rh	[1812-1830] 3'UTR
25	1294	GGGAAGAUGCAGAAGAAGA	1873	UCUUCUUCUGCAUCUCCCC	Rh,Rb,Rt	[1112-1130] ORF
26	1295	GGCCAUUGACAAGAACAAA	1874	UUUGUUCUUGUCA AUGGCC	Rh,D	[1213-1231] ORF
27	1296	3CCUUUGAGUUGGACACAA	1875	UUGUGUCCAACUCA AAGGC	Rh	[1295-1313] ORF
28	1297	AGCGGACCUUCCCAGCUAA	1876	UUAGCUGGGAAGGUCCGCU	Rh	[1636-1654] 3'UTR
29	1298	GAAGAAGGCUGUUGCCAUA	1877	UAUGGCAACAGCCUUCUUC	Rt	[1123-1141] ORF
30	1299	ACAAGAUGCAGACGAGUA	1878	UACUCGUCUCGCAUCUUGU	Rh,Rt	[1464-1482] ORF
31	1300	GAGGCGGAUUGAGAAGGAA	1879	UUCUUCUCAAUCCGCCUC		[1971-1989] 3'UTR
32	1301	GGACAACCGUGGCUUCAUA	1880	UAUGAAGCCACGGUUGUCC	Rh,M	[886-904] ORF
33	1302	CAUAUUUAUAGCCAGGUAA	1881	UUACCUGGCUAUAUAUAUG	Rh	[1788-1806] 3'UTR
34	1303	CGACGACGAGAAGGAAAAA	1882	UUUUUCCUUCUCGUCGUCG		[967-985] ORF
35	1304	CUCACCUGUGAGACCAAAA	1883	UUUUGGUCUCACAGGUGAG	Rh	[1810-1828] 3'UTR
36	1305	GCGGCUCCCUGCUAUUCA	1884	UUGAAUAGCAGGGAGCCGC		[1419-1437] ORF
37	1306	AGAACAUCCUGGUGUCACA	1885	UGUGACACCAGGAUGUUCU		[423-441] ORF
38	1307	CACACUGGGAUGAGAAUA	1886	UAUUUCUCAUCCCAGUGUG	Rh	[852-870] ORF
39	1308	GCUAGAAUUCACUCCACUA	1887	UAGUGGAGUGAAUUCUAGC	Rh	[1650-1668] 3'UTR
40	1309	CCUUAUCUUCUAGUGCA	1888	UGCACUAGGAAGAUGAAGG		[1389-1407] ORF
41	1310	UGCUAUCAAUCCAAGAACA	1889	UGUUCUUGGAUUGAUAGCA	Rh	[2126-2144] 3'UTR
42	1311	GGAAGAUGCAGAAGAAGGA	1890	UCCUUCUUCUGCAUCUCC	Rh,Rb,Rt	[1113-1131] ORF
43	1312	CAUGAGCCUUUGUUGCUA	1891	UUAGCAACAAGGCUCAUG	Rh	[2113-2131] 3'UTR
44	1313	GCGGAUUGAGAAGGAGCUA	1892	UAGCUCCUUCUCAUCCGC		[1974-1992] 3'UTR
45	1314	UGCAGUCCAUAACGAGUA	1893	UACUCGUUGAUGGACUGCA	Rh,Rt,M	[738-756] ORF
46	1315	GCACUGCGGAGAAGUUGAA	1894	UUCAACUUCUCCGCAGUGC		[321-339] ORF
47	1316	CCAGGCAGUGGAGAACAUA	1895	UAUGUUCUCCACUGCCUGG	Rh	[412-430] ORF
48	1317	3GCAAGAAGGACCUGUACA	1896	UGUACAGGUCCUUCUUGCC	Rh,D,M	[1253-1271] ORF
49	1318	CUCUACAACUACUACGACA	1897	UGUCGUAGUAGUUGAAG G	Rb	[953-971] ORF
50	1319	CUUCCAGCUAGAAUUCAA	1898	UUGAAUUCUAGCUGGGAAG	Rh	[1643-1661] 3'UTR
51	1320	AGGCGGAUUGAGAAGGAGA	1899	UCUCCUUCUCAAUCCGCCU		[1972-1990] 3'UTR
52	1321	GGUCCUAUACCGUGGGUGA	1900	UCACCCACGGUAUAGGACC	Rh	[912-930] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
53	1322	GCAAGAAGGACCGUACCA	1901	UGGUACAGGUCCUUCUUGC	Rh,D,M	[1254-1272] ORF
54	1323	CCGUGGGUGUCAUGAUGAA	1902	UUCAUCAUGACACCCACGG	Rh	[921-939] ORF
55	1324	GAUGCGAGACGAGUUUAUA	1903	UUUAACUCGUCUCGCAUC	Rh	[1468-1486] ORF
56	1325	3GCAGUGCUGAGCGCCGAA	1904	UUCGGCGCUCAGCACUGCC		[511-529] ORF
57	1326	CAGCUAGAAUUCACUCCAA	1905	UUGGAGUGAAUUCUAGCUG	Rh	[1648-1666] 3'UTR
58	1327	GAGCUUCGCUGAUGACUUA	1906	UAAGUCAUCAGCGAAGCUC	Rh	[649-667] ORF
59	1328	CUUUGAGUUGGACACAGAA	1907	UUCUGUGUCCAACUCAAAG	Rh	[1297-1315] ORF
60	1329	GGUGGACAACCGUGGCUUA	1908	UAAGCCACGGUUGUCCACC	Rh,M	[883-901] ORF
51	1330	GCCUCAUCAUCCUCAUGCA	1909	UGCAUGAGGAUGAUGAGGC	Rh,D,Rt, M	[1026-1044] ORF
62	1331	ACCAGGCAGUGGAGAACAA	1910	UUGUUCUCCACUGCCUGGU	Rh	[411-429] ORF
63	1332	CCUGCCUCAUUCAGUAUUA	1911	UAAUACUGAUUGAGGCAGG		[1770-1788] 3'UTR
64	1333	GAUCAAGCCUGCCUCAAUA	1912	UAUUGAGGCAGGCUUGAUC	Rh	[1763-1781] 3'UTR
65	1334	CAGACUCUGGUCAAGAAGA	1913	UCUUCUUGACCAGAGUCUG	Rh	[2011-2029] 3'UTR
66	1335	CGCGCUGCAGUCCAUA AAA	1914	UUUGAUGGACUGCAGCGCG	Rh,Rt	[733-751] ORF
67	1336	CUGGCACUGCGGAGAAGUA	1915	UACUUCUCCGAGUGCCAG		[318-336] ORF
68	1337	CCAGCUCUAUCCCAACCUA	1916	UAGGUUGGUAUAGACUG G		[1885-1903] 3'UTR
59	1338	AGGGUGUGGUGGAGGUGAA	1917	UUCACCUCCACACACCCU	Rh,D	[1152-1170] ORF
70	1339	AGUGAGGCGGAUUGAGAAA	1918	UUUCUCAUCCGCCUCACU		[1968-1986] 3'UTR
71	1340	CGGACAGGCCUCUACAACA	1919	UGUUGUAGAGGCCUGUCCG	Rh,Rb,Rt, P	[944-962] ORF
72	1341	CGACGAGAAGGAAAAGCUA	1920	UAGCUUUUCCUUCUCGUCG	Rh	[970-988] ORF
73	1342	AGGCCAAGGCAGUGCUGAA	1921	UUCAGCACUGCCUUGGCCU	Rh	[504-522] ORF
74	1343	GCCUCAGGGUGCACACAGA	1922	UCUGUGUGCACCCUGAGGC		[1488-1506] 3'UTR
75	1344	GGAUGAGAAUUCACCAA	1923	UUGGUGGAAUUCUCAUCC	Rh	[859-877] ORF
76	1345	AGAAGGAAAAGCUGCAAAA	1924	UUUUGCAGCUUUUCCUUCU	Rh	[975-993] ORF
77	1346	AGCUCUAUCCCAACCUCUA	1925	UAGAGGUUGGUAUAGAC U	Rh	[1887-1905] 3'UTR
78	1347	UGACAAGAUGCGAGACGAA	1926	UUCGUCUCGCAUCUUGUCA	Rh	[1462-1480] ORF
79	1348	AGAAGGAGCUCCAGGAGA	1927	UCUCCUGGGAGCUCCUUCU		[1982-2000] 3'UTR
80	1349	CCUUCUCACCUGUGAGACA	1928	UGUCUCACAGGUGAGAAGG	Rh	[1806-1824] 3'UTR
81	1350	GGCUUCUGGGCAGACUCUA	1929	UAGAGUCUGCCCAGAAGCC	Rh	[2001-2019] 3'UTR
82	1351	CCAGCCUCAUCAUCCUCAA	1930		Rh,D,Rt, M	[1023-1041]

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
				UUGAGGAUGAUGAGGCUG G		ORF
83	1352	CCAAAGGCUCCUGAGACAA	1931	UUGUCUCAGGAGCCUUUGG		[1521-1539] 3'UTR
84	1353	GGACCUGGGCCAUAGUCA	1932	UUGACUAUGGCCCAGGUCC		[1722-1740] 3'UTR
85	1354	GGGUGUCAUGAUGAUGCAA	1933	UUGCAUCAUCAUGACACCC	Rh	[925-943] ORF
86	1355	GUACCAGCCUUGGAUACUA	1934	UAGUAUCCAAGGCUGGUAC	Rh	[1572-1590] 3'UTR
87	1356	GGCUGUUGCCAUCUCCUUA	1935	UAAGGAGAUGGCAACAGCC		[1129-1147] ORF
88	1357	CGCAGUGAGGCGGAUUGAA	1936	UUCAAUCCGCCUCACUGCG		[1965-1983] 3'UTR
89	1358	CCAAGGACGUGGAGCGCAA	1937	UUGC GCUCCACGUCCUUGG	Rh,D	[795-813] ORF
90	1359	GGCUCCUGAGACACAUGGA	1938	UCCAUGUGUCUCAGGAGCC	D	[1526-1544] 3'UTR
91	1360	GCUGCAGUCCAUAACGAA	1939	UUCGUUGAUGGACUGCAGC	Rh,Rt	[736-754] ORF
92	1361	CCAGGUACCUUCUCACCUA	1940	UAGGUGAGAAGGUACCUGG	Rh	[1799-1817] 3'UTR
93	1362	GCAGCGCGCUGCAGUCCAA	1941	UUGGACUGCAGCGCGCUGC	Rh,Rt	[729-747] ORF
94	1363	GAGACCAAAUUGAGCUAGA	1942	UCUAGCUCAAUUUGGUCUC	Rh	[1819-1837] 3'UTR
95	1364	GCCGCCGAGGUGAAGAAAA	1943	UUUUCUUCACCUCGGCGGC		[281-299] ORF
96	1365	GCAGACUCUGGUCAAGAAA	1944	UUUCUUGACCAGAGUCUGC	Rh	[2010-2028] 3'UTR
97	1366	CUAGAAUUCACUCCACUUA	1945	UAAGUGGAGUGAAUUCUA G	Rh	[1651-1669] 3'UTR
98	1367	GCAGUGGAGAACAUCCUGA	1946	UCAGGAUGUUCUCCACUGC	Rh	[416-434] ORF
99	1368	CGCAUGUCAGGCAAGAAGA	1947	UCUUCUUGCCUGACAUGCG	Rh,D	[1244-1262] ORF
100	1369	CGGAUUGAGAAGGAGCUCA	1948	UGAGCUCCUUCUCAAUCCG		[1975-1993] 3'UTR
101	1370	AGGUGAGGUACCAGCCUUA	1949	UAAGGCUGGUACCUCACCU	Rh	[1565-1583] 3'UTR
102	1371	CCACACUGGGAUGAGAAAA	1950	UUUUCUCAUCCAGUGUGG	Rh	[851-869] ORF
103	1372	GCCAUUGACAAGAACAAGA	1951	UCUUGUUCUUGUCAAUUGC	Rh,D	[1214-1232] ORF
104	1373	GCGCUGCAGUCCAUAACA	1952	UGUUGAUGGACUGCAGCGC	Rh,Rt	[734-752] ORF
105	1374	CUCCCAACUAUAAAACUAA	1953	UUAGUUUUUAUAGUUGGA G	Rh	[1903-1921] 3'UTR
106	1375	GGUGACAAGAUGCGAGACA	1954	UGUCUCGCAUCUUGUCACC	Rh	[1460-1478] ORF
107	1376	GGCCGACUUGUCACGCAUA	1955	UAUGCGUGACAAGUCGGCC	Rh	[1231-1249] ORF
108	1377	CCUAAGGGUGACAAGAUGA	1956	UCAUCUUGUCACCCUUAGG	Rh	[1454-1472] ORF
109	1378	UGAGACACAUGGGUGCUAA	1957	UUAGCACCAUGUGUCUCA	Rh,D,Rt, M	[1532-1550] 3'UTR
110	1379	GGGUGGAAAAACAGACCGA	1958	UCGGUCUGUUUUUCCACCC		[1601-1619] 3'UTR
111	1380	GUGGAGGUGACCCAUGAA	1959	UUCAUGGGUCACCUCCACC	Rh,Rt,M	[1159-1177]

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
						ORF
112	1381	CUUUGACCAGGACAUCAA	1960	UUAGAUGUCCUGGUCAAAG	Rh,Rt	[1324-1342] ORF
113	1382	GAACAUGAGCCUUUGUUGA	1961	UCAACAAAGGCUCAUGUUC	Rh	[2110-2128] 3'UTR
114	1383	AGCCUUGGAUACUCCAUGA	1962	UCAUGGAGUAUCCAAGGCU	Rh	[1577-1595] 3'UTR
115	1384	GGAGGUGACCCAUGACCUA	1963	UAGGUCAUGGGUCACCUCC	Rh,Rt,M	[1162-1180] ORF
116	1385	AGAUAAGCCUGCCUCAA	1964	UUUGAGGCAGGCUUGAUCU	Rh	[1762-1780] 3'UTR
117	1386	GCCCAAGGGUGUGUGGAA	1965	UUCCACCACACCCUUGGGC	Rh,D	[1147-1165] ORF
118	1387	AGAACAAGGCCGACUUGUA	1966	UACAAGUCGGCCUUGUUCU	Rh	[1224-1242] ORF
119	1388	GUGGCUUCAUGGUGACUCA	1967	UGAGUCACCAUGAAGCCAC	Rh	[894-912] ORF
120	1389	CUCCUGAGACACAUGGGUA	1968	UACCCAUGUGUCUCAGGAG	D	[1528-1546] 3'UTR
121	1390	CAGCCUUGGAUACUCCAUA	1969	UAUGGAGUAUCCAAGGCUG	Rh	[1576-1594] 3'UTR
122	1391	AAGGCUCCUGAGACACUA	1970	UAUGUGUCUCAGGAGCCUU	D	[1524-1542] 3'UTR
123	1392	AGAAGAAGGCUGUUGCCAA	1971	UUGGCAACAGCCUUCUUCU	Rt	[1122-1140] ORF
124	1393	CUACUACGACGACGAGAAA	1972	UUUCUCGUCGUCGUAGUAG	Rb	[961-979] ORF
125	1394	CCUUUGUUGCUAUCAAUCA	1973	UGAUUGAUAGCAACAAAGG	Rh	[2119-2137] 3'UTR
126	1395	AGGCAGUGGAGAACAUCCA	1974	UGGAUGUUCUCCACUGCCU	Rh	[414-432] ORF
127	1396	CCAUCACGUGGAGCCUCUA	1975	UAGAGGCUCCACGUGAUGG	Rh	[1045-1063] ORF
128	1397	AGCUCUCCAGCCUCAUCAA	1976	UUGAUGAGGCUGGAGAGCU	Rh,D,Rt, M,P	[1017-1035] ORF
129	1398	GGCUCCCUGCUAUUCAUUA	1977	UAAUGAAUAGCAGGGAGCC	D	[1421-1439] ORF
130	1399	GGGAACAUGAGCCUUUGUA	1978	UACAAAGGCUCAUGUUCCC	Rh	[2108-2126] 3'UTR
131	1400	GGGCCAUAGUCAUUCUGCA	1979	UGCAGAAUGACUAUGGCCC		[1728-1746] 3'UTR
132	1401	CCAAAGAGCAGCUGAAGAA	1980	UUCUUCAGCUGCUCUUUGG	Rh,Rb,P	[1086-1104] ORF
133	1402	GACGAGAAGGAAAAGCUGA	1981	UCAGCUUUUCCUUCUCGUC	Rh	[971-989] ORF
134	1403	GGGCUUCUGGGCAGACUCA	1982	UGAGUCUGCCCAGAAGCCC	Rh	[2000-2018] 3'UTR
135	1404	CAAGGACCAGGCAGUGGAA	1983	UUCCACUGCCUGGUCCUUG	Rh	[406-424] ORF
136	1405	CUGUGAGACCAAUUGAGA	1984	UCUCAUUUUGGUCUCACAG	Rh	[1815-1833] 3'UTR
137	1406	GACUGAGGCCAUUGACAAA	1985	UUUGUCAUUGGCCUCAGUC	Rh	[1207-1225] ORF
138	1407	GACUUGUCACGCAUGUCAA	1986	UUGACAUGCGUGACAAGUC	Rh	[1235-1253] ORF
139	1408	GAGGUGAGGUACCAGCCUA	1987	UAGGCUGGUACCUCACCUC		[1564-1582] 3'UTR
140	1409	CAGAUACCAUGAUGCUGAA	1988	UUCAGCAUCAUGGUUUCUG	Rh	[1681-1699] 3'UTR
141	1410	AGGCAAGAAGGACCUGUAA	1989	UUACAGGUCCUUCUUGCCU	Rh,D	[1252-1270] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
142	1411	CUGGGAUGAGAAAUCCAA	1990	UUGGAAUUUCUCAUCCCAG	Rh	[856-874] ORF
143	1412	AGGUACCAGCCUUGGAUAA	1991	UUAUCCAAGGCUGGUACCU	Rh	[1570-1588] 3'UTR
144	1413	CAGCCAGCCCUCUUCUGAA	1992	UUCAGAAGAGGGCUGGCUG		[1843-1861] 3'UTR
145	1414	GUGUCAUGAUGAUGCACCA	1993	UGGUGCAUCAUCAUGACAC	Rh	[927-945] ORF
146	1415	CCUCUACAACUACUACGAA	1994	UUCGUAGUAGUUGUAGAGG	Rb,D	[952-970] ORF
147	1416	CCGCCGAGGUGAAGAAACA	1995	UGUUUCUUCACCUCGGCGG	Rh	[282-300] ORF
148	1417	GCUAUCAAUCCAAGAACUA	1996	UAGUUCUUGGAUUGAUAGC	Rh	[2127-2145] 3'UTR
149	1418	AGCCUGCCUCAUUCAGUAA	1997	UUACUGAUUAGAGGCAGGCU		[1768-1786] 3'UTR
150	1419	GUUCCGGCCUAAGGGUGAA	1998	UUCACCCUUAAGCCGGACC	Rh	[1447-1465] ORF
151	1420	3AAGGAAAAGCUGCAAAUA	1999	UAUUUGCAGCUUUUCCUUC	Rh	[976-994] ORF
152	1421	GCCUCUACAACUACUACA	2000	UGUAGUAGUUGUAGAGGCC	Rb,D	[950-968] ORF
153	1422	UGUUCUUAAGCCACACUA	2001	UAGUGUGGCUUGAAGAACA	Rh,Rb,D	[840-858] ORF
154	1423	GGCCAAGGCAGUGCUGAGA	2002	UCUCAGCACUGCCUUGGCC	Rh	[505-523] ORF
155	1424	AGAAAUUCCACCACAAGAA	2003	UUCUUGUGGUGGAAUUUCU	Rh	[864-882] ORF
156	1425	CUGCAGUCCAUCAACGAGA	2004	UCUCGUUGAUGGACUGCAG	Rh,Rt,M	[737-755] ORF
157	1426	CCAGCGUGUCCACGCCAA	2005	UUGGCGUGGAACACGCUGG		[1275-1293] ORF
158	1427	GCUCCCUCCUGCUUCUCAA	2006	UUGAGAAGCAGGAGGGAGC		[234-252] ORF
159	1428	CCGGACAGGCCUCUACAAA	2007	UUUGUAGAGGCCUGUCCGG	Rh,Rb,Rt, P	[943-961] ORF
160	1429	CCCAUCACGUGGAGCCUCA	2008	UGAGGCUCCACGUGAUGGG	Rh	[1044-1062] ORF
161	1430	CCGGCCUAAGGGUGACAAA	2009	UUUGUCACCCUUAAGCCGG	Rh	[1450-1468] ORF
162	1431	CCUAUACCGUGGGUGUCA	2010	UUGACACCCACGGUAUAGG	Rh,D,P	[915-933] ORF
163	1432	CAGUGGAGAACAUCUGGA	2011	UCCAGGAUGUUCUCCACUG	Rh	[417-435] ORF
164	1433	CACUGGGAUGAGAAAUUCA	2012	UGAAUUUCUCAUCCCAGUG	Rh	[854-872] ORF
165	1434	AUCCAAAGGCUCCUGAGAA	2013	UUCUCAGGAGCCUUGGAU		[1519-1537] 3'UTR
166	1435	UGAGAAAUUCCACCACAAA	2014	UUUGUGGUGGAAUUUCUCA	Rh	[862-880] ORF
167	1436	GGUGGAAAAACAGACCGGA	2015	UCCGGUCUGUUUUUCCACC		[1602-1620] 3'UTR
168	1437	GCUGGGCAGCCGACUGUAA	2016	UUACAGUCGGCUGCCCAGC		[616-634] ORF
169	1438	CCAUAUGUCAUUCUGCCUGA	2017	UCAGGCAGAAUGACUAUGG		[1731-1749] 3'UTR
170	1439	GCACCGGACAGGCCUCUAA	2018	UUAGAGGCCUGUCCGGUGC	Rh,Rb,Rt, P	[940-958] ORF
171	1440	GUUGGACACAGAUGGCAAA	2019	UUUGCCAUCUGUGUCCAAC		[1303-1321] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
172	1441	GCCUGCCUCAAU CAGUAUA	2020	UAUACUGAUUGAGGCAGGC		[1769-1787] 3'UTR
173	1442	GAUCAACUCCGCGACAAA	2021	UUUGUCGCGGAAGUUGAUC	D	[709-727] ORF
174	1443	GGCCGCAGUGAGGCGGAUA	2022	UAUCCGCCUCACUGCGGCC		[1962-1980] 3'UTR
175	1444	CUGCGGAGAAGUUGAGCCA	2023	UGGCUCAACUUCUCCGCAG		[324-342] ORF
176	1445	GCAUCCAAAGGCUCCUGAA	2024	UUCAGGAGCCUUUGGAUGC		[1517-1535] 3'UTR
177	1446	GCUUCUGGGCAGACUCUGA	2025	UCAGAGUCUGCCAGAAGC	Rh	[2002-2020] 3'UTR
178	1447	CCAGCCCUCUUCUGACACA	2026	UGUGUCAGAAGAGGGCUGG		[1846-1864] 3'UTR
179	1448	GCUCUAUCCCAACCUCUCA	2027	UGAGAGGUUGGGAUAGAG C	Rh	[1888-1906] 3'UTR
180	1449	GGACGUGGAGCGCACGGAA	2028	UUCCGUGCGCUCCACGUCC	Rh,D	[799-817] ORF
181	1450	CCAAGGCAGUGCUGAGCGA	2029	UCGCUCAGCACUGCCUUGG	Rh	[507-525] ORF
182	1451	3CAGAAGAAGGCUUUGCA	2030	UGCAACAGCCUUCUUCUGC	Rt	[1120-1138] ORF
183	1452	GACAUUUUGUUGGAGCGUA	2031	UACGCUCCAACAAAUGUC		[2183-2201] 3'UTR
184	1453	CGAGCACUCCAAGAUCAAA	2032	UUUGAUCUUGGAGUGCUCG	Rh,D	[697-715] ORF
185	1454	UCAUGAUGAUGCACCGGAA	2033	UUCCGGUGCAUCAUCAUGA	Rh	[930-948] ORF
186	1455	CCUGCUUCUCAGCGCCUUA	2034	UAAGGCGCUGAGAAGCAGG		[241-259] ORF
187	1456	CCCAACCUCUCCCAACUAA	2035	UUAGUUGGGAGAGGUUGG G	Rh	[1895-1913] 3'UTR
188	1457	UGGGCAGACUCUGGUCAAA	2036	UUUGACCAGAGUCUGCCCA	Rh	[2007-2025] 3'UTR
189	1458	CUCUGGUCAAGAAGCAUCA	2037	UGAUGCUUCUUGACCAGAG	Rh	[2015-2033] 3'UTR
190	1459	3AGCCUCUCGAGCGCCUUA	2038	UAAGGCGCUCGAGAGGCUC	ORF	[1055-1073] ORF
191	1460	AGAAGGCUGUUGCCAUCUA	2039	UAGAUGGCAACAGCCUUCU	Rt	[1125-1143] ORF
192	1461	CCCUGCUAGUCAACGCCAA	2040	UUGGCGUUGACUAGCAGGG	Rh	[822-840] ORF
193	1462	GCCUUCAGCUUGUACCAGA	2041	UCUGGUACAAGCUGAAGGC		[380-398] ORF
194	1463	GCUGCUAACCAAAGAGCAA	2042	UUGCUCUUUGGUUAGCAGC		[1078-1096] ORF
195	1464	CCCACAAGCUCUCCAGCCA	2043	UGGCUGGAGAGCUUGUGGG	Rh,D,P	[1011-1029] ORF
196	1465	GCUCCUGCUAUUCAUUGA	2044	UCAAUAGAAUAGCAGGGAGC	D	[1422-1440] ORF
197	1466	GUUCUUCAAAGAUAGGGAA	1045	UUCCCUAUCUUUGAAGAAC		[2083-2101] 3'UTR
198	1467	GUCAGCCAGCCCUCUUCUA	2046	UAGAAGAGGGCUGGCUGAC	Rh	[1841-1859] 3'UTR
199	1468	GCGGGACACCCAAAGCGGA	2047	UCCGCUUUGGGUGUCCCGC		[1405-1423] ORF
200	1469	AGCGCAGCGCGCUGCAGUA	2048	UACUGCAGCGCGCUGCGCU	Rh,Rt	[726-744] ORF
201	1470	CCGGAACUCCACAUCCUA	2049	UAGGAUGUGGAGUUUCCGG		[1701-1719]

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
						3'UTR
202	1471	CCAUUGACAAGAACAAGGA	2050	UCCUUGUUCUUGUCA AUGG	Rh,D	[1215-1233] ORF
203	1472	GACAUCUACGGGCGCGAA	2051	UUCGCGCCCGUAGAUGUCC	D	[1333-1351] ORF
204	1473	GACACAUGGGUGCUAUUGA	2052	UCAAUAGCACCCAUGUGUC	Rh,Rt,M	[1535-1553] 3'UTR
205	1474	CCUGGCACUGCGGAGAAGA	2053	UCUUCUCCGCAGUGCCAGG		[317-335] ORF
206	1475	GGGCCUGACUGAGGCCAUA	2054	UAUGGCCUCAGUCAGGCC	Rt	[1201-1219] ORF
207	1476	ACACUGGGAUGAGAAAUA	2055	UAAUUUCUCAUCCAGUGU	Rh	[853-871] ORF
208	1477	GGUCAGCCAGCCCUCUUA	2056	UGAAGAGGGCUGGCUGACC	Rh	[1840-1858] 3'UTR
209	1478	3UGAGGCGGAUUGAGAAGA	2057	UCUUCUCAUCCGCCUCAC		[1969-1987] 3'UTR
210	1479	UCACCUGUGAGACCAAUA	2058	UAUUUGGUCUCACAGGUGA	Rh	[1811-1829] 3'UTR
211	1480	AGCUGCAAUCGUGGAGAA	2059	UUCUCCACGAUUUGCAGCU	Rh	[984-1002] ORF
212	1481	GGUGCACACAGGAUGGCAA	2060	UUGCCAUCUGUGUGCACC	Rh	[1495-1513] 3'UTR
213	1482	GGGUGUGGUGGAGGUGACA	2061	UGUCACCUCCACCACACCC	Rh,D	[1153-1171] ORF
214	1483	CCAGCCUUGGAUACUCCAA	2062	UUGGAGUAUCCAAGGCUGG	Rh	[1575-1593] 3'UTR
215	1484	CCACAAGCUCUCCAGCCUA	2063	UAGGCUGGAGAGCUUGUGG	Rh,D,P	[1012-1030] ORF
216	1485	AAAGGCUCCUGAGACACAA	2064	UUGUGUCUCAGGAGCCUUU		[1523-1541] 3'UTR
217	1486	AGGAAAAGCUGCAAAUCGA	2065	UCGAUUUGCAGCUUUUCCU	Rh	[978-996] ORF
218	1487	CGCAGCAGCUCCUGGCACA	2066	UGUGCCAGGAGCUGCUGCG	ORF	[307-325] ORF
219	1488	GGUGUCAUGAUGAUGCACA	2067	UGUGCAUCAUCAUGACACC	Rh	[926-944] ORF
220	1489	CCUCUUCUGACACUAAAAA	2068	UUUUUAGUGUCAGAAGAG G		[1851-1869] 3'UTR
221	1490	AGCUAGAAUUCACUCCACA	2069	UGUGGAGUGAAUUCUAGCU	Rh	[1649-1667] 3'UTR
222	1491	CGCUGGGCGGCAAGGCGAA	2070	UUCGCCUUGCCGCCAGCG		[474-492] ORF
223	1492	GGCCUGGCCUUCAGCUUGA	2071	UCAAGCUGAAGGCCAGGCC		[374-392] ORF
224	1493	AGACACAUGGGUGCUAUUA	2072	UAAUAGCACCCAUGUGUCU	Rh,Rt,M	[1534-1552] 3'UTR
225	1494	CGUGGGUGUCAUGAUGAUA	2073	UAUCAUCAUGACACCCACG	Rh	[922-940] ORF
226	1495	GUGGGUGUCAUGAUGAUGA	2074	UCAUCAUCAUGACACCCAC	Rh	[923-941] ORF
227	1496	GAGAAGGAGCUCCAGGAA	2075	UUCUGGGAGCUCCUUCUC		[1981-1999] 3'UTR
228	1497	GACUCUGGUCAAGAAGCAA	2076	UUGCUUCUUGACCAGAGUC	Rh	[2013-2031] 3'UTR
229	1498	CACUAAAACACCUCAGCUA	2077	UAGCUGAGGUGUUUAGU G		[1861-1879] 3'UTR
230	1499	GGAGGCAUCCAAAGGCUCA	2078	UGAGCCUUUGGAUGCCUCC		[1513-1531] 3'UTR

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
231	1500	GACCCAGCUCAGUGAGCUA	2079	UAGCUCACUGAGCUCCCUC		[636-654] ORF
232	1501	CAUGACCUGCAGAAACAA	2080	UUGUUUCUGCAGGUCAUGG	Rh,Rt,M	[1171-1189] ORF
233	1502	AGAUGCAGAAGAAGGCUGA	2081	UCAGCCUUCUUCUGCAUCU	Rh,Rt,M	[1116-1134] ORF
234	1503	CAGCAAGCAGCACUACAAA	2082	UUUGUAGUGCUGCUUGCUG	Rh,D	[676-694] ORF
235	1504	CAAGCUCUCCAGCCUCAUA	2083	UAUGAGGCUGGAGAGCUUG	Rh,D,M,P	[1015-1033] ORF
236	1505	UGCAGAAGAAGGCUGUUGA	2084	UCAACAGCCUUCUUCUGCA	Rt	[1119-1137] ORF
237	1506	GGCGCGAGGAGCUGCGCAA	2085	UUGCGCAGCUCCUCGCGCC	Rh,D,M	[1344-1362] ORF
238	1507	GGUACCAGCCUUGGAUACA	2086	UGUAUCCAAGGCUGGUACC	Rh	[1571-1589] 3'UTR
239	1508	GCAGCCGACUGUACGGACA	2087	UGUCCGUACAGUCGGCUGC		[621-639] ORF
240	1509	CAGCCUCAUCAUCCUCAUA	2088	UAUGAGGAUGAUGAGGCU G	Rh,D,Rt, M	[1024-1042] ORF
241	1510	GCCACCGCCUUUGAGUUGA	2089	UCAACUCAAGGCGGUGGC	Rh	[1289-1307] ORF
242	1511	AGAAGGACCUGUACCUGGA	2090	UCCAGGUACAGGUCCUUCU	Rh,D	[1257-1275] ORF
243	1512	GGUGAAGAAACCGCAGCA	2091	UGCUGCAGGUUUCUUCACC	Rh	[289-307] ORF
244	1513	GUACCUUCUCACCUGUGAA	2092	UUCACAGGUGAGAAGGUAC	Rh	[1803-1821] 3'UTR
245	1514	GGCCAAGGACCAGGCAGUA	2093	UACUGCCUGGUCCUUGGCC	Rh	[403-421] ORF
246	1515	GGCGGCAAGGCGACCACGA	2094	UCGUGGUCGCCUUGCCGCC		[479-497] ORF
247	1516	AGCACUCCAAGAUAACUA	2095	UAGUUGAUCUUGGAGUGCU	Rh,D	[699-717] ORF
248	1517	AUAUUUAUAGCCAGGUACA	2096	UGUACCUGGCUAUAAAUAU	Rh	[1789-1807] 3'UTR
249	1518	GGCAGCCGACUGUACGGAA	2097	UUCCGUACAGUCGGCUGCC		[620-638] ORF
250	1519	GUCACGCAUGUCAGGCAAA	2098	UUUGCCUGACAUGCGUGAC	Rh,D	[1240-1258] ORF
251	1520	GACAGGCCUCUACAACUAA	2099	UUAGUUGUAGAGGCCUGUC	Rh,Rb,Rt, P	[946-964] ORF
252	1521	GAUGCAGAAGAAGGCUGUA	2100	UACAGCCUUCUUCUGCAUC	Rh,Rt,M	[1117-1135] ORF
253	1522	ACCCAUGACCUGCAGAAAA	2101	UUUUCUGCAGGUCAUGGGU	Rh,Rt,M	[1169-1187] ORF
254	1523	GGCUUCAUGGUGACUCGGA	2102	UCCGAGUCACCAUGAAGCC	Rh	[896-914] ORF
255	1524	UGCCUCAAUCAGUUUCAA	2103	UUGAAUACUGAUUGAGGCA		[1772-1790] 3'UTR
256	1525	GUUCUUAAGCCACACUGA	2104	UCAGUGUGGCUUGAAGAAC	Rh,Rb,D	[841-859] ORF
257	1526	ACUCCAAGAUAACUCCA	2105	UUGAAGUUGAUCUUGGAG U	Rh,D,Rt, M	[702-720] ORF
258	1527	GCUGUUCUACGCCGACCAA	2106	UUGGUCGGCGUAGAACAGC	Rh	[1369-1387] ORF
259	1528	UAGUCAACGCCAUGUUCUA	2107	UAGAACAUGGCGUUGACUA	Rh	[828-846] ORF
260	1529	CCGUGUGCCUGAGCGGACA	2108	UGUCCGCUCAGGCACACGG	Rh	[1625-1643] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
						3'UTR
261	1530	AGGCCUCUACAACUACUAA	2109	UUAGUAGUUGUAGAGGCCU	Rh,Rb,D, Rt,P	[949-967] ORF
262	1531	GCUUCAUGGUGACUCGGUA	2110	UACCGAGUCACCAUGAAGC	Rh	[897-915] ORF
263	1532	GGUCAAGAAGCAUCGUGUA	2111	UACACGAUGCUUCUUGACC	Rh	[2019-2037] 3'UTR
264	1533	CUGCGAGCACUCCAAGUA	2112	UAUCUUGGAGUGCUCGCAG	Rh,D	[694-712] ORF
265	1534	GUCCUAUACCGUGGGUGUA	2113	UACACCCACGGUAUAGGAC	Rh	[913-931] ORF
266	1535	GGCCUGACUGAGGCCAUUA	2114	UAAUGGCCUCAGUCAGGCC	Rh	[1202-1220] ORF
267	1536	CACUCCAAGAUCAACUUA	2115	UGAAGUUGAUUCUUGGAGU G	Rh,D,Rt, M	[701-719] ORF
268	1537	GCGUCGCAGGCCAAGGCAA	2116	UUGCCUUGGCCUGCGACGC		[497-515] ORF
269	1538	AAGGGUGACAAGAUGCGAA	2117	UUCGCAUCUUGUCACCCUU	Rh,D	[1457-1475] ORF
270	1539	CAAGCUGUUCUACGCCGAA	2118	UUCGGCGUAGAACAGCUUG	Rh	[1366-1384] ORF
271	1540	CCUGCUAGUCAACGCCAUA	2119	UAUGGCGUUGACUAGCAGG	Rh	[823-841] ORF
272	1541	CCAAGGGUGUGGUGGAGGA	2120	UCCUCCACCACACCCUUGG	Rh,D	[1149-1167] ORF
273	1542	CACACAGGAUGGCAGGAGA	2121	UCUCCUGCCAUCCUGUGUG	Rh	[1499-1517] 3'UTR
274	1543	UCCUGAGACACAUGGGUGA	2122	UCACCCAUGUGUCUCAGGA	D,Rt,M	[1529-1547] 3'UTR
275	1544	CUACAACUACUACGACGAA	2123	UUCGUCGUAGUAGUUGUAG	Rb	[955-973] ORF
276	1545	GACAAGAUGCGAGACGAGA	2124	UCUCGUCUCGCAUCUUGUC	Rh,Rt	[1463-1481] ORF
277	1546	CCUGGAAGCUGGGCAGCCA	2125	UGGCUGCCCAGCUUCCAGG		[609-627] ORF
278	1547	CUUCAAGCCACACUGGGAA	2126	UUCCCAGUGUGGCUUGAAG	Rh,Rb,D	[844-862] ORF
279	1548	GCGAGACGAGUUUAUAGGGA	2127	UCCCUAUAAACUCGUCUCGC	Rh	[1471-1489] ORF+3'UTR
280	1549	3AAGCUGGGCAGCCGACUA	2128	UAGUCGGCUGCCCAGCUUC		[613-631] ORF
281	1550	GUGCCUGAGCGGACCUUCA	2129	UGAAGGUCCGCUCAGGCAC	Rh	[1629-1647] 3'UTR
282	1551	GGUGACCCAUGACCUGCAA	2130	UUGCAGGUCAUGGGUCACC	Rh,Rt,M	[1165-1183] RF
283	1552	AUGAGCCUUUGUUGCUAUA	2131	UAUAGCAACAAAGGCUCAU	Rh	[2114-2132] 3'UTR
284	1553	CAACUACUACGACGACGAA	2132	UUCGUCGUCGUAGUAGUUG	Rb	[958-976] ORF
285	1554	GCUGCGCUCACUCAGCAAA	2133	UUUGCUGAGUGAGCGCAGC	Rh	[571-589] ORF
286	1555	GAGAACAUCCUGGUGUCAA	2134	UUGACACCAGGAUGUUCUC		[422-440] ORF
287	1556	CCCAAGCUGUUCUACGCCA	2135	UGGCGUAGAACAGCUUGGG	Rh	[1364-1382] ORF
288	1557	CAGCUCUAUCCCAACCUCA	2136	UGAGGUUGGGAUAGAGCU G		[1886-1904] 3'UTR
289	1558	UGAGCUUCGCUGAUGACUA	2137	UAGUCAUCAGCGAAGCUCA	Rh	[648-666] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
290	1559	CCCAAGGCGGCCACGCUUA	2138	UAAGCGUGGCCGCCUUGGG	Rh	[341-359] ORF
291	1560	CUAUACCGUGGGUGUCAUA	2139	UAUGACACCCACGGUAUAG	Rh	[916-934] ORF
292	1561	CAUUGACAAGAACAAGGCA	2140	UGCCUUGUUCUUGUCA AUG	Rh,D	[1216-1234] ORF
293	1562	GGACCCAGCUCAGUGAGCA	2141	UGCUCACUGAGCUGGGUCC	ORF	[635-653] ORF
294	1563	GACGACGAGAAGGAAAAGA	2142	UCUUUUCCUUCUCGUCGUC	Rh	[968-986] ORF
295	1564	GCGGCAAGGCGACCACGGA	2143	UCCGUGGUCGCCUUGCCGC		[480-498] ORF
296	1565	GGGACACCCAAAGCGGCUA	2144	UAGCCGCUUUGGGUGUCCC		[1407-1425] ORF
297	1566	GGGAGGUGAGGUACCAGCA	2145	UGCUGGUACCUCACCUC		[1562-1580] 3'UTR
298	1567	GCAGCACUACAACUGCGAA	2146	UUCGCAGUUGUAGUGCUGC	Rh,D	[682-700] ORF
299	1568	GCGCAACGUGACCUGGAAA	2147	UUUCCAGGUCACGUUGCGC	M	[598-616] ORF
300	1569	GGGCUGGGCCUGACUGAGA	2148	UCUCAGUCAGGCCAGCCC		[1196-1214] ORF
301	1570	CCUGAGCGGACCUUCCCAA	2149	UUGGGAAGGUCCGCUCAGG	Rh	[1632-1650] 3'UTR
302	1571	GCAGCUGAAGAUCUGGAUA	2150	UAUCCAGAUUCUACGUCG	Rh,D	[1093-1111] ORF
303	1572	AGUGGAGAACAUCUGGUA	2151	UACCAGGAUGUUCUCCACU	Rh	[418-436] ORF
304	1573	GCAAGCAGCACUACAACUA	2152	UAGUUGUAGUGCUGCUUGC	Rh,D	[678-696] ORF
305	1574	AGCUCAGUGAGCUUCGCUA	2153	UAGCGAAGCUCACUGAGCU		[641-659] ORF
306	1575	CCGACUUGUCACGCAUGUA	2154	UACAUGCGUGACAAGUCGG	Rh	[1233-1251] ORF
307	1576	CCGAGGUCACCAAGGACGA	2155	UCGUCCUUGGUGACCUCGG	Rh,D	[786-804] ORF
308	1577	GGAGCCUCUCGAGCGCCUA	2156	UAGGCGCUCGAGAGGCUC		[1054-1072] ORF
309	1578	GGCCGCGCAGACCACCGAA	2157	UUCGGUGGUCUGCGCGGCC		[757-775] ORF
310	1579	GGAAACUCCACAUCUGUA	2158	UACAGGAUGUGGAGUUUCC	Rh	[1703-1721] 3'UTR
311	1580	CAAAGCGGCUCCUGCUAA	2159	UUAGCAGGGAGCCGCUUUG		[1415-1433] ORF
312	1581	GCUCCUGAGACACAUGGGA	2160	UCCCAUGUGUCUCAGGAGC	D	[1527-1545] 3'UTR
313	1582	CCUGGGCCAUAGUCAUUA	2161	UGAAUGACUAUGGCCAGG		[1725-1743] 3'UTR
314	1583	CGUGGAGCCUCUCGAGCGA	2162	UCGCUCGAGAGGCUCCACG		[1051-1069] ORF
315	1584	CCUCCUGCUUCUCAOCGCA	2163	UGCGCUGAGAAGCAGGAGG		[238-256] ORF
316	1585	AGUCCCAGAUCAAGCCUGA	2164	UCAGGCUUGAUCUGGGACU	Rh	[1756-1774] 3'UTR
317	1586	UACCGUGGGUGUCAUGAUA	2165	UAUCAUGACACCCACGGUA	Rh	[919-937] ORF
318	1587	GCCAGCCCUCUUCUGACAA	2166	UUGUCAGAAGAGGGCUGGC		[1845-1863] 3'UTR
319	1588	CCGAGGUGAAGAAACCUGA	2167	UCAGGUUUUCUACCCUCGG	Rh,Rt	[285-303] ORF
320	1589	UCCUGGCACUGCGGAGAAA	2168	UUUCUCCGCAGUGCCAGGA		[316-334] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
321	1590	CCCGGAAACUCCACAUCCA	2169	UGGAUGUGGAGUUUCCGGG		[1700-1718] 3'UTR
322	1591	ACUCUGGUCAAGAAGCAUA	2170	UAUGCUUCUUGACCAGAGU	Rh	[2014-2032] 3'UTR
323	1592	CCCAGAUACCAUGAUGCUA	2171	UAGCAUCAUGGUAUCUGGG	Rh	[1679-1697] 3'UTR
324	1593	CCUGAGACACAUGGGUGCA	2172	UGCACCCAUGUGUCUCAGG	D,Rt,M	[1530-1548] 3'UTR
325	1594	GCACUACAACUGCGAGCAA	2173	UUGCUCGCAGUUGUAGUGC	Rh,D	[685-703] ORF
326	1595	CCACAAGAUGGUGGACAAA	2174	UUUGUCCACCAUCUUGUGG	Rh,Rb,M, P	[874-892] ORF
327	1596	GGACACAGAUGGCAACCCA	2175	UGGGUUGCCAUCUGUGUCC		[1306-1324] ORF
328	1597	GAAAAGCUGCUAACCAAAA	2176	UUUUGGUUAGCAGCUUUUC		[1073-1091] ORF
329	1598	ACUACAACUGCGAGCACUA	2177	UAGUGCUCGCAGUUGUAGU	Rh,D	[687-705] ORF
330	1599	GCACUCCAAGAUAACUUA	2178	UAAGUUGAUCUUGGAGUGC	Rh,D	[700-718] ORF
331	1600	GCCUUGAAAAGCUGCUAAA	2179	UUUAGCAGCUUUUCAAGGC		[1068-1086] ORF
332	1601	GUGACUCGGUCCUAUACCA	2180	UGGUUAUAGGACCGAGUCAC	Rh	[905-923] ORF
333	1602	UGGUGGAGGUGACCCAUA	2181	UAUGGGUCACCUCACCAC	Rh,Rb,Rt, M	[1157-1175] ORF
334	1603	AUGCGAGACGAGUUUAAGA	2182	UCUAUAACUCGUCUCGCAU	Rh	[1469-1487] - ORF+3'UTR
335	1604	ACCUUCCAGCUAGAAUUA	2183	UAAUUCUAGCUGGGAAGGU	Rh	[1641-1659] 3'UTR
336	1605	CCCAGCUAGAAUUCACUCA	2184	UGAGUGAAUUCUAGCUGGG	Rh	[1646-1664] 3'UTR
337	1606	GUCACCAAGGACGUGGAA	2185	UUCCACGUCCUUGGUGACC	Rh,D	[790-808] ORF
338	1607	GGCCUCAGGGUGCACACAA	2186	UUGUGUGCACCCUGAGGCC		[1487-1505] 3'UTR
339	1608	UGAGGUACCAGCCUUGGAA	2187	UUCCAAGGCUGGUACCUCU	Rh	[1568-1586] 3'UTR
340	1609	CAUGGUGACUCGGUCCUAA	2188	UUAGGACCGAGUCACCAUG	Rh	[901-919] ORF
341	1610	GGAGGUACCAGCCUUGA	2189	UCAAGGCUGGUACCUCACC	Rh	[1566-1584] 3'UTR
342	1611	GCCGAGGUGAAGAAACCUA	2190	UAGGUUUCUUCACCUCGGC	Rh,Rt	[284-302] ORF
343	1612	GUACGGACCCAGCUCAGUA	2191	UACUGAGCUGGGUCCGUAC		[631-649] ORF
344	1613	CAAGAAGGACCUGUACCUA	2192	UAGGUACAGGUCCUUCUUG	Rh,D,M	[1255-1273] ORF
345	1614	GAGCACUCCAAGAUAACA	2193	UGUUGAUCUUGGAGUGCUC	Rh,D	[698-716] ORF
346	1615	CAUGUUCUUAAGCCACAA	2194	UUGUGGCUUGAAGAACAUG	Rh,Rb,D	[838-856] ORF
347	1616	CCCUCUGCUUCUCAGCGA	2195	UCGCUGAGAAGCAGGAGGG		[237-255] ORF
348	1617	AUGUCAGGCAAGAAGGACA	2196	UGUCCUUCUUGCCUGACAU	Rh,D	[1247-1265] ORF
349	1618	CAAGAUAACUCCGCGAA	2197	UUCGCGGAAGUUGAUCUUG	D	[706-724] ORF
350	1619	GCGUGUCCACGCCACCGA	2198	UCGGUGGCGUGGAACACGC		[1278-1296] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
351	1620	CGGACCCAGCUCAGUGAGA	2199	UCUCACUGAGCUGGGUCCG		[634-652] ORF
352	1621	CCUUCAGCUUGUACCAGGA	2200	UCCUGGUACAAGCUGAAGG		[381-399] ORF
353	1622	GCUCUCCAGCCUCAUCAUA	2201	UAUGAUGAGGCUGGAGAGC	Rh,D,Rt, M,P	[1018-1036] ORF
354	1623	CCCUGGCCACACAAGCUCUA	2202	UAGAGCUUGUGGGCCAGGG	Rh,D,P	[1005-1023] ORF
355	1624	GCCCGAGGUCACCAAGGAA	2203	UUCUUGGUGACCUCGGGC	Rh,D	[784-802] ORF
356	1625	GUGGAGAACAUCUGGUGA	2204	UCACCAGGAUGUUCUCCAC	Rh	[419-437] ORF
357	1626	3CUCACUCAGCAACUCCAA	2205	UUGGAGUUGCUGAGUGAGC	Rh	[576-594] ORF
358	1627	ACGCCAUGUUCUUAAGCA	2206	UGCUUGAAGAACAUGGCGU	Rh,Rb,P	[834-852] ORF
359	1628	ACACAUGGGUGCUAUUGGA	2207	UCCAUAAGCACCCAUGUGU	Rh	[1536-1554] 3'UTR
360	1629	CCAGCUCAGUGAGCUUCGA	2208	UCGAAGCUCACUGAGCUGG		[639-657] ORF
361	1630	CCCAGCUCAGUGAGCUUCA	2209	UGAAGCUCACUGAGCUGGG		[638-656] ORF
362	1631	GGGCGGCAAGGCGACCACA	2210	UGUGGUCGCCUUGCCGCCC		[478-496] ORF
363	1632	CAGGGUGCACACAGGAUGA	2211	UCAUCCUGUGUGCACCCUG		[1492-1510] 3'UTR
364	1633	AGGUGAAGAAACCUGCAGA	2212	UCUGCAGGUUUCUUCACCU	Rh	[288-306] ORF
365	1634	CCUCUCCCAACUAUAAAAA	2213	UUUUUAUAGUUGGAGAG G I	Rh	[1900-1918] 3'UTR
366	1635	GACUGUACGGACCCAGCUA	2214	UAGCUGGGUCCGUACAGUC		[627-645] ORF
367	1636	3AAGGAGCUCCAGGAGGA	2215	UCCUCCUGGGAGCUCCUUC		[1983-2001] 3'UTR
368	1637	ACGCAUGUCAGGCAAGAAA	2216	UUUCUUGCCUGACAUGCGU	Rh,D	[1243-1261] ORF
369	1638	GACUCGGUCCUAUACCGUA	2217	UACGGUAUAGGACCGAGUC	Rh	[907-925] ORF
370	1639	CACUACAACUGCGAGCACA	2218	UGUGCUCGCAGUUGUAGUG	Rh,D	[686-704] ORF
371	1640	AGCUCCUGGCACUGCGGAA	2219	UUCGCGAGUGCCAGGAGCU		[313-331] ORF
372	1641	CUAAGGGUGACAAGAUGCA	2220	UGCAUCUUGUCACCCUAG	Rh	[1455-1473] ORF
373	1642	UGUGAGACCAAAUUGAGCA	2221	UGCUGAAUUUGGUCUCACA	Rh	[1816-1834] 3'UTR
374	1643	3CCGACUUGUCACGCAUGA	2222	UCAUGCGUGACAAGUCGGC	Rh	[1232-1250] ORF
375	1644	CAGGAUGGCAGGAGGCAUA	2223	UAUGCCUCCUGCCAUCUG		[1503-1521] 3'UTR
376	1645	ACAAGAACAAGGCCGACUA	2224	UAGUCGGCCUUGUUCUUGU	Rh	[1221-1239] ORF
377	1646	UGCGUCUCCUCCUGCUUCA	2225	UGAAGCAGGAGGGAGCGCA		[231-249] ORF
378	1647	GGCGAGCUGCUGCGCUCAA	2226	UUGAGCGCAGCAGCUCGCC	Rh	[563-581] ORF
379	1648	GAUGCACCGACAGGCCUA	2227	UAGGCCUGUCCGGUGCAUC	Rh,Rb,Rt, M,P	[937-955] ORF
380	1649	CGUGUCGCUGGGCGGCAAA	2228	UUUGCCGCCAGCGACACG		[469-487] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
381	1650	AUCCCAACCUUCUCCCAACA	2229	UGUUGGGAGAGGUUGGGAU	Rh	[1893-1911] 3'UTR
382	1651	UGUUCUACGCCGACCACCA	2230	UGGUGGUCGGCGUAGAACA	Rh	[1371-1389] ORF
383	1652	CGGCCUGGCCUUCAGCUUA	2231	UAAGCUGAAGGCCAGGCCG		[373-391] ORF
384	1653	GUCGCAGGCCAAGGCAGUA	2232	UACUGCCUUGGCCUGCGAC		[499-517] ORF
385	1654	AGUCAUUCUGCCUGCCCUA	2233	UAGGGCAGGCAGAAUGACU		[1735-1753] 3'UTR
386	1655	CCCAGAAUGACCUGGCCGA	2234	UCGGCCAGGUCAUUCUGGG		[1949-1967] 3'UTR
387	1656	ACAAGAUGGUGGACAACCA	2235	UGGUUGUCCACCAUCUUGU	Rh,Rb,M, P	[876-894] ORF
388	1657	GCUAGUCAACGCCAUGUUA	2236	UAACAUGGCGUUGACUAGC	Rh	[826-844] ORF
389	1658	ACGCCACCGCCUUUGAGUA	2237	UACUCAAGGCGGUGGCGU	Rh	[1287-1305] ORF
390	1659	3CCGCGCAGACCACCGACA	2238	UGUCGGUGGUCUGCGCGGC		[758-776] ORF
391	1660	3CUAUUCAUUGGGCGCCUA	2239	UAGGCGCCCAAUGAAUAGC	D	[1429-1447] ORF
392	1661	CUCAGUGAGCUUCGCUGAA	2240	UUCAGCGAAGCUCACUGAG	UUCAGCGAAGCUCACUGAG	[643-661] ORF
393	1662	GGAGGUGAGGUACCAGCCA	2241	UGGCUGGUACCUCACCUCC		[1563-1581] 3'UTR
394	1663	GCCAAGGCAGUGCUGAGCA	2242	UGCUCAGCACUGCCUUGGC	Rh	[506-524] ORF
395	1664	CUCUCCAGCCUCAUCAUA	2243	UGAUGAUGAGGCUUGGAGA	Rh,D,Rt, M,P	[1019-1037] ORF
396	1665	GAAUGACCUGGCCGCAGUA	2244	UACUGCGGCCAGGUCAUUC		[1953-1971] 3'UTR
397	1666	UGGUGACUCGGUCCUAUAA	2245	UUAUAGGACCGAGUCACCA	Rh	[903-921] ORF
398	1667	CAGGUACCUUCUACCUGA	2246	UCAGGUGAGAAGGUACCUG	Rh	[1800-1818] 3'UTR
399	1668	GUUCCACGCCACCGCCUUA	2247	UAAGGCGGUGGCGUGGAAC	D	[1282-1300] ORF
400	1669	CCGACUGUACGGACCCAGA	2248	UCUGGGUCCGUACAGUCGG		[625-643] ORF
401	1670	GCAGACCACCGACGGCAAA	2249	UUUGCCGUCGGUGGUCUGC	D,Rt	[763-781] ORF
402	1671	AAGAUGCGAGACGAGUUA	2250	UUAACUCGUCUCGCAUCUU	Rh	[1466-1484] ORF
403	1672	CAAAGAGCAGCUGAAGUA	2251	UAUCUUCAGCUGCUCUUUG	Rh	[1087-1105] ORF
404	1673	ACGACGAGAAGGAAAAGCA	2252	UGCUIUUUCCUUCGUCGU	Rh	[969-987] ORF
405	1674	CACUCCACUUGGACAUGGA	2253	UCCAUGUCCAAGUGGAGUG	Rh	[1659-1677] 3'UTR
406	1675	AGUCCAUCAACGAGUGGGA	2254	UCCCACUCGUUGAUGGACU	Rh,Rt,M	[741-759] ORF
407	1676	GCGCCGGCCUGGCCUUCAA	2255	UUGAAGGCCAGGCCGGCGC	Rh	[369-387] ORF
408	1677	GGAAAAGCUGCAAAUCGUA	2256	UACGAUUUGCAGCUUUUCC	Rh	[979-997] ORF
409	1678	ACAUUUUGUUGGAGCGUGA	2257	UCACGCUCCAACAAAUGU		[2184-2202] 3'UTR
410	1679	ACCGUGGCUUCAUGGUGAA	2258	UUCACCAUGAAGCCACGGU	Rh,Rt,M	[891-909]

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
						ORF
411	1680	CCCUUCAUCUCCUAGUGA	2259	UCACUAGGAAGAUGAAGGG		[1388-1406] ORF
412	1681	GAAAUUCCACCACAAGAU	2260	UAUCUUGUGGUGGAAUUUC	Rh	[865-883] ORF
413	1682	CUAUAAAACUAGGUGCUGA	2261	UCAGCACCUAGUUUUUAG	Rh	[1910-1928] 3'UTR
414	1683	GGAGGUGCACGCCGGCCUA	2262	UAGGCCGGCGUGCACCUC	ORF	[544-562] ORF
415	1684	GCAGGCCAAGGCAGUGCUGA	2263	UAGCACUGCCUUGGCCUGC		[502-520] ORF
416	1685	UGAGACCAAAUUGAGCUAA	2264	UUAGCUCAAUUUGGUCUCA	Rh	[1818-1836] 3'UTR
417	1686	GCCAUAGUCAUUCUGCCUA	2265	UAGGCAGAAUGACUAUGGC		[1730-1748] 3'UTR
418	1687	AGCUGAAGAUCUGGAUGGA	2266	UCCAUCCAGAUCUUCAGCU	Rh,D	[1095-1113] ORF
419	1688	CCAUCUCCUUGCCCAAGGA	2267	UCCUUGGGCAAGGAGAUGG	Rh	[1137-1155] ORF
420	1689	CCCAGAUAAGCCUGCCUA	2268	UAGGCAGGCUUGAUCUGGG	Rh	[1759-1777] 3'UTR
421	1690	GCUGUUGCCAUCUCCUUGA	2269	UCAAGGAGAUGGCAACAGC		[1130-1148] ORF
422	1691	CGAGGUCACCAAGGACGUA	2270	UACGUCCUUGGUGACCUCG	Rh,D	[787-805] ORF
423	1692	CAACUAUAAAACUAGGUGA	2271	UCACCUAGUUUUUAUAGUUG	Rh	[1907-1925] 3'UTR
424	1693	GAAGGCUGUUGCCAUCUCA	2272	UGAGAUGGCAACAGCCUUC	Rt	[1126-1144] ORF
425	1694	UGCGCAGAAGUUGAGCCCA	2273	UGGGCUCAACUUCUCCGCA		[325-343] ORF
426	1695	CUCCUUGCCCAAGGGUGUA	2274	UACACCCUUGGGCAAGGAG	Rh	[1141-1159] ORF
427	1696	GCCCUGAAAGUCCAGAU	2275	UAUCUGGGACUUUCAGGGC		[1748-1766] 3'UTR
428	1697	CAAGGGUGUGGUGGAGGUA	2276	UACCUCACCACACCCUUG	Rh,D	[1150-1168] ORF
429	1698	AAGAGCAGCUGAAGAUUA	2277	UAGAUCUUCAGCUGCUCUU	Rh	[1089-1107] ORF
430	1699	GAAGAUGCAGAAGAAGGCA	2278	UGCCUUCUUCUGCAUCUUC	Rh,Rb,Rt	[1114-1132] ORF
431	1700	CGGAAACUCCACAUCUGA	2279	UCAGGAUGUGGAGUUUCCG		[1702-1720] 3'UTR
432	1701	AGUCAACGCCAUGUUCUUA	2280	UAAGAACAUGGCGUUGACU	Rh	[829-847] ORF
433	1702	CGAGCGCCUUGAAAAGCUA	2281	UAGCUUUUCAAGGCGCUCG		[1063-1081] ORF
434	1703	AUACCGUGGGUGUCAUGAA	2282	UUCAUGACACCCACGGUAU	Rh	[918-936] ORF
435	1704	GACCUUGGCCAUAGUCAUA	2283	UAUGACUAUGGCCAGGUC		[1723-1741] 3'UTR
436	1705	CAUGUCAGGCAAGAAGGAA	2284	UUCUUCUUGCCUGACAUG	Rh,D	[1246-1264] ORF
437	1706	UGCGAGACGAGUUUAUAGGA	2285	UCCUAUAACUCGUCUCGCA	Rh	[1470-1488] ORF+3'UTR
438	1707	CGCAACGUGACCUGGAAGA	2286	UCUUCAGGUCACGUUGCG		[599-617] ORF
439	1708	AGCAAGCAGCACUACAACA	2287	UGUUGUAGUGCUGCUUGCU	Rh,D	[677-695] ORF
440	1709	GCUGCUGCGCUCACUCAGA	2288	UCUGAGUGAGCGCAGCAGC	Rh	[568-586] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
441	1710	UGAUGAUGCACCGGACAGA	2289	UCUGUCCGGUGCAUCAUCA	Rh	[933-951] ORF
442	1711	UUGUUGCUAUCAAUCCAAA	2290	UUUGGAUUGAUAGCAACAA	Rh	[2122-2140] 3'UTR
443	1712	CCUUGAAAAGCUGCUAACA	2291	UGUUAGCAGCUUUUCAAGG		[1069-1087] ORF
444	1713	CCCUUUGACCAGGACAUCA	2292	UGAUGUCCUGGUCAAAGGG	Rh,Rt	[1322-1340] ORF
445	1714	GAGGUGAAGAAACCUGCAA	2293	UUGCAGGUUUUCUACCUC	Rh	[287-305] ORF
446	1715	CCCAAGGGUGUGGUGGAGA	2294	UCUCCACCACACCCUUGGG	Rh,D	[1148-1166] ORF
447	1716	CCCUGCUAUUCAUUGGGCA	2295	UGCCCAAUGAAUAGCAGGG	D	[1425-1443] ORF
448	1717	CUGAAAGUCCCAGAUCAAA	2296	UUUGAUCUGGGACUUUCAG		[1751-1769] 3'UTR
449	1718	GCUGCAAAUCGUGGAGAU	2297	UAUCUCCACGAUUUGCAGC	Rh	[985-1003] ORF
450	1719	CAAGCCUGCCUCAAUCA	2298	UCUGAUUGAGCCAGGCUUG	Rh	[1766-1784] 3'UTR
451	1720	CGAGCAGCUGCGCGACGAA	2299	UUCGUCGCGCAGCUGCUCG		[526-544] ORF
452	1721	AGGCCGACUUGUCACGCAA	2300	UUGCGUGACAAGUCGGCCU	Rh	[1230-1248] ORF
453	1722	GCAGCAGCUCCUGGCACUA	2301	UAGUGCCAGGAGCUGCUGC		[308-326] ORF
454	1723	3GCCAUAGUCAUUCUGCCA	2302	UGGCAGAAUGACUAUGGCC		[1729-1747] 3'UTR
455	1724	CCCUGUGCCUGAGCGGAA	2303	UUCCGCUCAGGCACACGGG	Rh	[1624-1642] 3'UTR
456	1725	CAGCUGAAGAUCUGGAUGA	2304	UCAUCCAGAUUCUAGCUG	Rh,D	[1094-1112] ORF
457	1726	CAAGCCACACUGGGAUGAA	2305	UUCAUCCAGUGUGGCUUG	Rh,Rb	[847-865] ORF
458	1727	GAAUUCACUCCACUUGGAA	2306	UCCAAGUGGAGUGAAUUC	Rh	[1654-1672] 3'UTR
459	1728	CGGCGCCCUGCUAGUCAA	2307	UUUGACUAGCAGGGCGCCG	Rh	[817-835] ORF
460	1729	UGGAAGCUGGGCAGCCGAA	2308	UUCGGCUGCCCAGCUUCCA		[611-629] ORF
461	1730	GGCAAGGCGACCACGGCGA	2309	UCGCCGUGGUCGCCUUGCC	Rh	[482-500] ORF
462	1731	CACUGCGGAGAAGUUGAGA	2310	UCUCAACUUCUCCGCAGUG		[322-340] ORF
463	1732	GGCAGGAGGCAUCCAAAGA	2311	UCUUUGGAUGCCUCCUGCC		[1509-1527] 3'UTR
464	1733	GGUGACUCGGUCCUAUACA	2312	UGUAUAGGACCGAGUCACC	Rh	[904-922] ORF
465	1734	UUUAUAGCCAGGUACCUUA	2313	UAAGGUACCUGGCUAUAAA	Rh	[1792-1810] 3'UTR
466	1735	GGCCAUGGCCAAGGACCAA	2314	UUGGUCCUUGGCCAUGGCC	Rh,D	[397-415] ORF
467	1736	CAAAGAUAGGGAGGGAAGA	2315	UCUUCCCUCCCUAUCUUUG		[2089-2107] 3'UTR
468	1737	UCUUCUGACACUAAAACAA	2316	UUGUUUUAGUGCAGAAG A		[1853-1871] 3'UTR
469	1738	CUUCUGACACUAAAACACA	2317	UGUGUUUUAGUGCAGAA G		[1854-1872] 3'UTR
470	1739	UCACGUGGAGCCUCUCGAA	2318	UUCGAGAGGCUCACGUGA		[1048-1066]

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
						ORF
471	1740	CAGUCCAUCAACGAGUGGA	2319	UCCACUCGUUGAUGGACUG	Rh,Rt,M	[740-758] ORF
472	1741	AGACCAAUUGAGCUAGGA	2320	UCCUAGCUCAAUUUGGUCU		[1820-1838] 3'UTR
473	1742	GGGUUCCCGUGUGCCUGAA	2321	UUCAGGCACACGGGAACCC	Rh	[1619-1637] 3'UTR
474	1743	UUGCUAUCAAUCCAAGAAA	2322	UUUCUUGGAUUGAUAGCAA	Rh	[2125-2143] 3'UTR
475	1744	CAACCGUGGCUUCAUGGUA	2323	UACCAUGAAGCCACGGUUG	Rh,Rt,M	[889-907] ORF
476	1745	CUGUACGGACCCAGCUCAA	2324	UUGAGCUGGGUCCGUACAG		[629-647] ORF
477	1746	CAGCAGCAAGCAGCACUAA	2325	UUAGUGCUGCUUGCUGCUG	Rh,D	[673-691] ORF
478	1747	CUGCAGCCGCAGCAGCUA	2326	UAGCUGCUGCGGCUGCAGG		[299-317] ORF
479	1748	GACACUAAAACACCUCAGA	2327	UCUGAGGUGUUUUAGUGUC		[1859-1877] 3'UTR
480	1749	CAACUGCGAGCACUCCAAA	2328	UUUGGAGUGCUCGCAGUUG	Rh,D	[691-709] ORF
481	1750	ACUGCGGAGAAGUUGAGCA	2329	UGCUCACUUCUCCGCAGU	ORF	[323-341] ORF
482	1751	GCGCCUGCUAGUCAACGA	2330	UCGUUGACUAGCAGGGCGC	Rh	1819-8371 ORF
483	1752	GGAAGCUGGGCAGCCGACA	2331	UGUCGGCUGCCCAGCUUCC		[612-630] ORF
484	1753	AGGCUCCUGAGACACAUGA	2332	UCAUGUGUCUCAGGAGCCU	D	[1525-1543] 3'UTR
485	1754	CGACAAGCGCAGCGCGCUA	2333	UAGCGCGCUGCGCUUGUCG		[721-739] ORF
486	1755	UCAGUGAGCUUCGCUGAUA	2334	UAUCAGCGAAGCUCACUGA		[644-662] ORF
487	1756	UUGAGAAGGAGCUCCAGAA	2335	UCUGGGAGCUCCUUCUCAA		[1979-1997] 3'UTR
488	1757	ACUGCGAGCACUCCAAGAA	2336	UUCUUGGAGUGCUCGCAGU	Rh,D	[693-711] ORF
489	1758	CAUCCUGGUGUCACCCGUA	2337	UACGGGUGACACCAGGAUG		[427-445] ORF
490	1759	GUGCGCAGCAGCAAGCAGA	2338	UCUGCUUGCUGCUGCGCAC	Rh,D	[668-686] ORF
491	1760	CACGCCACCGCCUUUGAGA	2339	UCUCAAGGCGGUGGCGUG	Rh	[1286-1304] ORF
492	1761	UCUCGAGCGCCUUGAAAAA	2340	UUUUUCAAGGCGCUCGAGA	[1060-1078]	[1060-1078] ORF
493	1762	GCUUCGCUGAUGACUUCGA	2341	UCGAAGUCAUCAGCGAAGC	Rh	[651-669] ORF
494	1763	UCUCCUUGCCCAAGGGUGA	2342	UCACCCUUGGGCAAGGAGA	Rh	[1140-1158] ORF
495	1764	GCAGUCCAUCAACGAGUGA	2343	UCACUCGUUGAUGGACUGC	Rh,Rt,M	[739-757] ORF
496	1765	AGAUGGUGGACAACCGUGA	2344	UCACGGUUGUCCACCAUCU	Rh,M	[879-897] ORF
497	1766	CGGCUCCUGCUAUUCAUA	2345	UAUGAAUAGCAGGGAGCCG		[1420-1438] ORF
498	1767	AUACCAUGAUGCUGAGCCA	2346	UGGCUCAGCAUCAUGGUU		[1684-1702] 3'UTR
499	1768	AGCCAGGUACCUUCUCACA	2347	UGUGAGAAGGUACCUGGCU	Rh	[1797-1815] 3'UTR
500	1769	GAGCCCGGAAACUCCACAA	2348	UUGUGGAGUUUCCGGGCUC		[1697-1715] 3'UTR

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
501	1770	GCAGCUCCUGGCACUGCGA	2349	UCGCAGUGCCAGGAGCUGC		[311-329] ORF
502	1771	CCCAGGAGUACCAAGGACA	2350	UGUCCUUGGUGACCUCGGG	Rh,D	[785-803] ORF
503	1772	CCUGACUGAGGCCAUUGAA	2351	UUCAAUGGCCUCAGUCAGG	Rh	[1204-1222] ORF
504	1773	UGCUGAGCCCGGAAACUCA	2352	UGAGUUUCCGGGCUCAGCA		[1693-1711] 3'UTR
505	1774	GCCAUCUCCUUGCCCAAGA	2353	UCUUGGGCAAGGAGAUGGC	Rh	[1136-1154] ORF
506	1775	CAAGCAGCACUACAACUGA	2354	UCAGUUGUAGUGCUGCUUG	Rh,D	[679-697] ORF
507	1776	CAAGGCAGUGCUGAGCGCA	2355	UGCGCUCAGCACUGCCUUG	Rh	[508-526] ORF
508	1777	CAAUGACAUUUUGUUGGAA	2356	UUCCAACAAAUGUCAUUG		[2179-2197] 3'UTR
509	1778	AGUGAGCUUCGCUGAUGAA	2357	UUCAUCAGCGAAGCUCACU		[646-664] ORF
510	1779	AUGAUGAUGCACCGGACAA	2358	UUGUCCGGUGCAUCAUCAU	Rh	[932-950] ORF
511	1780	GAAACACCUGGCUGGGCUA	2359	UAGCCCAGCCAGGUGUUUC	D	[1183-1201] ORF
512	1781	CCUGCUAUUCAUUGGGCGA	2360	UCGCCCAAUGAAUAGCAGG	D	[1426-1444] ORF
513	1782	CGCCACCGCCUUUGAGUUA	2361	UAACUCAAAGGCGGUGGCG	Rh	[1288-1306] ORF
514	1783	GCUUCUCAGCGCCUUCUGA	2362	UCAGAAGGCGCUGAGAAGC		[244-262] ORF
515	1784	UGAUGCUGAGCCCGGAAAA	2363	UUUUCGGGGCUCAGCAUCA		[1690-1708] 3'UTR
516	1785	UGACCUGGCCGCGAGUGAGA	2364	UCUCACUGCGGCCAGGUCA		[1956-1974] 3'UTR
517	1786	UGCAGAAACACCUGGCUGA	2365	UCAGCCAGGUGUUUCUGCA		[1179-1197] ORF
518	1787	GCAGUGCUGAGCGCCGAGA	2366	UCUCGGCGCUCAGCACUGC		[512-530] ORF
519	1788	CGGCGCGCAACGUGACCUA	2367	UAGGUCACGUUGCGCGCCG		[594-612] ORF
520	1789	AGUGCUGAGCGCCGAGCAA	2368	UUGCUCGGCGCUCAGCACU		[514-532] ORF
521	1790	ACAGGCCUCUACAACUACA	2369	UGUAGUUGUAGAGGCCUGU	Rh,Rb,D, Rt,P	[947-965] ORF
522	1791	GCAGCUGCGCGACGAGGAA	2370	UUCCUCGUCGCGCAGCUGC	Rh,D	[529-547] ORF
523	1792	AUUGAGAAGGAGCUCCCAA	2371	UUGGGAGCUCCUUCUCAAU		[1978-1996] 3'UTR
524	1793	CGCGCAGACCACCGACGGA	2372	UCCGUCGGUGGUCUGCGCG		[760-778] ORF
525	1794	CCUGUACCUGGCCAGCGUA	2373	UACGCUGGCCAGGUACAGG	Rh	[1264-1282] ORF
526	1795	CUGAGCGGACCUUCCGAGA	2374	UCUGGGAAGGUCCGCUCAG	Rh	[1633-1651] 3'UTR
527	1796	GGCCUUCAGCUUGUACCAA	2375	UUGGUACAAGCUGAAGGCC		[379-397] ORF
528	1797	CACCCAAAGCGGCUCCCUA	2376	UAGGGAGCCGCUUUGGGUG		[1411-1429] ORF
529	1798	GCCAAGGACCAGGCAGUGA	2377	UCACUGCCUGGUCCUUGGC	Rh	[404-422] ORF
530	1799	CUCAGGGUGCACACAGGAA	2378	UUCCUGUGUGCACCUCAG		[1490-1508] 3'UTR
531	1800	CGAGCUGCUGCGCUCACUA	2379	UAGUGAGCGCAGCAGCUCG	Rh	[565-583] ORF

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
532	1801	GGCUGGGCCUGACUGAGGA	2380	UCCUCAGUCAGGCCAGCC		[1197-1215] ORF
533	1802	CCGCAGCAGCUCCUGGCAA	2381	UUGCCAGGAGCUGCUGCGG		[306-324] ORF
534	1803	UGUGGGACCUGGGCCAUAA	2382	UUAUGGCCCCAGGUCCCACA		[1718-1736] 3'UTR
535	1804	AAGAUGCAGAAGAAGGCUA	2383	UAGCCUUCUUCUGCAUCUU	Rh,Rt,M	[1115-1133] ORF
536	1805	CCACGGCGCGCAACGUGAA	2384	UUCACGUUGCGCGCCGUGG	Rh	[591-609] ORF
537	1806	ACCUUCUCACCUGUGAGAA	2385	UUCUCACAGGUGAGAAGGU	Rh	[1805-1823] 3'UTR
538	1807	UGAAGAAACCUGCAGCCGA	2386	UCGGCUGCAGGUUUCUUA		[291-309] ORF
539	1808	CAGCACUACAACUGCGAGA	2387	UCUCGCAGUUGUAGUCUG	Rh,D	[683-701] ORF
340	1809	GACGAAGCGCAGCGCGCA	2388	UGCGCGCUGCGCUUGUCGC		[720-738] ORF
541	1810	UAGAAUUCACUCCACUUGA	2389	UCAAGUGGAGUGAAUUCUA	Rh	[1652-1670] 3'UTR
542	1811	GUGGAAAAACAGACCGGGA	2390	UCCCGGUCUGUUUUUCCAC		[1603-1621] 3'UTR
543	1812	ACGUGGAGCCUCUCGAGCA	2391	UGCUCGAGAGGCUCCACGU	ORF	[1050-1068] ORF
544	1813	3GCGCGCAACGUGACCUGA	2392	UCAGGUCACGUUGCGCGCC		[595-613] ORF
545	1814	UGGACAACCGUGGCUUCAA	2393	UUGAAGCCACGGUUGUCCA	Rh,M	[885-903] ORF
546	1815	CUAGUCAACGCCAUGUUA	2394	UGAACAUGGCGUUGACUAG	Rh	[827-845] ORF
547	1816	AGAAUGACCUGGCCGAGA	2395	UCUGCGGCCAGGUCAUUCU		[1952-1970] 3'UTR
548	1817	AGCUGCUGCGCUCACUCAA	2396	UUGAGUGAGCGCAGCAGCU	Rh	[567-585] ORF
549	1818	CUCUAUCCCAACCUCUCCA	2397	UGGAGAGGUUGGGAUAGA G	Rh	[1889-1907] 3'UTR
550	1819	GCGAGCUGCUGCGCUCACA	2398	UGUGAGCGCAGCAGCUCGC	Rh	[564-582] ORF
551	1820	CGCAGCAGCAAGCAGCACA	2399	UGUGCUGCUUGCUGCUGCG	Rh,D	[671-689] ORF
552	1821	GCUGGGCUGGGCCUGACA	2400	UGUCAGGCCAGCCCAGCC		[1192-1210] ORF
553	1822	UCUCCAGCCUCAUCAUCCA	2401	UGGAUGAUGAGGCUAGAG A	Rh,D,Rt, M	[1020-1038] ORF
554	1823	CAACGCCAUGUUCUUCAAA	2402	UUUGAAGAACAUGGCGUUG	Rh,Rb,P	[832-850] ORF
555	1824	UGGCACUGCGGAGAAGUUA	2403	UAACUUCUCCGAGUGCCA		[319-337] ORF
556	1825	UUUGAGUUGGACACAGAU	2404	UAUCUGUGUCCAACUCAA		[1298-1316] ORF
557	1826	UGGGCGAGCUGCUGCGCUA	2405	UAGCGCAGCAGCUCGCCA	Rh	[561-579] ORF
558	1827	CUGCUAACCAAAGAGCAGA	2406	UCUGCUCUUUGGUUAGCAG		[1079-1097] ORF
559	1828	AACGUGACCUGGAAGCUGA	2407	UCAGCUUCCAGGUCACGUU		[602-620] ORF
560	1829	AUGACAUUUUGUUGGAGCA	2408	UGCUGCAACAAAUGUCAU		[2181-2199] 3'UTR
561	1830	CAGGAGGCAUCCAAAGGCA	2409	UGCCUUUGGAUGCCUCCUG		[1511-1529]

No.	SEQ ID	Sense siRNA	SEQ ID	AntiSense siRNA	Other Sp	human-32454740 ORF:230-1486
						3'UTR
562	1831	AUCUCCUUGCCCAAGGGUA	2410	UACCCUUGGGCAAGGAGAU	Rh	[1139-1157] ORF
563	1832	UGGGAUGAGAAAUCCACA	2411	UGUGGAAUUUCUCAUCCCA	Rh	[857-875] ORF
564	1833	AAAGCUGCUAACCAAAGAA	2412	UUCUUUGGUUAGCAGCUUU		[1075-1093] ORF
565	1834	AGGAGGCAUCCAAAGGCUA	2413	UAGCCUUUGGAUGCCUCCU		[1512-1530] 3'UTR
566	1835	CACCGCCUUUGAGUUGGAA	2414	UUCCAACUCAAGGCGGUG	Rh	[1291-1309] ORF
567	1836	CCAACUAUAAAACUAGGUA	2415	UACCUAGUUUUUAUAGUUGG	Rh	[1906-1924] 3'UTR
568	1837	CAAGAAGCAUCGUGUCUGA	2416	UCAGACACGAUGCUUCUUG	Rh	[2022-2040] 3'UTR
569	1838	AGCAGCUGAAGAUCUGGAA	2417	UUCCAGAUCUUCAGCUGCU	Rh,D	[1092-1110] ORF
570	1839	GCGCUCCCUCCUGCUUCUA	2418	UAGAAGCAGGAGGGAGCGC		[232-250] ORF
571	1840	UGCUAGUCAACGCCAUGUA	2419	UACAUGGCGUUGACUAGCA	Rh	[825-843] ORF
572	1841	CGCCGAGCAGCUGCGCGAA	2420	UUCGCGCAGCUGCUCGGCG		[523-541] ORF
573	1842	CCGCGCAGACCACCGACGA	2421	UCGUCGGUGGUCUGCGCGG		[759-777] ORF
574	1843	UAGCCAGGUACCUUCUCAA	2422	UUGAGAAGGUACCUGGCUA	Rh	[1796-1814] 3'UTR
575	1844	UGCUUCUCAGCGCCUUCUA	2423	UAGAAGGCGCUGAGAAGCA		[243-261] ORF
576	1845	CUCCCUCCUGCUUCUCAGA	2424	UCUGAGAAGCAGGAGGGAG		[235-253] ORF
577	1846	CGCAGGCCAAGGCAGUGCA	2425	UGCACUGCCUUGGCCUGCG		[501-519] ORF
578	1847	3CAAGGCGACCACGGCGUA	2426	UACGCCGUGGUCGCCUUGC	Rh	[483-501] ORF
579	1848	GCAGCCGCAGCAGCUCCUA	2427	UAGGAGCUGCUGCGGCUGC		[302-320] ORF

TABLE E: SERPINH1 Cross-Species 18+1-mer siRNAs

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
1	2428	UCACCAAGGACGUGGAGCA	2576	UGCUCCACGUCCUUGGU GA	Rh,D	[792-810] ORF
2	2429	CAGCGCGCUGCAGUCCAUA	2577	UAUGGACUGCAGCGCGC UG	Rh,Rt	[730-748] ORF
3	2430	CAUCUACGGGCGCGAGGAA	2578	UUCUCCGCGCCCGUAGA UG	D,M	[1336-1354] ORF
4	2431	CUCCAGCCUCAUCAUCCUA	2579	UAGGAUGAUGAGGCGUG AG	Rh,D,Rt,M	[1021-1039] ORF
5	2432	GACAUUCACGGGCGCGAGA	2580	UCUCGGGCGCGUAGAUG UC	D,M	[1334-1352] ORF
6	2433	CGUGCGCAGCAGCAAGCAA	2581	UUGCUUGCUUCUUGCGCA CG	Rh,D,M	[667-685] ORF
7	2434	GUCACCAAGGACGUGGAGA	2582	UCUCCACGUCCUUGGUG AC	Rh,D	[791-809] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
8	2435	CCGCGACAAGCGCAGCGCA	2583	UGCGCUGCGCUUGGCGC GG	D	[718-736] ORF
9	2436	GCGCAGCGCGCUGCAGUCA	2584	UGACUGCAGCGCGCUGC GC	Rh,Rt	[727-745] ORF
10	2437	3GCCACAAGCUCUCCAGA	2585	UCUGGAGAGCUUGUGGG CC	Rh,D,P	[1009-1027] ORF
11	2438	CAAGGACGUGGAGCGCACA	2586	UGUGGCGUCCACGUCCU UG	Rh,D	[796-814] ORF
12	2439	AGCCUCAUCAUCCUCAUGA	2587	UCAUGAGGAUGAUGAGG CU	Rh,D,Rt,M	[1025-1043] ORF
13	2440	GGUGUGGUGGAGGUGACCA	2588	UGGUCACCUCCACCACA CC	Rh,D	[1154-1172] ORF
14	2441	GCAAGCUGCCCAGGUCAA	2589	UUGACCUCGGCAGCUU GC	Rh,D	[777-795] ORF
15	2442	GUGGAGGUGACCCAUGACA	2590	UGUCAUGGGUACCUCC AC	Rh,Rt,M	[1160-1178] ORF
16	2443	CACAAGAUGGUGGACAACA	2591	UGUUGUCCACCAUCUUG UG	Rh,Rb,M,P	[875-893] ORF
17	2444	GCGAGGAGCUGCGCAGCCA	2592	UGGCGCGCAGCUCCUC GC	D,M	[1347-1365] ORF
18	2445	UACUACGACGACGAGAAGA	2593	UCUUCUGUGUCGUAG UA	Rb	[962-980] ORF
19	2446	GAGGUGACCCAUGACCUGA	2594	UCAGGUCAUGGUCACC UC	Rh,Rt,M	[1163-1181] ORF
20	2447	ACUUCCGCGACAAGCGCAA	2595	UUGCGCUUGUCGCGGAA GU	D	[714-732] ORF
21	2448	GCCCACAAGCUCUCCAGCA	2596	UGCUGGAGAGCUUGUGG GC	Rh,D,P	[1010-1028] ORF
22	2449	GCGCAGCAGCAAGCAGCAA	2597	UUGCUGCUUGCUGCUGC GC	Rh,D	[670-688] ORF
23	2450	CGAGGAGCUGCGCAGCCCA	2598	UGGGCUGCGCAGUCCU CG	D,M	[1348-1366] ORF
24	2451	AACGCCAUGUUCUUAAGA	2599	UCUUGAAGAACAUGGCG UU	Rh,Rb,P	[833-851] ORF
25	2452	GUCAGGCAAGAAGGACCUA	2600	UAGGUCCUUCUUGCCUG AC	Rh,D	[1249-1267] ORF
26	2453	GCCUGGGCGAGCUGCUGCA	2601	UGCAGCAGCUCGCCAG GC	Rh,D	[558-576] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
27	2454	GAUGAUGCACCGGACAGGA	2602	UCCUGUCCGGUGCAUCAUC	Rh,Rb,Rt,M	[934-952] ORF
28	2455	GGACCUUGUACCUGGCCAGA	2603	UCUGGCAGGUACAGGUCC	Rh,D	[1261-1279] ORF
29	2456	GCGACGAGGAGGUGCACGA	2604	UCUGGCACCUCCUCGUCGC	D	[537-555] ORF
30	2457	UGUGGUGGAGGUGACCCAA	2605	UUGGGUACCUCCACCA CA	Rh,D	[1156-1174] ORF
31	2458	UUCAAGCCACACUGGGAUA	2606	UAUCCAGUGUGGCUUGAA	Rh,Rb	[845-863] ORF
32	2459	CAAGAUGGUGGACAACCGA	2607	UCCGUUGUCCACCAUCUUG	Rh,Rb,M,P	[877-895] ORF
33	2460	UCAACUCCGCGACAAGCA	2608	UGCUUGUCGCGGAAGUUGA	D	[711-729] ORF
34	2461	AUUCAUUGGGCGCCUGGUA	2609	UACCAGGCGCCCAUUGAAU	D	[1432-1450] ORF
35	2462	CUCCAAGAUCAACUCCGA	2610	UCGGAAGUUGAUCUUGAG	Rh,D,Rt,M	[703-721] ORF
36	2463	CAGGCCAUGGCCAAGGACA	2611	UGUCCUUGGCCAUGGCCUG	Rh,D	[395-413] ORF
37	2464	GUACCAGGCCAUGGCCAAA	2612	UUUGGCCAUGGCCUGGUAC	Rh,D	[391-409] ORF
38	2465	UGUCAGGCAAGAAGGACCA	2613	UGGUCCUUCUUGCCUGACA	Rh,D	[1248-1266] ORF
39	2466	CUUCGUGCGCAGCAGCAAA	2614	UUUGCUGCUGCGCACGAAG	Rh,D,M	[664-682] ORF
40	2467	CAACUCCGCGACAAGCGA	2615	UCGCUUGUCGCGGAAGUUG	D	[712-730] ORF
41	2468	CCACCACAAGAUGGUGGAA	2616	UUCACCAUCUUGUGGUUG	Rh,Rb,D,P	[871-889] ORF
42	2469	GCGCGACGAGGAGGUGCAA	2617	UUGCACCUCCUCGUCGCGC	Rh,D	[535-553] ORF
43	2470	CUACAACUGCGAGCACUCA	2618	UGAGUGCUCGCAGUUGUAG	Rh,D	[688-706] ORF
44	2471	UGGAGGUGACCCAUGACCA	2619	UGGUCAUGGGUACCCUCA	Rh,Rt,M	[1161-1179] ORF
45	2472	GAGGUCACCAAGGACGUGA	2620	UCACGUCCUUGGUGACCUC	Rh,D	[788-806] ORF
46	2473	AAGAAGGACCUGUACCUGA	2621	UCAGGUACAGGUCCUUCUU	Rh,D	[1256-1274] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
47	2474	GACAACCGUGGCUUCAUGA	2622	UCAUGAAGCCACGGUUG UC	Rh,Rt,M	[887-905] ORF
48	2475	ACCAGGACAUCUACGGGCA	2623	UGCCCGUAGAU GUCCUG GU	D,Rt	[1329-1347] ORF
49	2476	GCUGCCCAGGUCACCAA	2624	UUUGGUGACCU CGGGCA GC	Rh,D	[781-799] ORF
50	2477	AUGCAGAAGAAGGCUGUUA	2625	UAACAGCCUUCUUCUGC AU	Rt	[1118-1136] ORF
51	2478	GGCCUGGGCGAGCUGCUGA	2626	UCAGCAGCUCGCCAGG CC	Rh,D	[557-575] ORF
52	2479	GAUGGUGGACAACCGUGGA	2627	UCCACGGUUGUCCACCA UC	Rh,M	[880-898] ORF
53	2480	CUCCCUGCUAUUCAUUGGA	2628	UCCAAUGAAUAGCAGGG AG	D	[1423-1441] ORF
54	2481	GAAGGACCUGUACCUGGCA	2629	UGCCAGGUACAGGUCCU UC	Rh,D	[1258-1276] ORF
55	2482	CCACCGACGGCAAGCUGCA	2630	UGCAGCUUGCCGUCGGU GG	D,Rt	[768-786] ORF
56	2483	UGCUAUUCAUUGGGCGCCA	2631	UGGCGCCCAUGAAUAG CA	D	[1428-1446] ORF
57	2484	AUGUUCUUAAGCCACACA	2632	UGUGUGGCUUGAAGAAC AU	Rh,Rb,D	[839-857] ORF
58	2485	CCAGGACAUCUACGGGCGA	2633	UCGCCCGUAGAU GUCCU GG	D,Rt	[1330-1348] ORF
59	2486	GCGCGAGGAGCUGCGCAGA	2634	UCUGCGCAGCUCCUGC GC	Rh,D,M	[1345-1363] ORF
50	2487	GAGCAGCUGCGCGACGAGA	2635	UCUCGUCGCGCAGCUGC UC	Rh,D	[527-545] ORF
61	2488	CUAUUCAUUGGGCGCCUGA	2636	UCAGGCGCCCAUGAAU AG	D	[1430-1448] ORF
62	2489	ACAAGCUCUCCAGCCUCAA	2637	UUGAGGCGGAGAGCUU GU	Rh,D,M,P	[1014-1032] ORF
63	2490	GCUGAAGAUCUGGAUGGGA	2638	UCCAUCCAGAUUUCA GC	Rh,D	[1096-1114] ORF
54	2491	GACCAGGACAUCUACGGGA	2639	UCCCGUAGAU GUCCUGG UC	D,Rt	[1328-1346] ORF
65	2492	CAAGCGCAGCGCGCUGCAA	2640	UUGCAGCGCGCUGCGCU UG	Rh,Rt	[724-742] ORF
66	2493	CCAUGGCCAAGGACCAGGA	2641	UCCUGGUCCUUGGCCAU GG	Rh,D	[399-417] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
67	2494	CACCAAGGACGUGGAGCGA	2642	U'CGUCCACGUCUUGG UG	Rh,D	[793-811] ORF
68	2495	CCGUGGCUUCAUGGUGACA	2643	UGUCAACGAAGCCAC GG	Rh,Rt,M	[892-910] ORF
69	2496	UGACCAGGACAUCUACGGA	2644	UCCGUAGAUUCCUGGU CA	Rt	[1327-1345] ORF
70	2497	AGACCACCGACGGCAAGCA	2645	UGCUUGCCGUCGGUGGU	D,Rt	[765-783] ORF
				CU		
71	2498	GACAAGCGCAGCGCGCUGA	2646	UCAGCGCGUCGCUUG UC	Rh,Rt	[722-740] ORF
72	2499	AGAAACACCUGGCUGGGCA	2647	UGCCAGCCAGGUGUUU CU	D	[1182-1200] ORF
73	2500	AAGAUGGUGGACAACCGUA	2648	U'ACGGUUGUCCACCAUC UU	Rh,M	[878-896] ORF
74	2501	CAGACCACCGACGGCAAGA	2649	U'CUUGCCGUCGGUGGUC UG	D,Rt	[764-782] ORF
75	2502	AGGACCUGUACCUGGCCAA	2650	UUGGCCAGGUACAGGUC CU	Rh,D	[1260-1278] ORF
76	2503	CUGCUAUUCAUUGGGCGCA	2651	UGCGCCCAAUGAAUAGC AG	D	[1427-1445] ORF
77	2504	GUCCAUAACGAGUGGGCA	2652	UGCCCAUCUGUUAUGG AC	Rh,Rt,M	[742-760] ORF
78	2505	CCAGGCCAUGGCCAAGGAA	2653	U'UCCUUGGCCAUGGCCU GG	Rh,D	[394-412] ORF
79	2506	AAGCAGCACUACAACUGCA	2654	U'GCAGUUGUAGUGCUGC UU	Rh,D	[680-698] ORF
80	2507	UGUUCCACGCCACCGCCUA	2655	UAGGCGGUGGCGUGGAA CA	D	[1281-1299] ORF
81	2508	UACAACUACUACGACGACA	2656	UGUCGUCGUAGUAGUUG UA	Rb	[956-974] ORF
82	2509	CCUCAUCAUCCUCAUGCCA	2657	UGGCAU'GAGGAUGAUGA' GG	Rh,D,Rt,M	[1027-1045] ORF
83	2510	UGGUGGACAACCGUGGCUA	2658	UAGCCACGGUUGUCCAC CA	Rh,M	[882-900] ORF
84	2511	GACCACCGACGGCAAGCUA	2659	UAGCUUGCCGUCGGUGG UC	D,Rt	[766-784] ORF
85	2512	AGCUGCGCGACGAGGAGGA	2660	UCCUCCUCGUCGCGCAG CU	Rh,D	[531-549] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
86	2513	CGGCAAGCUGCCCGAGGUA	2661	UACCUCGGGCAGCUUGC CG	Rh,D	[775-793] ORF
87	2514	UGGCCCACAAGCUCUCCAA	2662	UUGGAGAGCUUGUGGGC CA	Rh,D,P	[1008-1026] ORF
88	2515	CAGCUGCGCGACGAGGAGA	2663	UCUCCUCGUGCGCAGC UG	Rb,D	[530-548] ORF
89	2516	CUUCCGCGACAAGCGCAGA	2664	UCUGCGCUUGUCGCGGA AG	D	[715-733] ORF
90	2517	UGGGCCUGACUGAGGCCAA	2665	UUGGCCUCAGUCAGGCC CA	Rt	[1200-1218] ORF
91	2518	GCUGCGCGACGAGGAGGUA	2666	UACCUCUCGUGCGCA GC	Rh,D	[532-550] ORF
92	2519	CAGOACAUCUACGGGCGCA	2667	UGCGCCGUAGAUGUCC UG	D	[1331-1349] ORF
93	2520	GCCAUGGCCAAGGACCAGA	2668	UCUGGUCCUUGGCCAUG GC	Rh,D	[398-416] ORF
94	2521	UCCAAGAUCAACUCCGCA	2669	UGCGGAAGUUGAUCUUG GA	D	[704-722] ORF
95	2522	ACCACCGACGGCAAGCUGA	2670	UCAGCUUGCCGUCGGUG GU	D,Rt	[767-785] ORF
96	2523	AUCUACGGGCGCGAGGAGA	2671	UCUCCUCGCGCCGUAG AU	D,M	[1337-1355] ORF
97	2524	CUGCCCGAGGUCACCAAGA	2672	UCUUGGUGACCUCCGC AG	Rh,D	[782-800] ORF
98	2525	AUCAACUCCGCGACAAGA	2673	UCUUGUCGCGGAAGUUG AU	D	[710-728] ORF
99	2526	UCAUUGGGCGCCUGGUCCA	2674	UGGACCAAGCGCCCAAU GA	Rh,D	[1434-1452] ORF
100	2527	CAUUGGGCGCCUGGUCCGA	2675	UCGGACCAAGCGCCCAA UG	Rh,D	[1435-1453] ORF
101	2528	GUGUUCCACGCCACCGCCA	2676	UGGCGGUGGCUGGAAC AC	D	[1280-1298] ORF
102	2529	AUGAUGCACCGGACAGGCA	2677	UGCCUGUCCGGUGCAUC AU	Rh,Rb,Rt,M ,P	[935-953] ORF
103	2530	CGACGAGGAGGUGCACGCA	2678	UGCGUGCAUCCUCGU CG	D	[538-556] ORF
104	2531	CAGAAACACCUGGCUGGGA	2679	UCCAGCCAGGUGUUUC UG	D	[1181-1199] ORF
105	2532	UGAUGCACCGGACAGGCCA	2680	UGGCCUGUCCGGUGCAU CA	Rh,Rb,Rt,M ,P	[936-954] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
106	2533	AAGGCUGUUGCCAUCUCCA	2681	UGGAGAUGCCAACAGCC UU	D,Rt	[1127-1145] ORF
107	2534	AUGACUUCGUGCGCAGCAA	2682	UUGCUGCGCACGAAGUC AU	Rh,Rt,M	[660-678] ORF
108	2535	UCAGGCAAGAAGGACCUGA	2683	UCAGGUCCUUCUUGCCU GA	Rh,D	[1250-1268] ORF
109	2536	CUCAUCAUCCUCAUGCCCA	2684	UGGGCAUGAGGAUGAUG AG	Rh,Rt,M	[1028-1046] ORF
110	2537	CGCGACGAGGAGGUGCACA	2685	UGUGCAUCCUCCUGUCG CG	Rh,D	[536-554] ORF
111	2538	ACAACCGUGGCUUCAUGGA	2686	UCCAUGAAGCCAAGGUU GU	Rh,Rt,M	[888-906] ORF
112	2539	UUGACCAGGACAUCUACGA	2687	UCGUAGAUUGUCCUGGUC AA	Rt	[1326-1344] ORF
113	2540	CAAGCUGCCCCGAGGUCACA	2688	UGUGACCUCCGGGACGCU UG	Rh,D	[778-796] ORF
114	2541	UCCUGCUAUUCAUUGGGA	2689	UCCCAAUGAAUAGCAGG GA	D	[1424-1442] ORF
115	2542	UAUUCAUUGGGCGCCUGGA	2690	UCCAGGCGCCCAUGAA UA	D	[1431-1449] ORF
116	2543	CUGCGCGACGAGGAGGUGA	2691	UCACCUCCUCGUCGCGC AG	Rh,D	[533-551] ORF
117	2544	CUACGGGCGCGAGGAGCUA	2692	UAGCUCCUCGCGCCGU AG	D,M	[1339-1357] ORF
118	2545	CGCGAGGAGCUGCGCAGCA	2693	UGCUGGCGAGCUCCUG CG	D,M	[1346-1364] ORF
119	2546	ACACCUGGCUGGGCUGGGA	2694	UCCAGCCCAAGCAGGU GU	D	[1186-1204] ORF
120	2547	UCUACGGGCGCGAGGAGCA	2695	UGCUCUUCGCGCCCGUA GA	D,M	[1338-1356] ORF
121	2548	UUCUUAAGCCACACUGGA	2696	UCCAGUGUGGCUUGAAG AA	Rh,Rb,D	[842-860] ORF
122	2549	CCUGGGCGAGCUGCUGCGA	2697	UCGCAGCAGCUCGCCA GG	Rh,D	[559-577] ORF
123	2550	AAGAAGGCUGUUGCCAUCA	2698	UGAUGGCAACAGCCUUC UU	Rt	[1124-1142] ORF
124	2551	CGACGGCAAGCUGCCCGAA	2699	UUCGGGCGAGCUUGCCGU CG	D	[772-790] ORF
125	2552	GACGGCAAGCUGCCCGAGA	2700	UCUCGGGCGAGCUUGCCG UUC	Rh,D	[773-791] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
126	2553	UUCAUUGGGCGCCUGGUCA	2701	UGACCAGGCGCCAAUGAA	Rh,D	[1433-1451] ORF
127	2554	AAGCGCAGCGCGCUGCAGA	2702	UCUGCAGGCGCUGGCGC	Rh,Rt	[725-743] ORF
				UU		
128	2555	CCUGGCCCAACAAGCUCUCA	2703	UGAGAGCUUGUGGGCCAAG	Rh,D,P	[1006-1024] ORF
129	2556	ACGGCAAGCUGCCCGAGGA	2704	UCCUCGGGAGCUUGCCGU	Rh,D	[774-792] ORF
130	2557	UUUGACCAGGACAUCUACA	2705	UGUAGAUUGCCUGGUCAAA	Rt	[1325-1343] ORF
131	2558	UGACUUCGUGCGCAGCAGA	2706	UCUGCUGCGCACGAAGUCA	Rh,Rt,M	[661-679] ORF
132	2559	AAGGACGUGGAGCGCACGA	2707	UCGUGCGCUCCAGGUCCUU	Rh,D	[797-815] ORF
133	2560	UCCAUCAACGAGUGGGCCA	2708	UGGCCACUCUGUUGAUGGA	Rt,M	[743-761] ORF
134	2561	CACCGACGGCAAGCUGCCA	2709	UGGCAGCUUGCCGUGCGUG	D,Rt	[769-787] ORF
135	2562	ACGGGCGCGAGGAGCUGCA	2710	UGCAGCUCCUCGCGGCCGU	D,M	[1341-1359] ORF
136	2563	UCCGCGACAAGCGCAGCGA	2711	UCGCUGCGCUUGUCGCGGA	D	[717-735] ORF
137	2564	UUGGGCGCCUGGUCCGGCA	2712	UGCCGGACACGGCGCCCCAA	Rh,D	[1437-1455] ORF
138	2565	AUGGUGGACAACCGUGGCA	2713	UGCCACGGUUGUCCACC AU	Rh,M	[881-899] ORF
139	2566	AUUGGGCGCCUGGUCCGGA	2714	UCCGGACACGGCGCCCA AU	Rh,D	[1436-1454] ORF
140	2567	UACGGGCGCGAGGAGCUGA	2715	UCAGCUCCUCGCGCCCCUA	D,M	[1340-1358] ORF
141	2568	AUGCACCGGACAGGCCUCA	2716	UGAGGCCUGUCCGGUCC AU	Rh,Rb,Rt,P	[938-956] ORF
142	2569	UUCCACCACAAGAUGGUGA	2717	UCACCAUCUUGUGGUGGAA	Rh,Rb,D,P	[869-887] ORF
143	2570	UUCCGCGACAAGCGCAGCA	2718	UCUGCGCUUGUGCGCGGAA	D	[716-734] ORF
144	2571	UACCAGGCCAUGGCCAAGA	2719	UCUUGGCCAUGGCCUGGTTA	Rh,D	[392-410] ORF

No.	SEQ ID NO	Sense siRNA	SEQ ID NO	AntiSense siRNA	Other Species	human-32454740 ORF:230-1486
145	2572	AAACACCUGGCUGGGCUGA	2720	UCAGCCAGCCAGGUGU	D	[1184-1202] ORF
146	2573	ACCGACGGCAAGCUGCCCA	2721	UGGGCAGCUUGCCGUCG	D	[770-788] ORF
147	2574	AACACCUGGCUGGGCUGGA	2722	UCCAGCCAGCCAGGUGU	D	[1185-1203] ORF
148	2575	UUCGUGCGCAGCAGCAAGA	2723	UUCUGUGCUGCGCACGAA	Rh,D,M	[665-683] ORF

[0461] The most active sequences were selected from further assays. From Table 4 siRNA compounds SERPINH1_2, SERPINH1_6, SERPINH1_13, SERPINH1_45 SERPINH1_45a, SERPINH1_51, SERPINH1_51a, SERPINH1_52 and SERPINH1_86 were selected as preferred compounds (Tables 6-A and B).

TABLE 6-A: Select siRNAs

siRNA	SEQ ID SEN	SEQ ID AS	Activity 0.1nM	Activity 0.5nM	Activity 5nM	IC50 (nM)	Length
SERPINH1_2	60	127	65	48	7	.008	19
SERPINH1_6	3	130	164	39	5	.019	19
SERPINH1_11	68	135	119	54	6	.05	19
SERPINH1_13	69	136	91	24	4		19
SERPINH1_45	97	164	156	38	8	.07	19
SERPINH1_45a	98	165		(1)			19
SERPINH1_51	101	168	68	39	5	.05	19
SERPINH1_52	102	169	149	37	9	0.06	19
SERPINH1_86	123	190	121	61	0.27	0.27	19

TABLE 6B

siRNA	SEQ ID SEN	SEQ ID AS	Activity 0.026nM	Activity 0.077nM	Activity 0.23nM	Activity 0.69nM	(2) Activity 2.1nM	Activity 6.25nM	Activity 25 nM
SERPINH1_45	97	164	102	81	55	41	28	22	16
SERPINH1_45a	98	165	107	(3) 98	84	69	36	24	16

[0462] From Table 5 siRNA compounds SERPINH1_4, SERPINH1_12, SERPINH1_18, SERPINH1_30, SERPINH1_58 and SERPINH1_88 were selected as preferred compounds (Table 7).

TABLE 7: Select siRNAs

siRNA	SEQ ID NO SEN	SEQ ID NO AS	Activity 0.1nM	Activity 0.5nM	Activity 5nM	IC50 (nM)	Length
SERPINH1_4	195	220	60	35	5	.006	19
SERPINH1_12	196	221	54	42	8	.065	19
SERPINH1_18	197	222	139	43	9		19
SERPINH1_30	199	224	146	49	9	0.093	19
SERPINH1_58	208	233	na	na	8		19
SERPINH1_88	217	242	105	43	9		19

EXAMPLE 21: Animal model systems of fibrotic conditions

[0463] Testing the active siRNAs of the description may be done in predictive animal models. Rat diabetic and aging models of kidney fibrosis include Zucker diabetic fatty (ZDF) rats, aged fa/fa (obese Zucker) rats, aged Sprague-Dawley (SD) rats, and Goto Kakizaki (GK) rats; GK rats are an inbred strain derived from Wistar rats, selected for spontaneous development of NIDDM (diabetes type II). Induced models of kidney fibrosis include the permanent unilateral ureteral obstruction (UUO) model which is a model of acute interstitial fibrosis occurring in healthy non-diabetic animals; renal fibrosis develops within days following the obstruction. Another induced model of kidney fibrosis is 5/6 nephrectomy.

[0464] Two models of liver fibrosis in rats are the Bile Duct Ligation (BDL) with sham operation as controls, and CC14 poisoning, with olive oil fed

animals as controls, as described in the following references: Lotersztajn S, et al Hepatic Fibrosis: Molecular Mechanisms and Drug Targets. *Annu Rev Pharmacol Toxicol.* 2004 Oct 07; Uchio K, et al., Down-regulation of connective tissue growth factor and type I collagen mRNA expression by connective tissue growth factor antisense oligonucleotide during experimental liver fibrosis. *Wound Repair Regen.* 2004 Jan-Feb;12(1):60-6; Xu XQ, et al., Molecular classification of liver cirrhosis in a rat model by proteomics and bioinformatics *Proteomics.* 2004 Oct;4(10):3235-45.

[0465] Models for ocular scarring are well known in the art *e.g.* Sherwood MB et al., *J Glaucoma.* 2004 Oct;13(5):407-12. A new model of glaucoma filtering surgery in the rat; Miller MH et al., *Ophthalmic Surg.* 1989 May;20(5):350-7. Wound healing in an animal model of glaucoma fistulizing surgery in the Rb; vanBockxmeer FM *et al.*, *Retina.* 1985 Fall-Winter; 5(4): 239-52. Models for assessing scar tissue inhibitors; Wiedemann P et al., *J Pharmacol Methods.* 1984 Aug; 12(1): 69-78. Proliferative vitreoretinopathy: the Rb cell injection model for screening of antiproliferative drugs.

[0466] Models of cataract are described in the following publications: Zhou et al., 2002. *Invest Ophthalmol Vis Sci.*43:2293-300; Wang et al. 2004 *Curr Eye Res.* 29:51-58.

[0467] The compounds of Table 5 and Table 4 are tested in these models of fibrotic conditions, in which it is found that they are effective in treating liver fibrosis and other fibrotic conditions.

Model systems of glaucoma

[0468] Testing the active siRNA of the description for treating or preventing glaucoma is preformed in rat animal model for optic nerve crush described for example in: Maeda *et al.*, 2004 *Investigative Ophthalmology and visual Science (IOVS)*, 45:851. Specifically, for optic nerve transection the orbital optic nerve (ON) of anesthetized rats is exposed through a supraorbital approach, the meninges severed and all axons in the ON transected by crushing with forceps for 10 seconds, 2 mm from the lamina cribrosa.

[0469] Nucleic acid molecules as disclosed herein are tested in this animal model and the results show that these siRNA compounds are useful in treating and/or preventing glaucoma.

Rat Optic Nerve Crush (ONC) Model: intravitreal siRNA delivery and eye drop delivery

[0470] For optic nerve transection the orbital optic nerve (ON) of anesthetized rats is exposed through a supraorbital approach, the meninges severed and all axons in the ON transected by crushing with forceps for 10 seconds, 2 mm from the lamina cribrosa.

[0471] The siRNA compounds are delivered alone or in combination in 5uL volume (10 ug/uL) as eye drops. Immediately after optic nerve crush (ONC), 20 ug/10ul test siRNA or 10 ul PBS is administered to one or both eyes of adult Wistar rats and the levels of siRNA taken up into the dissected and snap frozen whole retinae at 5 h and 1 d, and later at 2 d, 4 d, 7 d, 14 d and 21 d post injection is determined. Similar experiments are performed in order to test activity and efficacy of siRNA administered via eye drops.

Model systems of ischemia reperfusion injury following lung transplantation in rats

[0472] Lung ischemia/reperfusion injury is achieved in a rat animal model as described in Mizobuchi et al., 2004 *J Heart Lung Transplantation*, 23 and in Kazuhiro et al., 2001 *Am. J. Respir. Cell Mol Biol*, 25:26-34.

[0473] Specifically, after inducing anesthesia with isoflurane, the trachea is cannulated with a 14-gauge Teflon catheter and the rat is mechanically ventilated with rodent ventilator using 100% oxygen, at a rate of 70 breaths per minute and 2 cm H₂O of positive end-respiratory pressure. The left pulmonary artery, veins and main stem bronchus are occluded with a Castaneda clamp. During the operation, the lung is kept moist with saline and the incision is covered to minimize evaporative losses. The period of ischemia is 60 minutes long. At the end of the ischemic period the clamp is removed and the lung is allowed to ventilate and perfuse for further 4h, 24h, and 5 d post induction of lung ischemia. At the end of the experiment, the lungs are gently harvested and either frozen for RNA extraction or fixed in glutaraldehyde cocktail for subsequent histological analysis.

The bleomycin animal model as a model for idiopathic pulmonary fibrosis (IPF)

[0474] Testing feasibility of lung and liver delivery of vitamin A-Coatsome formulated siRNA administered by intravenous injection and intratracheal administration of siRNA-vitaminA-Coatsome complex to a healthy mice and bleomycine-treated mice

[0475] Objective: To test two administration routes for feasibility of vitamin A-Coatsome formulated siRNA delivery to normal and fibrotic mouse lungs. The main hypothesis to be tested in the current study is whether systemic administration of vitamin A-Coatsome formulated modified siRNA provides efficient uptake and cell-specific distribution in the fibrotic and normal mouse lungs. Intratracheal route of vitaminA-Coatsome formulated modified siRNA will be tested in parallel. siRNA detection and cell-specific distribution in the lungs and liver will be performed by in situ hybridization (ISH)

[0476] The Bleomycin model of pulmonary fibrosis has been well developed and characterized over the last three decades (Moeller, et al. 2008 *Int J Biochem Cell Biol*, 40:362-82.; Chua et al., 2005 *Am J Respir Cell Mol Biol* 33:9-13.). Histological hallmarks, such as intra-alveolar buds, mural

incorporation of collagen and obliteration of alveolar space are present in BLM-treated animals similar to IPF patients. Early studies demonstrated that C57/B1 mice were consistently prone to BLM-induced lung fibrosis, whereas Balb/C mice were inheritantly resistant. Depending on the route of administration, different fibrotic pattern develops. Intratracheal instillation of BLM results in bronchiocentric accentuated fibrosis, whereas intravenous or intraperitoneal administration induces subpleural scarring similar to human disease (Chua *et al. ibid.*). A mouse model of usual interstitial pneumonia (UIP) is used. This model shows a heterogenous distribution of fibroproliferation, distributed mainly subpleurally, forming similar lesions to those observed in the lungs of patients with idiopathic pulmonary fibrosis (IPF) (Onuma, et al., 2001 J Exp Med 194:147-56, and Yamaguchi et al., 1988 Nature 336:244-46). UIP will be induced by intraperitoneal injection of bleomycin every other day for 7 days for a constant composition of subpleural fibroproliferation in the mouse lung (Swiderski et al. 1998 Am J Pathol 152: 821-28, and Shimizukawa et al., 2003 Am J Physiol Lung Cell Mol Physiol 284: L526-32.).

[0477] Vitamin A-loaded liposomes containing siRNA interact with retinol-binding protein (RBP) and provide efficient delivery to the HSC via RBP receptor. These liposomes are efficiently taken up by an RBP receptor-expressing activated myofibroblasts in the lungs of bleomycin-treated mice.

Study design

Mice - C57 B1 male

Starting N (BLM I.P.) - 40 (6 for the first pilot group, 34 for the study, taling in consideration anticipated 25% mortality)

Starting N (Total) - 60

Test siRNA: SERPINHI compounds disclosed herein.

TABLE 8: Groups:

No	BLM dose, mg/kg BW, in 0.1 ml saline	BLM adm. route	BLM regime	siRNA dose, mg/kg BW	siRNA adm route	siRNA regime	Termination post last siRNA adm	N (before siRNA administration)
1	0.75	I.P.	dd 0, 2, 4, 6	4.5	I.V.	2 daily	2h	4
2	0.75	I.P.	dd 0, 2, 4, 6	4.5	I.V.	2 daily	24 h	4
3	0.75	I.P.	dd 0, 2, 4, 6	2.25	I.T.	2 daily	2 h	4
4	0.75	I.P.	dd 0, 2, 4, 6	2.25	I.T.	2 daily	24 h	4
5		intact	n/a	4.5	I.V.	2 daily	2h	4
6		intact	n/a	4.5	I.V.	2 daily	24h	4
7		intact	n/a	2.25	I.T.	2 daily	2h	4
8		intact	n/a	2.25	I.T.	2 daily	24 h	4
9	0.75	I.P.	dd 0, 2, 4, 6	n/a	I.V. vehicle	2 daily	2h	3
10	0.75	I.P.	dd 0, 2, 4, 6	n/a	I/T/vehi cle	2 daily	24 h	3
11		Intact	n/a	n/a	intact	n/a	Any time	3

[0478] Bleomycin-induced pulmonary fibrosis. Pulmonary fibrosis of 12-wk-old female C57BL/6 mice will be induced by intraperitoneal instillation of bleomycin chlorate: 0.75 mg/kg body weight dissolved in 0.1 ml of saline every other day for 7 days, on days 0, 2, 4, and 6.

[0479] Pilot evaluation of the establishment of fibrosis. The mice (N=30) are subjected to BLM treatment in groups, to allow for a one week time interval between the first treated group (N=5) and the rest of the animals. On day 14, two mice from the first group are sacrificed and the lungs harvested for the fast HE stain and quick histopathological evaluation of fibrosis. When lung fibrosis is confirmed, the remaining rats are sorted into the groups and treated with siRNA on Day 14 after the first BLM treatment. In case that no sufficient fibrosis develops in the lungs by day 14, the remaining mice from the first treated group are sacrificed on day 21, followed by quick histopathology evaluation of fibrosis. The rest of the animals are treated with test siRNA complex starting from day 21 after the BLM treatment.

[0480] siRNA administration. On day 14 or day 21 after the first BLM administration (TBD during the study, based on pilot evaluation of establishment of fibrosis), the animals are group sorted, according to BW. The animals from groups 1 and 2 are administered intravenously (tail vein injection) with siRNA/vitA/Coatsome complex, at an siRNA concentration of 4.5 mg/kg BW. Intact animals of the same age (Groups 5 and 6) are treated in the same manner. BLM treated animals (Group 9) will be used as vitA-coatsome vehicle control). In 24 hours, the injection is repeated to all the animals, as above.

[0481] The BLM animals from the groups 3 and 4, and intact mice from groups 7 and 8 are anesthetized with isoflurane and subjected to intratracheal instillation of 2.25 mg/kg BW siRNA formulated in vitA-loaded liposomes. Mice from the BLM group 10 are administered with vitA/Coatsome vehicle only. The intratracheal instillation is repeated after 24 hours.

[0482] Study termination. The animals from the groups 1, 3, 5, 7, 9 are sacrificed at 2 hours after the second siRNA complex injection or instillation. The animals from the groups 2, 4, 6, 8, 10 are sacrificed at 24 hours after the second siRNA complex injection or instillation.

[0483] Upon animals sacrifice, the mice are perfused transcardially with 10% neutral buffered formalin. The lungs are inflated with 0.8-1.0 ml of 10% NBF, and the trachea ligated. The lungs are excised and fixed for 24 h in 10 % NBF. The liver is harvested from each animal and fixed in 10% NBF for 24 h.

[0484] *Sectioning and evaluation.* Consequent sections are prepared from the lungs and livers. First consequent section are stained with hematoxylin and eosin for assessment of lung and liver morphology, second section are stained with Sirius Red (trichrome) to identify collagen. The third consequent sections are subjected to in situ hybridization (ISH) for detection of siRNA.

EXAMPLE 22: *In vivo* anti-pulmonary-fibrosis activity of siRNA-containing VA-bound liposome

(1) Induction of pulmonary fibrosis and administration of drug

[0485] Male S-D rats (8 rats/group, 8 weeks old, Charles River Laboratories Japan, Inc.) were administered once with 0.45 mg bleomycin (BLM) dissolved in 0.1 mL of saline into the lung by intratracheal cannulation (MicroSprayer, Penn-Century, Inc.) under anesthesia, to produce a bleomycin pulmonary fibrosis model. With this method, a significant fibrosis occurs in the lung generally after approximately 2 weeks. The liposome formulation (1.5 mg/kg as an amount of siRNA, 1 mL/kg in volume, i.e., 200 μ L for a rat of 200 g) or PBS (1 mL/kg in volume) was administered to the rats via the tail vein, starting from the 2 weeks after the bleomycin administration, for total of ten times (every other day). The rats were sacrificed at two days post last treatment, histological investigation of the lung tissue was performed (see FIG. 30). One way ANOVA and Bonferroni multi comparison test was used for the evaluation of statistically-significant difference.

[0486] The composition of the liposome was HEDC/S-104/DOPE/Cholesterol/PEG-DMPE/diVA-PEG-diVA (20:20:30:25:5:2 Molar %). Details of siRNA were as follows:

S strand: 5'-idAB-rG-rA-rG-rA-rC-rA-rC-rA-rU-rG-rG-rU-rG-25rC-25rU-25rA-25rU-25rA-C3-P-3'

GS strand: 5'-mU-rA-mU-rA-mG-rC-25rA-rC-mC-rC-mA-rU-mG-rU-mG-rU-mC-rU-mC-C3-C3-3'

wherein: rX represents ribonucleotides; mX represents 2'-O-methyl ribonucleotides; 25rX represents ribonucleotides with 2'-5' linkages; C3 represents a 1,3-propanediol spacer; idAB represents inverted 1,2-dideoxy-D-ribose; P represents a phosphate group on the 3'-terminus. The 3'-terminus C3 is introduced by support-loaded 1,3-propanediol spacer. The 3'-terminus phosphate group (P) is introduced by the use of support-loaded diethyl sulfonyl (Pi) spacer.

(2) Histological investigation

[0487] A part of the removed lung was formalin-fixed in accordance with a routine method, and subjected to azan staining (azocarmine, aniline blue orange G solution). As shown by the results of the azan staining in FIG. 31, in the PBS administration group (Disease), a noticeable fibrotic image characterized by enlargement of interstice due to a large quantity of blue-stained collagenous fibrils was observed, whereas in the formulation administration group (Treatment), fibrosis were apparently suppressed.

[0488] As shown by the results of histological scoring (T. Ashcroft score) in FIG. 32, in the formulation administration group (Treatment), fibrosis score was significantly decreased.

EXAMPLE 23: *In-vivo* Reduction of HSP47 mRNA (DMN Model)

[0489] *In vivo* activity of the HSP47 liposomes of Example 22 was evaluated in the short-term liver damage model (referred to as the Quick Model). In this model, the short-term liver damage induced by treatment with a hepatotoxic agent such as dimethylnitrosamine (DMN) is accompanied by the elevation of HSP47 mRNA levels. To induce these changes, male Sprague-Dawley rats were injected intraperitoneally with DMN on six consecutive days. At the end of the DMN treatment period, the animals were randomized to groups based upon individual animal body weight. The liposome sample was administered as a single IV dose (0.375 or 0.75 mg/kg, reflecting the siRNA dose) one hour after the last injection of DMN. One day later, liver lobes were excised and both HSP47 and GAPDH mRNA levels were determined by quantitative RT-PCR (TaqMan) assay. HSP47 mRNA levels were normalized to GAPDH levels. As shown in Fig. 33, robust and dose-dependent mRNA reduction for HSP47 was detected in liver. After a single dose of 0.75 mg/kg of ND-L02-0101, 80% reduction of HSP47 mRNA was observed. Even at the lower dose of 0.375 mg/kg, significant knockdown of 40% was observed.

[0490] Applicants reserve the right to physically incorporate into this application any and all materials and information from any such articles, patents, patent applications, or other physical and electronic documents.

[0491] The present description teaches one skilled in the art to test various combinations and/or substitutions of chemical modifications described herein toward generating nucleic acid constructs with improved activity for mediating RNAi activity. Such improved activity can include improved stability, improved bioavailability, and/or improved activation of cellular responses mediating RNAi. Therefore, the specific embodiments described herein are not limiting and one skilled in the art can readily appreciate that specific combinations of the modifications described herein can be tested without undue experimentation toward identifying nucleic acid molecules with improved RNAi activity.

[0492] The descriptions illustratively described herein may suitably be practiced in the absence of any element or elements, limitation or limitations, not specifically disclosed herein. Thus, for example, the terms "a" and "an" and "the" and similar referents in the context of describing the description (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. The terms "comprising", "having", "including", "containing", etc. shall be read expansively and

without limitation (e.g., meaning "including, but not limited to,"). Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g., "such as") provided herein, is intended merely to better illuminate the description and does not pose a limitation on the scope of the description unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the description. Additionally, the terms and expressions employed herein have been used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but it is recognized that various modifications are possible within the scope of the description claimed. Thus, it should be understood that although the present description has been specifically disclosed by preferred embodiments and optional features, modification and variation of the descriptions embodied therein herein disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this description.

[0493] The description has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure also form part of the description. This includes the generic description of the description with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein. Other embodiments are within the following claims. In addition, where features or aspects of the description are described in terms of Markush groups, those skilled in the art will recognize that the description is also thereby described in terms of any individual member or subgroup of members of the Markush group.

[0494] The following items are also described:

1. 1. A pharmaceutical composition comprising a drug carrier and a double-stranded nucleic acid molecule, wherein the drug carrier comprises a stellate cell-specific amount of a retinoid, and wherein the double-stranded nucleic acid molecule comprises the structure :

5' (N)_x-Z 3' (antisense strand)

3' Z'-(N')_y-Z" 5' (sense strand)

wherein each of N and N' is a nucleotide which may be unmodified or modified, or an unconventional moiety;

wherein each of (N)_x and (N')_y is an oligonucleotide in which each consecutive N or N' is joined to the next N or N' by a covalent bond;

wherein each of Z and Z' is independently present or absent, but if present independently includes 1-5 consecutive nucleotides or non-nucleotide moieties or a combination thereof covalently attached at the 3'-terminus of the strand in which it is present;

wherein Z" may be present or absent, but if present is a capping moiety covalently attached at the 5'-terminus of (N')_y;

wherein each of x and y is independently an integer between 18 and 40;

wherein the sequence of (N')_y has complementary to the sequence of (N)_x; and

wherein (N)_x includes an antisense sequence to the mRNA coding sequence for human hsp47 exemplified by SEQ ID NO:1.

2. 2. The pharmaceutical composition of item 1, wherein the siRNA comprises oligonucleotide sequences selected from the group consisting of the sequences SEQ ID NOS: 60 and 127 (SERPINH1_2), SEQ ID NOS: 98 and 165 (SERPINH1_45a), and SEQ ID NOS: 101 and 168 (SERPINH1_51).
3. 3. The pharmaceutical composition according to item 1, wherein the retinoid consists of a molecule selected from the group consisting of vitamin A, retinoic acid, retinal, tretinoin, adapalene, retinol palmitate, and fenretinide (4-HPR).
4. 4. The pharmaceutical composition according to item 1, wherein the drug carrier further comprises a polyethylene glycol molecule-conjugated retinoid molecule.
5. 5. The pharmaceutical composition according to item 4, wherein the polyethylene glycol molecule-conjugated retinoid molecule is diVA-PEG-diVA.
6. 6. The pharmaceutical composition according to item 1, wherein the retinoid is at least partially exposed on the exterior of the drug carrier before the drug carrier reaches the stellate cell.
7. 7. The pharmaceutical composition according to item 1, wherein the drug carrier is in a form selected from the group consisting of a polymer micelle, a liposome, an emulsion, a microsphere, and nanosphere.
8. 8. The pharmaceutical composition according to item 7, wherein the drug carrier is in a form of a lipid vesicle comprising a bilayer of lipid molecules.
9. 9. The pharmaceutical composition according to item 8, wherein the retinoid is 0.2 wt% to 20 wt% of the lipid molecules.
10. 10. The pharmaceutical composition according to item 8, wherein the lipid molecules comprise HEDC.
11. 11. The pharmaceutical composition according to item 10, wherein the lipid molecules comprise S104.
12. 12. The pharmaceutical composition according to item 1, wherein the double-stranded nucleic acid molecule is exposed on the exterior surface of the lipid vesicle.
13. 13. The pharmaceutical composition according to item 1, wherein the double-stranded nucleic acid molecule is encapsulated by the lipid vesicle.
14. 14. The pharmaceutical composition according to item 13, wherein the double-stranded nucleic acid molecule is resistant to nucleases.
15. 15. The pharmaceutical composition according to any of items 1-14, wherein one strand of the double-stranded nucleic acid molecule comprises SEQ ID NO:127 and the second strand comprises SEQ ID NO:60.
16. 16. The pharmaceutical composition according to item 15, wherein the antisense strand comprises SEQ ID NO:127, and the sense strand comprises SEQ ID NO:60.
17. 17. The pharmaceutical composition according to item 16, wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides; a 2'-5'-ribonucleotide in at least one of positions 1, 5, 6, or 7; and a non-nucleotide moiety covalently attached to the 3'-terminus; and wherein the sense strand further comprises at least one 2'-5'-ribonucleotide or 2'-O-methyl modified ribonucleotide; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus.
18. 18. The pharmaceutical composition according to item 16, wherein the antisense strand further comprises 2'-O-methyl modified

- ribonucleotides at positions 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
19. 19. The pharmaceutical composition according to item 16, wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides at positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; and a C3C3 non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises a C3Pi non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
 20. 20. The pharmaceutical composition according to item 16, wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides at positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a C3Pi non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
 21. 21. The pharmaceutical composition according to item 16, wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides at positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises five consecutive 2'-5'-ribonucleotides in the 3'-terminal positions 15, 16, 17, 18, and 19; a C3Pi non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
 22. 22. The pharmaceutical composition according to item 16, wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 3, 5, 9, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 7, 13, 16 and 18; a 2'-5'-ribonucleotide at position 9; a C3 non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic moiety covalently attached at the 5'-terminus.
 23. 23. The pharmaceutical composition according to any of items 1-14, wherein one strand of the double-stranded nucleic acid molecule comprises SEQ ID NO:98 and the second strand comprises SEQ ID NO:165.
 24. 24. The pharmaceutical composition according to item 23, wherein the sense strand comprises SEQ ID NO:98 and the antisense strand comprises SEQ ID NO:165.
 25. 25. The pharmaceutical composition according to item 24, wherein the sense strand further comprises 2'-5'-ribonucleotides in positions at or near the 3'-terminus; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and the antisense strand further comprises a 2'-O-methyl modified ribonucleotide; a 2'-5'-ribonucleotide in at least one of positions 5, 6 or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus.
 26. 26. The pharmaceutical composition according to item 24, wherein the sense strand further comprises 2'-5'-ribonucleotides in positions 15, 16, 17, 18, and 19; a C3-OH 3' moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 2, 4, 6, 8, 11, 13, 15, 17, and 19; a 2'-5'-ribonucleotide in position 7; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.
 27. 27. The pharmaceutical composition according to any of items 1-14, wherein one strand of the double-stranded nucleic acid molecule comprises SEQ ID NO:101 and the second strand comprises SEQ ID NO:168.
 28. 28. The pharmaceutical composition according to item 1, wherein the sense strand comprises SEQ ID NO:101 and the antisense strand comprises SEQ ID NO:168.
 29. 29. The pharmaceutical composition according to item 28, wherein the sense strand further comprises 2'-O-methyl modified pyrimidine ribonucleotides; an optional 2'-5'-ribonucleotide in one of position 9 or 10; a non-nucleotide moiety covalently attached at the 3'-terminus; and a cap moiety covalently attached at the 5'-terminus; and the wherein antisense strand further comprises a 2'-O-methyl modified ribonucleotide; a 2'-5'-ribonucleotide in at least one of positions 5, 6, or 7; and a non-nucleotide moiety covalently attached at the 3'-terminus.
 30. 30. The pharmaceutical composition according to item 28, wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 4, 11, 13, and 17; a 2'-5'-ribonucleotide in position 9; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.
 31. 31. The pharmaceutical composition according to item 28, wherein sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 4, 11, 13, and 17; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 4, 8, 13 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH moiety covalently attached at the 3'-terminus.
 32. 32. The pharmaceutical composition according to item 28, wherein the sense strand further comprises 2'-O-methyl modified ribonucleotides in positions 2, 4, 11, 13, and 17; a C3OH non-nucleotide moiety covalently attached at the 3'-terminus; and an inverted abasic deoxyribonucleotide moiety covalently attached at the 5'-terminus; and wherein the antisense strand further comprises 2'-O-methyl modified ribonucleotides in positions 1, 4, 8, 11 and 15; a 2'-5'-ribonucleotide in position 6; and a C3Pi-C3OH non-nucleotide moiety covalently attached at the 3'-terminus.
 33. 33. The pharmaceutical composition according to any of items 1-32, wherein the double-stranded nucleic acid molecule reduces hsp47 expression in the stellate cell.
 34. 34. The pharmaceutical composition according to any of items 1-32, wherein the composition prevents or treats a disease associated with hsp47.
 35. 35. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats a disease selected from the group consisting of fibrosis of organs and tissues including liver, lung, kidney, pancreas, skin, heart, brain, intestine, colon, vocal cords, peritoneum, eye, muscle, bone marrow, bone, ovary, ureter, and uterus; and cancer and cancer-associated fibrotic tissues.
 36. 36. The pharmaceutical composition according to any of items 1-34, wherein the pharmaceutical composition prevents or treats liver fibrosis.
 37. 37. The pharmaceutical composition according to any of items 1-34, wherein the pharmaceutical composition prevents or treats pulmonary

- fibrosis.
38. 38. The pharmaceutical composition according to any of items 1-34, wherein the pharmaceutical composition prevents or treats kidney fibrosis.
39. 39. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats peritoneal fibrosis.
40. 40. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats chronic hepatic damage.
41. 41. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats fibrillogenesis.
42. 42. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats liver cirrhosis due to hepatitis C post-liver transplant or non-alcoholic steatohepatitis (NASH); idiopathic pulmonary fibrosis; radiation pneumonitis leading to pulmonary fibrosis; diabetic nephropathy; peritoneal sclerosis associated with continual ambulatory peritoneal dialysis (CAPD) or ocular cicatricial pemphigoid.
43. 43. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats liver cirrhosis due to hepatitis C post-liver transplant or non-alcoholic steatohepatitis (NASH).
44. 44. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats idiopathic pulmonary fibrosis.
45. 45. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats radiation pneumonitis leading to pulmonary fibrosis.
46. 46. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats diabetic nephropathy.
47. 47. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats peritoneal sclerosis associated with continual ambulatory peritoneal dialysis (CAPD).
48. 48. The pharmaceutical composition according to any of items 1-34, wherein the composition prevents or treats ocular cicatricial pemphigoid.
49. 49. A method for treating or preventing a disease, the method comprising administering an effective amount of the pharmaceutical composition according to any of items 1-48 to a subject in need thereof.
50. 50. The method of item 49 wherein the disease is selected from the group consisting of hepatitis, hepatic fibrosis, hepatic cirrhosis, liver cancer, pancreatitis, pancreatic fibrosis, pancreatic cancer, vocal cord scarring, vocal cord mucosal fibrosis, and laryngeal fibrosis.
51. 51. The method of item 49, wherein the disease is hepatitis or hepatic fibrosis.
52. 52. The method of item 49, wherein the disease associated with chronic hepatitis C virus (HCV) infection.
53. 53. The method according to item 49, wherein the preparation is parenterally administered.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US61497447B \[0001\]](#)
- [US61494832B \[0001\]](#)
- [EP1842557A \[0007\]](#)
- [WO0175164A \[0220\] \[0220\]](#)
- [WO0044895A \[0221\]](#)
- [WO0136646A \[0221\]](#)
- [WO9932619A \[0221\]](#)
- [WO0001846A \[0221\]](#)
- [WO0129058A \[0221\]](#)
- [WO9907409A \[0221\]](#)
- [WO0044914A \[0221\]](#)
- [WO0047599A \[0229\]](#)
- [WO9914226A \[0229\]](#)
- [WO9839952A \[0229\]](#)
- [WO04083430A \[0229\]](#)
- [WO2006047842A \[0232\]](#)
- [US6602858B \[0233\] \[0233\]](#)
- [EP0586520A \[0237\]](#)
- [EP0618925A \[0237\]](#)
- [US10427160B \[0246\] \[039.1\]](#)
- [US10201394B \[0246\]](#)
- [WO8902439A \[0248\]](#)
- [WO9506731A \[0248\]](#)
- [WO9511910A \[0248\]](#)
- [EP586520A \[0251\]](#)
- [EP618925A \[0251\]](#)

- [US7452987B \[0264\] \[0264\]](#)
- [US07081836W \[0295\]](#)
- [US20050244858A \[0297\]](#)
- [US4501729A \[0387\]](#)
- [US20040037780A \[0387\]](#)
- [US6592904E \[0387\]](#)
- [US6582728E \[0387\]](#)
- [US6565885E \[0387\]](#)
- [WO08132723A \[0387\]](#)
- [US6180613E \[0389\]](#)
- [WO04013280A \[0389\]](#)
- [US6586524E \[0390\]](#)
- [US6528631E \[0391\]](#)
- [US6335434E \[0391\]](#)
- [US6235886E \[0391\]](#)
- [US6153737A \[0391\]](#)
- [US5214136A \[0391\]](#)
- [US5138045A \[0391\]](#)
- [US5624803A \[0398\]](#)
- [WO9618736A \[0398\]](#)
- [WO9323669A \[0399\]](#)
- [WO9402595A \[0399\]](#)

Non-patent literature cited in the description

- [MADRO et al.](#) Med Sci Monit., 2004, vol. 10, RA166-70 [0003]
- [JASTER](#) Mol Cancer., 2004, vol. 6, 26- [0003]
- [FALLOWFIELD et al.](#) Expert Opin Ther Targets, 2004, vol. 8, 423-35 [0004]
- [PINZANI et al.](#) Dig Liver Dis., 2004, vol. 36, 231-42 [0004]
- [SATO et al.](#) Nature Biotechnology, 2008, vol. 26, 4431-442 [0006]
- [TURNER et al.](#) CSH Symp. Quant. Biol., 1987, vol. LII, 123-133 [0205]
- [FRIER et al.](#) Proc. Nat. Acad. Sci. USA, 1986, vol. 83, 9373-9377 [0205]
- [TURNER et al.](#) J. Am. Chem. Soc., 1987, vol. 109, 3783-3785 [0205]
- [TAKEI et al.](#) JBC, 2002, vol. 277, 2623800-06 [0214]
- [ZAMORE et al.](#) Cell, 2000, vol. 101, 25-33 [0218] [0219] [0219] [0221]
- [FIRE et al.](#) Nature, 1998, vol. 391, 806- [0218] [0220]
- [HAMILTON et al.](#) Science, 1999, vol. 286, 950-951 [0218]
- [LIN et al.](#) Nature, 1999, vol. 402, 128-129 [0218]
- [SHARP](#) Genes & Dev., 1999, vol. 13, 139-141 [0218]
- [STRAUSS](#) Science, 1999, vol. 286, 886- [0218]
- [BASS](#) Cell, 2000, vol. 101, 235- [0219]
- [BERSTEIN et al.](#) Nature, 2001, vol. 409, 363- [0219]
- [ELBASHIR et al.](#) Genes Dev., 2001, vol. 15, 188- [0219] [0219]
- [HUTVAGNER et al.](#) Science, 2001, vol. 293, 834- [0219]
- [BAHRAMIANZARBL](#) Molecular and Cellular Biology, 1999, vol. 19, 274-283 [0220]
- [WIANNYGOETZ](#) Nature Cell Biol., 1999, vol. 2, 70- [0220]
- [ELBASHIR et al.](#) Nature, 2001, vol. 411, 494- [0220]
- [ELBASHIR et al.](#) EMBO J., 2001, vol. 20, 6877- [0220]
- [BASS](#) Nature, 2001, vol. 411, 428-429 [0221]
- [ELBASHIR et al.](#) Nature, 2001, vol. 411, 494-498 [0221]
- [ALLSHIRE](#) Science, 2002, vol. 297, 1818-1819 [0221]
- [VOLPE et al.](#) Science, 2002, vol. 297, 1833-1837 [0221]
- [JENUWEIN](#) Science, 2002, vol. 297, 2215-2218 [0221]
- [HALL et al.](#) Science, 2002, vol. 297, 2232-2237 [0221]
- [HUTVAGNERZAMORE](#) Science, 2002, vol. 297, 2056-60 [0221]
- [MCMANUS et al.](#) RNA, 2002, vol. 8, 842-850 [0221]
- [REINHART et al.](#) Gene & Dev., 2002, vol. 16, 1616-1626 [0221]
- [REINHARTBARTEL](#) Science, 2002, vol. 297, 1831- [0221]
- [SANGER](#) Principles of Nucleic Acid Structure Springer-Verlag 1984 0000 [0223]
- [Nucleic Acids Symposium Series No. 52](#), 2008, 133-134 [0230]
- [ALLART et al.](#) Nucleosides & Nucleotides, 1998, vol. 17, 1523-1526 [0232]
- [PEREZ-PEREZ et al.](#) Bioorg. and Medicinal Chem Letters, 1996, vol. 6, 1457-1460 [0232]
- [GOLD et al.](#) Annu. Rev. Biochem., 1995, vol. 64, 763- [0247]
- [BRODYGOLDJ.](#) Biotechnol., 2000, vol. 74, 5- [0247]

- **SUN** Curr. Opin. Mol. Ther., 2000, vol. 2, 100- [\[0247\]](#)
- **KUSSERJ.** Biotechnol., 2000, vol. 74, 27- [\[0247\]](#)
- **HERMANNPATEL** Science, 2000, vol. 287, 820- [\[0247\]](#)
- **JAYASENA** Clinical Chemistry, 1999, vol. 45, 1628- [\[0247\]](#)
- **SEELAKAISERN** Nucleic Acids Res., 1990, vol. 18, 6353- [\[0248\]](#)
- Nucleic Acids Res., 1987, vol. 15, 3113- [\[0248\]](#)
- **CLOADSCHEPARTZ** J. Am. Chem. Soc., 1991, vol. 113, 6324- [\[0248\]](#)
- **RICHARDSONSCHEPARTZ** J. Am. Chem. Soc., 1991, vol. 113, 5109- [\[0248\]](#)
- **MA et al.** Nucleic Acids Res., 1993, vol. 21, 2585- [\[0248\]](#)
- Biochemistry, 1993, vol. 32, 1751- [\[0248\]](#)
- **DURAND et al.** Nucleic Acids Res., 1990, vol. 18, 6353- [\[0248\]](#)
- **MCCURDY et al.** Nucleosides & Nucleotides, 1991, vol. 10, 287- [\[0248\]](#)
- **JSCHKE et al.** Tetrahedron Lett., 1993, vol. 34, 301- [\[0248\]](#)
- **ONO et al.** Biochemistry, 1991, vol. 30, 9914- [\[0248\]](#)
- **FERENTZVERDINE** J. Am. Chem. Soc., 1991, vol. 113, 4000- [\[0248\]](#)
- **NAGATA** Trends Biochem Sci, 1996, vol. 21, 22-6 [\[0313\]](#)
- **RAZZAQUE et al.** Contrib Nephrol 2005, 2005, vol. 148, 57-69 [\[0313\]](#)
- **KOIDE et al.** J. Biol. Chem., 2006, vol. 281, 3432-38 [\[0313\]](#)
- **LEIVO et al.** Dev. Biol., 1980, vol. 76, 100-114 [\[0313\]](#)
- **MASUDA et al.** J. Clin. Invest., 1994, vol. 94, 2481-2488 [\[0313\]](#)
- **MASUDA et al.** Cell Stress Chaperones, 1998, vol. 3, 256-264 [\[0313\]](#)
- **KOIDE et al.** J Biol Chem, 1999, vol. 274, 34523-26 [\[0313\]](#)
- **MASUDA et al.** J Clin Invest, 1994, vol. 94, 2481-8 [\[0313\]](#)
- **RAZZAQUE et al.** Virchows Arch, 1998, vol. 432, 455-60 [\[0313\]](#)
- **KAKUGAWA et al.** Eur Respir J, 2004, vol. 24, 57-65 [\[0313\]](#)
- **MORIYAMA et al.** Kidney Int, 1998, vol. 54, 110-19 [\[0313\]](#)
- **SATO et al.** Nat Biotechnol, 2008, vol. 26, 431-442 [\[0314\]](#) [\[0324\]](#)
- **CHEN et al.** Br J Dermatol, 2007, vol. 156, 1188-1195 [\[0314\]](#) [\[0324\]](#)
- **WANG et al.** Plast. Reconstr Surg, 2003, vol. 111, 1980-7 [\[0314\]](#)
- **WANG et al.** Plast Reconstr Surg, 2003, vol. 111, 1980-7 [\[0324\]](#)
- **BERGERON et al.** Trends Biochem. Sci., 1994, vol. 19, 124-128 [\[0329\]](#)
- **HERBERT et al.** Cold Spring Harb. Symp. Quant. Biol., 1995, vol. 60, 405-415 [\[0329\]](#)
- **CORDEIRO MF et al.** Human anti-transforming growth factor-beta2 antibody: a new glaucoma anti-scarring agent Invest Ophthalmol Vis Sci., 1999, vol. 40, 102225-34 [\[0341\]](#)
- **LASKY et al.** Environ Health Perspect, 2000, vol. 108, 751-62 [\[0355\]](#)
- **MANABE et al.** Circ Res., 2002, vol. 13, 1103-13 [\[0358\]](#)
- **MOLINA V et al.** Harefuah, 2002, vol. 141, 973-81009- [\[0364\]](#)
- **YU et al.** Curr Opin Pharmacol., 2002, vol. 2, 2177-81 [\[0364\]](#)
- **KEANE et al.** Am J Kidney Dis., 2003, vol. 41, 22-5 [\[0364\]](#)
- **BOHLE et al.** Pathol Res Pract., 1989, vol. 185, 421-40 [\[0364\]](#)
- **KIKKAWA et al.** Kidney Int Suppl., 1997, vol. 62, 39-40 [\[0364\]](#)
- **BATALLER et al.** Semin Liver Dis., 2001, vol. 21, 437-51 [\[0364\]](#)
- **GROSS et al.** N Engl J Med., 2001, vol. 345, 517-25 [\[0364\]](#)
- **FROHLICH** Am J Hypertens, 2001, vol. 14, 194S-199S [\[0364\]](#)
- **FRIEDMAN** J Hepatol., 2003, vol. 38, 38-53 [\[0364\]](#)
- **ALBANIS et al.** Curr Gastroenterol Rep., 2003, vol. 5, 48-56 [\[0364\]](#)
- **WEBER** Curr Opin Cardiol., 2000, vol. 15, 264-72 [\[0364\]](#)
- **G. P. MOSS** Biochemical Nomenclature and Related Documents Portland Press 19920000247-251 [\[0366\]](#)
- **SOMMER et al.** Antisense Nuc. Acid Drug Dev., 1998, vol. 8, 75- [\[0388\]](#)
- **EPA et al.** Antisense Nuc. Acid Drug Dev., 2000, vol. 10, 469- [\[0388\]](#)
- **BROADDUS et al.** J. Neurosurg, 1998, vol. 88, 734- [\[0388\]](#)
- **KARLE et al.** Eur. J. Pharmacol., 1997, vol. 340, 153- [\[0388\]](#)
- **BANNAI et al.** Brain Research, 1998, vol. 784, 304- [\[0388\]](#)
- **RAJAKUMAR et al.** Synapse, 1997, vol. 26, 199- [\[0388\]](#)
- **WU-PONG et al.** BioPharm, 1999, vol. 12, 32- [\[0388\]](#)
- **BANNAI et al.** Brain Res. Protoc., 1998, vol. 3, 83- [\[0388\]](#)
- **SIMANTOV et al.** Neuroscience, 1996, vol. 74, 39- [\[0388\]](#)
- **OGRIS et al.** AAPA PharmSci, 2001, vol. 3, 1-11 [\[0390\]](#)
- **FURGESON et al.** Bioconjugate Chem., 2003, vol. 14, 840-847 [\[0390\]](#)
- **KUNATH et al.** Pharm Res, 2002, vol. 19, 810-17 [\[0390\]](#)
- **CHOI et al.** Bull. Korean Chem. Soc., 2001, vol. 22, 46-52 [\[0390\]](#)
- **BETTINGER et al.** Bioconjugate Chem., 1999, vol. 10, 558-561 [\[0390\]](#)
- **PETERSON et al.** Bioconjugate Chem., 2002, vol. 13, 845-54 [\[0390\]](#)
- **ERBACHER et al.** J Gene Med, 1999, vol. 1, 1-18 [\[0390\]](#)
- **GODBEY et al.** PNAS, 1999, vol. 96, 5177-81 [\[0390\]](#)
- **GODBEY et al.** J Controlled Release, 1999, vol. 60, 149-60 [\[0390\]](#)
- **DIEBOLD et al.** J Biol Chem, 1999, vol. 274, 19087-94 [\[0390\]](#)
- **THOMAS et al.** PNAS, 2002, vol. 99, 14640-45 [\[0390\]](#)

- PAUL et al.Nature Biotech, 2002, vol. 19, 505- [\[0394\]](#) [\[0396\]](#)
- MIYAGISHI et al.Nature Biotech, 2002, vol. 19, 497- [\[0394\]](#)
- LEE et al.Nature Biotech, 2002, vol. 19, 500- [\[0394\]](#) [\[0396\]](#)
- NOVINA et al.Nature Med, 2002, [\[0394\]](#) [\[0396\]](#)
- MIYAGISHITAIRA Nature Biotech, 2002, vol. 19, 497- [\[0396\]](#)
- ELROY et al.PNAS, 1990, vol. 87, 6743-47 [\[0398\]](#)
- GAO et al.Nucleic Acids Res, 1993, vol. 21, 2867-72 [\[0398\]](#)
- LIEBER et al.Methods Enzymol., 1993, vol. 217, 47-66 [\[0398\]](#)
- ZHOU et al.Mol. Cell. Biol., 1990, vol. 10, 4529-37 [\[0398\]](#)
- KASHANI-SABET et al.Antisense Res. Dev., 1992, vol. 2, 3-15 [\[0398\]](#) [\[0399\]](#)
- OJWANG et al.PNAS, 1992, vol. 89, 10802-06 [\[0398\]](#) [\[0399\]](#)
- CHEN et al.Nucleic Acids Res., 1992, vol. 20, 4581-89 [\[0398\]](#) [\[0399\]](#)
- YU et al.PNAS, 1993, vol. 90, 6340-44 [\[0398\]](#)
- L'HUILLIER et al.EMBO J., 1992, vol. 11, 4411-18 [\[0398\]](#)
- LISZIEWICZ et al.PNAS, 1993, vol. 90, 8000-04 [\[0398\]](#)
- THOMPSON et al.Nucleic Acids Res., 1995, vol. 23, 2259- [\[0398\]](#) [\[0399\]](#)
- SULLENGER et al.Science, 1993, vol. 262, 1566- [\[0398\]](#)
- NOONBERG et al.Nucleic Acid Res., 1994, vol. 22, 2830- [\[0398\]](#)
- GOOD et al.Gene Ther., 1997, vol. 4, 45- [\[0398\]](#)
- IZANTWEINTRAUB Science, 1985, vol. 229, 345- [\[0399\]](#)
- MCGARRY LINDQUIST PNAS, 1986, vol. 83, 399- [\[0399\]](#)
- SCANLON et al.PNAS, 1991, vol. 88, 10591-95 [\[0399\]](#)
- DROPULIC et al.J. Virol., 1992, vol. 66, 1432-41 [\[0399\]](#)
- WEERASINGHE et al.J. Virol., 1991, vol. 65, 5531-34 [\[0399\]](#)
- SARVER et al.Science, 1990, vol. 247, 1222-25 [\[0399\]](#)
- GOOD et al.Gene Therapy, 1997, vol. 4, 45- [\[0399\]](#)
- OHKAWA et al.Nucleic Acids Symp. Ser., 1992, vol. 27, 15-6 [\[0399\]](#)
- TAIRA et al.Nucleic Acids Res., 1991, vol. 19, 5125-30 [\[0399\]](#)
- VENTURA et al.Nucleic Acids Res., 1993, vol. 21, 3249-55 [\[0399\]](#)
- CHOWRIRA et al.J. Biol. Chem., 1994, vol. 269, 25856- [\[0399\]](#)
- PETERKOFISKY et al.Biochemistry, 1971, vol. 10, 988-94 [\[0423\]](#)
- JEZEQUEL et al.J Hepatol., 1987, vol. 5, 174-81 [\[0428\]](#)
- MCCAFFERY et al.Nature, 2002, vol. 418, 38-39 [\[0428\]](#)
- Hepatology, 1998, vol. 28, 1247-52 [\[0428\]](#)
- QI Z et al.PNAS, 1999, vol. 96, 2345-49 [\[0429\]](#)
- UEKI et al.Nat Med., 1999, vol. 5, 226-30 [\[0430\]](#)
- SATO Y. et al.Nature Biotech., 2008, vol. 26, 431- [\[0450\]](#)
- LOTERSZTAJN S et al.Hepatic Fibrosis: Molecular Mechanisms and Drug Targets Annu Rev Pharmacol Toxicol., 2004, [\[0464\]](#)
- UCHIO K et al.Down-regulation of connective tissue growth factor and type I collagen mRNA expression by connective tissue growth factor antisense oligonucleotide during experimental liver fibrosis Wound Repair Regen, 2004, vol. 12, 160-6 [\[0464\]](#)
- XU XQ et al.Molecular classification of liver cirrhosis in a rat model by proteomics and bioinformatics Proteomics, 2004, vol. 4, 103235-45 [\[0464\]](#)
- SHERWOOD MB et al.J Glaucoma., 2004, vol. 13, 5407-12 [\[0465\]](#)
- MILLER MH et al.A new model of glaucoma filtering surgery in the rat Ophthalmic Surg., 1989, vol. 20, 5350-7 [\[0465\]](#)
- RETINA Fall-Winter, 1985, vol. 5, 4239-52 [\[0465\]](#)
- WIEDEMANN P et al.J Pharmacol Methods, 1984, vol. 12, 169-78 [\[0465\]](#)
- ZHOU et al.Invest Ophthalmol Vis Sci., 2002, vol. 43, 2293-300 [\[0466\]](#)
- WANG et al.Curr Eye Res., 2004, vol. 29, 51-58 [\[0466\]](#)
- MIZOBUCHI et al.J Heart Lung Transplantation, 2004, 23- [\[0472\]](#)
- KAZUHIRO et al.Am. J. Respir. Cell Mol Biol, 2001, vol. 25, 26-34 [\[0472\]](#)
- MOELLER et al.Int J Biochem Cell Biol, 2008, vol. 40, 362-82 [\[0475\]](#)
- CHUA et al.Am J Respir Cell Mol Biol, 2005, vol. 33, 9-13 [\[0476\]](#)
- ONUMA et al.J Exp Med, 2001, vol. 194, 147-56 [\[0476\]](#)
- YAMAGUCHI et al.Nature, 1988, vol. 336, 244-46 [\[0476\]](#)
- SWIDERSKI et al.Am J Pathol, 1998, vol. 152, 821-28 [\[0476\]](#)
- SHIMIZUKAWA et al.Am J Physiol Lung Cell Mol Physiol, 2003, vol. 284, L526-32 [\[0476\]](#)

Patentkrav**1. Farmaceutisk sammensætning omfattende:**

- (i) et dobbeltstrenget nukleinsyremolekyle og lægemiddel-bærer
 5 omfattende en blanding af et retinoid og et lipid, hvor det dobbeltstrengede nukleinsyremolekyle omfatter strukturen (A1):

(A1) 5' (N)_x-Z 3' (antisense-streng) 3' Z'-(N')_y-z" 5' (sense-
 streng)

10

hvor hver af N og N' er et nukleotid, som kan være ikke-modificeret eller modificeret, eller en ukonventionel del;

15

hvor hver af (N)_x og (N')_y er et oligonukleotid i hvilket hvert på hinanden følgende N eller N' er forbundet til det næste N eller N' med en kovalent binding;

20

hvor hver af Z og Z' er uafhængigt til stede eller fraværende, men hvis til stede uafhængigt inkluderer 1-5 på hinanden følgende nukleotider eller ikke-nukleotiddele eller en kombination deraf kovalent fastgjort ved den 3'-terminale ende af strengen i hvilken den er til stede;

hvor z" kan være til stede eller fraværende, men hvis til stede er en capping-del kovalent fastgjort ved den 5'-terminale ende af (N')_y;


hvor hver af x og y er uafhængigt et heltal mellem 18 og 40;

hvor sekvensen af (N')_y er komplementær til sekvensen af (N)_x;

25

hvor (N)_x inkluderer en antisense-sekvens til den kodende mRNA-sekvens for human hsp47 eksemplificeret i SEQ ID NO:1; og


hvor lipidet omfatter et lipid valgt fra gruppen bestående af



HEDC


2

og




eller

- (ii) et dobbeltstrenget nukleinsyremolekyle omfattende en sense-streng og en antisense-streng, hvor sense- og antisense-strengene er valgt fra oligonukleotiderne beskrevet som SERPINH1_2 (SEQ ID NOS: 60 og 127), SERPINH1_45a (SEQ ID NOS: 98 og 165), og SERPINH1_51 (SEQ ID NOS: 101 og 168), og en lægemiddel-bærer omfattende en blanding af en lipid-vesikel og et retinoid eller et retinoid-konjugat, hvor et lipid af lipid-vesiklen omfatter et lipid valgt fra gruppen bestående af



og



2. Den farmaceutiske sammensætning ifølge krav 1, hvor retinoid-konjugatet er et PEG-konjugat.
3. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor antisense-strengen omfatter SEQ ID NO:127, og sense-strengen omfatter SEQ ID NO:60; hvor antisense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider; et 2'-5'-ribonukleotid i mindst en af positionerne 1, 5, 6, eller 7; og en ikke-nukleotiddel kovalent fastgjort til den 3'-terminale ende; og hvor sense-strengen yderligere omfatter mindst et 2'-5'-ribonukleotid eller 2'-O-methyl modificeret ribonukleotid; en ikke-nukleotiddel kovalent fastgjort ved

den 3'-terminale ende; og en cap-del kovalent fastgjort ved den 5'-terminale ende.

4. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2,
5 hvor antisense-strengen omfatter SEQ ID NO:127, og sense-strengen omfatter
SEQ ID NO:60; hvor antisense-strengen yderligere omfatter 2'-O-methyl
modificerede ribonukleotider i positionerne 3, 5, 9, 11, 13, 15, 17, og 19; et 2'-5'-
ribonukleotid i position 7; og en ikke-nukleotiddel kovalent fastgjort ved den 3'-
terminale ende; og hvor sense-strengen yderligere omfatter fem på hinanden
10 følgende 2'-5'-ribonukleotider i de 3'-terminale positioner 15, 16, 17, 18, og 19;
en ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og en
inverteret abasisk del kovalent fastgjort ved den 5'-terminale ende.

5. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2,
15 hvor antisense-strengen omfatter SEQ ID NO:127, og sense-strengen omfatter
SEQ ID NO:60; hvor antisense-strengen yderligere omfatter 2'-O-methyl
modificerede ribonukleotider i positionerne 1, 3, 5, 9, 11, 13, 15, 17, og 19; og en
C3C3 ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og hvor
sense-strengen yderligere omfatter en C3Pi ikke-nukleotiddel kovalent fastgjort
20 ved den 3'-terminale ende; og en inverteret abasisk del kovalent fastgjort ved den
5'-terminale ende.

6. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2,
hvor antisense-strengen omfatter SEQ ID NO:127, og sense-strengen omfatter
25 SEQ ID NO:60; hvor antisense-strengen yderligere omfatter 2'-O-methyl
modificerede ribonukleotider i positionerne 1, 3, 5, 9, 11, 13, 15, 17, og 19; et 2'-
5'-ribonukleotid i position 7; og en C3Pi-C3OH ikke-nukleotiddel kovalent fastgjort
ved den 3'-terminale ende; og hvor sense-strengen yderligere omfatter fem på
hinanden følgende 2'-5'-ribonukleotider i de 3'-terminale positioner 15, 16, 17,
30 18, og 19; en C3Pi ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende;
og en inverteret abasisk del kovalent fastgjort ved den 5'-terminale ende.

7. Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2,
hvor antisense-strengen omfatter SEQ ID NO:127, og sense-strengen omfatter

SEQ ID NO:60; hvor antisense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 1, 3, 5, 9, 11, 13, 15, 17, og 19; et 2'-5'-ribonukleotid i position 7; og en C3Pi-C3OH ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og hvor sense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 7, 13, 16 og 18; et 2'-5'-ribonukleotid i position 9; en C3 ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og en inverteret abasisk del kovalent fastgjort ved den 5'-terminale ende.

- 10 **8.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor sense-strengen omfatter SEQ ID NO:98 og antisense-strengen omfatter SEQ ID NO:165; hvor sense-strengen yderligere omfatter 2'-5'-ribonukleotider i positionerne ved eller nær den 3'-terminale ende; en ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og en cap-del kovalent fastgjort ved den 5'-terminale ende; og hvor antisense-strengen yderligere omfatter et 2'-O-methyl modificeret ribonukleotid; et 2'-5'-ribonukleotid i mindst en af positionerne 5, 6 eller 7; og en ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende.

- 20 **9.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor sense-strengen omfatter SEQ ID NO:98 og antisense-strengen omfatter SEQ ID NO:165; hvor sense-strengen yderligere omfatter 2'-5'-ribonukleotider i positionerne 15, 16, 17, 18, og 19; en C3-OH 3'-del kovalent fastgjort ved den 3'-terminale ende; og en inverteret abasisk deoxyribonukleotiddel kovalent fastgjort ved den 5'-terminale ende; og hvor antisense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 2, 4, 6, 8, 11, 13, 15, 17, og 19; et 2'-5'-ribonukleotid i position 7; og en C3Pi-C3OH del kovalent fastgjort ved den 3'-terminale ende.

- 30 **10.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor sense-strengen omfatter SEQ ID NO:101 og antisense-strengen omfatter SEQ ID NO:168; hvor sense-strengen yderligere omfatter 2'-O-methyl modificerede pyrimidin-ribonukleotider; et eventuelt 2'-5'-ribonukleotid i en af position 9 eller 10; en ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og en cap-del kovalent fastgjort ved den 5'-terminale ende; og hvor

antisense-strengen yderligere omfatter et 2'-O-methyl modificeret ribonukleotid; et 2'-5'-ribonukleotid i mindst en af positionerne 5, 6, eller 7; og en ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende.

- 5 **11.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor sense-strengen omfatter SEQ ID NO:101 og antisense-strengen omfatter SEQ ID NO:168; hvor sense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 4, 11, 13, og 17; et 2'-5'-ribonukleotid i position 9; en C3OH ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale
- 10 ende; og en inverteret abasisk deoxyribonukleotiddel kovalent fastgjort ved den 5'-terminale ende; og hvor antisense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 1, 4, 8, 11 og 15; et 2'-5'-ribonukleotid i position 6; og en C3Pi-C3OH del kovalent fastgjort ved den 3'-terminale ende.

15

- 12.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor sense-strengen omfatter SEQ ID NO:101 og antisense-strengen omfatter SEQ ID NO:168; hvor sense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 4, 11, 13, og 17; en C3OH ikke-
- 20 nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og en inverteret abasisk deoxyribonukleotiddel kovalent fastgjort ved den 5'-terminale ende; og hvor antisense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 1, 4, 8, 13 og 15; et 2'-5'-ribonukleotid i position 6; og en C3Pi-C3OH del kovalent fastgjort ved den 3'-terminale ende.

25

- 13.** Den farmaceutiske sammensætning ifølge et hvilket som helst af kravene 1-2, hvor sense-strengen omfatter SEQ ID NO:101 og antisense-strengen omfatter SEQ ID NO:168; hvor sense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 2, 4, 11, 13, og 17; en C3OH ikke-
- 30 nukleotiddel kovalent fastgjort ved den 3'-terminale ende; og en inverteret abasisk deoxyribonukleotiddel kovalent fastgjort ved den 5'-terminale ende; og hvor antisense-strengen yderligere omfatter 2'-O-methyl modificerede ribonukleotider i positionerne 1, 4, 8, 11 og 15; et 2'-5'-ribonukleotid i position 6; og en C3Pi-C3OH ikke-nukleotiddel kovalent fastgjort ved den 3'-terminale ende.

DRAWINGS




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5





FIG. 6






FIG. 8




FIG. 9




FIG. 10




FIG. 11





FIG. 12

Azan staining, 6 random positions imaged



Azan.x50 Azan.x100



Measured by picking up blue collagen area
using NIH Image

FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25




FIG. 26A




FIG. 26B




FIG. 26C




FIG. 26D




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32