



(86) Date de dépôt PCT/PCT Filing Date: 2009/01/02
(87) Date publication PCT/PCT Publication Date: 2009/07/16
(85) Entrée phase nationale/National Entry: 2010/06/30
(86) N° demande PCT/PCT Application No.: US 2009/030005
(87) N° publication PCT/PCT Publication No.: 2009/089082
(30) Priorité/Priority: 2008/01/04 (US61/019,021)

(51) Cl.Int./Int.Cl. *A61K 31/15* (2006.01),
A61K 45/06 (2006.01), *A61P 13/10* (2006.01),
A61P 19/02 (2006.01), *A61P 29/00* (2006.01)

(71) Demandeur/Applicant:
ABBOTT LABORATORIES, US



(72) Inventeurs/Inventors:
PERNER, RICHARD J., US;
KORT, MICHAEL E., US;
DIDOMENICO, STANLEY, JR., US;
CHEN, JUN, US;
VASUDEVAN, ANIL, US

(74) Agent: TORYS LLP

(54) Titre : ANTAGONISTES DE TRPA1

(54) Title: TRPA1 ANTAGONISTS

Figure 1



(57) Abrégé/Abstract:

Compounds of formula (I), wherein R^1 , R^2 , R^3 , m , and Y are defined in the specification are TRPA1 antagonists. Compositions comprising such compounds and methods for treating conditions and disorders using such compounds and compositions are also disclosed.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 July 2009 (16.07.2009)

PCT

(10) International Publication Number
WO 2009/089082 A1

(51) International Patent Classification:

A61K 31/15 (2006.01) A61P 29/00 (2006.01)
A61K 45/06 (2006.01) A61P 13/10 (2006.01)
A61P 19/02 (2006.01)

(74) Agent: SRIVASTAVA, Sonali, S.; Abbott Laboratories,
Dept. 0377, Bldg. AP6A-1A, 100 Abbott Park Road, Ab-
bott Park, Illinois 60064-6008 (US).

(21) International Application Number:

PCT/US2009/030005

(22) International Filing Date: 2 January 2009 (02.01.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

61/019,021 4 January 2008 (04.01.2008) US

(71) Applicant (for all designated States except US): ABBOTT
LABORATORIES [US/US]; 100 Abbott Park Road, Ab-
bott Park, Illinois 60064 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PERNER, Richard,
J. [US/US]; 704 Penny Lane, Gurnee, Illinois 60031 (US).
KORT, Michael, E. [US/US]; 507 North Green Avenue,
Lake Bluff, Illinois 60044 (US). DIDOMENICO, Stan-
ley, Jr. [US/US]; P.O. Box 84, Richmond, Illinois 60071
(US). CHEN, Jun [US/US]; 100 Abbott Park Road, Abbott
Park, Illinois 60064 (US). VASUDEVAN, Anil [US/US];
105 13th Avenue, Union Grove, Wisconsin 53182 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,
CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID,
IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK,
LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW,
MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT,
RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM,
ZW.



(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK,
MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ,
CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN,
TD, TG).

Published:

— with international search report

(54) Title: TRPA1 ANTAGONISTS

Figure 1



(57) Abstract: Compounds of formula (I), wherein R^1 , R^2 , R^3 , m , and Y are defined in the specification are TRPA1 antagonists. Compositions comprising such compounds and methods for treating conditions and disorders using such compounds and compositions are also disclosed.

WO 2009/089082 A1

TRPA1 ANTAGONISTS

5

TECHNICAL FIELD

The present invention generally relates to TRPA1 antagonists, compositions comprising such compounds, and methods of treating conditions and disorders using such compounds and compositions.

10

BACKGROUND


TRPA1 is a nonselective cation channel that belongs to the superfamily of Transient Receptor Potential (TRP) ion channels. Like other family members, functional TRPA1 channels are formed by tetramerization of 4 subunits, each containing six transmembrane domains, a pore loop between transmembrane domain 5 (S5) and 6 (S6), and intracellular N- and C-termini. TRPA1 is expressed in sensory neurons and co-localized with pain markers such as TRPV1, calcitonin gene-related peptide and bradykinin receptor (Nagata, K. et al., Journal of Neuroscience 2005, 25, 4052-4061; Story, G. M. et al., Cell 2003, 112, 819-829; Corey, D. P. et al., Nature 2004, 432, 723-730; Bautista, D. M. et al., Proceedings of the National Academy of Science U.S.A. 2005, 102, 12248-12252; Jaquemar, D. et al., Journal of Biological Chemistry 1999, 274, 7325-7333). In pain models, knockdown of TRPA1 expression by gene specific antisenses prevented and reversed cold hyperalgesia induced by inflammation and nerve injury (Obata, K. et al., Journal of Clinical Investigation 2005, 115, 2393-2401; Jordt, S. E. et al., Nature 2004, 427, 260-265; Katsura, H. et al., Exploratory Neurology 2006, 200, 112-123). Furthermore, TRPA1 gene knockout resulted in impaired sensory functions and deficits in bradykinin-evoked pain hypersensitivity (Kwan, K. Y. et al. Neuron 2006, 50, 277-289; Bautista, D. M. et al. Cell 2006, 124, 1269-1282). Collectively, these data suggest that TRPA1 plays an important role in sensory functions and pain states. As a ligand-gated channel, TRPA1 can be activated by a variety of stimuli, including noxious cold, intracellular Ca^{2+} , endogenous substances (e.g., bradykinin), pungent natural products (e.g., allyl isothiocyanate, or AITC), environmental irritants (e.g., acrolein), amphipathic molecules (e.g., trinitrophenol and chlorpromazine) and pharmacological agents (e.g., URB597) (Macpherson, L. J. et al., Current Biology 2005, 15, 929-934; Bandell, M. et al., Neuron 2004, 41, 849-857). Bradykinin activates TRPA1 indirectly through the

phospholipase C pathway following binding of bradykinin to its receptors . Trinitrophenol and chlorpromazine open TRPA1 by inducing curvature or crenation in the lipid bilayer membrane (Xu, H. et al., Nat. Neurosci. 2006, 9, 628-635; Hill, K. and Schaefer, M., J. Biol. Chem. 2007, 282, 7145-7153; Niforatos, W. et al., Molecular Pharmacology 2007, 71, 1209-1216). Most recently, it was shown that TRPA1 agonists could directly interact with the channel protein. AITC and cinnamaldehyde covalently modify several cysteine and lysine residues localized in the cytoplasmic N terminus and activate the channel (Hinman, A., Chuang, H. H., Bautista, D. M., and Julius, D. Proceedings of the National Academy of Science U.S.A. 2006, 103, 19564-19568; Macpherson, L. J., Dubin, A. E., Evans, M. J., Marr, F., Schultz, P. G., Cravatt, B. F., and Patapoutian, A., Nature 2007, 445, 541-545). In addition, intracellular Ca^{2+} binds to the N-terminus EF-hand domains and mediates channel opening (Zurborg, S. et al., Nature Neuroscience 2007, 10, 277-279). Together these findings have revealed potential physiological roles of TRPA1, and also indicate that TRPA1 channel gating may involve different mechanisms and molecular determinants.

Thus, modulation of TRPA1 can have many industrial and therapeutic applications. For example, TRPA1 antagonists may fulfill the need in the art for new analgesic pharmaceuticals suitable for the treatment and/or prophylaxis of nociceptive and neuropathic pain in mammals, especially in humans.

SUMMARY

Disclosed herein are methods for treating disorders that are ameliorated by inhibition of TRPA1. Such methods comprise of administering to the subject a therapeutically effective amount of one or more TRPA1 antagonists, alone, or in combination with one or more pharmaceutically acceptable carriers. The TRPA1 antagonists employed are selected from compounds of formula (I), or solvates, pharmaceutical salts, prodrugs, salts of prodrugs, or any combinations thereof,



wherein

R^1 represent optional substituents of the phenyl group and each R^1 is independently alkyl, alkenyl, alkynyl, -CN, halogen, -OR^a, -NO₂, -N(R^a)(R^b), -N(R^b)C(O)R^a, -N(R^b)S(O)₂R^{1a}, -N(R^b)C(O)OR^a, -N(R^b)C(O)N(R^a)(R^b), -N(R^b)S(O)₂N(R^a)(R^b), -C(O)R^a,

-C(O)OR^a, -C(O)N(R^a)(R^b), -SR^a, -SF₅, -S(O)R^{1a}, -S(O)₂R^{1a}, -S(O)₂OR^{1a}, -S(O)₂N(R^a)(R^b),
 -(CR^dR^e)_q-CN, haloalkyl, -(CR^dR^e)_q-OR^a, -(CR^dR^e)_q-NO₂, -(CR^dR^e)_q-N(R^a)(R^b),
 -(CR^dR^e)_q-N(R^b)C(O)R^a, -(CR^dR^e)_q-N(R^b)S(O)₂R^{1a}, -(CR^dR^e)_q-N(R^b)C(O)OR^a,
 -(CR^dR^e)_q-N(R^b)C(O)N(R^a)(R^b), -(CR^dR^e)_q-N(R^b)S(O)₂N(R^a)(R^b), -(CR^dR^e)_q-C(O)R^a,
 5 -(CR^dR^e)_q-C(O)OR^a, -(CR^dR^e)_q-C(O)N(R^a)(R^b), -(CR^dR^e)_q-S(O)₂R^{1a}, -(CR^dR^e)_q-S(O)₂OR^{1a},
 or -(CR^dR^e)_q-S(O)₂N(R^a)(R^b);

R² is hydrogen, C₁₋₆ alkyl, or haloalkyl;

R³ is C₁₋₆ alkyl, haloalkyl, or cyclopropyl, wherein the cyclopropyl is optionally substituted with 1, 2, 3, 4, or 5 substituents as represented by R^j;

10 R^j, at each occurrence, is independently C₁₋₄ alkyl, halogen, or haloalkyl;

Y is -OH, -O(C₁₋₄ alkyl), -N(R^f)(R^g), -N(R^f)C(O)R^g, -N(R^f)S(O)₂R^g, or
 -N(R^f)C(O)N(R^f)(R^g);

R^a and R^b, at each occurrence, are each independently hydrogen, alkyl, or haloalkyl;

R^{1a}, at each occurrence, is independently alkyl or haloalkyl;

15 R^d and R^e, at each occurrence, are each independently hydrogen, alkyl, halogen, or haloalkyl;

R^f and R^g, at each occurrence, are each independently hydrogen, alkyl, or haloalkyl;

m is 0, 1, 2, 3, 4, or 5; and

q is 1, 2, 3, or 4;

20 with the proviso that the TRPA1 antagonist is not (Z)-4-(4-chlorophenyl)-3-methylbut-3-en-2-oxime or (E)-4-(4-chlorophenyl)-3-methylbut-3-en-2-oxime.

Some of the methods are directed to treating subjects suffering from acute cerebral ischemia, chronic pain, neuropathic pain, inflammatory pain, post herpetic neuralgia, neuropathies, neuralgia, diabetic neuropathy, HIV-related neuropathy, nerve injury,
 25 rheumatoid arthritic pain, osteoarthritic pain, burns, back pain, visceral pain, cancer pain, dental pain, headache, migraine, carpal tunnel syndrome, fibromyalgia, neuritis, sciatica, pelvic hypersensitivity, pelvic pain, menstrual pain; bladder disease such as incontinence, micturition disorder, renal colic and cystitis; inflammation such as burns, rheumatoid arthritis and osteoarthritis; neurodegenerative disease such as stroke, post stroke pain and multiple
 30 sclerosis; pulmonary disease such as asthma, cough, chronic obstructive pulmonary disease (COPD) and broncho constriction; gastrointestinal disease such as gastroesophageal reflux disease (GERD), dysphagia, ulcer, irritable bowel syndrome (IBS), inflammatory bowel disease (IBD), colitis and Crohn's disease; and ischemia such as cerebrovascular ischemia; emesis such as cancer chemotherapy-induced emesis, or obesity. In some methods, the

subject being treated is human. In some methods, one or more second pain reducing agents can be co-administered with the TRPA1 antagonist(s) as set forth above, alone or with one or more pharmaceutically acceptable carrier(s).

A further aspect of the invention provides a use of TRPA1 antagonists, alone or in combination with one or more second pain reducing agent, in the manufacture of a medicament for treating conditions and disorders disclosed herein.

For example, the second pain reducing agent can be an analgesic agent selected from acetaminophen, or a nonsteroidal anti-inflammatory drug (NSAID), or combinations thereof. In some embodiments, the nonsteroidal anti-inflammatory drug is ibuprofen. The second pain reducing agent can also be opioids.


Pharmaceutical compositions comprising one or more compound of formula (I), or pharmaceutically acceptable salts thereof, alone, or in combination with one or more second pain reducing agents, are also provided.

BRIEF DESCRIPTION OF DRAWINGS

Figure 1 – depicts the effect of Example 1 on hindlimb grip force in osteoarthritic rats.

DETAILED DESCRIPTION

Compounds of formula (I) are disclosed in this invention,



wherein R^1 , R^2 , R^3 , Y, and m are defined above in the Summary of the Invention and below in the Detailed Description. Further, compositions comprising such compounds and methods for treating conditions and disorders using such compounds and compositions are also disclosed.

Compounds disclosed herein may contain one or more variable(s) that occur more than one time in any substituent or in the formulae herein. Definition of a variable on each occurrence is independent of its definition at another occurrence. Further, combinations of substituents are permissible only if such combinations result in stable compounds. Stable compounds are compounds, which can be isolated from a reaction mixture.

a. Definition

As used in the specification and the appended claims, unless specified to the contrary, the following terms have the meaning indicated:

The term "alkenyl" as used herein, means a straight or branched chain hydrocarbon containing from 2 to 10 carbons and containing at least one carbon-carbon double bond.

5 Representative examples of alkenyl include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl.

The term "alkyl" as used herein, means a straight or branched saturated hydrocarbon chain containing from 1 to 10 carbon atoms. The term "C₁₋₆ alkyl" means a straight or
10 branched saturated hydrocarbon chain containing from 1 to 6 carbon atoms. The term "C₁₋₄ alkyl" means a straight or branched saturated hydrocarbon chain containing from 1 to 4 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, n-heptyl, n-octyl, n-nonyl,
15 and n-decyl.

The term "alkynyl" as used herein, means a straight or branched chain hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond. Representative examples of alkynyl include, but are not limited, to acetylenyl, 1-propynyl, 2-propynyl, 3-butylnyl, 2-pentylnyl, and 1-butylnyl.

20 The term "halo" or "halogen" as used herein, means -Cl, -Br, -I or -F.

The term "haloalkyl" as used herein, means at least one halogen, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of haloalkyl include, but are not limited to, chloromethyl, 2-fluoroethyl, trifluoromethyl, pentafluoroethyl, 4,4,4-trifluorobutyl, and 2-chloro-3-
25 fluoropentyl.

The term "subject" includes mammals, especially humans, as well as other non-human animals, e.g. horses, dogs, cats, and fish.

b. Compounds

Compounds of the invention have the general formula (I) as described above.

30 Particular values of variable groups in compounds of formula (I) are as follows. Such values may be used where appropriate with any of the other values, definitions, claims or embodiments defined hereinbefore or hereinafter.

In compounds of formula (I), R² is hydrogen, C₁₋₆ alkyl, or haloalkyl. For example, R² is C₁₋₆ alkyl such as, but not limited to, methyl. In some embodiments, R² is hydrogen.

R^3 is C_{1-6} alkyl, haloalkyl, or optionally substituted cycloalkyl. For example, R^3 is C_{1-6} alkyl such as, but not limited to, methyl, ethyl, n-propyl, tert-butyl, n-butyl, and isopropyl.

R^1 is an optional substituent of the phenyl group in formula (I) as described in the Summary. In certain embodiments, R^1 is halogen, alkyl (for example, C_{1-4} alkyl such as, but
5 not limited thereto, methyl), haloalkyl (e.g. trifluoromethyl), or -CN.

Y is -OH, -O(C_{1-4} alkyl), -N(R^f)(R^g), -N(R^f)C(O) R^g , -N(R^f)S(O) $_2R^g$, or
-N(R^f)C(O)N(R^f)(R^g); wherein R^f and R^g are as described in the Summary. In certain
embodiments Y is -OH or -O(C_{1-4} alkyl), for example, Y is -OH or -O(methyl). Certain
embodiments of invention include compounds of formula (I) wherein Y is -N(R^f)(R^g),
10 -N(R^f)C(O) R^g , -N(R^f)S(O) $_2R^g$, or -N(R^f)C(O)N(R^f)(R^g); and R^f and R^g are as described in the
Summary.

It is appreciated that the present invention contemplates methods for treating
conditions and disorders described herein using compounds of formula (I), pharmaceutically
acceptable salts, prodrugs, salts of prodrugs, solvates, or any combinations thereof, with
15 combinations of the above embodiments, including particular, more particular and preferred
embodiments.

Accordingly, examples of a group of compounds of formula (I) include, but are not
limited to, those wherein R^2 is C_{1-6} alkyl, R^3 is C_{1-6} alkyl, and m, R^1 , and Y are as disclosed in
the Summary. For example, R^2 is methyl. R^3 , for example, include ethyl, n-propyl, tert-
20 butyl, n-butyl, and isopropyl.

Other examples of a group of compounds of formula (I) include, but are not limited
to, those wherein R^2 is hydrogen, R^3 is C_{1-6} alkyl, and m, R^1 , and Y are as disclosed in the
Summary. R^3 , for example, include ethyl, n-propyl, tert-butyl, n-butyl, and isopropyl.

Within the two groups of compounds of formula (I) as described herein above,
25 examples of a subgroup include those wherein Y is -OH.

Other examples of a subgroup include, but are not limited to, those wherein Y is
O(C_{1-4} alkyl), for example, Y is -O(methyl).

Examples of yet another subgroup of compounds of formula (I) include those wherein
Y is -N(R^f)(R^g), -N(R^f)C(O) R^g , -N(R^f)S(O) $_2R^g$, or -N(R^f)C(O)N(R^f)(R^g); and R^f and R^g are as
30 described in the Summary.


Of all the groups and subgroups of compounds of formula (I) disclosed in the
preceding paragraphs, R^1 and m are as described in the Summary. For example, R^1 is
halogen, alkyl (for example, C_{1-4} alkyl such as, but not limited thereto, methyl), haloalkyl

(e.g. trifluoromethyl), or -CN. In one embodiment, m is 0. In other embodiment, m is 1, 2, or 3.

Compounds of formula (I) may contain one or more asymmetrically substituted atoms. The present invention contemplates various individual stereoisomers (including enantiomers and diastereomers) and mixtures thereof. Individual stereoisomers of compounds of the present invention may be prepared synthetically from commercially available starting materials that contain asymmetric or chiral centers or by preparation of racemic mixtures followed by resolution of the individual stereoisomer using methods that are known to those of ordinary skill in the art. Examples of resolution are, for example, (i) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography, followed by liberation of the optically pure product; or (ii) separation of the mixture of enantiomers or diastereomers on chiral chromatographic columns.

The invention also contemplates the various geometric isomers and mixtures thereof resulting from the disposition of substituents around a carbon-carbon double bond, a carbon-nitrogen double bond, a cycloalkyl group, or a heterocycle group. Substituents around a carbon-carbon double bond or a carbon-nitrogen double bond are designated as being of Z or E configuration and substituents around a cycloalkyl or heterocycle are designated as being of cis or trans configuration.

Examples of some of the possible geometrical isomers of compounds of formula (I) include, but are not limited to, (Ia), (Ib), (Ic), and (Id).



wherein R^1 , m, R^2 , R^3 , and Y have meanings as described in the Summary and the Detailed Description. It is understood that embodiments for R^1 , R^2 , Y, m, and R^3 , and combinations of embodiments, including particular, and more particular embodiments as described for formula (I), are also contemplated for compounds of formula (Ia), (Ib), (Ic), and (Id).

Within the present invention it is to be understood that compounds disclosed herein may exhibit the phenomenon of tautomerism and all tautomeric isomers are included in the scope of the invention.

Thus, the formulae drawings within this specification can represent only one of the possible tautomeric, geometric, or stereoisomeric forms. It is to be understood that the

invention encompasses any tautomeric, geometric, or stereoisomeric form, and mixtures thereof, and is not to be limited merely to any one tautomeric, geometric, or stereoisomeric form utilized within the formulae drawings.

Exemplary TRPA1 antagonists to be employed for the treatment of conditions and disorders described herein include, but are not limited to:

- (1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime;
- (1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one O-methyloxime;
- (1E,3E)-1-(2-fluorophenyl)-2-methylpent-1-en-3-one oxime;
- (1E,3E)-1-(4-fluorophenyl)-2-methylhex-1-en-3-one oxime;
- (1E,3E)-1-(4-fluorophenyl)-2,4,4-trimethylpent-1-en-3-one oxime;
- (1E,3E)-1-(4-fluorophenyl)-2-methylhept-1-en-3-one oxime;
- (1E,3Z)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime;
- (1E,3E)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime;
- (1E,3E)-1-(3-chloro-4-fluorophenyl)-2-methylpent-1-en-3-one oxime;
- (1E,3Z)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime;
- (2E,3E)-4-(4-fluorophenyl)but-3-en-2-one oxime;
- (1E,3E)-1-(4-fluorophenyl)hex-1-en-3-one oxime;
- (1E,3E)-1-(4-fluorophenyl)pent-1-en-3-one oxime;
- (1E,3Z)-1-(4-fluorophenyl)pent-1-en-3-one oxime;
- (1E,3E)-1-(3,4-difluorophenyl)-2-methylpent-1-en-3-one oxime; and
- (1E,3E)-2-methyl-1-(3,4,5-trifluorophenyl)pent-1-en-3-one oxime;

or pharmaceutically acceptable salts, prodrugs, salts of prodrugs, solvates, or combinations thereof.

c. Biological Data

(i) In Vitro Data – Determination of Inhibition Potencies

Molecular biology and transient expression were performed as described in Chen, J. et al., Journal of Biomolecular Screening 2007, 12, 61-69. Briefly, TRPA1 full length cDNA was amplified from human dorsal root ganglia total RNA (BD Bioscience Clontech, Palo Alto, CA, USA) and was cloned into pcDNA3.1/ V5-His Topo vector (Invitrogen, Carlsbad, CA). Large scale transient transfection was performed using the FreeStyle™ 293 Expression System as recommended by the manufacturer (Invitrogen). HEK293-F cells were grown in suspension in flasks (cell volume 30 mL to 1 liter) or in a Wave Bioreactor (Wave Biotech, Somerset, NJ) (6 liters). To support high density, suspension culture and transfection, cells were cultured in FreeStyle 293 media, an optimized, serum free formulation. 293fectin™ (Invitrogen) was used as transfection reagent. In a transfection of 3×10^7 cells (30 mL

volume), 30 μ g of plasmid DNA and 40 μ L 293 fectin were used. For a larger volume transfection, each reagent was scaled up proportionally. Two days post-transfection, cells were harvested by centrifugation (1000xg, 5 minutes) and resuspended to a density of 1.5×10^7 cells/mL in freezing medium (Freestyle media/10% serum/10% DMSO). Cells were transferred into cryovials in 2mL aliquots, and these vials were placed in Nalgene Mr. Frosty slow-freeze devices (Sigma-Aldrich, St. Louis, MO) at -80°C . As needed, vials were removed from -80°C and quickly thawed in a 37°C water bath. Cells were aseptically transferred into conical tubes containing Freestyle media (10 mL/vial). After centrifugation at 1,000xg for 3-5 minutes, medium was removed by aspiration and cells were resuspended in Freestyle medium at desired densities (typically 1×10^6 cells/mL). Re-suspended cells were seeded into black-walled clear-bottom 96-well BiocoatTM poly-D-lysine assay plates (BD Biosciences, Bedford, MA) (10^5 cells/well) and incubated overnight at 37°C under a humidified 5% CO_2 atmosphere.

FLIPR based intracellular Ca^{2+} assay and membrane potential assay

Ca^{2+} influx was measured using a FLIPR calcium assay kit (R8033; Molecular Devices, Sunnyvale, CA). The Ca^{2+} indicator dye was dissolved in Hanks' balanced salt solution supplemented with 20 mM HEPES buffer (HBSS/HEPES) according to the manufacturer instructions. Prior to start of the assay, the medium was removed by aspiration, and cells were loaded with 100 μ L Ca^{2+} dye for 2 to 3 hours at room temperature. AITC (allyl isothiocyanate) was used to activate and open the channels. (4x) solutions of the test compounds were prepared in HBSS/HEPES, and 50 μ L were added to the cells at a delivery rate of 10 μ L/sec. Changes in fluorescence were measured over time in a fluorometric imaging plate reader (FLIPR, Molecular Devices). Two additions were made over the course of an experimental run. For agonist experiments, assay buffer was added at the 10 s time point, followed by addition of agonist at the 3 minute 10 sec time point. For antagonist experiments, the antagonist was added at the 10 sec time point, followed by addition of the agonist 3 minutes later. Final assay volume for both the agonist and antagonist experiments was 200 μ L. Total length of an experimental run was 6.5 minutes.

Data was analyzed with GraphPad Prism[®] software (GraphPad Software, San Diego, CA), using a four-parameter logistic Hill equation to curve-fit concentration-effect data and derive EC_{50} or IC_{50} values. Numerical values are reported as mean $\pm$ SEM (n = number of experiments). SEM is calculated by dividing the standard deviation by the square root of the sample size (minus 1).

Changes in membrane potential were measured in hTRPA1-expressing LSTT cells using a FLIPR membrane potential assay kit (R8034, Molecular Devices). The procedure for thawing and plating the LSTT cells was identical to the Ca^{2+} influx assay (see above). The membrane potential dye was dissolved in HBSS/HEPES buffer according to the manufacturer's instructions, and then cells were loaded with dye (100 μL /well) for 45 minutes to 2 hours at room temperature. The protocol for addition of chemical agents, and measurement of changes in fluorescence, was the same as the Ca^{2+} influx assay, except for the emission filter setting. The results are shown in Table 1.

Table 1

Example	Human TRPA1 IC_{50} (μM)
1	0.074
2	8.3
3	4.6
4	0.50
5	8.0
6	4.4
7	0.85
8	3.3
9	6.0
10	0.701
11	20.6
12	5.91
13	9.46
14	0.574
15	4.67
16	>100
17	>100
18	>100
19	>100

Certain compounds of the invention were tested in the assay described above and are effective TRPA1 antagonists with IC_{50} values from about 25 μM to about 100 nM, for example, from about 5 μM to about 80 nM.

(ii) Pharmacokinetic Data

Groups of three fasted male Sprague-Dawley rats received a 10 $\mu\text{mol/kg}$ (2 m/kg) oral dose of the test compound, administered by gavage. The compounds were prepared as a solution in a 10% DMSO/poly(ethylene glycol)-400 (v/v). Blood samples were obtained from each animal at various time points up to eight hours after dosing.

Test compound was selectively extracted from plasma and brain homogenate using liquid-liquid extraction with *tert*-butylmethylether. The compounds of interest were separated from co-extracted contaminants on a 50 x 3 mm, Luna CN column (Phenomenex), with an acetonitrile:0.1% aqueous trifluoroacetic acid mobile phase at a flow rate of 0.35 ml/min. Plasma concentrations were determined by HPLC-MS/MS on an API2000 with Turbo Ion Spray interface, with MRM detection in the positive ionization mode. To calibrate compound concentration, a separate portion of each tissue sample was spiked with a known quantity of test compound and analyzed simultaneously with the samples. The results are shown in Table 2.




Table 2

	$R^3 = \text{CH}_3$ (Compound A)	$R^3 = \text{CH}_2\text{CH}_3$ (Example 1)
Human TRPA1 IC ₅₀ (μM)	0.33	0.074
<i>F</i> , rat (oral bioavailability) @ 10 $\mu\text{mol/kg}$	4%	38%
C _{max} (μM) @ 10 $\mu\text{mol/kg}$	0.22	0.72
plasma/brain	0	1.2

As shown in Table 2, Example 1 exhibited better potency in the inhibition of TRPA1 and better bioavailability than Compound A. CNS penetration of Example 1 is also superior to Compound A, as demonstrated by the brain/plasma ratio after oral administration. These results suggest that compounds of the invention are of value as better therapeutic agents for the treatment of pain, particularly in pain mediated by central sensitization such as chronic inflammatory pain and osteoarthritic pain.

(iii) In Vivo Data – Effects on Grip Force in Osteoarthritic Rats

Unilateral knee joint osteoarthritis was induced in the rats by a single intra-articular (i.a.) injection of sodium monoiodoacetate (MIA) (Sigma, St. Louis, MO) (3 mg in 0.05 mL sterile isotonic saline) into the joint cavity using a 26G needle under light (2-4%) halothane (Halocarbon Laboratories, River Edge, NJ) anesthesia. Following injection, the animals were allowed to recover from the effects of anesthesia (usually 5-10 minutes) before returning them to their home cages. Animals were tested 21 days after MIA injection. Measurements of peak hind limb grip force were conducted by recording the maximum compressive force exerted on the hind limb strain gauge setup, in a commercially available grip force

Figure 1 – depicts the effect of Example 1 on hindlimb grip force in osteoarthritic rats measurement system (Columbus Instruments, Columbus, OH). In this experiment, ED₅₀ was 112 µmol/kg p.o. (95% CI, 79-158 µmol/kg). During testing, each rat was gently restrained by grasping around its rib cage and then allowed to grasp the wire mesh frame (10-12 cm²) attached to the strain gauge. The experimenter then moved the animal in a rostral-to-caudal direction until the grip was broken. Each rat was sequentially tested twice at approximately 2-minute interval to obtain a raw mean grip force (CFmax). This raw mean grip force data was in turn converted to a maximum hind limb compressive force (CFmax) (gram force) per kg body weight for each animal. A group of age matched naïve animals was added to each experiment and the data obtained from the different dose groups for the compound being tested were compared to the naïve group. Example 1 is fully efficacious in this model with an ED₅₀ of 112 µmol/kg. Example 1 exhibits excellent plasma and brain exposure with increasing oral dose. At the 100 µmol/kg dose in the osteoarthritis grip force model, the measured plasma concentration of Example 1 was 6.2 µM.

d. Methods of Using the Compounds

Compounds described herein are TRPA1 antagonist and are capable of interfering with the expression, modification, regulation, or activation of TRPA1, or down-regulates one or more of the normal biological activities of TRPA1 (e.g. its ion-channel).

One embodiment of the present invention provides a method for treating a disorder that may be ameliorated by inhibiting TRPA1 receptor in a subject in need of such treatment.

The method comprises administering a therapeutically effective amount of one or more compounds of formula (I), solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, or any combination thereof, alone or in combination with one or more pharmaceutically acceptable carriers.

Another embodiment of the present invention provides a method for treating pain (e.g. nociceptive pain, neuropathic pain, inflammatory pain, osteoarthritic pain, etc.) in a subject in need of such treatment. This method comprises administering a therapeutically effective amount of one or more compounds of formula (I), solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, or any combination thereof, alone or in combination with one or more pharmaceutically acceptable carriers.

Yet another embodiment of the present invention provides a method for prophylaxis or treatment of ischemia including acute cerebral ischemia, pain including chronic pain, neuropathic pain, nociceptive pain, allodynia, inflammatory pain, inflammatory hyperalgesia, post herpetic neuralgia, neuropathies, neuralgia, diabetic neuropathy, HIV-related neuropathy, nerve injury, rheumatoid arthritic pain, osteoarthritic pain, burns, back pain, visceral pain, cancer pain, dental pain, headache, migraine, carpal tunnel syndrome, fibromyalgia, neuritis, sciatica, pelvic hypersensitivity, pelvic pain, and menstrual pain; bladder disease such as incontinence and bladder overactivity, bladder cystitis, micturition disorder, and renal colic; inflammation such as burns, oral mucositis, rheumatoid arthritis and osteoarthritis; neurodegenerative disease such as stroke, post stroke pain and multiple sclerosis; pulmonary disease such as asthma, cough, chronic obstructive pulmonary disease (COPD) and bronchoconstriction; skin disorders such as psoriasis, eczema, and dermatitis; gastrointestinal disease such as gastroesophageal reflux disease (GERD), dysphagia, ulcer, irritable bowel syndrome (IBS), inflammatory bowel disease (IBD), colitis and Crohn's disease; ischemia such as cerebrovascular ischemia; emesis such as cancer chemotherapy-induced emesis, in mammals, especially humans. For example, the compounds of formula (I) are useful for the treatment of pain, particularly nociceptive and inflammatory pain, more particularly, osteoarthritic pain. This method comprises the step of administering to a subject in need thereof a therapeutically effective amount of one or more compounds of formula (I), solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, alone or in combination with one or more pharmaceutically acceptable carriers.

Compounds of formula (I), including but not limited to those specified in the examples, solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, can be used for prophylaxis or treatment of inflammatory, nociceptive, and neuropathic pain as demonstrated by Bautista, D. Cell 2006, 124, 1269-1282; Trevisani, M. et al. Proceedings of the National Academy of Sciences USA 2007, 104, 13519-13524; Dai, Y. et al. Journal of Clinical Investigation 2007, 117, 1979-1987; Diogenes, A. et al.

Journal of Dental Research 2007, 86, 550-555; Katsura, H. et al. Journal of Neurochemistry 2007, 102, 1614-1624; and McMahon, S. B. et al. Cell 2006, 124, 1123-1125.

Compounds of formula (I), including but not limited to those specified in the examples, solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, can be used for prophylaxis or treatment of colitis and Crohn's disease as demonstrated by Kimball, E. S. et al. Neurogastroenterology & Motility 2007, 19, 90-400.

Compounds of formula (I), including but not limited to those specified in the examples, solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, can be used for prophylaxis or treatment of respiratory hypersensitivity such as cough, asthma, chronic obstructive pulmonary disease (COPD) (Andre et al., Journal of Clinical Investigation 2008, 118, 2574-2582; Bessac et al., Journal of Clinical Investigation 2008, 118, 1899-1910; Simon and Liedtke, Journal of Clinical Investigation 2008, 118, 2383-2386).

Compounds of formula (I), including but not limited to those specified in the examples, solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, can be used for prophylaxis or treatment of gastrointestinal diseases such as irritable bowel syndrome (IBS) and inflammatory bowel disease (IBD) as demonstrated by Penuelas, A. et al. European Journal of Pharmacology 2007, 576, 143-150 and Hayashi, S. et al. Inflammopharmacology 2007, 15, 218-222.

Compounds of formula (I), including but not limited to those specified in the examples, solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, can be used for prophylaxis or treatment of cold hyperalgesia or cold sensitivity as demonstrated by Story, G. M. Current Neuropharmacology 2006, 4, 183-196 and Story, G. M. and Gereau, R. W. Neuron 2006, 50, 177-180.

Compounds of formula (I), including but not limited to those specified in the examples, solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or any combination thereof, can be used as depilatory agents to prevent or reverse hirsutism as demonstrated by Kwan, K. Y. et al. Neuron 2006, 50, 277-289.

Compounds described herein may be administered alone, or in combination with one or more other compounds of formula (I), or in combination (i.e. co-administered) with one or more additional pharmaceutical agents such as a second pain reducing agent. For example, the second pain reducing agent can be an analgesic agent such as, but not limited to, acetaminophen, or a nonsteroidal anti-inflammatory drug (NSAID), or combination thereof. Examples of nonsteroidal anti-inflammatory drug (NSAID) include, but not limited to,

aspirin, diclofenac, diflusal, etodolac, fenbufen, fenoprofen, flufenisal, flurbiprofen, ibuprofen, indomethacin, ketoprofen, ketorolac, meclofenamic acid, mefenamic acid, meloxicam, nabumetone, naproxen, nimesulide, nitroflurbiprofen, olsalazine, oxaprozin, phenylbutazone, piroxicam, sulfasalazine, sulindac, tolmetin and zomepirac. Yet other class
5 of second pain-reducing agent includes opioids. Other analgesic such as local anesthetics including anticonvulsants and antidepressants may also be used in the combination therapy. Administering one or more classes of drugs in addition to TRPA1 antagonists can provide more effective amelioration of pain. Combination therapy includes administration of a single pharmaceutical dosage formulation containing one or more of the compounds of formula (I)
10 and one or more additional pharmaceutical agents; as well as administration of the compounds of formula (I) and each additional pharmaceutical agent, in its own separate pharmaceutical dosage formulation. For example, one or more compounds of formula (I) and one or more additional pharmaceutical agents, may be administered to the patient together, in a single oral dosage composition having a fixed ratio of each active ingredient, such as a
15 tablet or capsule; or each agent may be administered in separate oral dosage formulations.

Where separate dosage formulations are used, each active ingredient may be administered at essentially the same time (e.g., concurrently) or at separately staggered times (e.g., sequentially).

Actual dosage levels of active ingredients in the pharmaceutical compositions of this
20 invention can be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated and the condition and prior medical history of the patient being treated. However, it is within the
25 skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

TRPA1 antagonists described herein can be administered alone, or as a pharmaceutical composition comprising a therapeutically effective amount of one or more of
30 the TRPA1 antagonists in combination with one or more pharmaceutically acceptable carriers, with or without one or more second pain reducing agent. The phrase "therapeutically effective amount" means a sufficient amount of TRPA1 antagonist to treat disorders, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the

invention will be decided by the attending physician within the scope of sound medical judgment. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound employed; the specific composition employed; 5 the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well-known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than required to 10 achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The total daily dose of the compounds of formula (I) administered to a human or lower animal range from about 0.10 µg/kg body weight to about 50 mg/kg body weight. More preferable doses can be in the range of from about 0.10 µg/kg body weight to about 10 15 mg/kg body weight. If desired, the effective daily dose can be divided into multiple doses for purposes of administration. Consequently, single dose compositions may contain such amounts or submultiples thereof to make up the daily dose.

e. Pharmaceutical Compositions

The present invention further provides pharmaceutical compositions that comprise 20 TRPA1 antagonists of the present invention thereof. The pharmaceutical compositions comprise compounds of formula (I), solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or combinations thereof that may be formulated together with one or more pharmaceutically acceptable carriers.

Another aspect of the present invention provides a pharmaceutical composition 25 comprising one or more compounds of formula (I), solvates, pharmaceutically acceptable salts, prodrugs, salts of prodrugs, or combinations thereof, and one or more pharmaceutically acceptable carriers, in combination with one or more second pain reducing agents. In certain embodiments, the second pain reducing agent is an analgesic such as acetaminophen, a nonsteroidal anti-inflammatory drug (NSAID), or combination thereof. In certain 30 embodiments, the nonsteroidal anti-inflammatory drug is ibuprofen. In other embodiments, the second pain reducing agent is an opioid. Other analgesics such as local anesthetics including anticonvulsants and antidepressants are also contemplated.

The pharmaceutical compositions of the invention can be used for the treatment of the disorders as described herein in mammals, including human.

The pharmaceutical compositions of this invention can be administered to humans and other mammals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments or drops), buccally or as an oral or nasal spray. The term "parenterally" as used herein, refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The term "pharmaceutically acceptable carrier" as used herein, means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as, but not limited to, lactose, glucose and sucrose; starches such as, but not limited to, corn starch and potato starch; cellulose and its derivatives such as, but not limited to, sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as, but not limited to, cocoa butter and suppository waxes; oils such as, but not limited to, peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols; such as propylene glycol; esters such as, but not limited to, ethyl oleate and ethyl laurate; agar; buffering agents such as, but not limited to, magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as, but not limited to, sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol and the like), vegetable oils (such as olive oil), injectable organic esters (such as ethyl oleate) and suitable mixtures thereof. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms can be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid and the like. It may also be desirable to include isotonic agents such as sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the inclusion of agents, which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally-administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microcapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound may be mixed with at least one inert, pharmaceutically acceptable excipient or carrier, such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol and silicic acid; b) binders such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting agents such as cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay and i)

lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

5 Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such carriers as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

10 The solid dosage forms of tablets, dragees, capsules, pills and granules can be prepared with coatings and shells such as enteric coatings and other coatings well-known in the pharmaceutical formulating art. They may optionally contain opacifying agents and may also be of a composition such that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned carriers.

15 Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, 20 corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan and mixtures thereof.

25 Besides inert diluents, the oral compositions may also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, tragacanth and mixtures thereof.

30 Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating carriers or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals which are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients and the like. The preferred lipids are natural and synthetic phospholipids and phosphatidyl cholines (lecithins) used separately or together.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers or propellants which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

The compounds of the present invention can be used in the form of pharmaceutically acceptable salts. The phrase "pharmaceutically acceptable salt" means those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like and are commensurate with a reasonable benefit/risk ratio.

Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al. describe pharmaceutically acceptable salts in detail in (*J. Pharmaceutical Sciences*, 1977, 66: 1 et seq).

The compounds of the invention may contain either a basic or an acidic functionality, or both, and can be converted to a pharmaceutically acceptable salt, when desired, by using a suitable acid or base. The salts can be prepared in situ during the final isolation and purification of the compounds of the invention.

Representative acid addition salts include, but are not limited to acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate (isothionate), lactate, malate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, palmitoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate,

tartrate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides such as, but not limited to, methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides
5 such as, but not limited to, decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; arylalkyl halides like benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained. Examples of acids which can be employed to form pharmaceutically acceptable acid addition salts include such inorganic acids as hydrochloric acid, hydrobromic acid, sulfuric acid, and phosphoric acid and such organic acids as acetic
10 acid, fumaric acid, maleic acid, 4-methylbenzenesulfonic acid, succinic acid and citric acid.

Basic addition salts can be prepared in situ during the final isolation and purification of compounds of this invention by reacting a carboxylic acid-containing moiety with a suitable base such as, but not limited to, the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia or an organic primary, secondary
15 or tertiary amine. Pharmaceutically acceptable salts include, but are not limited to, cations based on alkali metals or alkaline earth metals such as, but not limited to, lithium, sodium, potassium, calcium, magnesium and aluminum salts and the like and nontoxic quaternary ammonia and amine cations including ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine,
20 diethylamine, ethylamine and the like. Other representative organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine and the like.

The term "pharmaceutically acceptable prodrug" or "prodrug" as used herein, represents those prodrugs of the compounds of the present invention which are, within the
25 scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use.

The present invention contemplates compounds of formula (I) formed by synthetic means or formed by in vivo biotransformation of a prodrug.

30 The compounds of the invention can exist in unsolvated as well as solvated forms, including hydrated forms, such as hemi-hydrates. In general, the solvated forms, with pharmaceutically acceptable solvents such as water and ethanol among others are equivalent to the unsolvated forms for the purposes of the invention.

f. General Synthesis


This invention is intended to encompass compounds of the invention when prepared by synthetic processes or by metabolic processes. Preparation of the compounds of the invention by metabolic processes include those occurring in the human or animal body (*in vivo*) or processes occurring *in vitro*.

5 The synthesis of compounds of general formula (I) wherein the groups R^1 , R^2 , R^3 , R^f , R^g , m , and Y have the meanings as set forth in the summary section unless otherwise noted, is exemplified in the accompanying Schemes 1-5.

As used in the descriptions of the schemes and the examples, certain abbreviations are intended to have the following meanings: Dibal-H for diisobutylaluminum hydride, DMAP for (4-dimethylamino)pyridine, DMF for N,N-dimethylformamide, DBU for 1,8-diazabicyclo(5.4.0)undec-7-ene, EDCI for 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, HOBt for 1-hydroxybenzotriazole, NMO for 4-methylmorpholine N-oxide, NMP for N-methylpyrrolidinone, PDC for pyridinium dichromate, $PdCl_2(PPh_3)_2$ for bis(triphenylphosphine)palladium(II) dichloride, $Pd(PPh_3)_4$ for tetrakis(triphenylphosphine)palladium(0), Ph for phenyl, and TPAP for tetrapropylammonium perruthenate..

Compounds of general formula (I) wherein Y is $-OH$ or $-O(C_{1-4} \text{ alkyl})$ can be prepared using general procedures as illustrated in Scheme 1.

Scheme 1




20

Benzaldehydes of general formula (1) can be reacted with phosphorus ylides of general formula (2) wherein R'' is alkyl, in a solvent such as toluene at refluxing temperatures to provide unsaturated esters of general formula (3) (Adam, W. et al., Journal of Organic Chemistry 2002, 67, 8395-8399). The conversion of compound (3) to (4) is a stepwise reaction sequence comprising: (a) reduction of (3) to the corresponding aldehyde using a reagent such as diisobutylaluminum hydride or lithium aluminum hydride, (b) addition of an organometallic nucleophile in the form of a Grignard or organolithium reagent, and (c) conversion of the product from step (b) to the corresponding ketone derivative using an

25

oxidizing reagent such as tetrapropylammonium perruthenate (Ley, S. V. et al. Synthesis 1994, 7, 639-966) or pyridinium dichromate (PDC). Step (b) can be carried out in a solvent such as tetrahydrofuran or diethyl ether at a temperature ranging from about -78 °C to about 0 °C. The crude products from both steps (a) and (b) can be isolated and subjected to the subsequent step with or without purification. Step (c) can be carried out using a variety of oxidizing agents known to those skilled in the art in non-polar solvents such as dichloromethane typically at ambient temperatures. Condensation of ketone (4) with hydrochloride salt of hydroxylamine or alkoxyamine formula R^xONH_2 wherein R^x is hydrogen or C_{1-4} alkyl, in the presence of a base such as but not limited to triethylamine in a solvent such as benzene or tetrahydrofuran at a temperature ranging from about 25 °C to about 80 °C, provides oximes of general formula (5) or (6). Compounds (5) and (6) are geometrical isomers which may be separated and purified by standard chromatographic methods familiar to those skilled in the art.

Alternatively, ketone (4) can be prepared employing general procedures as illustrated in Scheme 2.




Unsaturated esters of general formula (3), prepared according to the methods of Scheme 1 or Scheme 3, can be reacted with N-methoxy-N-methylamine hydrochloride and trimethylaluminum (Lipton, M. F. et al. Organic Syntheses 1980, 59, 49-53) in a solvent such as toluene or dichloromethane at a temperature ranging from about -40 °C to about 25 °C to provide amides of general formula (7). Alternatively, (7) can be prepared by reacting (3) with N-methoxy-N-methylamine hydrochloride and isopropylmagnesium chloride (Williams, J.M. et al. Tetrahedron Letters 1995, 36, 5461-5464) in a solvent such as tetrahydrofuran at a temperature ranging from about -20 °C to about 0 °C.

Treatment of (7) with Grignard reagents of formula R^3MgX^{101} wherein X^{101} is Cl or Br, in a solvent such as tetrahydrofuran at a temperature ranging from about -78 °C to about 0 °C provides ketones of general formula (4).

Intermediate (7) can also be prepared according to the synthetic sequence as shown in Scheme 3. Ester (3) can be saponified to carboxylic acid (8) using reagents such as lithium or potassium hydroxide in aqueous solution at ambient temperature using a co-solvents


including but not limited to tetrahydrofuran and ethanol. Carboxylic acid (8) can be reacted with N-methoxy-N-methylamine hydrochloride (Basha, A. et al. Tetrahedron Letters 1977, 48, 4171-4174), using dehydrative coupling reagents such as EDCI in combination with HOBt or DMAP (Montalbetti, C. A. G. N. Tetrahedron 2005, 61 10827-10852), and a base such as but not limited to triethylamine. These reactions are carried out in a solvent such as toluene or dichloromethane at a temperature ranging from about 0 °C to about 25 °C to provide amides of general formula (7).

Scheme 3



Alternatively, unsaturated esters of formula (3) may be prepared according to the procedures as described in Scheme 4.


Scheme 4



Aryl halides of general formula (9) wherein the Z group is a chlorine, bromine, or iodine atom may be reacted with unsaturated esters of formula (10) in the presence of a palladium catalyst such as but not limited to palladium(II) acetate to generate (3) (Knowles, J. P. Organic & Biomolecular Chemistry 2007, 5, 31-44). These reactions are generally performed using an amine base such as but not limited to diisopropylethylamine or DBU, in a solvent such as acetonitrile, toluene, or DMF at temperatures ranging from about 50 °C to about 100 °C. A related approach for the synthesis of unsaturated esters of general formula (3) from (9) involves reaction with substituted alkenes of general formula (11) wherein R''' is alkyl. In this case, the substituted alkene (11) contains a trialkyltin moiety such as trimethylstannyl or tributylstannyl (Yin, L. et al. Chemical Reviews 2007, 107, 133-173). These reactions are generally performed in the presence of catalytic quantities of a palladium(II) or palladium (0) reagent, such as dichloropalladium bis(triphenylphosphine) or tetrakis(triphenylphosphine)palladium, respectively, in polar aprotic solvents such as DMF or NMP at temperatures ranging from about 25 °C to about 125 °C.

Compounds of general formula (I) wherein Y is NR^fR^g can be prepared using general procedures as illustrated in Scheme 5.

Scheme 5



Reaction of ketone (4), prepared according to the methods illustrated in Schemes 1 and 2 with substituted hydrazines of general formula $\text{R}^f\text{R}^g\text{NNH}_2$ in a solvent such as benzene or tetrahydrofuran at a temperature ranging from about 0 °C to about 80 °C provides hydrazones of general formula (12) or (13). Compounds (12) and (13) are geometrical isomers which may be separated and purified by standard chromatographic methods familiar to those skilled in the art.

It will be appreciated that the synthetic schemes and specific examples as illustrated in the Examples section are illustrative and are not to be read as limiting the scope of the invention as it is defined in the appended claims. All alternatives, modifications, and equivalents of the synthetic methods and specific examples are included within the scope of the claims.

Optimum reaction conditions and reaction times for each individual step may vary depending on the particular reactants employed and substituents present in the reactants used. Unless otherwise specified, solvents, temperatures and other reaction conditions may be readily selected by one of ordinary skill in the art. Specific procedures are provided in the Synthetic Examples section. Reactions may be worked up in the conventional manner, e.g. by eliminating the solvent from the residue and further purified according to methodologies generally known in the art such as, but not limited to, crystallization, distillation, extraction, trituration and chromatography. Unless otherwise described, the starting materials and reagents are either commercially available or may be prepared by one skilled in the art from commercially available materials using methods described in the chemical literature.

Routine experimentations, including appropriate manipulation of the reaction conditions, reagents and sequence of the synthetic route, protection of any chemical functionality that may not be compatible with the reaction conditions, and deprotection at a suitable point in the reaction sequence of the method are included in the scope of the invention. Suitable protecting groups and the methods for protecting and deprotecting different substituents using such suitable protecting groups are well known to those skilled in

the art; examples of which may be found in T. Greene and P. Wuts, Protecting Groups in Chemical Synthesis (3rd ed.), John Wiley & Sons, NY (1999), which is incorporated herein by reference in its entirety. Synthesis of the compounds of the invention may be accomplished by methods analogous to those described in the synthetic schemes described hereinabove and in specific examples.

Starting materials, if not commercially available, may be prepared by procedures selected from standard organic chemical techniques, techniques that are analogous to the synthesis of known, structurally similar compounds, or techniques that are analogous to the above described schemes or the procedures described in the synthetic examples section.

When an optically active form of a compound of the invention is required, it may be obtained by carrying out one of the procedures described herein using an optically active starting material (prepared, for example, by asymmetric induction of a suitable reaction step), or by resolution of a mixture of the stereoisomers of the compound or intermediates using a standard procedure (such as chromatographic separation, recrystallization or enzymatic resolution).

Similarly, when a pure geometric isomer of a compound of the invention is required, it may be obtained by carrying out one of the above procedures using a pure geometric isomer as a starting material, or by resolution of a mixture of the geometric isomers of the compound or intermediates using a standard procedure such as chromatographic separation.

Following Examples may be used for illustrative purposes and should not be deemed to narrow the scope of the invention.

g. Examples

Example 1

(1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime

Example 1A

(E)-ethyl 3-(4-fluorophenyl)-2-methylacrylate

4-Fluorobenzaldehyde (12 g, 97 mmol) was dissolved in toluene (500 mL). (Carboethoxyethylidene)triphenylphosphorane (42 g, 116 mmol) was added and the mixture was heated to reflux until the starting material was consumed. The solution was cooled, diluted with hexanes, and the solid was filtered. The filtrate was evaporated in vacuo. The remaining residue was dissolved in ether, and filtered through a plug of silica gel using ether as eluent. The solvents were evaporated in vacuo to afford the title compound (20 g, 99%

yield) as a yellow solid. ^1H NMR ($\text{DMSO}-d_6$) δ ppm: 7.64 (s, 1H), 7.38 (dd, $J = 8.65, 5.59$ Hz, 2H), 7.08 (t, $J = 8.65$ Hz, 2H), 4.27 (q, $J = 7.12$ Hz, 2H), 2.10 (d, $J = 1.36$ Hz, 3H), 1.27-1.38 (m, 3 H). MS (DCI^+) 209 ($\text{M}+\text{H}$).

5

Example 1B

(E)-3-(4-fluorophenyl)-2-methylacrylic acid

Example 1A (20 g, 0.1 mol) was dissolved in THF (400 mL) and 1M aq LiOH (200 mL) was added. The mixture was stirred at ambient temperature overnight. The homogeneous solution was diluted with water, and extracted with ethyl acetate. The organic
10 phase was washed with water and brine, dried over Na_2SO_4 , filtered and evaporated in vacuo to afford the title compound (16.1 g, 89% yield) as a white solid. ^1H NMR (300 MHz, CDCl_3) δ ppm 7.78 (s, 1H), 7.43 (dd, $J = 8.82, 5.42$ Hz, 2H), 7.11 (t, $J = 8.65$ Hz, 2H), 2.14 (s, 3H). MS (DCI^+) 198 ($\text{M}+\text{NH}_4$).

15

Example 1C

(E)-3-(4-fluorophenyl)-N-methoxy-N,2-dimethylacrylamide

Method A

A mixture of Example 1B (16.1 g, 89.4 mmol), *N,O*-dimethylhydroxylamine hydrochloride (9.15 g, 93.9 mmol), 1-hydroxybenzotriazole hydrate (12.7 g, 93.9 mmol), 1-
20 ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (17.9 g, 93.9 mmol), and diisopropylethyl amine (47.4 g, 370 mmol) was stirred in DMF (400 mL) at ambient temperature overnight. The solution was diluted with water and extracted with ethyl acetate. The organic phase was washed with saturated sodium bicarbonate solution, 1N HCl, water, brine, dried over Na_2SO_4 , and evaporated in vacuo. The resulting material was
25 chromatographed (SiO_2 , 20% ethyl acetate/hexanes to 50% ethyl acetate/hexanes) to yield the title compound (16.89g, 58% yield) as a yellow oil. ^1H NMR (300 MHz, CDCl_3) δ ppm 7.33 (dd, $J = 8.82, 5.43$ Hz, 2H), 7.00-7.14 (m, 2H), 6.79 (s, 1H), 3.70 (s, 3H), 3.29 (s, 3H), 2.11 (s, 3H). MS (DCI^+) 224 ($\text{M}+\text{H}$).

Method B

30

A solution of isopropylmagnesium chloride (2.0 M in THF, 59.1 mL, 118 mmol) was added dropwise via addition funnel to a mixture of Example 1A (5.47 g, 26.3 mmol) and *N,O*-dimethylhydroxylamine hydrochloride (5.12 g, 52.5 mmol) in THF (100 mL) at 0°C . Upon complete addition, the reaction was allowed to proceed 2 hr at 0°C . The reaction was

quenched by slow addition of saturated aq NH_4Cl solution (70 mL). The mixture was stirred 60 min, then partitioned between EtOAc (50 mL) and H_2O (100 mL). The separated aqueous phase was extracted with EtOAc, and the combined organic layer was washed with 1 N aq HCl and brine, dried (Na_2SO_4), filtered, and concentrated in vacuo. The crude product was
5 purified by chromatography on silica gel (Analogix[®] Intelliflash 280; SF65-400g column; 20% to 60% EtOAc/hexanes; 0-40 min) to give the title compound (5.06 g, 22.7 mmol, 86 % yield) as a colorless oil. ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ ppm 7.50–7.40 (m, 2H), 7.29–7.18 (m, 2H), 6.72 (s, 1H), 3.65 (s, 3H), 3.19 (s, 3H), 2.01 (d, $J = 1.5$ Hz, 3H). MS (DCI^+) 224 (M+H).

10

Example 1D

(E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one

Example 1C (7 g, 31.4 mmol) was dissolved in tetrahydrofuran (175 mL) and cooled to about -20°C . Ethylmagnesium bromide (1M in THF, 94 mL, 94 mmol) was added
15 dropwise. The solution was stirred while warming to about 0°C for 3 hr. The reaction was quenched with saturated NH_4Cl solution, and then diluted with water. The mixture was extracted with ethyl acetate. The combined organic phase was washed with water and brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The solid was chromatographed (SiO_2 , 10% ethyl acetate/hexanes) to afford the title compound as a yellow solid (4.59 g, 76% yield).
20 ^1H NMR (300 MHz, CDCl_3) δ ppm 7.48 (s, 1H), 7.40 (dd, $J = 8.82, 5.43$ Hz, 2H), 7.10 (t, $J = 8.65$ Hz, 2H), 2.83 (q, $J = 7.12$ Hz, 2H), 2.05 (d, $J = 1.36$ Hz, 3H), 1.17 (t, $J = 7.29$ Hz, 3H). MS (DCI^+) 193 (M+H).

Example 1E

(1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime

Example 1D (4.59 g, 23.9 mmol) was dissolved in pyridine (100 mL), followed by addition of hydroxylamine hydrochloride (2.5 g, 35.8 mmol). The reaction was stirred at ambient temperature for two hours. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and
30 brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The resulting solid was chromatographed (SiO_2 , 20% ethyl acetate/hexanes), and then recrystallized from hexanes to afford the title compound (3.25 grams, 66% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 8.69

(bs, 1H), 7.33–7.24 (m, 2H), 7.11–7.01 (m, 2H), 6.87 (s, 1H), 2.70 (q, $J = 7.6$ Hz, 2H), 2.04 (d, $J = 1.2$ Hz, 3H), 1.18 (t, $J = 7.6$ Hz, 3H). MS (DCI⁺) 208 (M+H).

Example 2

5 (1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one O-methyloxime

Example 1D (100 mg, 0.52 mmol) was dissolved in pyridine (5 mL), followed by addition of methoxylamine hydrochloride (65 mg, 0.78 mmol), and the reaction was stirred at ambient temperature overnight. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and
10 brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting solid was chromatographed (SiO₂, 20% ethyl acetate/hexanes) to afford the title compound (42 mg, 37% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.25-7.31 (m, 2H), 7.04 (t, 2H), 6.80 (s, 1H), 3.94 (s, 3H), 2.61 (q, $J = 7.46$ Hz, 2H), 2.04 (d, $J = 1.36$ Hz, 3H), 1.11 (t, $J = 7.46$ Hz, 3H). MS (ESI⁺) 221 (M+H).

15

Example 3

(1E,3E)-1-(2-fluorophenyl)-2-methylpent-1-en-3-one oxime

Example 3A

20 (E)-ethyl 3-(2-fluorophenyl)-2-methylacrylate

2-Fluorobenzaldehyde (5.0 g, 40.3 mmol) was dissolved in 150 mL toluene. (Carboethoxyethylidene)triphenylphosphorane (42 g; 116 mmol) was added and the mixture was heated to reflux overnight. The solution was cooled, diluted with hexanes, and the solid was filtered. The filtrate was evaporated in vacuo, then dissolved in ether and filtered
25 through a plug of silica gel using ether as eluent. The solvents were evaporated in vacuo to afford the title compound (10.57 g) as a yellow solid. ¹H NMR (300 MHz, CDCl₃) δ ppm 7.70 (s, 1H), 7.26-7.38 (m, 3H), 7.17 (m, 2H), 4.28 (q, $J = 7.12$ Hz, 2H), 2.04 (s, 3H), 1.35 (t, $J = 7.12$ Hz, 3H).

30

Example 3B

(E)-3-(2-fluorophenyl)-N-methoxy-N,2-dimethylacrylamide

N,O-dimethylhydroxylamine hydrochloride (7 g, 72 mmol) was slurried in CH₂Cl₂ (250 mL) and cooled to 0-5°C. A solution of trimethylaluminum (2M in toluene, 36 mL, 72

mmol) was added dropwise. The mixture was warmed to ambient temperature and stirred for 1 hr. A solution of Example 3A (5 g, 24.0 mmol) in CH₂Cl₂ (100 ml) was added and the solution was heated to reflux overnight. The reaction mixture was cooled in an ice bath and quenched carefully with ice cold 0.5 M HCl. The mixture was filtered through celite filter
5 aid and extracted with CH₂Cl₂. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The crude product was chromatographed (SiO₂, 100% hexanes to 20% ethyl acetate/hexanes) to afford the title compound (1.88 g, 35% yield) as a light yellow oil. ¹H NMR (300 MHz, CDCl₃) δ ppm 7.27-7.38 (m, 2H), 7.04-7.17 (m, 2H), 6.79 (s, 1H), 3.72 (s, 3H), 3.30 (s, 3H), 2.04 (s, 3H). MS (ESI⁺) 224
10 (M+H).

Example 3C

(E)-1-(2-fluorophenyl)-2-methylpent-1-en-3-one

Example 3B (870 mg, 3.91 mmol) was dissolved in tetrahydrofuran (20 mL) and
15 cooled to about -20°C. Ethylmagnesium bromide (1M in THF, 7.8 mL, 7.8 mmol) was added dropwise, and the reaction was stirred for 1.5 hr at about -20°C. The reaction was quenched with NH₄Cl solution and extracted with ethyl acetate. The combined organic extract was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting oil was chromatographed (SiO₂, 10% ethyl acetate/hexanes) to afford
20 the title compound as a yellow oil (240 mg, 32% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.55 (s, 1H), 7.29-7.41 (m, 2H), 7.05-7.20 (m, 3H), 2.85 (q, *J* = 7.46 Hz, 2H), 1.18 (q, *J* = 7.46 Hz, 3H). MS (DCI⁺) 193 (M+H).

Example 3D

(1E,3E)-1-(2-fluorophenyl)-2-methylpent-1-en-3-one oxime

Example 3C (100 mg, 0.52 mmol) was dissolved in pyridine (3 mL), followed by addition of hydroxylamine hydrochloride (78 mg, 1.12 mmol). The solution was stirred at ambient temperature for 1.5 hr. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and
30 brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting solid was recrystallized from hexanes to afford the title compound (52 mg, 48% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.27-7.34 (m, 2H), 7.04-7.17 (m, 2H), 6.89 (s, 1H), 2.72 (q, *J* = 7.46 Hz, 2H), 1.99 (s, 3H), 1.18 (t, *J* = 7.63 Hz, 3H). MS (DCI⁺) 208 (M+H).

Example 4

(1E,3E)-1-(4-fluorophenyl)-2-methylhex-1-en-3-one oxime

5

Example 4A

(E)-1-(4-fluorophenyl)-2-methylhex-1-en-3-one

Example 1C (2 g, 8.95 mmol) was dissolved in tetrahydrofuran (50 mL) and cooled to about -20°C . Propylmagnesium chloride (2M in Et_2O , 9 mL, 18 mmol) was added dropwise and the resulting mixture was stirred at about -20°C for 1 hr. The solution was quenched
10 with saturated NH_4Cl solution and extracted with ethyl acetate. The combined organic phase was washed with water and brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The resulting material was chromatographed (SiO_2 , 10% ethyl acetate/hexanes to 50% ethyl acetate/hexanes) to afford the title compound (900 mg, 49% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 7.47 (s, 1H), 7.36-7.43 (m, 2H), 7.06-7.14 (m, 2H), 2.77 (q, $J = 7.29$ Hz, 2H),
15 2.04 (d, $J = 1.36$ Hz, 3H), 1.65-1.78 (m, 2H), 0.98 (t, $J = 7.29$ Hz, 3H). MS (DCI^+) 207 (M+H).

Example 4B

(1E,3E)-1-(4-fluorophenyl)-2-methylhex-1-en-3-one oxime

20 Example 4A (200 mg, 0.97 mmol) was dissolved in pyridine (3 mL), followed by addition of hydroxylamine hydrochloride (101 mg, 1.45 mmol). After stirring at ambient temperature for 2 hr, the solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The resulting solid was recrystallized from
25 hexanes to afford the title compound (78 mg, 36% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 7.25-7.31 (m, 2H), 7.05 (m, 2H), 6.85 (s, 1H), 2.61-2.69 (m, 2H), 2.02 (s, 3H), 1.54-1.68 (m, 2H), 1.02 (t, $J = 7.29$ Hz, 3H).

Example 5

30

(1E,3E)-1-(4-fluorophenyl)-2,4,4-trimethylpent-1-en-3-one oxime

Example 5A

(E)-1-(4-fluorophenyl)-2,4,4-trimethylpent-1-en-3-one

Example 1C (2 g, 8.95 mmol) was dissolved in tetrahydrofuran (50 mL) and cooled to about -20°C . tert-Butylmagnesium chloride (1M in THF, 18 mL, 18 mmol) was added dropwise. After stirring at about -20°C for 1 hr, the reaction was quenched with saturated NH_4Cl solution and extracted with ethyl acetate. The organic phase was washed with water and brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The resulting material was chromatographed (SiO_2 , 10% ethyl acetate/hexanes to 50% ethyl acetate/hexanes) to afford the title compound (410 mg, 21% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 7.25-7.33 (m, 2H), 7.06 (m, 2H), 6.82 (s, 1H), 2.03 (d, $J = 1.70$ Hz, 3H), 1.31 (s, 9H). MS (DCI^+) 221 (M+H).

10

Example 5B

(1E,3E)-1-(4-fluorophenyl)-2,4,4-trimethylpent-1-en-3-one oxime

Example 5A (200 mg, 0.91 mmol) was dissolved in pyridine (3 mL), followed by addition of hydroxylamine hydrochloride (101 mg, 1.45 mmol). The solution was stirred at ambient temperature overnight. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The resulting solid was recrystallized from hexanes to afford the title compound (92 mg, 43% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 7.31 (m, 2H), 7.04 (m, 2H), 6.18 (s, 1H), 2.03 (d, $J = 1.36$ Hz, 3H), 1.22 (s, 9H).

20

Example 6

(1E,3E)-1-(4-fluorophenyl)-2-methylhept-1-en-3-one oxime

25

Example 6A

(E)-1-(4-fluorophenyl)-2-methylhept-1-en-3-one

Example 1C (2 g, 8.96 mmol) was dissolved in tetrahydrofuran (50 mL) and cooled to about -20°C . Butylmagnesium chloride (2M in Et_2O , 8.96 mL, 17.9 mmol) was added dropwise and the mixture was stirred at about -20°C for 1 hr. The reaction was quenched with saturated NH_4Cl solution and extracted with ethyl acetate. The combined organic extract was washed with water and brine, dried over Na_2SO_4 , filtered, and evaporated in vacuo. The resulting material was chromatographed (SiO_2 , 10% ethyl acetate/hexanes to 50% ethyl acetate/hexanes) to afford the title compound (1.2 g, 61% yield). ^1H NMR (300

30

MHz, CDCl₃) δ ppm 7.46 (s, 1H), 7.36-7.43 (m, 2H), 7.07-7.14 (m, 2H), 2.76-2.81 (m, 2H), 2.04 (d, J = 1.36 Hz, 3H), 1.66 (dt, J = 15.17, 7.50 Hz, 2H), 1.32-1.45 (m, 2H), 0.95 (t, J = 7.29 Hz, 3H). MS (DCI⁺) 221 (M+H).

5

Example 6B

(1E,3E)-1-(4-fluorophenyl)-2-methylhept-1-en-3-one oxime

Example 6A (250 mg, 1.14 mmol) was dissolved in pyridine (5 mL), followed by addition of hydroxylamine hydrochloride (119 mg, 1.70 mmol). The resulting mixture was stirred at ambient temperature for two hours. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting solid was chromatographed (SiO₂, 20% ethyl acetate/hexanes), then recrystallized from hexanes to afford the title compound as a white solid (110 mg, 38% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.25-7.31 (m, 2H), 7.01-7.10 (m, 2H), 6.84 (s, 1H), 2.63-2.70 (m, 2H), 2.02 (s, 3H), 1.49-1.58 (m, 2H), 1.39-1.47 (m, 2H), 0.96 (t, J = 7.29 Hz, 3H).

10

15

Example 7

(1E,3Z)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime

20

Example 7A

(E)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one

25

30

Example 1C (2 g, 8.96 mmol) was dissolved in tetrahydrofuran (50 mL) and cooled to about -20°C. Isopropylmagnesium bromide (2M in THF, 8.96 mL, 17.9 mmol) was added dropwise, and the resulting mixture was stirred while warming to 0°C for 2.5 hr. The reaction was quenched with saturated NH₄Cl solution and diluted with water. The mixture was extracted with ethyl acetate. The combined organic extract was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The solid was chromatographed (SiO₂, 10% ethyl acetate/hexanes) to afford the title compound as a yellow solid (590 mg, 32% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.46 (s, 1H), 7.37-7.43 (m, 3H), 7.10 (m, 2H), 3.38-3.51 (m, 1H), 2.05 (d, J = 1.36 Hz, 3H), 1.18 (s, 3H), 1.16 (s, 3H).

Example 7B

(1E,3Z)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime

Example 7A (250 mg, 1.21 mmol) was dissolved in pyridine (3 mL), followed by addition of hydroxylamine hydrochloride (101 mg, 1.45 mmol). The reaction was stirred at ambient temperature overnight. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting solid was chromatographed using 10% ethyl acetate/hexanes, and the earlier eluting component was isolated to yield the title compound (62 mg, 23% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.27-7.33 (m, 2H), 6.99-7.08 (m, 2H), 6.61 (s, 1H), 3.39 (m, 1H), 2.02 (s, 3H), 1.28 (s, 3H), 1.26 (s, 3H).

Example 8

(1E,3E)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime

Example 7A (250 mg, 1.21 mmol) was dissolved in pyridine (3 mL), followed by addition of hydroxylamine hydrochloride (101 mg, 1.45 mmol). The reaction was stirred at ambient temperature overnight. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting solid was chromatographed using 10% ethyl acetate/hexanes, and the later eluting component was isolated to yield the title compound (67 mg, 25% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.27-7.33 (m, 2H), 7.00-7.07 (m, 2H), 7.04 (s, 1H), 6.30 (s, 1H), 2.70 (m, 1H), 2.06 (s, 3H), 1.19 (s, 3H), 1.17 (s, 3H).

Example 9

(1E,3E)-1-(3-chloro-4-fluorophenyl)-2-methylpent-1-en-3-one oxime

Example 9A

(E)-ethyl 3-(3-chloro-4-fluorophenyl)-2-methylacrylate

3-Chloro-4-fluorobenzaldehyde (10 g, 63.1 mmol) was dissolved in 300 mL toluene. (Carboethoxyethylidene)triphenylphosphorane (25 g, 69.4 mmol) was added and the mixture was heated to reflux overnight. The solution was cooled, diluted with hexanes and the solid was filtered. The filtrate was evaporated in vacuo, and the residue was dissolved in ether and filtered through a plug of silica using ether as eluent. The solvent was evaporated in vacuo to afford the title compound (15.55 g) as a semi solid. ¹H NMR (300 MHz, CDCl₃) δ ppm 7.56

(s, 1H), 7.44 (m, 1H), 7.23-7.32 (m, 1H), 7.16 (m, 1H), 4.28 (q, $J = 7.12$ Hz, 2H), 2.09 (d, $J = 1.36$ Hz, 3H), 1.35 (t, $J = 7.36$ Hz, 3H). MS (DCI⁺) 243 (M+H).

Example 9B

5 (E)-3-(3-chloro-4-fluorophenyl)-2-methylacrylic acid

Example 9A (15.5 g, 63.9 mmol) was dissolved in tetrahydrofuran (300 mL) and 1M LiOH (128 mL) was added. The mixture was stirred at ambient temperature overnight. The homogeneous solution was diluted with water and extracted with ethyl acetate. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo
10 to afford the title compound (12 g, 88% yield) as an off-white solid. ¹H NMR (300 MHz, CDCl₃) δ ppm 7.70 (s, 1H), 7.48 (dd, $J = 7.12, 2.37$ Hz, 1H), 7.27-7.33 (m, 1H), 7.19 (t, $J = 8.65$ Hz, 1H), 2.12 (s, 3H).

Example 9C

15 (E)-3-(3-chloro-4-fluorophenyl)-N-methoxy-N,2-dimethylacrylamide

A mixture of Example 9B (11.9 g, 55.45 mmol), *N,O*-dimethylhydroxylamine hydrochloride (5.7 g, 58 mmol), 1-hydroxybenzotriazole hydrate (7.88 g, 53.8 mmol), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (11 g, 57.58 mmol) and diisopropylethyl amine (47.4 g, 63.9 mL, 370 mol) was stirred in DMF (400 mL) at ambient
20 temperature overnight. The solution was diluted with water and extracted with ethyl acetate. The combined organic extract was washed with saturated sodium bicarbonate solution, 1N HCl, water, brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting material was chromatographed (SiO₂, 20% ethyl acetate/hexanes to 50% ethyl acetate/hexanes) to leave the title compound (6.85 g, 48% yield). ¹H NMR (300 MHz,
25 CDCl₃) δ ppm 7.39 (dd, $J = 7.12, 2.37$ Hz, 1H), 7.10-7.21 (m, 2H), 6.71 (s, 1H), 3.69 (s, 3H), 3.28 (s, 3H), 2.10 (s, 3H). MS (DCI⁺) 258 (M+H).

Example 9D

(E)-1-(3-chloro-4-fluorophenyl)-2-methylpent-1-en-3-one

30 Example 9C (2 g, 7.76 mmol) was dissolved in tetrahydrofuran (50 mL) and cooled to about -20°C. Ethylmagnesium bromide (1M in THF, 15.5 mL, 15.5 mmol) was added dropwise and the resulting mixture was stirred at about -20°C for 1 hr. The reaction was quenched with saturated NH₄Cl solution and extracted with ethyl acetate. The organic phase

was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting material was chromatographed (SiO₂, 10% ethyl acetate/hexanes to 50% ethyl acetate/hexanes) to afford the title compound (930 mg, 53% yield) as a yellow waxy solid.

¹H NMR (300 MHz, CDCl₃) δ ppm 7.45 (dd, *J* = 6.95, 2.20 Hz, 1H), 7.39 (s, 1H), 7.20-7.30 (m, 1H), 7.18 (t, *J* = 8.65 Hz, 1H), 2.81 (q, *J* = 7.23 Hz, 2H), 2.04 (d, *J* = 1.36 Hz, 3H), 1.17 (t, *J* = 7.29 Hz, 3H). MS (DCI⁺) 227 (M+H).

Example 9E

(1E,3E)-1-(3-chloro-4-fluorophenyl)-2-methylpent-1-en-3-one oxime

Example 9D (400 mg, 1.76 mmol) was dissolved in pyridine (5 mL), followed by addition of hydroxylamine hydrochloride (184 mg, 2.65 mmol). The resulting mixture was stirred at ambient temperature two hours. The solvent was evaporated in vacuo and the residue partitioned between 1N HCl and ethyl acetate. The organic phase was washed with water and brine, dried over Na₂SO₄, filtered, and evaporated in vacuo. The resulting solid was chromatographed (SiO₂, 20% ethyl acetate/hexanes), then recrystallized from hexanes to afford the title compound as a white solid (221 mg, 52% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 7.36 (dd, *J* = 6.95, 1.86 Hz, 1H), 7.10-7.21 (m, 3H), 6.79 (s, 2H), 2.68 (q, *J* = 7.80 Hz, 2H), 2.03 (s, 1H), 1.16 (t, *J* = 7.63 Hz, 3H).

Example 10

(1E,3Z)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime

To a solution of Example 1D (1.96 g, 10.2 mmol) in pyridine (40 mL) was added hydroxylamine hydrochloride (1.06 g, 15.3 mmol) in one portion. The reaction was stirred at ambient temperature for 12 hr, and then concentrated in vacuo. The residue was partitioned between EtOAc and 1M aq HCl. The separated organic phase was washed with 1M aq HCl and brine, dried (Na₂SO₄), filtered, and concentrated in vacuo. The crude product was purified by chromatography on silica gel (Analogix Intelliflash 280; 5% to 30% EtOAc/hexanes; SF40-115g column) and the late eluting component was isolated to afford the title compound as a white solid (0.16 g, 7.6% yield). ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 10.34 (s, 1H), 7.44-7.34 (m, 2H), 7.25-7.15 (m, 2H), 6.36 (s, 1H), 2.36 (q, *J* = 7.4 Hz, 2H), 1.98 (d, *J* = 1.5 Hz, 3H), 1.05 (t, *J* = 7.4 Hz, 3H). MS (DCI⁺): 208 (M+H).

Example 11

(2E,3E)-4-(4-fluorophenyl)but-3-en-2-one oxime

Example 11A

(E)-methyl 3-(4-fluorophenyl)acrylate

A mixture of 1-bromo-4-fluorobenzene (10 g, 57 mmol), methyl acrylate (4.5 g, 52 mmol), tetraethylammonium chloride (8.6 g, 52 mmol), *N*-cyclohexyl-*N*-methylcyclohexanamine (15 g, 78 mmol), and palladium (II) acetate (0.35 g, 1.6 mmol) in dimethylacetamide (200 mL) was heated to 100°C under an argon atmosphere. The reaction was allowed to proceed until the aryl halide was consumed. The reaction mixture was then cooled to room temperature, diluted with Et₂O, and washed three times with water. The separated organic phase was dried over MgSO₄ and concentrated in vacuo. The crude product was purified by flash column chromatography on silica gel eluting with EtOAc/hexanes to afford 8.5 g (91% yield) of the title compound.

Example 11B

(E)-3-(4-fluorophenyl)-*N*-methoxy-*N*-methylacrylamide

The title compound was prepared according to the procedure of Example 1C method B, substituting Example 11A for Example 1A.

Example 11C

(E)-4-(4-fluorophenyl)but-3-en-2-one

The title compound was prepared according to the procedure of Example 1D, substituting Example 11B for Example 1C and methylmagnesium bromide for ethylmagnesium bromide.

Example 11D

(2E,3E)-4-(4-fluorophenyl)but-3-en-2-one oxime

The title compound was prepared according to the procedure of Example 1E, substituting Example 11C for Example 1D. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 11.13 (s, 1H), 7.61 (dd, *J* = 8.7, 5.6 Hz, 2H), 7.18 (dd, *J* = 12.2, 5.6 Hz, 2H), 6.95 (d, *J* = 16.6 Hz, 1H), 6.80 (d, *J* = 16.6 Hz, 1H), 1.98 (s, 3H). MS (DCI⁺): 180 (M+H).

Example 12

(1E,3E)-1-(4-fluorophenyl)hex-1-en-3-one oxime

Example 12A

(E)-1-(4-fluorophenyl)hex-1-en-3-one

The title compound was prepared according to the procedure of Example 1D, substituting Example 11B for Example 1C and propylmagnesium bromide for ethylmagnesium bromide.

Example 12B

(1E,3E)-1-(4-fluorophenyl)hex-1-en-3-one oxime

The title compound was prepared according to the procedure of Example 1E, substituting Example 12A for Example 1D. ¹H NMR (500 MHz, DMSO/D₂O) δ ppm 7.61 (dd, *J* = 8.7, 5.6 Hz, 2H), 7.22–7.15 (m, 2H), 6.95 (d, *J* = 16.7 Hz, 1H), 6.72 (d, *J* = 16.6 Hz, 1H), 2.55–2.49 (m, 2H), 1.55–1.44 (m, 2H), 0.94 (t, *J* = 7.1 Hz, 3H). MS (ESI⁺): 208 (M+H).

5

Example 13

(1E,3E)-1-(4-fluorophenyl)pent-1-en-3-one oxime and (1E,3E)-1-(4-fluorophenyl)pent-1-en-3-one oxime, about 1:1 mixture

Example 13A

(E)-1-(4-fluorophenyl)pent-1-en-3-one

10

The title compound was prepared according to the procedure of Example 1D, substituting Example 11B for Example 1C.

Example 13B

(1E,3E)-1-(4-fluorophenyl)pent-1-en-3-one oxime and (1E,3Z)-1-(4-fluorophenyl)pent-1-en-3-one oxime, about 1:1 mixture

15

The title compound was prepared according to the procedure of Example 1E, substituting Example 13A for Example 1D. Data for the mixture obtained: ¹H NMR (500 MHz, DMSO/D₂O) δ ppm 7.67–7.59 (m, 3H), 7.36–7.30 (m, 2H), 7.25–7.16 (m, 4H), 7.04 (d, *J* = 16.9 Hz, 1H), 6.96 (d, *J* = 16.7 Hz, 1H), 6.69 (d, *J* = 16.7 Hz, 1H), 2.56–2.45 (m, 4H), 1.11 (t, *J* = 7.5 Hz, 3H), 1.04 (t, *J* = 7.5 Hz, 3H). MS (ESI⁺): 176 (M-17).

20

Example 14

(1E,3E)-1-(3,4-difluorophenyl)-2-methylpent-1-en-3-one oxime

The title compound was prepared according to the procedure of Example 1, substituting 3,4-difluorobenzaldehyde for 4-fluorobenzaldehyde. ¹H NMR (300 MHz, CDCl₃) δ ppm 8.14 (bs, 1H), 7.21–7.09 (m, 2H), 7.07–6.98 (m, 1H), 6.80 (s, 1H), 2.68 (q, *J* = 7.6 Hz, 2H), 2.03 (d, *J* = 1.3 Hz, 3H), 1.16 (t, *J* = 7.6 Hz, 3H). MS (DCI⁺): 226 (M+H).

25

Example 15

(1E,3E)-2-methyl-1-(3,4,5-trifluorophenyl)pent-1-en-3-one oxime

The title compound was prepared according to the procedure of Example 1, substituting 3,4,5-trifluorobenzaldehyde for 4-fluorobenzaldehyde. ¹H NMR (300 MHz, CDCl₃) δ ppm 8.00 (bs, 1H), 6.97–6.88 (m, 2H), 6.72 (s, 1H), 2.66 (q, *J* = 7.6 Hz, 2H), 2.03 (d, *J* = 1.3 Hz, 3H), 1.15 (t, *J* = 7.6 Hz, 3H). MS (DCI⁺): 244 (M+H).

30

Example 16

(1E,3E)-1-(4-fluorophenyl)hept-1-en-3-one oxime

Example 16A

(E)-1-(4-fluorophenyl)hept-1-en-3-one

The title compound was prepared according to the procedure of Example 1D, substituting Example 11B for Example 1C and butylmagnesium bromide for ethylmagnesium bromide.

5

Example 16B

(1E,3E)-1-(4-fluorophenyl)hept-1-en-3-one oxime

The title compound was prepared according to the procedure of Example 1E, substituting Example 16A for Example 1D. ¹H NMR (500 MHz, DMSO/D₂O) δ ppm 7.61 (dd, *J* = 8.7, 5.6 Hz, 2H), 7.22–7.16 (m, 2H), 6.94 (d, *J* = 16.7 Hz, 1H), 6.71 (d, *J* = 16.6 Hz, 1H), 2.58–2.50 (m, 2H), 1.49–1.40 (m, 2H), 1.40–1.30 (m, 2H), 0.90 (t, *J* = 7.6 Hz, 3H). MS (ESI⁺): 222 (M+H).

10

Example 17

(1E,3E)-1-(4-fluoro-3-methylphenyl)-2-methylpent-1-en-3-one oxime

The title compound was prepared according to the procedure of Example 1, substituting 4-fluoro-3-methylbenzaldehyde for 4-fluorobenzaldehyde. ¹H NMR (300 MHz, CDCl₃) δ ppm 8.06 (bs, 1H), 7.17–7.07 (m, 2H), 7.03–6.94 (m, 1H), 6.83 (s, 1H), 2.69 (q, *J* = 7.6, 2H), 2.29 (d, *J* = 2.0, 3H), 2.03 (d, *J* = 1.3, 3H), 1.17 (t, *J* = 7.6, 3H). MS (DCI⁺): 222 (M+H).

15

Example 18

(1E,3E)-1-[4-fluoro-3-(trifluoromethyl)phenyl]-2-methylpent-1-en-3-one oxime

The title compound was prepared according to the procedure of Example 1, substituting 4-fluoro-3-(trifluoromethyl)benzaldehyde for 4-fluorobenzaldehyde. ¹H NMR (300 MHz, CDCl₃) δ ppm 7.86 (s, 1H), 7.58–7.43 (m, 2H), 7.24–7.15 (m, 1H), 6.84 (s, 1H), 2.69 (q, *J* = 7.6 Hz, 2H), 2.02 (d, *J* = 1.3 Hz, 3H), 1.17 (t, *J* = 7.6 Hz, 3H). MS (DCI⁺): 276 (M+H).

20

25

Example 19

2-fluoro-5-[(1E,3E)-3-(hydroxyimino)-2-methylpent-1-enyl]benzonitrile

The title compound was prepared according to the procedure of Example 1, substituting 2-fluoro-5-formylbenzonitrile for 4-fluorobenzaldehyde. ¹H NMR (300 MHz, CDCl₃) δ ppm 8.03 (bs, 1H), 7.59–7.48 (m, 2H), 7.22 (t, *J* = 8.6 Hz, 1H), 6.78 (s, 1H), 2.68 (q, *J* = 7.6 Hz, 2H), 2.01 (d, *J* = 1.3 Hz, 3H), 1.16 (t, *J* = 7.6 Hz, 3H). MS (DCI⁺): 233 (M+H).

30


It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention,

which is defined solely by the appended claims and their equivalents. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art. Such changes and modifications, including without limitation those relating to the chemical structures, substituents, derivatives, intermediates, syntheses, formulations and/or methods of use of the invention, may be made without departing from the spirit and scope thereof.

5

Claims:

1. A method for treating a disorder or condition that is ameliorated by the inhibition of TRPA1, comprising administering to the subject a therapeutically effective amount of a compound of formula (I) or a solvate, a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or any combination thereof, with or without one or more pharmaceutically acceptable carriers



wherein

R^1 represent optional substituents of the phenyl group and each R^1 is independently alkyl, alkenyl, alkynyl, -CN, halogen, -OR^a, -NO₂, -N(R^a)(R^b), -N(R^b)C(O)R^a, -N(R^b)S(O)₂R^{1a}, -N(R^b)C(O)OR^a, -N(R^b)C(O)N(R^a)(R^b), -N(R^b)S(O)₂N(R^a)(R^b), -C(O)R^a, -C(O)OR^a, -C(O)N(R^a)(R^b), -SR^a, -SF₅, -S(O)R^{1a}, -S(O)₂R^{1a}, -S(O)₂OR^{1a}, -S(O)₂N(R^a)(R^b), -(CR^dR^e)_q-CN, haloalkyl, -(CR^dR^e)_q-OR^a, -(CR^dR^e)_q-NO₂, -(CR^dR^e)_q-N(R^a)(R^b), -(CR^dR^e)_q-N(R^b)C(O)R^a, -(CR^dR^e)_q-N(R^b)S(O)₂R^{1a}, -(CR^dR^e)_q-N(R^b)C(O)OR^a, -(CR^dR^e)_q-N(R^b)C(O)N(R^a)(R^b), -(CR^dR^e)_q-N(R^b)S(O)₂N(R^a)(R^b), -(CR^dR^e)_q-C(O)R^a, -(CR^dR^e)_q-C(O)OR^a, -(CR^dR^e)_q-C(O)N(R^a)(R^b), -(CR^dR^e)_q-S(O)₂R^{1a}, -(CR^dR^e)_q-S(O)₂OR^{1a}, or -(CR^dR^e)_q-S(O)₂N(R^a)(R^b);

R^2 is hydrogen, C₁₋₆ alkyl, or haloalkyl;

R^3 is C₁₋₆ alkyl, haloalkyl, or cyclopropyl, wherein the cyclopropyl is optionally substituted with 1, 2, 3, 4, or 5 substituents as represented by R^j ;

R^j , at each occurrence, is independently C₁₋₄ alkyl, halogen, or haloalkyl;

Y is -OH, -O(C₁₋₄ alkyl), -N(R^f)(R^g), -N(R^f)C(O)R^g, -N(R^f)S(O)₂R^g, or -N(R^f)C(O)N(R^f)(R^g);

R^a and R^b , at each occurrence, are each independently hydrogen, alkyl, or haloalkyl;

R^{1a} , at each occurrence, is independently alkyl or haloalkyl;

R^d and R^e , at each occurrence, are each independently hydrogen, alkyl, halogen, or haloalkyl;

R^f and R^g , at each occurrence, are each independently hydrogen, alkyl, or haloalkyl;

m is 0, 1, 2, 3, 4, or 5; and

q is 1, 2, 3, or 4;


with the proviso that the compound is not (Z)-4-(4-chlorophenyl)-3-methylbut-3-en-2-oxime or (E)-4-(4-chlorophenyl)-3-methylbut-3-en-2-oxime.

2. The method of claim 1 wherein R^3 is C_{1-6} alkyl.
3. The method of claim 1 wherein R^3 is methyl, ethyl, n-propyl, tert-butyl, n-butyl, or isopropyl.
4. The method of claim 1 wherein
 R^3 is C_{1-6} alkyl; and
 R^2 is C_{1-6} alkyl.
5. The method of claim 1 wherein
 R^3 is C_{1-6} alkyl;
 R^2 is C_{1-6} alkyl; and
Y is OH.
6. The method of claim 1 wherein
 R^3 is C_{1-6} alkyl;
 R^2 is C_{1-6} alkyl; and
Y is $-O(C_{1-4}$ alkyl).
7. The method of claim 1 wherein
 R^3 is C_{1-6} alkyl; and
 R^2 is hydrogen.
8. The method of claim 1 wherein
 R^3 is C_{1-6} alkyl;
 R^2 is hydrogen; and
Y is OH.
9. The method of claim 1 where
 R^3 is C_{1-6} alkyl;
 R^2 is hydrogen; and

Y is $-\text{O}(\text{C}_{1-4} \text{ alkyl})$.

10. The method of claim 1 wherein Y is $-\text{N}(\text{R}^f)(\text{R}^g)$, $-\text{N}(\text{R}^f)\text{C}(\text{O})\text{R}^g$, $-\text{N}(\text{R}^f)\text{S}(\text{O})_2\text{R}^g$, or $-\text{N}(\text{R}^f)\text{C}(\text{O})\text{N}(\text{R}^f)(\text{R}^g)$.
11. The method of claim 1 wherein Y is
Y is $-\text{N}(\text{R}^f)(\text{R}^g)$, $-\text{N}(\text{R}^f)\text{C}(\text{O})\text{R}^g$, $-\text{N}(\text{R}^f)\text{S}(\text{O})_2\text{R}^g$, or $-\text{N}(\text{R}^f)\text{C}(\text{O})\text{N}(\text{R}^f)(\text{R}^g)$;
 R^3 is C_{1-6} alkyl; and
 R^2 is C_{1-6} alkyl.
12. The method of claim 1 wherein Y is
Y is $-\text{N}(\text{R}^f)(\text{R}^g)$, $-\text{N}(\text{R}^f)\text{C}(\text{O})\text{R}^g$, $-\text{N}(\text{R}^f)\text{S}(\text{O})_2\text{R}^g$, or $-\text{N}(\text{R}^f)\text{C}(\text{O})\text{N}(\text{R}^f)(\text{R}^g)$;
 R^3 is C_{1-6} alkyl; and
 R^2 is hydrogen.
13. The method of claim 1 wherein the compound is
(1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime;
(1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one O-methyloxime;
(1E,3E)-1-(2-fluorophenyl)-2-methylpent-1-en-3-one oxime;
(1E,3E)-1-(4-fluorophenyl)-2-methylhex-1-en-3-one oxime;
(1E,3E)-1-(4-fluorophenyl)-2,4,4-trimethylpent-1-en-3-one oxime;
(1E,3E)-1-(4-fluorophenyl)-2-methylhept-1-en-3-one oxime;
(1E,3Z)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime;
(1E,3E)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime;
(1E,3E)-1-(3-chloro-4-fluorophenyl)-2-methylpent-1-en-3-one oxime;
(1E,3Z)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one oxime;
(2E,3E)-4-(4-fluorophenyl)but-3-en-2-one oxime;
(1E,3E)-1-(4-fluorophenyl)hex-1-en-3-one oxime;
(1E,3E)-1-(4-fluorophenyl)pent-1-en-3-one oxime;
(1E,3Z)-1-(4-fluorophenyl)pent-1-en-3-one oxime;
(1E,3E)-1-(3,4-difluorophenyl)-2-methylpent-1-en-3-one oxime; or
(1E,3E)-2-methyl-1-(3,4,5-trifluorophenyl)pent-1-en-3-one oxime.
14. The method of claim 1 wherein the disorder or condition is pain.

15. The method of claim 1 wherein the disorder or condition is selected from the group consisting of acute cerebral ischemia, chronic pain, neuropathic pain, nociceptive pain, allodynia, inflammatory pain, post herpetic neuralgia, neuropathies, neuralgia, diabetic neuropathy, HIV-related neuropathy, nerve injury, rheumatoid arthritic pain, osteoarthritic pain, burns, back pain, visceral pain, cancer pain, dental pain, headache, migraine, carpal tunnel syndrome, fibromyalgia, neuritis, sciatica, pelvic hypersensitivity, pelvic pain, menstrual pain, bladder disease, micturition disorder, renal colic; inflammation, neurodegenerative disease, pulmonary disease, gastrointestinal disease, emesis, and obesity.
16. The method of claim 14 further comprising the step of co-administering with one or more second pain reducing agent.
17. The method of claim 16 wherein the second pain reducing agent is acetaminophen, a nonsteroidal anti-inflammatory drug, or combination thereof.
18. The method of claim 17 wherein the nonsteroidal anti-inflammatory drug is ibuprofen.
19. The method of claim 15 further comprising the step of co-administering with one or more second pain reducing agent.
20. A pharmaceutical composition comprising a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof



wherein

R^1 represent optional substituents of the phenyl group and each R^1 is independently alkyl, alkenyl, alkynyl, -CN, halogen, -OR^a, -NO₂, -N(R^a)(R^b), -N(R^b)C(O)R^a, -N(R^b)S(O)₂R^{1a}, -N(R^b)C(O)OR^a, -N(R^b)C(O)N(R^a)(R^b), -N(R^b)S(O)₂N(R^a)(R^b), -C(O)R^a, -C(O)OR^a, -C(O)N(R^a)(R^b), -SR^a, -SF₅, -S(O)R^{1a}, -S(O)₂R^{1a}, -S(O)₂OR^{1a}, -S(O)₂N(R^a)(R^b), -(CR^dR^e)_q-CN, haloalkyl, -(CR^dR^e)_q-OR^a, -(CR^dR^e)_q-NO₂, -(CR^dR^e)_q-N(R^a)(R^b),

$-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-N}(\text{R}^{\text{b}})\text{C}(\text{O})\text{R}^{\text{a}}$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-N}(\text{R}^{\text{b}})\text{S}(\text{O})_2\text{R}^{1\text{a}}$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-N}(\text{R}^{\text{b}})\text{C}(\text{O})\text{OR}^{\text{a}}$,
 $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-N}(\text{R}^{\text{b}})\text{C}(\text{O})\text{N}(\text{R}^{\text{a}})(\text{R}^{\text{b}})$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-N}(\text{R}^{\text{b}})\text{S}(\text{O})_2\text{N}(\text{R}^{\text{a}})(\text{R}^{\text{b}})$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-C}(\text{O})\text{R}^{\text{a}}$,
 $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-C}(\text{O})\text{OR}^{\text{a}}$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-C}(\text{O})\text{N}(\text{R}^{\text{a}})(\text{R}^{\text{b}})$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-S}(\text{O})_2\text{R}^{1\text{a}}$, $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-S}(\text{O})_2\text{OR}^{1\text{a}}$,
 or $-(\text{CR}^{\text{d}}\text{R}^{\text{e}})_q\text{-S}(\text{O})_2\text{N}(\text{R}^{\text{a}})(\text{R}^{\text{b}})$;

R^2 is hydrogen, C_{1-6} alkyl, or haloalkyl;

R^3 is C_{1-6} alkyl, haloalkyl, or cyclopropyl, wherein the cyclopropyl is optionally substituted with 1, 2, 3, 4, or 5 substituents as represented by R^{j} ;

R^{j} , at each occurrence, is independently C_{1-4} alkyl, halogen, or haloalkyl;

Y is $-\text{OH}$, $-\text{O}(\text{C}_{1-4} \text{ alkyl})$, $-\text{N}(\text{R}^{\text{f}})(\text{R}^{\text{g}})$, $-\text{N}(\text{R}^{\text{f}})\text{C}(\text{O})\text{R}^{\text{g}}$, $-\text{N}(\text{R}^{\text{f}})\text{S}(\text{O})_2\text{R}^{\text{g}}$, or
 $-\text{N}(\text{R}^{\text{f}})\text{C}(\text{O})\text{N}(\text{R}^{\text{f}})(\text{R}^{\text{g}})$;

R^{a} and R^{b} , at each occurrence, are each independently hydrogen, alkyl, or haloalkyl;

$\text{R}^{1\text{a}}$, at each occurrence, is independently alkyl or haloalkyl;

R^{d} and R^{e} , at each occurrence, are each independently hydrogen, alkyl, halogen, or haloalkyl;

R^{f} and R^{g} , at each occurrence, are each independently hydrogen, alkyl, or haloalkyl;

m is 0, 1, 2, 3, 4, or 5; and

q is 1, 2, 3, or 4;

with the proviso that the compound is not (Z)-4-(4-chlorophenyl)-3-methylbut-3-en-2-oxime or (E)-4-(4-chlorophenyl)-3-methylbut-3-en-2-oxime;
 and a pharmaceutically acceptable carrier.

21. The pharmaceutical composition of claim 20 further comprising one or more second pain reducing agent.

22. A compound that is

(1E,3E)-1-(4-fluorophenyl)-2-methylpent-1-en-3-one O-methyloxime;

(1E,3E)-1-(2-fluorophenyl)-2-methylpent-1-en-3-one oxime;

(1E,3E)-1-(4-fluorophenyl)-2-methylhex-1-en-3-one oxime;

(1E,3E)-1-(4-fluorophenyl)-2,4,4-trimethylpent-1-en-3-one oxime;

(1E,3E)-1-(4-fluorophenyl)-2-methylhept-1-en-3-one oxime;

(1E,3Z)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime;

(1E,3E)-1-(4-fluorophenyl)-2,4-dimethylpent-1-en-3-one oxime;

(1E,3E)-1-(3-chloro-4-fluorophenyl)-2-methylpent-1-en-3-one oxime;

(1E,3E)-1-(4-fluorophenyl)hex-1-en-3-one oxime;

WO 2009/089082**PCT/US2009/030005**

(1E,3E)-1-(3,4-difluorophenyl)-2-methylpent-1-en-3-one oxime; or
(1E,3E)-2-methyl-1-(3,4,5-trifluorophenyl)pent-1-en-3-one oxime;
or pharmaceutically acceptable salts thereof.

Figure 1




Figure 1

