

(19) **DANMARK**

(10) **DK/EP 4331584 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 31/428 (2006.01)** **A 61 P 1/00 (2006.01)** **A 61 P 1/16 (2006.01)**
A 61 P 11/00 (2006.01) **A 61 P 13/12 (2006.01)** **A 61 P 37/00 (2006.01)**
C 07 D 419/12 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2025-03-17**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2025-01-01**
- (86) Europæisk ansøgning nr.: **23201458.9**
- (86) Europæisk indleveringsdag: **2019-07-26**
- (87) Den europæiske ansøgnings publiceringsdag: **2024-03-06**
- (30) Prioritet: **2018-07-27 FR 1857021**
- (62) Stamansøgningsnr: **19759658.8**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Inventiva, 50 rue de Dijon, 21121 Daix, Frankrig**
- (72) Opfinder: **MONTALBETTI, Christian, , FONTAINE-LES-DIJON, Frankrig**
BOUBIA, Benaissa, , SAINT APOLLINAIRE, Frankrig
- (74) Fuldmægtig i Danmark: **Denнемeyer & Associates S.A, P.O. Box 700425, DE-81304 Munich, Tyskland**
- (54) Benævnelse: **TERAPEUTISKE ANVENDELSER AF DEUTEREREDE DERIVATER AF LANIFIBRANOR**
- (56) Fremdragne publikationer:
BOUBIA ET AL.: "DESIGN, SYNTHESIS AND EVALUATION OF A NOVEL SERIES OF INDOLE SULFONAMIDE PEROXISOME PROLIFERATOR ACTIVATED RECEPTOR (PPAR) ALPHA/GAMMA/DELTA TRIPLE ACTIVATORS: DISCOVERY OF LANIFIBRANOR A NEW ANTIFIBROTIC CLINICAL CANDIDATE", MEDICINAL CHEMISTRY, vol. 61, 15 February 2018 (2018-02-15), pages 2246 - 2265, XP002790345
KATSNELSON A.: "HEAVY DRUGS DRAW HEAVY INTEREST FROM PHARMA BACKERS", NATURE MEDICINE, vol. 19, no. 6, 1 June 2013 (2013-06-01), pages 656, XP002790346

THERAPEUTIC USES OF DEUTERATED DERIVATIVES OF LANIFIBRANOR

Field of the invention

The present invention relates to deuterated derivatives of lanifibranor, in particular for their use in therapy, in particular in the treatment of fibrotic diseases.

State of the art


Lanifibranor or 4-(1-(1,3-Benzothiazol-6-ylsulfonyl)-5-chloro-indol-2-yl)-butanoic acid is a PPAR receptor agonist useful in particular for its use in the treatment of systemic sclerosis (SSc) and in non-alcoholic steatohepatitis (NASH). Lanifibranor is in particular described with other indole derivatives in patent application WO 2007/026097, in particular for the prevention or treatment of hypertriglyceridemia, hypercholesterolemia and, more generally, for re-establishing normal parameters during disruption of lipid and carbohydrate metabolism, and also in the case of the treatment of endothelial dysfunction, inflammatory diseases or neurodegeneration.

The interest of using deuterated molecules is known as a means of modifying their metabolism by improving their metabolic stability and possibly their cell penetration properties (Maehr & al., J. Med. Chem. 2013, 56, 3878–3888; Kerekes et al., J. Med. Chem. 2011, 54, 201–210; Harbeson & Tung, Medchem News 2014, 2 9-22; Gant, J. Med. Chem. 2014, 57, 3595–3611; DeWitt & Maryanoff, Biochemistry 2018, 57, 472-473; WO 2017/136375; WO 2018/039521). The synthesis of deuterated drugs is also described in several articles (Modutwa & al., J. Label Compd. Radiopharm 2010, 53 686–692; Junk et al., J. Label Compd. Radiopharm 1997, 39 625-630).

However, it is recognized that the effect of hydrogen substitutions by deuteriums on the metabolism of chemicals remains unpredictable and for many authors deuteration of drugs is not considered a reliable means of modulation (Foster, AB, Adv Drug Res 1985, 14: 1-40; Fisher, MB et al, Curr Opin Drug Discov Devel, 2006, 9: 101-09; Katsnelson, Nature Medecine 2013, 19 6 656).

Description of the invention

The invention relates to a deuterated derivative of lanifibranor of formula (I):




wherein at least one of the groups R_1 to R_7 is a deuterium atom (D) and the other groups R_1 to R_7 are hydrogen atoms (H).

The invention also relates to a composition, in particular a pharmaceutical composition, which comprises at least one deuterated derivative of formula (I).

The invention also relates to the deuterated derivatives of formula (I) for their use in therapy, in particular for the treatment of fibrotic diseases.

Detailed description of the invention

The invention relates to a deuterated derivative of lanifibranor of formula (I):



wherein at least one of the groups R_1 to R_7 is a deuterium atom and the other groups R_1 to R_7 are hydrogen atoms, and its pharmaceutically acceptable salts and solvates.

According to a first embodiment of the invention, at least the group R_1 is D.

In particular, the deuterated derivative of lanifibranor is 4-(1-(2-deuterio-1,3-benzothiazol-6-yl)sulfonyl)-5-chloro-1H-indol-2-yl)butanoic acid.

According to another embodiment of the invention, at least one of the groups R_2 to R_7 is D. More preferably at least one of the groups R_2 and R_3 and/or at least one of the groups R_4 and R_5 and/or at least one of the groups R_6 and R_7 is D. Even more preferably, R_2 , R_3 , R_4 , R_5 , R_6 and R_7 are D.

In particular, the deuterated derivative of lanifibranor is 4-[1-(1,3-benzothiazol-6-ylsulfonyl)-5-chloro-indol-2-yl]-2,2,3,3,4,4-hexadeuteriobutanoic acid.

The pharmaceutically acceptable salts of the deuterated derivatives according to the invention are the same as the pharmaceutically acceptable salts of lanifibranor, in particular by combining the acid with a pharmaceutically acceptable non-toxic inorganic or organic base. Among the inorganic bases, it is possible to use, for example, the hydroxides of sodium, potassium, magnesium or calcium. Among the organic bases, use may be made, for example, of amines, amino alcohols, basic amino acids such as lysine or arginine or else compounds carrying a quaternary ammonium function such as, for example, betaine or choline.

The compounds according to the invention are prepared according to the usual methods for preparing lanifibranor, for example that described in patent application WO 2007/026097, the deuterated synthesis intermediates replacing the same isotopically unenriched intermediates used in these usual methods.

The invention also relates to a composition comprising a deuterated derivative according to the invention, or one of its salts or solvates, in particular a pharmaceutically acceptable salt or solvate, and a carrier suitable for its use, for the above-mentioned use.

For use in therapy, the composition according to the invention is advantageously a pharmaceutical composition, the carrier comprising usual excipients from the Pharmacopoeia, chosen according to the mode of administration envisaged.

Such compositions are known to those skilled in the art, and in particular described in patent applications WO 2007/026097 and WO 2015/189401.

For oral administration, for example in the form of a tablet, capsule, lozenge, gel, syrup, oral suspension, the pharmaceutical composition according to the invention advantageously comprises from 1 to 1000 mg of deuterated derivative of formula (I), for example 1 mg, 5 mg, 10 mg, 50 mg, 100 mg, 200 mg, 500 mg or 1000 mg.

The invention also relates to the deuterated derivatives of formula (I), or their pharmaceutically acceptable salts or solvates, for their use in therapy.

The different therapeutic uses of the deuterated derivatives according to the invention are those known for lanifibranor and its analogues, such as those described in patent applications WO 2007/026097 and WO 2015/189401.

The invention therefore relates to deuterated derivatives of formula (I), or their pharmaceutically acceptable salts or solvates, for their use to fight hypercholesterolemia, hyperlipidemia, hypertriglyceridemia, dyslipidemia, insulin resistance, diabetes or obesity as well as cardiovascular diseases which are the consequence of a serum lipoprotein

imbalance. The compounds according to the invention are also useful as active principles of medicaments intended to prevent or treat diseases associated with an endothelial dysfunction, atherosclerosis, myocardial infarction, hypertension, cerebrovascular problems, certain inflammatory diseases such as example rheumatoid arthritis, and neurodegeneration such as in particular Alzheimer's disease or Parkinson's disease.

The invention relates in particular to the deuterated derivatives of formula (I) for their use in the treatment of fibrotic diseases. In one embodiment, fibrotic disease is a condition affecting any organ which can develop fibrosis, such as the heart, the lung, the liver, the kidney, the gastrointestinal tract, the skin, muscles, etc. The fibrotic disease is in particular chosen from: liver fibrosis, liver steatosis, non-alcoholic steatohepatitis, chronic kidney disease, a pulmonary fibrotic disorder such as idiopathic pulmonary fibrosis, and systemic sclerosis.

The invention also relates to a method of treating a fibrotic disease as defined above in a patient awaiting such treatment which comprises the administration to this patient of a deuterated derivative of formula (I), in particular by means of a pharmaceutical composition comprising said deuterated derivative of formula (I) in a form suitable for the chosen mode of administration.

The need for patient treatment will have advantageously been previously determined by any appropriate method of diagnosis for the detection of the fibrotic disease to be treated, in particular by the analysis of the expression levels of PPAR receptors in the fibrotic tissues of the patients to treat.

Examples

Abbreviations

APCI = atmospheric-pressure chemical ionization

DCM = dichloromethane

DMSO = dimethyl sulfoxide

eq. = equivalent

EtOAc = ethyl acetate

EtOH = ethanol

h = hour

HPLC = high performance liquid chromatography

LCMS = liquid chromatography coupled with mass spectrometry

MeOH = methanol

min = minute

$\text{Pd(PPh}_3)_4$ = tetrakis(triphenylphosphine)palladium(0)

pTsOH = paratoluenesulfonic acid

NMR = nuclear magnetic resonance

THF = tetrahydrofuran

Example 1: 4-[5-Chloro-1-[(2-deuterio-1,3-benzothiazol-6-yl)sulfonyl]]indol-2-yl]butanoic acid

Preparation 1: 2-Deuterio-1,3-benzothiazole

n-BuLi (37.0 mL, 92.5 mmol, 2.5 eq.; 2.5 M solution in hexane) was added dropwise to 1,3-benzothiazole (5.0 g, 36.9 mmol, 1.0 eq.) dissolved in THF (80 mL), and cooled to -78°C. After complete addition, D₂O (4 mL, 222 mmol, 6.0 eq.) was added to the reaction mixture at -78°C, the whole was stirred for 30 min then brought to room temperature. The reaction was monitored by LCMS analysis. The reaction mixture was then treated with a saturated aqueous NH₄Cl solution and the whole was extracted 3 times with DCM. The organic phases were combined and dried over MgSO₄ then filtered and evaporated under reduced pressure to yield a crude product purified by flash chromatography (eluent: 0 to 40% EtOAc in *n*-heptane) to give the compound of **preparation 1** (3.22 g, 64% yield) as a brown oil. APCI MS *m/z* 137 [M+H]⁺; HPLC-MS (220-254nm) purity: 99%. ¹H NMR (300 MHz DMSO-*d*₆): δ 7.46-7.58 (2H, m); 8.08-8.19 (2H, m).

Preparation 2: 2-Deuterio-6-nitro-1,3-benzothiazole

2-Deuterio-1,3-benzothiazole (**preparation 1**, 2.9 g, 21.3 mmol, 1.0 eq.) was added slowly to a solution of sulfuric acid (13 mL) at 0°C, nitric acid (6.4 mL) was then added dropwise while maintaining the temperature below 0°C. The reaction mixture was then brought to room temperature and stirred for 12 h before being hydrolysed over a mixture of iced water. The yellow precipitate was filtered off, washed with water and crystallized from ethanol to yield the desired product (**preparation 2**, 1.4 g, yield 37%) in the form of yellow crystals. APCI MS *m/z* 182 [M+H]⁺; HPLC-MS (220-254nm) purity > 99%. ¹H NMR (300 MHz DMSO-*d*₆): δ 8.27-8.38 (2H, m); 9.25 (1H, d, *J* = 2.3Hz).

Preparation 3: 2-Deuterio-1,3-benzothiazol-6-amine

Tin chloride powder (7.3 g, 38.6 mmol, 3.5 eq.) was added to the compound of **preparation 2** (2.0 g, 11.0 mmol, 1.0 eq.) dissolved in a 1:1 EtOH - EtOAc mixture (60 mL), and the solution was stirred at room temperature for 4 h. At the end of the reaction, the reaction mixture was filtered through a Celite™ cartridge and extracted with a saturated Na₂CO₃ solution. The organic phases were combined and dried over MgSO₄ then filtered and evaporated under reduced pressure to yield a crude product purified by

flash chromatography (eluent: 0 to 20% MeOH in DCM) to obtain the desired compound (**preparation 3**, 970 mg, yield 58%) as brown crystals. APCI MS m/z 152 $[M+H]^+$; HPLC-MS (220-254nm) purity: 97%. ^1H NMR (300 MHz DMSO- d_6): δ 5.38 (2H, s); 6.8 (1H, dd, $J = 8.7\text{Hz}$ and $J = 2.2\text{Hz}$); 7.11 (1H, d, $J = 2.2\text{Hz}$); 7.7 (1H, d, $J = 8.7\text{Hz}$).

Preparation 4: 2-Deuterio-1,3-benzothiazol-6-sulfonyl chloride

An aqueous solution (4.5mL) of thionyl chloride (1.1 mL) was prepared at 0°C and stored overnight at 4°C. To this solution copper (I) chloride (0.05 eq.) was added at -10°C. The compound of **preparation 3** (450 mg, 2.9 mmol, 1.0 eq.) was dissolved in hydrochloric acid (3.5 mL) while maintaining the temperature below 25°C, the latter solution having then been cooled to -10°C and a solution of sodium nitrite (3.3 mmol, 0.7 mL, 1.1 eq.) was added without the temperature exceeding -2°C. The mixture thus obtained was stirred for 15 min at -2°C and then added dropwise to the first solution of thionyl chloride at -5°C. The reaction mixture was then stirred at -5°C for 3 h then the reaction was hydrolysed in water. The precipitate formed was filtered and then washed with water to yield the desired product (**preparation 4**, 255 mg, yield 37%) in the form of light brown crystals. This compound was used in the next step without any purification. HPLC-MS (220-254nm) purity: 94%.

Preparation 5: N-(4-Chloro-2-iodophenyl)-2-deuterio-1,3-benzothiazole-6-sulfonamide

Under argon, 4-chloro-2-iodoaniline (564 mg, 2.2 mmol, 1.0 eq.) was dissolved in anhydrous pyridine (8 mL) and the compound of **preparation 4** (600 mg, 2.5 mmol, 1.15 eq.) was added. The solution was then stirred at room temperature for 2 h. The solvent was evaporated and the reaction crude was purified by flash chromatography (eluent: 0 to 30% EtOAc in n-heptane) to yield the desired product (**preparation 5**, 0.73 g, yield 73%) in the form of brown crystals. APCI MS m/z 452 $[M+H]^+$; HPLC-MS (220-254nm) purity: 94%. ^1H NMR (300 MHz DMSO- d_6): δ 7.01 (1H, d, $J = 8.5\text{Hz}$); 7.4 (1H, dd, $J = 8.5\text{Hz}$ and $J = 2.3\text{Hz}$); 7.82 (1H, dd, $J = 8.7\text{Hz}$ and $J = 1.8\text{Hz}$); 7.88 (1H, d, $J = 2.4\text{Hz}$); 8.26 (1H, d, $J = 8.7\text{Hz}$); 8.6 (1H, d, $J = 1.5\text{Hz}$); 10.03 (1H, s).

Preparation 6: 4-[5-Chloro-1-[(2-deuterio-1,3-benzothiazol-6-yl)sulfonyl]indol-2-yl]butanoic acid

Under argon the compound of **preparation 5** (1.31 g, 2.9 mmol, 1.0 eq.) was dissolved in anhydrous DMF (30 mL), then 5-hexynoic acid (360 mg, 3.2 mmol, 1.1 eq.), copper iodide (56 mg, 0.3 mmol, 0.1 eq.), $Pd(PPh_3)_4$ (168 mg, 0.15 mmol, 0.05 eq.) and triethylamine (606 μL , 4.4 mmol, 1.5 eq.) were added and the reaction mixture was stirred at 80°C for 2 h. At the end of the reaction, the mixture was taken up in a 1N aqueous

hydrochloric acid solution and extracted twice with ethyl acetate. The organic phases were combined and dried over MgSO_4 then filtered and evaporated under reduced pressure to yield a crude product purified by flash chromatography (eluent: 0 to 5% MeOH in DCM) to yield the desired product (**Example 1**, 1.05 g, yield 83%) in the form of brown crystals. APCI MS m/z 452 $[\text{M}+\text{H}]^+$; HPLC-MS (220-254nm) purity: 94%. ^1H NMR (300 MHz DMSO-d_6): δ 1.86-2.01 (2H, m); 2.29-2.41 (2H, m); 3.08 (2H, t, $J = 7.2\text{Hz}$); 6.61 (1H, s); 7.31 (1H, dd, $J = 8.7\text{Hz}$ and $J = 2.0\text{Hz}$); 7.51 (1H, d, $J = 2.0\text{Hz}$); 7.84 (1H, dd, $J = 8.7\text{Hz}$ and $J = 1.8\text{Hz}$); 8.08 (1H, d, $J = 8.7\text{Hz}$); 8.19 (1H, d, $J = 8.7\text{Hz}$); 8.97 (1H, d, $J = 1.8\text{Hz}$); 12.13 (1H, s). Incorporation rate of deuterium assayed by 500 MHz NMR: 98%.

Example 2: 4-[1-(1,3-Benzothiazol-6-ylsulfonyl)-5-chloro-indol-2-yl]-2,2,3,3,4,4-hexadeuteriobutanoic acid

Preparation 7: 2-(1,1,2,2,3,3,4,4-octadeuterio-4-iodo-butoxy)tetra-hydropyrane

Trimethylsilyl iodide (25 g, 125 mmol, 1 eq.) was added at 0°C to THF- D_8 (10 g, 125 mmol, 1 eq.). After stirring for 2 h at 0°C , ether (80 mL) and water (20 mL) were added. The mixture was stirred for 3 h then decanted. The organic phase was then dried over MgSO_4 , filtered and concentrated under reduced pressure. The reaction crude thus obtained was taken up in DCM (100 mL) at 0°C , then dihydropyran (12.5 mL, 150 mmol, 1.2 eq.) and pTsOH (50 mg, 0.2 mmol, 0.002 eq.) were added. The reaction mixture was then stirred at 0°C overnight. The solution was washed successively with a saturated solution of NaHCO_3 ($3 \times 15\text{ mL}$), then with brine and then dried over Na_2SO_4 before being concentrated under reduced pressure. The reaction crude was purified by flash chromatography (eluent: 1:20 to 1:5 EtOAc:Hexane) to yield the desired product (**Preparation 7**, 23 g, 62% yield) in the form of a colourless oil which was used directly in the next step.

Preparation 8: 2-(1,1,2,2,3,3,4,4-octadeuteriohex-5-ynoxy)tetrahydropyran

To a solution of lithium acetylide (19.94 g, 186.3 mmol, 1 eq.) in freshly distilled DMSO (60mL) was added dropwise at 5°C a solution of the compound of **preparation 7** (30 g, 102.7 mmol, 0.55 eq.) in DMSO (30 mL). This solution thus obtained was stirred for 2 h then hydrolysed with a solution of NH_4Cl at 0°C . This mixture was then extracted with hexane and washed with a solution of copper sulfate (5%, 20 mL) then with brine (20 mL). The organic phases were combined and dried over Na_2SO_4 before being concentrated under reduced pressure, to yield the desired product (**Preparation 8**, 10.7 g, yield 55%) in the form of a yellow oil which was used directly in the next step without any purification.

Preparation 9: 1,1,2,2,3,3,4,4-octadeuteriohex-5-yn-1-ol

To a solution of the compound of **preparation 8** (10.7 g, 56.2 mmol, 1 eq.) in a mixture of THF (12.5 mL) and MeOH (350 mL) at 0°C, was added pTsOH in small portions (360 mg, 1.82 mmol, 0.03 eq.) and the reaction mixture thus obtained was stirred overnight. The mixture was then washed with a saturated solution of NaHCO₃ (3 × 15 mL) and brine (15 mL), then dried over Na₂SO₄ before being concentrated under reduced pressure. The reaction crude was purified by flash chromatography (eluent: 1:20 to 1:5 EtOAc:Hexane) to yield the desired product (**Preparation 9**, 5 g, 82% yield) in the form of a colourless oil which was used directly in the next step.

Preparation 10: 2,2,3,3,4,4-hexadeuteriohex-5-ynoic acid

To a solution of the compound of **preparation 9** (9.25 g, 87 mmol, 1 eq.) in acetone (87 mL) was added, under stirring, Jones's reagent (CrO₃ (17.45 g, 174.5 mmol, 2 eq.) in 10N H₂SO₄ (218 mL)) over 5 min at 0°C. The reaction mixture was then stirred for 1 h before being concentrated under reduced pressure. Then ether (130 mL) and water (4 mL) were added immediately. The solid obtained was filtered off and the filtrate was extracted with ether (6 × 100 mL). The organic phases were combined and washed with water and then dried over Na₂SO₄ before being filtered and concentrated under reduced pressure to yield the desired product (**Preparation 10**, 9.17 g, yield 89%) in the form of an orange oil which was used directly in the next step without any purification.

Preparation 11: 4-[1-(1,3-Benzothiazol-6-ylsulfonyl)-5-chloro-indol-2-yl]-2,2,3,3,4,4-hexadeuteriobutanoic acid

A mixture of benzothiazol-6-sulfonic acid (4-chloro-2-iodo-phenyl)-amide (16 g, 35.5 mmol, 1 eq.), hex-5-ynoic acid (**preparation 10**, 5.2 g, 44 mmol, 1.24 eq.), copper iodide (335 mg, 1.75 mmol, 0.05 eq.), dichlorobis(triphenylphosphine)palladium (1.24 g, 1.72 mmol, 0.05 eq.), triethylamine (130 mL) and N,N-dimethylformamide (130 mL) was stirred under nitrogen at 110°C for 1 h. The reaction mixture was then diluted with a 1M HCl solution and extracted with ethyl acetate, the insoluble matters were filtered off and the filtrate was dried over Na₂SO₄ before being concentrated under reduced pressure. The reaction crude was purified by flash chromatography and then recrystallized from DCM to yield the desired product (**Example 2**, 2.3 g, yield 15%) in the form of a white solid. Incorporation rate of deuterium assayed by 500 MHz NMR: > 99%. ¹H NMR (400 MHz, DMSO-d₆) δ 6.62 (1H, d, J = 0.6 Hz), 7.32 (1H, dd, J = 8.9 and 2.2 Hz), 7.57 (1H, d, J = 2.1 Hz), 7.85 (1H, dd, J = 8.7 and 2.1 Hz), 8.10 (1H, d, J = 8.9 Hz), 8.20 (1H, d, J = 8.7 Hz), 8.99 (1H, s), 9.66 (1H, s), 12.15 (1H, s).

Example 3: Pharmacological activity

The compounds of the invention were subjected to biological tests so as to evaluate their potential to treat or prevent certain pathologies. First, the ability of the compounds to behave as an activator of PPAR nuclear receptors was measured.

A transactivation test was used as the primary screening test. Cos-7 cells were transfected with a plasmid expressing a chimera of a murine or human receptor PPAR-Gal4 (receptor PPAR α -Gal4 or PPAR δ -Gal4 or PPAR γ -Gal4) and of a reporter plasmid 5Gal4pGL3 TK Luc. The transfections were carried out using a chemical agent (Jet PEI). Transfected cells were distributed into 384 well plates and allowed to stand for 24 hours. At 24 hours the culture medium was changed. The products to be tested were added (final concentration between 10^{-4} and 3.10^{-10} M) to the culture medium. After overnight incubation, luciferase expression was measured after addition of "SteadyGlo" according to the manufacturer's instructions (Promega).

Fenofibric acid at 10^{-5} M (PPAR α agonist), GW501516 at 10^{-8} M (PPAR δ agonist) and rosiglitazone at 10^{-6} M (PPAR γ agonist) were used as references.

The results are expressed as the induction rate (number of times) compared to the basal level as a percentage of activity of the adequate reference (reference = 100%). Effect-concentration curves and EC₅₀ were calculated using Assay Explorer (MDL) software.

Table 1

Human receptor	PPAR α		PPAR δ		PPAR γ	
Example	EC ₅₀ (nM)	Top(%)	EC ₅₀ (nM)	Top(%)	EC ₅₀ (nM)	Top(%)
1	2317	67	952	80	85	53
2	2280	62	1177	80	136	50

As can be seen from the table above, the deuterated derivatives of lanifibranor activate the three subtypes of PPAR receptors (PPAR α , PPAR γ and PPAR δ), with an EC₅₀ of less than 2.5 μ M for each of the sub-types. It can also be noted that the ratio between the EC₅₀ of two PPAR subtypes is either less than 100 or greater than 0.01.


References

1. DeWitt & Maryanoff, *Biochemistry* 2018, 57, 472–473
2. Fisher et al, *Curr Opin Drug Discov Devel*, 2006, 9: 101-09
3. Foster, *Adv Drug Res* 1985, 14: 1-40
4. Gant, *J. Med. Chem.* 2014, 57, 3595–3611
5. Harbeson & Tung, *Medchem News* 2014, 2 9-22
6. Junk et al., *J. Label Compd. Radiopharm* 1997, 39 625-630
7. Katsnelson, *Nature Medecine* 2013, 19 6 656
8. Kerekes et al., *J. Med. Chem.* 2011, 54, 201–210
9. Maehr et al., *J. Med. Chem.* 2013, 56, 3878–3888
10. Modutwa & al., *J. Label Compd. Radiopharm* 2010, 53 686–692
11. WO 2007/026097
12. WO 2015/189401
13. WO 2017/136375
14. WO 2018/039521

TERAPEUTISKE ANVENDELSER AF DEUTEREREDE DERIVATER AF LANIFIBRANOR

PATENTKRAV

1. Deutereret derivat af lanifibranor med formel (I):



hvor mindst en af grupperne R₁ til R₇ er et deuteriumatom og de andre grupper R₁ til R₇ er hydrogenatomer, og dets farmaceutisk acceptable salte og solvater til anvendelse ved behandling af hyperkolesterolæmi, hyperlipidæmi, hypertriglyceridæmi, dyslipidemier, insulinresistens, diabetes, fedme, hjerte-kar-sygdomme som konsekvens af en ubalance i serumlipoproteiner, sygdomme relateret til endoteldysfunktion, aterosklerose, myokardieinfarkt, hypertension, cerebrovaskulære problemer, inflammatoriske sygdomme som for eksempel reumatoid arthritis og neurodegenerationer såsom Alzheimers sygdom eller Parkinsons sygdom.

2. Deutereret derivat til anvendelse ifølge krav 1, hvori mindst gruppen R₁ er D i formel (I).
3. Deutereret derivat til anvendelse ifølge krav 2, som er syre 4-(1-(2-deuterio-1,3-benzothiazol-6-yl)sulfonyl)-5-chlor-1H-indol-2-yl)butansyre.
4. Deutereret derivat til anvendelse ifølge krav 1, hvori mindst en af grupperne R₂ til R₇ er D i formel (I).
5. Deutereret derivat til anvendelse ifølge krav 4, hvori mindst en af grupperne R₂ og R₃ og/eller mindst en af grupperne R₄ og R₅ og/eller mindst en af grupperne R₆ og R₇ er D i formel (I).

6. Deutereret derivat til anvendelse ifølge krav 5, som er syre 4-[1-(1,3-benzothiazol-6-ylsulfonyl)-5-chlor-indol-2-yl]-2,2,3,3,4,4-hexadeuteriobutansyre.