



(51) International Patent Classification:

C07D 403/12 (2006.01) *A61P 35/00* (2006.01)
A61K 31/506 (2006.01)

(21) International Application Number:

PCT/US2009/058212

(22) International Filing Date:

24 September 2009 (24.09.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/194,351 26 September 2008 (26.09.2008) US

(71) **Applicant** (for all designated States except US): **CONCERT PHARMACEUTICALS, INC.** [US/US]; 99 Hayden Avenue, Suite 500, Lexington, Massachusetts 02421 (US).

(72) **Inventor; and**

(75) **Inventor/Applicant** (for US only): **LIU, Julie, F.** [US/US]; 3 Whitman Circle, Lexington, Massachusetts 02420 (US).

(74) **Agents:** **DAVIS, Steven, G.** et al.; **Foley & Lardner LLP**, 111 Huntington Avenue, Boston, Massachusetts 02199 (US).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ,

CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

— with international search report (Art. 21(3))
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** PYRIDINEAMINE DERIVATIVES



(I)

(57) **Abstract:** This invention relates to novel compounds that are pyrimidineamine derivatives and pharmaceutically acceptable salts thereof. More specifically, this invention relates to novel pyrimidineamine derivatives that are derivatives of pazopanib. This invention also provides compositions comprising one or more compound of this invention and a carrier, and the use of the disclosed compounds and compositions in methods of treating diseases and conditions that are beneficially treated by administering a VEGFR (1-3) inhibitor, such as pazopanib.

PYRIDINEAMINE DERIVATIVES

CROSS-REFERENCE TO PRIOR APPLICATION

[1] This application claims priority under 35 USC 119 to United States Provisional Application Serial No. 61/194,351, filed September 26, 2008, the entire disclosure of which is hereby incorporated by reference.

BACKGROUND OF THE INVENTION

[2] Pazopanib, also known as 5-[4-[N-(2,3-Dimethyl-2H-indazol-6-yl)-N-methylamino]pyrimidin-2-ylamino]-2-methylbenzenesulfonamide hydrochloride, inhibits the VEGF (vascular epidermal growth factor)-1 (Flt-1) receptor, the VEGF-2 (FLK-1/KDR) receptor, and the VEGF-3 (Flt-4) receptor, thus blocking angiogenesis and tumor growth.

[3] Pazopanib is currently in clinical trials for renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, liver cancer, macular degeneration and psoriasis.

[4] The most common adverse events were nausea, diarrhea, fatigue, hypertension, anorexia, vomiting, and hair depigmentation (Sorbera, L et al, Drugs Fut, 2006, 31(7): 585).

[5] Despite the beneficial activities of pazopanib, there is a continuing need for new compounds to treat the aforementioned diseases and conditions.

SUMMARY OF THE INVENTION

[6] This invention relates to novel compounds that are pyrimidineamine derivatives and pharmaceutically acceptable salts thereof. More specifically, this invention relates to novel pyrimidineamine derivatives that are derivatives of pazopanib. This invention also provides compositions comprising one or more compound of this invention and a carrier, and the use of the disclosed compounds and compositions in methods of treating diseases and conditions that

are beneficially treated by administering a VEGFR (1-3) inhibitor, such as pazopanib.

DETAILED DESCRIPTION OF THE INVENTION

[7] The term “treat” means one or more of decrease, suppress, attenuate, diminish, arrest, or stabilize the development or progression of a disease (e.g., a disease or disorder delineated herein).

[8] “Disease” means any condition or disorder that damages or interferes with the normal function of a cell, tissue, or organ.

[9] It will be recognized that some variation of natural isotopic abundance occurs in a synthesized compound depending upon the origin of chemical materials used in the synthesis. Thus, a preparation of pazopanib will inherently contain small amounts of deuterated isotopologues. The concentration of naturally abundant stable hydrogen and carbon isotopes, notwithstanding this variation, is small and immaterial as compared to the degree of stable isotopic substitution of compounds of this invention. See, for instance, Wada E et al, Scikagaku 1994, 66:15; Gannes LZ et al, Comp Biochem Physiol Mol Integr Physiol 1998, 119:725. In a compound of this invention, when a particular position is designated as having deuterium, it is understood that the abundance of deuterium at that position is at least 3340 times greater than the natural abundance of deuterium, which is 0.015% (i.e., at least 50.1% incorporation of deuterium).

[10] The term “isotopic enrichment factor” as used herein means the ratio between the isotopic abundance and the natural abundance of a specified isotope.

[11] In other embodiments, a compound of this invention has an isotopic enrichment factor for each designated deuterium atom of at least 3500 (52.5% deuterium incorporation at each designated deuterium atom), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), at least 6333.3 (95% deuterium incorporation), at least 6466.7 (97% deuterium incorporation), at least 6600 (99% deuterium incorporation), or at least 6633.3 (99.5% deuterium incorporation).

[12] In the compounds of this invention any atom not specifically designated as a particular isotope is meant to represent any stable isotope of that atom. Unless otherwise stated, when a

position is designated specifically as "H" or "hydrogen", the position is understood to have hydrogen at its natural abundance isotopic composition.

[13] The term "isotopologue" refers to a species that differs from a specific compound of this invention only in the isotopic composition thereof.

[14] The term "compound," when referring to a compound of this invention, refers to a collection of molecules having an identical chemical structure, except that there may be isotopic variation among the constituent atoms of the molecules. Thus, it will be clear to those of skill in the art that a compound represented by a particular chemical structure containing indicated deuterium atoms, will also contain lesser amounts of isotopologues having hydrogen atoms at one or more of the designated deuterium positions in that structure. The relative amount of such isotopologues in a compound of this invention will depend upon a number of factors including the isotopic purity of deuterated reagents used to make the compound and the efficiency of incorporation of deuterium in the various synthesis steps used to prepare the compound. However, as set forth above the relative amount of such isotopologues *in toto* will be less than 49.9% of the compound.

[15] The invention also provides salts of the compounds of the invention.

[16] A salt of a compound of this invention is formed between an acid and a basic group of the compound, such as an amino functional group, or a base and an acidic group of the compound, such as a carboxyl functional group. According to another embodiment, the compound is a pharmaceutically acceptable acid addition salt.

[17] The term "pharmaceutically acceptable," as used herein, refers to a component that is, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and other mammals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any non-toxic salt that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention. A "pharmaceutically acceptable counterion" is an ionic portion of a salt that is not toxic when released from the salt upon administration to a recipient.

[18] Acids commonly employed to form pharmaceutically acceptable salts include inorganic acids such as hydrogen bisulfide, hydrochloric acid, hydrobromic acid, hydroiodic acid, sulfuric acid and phosphoric acid, as well as organic acids such as para-toluenesulfonic acid, salicylic

acid, tartaric acid, bitartaric acid, ascorbic acid, maleic acid, besylic acid, fumaric acid, gluconic acid, glucuronic acid, formic acid, glutamic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, lactic acid, oxalic acid, para-bromophenylsulfonic acid, carbonic acid, succinic acid, citric acid, benzoic acid and acetic acid, as well as related inorganic and organic acids. Such pharmaceutically acceptable salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caprate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, malate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, terephthalate, sulfonate, xylene sulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, β -hydroxybutyrate, glycolate, maleate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and other salts. In one embodiment, pharmaceutically acceptable acid addition salts include those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and especially those formed with organic acids such as maleic acid.

[19] The compounds of the present invention (e.g., compounds of Formula I), may contain an asymmetric carbon atom, for example, as the result of deuterium substitution or otherwise. As such, compounds of this invention can exist as either individual enantiomers, or mixtures of the two enantiomers. Accordingly, a compound of the present invention may exist as either a racemic mixture or a scalemic mixture, or as individual respective stereoisomers that are substantially free from another possible stereoisomer. The term "substantially free of other stereoisomers" as used herein means less than 25% of other stereoisomers, preferably less than 10% of other stereoisomers, more preferably less than 5% of other stereoisomers and most preferably less than 2% of other stereoisomers, or less than "X%" of other stereoisomers (wherein X is a number between 0 and 100, inclusive) are present. Methods of obtaining or synthesizing an individual enantiomer for a given compound are known in the art and may be applied as practicable to final compounds or to starting material or intermediates.

[20] Unless otherwise indicated, when a disclosed compound is named or depicted by a structure without specifying the stereochemistry and has one or more chiral centers, it is understood to represent all possible stereoisomers of the compound.


[21] The term “stable compounds,” as used herein, refers to compounds which possess stability sufficient to allow for their manufacture and which maintain the integrity of the compound for a sufficient period of time to be useful for the purposes detailed herein (e.g., formulation into therapeutic products, intermediates for use in production of therapeutic compounds, isolatable or storable intermediate compounds, treating a disease or condition responsive to therapeutic agents).

[22] “D” refers to deuterium. “Stereoisomer” refers to both enantiomers and diastereomers. “Tert”, “^t”, and “t-” each refer to tertiary. “US” refers to the United States of America.

[23] Throughout this specification, a variable may be referred to generally (e.g., “each R”) or may be referred to specifically (e.g., R¹, R², R³, etc.). Unless otherwise indicated, when a variable is referred to generally, it is meant to include all specific embodiments of that particular variable.

THERAPEUTIC COMPOUNDS

[24] The present invention provides a compound of Formula I:



wherein:

- each Y is independently selected from hydrogen and deuterium;
- each R is independently selected from CH₃, CH₂D, CHD₂ and CD₃; and
- when each R is CH₃, at least one Y is deuterium.

[25] One embodiment of this invention provides a compound of Formula I where each R is independently selected from CH₃ and CD₃.

[26] Another embodiment provides a compound of Formula I where R¹ is CD₃. In one aspect of this embodiment, each of R²-R⁴ is independently selected from CH₃ and CD₃.

[27] Another embodiment provides a compound where Y¹, Y² and Y³ are the same. In one aspect of this embodiment, each R is independently selected from CH₃ and CD₃. In another aspect, R¹ is CD₃ and each of R²-R⁴ is independently selected from CH₃ and CD₃. In another

aspect of this embodiment, Y^1 , Y^2 and Y^3 are all deuterium. In yet another aspect, Y^1 , Y^2 and Y^3 are all hydrogen.

[28] Another embodiment provides a compound where Y^6 , Y^7 and Y^8 are the same. In one aspect of this embodiment, each R is independently selected from CH_3 and CD_3 . In another aspect R^1 is CD_3 and each R^2 - R^4 is independently selected from CH_3 and CD_3 . In yet another aspect Y^1 , Y^2 and Y^3 are the same. In a particular aspect, Y^1 , Y^2 and Y^3 are all the same and each R is independently selected from CH_3 and CD_3 . In yet another aspect, Y^1 , Y^2 and Y^3 are the same, R^1 is CD_3 and each of R^2 - R^4 is independently selected from CH_3 and CD_3 . In another aspect, Y^6 , Y^7 and Y^8 are all deuterium. In another aspect, Y^6 , Y^7 and Y^8 are all hydrogen. In another aspect, R^2 is CD_3 . In another aspect, R^3 is CD_3 . In another aspect, R^2 and R^3 are both CD_3 .

[29] Another embodiment provides a compound where Y^5 is deuterium. In one aspect of this embodiment, each R is independently selected from CH_3 and CD_3 . In another aspect, R^1 is CD_3 and each R^2 - R^4 is independently selected from CH_3 and CD_3 . In yet another aspect, Y^1 , Y^2 and Y^3 are the same. In a further aspect Y^1 , Y^2 and Y^3 are the same and each R is independently selected from CH_3 and CD_3 . In another aspect, Y^6 , Y^7 and Y^8 are the same and each R is independently selected from CH_3 and CD_3 . In another aspect, Y^6 , Y^7 and Y^8 are the same, R^1 is CD_3 and each R^2 - R^4 is independently selected from CH_3 and CD_3 . In yet another aspect, Y^1 , Y^2 and Y^3 are the same and Y^6 , Y^7 and Y^8 are the same. In another aspect, Y^1 , Y^2 and Y^3 are the same, Y^6 , Y^7 and Y^8 are the same R^3 is CD_3 and each of R^2 - R^4 is independently selected from CH_3 and CD_3 . In another aspect of this embodiment, Y^4 is deuterium.

[30] In another embodiment, Y^1 , Y^2 and Y^3 are deuterium and R^1 is CD_3 . In one aspect of this embodiment, each of R^2 - R^4 is independently selected from CH_3 and CD_3 . In another aspect, Y^6 , Y^7 and Y^8 are the same. In yet another aspect, Y^6 , Y^7 and Y^8 are the same and each R^2 - R^4 is independently selected from CH_3 and CD_3 .

[31] Yet another embodiment provides a compound where R^4 is CD_3 .

[32] Another embodiment provides a compound where R^4 is CH_3 .

[33] Compounds of Formula I are comprised of three rings: a phenyl ring, a pyrimidine ring and an indazole ring, embodiments of which are described above. It will be apparent to one skilled in the art that features of a particular ring may be combined with features of one or more other rings to provide other embodiments of this invention.

[34] Examples of specific compounds of Formula I are set forth in Table 1 below.

Table 1: Examples of Specific Compounds of Formula I

Compound	Y ¹	R ¹	Y ²	Y ³	Y ⁴	Y ⁵	Y ⁶	Y ⁷	R ²	R ³	Y ⁸	R ⁴
100	D	CD ₃	D	D	D	D	D	D	CD ₃	CD ₃	D	CD ₃
101	H	CD ₃	H	H	D	D	D	D	CD ₃	CD ₃	D	CD ₃
102	D	CD ₃	D	D	H	D	D	D	CD ₃	CD ₃	D	CD ₃
103	D	CD ₃	D	D	D	H	D	D	CD ₃	CD ₃	D	CD ₃
104	D	CD ₃	D	D	H	H	D	D	CD ₃	CD ₃	D	CD ₃
105	D	CD ₃	D	D	D	D	H	H	CD ₃	CD ₃	H	CD ₃
106	H	CD ₃	H	H	H	H	H	H	CD ₃	CD ₃	H	CD ₃
107	H	CH ₃	H	H	H	H	H	H	CH ₃	CD ₃	H	CD ₃
108	H	CH ₃	H	H	H	H	H	H	CH ₃	CH ₃	H	CD ₃
109	H	CH ₃	H	H	H	H	H	H	CH ₃	CD ₃	H	CH ₃
110	H	CH ₃	H	H	H	H	H	H	CD ₃	CD ₃	H	CD ₃
111	H	CD ₃	H	H	H	H	H	H	CH ₃	CD ₃	H	CD ₃

[35] In another set of embodiments, any atom not designated as deuterium in any of the embodiments set forth above is present at its natural isotopic abundance.


[36] The synthesis of compounds of Formula I can be readily achieved by synthetic chemists of ordinary skill. Relevant procedures and intermediates are disclosed, for instance in Sorbera, L et al, *Drugs Fut*, 2006, 31(7):585; and PCT patent publications WO2006020564, WO2003106416 and WO2002059110.

[37] Such methods can be carried out utilizing corresponding deuterated and optionally, other isotope-containing reagents and/or intermediates to synthesize the compounds delineated herein, or invoking standard synthetic protocols known in the art for introducing isotopic atoms to a chemical structure. Certain intermediates can be used with or without purification (e.g., filtration, distillation, sublimation, crystallization, trituration, solid phase extraction, and chromatography).

EXEMPLARY SYNTHESIS


[38] The schemes shown below illustrate how compounds of Formula 1 may be prepared.

[39] Scheme 1. General Route to Compounds of Formula 1




[40] Scheme 1 shows a general route to preparing compounds of Formula 1. An appropriately-deuterated aniline **10** is treated with tert-butyl nitrite in acetic acid to afford cyclized indazole **11**. Alkylation with appropriately-deuterated dimethylsulfate provides heterocycle **12**. Reduction of the nitro group with tin chloride and HCl yields amine **13**. Reaction of amine **13** with appropriately-deuterated dichloropyrimidine **14** provides adduct **15**. Alkylation of the secondary amine with appropriately-deuterated methyl iodide yields compound **16**, and reaction with appropriately-deuterated aniline **17** in the presence of HCl affords desired

Formula I compound as the HCl salt.



[41] Known 5-deutero-2,4-dichloropyrimidine, (Ple, N et al, Tet Lett 1993, 34(10):1605-8.) may be used in Scheme 1 as intermediate **14** to provide a compound of Formula I wherein Y⁴ is hydrogen and Y⁵ is deuterium.


[42] Scheme 2. Preparation of Starting Material **10**




[43] Scheme 2 shows a route to making the deuterated anilines **10**, which are useful starting materials for Scheme 1. An appropriately-deuterated ethylbenzene **20** is treated with Bu₄NNO₃ and trifluoroacetic anhydride to provide nitrobenzene **21** according to the general procedure of Masci, B, Tetrahedron 1989, 45(9):2719-30. Hydrogenation of the nitro group affords aniline **22**, following the general procedure of Klyuev, MV et al, Neftekhimiya 2002, 42(1):32-35. Treatment of aniline **22** with KNO₃ and H₂SO₄ via the method of Ruiz-Caro J, et al, Bioorg Med Chem Lett 2006, 16(3):668-671 affords the desired aniline **10**.


[44] Commercially-available ethylbenzene-d₁₀ may be used in Scheme 2 as starting compound **20** to ultimately produce a compound of Formula I wherein Y⁶, Y⁷ and Y⁸ are deuterium, and R² is CD₃.

[45] Scheme 3. Preparation of Sulfonamide **17**



[46] Scheme 3 shows the preparation of deuterated sulfonamide **17** that is used in Scheme 1. An appropriately-deuterated nitrotoluene **30** is treated with chlorosulfonic acid, followed by ammonium hydroxide, to provide sulfonamide **31** according to the procedure of Chinese patent publication CN1869015A or that of Srikanth, K et al, *Bioorg Med Chem* 2002, 10(6):1841-1854. Reduction of the nitro group with either hydrogen and palladium (see Chinese patent publication CN1702064A) or iron and ammonium chloride (see Rockway, TW et al, *Bioorg Med Chem Lett* 2006, 16(14):3833-3838) yields desired sulfonamide **17**.



[47] Commercially-available p-nitrotoluene- α,α,α -d₃, D_3C , may be used in Scheme 3 as compound **30** to provide intermediate **17** and subsequently a compound of Formula I wherein R¹ is CD₃; and Y¹, Y² and Y³ are simultaneously hydrogen.

[48] Alternatively, p-nitrotoluene-d₇ (see Gorissen, II et al Editor(s): Pleiss, Ulrich; Voges, Rolf. *Synthesis and Applications of Isotopically Labelled Compounds*, Proceedings of the International Symposium, 7th, Dresden, Germany, June 18-22, 2000 (2001), Meeting Date 2000, 170-173; or Delgado, M et al, *J Am Chem Soc* 1986, 108(14):4135-8) may be used in Scheme 3 as compound **30** to provide intermediate **17** and a compound of Formula I wherein R¹ is CD₃ and Y¹, Y² and Y³ are simultaneously deuterium.

[49] The specific approaches and compounds shown above are not intended to be limiting. The chemical structures in the schemes herein depict variables that are hereby defined commensurately with chemical group definitions (moieties, atoms, etc.) of the corresponding position in the compound formulae herein, whether identified by the same variable name (i.e., R¹, R², R³, etc.) or not. The suitability of a chemical group in a compound structure for use in the synthesis of another compound is within the knowledge of one of ordinary skill in the art. Additional methods of synthesizing compounds of Formula I and their synthetic precursors, including those within routes not explicitly shown in schemes herein, are within the means of chemists of ordinary skill in the art. Methods for optimizing reaction conditions and, if necessary, minimizing competing by-products, are known in the art.

[50] Synthetic chemistry transformations and protecting group methodologies (protection and deprotection) useful in synthesizing the applicable compounds are known in the art and include, for example, those described in Larock R, *Comprehensive Organic Transformations*, VCH

Publishers (1989); Greene TW et al, *Protective Groups in Organic Synthesis*, 3rd Ed., John Wiley and Sons (1999); Fieser L et al, *Fieser and Fieser's Reagents for Organic Synthesis*, John Wiley and Sons (1994); and Paquette L, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons (1995) and subsequent editions thereof.

[51] Combinations of substituents and variables envisioned by this invention are only those that result in the formation of stable compounds.

COMPOSITIONS

[52] The invention also provides pyrogen-free compositions comprising an effective amount of a compound of Formula I (e.g., including any of the formulae herein), or a pharmaceutically acceptable salt of said compound; and an acceptable carrier. Preferably, a composition of this invention is formulated for pharmaceutical use ("a pharmaceutical composition"), wherein the carrier is a pharmaceutically acceptable carrier. The carrier(s) are "acceptable" in the sense of being compatible with the other ingredients of the formulation and, in the case of a pharmaceutically acceptable carrier, not deleterious to the recipient thereof in an amount used in the medicament.

[53] Pharmaceutically acceptable carriers, adjuvants and vehicles that may be used in the pharmaceutical compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[54] If required, the solubility and bioavailability of the compounds of the present invention in pharmaceutical compositions may be enhanced by methods well-known in the art. One method includes the use of lipid excipients in the formulation. See "Oral Lipid-Based Formulations: Enhancing the Bioavailability of Poorly Water-Soluble Drugs (Drugs and the Pharmaceutical Sciences)," David J. Hauss, ed. Informa Healthcare, 2007; and "Role of Lipid Excipients in Modifying Oral and Parenteral Drug Delivery: Basic Principles and Biological Examples."

Kishor M. Wasan, ed. Wiley-Interscience, 2006.

[55] Another known method of enhancing bioavailability is the use of an amorphous form of a compound of this invention optionally formulated with a poloxamer, such as LUTROL™ and PLURONIC™ (BASF Corporation), or block copolymers of ethylene oxide and propylene oxide. See United States patent 7,014,866; and United States patent publications 20060094744 and 20060079502.

[56] The pharmaceutical compositions of the invention include those suitable for oral, rectal, nasal, topical (including buccal and sublingual), vaginal or parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. In certain embodiments, the compound of the formulae herein is administered transdermally (e.g., using a transdermal patch or iontophoretic techniques). Other formulations may conveniently be presented in unit dosage form, e.g., tablets, sustained release capsules, and in liposomes, and may be prepared by any methods well known in the art of pharmacy. See, for example, Remington: The Science and Practice of Pharmacy, Lippincott Williams & Wilkins, Baltimore, MD (20th ed. 2000).

[57] Such preparative methods include the step of bringing into association with the molecule to be administered ingredients such as the carrier that constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers, liposomes or finely divided solid carriers, or both, and then, if necessary, shaping the product.

[58] In certain embodiments, the compound is administered orally. Compositions of the present invention suitable for oral administration may be presented as discrete units such as capsules, sachets, or tablets each containing a predetermined amount of the active ingredient; a powder or granules; a solution or a suspension in an aqueous liquid or a non-aqueous liquid; an oil-in-water liquid emulsion; a water-in-oil liquid emulsion; packed in liposomes; or as a bolus, etc. Soft gelatin capsules can be useful for containing such suspensions, which may beneficially increase the rate of compound absorption.

[59] In the case of tablets for oral use, carriers that are commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are administered orally, the active ingredient is combined with emulsifying

and suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

[60] Compositions suitable for oral administration include lozenges comprising the ingredients in a flavored basis, usually sucrose and acacia or tragacanth; and pastilles comprising the active ingredient in an inert basis such as gelatin and glycerin, or sucrose and acacia.

[61] Compositions suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampules and vials, and may be stored in a freeze dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets.

[62] Such injection solutions may be in the form, for example, of a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant.

[63] The pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a compound of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release

the active components. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

[64] The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art. See, e.g.: Rabinowitz JD and Zaffaroni AC, US Patent 6,803,031, assigned to Alexza Molecular Delivery Corporation.

[65] Topical administration of the pharmaceutical compositions of this invention is especially useful when the desired treatment involves areas or organs readily accessible by topical application. For topical application topically to the skin, the pharmaceutical composition should be formulated with a suitable ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax, and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active compound suspended or dissolved in a carrier. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol, and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-transdermal patches and iontophoretic administration are also included in this invention.

[66] Application of the subject therapeutics may be local, so as to be administered at the site of interest. Various techniques can be used for providing the subject compositions at the site of interest, such as injection, use of catheters, trocars, projectiles, pluronic gel, stents, sustained drug release polymers or other device which provides for internal access.

[67] Thus, according to yet another embodiment, the compounds of this invention may be incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents, or catheters. Suitable coatings and the general preparation of coated implantable devices are known in the art and are exemplified in US Patents

6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition. Coatings for invasive devices are to be included within the definition of pharmaceutically acceptable carrier, adjuvant or vehicle, as those terms are used herein.

[68] According to another embodiment, the invention provides a method of coating an implantable medical device comprising the step of contacting said device with the coating composition described above. It will be obvious to those skilled in the art that the coating of the device will occur prior to implantation into a mammal.

[69] According to another embodiment, the invention provides a method of impregnating an implantable drug release device comprising the step of contacting said drug release device with a compound or composition of this invention. Implantable drug release devices include, but are not limited to, biodegradable polymer capsules or bullets, non-degradable, diffusible polymer capsules and biodegradable polymer wafers.

[70] According to another embodiment, the invention provides an implantable medical device coated with a compound or a composition comprising a compound of this invention, such that said compound is therapeutically active.

[71] According to another embodiment, the invention provides an implantable drug release device impregnated with or containing a compound or a composition comprising a compound of this invention, such that said compound is released from said device and is therapeutically active.

[72] Where an organ or tissue is accessible because of removal from the patient, such organ or tissue may be bathed in a medium containing a composition of this invention, a composition of this invention may be painted onto the organ, or a composition of this invention may be applied in any other convenient way.

[73] In another embodiment, a composition of this invention further comprises a second therapeutic agent. The second therapeutic agent may be selected from any compound or therapeutic agent known to have or that demonstrates advantageous properties when administered with a compound having the same mechanism of action as pazopanib.

[74] Preferably, the second therapeutic agent is an agent useful in the treatment or prevention of a disease or condition selected from cancer including renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, and liver cancer; ocular neovascular disorders including macular degeneration, angioid streaks, uveitis and macular edema; and psoriasis.

[75] In one embodiment, the second therapeutic agent is selected from lapatinib, FOLFOLX6 (fluorouracil, oxaliplatin, leucovorin), CapeOx (capecitabine, oxaliplatin), paclitaxel, and carboplatin.

[76] In another embodiment, the invention provides separate dosage forms of a compound of this invention and one or more of any of the above-described second therapeutic agents, wherein the compound and second therapeutic agent are associated with one another. The term “associated with one another” as used herein means that the separate dosage forms are packaged together or otherwise attached to one another such that it is readily apparent that the separate dosage forms are intended to be sold and administered together (within less than 24 hours of one another, consecutively or simultaneously).

[77] In the pharmaceutical compositions of the invention, the compound of the present invention is present in an effective amount. As used herein, the term “effective amount” refers to an amount which, when administered in a proper dosing regimen, is sufficient to treat (therapeutically or prophylactically) the target disorder. For example, an effective amount is sufficient to reduce or ameliorate the severity, duration or progression of the disorder being treated, prevent the advancement of the disorder being treated, cause the regression of the disorder being treated, or enhance or improve the prophylactic or therapeutic effect(s) of another therapy.

[78] The interrelationship of dosages for animals and humans (based on milligrams per meter squared of body surface) is described in Freireich et al, (1966) Cancer Chemother. Rep 50: 219. Body surface area may be approximately determined from height and weight of the patient. See, e.g., Scientific Tables, Geigy Pharmaceuticals, Ardsley, NY, 1970, 537.

[79] In one embodiment, an effective amount of a compound of this invention can range from about 2 to 8000 mg per treatment. In more specific embodiments the range is from about 20 to 4000 mg, or from 40 to 1600 mg, or most specifically from 200 to 800 mg. Treatment typically is administered one to two times daily.

[80] Effective doses will also vary, as recognized by those skilled in the art, depending on the diseases treated, the severity of the disease, the route of administration, the sex, age and general health condition of the patient, excipient usage, the possibility of co-usage with other therapeutic treatments such as use of other agents and the judgment of the treating physician. For example, guidance for selecting an effective dose can be determined by reference to the prescribing information for pazopanib.

[81] For pharmaceutical compositions that comprise a second therapeutic agent, an effective amount of the second therapeutic agent is between about 20% and 100% of the dosage normally utilized in a monotherapy regime using just that agent. Preferably, an effective amount is between about 70% and 100% of the normal monotherapeutic dose. The normal monotherapeutic dosages of these second therapeutic agents are well known in the art. *See, e.g.,* Wells et al, eds., *Pharmacotherapy Handbook*, 2nd Edition, Appleton and Lange, Stamford, Conn. (2000); *PDR Pharmacopoeia*, Tarascon Pocket Pharmacopoeia 2000, Deluxe Edition, Tarascon Publishing, Loma Linda, Calif. (2000), each of which references are incorporated herein by reference in their entirety.

[82] It is expected that some of the second therapeutic agents referenced above will act synergistically with the compounds of this invention. When this occurs, it will allow the effective dosage of the second therapeutic agent and/or the compound of this invention to be reduced from that required in a monotherapy. This has the advantage of minimizing toxic side effects of either the second therapeutic agent or a compound of this invention, synergistic improvements in efficacy, improved ease of administration or use and/or reduced overall expense of compound preparation or formulation.

METHODS OF TREATMENT

[83] In another embodiment, the invention provides a method of inhibiting the activity of the VEGF (1-3) receptors in a cell, comprising contacting a cell with one or more compounds of Formula I herein.

[84] According to another embodiment, the invention provides a method of treating a disease that is beneficially treated by pazopanib in a patient in need thereof comprising the step of administering to said patient an effective amount of a compound of Formula I or pharmaceutically acceptable salt thereof or a composition of this invention. Such diseases are well known in the art and are disclosed in, but not limited to the following patents and published applications: WO 2002059110, WO 2006020564, WO 2007064752, and WO 2007064753. Such diseases include, but are not limited to, cancer including renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, and liver cancer; ocular neovascular disorders including macular degeneration, angioid streaks, uveitis and macular edema; and psoriasis.

[85] In one particular embodiment, the method of this invention is used to treat a disease or condition selected from renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, liver cancer, macular degeneration and psoriasis.

[86] Methods delineated herein also include those wherein the patient is identified as in need of a particular stated treatment. Identifying a patient in need of such treatment can be in the judgment of a patient or a health care professional and can be subjective (e.g. opinion) or objective (e.g. measurable by a test or diagnostic method).

[87] In another embodiment, any of the above methods of treatment comprises the further step of co-administering to said patient one or more second therapeutic agents. The choice of second therapeutic agent may be made from any second therapeutic agent known to be useful for co-administration with pazopanib. The choice of second therapeutic agent is also dependent upon the particular disease or condition to be treated. Examples of second therapeutic agents that may be employed in the methods of this invention are those set forth above for use in combination compositions comprising a compound of this invention and a second therapeutic agent.

[88] In particular, the combination therapies of this invention include co-administering a

compound of Formula I and a second therapeutic agent for treatment of the following conditions: breast cancer (lapatinib), colorectal cancer (FOLFOX6 [fluorouracil, oxaliplatin, leucovorin], CapeOx [capecitabine, oxaliplatin]), solid tumor cancer (lapatinib, paclitaxel, carboplatin), malignant glioma (lapatinib), and cervical cancer (lapatinib).

[89] The term “co-administered” as used herein means that the second therapeutic agent may be administered together with a compound of this invention as part of a single dosage form (such as a composition of this invention comprising a compound of the invention and an second therapeutic agent as described above) or as separate, multiple dosage forms. Alternatively, the additional agent may be administered prior to, consecutively with, or following the administration of a compound of this invention. In such combination therapy treatment, both the compounds of this invention and the second therapeutic agent(s) are administered by conventional methods. The administration of a composition of this invention, comprising both a compound of the invention and a second therapeutic agent, to a patient does not preclude the separate administration of that same therapeutic agent, any other second therapeutic agent or any compound of this invention to said patient at another time during a course of treatment.

[90] Effective amounts of these second therapeutic agents are well known to those skilled in the art and guidance for dosing may be found in patents and published patent applications referenced herein, as well as in Wells et al, eds., Pharmacotherapy Handbook, 2nd Edition, Appleton and Lange, Stamford, Conn. (2000); PDR Pharmacopoeia, Tarascon Pocket Pharmacopoeia 2000, Deluxe Edition, Tarascon Publishing, Loma Linda, Calif. (2000), and other medical texts. However, it is well within the skilled artisan’s purview to determine the second therapeutic agent’s optimal effective-amount range.

[91] In one embodiment of the invention, where a second therapeutic agent is administered to a subject, the effective amount of the compound of this invention is less than its effective amount would be where the second therapeutic agent is not administered. In another embodiment, the effective amount of the second therapeutic agent is less than its effective amount would be where the compound of this invention is not administered. In this way, undesired side effects associated with high doses of either agent may be minimized. Other potential advantages (including without limitation improved dosing regimens and/or reduced drug cost) will be apparent to those of skill in the art.

[92] In yet another aspect, the invention provides the use of a compound of Formula I alone or together with one or more of the above-described second therapeutic agents in the manufacture of a medicament, either as a single composition or as separate dosage forms, for treatment or prevention in a patient of a disease, disorder or symptom set forth above. Another aspect of the invention is a compound of Formula I or pharmaceutically acceptable salt thereof, or a composition comprising a compound of Formula I or pharmaceutically acceptable salt thereof, for use in the treatment or prevention in a patient of a disease, disorder or symptom thereof delineated herein.

[93] In one aspect, the compound of Formula I or pharmaceutically acceptable salt thereof, or a composition comprising a compound of Formula I or pharmaceutically acceptable salt thereof is for use in treating a disease or condition selected from renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, liver cancer, macular degeneration and psoriasis.

[94] In another aspect, the compound of Formula I or pharmaceutically acceptable salt thereof, or a composition comprising a compound of Formula I or pharmaceutically acceptable salt thereof is used in conjunction with a second therapeutic agent useful in the treatment of cancer, ocular neovascular disorders, or psoriasis.

[95] In another aspect, the compound of Formula I or pharmaceutically acceptable salt thereof, or a composition comprising a compound of Formula I or pharmaceutically acceptable salt thereof is for use in treating breast cancer, malignant glioma, cervical cancer and solid tumor cancer; and the compound or composition is used in conjunction with lapatinib.

[96] In another aspect, the compound of Formula I or pharmaceutically acceptable salt thereof, or a composition comprising a compound of Formula I or pharmaceutically acceptable salt thereof is for use in treating colorectal cancer; and the compound or composition is used in conjunction with a second therapeutic agent selected from a combination of fluorouracil, oxaliplatin, and leucovorin; and a combination of capecitabine and oxaliplatin.

[97] In another aspect, the compound of Formula I or pharmaceutically acceptable salt thereof, or a composition comprising a compound of Formula I or pharmaceutically acceptable salt

thereof is for use in treating a solid tumor cancer; and the compound or composition is used in conjunction with a second therapeutic agent selected from paclitaxel, or carboplatin.

[98] The term "used in conjunction with" as used herein means administered simultaneously with, or administered within 24 hours of the subject compound or composition.

PHARMACEUTICAL KITS

[99] The present invention also provides kits for use to treat renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, liver cancer, macular degeneration and psoriasis. These kits comprise (a) a pharmaceutical composition comprising a compound of Formula I or a salt thereof, wherein said pharmaceutical composition is in a container; and (b) instructions describing a method of using the pharmaceutical composition to treat renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, liver cancer, macular degeneration and psoriasis.

[100] The container may be any vessel or other sealed or sealable apparatus that can hold said pharmaceutical composition. Examples include bottles, ampules, divided or multi-chambered holders bottles, wherein each division or chamber comprises a single dose of said composition, a divided foil packet wherein each division comprises a single dose of said composition, or a dispenser that dispenses single doses of said composition. The container can be in any conventional shape or form as known in the art which is made of a pharmaceutically acceptable material, for example a paper or cardboard box, a glass or plastic bottle or jar, a re-sealable bag (for example, to hold a "refill" of tablets for placement into a different container), or a blister pack with individual doses for pressing out of the pack according to a therapeutic schedule. The container employed can depend on the exact dosage form involved, for example a conventional cardboard box would not generally be used to hold a liquid suspension. It is feasible that more than one container can be used together in a single package to market a single dosage form. For

example, tablets may be contained in a bottle, which is in turn contained within a box. In one embodiment, the container is a blister pack.

[101] The kits of this invention may also comprise a device to administer or to measure out a unit dose of the pharmaceutical composition. Such device may include an inhaler if said composition is an inhalable composition; a syringe and needle if said composition is an injectable composition; a syringe, spoon, pump, or a vessel with or without volume markings if said composition is an oral liquid composition; or any other measuring or delivery device appropriate to the dosage formulation of the composition present in the kit.

[102] In an embodiment of the kits of this invention, the composition comprising the second therapeutic agent may be in a vessel or container that is separate from the vessel containing the composition comprising a compound of Formula I.

Example 1. Evaluation of Metabolic Stability

[103] *Microsomal Assay:* Human liver microsomes (20 mg/mL) are obtained from Xenotech, LLC (Lenexa, KS). β -nicotinamide adenine dinucleotide phosphate, reduced form (NADPH), magnesium chloride (MgCl_2), and dimethyl sulfoxide (DMSO) are purchased from Sigma-Aldrich.

[104] *Determination of Metabolic Stability:* 7.5 mM stock solutions of test compounds are prepared in DMSO. The 7.5 mM stock solutions are diluted to 12.5-50 μM in acetonitrile (ACN). The 20 mg/mL human liver microsomes are diluted to 0.625 mg/mL in 0.1 M potassium phosphate buffer, pH 7.4, containing 3 mM MgCl_2 . The diluted microsomes are added to wells of a 96-well deep-well polypropylene plate in triplicate. A 10 μL aliquot of the 12.5-50 μM test compound is added to the microsomes and the mixture is pre-warmed for 10 minutes. Reactions are initiated by addition of pre-warmed NADPH solution. The final reaction volume is 0.5 mL and contains 0.5 mg/mL human liver microsomes, 0.25-1.0 μM test compound, and 2 mM NADPH in 0.1 M potassium phosphate buffer, pH 7.4, and 3 mM MgCl_2 . The reaction mixtures are incubated at 37 $^{\circ}\text{C}$, and 50 μL aliquots are removed at 0, 5, 10, 20, and 30 minutes and added to shallow-well 96-well plates which contain 50 μL of ice-cold ACN with internal standard to stop the reactions. The plates are stored at 4 $^{\circ}\text{C}$ for 20 minutes after which 100 μL of water is added to the wells of the plate before centrifugation to pellet precipitated proteins. Supernatants are transferred to another 96-well plate and analyzed for amounts of parent remaining by LC-

MS/MS using an Applied Bio-systems API 4000 mass spectrometer. The same procedure is followed for pazopanib and the positive control, 7-ethoxycoumarin (1 μ M). Testing is done in triplicate.

[105] **Data analysis:** The *in vitro* $t_{1/2}$ s for test compounds are calculated from the slopes of the linear regression of % parent remaining (ln) vs incubation time relationship.

$$\text{in vitro } t_{1/2} = 0.693/k$$


$$k = -[\text{slope of linear regression of \% parent remaining(ln) vs incubation time}]$$

[106] Data analysis is performed using Microsoft Excel Software.

[107] Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to those of ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. All the patents, journal articles and other documents discussed or cited above are herein incorporated by reference.

What is claimed is:

1. A compound of Formula I:



or a pharmaceutically acceptable salt thereof, wherein:

each Y is independently selected from hydrogen and deuterium;

each R is independently selected from CH₃, CH₂D, CHD₂ and CD₃; and

when each R is CH₃, at least one Y is deuterium.

2. The compound of claim 1, wherein each R is independently selected from CH₃ and CD₃.
3. The compound of claim 1 or claim 2, wherein R¹ is CD₃.
4. The compound of any one of claims 1, 2 or 3, wherein Y¹, Y² and Y³ are the same.
5. The compound of any one of claims 1 to 4, wherein Y⁶, Y⁷ and Y⁸ are the same.
6. The compound of any one of claims 1 to 5, wherein Y⁵ is deuterium.
7. The compound of claim 2, wherein Y¹, Y² and Y³ are simultaneously deuterium.
8. The compound of claim 1, selected from any one of the compounds set forth in the table below:

Compound	Y ¹	R ¹	Y ²	Y ³	Y ⁴	Y ⁵	Y ⁶	Y ⁷	R ²	R ³	Y ⁸	R ⁴
100	D	CD ₃	D	D	D	D	D	D	CD ₃	CD ₃	D	CD ₃
101	H	CD ₃	H	H	D	D	D	D	CD ₃	CD ₃	D	CD ₃
102	D	CD ₃	D	D	H	D	D	D	CD ₃	CD ₃	D	CD ₃
103	D	CD ₃	D	D	D	H	D	D	CD ₃	CD ₃	D	CD ₃
104	D	CD ₃	D	D	H	H	D	D	CD ₃	CD ₃	D	CD ₃
105	D	CD ₃	D	D	D	D	H	H	CD ₃	CD ₃	H	CD ₃
106	H	CD ₃	H	H	H	H	H	H	CD ₃	CD ₃	H	CD ₃
107	H	CH ₃	H	H	H	H	H	H	CH ₃	CD ₃	H	CD ₃
108	H	CH ₃	H	H	H	H	H	H	CH ₃	CH ₃	H	CD ₃
109	H	CH ₃	H	H	H	H	H	H	CH ₃	CD ₃	H	CH ₃
110	H	CH ₃	H	H	H	H	H	H	CD ₃	CD ₃	H	CD ₃
111	H	CD ₃	H	H	H	H	H	H	CH ₃	CD ₃	H	CD ₃

9. The compound of any one of claims 1 to 8, wherein any atom not designated as deuterium is present at its natural isotopic abundance.
10. A pyrogen-free pharmaceutical composition comprising a compound of claim 1 or pharmaceutically acceptable salt thereof and an acceptable carrier.
11. The composition of claim 10 additionally comprising a second therapeutic agent useful in the treatment or prevention of a disease or condition selected from cancer; ocular neovascular disorders; and psoriasis.
12. The composition of claim 11, wherein the second therapeutic agent is selected from lapatinib; a combination of fluorouracil, oxaliplatin, and leucovorin; a combination of capecitabine and oxaliplatin; paclitaxel; and carboplatin.
13. The compound of claim 1 or the composition of claim 10 or 11 for use in inhibiting the activity of one or more of a VEGF-1, VEGF-2, or VEGF-3 receptor in a cell.

14. The compound of claim 1 or the composition of claim 10 or 11 for use in treating a disease or condition selected from cancer; ocular neovascular disorders; and psoriasis.

15. The compound or composition of claim 14, wherein the disease or condition is selected from renal cell carcinoma, breast cancer, cervical cancer, non-small cell lung cancer, ovarian cancer, fallopian tube cancer, peritoneal cancer, prostate cancer, rhinopharyngeal cancer, solid tumor cancer, bladder cancer, urethral cancer, multiforme glioblastoma, malignant glioma, pleural mesothelioma, multiple myeloma, neuroendocrine cancer, sarcoma, colorectal cancer, liver cancer, macular degeneration and psoriasis.

16. The compound or composition of claim 15, wherein the composition is used in conjunction with a second therapeutic agent useful in the treatment of cancer, ocular neovascular disorders, or psoriasis.

17. The compound or composition of claim 16, wherein:

- a. the disease or condition is selected from breast cancer, malignant glioma, cervical cancer and solid tumor cancer; and the compound or composition is used in conjunction with lapatinib;
- b. the disease or condition is colorectal cancer; and the compound or composition is used in conjunction with a second therapeutic agent selected from a combination of fluorouracil, oxaliplatin, and leucovorin; and a combination of capecitabine and oxaliplatin; or
- c. the disease or condition is a solid tumor cancer; and the compound or composition is used in conjunction with a second therapeutic agent selected from paclitaxel, or carboplatin.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2009/058212

A. CLASSIFICATION OF SUBJECT MATTER
INV. C07D403/12 A61K31/506 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 02/059110 A1 (GLAXO GROUP LTD [GB]; BOLOOR AMOGH [US]; CHEUNG MUI [US]; DAVIS RONDA) 1 August 2002 (2002-08-01) cited in the application example 69 page 40, line 4 - line 26 -----	1-17
A	WO 95/26325 A2 (ISOTECHNIKA INC [CA]) 5 October 1995 (1995-10-05) claim 1 page 1, line 9 - line 15 page 19, line 8 - line 14 -----	1-17

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"&" document member of the same patent family

Date of the actual completion of the international search

18 December 2009

Date of mailing of the international search report

18/01/2010

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Miniejew, Catherine

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2009/058212

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 02059110	A1	01-08-2002	AT 430742 T 15-05-2009
		AU 2002246723 B2 14-07-2005	
		BR 0116452 A 30-09-2003	
		CA 2432000 A1 01-08-2002	
		CN 1549813 A 24-11-2004	
		CZ 20031748 A3 14-04-2004	
		DK 1343782 T3 24-08-2009	
		EP 1343782 A1 17-09-2003	
		ES 2324981 T3 21-08-2009	
		HU 0400691 A2 28-07-2004	
		JP 4253188 B2 08-04-2009	
		JP 2004517925 T 17-06-2004	
		MX PA03005696 A 06-10-2003	
		NO 20032831 A 15-08-2003	
		NZ 526542 A 28-01-2005	
		PL 363243 A1 15-11-2004	
		SI 1343782 T1 31-10-2009	
		US 2007015756 A1 18-01-2007	
		US 2004242578 A1 02-12-2004	
		ZA 200304482 A 09-09-2005	
WO 9526325	A2	05-10-1995	AT 372966 T 15-09-2007
		AU 707748 B2 22-07-1999	
		AU 1944195 A 17-10-1995	
		BR 9507200 A 16-09-1997	
		CA 2186371 A1 05-10-1995	
		CN 1148843 A 30-04-1997	
		DE 69535592 T2 12-06-2008	
		EP 0751926 A1 08-01-1997	
		ES 2293638 T3 16-03-2008	
		JP 3696884 B2 21-09-2005	
		JP 9510717 T 28-10-1997	
		US 5846514 A 08-12-1998	