



US 20050100913A1

(19) **United States**

(12) **Patent Application Publication**
Mello et al.

(10) **Pub. No.: US 2005/0100913 A1**

(43) **Pub. Date: May 12, 2005**

(54) **RNA INTERFERENCE PATHWAY GENES AS TOOLS FOR TARGETED GENETIC INTERFERENCE**

(75) Inventors: **Craig C. Mello**, Shrewsbury, MA (US);
Andrew Fire, Baltimore, MD (US);
Hiroaki Tabara, Worcester, MA (US);
Alla Grishok, Shrewsbury, MA (US)

Correspondence Address:
LAHIVE & COCKFIELD, LLP.
28 STATE STREET
BOSTON, MA 02109 (US)

(73) Assignee: **UNIVERSITY OF MASSACHUSETTS MEDICAL**, Boston, MA

(21) Appl. No.: **10/645,735**

(22) Filed: **Aug. 20, 2003**

Related U.S. Application Data

- (62) Division of application No. 09/689,992, filed on Oct. 13, 2000.
- (60) Provisional application No. 60/159,776, filed on Oct. 15, 1999. Provisional application No. 60/193,218, filed on Mar. 30, 2000.

Publication Classification

- (51) **Int. Cl.⁷** **C12Q 1/68**; C07H 21/02;
C12N 9/22
- (52) **U.S. Cl.** **435/6**; 435/69.1; 435/199;
435/320.1; 435/325; 536/23.1

(57) **ABSTRACT**

Genes involved in double-stranded RNA interference (RNAi pathway genes) are identified and used to investigate the RNAi pathway. The genes and their products are also useful for modulating RNAi pathway activity.




FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 3




FIG. 4A

RDE-1	203	VN-----EEIKVQAKNFVYDNNSLRVPESFHDENRFEOSLEAVPRLIEAWFGIIGIKLELGEPEVLN
F48F7.1	235	S-----AVRQFSLSLWVQNSLSTHEDAMEGRVROQPEEAVQAMVDEER--HEPSIKYIPVGRSEFSEFPVFNASGMAGSCPPQASGAV
eIF2C	77	-----GKDRIFKVSIKWVSCVSLQALHDALSGLPSPPTIQAIDVVMR--HEPSMRITPVGSRSEFTASEG-----CS
ZWILLE	226	GIINGPKERSYKVAIKVARANMHLGEFLAGKRADCPQEAVIDDIVER--EISVKRCPVGRSESESDI
Sting	189	-----RAGENIEIKKAVGVSQSTDAEQFVNLTPRAMEGDLKLYSYDYDEQAKIN
RDE-1	267	-----FAIVDKFYNAPKMSLLDYLLIVDPOSNDDDRKDKTKMLMAGKMTIROAARPRIR--OLLENLRKLCAE
F48F7.1	319	AGGAHSAGQYHAESKGGRENWEGFHQSVPRESQWQAMNIDVSAFAFRSMVTEFIIAFVIEFLPQALAEERRASDAQRVKPKETIRGL
eIF2C	144	-----N-----LGGRENWEGFHQSVPRESLMQAMNIDVSAFAFYKAQPVTEFVCEVIDFKSIEEQK--PLTDSQRVAKPTNLEKGL
ZWILLE	297	-----TPQRLGEGLESNCGEYQSIRPTOMGTSINIDMASAANFIEPFPVIEFVAQLHGKDYLSKPLS--DSDRVKIKKGLRQV
Sting	244	-----LENFRMQWPGYQTSIROHENDILICSEICHKVMRTETLYNILSDAIRSDDIQSTFA-----R-AY
RDE-1	336	VVDNEMSRLEERHLTFLDICEENSLVYKVTGKSD--RGRNAKAYDITLTKIYEENKAFIEFFHPLLVKVKSGAKKEYAVPMEHLEVHEKPO
F48F7.1	409	KIETRHGGVRRKRYVGNVTRRPAQOTRFPLOL-----ETGQITGCHVAKVFNKRYRQIKYPHLPCLGVGQCKHWTPEVCNIWV--GO
eIF2C	220	KVEITRHGGVRRKRYVGNVTRRPAHQTFPPLQ-----ESQVVEGVAQYKDRHKLIVRPHLPCLGVGQCKHWTPEVCNIWVA--GO
ZWILLE	372	KVEVTRRANVRESVYAGLITPQTRLEMFVD-----EN--CTMKSVEKFOEMVGTLOHTHLPCLGVGQCKHWTPEVCNIWVA--GO
Sting	305	MGMVILTDYNNKTYRIDDVDFQSTPLCKEKNINDGEISYVDYKRYNLIIRDLPVMSRETDKNIRGGNDQAIMIIPELARATGM--TD
RDE-1	424	RYKNIDLLVQDKFLKRAIRKPHDYKENTKMLKELDESSEELNEVEREGLCSKLOMIECPGKVKKEPMLVNSVNEQIKMTPVIRGFOEK
F48F7.1	494	RCURKUNDVQUSIVKNTARSAPEREREISNLYRKAEEFSAI--EENHEFGTINPAMTEVKGRVLSAPKMLIGGRTR--ATALNCGVNDM
eIF2C	305	RCURKUNDVQUSIVMIRATARSAPDROEETSKIMRSASENTD--EYVREFGIMVKDEMDVTGRVQPSLILYGGNK--AIAATVCGVNDM
ZWILLE	454	RYTKRENEKOITALKRVTCORAEGORNDITRTVQHNAYDQD--PYAKHEGNISEKASVEANILEAPMLKWHENGEKEDCLPQVGNM
Sting	394	AMRADFRTRAMSEHTLNPDRRIERLRMFKRRLKSKCKQSV--ETLNSWNTELDLPAVLEIPANVLEKILFQNKQIFVCDARADWTNTEF

K ne219

FIG. 4B-1

```

RDE-1      514  QLNVPKELCCAVFVNETAGNPCLENDVVKETVELIGGCKFRGIRIGANENRGAQSIMYDATKNEVAFYKNCNTLNTGIGRFEIAATE
F48F7.1    581  KKKQITGIDRNVNLAGNQQ-QDHYKENDIKRMEINQIQREINDAGMBIVENPCCKKVAVGEVDEVEPMKRYEKQNSYSG
eIF2C      392  RNKQFHIGTEKVAACAFAP-ORQCTEVHKSETEQIRKISRDAQMPIQGQPCPKYAQCADSVGPMRHRKNTYAG
ZWILLE     542  MNKKMINGMTVSRVACVNFSS-ASYQENVARGECEIIGQVCEVSCMEFNPEFVPIPIYSARPDQVEKALKHYHTSMNKTK
Sting      482  RTCSMFKNVHENRMYVITPS-----RNLR--EFOEFVQXCIRTASSMKMNICNPKEEIEPDDRNGTYSOAIDNANAN-----

RDE-1      604  AKNMFERLPDKEQVEMFIIISKRQLNAVCFVVKHYCDHTIGVANQHIITSETVTKALASLRHEKSGKRIFYQIALKINAKLGGINQELDWS
F48F7.1    658  ---KQVWVYVYNG-KWVYVNSVVKRYGDIYVIGIAQCYCAKNAIRIUTP-----QTLNQLCKMNYKLGCVNSHLLIPN
eIF2C      469  ---LQVAVVHILPE-KWVYVNSVVKRYGDIYVIGIAQCYVQMKNVORTTP-----QTLNQLCKMNYKLGCVNNEILLPQ
ZWILLE     620  ---GKEEELLLAILLDNNGSILYGDILKRIQEEIQLISQCLTRHVFRIKSK-----QYIADVSIKINNYGGRNTVIVDA
Sting      552  ---DPQIYVAVMRSNPNEEKYSCIKKRTCVDRPVSQVYVTLKVIAPRQQ-----KPTGLMSIATKQVIQMNAKLMGAP

RDE-1      694  EIAEISPEEKERKTMPLTMYVGLDVTHTPYSYCIDYSIAAVASINPGGT-IYRNMIVTQEECRPGERAVAHGRE---RTDILEAKFVK
F48F7.1    725  V-----RPEENEPVYVYVCGDITHPFAGDSRKPISIAAVVSYDNHPS-RYAATVRYOQHRQETIESDITY-----WURE
eIF2C      536  G-----RPPVFOQVTEHPCNDVTHPEAGDCKKPSIAAVVCSMDAHPN-RVCAWVRYOQHRQETIODELAA-----WVRE
ZWILLE     691  IS-----CRIPLVSDIPLTIFGADVTHPEENGEESSEPSIAAVASODWPEVTKVAGLYCAQAHRQFLIQDLYKTKWODPVRGTVSGGVIIRD
Sting      621  W-----QVVPLHGLNTVGFQDVVCHSP--KNKNKSYGAFVATMDQKESFRYFSTVNEHIKQGEIIEQMSVNV-----MAC

```

FIG. 4B-2

* ne300

780 LIREFAENNDNRAPAHIVVVRDGVSDSEMLRVSHDELRSIKSEVKQFMSERDGEDPEPKYTFEIVIQKRHNTRLRRMEKDKPQVWVKDLTP
 792 LIVCFYRNTR - FKPAP IIVVVRDGVSEGGFNMVQVELRARRKACMMIIRGKYQPG - - - I TFIAYQKRHFITRLRNVKDKDQ - - - V
 eIF2C LIHQEYKSTR - FKPPT IIVVVRDGVSEGGFNMVQVELRARRKACIKLEKDYQPG - - - I TFIAYQKRHFITRLRNVKDKDQ - - - V
 775 LISERKATG - QKPLR IIVVVRDGVSEGGFNMVQVELRARRKACASLEPNYQPP - - - V TFIAYQKRHFITRLRNVKDKDQ - - - V
 687 ALRSYQEQHR - SLPER IIVVVRDGVSDGOLGVNSEVNTLQORLDEIYKSAKQEGC - RMTFIIYSKEINSRYSYBTGHRNPV - - -

870 AETDVAAVKQWEEDMKE SKETGIVNPSGGTIVDKLIVSKYKEDFFLASHHGVLGTSRPGHYTVYDDKGM - SQDEVIYKQTYGIAFLSAR
 868 GKAYNIPFGITVVDVGIITHTTEFDFFLCSHAGLGGTSRPGHYTVYDDKGM - SQDEVIYKQTYGIAFLSAR
 679 QKSGNIPAGTIVDKLIVSKYKEDFFLCSHAGLGGTSRPGHYTVYDDKGM - SQDEVIYKQTYGIAFLSAR
 853 DRSSNIPAGTIVDKLIVSKYKEDFFLCSHAGLGGTSRPGHYTVYDDKGM - SQDEVIYKQTYGIAFLSAR
 766 B - GTVDDVITLERYDEFLVSONVRIGTIVTSKNIISDNMGENDKIOMESKMTHTMAYN

E ne297

960 CRKPISLPVPVHYAHILSCEKAKELRYRKYEHYIGDYAQPRURHEMEHFLQTNVKYPGMSFA - - - (SEQ ID NO: 13)
 F48F7.1 CTRSVSHPAPAYAHILVAFRARSYHLVDRHDSGEGSQPSGTSSEDTLLSNMARAVQVILAEINLVS I - - - (SEQ ID NO: 9)
 eIF2C CTRSVSHPAPAYAHILVAFRARSYHLVDRHDSGEGSQPSGTSSEDTLLSNMARAVQVILAEINLVS I - - - (SEQ ID NO: 10)
 921 CTRSVSHPAPAYAHILVAFRARSYHLVDRHDSGEGSQPSGTSSEDTLLSNMARAVQVILAEINLVS I - - - (SEQ ID NO: 6)
 827 YSGTIRVPAVCHYAHKLAELVAEISINRAPSAGLQNLQYFL - - - (SEQ ID NO: 7)

FIG. 4B-3

cagccacaagtgatgaaacatgtcctcgaattttcccgaattggaaaaaggattttatctctcctcgcgacccggta
tggatcattattagcagctataagatatataaagtttgatataataattatagggagatgaaatggcctggaggccaccctg
gtaaatgcgacggcaattctatgaggaagaagtctcttttggtaaatggttccaagttctcagcaaaaatttacgat
cgggaatactacgagatgaaagtgaaatgacaaaaggaaagtattgaaatagaaaaaccaggaaaacccttcccacaaaag
agaaattccaatgtaagtgcttgaattagcaaaaactaattttttcagtcctcgatcgtgcaaaaactcttctggc
aacatcttggcatggaagaagcagacagattttattctcgaagactatgtttttgatgaaaaaggacactgtttatagt
gtttgctgactgaacactgcacatcaaaaatgctggtttcggagaaagtagtaaaaaaggattcgggaaaaaagatg
aaaggattgggaaaaaaatcttatacacaatgatacttacctatcgtaaaaaaattcacctgaacttttagtcgagaaa
atccggaaaaagacggaagcgaatcggagttacaattcctgaaggtttatgaaaaaacacgcattataacaacaa
ttagcttcagaatgttatgaccagaaagttcgtacggccttttgtgaacggagagattaaggtgtgagttgcaata
ataataataatacctcaactattatataatttaagacaattcggaaaaattttgtgtacgataataatcaat

FIG. 5A

tctgcgagttcctgaatcgtttcagatccaaacagattcgaacaarcattagaagtagcaccaagaatcgaagcatggt
ttggaattacattggaatcaaagaattgttcgatgggaacctgtgctcaatttgcaagtaagttgagaaactgga
raaaaaatcatgtgattttgtgaagttgtcgataaaactattctacaatgcaccgaaaatgtctctctggattatct
ctcraattgtcgacccccagtcgtgtaacgatgatgtacgaaaagatcttaaaacaaaactgatggcgggaaaaatgac
aatcagacaagccgcgcccgaagaattcgacaattattgaaaaattgaaagctgaaatgcgacagaagttgggataac
aaaagttagtttaattatcaacaattaatatacaaaattgatttcaggctcgagattgacagaacgacatctgacatt
tctagatttgtgcgaggaaaaactctctgtttataaaagtcactggtaaatcggacagaggaagaatgcaaaaaagtagc
atactacattgttcaaaatctatgaggaaaaacaaaagttcattgagttccccacctaccactagcaaaagttaaaggt
ggagcaaaagaatacgtgtaccaatggaacatcttgaagttcatgagaagccacaagatacaagaatcgaattgate
ggtagtcaagacaagtttctaaagcgagctacacgaaaacctcagactacaagaaaataccctaaaaatgctgaaa
aattggatttctctctgaagagctaaatgttgaagattggattatgctccaaactcagatgatcgaatgtcca
ggaaagggtttgaaagagccaatgcttgaatagtgtaatgaacaaatataatgacaccagtgattcgtggatttca
agaaaaacaattgaaatgtggttcccgaaaaagaacttgtgtgctgtttttagtcaacgaaacagcgggaaatccat
gettagaagagaacgacgttgtgaagtgtttctacgtagattattccgaaatatttccagtaagttctacaccgaact
aattggtggttcaagttccgtggaatacgaattggtgccaatgaaaacagaggagcgcaatctattatgtacgacgca
cgaaaaatgaatatgccgtaagtttcagaaaattgaaagtttttaaatcatatttacagttctacaaaaattgtacac
taaataccggaatcggtagatttgaatatagccgcaacagaagcgaagaatatgtttgaacgttctccgataaagaaca
aaagcttaatgttcatcattatctccaaacgacaactgaatgcttacggttttgtgaaacattattgcgatcacacat
cgggttagctaatcagcatattacttctgaaacagtcacaaaagcttggcactcactaaggcagagaaaggatcaaaac
gaatctctatcaaatgcattgaaaatcaacgcgaaattaggaggattaccaggagcttactggtcagaaattgca
gaaatatcaccagaagaaaaagaagacggaaaacaatgccattaactatgtatgttgaattgatgtaactcatccaa
ctctacagtggaattgattattctatagcggctgtagtagcgagtatcaatccaggtggaactatctatcgaaatatga
tttgactcaagaagaatgtcgtcccggtagcgtgcagtggtcatggacgggaaagaacagatatttggagcaaaa
ttcgtgaaattgctcagagaattcgcagaagtgagttgtcttgagattttaaagatctctgggatttttaattttttg

FIG. 5B

taaaccttcagaacaacgacaatcgagcaccagcgcatatgtgtatgtctatcgagacggagttagcgattcgggagatgcta
cgtgtagtcatgatgagccttcgatcttttaaaaagcgaaagtaaaaacaattcatgtcggaacgggatgsgaagatccagga
gcccgaagtacacgcttcattgtgattcagaaaagacacaatacacgattgcttcgaaatgaaaaagataagccagtg
tcataaagatcttactcctcctgfgaacaagatgtcgtctgttgcctgctgctgtaacaatggsgagggatatgaaaagaaagc
aaagaaactggaattgtgaaccatcatccggacaactgtggataaacttatcgtttcgaatacaacaattcgaattttt
cttggcatctcatgtgcttggtagcatctcgtccaggacattacactgtttatgtatgacgataaaggaatgagcc
aagatgaagtctatgtaagcgttttgaatagcagttagcatttaggattttgtaataccgcatatagttattataaaaa
aatgttcagaaaaatgacctacggactgtctttctctcctctagatgtcgaaaaccatctcgttgcctgttccggttc
attaigtctattatcatgtgaaaagcgaagagcittatcgaaacttaacagggaacattacatcgggtgactatgcacag
ccaaggactcgacagaaatggaaacattttctccaactaacggaagtagccctggaaatgctgttcgcataacattttgc
aaaagtgccccgttcaatcaaatcttcaattgtagatattgtactacttttttaagccccggttcaaaaatt
cattccatgactaacgttttcataaattacttgaattt (SEQ ID NO:1)

FIG. 5C

CAGCCACAAAGTGATGAAAC- 5' UTR

1/1 31/11
 ATG TCC TCG AAT TTT CCC GAA TTG GAA AAA GGA TTT TAT CGT CAT TCT CTC GAT CCG GAG
 met ser ser asn phe pro glu leu glu lys gly phe tyr arg his ser leu asp pro glu

61/21 91/31
 ATG AAA TGG CTT GCG AGG CCC ACT GGT AAA TGC GAC GGC AAA TTC TAT GAG AAG AAA GTA
 met lys trp leu ala arg pro thr gly lys cys asp gly lys phe tyr glu lys lys val

121/41 151/51
 CTT CTT TTG GTA AAT TGG TTC AAG TTC TCC AGC AAA ATT TAC GAT CGG GAA TAC TAC GAG
 leu leu leu val asn trp phe lys phe ser ser lys ile tyr asp arg glu tyr tyr glu

181/61 211/71
 TAT GAA GTG AAA ATG ACA AAG GAA GTA TTG AAT AGA AAA CCA GGA AAA CCT TTC CCA AAA
 tyr glu val lys met thr lys glu val leu asn arg lys pro gly lys pro phe pro lys

241/81 271/91
 AAG ACA GAA ATT CCA ATT CCC GAT CGT GCA AAA CTC TTC TGG CAA CAT CTT CGG CAT GAG
 lys thr glu ile pro ile pro asp arg ala lys leu phe trp gln his leu arg his glu

301/101 331/111
 AAG AAG CAG ACA GAT TTT ATT CTC GAA GAC TAT GTT TTT GAT GAA AAG GAC ACT GTT TAT
 lys lys gln thr asp phe ile leu glu asp tyr val phe asp glu lys asp thr val tyr

361/121 391/131
 AGT GTT TGT CGA CTG AAC ACT GTC ACA TCA AAA ATG CTG GTT TCG GAG AAA GTA GTA AAA
 ser val cys arg leu asn thr val thr ser lys met leu val ser glu lys val val lys

421/141 451/151
 AAG GAT TCG GAG AAA AAA GAT GAA AAG GAT TTG GAG AAA AAA ATC TTA TAC ACA ATG ATA
 lys asp ser glu lys lys asp glu lys asp leu glu lys lys ile leu tyr thr met ile

481/161 511/171
 CTT ACC TAT CGT AAA AAA TTT CAC CTG AAC TTT AGT CGA GAA AAT CCG GAA AAA GAC GAA
 leu thr tyr arg lys lys phe his leu asn phe ser arg glu asn pro glu lys asp glu

541/181 571/191
 GAA GCG AAT CGG AGT TAC AAA TTC CTG AAG AAT GTT ATG ACC CAG AAA GTT CGC TAC GCG
 glu ala asn arg ser tyr lys phe leu lys asn val met thr gln lys val arg tyr ala

601/201 631/211
 CCT TTT GTG AAC GAG GAG ATT AAA GTA CAA TTC GCG AAA AAT TTT GTG TAC GAT AAT AAT
 pro phe val asn glu glu ile lys val gln phe ala lys asn phe val tyr asp asn asn

661/221 691/231
 TCA ATT CTG CGA GTT CCT GAA TCG TTT CAC GAT CCA AAC AGA TTC GAA CAA TCA TTA GAA
 ser ile leu arg val pro glu ser phe his asp pro asn arg phe glu gln ser leu glu

FIG. 6A

721/241	751/251
GTA GCA CCA AGA ATC GAA GCA TGG TTT GGA ATT TAC ATT GGA ATC AAA GAA TTG TTC GAT	
val ala pro arg ile glu ala trp phe gly ile tyr ile gly ile lys glu leu phe asp	
781/261	811/271
GGT GAA CCT GTG CTC AAT TTT GCA ATT GTC GAT AAA CTA TTC TAC AAT GCA CCG AAA ATG	
gly glu pro val leu asn phe ala ile val asp lys leu phe tyr asn ala pro lys met	
841/281	871/291
TCT CTT CTG GAT TAT CTT CTC CTA ATT GTC GAC CCC CAG TCG TGT AAC GAT GAT GTA CGA	
ser leu leu asp tyr leu leu leu ile val asp pro gln ser cys asn asp asp val arg	
901/301	931/311
AAA GAT CTT AAA ACA AAA CTG ATG GCG GGA AAA ATG ACA ATC AGA CAA GCC GCG CGG CCA	
lys asp leu lys thr lys leu met ala gly lys met thr ile arg gln ala ala arg pro	
961/321	991/331
AGA ATT CGA CAA TTA TTG GAA AAT TTG AAG CTG AAA TGC GCA GAA GTT TGG GAT AAC GAA	
arg ile arg gln leu leu glu asn leu lys leu lys cys ala glu val trp asp asn glu	
1021/341	1051/351
ATG TCG AGA TTG ACA GAA CGA CAT CTG ACA TTT CTA GAT TTG TGC GAG GAA AAC TCT CTT	
met ser arg leu thr glu arg his leu thr phe leu asp leu cys glu glu asn ser leu	
1081/361	1111/371
GTT TAT AAA GTC ACT GGT AAA TCG GAC AGA GGA AGA AAT GCA AAA AAG TAC GAT ACT ACA	
val tyr lys val thr gly lys ser asp arg gly arg asn ala lys lys tyr asp thr thr	
1141/381	1171/391
TTG TTC AAA ATC TAT GAG GAA AAC AAA AAG TTC ATT GAG TTT CCC CAC CTA CCA CTA GTC	
leu phe lys ile tyr glu glu asn lys lys phe ile glu phe pro his leu pro leu val	
1201/401	1231/411
AAA GTT AAA AGT GGA GCA AAA GAA TAC GCT GTA CCA ATG GAA CAT CTT GAA GTT CAT GAG	
lys val lys ser gly ala lys glu tyr ala val pro met glu his leu glu val his glu	
1261/421	1291/431
AAG CCA CAA AGA TAC AAG AAT CGA ATT GAT CTG GTG ATG CAA GAC AAG TTT CTA AAG CGA	
lys pro gln arg tyr lys asn arg ile asp leu val met gln asp lys phe leu lys arg	
1321/441	1351/451
GCT ACA CGA AAA CCT CAC GAC TAC AAA GAA AAT ACC CTA AAA ATG CTG AAA GAA TTG GAT	
ala thr arg lys pro his asp tyr lys glu asn thr leu lys met leu lys glu leu asp	
1381/461	1411/471
TTC TCT TCT GAA GAG CTA AAT TTT GTT GAA AGA TTT GGA TTA TGC TCC AAA CTT CAG ATG	
phe ser ser glu glu leu asn phe val glu arg phe gly leu cys ser lys leu gln met	
1441/481	1471/491
ATC GAA TGT CCA GGA AAG GTT TTG AAA GAG CCA ATG CTT GTG AAT AGT GTA AAT GAA CAA	
ile glu cys pro gly lys val leu lys glu pro met leu val asn ser val asn glu gln	
1501/501	1531/511
ATT AAA ATG ACA CCA GTG ATT CGT GGA TTT CAA GAA AAA CAA TTG AAT GTG GTT CCC GAA	
ile lys met thr pro val ile arg gly phe gln glu lys gln leu asn val val pro glu	

FIG. 6B

1561/521	1591/531
AAA GAA CTT TGC TGT GCT GTT TTT GTA GTC AAC GAA ACA GCG GGA AAT CCA TGC TTA GAA lys glu leu cys cys ala val phe val val asn glu thr ala gly asn pro cys leu glu	
1621/541	1651/551
GAG AAC GAC GTT GTT AAG TTC TAC ACC GAA CTA ATT GGT GGT TGC AAG TTC CGT GGA ATA glu asn asp val val lys phe tyr thr glu leu ile gly gly cys lys phe arg gly ile	
1681/561	1711/571
CGA ATT GGT GCC AAT GAA AAC AGA GGA GCG CAA TCT ATT ATG TAC GAC GCG ACG AAA AAT arg ile gly ala asn glu asn arg gly ala gln ser ile met tyr asp ala thr lys asn	
1741/581	1771/591
GAA TAT GCC TTC TAC AAA AAT TGT ACA CTA AAT ACC GGA ATC GGT AGA TTT GAA ATA GCC glu tyr ala phe tyr lys asn cys thr leu asn thr gly ile gly arg phe glu ile ala	
1801/601	1831/611
GCA ACA GAA GCG AAG AAT ATG TTT GAA CGT CTT CCC GAT AAA GAA CAA AAA GTC TTA ATG ala thr glu ala lys asn met phe glu arg leu pro asp lys glu gln lys val leu met	
1861/621	1891/631
TTC ATT ATC ATT TCC AAA CGA CAA CTG AAT GCT TAC GGT TTT GTG AAA CAT TAT TGC GAT phe ile ile ile ser lys arg gln leu asn ala tyr gly phe val lys his tyr cys asp	
1921/641	1951/651
CAC ACC ATC GGT GTA GCT AAT CAG CAT ATT ACT TCT GAA ACA GTC ACA AAA GCT TTG GCA his thr ile gly val ala asn gln his ile thr ser glu thr val thr lys ala leu ala	
1981/661	2011/671
TCA CTA AGG CAC GAG AAA GGA TCA AAA CGA ATT TTC TAT CAA ATT GCA TTG AAA ATC AAC ser leu arg his glu lys gly ser lys arg ile phe tyr gln ile ala leu lys ile asn	
2041/681	2071/691
GCG AAA TTA GGA GGT ATT AAC CAG GAG CTT GAC TGG TCA GAA ATT GCA GAA ATA TCA CCA ala lys leu gly gly ile asn gln glu leu asp trp ser glu ile ala glu ile ser pro	
2101/701	2131/711
GAA GAA AAA GAA AGA CGG AAA ACA ATG CCA TTA ACT ATG TAT GTT GGA ATT GAT GTA ACT glu glu lys glu arg arg lys thr met pro leu thr met tyr val gly ile asp val thr	
2161/721	2191/731
CAT CCA ACC TCC TAC AGT GGA ATT GAT TAT TCT ATA GCG GCT GTA GTA GCG AGT ATC AAT his pro thr ser tyr ser gly ile asp tyr ser ile ala ala val val ala ser ile asn	
2221/741	2251/751
CCA GGT GGA ACT ATC TAT CGA AAT ATG ATT GTG ACT CAA GAA GAA TGT CGT CCC GGT GAG pro gly gly thr ile tyr arg asn met ile val thr gln glu glu cys arg pro gly glu	
2281/761	2311/771
CGT GCA GTG GCT CAT GGA CGG GAA AGA ACA GAT ATT TTG GAA GCA AAG TTC GTG AAA TTG arg ala val ala his gly arg glu arg thr asp ile leu glu ala lys phe val lys leu	
2341/781	2371/791
CTC AGA GAA TTC GCA GAA AAC AAC GAC AAT CGA GCA CCA GCG CAT ATT GTA GTC TAT CGA leu arg glu phe ala glu asn asn asp asn arg ala pro ala his ile val val tyr arg	

FIG. 6C

2401/801	2431/811
GAC GGA GTT AGC GAT TCG GAG ATG CTA CGT GTT AGT CAT GAT GAG CTT CGA TCT TTA AAA	
asp gly val ser asp ser glu met leu arg val ser his asp glu leu arg ser leu lys	
2461/821	2491/831
AGC GAA GTA AAA CAA TTC ATG TCG GAA CGG GAT GGA GAA GAT CCA GAG CCG AAG TAC ACG	
ser glu val lys gln phe met ser glu arg asp gly glu asp pro glu pro lys tyr thr	
2521/841	2551/851
TTC ATT GTG ATT CAG AAA AGA CAC AAT ACA CGA TTG CTT CGA AGA ATG GAA AAA GAT AAG	
phe ile val ile gln lys arg his asn thr arg leu leu arg arg met glu lys asp lys	
2581/861	2611/871
CCA GTG GTC AAT AAA GAT CTT ACT CCT GCT GAA ACA GAT GTC GCT GTT GCT GCT GTT AAA	
pro val val asn lys asp leu thr pro ala glu thr asp val ala val ala ala val lys	
2641/881	2671/891
CAA TGG GAG GAG GAT ATG AAA GAA AGC AAA GAA ACT GGA ATT GTG AAC CCA TCA TCC GGA	
gln trp glu glu asp met lys glu ser lys glu thr gly ile val asn pro ser ser gly	
2701/901	2731/911
ACA ACT GTG GAT AAA CTT ATC GTT TCG AAA TAC AAA TTC GAT TTT TTC TTG GCA TCT CAT	
thr thr val asp lys leu ile val ser lys tyr lys phe asp phe phe leu ala ser his	
2761/921	2791/931
CAT GGT GTC CTT GGT ACA TCT CGT CCA GGA CAT TAC ACT GTT ATG TAT GAC GAT AAA GGA	
his gly val leu gly thr ser arg pro gly his tyr thr val met tyr asp asp lys gly	
2821/941	2851/951
ATG AGC CAA GAT GAA GTC TAT AAA ATG ACC TAC GGA CTT GCT TTT CTC TCT GCT AGA TGT	
met ser gln asp glu val tyr lys met thr tyr gly leu ala phe leu ser ala arg cys	
2881/961	2911/971
CGA AAA CCC ATC TCG TTG CCT GTT CCG GTT CAT TAT GCT CAT TTA TCA TGT GAA AAA GCG	
arg lys pro ile ser leu pro val pro val his tyr ala his leu ser cys glu lys ala	
2941/981	2971/991
AAA GAG CTT TAT CGA ACT TAC AAG GAA CAT TAC ATC GGT GAC TAT GCA CAG CCA CGG ACT	
lys glu leu tyr arg thr tyr lys glu his tyr ile gly asp tyr ala gln pro arg thr	
3001/1001	3031/1011
CGA CAC GAA ATG GAA CAT TTT CTC CAA ACT AAC GTG AAG TAC CCT GGA ATG TCG TTC GCA	
arg his glu met glu his phe leu gln thr asn val lys tyr pro gly met ser phe ala	
3061/1021	3091/1031
TAA CAT TTT GCA AAA GTG TCG CCC GTT TCA ATC AAA TTT TTC AAT TGT AGA TAT TGT ACT	
OCH (SEQ ID NO:3)	
3121/1041	3151/1051
TAC TTT TTT TTA AAG CCC GGT TTC AAA AAT TCA TTC CAT GAC TCC CGT TTT CAT AAA TTA	
3181/1061	
CTT GAA ATT TAA AAA AAA AAA AAA AAA (SEQ ID NO:2)	

FIG. 6D





FIG. 7B

Injected P0			F1	
♀	$\frac{\text{pos-1 dsRNA} \diagup \text{rde-1 unc-42}}{+}$	self X	<i>rde-1</i> (-) 11/24 <i>rde-1</i> (+) 9/72	
♀	$\frac{\text{rde-2 unc-13}}{+}$	→	<i>rde-2</i> (-) 0/39 <i>rde-2</i> (+) 23/78	
♀	$\frac{\text{mut-7 dpy-17}}{+}$	→	<i>mut-7</i> (-) 0/15 <i>mut-7</i> (+) 20/50	
♀	$\frac{\text{rde-4 unc-69}}{+}$	→	<i>rde-4</i> (-) 5/15 <i>rde-4</i> (+) 11/48	

FIG. 8A

P0		Injected F1	
♀	$\frac{\text{rde-1 unc-42}}{+}$	$\frac{\text{pos-1 dsRNA} \diagup \text{rde-1 unc-42}}{\text{rde-1 unc-42}}$	0/37
♀	$\frac{\text{rde-4 unc-69}}{+}$	$\frac{\text{rde-4 unc-69}}{\text{rde-4 unc-69}}$	0/37

FIG. 8B




FIG. 9A




FIG. 9B

10 20 30 40 50 60
 ATGGATTTAACCAAACCTAACGTTTTGAAAGCGTTTTTCGGTGGATCAGATGTTCCCTATGAAG
 M D L T K L T F E S J F G G S D V P M K

 70 80 90 100 110 120
 CCTTCCCGATCGGAGGATAACAAAACGCCAAGAAACAGAACAGATTTGGAGATGTTTCTG
 P S R S E D N K T P R N R T D L E M F L

 130 140 150 160 170 180
 AAGAAAACCTCCCTCATGGTACTAGAAGAGGCTGCTAAGGCTGTCTATCAAAGACGCCA
 K K T P L M V L E E A A K A V Y Q K T P

 190 200 210 220 230 240
 ACTTGGGGCACTGTCGAACCTCCTGAAGGCTTCGAGATGACGTTGATTCTGAATGAAATT
 T W G T V E L P E G F E M T L I L N E I

 250 260 270 280 290 300
 ACTGTAAAAGGCCAGGCAACAAGCAAGAAAGCTGCGAGACAAAAGGCTGCTGTTGAATAT
 T V K G Q A T S K K A A R Q K A A V E Y

 310 320 330 340 350 360
 TTACGCAAGGTTGTGGAGAAAGGAAAGCACGAAATCTTTTTTCATTCCTGGAACAACCAA
 L R K V V E K G K H E I F F I P G T T K

 370 380 390 400 410 420
 GAAGAAGCTCTTTTGAATATTGATCAAATATCGGATAAGGCTGAGGAATTGAAACGATCA
 E E A L S N I D Q I S D K A E E L K R S

 430 440 450 460 470 480
 ACTTCAGATGCTGTTTCAGGATAACGATAACGATGATTTCGATTCCTACAAGTGCTGAATTT
 T S D A V Q D N D N D D S I P T S A E F

 490 500 510 520 530 540
 CCACCTGGTATTTTCGCCAACCGAGAATTGGGTTCGGAAAGTTGCAGGAAAAATCTCAAAA
 P P G I S P T E N W V G K L Q E K S Q K

 550 560 570 580 590 600
 AGCAAGCTGCAAGCCCCAATCTATGAAGATTCCAAGAATGAGAGAACCGAGCGTTTTCTTG
 S K L Q A P I Y E D S K N E R T E R F L

 610 620 630 640 650 660
 GTTATATGCACGATGTGCAATCAAAAAACCGAGGAATCAGAAGTAAGAAGAAGGACGCA
 V I C T M C N Q K T R G I R S K K K D A

 670 680 690 700 710 720
 AAGAATCTTGCAAGCATGGTTGATGTGGAAGCGTTGGAAGACGGTATCGAATCTCTGGAA
 K N L A A W L M W K A L E D G I E S L E

 730 740 750 760 770 780
 TCATATGATATGGTTGATGTGATTTGAAAATTTGGAAGAAGCTGAACATTTACTCGAAATT
 S Y D M V D V I E N L E E A E H L L E I

FIG. 10A

790 800 810 820 830 840
CAGGATCAAGCATCCAAGATTAAAGACAAGCATTCCGCACTGATTGATATACTCTCGGAC
Q D Q A S K I K D K H S A L I D I L S D

850 860 870 880 890 900
AAGAAAAGATTTTCAGACTACAGCATGGATTTCAACGTATTATCAGTGAGCACAATGGGA
K K R F S D Y S M D F N W L S V S T M G

910 920 930 940 950 960
ATACATCAGGTGCTATTGGAAATCTCGTTCGGCGTCTAGTTTCTCCAGACCCCGACGAT
I H Q V L L E I S F R R L V S P D P D D

970 980 990 1000 1010 1020
TTGGAAATGGGAGCAGAACACACCCAGACTGAAGAAATTATGAAGGCTACTGCCGAGAAG
L E M G A E H T Q T E E I M K A T A E K

1030 1040 1050 1060 1070 1080
GAAAAGCTACGGAAGAAGAATATGCCAGATTCCGGGCGCTAGTGTTTGCTGGACATGGT
E K L R K K N M P D S G P L V F A G H G

1090 1100 1110 1120 1130 1140
TCATCGGCGGAAGAGGCTAAACAGTGTGCTTGTAATCGGCGATTATCCATTTCAACACC
S S A E E A K Q C A C K S A I I H F N T

1150 1160 1170 1180 1190 1200
TATGATTTACGGATTGAAAATATTATTGCGTATTCCTGAAAAATGAAGCGTCTGAATGA
Y D F T D * K Y Y C V F L K N E A S E *

1210 1220 1230
TTATAAAAAAAAAAAAAAAAAA (SEQ ID NO:4)
L * K K K K K (SEQ ID NO:5)

FIG. 10B

X1RBP P V G S L Q E L A V Q K G W R L - P E Y I V A Q E S G P P H K R E F T I I C R V E T F - V E T G S C I S K Q - V A K R V A A E K L L T K F (SEQ ID NO: 11)
HsPKR F M E E L N T Y R - Q K Q G V V - L K Y Q E L P N S G P P H D R R F T F Q V I I D G R E F P E G E C R S K K - E A K N A A A K L A V E I F (SEQ ID NO: 12)
RDE-4 W V C K L Q E K S - Q K S K L Q A P I Y E D S K N - - - E R T E R F L V I C T M C N Q - K T R G I - R S K K K D A K N I A A W L M W K A L

Q E Y H F S K Q K

CONSENSUS P - - - L - E - - - Q - - - - - Y - - - - - G P - H - - - F - - - V - - - G - - - - - G - G - S K K - - - A K - - - A A - - - A L - - - L (SEQ ID NO:8)





FIG. 11

Rescue of *rde-4*:

FIG. 12



RNA INTERFERENCE PATHWAY GENES AS TOOLS FOR TARGETED GENETIC INTERFERENCE

RELATED APPLICATION INFORMATION

[0001] This application claims priority from provisional application Ser. No. 60/159,776, filed Oct. 15, 1999, and 60/193,218, filed Mar. 30, 2000.

STATEMENT AS TO FEDERALLY SPONSORED RESEARCH

[0002] Funding for the work described herein was provided by the federal government (GM58800 and GM37706), which has certain rights in the invention.

FIELD OF THE INVENTION

[0003] This invention relates to the discovery of genes whose expression products are involved in mediation of genetic interference.

BACKGROUND OF THE INVENTION

[0004] All eukaryotic organisms share similar mechanisms for information transfer from DNA to RNA to protein. RNA interference represents an efficient mechanism for inactivating this transfer process for a specific targeted gene. Targeting is mediated by the sequence of the RNA molecule introduced to the cell. Double-stranded (ds) RNA can induce sequence-specific inhibition of gene function (genetic interference) in several organisms including the nematode, *C. elegans* (Fire, et al., 1998, *Nature* 391:806-811), plants, trypanosomes, *Drosophila*, and planaria (Waterhouse et al., 1998, *Proc. Natl. Acad. Sci. USA* 94:13959-13964; Ngo et al., 1998, *Proc. Natl. Acad. Sci. USA* 95:14687-14692; Kennerdell and Carthew, 1998, *Cell* 95:1017-1026; Misquitta and Patterson, 1999, *Proc. Natl. Acad. Sci. USA* 96:1451-1456; Sanchez-Alvorado and Newmark, 1999, *Proc. Natl. Acad. Sci. USA* 96:5049-5054). The discovery that dsRNA can induce genetic interference in organisms from several distinct phyla suggests a conserved mechanism and perhaps a conserved physiological role for the interference process. Although several models of RNAi have been proposed (Baulcombe, 1999, *Curr. Biol.* 9:R599-R601; Sharp, 1999, *Genes & Dev.* 13:139-141) the mechanisms of action of specific components of the pathway are not known.

[0005] Attempts to overexpress a gene (e.g., a transgene) often lead only to transient expression of the gene. Furthermore, the even more undesirable effect of "cosuppression" can occur in which a corresponding endogenous copy of the transgene becomes inactivated. In some cases, transgene silencing leads to problems with the commercial or therapeutic application of transgenic technology to alter the genetic makeup of a cell, organism, or human patient.

SUMMARY OF THE INVENTION

[0006] The present invention relates to the discovery of RNA interference (RNAi) pathway genes which are involved in mediating double-stranded RNA-dependent gene silencing (genetic interference). RNAi requires a set of conserved cellular factors to suppress gene expression. These factors are the components of the RNAi pathway. The RNAi pathway mutations and genes described herein (e.g., *rde-1*, *rde-2*, *rde-3*, *rde-4*, *rde-5*, *mut-2*, and *mut-7*), and their

protein products (e.g., RDE-1 and RDE-4) are useful tools for investigating the mechanisms involved in RNAi and developing methods of modulating the RNAi pathway. The sequences and methods described herein are useful for modulating the RNAi pathway and may be used in conjunction with other methods involving the use of genetic inhibition by dsRNA (e.g., see U.S. Ser. No. 09/215,257, filed Dec. 18, 1998, incorporated herein by reference in its entirety).

[0007] RNAi pathway components (e.g., RDE-1, RDE-4) provide activities necessary for interference. These activities may be absent or not sufficiently activated in many cell types, including those of organisms such as humans in which genetic interference may have potential therapeutic value. Components of the RNAi pathway in *C. elegans* may be sufficient when provided through transgenesis or as direct RNA:protein complexes to activate or directly mediate genetic interference in heterologous cells that are deficient in RNAi.

[0008] Nucleic acid sequences encoding RNAi pathway components (e.g., RDE-1, RDE-4) are useful, e.g., for studying the regulation of the RNAi pathway. Such sequences can also be used to generate knockout strains of animals such as *C. elegans*.

[0009] The nucleic acids of the invention include nucleic acids that hybridize, e.g., under stringent hybridization conditions (as defined herein), to all or a portion of the nucleotide sequence of SEQ ID NO:1 (FIG. 5A-C) or its complement; SEQ ID NO:2 (FIG. 6A-D) or its complement, or SEQ ID NO:4 or its complement. The hybridizing portion of the hybridizing nucleic acids are preferably 20, 30, 50, or 70 bases long. Preferably, the hybridizing portion of the hybridizing nucleic acid is 80%, more preferably 95%, or even 98% or 100% identical to the sequence of a portion or all of a nucleic acid encoding an RDE-1 polypeptide or an RDE-4 polypeptide. Hybridizing nucleic acids of the type described above can be used as a cloning probe, a primer (e.g., a PCR primer), or a diagnostic probe. Preferred hybridizing nucleic acids encode a polypeptide having some or all of the biological activities possessed by a naturally-occurring RDE-1 polypeptide or an RDE-4 polypeptide e.g., as determined in the assays described below.

[0010] Hybridizing nucleic acids may encode a protein that is shorter or longer than the RDE-1 protein or RDE-4 protein described herein. Hybridizing nucleic acids may also encode proteins that are related to RDE-1 or RDE-4 (e.g., proteins encoded by genes that include a portion having a relatively high degree of identity to the *rde-1* gene or *rde-4* gene described herein).

[0011] The invention also features purified or isolated RDE-1 polypeptides and RDE-4 polypeptides. RDE-1 and RDE-4 polypeptides are useful for generating and testing antibodies that specifically bind to an RDE-1 or an RDE-4. Such antibodies can be used, e.g., for studying the RNAi pathway in *C. elegans* and other organisms. As used herein, both "protein" and "polypeptide" mean any chain of amino acids, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation). Thus, the term "RNAi pathway polypeptide" includes a full-length, naturally occurring RNAi pathway polypeptide such as RDE-1 protein or RDE-4 protein, as well as recombinantly or synthetically produced polypeptides that correspond to a

full-length, naturally occurring RDE-1 protein, RDE-4 protein, or to particular domains or portions of a naturally occurring RNAi pathway protein.

[0012] RNAi pathway mutations and strains harboring those mutations (e.g., rde-1, rde-2, rde-3, rde-4, rde-5) are useful for studying the RNAi pathway, including identification of modulators of the RNAi pathway.

[0013] RNAi pathway components (e.g., those associated with mut-7 and rde-2) can be used to desilence or prevent silencing of transgenes. To facilitate this function, such RNAi pathway components are inhibited using specific inhibitors of an RNAi pathway gene or its product.

[0014] In one embodiment, the invention includes an isolated nucleic acid molecule comprising a nucleotide sequence encoding an RDE-1 polypeptide. The nucleic acid molecule hybridizes under high stringency conditions to the nucleic acid sequence of Genbank Accession No. AF180730 (SEQ ID NO:2) or its complement, or the sequence of SEQ ID NO:1 or its complement. In one embodiment, the isolated nucleic acid can complement an rde-1 mutation. The invention also encompasses an isolated nucleic acid whose nucleotide sequence encodes the amino acid sequence of SEQ ID NO:3.

[0015] The invention also encompasses a substantially pure RDE-1 polypeptide encoded by the isolated nucleic acids described herein.

[0016] The invention features an antibody that specifically binds to an RDE-1 polypeptide.

[0017] The invention also includes a method of enhancing the expression of a transgene in a cell, the method comprising decreasing activity of the RNAi pathway. In one embodiment of this invention, rde-2 expression or activity is decreased.

[0018] The invention also features an isolated nucleic acid molecule comprising a nucleotide sequence encoding an RDE-4 polypeptide, wherein the nucleic acid molecule hybridizes under high stringency conditions to the nucleic acid sequence of SEQ ID NO:4 or its complement. The invention also encompasses an isolated nucleic acid encoding an RDE-4 polypeptide, wherein the nucleic acid can complement an rde-4 mutation. The invention also encompasses an isolated nucleic acid encoding an RDE-4 polypeptide, in which the nucleotide sequence encodes the amino acid sequence of SEQ ID NO:5.

[0019] The invention also features a substantially pure RDE-4 polypeptide encoded by the isolated nucleic acids described herein.

[0020] In another embodiment the invention features an antibody that specifically binds to an RDE-4 polypeptide.

[0021] The invention also features a method of preparing an RNAi agent, the method includes incubating a dsRNA in the presence of an RDE-1 protein and an RDE-4 protein.

[0022] The invention also features a method of inhibiting the activity of a gene by introducing an RNAi agent into a cell, such that the dsRNA component of the RNAi agent is targeted to the gene. In another embodiment of the invention, the cell contains an exogenous RNAi pathway sequence. The exogenous RNAi pathway sequence can be an RDE-1 polypeptide or an RDE-4 polypeptide. In still

another embodiment, a dsRNA is introduced into a cell containing an exogenous RNAi pathway sequence such as nucleic acid sequence expressing an RDE-1 or RDE-4.

[0023] An RNAi pathway component is a protein or nucleic acid that is involved in promoting dsRNA-mediated genetic interference. A nucleic acid component can be an RNA or DNA molecule. A mutation in a gene encoding an RNAi pathway component may decrease or increase RNAi pathway activity.

[0024] An RNAi pathway protein is a protein that is involved in promoting dsRNA mediated genetic interference.

[0025] A "substantially pure DNA" is a DNA that is not immediately contiguous with (i.e., covalently linked to) both of the coding sequences with which it is immediately contiguous (i.e., one at the 5' end and one at the 3' end) in the naturally-occurring genome of the organism from which the DNA of the invention is derived. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote; or which exists as a separate molecule (e.g., a cDNA or a genomic or cDNA fragment produced by PCR (polymerase chain reaction) or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequences.

[0026] By "inhibited RNAi pathway" is meant decreased inhibitory activity of a dsRNA which results in at least two-fold less inhibition by a dsRNA relative to its ability to cause inhibition in a wild type cell. Techniques for measuring RNAi pathway activity are described herein. The pathway can be inhibited by inhibiting a component of the pathway (e.g., RDE-1) or mutating the component so that its function is reduced.

[0027] A "substantially pure polypeptide" is a polypeptide, e.g., an RNAi pathway polypeptide or fragment thereof, that is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, RNAi pathway polypeptide or fragment. A substantially pure RNAi pathway polypeptide or fragment thereof is obtained, for example, by extraction from a natural source; by expression of a recombinant nucleic acid encoding an RNAi pathway polypeptide or fragment thereof; or by chemically synthesizing the polypeptide or fragment. Purity can be measured by any appropriate method, e.g., column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

[0028] By "specifically binds" is meant a molecule that binds to a particular entity, e.g., an RNAi pathway polypeptide, but which does not substantially recognize or bind to other molecules in a sample, e.g., a biological sample, which includes the particular entity, e.g., RDE-1.

[0029] An RNAi agent is a dsRNA molecule that has been treated with those components of the RNAi pathway that are required to confer RNAi activity on the dsRNA. For example, treatment of a dsRNA under conditions that include RDE-1 and RDE-4 results in an RNAi agent. Injection of such an agent into an animal that is mutant for RDE-1

and RDE-4 will result in activation of the RNAi pathway with respect to a targeted gene. Typically, the dsRNA used to trigger the formation of the RNAi agent is selected to be an RNA corresponding to all or a portion of the nucleotide sequence of the targeted gene.

[0030] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

[0031] Other features and advantages of the invention will be apparent from the detailed description, and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

[0032] FIG. 1A illustrates the genetic scheme used to identify *rde* mutants.

[0033] FIG. 1B is an illustration summarizing data from the genetic mapping of *rde* and *mut* mutations. The vertical bars represent chromosomes; LGI, LGIII, and LGV. Reference genetic markers are indicated at the right of each chromosome and the relative genetic positions of the *rde* and *mut* alleles are indicated at the left.

[0034] FIG. 2A is a graphical representation of experiments investigating the sensitivity of *rde* and *mut* strains to RNAi by microinjection. The RNA species indicated above each graph was injected at high concentration (pos-1: 7 mg/ml, par-2: 3 mg/ml, sqt-3: 7 mg/ml). The strains receiving injection are indicated at the left and the horizontal bar graphs reflect the percent of progeny that exhibited genetic interference. The *Unc* marker mutants used are also indicated. The percent embryonic lethality of F1 progeny is plotted as shaded bars and the fraction of affected progeny is indicated at the right of each graph.

[0035] FIG. 2B is a graphical representation of experiments demonstrating that animals homozygous for *rde* and *mut* alleles are resistant to RNAi targeting maternally expressed genes, *pos-1* and *par-2*. The percent embryonic lethality of F1 progeny is plotted as shaded bars and the fraction of affected progeny is indicated at the right of each graph.

[0036] FIG. 3 is a schematic representation of homozygous *rde-1* (*ne219*) and *rde-4* (*ne299*) mutant mothers receiving injections of dsRNA targeting the body muscle structural gene *unc-22*.

[0037] FIG. 4A is a schematic representation of the physical map of the *rde-1* region. *C. elegans* YAC and cosmid DNA clones that were positive for rescue are indicated by an asterisk. A representation of the expanded interval showing a minimal, 25 kb, rescuing interval defined by the overlap between cosmids T10A5 and C₂₇H₆ is shown beneath the YAC and cosmid map. Predicted genes within this sequenced interval are illustrated above and below the hatch marked line. A single, rescuing, 4.5 kb PCR fragment

containing the KO8H.107 predicted gene is shown enlarged. Exon and intron (box/line) boundaries are shown as well as the positions of *rde-1* point mutation in the predicted coding sequences.

[0038] FIG. 4B is an illustration of the predicted sequence of RDE-1 and its alignment with four related proteins. The sequences are RDE-1 (*C. elegans*; Genbank Accession No. AF180730), F48F7.1 (*C. elegans*; Genbank Accession No. Z69661), eIF2C (rabbit; Genbank Accession No. AF005355), ZWILLE (*Arabidopsis*; Genbank Accession No. AJ223508), and Sting (*Drosophila*; Genbank Accession No. AF145680). Identities with RDE-1 are shaded in black, and identities among the homologs are shaded in gray.

[0039] FIGS. 5A-5C are an illustration of the genomic sequence from cosmid KO8H10 (Genbank accession Z83113.1; SEQ ID NO:1) corresponding to the *rde-1* gene from the first nucleotide of 5' untranslated region to the polyadenylation site.

[0040] FIGS. 6A-6D are an illustration of the cDNA sequence of *rde-1* (SEQ ID NO:2), including the first 20 nucleotides constituting the 5' untranslated sequence (5'UTR) and the predicted amino acid sequence encoded by *rde-1* (RDE-1; SEQ ID NO:3). The nucleotide sequence is numbered starting with the first nucleotide of the translated region.

[0041] FIG. 7A is an illustration of the protocol for injection of a wild-type hermaphrodite with dsRNA.

[0042] FIG. 7B is an illustration of a genetic scheme demonstrating extragenic inheritance of RNAi. The fraction shown represents the number of RNAi affected F2 hermaphrodites over the total number of cross progeny scored for each genotype class. Phenotypically uncoordinated (*Unc*).

[0043] FIGS. 8A-8B are illustrations of a genetic scheme to determine if the wild-type activities of *rde-1*, *rde-2*, *rde-4*, and *mut-7* are sufficient in the injected animal for interference among the F1 self progeny (A) illustrates crosses of heterozygous hermaphrodites; (B) illustrates crosses using homozygous F1 progeny from heterozygous mothers. The fraction shown represents the number of RNAi affected animals over the total number of cross progeny scored for each genotype class.

[0044] FIG. 9A depicts experiments of a the genetic scheme to determine if the wild-type activities of *rde-1*, *rde-2*, *rde-4*, and *mut-7* are sufficient in the injected animal for interference among the F1 self progeny. The fraction shown represents the number of RNAi affected animals over the total number of cross progeny scored for each genotype class.

[0045] FIG. 9B depicts experiments designed to determine the requirements for *rde-1*, *rde-2*, *rde-4*, and *mut-7* in F2 (FIG. 10A) and F1 (FIG. 10B) interference. The fraction shown represents the number of RNAi affected animals over the total number of cross progeny scored for each genotype class.

[0046] FIGS. 10A-10B are a depiction of the cDNA sequence of a wild type *rde-4* nucleic acid sequence (SEQ ID NO:4) and the predicted RDE-4 amino acid sequence (SEQ ID NO:5) of *C. elegans*. "*" indicates ambiguous base assignment.

[0047] FIG. 11 is a depiction of regions of homology between the predicted RDE-4 amino acid sequence, X1RBPA (SEQ ID NO:6), HsPKR (SEQ ID NO:7), and a consensus sequence (SEQ ID NO:8). A predicted secondary structure for RDE-4 is also shown illustrating predicted regions of α helix and β pleated sheet.

[0048] FIG. 12 illustrates a scheme for rescue of an rde-4.

DETAILED DESCRIPTION

[0049] Mutations have been discovered that identify genes involved in dsRNA-mediated genetic interference (RNAi). RNAi pathway genes encode products involved in genetic interference and are useful for mediating or enhancing genetic interference. These genes encode mediators of double-stranded RNA-mediated interference. The mediators can be nucleic acid or protein. RNAi pathway genes are also useful for mediating specific processes, e.g., a gene that mediates dsRNA uptake by cells may be useful for transporting other RNAs into cells or for facilitating entry of agents such as drugs into cells. The methods and examples described below illustrate the identification of RNAi pathway components, the uses of RNAi pathway components, mutants, genes and their products.

Identification of an RNAi-Deficient Mutants and an RNAi Pathway Gene rde-1

[0050] RNAi pathway genes were identified using screens for *C. elegans* strains mutant for RNAi (Examples 2 and 3). The mutations were further characterized for germline and somatic effects, effects on transposon mobilization, X chromosome loss and transgene silencing, and target tissue activity (Examples 4 and 5).

[0051] The rde-1 gene was identified using YACs (yeast artificial chromosomes) and cosmids to rescue rde-1 mutants. Based on the identified sequence, a cDNA sequence was identified in a *C. elegans* cDNA library and the complete cDNA sequence determined (Example 6).

Identification of RNAi Pathway Genes Homologous to rde-1, rde-2, rde-3, and rde-4

[0052] RNAi pathway genes from *C. elegans* (such as those described herein) and from other organisms (e.g. plant, mammalian, especially human) are useful for the elucidation of the biochemical pathways involved in genetic interference and for developing the uses of RNAi pathway genes described herein.

[0053] Several approaches can be used to isolate RNAi pathway genes including two-hybrid screens, complementation of *C. elegans* mutants by expression libraries of cloned heterologous (e.g., plant, mammalian, human) cDNAs, polymerase chain reactions (PCR) primed with degenerate oligonucleotides, low stringency hybridization screens of heterologous cDNA or genomic libraries with a *C. elegans* RNAi pathway gene, and database screens for sequences homologous to an RNAi pathway gene. Hybridization is performed under stringent conditions. Alternatively, a labeled fragment can be used to screen a genomic library derived from the organism of interest, again, using appropriately stringent conditions. Such stringent conditions are well known, and will vary predictably depending on the specific organisms from which the library and the labeled sequences are derived.

[0054] Nucleic acid duplex or hybrid stability is expressed as the melting temperature or T_m , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions. If sequences are to be identified that are related and substantially identical to the probe, rather than identical, then it is useful to first establish the lowest temperature at which only homologous hybridization occurs with a particular SSC or SSPE concentration. Then assume that 1% mismatching results in 1° C. decrease in the T_m and reduce the temperature of the final wash accordingly (for example, if sequences with $\geq 95\%$ identity with the probe are sought, decrease the final wash temperature by 5° C.). Note that this assumption is very approximate, and the actual change in T_m can be between 0.5° and 1.5° C. per 1% mismatch.

[0055] As used herein, high stringency conditions include hybridizing at 68° C. in 5×SSC/5× Denhardt solution/1.0% SDS, or in 0.5 M NaHPO₄ (pH 7.2)/1 mM EDTA/7% SDS, or in 50% formamide/0.25 M NaHPO₄ (pH 7.2)/0.25 M NaCl/1 mM EDTA/7% SDS; and washing in 0.2×SSC/0.1% SDS at room temperature or at 42° C., or in 0.1×SSC/0.1% SDS at 68° C., or in 40 mM NaHPO₄ (pH 7.2)/1 mM EDTA/5% SDS at 50° C., or in 40 mM NaHPO₄ (pH 7.2) 1 mM EDTA/1% SDS at 50° C. Moderately stringent conditions include washing in 3×SSC at 42° C. The parameters of salt concentration and temperature can be varied to achieve the desired level of identity between the probe and the target nucleic acid.

[0056] For guidance regarding such conditions see, for example, Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Springs Harbor Press, N.Y.; and Ausubel et al. (eds.), 1995, Current Protocols in Molecular Biology, (John Wiley & Sons, N.Y.) at Unit 2.10.

[0057] Methods of screening for and identifying homologs of *C. elegans* RNAi genes (e.g., rde-1) are known in the art. For example, complementation of mutants, described in the Examples can be performed using nucleic acid sequences from organisms other than *C. elegans*. Methods of inhibiting expression of a target gene in a cell using dsRNA are known in the art and are exemplified in U.S. Ser. No. 09/215,257, filed Dec. 18, 1998, which is incorporated herein by reference in its entirety.

[0058] Another method of screening is to use an identified RNAi pathway gene sequence to screen a cDNA or genomic library using low stringency hybridizations. Such methods are known in the art.

[0059] PCR with degenerate oligonucleotides is another method of identifying homologs of RNAi pathway genes (e.g., human rde-1). Homologs of an RNAi pathway gene identified in other species are compared to identify specific regions with a high degree of homology (as in the sequence comparison shown in FIG. 4). These regions of high homology are selected for designing PCR primers that maximize possible base-pairing with heterologous genes. Construction of such primers involves the use of oligonucleotide mixtures that account for degeneracy in the genetic code, i.e., allow for the possible base changes in an RNAi pathway gene that does not affect the amino acid sequence of the RNAi pathway protein. Such primers may be used to amplify and clone possible RNAi pathway gene fragments from DNA isolated from another organism (e.g., mouse or human). The latter are sequenced and those encoding protein fragments

with high degrees of homology to fragments of the RNAi pathway protein are used as nucleic acid probes in subsequent screens of genomic DNA and cDNA libraries (e.g., mouse or human). Full-length genes and cDNAs having substantial homology to the previously identified RNAi pathway gene are identified in these screens.

[0060] To produce an RNAi pathway gene product (e.g., RDE-1) a sequence encoding the gene is placed in an expression vector and the gene expressed in an appropriate cell type. The gene product is isolated from such cell lines using methods known to those in the art, and used in the assays and procedures described herein. The gene product can be a complete RNAi pathway protein (e.g., RDE-1) or a fragment of such a protein.

[0061] Methods of Expressing RNAi Pathway Proteins

[0062] Full-length polypeptides and polypeptides corresponding to one or more domains of a full-length RNAi pathway protein, e.g., the RNA-binding domain of RDE-4, are also within the scope of the invention. Also within the invention are fusion proteins in which a portion (e.g., one or more domains) of an RDE-1 or RDE-4 is fused to an unrelated protein or polypeptide (i.e., a fusion partner) to create a fusion protein. The fusion partner can be a moiety selected to facilitate purification, detection, or solubilization, or to provide some other function. Fusion proteins are generally produced by expressing a hybrid gene in which a nucleotide sequence encoding all or a portion of of an RNAi pathway protein is joined in-frame to a nucleotide sequence encoding the fusion partner. Fusion partners include, but are not limited to, the constant region of an immunoglobulin (IgFc). A fusion protein in which an RNAi pathway polypeptide is fused to IgFc can be more stable and have a longer half-life in the body than the polypeptide on its own.

[0063] In general, RNAi pathway proteins (e.g., RDE-1, RDE-4) according to the invention can be produced by transformation (transfection, transduction, or infection) of a host cell with all or part of an RNAi pathway protein-encoding DNA fragment (e.g., one of the cDNAs described herein) in a suitable expression vehicle. Suitable expression vehicles include: plasmids, viral particles, and phage. For insect cells, baculovirus expression vectors are suitable. The entire expression vehicle, or a part thereof, can be integrated into the host cell genome. In some circumstances, it is desirable to employ an inducible expression vector, e.g., the LACSWITCH™ Inducible Expression System (Stratagene; LaJolla, Calif.).

[0064] Those skilled in the field of molecular biology will understand that any of a wide variety of expression systems can be used to provide the recombinant protein. The precise host cell used is not critical to the invention. The RNAi pathway protein can be produced in a prokaryotic host (e.g., *E. coli* or *B. subtilis*) or in a eukaryotic host (e.g., *Saccharomyces* or *Pichia*; mammalian cells, e.g., COS, NIH 3T3 CHO, BHK, 293, or HeLa cells; or insect cells).

[0065] Proteins and polypeptides can also be produced in plant cells. For plant cells viral expression vectors (e.g., cauliflower mosaic virus and tobacco mosaic virus) and plasmid expression vectors (e.g., Ti plasmid) are suitable. Such cells are available from a wide range of sources (e.g., the American Type Culture Collection, Rockland, Md.; also, see, e.g., Ausubel et al., *Current Protocols in Molecular*

Biology, John Wiley & Sons, New York, 1994). The methods of transformation or transfection and the choice of expression vehicle will depend on the host system selected. Transformation and transfection methods are described, e.g., in Ausubel et al., supra; expression vehicles may be chosen from those provided, e.g., in *Cloning Vectors: A Laboratory Manual* (P. H. Pouwels et al., 1985, Supp. 1987).

[0066] The host cells harboring the expression vehicle can be cultured in conventional nutrient media adapted as need for activation of a chosen gene, repression of a chosen gene, selection of transformants, or amplification of a chosen gene.

[0067] One preferred expression system is the mouse 3T3 fibroblast host cell transfected with a pMAMneo expression vector (Clontech, Palo Alto, Calif.). pMAMneo provides an RSV-LTR enhancer linked to a dexamethasone-inducible MMTV-LTR promoter, an SV40 origin of replication which allows replication in mammalian systems, a selectable neomycin gene, and SV40 splicing and polyadenylation sites. DNA encoding an RNAi pathway protein would be inserted into the pMAMneo vector in an orientation designed to allow expression. The recombinant RNAi pathway protein would be isolated as described herein. Other preferable host cells that can be used in conjunction with the pMAMneo expression vehicle include COS cells and CHO cells (ATCC Accession Nos. CRL 1650 and CCL 61, respectively).

[0068] RNAi pathway polypeptides can be produced as fusion proteins. For example, the expression vector pUR278 (Ruther et al., *EMBO J.* 2:1791, 1983), can be used to create lacZ fusion proteins. The pGEX vectors can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can be easily purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

[0069] In an insect cell expression system, *Autographa californica* nuclear polyhidrosis virus (AcNPV), which grows in *Spodoptera frugiperda* cells, is used as a vector to express foreign genes. An RNAi pathway protein coding sequence can be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter, e.g., the polyhedrin promoter. Successful insertion of a gene encoding an RNAi pathway polypeptide or protein will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat encoded by the polyhedrin gene). These recombinant viruses are then used to infect *spodoptera frugiperda* cells in which the inserted gene is expressed (see, e.g., Smith et al., *J. Virol.* 46:584, 1983; Smith, U.S. Pat. No. 4,215, 051).

[0070] In mammalian host cells, a number of viral-based expression systems can be utilized. When an adenovirus is used as an expression vector, the RNAi pathway protein nucleic acid sequence can be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene can then be inserted into the adenovirus genome by in vitro or in vivo recombination. Insertion into a non-essential region of

the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing an RNAi pathway gene product in infected hosts (see, e.g., Logan, *Proc. Natl. Acad. Sci. USA* 81:3655, 1984).

[0071] Specific initiation signals may be required for efficient translation of inserted nucleic acid sequences. These signals include the ATG initiation codon and adjacent sequences. In cases where an entire native RNAi pathway protein gene or cDNA, including its own initiation codon and adjacent sequences, is inserted into the appropriate expression vector, no additional translational control signals may be needed. In other cases, exogenous translational control signals, including, perhaps, the ATG initiation codon, must be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators (Bittner et al., *Methods in Enzymol.* 153:516, 1987).

[0072] RNAi pathway polypeptides can be expressed directly or as a fusion with a heterologous polypeptide, such as a signal sequence or other polypeptide having a specific cleavage site at the N-and/or C-terminus of the mature protein or polypeptide. Included within the scope of this invention are RNAi pathway polypeptides with a heterologous signal sequence. The heterologous signal sequence selected should be one that is recognized and processed, i.e., cleaved by a signal peptidase, by the host cell. For prokaryotic host cells a prokaryotic signal sequence is selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion a yeast invertase, alpha factor, or acid phosphatase leaders may be selected. In mammalian cells, it is generally desirable to select a mammalian signal sequences.

[0073] A host cell may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in a specific, desired fashion. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product can be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, W138, and in particular, choroid plexus cell lines.

[0074] Alternatively, an RNAi pathway protein can be produced by a stably-transfected mammalian cell line. A number of vectors suitable for stable transfection of mammalian cells are available to the public, see e.g., Pouwels et al. (supra); methods for constructing such cell lines are also publicly available, e.g., in Ausubel et al. (supra). In one example, cDNA encoding an RNAi pathway protein (e.g., RDE-1 or RDE-4) is cloned into an expression vector that

includes the dihydrofolate reductase (DHFR) gene. Integration of the plasmid and, therefore, the RNAi pathway protein-encoding gene into the host cell chromosome is selected for by including 0.01-300 μ M methotrexate in the cell culture medium (as described in Ausubel et al., supra). This dominant selection can be accomplished in most cell types.

[0075] Recombinant protein expression can be increased by DHFR-mediated amplification of the transfected gene. Methods for selecting cell lines bearing gene amplifications are described in Ausubel et al. (supra); such methods generally involve extended culture in medium containing gradually increasing levels of methotrexate. DHFR-containing expression vectors commonly used for this purpose include pCVSEII-DHFR and pAdD26SV(A) (described in Ausubel et al., supra). Any of the host cells described above or, preferably, a DHFR-deficient CHO cell line (e.g., CHO DHFR⁻ cells, ATCC Accession No. CRL 9096) are among the host cells preferred for DHFR selection of a stably-transfected cell line or DHFR-mediated gene amplification.

[0076] A number of other selection systems can be used, including but not limited to the herpes simplex virus thymidine kinase, hypoxanthine-guanine phosphoribosyl-transferase, and adenine phosphoribosyltransferase genes can be employed in tk, hgp^rt, or apr^t cells, respectively. In addition, gpt, which confers resistance to mycophenolic acid (Mulligan et al., *Proc. Natl. Acad. Sci. USA*, 78:2072, 1981); neo, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., *J. Mol. Biol.*, 150:1, 1981); and hyg^r, which confers resistance to hygromycin (Santerre et al., *Gene*, 30:147, 1981), can be used.

[0077] Alternatively, any fusion protein can be readily purified by utilizing an antibody specific for the fusion protein being expressed. For example, a system described in Janknecht et al., *Proc. Natl. Acad. Sci. USA*, 88:8972 (1981), allows for the ready purification of non-denatured fusion proteins expressed in human cell lines. In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting of six histidine residues. Extracts from cells infected with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns, and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

[0078] Alternatively, an RNAi pathway protein or a portion thereof, can be fused to an immunoglobulin Fc domain. Such a fusion protein can be readily purified using a protein A column.

[0079] Antibodies that Recognize RNAi Pathway Proteins

[0080] Techniques for generating both monoclonal and polyclonal antibodies specific for a particular protein are well known. The invention also includes humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, and molecules produced using a Fab expression library.

[0081] Antibodies can be raised against a short peptide epitope of an RNAi pathway gene (e.g., rde-1), an epitope linked to a known immunogen to enhance immunogenicity, a long fragment of an RNAi pathway gene, or the intact protein. Such antibodies are useful for e.g., localizing RNAi pathway polypeptides in tissue sections or fractionated cell

preparations, determining whether an RNAi pathway gene is expressed (e.g., after transfection with an RNAi pathway gene), and evaluating the expression of an RNAi pathway gene in disorders (e.g., genetic conditions) where the RNAi pathway may be affected.

[0082] An isolated RNAi pathway protein (e.g., RDE-1), or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind to an RNAi pathway protein using standard techniques for polyclonal and monoclonal antibody preparation. The RNAi pathway immunogen can also be a mutant RNAi pathway protein or a fragment of a mutant RNAi pathway protein. A full-length RNAi pathway protein can be used or, alternatively, antigenic peptide fragments of RNAi pathway protein can be used as immunogens. The antigenic peptide of an RNAi pathway protein comprises at least 8 (preferably 10, 15, 20, or 30) amino acid residues. In the case of RDE-1, these residues are drawn from the amino acid sequence shown in SEQ ID NO:3 and encompass an epitope such that an antibody raised against the peptide forms a specific immune complex with RDE-1. Preferred epitopes encompassed by the antigenic peptide are regions of the protein that are located on the surface of the protein, e.g., hydrophilic regions.

[0083] An RNAi pathway protein immunogen typically is used to prepare antibodies by immunizing a suitable subject (e.g., rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed RNAi pathway protein or a chemically synthesized RNAi polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic RNAi pathway protein preparation induces a polyclonal anti-RNAi pathway protein antibody response.

[0084] Polyclonal antibodies that recognize an RNAi pathway protein ("RNAi pathway antibodies") can be prepared as described above by immunizing a suitable subject with an RNAi pathway protein immunogen. The RNAi pathway antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme-linked immunosorbent assay (ELISA) using immobilized RNAi pathway protein from which the immunogen was derived. If desired, the antibody molecules directed against the RNAi pathway protein can be isolated from the mammal (e.g., from the blood) and further purified by well-known techniques, such as protein A chromatography to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the RNAi pathway antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497, the human B cell hybridoma technique (Kozbor et al. (1983) *Immunol. Today* 4:72), the EBV-hybridoma technique (Cole et al. (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing hybridomas is well known (see generally *Current Protocols in Immunology* (1994) Coligan et al. (eds.) John Wiley & Sons, Inc., New York, N.Y.). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with an RNAi pathway immu-

nogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds to the RNAi pathway protein.

[0085] Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating a monoclonal antibody against an RNAi pathway protein (see, e.g., *Current Protocols in Immunology*, supra; Galfre et al., 1977, *Nature* 266:55052; R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, N.Y., 1980; and Lerner, 1981, *Yale J. Biol. Med.*, 54:387-402. Moreover, one in the art will appreciate that there are many variations of such methods which also would be useful. Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind to the RNAi pathway protein, e.g., using a standard ELISA assay.

[0086] Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal RNAi pathway antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with an RNAi pathway protein to thereby isolate immunoglobulin library members that bind to the RNAi pathway protein. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene SurfZAP™ Phage Display Kit, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening antibody display library can be found in, for example, U.S. Pat. No. 5,223,409; PCT Publication No. WO 92/18619; PCT Publication No. WO 91/17271; PCT Publication No. WO 92/20791; PCT Publication No. WO 92/15679; PCT Publication No. WO 93/01288; PCT Publication No. WO 92/01047; PCT Publication No. WO 92/09690; PCT Publication No. WO 90/02809; Fuchs et al., 1991, *Bio/Technology* 9:1370-1372; Hay et al., 1992, *Hum. Antibod. Hybridomas* 3:81-85; Huse et al., 1989, *Science* 246:1275-1281; Griffiths et al., 1993, *EMBO J.* 12:725-734.

[0087] Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851, 1984; Neuberger et al., *Nature*, 312:604, 1984; Takeda et al., *Nature*, 314:452, 1984) can be used to splice the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity. A chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

[0088] Alternatively, techniques described for the production of single chain antibodies (U.S. Pat. No. 4,946,778; and U.S. Pat. Nos. 4,946,778 and 4,704,692) can be adapted to produce single chain antibodies against an RNAi pathway protein or polypeptide. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide.

[0089] Antibody fragments that recognize and bind to specific epitopes can be generated by known techniques. For

example, such fragments can include but are not limited to F(ab')₂ fragments, which can be produced by pepsin digestion of the antibody molecule, and Fab fragments, which can be generated by reducing the disulfide bridges of F(ab')₂ fragments. Alternatively, Fab expression libraries can be constructed (Huse et al., *Science*, 246:1275, 1989) to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity.

[0090] Identification of RNAi Pathway Components

[0091] RNAi pathway components can be identified in *C. elegans* and other animals (e.g., a mammal) using the methods described in the Examples below. Pathway components can also be identified using methods known in the art and the information provided herein. Such components include those involved in protein:protein and protein:RNA interactions. Specifically, RDE-1 can be used to identify additional proteins and RNA molecules that bind to the RDE-1 protein and so facilitate genetic interference.

[0092] The RNAi pathway mutant strains described herein (e.g., rde-1, rde-2, rde-3, rde-4, and rde-5; also mut-2 and mut-7) can be used in genetic screens to identify additional RNAi pathway components. For example, a strain deficient for rde-1 activity can be mutagenized and screened for the recovery of genetic interference. This type of screen can identify allele-specific suppressors in other genes or second site mutations within the rde-1 gene that restore its activity. The resulting strains may define new genes that activate RNAi to overcome or bypass the rde-1 defect. The mutations identified by these methods can be used to identify their corresponding gene sequences.

[0093] Two-hybrid screens can also be used to identify proteins that bind to RNAi pathway proteins such as RDE-1. Genes encoding proteins that interact with RDE-1 or human homologs of the *C. elegans* RDE-1, are identified using the two-hybrid method (Fields and Song, 1989, *Nature* 340:245-246; Chien et al., 1991, *Proc. Natl. Acad. Sci. USA* 88:9578-9582; Fields and Stemglanz, 1994, *Trends Genet.* 10:286-292; Bartel and Fields, 1995, *Methods Enzymol.* 254:241-263). DNA encoding the RDE-1 protein is cloned and expressed from plasmids harboring GAL4 or lexA DNA-binding domains and co-transformed into cells harboring lacZ and HIS3 reporter constructs along with libraries of cDNAs that have been cloned into plasmids harboring the GAL4 activation domain. Libraries used for such co-transformation include those made from *C. elegans* or a vertebrate embryonic cell.

[0094] Mechanisms of Action of RNAi Pathway Components

[0095] Specific cellular functions associated with the RNAi pathway include the specific targeting of a nucleic acid by a dsRNA, uptake of dsRNA, transport of dsRNA, amplification of the dsRNA signal, and genetic interference. The mechanism of interference may involve translation inhibition, or interference with RNA processing. In addition, direct effects on the corresponding gene may contribute to interference. These mechanisms can be identified investigated using the methods described herein and methods known in the art.

[0096] Methods of Screening for Molecules that Inhibit the RNAi Pathway

[0097] The following assays are designed to identify compounds that are effective inhibitors of the RNAi pathway. Such inhibitors may act by, but are not limited to, binding to an RDE-1 polypeptide (e.g., from *C. elegans*, mouse, or human), binding to intracellular proteins that bind to an RNAi pathway component, compounds that interfere with the interaction between RNAi pathway components including between an RNAi pathway component and a dsRNA, and compounds that modulate the activity or expression of an RNAi pathway gene such as rde-1. An inhibitor of the RNAi pathway can also be used to promote expression of a transgene.

[0098] Assays can also be used to identify molecules that bind to RNAi pathway gene regulatory sequences (e.g., promoter sequences), thus modulating gene expression. See, e.g., Platt, 1994, *J. Biol. Chem.* 269:28558-28562, incorporated herein by reference in its entirety.

[0099] The compounds which may be screened by the methods described herein include, but are not limited to, peptides and other organic compounds (e.g., peptidomimetics) that bind to an RNAi pathway protein (e.g., that bind to an RDE-1), or inhibit its activity in any way.

[0100] Such compounds may include, but are not limited to, peptides; for example, soluble peptides, including but not limited to members of random peptide libraries; (see, e.g., Lam et al., 1991, *Nature* 354:82-94; Houghten et al., 1991, *Nature* 354:84-86), and combinatorial chemistry-derived molecular libraries made of D-and/or L-amino acids, phosphopeptides (including, but not limited to, members of random or partially degenerate, directed phosphopeptide libraries; see e.g., Songyang et al., 1993, *Cell* 72:767-778), and small organic or inorganic molecules.

[0101] Organic molecules are screened to identify candidate molecules that affect expression of an RNAi pathway gene (e.g., rde-1), e.g., by interacting with the regulatory region or transcription factors of a gene. Compounds are also screened to identify those that affect the activity of such proteins, (e.g., by inhibiting rde-1 activity) or the activity of a molecule involved in the regulation of, for example, rde-1.

[0102] Computer modeling or searching technologies are used to identify compounds, or identify modifications of compounds that modulate the expression or activity of an RNAi pathway protein. For example, compounds likely to interact with the active site of a protein (e.g., RDE-1) are identified. The active site of an RNAi pathway protein can be identified using methods known in the art including, for example, analysis of the amino acid sequence of a molecule, from a study of complexes of an RNAi pathway, with its native ligand (e.g., a dsRNA). Chemical or X-ray crystallographic methods can be used to identify the active site of an RNAi pathway protein by the location of a bound ligand such as a dsRNA.

[0103] The three-dimensional structure of the active site is determined. This can be done using known methods, including X-ray crystallography which may be used to determine a complete molecular structure. Solid or liquid phase NMR can be used to determine certain intra-molecular distances. Other methods of structural analysis can be used to determine partial or complete geometrical structures. Geometric

structure can be determined with an RNAi pathway protein bound to a natural or artificial ligand which may provide a more accurate active site structure determination.

[0104] Computer-based numerical modeling can also be used to predict protein structure (especially of the active site), or be used to complete an incomplete or insufficiently accurate structure. Modeling methods that may be used are, for example, parameterized models specific to particular biopolymers such as proteins or nucleic acids, molecular dynamics models based on computing molecular motions, statistical mechanics models based on thermal ensembles, or combined models. For most types of models, standard molecular force fields, representing the forces between constituent atoms and groups are necessary, and can be selected for the model from among the force fields known in physical chemistry. Information on incomplete or less accurate structures determined as above can be incorporated as constraints on the structures computed by these modeling methods.

[0105] Having determined the structure of the active site of an RNAi pathway protein (e.g., RDE-1), either experimentally, by modeling, or by a combination of methods, candidate modulating compounds can be identified by searching databases containing compounds along with information on their molecular structure. The compounds identified in such a search are those that have structures that match the active site structure, fit into the active site, or interact with groups defining the active site. The compounds identified by the search are potential RNAi pathway modulating compounds.

[0106] These methods may also be used to identify improved modulating compounds from an already known modulating compound or ligand. The structure of the known compound is modified and effects are determined using experimental and computer modeling methods as described above. The altered structure may be compared to the active site structure of an RNAi pathway protein (e.g., an RDE-1) to determine or predict how a particular modification to the ligand or modulating compound will affect its interaction with that protein. Systematic variations in composition, such as by varying side groups, can be evaluated to obtain modified modulating compounds or ligands of preferred specificity or activity.

[0107] Other experimental and computer modeling methods useful to identify modulating compounds based on identification of the active sites of an RNAi pathway protein and related transduction and transcription factors will be apparent to those of skill in the art.

[0108] Examples of molecular modeling systems are the QUANTA programs, e.g., CHARMM, MCSS/HOOK, and X-LIGAND, (Molecular Simulations, Inc., San Diego, Calif.). QUANTA analyzes the construction, graphic modeling, and analysis of molecular structure. CHARMM analyzes energy minimization and molecular dynamics functions. MCSS/HOOK characterizes the ability of an active site to bind a ligand using energetics calculated via CHARMM. X-LIGAND fits ligand molecules to electron density of protein-ligand complexes. It also allows interactive construction, modification, visualization, and analysis of the behavior of molecules with each other.

[0109] Articles reviewing computer modeling of compounds interacting with specific protein can provide addi-

tional guidance. For example, see Rotivinen et al., 1988, *Acta Pharmaceutica Fennica* 97:159-166; Ripka, *New Scientist* June 16, 1988 pp. 54-57; McKinaly and Rossmann, 1989, *Ann. Rev. Pharmacol. Toxicol.* 29:111-122; Perry and Davies. *OSAR Quantitative Structure-Activity Relationships in Drug Design* pp. 189-193 (Alan R. Liss, Inc., 1989); Lewis and Dean, 1989, *Proc. R. Soc. Lond.* 236:125-140, 141-152; and, regarding a model receptor for nucleic acid components, Askew et al., *Am. J. Chem. Soc.* 111: 1082-1090. Computer programs designed to screen and depict chemicals are available from companies such as MSI (supra), Allelix, Inc. (Mississauga, Ontario, Canada), and Hypercube, Inc. (Gainesville, Fla.).

[0110] These applications are largely designed for drugs specific to particular proteins; however, they can be adapted to the design of drugs specific to identified regions of DNA or RNA. Chemical libraries that can be used in the protocols described herein include those available, e.g., from ArQule, Inc. (Medford, Mass.) and Oncogene Science, Inc. (Uniondale, N.Y.).

[0111] In addition to designing and generating compounds that alter binding, as described above, libraries of known compounds, including natural products, synthetic chemicals, and biologically active materials including peptides, can be screened for compounds that are inhibitors or activators of the RNAi pathway components identified herein.

[0112] Compounds identified by methods described above can be used, for example, for elaborating the biological function of RNAi pathway gene products (e.g., an RDE-1), and to treat genetic disorders involving an RNAi pathway protein. Assays for testing the effectiveness of compounds such as those described herein are further described below.

[0113] In Vitro Screening Assays for Compounds that Bind to RNAi Pathway Proteins and Genes

[0114] In vitro systems can be used to identify compounds that interact with (e.g., bind to) RNAi pathway proteins or genes encoding those proteins (e.g., rde-1 and its protein product). Such compounds are useful, for example, for modulating the activity of these entities, elaborating their biochemistry, treating disorders in which a decrease or increase in dsRNA mediated genetic interference is desired. Such compounds may also be useful to treat diseases in animals, especially humans, involving nematodes, e.g., trichinosis, trichuriasis, and toxocariasis. Compounds such as those described herein may also be useful to treat plant diseases caused by nematodes. These compounds can be used in screens for compounds that disrupt normal function, or may themselves disrupt normal function.

[0115] Assays to identify compounds that bind to RNAi pathway proteins involve preparation of a reaction mixture of the protein and the test compound under conditions sufficient to allow the two components to interact and bind thus forming a complex which can be removed and/or detected.

[0116] Screening assays can be performed using a number of methods. For example, an RNAi pathway protein from an organism (e.g., RDE-1), peptide, or fusion protein can be immobilized onto a solid phase, reacted with the test compound, and complexes detected by direct or indirect labeling of the test compound. Alternatively, the test compound can be immobilized, reacted with the RNAi pathway molecule,

and the complexes detected. Microtiter plates may be used as the solid phase and the immobilized component anchored by covalent or noncovalent interactions. Non-covalent attachment may be achieved by coating the solid phase with a solution containing the molecule and drying. Alternatively, an antibody, for example, one specific for an RNAi pathway protein such as RDE-1 is used to anchor the molecule to the solid surface. Such surfaces may be prepared in advance of use, and stored.

[0117] In these screening assays, the non-immobilized component is added to the coated surface containing the immobilized component under conditions sufficient to permit interaction between the two components. The unreacted components are then removed (e.g., by washing) under conditions such that any complexes formed will remain immobilized on the solid phase. The detection of the complexes may be accomplished by a number of methods known to those in the art. For example, the nonimmobilized component of the assay may be pre-labeled with a radioactive or enzymatic entity and detected using appropriate means. If the non-immobilized entity was not pre-labeled, an indirect method is used. For example, if the non-immobilized entity is an RDE-1, an antibody against the RDE-1 is used to detect the bound molecule, and a secondary, labeled antibody used to detect the entire complex.

[0118] Alternatively, a reaction can be conducted in a liquid phase, the reaction products separated from unreacted components, and complexes detected (e.g., using an immobilized antibody specific for an RNAi pathway protein).

[0119] Cell-based assays can be used to identify compounds that interact with RNAi pathway proteins. Cell lines that naturally express such proteins or have been genetically engineered to express such proteins (e.g., by transfection or transduction of an *rde-1* DNA) can be used. For example, test compounds can be administered to cell cultures and the amount of mRNA derived from an RNAi pathway gene analyzed, e.g., by Northern analysis. An increase in the amount of RNA transcribed from such a gene compared to control cultures that did not contain the test compound indicates that the test compound is an inhibitor of the RNAi pathway. Similarly, the amount of a polypeptide encoded by an RNAi pathway gene, or the activity of such a polypeptide, can be analyzed in the presence and absence of a test compound. An increase in the amount or activity of the polypeptide indicates that the test compound is an inhibitor of the RNAi pathway.

[0120] Ectopic Expression of an RNAi Pathway Gene

[0121] Ectopic expression (i.e., expression of an RNAi pathway gene in a cell where it is not normally expressed or at a time when it is not normally expressed) of a mutant RNAi pathway gene (i.e., an RNAi pathway gene that suppresses genetic interference) can be used to block or reduce endogenous interference in a host organism. This is useful, e.g., for enhancing transgene expression in those cases where the RNAi pathway is interfering with expression of a transgene. Another method of accomplishing this is to knockout or down regulate an RNAi pathway gene using methods known in the art. These methods are useful in both plants and animals (e.g., in an invertebrate such as a nematode, a mouse, or a human).

[0122] Ectopic expression of an RNAi pathway gene, e.g., *rde-1* or *rde-4* can also be used to activate the RNAi

pathway. In some cases, targeting can be used to activate the pathway in specific cell types, e.g., tumor cells. For example, a non-viral RNAi pathway gene construct can be targeted *in vivo* to specific tissues or organs, e.g., the liver or muscle, in patients. Examples of delivery systems for targeting such constructs include receptor mediated endocytosis, liposome encapsulation (described below), or direct insertion of non-viral expression vectors.

[0123] An example of one such method is liposome encapsulation of nucleic acid. Successful *in vivo* gene transfer has been achieved with the injection of DNA, e.g., as a linear construct or a circular plasmid, encapsulated in liposomes (Ledley, Human Gene Therapy 6:1129-1144 (1995) and Farhood, et al., Ann. NY Acad. Sci. 716:23-35 (1994)). A number of cationic liposome amphiphiles are being developed (Ledley, Human Gene Therapy 6:1129-1144 (1995); Farhood, et al., Ann. NY Acad. Sci., 716:23-35 (1994) that can be used for this purpose.

[0124] Targeted gene transfer has been shown to occur using such methods. For example, intratracheal administration of cationic lipid-DNA complexes was shown to effect gene transfer and expression in the epithelial cells lining the bronchus (Brigham, et al., Am. J. Respir. Cell Mol. Biol. 8:209-213 (1993); and Canonico, et al., Am. J. Respir. Cell Mol. Biol. 10:24-29 (1994)). Expression in pulmonary tissues and the endothelium was reported after intravenous injection of the complexes (Brigham, et al., Am. J. Respir. Cell Mol. Biol. 8:209-213 (1993); Zhu, et al., Science, 261:209-211 (1993); Stewart, et al., Human Gene Therapy 3:267-275 (1992); Nabel, et al., Human Gene Therapy 3:649-656 (1992); and Canonico, et al., J. Appl. Physiol. 77:415-419 (1994)). An expression cassette for an RNAi pathway sequence in linear, plasmid or viral DNA forms can be condensed through ionic interactions with the cationic lipid to form a particulate complex for *in vivo* delivery (Stewart, et al., Human Gene Therapy 3:267-275 (1992)).

[0125] Other liposome formulations, for example, proteo-liposomes which contain viral envelope receptor proteins, i.e., virosomes, have been found to effectively deliver genes into hepatocytes and kidney cells after direct injection (Nicolau, et al., Proc. Natl. Acad. Sci. USA 80:1068-1072 (1993); Kaneda, et al., Science 243:375-378 (1989); Mannino, et al., Biotechniques 6:682 (1988); and Tomita, et al., Biochem. Biophys. Res. Comm. 186:129-134 (1992)).

[0126] Direct injection can also be used to administer an RNAi pathway nucleic acid sequence in a DNA expression vectors, e.g., into the muscle or liver, either as a solution or as a calcium phosphate precipitate (Wolff, et al., Science 247:1465-1468 (1990); Ascadi, et al., The New Biologist 3:71-81 (1991); and Benvenisty, et al., Proc. Natl. Acad. Sci. USA 83:9551-9555 (1986).

[0127] Preparation of RNAi Agents

[0128] RNAi pathway components can be used to prepare RNAi agents. Such agents are dsRNAs that have been treated with RNAi pathway components rendering the treated dsRNA capable of activity in the RNAi pathway and can be used as sequence-specific interfering agents useful for targeted genetic interference. Specifically, treating a dsRNA with an RDE-1 and RDE-4 is useful for making an RNAi agent. An RNAi agent can be produced by preincubating a dsRNA *in vitro* in the presence of RDE-1 and RDE-4.

[0129] Another method of preparing an RNAi agent is to activate the RNAi pathway in a target cell (i.e., a cell in which it is desirable to activate the RNAi pathway such as a tumor cell) by transgenesis of an rde-1 coding sequence and an rde-4 coding sequence into the target cell.

[0130] RNAi pathway polypeptides can be modified, e.g., to enhance their stability or cellular uptake, by attaching lipophilic or other helper groups to the polypeptide, by the formation of chimeras with proteins or other moieties that are taken up by cells, or by the use of liposomes or other techniques of drug delivery known in the art.

[0131] In *C. elegans*, RNAi agents appear to spread from cell to cell, thus, active RNAi agents can diffuse or be actively transported from conditioned media or serum directly into target cells. Alternatively, RNAi agents can be injected into an organism or cell. They may also be incorporated into a cell using liposomes or other such methods known in the art.

[0132] Such methods are useful for stimulating the RNAi pathway in *C. elegans* cells, and in heterologous cells including plants and vertebrate cells. Such methods are useful in mammalian, e.g., human cells.

[0133] Enhanced Delivery of a Cargo Compound

[0134] RNAi pathway components that mediate the transport of dsRNA into cells and tissues can be used to promote the entry of dsRNA into cells and tissues, including dsRNA that is linked to another compound. The method is accomplished by linking dsRNA to a cargo compound (e.g., a drug or DNA molecule), e.g., by a covalent bond. The endogenous RNAi pathway gene expressing dsRNA transport function is activated using methods known in the art. Alternatively, other methods can be used such as transfecting the target cell with the gene that affects transport thus permitting the cell or tissue to take up the dsDNA.

EXAMPLES

[0135] The invention is further described in the Examples below which describe methods of identifying mutations in the RNAi pathway and methods of identifying genes encoding components of the RNAi pathway.

EXAMPLE 1

Strains and Alleles

[0136] The Bristol strain N2 was used as standard wild-type strain. The marker mutations and deficiencies used are listed by chromosomes as follows: LGI: dpy-14(e188), unc-13(e51); LGIII: dpy-17(e164), unc-32(e189); LGV: dpy-11(e224), unc-42(e270), daf-11(m87), eDf1, mDf3, nDf31, sDf29, sDf35, unc-76(e911). The *C. elegans* strain DP13 was used to generate hybrids for STS linkage-mapping (Williams et al., 1992, *Genetics* 131:609-624).

[0137] Sensitivity to RNAi was tested in the following strains. MT3126: mut-2(r459) (obtained from John Collins, Department of Biochemistry & Molecular Biology, University of New Hampshire, Durham, N. H.); dpy-19(n1347), TW410: mut-2([459] sem-4(n1378), NL917: mut-7(pk204), SS552: mes-2(bn76) rol-1(e91)/mnC1 (obtained from S. Strome, Biology Dept., Indiana University), SS449: mes-3(bn88) dpy-5(e61) (from S. Strome, supra); hDp20, SS268:

dpy-11(e224) mes-4(bn23) unc-76(e91)/nT1, SS360: mes-6(bn66) dpy-20(e1282)/nT1, CB879: him-1 (e879). A non-Unc mut-6 strain used was derived from RW7096: mut-6(st702) unc-22(st192::Tc1), due to the loss of Tc1 insertion in unc-22.

[0138] Homozygous mutants of mut-6, mes-2, 3, 4, 6 and him-1 showed sensitivity to RNAi by injection of pos-1 dsRNA. The dose of injected RNA was about 0.7 mg/ml. This dose lies within the range where reduced concentration leads to reduced interference effects. The results of the injection of pos-1 dsRNA into these mutants (dead embryos/F1 progeny) were as follows: mut-6: 422/437, mes-2: 781/787, mes-3: 462/474, mes-4: 810/814, mes-6: 900/1,002, him-1: 241/248, N2 (control): 365/393.

[0139] To test mutator activity, a mutant that was caused by Tc4 transposon insertion was used; TR1175: unc-22(r765::Tc4). Strains TW410 and TR1175 were gifts from Q. Boese and J. Collins (Department of Biochemistry & Molecular Biology, University of New Hampshire, Durham, N.H.).

EXAMPLE 2

RNA Interference Assay

[0140] Genetic interference using RNAi administered by microinjection was performed as described in Fire et al., 1998, supra and Rocheleau et al., 1997, *Cell* 90:707-716. pos-1 cDNA clone yk61h1, par-2 cDNA clone yk96h7, sqt-3 cDNA clone yk75f2 were used to prepare dsRNA in vitro. These cDNA clones were obtained from the *C. elegans* cDNA project (Y. Kohara, Gene Network Lab, National Institute of Genetics, Mishima 411, Japan).

[0141] Genetic interference using RNAi administered by feeding was performed as described in Timmons and Fire, 1998, *Nature* 395:854. pos-1 cDNA was cloned into a plasmid that contains two T7 promoter sequences arranged in head-to-head configuration. The plasmid was transformed into an *E. coli* strain, BL21 (DE3), and the transformed bacteria were seeded on NGM (nematode growth medium) plates containing 60 µg/ml ampicillin and 80 µg/ml IPTG. The bacteria were grown overnight at room temperature to induce pos-1 dsRNA. Seeded plates (BL21(DE3)[dsRNA] plates) stored at 4° C. remained effective for inducing interference for up to two weeks. To test RNAi sensitivity, *C. elegans* larvae were transferred onto BL21 (DE3) [dsRNA] plates and embryonic lethality was assayed in the next generation.

[0142] Transgenic lines expressing interfering RNA for unc-22 were engineered using a mixture of three plasmids: pPD[L4218] (unc-22 antisense segment, driven by myo-3 promoter); pPD[L4218] (corresponding unc-22 sense segment, driven by myo-3 promoter); pRF4 (semidominant transformation marker). DNA concentrations in the injected mixture were 100 µg/ml each. Injections were as described (Mello et al., 1991, *EMBO J.* 10:3959; Mello and Fire, 1995, *Methods in Cell Biol.* 48:451-482).

EXAMPLE 3

Identification of RNAi-Deficient Mutants

[0143] A method of screening for mutants defective in the RNAi pathway was devised that would permit the large-

scale application of dsRNA to mutagenized populations. Feeding worms *E. coli* which express a dsRNA, or simply soaking worms in dsRNA solution, are both sufficient to induce interference in *C. elegans* (Timmons and Fire, 1998, supra; Tabara et al., 1998, *Science* 282:430-431). To carry out a selection, the feeding method was optimized to deliver interfering RNA for an essential gene, *pos-1*. *C. elegans* hermaphrodites that ingest bacteria expressing dsRNA corresponding to a segment of *pos-1* are themselves unaffected but produce dead embryos with the distinctive *pos-1* embryonic lethal phenotype.

[0144] To identify strains defective in the RNAi pathway, wild-type animals were mutagenized, backcrossed, and the F2 generation examined for rare individuals that were able to produce complete broods of viable progeny. Chemical mutagenesis was used to generate the mutations as well as spontaneous mutations arising in the mut-6 strain in which Tc1 transposons are activated (Mori et al., 1988, *Genetics* 120:397-407). To facilitate screens for mutations, an egg laying starting strain was used. In the absence of egg laying, the F3 progeny remained trapped within the mother's cuticle. Candidate mutants had internally hatched broods of viable embryos and were thus easily distinguished from the background population of individuals filled primarily with dead embryos (FIG. 1A). Candidates were then re-tested for resistance to injected dsRNA.

[0145] The genetic screen used to isolate RNAi pathway mutants was similar to one designed by James R. Preiss for the identification of maternal effect mutants (Kemphues et al., 1988, *Cell* 52:311-320). An Eg1 strain, *lin-2(e)309* was mutagenized with EMS and the F2 generation was cultured on a bacterial lawn expressing *pos-1* dsRNA. Mutagenized populations were then screened for rare individuals that were able to produce complete broods of viable progeny forming a distinctive "bag of worms" phenotype. To make sure that the animals were truly resistant to RNAi, candidate strains were next assayed for resistance to RNAi by injection. Independent EMS induced alleles of *rde-1* were found in two separate pools of mutagenized animals at a frequency of approximately one allele in 2,000 to 4,000 haploid genomes.

[0146] In addition, a search was made for spontaneous mutants using a mut-6 strain in which Tc1 transposons are activated (Mori et al., 1988). 100,000 mut-6; *lin-2* animals (Mello et al., 1994) were cultured on bacteria expressing *pos-1* dsRNA. After one generation of growth, surviving animals were transferred again to plates with bacteria expressing the dsRNA and screened for resistant mutants. Three resulting strains were genetically mapped. One of these strains (ne300) mapped to LGV and failed to complement *rde-1*(ne219). Two strains ne299 and ne301 mapped to LGIII and define the *rde-4* complementation group. Because the screen was clonal in nature and involved rounds of enrichment it is possible that both *rde-4* strains are related.

[0147] Seven mutant strains were selected for genetic mapping. These seven mutants defined four complementation groups; *rde-1*, with three alleles, *rde-4*, with two alleles, and *rde-2*, and *rde-3*, with one allele each (FIG. 1B).

[0148] To map the RNAi defective mutations, the RNAi resistant phenotype was assayed either by feeding bacteria expressing *pos-1* dsRNA or by injection of a dsRNA mixture of *pos-1* and *unc-22*. The same assays were used for comple-

mentation tests. In vivo expression of *unc-22* dsRNA was also used for mapping of *rde-1*. Mapping with visible marker mutations was performed as described in Brenner (1974, *Genetics*, 77:71-94) and mapping with STS marker was performed as described in Williams et al. (1992, supra).

[0149] ne219, ne297 and ne300 failed to complement each other, defining the *rde-1* locus. *rde-1* mutations mapped near *unc-42* V. Three factor mapping was used to locate *rde-1* (ne300) one eighth of the distance from *unc-42* in the *unc-42/daf-11* interval (3/24 Unc-non-Daf recombinants analyzed). The *rde-1* (ne300) allele complemented the chromosomal deficiency *sDf29* and failed to complement *eDf1*, *mDJ3*, *nDf31* and *sDf35*. *rde-2*(ne221) and *rde-3*(ne298) mapped near *unc-13* I. *rde-2* complemented *rde-3*. *rde-4*(ne299) and (ne301) mapped near *unc-69* III and failed to complement each other. ne299 complemented mut-7(pk204).

[0150] The *rde-1* (+) activity is sufficient maternally or zygotically. To test the maternal sufficiency, animals heterozygous for *rde-1* (ne219) were injected with dsRNA targeting the zygotic gene, *sqt-3*, and self progeny were assayed for the Sqt phenotype. 100% of the self progeny including *rde-1* homozygous progeny were found to exhibit the Sqt phenotype. Thus, maternally provided *rde-1*(+) activity is sufficient to mediate interference with a zygotic target gene. Zygotic sufficiency was assayed by injecting homozygous *rde-1* mothers with dsRNA targeting the zygotic *unc-22* gene (FIG. 3). Injected animals were allowed to produce self-progeny or instead were mated after 12 hours to wild-type males, to produce heterozygous *rde-1*/+ cross-progeny. Each class of progeny was scored for the *unc-22* twitching phenotype as indicated by the fraction shown in FIG. 3 (Unc progeny/total progeny). The injected animals were then mated with wild-type males. Self progeny from homozygous injected mothers were unaffected, however, 68% of the cross progeny were Unc. This result indicates that zygotically provided *rde-1*(+) activity is also sufficient. However both maternal and zygotic *rde-1* (+) activity contribute to zygotic interference as 100% of progeny from wild-type injected mothers exhibit *unc-22* interference (606/606). Thus, *rde-1*(+) and *rde-4*(+) activities are not needed for dsRNA uptake, transport or stability.

[0151] RNAi sensitivity of several existing *C. elegans* mutants was also examined. Most of these mutant strains were fully sensitive to RNAi. However, RNAi resistance was identified in two strains that had previously been shown to exhibit elevated levels of transposon mobilization (mutator strains): mut-2 (described in Collins et al., 1987, *Nature* 328:726-728) and mut-7 (described in Ketting et al., *Cell*, in press for release on Oct. 15, 1999). Another mutator strain, mut-6(st702), was fully sensitive to RNAi. Since mutator strains continually accumulate mutations, the resistance of mut-2 and mut-7 may have been due to the presence of secondary mutations. To test this possibility we examined the genetic linkage between the mutator and RNAi resistance phenotypes of mut-2 and mut-7. We found that independently outcrossed mut-2(r459) mutator strains TW410 and MT3126 both showed resistance to RNAi. We mapped the RNAi resistance phenotype of mut-7(pk204) to the

center of linkage group III (**FIG. 1B**), the position that had been defined for the mutator activity of mut-7(pk204) by Ketting et al. (supra). Together, these observations indicate that the RNAi resistance phenotypes of the mut-2 and mut-7 strains are genetically linked to their mutator activities. Animals heterozygous for the rde and mut alleles were generated by crossing wild-type males with Unc-Rde or Unc-Mut hermaphrodites. The rde and mut mutations appeared to be simple recessive mutations with the exception of mut-2([459]), which appeared to be weakly dominant (**FIG. 2A**).

[0152] These data demonstrate that some genes are non-essential (e.g., rde-1 and rde-4).

[0153] This method can be used to identify additional mutations in RNAi pathway genes.

EXAMPLE 4

Identification of Properties of RNAi-Deficient Mutants

[0154] Effects of rde Mutations in Germline and Somatic Tissues

[0155] Microinjection was used to assay the sensitivity of each rde strain to several distinct dsRNA species. The pos-1 and par-2 genes are expressed in the maternal germline and are required for proper embryonic development (Tabara et al., 1999, *Development* 126: 1-11; Boyd et al., 1996, *Development* 122:3075-3084). All rde-strains tested (as well as mut-2 and mut-7) showed significant resistance to dsRNA targeting of these germline-specific genes (**FIG. 2B**), as well as to several other germline specific genes tested. The rde-3 data (asterisk in **FIG. 2B**) includes a 10% non-specific embryonic lethality present in the rde-3 strain.

[0156] To examine the effect of these mutations on genetic interference of somatically expressed genes, cells were injected with dsRNA targeting the cuticle collagen gene sqt-3 and the body muscle structural gene unc-22. sqt-3 hypomorphic mutants exhibit a short, dumpy body shape (dpy; van der Keyl et al., 1994, *Dev. Dyn.* 201:86-94). unc-22 mutations exhibit severe paralysis with a distinctive body twitching phenotype (Moerman et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:2579-2583). rde-1, rde-3, rde-4 and mut-2 strains showed strong resistance to both sqt-3 and unc-22 dsRNA, while rde-2 and mut-7 strains showed partial resistance. Thus rde-2 and mut-7 appeared to be partially tissue- or gene-specific in that they were required for effective RNAi against germline but not somatically expressed genes. The rde-1, rde-3, rde-4, and mut-2 (+) activities appeared to be required for interference for all genes analyzed. The rde and mut strains differ from one another in sensitivity to sqt-2 dsRNA.

[0157] Effect of rde on Transposon Mobilization

[0158] The effect of rde mutations on transposon mobilization was examined. Two of the newly identified mutants, rde-2 and rde-3 exhibited a level of transposon activation similar to that of mut-7 (Table 1). In contrast, transposon mobilization was not observed in the presence of rde-1 or rde-4 (Table 1).

TABLE 1

TRANSPOSON MOBILIZATION AND MALE INCIDENCE IN rde AND mut STRAINS	
Percentage of Non-Unc	Revertants
unc-22 (r765::Tc4)	0 (0/2000)
rde-1 (ne219);	0 (0/4000)
unc-22 (r765::Tc4)	
rde-2 (ne221);	0.96 (8/830)
unc-22 (r765::Tc4)	
rde-3 (ne298);	1.6 (35/2141)
unc-22 (r765::Tc4)	
rde-4 (ne299);	0 (0/2885)
unc-22 (r765::Tc4)	
mut-7 (pk204);	1.0 (40/3895)
unc-22 (r765::Tc4)	
Percentage of Male Animals	
Wild type (n2)	0.21 (2/934)
rde-1 (ne219)	0.07 (1/1530)
rde-2 (ne221)	3.2 (25/788)
rde-3 (ne298)	7.8 (71/912)
rde-4 (ne299)	0.24 (5/2055)

[0159] X-Chromosome Loss

[0160] Mutator strains (including mut-2, mut-7) rde-2 and rde-3) exhibit a second phenotype: a high incidence of males reflecting an increased frequency of X-chromosome loss during meiosis (Collins et al., 1987, supra; Ketting et al., supra). This phenotype was observed in rde-2 and rde-3 strains, but not observed in the rde-1 and rde-4 strains which showed a wild-type incidence of males (Table 1).

[0161] A previously described gene-silencing process appears to act on transgenes in the germline of *C. elegans*. Although the silencing mechanisms are not well understood, they are known to depend on the products of the genes mes-2, 3, 4 and 6 (Kelly and Fire, 1998, *Development* 125:2451-2456). To examine the possibility that the RNAi and germline transgene-silencing might share common mechanistic features, we first asked if the mes mutants were resistant to RNAi. We found normal levels of RNA interference in each of these strains. We next asked if RNAi deficient strains were defective in transgene-silencing. Three strains were analyzed: mut-7(pk204), rde-1(ne219) and rde-2(ne221).

[0162] To analyze transgene silencing in mut-7 worms, homozygous mut-7 lines carrying various GFP reporters transgenes were generated as follows: N2 (Bristol strain) males were mated to mut-7 (pk204) unc-32 (e]89) hermaphrodites; cross progeny males were then mated to strains carrying the GFP transgenes. mut-7 unc-32/++ cross progeny from these matings were cloned, and mut-7 unc-32 homozygous animals carrying the transgenes were isolated from their self-progeny. After the GFP reporter transgenes were introduced into different genetic backgrounds, activation of GFP transgene expression in germ cells was assayed at 25°C by fluorescence microscopy. The tested GFP reporter transgenes were each active in some or all somatic tissues, but had become silenced in the germline. The plasmids used and transgene designations are as follows: 1) pBK48 which contains an in-frame insertion of GFP into a ubiquitously expressed gene, let-858 (Kelly, et al., 1997, *Genetics* 146:227-238). ccExPD7271 contains more than 100 copies of pBK48 in a high copy repetitive array that is carried extrachromosomally. 2) pJH3.92 is an in-frame

fusion of GFP with the maternal pie-1 gene (M. Dunn and G. Seydoux, Johns Hopkins University, Baltimore, Md.). jhEx1070 carries pJH3.92 in a low copy "complex" extra-chromosomal array generated by the procedure of Kelly et al. (1997, supra) pJKL380.4 is a fusion of GFP with the *C. elegans* nuclear laminin gene, lam-1, which is expressed in all tissues (J. Liu and A. Fire). ccIn4810 carries pJKL380.4 in a complex array that has been integrated into the X chromosome by gamma irradiation using standard techniques.

[0163] The mut-7 strain was analyzed most extensively and was found to exhibit desilencing of three different germline transgenes tested (Table 2). The rde-2 strain exhibited a similar level of desilencing for a single transgene. In contrast, no transgene desilencing was observed in rde-1 mutants (Table 2). Thus, mut-7 and rde-2 which differ from rde-1 in having transposon mobilization and a high incidence of X-chromosome loss also differ from rde-1 in their ability to partially reactivate silent germline transgenes.

TABLE 2

REACTIVATION OF SILENCED TRANSGENES IN THE GERMLINE OF mut-7(pk204)		
Genotype	Transgene Array	Percentage of Germline Desilencing
+/+	ccEx7271	8.3 (4/48)
mut-7/+	ccEx7271	14.5 (7/48)
mut-7/mut-7	ccEx7271	91.0 (71/78)
+/+	jhEx1070	3.9 (2/51)
mut-7/mut-7	jhEx1070	86.5 (32/37)
+/+	ccin4810	4.3 (2/46)
mut-7/mut-7	ccin4810	73.3 (33/45)
rde-1/rde-1	ccEx7271	0 (0/34)

EXAMPLE 5

Requirement for rde-1(+) and rde-4(+) Activities in
Target Tissue

[0164] The rde-1 and rde-4 mutants differ from other RNAi deficient strains identified herein in that they do not cause transposon mobilization nor do they cause chromosome loss. The role of these genes in upstream events such as dsRNA uptake, transport or stability was examined. Such events could be required for interference induced by exogenous trigger RNAs but might be dispensable for natural functions of RNAi. To evaluate these upstream events, rde-1 and rde-4 homozygotes were exposed to dsRNA. The next generation was scored for interference. dsRNA targeting the unc-22 gene was injected into the intestinal cells of homozygous rde-1 and rde-4 hermaphrodites and the injected animals were then mated to wild-type males (**FIG. 3**). The self-progeny for both strains exhibited no interference with the targeted gene. However, there was potent interference in the rde-1/+ and rde-4/+ cross progeny (**FIG. 3**). These observations indicated that rde-1 and rde-4 mutants have intact mechanisms for transporting the interference effect from the site of injection (the intestine) into the embryos of the injected animal and then into the tissues of the resulting progeny. The stability of the resulting interference also appeared to be normal in rde-1 and rde-4 as the homozygous injected mothers continued to produce affected cross progeny for several days after the time of injection.

[0165] To examine whether rde-1 and rde-4 mutants could block interference caused by dsRNA expressed directly in the target tissue, the muscle-specific promoter from the myo-3 gene (Dibb et al., 1989, *J. Mol. Biol.* 205:605-613) was used to drive the expression of both strands of the muscle structural gene unc-22 in the body wall muscles (Moerman et al., 1986, supra; Fire et al., 1991, *Development* 113:503-514). A mixture of three plasmids was injected: [myo-3 promoter::unc-22 antisense], [myo-3::unc-22 sense], and a marker plasmid (pRF4[rol-6(sul 006gf)] [Mello et al., 1991]). Frequencies of Unc transgenic animals were followed in F1 and F2 generations. The ^aUnc phenotype was weak. Wild-type animals bearing this transgene exhibit a strong twitching phenotype consistent with unc-22 interference. The twitching phenotype was strongly suppressed by both rde-1 and rde-4 mutants (Table 3). The mut-7 and rde-2 mutants which are both sensitive to unc-22(RNAi) by micro-injection were also sensitive to promoter driven unc-22 interference in the muscle (Table 3). Taken together these findings suggest that rde-1(+) and rde-4(+) activities are not necessary for uptake or stability of the interfering RNA and may function directly in the target tissue.

TABLE 3

SENSITIVITY OF rde AND mut STRAINS TO TRANSGENE-DRIVEN INTERFERING RNA		
	Unc Animals in Transgenic F1	Unc F2 Lines in Inherited Lines
Wild type (N2)	26/59	10/11
rde-1 (ne219)	0/25	0/3
rde-2 (ne221)	35/72	14/14
rde-3 (ne298)	1 ^a /38	1 ^a /9
rde-4 (ne299)	0/51	0/4
mut-7 (pk204)	9/13	3/3

EXAMPLE 6

Molecular Identification of the rde-1 Gene

[0166] The rde-1 gene was cloned using standard genetic mapping to define a physical genetic interval likely to contain the gene using YACs and cosmids that rescue rde-1 mutants. These were used to identify a cloned rde-1 cDNA sequence and a cloned rde-4 sequence. These methods can also be used to identify the genes for rde-2, rde-3, and rde-5 using the mutant strains provided herein.

[0167] To clone an rde-1 gene, yeast artificial chromosome clones (YACs) containing *C. elegans* DNA from this interval were used to rescue the rde-1 mutant phenotype. To facilitate this analysis candidate rescuing YACs were co-injected with plasmids designed to express unc-22(RNAi). YAC and cosmid clones that mapped near the rde-1 locus were obtained from A. Coulson. rde-1 (ne219) was rescued by YAC clones: Y97C12 and Y50B5. The two overlapping YAC clones provided rde-1 rescuing activity as indicated by unc-22 genetic interference with characteristic body paralysis and twitching in the F1 and F2 transgenic animals. In contrast a non-overlapping YAC clone failed to rescue resulting in 100% non-twitching transgenic strains (**FIG. 4A**).

[0168] The rescuing activity was further localized to two overlapping cosmid clones, cosmid C27H6 and T10A5, and finally to a single 4.5 kb genomic PCR fragment predicted to contain a single gene, designated K08H10.7 (SEQ ID NO:1; FIGS. 5A-5C) The K08H10.7 PCR product gave strong rescue when amplified from wild-type genomic DNA. This rescue was greatly diminished using a PCR fragment amplified from any of the three *rde-1* alleles and was abolished by a 4 bp insertion at a unique *NheI* site in the *rde-1* coding region. A wild-type PCR product from an adjacent gene $C_{27}H_{6,4}$, also failed to rescue.

[0169] The K08H10.7 gene from each of the *rde-1* mutant strains was sequenced, and distinct point mutations were identified that are predicted to alter coding sequences in K08H10.7 (FIG. 4A). Based on these findings *rde-1* can be identified as the K08H10.7 gene.

[0170] A full-length cDNA sequence was determined for *rde-1* using the cDNA clones, yk296b10 and yk595h5. cDNA clones for *rde-1* were obtained from Y. Kohara (Gene Network Lab, National Institute of Genetics, Mishima 411, Japan). The cDNA sequence of coding region and 3'UTR was determined on yk296b10 except that the sequence of 5'UTR was determined on yk595h5. The GenBank accession number for *rde-1* cDNA is AF180730 (SEQ ID NO:2). The *rde-1* cDNA sequence was used to generate a predicted translation product (SEQ ID NO:3), referred to as RDE-1, consisting of 1020 amino acids. The RDE-1 sequence was used to query Genbank and identify numerous related genes in *C. elegans* as well as other animals and plants. This gene family includes at least 23 predicted *C. elegans* genes, several of which appear to be members of conserved subfamilies. Within subfamilies, conservation extends throughout the protein and all family members have a carboxy-terminal region that is highly conserved (FIG. 4B). Besides the genes shown in FIG. 4B, other related genes include ARGONAUTE 1 (*Arabidopsis*), SPCC736.11 (*S. pombe*), and Piwi (*Drosophila*). A portion of the N terminal region of RDE-1 showed no significant similarity to any of the identified related genes. There are no defined functional motifs within this gene family, but members including RDE-1 are predicted to be cytoplasmic or nuclear by PSORT analysis (Nakai and Horton, 1999, *Trends Biochem. Sci.* 24:34-36). Furthermore, one family member named eIF2C has been identified as a component of a cytoplasmic protein fraction isolated from rabbit reticulocyte lysates. The RDE-1 protein is most similar to the rabbit eIF2C. However, two other *C. elegans* family members are far more similar to eIF2C than is RDE-1 (FIG. 4B). RDE-1 may provide sequence-specific inhibition of translation initiation in response to dsRNA.

[0171] The *rde-1* mutations appear likely to reduce or eliminate *rde-1*(+) activity. Two *rde-1* alleles ne219 and ne297 are predicted to cause amino acid substitutions within the RDE-1 protein and were identified at a frequency similar to that expected for simple loss-of-function mutations. The *rde-1*(ne219) lesion alters a conserved glutamate to a lysine (FIG. 4B). The *rde-1*(ne297) lesion changes a non-conserved glycine, located four residues from the end of the protein, to a glutamate (FIG. 4B). The third allele, ne300, contains the strongest molecular lesion and is predicted to cause a premature stop codon prior to the most highly conserved region within the protein (Q>Ochre in FIG. 4B). Consistent with the idea that *rde-1* (ne300) is a strong loss of function mutation, we found that when placed in trans to

a chromosomal deficiency the resulting deficiency trans-heterozyotes were RNAi deficient but showed no additional phenotypes. These observations suggest that *rde-1* alleles are simple loss-of-function mutations affecting a gene required for RNAi but that is otherwise non-essential.

[0172] Because of its upstream role RNA interference (see Examples 8-10 below), the RDE-1 protein and fragments thereof can be used to prepare dsRNA that is useful as an RNAi agent.

EXAMPLE 7

Maternal Establishment and Paternal Transmission of RNAi

[0173] To examine whether the interference effect induced by RNAi exhibited linkage to the target gene (e.g., was involved in a reversible alteration of the gene or associated chromatin), a strain was constructed such that the F1 males that carry the RNAi effect also bear a chromosomal deletion that removes the target gene (FIG. 7B). In the case of linkage to the target gene, the RNAi effect would be transmitted as a dominant factor.

[0174] In experiments testing the linkage of the interference effect to the target gene, three different species of dsRNA (*pos-1* dsRNA, *mom-2* dsRNA, or *sgg-1* dsRNA) were delivered into *C. elegans* in independent experiments. The dsRNA was delivered by injection through a needle inserted into the intestine. In general, dsRNA was synthesized in vitro using T3 and T7 polymerases. Template DNA was removed from the RNA samples by DNase treatment (30 minutes at 37°C.). Equal amounts of sense and antisense RNAs were then mixed and annealed to obtain dsRNA. dsRNA at a concentration of 1-5 mg/ml was injected into the intestine of animals. In control experiments, mixtures of linearized template DNA plasmids used for synthesizing RNA failed to induce interference in P0, F1, or F2 animals when injected into the intestine of hermaphrodites at a concentration of 0.2 mg/ml. FIG. 7A illustrates this experiment. The gonad of the parent (P0) hermaphrodite has symmetrical anterior and posterior U-shaped arms as shown in FIG. 7A. Several fertilized eggs are shown in FIG. 7A, centrally located in the uterus. The rectangular mature oocytes are cued up in the gonad arms most proximal to the uterus. The embryos present in P0 at the time of injection gave rise to unaffected F1 progeny. Oocytes in the proximal arms of the injected P0 gonad inherit the RNAi effect but also carry a functional maternal mRNA (F1 carriers of RNAi).

[0175] After a clearance period during which carrier and unaffected F1 progeny are produced, the injected P0 begins to exclusively produce dead F1 embryos with the phenotype corresponding to the inactivation of the gene targeted by the injected RNA (Tabara et al. 1999, *Development* 126:1; C. Rocheleau, 1997, *Cell* 90:707). Potential F1 and F2 carriers of the interference effect were identified within the brood of the injected animal. In the case of hermaphrodites, carriers were defined as "affected" if the animals produced at least 20% dead embryos with phenotypes corresponding to maternal loss of function for the targeted locus. In the case of males, carriers were defined as animals whose cross progeny included at least one affected F2 hermaphrodite. The total number of carriers identified in each generation for each of

three dsRNAs injected is shown in **FIG. 7A** as a fraction of the total number of animals assayed.

[0176] To examine the extragenic inheritance of RNAi, experiments were carried out investigating whether sperm that inherit the deletion and therefore have no copies of the target locus could carry the interference effect into the F2 generation. F1 males that carried both *pos-1* (RNAi) and a chromosomal deficiency for the *pos-1* locus were generated. The chromosome carrying the deficiency for *pos-1* also carried a deficiency for phenotypically uncoordinated (*unc*). F2 progeny of the carrier male includes two genotypes: phenotypically wild-type animals that inherit the (+) chromosome, and phenotypically uncoordinated (*Unc*) progeny that inherit the mDF3 chromosome. In these experiments, the deficiency bearing sperm were just as capable as wild-type sperm of transferring interference to the F2 hermaphrodite progeny (**FIG. 7B**). Thus, the target locus was not needed for inheritance of the interference effect.

[0177] Surprisingly, although males were sensitive to RNAi and could inherit and transmit RNAi acquired from their mothers, direct injections into males failed to cause transmission of RNAi to the F1 for several genes tested. In an example of this type of protocol, wild type males were injected with targeting dsRNA: body muscle structural gene *unc-22*, cuticle collagen gene *sqt-3*, maternal genes *pos-1* and *sgg-1*. Males of the *pes-10::gfp* strain (Seydoux, G. and Dunn, Mass., 1997, *Development* 124:2191-2201) were injected with *gfp* dsRNA. Injected males were affected by *unc-22* and *gfp* dsRNA to the same extent as injected hermaphrodites. No RNAi interference was detected in F1 progeny or injected males (40 to 200 F1 animals scored for each RNA tested). Therefore, the initial transmission of RNAi to F1 progeny may involve a mechanism active only in hermaphrodites while subsequent transmission to the F2 progeny appears to involve a distinct mechanism, active in both hermaphrodites and males. The hermaphrodite-specific step may indicate the existence of a maternal germline process that amplifies the RNAi agent. These data show that extracts from the maternal germline tissues of *C. elegans* may be used in conjunction with RDE-1 and RDE-4 activity to create and to then amplify RNAi agents.

[0178] In addition, the germline factors that amplify the RNAi agents can be identified by mutations that result in an RNAi deficient mutant phenotype. Such factors can be used as additional components of an in vitro system for the efficient amplification of RNAi agents.

EXAMPLE 8

Sufficiency of Wild-Type Activities of *rde-1*, *rde-2*, *mut-7*, and *rde-4* in Injected Animals for Interference Among F1 Self Progeny

[0179] To investigate whether the activities of *rde-1*, *rde-2*, *rde-4*, and *mut-7*, respectively, are sufficient in injected hermaphrodites for interference in the F1 and F2 generations, crosses were designed such that wild-type activities of these genes would be present in the injected animal but absent in the F1 or F2 generations. To examine inheritance in the F1 generation, (hermaphrodite) mothers heterozygous for each mutant (P0) were injected, allowed to produce self-progeny (F1) and the homozygous mutant progeny in the F1 generation were examined for genetic interference

(**FIG. 8A**). To do this, the heterozygous hermaphrodites from each genotype class, *rde-1*, *unc-42/+*; *rde-2*, *unc-13/+*; *mut-7*, *dpy-71/+*; and *rde-4*, *unc-69/+* (the following alleles were used in this study: *rde-1*(*ne300*) *unc-42*, *rde-1*(*ne219*), *rde-2*(*ne221*), *rde-4*(*ne299*), and *mut-7*(*pk2040*) were injected with *pos-1* dsRNA. In each case, two types of F1 self progeny, distinguished by the presence of the linked marker mutations, were scored for interference (**FIG. 8A**). In these experiments the *rde-1* and *rde-4* mutant F1 progeny exhibited robust interference, comparable to wild-type, while the *rde-2* and *mut-7* F1 progeny failed to do so. In control experiments, homozygous F1 progeny from heterozygous (uninjected) mothers were directly injected with *pos-1* dsRNA (**FIG. 8B**). Injection of dsRNA directly into the *rde-1* and *rde-4* mutant progeny of uninjected heterozygous mothers failed to result in interference. Thus, injection of dsRNA into heterozygous hermaphrodites resulted in an inherited interference effect that triggered gene silencing in otherwise RNAi resistant *rde-1* and *rde-4* mutant F1 progeny while *rde-2* and *mut-7* mutant F1 progeny remained resistant.

[0180] In this experiment, the expression of *rde-1*(+) and *rde-4*(+) in the injected animal was sufficient for interference in later generations.

[0181] These data suggest that treatment of a dsRNA with functional *rde-1* and *rde-4* gene products can produce an agent that activates the remainder of the RNAi pathway.

EXAMPLE 9

Requirements for *rde-1*, *rde-2*, *rde-4*, and *mut-7* in F1 and F2 interference

[0182] To examine the genetic requirements for RNAi genes in the F2 generation, F1 male progeny were generated that carry the interference effect as well as one mutant copy of each respective locus; *rde-1*, *rde-2*, and *mut-7* (**FIG. 9A**). Each of these males was then backcrossed with uninjected hermaphrodites homozygous for each corresponding mutant (**FIG. 9A**). The resulting cross progeny (F1) included 50% heterozygotes and 50% homozygotes that were distinguished by the presence of the linked marker mutations. The heterozygous siblings served as controls and in each case exhibited interference at a frequency similar to that seen in wild-type animals (**FIG. 9A**). In these crosses, *rde-2* and *mut-7* homozygous F2 progeny failed to exhibit interference, indicating that the activities of these two genes are required for interference in the F2 generation. In contrast, we found that homozygous *rde-1* F2 animals exhibited wild-type levels of F2 interference (**FIG. 9A**). Control *rde-1* homozygotes generated through identical crosses were completely resistant to *pos-1::RNAi* when challenged de novo with dsRNA in the F2 generation. In these experiments, 35 *rde-1* homozygous animals generated through crosses shown in **FIG. 9A** were tested by feeding bacteria expressing *pos-1* dsRNA, and 21 similar animals were tested by direct injections of *pos-1* dsRNA. All animals tested were resistant to *pos-1* (RNAi). Thus, *rde-1* activity in the preceding generations was sufficient to allow interference to occur in *rde-1* mutant F2 animals while the wild-type activities of *rde-2* and *mut-7* were required directly in the F2 animals for interference.

[0183] In this experiment, the expression of *rde-1*(+) and *rde-4*(+) in the injected animal was sufficient for interference

in later generations. The wild-type activities of the *rde-2* and *mut-7* genes were required for interference in all generations assayed. Thus, *rde-2* and *mut-7* might be required only downstream or might also function along with *rde-1* and *rde-4*.

[0184] These data lend additional support to the concept that an appropriately treated dsRNA could be used as an RNAi agent.

EXAMPLE 10

Sufficiency of *rde-1* Activity to Initiate RNA Interference in Injected Animals That Lack the Wild-Type Activities of *rde-2*, *mut-7*, or *rde-4*

[0185] To ask if *rde-2* and *mut-7* activities function along with or downstream of *rde-1*, genetic cross experiments were designed in which the activities of these genes were present sequentially (FIG. 9B). For example, *rde-1(+);rde-2(-)* animals were injected with *pos-1* dsRNA and then crossed to generate F1 hermaphrodites homozygous for *rde-1(-); rde-2(+)*. In these experiments *rde-1(+)* activity in the injected animals was sufficient for F1 interference even when the injected animals were homozygous for *rde-2* or *mut-7* mutations (FIG. 10B). In contrast, *rde-1(+)* activity in the injected animals was not sufficient when the injected animals were homozygous for *rde-4* mutant (FIG. 10B). Thus, *rde-1* can act independently of *rde-2* and *mut-7* in the injected animal, but *rde-1* and *rde-4* must function together. These findings are consistent with the model that *rde-1* and *rde-4* function in the formation of the inherited interfering agent (i.e., an RNAi agent) while *rde-2* and *mut-7* function at a later step necessary for interference.

[0186] In summary, the above Examples provide genetic evidence for the formation and transmission of extragenic interfering agents in the *C. elegans* germline. Two *C. elegans* genes, *rde-1*, and *rde-4*, appear to be necessary for the formation of these extragenic agents but not for interference mediated by them. In contrast, the activities of two other genes, *rde-2* and *mut-7*, are required only downstream for interference.

[0187] These examples provide evidence that the *rde-1* and *rde-4* gene products or their homologs (e.g., from a mammal) can be used to prepare agents effective in activating the RNAi pathway.

EXAMPLE 11

rde-4 Sequences

[0188] An *rde-4* gene was cloned using methods similar to those described in Example 6. The nucleic acid sequence (SEQ ID NO:4) and predicted amino acid sequence (SEQ ID NO:5) are illustrated in FIG. 10.

[0189] Analysis of the *rde-4* nucleic acid sequence shows that it encodes a protein (RDE-4) with similarities to dsRNA binding proteins. Examples of the homology to X1 RBP (SEQ ID NO:6; Swissprot: locus TRBP_XENLA, accession Q91836; Eckmann and Jantsch, 1997, J. Cell Biol. 138:239-253) and HSPKR (SEQ ID NO:7; AAF13156.1; Xu and Williams, 1998, J. Interferon Cytokine Res. 18:609-616), and a consensus sequence (SEQ ID NO:8) are shown in FIG. 11. Three regions have been identified within the predicted RDE-4 protein corresponding to conserved

regions found in all members of this dsRNA binding domain family. These regions appear to be important for proper folding of the dsRNA binding domain. Conserved amino acid residues, important for interactions with the backbone of the dsRNA helix, are found in all members of the protein family including RDE-4 (see consensus residues in FIG. 11). This motif is thought to provide for general non-sequence-specific interactions with dsRNA. The RDE-4 protein contains conserved protein folds that are thought to be important for the assembly of the dsRNA binding domain in this family of proteins. Conserved amino acid residues in RDE-4 are identical to those that form contacts with the dsRNA in the crystal structure of the X1 RBP dsRNA complex. These findings strongly suggest that RDE-4 is likely to have dsRNA binding activity.

[0190] Because RDE-4 contains a motif that is likely to bind in a general fashion to any dsRNA and because RDE-4 appears to function upstream in the generation of RNAi agents, the RDE-4 protein or fragments thereof can be used to convert any dsRNA into an RNAi agent. In addition to the dsRNA binding domain, RDE-4 contains other functional domains that may mediate the formation of RNAi agents. These domains may provide for interaction between RDE-4 and RDE-1 or for binding to enzymes such as nucleases that convert the dsRNA into the RNAi agent. Because of its RNA binding function in RNA interference, the RDE-4 protein and fragments thereof can be used to prepare dsRNA that is useful in preparing an RNAi agent.

EXAMPLE 12

Identification of Regions of RDE-1 and RDE-4 that are Required for Creating an RNAi Agent

[0191] In vivo and in vitro assays are used to identify regions in RDE-1 and RDE-4 that are important for the generation of RNAi agents. In the in vivo assay, *rde-1* and *rde-4* are introduced into the corresponding *C. elegans* mutant strains via transgenes (Tabara et al., Cell 99:123 (1999); and Example 13). Important functional domains in RDE-1 and RDE-4 are defined by systematically altering the proteins followed by reintroduction into mutant animals to test for rescue of the RNAi deficient phenotype. A series of nested deletions are analyzed for rescue activity for both *rde-1* and *rde-4*. Specific point mutations are used to analyze the importance of specific amino acids. Chimeras are produced between RDE proteins and related proteins and genes. For example, coding sequences from RDE-1 homologs from the worm or from human are tested for their ability to rescue *rde-1* mutants. Replacing the RDE-4 dsRNA binding motif with a distinct RNA binding motif, e.g., one that recognizes a specific viral dsRNA sequence or a ssRNA sequence will alter the specificity of the RNAi response perhaps causing sequence-specific or ssRNA-induced gene targeting. In one form of the in vitro assay, whole protein extracts from *rde-1* or *rde-4* deficient worm strains are used.

[0192] Recombinant RDE-1 or RDE-4 is then added back to reconstitute the extract. Altered RDE-1 and RDE-4 proteins (described above, including deletions, point mutants and chimeras) are made in vitro and then tested for their ability to function when added back to these extracts. RNAi activity is analyzed by injecting the reconstituted extracts directly into animals or by assaying for the destruction of an added in vitro synthesized target mRNA.

EXAMPLE 13

Rescue of rde-4 Animals

[0193] Rescue of animals (e.g., *C. elegans*) that are mutant for an RNAi pathway is a useful method for identifying sequences from RNAi pathway genes that encode functional polypeptides, e.g., polypeptides that can eliminate the mutant phenotype.

[0194] An example of such a method for identifying rde-4 mutant animals is as follows. PCR using primers located 1 kb upstream and 500 nucleotides downstream of the open reading frame (T20G5.11; illustrated in FIG. 12) are used to amplify the rde-4 gene from *C. elegans* genomic DNA. The resulting PCR product is then injected along with reporter constructs described in Tabara et al. (Cell 99:123 (1999); incorporated herein in its entirety by reference), and the progeny of the injected animal are assayed for rescue of the

RNAi deficient phenotype. The PCR product can also be cloned into a plasmid vector for site directed mutational analysis of RDE-4 (see Example 12). Co-injection of such a wild type RDE-4 plasmid and altered derivatives can be used to identify functional domains of rde-4. Similar methods can be used to identify functional domains of rde-1 and other RNAi pathway components.

Other Embodiments

[0195] It is to be understood that while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 14

<210> SEQ ID NO 1

<211> LENGTH: 3719

<212> TYPE: DNA

<213> ORGANISM: *Caenorhabditis elegans*

<400> SEQUENCE: 1

```

cagccacaaa gtgatgaaac atgctcctcga attttcccga attggaaaaa ggattttatc      60
gtcatttctct cgatccgcta tgatcaatta ttagcagcta taagatatat aagtttgata      120
ttaatattat aggagatgaa atggcttgcg aggccactg gtaaatgcga cggcaaattc      180
tatgagaaga aagtacttct tttggtaaat tggttcaagt tctccagcaa aatttacgat      240
cgggaatact acgagatgta agtgaaaatg acaaaggaag tattgaatag aaaaccagga      300
aaacctttcc caaaaaagac agaaattcca atgtaagtgc ttgtaaatta gtcaaaacta      360
attttatttt tcagtcccga tcgtgcaaaa ctcttctggc aacatcttcg gcatgagaag      420
aagcagacag attttattct cgaagactat gtttttgatg aaaaggacac tgtttatagt      480
gtttgtcgac tgaacactgt cacatcaaaa atgctggttt cggagaaagt agtaaaaaag      540
gattcggaga aaaaagatga aaaggatttg gagaaaaaaa tcttatacac aatgatactt      600
acctatcgta aaaaatttca cctgaacttt agtcgagaaa atccggaaaa agacgaagaa      660
gcgaatcggg gttacaaatt cctgaaggtt tatgaaaaac acgcattata acaaacaaaa      720
ttagctttca gaatgttatg acccagaaag ttcgctacgc gccttttttg aacgaggaga      780
ttaaagtgtg agttgcaata ataataataa taatcacctc aactcattta tatattttaa      840
gacaattcgc gaaaaatttt gtgtacgata ataattcaat tctgcgagtt cctgaatcgt      900
ttcacgatcc aaacagattc gaacaatcat tagaagtagc accaagaatc gaagcatggt      960
ttggaattta cattggaatc aaagaattgt tcgatggtga acctgtgctc aattttgcaa     1020
gtaagtttga gaaactgcga taaaaatca tgtgattttt gttgaagttg tcgataaact     1080
attctacaat gcaccgaaaa tgtctcttct ggattatcct ctccctaattg tcgaccccca     1140
gtcgtgtaac gatgatgtac gaaaagatct taaaacaaaa ctgatggcgg gaaaaatgac     1200
aatcagacaa gcccgcgggc caagaattcg acaattattg gaaaatttga agctgaaatg     1260

```


-continued

cgcagaagtt tgggataacg aatgttagt ttaaattatt caaacaatta atatacaaat	1320
tgatthtcag gtcgagattg acagaacgac atctgacatt tctagatttg tgcgaggaaa	1380
actctcttgt ttataaagtc actggtaaat cggacagagg aagaaatgca aaaaagtacg	1440
atactacatt gttcaaaatc tatgaggaaa acaaaaagtt cattgagttt cccacactac	1500
cactagtcaa agttaaaggt ggagcaaaag aatacgtgt accaatggaa catcttgaag	1560
ttcatgagaa gccacaaaga tacaagaatc gaattgatct ggtgatgcaa gacaagtthc	1620
taaagcgagc tacacgaaaa cctcacgact acaaaagaaa tacctaaaa atgctgaaag	1680
aattggattt ctctctgaa gagctaaatt ttgttgaaag atttgatta tgctccaaac	1740
ttcagatgat cgaatgtcca ggaaaggtht tgaaagagcc aatgcttggt aatagtgtaa	1800
atgaacaaat taaaatgaca ccagtgattc gtggatttca agaaaaacaa ttgaatgtgg	1860
ttcccgaaaa agaactttgc tgtgctgttt ttgtagtcaa cgaaacagcg ggaatccat	1920
gcttagaaga gaacgacgtht gtgtaagtgt thctacgta gattattccg aaataththc	1980
agtaagthct acaccgaact aattggthgt tgcaagthcc gtggaatagc aattgthgcc	2040
aatgaaaaca gaggagcgca atctattatg tacgacgcca cgaaaaatga atatgccgta	2100
agthtcagaa aattgaaagt thttaaatac cataththca gthctacaaa aattgtacac	2160
taaaaccgg aatcggtaga thtgaaatag ccgcaacaga agcgaagaat atgththgaa	2220
gtctthccga taaagaacaa aaagtcttaa tgttcattat cattthcaaa cgacaactga	2280
atgcttacgg ththtgaaa cattattgag atcacacat cggtgtagct aatcagcata	2340
thactthctga aacagtcaca aaagctthgg catcactaag gcacgagaaa ggatcaaac	2400
gaaththctta tcaaatgca thgaaatca acgcaaat agggagthatt aaccaggagc	2460
thgactgthc agaaatgca gaaatcac cagaagaaaa agaagacgg aaaacaatgc	2520
cattaactat gtatgthga athgatgtaa thcatccaac thctacagth ggaathgatt	2580
athctatagc ggctgtagta gcgagthaca atccagthgg aactatctat cgaaatagta	2640
thgtgactca agaagaatgt cgtccgthgt agcgtgagth ggctcatgga cgggaaagaa	2700
cagataththt ggaagcaaa thctgthaaat thctcagaga athctcagaa thgagthgthc	2760
thgagthattt aaaagathctc thggathththt aaththththt taaactthca gaacaacgac	2820
aatcagagcac cagcgcatat thgtagthctat cgagacggag thagcgathc ggagthgcta	2880
cgtgthtagthc atgathgagthc thgatctthta aaaagcgaag taaaacaath catgthcgga	2940
cgggathggag aagathccaga gccgaagthac acgthctathg thathctagaa aagacacath	3000
acagaththc thcgaagaat ggaaaaagat aagcagthgg tcaataaaga ththactctc	3060
gctgaaacag atgthctgtht thctgctgtht aaacaathggg aggagthath gaaagaaagc	3120
aaagaaactg gaaththgaa cccatcatcc ggaacaactg thgataaact thctgththc	3180
aaatacaaat thgatthththt ththggcatct catcatgthg thctthgthac atctcgtcca	3240
ggacaththca ctgththgta thgacgataaa ggaathgagcc aagathgaagth ctaththagc	3300
gthththgaa gcaaththagc athththgath ththgthaatcc gcatathgtht aththaaaa	3360
aathththcag aaththgacct acggaactthc ththctctctc gctagathgthc gaaacccat	3420
ctcgtthcct gththcggthc aththgthca thththctgtht gaaaaagcga aagagctthta	3480
thcgaactthac aaggaacath acathcggthg ctathgthcag ccacggaact gacacgaaat	3540

-continued

```

ggaacatttt ctccaaacta acgtgaagta ccctggaatg tcgttcgcat aacattttgc 3600
aaaagtgtcg cccggttcaa tcaaattttt caattgtaga tattgtactt actttttttt 3660
aaagcccggt ttcaaaaatt cattccatga ctaacgtttt cataaattac ttgaaattt 3719

```

```

<210> SEQ ID NO 2
<211> LENGTH: 3227
<212> TYPE: DNA
<213> ORGANISM: Caenorhabditis elegans
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (21)...(3080)

```

```

<400> SEQUENCE: 2

```

```

cagccacaaa gtgatgaaac atg tcc tcg aat ttt ccc gaa ttg gaa aaa gga 53
                Met Ser Ser Asn Phe Pro Glu Leu Glu Lys Gly
                1                5                10

ttt tat cgt cat tct ctc gat ccg gag atg aaa tgg ctt gcg agg ccc 101
Phe Tyr Arg His Ser Leu Asp Pro Glu Met Lys Trp Leu Ala Arg Pro
                15                20                25

act ggt aaa tgc gac ggc aaa ttc tat gag aag aaa gta ctt ctt ttg 149
Thr Gly Lys Cys Asp Gly Lys Phe Tyr Glu Lys Lys Val Leu Leu Leu
                30                35                40

gta aat tgg ttc aag ttc tcc agc aaa att tac gat cgg gaa tac tac 197
Val Asn Trp Phe Lys Phe Ser Ser Lys Ile Tyr Asp Arg Glu Tyr Tyr
                45                50                55

gag tat gaa gtg aaa atg aca aag gaa gta ttg aat aga aaa cca gga 245
Glu Tyr Glu Val Lys Met Thr Lys Glu Val Leu Asn Arg Lys Pro Gly
                60                65                70                75

aaa cct ttc cca aaa aag aca gaa att cca att ccc gat cgt gca aaa 293
Lys Pro Phe Pro Lys Lys Thr Glu Ile Pro Ile Pro Asp Arg Ala Lys
                80                85                90

ctc ttc tgg caa cat ctt cgg cat gag aag aag cag aca gat ttt att 341
Leu Phe Trp Gln His Leu Arg His Glu Lys Lys Gln Thr Asp Phe Ile
                95                100                105

ctc gaa gac tat gtt ttt gat gaa aag gac act gtt tat agt gtt tgt 389
Leu Glu Asp Tyr Val Phe Asp Glu Lys Asp Thr Val Tyr Ser Val Cys
                110                115                120

cga ctg aac act gtc aca tca aaa atg ctg gtt tcg gag aaa gta gta 437
Arg Leu Asn Thr Val Thr Ser Lys Met Leu Val Ser Glu Lys Val Val
                125                130                135

aaa aag gat tcg gag aaa aaa gat gaa aag gat ttg gag aaa aaa atc 485
Lys Lys Asp Ser Glu Lys Lys Asp Glu Lys Asp Leu Glu Lys Lys Ile
                140                145                150                155

tta tac aca atg ata ctt acc tat cgt aaa aaa ttt cac ctg aac ttt 533
Leu Tyr Thr Met Ile Leu Thr Tyr Arg Lys Lys Phe His Leu Asn Phe
                160                165                170

agt cga gaa aat ccg gaa aaa gac gaa gaa gcg aat cgg agt tac aaa 581
Ser Arg Glu Asn Pro Glu Lys Asp Glu Glu Ala Asn Arg Ser Tyr Lys
                175                180                185

ttc ctg aag aat gtt atg acc cag aaa gtt cgc tac gcg cct ttt gtg 629
Phe Leu Lys Asn Val Met Thr Gln Lys Val Arg Tyr Ala Pro Phe Val
                190                195                200

aac gag gag att aaa gta caa ttc gcg aaa aat ttt gtg tac gat aat 677
Asn Glu Glu Ile Lys Val Gln Phe Ala Lys Asn Phe Val Tyr Asp Asn
                205                210                215

aat tca att ctg cga gtt cct gaa tcg ttt cac gat cca aac aga ttc 725
Asn Ser Ile Leu Arg Val Pro Glu Ser Phe His Asp Pro Asn Arg Phe

```

-continued

220	225	230	235	
gaa caa tca tta gaa gta gca cca aga atc gaa gca tgg ttt gga att Glu Gln Ser Leu Glu Val Ala Pro Arg Ile Glu Ala Trp Phe Gly Ile 240 245 250				773
tac att gga atc aaa gaa ttg ttc gat ggt gaa cct gtg ctc aat ttt Tyr Ile Gly Ile Lys Glu Leu Phe Asp Gly Glu Pro Val Leu Asn Phe 255 260 265				821
gca att gtc gat aaa cta ttc tac aat gca ccg aaa atg tct ctt ctg Ala Ile Val Asp Lys Leu Phe Tyr Asn Ala Pro Lys Met Ser Leu Leu 270 275 280				869
gat tat ctt ctc cta att gtc gac ccc cag tcg tgt aac gat gat gta Asp Tyr Leu Leu Leu Ile Val Asp Pro Gln Ser Cys Asn Asp Asp Val 285 290 295				917
cga aaa gat ctt aaa aca aaa ctg atg gcg gga aaa atg aca atc aga Arg Lys Asp Leu Lys Thr Lys Leu Met Ala Gly Lys Met Thr Ile Arg 300 305 310 315				965
caa gcc gcg cgg cca aga att cga caa tta ttg gaa aat ttg aag ctg Gln Ala Ala Arg Pro Arg Ile Arg Gln Leu Leu Glu Asn Leu Lys Leu 320 325 330				1013
aaa tgc gca gaa gtt tgg gat aac gaa atg tcg aga ttg aca gaa cga Lys Cys Ala Glu Val Trp Asp Asn Glu Met Ser Arg Leu Thr Glu Arg 335 340 345				1061
cat ctg aca ttt cta gat ttg tgc gag gaa aac tct ctt gtt tat aaa His Leu Thr Phe Leu Asp Leu Cys Glu Glu Asn Ser Leu Val Tyr Lys 350 355 360				1109
gtc act ggt aaa tcg gac aga gga aga aat gca aaa aag tac gat act Val Thr Gly Lys Ser Asp Arg Gly Arg Asn Ala Lys Lys Tyr Asp Thr 365 370 375				1157
aca ttg ttc aaa atc tat gag gaa aac aaa aag ttc att gag ttt ccc Thr Leu Phe Lys Ile Tyr Glu Glu Asn Lys Lys Phe Ile Glu Phe Pro 380 385 390 395				1205
cac cta cca cta gtc aaa gtt aaa agt gga gca aaa gaa tac gct gta His Leu Pro Leu Val Lys Val Lys Ser Gly Ala Lys Glu Tyr Ala Val 400 405 410				1253
cca atg gaa cat ctt gaa gtt cat gag aag cca caa aga tac aag aat Pro Met Glu His Leu Glu Val His Glu Lys Pro Gln Arg Tyr Lys Asn 415 420 425				1301
cga att gat ctg gtg atg caa gac aag ttt cta aag cga gct aca cga Arg Ile Asp Leu Val Met Gln Asp Lys Phe Leu Lys Arg Ala Thr Arg 430 435 440				1349
aaa cct cac gac tac aaa gaa aat acc cta aaa atg ctg aaa gaa ttg Lys Pro His Asp Tyr Lys Glu Asn Thr Leu Lys Met Leu Lys Glu Leu 445 450 455				1397
gat ttc tct tct gaa gag cta aat ttt gtt gaa aga ttt gga tta tgc Asp Phe Ser Ser Glu Glu Leu Asn Phe Val Glu Arg Phe Gly Leu Cys 460 465 470 475				1445
tcc aaa ctt cag atg atc gaa tgt cca gga aag gtt ttg aaa gag cca Ser Lys Leu Gln Met Ile Glu Cys Pro Gly Lys Val Leu Lys Glu Pro 480 485 490				1493
atg ctt gtg aat agt gta aat gaa caa att aaa atg aca cca gtg att Met Leu Val Asn Ser Val Asn Glu Gln Ile Lys Met Thr Pro Val Ile 495 500 505				1541
cgt gga ttt caa gaa aaa caa ttg aat gtg gtt ccc gaa aaa gaa ctt Arg Gly Phe Gln Glu Lys Gln Leu Asn Val Val Pro Glu Lys Glu Leu 510 515 520				1589
tgc tgt gct gtt ttt gta gtc aac gaa aca gcg gga aat cca tgc tta Cys Cys Ala Val Phe Val Val Asn Glu Thr Ala Gly Asn Pro Cys Leu				1637

-continued

525	530	535	
gaa gag aac gac gtt gtt aag ttc tac acc gaa cta att ggt ggt tgc Glu Glu Asn Asp Val Val Lys Phe Tyr Thr Glu Leu Ile Gly Gly Cys 540 545 550 555			1685
aag ttc cgt gga ata cga att ggt gcc aat gaa aac aga gga gcg caa Lys Phe Arg Gly Ile Arg Ile Gly Ala Asn Glu Asn Arg Gly Ala Gln 560 565 570			1733
tct att atg tac gac gcg acg aaa aat gaa tat gcc ttc tac aaa aat Ser Ile Met Tyr Asp Ala Thr Lys Asn Glu Tyr Ala Phe Tyr Lys Asn 575 580 585			1781
tgt aca cta aat acc gga atc ggt aga ttt gaa ata gcc gca aca gaa Cys Thr Leu Asn Thr Gly Ile Gly Arg Phe Glu Ile Ala Ala Thr Glu 590 595 600			1829
gcg aag aat atg ttt gaa cgt ctt ccc gat aaa gaa caa aaa gtc tta Ala Lys Asn Met Phe Glu Arg Leu Pro Asp Lys Glu Gln Lys Val Leu 605 610 615			1877
atg ttc att atc att tcc aaa cga caa ctg aat gct tac ggt ttt gtg Met Phe Ile Ile Ile Ser Lys Arg Gln Leu Asn Ala Tyr Gly Phe Val 620 625 630 635			1925
aaa cat tat tgc gat cac acc atc ggt gta gct aat cag cat att act Lys His Tyr Cys Asp His Thr Ile Gly Val Ala Asn Gln His Ile Thr 640 645 650			1973
tct gaa aca gtc aca aaa gct ttg gca tca cta agg cac gag aaa gga Ser Glu Thr Val Thr Lys Ala Leu Ala Ser Leu Arg His Glu Lys Gly 655 660 665			2021
tca aaa cga att ttc tat caa att gca ttg aaa atc aac gcg aaa tta Ser Lys Arg Ile Phe Tyr Gln Ile Ala Leu Lys Ile Asn Ala Lys Leu 670 675 680			2069
gga ggt att aac cag gag ctt gac tgg tca gaa att gca gaa ata tca Gly Gly Ile Asn Gln Glu Leu Asp Trp Ser Glu Ile Ala Glu Ile Ser 685 690 695			2117
cca gaa gaa aaa gaa aga cgg aaa aca atg cca tta act atg tat gtt Pro Glu Glu Lys Glu Arg Arg Lys Thr Met Pro Leu Thr Met Tyr Val 700 705 710 715			2165
gga att gat gta act cat cca acc tcc tac agt gga att gat tat tct Gly Ile Asp Val Thr His Pro Thr Ser Tyr Ser Gly Ile Asp Tyr Ser 720 725 730			2213
ata gcg gct gta gta gcg agt atc aat cca ggt gga act atc tat cga Ile Ala Ala Val Val Ala Ser Ile Asn Pro Gly Gly Thr Ile Tyr Arg 735 740 745			2261
aat atg att gtg act caa gaa gaa tgt cgt ccc ggt gag cgt gca gtg Asn Met Ile Val Thr Gln Glu Glu Cys Arg Pro Gly Glu Arg Ala Val 750 755 760			2309
gct cat gga cgg gaa aga aca gat att ttg gaa gca aag ttc gtg aaa Ala His Gly Arg Glu Arg Thr Asp Ile Leu Glu Ala Lys Phe Val Lys 765 770 775			2357
ttg ctc aga gaa ttc gca gaa aac aac gac aat cga gca cca gcg cat Leu Leu Arg Glu Phe Ala Glu Asn Asn Asp Asn Arg Ala Pro Ala His 780 785 790 795			2405
att gta gtc tat cga gac gga gtt agc gat tcg gag atg cta cgt gtt Ile Val Val Tyr Arg Asp Gly Val Ser Asp Ser Glu Met Leu Arg Val 800 805 810			2453
agt cat gat gag ctt cga tct tta aaa agc gaa gta aaa caa ttc atg Ser His Asp Glu Leu Arg Ser Leu Lys Ser Glu Val Lys Gln Phe Met 815 820 825			2501
tcg gaa cgg gat gga gaa gat cca gag ccg aag tac acg ttc att gtg Ser Glu Arg Asp Gly Glu Asp Pro Glu Pro Lys Tyr Thr Phe Ile Val			2549

-continued

830	835	840	
att cag aaa aga cac aat aca cga ttg ctt cga aga atg gaa aaa gat			2597
Ile Gln Lys Arg His Asn Thr Arg Leu Leu Arg Arg Met Glu Lys Asp			
845	850	855	
aag cca gtg gtc aat aaa gat ctt act cct gct gaa aca gat gtc gct			2645
Lys Pro Val Val Asn Lys Asp Leu Thr Pro Ala Glu Thr Asp Val Ala			
860	865	870	875
gtt gct gct gtt aaa caa tgg gag gag gat atg aaa gaa agc aaa gaa			2693
Val Ala Ala Val Lys Gln Trp Glu Glu Asp Met Lys Glu Ser Lys Glu			
880		885	890
act gga att gtg aac cca tca tcc gga aca act gtg gat aaa ctt atc			2741
Thr Gly Ile Val Asn Pro Ser Ser Gly Thr Thr Val Asp Lys Leu Ile			
895	900		905
gtt tcg aaa tac aaa ttc gat ttt ttc ttg gca tct cat cat ggt gtc			2789
Val Ser Lys Tyr Lys Phe Asp Phe Phe Leu Ala Ser His His Gly Val			
910	915		920
ctt ggt aca tct cgt cca gga cat tac act gtt atg tat gac gat aaa			2837
Leu Gly Thr Ser Arg Pro Gly His Tyr Thr Val Met Tyr Asp Asp Lys			
925	930		935
gga atg agc caa gat gaa gtc tat aaa atg acc tac gga ctt gct ttt			2885
Gly Met Ser Gln Asp Glu Val Tyr Lys Met Thr Tyr Gly Leu Ala Phe			
940	945	950	955
ctc tct gct aga tgt cga aaa ccc atc tcg ttg cct gtt ccg gtt cat			2933
Leu Ser Ala Arg Cys Arg Lys Pro Ile Ser Leu Pro Val Pro Val His			
960		965	970
tat gct cat tta tca tgt gaa aaa gcg aaa gag ctt tat cga act tac			2981
Tyr Ala His Leu Ser Cys Glu Lys Ala Lys Glu Leu Tyr Arg Thr Tyr			
975	980		985
aag gaa cat tac atc ggt gac tat gca cag cca cgg act cga cac gaa			3029
Lys Glu His Tyr Ile Gly Asp Tyr Ala Gln Pro Arg Thr Arg His Glu			
990	995		1000
atg gaa cat ttt ctc caa act aac gtg aag tac cct gga atg tcg ttc			3077
Met Glu His Phe Leu Gln Thr Asn Val Lys Tyr Pro Gly Met Ser Phe			
1005	1010		1015
gca taacattttg caaaagtgtc gcccgtttca atcaaatttt tcaattgtag			3130
Ala			
1020			
atattgtact tacttttttt taaagcccg tttcaaaaat tcattccatg actaacgttt			3190
tcataaatta cttgaaattt aaaaaaaaa aaaaaaa			3227

<210> SEQ ID NO 3

<211> LENGTH: 1020

<212> TYPE: PRT

<213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 3

Met Ser Ser Asn Phe Pro Glu Leu Glu Lys Gly Phe Tyr Arg His Ser
 1 5 10 15

Leu Asp Pro Glu Met Lys Trp Leu Ala Arg Pro Thr Gly Lys Cys Asp
 20 25 30

Gly Lys Phe Tyr Glu Lys Lys Val Leu Leu Leu Val Asn Trp Phe Lys
 35 40 45

Phe Ser Ser Lys Ile Tyr Asp Arg Glu Tyr Tyr Glu Tyr Glu Val Lys
 50 55 60

Met Thr Lys Glu Val Leu Asn Arg Lys Pro Gly Lys Pro Phe Pro Lys
 65 70 75 80

-continued

Lys Thr Glu Ile Pro Ile Pro Asp Arg Ala Lys Leu Phe Trp Gln His
85 90 95

Leu Arg His Glu Lys Lys Gln Thr Asp Phe Ile Leu Glu Asp Tyr Val
100 105 110

Phe Asp Glu Lys Asp Thr Val Tyr Ser Val Cys Arg Leu Asn Thr Val
115 120 125

Thr Ser Lys Met Leu Val Ser Glu Lys Val Val Lys Lys Asp Ser Glu
130 135 140

Lys Lys Asp Glu Lys Asp Leu Glu Lys Lys Ile Leu Tyr Thr Met Ile
145 150 155 160

Leu Thr Tyr Arg Lys Lys Phe His Leu Asn Phe Ser Arg Glu Asn Pro
165 170 175

Glu Lys Asp Glu Glu Ala Asn Arg Ser Tyr Lys Phe Leu Lys Asn Val
180 185 190

Met Thr Gln Lys Val Arg Tyr Ala Pro Phe Val Asn Glu Glu Ile Lys
195 200 205

Val Gln Phe Ala Lys Asn Phe Val Tyr Asp Asn Asn Ser Ile Leu Arg
210 215 220

Val Pro Glu Ser Phe His Asp Pro Asn Arg Phe Glu Gln Ser Leu Glu
225 230 235 240

Val Ala Pro Arg Ile Glu Ala Trp Phe Gly Ile Tyr Ile Gly Ile Lys
245 250 255

Glu Leu Phe Asp Gly Glu Pro Val Leu Asn Phe Ala Ile Val Asp Lys
260 265 270

Leu Phe Tyr Asn Ala Pro Lys Met Ser Leu Leu Asp Tyr Leu Leu Leu
275 280 285

Ile Val Asp Pro Gln Ser Cys Asn Asp Asp Val Arg Lys Asp Leu Lys
290 295 300

Thr Lys Leu Met Ala Gly Lys Met Thr Ile Arg Gln Ala Ala Arg Pro
305 310 315 320

Arg Ile Arg Gln Leu Leu Glu Asn Leu Lys Leu Lys Cys Ala Glu Val
325 330 335

Trp Asp Asn Glu Met Ser Arg Leu Thr Glu Arg His Leu Thr Phe Leu
340 345 350

Asp Leu Cys Glu Glu Asn Ser Leu Val Tyr Lys Val Thr Gly Lys Ser
355 360 365

Asp Arg Gly Arg Asn Ala Lys Lys Tyr Asp Thr Thr Leu Phe Lys Ile
370 375 380

Tyr Glu Glu Asn Lys Lys Phe Ile Glu Phe Pro His Leu Pro Leu Val
385 390 395 400

Lys Val Lys Ser Gly Ala Lys Glu Tyr Ala Val Pro Met Glu His Leu
405 410 415

Glu Val His Glu Lys Pro Gln Arg Tyr Lys Asn Arg Ile Asp Leu Val
420 425 430

Met Gln Asp Lys Phe Leu Lys Arg Ala Thr Arg Lys Pro His Asp Tyr
435 440 445

Lys Glu Asn Thr Leu Lys Met Leu Lys Glu Leu Asp Phe Ser Ser Glu
450 455 460

Glu Leu Asn Phe Val Glu Arg Phe Gly Leu Cys Ser Lys Leu Gln Met
465 470 475 480

-continued

Ile	Glu	Cys	Pro	Gly	Lys	Val	Leu	Lys	Glu	Pro	Met	Leu	Val	Asn	Ser
				485					490					495	
Val	Asn	Glu	Gln	Ile	Lys	Met	Thr	Pro	Val	Ile	Arg	Gly	Phe	Gln	Glu
			500					505					510		
Lys	Gln	Leu	Asn	Val	Val	Pro	Glu	Lys	Glu	Leu	Cys	Cys	Ala	Val	Phe
		515					520					525			
Val	Val	Asn	Glu	Thr	Ala	Gly	Asn	Pro	Cys	Leu	Glu	Glu	Asn	Asp	Val
	530					535					540				
Val	Lys	Phe	Tyr	Thr	Glu	Leu	Ile	Gly	Gly	Cys	Lys	Phe	Arg	Gly	Ile
545					550					555					560
Arg	Ile	Gly	Ala	Asn	Glu	Asn	Arg	Gly	Ala	Gln	Ser	Ile	Met	Tyr	Asp
				565					570						575
Ala	Thr	Lys	Asn	Glu	Tyr	Ala	Phe	Tyr	Lys	Asn	Cys	Thr	Leu	Asn	Thr
			580						585					590	
Gly	Ile	Gly	Arg	Phe	Glu	Ile	Ala	Ala	Thr	Glu	Ala	Lys	Asn	Met	Phe
		595					600						605		
Glu	Arg	Leu	Pro	Asp	Lys	Glu	Gln	Lys	Val	Leu	Met	Phe	Ile	Ile	Ile
	610					615					620				
Ser	Lys	Arg	Gln	Leu	Asn	Ala	Tyr	Gly	Phe	Val	Lys	His	Tyr	Cys	Asp
625					630					635					640
His	Thr	Ile	Gly	Val	Ala	Asn	Gln	His	Ile	Thr	Ser	Glu	Thr	Val	Thr
				645					650					655	
Lys	Ala	Leu	Ala	Ser	Leu	Arg	His	Glu	Lys	Gly	Ser	Lys	Arg	Ile	Phe
			660					665					670		
Tyr	Gln	Ile	Ala	Leu	Lys	Ile	Asn	Ala	Lys	Leu	Gly	Gly	Ile	Asn	Gln
		675					680					685			
Glu	Leu	Asp	Trp	Ser	Glu	Ile	Ala	Glu	Ile	Ser	Pro	Glu	Glu	Lys	Glu
	690					695					700				
Arg	Arg	Lys	Thr	Met	Pro	Leu	Thr	Met	Tyr	Val	Gly	Ile	Asp	Val	Thr
705					710					715					720
His	Pro	Thr	Ser	Tyr	Ser	Gly	Ile	Asp	Tyr	Ser	Ile	Ala	Ala	Val	Val
				725					730					735	
Ala	Ser	Ile	Asn	Pro	Gly	Gly	Thr	Ile	Tyr	Arg	Asn	Met	Ile	Val	Thr
			740					745					750		
Gln	Glu	Glu	Cys	Arg	Pro	Gly	Glu	Arg	Ala	Val	Ala	His	Gly	Arg	Glu
		755					760					765			
Arg	Thr	Asp	Ile	Leu	Glu	Ala	Lys	Phe	Val	Lys	Leu	Leu	Arg	Glu	Phe
	770					775					780				
Ala	Glu	Asn	Asn	Asp	Asn	Arg	Ala	Pro	Ala	His	Ile	Val	Val	Tyr	Arg
785					790					795					800
Asp	Gly	Val	Ser	Asp	Ser	Glu	Met	Leu	Arg	Val	Ser	His	Asp	Glu	Leu
				805					810					815	
Arg	Ser	Leu	Lys	Ser	Glu	Val	Lys	Gln	Phe	Met	Ser	Glu	Arg	Asp	Gly
			820					825					830		
Glu	Asp	Pro	Glu	Pro	Lys	Tyr	Thr	Phe	Ile	Val	Ile	Gln	Lys	Arg	His
	835						840					845			
Asn	Thr	Arg	Leu	Leu	Arg	Arg	Met	Glu	Lys	Asp	Lys	Pro	Val	Val	Asn
	850					855					860				
Lys	Asp	Leu	Thr	Pro	Ala	Glu	Thr	Asp	Val	Ala	Val	Ala	Ala	Val	Lys
865					870					875					880
Gln	Trp	Glu	Glu	Asp	Met	Lys	Glu	Ser	Lys	Glu	Thr	Gly	Ile	Val	Asn

-continued

885										890					895				
Pro	Ser	Ser	Gly	Thr	Thr	Val	Asp	Lys	Leu	Ile	Val	Ser	Lys	Tyr	Lys				
			900					905					910						
Phe	Asp	Phe	Phe	Leu	Ala	Ser	His	His	Gly	Val	Leu	Gly	Thr	Ser	Arg				
		915					920					925							
Pro	Gly	His	Tyr	Thr	Val	Met	Tyr	Asp	Asp	Lys	Gly	Met	Ser	Gln	Asp				
	930					935					940								
Glu	Val	Tyr	Lys	Met	Thr	Tyr	Gly	Leu	Ala	Phe	Leu	Ser	Ala	Arg	Cys				
945					950					955					960				
Arg	Lys	Pro	Ile	Ser	Leu	Pro	Val	Pro	Val	His	Tyr	Ala	His	Leu	Ser				
				965					970						975				
Cys	Glu	Lys	Ala	Lys	Glu	Leu	Tyr	Arg	Thr	Tyr	Lys	Glu	His	Tyr	Ile				
			980					985					990						
Gly	Asp	Tyr	Ala	Gln	Pro	Arg	Thr	Arg	His	Glu	Met	Glu	His	Phe	Leu				
		995					1000					1005							
Gln	Thr	Asn	Val	Lys	Tyr	Pro	Gly	Met	Ser	Phe	Ala								
	1010						1015				1020								

<210> SEQ ID NO 4
 <211> LENGTH: 1222
 <212> TYPE: DNA
 <213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 4

```

atggatttaa ccaaactaac gtttgaaagc gttttcggcg gatcagatgt tcctatgaag    60
ccttcccgat cggaggataa caaaacgcca agaaacagaa cagatttggg gatgtttctg    120
aagaaaactc ccctcatggt actagaagag gctgctaagg ctgtctatca aaagacgcca    180
acttggggca ctgtcgaaact tcctgaagcc ttcgagatga cgttgattct gaatgaaatt    240
actgtaaaag gccaggcaac aagcaagaaa gctgcgagac aaaaggctgc tgttgaatat    300
ttacgcaagg ttgtggagaa aggaaagcac gaaatctttt tcattcctgg aacaacccaa    360
gaagaagctc tttcgaatat tgatcaaata tcggataagg ctgaggaatt gaaacgatca    420
acttcagatg ctgttcagga taacgataac gatgattcga ttcctacaag tgctgaattt    480
ccacctggta tttcgccaac cgagaattgg gtcggaaaagt tgcaggaaaa atctcaaaaa    540
agcaagctgc aagccccaat ctatgaagat tccaagaatg agagaaccga gcgtttcttg    600
ggtatatgca cgatgtgcaa tcaaaaaacc agaggatca gaagtaagaa gaaggacgca    660
aagaatcttg cagcatgggt gatgtggaaa gcgttgaag acggtatcga atctctggaa    720
tcatatgata tggttgatgt gattgaaaat ttggaagaag ctgaacattt actcgaatt    780
caggatcaag catccaagat taaagacaag cattccgcac tgattgatat actctcggac    840
aagaaaagat tttcagacta cagcatggat ttcaacgtat tatcagtga cacaatggga    900
atacatcagg tgctatttga aatctcgttc cggcgtctag tttctccaga ccccagcgat    960
ttggaaatgg gagcagaaca ccccagact gaagaaatta tgaaggctac tgcogagaag    1020
gaaaagctac ggaagaagaa tatgccagat tccgggccgc tagtgtttgc tggacatggt    1080
tcatcggcgg aagagctaa acagtgtgct tgtaaatcgg cgattatcca tttcaacacc    1140
tatgatttca cgattgaaa atattattgc gtattcctga aaaatgaagc gtctgaatga    1200
ttataaaaaa aaaaaaaaaa aa                                           1222
    
```


-continued

```

<210> SEQ ID NO 5
<211> LENGTH: 407
<212> TYPE: PRT
<213> ORGANISM: Caenorhabditis elegans
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)...(407)
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 5
Met Asp Leu Thr Lys Leu Thr Phe Glu Ser Val Phe Gly Gly Ser Asp
 1          5          10          15
Val Pro Met Lys Pro Ser Arg Ser Glu Asp Asn Lys Thr Pro Arg Asn
          20          25          30
Arg Thr Asp Leu Glu Met Phe Leu Lys Lys Thr Pro Leu Met Val Leu
          35          40          45
Glu Glu Ala Ala Lys Ala Val Tyr Gln Lys Thr Pro Thr Trp Gly Thr
          50          55          60
Val Glu Leu Pro Glu Gly Phe Glu Met Thr Leu Ile Leu Asn Glu Ile
          65          70          75          80
Thr Val Lys Gly Gln Ala Thr Ser Lys Lys Ala Ala Arg Gln Lys Ala
          85          90          95
Ala Val Glu Tyr Leu Arg Lys Val Val Glu Lys Gly Lys His Glu Ile
          100         105         110
Phe Phe Ile Pro Gly Thr Thr Lys Glu Glu Ala Leu Ser Asn Ile Asp
          115         120         125
Gln Ile Ser Asp Lys Ala Glu Glu Leu Lys Arg Ser Thr Ser Asp Ala
          130         135         140
Val Gln Asp Asn Asp Asn Asp Asp Ser Ile Pro Thr Ser Ala Glu Phe
          145         150         155         160
Pro Pro Gly Ile Ser Pro Thr Glu Asn Trp Val Gly Lys Leu Gln Glu
          165         170         175
Lys Ser Gln Lys Ser Lys Leu Gln Ala Pro Ile Tyr Glu Asp Ser Lys
          180         185         190
Asn Glu Arg Thr Glu Arg Phe Leu Val Ile Cys Thr Met Cys Asn Gln
          195         200         205
Lys Thr Arg Gly Ile Arg Ser Lys Lys Lys Asp Ala Lys Asn Leu Ala
          210         215         220
Ala Trp Leu Met Trp Lys Ala Leu Glu Asp Gly Ile Glu Ser Leu Glu
          225         230         235         240
Ser Tyr Asp Met Val Asp Val Ile Glu Asn Leu Glu Glu Ala Glu His
          245         250         255
Leu Leu Glu Ile Gln Asp Gln Ala Ser Lys Ile Lys Asp Lys His Ser
          260         265         270
Ala Leu Ile Asp Ile Leu Ser Asp Lys Lys Arg Phe Ser Asp Tyr Ser
          275         280         285
Met Asp Phe Asn Val Leu Ser Val Ser Thr Met Gly Ile His Gln Val
          290         295         300
Leu Leu Glu Ile Ser Phe Arg Arg Leu Val Ser Pro Asp Pro Asp Asp
          305         310         315         320
Leu Glu Met Gly Ala Glu His Thr Gln Thr Glu Glu Ile Met Lys Ala
          325         330         335
Thr Ala Glu Lys Glu Lys Leu Arg Lys Lys Asn Met Pro Asp Ser Gly

```

-continued

```

          340          345          350
Pro Leu Val Phe Ala Gly His Gly Ser Ser Ala Glu Glu Ala Lys Gln
   355          360          365

Cys Ala Cys Lys Ser Ala Ile Ile His Phe Asn Thr Tyr Asp Phe Thr
   370          375          380

Asp Xaa Lys Tyr Tyr Cys Val Phe Leu Lys Asn Glu Ala Ser Glu Xaa
   385          390          395          400

Leu Xaa Lys Lys Lys Lys Lys
   405

<210> SEQ ID NO 6
<211> LENGTH: 763
<212> TYPE: PRT
<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 6

Gly Ile Ile Asn Gly Pro Lys Arg Glu Arg Ser Tyr Lys Val Ala Ile
  1          5          10          15

Lys Phe Val Ala Arg Ala Asn Met His His Leu Gly Glu Phe Leu Ala
  20          25          30

Gly Lys Arg Ala Asp Cys Pro Gln Glu Ala Val Gln Ile Leu Asp Ile
  35          40          45

Val Leu Arg Glu Leu Ser Val Lys Arg Phe Cys Pro Val Gly Arg Ser
  50          55          60

Phe Phe Ser Pro Asp Ile Lys Thr Pro Gln Arg Leu Gly Glu Gly Leu
  65          70          75          80

Glu Ser Trp Cys Gly Phe Tyr Gln Ser Ile Arg Pro Thr Gln Met Gly
  85          90          95

Leu Ser Leu Asn Ile Asp Met Ala Ser Ala Ala Phe Ile Glu Pro Leu
  100         105         110

Pro Val Ile Glu Phe Val Ala Gln Leu Leu Gly Lys Asp Val Leu Ser
  115         120         125

Lys Pro Leu Ser Asp Ser Asp Arg Val Lys Ile Lys Lys Gly Leu Arg
  130         135         140

Gly Val Lys Val Glu Val Thr His Arg Ala Asn Val Arg Arg Lys Tyr
  145         150         155         160

Arg Val Ala Gly Leu Thr Thr Gln Pro Thr Arg Glu Leu Met Phe Pro
  165         170         175

Val Asp Glu Asn Cys Thr Met Lys Ser Val Ile Glu Tyr Phe Gln Glu
  180         185         190

Met Tyr Gly Phe Thr Ile Gln His Thr His Leu Pro Cys Leu Gln Val
  195         200         205

Gly Asn Gln Lys Lys Ala Ser Tyr Leu Pro Met Glu Ala Cys Lys Ile
  210         215         220

Val Glu Gly Gln Arg Tyr Thr Lys Arg Leu Asn Glu Lys Gln Ile Thr
  225         230         235         240

Ala Leu Leu Lys Val Thr Cys Gln Arg Ala Glu Gly Gln Arg Asn Asp
  245         250         255

Ile Leu Arg Thr Val Gln His Asn Ala Tyr Asp Gln Asp Pro Tyr Ala
  260         265         270

Lys Glu Phe Gly Met Asn Ile Ser Glu Lys Leu Ala Ser Val Glu Ala
  275         280         285

```

-continued

Arg Ile Leu Pro Ala Pro Trp Leu Lys Tyr His Glu Asn Gly Lys Glu
 290 295 300
 Lys Asp Cys Leu Pro Gln Val Gly Gln Trp Asn Met Met Asn Lys Lys
 305 310 315 320
 Met Ile Asn Gly Met Thr Val Ser Arg Trp Ala Cys Val Asn Phe Ser
 325 330 335
 Arg Ser Val Gln Glu Asn Val Ala Arg Gly Phe Cys Asn Glu Leu Gly
 340 345 350
 Gln Met Cys Glu Val Ser Gly Met Glu Phe Asn Pro Glu Pro Val Ile
 355 360 365
 Pro Ile Tyr Ser Ala Arg Pro Asp Gln Val Glu Lys Ala Leu Lys His
 370 375 380
 Val Tyr His Thr Ser Met Asn Lys Thr Lys Gly Lys Glu Leu Glu Leu
 385 390 395 400
 Leu Leu Ala Ile Leu Pro Asp Asn Asn Gly Ser Leu Tyr Gly Asp Leu
 405 410 415
 Lys Arg Ile Cys Glu Thr Glu Leu Gly Leu Ile Ser Gln Cys Cys Leu
 420 425 430
 Thr Lys His Val Phe Lys Ile Ser Lys Gln Tyr Leu Ala Asp Val Ser
 435 440 445
 Leu Lys Ile Asn Val Lys Met Gly Gly Arg Asn Thr Val Leu Val Asp
 450 455 460
 Ala Ile Ser Cys Arg Ile Pro Leu Val Ser Asp Ile Pro Thr Ile Ile
 465 470 475 480
 Phe Gly Ala Asp Val Thr His Pro Glu Asn Gly Glu Glu Ser Ser Pro
 485 490 495
 Ser Ile Ala Ala Val Val Ala Ser Gln Asp Trp Pro Glu Val Thr Lys
 500 505 510
 Tyr Ala Gly Leu Val Cys Ala Gln Ala His Arg Gln Glu Leu Ile Gln
 515 520 525
 Asp Leu Tyr Lys Thr Trp Gln Asp Pro Val Arg Gly Thr Val Ser Gly
 530 535 540
 Gly Met Ile Arg Asp Leu Leu Ile Ser Phe Arg Lys Ala Thr Gly Gln
 545 550 555 560
 Lys Pro Leu Arg Ile Ile Phe Tyr Arg Asp Gly Val Ser Glu Gly Gln
 565 570 575
 Phe Tyr Gln Val Leu Leu Tyr Glu Leu Asp Ala Ile Arg Lys Ala Cys
 580 585 590
 Ala Ser Leu Glu Pro Asn Tyr Gln Pro Pro Val Thr Phe Ile Val Val
 595 600 605
 Gln Lys Arg His His Thr Arg Leu Phe Ala Asn Asn His Arg Asp Lys
 610 615 620
 Asn Ser Thr Asp Arg Ser Gly Asn Ile Leu Pro Gly Thr Val Val Asp
 625 630 635 640
 Thr Lys Ile Cys His Pro Thr Glu Phe Asp Phe Tyr Leu Cys Ser His
 645 650 655
 Ala Gly Ile Gln Gly Thr Ser Arg Pro Ala His Tyr His Val Leu Trp
 660 665 670
 Asp Glu Asn Asn Phe Thr Ala Asp Gly Ile Gln Ser Leu Thr Asn Asn
 675 680 685
 Leu Cys Tyr Thr Tyr Ala Arg Cys Thr Arg Ser Val Ser Ile Val Pro

-continued

Trp Thr Asn Glu Phe Arg Thr Cys Ser Met Phe Lys Asn Val His Ile
 290 295 300
 Asn Arg Trp Tyr Val Ile Thr Pro Ser Arg Asn Leu Arg Glu Thr Gln
 305 310 315 320
 Glu Phe Val Gln Met Cys Ile Arg Thr Ala Ser Ser Met Lys Met Asn
 325 330 335
 Ile Cys Asn Pro Ile Tyr Glu Glu Ile Pro Asp Asp Arg Asn Gly Thr
 340 345 350
 Tyr Ser Gln Ala Ile Asp Asn Ala Ala Ala Asn Asp Pro Gln Ile Val
 355 360 365
 Met Val Val Met Arg Ser Pro Asn Glu Glu Lys Tyr Ser Cys Ile Lys
 370 375 380
 Lys Arg Thr Cys Val Asp Arg Pro Val Pro Ser Gln Val Val Thr Leu
 385 390 395 400
 Lys Val Ile Ala Pro Arg Gln Gln Lys Pro Thr Gly Leu Met Ser Ile
 405 410 415
 Ala Thr Lys Val Val Ile Gln Met Asn Ala Lys Leu Met Gly Ala Pro
 420 425 430
 Trp Gln Val Val Ile Pro Leu His Gly Leu Met Thr Val Gly Phe Asp
 435 440 445
 Val Cys His Ser Pro Lys Asn Lys Asn Lys Ser Tyr Gly Ala Phe Val
 450 455 460
 Ala Thr Met Asp Gln Lys Glu Ser Phe Arg Tyr Phe Ser Thr Val Asn
 465 470 475 480
 Glu His Ile Lys Gly Gln Glu Leu Ser Glu Gln Met Ser Val Asn Met
 485 490 495
 Ala Cys Ala Leu Arg Ser Tyr Gln Glu Gln His Arg Ser Leu Pro Glu
 500 505 510
 Arg Ile Leu Phe Phe Arg Asp Gly Val Gly Asp Gly Gln Leu Tyr Gln
 515 520 525
 Val Val Asn Ser Glu Val Asn Thr Leu Lys Asp Arg Leu Asp Glu Ile
 530 535 540
 Tyr Lys Ser Ala Gly Lys Gln Glu Gly Cys Arg Met Thr Phe Ile Ile
 545 550 555 560
 Val Ser Lys Arg Ile Asn Ser Arg Tyr Phe Thr Gly His Arg Asn Pro
 565 570 575
 Val Pro Gly Thr Val Val Asp Asp Val Ile Thr Leu Pro Glu Arg Tyr
 580 585 590
 Asp Phe Phe Leu Val Ser Gln Ala Val Arg Ile Gly Thr Val Ser Pro
 595 600 605
 Thr Ser Tyr Asn Val Ile Ser Asp Asn Met Gly Leu Asn Ala Asp Lys
 610 615 620
 Leu Gln Met Leu Ser Tyr Lys Met Thr His Met Tyr Tyr Asn Tyr Ser
 625 630 635 640
 Gly Thr Ile Arg Val Pro Ala Val Cys His Tyr Ala His Lys Leu Ala
 645 650 655
 Phe Leu Val Ala Glu Ser Ile Asn Arg Ala Pro Ser Ala Gly Leu Gln
 660 665 670
 Asn Gln Leu Tyr Phe Leu
 675

-continued

```

<210> SEQ ID NO 8
<211> LENGTH: 69
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Consensus sequence
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: (1)...(69)
<223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 8
Pro Xaa Xaa Xaa Leu Xaa Glu Xaa Xaa Xaa Gln Xaa Xaa Xaa Xaa Xaa
 1           5           10           15
Xaa Xaa Xaa Tyr Xaa Xaa Xaa Xaa Xaa Xaa Gly Pro Xaa His Xaa Xaa
 20           25           30
Xaa Phe Xaa Xaa Xaa Val Xaa Xaa Xaa Gly Xaa Xaa Xaa Xaa Xaa Gly
 35           40           45
Xaa Gly Xaa Ser Lys Lys Xaa Xaa Ala Lys Xaa Xaa Ala Ala Xaa Xaa
 50           55           60
Ala Leu Xaa Xaa Leu
 65

<210> SEQ ID NO 9
<211> LENGTH: 766
<212> TYPE: PRT
<213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 9
Ser Ala Val Glu Arg Gln Phe Ser Val Ser Leu Lys Trp Val Gly Gln
 1           5           10           15
Val Ser Leu Ser Thr Leu Glu Asp Ala Met Glu Gly Arg Val Arg Gln
 20           25           30
Val Pro Phe Glu Ala Val Gln Ala Met Asp Val Ile Leu Arg His Leu
 35           40           45
Pro Ser Leu Lys Tyr Thr Pro Val Gly Arg Ser Phe Phe Ser Pro Pro
 50           55           60
Val Pro Asn Ala Ser Gly Val Met Ala Gly Ser Cys Pro Pro Gln Ala
 65           70           75           80
Ser Gly Ala Val Ala Gly Gly Ala His Ser Ala Gly Gln Tyr His Ala
 85           90           95
Glu Ser Lys Leu Gly Gly Gly Arg Glu Val Trp Phe Gly Phe His Gln
 100          105          110
Ser Val Arg Pro Ser Gln Trp Lys Met Met Leu Asn Ile Asp Val Ser
 115          120          125
Ala Thr Ala Phe Tyr Arg Ser Met Pro Val Ile Glu Phe Ile Ala Glu
 130          135          140
Val Leu Glu Leu Pro Val Gln Ala Leu Ala Glu Arg Arg Ala Leu Ser
 145          150          155          160
Asp Ala Gln Arg Val Lys Phe Thr Lys Glu Ile Arg Gly Leu Lys Ile
 165          170          175
Glu Ile Thr His Cys Gly Gln Met Arg Arg Lys Tyr Arg Val Cys Asn
 180          185          190
Val Thr Arg Arg Pro Ala Gln Thr Gln Thr Phe Pro Leu Gln Leu Glu
 195          200          205
Thr Gly Gln Thr Ile Glu Cys Thr Val Ala Lys Tyr Phe Tyr Asp Lys

```

-continued

210					215					220					
Tyr	Arg	Ile	Gln	Leu	Lys	Tyr	Pro	His	Leu	Pro	Cys	Leu	Gln	Val	Gly
225					230					235					240
Gln	Glu	Gln	Lys	His	Thr	Tyr	Leu	Pro	Pro	Glu	Val	Cys	Asn	Ile	Val
				245					250					255	
Pro	Gly	Gln	Arg	Cys	Ile	Lys	Lys	Leu	Thr	Asp	Val	Gln	Thr	Ser	Thr
			260					265						270	
Met	Ile	Lys	Ala	Thr	Ala	Arg	Ser	Ala	Pro	Glu	Arg	Glu	Arg	Glu	Ile
		275					280					285			
Ser	Asn	Leu	Val	Arg	Lys	Ala	Glu	Phe	Ser	Ala	Asp	Pro	Phe	Ala	His
	290					295					300				
Glu	Phe	Gly	Ile	Thr	Ile	Asn	Pro	Ala	Met	Thr	Glu	Val	Lys	Gly	Arg
305					310					315					320
Val	Leu	Ser	Ala	Pro	Lys	Leu	Leu	Tyr	Gly	Gly	Arg	Thr	Arg	Ala	Thr
				325					330					335	
Ala	Leu	Pro	Asn	Gln	Gly	Val	Trp	Asp	Met	Arg	Gly	Lys	Gln	Phe	His
			340					345						350	
Thr	Gly	Ile	Asp	Val	Arg	Val	Trp	Ala	Ile	Ala	Cys	Phe	Ala	Gln	Gln
		355					360					365			
Gln	His	Val	Lys	Glu	Asn	Asp	Leu	Arg	Met	Phe	Thr	Asn	Gln	Leu	Gln
	370					375					380				
Arg	Ile	Ser	Asn	Asp	Ala	Gly	Met	Pro	Ile	Val	Gly	Asn	Pro	Cys	Phe
385					390					395					400
Cys	Lys	Tyr	Ala	Val	Gly	Val	Glu	Gln	Val	Glu	Pro	Met	Phe	Lys	Tyr
				405					410					415	
Leu	Lys	Gln	Asn	Tyr	Ser	Gly	Ile	Gln	Leu	Val	Val	Val	Val	Leu	Pro
			420					425						430	
Gly	Lys	Thr	Pro	Val	Tyr	Ala	Glu	Val	Lys	Arg	Val	Gly	Asp	Thr	Val
		435					440					445			
Leu	Gly	Ile	Ala	Thr	Gln	Cys	Val	Gln	Ala	Lys	Asn	Ala	Ile	Arg	Thr
	450					455					460				
Thr	Pro	Gln	Thr	Leu	Ser	Asn	Leu	Cys	Leu	Lys	Met	Asn	Val	Lys	Leu
465					470					475					480
Gly	Gly	Val	Asn	Ser	Ile	Leu	Leu	Pro	Asn	Val	Arg	Pro	Arg	Ile	Phe
				485					490					495	
Asn	Glu	Pro	Val	Ile	Phe	Phe	Gly	Cys	Asp	Ile	Thr	His	Pro	Pro	Ala
			500					505					510		
Gly	Asp	Ser	Arg	Lys	Pro	Ser	Ile	Ala	Ala	Val	Val	Gly	Ser	Met	Asp
		515					520					525			
Ala	His	Pro	Ser	Arg	Tyr	Ala	Ala	Thr	Val	Arg	Val	Gln	Gln	His	Arg
	530					535					540				
Gln	Glu	Ile	Ile	Ser	Asp	Leu	Thr	Tyr	Met	Val	Arg	Glu	Leu	Leu	Val
545					550					555					560
Gln	Phe	Tyr	Arg	Asn	Thr	Arg	Phe	Lys	Pro	Ala	Arg	Ile	Val	Val	Tyr
				565					570					575	
Arg	Asp	Gly	Val	Ser	Glu	Gly	Gln	Phe	Phe	Asn	Val	Leu	Gln	Tyr	Glu
			580					585					590		
Leu	Arg	Ala	Ile	Arg	Glu	Ala	Cys	Met	Met	Leu	Glu	Arg	Gly	Tyr	Gln
		595					600					605			
Pro	Gly	Ile	Thr	Phe	Ile	Ala	Val	Gln	Lys	Arg	His	His	Thr	Arg	Leu
	610					615					620				

-continued

```

Phe Ala Val Asp Lys Lys Asp Gln Val Gly Lys Ala Tyr Asn Ile Pro
625          630          635          640
Pro Gly Thr Thr Val Asp Val Gly Ile Thr His Pro Thr Glu Phe Asp
          645          650          655
Phe Tyr Leu Cys Ser His Ala Gly Ile Gln Gly Thr Ser Arg Pro Ser
          660          665          670
His Tyr His Val Leu Trp Asp Asp Asn Asn Leu Thr Ala Asp Glu Leu
          675          680          685
Gln Gln Leu Thr Tyr Gln Met Cys His Thr Tyr Val Arg Cys Thr Arg
          690          695          700
Ser Val Ser Ile Pro Ala Pro Ala Tyr Tyr Ala His Leu Val Ala Phe
705          710          715          720
Arg Ala Arg Tyr His Leu Val Asp Arg Glu His Asp Ser Gly Glu Gly
          725          730          735
Ser Gln Pro Ser Gly Thr Ser Glu Asp Thr Thr Leu Ser Asn Met Ala
          740          745          750
Arg Ala Val Gln Val Ile Leu Ala Phe Asn Leu Val Ser Ile
          755          760          765

```

```

<210> SEQ ID NO 10
<211> LENGTH: 737
<212> TYPE: PRT
<213> ORGANISM: Oryctolagus cuniculus

```

```

<400> SEQUENCE: 10

```

```

Gly Lys Asp Arg Ile Phe Lys Val Ser Ile Lys Trp Val Ser Cys Val
  1          5          10          15
Ser Leu Gln Ala Leu His Asp Ala Leu Ser Gly Arg Leu Pro Ser Val
          20          25          30
Pro Phe Glu Thr Ile Gln Ala Leu Asp Val Val Met Arg His Leu Pro
          35          40          45
Ser Met Arg Tyr Thr Pro Val Gly Arg Ser Phe Phe Thr Ala Ser Glu
          50          55          60
Gly Cys Ser Asn Pro Leu Gly Gly Gly Arg Glu Val Trp Phe Gly Phe
          65          70          75          80
His Gln Ser Val Arg Pro Ser Leu Trp Lys Met Met Leu Asn Ile Asp
          85          90          95
Val Ser Ala Thr Ala Phe Tyr Lys Ala Gln Pro Val Ile Glu Phe Val
          100          105          110
Cys Glu Val Leu Asp Phe Lys Ser Ile Glu Glu Gln Gln Lys Pro Leu
          115          120          125
Thr Asp Ser Gln Arg Val Lys Phe Thr Lys Glu Ile Lys Gly Leu Lys
          130          135          140
Val Glu Ile Thr His Cys Gly Gln Met Lys Arg Lys Tyr Arg Val Cys
          145          150          155          160
Asn Val Thr Arg Arg Pro Ala Ser His Gln Thr Phe Pro Leu Gln Gln
          165          170          175
Glu Ser Gly Gln Thr Val Glu Cys Thr Val Ala Gln Tyr Phe Lys Asp
          180          185          190
Arg His Lys Leu Val Leu Arg Tyr Pro His Leu Pro Cys Leu Gln Val
          195          200          205
Gly Gln Glu Gln Lys His Thr Tyr Leu Pro Leu Glu Val Cys Asn Ile

```


-continued

210		215		220											
Val	Ala	Gly	Gln	Arg	Cys	Ile	Lys	Lys	Leu	Thr	Asp	Asn	Gln	Thr	Ser
225					230					235					240
Thr	Met	Ile	Arg	Ala	Thr	Ala	Arg	Ser	Ala	Pro	Asp	Arg	Gln	Glu	Glu
				245					250					255	
Ile	Ser	Lys	Leu	Met	Arg	Ser	Ala	Ser	Phe	Asn	Thr	Asp	Pro	Tyr	Val
			260					265					270		
Arg	Glu	Phe	Gly	Ile	Met	Val	Lys	Asp	Glu	Met	Thr	Asp	Val	Thr	Gly
		275					280					285			
Arg	Val	Leu	Gln	Pro	Pro	Ser	Ile	Leu	Tyr	Gly	Gly	Arg	Asn	Lys	Ala
	290					295					300				
Ile	Ala	Thr	Pro	Val	Gln	Gly	Val	Trp	Asp	Met	Arg	Asn	Lys	Gln	Phe
305					310					315					320
His	Thr	Gly	Ile	Glu	Ile	Lys	Val	Trp	Ala	Ile	Ala	Cys	Phe	Ala	Pro
				325					330					335	
Gln	Arg	Gln	Cys	Thr	Glu	Val	His	Leu	Lys	Ser	Phe	Thr	Glu	Gln	Leu
			340					345					350		
Arg	Lys	Ile	Ser	Arg	Asp	Ala	Gly	Met	Pro	Ile	Gln	Gly	Gln	Pro	Cys
		355					360					365			
Phe	Cys	Lys	Tyr	Ala	Gln	Gly	Ala	Asp	Ser	Val	Gly	Pro	Met	Phe	Arg
	370					375					380				
His	Leu	Lys	Asn	Thr	Tyr	Ala	Gly	Leu	Gln	Leu	Val	Val	Val	Ile	Leu
385					390					395					400
Pro	Gly	Lys	Thr	Pro	Val	Tyr	Ala	Glu	Val	Lys	Arg	Val	Gly	Asp	Thr
				405					410					415	
Val	Leu	Gly	Met	Ala	Thr	Gln	Cys	Val	Gln	Met	Lys	Asn	Val	Gln	Arg
			420					425					430		
Thr	Thr	Pro	Gln	Thr	Leu	Ser	Asn	Leu	Cys	Leu	Lys	Ile	Asn	Val	Lys
		435					440					445			
Leu	Gly	Gly	Val	Asn	Asn	Ile	Leu	Leu	Pro	Gln	Gly	Arg	Pro	Pro	Val
	450					455					460				
Phe	Gln	Gln	Pro	Val	Ile	Phe	Leu	Gly	Ala	Asp	Val	Thr	His	Pro	Pro
465					470					475					480
Ala	Gly	Asp	Gly	Lys	Lys	Pro	Ser	Ile	Ala	Ala	Val	Val	Gly	Ser	Met
				485					490					495	
Asp	Ala	His	Pro	Asn	Arg	Tyr	Cys	Ala	Thr	Val	Arg	Val	Gln	Gln	His
			500					505					510		
Arg	Gln	Glu	Ile	Ile	Gln	Asp	Leu	Ala	Ala	Met	Val	Arg	Glu	Leu	Leu
		515					520					525			
Ile	Gln	Phe	Tyr	Lys	Ser	Thr	Arg	Phe	Lys	Pro	Thr	Arg	Ile	Ile	Phe
	530					535					540				
Tyr	Arg	Asp	Gly	Val	Ser	Glu	Gly	Gln	Phe	Gln	Gln	Val	Leu	His	His
545					550					555					560
Glu	Leu	Leu	Ala	Ile	Arg	Glu	Ala	Cys	Ile	Lys	Leu	Glu	Lys	Asp	Tyr
				565					570					575	
Gln	Pro	Gly	Ile	Thr	Phe	Ile	Val	Val	Gln	Lys	Arg	His	His	Thr	Arg
			580					585					590		
Leu	Phe	Cys	Thr	Asp	Lys	Asn	Glu	Arg	Val	Gly	Lys	Ser	Gly	Asn	Ile
		595					600					605			
Pro	Ala	Gly	Thr	Thr	Val	Asp	Thr	Lys	Ile	Thr	His	Pro	Thr	Glu	Phe
	610					615					620				

-continued

```

Asp Phe Tyr Leu Cys Ser His Ala Gly Ile Gln Gly Thr Ser Arg Pro
625                630                635                640
Ser His Tyr His Val Leu Trp Asp Asp Asn Arg Phe Ser Ser Asp Glu
645                650                655
Leu Gln Ile Leu Thr Tyr Gln Leu Cys His Thr Tyr Val Arg Cys Thr
660                665                670
Arg Ser Val Ser Ile Pro Ala Pro Ala Tyr Tyr Ala His Leu Val Ala
675                680                685
Phe Arg Ala Arg Tyr His Leu Val Asp Lys Glu His Asp Ser Ala Glu
690                695                700
Gly Ser His Thr Ser Gly Gln Ser Asn Gly Arg Asp His Gln Ala Leu
705                710                715                720
Ala Lys Ala Val Gln Val His Gln Asp Thr Leu Arg Thr Met Tyr Phe
725                730                735

```

Ala

```

<210> SEQ ID NO 11
<211> LENGTH: 66
<212> TYPE: PRT
<213> ORGANISM: Xenopus laevis

```

<400> SEQUENCE: 11

```

Pro Val Gly Ser Leu Gln Glu Leu Ala Val Gln Lys Gly Trp Arg Leu
  1                5                10                15
Pro Glu Tyr Thr Val Ala Gln Glu Ser Gly Pro Pro His Lys Arg Glu
                20                25                30
Phe Thr Ile Thr Cys Arg Val Glu Thr Phe Val Glu Thr Gly Ser Gly
  35                40                45
Thr Ser Lys Gln Val Ala Lys Arg Val Ala Ala Glu Lys Leu Leu Thr
  50                55                60
Lys Phe
  65

```

```

<210> SEQ ID NO 12
<211> LENGTH: 66
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

<400> SEQUENCE: 12

```

Phe Met Glu Glu Leu Asn Thr Tyr Arg Gln Lys Gln Gly Val Val Leu
  1                5                10                15
Lys Tyr Gln Glu Leu Pro Asn Ser Gly Pro Pro His Asp Arg Arg Phe
                20                25                30
Thr Phe Gln Val Ile Ile Asp Gly Arg Glu Phe Pro Glu Gly Glu Gly
  35                40                45
Arg Ser Lys Lys Glu Ala Lys Asn Ala Ala Ala Lys Leu Ala Val Glu
  50                55                60
Ile Leu
  65

```

```

<210> SEQ ID NO 13
<211> LENGTH: 818
<212> TYPE: PRT
<213> ORGANISM: Caenorhabditis elegans

```

-continued

<400> SEQUENCE: 13

```

Val Asn Glu Glu Ile Lys Val Gln Phe Ala Lys Asn Phe Val Tyr Asp
  1          5          10          15
Asn Asn Ser Ile Leu Arg Val Pro Glu Ser Phe His Asp Pro Asn Arg
  20          25          30
Phe Glu Gln Ser Leu Glu Val Ala Pro Arg Ile Glu Ala Trp Phe Gly
  35          40          45
Ile Tyr Ile Gly Ile Lys Glu Leu Phe Asp Gly Glu Pro Val Leu Asn
  50          55          60
Phe Ala Ile Val Asp Lys Leu Phe Tyr Asn Ala Pro Lys Met Ser Leu
  65          70          75          80
Leu Asp Tyr Leu Leu Leu Ile Val Asp Pro Gln Ser Cys Asn Asp Asp
  85          90          95
Val Arg Lys Asp Leu Lys Thr Lys Leu Met Ala Gly Lys Met Thr Ile
  100         105         110
Arg Gln Ala Ala Arg Pro Arg Ile Arg Gln Leu Leu Glu Asn Leu Lys
  115         120         125
Leu Lys Cys Ala Glu Val Trp Asp Asn Glu Met Ser Arg Leu Thr Glu
  130         135         140
Arg His Leu Thr Phe Leu Asp Leu Cys Glu Glu Asn Ser Leu Val Tyr
  145         150         155         160
Lys Val Thr Gly Lys Ser Asp Arg Gly Arg Asn Ala Lys Lys Tyr Asp
  165         170         175
Thr Thr Leu Phe Lys Ile Tyr Glu Glu Asn Lys Lys Phe Ile Glu Phe
  180         185         190
Pro His Leu Pro Leu Val Lys Val Lys Ser Gly Ala Lys Glu Tyr Ala
  195         200         205
Val Pro Met Glu His Leu Glu Val His Glu Lys Pro Gln Arg Tyr Lys
  210         215         220
Asn Arg Ile Asp Leu Val Met Gln Asp Lys Phe Leu Lys Arg Ala Thr
  225         230         235         240
Arg Lys Pro His Asp Tyr Lys Glu Asn Thr Leu Lys Met Leu Lys Glu
  245         250         255
Leu Asp Phe Ser Ser Glu Glu Leu Asn Phe Val Glu Arg Phe Gly Leu
  260         265         270
Cys Ser Lys Leu Gln Met Ile Glu Cys Pro Gly Lys Val Leu Lys Glu
  275         280         285
Pro Met Leu Val Asn Ser Val Asn Glu Gln Ile Lys Met Thr Pro Val
  290         295         300
Ile Arg Gly Phe Gln Glu Lys Gln Leu Asn Val Val Pro Glu Lys Glu
  305         310         315         320
Leu Cys Cys Ala Val Phe Val Val Asn Glu Thr Ala Gly Asn Pro Cys
  325         330         335
Leu Glu Glu Asn Asp Val Val Lys Phe Tyr Thr Glu Leu Ile Gly Gly
  340         345         350
Cys Lys Phe Arg Gly Ile Arg Ile Gly Ala Asn Glu Asn Arg Gly Ala
  355         360         365
Gln Ser Ile Met Tyr Asp Ala Thr Lys Asn Glu Tyr Ala Phe Tyr Lys
  370         375         380
Asn Cys Thr Leu Asn Thr Gly Ile Gly Arg Phe Glu Ile Ala Ala Thr
  385         390         395         400

```

-continued

Glu Ala Lys Asn Met Phe Glu Arg Leu Pro Asp Lys Glu Gln Lys Val
 405 410 415
 Leu Met Phe Ile Ile Ile Ser Lys Arg Gln Leu Asn Ala Tyr Gly Phe
 420 425 430
 Val Lys His Tyr Cys Asp His Thr Ile Gly Val Ala Asn Gln His Ile
 435 440 445
 Thr Ser Glu Thr Val Thr Lys Ala Leu Ala Ser Leu Arg His Glu Lys
 450 455 460
 Gly Ser Lys Arg Ile Phe Tyr Gln Ile Ala Leu Lys Ile Asn Ala Lys
 465 470 475 480
 Leu Gly Gly Ile Asn Gln Glu Leu Asp Trp Ser Glu Ile Ala Glu Ile
 485 490 495
 Ser Pro Glu Glu Lys Glu Arg Arg Lys Thr Met Pro Leu Thr Met Tyr
 500 505 510
 Val Gly Ile Asp Val Thr His Pro Thr Ser Tyr Ser Gly Ile Asp Tyr
 515 520 525
 Ser Ile Ala Ala Val Val Ala Ser Ile Asn Pro Gly Gly Thr Ile Tyr
 530 535 540
 Arg Asn Met Ile Val Thr Gln Glu Glu Cys Arg Pro Gly Glu Arg Ala
 545 550 555 560
 Val Ala His Gly Arg Glu Arg Thr Asp Ile Leu Glu Ala Lys Phe Val
 565 570 575
 Lys Leu Leu Arg Glu Phe Ala Glu Asn Asp Asn Arg Ala Pro Ala
 580 585 590
 His Ile Val Val Tyr Arg Asp Gly Val Ser Asp Ser Glu Met Leu Arg
 595 600 605
 Val Ser His Asp Glu Leu Arg Ser Leu Lys Ser Glu Val Lys Gln Phe
 610 615 620
 Met Ser Glu Arg Asp Gly Glu Asp Pro Glu Pro Lys Tyr Thr Phe Ile
 625 630 635 640
 Val Ile Gln Lys Arg His Asn Thr Arg Leu Arg Arg Met Glu Lys
 645 650 655
 Asp Lys Pro Val Val Asn Lys Asp Leu Thr Pro Ala Glu Thr Asp Val
 660 665 670
 Ala Val Ala Ala Val Lys Gln Trp Glu Glu Asp Met Lys Glu Ser Lys
 675 680 685
 Glu Thr Gly Ile Val Asn Pro Ser Ser Gly Thr Thr Val Asp Lys Leu
 690 695 700
 Ile Val Ser Lys Tyr Lys Phe Asp Phe Phe Leu Ala Ser His His Gly
 705 710 715 720
 Val Leu Gly Thr Ser Arg Pro Gly His Tyr Thr Val Met Tyr Asp Asp
 725 730 735
 Lys Gly Met Ser Gln Asp Glu Val Tyr Lys Met Thr Tyr Gly Leu Ala
 740 745 750
 Phe Leu Ser Ala Arg Cys Arg Lys Pro Ile Ser Leu Pro Val Pro Val
 755 760 765
 His Tyr Ala His Leu Ser Cys Glu Lys Ala Lys Glu Leu Tyr Arg Thr
 770 775 780
 Tyr Lys Glu His Tyr Ile Gly Asp Tyr Ala Gln Pro Arg Thr Arg His
 785 790 795 800

-continued

Glu Met Glu His Phe Leu Gln Thr Asn Val Lys Tyr Pro Gly Met Ser
 805 810 815

Phe Ala

<210> SEQ ID NO 14
 <211> LENGTH: 63
 <212> TYPE: PRT
 <213> ORGANISM: Caenorhabditis elegans

<400> SEQUENCE: 14

Trp Val Gly Lys Leu Gln Phe Lys Ser Gln Lys Ser Lys Leu Gln Ala
 1 5 10 15
 Asp Ile Tyr Glu Asp Ser Lys Asn Glu Arg Thr Glu Phe Thr Leu Val
 20 25 30
 Ile Cys Thr Met Cys Asn Gln Lys Thr Arg Gly Ile Thr Ser Lys Gln
 35 40 45
 Lys Asp Ala Lys Asn Leu Ala Ala Trp Leu Met Trp Lys Ala Leu
 50 55 60

What is claimed is:

1-10. (canceled)

11. A substantially pure polypeptide encoded by a nucleic acid molecule which hybridizes under high stringency conditions to a complement of the nucleic acid molecule set forth in SEQ ID NO: 4

12-16. (canceled)

17. The polypeptide of claim 11, which hybridizes under conditions of hybridization at 68° C. in 5×SSC/5× Denhardt solution/1.0% SDS, followed by washing in 0.2×SSC/0.1% SDS at room temperature.

18. A substantially pure polypeptide encoded by a nucleic acid molecule having at least 80% sequence identity with the nucleic acid molecule set forth as SEQ ID No:4.

19. The polypeptide of claim 18, which is encoded by a nucleic acid molecule having at least 95% identity with the nucleic acid molecule set forth as SEQ ID NO:4.

20. The polypeptide of claim 18, which is encoded by a nucleic acid molecule having at least 98% identity with the nucleic acid molecule set forth as SEQ ID NO:4.

21. The polypeptide of claim 11 or **18**, which polypeptide has a dsRNA binding activity.

22. The polypeptide of claim 11 or **18**, which polypeptide has the dsRNA binding motif of SEQ ID NOS:8.

23. A substantially pure polypeptide fragment comprising at least 30 contiguous amino acids of SEQ ID NO:5.

24. A substantially pure protein encoded by the nucleic acid molecule set forth in SEQ ID NO:4.

25. A substantially pure protein comprising the amino acid sequence of SEQ ID NO:5.

26. A fragment of claim 24, wherein said fragment comprises amino acids 150 to 212 of SEQ ID NO:5.

27. A substantially pure RDE-4 polypeptide encoded by a nucleic acid molecule which can complement a RDE-4 mutation.

28. A fusion protein comprising a fragment of the polypeptide of any one of the preceding claims and a heterologous polypeptide.

29. The fusion protein of claim 28, wherein the heterologous polypeptide is selected from the group consisting of an immunoglobulin Fc (IgFc) polypeptide, a lacZ polypeptide, a glutathione S-transferase (GST) polypeptide, a six histidine tag polypeptide and a signal sequence polypeptide.

30. A method of preparing an RNAi agent, the method comprising incubating a dsRNA in the presence of an RDE-4 protein.

* * * * *