(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
30 April 2009 (30.04.2009)

PCT

(10) International Publication Number
WO 2009/055053 A2

(51) International Patent Classification:
C07D 233/86 (2006.01) *A61K 31/4166* (2006.01)
A61P 35/00 (2006.01)

(21) International Application Number:
PCT/US2008/012149

(22) International Filing Date: 24 October 2008 (24.10.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/996,076 26 October 2007 (26.10.2007) US

(71) Applicant *(for all designated States except US)*: **THE REGENTS OF THE UNIVERSITY OF CALIFORNIA** [US/US]; 1111 Franklin Street, Oakland, California 94607-5200 (US).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: **JUNG, Michael E.** [US/US]; UCLA Chem & Biochem, Box 951569, 3505A, Mol. Sci. Bldg., Los Angeles, California 90095-1569 (US). **YOO, Dongwon** [KR/US]; UCLA Chem & Biochem, Box 951569, 3505A, Mol. Sci. Bldg., Los Angeles, California 90095-1569 (US). **SAWYERS, Charles L.** [US/US]; HOPP, MSKCC, 1275 York Avenue, Box 20, New York, New York 10021 (US). **TRAN, Chris** [US/US]; HOPP, MSKCC, 1275 York Avenue, Box 20, New York, New York 10021 (US).

(74) Agents: **GENIESER, Lars H.** et al.; Venable LLP, P.O. Box 34385, Washington, District Of Columbia 20043-9998 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:
— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
— *of inventorship (Rule 4.17(iv))*

Published:
— *without international search report and to be republished upon receipt of that report*

WO 2009/055053 A2

(54) Title: DIARYLHYDANTOIN COMPOUNDS

(57) Abstract: The present invention relates to diarylhydantoin compounds and methods for synthesizing them and using them in the treatment of hormone refractory prostate cancer.

DIARYLHYDANTOIN COMPOUNDS

FIELD OF THE INVENTION

[0001] The present invention relates to diarylhydantoin compounds including diarylthiohydantoins, and methods for synthesizing them and using them in the treatment of hormone refractory prostate cancer. This application claims the benefit of U.S. Provisional Application No. 60/996,076, filed October 26, 2007, the specification of which is hereby incorporated by reference.

BACKGROUND OF THE INVENTION

[0002] Prostate cancer is the most common incidence of cancer and the second leading cause of cancer death in Western men. When the cancer is confined locally, the disease can be cured by surgery or radiation. However, 30% of such cancer relapses with distant metastatic disease and others have advanced disease at diagnoses. Advanced disease is treated by castration and/or administration of antiandrogens, the so-called androgen deprivation therapy. Castration lowers the circulating levels of androgens and reduces the activity of androgen receptor (AR). Administration of antiandrogens blocks AR function by competing away androgen binding, therefore, reducing the AR activity. Although initially effective, these treatments quickly fail and the cancer becomes hormone refractory.

[0003] Nonsteroidal anti-androgens, such as bicalutamide, have been preferred over steroidal compounds for prostate cancer because they are more selective and have fewer side effects. This class of compounds has been described in patents such as U.S. Patent Number 4,097,578, U.S. Pat. No. 5,411,981, U.S. Pat. No. 5,705,654, PCT International Applications WO 97/00071 and WO 00/17163, and U.S. Published Patent Application Number 2004/0009969, all of which are hereby incorporated by reference. Bicalutamide (brand name: Casodex) is the most commonly used anti-androgen. While it has an inhibitory effect on AR in hormone sensitive prostate cancer, it fails to suppress AR when cancer becomes hormone refractory.

[0004] U.S. Patent No. 5,434,176 includes broad claims which encompass a very large number of compounds, but synthetic routes are only presented for a small fraction of these compounds and pharmacological data are only presented for two of them, and one skilled in the

art could not readily envision other specific compounds. U.S. Patent No. 5,434,176 is hereby incorporated by reference.

SUMMARY OF THE INVENTION

[0005] The invention provides a series of compounds having strong antagonistic activities with minimal agonistic activities against androgen receptor (AR). These compounds inhibit the growth of hormone refractory prostate cancer.

[0006] The invention includes a compound having the formula



R_1 and R_2 together can include eight or fewer carbon atoms and can be selected from the group consisting of alkyl, substituted alkyl, and, together with the carbon to which they are linked, a cycloalkyl or substituted cycloalkyl group. R_3 can be hydrogen, cyano, formyl,



R_4 can be hydrogen, F, Cl, Br, or I. R_{11} and R_{12} can be the same or different and are hydrogen or methyl. R_{13} can be hydrogen or $-NR_{14}R_{15}$. R_{14} and R_{15} can be the same or different and are hydrogen or methyl.

[0007] For example, R_1 and R_2 can be independently methyl or, together with the carbon to which they are linked, cyclobutyl or cyclopentyl. For example, R_{11} and R_{12} can be both hydrogen or both methyl. For example, R_{13} can be $-NH(CH_3)$ or $-N(CH_3)_2$. For example, when R_4, R_{11}, and R_{12} are each hydrogen and when R_1 and R_2 together with the carbon to which they

are linked are cyclobutyl, then R_3 can be other than cyano and  , with R_{13} hydrogen, $-NH_2$, $-NH(CH_3)$, or $-N(CH_3)_2$.

[0008] The invention provides a pharmaceutical composition comprising a therapeutically effective amount of a compound according to any of the preceding compounds or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

[0009] The invention encompasses a method for treating a hyperproliferative disorder comprising administering such a pharmaceutical composition to a subject in need of such treatment, thereby treating the hyperproliferative disorder. The hyperproliferative disorder may be hormone refractory prostate cancer. The dosage may be in the range of from about 0.001 mg per kg body weight per day to about 100 mg per kg body weight per day, about 0.01 mg per kg body weight per day to about 100 mg per kg body weight per day, about 0.1 mg per kg body weight per day to about 10 mg per kg body weight per day, or about 1 mg per kg body weight per day.

[0010] The compound may be administered by intravenous injection, by injection into tissue, intraperitoneally, orally, or nasally. The composition may have a form selected from the group consisting of a solution, dispersion, suspension, powder, capsule, tablet, pill, time release capsule, time release tablet, and time release pill.

[0011] The invention provides a method of synthesizing a diaryl compound of formula:



The method includes mixing Compound I



Compound I

with Compound II



Compound II

in a first polar solvent to form a mixture. The method further includes the following: adding a second polar solvent, the same as or different from the first polar solvent, and an aqueous acid to the mixture; refluxing the mixture; cooling the mixture and combining with water; and separating the diaryl compound from the mixture. R_{31} is cyano, carboxy,



(methylcarbamoyl),



(dimethylcarbamoyl), or



(Boc is *t*-butoxycarbonyl).

R_{32} is



or



. R_1 and R_2 together include eight or fewer carbon atoms and are alkyl, substituted alkyl, or, together with the carbon to which they are linked, a cycloalkyl or substituted cycloalkyl group. R_3 is hydrogen, cyano, formyl,



,



,



, or



.

R_4 is hydrogen, F, Cl, Br, or I. R_{11} and R_{12} can be the same or different and are hydrogen or methyl. R_{13} is hydrogen or $-NR_{14}R_{15}$. R_{14} and R_{15} can be the same or different and are hydrogen or methyl.

[0012] The compounds presented are expected to have substantial androgen receptor antagonist activity and no substantial agonist activity on hormone refractory prostate cancer cells.

[0013] The invention encompasses a method comprising providing at least one such compound, measuring inhibition of androgen receptor activity for the compound and determining if the inhibition is above a first predetermined level, measuring stimulation of androgen receptor activity in hormone refractory cancer cells for the compound and determining if the stimulation is below a second predetermined level, and selecting the compound if the inhibition is above the first predetermined level and the stimulation is below the second predetermined level. The predetermined levels may be those of bicalutamide. The step of measuring inhibition may comprise measuring inhibitory concentration (IC50) in an AR response reporter system or a prostate specific antigen secreting system. The step of measuring stimulation may comprise measuring fold induction by increasing concentrations in an AR response reporter system or a prostate specific antigen secreting system. The method of measuring inhibition and/or stimulation may comprise measuring an effect of the compound on tumor growth in an animal.

DETAILED DESCRIPTION

[0014] Embodiments of the invention are discussed in detail below. In describing embodiments, specific terminology is employed for the sake of clarity. However, the invention is not intended to be limited to the specific terminology so selected. A person skilled in the relevant art will recognize that other equivalent parts can be employed and other methods developed without parting from the spirit and scope of the invention. All references cited herein are incorporated by reference as if each had been individually incorporated.

[0015] Recently, overexpression of AR has been identified and validated as a cause of hormone refractory prostate cancer. See Chen, C.D., Welsbie, D.S., Tran, C., Baek, S.H., Chen, R., Vessella, R., Rosenfeld, M.G., and Sawyers, C.L., Molecular determinants of resistance to antiandrogen therapy, Nat. Med., 10: 33-39, 2004, which is hereby incorporated by reference. Overexpression of AR is sufficient to cause progression from hormone sensitive to hormone refractory prostate cancer, suggesting that better AR inhibitors than the current drugs can slow the progression of prostate cancer. It was demonstrated that AR and its ligand binding are necessary for growth of hormone refractory prostate cancer, indicating that AR is still a target for this disease. It was also demonstrated that overexpression of AR converts anti-androgens from antagonists to agonists in hormone refractory prostate cancer (an AR antagonist inhibits AR activity and an AR agonist stimulates AR activity). Data from this work explains why castration and anti-androgens fail to prevent prostate cancer progression and reveals unrecognized properties of hormone refractory prostate cancer.

[0016] Two weaknesses of current antiandrogens are blamed for the failure to prevent prostate cancer progression from the hormone sensitive stage to the hormone refractory disease and to effectively treat hormone refractory prostate cancer. One is their weak antagonistic activities and the other is their strong agonistic activities when AR is overexpressed in hormone refractory prostate cancer. Better AR inhibitors with more potent antagonistic activities and minimal agonistic activities are needed to delay disease progression and to treat the fatal hormone refractory prostate cancer.

[0017] Some new properties of hormone refractory prostate cancer are reported in PCT applications US04/42221 and US05/05529, which are hereby incorporated by reference. PCT

International Application US05/05529 presented a methodology for identifying androgen receptor antagonist and agonist characteristics of compounds.

Synthesis of Diarylhydantoin Compounds

[0018] The invention provides for synthesis of diarylthiohydantoin compounds having the formula



R1 and R2 together can comprise eight or fewer carbon atoms and can be alkyl, substituted alkyl, or, together with the carbon to which they are linked, a cycloalkyl or substituted cycloalkyl group. R3 can be hydrogen, cyano, formyl,



,


,


, or


. R_4 can be hydrogen, F, Cl, Br, and I. R_{11} and R_{12} can be the same or different and can be hydrogen or methyl. R_{13} can be hydrogen or $-NR_{14}R_{15}$. R_{14} and R_{15} can be the same or different and can be hydrogen or methyl.

Definitions

[0019] As used herein, the term "alkyl" denotes branched or unbranched hydrocarbon

chains, preferably having about 1 to about 8 carbons, such as, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, 2-methylpentyl pentyl, hexyl, isohexyl, heptyl, 4,4-dimethyl pentyl, octyl, 2,2,4-trimethylpentyl and the like. "Substituted alkyl" includes an alkyl group optionally substituted with one or more functional groups which may be attached to such chains, such as, hydroxyl, bromo, fluoro, chloro, iodo, mercapto or thio, cyano, alkylthio, heterocyclyl, aryl, heteroaryl, carboxyl, carbalkoyl, alkyl, alkenyl, nitro, amino, alkoxyl, amido, and the like to form alkyl groups such as trifluoro methyl, 3-hydroxyhexyl, 2-carboxypropyl, 2-fluoroethyl, carboxymethyl, cyanobutyl and the like.

[0020] Unless otherwise indicated, the term "cycloalkyl" as employed herein alone or as part of another group includes saturated or partially unsaturated (containing 1 or more double bonds) cyclic hydrocarbon groups containing 1 to 3 rings, including monocyclicalkyl, bicyclicalkyl and tricyclicalkyl, containing a total of 3 to 20 carbons forming the rings, preferably 3 to 10 carbons, forming the ring and which may be fused to 1 or 2 aromatic rings as described for aryl, which include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclodecyl and cyclododecyl, cyclohexenyl. "Substituted cycloalkyl" includes a cycloalkyl group optionally substituted with 1 or more substituents such as halogen, alkyl, alkoxy, hydroxy, aryl, aryloxy, arylalkyl, cycloalkyl, alkylamido, alkanoylamino, oxo, acyl, arylcarbonylamino, amino, nitro, cyano, thiol and/or alkylthio and/or any of the substituents included in the definition of "substituted alkyl." For example,



and the like.

[0021] Unless otherwise indicated, the term "alkenyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons, and more preferably 2 to 8 carbons in the normal chain, which include one or more double bonds in the normal chain, such as vinyl, 2-propenyl, 3-butenyl, 2-butenyl, 4-pentenyl, 3-pentenyl, 2-hexenyl, 3-hexenyl, 2-heptenyl, 3-heptenyl, 4-heptenyl, 3-octenyl, 3-nonenyl, 4-decenyl, 3-undecenyl, 4-dodecenyl, 4,8,12-tetradecatrienyl, and the like. "Substituted alkenyl"

includes an alkenyl group optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."

[0022] Unless otherwise indicated, the term "alkynyl" as used herein by itself or as part of another group refers to straight or branched chain radicals of 2 to 20 carbons, preferably 2 to 12 carbons and more preferably 2 to 8 carbons in the normal chain, which include one or more triple bonds in the normal chain, such as 2-propynyl, 3-butynyl, 2-butynyl, 4-pentynyl, 3-pentynyl, 2-hexynyl, 3-hexynyl, 2-heptynyl, 3-heptynyl, 4-heptynyl, 3-octynyl, 3-nonynyl, 4-decynyl, 3-undecynyl, 4-dodecynyl and the like. "Substituted alkynyl" includes an alkynyl group optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."

[0023] The terms "arylalkyl", "arylalkenyl" and "arylalkynyl" as used alone or as part of another group refer to alkyl, alkenyl and alkynyl groups as described above having an aryl substituent. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, phenethyl, benzhydryl and naphthylmethyl and the like. "Substituted arylalkyl" includes arylalkyl groups wherein the aryl portion is optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."

[0024] The terms "arylalkyl", "arylalkenyl" and "arylalkynyl" as used alone or as part of another group refer to alkyl, alkenyl and alkynyl groups as described above having an aryl substituent. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, phenethyl, benzhydryl and naphthylmethyl and the like. "Substituted arylalkyl" includes arylalkyl groups wherein the aryl portion is optionally substituted with one or more substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl."

[0025] The term "halogen" or "halo" as used herein alone or as part of another group refers to chlorine, bromine, fluorine, and iodine.

[0026] The terms “halogenated alkyl”, “halogenated alkenyl” and “alkynyl” as used herein alone or as part of another group refers to “alkyl”, “alkenyl” and “alkynyl” which are substituted by one or more atoms selected from fluorine, chlorine, bromine, and iodine.

[0027] Unless otherwise indicated, the term "aryl" or "Ar" as employed herein alone or as part of another group refers to monocyclic and polycyclic aromatic groups containing 6 to 10 carbons in the ring portion (such as phenyl or naphthyl including l-naphthyl and 2-naphthyl) and may optionally include one to three additional rings fused to a carbocyclic ring or a heterocyclic ring (such as aryl, cycloalkyl, heteroaryl or cycloheteroalkyl rings).

[0028] "Substituted aryl" includes an aryl group optionally substituted with one or more functional groups, such as halo, haloalkyl, alkyl, haloalkyl, alkoxy, haloalkoxy, alkenyl, trifluoromethyl, trifluoromethoxy, alkynyl, cycloalkyl-alkyl, cycloheteroalkyl, cycloheteroalkylalkyl, aryl, heteroaryl, arylalkyl, aryloxy, aryloxyalkyl, arylalkoxy, alkoxycarbonyl, arylcarbonyl, arylalkenyl, aminocarbonylaryl, arylthio, arylsulfinyl, arylazo, heteroarylalkyl, heteroarylalkenyl, heteroarylheteroaryl, heteroaryloxy, hydroxy, nitro, cyano, amino, substituted amino wherein the amino includes 1 or 2 substituents (which are alkyl, aryl or any of the other aryl compounds mentioned in the definitions), thiol, alkylthio, arylthio, heteroarylthio, arylthioalkyl, alkoxyarylthio, alkylcarbonyl, arylcarbonyl, alkylaminocarbonyl, arylaminocarbonyl, alkoxycarbonyl, aminocarbonyl, alkylcarbonyloxy, arylcarbonyloxy, alkylcarbonylamino, arylcarbonylamino, arylsulfinyl, arylsulfinylalkyl, arylsulfonylamino or arylsulfonaminocarbonyl and/or any of the alkyl substituents set out herein.

[0029] Unless otherwise indicated,, the term “heterocyclic” or “heterocycle”, as used herein, represents an unsubstituted or substituted stable 5- to 10-membered monocyclic ring system which may be saturated or unsaturated, and which consists of carbon atoms and from one to four heteroatoms selected from N, O or S, and wherein the nitrogen and sulfur heteroatoms may optionally be oxidized, and the nitrogen heteroatom may optionally be quaternized. The heterocyclic ring may be attached at any heteroatom or carbon atom which results in the creation of a stable structure. Examples of such heterocyclic groups include, but is not limited to, piperidinyl, piperazinyl, oxopiperazinyl, oxopiperidinyl, oxopyrrolidinyl, oxoazepinyl, azepinyl, pyrrolyl, pyrrolidinyl, furanyl, thienyl, pyrazolyl, pyrazolidinyl, imidazolyl, imidazolinyl, imidazolidinyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, oxazolyl, oxazolidinyl, isooxazolyl, isoxazolidinyl, morpholinyl, thiazolyl, thiazolidinyl, isothiazolyl, thiadiazolyl, tetrahydropyranyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, and oxadiazolyl. The term “heterocyclic aromatic” as used here in alone or as part of another group refers to a 5- or 7-membered aromatic ring which includes 1, 2, 3 or 4 hetero atoms such as

nitrogen, oxygen or sulfur and such rings fused to an aryl, cycloalkyl, heteroaryl or heterocycloalkyl ring (e.g. benzothiophenyl, indolyl), and includes possible N-oxides. "Substituted heteroaryl" includes a heteroaryl group optionally substituted with 1 to 4 substituents, such as the substituents included above in the definition of "substituted alkyl" and "substituted cycloalkyl." Examples of heteroaryl groups include the following:



and the like.

[0030] Materials were obtained from commercial suppliers and were used without further purification. Air or moisture sensitive reactions were conducted under argon atmosphere using oven-dried glassware and standard syringe/septa techniques. The reactions were monitored with a silica gel TLC plate under UV light (254 nm) followed by visualization with a *p*-anisaldehyde or ninhydrin staining solution. Column chromatography was performed on silica gel 60. ^{1}H NMR spectra were measured at 400 MHz in $CDCl_3$ unless stated otherwise and data were reported as follows in ppm (δ) from the internal standard (TMS, 0.0 ppm): chemical shift (multiplicity, integration, coupling constant in Hz.).

Synthesis of ND-1

4-[4-(*t*-Butoxycarbonylamino)phenyl]butanoic acid (100)

[0031] Di-tert-butyl dicarbonate (0.73 g, 3.35 mmol) was added to a solution of 4-(4-aminophenyl)butyric acid (0.5 g, 2.79 mmol) and sodium hydroxide (0.14 g, 3.35 mmol) in tert-butanol (5 mL) and water (5 mL) at 0 °C. The mixture was warmed to room temperature and stirred for 9 h. The mixture was partitioned with diethyl ether (20 mL) and water (20 mL) and then the aqueous layer was acidified to pH 2-3 by 1 N $KHSO_4$ solution. The aqueous mixture extracted with ethyl acetate (3 x 20 mL) and the organic layer was dried over $MgSO_4$,

concentrated to give crude 4-[4-(t-Butoxycarbonylamino)phenyl]butanoic acid **(100)** (0.73 g, 94%) which was used without further purification.



^{1}H NMR δ 7.26 (d, 2H, J = 8.5 Hz), 7.10 (d, 2H, J = 8.5 Hz), 6.48 (br s, 1H), 2.62 (t, 2H, J = 7.5 Hz), 2.33 (t, 2H, J = 7.5 Hz), 1.93 (p, 2H, J = 7.5 Hz).

4-[4-(*t*-Butoxycarbonylamino)phenyl]butanamide (99)

[0032] Thionyl chloride (0.22 mL, 3.01 mmol) was added slowly to a solution of 4-[4-(t-Butoxycarbonylamino)phenyl]butanoic acid **(100)** (0.70 g, 2.51 mmol) in DMF (5 mL) cooled at -5 °C. The mixture was stirred for an additional 1 h at -5 °C. Excess ammonia (freshly distilled from its aqueous solution) was added to the reaction medium. The second mixture was stirred for an additional 1 h. Ethyl acetate (50 mL) was added to the mixture, which was washed with brine (2 x 50 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified by silica gel column chromatography (dichloromethane:acetone, 9:1) to give 4-[4-(*t*-Butoxycarbonylamino)phenyl]butanamide **(99)** (0.57 g, 82%) as a white solid.



^{1}H NMR 6 7.26 δ, 2H, J = 8.4 Hz), 7.09 (d, 2H, J = 8.4 Hz), 6.48 (br s, 1H), 5.47 (br s, 2H), 2.62 (t, 2H, J = 7.4 Hz), 2.20 (t, 2H, J = 7.4 Hz), 1.94 (p, 2H, J = 7.4 Hz), 1.51 (s, 9H).

4-[4-(*t*-Butoxycarbonylamino)phenyl]butanenitrile (98)

[0033] A solution of DMSO (0.13 mL, 1.84 mmol) in dichloromethane (2 mL) was added to a stirred solution of oxalyl chloride (0.12 mL, 1.38 mmol) in dichloromethane (2 mL) at -78 °C. After 15 min, a dichloromethane (1 mL) solution of 2 (0.32 g, 1.15 mmol) was added to the reaction mixture. Stirring was continued for 20 min at -78 °C, and then triethylamine (0.48 mL, 3.45 mmol) was added. After 30 min, the reaction mixture was warmed to room temperature and then reaction was quenched with saturated aq. NH_4Cl solution. The mixture was partitioned with diethyl ether (30 mL) and water (20 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified by silica gel column chromatography (hexane:ethyl acetate, 4:1) to give 4-[4-(*t*-Butoxycarbonylamino)phenyl]butanenitrile **(98)**

(0.22 g, 73%) as a white solid.



^{1}H NMR δ 7.30 (d, 2H, J = 8.4 Hz), 6.10 (d, 2H, J = 8.4 Hz), 6.42 (br s, 1H), 2.73 (t, 2H, J = 7.3 Hz), 2.30 (t, 2H, J = 7.3 Hz), 1.95 (p, 2H, J = 7.3 Hz), 1.52 (s, 9H).

4-(4-Aminophenyl)butanenitrile (97)

[0034] A 0.25 M solution of trifluoroacetic acid in dichloromethane (5 mL, 1.25 mmol) was added to 4-[4-(*t*-Butoxycarbonylamino)phenyl]butanenitrile **(98)** (0.22 g, 0.85 mmol). After 30 min, reaction was quenched with 1 N NaOH solution. The mixture was partitioned with ethyl acetate (30 mL) and water (20 mL). The organic layer was dried over $MgSO_4$, concentrated to give 4-(4-Aminophenyl)butanenitrile **(97)** (0.16 g, 99%) which was used without further purification.



^{1}H NMR δ 6.97 (d, 2H, J = 8.5 Hz), 6.64 (d, 2H, J = 8.5 Hz), 3.59 (br s, 2H), 2.67 (t, 2H, J = 7.3 Hz), 2.29 (t, 2H, J = 7.3 Hz), 1.92 (p, 2H, J = 7.3 Hz).

4-Isothiocyanato-2-trifluoromethylbenzonitrile (96)

[0035] 4-Amino-2-trifluoromethylbenzonitrile (2.23 g, 12 mmol) was added portionwise over 15 min into a well-stirred heterogeneous mixture of thiophosgene (1 mL, 13 mmol) in water (22 mL) at room temperature. Stirring was continued for an additional 1 h. The reaction medium was extracted with chloroform (3 x 15 mL). The combined organic phase was dried over $MgSO_4$ and evaporated to dryness under reduced pressure to yield desired product 4-Isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (2.72 g, 11.9 mmol, 99%) as brownish solid and was used without further purification.



^{1}H NMR δ 7.84 (d, 1H, J = 8.3 Hz), 7.59 (d, 1H, J = 2.1 Hz), 7.49 (dd, 1H, J =8.3, 2.1 Hz).

4-[4-(1-Cyanodimethylamino)phenyl]butanenitrile (95)

[0036] A mixture of 4-(4-Aminophenyl)butanenitrile **(97)** (50 mg, 0.26 mmol), acetone cyanohydrin (0.15 mL, 1.58 mmol) was heated to 80 °C and stirred for 12 h. To the medium was added ethyl acetate (20 mL) and then washed with water (2 x 20 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified by silica gel column chromatography (hexane:ethyl acetate, 1:1) to give 4-[4-(1-Cyanodimethylamino)phenyl]butanenitrile **(95)** (52 mg, 87%) as a white solid.



^{1}H NMR δ 7.07 (d, 2H, J = 8.3 Hz), 6.87 (d, 2H, J = 8.3 Hz), 3.68 (br s, 1H), 2.70 (t, 2H, J = 7.3 Hz), 2.31 (t, 2H, J = 7.3 Hz), 1.94 (p, 2H, J = 7.3 Hz), 1.69 (s, 6H).

4-(3-(4-(3-Cyanopropyl)phenyl)-4,4-dimethyl-5-oxo-2-thioxo-imidazolidin-1-yl)-2-(trifluoromethyl)benzonitrile (94) [ND-1]

[0037] A mixture of 4-Isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (32 mg, 0.14 mmol) and 4-[4-(1-Cyanodimethylamino)phenyl]butanenitrile **(95)** (16 mg, 0.07 mmol) in DMF (1 mL) was heated under microwave irradiation at 80 °C for 6 h. To this mixture was added methanol (10 mL) and aq. 1 N HCl (3 mL). The second mixture was refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture was poured into cold water (20 mL) and extracted with ethyl acetate (30 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified by silica gel column chromatography (hexane:ethyl acetate, 2:1) to give 4-(3-(4-(3-Cyanopropyl)phenyl)-4,4-dimethyl-5-oxo-2-thioxo-imidazolidin-1-yl)-2-(trifluoromethyl) benzonitrile **(94)** [ND-1] (20 mg, 62%) as a white solid.



^{1}H NMR δ 7.98 (d, 1H, J = 8.3 Hz), 7.97 (d, 1H, J = 1.8 Hz), 7.85 (dd, 1H, J = 8.3, 1.8 Hz), 7.37 (d, 2H, J = 8.3 Hz), 7.25 (d, 2H, J = 8.3 Hz), 2.87 (t, 2H, J = 7.0 Hz), 2.40 (t, 2H, J = 7.0 Hz), 2.05 (p, 2H, J = 7.0 Hz), 1.59 (s, 6H).

Synthesis of ND-2

4-[4-(1-Cyanocyclopentyl amino)phenyl]butanenitrile (93)

[0038] A mixture of 4-(4-Aminophenyl)butanenitrile **(97)** (52 mg, 0.27 mmol), cyclopentanone (0.07 mL, 0.55 mmol) and TMSCN (0.05 mL, 0.55 mmol) was heated to 80 °C and stirred for 13 h. To the medium was added ethyl acetate (2 x 20 mL) and then washed with water (2 x 20 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (hexane: ethyl acetate, 1:1) to give 4-[4-(1-Cyanocyclopentyl amino)phenyl]butanenitrile **(93)** (70 mg, quant.) as a white solid.



^{1}H NMR δ 7.06 (d, 2H, J = 8.3 Hz), 6.78 (d, 2H, J = 8.3 Hz), 3.80 (br s, 1H), 2.70 (t, 2H, J = 7.3 Hz), 2.34-2.42 (m, 2H), 2.31 (t, 2H, J = 7.3 Hz), 2.09-2.18 (m, 2H), 1.94 (p, 2H, J = 7.3 Hz), 1.86-1.91 (m, 4H).

4-(1-(4-(3-Cyanopropyl)phenyl)-4-oxo-2-thioxo-1,3-diazaspiro[4.4]non-3-yl)-2-trifluoromethylbenzonitrile (92) [ND-2]

[0039] A mixture of 4-Isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (36 mg, 0.16 mmol) and 4-[4-(1-Cyanocyclopentyl amino)phenyl]butanenitrile **(93)** (20 mg, 0.08 mmol) in DMF (1 mL) was heated under microwave irradiation at 80 °C for 6 h. To this mixture was added methanol (10 mL) and aq. 1 N HCl (3 mL). The second mixture was refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture was poured into cold water (20 mL) and extracted with ethyl acetate (30 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified by silica gel column chromatography (hexane:ethyl acetate, 2:1) to give 4-(1-(4-(3-Cyanopropyl)phenyl)-4-oxo-2-thioxo-1,3-diazaspiro[4.4]non-3-yl)-2-trifluoromethylbenzonitrile **(92)** [ND-2]



(25 mg, 65%) as a white solid. ^{1}H NMR δ 7.98 (d, 1H, J = 1.8 Hz), 7.97 (d, 1H, J = 8.3 Hz), 7.86 (dd, 1H, J = 8.3, 1.8 Hz), 7.37 (d, 2H, J = 8.3 Hz), 7.27 (d, 2H, J = 8.3 Hz), 2.87 (t, 2H, J =

7.3 Hz), 2.40 (t, 2H, J = 7.3 Hz), 2.28-2.35 (m, 2H), 2.14-2.23 (m, 2H), 2.05 (p, 2H, J = 7.3 Hz), 1.85-1.92 (m, 2H), 1.48-1.55 (m, 2H).



Synthesis of ND-14

4-(3-(4-(3-Cyanopropyl)-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-thioxo-imidazolidin-1-yl)-2-(trifluoromethyl) benzonitrile (103)

[0040] 4-(3-(4-(3-Cyanopropyl)-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-thioxo-imidazolidin-1-yl)-2-(trifluoromethyl) benzonitrile **(103)** [ND-14] can be synthesized in a manner similar as to that for synthesizing **(92)** [ND-2]. A mixture of 4-Isothiocyanato-2-

trifluoromethylbenzonitrile **(96)** and 4-(4-(2-cyanopropan-2-ylamino)-2-fluorophenyl)butanenitrile **(101)** in solvent, for example, in DMF, is heated under microwave irradiation at 80 °C for 6 h.



101

To this mixture is added alcohol, e.g., methanol, and acid, e.g., aqueous hydrochloric acid. The second mixture is refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture is poured into cold water and extracted, for example, with ethyl acetate. The organic layer is dried, e.g., dried over $MgSO_4$, concentrated, and the residue is purified, for example, by silica gel column chromatography using hexane:ethyl acetate (2:1), to give 4-(3-(4-(3-Cyanopropyl)-3-fluorophenyl)-4,4-dimethyl-5-oxo-2-thioxo-imidazolidin-1-yl)-2-(trifluoromethyl) benzonitrile **(103)** [ND-14].



103 [ND-14]

Synthesis of ND-3

4-[4-(1-Cyanocyclobutylamino)-phenyl]-butyric acid (91)

[0041] Trimethylsilyl cyanide (0.50 g, 5 mmol) was added dropwise to a mixture of 4-(4-aminophenyl)-butyric acid (0.537 g, 3 mmol), cyclobutanone (0.35 g, 5 mmol) and sodium sulfate (1 g) in 1,4-dioxane (10 ml). The mixture was stirred for 15 hours. After filtration to

eliminate sodium sulfate, the medium was concentrated under vacuum to obtain a brown liquid which was subjected to chromatography (dichloromethane:acetone, 50:50) to yield 4-[4-(1-Cyanocyclobutylamino)-phenyl]-butyric acid **(91)** (0.665 g, 2.58 mmol, 86%) as a yellowish solid.

4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-5-yl]-phenyl}-butyric acid methyl ester (90) [ND-4]

[0042] A mixture of 4-isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (0.547 g, 2.4 mmol) and 4-[4-(1-Cyanocyclobutylamino)-phenyl]-butyric acid **(91)** (0.342 g, 1.5 mmol) in dry DMF (2 ml) was stirred at room temperature for 15 hours. To this mixture were added methanol (10 ml) and HCl aq. (5 ml, 2M). The second mixture was refluxed for 3 h. After being cooled to room temperature, the reaction mixture was poured into cold water (10 ml) and extracted with ethyl acetate (3 × 30 ml). The organic layer was dried over $MgSO_4$, concentrated and chromatographed (dichloromethane) to yield 4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-5-yl]-phenyl}-butyric acid methyl ester **(90)** [ND-4] (0.594 g, 1.18 mmol, 79%) as a white powder.



^{1}H NMR ($CDCl_3$, 400 MHz) δ 1.60-1.70 (m, 1H), 1.98-2.07 (m, 2H), 2.14-2.26 (m, 1H), 2.40 (t, J = 7.4 Hz, 2H), 2.52-2.60 (m, 2H), 2.62-2.68 (m, 2H), 2.74 (t, J = 7.4 Hz, 2H), 3.68 (s, 3H), 7.22 (d, J = 8.2 Hz, 2H), 7.38 (d, J = 8.2 Hz, 2H), 7.86 (dd, J_1 = 8.3 Hz, J_2 = 1.8 Hz, 1H), 7.95 (d, J = 8.3 Hz, 1H), 7.98 (d, J = 1.8 Hz, 1H); ^{13}C NMR ($CDCl_3$, 100 MHz) δ 13.7, 26.1, 31.4, 33.5, 34.8, 51.7, 67.5, 109.9, 114.9, 121.9 (q, J = 272.7 Hz), 127.1 (q, J = 4.7 Hz), 129.7, 130.1, 132.3, 133.0, 133.3 (q, J = 33.2 Hz), 135.2, 137.2, 143.5, 173.8, 175.0, 179.9.

4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]oct-5-yl]-phenyl}-butyric acid (89) [ND-5]

[0043] A mixture of 4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-5-yl]-phenyl}-butyric acid methyl ester **(90)** [ND-4] (0.501 g, 1 mmol) in methanol (10 ml) and solution of sodium hydroxide (10 ml, 2M) was stirred at room temperature for 5 hours. The methanol was evaporated. The residue was adjusted to pH = 5 by HCl aq. (2M)

and then, the medium was extracted with ethyl acetate (3 ×50 ml). The organic layer was dried over $MgSO_4$ and concentrated to dryness to obtain 4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]oct-5-yl]-phenyl}-butyric acid **(89)** [ND-5] (0.482 g, 0.99 mmol, 99%), the structure of which is illustrated in Formula 89.



Formula 89

^{1}H NMR ($CDCl_3$, 400 MHz) δ 1.60-1.70 (m, 1H), 1.98-2.07 (m, 2H), 2.14-2.26 (m, 1H), 2.45 (t, J = 7.3 Hz, 2H), 2.51-2.59 (m, 2H), 2.62-2.68 (m, 2H), 2.77 (t, J = 7.3 Hz, 2H), 7.23 (d, J = 8.1 Hz, 2H), 7.40 (d, J = 8.1 Hz, 2H), 7.85 (dd, J = 8.3, 1.8 Hz, 1H), 7.95 (d, J = 8.3 Hz, 1H), 7.97 (d, J = 1.8 Hz, 1H); ^{13}C NMR ($CDCl_3$, 100 MHz) δ 13.7, 25.9, 31.4, 33.4, 34.7, 67.5, 109.9, 114.9, 121.9 (q, J = 272.6 Hz), 127.1 (q, J = 4.7 Hz), 129.8, 130.1, 132.3, 133.0, 133.4 (q, J = 33.1 Hz), 135.2, 137.2, 143.3, 174.9, 178.9, 179.9.

42-5) 4-{4-[7-(4-Cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]oct-5-yl]-phenyl}-N-methyl-butyramide (88) [ND-6]

[0044] To a suspension of 4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]oct-5-yl]-phenyl}-butyric acid **(89)** (0.097 g, 0.2 mmol) in THF (10 ml) at -5 °C was added thionyl chloride (0.019 ml, 0.26 mmol). The medium was stirred at -5 °C for one hour. Then methylamine was bubbled into the mixture at -5 °C for 30 minutes. The medium was filtered. The filtrate was concentrated and chromatographed (dichloromethane:acetone, 75:25) to yield 4-{4-[7-(4-Cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]oct-5-yl]-phenyl}-N-methyl-butyramide **(88)** [ND-6] (0.095 g, 0.19 mmol, 95%) as an off-white powder.



^{1}H NMR ($CDCl_3$, 400 MHz) δ 1.52-1.64 (m, 1H), 1.94-2.01 (m, 2H), 2.10-2.17 (m, 1H), 2.20 (t, J = 7.3 Hz, 2H), 2.46-2.62 (m, 4H), 2.69 (t, J = 7.3 Hz, 2H), 2.73 (d, J = 4.7 Hz, 3H), 6.09 (bs, 1H), 7.16 (d, J = 8.2 Hz, 2H), 7.33 (d, J = 8.2 Hz, 2H), 7.82 (dd, J_1 = 8.3 Hz, J_2 = 1.8 Hz, 1H),

7.91 (d, J = 8.3 Hz, 1H), 7.94 (d, J = 1.8 Hz, 1H); ^{13}C NMR ($CDCl_3$, 100 MHz) δ 13.7, 26.2, 26.8, 31.4, 35.0, 35.7, 67.5, 109.7, 114.9, 121.9 (q, J = 272.7 Hz), 127.1 (q, J = 4.7 Hz), 129.7, 130.0, 132.3, 133.8, 133.3 (q, J = 33.2 Hz), 135.2, 137.3, 143.7, 173.3, 174.9, 179.8.

4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)phenyl)-*N*-methylbutanimidamide (87) [ND-3]

[0045] To a solution of 4-{4-[7-(4-Cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]oct-5-yl]-phenyl}-N-methyl-butyramide **(88)** [ND-6] (4.0 mg, 0.008 mmol) and pyridine (1.94 μL, 0.02 mmol) in dichloromethane (3 mL) at -40 °C was slowly added triflic anhydride (Tf_2O, 1.75 μL, 0.01 mmol). The mixture was allowed to warm to 0 °C over 3 h. The solution was then cooled to -40 °C and ammonia was introduced by bubbling. The reaction was then warmed to room temperature and stirred overnight. Without aqueous work up, flash chromatography using 10% methanol in ethyl acetate afforded 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)phenyl)-N-methylbutanimidamide **(87)** [ND-3] (2.9 mg, 72%) of as a colorless oil: 1H NMR (CD_3CN) δ 8.10 (d, 1H, J = 8.2 Hz), 8.04 (s, 1H), 7.92 (d, 1H, J = 8.2 Hz), 7.49 (br s, 2H), 7.43 (d, 2H, J = 8.3 Hz), 7.30 (d, 2H, J = 8.3 Hz), 3.26 (d, 3H, J = 5.4 Hz), 2.77 (t, 2H, J = 8.0 Hz), 2.56-2.65 (m, 2H), 2.52 (t, 2H, J = 7.7 Hz), 2.42-2.52 (m, 2H), 1.95-2.12 (m, 3H), 1.47-1.62 (m, 1H).



ND-6 (88) ND-3 (87)

Synthesis of ND-7 and ND-8

Dimethyl 2-(2-fluoro-4-nitrophenyl)malonate (86)

[0046] To a suspension of sodium hydride (NaH, 60%, 0.40 g, 10.0 mmol) in dry DMF (10 mL) under ice cooling was added dimethyl malonate (1.04 mL, 9.1 mmol) dropwise followed by a solution of 1-bromo-2-fluoro-4-nitrobenzene (1.00 g, 4.55 mmol) in dry DMF (3 mL) under an argon atmosphere. The resulting mixture was stirred at 70 °C overnight and then allowed to cool to 21 °C. The reaction mixture was quenched with saturated NH_4Cl and

extracted with ethyl acetate (2 × 50 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (hexane:ethyl acetate, 2:1) to give Dimethyl 2-(2-fluoro-4-nitrophenyl)malonate **(86)** (0.90 g, 73%) as a light yellowish solid: 1H NMR δ 8.07 (dd, 1H, J = 8.6, 2.2 Hz), 7.98 (dd, 1H, J = 9.3, 2.2 Hz), 7.74 (dd, 1H, J = 8.6, 7.1 Hz), 5.08 (s, 1H), 3.81 (s, 6H).

Trimethyl 1-(2-fluoro-4-nitrophenyl)propane-1,1,3-tricarboxylate (85)

[0047] To a solution of the nitro diester, Dimethyl 2-(2-fluoro-4-nitrophenyl)malonate **(86)** (0.44 g, 1.62 mmol) and methyl acrylate (0.22 mL, 2.43 mmol) in absolute methanol (5 mL) was added a catalytic amount of sodium methoxide at 21 °C under argon. The reaction mixture was stirred for 40 h at the same temperature and then diluted with dichloromethane (50 mL). The resulting mixture was washed with water, brine and dried. The residue obtained upon evaporation of the solvents was purified on a silica gel (hexane:ethyl acetate, 8:1) to give Trimethyl 1-(2-fluoro-4-nitrophenyl)propane-1,1,3-tricarboxylate **(85)** (0.49 g, 85%): 1H NMR δ 8.04 (dd, 1H, J = 8.7, 2.3 Hz), 7.95 (dd, 1H, J = 10.9, 2.3 Hz), 7.57 (dd, 1H, J = 8.7, 7.5 Hz), 3.81 (s, 6H), 3.62 (s, 3H), 2.64-2.69 (m, 2H), 2.35-2.40 (m, 2H).

Methyl 4-(2-fluoro-4-nitrophenyl)butanoate (84)

[0048] A solution of compound Trimethyl 1-(2-fluoro-4-nitrophenyl)propane-1,1,3-tricarboxylate **(85)** (0.23 g, 0.63 mmol), sodium chloride (0.11 g, 1.90 mmol) and water (0.15 mL) in distilled dimethylsulfoxide (4 mL) was heated to 155 °C overnight. The reaction mixture was allowed to cool to 21 °C and then worked up by adding water and extracting with ethyl acetate (2 × 50 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (hexane:ethyl acetate, 8:1) to give desired Methyl 4-(2-fluoro-4-nitrophenyl)butanoate **(84)** (69 mg, 45%) and dimethyl 2-(2-fluoro-4-nitrophenyl)pentanedioate **(83)** (72 mg, 38%): 1H NMR of (10) δ 8.04 (dd, 1H, J = 8.5, 2.2 Hz), 7.95 (dd, 1H, J = 9.5, 2.2 Hz), 7.53 (dd, 1H, J = 8.5, 7.1 Hz), 4.08 (t, 1H, J = 7.6 Hz), 3.71 (s, 3H), 3.66 (s, 3H), 2.43-2.52 (m, 1H), 2.31-2.35 (m, 2H), 2.06-2.14 (m, 1H); 1H NMR of **(84)** δ 7.98 (dd, 1H, J = 8.4, 2.2 Hz), 7.90 (dd, 1H, J = 9.5, 2.2 Hz), 7.38 (dd, 1H, J = 8.4, 7.3 Hz), 3.68 (s, 3H), 2.79 (t, 2H, J = 7.7 Hz), 2.38 (t, 2H, J = 7.3 Hz), 1.94-2.02 (m, 2H).

4-(2-Fluoro-4-nitrophenyl)butanoic acid (82)

[0049] To a solution of Methyl 4-(2-fluoro-4-nitrophenyl)butanoate **(84)** (43 mg, 0.18 mmol) in methanol (1 mL) and water (3 mL) was added sodium hydroxide (0.18 g, 4.50 mmol). The reaction mixture was stirred at 21 °C overnight. The reaction mixture was quenched with 1 N HCl solution and extracted with ethyl acetate (2 × 30 mL). The organic layer was dried over $MgSO_4$, concentrated to give 4-(2-Fluoro-4-nitrophenyl)butanoic acid **(82)** (40 mg, 98%) and the residue was used without further purification.

4-(2-Fluoro-4-nitrophenyl)-*N*-methylbutanamide (81)

[0050] Thionyl chloride (0.01 mL, 0.11 mmol) was added slowly to a solution of 4-(2-Fluoro-4-nitrophenyl)butanoic acid **(82)** (20 mg, 0.09 mmol) in DMF (3 mL) cooled at -5 °C. The mixture was stirred for an additional 1 h at -5 °C. Excess methylamine (freshly distilled from its 40% aqueous solution) was added to the reaction medium. The second mixture was stirred for an additional 1 h. Ethyl acetate (30 mL) was added to the mixture, which was washed with brine (2 × 30 mL). The organic layer was dried over $MgSO_4$, and concentrated to yield 4-(2-Fluoro-4-nitrophenyl)-N-methylbutanamide **(81)** (18 mg, 85%): 1H NMR δ 7.97 (dd, 1H, J = 8.4, 2.2 Hz), 7.89 (dd, 1H, J = 9.5, 2.2 Hz), 7.40 (dd, 1H, J = 8.4, 7.3 Hz), 5.44 (br s, 1H), 2.81 (d, 3H, J = 4.9 Hz), 2.79 (t, 2H, J = 7.6 Hz), 2.22 (t, 2H, J = 7.3 Hz), 1.96-2.04 (m, 2H).

4-(4-Amino-2-fluorophenyl)-*N*-methylbutanamide (80)

[0051] A solution of compound 4-(2-Fluoro-4-nitrophenyl)-N-methylbutanamide **(81)** (18 mg, 0.07 mmol), Fe (30 mg, 0.52 mmol) and AcOH (1 mL) in ethyl acetate (3 mL) was heated under reflux for 2 h. The reaction mixture was allowed to cool to 21 °C and then filtered. The organic layer was concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give desired 4-(4-Amino-2-fluorophenyl)-N-methylbutanamide **(80)** (14 mg, 86%): 1H NMR δ 6.92 (dd, 1H, J = 8.3, 8.2 Hz), 6.39 (dd, 1H, J = 8.3, 2.0 Hz), 6.33 (dd, 1H, J = 13.3, 2.0 Hz), 5.48 (br s, 1H), 3.69 (br s, 2H), 2.79 (d, 3H, J = 4.8 Hz), 2.55 (t, 2H, J = 7.4 Hz), 2.16 (t, 2H, J = 7.5 Hz), 1.85-1.94 (m, 2H).

4-(4-(1-Cyanocyclobutylamino)-2-fluorophenyl)-*N*-methylbutanamide (79)

[0052] A mixture of 4-(4-Amino-2-fluorophenyl)-N-methylbutanamide **(80)** (8 mg, 0.04 mmol), cyclobutanone (5 mg, 0.08 mmol) and trimethylsilyl cyanide (TMSCN, 8 mg,

0.08 mmol) was heated to 80 °C and stirred for 15 h. To the medium was added ethyl acetate (2 × 20 mL) and then washed with water (2 × 20 mL). The organic layer was dried over $MgSO_4$ and concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give 4-(4-(1-Cyanocyclobutylamino)-2-fluorophenyl)-N-methylbutanamide **(79)** (10 mg, 92%).

4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-*N*-methylbutanamide (78) [ND-7]

[0053] A mixture of 4-(4-(1-Cyanocyclobutylamino)-2-fluorophenyl)-N-methylbutanamide **(79)** (7 mg, 0.02 mmol) and 4-isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (12 mg, 0.05 mmol) in DMF (1 mL) was heated to 80 °C using microwave for 16 h. To this mixture was added methanol (3 mL) and aq. 1 N HCl (3 mL). The second mixture was refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture was poured into cold water (30 mL) and extracted with ethyl acetate (30 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 95:5) to give 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-N-methylbutanamide **(78)** [ND-7] (8 mg, 62%) as a pale yellowish solid: ^{1}H NMR δ 7.98 (d, 1H, J = 8.2 Hz), 7.97 (d, 1H, J = 2.0 Hz), 7.84 (dd, 1H, J = 8.2, 2.0 Hz), 7.43 (dd, 1H, J = 8.0, 8.0 Hz), 7.06 (dd, 1H, J = 8.0, 2.0 Hz), 7.02 (dd, 1H, J = 9.7, 2.0 Hz), 2.83 (d, 3H, J = 4.8 Hz), 2.78 (t, 2H, J = 7.7 Hz), 2.63-2.71 (m, 2H), 2.51-2.62 (m, 2H), 2.27 (t, 2H, J = 7.3 Hz), 2.18-2.27 (m, 1H), 2.00-2.09 (m, 2H), 1.66-1.76 (m, 1H); ^{13}C NMR δ 179.9, 174.7, 172.8, 161.2 (d, J = 247 Hz), 137.0, 135.2, 134.1 (d, J = 9.6 Hz), 133.6 (q, J = 33.7 Hz), 132.1, 131.9 (d, J = 5.6 Hz), 130.8 (d, J = 15.2 Hz), 127.1, 125.7 (d, J = 3.9 Hz), 121.9 (q, J = 272 Hz), 117.3 (d, J = 22.3 Hz), 114.8, 110.0, 67.4, 35.8, 31.5, 28.3, 26.4, 25.6, 13.7.

4-(2-Fluoro-4-nitrophenyl)-*N,N*-dimethylbutanamide (77)

[0054] Thionyl chloride (0.01 mL, 0.11 mmol) was added slowly to a solution of 4-(2-Fluoro-4-nitrophenyl)butanoic acid **(82)** (18 mg, 0.08 mmol) in DMF (3 mL) cooled at -5 °C. The mixture was stirred for an additional 1 h at -5 °C. Excess dimethylamine (freshly distilled from its 40% aqueous solution) was added to the reaction medium. The second mixture was stirred for an additional 1 h. Ethyl acetate (30 mL) was added to the mixture, which was washed

with brine (2 × 30 mL). The organic layer was dried over $MgSO_4$, and concentrated to yield 4-(2-Fluoro-4-nitrophenyl)-N,N-dimethylbutanamide **(77)** (18 mg, 87%): ^{1}H NMR δ 7.98 (dd, 1H, J = 8.3, 2.1 Hz), 7.89 (dd, 1H, J = 9.5, 2.1 Hz), 7.42 (dd, 1H, J = 8.3, 7.4 Hz), 2.98 (s, 3H), 2.95 (s, 3H), 2.81 (t, 2H, J = 7.6 Hz), 2.36 (t, 2H, J = 7.2 Hz), 1.96-2.04 (m, 2H).

4-(4-Amino-2-fluorophenyl)-*N,N*-dimethylbutanamide (76)

[0055] A solution of compound 4-(2-Fluoro-4-nitrophenyl)-N,N-dimethylbutanamide **(77)** (15 mg, 0.06 mmol), Fe (20 mg, 0.37 mmol) and acetic acid (1 mL) in ethyl acetate (3 mL) was heated under reflux for 2 h. The reaction mixture was allowed to cool to 21 °C and then filtered. The organic layer was concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give desired 4-(4-Amino-2-fluorophenyl)-N,N-dimethylbutanamide **(76)** (12 mg, 87%): ^{1}H NMR δ 6.95 (dd, 1H, J = 8.3, 8.2 Hz), 6.40 (dd, 1H, J = 8.3, 2.2 Hz), 6.35 (dd, 1H, J = 11.6, 2.2 Hz), 3.66 (br s, 2H), 2.95 (s, 3H), 2.93 (s, 3H), 2.58 (t, 2H, J = 7.4 Hz), 2.30 (t, 2H, J = 7.6 Hz), 1.85-1.95 (m, 2H).

4-(4-(2-Cyanopropan-2-ylamino)-2-fluorophenyl)-*N,N*-dimethylbutanamide (75)

[0056] A mixture of 4-(4-Amino-2-fluorophenyl)-*N,N*-dimethylbutanamide **(76)** (10 mg, 0.05 mmol), cyclobutanone (6 mg, 0.09 mmol) and trimethylsilyl cyanide (TMSCN, 9 mg, 0.09 mmol) was heated to 80 °C and stirred for 15 h. To the medium was added ethyl acetate (2 × 20 mL) and then washed with water (2 × 20 mL). The organic layer was dried over $MgSO_4$ and concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give 4-(4-(2-Cyanopropan-2-ylamino)-2-fluorophenyl)-*N,N*-dimethylbutanamide **(75)** (12 mg, 89%): ^{1}H NMR δ 7.04 (dd, 1H, J = 8.0, 7.8 Hz), 6.36 (dd, 1H, J = 8.0, 2.3 Hz), 6.32 (dd, 1H, J = 11.6, 2.3 Hz), 4.08 (br s, 1H), 2.96 (s, 3H), 2.93 (s, 3H), 2.77-2.81 (m, 2H), 2.61 (t, 2H, J = 7.4 Hz),2.35-2.38 (m, 2H), 2.31 (t, 2H, J = 7.6 Hz), 2.10-2.37 (m, 2H), 1.87-1.95 (m, 2H).

4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-*N,N*-dimethylbutanamide (74) [ND-8]

[0057] A mixture of 4-(4-(2-Cyanopropan-2-ylamino)-2-fluorophenyl)-*N,N*-dimethylbutanamide **(75)** (7 mg, 0.02 mmol) and 4-isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (12 mg, 0.05 mmol) in DMF (1 mL) was heated to 80 °C using microwave for 16 h. To this

mixture was added methanol (3 mL) and aq. 1 N HCl (3 mL). The second mixture was refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture was poured into cold water (30 mL) and extracted with ethyl acetate (30 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 95:5) to give 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-*N*,*N*-dimethylbutanamide **(74)** [ND-8] (8 mg, 65%) as a pale yellowish solid: ^{1}H NMR δ 7.98 (d, 1H, J = 8.2 Hz), 7.97 (d, 1H, J = 2.1 Hz), 7.84 (dd, 1H, J = 8.2, 2.1 Hz), 7.46 (dd, 1H, J = 8.0, 8.0 Hz), 7.05 (dd, 1H, J = 8.0, 2.2 Hz), 7.02 (dd, 1H, J = 9.6, 2.2 Hz), 3.01 (s, 3H), 2.97 (s, 3H), 2.80 (t, 2H, J = 7.8 Hz), 2.63-2.71 (m, 2H), 2.52-2.62 (m, 2H), 2.42 (t, 2H, J = 7.4 Hz), 2.20-2.31 (m, 1H), 2.00-2.08 (m, 2H), 1.65-1.75 (m, 1H); ^{13}C NMR δ 179.9, 174.7 (2 C's), 161.3 (d, J = 248 Hz), 137.0, 135.2, 134.1 (d, J = 10.3 Hz), 133.6 (q, J = 33.3 Hz), 132.1, 131.9 (d, J = 5.7 Hz), 131.2 (d, J = 16.2 Hz), 127.1, 125.7 (d, J = 4.3 Hz), 121.9 (q, J = 272 Hz), 117.2 (d, J = 25.1 Hz), 114.8, 110.2, 67.5, 37.2, 35.5, 32.7, 31.6, 28.5, 25.2, 13.7.



Synthesis of ND-9

Dimethyl 2-(2-cyanoethyl)-2-(2-fluoro-4-nitrophenyl)malonate (73)

[0058] To a solution of the nitro diester, Dimethyl 2-(2-fluoro-4-nitrophenyl)malonate **(86)** (0.4 g, 1.47 mmol) and acrylonitrile (0.11 mL, 1.62 mmol) in absolute methanol (5 mL) was added a catalytic amount of sodium methoxide at 21 °C under argon. The reaction mixture was stirred for 40 h at the same temperature and then diluted with dichloromethane (50 mL). The resulting mixture was washed with water, brine and dried. The residue obtained upon evaporation of the solvents was purified on a silica gel (hexane:ethyl acetate, 8:1) to give Dimethyl 2-(2-cyanoethyl)-2-(2-fluoro-4-nitrophenyl)malonate **(73)** (0.25 g, 52%): ^{1}H NMR δ 8.07 (dd, 1H, J = 8.7, 2.3 Hz), 7.99 (dd, 1H, J = 10.9, 2.3 Hz), 7.47 (dd, 1H, J = 8.7, 7.3 Hz), 3.85 (s, 6H), 2.65-2.70 (m, 2H), 2.47-2.51 (m, 2H).

4-(2-Fluoro-4-nitrophenyl)butanenitrile (72)

[0059] A solution of compound Dimethyl 2-(2-cyanoethyl)-2-(2-fluoro-4-nitrophenyl)malonate **(73)** (0.19 g, 0.59 mmol), sodium chloride (0.10 g, 1.76 mmol) and water (0.15 mL) in distilled dimethylsulfoxide (DMSO, 4 mL) was heated to 155 °C overnight. The reaction mixture was allowed to cool to 21 °C and then worked up by adding water and extracting with ethyl acetate (2 × 50 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (hexane:ethyl acetate, 8:1) to give desired 4-(2-Fluoro-4-nitrophenyl)butanenitrile **(72)** (79 mg, 65%): ^{1}H NMR δ 8.02 (dd, 1H, J = 8.3, 2.2 Hz), 7.94 (dd, 1H, J = 9.5, 2.2 Hz), 7.42 (dd, 1H, J = 8.3, 7.4 Hz), 2.93 (t, 2H, J = 7.7 Hz), 2.41 (t, 2H, J = 7.0 Hz), 2.01-2.07 (m, 2H).

4-(4-Amino-2-fluorophenyl)butanenitrile (71)

[0060] A solution of compound 4-(2-Fluoro-4-nitrophenyl)butanenitrile **(72)** (47 mg, 0.23 mmol), Fe (78 mg, 1.40 mmol) and acetic acid (1 mL) in ethyl acetate (3 mL) was heated under reflux for 2 h. The reaction mixture was allowed to cool to 21 °C and then filtered. The organic layer was concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give desired 4-(4-Amino-2-fluorophenyl)butanenitrile **(71)** (33 mg, 83%): ^{1}H NMR δ 6.98-7.01 (m, 1H), 6.46-6.52 (m, 2H), 2.70 (t, 2H, J = 7.6 Hz), 2.32 (t, 2H, J = 7.2 Hz), 1.89-1.98 (m, 2H).

1-(4-(3-Cyanopropyl)-3-fluorobenzyl)cyclobutanecarbonitrile (70)

[0061] A mixture of 4-(4-Amino-2-fluorophenyl)butanenitrile **(71)** (30 mg, 0.17 mmol), cyclobutanone (24 mg, 0.34 mmol) and trimethylsilyl cyanide (TMSCN, 33 mg, 0.34 mmol) was heated to 80 °C and stirred for 15 h. To the medium was added ethyl acetate (2 × 20 mL) and then washed with water (2 × 20 mL). The organic layer was dried over $MgSO_4$ and concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give 1-(4-(3-Cyanopropyl)-3-fluorobenzyl) cyclobutanecarbonitrile **(70)** (40 mg, 92%): ^{1}H NMR δ 7.01 (dd, 1H, J = 8.0, 7.5 Hz), 6.37 (dd, 1H, J = 8.0, 2.4 Hz), 6.34 (dd, 1H, J = 11.8, 2.4 Hz), 4.18 (br s, 1H), 2.76-2.81 (m, 2H), 2.70 (t, 2H, J = 7.3 Hz),2.33-2.39 (m, 2H), 2.33 (t, 2H, J = 7.1 Hz), 2.12-2.30 (m, 2H), 1.90-1.95 (m, 2H).

4-(5-(4-(3-Cyanopropyl)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile (69) [ND-9]

[0062] A mixture of 1-(4-(3-Cyanopropyl)-3-fluorobenzyl)cyclobutanecarbonitrile **(70)** (32 mg, 0.12 mmol) and 4-isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (62 mg, 0.27 mmol) in DMF (1 mL) was heated to 80 °C using microwave for 16 h. To this mixture was added methanol (3 mL) and aq. 1 N HCl (3 mL). The second mixture was refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture was poured into cold water (30 mL) and extracted with ethyl acetate (30 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 95:5) to give 4-(5-(4-(3-Cyanopropyl)-3-fluorophenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile **(69)** [ND-9] (48 mg, 80%) as a pale yellowish solid: ^{1}H NMR δ 7.98 (d, 1H, J = 8.2 Hz), 7.97 (d, 1H, J = 2.0 Hz), 7.84 (dd, 1H, J = 8.2, 2.0 Hz), 7.44 (dd, 1H, J = 8.0, 8.0 Hz), 7.10 (dd, 1H, J = 8.0, 2.0 Hz), 7.07 (dd, 1H, J = 10.2, 2.0 Hz), 2.92 (t, 2H, J = 7.6 Hz), 2.64-2.71 (m, 2H), 2.51-2.61 (m, 2H), 2.45 (t, 2H, J = 7.1 Hz), 2.20-2.31 (m, 1H), 2.03-2.11 (m, 2H), 1.64-1.75 (m, 1H); ^{13}C NMR δ 179.9, 174.6, 161.3 (d, J = 248 Hz), 137.0, 135.2, 134.9 (d, J = 10.0 Hz), 133.6 (q, J = 33.2 Hz), 132.2, 131.9 (d, J = 5.8 Hz), 129.0 (d, J = 15.7 Hz), 127.1, 126.0 (d, J = 3.6 Hz), 121.9 (q, J = 273 Hz), 119.1, 117.6 (d, J = 23.4 Hz), 114.8, 110.1, 67.4, 31.6 (2 C's), 28.1, 25.4, 16.8, 13.7.



Synthesis of ND-11 and ND-10

4-(8-Oxo-5-(4-(4-oxobutyl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile (68) [ND-10]

[0063] To a stirred solution of 4-{4-[7-(4-cyano-3-trifluoromethylphenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-5-yl]-phenyl}-butyric acid methyl ester **(67)** [ND-4] (61 mg, 0.12 mmol) in dichloromethane (5 mL), 1M diisobutylaluminum hydride (DIBAL) solution in hexane (0.16 mL, 0.16 mmol) was added at -78 °C. After 30 min, the reaction mixture was quenched with saturated Rochelle's salt solution. The resulting mixture was stirred at 21 °C until both phases were clearly separated and the organic layer was clear. After extraction, the separated organic layer was dried over $MgSO_4$, filtered, and concentrated under vacuum. The crude mixture of **(66)** and **(68)** was purified by flash column chromatography (hexane:ethyl acetate, 4:1) to give 4-(8-hydroxy-5-(4-(4-oxobutyl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl) **(66)** (20 mg, 35%) and 4-(8-Oxo-5-(4-(4-oxobutyl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile **(68)** [ND-10] (23 mg, 40%): ^{1}H NMR of **(68)** δ 9.81 (s, 1H), 7.98 (d, 1H, J = 2.0 Hz), 7.97 (d, 1H, J = 8.0 Hz), 7.86 (dd, 1H, J = 8.2, 2.0 Hz), 7.40 (d, 1H, J = 8.3 Hz), 7.24 (d, 1H, J = 8.3 Hz), 2.76 (t, 2H, J = 7.5 Hz), 2.63-2.67 (m, 2H), 2.57-2.63 (m, 2H), 2.55 (t, 2H, J = 7.2 Hz), 2.13-2.31 (m, 1H), 2.01-2.07 (m, 2H), 1.57-1.77 (m, 1H).

4-(5-(4-(3-(4,5-Dihydro-1H-imidazol-2-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile (65) [ND-11]

[0064] The mixture of 4-(8-Oxo-5-(4-(4-oxobutyl)phenyl)-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile **(68)** [ND-10] (15 mg, 0.03 mmol) and ethylene diamine (2 μL, 0.04 mmol) in dry dichloromethane (3 mL) was stirred at 0 °C for 30 min under argon. N-Bromosuccinimide (NBS, 6 mg, 0.04 mmol) was added to the mixture and the resulting solution was stirred overnight at 21 °C. Reaction was quenched by the addition of saturated $NaHCO_3$ solution. The mixture was extracted with dichloromethane. The organic layer was dried over $MgSO_4$, and evaporated in vacuo. The residue was purified by flash column chromatography (ethanol:ethyl acetate, 1:4) to give 4-(5-(4-(3-(4,5-Dihydro-1H-imidazol-2-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diaza-spiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile **(65)** [ND-11] (6 mg, 35%): ^{1}H NMR δ 7.98 (d, 1H, J = 1.9 Hz), 7.97 (d, 1H, J = 8.3 Hz), 7.85 (dd, 1H, J = 8.3, 1.9 Hz), 7.46 (d, 1H, J = 8.2 Hz), 7.25 (d, 1H, J = 8.2 Hz), 4.72 (br s, 1H), 3.67-3.80 (m, 4H), 2.85-3.05 (m, 2H), 2.57-2.70 (m, 2H), 2.43-2.57 (m, 4H), 2.15-2.30 (m, 1H), 1.63-1.80 (m, 3H).



Synthesis of ND-12

4-(4-Nitrophenyl)butanal (64)

[0065] To a stirred solution of methyl 4-(4-nitrophenyl)butanoate **(63)** (0.45 g, 2.02 mmol) in dichloromethane (30 mL), 1M diisobutylaluminum hydride (DIBAL) solution in

hexane (2.62 mL, 2.62 mmol) was added at -78 °C. After 30 min, the reaction mixture was quenched with saturated Rochelle's salt solution. The resulting mixture was stirred at 21 °C until both phases were clearly separated and the organic layer was clear. After extraction, the separated organic layer was dried over $MgSO_4$, filtered, and concentrated under vacuum. The crude 4-(4-Nitrophenyl)butanal **(64)** was purified by flash column chromatography (hexane:ethyl acetate, 8:1) to give 4-(4-Nitrophenyl)butanal **(64)** (0.28 g, 72%): ^{1}H NMR δ 9.79 (s, 1H), 8.16 (d, 2H, J = 8.7 Hz), 7.34 (d, 2H, J = 8.7 Hz), 2.77 (t, 2H, J = 7.7 Hz), 2.51 (t, 2H, J = 7.1 Hz), 1.95-2.04 (m, 2H).

2-(3-(4-Nitrophenyl)propyl)-4,5-dihydro-1H-imidazole (62)

[0066] The mixture of 4-(4-Nitrophenyl)butanal **(64)** (0.28 g, 1.45 mmol) and ethylene diamine (0.1 mL, 1.59 mmol) in dry dichloromethane (10 mL) was stirred at 0 °C for 30 min under argon. NBS (0.26 g, 1.59 mmol) was added to the mixture and the resulting solution was stirred overnight at 21 °C. Reaction was quenched by the addition of saturated $NaHCO_3$ solution. The mixture was extracted with dichloromethane. The organic layer was dried over $MgSO_4$, and evaporated in vacuo. The residue was purified by flash column chromatography (ethanol:ethyl acetate:triethylamine, 1:1:0.2) to give 2-(3-(4-Nitrophenyl)propyl)-4,5-dihydro-1H-imidazole **(62)** (0.26 g, 76%): ^{1}H NMR δ 8.14 (d, 2H, J = 8.7 Hz), 7.35 (d, 2H, J = 8.7 Hz), 3.59 (s, 4H), 2.79 (t, 2H, J = 7.7 Hz), 2.26 (t, 2H, J = 7.4 Hz), 1.96-2.05 (m, 2H).

[0067] An alternative route for synthesizing 2-(3-(4-Nitrophenyl)propyl)-4,5-dihydro-1H-imidazole **(62)** from methyl 4-(4-nitrophenyl)butanoate **(63)** was also used and is as follows. Ethylenediamine (0.1 mL, 1.59 mmol) was added dropwise to a stirred solution of trimethylaluminum (1.59 mmol) in 2 mL of toluene, so that the temperature did not exceed 10 °C. At the end of methane evolution the ester **(63)** (0.22 g, 1.00 mmol) was gradually added at room temperature. The reaction mixture was refluxed for 3 h. After cooling, the solution was treated dropwise with 1 mL of water, diluted with 3 mL of methanol and 3 mL of methylene chloride, and refluxed on a steam bath for 15 min. After filtration over $MgSO_4$ and solvent evaporation the residue was purified by flash column chromatography (ethanol:ethyl acetate:triethylamine, 1:1:0.2) to give 2-(3-(4-Nitrophenyl)propyl)-4,5-dihydro-1H-imidazole **(62)** (0.10 g, 45%): ^{1}H NMR δ 8.14 (d, 2H, J = 8.7 Hz), 7.35 (d, 2H, J = 8.7 Hz), 3.59 (s, 4H), 2.79 (t, 2H, J = 7.7 Hz), 2.26 (t, 2H, J = 7.4 Hz), 1.96-2.05 (m, 2H).

***tert*-Butyl 2-(3-(4-nitrophenyl)propyl)-1H-imidazole-1-carboxylate (61)**

[0068] To a solution of dichloromethane (5 mL) and dimethylsulfoxide (0.06 mL, 0.79 mmol) was added oxalyl chloride (0.07 mL, 0.79 mmol) at -78 °C under an argon atmosphere. After stirring for 20 min, a solution of 2-(3-(4-Nitrophenyl)propyl)-4,5-dihydro-1H-imidazole **(62)** (74 mg, 0.32 mmol) in dichloromethane was added to the reaction mixture. After stirring for 50 min, triethylamine (0.22 mL, 1.59 mmol) was added and then the reaction mixture was warmed to room temperature. After stirring for 50 min, aqueous ammonia solution (10 mL) was added and the resulting mixture was extracted with chloroform (20 mL). The combined organic layer was washed with brine, dried, filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography (dichloromethane:methanol, 10:1) to give the corresponding imidazole (61 mg, 83%). To a solution of the imidazole (50 mg, 0.22 mmol) in dichloromethane (5 mL) was added triethylamine (0.04 mL, 0.26 mmol) and *tert*-butoxycarbonyl anhydride (Boc_2O, 57 mg, 0.26 mmol). The reaction mixture was stirred at 21 °C overnight. The reaction mixture was extracted with dichloromethane (20 mL). The organic layer was washed with brine, dried, filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography (dichloromethane:methanol, 20:1) to give *tert*-Butyl 2-(3-(4-nitrophenyl)propyl)-1H-imidazole-1-carboxylate **(61)** (72 mg, quant.): ^{1}H NMR δ 8.14 (d, 2H, J = 8.7 Hz), 7.37 (d, 2H, J = 8.7 Hz), 7.30 (d, 1H, J = 1.7 Hz), 6.86 (d, 1H, J = 1.7 Hz), 3.05 (t, 2H, J = 7.6 Hz), 2.85 (t, 2H, J = 7.7 Hz), 2.11-2.19 (m, 2H), 1.60 (s, 9H).

***tert*-Butyl 2-(3-(4-(1-cyanocyclobutylamino)phenyl)propyl)-1H-imidazole-1-carboxylate (60)**

[0069] To a solution of *tert*-Butyl 2-(3-(4-nitrophenyl)propyl)-1H-imidazole-1-carboxylate **(61)** (72 mg, 0.22 mmol) in ethyl acetate (5 mL) was introduced hydrogen gas in the presence of a catalytic amount of Pd/C. After completion of the reaction, the reaction mixture was filtered, concentrated and then purified by flash column chromatography (dichloromethane:methanol, 10:1) to give the corresponding amine (59 mg, 90%): ^{1}H NMR δ 7.30 (d, 1H, J = 1.7 Hz), 7.00 (d, 2H, J = 8.3 Hz), 6.85 (d, 1H, J = 1.7 Hz), 6.62 (d, 2H, J = 8.3 Hz), 3.54 (br s, 2H), 3.01 (t, 2H, J = 7.8 Hz), 2.62 (t, 2H, J = 7.7 Hz), 2.01-2.08 (m, 2H), 1.65 (s, 9H). A mixture of the amine (55 mg, 0.18 mmol), cyclobutanone (26 mg, 0.36 mmol) and trimethylsilyl cyanide (TMSCN, 36 mg, 0.36 mmol) was heated to 80 °C and stirred for 15 h. To the medium was added ethyl acetate (2 × 20 mL) and then washed with water (2 ×

20 mL). The organic layer was dried over $MgSO_4$ and concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give *tert*-Butyl 2-(3-(4-(1-cyanocyclobutylamino)phenyl)propyl)-1H-imidazole-1-carboxylate **(60)** (57 mg, 82%): ^{1}H NMR δ 7.30 (d, 1H, J = 1.7 Hz), 7.09 (d, 2H, J = 8.4 Hz), 6.85 (d, 1H, J = 1.7 Hz), 6.58 (d, 2H, J = 8.4 Hz), 3.92 (br s, 1H), 3.01 (t, 2H, J = 7.7 Hz), 2.74-2.80 (m, 2H), 2.64 (t, 2H, J = 7.6 Hz), 2.29-2.42 (m, 2H), 2.10-2.27 (m, 2H), 2.01-2.09 (m, 2H), 1.60 (s, 9H).

4-(5-(4-(3-(1H-Imidazol-2-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile (59) [ND-12]

[0070] A mixture of *tert*-Butyl 2-(3-(4-(1-cyanocyclobutylamino)phenyl)propyl)-1H-imidazole-1-carboxylate **(60)** (22 mg, 0.06 mmol) and 4-isothiocyanato-2-trifluoromethylbenzonitrile **(96)** (26 mg, 0.12 mmol) in DMF (1 mL) was heated to 80 °C using microwave for 16 h. To this mixture was added methanol (3 mL) and aq. 1 N HCl (3 mL). The second mixture was refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture was poured into cold water (30 mL), treated with saturated $NaHCO_3$ solution and extracted with ethyl acetate (50 mL). The organic layer was dried over $MgSO_4$, concentrated and the residue was purified with silica gel column chromatography (dichloromethane:acetone, 9:1) to give 4-(5-(4-(3-(1H-Imidazol-2-yl)propyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]octan-7-yl)-2-(trifluoromethyl)benzonitrile **(59)** [ND-12] (29 mg, 52%) as a pale yellowish solid: ^{1}H NMR δ 8.65 (br s, 1H), 7.97 (d, 1H, J = 2.0 Hz), 7.96 (d, 1H, J = 8.4 Hz), 7.84 (dd, 1H, J = 8.4, 2.0 Hz), 7.34 (d, 2H, J = 8.2 Hz), 7.19 (d, 2H, J = 8.2 Hz), 3.00 (t, 2H, J = 7.5 Hz), 2.73 (t, 2H, J = 7.7 Hz), 2.47-2.77 (m, 4H), 2.13-2.25 (m, 3H), 1.51-1.71 (m, 1H).



[0071] One skilled in the art could modify and/or combine the syntheses described herein to make other diarylhydantoin compounds.

Synthesis of ND-13

4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-2,2-dimethyl-*N*-methylbutanamide (113)

[0072] Another compound envisioned is 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-2,2-dimethyl-*N*-methylbutanamide **(113)** [ND-13].



[ND-13]

[0073] An example of a synthetic route for making 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-2,2-dimethyl-*N*-methylbutanamide **(113)** [ND-13] is below.



ND-13

[0074] Alternatively, 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-2,2-dimethyl-*N*-methylbutanamide **(113)** [ND-13] can be synthesized in a manner similar as to that for synthesizing **(92)** [ND-2]. A mixture of 4-Isothiocyanato-2-trifluoromethylbenzonitrile **(96)** and 4-(4-(1-cyanocyclobutylamino)phenyl)-N,2,2-trimethylbutanamide **(111)** in solvent, for example, in DMF, is heated under microwave irradiation at 80 °C for 6 h.



111

To this mixture is added alcohol, e.g., methanol, and acid, e.g., aqueous hydrochloric acid. The second mixture is refluxed for 1.5 h. After being cooled to room temperature, the reaction mixture is poured into cold water and extracted, for example, with ethyl acetate. The organic layer is dried, e.g., dried over $MgSO_4$, concentrated, and the residue is purified, for example, by

silica gel column chromatography using hexane:ethyl acetate (2:1), to give 4-(4-(7-(4-Cyano-3-(trifluoromethyl)phenyl)-8-oxo-6-thioxo-5,7-diazaspiro[3.4]oct-an-5-yl)-2-fluorophenyl)-2,2-dimethyl-*N*-methylbutanamide **(113)** [ND-13].



(113) [ND-13]

[0075] Inventive compounds also include those with the following formulas.



[0076] R_1 and R_2 together can comprise eight or fewer carbon atoms and can be alkyl, substituted alkyl, or, together with the carbon to which they are linked, a cycloalkyl or substituted cycloalkyl group. R_3 can be hydrogen, cyano, formyl,



,


,


, or


. R_4 can be hydrogen, F, Cl, Br, and I. R_{11} and R_{12} can be the same or different and can be hydrogen or methyl. R_{13} can be hydrogen or $-NR_{14}R_{15}$. R_{14} and R_{15} can be the same or different and can be hydrogen or methyl.

Pharmacological examination of the compounds

[0077] Compounds for which synthetic routes are described above can be evaluated through screening on hormone refractory prostate cancer cells for antagonistic and agonistic activities against AR utilizing screening procedures similar to those in PCT applications bearing numbers US04/42221, US05/05529, and US06/11417 and U.S. Application Serial Number 11/433,829, which are hereby incorporated by reference.

In vitro biological assay

Effect of compounds on AR by a reporter assay

[0078] For example, the compounds can be subjected to tests using an artificial androgen receptor (AR) response reporter system in a hormone refractory prostate cancer cell line. The prostate cancer LNCaP cells are engineered to stably express about 5-fold higher level of AR than endogenous level. The exogenous AR has similar properties to endogenous AR in that both are stabilized by a synthetic androgen R1881. The AR-over expressed cells are also engineered to stably incorporate an AR response reporter and the reporter activity of these cells shows features of hormone refractory prostate cancer. It responds to low concentration of a synthetic androgen R1881, is inhibited only by high concentrations of bicalutamide, and displays agonistic activity with bicalutamide. Bicalutamide inhibits AR response reporter and does not have agonistic activity in hormone sensitive prostate cancer cells.

[0079] The antagonistic activity of the compounds for which the synthesis is described above can be examined in the presence of 100 pM of R1881. Engineered LNCaP cells (LNCaP-AR, also abbreviated LN-AR) are maintained in Iscove's medium containing 10% fetal bovine

serum (FBS). Two days prior to drug treatment, the cells are grown in Iscove's medium containing 10% charcoal-stripped FBS (CS-FBS) to deprive of androgens. The cells are split and grown in Iscove's medium containing 10% CS-FBS with 100 pM of R1881 and increasing concentrations of test compounds. After two days of incubation, reporter activities are assayed. Bicalutamide is used as a control substance.

[0080] One previously unrecognized property of AR overexpression in hormone refractory prostate cancer is its ability to switch antagonists to agonists. Therefore, only those compounds with minimal or no agonistic activities are qualified to be anti-androgens for this disease. To determine agonistic activities of different compounds, the stimulating activities on androgen receptor (AR) using the AR response reporter as the measure in the LN-AR system in the absence of R1881 can be examined. Bicalutamide can activate AR in hormone refractory prostate cancer. RU59063 and other anti-androgenic compounds listed as examples in US Patent Number 5,705,654 can activate AR in hormone refractory prostate cancer.

[0081] To examine the specificity of AR inhibitors, compounds can be tested in LNCaP cells with an over expression of glucocorticoid receptor (GR), the closest member of AR in the nuclear receptor family. These cells also carry a GR response reporter and the reporter activity can be induced by dexamethasone, a GR agonist, and the induction can be blocked by RU486, a GR inhibitor.

Effect of compounds on AR by measuring secreted levels of prostate specific antigen (PSA)

[0082] PSA levels are indicators of androgen receptor (AR) activities in prostate cancer. To examine if the compounds affect AR function in a physiological environment, secreted levels of endogenous PSA induced by R1881 in the AR-overexpressed LNCaP cells (LNCaP-AR, also abbreviated LN-AR) can be determined. The LNCaP-AR cells are a line of lymph node carcinoma of prostate cells transduced with a plasmid that makes express androgen receptors. LNCaP-AR cells are maintained in Iscove's medium containing 10% FBS. Two days prior to drug treatment, the cells are grown in Iscove's medium containing 10% CS-FBS to deprive of androgens. The cells are split and grown in Iscove's medium containing 10% CS-FBS with appropriate concentrations of R1881 and the test compounds. After four days incubation,

secreted PSA levels are assayed using PSA ELISA kits (American Qualex, San Clemente, CA)

[0083] The secreted PSA level of LNCaP-AR cells are strongly induced by 25 pM of R1881. In contrast, PSA is not induced in the parental LNCaP cells until concentration of R1881 reached 100 pM. Thus, the AR in hormone refractory prostate cancer is hyper-sensitive to androgens. A dose-dependent inhibition on AR activity is carried out to determine the IC50s of different compounds in inhibiting PSA expression.

[0084] Agonistic activities of selective compounds on AR in hormone refractory prostate cancer can be examined using secreted PSA as the surrogate marker. To do this, androgen-starved AR over expressed LNCaP cells are incubated with increasing concentrations of the compounds for which a synthesis is described above in the absence of R1881 and secreted PSA in the culture medium are measured 4 days later.

[0085] RU59063 and other antiandrogenic compounds listed as examples in US Patent No. 5,705,654 can stimulate PSA expression in hormone refractory prostate cancer.

Effect of compounds on AR mitochondrial activity by MTS assay

[0086] LNCaP-AR cells can be maintained in Iscove's medium containing 10% FBS. The compounds are examined for their effect on growth of hormone refractory prostate cancer cells. Overexpressed LNCaP cells are used because these cells behave as hormone refractory prostate cancer cells in vitro and in vivo. Mitochondria activity by MTS assay is measured, a surrogate for growth. LNCaP cells with overexpressed AR (LN-AR) are maintained in Iscove's medium containing 10% FBS. Two days prior to drug treatment, the cells are grown in Iscove's medium containing 10% CS-FBS to deprive of androgens. The cells are then split and grown in Iscove's medium containing 10% CS-FBS with appropriate concentrations of R1881 and increasing concentrations of the test compounds. After four days incubation, cell growth is monitored by MTS (Promega, Madison, WI).

[0087] Consistent with the reporter assay and PSA assay, growth of the AR-overexpressed LNCaP is stimulated by 25 microM of R1881, but the parental cells are not stimulated until R1881 concentration reaches 100 microM. The inhibitory effect of compounds

on growth of hormone refractory prostate cancer in the presence of 100 pM of R1881 is measured. Bicalutamide does not inhibit hormone refractory prostate cancer.

[0088] To examine whether growth inhibition in the MTS assay occurs by targeting AR, compounds can be tested in DU-145 cells, a prostate cancer cell line that lacks AR expression. The compounds can be tested for their ability to inhibit cells other than AR-expressed prostate cancer cells, such as MCF7 and SkBr3, two commonly used breast cancer cells, or 3T3, a normal mouse fibroblast cell line.

[0089] Based on the observations with various assays, the compounds can be ranked in order of their activity.

Inhibitory effect on hormone refractory prostate cancer xenograft tumors.

[0090] The in vivo effects of compounds on hormone refractory prostate cancer can be examined. The effect of compounds on xenograft tumors established from AR-overexpressed LNCaP cells can be examined. The engineered cells in Matrigel (Collaborative Biomedical) are injected subcutaneously into the flanks of the castrated male SCID mice. Tumor size is measured weekly in three dimensions using calipers. After xenograft tumors become established (for example, with a tumor size of at least 40 mm^3), mice with tumors are randomized and treated with different doses of compounds orally once daily. Bicalutamide does not inhibit growth of hormone refractory prostate cancer, the same as vehicle.

[0091] Compounds can also be tested in another xenograft model of hormone refractory prostate cancer, hormone refractory LAPC4. This model is established from passaging of hormone sensitive prostate cancer in castrated mice, which mimics the clinical progression of prostate cancer. Bicalutamide does not inhibit growth and PSA expression in hormone refractory LAPC4 xenograft model, the same as vehicle.

Inhibitory effect on growth of hormone sensitive prostate cancer cells.

[0092] To determine if compounds inhibit hormone sensitive prostate cancer cells, the effect of the compounds on growth of LNCaP cells can be examined by measuring MTS of mitochondria activities. Bicalutamide mildly inhibits hormone sensitive LNCaP cells in a dose-dependent manner.

In vivo biological assay

[0093] Animal experiments are performed in compliance with the guidelines of the Animal Research Committee of the University of California at Los Angeles. Animals are bought from Taconic and maintained in a laminar flow tower in a defined flora colony. LNCaP-AR and LNCaP-vector cells are maintained in RPMI medium supplemented with 10% FBS. 10^6 cells in 100 μl of 1:1 Matrigel to RPMI medium are injected subcutaneously into the flanks of intact or castrated male SCID mice. Tumor size is measured weekly in three dimensions (length x width x depth) using calipers. Mice are randomized to treatment groups when tumor size reaches approximately 100 mm^3. Drugs are given orally every day at 10 mg/kg and 50 mg/kg. To obtain pharmacodynamic readout, the animals are imaged via an optical CCD camera, 3 hours after last dose of the treatment. An ROI is drawn over the tumor for luciferase activity measurement in photon/second.

[0094] The pharmacokinetics of bicalutamide and compounds being tested is evaluated in vivo using 8 week-old FVB mice which are purchased from Charles River Laboratories. Mice are divided into groups of three for each time points. Two mice are not treated with drug and two other mice are treated with vehicle solution. Each group is treated with 10 mg per kilogram of body weight.

[0095] The drug is dissolved in a mixture 1:5:14 of DMSO : PEG400 : H_2O. (Vehicle solution) and is administered into mice through the tail vein. The animals are warmed under a heat lamp for approximately 20 minutes prior to treatment to dilate their tail vein. Each mouse is placed into a mouse restrainer (Fisher Sci. Cat# 01-288-32A) and is injected with 200 μl of drug in vehicle solution into the dilated tail vein. After drug administration, the animals are euthanized via CO_2 inhalation at different timepoints: 5 mn, 30 mn, 2 h, 6 h, 16 h. Animals are immediately bled after exposure to CO_2 via cardiac puncture (1 ml BD syringe + 27G 5/8 needle). For oral dosage, the drug is dissolved in a mixture 50:10:1:989 of DMSO : Carboxymethylcellulose : Tween80:H_2O before oral administration via a feeding syringe.

[0096] The serum samples are analyzed to determine the drug's concentration by the HPLC which (Waters 600 pump, Waters 600 controller and Waters 2487 detector) is equipped with an Alltima C18 column (3μ, 150 mm×4.6 mm). For example, the compounds being tested can be detected at 254 nm wave length and bicalutamide can be detected at 270 nm wave length.

[0097] The samples for HPLC analysis are prepared according to the following procedure:

- Blood cells are separated from serum by centrifugation.

- To 400 μl of serum are added 80 μl of a 10 μM solution of an internal standard and 520 μl of acetonitrile. Precipitation is watched for.

- The mixture is vortexed for 3 minutes and then placed under ultrasound for 30 minutes.

- The solid particles are filtered off or are separated by centrifugation.

- The filtrate is dried under an argon flow to dryness. The sample is reconstructed to 80 μl with acetonitrile before analyzing by HPLC to determine the drug concentration.

- Standard curve of drug is used to improve accuracy.

[0098] The steady state concentration (Css) of a compound can be determined and compared with that of bicalutamide.

Ranking of Compounds

[0099] To rank the compounds, the following data can be considered: in vitro assays (AR response reporter system in LNCaP cell line, PSA level measurement, MTS mitochondrial assay) and in vivo experiments (tumor size measured directly or by emission induced by luciferase reporter gene, pharmacokinetic assays based on blood plasma levels). Characteristics considered in establishing a ranking can include androgen receptor (AR) antagonism activity, lack of AR agonism in hormone refractory cells, prevention of tumor growth, tumor shrinkage, and pharmacokinetic behavior, with a longer residence time in blood being advantageous.

[00100] Compounds that are highly ranked can be advantageous for use as AR antagonists, and as therapeutic agents for hormone refractory prostate cancer. They may be useful to treat other AR related diseases or conditions such as benign prostate hyperplasia, hair loss, and acne. Highly ranked compounds may also be useful as modulators of other nuclear receptors, such as glucocorticoid receptor, estrogen receptor, and peroxisome proliferator-

activated receptor, and as therapeutic agents for diseases in which nuclear receptors play a role, such as breast cancer, ovarian cancer, diabetes, cardiac diseases, and metabolism related diseases. They may be useful in assays, e.g., as standards, or as intermediates or prodrugs.

[00101] The compounds presented in this application can be superior to bicalutamide in treating prostate cancer.

Pharmaceutical Compositions and Administration

[00102] The compounds of the invention are useful as pharmaceutical compositions prepared with a therapeutically effective amount of a compound of the invention, as defined herein, and a pharmaceutically acceptable carrier or diluent.

[00103] The diarylhydantoin compounds of the invention can be formulated as pharmaceutical compositions and administered to a subject in need of treatment, for example a mammal, such as a human patient, in a variety of forms adapted to the chosen route of administration, for example, orally, nasally, intraperitoneally, or parenterally, by intravenous, intramuscular, topical or subcutaneous routes, or by injection into tissue.

[00104] Thus, diarylhydantoin compounds of the invention may be systemically administered, e.g., orally, in combination with a pharmaceutically acceptable vehicle such as an inert diluent or an assimilable edible carrier, or by inhalation or insufflation. They may be enclosed in hard or soft shell gelatin capsules, may be compressed into tablets, or may be incorporated directly with the food of the patient's diet. For oral therapeutic administration, the diarylhydantoin compounds may be combined with one or more excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. The diarylhydantoin compounds may be combined with a fme inert powdered carrier and inhaled by the subject or insufflated. Such compositions and preparations should contain at least 0.1% diarylhydantoin compounds. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 2% to about 60% of the weight of a given unit dosage form. The amount of diarylhydantoin compounds in such therapeutically useful compositions is such that an effective dosage level will be obtained.

[00105] The tablets, troches, pills, capsules, and the like may also contain the following: binders such as gum tragacanth, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, fructose, lactose or aspartame or a flavoring agent such as peppermint, oil of wintergreen, or cherry flavoring may be added. When the unit dosage form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier, such as a vegetable oil or a polyethylene glycol. Various other materials may be present as coatings or to otherwise modify the physical form of the solid unit dosage form. For instance, tablets, pills, or capsules may be coated with gelatin, wax, shellac or sugar and the like. A syrup or elixir may contain the active compound, sucrose or fructose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavoring such as cherry or orange flavor. Of course, any material used in preparing any unit dosage form should be pharmaceutically acceptable and substantially non-toxic in the amounts employed. In addition, the diarylhydantoin compounds may be incorporated into sustained-release preparations and devices. For example, the diarylhydantoin compounds may be incorporated into time release capsules, time release tablets, and time release pills.

[00106] The diarylhydantoin compounds may also be administered intravenously or intraperitoneally by infusion or injection. Solutions of the diarylhydantoin compounds can be prepared in water, optionally mixed with a nontoxic surfactant. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, triacetin, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations can contain a preservative to prevent the growth of microorganisms.

[00107] The pharmaceutical dosage forms suitable for injection or infusion can include sterile aqueous solutions or dispersions or sterile powders comprising the diarylhydantoin compounds which are adapted for the extemporaneous preparation of sterile injectable or infusible solutions or dispersions, optionally encapsulated in liposomes. In all cases, the ultimate dosage form should be sterile, fluid and stable under the conditions of manufacture and storage. The liquid carrier or vehicle can be a solvent or liquid dispersion medium comprising, for example, water, ethanol, a polyol (for example, glycerol, propylene glycol, liquid polyethylene glycols, and the like), vegetable oils, nontoxic glyceryl esters, and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the formation of

liposomes, by the maintenance of the required particle size in the case of dispersions or by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, buffers or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[00108] Sterile injectable solutions are prepared by incorporating the diarylhydantoin compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filter sterilization. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze drying techniques, which yield a powder of the active ingredient plus any additional desired ingredient present in the previously sterile-filtered solutions.

[00109] For topical administration, the diarylhydantoin compounds may be applied in pure form. However, it will generally be desirable to administer them to the skin as compositions or formulations, in combination with a dermatologically acceptable carrier, which may be a solid or a liquid.

[00110] Useful solid carriers include finely divided solids such as talc, clay, microcrystalline cellulose, silica, alumina and the like. Other solid carriers include nontoxic polymeric nanoparticles or microparticles. Useful liquid carriers include water, alcohols or glycols or water/alcohol/glycol blends, in which the diarylhydantoin compounds can be dissolved or dispersed at effective levels, optionally with the aid of non-toxic surfactants. Adjuvants such as fragrances and additional antimicrobial agents can be added to optimize the properties for a given use. The resultant liquid compositions can be applied from absorbent pads, used to impregnate bandages and other dressings, or sprayed onto the affected area using pump-type or aerosol sprayers.

[00111] Thickeners such as synthetic polymers, fatty acids, fatty acid salts and esters, fatty alcohols, modified celluloses or modified mineral materials can also be employed with liquid carriers to form spreadable pastes, gels, ointments, soaps, and the like, for application directly to the skin of the user.

[00112] Examples of useful dermatological compositions which can be used to deliver the diarylhydantoin compounds to the skin are known to the art; for example, see Jacquet et al. (U.S. Pat. No. 4,608,392), Geria (U.S. Pat No. 4,992,478), Smith et al. (U.S. Pat. No. 4,559,157) and Wortzman (U.S. Pat. No. 4,820,508), all of which are hereby incorporated by reference.

[00113] Useful dosages of the compounds of formula I can be determined by comparing their in vitro activity, and in vivo activity in animal models. Methods for the extrapolation of effective dosages in mice, and other animals, to humans are known to the art; for example, see U.S. Pat. No. 4,938,949, which is hereby incorporated by reference.

[00114] For example, the concentration of the diarylhydantoin compounds in a liquid composition, such as a lotion, can be from about 0.1-25% by weight, or from about 0.5-10% by weight. The concentration in a semi-solid or solid composition such as a gel or a powder can be about 0.1-5% by weight, or about 0.5-2.5% by weight.

[00115] The amount of the diarylhydantoin compounds required for use in treatment will vary not only with the particular salt selected but also with the route of administration, the nature of the condition being treated and the age and condition of the patient and will be ultimately at the discretion of the attendant physician or clinician.

[00116] Effective dosages and routes of administration of agents of the invention are conventional. The exact amount (effective dose) of the agent will vary from subject to subject, depending on, for example, the species, age, weight and general or clinical condition of the subject, the severity or mechanism of any disorder being treated, the particular agent or vehicle used, the method and scheduling of administration, and the like. A therapeutically effective dose can be determined empirically, by conventional procedures known to those of skill in the art. See, *e.g.*, *The Pharmacological Basis of Therapeutics*, Goodman and Gilman, eds., Macmillan Publishing Co., New York. For example, an effective dose can be estimated initially either in cell culture assays or in suitable animal models. The animal model may also be used to determine the appropriate concentration ranges and routes of administration. Such information can then be used to determine useful doses and routes for administration in humans. A therapeutic dose can also be selected by analogy to dosages for comparable therapeutic agents.

[00117] The particular mode of administration and the dosage regimen will be selected by the attending clinician, taking into account the particulars of the case *(e.g.,* the subject, the disease, the disease state involved, and whether the treatment is prophylactic). Treatment may involve daily or multi-daily doses of compound(s) over a period of a few days to months, or even years.

[00118] In general, however, a suitable dose will be in the range of from about 0.001 to about 100 mg/kg, e.g., from about 0.01 to about 100 mg/kg of body weight per day, such as above about 0.1 mg per kilogram, or in a range of from about 1 to about 10 mg per kilogram body weight of the recipient per day. For example, a suitable dose may be about 0.1 mg/kg, 1 mg/kg, 10 mg/kg, or 50 mg/kg of body weight per day.

[00119] The diarylhydantoin compounds are conveniently administered in unit dosage form; for example, containing 0.05 to 10000 mg, 0.5 to 10000 mg, 5 to 1000 mg, or about 100 mg of active ingredient per unit dosage form.

[00120] The diarylhydantoin compounds can be administered to achieve peak plasma concentrations of, for example, from about 0.5 to about 75 μM, about 1 to 50 μM, about 2 to about 30 μM, or about 5 to about 25 μM. Exemplary desirable plasma concentrations include at least or no more than 0.01, 0.025, 0.05, 0.1, 0.25, 0.5, 1, 5, 10, 25, 50, 75, 100 or 200 μM. For example, plasma levels may be from about 1 to 100 micromolar or from about 10 to about 25 micromolar. This may be achieved, for example, by the intravenous injection of a 0.05 to 5% solution of the diarylhydantoin compounds, optionally in saline, or orally administered as a bolus containing about 1-100 mg of the diarylhydantoin compounds. Desirable blood levels may be maintained by continuous infusion to provide about 0.00005 - 5 mg per kg body weight per hour, for example at least or no more than 0.00005, 0.0005, 0.005, 0.05, 0.5, or 5 mg/kg/hr. Alternatively, such levels can be obtained by intermittent infusions containing about 0.0002 - 20 mg per kg body weight, for example, at least or no more than 0.0002, 0.002, 0.02, 0.2, 2, 20, or 50 mg of the diarylhydantoin compounds per kg of body weight.

[00121] The diarylhydantoin compounds may conveniently be presented in a single dose or as divided doses administered at appropriate intervals, for example, as two, three, four or more sub-doses per day. The sub-dose itself may be further divided, e.g., into a number of discrete loosely spaced administrations; such as multiple inhalations from an insufflator.

[00122] A number of the above-identified compounds exhibit little or no agonistic activities with respect to hormone refractory prostate cancer cells. Because these compounds are strong androgen receptor (AR) inhibitors, they can be used not only in treating prostate cancer, but also in treating other AR related diseases or conditions such as benign prostate hyperplasia, hair loss, and acne. Because AR belongs to the family of nuclear receptors, these compounds may serve as scaffolds for drug synthesis targeting other nuclear receptors, such as estrogen receptor and peroxisome proliferator-activated receptor. Therefore, they may be further developed for other diseases such as breast cancer, ovarian cancer, diabetes, cardiac diseases, and metabolism related diseases, in which nuclear receptors play a role.

[00123] The embodiments illustrated and discussed in this specification are intended only to teach those skilled in the art the best way known to the inventors to make and use the invention. Nothing in this specification should be considered as limiting the scope of the present invention. All examples presented are representative and non-limiting. The above-described embodiments of the invention may be modified or varied, without departing from the invention, as appreciated by those skilled in the art in light of the above teachings. It is therefore to be understood that, within the scope of the claims and their equivalents, the invention may be practiced otherwise than as specifically described.

WE CLAIM:

1. A compound having the formula



wherein R_1 and R_2 together comprise eight or fewer carbon atoms and are selected from the group consisting of alkyl, substituted alkyl, and, together with the carbon to which they are linked, a cycloalkyl or substituted cycloalkyl group,

wherein R_3 is selected from the group consisting of hydrogen, cyano, formyl,


, 
, 
, and 
,

wherein R_4 is selected from the group consisting of hydrogen, F, Cl, Br, and I,

wherein R_{11} and R_{12} are independently selected from the group consisting of hydrogen and methyl,

wherein R_{13} is selected from the group consisting of hydrogen and $-NR_{14}R_{15}$, and

wherein R_{14} and R_{15} are independently selected from the group consisting of hydrogen and methyl.

2. The compound of claim 1,

wherein R_1 and R_2 are independently methyl or, together with the carbon to which they are linked, cyclobutyl or cyclopentyl,

wherein R_3 is selected from the group consisting of cyano, formyl,



,


,


, and


,

wherein R_4 is selected from the group consisting of hydrogen and fluorine,

wherein R_{11} and R_{12} are both hydrogen or both methyl, and

wherein R_{13} is selected from the group consisting of $-NH(CH_3)$ and $-N(CH_3)_2$.

3. The compound of claim 2,

wherein R_{11} and R_{12} are both hydrogen and

provided that when R_4 is hydrogen and when R_1 and R_2 together with the carbon to which they are linked are cyclobutyl, then R_3 is not cyano and is not 

.

4. The compound of claim 1,

provided that when R_4, R_{11}, and R_{12} are all hydrogen and when R_1 and R_2 together with the carbon to which they are linked are cyclobutyl, then R_3 is not cyano and is not



.

5. The compound of claim 1,

wherein R_1 and R_2 together comprise four or fewer carbon atoms and are selected from the group consisting of alkyl and, together with the carbon to which they are linked, cycloalkyl.

6. The compound of claim 1,

wherein R_1 and R_2 are independently selected from the group consisting of hydrogen and methyl.

7. The compound of claim 1,

wherein R_1 and R_2, together with the carbon to which they are linked, are selected from the group consisting of cyclobutyl and cyclopropyl.

8. The compound of claim 1,

wherein R_3 is selected from the group consisting of cyano and formyl.

9. The compound of claim 1,

wherein R_3 is selected from the group consisting of



,


, and


.

10. The compound of claim 1,

wherein R_3 is selected from the group consisting of



and


.

11. The compound of claim 1, wherein R_4 is hydrogen.

12. The compound of claim 1, wherein R_4 is fluorine.

13. The compound of claim 1, wherein R_4 is chlorine.

14. The compound of claim 1, having the formula



[ND-1].

15. The compound of claim 1, having the formula



[ND-2].

16. The compound of claim 1, having the formula



wherein R_{21} is selected from the group consisting of



[ND-3] and



[ND-13].

17. The compound of claim 1, having the formula



wherein R_{21} is selected from the group consisting of



[ND-11] and



[ND-12].

18. The compound of claim 1, having the formula



[ND-10].

19. The compound of claim 1, having the formula



wherein R_{22} is hydrogen [ND-7] or methyl [ND-8].

20. The compound of claim 1, having the formula



[ND-9].

21. The compound of claim 1, having the formula



[ND-14].

22. A pharmaceutical composition comprising a therapeutically effective amount of a compound according to claim 1 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or diluent.

23. A method for treating a hyperproliferative disorder comprising administering a pharmaceutical composition of claim 22 to a subject in need of such treatment, thereby treating the hyperproliferative disorder.

24. The method of claim 23, wherein the composition is administered at a dosage of the compound in the range of from about 0.001 mg per kg body weight per day to about 100 mg per kg body weight per day.

25. The method of claim 23, wherein the composition is administered at a dosage of the compound in the range of from about 0.01 mg per kg body weight per day to about 100 mg per kg body weight per day.

26. The method of claim 23, wherein the composition is administered at a dosage of the compound in the range of from about 0.1 mg per kg body weight per day to about 10 mg per kg body weight per day.

27. The method of claim 23, wherein the composition is administered at a dosage of the compound of about 1 mg per kg body weight per day.

28. The method of claim 23, wherein the hyperproliferative disorder is hormone refractory prostate cancer.

29. The method of claim 23, wherein the composition is administered by intravenous injection, by injection into tissue, intraperitoneally, orally, or nasally.

30. The method of claim 23, wherein the composition is administered orally.

31. The method of claim 22, wherein the composition has a form selected from the group consisting of a solution, dispersion, suspension, powder, capsule, tablet, pill, time release capsule, time release tablet, and time release pill.

32. The method of claim 22, wherein the composition has a form selected from the group consisting of a capsule, tablet, and pill.

33. A method of synthesizing a diaryl compound of formula:



comprising mixing Compound I



Compound I

with Compound II



Compound II

in a first polar solvent to form a mixture,

adding a second polar solvent, the same as or different from the first polar solvent, and an aqueous acid to the mixture,

refluxing the mixture,

cooling the mixture and combining with water, and

separating the diaryl compound from the mixture,

wherein R_{31} is selected from the group consisting of cyano, carboxy,


(methylcarbamoyl),

(dimethylcarbamoyl), and


,

wherein Boc is *t*-butoxycarbonyl,

wherein R_{32} is selected from the group consisting of


and

,

wherein R_1 and R_2 together comprise eight or fewer carbon atoms and are selected from the group consisting of alkyl, substituted alkyl, and, together with the carbon to which they are linked, a cycloalkyl or substituted cycloalkyl group,

wherein R_3 is selected from the group consisting of hydrogen, cyano, formyl,


,

,

, and

,

wherein R_4 is selected from the group consisting of hydrogen, F, Cl, Br, and I,

wherein R_{11} and R_{12} are independently selected from the group consisting of hydrogen and methyl,

wherein R_{13} is selected from the group consisting of hydrogen and $-NR_{14}R_{15}$, and

wherein R_{14} and R_{15} are independently selected from the group consisting of hydrogen and methyl.

34. The compound of claim 1, wherein the compound has substantial androgen receptor antagonist activity and no substantial agonist activity on hormone refractory prostate cancer cells.

35. A method comprising:

providing at least one compound according to claim 1;

measuring inhibition of androgen receptor activity for the compound and determining if the inhibition is above a first predetermined level,

measuring stimulation of androgen receptor activity in hormone refractory cancer cells for the compound and determining if the stimulation is below a second predetermined level,

selecting the compound if the inhibition is above the first predetermined level and the stimulation is below the second predetermined level.

36. The method of claim 35, wherein the predetermined levels are those of bicalutamide.

37. The method of claim 35, wherein the step of measuring inhibition comprises measuring inhibitory concentration (IC50) in an AR response reporter system or a prostate specific antigen secreting system.

38. The method of claim 35, wherein the step of measuring stimulation comprises measuring fold induction by increasing concentrations in an AR response reporter system or a prostate specific antigen secreting system.

39. The method of claim 35, wherein the steps of measuring inhibition and/or stimulation comprise measuring an effect of the compound on tumor growth in an animal.