

**(12) PATENT**  
**(19) AUSTRALIAN PATENT OFFICE**

**(11)** Application No. **AU 199961999 B2**  
**(10)** Patent No. **756233**

(54) Title  
Pharmaceutical compositions based on alpha-cyclodextrin for the oral administration of LH-RH analogues

(51)<sup>6</sup> International Patent Classification(s)  
A61K 038/09 A61K 047/40

(21) Application No: 199961999 (22) Application Date: 1999.09.23

(87) WIPO No: W000/18423

(30) Priority Data

(31) Number	(32) Date	(33) Country
98402403	1998.09.30	EP

(43) Publication Date : 2000.04.17

(43) Publication Journal Date : 2000.06.08

(44) Accepted Journal Date : 2003.01.09

(71) Applicant(s)  
Laboratoire Theramex

(72) Inventor(s)  
Remi Delansorne; Paule Bonnet; Jacques Paris

(74) Agent/Attorney  
SPRUSON and FERGUSON,GPO Box 3898,SYDNEY NSW 2001

61774/99



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>A61K 38/09, 47/40</b>		A1	(11) International Publication Number: <b>WO 00/18423</b>
			(43) International Publication Date: 6 April 2000 (06.04.00)
(21) International Application Number: PCT/EP99/07389 (22) International Filing Date: 23 September 1999 (23.09.99) (30) Priority Data: 98402403.4 30 September 1998 (30.09.98) EP (71) Applicant (for all designated States except US): LABO- RATOIRE THERAMEX [MC/MC]; 6, avenue Prince Héréditaire Albert, MC-98000 Monaco (MC). (72) Inventors; and (75) Inventors/Applicants (for US only): DELANSORNE, Rémi [FR/FR]; 7bis, avenue Edith Cavell, F-06000 Nice (FR). BONNET, Paule [FR/FR]; Chemin Gliena, 37, route de Sospel, F-06500 Menton (FR). PARIS, Jacques [FR/FR]; 31, avenue Cap de Croix, Bât. E1, F-06100 Nice (FR). (74) Agents: NEVANT, Marc et al.; Cabinet Beau de Loménie, 158, rue de l'Université, F-75340 Paris Cedex 07 (FR).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: PHARMACEUTICAL COMPOSITIONS BASED ON ALPHA-CYCLODEXTRIN FOR THE ORAL ADMINISTRATION OF LH-RH ANALOGUES			
(57) Abstract			
The invention relates to the use of $\alpha$ -cyclodextrin or a derivative thereof for the preparation of pharmaceutical compositions for the oral administration of LH-RH peptide analogues. The invention also relates to oral pharmaceutical compositions containing LH-RH peptide analogues in combination with $\alpha$ -cyclodextrin or a derivative thereof.			

**Pharmaceutical compositions based on alpha-cyclodextrin for the oral administration of LH-RH analogues**

The present invention relates to the pharmaceutical field. More specifically, the invention relates to the use of  $\alpha$ -cyclodextrin or derivatives thereof for the preparation of pharmaceutical compositions for the oral administration of LH-RH (luteinizing hormone - releasing hormone) peptide analogues. The invention also relates to oral pharmaceutical compositions containing LH-RH peptide analogues in combination with  $\alpha$ -cyclodextrin.

Natural and modified cyclodextrins (CDs) are well known ingredients used in a large variety of pharmaceutical preparations taking advantage of one or several of their properties relating to drug solubilization and stabilization (Loftsson and Brewster, 1996, *J. Pharm. Sci.*, **85**(10): 1017-1025) or to overall improvement of *in vivo* drug delivery (Rajewski and Stella, 1996, *J. Pharm. Sci.*, **85**(11): 1142-1169). CDs are cyclic oligosaccharides containing at least 6  $\alpha$ -D-(+)-glucopyranose units attached by  $\alpha$ (1-4) glucoside bonds (Nash, *Handbook of Pharmaceutical Excipients*, ed. by Wade and Weller, 1994, American Pharmaceutical Association, Washington, and The Pharmaceutical Press, London, pp 145-148); the three most common CDs are  $\alpha$ -,  $\beta$ - and  $\gamma$ -CD which consist of 6, 7 and 8 sugar units, respectively. Numerous derivatives of each type of CD can be obtained by random or controlled modifications of one, several or all free hydroxyl groups of the sugar moieties.

LH-RH is a neurohormone produced by hypothalamic neurons and secreted in the pituitary portal vasculature to stimulate the release of luteinizing hormone (LH) and follicle stimulating hormone (FSH) by the pituitary gland. In turn, LH and FSH regulate the endocrine and germinal functions of the ovary in the female and of the testis in the male. LH-RH is a peptide of the following structure: pGlu-His-Trp-Ser-Tyr-Gly-Leu-Arg-Pro-Gly-NH<sub>2</sub>. Numerous normal or reduced-size, linear or cyclic peptide analogues of LH-RH incorporating natural, unusual or chemically-modified amino-acids have been synthesized over the years to yield potent agonist or antagonistic properties (Karten and Rivier, 1986, *Endocr. Rev.*, **7**(1): 44-66; Dutta, 1988, *Drugs of the Future*, **13**(8): 761-787; Kutscher et al., 1997, *Angew. Chem. Int. Ed. Engl.*, **36**: 2148-2161). Due to their total or partial peptide structure, however, all these analogues show poor oral bioavailability and bioactivity.

To date, only non-oral administration of LH-RH peptide analogues, has been reported. For example, Matsubara et al. (1996, *J. Pharm. Sci.*, **84**(11) : 1295-1300) describe a nasal formulation of buserelin, based on dimethyl- $\beta$ -CD, with improved bioavailability.


There is therefore a need, for the patients' comfort, to provide formulations which enable the oral administration of LH-RH peptide analogues.

It has now surprisingly been found that  $\alpha$ -CD or its derivatives, enhance the biological activity of LH-RH peptide analogues when orally administered.

5 Thus, according to one of its feature, the invention relates to the use of  $\alpha$ -cyclodextrin or derivatives thereof for the preparation of pharmaceutical compositions for the oral administration of LH-RH peptide analogues.

Examples of LH-RH peptide analogues which can be used within the scope of the invention include those described in International patent applications  
10 WO 98/21229 and WO 98/55505, the content of which is incorporated by reference, as well as standard agonists and antagonists of LH-RH, such as for example buserelin, nafarelin, leuprorelin, goserelin, histrelin, triptorelin, deslorelin, lutrelin, avorelin, cetorelix, antide, ganirelix, azaline B, antarelix, detirelix, ramorelix, teverelix or abarelix.

15 Preferably, these peptide analogues have the formula (SEQ ID N° : 1) :



in which :

- A1 is pGlu ; D-pGlu ; Sar ; AcSar ; Pro or a derivative thereof such as AcPro, ForPro, OH-Pro, Ac-OH-Pro, dehydro-Pro or Ac-dehydro-Pro ; Ser ; D-Ser ; Ac-D-Ser ; Thr ; D-Thr ; Ac-D-Thr ; or an aromatic D-amino acid which may be acylated, such as D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-diphenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe, D-HPhe, D-Tyr, D-HTyr, and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;
- 25 - A2 is a direct bond ; His ; or an aromatic D-amino acid such as D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-diphenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe, D-HPhe, D-Tyr, D-HTyr and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;
- 30 - A3 is an aromatic L- or D-amino acid such as Phe, HPhe, Tyr, HTyr, Trp, 2MeTrp, Nal, 1Nal, diphenyl-Ala, Bal, Pal, 4Pal or Qal, where Phe, HPhe, Tyr, HTyr and Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;

- A5 is an aromatic L-amino acid such as Phe, HPhe, Tyr, HTyr, Trp, 2MeTrp, Nal, 1Nal, diphenyl-Ala, Bal, Pal, 4Pal or Qal, where Phe, HPhe, Tyr, HTyr and Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups and/or N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms; or a basic L- or D-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HCit, APhe or ACha, where Arg and HArg may be N-substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group on one or both nitrogen atoms, and where Orn, Lys, HLys, APhe and ACha may be N-substituted by one or two (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by an aminotriazolyl or a nicotinoyl, isonicotinoyl, 6-methyl-nicotinoyl, glycyl-nicotinoyl, nicotinyl-azaglycyl, furyl, glycyl-furyl, furyl-azaglycyl, pyrazinyl, pyrazinyl-carbonyl, picolinoyl, 6-methyl-picolinoyl, shikimyl, shikimyl-glycyl, Fmoc or Boc group;
- A6 is Gly; (S)-spirolactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>t</sup>); D-Asp(OBu<sup>t</sup>); D-Glu(OBu<sup>t</sup>); D-Thr(OBu<sup>t</sup>); D-Cys(OBu<sup>t</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid such as azaGly or azaAla; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain such as D-Ala, D-Abu, D-Aib, D-3Aib, D-Val, D-Nva, D-Leu, D-Ile, D-Tle, D-Nle, D-Hol, D-Npg, D-CPa, D-Cpa, D-Cba or D-Cha; an aromatic D-amino acid such as D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-diphenyl-Ala, D-anthryl-Ala, D-phenanthryl-Ala, D-benzhydryl-Ala, D-fluorenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe, D-HPhe, D-Tyr, D-HTyr and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HCit, APhe or ACha, where Arg and HArg may be N-substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group on one or both nitrogen atoms, and where Orn, Lys, HLys, APhe and ACha may be N-substituted by one or two (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by an aminotriazolyl or a nicotinoyl, isonicotinoyl, 6-methyl-nicotinoyl, glycyl-nicotinoyl, nicotinyl-azaglycyl, furyl, glycyl-furyl, furyl-azaglycyl, pyrazinyl, pyrazinyl-carbonyl, picolinoyl, 6-methyl-picolinoyl, shikimyl, shikimyl-glycyl, Fmoc or Boc group;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms such as Ala, Abu, Aib, 3Aib, Val, Nva, Leu, Ile, Tle, Nle, Hol, Npg, CPa, Cpa,

Cba, Cha or Ada, which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms ;

- A8 is a basic L- or D-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HCit, APhe or ACha, where Arg or HArg may be N-substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group on one or both nitrogen atoms, and where Orn, Lys, HLys, APhe or ACha may be N-substituted by one or two (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by an aminotriazolyl or a nicotinoyl, isonicotinoyl, 6-methyl-nicotinoyl, glycylic-nicotinoyl, nicotinyl-azaglycyl, furyl, glycylic-furyl, furyl-azaglycyl, pyrazinyl, pyrazinyl-carbonyl, picolinoyl, 6-methyl-picolinoyl, shikimyl, shikimyl-glycyl, Fmoc or Boc group;

- Z is GlyNH<sub>2</sub> ; D-AlaNH<sub>2</sub> ; azaGlyNH<sub>2</sub> ; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms ; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl;

- as well as their pharmaceutically acceptable salts.

In the present description the term "(C<sub>1</sub>-C<sub>4</sub>)alkyl" denotes methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl and t-butyl groups.

The term "(C<sub>1</sub>-C<sub>6</sub>)alkyl" denotes methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, s-pentyl, t-pentyl and hexyl groups.

- The term "(C<sub>1</sub>-C<sub>8</sub>)alkyl" denotes methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, i-pentyl, s-pentyl, t-pentyl, hexyl, heptyl and octyl groups ;

The term "(C<sub>1</sub>-C<sub>4</sub>)alkoxy" denotes a group -OR where R is a (C<sub>1</sub>-C<sub>4</sub>)alkyl.


The term "(C<sub>2</sub>-C<sub>7</sub>)acyl" denotes a group -COR where R is a (C<sub>1</sub>-C<sub>6</sub>)alkyl.

- The term "(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl" denotes cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl groups.

The term "sugar moiety" denotes D- or L-pentoses or hexoses and their amino-derivatives.

- The term "LH-RH analogues" denotes peptides in which at least one amino acid has been modified in the sequence of LH-RH.

The term "(S)spirolactam-Pro" denotes the residue of the formula :



The term "oral administration" denotes the delivery of the peptide analogues of the invention to the gastrointestinal tract by means of an oral formulation or composition.


- 5 Peptidomimetic analogues of LH-RH defined by the absence of at least one peptide amide bond, as exemplified in the latest review by Kutscher et al. (1997, *Angew. Chem. Int. Ed. Engl.*, **36**: 2148-2161), are not considered within the scope of the present invention.

In the present description and in the claims, the following abbreviations are  
10 used :

Abu : 2-aminobutyric acid	Ac : acetyl
ACha : aminocyclohexylalanine	Aib : 2-aminoisobutyric acid
3Aib : 3-aminoisobutyric acid	Ala : alanine
AlaNH <sub>2</sub> : alaninamide	APhe : p-aminophenylalanine
Arg : arginine	Asp : aspartic acid
azaAla : aza-alanine	azaGly : aza-glycine
azaGlyNH <sub>2</sub> : azaglycinamide	Bal : benzothienylalanine
Boc : <i>tert</i> -butoxycarbonyl	Cba : cyclobutylalanine
Cha : cyclohexylalanine	Cit : citrulline
CPa : cyclopropylalanine	Cpa : cyclopentylalanine
Fmoc : fluorenylmethoxycarbonyl	For : formyl
Glu : glutamic acid	Gly : glycine
GlyNH <sub>2</sub> : glycineamide	HArg : homoarginine
HCit : homocitrulline	His : histidine
HLys : homolysine	Hol : homoleucine
Ile : isoleucine	IprLys : N <sup>ε</sup> -isopropyllysine
Leu : leucine	Lys : lysine
MeSer : N-methylserine	Met : methionine
Nal : 3-(2-naphtyl)alanine	1Nal : 3-(1-naphtyl)alanine
NEt : N-ethylamide	NicLys : N <sup>ε</sup> -nicotinoyllysine

Nle : norleucine	Npg : neopentylglycine
Nva : norvaline	OBu <sup>t</sup> : <i>tert</i> -butoxy
OBzl : benzyl ester	Orn : ornithine
Pal : 3-(3-pyridyl)alanine	pClPhe : 3-(4-chlorophenyl)alanine
Pen : penicillamine	pGlu : pyroglutamic acid
Phe : phenylalanine	Pro : proline
Qal : 3-(3-quinolyl)alanine	Sar : sarcosine
Ser : serine	(S-Me)Pen : S-methyl-penicillamine
(S-Et)Pen : S-ethyl-penicillamine	Thr : threonine
Tle : <i>tert</i> -leucine	Trp : tryptophan
Tyr : tyrosine	Val : valine
Ada : adamantylalanine	HPhe : homophenylalanine
MeNpg : N-methylneopentylglycine	4Pal : 3-(4-pyridyl)alanine
HTyr : homotyrosine	2MeTrp : 2-methyltryptophan
Bzl : benzyl	SPL : (S)spiro lactam-Pro
Asn : asparagine	MeLeu : N-methylleucine
MeTyr : N-methyltyrosine	MeHTyr : N-methylhomotyrosine

A preferred group of peptide analogues (A) comprises the peptides of the formula (SEQ ID N° : 2) :



5 in which:

- A1 is pGlu, Sar or AcSar;

- A3 is an aromatic L-amino acid such as Phe, HPhe, Tyr, HTyr, Trp, 2MeTrp, Nal, 1Nal, diphenyl-Ala, Bal, Pal, 4Pal or Qal, where Phe, HPhe, Tyr, HTyr and Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;

10

- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;

- A5 is an aromatic L-amino acid such as Phe, HPhe, Tyr, HTyr, Trp, 2MeTrp, Nal, 1Nal, diphenyl-Ala, Bal, Pal, 4Pal or Qal, where Phe, HPhe, Tyr, HTyr and Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or

15 trifluoromethyl groups;



- A6 is Gly ; D-Pro ; (S)-spirolactam-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>t</sup>) ; D-Asp(OBu<sup>t</sup>) ; D-Glu(OBu<sup>t</sup>) ; D-Thr(OBu<sup>t</sup>) ; D-Cys(OBu<sup>t</sup>) ; D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aza-amino acid such as azaGly or azaAla ; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain such as D-Ala, D-Abu, D-Aib, D-3Aib, D-Val, D-Nva, D-Leu, D-Ile, D-Tle, D-Nle, D-Hol, D-Npg, D-CPa, D-Cpa, D-Cba or D-Cha ; an aromatic D-amino acid such as D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-diphenyl-Ala, D-anthryl-Ala, D-phenanthryl-Ala, D-benzhydryl-Ala, D-fluorenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe, DHPhe, D-Tyr, D-HTyr and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups ; D-cyclohexadienyl-Gly ; D-perhydronaphtyl-Ala ; D-perhydrodiphenyl-Ala ; or a basic D-amino acid such as D-Arg, D-HArg, D-Orn, D-Lys, D-HLys, D-Cit, D-HCit, D-APhe optionally substituted by an aminotriazolyl group or D-ACha, where D-Arg and D-HArg may be substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by a Fmoc or Boc group;


- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms such as Ala, Abu, Aib, 3Aib, Val, Nva, Leu, Ile, Tle, Nle, Hol, Npg, CPa, Cpa, Cba, Cha or Ada, which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is a basic L-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HCit, APhe optionally substituted by an aminotriazolyl group, or ACha;

- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> ; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms ; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl;

as well as their pharmaceutically acceptable salts.

Among the peptide analogues of formula (I), those of the formula (SEQ ID N° 3) :



in which:

- A3 and A5 are aromatic L-amino acids as defined for (I) ;
- A6 is as defined for (I) ;

- A7 is Leu, Tie, Nle, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several fluorine atoms;


- Z is as defined for (I) ;

as well as their pharmaceutically acceptable salts,

5 are preferred.

Especially preferred are the peptide analogues of the formula (SEQ ID N° ;

4) :



in which:


10 - A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Bal, Pal, 4Pal, or pClPhe ;

- A6 is (S)-spiro lactam-Pro ; Gly; D-Pro ; D-Ser(OBu<sup>t</sup>) ; D-Asp(OBu<sup>t</sup>) ; D-Glu(OBu<sup>t</sup>) ; D-Thr(OBu<sup>t</sup>) ; D-Cys(OBu<sup>t</sup>) ; D-His or D-His(Bzl) ; D-Ala, D-Leu, D-Tie, D-Nle, D-Hol, D-Npg or D-Cha ; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp,  
15 D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydrodiphenyl-Ala ; or D-APhe optionally substituted by an aminotriazolyl group ;

- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;

- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

20 Also especially preferred are the peptide analogues of the formula (SEQ ID N° : 5) :



in which:

- A6 is (S)-spiro lactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>t</sup>) or D-Trp;  
25

- A7 is Leu, MeLeu, Npg or MeNpg;

- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

The peptide analogues of formula (I) to (IV) in which A7 is Npg are especially preferred.

30 Representative peptide analogues of formula (I) to (IV) include leuporelin, [Npg<sup>7</sup>]-leuporelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, buserelin and [Npg<sup>7</sup>]-buserelin.

Another preferred group of peptide analogues (A) comprises the peptides of the formula (SEQ ID N° : 6) :

## A1-A2-A3-A4-A5-A6-A7-A8-Pro-Z (I')

in which:

- A1 is pGlu ; D-pGlu ; Sar ; AcSar ; Pro or a derivative thereof such as AcPro, ForPro, OH-Pro, Ac-OH-Pro, dehydro-Pro or Ac-dehydro-Pro ; Ser ; D-Ser ; Ac-D-Ser ; Thr ; D-Thr ; Ac-D-Thr ; or an aromatic D-amino acid which may be acylated, preferably acetylated, such as D-Phe, D-HPhe, D-Tyr, D-Trp, D-Nal, D-1Nal, D-diphenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;
- A2 is a direct bond or an aromatic D-amino acid such as D-Phe, D-HPhe, D-Tyr, D-Trp, D-Nal, D-1Nal, D-diphenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;
- A3 is an aromatic L- or D-amino acid such as Phe, HPhe, Tyr, Trp, Nal, 1Nal, diphenyl-Ala, Bal, Pal, 4Pal or Qal, where Phe and Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid such as Phe, HPhe, Tyr, HTyr, Trp, Nal, 1Nal, diphenyl-Ala, Bal, Pal, 4Pal or Qal, where Phe, Tyr, HTyr and Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups and/or N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms; or a basic L- or D-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HCit, APhe or ACha, where Arg and HArg may be N-substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group on one or both nitrogen atoms, and where Orn, Lys, HLys, APhe and ACha may be N-substituted by one or two (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by a nicotinoyl, isonicotinoyl, 6-methyl-nicotinoyl, glycyl-nicotinoyl, nicotinyl-azaglycyl, furyl, glycyl-furyl, furyl-azaglycyl, pyrazinyl, pyrazinyl-carbonyl, picolinoyl, 6-methyl-picolinoyl, shikimyl, shikimyl-glycyl, Fmoc or Boc group;
- A6 is Gly ; (S)-spirolactam-Pro ; D-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Asn ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>t</sup>) ; D-Asp(OBu<sup>t</sup>) ; D-Glu(OBu<sup>t</sup>) ; D-Thr(OBu<sup>t</sup>) ; D-Cys(OBu<sup>t</sup>) ; D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain such as D-Ala, D-Abu, D-Aib, D-3Aib, D-Val, D-Nva, D-Leu, D-Ile, D-Tle, D-Nle, D-Hol, D-

Npg, D-CPa, D-Cpa, D-Cba or D-Cha ; an aromatic D-amino acid such as D-Phe, D-HPhe, D-Tyr, D-Trp, D-Nal, D-1Nal, D-diphenyl-Ala, D-anthryl-Ala, D-phenanthryl-Ala, D-benzhydryl-Ala, D-fluorenyl-Ala, D-Bal, D-Pal, D-4Pal or D-Qal, where D-Phe and D-Trp may be substituted by one or more halogens, (C<sub>1</sub>-C<sub>4</sub>)alkyl, (C<sub>1</sub>-C<sub>4</sub>)alkoxy, nitro or trifluoromethyl groups ; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydrodiphenyl-Ala ; or a basic L- or D-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HClit, APhe or ACha, where Arg and HArg may be N-substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group on one or both nitrogen atoms, and where Orn, Lys, HLys, APhe and ACha may be N-substituted by one or two (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by a nicotinoyl, isonicotinoyl, 6-methyl-nicotinoyl, glycyl-nicotinoyl, nicotinyl-azaglycyl, furyl, glycyl-furyl, furyl-azaglycyl, pyrazinyl, pyrazinyl-carbonyl, picolinoyl, 6-methyl-picolinoyl, shikimyl, shikimyl-glycyl, Fmoc or Boc group;

- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms such as Ala, Abu, Aib, 3Aib, Val, Nva, Leu, Ile, Tle, Nle, Hol, Npg, Cpa, Cba, Cha or Ada, which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms ;

- A8 is a basic L- or D-amino acid such as Arg, HArg, Orn, Lys, HLys, Cit, HClit, APhe or ACha, where Arg and HArg may be N-substituted by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group on one or both nitrogen atoms, and where Orn, Lys, HLys, APhe and ACha may be N-substituted by one or two (C<sub>1</sub>-C<sub>6</sub>)alkyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl groups, or by a nicotinoyl, isonicotinoyl, 6-methyl-nicotinoyl, glycyl-nicotinoyl, nicotinyl-azaglycyl, furyl, glycyl-furyl, furyl-azaglycyl, pyrazinyl, pyrazinyl-carbonyl, picolinoyl, 6-methyl-picolinoyl, shikimyl, shikimyl-glycyl, Fmoc or Boc group;

- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>;

as well as their pharmaceutically acceptable salts.

Among the peptides of formula (I'), those of the formula (SEQ ID N° : 7):

Ac-D-Nal-D-pClPhe-D-Pal-Ser-A5-A6-A7-A8-Pro-D-AlaNH<sub>2</sub> (II')

in which :

- A5 is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys;

- A6 is (S)-spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HClit or D-Asn;

- A7 is Leu, MeLeu, Npg or MeNpg;

- A8 is Arg, NicLys or IprLys;  
and their pharmaceutically acceptable salts,  
are preferred.

5 The peptide analogues of formula (I') and (II') in which A7 is Npg are especially preferred.

Representative peptide analogues of formula (I') and (II') include antide, [Npg<sup>7</sup>]-antide, cetorelix, [Npg<sup>7</sup>]-cetorelix, abarelix and [Npg<sup>7</sup>]-abarelix.

Further preferred peptide analogues comprise those of formula (A) where A6 is as defined therein except D-Asn.

10 Examples of the salts with pharmaceutically acceptable acids are those with mineral acids, such as for example the hydrochloride, hydrobromide, sulfate, phosphate, borate, hydrogensulfate, dihydrogenphosphate or nitrate, and those with organic acids, such as for example the acetate, oxalate, tartrate, succinate, maleate, fumarate, gluconate, citrate, pamoate, malate, ascorbate, benzoate, p-  
15 toluenesulfonate or naphthalenesulfonate.

Examples of the salts with pharmaceutically acceptable bases are those with alkali or alkaline earth metals such as sodium, potassium, calcium or magnesium, and those with organic bases such as amines, trometamol, N-methylglutamine, and the like.

20 The peptides used in the present invention can be prepared by the well-known techniques of peptide chemistry such as for example peptide synthesis in solution or solid phase peptide synthesis. In general, these techniques involve the stepwise addition of one or more amino acids -which may be suitably protected- to a forming peptide chain. Reference can for example be made to *Synthetic Peptides: a user's*  
25 *guide*, ed. by G.A. Grant, 1992, UWBC Biotechnical Resource Series, Washington University Press, Saint-Louis, USA.

Each molecule of  $\alpha$ -CD bears 6 primary hydroxyl groups and 12 secondary hydroxyl groups, respectively corresponding to the 6-OH and to the 2- and 3-OH groups of each of the 6 glucopyranose units. Another general aspect of the present  
30 invention concerns  $\alpha$ -CD and its derivatives which are defined as the result of chemical or biochemical modifications involving a precise or average number between 1 and 18 hydroxyl groups of the  $\alpha$ -CD molecule, in a random or regioselective fashion, with one or several different types of reactions such as oxidation, reduction, alkylation, hydroxyalkylation, esterification with organic or

mineral acids, intramolecular dehydration, tosylation followed by reductive amination or halogen substitution, sugar branching or further polymerization, and their different possible combinations and mixtures. Examples of  $\alpha$ -CD derivatives include  $\alpha$ -CD modified with one or more groups selected from methyl, carboxymethyl, ethyl, butyl, octyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, acetyl, propionyl, butyryl, succinyl, benzoyl, palmityl, sulfonyl, toluenesulfonyl, amino, aminopropyl, glucosyl, maltosyl, dimaltosyl, carboxymethyl ether, sulfobutylether, and phosphate ester.

Preferred  $\alpha$ -CD derivatives according to the invention comprise methylated  $\alpha$ -CD ; hexakis(2,3,6-tri-O-methyl)- $\alpha$ -CD, also known as "permethylated"  $\alpha$ -CD ; carboxymethylated  $\alpha$ -CD and phosphated  $\alpha$ -CD.  $\alpha$ -CD and hexakis(2,3,6-tri-O-methyl)- $\alpha$ -CD are especially advantageous when used in the preparation of the pharmaceutical compositions of the invention.

As mentioned above,  $\alpha$ -CD or its derivatives enhance the biological activity of LH-RH peptide analogues in oral pharmaceutical compositions.

Thus, according to another feature, the invention relates to oral pharmaceutical compositions which comprise as the active principle a LH-RH peptide analogue as defined above in the form of a combination with  $\alpha$ -CD or a derivative thereof, said compositions being intended to be delivered to the gastrointestinal tract.

The peptides according to the general formula (I) exert an agonist activity upon the LH-RH receptors *in vivo*, resulting in the stimulation of LH secretion by the pituitary, which, in males, stimulates the secretion of testosterone by the testis.

Adult male Sprague-Dawley rats were orally administered by gavage an oral formulation comprising leuprorelin (LEU, Bachem), triptorelin (TRI, Bachem), deslorelin (DES, Saxon Biochemicals), goserelin (GOS, Saxon Biochemicals) or the other following example analogues: example 1 ((S)spiro lactam(Pro<sup>6</sup>, Npg<sup>7</sup>), desGly<sup>10</sup>-ProNet<sup>9</sup>)LH-RH), example 2 ([D-Ala<sup>6</sup>, Npg<sup>7</sup>, desGly<sup>10</sup>-ProNet<sup>9</sup>)LH-RH), example 3 ([Npg<sup>7</sup>]leuprorelin), example 4 ([D-Phe<sup>6</sup>, Npg<sup>7</sup>, desGly<sup>10</sup>-ProNet<sup>9</sup>)LH-RH), example 5 ([Npg<sup>7</sup>]triptorelin) and example 6 ([D-Ala<sup>6</sup>, desGly<sup>10</sup>-ProNet<sup>9</sup>)LH-RH, Bachem), in combination with  $\alpha$ -CD (Sigma or Wacker Chemie). As a comparison, the same agonists have been orally administered by gavage in a standard aqueous vehicle not comprising  $\alpha$ -CD (comparative examples). For screening purposes, blood samples were drawn 2 hours after oral administration of a common dose of 2 nmoles/rat of LH-RH peptide agonists in an aqueous solution containing 10 or 100

mM of  $\alpha$ -CD (Tables 1 and 2; Figures 1, 4 and 6). For kinetic purposes, the effects of 2 nmoles/rat of example 3 with or without 100 mM of  $\alpha$ -CD were tested between 0.5 and 8 hours on plasma LH and testosterone levels (Figures 2 and 5). The influence of increasing concentrations of  $\alpha$ -CD (5%, 10% or 14%) was tested with example 2 at the dose of 5  $\mu$ g/kg 2 hours after oral administration (Figure 3). Total plasma testosterone (Diagnostic System Laboratories) and LH (Amersham Pharmacia Biotech) determinations were performed by radioimmunoassay. In screening 2-hour experiments, each group comprised between 6 and 8 rats; each time point of the kinetic study was studied on four animals.

10

Table 1 : stimulation of testosterone secretion

Compound without $\alpha$ -CD	total plasma testosterone (nmol/l) (m $\pm$ sem)	Compound with $\alpha$ -CD	total plasma testosterone (nmol/l) (m $\pm$ sem)
Control	8.6 $\pm$ 3.32	Control ( $\alpha$ -CD)	3.8 $\pm$ 0.66
Triptorelin	25.8 $\pm$ 3.14	Triptorelin ( $\alpha$ -CD)	61.9 $\pm$ 6.01
Leuprorelin	26.3 $\pm$ 5.77	Leuprorelin ( $\alpha$ -CD)	70.7 $\pm$ 4.06
Goserelin	21.4 $\pm$ 5.99	Goserelin ( $\alpha$ -CD)	66.5 $\pm$ 6.19
Deslorelin	9.7 $\pm$ 2.41	Deslorelin ( $\alpha$ -CD)	48.2 $\pm$ 7.29
C. ex 1	40.1 $\pm$ 6.78	Example 1	58.0 $\pm$ 8.75
C. ex 2	23.0 $\pm$ 8.54	Example 2	72.8 $\pm$ 4.64
C. ex 3	39.7 $\pm$ 8.11	Example 3	69.7 $\pm$ 3.6
C. ex 4	30.1 $\pm$ 5.86	Example 4	67.9 $\pm$ 9.11
C. ex 5	15.8 $\pm$ 4.24	Example 5	52.1 $\pm$ 6.99
C. ex 6	28.3 $\pm$ 4.56	Example 6	61.8 $\pm$ 5.10

As can be seen from the above results as well as from Figures 1-4, oral formulations with  $\alpha$ -CD significantly enhance the stimulation of testosterone secretion induced by LH-RH analogues. Especially, deslorelin alone was inactive at this threshold dose of 2 nmoles/rat, but showed a marked potency when formulated with  $\alpha$ -CD. The crucial role played by  $\alpha$ -CD is demonstrated by the concentration-dependance of its effect : combined with 10 mM  $\alpha$ -CD (0.972%), the oral activity of example 3 was not significantly improved (Figure 1) ; at 5% (51.4 mM),  $\alpha$ -CD did

15

enhance the stimulation of testosterone secretion induced by example 2 by oral administration, although not to the maximal level achieved with 10% (103 mM) as well as 14% (144 mM) (Figure 3).

It is also worth noting (see figures 1 and 3) that  $\beta$ -CD, hydroxypropyl- $\beta$ -CD (HP- $\beta$ -CD) and  $\gamma$ -CD have no potentiating effect on the LH-RH analogue-induced stimulation of testosterone secretion.

Table 2 : stimulation of LH secretion

Compound without $\alpha$ -CD	total plasma LH (ng/ml) (m $\pm$ sem)	Compound with $\alpha$ -CD	total plasma LH (ng/ml) (m $\pm$ sem)
Control	1.2 $\pm$ 0.11	Control ( $\alpha$ -CD)	1.1 $\pm$ 0.10
Triptorelin	1.4 $\pm$ 0.10	Triptorelin ( $\alpha$ -CD)	10.1 $\pm$ 2.54
Leuprorelin	1.2 $\pm$ 0.14	Leuprorelin ( $\alpha$ -CD)	12.3 $\pm$ 2.03
C. ex 1	1.5 $\pm$ 0.19	Example 1	7.1 $\pm$ 1.68
C. ex 2	1.6 $\pm$ 0.14	Example 2	19.7 $\pm$ 3.70
C. ex 3	2.2 $\pm$ 0.58	Example 3	10.9 $\pm$ 1.66
C. ex 4	1.4 $\pm$ 0.17	Example 4	16.1 $\pm$ 5.22
C. ex 5	1.4 $\pm$ 0.10	Example 5	3.2 $\pm$ 0.56

As can be seen from the above results as well as from Figures 5-6, the potentiating effect of  $\alpha$ -CD in oral formulations containing LH-RH analogues on LH release is even more pronounced than on testosterone secretion : all tested LH-RH analogues were inactive when administered alone at the same dose of 2 nmoles/rat, whereas, depending on the analogue, they induced a 3- to 16-fold increase above control levels when administered in combination with  $\alpha$ -CD.

Similar or even better results were obtained with  $\alpha$ -CD derivatives such as methylated  $\alpha$ -CD, hexakis(2,3,6-tri-O-methyl)- $\alpha$ -CD, carboxymethylated  $\alpha$ -CD or phosphated  $\alpha$ -CD. All  $\alpha$ -CD derivatives were purchased from Cyclolab (Budapest, Hungary). The influence of  $\alpha$ -CD derivatives were compared with that of  $\alpha$ -CD itself on the potentiation of LH-RH agonist activity of example 3 when administered by gavage to rats at the low dose of 5  $\mu$ g/kg p.o. : total testosterone plasma levels were measured 2 hours after administration (Table 3).



Table 3 : Stimulation of testosterone secretion by example 3

dose ( $\mu\text{g/kg p.o.}$ )	cyclodextrin (CD) type (concentration)	testosterone levels (ng/ml) ; ( $m \pm \text{sem}$ )	n rats
0 (control)	none	$1.0 \pm 0.17$	24
5	none	$3.4 \pm 0.79$	16
5	carboxymethylated $\alpha$ -CD (50%)	$6.9 \pm 1.62$	10
5	methylated $\alpha$ -CD (30%)	$7.1 \pm 1.59$	10
5	phosphated $\alpha$ -CD (30%)	$7.4 \pm 2.00$	10
5	$\alpha$ -CD (10%)	$10.0 \pm 1.22$	24
5	permethylated $\alpha$ -CD (15%)	$12.9 \pm 1.10$	10

The  $\alpha$ -CD derivatives tested above at least doubled the effect of example 3 by oral administration. Native  $\alpha$ -CD and its permethylated derivative appeared to be especially favorable with respectively a 2.9- and 3.8- fold enhancement of agonist activity of example 3 on testosterone at this dose level of  $5 \mu\text{g/kg p.o.}$

In a further experiment, example 3 was tested with  $\alpha$ -CD or permethylated  $\alpha$ -CD at an equal concentration of 10%. Two hours after administration, plasma LH levels were measured on eight rats per dose group (Figure 7). The  $5 \mu\text{g/kg p.o.}$  dose of example 3 alone was inactive on LH levels at this time point, and 10 and  $20 \mu\text{g/kg p.o.}$  were clearly threshold doses in these experimental conditions.

Combination of example 3 with 10%  $\alpha$ -CD resulted in slight but significant stimulations at 5 and  $10 \mu\text{g/kg p.o.}$ , and in a much greater effect at  $20 \mu\text{g/kg p.o.}$  when compared with example 3 alone (over 5-fold enhancement of LH-releasing activity). Moreover, combination of example 3 with 10% permethylated  $\alpha$ -CD resulted in an even higher potentiation : the doses of 2.5, 5 and  $10 \mu\text{g/kg p.o.}$ , which remained inactive when example 3 was given alone, yielded a sharp dose-related stimulatory response (Figure 7).

The peptides according to the general formula (I') exert an antagonist activity upon the LH-RH receptors *in vivo*, resulting in particular in the inhibition of ovulation.

The influence of  $\alpha$ -,  $\beta$ - and  $\gamma$ -CD was tested on the activity of antide (an example of LH-RH peptide antagonist) when orally administered by gavage to normally cycling adult female Wistar rats between 1:30 and 3:00 p.m. on the day of proestrus, after at least two full regular estrous cycles as monitored by daily vaginal

smears. The antioviulatory efficacy was checked the next morning, on the day of expected estrus, by looking for ova in the oviduct of treated females. The presence of at least one ovum attested that some degree of spontaneous ovulation did occur, and only the total absence of ovum was considered as effective LH-RH antagonist-induced inhibition of ovulation. Antide was solubilized in a vehicle consisting of 20% (vol/vol) propylene glycol in water already containing 1% bovine albumin, to which 10% (wt/vol) of either  $\alpha$ -,  $\beta$ - or  $\gamma$ -CD was then added. The results of the experiments are summarized in the following Table 4.

Table 4 : Inhibition of ovulation by oral administration of antide

oral formulation	antide ( $\mu\text{g}/\text{rat p.o.}$ )	n ovulations/ N treated rats	percentage of inhibition
vehicle only	0	24/24	0%
vehicle + $\alpha$ -CD (10%)	0	8/8	0%
vehicle only	200	8/8	0%
	400	19/22	14%
	600	5/8	44%
vehicle + $\alpha$ -CD (10%)	200	6/8	25%
	400	6/22	73%
	600	2/8	75%
vehicle + $\beta$ -CD (10%)	400	7/7	0%
vehicle + $\gamma$ -CD (10%)	400	7/7	0%

The vehicle with or without  $\alpha$ -CD had no effect by itself. The threshold effective dose of antide by oral administration was 400  $\mu\text{g}/\text{kg p.o.}$  with only 3 animals out of 22 showing inhibition of ovulation. Beta- and  $\gamma$ -CD had no influence on the minimal activity of antide at this dose level.

However,  $\alpha$ -CD significantly enhanced antide potency from 14% to 73% of inhibition at 400  $\mu\text{g}/\text{kg}$ . The twice lower dose of 200  $\mu\text{g}/\text{kg}$  was even slightly effective (25% of inhibition) in combination with  $\alpha$ -CD 10%. Therefore,  $\alpha$ -CD was able to potentiate the activity of a LH-RH peptide antagonist by oral administration, but not  $\beta$ - or  $\gamma$ -CD.

The oral formulations of the invention can be prepared by methods well known to those skilled in the art, generally as follows : a known amount of a drug is added to an aqueous cyclodextrin solution in sufficient concentration; the drug-cyclodextrin interaction can take place in solution or suspension within minutes or after stirring for up to 1 week at the desired temperature with or without sonication, depending on the nature of the drug and of the cyclodextrin, and on their respective concentrations. Then, the resulting drug-cyclodextrin combination or complex can be further obtained in a dry form by filtration, centrifugation, evaporation or sublimation.

By way of illustration, the example combinations of LH-RH analogues with  $\alpha$ -CD hereafter describe one basic method for the preparation of the formulations according to the invention in solution, notwithstanding their further processing to any appropriate dry form that will take advantage of the same potentiating properties.

Such formulations may further comprise, one or several other pharmaceutically appropriate excipients for oral administration such as lactose, fructose, glucose, sucrose, compressible sugar, saccharin, povidone, crospovidone, magnesium stearate, kaolin, bentonite, colloidal silica, mannitol, sorbitol, starch and its derivatives, microcrystalline or powdered cellulose, methylcellulose, carboxymethylcellulose, ethylcellulose or other chemically modified celluloses, other cyclodextrins, maltodextrin, - dextrates, dextrin, dextrose, alginates, pectins, pectates, sorbitan esters, polysorbate 80, chitosan, guar or xanthan gums, mono-, di- or tri-ethanolamine, oleic acid or ethyl oleate, stearic acid, water, liquid glucose, propylene glycol, lactic acid, malic acid, ethanol, isopropyl myristate or palmitate, glycerin, glyceryl monooleate, glyceryl monostearate, glyceryl palmitostearate, lecithin, medium or short chain triglycerides, various oils from corn, cottonseed, olive, peanut, sesame or soybean, and the like. These formulations are administered by mouth (or naso-gastric tubing) in various aqueous or non-aqueous solutions or suspensions such as true solutions, syrups, elixirs, mucilages, jellies, gels, milks, magmas, macro-, micro- or nano-emulsions, or in various solid forms such as compressed, coated, buccal, sublingual, effervescent or molded tablets, hard or soft capsules, pills, troches or cachets. Enteric coatings of usual solid oral dosage forms or of soft capsules containing liquid formulations, and sustained, delayed or programmed gastric, enteric or colonic release forms or devices are preferred means to deliver the active principle.

The main target of LH-RH peptide agonists according to formula (I) is the pituitary gland, but direct actions have been reported on the gonads themselves (testis and ovary), on the thymus and some lymphoid cell lines, and on breast, prostate, pancreatic or nervous system tumors. They exert on any LH-RH sensitive target, either a stimulatory activity by short-term acute or pulsatile administration, or an inhibitory effect by repeated or continuous administrations that induce the desensitization and the down-regulation of LH-RH receptors. In the case of the hypothalamo-pituitary-gonadal axis, prolonged administration results in a so-called "chemical" castration.

The main target of LH-RH peptide antagonists according to formula (I') is also the pituitary gland, where they bind to the LH-RH receptors and prevent the activity of endogenous LH-RH. By this mechanism, the pituitary-gonadal axis can be inhibited. The programmed use of LH-RH antagonists can also be taken advantage of to obtain a spontaneous stimulation of the pituitary-gonadal axis at any given time by stopping their administration at an appropriate earlier time point.

Therefore, LH-RH agonists or antagonists according to formula (A) are useful in all situations where the actions of LH-RH must be either inhibited, prevented or stimulated. Especially, the peptide analogues of the invention are useful in the treatment of LH-RH-sensitive diseases, namely the diseases where a LH-RH agonist or antagonist action is required.

Accordingly, the oral pharmaceutical compositions of the invention can find an appropriate therapeutic use in humans as well as in animals, depending on doses and treatment regimens, in reproductive endocrinology and in the treatment or prevention of sex hormone-dependent benign or malignant tumors ; said treatment or prevention may involve parallel and/or sequential supplementary curative or preventive regimens based on other hormonal or antitumoral agents. LH-RH sensitive sex hormone-independent benign or malignant tumors can also regress upon treatment with the oral pharmaceutical compositions according to the invention, alone or associated with other parallel and/or sequential antitumoral treatments. Immune mechanisms can also be modified by the oral pharmaceutical compositions according to the invention, alone or associated with other parallel and/or sequential treatments based on immuno-modulating or -suppressive agents such as glucocorticoids, cyclosporin, rapamycin, tacrolimus, their derivatives, and the like. The oral pharmaceutical compositions according to the invention are

therefore very valuable in the treatment and prevention of autoimmune diseases, graft rejection or atopic diseases, and in the treatment of benign or malignant lymphoproliferative disorders.

The oral pharmaceutical compositions according to the invention are especially useful in the inhibition, planning and triggering of ovulation in *in vitro* fertilization programs, and in the treatment of male and female infertility or hypogonadic states. Conversely, they can also be used in male or female contraception or treatment of hypergonadic states. In both cases, said treatments may involve other parallel and/or sequential treatments with sex steroids and/or gonadotrophins. This applies to men and women, but also to wild or domestic animals in uses such as improvement or control of reproductive performance, or as a tool to optimize breeding strategies.

The oral pharmaceutical compositions according to the invention are also especially useful in men to treat advanced prostate cancer, but can also be used as a first line therapy in this indication and in benign prostatic hypertrophy ; in both cases, said treatments may also involve additional parallel and/or sequential treatments based on inhibitors of androgen action, i.e. antiandrogens such as cyproterone acetate, osaterone acetate, chlormadinone acetate, flutamide, nilutamide or bicalutamide and the like, and/or on 5 $\alpha$ -reductase inhibitors such as finasteride, epristeride or turosteride and the like, and/or on C<sub>17-20</sub> lyase inhibitors such as abiraterone and the like.

The oral pharmaceutical compositions according to the invention are also especially useful in the treatment or prevention of breast cancer in women and in men, especially estrogen receptor positive tumors ; said treatment or prevention may involve parallel or sequential supplementary curative or preventive regimens based on antiestrogens such as tamoxifen, raloxifen or droloxifen and the like, and/or on aromatase inhibitors such as atamestane, formestane, letrozole, anastrozole and the like, and/or on C<sub>17-20</sub> lyase inhibitors such as abiraterone and the like. The oral pharmaceutical compositions according to the invention are also very useful in the treatment or prevention of certain estrogen receptor negative tumors that respond to the direct effects of LH-RH analogues or indirectly to their gonadal suppressive activity.


Other gynecological conditions, such as endometrial hyperplasia, leiomyoma, adenomyoma, endometriosis, polycystic ovary syndrome, hirsutism and benign

breast disease (pain, cysts or fibrosis), can also be prevented by or benefit from treatment with the oral pharmaceutical compositions according to the invention; said treatment or prevention may involve additional parallel and/or sequential curative or preventive treatments based on antiestrogens (cited above), progestins such as cyproterone acetate, osaterone acetate, chlormadinone acetate, norgestrol acetate, promegestone, demegestone, trimegestone and the like, and/or their contraceptive or post-menopausal replacement combination formulations with estrogens such as estradiol or ethynylestradiol. The oral compositions of the invention can also interfere with gestation by inducing abortion or by triggering labour; in this case they may also be used in parallel or in sequence with treatments based on estrogens (cited above), antiprogestins such as mifepristone and/or prostaglandin analogues such as sulprostone.

Similar indications can be encountered in veterinary medicine for male or female domestic or wild animals that may require the use of pharmaceutical compositions according to the invention.

A further aspect of the invention relates to a method of treating and/or preventing the above diseases which comprises orally administering to patients or animals in need thereof a pharmaceutical composition according to the invention, said composition comprising an effective amount of a LH-RH peptide analogue as previously defined in combination with  $\alpha$ -cyclodextrin or a derivative thereof. Said method may comprise the further administration of at least one of the active principles mentioned above such as for example a hormonal agent, an antitumoral agent, an immuno-modulating or suppressive agent, a sex steroid, a gonadotrophin, an inhibitor of androgen action, a  $5\alpha$ -reductase inhibitor, a  $C_{17-20}$  lyase inhibitor, an antiestrogen, an aromatase inhibitor, a progestin, an estrogen, an antiprogestin or a prostaglandin analogue, said further administration being parallel, sequential or over a period of time.

In a further aspect, the invention relates to a method of treating a disease wherein a LH-RH agonist or antagonist action is required which comprises orally administering to a patient in need thereof a therapeutically effective amount of a LH-RH peptide analogue in combination with  $\alpha$ -cyclodextrin or a derivative thereof, wherein said peptide analogue has the formula (SEQ ID No: 1):



in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond; His; or an aromatic D-amino acid;




[R:\LIBH\03736.doc:aak

- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; (S)-spirolactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn;
- 5 D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala;
- 10 D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;
- 15 - Z is GlyNH<sub>2</sub>; D-AlaNH<sub>2</sub>; azaGlyNH<sub>2</sub> or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

In another aspect, the invention relates to a method of enhancing the biological activity of a LH-RH peptide analogue which comprises orally administering to a patient in need thereof a therapeutically effective amount of said analogue in combination with  $\alpha$ -cyclodextrin or a derivative thereof, or a pharmaceutical composition according to the invention.

According to another aspect the invention relates to  $\alpha$ -cyclodextrin or a derivative thereof when used in combination with a therapeutically effective amount of an LH-RH peptide analogue for enhancing the biological activity of said LH-RH peptide analogue.

In a further aspect, the invention relates to  $\alpha$ -cyclodextrin or a derivative thereof when used in combination with a therapeutically effective amount of an LH-RH peptide analogue for treating a disease where a LH-RH agonist or antagonist action is required, wherein said peptide analogue has the formula (SEQ ID No: 1):



in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond; His; or an aromatic D-amino acid;



[R:\LIBH\03736.doc:aak

- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; (S)-spiro lactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn;
- 5 D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>t</sup>); D-Asp(OBu<sup>t</sup>); D-Glu(OBu<sup>t</sup>);  
D-Thr(OBu<sup>t</sup>); D-Cys(OBu<sup>t</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid;  
D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or  
a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side  
chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala;
- 10 D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon  
atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted  
by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;
- 15 - Z is GlyNH<sub>2</sub>; D-AlaNH<sub>2</sub>; azaGlyNH<sub>2</sub> or a group -NHR<sub>2</sub> where R<sub>2</sub> is a  
(C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a  
(C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and  
piperidyl.

The unit dose of oral administration of LH-RH peptide analogues according to  
20 formula (A) may range from 0.1 to 100 mg per human patient, from one to 16 times per  
day (in the case of pulsatile administration), in combination with at least an equimolar  
amount of  $\alpha$ -CD or its derivatives and up to the total remaining part of the oral  
formulation.



[R:\LIBH\03736.doc:aak]



All the above-mentioned oral pharmaceutical compositions may additionally contain one or several proteases inhibitors, and/or one or several other absorption enhancers.

Examples of preparations of leuporelin, triptorelin, goserelin, deslorelin and examples 1 to 6 in combination with 100 mM  $\alpha$ -CD in solution

On each experimental day, solutions of  $\alpha$ -CD were freshly prepared by dissolving 9.72 g in 100 ml of pure water, or 4.86 g in 50 ml, for 1 hour at room temperature with gentle magnetic stirring; meanwhile, an appropriate volume of each LH-RH analogue was taken from thawed individual stock vials containing 50  $\mu$ g of net peptide in 50  $\mu$ l of phosphate-buffered saline containing 0.1% bovine serum albumin, to make 20 nmoles (24.2  $\mu$ l for LEU, 26.2  $\mu$ l for TRI, 23.4  $\mu$ l for GOS, 25.4  $\mu$ l for DES, 24.7  $\mu$ l for example 1, 23.6  $\mu$ l for example 2, 24.5  $\mu$ l for example 3, 25.1  $\mu$ l for example 4, 26.5  $\mu$ l for example 5 and 23.4  $\mu$ l for example 6) and put in a 10 ml gauged flask. Then, the  $\alpha$ -CD solution was added to fill the flask up to 10 ml to make a 2 nmol/ml solution of which 1ml was administered by oral gavage to each rat.

Examples of preparation of formulations of example 2 in combination with 5%, 10% or 14%  $\alpha$ -CD solutions

On each experimental day, 45  $\mu$ l of one thawed individual vial containing 50  $\mu$ g of net example 2 in 50  $\mu$ l of phosphate-buffered saline containing 0.1 % bovine serum albumin, were diluted in 36 ml of distilled water to give a 1.25  $\mu$ g/ml solution from which three fractions of 3.8 ml were taken; then, 190, 380 or 532 mg of  $\alpha$ -CD were added to each fraction to give a concentration of 5%, 10% or 14%, respectively. After overnight magnetic stirring at room temperature, each solution was given to rats by oral gavage in a 4 ml/kg volume to administer the same dose of 5  $\mu$ g/kg of example 2 without or with increasing concentrations of  $\alpha$ -CD.

Examples of preparation of formulations for oral administration of example 3 in combination with  $\alpha$ -CD derivatives

On each experimental day, frozen vials containing 50  $\mu$ g of net example 3 in 50  $\mu$ l of phosphate-buffered saline containing 0.1% bovine serum albumin were thawed and diluted by half with an equal volume of the same fresh bovine serum albumin solution. Then, 12.5  $\mu$ l of this 0.5  $\mu$ g/ $\mu$ l solution were added to 5 ml of aqueous vehicle for oral administration (with or without  $\alpha$ -CD derivative) to give a

final formulation containing 1.25 µg/ml of example 3 to be administered by gavage under a volume of 4 ml/kg, after gentle magnetic stirring overnight.

The solutions of α-CD derivatives were prepared by weighing the appropriate amount to put in 10 ml gauged flasks to fill up with water : 5 g of carboxymethylated  
5 α-CD (50%), 3 g of methylated α-CD (30%), 3 g of phosphated α-CD (30%) or 1 or 1.5 g of permethylated α-CD (10 or 15%).

Appropriate volumes of convenient dilutions of the 50 µg/50 µl stock vials of net example 3 were added to 10% solutions of α-CD or permethylated α-CD in water to obtain the dose range described in figure 7.

10 Examples of preparations of formulations for oral administration of antide in combination with α-, β- or γ-CD

Each 5 mg powder vial of antide (from Bachem, Bubendorf, Switzerland) containing 4.2434 mg net peptide was dissolved in a mixture of 2.122 ml propylene glycol with 8.487 ml water containing 0.1% bovine albumin. To each 10.609 ml  
15 solution of antide (400 µg/ml), 1.061 g of either α-, β- or γ-CD was directly added to obtain a 10% concentration. Each female rat received the same volume of 1 ml of test formulation by gavage.


The same volume of administration was used for the doses of 200 and 600 µg/rat. The appropriate concentrations of antide to be administered (200 and 600  
20 µg/ml) were respectively obtained by diluting by half the 400 µg/ml solution with the same [20% vol. propylene glycol/80% vol. albuminated water/10% wt α-CD] mixture, and by dissolving other 5 mg powder vials in 1.415 ml propylene glycol with 5.658 ml water already containing 0.1% bovine albumin to which 0.707 mg of α-CD was finally added.

25

## CLAIMS

1. Use of  $\alpha$ -cyclodextrin or a derivative thereof for the preparation of a pharmaceutical composition for the oral administration of a LH-RH peptide analogue or one of its pharmaceutically acceptable salt.


2. Use according to claim 1 wherein said peptide analogue has the formula (SEQ ID N° : 1) :



in which :

- 10 - A1 is pGlu ; D-pGlu ; Sar ; AcSar ; Pro or a derivative thereof ; Ser ; D-Ser ; Ac-D-Ser ; Thr ; D-Thr ; Ac-D-Thr ; or an optionally substituted and/or acylated aromatic D-amino acid;
- A2 is a direct bond ; His ; or an optionally substituted aromatic D-amino acid;
- A3 is an optionally substituted aromatic L- or D-amino acid;
- 15 - A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;
- A5 is an optionally substituted aromatic L-amino acid or an optionally substituted basic L- or D-amino acid ;
- A6 is Gly ; (S)-spiro lactam-Pro ; D-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Asn ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>t</sup>) ; D-Asp(OBu<sup>t</sup>) ; D-Glu(OBu<sup>t</sup>) ; D-Thr(OBu<sup>t</sup>) ; D-Cys(OBu<sup>t</sup>) ; D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ;
- 20 an aza-amino acid ; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain ; an optionally substituted aromatic D-amino acid ; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydrodiphenyl-
- 25 Ala ; or an optionally substituted basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms ;
- A8 is an optionally substituted basic L- or D-amino acid;
- 30 - Z is GlyNH<sub>2</sub> ; D-AlaNH<sub>2</sub> ; azaGlyNH<sub>2</sub> ; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms ; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.


3. Use according to claim 2 wherein said peptide analogue has the formula (SEQ ID N° : 2) :



in which:


- 5 - A1 is pGlu, Sar or AcSar;
- A3 is an optionally substituted aromatic L-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>h</sup>), Ser(OBzl) or Thr;
- A5 is an optionally substituted aromatic L-amino acid ;
- A6 is Gly ; D-Pro ; (S)-spirolactam-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>h</sup>) ; D-Asp(OBu<sup>h</sup>) ; D-Glu(OBu<sup>h</sup>) ; D-Thr(OBu<sup>h</sup>) ; D-Cys(OBu<sup>h</sup>) ; D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aza-amino acid ; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain ; an optionally substituted aromatic D-amino acid ; D-cyclohexadienyl-Gly ;
- 10 D-perhydronaphtyl-Ala ; D-perhydrodiphenyl-Ala ; or an optionally substituted basic D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- 20 - A8 is an optionally substituted basic L-amino acid;
- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> ; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms ; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

25 4. Use according to claim 3 wherein said peptide analogue has the formula (SEQ ID N° : 3) :



in which A7 is Leu, Tle, Nle, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several  
30 fluorine atoms.

5. Use according to claim 3 wherein said peptide analogue has the formula (SEQ ID N° : 4) :



in which:


- A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Bal, Pal, 4Pal, or pClPhe ;

- A6 is (S)-spiro lactam-Pro ; Gly, D-Pro ; D-Ser(OBu<sup>t</sup>) ; D-Asp(OBu<sup>t</sup>) ; D-Glu(OBu<sup>t</sup>) ; D-Thr(OBu<sup>t</sup>) ; D-Cys(OBu<sup>t</sup>) ; D-His or D-His(Bzl) ; D-Ala, D-Leu, D-Tle, D-Nle, D-Hol, D-Npg or D-Cha ; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe ; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydrodiphenyl-Ala ; or D-APhe optionally substituted by an aminotriazolyl group ;

- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;

- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

6. Use according to claim 3 wherein said peptide analogue has the formula (SEQ ID N° : 5) :



in which:


- A6 is (S)-spiro lactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>t</sup>) or D-Trp;

- A7 is Leu, MeLeu, Npg or MeNpg;

- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

7. Use according to one of claims 3 to 6 wherein the peptide analogue is selected from the group consisting of leuporelin, [Npg<sup>7</sup>]-leuporelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, buserelin and [Npg<sup>7</sup>]-buserelin.

8. Use according to claim 2 wherein said peptide analogue has the formula (SEQ ID N° : 6) :



in which:

- A1 is pGlu ; D-pGlu ; Sar ; AcSar ; Pro or a derivative thereof ; Ser ; D-Ser ; Ac-D-Ser ; Thr ; D-Thr ; Ac-D-Thr ; or an optionally substituted and/or acylated aromatic D-amino acid;

- A2 is a direct bond or an optionally substituted aromatic D-amino acid;

- A3 is an optionally substituted aromatic L- or D-amino acid;

- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;

- A5 is an optionally substituted aromatic L-amino acid or an optionally substituted basic L- or D-amino acid;

- A6 is Gly ; (S)-spirolactam-Pro ; D-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Asn ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>1</sup>) ; D-Asp(OBu<sup>1</sup>) ; D-Glu(O-Bu<sup>1</sup>) ; D-Thr(O-Bu<sup>1</sup>) ; D-Cys(O-Bu<sup>1</sup>) ; D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain ; an optionally substituted aromatic D-amino acid ; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydrodiphenyl-Ala ; or an optionally substituted basic L- or D-amino acid;

- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is an optionally substituted basic L- or D-amino acid;

- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>.

9. Use according to claim 8 wherein the peptide analogue has the formula (SEQ ID N° : 7):

15 Ac-D-Nal-D-pCIPhe-D-Pal-Ser-A5-A6-A7-A8-Pro-D-AlaNH<sub>2</sub> (II')

in which :

- A5 is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys;

- A6 is (S)-spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HCit or D-Asn;

20 - A7 is Leu, MeLeu, Npg or MeNpg;

- A8 is Arg, NicLys or IprLys.

10. Use according to claim 8 or 9 wherein the peptide analogue is selected from the group consisting of antide, [Npg<sup>7</sup>]-antide, cetorelix, [Npg<sup>7</sup>]-cetorelix, abarelix and [Npg<sup>7</sup>]-abarelix.

25 11. Use according to one of claims 1 to 10 wherein the  $\alpha$ -cyclodextrin derivative is selected from the group consisting of methylated  $\alpha$ -cyclodextrin, hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin, carboxymethylated  $\alpha$ -cyclodextrin and phosphated  $\alpha$ -cyclodextrin.

12. Use according to claim 11, wherein the  $\alpha$ -cyclodextrin derivative is 30 hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin.


13. Use according to one of claims 1 to 12 wherein the pharmaceutical composition is intended to be delivered to the gastrointestinal tract.



14. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is intended for the treatment of infertility, hypogonadic or hypergonadic states.
15. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is a contraceptive agent.
16. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is intended for the treatment or prevention of prostate cancer or benign prostatic hypertrophy.
17. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is intended for the treatment or prevention of breast cancer.
18. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is intended for the treatment or prevention of sex hormone-related benign or malignant tumors.
19. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is intended for the treatment or prevention of sex hormone-independent but LH-RH sensitive benign or malignant tumors.
20. Use according to one of claims 1 to 13 wherein the pharmaceutical composition is intended for the treatment or prevention of benign or malignant lymphoproliferative disorders.
21. A pharmaceutical composition for the gastrointestinal delivery by oral administration of a LH-RH peptide analogue which comprises a therapeutically effective amount of said peptide analogue in combination with  $\alpha$ -cyclodextrin or a derivative thereof.
22. The pharmaceutical composition according to claim 21 which further comprises excipients suitable for the gastrointestinal delivery of the peptide analogue.
23. The pharmaceutical composition according to claim 21 or 22 wherein said peptide analogue has the formula (SEQ ID N° : 1) :
- A1-A2-A3-A4-A5-A6-A7-A8-Pro-Z (A)
- in which :
- A1 is pGlu ; D-pGlu ; Sar ; AcSar ; Pro or a derivative thereof ; Ser ; D-Ser ; Ac-D-Ser ; Thr ; D-Thr ; Ac-D-Thr ; or an optionally substituted and/or acylated aromatic D-amino acid;
  - A2 is a direct bond ; His ; or an optionally substituted aromatic D-amino acid;

- A3 is an optionally substituted aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>h</sup>), Ser(OBzl) or Thr;
- A5 is an optionally substituted aromatic L-amino acid or an optionally substituted basic L- or D-amino acid;
- 5     - A6 is Gly ; (S)-spirolactam-Pro ; D-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Asn ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>h</sup>) ; D-Asp(OBu<sup>h</sup>) ; D-Glu(OBu<sup>h</sup>) ; D-Thr(OBu<sup>h</sup>) ; D-Cys(OBu<sup>h</sup>) ; D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aza-amino acid ; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain ; an optionally substituted aromatic D-amino acid ; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydropiphenyl-Ala ; or an optionally substituted basic L- or D-amino acid;
- 10     - A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- 15     - A8 is an optionally substituted basic L- or D-amino acid;
- Z is GlyNH<sub>2</sub> ; D-AlaNH<sub>2</sub> ; azaGlyNH<sub>2</sub> ; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms ; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.
- 20

24. The pharmaceutical composition according to claim 23 wherein said peptide analogue has the formula (SEQ ID N° : 2) :



in which:

- 25     - A1 is pGlu, Sar or AcSar;
- A3 is an optionally substituted aromatic L-amino acid ;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>h</sup>), Ser(OBzl) or Thr;
- A5 is an optionally substituted aromatic L-amino acid ;
- A6 is Gly ; D-Pro ; (S)-spirolactam-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>h</sup>) ; D-Asp(OBu<sup>h</sup>) ; D-Glu(OBu<sup>h</sup>) ; D-Thr(OBu<sup>h</sup>) ; D-Cys(OBu<sup>h</sup>) ; D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aza-amino acid ; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain ; an optionally substituted aromatic D-amino acid ; D-cyclohexadienyl-Gly ;
- 30




D-perhydronaphtyl-Ala ; D-perhydrodiphenyl-Ala ; or an optionally substituted basic D-amino acid ;

- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is an optionally substituted basic L-amino acid;


- Z is GlyNH<sub>2</sub> ; azaGlyNH<sub>2</sub> ; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms ; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl ; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

25. The pharmaceutical composition according to claim 24 wherein said peptide analogue has the formula (SEQ ID N° : 3) :



in which A7 is Leu, Tie, Nie, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several fluorine atoms.

26. The pharmaceutical composition according to claim 24 wherein said peptide analogue has the formula (SEQ ID N° : 4) :



in which:


- A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Bal, Pal, 4Pal, or pClPhe ;

- A6 is (S)-spirolactam-Pro ; Gly ; D-Pro ; D-Ser(OBu<sup>t</sup>) ; D-Asp(OBu<sup>t</sup>) ; D-Glu(OBu<sup>t</sup>) ; D-Thr(OBu<sup>t</sup>) ; D-Cys(OBu<sup>t</sup>) ; D-His or D-His(Bzl) ; D-Ala, D-Leu, D-Tie, D-Nie, D-Hol, D-Npg or D-Cha ; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe; D-cyclohexadienyl-Gly ; D-perhydronaphtyl-Ala ; D-perhydrodiphenyl-Ala or D-APhe optionally substituted by an aminotriazolyl group;

- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;

- Z is GlyNH<sub>2</sub>, azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

27. The pharmaceutical composition according to claim 24 wherein said peptide analogue has the formula (SEQ ID N° : 5) :



in which:


- A6 is (S)-spirolactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>1</sup>) or D-Trp;

- A7 is Leu, MeLeu, Npg or MeNpg;

- Z is GlyNH<sub>2</sub>, azaGlyNH<sub>2</sub> or -NC<sub>3</sub>H<sub>5</sub>.

5 28. The pharmaceutical composition according to one of claims 24 to 27 wherein the peptide analogue is selected from the group consisting of leuprorelin, [Npg<sup>7</sup>]-leuprorelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, busarelin and [Npg<sup>7</sup>]-busarelin.

10 29. The pharmaceutical composition according to claim 23 wherein said peptide analogue has the formula (SEQ ID N° : 6) :



in which:

- A1 is pGlu ; D-pGlu ; Sar ; AcSar ; Pro or a derivative thereof ; Ser ; D-Ser ; Ac-D-Ser ; Thr ; D-Thr ; Ac-D-Thr ; or an optionally substituted and/or acylated aromatic D-amino acid;

- A2 is a direct bond or an optionally substituted aromatic D-amino acid;

- A3 is an optionally substituted aromatic L- or D-amino acid;

- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;

20 - A5 is an optionally substituted aromatic L-amino acid or an optionally substituted basic L- or D-amino acid;

25 - A6 is Gly ; (S)-spirolactam-Pro ; D-Pro ; D-Ser ; D-Thr ; D-Cys ; D-Met ; D-Asn ; D-Pen ; D-(S-Me)Pen ; D-(S-Et)Pen ; D-Ser(OBu<sup>1</sup>) ; D-Asp(OBu<sup>1</sup>) ; D-Glu(OBu<sup>1</sup>) ; D-Thr(OBu<sup>1</sup>) ; D-Cys(OBu<sup>1</sup>) ; D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety ; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain ; an optionally substituted aromatic D-amino acid ; D-cyclohexadienyl-Gly ; D-perhydronaphthyl-Ala ; D-perhydrodiphenyl-Ala ; or an optionally substituted basic L- or D-amino acid;

30 - A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is an optionally substituted basic L- or D-amino acid;

- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>.

30. The pharmaceutical composition according to claim 29 wherein the peptide analogue has the formula (SEQ ID N° : 7):



Ac-D-Nal-D-pCIPhe-D-Pal-Ser-A5-A6-A7-A8-Pro-D-AlaNH<sub>2</sub> (II') in which:

- A5 is Tyr, Htyr, MeTyr, MeHTyr, NicLys or IprLys;
- A6 is (S)-spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HCit or D-Asn;
- 5     - A7 is Leu, MeLeu, Npg or MeNpg;
- A8 is Arg, NicLys or IprLys.

31. The pharmaceutical composition according to claim 29 or 30 wherein the peptide analogue is selected from the group consisting of antide, [Npg<sup>7</sup>]-antide, cetrorelix, [Npg<sup>7</sup>]-cetrorelix, abarelix and [Npg<sup>7</sup>]-abarelix.

10     32. The pharmaceutical composition according to any one of claims 21 to 31 wherein the  $\alpha$ -cyclodextrin derivative is selected from the group consisting of methylated  $\alpha$ -cyclodextrin, hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin, carboxymethylated  $\alpha$ -cyclodextrin and phosphated  $\alpha$ -cyclodextrin.


15     33. The pharmaceutical composition according to claim 32, wherein the  $\alpha$ -cyclodextrin derivative is hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin.

34. The pharmaceutical composition according to any one of claims 21 to 33 which further comprises a protease inhibitor and/or an absorption enhancer.

35. A pharmaceutical composition for the gastrointestinal delivery by oral administration of a LH-RH peptide analogue, substantially as hereinbefore described with reference to any one of the examples but excluding the comparative examples.

20     36. A method of enhancing the biological activity of a LH-RH peptide analogue which comprises orally administering to a patient in need thereof a therapeutically effective amount of said analogue in combination with  $\alpha$ -cyclodextrin or a derivative thereof, or a pharmaceutical composition according to any one of claims 21 to 34.

25     37. The method according to claim 36, wherein said peptide analogue has the formula (SEQ ID No: 1):



in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser;
- 30     Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond; His; or an aromatic D-amino acid;
- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;



[R:\LIBH\03736.doc:aak


- A6 is Gly; (S)-spirolactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;

- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is a basic L- or D-amino acid;

- Z is GlyNH<sub>2</sub>; D-AlaNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

38. The method according to claim 37 wherein said peptide analogue has the formula (SEQ ID No: 2):



in which:

- Al is pGlu, Sar or AcSar;

- A3 is an aromatic L-amino acid;

- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;

- A5 is an aromatic L-amino acid;

- A6 is Gly; D-Pro; (S)-spirolactam-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic D-amino acid;


- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is a basic L-amino acid;




- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

39. The method according to claim 38 wherein said peptide analogue has the  
5 formula (SEQ ID No: 3):



in which A7 is Leu, Tle, Nle, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several fluorine atoms.

40. The method according to claim 38 wherein said peptide analogue has the  
10 formula (SEQ ID No: 4):



in which:


- A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Bal, Pal, 4Pal, or pClPhe;

15 - A6 is (S)-spirolactam-Pro; Gly; D-Pro; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-His or D-His(Bzl); D-Ala, D-Leu, D-Tle, D-Nle, D-Hol, D-Npg or D-Cha; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or D-APhe optionally substituted by  
20 an aminotriazolyl group;

- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;

- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

41. The method according to claim 38 wherein said peptide analogue has the  
formula (SEQ ID No: 5):



in which:

- A6 is (S)-spirolactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>1</sup>) or D-Trp;

- A7 is Leu, MeLeu, Npg or MeNpg;

30 - Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

42. The method according to any one of claims 38 to 41 wherein the peptide analogue is selected from the group consisting of leuporelin, [Npg<sup>7</sup>]-leuporelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, buserelin and [Npg<sup>7</sup>]-buserelin.

43. The method according to claim 37 wherein said peptide analogue has the  
formula (SEQ ID No: 6):




[R:\LIBH\03736.doc:aak

## A1-A2-A3-A4-A5-A6-A7-A8-Pro-Z (I')

in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- 5 - A2 is a direct bond or an aromatic D-amino acid;
- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(O-Bu<sup>1</sup>); D-Thr(O-Bu<sup>1</sup>); D-Cys(O-Bu<sup>1</sup>); D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- 10 - A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;
- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>.

- 20 44. The method according to claim 43 wherein the peptide analogue has the formula (SEQ ID No: 7):



in which:

- A5 is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys;
- 25 - A6 is (S)spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HCit or D-Asn;
- A7 is Leu, MeLeu, Npg or MeNpg;
- A8 is Arg, NicLys or IprLys.

- 30 45. The method according to claim 43 or 44 wherein the peptide analogue is selected from the group consisting of antide, [Npg<sup>7</sup>]-antide, cetorelix, [Npg<sup>7</sup>]-cetorelix, abarelix and [Npg<sup>7</sup>]-abarelix.


46. The method according to any one of claims 36 to 45 wherein the  $\alpha$ -cyclodextrin derivative is selected from the group consisting of methylated  $\alpha$ -cyclodextrin hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin, carboxymethylated  $\alpha$ -cyclodextrin and phosphated  $\alpha$ -cyclodextrin.



[RALIBH]03736.doc:aak

47. The method according to claim 46 wherein the  $\alpha$ -cyclodextrin derivative is hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin.


48. A method of treating a disease wherein a LH-RH agonist or antagonist action is required which comprises orally administering to a patient in need thereof a therapeutically effective amount of a LH-RH peptide analogue in combination with  $\alpha$ -cyclodextrin or a derivative thereof, wherein said peptide analogue has the formula (SEQ ID No: 1);



in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond; His; or an aromatic D-amino acid;
- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; (S)-spirolactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>t</sup>); D-Asp(OBu<sup>t</sup>); D-Glu(OBu<sup>t</sup>); D-Thr(OBu<sup>t</sup>); D-Cys(OBu<sup>t</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;
- Z is GlyNH<sub>2</sub>; D-AlaNH<sub>2</sub>; azaGlyNH<sub>2</sub> or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

49. The method according to claim 48 wherein said peptide analogue has the formula (SEQ ID No: 2):



in which:


- A1 is pGlu, Sar or AcSar;



[R:\LIBH\03736.doc:aak


- A3 is an aromatic L-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid;
- A6 is Gly; D-Pro; (S)-spirolactam-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L-amino acid;
- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

50. The method according to claim 49 wherein said peptide analogue has the formula (SEQ ID No: 3):



in which A7 is Leu, Tle, Nle, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several fluorine atoms.

51. The method according to claim 49 wherein said peptide analogue has the formula (SEQ ID No: 4):



in which:

- A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Bal, Pal, 4Pal, or pClPhe;
- A6 is (S)-spirolactam-Pro; Gly; D-Pro; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-His or D-His(Bzl); D-Ala, D-Leu, D-Tle, D-Nle, D-Hol, D-Npg or D-Cha; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or D-APhe optionally substituted by an aminotriazolyl group;
- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;




[R:\LIBH\03736.doc:ask



- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or -NC<sub>2</sub>H<sub>5</sub>.

52. The method according to claim 49 wherein said peptide analogue has the formula (SEQ ID No: 5):




5 in which:

- A6 is (S)-spirolactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>1</sup>) or D-Trp;
- A7 is Leu, MeLeu, Npg or MeNpg;
- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or -NC<sub>2</sub>H<sub>5</sub>.

10 53. The method according to any one of claims 49 to 52 wherein the peptide analogue is selected from the group consisting of leuporelin, [Npg<sup>7</sup>]-leuporelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, buserelin and [Npg<sup>7</sup>]-buserelin.

54. The method according to claim 48 wherein said peptide analogue has the formula (SEQ ID No: 6):




15 in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond or an aromatic D-amino acid;
- 20 - A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(O-Bu<sup>1</sup>); D-Thr(O-Bu<sup>1</sup>); D-Cys(O-Bu<sup>1</sup>); D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- 30 - A8 is a basic L- or D-amino acid;
- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>.

55. The method according to claim 54 wherein the peptide analogue has the formula (SEQ ID No: 7):



[R:\LIBH\03736.doc:aak



in which:

- A5 is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys;
- A6 is (S)spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HCit or  
5 D-Asn;
- A7 is Leu, MeLeu, Npg or MeNpg;
- A8 is Arg, NicLys or IprLys.

56. The method according to claim 54 or 55 wherein the peptide analogue is selected from the group consisting of antide, [Npg<sup>7</sup>]-antide, cetorelix, [Npg<sup>7</sup>]-cetorelix,  
10 abarelix and [Npg<sup>7</sup>]-abarelix.

57. The method according to any one of claims 48 to 56 wherein the  $\alpha$ -cyclodextrin derivative is selected from the group consisting of methylated  $\alpha$ -cyclodextrin, hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin, carboxymethylated  $\alpha$ -cyclodextrin and phosphated  $\alpha$ -cyclodextrin.

15 58. The method according to claim 57 wherein the  $\alpha$ -cyclodextrin derivative is hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin.

59. The method according to any one of claims 48 to 58 for the treatment or prevention of breast cancer.

20 60. The method according to claim 59 which further comprises the sequential, parallel or over a period of time administration of at least one compound selected from the group consisting of an antiestrogen, an aromatase inhibitor and a C<sub>17</sub>-C<sub>20</sub> lyase inhibitor.

61. The method according to any one of claims 48 to 58 for the treatment or prevention of prostate cancer or benign prostatic hypertrophy.

25 62. The method according to claim 61 which further comprises the sequential, parallel or over a period of time administration of at least one compound selected from the group consisting of an antiandrogen, a 5 $\alpha$ -reductase inhibitor and a C<sub>17</sub>-C<sub>20</sub> lyase inhibitor.

30 63. The method according to any one of claims 48 to 62 wherein the peptide analogue is delivered to the gastrointestinal tract of the patient.

64.  $\alpha$ -Cyclodextrin or a derivative thereof when used in combination with a therapeutically effective amount of an LH-RH peptide analogue for enhancing the biological activity of said LH-RH peptide analogue.

65.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 64, wherein said peptide analogue has the formula (SEQ ID No: 1):



[RALIBH]03736.doc:aak

## Al-A2-A3-A4-A5-A6-A7-A8-Pro-Z (A)

in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- 5 - A2 is a direct bond; His; or an aromatic D-amino acid;
- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; (S)-spirolactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn;
- 10 D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala;
- 15 D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;
- 20 - Z is GlyNH<sub>2</sub>; D-AlaNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

66. α-Cyclodextrin when used in combination an LH-RH peptide analogue  
25 according to claim 65, wherein said peptide analogue has the formula (SEQ ID No: 2):

## Al-His-A3-A4-A5-A6-A7-A8-Pro-Z (I)

in which:

- A1 is pGlu, Sar or AcSar;
- A3 is an aromatic L-amino acid;
- 30 - A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid;
- A6 is Gly; D-Pro; (S)-spirolactam-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group; an aliphatic



[R:\LIBH\03736.doc:aak


D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic D-amino acid;

- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is a basic L-amino acid;


- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

67. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 66, wherein said peptide analogue has the formula (SEQ ID No: 3):



in which A7 is Leu, Tle, Nle, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several fluorine atoms.

68. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 66 wherein said peptide analogue has the formula (SEQ ID No: 4):



in which:


- A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Bal, Pal, 4Pal, or pClPhe;

- A6 is (S)-spirolactam-Pro; Gly; D-Pro; D-Ser(OBu<sup>t</sup>); D-Asp(OBu<sup>t</sup>); D-Glu(OBu<sup>t</sup>); D-Thr(OBu<sup>t</sup>); D-Cys(OBu<sup>t</sup>); D-His or D-His(Bzl); D-Ala, D-Leu, D-Tle, D-Nle, D-Hol, D-Npg or D-Cha; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or D-APhe optionally substituted by an aminotriazolyl group;

- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;

- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

69. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 66 wherein said peptide analogue has the formula (SEQ ID No: 5):



in which:

- A6 is (S)-spirolactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>t</sup>) or D-Trp;




[R:\LIBH\03736.doc:aak

- A7 is Leu, MeLeu, Npg or MeNpg;
- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub> or -NC<sub>2</sub>H<sub>5</sub>.

70. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 66 to 69 wherein the peptide analogue is selected from the group consisting of leuprorelin, [Npg<sup>7</sup>]-leuprorelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, buserelin and [Npg<sup>7</sup>]-buserelin.


71. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 65 wherein said peptide analogue has the formula (SEQ ID No: 6):



in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond or an aromatic D-amino acid;
- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(O-Bu<sup>1</sup>); D-Thr(O-Bu<sup>1</sup>); D-Cys(O-Bu<sup>1</sup>); D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;
- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>.

72. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 71 wherein the peptide analogue has the formula (SEQ ID No: 7):



in which:

- A5 is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys;
- A6 is (S)spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HCit or D-Asn;
- A7 is Leu, MeLeu, Npg or MeNpg;



[R:\L\IBH\03736.doc:aak


- A8 is Arg, NicLys or IprLys.

73.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 71 or 72 wherein the peptide analogue is selected from the group consisting of antide, [Npg<sup>7</sup>]-antide, cetorelix, [Npg<sup>7</sup>]-cetorelix, abarelix and [Npg<sup>7</sup>]-abarelix.

74.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 64 to 73 wherein the  $\alpha$ -cyclodextrin derivative is selected from the group consisting of methylated  $\alpha$ -cyclodextrin hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin, carboxymethylated  $\alpha$ -cyclodextrin and phosphated  $\alpha$ -cyclodextrin.

75.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 74 wherein the  $\alpha$ -cyclodextrin derivative is hexakis(2,3,6-tri-O-methyl)- $\alpha$ -cyclodextrin.

76.  $\alpha$ -Cyclodextrin or a derivative thereof when used in combination with a therapeutically effective amount of an LH-RH peptide analogue for treating a disease where a LH-RH agonist or antagonist action is required, wherein said peptide analogue has the formula (SEQ ID No: 1):



in which:


- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;
- A2 is a direct bond; His; or an aromatic D-amino acid;
- A3 is an aromatic L- or D-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;
- A6 is Gly; (S)-spirolactam-Pro; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl, a (C<sub>2</sub>-C<sub>7</sub>)acyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydropiphenyl-Ala; or a basic L- or D-amino acid;
- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L- or D-amino acid;



[RALIBH]03736.doc:aaak

- Z is GlyNH<sub>2</sub>; D-AlaNH<sub>2</sub>; azaGlyNH<sub>2</sub> or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.


5 77. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 76 wherein said peptide analogue has the formula (SEQ ID No: 2):



in which:


- Al is pGlu, Sar or AcSar;
- 10 - A3 is an aromatic L-amino acid;
- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>t</sup>), Ser(OBzl) or Thr;
- A5 is an aromatic L-amino acid;
- A6 is Gly; D-Pro; (S)-spirolactam-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>t</sup>); D-Asp(OBu<sup>t</sup>); D-Glu(OBu<sup>t</sup>); D-Thr(OBu<sup>t</sup>);
- 15 D-Cys(OBu<sup>t</sup>); D-Ser(OR<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aza-amino acid; D-His which may be substituted on the imidazole ring by a (C<sub>1</sub>-C<sub>6</sub>)alkyl or a benzyl group; an aliphatic D-amino acid with a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic D-amino acid;
- 20 - A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;
- A8 is a basic L-amino acid;
- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or a group -NHR<sub>2</sub> where R<sub>2</sub> is a (C<sub>1</sub>-C<sub>4</sub>)alkyl which
- 25 may be substituted by an hydroxy or one or several fluorine atoms; a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl; or a heterocyclic radical selected from morpholinyl, pyrrolidinyl and piperidyl.

78. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 77 wherein said peptide analogue has the formula (SEQ ID No: 3):



30 in which A7 is Leu, Ile, Nle, Hol, Npg, Cha or Ada, which may be N-alpha-substituted by a methyl or ethyl group optionally substituted by one or several fluorine atoms.

79. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 77 wherein said peptide analogue has the formula (SEQ ID No: 4):



in which:



[R:\LIBH\03736.doc:aak


- A3 and A5 are each independently Phe, Tyr, Trp, 2MeTrp, HPhe, HTyr, Nal, 1Nal, Ba1, Pal, 4Pal, or pClPhe;

- A6 is (S)-spirolactam-Pro; Gly; D-Pro; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(OBu<sup>1</sup>); D-Thr(OBu<sup>1</sup>); D-Cys(OBu<sup>1</sup>); D-His or D-His(Bzl); D-Ala, D-Leu, D-Tle, D-Nle, D-Hol, D-Npg or D-Cha; D-Phe, D-HPhe, D-Tyr, D-HTyr, D-Trp, D-2MeTrp, D-Nal, D-1Nal, D-Bal, D-Pal, D-4Pal, or D-pClPhe; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or D-APhe optionally substituted by an aminotriazolyl group;

- A7 is Leu, Npg or Cha, which may be N-alpha-substituted by a methyl group;

- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or -NC<sub>2</sub>H<sub>5</sub>.

80. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 77 wherein said peptide analogue has the formula (SEQ ID No: 5):



in which:


- A6 is (S)-spirolactam-Pro, D-Leu, D-Ala, D-Nal, D-Phe, D-Ser(OBu<sup>1</sup>) or D-Trp;

- A7 is Leu, MeLeu, Npg or MeNpg;

- Z is GlyNH<sub>2</sub>; azaGlyNH<sub>2</sub>; or -NC<sub>2</sub>H<sub>5</sub>.

81. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 77 to 80 wherein the peptide analogue is selected from the group consisting of leuprorelin, [Npg<sup>7</sup>]-leuprorelin, triptorelin, [Npg<sup>7</sup>]-triptorelin, goserelin, [Npg<sup>7</sup>]-goserelin, buserelin and [Npg<sup>7</sup>]-buserelin.

82. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 76 wherein said peptide analogue has the formula (SEQ ID No: 6):



in which:

- A1 is pGlu; D-pGlu; Sar; AcSar; Pro or a derivative thereof; Ser; D-Ser; Ac-D-Ser; Thr; D-Thr; Ac-D-Thr; or an aromatic D-amino acid which may be acylated;

- A2 is a direct bond or an aromatic D-amino acid;

- A3 is an aromatic L- or D-amino acid;

- A4 is Ala, Ser, D-Ser, MeSer, Ser(OBu<sup>1</sup>), Ser(OBzl) or Thr;

- A5 is an aromatic L-amino acid or a basic L- or D-amino acid;

- A6 is Gly; D-Pro; D-Ser; D-Thr; D-Cys; D-Met; D-Asn; D-Pen; D-(S-Me)Pen; D-(S-Et)Pen; D-Ser(OBu<sup>1</sup>); D-Asp(OBu<sup>1</sup>); D-Glu(O-Bu<sup>1</sup>); D-Thr(O-Bu<sup>1</sup>); D-Cys(O-Bu<sup>1</sup>); D-Ser(O-R<sub>1</sub>) where R<sub>1</sub> is a sugar moiety; an aliphatic D-amino acid with



[R:\LIBH\03736.doc:aak




a (C<sub>1</sub>-C<sub>8</sub>)alkyl or a (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl side chain; an aromatic D-amino acid; D-cyclohexadienyl-Gly; D-perhydronaphthyl-Ala; D-perhydrodiphenyl-Ala; or a basic L- or D-amino acid;

- A7 is a linear, branched or cyclic aliphatic L-amino acid of 3 to 20 carbon atoms which may be N-alpha-substituted by a (C<sub>1</sub>-C<sub>4</sub>)alkyl group optionally substituted by one or several fluorine atoms;

- A8 is a basic L- or D-amino acid;

- Z is GlyNH<sub>2</sub> or D-AlaNH<sub>2</sub>.

83. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 82 wherein the peptide analogue has the formula (SEQ ID No: 7):



in which:

- A5 is Tyr, HTyr, MeTyr, MeHTyr, NicLys or IprLys;
- A6 is (S)spirolactam-Pro, D-Arg, D-NicLys, D-IprLys, D-Cit, D-HCit or D-Asn;
- A7 is Leu, MeLeu, Npg or MeNpg;
- A8 is Arg, NicLys or IprLys.

84. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 82 or 83 wherein the peptide analogue is selected from the group consisting of antide, [Npg<sup>7</sup>]-antide, cetorelix, [Npg<sup>7</sup>]-cetorelix, abarelix and [Npg<sup>7</sup>]-abarelix.

85. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 76 to 84 wherein the α-cyclodextrin derivative is selected from the group consisting of methylated α-cyclodextrin, hexakis(2,3,6-tri-O-methyl)-α-cyclodextrin, carboxymethylated α-cyclodextrin and phosphated α-cyclodextrin.

86. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 85 wherein the α-cyclodextrin derivative is hexakis(2,3,6-tri-O-methyl)-α-cyclodextrin.

87. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 76 to 86 for the treatment or prevention of breast cancer.

88. α-Cyclodextrin when used in combination an LH-RH peptide analogue according to claim 87 which further comprises the sequential, parallel or over a period of time administration of at least one compound selected from the group consisting of an antiestrogen, an aromatase inhibitor and a C<sub>17</sub>-C<sub>20</sub> lyase inhibitor.



[R:\LIBH\03736.doc:aak

89.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 76 to 86 for the treatment or prevention of prostate cancer or benign prostatic hypertrophy.

90.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue  
5 according to claim 89 which further comprises the sequential, parallel or over a period of time administration of at least one compound selected from the group consisting of an antiandrogen, a  $5\alpha$ -reductase inhibitor and a  $C_{17}$ - $C_{20}$  lyase inhibitor.


91.  $\alpha$ -Cyclodextrin when used in combination an LH-RH peptide analogue according to any one of claims 76 to 90 wherein the peptide analogue is delivered to the  
10 gastrointestinal tract of the patient.

**Dated 18 October, 2002**

**Laboratoire Theramex**

**Patent Attorneys for the Applicant/Nominated Person**

**SPRUSON & FERGUSON**




15



[R\LIBH\03736.doc:aak

1/7

FIG.1



\*p &lt; 0.05 vs C. ex.3 in vehicle

2/7





FIG. 2

3/7




Concentration of cyclodextrins

FIG.3

\*  $p < 0.05$ , \*\*\*  $p < 0.001$  vs control group  
 o  $p < 0.05$ , ooo  $p < 0.001$  vs C. ex.2 in vehicle

4/7

\*  $p < 0.05$ , \*\*  $p < 0.01$  vs control without  $\alpha$ -CD\*\*\*  $p < 0.001$  vs control  $\alpha$ -CD

5/7





FIG.5

6/7




\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$  vs control  $\alpha$ -CD



7/7

FIG.7



(\*\*p&lt;0.01;\*\*\*p&lt;0.001 vs. control unstimulated LH levels)