

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 200189257 B2**
(10) Patent No. **776101**

(54) Title
Production method of 2-cyclohexyl-2-hydroxy 2-phenylacetic acid
intermediate therefor and production method thereof

(51)⁶ International Patent Classification(s)
C07C 069/732 C07C 059/54
C07C 051/09

(21) Application No: 200189257 (22) Application Date: 2001.11.07

(30) Priority Data

(31) Number (32) Date (33) Country
2000-339437 2000.11.07 JP

(43) Publication Date : 2002.05.09
(43) Publication Journal Date : 2002.05.09
(44) Accepted Journal Date : 2004.08.26

(71) Applicant(s)
Sumika Fine Chemicals Co., Ltd.

(72) Inventor(s)
Tetsuya Ikemoto; Wei-Guo Gao; Mitsuhiro
Takeda; Masami Igi

(74) Agent/Attorney
Griffith Hack,GPO Box 4164,SYDNEY NSW 2001

(56) Related Art
US 6180823

ABSTRACT OF THE DISCLOSURE

The present invention relates to production of 2-cyclohexyl-2-hydroxy-2-phenylacetic acid useful as an intermediate for pharmaceutical products, by an industrial means, economically, safely in a good yield. Novel 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid ester obtained by reacting cyclohexene and benzoylformic acid ester in the presence of a Lewis acid is hydrolyzed and reduced to give 2-cyclohexyl-2-hydroxy-2-phenylacetic acid.

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
221

AUSTRALIA
Patents Act 1990

COMPLETE SPECIFICATION
STANDARD PATENT

Applicant(s):

SUMIKA FINE CHEMICALS CO., LTD.

Invention Title:

PRODUCTION METHOD OF 2-CYCLOHEXYL-2-HYDROXY-2-
PHENYLACETIC ACID INTERMEDIATE THEREFOR AND
PRODUCTION METHOD THEREOF

1
2
3
4
5
6
7
8
9
10

The following statement is a full description of this
invention, including the best method of performing it known to
me/us:

SPECIFICATION

PRODUCTION METHOD OF 2-CYCLOHEXYL-2-HYDROXY-2-PHENYLACETIC ACID INTERMEDIATE THEREFOR AND PRODUCTION METHOD THEREOF

TECHNICAL FIELD OF THE INVENTION

5 The present invention relates to a production method of
2-cyclohexyl-2-hydroxy-2-phenylacetic acid useful as an
intermediate for pharmaceutical products and the like, an
intermediate therefor and a production method thereof.

BACKGROUND OF THE INVENTION

10 2-Cyclohexyl-2-hydroxy-2-phenylacetic acid and ester
thereof are useful as intermediates for pharmaceutical products,
such as oxybutynin, which is a therapeutic agent for
pollakiuria, and the like. There have been proposed various
production methods of 2-cyclohexyl-2-hydroxy-2-phenylacetic
15 acid and ester thereof.

In J. Org. Chem., Vol. 42, No. 17, 2948-2949 (1977), 2-
cyclohexyl-2-hydroxy-2-phenylacetic acid ester is obtained by
reacting alkyl mandelate and lithium diisopropylamide, and then
reacting the obtained reaction product with cyclohexyl iodide.
20 According to this method, expensive lithium diisopropylamide is
used in 2 equivalents or more relative to alkyl mandelate and
expensive and unstable cyclohexyl iodide is used, which makes
this method industrially disadvantageous.

According to JP-A-11-193271, ethyl benzoylformate and
25 cyclohexylmagnesium bromide are condensed to give ethyl 2-
cyclohexyl-2-hydroxy-2-phenylacetate, which is hydrolyzed to
give 2-cyclohexyl-2-hydroxy-2-phenylacetic acid. In this
condensation reaction, diethyl ether having a boiling point of
35°C is used in an amount of about 12-fold volume relative to
30 ethyl benzoylformate, manipulation of dangerous reaction using
the boiling point of diethyl ether is required and the yield is
as low as 53.3%. In an attempt to improve this method into an
industrially safe one, the present inventors used
tetrahydrofuran instead of diethyl ether as a solvent and

reproduced the reaction. As a result, there occurred reduction due to Grignard reagent to produce ethyl mandelate as a by-product, as well as addition of the Grignard reagent to ester bond. The proportion of the starting material that became a by-product was 42% and the yield of the objective product was as low as 58%. After hydrolysis of the obtained crude ethyl 2-cyclohexyl-2-hydroxy-2-phenylacetate, it was subjected to recrystallization and the like to increase the purity but the yield from ethyl benzoylformate of 2-cyclohexyl-2-hydroxy-2-phenylacetic acid did not exceed 43%.

Therefore, it is concluded that none of the above methods produces 2-cyclohexyl-2-hydroxy-2-phenylacetic acid or ester thereof industrially, economically and safely in a good yield.

There have been also proposed various production methods of optically active 2-cyclohexyl-2-hydroxy-2-phenylacetic acid and ester thereof.

In WO00/23414, a method for optical resolution of a racemate of 2-cyclohexyl-2-hydroxy-2-phenylacetic acid using optically active amine is described. This method, nevertheless, is not entirely efficient because it produces undesirable enantiomer in a half amount.

In WO00/27786, moreover, (i) optically active mandelic acid is converted to 2-tert-butyl-5-phenyl-4-oxo-1,3-dioxolane protected by tert-butyl at the 2-position, by the use of pivalaldehyde, (ii) the resulting compound is reacted with cyclohexanone in the presence of lithium bis(trimethylsilyl)amide at -78°C , and (iii) the reaction product is subjected to reduction and hydrolysis to give optically active 2-cyclohexyl-2-hydroxy-2-phenylacetic acid. This method is industrially disadvantageous in that lithium bis(trimethylsilyl)amide is expensive, the reaction needs to be carried out at an extremely low temperature of -78°C , a number of steps are required and the like.

In J. Med. Chem., 40, 117-124 (1997), moreover, as in WO00/27786, 2-tert-butyl-5-phenyl-4-oxo-1,3-dioxolane protected by tert-butyl at the 2-position is reacted with 3-cyclohexenyl bromide in the presence of lithium diisopropylamide at -80°C and subjected to reduction and hydrolysis to give optically active 2-cyclohexyl-2-hydroxy-2-phenylacetic acid. This method is industrially disadvantageous in that lithium diisopropylamide and 3-cyclohexenyl bromide are expensive, the reaction needs to be carried out at an extremely low temperature of -80°C and the like.

From the foregoing, it follows that none of the above methods produces optically active 2-cyclohexyl-2-hydroxy-2-phenylacetic acid and ester thereof industrially, economically and safely in a good yield.

SUMMARY OF THE INVENTION

It would be advantageous if at least preferred embodiments of the present invention provide an industrial method of producing 2-cyclohexyl-2-hydroxy-2-phenylacetic acid, optically active form thereof and intermediates therefor, economically and safely in a good yield.

To solve the above-mentioned problems, the present inventors considered that novel 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid ester (compound of the formula [II] below) obtained by reacting cyclohexene and benzoylformic acid ester may be usable as a precursor of 2-cyclohexyl-2-hydroxy-2-phenylacetic acid and ester thereof, which are useful as intermediates for pharmaceutical products.


The reaction between cyclohexene and benzoylformic acid ester is called an ene reaction. In general, it is considered that a sterically complicated compound, such as benzoylformic acid ester, has poor reactivity and the reaction with olefin having a double bond in a ring, such as cyclohexene, is difficult to the degree that a reaction in a good yield is remotely available.

The present inventors have conducted this reaction in

the presence of a Lewis acid, and surprisingly found that highly pure 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid ester could be obtained in a good yield. In addition, this reaction proceeds by an extremely simple manipulation of stirring at room temperature. It was found, therefore, that 2-cyclohexyl-2-hydroxy-2-phenylacetic acid and ester thereof could be obtained by a strategy completely different from conventional production, by the use of 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid ester as a precursor, which is subjected to reduction and hydrolysis.

It was also found that, while an environmentally necessarily preferable solvent, such as methylene chloride, is generally used for the ene reaction, incineratable monochlorobenzene can be used as an alternative solvent. Further, it was found that by converting benzoylformic acid to optically active benzoylformic acid ester having an asymmetric carbon atom in the ester moiety thereof (optically active form of compound of the formula [I] below) and then carrying out the aforementioned ene reaction, asymmetry is induced to give optically active 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid ester, and by subjecting this ester to hydrolysis and reduction, optically active 2-cyclohexyl-2-hydroxy-2-phenylacetic acid could be obtained.

Accordingly, the present invention provides (1) a compound of the formula [II]



wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α-(2-cyclohexen-1-yl)-α-hydroxy-


benzyl)carbonyloxy, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl,

5 or an optically active form thereof,

(2) the compound of the above-mentioned (1) wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl,


10 cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl, or an optically active form thereof,

15 (3) a compound of the formula [V]




, an optically active form thereof or a salt thereof,

(4) a method for producing a compound of the formula [II]



20 wherein R' is as defined in the above-mentioned (1), or an optically active form thereof, which method comprising reacting a compound the formula [I]



wherein R is linear or branched chain alkyl having 1 to 15

25 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and benzoylcarbonyloxy, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at

least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl or an optically active form thereof, with cyclohexene in the presence of a Lewis acid,

- 5 (5) the production method of the above-mentioned (4) wherein R and R' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, 10 cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl,

- (6) the production method of the above-mentioned (4), wherein R 15 and R' are each a group having an asymmetric carbon atom,


(7) the production method of the above-mentioned (4), wherein the Lewis acid is an optically active Lewis acid having an asymmetric ligand,

- (8) the production method of the above-mentioned (4), wherein 20 the Lewis acid is titanium tetrachloride,

(9) the production method of any of the above-mentioned (4) -


(8), wherein the reaction is carried out in monochlorobenzene,

(10) a method for producing a compound of the formula [III]



- 25 wherein R'' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α-cyclohexyl-α-hydroxybenzyl)carbonyloxy, 30 or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15


carbon atom(s) and phenyl or an optically active form, which method comprising reducing a compound of the formula [II]




wherein R' is as defined in (1) above, or an optically active form thereof,

(11) the production method of the above-mentioned (10), wherein R' and R'' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl,

(12) a method for producing a compound of the formula [V]



, an optically active form thereof or a salt thereof, which method comprising hydrolyzing a compound of the formula [II]




wherein R' is as defined in the above-mentioned (1), or an optically active form thereof,

(13) the production method of the above-mentioned (12), wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of


linear or branched chain alkyl having 1 to 15 carbon atom(s)
and phenyl,

(14) a method for producing 2-cyclohexyl-2-hydroxy-2-phenylacetic acid of the formula [IV]




5

, an optically active form thereof or a salt thereof, which
method comprising reducing a compound of the formula [V]




, an optically active form thereof or a salt thereof,

10 (15) a method for producing 2-cyclohexyl-2-hydroxy-2-phenylacetic acid of the formula [IV]



, an optically active form thereof or a salt thereof, which
method comprising subjecting a compound of the formula [II]



15


wherein R' is as defined in (1) above, or an optically active
form thereof, to hydrolysis and reduction,

(16) the production method of the above-mentioned (15), wherein
R' is linear or branched chain alkyl having 1 to 15 carbon


20

atom(s), which is optionally substituted by at least one
substituent selected from the group consisting of phenyl,
naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl,
cyclopentyl or norbornyl, which is optionally substituted by at
least one substituent selected from the group consisting of
25 linear or branched chain alkyl having 1 to 15 carbon atom(s)
and phenyl,


- (17) the production method of the above-mentioned (15), which comprises simultaneous hydrolysis and reduction,
 (18) the production method of the above-mentioned (15), which comprises hydrolysis after reduction,
 5 (19) the production method of the above-mentioned (15), which comprises reduction after hydrolysis,
 (20) a method for producing a compound of the formula [III]



wherein R'' is as defined in (10) above, or an optically active
 10 form thereof, which method comprising reacting a compound the formula [I]




wherein R is as defined in (4) above, or an optically active
 form thereof, with cyclohexene in the presence of an Lewis acid
 15 to give a compound of the formula [II]




wherein R' is as defined in (1) above, or an optically active
 form thereof, and reducing the same,


- (21) the production method of the above-mentioned (20) wherein
 20 R' and R'' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at
 25 least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl,
 (22) a method for producing a compound of the formula [V]




, an optically active form thereof or a salt thereof, which method comprising reacting a compound of the formula [I]




5 wherein R is as defined in (4) above, or an optically active form thereof, with cyclohexene in the presence of a Lewis acid to give a compound of the formula [II]



wherein R' is as defined in (1) above, or an optically active
10 form thereof, and hydrolyzing the same,
(23) the production method of the above-mentioned (22), wherein R and R' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl,
15 naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl,
20 (24) a method of producing 2-cyclohexyl-2-hydroxy-2-phenylacetic acid of the formula [IV]




, an optically active form thereof or a salt thereof, which method comprising reacting a compound of the formula [I]



25 wherein R is as defined in (4) above, or an optically active

form thereof, with cyclohexene in the presence of an Lewis acid to give a compound of the formula [II]



- wherein R' is as defined in (1) above, or an optically active form thereof, and subjecting the same to hydrolysis and reduction,
- (25) the production method of the above-mentioned (24), wherein R and R' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl,
- (26) the production method of the above-mentioned (24), which comprises simultaneous hydrolysis and reduction,
- (27) the production method of the above-mentioned (24), which comprises hydrolysis after reduction, and
- (28) the production method of the above-mentioned (24), which comprises reduction after hydrolysis.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is explained in detail in the following.

- Each symbol used in this specification is defined in the following.

- The alkyl in the present invention is linear as long as it does not have a prefix (e.g., iso, neo, sec-, tert- etc.). For example, a simple propyl means linear propyl.
- The "linear or branched chain alkyl having 1 to 15 carbon atom(s)" of the "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at

least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl" is exemplified by methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5 hexyl, heptyl, decyl, pentadecyl and the like, with preference given to alkyl having 1 to 6 carbon atoms, which is more preferably methyl, ethyl, isopropyl and butyl.

The "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one 10 substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl" is exemplified by methyl, ethyl, isopropyl, butyl, 1-phenylethyl, 2-phenylethyl, 1-naphthylethyl, 2-naphthylethyl, cyclohexylmethyl, cyclopentylmethyl, 2-norbornylmethyl, 2-methylbutyl, 1- 15 methylpropyl, 1-methylbutyl, 1-methylpentyl, 1-methylhexyl, 1-methylheptyl, 1-methyloctyl, 1-methylnonyl, 1-methyldecyl and the like, with preference given to methyl, ethyl, isopropyl, butyl and 2-phenylethyl.

The "linear or branched chain alkyl having 1 to 15 20 carbon atom(s)" in the "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and benzoylcarbonyloxy", 25 "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α -(2-cyclohexen-1-yl)- α -hydroxybenzyl)- 30 carbonyloxy" and "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α -cyclohexyl- α -hydroxybenzyl)carbonyloxy"

is as defined for the above-mentioned "linear or branched chain alkyl having 1 to 15 carbon atom(s)".

Examples of the "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and benzoylcarbonyloxy" include examples recited for the above-mentioned "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl", 1-(methoxycarbonyl)ethyl, 1,2-bis(ethoxycarbonyl)-2-(benzoylcarbonyloxy)ethyl and the like. Of these, methyl, ethyl, isopropyl, butyl, 2-phenylethyl, 1-(methoxycarbonyl)ethyl and 1,2-bis(ethoxycarbonyl)-2-(benzoylcarbonyloxy)ethyl are preferable.

Examples of the "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α -(2-cyclohexen-1-yl)- α -hydroxybenzyl)carbonyloxy" include examples recited for the above-mentioned "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl", 1-(methoxycarbonyl)ethyl, 1,2-bis(ethoxycarbonyl)-2-(α -(2-cyclohexen-1-yl)- α -hydroxybenzyl)carbonyloxy)ethyl and the like. Of these, methyl, ethyl, isopropyl, butyl, 2-phenylethyl, 1-(methoxycarbonyl)ethyl and 1,2-bis(ethoxycarbonyl)-2-(α -(2-cyclohexen-1-yl)- α -hydroxybenzyl)carbonyloxy)ethyl are preferable.

Examples of the "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at

least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α -cyclohexyl- α -hydroxybenzyl)carbonyloxy" include examples recited for the above-mentioned "linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl", 1-(methoxycarbonyl)ethyl, 1,2-bis(ethoxycarbonyl)-2-(α -cyclohexyl- α -hydroxybenzyl)carbonyloxyethyl and the like. Of these, methyl, ethyl, isopropyl, butyl, 2-phenylethyl, 1-(methoxycarbonyl)ethyl and 1,2-bis(ethoxycarbonyl)-2-(α -cyclohexyl- α -hydroxybenzyl)carbonyloxyethyl are preferable.

The "linear or branched chain alkyl having 1 to 15 carbon atom(s)" of the "cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl" is as defined for the above-mentioned "linear or branched chain alkyl having 1 to 15 carbon atom(s)".

Examples of the "cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl" include cyclohexyl, 2-methylcyclohexyl, 4-methylcyclohexyl, 4-*t*-butylcyclohexyl, cyclopentyl, 2-methylcyclopentyl, 2-norbornyl, isobornyl, bornyl, menthyl, isomenthyl, neomenthyl, 8-phenylmenthyl and the like, with preference given to cyclohexyl, 2-methylcyclohexyl and cyclopentyl.

In view of the availability of the starting material compound (compound [I]) and the reaction speed in each step, R is particularly preferably methyl or ethyl.

As R, one having an asymmetric carbon atom can be also used, such as menthyl, isomenthyl, neomenthyl, bornyl, 8-

phenylmenthyl and norbornyl having an asymmetric carbon atom;
 1-phenylethyl, 1-naphthylethyl, 2-methylcyclohexyl, 2-
 methylbutyl, 1-methylethyl, 1-methylpropyl, 1-methylbutyl, 1-
 methylpentyl, 1-methylhexyl, 1-methylheptyl, 1-methyloctyl, 1-
 5 methylnonyl, 1-methyldecyl, 1-(methoxycarbonyl)ethyl and 1,2-
 bis(ethoxycarbonyl)-2-(benzoylcarbonyloxy)ethyl, which are R-
 compounds and S-compounds; and the like.

As R', one having an asymmetric carbon atom can be also
 used, such as menthyl, isomenthyl, neomenthyl, bornyl, 8-
 10 phenylmenthyl and norbornyl having an asymmetric carbon atom;
 1-phenylethyl, 1-naphthylethyl, 2-methylcyclohexyl, 2-
 methylbutyl, 1-methylethyl, 1-methylpropyl, 1-methylbutyl, 1-
 methylpentyl, 1-methylhexyl, 1-methylheptyl, 1-methyloctyl, 1-
 methylnonyl, 1-methyldecyl, 1-(methoxycarbonyl)ethyl and 1,2-
 15 bis(ethoxycarbonyl)-2-((α -(2-cyclohexen-1-yl)- α -hydroxy-
 benzyl)carbonyloxy)ethyl, which are R-compounds and S-
 compounds; and the like.


As R'', one having an asymmetric carbon atom can be also
 used, such as menthyl, isomenthyl, neomenthyl, bornyl, 8-
 20 phenylmenthyl and norbornyl having an asymmetric carbon atom;
 1-phenylethyl, 1-naphthylethyl, 2-methylcyclohexyl, 2-
 methylbutyl, 1-methylethyl, 1-methylpropyl, 1-methylbutyl, 1-
 methylpentyl, 1-methylhexyl, 1-methylheptyl, 1-methyloctyl, 1-
 methylnonyl, 1-methyldecyl, 1-(methoxycarbonyl)ethyl, 1,2-
 25 bis(ethoxycarbonyl)-2-((α -cyclohexyl- α -hydroxybenzyl)-
 carbonyloxy)ethyl, which are R-compounds and S-compounds; and
 the like.

In the present invention, the compound [II] - compound
 [V] have an asymmetric carbon atom, wherein, for example,
 30 compound [II] - compound [V] have an asymmetric carbon atom at
 the 2-position of the acetic acid skeleton, compound [II] and
 compound [V] have an asymmetric carbon atom at the binding site
 of cyclohexene, and compound [I], compound [II] and compound
 [III] may have one or more asymmetric carbon atoms at

substituent R. The compound [I] - compound [V] of the present invention encompass any imaginable optically active forms and mixtures thereof (e.g., racemate, enantiomer mixture, diastereomer mixture and the like).

5 The compound [IV] and compound [V] have a carboxyl group and may be in the form of a salt. Examples of the salts of compound [IV] and compound [V] include salts with alkali metal (e.g., sodium, potassium, lithium etc.), amines (e.g., ammonia, methylamine, dimethylamine, triethylamine etc.) and the like.

10 The production method of the present invention is shown in the following reaction scheme.



15 wherein each symbol is as defined above.

1) Production method of compound [II]

The compound [II] is a novel compound, which can be obtained by, for example, reacting compound [I] with cyclohexene in a solvent in the presence of a Lewis acid. The

order of addition of reagents is not particularly limited. For example, compound [I] and cyclohexene are added to a solvent and then a Lewis acid may be added, or compound [I] and a Lewis acid are added to a solvent and then cyclohexene may be added, 5 or a Lewis acid is added to a solvent and then compound [I] and cyclohexene may be added, or cyclohexene and a Lewis acid are added to a solvent and then compound [I] may be added.

The Lewis acid to be used in 1) may be, for example, titanium tetrachloride, titanium tetrabromide, tin 10 tetrachloride, silicon tetrachloride, zirconium tetrachloride, aluminum chloride, dimethylaluminum chloride, methylaluminum dichloride, aluminum bromide, ferric chloride, zinc chloride, zinc bromide, zinc iodide, zinc trifluoromethanesulfonate, magnesium chloride, magnesium bromide, magnesium iodide, 15 magnesium trifluoromethanesulfonate, boron trifluoride, hafnium trifluoromethanesulfonate, lanthanum trifluoromethanesulfonate, scandium trifluoromethanesulfonate, ytterbium trifluoromethanesulfonate and the like, preferably titanium tetrachloride. An optically active Lewis acid having an 20 asymmetric ligand may be used for obtaining an optically active form. Examples of the asymmetric ligand include R- or S-1,1'-bi-2-naphthol and the like.

The amount of use of the Lewis acid is generally 0.005 mol - 5 mol, preferably 0.1 mol - 2 mol, more preferably 1 mol 25 - 1.5 mol, per 1 mol of compound [I].

The amount of cyclohexene used in 1) is generally 0.8 mol - 5 mol, preferably 1 mol - 3 mol, more preferably 1.2 mol - 2.2 mol, per 1 mol of compound [I].

Examples of the solvent to be used in 1) include 30 methylene chloride, 1,2-dichloroethane, monochlorobenzene, 1,2-dichlorobenzene, 2-chlorotoluene, 3-chlorotoluene, 4-chlorotoluene, 2-chloro-m-xylene, 2-chloro-p-xylene, 4-chloro-o-xylene, 2,3-dichlorotoluene, 2,4-dichlorotoluene, 2,5-dichlorotoluene, 2,6-dichlorotoluene, 3,4-dichlorotoluene,

monofluorobenzene, nitrobenzene, carbon disulfide, toluene, acetonitrile, propionitrile, nitromethane, nitroethane, water, methanol, ethanol and the like, with preference given to methylene chloride, 1,2-dichloroethane, monochlorobenzene, 1,2-dichlorobenzene, 2-chlorotoluene, 3-chlorotoluene and 4-chlorotoluene. In view of the yield and incineratability of compound [II], monochlorobenzene, 2-chlorotoluene, 4-chlorotoluene and nitrobenzene are more preferable, which is most preferably monochlorobenzene.

10 The amount of use of the solvent is generally 1 L - 50 L, preferably 3 L - 30 L, more preferably 5 L - 20 L, per 1 kg of compound [I].

While the reaction conditions in 1) depend on the reagent to be used and the like, the reaction is generally 15 carried out at -30°C to 150°C, preferably 0°C to 80°C, more preferably 15°C to 60°C, generally for 1 h - 24 h, preferably 3 h - 15 h.

The compound [II] can be separated and purified by conventional methods. For example, a reaction mixture is 20 poured into water, and after partitioning, an organic layer is washed and filtrated, and the obtained filtrate is washed, dried and concentrated under reduced pressure to separate compound [II]. After separation, silica gel column chromatography may be applied for purification. When compound 25 [II] is obtained in the form of a mixture of an optically active form, it can be resolved into each optically active form by conventional methods such as optical resolution and the like.

The compound [I], which is a starting material in 1), may be commercially available. Alternatively, one obtained by 30 transesterification of methyl benzoylformate and a compound of ROH (R is as defined above), by esterification of benzoylformic acid and a compound of ROH (R is as defined above), or by reaction of a reactive derivative of benzoylformic acid and a compound of ROH (R is as defined above) can be used. An

optically active compound [I] wherein R has an asymmetric carbon atom can be obtained using optically active ROH wherein R has an asymmetric carbon atom (e.g., compound such as lactic acid ester (e.g., methyl lactate etc.) and tartaric acid ester (e.g., diethyl tartrate etc.) and the like). When an optically active compound [I] wherein R has an asymmetric carbon atom is used as the starting material for the "production method of compound [II]", the carbon atom at the 2-position of the acetic acid skeleton induces chirality to produce an optically active compound [II].

2) Production method of compound [IV]

The compound [IV] can be produced by subjecting compound [II] to hydrolysis and reduction. The order of hydrolysis and reduction is not particularly limited. For example,

- Route 1: hydrolysis after reduction of compound [II],
Route 2: reduction after hydrolysis of compound [II], or
Route 3: simultaneous hydrolysis and reduction of compound [II] is employed to give compound [IV]. The compound [II], which is a starting material used for Routes 1 - 3, can be produced according to the method of the above-mentioned 1). In the following, each Route is explained in detail.

2-1) Route 1

- In Route 1, compound [II] is reduced to give compound [III], which is hydrolyzed to give compound [IV]. The production method of compound [III], namely, reduction of compound [II] is described in detail.

2-1-1) Production method of compound [III]

- The compound [III] can be obtained by reduction of compound [II]. The reduction can be conducted according to a conventional method. For example, compound [II] is reacted with hydrogen in a solvent in the presence of a reduction catalyst. It is also possible to add active charcoal simultaneously with the reduction catalyst. When an optically active compound [II] is used as a starting material, optically

active compound [III] retaining the steric configuration of the carbon atom at the 2-position of the acetic acid skeleton of the starting material can be obtained.

Examples of the reduction catalyst to be used in 2-1-1) include palladium carbon, palladium, palladium hydroxide, platinum oxide, platinum, platinum carbon, ruthenium oxide, ruthenium, ruthenium carbon and the like, with preference given to palladium carbon and platinum oxide, and more preference given to palladium carbon. The amount of use of the reduction catalyst is generally 0.00001 mol - 0.2 mol, preferably 0.0001 mol - 0.1 mol, per 1 mol of compound [II].

Examples of the solvent to be used in 2-1-1) include water, methanol, ethanol, isopropyl alcohol, acetic acid, ethyl acetate and the like, and mixed solvents thereof, with preference given to water, methanol, ethanol, a mixed solvent of methanol and water and a mixed solvent of ethanol and water. The residual solvent, by-product and the like from the production process of compound [II], which is the starting material, may cause production of a slight amount of an acid due to the simultaneous reduction during reduction of compound [II]. In such a case, an appropriate amount of a base (e.g., triethylamine, sodium acetate and the like) to neutralize the acid produced during this reduction step is preferably added.

The amount of use of the solvent is generally 0.1 L - 20 L, preferably 0.5 L - 8 L, more preferably 1 L - 6 L, relative to 1 kg of compound [II].

The amount of use of hydrogen in 2-1-1) is not particularly limited as long as it can reduce compound [II].

While the reaction conditions in 2-1-1) depend on the kind and the amount of the reduction catalyst, the starting material to be used and the like, the reaction is carried out at generally -20°C to 150°C, preferably 0°C to 100°C, more preferably 15°C to 60°C, generally for 1 h - 24 h, preferably 3 h - 15 h.

The reaction pressure in 2-1-1) is generally 101 kPa - 3040 kPa, preferably 101 kPa - 2020 kPa, more preferably 202 kPa - 1530 kPa.

The compound [III] may be subjected to hydrolysis
5 without separation or purification. For separation and purification, conventional methods can be used. When compound [III] is obtained as a mixture of an optically active form, it can be resolved into each optically active form according to a conventional method such as optical resolution and the like.

10 2-1-2) Hydrolysis of compound [III]

The hydrolysis of compound [III] is now described in detail.

The hydrolysis of compound [III] is conducted according to a conventional method. For example, by (a) reaction of
15 compound [III] and alkali metal compound in a solvent, followed by neutralization with an acid, where necessary, or by (b) reaction of compound [III] with an inorganic acid in a solvent, compound [IV] or a salt thereof can be obtained. When an optically active compound [III] is used as a starting material,
20 optically active compound [IV] retaining the steric configuration of the starting material can be obtained.

As the solvent to be used in (a) and (b), there are mentioned, for example, methanol, ethanol, water and the like and mixed solvents thereof, with preference given to a mixed
25 solvent of methanol and water and a mixed solvent of ethanol and water. As the solvent, the solvent used for reduction of compound [II] can be used as it is for this hydrolysis. When the solvent used for reduction of compound [II] is utilized, evaporation of the solvent after production of compound [III]
30 is not necessary, which is economical and shortens the step. The amount of use of this solvent is generally 1 L - 50 L, preferably 5 L - 20 L, in the case of (a), and generally 1 L - 30 L, preferably 5 L - 15 L, in the case of (b), per 1 kg of compound [III] (per 1 kg of compound [II] when subjected to

hydrolysis without separation of compound [III] after reduction of compound [II]).

As the alkali metal compound to be used in (a), for example, sodium hydroxide and potassium hydroxide are mentioned, 5 with preference given to sodium hydroxide. The amount of use of this alkali metal compound is generally 1 mol - 3.3 mol, preferably 1 mol - 2.2 mol, per 1 mol of compound [III] (per 1 mol of compound [II] when subjected to hydrolysis without separation of compound [III] after reduction of compound [II]). 10 While the amount varies depending on the kind of alkali metal compound and solvent, alkali metal compound is used in an amount to make the concentration in the reaction system generally 0.1 wt% - 50 wt%, preferably 3 wt% - 25 wt%. The alkali metal compound may be added to the reaction system as it 15 is, but addition in the form of an aqueous solution or alcohol solution is preferable. The amount of use of the solvent to dissolve the alkali metal compound is included in the amount of use of the above-mentioned solvent.

Examples of the inorganic acid to be used in (b) include 20 hydrochloric acid, sulfuric acid, phosphoric acid and the like, with preference given to hydrochloric acid and sulfuric acid. The amount of use of the inorganic acid is generally 0.01 mol - 5 mol, preferably 0.1 mol - 2 mol, per 1 mol of compound [III] (per 1 mol of compound [II] when subjected to hydrolysis 25 without separation of compound [III] after reduction of compound [II]).

The acid to be used for neutralization in (a) is exemplified by those mentioned with regard to the "inorganic acid to be used in (b)" above. The acid is used in an amount 30 that makes the pH of the reaction mixture generally not more than 7, preferably not more than 4. When neutralization with acid is not performed in (a), an alkali metal salt of compound [IV] is obtained and when the starting material (compound [III]) is in an optically active form, an alkali metal salt of

optically active compound [IV] is obtained.

In (a), while the conditions of the reaction between compound [III] and alkali metal compound depend on the kind and the amount of use of compound [III] and alkali metal compound, 5 and the like, the reaction completes generally at a temperature of not less than 0°C, preferably not less than 60°C, and generally not higher than 180°C, preferably not higher than 150°C, more preferably not higher than 120°C, and most preferably not higher than 100°C, generally for not less than 10 30 min, preferably not less than 2 h; and generally within 15 h, preferably within 8 h, more preferably within 6 h.

The reaction conditions in (b) depend on the kind and the amount of use of compound [III] and inorganic acid to be used, and the like, but the reaction generally ends at 0°C - 15 120°C, preferably 60°C - 100°C, generally for 1 h - 24 h, preferably 4 h - 10 h.

2-2) Route 2

In Route 2, compound [II] is hydrolyzed to give novel compound [V], which is reduced to give compound [IV]. First, 20 the production method of compound [V], namely, hydrolysis of compound [II] is described in detail.

2-2-1) Production method of compound [V]

The compound [V] is a novel compound and can be obtained by hydrolysis of compound [II]. The compound [II] can be 25 hydrolyzed according to completely the same method as in the above-mentioned "2-1-2) Hydrolysis of compound [III]", and the alkali metal compound, inorganic acid and solvent to be used are the same. The amounts of alkali metal compound, inorganic acid and solvent to be used in 2-2-1) are those obtained by 30 changing the standard from compound [III] to compound [II].

However, when compound [II] is hydrolyzed according to the method of (a) in the above-mentioned "2-1-2) Hydrolysis of compound [III]", neutralization with an acid after hydrolysis is not necessarily essential, and compound [V] can be subjected

to reduction as it is.

When an optically active compound [II] is used as a starting material, optically active compound [V] retaining the steric configuration of the starting material can be obtained.

5 It is needless to say that the method of (a) without neutralization after hydrolysis of compound [II] affords an alkali metal salt of optically active compound [V].

The compound [V] can be separated and purified by a conventional method. For example, the crystals obtained by
10 reaction are aged where necessary, and subjected to filtration, washing and drying to separate and purify compound [V]. The compound [V] can be also used for the next step without separation. When compound [V] is obtained in the form of a mixture of an optically active form, it can be resolved into
15 each optically active form by conventional methods such as optical resolution and the like.

2-2-2) Reduction of compound [V]

The compound [V] can be reduced according to the method completely the same as the method of the above-mentioned "2-1-
20 1) Production method of compound [III]", and the reduction catalyst and solvent to be used are the same. The amounts of reduction catalyst and solvent to be used in 2-2-2) are those obtained by changing the standard from compound [II] to compound [V]. When the compound [V] is reduced without
25 separation after hydrolysis of compound [II], the standard is compound [II]. As in Route 1, the use of the same solvent for hydrolysis and reduction in Route 2 is preferable from the aspect of the number of steps and economical production.

When an optically active compound [V] is used as a
30 starting material, optically active compound [IV] retaining the steric configuration of the carbon atom at the 2-position of the acetic acid skeleton of the starting material can be obtained.

2-3) Route 3

In Route 3, since hydrolysis and reduction of compound [II] simultaneously proceed, the reaction system is considered to contain compound [III] and compound [V] as intermediates. According to Route 3, the reagents necessary for reduction and
5 hydrolysis and compound [II] are simultaneously added to the solvent and reacted to give compound [IV]. When an optically active compound [II] is used as a starting material, optically active compound [IV] retaining the steric configuration of the carbon atom at the 2-position of the acetic acid skeleton of
10 the starting material can be obtained.

The solvent to be used in Route 3 may be the solvent used for "2-1-2) Hydrolysis of compound [III]", and the amount of use of the solvent is the same as that in the case of "2-1-2) Hydrolysis of compound [III]" (standard being not compound
15 [III] but compound [II]).

The reduction in Route 3 can be carried out using a reduction catalyst and hydrogen in the same manner as in "2-1-1) Production method of compound [III]". The reduction catalyst is similar to that used for "2-1-1) Production method
20 of compound [III]". The amounts of use of hydrogen and reduction reagent in Route 3 are the same as those in the case of "2-1-1) Production method of compound [III]".

The hydrolysis in Route 3 can be carried out according to two methods using (A) alkali metal compound (where necessary,
25 an acid) or (B) inorganic acid, as in the "2-1-2) Hydrolysis of compound [III]". Each reagent may be that used for the "2-1-2) Hydrolysis of compound [III]" and the amount of use of each reagent is the same as that in the "2-1-2) Hydrolysis of compound [III]" (standard being not compound [III] but compound
30 [II]).

The Route 3 proceeds at a reaction pressure of generally 101 kPa - 3040 kPa, preferably 101 kPa - 2020 kPa, more preferably 202 kPa - 1530 kPa, at generally 0 - 150°C, preferably 50 - 100°C, for generally 1 h - 24 h, preferably 5 h

- 15 h.

The compound [IV] obtained in Routes 1 - 3 can be separated and purified by conventional methods. For example, a reaction mixture is partitioned, the obtained organic layer is cooled to allow precipitation of crystals, and the crystals are aged, filtrated, washed and dried for separation and purification. It is also possible to add an organic solvent, where necessary, before partitioning. For efficient removal of by-produced mandelic acid, benzoylformic acid and the like, the reaction mixture is preferably washed with a base, such as ammonia and the like, before partitioning (when organic solvent is added before partitioning, after the addition)) to make the pH of the washing 4 - 7 (preferably 4.5 - 6).

When compound [IV] or a salt thereof obtained by Routes 1 - 3 is a mixture of an optically active form, it can be resolved into each optically active form by conventional methods such as optical resolution and the like.

The compound [IV] can be introduced into oxybutynin useful as a therapeutic agent for pollakiuria, according to the method described in, for example, WO00/23414.

Examples

The present invention is explained in detail by referring to examples. The present invention is not limited by these examples in any way.

Example 1

Methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate (compound [II])

Methyl benzoylformate (41.0 g, 0.25 mol) and cyclohexene (41.0 g, 0.50 mol) were dissolved in methylene chloride (287 ml) under a nitrogen atmosphere, and titanium tetrachloride (60.7 g, 0.32 mol) was added dropwise thereto at 15 - 25°C over 30 min, which was followed by stirring at 25 - 30°C for 5 h. The reaction mixture was poured into water (287 ml) and partitioned. The organic layer was washed with 10 wt% aqueous

sodium carbonate solution (287 g) and filtrated. The organic layer was washed with water (287 ml) again and dried over anhydrous magnesium sulfate (10 g). The organic layer is concentrated under reduced pressure to give a crude product
5 (62.6 g) of the title compound. The crude product is purified by silica gel column chromatography to give the title compound (44.9 g, yield: 72.9%) as a diastereomer mixture of (2S*,1'S*) compound and (2S*,1'R*) compound.

¹H-NMR(400MHz, CDCl₃) δ=1.24-1.77(4H, m), 1.92-2.06(2H, m),
10 3.14-3.21(1H, m), 3.44(1H, s), 3.80(0.45H, s), 3.81(2.55H, s), 5.13-5.18(0.15H, m), 5.40-5.45(0.85H, m), 5.75-5.80(0.15H, m), 5.92-5.98(0.85H, m), 7.26-7.32(1H, m), 7.33-7.39(2H, m), 7.62-7.69(2H, m) ppm

Example 2

15 2-(2'-Cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid (compound [V])

A diastereomer mixture (44.0 g) of (2S*,1'S*) compound and (2S*,1'R*) compound of methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate reacted and purified by the same method
20 as in Example 1 was dissolved in methanol (440 ml) and 48.5 wt% aqueous sodium hydroxide solution (30.1 g) was added. The mixture was stirred at 65 - 70°C for 1 h. After confirmation of disappearance of ester compound (methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate) by HPLC, methanol was
25 evaporated under reduced pressure. The pH was adjusted to 1 with 6N hydrochloric acid and the resulted crystals were filtrated, washed with water and dried under reduced pressure to give the title compound (40.5 g, yield :97.5%) as a diastereomer mixture of (2S*,1'S*) compound and (2S*,1'R*)
30 compound.

¹H-NMR(400MHz, CDCl₃) δ=1.18-1.76(4H, m), 1.84-1.91(0.3H, m), 1.93-2.07(1.7H, m), 3.00-3.40(1H, br), 3.23-3.37(1H, m), 5.11-5.16(0.15H, m), 5.52-5.57(0.85H, m), 5.85-5.91(0.15H, m), 6.04-6.11(0.85H, m), 7.28-7.40(3H, m), 7.64-7.70(2H, m) ppm

Example 3

2-(2'-Cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid (compound [V])

Crude methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate (62.0 g) obtained on a similar scale to and by the method of Example 1 was dissolved in methanol (410 ml) and 48.5 wt% aqueous sodium hydroxide solution (41.2 g) was added. The mixture was stirred at 65 - 70°C for 1 h. After confirmation of disappearance of ester compound (methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate) by HPLC, methanol was evaporated at 65 - 85°C under reduced pressure. Toluene (300 ml) and water (100 ml) were in poured, and the aqueous layer was adjusted to pH 1 or below at 60 - 70°C with 6N hydrochloric acid. The mixture was partitioned by allowing to stand at 60 - 70°C and the organic layer was cooled to 0 - 5°C. The precipitated crystals were aged for 1 h at the same temperature, filtrated, washed with cooled toluene (50 ml) and dried under reduced pressure at 65 - 75°C to give the title compound as a diastereomer mixture (34.0 g) of (2S*,1'S*) compound and (2S*,1'R*) compound.
yield from methyl benzoylformate: 58.5%
purity: 98.8%.

¹H-NMR(400MHz, CDCl₃) δ=1.18-1.76(4H, m), 1.84-1.91(0.2H, m), 1.93-2.07(1.8H, m), 3.00-3.40(1H, br), 3.23-3.37(1H, m), 5.11-5.16(0.1H, m), 5.52-5.57(0.9H, m), 5.85-5.91(0.1H, m), 6.04-6.11(0.9H, m), 7.28-7.40(3H, m), 7.64-7.70(2H, m) ppm

Example 4

2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (compound [IV], Route 1)

Crude methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate (62.6 g) obtained on a similar scale to and by the method of Example 1 was dissolved in methanol (287 ml) and 5% palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 0.82 g) was added. The mixture was stirred in a hydrogen gas

(350 kPa) for 8 h and the catalyst was filtered off. A 48.5 wt% aqueous sodium hydroxide solution (41.5 g) was added and the mixture was stirred at 75 - 80°C for 4 h. After confirmation of disappearance of ester compound (methyl 2-cyclohexyl-2-hydroxy-2-phenylacetate) by HPLC, methanol was evaporated under reduced pressure at 65 - 85°C. Toluene (164 ml) was in poured, and the aqueous layer was adjusted to pH 1 or below with 6N hydrochloric acid at 60 - 70°C. The mixture was partitioned by allowing to stand at 60 - 70°C and the organic layer was cooled to 0 - 5°C. The organic layer was aged for 1 h at the same temperature, filtrated, washed with cooled toluene (50 ml) and dried under reduced pressure at 65 - 75°C to give the title compound (36.9 g).
yield from methyl benzoylformate: 63.1%

15 purity: 99.3%.

¹H-NMR(400MHz, CDCl₃) δ=0.99-1.48(6H, m), 1.58-1.68(3H, m), 1.78-1.83(1H, m), 2.22-2.30(1H, m), 3.25-3.60(1H, br), 7.26-7.37(3H, m), 7.63-7.66(2H, m) ppm

Example 5

20 2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (compound [IV], Route 2)

Crude methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate (64.1 g) obtained on a similar scale to and by the method of Example 1 was dissolved in methanol (287 ml) and 48.5 wt% aqueous sodium hydroxide solution (41.5 g) was added. The mixture was stirred at 75 - 80°C for 4 h. After confirmation of disappearance of ester compound (methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate) by HPLC, the aqueous layer was neutralized with 6N hydrochloric acid to pH 7. The resulting salt was filtrated, and 5% palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 0.82 g) was added. The mixture was stirred in a hydrogen gas (350 kPa) for 8 h. The catalyst was filtered off and the residue was adjusted to pH 1 with 6N hydrochloric acid. Methanol was evaporated under

reduced pressure and toluene (164 ml) was added. The mixture was partitioned by allowing to stand at 60 - 70°C and the organic layer was cooled to 0 - 5°C. The organic layer was aged at the same temperature for 1 h, filtrated, washed with
5 cooled toluene (50 ml) and dried under reduced pressure at 65 - 75°C to give the title compound (36.4 g).
yield from methyl benzoylformate: 62.2%,
purity: 99.0%,

The ¹H-NMR spectrum data were the same as those obtained
10 in Example 4.

Comparative Example 1

Synthesis of 2-cyclohexyl-2-hydroxy-2-phenylacetic acid by Grignard method

Magnesium (15.8 g) and iodine (200 mg) were added to
15 tetrahydrofuran (73 g) under a nitrogen atmosphere and the mixture was stirred at about 25°C for 30 min. Cyclohexyl bromide (4.1 g) was added dropwise, tetrahydrofuran (291.9 g) was added and cyclohexyl chloride (86 g) was added dropwise at 60 - 70°C. The mixture was stirred at 60 - 75°C for 2 h and
20 analyzed by gas chromatography. As a result, the remainder of cyclohexyl chloride was 0.1%. The mixture was cooled to 20 - 30°C and Grignard solution was added dropwise to a mixture of methyl benzoylformate (82.1 g) and tetrahydrofuran (82 ml) at 6 - 14°C. The dropwise addition ended in 1 h. The mixture was
25 stirred at the same temperature for 1 h and analyzed by HPLC. As a result, methyl benzoylformate was not detected. Tetrahydrofuran was evaporated under reduced pressure at 65 - 80°C. The evaporated amount was 345 ml. Toluene (164 ml) was added and the mixture was added dropwise at 5 - 38°C to 7N
30 hydrochloric acid (215 ml) and partitioned by allowing to stand. The pure yield of methyl 2-cyclohexyl-2-hydroxy-2-phenylacetate in the organic layer was calculated by HPLC external standard method using a reference standard. As a result, yield was 58% (42% was by-product other than the objective product).

Methanol (82 ml) was added to the organic layer and 27 wt% aqueous sodium hydroxide solution (242 g) was added dropwise at 65 - 75°C, and the mixture was stirred at the same temperature for 3 h. After confirmation of disappearance of ester compound (methyl 2-cyclohexyl-2-hydroxy-2-phenylacetate) by HPLC, tetrahydrofuran and methanol were evaporated under reduced pressure at 65 - 85°C. Toluene (410 ml) was in-poured, and the aqueous layer was adjusted to pH 1 or below with 7N hydrochloric acid at 60 - 70°C. The mixture was partitioned by allowing to stand at 70 - 80°C and the organic layer was cooled to 0 - 5°C. The organic layer was aged at the same temperature for 2 h, filtrated, washed with cooled toluene (200 ml) and dried under reduced pressure at 65 - 75°C to give the title compound (51.6 g). yield 44.0%, purity 98.6%.

The ¹H-NMR spectrum data were the same as those obtained in Example 4.

Example 6

2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (compound [IV], Route 3)

Crude methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate (61.5 g) obtained on a similar scale to and by the method of Example 1 was dissolved in methanol (287 ml) and 5% palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 0.82 g) and 25 wt% aqueous sodium hydroxide solution (50.5 g) were added. The mixture was stirred in a hydrogen gas (1013 kPa) at 60 - 65°C for 6 h. 5% Palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 0.82 g) was added and the mixture was further stirred under the same conditions for 4 h. The catalyst was filtered off and methanol was evaporated under reduced pressure. The residue was adjusted to pH 1 with 3N hydrochloric acid and toluene (164 ml) was added. The mixture was partitioned by allowing to stand at 60 - 70°C and the organic layer was cooled to 0 - 5°C. The organic layer was aged for 1 h at the same temperature, filtrated, washed with

cooled toluene (50 ml) and dried under reduced pressure at 65 - 75°C to give the title compound (32.5 g).

yield from methyl benzoylformate: 55.5%

purity:98.8%,

5 The ¹H-NMR spectrum data were the same as those obtained in Example 4.

Example 7

Methyl 2-cyclohexyl-2-hydroxy-2-phenylacetate (compound [III])

Crude methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate (62.8 g) obtained on a similar scale to and by the method of Example 1 was dissolved in methanol (287 ml) and triethylamine (28.7 ml) and 10% palladium carbon (Degussa Japan E106XNN/W, 1.24 g) were added. The mixture was stirred at 40°C in a hydrogen gas (1520 kPa) for 10 h. After the catalyst was filtered off, methanol was evaporated under reduced pressure and the residue was purified by silica gel column chromatography to give the title compound (40.2 g, yield :64.8%). This was used as the reference standard in Examples 9 - 16.

20 ¹H-NMR(400MHz, CDCl₃) δ=1.02-1.48(7H, m), 1.58-1.70(2H, m), 1.75-1.83(1H, m), 2.18-2.27(1H, m), 3.68(1H, s), 3.77(3H, s), 7.23-7.28(1H, m), 7.30-7.36(2H, m), 7.62-7.65(2H, m) ppm

Example 8

2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (compound [IV])

25 2-(2'-Cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid (20.2 g) obtained in Example 2 was dissolved in methanol (200 ml) and 5% palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 1.0 g) was added. The mixture was stirred at 25 - 30°C in a hydrogen gas (206 kPa) for 13 h and the catalyst was filtered off. Methanol was evaporated under reduced pressure to give the title compound (20.1 g, yield: 98.5%).

The ¹H-NMR spectrum data were the same as those obtained in Example 4.

Examples 9 - 16

Methyl 2-cyclohexyl-2-hydroxy-2-phenylacetate (compound [III])

Methyl benzoylformate (1.64 g, 10 mmol) and cyclohexene (1.64 g, 20 mmol) were dissolved in a solvent under a nitrogen atmosphere and a Lewis acid was added. The mixture was stirred
5 at the reaction temperature for the reaction time. The solvent, Lewis acid, reaction temperature and reaction time are as shown in Table 1.

The reaction mixture was poured into 6N aqueous hydrochloric acid (12 ml) and partitioned. The organic layer
10 was washed with 10 wt% aqueous sodium carbonate solution (12 ml) and the resulting precipitate was collected by filtration. The organic layer was washed with water (12 ml) and dried over anhydrous magnesium sulfate (0.5 g). The organic layer was concentrated under reduced pressure to give methyl 2-(2'-
15 cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate. The ¹H-NMR spectrum data were the same as those obtained in Example 1.

There to was added methanol (12 ml) and 5% palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 0.04 g) was added and the mixture was stirred in a hydrogen gas (350 kPa)
20 for 10 h. The yield of methyl 2-cyclohexyl-2-hydroxy-2-phenylacetate in the solution was calculated by HPLC external standard method using the reference standard (compound obtained in Example 7). The results are as shown in Table 1.

Table 1

	Lewis acid	solvent	reaction temperature (°C)	Yield (g)
	kind amount of use (mmol)	kind amount of use (ml)	reaction time (h)	yield (%)
Ex. 9	titanium tetrachloride	Monochlorobenzene	25-40	1.61
	15	16	9	69.1
Ex. 10	titanium tetrachloride	2-chlorotoluene	60-70	1.09
	15	16	9	46.8
Ex. 11	titanium tetrachloride	4-chlorotoluene	60-70	1.01
	15	16	9	43.5
Ex. 12	titanium tetrachloride	nitrobenzene	25-30	1.21
	15	16	9	52.2
Ex. 13	aluminum chloride	methylene chloride	25-30	0.75
	15	12	15	32.2
Ex. 14	aluminum chloride	Monochlorobenzene	25-30	0.37
	15	16	15	15.8
Ex. 15	ferric chloride	methylene chloride	25-30	0.28
	15	12	15	12.1
Ex. 16	zirconium tetrachloride	Monochlorobenzene	25-30	0.05
	15	12	9	2.2

Example 17

5 2-Cyclohexyl-2-hydroxy-2-phenylacetic acid (compound [IV],
Route 3)

To a solution of titanium tetrachloride (11.5 kg, 61 mol) in monochlorobenzene (28.5 L) was added dropwise at 35 - 45°C over 5 h under a nitrogen atmosphere, a mixture of methyl
10 benzoylformate (9.5 kg, 58 mol) and cyclohexene (9.5 kg, 116 mol). The reaction mixture was stirred at 35 - 45°C for 9 h and added dropwise to 11% hydrochloric acid (20.9 kg) at 15 -

40°C. The mixture after hydrolysis was heated to 55°C and the aqueous layer was separated. After washing with 35% hydrochloric acid (10.2 kg), the aqueous layer was separated. The organic layer was concentrated under reduced pressure until
5 the inert temperature reached 85°C at 4 kPa. Water (4.8 kg) was added and evaporated again under reduced pressure. To the residue was added methanol (14.3 L), and then was added 26% aqueous sodium hydroxide solution (0.2 kg) to retain a 40% aqueous methanol solution (25.8 kg) of methyl 2-(2'-cyclohexen-
10 1'-yl)-2-hydroxy-2-phenylacetate.

A half amount (12.9 kg) of the 40% aqueous methanol solution (25.8 kg) of methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetate thus obtained was measured, and methanol (2.4 L) was added thereto. The mixture was heated to 67°C, and 26%
15 aqueous sodium hydroxide solution (8.8 kg) was added dropwise at 65 - 80°C. After the completion of the dropwise addition, the mixture was heated as it was in an autoclave, and stirred at 130°C, 0.45 Mpa for 3 h. The reaction mixture was cooled to 60°C and adjusted to pH 5.5 with 30% sulfuric acid (4.7 kg).
20 The reaction mixture was heated to 60°C. Active charcoal (0.5 kg) and 5% palladium carbon (N.E. CHEMCAT CORPORATION NX-Type, wet amount 0.4 kg, dry amount 0.2 kg) were added and the inside of the system was thoroughly substituted with nitrogen. Then, the inside of the system was substituted with hydrogen and
25 hydrogen was fed according to absorption, thereby raising the final pressure from 101 kPa to 800 kPa, and the mixture was maintained further at 800 kPa, 60°C for 2 h. The inside of the system was substituted with nitrogen, and the catalyst and active charcoal were filtered off. The residue was washed with
30 a mixture (11 kg) of methanol-water (2:1). The filtrate and washing were combined to give a 30% aqueous methanol solution (36.8 kg) of sodium 2-cyclohexyl-2-hydroxy-2-phenylacetate.

The remaining half amount of the 40% aqueous methanol solution of methyl 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-

phenylacetate was treated in the same manner to give a 30% aqueous methanol solution (37.4 kg) of sodium 2-cyclohexyl-2-hydroxy-2-phenylacetate. The solvent was evaporated from the thus-obtained aqueous methanol solution (74.2 kg) of sodium 2-cyclohexyl-2-hydroxy-2-phenylacetate at atmospheric pressure at an inert temperature of 86°C. To the residue was added toluene (57 L), and then was added 18% sulfuric acid (15.8 kg) at around 70°C to adjust the mixture to pH 1. After stirring and standing of the mixture, the aqueous layer was removed by partitioning. To the organic layer were added toluene (9.5 L), warm water (19 L) and 28% aqueous ammonia (0.4 kg), and after stirring and standing, the aqueous layer was removed by partitioning. The organic layer was partitioned twice and washed with warm water (19 L) at 70°C twice. Toluene (ca. 10 L) was evaporated under reduced pressure at an inert temperature of 85 - 95°C. The mixture was heated to 95°C to completely dissolve crystals. The solution was cooled to 5°C over about 8 h, and the resulting crystals were collected by filtration and dried to give the title compound (yield: 7.72 kg, yield :58.5%, purity: 99.8%).

Example 18

Optically active 2-cyclohexyl-2-hydroxy-2-phenylacetic acid

To a solution of benzoylformic acid (9.0 g) in toluene (135 ml) was added thionyl chloride (8.2 g) and the reaction mixture was stirred at 75°C for 2 h. A solution, from which about 40 ml of the solvent had been evaporated under reduced pressure, was added dropwise to a solution of (S)-methyl lactate (7.1 g) and triethylamine (6.6 g) in toluene (60 ml) at 2 - 5°C and the mixture was stirred at 20°C for 14 h. Water (135 ml) was added to the mixture and the mixture was partitioned. The organic layer was washed successively with 10% aqueous sodium carbonate solution, water and saturated brine, and the organic solvent was evaporated to give methyl (S)-2-(benzoylformyloxy)propionate (7.6 g).

The obtained methyl (S)-2-(benzoylformyloxy)propionate (7.4 g) was dissolved in monochlorobenzene (74 ml), and cyclohexene (5.2 g) and titanium tetrachloride (7.8 g) were added. The reaction mixture was stirred at 20 - 25°C for 15 h. The reaction mixture was poured into 6N hydrochloric acid (74 ml) and extracted with toluene (37 ml). The extract was washed with 6N hydrochloric acid (37 ml). To the organic layer were added 25% aqueous sodium hydroxide solution (20 g) and methanol (74 ml) and the reaction mixture was stirred at 60 - 65°C for 2 h. Water (74 ml) and toluene (74 ml) were added to the reaction mixture and the partitioned organic layer was combined with the extract obtained by extracting the aqueous layer again with toluene (74 ml) and concentrated. 6N Hydrochloric acid (29 g) was added to the concentrated residue and the mixture was extracted with ethyl acetate (74 ml). The organic layer was washed with water and concentrated to give a diastereomer mixture (5.2 g) of crude optically active 2-(2'-cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid. This (4.1 g) was dissolved in methanol (40 ml) and 5% palladium carbon (Kawaken Fine Chemicals Co., Ltd., type M, 3 g) was added. The mixture was stirred at atmospheric pressure in a hydrogen gas for 15 min. The catalyst was filtered off and the solvent was evaporated. The residue was purified by silica gel column chromatography (eluent: toluene/ethyl acetate) to give the title compound (3.3 g, yield from benzoylformic acid: 31%). This was amidated with (S)-phenylethylamine by a conventional method and the optical purity was measured by HPLC, which was found to be 22% e.e.

Example 19

In the same manner as in Example 18 except that (L)-(+)-diethyl tartrate was used at 1/2 molar ratio instead of (S)-methyl lactate, (2R,3R)-diethyl bis(benzoylformyloxy)-butanedionate was synthesized, from which the title compound was obtained (yield from benzoylformic acid: 20%). This was amidated with (S)-phenylethylamine by a conventional method and

the optical purity was measured by HPLC, which was found to be 36% e.e.

According to the present invention, novel 2-(2'-
5 cyclohexen-1'-yl)-2-hydroxy-2-phenylacetic acid ester (compound [II]), obtained by reacting cyclohexene and benzoylformic acid ester in the presence of a Lewis acid, is used as an intermediate and hydrolyzed and reduced to give 2-cyclohexyl-2-hydroxy-2-phenylacetic acid useful as an intermediate for
10 pharmaceutical products, by an industrial method, economically, safely and in a good yield.

This application is based on a patent application No. 2000-339437 filed in Japan, the contents of which are hereby
15 incorporated by reference.


A reference herein to a prior art document does not constitute an admission that the document forms part of the common general knowledge in the art in Australia.

20

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as
25 "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

The claims defining the invention are as follows:

1. A compound of the formula [II]




5 wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α-(2-cyclohexen-1-yl)-α-hydroxy-
10 benzyl)carbonyloxy, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl, or an optically active form thereof.

15

2. The compound of claim 1, wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and
20 norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl, or an optically active form thereof.


25

3. A compound of the formula [V]




, an optically active form thereof or a salt thereof.

30 4. A method for producing a compound of the formula [II]



wherein R' is as defined in claim 1, or an optically active form thereof, which method comprising reacting a compound the formula [I]



5 wherein R is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, 10 ethoxycarbonyl and benzoylcarbonyloxy, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl or an optically active form thereof, with 15 cyclohexene in the presence of a Lewis acid.

5. The production method of claim 4, wherein R and R' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent 20 selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) 25 and phenyl.


6. The production method of claim 4, wherein R and R' are each a group having an asymmetric carbon atom.

30 7. The production method of claim 4, wherein the Lewis acid is an optically active Lewis acid having an asymmetric ligand.


8. The production method of claim 4, wherein the Lewis acid is titanium tetrachloride.

9. The production method of any of claims 4 to 8, wherein the reaction is carried out in monochlorobenzene.

10. A method for producing a compound of the formula [III]



10 wherein R'' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl, norbornyl, methoxycarbonyl, ethoxycarbonyl and (α-cyclohexyl-α-hydroxybenzyl)carbonyloxy, 15 or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl or an optically active form, which method comprising reducing a compound of the formula [II]




20 wherein R' is as defined in claim 1, or an optically active form thereof.


11. The production method of claim 10, wherein R' and R'' are 25 each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at 30 least one substituent selected from the group consisting of

linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl.

12. A method for producing a compound of the formula [V]



5 , an optically active form thereof or a salt thereof, which method comprising hydrolyzing a compound of the formula [II]




wherein R' is as defined in claim 1,
10 or an optically active form thereof.


13. The production method of claim 12, wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected
15 from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl.

20

14. A method for producing 2-cyclohexyl-2-hydroxy-2-phenylacetic acid of the formula [IV]




, an optically active form thereof or a salt thereof, which
25 method comprising reducing a compound of the formula [V]




, an optically active form thereof or a salt thereof.

15. A method for producing 2-cyclohexyl-2-hydroxy-2-phenylacetic acid of the formula [IV]



5 , an optically active form thereof or a salt thereof, which method comprising subjecting a compound of the formula [II]



10 wherein R' is as defined in claim 1, or an optically active form thereof, to hydrolysis and reduction.


16. The production method of claim 15, wherein R' is linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected
15 from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl.

20 17. The production method of claim 15, which comprises simultaneous hydrolysis and reduction.


18. The production method of claim 15, which comprises
25 hydrolysis after reduction.

19. The production method of claim 15, which comprises reduction after hydrolysis.


30 20. A method for producing a compound of the formula [III]



wherein R'' is as defined in claim 10, or an optically active form thereof, which method comprising reacting a compound the formula [I]




5 wherein R is as defined in claim 4, or an optically active form thereof, with cyclohexene in the presence of a Lewis acid to give a compound of the formula [II]




10 wherein R' is as defined in claim 1, or an optically active form thereof, and reducing the same.

21. The production method of claim 20, wherein R' and R'' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of
20 linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl.


22. A method for producing a compound of the formula [V]



25 , an optically active form thereof or a salt thereof, which method comprising reacting a compound of the formula [I]




wherein R is as defined in claim 4, or an optically active form thereof, with cyclohexene in the presence of a Lewis acid to give a compound of the formula [II]




5 wherein R' is as defined in claim 1, or an optically active form thereof, and hydrolyzing the same.

23. The production method of claim 22, wherein R and R' are
 10 each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at
 15 least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl.


24. A method of producing 2-cyclohexyl-2-hydroxy-2-phenylacetic
 20 acid of the formula [IV]



, an optically active form thereof or a salt thereof, which method comprising reacting a compound of the formula [I]



25 wherein R is as defined in claim 4, or an optically active form thereof, with cyclohexene in the presence of a Lewis acid to give a compound of the formula [II]



wherein R' is as defined in claim 1, or an optically active form thereof, and subjecting the same to hydrolysis and reduction.

5

25. The production method of claim 24, wherein R and R' are each linear or branched chain alkyl having 1 to 15 carbon atom(s), which is optionally substituted by at least one substituent selected from the group consisting of phenyl, 10 naphthyl, cyclohexyl, cyclopentyl and norbornyl, or cyclohexyl, cyclopentyl or norbornyl, which is optionally substituted by at least one substituent selected from the group consisting of linear or branched chain alkyl having 1 to 15 carbon atom(s) and phenyl.

15

26. The production method of claim 24, which comprises simultaneous hydrolysis and reduction.

27. The production method of claim 24, which comprises 20 hydrolysis after reduction.

28. The production method of claim 24, which comprises reduction after hydrolysis.

29. A compound according to claim 1 or 3 substantially as herein described with reference to any one of Examples 1 to 19.

5

30. A method according to any one of claims 4, 10, 12, 14 15, 20, 22 or 24 substantially as herein described with reference to any one of Examples 1 to 19.

10 Dated this 7th day of November 2001

SUMIKA FINE CHEMICALS CO., LTD.

By its Patent Attorneys

GRIFFITH HACK

5
5
5
5
5
5
5
5