



(72) MCCARTHY, SEAN A., US

(71) MILLENNIUM PHARMACEUTICALS, INC., US

(51) Int.Cl.⁶ C07H 21/04, C12N 15/86, C12N 15/85, C07K 16/00,
C07K 14/435, C12N 1/21

(30) 1998/05/29 (09/087,121) US

(30) 1998/05/29 (09/086,892) US

(54) **NOUVELLES PROTEINES SECRETEES ET ASSOCIEES A UNE
MEMBRANE ET LEURS UTILISATIONS**

(54) **NOVEL SECRETED AND MEMBRANE-ASSOCIATED
PROTEINS AND USES THEREFOR**

(57) L'invention porte sur de nouveaux polypeptides, de nouvelles protéines et molécules d'acides nucléique, tous étant sécrétés et associés à une membrane (telles que des molécules BDSF et STMST). Outre les protéines isolées, pleine longueur, l'invention porte également sur des protéines de fusion isolées, des peptides antigéniques et des anticorps. L'invention porte notamment sur des molécules d'acide nucléique, des vecteurs d'expression de recombinaison contenant une molécule d'acide nucléique de l'invention, des cellules hôtes dans lesquelles ont été introduits les vecteurs d'expression, et des animaux transgéniques dans lesquels un gène BDSF ou STMST a été introduit ou dissocié. Des procédés de diagnostic, criblage et thérapeutiques utilisant les compositions de cette invention sont également décrits.

(57) Novel secreted and membrane-associated polypeptides, proteins, and nucleic acid molecules are disclosed (e.g., BDSF and STMST molecules). In addition to isolated, full-length proteins, the invention further provides isolated fusion proteins, antigenic peptides and antibodies. The invention also provides nucleic acid molecules, recombinant expression vectors containing a nucleic acid molecule of the invention, host cells into which the expression vectors have been introduced and non-human transgenic animals in which a BDSF or STMST gene has been introduced or disrupted. Diagnostic, screening and therapeutic methods utilizing compositions of the invention are also provided.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07H 21/04, C12N 15/85, 15/86, 1/21, C07K 14/435, 16/00, 39 /395, C07H 19 /34	A1	(11) International Publication Number: WO 99/61463 (43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/US99/11842 (22) International Filing Date: 28 May 1999 (28.05.99) (30) Priority Data: 09/086,892 29 May 1998 (29.05.98) US 09/087,121 29 May 1998 (29.05.98) US (71) Applicant: MILLENNIUM BIOTHERAPEUTICS, INC. [US/US]; 620 Memorial Drive, Cambridge, MA 02319 (US). (72) Inventor: MCCARTHY, Sean, A.; Apartment #4, 62 Com- monwealth Avenue, Boston, MA 02116 (US). (74) Agents: MANDRAGOURAS, Amy, E.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US) et al.		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: NOVEL SECRETED AND MEMBRANE-ASSOCIATED PROTEINS AND USES THEREFOR (57) Abstract Novel secreted and membrane-associated polypeptides, proteins, and nucleic acid molecules are disclosed (e.g., BDSF and STMST molecules). In addition to isolated, full-length proteins, the invention further provides isolated fusion proteins, antigenic peptides and antibodies. The invention also provides nucleic acid molecules, recombinant expression vectors containing a nucleic acid molecule of the invention, host cells into which the expression vectors have been introduced and non-human transgenic animals in which a BDSF or STMST gene has been introduced or disrupted. Diagnostic, screening and therapeutic methods utilizing compositions of the invention are also provided.		

NOVEL SECRETED AND MEMBRANE-ASSOCIATED PROTEINS AND USES THEREFOR

Background of the Invention

5 There is considerable medical interest in secreted and membrane-associated mammalian proteins. Many such proteins, for example, signaling factors and/or cell-surface receptors, are important in the regulation of growth and/or differentiation of cells of for triggering one or more specific cellular responses.

 Signaling factors play an important role in the development and functioning of
10 different cell types by allowing for communication between interacting cells. Such factors provide a signal between cells which can cause cells which recognize the signal to perform specialized tasks, such as cell growth, differentiation and/or proliferation.

 For example, cells of the immune system characteristically express a variety of signaling proteins which are crucial to proper functioning of the immune system. Such
15 proteins include secreted immunoglobulins and non-immunoglobulin molecules which interact with cellular adhesion molecules, as well as other selected target molecules. Many of these proteins are members of the immunoglobulin (Ig) superfamily of proteins, characterized by the existence of at least one immunoglobulin (Ig)-like domain. Such proteins function in a variety of immune cell functions ranging from immune cell
20 development and differentiation, antigen recognition, antibody production, cellular signal transduction, and cellular homing of immune responsive cells from the circulation to sites of increased antigen concentration.

 Cell surface receptors likewise play an important role in the development and functioning of different cell types by allowing for communication between interacting
25 cells or between a cell and a soluble ligand in the intracellular milieu. For example, the G protein-coupled receptors ("GPCRs") form one of the largest receptor superfamilies found in nature, and it is estimated that greater than 1000 different such receptors exist in mammals. Upon binding of extracellular ligands, GPCRs interact with a specific subset of heterotrimeric G-proteins that can then, in their activated forms, inhibit or
30 activate various effector enzymes and/or ion channels. The ligands for many of these receptors are known although there exists an ever-increasing number of GPCRs which

- 2 -

have been identified in the sequencing of the human genome for which no ligands have yet been identified. This latter subfamily of GPCRs is called the orphan family of GPCRs. In addition to both GPCRs with known ligands, as well as orphan GPCRs, there exist a family of GPCR-like molecules which share significant homology as well as many of the structural properties of the GPCR superfamily. For example, a family of GPCR-like proteins which arises from three alternatively-spliced forms of a gene occurring between the CD4 and triosephosphate isomerase genes at human chromosome 12p13, has been recently identified (including protein A-1, A-2, and A-3). Ansari-Lari *et al.* (1996) *Genome Res.* 6(4):314-326. Comparative sequence analysis of the syntenic region in mouse chromosome 6 has further revealed a murine homologue of at least the A-2 splice product. Ansari-Lari *et al.* (1998) *Genome Res.* 8(1):29-40.

The fundamental knowledge that GPCRs play a role in regulating that activity of virtually every cell in the human body has fostered an extensive search for modulators of such receptors for use as human therapeutics. In fact, the superfamily of GPCRs has proven to be among the most successful drug targets. Consequently, it has been recognized that the newly isolated orphan GPCRs, as well as the GPCR-like proteins, have great potential for drug discovery.

Given the importance of such secreted proteins (*e.g.*, signaling factors) and membrane-associated proteins (*e.g.*, G-protein coupled receptors) in the proper functioning of a variety of cellular processes, there exists a need to identify novel signaling factors and/or receptors as well as for modulators of such molecules for use in regulating a variety of cellular responses and for use in the design and development of new therapies. Moreover, with the identification of each new GPCR or GPCR-like protein, there exists a need for identifying the surrogate ligands for such molecules.

25

Summary of the Invention

The present invention is based, at least in part, on the discovery of novel secreted proteins and membrane-associated proteins. In one aspect, the present invention involves novel signaling molecules, referred to herein as Brain-Derived Signaling Factor ("BDSF") molecules, as well as the nucleic acids encoding them. The BDSF molecules of the present invention are useful as modulating agents in regulating a variety of

30

- 3 -

cellular processes. Accordingly, in one aspect, this invention provides isolated nucleic acid molecules encoding BDSF proteins or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of BDSF-encoding nucleic acids.

5 In one embodiment, a BDSF nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or complements thereof. In a preferred embodiment, the isolated nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:1,
10 SEQ ID NO:3, SEQ ID NO:4, or a complement thereof. In yet another preferred embodiment, an isolated nucleic acid molecule has the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a complement thereof. In another embodiment, the nucleic acid molecule further comprises nucleotides 244-701 of SEQ ID NO:1. In another embodiment, the nucleic
15 acid molecule further comprises nucleotides 31-487 of SEQ ID NO:4

In yet another embodiment, a BDSF nucleic acid molecule is 60% homologous to the nucleotide sequence shown in SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or a complement thereof. In a preferred embodiment, an isolated nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:6, SEQ ID NO: 8, SEQ ID NO:9, or a
20 complement thereof.

In another embodiment, a BDSF nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:5. In another embodiment, a BDSF nucleic acid molecule includes a nucleotide sequence encoding a protein having
25 an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:7 or SEQ ID NO:10. In a preferred embodiment, a BDSF nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:10. In another preferred embodiment, an isolated nucleic
30 acid molecule encodes the amino acid sequence of human BDSF. In yet another preferred embodiment, the nucleic acid molecule includes a nucleotide sequence

- 4 -

encoding a protein having the amino acid sequence of SEQ ID NO: 2 or SEQ ID NO:5. In another preferred embodiment, an isolated nucleic acid molecule encodes the amino acid sequence of murine BDSF. In yet another preferred embodiment, the nucleic acid molecule includes a nucleotide sequence encoding a protein having the amino acid
5 sequence of SEQ ID NO: 7 or SEQ ID NO:10.

In another embodiment, an isolated nucleic acid molecule of the present invention encodes a protein, preferably a BDSF protein, which includes an immunoglobulin-like domain. In another embodiment, an isolated nucleic acid molecule of the present invention encodes a protein, preferably a BDSF protein, which includes a
10 signal sequence, an immunoglobulin-like domain, and, preferably, is secreted. In yet another embodiment, a BDSF nucleic acid molecule encodes a BDSF protein and is a naturally occurring nucleotide sequence.

Another embodiment of the invention features nucleic acid molecules, preferably BDSF nucleic acid molecules, which specifically detect BDSF nucleic acid molecules
15 relative to nucleic acid molecules encoding non-BDSF proteins. For example, in one embodiment, such a nucleic acid molecule is at least 450, preferably 500-700, more preferably 700-900, more preferably 900-1100, and even more preferably 1100-1120 nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:6,
20 the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a complement thereof. Another embodiment of the invention provides an isolated nucleic acid molecule which is antisense to the coding strand of a BDSF nucleic acid.

Another aspect of the invention provides a vector comprising a BDSF nucleic
25 acid molecule. In certain embodiments, the vector is a recombinant expression vector. In another embodiment, the invention provides a host cell containing a vector of the invention. The invention also provides a method for producing a protein, preferably a BDSF protein, by culturing in a suitable medium, a host cell of the invention containing a recombinant expression vector such that the protein is produced.

- 5 -

Another aspect of this invention features isolated or recombinant BDSF proteins and polypeptides. In one embodiment, an isolated protein, preferably a BDSF protein, includes an immunoglobulin-like domain. In another embodiment, an isolated protein, preferably a BDSF protein, includes a signal sequence, an immunoglobulin-like domain, and is, preferably, secreted. In another embodiment, an isolated protein, preferably a BDSF protein, has an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:5. In another embodiment, an isolated protein, preferably a BDSF protein, has an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:7 or SEQ ID NO:10. In a preferred embodiment, a protein, preferably a BDSF protein, has an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:10. In another embodiment, the invention features fragments of the proteins having the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:5, wherein the fragment comprises at least 15 contiguous amino acids of the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with the ATCC as Accession No. 98756. In another embodiment, the invention features fragments of the proteins having the amino acid sequence of SEQ ID NO:7 or SEQ ID NO:10, wherein the fragment comprises at least 15 contiguous amino acids of the amino acid sequence of SEQ ID NO:7 or SEQ ID NO:10. In another embodiment, a protein, preferably a BDSF protein, has the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:10.

Another embodiment of the invention features an isolated protein, preferably a BDSF protein, which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, or a complement thereof. This invention further features an isolated protein, preferably a BDSF protein, which is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, or a complement thereof.

Yet another embodiment of the invention features an isolated protein, preferably a BDSF protein, which is encoded by a nucleic acid molecule having a nucleotide

- 6 -

sequence at least about 60% homologous to a nucleotide sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or a complement thereof. This invention further features an isolated protein, preferably a BDSF protein, which is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization
5 conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or a complement thereof.

In another aspect, the present invention involves proteins of a novel family of G protein-coupled receptor-like proteins, referred to herein as the Seven Transmembrane Signal Transducer ("STMST" family or "STMST proteins"), as well as the nucleic acids
10 encoding them. The STMST molecules of the present invention as well as STMST ligands and/or STMST modulators, are useful in regulating a variety of cellular processes. Accordingly, in one aspect, this invention provides isolated nucleic acid molecules encoding STMST proteins or biologically active portions thereof, as well as nucleic acid fragments suitable as primers or hybridization probes for the detection of
15 STMST-encoding nucleic acids.

In one embodiment, an STMST nucleic acid molecule is 75% homologous to the nucleotide sequence shown in SEQ ID NO:14, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or complement thereof. In another embodiment, an STMST nucleic acid molecule is 80% homologous
20 to the nucleotide sequence shown in SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof. In a preferred embodiment, an isolated STMST nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:16, or a complement thereof. In another embodiment, an STMST nucleic acid molecule further comprises nucleotides
25 1-403 of SEQ ID NO:1. In another embodiment, an STMST nucleic acid molecule further comprises nucleotides 1295-2915 of SEQ ID NO:14. In another preferred embodiment, an isolated STMST nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:1. In yet another preferred embodiment, an isolated nucleic acid molecule has the nucleotide sequence of the DNA insert of the plasmid deposited with
30 ATCC as Accession Number _____, or a complement thereof.

- 7 -

In another preferred embodiment, an isolated STMST nucleic acid molecule has the nucleotide sequence shown SEQ ID NO:19, or a complement thereof. In another embodiment, an STMST nucleic acid molecule further comprises nucleotides 1-333 of SEQ ID NO:17. In another embodiment, an STMST nucleic acid molecule further
5 comprises nucleotides 2161-4166 of SEQ ID NO:17. In another preferred embodiment, an isolated STMST nucleic acid molecule has the nucleotide sequence shown in SEQ ID NO:17. In yet another preferred embodiment, an isolated nucleic acid molecule has the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof.

10 In another embodiment, an STMST nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:15, the amino acid sequence of SEQ ID NO:18, an amino acid or an amino acid sequence encoded by the DNA insert of the plasmid deposited with the ATCC as Accession No. _____, or an amino acid or an amino acid
15 sequence encoded by the DNA insert of the plasmid deposited with the ATCC as Accession No. _____. In another preferred embodiment, an STMST nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 75% homologous to the amino acid sequence of SEQ ID NO:15 or an amino acid or an amino acid sequence encoded by the DNA insert of the plasmid
20 deposited with the ATCC as Accession No. _____. In yet another preferred embodiment, an STMST nucleic acid molecule includes a nucleotide sequence encoding a protein having an amino acid sequence at least 60% homologous to the amino acid sequence of SEQ ID NO:18 or an amino acid or an amino acid sequence encoded by the DNA insert of the plasmid deposited with the ATCC as Accession No. _____.

25 In another embodiment, an isolated nucleic acid molecule of the present invention encodes an STMST protein which includes at least one transmembrane domain. In another embodiment, an isolated nucleic acid molecule of the present invention encodes a protein which includes a 7 transmembrane receptor profile. In another embodiment, an isolated nucleic acid molecule of the present invention encodes
30 a protein which includes a spectrin α -chain motif. In yet another embodiment, an

- 8 -

STMST nucleic acid molecule encodes an STMST protein and is a naturally occurring nucleotide sequence.

Another embodiment of the invention features STMST nucleic acid molecules which specifically detect STMST nucleic acid molecules relative to nucleic acid
5 molecules encoding non-STMST proteins. For example, in one embodiment, an STMST nucleic acid molecule is at least 350 nucleotides in length and hybridizes under stringent conditions to a nucleic acid molecule comprising the nucleotide sequence shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the nucleotide sequence
10 of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof.

Another embodiment of the invention provides an isolated nucleic acid molecule which is antisense to the coding strand of an STMST nucleic acid.

Another aspect of the invention provides a vector comprising an STMST nucleic
15 acid molecule. In certain embodiments, the vector is a recombinant expression vector. In another embodiment, the invention provides a host cell containing a vector of the invention. The invention also provides a method for producing an STMST protein by culturing in a suitable medium, a host cell of the invention containing a recombinant expression vector such that an STMST protein is produced.

20 Another aspect of this invention features isolated or recombinant STMST proteins and polypeptides. In one embodiment, an isolated STMST protein includes at least one transmembrane domain. In another embodiment, an isolated STMST protein includes at least six transmembrane domains. In another embodiment, an isolated STMST protein includes seven transmembrane domains. In another embodiment, an
25 isolated STMST protein includes a 7 transmembrane receptor profile. In another embodiment, an isolated STMST protein includes a spectrin α -chain profile. In another embodiment, an isolated STMST protein has an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:15 or SEQ ID NO:18. In a preferred embodiment, an STMST protein has an amino acid sequence at least about
30 75% homologous to the amino acid sequence of SEQ ID NO:15. In another preferred embodiment, an STMST protein has an amino acid sequence at least about 60%

- 9 -

homologous to the amino acid sequence of SEQ ID NO:18. In another embodiment, an STMST protein has the amino acid sequence of SEQ ID NO:15 or SEQ ID NO:18.

Another embodiment of the invention features an isolated STMST protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 75% homologous to a nucleotide sequence of SEQ ID NO:14, or a complement thereof.

Another embodiment of the invention features an isolated STMST protein which is encoded by a nucleic acid molecule having a nucleotide sequence at least about 80% homologous to a nucleotide sequence of SEQ ID NO:17, or a complement thereof. This invention further features an isolated STMST protein which is encoded by a nucleic acid molecule having a nucleotide sequence which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:14, SEQ ID NO:17, or a complement thereof.

The proteins of the present invention, preferably BDSF or STMST proteins or biologically active portions thereof, can be operatively linked to a non-BDSF or non-STMST polypeptide to form fusion proteins. The invention further features antibodies that specifically bind BDSF or STMST proteins, such as monoclonal or polyclonal antibodies. In addition, the BDSF or STMST proteins or biologically active portions thereof can be incorporated into pharmaceutical compositions, which optionally include pharmaceutically acceptable carriers.

In another aspect, the present invention provides a method for detecting BDSF or STMST expression in a biological sample by contacting the biological sample with an agent capable of detecting a BDSF or STMST nucleic acid molecule, protein or polypeptide such that the presence of a BDSF or STMST nucleic acid molecule, protein or polypeptide is detected in the biological sample.

In another aspect, the present invention provides a method for detecting the presence of BDSF or STMST activity in a biological sample by contacting the biological sample with an agent capable of detecting an indicator of BDSF or STMST activity such that the presence of BDSF or STMST activity is detected in the biological sample.

- 10 -

In another aspect, the invention provides a method for modulating BDSF or STMST activity comprising contacting the cell with an agent that modulates BDSF or STMST activity such that BDSF or STMST activity in the cell is modulated. In one embodiment, the agent inhibits BDSF or STMST activity. In another embodiment, the agent stimulates BDSF or STMST activity. In one embodiment, the agent is an antibody that specifically binds to a BDSF or STMST protein. In another embodiment, the agent modulates expression of BDSF or STMST by modulating transcription of a BDSF or STMST gene or translation of a BDSF or STMST mRNA. In yet another embodiment, the agent is a nucleic acid molecule having a nucleotide sequence that is antisense to the coding strand of a BDSF or STMST mRNA or a BDSF or STMST gene.

In one embodiment, the methods of the present invention are used to treat a subject having a disorder characterized by aberrant BDSF or STMST protein or nucleic acid expression or activity by administering an agent which is a BDSF or STMST modulator to the subject. In one embodiment, the BDSF or STMST modulator is a BDSF or STMST protein, respectively. In another embodiment, the BDSF or STMST modulator is a BDSF or STMST nucleic acid molecule, respectively. In a preferred embodiment, the STMST modulator is an STMST ligand. In yet another embodiment, the BDSF or STMST modulator is a peptide, peptidomimetic, or other small molecule. In a preferred embodiment, the disorder characterized by aberrant BDSF protein or nucleic acid expression is a proliferative or differentiative disorder. In another preferred embodiment, the disorder characterized by aberrant STMST protein or nucleic acid expression is a developmental, differentiative, proliferative disorder, an inflammatory disorder, a respiratory disorder (*e.g.*, asthma), or cell death.

The present invention also provides a diagnostic assay for identifying the presence or absence of a genetic alteration characterized by at least one of (i) aberrant modification or mutation of a gene encoding a STMST protein; (ii) mis-regulation of said gene; and (iii) aberrant post-translational modification of a BDSF or STMST protein, wherein a wild-type form of said gene encodes an protein with a BDSF or STMST activity, respectively.

- 11 -

In another aspect the invention provides a method for identifying a compound that binds to or modulates the activity of a BDSF or STMST protein. In one embodiment, the invention provides a method for identifying a compound which binds to a BDSF or STMST protein which involves contacting the BDSF or STMST protein, or a cell expressing the BDSF or STMST protein with a test compound and determining whether the BDSF or STMST protein binds to the test compound. In another embodiment, the invention provides a method for identifying a compound which modulates the activity of a BDSF or STMST protein which involves contacting a BDSF or STMST protein with a test compound, and determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the BDSF or STMST protein.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1 depicts the cDNA sequence and predicted amino acid sequence of human BDSF-1. The nucleotide sequence corresponds to nucleic acids 1 to 1119 of SEQ ID NO:1. The amino acid sequence corresponds to amino acids 1 to 244 of SEQ ID NO:2.

Figure 2 depicts the cDNA sequence and predicted amino acid sequence of murine BDSF-1. The nucleotide sequence corresponds to nucleic acids 1 to 3196 of SEQ ID NO:6. The amino acid sequence corresponds to amino acids 1 to 251 of SEQ ID NO:7.

Figure 3 depicts an alignment of the amino acid sequence of human BDSF-1, corresponding to SEQ ID NO:2, with the amino acid sequence of murine BDSF-1, corresponding to SEQ ID NO:7. The immunoglobulin-like domains are underlined. The conserved cysteine residues of the immunoglobulin-like domain of human and murine BDSF are indicated with an asterix. The alignment was performed using the ALIGN program with a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4.

- 12 -

Figure 4 depicts the cDNA sequence and predicted amino acid sequence of human STMST-1. The nucleotide sequence corresponds to nucleic acids 1 to 2915 of SEQ ID NO:14. The amino acid sequence corresponds to amino acids 1 to 297 of SEQ ID NO:15.

5 *Figure 5* depicts the cDNA sequence and predicted amino acid sequence of human STMST-2. The nucleotide sequence corresponds to nucleic acids 1 to 4166 of SEQ ID NO:17. The amino acid sequence corresponds to amino acids 1 to 609 of SEQ ID NO:18.

10 *Figure 6* depicts an alignment of the amino acid sequences of human STMST-1 (SEQ ID NO:15), human STMST-2 (SEQ ID NO:18), human protein A-2 (Accession No. U47928, SEQ ID NO:29), and human protein A-3 (Accession No. U47929, SEQ ID NO:30). The 7 transmembrane receptor profile is indicated in italics. The transmembrane domains are underlined. The spectrin α -chain profile is indicated in bold.

15

Detailed Description of the Invention

The present invention is based on the discovery of novel molecules, referred to herein as BDSF protein and nucleic acid molecules, which comprise a family of molecules having certain conserved structural and functional features. The present invention is also based on the discovery of novel molecules, referred to herein as STMST protein and nucleic acid molecules, which comprise a second family of molecules having certain conserved structural and functional features.

20 The term "family" when referring to the protein and nucleic acid molecules of the invention is intended to mean two or more proteins or nucleic acid molecules having a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology as defined herein. Such family members can be naturally occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can contain homologues of non-human origin. Members of a family may also have common functional characteristics.

25

30

- 13 -

Members of a family may also share sufficient sequence homology with other members of the same family. For example, isolated proteins of the present invention, preferably BDSF proteins, have an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:7 or SEQ ID NO:10 or are encoded by a nucleotide sequence sufficiently homologous to SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:6 or SEQ ID NO:9. In another embodiment, isolated proteins of the present invention, preferably STMST proteins, have an amino acid sequence sufficiently homologous to the amino acid sequence of SEQ ID NO:15 or SEQ ID NO:18. As used herein, the term "sufficiently homologous" refers to a first amino acid or nucleotide sequence which contains a sufficient or minimum number of identical or equivalent (*e.g.*, an amino acid residue which has a similar side chain) amino acid residues or nucleotides to a second amino acid or nucleotide sequence such that the first and second amino acid or nucleotide sequences share common structural domains or motifs and/or a common functional activity. For example, amino acid or nucleotide sequences which share common structural domains have at least about 30-40% homology, preferably 40-50% homology, more preferably 50-60%, and even more preferably 60-70%, 70-80%, or 80-90% homology across the amino acid sequences of the domains and contain at least one and preferably two structural domains or motifs, are defined herein as sufficiently homologous. Furthermore, amino acid or nucleotide sequences which share at least 30-40%, preferably 40-50%, more preferably 50-60%, 60-70%, 70-80%, or 80-90% homology and share a common functional activity are defined herein as sufficiently homologous.

The BDSF Protein Family

For example, the BDSF proteins of the present invention belong to a family of signaling proteins having common structural and functional characteristics. In one embodiment, the isolated proteins of the present invention, preferably BDSF proteins, are proteins having an amino acid sequence of about 150-340 amino acid residues in length, preferably about 170-320, more preferably about 190-300, more preferably about 210-280, or about 230-260 amino acid residues in length. In one embodiment, an isolated protein of the present invention, preferably a BDSF protein, includes an

- 14 -

immunoglobulin (Ig)-like domain. As used herein, the term an "immunoglobulin-like domain" includes an amino acid sequence having about 65-115, preferably about 70-110, more preferably about 80-100 amino acid residues, and even more preferably at least about 85-95 amino acids in length and having a bit score for the alignment of the
 5 sequence to the Ig family Hidden Markov Model (HMM) of at least 10, preferably 10-15, more preferably 15-20, more preferably 20-25, even more preferably 25-35, 35-55, 55-100 or greater. The Ig family HMM has been assigned the PFAM Accession PF00047 (<http://genome.wustl.edu/Pfam/WWWdata/ig.html>).

To identify the presence of an Ig-like domain in a BDSF family member, the
 10 amino acid sequence of the family member is searched against a database of HMMs (e.g., the Pfam database, release 2.1) using the default parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). For example, the hmmsf program, which is available as part of the HMMER package of search programs, is a family specific default program for PF00047 having a score of 15 as the default
 15 threshold score for determining a hit. For example, a search using the amino acid sequence of SEQ ID NO:2 was performed against the HMM database resulting in the identification of an Ig-like domain in the amino acid sequence of SEQ ID NO:2 and a score of 22.43 against the Ig family HMM Accession PF00047. The results of the search are set forth below.

20

Score: 22.43 SEQ ID NO:2: aa41-129 HMM: aa1-47 (SEQ ID NO:13)

25

GqsVTLTCmVs.fhPpdYt.IwWY.rNgqpi.....
 41 GQNVEMSCAFQSGSASVYLEIQWWFLRGPEDLDPGAEGAGAQVELLPDR
tLtInsWqyEDsGtYwCmV
 90 DPDSGDKISTVKVQGNDISHKLQISKVRKKDEGLYECRV

- 15 -

In another example, a search was performed using the amino acid sequence of SEQ ID NO:7 against the HMM database resulting in the identification of an Ig-like domain in the amino acid sequence of SEQ ID NO:7 and a score of 22.43 against the Ig family HMM Accession PF00047. The results of the search are set forth below.

5

Score: 22.43 SEQ ID NO:2: aa40-128 HMM: aa1-47 (SEQ ID NO:13)

10

GqsVTLTCmVs.fhPpdYt.IwWY.rNgqpi.....
 40 GQNVEMSCAFQSGSASVYLEIQWWFLRGPEdleQGTEAAGSQVELLPDR
tLtInsWqyEDsGtYwCmV
 89 DPDNDGTKISTVKVGGNDISHKLQISKVRKKDEGLYECRV

Accordingly, in one embodiment of the invention, a BDSF protein is a human
 15 BDSF-1 protein having an Ig-like domain at about amino acids 41-129 of SEQ ID NO:2.
 Such an Ig-like domain has the amino acid sequence:

GQNVEMSCAFQSGSASVYLEIQWWFLRGPELDPGAEGAGAQVELLPDR
 DPDSGDKISTVKVQGGNDISHKLQISKVRKKDEGLYECRV (SEQ ID
 20 NO:11)

Accordingly, BDSF family members having at least 50-60% homology,
 preferably about 60-70%, more preferably about 70-80%, or about 80-90% homology
 with the Ig-like domain of human BDSF-1 (*e.g.*, SEQ ID NO:11) are within the scope of
 the invention.

25 In yet another embodiment of the invention, a BDSF protein is a murine BDSF-1
 protein having an Ig-like domain at about amino acids 40-128 of SEQ ID NO:7. Such
 an Ig-like domain has the amino acid sequence:

GQNVEMSCAFQSGSASVYLEIQWWFLRGPEdleQGTEAAGSQVELLPDR
 30 DPDNDGTKISTVKVGGNDISHKLQISKVRKKDEGLYECRV (SEQ ID
 NO:12)

- 16 -

Accordingly, a BDSF family member having at least 50-60% homology, preferably about 60-70%, more preferably about 70-80%, or about 80-90% homology with the Ig-like domain of murine BDSF-1 (*e.g.*, SEQ ID NO:12) is within the scope of the invention. Description of the Pfam database can be found in Sonhammer *et al.* (1997) *Proteins* 28(3)405-420, and description of HMMs can be found, for example, in Gribskov *et al.* (1990) *Meth. Enzymol.* 183:146-159; Gribskov *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh *et al.* (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz *et al.* (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated by reference.

10 An Ig-like domain further contains at least one, preferably two, cysteine residues which are conserved between BDSF molecules. Preferably, the Ig-like domain of a protein, preferably a BDSF protein, has cysteine residues which are located in the same or similar positions as cysteine residues in other BDSF protein family members. For example, when a BDSF protein of the invention is aligned with a BDSF family member
15 for purposes of comparison (see *e.g.*, Fig. 3) preferred cysteine residues of the invention are those in which cysteine residues in the amino acid sequence of BDSF are located in the same or similar position as the cysteine residues in other BDSF family members. As an illustrative embodiment, Fig. 3 shows cysteine residues located in the same or similar positions of the human BDSF protein (corresponding to SEQ ID NO:2) and murine
20 BDSF protein (corresponding to SEQ ID NO:7) in the following locations: amino acid number 48 of human BDSF and amino acid number 47 of murine BDSF; and amino acid number 127 of human BDSF and amino acid number 126 of murine BDSF.

In another embodiment of the invention, a BDSF protein has an Ig-like domain and a signal sequence. As used herein, a "signal sequence" refers to a peptide of about
25 20-30 amino acid residues in length which occurs at the N-terminus of secretory and integral membrane proteins and which contains a majority of hydrophobic amino acid residues. For example, a signal sequence contains at least about 15-45 amino acid residues, preferably about 20-40 amino acid residues, more preferably about 20-30 amino acid residues, and more preferably about 24-28 amino acid residues, and has at
30 least about 40-70%, preferably about 50-65%, and more preferably about 55-60% hydrophobic amino acid residues (*e.g.*, Alanine, Valine, Leucine, Isoleucine,

- 17 -

Phenylalanine, Tyrosine, Tryptophan, or Proline). Such a "signal sequence", also referred to in the art as a "signal peptide", serves to direct a protein containing such a sequence to a lipid bilayer. For example, in one embodiment, a BDSF protein contains a signal sequence of about amino acids 1-25 of SEQ ID NO:2, or a signal sequence of
5 about amino acids 1-24 of SEQ ID NO:7.

As used interchangeably herein, a "BDSF activity", "biological activity of BDSF" or "functional activity of BDSF", refers to an activity exerted by a BDSF protein, polypeptide or nucleic acid molecule as determined *in vivo*, or *in vitro*, according to standard techniques. In one embodiment, a BDSF activity is a direct
10 activity, such as an association with a BDSF-target molecule. As used herein, a "target molecule" is a molecule with which a BDSF protein binds or interacts in nature (*e.g.*, a BDSF receptor), such that BDSF-mediated function is achieved. A BDSF target molecule can be a BDSF protein or polypeptide of the present invention or a non-BDSF molecule. Alternatively, a BDSF activity is an indirect activity, such as an activity
15 mediated by interaction of the BDSF protein with a BDSF target molecule such that the target molecule modulates a downstream cellular activity (*e.g.*, interaction of an BDSF molecule with a BDSF target molecule can modulate the activity of that target molecule on an intracellular signaling pathway). In a preferred embodiment, a BDSF activity is at least one or more of the following activities: (i) interaction of a BDSF protein in the
20 extracellular milieu with a non-BDSF protein molecule on the surface of the same cell which secreted the BDSF protein molecule; (ii) interaction of a BDSF protein in the extracellular milieu with a non-BDSF protein molecule on the surface of a different cell from that which secreted the BDSF protein molecule; (iii) complex formation between a BDSF protein and a BDSF receptor; (iv) complex formation between a BDSF protein
25 and non-BDSF receptor; and (v) interaction of a BDSF protein with a second protein in the extracellular milieu. In yet another preferred embodiment, a BDSF activity is at least one or more of the following activities: (1) modulation of cellular signal transduction, either *in vitro* or *in vivo*; (2) modulation of protein:protein interactions, either *in vitro* or *in vivo*; (3) regulation of cellular proliferation; or (4) regulation of
30 cellular differentiation.

- 18 -

Accordingly, another embodiment of the invention features isolated BDSF proteins and polypeptides having a BDSF activity. Preferred proteins are BDSF proteins having an Ig-like domain, and, preferably, a BDSF activity. In another preferred embodiment, the isolated protein further comprises a signal sequence. In still another preferred embodiment, the isolated protein is a BDSF protein having an Ig-like domain, a BDSF activity, preferably an amino acid sequence sufficiently homologous to an amino acid sequence of SEQ ID NO:2 or SEQ ID NO:7, and optionally a signal sequence and/or propeptide.

The human BDSF-1 cDNA, which is approximately 1119 nucleotides in length, encodes a protein which is approximately 244 amino acid residues in length. The human BDSF-1 protein has an Ig-like domain. An Ig-like domain includes, for example, about amino acids 41-129 of SEQ ID NO:2. The Ig-like domain further contains at least about two conserved cysteine residues. Cysteine residues can be found at least about at amino acids 48 and 127 of SEQ ID NO:2. The human BDSF-1 protein is predicted to be a secreted protein which contains a signal sequence at about amino acids 1-25 of SEQ ID NO:2. The prediction of such a signal peptide can be made, for example, utilizing the computer algorithm SIGNALP (Henrik, *et al.* (1997) *Protein Engineering* 10:1-6).

The murine BDSF-1 cDNA, which is approximately 3196 nucleotides in length, encodes a protein which is approximately 251 amino acid residues in length. The murine BDSF-1 protein has an Ig-like domain. An Ig-like domain includes, for example, about amino acids 40-128 of SEQ ID NO:7. The Ig-like domain further contains at least about two conserved cysteine residues. Cysteine residues can be found at least about at amino acids 47 and 126 of SEQ ID NO:7. The murine BDSF-1 protein is predicted to be a secreted protein which contains a signal sequence at about amino acids 1-24 of SEQ ID NO:7.

Analysis of human BDSF expression demonstrated highest levels of BDSF mRNA in adult brain tissue, in particular, in the sub-regions of the brain including amygdala, caudate nucleus, hippocampus, substantia nigra, sub-thalamate nucleus and thalamus, but not in corpus callosum. Human BDSF maps to hu7p12-14 between markers WI-967 and WI-4253 close to the CMT2D (Charcot-Marie-Tooth neuropathy) locus. The syntenic chromosome in mouse, mo11, is in close proximity with the mouse

- 19 -

known genes: egfr (epidermal growth factor receptor), ddc (dopa decarboxylase) and cobl (cordon bleu)). BDSF also maps close to the following human genes:

ADCYAP1R1 (adenylate cyclase activating polypeptide 1), AMPH (amphiphysin), BLVRA (biliverdin reductase A), OGDH (oxoglutarate dehydrogenase), OCM
 5 (oncomodulin) and EGFR (epidermal growth factor receptor).

In situ analysis of adult mouse brain demonstrated high punctate expression in the cortex, hypothalamus, hippocampus (including, but not restricted to, granule cells) and hind brain. Expression in the cerebellum is limited to purkinje cells. Embryonic sagittal sections (day 14.5 - 15.5) showed expression in developing brain and spinal cord
 10 (*e.g.*, in the ependyma of the brain and the primordium of the lower incisor tooth).

Accordingly, it is postulated that regulation and/or modulation of BDSF (*e.g.*, using BDSF nucleic acid molecules, polypeptides, antibodies and/or BDSF modulators) can play an important role in the following: (1) regulation of neuronal proliferation and/or differentiation; (2) modulation of neuronal signaling; (3) regulation of
 15 neurodegeneration (*e.g.*, modulation of apoptotic degeneration and/or neuron atrophy); and (4) regulation of neurotoxicity. Moreover, BDSF molecules and/or modulators can provide novel therapeutic approaches for treatment of disorders and/or diseases including (1) neurodegenerative diseases (*e.g.*, Alzheimer's disease, dementias related to Alzheimer's disease (such as Pick's disease), Huntington's Disease, Parkinson's and
 20 other Lewy diffuse body diseases, multiple sclerosis, amyotrophic lateral sclerosis, progressive supranuclear palsy, epilepsy, Jakob-Creutzfeldt disease, or AIDS related dementia; (2) peripheral neuropathies and/or demyelinopathies; and (3) nervous system-related disorders and/or diseases including cognitive disorders, *e.g.*, memory and learning disorders, such as amnesia, apraxia, agnosia, amnesic dysnomia, amnesic
 25 spatial disorientation, Kluver-Bucy syndrome, Alzheimer's related memory loss (Eglen R.M. (1996) *Pharmacol. and Toxicol.* 78(2):59-68; Perry E.K. (1995) *Brain and Cognition* 28(3):240-58) and learning disability; disorders affecting consciousness, *e.g.*, visual hallucinations, perceptual disturbances, or delerium associated with Lewy body dementia; schitzo-effective disorders (Dean B. (1996) *Mol. Psychiatry* 1(1):54-8),
 30 schizophrenia with mood swings (Bymaster F.P. (1997) *J. Clin. Psychiatry* 58 (suppl.10):28-36; Yeomans J.S. (1995) *Neuropharmacol.* 12(1):3-16; Reimann D.

- 20 -

(1994) *J. Psychiatric Res.* 28(3):195-210), depressive illness (primary or secondary); affective disorders (Janowsky D.S. (1994) *Am. J. Med. Genetics* 54(4):335-44); sleep disorders (Kimura F. (1997) *J. Neurophysiol.* 77(2):709-16), *e.g.*, REM sleep abnormalities in patients suffering from, for example, depression (Riemann D. (1994) *J. Psychosomatic Res.* 38 Suppl. 1:15-25; Bourgin P. (1995) *Neuroreport* 6(3): 532-6), paradoxical sleep abnormalities (Sakai K. (1997) *Eur. J. Neuroscience* 9(3):415-23), sleep-wakefulness, and body temperature or respiratory depression abnormalities during sleep (Shuman S.L. (1995) *Am. J. Physiol.* 269(2 Pt 2):R308-17; Mallick B.N. (1997) *Brain Res.* 750(1-2):311-7). Other examples of nervous system related disorders include disorders affecting pain generation mechanisms, *e.g.*, pain related to irritable bowel syndrome (Mitch C.H. (1997) *J. Med. Chem.* 40(4):538-46; Shannon H.E. (1997) *J. Pharmac. and Exp. Therapeutics* 281(2):884-94; Bouaziz H. (1995) *Anesthesia and Analgesia* 80(6):1140-4; or Guimaraes A.P. (1994) *Brain Res.* 647(2):220-30) or chest pain; movement disorders (Monassi C.R. (1997) *Physiol. and Behav.* 62(1):53-9), *e.g.*, Parkinson's disease related movement disorders (Finn M. (1997) *Pharmacol. Biochem. & Behavior* 57(1-2):243-9; Mayorga A.J. (1997) *Pharmacol. Biochem. & Behavior* 56(2):273-9); eating disorders, *e.g.*, insulin hypersecretion related obesity (Maccario M. (1997) *J. Endocrinol. Invest.* 20(1):8-12; Premawardhana L.D. (1994) *Clin. Endocrinol.* 40(5): 617-21); or drinking disorders, *e.g.*, diabetic polydipsia (Murzi E. (1997) *Brain Res.* 752(1-2):184-8; Yang X. (1994) *Pharmacol. Biochem. & Behavior* 49(1):1-6).

The GPCR Family

The family of G protein-coupled receptors (GPCRs), to which the STMST proteins of the present invention bear significant homology, comprise an N-terminal extracellular domain, seven transmembrane domains (also referred to as membrane-spanning domains), three extracellular domains (also referred to as extracellular loops), three cytoplasmic domains (also referred to as cytoplasmic loops), and a C-terminal cytoplasmic domain (also referred to as a cytoplasmic tail). Members of the GPCR family also share certain conserved amino acid residues, some of which have been determined to be critical to receptor function and/or G protein signaling. For example, GPCRs contain the following features: a conserved asparagine residue in the first

2345678901234567890123456789012

WO 99/61463

PCT/US99/11842

- 22 -

TM2

* *

	6.	138	VVYMLHLATADVLFVSVLPFKISYYFSG	165
5	19.	73	NYILLNLAVADLFMVGGFTSTLYTSLH	100
	21.	61	NYFLLSLACADLIIGTFSMNLYTTYLLM	88
	30.	75	DVYLLNLALADLLFALTLPPIWAASKVNG	102
	40.	145	NFFIVSLAVADLTVALLVLPFNVAYSIL	172

10

22222222222222222222222222222222
44444444444555555555566666666666
0123456789012345678901234567

15 TM3

*

	6.	176	RFVTAIFYCNMYASILLMTVISIDR	200
	19.	111	NLEGFFATLGGEIALWSLVLAIER	135
	21.	99	DLWLALDYVASNASVMNLLISFDR	123
20	30.	111	KVVSLLEKVNIFYSGILLACISVDR	135
	40.	183	KLWLTCDVLCCTSSILNLCAIALDR	207

25

33333333333333333333333333333333
2222333333333334444444444445
6789012345678901234567890

TM4

* *

	6.	215	TLGRASFTCLAIWALAIAGVVPLVLKE	241
30	19.	149	GENHAIMGVAFTWVMALACAAPPLAGW	175
	21.	138	TPRRAALMIGLAWLVSFVLWAPAILFW	164
	30.	149	KRHLVKFVCLGCWGLSMNLSLPFFLFR	175
	40.	222	TVGRVLLLLISGVWLLSLLISSPPLIGW	248

35

44444444444444444444444444444444
3344444444444555555555566666
890123456789012345678901234

TM5

* * *

6.	268	YYYFSAFSAVFFFVPLIISTVCYVSIIRC	296	
5	19.	201	ESFVIYMFVVHFTIPMIIFFCYGQLVFT	229
	21.	186	PIITFGTAMAAFYLPVTVMCTLYWRIYRE	214
	30.	200	MVLRILPHTFGFIVPLFVMLFCYGFTLRT	228
	40.	267	RGYVIYSSLGSGFFIPLAIMTIVYIEIFVA	295

1

10

55555555555555555555555555555555
33334444444444445555555555666666
67890123456789012345678901234

TM6

15

* * *

6.	313	FLSAAVFCIFIICFGPTNVLLIAHYSFL	340	
19.	252	RMVIIMVIAFLICWVPYASVAFYIFTHQ	279	
21.	365	RTLSAILLAFILTWTPTYNIMVLVSTFCK	397	
30.	242	RVIFAVVLIFLLCWLPLYNLVLLADTLMR	269	
20	40.	529	RTLGIIMGVFVICWLPFFLMYVILPFCQ	556

1

66666666666666666666666666666666
333334444444444455555555556666
5678901234567890123456789012

25

TM7

* * *

6.	347	EAAYFAYLLCVCVSSISSCIDPLIYYYASSECO	379	
19.	282	NFGPIFMTIPAFFAKSAAIYNPVIYIMMNKQFR	314	
30	21.	394	CVPETLWELGYWLCYVNSTVNPMCYALCNKAFR	426
30.	281	NNIGRALDATEILGFLHSCLNPPIIYAFIGQNFR	313	
40.	559	CPTNKFKNFITWLGYINSGLNPVIYTI FNLDYR	591	

1

```
777777777777777777777777777777777777  
35 233333333334444444444455555555566  
901234567890123456789012345678901
```

- 24 -

The amino acid sequences of thrombin receptor (Accession No. P25116), rhodopsin receptor (Accession No. P08100), mACh receptor (Accession No. P08482), IL-8A receptor (Accession No. P25024), octopamine receptor (Accession No. P22270), are set forth as SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, and
 5 SEQ ID NO:145, respectively. Accordingly, GPCR-like proteins such as the STMST proteins of the present invention contain a significant number of structural characteristics of the GPCR family. For instance, the STMSTs of the present invention contain conserved cysteines found in the first 2 extracellular loops (prior to the third and fifth transmembrane domains) of most GPCRs (cys 83 and cys 161 of SEQ ID NO:15 or
 10 SEQ ID NO:18). A highly conserved asparagine residue in the first transmembrane domain is present (asn25 in SEQ ID NO:15 or SEQ ID NO:18). Transmembrane domain two of the STMST proteins contains a highly conserved leucine (leu49 of SEQ ID NO:15 or SEQ ID NO:18). The two cysteine residues are believed to form a disulfide bond that stabilizes the functional protein structure. A highly conserved
 15 tryptophan and proline in the fourth transmembrane domain of the STMST proteins is present (trp135 and pro 145 of SEQ ID NO:15 or SEQ ID NO:18). The third cytoplasmic loop contains 49 amino acid residues and is thus the longest cytoplasmic loop of the three, characteristic of G protein coupled receptors. Moreover, a highly conserved proline in the sixth transmembrane domain is present (pro260 of SEQ ID
 20 NO:15 and SEQ ID NO:18). The proline residues in the fourth, fifth, sixth, and seventh transmembrane domains are thought to introduce kinks in the alpha-helices and may be important in the formation of the ligand binding pocket. Furthermore, the conserved (in the second cytoplasmic loop) HRM motif found in almost all Rhodopsin family GPCRs is present in the STMST proteins of the instant invention (his107, arg108, met109 of
 25 SEQ ID NO:15 or SEQ ID NO:18). (The arginine of the HRM sequence is thought to be the most important amino acid in GPCRs and is invariant). Moreover, an almost invariant proline is present in the seventh transmembrane domain of STMST-2 (pro294 of SEQ ID NO:18).

In one embodiment, the STMST proteins of the present invention are proteins
 30 having an amino acid sequence of about 150-450, preferably about 200-400, more preferably about 225-375, more preferably about 250-350, or about 275-325 amino acids

- 25 -

in length. In another embodiment, the STMST proteins of the present invention are proteins having an amino acid sequence of about 450-750, preferably about 500-700, more preferably about 525-675, even more preferably about 550-650, and even more preferably about 575-625 amino acid residues in length. In one embodiment, the

5 STMST proteins of the present invention contain at least one transmembrane domain. As used herein, the term "transmembrane domain" includes an amino acid sequence having at least about 10, preferably about 13, preferably about 16, more preferably about 19, 21, 23, 25, 30, 35 or 40 amino acid residues, of which at least about 50-60%, 60-70%, preferably about 70-80% more preferably about 80-90%, or about 90-95% of the

10 amino acid residues contain non-polar side chains, for example, alanine, valine, leucine, isoleucine, proline, phenylalanine, tryptophan, and methionine. A transmembrane domain is lipophilic in nature. For example, a transmembrane domain can be found at about amino acids 11-34 of SEQ ID NO:15 or SEQ ID NO:18. In a preferred embodiment, an STMST protein of the present invention has more than one

15 transmembrane domain, preferably 2, 3, 4, 5, 6, or 7 transmembrane domains. For example, transmembrane domains can be found at about amino acids 11-34, 44-67, 85-106, 127-149, 172-196, and 244-262 of SEQ ID NO:15 as well as at 11-34, 44-67, 85-106, 127-149, 172-196, 245-269, and 277-300 of SEQ ID NO:18. In a particularly preferred embodiment, an STMST protein of the present invention has 7 transmembrane

20 domains.

In another embodiment, an STMST family member is identified based on the presence of at least one cytoplasmic loop, also referred to herein as a cytoplasmic domain. In another embodiment, an STMST family member is identified based on the presence of at least one extracellular loop. As defined herein, the term "loop" includes

25 an amino acid sequence having a length of at least about 4, preferably about 5-10, preferably about 10-20, and more preferably about 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, 90-100, or 100-150 amino acid residues, and has an amino acid sequence that connects two transmembrane domains within a protein or polypeptide.

Accordingly, the N-terminal amino acid of a loop is adjacent to a C-terminal amino acid

30 of a transmembrane domain in a naturally-occurring GPCR or GPCR-like molecule,

- 26 -

and the C-terminal amino acid of a loop is adjacent to an N-terminal amino acid of a transmembrane domain in a naturally-occurring GPCR or GPCR-like molecule.

As used herein, a "cytoplasmic loop" includes an amino acid sequence located within a cell or within the cytoplasm of a cell. For example, a cytoplasmic loop is found
5 at about amino acids 35-43, 107-126, and 197-243 of SEQ ID NO:15, or alternatively, at about amino acid residues 35-43, 107-126, and 197-244 of SEQ ID NO:18. Also as used herein, an "extracellular loop" includes an amino acid sequence located outside of a cell, or extracellularly. For example, an extracellular loop can be found at about amino acid residues 68-84 and 150-171 of SEQ ID NO:15, or alternatively, at about amino acid
10 residues 86-84, 150-171, or 270-276 of SEQ ID NO:18.

In another embodiment of the invention, an STMST family member is identified based on the presence of a "C-terminal cytoplasmic domain", also referred to herein as a C-terminal cytoplasmic tail, in the sequence of the protein. As used herein, a "C-terminal cytoplasmic domain" includes an amino acid sequence having a length of at
15 least about 10, preferably about 10-25, more preferably about 25-50, more preferably about 50-75, even more preferably about 75-100, 100-150, 150-200, 200-250, 250-300, 300-400, 400-500, or 500-600 amino acid residues and is located within a cell or within the cytoplasm of a cell. Accordingly, the N-terminal amino acid residue of a "C-terminal cytoplasmic domain" is adjacent to a C-terminal amino acid residue of a
20 transmembrane domain in a naturally-occurring GPCR or GPCR-like protein. For example, a C-terminal cytoplasmic domain is found at about amino acid residues 301-609 of SEQ ID NO:18.

In another embodiment, an STMST family member is identified based on the presence of an "N-terminal extracellular domain", also referred to herein as an N-
25 terminal extracellular loop in the amino acid sequence of the protein. As used herein, an "N-terminal extracellular domain" includes an amino acid sequence having about 1-500, preferably about 1-400, more preferably about 1-300, more preferably about 1-200, even more preferably about 1-100, and even more preferably about 1-50, 1-25, or 1-10 amino acid residues in length and is located outside of a cell or extracellularly. The C-terminal
30 amino acid residue of a "N-terminal extracellular domain" is adjacent to an N-terminal amino acid residue of a transmembrane domain in a naturally-occurring GPCR or

- 27 -

GPCR-like protein. For example, an N-terminal cytoplasmic domain is found at about amino acid residues 1-10 of SEQ ID NO:15 or SEQ ID NO:18.

Accordingly in one embodiment of the invention, an STMST family member includes at least one, preferably 6 or 7, transmembrane domains and and/or at least one
5 cytoplasmic loop, and/or at least one extracellular loop. In another embodiment, the STMST family member further includes an N-terminal extracellular domain and/or a C-terminal cytoplasmic domain. In another embodiment, the STMST family member can include six transmembrane domains, three cytoplasmic loops, and two extracellular loops, or can include six transmembrane domains, three extracellular loops, and 2
10 cytoplasmic loops. The former embodiment can further include an N-terminal extracellular domain. The latter embodiment can further include a C-terminal cytoplasmic domain. In another embodiment, the STMST family member can include seven transmembrane domains, three cytoplasmic loops, and three extracellular loops and can further include an N-terminal extracellular domain or a C-terminal cytoplasmic
15 domain.

In another embodiment, an STMST family member is identified based on the presence of at least one "7 transmembrane receptor profile", also referred to as a "Rhodopsin family sequence profile", in the protein or corresponding nucleic acid molecule. As used herein, the term "7 transmembrane receptor profile" includes an
20 amino acid sequence having at least about 100-400, preferably about 150-350, more preferably about 200-300 amino acid residues, or at least about 250-275 amino acids in length and having a bit score for the alignment of the sequence to the 7tm_1 family Hidden Markov Model (HMM) of at least 20, preferably 20-30, more preferably 30-40, more preferably 40-50, 50-75, 75-100, 100-200 or greater. The 7tm_1 family HMM has
25 been assigned the PFAM Accession PF00001
(http://genome.wustl.edu/Pfam/WWWdata/7tm_1.html).

To identify the presence of a 7 transmembrane receptor profile in an STMST family member, the amino acid sequence of the protein family member is searched against a database of HMMs (*e.g.*, the Pfam database, release 2.1) using the default
30 parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). For example, the hmmsf program, which is available as part of the HMMER package of search programs,

- 28 -

is a family specific default program for PF00001 and score of 15 is the default threshold score for determining a hit. For example, a search using the amino acid sequence of SEQ ID NO:15 was performed against the HMM database resulting in the identification of a 7 TM receptor profile in the amino acid sequence of SEQ ID NO:15. The results of
 5 the search are set forth below.

Score:44.14 SEQ ID NO:15: aa24-191 HMM: aa1-174

GNiLVIWvIcRyRRMRTPMNYFIvNLAvADLLFs1ft.MPFWMvYyvMq
 10 24 ANAWGILSVGAKQKKWKPLEFLLCTLAATHMLN-VAVPIATYSVVQLRR

gRWpFGdfMcrIWmYFDYMNMYASIFfLTcISIDRYLWAICHPMrYmR
 72 QRPDFEWNEGLCKVFVSTFYTLTLATCFSVTSLSYHRMWMVCWVPVNYRL

WMTpRHRAWvMIiiIWvMSFlISMPPFLMFrWstyrDEneWNmTWCmIyD
 15 121 SNAKK-QAVHTVMGIWMVSFILSALPA-VG-W-HDTSERFYTHG-CRFIV

WPewMWrWYvILmtiimgFYIPMiIMlF
 166 AEIGLGFGVCFLLLV-GGSVA-MGVICT
 20

Likewise, a search using the amino acid sequence of SEQ ID NO:18 results in an identical hit with a score of 44.14 against the 7tm_1 family HMM Accession PF00001. Accordingly, in one embodiment of the invention, an STMST protein is a human STMST-1 or a human STMST-2 protein having a 7 transmembrane receptor profile at
 25 about amino acids 24-191 of SEQ ID NO:15 or SEQ ID NO:18, respectively. Such a 7 transmembrane receptor profile has the amino acid sequence:

ANAWGILSVGAKQKKWKPLEFLLCTLAATHMLNVAVPIATYSVVQLRRQR
 PDFEWNEGLCKVFVSTFYTLTLATCFSVTSLSYHRMWMVCWVPVNYRLSNA
 KKQAVHTVMGIWMVSFILSALPAVGWHDTSERFYTHGCRFIVAEIGLGFG
 30 VCFLLLVGGSVAMGVICT. (SEQ ID NO:22)

- 29 -

Accordingly, STMST family members having at least 50-60% homology, preferably about 60-70%, more preferably about 70-80%, or about 80-90% homology with the 7 transmembrane receptor profile of human STMST-1 or STMST-2 (*e.g.*, SEQ ID NO:22) are within the scope of the invention.

5 In another embodiment, an STMST family member is identified based on the presence of a "spectrin α -chain profile" in the protein or corresponding nucleic acid molecule. As used herein, the term "spectrin α -chain profile" includes a protein domain having an amino acid sequence of about 50-250, preferably about 75-225, more preferably about 100-200 amino acid residues, or about 125-175 amino acids and having
10 a bit score for the alignment of the sequence to the spectrin family (HMM) of at least 7, preferably 8-10, more preferably 10-30, more preferably 30-50, even more preferably 50-75, 75-100, 100-200 or greater. The spectrin family HMM has been assigned the PFAM Accession PF00435 (<http://genome.wustl.edu/Pfam/WWWdata/spectrin.html>).

To identify the presence of a spectrin alpha chain profile in a STMST family
15 member, make the determination that a protein of interest has a particular profile, the amino acid sequence of the protein is searched against a database of HMMs (*e.g.*, the Pfam database, release 2.1) using the default parameters (http://www.sanger.ac.uk/Software/Pfam/HMM_search). For example, the hmmsf program, which is available as part of the HMMER package of search programs, is a
20 family specific default program for PF00435 and a score of 15 is the default threshold score for determining a hit. Alternatively, the threshold score for determining a hit can be lowered (*e.g.*, to 8 bits). A description of the Pfam database can be found in Sonhammer *et al.* (1997) *Proteins* 28(3)405-420 and a detailed description of HMMs can be found, for example, in Gribskov *et al.* (1990) *Meth. Enzymol.* 183:146-159;
25 Gribskov *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4355-4358; Krogh *et al.* (1994) *J. Mol. Biol.* 235:1501-1531; and Stultz *et al.* (1993) *Protein Sci.* 2:305-314, the contents of which are incorporated herein by reference. A search was performed against the HMM database resulting in the identification of a spectrin alpha chain profile in the amino acid sequence of SEQ ID NO:15. The results of the search are set forth below.

WO 99/61463

PCT/US99/11842

- 30 -

Score:8.78 SEQ ID NO:15: aa266-372 HMM: aa1-106

IqeRMnElndRWerLkelMeqRRQMledSmrlQQFfRDmDEeEsWInEK
 266 FSSLRADASAPWMALCVLWCSVAQALLLPVFLWACDRYRADLKAVREKC
 5
 EqilnSDDYGkDLtsVQnLlkKHQaFEaDIaAHE.dRIqalnefaqqLIq
 315 MALMANDEESDDETSLEGGISPDLVLERSLDYGYGGDFVALDRMAKYEIS
 enHYasEe
 10 365 ALEGGLPQ

All amino acids are described using universal single letter abbreviations according to these motifs.

Accordingly, in one embodiment, an STMST protein is human STMST-2 protein which includes a spectrin α -chain profile at about amino acids 266-372 of SEQ ID NO:18. Such a spectrin α -chain profile has the amino acid sequence:

FSSLRADASAPWMALCVLWCSVAQALLLPVFLWACDRYRADLKAV
 REKCMALMANDEESDDETSLEGGISPDLVLERSLDYGYGGDFVAL
 DRMAKYEISALEGGLPQ (SEQ ID NO:23).

Accordingly, STMST family members having at least 50-60% homology, preferably about 60-70%, more preferably about 70-80%, or about 80-90% homology with a spectrin α -chain profile of human STMST-2 (e.g., SEQ ID NO:23) are within the scope of the invention.

In another embodiment, an STMST protein includes at least a spectrin α -chain profile. In another embodiment, an STMST protein includes a spectrin α -chain profile and a 7 transmembrane receptor profile. In another embodiment, an STMST protein is human STMST-2 which includes a spectrin α -chain profile having about amino acids 266-372 of SEQ ID NO:18. In yet another embodiment, an STMST protein is human STMST-2 which includes a 7 transmembrane receptor profile having about amino acids 24-191 of SEQ ID NO:18 and a spectrin α -chain profile having about amino acids 266-372 of SEQ ID NO:18.

- 31 -

As used interchangeably herein, an "STMST activity", "biological activity of STMST" or "functional activity of STMST", refers to an activity exerted by an STMST protein, polypeptide or nucleic acid molecule on an STMST responsive cell as determined *in vivo*, or *in vitro*, according to standard techniques. In one embodiment, an STMST activity is a direct activity, such as an association with an STMST-target molecule. As used herein, a "target molecule" or "binding partner" is a molecule with which an STMST protein binds or interacts in nature, such that STMST-mediated function is achieved. An STMST target molecule can be a non-STMST molecule or an STMST protein or polypeptide of the present invention. In an exemplary embodiment, an STMST target molecule is an STMST ligand. Alternatively, an STMST activity is an indirect activity, such as a cellular signaling activity mediated by interaction of the STMST protein with an STMST ligand.

In a preferred embodiment, an STMST activity is at least one or more of the following activities: (i) interaction of an STMST protein with soluble STMST ligand; (ii) interaction of an STMST protein with a membrane-bound non-STMST protein; (iii) interaction of an STMST protein with an intracellular protein (e.g., an intracellular enzyme or signal transduction molecule); and (iv) indirect interaction of an STMST protein with an intracellular protein (e.g., a downstream signal transduction molecule).

In yet another preferred embodiment, an STMST activity is at least one or more of the following activities: (1) modulation of cellular signal transduction, either *in vitro* or *in vivo*; (2) regulation of gene transcription in a cell expressing an STMST protein; (3) regulation of gene transcription in a cell expressing an STMST protein, wherein said cell is involved inflammation; (4) regulation of cellular proliferation; (5) regulation of cellular differentiation; (6) regulation of development; (7) regulation of cell death; (8) regulation of regulation of inflammation; (9) regulation of respiratory cell function (e.g., asthma); (10) regulation of actin binding; (11) regulation of cytoskeletal attachment; and (12) regulation of chemotaxis, trafficking and/or migration.

Accordingly, another embodiment of the invention features isolated STMST proteins and polypeptides having an STMST activity. Preferred STMST proteins have at least one transmembrane domain and an STMST activity. In a preferred embodiment, an STMST protein has a 7 transmembrane receptor profile and an STMST activity. In another preferred embodiment, an STMST protein has a spectrin α -chain profile and an

- 32 -

STMST activity. In still another preferred embodiment, an STMST protein has a 7 transmembrane receptor profile, a spectrin α -chain profile, and STMST activity. In still another preferred embodiment, an STMST protein has a 7 transmembrane receptor profile, a spectrin α -chain profile, an STMST activity, and an amino acid sequence
5 sufficiently homologous to an amino acid sequence of SEQ ID NO:15, SEQ ID NO:18, SEQ ID NO:21, or SEQ ID NO:24.

The human STMST-1 cDNA, which is approximately 2915 nucleotides in length, encodes a protein which is approximately 297 amino acid residues in length. The human STMST-1 protein contains 6 transmembrane domains at about amino acids
10 11-34, 44-67, 85-106, 127-149, 172-196, and 244-262 of SEQ ID NO:15. The human STMST-1 protein further contains a 7 transmembrane receptor profile. The 7 transmembrane receptor profile can be found at least, for example, from about amino acids 24-191 of SEQ ID NO:15.

The human STMST-2 cDNA, which is approximately 4166 nucleotides in
15 length, encodes approximately 609 amino acid residues of the human STMST-1 protein. The human STMST-2 protein contains 7 transmembrane domains at about amino acids 11-34, 44-67, 85-106, 127-149, 172-196, 245-269, and 277-300 of SEQ ID NO:18. The human STMST-2 protein further contains a 7 transmembrane receptor profile. The 7 transmembrane receptor profile can be found at least, for example, from about amino
20 acids 24-191 of SEQ ID NO:18. Moreover, the human STMST protein contains a spectrin α -chain profile from about amino acids 266-372 of SEQ ID NO:5.

Various aspects of the invention are described in further detail in the following subsections:

25 **I. Isolated Nucleic Acid Molecules**

One aspect of the invention pertains to isolated nucleic acid molecules that encode BDSF proteins or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes to identify BDSF-encoding nucleic acids (*e.g.*, BDSF mRNA) and fragments for use as PCR primers for the amplification or
30 mutation of BDSF nucleic acid molecules. Another aspect of the invention pertains to

- 33 -

isolated nucleic acid molecules that encode STMST proteins or biologically active portions thereof, as well as nucleic acid fragments sufficient for use as hybridization probes to identify STMST-encoding nucleic acids (*e.g.*, STMST mRNA) and fragments for use as PCR primers for the amplification or mutation of STMST nucleic acid
5 molecules.

As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (*e.g.*, cDNA or genomic DNA) and RNA molecules (*e.g.*, mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded
10 DNA.

An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the
15 organism from which the nucleic acid is derived. For example, in various embodiments, the isolated BDSF nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free
20 of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, preferably a BDSF or STMST nucleic acid molecule, or a portion thereof, can be isolated using standard molecular
25 biology techniques and the sequence information provided herein. Using all or portion of the nucleic acid sequence of a BDSF or STMST nucleic acid molecule of the present invention, as a hybridization probe, BDSF or STMST nucleic acid molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*.
30 *2nd, ed.*, Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

- 34 -

Moreover, a nucleic acid molecule encompassing all or a portion of a BDSF or STMST nucleic acid molecule, can be isolated by the polymerase chain reaction (PCR) using synthetic oligonucleotide primers designed based upon the sequence of the BDSF or STMST nucleic acid molecules described herein.

5 A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to BDSF or STMST nucleotide sequences
10 can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

A. BDSF Nucleic Acid Molecules

In a preferred embodiment, an isolated nucleic acid molecule of the invention
15 comprises the nucleotide sequence shown in SEQ ID NO:1. The sequence of SEQ ID NO:1 corresponds to the human BDSF cDNA. This cDNA comprises sequences encoding the human BDSF protein (*i.e.*, "the coding region", from nucleotides 140-871), as well as 5' untranslated sequences (nucleotides 1-139) and 3' untranslated sequences (nucleotides 872-1119). Alternatively, the nucleic acid molecule can comprise only the
20 coding region of SEQ ID NO:1 (*e.g.*, nucleotides 140-871, corresponding to SEQ ID NO:3).

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:6. The sequence of SEQ ID NO:6 corresponds to the murine BDSF cDNA. This cDNA comprises
25 sequences encoding the murine BDSF protein (*i.e.*, "the coding region", from nucleotides 268-1020), as well as 5' untranslated sequences (nucleotides 1-267) and 3' untranslated sequences (nucleotides 1021-3196). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:6 (*e.g.*, nucleotides 268-1020, corresponding to SEQ ID NO:8).

- 35 -

In yet another preferred embodiment, an isolated nucleic acid molecule of the invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a portion of any of these nucleotide sequences. A nucleic acid molecule which is complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, is one which is sufficiently complementary to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, such that it can hybridize to the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, thereby forming a stable duplex.

In still another preferred embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 30-35%, preferably about 35-40%, more preferably at least about 40-45%, more preferably at least about 45-50%, and even more preferably at least about 50-55%, 55-60%, 60-65%, 65-70%, 70-75%, 75-80%, 80-85%, 85-90%, or 90-95% or more homologous to the nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a portion of any of these nucleotide sequences.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of a BDSF protein. The nucleotide sequence determined from the cloning of the

- 36 -

BDSF genes allows for the generation of probes and primers designed for use in identifying and/or cloning other BDSF family members, as well as BDSF homologues from other species. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide
5 sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, of an anti-sense sequence of SEQ ID NO:1, SEQ
10 ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or of a naturally occurring mutant of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756. In an
15 exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide sequence which is about 450, preferably 450-750, more preferably 750-950, more preferably 950-1100, and even more preferably 1100-1150 nucleotides in length and hybridizes under stringent hybridization conditions to a nucleic acid molecule of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID
20 NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756.

Probes based on the BDSF nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred
embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label
25 group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress a BDSF protein, such as by measuring a level of a BDSF-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting BDSF mRNA levels or determining whether a genomic BDSF gene has been mutated or deleted.

- 37 -

A nucleic acid fragment encoding a "biologically active portion of a BDSF protein" can be prepared by isolating a portion of the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
5 Accession Number 98756, which encodes a polypeptide having a BDSF biological activity (the biological activities of the BDSF proteins have previously been described), expressing the encoded portion of the BDSF protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the BDSF protein.

The invention further encompasses nucleic acid molecules that differ from the
10 nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, due to degeneracy of the genetic code and thus encode the same BDSF proteins as those encoded by the nucleotide sequence shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID
15 NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:7.

20 In addition to the BDSF nucleotide sequences shown in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the BDSF proteins
25 may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the BDSF genes may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding a BDSF protein (or STMST protein), preferably a mammalian BDSF protein (or STMST protein). Such
30 natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of a BDSF gene (or STMST gene). Any and all such nucleotide variations and resulting

- 38 -

amino acid polymorphisms in BDSF genes (and STMST genes) that are the result of natural allelic variation and that do not alter the functional activity of a BDSF protein (or STMST protein, respectively) are intended to be within the scope of the invention.

Moreover, nucleic acid molecules encoding other BDSF family members, and
5 thus which have a nucleotide sequence which differs from the BDSF sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, are intended to be within the scope of the invention. For example, a BDSF cDNA can be identified based on the nucleotide sequence of human
10 BDSF or the nucleotide sequence of murine BDSF. Moreover, nucleic acid molecules encoding BDSF proteins from different species, and thus which have a nucleotide sequence which differs from the human BDSF sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or which differs from the murine
15 BDSF sequences of SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:9 are intended to be within the scope of the invention. For example, a rat BDSF cDNA can be identified based on the nucleotide sequence of a human or murine BDSF.

Nucleic acid molecules corresponding to natural allelic variants and homologues of the BDSF cDNAs of the invention can be isolated based on their homology to the
20 BDSF nucleic acids disclosed herein using the cDNAs disclosed herein, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions.

Accordingly, in another embodiment, an isolated nucleic acid molecule of the invention is at least 15 nucleotides in length and hybridizes under stringent conditions to
25 the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756. In other embodiment, the nucleic acid is at least 30, 50, 100, 250 or 500 nucleotides in length. As used herein, the term "hybridizes under stringent conditions"
30 is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 60% homologous to each other typically remain hybridized to each

- 39 -

other. Preferably, the conditions are such that sequences at least about 70%, more preferably at least about 80%, even more preferably at least about 85% or 90% homologous to each other typically remain hybridized to each other. Such stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. A preferred, non-limiting example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2 X SSC, 0.1% SDS at 50-65°C. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:1 or SEQ ID NO:6 corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In addition to naturally-occurring allelic variants of the BDSF sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, thereby leading to changes in the amino acid sequence of the encoded BDSF proteins, without altering the functional ability of the BDSF proteins. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of BDSF (*e.g.*, the sequence of SEQ ID NO:2 or SEQ ID NO:7) without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the BDSF proteins of the present invention, are predicted to be particularly unamenable to alteration (*e.g.*, the conserved cysteines set forth in Figure 3). Moreover, amino acid residues that are defined by the Ig-like domain profile sequence are particularly unamenable to alteration.

- 40 -

Furthermore, additional amino acid residues that are conserved between the BDSF proteins of the present invention and other BDSF family members.

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding BDSF proteins that contain changes in amino acid residues that are not
5 essential for activity. Such BDSF proteins differ in amino acid sequence from SEQ ID NO:2 or SEQ ID NO:7 yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:7. Preferably, the protein encoded by the
10 nucleic acid molecule is at least about 65-70% homologous to SEQ ID NO:2 or SEQ ID NO:7, more preferably at least about 75-80% homologous to SEQ ID NO:2 or SEQ ID NO:7, even more preferably at least about 85-90% homologous to SEQ ID NO:2 or SEQ ID NO:7, and most preferably at least about 95% homologous to SEQ ID NO:2 or SEQ ID NO:7.

15 An isolated nucleic acid molecule encoding a BDSF protein homologous to the protein of SEQ ID NO:2 or SEQ ID NO:7 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
20 Accession Number 98756, such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, by standard techniques, such as site-directed
25 mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. For example, sequence comparisons of the human BDSF-1 and murine BDSF-1 protein are 90.5% identical over the first 211 amino acids of the protein which includes the Ig-like domain. This region, therefore, suggests a highly conserved function and thus, indicates
30 an "essential region" of human BDSF-1 and murine BDSF-1. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid

- 41 -

residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (*e.g.*, lysine, arginine, histidine), acidic side chains (*e.g.*, aspartic acid, glutamic acid), uncharged polar side chains (*e.g.*, glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (*e.g.*, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (*e.g.*, threonine, valine, isoleucine) and aromatic side chains (*e.g.*, tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in a BDSF protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of a BDSF coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for BDSF biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

In a preferred embodiment, a mutant BDSF protein can be assayed for the ability to (1) modulate cellular signal transduction; (2) modulate protein:protein interactions; (3) regulate cellular proliferation; or (4) regulate cellular differentiation.

B. STMST Nucleic Acid Molecules

A nucleic acid molecule of the present invention, *e.g.*, a nucleic acid molecule having the nucleotide sequence of SEQ ID NO:14, the nucleotide sequence of SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. Using all or portion of the nucleic acid sequence of SEQ ID NO:14, the nucleotide sequence of SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence

- 42 -

of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ as a hybridization probe, STMST nucleic acid molecules can be isolated using standard hybridization and cloning techniques (*e.g.*, as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989*).

Moreover, a nucleic acid molecule encompassing all or a portion of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ can be isolated by the polymerase chain reaction (PCR) using synthetic oligonucleotide primers designed based upon the sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

A nucleic acid of the invention can be amplified using cDNA, mRNA or alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to STMST nucleotide sequences can be prepared by standard synthetic techniques, *e.g.*, using an automated DNA synthesizer.

In a preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:14. The sequence of SEQ ID NO:14 corresponds to the human STMST-1 cDNA. This cDNA comprises sequences encoding the human STMST-1 protein (*i.e.*, "the coding region", from nucleotides 404-1294), as well as 5' untranslated sequences (nucleotides 1-403) and 3' untranslated sequences (nucleotides 1295-2915). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:14 (*e.g.*, nucleotides 404-1294, corresponding to SEQ ID NO:16).

- 43 -

In another preferred embodiment, an isolated nucleic acid molecule of the invention comprises the nucleotide sequence shown in SEQ ID NO:17. The sequence of SEQ ID NO:14 corresponds to the human STMST-2 cDNA. This cDNA comprises sequences encoding the human STMST-2 protein (*i.e.*, "the coding region", from
5 nucleotides 334-2160), as well as 5' untranslated sequences (nucleotides 1-333) and 3' untranslated sequences (nucleotides 2161-4166). Alternatively, the nucleic acid molecule can comprise only the coding region of SEQ ID NO:14 (*e.g.*, nucleotides 334-2160, corresponding to SEQ ID NO:19).

In another preferred embodiment, an isolated nucleic acid molecule of the
10 invention comprises a nucleic acid molecule which is a complement of the nucleotide sequence shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences. A nucleic
15 acid molecule which is complementary to the nucleotide sequence shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ is one which is sufficiently complementary to the nucleotide sequence shown in SEQ ID
20 NO:14, SEQ ID NO:17, or such that it can hybridize to the nucleotide sequence shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby forming a stable duplex.

25 In still another preferred embodiment, an isolated nucleic acid molecule of the present invention comprises a nucleotide sequence which is at least about 60-65%, preferably at least about 70-75%, more preferable at least about 80-85%, and even more preferably at least about 90-95% or more homologous to the nucleotide sequences shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of
30 the plasmid deposited with ATCC as Accession Number _____, or the nucleotide

- 44 -

sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a portion of any of these nucleotide sequences.

Moreover, the nucleic acid molecule of the invention can comprise only a portion of the nucleic acid sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide
5 sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of an STMST protein. The nucleotide sequence determined from the cloning of the STMST-1 genes
10 allows for the generation of probes and primers designed for use in identifying and/or cloning other STMST family members, as well as STMST homologues from other species. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, preferably about 25, more preferably
15 about 40, 50 or 75 consecutive nucleotides of a sense sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, of an anti-sense sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the
20 plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or of a naturally occurring mutant of SEQ ID NO:14 or SEQ ID NO:17. In an exemplary embodiment, a nucleic acid molecule of the present invention comprises a nucleotide sequence which is greater than 350, 351-450, 451-550, 551-650, 651-750, or 751-850,
25 851-950, 951-1050, 1051-1150, or 1151-1250 nucleotides in length and hybridizes under stringent hybridization conditions to a nucleic acid molecule of SEQ ID NO:14 or SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

- 45 -

Probes based on the STMST nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, *e.g.*, the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which misexpress an STMST protein, such as by measuring a level of an STMST-encoding nucleic acid in a sample of cells from a subject *e.g.*, detecting STMST mRNA levels or determining whether a genomic STMST gene has been mutated or deleted.

10 A nucleic acid fragment encoding a "biologically active portion of an STMST protein" can be prepared by isolating a portion of the nucleotide sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, which
15 encodes a polypeptide having an STMST biological activity (the biological activities of the STMST proteins have previously been described), expressing the encoded portion of the STMST protein (*e.g.*, by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the STMST protein.

The invention further encompasses nucleic acid molecules that differ from the
20 nucleotide sequence shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, due to degeneracy of the genetic code and thus encode the same STMST proteins as those encoded by the nucleotide sequence shown in SEQ ID
25 NO:14 or SEQ ID NO:17. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID NO:15 or SE ID NO:5.

In addition to the STMST nucleotide sequences shown in SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with
30 ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, it will be appreciated by

- 46 -

those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the STMST proteins may exist within a population (*e.g.*, the human population). Such genetic polymorphism in the STMST genes may exist among individuals within a population due to natural allelic variation.

5 Moreover, nucleic acid molecules encoding other STMST family members and thus which have a nucleotide sequence which differs from the STMST-1 sequences of SEQ ID NO:14 or SEQ ID NO:17 are intended to be within the scope of the invention. For example, an STMST-3 cDNA can be identified based on the nucleotide sequence of human STMST-1 or STMST-2. Moreover, nucleic acid molecules encoding STMST
10 proteins from different species, and thus which have a nucleotide sequence which differs from the STMST sequences of SEQ ID NO:14 or SEQ ID NO:17 are intended to be within the scope of the invention. For example, an mouse STMST cDNA can be identified based on the nucleotide sequence of a human STMST.

 Nucleic acid molecules corresponding to natural allelic variants and homologues
15 of the STMST cDNAs of the invention can be isolated based on their homology to the STMST nucleic acids disclosed herein using the cDNAs disclosed herein, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions.

 Accordingly, in another embodiment, an isolated nucleic acid molecule of the
20 invention is at least 15 nucleotides in length and hybridizes under stringent conditions to the nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. In another embodiment,
25 the nucleic acid is at least 30, 50, 100, 250 or 500 nucleotides in length. Preferably, an isolated nucleic acid molecule of the invention that hybridizes under stringent conditions to the sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as
30 Accession Number _____, corresponds to a naturally-occurring nucleic acid molecule. As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA

- 47 -

molecule having a nucleotide sequence that occurs in nature (*e.g.*, encodes a natural protein).

In addition to naturally-occurring allelic variants of the STMST sequences that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequences of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, thereby leading to changes in the amino acid sequence of the encoded STMST proteins, without altering the functional ability of the STMST proteins. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in the sequence of SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____. A "non-essential" amino acid residue is a residue that can be altered from the wild-type sequence of STMST (*e.g.*, the sequence of SEQ ID NO:15) without altering the biological activity, whereas an "essential" amino acid residue is required for biological activity. For example, amino acid residues that are conserved among the STMST proteins of the present invention, are predicted to be particularly unamenable to alteration. Moreover, amino acid residues that are defined by the 7 transmembrane signature profile and the spectrin α -chain, repeated domain signature profile are particularly unamenable to alteration. Furthermore, additional amino acid residues that are conserved between the STMST proteins of the present invention and members of the G protein coupled receptor protein family are not likely to be amenable to alteration (*e.g.*, the conserved asn residue within the first TM domain, asn25 of SEQ ID NO:15 or SEQ ID NO:18; the conserved cys in the first extracellular loop, cys83 of SEQ ID NO:15 or SEQ ID NO:18; the conserved arg at the interface of the third TM domain and the first cytoplasmic loop, arg108 of SEQ ID NO:2 or SEQ ID NO:18; the conserved trp and pro in the fourth TM domain, trp135 and pro145 of SEQ ID NO:15 or SEQ ID NO:18; the conserved cys residue in the second extracellular domain, cys161 of SEQ ID NO:15 or SEQ ID NO:18; the conserved phe residue in the fifth TM domain, phe251 of

- 48 -

SEQ ID NO:15 or SEQ ID NO:18; or the conserved pro in the seventh TM domain, pro294 of SEQ ID NO:18).

Accordingly, another aspect of the invention pertains to nucleic acid molecules encoding STMST proteins that contain changes in amino acid residues that are not essential for activity. Such STMST proteins differ in amino acid sequence from SEQ ID NO:15 and SEQ ID NO:18 yet retain biological activity. In one embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:15. Preferably, the protein encoded by the nucleic acid molecule is at least about 75-80% homologous to SEQ ID NO:15, more preferably at least about 80-85% homologous to SEQ ID NO:15, even more preferably at least about 85-90% homologous to SEQ ID NO:15, and even more preferably at least about 90-95% homologous to SEQ ID NO:15. In another embodiment, the isolated nucleic acid molecule comprises a nucleotide sequence encoding a protein, wherein the protein comprises an amino acid sequence at least about 65% homologous to the amino acid sequence of SEQ ID NO:18. Preferably, the protein encoded by the nucleic acid is at least about 65-70% homologous to SEQ ID NO:15, more preferably at least about 70-75% homologous to SEQ ID NO:18, even more preferably at least about 75-80%, and even more preferably at least about 80-85%, 85-90%, or 90-95% homologous to SEQ ID NO:18.

An isolated nucleic acid molecule encoding an STMST protein homologous to the protein of SEQ ID NO:15 or SEQ ID NO:18 can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence of SEQ ID NO:14, SEQ ID NO:17, or such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced into SEQ ID NO:14, SEQ ID NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____, by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. Thus, a predicted

- 49 -

nonessential amino acid residue in an STMST protein is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an STMST coding sequence, such as by saturation mutagenesis, and the resultant mutants can be
 5 screened for STMST biological activity to identify mutants that retain activity. Following mutagenesis of SEQ ID NO:14, SEQ ID NO:17, or the encoded protein can be expressed recombinantly and the activity of the protein can be determined.

In a preferred embodiment, a mutant STMST protein can be assayed for the ability to (1) modulation of cellular signal transduction, either *in vitro* or *in vivo*; (2)
 10 regulation of gene transcription in a cell expressing an STMST protein; (3) regulation of gene transcription in a cell expressing an STMST protein, wherein said cell is involved inflammation; (4) regulation of cellular proliferation; (5) regulation of cellular differentiation; (6) regulation of development; (7) regulation of cell death; (8) regulation of inflammation; and (9) regulation of respiratory cell function.

15

C. Antisense Nucleic Acid Molecules

In addition to the nucleic acid molecules encoding BDSF and STMST proteins described above, another aspect of the invention pertains to isolated nucleic acid molecules which are antisense thereto. An "antisense" nucleic acid comprises a
 20 nucleotide sequence which is complementary to a "sense" nucleic acid encoding a protein, *e.g.*, complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid can hydrogen bond to a sense nucleic acid. The antisense nucleic acid can be complementary to an entire BDSF or STMST coding strand, or to only a portion thereof.
 25 In one embodiment, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence encoding a BDSF or STMST protein. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues (*e.g.*, the coding region of human BDSF corresponds to SEQ ID NO:3, the coding region of murine BDSF corresponds to SEQ
 30 ID NO:8, the coding region of human STMST-1 corresponds to SEQ ID NO:16 and the coding region of human STMST-2 corresponds to SEQ ID NO:19). In another

- 50 -

embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding BDSF. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into amino acids (*i.e.*, also referred to as 5' and 3' untranslated regions).

5 Given the coding strand sequences encoding BDSF and STMST proteins disclosed herein (*e.g.*, SEQ ID NO:3, SEQ ID NO:8, SEQ ID NO:16 and SEQ ID NO:19), antisense nucleic acids of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of BDSF or STMST mRNA, but more
10 preferably is an oligonucleotide which is antisense to only a portion of the coding or noncoding region of BDSF or STMST mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of BDSF or STMST mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid of
15 the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid (*e.g.*, an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex
20 formed between the antisense and sense nucleic acids, *e.g.*, phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-
25 carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-
30 N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-

- 51 -

methylyluracil, uracil-5- oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (*i.e.*, RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a subject or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a BDSF or STMST protein to thereby inhibit expression of the protein, *e.g.*, by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. An example of a route of administration of antisense nucleic acid molecules of the invention include direct injection at a tissue site. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then administered systemically. For example, for systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, *e.g.*, by linking the antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. The antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong pol II or pol III promoter are preferred.

In yet another embodiment, the antisense nucleic acid molecule of the invention is an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gaultier *et al.* (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methylribonucleotide (Inoue *et al.* (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue *et al.* (1987) *FEBS Lett.* 215:327-330).

- 52 -

In still another embodiment, an antisense nucleic acid of the invention is a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (*e.g.*, hammerhead ribozymes described in Haselhoff and Gerlach (1988) *Nature* 334:585-591) can be used to catalytically cleave BDSF or STMST mRNA transcripts to thereby inhibit translation of BDSF or STMST mRNA, respectively. A ribozyme having specificity for a BDSF-encoding nucleic or STMST-encoding nucleic acid can be designed based upon the nucleotide sequence of a BDSF cDNA or STMST mRNA disclosed herein. For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a BDSF-encoding mRNA or STMST-encoding mRNA. See, *e.g.*, Cech *et al.* U.S. Patent No. 4,987,071; and Cech *et al.* U.S. Patent No. 5,116,742. Alternatively, BDSF mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, *e.g.*, Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

Alternatively, BDSF or STMST gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of the BDSF or STMST genes (*e.g.*, promoter and/or enhancer sequences) to form triple helical structures that prevent transcription of the BDSF or STMST genes in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. *et al.* (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J. (1992) *Bioassays* 14(12):807-15.

D. Peptide Nucleic Acids

In yet another embodiment, the nucleic acid molecules of the present invention can be modified at the base moiety, sugar moiety or phosphate backbone to improve, *e.g.*, the stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of the nucleic acid molecules can be modified to generate peptide nucleic acids (see Hyrup B. *et al.* (1996) *Bioorganic & Medicinal Chemistry* 4 (1): 5-23). As used herein, the terms "peptide nucleic acids" or "PNAs" refer to nucleic acid mimics, *e.g.*, DNA mimics, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases

- 53 -

are retained. The neutral backbone of PNAs has been shown to allow for specific hybridization to DNA and RNA under conditions of low ionic strength. The synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described in Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *et al.* PNAS 93:
5 14670-675.

PNAs of BDSF or STMST nucleic acid molecules can be used in therapeutic and diagnostic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNAs of BDSF or STMST
10 nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (*e.g.*, by PNA-directed PCR clamping); as 'artificial restriction enzymes' when used in combination with other enzymes, (*e.g.*, S1 nucleases (Hyrup B. (1996) *supra*)); or as probes or primers for DNA sequencing or hybridization (Hyrup B. *et al.* (1996) *supra*; Perry-O'Keefe *supra*).

15 In another embodiment, PNAs can be modified, (*e.g.*, to enhance their stability or cellular uptake), by attaching lipophilic or other helper groups to PNA, by the formation of PNA-DNA chimeras, or by the use of liposomes or other techniques of drug delivery known in the art. For example, PNA-DNA chimeras can be generated which may combine the advantageous properties of PNA and DNA. Such chimeras
20 allow DNA recognition enzymes, (*e.g.*, RNase H and DNA polymerases), to interact with the DNA portion while the PNA portion would provide high binding affinity and specificity. PNA-DNA chimeras can be linked using linkers of appropriate lengths selected in terms of base stacking, number of bonds between the nucleobases, and orientation (Hyrup B. (1996) *supra*). The synthesis of PNA-DNA chimeras can be
25 performed as described in Hyrup B. (1996) *supra* and Finn P.J. *et al.* (1996) *Nucleic Acids Res.* 24 (17): 3357-63. For example, a DNA chain can be synthesized on a solid support using standard phosphoramidite coupling chemistry and modified nucleoside analogs, *e.g.*, 5'-(4-methoxytrityl)amino-5'-deoxy-thymidine phosphoramidite, can be used as a between the PNA and the 5' end of DNA (Mag, M. *et al.* (1989) *Nucleic Acid*
30 *Res.* 17: 5973-88). PNA monomers are then coupled in a stepwise manner to produce a chimeric molecule with a 5' PNA segment and a 3' DNA segment (Finn P.J. *et al.* (1996)

- 54 -

supra). Alternatively, chimeric molecules can be synthesized with a 5' DNA segment and a 3' PNA segment (Peterser, K.H. *et al.* (1975) *Bioorganic Med. Chem. Lett.* 5: 1119-11124).

In other embodiments, the oligonucleotide may include other appended groups such as peptides (*e.g.*, for targeting host cell receptors *in vivo*), or agents facilitating transport across the cell membrane (see, *e.g.*, Letsinger *et al.* (1989) *Proc. Natl. Acad. Sci. US.* 86:6553-6556; Lemaitre *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:648-652; PCT Publication No. W088/09810, published December 15, 1988) or the blood-brain barrier (see, *e.g.*, PCT Publication No. W089/10134, published April 25, 1988). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, *e.g.*, Krol *et al.* (1988) *BioTechniques* 6:958-976) or intercalating agents. (See, *e.g.*, Zon (1988) *Pharm. Res.* 5:539-549). To this end, the oligonucleotide may be conjugated to another molecule, (*e.g.*, a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

15

E. STMST Ligands and Modulators

Furthermore, given the fact that an important use for the STMST molecules of the present invention is in the screening for STMST ligands (*e.g.*, surrogate ligands) and/or STMST modulators, it is intended that the following are also within the scope of the present invention: isolated nucleic acids which encode and STMST ligands or STMST modulators, probes and/or primers useful for identifying STMST ligands or STMST modulators based on the sequences of nucleic acids which encode and STMST ligands or STMST modulators, isolated nucleic acid molecules which are complementary or antisense to the sequences of nucleic acids which encode and STMST ligands or STMST modulators, isolated nucleic acid molecules which are at least about 60-65%, preferably at least about 70-75%, more preferable at least about 80-85%, and even more preferably at least about 90-95% or more homologous to the sequences of nucleic acids which encode and STMST ligands or STMST modulators, portions of nucleic acids which encode and STMST ligands or STMST modulators (*e.g.*, biologically-active portions), naturally-occurring allelic variants of nucleic acids which encode and STMST ligands or STMST modulators, nucleic acid molecules which

30

- 55 -

hybridize under stringent hybridization conditions to nucleic acids which encode and STMST ligands or STMST modulators, functionally-active mutants of nucleic acids which encode and STMST ligands or STMST modulators, PNAs of nucleic acids which encode and STMST ligands or STMST modulators, as well as vectors containing a
5 nucleic acid encoding an STMST ligand or STMST modulator, described herein, host cells into which an expression vector encoding an STMST ligand or STMST modulator has been introduced, and homologous recombinant animal which express STMST ligands or STMST modulators.

10 **II. Isolated BDSF and STMST Proteins, Fragments Thereof and Antibodies Thereeto**

One aspect of the invention pertains to isolated BDSF proteins, and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise anti-BDSF antibodies. Another aspect of the invention pertains to isolated
15 STMST proteins, and biologically active portions thereof, as well as polypeptide fragments suitable for use as immunogens to raise anti-STMST antibodies. In one embodiment, native proteins (*e.g.*, BDSF or STMST proteins) can be isolated from cells or tissue sources by an appropriate purification scheme using standard protein purification techniques. In another embodiment, proteins (*e.g.*, BDSF or STMST
20 proteins) are produced by recombinant DNA techniques. Alternative to recombinant expression, a BDSF or STMST protein or polypeptide can be synthesized chemically using standard peptide synthesis techniques.

An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or
25 tissue source from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language "substantially free of cellular material" includes preparations of protein (*e.g.*, BDSF or STMST proteins) in which the protein is separated from cellular components of the cells from which it is isolated or recombinantly produced. In one embodiment, the language
30 "substantially free of cellular material" includes preparations of protein having less than about 30% (by dry weight) of non-BDSF protein or non-STMST protein (also referred

- 56 -

to herein as a "contaminating protein"), more preferably less than about 20% of non-BDSF protein or non-STMST protein, still more preferably less than about 10% of non-BDSF protein or non-STMST protein, and most preferably less than about 5% non-BDSF protein or non-STMST protein. When the protein or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, *i.e.*, culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

The language "substantially free of chemical precursors or other chemicals" includes preparations of protein (*e.g.*, BDSF or STMST proteins) in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one embodiment, the language "substantially free of chemical precursors or other chemicals" includes preparations of protein having less than about 30% (by dry weight) of chemical precursors or non-BDSF chemicals (or non-BDSF chemicals), more preferably less than about 20% chemical precursors or non-BDSF chemicals (or non-BDSF chemicals), still more preferably less than about 10% chemical precursors or non-BDSF chemicals (or non-BDSF chemicals), and most preferably less than about 5% chemical precursors or non-BDSF chemicals (or non-BDSF chemicals).

20

A. BDSF Proteins and Biologically-Active Portions Thereof

In a preferred embodiment, the BDSF protein has an amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:7. In other embodiments, the BDSF protein is substantially homologous to SEQ ID NO:2 or SEQ ID NO:7, and retains the functional activity of the protein of SEQ ID NO:2 or SEQ ID NO:7, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the BDSF protein is a protein which comprises an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:7 and retains the functional activity of the BDSF proteins of SEQ ID NO:2 or SEQ ID NO:7. Preferably, the protein is at least about 35-40% homologous to SEQ ID NO:2 or SEQ ID NO:7, more preferably at least

- 57 -

about 40-45% homologous to SEQ ID NO:2 or SEQ ID NO:7, even more preferably at least about 45-50% homologous to SEQ ID NO:2 or SEQ ID NO:7, and even more preferably at least about 50-55%, 55-60%, 60-65%, 65-70%, 70-75%, 75-80%, 80-85%, 85-90%, or 90-95% or more homologous to SEQ ID NO:2 or SEQ ID NO:7.

5 To determine the percent homology of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence and non-homologous sequences can be disregarded for comparison purposes). In a preferred embodiment, the length of a
10 reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, even more preferably at least 60%, and even more preferably at least 70%, 80%, or 90% of the length of the reference sequence (*e.g.*, when aligning a second sequence to the BDSF amino acid sequence of SEQ ID NO:2 or SEQ
ID NO:7 having 293 amino acid residues, at least 88, preferably at least 117, more
15 preferably at least 147, even more preferably at least 176, and even more preferably at least 205, 234 or 264 amino acid residues are aligned). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid
residue or nucleotide as the corresponding position in the second sequence, then the
20 molecules are homologous at that position (*i.e.*, as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The percent homology between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100).

25 The comparison of sequences and determination of percent homology between two sequences can be accomplished using a mathematical algorithm. A preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264-68, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-77. Such
30 an algorithm is incorporated into the NBLAST and XBLAST programs (version 2.0) of Altschul, *et al.* (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be

- 58 -

performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to BDSF nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to BDSF protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) Nucleic Acids Research 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Another preferred, non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

Biologically active portions of a BDSF protein include peptides comprising amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the BDSF protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:7, which include less amino acids than the full length BDSF proteins, and exhibit at least one activity of a BDSF protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the BDSF protein. A biologically active portion of a BDSF protein can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

In one embodiment, a biologically active portion of a BDSF protein comprises an Ig-like domain. In another embodiment, a biologically active portion of a BDSF protein comprises at least an Ig-like domain and a signal sequence.

It is to be understood that a preferred biologically active portion of a BDSF protein of the present invention may contain at least one of the above-identified structural domains. A more preferred biologically active portion of a BDSF protein may contain at least two of the above-identified structural domains. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be

- 59 -

prepared by recombinant techniques and evaluated for one or more of the functional activities of a native BDSF protein.

B. STMST Proteins and Biologically-Active Portions Thereof

5 In a preferred embodiment, the STMST protein has an amino acid sequence shown in SEQ ID NO:15. In other embodiments, the STMST protein is substantially homologous to SEQ ID NO:15, and retains the functional activity of the protein of SEQ ID NO:15, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another
10 embodiment, the STMST protein is a protein which comprises an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:15 and retains the functional activity of the STMST proteins of SEQ ID NO:15, respectively. Preferably, the protein is at least about 75-80%% homologous to SEQ ID NO:15, more preferably at least about 80-85% homologous to SEQ ID NO:15, even more preferably
15 at least about 85-90% homologous to SEQ ID NO:15, and most preferably at least about 90-95% or more homologous to SEQ ID NO:15.

 In a preferred embodiment, the STMST protein has an amino acid sequence shown in SEQ ID NO:18. In other embodiments, the STMST protein is substantially homologous to SEQ ID NO:18, and retains the functional activity of the protein of SEQ
20 ID NO:18, yet differs in amino acid sequence due to natural allelic variation or mutagenesis, as described in detail in subsection I above. Accordingly, in another embodiment, the STMST protein is a protein which comprises an amino acid sequence at least about 65% homologous to the amino acid sequence of SEQ ID NO:15 and retains the functional activity of the STMST proteins of SEQ ID NO:15, respectively.
25 Preferably, the protein is at least about 65-70%% homologous to SEQ ID NO:15, more preferably at least about 70-75% homologous to SEQ ID NO:15, even more preferably at least about 75-80% homologous to SEQ ID NO:15, and most preferably at least about 80-85%, 85-90%, or 90-95% homologous to SEQ ID NO:15.

 Biologically active portions of an STMST protein include peptides comprising
30 amino acid sequences sufficiently homologous to or derived from the amino acid sequence of the STMST protein, *e.g.*, the amino acid sequence shown in SEQ ID NO:15

- 60 -

or the amino acid sequence shown in SEQ ID NO:18, which include less amino acids than the full length STMST proteins, and exhibit at least one activity of an STMST protein. Typically, biologically active portions comprise a domain or motif with at least one activity of the STMST protein. A biologically active portion of an STMST protein
5 can be a polypeptide which is, for example, 10, 25, 50, 100 or more amino acids in length.

In one embodiment, a biologically active portion of an STMST protein comprises at least a transmembrane domain. In another embodiment, a biologically active portion of an STMST protein comprises at least one 7 transmembrane receptor
10 profile. In another embodiment, a biologically active portion of an STMST protein comprises at least a spectrin α -chain, repeated domain profile. In another embodiment a biologically active portion of an STMST protein comprises at least a 7 transmembrane receptor profile and a spectrin α -chain profile.

It is to be understood that a preferred biologically active portion of an STMST
15 protein of the present invention may contain at least one of the above-identified structural domains and/or profiles. A more preferred biologically active portion of an STMST protein may contain at least two of the above-identified structural domains and/or profiles. Moreover, other biologically active portions, in which other regions of the protein are deleted, can be prepared by recombinant techniques and evaluated for one
20 or more of the functional activities of a native STMST protein.

C. Chimeric/Fusion Proteins

The invention also provides BDSF and STMST chimeric or fusion proteins. As used herein, a "chimeric protein" or "fusion protein" comprises a BDSF or STMST
25 polypeptide operatively linked to a non-BDSF polypeptide or non-STMST polypeptide, respectively. A "BDSF polypeptide" refers to a polypeptide having an amino acid sequence corresponding to BDSF, whereas a "non-BDSF polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the BDSF protein, *e.g.*, a protein which is different from the
30 BDSF protein and which is derived from the same or a different organism. Likewise, a "STMST polypeptide" refers to a polypeptide having an amino acid sequence

- 61 -

corresponding to STMST, whereas a "non-STMST polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the STMST protein, *e.g.*, a protein which is different from the STMST protein and which is derived from the same or a different organism.

5 Within a BDSF or STMST fusion protein the BDSF or STMST polypeptide can correspond to all or a portion of a BDSF or STMST protein, respectively. In a preferred embodiment, a fusion protein comprises at least one biologically active portion of a BDSF or STMST protein. In another preferred embodiment, a fusion protein comprises at least two biologically active portions of a BDSF or STMST protein. Within the
10 fusion protein, the term "operatively linked" is intended to indicate that the BDSF or STMST polypeptide and the non-BDSF polypeptide or non-STMST polypeptide are fused in-frame to each other. The non-BDSF polypeptide or non-STMST polypeptide can be fused to the N-terminus or C-terminus of the BDSF polypeptide or STMST polypeptide, respectively.

15 For example, in one embodiment, the fusion protein is a GST-BDSF fusion protein in which the BDSF sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant BDSF. In another embodiment, the fusion protein is a GST-STMST fusion protein in which the STMST sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can
20 facilitate the purification of recombinant STMST.

 In another embodiment, the fusion protein is a BDSF protein containing a heterologous signal sequence at its N-terminus. For example, the native BDSF signal sequence (*i.e.*, about amino acids 1 to 25 of SEQ ID NO:2 or about amino acids 1 to 24 of SEQ ID NO:7) can be removed and replaced with a signal sequence from another
25 protein. In another embodiment, the fusion protein is an STMST protein containing a heterologous signal sequence at its N-terminus. In certain host cells (*e.g.*, mammalian host cells), expression and/or secretion of BDSF can be increased through use of a heterologous signal sequence.

 The fusion proteins of the invention can be incorporated into pharmaceutical
30 compositions and administered to a subject *in vivo*. The fusion proteins can be used to affect the bioavailability of a BDSF or STMST target molecule. Use of BDSF fusion

- 62 -

proteins may be useful therapeutically for the treatment of proliferative disorders (*e.g.*, cancer). Use of STMST fusion proteins may be useful therapeutically for the treatment of respiratory disorders (*e.g.*, asthma). Moreover, the fusion proteins of the invention can be used as immunogens to produce antibodies in a subject, to purify ligands and/or target molecules and in screening assays to identify molecules which inhibit the interaction of BDSF with a BDSF target molecule or the interaction of STMST with an STMST ligand.

Preferably, a chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel *et al.* John Wiley & Sons: 1992). Moreover, many expression vectors are commercially available that already encode a fusion moiety (*e.g.*, a GST polypeptide). A BDSF-encoding nucleic acid or STMST-encoding nucleic acid can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the BDSF protein or STMST protein, respectively.

25

D. Agonist/Anatgonist Variant Proteins

The present invention also pertains to variants of the BDSF proteins which function as either BDSF agonists (mimetics) or as BDSF antagonists and to variants of the STMST proteins which function as either STMST agonists (mimetics) or as STMST antagonists. Variants of the BDSF and BDSF proteins can be generated by mutagenesis, *e.g.*, discrete point mutation or truncation of a BDSF or STMST protein, respectively.

30

- 63 -

An agonist of the BDSF or STMST proteins can retain substantially the same, or a subset, of the biological activities of the naturally occurring form of a BDSF protein or STMST protein, respectively. An antagonist can inhibit one or more of the activities of the naturally occurring form of the protein by, for example, competitively inhibiting the activity of the BDSF or STMST protein. Thus, specific biological effects can be elicited by treatment with a variant of limited function. In one embodiment, treatment of a subject with a variant having a subset of the biological activities of the naturally occurring form of the protein has fewer side effects in a subject relative to treatment with the naturally occurring form of the protein.

10 In one embodiment, variants of a BDSF or STMST protein which function as either agonists (mimetics) or as antagonists can be identified by screening combinatorial libraries of mutants, *e.g.*, truncation mutants, of a BDSF or STMST protein for agonist or antagonist activity. In one embodiment, a variegated library of BDSF or STMST variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library can be produced by, for example, enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential BDSF or STMST sequences is expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (*e.g.*, for phage display) containing the set of BDSF or STMST sequences therein.

15 20 There are a variety of methods which can be used to produce libraries of variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential BDSF or STMST sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, *e.g.*, Narang, S.A. (1983) *Tetrahedron* 39:3; Itakura *et al.* (1984) *Annu. Rev. Biochem.* 53:323; Itakura *et al.* (1984) *Science* 198:1056; Ike *et al.* (1983) *Nucleic Acid Res.* 11:477.

In addition, libraries of fragments of a BDSF or STMST protein coding sequence can be used to generate a variegated population of fragments for screening and subsequent selection of protein variants. In one embodiment, a library of coding

30

- 64 -

sequence fragments can be generated by treating a double stranded PCR fragment of a coding sequence with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of a BDSF or STMST protein.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis procedures described herein. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a new technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify BDSF or STMST variants (Arkin and Yourvan (1992) *PNAS* 89:7811-7815; Delgrave *et al.* (1993) *Protein Engineering* 6(3):327-331).

In one embodiment, cell based assays can be exploited to analyze a variegated library. For example, a library of expression vectors can be transfected into a cell line which ordinarily synthesizes and secretes BDSF or STMST. The transfected cells are then cultured such that BDSF and a particular mutant BDSF (or STMST and a particular mutant) are secreted and the effect of expression of the mutant on BDSF or STMST activity in cell supernatants can be detected, *e.g.*, by any of a number of enzymatic or binding assays. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of BDSF or STMST activity, and the individual clones further characterized.

- 65 -

E. Protein Fragments, Immunogenic Peptides and Antibodies

An isolated BDSF or STMST protein, or a portion or fragment thereof, can be used as an immunogen to generate antibodies that bind BDSF or STMST using standard techniques for polyclonal and monoclonal antibody preparation. A full-length protein
5 can be used or, alternatively, the invention provides antigenic peptide fragments or use as immunogens. For example, an antigenic peptide of BDSF comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:2 or SEQ ID NO:7 and encompasses an epitope of BDSF such that an antibody raised against the peptide forms a specific immune complex with BDSF. Likewise, an antigenic peptide of STMST
10 comprises at least 8 amino acid residues of the amino acid sequence shown in SEQ ID NO:15 and encompasses an epitope of STMST such that an antibody raised against the peptide forms a specific immune complex with STMST. Preferably, the antigenic peptide comprises at least 10 amino acid residues, more preferably at least 15 amino acid residues, even more preferably at least 20 amino acid residues, and most preferably at
15 least 30 amino acid residues.

Preferred epitopes encompassed by the antigenic peptide are regions of BDSF that are located on the surface of the protein, *e.g.*, hydrophilic regions.

A BDSF or STMST immunogen typically is used to prepare antibodies by immunizing a suitable subject, (*e.g.*, rabbit, goat, mouse or other mammal) with the
20 immunogen. An appropriate immunogenic preparation can contain, for example, recombinantly expressed protein or a chemically synthesized polypeptide. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic BDSF preparation induces a polyclonal anti-BDSF antibody response.
25 Likewise, immunization of a suitable subject with an immunogenic STMST preparation induces a polyclonal anti-STMST antibody response.

Accordingly, another aspect of the invention pertains to anti-BDSF antibodies or anti-STMST antibodies. The term "antibody" as used herein refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, *i.e.*,
30 molecules that contain an antigen binding site which specifically binds (immunoreacts with) an antigen, such as BDSF or STMST. Examples of immunologically active

- 66 -

portions of immunoglobulin molecules include F(ab) and F(ab')₂ fragments which can be generated by treating the antibody with an enzyme such as pepsin. The invention provides polyclonal and monoclonal antibodies that bind BDSF or STMST. The term "monoclonal antibody" or "monoclonal antibody composition", as used herein, refers to
5 a population of antibody molecules that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of BDSF or STMST. A monoclonal antibody composition thus typically displays a single binding affinity for a particular BDSF protein with which it immunoreacts.

Polyclonal antibodies can be prepared as described above by immunizing a
10 suitable subject with a BDSF or STMST immunogen. The specific antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized BDSF or STMST. If desired, the antibody molecules can be isolated from the mammal (*e.g.*, from the blood) and further purified by well known techniques, such as protein A chromatography to
15 obtain the IgG fraction. At an appropriate time after immunization, *e.g.*, when the specific antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique originally described by Kohler and Milstein (1975) *Nature* 256:495-497) (see also, Brown *et al.* (1981) *J. Immunol.* 127:539-46; Brown *et al.*
20 (1980) *J. Biol. Chem.* 255:4980-83; Yeh *et al.* (1976) *PNAS* 76:2927-31; and Yeh *et al.* (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor *et al.* (1983) *Immunol Today* 4:72), the EBV-hybridoma technique (Cole *et al.* (1985), *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is
25 well known (see generally R. H. Kenneth, in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, New York (1980); E. A. Lerner (1981) *Yale J. Biol. Med.*, 54:387-402; M. L. Gefter *et al.* (1977) *Somatic Cell Genet.* 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with a BDSF or
30 STMST immunogen as described above, and the culture supernatants of the resulting

- 67 -

hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds BDSF or STMST, respectively.

Any of the many well known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating monoclonal antibodies (see, *e.g.*, G. Galfre *et al.* (1977) *Nature* 266:55052; Gefter *et al.* *Somatic Cell Genet.*, cited *supra*; Lerner, *Yale J. Biol. Med.*, cited *supra*; Kenneth, *Monoclonal Antibodies*, cited *supra*). Moreover, the ordinarily skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (*e.g.*, a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, *e.g.*, the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or Sp2/O-Ag14 myeloma lines. These myeloma lines are available from ATCC. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind BDSF or STMST, *e.g.*, using a standard ELISA assay.

Alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal anti-BDSF or anti-STMST antibody can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (*e.g.*, an antibody phage display library) with BDSF or STMST, respectively, to thereby isolate immunoglobulin library members that bind BDSF or STMST. Kits for generating and screening phage display libraries are commercially available (*e.g.*, the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents

- 68 -

particularly amenable for use in generating and screening antibody display library can be found in, for example, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* PCT International Publication No. WO 92/18619; Dower *et al.* PCT International Publication No. WO 91/17271; Winter *et al.* PCT International Publication WO 92/20791; Markland
 5 *et al.* PCT International Publication No. WO 92/15679; Breitling *et al.* PCT International Publication WO 93/01288; McCafferty *et al.* PCT International Publication No. WO 92/01047; Garrard *et al.* PCT International Publication No. WO 92/09690; Ladner *et al.* PCT International Publication No. WO 90/02809; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; Hay *et al.* (1992) *Hum. Antibod. Hybridomas* 3:81-85;
 10 Huse *et al.* (1989) *Science* 246:1275-1281; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J. Mol. Biol.* 226:889-896; Clarkson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc. Acid Res.* 19:4133-4137; Barbas *et al.* (1991) *PNAS* 88:7978-7982; and McCafferty *et al.* *Nature* (1990) 348:552-554.

15 Additionally, recombinant antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in
 20 Robinson *et al.* International Application No. PCT/US86/02269; Akira, *et al.* European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison *et al.* European Patent Application 173,494; Neuberger *et al.* PCT International Publication No. WO 86/01533; Cabilly *et al.* U.S. Patent No. 4,816,567; Cabilly *et al.* European Patent Application 125,023; Better *et al.* (1988) *Science*
 25 240:1041-1043; Liu *et al.* (1987) *PNAS* 84:3439-3443; Liu *et al.* (1987) *J. Immunol.* 139:3521-3526; Sun *et al.* (1987) *PNAS* 84:214-218; Nishimura *et al.* (1987) *Canc. Res.* 47:999-1005; Wood *et al.* (1985) *Nature* 314:446-449; and Shaw *et al.* (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi *et al.* (1986) *BioTechniques* 4:214; Winter U.S. Patent 5,225,539; Jones *et al.* (1986) *Nature*
 30 321:552-525; Verhoeyan *et al.* (1988) *Science* 239:1534; and Beidler *et al.* (1988) *J. Immunol.* 141:4053-4060.

- 69 -

An anti-BDSF or anti-STMST antibody (*e.g.*, monoclonal antibody) can be used to isolate BDSF or STMST by standard techniques, such as affinity chromatography or immunoprecipitation. An anti-BDSF or anti-STMST antibody can facilitate the purification of natural BDSF or STMST from cells and of recombinantly produced BDSF or STMST expressed in host cells. Moreover, an anti-BDSF or anti-STMST antibody can be used to detect BDSF protein or STMST (*e.g.*, in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the BDSF or STMST protein. Anti-BDSF or anti-STMST antibodies can be used diagnostically to monitor protein levels in tissue as part of a clinical testing procedure, *e.g.*, to, for example, determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (*i.e.*, physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S or ^3H .

III. Recombinant Expression Vectors and Host Cells

Another aspect of the invention pertains to vectors, preferably expression vectors, containing a nucleic acid encoding a BDSF or STMST protein (or a portion thereof). As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. Another type of vector is a viral vector, wherein additional DNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (*e.g.*,

- 70 -

bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (*e.g.*, non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (*e.g.*, replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The recombinant expression vectors of the invention comprise a nucleic acid of the invention in a form suitable for expression of the nucleic acid in a host cell, which means that the recombinant expression vectors include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operatively linked to the nucleic acid sequence to be expressed. Within a recombinant expression vector, "operably linked" is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleotide sequence (*e.g.*, in an *in vitro* transcription/translation system or in a host cell when the vector is introduced into the host cell). The term "regulatory sequence" is intended to include promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells (*e.g.*, tissue-specific regulatory sequences). It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce

- 71 -

proteins or peptides, including fusion proteins or peptides, encoded by nucleic acids as described herein (*e.g.*, BDSF proteins, mutant forms of BDSF proteins, STMST proteins, mutant forms of STMST proteins, fusion proteins, etc.).

The recombinant expression vectors of the invention can be designed for
5 expression of BDSF or STMST proteins in prokaryotic or eukaryotic cells. For example, BDSF or STMST proteins can be expressed in bacterial cells such as *E. coli*, insect cells (using baculovirus expression vectors) yeast cells or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the
10 recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein
15 encoded therein, usually to the amino terminus of the recombinant protein. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion
20 moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA)
25 and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding protein, or protein A, respectively, to the target recombinant protein.

Purified fusion proteins can be utilized in BDSF or STMST activity assays, (*e.g.*, direct assays or competitive assays described in detail below), or to generate antibodies specific for BDSF or STMST proteins, for example. In a preferred
30 embodiment, a BDSF or STMST fusion protein expressed in a retroviral expression vector of the present invention can be utilized to infect bone marrow cells which are

- 72 -

subsequently transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g six (6) weeks).

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann *et al.*, (1988) *Gene* 69:301-315) and pET 11d (Studier *et al.*, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89). Target gene expression from the pTrc vector relies on host RNA polymerase transcription from a hybrid trp-lac fusion promoter. Target gene expression from the pET 11d vector relies on transcription from a T7 gn10-lac fusion promoter mediated by a coexpressed viral RNA polymerase (T7 gn1). This viral polymerase is supplied by host strains BL21(DE3) or HMS174(DE3) from a resident prophage harboring a T7 gn1 gene under the transcriptional control of the lacUV 5 promoter.

One strategy to maximize recombinant protein expression in *E. coli* is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in *E. coli* (Wada *et al.*, (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

In another embodiment, the BDSF or STMST expression vector is a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, *et al.*, (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz *et al.*, (1987) *Gene* 54:113-123), pYES2 (Invitrogen Corporation, San Diego, CA), and picZ (InVitrogen Corp, San Diego, CA).

Alternatively, BDSF or STMST proteins can be expressed in insect cells using baculovirus expression vectors. Baculovirus vectors available for expression of proteins in cultured insect cells (e.g., Sf 9 cells) include the pAc series (Smith *et al.* (1983) *Mol. Cell Biol.* 3:2156-2165) and the pVL series (Lucklow and Summers (1989) *Virology* 170:31-39).

- 73 -

In yet another embodiment, a nucleic acid of the invention is expressed in mammalian cells using a mammalian expression vector. Examples of mammalian expression vectors include pCDM8 (Seed, B. (1987) *Nature* 329:840) and pMT2PC (Kaufman *et al.* (1987) *EMBO J.* 6:187-195). When used in mammalian cells, the
5 expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. For other suitable expression systems for both prokaryotic and eukaryotic cells see chapters 16 and 17 of Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual. 2nd, ed., Cold Spring*
10 *Harbor Laboratory*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989.

In another embodiment, the recombinant mammalian expression vector is capable of directing expression of the nucleic acid preferentially in a particular cell type (*e.g.*, tissue-specific regulatory elements are used to express the nucleic acid). Tissue-
15 specific regulatory elements are known in the art. Non-limiting examples of suitable tissue-specific promoters include the albumin promoter (liver-specific; Pinkert *et al.* (1987) *Genes Dev.* 1:268-277), lymphoid-specific promoters (Calame and Eaton (1988) *Adv. Immunol.* 43:235-275), in particular promoters of T cell receptors (Winoto and Baltimore (1989) *EMBO J.* 8:729-733) and immunoglobulins (Banerji *et al.* (1983) *Cell*
20 33:729-740; Queen and Baltimore (1983) *Cell* 33:741-748), neuron-specific promoters (*e.g.*, the neurofilament promoter; Byrne and Ruddle (1989) *PNAS* 86:5473-5477), pancreas-specific promoters (Edlund *et al.* (1985) *Science* 230:912-916), and mammary gland-specific promoters (*e.g.*, milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters
25 are also encompassed, for example the murine hox promoters (Kessel and Gruss (1990) *Science* 249:374-379) and the α -fetoprotein promoter (Campes and Tilghman (1989) *Genes Dev.* 3:537-546).

The invention further provides a recombinant expression vector comprising a DNA molecule of the invention cloned into the expression vector in an antisense
30 orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA molecule) of an

- 74 -

RNA molecule which is antisense to BDSF or STMST mRNA. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be
5 chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acids are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the
10 regulation of gene expression using antisense genes see Weintraub, H. *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) 1986.

Another aspect of the invention pertains to host cells into which a recombinant expression vector of the invention has been introduced. The terms "host cell" and
15 "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used
20 herein.

A host cell can be any prokaryotic or eukaryotic cell. For example, a BDSF or STMST protein can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

25 Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (*e.g.*, DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-mediated
30 transfection, lipofection, or electroporation. Suitable methods for transforming or transfecting host cells can be found in Sambrook, *et al.* (*Molecular Cloning: A*

- 75 -

Laboratory Manual. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989), and other laboratory manuals.

For stable transfection of mammalian cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may
5 integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (*e.g.*, resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate. Nucleic acid encoding a selectable marker can be
10 introduced into a host cell on the same vector as that encoding a BDSF or STMST protein or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by drug selection (*e.g.*, cells that have incorporated the selectable marker gene will survive, while the other cells die).

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in
15 culture, can be used to produce (*i.e.*, express) a BDSF or STMST protein. Accordingly, the invention further provides methods for producing a BDSF or STMST protein using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention (into which a recombinant expression vector encoding a BDSF or STMST protein has been introduced) in a suitable medium such that a BDSF or STMST
20 protein is produced. In another embodiment, the method further comprises isolating a BDSF or STMST protein from the medium or the host cell.

The host cells of the invention can also be used to produce nonhuman transgenic animals. For example, in one embodiment, a host cell of the invention is a fertilized oocyte or an embryonic stem cell into which BDSF-coding sequences or STMST-
25 coding sequences have been introduced. Such host cells can then be used to create non-human transgenic animals in which exogenous BDSF or STMST sequences have been introduced into their genome or homologous recombinant animals in which endogenous BDSF or STMST sequences have been altered, respectively. Such animals are useful for studying the function and/or activity of a BDSF or STMST and for identifying and/or
30 evaluating modulators of BDSF or STMST activity.

- 76 -

As used herein, a "transgenic animal" is a non-human animal, preferably a mammal, more preferably a rodent such as a rat or mouse, in which one or more of the cells of the animal includes a transgene. Other examples of transgenic animals include non-human primates, sheep, dogs, cows, goats, chickens, amphibians, etc. A transgene
5 is exogenous DNA which is integrated into the genome of a cell from which a transgenic animal develops and which remains in the genome of the mature animal, thereby directing the expression of an encoded gene product in one or more cell types or tissues of the transgenic animal. As used herein, a "homologous recombinant animal" is a non-human animal, preferably a mammal, more preferably a mouse, in which an endogenous
10 BDSF or STMST gene has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal, *e.g.*, an embryonic cell of the animal, prior to development of the animal.

A transgenic animal of the invention can be created by introducing a BDSF- or STMST-encoding nucleic acid into the male pronuclei of a fertilized oocyte, *e.g.*, by
15 microinjection, retroviral infection, and allowing the oocyte to develop in a pseudopregnant female foster animal. The BDSF cDNA sequence of SEQ ID NO:1 or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number 98756 can be introduced as a transgene into the genome of a non-human animal. Likewise, the STMST cDNA sequence of SEQ ID NO:14, SEQ ID
20 NO:17, the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ or the nucleotide sequence of the DNA insert of the plasmid deposited with ATCC as Accession Number _____ can be introduced as a transgene into the genome of a non-human animal. Alternatively, a nonhuman homologue of a human BDSF gene or STMST gene, such as a mouse BDSF or STMST
25 gene, can be used as a transgene. Alternatively, a BDSF or STMST gene homologue can be isolated based on hybridization to a BDSF or STMST cDNA sequences and used as a transgene.

Intronic sequences and polyadenylation signals can also be included in the transgene to increase the efficiency of expression of the transgene. A tissue-specific
30 regulatory sequence(s) can be operably linked to a BDSF or STMST transgene to direct expression of a BDSF or STMST protein to particular cells. Methods for generating

- 77 -

transgenic animals *via* embryo manipulation and microinjection, particularly animals such as mice, have become conventional in the art and are described, for example, in U.S. Patent Nos. 4,736,866 and 4,870,009, both by Leder *et al.*, U.S. Patent No. 4,873,191 by Wagner *et al.* and in Hogan, B., *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986). Similar methods are used for production of other transgenic animals. A transgenic founder animal can be identified based upon the presence of a BDSF or STMST transgene in its genome and/or expression of BDSF or STMST mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene.

Moreover, transgenic animals carrying a transgene encoding a BDSF or STMST protein can further be bred to other transgenic animals carrying other transgenes.

To create a homologous recombinant animal, a vector is prepared which contains at least a portion of a BDSF or STMST gene into which a deletion, addition or substitution has been introduced to thereby alter, *e.g.*, functionally disrupt, the BDSF or STMST gene. The BDSF or STMST gene can be a human gene (*e.g.*, the cDNA of SEQ ID NO:1 or the cDNA sequence of SEQ ID NO:14 or SEQ ID NO:17), but more preferably, is a non-human homologue of a human BDSF or STMST gene such as a murine BDSF or STMST gene. (For example, the mouse BDSF gene of SEQ ID NO:6 can be used to construct a homologous recombination vector suitable for altering an endogenous BDSF gene in the mouse genome.) In a preferred embodiment, the vector is designed such that, upon homologous recombination, the endogenous BDSF or STMST gene is functionally disrupted (*i.e.*, no longer encodes a functional protein; also referred to as a "knock out" vector). Alternatively, the vector can be designed such that, upon homologous recombination, the endogenous BDSF or STMST gene is mutated or otherwise altered but still encodes functional protein (*e.g.*, the upstream regulatory region can be altered to thereby alter the expression of the endogenous BDSF or STMST protein).

In the homologous recombination vector, the altered portion of the BDSF or STMST gene is flanked at its 5' and 3' ends by additional nucleic acid sequence of the BDSF or STMST gene to allow for homologous recombination to occur between the exogenous BDSF or STMST gene carried by the vector and an endogenous BDSF or

- 78 -

STMST gene in an embryonic stem cell. The additional flanking BDSF or STMST nucleic acid sequence is of sufficient length for successful homologous recombination with the endogenous gene. Typically, several kilobases of flanking DNA (both at the 5' and 3' ends) are included in the vector (see *e.g.*, Thomas, K.R. and Capecchi, M. R. (1987) *Cell* 51:503 for a description of homologous recombination vectors). The vector is introduced into an embryonic stem cell line (*e.g.*, by electroporation) and cells in which the introduced BDSF or STMST gene has homologously recombined with the endogenous BDSF or STMST gene are selected (see *e.g.*, Li, E. *et al.* (1992) *Cell* 69:915). The selected cells are then injected into a blastocyst of an animal (*e.g.*, a mouse) to form aggregation chimeras (see *e.g.*, Bradley, A. in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E.J. Robertson, ed. (IRL, Oxford, 1987) pp. 113-152). A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term. Progeny harboring the homologously recombined DNA in their germ cells can be used to breed animals in which all cells of the animal contain the homologously recombined DNA by germline transmission of the transgene. Methods for constructing homologous recombination vectors and homologous recombinant animals are described further in Bradley, A. (1991) *Current Opinion in Biotechnology* 2:823-829 and in PCT International Publication Nos.: WO 90/11354 by Le Mouellec *et al.*; WO 91/01140 by Smithies *et al.*; WO 92/0968 by Zijlstra *et al.*; and WO 93/04169 by Berns *et al.*

In another embodiment, transgenic non-humans animals can be produced which contain selected systems which allow for regulated expression of the transgene. One example of such a system is the *cre/loxP* recombinase system of bacteriophage P1. For a description of the *cre/loxP* recombinase system, see, *e.g.*, Lakso *et al.* (1992) *PNAS* 89:6232-6236. Another example of a recombinase system is the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman *et al.* (1991) *Science* 251:1351-1355. If a *cre/loxP* recombinase system is used to regulate expression of the transgene, animals containing transgenes encoding both the *Cre* recombinase and a selected protein are required. Such animals can be provided through the construction of "double" transgenic animals, *e.g.*, by mating two transgenic animals, one containing a transgene encoding a selected protein and the other containing a transgene encoding a recombinase.

- 79 -

Clones of the non-human transgenic animals described herein can also be produced according to the methods described in Wilmut, I. *et al.* (1997) *Nature* 385:810-813. In brief, a cell, *e.g.*, a somatic cell, from the transgenic animal can be isolated and induced to exit the growth cycle and enter G₀ phase. The quiescent cell can then be
5 fused, *e.g.*, through the use of electrical pulses, to an enucleated oocyte from an animal of the same species from which the quiescent cell is isolated. The reconstructed oocyte is then cultured such that it develops to morula or blastocyte and then transferred to pseudopregnant female foster animal. The offspring borne of this female foster animal will be a clone of the animal from which the cell, *e.g.*, the somatic cell, is isolated.

10 **IV. Pharmaceutical Compositions**

The BDSF nucleic acid molecules, BDSF proteins, anti-BDSF antibodies, STMST nucleic acid molecules, STMST proteins and anti-STMST antibodies (also referred to herein as "active compounds") of the invention can be incorporated into pharmaceutical compositions suitable for administration. Such compositions typically
15 comprise the nucleic acid molecule, protein, or antibody and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically
20 active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition of the invention is formulated to be compatible
25 with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils,
30 polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as

- 80 -

ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be
5 enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For
10 intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as
15 bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of
20 surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be
25 brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound (*e.g.*, a BDSF protein, anti-BDSF antibody, STMST protein or anti-STMST antibody) in the required amount in an appropriate solvent with one or a combination of
30 ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle

- 81 -

which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient
5 from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared
10 using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as
15 microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

20 For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be
25 permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in
30 the art.

- 82 -

The compounds can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, the active compounds are prepared with carriers that will
5 protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art.
10 The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

15 It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required
20 pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Toxicity and therapeutic efficacy of such compounds can be determined by
25 standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While
30 compounds that exhibit toxic side effects may be used, care should be taken to design a

- 83 -

delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

The nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see *e.g.*, Chen *et al.* (1994) *PNAS* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

V. Uses and Methods of the Invention

The nucleic acid molecules, proteins, protein homologues, and antibodies described herein can be used in one or more of the following methods: a) screening assays; b) predictive medicine (*e.g.*, diagnostic assays, prognostic assays, monitoring

- 84 -

clinical trials, and pharmacogenetics); and c) methods of treatment (*e.g.*, therapeutic and prophylactic).

As described herein, a BDSF protein of the invention has one or more of the following activities: (i) interaction of a BDSF protein in the extracellular milieu with a non-BDSF protein molecule on the surface of the same cell which secreted the BDSF protein molecule; (ii) interaction of a BDSF protein in the extracellular milieu with a non-BDSF protein molecule on the surface of a different cell from that which secreted the BDSF protein molecule; (iii) complex formation between a BDSF protein and a BDSF receptor; (iv) complex formation between a BDSF protein and non-BDSF receptor; and (v) interaction of a BDSF protein with a second protein in the extracellular milieu, and can thus be used in, for example, (1) modulation of cellular signal transduction, either *in vitro* or *in vivo*; (2) modulation of protein:protein interaction, either *in vitro* or *in vivo*; (3) regulation of cellular proliferation; or (4) regulation of cellular differentiation.

As described herein, an STMST protein of the invention has one or more of the following activities: (i) interaction of an STMST protein with soluble STMST ligand; (ii) interaction of an STMST protein with a membrane-bound non-STMST protein; (iii) interaction of an STMST protein with an intracellular protein (*e.g.*, an intracellular enzyme or signal transduction molecule); and (iv) indirect interaction of an STMST protein with an intracellular protein (*e.g.*, a downstream signal transduction molecule, and can thus be used in, for example, (1) modulation of cellular signal transduction, either *in vitro* or *in vivo*; (2) regulation of gene transcription in a cell expressing an STMST protein; (3) regulation of gene transcription in a cell expressing an STMST protein, wherein said cell is involved inflammation; (4) regulation of cellular proliferation; (5) regulation of cellular differentiation; (6) regulation of development; (7) regulation of cell death; (8) regulation of inflammation; and (9) regulation of respiratory cell function.

Accordingly, the isolated nucleic acid molecules of the invention can be used, for example, to express BDSF or STMST protein (*e.g.*, *via* a recombinant expression vector in a host cell in gene therapy applications), to detect BDSF or STMST mRNA (*e.g.*, in a biological sample) or a genetic alteration in a BDSF or STMST gene, and to modulate

- 85 -

BDSF or STMST activity, as described further below. The BDSF or STMST proteins can be used to treat disorders characterized by insufficient or excessive production of a BDSF, STMST, BDSF target molecule, or STMST target molecule. In addition, the BDSF or STMST proteins can be used to screen for naturally occurring BDSF or STMST target molecules, to screen for drugs or compounds which modulate BDSF or STMST activity, as well as to treat disorders characterized by insufficient or excessive production of BDSF or STMST protein or production of BDSF or STMST protein forms which have decreased or aberrant activity compared to BDSF or STMST wild type protein. Moreover, the anti-BDSF antibodies or anti-STMST antibodies of the invention can be used to detect and isolate BDSF or STMST proteins, regulate the bioavailability of BDSF or STMST proteins, and modulate BDSF or STMST activity.

Accordingly one embodiment of the present invention involves a method of use (*e.g.*, a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (*e.g.*, a BDSF protein, BDSF nucleic acid, BDSF modulator, STMST protein, STMST nucleic acid, STMST ligand or STMST modulator) is used, for example, to diagnose, prognose and/or treat a disease and/or condition in which any of the aforementioned activities is indicated. In another embodiment, the present invention involves a method of use (*e.g.*, a diagnostic assay, prognostic assay, or a prophylactic/therapeutic method of treatment) wherein a molecule of the present invention (*e.g.*, a BDSF protein, BDSF nucleic acid, BDSF modulator, STMST protein, STMST nucleic acid, STMST ligand or STMST modulator) is used, for example, for the diagnosis, prognosis, and/or treatment of subjects, preferably a human subject, in which any of the aforementioned activities is pathologically perturbed. In a preferred embodiment, the methods of use (*e.g.*, diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a subject, preferably a human subject, a molecule of the present invention (*e.g.*, a BDSF protein, BDSF nucleic acid, a BDSF modulator, STMST protein, STMST nucleic acid, STMST ligand or STMST modulator) for the diagnosis, prognosis, and/or therapeutic treatment. In another embodiment, the methods of use (*e.g.*, diagnostic assays, prognostic assays, or prophylactic/therapeutic methods of treatment) involve administering to a human subject a molecule of the present invention (*e.g.*, a BDSF protein, BDSF nucleic acid,

- 86 -

BDSF modulator, STMST protein, STMST nucleic acid, STMST ligand or STMST modulator).

A. Screening Assays:

5 The invention provides a method (also referred to herein as a "screening assay") for identifying modulators, *i.e.*, candidate or test compounds or agents (*e.g.*, peptides, peptidomimetics, small molecules or other drugs) which bind to BDSF or STMST proteins, have a stimulatory or inhibitory effect on, for example, BDSF or STMST expression or BDSF or STMST activity, or have a stimulatory or inhibitory effect on,
10 for example, the activity of an BDSF or STMST target molecule.

 In one embodiment, the invention provides assays for screening candidate or test compounds which are target molecules of a BDSF or STMST protein or polypeptide or biologically active portion thereof. In another embodiment, the invention provides assays for screening candidate or test compounds which bind to or modulate the activity
15 of a BDSF or STMST protein or polypeptide or biologically active portion thereof. The test compounds of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the 'one-bead one-compound' library method;
20 and synthetic library methods using affinity chromatography selection. The biological library approach is limited to peptide libraries, while the other four approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, K.S. (1997) *Anticancer Drug Des.* 12:145).

 Examples of methods for the synthesis of molecular libraries can be found in the
25 art, for example in: DeWitt *et al.* (1993) *Proc. Natl. Acad. Sci. U.S.A.* 90:6909; Erb *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann *et al.* (1994). *J. Med. Chem.* 37:2678; Cho *et al.* (1993) *Science* 261:1303; Carrell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell *et al.* (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop *et al.* (1994) *J. Med. Chem.* 37:1233.

- 87 -

Libraries of compounds may be presented in solution (*e.g.*, Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner USP 5,223,409), spores (Ladner USP '409), plasmids (Cull *et al.* (1992) *Proc Natl Acad Sci USA* 89:1865-1869) or on phage
5 (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla *et al.* (1990) *Proc. Natl. Acad. Sci.* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner *supra.*).

1. Cell-Based Assays

10 In one embodiment, an assay is a cell-based assay in which a cell which expresses a BDSF protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate BDSF activity determined. Determining the ability of the test compound to modulate BDSF activity can be accomplished by monitoring the bioactivity of the BDSF protein or biologically active
15 portion thereof. The cell, for example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to modulate BDSF activity can be accomplished, for example, by coupling the BDSF protein or biologically active portion thereof with a radioisotope or enzymatic label such that binding of the BDSF protein or biologically active portion thereof to its cognate target molecule can be determined by
20 detecting the labeled BDSF protein or biologically active portion thereof in a complex. For example, compounds (*e.g.*, BDSF protein or biologically active portion thereof) can be labeled with ¹²⁵I, ³⁵S, ¹⁴C, or ³H, either directly or indirectly, and the radioisotope detected by direct counting of radioemmission or by scintillation counting. Alternatively, compounds can be enzymatically labeled with, for example, horseradish
25 peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an appropriate substrate to product.

It is also within the scope of this invention to determine the ability of a compound (*e.g.*, BDSF protein or biologically active portion thereof) to interact with its cognate target molecule without the labeling of any of the interactants. For example, a
30 microphysiometer can be used to detect the interaction of a compound with its cognate target molecule without the labeling of either the compound or the receptor. McConnell,

- 88 -

H. M. *et al.* (1992) *Science* 257:1906-1912. As used herein, a "microphysiometer" (e.g., Cytosensor) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between
5 compound and receptor.

In a preferred embodiment, the assay comprises contacting a cell which expresses a BDSF protein or biologically active portion thereof, with a target molecule to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to modulate the activity of the BDSF
10 protein or biologically active portion thereof, wherein determining the ability of the test compound to modulate the activity of the BDSF protein or biologically active portion thereof, comprises determining the ability of the test compound to modulate a biological activity of the BDSF expressing cell (e.g., determining the ability of the test compound to modulate signal transduction or protein:protein interactions).

15 In another preferred embodiment, the assay comprises contacting a cell which is responsive to a BDSF protein or biologically active portion thereof, with a BDSF protein or biologically-active portion thereof, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to modulate the activity of the BDSF protein or biologically active portion thereof, wherein
20 determining the ability of the test compound to modulate the activity of the BDSF protein or biologically active portion thereof comprises determining the ability of the test compound to modulate a biological activity of the BDSF-responsive cell (e.g., determining the ability of the test compound to modulate signal transduction or protein:protein interactions).

25 In another embodiment, an assay is a cell-based assay comprising contacting a cell expressing a BDSF target molecule with a test compound and determining the ability of the test compound to modulate (e.g. stimulate or inhibit) the activity of the BDSF target molecule. Determining the ability of the test compound to modulate the activity of a BDSF target molecule can be accomplished, for example, by determining
30 the ability of the BDSF protein to bind to or interact with the BDSF target molecule.

- 89 -

Determining the ability of the BDSF protein to bind to or interact with a BDSF target molecule can be accomplished by one of the methods described above for determining direct binding. In a preferred embodiment, determining the ability of the BDSF protein to bind to or interact with a BDSF target molecule can be accomplished
5 by determining the activity of the target molecule. For example, the activity of the target molecule can be determined by detecting induction of a cellular second messenger of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (comprising a target-responsive regulatory element operatively linked
10 to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a target-regulated cellular response, for example, signal transduction or protein:protein interactions.

In another embodiment, an assay is a cell-based assay in which a cell which expresses an STMST protein on the cell surface is contacted with a test compound and
15 the ability of the test compound to bind to the STMST protein determined. The cell, for example, can be of mammalian origin or a yeast cell. Determining the ability of the test compound to bind to an STMST protein can be accomplished, for example, by coupling the test compound with a radioisotope or enzymatic label such that binding of the test compound to the STMST protein can be determined by detecting the labeled compound
20 in a complex. For example, test compounds can be labeled with ^{125}I , ^{35}S , ^{14}C , or ^3H , either directly or indirectly, and the radioisotope detected by direct counting of radioemission or by scintillation counting. Alternatively, test compounds can be enzymatically labeled with, for example, horseradish peroxidase, alkaline phosphatase, or luciferase, and the enzymatic label detected by determination of conversion of an
25 appropriate substrate to product.

It is also within the scope of this invention to determine the ability of a test compound to interact with an STMST protein without the labeling of any of the interactants. For example, a microphysiometer can be used to detect the interaction of a test compound with an STMST protein without the labeling of either the test compound
30 or the receptor, as described above.

- 90 -

In a preferred embodiment, the assay comprises contacting a cell which expresses an STMST protein or biologically active portion thereof, on the cell surface with an STMST ligand, to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the STMST protein or biologically active portion thereof, wherein determining the ability of the test compound to interact with the STMST protein or biologically active portion thereof, comprises determining the ability of the test compound to preferentially bind to the STMST protein or biologically active portion thereof, as compared to the ability of the STMST ligand to bind to the STMST protein or biologically active portion thereof.

Determining the ability of the STMST ligand or STMST modulator to bind to or interact with an STMST protein or biologically active portion thereof, can be accomplished by one of the methods described above for determining direct binding. In a preferred embodiment, determining the ability of the STMST ligand or modulator to bind to or interact with an STMST protein or biologically active portion thereof, can be accomplished by determining the activity of an STMST protein or of a downstream STMST target molecule. For example, the target molecule can be a cellular second messenger, and the activity of the target molecule can be determined by detecting induction of the target (*i.e.* intracellular Ca^{2+} , diacylglycerol, IP_3 , etc.), detecting catalytic/enzymatic activity of the target on an appropriate substrate, detecting the induction of a reporter gene (comprising an STMST-responsive regulatory element operatively linked to a nucleic acid encoding a detectable marker, *e.g.*, luciferase), or detecting a cellular response, for example, a proliferative response or an inflammatory response. Accordingly, in one embodiment the present invention involves a method of identifying a compound which modulates the activity of an STMST protein, comprising contacting a cell which expresses an STMST protein with a test compound, determining the ability of the test compound to modulate the activity the STMST protein, and identifying the compound as a modulator of STMST activity. In another embodiment, the present invention involves a method of identifying a compound which modulates the activity of an STMST protein, comprising contacting a cell which expresses an STMST protein with a test compound, determining the ability of the test compound to modulate

- 91 -

the activity of a downstream STMST target molecule, and identifying the compound as a modulator of STMST activity.

2. Cell-Free Assays

5 In yet another embodiment, an assay of the present invention is a cell-free assay in which a BDSF or STMST protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the BDSF or STMST protein or biologically active portion thereof is determined. Binding of the test compound to the BDSF or STMST protein can be determined either directly or
10 indirectly as described above. In a preferred embodiment, the assay includes contacting the BDSF or STMST protein or biologically active portion thereof with a known compound which binds BDSF or STMST (*e.g.*, a BDSF or STMST target molecule) to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with a BDSF or STMST protein,
15 wherein determining the ability of the test compound to interact with a BDSF or STMST protein comprises determining the ability of the test compound to preferentially bind to BDSF or STMST or biologically active portion thereof as compared to the known compound.

 In another embodiment, the assay is a cell-free assay in which a BDSF protein or
20 biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the BDSF protein or biologically active portion thereof is determined. Determining the ability of the test compound to modulate the activity of a BDSF protein can be accomplished, for example, by determining the ability of the BDSF protein to bind to a BDSF target
25 molecule by one of the methods described above for determining direct binding. Determining the ability of the BDSF protein to bind to a BDSF target molecule can also be accomplished using a technology such as real-time Biomolecular Interaction Analysis (BIA). Sjolander, S. and Urbaniczky, C. (1991) *Anal. Chem.* 63:2338-2345 and Szabo
et al. (1995) *Curr. Opin. Struct. Biol.* 5:699-705. As used herein, "BIA" is a technology
30 for studying biospecific interactions in real time, without labeling any of the interactants

- 92 -

(*e.g.*, BIAcore). Changes in the optical phenomenon of surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In an alternative embodiment, determining the ability of the test compound to modulate the activity of a BDSF protein can be accomplished by determining the ability
5 of the BDSF protein to further modulate the activity of a downstream effector (*e.g.*, a growth factor mediated signal transduction pathway component) of a BDSF target molecule. For example, the activity of the effector molecule on an appropriate target can be determined or the binding of the effector to an appropriate target can be determined as previously described.

10 In yet another embodiment, the cell-free assay involves contacting a BDSF protein or biologically active portion thereof with a known compound which binds the BDSF protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the BDSF protein, wherein determining the ability of the test compound to interact with the BDSF
15 protein comprises determining the ability of the BDSF protein to preferentially bind to or modulate the activity of a BDSF target molecule.

In yet another embodiment, an assay of the present invention is a cell-free assay in which an STMST protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to the STMST protein or
20 biologically active portion thereof is determined. Binding of the test compound to the STMST protein can be determined either directly or indirectly or using BIA as described above.

In a preferred embodiment, the assay includes contacting the STMST protein or biologically active portion thereof with a known ligand which binds STMST to form an
25 assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with an STMST protein, wherein determining the ability of the test compound to interact with an STMST protein comprises determining the ability of the test compound to preferentially bind to STMST or biologically active portion thereof as compared to the known ligand.

- 93 -

In another embodiment, the assay is a cell-free assay in which an STMST protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to modulate (*e.g.*, stimulate or inhibit) the activity of the STMST protein or biologically active portion thereof is determined. Determining the ability of the test compound to modulate the activity of an STMST protein can be accomplished, for example, by determining the ability of the STMST protein to modulate the activity of a downstream STMST target molecule by one of the methods described above for cell-based assays. For example, the catalytic/enzymatic activity of the target molecule on an appropriate substrate can be determined as previously described.

In yet another embodiment, the cell-free assay involves contacting an STMST protein or biologically active portion thereof with a known ligand which binds the STMST protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to interact with the STMST protein, wherein determining the ability of the test compound to interact with the STMST protein comprises determining the ability of the test compound to preferentially bind to or modulate the activity of an STMST target molecule, as compared to the known ligand.

The cell-free assays of the present invention are amenable to use of both soluble and/or membrane-bound forms of isolated proteins (*e.g.* BDSF or STMST proteins or biologically active portions thereof or receptors to which BDSF targets bind or STMST proteins or biologically active portions thereof or STMST proteins). In the case of cell-free assays in which a membrane-bound form of an isolated protein is used (*e.g.*, a cell surface receptor) it may be desirable to utilize a solubilizing agent such that the membrane-bound form of the isolated protein is maintained in solution. Examples of such solubilizing agents include non-ionic detergents such as n-octylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton[®] X-100, Triton[®] X-114, Thesit[®], Isotridecypoly(ethylene glycol ether)_n, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPS), 3-[(3-cholamidopropyl)dimethylamminio]-2-hydroxy-1-propane sulfonate (CHAPSO), or N-dodecyl=N,N-dimethyl-3-ammonio-1-propane sulfonate.

- 94 -

In more than one embodiment of the above assay methods of the present invention, it may be desirable to immobilize either BDSF or STMST or their target molecule to facilitate separation of complexed from uncomplexed forms of one or both of the proteins, as well as to accommodate automation of the assay. Binding of a test
5 compound to a BDSF or STMST protein, or interaction of a BDSF or STMST protein with a target molecule in the presence and absence of a candidate compound, can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtitre plates, test tubes, and micro-centrifuge tubes. In one
10 embodiment, a fusion protein can be provided which adds a domain that allows one or both of the proteins to be bound to a matrix. For example, GST/ BDSF fusion proteins, GST/STMST fusion proteins or GST/target fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivatized microtitre plates, which are then combined with the test compound or the test compound and either the non-adsorbed target protein, BDSF or STMST protein, and the mixture
15 incubated under conditions conducive to complex formation (*e.g.*, at physiological conditions for salt and pH). Following incubation, the beads or microtitre plate wells are washed to remove any unbound components, the matrix immobilized in the case of beads, complex determined either directly or indirectly, for example, as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of BDSF
20 or STMST binding or activity determined using standard techniques.

Other techniques for immobilizing proteins on matrices can also be used in the screening assays of the invention. For example, a BDSF protein. STMST fusion protein, a BDSF target molecule or STMST target molecule can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated BDSF or STMST protein or target
25 molecules can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques well known in the art (*e.g.*, biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies reactive with BDSF protein, STMST proteins or target molecules but which do not interfere with binding of the BDSF or STMST protein to its
30 target molecule can be derivatized to the wells of the plate, and unbound target or BDSF or STMST protein trapped in the wells by antibody conjugation. Methods for detecting

- 95 -

such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the BDSF protein, STMST protein or target molecules, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the BDSF protein,
5 STMST protein or respective target molecule.

In another embodiment, modulators of BDSF or STMST expression are identified in a method wherein a cell is contacted with a candidate compound and the expression of BDSF or STMST mRNA or protein in the cell is determined. The level of expression of BDSF or STMST mRNA or protein in the presence of the candidate
10 compound is compared to the level of expression of BDSF or STMST mRNA or protein in the absence of the candidate compound. The candidate compound can then be identified as a modulator of BDSF or STMST expression based on this comparison. For example, when expression of BDSF or STMST mRNA or protein is greater (statistically significantly greater) in the presence of the candidate compound than in its absence, the
15 candidate compound is identified as a stimulator of BDSF or STMST mRNA or protein expression. Alternatively, when expression of BDSF or STMST mRNA or protein is less (statistically significantly less) in the presence of the candidate compound than in its absence, the candidate compound is identified as an inhibitor of BDSF or STMST mRNA or protein expression. The level of BDSF or STMST mRNA or protein
20 expression in the cells can be determined by methods described herein for detecting BDSF or STMST mRNA or protein.

In yet another aspect of the invention, the BDSF or STMST proteins can be used as "bait proteins" in a two-hybrid assay or three-hybrid assay (see, *e.g.*, U.S. Patent No. 5,283,317; Zervos *et al.* (1993) *Cell* 72:223-232; Madura *et al.* (1993) *J. Biol. Chem.*
25 268:12046-12054; Bartel *et al.* (1993) *Biotechniques* 14:920-924; Iwabuchi *et al.* (1993) *Oncogene* 8:1693-1696; and Brent WO94/10300), to identify other proteins, which bind to or interact with BDSF or STMST ("BDSF-binding proteins", "BDSF-bp", "STMST-binding proteins" or "STMST-bp") and are involved in BDSF or STMST activity. Such binding proteins are also likely to be involved in the propagation of
30 signals by the BDSF proteins, STMST proteins or respective targets as, for example, downstream elements of a BDSF- or STMST-mediated signaling pathway.

- 96 -

Alternatively, such BDSF-binding proteins or STMST-binding proteins are likely to be BDSF or STMST inhibitors.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the
5 assay utilizes two different DNA constructs. In one construct, the gene that codes for a BDSF or STMST protein is fused to a gene encoding the DNA binding domain of a known transcription factor (*e.g.*, GAL-4). In the other construct, a DNA sequence, from a library of DNA sequences, that encodes an unidentified protein ("prey" or "sample") is fused to a gene that codes for the activation domain of the known transcription factor. If
10 the "bait" and the "prey" proteins are able to interact, *in vivo*, forming a BDSF-dependent complex or STMST-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (*e.g.*, LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the
15 reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to obtain the cloned gene which encodes the protein which interacts with the BDSF or STMST protein.

This invention further pertains to novel agents identified by the above-described screening assays and to processes for producing such agents by use of these assays.
20 Accordingly, in one embodiment, the present invention includes a compound or agent obtainable by a method comprising the steps of any one of the aforementioned screening assays (*e.g.*, cell-based assays or cell-free assays). For example, in one embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a BDSF or STMST target molecule with a test compound and the
25 determining the ability of the test compound to bind to, or modulate the activity of, the BDSF or STMST target molecule. In another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a cell which expresses a BDSF or STMST target molecule with a BDSF or STMST protein or biologically-active portion thereof, to form an assay mixture, contacting the assay
30 mixture with a test compound, and determining the ability of the test compound to interact with, or modulate the activity of, the BDSF or STMST target molecule. In

- 97 -

another embodiment, the invention includes a compound or agent obtainable by a method comprising contacting a BDSF or STMST protein or biologically active portion thereof with a test compound and determining the ability of the test compound to bind to, or modulate (*e.g.*, stimulate or inhibit) the activity of, the BDSF or STMST protein or
5 biologically active portion thereof. In yet another embodiment, the present invention included a compound or agent obtainable by a method comprising contacting a BDSF or STMST protein or biologically active portion thereof with a known compound which binds the BDSF or STMST protein to form an assay mixture, contacting the assay mixture with a test compound, and determining the ability of the test compound to
10 interact with, or modulate the activity of the BDSF or STMST protein.

Accordingly, it is within the scope of this invention to further use an agent identified as described herein in an appropriate animal model. For example, an agent identified as described herein (*e.g.*, a BDSF or STMST modulating agent, an antisense BDSF or STMST nucleic acid molecule, a BDSF-specific or STMST-specific antibody,
15 or a BDSF- or STMST-binding partner) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as
20 described herein.

The present invention also pertains to uses of novel agents identified by the above-described screening assays for diagnoses, prognoses, and treatments as described herein. Accordingly, it is within the scope of the present invention to use such agents in the design, formulation, synthesis, manufacture, and/or production of a drug or
25 pharmaceutical composition for use in diagnosis, prognosis, or treatment, as described herein. For example, in one embodiment, the present invention includes a method of synthesizing or producing a drug or pharmaceutical composition by reference to the structure and/or properties of a compound obtainable by one of the above-described screening assays. For example, a drug or pharmaceutical composition can be
30 synthesized based on the structure and/or properties of a compound obtained by a method in which a cell which expresses a BDSF or STMST target molecule is contacted

- 98 -

with a test compound and the ability of the test compound to bind to, or modulate the activity of, the BDSF or STMST target molecule is determined. In another exemplary embodiment, the present invention includes a method of synthesizing or producing a drug or pharmaceutical composition based on the structure and/or properties of a compound obtainable by a method in which a BDSF or STMST protein or biologically active portion thereof is contacted with a test compound and the ability of the test compound to bind to, or modulate (*e.g.*, stimulate or inhibit) the activity of, the BDSF or STMST protein or biologically active portion thereof is determined.

10 B. Detection Assays

Portions or fragments of the cDNA sequences identified herein (and the corresponding complete gene sequences) can be used in numerous ways as polynucleotide reagents. For example, these sequences can be used to: (i) map their respective genes on a chromosome; and, thus, locate gene regions associated with genetic disease; (ii) identify an individual from a minute biological sample (tissue typing); and (iii) aid in forensic identification of a biological sample. These applications are described in the subsections below.

1. Chromosome Mapping

20 Once the sequence (or a portion of the sequence) of a gene has been isolated, this sequence can be used to map the location of the gene on a chromosome. This process is called chromosome mapping. Accordingly, portions or fragments of the BDSF or STMST nucleotide sequences, described herein, can be used to map the location of the BDSF or STMST genes on a chromosome. The mapping of the BDSF or STMST sequences to chromosomes is an important first step in correlating these sequences with genes associated with disease.

Briefly, BDSF or STMST genes can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp in length) from the BDSF or STMST nucleotide sequences. Computer analysis of the BDSF or STMST sequences can be used to predict primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers can then be used for PCR screening of somatic

- 99 -

cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the BDSF or STMST sequences will yield an amplified fragment.

Somatic cell hybrids are prepared by fusing somatic cells from different mammals (*e.g.*, human and mouse cells). As hybrids of human and mouse cells grow and divide, they gradually lose human chromosomes in random order, but retain the mouse chromosomes. By using media in which mouse cells cannot grow, because they lack a particular enzyme, but human cells can, the one human chromosome that contains the gene encoding the needed enzyme, will be retained. By using various media, panels of hybrid cell lines can be established. Each cell line in a panel contains either a single human chromosome or a small number of human chromosomes, and a full set of mouse chromosomes, allowing easy mapping of individual genes to specific human chromosomes. (D'Eustachio P. *et al.* (1983) *Science* 220:919-924). Somatic cell hybrids containing only fragments of human chromosomes can also be produced by using human chromosomes with translocations and deletions.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular sequence to a particular chromosome. Three or more sequences can be assigned per day using a single thermal cycler. Using the BDSF or STMST nucleotide sequences to design oligonucleotide primers, sublocalization can be achieved with panels of fragments from specific chromosomes. Other mapping strategies which can similarly be used to map a 9o, 1p, or 1v sequence to its chromosome include *in situ* hybridization (described in Fan, Y. *et al.* (1990) *PNAS*, 87:6223-27), pre-screening with labeled flow-sorted chromosomes, and pre-selection by hybridization to chromosome specific cDNA libraries.

Fluorescence *in situ* hybridization (FISH) of a DNA sequence to a metaphase chromosomal spread can further be used to provide a precise chromosomal location in one step. Chromosome spreads can be made using cells whose division has been blocked in metaphase by a chemical such as colcemid that disrupts the mitotic spindle. The chromosomes can be treated briefly with trypsin, and then stained with Giemsa. A pattern of light and dark bands develops on each chromosome, so that the chromosomes can be identified individually. The FISH technique can be used with a DNA sequence

- 100 -

as short as 500 or 600 bases. However, clones larger than 1,000 bases have a higher likelihood of binding to a unique chromosomal location with sufficient signal intensity for simple detection. Preferably 1,000 bases, and more preferably 2,000 bases will suffice to get good results at a reasonable amount of time. For a review of this
5 technique, see Verma *et al.*, Human Chromosomes: A Manual of Basic Techniques (Pergamon Press, New York 1988).

Reagents for chromosome mapping can be used individually to mark a single chromosome or a single site on that chromosome, or panels of reagents can be used for marking multiple sites and/or multiple chromosomes. Reagents corresponding to
10 noncoding regions of the genes actually are preferred for mapping purposes. Coding sequences are more likely to be conserved within gene families, thus increasing the chance of cross hybridizations during chromosomal mapping.

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map
15 data. (Such data are found, for example, in V. McKusick, Mendelian Inheritance in Man, available on-line through Johns Hopkins University Welch Medical Library). The relationship between a gene and a disease, mapped to the same chromosomal region, can then be identified through linkage analysis (co-inheritance of physically adjacent genes), described in, for example, Egeland, J. *et al.* (1987) *Nature*, 325:783-787.

Moreover, differences in the DNA sequences between individuals affected and
20 unaffected with a disease associated with the BDSF or STMST gene, can be determined. If a mutation is observed in some or all of the affected individuals but not in any unaffected individuals, then the mutation is likely to be the causative agent of the particular disease. Comparison of affected and unaffected individuals generally involves
25 first looking for structural alterations in the chromosomes, such as deletions or translocations that are visible from chromosome spreads or detectable using PCR based on that DNA sequence. Ultimately, complete sequencing of genes from several individuals can be performed to confirm the presence of a mutation and to distinguish mutations from polymorphisms.

2. Tissue Typing

The BDSF or STMST sequences of the present invention can also be used to identify individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for
5 identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identification. This method does not suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The sequences of the present invention are useful as additional DNA markers
10 for RFLP (described in U.S. Patent 5,272,057).

Furthermore, the sequences of the present invention can be used to provide an alternative technique which determines the actual base-by-base DNA sequence of selected portions of an individual's genome. Thus, the BDSF or STMST nucleotide sequences described herein can be used to prepare two PCR primers from the 5' and 3'
15 ends of the sequences. These primers can then be used to amplify an individual's DNA and subsequently sequence it.

Panels of corresponding DNA sequences from individuals, prepared in this manner, can provide unique individual identifications, as each individual will have a unique set of such DNA sequences due to allelic differences. The sequences of the
20 present invention can be used to obtain such identification sequences from individuals and from tissue. The BDSF or STMST nucleotide sequences of the invention uniquely represent portions of the human genome. Allelic variation occurs to some degree in the coding regions of these sequences, and to a greater degree in the noncoding regions. It is estimated that allelic variation between individual humans occurs with a frequency of
25 about once per each 500 bases. Each of the sequences described herein can, to some degree, be used as a standard against which DNA from an individual can be compared for identification purposes. Because greater numbers of polymorphisms occur in the noncoding regions, fewer sequences are necessary to differentiate individuals. The noncoding sequences of SEQ ID NO:1, SEQ ID NO:6, SEQ ID NO:14 or SEQ ID
30 NO:17, can comfortably provide positive individual identification with a panel of perhaps 10 to 1,000 primers which each yield a noncoding amplified sequence of 100

- 102 -

bases. If predicted coding sequences, such as those in SEQ ID NO:3, SEQ ID NO:8, SEQ ID NO:16 or SEQ ID NO:19 are used, a more appropriate number of primers for positive individual identification would be 500-2,000.

If a panel of reagents from BDSF or STMST nucleotide sequences described herein is used to generate a unique identification database for an individual, those same reagents can later be used to identify tissue from that individual. Using the unique identification database, positive identification of the individual, living or dead, can be made from extremely small tissue samples.

10 **3. Use of Partial BDSF or STMST Sequences in Forensic Biology**

DNA-based identification techniques can also be used in forensic biology. Forensic biology is a scientific field employing genetic typing of biological evidence found at a crime scene as a means for positively identifying, for example, a perpetrator of a crime. To make such an identification, PCR technology can be used to amplify
15 DNA sequences taken from very small biological samples such as tissues, *e.g.*, hair or skin, or body fluids, *e.g.*, blood, saliva, or semen found at a crime scene. The amplified sequence can then be compared to a standard, thereby allowing identification of the origin of the biological sample.

The sequences of the present invention can be used to provide polynucleotide
20 reagents, *e.g.*, PCR primers, targeted to specific loci in the human genome, which can enhance the reliability of DNA-based forensic identifications by, for example, providing another "identification marker" (*i.e.* another DNA sequence that is unique to a particular individual). As mentioned above, actual base sequence information can be used for identification as an accurate alternative to patterns formed by restriction enzyme
25 generated fragments. Sequences targeted to noncoding regions of SEQ ID NO:1 are particularly appropriate for this use as greater numbers of polymorphisms occur in the noncoding regions, making it easier to differentiate individuals using this technique. Examples of polynucleotide reagents include the BDSF or STMST nucleotide sequences or portions thereof, *e.g.*, fragments derived from the noncoding regions of SEQ ID
30 NO:1, SEQ ID NO:6, SEQ ID NO:14 or SEQ ID NO:17 having a length of at least 20 bases, preferably at least 30 bases.

- 103 -

The BDSF or STMST nucleotide sequences described herein can further be used to provide polynucleotide reagents, *e.g.*, labeled or labelable probes which can be used in, for example, an *in situ* hybridization technique, to identify a specific tissue, *e.g.*, brain tissue. This can be very useful in cases where a forensic pathologist is presented
5 with a tissue of unknown origin. Panels of such BDSF or STMST probes can be used to identify tissue by species and/or by organ type.

In a similar fashion, these reagents, *e.g.*, BDSF or STMST primers or probes can be used to screen tissue culture for contamination (*i.e.* screen for the presence of a mixture of different types of cells in a culture).

10

C. Predictive Medicine:

The present invention also pertains to the field of predictive medicine in which diagnostic assays, prognostic assays, and monitoring clinical trials are used for prognostic (predictive) purposes to thereby treat an individual prophylactically.
15 Accordingly, one aspect of the present invention relates to diagnostic assays for determining BDSF or STMST protein and/or nucleic acid expression as well as BDSF or STMST activity, in the context of a biological sample (*e.g.*, blood, serum, cells, tissue) to thereby determine whether an individual is afflicted with a disease or disorder, or is at risk of developing a disorder, associated with aberrant BDSF or STMST expression or
20 activity. The invention also provides for prognostic (or predictive) assays for determining whether an individual is at risk of developing a disorder associated with BDSF or STMST protein, nucleic acid expression or activity. For example, mutations in a BDSF or STMST gene can be assayed in a biological sample. Such assays can be used for prognostic or predictive purpose to thereby prophylactically treat an individual prior
25 to the onset of a disorder characterized by or associated with BDSF or STMST protein, nucleic acid expression or activity.

Another aspect of the invention pertains to monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of BDSF in clinical trials.

These and other agents are described in further detail in the following sections.

30

1. Diagnostic Assays

An exemplary method for detecting the presence or absence of BDSF or STMST protein or nucleic acid in a biological sample involves obtaining a biological sample from a test subject and contacting the biological sample with a compound or an agent
5 capable of detecting BDSF or STMST protein or nucleic acid (*e.g.*, mRNA, genomic DNA) that encodes BDSF or STMST protein such that the presence of BDSF or STMST protein or nucleic acid is detected in the biological sample. A preferred agent for detecting BDSF or STMST mRNA or genomic DNA is a labeled nucleic acid probe capable of hybridizing to BDSF or STMST mRNA or genomic DNA. The nucleic acid
10 probe can be, for example, a full-length BDSF nucleic acid, or a portion thereof, such as an oligonucleotide of at least 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to BDSF or STMST mRNA or genomic DNA. Other suitable probes for use in the diagnostic assays of the invention are described herein.

15 A preferred agent for detecting BDSF or STMST protein is an antibody capable of binding to BDSF or STMST protein, preferably an antibody with a detectable label. Antibodies can be polyclonal, or more preferably, monoclonal. An intact antibody, or a fragment thereof (*e.g.*, Fab or F(ab')₂) can be used. The term "labeled", with regard to the probe or antibody, is intended to encompass direct labeling of the probe or antibody
20 by coupling (*i.e.*, physically linking) a detectable substance to the probe or antibody, as well as indirect labeling of the probe or antibody by reactivity with another reagent that is directly labeled. Examples of indirect labeling include detection of a primary antibody using a fluorescently labeled secondary antibody and end-labeling of a DNA probe with biotin such that it can be detected with fluorescently labeled streptavidin.
25 The term "biological sample" is intended to include tissues, cells and biological fluids isolated from a subject, as well as tissues, cells and fluids present within a subject. That is, the detection method of the invention can be used to detect BDSF or STMST mRNA, protein, or genomic DNA in a biological sample *in vitro* as well as *in vivo*. For example, *in vitro* techniques for detection of BDSF or STMST mRNA include Northern
30 hybridizations and *in situ* hybridizations. *In vitro* techniques for detection of BDSF or STMST protein include enzyme linked immunosorbent assays (ELISAs), Western blots,

- 105 -

immunoprecipitations and immunofluorescence. *In vitro* techniques for detection of BDSF or STMST genomic DNA include Southern hybridizations. Furthermore, *in vivo* techniques for detection of BDSF or STMST protein include introducing into a subject a labeled anti-BDSF or anti-STMST antibody. For example, the antibody can be labeled
5 with a radioactive marker whose presence and location in a subject can be detected by standard imaging techniques.

In one embodiment, the biological sample contains protein molecules from the test subject. Alternatively, the biological sample can contain mRNA molecules from the test subject or genomic DNA molecules from the test subject. A preferred biological
10 sample is a serum sample isolated by conventional means from a subject.

In another embodiment, the methods further involve obtaining a control biological sample from a control subject, contacting the control sample with a compound or agent capable of detecting BDSF or STMST protein, mRNA, or genomic DNA, such that the presence of BDSF or STMST protein, mRNA or genomic DNA is
15 detected in the biological sample, and comparing the presence of BDSF or STMST protein, mRNA or genomic DNA in the control sample with the presence of BDSF or STMST protein, mRNA or genomic DNA in the test sample.

The invention also encompasses kits for detecting the presence of BDSF or STMST in a biological sample. For example, the kit can comprise a labeled compound
20 or agent capable of detecting BDSF or STMST protein or mRNA in a biological sample; means for determining the amount of BDSF or STMST in the sample; and means for comparing the amount of BDSF or STMST in the sample with a standard. The compound or agent can be packaged in a suitable container. The kit can further comprise instructions for using the kit to detect BDSF or STMST protein or nucleic
25 acid.

2. Prognostic Assays

The diagnostic methods described herein can furthermore be utilized to identify subjects having or at risk of developing a disease or disorder associated with aberrant
30 BDSF or STMST expression or activity. For example, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a

- 106 -

subject having or at risk of developing a disorder associated with BDSF protein, nucleic acid expression or activity such as a proliferative disorder (*e.g.*, cancer). Alternatively, the assays described herein, such as the preceding diagnostic assays or the following assays, can be utilized to identify a subject having or at risk of developing a disorder

5 associated with BDSF protein, nucleic acid expression or activity such as a proliferative disorder. Alternatively, the prognostic assays can be utilized to identify a subject having or at risk for developing a differentiative disorder. Thus, the present invention provides a method for identifying a disease or disorder associated with aberrant BDSF or STMST expression or activity in which a test sample is obtained from a subject and BDSF or

10 STMST protein or nucleic acid (*e.g.*, mRNA, genomic DNA) is detected, wherein the presence of BDSF or STMST protein or nucleic acid is diagnostic for a subject having or at risk of developing a disease or disorder associated with aberrant BDSF or STMST expression or activity. As used herein, a "test sample" refers to a biological sample obtained from a subject of interest. For example, a test sample can be a biological fluid

15 (*e.g.*, serum), cell sample, or tissue.

Furthermore, the prognostic assays described herein can be used to determine whether a subject can be administered an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate) to treat a disease or disorder associated with aberrant BDSF or STMST expression or

20 activity. For example, such methods can be used to determine whether a subject can be effectively treated with an agent for a proliferative disorder (*e.g.*, cancer), a differentiative disorder or an immune disorder. Thus, the present invention provides methods for determining whether a subject can be effectively treated with an agent for a disorder associated with aberrant BDSF or STMST expression or activity in which a test

25 sample is obtained and BDSF or STMST protein or nucleic acid expression or activity is detected (*e.g.*, wherein the abundance of BDSF or STMST protein or nucleic acid expression or activity is diagnostic for a subject that can be administered the agent to treat a disorder associated with aberrant BDSF or STMST expression or activity.)

The methods of the invention can also be used to detect genetic alterations in a

30 BDSF or STMST gene, thereby determining if a subject with the altered gene is at risk for a disorder characterized by an aberrant proliferative response. In preferred

- 107 -

embodiments, the methods include detecting, in a sample of cells from the subject, the presence or absence of a genetic alteration characterized by at least one of an alteration affecting the integrity of a gene encoding a BDSF or STMST-protein, or the mis-expression of the BDSF or STMST gene. For example, such genetic alterations can be
5 detected by ascertaining the existence of at least one of 1) a deletion of one or more nucleotides from a BDSF or STMST gene; 2) an addition of one or more nucleotides to a BDSF or STMST gene; 3) a substitution of one or more nucleotides of a BDSF or STMST gene, 4) a chromosomal rearrangement of a BDSF or STMST gene; 5) an alteration in the level of a messenger RNA transcript of a BDSF or STMST gene, 6)
10 aberrant modification of a BDSF or STMST gene, such as of the methylation pattern of the genomic DNA, 7) the presence of a non-wild type splicing pattern of a messenger RNA transcript of a BDSF or STMST gene, 8) a non-wild type level of a BDSF or STMST-protein, 9) allelic loss of a BDSF or STMST gene, and 10) inappropriate post-translational modification of a BDSF or STMST-protein. As described herein, there are
15 a large number of assay techniques known in the art which can be used for detecting alterations in a BDSF or STMST gene. A preferred biological sample is a tissue or serum sample isolated by conventional means from a subject.

In certain embodiments, detection of the alteration involves the use of a probe/primer in a polymerase chain reaction (PCR) (see, *e.g.*, U.S. Patent Nos.
20 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligation chain reaction (LCR) (see, *e.g.*, Landegran *et al.* (1988) *Science* 241:1077-1080; and Nakazawa *et al.* (1994) *PNAS* 91:360-364), the latter of which can be particularly useful for detecting point mutations in the BDSF or STMST-gene (see Abravaya *et al.* (1995) *Nucleic Acids Res.* 23:675-682). This method can include the steps of collecting
25 a sample of cells from a patient, isolating nucleic acid (*e.g.*, genomic, mRNA or both) from the cells of the sample, contacting the nucleic acid sample with one or more primers which specifically hybridize to a BDSF or STMST gene under conditions such that hybridization and amplification of the BDSF or STMST-gene (if present) occurs, and detecting the presence or absence of an amplification product, or detecting the size
30 of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification

- 108 -

step in conjunction with any of the techniques used for detecting mutations described herein.

Alternative amplification methods include: self sustained sequence replication (Guatelli, J.C. *et al.*, 1990, Proc. Natl. Acad. Sci. USA 87:1874-1878), transcriptional
5 amplification system (Kwoh, D.Y. *et al.*, 1989, Proc. Natl. Acad. Sci. USA 86:1173-1177), Q-Beta Replicase (Lizardi, P.M. *et al.*, 1988, Bio/Technology 6:1197), or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such
10 molecules are present in very low numbers.

In an alternative embodiment, mutations in a BDSF or STMST gene from a sample cell can be identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel
15 electrophoresis and compared. Differences in fragment length sizes between sample and control DNA indicates mutations in the sample DNA. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

20 In other embodiments, genetic mutations in BDSF can be identified by hybridizing a sample and control nucleic acids, *e.g.*, DNA or RNA, to high density arrays containing hundreds or thousands of oligonucleotides probes (Cronin, M.T. *et al.* (1996) *Human Mutation* 7: 244-255; Kozal, M.J. *et al.* (1996) *Nature Medicine* 2: 753-759). For example, genetic mutations in BDSF or STMST can be identified in two
25 dimensional arrays containing light-generated DNA probes as described in Cronin, M.T. *et al. supra*. Briefly, a first hybridization array of probes can be used to scan through long stretches of DNA in a sample and control to identify base changes between the sequences by making linear arrays of sequential overlapping probes. This step allows the identification of point mutations. This step is followed by a second hybridization array
30 that allows the characterization of specific mutations by using smaller, specialized probe arrays complementary to all variants or mutations detected. Each mutation array is

- 109 -

composed of parallel probe sets, one complementary to the wild-type gene and the other complementary to the mutant gene.

In yet another embodiment, any of a variety of sequencing reactions known in the art can be used to directly sequence the BDSF or STMST gene and detect mutations
5 by comparing the sequence of the sample BDSF or STMST with the corresponding wild-type (control) sequence. Examples of sequencing reactions include those based on techniques developed by Maxim and Gilbert ((1977) *PNAS* 74:560) or Sanger ((1977) *PNAS* 74:5463). It is also contemplated that any of a variety of automated sequencing procedures can be utilized when performing the diagnostic assays ((1995) *Biotechniques*
10 19:448), including sequencing by mass spectrometry (see, *e.g.*, PCT International Publication No. WO 94/16101; Cohen *et al.* (1996) *Adv. Chromatogr.* 36:127-162; and Griffin *et al.* (1993) *Appl. Biochem. Biotechnol.* 38:147-159).

Other methods for detecting mutations in the BDSF or STMST gene include methods in which protection from cleavage agents is used to detect mismatched bases in
15 RNA/RNA or RNA/DNA heteroduplexes (Myers *et al.* (1985) *Science* 230:1242). In general, the art technique of "mismatch cleavage" starts by providing heteroduplexes of formed by hybridizing (labeled) RNA or DNA containing the wild-type BDSF or STMST sequence with potentially mutant RNA or DNA obtained from a tissue sample. The double-stranded duplexes are treated with an agent which cleaves single-stranded
20 regions of the duplex such as which will exist due to basepair mismatches between the control and sample strands. For instance, RNA/DNA duplexes can be treated with RNase and DNA/DNA hybrids treated with S1 nuclease to enzymatically digesting the mismatched regions. In other embodiments, either DNA/DNA or RNA/DNA duplexes can be treated with hydroxylamine or osmium tetroxide and with piperidine in order to
25 digest mismatched regions. After digestion of the mismatched regions, the resulting material is then separated by size on denaturing polyacrylamide gels to determine the site of mutation. See, for example, Cotton *et al.* (1988) *Proc. Natl Acad Sci USA* 85:4397; Saleeba *et al.* (1992) *Methods Enzymol.* 217:286-295. In a preferred embodiment, the control DNA or RNA can be labeled for detection.

- 110 -

In still another embodiment, the mismatch cleavage reaction employs one or more proteins that recognize mismatched base pairs in double-stranded DNA (so called "DNA mismatch repair" enzymes) in defined systems for detecting and mapping point mutations in BDSF or STMST cDNAs obtained from samples of cells. For example, the
5 mutY enzyme of *E. coli* cleaves A at G/A mismatches and the thymidine DNA glycosylase from HeLa cells cleaves T at G/T mismatches (Hsu *et al.* (1994) *Carcinogenesis* 15:1657-1662). According to an exemplary embodiment, a probe based on a BDSF sequence, *e.g.*, a wild-type BDSF or STMST sequence, is hybridized to a cDNA or other DNA product from a test cell(s). The duplex is treated with a DNA
10 mismatch repair enzyme, and the cleavage products, if any, can be detected from electrophoresis protocols or the like. See, for example, U.S. Patent No. 5,459,039.

In other embodiments, alterations in electrophoretic mobility will be used to identify mutations in BDSF or STMST genes. For example, single strand conformation polymorphism (SSCP) may be used to detect differences in electrophoretic mobility
15 between mutant and wild type nucleic acids (Orita *et al.* (1989) *Proc Natl. Acad. Sci USA*: 86:2766, see also Cotton (1993) *Mutat Res* 285:125-144; and Hayashi (1992) *Genet Anal Tech Appl* 9:73-79). Single-stranded DNA fragments of sample and control BDSF or STMST nucleic acids will be denatured and allowed to renature. The secondary structure of single-stranded nucleic acids varies according to sequence, the
20 resulting alteration in electrophoretic mobility enables the detection of even a single base change. The DNA fragments may be labeled or detected with labeled probes. The sensitivity of the assay may be enhanced by using RNA (rather than DNA), in which the secondary structure is more sensitive to a change in sequence. In a preferred embodiment, the subject method utilizes heteroduplex analysis to separate double
25 stranded heteroduplex molecules on the basis of changes in electrophoretic mobility (Keen *et al.* (1991) *Trends Genet* 7:5).

In yet another embodiment, the movement of mutant or wild-type fragments in polyacrylamide gels containing a gradient of denaturant is assayed using denaturing gradient gel electrophoresis (DGGE) (Myers *et al.* (1985) *Nature* 313:495). When
30 DGGE is used as the method of analysis, DNA will be modified to insure that it does not completely denature, for example by adding a GC clamp of approximately 40 bp of

- 111 -

high-melting GC-rich DNA by PCR. In a further embodiment, a temperature gradient is used in place of a denaturing gradient to identify differences in the mobility of control and sample DNA (Rosenbaum and Reissner (1987) *Biophys Chem* 265:12753).

Examples of other techniques for detecting point mutations include, but are not limited to, selective oligonucleotide hybridization, selective amplification, or selective primer extension. For example, oligonucleotide primers may be prepared in which the known mutation is placed centrally and then hybridized to target DNA under conditions which permit hybridization only if a perfect match is found (Saiki *et al.* (1986) *Nature* 324:163); Saiki *et al.* (1989) *Proc. Natl Acad. Sci USA* 86:6230). Such allele specific oligonucleotides are hybridized to PCR amplified target DNA or a number of different mutations when the oligonucleotides are attached to the hybridizing membrane and hybridized with labeled target DNA.

Alternatively, allele specific amplification technology which depends on selective PCR amplification may be used in conjunction with the instant invention. Oligonucleotides used as primers for specific amplification may carry the mutation of interest in the center of the molecule (so that amplification depends on differential hybridization) (Gibbs *et al.* (1989) *Nucleic Acids Res.* 17:2437-2448) or at the extreme 3' end of one primer where, under appropriate conditions, mismatch can prevent, or reduce polymerase extension (Prossner (1993) *Tibtech* 11:238). In addition it may be desirable to introduce a novel restriction site in the region of the mutation to create cleavage-based detection (Gasparini *et al.* (1992) *Mol. Cell Probes* 6:1). It is anticipated that in certain embodiments amplification may also be performed using Taq ligase for amplification (Barany (1991) *Proc. Natl. Acad. Sci USA* 88:189). In such cases, ligation will occur only if there is a perfect match at the 3' end of the 5' sequence making it possible to detect the presence of a known mutation at a specific site by looking for the presence or absence of amplification.

The methods described herein may be performed, for example, by utilizing pre-packaged diagnostic kits comprising at least one probe nucleic acid or antibody reagent described herein, which may be conveniently used, *e.g.*, in clinical settings to diagnose patients exhibiting symptoms or family history of a disease or illness involving a BDSF or STMST gene.

Furthermore, any cell type or tissue in which BDSF is expressed may be utilized in the prognostic assays described herein.

3. Monitoring of Effects During Clinical Trials

5 Monitoring the influence of agents (*e.g.*, drugs, compounds) on the expression or activity of a BDSF or STMST protein (*e.g.*, modulation of angiogenesis or of an inflammatory response) can be applied not only in basic drug screening, but also in clinical trials. For example, the effectiveness of an agent determined by a screening assay as described herein to increase BDSF or STMST gene expression, protein levels, 10 or upregulate BDSF or STMST activity, can be monitored in clinical trials of subjects exhibiting decreased BDSF or STMST gene expression, protein levels, or downregulated BDSF or STMST activity. Alternatively, the effectiveness of an agent determined by a screening assay to decrease BDSF or STMST gene expression, protein levels, or downregulate BDSF or STMST activity, can be monitored in clinical trials of 15 subjects exhibiting increased BDSF or STMST gene expression, protein levels, or upregulated BDSF or STMST activity. In such clinical trials, the expression or activity of a BDSF or STMST gene, and preferably, other genes that have been implicated in, for example, a proliferative disorder can be used as a "read out" or markers of the phenotype of a particular cell.

20 For example, and not by way of limitation, genes, including BDSF or STMST, that are modulated in cells by treatment with an agent (*e.g.*, compound, drug or small molecule) which modulates BDSF or STMST activity (*e.g.*, identified in a screening assay as described herein) can be identified. Thus, to study the effect of agents on proliferative disorders, for example, in a clinical trial, cells can be isolated and RNA 25 prepared and analyzed for the levels of expression of BDSF or STMST and other genes implicated in the proliferative disorder, respectively. The levels of gene expression (*i.e.*, a gene expression pattern) can be quantified by Northern blot analysis or RT-PCR, as described herein, or alternatively by measuring the amount of protein produced, by one of the methods as described herein, or by measuring the levels of activity of BDSF or 30 STMST or other genes. In this way, the gene expression pattern can serve as a marker, indicative of the physiological response of the cells to the agent. Accordingly, this

- 113 -

response state may be determined before, and at various points during treatment of the individual with the agent.

In a preferred embodiment, the present invention provides a method for monitoring the effectiveness of treatment of a subject with an agent (*e.g.*, an agonist, antagonist, peptidomimetic, protein, peptide, nucleic acid, small molecule, or other drug candidate identified by the screening assays described herein) comprising the steps of (i) obtaining a pre-administration sample from a subject prior to administration of the agent; (ii) detecting the level of expression of a BDSF or STMST protein, mRNA, or genomic DNA in the preadministration sample; (iii) obtaining one or more post-administration samples from the subject; (iv) detecting the level of expression or activity of the BDSF or STMST protein, mRNA, or genomic DNA in the post-administration samples; (v) comparing the level of expression or activity of the BDSF or STMST protein, mRNA, or genomic DNA in the pre-administration sample with the BDSF or STMST protein, mRNA, or genomic DNA in the post administration sample or samples; and (vi) altering the administration of the agent to the subject accordingly. For example, increased administration of the agent may be desirable to increase the expression or activity of BDSF or STMST to higher levels than detected, *i.e.*, to increase the effectiveness of the agent. Alternatively, decreased administration of the agent may be desirable to decrease expression or activity of BDSF or STMST to lower levels than detected, *i.e.* to decrease the effectiveness of the agent. According to such an embodiment, BDSF expression or activity may be used as an indicator of the effectiveness of an agent, even in the absence of an observable phenotypic response.

C. Methods of Treatment:

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) a disorder or having a disorder associated with aberrant BDSF or STMST expression or activity. With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in

- 114 -

clinical development and on the market. More specifically, the term refers the study of how a patient's genes determine his or her response to a drug (*e.g.*, a patient's "drug response phenotype", or "drug response genotype".) Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the BDSF or STMST molecules of the present invention or BDSF or STMST modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

10

1. Prophylactic Methods

In one aspect, the invention provides a method for preventing in a subject, a disease or condition associated with an aberrant BDSF or STMST expression or activity, by administering to the subject a BDSF or STMST or an agent which modulates BDSF or STMST expression or at least one BDSF or STMST activity. Subjects at risk for a disease which is caused or contributed to by aberrant BDSF or STMST expression or activity can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the BDSF or STMST aberrancy, such that a disease or disorder is prevented or, alternatively, delayed in its progression. Depending on the type of BDSF or STMST aberrancy, for example, a BDSF or STMST, BDSF or STMST agonist or BDSF or STMST antagonist agent can be used for treating the subject. The appropriate agent can be determined based on screening assays described herein. The prophylactic methods of the present invention are further discussed in the following subsections.

25

2. Therapeutic Methods

Another aspect of the invention pertains to methods of modulating BDSF or STMST expression or activity for therapeutic purposes. Accordingly, in an exemplary embodiment, the modulatory method of the invention involves contacting a cell with a BDSF or STMST or agent that modulates one or more of the activities of BDSF or

30

- 115 -

STMST protein activity associated with the cell. An agent that modulates BDSF or STMST protein activity can be an agent as described herein, such as a nucleic acid or a protein, a naturally-occurring target molecule of a BDSF or STMST protein, a BDSF or STMST antibody, a BDSF or STMST agonist or antagonist, a peptidomimetic of a BDSF or STMST agonist or antagonist, or other small molecule. In one embodiment, the agent stimulates one or more BDSF or STMST activities. Examples of such stimulatory agents include active BDSF or STMST protein and a nucleic acid molecule encoding BDSF or STMST that has been introduced into the cell. In another embodiment, the agent inhibits one or more BDSF or STMST activities. Examples of such inhibitory agents include antisense BDSF or STMST nucleic acid molecules, anti-BDSF or STMST antibodies, and BDSF or STMST inhibitors. These modulatory methods can be performed *in vitro* (e.g., by culturing the cell with the agent) or, alternatively, *in vivo* (e.g., by administering the agent to a subject). As such, the present invention provides methods of treating an individual afflicted with a disease or disorder characterized by aberrant expression or activity of a BDSF or STMST protein or nucleic acid molecule. In one embodiment, the method involves administering an agent (e.g., an agent identified by a screening assay described herein), or combination of agents that modulates (e.g., upregulates or downregulates) BDSF or STMST expression or activity. In another embodiment, the method involves administering a BDSF or STMST protein or nucleic acid molecule as therapy to compensate for reduced or aberrant BDSF or STMST expression or activity.

Stimulation of BDSF activity is desirable in situations in which BDSF or STMST is abnormally downregulated and/or in which increased BDSF or STMST activity is likely to have a beneficial effect. For example, stimulation of BDSF or STMST activity is desirable in situations in which a BDSF or STMST is downregulated and/or in which increased BDSF or STMST activity is likely to have a beneficial effect. Likewise, inhibition of BDSF or STMST activity is desirable in situations in which BDSF or STMST is abnormally upregulated and/or in which decreased BDSF or STMST activity is likely to have a beneficial effect.

- 116 -

3. Pharmacogenomics

The BDSF or STMST molecules of the present invention, as well as agents, or modulators which have a stimulatory or inhibitory effect on BDSF or STMST activity (*e.g.*, BDSF or STMST gene expression) as identified by a screening assay described
5 herein can be administered to individuals to treat (prophylactically or therapeutically) disorders (*e.g.*, cancer) associated with aberrant BDSF or STMST activity. In conjunction with such treatment, pharmacogenomics (*i.e.*, the study of the relationship between an individual's genotype and that individual's response to a foreign compound or drug) may be considered. Differences in metabolism of therapeutics can lead to
10 severe toxicity or therapeutic failure by altering the relation between dose and blood concentration of the pharmacologically active drug. Thus, a physician or clinician may consider applying knowledge obtained in relevant pharmacogenomics studies in determining whether to administer a BDSF or STMST molecule or BDSF or STMST modulator as well as tailoring the dosage and/or therapeutic regimen of treatment with a
15 BDSF or STMST molecule or BDSF or STMST modulator.

Pharmacogenomics deals with clinically significant hereditary variations in the response to drugs due to altered drug disposition and abnormal action in affected persons. See *e.g.*, Eichelbaum, M., *Clin Exp Pharmacol Physiol*, 1996, 23(10-11) :983-985 and Linder, M.W., *Clin Chem*, 1997, 43(2):254-266. In general, two types of
20 pharmacogenetic conditions can be differentiated. Genetic conditions transmitted as a single factor altering the way drugs act on the body (altered drug action) or genetic conditions transmitted as single factors altering the way the body acts on drugs (altered drug metabolism). These pharmacogenetic conditions can occur either as rare genetic defects or as naturally-occurring polymorphisms. For example, glucose-6-phosphate
25 dehydrogenase deficiency (G6PD) is a common inherited enzymopathy in which the main clinical complication is haemolysis after ingestion of oxidant drugs (anti-malarials, sulfonamides, analgesics, nitrofurans) and consumption of fava beans.

One pharmacogenomics approach to identifying genes that predict drug response, known as "a genome-wide association", relies primarily on a high-resolution
30 map of the human genome consisting of already known gene-related markers (*e.g.*, a "bi-allelic" gene marker map which consists of 60,000-100,000 polymorphic or variable

- 117 -

sites on the human genome, each of which has two variants.) Such a high-resolution genetic map can be compared to a map of the genome of each of a statistically significant number of patients taking part in a Phase II/III drug trial to identify markers associated with a particular observed drug response or side effect. Alternatively, such a high resolution map can be generated from a combination of some ten-million known single nucleotide polymorphisms (SNPs) in the human genome. As used herein, a "SNP" is a common alteration that occurs in a single nucleotide base in a stretch of DNA. For example, a SNP may occur once per every 1000 bases of DNA. A SNP may be involved in a disease process, however, the vast majority may not be disease-associated. Given a genetic map based on the occurrence of such SNPs, individuals can be grouped into genetic categories depending on a particular pattern of SNPs in their individual genome. In such a manner, treatment regimens can be tailored to groups of genetically similar individuals, taking into account traits that may be common among such genetically similar individuals.

Alternatively, a method termed the "candidate gene approach", can be utilized to identify genes that predict drug response. According to this method, if a gene that encodes a drug's target is known (*e.g.*, a BDSF or STMST protein or a BDSF receptor of the present invention), all common variants of that gene can be fairly easily identified in the population and it can be determined if having one version of the gene versus another is associated with a particular drug response.

As an illustrative embodiment, the activity of drug metabolizing enzymes is a major determinant of both the intensity and duration of drug action. The discovery of genetic polymorphisms of drug metabolizing enzymes (*e.g.*, N-acetyltransferase 2 (NAT 2) and cytochrome P450 enzymes CYP2D6 and CYP2C19) has provided an explanation as to why some patients do not obtain the expected drug effects or show exaggerated drug response and serious toxicity after taking the standard and safe dose of a drug. These polymorphisms are expressed in two phenotypes in the population, the extensive metabolizer (EM) and poor metabolizer (PM). The prevalence of PM is different among different populations. For example, the gene coding for CYP2D6 is highly polymorphic and several mutations have been identified in PM, which all lead to the absence of functional CYP2D6. Poor metabolizers of CYP2D6 and CYP2C19 quite frequently

- 118 -

experience exaggerated drug response and side effects when they receive standard doses. If a metabolite is the active therapeutic moiety, PM show no therapeutic response, as demonstrated for the analgesic effect of codeine mediated by its CYP2D6-formed metabolite morphine. The other extreme are the so called ultra-rapid metabolizers who
5 do not respond to standard doses. Recently, the molecular basis of ultra-rapid metabolism has been identified to be due to CYP2D6 gene amplification.

Alternatively, a method termed the "gene expression profiling", can be utilized to identify genes that predict drug response. For example, the gene expression of an animal dosed with a drug (*e.g.*, a BDSF or STMST molecule or BDSF or STMST
10 modulator of the present invention) can give an indication whether gene pathways related to toxicity have been turned on.

Information generated from more than one of the above pharmacogenomics approaches can be used to determine appropriate dosage and treatment regimens for prophylactic or therapeutic treatment an individual. This knowledge, when applied to
15 dosing or drug selection, can avoid adverse reactions or therapeutic failure and thus enhance therapeutic or prophylactic efficiency when treating a subject with a BDSF or STMST molecule or BDSF modulator, such as a modulator identified by one of the exemplary screening assays described herein.

This invention is further illustrated by the following examples which should not
20 be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference. References throughout the instant specification to websites maintained as part of the World Wide Web are referred to herein by the prefix <http://>. The information contained in such websites is publically-avialable and can be accessed elctronically by contacting
25 the cited address.

EXAMPLES

The novel genes exemplified herein were isolated utilizing a methodology which takes advantage of the fact that a majority of secreted and membrane-associated
30 proteins posess at their amino termini a signal sequence, as defined herein. The methodology identifies such secreted and/or membrane-associated proteins by virtue of

- 119 -

their ability to direct export of a reporter protein, alkaline phosphatase (AP), from mammalian cells. The above described methodology, referred to herein as "signal trapping" or "signal sequence trapping" is described in detail in PCT/US97/20201 (WO98/22491 published May 28, 1998), the content of which is incorporated herein in its entirety. The present methodology is further combined with an improved method for cDNA library construction in which directional random primed cDNA libraries are prepared.

Example 1: Identification and Characterization of Human and Murine BDSF

cDNA

In this example, the identification and characterization of the genes encoding human BDSF-1 (also referred to as "TANGO 122" and "hT122") and murine BDSF-1 (also referred to as "mT122") are described.

Isolation of the human BDSF cDNA

The invention is based, at least in part, on the discovery of a human gene encoding a novel brain-derived signaling factor protein, referred to herein as BDSF.

The methodology used to isolate the human BDSF-1 gene takes advantage of the fact that molecules such as BDSF-1 have an amino terminal signal sequence which directs certain secreted and membrane-bound proteins through the cellular secretory apparatus.

The human BDSF-1 mRNA was identified by screening of a human fetal brain cDNA library. This library was prepared using mRNA purchased from Clontech, Palo Alto (Cat. no, 6573-1). A signal trap cDNA library was prepared by ligating random primed double stranded cDNA into the expression vector, ptrAP1, resulting in fusions of cDNAs to the reporter, alkaline phosphatase (AP). DNAs from individual clones from this library were prepared by standard techniques and transfected into human embryonic kidney fibroblasts (293T cells). After 48 hours, cell supernatants were collected and assayed for AP activity. Clones giving rise to detectable AP activity in the supernatants of transfected cells were analyzed further by DNA sequencing and the novel clones subjected to further DNA sequencing.

- 120 -

The nucleotide sequence encoding the human BDSF-1 protein is shown in Figure 1 and is set forth as SEQ ID NO: 1. The full length protein encoded by this nucleic acid is comprised of about 244 amino acids and has the amino acid sequence shown in Figure 1 and set forth as SEQ ID NO:2. The coding portion (open reading frame) of SEQ ID NO:1 is set forth as SEQ ID NO:3. Notable features of the human BDSF-1 protein include a signal peptide (about amino acids 1-25 of SEQ ID NO:2), an Ig-like domain (about amino acids 41-129 of SEQ ID NO:2) and two conserved cysteine residues (about amino acids 48 and 127 of SEQ ID NO:2). A clone, comprising the entire coding region of human BDSF-1 has been deposited with the American Type Culture Collection (ATCC), Manassas, Virginia, on May 15, 1998 and assigned Accession Number 98756.

Isolation of the murine BDSF cDNA

The murine BDSF-1 gene was identified from a murine choroid plexus cDNA library. More specifically, a murine choroid plexus cDNA library was plated out and colonies picked into 96 well plates. The colonies were cultured, plasmids were prepared from each well, and several of the inserts were sequenced. The nucleotide sequences were compared against the human BDSF-1 nucleotide sequence. Upon review of the results from this sequence comparison, a murine BDSF-1 gene was obtained.

The nucleotide sequence encoding the murine BDSF-1 protein is shown in Figure 2 and is set forth as SEQ ID NO: 6. The full length protein encoded by this nucleic acid is comprised of about 251 amino acids and has the amino acid sequence shown in Figure 2 and set forth as SEQ ID NO:7. The coding portion (open reading frame) of SEQ ID NO:6 is set forth as SEQ ID NO:8. Notable features of the murine BDSF-1 protein include a signal peptide (about amino acids 1-24 of SEQ ID NO:7), an Ig-like domain (about amino acids 40-128 of SEQ ID NO:7) and two conserved cysteine residues (about amino acids 47 and 126 of SEQ ID NO:7).

Analysis of Human and Murine BDSF

A BLAST search (Altschul *et al.* (1990) J. Mol. Biol. 215:403) of the nucleotide and protein sequences of human BDSF and murine BDSF has revealed that BDSF does not display significant homology to other proteins except for distant similarity to other

- 121 -

Ig-like domain containing proteins. An alignment of human BDSF and murine BDSF, however, indicated that these proteins are 77.4% identical overall and 90.5% identical over the first 211 amino acids. The alignment of human BDSF-1 and murine BDSF-1 is presented in Figure 3. The alignment of human BDSF-1 and murine BDSF-1 was
5 performed using the ALIGN program. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 was used.

Expression of human BDSF

10 The expression of human BDSF was analyzed using Northern blot hybridization and a probe specific for human BDSF. The DNA was radioactively labeled with ³²P-dCTP using the Prime-It-kit (Stratagene, La Jolla, CA) according to the instructions of the supplier. Filters containing human mRNA (Multi-Tissue Northern I, Multi-Tissue Northern II and Multi-Tissue Northern III from Clontech, Palo Alto, CA) were probed in
15 ExpressHyb hybridization solution (Clontech) and washed at high stringency according to manufacturer's recommendations. In addition, filters containing human brain mRNA (Brain-Subregion Blot from Clontech) were also probed for human BDSF-1 expression.

Results of Northern blot hybridization indicate that human BDSF is expressed as an approximately 5.0 kilobase transcript in all tissues (spleen, lymph node, thymus,
20 peripheral blood leukocytes, bone marrow, stomach, thyroid, spinal cord, trachea, adrenal, testis, small intestine, heart, placenta, lung, liver, kidney and pancreas). BDSF mRNA expression was also observed in human fetal liver.

The highest level of human BDSF expression was found in the adult brain where the pattern of transcripts was also different than for the other tissues. In the brain, major
25 transcripts of 2.6 kb, 3.2 kb and 6.5 kb were observed. This pattern of BDSF mRNA transcripts in the brain is found in the sub-regions of the brain including amygdala, caudate nucleus, hippocampus, substantia nigra, sub-thalamate nucleus and thalamus, but not in corpus callosum.

Example 2: Expression of Recombinant BDSF Protein in Bacterial Cells

BDSF is expressed as a recombinant glutathione-S-transferase (GST) fusion polypeptide in *E. coli* and the fusion polypeptide isolated and characterized. Specifically, human or murine BDSF is fused to GST and this fusion polypeptide is
5 expressed in *E. coli*, e.g., strain PEB199. Expression of the GST-BDSF fusion protein in PEB199 is induced with IPTG. The recombinant fusion polypeptide is purified from crude bacterial lysates of the induced PEB199 strain by affinity chromatography on glutathione beads. Using polyacrylamide gel electrophoretic analysis of the polypeptide purified from the bacterial lysates, the molecular weight of the resultant
10 fusion polypeptide is determined.

Example 3: Expression of Recombinant BDSF Protein in COS Cells

To express the human or murine BDSF gene in COS cells, the pcDNA/Amp vector by Invitrogen Corporation (San Diego, CA) is used. This vector contains an
15 SV40 origin of replication, an ampicillin resistance gene, an *E. coli* replication origin, a CMV promoter followed by a polylinker region, and an SV40 intron and polyadenylation site. A DNA fragment encoding the entire BDSF protein and a HA tag (Wilson *et al.* (1984) *Cell* 37:767) fused in-frame to its 3' end of the fragment is cloned into the polylinker region of the vector, thereby placing the expression of the
20 recombinant protein under the control of the CMV promoter.

To construct the plasmid, the BDSF DNA sequence is amplified by PCR using two primers. The 5' primer contains the restriction site of interest followed by approximately twenty nucleotides of the BDSF coding sequence starting from the initiation codon; the 3' end sequence contains complementary sequences to the other
25 restriction site of interest, a translation stop codon, the HA tag and the last 20 nucleotides of the BDSF coding sequence. The PCR amplified fragment and the pCDNA/Amp vector are digested with the appropriate restriction enzymes and the vector is dephosphorylated using the CIAP enzyme (New England Biolabs, Beverly, MA). Preferably the two restriction sites chosen are different so that the BDSF gene is
30 inserted in the correct orientation. The ligation mixture is transformed into *E. coli* cells (strains HB101, DH5a, SURE, available from Stratagene Cloning Systems, La Jolla,

- 123 -

CA, can be used), the transformed culture is plated on ampicillin media plates, and resistant colonies are selected. Plasmid DNA is isolated from transformants and examined by restriction analysis for the presence of the correct fragment.

COS cells are subsequently transfected with the BDSF-pcDNA/Amp plasmid DNA using the calcium phosphate or calcium chloride co-precipitation methods, DEAE-dextran-mediated transfection, lipofection, or electroporation. Other suitable methods for transfecting host cells can be found in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989. The expression of the BDSF protein is detected by radiolabelling (^{35}S -methionine or ^{35}S -cysteine available from NEN, Boston, MA, can be used) and immunoprecipitation (Harlow, E. and Lane, D. *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1988) using an HA specific monoclonal antibody. Briefly, the cells are labeled for 8 hours with ^{35}S -methionine (or ^{35}S -cysteine). The culture media are then collected and the cells are lysed using detergents (RIPA buffer, 150 mM NaCl, 1% NP-40, 0.1% SDS, 0.5% DOC, 50 mM Tris, pH 7.5). Both the cell lysate and the culture media are precipitated with an HA specific monoclonal antibody. Precipitated proteins are then analyzed by SDS-PAGE.

Alternatively, DNA containing the BDSF coding sequence is cloned directly into the polylinker of the pCDNA/Amp vector using the appropriate restriction sites. The resulting plasmid is transfected into COS cells in the manner described above, and the expression of the BDSF protein is detected by radiolabelling and immunoprecipitation using a BDSF specific monoclonal antibody

25 **Example 4: Retroviral Delivery of BDSF**

The entire open reading frame of BDSF is subcloned into the retroviral vector MSCVneo, described in Hawley *et al.* (1994) *Gene Therapy* 1:136-138. Cells (293Ebna, Invitrogen) are then transiently transfected with the BDSF construct and with constructs containing viral regulatory elements, to produce high titre retrovirus containing the BDSF gene. The virus is then used to transfect mice. These mice are

- 124 -

then tested for any gross pathology and for changes in biological response, *e.g.*, cell proliferation and/or differentiation, using standard assays.

Example 5: Chromosomal Location of BDSF

5 Human BDSF maps to hu7p12-14 between markers WI-967 and WI-4253 close to the CMT2D (Charcot-Marie-Tooth neuropathy) locus. The syntenic chromosome in mouse, mol1, is in close proximity with the mouse known genes: egfr (epidermal growth factor receptor), ddc (dopa decarboxylase) and cob1 (cordon bleu). BDSF also mapped close to the following human genes: ADCYAP1R1 (adenylate cyclase activating
10 polypeptide 1), AMPH (amphiphysin), BLVRA (biliverdin reductase A), OGDH (oxoglutarate dehydrogenase), OCM (oncomodulin) and EGFR (epidermal growth factor receptor).

Example 6: *In situ* Expression of BDSF

15 Mouse tissues were analysed *in situ* for expression of BDSF. The *in situ* data (a 6 day film exposure of coronal brain sections from adult mouse brain) demonstrated high punctate expression in the cortex, hypothalamus, hippocampus, and hind brain, indicating that a subset of cells within these regions is positive. Expression in the hippocampus includes, but is not restricted to granule cells. Expression in the
20 cerebellum is limited to purkinje cells. Embryonic sagittal sections (day 14.5 - 15.5) had expression in developing brain and spinal cord. For embryonic mouse expression (embryonic sagittal sections, day 14.5 and 15.5), signal was detected in the ependyma of the brain and the primordium of the lower incisor tooth.

25 **Example 7: Identification And Characterization of STMST-1 cDNAs**

In this example, the identification and characterization of the genes encoding human STMST-1 and STMST-2 (also referred to as "TANGO123a" (clone Fmhu123a) and "TANGO 123c" (clone Fmhu123c), respectively) is described.

Isolation of the human STMST cDNAs

The 'signal sequence trapping' method described above was utilized to analyse the sequences of several cDNAs of a cDNA library derived from bronchial epithelial cells which had been stimulated with the cytokine, TNF α . This analysis identified a
5 partial human clone having an insert of approximately 231 kb containing a protein-encoding sequence of approximately 225 nucleotides capable of encoding approximately 75 amino acids of STMST (*e.g.*, the start met through residue 74 of, for example, SEQ ID NO:15). This cDNA was used to re-screen the library. Two full-length cDNA clones were isolated. Sequencing of these clones revealed the nucleotide sequences of
10 human STMST-1 and STMST-2.

The nucleotide sequence encoding the human STMST-1 protein is shown in Figure 4 and is set forth as SEQ ID NO:14. The full length protein encoded by this nucleic acid is comprised of about 297 amino acids and has the amino acid sequence shown in Figure 4 and set forth as SEQ ID NO:15. The coding portion (open reading
15 frame) of SEQ ID NO:14 is set forth as SEQ ID NO:16.

The nucleotide sequence encoding the human STMST-2 protein is shown in Figure 5 and is set forth as SEQ ID NO:17. The full length protein encoded by this nucleic acid is comprised of about 609 amino acids and has the amino acid sequence shown in Figure 5 and set forth as SEQ ID NO:18. The coding portion (open reading
20 frame) of SEQ ID NO:17 is set forth as SEQ ID NO:19.

Analysis of Human STMST-1 and STMST-2

Examination of the cDNA sequences depicted in Figures 3 and 4 showed that they were likely encoded by alternatively spliced mRNAs derived from the same gene.
25 Thus, the amino acid sequence of STMST-1 diverges from that of STMST-2 at about amino acid residue 263 of SEQ ID NO:15 or SEQ ID NO:18. The amino acid sequence of STMST-1 lacks the extensive cytoplasmic domain of STMST-2.

A BLAST search (Altschul *et al.* (1990) J. Mol. Biol. 215:403) of the nucleotide sequence of human STMST-2 has revealed that STMST-2 is significantly similar to a
30 protein identified as protein A-2 (human A-2, Accession No. U47928; murine A-2, Accession No. AC002393) which were sequenced as part of the sequencing of human

- 126 -

chromosome 12p13 and mouse chromosome 6, respectively. The human A-2 protein appears to be one of a family of alternatively-spliced gene products which further includes protein A-1 (Accession No. U47925) as well as A-3 (Accession No. U47929). The A-2 proteins, like the STMST proteins of the present invention, include many
5 features indicative of the G protein-coupled receptor family of proteins.

For instance, the STMSTs of the present invention contain conserved cysteines found in the first 2 extracellular loops (prior to the third and fifth transmembrane domains) of most GPCRs (cys 83 and cys 161 of SEQ ID NO:15 or SEQ ID NO:18). A highly conserved asparagine residue in the first transmembrane domain is present (asn25
10 in SEQ ID NO:15 or SEQ ID NO:18). Transmembrane domain two of the STMST proteins contains a highly conserved leucine (leu49 of SEQ ID NO:15 or SEQ ID NO:18). The two cysteine residues are believed to form a disulfide bond that stabilizes the functional protein structure. A highly conserved tryptophan and proline in the fourth transmembrane domain of the STMST proteins is present (trp135 and pro 145 of SEQ
15 ID NO:15 or SEQ ID NO:18). The third cytoplasmic loop contains 49 amino acid residues and is thus the longest cytoplasmic loop of the three, characteristic of G protein coupled receptors. Moreover, a highly conserved proline in the sixth transmembrane domain is present (pro260 of SEQ ID NO:15 and SEQ ID NO:18). The proline residues in the fourth, fifth, sixth, and seventh transmembrane domains are thought to introduce
20 kinks in the alpha-helices and may be important in the formation of the ligand binding pocket. Furthermore, the conserved (in the second cytoplasmic loop) HRM motif found in almost all Rhodopsin family GPCRs is present in the STMST proteins of the instant invention (his107, arg108, met109 of SEQ ID NO:15 or SEQ ID NO:18). (The arginine of the HRM sequence is thought to be the most important amino acid in GPCRs and is
25 invariant). Moreover, an almost invariant proline is present in the seventh transmembrane domain of STMST-2 (pro294 of SEQ ID NO:18).

As such, the STMST family of proteins, like the A-2 family of proteins, are referred to herein as G protein-coupled receptor-like proteins.

STMST-1 is also predicted to contain the following sites: cAMP and cGMP-
30 dependent protein kinase phosphorylation site at aa 225-228 (KRRS); Protein kinase C phosphorylation sites at aa 153-155 (SER) and at aa 290-292 (SSR); Casein kinase II

- 127 -

phosphorylation sites at aa 228-231 (SSID) and at aa 291-294 (SRQD); N-myristoylation sites at aa 9-14 (GSAVGW), aa 169-174 (GLGFGV), aa 181-186 (GGSVAM), aa 187-192 (GVICTA), aa 232-237 (GSEPAK), and at aa 244-249 (GLVTTI); Amidation site at aa 223-226 (QGKR).

- 5 Likewise, STMST is predicted to contain the following sites: cAMP- and cGMP-dependent protein kinase phosphorylation sites at aa 225-228 (KRRS), aa 393-396 (RRFS), aa 436-439 (RRAS), and at aa 453-456 (RRRS); Protein kinase C phosphorylation sites at aa 253-255 (SER), aa 268-270 (SLR), aa 392-394 (TRR), aa 462-464 (SLR), aa 482-484 (SPR), and at aa 560-562 (SLR); Casein kinase II
- 10 phosphorylation sites at aa 228-231 (SSID), aa 324-327 (SDDE), aa 328-331 (TSLE), aa 364-367 (SALE), aa 396-399 (SHDD), aa 417-420 (SGED), aa 466-469 (SALD), aa 506-509 (TAFE), aa 568-571 (SWGE), and at aa 590-593 (SPSE); Tyrosine kinase phosphorylation site at aa 342-348 (RSLDYG Y); N-myristoylation sites at aa 9-14 (GSAVGW), aa 169-174 (GLGFGV), aa 181-186 (GGSVAM), aa 187-192 (GVICTA),
- 15 aa 232-237 (GSEPAK), aa 244-249 (GLVTTI), aa 531-536 (GADPGE), aa 564-569 (GLSASW), aa 573-578 (GGLRAA), and at aa 579-584 (GGGGST); Amidation site at aa 223-226 (QGKR).

Tissue Distribution of STMST-1 mRNA

- 20 This Example describes the tissue distribution of STMST mRNA, as determined by Northern blot hybridization.

Northern blot hybridizations with the various RNA samples were performed (Clontech Multi-tissue Northern I and human fetal tissue northern) under standard conditions and washed under stringent conditions. A 4.5 Kb mRNA transcript was

25 detected in heart, brain, placenta, lung, liver, skeletal muscle, fetal brain, fetal lung, and fetal kidney. Expression was highest in fetal brain.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following

5 claims.

-129-

SEQUENCE LISTING

<110> Millennium Biotherapeutics, Inc.

<120> NOVEL SECRETED AND MEMBRANE-ASSOCIATED PROTEINS AND
USES THEREFOR

<130> PAT 48023W-1

<140> PCT/US99/11842

<141> 1999-05-28

<150> USSN 09/086,892

<151> 1998-05-29

<150> USSN 09/087,121

<151> 1998-05-29

<160> 30

<170> PatentIn Ver. 2.0

<210> 1

<211> 1119

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (140)..(871)

<400> 1

gtcgacccac gcg tccggca ggatgtttgc agtgtcgcgc ccagggctct gagactgagc 60

ctgccatcca ctgcacgcc tttctttcag ggcttttcgg ctgttggtta cactgatgtg 120

acccccctcc cttttttgga atg atg ggg atc ttt ttg gtg tat gtt gga ttt 172

Met Met Gly Ile Phe Leu Val Tyr Val Gly Phe
1 5 10

gtt ttc ttt tcc gtt tta tat gta caa caa ggg ctt tct tct caa gca 220

Val Phe Phe Ser Val Leu Tyr Val Gln Gln Gly Leu Ser Ser Gln Ala
15 20 25

aaa ttt acc gag ttt ccg cgg aac gtg acg gcg acc gag ggg cag aat 268

Lys Phe Thr Glu Phe Pro Arg Asn Val Thr Ala Thr Glu Gly Gln Asn
30 35 40

gtg gag atg tcc tgc gcc ttc cag agc ggc tcc gcc tcg gtg tat ctg 316

Val Glu Met Ser Cys Ala Phe Gln Ser Gly Ser Ala Ser Val Tyr Leu
45 50 55

-130-

gag atc caa tgg tgg ttc ctg cgg ggg ccg gag gac ctg gat ccc ggg	364
Glu Ile Gln Trp Trp Phe Leu Arg Gly Pro Glu Asp Leu Asp Pro Gly	
60 65 70 75	
gcc gag ggg gcc ggc gcg cag gtg gag ctc ttg ccc gac aga gac ccg	412
Ala Glu Gly Ala Gly Ala Gln Val Glu Leu Leu Pro Asp Arg Asp Pro	
80 85 90	
gac agc gac ggg acc aag atc agc aca gtg aaa gtc caa ggc aat gac	460
Asp Ser Asp Gly Thr Lys Ile Ser Thr Val Lys Val Gln Gly Asn Asp	
95 100 105	
atc tcc cac aag ctt cag att tcc aaa gtg agg aaa aag gat gaa ggc	508
Ile Ser His Lys Leu Gln Ile Ser Lys Val Arg Lys Lys Asp Glu Gly	
110 115 120	
tta tat gag tgc agg gtg act gat gcc aac tac ggg gag ctt cag gaa	556
Leu Tyr Glu Cys Arg Val Thr Asp Ala Asn Tyr Gly Glu Leu Gln Glu	
125 130 135	
cac aag gcc cag gcc tat ctg aaa gtc aat gcc aac agc cat gcc cgc	604
His Lys Ala Gln Ala Tyr Leu Lys Val Asn Ala Asn Ser His Ala Arg	
140 145 150 155	
aga atg cag gcc ttc gaa gcc tcg ccc atg tgg ctg cag gat atg aag	652
Arg Met Gln Ala Phe Glu Ala Ser Pro Met Trp Leu Gln Asp Met Lys	
160 165 170	
ccc cgc aag aac gtc tcc gca gcc atc ccc agc agc atc cat ggc tct	700
Pro Arg Lys Asn Val Ser Ala Ala Ile Pro Ser Ser Ile His Gly Ser	
175 180 185	
gcc aac caa cga acg cac tcc acc tcc agc cct caa gtg gta gcc aaa	748
Ala Asn Gln Arg Thr His Ser Thr Ser Ser Pro Gln Val Val Ala Lys	
190 195 200	
atc ccc aaa caa agt cca caa tca ggt atg gaa acc cat ttc gag cct	796
Ile Pro Lys Gln Ser Pro Gln Ser Gly Met Glu Thr His Phe Glu Pro	
205 210 215	
ttt att tta cca ctc aca aac gct cca cag aaa ggt cag tcg tat aga	844
Phe Ile Leu Pro Leu Thr Asn Ala Pro Gln Lys Gly Gln Ser Tyr Arg	
220 225 230 235	
gta gac aga ttt atg aat ggt gat ttt taaaatcgga gacctagtct	891
Val Asp Arg Phe Met Asn Gly Asp Phe	
240	
agtgcaagtg attatgagag gtgagcactg agcctgcacc aattcactca gagctcaaag	951
catgtgggtg caccctgtca gtcccctagt ggtgcttcat ttccagggca tctgagagct	1011
ggactctggt ttttatacctt tctgtattta cacattataa gaacaataaa tcatgtaatg	1071
ttggttacat tacaaaaaaaa aaaaaaaaaa aaaaaaaagg gcggccgc	1119

-131-

<210> 2

<211> 244

<212> PRT

<213> Homo sapiens

<400> 2

Met	Met	Gly	Ile	Phe	Leu	Val	Tyr	Val	Gly	Phe	Val	Phe	Phe	Ser	Val
1				5					10					15	

Leu	Tyr	Val	Gln	Gln	Gly	Leu	Ser	Ser	Gln	Ala	Lys	Phe	Thr	Glu	Phe
			20					25					30		

Pro	Arg	Asn	Val	Thr	Ala	Thr	Glu	Gly	Gln	Asn	Val	Glu	Met	Ser	Cys
		35					40					45			

Ala	Phe	Gln	Ser	Gly	Ser	Ala	Ser	Val	Tyr	Leu	Glu	Ile	Gln	Trp	Trp
	50					55					60				

Phe	Leu	Arg	Gly	Pro	Glu	Asp	Leu	Asp	Pro	Gly	Ala	Glu	Gly	Ala	Gly
65					70					75					80

Ala	Gln	Val	Glu	Leu	Leu	Pro	Asp	Arg	Asp	Pro	Asp	Ser	Asp	Gly	Thr
				85					90					95	

Lys	Ile	Ser	Thr	Val	Lys	Val	Gln	Gly	Asn	Asp	Ile	Ser	His	Lys	Leu
			100					105					110		

Gln	Ile	Ser	Lys	Val	Arg	Lys	Lys	Asp	Glu	Gly	Leu	Tyr	Glu	Cys	Arg
		115					120					125			

Val	Thr	Asp	Ala	Asn	Tyr	Gly	Glu	Leu	Gln	Glu	His	Lys	Ala	Gln	Ala
	130					135					140				

Tyr	Leu	Lys	Val	Asn	Ala	Asn	Ser	His	Ala	Arg	Arg	Met	Gln	Ala	Phe
145					150					155					160

Glu	Ala	Ser	Pro	Met	Trp	Leu	Gln	Asp	Met	Lys	Pro	Arg	Lys	Asn	Val
				165					170					175	

Ser	Ala	Ala	Ile	Pro	Ser	Ser	Ile	His	Gly	Ser	Ala	Asn	Gln	Arg	Thr
			180					185					190		

His	Ser	Thr	Ser	Ser	Pro	Gln	Val	Val	Ala	Lys	Ile	Pro	Lys	Gln	Ser
		195					200					205			

Pro	Gln	Ser	Gly	Met	Glu	Thr	His	Phe	Glu	Pro	Phe	Ile	Leu	Pro	Leu
	210					215					220				

Thr	Asn	Ala	Pro	Gln	Lys	Gly	Gln	Ser	Tyr	Arg	Val	Asp	Arg	Phe	Met
225					230					235					240

Asn Gly Asp Phe

-132-

<210> 3
 <211> 732
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(732)

<400> 3

atg atg ggg atc ttt ttg gtg tat gtt gga ttt gtt ttc ttt tcc gtt	48
Met Met Gly Ile Phe Leu Val Tyr Val Gly Phe Val Phe Phe Ser Val	
1 5 10 15	
tta tat gta caa caa ggg ctt tct tct caa gca aaa ttt acc gag ttt	96
Leu Tyr Val Gln Gln Gly Leu Ser Ser Gln Ala Lys Phe Thr Glu Phe	
20 25 30	
ccg cgg aac gtg acg gcg acc gag ggg cag aat gtg gag atg tcc tgc	144
Pro Arg Asn Val Thr Ala Thr Glu Gly Gln Asn Val Glu Met Ser Cys	
35 40 45	
gcc ttc cag agc ggc tcc gcc tcg gtg tat ctg gag atc caa tgg tgg	192
Ala Phe Gln Ser Gly Ser Ala Ser Val Tyr Leu Glu Ile Gln Trp Trp	
50 55 60	
ttc ctg cgg ggg ccg gag gac ctg gat ccc ggg gcc gag ggg gcc ggc	240
Phe Leu Arg Gly Pro Glu Asp Leu Asp Pro Gly Ala Glu Gly Ala Gly	
65 70 75 80	
gcg cag gtg gag ctc ttg ccc gac aga gac ccg gac agc gac ggg acc	288
Ala Gln Val Glu Leu Leu Pro Asp Arg Asp Pro Asp Ser Asp Gly Thr	
85 90 95	
aag atc agc aca gtg aaa gtc caa ggc aat gac atc tcc cac aag ctt	336
Lys Ile Ser Thr Val Lys Val Gln Gly Asn Asp Ile Ser His Lys Leu	
100 105 110	
cag att tcc aaa gtg agg aaa aag gat gaa ggc tta tat gag tgc agg	384
Gln Ile Ser Lys Val Arg Lys Lys Asp Glu Gly Leu Tyr Glu Cys Arg	
115 120 125	
gtg act gat gcc aac tac ggg gag ctt cag gaa cac aag gcc cag gcc	432
Val Thr Asp Ala Asn Tyr Gly Glu Leu Gln Glu His Lys Ala Gln Ala	
130 135 140	
tat ctg aaa gtc aat gcc aac agc cat gcc cgc aga atg cag gcc ttc	480
Tyr Leu Lys Val Asn Ala Asn Ser His Ala Arg Arg Met Gln Ala Phe	
145 150 155 160	
gaa gcc tcg ccc atg tgg ctg cag gat atg aag ccc cgc aag aac gtc	528
Glu Ala Ser Pro Met Trp Leu Gln Asp Met Lys Pro Arg Lys Asn Val	
165 170 175	

-133-

tcc gca gcc atc ccc agc agc atc cat ggc tct gcc aac caa cga acg	576
Ser Ala Ala Ile Pro Ser Ser Ile His Gly Ser Ala Asn Gln Arg Thr	
180 185 190	

cac tcc acc tcc agc cct caa gtg gta gcc aaa atc ccc aaa caa agt	624
His Ser Thr Ser Ser Pro Gln Val Val Ala Lys Ile Pro Lys Gln Ser	
195 200 205	

cca caa tca ggt atg gaa acc cat ttc gag cct ttt att tta cca ctc	672
Pro Gln Ser Gly Met Glu Thr His Phe Glu Pro Phe Ile Leu Pro Leu	
210 215 220	

aca aac gct cca cag aaa ggt cag tcg tat aga gta gac aga ttt atg	720
Thr Asn Ala Pro Gln Lys Gly Gln Ser Tyr Arg Val Asp Arg Phe Met	
225 230 235 240	

aat ggt gat ttt	732
Asn Gly Asp Phe	

<210> 4
 <211> 660
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(657)

<400> 4

caa gca aaa ttt acc gag ttt ccg cgg aac gtg acg gcg acc gag ggg	48
Gln Ala Lys Phe Thr Glu Phe Pro Arg Asn Val Thr Ala Thr Glu Gly	
1 5 10 15	

cag aat gtg gag atg tcc tgc gcc ttc cag agc ggc tcc gcc tcg gtg	96
Gln Asn Val Glu Met Ser Cys Ala Phe Gln Ser Gly Ser Ala Ser Val	
20 25 30	

tat ctg gag atc caa tgg tgg ttc ctg cgg ggg ccg gag gac ctg gat	144
Tyr Leu Glu Ile Gln Trp Trp Phe Leu Arg Gly Pro Glu Asp Leu Asp	
35 40 45	

ccc ggg gcc gag ggg gcc ggc gcg cag gtg gag ctc ttg ccc gac aga	192
Pro Gly Ala Glu Gly Ala Gly Ala Gln Val Glu Leu Leu Pro Asp Arg	
50 55 60	

gac ccg gac agc gac ggg acc aag atc agc aca gtg aaa gtc caa ggc	240
Asp Pro Asp Ser Asp Gly Thr Lys Ile Ser Thr Val Lys Val Gln Gly	
65 70 75 80	

aat gac atc tcc cac aag ctt cag att tcc aaa gtg agg aaa aag gat	288
Asn Asp Ile Ser His Lys Leu Gln Ile Ser Lys Val Arg Lys Lys Asp	
85 90 95	

-134-

gaa ggc tta tat gag tgc agg gtg act gat gcc aac tac ggg gag ctt 336
 Glu Gly Leu Tyr Glu Cys Arg Val Thr Asp Ala Asn Tyr Gly Glu Leu
 100 105 110

cag gaa cac aag gcc cag gcc tat ctg aaa gtc aat gcc aac agc cat 384
 Gln Glu His Lys Ala Gln Ala Tyr Leu Lys Val Asn Ala Asn Ser His
 115 120 125

gcc cgc aga atg cag gcc ttc gaa gcc tcg ccc atg tgg ctg cag gat 432
 Ala Arg Arg Met Gln Ala Phe Glu Ala Ser Pro Met Trp Leu Gln Asp
 130 135 140

atg aag ccc cgc aag aac gtc tcc gca gcc atc ccc agc agc atc cat 480
 Met Lys Pro Arg Lys Asn Val Ser Ala Ala Ile Pro Ser Ser Ile His
 145 150 155 160

ggc tct gcc aac caa cga acg cac tcc acc tcc agc cct caa gtg gta 528
 Gly Ser Ala Asn Gln Arg Thr His Ser Thr Ser Ser Pro Gln Val Val
 165 170 175

gcc aaa atc ccc aaa caa agt cca caa tca ggt atg gaa acc cat ttc 576
 Ala Lys Ile Pro Lys Gln Ser Pro Gln Ser Gly Met Glu Thr His Phe
 180 185 190

gag cct ttt att tta cca ctc aca aac gct cca cag aaa ggt cag tcg 624
 Glu Pro Phe Ile Leu Pro Leu Thr Asn Ala Pro Gln Lys Gly Gln Ser
 195 200 205

tat aga gta gac aga ttt atg aat ggt gat ttt taa 660
 Tyr Arg Val Asp Arg Phe Met Asn Gly Asp Phe
 210 215

<210> 5
 <211> 219
 <212> PRT
 <213> Homo sapiens

<400> 5

Gln Ala Lys Phe Thr Glu Phe Pro Arg Asn Val Thr Ala Thr Glu Gly
 1 5 10 15

Gln Asn Val Glu Met Ser Cys Ala Phe Gln Ser Gly Ser Ala Ser Val
 20 25 30

Tyr Leu Glu Ile Gln Trp Trp Phe Leu Arg Gly Pro Glu Asp Leu Asp
 35 40 45

Pro Gly Ala Glu Gly Ala Gly Ala Gln Val Glu Leu Leu Pro Asp Arg
 50 55 60

Asp Pro Asp Ser Asp Gly Thr Lys Ile Ser Thr Val Lys Val Gln Gly
 65 70 75 80

-135-

Asn	Asp	Ile	Ser	His	Lys	Leu	Gln	Ile	Ser	Lys	Val	Arg	Lys	Lys	Asp
				85					90					95	
Glu	Gly	Leu	Tyr	Glu	Cys	Arg	Val	Thr	Asp	Ala	Asn	Tyr	Gly	Glu	Leu
			100					105					110		
Gln	Glu	His	Lys	Ala	Gln	Ala	Tyr	Leu	Lys	Val	Asn	Ala	Asn	Ser	His
		115					120					125			
Ala	Arg	Arg	Met	Gln	Ala	Phe	Glu	Ala	Ser	Pro	Met	Trp	Leu	Gln	Asp
	130					135					140				
Met	Lys	Pro	Arg	Lys	Asn	Val	Ser	Ala	Ala	Ile	Pro	Ser	Ser	Ile	His
145					150					155					160
Gly	Ser	Ala	Asn	Gln	Arg	Thr	His	Ser	Thr	Ser	Ser	Pro	Gln	Val	Val
				165					170					175	
Ala	Lys	Ile	Pro	Lys	Gln	Ser	Pro	Gln	Ser	Gly	Met	Glu	Thr	His	Phe
			180					185					190		
Glu	Pro	Phe	Ile	Leu	Pro	Leu	Thr	Asn	Ala	Pro	Gln	Lys	Gly	Gln	Ser
		195					200					205			
Tyr	Arg	Val	Asp	Arg	Phe	Met	Asn	Gly	Asp	Phe					
	210					215									

<210> 6

<211> 3196

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (268)..(1020)

<400> 6

cacgcgtccg cagccagccg gcggcggaga cacttcacgg cgtggcaacc cgggtctgtg	60
ccttgaagcc tccggatcgc agccagctcg gtccatccct cactagtcgc aatcccctgt	120
gtccaagcta ctctttgcta tgagcggcag catgcgtgca gtatcgcgcc ccaggctctg	180
agagcagcct gcggacacgc ttgcctatct gtcttttttag gttttggggc tctgggctac	240
acggatgtgc cccactccct tggcatg atg ggg atc ttt ttg gcg tct gtt gga	294
Met Gly Ile Phe Leu Ala Ser Val Gly	
1 5	
ttt atg ttc ttt tcc gtg tta tat gta caa caa ggg ctt tct tct caa	342
Phe Met Phe Phe Ser Val Leu Tyr Val Gln Gln Gly Leu Ser Ser Gln	
10 15 20 25	

-136-

gca aaa ttt acc gag ttg ccg aga aat gtg act gct acc gaa ggg caa	390
Ala Lys Phe Thr Glu Leu Pro Arg Asn Val Thr Ala Thr Glu Gly Gln	
30 35 40	
aat gtg gag atg tcc tgt gct ttc caa agc ggc tct gct tca gtg tac	438
Asn Val Glu Met Ser Cys Ala Phe Gln Ser Gly Ser Ala Ser Val Tyr	
45 50 55	
ctg gag atc cag tgg tgg ttc ctt cgg ggg cca gag gac ctg gag caa	486
Leu Glu Ile Gln Trp Trp Phe Leu Arg Gly Pro Glu Asp Leu Glu Gln	
60 65 70	
ggc acg gag gct gca ggc tcg cag gtg gag ctc tta ccc gac aga gac	534
Gly Thr Glu Ala Ala Gly Ser Gln Val Glu Leu Leu Pro Asp Arg Asp	
75 80 85	
ccg gac aac gat ggg acc aag att agt aca gtg aaa gtc caa ggc aat	582
Pro Asp Asn Asp Gly Thr Lys Ile Ser Thr Val Lys Val Gln Gly Asn	
90 95 100 105	
gat atc tcc cac aag ctt cag ata tcc aaa gtg aga aaa aag gat gaa	630
Asp Ile Ser His Lys Leu Gln Ile Ser Lys Val Arg Lys Lys Asp Glu	
110 115 120	
ggt tta tac gag tgc agg gtg act gac gct aac tac ggg gag ctt cag	678
Gly Leu Tyr Glu Cys Arg Val Thr Asp Ala Asn Tyr Gly Glu Leu Gln	
125 130 135	
gaa cac aag gcc cag gcc tat ctg aaa gtc aat gcc aac agc cat gct	726
Glu His Lys Ala Gln Ala Tyr Leu Lys Val Asn Ala Asn Ser His Ala	
140 145 150	
cgg agg atg cag gcc ttt gaa gcc tca cct atg tgg ctg caa gac acg	774
Arg Arg Met Gln Ala Phe Glu Ala Ser Pro Met Trp Leu Gln Asp Thr	
155 160 165	
aag cct cga aag aac gca tca tcg gtg gtt ccc agc agc gtc cac aac	822
Lys Pro Arg Lys Asn Ala Ser Ser Val Val Pro Ser Ser Val His Asn	
170 175 180 185	
tct gcc aac caa cga atg cac tcc acc tcc agc cct caa gcg gta gcc	870
Ser Ala Asn Gln Arg Met His Ser Thr Ser Ser Pro Gln Ala Val Ala	
190 195 200	
aaa atc ccc aag caa agt cca caa tca gca aag agc aaa tcg cct gta	918
Lys Ile Pro Lys Gln Ser Pro Gln Ser Ala Lys Ser Lys Ser Pro Val	
205 210 215	
aaa tct acg gag cgg aca gca aag ttg acc cta tac tcc aag cac cat	966
Lys Ser Thr Glu Arg Thr Ala Lys Leu Thr Leu Tyr Ser Lys His His	
220 225 230	
tct gca ccc ctg tac tct agt tat cta cac aag gag cat cag ctt ccg	1014
Ser Ala Pro Leu Tyr Ser Ser Tyr Leu His Lys Glu His Gln Leu Pro	
235 240 245	

-137-

gaa gca taagtgaaga cactgtcaca cgctttattg ataatatattt ctttgggaag 1070
 Glu Ala
 250

ttgctgatct tttattttcaa gagaattaat gggaagagat aggacatttt ccaattacaa 1130
 gaccaatttt tttcctttta tttcaacaaa taaaacctgc atttcactga ctgctcagga 1190
 gttggcctga atgacatcag tataactaaat atttccatgg attccaccaa tttcctaacg 1250
 aggacacct aatcttcaag aagcaaacaa agatggaaaa cctaagaacc acaaactgtc 1310
 tcatacagca cccagctga ggaacaaaac aaatagctaa atgctgacca tggcaaatca 1370
 acatcagaca actttatttt acatatggaa taatcaaaga aagttttttt tttacttcct 1430
 ttttgcccc tggaatttat cttggagttt cccttttttc cttgattgcc gttttcgttc 1490
 aatggtagca agtgccaatt atggccaatc cttgtcaatc ctggaagggt tatattcata 1550
 tacattgagt gtggtatata tcaatgtatt ttaattcatt tggcaatttc tgtataggca 1610
 aacctggcaa attctgtaaa ttgcttatag tatgtgtgat atgacttcaa ggtagatagg 1670
 ctatgatgct catgcaagct gactttcttc attctatata caaatatatt catgagcata 1730
 tattaggcca ccaacttctt ttcctaaaga attatttttc atttgtacct catgtatttt 1790
 gtgaattttg tagtatattt ctctgttcca ctagtttgac cgctacagtt tgtctctggt 1850
 gtcctctact tccttctgga aaaattttaa attgtgtatg tctctgataa atgaattaat 1910
 tttgttgtgt gtagctatg ttggaatttg ctgtgttctt ttaaacaatgt atttattaag 1970
 gtttggggat cttgagttga gtctgaagaa tgcacacctg gtttttgaca gagttcctca 2030
 tgttaccaat attctatctc agagaaagaa agacaccaag tgggaaaact aagaagacat 2090
 tttgacttcc caagatcctg gaagagcact tcacactctg actaaataat gttgcttttt 2150
 ttgttcttca agactttttt gtagctttgt ctttctgtta gttgctgcta attatatattt 2210
 aatgtctact aattaaaaat taaaatgtga ttgttggctg aatacaatat gcaaactgact 2270
 gcaaagccca tactgaagaa aatagatggt taatcttcac tcaataatta taatttttaa 2330
 tagttcatca ttattttttg accttatgat attttgttta gacctgttct aattacatct 2390
 ttctctggca aagaaagata gaacaatcaa tacattccct cttacagtat ggaatgggtg 2450
 tggcttaaga aagaatgcat ccagatgggc ttccagagag attattttat tttcattata 2510
 aaaccagaaa ccatatatgt aggaatgggt cattcctaata gtaaggccat aaattgtagc 2570
 ttgaaggcaa ggaatacatt tgttttttta tggtaaagga ctggcctctg acatgcactt 2630

-138-

ataagcaatg tgaatatttt cataatatgc ttgacattct cctttaacaa atattgtttt 2690
 atggtaaato tttccttgcc atttttcttc tttcatttga ttcattattt cattctaag 2750
 aagaaaataa aggtttaatt atgatacttt attaacatac aaatgtattt tctttctaag 2810
 ttaaatatct gaaagttgta taaaatgatg gtagagaaat attactcatt cggtttcttt 2870
 gagctttaag aatcccatatc attgcagtat atattagaat actgatttaa catcaaactg 2930
 ggggggaaaa tcatgtatta tacttttact caatgtctag gtaatggatt cagctaattt 2990
 tacagcaagc caaatgtgta cccgtatcag taatgttcac catgcttgta ataaaagggc 3050
 atatgctagt tttggaagaa tgctcattag attcattgta tcagtgtcca aaataataaa 3110
 gacctgttta tcactgtgaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3170
 aaaaaaaaaa aaaaaagggc ggccgc 3196

<210> 7

<211> 251

<212> PRT

<213> Mus musculus

<400> 7

Met	Gly	Ile	Phe	Leu	Ala	Ser	Val	Gly	Phe	Met	Phe	Phe	Ser	Val	Leu
1				5				10						15	
Tyr	Val	Gln	Gln	Gly	Leu	Ser	Ser	Gln	Ala	Lys	Phe	Thr	Glu	Leu	Pro
		20						25					30		
Arg	Asn	Val	Thr	Ala	Thr	Glu	Gly	Gln	Asn	Val	Glu	Met	Ser	Cys	Ala
		35					40					45			
Phe	Gln	Ser	Gly	Ser	Ala	Ser	Val	Tyr	Leu	Glu	Ile	Gln	Trp	Trp	Phe
	50					55					60				
Leu	Arg	Gly	Pro	Glu	Asp	Leu	Glu	Gln	Gly	Thr	Glu	Ala	Ala	Gly	Ser
65					70					75					80
Gln	Val	Glu	Leu	Leu	Pro	Asp	Arg	Asp	Pro	Asp	Asn	Asp	Gly	Thr	Lys
			85					90						95	
Ile	Ser	Thr	Val	Lys	Val	Gln	Gly	Asn	Asp	Ile	Ser	His	Lys	Leu	Gln
			100					105					110		
Ile	Ser	Lys	Val	Arg	Lys	Lys	Asp	Glu	Gly	Leu	Tyr	Glu	Cys	Arg	Val
		115					120					125			
Thr	Asp	Ala	Asn	Tyr	Gly	Glu	Leu	Gln	Glu	His	Lys	Ala	Gln	Ala	Tyr
	130						135					140			

-139-

Leu Lys Val Asn Ala Asn Ser His Ala Arg Arg Met Gln Ala Phe Glu
 145 150 155 160

Ala Ser Pro Met Trp Leu Gln Asp Thr Lys Pro Arg Lys Asn Ala Ser
 165 170 175

Ser Val Val Pro Ser Ser Val His Asn Ser Ala Asn Gln Arg Met His
 180 185 190

Ser Thr Ser Ser Pro Gln Ala Val Ala Lys Ile Pro Lys Gln Ser Pro
 195 200 205

Gln Ser Ala Lys Ser Lys Ser Pro Val Lys Ser Thr Glu Arg Thr Ala
 210 215 220

Lys Leu Thr Leu Tyr Ser Lys His His Ser Ala Pro Leu Tyr Ser Ser
 225 230 235 240

Tyr Leu His Lys Glu His Gln Leu Pro Glu Ala
 245 250

<210> 8

<211> 753

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1)..(753)

<400> 8

atg ggg atc ttt ttg gcg tct gtt gga ttt atg ttc ttt tcc gtg tta 48
 Met Gly Ile Phe Leu Ala Ser Val Gly Phe Met Phe Phe Ser Val Leu
 1 5 10 15

tat gta caa caa ggg ctt tct tct caa gca aaa ttt acc gag ttg ccg 96
 Tyr Val Gln Gln Gly Leu Ser Ser Gln Ala Lys Phe Thr Glu Leu Pro
 20 25 30

aga aat gtg act gct acc gaa ggg caa aat gtg gag atg tcc tgt gct 144
 Arg Asn Val Thr Ala Thr Glu Gly Gln Asn Val Glu Met Ser Cys Ala
 35 40 45

ttc caa agc ggc tct gct tca gtg tac ctg gag atc cag tgg tgg ttc 192
 Phe Gln Ser Gly Ser Ala Ser Val Tyr Leu Glu Ile Gln Trp Trp Phe
 50 55 60

ctt cgg ggg cca gag gac ctg gag caa ggc acg gag gct gca ggc tcg 240
 Leu Arg Gly Pro Glu Asp Leu Glu Gln Gly Thr Glu Ala Ala Gly Ser
 65 70 75 80

-140-

cag gtg gag ctc tta ccc gac aga gac ccg gac aac gat ggg acc aag	288
Gln Val Glu Leu Leu Pro Asp Arg Asp Pro Asp Asn Asp Gly Thr Lys	
85 90 95	
att agt aca gtg aaa gtc caa ggc aat gat atc tcc cac aag ctt cag	336
Ile Ser Thr Val Lys Val Gln Gly Asn Asp Ile Ser His Lys Leu Gln	
100 105 110	
ata tcc aaa gtg aga aaa aag gat gaa ggt tta tac gag tgc agg gtg	384
Ile Ser Lys Val Arg Lys Lys Asp Glu Gly Leu Tyr Glu Cys Arg Val	
115 120 125	
act gac gct aac tac ggg gag ctt cag gaa cac aag gcc cag gcc tat	432
Thr Asp Ala Asn Tyr Gly Glu Leu Gln Glu His Lys Ala Gln Ala Tyr	
130 135 140	
ctg aaa gtc aat gcc aac agc cat gct cgg agg atg cag gcc ttt gaa	480
Leu Lys Val Asn Ala Asn Ser His Ala Arg Arg Met Gln Ala Phe Glu	
145 150 155 160	
gcc tca cct atg tgg ctg caa gac acg aag cct cga aag aac gca tca	528
Ala Ser Pro Met Trp Leu Gln Asp Thr Lys Pro Arg Lys Asn Ala Ser	
165 170 175	
tcg gtg gtt ccc agc agc gtc cac aac tct gcc aac caa cga atg cac	576
Ser Val Val Pro Ser Ser Val His Asn Ser Ala Asn Gln Arg Met His	
180 185 190	
tcc acc tcc agc cct caa gcg gta gcc aaa atc ccc aag caa agt cca	624
Ser Thr Ser Ser Pro Gln Ala Val Ala Lys Ile Pro Lys Gln Ser Pro	
195 200 205	
caa tca gca aag agc aaa tcg cct gta aaa tct acg gag cgg aca gca	672
Gln Ser Ala Lys Ser Lys Ser Pro Val Lys Ser Thr Glu Arg Thr Ala	
210 215 220	
aag ttg acc cta tac tcc aag cac cat tct gca ccc ctg tac tct agt	720
Lys Leu Thr Leu Tyr Ser Lys His His Ser Ala Pro Leu Tyr Ser Ser	
225 230 235 240	
tat cta cac aag gag cat cag ctt ccg gaa gca	753
Tyr Leu His Lys Glu His Gln Leu Pro Glu Ala	
245 250	

<210> 9

<211> 681

<212> DNA

<213> Mus musculus

<220>

<221> CDS

<222> (1)..(681)

<400> 9

-141-

caa gca aaa ttt acc gag ttg ccg aga aat gtg act gct acc gaa ggg	48
Gln Ala Lys Phe Thr Glu Leu Pro Arg Asn Val Thr Ala Thr Glu Gly	
1 5 10 15	
caa aat gtg gag atg tcc tgt gct ttc caa agc ggc tct gct tca gtg	96
Gln Asn Val Glu Met Ser Cys Ala Phe Gln Ser Gly Ser Ala Ser Val	
20 25 30	
tac ctg gag atc cag tgg tgg ttc ctt cgg ggg cca gag gac ctg gag	144
Tyr Leu Glu Ile Gln Trp Trp Phe Leu Arg Gly Pro Glu Asp Leu Glu	
35 40 45	
caa ggc acg gag gct gca ggc tcg cag gtg gag ctc tta ccc gac aga	192
Gln Gly Thr Glu Ala Ala Gly Ser Gln Val Glu Leu Leu Pro Asp Arg	
50 55 60	
gac ccg gac aac gat ggg acc aag att agt aca gtg aaa gtc caa ggc	240
Asp Pro Asp Asn Asp Gly Thr Lys Ile Ser Thr Val Lys Val Gln Gly	
65 70 75 80	
aat gat atc tcc cac aag ctt cag ata tcc aaa gtg aga aaa aag gat	288
Asn Asp Ile Ser His Lys Leu Gln Ile Ser Lys Val Arg Lys Lys Asp	
85 90 95	
gaa ggt tta tac gag tgc agg gtg act gac gct aac tac ggg gag ctt	336
Glu Gly Leu Tyr Glu Cys Arg Val Thr Asp Ala Asn Tyr Gly Glu Leu	
100 105 110	
cag gaa cac aag gcc cag gcc tat ctg aaa gtc aat gcc aac agc cat	384
Gln Glu His Lys Ala Gln Ala Tyr Leu Lys Val Asn Ala Asn Ser His	
115 120 125	
gct cgg agg atg cag gcc ttt gaa gcc tca cct atg tgg ctg caa gac	432
Ala Arg Arg Met Gln Ala Phe Glu Ala Ser Pro Met Trp Leu Gln Asp	
130 135 140	
acg aag cct cga aag aac gca tca tcg gtg gtt ccc agc agc gtc cac	480
Thr Lys Pro Arg Lys Asn Ala Ser Ser Val Val Pro Ser Ser Val His	
145 150 155 160	
aac tct gcc aac caa cga atg cac tcc acc tcc agc cct caa gcg gta	528
Asn Ser Ala Asn Gln Arg Met His Ser Thr Ser Ser Pro Gln Ala Val	
165 170 175	
gcc aaa atc ccc aag caa agt cca caa tca gca aag agc aaa tcg cct	576
Ala Lys Ile Pro Lys Gln Ser Pro Gln Ser Ala Lys Ser Lys Ser Pro	
180 185 190	
gta aaa tct acg gag cgg aca gca aag ttg acc cta tac tcc aag cac	624
Val Lys Ser Thr Glu Arg Thr Ala Lys Leu Thr Leu Tyr Ser Lys His	
195 200 205	
cat tct gca ccc ctg tac tct agt tat cta cac aag gag cat cag ctt	672
His Ser Ala Pro Leu Tyr Ser Ser Tyr Leu His Lys Glu His Gln Leu	
210 215 220	

681

```
<210> 10
<211> 227
<212> PRT
<213> Mus musculus
```

<400> 10

Gln 1	Ala	Lys	Phe	Thr 5	Glu	Leu	Pro	Arg	Asn 10	Val	Thr	Ala	Thr	Glu	Gly 15
Gln	Asn	Val	Glu 20	Met	Ser	Cys	Ala	Phe 25	Gln	Ser	Gly	Ser	Ala 30	Ser	Val
Tyr	Leu	Glu 35	Ile	Gln	Trp	Trp	Phe 40	Leu	Arg	Gly	Pro	Glu 45	Asp	Leu	Glu
Gln	Gly 50	Thr	Glu	Ala	Ala	Gly 55	Ser	Gln	Val	Glu	Leu 60	Leu	Pro	Asp	Arg
Asp 65	Pro	Asp	Asn	Asp	Gly 70	Thr	Lys	Ile	Ser	Thr 75	Val	Lys	Val	Gln	Gly 80
Asn	Asp	Ile	Ser	His 85	Lys	Leu	Gln	Ile	Ser 90	Lys	Val	Arg	Lys	Lys 95	Asp
Glu	Gly	Leu	Tyr 100	Glu	Cys	Arg	Val	Thr 105	Asp	Ala	Asn	Tyr	Gly 110	Glu	Leu
Gln	Glu 115	His	Lys	Ala	Gln	Ala	Tyr 120	Leu	Lys	Val	Asn 125	Ala	Asn	Ser	His
Ala 130	Arg	Arg	Met	Gln	Ala	Phe 135	Glu	Ala	Ser	Pro	Met 140	Trp	Leu	Gln	Asp
Thr 145	Lys	Pro	Arg	Lys	Asn 150	Ala	Ser	Ser	Val	Val 155	Pro	Ser	Ser	Val	His 160
Asn	Ser	Ala	Asn	Gln 165	Arg	Met	His	Ser	Thr 170	Ser	Ser	Pro	Gln	Ala 175	Val
Ala	Lys	Ile	Pro 180	Lys	Gln	Ser	Pro	Gln 185	Ser	Ala	Lys	Ser	Lys 190	Ser	Pro
Val	Lys 195	Ser	Thr	Glu	Arg	Thr	Ala 200	Lys	Leu	Thr	Leu 205	Tyr	Ser	Lys	His
His 210	Ser	Ala	Pro	Leu	Tyr	Ser 215	Ser	Tyr	Leu	His	Lys 220	Glu	His	Gln	Leu

<400> 11

[illegible]

```
<210> 12
<211> 89
<212> PRT
<213> Mus musculus
```

<400> 12

Gly 1	Gln	Asn	Val	Glu 5	Met	Ser	Cys	Ala	Phe 10	Gln	Ser	Gly	Ser	Ala 15	Ser
Val	Tyr	Leu	Glu 20	Ile	Gln	Trp	Trp	Phe 25	Leu	Arg	Gly	Pro	Glu 30	Asp	Leu
Glu	Gln	Gly 35	Thr	Glu	Ala	Ala	Gly 40	Ser	Gln	Val	Glu 45	Leu	Leu	Pro	Asp
Arg 50	Asp	Pro	Asp	Asn	Asp	Gly 55	Thr	Lys	Ile	Ser 60	Thr	Val	Lys	Val	Gln
Gly 65	Asn	Asp	Ile	Ser	His 70	Lys	Leu	Gln	Ile	Ser 75	Lys	Val	Arg	Lys	Lys 80
Asp	Glu	Gly	Leu	Tyr 85	Glu	Cys	Arg	Val							

-144-

<210> 13

<211> 47

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Hidden Markov Model

<400> 13

Gly	Gln	Ser	Val	Thr	Leu	Thr	Cys	Met	Val	Ser	Phe	His	Pro	Pro	Asp
1				5					10					15	
Tyr	Thr	Ile	Trp	Trp	Tyr	Arg	Asn	Gly	Gln	Pro	Ile	Thr	Leu	Thr	Ile
			20					25					30		
Asn	Ser	Trp	Gln	Tyr	Glu	Asp	Ser	Glu	Thr	Tyr	Trp	Cys	Met	Val	
		35					40					45			

<210> 14

<211> 2915

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (404) .. (1294)

<400> 14

gaattcccgg	gtcgacccac	gcgtccgccc	acgcgtccgc	ccacgcgtcc	ggctgagcga	60
agcgcggcgg	cggcggcggc	gcctagggga	gggaggggcg	gcggggccga	gcccacctag	120
cggagcgcgc	cggccgcccgg	tggccgcccgc	cagcatgccc	cggcccgcgg	gccgctccgc	180
cgccagccac	ccccgcggcc	ctcggcggcc	tgcgctcggc	ccggggggcgc	gggaaccgca	240
gccggagccg	gaggcgggag	cagcgagccg	gagccccggg	cgctcgaatg	caggatgctc	300
gtggtcccca	gcatecttga	gccaccagga	gtgagggctg	ctgctccctg	agacctggct	360
ccaaggagga	tgccacagcc	gcctgccagc	tccggtctgc	acc atg agt gat gag		415
				Met Ser Asp Glu		
				1		
cgg cgg ctg cct ggc agt gca gtg ggc tgg ctg gta tgt ggg ggc ctc	463					
Arg Arg Leu Pro Gly Ser Ala Val Gly Trp Leu Val Cys Gly Gly Leu						
5 10 15 20						
tcc ctg ctg gcc aat gcc tgg ggc atc ctc agc gtt ggc gcc aag cag	511					
Ser Leu Leu Ala Asn Ala Trp Gly Ile Leu Ser Val Gly Ala Lys Gln						
25 30 35						

-145-

aag aag tgg aag ccc ttg gag ttc ctg ctg tgt acg ctc gcg gcc acc	559
Lys Lys Trp Lys Pro Leu Glu Phe Leu Leu Cys Thr Leu Ala Ala Thr	
40 45 50	
cac atg cta aat gtg gcc gtg ccc atc gcc acc tac tcc gtg gtg cag	607
His Met Leu Asn Val Ala Val Pro Ile Ala Thr Tyr Ser Val Val Gln	
55 60 65	
ctg cgg cgg cag cgc ccc gac ttc gag tgg aat gag ggt ctc tgc aag	655
Leu Arg Arg Gln Arg Pro Asp Phe Glu Trp Asn Glu Gly Leu Cys Lys	
70 75 80	
gtc ttc gtg tcc acc ttc tac acc ctc acc ctg gcc acc tgt ttc tct	703
Val Phe Val Ser Thr Phe Tyr Thr Leu Thr Leu Ala Thr Cys Phe Ser	
85 90 95 100	
gtc acc tcc ctc tcc tac cac cgc atg tgg atg gtc tgc tgg cct gtc	751
Val Thr Ser Leu Ser Tyr His Arg Met Trp Met Val Cys Trp Pro Val	
105 110 115	
aac tac cgg ctg agc aat gcc aag aag cag gcg gtg cac aca gtc atg	799
Asn Tyr Arg Leu Ser Asn Ala Lys Lys Gln Ala Val His Thr Val Met	
120 125 130	
ggt atc tgg atg gtg tcc ttc atc ctg tcg gcc ctg cct gcc gtt ggc	847
Gly Ile Trp Met Val Ser Phe Ile Leu Ser Ala Leu Pro Ala Val Gly	
135 140 145	
tgg cac gac acc agc gag cgc ttc tac acc cat ggc tgc cgc ttc atc	895
Trp His Asp Thr Ser Glu Arg Phe Tyr Thr His Gly Cys Arg Phe Ile	
150 155 160	
gtg gct gag atc ggc ctg ggc ttt ggc gtc tgc ttc ctg ctg ctg gtg	943
Val Ala Glu Ile Gly Leu Gly Phe Gly Val Cys Phe Leu Leu Leu Val	
165 170 175 180	
ggc ggc agc gtg gcc atg ggc gtg atc tgc aca gcc atc gcc ctc ttc	991
Gly Gly Ser Val Ala Met Gly Val Ile Cys Thr Ala Ile Ala Leu Phe	
185 190 195	
cag acg ctg gcc gtg cag gtg ggg cgc cag gcc gac cac cgc gcc ttc	1039
Gln Thr Leu Ala Val Gln Val Gly Arg Gln Ala Asp His Arg Ala Phe	
200 205 210	
acc gtg ccc acc atc gtg gtg gag gac gcg cag ggc aag cgg cgc tcc	1087
Thr Val Pro Thr Ile Val Val Glu Asp Ala Gln Gly Lys Arg Arg Ser	
215 220 225	
tcc atc gat ggc tcg gag ccc gcc aaa acc tct ctg cag acc acg ggc	1135
Ser Ile Asp Gly Ser Glu Pro Ala Lys Thr Ser Leu Gln Thr Thr Gly	
230 235 240	
ctc gtg acc acc ata gtc ttc atc tac gac tgc ctc atg ggc ttc cct	1183
Leu Val Thr Thr Ile Val Phe Ile Tyr Asp Cys Leu Met Gly Phe Pro	
245 250 255 260	

-146-

gtg ctg gac tct acg ccc atc ccc gaa agg tct gca gtg aga cag gga	1231
Val Leu Asp Ser Thr Pro Ile Pro Glu Arg Ser Ala Val Arg Gln Gly	
265 270 275	
gag gac tgg ggc aaa gac cag cct gag ggg ttt cat cca agc agc agg	1279
Glu Asp Trp Gly Lys Asp Gln Pro Glu Gly Phe His Pro Ser Ser Arg	
280 285 290	
caa gac tgc ctt ccc tgagccattg caggacatga ggacatgagc tccagaatgg	1334
Gln Asp Cys Leu Pro	
295	
tgccaggccg agccctgtgc ccacaggtgg tgagcttcag cagcctgcgg gccgacgcct	1394
cagcgccctg gatggcactc tgcgtgctgt ggtgctccgt ggcccaggcc ctgcttccga	1454
cctgtcctgc ctcagtttcc ccatctgtga tgagcaggtg accacgttaa ctctcagggc	1514
tgtttgaagt ctcttggttg taggcctcgc ccactgagtg gcccaggtgt gagaggtagt	1574
ctagagccct ctgcggcctt tgtggaggtc cgttctcagc atgtggcctg gtgtggctcc	1634
ctaggcctga ggcctccac tctcagtgcc ccctgcccc ttgggaacct aactccacc	1694
ccagctaagc acaacactgt ggaccaccac caatggcacc tgagccacct tgaccaccat	1754
tagtgtcccc accctcatta ctctgccgtc gttgcctgtc cccaccacat catctccctc	1814
catggctccc accatgacgc tggcaggtgc aagcggagag cacgagggca gcagggcctg	1874
gccttccgtg tccctgcgtc atccccagcc tcaactccca cccgtgcagg cctggccagg	1934
aagggatcct ggcaggggct tccaggttct cagctcaagg cctgggtccc ggcagggcgtc	1994
caacccttg gagcaatgta tttctttgcc ttccatcctg ggcagacccc ttacaggcct	2054
gggcattgcc atgggccctg ggtcttccca ggctaaggag aaccaggaac agctataacc	2114
ttgagctagt gaaatagggt agatgagaag gctgtctcct ccagaccct acccctacac	2174
agtggcccca caatatgaag acctggggta attccaaggt gagcatagag cctgcctgtg	2234
cccagttcct tctggccctc aggtggccaa gccatctct tcatccttca gatagggctc	2294
cactcccaga agaagctgct ggggtggggg tgggaggctg cctaagcctg tctgtgcttc	2354
agaggccct ccagtcctg gctgtgggggt aactgggggt atgagctgtg gccacaggtg	2414
agcaaggcag ggaactgcaa tccagccctg gccgcgggag gggccatctc tggccaatgc	2474
tgctgtgct tcaaggactg acaagttacg taggggcaga ggtcgccagc tagccagtgt	2534
ctcctccatc tggggggcgt ctgtccactt gtcaccttag gttttcactc atttgtcacc	2594
ttggggtttt gctctgtgtg tttcatatcc aacggcaata cttgcagggg gacagagtcc	2654

-147-

tctaaatact ccaatcctgc ggttttttaca aacataaagg gggagacccc aagtggagga 2714
 ccctgggcct ggagctccct cccaaacttt gtccagcatc cagcctgttc cctgggctca 2774
 ctggggaggg agttgtcttc atagcacacc cagagccagg gatccctttg tagttttttg 2834
 acaacggagc atttctcttc tgtacaggac ccaataaaaa cttccttatg atttgcaaaa 2894
 aaaaaaaaaa agggcggccg c 2915

<210> 15

<211> 297

<212> PRT

<213> Homo sapiens

<400> 15

Met Ser Asp Glu Arg Arg Leu Pro Gly Ser Ala Val Gly Trp Leu Val
 1 5 10 15

Cys Gly Gly Leu Ser Leu Leu Ala Asn Ala Trp Gly Ile Leu Ser Val
 20 25 30

Gly Ala Lys Gln Lys Lys Trp Lys Pro Leu Glu Phe Leu Leu Cys Thr
 35 40 45

Leu Ala Ala Thr His Met Leu Asn Val Ala Val Pro Ile Ala Thr Tyr
 50 55 60

Ser Val Val Gln Leu Arg Arg Gln Arg Pro Asp Phe Glu Trp Asn Glu
 65 70 75 80

Gly Leu Cys Lys Val Phe Val Ser Thr Phe Tyr Thr Leu Thr Leu Ala
 85 90 95

Thr Cys Phe Ser Val Thr Ser Leu Ser Tyr His Arg Met Trp Met Val
 100 105 110

Cys Trp Pro Val Asn Tyr Arg Leu Ser Asn Ala Lys Lys Gln Ala Val
 115 120 125

His Thr Val Met Gly Ile Trp Met Val Ser Phe Ile Leu Ser Ala Leu
 130 135 140

Pro Ala Val Gly Trp His Asp Thr Ser Glu Arg Phe Tyr Thr His Gly
 145 150 155 160

Cys Arg Phe Ile Val Ala Glu Ile Gly Leu Gly Phe Gly Val Cys Phe
 165 170 175

Leu Leu Leu Val Gly Gly Ser Val Ala Met Gly Val Ile Cys Thr Ala
 180 185 190

-148-

Ile	Ala	Leu	Phe	Gln	Thr	Leu	Ala	Val	Gln	Val	Gly	Arg	Gln	Ala	Asp	
		195					200					205				
His	Arg	Ala	Phe	Thr	Val	Pro	Thr	Ile	Val	Val	Glu	Asp	Ala	Gln	Gly	
	210					215					220					
Lys	Arg	Arg	Ser	Ser	Ile	Asp	Gly	Ser	Glu	Pro	Ala	Lys	Thr	Ser	Leu	
225					230					235					240	
Gln	Thr	Thr	Gly	Leu	Val	Thr	Thr	Ile	Val	Phe	Ile	Tyr	Asp	Cys	Leu	
				245					250					255		
Met	Gly	Phe	Pro	Val	Leu	Asp	Ser	Thr	Pro	Ile	Pro	Glu	Arg	Ser	Ala	
			260					265					270			
Val	Arg	Gln	Gly	Glu	Asp	Trp	Gly	Lys	Asp	Gln	Pro	Glu	Gly	Phe	His	
	275						280					285				
Pro	Ser	Ser	Arg	Gln	Asp	Cys	Leu	Pro								
	290					295										

<210> 16
 <211> 891
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(891)

<400> 16

atg	agt	gat	gag	cgg	cgg	ctg	cct	ggc	agt	gca	gtg	ggc	tgg	ctg	gta	48
Met	Ser	Asp	Glu	Arg	Arg	Leu	Pro	Gly	Ser	Ala	Val	Gly	Trp	Leu	Val	
1				5				10					15			
tgt	ggg	ggc	ctc	tcc	ctg	ctg	gcc	aat	gcc	tgg	ggc	atc	ctc	agc	gtt	96
Cys	Gly	Gly	Leu	Ser	Leu	Leu	Ala	Asn	Ala	Trp	Gly	Ile	Leu	Ser	Val	
			20					25					30			
ggc	gcc	aag	cag	aag	aag	tgg	aag	ccc	ttg	gag	ttc	ctg	ctg	tgt	acg	144
Gly	Ala	Lys	Gln	Lys	Lys	Trp	Lys	Pro	Leu	Glu	Phe	Leu	Leu	Cys	Thr	
		35					40					45				
ctc	gcg	gcc	acc	cac	atg	cta	aat	gtg	gcc	gtg	ccc	atc	gcc	acc	tac	192
Leu	Ala	Ala	Thr	His	Met	Leu	Asn	Val	Ala	Val	Pro	Ile	Ala	Thr	Tyr	
	50					55					60					
tcc	gtg	gtg	cag	ctg	cgg	cgg	cag	cgc	ccc	gac	ttc	gag	tgg	aat	gag	240
Ser	Val	Val	Gln	Leu	Arg	Arg	Gln	Arg	Pro	Asp	Phe	Glu	Trp	Asn	Glu	
65					70				75						80	

-149-

ggt ctc tgc aag gtc ttc gtg tcc acc ttc tac acc ctc acc ctg gcc	288
Gly Leu Cys Lys Val Phe Val Ser Thr Phe Tyr Thr Leu Thr Leu Ala	
85 90 95	
acc tgt ttc tct gtc acc tcc ctc tcc tac cac cgc atg tgg atg gtc	336
Thr Cys Phe Ser Val Thr Ser Leu Ser Tyr His Arg Met Trp Met Val	
100 105 110	
tgc tgg cct gtc aac tac cgg ctg agc aat gcc aag aag cag gcg gtg	384
Cys Trp Pro Val Asn Tyr Arg Leu Ser Asn Ala Lys Lys Gln Ala Val	
115 120 125	
cac aca gtc atg ggt atc tgg atg gtg tcc ttc atc ctg tcg gcc ctg	432
His Thr Val Met Gly Ile Trp Met Val Ser Phe Ile Leu Ser Ala Leu	
130 135 140	
cct gcc gtt ggc tgg cac gac acc agc gag cgc ttc tac acc cat ggc	480
Pro Ala Val Gly Trp His Asp Thr Ser Glu Arg Phe Tyr Thr His Gly	
145 150 155 160	
tgc cgc ttc atc gtg gct gag atc ggc ctg ggc ttt ggc gtc tgc ttc	528
Cys Arg Phe Ile Val Ala Glu Ile Gly Leu Gly Phe Gly Val Cys Phe	
165 170 175	
ctg ctg ctg gtg ggc ggc agc gtg gcc atg ggc gtg atc tgc aca gcc	576
Leu Leu Leu Val Gly Gly Ser Val Ala Met Gly Val Ile Cys Thr Ala	
180 185 190	
atc gcc ctc ttc cag acg ctg gcc gtg cag gtg ggg cgc cag gcc gac	624
Ile Ala Leu Phe Gln Thr Leu Ala Val Gln Val Gly Arg Gln Ala Asp	
195 200 205	
cac cgc gcc ttc acc gtg ccc acc atc gtg gtg gag gac gcg cag ggc	672
His Arg Ala Phe Thr Val Pro Thr Ile Val Val Glu Asp Ala Gln Gly	
210 215 220	
aag cgg cgc tcc tcc atc gat ggc tcg gag ccc gcc aaa acc tct ctg	720
Lys Arg Arg Ser Ser Ile Asp Gly Ser Glu Pro Ala Lys Thr Ser Leu	
225 230 235 240	
cag acc acg ggc ctc gtg acc acc ata gtc ttc atc tac gac tgc ctc	768
Gln Thr Thr Gly Leu Val Thr Thr Ile Val Phe Ile Tyr Asp Cys Leu	
245 250 255	
atg ggc ttc cct gtg ctg gac tct acg ccc atc ccc gaa agg tct gca	816
Met Gly Phe Pro Val Leu Asp Ser Thr Pro Ile Pro Glu Arg Ser Ala	
260 265 270	
gtg aga cag gga gag gac tgg ggc aaa gac cag cct gag ggg ttt cat	864
Val Arg Gln Gly Glu Asp Trp Gly Lys Asp Gln Pro Glu Gly Phe His	
275 280 285	
cca agc agc agg caa gac tgc ctt ccc	891
Pro Ser Ser Arg Gln Asp Cys Leu Pro	
290 295	

-150-

<210> 17
 <211> 4166
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (334) .. (2160)

<400> 17

```

gaattccccgg gtcgacccac gcgtccggcg gcggggccga gcccacctag cggagcgcgc      60
cggccgcgcgg tggccgcgcg cagcatgccc cggcccgcgg gccgctccgc cgccagccac      120
ccccgcggcc ctcggcgggc tgcgctcggc ccggggggcg gggaaccgca gccggagccg      180
gaggcgggag cagcgagccg gagccccggg cgctcgaatg caggatgctc gtgggtcccca      240
gcatccttga gccaccagga gtgagggctg ctgctccctg agacctggct ccaaggagga      300
tgccacagcc gcctgccagc tccgggtctgc acc atg agt gat gag cgg cgg ctg      354
                               Met Ser Asp Glu Arg Arg Leu
                               1                               5

cct ggc agt gca gtg ggc tgg ctg gta tgt ggg ggc ctc tcc ctg ctg      402
Pro Gly Ser Ala Val Gly Trp Leu Val Cys Gly Gly Leu Ser Leu Leu
          10                      15                      20

gcc aat gcc tgg ggc atc ctc agc gtt ggc gcc aag cag aag aag tgg      450
Ala Asn Ala Trp Gly Ile Leu Ser Val Gly Ala Lys Gln Lys Lys Trp
          25                      30                      35

aag ccc ttg gag ttc ctg ctg tgt acg ctc gcg gcc acc cac atg cta      498
Lys Pro Leu Glu Phe Leu Leu Cys Thr Leu Ala Ala Thr His Met Leu
          40                      45                      50                      55

aat gtg gcc gtg ccc atc gcc acc tac tcc gtg gtg cag ctg cgg cgg      546
Asn Val Ala Val Pro Ile Ala Thr Tyr Ser Val Val Gln Leu Arg Arg
                      60                      65                      70

cag cgc ccc gac ttc gag tgg aat gag ggt ctc tgc aag gtc ttc gtg      594
Gln Arg Pro Asp Phe Glu Trp Asn Glu Gly Leu Cys Lys Val Phe Val
                      75                      80                      85

tcc acc ttc tac acc ctc acc ctg gcc acc tgt ttc tct gtc acc tcc      642
Ser Thr Phe Tyr Thr Leu Thr Leu Ala Thr Cys Phe Ser Val Thr Ser
          90                      95                      100

ctc tcc tac cac cgc atg tgg atg gtc tgc tgg cct gtc aac tac cgg      690
Leu Ser Tyr His Arg Met Trp Met Val Cys Trp Pro Val Asn Tyr Arg
          105                      110                      115

```

-151-

ctg agc aat gcc aag aag cag gcg gtg cac aca gtc atg ggt atc tgg	738
Leu Ser Asn Ala Lys Lys Gln Ala Val His Thr Val Met Gly Ile Trp	
120 125 130 135	
atg gtg tcc ttc atc ctg tcg gcc ctg cct gcc gtt ggc tgg cac gac	786
Met Val Ser Phe Ile Leu Ser Ala Leu Pro Ala Val Gly Trp His Asp	
140 145 150	
acc agc gag cgc ttc tac acc cat ggc tgc cgc ttc atc gtg gct gag	834
Thr Ser Glu Arg Phe Tyr Thr His Gly Cys Arg Phe Ile Val Ala Glu	
155 160 165	
atc ggc ctg ggc ttt ggc gtc tgc ttc ctg ctg ctg gtg ggc ggc agc	882
Ile Gly Leu Gly Phe Gly Val Cys Phe Leu Leu Leu Val Gly Gly Ser	
170 175 180	
gtg gcc atg ggc gtg atc tgc aca gcc atc gcc ctc ttc cag acg ctg	930
Val Ala Met Gly Val Ile Cys Thr Ala Ile Ala Leu Phe Gln Thr Leu	
185 190 195	
gcc gtg cag gtg ggg cgc cag gcc gac cac cgc gcc ttc acc gtg ccc	978
Ala Val Gln Val Gly Arg Gln Ala Asp His Arg Ala Phe Thr Val Pro	
200 205 210 215	
acc atc gtg gtg gag gac gcg cag ggc aag cgg cgc tcc tcc atc gat	1026
Thr Ile Val Val Glu Asp Ala Gln Gly Lys Arg Arg Ser Ser Ile Asp	
220 225 230	
ggc tcg gag ccc gcc aaa acc tct ctg cag acc acg ggc ctc gtg acc	1074
Gly Ser Glu Pro Ala Lys Thr Ser Leu Gln Thr Thr Gly Leu Val Thr	
235 240 245	
acc ata gtc ttc atc tac gac tgc ctc atg ggc ttc cct gtg ctg gtg	1122
Thr Ile Val Phe Ile Tyr Asp Cys Leu Met Gly Phe Pro Val Leu Val	
250 255 260	
gtg agc ttc agc agc ctg cgg gcc gac gcc tca gcg ccc tgg atg gca	1170
Val Ser Phe Ser Ser Leu Arg Ala Asp Ala Ser Ala Pro Trp Met Ala	
265 270 275	
ctc tgc gtg ctg tgg tgc tcc gtg gcc cag gcc ctg ctg ctg cct gtg	1218
Leu Cys Val Leu Trp Cys Ser Val Ala Gln Ala Leu Leu Leu Pro Val	
280 285 290 295	
ttc ctc tgg gcc tgc gac cgc tac cgg gct gac ctc aaa gct gtc cgg	1266
Phe Leu Trp Ala Cys Asp Arg Tyr Arg Ala Asp Leu Lys Ala Val Arg	
300 305 310	
gag aag tgc atg gcc ctc atg gcc aac gac gag gag tca gac gat gag	1314
Glu Lys Cys Met Ala Leu Met Ala Asn Asp Glu Glu Ser Asp Asp Glu	
315 320 325	
acc agc ctg gaa ggt ggc atc tcc ccg gac ctg gtg ttg gag cgc tcc	1362
Thr Ser Leu Glu Gly Gly Ile Ser Pro Asp Leu Val Leu Glu Arg Ser	
330 335 340	

-152-

ctg gac tat ggc tat gga ggt gat ttt gtg gcc cta gat agg atg gcc Leu Asp Tyr Gly Tyr Gly Gly Asp Phe Val Ala Leu Asp Arg Met Ala 345 350 355	1410
aag tat gag atc tcc gcc ctg gag ggg ggc ctg ccc cag ctc tac cca Lys Tyr Glu Ile Ser Ala Leu Glu Gly Gly Leu Pro Gln Leu Tyr Pro 360 365 370 375	1458
ctg cgg ccc ttg cag gag gac aag atg caa tac ctg cag gtc ccg ccc Leu Arg Pro Leu Gln Glu Asp Lys Met Gln Tyr Leu Gln Val Pro Pro 380 385 390	1506
acg cgg cgc ttc tcc cac gac gat gcg gac gtg tgg gcc gcc gtc ccg Thr Arg Arg Phe Ser His Asp Asp Ala Asp Val Trp Ala Ala Val Pro 395 400 405	1554
ctg ccc gcc ttc ctg ccg cgc tgg ggc tcc ggc gag gac ctg gcc gcc Leu Pro Ala Phe Leu Pro Arg Trp Gly Ser Gly Glu Asp Leu Ala Ala 410 415 420	1602
ctg gcg cac ctg gtg ctg cct gcc ggg ccc gag cgg cgc cgc gcc agc Leu Ala His Leu Val Leu Pro Ala Gly Pro Glu Arg Arg Arg Ala Ser 425 430 435	1650
ctc ctg gcc ttc gcg gag gac gca cca ccg tcc cgc gcg cgc cgc cgc Leu Leu Ala Phe Ala Glu Asp Ala Pro Pro Ser Arg Ala Arg Arg Arg 440 445 450 455	1698
tcg gcc gag agc ctg ctg tcg ctg cgg acc tcg gcc ctg gat agc ggc Ser Ala Glu Ser Leu Leu Ser Leu Arg Thr Ser Ala Leu Asp Ser Gly 460 465 470	1746
ccg cgg gga gcc cgc gac tcg ccc ccc ggc agc ccg cgc cgc cgc ccc Pro Arg Gly Ala Arg Asp Ser Pro Pro Gly Ser Pro Arg Arg Arg Pro 475 480 485	1794
ggg ccc ggc ccc cgc tcc gcc tcg gcc tcg ctg ctg ccc gac gcc ttc Gly Pro Gly Pro Arg Ser Ala Ser Ala Ser Leu Leu Pro Asp Ala Phe 490 495 500	1842
gcc ctg acc gcc ttc gag tgc gag cca cag gcc ctg cgc cgc ccg ccc Ala Leu Thr Ala Phe Glu Cys Glu Pro Gln Ala Leu Arg Arg Pro Pro 505 510 515	1890
ggg ccc ttc ccc gct gcg ccc gcc gcc ccc gac ggc gca gat ccc gga Gly Pro Phe Pro Ala Ala Pro Ala Ala Pro Asp Gly Ala Asp Pro Gly 520 525 530 535	1938
gag gcc ccg acg ccc cca agc agc gcc cag cgg agc cca ggg cca cgc Glu Ala Pro Thr Pro Pro Ser Ser Ala Gln Arg Ser Pro Gly Pro Arg 540 545 550	1986

-153-

ccc tct gcg cac tcg cac gcc ggc tct ctg cgc ccc ggc ctg agc gcg Pro Ser Ala His Ser His Ala Gly Ser Leu Arg Pro Gly Leu Ser Ala 555 560 565	2034
tcg tgg ggc gag ccc ggg ggg ctg cgc gcg gcg ggc ggc ggc ggc agc Ser Trp Gly Glu Pro Gly Gly Leu Arg Ala Ala Gly Gly Gly Gly Ser 570 575 580	2082
acc agc agc ttc ctg agt tcc ccc tcc gag tcc tcg ggc tac gcc acg Thr Ser Ser Phe Leu Ser Ser Pro Ser Glu Ser Ser Gly Tyr Ala Thr 585 590 595	2130
ctg cac tcg gac tcg ctg ggc tcc gcg tcc taggaccgcc ggcgcctccc Leu His Ser Asp Ser Leu Gly Ser Ala Ser 600 605	2180
cacggacgcc aggcaggcca ggccgctctc cggggccgca gcaccaaaga cgcccgcctc	2240
ccccccgcgc gcagacatgc gccacccctc ccaggggtga gggggcgcttg gcctcagcgt	2300
ttgtcttccg gctcctccca gctggccttg tcccaggggc gacggctgcc ccggacgact	2360
gcgctgggca ccgcatgtcc cgggccgagt gaggtcgggc ctggggagct gaggacatc	2420
ccaagcttgg gcttgggtag tgagtacat gtgcacacgt ccagctgcgc catcaccagc	2480
cctgcgcaac aggacgtcgg gagcaggga cctgagacag ggccactgcg ggatcggaca	2540
aagccccgct ttggagaggc tgagctggag ccattggcct cccaggggc tttccacca	2600
cactgcacca taaccgccac acccttcggg gggggggagg gtacagaggg tctctaagca	2660
caggggtggt cagagcccga acaagctttg atcaggtttc cctgcttccg acctgtcctg	2720
cctcagtttc cccatctgtg atgagcaggt gaccacgtta actctcaggg ctgtttgaag	2780
tctcttggtt gtaggccctc gccactgagt ggcccagggtg tgagaggtag tctagagccc	2840
tctgcggcct ttgtggaggt ccgttctcag catgtggcct ggtgtggctc cctaggcctg	2900
aggcctccca ctctcagtgc cccctgcccc cttgggaacc cacactccac ccagctaag	2960
cacaacactg tggaccacca ccaatggcac ctgagccacc ttgaccacca ttagtgtccc	3020
caccctcatt actctgccgt cgttgccctg cccaccaca tcatctccct ccatggctcc	3080
cacatgacg ctggcaggtg caagcggaga gcacgagggc agcagggcct ggccttccgt	3140
gtccctgcgt catccccagc ctcaactcccc acccgtgcag gcctggccag gaagggatcc	3200
tggcaggggc ttccaggttc tcagctcaag gcctgggtccc gggcaggcgt ccaaccctg	3260
ggagcaatgt atttctttgc ctccatcct gggcagaccc cttacaggcc tgggcattgc	3320
catgggccct gggcttccc aggctaagga gaaccaggaa cagctataac cttgagctag	3380

-154-

```

tgaaataggg tagatgagaa ggctgtctcc tccagacccc tacccttaca cagtggcccc 3440
acaatatgaa gacctggggg aattccaagg tgagcataga gcctgcctgt gccagttcc 3500
ttctggccct caggtggcca agcccatctc ttcacccctc agataggggc ccactcccag 3560
aagaagctgc tgggggtgggg gtgggaggct gcctaagcct gtctgtgctt cagaggcccc 3620
tccagtcctt ggctgtgggg taactggggg tatgagctgt ggccacaggt gagcaaggca 3680
gggaactgca atccagccct ggccgcggga ggggccatct ctggccaatg ctgctgtgcc 3740
ttcaaggact gacaagttac gtaggggcag aggtcgccag ctagccagtg tctcctccat 3800
ctggggggcg tctgtccact tgtcacctta ggttttccact catttgtcac cttgggggtt 3860
tgctctgtgt gtttcatatc caacggcaat acttgcaggg ggacagagtc ctctaaatac 3920
tccaatcctg cggttttttac aaacataaag ggggagaccc caagtggagg accctggggc 3980
tggagctccc tcccaaactt tgtccagcat ccagcctgtt ccttgggctc actggggagg 4040
gagttgtctt catagcacac ccagagccag ggatcccttt gtagtttttt gacaacggag 4100
catttctctt ctgtacagga cccaataaaa acttccttat gaaaaaaaaa aaaaaagggc 4160
ggccgc 4166

```

<210> 18

<211> 609

<212> PRT

<213> Homo sapiens

<400> 18

```

Met Ser Asp Glu Arg Arg Leu Pro Gly Ser Ala Val Gly Trp Leu Val
 1             5             10             15
Cys Gly Gly Leu Ser Leu Leu Ala Asn Ala Trp Gly Ile Leu Ser Val
          20             25             30
Gly Ala Lys Gln Lys Lys Trp Lys Pro Leu Glu Phe Leu Leu Cys Thr
          35             40             45
Leu Ala Ala Thr His Met Leu Asn Val Ala Val Pro Ile Ala Thr Tyr
          50             55             60
Ser Val Val Gln Leu Arg Arg Gln Arg Pro Asp Phe Glu Trp Asn Glu
          65             70             75             80
Gly Leu Cys Lys Val Phe Val Ser Thr Phe Tyr Thr Leu Thr Leu Ala
          85             90             95

```

Thr	Cys	Phe	Ser	Val	Thr	Ser	Leu	Ser	Tyr	His	Arg	Met	Trp	Met	Val	
			100					105					110			
Cys	Trp	Pro	Val	Asn	Tyr	Arg	Leu	Ser	Asn	Ala	Lys	Lys	Gln	Ala	Val	
		115					120					125				
His	Thr	Val	Met	Gly	Ile	Trp	Met	Val	Ser	Phe	Ile	Leu	Ser	Ala	Leu	
	130					135					140					
Pro	Ala	Val	Gly	Trp	His	Asp	Thr	Ser	Glu	Arg	Phe	Tyr	Thr	His	Gly	
145					150					155					160	
Cys	Arg	Phe	Ile	Val	Ala	Glu	Ile	Gly	Leu	Gly	Phe	Gly	Val	Cys	Phe	
			165						170					175		
Leu	Leu	Leu	Val	Gly	Gly	Ser	Val	Ala	Met	Gly	Val	Ile	Cys	Thr	Ala	
			180					185					190			
Ile	Ala	Leu	Phe	Gln	Thr	Leu	Ala	Val	Gln	Val	Gly	Arg	Gln	Ala	Asp	
		195					200					205				
His	Arg	Ala	Phe	Thr	Val	Pro	Thr	Ile	Val	Val	Glu	Asp	Ala	Gln	Gly	
	210					215					220					
Lys	Arg	Arg	Ser	Ser	Ile	Asp	Gly	Ser	Glu	Pro	Ala	Lys	Thr	Ser	Leu	
225					230					235					240	
Gln	Thr	Thr	Gly	Leu	Val	Thr	Thr	Ile	Val	Phe	Ile	Tyr	Asp	Cys	Leu	
			245						250					255		
Met	Gly	Phe	Pro	Val	Leu	Val	Val	Ser	Phe	Ser	Ser	Leu	Arg	Ala	Asp	
			260					265					270			
Ala	Ser	Ala	Pro	Trp	Met	Ala	Leu	Cys	Val	Leu	Trp	Cys	Ser	Val	Ala	
		275					280					285				
Gln	Ala	Leu	Leu	Leu	Pro	Val	Phe	Leu	Trp	Ala	Cys	Asp	Arg	Tyr	Arg	
	290					295					300					
Ala	Asp	Leu	Lys	Ala	Val	Arg	Glu	Lys	Cys	Met	Ala	Leu	Met	Ala	Asn	
305					310					315					320	
Asp	Glu	Glu	Ser	Asp	Asp	Glu	Thr	Ser	Leu	Glu	Gly	Gly	Ile	Ser	Pro	
			325						330					335		
Asp	Leu	Val	Leu	Glu	Arg	Ser	Leu	Asp	Tyr	Gly	Tyr	Gly	Gly	Asp	Phe	
			340					345					350			
Val	Ala	Leu	Asp	Arg	Met	Ala	Lys	Tyr	Glu	Ile	Ser	Ala	Leu	Glu	Gly	
		355					360					365				
Gly	Leu	Pro	Gln	Leu	Tyr	Pro	Leu	Arg	Pro	Leu	Gln	Glu	Asp	Lys	Met	
	370					375					380					

-156-

Gln Tyr Leu Gln Val Pro Pro Thr Arg Arg Phe Ser His Asp Asp Ala
 385 390 395 400
 Asp Val Trp Ala Ala Val Pro Leu Pro Ala Phe Leu Pro Arg Trp Gly
 405 410 415
 Ser Gly Glu Asp Leu Ala Ala Leu Ala His Leu Val Leu Pro Ala Gly
 420 425 430
 Pro Glu Arg Arg Arg Ala Ser Leu Leu Ala Phe Ala Glu Asp Ala Pro
 435 440 445
 Pro Ser Arg Ala Arg Arg Arg Ser Ala Glu Ser Leu Leu Ser Leu Arg
 450 455 460
 Thr Ser Ala Leu Asp Ser Gly Pro Arg Gly Ala Arg Asp Ser Pro Pro
 465 470 475 480
 Gly Ser Pro Arg Arg Arg Pro Gly Pro Gly Pro Arg Ser Ala Ser Ala
 485 490 495
 Ser Leu Leu Pro Asp Ala Phe Ala Leu Thr Ala Phe Glu Cys Glu Pro
 500 505 510
 Gln Ala Leu Arg Arg Pro Pro Gly Pro Phe Pro Ala Ala Pro Ala Ala
 515 520 525
 Pro Asp Gly Ala Asp Pro Gly Glu Ala Pro Thr Pro Pro Ser Ser Ala
 530 535 540
 Gln Arg Ser Pro Gly Pro Arg Pro Ser Ala His Ser His Ala Gly Ser
 545 550 555 560
 Leu Arg Pro Gly Leu Ser Ala Ser Trp Gly Glu Pro Gly Gly Leu Arg
 565 570 575
 Ala Ala Gly Gly Gly Gly Ser Thr Ser Ser Phe Leu Ser Ser Pro Ser
 580 585 590
 Glu Ser Ser Gly Tyr Ala Thr Leu His Ser Asp Ser Leu Gly Ser Ala
 595 600 605

Ser

<210> 19
 <211> 1827
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1) .. (1827)

-157-

<400> 19

atg agt gat gag cgg cgg ctg cct ggc agt gca gtg ggc tgg ctg gta	48
Met Ser Asp Glu Arg Arg Leu Pro Gly Ser Ala Val Gly Trp Leu Val	
1 5 10 15	
tgt ggg ggc ctc tcc ctg ctg gcc aat gcc tgg ggc atc ctc agc gtt	96
Cys Gly Gly Leu Ser Leu Leu Ala Asn Ala Trp Gly Ile Leu Ser Val	
20 25 30	
ggc gcc aag cag aag aag tgg aag ccc ttg gag ttc ctg ctg tgt acg	144
Gly Ala Lys Gln Lys Lys Trp Lys Pro Leu Glu Phe Leu Leu Cys Thr	
35 40 45	
ctc gcg gcc acc cac atg cta aat gtg gcc gtg ccc atc gcc acc tac	192
Leu Ala Ala Thr His Met Leu Asn Val Ala Val Pro Ile Ala Thr Tyr	
50 55 60	
tcc gtg gtg cag ctg cgg cgg cag cgc ccc gac ttc gag tgg aat gag	240
Ser Val Val Gln Leu Arg Arg Gln Arg Pro Asp Phe Glu Trp Asn Glu	
65 70 75 80	
ggc ctc tgc aag gtc ttc gtg tcc acc ttc tac acc ctc acc ctg gcc	288
Gly Leu Cys Lys Val Phe Val Ser Thr Phe Tyr Thr Leu Thr Leu Ala	
85 90 95	
acc tgt ttc tct gtc acc tcc ctc tcc tac cac cgc atg tgg atg gtc	336
Thr Cys Phe Ser Val Thr Ser Leu Ser Tyr His Arg Met Trp Met Val	
100 105 110	
tgc tgg cct gtc aac tac cgg ctg agc aat gcc aag aag cag gcg gtg	384
Cys Trp Pro Val Asn Tyr Arg Leu Ser Asn Ala Lys Lys Gln Ala Val	
115 120 125	
cac aca gtc atg ggt atc tgg atg gtg tcc ttc atc ctg tcg gcc ctg	432
His Thr Val Met Gly Ile Trp Met Val Ser Phe Ile Leu Ser Ala Leu	
130 135 140	
cct gcc gtt ggc tgg cac gac acc agc gag cgc ttc tac acc cat ggc	480
Pro Ala Val Gly Trp His Asp Thr Ser Glu Arg Phe Tyr Thr His Gly	
145 150 155 160	
tgc cgc ttc atc gtg gct gag atc ggc ctg ggc ttt ggc gtc tgc ttc	528
Cys Arg Phe Ile Val Ala Glu Ile Gly Leu Gly Phe Gly Val Cys Phe	
165 170 175	
ctg ctg ctg gtg ggc ggc agc gtg gcc atg ggc gtg atc tgc aca gcc	576
Leu Leu Leu Val Gly Gly Ser Val Ala Met Gly Val Ile Cys Thr Ala	
180 185 190	
atc gcc ctc ttc cag acg ctg gcc gtg cag gtg ggg cgc cag gcc gac	624
Ile Ala Leu Phe Gln Thr Leu Ala Val Gln Val Gly Arg Gln Ala Asp	
195 200 205	

-158-

cac cgc gcc ttc acc gtg ccc acc atc gtg gtg gag gac gcg cag ggc	672
His Arg Ala Phe Thr Val Pro Thr Ile Val Val Glu Asp Ala Gln Gly	
210 215 220	
aag cgg cgc tcc tcc atc gat ggc tcg gag ccc gcc aaa acc tct ctg	720
Lys Arg Arg Ser Ser Ile Asp Gly Ser Glu Pro Ala Lys Thr Ser Leu	
225 230 235 240	
cag acc acg ggc ctc gtg acc acc ata gtc ttc atc tac gac tgc ctc	768
Gln Thr Thr Gly Leu Val Thr Thr Ile Val Phe Ile Tyr Asp Cys Leu	
245 250 255	
atg ggc ttc cct gtg ctg gtg gtg agc ttc agc agc ctg cgg gcc gac	816
Met Gly Phe Pro Val Leu Val Val Ser Phe Ser Ser Leu Arg Ala Asp	
260 265 270	
gcc tca gcg ccc tgg atg gca ctc tgc gtg ctg tgg tgc tcc gtg gcc	864
Ala Ser Ala Pro Trp Met Ala Leu Cys Val Leu Trp Cys Ser Val Ala	
275 280 285	
cag gcc ctg ctg ctg cct gtg ttc ctc tgg gcc tgc gac cgc tac cgg	912
Gln Ala Leu Leu Leu Pro Val Phe Leu Trp Ala Cys Asp Arg Tyr Arg	
290 295 300	
gct gac ctc aaa gct gtc cgg gag aag tgc atg gcc ctc atg gcc aac	960
Ala Asp Leu Lys Ala Val Arg Glu Lys Cys Met Ala Leu Met Ala Asn	
305 310 315 320	
gac gag gag tca gac gat gag acc agc ctg gaa ggt ggc atc tcc ccg	1008
Asp Glu Glu Ser Asp Asp Glu Thr Ser Leu Glu Gly Gly Ile Ser Pro	
325 330 335	
gac ctg gtg ttg gag cgc tcc ctg gac tat ggc tat gga ggt gat ttt	1056
Asp Leu Val Leu Glu Arg Ser Leu Asp Tyr Gly Tyr Gly Gly Asp Phe	
340 345 350	
gtg gcc cta gat agg atg gcc aag tat gag atc tcc gcc ctg gag ggg	1104
Val Ala Leu Asp Arg Met Ala Lys Tyr Glu Ile Ser Ala Leu Glu Gly	
355 360 365	
ggc ctg ccc cag ctc tac cca ctg cgg ccc ttg cag gag gac aag atg	1152
Gly Leu Pro Gln Leu Tyr Pro Leu Arg Pro Leu Gln Glu Asp Lys Met	
370 375 380	
caa tac ctg cag gtc ccg ccc acg cgg cgc ttc tcc cac gac gat gcg	1200
Gln Tyr Leu Gln Val Pro Pro Thr Arg Arg Phe Ser His Asp Asp Ala	
385 390 395 400	
gac gtg tgg gcc gcc gtc ccg ctg ccc gcc ttc ctg ccg cgc tgg ggc	1248
Asp Val Trp Ala Ala Val Pro Leu Pro Ala Phe Leu Pro Arg Trp Gly	
405 410 415	
tcc ggc gag gac ctg gcc gcc ctg gcg cac ctg gtg ctg cct gcc ggg	1296
Ser Gly Glu Asp Leu Ala Ala Leu Ala His Leu Val Leu Pro Ala Gly	
420 425 430	

-159-

ccc	gag	cgg	cgc	cgc	gcc	agc	ctc	ctg	gcc	ttc	gcg	gag	gac	gca	cca	1344
Pro	Glu	Arg	Arg	Arg	Ala	Ser	Leu	Leu	Ala	Phe	Ala	Glu	Asp	Ala	Pro	
		435					440					445				

ccg	tcc	cgc	gcg	cgc	cgc	cgc	tcg	gcc	gag	agc	ctg	ctg	tcg	ctg	cgg	1392
Pro	Ser	Arg	Ala	Arg	Arg	Arg	Ser	Ala	Glu	Ser	Leu	Leu	Ser	Leu	Arg	
	450					455					460					

acc	tcg	gcc	ctg	gat	agc	ggc	ccg	cgg	gga	gcc	cgc	gac	tcg	ccc	ccc	1440
Thr	Ser	Ala	Leu	Asp	Ser	Gly	Pro	Arg	Gly	Ala	Arg	Asp	Ser	Pro	Pro	
465					470					475					480	

ggc	agc	ccg	cgc	cgc	cgc	ccc	ggg	ccc	ggc	ccc	cgc	tcc	gcc	tcg	gcc	1488
Gly	Ser	Pro	Arg	Arg	Arg	Pro	Gly	Pro	Gly	Pro	Arg	Ser	Ala	Ser	Ala	
			485					490						495		

tcg	ctg	ctg	ccc	gac	gcc	ttc	gcc	ctg	acc	gcc	ttc	gag	tgc	gag	cca	1536
Ser	Leu	Leu	Pro	Asp	Ala	Phe	Ala	Leu	Thr	Ala	Phe	Glu	Cys	Glu	Pro	
			500					505					510			

cag	gcc	ctg	cgc	cgc	ccg	ccc	ggg	ccc	ttc	ccc	gct	gcg	ccc	gcc	gcc	1584
Gln	Ala	Leu	Arg	Arg	Pro	Pro	Gly	Pro	Phe	Pro	Ala	Ala	Pro	Ala	Ala	
		515					520				525					

ccc	gac	ggc	gca	gat	ccc	gga	gag	gcc	ccg	acg	ccc	cca	agc	agc	gcc	1632
Pro	Asp	Gly	Ala	Asp	Pro	Gly	Glu	Ala	Pro	Thr	Pro	Pro	Ser	Ser	Ala	
	530					535					540					

cag	cgg	agc	cca	ggg	cca	cgc	ccc	tct	gcg	cac	tcg	cac	gcc	ggc	tct	1680
Gln	Arg	Ser	Pro	Gly	Pro	Arg	Pro	Ser	Ala	His	Ser	His	Ala	Gly	Ser	
545					550					555					560	

ctg	cgc	ccc	ggc	ctg	agc	gcg	tcg	tgg	ggc	gag	ccc	ggg	ggg	ctg	cgc	1728
Leu	Arg	Pro	Gly	Leu	Ser	Ala	Ser	Trp	Gly	Glu	Pro	Gly	Gly	Leu	Arg	
				565				570						575		

gcg	gcg	ggc	ggc	ggc	ggc	agc	acc	agc	agc	ttc	ctg	agt	tcc	ccc	tcc	1776
Ala	Ala	Gly	Gly	Gly	Gly	Ser	Thr	Ser	Ser	Phe	Leu	Ser	Ser	Pro	Ser	
			580					585					590			

gag	tcc	tcg	ggc	tac	gcc	acg	ctg	cac	tcg	gac	tcg	ctg	ggc	tcc	gcg	1824
Glu	Ser	Ser	Gly	Tyr	Ala	Thr	Leu	His	Ser	Asp	Ser	Leu	Gly	Ser	Ala	
		595				600						605				

tcc																1827
Ser																

<210> 20

<211> 177

<212> PRT

<213> Homo sapiens

<220>

<223> Xaa's at postions 38, 55 and 56 may be any amino acid

-160-

<400> 20

Gly	Asn	Ile	Leu	Val	Ile	Trp	Val	Ile	Cys	Arg	Tyr	Arg	Arg	Met	Arg
1				5					10					15	
Thr	Pro	Met	Asn	Tyr	Phe	Ile	Val	Asn	Leu	Ala	Val	Ala	Asp	Leu	Leu
			20					25					30		
Phe	Ser	Leu	Phe	Thr	Xaa	Met	Pro	Phe	Trp	Met	Val	Tyr	Tyr	Val	Met
		35					40					45			
Gln	Gly	Arg	Trp	Pro	Phe	Xaa	Xaa	Gly	Asp	Phe	Met	Cys	Arg	Ile	Trp
	50					55					60				
Met	Tyr	Phe	Asp	Tyr	Met	Asn	Met	Tyr	Ala	Ser	Ile	Phe	Phe	Leu	Thr
65					70					75					80
Cys	Ile	Ser	Ile	Asp	Arg	Tyr	Leu	Trp	Ala	Ile	Cys	His	Pro	Met	Arg
				85					90					95	
Tyr	Met	Arg	Trp	Met	Thr	Pro	Arg	His	Arg	Ala	Trp	Val	Met	Ile	Ile
			100					105					110		
Ile	Ile	Trp	Val	Met	Ser	Phe	Leu	Ile	Ser	Met	Pro	Pro	Phe	Leu	Met
		115					120					125			
Phe	Arg	Trp	Ser	Thr	Tyr	Arg	Asp	Glu	Asn	Glu	Trp	Asn	Met	Thr	Trp
	130					135					140				
Cys	Met	Ile	Tyr	Asp	Trp	Pro	Glu	Trp	Met	Trp	Arg	Trp	Tyr	Val	Ile
145					150					155					160
Leu	Met	Thr	Ile	Ile	Met	Gly	Phe	Tyr	Ile	Pro	Met	Ile	Ile	Met	Leu
				165					170					175	

Phe

<210> 21

<211> 107

<212> PRT

<213> Homo sapiens

<220>

<223> Xaa at postion 84 may be any amino
acid

<400> 21

Ile	Gln	Glu	Arg	Met	Asn	Glu	Leu	Asn	Asp	Arg	Trp	Glu	Arg	Leu	Lys
1				5					10					15	
Glu	Leu	Met	Glu	Gln	Arg	Arg	Gln	Met	Leu	Glu	Asp	Ser	Met	Arg	Leu
			20					25					30		

-161-

Gln Gln Phe Phe Arg Asp Met Asp Glu Glu Glu Ser Trp Ile Asn Glu
35 40 45

Lys Glu Gln Ile Leu Asn Ser Asp Asp Tyr Gly Lys Asp Leu Thr Ser
50 55 60

Val Gln Asn Leu Leu Lys Lys His Gln Ala Phe Glu Ala Asp Ile Ala
65 70 75 80

Ala His Glu Xaa Asp Arg Ile Gln Ala Leu Asn Glu Phe Ala Gln Gln
85 90 95

Leu Ile Gln Glu Asn His Tyr Ala Ser Glu Glu
100 105

```
<210> 22
<211> 168
<212> PRT
<213> Homo sapiens
```

<400> 22

Ala Asn Ala Trp Gly Ile Leu Ser Val Gly Ala Lys Gln Lys Lys Trp
1 5 10 15

Lys Pro Leu Glu Phe Leu Leu Cys Thr Leu Ala Ala Thr His Met Leu
20 25 30

Asn Val Ala Val Pro Ile Ala Thr Tyr Ser Val Val Gln Leu Arg Arg
35 40 45

Gln Arg Pro Asp Phe Glu Trp Asn Glu Gly Leu Cys Lys Val Phe Val
50 55 60

Ser Thr Phe Tyr Thr Leu Thr Leu Ala Thr Cys Phe Ser Val Thr Ser
65 70 75 80

Leu Ser Tyr His Arg Met Trp Met Val Cys Trp Pro Val Asn Tyr Arg
85 90 95

Leu Ser Asn Ala Lys Lys Gln Ala Val His Thr Val Met Gly Ile Trp
100 105 110

Met Val Ser Phe Ile Leu Ser Ala Leu Pro Ala Val Gly Trp His Asp
115 120 125

Thr Ser Glu Arg Phe Tyr Thr His Gly Cys Arg Phe Ile Val Ala Glu
130 135 140

Ile Gly Leu Gly Phe Gly Val Cys Phe Leu Leu Leu Val Gly Gly Ser
145 150 155 160

-162-

Val Ala Met Gly Val Ile Cys Thr
165

<210> 23
<211> 107
<212> PRT
<213> Homo sapiens

<400> 23

Phe Ser Ser Leu Arg Ala Asp Ala Ser Ala Pro Trp Met Ala Leu Cys
1 5 10 15
Val Leu Trp Cys Ser Val Ala Gln Ala Leu Leu Leu Pro Val Phe Leu
20 25 30
Trp Ala Cys Asp Arg Tyr Arg Ala Asp Leu Lys Ala Val Arg Glu Lys
35 40 45
Cys Met Ala Leu Met Ala Asn Asp Glu Glu Ser Asp Asp Glu Thr Ser
50 55 60
Leu Glu Gly Gly Ile Ser Pro Asp Leu Val Leu Glu Arg Ser Leu Asp
65 70 75 80
Tyr Gly Tyr Gly Gly Asp Phe Val Ala Leu Asp Arg Met Ala Lys Tyr
85 90 95
Glu Ile Ser Ala Leu Glu Gly Gly Leu Pro Gln
100 105

<210> 24
<211> 425
<212> PRT
<213> Homo sapiens

<400> 24

Met Gly Pro Arg Arg Leu Leu Leu Val Ala Ala Cys Phe Ser Leu Cys
1 5 10 15
Gly Pro Leu Leu Ser Ala Arg Thr Arg Ala Arg Arg Pro Glu Ser Lys
20 25 30
Ala Thr Asn Ala Thr Leu Asp Pro Arg Ser Phe Leu Leu Arg Asn Pro
35 40 45
Asn Asp Lys Tyr Glu Pro Phe Trp Glu Asp Glu Glu Lys Asn Glu Ser
50 55 60
Gly Leu Thr Glu Tyr Arg Leu Val Ser Ile Asn Lys Ser Ser Pro Leu
65 70 75 80

-163-

Gln	Lys	Gln	Leu	Pro	Ala	Phe	Ile	Ser	Glu	Asp	Ala	Ser	Gly	Tyr	Leu	85	90	95
Thr	Ser	Ser	Trp	Leu	Thr	Leu	Phe	Val	Pro	Ser	Val	Tyr	Thr	Gly	Val	100	105	110
Phe	Val	Val	Ser	Leu	Pro	Leu	Asn	Ile	Met	Ala	Ile	Val	Val	Phe	Ile	115	120	125
Leu	Lys	Met	Lys	Val	Lys	Lys	Pro	Ala	Val	Val	Tyr	Met	Leu	His	Leu	130	135	140
Ala	Thr	Ala	Asp	Val	Leu	Phe	Val	Ser	Val	Leu	Pro	Phe	Lys	Ile	Ser	145	150	155
Tyr	Tyr	Phe	Ser	Gly	Ser	Asp	Trp	Gln	Phe	Gly	Ser	Glu	Leu	Cys	Arg	165	170	175
Phe	Val	Thr	Ala	Ala	Phe	Tyr	Cys	Asn	Met	Tyr	Ala	Ser	Ile	Leu	Leu	180	185	190
Met	Thr	Val	Ile	Ser	Ile	Asp	Arg	Phe	Leu	Ala	Val	Val	Tyr	Pro	Met	195	200	205
Gln	Ser	Leu	Ser	Trp	Arg	Thr	Leu	Gly	Arg	Ala	Ser	Phe	Thr	Cys	Leu	210	215	220
Ala	Ile	Trp	Ala	Leu	Ala	Ile	Ala	Gly	Val	Val	Pro	Leu	Val	Leu	Lys	225	230	235
Glu	Gln	Thr	Ile	Gln	Val	Pro	Gly	Leu	Asn	Ile	Thr	Thr	Cys	His	Asp	245	250	255
Val	Leu	Asn	Glu	Thr	Leu	Leu	Glu	Gly	Tyr	Tyr	Ala	Tyr	Tyr	Phe	Ser	260	265	270
Ala	Phe	Ser	Ala	Val	Phe	Phe	Phe	Val	Pro	Leu	Ile	Ile	Ser	Thr	Val	275	280	285
Cys	Tyr	Val	Ser	Ile	Ile	Arg	Cys	Leu	Ser	Ser	Ser	Ala	Val	Ala	Asn	290	295	300
Arg	Ser	Lys	Lys	Ser	Arg	Ala	Leu	Phe	Leu	Ser	Ala	Ala	Val	Phe	Cys	305	310	315
Ile	Phe	Ile	Ile	Cys	Phe	Gly	Pro	Thr	Asn	Val	Leu	Leu	Ile	Ala	His	325	330	335
Tyr	Ser	Phe	Leu	Ser	His	Thr	Ser	Thr	Thr	Glu	Ala	Ala	Tyr	Phe	Ala	340	345	350
Tyr	Leu	Leu	Cys	Val	Cys	Val	Ser	Ser	Ile	Ser	Ser	Cys	Ile	Asp	Pro	355	360	365

-164-

Leu Ile Tyr Tyr Tyr Ala Ser Ser Glu Cys Gln Arg Tyr Val Tyr Ser
 370 375 380

Ile Leu Cys Cys Lys Glu Ser Ser Asp Pro Ser Ser Tyr Asn Ser Ser
 385 390 395 400

Gly Gln Leu Met Ala Ser Lys Met Asp Thr Cys Ser Ser Asn Leu Asn
 405 410 415

Asn Ser Ile Tyr Lys Lys Leu Leu Thr
 420 425

<210> 25

<211> 348

<212> PRT

<213> Homo sapiens

<400> 25

Met Asn Gly Thr Glu Gly Pro Asn Phe Tyr Val Pro Phe Ser Asn Ala
 1 5 10 15

Thr Gly Val Val Arg Ser Pro Phe Glu Tyr Pro Gln Tyr Tyr Leu Ala
 20 25 30

Glu Pro Trp Gln Phe Ser Met Leu Ala Ala Tyr Met Phe Leu Leu Ile
 35 40 45

Val Leu Gly Phe Pro Ile Asn Phe Leu Thr Leu Tyr Val Thr Val Gln
 50 55 60

His Lys Lys Leu Arg Thr Pro Leu Asn Tyr Ile Leu Leu Asn Leu Ala
 65 70 75 80

Val Ala Asp Leu Phe Met Val Leu Gly Gly Phe Thr Ser Thr Leu Tyr
 85 90 95

Thr Ser Leu His Gly Tyr Phe Val Phe Gly Pro Thr Gly Cys Asn Leu
 100 105 110

Glu Gly Phe Phe Ala Thr Leu Gly Gly Glu Ile Ala Leu Trp Ser Leu
 115 120 125

Val Val Leu Ala Ile Glu Arg Tyr Val Val Val Cys Lys Pro Met Ser
 130 135 140

Asn Phe Arg Phe Gly Glu Asn His Ala Ile Met Gly Val Ala Phe Thr
 145 150 155 160

Trp Val Met Ala Leu Ala Cys Ala Ala Pro Pro Leu Ala Gly Trp Ser
 165 170 175

Arg Tyr Ile Pro Glu Gly Leu Gln Cys Ser Cys Gly Ile Asp Tyr Tyr
 180 185 190

-165-

Thr Leu Lys Pro Glu Val Asn Asn Glu Ser Phe Val Ile Tyr Met Phe
 195 200 205
 Val Val His Phe Thr Ile Pro Met Ile Ile Ile Phe Phe Cys Tyr Gly
 210 215 220
 Gln Leu Val Phe Thr Val Lys Glu Ala Ala Ala Gln Gln Gln Glu Ser
 225 230 235 240
 Ala Thr Thr Gln Lys Ala Glu Lys Glu Val Thr Arg Met Val Ile Ile
 245 250 255
 Met Val Ile Ala Phe Leu Ile Cys Trp Val Pro Tyr Ala Ser Val Ala
 260 265 270
 Phe Tyr Ile Phe Thr His Gln Gly Ser Asn Phe Gly Pro Ile Phe Met
 275 280 285
 Thr Ile Pro Ala Phe Phe Ala Lys Ser Ala Ala Ile Tyr Asn Pro Val
 290 295 300
 Ile Tyr Ile Met Met Asn Lys Gln Phe Arg Asn Cys Met Leu Thr Thr
 305 310 315 320
 Ile Cys Cys Gly Lys Asn Pro Leu Gly Asp Asp Glu Ala Ser Ala Thr
 325 330 335
 Val Ser Lys Thr Glu Thr Ser Gln Val Ala Pro Ala
 340 345

<210> 26
 <211> 460
 <212> PRT
 <213> Homo sapiens

<400> 26

Met Asn Thr Ser Val Pro Pro Ala Val Ser Pro Asn Ile Thr Val Leu
 1 5 10 15
 Ala Pro Gly Lys Gly Pro Trp Gln Val Ala Phe Ile Gly Ile Thr Thr
 20 25 30
 Gly Leu Leu Ser Leu Ala Thr Val Thr Gly Asn Leu Leu Val Leu Ile
 35 40 45
 Ser Phe Lys Val Asn Thr Glu Leu Lys Thr Val Asn Asn Tyr Phe Leu
 50 55 60
 Leu Ser Leu Ala Cys Ala Asp Leu Ile Ile Gly Thr Phe Ser Met Asn
 65 70 75 80
 Leu Tyr Thr Thr Tyr Leu Leu Met Gly His Trp Ala Leu Gly Thr Leu
 85 90 95

-166-

Ala	Cys	Asp	Leu	Trp	Leu	Ala	Leu	Asp	Tyr	Val	Ala	Ser	Asn	Ala	Ser	100	105	110	
Val	Met	Asn	Leu	Leu	Leu	Ile	Ser	Phe	Asp	Arg	Tyr	Phe	Ser	Val	Thr	115	120	125	
Arg	Pro	Leu	Ser	Tyr	Arg	Ala	Lys	Arg	Thr	Pro	Arg	Arg	Ala	Ala	Leu	130	135	140	
Met	Ile	Gly	Leu	Ala	Trp	Leu	Val	Ser	Phe	Val	Leu	Trp	Ala	Pro	Ala	145	150	155	160
Ile	Leu	Phe	Trp	Gln	Tyr	Leu	Val	Gly	Glu	Arg	Thr	Val	Leu	Ala	Gly	165	170	175	
Gln	Cys	Tyr	Ile	Gln	Phe	Leu	Ser	Gln	Pro	Ile	Ile	Thr	Phe	Gly	Thr	180	185	190	
Ala	Met	Ala	Ala	Phe	Tyr	Leu	Pro	Val	Thr	Val	Met	Cys	Thr	Leu	Tyr	195	200	205	
Trp	Arg	Ile	Tyr	Arg	Glu	Thr	Glu	Asn	Arg	Ala	Arg	Glu	Leu	Ala	Ala	210	215	220	
Leu	Gln	Gly	Ser	Glu	Thr	Pro	Gly	Lys	Gly	Gly	Gly	Ser	Ser	Ser	Ser	225	230	235	240
Ser	Glu	Arg	Ser	Gln	Pro	Gly	Ala	Glu	Gly	Ser	Pro	Glu	Ser	Pro	Pro	245	250	255	
Gly	Arg	Cys	Cys	Arg	Cys	Cys	Arg	Ala	Pro	Arg	Leu	Leu	Gln	Ala	Tyr	260	265	270	
Ser	Trp	Lys	Glu	Glu	Glu	Glu	Glu	Asp	Glu	Gly	Ser	Met	Glu	Ser	Leu	275	280	285	
Thr	Ser	Ser	Glu	Gly	Glu	Glu	Pro	Gly	Ser	Glu	Val	Val	Ile	Lys	Met	290	295	300	
Pro	Met	Val	Asp	Ser	Glu	Ala	Gln	Ala	Pro	Thr	Lys	Gln	Pro	Pro	Lys	305	310	315	320
Ser	Ser	Pro	Asn	Thr	Val	Lys	Arg	Pro	Thr	Lys	Lys	Gly	Arg	Asp	Arg	325	330	335	
Gly	Gly	Lys	Gly	Gln	Lys	Pro	Arg	Gly	Lys	Glu	Gln	Leu	Ala	Lys	Arg	340	345	350	
Lys	Thr	Phe	Ser	Leu	Val	Lys	Glu	Lys	Lys	Ala	Ala	Arg	Thr	Leu	Ser	355	360	365	
Ala	Ile	Leu	Leu	Ala	Phe	Ile	Leu	Thr	Trp	Thr	Pro	Tyr	Asn	Ile	Met	370	375	380	

-167-

Val Leu Val Ser Thr Phe Cys Lys Asp Cys Val Pro Glu Thr Leu Trp
 385 390 395 400

Glu Leu Gly Tyr Trp Leu Cys Tyr Val Asn Ser Thr Val Asn Pro Met
 405 410 415

Cys Tyr Ala Leu Cys Asn Lys Ala Phe Arg Asp Thr Phe Arg Leu Leu
 420 425 430

Leu Leu Cys Arg Trp Asp Lys Arg Arg Trp Arg Lys Ile Pro Lys Arg
 435 440 445

Pro Gly Ser Val His Arg Thr Pro Ser Arg Gln Cys
 450 455 460

<210> 27

<211> 350

<212> PRT

<213> Rattus norvegicus

<400> 27

Met Ser Asn Ile Thr Asp Pro Gln Met Trp Asp Phe Asp Asp Leu Asn
 1 5 10 15

Phe Thr Gly Met Pro Pro Ala Asp Glu Asp Tyr Ser Pro Cys Met Leu
 20 25 30

Glu Thr Glu Thr Leu Asn Lys Tyr Val Val Ile Ile Ala Tyr Ala Leu
 35 40 45

Val Phe Leu Leu Ser Leu Leu Gly Asn Ser Leu Val Met Leu Val Ile
 50 55 60

Leu Tyr Ser Arg Val Gly Arg Ser Val Thr Asp Val Tyr Leu Leu Asn
 65 70 75 80

Leu Ala Leu Ala Asp Leu Leu Phe Ala Leu Thr Leu Pro Ile Trp Ala
 85 90 95

Ala Ser Lys Val Asn Gly Trp Ile Phe Gly Thr Phe Leu Cys Lys Val
 100 105 110

Val Ser Leu Leu Lys Glu Val Asn Phe Tyr Ser Gly Ile Leu Leu Leu
 115 120 125

Ala Cys Ile Ser Val Asp Arg Tyr Leu Ala Ile Val His Ala Thr Arg
 130 135 140

Thr Leu Thr Gln Lys Arg His Leu Val Lys Phe Val Cys Leu Gly Cys
 145 150 155 160

Trp Gly Leu Ser Met Asn Leu Ser Leu Pro Phe Phe Leu Phe Arg Gln
 165 170 175

-168-

Ala	Tyr	His	Pro	Asn	Asn	Ser	Ser	Pro	Val	Cys	Tyr	Glu	Val	Leu	Gly
			180					185					190		
Asn	Asp	Thr	Ala	Lys	Trp	Arg	Met	Val	Leu	Arg	Ile	Leu	Pro	His	Thr
		195					200					205			
Phe	Gly	Phe	Ile	Val	Pro	Leu	Phe	Val	Met	Leu	Phe	Cys	Tyr	Gly	Phe
	210					215					220				
Thr	Leu	Arg	Thr	Leu	Phe	Lys	Ala	His	Met	Gly	Gln	Lys	His	Arg	Ala
225					230					235					240
Met	Arg	Val	Ile	Phe	Ala	Val	Val	Leu	Ile	Phe	Leu	Leu	Cys	Trp	Leu
				245					250					255	
Pro	Tyr	Asn	Leu	Val	Leu	Leu	Ala	Asp	Thr	Leu	Met	Arg	Thr	Gln	Val
			260					265					270		
Ile	Gln	Glu	Thr	Cys	Glu	Arg	Arg	Asn	Asn	Ile	Gly	Arg	Ala	Leu	Asp
		275					280					285			
Ala	Thr	Glu	Ile	Leu	Gly	Phe	Leu	His	Ser	Cys	Leu	Asn	Pro	Ile	Ile
	290					295					300				
Tyr	Ala	Phe	Ile	Gly	Gln	Asn	Phe	Arg	His	Gly	Phe	Leu	Lys	Ile	Leu
305					310					315					320
Ala	Met	His	Gly	Leu	Val	Ser	Lys	Glu	Phe	Leu	Ala	Arg	His	Arg	Val
				325					330					335	
Thr	Ser	Tyr	Thr	Ser	Ser	Ser	Val	Asn	Val	Ser	Ser	Asn	Leu		
			340					345					350		

<210> 28

<211> 601

<212> PRT

<213> Drosophila melanogaster

<400> 28

Met	Pro	Ser	Ala	Asp	Gln	Ile	Leu	Phe	Val	Asn	Val	Thr	Thr	Thr	Val
1				5					10					15	
Ala	Ala	Ala	Ala	Leu	Thr	Ala	Ala	Ala	Ala	Val	Ser	Thr	Thr	Lys	Ser
			20					25					30		
Gly	Asn	Gly	Asn	Ala	Ala	Arg	Gly	Tyr	Thr	Asp	Ser	Asp	Asp	Asp	Ala
		35					40					45			
Gly	Met	Gly	Thr	Glu	Ala	Val	Ala	Asn	Ile	Ser	Gly	Ser	Leu	Val	Glu
	50					55					60				
Gly	Leu	Thr	Thr	Val	Thr	Ala	Ala	Leu	Ser	Thr	Ala	Gln	Ala	Asp	Lys
65					70					75					80

-169-

Asp	Ser	Ala	Gly	Glu	Cys	Glu	Gly	Ala	Val	Glu	Glu	Leu	His	Ala	Ser	85	90	95	
Ile	Leu	Gly	Leu	Gln	Leu	Ala	Val	Pro	Glu	Trp	Glu	Ala	Leu	Leu	Thr	100	105	110	
Ala	Leu	Val	Leu	Ser	Val	Ile	Ile	Val	Leu	Thr	Ile	Ile	Gly	Asn	Ile	115	120	125	
Leu	Val	Ile	Leu	Ser	Val	Phe	Thr	Tyr	Lys	Pro	Leu	Arg	Ile	Val	Gln	130	135	140	
Asn	Phe	Phe	Ile	Val	Ser	Leu	Ala	Val	Ala	Asp	Leu	Thr	Val	Ala	Leu	145	150	155	160
Leu	Val	Leu	Pro	Phe	Asn	Val	Ala	Tyr	Ser	Ile	Leu	Gly	Arg	Trp	Glu	165	170	175	
Phe	Gly	Ile	His	Leu	Cys	Lys	Leu	Trp	Leu	Thr	Cys	Asp	Val	Leu	Cys	180	185	190	
Cys	Thr	Ser	Ser	Ile	Leu	Asn	Leu	Cys	Ala	Ile	Ala	Leu	Asp	Arg	Tyr	195	200	205	
Trp	Ala	Ile	Thr	Asp	Pro	Ile	Asn	Tyr	Ala	Gln	Lys	Arg	Thr	Val	Gly	210	215	220	
Arg	Val	Leu	Leu	Leu	Ile	Ser	Gly	Val	Trp	Leu	Leu	Ser	Leu	Leu	Ile	225	230	235	240
Ser	Ser	Pro	Pro	Leu	Ile	Gly	Trp	Asn	Asp	Trp	Pro	Asp	Glu	Phe	Thr	245	250	255	
Ser	Ala	Thr	Pro	Cys	Glu	Leu	Thr	Ser	Gln	Arg	Gly	Tyr	Val	Ile	Tyr	260	265	270	
Ser	Ser	Leu	Gly	Ser	Phe	Phe	Ile	Pro	Leu	Ala	Ile	Met	Thr	Ile	Val	275	280	285	
Tyr	Ile	Glu	Ile	Phe	Val	Ala	Thr	Arg	Arg	Arg	Leu	Arg	Glu	Arg	Ala	290	295	300	
Arg	Ala	Asn	Lys	Leu	Asn	Thr	Ile	Ala	Leu	Lys	Ser	Thr	Glu	Leu	Glu	305	310	315	320
Pro	Met	Ala	Asn	Ser	Ser	Pro	Val	Ala	Ala	Ser	Asn	Ser	Gly	Ser	Lys	325	330	335	
Ser	Arg	Leu	Leu	Ala	Ser	Trp	Leu	Cys	Cys	Gly	Arg	Asp	Arg	Ala	Gln	340	345	350	
Phe	Ala	Thr	Pro	Met	Ile	Gln	Asn	Asp	Gln	Glu	Ser	Ile	Ser	Ser	Glu	355	360	365	

-170-

Thr His Gln Pro Gln Asp Ser Ser Lys Ala Gly Pro His Gly Asn Ser
 370 375 380
 Asp Pro Gln Gln Gln His Val Val Val Leu Val Lys Lys Ser Arg Arg
 385 390 395 400
 Ala Lys Thr Lys Asp Ser Ile Lys His Gly Lys Thr Arg Gly Gly Arg
 405 410 415
 Lys Ser Gln Ser Ser Ser Thr Cys Glu Pro His Gly Glu Gln Gln Leu
 420 425 430
 Leu Pro Ala Gly Gly Asp Gly Gly Ser Cys Gln Pro Gly Gly Gly His
 435 440 445
 Ser Gly Gly Gly Lys Ser Asp Ala Glu Ile Ser Thr Glu Ser Gly Ser
 450 455 460
 Asp Pro Lys Gly Cys Ile Gln Val Cys Val Thr Gln Ala Asp Glu Gln
 465 470 475 480
 Thr Ser Leu Lys Leu Thr Pro Pro Gln Ser Ser Thr Gly Val Ala Ala
 485 490 495
 Val Ser Val Thr Pro Leu Gln Lys Lys Thr Ser Gly Val Asn Gln Phe
 500 505 510
 Ile Glu Glu Lys Gln Lys Ile Ser Leu Ser Lys Glu Arg Arg Ala Ala
 515 520 525
 Arg Thr Leu Gly Ile Ile Met Gly Val Phe Val Ile Cys Trp Leu Pro
 530 535 540
 Phe Phe Leu Met Tyr Val Ile Leu Pro Phe Cys Gln Thr Cys Cys Pro
 545 550 555 560
 Thr Asn Lys Phe Lys Asn Phe Ile Thr Trp Leu Gly Tyr Ile Asn Ser
 565 570 575
 Gly Leu Asn Pro Val Ile Tyr Thr Ile Phe Asn Leu Asp Tyr Arg Arg
 580 585 590
 Ala Phe Lys Arg Leu Leu Gly Leu Asn
 595 600

<210> 29

<211> 588

<212> PRT

<213> Homo sapiens

<400> 29

Met Ala Arg Gly Gly Ala Gly Ala Glu Glu Ala Ser Leu Arg Ser Asn
 1 5 10 15

-171-

Ala	Leu	Ser	Trp	Leu	Ala	Cys	Gly	Leu	Leu	Ala	Leu	Leu	Ala	Asn	Ala	
			20					25					30			
Trp	Ile	Ile	Leu	Ser	Ile	Ser	Ala	Lys	Gln	Gln	Lys	His	Lys	Pro	Leu	
		35					40					45				
Glu	Leu	Leu	Leu	Cys	Phe	Leu	Ala	Gly	Thr	His	Ile	Leu	Met	Ala	Ala	
	50					55					60					
Val	Pro	Leu	Thr	Thr	Phe	Ala	Val	Val	Gln	Leu	Arg	Arg	Gln	Ala	Ser	
65					70					75					80	
Ser	Asp	Tyr	Asp	Trp	Asn	Glu	Ser	Ile	Cys	Lys	Val	Phe	Val	Ser	Thr	
				85					90					95		
Tyr	Tyr	Thr	Leu	Ala	Leu	Ala	Thr	Cys	Phe	Thr	Val	Ala	Ser	Leu	Ser	
			100					105					110			
Tyr	His	Arg	Met	Trp	Met	Val	Arg	Trp	Pro	Val	Asn	Tyr	Arg	Leu	Ser	
		115					120					125				
Asn	Ala	Lys	Lys	Gln	Ala	Leu	His	Ala	Val	Met	Gly	Ile	Trp	Met	Val	
	130					135					140					
Ser	Phe	Ile	Leu	Ser	Thr	Leu	Pro	Ser	Ile	Gly	Trp	His	Asn	Asn	Gly	
145					150					155					160	
Glu	Arg	Tyr	Tyr	Ala	Arg	Gly	Cys	Gln	Phe	Ile	Val	Ser	Lys	Ile	Gly	
				165					170					175		
Leu	Gly	Phe	Gly	Val	Cys	Phe	Ser	Leu	Leu	Leu	Leu	Gly	Gly	Ile	Val	
			180					185					190			
Met	Gly	Leu	Val	Cys	Val	Ala	Ile	Thr	Phe	Tyr	Gln	Thr	Leu	Trp	Ala	
		195					200					205				
Arg	Pro	Arg	Arg	Ala	Arg	Gln	Ala	Arg	Arg	Val	Gly	Gly	Gly	Gly	Gly	
	210					215					220					
Thr	Lys	Ala	Gly	Gly	Pro	Gly	Ala	Leu	Gly	Thr	Arg	Pro	Ala	Phe	Glu	
225					230					235					240	
Val	Pro	Ala	Ile	Val	Val	Glu	Asp	Ala	Arg	Gly	Lys	Arg	Arg	Ser	Ser	
				245					250					255		
Leu	Asp	Gly	Ser	Glu	Ser	Ala	Lys	Thr	Ser	Leu	Gln	Val	Thr	Asn	Leu	
			260					265					270			
Val	Ser	Ala	Ile	Val	Phe	Leu	Tyr	Asp	Ser	Leu	Thr	Gly	Val	Pro	Ile	
		275					280					285				
Leu	Val	Val	Ser	Phe	Phe	Ser	Leu	Lys	Ser	Asp	Ser	Ala	Pro	Pro	Trp	
	290					295						300				

-172-

Met Val Leu Ala Val Leu Trp Cys Ser Met Ala Gln Thr Leu Leu Leu
 305 310 315 320

Pro Ser Phe Ile Trp Ser Cys Glu Arg Tyr Arg Ala Asp Val Arg Thr
 325 330 335

Val Trp Glu Gln Cys Val Ala Ile Met Ser Glu Glu Asp Gly Asp Asp
 340 345 350

Asp Gly Gly Cys Asp Asp Tyr Ala Glu Gly Arg Val Cys Lys Val Arg
 355 360 365

Phe Asp Ala Asn Gly Ala Thr Gly Pro Gly Ser Arg Asp Pro Ala Gln
 370 375 380

Val Lys Leu Leu Pro Gly Arg His Met Leu Phe Pro Pro Leu Glu Arg
 385 390 395 400

Val His Tyr Leu Gln Val Pro Leu Ser Arg Arg Leu Ser His Asp Glu
 405 410 415

Thr Asn Ile Phe Ser Thr Pro Arg Glu Pro Gly Ser Phe Leu His Lys
 420 425 430

Trp Ser Ser Ser Asp Asp Ile Arg Val Leu Pro Ala Gln Ser Arg Ala
 435 440 445

Leu Gly Gly Pro Pro Glu Tyr Leu Gly Gln Arg His Arg Leu Glu Asp
 450 455 460

Glu Glu Asp Glu Glu Glu Ala Glu Gly Gly Gly Leu Ala Ser Leu Arg
 465 470 475 480

Gln Phe Leu Glu Ser Gly Val Leu Gly Ser Gly Gly Gly Pro Pro Arg
 485 490 495

Gly Pro Gly Phe Phe Arg Glu Glu Ile Thr Thr Phe Ile Asp Glu Thr
 500 505 510

Pro Leu Pro Ser Pro Thr Ala Ser Pro Gly His Ser Pro Arg Arg Pro
 515 520 525

Arg Pro Leu Gly Leu Ser Pro Arg Arg Leu Ser Leu Gly Ser Pro Glu
 530 535 540

Ser Arg Ala Val Gly Leu Pro Leu Gly Leu Ser Ala Gly Arg Arg Cys
 545 550 555 560

Ser Leu Thr Gly Gly Glu Glu Ser Ala Arg Ala Trp Gly Gly Ser Trp
 565 570 575

Gly Pro Gly Asn Pro Ile Phe Pro Gln Leu Thr Leu
 580 585

-173-

<210> 30
 <211> 227
 <212> PRT
 <213> Homo sapiens

<400> 30

Ile	Thr	Phe	Tyr	Gln	Thr	Leu	Trp	Ala	Arg	Pro	Arg	Arg	Ala	Arg	Gln	1	5	10	15
Ala	Arg	Arg	Val	Gly	Gly	Gly	Gly	Gly	Thr	Lys	Ala	Gly	Gly	Pro	Gly	20	25	30	
Ala	Leu	Gly	Thr	Arg	Pro	Ala	Phe	Glu	Val	Pro	Ala	Ile	Val	Val	Glu	35	40	45	
Asp	Ala	Arg	Gly	Lys	Arg	Arg	Ser	Ser	Leu	Asp	Gly	Ser	Glu	Ser	Ala	50	55	60	
Lys	Thr	Ser	Leu	Gln	Val	Thr	Asn	Leu	Val	Val	Ser	Phe	Phe	Ser	Leu	65	70	75	80
Lys	Ser	Asp	Ser	Ala	Pro	Pro	Trp	Met	Val	Leu	Ala	Val	Leu	Trp	Cys	85	90	95	
Ser	Met	Ala	Gln	Thr	Leu	Leu	Leu	Pro	Ser	Phe	Ile	Trp	Ser	Cys	Glu	100	105	110	
Arg	Tyr	Arg	Ala	Asp	Val	Arg	Thr	Val	Trp	Glu	Gln	Cys	Val	Ala	Ile	115	120	125	
Met	Ser	Glu	Glu	Asp	Gly	Asp	Asp	Asp	Gly	Gly	Cys	Asp	Asp	Tyr	Ala	130	135	140	
Glu	Gly	Arg	Val	Cys	Lys	Val	Arg	Phe	Asp	Ala	Asn	Gly	Ala	Thr	Gly	145	150	155	160
Pro	Gly	Ser	Arg	Asp	Pro	Ala	Gln	Val	Lys	Leu	Leu	Pro	Gly	Arg	His	165	170	175	
Met	Leu	Phe	Pro	Pro	Leu	Glu	Arg	Val	His	Tyr	Leu	Gln	Leu	Lys	Lys	180	185	190	
Leu	Asp	Leu	Ala	Ala	Ala	Ala	Ala	His	Thr	Phe	Phe	Val	Ala	Asn	Pro	195	200	205	
Met	His	Leu	Gln	Met	Arg	Glu	Asp	Met	Ala	Lys	Tyr	Arg	Arg	Met	Ser	210	215	220	
Gly	Val	Arg	225																

What is claimed:

1. An isolated nucleic acid molecule selected from the group consisting of:
 - a) a nucleic acid molecule comprising a nucleotide sequence which is at
5 least 60% homologous to a nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a complement thereof;
 - b) a nucleic acid molecule comprising a nucleotide sequence which is at
10 least 60% homologous to a nucleotide sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or a complement thereof;
 - c) a nucleic acid molecule comprising a fragment of at least 450 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a complement thereof;
 - 15 d) a nucleic acid molecule comprising a fragment of at least 450 nucleotides of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or a complement thereof;
 - e) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 60% homologous to the amino acid sequence of SEQ
20 ID NO:2, SEQ ID NO:7, or amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756;
 - f) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:7, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession
25 Number 98756, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, SEQ ID NO:7, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756;
 - g) a nucleic acid molecule which encodes a naturally occurring allelic
30 variant of a polypeptide comprising the amino acid sequence of SEQ ID NO: 2 or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as

- 130 -

Accession Number 98756, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1, SEQ ID NO:3 or SEQ ID NO:4 under stringent conditions;

- h) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO: 7, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:6, SEQ ID NO:8 or SEQ ID NO:9 under stringent conditions;
- i) a nucleic acid molecule comprising a nucleotide sequence which is at least 75% homologous to the nucleotide sequence of SEQ ID NO:1, the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
- j) a nucleic acid molecule comprising a nucleotide sequence which is at least 80% homologous to the nucleotide sequence of SEQ ID NO:4, the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
- of a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof;
- l) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 75% homologous to the amino acid sequence of SEQ ID NO:2 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- m) a nucleic acid molecule which encodes a polypeptide comprising an amino acid sequence at least about 65% homologous to the amino acid sequence of SEQ ID NO:5 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- n) a nucleic acid molecule which encodes a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the

- 131 -

DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acid residues of the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession
5 Number _____, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and

o) a nucleic acid molecule which encodes a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA
10 insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the nucleic acid molecule hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or SEQ ID NO:4 under stringent conditions.

15 2. The isolated nucleic acid molecule of claim 1 which is selected from the group consisting of:

- a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, or a complement thereof;
- 20 b) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:9, or a complement thereof; and
- c) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:7 or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number
25 98756.

- 132 -

3. The isolated nucleic acid molecule of claim 1 which is selected from the group consisting of:

a) a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:4, the DNA insert of the plasmid deposited with ATCC as Accession Number _____, the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or a complement thereof; and

b) a nucleic acid molecule which encodes a polypeptide comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

4. The nucleic acid molecule of claim 1 further comprising vector nucleic acid sequences.

15

5. The nucleic acid molecule of claim 1 further comprising nucleic acid sequences encoding a heterologous polypeptide.

6. A host cell which contains the nucleic acid molecule of claim 1.

20

7. The host cell of claim 5 which is a mammalian host cell.

8. A non-human mammalian host cell containing the nucleic acid molecule of claim 1.

25

- 133 -

9. An isolated polypeptide selected from the group consisting of:
- a) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, wherein the fragment
5 comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756;
 - b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:7 or SEQ ID NO:10, wherein the fragment comprises at least 15 contiguous
10 amino acids of SEQ ID NO:7 or SEQ ID NO:10;
 - c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid
15 molecule comprising SEQ ID NO:1 or SEQ ID NO:3 under stringent conditions;
 - d) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:7, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:6 or SEQ ID NO:8 under stringent conditions;
 - 20 e) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 60% homologous to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, or the DNA insert of the plasmid deposited with ATCC as Accession Number 98756;
 - f) a polypeptide which is encoded by a nucleic acid molecule comprising a
25 nucleotide sequence which is at least 60% homologous to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:6, SEQ ID NO:8, or SEQ ID NO:9;
 - g) a polypeptide comprising an amino acid sequence which is at least 60% homologous to the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:10 or the polypeptide encoded by the DNA insert of the plasmid deposited
30 with ATCC as Accession Number 98756;

- 134 -

- h) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as
- 5 Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- 10 i) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, wherein the polypeptide is
- 15 encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or SEQ ID NO:4 under stringent conditions; and
- j) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 75% homologous to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:1 or the DNA insert of the plasmid deposited with
- 20 ATCC as Accession Number _____;
- k) a polypeptide which is encoded by a nucleic acid molecule comprising a nucleotide sequence which is at least 80% homologous to a nucleic acid comprising the nucleotide sequence of SEQ ID NO:4 or the DNA insert of the plasmid deposited with ATCC as Accession Number _____;
- 25 l) a polypeptide comprising an amino acid sequence which is at least 75% homologous to the amino acid sequence of SEQ ID NO:2 or the polypeptide encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____; and
- m) a polypeptide comprising an amino acid sequence which is at least 65% homologous to the amino acid sequence of SEQ ID NO:5 or the polypeptide encoded by
- 30 the DNA insert of the plasmid deposited with ATCC as Accession Number _____.

- 135 -

10. The isolated polypeptide of claim 8 comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:7 or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756.

5

11. The isolated polypeptide of claim 8 comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with
10 ATCC as Accession Number _____.

12. The polypeptide of claim 9 further comprising heterologous amino acid sequences.

15 13. An antibody which selectively binds to a polypeptide of claim 9.

14. A method for producing a polypeptide selected from the group consisting of:
a) a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:10 or the amino acid sequence encoded by
20 the DNA insert of the plasmid deposited with ATCC as Accession Number 98756;
b) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:10, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID
25 NO:2, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:10, or the amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number 98756;
c) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2 or the amino acid sequence encoded by the DNA
30 insert of the plasmid deposited with ATCC as Accession Number 98756, wherein the

- 136 -

polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or SEQ ID NO:3 under stringent conditions;

d) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:7, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:6 or
5 SEQ ID NO:8 under stringent conditions;

e) a polypeptide comprising the amino acid sequence of SEQ ID NO: 2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino
10 acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____;

f) a fragment of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the DNA insert of the plasmid deposited with
15 ATCC as Accession Number _____, wherein the fragment comprises at least 15 contiguous amino acids of SEQ ID NO:2, SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____. or an amino acid sequence encoded by the DNA insert of the plasmid deposited
20 with ATCC as Accession Number _____; and

g) a naturally occurring allelic variant of a polypeptide comprising the amino acid sequence of SEQ ID NO:2, the amino acid sequence of SEQ ID NO:5, an amino acid sequence encoded by the DNA insert of the plasmid deposited with ATCC as Accession Number _____, or an amino acid sequence encoded by the DNA insert of the
25 plasmid deposited with ATCC as Accession Number _____, wherein the polypeptide is encoded by a nucleic acid molecule which hybridizes to a nucleic acid molecule comprising SEQ ID NO:1 or SEQ ID NO:4 under stringent conditions;

comprising culturing the host cell of claim 6 under conditions in which the nucleic acid molecule is expressed.

- 137 -

15. A method for detecting the presence of a polypeptide of claim 9 in a sample comprising:

a) contacting the sample with a compound which selectively binds to the polypeptide; and

5 b) determining whether the compound binds to the polypeptide in the sample to thereby detect the presence of a polypeptide of claim 8 in the sample.

16. The method of claim 15, wherein the compound which binds to the polypeptide is an antibody.

10

17. A kit comprising a compound which selectively binds to a polypeptide of claim 9 and instructions for use.

18. A method for detecting the presence of a nucleic acid molecule in claim 1 in a
15 sample comprising:

a) contacting the sample with a nucleic acid probe or primer which selectively hybridizes to the nucleic acid molecule; and

b) determining whether the nucleic acid probe or primer binds to a nucleic acid molecule in the sample to thereby detect the presence of a nucleic acid molecule of
20 claim 1 in the sample.

19. The method of claim 18, wherein the sample comprises mRNA molecules and is contacted with a nucleic acid probe.

25 20. A kit comprising a compound which selectively hybridizes to a nucleic acid molecule of claim 1 and instructions for use.

- 138 -

21. A method for identifying a compound which binds to a polypeptide of claim 9 comprising:
- a) contacting the polypeptide, or a cell expressing the polypeptide with a test compound; and
 - 5 b) determining whether the polypeptide binds to the test compound.
22. The method of claim 21, wherein the binding of the test compound to the polypeptide is detected by a method selected from the group consisting of:
- a) detection of binding by direct detection of test compound/polypeptide
 - 10 binding;
 - b) detection of binding using a competition binding assay; and
 - c) detection of binding using an assay for BDSF activity.
23. A method of modulating the activity of a polypeptide of claim 9 comprising
- 15 contacting the polypeptide or a cell expressing the polypeptide with a compound which binds to the polypeptide in a sufficient concentration to modulate the activity of the polypeptide.
24. A method for identifying a compound which modulates the activity of a
- 20 polypeptide of claim 9 comprising:
- a) contacting a polypeptide of claim 9 with a test compound; and
 - b) determining the effect of the test compound on the activity of the polypeptide to thereby identify a compound which modulates the activity of the polypeptide.

GTCGACCCACGCGTCCGCGCAGGATGTTTGAGTGTCGCGCCAGGGCTCTGAGACTGAGCCTGCCATCCACTCGCACGC	79
CTTTCTTTCAGGGCTTTTCGGCTGTTGGCTACACTGATGTGACCCCTCCCTTTTGGGA	151
M M G I	
4	
F L V Y V G F V F F S V L Y V Q Q G L S	24
TTT TTG GTG TAT GTT GGA TTT GTT TTC TTT TCC GTT TTA TAT GTA CAA CAA GGG CTT TCT	211
S Q A K F T E F P R N V T A T E G Q N V	44
TCT CAA GCA AAA TTT ACC GAG TTT CCG CGG AAC GTG ACG GCG ACC GAG GGG CAG AAT GTG	271
E M S C A F Q S G S A S V Y L E I Q W	64
GAG ATG TCC TGC GCC TTC CAG AGC GGC TCC GCC TCG GTG TAT CTG GAG ATC CAA TGG TGG	331
F L R G P E D L D P G A E G A Q V E	84
TTC CTG CGG GGG CCG GAG GAC CTG GAT CCC GGG GCC GAG GGG GCC GGC CAG GTG GAG	391
L L P D R D P D S D G T K I S T V K V Q	104
CTC TTG CCC GAC AGA GAC CCG GAC AGC GAC GGC ACC AAG ATC AGC ACA GTG AAA GTC CAA	451
G N D I S H K L Q I S K V R K K D E G L	124
GGC AAT GAC ATC TCC CAC AAG CTT CAG ATT TCC AAA GTG AGG AAA AAG GAT GAA GGC TTA	511
Y E C R V T D A N Y G E L Q E H K A Q A	144
TAT GAG TGC AGG GTG ACT GAT GCC AAC TAC GGG GAG CTT CAG GAA CAC AAG GCC CAG GCC	571

FIG. 1

Y	L	K	V	N	A	N	S	H	A	R	R	M	Q	A	F	E	A	S	P	164
TAT	CTG	AAA	GTC	AAT	GCC	AAC	AGC	CAT	GCC	CGC	AGA	ATG	CAG	GCC	TTC	GAA	GCC	TCG	CCC	631
M	W	L	Q	D	M	K	P	R	K	N	V	S	A	A	I	P	S	S	I	184
ATG	TGG	CTG	CAG	GAT	ATG	AAG	CCC	CGC	AAG	AAC	GTC	TCC	GCA	GCC	ATC	CCC	AGC	AGC	ATC	691
H	G	S	A	N	Q	R	T	H	S	T	S	S	P	Q	V	V	A	K	I	204
CAT	GGC	TCT	GCC	AAC	CAA	CGA	ACG	CAC	TCC	ACC	TCC	AGC	CCT	CAA	GTG	GTA	GCC	AAA	ATC	751
P	K	Q	S	P	Q	S	G	M	E	T	H	F	E	P	F	I	L	P	L	224
CCC	AAA	CAA	AGT	CCA	CAA	TCA	GGT	ATG	GAA	ACC	CAT	TTC	GAG	CCT	TTT	ATT	TTA	CCA	CTC	811
T	N	A	P	Q	K	G	Q	S	Y	R	V	D	R	F	M	N	G	D	F	244
ACA	AAC	GCT	CCA	CAG	AAA	GGT	CAG	TCG	TAT	AGA	GTA	GAC	AGA	TTT	ATG	AAT	GGT	GAT	TTT	871
*																				244
TAA																				874
AATCGGAGACCTAGTT	CAGTGCAAGTGATTATGAGAGGTGAGCACTGAGCCTGCACCAATTCACTCAGAGCTCAAAGCA																			953
TGTGGGTGCACCCCGTCAGTCCCCCTAGTGGTGCTTTCATTTCCAGGGCATCTGAGAGCTGGACTCTGGTTTTTATCCTTT																				1032
CTGTATTTACACATTATAAGAAACAATAAATCATGTAAATGTTGTTACATTACAAAAAATAAAAAAAGG																				1111
GCGGCCGC																				1119

FIG. 1
(continued)

CACGCGTCCGCAGCCAGCGCGGGGAGACACTTCACGGCGTGGCAACCCGGGTCTGTGCCCTTGAAGCCTCCGGATCG79

CAGCCAGCTCGGTCCATCCCTCACTAGTCGCAATCCCTGTGTCCAAAGCTACTCTTTTGCTATGAGCGGCAGCATGCGTG158

CAGTATCGCGCCCGAGGCTCTGAGAGCAGCCTGCGGACACGCTTGCCCTATCTGTCTTTTAGGTTTGGGGCTCTGGGC237

M

G

I

F

L

A

S

V

G

F

M

F

12

TACACGGATGTGCCCCACTCCCTTGGCATGATGGGATCTTTTGTGCGCTCTGTTGGA TTT ATG TTC303

F

S

V

L

Y

V

Q

Q

G

L

S

S

Q

A

K

F

T

E

L

P

32

TTT TCC GTG TTA TAT GTA CAA CAA GGG CTT TCT TCT CAA GCA AAA TTT ACC GAG TTG CCG363

R

N

V

T

A

T

E

G

Q

N

V

E

M

S

C

A

F

Q

S

G

52

AGA AAT GTG ACT GCT ACC GAA GGG CAA AAT GTG GAG ATG TCC TGT GCT TTC CAA AGC GGC423

S

A

S

V

Y

L

E

I

Q

W

W

F

L

R

G

P

E

D

L

E

72

TCT GCT TCA GTG TAC CTG GAG ATC CAG TGG TGG TTC CTT CGG GGG CCA GAG GAC CTG GAG483

Q

G

T

E

A

A

G

S

Q

V

E

L

L

P

D

R

D

P

D

N

92

CAA GGC ACG GAG GCT GCA GGC TCG CAG GTG GAG CTC TTA CCC GAC AGA GAC CCG GAC AAC543

D

G

T

K

I

S

T

V

K

V

Q

G

N

D

I

S

H

K

L

Q

112

GAT GGG ACC AAG ATT AGT ACA GTG AAA GTC CAA GGC AAT GAT ATC TCC CAC AAG CTT CAG603

I

S

K

V

R

K

K

D

E

G

L

Y

E

C

R

V

T

D

A

N

132

ATA TCC AAA GTG AGA AAA AAG GAT GAA GGT TTA TAC GAG TGC AGG GTG ACT GAC GCT AAC663

Fig. 2

Y	G	E	L	Q	E	H	K	A	Q	A	Y	L	K	V	N	A	N	S	H	152
TAC	GGG	GAG	CTT	CAG	GAA	CAC	AAG	GCC	CAG	GCC	TAT	CTG	AAA	GTC	AAT	GCC	AAC	AGC	CAT	723
A	R	R	M	Q	A	F	E	A	S	P	M	W	L	Q	D	T	K	P	R	172
GCT	CGG	AGG	ATG	CAG	GCC	TTT	GAA	GCC	TCA	CCT	ATG	TGG	CTG	CAA	GAC	ACG	AAG	CCT	CGA	783
K	N	A	S	S	V	V	P	S	S	V	H	N	S	A	N	Q	R	M	H	192
AAG	AAC	GCA	TCA	TCG	GTG	GTT	CCC	AGC	AGC	GTC	CAC	AAC	TCT	GCC	AAC	CAA	CGA	ATG	CAC	843
S	T	S	S	P	Q	A	V	A	K	I	P	K	Q	S	P	Q	S	A	K	212
TCC	ACC	TCC	AGC	CCT	CAA	GCG	GTA	GCC	AAA	ATC	CCC	AAG	CAA	AGT	CCA	CAA	TCA	GCA	AAG	903
S	K	S	P	V	K	S	T	E	R	T	A	K	L	T	L	Y	S	K	H	232
AGC	AAA	TCG	CCT	GTA	AAA	TCT	ACG	GAG	CGG	ACA	GCA	AAG	TTG	ACC	CTA	TAC	TCC	AAG	CAC	963
H	S	A	P	L	Y	S	S	Y	L	H	K	E	H	Q	L	P	E	A	*	251
CAT	TCT	GCA	CCC	CTG	TAC	TCT	AGT	TAT	CTA	CAC	AAG	GAG	CAT	CAG	CTT	CCG	GAA	GCA	TAA	1023
GTGAAGACACTGT	CAC	GCCTTTATTGATAAATATTTCTTTGGGAAGTTGCTGATCTTTTATTTC	CAAGAGAAATTAATGG																	1102
GAAGAGATAGGACA	TTTTCCCAATTACAGACCAATTTTTTCTTTTATTTCAACAAATAAAACCTG	CATTTCAC																		1181
TGCTCAGGAGTTGG	CCTGAATGACATCAGTATACTAAATATTTCCATGGATTCACCAATTTCTTAACGAGGACACCT																			1260
AATCTTCAAGAAGCA	AAACAAAGATGGAAACCTAAGAACCAACAACTGTCTCATACAGCACCCAGCTGAGGAACAAA																			1339

Fig. 2
(continued)

CAAATAGCTAAATGCTGACCATGGCAAATCAACATCAGACAACTTTATTTTACATATGGAATAATCAAAGAAAGTTTTT 1418

TTTTTACTTCCTTTTGTGCCCCCTGGAATTTATCTTGGAGTTTCCCTTTTTCCTTGATTGCCGTTTTTCGTTCAATGGTA 1497

GCAAGTGCCAATTATGGCCAATCCTTGTCAATCCTGGAAGTTTATATTCATATACATTGAGTGTGGTATATATCAATG 1576

TAATTTAATTCAATTTGGCAATTTCTGTATAGGCAAAACCTGGCAAATTCGTAAATTGCTTATAGTATGTGTGATAIGAC 1655

TTCAAGGTAGATAGGCTATGATGCTCATGCAAGCTGACTTCTTCAATTCATATACAAATATATTCATGAGCATATATT 1734

AGGCCACCAACTTCTTTTCCCTAAAGAAATTAATTTTCATTTGTACCTCATGTATTTTGTGAAATTTTGTAGTATATTCTC 1813

TGTTCCACTAGTTTGACCGCTACAGTTTGTCTGTCTGTTCCCTCTACTTCCCTCTGGAAAAATTTAAAAATTGTGTATGTC 1892

TCTGATAAAATGAATTAATTTTGTGTGTATGCTATGTTGGAATTTGCTGTGTTCTTTTAAACATGTATTTATTAAGG 1971

TTTGGGGATCTTGAGTTGAGTCTGAAGAAATGCACACCTGGTTTGTGACAGAGTTCCTCATGTTACCAATATTCATCTC 2050

AGAGAAAGAAAGACACCAAGTGGGAAAACTAAGAAGACATTTTGACTTCCCAAGATCCTGGAAGAGCACTTCACACTCT 2129

GACTAAATAATGTTGCTTTTGTGTTCTTCAAGACTTTTGTAGCTTTTGTCTTTCTGTAGTTGCTGCTAATTATATT 2208

TTAATGTCTACTAATTAAAAATTAATAATGTGATTGTTGGCTGAATACAATATGCAAAATGACTGCAAAAGCCCATACTGAA 2287

GAAAAATAGATGTTTAATCTTCACTCAATAATTATAATTTTAAATAGTTCATCATTTATTTTGTGACCTTATGATATTTTG 2366

Fig. 2
(continued)

TTTAGACCTGTTCTAATTACATCTTTCTCTGGCAAGAAAGATAGAACAAATCAATACATTCCCTCTTACAGTATGGAAT	2445
GGTTGTGGCTTAAGAAAGAAATGCATCCAGATGGTCTTCCAGAGAGATTATTTTATTTTCAATTATAAAACAGAAACCAT	2524
ATATGTAGGAATGGTTCATTCCCTAATGTAAGGCCATATAAATTGTAGCTTGAAGGCAAGGAATACATTTGTGTTTTTTTATGG	2603
TAAAGGACTGGCCTCTGACATGCACTTATAAGCAATGTGAATATTTTCATAAATATGCTTGACATTCCTCTTTAAACAAAT	2682
ATTGTTTTATGGTAAATCTTTCCTTGCCATTTTCTCTTTCATTGATTTCATTATTTTCATTCTAATGAAGAAAAATAAA	2761
GGTTTAATTATGATACTTTTATTAAACATACAAATGTATTTTCTTCTAAGTTAAATATCTGAAAGTTGTATAAAATGATG	2840
GTAGAGAAATATTACTCATTCGGTTTCTTTGAGCTTTAAGAAATCCCATACATTGCAGTATATATTAGAAATACTGATTTA	2919
ACATCAAACCTGGGGGGAAATCATGTATTATACTTTTACTCAATGTCTAGGTAATGGATTTCAGCTAATTTTACAGCAA	2998
GCCAAATGTGTACCCGTATCAGTAATGTTCAACCATGCTTGTATAAAAGGGCATATGCTAGTTTTTGGAAAGAAATGCTCAT	3077
TAGATTTCATTGTATCAGTGTCCAAAATAATAAGACCTGTTTATCACTGTGAAAAAATAAAAAAAAAAAAAA	3156
AAAAAAAAAAAAAAAAAAAAAAAAAAGGGCGGCCGC	3196

Fig. 2
(continued)

7/19

```

>ht122                                     244 aa vs.
>mt122                                     251 aa
scoring matrix: , gap penalties: -12/-2
77.4% identity;           Global alignment score: 1236

      10      20      30      40      *50      60
ht122  MMGIFLVYVGFVFFSVLYVQOGLSSQAKFTEFPRNVTATEGQNVEMSCAFQSGSASVYLE
      :  ::::.  ::::::::::::::::::::::::::::::::::::::::::::::::::::::
mt122  M-GIFLASVGFMFSSVLYVQOGLSSQAKFTELPRNVTATEGQNVEMSCAFQSGSASVYLE
      10      20      30      40      50

      70      80      90      100     110     120
ht122  IQWWFLRGPEDLDPGAEGAGAQVELLPDRDPDSGTKISTVKVQGNDISHKLQISKVRKK
      ::::::::::.  ::::::::::::::::::::::::::::::::::::::::::::::::::::::
mt122  IQWWFLRGPEDLEQGTEAAGSQVELLPDRDPDNDGTKISTVKVQGNDISHKLQISKVRKK
  60      70      80      90      100     110

      *130     140     150     160     170     180
ht122  DEGLYECRVTDANYGELQEHKAQAYLKVNANSHARRMQAFEASPMWLQDMKPRKNVSAAI
      ::::::::::.  ::::::::::::::::::::::::::::::::::::::::::::::::::::::
mt122  DEGLYECRVTDANYGELQEHKAQAYLKVNANSHARRMQAFEASPMWLQDTKPRKNASSV
  120     130     140     150     160     170

      190     200     210           220     230
ht122  PSSIHGSANQORTHSTSSPQVVAKIPKQSPQSG-----METHFEPFILPLTNAPQKG---Q
      ::::::::::.  ::::::::::::::::::::::::::.  ...  .  :  :  .  ...  .
mt122  PSSVHNSANQORMHSTSSPQAVAKIPKQSPQSAKSKSPVKSTERTAKLTLYSKHHSAPLYS
  180     190     200     210           220     230

      240
ht122  SYRVDRFMNGDF
      ::  .  .  .
mt122  SYLHKEHQQLPEA
  240     250

```

Fig.3

GAATTCCGGGTCGACCCACGCGTCCGCCACGCGTCCGCCACGCGTCCGGCTGAGCGAAGCGCGCGGGCGGCGG 79

CGCCTAGGGAGGAGGGCGGGCCGAGCCACCTAGCGGAGCGCGCGCGCGTGGCCCGCCAGCATGC 158

CCCGGCCCGGGCGGCTCCGCCCGCAGCCACCCCGGGCCCTCGCGGCTCGGCTCGGCGCGGGCGCGGAACC 237

GCAGCCGAGCCGAGCGGGAGCAGCGAGCCCGGGCGGCTCGAATGCAGGATGCTCGTGGTCCCCAGCATCC 316

TTGAGCCACCAGGAGTGAGGGCTGTCTCCCTGAGACCTGGCTCCAAGGAGGATGCCACAGCCGCTGCCAGCTCCGG 395

M S D E R R L P G S A V G W L V C 17

TCTGCACC ATG AGT GAT GAG CGG CTG CCT GGC AGT GCA GTG GGC TGG CTG GTA TGT 454

G G L S L L A N A W G I L S V G A K Q K 37

GGG GGC CTC TCC CTG CTG GCC AAT GCC TGG GGC ATC CTC AGC GTT GGC GCC AAG CAG AAG 514

K W K P L L E F L L C T L A A T H M L N V 57

AAG TGG AAG CCC TTG GAG TTC CTG CTG TGT ACG CTC GCG GCC ACC CAC ATG CTA AAT GTG 574

A V P I A T Y S V V Q L R R Q R P D F E 77

GCC GTG CCC ATC GCC ACC TAC TCC GTG GTG CAG CTG CGG CAG CGC CCC GAC TTC GAG 634

W N E G L C K V F V S T F Y T L T L A T 97

TGG AAT GAG GGT CTC TGC AAG GTC TTC GTG TCC ACC TTC TAC ACC CTC ACC CTG GCC ACC 694

Fig.4

C	F	S	S	V	T	S	S	L	S	Y	H	R	M	W	V	C	W	P	V	N	117	
TGT	TTC	TCT	TCT	GTC	ACC	TCC	CTC	CTC	TCC	TAC	CAC	CGC	ATG	TGG	ATG	GTC	TGC	TGG	CCT	GTC	AAC	754
Y	R	L	S	S	N	A	K	K	K	Q	A	V	H	T	V	M	G	I	W	M	V	137
TAC	CGG	CTG	AGC	AGC	AAT	GCC	AAG	AAG	CAG	CAG	GCG	GTG	CAC	ACA	GTC	ATG	GGT	ATC	TGG	ATG	GTG	814
S	F	I	L	S	S	A	L	P	A	A	V	G	W	H	D	T	S	E	R	F	Y	157
TCC	TTC	ATC	CTG	TCG	TCG	GCC	CTG	CCT	GCC	GCT	GTT	GGC	TGG	CAC	GAC	ACC	AGC	GAG	CGC	TTC	TAC	874
T	H	G	C	C	R	F	I	V	A	A	E	I	G	L	G	F	G	V	C	F	L	177
ACC	CAT	GGC	TGC	TGC	CGC	TTC	ATC	GTG	GCT	GCT	GAG	ATC	GGC	CTG	GGC	TTT	GGC	GTC	TGC	TTC	CTG	934
L	L	V	G	G	G	S	V	A	M	A	G	V	I	C	T	A	I	A	L	F	Q	197
CTG	CTG	GTG	GGC	GGC	GGC	AGC	GTG	GCC	ATG	ATG	GGC	GTG	ATC	TGC	ACA	GCC	ATC	GCC	CTC	TTC	CAG	994
T	L	A	V	V	Q	V	G	G	R	Q	A	D	H	R	A	F	T	V	P	T	I	217
ACG	CTG	GCC	GTG	CAG	GTG	GGG	CGC	CGC	CAG	CAG	GCC	GAC	CAC	CGC	GCC	TTC	ACC	GTG	CCC	ACC	ATC	1054
V	V	E	D	A	A	Q	G	G	K	R	R	S	S	I	D	G	S	E	P	A	K	237
GTG	GTG	GAG	GAC	GAC	GCG	CAG	GGC	GGC	AAG	CGG	CGC	TCC	TCC	ATC	GAT	GGC	TCG	GAG	CCC	GCC	AAA	1114
T	S	L	Q	T	T	T	G	L	V	V	T	T	I	V	F	I	Y	D	C	L	M	257
ACC	TCT	CTG	CAG	ACC	ACG	GGC	CTC	CTC	GTG	GTG	ACC	ACC	ATA	GTC	TTC	ATC	TAC	GAC	TGC	CTC	ATG	1174

Fig.4
(continued)

G F P V L D S T P I P E R S A V R Q G E 277
GGC TTC CCT GTG CTG GAC TCT ACG CCC ATC CCC GAA AGG TCT GCA GTG AGA CAG GGA GAG 1234

D W G K D Q P E G F H P S S R Q D C L P 297
GAC TGG GGC AAA GAC CAG CCT GAG GGG TTT CAT CCA AGC AGC AGG CAA GAC TGC CTT CCC 1294

*
TGA 1297

GCCATTGCAGGACATGAGGACATGAGCTCCAGAAATGGTGCCAGGCCGAGCCCTGTGCCACAGGTGGTGAGCTTCAGCA 1376

GCCTGCGGGCCGACGCCCTCAGCGCCCTGGATGGCACTCTGCGTGCTGTGTGCTCCGTGGCCAGGCCCTGCTTCCGAC 1455

CTGTCCCTGCCTCAGTTTCCCCCATCTGTGATGAGCAGGTGACCACGTTAACTCTCAGGGCTGTTTGAAGTCTCTTGGTTG 1534

TAGGCCCTCGCCACTGAGTGGCCAGGTGTGAGAGGTAGTCTAGAGCCCTCTGCGGCCCTTTGTGGAGGTCCGTTCTCAG 1613

CATGTGCCCTGGTGGCTCCCTAGGCCCTGAGGCCCTCCCACTCTCAGTGCCCCCTGTGCCCTTGGGAACCCACACTCCA 1692

CCCCAGCTAAGCACAACTGTGGACCACCACCAATGGCACCTGAGCCACCTTGACCACCATTAGTGTCCCCACCCCTCA 1771

TTACTCTGCCGTGTTGCCCTGTCCCCACCAACATCATCTCCCTCCATGGCTCCACCATGACGCTGGCAGGTGCAAGCGG 1850

AGAGCACGAGGGCAGGGCCCTGGCCCTTCCGTGTCCCTGCGTCACTCCCCAGCCCTCACTCCCCACCCGTGCAGGCCCTGG 1929

Fig.4
(continued)

CCAGGAAGGGATCCTGGCAGGGGCTTCCAGGTTCTCAGCTCAAGGCCCTGGTCCCGGCAGGCGTCCAACCCCTGGGAGC 2008
AATGTATTCTTTGCCCTTCCATCCTGGGCAGACCCCTTACAGGCCCTGGGCATTGCCATGGGCCCTGGGTCTTCCCAGGC 2087
TAAGGAGAACCAAGAACAGCTATAACCTTGAGCTAGTGAATAGGGTAGATGAGAAGGCTGTCTCCTCCAGACCCCTAC 2166
CCCTACACAGTGGCCCCACAATATGAAGACCTGGGGTAATTCCAAGGTGAGCATAGAGCCTGCCTGTGCCCAGTTCCTT 2245
CTGGCCCTCAGGTGGCCAAGCCCATCTCTTCATCCTTCAGATAGGTTCCCACTCCCAAGAAAGCTGTCTGGGTGGGG 2324
TGGAGGCTGCCCTAAGCCTGTCTGTGCTTCAGAGGCCCTCCAGTCCCTGGCTGTGGGGTAACCTGGGGGTATGAGCTGT 2403
GGCCACAGGTGAGCAAGGCAGGAACTGCAATCCAGCCCTGGCCGCGGAGGGGCCATCTCTGGCCAAATGCTGCTGTGC 2482
CTTCAAGGACTGACAAGTTACGTAGGGGCAGAGGTCGCCAGCTAGCCAGTGTCTCTCCATCTGGGGGCTGTGTCCA 2561
CTTGTCACCTTAGGTTTTTCACTCATTTGTACCTTGGGGTTTTTGCTCTGTGTGTTTCATATCCAACGGCAATACTTGCA 2640
GGGGACAGAGTCCTCTAAATACTCCAATCCTGCGGTTTTTACAAACATAAAGGGGAGACCCCAAGTGGAGGACCCTG 2719
GGCCTGGAGCTCCCTCCCAACTTTGTCCAGCATCCAGCCTGTTCCCTGGGCTCACTGGGGAGGAGTTGTCTTCATAG 2798
CACACCCAGAGCCAGGGATCCCTTTGTAGTTTTTTGACAAACGGAGCATTTCTCTGTACAGGACCCCAATAAAACTT 2877
CCTTATGATTTGCAAAAAAAGGGCGGCCGC 2915

Fig. 4
(continued)

GAATTCCCGGTGACCCACGCGTCCGGCGGGGCGAGCCACCTAGCGGAGCGCGCGCGGTGGCCGCCG 79

CCAGCATGCCCCCGGGCGGCTCCGCCCGCAGCCACCCCCCGGCCCTCGGGGCCCTGGCTCGGCCCGGGGC 158

GCGGGAACCGCAGCCGGAGCGGAGCGGAGCAGCCGGAGCCCCGGGGCTCGAATGCAGGATGCTCGTGGTCC 237

CCAGCATCCTTGAGCCACCAGAGTGAGGGCTGCTGCTCCCTGAGACCTGGCTCCAAGGAGGATGCCACAGCCGCTGC 316

M S D E R R L P G S A V G W L 15

CAGCTCCGGTCTGCACC ATG AGT GAT GAG CGG CGG CTG CCT GGC AGT GCA GTG GGC TGG CTG 378

V C G G L S L L A N A W G I L S V G A K 35

GTA TGT GGG GGC CTC TCC CTG CTG GCC AAT GCC TGG GGC ATC CTC AGC GTT GGC GCC AAG 438

Q K K W K P L L E F L L C T L A A T H M L 55

CAG AAG AAG TGG AAG CCC TTG GAG TTC CTG CTG TGT ACG CTC GCG GCC ACC CAC ATG CTA 498

N V A V P I A T Y S V V Q L R R Q R P D 75

AAT GTG GCC GTG CCC ATC GCC ACC TAC TCC GTG GTG CAG CTG CGG CAG CGC CCC GAC 558

F E W N E G L C K V F V S T F Y T L T L 95

TTC GAG TGG AAT GAG GGT CTC TGC AAG GTC TTC TCC ACC TTC TAC ACC CTC ACC CTG 618

A T C F S V T S L S Y H R M W M V C W P 115

GCC ACC TGT TTC TCT GTC ACC TCC CTC TCC TAC CAC CGC ATG TGG ATG GTC TGC TGG CCT 678

Fig. 5

V	N	Y	R	L	S	N	A	K	K	Q	A	V	H	T	V	M	G	I	W	135
GTC	AAC	TAC	CGG	CTG	AGC	AAT	GCC	AAG	AAG	CAG	GCG	GTG	CAC	ACA	GTC	ATG	GGT	ATC	TGG	738
M	V	S	F	I	L	S	A	L	P	A	V	G	W	H	D	T	S	E	R	155
ATG	GTG	TCC	TTC	ATC	CTG	TCG	GCC	CTG	CCT	GCC	GTT	GGC	TGG	CAC	GAC	ACC	AGC	GAG	CGC	798
F	Y	T	H	G	C	R	F	I	V	A	E	I	G	L	G	F	G	V	C	175
TTC	TAC	ACC	CAT	GGC	TGC	CGC	TTC	ATC	GTG	GCT	GAG	ATC	GGC	CTG	GGC	TTT	GGC	GTC	TGC	858
F	L	L	L	V	G	G	S	V	A	M	G	V	I	C	T	A	I	A	L	195
TTC	CTG	CTG	CTG	GTG	GGC	GGC	AGC	GTG	GCC	ATG	GGC	GTG	ATC	TGC	ACA	GCC	ATC	GCC	CTC	918
F	Q	T	L	A	V	Q	V	G	R	Q	A	D	H	R	A	F	T	V	P	215
TTC	CAG	ACG	CTG	GCC	GTG	CAG	GTG	GGG	CGC	CAG	GCC	GAC	CAC	CGC	GCC	TTC	ACC	GTG	CCC	978
T	I	V	V	E	D	A	Q	G	K	R	R	S	S	I	D	G	S	E	P	235
ACC	ATC	GTG	GTG	GAG	GAC	GCG	CAG	GGC	AAG	CGG	CGC	TCC	TCC	ATC	GAT	GGC	TCG	GAG	CCC	1038
A	K	T	S	L	Q	T	T	G	L	V	T	T	I	V	F	I	Y	D	C	255
GCC	AAA	ACC	TCT	CTG	CAG	ACC	ACG	GGC	CTC	GTG	ACC	ACC	ATA	GTC	TTC	ATC	TAC	GAC	TGC	1098
L	M	G	F	P	V	L	V	V	S	F	S	S	L	R	A	D	A	S	A	275
CTC	ATG	GGC	TTC	CCT	GTG	CTG	GTG	GTG	AGC	TTC	AGC	AGC	CTG	CGG	GCC	GAC	GCC	TCA	GCG	1158
P	W	M	A	L	C	V	L	W	C	S	V	A	Q	A	L	L	L	P	V	295
CCC	TGG	ATG	GCA	CTC	TGC	GTG	CTG	TGG	TGC	TCC	GTG	GCC	CAG	GCC	CTG	CTG	CCT	GTG		1218

Fig.5
(continued)

14/19

F	L	W	A	C	D	R	Y	R	A	D	L	K	A	V	R	E	K	C	M	315
TTC	CTC	TGG	GCC	TGC	GAC	CGC	TAC	CGG	GCT	GAC	CTC	AAA	GCT	GTC	CGG	GAG	AAG	TGC	ATG	1278
A	L	M	A	N	D	E	E	S	D	D	E	T	S	L	E	G	G	I	S	335
GCC	CTC	ATG	GCC	AAC	GAC	GAG	GAG	TCA	GAC	GAT	GAG	ACC	AGC	CTG	GAA	GGT	GGC	ATC	TCC	1338
P	D	L	V	L	E	R	S	L	D	Y	G	Y	G	G	D	F	V	A	L	355
CCG	GAC	CTG	GTG	TTG	GAG	CGC	TCC	CTG	GAC	TAT	GGC	TAT	GGA	GGT	GAT	TTT	GTG	GCC	CTA	1398
D	R	M	A	K	Y	E	I	S	A	L	E	G	G	L	P	Q	L	Y	P	375
GAT	AGG	ATG	GCC	AAG	TAT	GAG	ATC	TCC	GCC	CTG	GAG	GGG	GGC	CTG	CCC	CAG	CTC	TAC	CCA	1458
L	R	P	L	Q	E	D	K	M	Q	Y	L	Q	V	P	P	T	R	R	F	395
CTG	CGG	CCC	TTG	CAG	GAG	GAC	AAG	ATG	CAA	TAC	CTG	CAG	GTC	CCG	CCC	ACG	CGG	CGC	TTC	1518
S	H	D	D	A	D	V	W	A	A	V	P	L	P	A	F	L	P	R	W	415
TCC	CAC	GAC	GAT	GCG	GAC	GTG	TGG	GCC	GCC	GTC	CCG	CTG	CCC	GCC	TTC	CTG	CCG	CGC	TGG	1578
G	S	G	E	D	L	A	A	L	A	H	L	V	L	P	A	G	P	E	R	435
GGC	TCC	GGC	GAG	GAC	CTG	GCC	GCC	CTG	GCG	CAC	CTG	GTG	CTG	CCT	GCC	GGG	CCC	GAG	CGG	1638
R	R	A	S	L	L	A	F	A	E	D	A	P	P	S	R	A	R	R	R	455
CGC	CGC	GCC	AGC	CTC	CTG	GCC	TTC	GCG	GAG	GAC	GCA	CCA	CCG	TCC	CGC	GCG	CGC	CGC	CGC	1698
S	A	E	S	L	L	S	L	R	T	S	A	L	D	S	G	P	R	G	A	475
TCG	GCC	GAG	AGC	CTG	CTG	TCG	TCG	CGG	ACC	TCG	GCC	CTG	GAT	AGC	GGC	CCG	CGG	GGA	GCC	1758

Fig.5
(continued)

15/19

R	D	S	P	P	G	S	P	R	R	R	P	G	P	G	P	R	S	A	S	495
CGC	GAC	TCG	CCC	CCC	GGC	AGC	CCG	CGC	CGC	CGC	CCC	GGG	CCC	GGC	CCC	CGC	TCC	GCC	TCG	1818
A	S	L	L	P	D	A	F	A	L	T	A	F	E	C	E	P	Q	A	L	515
GCC	TCG	CTG	CTG	CCC	GAC	GCC	TTC	GCC	CTG	ACC	GCC	TTC	GAG	TGC	GAG	CCA	CAG	GCC	CTG	1878
R	R	P	P	G	P	F	P	A	A	P	A	A	P	D	G	A	D	P	G	535
CGC	CGC	CCG	CCC	GGG	CCC	TTC	CCC	GCT	GCG	CCC	GCC	GCC	GAC	GGC	GCA	GAT	CCC	GGA		1938
E	A	P	T	P	P	S	S	A	Q	R	S	P	G	P	R	P	S	A	H	555
GAG	GCC	CCG	ACG	CCC	CCA	AGC	AGC	GCC	CAG	CGG	AGC	CCA	GGG	CCA	CGC	CCC	TCT	GCG	CAC	1998
S	H	A	G	S	L	R	P	G	L	S	A	S	W	G	E	P	G	G	L	575
TCG	CAC	GCC	GGC	TCT	CTG	CGC	CCC	GGC	CTG	AGC	GCG	TCG	TGG	GGC	GAG	CCC	GGG	GGG	CTG	2058
R	A	A	G	G	G	G	S	T	S	S	F	L	S	S	P	S	E	S	S	595
CGC	GCG	GCG	GGC	GGC	GGC	AGC	AGC	ACC	AGC	AGC	TTC	CTG	AGT	TCC	CCC	TCC	GAG	TCC	TCG	2118
G	Y	A	T	L	H	S	D	S	L	G	S	A	S	*						610
GGC	TAC	GCC	ACG	CTG	CAC	TCG	GAC	TCG	CTG	GGC	TCC	GCG	TCC	TAG						2163
GACCGCGCGCCTCCCCACGGACGCCAGGCAGGCCGCTCTCCGGGGCCGACACCAAGACGCCCGCTCCC																				2242
CCCCGCGCAGACATGCGCCACCCCTCCAGGGGTGAGGGGCGTTGGCCCTCAGCGTTTGTCTTCCGGCTCCTCCCAG																				2321
CTGGCCTTGTCCAGGGCGACGGCTGCCCGGACGACTGCGCTGGGCACCGCATGTCCGGGCCGAGTGAGGTCGGGC																				2400

Fig.5
(continued)

16/19

CTGGGAGCTGAGTGACATCCCAAGCTTGGGCTTGGGTAGTGAGTGACATGTGCACACGTCCAGCTGCGCCATCACCAG	2479
CCCTGCGCAACAGGACGTCGGGAGCAGGGAACCTGAGACAGGGCCACTGCGGGATCGGACAAAGCCCCGCTTTTGGAGAG	2558
GCTGAGCTGGAGCCAATTGGCCTCCCCAGGGGCTTTCACCCACACTGCACCATAAACCGCCACACCCCTTCGGGGGGGGGG	2637
AGGGTACAGAGGGTCTCTAAGCACAGGGGTGTTCAAGAGCCCGAACAAGCTTTGATCAGGTTTCCCTGCTTCCGACCTGT	2716
CCTGCCCTCAGTTTCCCCCATCTGTGATGAGCAGGTGACCACGTTAACCTCAGGGCTGTTTGAAGTCTCTTGTTGTAGG	2795
CCCTCGCCACTGAGTGGCCCAAGGTGTGAGAGGTAGTCTAGAGCCCTCTGCGGCCCTTTGTGGAGGTCCGTTCTCAGCATG	2874
TGGCCTGGTGGCTCCCTAGGCCTGAGGCCTCCCACTCTCAGTGCCCCCTGCCCCCTTGGGAACCCACACTCCACCCC	2953
AGCTAAGCACAACTGTGGACCACCACCAATGGCACCTGAGCCACCTTGACCACCATTAGTGTCCTCCACCCCTCATTAC	3032
TCTGCCGTGCGTTGCCCTGTCCCCACCACATCATCTCCCTCCATGGCTCCACCATGACGCTGGCAGGTGCAAGCGGAGAG	3111
CACGAGGGCAGCAGGGCCTGGCCCTTCCGTGTCCTGCGTCATCCCAGCCTCACTCCCCACCCGTGCAGGCCTGGCCAG	3190
GAAGGGATCCTGGCAGGGGCTTCCAGGTTCTCAGCTCAAGGCCTGGTCCCGGGCAGGCGTCCAACCCCTGGGAGCAATG	3269
TATTTCTTTGCCCTTCCATCCTGGGCAGACCCCTTACAGGCCCTGGGCATTGCCATGGGCCCTGGGTCTTCCCAGGCTAAG	3348
GAGAACCGAACAGCTATAACCTTGAGCTAGTGAAATAGGGTAGATGAGAAGGCTGTCTCCTCCAGACCCCTACCCCT	3427

Fig.5
(continued)

ACACAGTGGCCCCACAATATGAAGACCTGGGGTAATTCCAAGTGAGCATAGAGCCCTGCCTGTGCCCAGTTCTCTGG	3506
CCCTCAGGTGGCCAAGCCCATCTCTTCATCCTTCAGATAGGTCCCACTCCCAGAAAGCTGCTGGGGTGGGGTGGG	3585
AGGCTGCCTAAGCCTGTCTGTGCTTCAGAGGCCCTCCAGTCCCTGGCTGTGGGTAAGTGGGGTATGAGCTGTGGCC	3664
ACAGGTGAGCAAGGCAGGAACTGCAATCCAGCCCTGGCCGCGGAGGGCCATCTCTGGCCAATGCTGCTGTCCTTC	3743
AAGGACTGACAAAGTTACGTAGGGGCAGAGGTCGCCAGCTAGCCAGTGTCTCCTCCATCTGGGGGCGTCTGTCCACTTG	3822
TCACCCTTAGGTTTTCACTCATTTGTCAACCTTGGGGTTTTTGCTCTGTGTGTTTCATATCCAACGGCAATACTTGCAGGGG	3901
GACAGAGTCCCTCTAAATACTCCAATCCCTGCGGTTTTTACAAACATAAAGGGGGAGACCCCAAGTGGAGGACCCCTGGGCC	3980
TGGAGCTCCCTCCCAAACCTTGTCCAGCATCCAGCCTGTTCCTGGGCTCACTGGGGAGGGAGTTGTCTTCATAGCACA	4059
CCCAGAGCCAGGATCCCTTTGTAGTTTTTTTGACAAACGGAGCATTTCTCTGTACAGGACCCCAATAAAACTTCCCTT	4138
ATGAAAAAATAAAAGGGCGGCGC	4166

Fig.5
(continued)

18/19

	1	60
SSTM-1	MS-----DERRLPGS AVGWL VC GGLS LLANAWGILSV GAKQ KKW KPLEF LLCTLAATHM	
SSTM-2	MS-----DERRLPGS AVGWL VC GGLS LLANAWGILSV GAKQ KKW KPLEF LLCTLAATHM	
proteinA-2	MARGGAGAE EASLR SNALSWLACGLLALLANAWIILSISAKQ QKH KPLELLCFLAGTHI	
proteinA-3	-----	
	61	120
SSTM-1	<u>LNVAVPIATYSVVQLRRQ-RPDFEWNEGLCKVFVSTFYTLTLATCF</u> SVTSLSYHRMWMVC	
SSTM-2	<u>LNVAVPIATYSVVQLRRQ-RPDFEWNEGLCKVFVSTFYTLTLATCF</u> SVTSLSYHRMWMVC	
proteinA-2	LMAAVPLTTFAVVQLRRQASSDYDWNESICKVFVSTYYTLALATCF TVAS LSYHRMWMVR	
proteinA-3	-----	
	121	180
SSTM-1	<u>WPVNYRLSNAKKQAVHTVMGIWMVSFILSALPAVGW</u> HDTSE RFYTHG CRFIVAEIGLGFG	
SSTM-2	<u>WPVNYRLSNAKKQAVHTVMGIWMVSFILSALPAVGW</u> HDTSE RFYTHG CRFIVAEIGLGFG	
proteinA-2	WPVNYRLSNAKKQALHAVMGIWMVSFILSTLPSIGWHNNGERY YARGCQ FIVSKIGLGFG	
proteinA-3	-----	
	181	240
SSTM-1	<u>VCFLLLVGGSVAMGVICTAIALFQTL</u> -----AVQVGRQADHRAFT	
SSTM-2	<u>VCFLLLVGGSVAMGVICTAIALFQTL</u> -----AVQVGRQADHRAFT	
proteinA-2	VCFSLLLGGIVMGLVCVAITFYQTLWARPRRARQARRVGGGGG TKAGG PGALGTRPAFE	
proteinA-3	-----ITFYQTLWARPRRARQARRVGGGGG TKAGG PGALGTRPAFE	
	241	300
SSTM-1	VPTIVVEDAQGKRRSSIDGSEPAKTS LQTTGL VTTIVFIYDCLMGFPVL-----	
SSTM-2	VPTIVVEDAQGKRRSSIDGSEPAKTS LQTTGL VTTIVFIYDCLMGFPVL VVSFSSLRADA	
proteinA-2	VPAIVVEDARGKRRSSLDGSESAKTS LQVTNLV SAIVFLYDSL TGVPIL VVSFFSLKSDS	
proteinA-3	VPAIVVEDARGKRRSSLDGSESAKTS LQVTNLV -----VSFFSLKSDS	
	301	360
SSTM-1	-----DSTPI---	
SSTM-2	SAPWMALCVLWCSVAQALLPVFLWACDRYRADLKAVREKCMALMANDEESDDETSLEGG	
proteinA-2	APPWMVLAVLWCSMAQTLLLPSFIWSCERYRADVRTVWEQCVAIMSEEDGDDD-----G	
proteinA-3	APPWMVLAVLWCSMAQTLLLPSFIWSCERYRADVRTVWEQCVAIMSEEDGDDD-----G	
	361	420
SSTM-1	-----P	
SSTM-2	ISPDLVLER--SLDYG YG GDFVALDRMAKYEISA LEG GLPQ LYPLRPLQEDKM QYLQVPP	
proteinA-2	GCDDYAEGRVCKVRFDANGATGPGSRDPA-QVKLLPGR-HMLFP--PL--ERVHYLQVPL	
proteinA-3	GCDDYAEGRVCKVRFDANGATGPGSRDPA-QVKLLPGR-HMLFP--PL--ERVHYLQ LK-	

Fig.6

19/19

	421		480
SSTM-1	ERSAVRQGED-----WGKDQ-----		
SSTM-2	TRRFSHDDADVWAAVPLPA-FLPRWGSGEDLAALAHLVLP-AGPERRRASLLAFAEDAPP		
proteinA-2	SRRLSHDETNI FSTPREPGSFLHKWSSDDIRVLPAQSRALGGPPEYLGQRHRLEDEEDE		
proteinA-3	KLDLAAAAAHTF-----FVANPMHLQ-----		
	481		540
SSTM-1	-----PEGFH-----		
SSTM-2	SRARRRSAESLLSLRTSALDSGPRGARDSPPGSPRRRPGPGPRSASASLLPDAFALTAFE		
proteinA-2	EEA---EGGGLASLRQF-LESGVLGSGGGPP-----RGPG-----FFRE--EITTF-		
proteinA-3	-----MRE--DMAKY-		
	541		600
SSTM-1	-----PSSRQ-----		
SSTM-2	CEPQALRRPPGPFPAAPAPDGADPGEAPTPPSSAQRSPGPRP--SAHSHAGSLRPGLSA		
proteinA-2	-----IDETPLPSPTASPGHSPRRPRPLGLSPRRLSLGSPESRAVGLPLGLSA		
proteinA-3	-----RRMS-----		
	601		642
SSTM-1	-----DCL---P		
SSTM-2	SWGEPGGLRAAGGGGSTSSFLSSPSESSGYATLHSDSLGSAS		
proteinA-2	-----GRRCSLTGGEESARAWGGSWGPGNPIFPQLTL----		
proteinA-3	-----GVR-----		

Fig. 6
(continued)