




(86) Date de dépôt PCT/PCT Filing Date: 2005/06/07
(87) Date publication PCT/PCT Publication Date: 2005/12/29
(85) Entrée phase nationale/National Entry: 2006/12/15
(86) N° demande PCT/PCT Application No.: EP 2005/006085
(87) N° publication PCT/PCT Publication No.: 2005/122772
(30) Priorité/Priority: 2004/06/18 (EP04014307.5)

(51) Cl.Int./Int.Cl. *A01N 43/40* (2006.01),
A01N 43/50 (2006.01), *A01N 43/54* (2006.01),
A01N 43/653 (2006.01), *A01N 47/38* (2006.01),
A01N 55/00 (2006.01)
(71) Demandeur/Applicant:
BAYER CROPS SCIENCE AG, DE
(72) Inventeurs/Inventors:
KEMPER, KONRAD, BR;
HAEUSER-HAHN, ISOLDE, DE;
REINECKE, PAUL, DE
(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : PRODUITS DE TRAITEMENT DES SEMENCES POUR LE SOJA
(54) Title: SOY SEED DRESSING

(57) **Abrégé/Abstract:**

Disclosed is the use of one or several DMI fungicides from the group comprising: a) triazoles: a.1. azaconazoles, a.2. bitertanol, a.3. bromuconazoles, a.4. cyproconazoles, a.5. difenoconazoles, a.6. diniconazoles, a.7. epoxiconazoles, a.8. fenbuconazoles, a.9. fluquinconazoles, a.10. flusilazoles, a.11. flutriafol, a.12. hexaconazoles, a.13. imibenconazoles, a.14. ipconazoles, a.15. metconazoles, a.16. myclobutanil, a.17. paclobutrazol, a.18. penconazoles, a.19. propiconazoles, a.20. prothioconazoles, a.21. simeconazoles, a.22. tebuconazoles, a.23. tetraconazoles, a.24. triadimenol, a.25. triticonazoles; b) pyrimidines: b.1. fenarimol, b.2. nuarimol; c) pyridines: c.1. pyrifenoxy; and d) imidazoles: d.1. imazalil, d.2. oxpoconazole fumarate, d.3. pefurazoate, d.4. prochloraz, d.5. triflumizoles, as a soy seed dressing against soybean rust.



Seed dressing for soya beans**A b s t r a c t**

The use of one or more DMI fungicides from the group consisting of:

a) triazoles:

a.1. azaconazole, a.2. bitertanol, a.3. bromuconazole, a.4. cyproconazole, a.5. difenoconazole, a.6. diniconazole, a.7. epoxiconazole, a.8. fenbuconazole, a.9. fluquinconazole, a.10. flusilazole, a.11. flutriafol, a.12. hexaconazole, a.13. imibenconazole, a.14. ipconazole, a.15. metconazole, a.16. myclobutanil, a.17. paclobutrazol, a.18. penconazole, a.19. propiconazole, a.20. prothioconazole, a.21. simeconazole, a.22. tebuconazole, a.23. tetraconazole, a.24. triadimenol, a.25. triticonazole;

b) pyrimidines:

b.1. fenarimol, b.2. nuarimol;

c) pyridiness:

c.1. pyrifenox;

and

d) imidazoles:

d.1. imazalil, d.2. oxpoconazole fumarate, d.3. peforazoate, d.4. prochloraz, d.5. triflumizole,

as seed dressing for soya beans against soya bean rust.

Seed dressing for soya bean

The invention relates to the use of fungicides acting as sterol biosynthesis inhibitors as seed dressings, to a method for controlling phytopathogenic fungi by treating seed with these fungicides, and also to seed treated with such fungicides.

- 5 Inhibitors of the sterol biosynthesis in fungi which block the demethylation in position 14 in the ergosteroid biosynthesis (demethylation inhibitors, DMI) have attained great practical importance as fungicides (see, for example, C.D.S. Tomlin (Publ.), The Pesticide Manual, 13th edition, The British Crop Protection Council, Farnham 2003).

Some of these fungicides are already known as seed dressing for seed of, inter alia, soya
10 beans (see, for example, WO-A 03/09 90 10, Fitopatologia Brasileira (1998) 23(2) 127-131). Also, in a few cases, the action of such a seed dressing against soya bean rust has been described (Fitopatologia Brasileira (1984) 9(1), 13 and 119).

US 2003/0060371 proposes the dressing of soya bean seed with, inter alia, DMI fungicides, to improve yield and strength of the plants. There is no indication of any action against soya
15 bean rust.

However, since the ecological and economic requirements imposed on modern-day fungicides are continually increasing, with regard, for example, to the activity spectrum, toxicity, selectivity, application rate, formation of residues, and favourable preparability, and since, furthermore, there may be problems, for example, with resistances, a constant
20 task is to develop new fungicides which in some areas at least have advantages over their known counterparts.

Surprisingly, it has now been found that fungicides from the group of the DMI, used as seed dressing for soya bean seed, are particularly effective against soya bean rust (*Phakopsora*, in particular *Phakopsora pachyrhizi* and *Phakopsora meibomia*).

- 25 Accordingly, the invention provides the use of one or more DMI fungicides from the group consisting of:

a) triazoles:

- a.1. azaconazole, a.2. bitertanol, a.3. bromuconazole, a.4. cyproconazole,
a.5. difenoconazole, a.6. diniconazole, a.7. epoxiconazole, a.8. fenbuconazole,
30 a.9. fluquinconazole, a.10. flusilazole, a.11. flutriafol, a.12. hexaconazole,

a.13. imibenconazole, a.14. ipconazole, a.15. metconazole, a.16. myclobutanil,
a.17. paclobutrazol, a.18. penconazole, a.19. propiconazole, a.20. prothioconazole,
a.21. simeconazole, a.22. tebuconazole, a.23. tetraconazole, a.24. triadimenol,
a.25. triticonazole;

5 b) pyrimidines:

 b.1. fenarimol, b.2. nuarimol;

 c) pyridines:

 c.1. pyrifenox;

and

10 d) imidazoles:

 d.1. imazalil, d.2. oxpoconazole fumarate, d.3. peforazoate, d.4. prochloraz,
 d.5. triflumizole,

as seed dressing for soya beans against soya bean rust.

15 The invention furthermore provides a method for protecting soya bean plants against soya
bean rust, which method comprises treating the seed of the plants with one or more DMI
fungicides from the group a-d.

Some of the DMI fungicides mentioned have hitherto not been used for dressing soya bean
seed.

20 Accordingly, the invention also provides the use of one or more DMI fungicides from the
groups a-d for dressing the seed of soya beans, a method for protecting soya bean plants
against phytopathogenic fungi, which method comprises treating the seeds of the plants with
one or more DMI fungicides from the groups a-d, and also soya bean seed, treated and/or
coated with one or more DMI fungicides from the groups a-d.

25 In particular, the invention also provides the use of one or more DMI fungicides from the
group consisting of:

 a) triazoles:

 a.20. prothioconazole;

 b) pyrimidines:

- 3 -

b.1. fenarimol, b.2. nuarimol;

and

c) pyridines:

c.1. pyrifenox,

5 for the dressing of soya bean seeds.

In particular, the invention furthermore provides a method for protecting soya bean plants against phytopathogenic fungi, which method comprises treating the seed of the plants with one or more DMI fungicides from the group consisting of

a) triazoles:

10 a.20. prothioconazole;

b) pyrimidines:

b.1. fenarimol, b.2. nuarimol;

and

c) pyridines:

15 c.1. pyrifenox.

The invention also provides soya bean seed, treated and/or coated with one or more DMI fungicides from the group consisting of

a) triazoles:

a.20. prothioconazole;

20 b) pyrimidines:

b.1. fenarimol, b.2. nuarimol;

and

c) pyridines:

c.1. pyrifenox.

Surprisingly, the dressing according to the invention is particularly effective against soya bean rust, in spite of the fact that this particularly aggressive plant disease is not soil-borne but wind-borne.

- 5 Moreover, using the dressing according to the invention it is possible to control fungal diseases in soya beans in an economically and ecologically advantageous manner, and also to improve plant health and vitality.

The DMI fungicides of the group a-e are known and commercially available. Details concerning the individual substances and their purchase can be found in "The Pesticide
10 Manual", 13th edition.

Preference is given to the DMI fungicides of group a and to prochloraz of group e.

Particular preference is given to fluquinconazole, flutriafol, ipconazole, prothioconazole, tebuconazole, triticonazole, metconazole, cyproconazole, bitertanol, triadimefon and triadimenol.

- 15 Especially preferred are fluquinconazole, flutriafol, ipconazole, prothioconazole and triticonazole.

Furthermore, it is preferred to use mixtures of two or more, in particular one, of the DMI fungicides mentioned.

Particular preference is given to the mixtures of the compounds fluquinconazole and
20 prothioconazole.

In the above-mentioned mixtures of two active compounds, the weight ratio of the DMI fungicides in question is generally 1:0.01-100, preferably 1:0.2-20, particularly preferably 1:0.1-10.

The DMI fungicides used according to the invention, as such or in their formulations,
25 can also be used as a mixture with other known fungicides, bactericides, acaricides, nematocides or insecticides, for example in order to broaden the activity spectrum or to prevent the development of resistances in this way. In many cases, synergistic effects result, i.e. the effectiveness of the mixture exceeds the effectiveness of the

individual components.

Compounds which are suitable as mixing partners are, for example, the following:

Fungicides:

- 2-Phenylphenol; 8-hydroxyquinoline sulphate; acibenzolar-S-methyl; actinovate; aldimorph;
5 amidoflumet; ampropylfos; ampropylfos-potassium; andoprim; anilazine; azoxystrobin;
benalaxyl; benodanil; benomyl; benthiavalicarb-isopropyl; benzamacril; benzamacril-iso-
butyl; bilanafos; binapacryl; biphenyl; blasticidin-S; boscalid; bupirimate; buthiobate;
butylamine; calcium polysulphide; capsimycin; captafol; captan; carbendazim; carboxin;
carpropamid; carvone; chinomethionat; chlobenthiazole; chlorfenazole; chloroneb;
10 chlorothalonil; chlōzolate; cis-1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)-cycloheptanol;
clozylacon; cyazofamid; cyflufenamid; cymoxanil; cyprodinil; cyprofuram; Dagger G;
debacarb; dichlofluanid; dichlone; dichlorophen; diclocymet; diclomezine; dicloran;
diethofencarb; diflumentorim; dimethirimol; dimethomorph; dimoxystrobin; dinocap;
diphenylamine; dipyrithione; ditalimfos; dithianon; dodine; drazoxolon; edifenphos;
15 ethaboxam; ethirimol; etridiazole; famoxadone; fenamidone; fenapanil; fenfuram;
fenhexamid; fenitropan; fenoxanil; fencpiclonil; fenpropidin; fenpropimorph; ferbam;
fluazinam; flubenzimine; fludioxonil; flumetover; flumorph; fluoromide; fluoxastrobin;
flurprimidol; flusulfamide; flutolanil; folpet; fosetyl-Al; fosetyl-sodium; fuberidazole;
furalaxyl; furametpyr; furcarbanil; furmecyclox; guazatine; hexachlorobenzene; hymexazol;
20 iminoctadine triacetate; iminoctadine tris(albesilate); iodocarb; iprobenfos; iprodione;
iprovalicarb; irumamycin; isoprothiolane; isovaledione; kasugamycin; kresoxim-methyl;
mancozeb; maneb; meferimzone; mepanipyrim; mepronil; metalaxyl; metalaxyl-M;
methasulfocarb; methfuroxam; methyl 1-(2,3-dihydro-2,2-dimethyl-1H-inden-1-yl)-1H-
imidazole-5-carboxylate; methyl 2-[[[cyclopropyl[(4-methoxyphenyl)imino]methyl]thio]-
25 methyl]-α-(methoxymethylene)benzeneacetate; methyl 2-[2-[3-(4-chlorophenyl)-1-methyl-
allylideneaminooxymethyl]phenyl]-3-methoxyacrylate; metiram; metominostrobin;
metrafenone; metsulfovax; mildiomicin; monopotassium carbonate; myclozolin; N-(3-ethyl-
3,5,5-trimethylcyclohexyl)-3-formylamino-2-hydroxybenzamide; N-(6-methoxy-3-pyridin-
yl)cyclopropanecarboxamide; N-butyl-8-(1,1-dimethylethyl)-1-oxaspiro[4.5]decan-3-amine;
30 natamycin; nitrothal-isopropyl; noviflumuron; ofurace; orysastrobin; oxadixyl; oxolinic
acid; oxycarboxin; oxyfenthiin; pencycuron; penthiopyrad; phosdiphen; phthalide;
picobenzamid; picoxystrobin; piperalin; polyoxins; polyoxorim; procymidone;
propamocarb; propanosine-sodium; propineb; proquinazid; pyraclostrobin; pyrazophos;

pyrimethanil; pyroquilon; pyroxyfur; pyrrolnitrine; quinconazole; quinoxifen; quintozone;
 silthiofam; sodium tetrathiocarbonate; spiroxamine; sulphur; tecloftalam; tecnazene;
 tetcyclacis; thicyofen; thifluzamide; thiophanate-methyl; thiram; tiadinil; tioxyamid;
 tolclofos-methyl; tolylfluanid; triazbutil; triazoxide; tricyclamide; tricyclazole; tridemorph;
 5 trifloxystrobin; validamycin A; vinclozolin; zineb; ziram; zoxamide; (2S)-N-[2-[4-[[3-(4-
 chlorophenyl)-2-propynyl]oxy]-3-methoxyphenyl]ethyl]-3-methyl-2-[(methylsulphonyl)-
 amino]butanamide; 1-(1-naphthalenyl)-1H-pyrrole-2,5-dione; 2,3,5,6-tetrachloro-4-(methyl-
 sulphonyl)pyridine; 2,4-dihydro-5-methoxy-2-methyl-4-[[[1-[3-(trifluoromethyl)phenyl]-
 ethylidene]amino]oxy]methyl]phenyl]-3H-1,2,3-triazol-3-one; 2-amino-4-methyl-N-phenyl-
 10 5-thiazolecarboxamide; 2-chloro-N-(2,3-dihydro-1,1,3-trimethyl-1H-inden-4-yl)-3-pyridine-
 carboxamide; 3,4,5-trichloro-2,6-pyridinedicarbonitrile; 3-[(3-bromo-6-fluoro-2-methyl-1H-
 indol-1-yl)sulphonyl]-N,N-dimethyl-1H-1,2,4-triazole-1-sulphonamide;

and copper salts and preparations, such as Bordeaux mixture; copper hydroxide;
 copper naphthenate; copper oxychloride; copper sulphate; cufraneb; cuprous oxide;
 15 mancopper; oxine-copper:

Bactericides:

Bronopol, dichlorophen, nitrapyrin, nickel dimethyldithiocarbamate, kasugamycin,
 oethilidon, furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam,
 copper sulphate and other copper preparations.

20 Insecticides/acaricides/nematicides:

Acetylcholine esterase (AChE) inhibitors

1.1 Carbamates,

~~for example alanycarb, aldicarb, aldoxycarb, allyxycarb, aminocarb,~~
 bendiocarb, benfuracarb, bufencarb, butacarb, butocarboxim,
 25 butoxycarboxim, carbaryl, carbofuran, carbosulfan, cloethocarb, dimetilan,
 ethiofencarb, fenobucarb, fenothiocarb, formetanate, furathiocarb, isoprocarb,
 metam-sodium, methiocarb, methomyl, metolcarb, oxamyl, pirimicarb,
 promecarb, propoxur, thiodicarb, thiofanox, trimethacarb, XMC, xylylcarb,
 triazamate

30 1.2 Organophosphates,

for example acephate, azamethiphos, azinphos (-methyl, -ethyl), bromophos-ethyl, bromfeninfos (-methyl), butathiofos, cadusafos, carbophenothion, chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos (-methyl/-ethyl), coumaphos, cyanofenphos, cyanophos, chlorfenvinphos, demeton-S-methyl, demeton-S-methylsulphon, dialifos, diazinon, dichlofenthion, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos, dioxabenzofos, disulfoton, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion, fensulfothion, fenthion, flupyrazofos, fonofos, formothion, fosmethilan, fosthiazate, heptenophos, iodofenphos, iprobenfos, isazofos, isofenphos, isopropyl O-salicylate, isoxathion, malathion, mecarbam, methacrifos, methamidophos, methidathion, mevinphos, monocrotophos, naled, omethoate, oxydemeton-methyl, parathion (-methyl/-ethyl), phenthoate, phorate, phosalone, phosmet, phosphamidon, phosphocarb, phoxim, pirimiphos (-methyl/-ethyl), profenofos, propaphos, propetamphos, prothiofos, prothoate, pyraclofos, pyridaphenthion, pyridathion, quinalphos, sebufos, sulfotep, sulprofos, tebupirimfos, temephos, terbufos, tetrachlorvinphos, thiometon, triazophos, triclofon, vamidothion

Sodium channel modulators / voltage-dependent sodium channel blockers

2.1 Pyrethroids,

for example acrinathrin, allethrin (d-cis-trans, d-trans), beta-cyfluthrin, bifenthrin, bioallethrin, bioallethrin-S cyclopentyl isomer, bioethanomethrin, biopermethrin, bioresmethrin, chlovaporthrin, cis-cypermethrin, cis-resmethrin, cis-permethrin, clocythrins, cycloprothrin, cyfluthrin, cyhalothrin, cypermethrin (alpha-, beta-, theta-, zeta-), cyphenothrin, deltamethrin, empenthrin (1R isomer), esfenvalerate, etofenprox, fenfluthrin, fenpropathrin, fenpyrithrin, fenvalerate, flubrocycrin, flucythrinate, flufenprox, flumethrin, fluvalinate, fubfenprox, gamma-cyhalothrin, imiprothrin, kadethrin, lambda-cyhalothrin, metofluthrin, permethrin (cis-, trans-), phenothrin (1R trans-isomer), prallethrin, profluthrin, protrifenbute, pyresmethrin, resmethrin, RU 15525, silafluofen, tau-fluvalinate, tefluthrin, terallethrin, tetramethrin (1R isomer), tralomethrin, transfluthrin, ZXI 8901,

pyrethrins (pyrethrum)

DDT

2.2 Oxadiazines,
for example indoxacarb

5 Acetylcholine receptor agonists/antagonists

3.1 Chloronicotinyis,
for example acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram,
nithiazine, thiacloprid, thiamethoxam

3.2 Nicotines, bensultap, cartap

10 Acetylcholine receptor modulators

4.1 Spinosyns,
for example spinosad

GABA-controlled chloride channel antagonists

15 5.1 Organochlorines,
for example camphechlor, chlordane, endosulfan, gamma-HCH, HCH,
heptachlor, lindane, methoxychlor

5.2 Fiproles,
for example acetoprole, ethiprole, fipronil, pyrafluprole, pyriprole, vaniliprole

Chloride channel activators

20 6.1 Mectins,
for example avermectin, emamectin, emamectin-benzoate, ivermectin, milbemycin

Juvenile hormone mimetics,

for example diofenolan, epofenonane, fenoxycarb, hydroprone, kinoprene,
methoprene, pyriproxifen, triprene

25 Ecdyson agonists/disruptors

- 8.1 Diacylhydrazines,
for example chromafenozide, halofenozide, methoxyfenozide, tebufenozide

Chitin biosynthesis inhibitors

- 9.1 Benzoylureas,
5 for example bistrifluron, chlofluazuron, diflubenzuron, fluazuron,
flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron,
noviflumuron, penfluron, teflubenzuron, triflumuron

- 9.2 Buprofezin

- 9.3 Cyromazine

10 Oxidative phosphorylation inhibitors, ATP disruptors

- 10.1 Diafenthiuron

- 10.2 Organotin,
for example azocyclotin, cyhexatin, fenbutatin-oxide

Oxidative phosphorylation decouplers acting by interrupting the H-proton gradient

- 15 11.1 Pyrroles,
for example chlorfenapyr

- 11.2 Dinitrophenols,
for example binapacyrl, dinobuton, dinocap, DNOC

Site-I electron transport inhibitors

- 20 12.1 METI's,
for example fenazaquin, fenpyroximate, pyrimidifen, pyridaben, tebufenpyrad,
tolfenpyrad

- 12.2 Hydramethylnon

- 12.3 Dicofol

25 Site-II electron transport inhibitors

Rotenone

Site-III electron transport inhibitors

Acequinocyl, fluacrypyrim

Microbial disruptors of the insect gut membrane

5 Bacillus thuringiensis strains

Fat synthesis inhibitors

tetronic acids,

for example spirodiclofen, spiromesifen

tetramic acids,

10 for example spirotetramat (CAS Reg. No.: 203313-25-1) and 3-(2,5-dimethyl-phenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl carbonate (aka: carbonic acid, 3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl-ethyl ester, CAS Reg. No.: 382608-10-8)

carboxamides,

15 for example flonicamid

octopaminergic agonists,

for example amitraz

Inhibitors of magnesium-stimulated ATPase,

propargite

20 benzoic acid dicarboxamides,
for example flubendiamide

nereistoxin analogues

for example thiocyclam hydrogen oxalate, thiosultap-sodium

Biologicals, hormones or pheromones

25 azadirachtin, Bacillus spec., Beauveria spec., codlemone, Metarrhizium spec.,

Paecilomyces spec., *thuringiensin*, *Verticillium spec.*

Active compounds with unknown or unspecific mechanisms of action

- 23.1 Fumigants,
for example aluminium phosphide, methyl bromide, sulphuryl fluoride
 - 5 23.2 Antifeedants,
for example cryolite, flonicamid, pymetrozine
 - 23.3 Mite growth inhibitors,
for example clofentezine, etoxazole, hexythiazox
 - 10 23.4 Amidoflumet, benclothiaz, benzoximate, bifenazate, bromopropylate, buprofezin, quinomethionate, chlordimeform, chlorobenzilate, chloropicrin, clothiazoben, cycloprene, cyflumetofen, dicyclanil, fenoxacrim, fentrifanil, flubenzimine, flufenimer, flutenzin, gossyplure, hydramethylnone, japonilure, metoxadiazon, petroleum, piperonyl butoxide, potassium oleate, pyridalyl, sulfluramid, tetradifon, tetrasul, triarathene, verbutin
 - 15 A mixture with other known active compounds, such as herbicides, or with fertilizers and growth regulators, safeners and/or semiochemicals is also possible.
- Preference is given to mixtures of the above-mentioned triazoles, in particular fluquinconazole, flutriafol, ipconazole, triticonazole and prothioconazole, with tolylfluanid or strobilurins, in particular fluoxastrobin, trifloxystrobin, azoxystrobin and pyraclostrobin.
- 20 Particular preference is given to the mixtures fluquinconazole/tolylfluanid, fluquinconazole/fluoxastrobin, fluquinconazole/trifloxystrobin, fluquinconazole/azoxystrobin, fluquinconazole/pyraclostrobin, prothioconazole/tolylfluanid, prothioconazole/fluoxastrobin, prothioconazole/trifloxystrobin, prothioconazole/azoxystrobin and prothioconazole/pyraclostrobin.
 - 25 Preference is given to the weight ratios mentioned above.

According to the invention, the DMI fungicides are converted into the customary seed dressing formulations, such as solutions, emulsions, suspensions, powders, foams, slurries or other coating materials for seed, and also ULV formulations.

Dry treatments (preferably with addition of tackifiers, such as, for example, liquid paraffin or talc, and, if appropriate, colorants),

slurry treatments (preferably with addition of wetting agents, dispersants, emulsifiers, adhesives, inert fillers and colorants),

5 aqueous liquid treatments (preferably with addition of emulsifiers, dispersants, thickeners, antifreeze agents, polymers, adhesives and colorants),

solvent-based liquid treatments (with addition of solvents and colorants),

emulsion treatments (with addition of emulsifiers, solvents and colorants).

10 These formulations are prepared in a known manner by mixing the active compounds or active compound combinations with customary additives, such as, for example, customary extenders and also solvents or diluents, colorants, wetting agents, dispersants, emulsifiers, defoamers, preservatives, secondary thickeners, adhesives, gibberellins and water as well.

15 Suitable colorants that may be present in the seed dressing formulations to be used according to the invention include all colorants customary for such purposes. Use may be made both of pigments, of sparing solubility in water, and of dyes, which are soluble in water. Examples that may be mentioned include the colorants known under the designations Rhodamin B, C.I. Pigment Red 112 and C.I. Solvent Red 1.

20 Suitable wetting agents that may be present in the seed dressing formulations to be used according to the invention include all substances which promote wetting and are customary in the formulation of agrochemically active compounds. Preference is given to using alkyl-naphthalenesulphonates, such as diisopropyl- or diisobutyl-naphthalenesulphonates.

25 Suitable dispersants and/or emulsifiers that may be present in the seed dressing formulations to be used according to the invention include all nonionic, anionic and cationic dispersants which are customary in the formulation of agrochemically active compounds. Preference is given to using nonionic or anionic dispersants or mixtures of nonionic and anionic dispersants. Particularly suitable nonionic dispersants are ethylene oxide/propylene oxide block polymers, alkylphenol polyglycol ethers, and also tristyrylphenol polyglycol ethers and their phosphated or sulphated derivatives. Particularly suitable anionic dispersants are lignosulphonates, polyacrylic acid salts and arylsulphonate/formaldehyde condensates.

30 Defoamers that may be present in the seed dressing formulations to be used according to the invention include all foam-inhibiting compounds which are customary in the formulation of agrochemically active compounds. Preference is given to using silicone defoamers and

magnesium stearate.

Preservatives that may be present in the seed dressing formulations to be used according to the invention include all compounds which can be used for such purposes in agrochemical compositions. By way of example, mention may be made of dichlorophen and benzyl alcohol
5 hemiformal.

Suitable secondary thickeners that may be present in the seed dressing formulations to be used according to the invention include all compounds which can be used for such purposes in agrochemical compositions. Preference is given to cellulose derivatives, acrylic acid derivatives, xanthan, modified clays and finely divided silica.


10 Solvents suitable for combination with the DMI fungicides are all organic solvents which can be used in agrochemical compositions. Preference is given to ketones, such as methyl isobutyl ketone and cyclohexanone, furthermore amides, such as dimethylformamide, furthermore cyclic compounds, such as N-methylpyrrolidone, N-octylpyrrolidone, N-dodecylpyrrolidone, N-octylcaprolactam, N-dodecylcaprolactam and γ -butyrolactone, in
15 addition strongly polar solvents, such as dimethyl sulphoxide, furthermore aromatic hydrocarbons, such as xylene, moreover esters, such as propylene glycol monomethyl ether acetate, dibutyl adipate, hexyl acetate, heptyl acetate, tri-n-butyl citrate, diethyl phthalate and di-n-butyl phthalate, and furthermore alcohols, such as ethanol, n- and i-propanol, n- and i-butanol, n- and i-amyl alcohol, benzyl alcohol and 1-methoxy-2-propanol.

20 Suitable polymeric additive components are customary biodegradable natural and synthetic compounds. Preference is given to polyesters, polyether esters, copolyesters, polyanhydrides, polyesterurethanes, thermoplastic polysaccharides or polysaccharide derivatives, and also polyesters, polyether esters and polyesteramides containing aliphatic and aromatic ester groupings.

25 Suitable adhesives that may be present in the seed dressing formulations to be used according to the invention include all customary binders which can be used in seed dressings. Polyvinylpyrrolidone, polyvinyl acetate, polyvinyl alcohol and tylose may be mentioned as being preferred.

Suitable gibberellins that may be present in the seed dressing formulations to be used
30 according to the invention preferably include compounds of the formula

- 14 -






in which


R represents a hydrogen atom or a hydroxyl group and

the dashed line indicates that at the position of the ring either a C-C single bond or a C=C double bond is present.

Examples that may be mentioned of gibberellins of the formula (I) include the following:

gibberellin A₁gibberellin A₃
(= gibberellic acid)gibberellin A₄

and



gibberellin A₇

(I-4)

Particular preference is given to gibberellic acid of formula (I-2).

The gibberellins of the formula (I) are known (cf. R. Wegler "Chemie der Pflanzenschutz-
5 und Schädlingsbekämpfungsmittel", volume 2, Springer Verlag, Berlin-Heidelberg-New
York, 1970, pages 401 - 412).

The seed dressing formulations to be used according to the invention are used either directly
or after prior dilution with water for the treatment of soya bean seed. The seed dressing
formulations to be used according to the invention or dilute preparations thereof can also be
10 used for dressing seed of transgenic soya bean plants. In this context, additional synergistic
effects may also arise in interaction with the substances formed by expression.

Suitable mixing equipment for treating soya bean seed with the seed dressing formulations
to be used according to the invention or the preparations prepared therefrom by addition of
water includes all mixing equipment which can commonly be used for dressing. The specific
15 procedure adopted when dressing comprises introducing the seed into a mixer, adding the
particular desired amount of seed dressing formulations, either as such or after prior dilution
with water, and carrying out mixing until the formulation is uniformly distributed on the seed.
If appropriate, this is followed by a drying operation.

The application rate of the seed dressing formulations to be used according to the invention
20 may be varied within a relatively wide range. It depends on the respective content of the active
compounds in the formulation and on the seed. In general, the application rates of DMI
fungicides or combination thereof are between 0.001 and 50 g per kilogram of seed, preferably
between 0.01 and 15 g per kilogram of seed.

In addition to the control of *Phakopsora*, in particular *Phakopsora pachyrhizi* and
25 *Phakopsora meibomia*, the seed dressings used according to the invention can be employed,
for example, for controlling Plasmodiophoromycetes, Oomycetes, Chytridiomycetes,
Zygomycetes, Ascomycetes, Basidiomycetes, Deuteromycetes, Pseudomonadaceae,
Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomycetaceae.

Some pathogens causing fungal and bacterial diseases which come under the generic names listed above may be mentioned by way of example, but not by way of limitation:

- Xanthomonas species, such as, for example, Xanthomonas campestris pv. oryzae;
Pseudomonas species, such as, for example, Pseudomonas syringae pv. lachrymans;
5 Erwinia species, such as, for example, Erwinia amylovora;
Phytophthora species, such as, for example, Phytophthora sojae;
Pseudoperonospora species, such as, for example, Pseudoperonospora humuli or
Pseudoperonospora cubensis;
Peronospora species, such as, for example, Peronospora brassicae or Peronospora
10 manshurica;
Erysiphe species, such as, for example, Erysiphe polygoni;
Podosphaera species, such as, for example, Podosphaera leucotricha;
Puccinia species, such as, for example, Puccinia recondita;
Sclerotinia species, such as, for example, Sclerotinia sclerotiorum;
15 Ustilago species, such as, for example, Ustilago nuda or Ustilago avenae;
Pellicularia species, such as, for example, Pellicularia sasakii;
Fusarium species, such as, for example, Fusarium culmorum;
Septoria species, such as, for example, Septoria nodorum or Septoria glycines;
Leptosphaeria species, such as, for example, Leptosphaeria nodorum;
20 Cercospora species, such as, for example, Cercospora kikuchii;
Alternaria species, such as, for example, Alternaria brassicae.

Phomopsis species, such as, for example, Phomopsis longicola; Colletotrichum species, such as, for example, Colletotrichum truncatum;

Rhizoctonia species, such as, for example, Rhizoctonia solani;

- 25 and in particular

Microsphaera species, such as Microsphaera manshurica.

The seed dressings used according to the invention are particularly suitable for controlling soya bean rust.

- 30 The dressings according to the invention also exhibit a very strong invigorating action in plants. Accordingly, they are suitable for mobilizing the internal defences of the plant against attack by unwanted microorganisms.

In the present context, plant-invigorating (resistance-inducing) compounds are to be understood as meaning substances which are capable of stimulating the defence system of plants such that, when the treated plants are subsequently inoculated with unwanted microorganisms, they display substantial resistance to these microorganisms.

- 5 In the present case, unwanted microorganisms are to be understood as meaning phyto-pathogenic fungi and bacteria. The compounds according to the invention can thus be used to protect plants within a certain period of time after treatment against attack by the pathogens mentioned. The period of time for which this protection is achieved generally extends for 1 to 70 days, preferably 1 to 35 days, from sowing.
- 10 The invention is illustrated in more detail by the examples, without being limited to these.

Example 1

Action of fluquinconazole against *Phakopsora pachyrhizi*

Seed dressing: Fluquinconazole 25 WPC (wetable powder)

For dressing, the seed was shaken for 1 minute with the seed dressing in a closed plastic
5 bag.

The seed was sown in a field (4 plots, plot size in each case 10 m², 40 seeds per m²) and, one
day after sowing, artificially irrigated.

The natural degree of infection in the control was, on average, 75%.

The efficacy is expressed in percent. 0% means an efficacy which corresponds with that of
10 the control, whereas an efficacy of 100% means that no infection is observed.

Application rate [g of fluquinconazole / 100 kg of seed]	Duration [days]	Efficacy [%]
50	22	100
50	33	100
50	37	100
5	28	100

Example 2

Action of prothioconazole against *Phakopsora pachyrhizi*

Seed dressing: Prothioconazole

For dressing, the seed was shaken for 1 minute with the seed dressing in a closed plastic
5 bag.

The seed was sown in a field (4 plots, plot size in each case 10 m², 40 seeds per m²) and, one day after sowing, artificially irrigated.

The natural degree of infection in the control was, on average, 75%.

The efficacy is expressed in percent. 0% means an efficacy which corresponds with that of
10 the control, whereas an efficacy of 100% means that no infection is observed.

Application rate [g of prothioconazole / 100 kg of seed]	Duration [days]	Efficacy [%]
5	28	100
10	28	100

Example 3

Action of flutriafol against *Phakopsora pachyrhizi*

Seed dressing: Flutriafol / carbendazim / imidacloprid

For dressing, the seed was shaken for 1 minute with the seed dressing in a closed plastic
5 bag.

The seed was sown in a field (4 plots, plot size in each case 10 m², 40 seeds per m²) and, one day after sowing, artificially irrigated.

Soy bean plants the seeds of which had been treated with a mixture of the fungicide carbendazim (100 g / 100 kg of seed) and the insecticide imidacloprid (120 g / 100 kg of
10 seed) were used as control.

The natural degree of infection in the control was, on average, 75%.

The efficacy is expressed in percent according to Abbot's formula ($[(\text{efficacy control} - \text{efficacy inventive example}) / \text{efficacy control}] * 100$). 0% means an efficacy which
15 corresponds with that of the control, whereas an efficacy of 100% means that no infection is observed.

Application rate [g flutriafol / g carbendazim / g Imidacloprid / 100 kg of seed]	Duration [days]	Efficacy [%]
7.5 / 100 / 120	19	99.8
7.5 / 100 / 120	25	100

Patent claims

1. Use of one or more demethylation inhibitor (DMI) fungicides from the group consisting of:
 - a) triazoles:
 - a.1. azaconazole, a.2. bitertanol, a.3. bromuconazole, a.4. cyproconazole,
 - 5 a.5. difenoconazole, a.6. diniconazole, a.7. epoxiconazole, a.8. fen-
 - buconazole, a.9. fluquinconazole, a.10. flusilazole, a.11. flutriafol,
 - a.12. hexaconazole, a.13. imibenconazole, a.14. ipconazole, a.15. met-
 - conazole, a.16. myclobutanil, a.17. paclobutrazol, a.18. penconazole,
 - a.19. propiconazole, a.20. prothioconazole, a.21. simeconazole,
 - 10 a.22. tebuconazole, a.23. tetraconazole, a.24. triadimenol, a.25. tri-
 - triconazole;
 - b) pyrimidines:
 - b.1. fenarimol, b.2. nuarimol;
 - c) pyridines:
 - 15 c.1. pyrifenox;

and

 - d) imidazoles:
 - d.1. imazalil, d.2. oxpoconazole fumarate, d.3. peforazoate, d.4. prochloraz,
 - d.5. triflumizole,

20 as seed dressing for soya beans against soya bean rust.
2. Use according to Claim 1 using a DMI fungicide from the group consisting of fluquinconazole, flutriafol, ipconazole, prothioconazole and triticonazole.
3. Method for protecting soya bean plants against soya bean rust, which method comprises treating seed of the plants with one or more DMI fungicides from the

25 group a – e according to Claim 1.
4. Method according to Claim 3 where one or more DMI fungicides from the group consisting of fluquinconazole, flutriafol, ipconazole, prothioconazole and triticonazole is used.

5. Use of one or more DMI fungicides from the group consisting of:
- a) triazoles:
 - a.20. prothioconazole;
 - b) pyrimidines:
 - b.1. fenarimol, b.2. nuarimol;
- 5 and
- c) pyridines:
 - c.1. pyrifenox,
- as seed dressing for soya beans.
- 10 6. Use according to Claim 5 using the DMI fungicide prothioconazole.
7. Method for protecting soya bean plants against phytopathogenic fungi, which method comprises treating seed of the plants with one or more DMI fungicides from the group a – c according to Claim 5.
8. Method according to Claim 7 where the DMI fungicide used is prothioconazole.
- 15 9. Soya bean seed, treated and/or coated with one or more DMI fungicides from the group a – c according to Claim 5.
10. Soya bean seed according to Claim 9, treated and/or coated with prothioconazole.