

AUSTRALIA

PATENTS ACT 1990

NOTICE OF ENTITLEMENT

We, Indiana University Foundation, Showalter House, PO Box 500, Bloomington, Indiana, 47402, United States of America, being the applicant and nominated person in respect of Application No. 77294/94 state the following:-

The person nominated for the grant of the patent has entitlement from the actual inventor by assignment.

The person nominated for the grant of the patent has entitlement from the applicant listed in the declaration under Article 8 of the PCT by assignment. The basic application listed in the declaration made under Article 8 of the PCT is the first application made in a Convention country in respect of the invention.

The person nominated for the grant of the patent has the consent of Byoung S Kwon, the depositor with Agricultural Research Service Culture Collection of the deposit listed hereafter, to rely on that deposit:

Deposited material	-	<i>Escherichia coli</i> Human 4-1BBcDNA(pH4-1BB)
Deposit number	-	NRRL B-21131
Deposit date	-	25 August 1993

INDIANA UNIVERSITY FOUNDATION
By their Patent Attorneys
CULLEN & CO.


Registered Patent Attorney

Date: 1 April 1996



AU9477294

(12) PATENT ABRIDGMENT (11) Document No. AU-B-77294/94
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 697482

(54) Title
HUMAN RECEPTOR H4-1BB

(51)⁶ International Patent Classification(s)
C12N 015/12 C07K 014/71 C07K 016/28 C12N 015/62
G01N 033/68 C07K 014/715

(21) Application No. : 77294/94 (22) Application Date : 15.09.94

(87) PCT Publication Number : WO95/07984

(30) Priority Data

(31) Number (32) Date (33) Country
08/122796 16.09.93 US UNITED STATES OF AMERICA

(43) Publication Date : 03.04.95

(44) Publication Date of Accepted Application : 08.10.98

(71) Applicant(s)
INDIANA UNIVERSITY FOUNDATION

(72) Inventor(s)
BYOUNG S. KWON

(74) Attorney or Agent
CULLEN & CO , GPO Box 1074, BRISBANE QLD 4001

(57) Claim

1. A cDNA encoding for human receptor protein H4-1BB having a nucleotide sequence as shown in SEQ ID NO: 1.
3. An isolated DNA, wherein said DNA comprises a nucleotide sequence selected from:
 - a) nucleotides of H4-1BBFI, herein defined;
 - b) nucleotides of H4-1BBFII, herein defined;
 - c) nucleotides of H4-1BBRI, herein defined;
 - d) nucleotides of H4-1BBRII, herein defined;
 - e) an isolated DNA encoding a soluble human 4-1BB polypeptide comprising the extracellular domain of human 4-1BB (amino acids 1-163 of SEQ ID NO:1) or a fragment thereof capable of binding a 4-1BB-L;
 - f) an isolated DNA of subparagraph e) wherein said DNA additionally encodes a polypeptide fused to the C-terminus of said soluble human 4-1BB polypeptide; or
 - g) derivatives and degeneracies of the above that encode for the same amino acid sequence as in any of the nucleotide sequences above.

(11) AU-B-77294/94
(10) 697482

-2-

8. A monoclonal antibody against H4-1BB which specifically recognizes receptor protein H4-1BB.

14. A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB, comprising:
 - a) at least a portion of said receptor protein H4-1BB corresponding to the extracellular portion of said receptor protein H4-1BB such that said portion of said receptor protein H4-1BB binds to said cell membrane ligands; and
 - b) a detection protein bound to said portion of said receptor protein H4-1BB such that ligand binding can be detected by relative activity assays for said detection protein.

OPI DATE 03/04/95 APPLN. ID 77294/94
 AOJP DATE 25/05/95 PCT NUMBER PCT/US94/10457



AU9477294


1)

(51) International Patent Classification ⁶ : C12N 15/12, 15/62, C07K 14/71, 14/715, 16/28, G01N 33/68		A1	(11) International Publication Number: WO 95/07984
			(43) International Publication Date: 23 March 1995 (23.03.95)
(21) International Application Number: PCT/US94/10457		(81) Designated States: AT, AU, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KR, LT, LU, LV, NL, NO, NZ, RU, SE, SI, SK, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(22) International Filing Date: 15 September 1994 (15.09.94)			
(30) Priority Data: 08/122,796 16 September 1993 (16.09.93) US		Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(71) Applicant: INDIANA UNIVERSITY FOUNDATION [US/US]; Showalter House, P.O. Box 500, Bloomington, IN 47402 (US).			
(72) Inventor: KWON, Byoung, S.; Indiana School of Medicine, Medical Science, Room 255, 63 Barnhill Drive, Indianapolis, IN 46223 (US).			
(74) Agent: MICHAELS, Christopher, A.; Barnard, Brown & Michaels, Suite 220, 306 E. State Street, Ithaca, NY 14850 (US).			

(54) Title: HUMAN RECEPTOR H4-1BB

(57) Abstract

The human receptor H4-1BB has been isolated, sequenced and disclosed herein. The cDNA of the human receptor H4-1BB is about 65 % homologous to the mouse cDNA 4-1BB and was isolated by using probes derived from cDNA 4-1BB. A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB was developed. It comprises the extracellular portion of the receptor protein H4-1BB and a detection protein (alkaline phosphatase) bound to the portion of the receptor protein H4-1BB. B-cells that have expressed a ligand to receptor protein H4-1BB can be treated with cells that have expressed receptor protein H4-1BB and B-cell proliferation may be induced. The use of H4-1BB to block H4-1BB ligand binding has practical application in the suppression of the immune system during organ transplantation. A monoclonal antibody against H4-1BB can be used to enhance T-cell proliferation by treating T-cells that have expressed receptor protein H4-1BB with the anti H4-1BB monoclonal antibody. Tumors transfected with H4-1BBL may be capable of delivering antigen-specific signals as well as the co-stimulatory signals and can be killed by human cytotoxic T lymphocytes.



HUMAN RECEPTOR H4-1BB

5

This application is a continuation-in-part of co-pending application Serial No. 08/012,269 filed February 1, 1993, which is a continuation-in-part of co-pending application Serial No. 07/922,996 filed July 30, 1992,
10 which is a continuation-in-part of copending application Serial No. 07/267,577 filed November 7, 1988.

The subject matter described herein was in part a subject invention of NIH Grants Nos. IR23AI23058-03, RO1 AI28175 and P60 KD20542 of which the present inventor was
15 the Principal Investigator and either the Donald Guthrie Foundation for Medical Research Inc. of Guthrie Square, Sayre, Pennsylvania 18849-1669 or Indiana University School of Medicine of Indianapolis, Indiana 46202, was the Grantee.

20

FIELD OF THE PRESENT INVENTION

The present invention relates to a previously unknown
25 human receptor protein, H4-1BB, which was isolated and identified based upon work with a homologous murine (mouse) receptor protein, 4-1BB, which was isolated and identified by specific expression of the T cell genes by the present inventor.

30

BACKGROUND OF THE PRESENT INVENTION

The immune system of humans and other species requires that white blood cells be made in the bone marrow, which
35 white blood cells include phagocytes, lymphocytes and B cells. As presently understood, the phagocytes include macrophage cells which scavenge unwanted materials such as

virus protein from the system. The lymphocytes include helper T cells and killer T cells and B cells as well as other cells, including those categorized as suppressor T cells. The B cells produce the antibodies. The killer T cells physically pierce the cell and the helper T cells facilitate the whole process. In any event, the immune process is facilitated by lymphokines.

Lymphokines are the proteins by which the immune cells communicate with each other. Scientists produce them in sufficient quantities for therapeutic use against immunologic diseases. There are many known lymphokine proteins and they include the interferons, interleukin-1,2,3,4,5,6,7, colony-stimulating factors, lymphotoxin, tumor necrosis factor and erythropoietin, as well as others.

Interleukin 1, secreted from macrophages activate the helper T cells and raise the body temperature causing fever which enhances the activity of the immune cells. The activated helper T Cells produce Interleukin 2 and Interleukin 2 stimulates the helper and killer T cells to grow and divide. The helper T cells also produce another lymphokine, B cell growth factor (BCGF), which causes B cells to multiply. As the number of B cells increases, the helper T cells produce another lymphokine known as the B cell differentiating factor (BCDF), which instructs some of the B cells to stop replicating and start producing antibodies. T cells also produce a lymphokine, gamma interferon (IF), which has multiple effects like Interleukin 2. Interferon helps activate killer T cells, enabling them to attack the invading organisms. Like BCGF, interferon increases the ability of the B cells to produce antibodies. Interferon also affects the macrophages to keep them at the site of the infection and help the macrophages to digest the cells they have engulfed. Gathering momentum with each kind of lymphokine signal between the macrophages and the T cells, the lymphokines amplify the immune system response and the virus protein or

other foreign matter on the infected cells is overwhelmed. There are many other lymphokines, maybe a hundred or more, which participate in the immune process. Many lymphokines are known and many are not.

5 Lymphokines are sometimes called intercellular peptide signals. Among scientists there is widespread use of cloned cell lines as lymphokine producers and the isolation of lymphokine mRNA has become a common technique. The mouse receptor protein, 4-1BB, was isolated and identified based
10 on specific expression of the T cell genes using a technique identified by the present inventor in a publication (Proc. Natl. Acad. Sci. USA. 84, 2896-2900, May 1987, Immunology). The protocol reported in this publication can be used by scientists to detect virtually
15 all of the lymphokines. The method is designed to detect virtually all mRNA expressed differentially and the mRNA sequences of the immune cells are expressed differentially (as they relate to the T cells and the killer T cells) even though the level of expression is low and the quantity of
20 the secreted lymphokine protein is low. The present inventor believes that the analysis described in the above identified publication can reveal biologically important molecules such as lymphokines because there are many indications that biologically important or active molecules
25 are coded by the most scarce messages. An example is a transforming growth factor (TGF) which is present as only one of a million clones.

Most T cell factors have been classically identified by recognizing biologic activities in assays, purifying the
30 protein information. An alternative approach is to isolate putative T cell genes based upon specific expression and then demonstrate the function of the unknown molecule. Using the aforesaid modified differential screening procedure, the present inventor cloned a series of T cell
35 subset-specific cDNAs from cloned helper T (HTL) L2 and cloned cytolytic T lymphocyte (CTL) L3.

A series of T-cell subset-specific cDNAs were isolated from cloned murine T-cells by employing a modified differential screening procedure. The nucleotide sequence and expression properties of some of the cDNA species have been reported. One of the genes not previously characterized, that encodes mouse receptor protein 4-1BB, was studied further. These studies have led to the isolation of the human homologue to 4-1BB, H4-1BB.

10

SUMMARY OF THE PRESENT INVENTION

The present invention includes the human receptor protein H4-1BB and the cDNA gene encoding for human receptor protein H4-1BB. The nucleotide sequence of the isolated cDNA is disclosed herein along with the deduced amino acid sequence. The cDNA gene identified as pH4-1BB was deposited at the Agricultural Research Service Culture Collection ^{on 25 August 1993} and assigned the accession number: NRRL B21131

The cDNA, and fragments and derivatives thereof, can be used as a probe to isolate DNA sequences encoding for proteins similar to the receptor protein encoded by the cDNA. The cDNA of the human receptor H4-1BB is about 65% homologous to the mouse cDNA 4-1BB and was isolated by using probes derived from cDNA 4-1BB. The cDNA gene identified as p4-1BB was deposited at the American Type Culture Collection at 12301 Parklawn Drive, Rockville, Maryland 20852 under ATCC No.: 67825.

The human receptor protein H4-1BB can be produced by:

- 1) inserting the cDNA of H4-1BB into an appropriate expression vector,
- 2) transfecting the expression vector into an appropriate transfection host,
- c) growing the transfected hosts in appropriate culture media and
- d) purifying the receptor protein from the culture media.

The protein and fragments and derivatives can be used: 1) as a probe to isolate ligands to human receptor protein H4-



1BB, 2) to stimulate proliferation of B-cells expressing H4-1BB ligands, or 3) to block H4-1BB ligand binding.

B-cell proliferation can be induced by treating B-cells that have expressed a ligand to receptor protein H4-1BB with cells that have expressed receptor protein H4-1BB. The use of H4-1BB to block H4-1BB ligand binding has practical application in the suppression of the immune system during organ transplantation. A similar costimulatory immune system pathway is being analyzed for this type of application. See "Mounting a Targeted Strike on Unwanted Immune Responses", Jon Cohen, Science, Vol. 257, 8-7-92; "Long Term Survival of Xenogeneic Pancreatic Islet Grafts Induced by CTLA4Ig", Lenschow et al, Science Vol. 257, 7-8-92; and "Immunosuppression in Vivo by a Soluble Form of the CTLA-4 T Cell Activation Molecule", Linsley et al, Science Vol. 257 7-8-92.

A monoclonal antibody against H4-1BB can be used to enhance T-cell proliferation by treating T-cells that have expressed receptor protein H4-1BB with the anti H4-1BB monoclonal antibody. Some tumors are potentially immunogenic but do not stimulate an effective anti-immune response in vivo. Tumors may be capable of delivering antigen-specific signals to T cells, but may not deliver the co-stimulatory signals necessary for full activation of T cells. Expression of the co-stimulatory ligand B7 on melanoma cells was found to induce the rejection of a murine melanoma in vivo. ("Tumor Rejection After Direct Co-Stimulation of CD8⁺ T Cells by B7-Transfected Melanoma Cells", Sarah E. Townsend and James P. Allison, Science Vol. 259, 1-5-93.) A monoclonal antibody against H4-1BB may be capable of the same effect as it is now known to induce T cell proliferation and activation.

A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB was developed. It comprises the extracellular portion of the receptor protein H4-1BB and a detection protein (alkaline phosphatase) bound to the portion of the receptor protein H4-1BB. The portion

of the receptor protein H4-1BB binds to the cell membrane ligands and binding can be detected by relative activity assays for the detection protein. The fusion protein is placed in the presence of a cell suspected to express the
5 receptor protein H4-1BB. Then the cell is washed of any fusion protein not bound to the cell membrane ligands. Once the washed cells are placed in the presence of a substrate for the detection protein and the relative activity of the detection protein can be measured.

10 The primary object of the present invention is the identification of the new human receptor, H4-1BB as identified herein by its sequence.

Another object of the present invention is to teach a fusion protein comprising the extracellular portion of H4-
15 1BB and a detection protein.

Still another object of the present invention is to teach methods of using the cDNA H4-1BB, the receptor protein H4-1BB, the monoclonal antibody and the legend for H4-1BB.

20

BRIEF DESCRIPTIONS OF THE FIGURES

Figure 1 shows the sequence for the cDNA of mouse
25 receptor protein 4-1BB and the regions used as PCR primers to obtain the human homologue H4-1BB.

Figures 2a and 2b show the nucleotide sequence and the deduced amino acid sequence of human receptor H4-1BB respectively.

30 Figures 3a and 3b illustrate the molecules involved in T-cell activation.

Figures 4a, 4b, and 4c illustrate a normal T-cell activation pathway.

Figures 5a, 5b, and 5c illustrate CTLA4-Ig alone, 4-
35 1BB/AP and CTLA4-Ig together and 4-1BB/AP alone respectively being used to block steps in the T-cell activation pathway.

DETAILED DESCRIPTION

In the following detailed description references are made to known procedures and studies, as well as published work of the applicant. These publications are incorporated herein by reference for clarity and listed in an appendix included at the end of this detailed description.

10 Isolation and characterization of mouse receptor 4-1BB

Figure 1 shows the nucleotide sequence and the deduced amino acid sequence of the mouse receptor 4-1BB. The nucleotides of the message strand are numbered in the 5' to 3' direction and numbers are shown on both sides of the sequence. Nucleotide residue 1 is the A of the initiation codon ATG, and the nucleotides on the 5' side of residue 1 are indicated by negative numbers. The predicted amino acid sequence is shown below the nucleotide sequence. Putative signal peptide is underlined. Stop codon is indicated by (---). Cysteine residues are highlighted by the dots. An unusual feature of 4-1BB sequence is that there is a potential polyadenylation signal of AATAAA at nucleotides 1158-1163 (Fig. 1 boxed). It was believed that this signal was functional because this gene produces at least two different sizes of mRNA.

The transcript of 4-1BB was inducible by concanavalin A in mouse splenocytes, T-cell clones, and hybridomas. The expression of 4-1BB transcripts was inhibited by cyclosporin A. The 4-1BB mRNA was inducible by antigen receptor stimulation but was not inducible by IL-2 stimulation in the cloned T-cells (1). The 4-1BB cDNA encodes a peptide of 256 amino acids containing a putative leader sequence, a potential membrane anchor segment, and other features of known receptor proteins. Therefore, the expression pattern of 4-1BB resembles those of lymphokine mRNAs while the sequence appeared consistent with those of receptor proteins.

The major species of 4-1BB on the cell surface appears to be a 55-kDa dimer. 4-1BB also appears to exist as a 30-kDa monomer and possibly as a 110-kDa tetramer. Since these 4-1BB species were immunoprecipitated from a homogenous population of cells (T cell clone F1), all forms potentially co-exist on each cell. A comparison of peptide digests from the 4-1BB monomer and dimer will be needed to determine whether 4-1BB exists as a homodimer on the cell surface. A variety of cell surface receptors such as the insulin receptor (2), the B cell surface immunoglobulin receptor (3), the T cell Ag receptor (4), the CD28 costimulatory receptor (5), and the CD27 T cell antigen (6) are composed of disulfide-bonded subunits. Receptor dimerization may be required for ligand binding and subsequent biochemical signaling.

4-1BB is not expressed on resting T cells but is inducible by activators which deliver a complete growth stimulus to the T cell. The combination of PMA and ionomycin is capable of mimicing those signals required for T cell proliferation. Although PMA or ionomycin alone induced 4-1BB mRNA, the combination of PMA and ionomycin resulted in optimal 4-1BB expression. Furthermore, the expression of 4-1BB was not transient. When purified splenic T cells were stimulated with immobilized anti-CD3, 4-1BB mRNA was expressed and this expression was maintained for up to 96 hrs poststimulation. Cell cycle analysis will be required to confirm that 4-1BB is expressed throughout cell cycle progression.

4-1BB is structurally related to members of the nerve growth factor receptor super-family. Although these receptors possess structurally similar ligand-binding properties (cysteine-rich regions), the cytoplasmic domains of these proteins are nonconserved which could allow for diversity in transmembrane signaling. Some members of this family are involved in the T or B cell activation process. There are in vitro functional data on the OX-40, CD40 and CD27 antigens. Antibodies against the OX-40 augment the T

cell response in a mixed lymphocyte reaction (7) and antibodies against CD40 enhance B-cell proliferation in the presence of a coactivator, such as PMA or CD20 antibodies, and synergize with IL-4 in vitro to induce B-cell differentiation and to generate long-term normal B cell lines (8). One monoclonal antibody, anti-1A4, which recognizes an epitope on the CD27 molecule inhibited calcium mobilization, IL-2 secretion, helper T cell function, and T cell proliferation. On the other hand, CLB-CD27/1, another anti-CD27 mAb enhanced proliferation of human T cells stimulated with PHA or anti-CD3 mAb (6). These results indicate that the CD27 molecule plays an important role in T cell activation. Except for TNFRs, NCFR and CD40, the ligands or cell surface molecules to which the members of the superfamily bind are not yet identified. Identification and characterization of the ligands to which the receptors bind will be helpful in better defining the physiologic role of 4-1BB.

To ascertain whether cell surface 4-1BB could contribute to T cell activation, the anti-4-1BB 53A2 was used as an antagonist to 4-1BB. These data suggested that 4-1BB does in fact have the potential to function as an accessory signaling molecule during T cell activation and proliferation. The addition of soluble 53A2 to purified splenic T cells stimulated with immobilized anti-CD3 resulted in an amplification of ³H thymidine incorporation compared to T cells stimulated with anti-CD3 alone. This pattern of enhancement ranged from 2- to 10- fold in three independent experiments.

In the original two signal model of Bretcher and Cohn, they proposed that signal 1, the occupancy of the T cell antigen receptor (TCR), resulted in inactivation of the T cell in the absence of signal 2, which is provided by accessory cells. This has since been confirmed by a variety of studies (9). The identification of the accessory cell CD28 as a potent costimulatory receptor on T cells was a significant contribution in beginning to characterize the

accessory signal(s) required for optimal T cell proliferation (10). It is possible that other cell surface molecules may contribute to these costimulatory activation requirements (11).

5 The biochemical signals delivered through 4-1BB are not completely known. One possibility considered was the observation that 4-1BB contains a putative p56^{lck} tyrosine kinase binding domain in its cytoplasmic tail. It was later determined that p56^{lck} tyrosinase kinase binds to
10 4-1BB. It will also be worthwhile to determine if 4-1BB-mediated signaling can regulate genes such as IL-2 and IL-2 receptor, whose expression is required for T cell activation and subsequent proliferation.

Although the precise functions of members of the Nerve
15 Growth Factor Receptor (NGFR) family appear to be diverse, an emerging theme is one in which these molecules may contribute in various ways to a maintenance of responsiveness or viability of the particular cell type in which they are expressed. For instance, NGF is absolutely
20 required for viability of neurons in vitro and in vivo (12). The crosslinking of CD40 by soluble antiCD40 monoclonal antibody blocks germinal center centrocytes from undergoing apoptosis in vitro (13). Signals delivered through CD40 may also aid in maintenance of responsiveness
25 to differentiation factors. The ligation of CD40 with anti-CD40 F(ab')₂ fragments in the presence of IL-4 induced large increases IgE synthesis (14). Also, anti-CD40 activated naive B cells treated with IL-10 and transforming growth factor- β became committed to IgA secretion (15).

30 In addition to sharing the molecular characteristics with the NGFR superfamily, it was noted that the 4-1BB contained a putative zinc finger structure of the yeast eIF-2 β protein (16). 4-1BB also shares a conserved region with the *sina* seven in absentia of *Drosophila*, which is
35 required for correct photoreceptor cell development (17). That particular region is also similar to the protein

product of the DGL7 gene of *Dictyostelium*, whose expression is specifically induced during aggregation by cAMP (18).

This region forms the pattern of C-X₂-C-X₃-C-X₃-H-X₃-C-X-C; and the cysteines and histidine are conserved in a similar space in 4-1BB, *sina*, and DGL7 proteins. Ten of 24 amino acids between the 4-1BB and *sina* proteins are identical, and 3 of 24, are conservative substitutes. The conserved pattern suggests that these amino acids are functionally important. The *sina* protein is localized in the nucleus, suggesting that it has a regulatory function in cells. The fact that the amino acid sequence of 4-1BB contains features like a zinc finger motif, a nuclear protein, and a receptor domain suggests that 4-1BB may play diverse roles during cellular proliferation and differentiation.

4-1BB may represent another cell-surface molecule involved in T cell-APC interactions. The 4-1BB-AP fusion protein specifically bound to mature B-cell lines, anti- μ -activated primary B cells, and mature macrophage-cell lines. 4-1BB-AP bound at low or insignificant levels to immature B- and macrophage-cell lines, T-cell clones, T-cell lines, primary culture T cells, and various nonlymphoid-cell lines. Since 4-1BB-AP binds to mature B cells and macrophages, it is possible that signals delivered upon 4-1BB binding may modulate APC functions in some way. This possibility remains to be explored.

Chalupny and colleagues (19) have proposed that 4-1BB Rg, a fusion protein consisting of the extracellular domain of 4-1BB and the Fc region of human IgG, bound to the extracellular matrix (ECM). The highest level of 4-1BB Rg binding was to human vitronectin. In data not shown, an ELISA was performed using 4-1BB-AP and human vitronectin (Yelios Pharmaceuticals/GIBCO-BRL, Grand Island, NY.) immobilized at 0.007 μ g-10 μ g per well on microtiter plates. No binding of 4-1BB-AP based on AP activity was observed. To rule out the possibility that 4-1BB-AP was binding to proteins extrinsically attached to the cell

surface (possible extracellular matrix components), B-cell lymphomas were washed in acid conditions prior to the binding assay. 4-1BB-AP still bound specifically to mature B-cell lymphomas. It is still to be determined whether a 4-1BB-ligand specifically expressed on B cells and macrophages exists, and whether 4-1BB-AP may bind to the ECM under particular binding conditions. It is possible that the ECM could facilitate the binding of 4-1BB to a specific cell-surface ligand.

10 B cells and helper T cells interact with each other through receptors on B cells binding to their specific counter-receptors on T cells. It is thought that this interaction results in a cascade of biochemical signaling relays between these two cell types (20). As this
15 interaction proceeds, these cells become committed to enter the S phase of the cell cycle. Initial interactions between TCR and CD4 on T cells, and processed antigen-MHC II on B cells, do not result in B cells capable of entering the cell cycle (21). However, studies from in vitro systems
20 suggest that once T-cells are stimulated, they express newly synthesized or modified cell-surface molecules capable of inducing B cells to enter the cell cycle (22, 23). This T-cell function is not antigen-specific or MHC-restricted (24). In addition, soluble factors are not
25 required for the activated Th induction of B-cell activation (25). Once B cells enter the cell cycle, IL-4 induces B cells to progress from G₁ to S phase. The ability of activated T cells or T-cell membranes to promote the entry of B cells into the cell cycle can be blocked by
30 either cycloheximide or cyclosporin A treatment (26, 27). These newly expressed membrane proteins appear to be "lymphokine-like" in their induction characteristics.

4-1BB has expression properties which meet the requirements of a B-cell costimulator. 4-1BB is inducible
35 by anti-CD3 or TCR-mediated T-cell stimulation, and its expression is sensitive to cyclosporin A as well as cycloheximide treatment (28). Interestingly,

paraformaldehyde-fixed SF21-4-1BB cells, synergized with anti- μ in inducing B-cell proliferation. The costimulation of splenic B cells by SF21-4-1BB occurred at optimal (10 $\mu\text{g/ml}$) and suboptimal (1.0-0.1 $\mu\text{g/ml}$) doses of anti- μ . The addition of SF21-4-1BB cells to resting B cells, did not result in significant B-cell proliferation. SF21-4-1BB cells did not synergize with TPA or ionomycin, or suboptimal concentrations of LPS in inducing B-cell proliferation.

Although the baculovirus system has been used to express large amounts of recombinant soluble proteins, this system may be utilized for the expression of recombinant cell-surface proteins. The baculovirus infection provides a convenient means to express uniformity high levels of recombinant protein on a per cell basis. It is noteworthy, that the addition of SF21 cells alone did not result in significant levels of costimulation. This can be a potential problem when using cos- or L- cell lines which can exhibit strong costimulator activity on their own.

Another member of the NGFR superfamily, CD40, is expressed on B cells and interacts with gp39, a molecule expressed on activated T cells. The cDNAs encoding the murine (29) and human (30) gp39 proteins have been cloned; this cell surface molecule is a type II membrane protein with homology to tumor necrosis factor. Noelle et al. (31) found that a CD40-inununoglobulin fusion protein, is capable of blocking T cell-induced B-cell proliferation and differentiation in a dose-dependent manner. Armitage et al. have isolated a cDNA for murine gp39 and showed that gp39 could induce B-cell proliferation in the absence of co-stimuli, and result in IgE production in the presence of IL-4-. Hollenbaugh et al. (32) have shown that COS cells transfected with human gp 39 can synergize with either TPA or anti-CD20 in inducing human B-cell proliferation and is able to stimulate B cells without a costimulator only at low levels. These data indicate that CD40 may be one of the

B-cell-surface molecules that transmit signals during physical contact with T cells.

Cell-surface receptors communicate with their external milieu by interacting either with soluble factors or other cell surface molecules expressed on neighboring cells. The role of biochemical signals delivered by cell-cell contact versus those delivered by soluble factors interacting with cell surface receptors is not clear. The NGFR superfamily is unusual for the TNFR I and II as well as the NGFR bind to more than one ligand. The TNFRs I and II both bind to TNF- α and TNF-R (33). The NGFR binds to NGF, brain-derived neurotrophic factor, and neurotrophin-3 (34).

In addition, one ligand may function as both a cell surface and soluble ligand. Recent evidence on the CD4-0 ligand, gp39, suggests that this ligand can exist as a membrane bound as well as a soluble ligand (35). It may be possible that 4-1BB is secreted and interacts with B cells in a soluble form as well as a membrane bound form. A member of the NGFR receptor family, CD27, which is expressed on T cells, is secreted in addition to being expressed on the cell surface (36). It is also possible that more than one ligand (soluble and cell surface) may bind to 4-1BB.

25 Isolation of the human homologue, H4-1BB

In order to isolate the human homologue (H4-1BB) of mouse 4-1BB two sets of polymerase chain reaction (PCR) primers were designed. To design the PCR primers, the amino acid sequence among the members of nerve growth factor receptor (NGFR) superfamily were compared because 4-1BB is a member of the superfamily (37). The amino acid sequences employed were mouse 4-1BB (38), human NGFR (39), human tumor necrosis factor receptors (33), human CD40 (40), and human CD27 (6). The areas of sequence conservation among the NGFR superfamily were chosen.

Forward primer I (H4-1BBFI) spans from amino acids 36 to 41 and forward primer II (H4-1BBFII) spans from amino

acids 52 to 58 of the mouse 4-1BB. Reverse primer I (H4-1BBRI) spans from amino acids 116 to 121 and reverse primer II (H4-1BBRII) spans from amino acids 122 to 128 of mouse 4-1BB. The regions used as PCR primers in mouse 4-1BB are indicated in Fig. 1.

The degenerative oligonucleotide sequence of each primer is as follows:

```

10      H4-1BBFI:      5' TTC TGT CGA AAA TAT AAT CC 3'
                        T   C A       G   C   C

      H4-1BBFII:      5' TTC TCC TCC ATT GGT GGA CA 3'
                        T   G   G       C
                                A

15      H4-1BBRI:      5' CC CAA CGA ACA GGT TTT ACA 3'
                        G   CT G       C   G

      H4-1BBRII:      5' TT TTG ATC ATT AAA TGT TCC 3'
                        C   G   G   G
  
```

Peripheral blood lymphocytes from normal healthy individuals were isolated and activated with PMA (10 ng/ml) and ionomycin (1 μ M). mRNA from the lymphocytes was isolated. Using reverse transcriptase the human lymphocyte mRNA was converted to single-stranded cDNA. The cDNA was then amplified with Taq polymerase with combination of the primers. The combination of primers was as follows: H4-1BBFI vs H4-1BBRI; H4-1BBFI vs H4-1BBRII; H4-1BBFII vs H4-1BBRI; and H4-1BBFII vs H4-1BBRII.

The primer set of H4-1BBFII and H4-1BBRII produced a specific band of ~240 bp. The 240 bp is an expected size of human 4-1BB if the human homologue protein is similar to mouse 4-1BB in size. The PCR product (240 bp) was cloned in PGEM3 vector and sequenced. One open reading frame of the PCR product was ~65% identical to mouse 4-1BB. Therefore, it was concluded that the 240 bp PCR product is the human homologue of mouse 4-1BB. The 240 bp PCR product was used to screen λ gt11 cDNA library of activated human T lymphocytes. An ~0.85 kb cDNA was isolated. The sequence of the cDNA is shown in Figure 2 and the predicted amino acid sequence is shown in Figure 2b. The same information



16

is shown is the sequence listing attached to this specification in sequence id. 1.

An expression plasmid to produce H4-1BB-AP fusion protein was constructed. The 5' portion of the H4-1BB cDNA including sequences encoding the signal sequence and the entire extracellular domain, was amplified by PCR. For correctly oriented cloning, a Hind III site on the 5' end of the forward primer and a Bgl II site on the 5' end of the reverse primer were created.

The Hind III - Bgl II H4-1BB fragment was inserted into the mammalian expression vector APTaq-1, upstream of the coding sequence for human placental alkaline phosphatase (AP). The oligonucleotides PCR primers used for the amplification of 5' portion of H4-1BB are as follows:

Forward

primer: 5' AAT AAG CTT TGC TAG TAT CAT ACC T 3'

Reverse

primer: 5' TTA AGA TCT CTG CGG AGA GTG TCC TGG CTC 3'

H4-1BB-AP will be used to identify cells and tissues that express ligand for human 4-1BB (i.e. H4-1BBL). The studies with mouse 4-1BB indicated that the ligand for 4-1BB is on the cell surface. B cells and macrophages were major cells that express 4-1BBL. It is expected that H4-1BBL also expresses on human B cells and macrophages.

A mammalian expression cDNA library will be generated from human cell lines that express H4-1BBL. The library will be screened by [¹²⁵] I-labeled H4-1BB-AP. cDNA for H4-1BBL will then be isolated and characterized. Soluble recombinant H4-1BBL will then be produced. Both H4-1BB-AP and H4-1BBL will be used to suppress or enhance immune responses as described below. Monoclonal antibody to H4-1BB and H4-1BBL will be produced.

According to studies with mouse 4-1BB, 4-1BB acts as a costimulatory signal. It is expected that H4-1BB will

act as a costimulatory signal for T cell activation. Mouse 4-1BB helped B cells with proliferation and differentiation. It is expected that H4-1BB will do the same. H4-1BB-AP, H4-1BBL and monoclonal antibody can be used to suppress or enhance human immune responses.

Figures 3a and 3b illustrate the molecules involved in T-cell activation. During early T-cell activation (cognitive phase), resting T cells express the TCR/CD3 complex and other "accessory" molecules. Among these constitutively expressed molecules, CD4 (or CD8), LFA-1 and CD28 are probably the ones to receive costimulatory signals. Initial interaction with the TCR/CD3 complex in combination with these 'accessory' costimulatory signals leads to subsequent expression of additional receptor molecules such as CD28, CTLA4, and 4-1BB. These newly expressed molecules are probably going to receive additional important costimulatory signals at later stages of T-cell activation (clonal expansion).

20 Suppression of immune responses.

Figures 4a-c illustrate a normal T-cell activation pathway. Figures 5a-c illustrate the blocking of immune responses with soluble chimera of 4-1BB. If 4-1BB plays a role in T-cell activation, blocking of the interaction to its ligand on antigen-presenting cells should result in suppression of T-cell dependent immune responses. It is well documented that blocking of the interaction of CD28 to its counter-receptor B7 suppresses in varying degrees, both in vivo antibody production and cell-mediated immune responses. Blocking of both interactions should result in a more effective immunosuppression; since 4-1BB is induced during T-cell activation. Blocking of the interaction of 4-1BB to its ligand may be of importance at later stages of the activation process where the CD28/B7 interaction may no longer be of relevance.

As illustrated with mouse receptor 4-1BB and mouse ligand 4-1BBL above, addition of H4-1BB-AP will coat the H4-

1BBL expressing cells and block the normal interaction between H4-1BB and H4-1BBL. This will lead to immunosuppression. This type of immunosuppression is antigen-specific. Therefore it avoids the generalized immunosuppression produced by antiCD3 or cyclosporin A treatments. H4-1BB-AP treatment can be used to treat certain autoimmune diseases and to facilitate organ transplantation.

10 Immune enhancement.

H4-1BB may function at the late stage of T cell activation and may be a critical molecule for completion of T cell activation. Most tumors display tumor-specific antigens. One reason, however, why immunogenic tumors can escape host immunity is that tumor-reactive T cells receive inadequate costimulation. The introduction of the costimulatory molecules, such as H4-1BB into the tumor, therefore, could enhance the antitumor immunity of cytotoxic T cells (CTL). H4-1BBL can be expressed in cell-specific fashion. For example, the H4-1BBL can be expressed in melanoma using melanocyte-specific promoter such as tyrosinase promoters. The H4-1BBL-expressing melanoma will stimulate cytotoxic T cells through H4-1BB and activate the melanoma-specific CTL. The activated melanoma-specific CTL can destroy melanoma.

Appendix to References Incorporated by Reference

1. Smith, C.A., Davis, T., Anderson, D., Solam, L., Beckmann, M.P., Jerzy, R., Dower, S.K., Cosman, D., and Goodwin, R.G. 1990. A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. Science 248:1019-1023.
2. Ebina, Y., L. Ellis, K. Jaruagin, M. Edery, L. Graf, E. Clauser, J. On, F. Marizrz, Y.W. Kan, J.D. Goldfine, R.A. Roth and W.J. Rutter, 1985, The human insulin receptor cDNA: the structural basis for hormone-activated transmembrane signalling, Cell 40:747.
3. Vassali, R., R. Tedghi, B. Listowska-Bernstein, A. Tartakoff and J.C. Jaton, 1979, Evidence for hydrophobic

region within heavy chains of mouse B lymphocyte membrane-bound IgM, Proc. Natl. Acad. Sci. USA 76:5515.

4. Haskins, K., R. Kubo, J. White, M. Pigeon, J. Kappler
5 and P. Marrack, 1983, The major histocompatibility complex-restricted antigen receptor on T cells I Isolation with monoclonal antibody, J. Exp. Med. 157:1149.
5. Lesslaver, W. and H. Gmunder, 1986, Biochemical
10 characterization of the 9.3 antigens of human T-cells: simultaneous expression of disulfide-bonded 90-Kiladalton dimers and free subunits at the cell surface, Mol. Immunol. 23:271.
- 15 6. Van Lier, R., J. Borst, T. Vroom, H. Klein, P. Mourik, W. Zeijlemaker and C. Melife, 1987, Tissue distribution and biochemical and functional properties of Tp55 (CD27) a novel T cell differentiation antigen, J. Immunol. 139:1589.
- 20 7. Mallett, S., S. Fossum and A. Barclay, 1990, Characterization of the MRC OX40 antigen of activated CD4 positive T lymphocytes-a molecule related to nerve growth factor receptor, EMBO J. 9:1603.
- 25 8. Banchereau, J., P. Paoli, A., Valle, E. Garcia and F. Roussel, 1991, Long-term human B cell lines dependent on interleukin-4 and antibody to CD40, Science 251:70.
- 30 9. Moeller, D.L., M.K. Jenkins and R.H. Schwartz, 1989, Clonal expansion versus functional clonal inactivation: a co-stimulatory signalling pathway determines the outcome of T cell antigen receptor occupancy, Ann. Rev. Immunol. 7:445.
- 35 10. June, D.H., J.A. Ledbetter, P.S. Linsley and C.B. Thompson, 1989, Role of CD28 receptor in T cell activation, Immunol. Today 11:211.
- 40 11. Yang, L., B. Jones, A. Aruffo, K.M. Sullivan, P.S. Linsley and C.A. Janeway, Jr., 1992, Heat stable antigen is a co-stimulatory molecule for CD4 T cell growth, J. Exp. Med. 175:437.
- 45 12. Yamori, T., 1992, Molecular mechanisms for generation of neural diversity and specificity: roles of polypeptide factors in development of post-mitotic neurons, Neurosci. Res. 12:545.
- 50 13. Liu, Y.J., D.E. Joshua, G.T. Williams, C.A. Smith, J. Gordon and I.C. M. MacLennan, 1989, Mechanism of antigen-driven selection in germinal centres, Nature, 342:929.
14. Jabara, H. H., S. M. Fu, R.S. Geha and D. Vercelli, 1990, CD40 and IgE: synergism between anti-CD40 monoclonal
55 antibody and interleukin 4 in the induction of IgE

- synthesis by highly purified human B cells, J. Exp. Med. 172:1861.
15. Defrance, R., B. Vanbervliet, F. Briere, I. Durnad, F. Roussle and J. Banchereau, 1992, Interleukin 10 and transforming growth factor β cooperate to induce anti-CD40 activated naive human B cells to secrete immunoglobulin A, J. Exp. Med. 175:671.
- 10 16. Donahue, T., Cigan, A., Pahich, E. and Valavicius, B., Mutations at a Zn(II) finger motif in the yeast eIF-2 β gene alter ribosomal start-site selection during the scanning process, Cell 54 (1988) 621-632).
- 15 17. Carthew, R.W and Rubin, G.M., seven in absentia, a gene required for specification of R7 cell fate in the Drosophila eye, Cell, 63 (1990) 561-577.
- 20 18. Driscoll, D.M. and Williams, J.G., Two divergently transcribed genes of Dictyostelium discoideum are cyclic AMP-inducible and coregulated during development, Mol. and Cell. Biol. 7 (1987) 4482-4489.
- 25 19. Chalupny, N.J., Peach, R., Hollenbaugh, D., Ledbetter, J.A., Farr, A.G. and Aruffo, A., 1992, Proc. Natl. Acad. Sci USA 89:10360-10364.
- 30 20. Noelle, R.J., and Snow, E.C., 1991, The FASEB J. 5:2770-2776.
21. Noelle, R. and Snow, E., 1990, Immunol. Today 11:361-368.
- 35 22. Zurawski, G., Benedik, M., Kamb, B.J., Abrams, J.S., Zurawski, S.M. and Lee, F.D. (1986) Science 232:772-775.
23. Kinachi, T. (1986) Nature 325,70-73.
- 40 24. Gershenfeld, H.K. and Weissman, I.L. Science (1986) 232,854-858.
25. Biggin, M., Gison, T. and Hung, G. (1983 Proc. Natl. Acad. Sci. USA 80,3963-3965.
- 45 26. Hodgkin, P.D., Yamashita, L.C., Coffman, R.L. and Kehry, M.R., 1990, J. Immunol. 145:2025-2034.
27. Barlett, W.C., McCann, J., Shephaer, D.M., Roy, M. and Noelle, R.J., 1990, J. Immunol. 145:3956-3962.
- 50 28. Kwon, B.S., Kestler, D.P., Eshhar, Z., Oh, K., and Wakulchik, M. 1989. Expression characteristics of two potential T cell mediator genes. Cell. Immunol. 121:414-422.
- 55

29. Armitage, R., Fanslow, W., Strockbine, L., Sato, T., Clifford, K., MacDuff, B., Anderson, D., Gimpel, S., Davis-Smith, T., Maliszewski, C., Clark, E., Smith, C., Grabstein, K., Cosman, D. and Spriggs, M., 191, Nature 5 357:80-82.
30. Kwon, B., Kestler, D., Lee, E., Wakulchik, M. and Young J. (1988) J. Exp. Med (1988) (In press).
- 10 31. Noelle, R.J., Roy, M., Shepherd, D.M., Stamenkovic, I., Ledbetter, J.A. and Aruffo, A., 1992, Proc. Natl. Acad. Sci. USA 89:6550-6554.
- 15 32. Hollenbaugh, D., Grosmaier, L.S., Kullas, C.D., Chalupny, N.J., Braesch-Andersen, S., Noelle, R.J., Stamenkovic, I., Ledbetter, J.A. and Aruffo, A., 1992, EMBO 11:4314-4321.
- 20 33. Schall, T.J., M. Lewis, K.J. Koller, A. Lee, G.C. Rice, G.H. W. Wong, T. Gatanaga, G.A. Granger, R. Lentz, H. Raab, W.J. Kohr and D.V. Goeddel, 1990, Molecular cloning and expression of a receptor for human tumor necrosis factor, Cell 61:361.
- 25 34. Klein, R., Nanduri, V., Jing, S., Lamballe, F., Tapley, P., Bryant, S., Cordon-Cardo, C., Jones, K.R., Reichardt, L.F. and Barbacid, M., 1991, Cell 66:395-403.
- 30 35. Armitage, R.J., Sato, T.A., Macduff, B.M., Clifford, K.N., Alpert, A.R., Smith, C.A. and Fanslow, W.C., 1992, Eur. J. Immunol. 22:2071-2076.
- 35 36. Hintzen, R.Q., deJong, R., Hack, E.E., Chamuleau, M., de Vries, E.F.R., ten Berge, I.J.M., Borst, J. and van Lier, R.A.W., 1991, J. Immunol. 147:29-35.
37. Mallett, S., and Barclay, A.N. 1991. A new super-family of cell surface proteins related to the nerve growth factor receptor. Immunol. Today. 12:220-223.
- 40 38. Kwon, B.S., and Weissman, S.M. 1989. cDNA sequences of two inducible T-cell genes. Proc. Natl. Acad. Sci. USA. 86:1963-1967.
- 45 39. Johnson, D., Lanahan, A., Buck C.R., Sehgal, A., Morgan, C., Mercer, E., Bothwell, M., and Chao, M. 1986. Expression and structure of the human NGF receptor. Cell 47:545-554.
- 50 40. Stamenkovic, I., Clark, E., and Seed, B. 1989. A B-lymphocyte activation molecule related to the nerve growth factor receptor and induced by cytokines in carcinomas. EMBO. J. 8:1403-1408.

The foregoing description has been directed to particular embodiments of the invention in accordance with the requirements of the Patent Statutes for the purposes of illustration and explanation. It will be apparent, 5 however, to those skilled in this art that many modifications and changes will be possible without departure from the scope and spirit of the invention. It is intended that the following claims be interpreted to embrace all such modifications.

SEQUENCE LISTING

- (1) GENERAL INFORMATION:
(i) APPLICANT: Byoung Se Kwon
(ii) TITLE OF INVENTION: New Human Receptor and Related Products and Methods
(iii) NUMBER OF SEQUENCES: 1
(iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Barnard & Brown
(B) STREET: 306 E. State St., Suite 220
(C) CITY: Ithaca
(D) STATE: New York
(E) COUNTRY: United States
(F) ZIP: 14850
(v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Diskette, 3.50 inch, 1.4Mb storage
(B) COMPUTER: IBM AT Compatible
(C) OPERATING SYSTEM: MS DOS, Version 5.0
(D) SOFTWARE: Special QBasic program
(vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:
(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/012,269
(B) FILING DATE: 2/1/93
(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 07/922,996
(B) FILING DATE: 7/30/92
(vii) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 07/267,577
(B) FILING DATE: 11/7/88
(viii) ATTORNEY/AGENT INFORMATION:
(A) NAME: Michaels, Christopher A.
(B) REGISTRATION NUMBER: 34,390
(C) REFERENCE/DOCKET NUMBER: kwnh41bb
(ix) TELECOMMUNICATION INFORMATION:
(A) TELEPHONE: 607-273-1711
(B) TELEFAX: 607-273-2609
(C) TELEX:
(2) INFORMATION FOR SEQ ID NO: 1 :
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 838
(B) TYPE: nucleic acid
(C) STRANDEDNESS: double stranded
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: cDNA to mRNA
(iii) HYPOTHETICAL: No
(iv) ANTI-SENSE: No
(v) FRAGMENT TYPE: n/a
(vi) ORIGINAL SOURCE:
(A) ORGANISM: Human
(B) STRAIN:
(C) INDIVIDUAL ISOLATE: H4-1BB #1
(D) DEVELOPMENTAL STAGE: Differentiated T-cell
(E) HAPLOTYPE:
(F) TISSUE TYPE:
(G) CELL TYPE: Lymphocytes
(H) CELL LINE:
(I) ORGANELLE:
(vii) IMMEDIATE SOURCE:
(A) LIBRARY: cDNA library
(B) CLONE:
(viii) POSITION IN GENOME:
(A) CHROMOSOME/SEGMENT:
(B) MAP POSITION:
(C) UNITS:
(ix) FEATURE:
(A) NAME/KEY: H4-1BB

(B) LOCATION:
 (C) IDENTIFICATION METHOD: Similarity to mouse 4-1BB and other members of NGFR superfamily
 (D) OTHER INFORMATION:
 (x) PUBLICATION INFORMATION:
 (A) AUTHORS: Kwon, B.S., and Weissman, S.M.
 (B) TITLE: cDNA sequences of two inducible T-cell genes
 (C) JOURNAL: Proc. Natl. Acad. Sci. USA
 (D) VOLUME: 86
 (E) ISSUE:
 (F) PAGES: 1963-1967
 (G) RELEVANT RESIDUES: all

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:
 AATCAGCTTT GCTAGTATCA TACCTGTGCC AGATTTTCATC

																	40
ATG	GGA	AAC	AGC	TGT	TAC	AAC	ATA	GTA	GCC	ACT	CTG	TTG	CTG	GTC			85
Met	Gly	Asn	Ser	Cys	Tyr	Asn	Ile	Val	Ala	Thr	Leu	Leu	Leu	Val			
1				5					10					15			
CTC	AAC	TTT	GAG	AGG	ACA	AGA	TCA	TTG	CAG	GAT	CCT	TGT	AGT	AAC			130
Leu	Asn	Phe	Glu	Arg	Thr	Arg	Ser	Leu	Gln	Asp	Pro	Cys	Ser	Asn			
				20					25					30			
TGC	CCA	GCT	GGT	ACA	TTC	TGT	GAT	AAT	AAC	AGG	AAT	CAG	ATT	TGC			175
Cys	Pro	Ala	Gly	Thr	Phe	Cys	Asp	Asn	Asn	Arg	Asn	Gln	Ile	Cys			
				35					40					45			
AGT	CCC	TGT	CCT	CCA	AAT	AGT	TTC	TCC	AGC	GCA	GGT	GGA	CAA	AGG			220
Ser	Pro	Cys	Pro	Pro	Asn	Ser	Phe	Ser	Ser	Ala	Gly	Gly	Gln	Arg			
				50					55					60			
ACC	TGT	GAC	ATA	TGC	AGG	CAG	TGT	AAA	GGT	GTT	TTC	AGG	ACC	AGG			265
Thr	Cys	Asp	Ile	Cys	Arg	Gln	Cys	Lys	Gly	Val	Phe	Arg	Thr	Arg			
				65					70					75			
AAG	GAG	TGT	TCC	TCC	ACC	AGC	AAT	GCA	GAG	TGT	GAC	TGC	ACT	CCA			310
Lys	Glu	Cys	Ser	Ser	Thr	Ser	Asn	Ala	Glu	Cys	Asp	Cys	Thr	Pro			
				80					85					90			
GGG	TTT	CAC	TGC	CTG	GGG	GCA	GGA	TGC	AGC	ATG	TGT	GAA	CAG	GAT			355
Gly	Phe	His	Cys	Leu	Gly	Ala	Gly	Cys	Ser	Met	Cys	Glu	Gln	Asp			
				95					100					105			
TGT	AAA	CAA	GGT	CAA	GAA	CTG	ACA	AAA	AAA	GGT	TGT	AAA	GAC	TGT			400
Cys	Lys	Gln	Gly	Gln	Glu	Leu	Thr	Lys	Lys	Gly	Cys	Lys	Asp	Cys			
				110					115					120			
TGC	TTT	GGG	ACA	TTT	AAC	GAT	CAG	AAA	CGT	GGC	ATC	TGT	CGA	CCC			445
Cys	Phe	Gly	Thr	Phe	Asn	Asp	Gln	Lys	Arg	Gly	Ile	Cys	Arg	Pro			
				125					130					135			
TGG	ACA	AAC	TGT	TCT	TTG	GAT	GGA	AAG	TCT	GTG	CTT	GTG	AAT	GGG			490
Trp	Thr	Asn	Cys	Ser	Leu	Asp	Gly	Lys	Ser	Val	Leu	Val	Asn	Gly			
				140					145					150			
ACG	AAG	GAG	AGG	GAC	GTG	GTC	TGT	GGA	CCA	TCT	CCA	GCT	GAC	CTC			535
Thr	Lys	Glu	Arg	Asp	Val	Val	Cys	Gly	Pro	Ser	Pro	Ala	Asp	Leu			
				155					160					165			
TCT	CCG	GGA	GCA	TCC	TCT	GTG	ACC	CCG	CCT	GCC	CCT	GCG	AGA	GAG			580
Ser	Pro	Gly	Ala	Ser	Ser	Val	Thr	Pro	Pro	Ala	Pro	Ala	Arg	Glu			
				170					175					180			
CCA	GGA	CAC	TCT	CCG	CAG	ATC	ATC	TCC	TTC	TTT	CTT	GCG	CTG	ACG			625
Pro	Gly	His	Ser	Pro	Gln	Ile	Ile	Ser	Phe	Phe	Leu	Ala	Leu	Thr			
				185					190					195			

TCG	ACT	GCG	TTG	CTC	TTC	CTG	CTG	TTC	TTC	CTC	ACG	CTC	CGT	TTC	670
Ser	Thr	Ala	Leu	Leu	Phe	Leu	Leu	Phe	Phe	Leu	Thr	Leu	Arg	Phe	
				200					205					210	
TCT	GTT	GTT	AAA	CGG	GGC	AGA	AAG	AAA	CTC	CTG	TAT	ATA	TTC	AAA	715
Ser	Val	Val	Lys	Arg	Gly	Arg	Lys	Lys	Leu	Leu	Tyr	Ile	Phe	Lys	
				215					220					225	
CAA	CCA	TTT	ATG	AGA	CCA	GTA	CAA	ACT	ACT	CAA	GAG	GAA	GAT	GGC	760
Gln	Pro	Phe	Met	Arg	Pro	Val	Gln	Thr	Thr	Gln	Glu	Glu	Asp	Gly	
				230					235					240	
TGT	AGC	TGC	CGA	TTT	CCA	GAA	GAA	GAA	GAA	GGA	GGA	TGT	GAA	CTG	805
Cys	Ser	Cys	Arg	Phe	Pro	Glu	Glu	Glu	Glu	Gly	Gly	Cys	Glu	Leu	
				245					250					255	
TGAAATGGAA	GTCAATAGGG	CTGTTGGGAC	TTT												838

□

The claims defining the invention are as follows:

1. A cDNA encoding for human receptor protein H4-1BB having a nucleotide sequence as shown in SEQ ID NO: 1.
2. The cDNA of claim 1, identified as pH4-1BB deposited at the Agricultural Research Service Culture Collection with the accession number NRRL B21131.
3. An isolated DNA, wherein said DNA comprises a nucleotide sequence selected from:
 - a) nucleotides of H4-1BBFI, herein defined;
 - b) nucleotides of H4-1BBFII, herein defined;
 - c) nucleotides of H4-1BBRI, herein defined;
 - d) nucleotides of H4-1BBRII, herein defined;
 - e) an isolated DNA encoding a soluble human 4-1BB polypeptide comprising the extracellular domain of human 4-1BB (amino acids 1-163 of SEQ ID NO:1) or a fragment thereof capable of binding a 4-1BB-L;
 - f) an isolated DNA of subparagraph e) wherein said DNA additionally encodes a polypeptide fused to the C-terminus of said soluble human 4-1BB polypeptide; or
 - g) derivatives and degeneracies of the above that encode for the same amino acid sequence as in any of the nucleotide sequences above.
4. An isolated DNA encoding a human 4-1BB polypeptide, wherein said DNA comprises a nucleotide sequence selected from:
 - a) nucleotides 41-805 of SEQ ID NO:1;
 - b) nucleotides 110-805 of SEQ ID NO: 1; or
 - c) a nucleotide sequence that is degenerate as a result of the genetic code to a nucleotide sequence of (a) or (b).
5. The receptor protein of H4-1BB produced by
 - a) transfecting an expression vector comprising at least nucleotides 110-805 of SEQ ID NO:1 into an appropriate



transfection host;

b) growing said transfected host in an appropriate culture medium; and

c) purifying the receptor protein from said culture medium.

5

6. A protein having the amino acid sequence shown in Figure 2b.

10

7. The protein of claim 6 and fragments and derivatives thereof, wherein said fragments and derivatives:

a) can be used as a probe to identify ligands to receptor protein H4-1BB;

b) can be used to stimulate proliferation of B-cells expressing H4-1BB ligands; or

c) can be used to block H4-1BB ligand binding.

15

8. A monoclonal antibody against H4-1BB which specifically recognizes receptor protein H4-1BB.

9. A hybridoma capable of producing a monoclonal antibody against H4-1BB which specifically recognizes receptor protein H4-1BB.

20

10. A method of enhancing T-cell proliferation comprising the step of treating T-cells that have expressed receptor protein H4-1BB with the monoclonal antibody of claim 8.

11. The method of claim 10 further comprising the step of conducting said treatment in the presence of protein tyrosinase kinase.

25

12. A method of enhancing T-cell activation comprising the step of treating T-cells that have expressed receptor protein H4-1BB with the monoclonal antibody of claim 8.

30

13. The method of claim 12 further comprising the step of conducting said treatment in the presence of protein tyrosinase kinase.



14. A fusion protein for detecting cell membrane ligands to human receptor protein H4-1BB, comprising:

- 5 a) at least a portion of said receptor protein H4-1BB corresponding to the extracellular portion of said receptor protein H4-1BB such that said portion of said receptor protein H4-1BB binds to said cell membrane ligands; and
- 10 b) a detection protein bound to said portion of said receptor protein H4-1BB such that ligand binding can be detected by relative activity assays for said detection protein.

15. The fusion protein of claim 14 wherein said detection protein is alkaline phosphatase.

16. A method of detecting cell membrane ligands to human receptor protein H4-1BB, comprising:

- 15 a) placing a fusion protein in the presence of a cell suspected to express said receptor protein H4-1BB, wherein said fusion protein comprises:
- 20 i) at least a portion of said receptor protein H4-1BB corresponding to the extracellular portion of said receptor protein H4-1BB such that said portion of said receptor protein H4-1BB binds to said cells membrane ligands, and
- 25 ii) a detection protein bound to said portion of said receptor protein H4-1BB such that ligand binding can be detected by relative activity assays for said detection protein;
- 30 b) washing said cell of any fusion protein not bound to said cell membrane ligands; and
- c) placing said washed cells in the presence of a substrate for said detection protein and measuring relative activity of said detection protein.

17. The method of claim 16 wherein said detection protein is alkaline phosphatase.



18. A method of inducing B-cell proliferation comprising the step of treating B-cells that have expressed a ligand to human receptor protein H4-1BB with cells that have expressed receptor protein H4-1BB.
- 5 19. A recombinant expression vector comprising a DNA sequence according to claim 1 operably linked to regulatory sequences suitable for expression of said DNA sequence in a host cell.
- 10 20. A recombinant expression vector comprising a DNA sequence according to claim 3 operably linked to regulatory sequences suitable for expression of said DNA sequence in a host cell.
- 15 21. A recombinant expression vector comprising a DNA sequence according to claim 4 operably linked to regulatory sequences suitable for expression of said DNA sequence in a host cell.



22. A purified human 4-lBB polypeptide comprising the N-terminal amino acid sequence Leu-Gln-Asp-Pro-Cys-Ser-Asn-Cys-Pro-Ala-Gly-Thr-.
23. A purified 4-lBB according to claim 22, comprising an amino acid sequence selected from the group consisting of amino acids 24- 255 of SEQ ID NO:1 and amino acids 24-186 of SEQ ID NO:1.
24. A purified 4-lBB according to claim 22, comprising an amino acid sequence that is identical to a sequence selected from the group consisting of amino acids 24-255 of SEQ ID NO:1 and amino acids 24-186 of SEQ ID NO:1, except for conservative amino acid substitution(s).
25. A purified soluble human 4-lBB polypeptide, wherein said polypeptide comprises the extracellular domain of human 4-lBB (amino acids 24-186 of SEQ ID NO:1) or a fragment thereof capable of binding a 4-lBB ligand.
26. A purified soluble human 4-lBB polypeptide of claim 25, additionally comprising a polypeptide fused to the C-terminus of said human 4-lBB polypeptide.
27. A pharmaceutical composition comprising an effective amount of a soluble human 4-lBB of claim 22 in admixture with a suitable diluent, carrier, or excipient.
28. An antibody that is immunoreactive with a human 4-lBB polypeptide of claim 22.

Dated this thirtieth day of July 1998

INDIANA UNIVERSITY FOUNDATION

By their Patent Attorneys

CULLEN & CO.



Fig. 1

ATGTCCATGA	ACTGCTGAGT	GGATAAACAG	CACGGGATAT	CTCTGTCTAA	- 96
AGGAATATTA	CTACACCAGG	AAAAGGACAC	ATTCGACAAC	AGGAAAGGAG	- 46
CCTGTCACAG	AAAACCACAG	TGTCCTGTGC	ATGTGACATT	TCGCC	- 1
ATG GGA AAC AAC TGT TAC AAC GTG GTG GTC ATT GTG CTG CTG CTA	45				
Met Gly Asn Asn Cys Tyr Asn Val Val Val Ile Val Leu Leu Leu					
GTG GGC TGT GAG AAG GTG GGA GCC GTG CAG AAC TCC TGT GAT AAC	90				
Val Gly Cys Glu Lys Val Gly Ala Val Glu Asn Ser Cys Asp Asn					
TGT CAG CCT GGT ACT TTC TGC AGA AAA TAC AAT CCA GTC TGC AAG	135				
Cys Glu Pro Gly Thr Phe Cys Arg Lys Tyr Asn Pro Val Cys Lys					
● H4-1BB FI ●					
AGC TGC CCT CCA AGT ACC TTC TCC AGC ATA GGT GGA CAG CCG AAC	180				
Ser Cys Pro Pro Ser Thr Phe Ser Ser Ile Gly Glu Glu Pro Asn					
H4-1BB FI					
TGT AAC ATC TGC AGA GTG TGT GCA GGC TAT TTC AGG TTC AAG AAG	225				
Cys Asn Ile Cys Arg Val Cys Ala Gly Tyr Phe Arg Phe Lys Lys					
TTT TGC TCC TCT ACC CAC AAC GCG GAG TGT GAG TGC ATT GAA GGA	270				
Phe Cys Ser Ser Thr His Asn Ala Glu Cys Glu Cys Ile Glu Gly					
TTC CAT TGC TTG GGG CCA CAG TGC ACC AGA TGT GAA AAG GAC TGC	315				
Phe His Cys Leu Gly Pro Glu Cys Thr Arg Cys Glu Lys Asp Cys					
AGG CCT GGC CAG GAG CTA ACG AAG CAG GGT TGC AAA ACC TGT AGC	360				
Arg Pro Gly Glu Glu Leu Thr Lys Glu Gly Cys Lys Thr Cys Ser					
● H4-1BB RI ●					
TTG GGA ACA TTT AAT GAC CAG AAC GGT ACT GGC GTC TGT CGA CCC	405				
Leu Gly Thr Phe Asn Asp Glu Asn Gly Thr Glu Val Cys Arg Pro					
H4-1BB RI					
TGG ACG AAC TGC TCT CTA GAC GGA AGG TCT GTG CTT AAG ACC GGG	450				
Trp Thr Asn Cys Ser Leu Asp Gly Arg Ser Val Leu Lys Thr Gly					
ACC ACG GAG AAG GAC GTG GTG TGT GGA CCC CCT GTG GTG AGC TTC	495				
Thr Thr Glu Lys Asp Val Val Cys Gly Pro Pro Val Val Ser Phe					
TCT CCC AGT ACC ACC ATT TCT GTG ACT CCA GAG GGA GGA CCA GGA	540				
Ser Pro Ser Thr Thr Ile Ser Val Thr Pro Glu Gly Gly Pro Gly					
GGG CAC TCC TTG CAG GTC CTT ACC TTG TTC CTG GCG CTG ACA TCG	585				
Gly His Ser Leu Glu Val Leu Thr Leu Phe Leu Ala Leu Thr Ser					
GCT TTG CTG CTG GCC CTG ATC TTC ATT ACT CTC CTG TTC TCT GTG	630				
Ala Leu Leu Leu Ala Leu Ile Phe Ile Thr Leu Leu Phe Ser Val					
CTC AAA TGG ATC AGG AAA AAA TTC CCC CAC ATA TTC AAG CAA CCA	675				
Leu Lys Trp Ile Arg Lys Lys Phe Pro His Ile Phe Lys Glu Pro					
TTT AAG AAG ACC ACT GGA GCA GCT CAA GAG GAA GAT GCT TGT AGC	720				
Phe Lys Lys Thr Thr Gly Ala Ala Glu Glu Glu Asp Ala Cys Ser					

Fig.1 (cont'd)

TGC CGA TGT CCA CAG GAA GAA GAA GGA GGA GGA GGA GGC TAT GAG 765
 Cys Arg Cys Pro Gl n Gl u Gl u Gl u Gl y Gl y Gl y Gl y Gl y Tyr Gl u

CTG TGA
 Leu ---

771

TGTA	CTATCC	TAGGAGATGT	GTGGGCCCGAA	ACCGAGAAGC	ACTAGGACCC	821
CACCATCCTG	TGGAACAGCA	CAAGCAACCC	CACCACCCTG	TTCTTACACA	871	
TCATCCTAGA	TGATGTGTGG	GCGCGCACCT	CATCCAAGTC	TCTTCTAACG	921	
CTAACATATT	TGTCTTTACC	TTTTTTAAAT	CTTTTTTTAA	ATTTAAATTT	971	
TATGTGTGTG	AGTGTTTTGC	CTGCCTGTAT	GCACACGTGT	GTGTGTGTGT	1021	
GTGTGTGACA	CTCCTGATGC	CTGAGGAGGT	CAGAAGACAA	AGGGTTGGTT	1071	
CCATAAGAAC	TGGAGTTATG	GATGGCTGTG	AGCCGGNNNG	ATAGGTCGGG	1121	
ACGGAGACCT	GTCTTCTTAT	TTTAACGTGA	CTGTATAATA	AAAAAAAAT	1171	
GATATTTTCGG	GAATTGTAGA	GATTGTCCTG	ACACCCTTCT	AGTTAATGAT	1221	
CTAAGAGGAA	TTGTTGATAC	GTAGTATACT	GTATATGTGT	ATGTATATGT	1271	
ATATGTATAT	ATAAGACTCT	TTTACTGTCA	AAGTCAACCT	AGAGTGTCTG	1321	
GTTACCAGGT	CAATTTTATT	GGACATTTTA	CGTCACACAC	ACACACACAC	1371	
ACACACACAC	ACGTTTATAC	TACGTACTGT	TATCGGTATT	CTACGTCATA	1421	
TAATGGGATA	GGGTAAAAGG	AAACCAAAGA	GTGAGTGATA	TTATTGTGGA	1471	
GGTGACAGAC	TACCCCTTCT	GGGTACGTAG	GGACAGACCT	CCTTCGGA	1521	
GTCTAAAAC	CCCCTTAGAA	GTCTCGTCAA	GTTCCCGGAC	GAAGAGGACA	1571	
GAGGAGACAC	AGTCCGAAAA	GTTATTTTTT	CGGCAAATCC	TTTCCCTGTT	1621	
TCGTGACACT	CCACCCCTTG	TGGACACTTG	AGTGTATCC	TTGCGCCGGA	1671	
AGGTCAGGTG	GTACCCGTCT	GTAGGGGCGG	GGAGACAGAG	CCGCGGGGGA	1721	
GCTACGAGAA	TCGACTCACA	GGGCGCCCCG	GGCTTCGCAA	ATGAAACTTT	1771	
TTTAATCTCA	CAAGTTTCGT	CCGGGCTCGG	CGGACCTATG	GCGTCGATCC	1821	
TTATTACCTT	ATCCTGGCGC	CAAGATAAAA	CAACCAAAAG	CCTTGACTCC	1871	
GGTACTAATT	CTCCCTGCCG	GCCCCCGTAA	GCATAACGCG	GCGATCTCCA	1921	
CTTTAAGAAC	CTGGCCGCGT	TCTGCCTGGT	CTCGCTTTTC	TAAACGGTTC	1971	
TTACAAAAGT	AATTAGTTCT	TGCTTTCAGC	CTCCAAGCTT	CTGCTAGTCT	2021	
ATGGCAGCAT	CAAGGCTGGT	ATTTGCTACG	GCTGACCGCT	ACGCCGCCGC	2071	
AATAAGGGTA	CTGGGCGGCC	CGTCGAAGGC	CCTTTGGTTT	CAGAAACCCA	2121	
AGGCCCCCT	CATACCAACG	TTTCGACTTT	GATTCTTGCC	GGTACGTGGT	2171	
GGTGGGTGCC	TTAGCTCTTT	CTCGATAGTT	AGAC		2205	

Fig. 2a

human homologue of mouse 4-1bb
h4-1bb Length 838

```

1  AATCAGCTTT GCTAGTATCA TACCTGTGCC AGATTTTCATC ATGGGAAACA
51 GCTGTTACAA CATAGTAGCC ACTCTGTTGC TGGTCCTCAA CTTTGAGAGG
101 ACAAGATCAT TGCAGGATCC TTGTAGTAAC TGCCCAGCTG GTACATTCTG
151 TGATAATAAC AGGAATCAGA TTTGCAGTCC CTGTCCTCCA AATAGTTTCT
201 CCAGCGCAGG TGGACAAAGG ACCTGTGACA TATGCAGGCA GTGTAAAGGT
251 GTTTTTCAGGA CCAGGAAGGA GTGTTCTCC ACCAGCAATG CAGAGTGTGA
301 CTGCACTCCA GGGTTTCACT GCCTGGGGGC AGGATGCAGC ATGTGTGAAC
351 AGGATTGTAA ACAAGGTCAA GAACTGACAA AAAAAGGTTG TAAAGACTGT
401 TGCTTTGGGA CATTTAACGA TCAGAAACGT GGCATCTGTC GACCCTGGAC
451 AAAGTGTCTT TTGGATGGAA AGTCTGTGCT TGTGAATGGG ACGAAGGAGA
501 GGGACGTGGT CTGTGGACCA TCTCCAGCTG ACCTCTCTCC GGGAGCATCC
551 TCTGTGACCC CGCCTGCCCC TGCGAGAGAG CCAGGACACT CTCCGCAGAT
601 CATCTCCTTC TTTCTTGCGC TGACGTCGAC TGCCTTGCTC TTCTTGCTGT
651 TCTTCCTCAC GCTCCGTTTC TCTGTTGTTA AACGGGGCAG AAAGAACTC
701 CTGTATATAT TCAAACAACC ATTTATGAGA CCAGTACAAA CTACTCAAGA
751 GGAAGATGGC TGTAGCTGCC GATTTCCAGA AGAAGAAGAA GGAGGATGTG
801 AACTGTGAAA TGGAAGTCAA TAGGGCTGTT GGGACTTT


```

Fig. 2b

```


1  MGNSCYNIVA TLLLVLNFER TRSLQDPCSN CPAGTFCDNN RNQICSPCPP
51 NSFSSAGGQR TCDICRQCKG VFRTRKECSS TSNAECDCTP GFHCLGAGCS
101 MCEQDCKQGQ ELTKKGCKDC CFGTFNDQKR GICRPWTNCS LDGKSVLVNG
151 TKERDVVCGP SPADLSPGAS SVTPPAPARE PGHSPQIISF FLALTSTALL
201 FLLFFLTLRF SVVKRGRKKL LYIFKQPFMR PVQTTQEEDG CSCRFPEEEE
251 GGCEL

```



COGNITIVE PHASE
early activation




Fig. 3a






PROLIFERATION
CLONAL EXPANSION
late activation

Fig. 3b

NORMAL T-CELL ACTIVATION PATHWAY

*Fig. 4a**Fig. 4b**Fig. 4c*

BLOCKING STEPS IN T-CELL ACTIVATION PATHWAY

*Fig. 5a**Fig. 5b**Fig. 5c*

INTERNATIONAL SEARCH REPORT

Inter. nal Application No
PCT/US 94/10457

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C12N15/62 C07K14/71 C07K14/715 C07K16/28
G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF IMMUNOLOGY., vol.150, no.3, 1 February 1993, BALTIMORE US pages 771 - 781 POLLOK KE;KIM YJ;ZHOU Z;HURTADO J;KIM KK;PICKARD RT;KWON BS; 'Inducible T cell antigen 4-1BB. Analysis of expression and function.' see the whole document ---	1-18
A	JOURNAL OF IMMUNOLOGY., vol.151, no.3, August 1993, BALTIMORE US pages 1255 - 1262 KIM YJ;POLLOK KE;ZHOU Z;SHAW A;BOHLEN JB;FRASER M;KWON BS; 'Novel T cell antigen 4-1BB associates with the protein tyrosine kinase p56lck1.' see the whole document ---	1-18
-/--		

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

20 January 1995

Date of mailing of the international search report

15 -02- 1995

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Nauche, S

INTERNATIONAL SEARCH REPORT

Inter. nal Application No
PCT/US 94/10457

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA., vol.86, March 1989, WASHINGTON US pages 1963 - 1967 KWON, B.S. ET AL.; 'cDNA sequences of two inducible T-cell genes' see the whole document ---	1-18
P,X	GENE., vol.134, 8 December 1993, AMSTERDAM NL pages 295 - 298 SCHWARZ H;TUCKWELL J;LOTZ M; 'A receptor induced by lymphocyte activation (ILA): a new member of the human nerve-growth-factor/tumor-necrosis-factor receptor family.' see the whole document -----	1-18