




(86) **Date de dépôt PCT/PCT Filing Date:** 2008/10/17  
(87) **Date publication PCT/PCT Publication Date:** 2009/04/23  
(45) **Date de délivrance/Issue Date:** 2016/05/03  
(85) **Entrée phase nationale/National Entry:** 2010/04/15  
(86) **N° demande PCT/PCT Application No.:** US 2008/011911  
(87) **N° publication PCT/PCT Publication No.:** 2009/051822  
(30) **Priorités/Priorities:** 2007/10/19 (US60/981,432);  
2008/05/09 (US61/052,002)

(51) **Cl.Int./Int.Cl. C07D 401/12** (2006.01),  
**A61K 31/505** (2006.01), **A61K 31/506** (2006.01),  
**A61P 25/28** (2006.01), **A61P 29/00** (2006.01),  
**A61P 35/00** (2006.01), **A61P 37/00** (2006.01),  
**C07D 239/48** (2006.01), **C07D 403/04** (2006.01),  
**C07D 403/12** (2006.01), **C07D 405/14** (2006.01),  
**C07D 413/12** (2006.01), **C07D 417/14** (2006.01),  
**C07D 471/04** (2006.01)

(72) **Inventeurs/Inventors:**  
TESTER, RICHLAND W., US;  
SINGH, JUSWINDER, US;  
GHOSH, SHOMIR, US;  
KLUGE, ARTHUR F., US; ...

(54) **Titre : COMPOSES HETEROARYLES ET LEURS UTILISATIONS**  
(54) **Title: HETEROARYL COMPOUNDS AND USES THEREOF**



(57) **Abrégé/Abstract:**

The present invention provides compounds, pharmaceutically acceptable compositions thereof, and methods of using the same.



(72) **Inventeurs(suite)/Inventors(continued):** PETTER, RUSSELL C., US

(73) **Propriétaires(suite)/Owners(continued):** CELGENE AVILOMICS RESEARCH, INC., US

(74) **Agent:** TORYS LLP

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 April 2009 (23.04.2009)

PCT

(10) International Publication Number  
**WO 2009/051822 A1**

(51) International Patent Classification:  
*A01N 43/54* (2006.01)

(21) International Application Number:

PCT/US2008/011911

(22) International Filing Date: 17 October 2008 (17.10.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/981,432 19 October 2007 (19.10.2007) US  
61/052,002 9 May 2008 (09.05.2008) US

(71) Applicant (for all designated States except US): **AVILA THERAPEUTICS, INC.** [US/US]; 100 Beaver Street, First Floor, Waltham, MA 02453-8425 (US).

(71) Applicant and

(72) Inventor: **TESTER, Richland, W.** [US/US]; 23 Gregoire Drive, Marlborough, MA 01752 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SINGH, Juswinder** [GB/US]; 94 Heritage Ave., Ashland, MA 01721 (US). **GHOSH, Shomir** [US/US]; 134 Sewall Avenue, Unit C, Brookline, MA 02446 (US). **KLUGE, Arthur, F.** [US/US]; 111 Old County Road, Lincoln, MA 01773 (US). **PETTER, Russell, C.** [US/US]; 22 Robinwood Lane, Stow, MA 01775 (US).

(74) Agent: **ROBIDOUX, Andrea, L.C.**; Choate, Hall & Stewart LLP, Two International Place, Boston, MA 02110 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(54) Title: HETEROARYL COMPOUNDS AND USES THEREOF




Figure 1

(57) Abstract: The present invention provides compounds, pharmaceutically acceptable compositions thereof, and methods of using the same.

WO 2009/051822 A1

**HETEROARYL COMPOUNDS AND USES THEREOF**

[0001]

**TECHNICAL FIELD OF THE INVENTION**

[0002] The present invention relates to compounds useful as inhibitors of protein kinases. The invention also provides pharmaceutically acceptable compositions comprising compounds of the present invention and methods of using said compositions in the treatment of various disorders.

**BACKGROUND OF THE INVENTION**

[0003] The search for new therapeutic agents has been greatly aided in recent years by a better understanding of the structure of enzymes and other biomolecules associated with diseases. One important class of enzymes that has been the subject of extensive study is protein kinases.

[0004] Protein kinases constitute a large family of structurally related enzymes that are responsible for the control of a variety of signal transduction processes within the cell. Protein kinases are thought to have evolved from a common ancestral gene due to the conservation of their structure and catalytic function. Almost all kinases contain a similar 250-300 amino acid catalytic domain. The kinases may be categorized into families by the substrates they phosphorylate (e.g., protein-tyrosine, protein-serine/threonine, lipids, etc.).


[0005] In general, protein kinases mediate intracellular signaling by effecting a phosphoryl transfer from a nucleoside triphosphate to a protein acceptor that is involved in a signaling pathway. These phosphorylation events act as molecular on/off switches that can modulate or regulate the target protein biological function. These phosphorylation events are ultimately triggered in response to a variety of extracellular and other stimuli. Examples of such stimuli include environmental and chemical stress signals (e.g., osmotic shock, heat shock, ultraviolet radiation, bacterial endotoxin, and  $H_2O_2$ ), cytokines (e.g., interleukin-1 (IL-1) and tumor necrosis factor  $\alpha$  (TNF- $\alpha$ )), and growth factors (e.g., granulocyte macrophage-colony-stimulating factor

(GM-CSF), and fibroblast growth factor (FGF)). An extracellular stimulus may affect one or more cellular responses related to cell growth, migration, differentiation, secretion of hormones, activation of transcription factors, muscle contraction, glucose metabolism, control of protein synthesis, and regulation of the cell cycle.

[0006] Many diseases are associated with abnormal cellular responses triggered by protein kinase-mediated events as described above. These diseases include, but are not limited to, autoimmune diseases, inflammatory diseases, bone diseases, metabolic diseases, neurological and neurodegenerative diseases, cancer, cardiovascular diseases, allergies and asthma, Alzheimer's disease, and hormone-related diseases. Accordingly, there remains a need to find protein kinase inhibitors useful as therapeutic agents.

#### SUMMARY OF THE INVENTION

[0007] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are effective as inhibitors of one or more protein kinases. Such compounds have the general formula I:



or a pharmaceutically acceptable salt thereof, wherein Ring A, Ring B, m,  $R^x$ ,  $R^y$ , W, and  $R^1$  are as defined herein.

[0008] Compounds of the present invention, and pharmaceutically acceptable compositions thereof, are useful for treating a variety of diseases, disorders or conditions, associated with abnormal cellular responses triggered by protein kinase-mediated events. Such diseases, disorders, or conditions include those described herein.

[0009] Compounds provided by this invention are also useful for the study of kinases in biological and pathological phenomena; the study of intracellular signal transduction pathways mediated by such kinases; and the comparative evaluation of new kinase inhibitors.

### BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1** depicts the EGF inhibiting activity of compound **I-1**.

**Figure 2** depicts the results of compound **I-1** in a “washout” experiment as compared with compound **I-93**.

**Figure 3** depicts dose response inhibition of EGFR phosphorylation and p42/p44 Erk phosphorylation with compounds **I-16** and **I-17** in A431 cells.

**Figure 4** depicts dose response inhibition of EGFR phosphorylation and p42/p44 Erk phosphorylation with compound **I-19** in A431 cells.

**Figure 5** depicts dose response inhibition of EGFR phosphorylation with compound **I-1** in A431 cells as compared with its “reversible control” compound (**I<sup>R</sup>-3**).

**Figure 6** depicts the results of compound **I-1** in a “washout” experiment as compared with its “reversible control” compound (**I<sup>R</sup>-3**).

**Figure 7** depicts MS analysis confirming covalent modification of ErbB4 by compound **I-1**.

**Figure 8** depicts MS analysis confirming covalent modification of ErbB1 at Cys797 by compound **I-1**.

**Figure 9** depicts the inhibition of BTK signaling in Ramos cells by compound **I-13**.

**Figure 10** depicts the results of compound **I-13** in a “washout” experiment with BTK in Ramos cells.

**Figure 11** depicts MS analysis of the tryptic digests confirming covalent modification of TEC kinase by compound **I-13**.

**Figure 12** depicts MS analysis confirming covalent modification of BTK by compound **I-63**.

**Figure 13** depicts MS analysis confirming covalent modification of BTK by compound **I-66**.

**Figure 14** depicts an amino acid sequence for BTK (SEQ ID 1).

**Figure 15** depicts an amino acid sequence for TEC (SEQ ID 2).

**Figure 16** depicts an amino acid sequence for ITK (SEQ ID 3).

**Figure 17** depicts an amino acid sequence for BMX (SEQ ID 4).


**Figure 18** depicts an amino acid sequence for JAK3 (SEQ ID 5).

**Figure 19** depicts an amino acid sequence for TXK (SEQ ID 6).

# DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

## 1. General Description of Compounds of the Invention

[0010] In certain embodiments, the present invention provides a compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted group selected from phenyl, an 8-10 membered bicyclic partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Ring B is phenyl, a 5-6 membered heteroaryl ring having 1-3 heteratoms independently selected from N, O or S, a 5-6 membered saturated heterocyclic ring having 1-2 heteroatoms independently selected from N, O or S, or an 8-10 membered bicyclic partially unsaturated or aryl ring having 1-3 heteroatoms independently selected from N, O or S;

R<sup>1</sup> is a warhead group;

R<sup>y</sup> is hydrogen, halogen, CN, lower alkyl, or lower haloalkyl;

W is a bivalent C<sub>1-3</sub> alkylene chain wherein one methylene unit of W is optionally replaced by – NR<sup>2</sup>–, –N(R<sup>2</sup>)C(O)–, –C(O)N(R<sup>2</sup>)–, –N(R<sup>2</sup>)SO<sub>2</sub>–, –SO<sub>2</sub>N(R<sup>2</sup>)–, –O–, –C(O)–, –OC(O)–, –C(O)O–, –S–, –SO– or –SO<sub>2</sub>–;

R<sup>2</sup> is hydrogen or optionally substituted C<sub>1-6</sub> aliphatic, or:

R<sup>2</sup> and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated ring, or:

R<sup>2</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a 4-7 membered carbocyclic ring;


m is 0-4;

each  $R^x$  is independently selected from -R, halogen, -OR, -CN, -NO<sub>2</sub>, -SO<sub>2</sub>R, -SOR, -C(O)R, -CO<sub>2</sub>R, -C(O)N(R)<sub>2</sub>, -NRC(O)R, -NRC(O)NR<sub>2</sub>, -NRSO<sub>2</sub>R, or -N(R)<sub>2</sub>; or:

$R^x$  and  $R^1$  are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, CN, or C<sub>1-6</sub> aliphatic; and

each R group is independently hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0011] In certain embodiments, the present invention provides a compound of formula II:



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted group selected from phenyl, an 8-10 membered bicyclic partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

$R^1$  is a warhead group;

$R^y$  is hydrogen, halogen, CN, lower alkyl, or lower haloalkyl;

G is CH, or N;

W is -NR<sup>2</sup>-, -S-, or -O-;

$R^2$  is hydrogen or optionally substituted C<sub>1-6</sub> aliphatic, or:

$R^2$  and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated ring;

m is 0-4;

each R<sup>x</sup> is independently selected from -R, halogen, -OR, -CN, -NO<sub>2</sub>, -SO<sub>2</sub>R, -SOR, -C(O)R, -CO<sub>2</sub>R, -C(O)N(R)<sub>2</sub>, -NRC(O)R, -NRC(O)NR<sub>2</sub>, -NRSO<sub>2</sub>R, or -N(R)<sub>2</sub>; or:

R<sup>x</sup> and R<sup>1</sup> are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, CN, or C<sub>1-6</sub> aliphatic; and

each R group is independently hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

## 2. Compounds and Definitions

[0012] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75<sup>th</sup> Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5<sup>th</sup> Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001.

[0013] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-6 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-5 aliphatic carbon atoms. In other embodiments, aliphatic groups

contain 1-4 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-3 aliphatic carbon atoms, and in yet other embodiments, aliphatic groups contain 1-2 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C<sub>3</sub>-C<sub>6</sub> hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.

[0014] The term "lower alkyl" refers to a C<sub>1-4</sub> straight or branched alkyl group. Exemplary lower alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, and tert-butyl.

[0015] The term "lower haloalkyl" refers to a C<sub>1-4</sub> straight or branched alkyl group that is substituted with one or more halogen atoms.

[0016] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR<sup>+</sup> (as in N-substituted pyrrolidinyl)).


[0017] The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation.

[0018] As used herein, the term "bivalent C<sub>1-8</sub> (or C<sub>1-6</sub>) saturated or unsaturated, straight or branched, hydrocarbon chain", refers to bivalent alkylene, alkenylene, and alkynylene chains that are straight or branched as defined herein.

[0019] The term "alkylene" refers to a bivalent alkyl group. An "alkylene chain" is a polymethylene group, i.e., -(CH<sub>2</sub>)<sub>n</sub>-, wherein n is a positive integer, preferably from 1 to 6, from 1 to 4, from 1 to 3, from 1 to 2, or from 2 to 3. A substituted alkylene chain is a polymethylene group in which one or more methylene hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.

[0020] The term "alkenylene" refers to a bivalent alkenyl group. A substituted alkenylene chain is a polymethylene group containing at least one double bond in which one or more hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.

[0021] As used herein, the term “cyclopropylenyl” refers to a bivalent cyclopropyl group of

the following structure: .

[0022] The term “halogen” means F, Cl, Br, or I.

[0023] The term “aryl” used alone or as part of a larger moiety as in “aralkyl”, “aralkoxy”, or “aryloxyalkyl”, refers to monocyclic and bicyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains three to seven ring members. The term “aryl” may be used interchangeably with the term “aryl ring”. In certain embodiments of the present invention, “aryl” refers to an aromatic ring system which includes, but not limited to, phenyl, biphenyl, naphthyl, anthracyl and the like, which may bear one or more substituents. Also included within the scope of the term “aryl”, as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings, such as indanyl, phthalimidyl, naphthimidyl, phenanthridinyl, or tetrahydronaphthyl, and the like.

[0024] The terms “heteroaryl” and “heteroar-”, used alone or as part of a larger moiety, e.g., “heteroaralkyl”, or “heteroaralkoxy”, refer to groups having 5 to 10 ring atoms, preferably 5, 6, or 9 ring atoms; having 6, 10, or 14  $\pi$  electrons shared in a cyclic array; and having, in addition to carbon atoms, from one to five heteroatoms. The term “heteroatom” refers to nitrogen, oxygen, or sulfur, and includes any oxidized form of nitrogen or sulfur, and any quaternized form of a basic nitrogen. Heteroaryl groups include, without limitation, thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolizinyl, purinyl, naphthyridinyl, and pteridinyl. The terms “heteroaryl” and “heteroar-”, as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl, cycloaliphatic, or heterocyclyl rings, where the radical or point of attachment is on the heteroaromatic ring. Nonlimiting examples include indolyl, isoindolyl, benzothienyl, benzofuranyl, dibenzofuranyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, isoquinolyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, 4*H*-quinolizinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, tetrahydroquinolyl, tetrahydroisoquinolyl, and pyrido[2,3-*b*]-1,4-oxazin-3(4*H*)-one. A heteroaryl group may be mono- or bicyclic. The term “heteroaryl” may be used interchangeably with the terms “heteroaryl ring”, “heteroaryl group”, or “heteroaromatic”, any of

which terms include rings that are optionally substituted. The term “heteroaralkyl” refers to an alkyl group substituted by a heteroaryl, wherein the alkyl and heteroaryl portions independently are optionally substituted.

[0025] As used herein, the terms “heterocycle”, “heterocyclyl”, “heterocyclic radical”, and “heterocyclic ring” are used interchangeably and refer to a stable 5- to 7-membered monocyclic or 7-10-membered bicyclic heterocyclic moiety that is either saturated or partially unsaturated, and having, in addition to carbon atoms, one or more, preferably one to four, heteroatoms, as defined above. When used in reference to a ring atom of a heterocycle, the term “nitrogen” includes a substituted nitrogen. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl), or <sup>+</sup>NR (as in *N*-substituted pyrrolidinyl).

[0026] A heterocyclic ring can be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure and any of the ring atoms can be optionally substituted. Examples of such saturated or partially unsaturated heterocyclic radicals include, without limitation, tetrahydrofuranyl, tetrahydrothiophenyl pyrrolidinyl, piperidinyl, pyrrolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, oxazolidinyl, piperazinyl, dioxanyl, dioxolanyl, diazepinyl, oxazepinyl, thiazepinyl, morpholinyl, and quinuclidinyl. The terms “heterocycle”, “heterocyclyl”, “heterocyclyl ring”, “heterocyclic group”, “heterocyclic moiety”, and “heterocyclic radical”, are used interchangeably herein, and also include groups in which a heterocyclyl ring is fused to one or more aryl, heteroaryl, or cycloaliphatic rings, such as indolinyl, 3*H*-indolyl, chromanyl, phenanthridinyl, or tetrahydroquinolinyl, where the radical or point of attachment is on the heterocyclyl ring. A heterocyclyl group may be mono- or bicyclic. The term “heterocyclylalkyl” refers to an alkyl group substituted by a heterocyclyl, wherein the alkyl and heterocyclyl portions independently are optionally substituted.

[0027] As used herein, the term “partially unsaturated” refers to a ring moiety that includes at least one double or triple bond. The term “partially unsaturated” is intended to encompass rings having multiple sites of unsaturation, but is not intended to include aryl or heteroaryl moieties, as herein defined.

[0028] As described herein, compounds of the invention may contain “optionally substituted” moieties. In general, the term “substituted”, whether preceded by the term “optionally” or not, means that one or more hydrogens of the designated moiety are replaced

with a suitable substituent. Unless otherwise indicated, an “optionally substituted” group may have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term “stable”, as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and use for one or more of the purposes disclosed herein.

[0029] Suitable monovalent substituents on a substitutable carbon atom of an “optionally substituted” group are independently halogen;  $-(CH_2)_{0-4}R^\circ$ ;  $-(CH_2)_{0-4}OR^\circ$ ;  $-O(CH_2)_{0-4}R^\circ$ ;  $-O-(CH_2)_{0-4}C(O)OR^\circ$ ;  $-(CH_2)_{0-4}CH(OR^\circ)_2$ ;  $-(CH_2)_{0-4}SR^\circ$ ;  $-(CH_2)_{0-4}Ph$ , which may be substituted with  $R^\circ$ ;  $-(CH_2)_{0-4}O(CH_2)_{0-1}Ph$  which may be substituted with  $R^\circ$ ;  $-CH=CHPh$ , which may be substituted with  $R^\circ$ ;  $-(CH_2)_{0-4}O(CH_2)_{0-1}$ -pyridyl which may be substituted with  $R^\circ$ ;  $-NO_2$ ;  $-CN$ ;  $-N_3$ ;  $-(CH_2)_{0-4}N(R^\circ)_2$ ;  $-(CH_2)_{0-4}N(R^\circ)C(O)R^\circ$ ;  $-N(R^\circ)C(S)R^\circ$ ;  $-(CH_2)_{0-4}N(R^\circ)C(O)NR^\circ_2$ ;  $-N(R^\circ)C(S)NR^\circ_2$ ;  $-(CH_2)_{0-4}N(R^\circ)C(O)OR^\circ$ ;  $-N(R^\circ)N(R^\circ)C(O)R^\circ$ ;  $-N(R^\circ)N(R^\circ)C(O)NR^\circ_2$ ;  $-N(R^\circ)N(R^\circ)C(O)OR^\circ$ ;  $-(CH_2)_{0-4}C(O)R^\circ$ ;  $-C(S)R^\circ$ ;  $-(CH_2)_{0-4}C(O)OR^\circ$ ;  $-(CH_2)_{0-4}C(O)SR^\circ$ ;  $-(CH_2)_{0-4}C(O)OSiR^\circ_3$ ;  $-(CH_2)_{0-4}OC(O)R^\circ$ ;  $-OC(O)(CH_2)_{0-4}SR-$ ,  $SC(S)SR^\circ$ ;  $-(CH_2)_{0-4}SC(O)R^\circ$ ;  $-(CH_2)_{0-4}C(O)NR^\circ_2$ ;  $-C(S)NR^\circ_2$ ;  $-C(S)SR^\circ$ ;  $-SC(S)SR^\circ$ ;  $-(CH_2)_{0-4}OC(O)NR^\circ_2$ ;  $-C(O)N(OR^\circ)R^\circ$ ;  $-C(O)C(O)R^\circ$ ;  $-C(O)CH_2C(O)R^\circ$ ;  $-C(NOR^\circ)R^\circ$ ;  $-(CH_2)_{0-4}SSR^\circ$ ;  $-(CH_2)_{0-4}S(O)_2R^\circ$ ;  $-(CH_2)_{0-4}S(O)_2OR^\circ$ ;  $-(CH_2)_{0-4}OS(O)_2R^\circ$ ;  $-S(O)_2NR^\circ_2$ ;  $-(CH_2)_{0-4}S(O)R^\circ$ ;  $-N(R^\circ)S(O)_2NR^\circ_2$ ;  $-N(R^\circ)S(O)_2R^\circ$ ;  $-N(OR^\circ)R^\circ$ ;  $-C(NH)NR^\circ_2$ ;  $-P(O)_2R^\circ$ ;  $-P(O)R^\circ_2$ ;  $-OP(O)R^\circ_2$ ;  $-OP(O)(OR^\circ)_2$ ;  $SiR^\circ_3$ ;  $-(C_{1-4}$  straight or branched alkylene) $O-N(R^\circ)_2$ ; or  $-(C_{1-4}$  straight or branched alkylene) $C(O)O-N(R^\circ)_2$ , wherein each  $R^\circ$  may be substituted as defined below and is independently hydrogen,  $C_{1-6}$  aliphatic,  $-CH_2Ph$ ,  $-O(CH_2)_{0-1}Ph$ ,  $-CH_2$ -(5-6 membered heteroaryl ring), or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of  $R^\circ$ , taken together with their intervening atom(s), form a 3-12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0-4

heteroatoms independently selected from nitrogen, oxygen, or sulfur, which may be substituted as defined below.

**[0030]** Suitable monovalent substituents on  $R^\circ$  (or the ring formed by taking two independent occurrences of  $R^\circ$  together with their intervening atoms), are independently halogen,  $-(CH_2)_{0-2}R^\bullet$ ,  $-(haloR^\bullet)$ ,  $-(CH_2)_{0-2}OH$ ,  $-(CH_2)_{0-2}OR^\bullet$ ,  $-(CH_2)_{0-2}CH(OR^\bullet)_2$ ;  $-O(haloR^\bullet)$ ,  $-CN$ ,  $-N_3$ ,  $-(CH_2)_{0-2}C(O)R^\bullet$ ,  $-(CH_2)_{0-2}C(O)OH$ ,  $-(CH_2)_{0-2}C(O)OR^\bullet$ ,  $-(CH_2)_{0-2}SR^\bullet$ ,  $-(CH_2)_{0-2}SH$ ,  $-(CH_2)_{0-2}NH_2$ ,  $-(CH_2)_{0-2}NHR^\bullet$ ,  $-(CH_2)_{0-2}NR^\bullet_2$ ,  $-NO_2$ ,  $-SiR^\bullet_3$ ,  $-OSiR^\bullet_3$ ,  $-C(O)SR^\bullet$ ,  $-(C_{1-4}$  straight or branched alkylene) $C(O)OR^\bullet$ , or  $-SSR^\bullet$  wherein each  $R^\bullet$  is unsubstituted or where preceded by "halo" is substituted only with one or more halogens, and is independently selected from  $C_{1-4}$  aliphatic,  $-CH_2Ph$ ,  $-O(CH_2)_{0-1}Ph$ , or a 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents on a saturated carbon atom of  $R^\circ$  include  $=O$  and  $=S$ .

**[0031]** Suitable divalent substituents on a saturated carbon atom of an "optionally substituted" group include the following:  $=O$ ,  $=S$ ,  $=NNR^\bullet_2$ ,  $=NNHC(O)R^\bullet$ ,  $=NNHC(O)OR^\bullet$ ,  $=NNHS(O)_2R^\bullet$ ,  $=NR^\bullet$ ,  $=NOR^\bullet$ ,  $-O(C(R^\bullet_2))_{2-3}O-$ , or  $-S(C(R^\bullet_2))_{2-3}S-$ , wherein each independent occurrence of  $R^\bullet$  is selected from hydrogen,  $C_{1-6}$  aliphatic which may be substituted as defined below, or an unsubstituted 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Suitable divalent substituents that are bound to vicinal substitutable carbons of an "optionally substituted" group include:  $-O(CR^\bullet_2)_{2-3}O-$ , wherein each independent occurrence of  $R^\bullet$  is selected from hydrogen,  $C_{1-6}$  aliphatic which may be substituted as defined below, or an unsubstituted 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

**[0032]** Suitable substituents on the aliphatic group of  $R^\bullet$  include halogen,  $-R^\bullet$ ,  $-(haloR^\bullet)$ ,  $-OH$ ,  $-OR^\bullet$ ,  $-O(haloR^\bullet)$ ,  $-CN$ ,  $-C(O)OH$ ,  $-C(O)OR^\bullet$ ,  $-NH_2$ ,  $-NHR^\bullet$ ,  $-NR^\bullet_2$ , or  $-NO_2$ , wherein each  $R^\bullet$  is unsubstituted or where preceded by "halo" is substituted only with one or more halogens, and is independently  $C_{1-4}$  aliphatic,  $-CH_2Ph$ ,  $-O(CH_2)_{0-1}Ph$ , or a 5–6–membered saturated, partially unsaturated, or aryl ring having 0–4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0033] Suitable substituents on a substitutable nitrogen of an "optionally substituted" group include  $-R^{\dagger}$ ,  $-NR^{\dagger}_2$ ,  $-C(O)R^{\dagger}$ ,  $-C(O)OR^{\dagger}$ ,  $-C(O)C(O)R^{\dagger}$ ,  $-C(O)CH_2C(O)R^{\dagger}$ ,  $-S(O)_2R^{\dagger}$ ,  $-S(O)_2NR^{\dagger}_2$ ,  $-C(S)NR^{\dagger}_2$ ,  $-C(NH)NR^{\dagger}_2$ , or  $-N(R^{\dagger})S(O)_2R^{\dagger}$ ; wherein each  $R^{\dagger}$  is independently hydrogen,  $C_{1-6}$  aliphatic which may be substituted as defined below, unsubstituted  $-OPh$ , or an unsubstituted 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of  $R^{\dagger}$ , taken together with their intervening atom(s) form an unsubstituted 3-12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0034] Suitable substituents on the aliphatic group of  $R^{\dagger}$  are independently halogen,  $-R^{\bullet}$ ,  $-(haloR^{\bullet})$ ,  $-OH$ ,  $-OR^{\bullet}$ ,  $-O(haloR^{\bullet})$ ,  $-CN$ ,  $-C(O)OH$ ,  $-C(O)OR^{\bullet}$ ,  $-NH_2$ ,  $-NHR^{\bullet}$ ,  $-NR^{\bullet}_2$ , or  $-NO_2$ , wherein each  $R^{\bullet}$  is unsubstituted or where preceded by "halo" is substituted only with one or more halogens, and is independently  $C_{1-4}$  aliphatic,  $-CH_2Ph$ ,  $-O(CH_2)_{0-1}Ph$ , or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0035] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1-19.

Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide,

2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

[0036] Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and  $N^+(C_{1-4}alkyl)_4$  salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0037] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, Z and E double bond isomers, and Z and E conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures including the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a  $^{13}C$ - or  $^{14}C$ -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools, as probes in biological assays, or as therapeutic agents in accordance with the present invention. In some embodiments, the  $R^1$  group of formula I comprises one or more deuterium atoms. Such compounds include compound I-61 as depicted in Table 5.

[0038] As used herein, the term “irreversible” or “irreversible inhibitor” refers to an inhibitor (i.e. a compound) that is able to be covalently bonded to a target protein kinase in a substantially non-reversible manner. That is, whereas a reversible inhibitor is able to bind to (but is generally unable to form a covalent bond) the target protein kinase, and therefore can become dissociated from the target protein kinase, an irreversible inhibitor will remain substantially bound to the

target protein kinase once covalent bond formation has occurred. Irreversible inhibitors usually display *time dependency*, whereby the degree of inhibition increases with the time with which the inhibitor is in contact with the enzyme. Methods for identifying if a compound is acting as an irreversible inhibitor are known to one of ordinary skill in the art. Such methods include, but are not limited to, enzyme kinetic analysis of the inhibition profile of the compound with the protein kinase target, the use of mass spectrometry of the protein drug target modified in the presence of the inhibitor compound, discontinuous exposure, also known as “washout,” experiments, and the use of labeling, such as radiolabelled inhibitor, to show covalent modification of the enzyme, as well as other methods known to one of skill in the art.

[0039] One of ordinary skill in the art will recognize that certain reactive functional groups can act as “warheads.” As used herein, the term “warhead” or “warhead group” refers to a functional group present on a compound of the present invention wherein that functional group is capable of covalently binding to an amino acid residue (such as cysteine, lysine, histidine, or other residues capable of being covalently modified) present in the binding pocket of the target protein, thereby irreversibly inhibiting the protein. It will be appreciated that the -L-Y group, as defined and described herein, provides such warhead groups for covalently, and irreversibly, inhibiting the protein.

[0040] As used herein, the term “inhibitor” is defined as a compound that binds to and /or inhibits the target protein kinase with measurable affinity. In certain embodiments, an inhibitor has an IC<sub>50</sub> and/or binding constant of less about 50  $\mu$ M, less than about 1  $\mu$ M, less than about 500 nM, less than about 100 nM, or less than about 10 nM.

[0041] A compound of the present invention may be tethered to a detectable moiety. One of ordinary skill in the art will recognize that a detectable moiety may be attached to a provided compound *via* a suitable substituent on Ring A or a suitable R<sup>x</sup> substituent. As used herein, the term “suitable substituent” refers to a moiety that is capable of covalent attachment to a detectable moiety. Such moieties are well known to one of ordinary skill in the art and include groups containing, e.g., a carboxylate moiety, an amino moiety, a thiol moiety, or a hydroxyl moiety, to name but a few. It will be appreciated that such moieties may be directly attached to a provided compound (i.e. directly attached to Ring A or the aniline ring) or *via* a tethering group, such as a bivalent saturated or unsaturated hydrocarbon chain.

[0042] As used herein, the term “detectable moiety” is used interchangeably with the term “label” and relates to any moiety capable of being detected, e.g., primary labels and secondary labels. Primary labels, such as radioisotopes (e.g., tritium,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$ , or  $^{14}\text{C}$ ), mass-tags, and fluorescent labels are signal generating reporter groups which can be detected without further modifications.

[0043] The term “secondary label” as used herein refers to moieties such as biotin and various protein antigens that require the presence of a second intermediate for production of a detectable signal. For biotin, the secondary intermediate may include streptavidin-enzyme conjugates. For antigen labels, secondary intermediates may include antibody-enzyme conjugates. Some fluorescent groups act as secondary labels because they transfer energy to another group in the process of nonradiative fluorescent resonance energy transfer (FRET), and the second group produces the detected signal.

[0044] The terms “fluorescent label”, “fluorescent dye”, and “fluorophore” as used herein refer to moieties that absorb light energy at a defined excitation wavelength and emit light energy at a different wavelength. Examples of fluorescent labels include, but are not limited to: Alexa<sup>TM</sup> Fluor dyes (Alexa<sup>TM</sup> Fluor 350, Alexa<sup>TM</sup> Fluor 488, Alexa<sup>TM</sup> Fluor 532, Alexa<sup>TM</sup> Fluor 546, Alexa<sup>TM</sup> Fluor 568, Alexa<sup>TM</sup> Fluor 594, Alexa<sup>TM</sup> Fluor 633, Alexa<sup>TM</sup> Fluor 660 and Alexa<sup>TM</sup> Fluor 680), AMCA, AMCA-S, BODIPY dyes (BODIPY FL, BODIPY R6G, BODIPY TMR, BODIPY TR, BODIPY 530/550, BODIPY 558/568, BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY 630/650, BODIPY 650/665), Carboxyrhodamine 6G, carboxy-X-rhodamine (ROX), Cascade Blue, Cascade Yellow, Coumarin 343, Cyanine dyes (Cy3, Cy5, Cy3.5, Cy5.5), Dansyl, Dapoxyl, Dialkylaminocoumarin, 4',5'-Dichloro-2',7'-dimethoxy-fluorescein, DM-NERF, Eosin, Erythrosin, Fluorescein, FAM, Hydroxycoumarin, IRDyes (IRD40, IRD 700, IRD 800), JOE, Lissamine rhodamine B, Marina Blue, Methoxycoumarin, Naphthofluorescein, Oregon Green 488, Oregon Green 500, Oregon Green 514, Pacific Blue, PyMPO, Pyrene, Rhodamine B, Rhodamine 6G, Rhodamine Green, Rhodamine Red, Rhodol Green, 2',4',5',7'-Tetra-bromosulfone-fluorescein, Tetramethyl-rhodamine (TMR), Carboxytetramethylrhodamine (TAMRA), Texas Red, Texas Red-X.


[0045] The term “mass-tag” as used herein refers to any moiety that is capable of being uniquely detected by virtue of its mass using mass spectrometry (MS) detection techniques. Examples of mass-tags include electrophore release tags such as N-[3-[4'-(p-

Methoxytetrafluorobenzyl)oxy]phenyl]-3-methylglyceronyl]isonipecotic Acid, 4'-[2,3,5,6-Tetrafluoro-4-(pentafluorophenoxy)]methyl acetophenone, and their derivatives. The synthesis and utility of these mass-tags is described in United States Patents 4,650,750, 4,709,016, 5,360,8191, 5,516,931, 5,602,273, 5,604,104, 5,610,020, and 5,650,270. Other examples of mass-tags include, but are not limited to, nucleotides, dideoxynucleotides, oligonucleotides of varying length and base composition, oligopeptides, oligosaccharides, and other synthetic polymers of varying length and monomer composition. A large variety of organic molecules, both neutral and charged (biomolecules or synthetic compounds) of an appropriate mass range (100-2000 Daltons) may also be used as mass-tags.

[0046] The terms “measurable affinity” and “measurably inhibit,” as used herein, means a measurable change in at least one of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3 activity between a sample comprising a compound of the present invention, or composition thereof, and at least one of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, and an equivalent sample comprising at least one of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, in the absence of said compound, or composition thereof.

### 3. Description of Exemplary Compounds

[0047] According to one aspect, the present invention provides a compound of formula I,



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted group selected from phenyl, an 8-10 membered bicyclic partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Ring B is phenyl, a 5-6 membered heteroaryl ring having 1-3 heteratoms independently selected from N, O or S, a 5-6 membered saturated heterocyclic ring having 1-2 heteratoms independently selected from N, O or S, or an 8-10 membered bicyclic partially unsaturated or aryl ring having 1-3 heteratoms independently selected from N, O or S;

R<sup>1</sup> is -L-Y, wherein:

L is a covalent bond or a bivalent C<sub>1-8</sub> saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one, two, or three methylene units of L are optionally and independently replaced by cyclopropylene, -NR-, -N(R)C(O)-, -C(O)N(R)-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -S-, -SO-, -SO<sub>2</sub>-, -C(=S)-, -C(=NR)-, -N=N-, or -C(=N<sub>2</sub>)-;

Y is hydrogen, C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteratoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with 1-4 R<sup>e</sup> groups; and

each R<sup>e</sup> is independently selected from -Q-Z, oxo, NO<sub>2</sub>, halogen, CN, a suitable leaving group, or a C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, wherein:

Q is a covalent bond or a bivalent C<sub>1-6</sub> saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by -N(R)-, -S-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -SO-, or -SO<sub>2</sub>-, -N(R)C(O)-, -C(O)N(R)-, -N(R)SO<sub>2</sub>-, or -SO<sub>2</sub>N(R)-; and

Z is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN;

R<sup>y</sup> is hydrogen, halogen, CN, lower alkyl, or lower haloalkyl;

W is a bivalent C<sub>1-3</sub> alkylene chain wherein one methylene unit of W is optionally replaced by -NR<sup>2</sup>-, -N(R<sup>2</sup>)C(O)-, -C(O)N(R<sup>2</sup>)-, -N(R<sup>2</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>2</sup>)-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -S-, -SO- or -SO<sub>2</sub>-;

R<sup>2</sup> is hydrogen or optionally substituted C<sub>1-6</sub> aliphatic, or:

R<sup>2</sup> and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated ring, or:

R<sup>2</sup> and R<sup>y</sup> are taken together with their intervening atoms to form a 4-7 membered carbocyclic ring;


m is 0-4;

each  $R^x$  is independently selected from -R, halogen, -OR, -CN, -NO<sub>2</sub>, -SO<sub>2</sub>R, -SOR, -C(O)R, -CO<sub>2</sub>R, -C(O)N(R)<sub>2</sub>, -NRC(O)R, -NRC(O)NR<sub>2</sub>, -NRSO<sub>2</sub>R, or -N(R)<sub>2</sub>; or:

$R^x$  and  $R^1$  are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, CN, or C<sub>1-6</sub> aliphatic; and

each R group is independently hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0048] In certain embodiments, the present invention provides a compound of formula II:



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is an optionally substituted group selected from phenyl, an 8-10 membered bicyclic partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

$R^1$  is -L-Y, wherein:

L is a covalent bond or a bivalent C<sub>1-8</sub> saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one, two, or three methylene units of L are optionally and independently replaced by cyclopropylene, -NR-, -N(R)C(O)-, -C(O)N(R)-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -S-, -SO-, -SO<sub>2</sub>-, -C(=S)-, -C(=NR)-, -N=N-, or -C(=N<sub>2</sub>)-

Y is hydrogen, C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with 1-4 R<sup>e</sup> groups; and

each R<sup>e</sup> is independently selected from -Q-Z, oxo, NO<sub>2</sub>, halogen, CN, a suitable leaving group, or a C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, wherein:

Q is a covalent bond or a bivalent C<sub>1-6</sub> saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by -N(R)-, -S-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -SO-, or -SO<sub>2</sub>-, -N(R)C(O)-, -C(O)N(R)-, -N(R)SO<sub>2</sub>-, or -SO<sub>2</sub>N(R)-; and

Z is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN;

R<sup>y</sup> is hydrogen, halogen, CN, lower alkyl, or lower haloalkyl;

G is CH, or N;

W is -NR<sup>2</sup>-, -S-, or -O-;

R<sup>2</sup> is hydrogen or optionally substituted C<sub>1-6</sub> aliphatic, or:

R<sup>2</sup> and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated ring;


m is 0-4;

each R<sup>x</sup> is independently selected from -R, halogen, -OR, -CN, -NO<sub>2</sub>, -SO<sub>2</sub>R, -SOR, -C(O)R, -CO<sub>2</sub>R, -C(O)N(R)<sub>2</sub>, -NRC(O)R, -NRC(O)NR<sub>2</sub>, -NRSO<sub>2</sub>R, or -N(R)<sub>2</sub>; or:

R<sup>x</sup> and R<sup>1</sup> are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, CN, or C<sub>1-6</sub> aliphatic; and

each R group is independently hydrogen or an optionally substituted group selected from C<sub>1-6</sub> aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

**[0049]** According to one aspect, the present invention provides a compound of formula **II-a** or **II-b**:



or a pharmaceutically acceptable salt thereof, wherein each of Ring A, W, R<sup>1</sup>, G, R<sup>y</sup>, R<sup>x</sup> and m are as defined above for formula II and as described herein.

**[0050]** In certain embodiments, the present invention provides a compound of formula **II-b** where in said compound is other than N<sup>6</sup>-m-tolyl-N<sup>4</sup>-p-tolylpyrimidine-4,6-diamine.

**[0051]** In certain embodiments, the present invention provides a compound of formula **II-a** wherein said compound is other than N<sup>4</sup>-(3-aminophenyl)-N<sup>6</sup>-(3-bromophenyl)pyrimidine-4,6-diamine, N-(3-(6-(3-(trifluoromethyl)phenylamino)pyrimidin-4-ylamino)phenyl)cyclopropanecarboxamide, N-(3-(6-(3-bromophenylamino)pyrimidin-4-ylamino)phenyl)propionamide, N<sup>4</sup>-(3-aminophenyl)-N<sup>6</sup>-m-tolylpyrimidine-4,6-diamine, or N<sup>4</sup>-(3-aminophenyl)-N<sup>6</sup>-methyl-N<sup>6</sup>-phenylpyrimidine-4,6-diamine.


**[0052]** As defined generally above, the Ring A group of formulae **I** and **II** is an optionally substituted group selected from phenyl, an 8-10 membered bicyclic partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In certain embodiments, Ring A is an optionally substituted phenyl group. In some embodiments, Ring A is an optionally substituted naphthyl ring or a bicyclic 8-10 membered heteraryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. In some embodiments, Ring A is an optionally substituted diphenyl ether. In some embodiments, Ring A is an optionally substituted phenyl benzyl ether. In other embodiments, Ring A is an optionally substituted pyridine methoxy phenyl group.




















**[0053]** In certain embodiments, the Ring A group of formulae **I** and **II** is substituted as defined herein. In some embodiments, Ring A is substituted with one, two, or three groups independently selected from halogen, R<sup>0</sup>, or -(CH<sub>2</sub>)<sub>0-4</sub>OR<sup>0</sup>, or -O(CH<sub>2</sub>)<sub>0-4</sub>R<sup>0</sup>, wherein each R<sup>0</sup> is









as defined herein. Exemplary substituents on Ring A include Br, I, Cl, methyl,  $-\text{CF}_3$ ,  $-\text{C}\equiv\text{CH}$ ,  $-\text{OCH}_2\text{phenyl}$ ,  $-\text{OCH}_2(\text{fluorophenyl})$ , or  $-\text{OCH}_2\text{pyridyl}$ .

[0054] Exemplary Ring A groups of formulae I and II are set forth in Table 1.

**Table 1. Exemplary Ring A Groups**



*xviii**xix**xx**xxi**xxii**xxiii**xxiv**xxv**xxvi**xxvii**xxviii**xxix**xxx**xxxi**xxxii**xxxiii**xxxiv**xxxv**xxxvi*

*xxxvii**xxxviii**xxxix**xl**xli**xlii**xliii**xliv**xlvi**xlvi**xlvi**xlvi**xlvi**l**li**lii**liii**liv**lv**lvi*


[0055] As defined generally above, the Ring B group of formula I is phenyl, a 5-6 membered heteroaryl ring having 1-3 heteratoms independently selected from N, O or S, a 5-6 membered saturated heterocyclic ring having 1-2 heteratoms independently selected from N, O or S, or an 8-10 membered bicyclic partially unsaturated or aryl ring having 1-3 heteratoms independently selected from N, O or S.


[0056] In some embodiments, the Ring B group of formula I is phenyl. In some embodiments, Ring B is a 6-membered heteroaryl ring having 1-3 nitrogens. In some embodiments, Ring B is a 5-membered heteroaryl ring having 1 or 2 or 3 heteratoms independently selected from N, O or S.

[0057] In some embodiments, the Ring B group of formula I is a 5-6 membered saturated heterocyclic ring having 1 nitrogen. In some embodiments, Ring B is a 9-10 membered bicyclic partially saturated heteroaryl ring having 1-3 nitrogens. In some embodiments, Ring B is a 9-10 membered bicyclic partially saturated heteroaryl ring having 1 nitrogen.

[0058] Exemplary Ring B groups are set forth in Table 2.

**Table 2. Ring B Groups**





[0059] In some embodiments, the m moiety of formula I, II, IIa, or IIb is 1, 2, 3 or 4. In some embodiments, m is 1. In other embodiments, m is 0.

[0060] As defined generally above, each  $R^x$  group of formula I or II is independently selected from -R, halogen, -OR, -CN, -NO<sub>2</sub>, -SO<sub>2</sub>R, -SOR, -C(O)R, -CO<sub>2</sub>R, -C(O)N(R)<sub>2</sub>, -NRC(O)R, -NRC(O)NR<sub>2</sub>, -NRSO<sub>2</sub>R, or -N(R)<sub>2</sub>, or

$R^x$  and  $R^1$  are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group, wherein the warhead group is -Q-Z, and said ring is further substituted with 0-3 groups independently selected from oxo, halogen, CN, or C<sub>1-6</sub> aliphatic.


[0061] In some embodiments, each instance of  $R^x$  is independently selected from -R, -OR or halogen. In certain embodiments,  $R^x$  is lower alkyl, lower alkoxy, or halogen. Exemplary  $R^x$  groups include methyl, methoxy, and chloro. In some embodiments,  $R^x$  is hydrogen.

[0062] In some embodiments, the G group of any of formula II, II-a, or II-b is CH. In other embodiments, the G group of any of formula II, II-a, or II-b is N.

[0063] As defined generally above, the W group of formula I is a bivalent C<sub>1-3</sub> alkylene chain wherein one methylene unit of W is optionally replaced by -NR<sup>2</sup>-, -N(R<sup>2</sup>)C(O)-, -C(O)N(R<sup>2</sup>)-, -N(R<sup>2</sup>)SO<sub>2</sub>-, -SO<sub>2</sub>N(R<sup>2</sup>)-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -S-, -SO- or -SO<sub>2</sub>-.


[0064] In certain embodiments, the W group of formula I is -NH-, -S-, or -O-. In some embodiments, the W group of formula I is -CH<sub>2</sub>O-, -CH<sub>2</sub>S-, or -CH<sub>2</sub>NH-. In some aspects, W is -OCH<sub>2</sub>-, -SCH<sub>2</sub>-, -NHCH<sub>2</sub>-, or -CH<sub>2</sub>CH<sub>2</sub>-.

[0065] In some embodiments, the W group of formula I is -O- thus forming a compound of formula I-i:

**I-i**


or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.

[0066] In some embodiments, W is  $-NR^2-$  thus forming a compound of formula **I-ii**:

**I-ii**


or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^2$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.

[0067] In some embodiments, W is  $-S-$  thus forming a compound of formula **I-iii**:

**I-iii**


or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.

[0068] In some embodiments, the W group of formula **II** is  $-O-$  thus forming a compound of formula **II-i**:

**II-i**

or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.


[0069] In some embodiments, the W group of formula II is  $-NR^2-$  thus forming a compound of formula II-ii:

**II-ii**

or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^2$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.


[0070] In certain embodiments,  $R^2$  is hydrogen. In some embodiments,  $R^2$  is methyl. In still other embodiments,  $R^2$  is lower alkyl.

[0071] In some embodiments, the W group of formula II is  $-S-$  thus forming a compound of formula II-iii:

**II-iii**

or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.


[0072] In certain embodiments, the G group of formula II is CH, thus forming a compound of formula III:




III

or a pharmaceutically acceptable salt thereof, wherein each of Ring A, W, R<sup>1</sup>, R<sup>x</sup>, R<sup>y</sup>, and m are as defined above and described in classes and subclasses above and herein.


[0073] In certain embodiments, the compound of formula III is of formula III-a-i, III-b-i, III-a-ii, III-b-ii, III-a-iii, or III-b-iii:




III-a-i





III-b-i



III-a-ii




III-b-ii

**III-a-iii****III-b-iii**



or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^2$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.





[0074] In certain embodiments, the G group of formula II is N, thus forming a compound of formula IV:

**IV**

or a pharmaceutically acceptable salt thereof, wherein each of Ring A, W,  $R^1$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.



[0075] In certain embodiments, the compound of formula IV is of formula IV-a-i, IV-b-i, IV-a-ii, IV-b-ii, IV-a-iii, or IV-b-iii:

**IV-a-i****IV-b-i**

**IV-a-ii****IV-b-ii****IV-a-iii****IV-b-iii**


or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^2$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.

[0076] According to some aspects,  $R^2$  and a substituent on Ring A are taken together with their intervening atoms to form a 4-7 membered saturated or partially unsaturated ring, thus forming a compound of formula **II-a-iv** or **II-b-iv**:

**II-a-iv****II-b-iv**


or a pharmaceutically acceptable salt thereof, wherein each of Ring A,  $R^1$ ,  $R^x$ , and  $m$  are as defined above and described in classes and subclasses above and herein.

[0077] In certain embodiments, the present invention provides a compound of formula **II-a-v** or **II-b-v** wherein Ring A is phenyl and said compound is of formula **II-a-v** or **II-b-v**:



or a pharmaceutically acceptable salt thereof, wherein the Ring A phenyl moiety is optionally substituted and each of  $R^1$ ,  $R^x$ ,  $R^y$ , and  $m$  are as defined above and described in classes and subclasses above and herein.

[0078] In some embodiments,  $R^2$  is hydrogen. In other embodiments,  $R^2$  and  $R^y$  are taken together thereby forming a compound of formula **II-a-vi** or **II-b-vi**:



[0079] As defined generally above, the  $R^1$  group of formulae **I** and **II** is  $-L-Y$ , wherein:

$L$  is a covalent bond or a bivalent  $C_{1-8}$  saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one, two, or three methylene units of  $L$  are optionally and independently replaced by cyclopropylene,  $-NR-$ ,  $-N(R)C(O)-$ ,  $-C(O)N(R)-$ ,  $-N(R)SO_2-$ ,  $-SO_2N(R)-$ ,  $-O-$ ,  $-C(O)-$ ,  $-OC(O)-$ ,  $-C(O)O-$ ,  $-S-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-C(=S)-$ ,  $-C(=NR)-$ ,  $-N=N-$ , or  $-C(=N_2)-$ ;

$Y$  is hydrogen,  $C_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $NO_2$ , or  $CN$ , or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with 1-4  $R^e$  groups; and

each  $R^e$  is independently selected from  $-Q-Z$ , oxo,  $NO_2$ , halogen,  $CN$ , a suitable leaving group, or a  $C_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $NO_2$ , or  $CN$ , wherein:

Q is a covalent bond or a bivalent C<sub>1-6</sub> saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by -N(R)-, -S-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -SO-, or -SO<sub>2</sub>-, -N(R)C(O)-, -C(O)N(R)-, -N(R)SO<sub>2</sub>-, or -SO<sub>2</sub>N(R)-; and

Z is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.

[0080] In certain embodiments, L is a covalent bond.

[0081] In certain embodiments, L is a bivalent C<sub>1-8</sub> saturated or unsaturated, straight or branched, hydrocarbon chain. In certain embodiments, L is -CH<sub>2</sub>-.

[0082] In certain embodiments, L is a covalent bond, -CH<sub>2</sub>-, -NH-, -CH<sub>2</sub>NH-, -NHCH<sub>2</sub>-, -NHC(O)-, -NHC(O)CH<sub>2</sub>OC(O)-, -CH<sub>2</sub>NHC(O)-, -NHSO<sub>2</sub>-, -NHSO<sub>2</sub>CH<sub>2</sub>-, -NHC(O)CH<sub>2</sub>OC(O)-, or -SO<sub>2</sub>NH-.

[0083] In some embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and one or two additional methylene units of L are optionally and independently replaced by -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, -C(O)O-, cyclopropylene, -O-, -N(R)-, or -C(O)-.

[0084] In certain embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-.

[0085] In some embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-.

[0086] As described above, in certain embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond. One of ordinary skill in the art will recognize that such a double bond may exist within the hydrocarbon chain backbone or may be "exo" to the backbone chain and thus forming an alkylidene group. By way of example, such an L group having an alkylidene branched chain includes -CH<sub>2</sub>C(=CH<sub>2</sub>)CH<sub>2</sub>-. Thus, in some embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one alkylidenyl double bond. Exemplary L groups include -NHC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-.

[0087] In certain embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-. In certain embodiments, L is -C(O)CH=CH(CH<sub>3</sub>)-, -C(O)CH=CHCH<sub>2</sub>NH(CH<sub>3</sub>)-, -C(O)CH=CH(CH<sub>3</sub>)-, -C(O)CH=CH-, -CH<sub>2</sub>C(O)CH=CH-, -CH<sub>2</sub>C(O)CH=CH(CH<sub>3</sub>)-, -CH<sub>2</sub>CH<sub>2</sub>C(O)CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>C(O)CH=CHCH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>C(O)CH=CHCH<sub>2</sub>NH(CH<sub>3</sub>)-, or -CH<sub>2</sub>CH<sub>2</sub>C(O)CH=CH(CH<sub>3</sub>)-, or -CH(CH<sub>3</sub>)OC(O)CH=CH-.

[0088] In certain embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -OC(O)-.

[0089] In some embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-. In some embodiments, L is -CH<sub>2</sub>OC(O)CH=CHCH<sub>2</sub>-, -CH<sub>2</sub>-OC(O)CH=CH-, or -CH(CH=CH<sub>2</sub>)OC(O)CH=CH-.

[0090] In certain embodiments, L is -NRC(O)CH=CH-, -NRC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NRC(O)CH=CHCH<sub>2</sub>O-, -CH<sub>2</sub>NRC(O)CH=CH-, -NRSO<sub>2</sub>CH=CH-, -NRSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NRC(O)(C=N<sub>2</sub>)C(O)-, -NRC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NRSO<sub>2</sub>CH=CH-, -NRSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NRC(O)CH=CHCH<sub>2</sub>O-, -NRC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>NRC(O)-, -CH<sub>2</sub>NRC(O)CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>NRC(O)-, or -CH<sub>2</sub>NRC(O)cyclopropylene-, wherein each R is independently hydrogen or optionally substituted C<sub>1-6</sub> aliphatic.

[0091] In certain embodiments, L is -NHC(O)CH=CH-, -NHC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NHC(O)CH=CHCH<sub>2</sub>O-, -CH<sub>2</sub>NHC(O)CH=CH-, -NHSO<sub>2</sub>CH=CH-, -NHSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NHC(O)(C=N<sub>2</sub>)C(O)-, -NHC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NHSO<sub>2</sub>CH=CH-, -NHSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NHC(O)CH=CHCH<sub>2</sub>O-, -NHC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>NHC(O)-, -CH<sub>2</sub>NHC(O)CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>NHC(O)-, or -CH<sub>2</sub>NHC(O)cyclopropylene-.

[0092] In some embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one triple bond. In certain embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one triple bond and one or two additional methylene units of L are optionally and independently replaced by -NRC(O)-, -C(O)NR-, -S-, -S(O)-, -SO<sub>2</sub>-, -C(=S)-, -C(=NR)-, -O-, -N(R)-, or -C(O)-. In some embodiments, L has at least

one triple bond and at least one methylene unit of L is replaced by -N(R)-, -N(R)C(O)-, -C(O)-, -C(O)O-, or -OC(O)-, or -O-.

[0093] Exemplary L groups include -C≡C-, -C≡CCH<sub>2</sub>N(isopropyl)-, -NHC(O)C≡CCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-, -C≡CCH<sub>2</sub>O-, -CH<sub>2</sub>C(O)C≡C-, -C(O)C≡C-, or -CH<sub>2</sub>OC(=O)C≡C-.

[0094] In certain embodiments, L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein one methylene unit of L is replaced by cyclopropylene and one or two additional methylene units of L are independently replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, or -SO<sub>2</sub>N(R)-. Exemplary L groups include -NHC(O)-cyclopropylene-SO<sub>2</sub>- and -NHC(O)-cyclopropylene-.

[0095] As defined generally above, Y is hydrogen, C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with at 1-4 R<sup>e</sup> groups, each R<sup>e</sup> is independently selected from -Q-Z, oxo, NO<sub>2</sub>, halogen, CN, a suitable leaving group, or C<sub>1-6</sub> aliphatic, wherein Q is a covalent bond or a bivalent C<sub>1-6</sub> saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by -N(R)-, -S-, -O-, -C(O)-, -OC(O)-, -C(O)O-, -SO-, or -SO<sub>2</sub>-, -N(R)C(O)-, -C(O)N(R)-, -N(R)SO<sub>2</sub>-, or -SO<sub>2</sub>N(R)-; and, Z is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.

[0096] In certain embodiments, Y is hydrogen.

[0097] In certain embodiments, Y is C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN. In some embodiments, Y is C<sub>2-6</sub> alkenyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN. In other embodiments, Y is C<sub>2-6</sub> alkynyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN. In some embodiments, Y is C<sub>2-6</sub> alkenyl. In other embodiments, Y is C<sub>2-4</sub> alkynyl.




[0098] In other embodiments, Y is C<sub>1-6</sub> alkyl substituted with oxo, halogen, NO<sub>2</sub>, or CN. Such Y groups include -CH<sub>2</sub>F, -CH<sub>2</sub>Cl, -CH<sub>2</sub>CN, and -CH<sub>2</sub>NO<sub>2</sub>.

[0099] In certain embodiments, Y is a saturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein Y is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein.

[00100] In some embodiments, Y is a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2 R<sup>e</sup> groups,

wherein each  $R^e$  is as defined above and described herein. Exemplary such rings are epoxide and oxetane rings, wherein each ring is substituted with 1-2  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein.

[00101] In other embodiments, Y is a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein. Such rings include piperidine and pyrrolidine, wherein each ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined


above and described herein. In certain embodiments, Y is , , or , wherein each R, Q, Z, and  $R^e$  is as defined above and described herein.

[00102] In some embodiments, Y is a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein. In certain embodiments, Y is cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl, wherein each ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein..


In certain embodiments, Y is , wherein  $R^e$  is as defined above and described herein. In certain embodiments, Y is cyclopropyl optionally substituted with halogen, CN or  $NO_2$ .

[00103] In certain embodiments, Y is a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein.

[00104] In some embodiments, Y is a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein. In some embodiments, Y is cyclopropenyl, cyclobutenyl, cyclopentenyl, or cyclohexenyl wherein each ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined

above and described herein. In certain embodiments, Y is , wherein each  $R^e$  is as defined above and described herein.


[00105] In certain embodiments, Y is a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein. In certain embodiments, Y is selected from:



wherein each R and R<sup>e</sup> is as defined above and described herein.


[00106] In certain embodiments, Y is a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> group is as defined above and described herein. In certain embodiments, Y is phenyl, pyridyl, or pyrimidinyl, wherein each ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein.


[00107] In some embodiments, Y is selected from:



wherein each R<sup>e</sup> is as defined above and described herein.

[00108] In other embodiments, Y is a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3 R<sup>e</sup> groups, wherein each R<sup>e</sup> group is as defined above and described herein. In some embodiments, Y is a 5 membered partially unsaturated or aryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, and sulfur, wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> group is as defined above and described herein. Exemplary such rings are isoxazolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, pyrrolyl, furanyl, thienyl, triazole, thiadiazole, and oxadiazole, wherein each ring is substituted with 1-3 R<sup>e</sup> groups, wherein each R<sup>e</sup> group is as defined above and described herein. In certain embodiments, Y is selected from:





wherein each R and  $R^e$  is as defined above and described herein.

[00109] In certain embodiments, Y is an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein. According to another aspect, Y is a 9-10 membered bicyclic, partially unsaturated, or aryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein. Exemplary such bicyclic rings include 2,3-dihydrobenzo[d]isothiazole, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein.

[00110] As defined generally above, each  $R^e$  group is independently selected from  $-Q-Z$ , oxo,  $NO_2$ , halogen, CN, a suitable leaving group, or  $C_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $NO_2$ , or CN, wherein Q is a covalent bond or a bivalent  $C_{1-6}$  saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by  $-N(R)-$ ,  $-S-$ ,  $-O-$ ,  $-C(O)-$ ,  $-OC(O)-$ ,  $-C(O)O-$ ,  $-SO-$ , or  $-SO_2-$ ,  $-N(R)C(O)-$ ,  $-C(O)N(R)-$ ,  $-N(R)SO_2-$ , or  $-SO_2N(R)-$ ; and Z is hydrogen or  $C_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $NO_2$ , or CN.

[00111] In certain embodiments,  $R^e$  is  $C_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $NO_2$ , or CN. In other embodiments,  $R^e$  is oxo,  $NO_2$ , halogen, or CN.

[00112] In some embodiments,  $R^e$  is  $-Q-Z$ , wherein Q is a covalent bond and Z is hydrogen (i.e.,  $R^e$  is hydrogen). In other embodiments,  $R^e$  is  $-Q-Z$ , wherein Q is a bivalent  $C_{1-6}$  saturated or unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by  $-NR-$ ,  $-NRC(O)-$ ,  $-C(O)NR-$ ,  $-S-$ ,  $-O-$ ,  $-C(O)-$ ,  $-SO-$ , or  $-SO_2-$ . In other embodiments, Q is a bivalent  $C_{2-6}$  straight or branched, hydrocarbon

chain having at least one double bond, wherein one or two methylene units of Q are optionally and independently replaced by  $\text{-NR-}$ ,  $\text{-NRC(O)-}$ ,  $\text{-C(O)NR-}$ ,  $\text{-S-}$ ,  $\text{-O-}$ ,  $\text{-C(O)-}$ ,  $\text{-SO-}$ , or  $\text{-SO}_2\text{-}$ . In certain embodiments, the Z moiety of the  $\text{R}^e$  group is hydrogen. In some embodiments,  $\text{-Q-Z}$  is  $\text{-NHC(O)CH=CH}_2$  or  $\text{-C(O)CH=CH}_2$ .

[00113] In certain embodiments, each  $\text{R}^e$  is independently selected from from oxo,  $\text{NO}_2$ , CN, fluoro, chloro,  $\text{-NHC(O)CH=CH}_2$ ,  $\text{-C(O)CH=CH}_2$ ,  $\text{-CH}_2\text{CH=CH}_2$ ,  $\text{-C}\equiv\text{CH}$ ,  $\text{-C(O)OCH}_2\text{Cl}$ ,  $\text{-C(O)OCH}_2\text{F}$ ,  $\text{-C(O)OCH}_2\text{CN}$ ,  $\text{-C(O)CH}_2\text{Cl}$ ,  $\text{-C(O)CH}_2\text{F}$ ,  $\text{-C(O)CH}_2\text{CN}$ , or  $\text{-CH}_2\text{C(O)CH}_3$ .


[00114] In certain embodiments,  $\text{R}^e$  is a suitable leaving group, ie a group that is subject to nucleophilic displacement. A "suitable leaving" is a chemical group that is readily displaced by a desired incoming chemical moiety such as the thiol moiety of a cysteine of interest. Suitable leaving groups are well known in the art, e.g., see, "Advanced Organic Chemistry," Jerry March, 5<sup>th</sup> Ed., pp. 351-357, John Wiley and Sons, N.Y. Such leaving groups include, but are not limited to, halogen, alkoxy, sulphonyloxy, optionally substituted alkylsulphonyloxy, optionally substituted alkenylsulphonyloxy, optionally substituted arylsulphonyloxy, acyl, and diazonium moieties. Examples of suitable leaving groups include chloro, iodo, bromo, fluoro, acetoxy, methanesulphonyloxy (mesyloxy), tosyloxy, triflyloxy, nitro-phenylsulphonyloxy (nosyloxy), and bromo-phenylsulphonyloxy (brosyloxy).

[00115] In certain embodiments, the following embodiments and combinations of  $\text{-L-Y}$  apply:

- (a) L is a bivalent  $\text{C}_{2-8}$  straight or branched, hydrocarbon chain wherein L has at least one double bond and one or two additional methylene units of L are optionally and independently replaced by  $\text{-NRC(O)-}$ ,  $\text{-C(O)NR-}$ ,  $\text{-N(R)SO}_2\text{-}$ ,  $\text{-SO}_2\text{N(R)-}$ ,  $\text{-S-}$ ,  $\text{-S(O)-}$ ,  $\text{-SO}_2\text{-}$ ,  $\text{-OC(O)-}$ ,  $\text{-C(O)O-}$ , cyclopropylene,  $\text{-O-}$ ,  $\text{-N(R)-}$ , or  $\text{-C(O)-}$ ; and Y is hydrogen or  $\text{C}_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or
- (b) L is a bivalent  $\text{C}_{2-8}$  straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by  $\text{-C(O)-}$ ,  $\text{-NRC(O)-}$ ,  $\text{-C(O)NR-}$ ,  $\text{-N(R)SO}_2\text{-}$ ,  $\text{-SO}_2\text{N(R)-}$ ,  $\text{-S-}$ ,  $\text{-S(O)-}$ ,  $\text{-SO}_2\text{-}$ ,  $\text{-OC(O)-}$ , or  $\text{-C(O)O-}$ , and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene,  $\text{-O-}$ ,  $\text{-N(R)-}$ , or  $\text{-C(O)-}$ ; and Y is hydrogen or  $\text{C}_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or

- (c) L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (d) L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (e) L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -OC(O)-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (f) L is -NRC(O)CH=CH-, -NRC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NRC(O)CH=CHCH<sub>2</sub>O-, -CH<sub>2</sub>NRC(O)CH=CH-, -NRSO<sub>2</sub>CH=CH-, -NRSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NRC(O)(C=N<sub>2</sub>)-, -NRC(O)(C=N<sub>2</sub>)C(O)-, -NRC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NRSO<sub>2</sub>CH=CH-, -NRSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NRC(O)CH=CHCH<sub>2</sub>O-, -NRC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>NRC(O)-, -CH<sub>2</sub>NRC(O)CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>NRC(O)-, or -CH<sub>2</sub>NRC(O)cyclopropylene-; wherein R is H or optionally substituted C<sub>1-6</sub> aliphatic; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (g) L is -NHC(O)CH=CH-, -NHC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NHC(O)CH=CHCH<sub>2</sub>O-, -CH<sub>2</sub>NHC(O)CH=CH-, -NHSO<sub>2</sub>CH=CH-, -NHSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NHC(O)(C=N<sub>2</sub>)-, -NHC(O)(C=N<sub>2</sub>)C(O)-, -NHC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NHSO<sub>2</sub>CH=CH-, -NHSO<sub>2</sub>CH=CHCH<sub>2</sub>-, -NHC(O)CH=CHCH<sub>2</sub>O-, -NHC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-, -CH<sub>2</sub>NHC(O)-, -CH<sub>2</sub>NHC(O)CH=CH-, -CH<sub>2</sub>CH<sub>2</sub>NHC(O)-, or -CH<sub>2</sub>NHC(O)cyclopropylene-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (h) L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one alkylidenyl double bond and at least one methylene unit of L is replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or


- (i) L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one triple bond and one or two additional methylene units of L are optionally and independently replaced by -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (j) L is -C≡C-, -C≡CCH<sub>2</sub>N(isopropyl)-, -NHC(O)C≡CCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-, -C≡CCH<sub>2</sub>O-, -CH<sub>2</sub>C(O)C≡C-, -C(O)C≡C-, or -CH<sub>2</sub>OC(=O)C≡C-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (k) L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein one methylene unit of L is replaced by cyclopropylene and one or two additional methylene units of L are independently replaced by -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (l) L is a covalent bond and Y is selected from:
- (i) C<sub>1-6</sub> alkyl substituted with oxo, halogen, NO<sub>2</sub>, or CN;
  - (ii) C<sub>2-6</sub> alkenyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
  - (iii) C<sub>2-6</sub> alkynyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
  - (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein; or
  - (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein; or




- (vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein; or
- (viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is

substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




wherein each R and  $R^e$  is as defined above and described herein; or


- (xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or



wherein each  $R^e$  is as defined above and described herein; or

- (xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or





wherein each R and  $R^e$  is as defined above and described herein; or

(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein;

(m) L is  $-C(O)-$  and Y is selected from:

- (i)  $C_{1-6}$  alkyl substituted with oxo, halogen,  $NO_2$ , or CN; or
- (ii)  $C_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iii)  $C_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or


(vi) , wherein each R, Q, Z, and  $R^e$  is as defined above and described herein; or

The image shows three chemical structures of saturated 5-6 membered heterocyclic rings with one nitrogen atom. The first structure is a 5-membered ring with a nitrogen atom bonded to a  $-Q-Z$  group. The second structure is a 5-membered ring with a nitrogen atom bonded to an  $-NR$  group. The third structure is a 6-membered ring with a nitrogen atom bonded to a wavy line. Each structure has wavy lines indicating attachment points and substituents labeled as  $(R^e)_{1-2}$ .


- (vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or
- (viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is

substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




wherein each R and  $R^e$  is as defined above and described herein; or


- (xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or



wherein each  $R^e$  is as defined above and described herein; or

- (xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or





wherein each R and  $R^e$  is as defined above and described herein; or

(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein;

(n) L is  $-N(R)C(O)-$  and Y is selected from:

- (i)  $C_{1-6}$  alkyl substituted with oxo, halogen,  $NO_2$ , or CN; or
- (ii)  $C_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iii)  $C_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or


(vi) , wherein each R, Q, Z, and  $R^e$  is as defined above and described herein; or

(vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or


(viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is

substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




wherein each R and  $R^e$  is as defined above and described herein; or


- (xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or



wherein each  $R^e$  is as defined above and described herein; or

- (xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or






wherein each R and R<sup>e</sup> is as defined above and described herein; or

(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein R<sup>e</sup> is as defined above and described herein;

(o) L is a bivalent C<sub>1-8</sub> saturated or unsaturated, straight or branched, hydrocarbon chain; and

Y is selected from:

- (i) C<sub>1-6</sub> alkyl substituted with oxo, halogen, NO<sub>2</sub>, or CN;
- (ii) C<sub>2-6</sub> alkenyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (iii) C<sub>2-6</sub> alkynyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein; or


(vi) , wherein each R, Q, Z, and R<sup>e</sup> is as defined above and described herein; or

(vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4 R<sup>e</sup> groups, wherein each R<sup>e</sup> is as defined above and described herein; or


(viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is

substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




wherein each R and  $R^e$  is as defined above and described herein; or


- (xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or



wherein each  $R^e$  is as defined above and described herein; or

- (xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or





wherein each R and  $R^e$  is as defined above and described herein; or

(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein;

(p) L is a covalent bond,  $-\text{CH}_2-$ ,  $-\text{NH}-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{CH}_2\text{NH}-$ ,  $-\text{NHCH}_2-$ ,  $-\text{NHC}(\text{O})-$ ,  $-\text{NHC}(\text{O})\text{CH}_2\text{OC}(\text{O})-$ ,  $-\text{CH}_2\text{NHC}(\text{O})-$ ,  $-\text{NH}\text{SO}_2-$ ,  $-\text{NH}\text{SO}_2\text{CH}_2-$ ,  $-\text{NHC}(\text{O})\text{CH}_2\text{OC}(\text{O})-$ , or  $-\text{SO}_2\text{NH}-$ ; and Y is selected from:


(i)  $\text{C}_{1-6}$  alkyl substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or

(ii)  $\text{C}_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or

(iii)  $\text{C}_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or

(iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or


(v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

(vi) , wherein each R, Q, Z, and  $R^e$  is as defined above and described herein; or


(vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

(viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or

(ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or


(x) , wherein each  $R^e$  is as defined above and described herein; or

(xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  is as defined above and described herein; or




wherein each R and  $R^e$  is as defined above and described herein; or


(xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or



wherein each  $R^e$  is as defined above and described herein; or

(xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups, wherein each  $R^e$  group is as defined above and described herein; or







wherein each R and  $R^e$  is as defined above and described herein; or


(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups, wherein  $R^e$  is as defined above and described herein.


[00116] In certain embodiments, the Y group of formula I is selected from those set forth in Table 3, below, wherein each wavy line indicates the point of attachment to the rest of the molecule.

**Table 3. Exemplary Y groups:**











wherein each  $R^e$  is independently a suitable leaving group,  $\text{NO}_2$ ,  $\text{CN}$ , or oxo.































[00117] In certain embodiments,  $R^1$  is  $-\text{C}\equiv\text{CH}$ ,  $-\text{C}\equiv\text{CCH}_2\text{NH}(\text{isopropyl})$ ,  $-\text{NHC}(\text{O})\text{C}\equiv\text{CCH}_2\text{CH}_3$ ,  $-\text{CH}_2-\text{C}\equiv\text{C}-\text{CH}_3$ ,  $-\text{C}\equiv\text{CCH}_2\text{OH}$ ,  $-\text{CH}_2\text{C}(\text{O})\text{C}\equiv\text{CH}$ ,  $-\text{C}(\text{O})\text{C}\equiv\text{CH}$ , or  $-\text{CH}_2\text{OC}(\text{=O})\text{C}\equiv\text{CH}$ . In some embodiments,  $R^1$  is selected from  $-\text{NHC}(\text{O})\text{CH}=\text{CH}_2$ ,  $-\text{NHC}(\text{O})\text{CH}=\text{CHCH}_2\text{N}(\text{CH}_3)_2$ , or  $-\text{CH}_2\text{NHC}(\text{O})\text{CH}=\text{CH}_2$ .


[00118] In certain embodiments,  $R^1$  is selected from those set forth in Table 4, below, wherein each wavy line indicates the point of attachment to the rest of the molecule.


**Table 4: Exemplary  $R^1$  Groups**





**xx****yy****zz****aaa****bbb****ccc****ddd****eee****fff****ggg****hhh****iii****jjj****kkk****lll****mmm****nnn****ooo****ppp****qqq****vvv****rrr****sss****ttt****uuu****www****xxx****yyy****zzz****aaaa**





wherein each R<sup>e</sup> is independently a suitable leaving group, NO<sub>2</sub>, CN, or oxo.

[00119] As defined generally above, R<sup>1</sup> is a warhead group, or, when R<sup>1</sup> and R<sup>x</sup> form a ring, then -Q-Z is a warhead group. Without wishing to be bound by any particular theory, it is

believed that such R<sup>1</sup> groups, i.e. warhead groups, are particularly suitable for covalently binding to a key cysteine residue in the binding domain of certain protein kinases. Protein kinases having a cysteine residue in the binding domain are known to one of ordinary skill in the art and include ErbB1, ErbB2, and ErbB4, or a mutant thereof. In certain embodiments, compounds of the present invention have a warhead group characterized in that inventive compounds target one or more of the following cysteine residues:


|       |                                      |
|-------|--------------------------------------|
| ERBB1 | SEQ ID 7: ITQLMPFG <u>C</u> LLDYVREH |
| ERBB2 | SEQ ID 8: VTQLMPYG <u>C</u> LLDHVREN |
| ERBB4 | SEQ ID 9: VTQLMPHG <u>C</u> LLEYVHEH |

[00120] Thus, in some embodiments, R<sup>1</sup> is characterized in that the -L-Y moiety is capable of covalently binding to a cysteine residue thereby irreversibly inhibiting the enzyme. In certain embodiments, the cysteine residue is Cys797 of ErbB1, Cys805 of ErbB2 and Cys803 of ErbB4, or a mutant thereof, where the provided residue numbering is in accordance with Uniprot (code P00533 for ErbB1; code P04626 for ErbB2, and Q15303 for ErbB4). It will be understood that the Cys of ErbB1 (EGFR) is variably called 773 or 797 depending on whether the parent sequence contains the signal peptide or not. Thus, in accordance with the present invention, the relevant cysteine residue of ErbB1 may be described as Cys 773 or Cys 797 and these terms are used interchangeably.

[00121] One of ordinary skill in the art will recognize that a variety of warhead groups, as defined herein, are suitable for such covalent bonding. Such R<sup>1</sup> groups include, but are not limited to, those described herein and depicted in Table 4, *infra*. One of ordinary skill in the art will recognize that ErbB3 has no corresponding residue and, as recognized in the relevant art, is not catalytically active.

[00122] As depicted in Formula I *supra*, the R<sup>1</sup> warhead group can be in an ortho-, meta-, or para-position. In certain embodiments, the R<sup>1</sup> warhead group is in a meta-position of the phenyl ring relative to the rest of the molecule. Without wishing to be bound by any particular theory, it is believed that when R<sup>1</sup> is in such a meta-position, the warhead group is better positioned for covalent modification of the cysteine residue thus effecting irreversible inhibition of the enzyme. Indeed, it has been surprisingly found that a compound having a warhead group at a meta-position (compound I-1) irreversibly binds to ErbB1 whereas a compound having a warhead

group at a para-position (compound **I-93**) reversibly binds to ErbB1. These compounds have the following structures:



[00123] This phenomenon was determined by performing a washout experiment using the protocol described in detail in Example 42, *infra*. The results of this experiment are depicted in **Figure 2** where it is shown that compound **I-1** maintains enzyme inhibition after “washout” whereas compound **I-93** was washed away in the experiment thereby resulting in reactivated enzyme activity.

[00124] In certain embodiments,  $R^1$  is characterized in that the  $-L-Y$  moiety is capable of covalently binding to a cysteine residue of TEC, thereby irreversibly inhibiting the enzyme. In some embodiments, the cysteine residue is Cys 449.

[00125] In certain embodiments,  $R^1$  is characterized in that the  $-L-Y$  moiety is capable of covalently binding to a cysteine residue of BTK, thereby irreversibly inhibiting the enzyme. In some embodiments, the cysteine residue is Cys 481.

[00126] In certain embodiments,  $R^1$  is characterized in that the  $-L-Y$  moiety is capable of covalently binding to a cysteine residue of ITK, thereby irreversibly inhibiting the enzyme. In some embodiments, the cysteine residue is Cys 442.

[00127] In certain embodiments,  $R^1$  is characterized in that the  $-L-Y$  moiety is capable of covalently binding to a cysteine residue of BMX, thereby irreversibly inhibiting the enzyme. In some embodiments, the cysteine residue is Cys 496.


[00128] In certain embodiments,  $R^1$  is characterized in that the  $-L-Y$  moiety is capable of covalently binding to a cysteine residue of JAK3, thereby irreversibly inhibiting the enzyme. In some embodiments, the cysteine residue is Cys 909.













[00129] In certain embodiments,  $R^1$  is characterized in that the  $-L-Y$  moiety is capable of covalently binding to a cysteine residue of TXK, thereby irreversibly inhibiting the enzyme. In some embodiments, the cysteine residue is Cys 350.










[00130] One of ordinary skill in the art will recognize that a variety of warhead groups, as defined herein, are suitable for such covalent bonding. Such R<sup>1</sup> groups include, but are not limited to, those described herein and depicted in Table 3, *infra*.


[00131] Exemplary compounds of formula I are set forth in Table 5 below.

**Table 5. Exemplary Compounds of Formula I**



**I-10****I-11****I-12****I-13****I-14****I-15****I-16****I-17****I-18****I-19****I-20****I-21**


**I-22****I-23****I-24****I-25****I-26****I-27****I-28****I-29****I-30**



I-31

I-32


I-33



I-34

I-35


I-36



I-37

I-38
















I-39















I-40

I-41


I-42

**I-43****I-44****I-45****I-46****I-47****I-48****I-49****I-50****I-51****I-52****I-53****I-54****I-55****I-56****I-57**


**I-58****I-59****I-60****I-61****I-62****I-63****I-64****I-65****I-66****I-67****I-68**




**I-69**




**I-70**




**I-71**




**I-72**




**I-73**




I-74




I-75




I-76




**I-77**



**I-78**




**I-79**




I-80


I-81




I-82




I-83




I-84




I-85




I-86




I-87




I-88




I-89




I-90




I-91




I-92




I-93



I-94




I-95




I-96

[00132] In certain embodiments, the present invention provides any compound depicted in Table 5, above, or a pharmaceutically acceptable salt thereof.


[00133] In certain embodiments, the present invention provides a compound selected from:








I-16



I-17





I-18

**I-19****I-6****I-66****I-82****I-89**

or a pharmaceutically acceptable salt thereof.

**[00134]** As described herein, compounds of the present invention are irreversible inhibitors of at least one of ErbB1, ErbB2, ErbB3 and ErbB4, or a mutant thereof. In some embodiments, provided compounds are irreversible inhibitors of a TEC-kinase (e.g. BTK) and JAK3. One of ordinary skill in the art will recognize that certain compounds of the present invention are reversible inhibitors. In certain embodiments, such compounds are useful as assay comparator compounds. In other embodiments, such reversible compounds are useful as inhibitors of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, and therefore useful for treating one or disorders as described herein. Exemplary reversible compounds of the present invention are set forth in Table 6, below.

**Table 6. Reversible Inhibitors**



or a pharmaceutically acceptable salt thereof.

#### 4. Uses, Formulation and Administration

##### *Pharmaceutically Acceptable Compositions*

[00135] According to another embodiment, the invention provides a composition comprising a compound of this invention or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier, adjuvant, or vehicle. The amount of compound in compositions of this invention is such that is effective to measurably inhibit a protein kinase, particularly at least one of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, in a biological sample or in a patient. In certain embodiments, the amount of compound in compositions of this invention is such that is effective to measurably inhibit at least one of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, in a biological sample or in a patient. In certain embodiments, a composition of this invention is formulated for administration to a patient in need of such composition. In some embodiments, a composition of this invention is formulated for oral administration to a patient.

[00136] The term “patient”, as used herein, means an animal, preferably a mammal, and most preferably a human.

[00137] The term “pharmaceutically acceptable carrier, adjuvant, or vehicle” refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or

vehicles that may be used in the compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[00138] A "pharmaceutically acceptable derivative" means any non-toxic salt, ester, salt of an ester or other derivative of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

[00139] As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of at least one of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof.

[00140] Compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

[00141] For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or

castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens<sup>TM</sup>, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[00142] Pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[00143] Alternatively, pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[00144] Pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[00145] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[00146] For topical applications, provided pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol,

polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, provided pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[00147] For ophthalmic use, provided pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

[00148] Pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[00149] Most preferably, pharmaceutically acceptable compositions of this invention are formulated for oral administration.

[00150] The amount of compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, provided compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

[00151] It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

***Uses of Compounds and Pharmaceutically Acceptable Compositions***

[00152] Compounds and compositions described herein are generally useful for the inhibition of protein kinase activity of one or more enzymes.

[00153] Drug resistance is emerging as a significant challenge for targeted therapies. For example, drug resistance has been reported for Gleevec® and Iressa®, as well as several other kinase inhibitors in development. In addition, drug resistance has been reported for the cKit and PDGFR receptors. It has been reported that irreversible inhibitors may be effective against drug resistant forms of protein kinases (Kwak, E. L., R. Sordella, et al. (2005). "Irreversible inhibitors of the EGF receptor may circumvent acquired resistance to gefitinib." PNAS 102(21): 7665-7670.) Without wishing to be bound by any particular theory, it is believed that compounds of the present invention may be effective inhibitors of drug resistant forms of protein kinases.

[00154] As used herein, the term "clinical drug resistance" refers to the loss of susceptibility of a drug target to drug treatment as a consequence of mutations in the drug target.

[00155] As used herein, the term "resistance" refers to changes in the wild-type nucleic acid sequence coding a target protein, and/or the protein sequence of the target, which change, decrease or abolish the inhibitory effect of the inhibitor on the target protein.

[00156] Examples of kinases that are inhibited by the compounds and compositions described herein and against which the methods described herein are useful include ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof.

[00157] The activity of a compound utilized in this invention as an inhibitor of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, may be assayed *in vitro*, *in vivo* or in a cell line. *In vitro* assays include assays that determine inhibition of either the phosphorylation activity and/or the subsequent functional consequences, or ATPase activity of activated ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof. Alternate *in vitro* assays quantitate the ability of the inhibitor to bind to ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3. Inhibitor binding may be measured by radiolabelling the inhibitor prior to binding, isolating the inhibitor/ErbB1, inhibitor/ErbB2, inhibitor/ErbB3, inhibitor/ErbB4, inhibitor/TEC-kinase, or inhibitor/JAK3 complex and determining the amount of radiolabel bound. Alternatively, inhibitor binding may be determined by running a competition experiment where new inhibitors are incubated with ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3 bound to known radioligands. Detailed conditions for assaying a

compound utilized in this invention as an inhibitor of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, are set forth in the Examples below.

**[00158]** Protein tyrosine kinases are a class of enzymes that catalyze the transfer of a phosphate group from ATP or GTP to a tyrosine residue located on a protein substrate. Receptor tyrosine kinases act to transmit signals from the outside of a cell to the inside by activating secondary messaging effectors via a phosphorylation event. A variety of cellular processes are promoted by these signals, including proliferation, carbohydrate utilization, protein synthesis, angiogenesis, cell growth, and cell survival.

**[00159]** ErbB receptors, a major family of receptor tyrosine kinases, are composed of an extracellular ligand binding domain, a single transmembrane domain, and an intracellular domain with tyrosine kinase activity. The ErbB family comprises ErbB1 (commonly known as EGFR), ErbB2 (commonly known as HER2 or *neu*), ErbB3 (commonly known as HER3), and ErbB4 (commonly known as HER4). More than 10 ligands (including EGF, TGF $\alpha$ , AR, BTC, EPR, HB-EGF, NRG-1, NRG-2, NRG-3, NRG-4) have been identified for the various receptor family members. Upon ligand binding the extracellular domain undergoes conformational change, allowing the formation of homodimers or heterodimers with other members of the ErbB family. Dimerization induces tyrosine phosphorylation of specific residues in the intracellular domain that serve as docking sites for adaptor proteins and downstream effectors. In some contexts, activation of phosphatidyl-inositol 3-kinase (PI3K) and mitogen-activated protein kinase pathways occur, leading to cell proliferation and survival (Lin, N. U.; Winer, E. P., *Breast Cancer Res* 6: 204-210, 2004).

**[00160]** Interaction between family members is necessitated by deficiencies in ErbB2, which has no known ligand, and ErbB3, which is kinase dead. EGFR, ErbB3, and ErbB4 bind ligand to induce ErbB receptor homodimerization or heterodimerization, whereas ErbB2 functions as the preferred dimerization partner. The composition of the pairwise combinations is important for signal diversification, as dimer identity determines which downstream pathways are activated. Representative downstream gene products in the ErbB signal transduction pathway include Shc, Grb2, SOS1, Ras, Raf1, Mek, ERK1, ERK2, ER $\alpha$ , Akt, mTOR, FKHR, p27, Cyclin D1, FasL, GSK-3, Bad, and STAT3.

**[00161]** There is strong precedent for involvement of the EGFR and other members of the ErbB family in human cancer because over 60% of all solid tumors overexpress at least one of

these proteins or their ligands. Constitutively active, tumorigenic EGFR vIII, a mutant possessing a truncated extracellular domain, has been reported to be present in up to 78% of breast carcinomas and has also been found in glioblastomas. Overexpression of EGFR is commonly found in breast, lung, head and neck, bladder tumors, while ErbB2 expression is frequently elevated in human tumors of epithelial origin. Activating mutations in the tyrosine kinase domain have been identified in patients with non-small cell lung cancer (Lin, N. U.; Winer, E. P., *Breast Cancer Res* 6: 204-210, 2004). ErbB1 and/or ErbB2 amplification has also been implicated in squamous cell carcinomas, salivary gland carcinomas, ovarian carcinomas, and pancreatic cancers (Cooper, G.C. *Oncogenes*. 2<sup>nd</sup> ed. Sudbury: Jones and Barlett, 1995; Zhang, Y., et al., *Cancer Res* 66 : 1025-32, 2006). Overexpression of ErbB2 has potent transforming activity, likely due to its ability to cooperate with other ErbB receptors (Sherman, L., et al., *Oncogene* 18: 6692-99, 1999). In fact, some human cancers that overexpress both EGFR and ErbB2 have a poorer prognosis than cancers that overexpress either receptor alone.

[00162] The ErbB signaling network is often a key component in the pathogenesis of breast cancer. Amplification of ErbB2 is associated with an aggressive tumor phenotype that is characterized by relatively rapid tumor growth, metastatic spread to visceral sites, and drug resistance. ErbB2 has been shown to be amplified in 20% of axillary node-negative ("ANN") breast cancer cases, and this amplification has been identified as an independent prognostic factor for risk of recurrence in ANN breast cancer. (Andrulis, I.L., et al., *J Clin Oncol* 16: 1340-9, 1998).

[00163] Targeted blockade of ErbB signaling with trastuzumab (Herceptin), a monoclonal antibody directed at ErbB2, has been shown to improve survival in women with ErbB2-positive, advanced breast cancer. Other monoclonal antibodies directed against ErbB receptors include cetuximab (Erbix<sup>TM</sup>) and panitumumab (Vectibix<sup>TM</sup>).

[00164] Several small molecule tyrosine kinase inhibitors (TKIs) have been found to act selectively upon ErbB family members. Notable examples include gefitinib (Iressa) and erlotinib (Tarceva<sup>TM</sup>), both of which target the EGFR. These small molecules compete with ATP for binding to the kinase domain of the receptor. Compared to monoclonal antibodies, TKIs have several advantages in that they are orally bioavailable, well-tolerated, and appear to be active against truncated forms of ErbB2 and EGFR receptors (e.g., EGFR vIII) *in vitro*. In addition, the small size of small molecule TKIs may allow them to penetrate sanctuary sites such

as the central nervous system. Finally, the homology between kinase domains of ErbB receptors allows for development of TKIs that target more than one member of the ErbB family simultaneously, the advantages of which are described herein.

[00165] Although certain malignancies have been linked to the overexpression of individual receptors, efficient signal transduction relies on the coexpression of ErbB receptor family members. This cooperation of ErbB receptor family members in signal transduction and malignant transformation may limit the success of agents that target individual receptors in the treatment of cancer; a potential mechanism of resistance to agents targeting a single ErbB receptor is upregulation of other members of the receptor family (Britten, C. D., Mol Cancer Ther 3: 1335-42, 2004).

[00166] Agents that target two or more ErbB receptors are called pan-ErbB regulators. ERRP is a pan-ErbB negative regulator that is expressed in most benign pancreatic ductal epithelium and islet cells. Tumors have been found to experience a progressive loss in ERRP expression. Pan-ErbB regulators may be more successful in treating tumors than compounds that only target one ErbB receptor. Erbitux and Herceptin show success in a limited patient base (tumors having increased expression of EGFR or ErbB2), which could be partly due to lack of pan-ErbB activity.

[00167] In both *in vitro* and *in vivo* models, strategies that employ a dual ErbB approach seem to have greater antitumor activity than agents targeting a single ErbB receptor. Thus, agents that target multiple members of the ErbB family are likely to provide therapeutic benefit to a broader patient population (Zhang, Y., et al., Cancer Res 66: 1025-32, 2006). In certain embodiments, provided compounds inhibit one or more of ErbB1, ErbB2, ErbB3, and ErbB4. In some embodiments, provided compounds inhibit two or more of ErbB1, ErbB2, ErbB3, and ErbB4, or a mutant thereof, and are therefore pan-ErbB inhibitors.

[00168] Clearly, there is growing evidence to support the concurrent inhibition of two or more ErbB (i.e., pan-erbB) receptors in cancer therapy. Possible pan-ErbB approaches with small molecules include using combinations of agents that target individual ErbB receptors, using single agents that target multiple ErbB receptors, or using agents that interfere with ErbB receptor interactions (e.g., dimerization). Additional strategies include therapies utilizing a small molecule in combination with antibodies, or chemoprevention therapies (Lin, N. U.; Winer, E. P., Breast Cancer Res 6: 204-210, 2004).

[00169] An example of small molecule pan-ErbB inhibition is CI-1033, an irreversible pan-ErbB inhibitor that covalently binds to the ATP binding site of the intracellular kinase domain. Another irreversible pan-ErbB receptor tyrosine kinase inhibitor is HKI-272, which inhibits the growth of tumor cells that express ErbB-1 (EGFR) and ErbB-2 (HER-2) in culture and xenografts, and has antitumor activity in HER-2-positive breast cancer (Andrulis, I.L., et al., J Clin Oncol 16: 1340-9, 1998). Irreversible inhibitors have demonstrated superior antitumor activity in comparison with reversible inhibitors.

[00170] Neurofibromatosis type I (NF1) is a dominantly inherited human disease affecting one in 2500-3500 individuals. Several organ systems are affected, including bones, skin, iris, and the central nervous system, as manifested in learning disabilities and gliomas. A hallmark of NF1 is the development of benign tumors of the peripheral nervous system (neurofibromas), which vary greatly in both number and size among patients. Neurofibromas are heterogeneous tumors composed of Schwann cells, neurons, fibroblasts and other cells, with Schwann cells being the major (60-80%) cell type.

[00171] Abberant expression of the EGFR is associated with tumor development in NF1 and in animal models of NF1, suggesting a role in pathogenesis and representing a novel potential therapeutic target. EGFR expression affects the growth of tumor cell lines derived from NF1 patients under conditions where EGF is not the primary factor driving growth of the cells. These data suggest that EGFR may play an important role in NF1 tumorigenesis and Schwann cell transformation (DeClue, J. E., et al., J Clin Invest 105: 1233-41, 2000).

[00172] Patients with NF1 develop aggressive Schwann cell neoplasms known as malignant peripheral nerve sheath tumors (MPNSTs). Schwann cells are the major supportive cell population in the peripheral nervous system. Neoplastic Schwann cells within these neoplasms variably express the ErbB tyrosine kinases mediating NRG-1 responses (ErbB2, ErbB3, ErbB4). Neuregulin-1 (NRG-1) proteins promote the differentiation, survival, and/or proliferation of many cell types in the developing nervous system, and overexpression of NRG-1 in myelinating Schwann cells induces the formation of malignant peripheral nerve sheath tumors (MPNSTs) (Fallon, K. B., et al., J Neuro Oncol 66: 273-84, 2004).

[00173] Deregulation of Schwann cell growth is a primary defect driving the development of both benign neurofibromas and MPNST in neurofibromatosis type I (NF1) patients. Growth of MPNSTs and transformed mouse Schwann cells *in vitro* is highly EGF-dependent and can be

blocked by EGFR inhibitors under conditions where EGF is the primary growth factor. Some human MPNST cell lines have been found to demonstrate constitutive ErbB phosphorylation. While treatment with ErbB inhibitors abolishes ErbB phosphorylation and reduces DNA synthesis in these lines, effective chemotherapeutic regimens for MPNST remain elusive (Stonecypher, M. S., et al., *Oncogene* 24: 5589-5605, 2005).

[00174] Schwannomas are peripheral nerve tumors comprised almost entirely of Schwann-like cells, and typically have mutations in the neurofibromatosis type II (NF2) tumor suppressor gene. Ninety percent of NF2 patients develop bilateral vestibular schwannomas and/or spinal schwannomas. Enlarging schwannomas can compress adjacent structures, resulting in deafness and other neurologic problems. Surgical removal of these tumors is difficult, often resulting in increased patient morbidity.

[00175] Both normal human Schwann cells and schwannoma cells express neuregulin receptors (i.e., ErbB receptors), and schwannoma cells proliferate in response to neuregulin. It is possible that aberrant neuregulin production or response contributes to aberrant schwannoma cell proliferation (Pelton, P. D., et al., *Oncogene* 17: 2195-2209, 1998).

[00176] The NF2 tumor suppressor, Merlin, is a membrane/cytoskeleton-associated protein implicated in the regulation of tyrosine kinase activity. Genetic interactions between a Merlin mutation and EGFR pathway mutations have been documented in *Drosophila* (LaJeunesse, D. R., et al., *Genetics* 158: 667-79, 2001). Other evidence suggests Merlin can inhibit EGFR internalization and signaling upon cell-cell contact by restraining the EGFR into a membrane compartment from which it can neither signal nor be internalized (McClatchey, A. I., et al., *Genes and Development* 19: 2265-77, 2005; Curto, M. C., et al., *J Cell Biol* 177: 893-903, 2007).

[00177] As used herein, the terms "treatment," "treat," and "treating" refer to reversing, alleviating, delaying the onset of, or inhibiting the progress of a disease or disorder, or one or more symptoms thereof, as described herein. In some embodiments, treatment may be administered after one or more symptoms have developed. In other embodiments, treatment may be administered in the absence of symptoms. For example, treatment may be administered to a susceptible individual prior to the onset of symptoms (e.g., in light of a history of symptoms and/or in light of genetic or other susceptibility factors). Treatment may also be continued after symptoms have resolved, for example to prevent or delay their recurrence.

[00178] Provided compounds are inhibitors of one or more of ErbB1, ErbB2, ErbB3, and ErbB4 and are therefore useful for treating one or more disorders associated with activity of one or more of ErbB1, ErbB2, ErbB3, and ErbB4. Thus, in certain embodiments, the present invention provides a method for treating an ErbB1-mediated, an ErbB2-mediated, an ErbB3-mediated, and/or ErbB4-mediated disorder comprising the step of administering to a patient in need thereof a compound of the present invention, or pharmaceutically acceptable composition thereof.

[00179] As used herein, the terms “ErbB1-mediated”, “ErbB2-mediated,” “ErbB3-mediated,” and/or “ErbB4-mediated” disorders or conditions as used herein means any disease or other deleterious condition in which one or more of ErbB1, ErbB2, ErbB3, and/or ErbB4, or a mutant thereof, are known to play a role. Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which one or more of ErbB1, ErbB2, ErbB3, and/or ErbB4, or a mutant thereof, are known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from a proliferative disorder, wherein said method comprises administering to a patient in need thereof a compound or composition according to the present invention.

[00180] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more disorders selected from a cancer. In some embodiments, the cancer is associated with a solid tumor. In certain embodiments, the cancer is breast cancer, glioblastoma, lung cancer, cancer of the head and neck, colorectal cancer, bladder cancer, or non-small cell lung cancer. In some embodiments, the present invention provides a method for treating or lessening the severity of one or more disorders selected from squamous cell carcinoma, salivary gland carcinoma, ovarian carcinoma, or pancreatic cancer.

[00181] In certain embodiments, the present invention provides a method for treating or lessening the severity of neurofibromatosis type I (NF1), neurofibromatosis type II (NF2) Schwann cell neoplasms (e.g. MPNST's), or Schwannomas.

[00182] The TEC family of non-receptor tyrosine kinases, referred to herein as “TEC-kinases,” plays a central role in signaling through antigen-receptors such as the TCR, BCR and Fee receptors (reviewed in Miller A, et al. Current Opinion in Immunology 14; 331-340 (2002). TEC-kinases are essential for T cell activation. Three members of the family, Itk, Rlk and, are activated downstream of antigen receptor engagement in T cells and transmit signals to

downstream effectors, including PLC-g. Combined deletion of Itk and Rlk in mice leads to a profound inhibition of TCR responses including proliferation, cytokine production and immune responses to an intracellular parasite (*Toxoplasma gondii*) (Schaeffer et al, Science 284; 638-641 (1999) ). Intracellular signalling following TCR engagement is effected in ITK/RLK deficient T cells; inositol triphosphate production, calcium mobilization and MAP kinase activation are all reduced. Tec-kinases are also essential for B cell development and activation.

[00183] TEC-kinases include five family members, which are expressed primarily in hematopoietic cells: TEC, BTK, ITK (also known as TSK and EMT), RLK (also known as TXK), and BMX (also known as ETK). Additional related TEC-kinases have been found in *Drosophila melanogaster*, zebrafish (*Danio rerio*), skate (*Raja eglanteria*), and sea urchin (*Anthocidaris crassispina*).

[00184] Provided compounds are inhibitors of one or more TEC-kinases and are therefore useful for treating one or more disorders associated with activity of one or more TEC-kinases. Thus, in certain embodiments, the present invention provides a method for treating a TEC-mediated disorder comprising the step of administering to a patient in need thereof a compound of the present invention, or pharmaceutically acceptable composition thereof.

[00185] The term "TEC-mediated condition", as used herein means any disease or other deleterious condition in which TEC-kinases are known to play a role. Such conditions include those described herein and in Melcher, M *et al.*, "The Role of TEC Family Kinases in Inflammatory Processes", *Anti-Inflammatory & Anti-Allergy Agents in Medicinal Chemistry*, Vol. 6, No. 1, pp. 61-69 (Feb. 2007). Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which TEC-kinases are known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from autoimmune, inflammatory, proliferative, and hyperproliferative diseases and immunologically-mediated diseases including rejection of transplanted organs or tissues and Acquired Immunodeficiency Syndrome (AIDS), wherein said method comprises administering to a patient in need thereof a composition of the present invention.

[00186] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with TEC-kinases including diseases of the respiratory tract including, without limitation, reversible obstructive

airways diseases including asthma, such as bronchial, allergic, intrinsic, extrinsic and dust asthma, particularly chronic or inveterate asthma (e. g. late asthma airways hyper-responsiveness) and bronchitis. In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with Tec-kinases including those conditions characterized by inflammation of the nasal mucus membrane, including acute rhinitis, allergic, atrophic rhinitis and chronic rhinitis including rhinitis caseosa, hypertrophic rhinitis, rhinitis purulenta, rhinitis sicca and rhinitis medicamentosa; membranous rhinitis including croupous, fibrinous and pseudomembranous rhinitis and scrofulous rhinitis, seasonal rhinitis including rhinitis nervosa (hay fever) and vasomotor rhinitis, sarcoidosis, farmer's lung and related diseases, fibroid lung and idiopathic interstitial pneumonia.

[00187] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with TEC-kinases including diseases of the bone and joints including, without limitation, (pannus formation in) rheumatoid arthritis, seronegative spondyloarthropathis (including ankylosing spondylitis, psoriatic arthritis and Reiter's disease), Behcet's disease, Sjogren's syndrome, and systemic sclerosis.

[00188] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with TEC-kinases including diseases and disorders of the skin, including, without limitation, psoriasis, systemic sclerosis, atopic dermatitis, contact dermatitis and other eczematous dermatitis, seborrheic dermatitis, lichen planus, pemphigus, bullous pemphigus, epidermolysis bullosa, urticaria, angiodermas, vasculitides, erythemas, cutaneous eosinophilias, uveitis, Alopecia, areata and vernal conjunctivitis.

[00189] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with TEC-kinases including diseases and disorders of the gastrointestinal tract, including, without limitation, celiac disease, proctitis, eosinophilic gastro-enteritis, mastocytosis, pancreatitis, Crohn's disease, ulcerative colitis, food-related allergies which have effects remote from the gut, *e.g.*, migraine, rhinitis and eczema.

[00190] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with TEC-kinases

including those diseases and disorders of other tissues and systemic disease, including, without limitation, multiple sclerosis, arteriosclerosis, lupus erythematosus, systemic lupus erythematosus, Hashimoto's thyroiditis, myasthenia gravis, type I diabetes, nephrotic syndrome, eosinophilia fascitis, hyper IgE syndrome, lepromatous leprosy, sezary syndrome and idiopathic thrombocytopenia purpura, restenosis following angioplasty, tumours (for example leukemia, lymphomas including prostate cancers), arteriosclerosis, and systemic lupus erythematosus.

[00191] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with TEC-kinases including allograft rejection including, without limitation, acute and chronic allograft rejection following for example transplantation of kidney, heart, liver, lung, bone marrow, skin and cornea; and chronic graft versus host disease.

[00192] In some embodiments, the present invention relates to a method of treating or lessening the severity of one or more of the diseases or conditions associated with TEC-kinases, as recited above, wherein said method comprises administering to a patient in need thereof a compound or composition according to the present invention.

[00193] Bruton's tyrosine kinase ("BTK"), a member of TEC-kinases, is a key signaling enzyme expressed in all hematopoietic cells types except T lymphocytes and natural killer cells. BTK plays an essential role in the B-cell signaling pathway linking cell surface B-cell receptor (BCR) stimulation to downstream intracellular responses.

[00194] BTK is a key regulator of B-cell development, activation, signaling, and survival (Kurosaki, Curr Op Imm, 2000, 276-281; Schaeffer and Schwartzberg, Curr Op Imm 2000, 282-288). In addition, BTK plays a role in a number of other hematopoietic cell signaling pathways, e.g., Toll like receptor (TLR) and cytokine receptor—mediated TNF- $\alpha$  production in macrophages, IgE receptor (Fc $\epsilon$ silonRI) signaling in mast cells, inhibition of Fas/APO-1 apoptotic signaling in B-lineage lymphoid cells, and collagen-stimulated platelet aggregation. See, e.g., C. A. Jeffries, et al., (2003), Journal of Biological Chemistry 278:26258-26264; N. J. Horwood, et al., (2003), The Journal of Experimental Medicine 197: 1603- 1611 ; Iwaki et al. (2005), Journal of Biological Chemistry 280(48):40261 -40270; Vassilev et al. (1999), Journal of Biological Chemistry 274(3): 1646-1656, and Quek et al. (1998), Current Biology 8(20): 1137-1140.

[00195] Patients with mutations in BTK have a profound block in B cell development, resulting in the almost complete absence of B lymphocytes and plasma cells, severely reduced Ig levels and a profound inhibition of humoral response to recall antigens (reviewed in Vihinen et al Frontiers in Bioscience 5 : d917-928). Mice deficient in BTK also have a reduced number of peripheral B cells and greatly decreased levels of IgM and IgG3. BTK deletion in mice has a profound effect on B cell proliferation induced by anti-IgM, and inhibits immune responses to thymus-independent type II antigens (Ellmeier et al, J Exp Med 192: 1611-1623 (2000) ). BTK also plays a crucial role in mast cell activation through the high-affinity IgE receptor (FceRI). BTK deficient murine mast cells have reduced degranulation and decreased production of proinflammatory cytokines following FceRI cross-linking (Kawakami et al. Journal of leukocyte biology 65: 286-290).

[00196] Provided compounds are inhibitors of BTK and are therefore useful for treating one or more disorders associated with activity of BTK. Thus, in some embodiments, the present invention provides a method for treating an BTK-mediated disorder comprising the step of administering to a patient in need thereof a compound of the present invention, or pharmaceutically acceptable composition thereof.

[00197] As used herein, the term "BTK-mediated" disorders or conditions as used herein means any disease or other deleterious condition in which BTK, or a mutant thereof, is known to play a role. Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which BTK, or a mutant thereof, is known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from a proliferative disorder, wherein said method comprises administering to a patient in need thereof a compound or composition according to the present invention.

[00198] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK. In some embodiments, the disease or condition is an autoimmune disease, e.g., inflammatory bowel disease, arthritis, lupus, rheumatoid arthritis, psoriatic arthritis, osteoarthritis, Still's disease, juvenile arthritis, diabetes, myasthenia gravis, Hashimoto's thyroiditis, Ord's thyroiditis, Graves' disease, Sjogren's syndrome, multiple sclerosis, Guillain-Barre syndrome, acute disseminated encephalomyelitis, Addison's disease, opsoclonus-myoclonus syndrome, ankylosing

spondylosis, antiphospholipid antibody syndrome, aplastic anemia, autoimmune hepatitis, coeliac disease, Goodpasture's syndrome, idiopathic thrombocytopenic purpura, optic neuritis, scleroderma, primary biliary cirrhosis, Reiter's syndrome, Takayasu's arteritis, temporal arteritis, warm autoimmune hemolytic anemia, Wegener's granulomatosis, psoriasis, alopecia universalis, Behcet's disease, chronic fatigue, dysautonomia, endometriosis, interstitial cystitis, neuromyotonia, scleroderma, or vulvodynia.

[00199] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK, wherein the disease or condition is selected from heteroimmune conditions or diseases, which include, but are not limited to graft versus host disease, transplantation, transfusion, anaphylaxis, allergies (e.g., allergies to plant pollens, latex, drugs, foods, insect poisons, animal hair, animal dander, dust mites, or cockroach calyx), type I hypersensitivity, allergic conjunctivitis, allergic rhinitis, and atopic dermatitis.

[00200] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK, wherein the disease or condition is selected from an inflammatory disease, e.g., asthma, appendicitis, blepharitis, bronchiolitis, bronchitis, bursitis, cervicitis, cholangitis, cholecystitis, colitis, conjunctivitis, cystitis, dacryoadenitis, dermatitis, dermatomyositis, encephalitis, endocarditis, endometritis, enteritis, enterocolitis, epicondylitis, epididymitis, fasciitis, fibrositis, gastritis, gastroenteritis, hepatitis, hidradenitis suppurativa, laryngitis, mastitis, meningitis, myelitis myocarditis, myositis, nephritis, oophoritis, orchitis, osteitis, otitis, pancreatitis, parotitis, pericarditis, peritonitis, pharyngitis, pleuritis, phlebitis, pneumonitis, pneumonia, proctitis, prostatitis, pyelonephritis, rhinitis, salpingitis, sinusitis, stomatitis, synovitis, tendonitis, tonsillitis, uveitis, vaginitis, vasculitis, or vulvitis.

[00201] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK, wherein the disease or condition is selected from a cancer. In one embodiment, the cancer is a B-cell proliferative disorder, e.g., diffuse large B cell lymphoma, follicular lymphoma, chronic lymphocytic lymphoma, chronic lymphocytic leukemia, B-cell prolymphocytic leukemia, lymphoplasmacytic lymphoma/Waldenstrom macroglobulinemia, splenic marginal zone lymphoma, plasma cell myeloma, plasmacytoma, extranodal marginal zone B cell lymphoma,

nodal marginal zone B cell lymphoma, mantle cell lymphoma, mediastinal (thymic) large B cell lymphoma, intravascular large B cell lymphoma, primary effusion lymphoma, burkitt lymphoma/leukemia, or lymphomatoid granulomatosis. In some embodiments, the cancer is breast cancer or prostate cancer.

[00202] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK, wherein the disease or condition is selected from a thromboembolic disorder, e.g., myocardial infarct, angina pectoris, reocclusion after angioplasty, restenosis after angioplasty, reocclusion after aortocoronary bypass, restenosis after aortocoronary bypass, stroke, transitory ischemia, a peripheral arterial occlusive disorder, pulmonary embolism, or deep venous thrombosis.

[00203] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK, including infectious and noninfectious inflammatory events and autoimmune and other inflammatory diseases. These autoimmune and inflammatory diseases, disorders, and syndromes include inflammatory pelvic disease, urethritis, skin sunburn, sinusitis, pneumonitis, encephalitis, meningitis, myocarditis, nephritis, osteomyelitis, myositis, hepatitis, gastritis, enteritis, dermatitis, gingivitis, appendicitis, pancreatitis, cholecystitis, agammaglobulinemia, psoriasis, allergy, Crohn's disease, irritable bowel syndrome, ulcerative colitis, Sjogren's disease, tissue graft rejection, hyperacute rejection of transplanted organs, asthma, allergic rhinitis, chronic obstructive pulmonary disease (COPD), autoimmune polyglandular disease (also known as autoimmune polyglandular syndrome), autoimmune alopecia, pernicious anemia, glomerulonephritis, dermatomyositis, multiple sclerosis, scleroderma, vasculitis, autoimmune hemolytic and thrombocytopenic states, Goodpasture's syndrome, atherosclerosis, Addison's disease, Parkinson's disease, Alzheimer's disease, Type I diabetes, septic shock, systemic lupus erythematosus (SLE), rheumatoid arthritis, psoriatic arthritis, juvenile arthritis, osteoarthritis, chronic idiopathic thrombocytopenic purpura, Waldenstrom macroglobulinemia, myasthenia gravis, Hashimoto's thyroiditis, atopic dermatitis, degenerative joint disease, vitiligo, autoimmune hypopituitarism, Guillain-Barre syndrome, Behcet's disease, scleroderma, mycosis fungoides, acute inflammatory responses (such as acute respiratory distress syndrome and ischemia/reperfusion injury), and Graves' disease.

[00204] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with BTK, selected from rheumatoid arthritis, multiple sclerosis, B-cell chronic lymphocytic leukemia, hairy cell leukemia, non-Hodgkin's lymphoma, irritable bowel syndrome, Crohn's Disease, lupus and renal transplant.

[00205] Interleukin-2 tyrosine kinase ("ITK") is expressed in T cells, mast cells and natural killer cells. It is activated in T cells upon stimulation of the T cell receptor (TCR), and in mast cells upon activation of the high affinity IgE receptor. Following receptor stimulation in T cells, Lck, a src tyrosine kinase family member, phosphorylates Y511 in the kinase domain activation loop of Itk (S. D. Hyeck et al., 1997, J. Biol. Chem, 272, 25401-25408). Activated Itk, together with Zap-70 is required for phosphorylation and activation of PLC- $\gamma$ . (S. C. Bunnell et al., 2000, J. Biol. Chem., 275, 2219-2230). PLC- $\gamma$  catalyzes the formation of inositol 1,4,5-triphosphate and diacylglycerol, leading to calcium mobilization and PKC activation, respectively. These events activate numerous downstream pathways and lead ultimately to degranulation (mast cells) and cytokine gene expression (T cells) (Y. Kawakami et al., 1999, J. Leukocyte Biol., 65, 286-290).

[00206] The role of ITK in T cell activation has been confirmed in ITK knockout mice. CD4<sup>+</sup> T cells from ITK knockout mice have a diminished proliferative response in a mixed lymphocyte reaction or upon Con A or anti-CD3 stimulation. (X. C. Liao and D. R. Littman, 1995, Immunity, 3, 757-769). Also, T cells from ITK knockout mice produced little IL-2 upon TCR stimulation resulting in reduced proliferation of these cells. In another study, ITK deficient CD4<sup>+</sup> T cells produced reduced levels of cytokines including IL-4, IL-5 and IL-13 upon stimulation of the TCR, even after priming with inducing conditions. (D. J. Fowell, 1999, Immunity, 11, 399-409).

[00207] The role of ITK in PLC- $\gamma$  activation and in calcium mobilization was also confirmed in the T cells of these knockout mice, which had severely impaired IP<sub>3</sub> generation and no extracellular calcium influx upon TCR stimulation (K. Liu et al., 1998, J. Exp. Med. 187, 1721-1727). Such studies support a key role for ITK in activation of T cells and mast cells. Thus an inhibitor of ITK would be of therapeutic benefit in diseases mediated by inappropriate activation of these cells.

[00208] It has been well established that T cells play an important role in regulating the immune response (Powrie and Coffman, 1993, Immunology Today, 14, 270-274). Indeed,

activation of T cells is often the initiating event in immunological disorders. Following activation of the TCR, there is an influx of calcium that is required for T cell activation. Upon activation, T cells produce cytokines, including IL-2, 4, 5, 9, 10, and 13 leading to T cell proliferation, differentiation, and effector function. Clinical studies with inhibitors of IL-2 have shown that interference with T cell activation and proliferation effectively suppresses immune response in vivo (Waldmann, 1993, *Immunology Today*, 14, 264-270). Accordingly, agents that inhibit T lymphocyte activation and subsequent cytokine production, are therapeutically useful for selectively suppressing the immune response in a patient in need of such immunosuppression.

[00209] Mast cells play a critical roll in asthma and allergic disorders by releasing pro-inflammatory mediators and cytokines. Antigen-mediated aggregation of Fc.epsilon.RI, the high-affinity receptor for IgE results in activation of mast cells (D. B. Corry et al., 1999, *Nature*, 402, B18-23). This triggers a series of signaling events resulting in the release of mediators, including histamine, proteases, leukotrienes and cytokines (J. R. Gordon et al., 1990, *Immunology Today*, 11, 458-464.) These mediators cause increased vascular permeability, mucus production, bronchoconstriction, tissue degradation and inflammation thus playing key roles in the etiology and symptoms of asthma and allergic disorders.

[00210] Published data using ITK knockout mice suggests that in the absence of ITK function, increased numbers of memory T cells are generated (A. T. Miller et al., 2002 *The Journal of Immunology*, 168, 2163-2172). One strategy to improve vaccination methods is to increase the number of memory T cells generated (S. M. Kaech et al., *Nature Reviews Immunology*, 2, 251-262). In addition, deletion of ITK in mice results in reduced T cell receptor (TCR)-induced proliferation and secretion of the cytokines IL-2, IL-4, IL-5, IL-10 and IFN- $\gamma$  (Schaeffer et al, *Science* 284; 638-641 (1999) ), Fowell et al, *Immunology* 11, 399-409 (1999), Schaeffer et al, *Nature Immunology* 2 (12): 1183-1188 (2001) ). The immunological symptoms of allergic asthma are attenuated in ITK<sup>-/-</sup>-mice. Lung inflammation, eosinophil infiltration and mucous production are drastically reduced in ITK<sup>-/-</sup>-mice in response to challenge with the allergen OVA (Mueller et al, *Journal of Immunology* 170: 5056-5063 (2003)). ITK has also been implicated in atopic dermatitis. This gene has been reported to be more highly expressed in peripheral blood T cells from patients with moderate and/or severe atopic dermatitis than in controls or patients with mild atopic dermatitis (Matsumoto et al, *International Archives of Allergy and Immunology* 129: 327-340 (2002)).

[00211] Splenocytes from RLK<sup>-/-</sup> mice secrete half the IL-2 produced by wild type animals in response to TCR engagement (Schaeffer et al, Science 284: 638-641 (1999) ), while combined deletion of ITK and RLK in mice leads to a profound inhibition of TCR-induced responses including proliferation and production of the cytokines IL-2, IL-4, IL-5 and IFN- $\gamma$  (Schaeffer et al, Nature Immunology 2 (12): 1183-1188 (2001) ), Schaeffer et al, Science 284: 638-641 (1999)). Intracellular signalling following TCR engagement is effected in ITK/RLK deficient T cells; inositol triphosphate production, calcium mobilization, MAP kinase activation, and activation of the transcription factors NFAT and AP-1 are all reduced (Schaeffer et al, Science 284: 638-641 (1999), Schaeffer et al, Nature Immunology 2 (12): 1183-1188 (2001)).

[00212] Provided compounds are inhibitors of ITK and are therefore useful for treating one or more disorders associated with activity of ITK. Thus, in some embodiments, the present invention provides a method for treating an ITK-mediated disorder comprising the step of administering to a patient in need thereof a compound of the present invention, or pharmaceutically acceptable composition thereof.

[00213] As used herein, the term "ITK-mediated" disorders or conditions as used herein means any disease or other deleterious condition in which ITK, or a mutant thereof, is known to play a role. Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which ITK, or a mutant thereof, is known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from a mast cell-mediated condition, a basophil-mediated disorder, an immune or allergic disorder, wherein said method comprises administering to a patient in need thereof a compound or composition according to the present invention.

[00214] In some embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with ITK, wherein the disease or condition is an immune disorder, including inflammatory diseases, autoimmune diseases, organ and bone marrow transplant rejection and other disorders associated with T cell-mediated immune response or mast cell-mediated immune response.

[00215] In certain embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with ITK, wherein the disease or condition is acute or chronic inflammation, an allergy, contact dermatitis, psoriasis, rheumatoid arthritis, multiple sclerosis, type 1 diabetes, inflammatory bowel disease, Guillain-

Barre syndrome, Crohn's disease, ulcerative colitis, cancer, graft versus host disease (and other forms of organ or bone marrow transplant rejection) or lupus erythematosus.

[00216] In certain embodiments, the present invention provides a method for treating or lessening the severity of one or more diseases and conditions associated with ITK, wherein the disease or condition is a mast cell driven conditions, a basophil-mediated disorder, reversible obstructive airway disease, asthma, rhinitis, chronic obstructive pulmonary disease (COPD), peripheral T-cell lymphomas or HIV. Such conditions include those described in Readinger, J.A. et al., "Selective Targeting of ITK Blocks Multiple Steps of HIV Replication", PNAS 2008, Vol. 105, No. 18 (May 6, 2008).

[00217] The Janus kinases (JAK) are a family of tyrosine kinases consisting of JAK1, JAK2, JAK3 and TYK2. The JAKs play a critical role in cytokine signaling. The downstream substrates of the JAK family of kinases include the signal transducer and activator of transcription (STAT) proteins. JAK/STAT signaling has been implicated in the mediation of many abnormal immune responses such as allergies, asthma, autoimmune diseases such as transplant rejection, rheumatoid arthritis, amyotrophic lateral sclerosis and multiple sclerosis as well as in solid and hematologic malignancies such as leukemias and lymphomas. The pharmaceutical intervention in the JAK/STAT pathway has been reviewed [Frank Mol. Med. 5 : 432-456 (1999) & Seidel, et al, Oncogene 19 : 2645-2656 (2000)].

[00218] JAK1, JAK2, and TYK2 are ubiquitously expressed, while JAK3 is predominantly expressed in hematopoietic cells. JAK3 binds exclusively to the common cytokine receptor gamma chain (yc) and is activated by IL-2, IL-4, IL-7, IL-9, and IL-15.

[00219] The proliferation and survival of murine mast cells induced by IL-4 and IL-9 have, in fact, been shown to be dependent on JAK3-and yc-signaling [Suzuki et al, Blood 96 : 2172-2180 (2000) ].

[00220] Cross-linking of the high-affinity immunoglobulin (Ig) E receptors of sensitized mast cells leads to a release of proinflammatory mediators, including a number of vasoactive cytokines resulting in acute allergic, or immediate (type I) hypersensitivity reactions [Gordon et al, Nature 346 : 274-276 (1990) & Galli, N. Engl. J. Med., 328 : 257- 265 (1993) ]. A crucial role for JAK3 in IgE receptor-mediated mast cell responses in vitro and in vivo has been established [Malaviya, et al, Biochem. Biophys. Res. Commun. 257 : 807-813 (1999) ]. In addition, the prevention of type I hypersensitivity reactions, including anaphylaxis, mediated by

mast cell-activation through inhibition of JAK3 has also been reported [Malaviya et al, J. Biol. Chem. 274 : 27028-27038 (1999) ]. Targeting mast cells with JAK3 inhibitors modulated mast cell degranulation in vitro and prevented IgE receptor/antigen-mediated anaphylactic reactions in vivo.

[00221] A recent study described the successful targeting of JAK3 for immune suppression and allograft acceptance. The study demonstrated a dose-dependent survival of Buffalo heart allograft in Wistar Furth recipients upon administration of inhibitors of JAK3 indicating the possibility of regulating unwanted immune responses in graft versus host disease [Kirken, transpl. proc. 33 : 3268-3270 (2001) ].

[00222] IL-4-mediated STAT-phosphorylation has been implicated as the mechanism involved in early and late stages of rheumatoid arthritis (RA). Up-regulation of proinflammatory cytokines in RA synovium and synovial fluid is a characteristic of the disease. It has been demonstrated that IL-4-mediated activation of IL-4/STAT pathway is mediated through the Janus kinases (JAK 1 & 3) and that IL-4-associated JAK kinases are expressed in the RA synovium [Muller-Ladner, et al, J. Immunol. 164 : 3894-3901 (2000)].

[00223] Familial amyotrophic lateral sclerosis (FALS) is a fatal neurodegenerative disorder affecting about 10% of ALS patients. The survival rates of FALS mice were increased upon treatment with a JAK3 specific inhibitor. This confirmed that JAK3 plays a role in FALS [Trieu, et al, Biochem. Biophys. Res. Commun. 267 : 22-25 (2000)].

[00224] Signal transducer and activator of transcription (STAT) proteins are activated by, among others, the JAK family kinases. Results from a recent study suggested the possibility of intervention in the JAK/STAT signaling pathway by targeting JAK family kinases with specific inhibitors for the treatment of leukemia [Sudbeck, et al, Clin. Cancer Res. 5 : 1569-1582 (1999) ]. JAK3 specific compounds were shown to inhibit the clonogenic growth of JAK3-expressing cell lines DAUDI, RAMOS, LC1 ; 19, NALM-6, MOLT-3 and HL-60. Inhibition of JAK3 and TYK 2 abrogated tyrosine phosphorylation of STAT3, and inhibited cell growth of mycosis fungoides, a form of cutaneous T cell lymphoma.

[00225] According to another embodiment, the invention provides a method for treating or lessening the severity of a JAK3-mediated disease or condition in a patient comprising the step of administering to said patient a composition according to the present invention.

[00226] The term “JAK3-mediated disease”, as used herein means any disease or other deleterious condition in which a JAK3 kinase is known to play a role. Accordingly, another embodiment of the present invention relates to treating or lessening the severity of one or more diseases in which JAK3 is known to play a role. Specifically, the present invention relates to a method of treating or lessening the severity of a disease or condition selected from immune responses such as allergic or type I hypersensitivity reactions, asthma, autoimmune diseases such as transplant rejection, graft versus host disease, rheumatoid arthritis, amyotrophic lateral sclerosis, and multiple sclerosis, neurodegenerative disorders such as familial amyotrophic lateral sclerosis (FALS), as well as in solid and hematologic malignancies such as leukemias and lymphomas, wherein said method comprises administering to a patient in need thereof a composition according to the present invention.

[00227] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of cancer, an autoimmune disorder, a neurodegenerative or neurological disorder, schizophrenia, a bone-related disorder, liver disease, or a cardiac disorder. The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. Compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term “patient”, as used herein, means an animal, preferably a mammal, and most preferably a human.

[00228] Pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[00229] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[00230] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[00231] Injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[00232] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[00233] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[00234] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[00235] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[00236] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[00237] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper

medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

**[00238]** According to one embodiment, the invention relates to a method of inhibiting protein kinase activity in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a composition comprising said compound.

**[00239]** According to another embodiment, the invention relates to a method of inhibiting ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, activity in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a composition comprising said compound. In certain embodiments, the invention relates to a method of irreversibly inhibiting ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3,, or a mutant thereof, activity in a biological sample comprising the step of contacting said biological sample with a compound of this invention, or a composition comprising said compound.

**[00240]** The term “biological sample”, as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

**[00241]** Inhibition of protein kinase, or a protein kinase selected from ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3,, or a mutant thereof, activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, biological specimen storage, and biological assays.

**[00242]** Another embodiment of the present invention relates to a method of inhibiting protein kinase activity in a patient comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound.

**[00243]** According to another embodiment, the invention relates to a method of inhibiting one or more of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3,, or a mutant thereof, activity in a patient comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound. According to certain embodiments, the invention relates to a method of irreversibly inhibiting one or more of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3,, or a mutant thereof, activity in a patient

comprising the step of administering to said patient a compound of the present invention, or a composition comprising said compound. In other embodiments, the present invention provides a method for treating a disorder mediated by one or more of ErbB1, ErbB2, ErbB3, ErbB4, a TEC-kinase, and/or JAK3, or a mutant thereof, in a patient in need thereof, comprising the step of administering to said patient a compound according to the present invention or pharmaceutically acceptable composition thereof. Such disorders are described in detail herein.

[00244] Depending upon the particular condition, or disease, to be treated, additional therapeutic agents, which are normally administered to treat that condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat a particular disease, or condition, are known as “appropriate for the disease, or condition, being treated.”

[00245] For example, compounds of the present invention, or a pharmaceutically acceptable composition thereof, are administered in combination with chemotherapeutic agents to treat proliferative diseases and cancer. Examples of known chemotherapeutic agents include, but are not limited to, Adriamycin, dexamethasone, vincristine, cyclophosphamide, fluorouracil, topotecan, taxol, interferons, platinum derivatives, taxane (e.g., paclitaxel), vinca alkaloids (e.g., vinblastine), anthracyclines (e.g., doxorubicin), epipodophyllotoxins (e.g., etoposide), cisplatin, an mTOR inhibitor (e.g., a rapamycin), methotrexate, actinomycin D, dolastatin 10, colchicine, emetine, trimetrexate, metoprine, cyclosporine, daunorubicin, teniposide, amphotericin, alkylating agents (e.g., chlorambucil), 5-fluorouracil, camptothecin, cisplatin, metronidazole, and Gleevec<sup>TM</sup>, among others. In other embodiments, a compound of the present invention is administered in combination with a biologic agent, such as Avastin<sup>TM</sup> or VECTIBIX<sup>TM</sup>.

[00246] In certain embodiments, compounds of the present invention, or a pharmaceutically acceptable composition thereof, are administered in combination with an antiproliferative or chemotherapeutic agent selected from any one or more of Abarelix, aldesleukin, Aldesleukin, Alemtuzumab, Alitretinoin, Allopurinol, Altretamine, Amifostine, Anastrozole, Arsenic trioxide, Asparaginase, Azacitidine, BCG Live, Bevacuzimab, Fluorouracil, Bexarotene, Bleomycin, Bortezomib, Busulfan, Calusterone, Capecitabine, Camptothecin, Carboplatin, Carmustine, Celecoxib, Cetuximab, Chlorambucil, Cladribine, Clofarabine, Cyclophosphamide, Cytarabine, Dactinomycin, Darbepoetin alfa, Daunorubicin, Denileukin, Dexrazoxane, Docetaxel, Doxorubicin (neutral), Doxorubicin hydrochloride, Dromostanolone Propionate, Epirubicin,

Epoetin alfa, Erlotinib, Estramustine, Etoposide Phosphate, Etoposide, Exemestane, Filgrastim, floxuridine fludarabine, Fulvestrant, Gefitinib, Gemcitabine, Gemtuzumab, Goserelin Acetate, Histrelin Acetate, Hydroxyurea, Ibritumomab, Idarubicin, Ifosfamide, Imatinib Mesylate, Interferon Alfa-2a, Interferon Alfa-2b, Irinotecan, Lenalidomide, Letrozole, Leucovorin, Leuprolide Acetate, Levamisole, Lomustine, Megestrol Acetate, Melphalan, Mercaptopurine, 6-MP, Mesna, Methotrexate, Methoxsalen, Mitomycin C, Mitotane, Mitoxantrone, Nandrolone, Nelarabine, Nofetumomab, Oprelvekin, Oxaliplatin, Paclitaxel, Palifermin, Pamidronate, Pegademase, Pegaspargase, Pegfilgrastim, Pemetrexed Disodium, Pentostatin, Pipobroman, Plicamycin, Porfimer Sodium, Procarbazine, Quinacrine, Rasburicase, Rituximab, Sargramostim, Sorafenib, Streptozocin, Sunitinib Maleate, Talc, Tamoxifen, Temozolomide, Teniposide, VM-26, Testolactone, Thioguanine, 6-TG, Thiotepa, Topotecan, Toremifene, Tositumomab, Trastuzumab, Tretinoin, ATRA, Uracil Mustard, Valrubicin, Vinblastine, Vincristine, Vinorelbine, Zoledronate, or Zoledronic acid.

[00247] Other examples of agents the inhibitors of this invention may also be combined with include, without limitation: treatments for Alzheimer's Disease such as Aricept® and Exelon®; treatments for Parkinson's Disease such as L-DOPA/carbidopa, entacapone, ropinrole, pramipexole, bromocriptine, pergolide, trihexephendyl, and amantadine; agents for treating Multiple Sclerosis (MS) such as beta interferon (e.g., Avonex® and Rebif®), Copaxone®, and mitoxantrone; treatments for asthma such as albuterol and Singulair®; agents for treating schizophrenia such as zyprexa, risperdal, seroquel, and haloperidol; anti-inflammatory agents such as corticosteroids, TNF blockers, IL-1 RA, azathioprine, cyclophosphamide, and sulfasalazine; immunomodulatory and immunosuppressive agents such as cyclosporin, tacrolimus, rapamycin, mycophenolate mofetil, interferons, corticosteroids, cyclophosphamide, azathioprine, and sulfasalazine; neurotrophic factors such as acetylcholinesterase inhibitors, MAO inhibitors, interferons, anti-convulsants, ion channel blockers, riluzole, and anti-Parkinsonian agents; agents for treating cardiovascular disease such as beta-blockers, ACE inhibitors, diuretics, nitrates, calcium channel blockers, and statins; agents for treating liver disease such as corticosteroids, cholestyramine, interferons, and anti-viral agents; agents for treating blood disorders such as corticosteroids, anti-leukemic agents, and growth factors; and agents for treating immunodeficiency disorders such as gamma globulin.

[00248] In certain embodiments, compounds of the present invention, or a pharmaceutically acceptable composition thereof, are administered in combination with a monoclonal antibody or an siRNA therapeutic.

[00249] Those additional agents may be administered separately from an inventive compound-containing composition, as part of a multiple dosage regimen. Alternatively, those agents may be part of a single dosage form, mixed together with a compound of this invention in a single composition. If administered as part of a multiple dosage regime, the two active agents may be submitted simultaneously, sequentially or within a period of time from one another normally within five hours from one another.

[00250] As used herein, the term “combination,” “combined,” and related terms refers to the simultaneous or sequential administration of therapeutic agents in accordance with this invention. For example, a compound of the present invention may be administered with another therapeutic agent simultaneously or sequentially in separate unit dosage forms or together in a single unit dosage form. Accordingly, the present invention provides a single unit dosage form comprising a compound of formula I, an additional therapeutic agent, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

[00251] The amount of both, an inventive compound and additional therapeutic agent (in those compositions which comprise an additional therapeutic agent as described above)) that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Preferably, compositions of this invention should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of an inventive can be administered.

[00252] In those compositions which comprise an additional therapeutic agent, that additional therapeutic agent and the compound of this invention may act synergistically. Therefore, the amount of additional therapeutic agent in such compositions will be less than that required in a monotherapy utilizing only that therapeutic agent. In such compositions a dosage of between 0.01 – 1,000 µg/kg body weight/day of the additional therapeutic agent can be administered.

[00253] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of

the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[00254] The compounds of this invention, or pharmaceutical compositions thereof, may also be incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents and catheters. Vascular stents, for example, have been used to overcome restenosis (re-narrowing of the vessel wall after injury). However, patients using stents or other implantable devices risk clot formation or platelet activation. These unwanted effects may be prevented or mitigated by pre-coating the device with a pharmaceutically acceptable composition comprising a kinase inhibitor. Implantable devices coated with a compound of this invention are another embodiment of the present invention.


### EXEMPLIFICATION

[00255] As depicted in the Examples below, in certain exemplary embodiments, compounds are prepared according to the following general procedures. It will be appreciated that, although the general methods depict the synthesis of certain compounds of the present invention, the following general methods, and other methods known to one of ordinary skill in the art, can be applied to all compounds and subclasses and species of each of these compounds, as described herein.

#### EXAMPLE 1

##### SCHEME 1a

##### Synthesis of Compounds of Formula II-a



[00256] **Synthesis of (6-chloro-pyrimidin-4-yl)-(3-bromo-phenyl)-amine:** A solution of 4,6-dichloropyrimidine (5 g, 33.6 mmol), 3-bromoaniline (5.8 g, 33.7 mmol) and *N,N*-diisopropylethylamine (DIEA) (5.2 g, 40.2 mmol) in ethanol (40 mL) was heated at 80°C for 16 hr. The reaction mixture was cooled to ambient temperature, diethylether (35 mL) was added

while the mixture was being stirred. The product was precipitated, filtered, washed with water and dried to afford 5.9 g (62% yield) of a light colored solid. MS (m/z):  $MH^+$  = 284, 286, 288.

**[00257] Synthesis of 3-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]-phenylamine:** A mixture of (6-chloro-pyrimidin-4-yl)-(3-bromo-phenyl)-amine (300 mg, 1.1 mmol) and benzene-1,3-diamine (300mg, 2.75 mmol) in 3 mL of *n*-BuOH was heated in a sealed tube to 150°C for 16 hr. Solvent was removed by vacuum evaporation and the crude product was purified by flash chromatography on silica gel with EtOAc/DCM solvent system to afford 255 mg (65% yield) of the title compound as a yellow solid. MS (m/z):  $MH^+$  = 356, 358.

**[00258]** The following compounds were prepared in a manner substantially similar to that described in Scheme 1a and the Examples above:

- (a) 3-Bromoaniline and benzene-1,4-diamine gave 4-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]phenylamine: MS (m/z):  $MH^+$  = 356, 358.
- (b) 3-Chloro-4-fluoroaniline and benzene-1,3-diamine gave 3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**). MS (m/z):  $MH^+$  = 330, 332.
- (c) 3-Methylaniline and benzene-1,3-diamine gave 3-[6-(3-methylphenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-5**). MS (m/z):  $MH^+$  = 292.
- (d) 3-Chloro-4-(3-fluorophenyl)methoxyaniline and benzene-1,3-diamine gave 3-{6-[3-chloro-4-(3-fluorophenyl)methoxyphenylamino]-pyrimidin-4-ylamino}phenylamine (**I<sup>R</sup>-6**). MS (m/z):  $MH^+$  = 436, 438.
- (e) 3-Bromoaniline and 3-ethynylaniline gave *N*<sup>4</sup>-(3-Bromophenyl)-*N*<sup>6</sup>-(3-ethynylphenyl)pyrimidine-4,6-diamine (**I-12**). MS (*M* + *H*<sup>+</sup>) 365, 367; <sup>1</sup>H NMR (400 MHz, d<sup>6</sup>-DMSO) δ 9.73 (s, 1H), 9.60 (s, 1H), 8.67 (s, 1H), 7.95 (t, 1H), 7.76 (s, 1H), 7.50 (d, 1H), 7.46 (d, 1H), 7.55 (t, 1H), 7.40 (t, 1H), 7.20 (d, 1H), 7.10 (d, 1H), 6.30 (s, 1H), 4.25 (s, 1H) ppm.
- (f) 3-Ethynylaniline gave *N*<sup>4</sup>,*N*<sup>6</sup>-bis(3-ethynylphenyl)pyrimidine-4,6-diamine (**I-11**). MS (*M* + *H*<sup>+</sup>) 311, 312; <sup>1</sup>H NMR (400 MHz, d<sup>6</sup>-DMSO) δ 9.24 (s, 2H), 8.32 (s, 1H), 7.78 (s, 2H), 7.52 (d, 2H), 7.25 (t, 2H), 7.02 (d, 2H), 6.13 (s, 1H), 4.12 (s, 2H) ppm.


(g) 4-Phenoxyaniline and 2,6-diaminopyridine gave 2-[6-(4-phenoxyphenyl)-aminopyrimidin-4-yl]amino-6-aminopyridine (**I<sup>R</sup>-11**) MS (*m/z*):  $MH^+ = 372, 371$ .

(h) 4-Phenoxyaniline and 1,4-diaminobenzene gave 4-[6-(4-phenoxyphenyl)amino-pyrimidin-4-yl]amino-aminobenzene (**I<sup>R</sup>-14**) MS (*m/z*):  $MH^+ = 370$

## EXAMPLE 2

### SCHEME 2a

Sequence A for Synthesis of 3-(6-mono- or disubstituted-amino)pyrimidin-4-ylamino)-phenylamines



[00259] Although a BOC protecting group is depicted in Scheme 2a above and the ensuing schemes below, one of ordinary skill in the art will recognize that other amine protecting groups are amenable for use in preparing compounds of the present invention. Accordingly, a variety of amine protecting groups is contemplated.

[00260] **Synthesis of tert-butyl 3-(6-chloropyrimidin-4-ylamino)phenylcarbamate** 4,6-Dichloropyrimidine (1.5 g, 10 mmol), *tert*-butyl 3-aminophenylcarbamate (2.1 g, 10 mmol) and triethylamine (2.2 g, 20 mmol) were mixed in ethanol (20 mL), heated at 80 °C for 16 hr. The reaction mixture was cooled to RT and solvent was removed in vacuo. The gummy crude product was mixed with 20 mL of DCM and was stirred at RT to give an off-white solid, which was filtered and dried under vacuum (1.6 g, 5 mmol). MS (*m/z*):  $MH^+ = 321, 323$ .

[00261] **Synthesis of tert-butyl 3-(6-[*N*-methyl-*N*-phenylamino]pyrimidin-4-ylamino)phenyl-carbamate** A mixture of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)phenylcarbamate (1.6g, 5 mmol) and *N*-methylaniline (1.07 g, 10 mmol) was heated at 120°C in a sealed tube for 2 hr. The reaction mixture was cooled to RT, mixed with 1 mL of 1N

NaOH and 5 mL of DCM, stirred for 30 min, the product was filtered and dried under vacuum to give a solid product. MS (m/z):  $MH^+ = 392$ .

**[00262] Synthesis of 3-[6-(*N*-methyl-*N*-phenylamino)-pyrimidin-4-ylamino]phenylamine (*I<sup>R</sup>*-8)** A mixture of tert-butyl 3-(6-[*N*-methyl-*N*-phenylamino]pyrimidin-4-ylamino)phenylcarbamate (1.17 g, 3 mmol) and TFA (50% in DCM, 10 mL) was stirred at RT for 4 hr. The reaction solvent was removed under vacuum. The crude product was mixed with 1 mL of 2N NaOH and 5 mL of EtOAc, was stirred for 30 min, was filtered, and was dried under vacuum to give 0.65g (75%) of the title compound, an off-white solid. (*I<sup>R</sup>*-8) MS (m/z):  $MH^+ = 292$ .

**[00263]** The following compounds were prepared in a manner substantially similar to that described in Scheme 2a:


(a) 4-Phenoxyphenyl amine gave 3-[6-(4-phenoxyphenylamino)-pyrimidin-4-ylamino]phenylamine. (*I<sup>R</sup>*-7) MS (m/z):  $MH^+ = 370$ .

(b) 3-Chloro-4-(2-pyridyl)methoxyaniline gave 3-[[6-(3-chloro-4-(2-pyridyl)methoxyphenylamino)-pyrimidin-4-ylamino]phenylamine. (*I<sup>R</sup>*-9) MS (m/z):  $MH^+ = 419, 421$ .

(c) 3-Chloro-4-(3-fluorobenzyloxy)aniline gave 3-[6-(3-chloro-4-{3-fluorobenzyloxy}phenylamino)-pyrimidin-4-ylamino]-phenyl amine. (*I<sup>R</sup>*-6) MS (m/z):  $M^+ = 436, 438$ .


(d) Aniline gave 3-[6-(phenylamino)-pyrimidin-4-yl]aminophenylamine. (*I<sup>R</sup>*-15) MS (m/z):  $MH^+ = 278$ .

## SCHEME 2b

Sequence B for Synthesis of 3-(6-mono- or disubstituted-aminopyrimidin-4-yl)amino-phenylamines

[00264] **Synthesis of 3-[6-(4-bromo-2-fluorophenylamino)-pyrimidin-4-yl-amino]-phenylamine ( $I^R$ -17).** A solution of 4-bromo-2-fluoroaniline (701 mg, 3.69 mmol), diisopropylethylamine (0.70 mL, 4.03 mmol), and 4,6-dichloropyrimidine (500 mg, 3.36 mmol) in EtOH (10 mL) was heated in an 85 °C oil bath for 5 d. Flash chromatography (2% MeOH / CHCl<sub>3</sub>) of the residue gave 380 mg (37%) of *N*-(4-bromo-2-fluorophenyl)-6-chloropyrimidin-4-amine as a pale yellow solid. MS (*m/z*): 304, 302.. A suspension of *N*-(4-bromo-2-fluorophenyl)-6-chloropyrimidin-4-amine (0.37 g, 1.22 mmol) and *tert*-butyl 3-aminophenylcarbamate (0.28 g, 1.35 mmol) in *n*-BuOH (4 mL) was heated in an oil bath at 120-130 °C for 6 h. The reaction mixture was cooled and concentrated to give 0.68 g of a brown foam. Flash chromatography gave 0.16 g (28%) *tert*-butyl 3-(6-[4-bromo-2-fluorophenyl]aminopyrimidin-4-yl)aminophenylcarbamate as a white solid. MS (*m/z*): (*M* + *H*) 476, 474. *tert*-Butyl 3-(6-[4-bromo-2-fluorophenyl]aminopyrimidin-4-yl)aminophenylcarbamate (0.15 g, 0.32 mmol) was taken up in 4M HCl in dioxane (5 mL). After several minutes a white solid began to precipitate. The mixture was allowed to stand for 3 h and was concentrated via rotary evaporation. Saturated aqueous sodium bicarbonate (5 mL) was added and the mixture was sonicated for several minutes. The solid was collected by filtration, was washed with water (5 mL), and was dried overnight under vacuum to give 0.12 g (100%) of 3-[6-(4-bromo-2-fluorophenylamino)-pyrimidin-4-yl-amino]-phenylamine ( $I^R$ -17) as a white powder. MS (*m/z*): (*M* + *H*) 376, 374.

## SCHEME 2c

Synthesis of 3-(6-mono or disubstituted-aminopyrimidin-4-yl)amino-N-substituted phenylamines


**[00265] Synthesis of *tert*-butyl N-3-(6-chloropyrimidin-4-ylamino)phenyl-N-methylcarbamate** To a stirring solution of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)phenyl carbamate (2.000 g, 6.235 mmol) in 10 mL of THF at 0 °C under N<sub>2</sub> was added drop-wise a solution of 1.0 M lithium bis(trimethylsilyl)amide in *tert*-butyl methyl ether (6.23 mL, 6.23 mmol). The light yellow solution was stirred at 0 °C for 30 min then iodomethane (0.43 mL, 6.892 mmol, 1.1 eq) was added. The solution was allowed to slowly warm to room temperature overnight, was concentrated, and was then partitioned between EtOAc and saturated KH<sub>2</sub>PO<sub>4</sub> solution. The organic extract was washed with brine solution, dried (MgSO<sub>4</sub>), was filtered, was concentrated in vacuo and was chromatographed (silica gel, 2% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to give 0.904 g of *tert*-butyl N-3-(6-chloropyrimidin-4-ylamino)phenyl-N-methylcarbamate as an off-white solid. MS (m/z) M+1 = 335/337 (100/44%), <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.48 (s, 1H), 7.49 (bs, 1H), 7.38 (m, 1H), 7.24 (m, 1H), 6.91 (m, 1H), 6.71 (bs, 1H), 6.30 (s, 1H), 3.48 (s, 3H), 1.53 (s, 9H).

**[00266] Synthesis of N-3-[6-(3-Chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl-N-methylamine (Int-C)** A solution of *tert*-butyl N-3-(6-chloropyrimidin-4-ylamino)phenyl-N-methylcarbamate (0.486 g, 1.095 mmol) in 25 mL of CH<sub>2</sub>Cl<sub>2</sub> was treated with trifluoroacetic acid (5 mL). The solution was stirred at room temperature under N<sub>2</sub> for 2 h, was concentrated and was partitioned between CHCl<sub>3</sub> and 10% aq. NH<sub>4</sub>OH solution. The organic extract was washed with brine solution, was dried (MgSO<sub>4</sub>), was filtered and was concentrated to give 0.630 g of N-3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl-N-methylamine (**Int-C**) as yellow oil. MS (m/z): 344/346 (M+1, 100/68%). TLC (SiO<sub>2</sub>, 10% MeOH in CHCl<sub>3</sub>): R<sub>f</sub> 0.44.

## EXAMPLE 3

## SCHEME 3a

Sequence B for Synthesis of 3-(6-mono- or disubstituted-aminopyrimidin-4-yl)amino-phenylamines



wherein L' is a subset of L, as defined herein, such that Y-L'C(O)Cl results in formation of provided compounds wherein R<sup>1</sup> is -L-Y wherein a terminal methylene unit of L is replaced with -NHC(O)-.

**[00267] Synthesis of *N*-{3-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]-phenyl}-2-propenamide (I-1)** A solution of 3-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]phenylamine (250 mg, 0.7 mmol) and triethylamine (180 mg, 1.75 mmol) in 5 mL of THF was stirred at RT. Acryloyl chloride (80 mg, 0.9 mmol) was added into the reaction mixture and it was stirred at RT for 1h. The solvent was removed by vacuum evaporation and the crude product was purified by flash chromatography on silica gel with EtOAc/DCM solvent system to afford 115mg (40% yield) of the title compound as a light colored solid. MS (m/z): MH<sup>+</sup> = 410, 412. <sup>1</sup>H NMR (DMSO) : 10.15 (s, 1H), 9.36 (s, 1H), 9.28 (s, 1H), 8.34 (s, 1H), 8.02 (s, 1H), 8.00 (s, 1H), 7.51-7.11 (m, 5H), 6.47 (dd, 1H, J<sub>1</sub> = 10.1 Hz, J<sub>2</sub> = 17.0 Hz), 6.27 (dd, 1H, J<sub>1</sub> = 1.9 Hz, J<sub>2</sub> = 17.0 Hz), 6.20 (s, 1H), 5.76 (dd, 1H, J<sub>1</sub> = 10.1 Hz, J<sub>2</sub> = 1.9 Hz) ppm.

**[00268]** The following compounds were prepared in a manner substantially similar to those described in Schemes 3a:

- (a) 4-[6-(3-Bromophenylamino)-pyrimidin-4-ylamino]phenylamine and acryoyl chloride gave *N*-{4-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]phenyl}-2-propenamide (**I-93**). MS (m/z): MH<sup>+</sup> = 410, 412. <sup>1</sup>H NMR (DMSO): 10.10 (s, 1H), 9.33 (s, 1H), 9.17 (s, 1H), 8.30 (s, 1H), 8.02 (s, 1H), 7.62 (m, 2H), 7.46 (m, 3H), 7.22 (t, 1H, J = 8.0 Hz), 7.10 (d, 1H, J = 8.0 Hz), 6.43 (dd, 1H, J<sub>1</sub> = 10.0 Hz, J<sub>2</sub> = 17.0 Hz), 6.24 (d, 1H, J = 17.0 Hz), 6.13 (s, 1H), 5.73 (d, 1H, J = 10.0 Hz) ppm.

- (b) 3-[6-(3-Bromophenylamino)-pyrimidin-4-ylamino]phenylamine and propionyl chloride gave *N*-{3-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]phenyl}-propionamide (**I<sup>R</sup>-3**). MS (m/z):  $MH^+$  = 412, 414.  $^1H$  NMR (DMSO): 9.78 (s, 1H), 9.28 (s, 1H), 9.16 (s, 1H), 8.26 (s, 1H), 7.96 (s, 1H), 7.76 (s, 1H), 7.43 (d, 1H,  $J = 8.2$  Hz), 7.20 (m, 2H), 7.05 (m, 3H), 6.12 (s, 1H), 2.27 (q, 2H,  $J = 7.6$  Hz), 1.03 (t, 3H,  $J = 7.6$  Hz) ppm.
- (c) 3-[6-(3-Bromophenylamino)-pyrimidin-4-ylamino]phenylamine and (*E*)-2-butenoylchloride gave (*E*)-*N*-{3-[6-(3-bromophenylamino)-pyrimidin-4-ylamino]phenyl}-2-butenamide (**I-8**). MS (m/z):  $M+H^+$  = 424, 426.  $^1H$  NMR (DMSO): 9.87 (s, 1H), 9.29 (s, 1H), 9.18 (s, 1H), 8.27 (s, 1H), 7.96 (s, 1H), 7.82 (s, 1H), 7.44 (d, 1H,  $J = 8.2$  Hz), 7.20 (m, 4H), 7.05 (dd, 1H,  $J_1 = 1.0$  Hz,  $J_2 = 7.8$  Hz), 6.75 (m, 1H), 6.15 (m, 2H), 1.81 (d, 3H,  $J = 7.8$  Hz) ppm.
- (d) 3-[6-(3-Chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**) and acryoyl chloride gave *N*-{3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl}-2-propenamide (**I-2**). MS (m/z):  $MH^+$  = 384, 386.  $^1H$  NMR (DMSO): 9.29 (s, 1H), 9.20 (s, 1H), 8.27 (s, 1H), 7.93 (m, 1H), 7.86 (s, 1H), 7.41 (m, 1H), 7.25 (m, 5H), 6.42 (dd, 1H,  $J_1 = 10.1$  Hz,  $J_2 = 17.0$  Hz), 6.22 (dd, 1H,  $J_1 = 1.9$  Hz,  $J_2 = 17.0$  Hz), 6.09 (d, 1H,  $J_1 = 0.7$  Hz), 5.69 (dd, 1H,  $J_1 = 10.1$  Hz,  $J_2 = 1.9$  Hz) ppm.
- (e) 3-[6-(3-Methylphenylamino)-pyrimidin-4-ylamino]phenylamine and acryolyl chloride gave *N*-{3-[6-(3-methylphenylamino)-pyrimidin-4-ylamino]phenyl}-2-propenamide (**I-4**). MS (m/z):  $M+H^+$  = 346.  $^1H$  NMR (DMSO): 9.11 (s, 1H), 9.00 (s, 1H), 8.21 (s, 1H), 7.86 (d, 1H,  $J = 1.7$  Hz), 7.31-7.05 (m, 7H), 6.74 (d, 1H,  $J = 7.6$  Hz), 6.41 (dd, 1H,  $J_1 = 10.0$  Hz,  $J_2 = 17.0$  Hz), 6.20 (dd, 1H,  $J_1 = 2.0$  Hz,  $J_2 = 17.0$  Hz), 6.14 (s, 1H), 5.69 (dd, 1H,  $J_1 = 10.1$  Hz,  $J_2 = 2.0$  Hz), 2.23 (s, 3H) ppm.
- (f) 3-{6-[3-Chloro-4-(3-fluorophenyl)methoxyphenylamino]-pyrimidin-4-ylamino}phenylamine (**I<sup>R</sup>-6**) and acryolyl chloride gave *N*-{3-[6-(3-chloro-4-(3-fluorophenyl)methoxyphenylamino)-pyrimidin-4-ylamino]phenyl}-2-propenamide (**II-6**). MS (m/z):  $M+H^+$  = 490, 492.  $^1H$  NMR (DMSO): 9.13 (s, 1H), 9.07 (s, 1H), 8.22 (s, 1H), 7.85 (d, 1H,  $J = 1.5$  Hz), 7.40-7.10 (m, 10H), 6.41 (dd, 1H,  $J_1 = 10.0$  Hz,  $J_2 = 17.0$  Hz),

6.20 (dd, 1H,  $J_1 = 2.0$  Hz,  $J_2 = 17.0$  Hz), 6.05 (s, 1H), 5.70 (dd, 1H,  $J_1 = 10.1$  Hz,  $J_2 = 2.0$  Hz), 5.14 (s, 2H) ppm.

- (g) 4-Phenoxyphenyl amine gave 3-[6-(4-phenoxyphenylamino)-pyrimidin-4-ylamino]phenylamine. (**I<sup>R</sup>-7**) and acryloyl chloride gave *N*-{3-[6-(4-phenoxyphenylamino)-pyrimidin-4-ylamino]phenyl}-2 propenamide (**I-13**). MS (m/z):  $MH^+ = 424$ .  $^1H$  NMR (DMSO): 9.14 (s, 1H), 9.10 (s, 1H), 8.22 (s, 1H), 7.89 (s, 1H), 7.52 (d, 2H,  $J = 9.0$  Hz), 7.35 – 6.92 (m, 11H), 6.42 (dd, 1H,  $J_1 = 10.1$  Hz,  $J_2 = 16.9$  Hz), 6.22 (dd, 1H,  $J_1 = 1.9$  Hz,  $J_2 = 16.9$  Hz), 6.12 (s, 1H), 5.70 (dd, 1H,  $J_1 = 1.9$  Hz,  $J_2 = 10.1$  Hz) ppm.
- (h) 3-[6-(*N*-methyl-*N*-phenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-8**) and acryoyl chloride gave *N*-{3-[6-(*N*-methyl-*N*-phenylamino)-pyrimidin-4-ylamino]phenyl}-2 propenamide (**I-15**): an off-white solid; 70mg; 20%; MS (m/z):  $MH^+ = 346$ .  $^1H$  NMR (DMSO): 10.02 (s, 1H), 9.00 (s, 1H), 8.25 (s, 1H), 7.85 (s, 1H), 7.45 – 7.12 (m, 6H), 6.45 (dd, 1H), 6.20 (d, 1H), 5.70 (m, 1H), 3.35 (s, 3H) ppm.
- (i) 3-{[6-(3-Chloro-4-(2-pyridyl)methoxyphenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-9**) and acryloyl chloride gave *N*-(3-(6-(3-chloro-4-(pyridin-2-ylmethoxy)phenylamino)pyrimidin-4-ylamino)phenyl)-2-propenamide (**I-16**). MS (m/z):  $MH^+ = 473, 475(3:1)$ .  $^1H$  NMR (DMSO): 10.10 (s, 1H), 9.17 (s, 1H), 9.08 (s, 1H), 8.55 (m, 1H), 8.24 (s, 1H), 7.92 – 7.73 (m, 4H), 7.57 (d, 1H), 7.38 – 7.06 (m, 5H), 6.45 (dd, 1H), 6.23 (dd, 1H), 6.08 (s, 1H), 5.72 (dd, 1H), 5.20 (s, 2H) ppm.
- (j) 3-[6-(3-Chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**) and 1-cyanocyclopropanecarbonyl chloride gave *N*-[3-(6-{3-chloro-4-fluorophenylamino}pyrimidin-4-yl)amino]phenyl-1-cyanocyclopropanecarboxamide (**I-47**). MS (m/z):  $M+1 = 423/425$ ,  $^1H$  NMR (DMSO- $d_6$ )  $\delta$  10.04 (s, 1H), 9.18 (s, 1H), 9.12 (s, 1H), 8.27 (d, 1H), 7.88 (s, 1H), 7.8 (s, 1H), 7.47-7.16 (m, 9H), 6.74 (b, 1H), 6.28 (d, 1H), 6.1 (s, 1H), 5.19 (s, 2H), 3.05 (s, 2H), 2.18 (s, 6H).
- (k) 3-[6-(3-Chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**) and chloroacetyl chloride gave 2-chloro-*N*-{3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]-phenyl}-acetamide (**I-49**). MS (ES $^+$ ):  $(M + 1)^+ = 406$  (100%),  $(M + 3)^+ =$


- 408 (75%). <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, δ 10.31 (s, 1H), 9.37 (s, 1H), 9.30 (s, 1H), 8.33 (s, 1H), 8.00 (m, 1H), 7.87 (s, 1H), 7.47-7.24 (m, 5H), 6.15 (s, 1H), 4.26 (s, 2H).
- (l) 3-[6-(3-Chloro-4-fluorophenyl)aminopyrimidin-4-yl]aminophenylamine (**I<sup>R</sup>-4**) and 2-chloropropionyl chloride gave 2-chloro-*N*-[3-(6-{3-chloro-4-fluorophenyl}aminopyrimidin-4-yl)aminophenyl]propionamide (**I-50**). MS (ES<sup>+</sup>): (M + 1)<sup>+</sup> = 420 (100%), (M + 3)<sup>+</sup> = 422 (75%). <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, δ 10.32 (s, 1H), 9.37 (s, 1H), 9.28 (s, 1H), 8.33 (s, 1H), 8.00-7.98 (m, 1H), 7.87 (s, 1H), 7.47-7.26 (m, 5H), 6.14 (s, 1H), 4.69 (quartet, 1H), 1.61 (d, 3H).
- (m) 3-[6-(3-Chloro-4-fluorophenylamino)pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**) and 1-trifluoromethylcyclopropanecarbonyl chloride gave *N*-[3-(6-{3-chloro-4-fluorophenyl}aminopyrimidin-4-yl)aminophenyl]-1-trifluoromethylcyclopropanecarboxamide (**I-51**). MS (ES) (m/z): 466/468 [M+1, 100/45%]. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.61 (m, 1H), 9.07-9.18 (m, 2H), 8.14 (m, 1H), 7.63-7.8 (m, 2H), 7.03-7.21 (m, 5H), 5.94 (bs, 1H), 1.12-1.27 (m, 4H).
- (n) *N*-3-[6-(3-Chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl-*N*-methylamine (**Int-D**) and acryloyl chloride gave *N*-3-[6-(3-chloro-4-fluorophenylamino)pyrimidin-4-yl]aminophenyl-*N*-methyl-2-propenamide (**I-53**). MS (m/z): 398/400 (M+1, 100/63%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 10.32 (s, 1H), 9.21 (s, 1H), 8.33 (s, 1H), 7.99 (bs, 1H), 7.25-7.68 (m, 5H), 7.06 (bs, 1H), 6.41-6.47 (m, 1H), 6.26-6.29 (m, 1H), 5.71-5.80 (m, 2H).
- (o) 3-[6-(4-Bromo-2-fluorophenyl)aminopyrimidin-4-yl]aminophenyl amine (**I<sup>R</sup>-17**) and acryloyl chloride gave *N*-3-[6-(4-bromo-2-fluorophenylamino)pyrimidin-4-yl]aminophenyl]-2-propenamide (**I-55**). MS (M + H<sup>+</sup>) 430, 428. <sup>1</sup>H NMR (d<sup>6</sup>-DMSO) δ 10.13 (s, 1H), 9.25 (s, 1H), 9.02 (s, 1H), 8.26 (s, 1H), 7.91 (m, 2H), 7.57 (d, J = 11 Hz, 1H), 7.45-7.15 (m, 4H), 6.46 (dd, J = 12 and 10 Hz, 1H), 6.26 (d, J = 12 Hz, 1H), 6.23 (s, 1H), 5.75 (d, J = 10 Hz, 1H).
- (p) 2-[6-(4-phenoxyphenyl)amino-pyrimidin-4-yl]amino-6-aminopyridine (**I<sup>R</sup>-11**) and acryloyl chloride gave *N*-6-[6-(4-phenoxyphenyl)amino-pyrimidin-4-yl]aminopyridin-2-ylpropenamide (**I-63**) <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 5.75 (d, J = 10.20 Hz, 1H), 6.29 (d, J = 17.04 Hz, 1H), 6.62 (dd, J = 10.08 & 16.92 Hz, 1H), 6.96-7.00 (m, 4H), 7.07-7.11 (m, 1H), 7.18 (bd, J = 3.28 Hz, 1H), 7.33-7.38 (m, 3H), 7.62-7.69 (m, 4H), 8.29 (s, 1H), 9.13 (s, 1H), 9.66 (s, 1H), 10.15 (s, 1H); MS: m/z 425.3 (M+1).

- (q) 2-[6-(3-methylphenoxy)-pyrimidin-4-yl]amino-6-aminopyridine (**I<sup>R</sup>-12**) and acryloyl chloride gave *N*-6-[6-(3-methylphenoxy)-pyrimidin-4-yl]aminopyridin-2-ylpropenamide (**I-65**) <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ ppm: 2.40 (s, 3H), 5.86 (d, *J* = 9.44 Hz, 1H), 6.49-6.61 (m, 2H), 6.83 (bs, 1H), 6.95-7.05 (m, 3H), 7.15 (d, *J* = 7.48 Hz, 1H), 7.34 (t, *J* = 15.08 Hz, 2H), 7.54 (bd, *J* = 5.56 Hz, 1H), 8.78 (s, 1H), 10.78 (s, 1H); LCMS: *m/z* 348.8 (M+1).
- (r) 2-[6-(4-phenoxyphenoxy)-pyrimidin-4-yl]amino-6-aminopyridine (**I<sup>R</sup>-13**) and acryloyl chloride gave *N*-6-[6-(4-phenoxyphenoxy)-pyrimidin-4-yl]aminopyridin-2-ylpropenamide (**I-66**) <sup>1</sup>H NMR (MeOD) δ ppm: 5.92 (dd, *J* = 11.60, Hz, 1H), 6.50-6.54 (m, 2H), 6.75-7.28 (m, 10H), 7.37-7.41 (m, 2H), 7.91 (t, *J* = 16.08 Hz, 1H), 8.63 (s, 1H); LCMS: *m/z* 426 (M+1).
- (s) 3-[6-(phenylamino)-pyrimidin-4-ylamino]phenylamine. (**I<sup>R</sup>-15**) acryloyl chloride gave *N*-3-[6-(phenylamino)-pyrimidin-4-yl]aminophenylpropenamide (**I-67**) <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm: 5.72 (dd, *J* = 2.04 & 10.04 Hz, 1H), 6.19 (s, 1H), 6.25 (dd, *J* = 2 & 16.92 Hz, 1H), 6.46 (dd, *J* = 10.04 & 16.88 Hz, 1H), 6.96 (t, *J* = 7.36 Hz, 1H), 7.19-7.31 (m, 4H), 7.54 (d, *J* = 7.68 Hz, 2H), 7.91 (s, 1H), 8.26 (s, 1H), 9.13 (s, 1H), 9.18 (s, 1H), 10.11 (s, 1H); MS: *m/z* 332.8 (M+1).

## EXAMPLE 4


## SCHEME 4a

Sequence A for Synthesis of *N*-3-(6-mono or disubstituted-aminopyrimidin-4-yl)amino-*N*-mono- or unsubstituted-phenyl-4-amino-substituted-2-butenamides



## SCHEME 4b

Sequence B for Synthesis of *N*-3-(6-mono or disubstituted-aminopyrimidin-4-yl)amino-*N*-mono- or unsubstituted-phenyl-4-amino-substituted-2-butenamides



[00269] Synthesis of (*E*)-*N*-(3-(6-(3-chloro-4-(pyridin-2-ylmethoxy)phenylamino)-pyrimidin-4-ylamino)phenyl)-4-(dimethylamino)but-2-enamide (I-19). 3-{[6-(3-chloro-4-(2-pyridyl)methoxyphenylamino)-pyrimidin-4-ylamino]phenylamine (I<sup>R</sup>-9) was dissolved in *N*-methylpyrrolidinone (1.2 mL) and added dropwise over 10 minutes to the ice-cold solution of (*E*)-4-(dimethylamino)but-2-enoyl chloride hydrochloride in acetonitrile. The reaction was stirred in an ice bath for 2 hr. To the mixture was added sodium bicarbonate to pH greater than 9. The oil that formed was extracted with EtOAc (3 x 25 mL). A portion of the material was insoluble and set aside. The organic layer was dried (MgSO<sub>4</sub>) filtered and evaporated to a dark red oil. Both oils contained substantial amount of product as shown by TLC: SiO<sub>2</sub> CHCl<sub>3</sub>:MeOH/NH<sub>4</sub>OH (8:1) 19:1. The oils were combined and purified by flash column chromatography silica gel (25X300 mm) eluted first with 5% methanol/ammonium hydroxide 8/1 to remove non-polar impurities followed by 10% methanol/ammonium hydroxide 8/1 to elute 59 mg of product. The sample was further purified by a second flash column chromatography silica gel (25x250 mm) eluted with 10% methanol/ammonium hydroxide 8/1 to give the title compound (16.3 mg, 0.03 mmol, 6.4% yield). MS (ES<sup>+</sup>) 530(M<sup>+</sup>): 552, (M+Na); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>, 500 MHz) δ (ppm): 10.04 (s, 1H), 9.19 (s, 1H), 9.14(s, 1H), 8.60 (s, 1H), 8.27 (s, 1H), 7.88 (s, 2H), 7.57 (s, 1H), 7.36 (d, 1H, *J* = 7.4 Hz), 7.29 (d, 2H, *J* = 7.8 Hz), 7.22 (d, 2H, *J* = 7.9 Hz), 7.18 (m, 2H), 6.74 (m, 1H), 6.30 (d, 1H, *J* = 15 Hz) 6.10 (s, 1H), 5.24 (s, 2H), 3.06 (s, 2H), 2.18 (s, 6H) ppm; HPLC: *t*<sub>R</sub> = 5.15 min, 97.5% (YMC-Pack ODS-A 4.6x100 mm, 80% water/20% acetonitrile to 5% water/95% acetonitrile over 5.5 min, hold to 9 min.).

[00270] Synthesis of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-bromo-2-butenamide (Int-D) To a stirring solution of 4-bromo-but-2-enoic acid (0.72 g) at 0 °C under nitrogen atmosphere and triethylamine (0.61 mL, 4.38 mmol) in 5 mL of THF was added *iso*-butyl chloroformate (0.56 mL, 4.32 mmol). The mixture was stirred for 15 min followed by dropwise addition of a solution of *N*-(3-amino-phenyl)-*N*'-3-methylphenylamino-

pyrimidine-4,6-diamine (1.015 g, 3.483 mmol) in 50 mL of THF. The reaction was allowed to warm to room temperature overnight. The sample was concentrated then partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed with brine solution, dried (MgSO<sub>4</sub>), filtered and was concentrated. The resultant brown foamy solid was washed with diethyl ether and vacuum dried to give 0.960 g of crude (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-bromo-2-butenamide (**Int-D**) as brick-brown solid. MS (*m/z*): (*M*+1) 438/440 (71/75%).

[00271] **Synthesis of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-(methyl-prop-2-ynyl)amino-2-butenamide (I-58)** To a stirring solution at 0 °C under N<sub>2</sub> of *N*-[3-(6-{3-methylphenyl}amino-pyrimidin-4-ylamino)-4-bromo-2-butenamide (**Int-D**) (0.492 g, 1.122 mmol) and triethylamine (0.20 mL, 1.44 mmol) in 10 mL of THF was added (via syringe) *N*-methyl-propargylamine (0.11 mL, 1.173 mmol). The solution was allowed to warm to room temperature overnight, was concentrated and was then partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed with brine solution, was dried (MgSO<sub>4</sub>), was filtered and was concentrated. The residue was chromatographed (silica gel, 10% MeOH in CHCl<sub>3</sub>) to give 0.110 g of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-(methyl-prop-2-ynyl)amino-2-butenamide (**I-58**). MS (APCI) *m/z* 427 (*M*+1, 100%). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.06 (s, 1H), 9.07 (s, 1H), 9.17 (s, 1H), 8.27 (s, 1H), 7.90 (s, 1H), 7.16-7.36 (m, 6H), 6.70-6.73 (m, 1H), 6.79 (d, 1H), 6.20 (s, 1H), 6.32 (d, 1H), 3.30-3.49 (m), 3.14-3.27 (m, 3H), 2.18-2.39 (m, 7H which contain singlets at δ 2.25 [3H] and δ 2.29 [3H]).

[00272] **Synthesis of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-piperazinyl-2-butenamide (I-57)** To a stirring solution at 0 °C under N<sub>2</sub> of *N*-[3-(6-{3-methylphenyl}amino-pyrimidin-4-ylamino)-4-bromo-2-butenamide (**Int-D**) (2.15 g, 4.91 mmol) and triethylamine (0.86 mL, 6.17 mmol) in 10 mL of THF was added dropwise a solution of 1-Boc-piperazine (0.92 g, 4.91 mmol) in 10 mL of THF. The solution was allowed to warm to room temperature overnight, was concentrated and was then partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed with brine solution, was dried (MgSO<sub>4</sub>), was filtered and was concentrated to give 2.76 g of 4-{3-[3-(6-{3-methylphenyl}amino-pyrimidin-4-ylamino)-phenylcarbonyl]-allyl}-piperazine-1-carboxylic acid *tert*-butyl ester as gummy brown solid. This material was dissolved in 100 mL of CH<sub>2</sub>Cl<sub>2</sub>. 20 mL of trifluoroacetic acid was added and the mixture was stirred at room temperature under N<sub>2</sub> for 2 h. The mixture was

concentrated in vacuo, was basified with sat. NaHCO<sub>3</sub> solution and was extracted with EtOAc (2x 100 ml). The combined organic extract was dried (MgSO<sub>4</sub>), was filtered and was concentrated in vacuo. The residue was chromatographed (silica gel, 5% MeOH in CHCl<sub>3</sub> [500 mL] then 1% NH<sub>4</sub>OH-10% MeOH in CHCl<sub>3</sub>) to give 0.404 g of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-piperazinyl-2-butenamide. (**I-57**) MS (*m/z*): 444 (*M*+1, 100%). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 10.03 (s, 1H), 9.17 (s, 1H), 9.07 (s, 1H), 8.27 (s, 1H), 7.90 (s, 1H), 7.17-7.36 (m, 6H), 6.79-6.80 (d, 1H), 6.72-6.75 (m, 1H), 6.29 (d, 1H), 6.20 (s, 1H), 3.08-3.39 (m, containing water and ~3H), 2.70-2.72 (m, 4H), 2.30-2.42 (m, 4H), 2.29 (s, 3H).

[00273] The following compounds were prepared in a manner substantially similar to those described in Scheme 4a, above:

- (a) 3-(6-[3-chloro-4-{3-fluorobenzyloxy}]phenylamino-pyrimidin-4-yl)aminophenylamine (**I<sup>R</sup>-6**) and 4-dimethylamino-2-butenoyl chloride gave (*E*)-*N*-3-([6-(3-chloro-4-{3-fluorobenzyloxy}]phenylamino)-pyrimidin-4-yl)aminophenyl-4-(dimethylamino)but-2-enamide (**I-46**). MS (*m/z*): *M*<sup>+</sup> = 547, 549(3:1), H-NMR (DMSO-*d*<sub>6</sub>) δ 10.04 (s, 1H), 9.18 (s, 1H), 9.12 (s, 1H), 8.27 (d, 1H), 7.88 (s, 1H), 7.8 (s, 1H), 7.47-7.16 (m, 9H), 6.74 (b, 1H), 6.28 (d, 1H), 6.1 (s, 1H), 5.19 (s, 2H), 3.05 (s, 2H), 2.18 (s, 6H).
- (b) 3-[6-(3-methylphenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-5**) and 4-(dimethylamino)-2-butenoyl chloride gave (*E*)-*N*-{3-[6-(3-methylphenylamino)-pyrimidin-4-ylamino]phenyl}-4-dimethylamino-2-butenamide (**I-17**). MS (*m/z*): *MH*<sup>+</sup> = 403.
- (c) 3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**) and 4-(dimethylamino)-2-butenoyl chloride gave (*E*)-*N*-{3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl}-4-dimethylamino-2-butenamide (**I-18**). MS (*M* + *H*<sup>+</sup>) 441, 443; <sup>1</sup>H NMR (400 MHz, *d*<sup>6</sup>-DMSO) δ 10.01 (s, 1H), 9.30 (s, 1H), 9.20 (s, 1H), 8.28 (s, 1H), 7.95 (dd, *J* = 7 and 3 Hz, 1H), 7.86 (s, 1H), 7.41 (m, 1H), 7.35-7.10 (m, 3H), 6.69 (dt, *J* = 15 and 5 Hz, 1H), 6.26 (d, *J* = 15 Hz, 1H), 6.11 (s, 1H), 3.02 (d, *J* = 5 Hz, 2H), 2.14 (s, 6H) ppm.
- (d) *N*-3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl-*N*-methylamine (**Int-C**) and 4-dimethylamino-2-butenoyl chloride gave (*E*)-*N*-(3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl)-*N*-methyl-4-(dimethylamino)but-2-


enamide (**I-48**) MS (ES) ( $m/z$ ): 455/457 ( $M+1$ , 37/13%) and 228 (100%).  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  10.22 (s, 1H), 9.20 (s, 1H), 8.33 (s, 1H), 7.99 (bs, 1H), 7.25-7.68 (m, 5H), 7.05 (bs, 1H), 6.73-6.76 (m, 1H), 6.26-6.29 (m, 1H), 5.71 (s, 1H), 3.06 (bs, 2H), 1.99 (s, 6H).

- (e) 3-[6-(4-phenoxyphenylamino)-pyrimidin-4-yl]aminophenylamine (**I<sup>R</sup>-7**) and 4-dimethylamino-2-butenoyl chloride gave (*E*)-*N*-3-[6-(4-phenoxyphenyl)amino-pyrimidin-4-yl]aminophenyl-4-(dimethylamino)but-2-enamide (**I-62**)  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  ppm: 2.19 (s, 6H), 3.07 (d,  $J = 5.36$  Hz, 2H), 6.16 (s, 1H), 6.29 (d,  $J = 15.4$  Hz, 1H), 6.68-6.75 (m, 1H), 6.95-6.98 (m, 4H), 7.06-7.10 (m, 1H), 7.20 (dd,  $J = 7.88$  & 8.12 Hz, 1H), 7.24 (d,  $J = 8.32$  Hz, 2H), 7.36 (dd,  $J = 7.52$  & 7.84 Hz, 2H), 7.55 (d,  $J = 8.76$  Hz, 2H), 7.90 (s, 1H), 8.24 (s, 1H), 9.15 (d,  $J = 8.84$  Hz, 2H), 10.04 (s, 1H); LCMS:  $m/z$  481 ( $M+1$ ).

## EXAMPLE 5

### SCHEME 5a

Synthesis of *N*-{3-[6-(arylamino)-pyrimidin-4-ylamino]-phenyl}-ethensulfonamides



[00274] wherein  $L'$  is a subset of  $L$ , as defined herein, such that  $Y-L'SO_2Cl$  results in formation of provided compounds wherein  $R^1$  is  $-L-Y$  wherein a terminal methylene unit of  $L$  is replaced with  $-NHSO_2-$ .

[00275] **Synthesis of *N*-{3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl}-ethenesulfonamide (**I-3**)** A solution of 3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-4**) (300 mg, 0.9 mmol) and triethylamine (500 mg, 5 mmol) in 10 mL of THF was stirred at RT. 2-Chloroethanesulfonyl chloride (360 mg, 2.25 mmol) was added into the reaction mixture and stirring was continued at RT for 1 hr. The crude product was purified by flash chromatography on silica gel with EtOAc/heptane solvent system to afford 35 mg (9%) of the title compound, a brown colored solid. MS ( $m/z$ ):  $MH^+ = 420, 422$ .

$^1\text{H}$  NMR (DMSO): 9.92 (s, 1H), 9.29 (s, 1H), 9.21 (s, 1H), 8.26 (s, 1H), 7.92 (m, 1H), 7.40-7.22 (m, 4H), 7.14 (m, 1H), 6.20 (m, 2H), 6.05 (m, 3H) ppm. .


[00276] The following compounds were prepared in a manner substantially similar to Schemes 5a:

- (a) 3-{6-[3-chloro-4-(3-fluorophenyl)methoxyphenylamino]-pyrimidin-4-ylamino}phenylamine (**I<sup>R</sup>-6**) gave *N*-{3-[6-(3-chloro-4-(3-fluorophenyl)methoxyphenylamino)-pyrimidin-4-ylamino]phenyl}-ethanesulfonamide (**I-7**). MS (*m/z*):  $\text{M}+\text{H}^+$  = 526, 528 (2:1). MS (*m/z*):  $\text{M}+\text{H}^+$  = 490, 492 (2:1).  $^1\text{H}$  NMR (DMSO): 9.91 (s, 1H), 9.16 (s, 1H), 9.07 (s, 1H), 8.22 (s, 1H), 7.73 (s, 1H), 7.40-7.10 (m, 9H), 6.70 (m, 2H), 6.12 (d, 1H,  $J = 17.0$  Hz), 6.02 (m, 2H), 5.14 (s, 2H) ppm.
- (b) 3-[6-(3-methylphenylamino)-pyrimidin-4-ylamino]phenylamine (**I<sup>R</sup>-5**) gave *N*-{3-[6-(3-methylphenylamino)-pyrimidin-4-ylamino]phenyl}-ethanesulfonamide (**II-5**). MS (*m/z*):  $\text{M}+\text{H}^+$  = 382.  $^1\text{H}$  NMR (DMSO): 9.90 (s, 1H), 9.12 (s, 1H), 9.00 (s, 1H), 8.21 (s, 1H), 7.37 (d, 1H,  $J = 1.7$  Hz), 7.26 (m, 3H), 7.13 (m, 2H), 6.70 (m, 3H), 6.13 (m, 2H), 6.00 (d, 1H,  $J = 10.0$  Hz), 2.24 (s, 3H) ppm.

### EXAMPLE 6

#### SCHEME 6a

Synthesis of *N*<sup>2</sup>-acylated 6-(6-mono or disubstituted-aminopyrimidin-4-yl)amino-2-aminopyridines



[00277] **Synthesis of *N*-(6-chloro-pyrimidin-4-yl)-pyridine-2,6-diamine** A mixture of 2,6-diaminopyridine (1.530 g, 14.020 mmol) and 4,6-dichloropyrimidine (2.610 g, 17.519 mmol) in 15 mL of *n*-butanol in a sealed vial was heated at 100 °C for 72 h. The dark brown sample was cooled, was concentrated to remove most of the *n*-butanol, and was then partitioned between

EtOAc and sat. NaHCO<sub>3</sub> solution. An emulsion formed, the sample was filtered through a pad of Celite and the layers were separated. The organic extract was washed with sat. KH<sub>2</sub>PO<sub>4</sub> and brine solutions, was dried (MgSO<sub>4</sub>), was filtered and was concentrated to brown oily-solid. The sample was suspended into 50 mL of CH<sub>2</sub>Cl<sub>2</sub>, was cooled and was filtered to give 1.017 g of *N*-(6-chloro-pyrimidin-4-yl)-pyridine-2,6-diamine as yellow-orange solid. MS (ES) (m/z) 222/224 (M+1, 100/63%). TLC (SiO<sub>2</sub>, 50% EtOAc in hexanes): R<sub>f</sub> 0.33.

**[00278] Synthesis of *N*-(6-amino-pyridin-2-yl)-*N'*-(3-chloro-4-fluorophenyl)-pyrimidine-4,6-diamine (Int-E)** A mixture of *N*-(6-chloro-pyrimidin-4-yl)-pyridine-2,6-diamine (1.000 g, 4.512 mmol) and 3-chloro-4-fluoroaniline (1.380 mmol) in 10 mL of n-butanol in a sealed vial was heated at 120 °C for 24 h. The sample was cooled, was concentrated to remove most of the n-butanol, and was then diluted with EtOAc and sat. NaHCO<sub>3</sub> solution. The sample was stirred at room temperature for 30 min and was filtered. The solid was washed with fresh EtOAc and water and vacuum dried to give 1.063 g of *N*-(6-amino-pyridin-2-yl)-*N'*-(3-chloro-4-fluorophenyl)-pyrimidine-4,6-diamine (**Int-E**) as tan solid. MS (ES) (m/z) 331/333 (M+1, 100/65%). TLC (SiO<sub>2</sub>, 10% MeOH in CHCl<sub>3</sub>): R<sub>f</sub> 0.25.

**[00279] Synthesis of (*E*)-*N*-[6-(3-chloro-4-fluorophenylamino)pyrimidin-4-ylaminopyridin-2-yl]-4-(dimethylamino)but-2-enamide (I-52).**


To a stirring suspension at 0 °C under N<sub>2</sub> of 4-dimethylamino-but-2-enoic acid, hydrochloride (0.500 g, 3.019 mmol) in 15 mL of THF containing 5 drops of DMF was added drop-wise (via syringe) oxalyl chloride (0.28 mL, 3.210 mmol). Gas formation started immediately. The sample was stirred at 0 °C for ~30 min, room temperature for ~2 h, re-cooled to 0 °C, then treated with drop-wise addition of a solution of *N*-(6-amino-pyridin-2-yl)-*N'*-(3-chloro-4-fluorophenyl)-pyrimidine-4,6-diamine (**Int-E**) (0.500 g, 1.512 mmol) in 15 mL of THF and 3 mL of NMP. The ice-bath was removed, the sample was stirred at room temperature for 2 h then was partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed with brine solution, dried (MgSO<sub>4</sub>), filtered and concentrated to a yellow solid. The solid was suspended into EtOAc (~25 mL), stirred at room temperature for ~12 h, filtered and vacuum dried to give 0.459 g (69%) of (*E*)-*N*-[6-(3-chloro-4-fluorophenylamino)pyrimidin-4-ylaminopyridin-2-yl]-4-(dimethylamino)but-2-enamide (**I-52**) as light yellow solid. MS (m/z): 442/444 (M+1, 100/37%). <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 10.19 (s, 1H), 9.78 (s, 1H), 9.43 (s, 1H), 8.36

(s, 1H), 8.05-8.07 (m, 1H), 7.54-7.72 (m, 4H), 7.32-7.35 (m, 1H), 7.10 (bs, 1H), 6.79-6.82 (m, 1H), 6.54-6.57 (m, 1H), 3.14 (bs, 2H), 2.23 (bs, 6H).

### EXAMPLE 7

#### SCHEME 7a

#### Synthesis of *N*-acylated 3-(6-mono or disubstituted-aminopyrimidin-4-yl)amino-mono-substituted-phenylamines



[00280] **Synthesis of *N*-(5-Amino-2-methylphenyl)-*N'*-(3-chloro-4-fluorophenyl)pyrimidine-4,6-diamine (Int-F)** A mixture of 4-(3-chloro-4-fluorophenyl)-6-chloropyrimidin-4-ylamine (**I<sup>R</sup>-4**) (1.2 g, 4.6 mmol), 2-methyl-5-nitroaniline (0.85 g, 5.5 mmol) and 1 mL of concentrated HCl in 10 mL of n-butanol was heated at 120 °C for 16 h. With stirring, 5 mL of EtOAc was added to the reaction mixture. The light yellow colored product that precipitated was filtered and dried under vacuum to give *N*-(3-chloro-4-fluorophenyl)-*N'*-(2-methyl-5-nitrophenyl)pyrimidine-4,6-diamine. MS (APCI) (*m/z*): 374/376 (*M*+1). A mixture of *N*-(3-chloro-4-fluorophenyl)-*N'*-(2-methyl-5-nitrophenyl)pyrimidine-4,6-diamine (0.55 g, 1.5 mmol) and iron powder (35 mesh, 0.5 g, 5 eq) in 5 mL of HOAc and 2 mL of MeOH was heated at reflux for 2 h. The solvent was removed in vacuo and the dark colored residue was mixed with 150 mL of CH<sub>2</sub>Cl<sub>2</sub> and 15 mL of sat. K<sub>2</sub>CO<sub>3</sub> solution and was stirred at room temperature for 30 min. The organic layer was dried (MgSO<sub>4</sub>), was concentrated, and was then purified by flash chromatography (silica gel, MeOH/NH<sub>4</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>) to afford 0.115 g of *N*-(5-amino-2-methylphenyl)-*N'*-(3-chloro-4-fluorophenyl)pyrimidine-4,6-diamine as a light colored solid. (**Int-F**) MS (*m/z*): 344/346 (*M*+1).

[00281] In a manner substantially similar to that described above 4-(3-chloro-4-fluorophenylamino)-6-chloropyrimidine and 2-methoxy-5-nitroaniline gave *N*-(5-amino-2-methoxyphenyl)-*N'*-(3-chloro-4-fluorophenyl)pyrimidine-4,6-diamine. (Int-G) MS (m/z): 360/362 (M+1).

[00282] Synthesis of (*E*)-*N*-[3-(6-{3-Chloro-4-fluorophenyl}amino-pyrimidin-4-ylamino)-4-methylphenyl]-4-(dimethylamino)but-2-enamide (I-54) *N*-(5-amino-2-methylphenyl)-*N'*-(3-chloro-4-fluorophenyl)pyrimidine-4,6-diamine (Int-F) and 4-dimethylamino-2-butenoyl chloride were combined in a manner similar to that described in Scheme 4a to give (*E*)-*N*-[3-(6-{3-chloro-4-fluorophenyl}amino-pyrimidin-4-ylamino)-4-methylphenyl]-4-(dimethylamino)but-2-enamide (I-54). MS (m/z): 455/457 (M+1, 100/39%). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 11.02 (bs, 1H), 10.66 (bs, 1H), 9.92 (bs, 1H), 9.39 (bs, 1H), 7.92-7.94 (m, 1H), 7.75 (bs, 1H), 7.27-7.56 (m, 4H), 6.80 (m, 1H), 6.54 (d, 1H), 5.89 (bs, 1H), 3.92 (bs, 2H), 2.75 (bs, 6H), 2.17 (bs, 3H).

[00283] In a manner substantially similar to that described for the synthesis of (I-54) *N*-(5-Amino-2-methoxyphenyl)-*N'*-(3-chloro-4-fluorophenyl)pyrimidine-4,6-diamine (Int-G) and 4-dimethylamino-2-butenoyl chloride gave (*E*)-*N*-[3-(6-[3-chloro-4-fluorophenyl]-aminopyrimidin-4-ylamino-4-methoxyphenyl)-4-(dimethylamino)but-2-enamide (I-59). MS (m/z): 471/473 (M+1, 100/41%). <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ 9.97 (s, 1H), 9.28 (s, 1H), 8.46 (s, 1H), 8.26 (s, 1H), 7.93-7.98 (m, 2H), 7.44-7.51 (m, 2H), 7.30-7.33 (m, 1H), 7.02 (d, 1H), 6.69-6.72 (m, 1H), 6.26 (d, 1H), 6.03 (s, 1H), 3.80 (s, 3H), 3.05 (m, 2H), 2.17 (s, 6H).

#### EXAMPLE 8

[00284] Synthesis of 2-[6-(3-methylphenoxy)-pyrimidin-4-yl]amino-6-aminopyridine (I<sup>R</sup>-12)

A mixture of 2,6-diaminopyridine (1.530 g, 14.020 mmol) and 4,6-dichloropyrimidine (2.610 g, 17.519 mmol) in 15 mL of *n*-butanol in a sealed vial was heated at 100 °C for 72 h. The dark brown sample was cooled and was concentrated at reduced pressure to remove most of the *n*-butanol. The residue was then partitioned between EtOAc and saturated NaHCO<sub>3</sub> solution. An emulsion formed, the sample was filtered through a pad of Celite<sup>TM</sup> and the layers were separated. The organic extract was washed with sat. KH<sub>2</sub>PO<sub>4</sub> and brine solutions, dried (MgSO<sub>4</sub>), filtered and concentrated to brown oily-solid. The sample was suspended into ~50 mL of CH<sub>2</sub>Cl<sub>2</sub>, cooled and filtered to give 1.017 g (36%) of *N*-(6-chloro-pyrimidin-4-yl)-pyridine-2,6-diamine as

yellow-orange solid. MS:  $m/z$  222/224 ( $M+1$ , 100/63%). To a solution of *N*-(6-chloro-pyrimidin-4-yl)-pyridine-2,6-diamine (100 mg, 0.45 mmol) in DMF (1 mL) was added 3-methylphenol (88 mg, 0.8 mmol) and anhydrous  $K_2CO_3$  (93 mg, 0.6 mmol). The reaction mixture was heated at 145 °C for 16 h. It was then cooled and DMF was removed under reduced pressure to get a yellow gummy residue. The residue was taken in EtOAc (20 mL), was washed sequentially with water (5 mL) and brine (5 mL) and was dried over  $Na_2SO_4$ . Filtration followed by concentration under reduced pressure afforded a yellow gum which was further purified by column chromatography ( $SiO_2$ , 60-120, EtOAc/hexane, 5/5) to give 100 mg of 2-[6-(3-methylphenoxy)-pyrimidin-4-yl]amino-6-aminopyridine (**I<sup>R</sup>-12**) as a pale yellow solid. MS:  $m/z$  294 ( $M+H$ )

[00285] The following compounds were prepared in a manner substantially similar to the procedure described above:

- (a) 4-Phenoxyphenol and 2,6-diaminopyridine gave 2-[6-(4-phenoxyphenoxy)-pyrimidin-4-yl]amino-6-aminopyridine (**I<sup>R</sup>-13**) MS ( $m/z$ ):  $MH^+ = 372$ .
- (b) 4-nitrophenol and 1,3-diaminobenzene gave 3-[6-(4-nitrophenoxy)-pyrimidin-4-yl]amino-aminobenzene (**I<sup>R</sup>-16**) MS ( $m/z$ ):  $MH^+ = 324$

#### EXAMPLE 9

[00286] **Synthesis of *N*-3-[6-(3-ethynylphenylamino)-pyrimidin-4-yl]aminophenyl-2-propenamide (I-68)** To a stirred solution of **I<sup>R</sup>-1** (150 mg, 0.42 mmol) in dry DMF (4.0 mL) under  $N_2$  was added  $Pd(PPh_3)_2Cl_2$  (14.7 mg, 0.021 mmol), CuI (3.9 mg, 0.021 mmol),  $PPh_3$  (22.07 mg, 0.08 mmol), and diethylamine (94.8 mg, 6.3 mmol). The reaction mixture was purged with  $N_2$  for additional 10 min, trimethylsilylacetylene (45.5 mg, 0.46 mmol) was added, and it was then subjected to microwave irradiation at 120 °C for 30 min. The mixture was cooled, was diluted with 5 mL water, was filtered through celite® and was extracted with EtOAc (2x10 mL). The combined EtOAc extract was washed with water, then with brine and was dried over  $Na_2SO_4$ . Concentration under reduced pressure gave a crude product that was purified by column chromatography ( $SiO_2$ , MeOH/ $CHCl_3$ , 1/99) to give 100 mg of a brown solid. A solution of this material in 2 mL of dry methanol under nitrogen atmosphere containing anhydrous  $K_2CO_3$  (73.9 mg, 0.53 mmol) was stirred at rt for 16 h. The reaction mixture was filtered and the filtrate was concentrated under reduced pressure to give a residue, which was further purified by column

chromatography (SiO<sub>2</sub>, 230-400, MeOH/CHCl<sub>3</sub>, 1/99) to give 70 mg of a light brown solid. To a stirred solution of this material in NMP (1.0 mL) at 0 °C was added acryloyl chloride (105 mg, 1.16 mmol), and the reaction mixture was stirred at 0 °C for 1 h. The reaction mixture was then quenched with water, was basified with 10% NaHCO<sub>3</sub> sol and was extracted with EtOAc. The combined EtOAc extract was washed sequentially with water and brine, was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The residue obtained was further purified by preparative HPLC to give 8 mg of *N*-3-[6-(3-ethynylphenyl)aminopyrimidin-4-yl]amino-phenyl-2-propenamide (**I-68**) as an off white solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 4.15 (s, 1H), 5.73-5.76 (m, 1H), 6.19 (s, 1H), 6.25 (dd, *J* = 1.96 & 17.00 Hz, 2H), 6.46 (dd, *J* = 10.2 & 16.96 Hz, 1H), 7.05 (d, *J* = 7.64 Hz, 1H), 7.21-7.33 (m, 3H), 7.54 (d, *J* = 7.96 Hz, 1H), 7.81 (s, 1H), 7.91 (s, 1H), 8.32 (s, 1H), 9.28 (d, *J* = 11.08 Hz, 2H), 10.13 (s, 1H); MS: *m/z* 356.8 (M+1).

#### EXAMPLE 10

[00287] Synthesis of *N*-3-[6-(4-[4-[[4-chloro-3-(trifluoromethyl)phenyl]amino]-carbonyl]amino)phenoxy pyrimidin-4-yl]aminophenylchloroacetamide (**I-69**)

[00288] Step 1: To a solution of 3-[6-(4-nitrophenoxy)-pyrimidin-4-yl]amino-aminobenzene (**I<sup>R</sup>-16**) (650 mg, 2.01 mmol) in THF (10 mL) was added Et<sub>3</sub>N (305 mg, 3.01 mmol) and (Boc)<sub>2</sub>O (525 mg, 2.4 mmol) under N<sub>2</sub> atmosphere. The reaction mixture was further heated at 60 °C for 16 h. A residue was obtained after cooling to room temperature and removal of solvent under vacuum. The residue was dissolved in EtOAc (10 mL). The EtOAc extract was washed sequentially with water (5 mL) and brine (2 mL), was dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration under reduced pressure followed by purification by column chromatography (SiO<sub>2</sub>, 60-120, CHCl<sub>3</sub>/MeOH, 9/1) gave 400 mg of the Boc derivative as a yellow solid.

[00289] Step 2: The material from Step 1 was dissolved in MeOH (8 mL) and 10% Pd/C (40 mg) was added under N<sub>2</sub> atmosphere. The reaction mixture was hydrogenated in a Parr apparatus (H<sub>2</sub>, 3 Kg, rt, 16 h). The reaction mixture was filtered through celite® and the solvent was removed under reduced pressure to give 250 mg, of the amine as a yellow solid.

[00290] Step 3: To the material from Step 2 was added a toluene solution of 4-chloro-3-trifluoromethylphenylisocyanate prepared by reacting under nitrogen atmosphere at 0 °C 24 mg of 4-chloro-3-trifluoromethylaniline in 10 mL toluene with 0.08 mL of a 20%-solution of phosgene in toluene followed by addition of Et<sub>3</sub>N (0.07 mL) and at 110 °C for 16 h. The reaction

mixture of the amine and the isocyanate was further heated at 110 °C for 4 h and then was quenched with water (1 mL) and was extracted with EtOAc (2x20 mL). The EtOAc extract was washed with water (5 mL), brine (2 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration followed by concentration under vacuum offered a residue which was purified by a column chromatography (SiO<sub>2</sub>, 230-400, hexane/EtOAc, 9/1) to 20 mg of the Boc/urea intermediate as a yellow solid.

[00291] **Step 4:** To a solution of 45 mg of this intermediate in CH<sub>2</sub>Cl<sub>2</sub> (2 mL) at 0 °C was added TFA (0.01 mL) under N<sub>2</sub> atmosphere. The reaction mixture stirred at room temperature for 4 h, was washed sequentially with 10% NaHCO<sub>3</sub> solution (1 mL) and brine (1 mL) and was dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration under reduced pressure gave 25 mg of the amine/urea intermediate as a brownish solid.

[00292] **Step 5:** To a solution of the amine/urea intermediate from **Step 4** (45 mg, 0.05 mmol), in THF (2 mL) and Et<sub>3</sub>N (10 mg, 0.1 mmol) at 0 °C under N<sub>2</sub> atmosphere was added chloroacetyl chloride (55 mg, 0.1 mmol). The reaction mixture was allowed to come to rt and with stirring for 2 h. The reaction mixture was concentrated under reduced pressure and the residue was partitioned between EtOAc (4 mL) and water (1 mL). The EtOAc layer was separated, was washed with brine (2 mL) and was dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration followed by concentration under reduced pressure gave a residue which was further purified by column chromatography (SiO<sub>2</sub>, 230-400, CHCl<sub>3</sub>/MeOH, 9/1) to give *N*-3-[6-(4-[4-[[[4-chloro-3-(trifluoromethyl)phenyl]amino]carbonyl]amino]phenoxy]phenyl]amino-pyrimidin-4-yl]aminophenylchloroacetamide (**I-69**) as a pale yellow solid. <sup>1</sup>H NMR (MeOD) δ ppm: 4.19 (s, 2H), 6.61 (s, 1H), 7.13 (d, *J* = 6.92 Hz, 2H), 7.14-7.23 (m, 3H), 7.50-7.55 (m, 3H), 7.63-7.66 (m, 1H), 7.95 (s, 1H), 8.00 (s, 1H), 8.30 (s, 1H); MS: *m/z* 593 (M+1).

#### EXAMPLE 11

[00293] **Synthesis of (*E*)-*N*-3-(6-[3-methylphenylamino]-pyrimidin-4-yl)aminophenyl-4-(4-acetylpiperazin-1yl)-2-butenamide (**I-60**)**

To a stirring solution at 0 °C under N<sub>2</sub> of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-piperazinyl-2-butenamide (**I-57**) (0.291 g, 0.655 mmol) and triethylamine (0.14 mL, 1.004 mmol) in 10 mL of THF was added (via syringe) acetyl chloride (0.05 mL, 0.70 mmol). The sample was allowed to warm to room temperature overnight, was concentrated and was then partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed

with brine solution, was dried (MgSO<sub>4</sub>), was filtered, was concentrated and was chromatographed (silica gel, 1% NH<sub>4</sub>OH-10% MeOH in CHCl<sub>3</sub>) to give 0.0705 g of (*E*)-*N*-3-(6-[3-methylphenyl]aminopyrimidin-4-yl)aminophenyl-4-(4-acetylpiperazin-1-yl)-2-butenamide (**I-60**), a white solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 2.00 (s, 3H), 2.29 (s, 3H), 2.35-2.41 (m, 4H), 3.15-3.16 (m, 2H), 3.31-3.46 (m, containing water and ~ 4H), 6.19 (s, 1H), 6.32 (d, 1H), 6.73-6.80 (m, 2H), 7.16-7.35 (m, 6H), 7.90 (s, 1H), 8.27 (s, 1H), 9.07 (s, 1H), 9.17 (s, 1H) and 10.05 (s, 1H); MS: *m/z* 486 (M+1, 100%).


### EXAMPLE 12

#### [00294] Synthesis of (*E*)-*N*-(3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-ylamino]phenyl)-*N*-methyl-4-(dimethyl-d<sub>6</sub>-amino)but-2-enamide (**I-61**)

To a stirring solution of 4-bromo-but-2-enoic acid (0.28 g) at 0 °C under nitrogen atmosphere and triethylamine (0.25 mL) in 3 mL of THF was added *iso*-butyl chloroformate (0.22 mL). The mixture was stirred for 15 min followed by dropwise addition of a solution of 3-[6-(3-chloro-4-fluorophenyl)aminopyrimidine-4-yl]aminophenylamine (**I<sup>R</sup>-4**) (0.46 g) in 50 mL of THF. The reaction was allowed to warm to room temperature overnight. The sample was concentrated then partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed with brine solution, dried (MgSO<sub>4</sub>), filtered and was concentrated. The resultant brown foamy solid was washed with diethyl ether and vacuum dried to give 0.378 g of (*E*)-*N*-3-(6-[3-chloro-4-fluorophenylamino]pyrimidin-4-ylaminophenyl)-4-bromo-2-butenamide (**Int-E**). MS (*m/z*): 480, 478, 476 (25/100/80 %). To a stirring solution at 0 °C under N<sub>2</sub> of **Int-E** (0.378 g, 0.794 mmol) and triethylamine (0.28 mL, 2.01 mmol) in 10 mL of THF was added in one portion dimethyl-d<sub>6</sub>-amine hydrochloride (0.070 g, 0.799 mmol). The sample was allowed to warm to room temperature overnight and then was partitioned between EtOAc and sat. NaHCO<sub>3</sub> solution. The organic extract was washed with brine solution, was dried (MgSO<sub>4</sub>), was filtered and was concentrated. The residue was chromatographed (silica gel, 1% NH<sub>4</sub>OH-10% MeOH in CHCl<sub>3</sub>) to give 0.0582 g (16%) of (*E*)-*N*-(3-[6-(3-chloro-4-fluorophenylamino)-pyrimidin-4-yl]aminophenyl)-4-(dimethyl-d<sub>6</sub>-amino)but-2-enamide (**I-61**), a tan solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 3.05-3.07 (m, 2H), 6.16 (s, 1H), 6.29-6.32 (m, 1H), 6.72-6.75 (m, 1H), 7.21-7.47 (m, 5H), 7.90 (s, 1H), 7.92-8.01 (m, 1H), 8.32 (s, 1H), 9.26 (s, 1H), 9.36 (s, 1H) and 10.06 (s, 1H); MS: *m/z* 447/449 (M+1, 100/49%).

## EXAMPLE 13


[00295] **I-80** may be prepared in a manner substantially similar to that described in Scheme 4a, above:

**I-80**

4-[6-(4-phenoxyphenyl)amino-pyrimidin-4-yl]amino-aminobenzene (**I<sup>R</sup>-14**) and acryloyl chloride gave *N*-4-[6-(4-phenoxyphenyl)amino-pyrimidin-4-yl]aminophenylpropenamide (**I-80**), an off white solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm : 5.73 (dd, *J* = 1.6 & 10.0 Hz, 1H), 6.08 (d, *J* = 5.6 Hz, 1H), 6.24 (dd, *J* = 2 & 16.8 Hz, 1H), 6.43 (dd, *J* = 10 & 16.8 Hz, 1H), 6.95-6.70 (m, 4H), 7.09 (t, *J* = 7.6 Hz, 1H), 7.35-7.39 (m, 2H), 7.45-7.49 (m, 2H), 7.55-7.61 (m, 4H), 8.23 (s, 1H), 9.08 (s, 1H), 9.1 (s, 1H), 10.1 (s, 1H); LCMS : *m/e* 423.8 (M<sup>+</sup>).

## EXAMPLE 14

## SCHEME 14a



[00296] **Synthesis of *N*-6-(6-(phenylamino)pyrimidin-4-ylamino)pyridine-2-yl)acrylamide (I-72)**


**Step-1** A solution of *N*<sup>2</sup>-(6-chloropyrimidin-4-yl)pyridine-2,6-diamine (**1**) (0.25 g, 1.1 mmol) and aniline (**2**) (0.16 g, 1.7 mmol) in *n*-BuOH (25 mL) was heated at 120 °C for 12 h in a pressure tube. The reaction mixture was cooled, was dissolved in methanol (10 mL) and was

concentrated under reduced pressure. The residue was dissolved in ethyl acetate (35 mL) and was washed successively with 10% sodium bicarbonate solution (20 mL), water (20 mL), and saturated brine (20 mL). The ethyl acetate extract was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to give a residue, which was triturated with diethyl ether to give (260 mg, 83%) *N*<sup>4</sup>-(6-aminopyridin-2-yl)-*N*<sup>6</sup>-phenylpyrimidine-4,6-diamine (**3**) as an off-white solid.

**Step-2** To a stirred solution of **3** (0.2 g, 0.7 mmol) in NMP (10 mL) was added acryloyl chloride (0.097 g, 1 mmol), drop wise at 0 °C. The reaction mixture was allowed to stir at the same temperature for 20 min and then warmed to rt for 1.5 h. It was quenched with 10% sodium bicarbonate solution (4 mL) and was extracted with ethyl acetate (2×35 mL). The combined ethyl acetate layer was washed with water (20 mL), saturated brine (20 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The residue was further purified by column chromatography (SiO<sub>2</sub>, 60-120, Petroleum ether/EtOAc : 90/10) to give *N*-6-(6-(phenylamino)pyrimidin-4-ylamino)pyridine-2-yl)acrylamide (**I-72**) as a brown solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm : 5.80 (dd, *J* = 1.84 & 10.12 Hz, 1H), 6.31 (dd, *J* = 1.8 & 16.96 Hz, 1H), 6.64 (dd, *J* = 10.08 & 16.88 Hz, 1H), 6.96 (t, *J* = 7.32 Hz, 1H), 7.15-7.20 (m, 1H), 7.28 (t, *J* = 7.52 Hz, 2H), 7.34 (s, 1H), 7.60-7.70 (m, 4H), 8.30 (s, 1H), 9.08 (s, 1H), 9.66 (s, 1H), 10.06 (s, 1H); LCMS : *m/e* 332.6 (M<sup>+</sup>).

### EXAMPLE 15

#### SCHEME 15a




#### [00297] Synthesis of (*E*)-4-(dimethylamino)-*N*-(6-(6-(phenylamino)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-73**)

To a stirred solution of acetonitrile (20 mL) and DMF (0.05 mL) under N<sub>2</sub> was added *N,N*-dimethylamino crotonic acid hydrochloride (0.47 g, 2.8 mmol). After 10 min this solution was

cooled to 0-5 °C. Oxalyl chloride (0.44 g, 3.5 mmol) was added and the reaction mixture was maintained at 0-5 °C for 30 min. It was allowed to warm to rt and stirring was continued for 2 h. It was then heated to 40 °C for 5 min and again brought to rt and stirred for 10 min to get a light greenish colored solution of dimethylaminocrotonyl chloride that was used as such for next step. To a stirred solution of *N*<sup>4</sup>-(6-aminopyridin-2-yl)-*N*<sup>6</sup>-phenylpyrimidine-4,6-diamine (0.2 g, 0.7 mmol) in NMP (10 mL) was added dropwise under N<sub>2</sub> atmosphere at 0 °C the solution of dimethylaminocrotonyl chloride. The reaction mixture was maintained at this temperature for 30 min and was warmed to rt and stirred for 2 h. The mixture was quenched with sodium bicarbonate solution (1 mL) and was extracted with EtOAc (2×35 mL). The combined ethyl acetate extract was washed with water (20 mL) and brine (20 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The residue was purified by column chromatography (SiO<sub>2</sub>, 60-120, product was eluted at 4-6% methanol in chloroform) to give (*E*)-4-(dimethylamino)-*N*-(6-(6-(phenylamino)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-73**) as a yellowish solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm : 2.24 (s, 6H), 3.13 (d, *J* = 5.76 Hz, 2H), 6.56 (d, *J* = 15.52 Hz, 1H), 6.80 (d, *J* = 15.36 Hz, 1H), 6.95 (t, *J* = 7.36 Hz, 1H), 7.11 (t, *J* = 1.16 Hz, 1H), 7.29 (t, *J* = 7.56 Hz, 2H), 7.58 (s, 1H), 7.65-7.7 (m, 4H), 8.31 (d, *J* = 2.44 Hz, 1H), 9.23 (s, 1H), 9.68 (s, 1H), 10.16 (s, 1H); LCMS : *m/e* 390.3 (M+1).

## EXAMPLE 16

## SCHEME 16a




**[00298] Synthesis of (*E*)-4-(dimethylamino)-*N*-(6-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-64**)**

A solution of dimethylaminocrotonyl chloride was prepared by reaction of dimethylaminocrotonic acid hydrochloride (0.36 g, 2.16 mmol) in CH<sub>3</sub>CN (4 mL) containing DMF (1 drop) with oxalyl chloride (0.34 g, 2.70 mmol) according to the procedure in **Example 15a**. This acid chloride was added drop wise at 0 °C in to a stirred solution of *N*<sup>4</sup>-(6-

aminopyridin-2-yl)-*N*<sup>6</sup>-4-phenoxyphenylpyrimidine-4,6-diamine (**I<sup>R</sup>-11**) (0.2 g, 0.54 mmol) in NMP (8 mL). The reaction was stirred at 0 °C for 1 h, was diluted with EtOAc (5 mL), and was washed with 10% NaHCO<sub>3</sub> (2 mL), water (2 mL) and brine (2 mL). Drying over Na<sub>2</sub>SO<sub>4</sub> followed by concentration under reduced pressure offered a residue which was further purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to give (*E*)-4-(dimethylamino)-*N*-(6-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-64**) as a yellow solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm : 2.18 (s, 6H), 3.05 (d, *J* = 5.2 Hz, 2H), 6.48 (d, *J* = 15.2 Hz, 1H), 6.79 (d, *J* = 6 & 15.6 Hz, 1H), 6.96-6.99 (m, 4H), 7.07-7.15 (m, 2H), 7.36 (t, *J* = 7.6 Hz, 2H), 7.42 (s, 1H), 7.64-7.7 (m, 4H), 8.30 (s, 1H), 9.12 (s, 1H), 9.69 (s, 1H), 10.07 (s, 1H); LCMS : *m/e* 482.4 (M+1).

### EXAMPLE 17

#### SCHEME 17a



**[00299] Synthesis of (*E*)-4-(dimethylamino)-*N*-(6-(6-(3-methylphenoxy)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-74**)**

#### Step 1


To a solution of *N*<sup>2</sup>-(6-chloropyrimidin-4-yl)pyridine-2,6-diamine (100 mg, 0.45 mmol) in DMF (1 mL) was added 3-methylphenol (88 mg, 0.8 mmol) and anhydrous K<sub>2</sub>CO<sub>3</sub> (93 mg, 0.6 mmol). The reaction mixture was heated at 145 °C for 16 h. It was then cooled and DMF was removed under reduced pressure to get a yellow gummy residue. The residue was taken in EtOAc (20 mL) and washed with water (5 mL), brine (5 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration followed by concentration under reduced pressure afforded a yellow gum which was further purified by column chromatography (SiO<sub>2</sub>, 60-120, EtOAc/hexane, 5/5) to give *N*<sup>2</sup>-(6-(3-methylphenoxy)pyrimidin-4-yl)pyridine-2,6-diamine (**I<sup>R</sup>-12**) (100 mg, 75%) as a pale yellow solid.

#### Step 2

To a solution of **I<sup>R</sup>-12** (75 mg, 0.25 mmol) in NMP (2 mL) was added dimethylaminocrotonyl chloride (168 mg, 1.02 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 1 h and was then quenched with NaHCO<sub>3</sub> solution (2 mL). The mixture was extracted with EtOAc (3x5 mL) and the combined EtOAc extract was washed with water (5 mL), brine (5 mL) and was dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration under reduced pressure afforded a yellow oil, which was further purified by column chromatography (SiO<sub>2</sub>, 60-120, Chloroform/Methanol, 9/1) to give (*E*)-4-(dimethylamino)-N-(6-(6-(3-methylphenoxy)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-74**) as a light brown solid. <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ ppm: 2.35 (s, 6H), 2.38 (s, 3H), 3.25 (dd, *J* = 1.2 & 6.6 Hz, 2H), 6.40 (d, *J* = 13.16 Hz, 1H), 6.92-7.01 (m, 3H), 7.12 (t, *J* = 8.12 Hz, 2H), 7.34 (t, *J* = 7.84 Hz, 1H), 7.54 (s, 1H), 7.69 (t, *J* = 6.24 Hz, 1H), 7.79 (d, *J* = 8 Hz, 1H), 8.30 (s, 1H); LCMS: *m/e* 404.8 (M<sup>+</sup>).

## EXAMPLE 18

## SCHEME 18a



**[00300] Synthesis of *N*-(3-(6-(3-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)acrylamide (**I-75**)**

**Step 1**


A solution of 4,6-dichloropyrimidine (0.5 g, 3.3 mmol), 3-phenoxyaniline (0.75 g, 4 mmol) and DIPEA (0.65 g, 5 mmol) in *n*-butanol (5 mL) was subjected to microwave irradiation (110 °C, 30 min). The reaction mixture was cooled, concentrated under reduced pressure and the residue was dissolved in EtOAc (10 mL). This solution was washed with water (5 mL) and brine (5 mL), and was dried over Na<sub>2</sub>SO<sub>4</sub>. Concentrated under reduced pressure gave a residue that was purified by column chromatography (SiO<sub>2</sub>, 60-120, hexane/ethylacetate, 8/2) to give 6-chloro-*N*-(3-phenoxyphenyl)pyrimidine-4-amine (0.56 g, 56%) as an off-white solid.

**Step 2**

A solution of 6-chloro-*N*-(3-phenoxyphenyl)pyrimidine-4-amine (0.25 g, 0.8 mmol), 1,3-diaminobenzene (0.36 g, 3.3 mmol), *n*-butanol (10 mL) and conc. HCl (61 mg, 1.6 mmol) was subjected to microwave irradiation (160 °C, 15 min). The reaction mixture was cooled, was concentrated under reduced pressure and the residue was taken in EtOAc (10 mL). The EtOAc solution was washed with water (5 mL) and brine (5 mL), and was dried over Na<sub>2</sub>SO<sub>4</sub>. Concentration under reduced pressure gave a residue that was purified by column chromatography (SiO<sub>2</sub>, 60-120, hexane/ethyl acetate, 6/4) to give *N*<sup>4</sup>-(3-aminophenyl)-*N*<sup>6</sup>-(3-phenoxyphenyl)pyrimidine-4,6-diamine (0.1 g, 32%) as a brown solid.

**Step 3**

To a stirred solution of *N*<sup>4</sup>-(3-aminophenyl)-*N*<sup>6</sup>-(3-phenoxyphenyl)pyrimidine-4,6-diamine (40 mg, 0.1 mmol), Et<sub>3</sub>N (0.03 mL, 0.2 mmol) and NMP (0.4 mL) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL), at 0 °C, was added acryloyl chloride (**6**) (29 mg, 0.3 mmol). The reaction mixture was allowed to come to rt and stirred at this temperature for 3 h. It was washed with 10% NaHCO<sub>3</sub> solution (2 mL), water (2 mL), brine (2 mL) and dried over Na<sub>2</sub>SO<sub>4</sub>. Filtration followed by concentration under reduced pressure offered a residue which was purified by column chromatography (SiO<sub>2</sub>, 230-400, chloroform/methanol, 9/1) to gave *N*-(3-(6-(3-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)acrylamide (**I-75**) as a light brown solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm : 5.73 (dd, *J* = 1.72 & 10 Hz, 1H), 6.18 (s, 1H), 6.24 (dd, *J* = 1.92 & 17 Hz, 1H), 6.45 (dd, *J* = 10.04 & 16.88 Hz, 1H), 6.54-6.57 (m, 1H), 7.01-7.05 (m, 2H), 7.11-7.15 (dd, *J* = 0.88 & 7.48 Hz, 1H), 7.19-7.32 (s, 4H), 7.35-7.41 (m, 4H), 7.89 (s, 1H), 8.26 (s, 1H), 9.2 (s, 1H), 9.25 (s, 1H), 10.26 (s, 1H); LCMS : *m/e* 423.8 (M+1).


**EXAMPLE 19****SCHEME 19a**

[00301] Compound **I-76** can be prepared according to Scheme 19a by coupling intermediate **1** with 3-bromophenol, followed by boronic acid coupling of intermediates **3** and **4** to give

intermediate 5. Intermediate 5 can then be treated with acryloyl chloride using a protocol similar to that in Example 6 to give compound I-76.

### EXAMPLE 20


#### SCHEME 20a



[00302] Compound I-77 can be prepared according to Scheme 20a by treatment of intermediate 5 with (E)-4-(dimethylamino)but-2-enoyl chloride using a protocol similar to that in Example 6.

### EXAMPLE 21

#### SCHEME 21a



[00303] Synthesis of *N*-methyl-*N*-(6-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyridin-2-yl)acrylamide (I-78)

**Step 1**

To a solution of 2,6-diaminopyridine (10 g, 91.63 mmol) in dry THF (100 mL) was added  $K_2CO_3$  (18.8 g, 136.23 mmol) and  $CH_3I$  (13 g, 91.63 mmol) and the reaction mixture was stirred at room temperature for 16 h. Water was added (10 mL) and the mixture was extracted with EtOAc (100 mL). The EtOAc layer was dried and was concentrated under reduced pressure. The residue was further purified by column chromatography ( $SiO_2$ , 60-120, chloroform) to give 2-methylamino-6-aminopyridine (1.1 g, 10%) as a brown solid.

**Step 2**

A mixture of 2-methylamino-6-aminopyridine (0.5 g, 4.04 mmol), 4,6-dichloropyrimidine (1.51 g, 10.13 mmol), DIPEA (1.5 g, 12.17 mmol) in n-butanol (5 mL) was heated at 120 °C for 16 h. The reaction mixture was cooled, was concentrated under reduced pressure and the residue was taken in dichloromethane (25 mL). The dichloromethane solution was washed with  $NaHCO_3$  solution (2 mL), water (2 mL) and brine (2 mL), was dried over  $Na_2SO_4$  and was concentrated under reduced pressure. The residue was purified by column chromatography ( $SiO_2$ , 60-120, petroleum ether/ethyl acetate, 6/4) to give  $N^2$ -(6-chloropyrimidin-4-yl)- $N^6$ -methylpyridine-2,6,diamine (0.3 g, 33%) as a yellow solid.

**Step 3**

A solution of  $N^2$ -(6-chloropyrimidin-4-yl)- $N^6$ -methylpyridine-2,6,diamine (0.3 g 1.27 mmol), 4-phenoxy aniline (0.28 g, 1.52 mmol) and conc. HCl (2 drops) in n-butanol (2 mL) was subjected to microwave irradiation (120 °C, 1 h). The reaction mixture was cooled, was concentrated under reduced pressure, and the residue was diluted with  $CH_2Cl_2$  (5 mL). The dichloromethane solution was washed with  $NaHCO_3$  (2 mL), water (2 mL), and brine (2 mL), and was dried over  $Na_2SO_4$ . Filtration followed by concentration under reduced pressure gave a residue which was purified by column chromatography ( $SiO_2$ , 60-120, chloroform/methanol, 9/1) to give  $N^4$ -(6-(methylamino)pyridine-2-yl)- $N^6$ -(4-phenoxyphenyl)pyrimidine-4,6-diamine (0.2 g, 41%) as a light brown solid.


**Step 4**

To a solution of  $N^4$ -(6-(methylamino)pyridine-2-yl)- $N^6$ -(4-phenoxyphenyl)pyrimidine-4,6-diamine (0.07 g, 0.18 mmol) in NMP (1 mL) was added acryloyl chloride (0.032 g, 0.36 mmol) at 0 °C and the reaction mixture was stirred at rt for 1 h. The reaction mixture was diluted with dichloromethane (2 mL), was washed with  $NaHCO_3$  (1 mL), water (1 mL), and brine (1 mL). The dichloromethane solution was dried over  $Na_2SO_4$  and was concentrated under reduced

pressure. The residue was purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to give *N*-methyl-*N*-(6-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyridin-2-yl)acrylamide (**I-78**) as a yellow solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 3.22 (s, 3H), 5.60 (dd, *J* = 2.56 & 9.76 Hz, 1H), 6.12-6.16 (m, 2H), 6.79 (d, *J* = 7.56 Hz, 1H), 6.96 (d, *J* = 8.64 Hz, 5H), 7.09 (t, *J* = 7.32 Hz, 1H), 7.23 (s, 1H), 7.33-7.37 (m, 4H), 7.45 (d, *J* = 8.2 Hz, 2H), 7.32 (t, *J* = 7.8 Hz, 1H), 8.24 (s, 1H); LCMS: *m/e* 439.3 (M+1).

## EXAMPLE 22

## SCHEME 22a




**[00304] Synthesis of (*E*)-4-(dimethylamino)-*N*-methyl-*N*-(6-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-79**)**

To a solution of dimethylaminocrotonic acid hydrochloride (0.120 g, 0.72 mmol) in CH<sub>3</sub>CN (1.4 mL) was added DMF (1 drop) followed by oxalyl chloride (0.07 mL, 0.91 mmol) at 0 °C under nitrogen atmosphere. The reaction was allowed to stir at this temperature for 30 min and then at rt for 2 h. This acid chloride was added, drop wise, at 0 °C in to a stirred solution of *N*<sup>4</sup>-(6-(methylamino)pyridine-2-yl)-*N*<sup>6</sup>-(4-phenoxyphenyl)pyrimidine-4,6-diamine (0.07g, 0.18 mmol) in NMP (2.8 mL). The reaction was allowed to stir at 0 °C for 1 h, diluted with EtOAc (5 mL), and washed with 10% NaHCO<sub>3</sub> (2 mL), water (2 mL) and brine (2 mL). drying over Na<sub>2</sub>SO<sub>4</sub> followed by concentration under reduced pressure gave a residue which was further purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to give (*E*)-4-(dimethylamino)-*N*-methyl-*N*-(6-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyridine-2-yl)but-2-enamide (**I-79**) as a yellow solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 2.05 (s, 6H), 2.91 (d, *J* = 6 Hz, 2H), 3.27 (s, 3H), 6.08 (d, *J* = 15.2 Hz, 1H), 6.65 (dd, *J* = 5.6 & 14.8 Hz, 1H), 6.82 (d, *J* = 7.6 Hz, 1H), 6.97-7.00 (m, 4H), 7.10 (t, *J* = 7.2 Hz, 1H), 7.20 (s, 1H), 7.35-7.39 (m, 2H), 7.46

(d,  $J = 8$  Hz, 1H), 7.53 (d,  $J = 8.8$  Hz, 2H), 7.75 (t,  $J = 8$  Hz, 1H), 8.30 (s, 1H), 9.30 (s, 1H), 9.95 (s, 1H); LCMS:  $m/e$  496 (M+1).

### EXAMPLE 23

#### Scheme 23a




**[00305] Synthesis of (*E*)-4-dimethylamino)-*N*-(6-(6-(4-phenoxyphenoxy)pyrimidin-4-ylamino)pyridin-2-yl)but-2-enamide (I-82)**

To a stirred solution of *N*<sup>2</sup>-(6-(4-phenoxyphenoxy)pyrimidin-4-yl)pyridine-2,6-diamine (0.65 g, 1.75 mmol) in NMP (10 mL) was added dimethylaminocrotonyl chloride (1.026 g, 7 mmol) at 0 °C. The reaction mixture was allowed to come to room temperature and kept at it for 1 h. It was diluted with dichloromethane (10 mL), was washed with NaHCO<sub>3</sub> solution (2 mL) and water (2 mL), and dried over Na<sub>2</sub>SO<sub>4</sub>. The dichloromethane solution was filtered and was concentrated under reduced pressure to give a residue that was purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to give (*E*)-4-dimethylamino)-*N*-(6-(6-(4-phenoxyphenoxy)pyrimidin-4-ylamino)pyridin-2-yl)but-2-enamide (**I-82**) as an off white solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm : 2.19 (s, 6H), 3.08 (d,  $J = 5.52$  Hz, 2H), 6.50 (d,  $J = 15.4$  Hz, 1H), 6.77 (td,  $J = 5.92$  & 15.4 Hz, 1H), 7.00-7.07 (m, 5H), 7.15 (t,  $J = 7.36$  Hz, 1H), 7.21 (dd,  $J = 2.2$  & 8.92 Hz, 2H), 7.41 (t,  $J = 7.52$  Hz, 2H), 7.68 (t,  $J = 7.96$  Hz, 1H), 7.77 (d,  $J = 7.96$  Hz, 1H), 7.95 (s, 1H), 8.35 (s, 1H), 10.20 (s, 1H), 10.40 (s, 1H); LCMS :  $m/e$  483 (M+1).

## EXAMPLE 24

## Scheme 24a



[00306] **Synthesis of 2-(hydroxy(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)methyl)acrylamide (I-84)**


**Step 1**

To a stirred solution of *N*-methoxy-*N*-methyl-3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzamide (0.75 g, 1.7 mmol) in THF (10 mL) was added LAH (3.4 mL, 3.4 mmol, 1 M solution in THF) at  $-60^{\circ}\text{C}$ . The reaction mixture was stirred at  $-60^{\circ}\text{C}$  for 1 h, was quenched with  $\text{Na}_2\text{SO}_4$  solution (2 mL) and was extracted with ethyl acetate (10 mL). The organic layer was separated and washed with water (2 mL) and with brine solution (2 mL) and was dried over anhydrous  $\text{Na}_2\text{SO}_4$ . Filtration followed by concentration under reduced pressure gave a residue that was purified by column chromatography ( $\text{SiO}_2$ , 60-120, chloroform/methanol, 9/1) to give 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzaldehyde (0.6 g, 92%) as an off yellow solid.

**Step 2**

To a stirred solution of 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzaldehyde (100 mg, 0.26 mmol) and acrylonitrile (36 mg, 0.52 mmol) in 1,4- dioxane/ $\text{H}_2\text{O}$  (0.5 mL / 0.5 mL) was added DABCO (29 mg, 0.26 mmol) at rt. Stirring was continued at room temperature for 48 h after which time the reaction mixture was concentrated under reduced pressure. The residue obtained was further purified by column chromatography ( $\text{SiO}_2$ , 60-120, pet ether/ethyl acetate, 6/4) to give 2-(hydroxy(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)methyl)acrylamide (**I-84**) as a white solid.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  ppm : 5.33 (s, 1H), 6.06 (s, 1H), 6.20 (s, 1H), 6.25 (s, 1H), 6.80 (s, 1H), 6.88 (s, 1H), 6.95-7.05 (m, 4H),

7.09-7.13 (m, 2H), 7.16 (bd,  $J = 7.92$  Hz, 1H), 7.32-7.39 (m, 5H), 7.47 (s, 1H), 8.31 (s, 1H);  
LCMS :  $m/e$  436 (M+1).

**EXAMPLE 25****Scheme 25a**

[00307] **Synthesis of 1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyrrolidin-1-yl)prop-2-en-1-one (I-85)**

**Step 1**

A solution of 4,5-dichloropyrimidine (0.6 g, 4.02 mmol), 3-amino-Boc-pyrrolidine (0.5 g, 2.6 mmol) and DIPEA (1.73 g, 13.3 mmol) in n-butanol (5.0 mL) was heated in a pressure tube (120 °C, 12 h). It was cooled, quenched with water (10 mL) and was extracted with EtOAc (2x25 mL). The combined EtOAc extract was washed with water (5 mL), brine (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure to afford tert-butyl 3-(6-chloropyrimidin-4-ylamino)pyrrolidine-1-carboxylate (0.4 g, 50%) as a yellow solid.

**Step 2**

To a stirring solution of tert-butyl 3-(6-chloropyrimidin-4-ylamino)pyrrolidin-1-carboxylate (0.5 g, 1.6 mmol) and 4-phenoxy aniline (0.309 g, 1.6 mmol) in ethanol (4 mL) was added acetic acid (0.1 mL) and the reaction mixture was heated at 100 °C for 36 h. The reaction mixture was cooled, ethanol was removed under reduced pressure and the residue was taken in ethyl acetate (10 mL). It was washed with NaHCO<sub>3</sub> solution (2 mL), brine (2 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The residue was further purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to yield tert-butyl 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyrrolidine-1-carboxylate (0.3 g, 42.8%) as a white solid.


**Step 3**

To a stirred solution of tert-butyl 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyrrolidin-1-carboxylate (0.1 g, 0.2 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (2.0 mL) at 0 °C was added CF<sub>3</sub>COOH (2 mL, 20 vol.) and the reaction mixture was kept at this temperature for 30 min. It was allowed to come to rt and stir at this temperature for 3 h. The reaction mixture was concentrated under reduced pressure and the residue was quenched with water (2 mL), basified with NaHCO<sub>3</sub> solution, and was extracted with ethyl acetate (2x8 mL). The combined ethyl acetate extract was washed with water (2 mL) and brine (2 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to give *N*<sup>4</sup>-(4-phenoxyphenyl)-*N*<sup>6</sup>-(pyrrolidin-3-yl)pyrimidine-4,6-diamine (0.025 g, 32.4%) as a light a brown solid.

**Step 4**

To a stirred solution of *N*<sup>4</sup>-(4-phenoxyphenyl)-*N*<sup>6</sup>-(pyrrolidin-3-yl)pyrimidine-4,6-diamine (0.13 g, 0.3 mmol) in THF (1.5 mL) at -60 °C were added DIPEA (0.07 g, 0.5 mmol) and acryloyl chloride (1 M solution in THF, 0.3 mL, 0.3 mmol), and the reaction mixture was stirred at -60 °C for 5 min. The reaction mixture was quenched by adding water and it was extracted with EtOAc (2x5 mL). The combined EtOAc extract was washed with brine (3 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The residue obtained was purified by column chromatography (SiO<sub>2</sub>, 230-400, chloroform/methanol: 98/2) to give 1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)pyrrolidin-1-yl)prop-2-en-1-one (**I-85**) as a light green solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 1.83-1.86 & 1.90-1.94 (m, 1H), 2.07-3.0 & 2.16-2.20 (m, 1H), 3.38-3.86 (m, 4H), 4.2-4.75 & 4.35-4.5 (bs, 1H), 5.65 (dt, *J* = 2 & 10 Hz, 1H), 5.78 (d,

$J = 3.6$  Hz, 1H), 6.11 & 6.15 (dd,  $J = 2.4$  & 7.0 Hz & dd,  $J = 2.4$  & 7.2 Hz, 1H), 6.51-6.64 (m, together 1H), 6.93-7.00 (m, 4H), 7.07-7.18 (m, 2H), 7.36 (t,  $J = 8$  Hz, 2H), 7.51 (d,  $J = 8$  Hz, 2H), 8.12 (s, 1H), 8.93 (s, 1H); LCMS :  $m/e$  401.8 (M+1).

**EXAMPLE 26****Scheme 26a**

[00308] **Synthesis of 1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)piperidin-1-yl)prop-2-en-1-one (I-86)**

**Step 1**

A solution of 4,6-dichloropyrimidine (0.1 g, 0.671 mmol), 3-amino-Boc-piperidine (0.16 g, 0.80 mmol) and DIPEA (0.086 g, 6.71 mmol) in *n*-butanol (5.0 mL) was heated in a pressure tube

(120 °C, 12 h). The solution was cooled, was quenched with water (2 mL) and was extracted with EtOAc (2x15 mL). The combined EtOAc extract was washed with water (5 mL), brine (5 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to give *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)piperidine-1-carboxylate, which was dried under high vacuum and was used as such for next step without further purification.

### **Step 2**

A solution of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)piperidine-1-carboxylate (0.15 g, 0.48 mmol), 4-phenoxyaniline (0.089 g, 0.48 mmol), Pd(OAc)<sub>2</sub> (0.010 g, 0.048 mmol), BINAP (0.014 g, 0.024 mmol) and Cs<sub>2</sub>CO<sub>3</sub> (0.39 g, 1.2 mmol) in degassed toluene (toluene was purged with N<sub>2</sub> for 15 min) was heated for 12 h at 100 °C under N<sub>2</sub> atmosphere. The reaction mixture was cooled, was diluted with EtOAc (20 mL) and was washed with water (4 mL) and brine (2 mL) and was dried over Na<sub>2</sub>SO<sub>4</sub>. The crude product obtained was purified by column chromatography (SiO<sub>2</sub>, 230-400, chloroform/methanol : 99/1) to give *tert*-butyl 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)piperidine-1-carboxylate (90 mg, 40.9%) as a light yellow solid.

### **Step 3**

To a stirred solution of *tert*-butyl 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)piperidine-1-carboxylate (110 mg, 0.238 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL) at 0 °C was added CF<sub>3</sub>COOH (0.5 mL, 5 vol) and the reaction mixture was kept at this temperature for 30 min. It was allowed to come to rt and to stir at this temperature for 3 h. The reaction mixture was concentrated under reduced pressure and the residue was quenched with water (2 mL), was basified with NaHCO<sub>3</sub> solution, and was extracted with ethyl acetate (2x8 mL). The combined ethyl acetate extract was washed with water (2 mL), brine (2 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure to give *N*<sup>4</sup>-(4-phenoxyphenyl)-*N*<sup>6</sup>-(piperidin-3-yl)pyrimidine-4,6-diamine (0.07 g, 81%) as a light yellow solid.


### **Step 4**

To a stirred solution of *N*<sup>4</sup>-(4-phenoxyphenyl)-*N*<sup>6</sup>-(piperidin-3-yl)pyrimidine-4,6-diamine (0.025 g, 0.069 mmol) in NMP (0.5 mL) at 0 °C was added acryloyl chloride (0.007 g, 0.083 mmol), and the reaction mixture was stirred at 0 °C for 5 min. The reaction mixture was quenched by adding 10% NaHCO<sub>3</sub> solution and it was extracted with EtOAc (2x5 mL). The combined EtOAc extract was washed with water (3 mL) and brine (3 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was

concentrated under reduced pressure. The residue obtained was purified by column chromatography (SiO<sub>2</sub>, 230-400, chloroform/methanol: 98/2) to give 1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)piperidin-1-yl)prop-2-en-1-one (**I-86**) as an off white solid. <sup>1</sup>H NMR (MeOD) δ ppm: 1.5-1.75 (m, 2H), 1.80-2.00 (m, 1H), 2.0-2.20 (m, 1H), 2.70-2.90 (m, 1H), 2.90-3.05 (m, 1H), 3.80-4.00 (m, 2H), 4.30-4.45 (m, 1H), 5.65 & 5.75 (d, *J* = 10.8 Hz & d, *J* = 10.8 Hz respectively, together 1H), 5.81 (d, *J* = 10.8 Hz, 1H), 6.14 & 6.18 (d, *J* = 17.2 Hz & d, *J* = 19.2 Hz respectively, together 1H), 6.60-6.70 & 6.70-6.85 (m, together 1H), 6.97-7.00 (m, 4H), 7.04 (t, *J* = 7.6 Hz, 1H), 7.32-7.38 (m, 4H), 8.04 & 8.07 (s, together 1H); LCMS: *m/e* 416.1 (M+1).

## EXAMPLE 27

## SCHEME 27a



[00309] Synthesis of 3-methyl-1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)but-2-en-1-one (**I-87**)

**Step 1**

To a stirred solution of 4,6-dichloropyrimidine (0.5 g, 3.7 mmol) in *n*-butanol (10 mL) was added methyl 3-aminobenzoate (0.498 g, 3.7 mmol) and DIPEA (0.65 g, 5.0 mmol) and the

reaction mixture was heated at 110 °C for 12 h. It was cooled and excess n-butanol was removed under reduced pressure. The residue was extracted with EtOAc (2x30 mL) and the combined EtOAc extract was washed with water (5 mL), brine (2.5 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated under reduced pressure. The residue was stirred with pet ether (30 mL) for 30 min, the pet ether removed by decantation and the solid obtained was dried under high vacuum to give methyl 3-(6-chloropyrimidin-4-ylamino)benzoate (0.3 g, 34%) as light brown solid.

### **Step 2**

To a stirring solution of methyl 3-(6-chloropyrimidin-4-ylamino)benzoate (1.0 g, 3.8 mmol) and 4-phenoxy aniline (0.703 g, 3.8 mmol) in ethanol (5 mL) was added acetic acid (0.22 mL) and the reaction mixture was heated at 100 °C for 48 h. The reaction mixture was cooled, ethanol was removed under reduced pressure and the residue was dissolved in ethyl acetate (50 mL). It was washed with NaHCO<sub>3</sub> solution (5 mL) and brine (5 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The residue was purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to yield methyl 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzoate (1 g, 66%) as an off-white solid.

### **Step 3**


To a stirred solution of methyl 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzoate (0.3 g, 0.72 mmol) in methanol/THF (2/2, 4 mL) was added LiOH (0.122 g, 2.9 mmol) in H<sub>2</sub>O (4 mL) and the reaction mixture was stirred at rt for 2 h. It was concentrated under reduced pressure. the residue was diluted with water (2 mL) and was extracted with dichloromethane (5 mL). The aqueous layer was separated and was acidified with 1.5 N HCl (pH ~5-6) to get a white precipitate, which was collected by filtration and dried under vacuum to give 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzoic acid (0.2 g 69%) as a white solid.

### **Step 4**

To a stirred solution of 3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzoic acid (0.05 g, 0.12 mmol) in DMF (2 mL) were added MeNH-OMe.HCl (0.0084 g, 0.12 mmol), EDCI.HCl (0.0361 g, 0.18 mmol), HOBT (0.0084 g, 0.062 mmol) and DIPEA (0.023 g, 0.18 mmol). The reaction mixture was stirred at room temperature for 1 h and was quenched with water. A white solid was isolated by filtration and dried under vacuum to give *N*-methoxy-*N*-methyl-3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzamide (0.025 g, 44.5%) as a white solid.

**Step 5**

To a stirred solution of *N*-methoxy-*N*-methyl-3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzamide (50 mg, 0.11 mmol) in THF (0.5 mL) at 0 °C was added 2-methylpropenylmagnesium bromide (1.1 mL, 0.55 mmol, 0.5 M in THF). The reaction mixture was allowed to stir at room temperature for 30 min. It was quenched with sat. NH<sub>4</sub>Cl solution (0.5 mL) and was extracted with EtOAc (3x2 mL). The combined organic layer was washed with brine, was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, was filtered and was concentrated under reduced pressure to give a white solid, which was further purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to give 3-methyl-1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)but-2-en-1-one (**I-87**) as an off white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ ppm: 2.01 (s, 3H), 2.20 (s, 3H), 6.14 (s, 1H), 6.71 (s, 1H), 6.95 (s, 1H), 6.99-7.02 (m, 5H), 7.11 (t, *J* = 7.36 Hz, 1H), 7.26-7.28 (m, 2H), 7.34 (t, *J* = 7.56 Hz, 2H), 7.40-7.53 (m, 2H), 7.65 (d, *J* = 7 Hz, 1H), 7.86 (s, 1H), 8.32 (s, 1H); LCMS : *m/e* 437.2 (M+1).

**EXAMPLE 28****Scheme 28a**


**[00310] Synthesis of 1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)but-2-yn-1-one (**I-88**)**

To a stirred solution of *N*-methoxy-*N*-methyl-3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzamide (50 mg, 0.11 mmol) in THF (0.5 mL) at 0 °C was added a THF solution of 1-butyne (1.1 mL, 1.1 mmol). The reaction mixture was allowed to come to rt and was stirred at rt for 30 min. The reaction mixture was quenched with saturated NH<sub>4</sub>Cl solution (0.5 mL) and was extracted with EtOAc (2x3 mL). The combined EtOAc layer was

washed with brine, was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , was filtered and was concentrated under reduced pressure to give a white solid, which h was further purified by column chromatography ( $\text{SiO}_2$ , 60-120, chloroform/methanol, 9/1) to give 1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)but-2-yn-1-one (**I-88**) as an off-yellow solid.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  ppm: 2.22 (s, 3H), 6.16 (s, 1H), 6.97 (d,  $J = 8.6$  Hz, 2H), 7.01 (d,  $J = 8.8$  Hz, 2H), 7.1 (t,  $J = 7.2$  Hz, 1H), 7.35-7.39 (m, 2H), 7.48 (t,  $J = 8$  Hz, 1H), 7.56 (d,  $J = 8.8$  Hz, 2H), 7.66 (d,  $J = 7.2$  Hz, 1H), 7.95 (d,  $J = 7.6$  Hz, 1H), 8.32 (s, 1H), 8.39 (s, 1H), 9.22 (s, 1H), 9.47 (s, 1H); LCMS :  $m/e$  421.1 (M+1).

### EXAMPLE 29

#### Scheme 29a




#### [00311] Synthesis of (*E,Z*)-1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)but-2-en-1-one (**I-89**)

To a stirred solution of *N*-methoxy-*N*-methyl-3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzamide (50 mg, 0.11 mmol) in THF (0.5 mL) at 0 °C was added a THF solution of propenylmagnesium bromide (2.2 mL, 1.1 mL, 0.5 M soln. in THF). The reaction mixture was allowed to stir at room temperature for 30 min. It was quenched with saturated  $\text{NH}_4\text{Cl}$  solution (0.5 mL) and was extracted with EtOAc (2x3 mL). The combined organic layer was washed with brine, was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , was filtered and was concentrated under reduced pressure to get a white solid, which was further purified by column chromatography ( $\text{SiO}_2$ , 60-120, chloroform/methanol, 9/1) to give (*E,Z*)-1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)but-2-en-1-one (**I-89**) as a pale yellow solid.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  ppm: 1.98 (dd,  $J = 1.6$  & 6.8 Hz, 3H) & 2.13 (dd,  $J = 1.6$  & 7.2 Hz, 3H), 6.10-6.13 (m, 1H), 6.75-6.90 (m,

1H), 6.90-7.13 (m, 7H), 7.25-7.27 (m, 1H), 7.32-7.34 (m, 2H), 7.34-7.50 (m, 2H), 7.63-7.65 (m, 1H), 7.86-7.88 (m, 1H), 8.32 (s, 1H) ; LCMS :  $m/e$  423 (M+1).


### EXAMPLE 30

#### Scheme 30a



#### [00312] Synthesis of 2-methyl-1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)prop-2-en-1-one (I-83)

To a *N*-methoxy-*N*-methyl-3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)benzamide (0.150 g, 0.340 mmol) at 0 °C was added 2-methylpropenylmagnesium bromide (6.8 mL, 3.4 mmol, 0.5 M solution in THF). The reaction mixture was allowed to stir at room temperature for 30 min. It was quenched with saturated NH<sub>4</sub>Cl solution (0.5 mL) and was extracted with EtOAc (2x3 mL). The combined organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure to get a white solid, which was further purified by column chromatography (SiO<sub>2</sub>, 60-120, product getting eluted in methanol/chloroform: 2/98) to give 2-methyl-1-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)prop-2-en-1-one (**I-83**) as a white solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm: 1.99 (s, 3H), 5.64 (s, 1H), 6.03 (s, 1H), 6.14 (s, 1H), 6.96-7.02 (m, 4H), 7.10 (t, *J* = 7.6 Hz, 1H), 7.25 (d, *J* = 7.6 Hz, 1H), 7.35-7.43 (m, 3H), 7.56 (d, *J* = 8.8 Hz, 2H), 7.88 (d, *J* = 8 Hz, 1H), 7.92 (s, 1H), 8.29 (s, 1H), 9.21 (s, 1H), 9.38 (s, 1H); LCMS : 423  $m/e$  (M+1).

**EXAMPLE 31****Scheme 31a**

**[00313] Synthesis of *N*-(6-(6-(3-methoxyphenoxy)pyrimidin-4-ylamino)pyridine-2-yl)acrylamide (I-90)**

**Step 1**

To a solution of *N*<sup>2</sup>-(6-chloropyrimidin-4-yl)pyridine-2,6-diamine (200 mg, 0.90 mmol) in dry DMF (2 mL) was added 3-methoxyphenol (112 mg, 0.90 mmol) and anhydrous K<sub>2</sub>CO<sub>3</sub> (186 mg, 1.353 mmol). The reaction mixture was heated at 100 °C for 16 h under N<sub>2</sub> atmosphere. It was then cooled and DMF removed under reduced pressure to give a yellowish gummy residue that was taken in EtOAc (10 mL). It was washed with water (5 mL), brine (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub> and then concentrated under reduced pressure to give a crude product. This was further purified by column chromatography (SiO<sub>2</sub>, 60-120, hexane/EtOAc, 5/5) to give *N*<sup>2</sup>-(6-(3-methoxyphenoxy)pyrimidin-4-yl)pyridine-2,6-diamine (110 mg, 40.7%) as a pale yellow solid.


**Step 2**

To a stirred solution of *N*<sup>2</sup>-(6-(3-methoxyphenoxy)pyrimidin-4-yl)pyridine-2,6-diamine (100 mg, 0.323 mmol) in THF/NMP (1 mL / 0.5 mL) was added acryloyl chloride (0.029 g, 0.3 mmol) under N<sub>2</sub> atmosphere at -10 °C. The stirring was continued at the same temperature for 2 h and the reaction mixture was concentrated under reduced pressure to give a residue that was further purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol) to give *N*-(6-(6-(3-methoxyphenoxy)pyrimidin-4-ylamino)pyridine-2-yl)acrylamide (**I-90**) as a pale yellow solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 3.75 (s, 3H), 5.79 (dd, *J* = 1.8 & 10.08 Hz, 1H), 6.30 (dd, *J* = 1.8 & 16.96 Hz, 1H), 6.65 (dd, *J* = 10.12 & 16.96 Hz, 1H), 6.74-6.76 (m, 2H), 6.82 (td, *J* = 1.52 & 9.24 Hz, 1H), 7.04 (d, *J* = 7.92 Hz, 1H), 7.33 (t, *J* = 8.28 Hz, 1H), 7.70 (t, *J* = 7.96 Hz, 1H),

7.76 (d,  $J = 8$  Hz, 1H), 7.90 (s, 1H), 8.34 (s, 1H), 10.20 (s, 1H), 10.48 (s, 1H); LCMS :  $m/e$  364 (M+1).

### EXAMPLE 32

#### Scheme 32a



#### [00314] Synthesis of *N*²-(6-(6-(4-methoxyphenoxy)pyrimidin-4-ylamino)pyridine-2-yl)acrylamide (I-91)

##### Step 1

To a solution of *N*²-(6-chloropyrimidin-4-yl)pyridine-2,6-diamine (200 mg, 0.90 mmol) in dry DMF (2 mL) was added 4-methoxyphenol (168 mg, 1.3 mmol) and anhydrous  $K_2CO_3$  (179 mg, 1.3 mmol). The reaction mixture was heated at 100 °C for 16 h under  $N_2$  atmosphere. It was then cooled and DMF removed under reduced pressure to give a yellowish gummy residue that was taken in EtOAc (10 mL). The solution was washed with water (5 mL) and brine (5 mL), was dried over  $Na_2SO_4$  and was then concentrated under reduced pressure to get crude product. This was further purified by column chromatography ( $SiO_2$ , 60-120, hexane/EtOAc, 5/5) to give *N*²-(6-(4-methoxyphenoxy)pyrimidin-4-yl)pyridine-2,6-diamine (160 mg, 59.2 %) as a pale yellow solid.


##### Step 2

To a stirred solution of *N*²-(6-(4-methoxyphenoxy)pyrimidin-4-yl)pyridine-2,6-diamine (150 mg, 0.474 mmol) in THF/NMP (1 mL / 0.5 mL) was added acryloyl chloride (64 mg, 0.712 mmol) at -10 °C under  $N_2$  atmosphere. After stirring at this temperature for 30 min, the reaction was stopped and the reaction mixture was slowly added to  $NaHCO_3$  solution (10 mL). A white solid was precipitated which was isolated by filtration and was dissolved in a mixture of ethyl acetate (5 mL) and  $Et_3N$  (0.5 mL). The solution was washed with water (2 mL) and brine (2 mL). Drying over  $Na_2SO_4$  followed by filtration and concentration under reduced pressure afforded a

yellow solid. It was further purified by column chromatography (SiO<sub>2</sub>, 60-120, chloroform/methanol, 9/1) to give N-(6-(6-(4-methoxyphenoxy)pyrimidin-4-ylamino)pyridine-2-yl)acrylamide (**I-91**) as an off white solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ ppm: 3.76 (s, 3H), 5.79 (dd, *J* = 1.84 & 10.16 Hz, 1H), 6.30 (dd, *J* = 1.84 & 17 Hz, 1H), 6.65 (dd, *J* = 10.12 & 16.92 Hz, 1H), 6.96-6.99 (m, 2H), 7.02 (d, *J* = 7.88 Hz, 1H), 7.09-7.13 (m, 2H), 7.69 (t, *J* = 7.96 Hz, 1H), 7.76 (d, *J* = 7.44 Hz, 1H), 7.86 (s, 1H), 8.30 (d, *J* = 0.88 Hz, 1H), 10.17 (s, 1H), 10.17 (s, 1H); LCMS : *m/e* 364 (M+1).

### EXAMPLE 33

#### Scheme 33a



#### [00315] Synthesis of *N*-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)propionamide (**I<sup>R</sup>-10**)

##### Step 1

A solution of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)phenylcarbamate (200 mg, 0.6 mmol), 4-phenoxyaniline (346 mg, 1.8 mmol) and concentrated HCl (45 mg, 1.2 mmol) in *n*-butanol (8 mL) was subjected to microwave irradiation (160 °C, 20 min). The reaction mixture was quenched with NaHCO<sub>3</sub> solution (2 mL) and extracted with EtOAc (2x10 mL). The combined EtOAc extract was washed with water (5 mL), brine (5 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure. The residue was further purified by column chromatography (SiO<sub>2</sub>, 60-120, Chloroform/methanol, 9/1) to give *N*<sup>4</sup>-(3-aminophenyl)-*N*<sup>6</sup>-(4-phenoxyphenyl)pyrimidine-4,6-diamine (87 mg, 37.8%) as a brown solid.


##### Step 2

To a solution of propionic acid (12 mg, 0.1 mmol) in DMF (0.6 mL) was added HATU (92 mg, 0.2 mmol) and the reaction mixture was stirred at room temperature for 30 min. To it was added *N*<sup>4</sup>-(3-aminophenyl)-*N*<sup>6</sup>-(4-phenoxyphenyl)pyrimidine-4,6-diamine (60 mg, 0.1 mmol) followed by DIPEA (41 mg, 0.3 mmol) and the reaction mixture was stirred at this temperature for 16 h. It

was concentrated under reduced pressure. The residue was diluted with  $\text{CH}_2\text{Cl}_2$  (5 mL) and was washed with  $\text{NaHCO}_3$  solution (2 mL), water (2 mL), and brine (2 mL). Drying over  $\text{Na}_2\text{SO}_4$  followed by concentration under reduced pressure offered a residue that was purified by column chromatography ( $\text{SiO}_2$ , 230-400, chloroform/methanol: 9/1) to give *N*-(3-(6-(4-phenoxyphenylamino)pyrimidin-4-ylamino)phenyl)propionamide (**I<sup>R</sup>-10**) as a brown solid.  $^1\text{H}$  NMR ( $\text{DMSO}-d_6$ )  $\delta$  ppm: 1.06 (t,  $J = 7.56$  Hz, 3H), 2.30 (q,  $J = 7.48$  Hz, 2H), 6.13 (s, 1H), 6.94-6.99 (m, 4H), 7.08 (t,  $J = 7.36$  Hz, 1H), 7.16-7.24 (m, 3H), 7.36 (t,  $J = 7.44$  Hz, 2H), 7.54 (d,  $J = 8.88$  Hz, 2H), 7.81 (s, 1H), 8.23 (s, 1H), 9.12 (s, 2H), 9.81 (s, 1H); LCMS:  $m/e$  426.3 ( $\text{M}+1$ ).

### EXAMPLE 34

#### Scheme 34a



#### [00316] Synthesis of (*E*)-*N*-(3-(6-(3-chloro-4-fluorophenoxy)pyrimidin-4-ylamino)phenyl)-4-(dimethylamino)but-2-enamide (**I-92**)

##### Step 1

A solution of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)phenylcarbamate (1.6g, 5 mmol), 3-chloro-4-fluorophenol (1.4g, 10 mmol) and potassium carbonate (1.4g, 10 mmol) in 15 mL of DMF was heated to  $120^\circ\text{C}$  for 16h. The reaction mixture was mixed in 15 mL of water, the crude product was precipitated, filtered, purified by flash chromatography on silica gel with MeOH/DCM solvent system to afford 750 mg (45% yield) of *N*<sup>1</sup>-(6-(3-chloro-4-fluorophenoxy)pyrimidin-4-yl)benzene-1,3-diamine as an off-white solid. MS ( $m/z$ ):  $\text{MH}^+ = 331$ .


##### Step 2

Oxalyl chloride (155 mg, 1.2 mmol) was added dropwise to a mixture of 4-*N,N*-dimethylaminocrotonic acid HCl salt (200mg, 1.2 mmol) in 5 mL of THF at  $0^\circ\text{C}$ . To this mixture was added 3 drops of DMF/THF solution (made from 5 drops of DMF in 1 mL of THF). The reaction mixture was stirred at RT for 2 h, then was cooled to  $0^\circ\text{C}$  ice bath. A solution of *N*<sup>1</sup>-(6-(3-chloro-4-fluorophenoxy)pyrimidin-4-yl)benzene-1,3-diamine (200mg, 0.6 mmol) 2 mL of NMP) was added to the dimethylaminocrotonyl chloride solution and the resulting mixture was

stirred for 3 h at 0 °C. The reaction was quenched with 3 mL of 1N NaOH, and was extracted with EtOAc (2 x 25 mL). The crude product mixture was purified by flash chromatography on silica gel with MeOH/NH<sub>4</sub>OH/DCM solvent system to afford (*E*)-*N*-(3-(6-(3-chloro-4-fluorophenoxy)pyrimidin-4-ylamino)phenyl)-4-(dimethylamino)but-2-enamide (**I-92**) as a light colored solid. MS (m/z): MH<sup>+</sup> = 442, 444(3:1), H-NMR (DMSO) δ 10.08 (s, 1H), 9.67 (s, 1H), 8.36 (s, 1H), 7.98 (s, 1H), 7.59 (d, 1H), 7.57 (t, 1H), 7.53-7.24 (m, 4H), 6.29 (b, 1H), 6.23 (s, 1H), 3.51 (d, 2H), 2.21 (s, 6H).

### EXAMPLE 35

#### Scheme 35a



#### [00317] Synthesis of 1-(3-(6-(3-chloro-4-fluorophenyamino)pyrimidin-4-ylamino)pyrrolidin-1-yl)prop-2-en-1-one (**I-70**)

##### Step 1

A neat mixture of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)pyrrolidin-1-carboxylate (354 mg, 1.18 mmol) and 3-chloro-4-fluoroaniline (3.03 g, 20.8 mmol) was heated at 140 °C for 21 hours. Upon cooling to ambient temperature, the melt was diluted with EtOAc and the mixture was stirred for 1 hour. A beige, amorphous precipitate was collected, washed with water and dried *in vacuo* at 50-60 °C giving 292 mg (80%) of *N*<sup>4</sup>-(3-chloro-4-fluorophenyl)-*N*<sup>6</sup>-(pyrrolidin-3-yl)pyrimidine-4,6-diamine. MS (APCI): (M + 1) = 308, (M - 1) = 306.


##### Step 2

To a solution of *N*<sup>4</sup>-(3-chloro-4-fluorophenyl)-*N*<sup>6</sup>-(pyrrolidin-3-yl)pyrimidine-4,6-diamine (287 mg, 0.93 mmol) and triethylamine (0.32 ml, 2.33 mmol) in anhydrous THF (5 ml) under nitrogen was added acryloyl chloride (91 µl, 1.12 mmol). The reaction mixture was stirred at room temperature for 1 hour and was concentrated under reduced pressure. The residue was eluted through a flash column (silica gel 60, 230-400 mesh, 5% MeOH in EtOAc to 10% MeOH in EtOAc) to give two products. The less polar product (*R*<sub>f</sub> = 0.24 in 1 : 9 MeOH : EtOAc) was

found to be a diacrylated analog. The more polar product ( $R_f = 0.14$  in 1 : 9 MeOH : EtOAc) was 1-(3-(6-(3-chloro-4-fluorophenyamino)pyrimidin-4-ylamino)piperidin-1-yl)prop-2-en-1-one (I-70). MS (APCI)  $(M + 1)^+ = 362$ ,  $(M - 1)^+ = 360$ :  $^1\text{H-NMR}$  DMSO- $d_6$ )  $\delta$  9.11 (s, 1H), 8.14 (s, 1H), 7.92 (d, 1H), 7.38-7.23 (m, 3H), 6.58-6.48 (m, 1H), 6.13-6.08 (m, 1H), 5.75 (d, 1H), 5.63-5.59 (m, 1H), 3.64-3.59 (m, 2H), 2.17-1.81 (m, 6H).

### EXAMPLE 36


#### Scheme 36a



[00318] **Synthesis of 1-(3-(6-(3-chloro-4-fluorophenyamino)pyrimidin-4-ylamino)piperidin-1-yl)prop-2-en-1-one (I-71)**

#### Step 1

*N*-(3-Chloro-4-fluorophenyl)-*N'*-piperidin-3-yl pyrimidine-4,6-diamine.



A mixture of *tert*-butyl 3-(6-chloropyrimidin-4-ylamino)piperidine-1-carboxylate (295 mg, 0.94 mmol) and 3-chloro-4-fluoroaniline (2.83 g, 19.4 mmol) was heated neat at 140 °C for 19 hours. Upon cooling, the melt was stirred in EtOAc and allowed to stand at room temperature for 1 hour. The precipitate was collected, washed with EtOAc and dried to give 256 mg (85%) of *N*<sup>4</sup>-(3-chloro-4-fluorophenyl)-*N*<sup>6</sup>-(piperidin-3-yl)pyrimidine-4,6-diamine, a grayish-violet amorphous solid. MS (APCI):  $(M + 1)^+ = 322$ .


#### Step 2

To a solution of *N*<sup>4</sup>-(3-chloro-4-fluorophenyl)-*N*<sup>6</sup>-(piperidin-3-yl)pyrimidine-4,6-diamine (251 mg, 0.78 mmol) and triethylamine (0.27 ml, 1.95 mmol) in anhydrous THF (7 ml) under nitrogen was added acryloyl chloride (76  $\mu\text{l}$ , 0.94 mmol). The reaction mixture was stirred at room

temperature for 1 hour and was concentrated under reduced pressure. The residue was eluted through a flash column (silica gel 60, 230-400 mesh with 5% MeOH in EtOAc to give two products. The less polar product ( $R_f = 0.33$  in 1 : 9 MeOH : EtOAc) was found to be a diacrylated analog. The more polar product ( $R_f = 0.24$  in 1 : 9 MeOH : EtOAc) was 1-(3-(6-(3-chloro-4-fluorophenyamino)pyrimidin-4-ylamino)piperidin-1-yl)prop-2-en-1-one (I-71). MS (APCI) ( $M + 1$ )<sup>+</sup> = 376, ( $M - 1$ )<sup>+</sup> = 374: <sup>1</sup>H-NMR DMSO-*d*<sub>6</sub>,  $\delta$  9.15 (br s, 1H), 8.18 (d, 1H), 7.95 (br s, 1H), 7.43-7.20 (m, 2H), 7.03-6.48 (m, 3H), 6.26-5.97 (m, 2H), 5.86-5.51 (m, 3H), 3.90-3.66 (m, 2H), 1.98-1.66 (m, 2H), 1.59-1.12 (m, 2H).

### EXAMPLE 37

#### Scheme 37a



#### [00319] Synthesis of *N*-(6-(6-(2,3-dihydrobenzo[*b*][1,4]dioxin-6-yloxy)pyrimidin-4-ylamino)pyridin-2-yl)acrylamide (I-95)

##### Step 1

To a stirred solution of 2,3-dihydrobenzo[*b*][1,4]dioxine-6-carbaldehyde (1 g, 6.09 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (16 mL) was added *m*-CPBA (4.204 g, 24.36 mmol). The suspension was heated at 50 °C for 2 days, was cooled to rt, was quenched with saturated NaHCO<sub>3</sub> soln. and was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x10 mL). The combined extract was concentrated under reduced pressure, and then was dissolved in MeOH containing NaOH. The solution was stirred at rt for 2 h, was acidified with HCl and was extracted with ethyl acetate (3x10 mL). The combined extract was washed with saturated NaHCO<sub>3</sub> solution and brine, was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, was filtered and was concentrated under reduced pressure. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and was filtered. The DCM solution was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, was filtered and was concentrated under reduced pressure to give 2,3-dihydrobenzo[*b*][1,4]dioxin-6-ol (0.591 g, 63%) as a reddish brown oily liquid.

**Step 2**

A stirred mixture of 2,3-dihydrobenzo[*b*][1,4]dioxin-6-ol (0.137 g, 0.90 mmol), Cs<sub>2</sub>CO<sub>3</sub> (0.734 g, 2.25 mmol), CuI (0.02 g, 10% w/w), and *N*<sup>2</sup>-(6-chloropyrimidin-4-yl)pyridine-2,6-diamine (0.29 g, 0.90 mmol) in NMP (1 mL) was heated at 100 °C for 16 h. The reaction mixture was cooled to rt and slowly added to demineralised water. The solid that precipitated was collected by filtration and was further purified by column chromatography (SiO<sub>2</sub>, 60-120, pet ether/ethyl acetate, 7/3) to give *N*<sup>2</sup>-(6-(2,3-dihydrobenzo[*b*][1,4]dioxin-6-yloxy)pyrimidin-4-yl)pyridine-2,6-diamine (0.088 g, 29%) as yellow solid.

**Step 3**

To a stirred solution of *N*<sup>2</sup>-(6-(2,3-dihydrobenzo[*b*][1,4]dioxin-6-yloxy)pyrimidin-4-yl)pyridine-2,6-diamine (0.080 g, 0.23 mmol) and potassium carbonate (0.065 g, 0.47 mmol) in NMP (0.8 mL) at 0 °C was added acryloyl chloride (0.026 g, 0.29 mmol) and the reaction mixture was stirred at 0 °C for 30 min. The reaction mixture was added dropwise to a cold, stirring solution of 10% NaHCO<sub>3</sub> and was stirred at 0 °C for 30 min. A white solid was isolated by filtration. The solid was washed with cold water and hexane and dissolved in a 2 mL of methanol/dichloromethane (1/1). This solution was concentrated under reduced pressure. The residue obtained was suspended in cold water (5 mL), Et<sub>3</sub>N was added to it and it was extracted with ethyl acetate (2 x 5 mL). The combined ethyl acetate extract was washed with water (2 mL) and brine (2 mL), was dried over Na<sub>2</sub>SO<sub>4</sub> and was concentrated under reduced pressure to obtain *N*-(6-(6-(2,3-dihydrobenzo[*b*][1,4]dioxin-6-yloxy)pyrimidin-4-ylamino)pyridin-2-yl)acrylamide (**I-95**) as a light yellow solid. <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>) δ ppm: 4.25 (s, 4H), 5.79 (dd, *J* = 1.84 & 10.08 Hz, 1H), 6.30 (dd, *J* = 1.84 & 16.96 Hz, 1H), 6.62-6.72 (m, 3H), 6.88 (d, *J* = 8.72 Hz, 1H), 7.03 (d, *J* = 7.84 Hz, 1H), 7.69 (t, *J* = 7.96 Hz, 1H), 7.70 (d, *J* = 7.84 Hz, 1H), 7.85 (s, 1H), 8.32 (d, *J* = 0.4 Hz, 1H), 10.16 (s, 1H), 10.47 (s, 1H); LCMS: *m/e* 392.

[00320] Described below are assays used to measure the biological activity of provided compounds as inhibitors of ErbB1 (EGFR), ErbB2, ErbB4, TEC, BTK, ITK, BMX, and JAK3.

## EXAMPLE 38

**Cloning, Expression and Purification of EGFR-WT and EGFR C797S Mutant Using Baculovirus and Insect Cells****(i) Subcloning of EGFR-WT and mutant kinase domains**

[00321] Amino acids 696 to 1022 of the EGFR-WT kinase domain (NM\_005228, NP\_005219.2) was subcloned into the NcoI and HindIII sites of the pFastHTa vector (Invitrogen<sup>TM</sup>, Carlsbad, CA). To make the EGFR-mutant protein, the cysteine at position 797 was changed to a serine using the Stratagene QuikChange kit (Stratagene, Cedar Creek, TX), according to manufacturer's instructions.

**(ii) Expression**

[00322] P1 baculovirus stocks were generated in SF9 cells via Blue Sky Biotech's suspension transfection protocol (Worcester, MA). Expression analysis was conducted in 125 ml culture of SF21 insect cells ((grown in SF900I SFM (Invitrogen cat # 10902-088), supplemented with 10mg/L gentamicin (Invitrogen, Carlsbad, CA, cat# 15710-064)) using a viral load of 0.1ml of virus per 100 ml of cell suspension. Expression was optimized using Blue Sky Biotech's Infection Kinetics Monitoring system (Worcester, MA).

**(iii) Purification**

[00323] Infected insect cells were pelleted. Cell pellets were resuspended in Blue Sky Biotech's lysis buffer (Worcester, MA, 1X WX; solubilization buffer, containing a protease inhibitor cocktail of leupeptin, pepstatin, PMSF, aprotinin and EDTA) at a ratio of 10 ml per gram of wet cell paste. Cells were lysed by sonication and the lysate was clarified by centrifugation at 9,000 RPM for 30 minutes in a GSA rotor. 500 µl bed volume of NiNTA resin (Qiagen, Valencia, CA) was added to the supernatants and batch bound for two hours with constant agitation. The material was transferred by gravity into an empty 2ml column. The column was washed with 2 ml of wash buffer (Blue Sky Biotech, Worcester, MA, 1X WX, 25mM imidazole ).The protein was eluted with 1X WX + imidazole at varying concentrations: Elution 1: 75 mM imidazole ( 2 fractions, 1 column volume); Elution 2 : 150 mM imidazole (2 fractions, 1 column volume); Elution 3 : 300 mM imidazole (2 fractions, 1 column volume). All the elution fractions were analyzed by SDS page followed by Coomassie staining and Western Blotting using anti-penta-his antibody (Qiagen, Valencia, CA). The carboxy-terminal six-histidine "tag" was removed from some of the purified protein using AcTEV Protease kit (

Invitrogen, Carlsbad, CA, Cat# 12575-015), following manufacturer's instructions. All the samples (pre- and post- Tev cut) were analyzed by SDS page followed by Coomassie staining and Western Blotting, as described above.

#### EXAMPLE 39

##### Mass Spectrometry for EGFR

[00324] EGFR wild type and EGFR mutant (C797S) was incubated with 10-fold excess of compound I-1 for 1 hr and 3 hrs. 1 ul aliquots of the samples (total volume 5-8 ul) were diluted with 10 ul of 0.1% TFA prior to micro C4 ZipTipping directly onto the MALDI target using Sinapinic acid as the desorption matrix (10 mg/ml in 0.1%TFA:Acetonitrile 50:50). Intact mass measurement reveals that the wild type has a nominal mass of about 37557 and the mutant slightly lower at 37500. Reactivity was only observed for the wild type EGFR with a new peak appearing at a mass consistent with a single site covalent modification with compound I-1 which has a mass of 410 Da. (See Figure 8). The mutant EGFR (C797S) showed no significant reactivity even after 3 hrs, confirming modification of the cysteine of interest, Cys797.

#### EXAMPLE 40

##### Omnia Assay Protocol for Potency Assessment Against EGFR (WT) and EGFR (T790M/L858R) Active Enzymes

[00325] The protocol below describes continuous-read kinase assays to measure inherent potency of compounds against active forms of EGFR (WT) and EGFR (T790M/L858R) enzymes. Description of the mechanics of the assay platform may be obtained from the vendor (Invitrogen, Carlsbad, CA).

[00326] Briefly, 10X stocks of EGFR-WT (PV3872) from Invitrogen and EGFR-T790M/L858R (40350) from BPS Bioscience, San Diego, CA, 1.13X ATP (AS001A) and appropriate Tyr-Sox conjugated peptide substrates (KCZ1001) were prepared in 1X kinase reaction buffer consisting of 20 mM Tris, pH 7.5, 5 mM MgCl<sub>2</sub>, 1 mM EGTA, 5 mM β-

glycerophosphate, 5% glycerol (10X stock, KB002A) and 0.2 mM DTT (DS001A). 5  $\mu$ L of each enzyme were pre-incubated in a Corning<sup>TM</sup> (#3574) 384-well, white, non-binding surface microtiter plate (Corning, NY) for 30 min. at 27°C with a 0.5  $\mu$ L volume of 50% DMSO and serially diluted compounds prepared in 50% DMSO. Kinase reactions were started with the addition of 45  $\mu$ L of the ATP/Tyr-Sox peptide substrate mix and monitored every 30-90 seconds for 60 minutes at  $\lambda_{ex}$ 360/ $\lambda_{em}$ 485 in a Synergy<sup>4</sup> plate reader from BioTek (Winooski, VT). At the conclusion of each assay, progress curves from each well were examined for linear reaction kinetics and fit statistics ( $R^2$ , 95% confidence interval, absolute sum of squares). Initial velocity (0 minutes to ~30 minutes) from each reaction was determined from the slope of a plot of relative fluorescence units vs time (minutes) and then plotted against inhibitor concentration to estimate  $IC_{50}$  from log[Inhibitor] vs Response, Variable Slope model in GraphPad Prism from GraphPad Software (San Diego, CA).

[00327] The EGFR-WT- and EGFR T790M/L858R-modified optimized reagent conditions are:

[EGFR-WT] = 5 nM, [ATP] = 15 mM, [Y12-Sox] = 5 mM (ATP  $K_{Mapp}$  ~ 12 mM); and  
[EGFR-T790M/L858R] = 3 nM, [ATP] = 50 mM, [Y12-Sox] = 5 mM (ATP  $K_{Mapp}$  ~ 45 mM).

#### EXAMPLE 41

[00328] Table 7 shows the activity of selected compounds of this invention in the EGFR inhibition assay. The compound numbers correspond to the compound numbers in Table 5. Compounds having an activity designated as "A" provided an  $IC_{50} \leq 10$  nM; compounds having an activity designated as "B" provided an  $IC_{50}$  10-100 nM; compounds having an activity designated as "C" provided an  $IC_{50}$  of 100-1000 nM; compounds having an activity designated as "D" provided an  $IC_{50}$  of 1000-10,000 nM; and compounds having an activity designated as "E" provided an  $IC_{50} \geq 10,000$  nM.

**Table 7. EGFR Wild Type and EGFR (mutant C797S) Inhibition Data**

| Compound # | EGFR inhibition | EGFR (T790M/L858R) inhibitor |
|------------|-----------------|------------------------------|
| I-1        | A               | A                            |
| I-2        | A               | A                            |

| <b>Compound #</b> | <b>EGFR<br/>inhibition</b> | <b>EGFR<br/>(T790M/L858R)<br/>inhibitor</b> |
|-------------------|----------------------------|---------------------------------------------|
| <b>I-3</b>        | A                          | A                                           |
| <b>I-4</b>        | A                          | B                                           |
| <b>I-5</b>        | A                          | A                                           |
| <b>I-6</b>        | B                          | B                                           |
| <b>I-7</b>        | A                          | B                                           |
| <b>I-8</b>        | A                          | D                                           |
| <b>I-9</b>        | A                          | C                                           |
| <b>I-10</b>       | A                          | C                                           |
| <b>I-11</b>       | B                          | E                                           |
| <b>I-12</b>       | B                          | E                                           |
| <b>I-13</b>       | A                          | B                                           |
| <b>I-14</b>       | A                          | B                                           |
| <b>I-15</b>       | A                          | B                                           |
| <b>I-16</b>       | A                          | B                                           |
| <b>I-17</b>       | A                          | B                                           |
| <b>I-18</b>       | A                          | A                                           |
| <b>I-19</b>       | A                          | A                                           |
| <b>I-46</b>       | A                          | A                                           |
| <b>I-47</b>       | A                          | E                                           |
| <b>I-48</b>       | C                          | C                                           |
| <b>I-49</b>       | A                          | A                                           |
| <b>I-50</b>       | A                          | B                                           |
| <b>I-51</b>       | A                          | D                                           |
| <b>I-52</b>       | A                          | A                                           |
| <b>I-53</b>       | B                          | C                                           |
| <b>I-54</b>       | A                          | A                                           |
| <b>I-55</b>       | A                          | A                                           |
| <b>I-56</b>       | A                          | A                                           |
| <b>I-57</b>       | -                          | C                                           |
| <b>I-58</b>       | -                          | C                                           |
| <b>I-59</b>       | A                          | A                                           |
| <b>I-60</b>       | D                          | C                                           |
| <b>I-61</b>       | -                          | A                                           |
| <b>I-63</b>       | A                          | A                                           |
| <b>I-65</b>       | A                          | B                                           |
| <b>I-66</b>       | A                          | C                                           |
| <b>I-67</b>       | A                          | A                                           |
| <b>I-68</b>       | A                          | -                                           |
| <b>I-69</b>       | B                          | -                                           |
| <b>I-70</b>       | A                          | -                                           |
| <b>I-71</b>       | A                          | -                                           |
| <b>I-73</b>       | A                          | -                                           |

| Compound # | EGFR<br>inhibition | EGFR<br>(T790M/L858R)<br>inhibitor |
|------------|--------------------|------------------------------------|
| I-75       | A                  | -                                  |
| I-78       | A                  | -                                  |
| I-79       | A                  | -                                  |
| I-80       | C                  | -                                  |
| I-81       | A                  | -                                  |
| I-82       | A                  | -                                  |
| I-83       | B                  | -                                  |
| I-84       | B                  | -                                  |
| I-85       | B                  | -                                  |
| I-86       | B                  | -                                  |
| I-87       | B                  | -                                  |
| I-88       | A                  | -                                  |
| I-89       | A                  | -                                  |
| I-90       | A                  | -                                  |
| I-91       | A                  | -                                  |
| I-92       | A                  | -                                  |
| I-93       | A                  | -                                  |
| I-94       | A                  | A                                  |

**EXAMPLE 42****Cellular Assays for EGFR Activity**

[00329] Compounds were assayed in A431 human epidermoid carcinoma cells using a method substantially similar to that described in Fry, et al., *Proc. Natl. Acad. Sci. USA* Vol 95, pp 12022-12027, 1998. Specifically, A431 human epidermoid carcinoma cells were grown in 6-well plates to 90% confluence and then incubated in serum-free media for 18 hr. Duplicate sets of cells were treated with 1  $\mu$ M designated compound for 2, 5, 10, 30, or 60 min. Cells were washed free of the compound with warmed serum-free medium, incubated for 2 hr, washed again, incubated another 2 hr, washed again, and then incubated another 2 hr washed again and incubated for additional 2 hr and then stimulated with 100 ng/ml EGF for 5 min. Extracts were made as described Fry, et al. **Figure 1** depicts the EGFR inhibiting activity of compound I-1.

[00330] Compounds were assayed in A431 human epidermoid carcinoma cells using a method substantially similar to that described in Fry, et al. Specifically, A431 human epidermoid carcinoma cells were grown in 6-well plates to 90% confluence and then incubated in serum-free media for 18 hr. Cells were then treated with 10, 1, 0.1, 0.01, or 0.001  $\mu$ M test compound for 1 hr. Cells were then stimulated with 100 ng/ml EGF for 5 min, and extracts were

made as described in Fry, et al. 20ug total protein from lysates were loaded on gel and blots were probed for either EGFR phosphorylation or p42/p44 Erk phosphorylation.

[00331] Dose response inhibition of EGFR phosphorylation and p42/p44 Erk phosphorylation with compound **I-16** and **I-17** in A431 cells is depicted in **Figure 3**. Dose response inhibition of EGFR phosphorylation and p42/p44 Erk phosphorylation with compound **I-19** in A431 cells is depicted in **Figure 4**. Dose response inhibition of EGFR phosphorylation with compound **I-1** in A431 cells, as compared with its “reversible control” compound (**I<sup>R</sup>-3**), is depicted in **Figure 5**.

#### EXAMPLE 43

##### Washout Experiment for EGFR Activity

[00332] A431 human epidermoid carcinoma cells were grown in 6-well plates to '90% confluence and then incubated in serum-free media for 18 hr. Duplicate sets of cells were treated with 1μM designated compound for 1 hr. One set of cells was then stimulated with 100 ng/ml EGF for 5 min, and extracts were made as described. The other set of cells was washed free of the compound with warmed compound-free medium, incubated for 2 hr, washed again, incubated another 2 hr, washed again, and then incubated another 2hr washed again and incubated for additional 2 hr and then stimulated with EGF. The results of this experiment are depicted in **Figure 6** where it is shown that compound **I-1** maintains enzyme inhibition after “washout” whereas its “reversible control” compound (**I<sup>R</sup>-3**) was washed away in the experiment thereby resulting in reactivated enzyme activity.

#### EXAMPLE 44

##### Mass Spectrometry for ErbB4

[00333] ErbB4 kinase domain (Upstate) was incubated with compound for 90 minutes at 10-fold excess of compound **I-1** to protein. 1 ul aliquots of the samples (total volume of 4.24ul) were diluted with 10 ul of 0.1% TFA prior to micro C4 ZipTipping directly onto the MALDI target using Sinapinic acid as the desorption matrix (10 mg/ml in 0.1%TFA:Acetonitrile 50:50). For intact protein mass measurement the instrument was set in Linear mode using a pulsed extraction setting of 16,952 for the myoglobin standard used to calibrate the instrument (Shimadzu Axima TOF<sup>2</sup>).

[00334] Intact ErbB4 protein occurs at MH<sup>+</sup> of 35850 with corresponding sinapinic (matrix) adducts occurring about 200 Da higher. A stoichiometric incorporation of the compound I-1 (Mw of 410 Da) produced a new mass peak which is approximately 410 Da higher (MH<sup>+</sup> of 36260). This is consistent with covalent modification of ErbB4 with compound I-1 as depicted in Figure 7.

#### EXAMPLE 45

##### ErbB1, ErbB2 and/or ErbB4 Kinase Inhibition

[00335] Compounds of the present invention were assayed as inhibitors of one or more of ErbB1, ErbB2, and/or ErbB4 in a manner substantially similar to the method described by Invitrogen Corp (Invitrogen Corporation, 1600 Faraday Avenue, Carlsbad, California, CA; [http://www.invitrogen.com/downloads/Z-LYTE\\_Brochure\\_1205.pdf](http://www.invitrogen.com/downloads/Z-LYTE_Brochure_1205.pdf)) using the Z'-LYTE™ biochemical assay procedure or similar biochemical assay. The Z'-LYTE™ biochemical assay employs a fluorescence-based, coupled-enzyme format and is based on the differential sensitivity of phosphorylated and non-phosphorylated peptides to proteolytic cleavage.

#### EXAMPLE 46

[00336] Table 8 shows the activity of selected compounds of this invention in the ErbB inhibition assay. The compound numbers correspond to the compound numbers in Table 5.

**Table 8. ErbB1, ErbB2, and/or ErbB4 Inhibition Data**

| Compound # | ErbB1<br>% Inhibition        | ErbB2<br>% Inhibition      | ErbB4<br>% Inhibition      |
|------------|------------------------------|----------------------------|----------------------------|
| I-1        | 99% @ 10 nM                  | 76% @ 1 μM<br>61% @ 100 nM | 86% @ 1 μM<br>64% @ 100 nM |
| I-2        | 98% @ 100 nM<br>75% @ 10 nM  | 96% @ 1 μM<br>39% @ 100 nM | 75% @ 1 μM                 |
| I-3        | 96% @ 100nM<br>56% @ 10 nM   | 89% @ 1 μM                 | 95% @ 1 μM                 |
| I-4        | 100% @ 10 nM                 | 86% @ 1 μM                 | 78% @ 1 μM                 |
| I-5        | 100% @ 10 nM                 | 86% @ 1 μM                 | 95% @ 1 μM                 |
| I-6        | 100% @ 100 nM<br>49% @ 10 nM | 84% @ 1 μM                 | 97% @ 1 μM                 |
| I-7        | 100% @ 100 nM<br>53% @ 10 nM | 89% @ 1 μM                 | 100% @ 1 μM                |
| I-8        | 83% @ 10 nM                  | -                          | 57% @ 1 μM                 |
| I-9        | 100% @ 1 μM                  | -                          | 75% @ 1 μM                 |
| I-10       | 96% @ 1 μM                   | -                          | -                          |

| Compound #  | ErbB1<br>% Inhibition | ErbB2<br>% Inhibition       | ErbB4<br>% Inhibition       |
|-------------|-----------------------|-----------------------------|-----------------------------|
| <b>I-11</b> | 79% @ 1 $\mu$ M       | -                           | -                           |
| <b>I-12</b> | 82% @ 1 $\mu$ M       | -                           | -                           |
| <b>I-13</b> | 92% @ 1 $\mu$ M       | 98% @ 1 $\mu$ M             | -                           |
| <b>I-14</b> | 96% @ 1 $\mu$ M       | -                           | -                           |
| <b>I-15</b> | 98% @ 1 $\mu$ M       | -                           | -                           |
| <b>I-16</b> | 98% @ 1 $\mu$ M       | 87% @ 100 nM<br>24% @ 10 nM | 93% @ 100 nM<br>26% @ 10 nM |
| <b>I-17</b> | 95% @ 1 $\mu$ M       | 89% @ 1 $\mu$ M             | 94% @ 1 $\mu$ M             |
| <b>I-18</b> | -                     | 91% @ 1 $\mu$ M             | 94% @ 1 $\mu$ M             |
| <b>I-19</b> | 96% @ 1 $\mu$ M       | 98% @ 1 $\mu$ M             | 97% @ 1 $\mu$ M             |

**EXAMPLE 47****Mass Spectrometry for TEC Kinase**

[00337] TEC kinase (45 pmols; Invitrogen) was incubated with (**I-13**) (450 pmols) for 3hrs at 10X excess prior to tryptic digestion. Iodoacetamide was used as the alkylating agent after compound incubation. A control sample (45 pmols) was also prepared which did not have the addition of (**I-13**). For tryptic digests a 5ul aliquot (7.5 pmols) was diluted with 15 ul of 0.1% TFA prior to micro C18 Zip Tipping directly onto the MALDI target using alpha cyano-4-hydroxy cinnamic acid as the matrix (5mg/ml in 0.1%TFA:Acetonitrile 50:50).

[00338] As depicted in **Figure 11**, the expected peptide (GCLLNFLR) to be modified was immediately evident at MH<sup>+</sup> of 1358.65. This is the mass to be expected when compound **I-13**, with an adduct mass of 423.17, is added to the peptide mass of 935.51. The peptide was also quite evident in the control sample as modified by Iodoacetamide at MH<sup>+</sup> of 992.56. Interestingly the Iodoacetamide modified peptide was not evident in the digest reacted with compound **I-13** indicating that the reaction was complete. There was no evidence of any other modified peptides.

[00339] Evidence of compound **I-13** was observed at MH<sup>+</sup> of 424.20 in the low mass range of the spectra. The fragmentation spectra of the 424.20 peak did show many diagnostic fragments that were apparent in the PSD spectra of the modified peptide at 1358.65 (see **Figure 11**).

[00340] To further verify the presence of the modified peptide with compound **I-13**, the peptide at MH<sup>+</sup> of 1358.65 was subjected to PSD (MS/MS) analysis. A correlational analysis with the homospaien database identified the correct peptide modified by **I-13**.

**Instrumental:**

[00341] For tryptic digests the instrument was set in Reflectron mode with a pulsed extraction setting of 2200. Calibration was done using the Laser Biolabs Pep Mix standard (1046.54, 1296.69, 1672.92, 2093.09, 2465.20). For CID/PSD analysis the peptide was selected using cursors to set ion gate timing and fragmentation occurred at a laser power about 20% higher and He was used as the collision gas for CID. Calibration for fragments was done using the P14R fragmentation calibration for the Curved field Reflectron.

**EXAMPLE 48****Omnia Assay Protocol for Potency Assessment Against BTK**

[00342] The Omnia Assay Protocol for potency assessment against BTK is performed in a substantially similar manner as that described in Example 25 above except that the modified BTK-optimized reagent conditions are:

[BTK] = 5 nM, [ATP] = 40 mM, [Y5-Sox] = 10 mM (ATP K<sub>M</sub>app ~ 36 mM).

**EXAMPLE 49**

[00343] Table 9 shows the activity of selected compounds of this invention in the BTK inhibition assay. The compound numbers correspond to the compound numbers in Table 5. Compounds having an activity designated as "A" provided an IC<sub>50</sub> ≤ 10 nM; compounds having an activity designated as "B" provided an IC<sub>50</sub> 10-100 nM; compounds having an activity designated as "C" provided an IC<sub>50</sub> of 100-1000 nM; compounds having an activity designated as "D" provided an IC<sub>50</sub> of 1000-10,000 nM; and compounds having an activity designated as "E" provided an IC<sub>50</sub> ≥ 10,000 nM.

**Table 9. BTK Inhibition Data**

| Compound # | BTK Inhibition |
|------------|----------------|
| I-1        | B              |
| I-2        | B              |
| I-3        | B              |
| I-4        | A              |
| I-5        | A              |
| I-6        | C              |
| I-7        | B              |
| I-8        | E              |
| I-9        | C              |

| Compound # | BTK Inhibition |
|------------|----------------|
| I-10       | C              |
| I-11       | E              |
| I-12       | E              |
| I-13       | A              |
| I-14       | C              |
| I-15       | B              |
| I-16       | B              |
| I-17       | A              |
| I-18       | B              |
| I-19       | B              |
| I-46       | C              |
| I-47       | E              |
| I-48       | B              |
| I-49       | A              |
| I-50       | D              |
| I-51       | E              |
| I-52       | A              |
| I-53       | B              |
| I-54       | C              |
| I-55       | B              |
| I-56       | B              |
| I-57       | D              |
| I-58       | D              |
| I-59       | C              |
| I-60       | D              |
| I-61       | B              |
| I-62       | A              |
| I-63       | A              |
| I-64       | A              |
| I-65       | A              |
| I-66       | A              |
| I-67       | B              |
| I-68       | B              |
| I-69       | C              |
| I-70       | C              |
| I-71       | D              |
| I-73       | A              |
| I-75       | A              |
| I-78       | A              |
| I-79       | A              |
| I-80       | D              |
| I-81       | A              |
| I-82       | A              |
| I-83       | B              |

| Compound # | BTK Inhibition |
|------------|----------------|
| I-84       | D              |
| I-85       | B              |
| I-86       | B              |
| I-87       | E              |
| I-88       | A              |
| I-89       | A              |
| I-90       | A              |
| I-91       | A              |
| I-92       | B              |
| I-93       | D              |
| I-94       | A              |

#### EXAMPLE 50

##### **BTK Ramos Cellular Assay**

[00344] Compounds I-13 and (I-52) were assayed in Ramos human Burkitt lymphoma cells. Ramos cells were grown in suspension in T225 flasks, spun down, resuspended in 50mls serum-free media and incubated for 1 hour. Compound was added to Ramos cells in serum free media to a final concentration of 1, 0.1, 0.01, or 0.001  $\mu$ M. Ramos cells were incubated with compound for 1 hour, washed again and resuspended in 100ul serum-free media. Cells were then stimulated with 1  $\mu$ g of goat F(ab')<sub>2</sub> Anti-Human IgM and incubated on ice for 10 minutes to activate B cell receptor signaling pathways. After 10 minutes, the cells were washed once with PBS and then lysed on ice with Invitrogen Cell Extraction buffer. 16  $\mu$ g total protein from lysates was loaded on gels and blots were probed for phosphorylation of the BTK substrate PLC $\gamma$ 2. At 1  $\mu$ M I-13 showed 85 % inhibition and (I-52) showed 50% inhibition of BTK signaling in Ramos cells. Additional dose response inhibition of BTK signaling with I-13 is depicted in **Figure 9**.

[00345] Table 10 provides inhibition data for selected compounds in Ramos cells. The compound numbers correspond to the compound numbers in Table 5. Compounds having an activity designated as "A" provided an IC<sub>50</sub>  $\leq$  10 nM; compounds having an activity designated as "B" provided an IC<sub>50</sub> of 10-100 nM; compounds having an activity designated as "C" provided an IC<sub>50</sub> of 100-1000 nM; compounds having an activity designated as "D" provided an IC<sub>50</sub> of  $\geq$  1000 nM.

**Table 10. BTK Ramos Cellular Inhibition Data**

| Compound # | BTK Inhibition (nM) |
|------------|---------------------|
| I-5        | C                   |
| I-13       | B                   |
| I-17       | D                   |
| I-63       | A                   |
| I-64       | B                   |
| I-65       | A                   |
| I-66       | A                   |
| I-78       | A                   |
| I-79       | A                   |
| I-81       | A                   |
| I-82       | A                   |

**EXAMPLE 51****Washout Experiment with Ramos cells for BTK Activity**

[00346] Ramos cells were serum starved for one hour in RPMI media +1% glutamine at 37°C. After starvation, Ramos cells were treated with 100nM compound diluted in serum free RPMI media for 1 hour. After compound treatment, the media was removed and cells were washed with compound-free media. Subsequently, Ramos cells were washed every 2 hours and resuspended in fresh compound-free media. Cells were collected at specified timepoints, treated with 1ug anti-human IgM (Southern Biotech cat # 2022-01) for 10 minutes on ice to induce BCR signaling and then washed in PBS. Ramos cells were then lysed in Cell Extraction Buffer (Invitrogen FNN0011) supplemented with Roche complete protease inhibitor tablets (Roche 11697498001) and phosphatase inhibitors (Roche 04 906 837 001) and 18 ug total protein lysate was loaded in each lane. Inhibition of Btk kinase activity was assayed by measuring its substrate (PLC $\gamma$ 2) phosphorylation by western blot with phospho-specific antibodies from Cell Signaling Technologies cat# 3871. **Figure 10** depicts the results of compound I-13 in the washout experiment at 0 hour, 4 hour, 6 hour, and 8 hour time points. As shown in Figure 10, compound I-13 maintains inhibition of BTK for 8 hours.

[00347] Table 11 provides data for selected compounds in the Ramos washout assay.

**Table 11. BTK Washout Data**

| Compound # | BTK Inhibition Type |
|------------|---------------------|
| I-13       | irreversible        |
| I-63       | irreversible        |
| I-64       | irreversible        |

| Compound # | BTK Inhibition Type |
|------------|---------------------|
| I-65       | reversible          |
| I-66       | irreversible        |
| I-82       | irreversible        |

**EXAMPLE 52****Mass Spectrometry for BTK**

[00348] Intact BTK was incubated for 1 hr at a 10x fold excess of compound I-63 or I-66 to protein. Aliquots (2 ul) of the samples were diluted with 10ul of 0.1% TFA prior to micro C4 ZipTipping directly onto the MALDI target using Sinapinic acid as the desorption matrix (10mg/ml in 0.1%TFA:Acetonitrile 20:80). Mass spectrometry traces are shown in **Figure 12** and **Figure 13**. The top panels of Figures 12 and 13 show the mass spec trace of the intact BTK protein ( $m/z$  81,169 Da). The bottom panels show the mass spec trace when BTK was incubated with compound I-63 (mw 424.5) or compound I-66 (mw 425.5) in Figures 12 and 13, respectively. The centroid mass ( $m/z$  81,593 kDa) in the bottom panel of Figure 12 shows a positive shift of about 424 Da, indicating complete modification of BTK by compound I-63. The centroid mass ( $m/z$  81,593 kDa) in the bottom panel of Figure 13 shows a positive shift of about 407 Da, indicating complete modification of BTK by compound I-66.

**EXAMPLE 53****Omnia Assay Protocol for Potency Assessment Against Active Forms of ITK Kinase**

[00349] This example describes continuous-read kinase assays to measure inherent potency of compound against active forms of ITK enzymes as described in Example 10 above except that the modified ITK-optimized reagent conditions are:

$$[\text{ITK}] = 10 \text{ nM}, [\text{ATP}] = 25 \text{ }\mu\text{M}, [\text{Y6-Sox}] = 10 \text{ }\mu\text{M} (\text{ATP } K_{\text{Mapp}} = 33 \text{ }\mu\text{M}).$$

**EXAMPLE 54**

[00350] Table 12 shows the activity of selected compounds of this invention in the ITK inhibition assay. The compound numbers correspond to the compound numbers in Table 5. Compounds having an activity designated as "A" provided an  $\text{IC}_{50} \leq 10 \text{ nM}$ ; compounds having an activity designated as "B" provided an  $\text{IC}_{50}$  10-100 nM; compounds having an activity designated as "C" provided an  $\text{IC}_{50}$  of 100-1000 nM; compounds having an activity designated

as "D" provided an  $IC_{50}$  of 1000-10,000 nM; and compounds having an activity designated as "E" provided an  $IC_{50} \geq 10,000$  nM.

**Table 12. ITK Inhibition Data**

| Compound # | ITK inhibition |
|------------|----------------|
| I-10       | E              |
| I-13       | D              |
| I-14       | D              |
| I-15       | D              |
| I-63       | D              |
| I-64       | B              |
| I-65       | B              |
| I-66       | D              |
| I-78       | B              |
| I-79       | A              |
| I-81       | B              |
| I-88       | B              |
| I-89       | C              |
| I-90       | C              |
| I-94       | B              |

#### EXAMPLE 55

##### **Omnia Assay Protocol for Potency Assessment Against Active Forms of BMX Kinase**

[00351] This example describes continuous-read kinase assays to measure inherent potency of compound against active forms of BMX enzymes as described in Example 10 above except that the modified BMX-optimized reagent conditions are:

$$[BMX] = 2.5 \text{ nM}, [ATP] = 100 \text{ } \mu\text{M}, [Y5\text{-Sox}] = 7.5 \text{ } \mu\text{M} \text{ (ATP } K_{Mapp} = 107 \text{ } \mu\text{M)}.$$

#### EXAMPLE 56

[00352] Table 13 shows the activity of selected compounds of this invention in the BMX inhibition assay. The compound numbers correspond to the compound numbers in Table 5. Compounds having an activity designated as "A" provided an  $IC_{50} \leq 10$  nM; compounds having an activity designated as "B" provided an  $IC_{50}$  10-100 nM; compounds having an activity designated as "C" provided an  $IC_{50}$  of 100-1000 nM; compounds having an activity designated as "D" provided an  $IC_{50}$  of 1000-10,000 nM; and compounds having an activity designated as "E" provided an  $IC_{50} \geq 10,000$  nM.

**Table 13. BMX Inhibition Data**

| Compound # | BMX inhibition |
|------------|----------------|
| I-10       | -              |
| I-13       | A              |
| I-14       | -              |
| I-15       | -              |
| I-63       | A              |
| I-64       | B              |
| I-65       | A              |
| I-66       | A              |
| I-78       | B              |
| I-79       | A              |
| I-81       | A              |
| I-94       | A              |

**EXAMPLE 57****Omnia Assay Protocol for Potency Assessment Against the Active Form of Janus-3 Kinase (JAK3):**

[00353] The Omnia Assay Protocol for potency assessment against JAK3 is performed in a substantially similar manner as that described in Example 25 above except that the modified JAK3-optimized reagent conditions are:

[JAK3] = 5 nM, [ATP] = 5  $\mu$ M, [Y12-Sox] = 5  $\mu$ M (ATP  $K_{Mapp}$  ~5  $\mu$ M).

**EXAMPLE 58**

[00354] Table 14 shows the activity of selected compounds of this invention in the JAK3 inhibition assay. The compound numbers correspond to the compound numbers in Table 5. Compounds having an activity designated as "A" provided an  $IC_{50} \leq 10$  nM; compounds having an activity designated as "B" provided an  $IC_{50}$  10-100 nM; compounds having an activity designated as "C" provided an  $IC_{50}$  of 100-1000 nM; compounds having an activity designated as "D" provided an  $IC_{50}$  of 1000-10,000 nM; and compounds having an activity designated as "E" provided an  $IC_{50} \geq 10,000$  nM.

**Table 14. JAK3 Inhibition Data**

| Compound # | JAK3 Inhibition |
|------------|-----------------|
| I-1        | A               |
| I-2        | A               |

| Compound # | JAK3<br>Inhibition |
|------------|--------------------|
| I-3        | A                  |
| I-4        | A                  |
| I-5        | A                  |
| I-6        | D                  |
| I-7        | C                  |
| I-8        | D                  |
| I-9        | C                  |
| I-10       | C                  |
| I-11       | E                  |
| I-12       | E                  |
| I-13       | C                  |
| I-14       | A                  |
| I-15       | B                  |
| I-16       | B                  |
| I-17       | B                  |
| I-18       | B                  |
| I-19       | C                  |
| I-46       | D                  |
| I-47       | E                  |
| I-48       | E                  |
| I-49       | A                  |
| I-50       | C                  |
| I-51       | E                  |
| I-52       | A                  |
| I-53       | C                  |
| I-54       | D                  |
| I-55       | B                  |
| I-56       | B                  |
| I-57       | E                  |
| I-58       | C                  |
| I-59       | D                  |
| I-60       | D                  |
| I-61       | B                  |
| I-62       | C                  |
| I-63       | B                  |
| I-64       | A                  |
| I-65       | A                  |
| I-66       | C                  |
| I-67       | A                  |
| I-68       | A                  |
| I-69       | C                  |
| I-70       | C                  |
| I-71       | D                  |


| <b>Compound #</b> | <b>JAK3<br/>Inhibition</b> |
|-------------------|----------------------------|
| <b>I-73</b>       | A                          |
| <b>I-75</b>       | C                          |
| <b>I-78</b>       | A                          |
| <b>I-79</b>       | A                          |
| <b>I-80</b>       | E                          |
| <b>I-81</b>       | A                          |
| <b>I-82</b>       | B                          |
| <b>I-83</b>       | B                          |
| <b>I-84</b>       | E                          |
| <b>I-85</b>       | D                          |
| <b>I-86</b>       | E                          |
| <b>I-92</b>       | B                          |
| <b>I-93</b>       | E                          |
| <b>I-94</b>       | A                          |

[00355] While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds and methods of this invention. Therefore, it will be appreciated that the scope of this invention is to be defined by the appended claims rather than by the specific embodiments that have been represented by way of example.

## CLAIMS


## We claim:

1. A compound of formula I:



or a pharmaceutically acceptable salt thereof, wherein:

Ring A is selected from:




Ring A is an optionally substituted group selected from phenyl, an 8-10 membered bicyclic partially unsaturated or aryl ring, a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-10 membered bicyclic heteroaryl ring having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Ring B is phenyl, a 5-6 membered heteroaryl ring having 1-3 heteroatoms independently selected from N, O or S, a 5-6 membered saturated heterocyclic ring having 1-2 heteroatoms independently selected from N, O or S, or an 8-10 membered bicyclic

partially unsaturated or aryl ring having 1-3 heteroatoms independently selected from N, O or S;

R<sup>1</sup> is selected from the group consisting of:



R<sup>1</sup> is a warhead group -L-Y, wherein:

L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and one or two methylene units of L are optionally and independently replaced by -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, -C(O)O-, cyclopropylene, -O-, -N(R)-, or -C(O)-; or

L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-; or

L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-; or

L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one alkylidenyl double bond and at least one methylene unit of L is replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one or two additional methylene units of L are optionally and independently replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-; or

L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one triple bond and one or two methylene units of L are optionally and independently replaced by –C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, –S-, -S(O)-, -SO<sub>2</sub>-, –O-, -N(R)-, -OC(O)-, or -C(O)O-; and


Y is hydrogen, C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or

L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein one methylene unit of L is replaced by cyclopropylene and one or two additional methylene units of L are independently replaced by –C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, –S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-; and

Y is C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN, or a 3-10 membered monocyclic or bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or


L is a covalent bond and Y is selected from:

- (ii) C<sub>2-6</sub> alkenyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (iii) C<sub>2-6</sub> alkynyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2 R<sup>e</sup> groups; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatoms selected from oxygen or nitrogen wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or




- (vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or
- (viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or


- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or



- (xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups; or



- (xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups; or




- (xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

L is  $-\text{C}(\text{O})-$  and Y is selected from:

- (i)  $\text{C}_{1-6}$  alkyl substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or
- (ii)  $\text{C}_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or
- (iii)  $\text{C}_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $\text{NO}_2$ , or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $\text{R}^e$  groups; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatoms selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $\text{R}^e$  groups; or

(vi)




- (vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $\text{R}^e$  groups; or

- (viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $\text{R}^e$  groups; or


- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $\text{R}^e$  groups; or

(x)




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $\text{R}^e$  groups; or

(xii)




- (xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $\text{R}^e$  groups; or

(xiv)



(xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups; or

(xvi)




(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

L is  $-N(R)C(O)-$  and Y is selected from:

- (i)  $C_{1-6}$  alkyl substituted with oxo, halogen,  $NO_2$ , or CN; or
- (ii)  $C_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iii)  $C_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $R^e$  groups; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatoms selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups; or

(vi)




(vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or

(viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or


(ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or

(x)




(xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

(xii)




(xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups; or

(xiv)



(xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups; or

(xvi)






- (xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

L is a bivalent  $C_{1-8}$  saturated or unsaturated, straight or branched, hydrocarbon chain; and Y is selected from:


- (ii)  $C_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iii)  $C_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $NO_2$ , or CN; or
- (iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $R^e$  groups; or
- (v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatoms selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups; or

(vi)




- (vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or
- (viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or
- (ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or

(x)




- (xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

(xii)




(xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups; or

(xiv)



(xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups; or

(xvi)



(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

L is  $-\text{CH}_2-$ ,  $-\text{NH}-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{CH}_2\text{NH}-$ ,  $-\text{NHCH}_2-$ ,  $-\text{NHC}(\text{O})-$ ,  $-\text{NHC}(\text{O})\text{CH}_2\text{OC}(\text{O})-$ ,  $-\text{CH}_2\text{NHC}(\text{O})-$ ,  $-\text{NHSO}_2-$ ,  $-\text{NHSO}_2\text{CH}_2-$ , or  $-\text{SO}_2\text{NH}-$ ; and Y is selected from:


(ii)  $\text{C}_{2-6}$  alkenyl optionally substituted with oxo, halogen,  $\text{NO}_2$ , or  $\text{CN}$ ; or

(iii)  $\text{C}_{2-6}$  alkynyl optionally substituted with oxo, halogen,  $\text{NO}_2$ , or  $\text{CN}$ ; or

(iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2  $R^e$  groups; or

(v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatoms selected from oxygen or nitrogen wherein said ring is substituted with 1-4  $R^e$  groups; or

(vi)




(vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or

(viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or


(ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4  $R^e$  groups; or

(x)




(xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups; or

(xii)




(xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4  $R^e$  groups; or


(xiv)



(xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups; or

(xvi)





(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups;

each  $R^e$  is independently selected from  $-Q-Z$ , oxo,  $NO_2$ , halogen, CN, diazonium,  $C_{1-6}$  aliphatic substituted with oxo, halogen,  $NO_2$ , or CN, or a group selected from  $C_{1-6}$  alkoxy, sulphonyloxy,  $C_{1-6}$  alkylsulphonyloxy,  $C_{2-6}$  alkenylsulphonyloxy, 5 to 14-membered arylsulphonyloxy, optionally substituted with oxo, halogen,  $NO_2$ , or CN, wherein:

Q is a bivalent  $C_{2-6}$  unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by  $-N(R)-$ ,  $-S-$ ,  $-O-$ ,  $-C(O)-$ ,  $-OC(O)-$ ,  $-C(O)O-$ ,  $-SO-$ ,  $-SO_2-$ ,  $-N(R)C(O)-$ ,  $-C(O)N(R)-$ ,  $-N(R)SO_2-$ , or  $-SO_2N(R)-$ ; and

Z is hydrogen or  $C_{1-6}$  aliphatic optionally substituted with oxo, halogen,  $NO_2$ , or CN;

$R^y$  is hydrogen, halogen, CN, lower alkyl, or lower haloalkyl;

W is a bivalent  $C_{1-3}$  alkylene chain wherein one methylene unit of W is optionally replaced by  $-NR^2-$ ,  $-N(R^2)C(O)-$ ,  $-C(O)N(R^2)-$ ,  $-N(R^2)SO_2-$ ,  $-SO_2N(R^2)-$ ,  $-O-$ ,  $-C(O)-$ ,  $-OC(O)-$ ,  $-C(O)O-$ ,  $-S-$ ,  $-SO-$  or  $-SO_2-$ ;

$R^2$  is hydrogen or optionally substituted  $C_{1-6}$  aliphatic, or:

$R^2$  and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated ring, or:

$R^2$  and  $R^y$  are taken together with their intervening atoms to form a 4-7 membered carbocyclic ring;


m is 0-4;

each  $R^x$  is independently selected from  $-R$ , halogen,  $-OR$ ,  $-CN$ ,  $-NO_2$ ,  $-SO_2R$ ,  $-SOR$ ,  $-C(O)R$ ,  $-CO_2R$ ,  $-C(O)N(R)_2$ ,  $-NRC(O)R$ ,  $-NRC(O)NR_2$ ,  $-NRSO_2R$ , or  $-N(R)_2$ ; or:

$R^x$  and  $R^1$  are taken together with their intervening atoms to form a 5-7 membered saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with a warhead group and 0-3 groups independently selected from oxo, halogen, CN, or  $C_{1-6}$  aliphatic; and

each R group is independently hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

2. The compound or pharmaceutically acceptable salt according to claim 1, wherein the compound is of formula **II**:



wherein:























W is  $-NR^2-$ ,  $-S-$ , or  $-O-$ ;

$R^2$  is hydrogen or optionally substituted  $C_{1-6}$  aliphatic, or:





















$R^2$  and a substituent on Ring A are taken together with their intervening atoms to form a 4-6 membered saturated ring; and

G is CH or N.














3. The compound or pharmaceutically acceptable salt according to claim 2, wherein Ring A is selected from:

*i**ii**iii**iv**v**vi**vii**viii**ix**x**xi**xii**xiii**xiv**xv**xvi**xvii**xviii**xix**xx**xxi**xxii*


182

*xxiii**xxiv**xxv**xxvi**xxvii**xxviii**xxix**xxx**xxxi**xxxii**xxxiii**xxxiv**xxxv**xxxvi**xxxvii**xxxviii**xxxix**xli**xlii**xlii*


183

*xliii**xliv**xlvi**xlvii**xlviii**l**li**lii**liii**liv**lv**lvii**lviii*

and


*lvi*

4. The compound or pharmaceutically acceptable salt according to claim 1, wherein Ring B is selected from:




5. The compound or pharmaceutically acceptable salt according to claim 2, wherein  $m$  is 1, 2, 3, or 4.

6. The compound according to claim 1, wherein said compound is of formula **III** or **IV**:




or a pharmaceutically acceptable salt thereof.

7. The compound according to claim 1, wherein said compound is of formula **I-i**, **I-ii**, or **I-iii**:




or a pharmaceutically acceptable salt thereof.





8. The compound according to claim 1, wherein said compound is of formula **II-i**, **II-ii**, or **II-iii**:



or a pharmaceutically acceptable salt thereof; wherein  
G is CH or N.





9. The compound according to claim 1, wherein said compound is of formula **III-a-i**, **III-b-i**, **III-a-ii**, **III-b-ii**, **III-a-iii**, or **III-b-iii**:





**III-a-i****III-b-i****III-a-ii****III-b-ii****III-a-iii****III-b-iii**

or a pharmaceutically acceptable salt thereof.







10. The compound according to claim 1, wherein said compound is of formula **IV-a-i**, **IV-b-i**, **IV-a-ii**, **IV-b-ii**, **IV-a-iii**, or **IV-b-iii**:

**IV-a-i****IV-b-i****IV-a-ii****IV-b-ii**

**IV-a-iii****IV-b-iii**

or a pharmaceutically acceptable salt thereof.

11. The compound according to claim 1, wherein said compound is of formula **II-a-iv**, **II-b-iv**, **II-a-v**, **II-b-v**, **II-a-vi**, or **II-b-vi**:

**II-a-iv****II-b-iv****II-a-v****II-b-v****II-a-vi****II-b-vi**

or a pharmaceutically acceptable salt thereof; wherein

G is CH or N.

12. The compound or pharmaceutically acceptable salt according to claim 1, wherein:  
L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and one or two methylene units of L are optionally and independently replaced by -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, -C(O)O-, cyclopropylene, -O-, -N(R)-, or -C(O)-.
13. The compound or pharmaceutically acceptable salt according to claim 12, wherein:  
L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one additional methylene unit of L is optionally replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-; and  
Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.
14. The compound or pharmaceutically acceptable salt according to claim 13, wherein L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -C(O)-, and one additional methylene unit of L is optionally replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-.
15. The compound or pharmaceutically acceptable salt according to claim 13, wherein L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one double bond and at least one methylene unit of L is replaced by -OC(O)-.
16. The compound or pharmaceutically acceptable salt according to claim 12, wherein L is -NRC(O)CH=CH-, -NRC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NRC(O)CH=CHCH<sub>2</sub>O-, -CH<sub>2</sub>NRC(O)CH=CH-, -NRSO<sub>2</sub>CH=CH-, -NRSO<sub>2</sub>CH=CHCH<sub>2</sub>- or -NRC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-; wherein the R group of L is H or optionally substituted C<sub>1-6</sub> aliphatic; and Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.
17. The compound or pharmaceutically acceptable salt according to claim 16, wherein L is -NHC(O)CH=CH-, -NHC(O)CH=CHCH<sub>2</sub>N(CH<sub>3</sub>)-, -NHC(O)CH=CHCH<sub>2</sub>O-

, -CH<sub>2</sub>NHC(O)CH=CH-, -NHSO<sub>2</sub>CH=CH-, -NHSO<sub>2</sub>CH=CHCH<sub>2</sub>- or -NHC(O)C(=CH<sub>2</sub>)CH<sub>2</sub>-

18. The compound or pharmaceutically acceptable salt according to claim 12, wherein L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one alkylidenyl double bond and at least one methylene unit of L is replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-, and one additional methylene unit of L is optionally replaced by cyclopropylene, -O-, -N(R)-, or -C(O)-.

19. The compound or pharmaceutically acceptable salt according to claim 1:  
L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein L has at least one triple bond and one or two methylene units of L are optionally and independently replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -O-, -N(R)-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-.

20. The compound or pharmaceutically acceptable salt according to claim 19, wherein Y is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.

21. The compound or pharmaceutically acceptable salt according to claim 20, wherein L is -C≡C-, -C≡CCH<sub>2</sub>N(isopropyl)-, -NHC(O)C≡CCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-, -C≡CCH<sub>2</sub>O-, -CH<sub>2</sub>C(O)C≡C-, -C(O)C≡C-, or -CH<sub>2</sub>OC(=O)C≡C-.

22. The compound or pharmaceutically acceptable salt according to claim 1, wherein:  
L is a bivalent C<sub>2-8</sub> straight or branched, hydrocarbon chain wherein one methylene unit of L is replaced by cyclopropylene and one or two additional methylene units of L are independently replaced by -C(O)-, -NRC(O)-, -C(O)NR-, -N(R)SO<sub>2</sub>-, -SO<sub>2</sub>N(R)-, -S-, -S(O)-, -SO<sub>2</sub>-, -OC(O)-, or -C(O)O-.

23. The compound or pharmaceutically acceptable salt according to claim 22, wherein Y is C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.

24. The compound or pharmaceutically acceptable salt according to claim 1, wherein:

L is a covalent bond, -C(O)-, -N(R)C(O)-, or a bivalent C<sub>1-8</sub> saturated or unsaturated, straight or branched, hydrocarbon chain; and


Y is selected from the following (ii) through (xvii):

(ii) C<sub>2-6</sub> alkenyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or

(iii) C<sub>2-6</sub> alkynyl optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN; or

(iv) a saturated 3-4 membered heterocyclic ring having 1 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-2 R<sup>e</sup> groups; or


(v) a saturated 5-6 membered heterocyclic ring having 1-2 heteroatom selected from oxygen or nitrogen wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or




(vii) a saturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or

(viii) a partially unsaturated 3-6 membered monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or


(ix) a partially unsaturated 3-6 membered carbocyclic ring, wherein said ring is substituted with 1-4 R<sup>e</sup> groups;



(xi) a partially unsaturated 4-6 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or




(xiii) a 6-membered aromatic ring having 0-2 nitrogens wherein said ring is substituted with 1-4 R<sup>e</sup> groups; or



; or

(xv) a 5-membered heteroaryl ring having 1-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-3  $R^e$  groups; or



(xvii) an 8-10 membered bicyclic, saturated, partially unsaturated, or aryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is substituted with 1-4  $R^e$  groups;

wherein:

each R group is independently hydrogen or an optionally substituted group selected from  $C_{1-6}$  aliphatic, phenyl, a 4-7 membered heterocyclic ring having 1-2 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

each  $R^e$  is independently selected from  $-Q-Z$ , oxo,  $NO_2$ , halogen, CN, diazonium,  $C_{1-6}$  aliphatic substituted with oxo, halogen,  $NO_2$ , or CN, or a group selected from  $C_{1-6}$  alkoxy,  $C_{1-6}$  sulphonyloxy,  $C_{1-6}$  alkylsulphonyloxy,  $C_{2-6}$  alkenylsulphonyloxy, 5 to 14-membered arylsulphonyloxy, optionally substituted with oxo, halogen,  $NO_2$ , or CN, wherein:


Q is a bivalent  $C_{2-6}$  unsaturated, straight or branched, hydrocarbon chain, wherein one or two methylene units of Q are optionally and independently replaced by  $-N(R)-$ ,  $-S-$ ,  $-O-$ ,  $-C(O)-$ ,  $-OC(O)-$ ,  $-C(O)O-$ ,  $-SO-$ , or  $-SO_2-$ ,  $-N(R)C(O)-$ ,  $-C(O)N(R)-$ ,  $-N(R)SO_2-$ , or  $-SO_2N(R)-$ ; and







































Z is hydrogen or C<sub>1-6</sub> aliphatic optionally substituted with oxo, halogen, NO<sub>2</sub>, or CN.































25. The compound or pharmaceutically acceptable salt according to claim 1, wherein L is a covalent bond, -CH<sub>2</sub>-, -NH-, -C(O)-, -CH<sub>2</sub>NH-, -NHCH<sub>2</sub>-, -NHC(O)-, -NHC(O)CH<sub>2</sub>OC(O)-, -CH<sub>2</sub>NHC(O)-, -NHSO<sub>2</sub>-, -NHSO<sub>2</sub>CH<sub>2</sub>-, or -SO<sub>2</sub>NH-.

26. The compound or pharmaceutically acceptable salt according to claim 25, wherein L is a covalent bond.


27. The compound or pharmaceutically acceptable salt according to any one of claims 24, 25, and 26, wherein Y is selected from:



***z******aa******bb******cc******dd******ee******ff******gg******hh******ii******jj******kk******ll******mm******nn******oo******pp******qq******rr******ss******tt******uu******vv******ww******xx******yy******zz******aaa******bbb******ccc******ddd******eee******fff******ggg******hhh******iii******jjj******kkk***


*lll**mmm**nnn**ooo**ppp**qqq**rrr**sss**ttt**uuu**vvv**qqq**www**xxx**yyy**zzz**aaaa**bbbb**cccc**dddd**eeee**ffff**gggg**hhhh**iiii**jjjj**kkkk**llll**mmmm**nnnn*


195







































wherein each  $R^e$  is independently halogen.


28. The compound or pharmaceutically acceptable salt according to claim 1, wherein  $R^1$  is selected from:

















*jji**kkk**lll**mmm**nnn**ooo**ppp**qqq**sss**ttt**uuu**vvv**www**xxx**yyy**zzz**aaaa**bbbb**cccc**dddd**eeee**ffff**gggg**hhhh**iii**jjj**kkkk*

*llll**mmmm**nnnn**oooo**pppp**qqqq**rrrr**ssss**tttt**uuuu**vvvv**wwww**xxxx**yyyy**zzzz**aaaaa**bbbbb**ccccc**ddddd**eeeee**ffffff**ggggg**hhhhh**iiiii**jjjjj**kkkkk**lllll**mmmmm**nnnnn**ooooo**ppppp**qqqqq**rrrrr**sssss**ttttt**uuuuu*















wherein each R<sup>e</sup> is independently a suitable leaving group selected from C<sub>1-6</sub> alkoxy, sulphonyloxy, optionally substituted C<sub>1-6</sub> alkylsulphonyloxy, optionally substituted C<sub>2-6</sub> alkenylsulphonyloxy, optionally substituted 5 to 14-membered arylsulphonyloxy, acyl, diazonium, NO<sub>2</sub>, CN, or oxo.













29. A compound that is:

**I-1****I-2****I-3****I-4****I-5****I-6****I-7****I-8****I-9****I-10****I-11****I-12**
















201

**I-13****I-14****I-15****I-16****I-17****I-18****I-19****I-20****I-21****I-22****I-23****I-24**






202












**I-25****I-26****I-27****I-28****I-29****I-30****I-31****I-32****I-33****I-34****I-35****I-36**


203

**I-37****I-38****I-39****I-40****I-41****I-42****I-43****I-44****I-45****I-46****I-47****I-48****I-49****I-50****I-51**

204




**I-66****I-67****I-68****I-69****I-70****I-71****I-72****I-73****I-74****I-75****I-76**



I-77


I-78

I-79




I-80


I-81




I-82



I-83













I-84




I-85

207

**I-86****I-87****I-88****I-89****I-90****I-91****I-92****I-93****I-94****I-95**


or

208

**I-96**

or a pharmaceutically acceptable salt thereof.


30. A compound that is:



**I-16**

or a pharmaceutically acceptable salt thereof.


31. A compound that is:



**I-17**

or a pharmaceutically acceptable salt thereof.


32. A compound that is:



**I-18**

or a pharmaceutically acceptable salt thereof.


33. A compound that is:



**I-19**

or a pharmaceutically acceptable salt thereof.


34. A compound that is:



**I-6**

or a pharmaceutically acceptable salt thereof.


35. A compound that is:



**I-66**

or a pharmaceutically acceptable salt thereof.


36. A compound that is:



**I-82**

or a pharmaceutically acceptable salt thereof.

37. A compound that is:



**I-89**

or a pharmaceutically acceptable salt thereof.

38. A composition comprising a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable adjuvant, carrier, or vehicle.

39. The composition according to claim 38, in combination with an additional therapeutic agent.

40. The composition according to claim 39, wherein the additional therapeutic agent is a chemotherapeutic agent.

41. The composition according to claim 38, 39, or 40 for inhibiting one or more of ErbB1, ErbB2, ErbB3, or ErbB4 kinases, or a mutant thereof, activity in a biological sample.

42. The composition according to claim 41, wherein the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity is inhibited irreversibly.

43. The composition according to claim 41 or 42, wherein the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys797 of ErbB1, Cys805 of ErbB2 or Cys803 of ErbB4.
44. The composition according to claim 38, 39, or 40 for inhibiting one or more of ErbB1, ErbB2, ErbB3, or ErbB4, or a mutant thereof, activity in a patient.
45. The composition according to claim 44 for irreversibly inhibiting the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity.
46. The composition according to claim 44 or 45 for irreversibly inhibiting the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity by covalent modification of Cys797 of ErbB1, Cys805 of ErbB2 or Cys803 of ErbB4.
47. The composition according to claim 38, 39, or 40 for treating an ErbB1-, ErbB2-, ErbB3-, or ErbB4-mediated disorder in a patient in need thereof.
48. The composition according to claim 47, wherein the disorder is a carcinoma selected from breast cancer, glioblastoma, lung cancer, cancer of the head and neck, colorectal cancer, bladder cancer, non-small cell lung cancer, squamous cell carcinoma, salivary gland carcinoma, ovarian carcinoma, or pancreatic cancer.
49. The composition according to claim 47, wherein the disorder is selected from neurofibromatosis type I (NF1), neurofibromatosis type II (NF2) Schwann cell neoplasms or a Schwannoma.
50. The composition according to claim 49, wherein the disorder is Schwann cell neoplasm and the Schwann cell neoplasm comprises malignant peripheral nerve sheath tumors (MPNST's).

51. The composition according to claim 38, 39, or 40 for inhibiting one or more TEC-kinases, or a mutant thereof, activity in a biological sample.
52. The composition according to claim 51, wherein the TEC-kinase, or a mutant thereof, activity is inhibited irreversibly.
53. The composition according to claim 51 or 52, where the one or more of TEC, ITK or BMX activity is inhibited irreversibly by covalently modifying Cys 449 of TEC, Cys 442 of ITK or Cys 496 of BMX.
54. The composition according to claim 38, 39, or 40 for inhibiting one or more TEC-kinases, or a mutant thereof, activity in a patient.
55. The composition according to claim 54 for irreversibly inhibiting the TEC-kinase, or a mutant thereof, activity.
56. The composition according to claim 54 or 55 for irreversibly inhibiting the one or more of TEC, ITK or BMX activity by covalent modification of Cys 449 of TEC, Cys 442 of ITK or Cys 496 of BMX.
57. The composition according to any one of claims 51 to 56 wherein the TEC-kinase is selected from one or more of TEC, ITK or BMX.
58. The composition according to claim 38, 39, or 40 for treating a TEC-kinase-mediated disorder in a patient in need thereof.
59. The composition according to claim 58, wherein the disorder is an autoimmune disorder, an inflammatory disorder, a proliferative disorder, a hyperproliferative disease, an immunological-mediated diseases, a disease of the respiratory tract, a disease of the bone and joints, a skin disorder, a gastrointestinal disorder, a systemic disease, or allograft rejection.

60. The composition according to claim 38, 39, or 40 for inhibiting BTK, or a mutant thereof, activity in a biological sample.
61. The composition according to claim 60, wherein the BTK, or a mutant thereof, activity is inhibited irreversibly.
62. The composition according to claim 60 or 61, wherein the BTK, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys 481 of BTK.
63. The composition according to claim 38, 39, or 40 for inhibiting BTK, or a mutant thereof, activity in a patient.
64. The composition according to claim 63 for irreversibly inhibiting the BTK, or a mutant thereof, activity.
65. The composition according to claim 63 or 64 for irreversibly inhibiting the BTK, or a mutant thereof, activity by covalent modification of Cys 481 of BTK.
66. The composition according to claim 38, 39, or 40 for treating a BTK-mediated disorder in a patient in need thereof.
67. The composition according to claim 66, wherein the disorder is an autoimmune disease, a heteroimmune disease, an inflammatory disease, a cancer, or a thromboembolic disorder.
68. The composition according to claim 67, wherein the disorder is selected from rheumatoid arthritis, multiple sclerosis, B-cell chronic lymphocytic leukemia, hairy cell leukemia, non-Hodgkin's lymphoma, irritable bowel syndrome, Crohn's disease, lupus or disorders associated with renal transplant.
69. The composition according to claim 38, 39, or 40 for inhibiting JAK3, or a mutant thereof, activity in a biological sample.

70. The composition according to claim 69, wherein the JAK3, or a mutant thereof, activity is inhibited irreversibly.
71. The composition according to claim 69 or 70, wherein the JAK3, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys 909 of JAK3.
72. The composition according to claim 38, 39, or 40 for inhibiting JAK3, or a mutant thereof, activity in a patient.
73. The composition according to claim 72 for irreversibly inhibiting the JAK3, or a mutant thereof, activity.
74. The composition according to claim 72 or 73 for irreversibly inhibiting the JAK3, or a mutant thereof, activity by covalent modification of Cys 909 of JAK3.
75. The composition according to claim 38, 39, or 40 for treating a JAK3-mediated disorder in a patient in need thereof.
76. The composition according to claim 75, wherein the disorder is selected from an autoimmune disorder, an inflammatory disorder, a neurodegenerative disorder, or a solid or hematologic malignancy.
77. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting one or more of ErbB1, ErbB2, ErbB3, or ErbB4 kinases, or a mutant thereof, activity in a biological sample.
78. The use according to claim 77, wherein the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity is inhibited irreversibly.

79. The use according to claim 77 or 78, wherein the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys797 of ErbB1, Cys805 of ErbB2 or Cys803 of ErbB4.

80. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting one or more of ErbB1, ErbB2, ErbB3, or ErbB4, or a mutant thereof, activity in a patient.

81. The use according to claim 80 for irreversibly inhibiting the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity.

82. The use according to claim 80 or 81 for irreversibly inhibiting the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity by covalent modification of Cys797 of ErbB1, Cys805 of ErbB2 or Cys803 of ErbB4.

83. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for treating an ErbB1-, ErbB2-, ErbB3-, or ErbB4-mediated disorder in a patient in need thereof.

84. The use according to claim 83, wherein the disorder is a carcinoma selected from breast cancer, glioblastoma, lung cancer, cancer of the head and neck, colorectal cancer, bladder cancer, non-small cell lung cancer, squamous cell carcinoma, salivary gland carcinoma, ovarian carcinoma, or pancreatic cancer.

85. The use according to claim 83, wherein the disorder is selected from neurofibromatosis type I (NF1), neurofibromatosis type II (NF2) Schwann cell neoplasms or a Schwannoma.

86. The use according to claim 85, wherein the disorder is Schwann cell neoplasm and the Schwann cell neoplasm comprises malignant peripheral nerve sheath tumors (MPNST's).

87. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting one or more TEC-kinases, or a mutant thereof, activity in a biological sample.

88. The use according to claim 87, wherein the TEC-kinase, or a mutant thereof, activity is inhibited irreversibly.

89. The use according to claim 87 or 88, where the one or more of TEC, ITK or BMX activity is inhibited irreversibly by covalently modifying Cys 449 of TEC, Cys 442 of ITK or Cys 496 of BMX.

90. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting one or more TEC-kinases, or a mutant thereof, activity in a patient.

91. The use according to claim 90 for irreversibly inhibiting the TEC-kinase, or a mutant thereof, activity.

92. The use according to claim 90 or 91 for irreversibly inhibiting the one or more of TEC, ITK or BMX activity by covalent modification of Cys 449 of TEC, Cys 442 of ITK or Cys 496 of BMX.

93. The use according to any one of claims 87 to 92 wherein the TEC-kinase is selected from one or more of TEC, ITK or BMX.

94. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for treating a TEC-kinase-mediated disorder in a patient in need thereof.

95. The use according to claim 94, wherein the disorder is an autoimmune disorder, an inflammatory disorder, a proliferative disorder, a hyperproliferative disease, an

immunological-mediated diseases, a disease of the respiratory tract, a disease of the bone and joints, a skin disorder, a gastrointestinal disorder, a systemic disease, or allograft rejection.

96. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting BTK, or a mutant thereof, activity in a biological sample.

97. The use according to claim 96, wherein the BTK, or a mutant thereof, activity is inhibited irreversibly.

98. The use according to claim 96 or 97, wherein the BTK, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys 481 of BTK.

99. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting BTK, or a mutant thereof, activity in a patient.

100. The use according to claim 99 for irreversibly inhibiting the BTK, or a mutant thereof, activity.

101. The use according to claim 99 or 100 for irreversibly inhibiting the BTK, or a mutant thereof, activity by covalent modification of Cys 481 of BTK.

102. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for treating a BTK-mediated disorder in a patient in need thereof.

103. The use according to claim 102, wherein the disorder is an autoimmune disease, a heteroimmune disease, an inflammatory disease, a cancer, or a thromboembolic disorder.

104. The use according to claim 103, wherein the disorder is selected from rheumatoid arthritis, multiple sclerosis, B-cell chronic lymphocytic leukemia, hairy cell leukemia, non-Hodgkin's lymphoma, irritable bowel syndrome, Crohn's disease, lupus or disorders associated with renal transplant.

105. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting JAK3, or a mutant thereof, activity in a biological sample.

106. The use according to claim 105, wherein the JAK3, or a mutant thereof, activity is inhibited irreversibly.

107. The use according to claim 105 or 106, wherein the JAK3, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys 909 of JAK3.

108. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for inhibiting JAK3, or a mutant thereof, activity in a patient.

109. The use according to claim 108 for irreversibly inhibiting the JAK3, or a mutant thereof, activity.

110. The use according to claim 108 or 109 for irreversibly inhibiting the JAK3, or a mutant thereof, activity by covalent modification of Cys 909 of JAK3.

111. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, for treating a JAK3-mediated disorder in a patient in need thereof.

112. The use according to claim 111, wherein the disorder is selected from an autoimmune disorder, an inflammatory disorder, a neurodegenerative disorder, or a solid or hematologic malignancy.

113. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for inhibiting one or more of ErbB1, ErbB2, ErbB3, or ErbB4, or a mutant thereof, activity in a patient.

114. The use according to claim 113, wherein the medicament is for irreversibly inhibiting the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity.

115. The use according to claim 113 or 114, wherein the medicament is for irreversibly inhibiting the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity by covalent modification of Cys797 of ErbB1, Cys805 of ErbB2 or Cys803 of ErbB4.

116. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating an ErbB1-, ErbB2-, ErbB3-, or ErbB4-mediated disorder in a patient in need thereof.

117. The use according to claim 116, wherein the disorder is a carcinoma selected from breast cancer, glioblastoma, lung cancer, cancer of the head and neck, colorectal cancer, bladder cancer, non-small cell lung cancer, squamous cell carcinoma, salivary gland carcinoma, ovarian carcinoma, or pancreatic cancer.

118. The use according to claim 116, wherein the disorder is selected from neurofibromatosis type I (NF1), neurofibromatosis type II (NF2) Schwann cell neoplasms or a Schwannoma.

119. The use according to claim 118, wherein the disorder is Schwann cell neoplasm and the Schwann cell neoplasm comprises malignant peripheral nerve sheath tumors (MPNST's).

120. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for inhibiting one or more TEC-kinases, or a mutant thereof, activity in a patient.

121. The use according to claim 120, wherein the medicament is for irreversibly inhibiting the TEC-kinase, or a mutant thereof, activity.

122. The use according to claim 120 or 121, wherein the medicament is for irreversibly inhibiting the one or more of TEC, ITK or BMX activity by covalent modification of Cys 449 of TEC, Cys 442 of ITK or Cys 496 of BMX.

123. The use according to any one of claims 120 to 122 wherein the TEC-kinase is selected from one or more of TEC, ITK or BMX.

124. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a TEC-kinase-mediated disorder in a patient in need thereof.

125. The use according to claim 124, wherein the disorder is an autoimmune disorder, an inflammatory disorder, a proliferative disorder, a hyperproliferative disease, an immunological-mediated diseases, a disease of the respiratory tract, a disease of the bone and joints, a skin disorder, a gastrointestinal disorder, a systemic disease, or allograft rejection.

126. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for inhibiting BTK, or a mutant thereof, activity in a patient.

127. The use according to claim 126, wherein the medicament is for irreversibly inhibiting the BTK, or a mutant thereof, activity.

128. The use according to claim 126 or 127, wherein the medicament is for irreversibly inhibiting the BTK, or a mutant thereof, activity by covalent modification of Cys 481 of BTK.

129. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a BTK-mediated disorder in a patient in need thereof.

130. The use according to claim 129, wherein the disorder is an autoimmune disease, a heteroimmune disease, an inflammatory disease, a cancer, or a thromboembolic disorder.

131. The use according to claim 130, wherein the disorder is selected from rheumatoid arthritis, multiple sclerosis, B-cell chronic lymphocytic leukemia, hairy cell leukemia, non-Hodgkin's lymphoma, irritable bowel syndrome, Crohn's disease, lupus or disorders associated with renal transplant.

132. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for inhibiting JAK3, or a mutant thereof, activity in a patient.

133. The use according to claim 132, wherein the medicament is for irreversibly inhibiting the JAK3, or a mutant thereof, activity.

134. The use according to claim 132 or 133, wherein the medicament is for irreversibly inhibiting the JAK3, or a mutant thereof, activity by covalent modification of Cys 909 of JAK3.

135. A use of a compound according to any one of claims 1 to 37, or a pharmaceutically acceptable salt thereof, in the preparation of a medicament for treating a JAK3-mediated disorder in a patient in need thereof.

136. The use according to claim 135, wherein the disorder is selected from an autoimmune disorder, an inflammatory disorder, a neurodegenerative disorder, or a solid or hematologic malignancy.

137. A method for inhibiting one or more of ErbB1, ErbB2, ErbB3, or ErbB4 kinases, or a mutant thereof, activity in a biological sample comprising the step of contacting said biological sample with a compound according to any one of claims 1 to 37 or a pharmaceutically acceptable salt thereof.

138. The method according to claim 137, wherein the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity is inhibited irreversibly.

139. The method according to claim 137 or 138, wherein the one or more of ErbB1, ErbB2, or ErbB4, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys797 of ErbB1, Cys805 of ErbB2 or Cys803 of ErbB4.

140. A method for inhibiting one or more TEC-kinases, or a mutant thereof, activity in a biological sample comprising the step of contacting said biological sample with a compound according to any one of claims 1 to 37 or a pharmaceutically acceptable salt thereof.

141. The method according to claim 140, wherein the TEC-kinase, or a mutant thereof, activity is inhibited irreversibly.

142. The method according to claim 140 or 141, where the one or more of TEC, ITK or BMX activity is inhibited irreversibly by covalently modifying Cys 449 of TEC, Cys 442 of ITK or Cys 496 of BMX.

143. The method according to any one of claims 140 to 142 wherein the TEC-kinase is selected from one or more of TEC, ITK or BMX.

144. A method for inhibiting BTK, or a mutant thereof, activity in a biological sample comprising the step of contacting said biological sample with a compound according to any one of claims 1 to 37 or a pharmaceutically acceptable salt thereof.

145. The method according to claim 144, wherein the BTK, or a mutant thereof, activity is inhibited irreversibly.

146. The method according to claim 144 or 145, wherein the BTK, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys 481 of BTK.

147. A method for inhibiting JAK3, or a mutant thereof, activity in a biological sample comprising the step of contacting said biological sample with a compound according to any one of claims 1 to 37 or a pharmaceutically acceptable salt thereof.

148. The method according to claim 147, wherein the JAK3, or a mutant thereof, activity is inhibited irreversibly.

149. The method according to claim 147 or 148, wherein the JAK3, or a mutant thereof, activity is inhibited irreversibly by covalently modifying Cys 909 of JAK3.











Figure 1

**Figure 2**

**Dose Response with I-16 and I-17 in A431 cells****EGFR Y1068 Phosphorylation****P42/p44 Erk phosphorylation****Figure 3**

**Dose Response with I-19 in A431 cells****I-19 Dose Response EGFR  
Phospho Y1068****I-19 Dose Response p42.p44 Erk  
phosphorylation****Figure 4**

**Figure 5**

**EGFR (Y1064) Phosphorylation in A431 Cells****Figure 6**




Figure 7




Figure 8

I-13 Shows Potent Inhibition of BTK Signaling in Ramos Cells




Figure 9





Figure 10

**TEC Kinase (I-13) Tryptic Digest Results****Figure 11**




Figure 12




Figure 13

SEQ ID 1: FULL LENGTH BTK PROTEIN:

MAAVILESIFLKRSQQKKKTSPLNFKKRLFLLTVHKLSYYEYDFERGRRGSKKGSIDVEK  
ITCVETVVPEKNPPPERQIPRRGEESSEMEQISIIERFPYPFQVVYDEGPLYVFSPTTEL  
RKRWIHQLKNVIRYNSDLVQKYHPCFWIDGQYLCCSQTAKNAMGCQILENRNGSLKPG  
SSHRKTKKPLPPTPEEDQILKKPLPPEPAAAPVSTSELKKVVALYDYMPMNANDLQLRK  
GDEYFILEESNLPWWRARDKNGQEGYIPSNYVTEAEDSIEMYEWYSKHMTRSQAELL  
KQEGKEGGFIVRDSSKAGKYTVSVFAKSTGDPQGVIRHYVVCSTPQSQYYLAEKHLFST  
IPELINYHQHNSAGLISRLKYPVSQQNKNAPSTAGLGYGSWEIDPKDLTFLKELGTGQFG  
VVKYGKWRGQYDVAIKMIKEGSMSEDEFIEEAKVMMNLSHEKLVQLYGVCTKQRPIFI  
TEYMANGCLLNLYLREMRHRFQTQQLLEMCKDVCEAMEYLESKQFLHRDLAARNCLVN  
DQGVVKVSDF

Cys=481

**Figure 14**

SEQ ID 2: FULL LENGTH TEC PROTEIN NP\_003206 631 aa

MNFNTILEEILIKRSQQKKKTSPLNYKERLFVLTKSMLTYYEGRAEKKYRKGFIDVSKIK  
CVEIVKNDDGVIPCQNKYPFQVVHDANTLYIFAPSPQSRDLWVKKLKEEIKNNNNNIMIK  
YHPKFWTDGSYQCCRQTEKLAPGCEKYNLFESSIRKALPPAPETKKRRPPPIPLEEEDNS  
EEIVVAMYDFQAAEGHDLRLERGQEYLILEKNDVHWWRARDKYGNEGYIPSNYVTGK  
KSNNLDQYEWYCRNMNRSKAEQLLRSEDKEGGFMVRDSSQPGLYTVSLYTKFGGEGS  
SGFRHYHIKETTTSPKKYYLAEKHAFGSIPEIIEYHKHNAAGLVTRLRYPVSVK GK  
NAPTTAGFSYEKWEINPSELTFMRELGSGLFGVVRLGKWRAQYKVAIKAIREGAMCEE  
DFIEEAKVMMKLTHPKLVQLYGVCTQQKPIYIVTEFMERGCLLNFLRQRQGHFSRDVLL  
SMCQDVCEGMEYLERNSFIHRDLAARNCLVSEAGVVKVSDFGMARYVLDDQYTSSSG  
AKFPVKWCPPEVFNYSRFSSKSDVWSFGVLMWEVFTGRMPFEKYTNYEVVTMVTRG  
HRLYQPKLASNYVYEVMLRCWQEKPEGRPSFEDLLRTIDELVECEETFGR

Cys=449

**Figure 15**

SEQ ID 3: FULL LENGTH ITK PROTEIN NP 005537 620aa

MNNFILLEEQLIKKSQQKRRTSPSNFKVRFFVLTKASLAYFEDRHGKKRTLKGSIELSRIKCVE  
IVKSDISIPCHYKYPFQVVHDNYLLYVFAPDRESRQRWVLALKEETRNNNSLVPKYHPNFWMDG  
KWRCCSQLEKLATGCAQYDPTKNASKKPLPPTPEDNRRPLWEPEETVVIALYDYQTNDPQELAL  
RRNEEYCLLDSSEIHWVRVQDRNGHEGYVPSSYLVEKSPNNLETYEWYNKSI SRDKAEKLLLD  
GKEGAFMVRDSRTAGTYTVSVFTKAVVSENNPCIKHYHIKETNDNPKRYYVAEKYVFDSIPLLI  
NYHQHNGGGLVTRLRYPVCFGRQKAPVTAGLRYGKWVIDPSELTFVQEIGSGQFGLVHLGYWLN  
KDKVAIKTIREGAMSEEDFIEEAEVMMKLSHPKLVQLYGVCLEQAPICLVFEFMEHGCLSDYLR  
TQRGLFAAETLLGMCLDVCEGMAYLEEACVIHRDLAARNCLVGENQVIKVSDFGMTRFVLDDQY  
TSSTGTKFPVKWASPEVFSFSRYSSKSDVWSFGVLMWEVFSEGKIPYENRSNSEVVEDI STGFR  
LYKPRLASTHVYQIMNHCWKERPEDRPAFSRLRLRLAEIAESGL

Cys=442

**Figure 16**

SEQ ID 4: FULL LENGTH BMX PROTEIN NP 001712 675aa

MDTKSILEELLLKRSQQKKKMSPNNYKERLFVLTKTNLSYYEYDKMKRGSRKGSIEIKK  
IRCVEKVNLEEQTPVERQYPFQIVYKDGLLYVYASNEESRSQWLKALQKEIRGNPHLLV  
KYHSGFFVDGKFLCCQQSCKAAPGCTLWEAYANLHTAVNEEKHRVPTFPDRVLKIPRA  
VPVLKMDAPSSSTTLAQYDNESKKNYGSQPPSSSTSLAQYDSNSKKIYGSQPNFNMQYIP  
REDFPDWWQVRKLKSSSSSEDEVASSNQKERNVNHTTSKISWEFPESSSSEEEENLDDYD  
WFAGNISRSQSEQLLRQKGKEGAFMVRNSSQVGMVTVSLFSKAVNDKKGTVKHYYH  
VHTNAENKLYLAENYCFDSIPKLIHYHQHNSAGMITRLRHPVSTKANKVPDSVSLGNGI  
WELKREEITLLKELGSGQFGVVQLGKWKGQYDVAVKMIKEGSMSEDEFFQEAQTMMK  
LSHPKLVKFYGVCSKEYPIYIVTEYISNGCLLNYLRSHGKGLEPSQLLEM CYDVCEGMA  
FLESHQFIHRDLAARNCLVDRDLCVKVSDFGMTRYVLDDQYVSSVGTKFPVKWSAPEV  
FHYFKYSSKSDVWAFGILMWEVFSLGKQPYDLYDNSQVVLKVSQGHRLYRPHLASDTI  
YQIMYSCWHELPEKRPTFQQLSSIEPLREKDKH

Cys=496

**Figure 17**

SEQ ID 5: FULL LENGTH JAK3 PROTEIN NP 000206 1124 aa

MAPPSEETPLIPQRSCSLLSTEAGALHVLLPARGPGPPQRLSFSFGDHLAEDLCVQAAKASGIL  
PVIHSLFALATEDLSCWFPPSHIFSVEDASTQVLLYRIRFYFPNWFGLEKCHRFGLRKDLASAI  
LDLPVLEHLFAQHRSDLVSGRLPVGLSLKEQGECLESLAVLDLARMAREQAQRPGELLKTVSYKA  
CLPPSLRDLIQGLSFVTRRRIRRTVRRALRRVAACQADRHSIMAKYIMDLERLDPAGAAETFHV  
GLPGALGGHDGLGLLRVAGDGGIAWTQGEQEVLPFCDFPEIVDISIKQAPRVGPAGEHRLVTV  
TRTDNQILEAEFPGLPEALSFVALVDGYFRLTTDSQHFFCKEVAPPRLLEEVAEQCHGPITLDF  
AINKLKTGGSRPGSYVLRSPQDFDSFLLTVCVQNPLGPDYKGCLIRRSPTGTFLLVGLSRPHS  
SLRELLATCWDGGLHVDGVAVTLTSCCIPRPKEKSNLIVVQRGHSPTSSLVQPQSQYQLSQMT  
FHKIPADSLEWHENLGHGSFTKIYRGCRHEVVDGEARKTEVLLKVMDAKHKNCMESFLEAASLM  
SQVSYRHLVLLHGVCMAGDSTMVQEFVHLGAIDMYLRKRGHLVPASWKLQVVKQLAYALNYLED  
KGLPHGNVSARKVLLAREGADGSPFFIKLSDPGVSPAVLSLEMLTDRI PWVAPECLREAQTLSL  
EADKWGFGATVWEVFSGVTMPI SALDPAKKLQFYEDRQQLPAPKWTELALLIQQCMAYEPVQRP  
SFRAVIRDLNSLISSDYELLSDPTPGALAPRDGLWNGAQLYACQDPTIFEERHLKYISQLGKGN  
FGSVELCRYDPLGDNTGALVAVKQLQHSGPDQQRDFQREIQILKALHSDFIVKYRGVSYGPGRQ  
SLRLVMEYLPSCCLRDFLQRHRARLDASRLLLYSSQICKGMEYLGSRRCVHRDLAARNILVESE  
AHVKIADFGLAKLLPLDKDYYVVREPGQSPIFWYAPESLSDNIFSRQSDVWSFGVVLYELFTYC  
DKSCSPSAEFLRMMGCERDVPALCRLLELLEEGQRLPAPPACPAEVHELMKLCWAPSPQDRPSF  
SALGPQLDMLWSGSRGCETHAFTAHPGKHHSLSFS

Cys=909


**Figure 18**

SEQ ID 6: FULL LENGTH TXK PROTEIN NP\_003319 527 aa

MILSSYNTIQSVFCCCCCSCVQKRQMRTQISLSTDEELPEKYTQRRRPWLSQLSNKKQSN  
TGRVQPSKRKPLPPLPPSEVAEEKIQVKALYDFLPREPCNLALRRAEEYLILEKYNPHWW  
KARDRLGNEGLIPSNYVTENKITNLEIYEWYHRNITRNQAEHLLRQESKEGAFIVRDSRH  
LGSYTISVFMGARRSTEAAIKHYQIKKND SGQWYVAERHAFQSIPELIWYHQHNAAGL  
MTRLRYPVGLMGSCLPATAGFSYEKWEIDPSELAFIKEIGSGQFGVVHLGEWRSHIQVAI  
KAINEGSMSEEDFIEEAKVMMKLSHSLVQLYGVCIQRKPLYIVTEFMENGCLLNYLRE  
NKGKLRKEMLLSVCQDICEGMEYLERNGYIHRDLAARNCLVSSTCIVKISDFGMTRYVL  
DDEYVSSFGAKFPIKWSPPEVFLFNKYSSKSDVWSFGVLMWEVFTEGKMPFENKSNLQ  
VVEAISEGFRLYRPHLAPMSIYEV MYSCWHEKPEGRPTFAELLRAVTEIAETW

Cys 350

**Figure 19**



**I**