




(86) Date de dépôt PCT/PCT Filing Date: 2011/03/30
 (87) Date publication PCT/PCT Publication Date: 2011/10/06
 (45) Date de délivrance/Issue Date: 2021/11/16
 (85) Entrée phase nationale/National Entry: 2012/09/28
 (86) N° demande PCT/PCT Application No.: US 2011/030598
 (87) N° publication PCT/PCT Publication No.: 2011/123572
 (30) Priorités/Priorities: 2010/03/31 (US61/319,494);
 2010/10/15 (US61/393,724); 2010/10/26 (US61/406,892)

(51) Cl.Int./Int.Cl. *C12N 5/071* (2010.01),
A61K 35/12 (2015.01), *C12N 5/00* (2006.01),
C12N 5/0735 (2010.01), *C12N 9/12* (2006.01)
 (72) Inventeurs/Inventors:
 ZHU, SAIYONG, US;
 DING, SHENG, US
 (73) Propriétaire/Owner:
 THE SCRIPPS RESEARCH INSTITUTE, US
 (74) Agent: SMART & BIGGAR IP AGENCY CO.

(54) Titre : NOUVELLE PROGRAMMATION DE CELLULES
 (54) Title: REPROGRAMMING CELLS



(57) Abrégé/Abstract:

The present invention provides for methods, compositions, and kits for producing an induced pluripotent stem cell from a non-pluripotent mammalian cell using a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator or a compound that promotes glycolytic metabolism as well as other small molecules.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 October 2011 (06.10.2011)

(10) International Publication Number
WO 2011/123572 A1

- (51) **International Patent Classification:**
C12N 5/00 (2006.01)
- (21) **International Application Number:**
PCT/US2011/030598
- (22) **International Filing Date:**
30 March 2011 (30.03.2011)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/319,494 31 March 2010 (31.03.2010) US
61/393,724 15 October 2010 (15.10.2010) US
61/406,892 26 October 2010 (26.10.2010) US
- (71) **Applicant (for all designated States except US):** THE SCRIPPS RESEARCH INSTITUTE [US/US]; 10550 North Torrey Pines Road, Mail TPC-8, La Jolla, California 92037 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** ZHU, Saiyong [CN/US]; 10550 North Torrey Pines Road, Mail TPC-8, La Jolla, California 92037 (US). DING, Sheng [CN/US]; 10550 North Torrey Pines Road, Mail TPC-8, La Jolla, California 92037 (US).
- (74) **Agents:** FLANAGAN, Lisa Dornbach et al.; Kilpatrick Townsend & Stockton LLP, Two Embarcadero Center, Eighth Floor, San Francisco, California 94111-3834 (US).

- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) **Title:** REPROGRAMMING CELLS




Fig. 1

(57) **Abstract:** The present invention provides for methods, compositions, and kits for producing an induced pluripotent stem cell from a non-pluripotent mammalian cell using a 3'-phosphoinositide-dependent kinase-1 (PKD1) activator or a compound that promotes glycolytic metabolism as well as other small molecules.

WO 2011/123572 A1

REPROGRAMMING CELLS

CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] The present application claims benefit of priority to U.S. Provisional Application No. 61/319,494, filed March 31, 2010, U.S. Provisional Application No. 61/393,724, filed October 15, 2010, and U.S. Provisional Application No. 61/406,892, filed October 26, 2010.

BACKGROUND

[0002] Induced pluripotent stem cell (iPSC) technology, i.e. reprogramming somatic cells into pluripotent cells that closely resemble embryonic stem cells (ESCs) by introduction of defined factors, holds great potential in biomedical research and regenerative medicine (Takahashi, K., and Yamanaka, S., *Cell* 126, 663-676 (2006); Takahashi et al., *Cell* 131, 861-872 (2007); Yu et al., *Science* 318, 1917-1920 (2007); Zhou et al., *Cell Stem Cell* 4, 381-384 (2009); Kim et al., *Cell Stem Cell* 4, 472-476 (2009); Maherali, N., and Hochedlinger, K., *Cell Stem Cell* 3, 595-605 (2009a); Daley et al., *Cell Stem Cell* 4, 200-201 (2009)). Various strategies have been developed to generate iPSCs with less or no exogenous genetic manipulations, which represent a major hurdle for iPSC applications (Yamanaka et al., 2009; Saha, K., Jaenisch, R., *Cell Stem Cell* 5, 584-595 (2009)). Toward an ultimate goal of generating iPSCs with a defined small molecule cocktail that would offer significant advantages over genetic manipulations or more difficult-to-manufacture/use biologics, substantial efforts have been made in identifying chemical compounds that can functionally replace exogenous reprogramming transcription factors (TFs) and/or enhance reprogramming efficiency and kinetics (Shi et al., *Cell Stem Cell* 2, 525-528 (2008a); Shi et al., *Cell Stem Cell* 3, 568-574 (2008b); Huangfu et al., *Nat Biotechnol* 26, 795-797 (2008a); Huangfu et al., *Nat Biotechnol* 26, 1269-1275 (2008b); Silva et al., *Plos Bio* 6, e253. doi: 10.1371/journal.pbio.0060253 (2008); Lyssiotis et al., *PNAS* 106, 8912-8917 (2009); Ichida et al., *Cell Stem Cell* 5, 491-503 (2009); Maherali, N., Hochedlinger, K., *Curr Biol* 19, 1718-1723 (2009b); Esteban et al., *Cell Stem Cell* 6, 71-79 (2010); Feng et al., *Cell Stem Cell* 4, 301-312 (2009)). However, further reducing the number of exogenous TFs has been extraordinarily challenging as (1) most reprogramming enabling or enhancing conditions (e.g., exploiting a specific cell type or using small molecules) are context dependent, i.e., such specific conditions (e.g., a reprogramming small molecule) typically would be much less effective or even harmful in a

different cell type with different exogenous factors and used in a different window of treatment; and (2) high throughput screening is technically challenging when the reprogramming efficiency and speed further decrease exponentially due to fewer exogenous TFs used. To date, only neural stem cells (NSCs) that endogenously express Sox2 and cMyc at a high level were shown to be reprogrammed to iPSCs by exogenous expression of only Oct4 (Kim et al., Cell 136, 411-419 (2009a); Kim et al., Nature 461, 643-649 (2009b)). However, human fetal NSCs are rare and practically difficult to obtain (Nunes et al., Nat Med 9, 439-447 (2003)). Consequently, it would be beneficial to develop chemical reprogramming conditions applicable to other more accessible and abundant somatic cells.

BRIEF SUMMARY

[0003] The present disclosure provides for a method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell. In some embodiments, the method comprises contacting the non-pluripotent cell with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient to induce the cell to become a pluripotent stem cell. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, e.g., (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid ("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z").

[0004] In some embodiments, the method further comprises contacting the non-pluripotent cell with a TGF β receptor/ALK5 inhibitor, e.g., A-83-01. In some embodiments, the method further comprises contacting the non-pluripotent cell with a MEK inhibitor, e.g., PD0325901. In some embodiments, the method further comprises contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, e.g., sodium butyrate (NaB), or valproic acid (VPA).

[0005] In some embodiments, the method comprises contacting the non-pluripotent cell with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient to induce the cell to become a pluripotent stem cell. In some embodiments, the conditions comprise introducing at least one exogenous transcription factor into the non-pluripotent cell. In some embodiments, the exogenous transcription factor comprises a polypeptide. In some

embodiments, the exogenous transcription factor comprises an Oct polypeptide. In some
embodiments, the exogenous transcription factor comprises a protein selected from the group
consisting of an Oct polypeptide and a Klf polypeptide. In some embodiments, the
exogenous transcription factor comprises a protein selected from the group consisting of an
5 Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some
embodiments, the condition comprises introducing at least two, three or four exogenous
transcription factors into the non-pluripotent cell, wherein the exogenous transcription factors
each comprise a different protein selected from the group consisting of an Oct polypeptide, a
Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the
10 exogenous transcription factor is introduced by introducing a polynucleotide into the non-
pluripotent cell, wherein the polynucleotide encodes the exogenous transcription factor,
thereby expressing the transcription factor(s) in the non-pluripotent cell. In some
embodiments, the exogenous transcription factor is introduced by contacting the exogenous
transcription factor to the non-pluripotent cell. In some embodiments, the exogenous
15 transcription factor comprises an amino acid sequence that enhances transport across cell
membranes.

[0006] In some embodiments, the non-pluripotent cell is a human cell. In some
embodiments, the PDK1 activator is present in a concentration sufficient to improve by at
least 10% the efficiency of induction of the non-pluripotent cell into an induced pluripotent
20 stem cell, under conditions sufficient to induce conversion of the non-pluripotent cell into the
induced pluripotent stem cell.

[0007] In some embodiments, the method comprises contacting the non-pluripotent cell
with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient
to induce the cell to become a pluripotent stem cell. In some embodiments, the method
25 comprises contacting the non-pluripotent cell with a PDK1 activator in the absence of a MEK
inhibitor, followed by contacting the non-pluripotent cell with a PDK1 activator and a MEK
inhibitor. In some embodiments, the method comprises contacting the non-pluripotent cell
with a PDK1 activator, a TGF β receptor/ALK5 inhibitor, and a histone deacetylase (HDAC)
inhibitor in the absence of a MEK inhibitor, followed by contacting the non-pluripotent cell
30 with a PDK1 activator, a TGF β receptor/ALK5 inhibitor, a histone deacetylase (HDAC)
inhibitor and a MEK inhibitor.

[0008] In some embodiments, the method further comprises purifying the pluripotent cells
to generate a homogenous population of the pluripotent cells. In some embodiments, wherein

a plurality of pluripotent stem cells are induced, the method further comprises purifying the pluripotent stem cells to generate a homogenous population of pluripotent stem cells.

[0009] In another aspect, the present disclosure provides for a mixture comprising: mammalian cells, a PDK1 activator, and one or more of (1) a TGF β receptor/ALK5 inhibitor; (2) a MEK inhibitor; (3) a histone deacetylase (HDAC) inhibitor; or (4) one or more exogenous transcription factors selected from the group consisting of an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide.

[0010] In some embodiments, at least 99% of the cells in the mixture are non-pluripotent cells. In some embodiments, essentially all of the cells are non-pluripotent cells. In some embodiments, the cells are human cells. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, e.g., (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid ("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z"). In some embodiments, the mixture further comprises a TGF β receptor/ALK5 inhibitor, e.g., A-83-01. In some embodiments, the mixture further comprises a MEK inhibitor, e.g., PD0325901. In some embodiments, the mixture further comprises a histone deacetylase (HDAC) inhibitor, e.g., sodium butyrate (NaB), or valproic acid (VPA).

[0011] In some embodiments, the mixture further comprises an exogenous transcription factor selected from an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the exogenous transcription factor comprises an amino acid sequence that enhances transport across cell membranes. In some embodiments, the PDK1 activator in the mixture is present in a concentration sufficient to improve by at least 10% the efficiency of induction of non-pluripotent cells in the mixture into induced pluripotent stem cells under conditions sufficient to induce conversion of the cells into induced pluripotent stem cells.

[0012] In still another aspect, the present disclosure provides for a kit for inducing pluripotency in a non-pluripotent mammalian cell, the kit comprising a PDK1 activator, and one or more of (1) a TGF β receptor/ALK5 inhibitor; (2) a MEK inhibitor; (3) a histone deacetylase (HDAC) inhibitor; or (4) one or more transcription factors selected from the group consisting of an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, e.g., (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid

("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z"). In some embodiments, the kit further comprises a TGF β receptor/ALK5 inhibitor, e.g., A-83-01. In some embodiments, the kit further comprises a MEK inhibitor, e.g., PD0325901. In some embodiments, the kit further comprises a histone deacetylase (HDAC) inhibitor, e.g., sodium butyrate (NaB), or valproic acid (VPA).

[0013] In some embodiments, the kit further comprises an exogenous transcription factor selected from an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the exogenous transcription factor comprises an amino acid sequence that enhances transport across cell membranes.

[0014] In yet another aspect, the present disclosure provides a method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell. In some embodiments, the method comprises contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism under conditions sufficient to induce the cell to become a pluripotent stem cell, thereby inducing the non-pluripotent mammalian cell into an induced pluripotent stem cell. In some embodiments, the compound that promotes glycolytic metabolism is a PDK1 activator. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, e.g., PS48, PS08, 12Z, or 13Z. In some embodiments, the compound that promotes glycolytic metabolism is a glycolysis activator, e.g., fructose 2, 6-bisphosphate. In some embodiments, the compound that promotes glycolytic metabolism is a substrate for glycolysis, e.g., fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glycolytic intermediate or its metabolic precursors, e.g., nicotinic acid, NADH, or fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glucose uptake transporter activator. In some embodiments, the compound that promotes glycolytic metabolism is a mitochondrial respiration modulator. In some embodiments, the mitochondrial respiration modulator is an oxidative phosphorylation inhibitor, e.g., 2, 4-dinitrophenol, or 2-hydroxyglutaric acid. In some embodiments, the compound that promotes glycolytic metabolism is a hypoxia-inducible factor activator, e.g., N-oxaloylglycine, or quercetin. In some embodiments, the method further comprises contacting the non-pluripotent cell with a TGF β receptor/ALK5 inhibitor, e.g., A-83-01. In some embodiments, the method further comprises contacting the non-pluripotent cell with a MEK inhibitor, e.g., PD0325901. In some embodiments, the method further comprises

contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, *e.g.*, sodium butyrate (NaB), or valproic acid (VPA).

[0015] In some embodiments, the method comprises contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism under conditions sufficient to induce the cell to become a pluripotent stem cell. In some embodiments, the conditions comprise introducing at least one exogenous transcription factor into the non-pluripotent cell. In some embodiments, the exogenous transcription factor comprises a polypeptide. In some embodiments, the exogenous transcription factor comprises an Oct polypeptide. In some embodiments, the exogenous transcription factor comprises a protein selected from the group consisting of an Oct polypeptide and a Klf polypeptide. In some embodiments, the exogenous transcription factor comprises a protein selected from the group consisting of an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the condition comprises introducing at least two, three or four exogenous transcription factors into the non-pluripotent cell, wherein the exogenous transcription factors each comprise a different protein selected from the group consisting of an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the exogenous transcription factor is introduced by introducing a polynucleotide into the non-pluripotent cell, wherein the polynucleotide encodes the exogenous transcription factor, thereby expressing the transcription factor(s) in the non-pluripotent cell. In some embodiments, the exogenous transcription factor is introduced by contacting the exogenous transcription factor to the non-pluripotent cell. In some embodiments, the exogenous transcription factor comprises an amino acid sequence that enhances transport across cell membranes.

[0016] In some embodiments, the non-pluripotent cell is a human cell. In some embodiments, the compound that promotes glycolytic metabolism is present in a concentration sufficient to improve by at least 10% the efficiency of induction of the non-pluripotent cell into an induced pluripotent stem cell, under conditions sufficient to induce conversion of the non-pluripotent cell into the induced pluripotent stem cell.

[0017] In some embodiments, the method comprises contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism under conditions sufficient to induce the cell to become a pluripotent stem cell. In some embodiments, the method comprises contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism and a MEK inhibitor. In some embodiments, the method comprises contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism in the absence of a

MEK inhibitor, followed by contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism and a MEK inhibitor. In some embodiments, the method comprises contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism, a TGF β receptor/ALK5 inhibitor, and a histone deacetylase (HDAC) inhibitor in the absence of a MEK inhibitor, followed by contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism, a TGF β receptor/ALK5 inhibitor, a histone deacetylase (HDAC) inhibitor and a MEK inhibitor.

[0018] In some embodiments, the method further comprises purifying the pluripotent cells to generate a homogenous population of the pluripotent cells. In some embodiments, wherein a plurality of pluripotent stem cells are induced, the method further comprises purifying the pluripotent stem cells to generate a homogenous population of pluripotent stem cells.

[0019] In still another aspect, the present disclosure provides for a mixture comprising: mammalian cells, a compound that promotes glycolytic metabolism, and one or more of (1) a TGF β receptor/ALK5 inhibitor; (2) a MEK inhibitor; (3) a histone deacetylase (HDAC) inhibitor; or (4) one or more exogenous polypeptides selected from the group consisting of an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, at least 99% of the cells in the mixture are initially non-pluripotent cells. In some embodiments, essentially all of the cells are initially non-pluripotent cells. In some embodiments, the cells are human cells. In some embodiments, the compound that promotes glycolytic metabolism is a PDK1 activator. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, e.g., PS48, PS08, 12Z, or 13Z. In some embodiments, the compound that promotes glycolytic metabolism is a glycolysis activator, e.g., fructose 2, 6-bisphosphate. In some embodiments, the compound that promotes glycolytic metabolism is a substrate for glycolysis, e.g., fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glycolytic intermediate or its metabolic precursors, e.g., nicotinic acid, NADH, or fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glucose uptake transporter activator. In some embodiments, the compound that promotes glycolytic metabolism is a mitochondrial respiration modulator. In some embodiments, the mitochondrial respiration modulator is an oxidative phosphorylation inhibitor, e.g., 2, 4-dinitrophenol, or 2-hydroxyglutaric acid. In some embodiments, the compound that promotes glycolytic metabolism is a hypoxia-inducible factor activator, e.g., N-oxaloylglycine, or quercetin. In some embodiments, the mixture further

comprises a TGF β receptor/ALK5 inhibitor, e.g., A-83-01. In some embodiments, the mixture further comprises a MEK inhibitor, e.g., PD0325901. In some embodiments, the mixture further comprises a histone deacetylase (HDAC) inhibitor, e.g., sodium butyrate (NaB), or valproic acid (VPA).

[0020] In some embodiments, the exogenous transcription factor comprises an amino acid sequence that enhances transport across cell membranes. In some embodiments, the compound that promotes glycolytic metabolism in the mixture is present in a concentration sufficient to improve by at least 10% the efficiency of induction of non-pluripotent cells in the mixture into induced pluripotent stem cells under conditions sufficient to induce conversion of the cells into induced pluripotent stem cells.

[0021] In yet another aspect, the present disclosure provides for a kit for inducing pluripotency in a non-pluripotent mammalian cell, the kit comprising a compound that promotes glycolytic metabolism, and one or more of (1) a TGF β receptor/ALK5 inhibitor; (2) a MEK inhibitor; (3) a histone deacetylase (HDAC) inhibitor; or (4) one or more transcription factors selected from the group consisting of an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide; or a polynucleotide encoding a transcription factor selected from an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the compound that promotes glycolytic metabolism is a PDK1 activator. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, e.g., PS48, PS08, 12Z, or 13Z. In some embodiments, the compound that promotes glycolytic metabolism is a glycolysis activator, e.g., fructose 2,6-bisphosphate. In some embodiments, the compound that promotes glycolytic metabolism is a substrate for glycolysis, e.g., fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glycolytic intermediate or its metabolic precursors, e.g., nicotinic acid, NADH, or fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glucose uptake transporter activator. In some embodiments, the compound that promotes glycolytic metabolism is a mitochondrial respiration modulator. In some embodiments, the mitochondrial respiration modulator is an oxidative phosphorylation inhibitor, e.g., 2, 4-dinitrophenol, or 2-hydroxyglutaric acid. In some embodiments, the compound that promotes glycolytic metabolism is a hypoxia-inducible factor activator, e.g., N-oxaloylglycine, or quercetin. In some embodiments, the kit further comprises a TGF β receptor/ALK5 inhibitor, e.g., A-83-01. In some embodiments, the kit further comprises a MEK inhibitor, e.g., PD0325901.

In some embodiments, the kit further comprises a histone deacetylase (HDAC) inhibitor, e.g., sodium butyrate (NaB), or valproic acid (VPA). In some embodiments, the exogenous transcription factor comprises an amino acid sequence that enhances transport across cell membranes.

[0021a] Various embodiments of the claimed invention pertain to an *in vitro* method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell, the method comprising: (a) contacting the non-pluripotent cell with an Oct4 polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding an Oct4 polypeptide; (b) contacting the non-pluripotent cell with (i) a TGF β receptor/ALK5 inhibitor or a MEK inhibitor, or (ii) a TGF β receptor/ALK5 inhibitor and a MEK inhibitor; (c) contacting the non-pluripotent cell with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient to induce the cell to become a pluripotent stem cell, thereby inducing the non-pluripotent mammalian cell into an induced pluripotent stem cell; and (d) one or more of (i) contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, (ii) contacting the non-pluripotent cell with a Klf polypeptide or (iii) introducing into the non-pluripotent cell a polynucleotide encoding a Klf polypeptide.

[0021b] Various embodiments of the claimed invention also pertain to a mixture comprising: mammalian cells, a PDK1 activator, an Oct4 polypeptide, and: (a) a TGF β receptor/ ALK5 inhibitor, a MEK inhibitor, or a combination thereof; and (b) one or more of (i) a histone deacetylase (HDAC) inhibitor, (ii) a Klf polypeptide, or (iii) a polynucleotide encoding a Klf polypeptide.

[0021c] Various embodiments of the claimed invention also pertain to an *in vitro* method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell, the method comprising: (a) contacting the non-pluripotent cell with an Oct4 polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding an Oct4 polypeptide; (b) contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism, a small molecule TGF β receptor/ALK5 inhibitor, and a small molecule MEK inhibitor, thereby inducing the non-pluripotent mammalian cell to reprogram into an induced pluripotent stem cell, wherein the compound that promotes glycolytic metabolism comprises F2,6P (fructose 2,6-bisphosphate), F6P (fructose 6-phosphate), DNP (2,4-dinitrophenol), NOG (N-oxalylglycine), QC (quercetin), 2-HA (2-hydroxyglutaric acid), NA (nicotinic acid), or a PDK1 activator; and (c) one or more of (i) contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, (ii) contacting the non-pluripotent cell with a Klf polypeptide or (iii) introducing into the non-pluripotent cell a polynucleotide encoding a Klf polypeptide.

[0021d] Various aspects of the disclosure pertain to a mixture comprising non-pluripotent mammalian cells, a compound that promotes glycolytic metabolism, one or more exogenous transcription factors, a small molecule TGF β receptor/ALK5 inhibitor, and a small molecule MEK inhibitor, wherein: (a) the compound that promotes glycolytic metabolism comprises F2,6P (fructose 2,6-bisphosphate), F6P (fructose 6-phosphate), DNP (2,4-dinitrophenol), NOG (N-oxalylglycine), QC (quercetin), 2-HA (2-hydroxyglutaric acid), NA (nicotinic acid), or a PDK1 activator; (b) the one or more exogenous transcription factors comprises an Oct4 polypeptide; and (c) (i) the one or more exogenous transcription factors comprises a Klf polypeptide, a Sox polypeptide, a Myc polypeptide, or a combination thereof; or (ii) the mixture comprises an HDAC inhibitor.

[0021e] Various embodiments of the claimed invention also pertain to a mixture comprising non-pluripotent mammalian cells, a compound that promotes glycolytic metabolism, one or more exogenous transcription factors, a small molecule TGF β receptor/ALK5 inhibitor, and a small molecule MEK inhibitor, wherein: (a) the compound that promotes glycolytic metabolism comprises F2,6P (fructose 2,6-bisphosphate), F6P (fructose 6-phosphate), DNP (2,4-dinitrophenol), NOG (N-oxalylglycine), QC (quercetin), 2-HA (2-hydroxyglutaric acid), NA (nicotinic acid), or a PDK1 activator; (b) the one or more exogenous transcription factors comprises an Oct4 polypeptide; and (c) (i) the one or more exogenous transcription factors further comprises a Klf polypeptide; or (ii) the mixture comprises an HDAC inhibitor.

[0021f] Various embodiments of the claimed invention also pertain to an *in vitro* method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell, the method comprising: (a) contacting the non-pluripotent cell with an Oct4 polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding an Oct4 polypeptide, wherein no c-Myc polypeptide and no polynucleotide encoding c-Myc are introduced into the non-pluripotent cell; (b) contacting the non-pluripotent cell with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient to induce the cell to become a pluripotent stem cell, thereby inducing the non-pluripotent mammalian cell into an induced pluripotent stem cell; and (c) one or more of (i) contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, (ii) contacting the non-pluripotent cell with a Klf polypeptide, or (iii) introducing into the non-pluripotent cell a polynucleotide encoding a Klf polypeptide.

[0021g] Various aspects of the disclosure also pertain to an induced pluripotent stem cell obtain by a method as claimed herein. Also claimed is a composition comprising such an induced pluripotent stem cell and a carrier and a therapeutic composition comprising such an induced pluripotent stem cell and a pharmaceutically acceptable carrier.

[0021h] Various embodiments of the claimed invention also pertain to a small molecule for use in manufacturing a pluripotent stem cell from a non-pluripotent cell for application in regenerative medicine or tissue regeneration, wherein the small molecule enhances reprogramming of a mammalian cell, wherein the small molecule comprises a PDK1 (3'-phosphoinositide-dependent kinase-1) activator, and wherein the PDK1 activator comprises 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ('12Z'), (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ('13Z'), or a 3,5-diphenylpent-2-enoic acid.

[0021i] Various embodiments of the claimed invention also pertain to a method of making a therapeutic composition comprising an induced pluripotent stem cell (iPSC) for tissue regeneration or regenerative medicine in a subject in need thereof, the method comprising: (a) obtaining an iPSC induced according to a method as defined in any one of claims 1 to 14, 28 to 38 or 49 to 59, and (b) making the therapeutic composition comprising the iPSC for tissue regeneration or regenerative medicine in the subject in need thereof.

[0021j] Various aspects of the disclosure also pertain to a kit comprising a mixture as claimed herein.

DEFINITIONS

[0022] An "Oct polypeptide" refers to any of the naturally-occurring members of Octamer family of transcription factors, or variants thereof that maintain transcription factor activity, 5 *e.g.*, within at least 50%, 80%, or 90% activity compared to the closest related naturally occurring family member, or polypeptides comprising at least the DNA-binding domain of the naturally occurring family member, and can further comprise a transcriptional activation domain. Exemplary Oct polypeptides include, Oct-1, Oct-2, Oct-3/4, Oct-6, Oct-7, Oct-8, Oct-9, and Oct-11. *e.g.* Oct3/4 (referred to herein as "Oct4") contains the POU domain, a 10 150 amino acid sequence conserved among Pit-1, Oct-1, Oct-2, and uric-86. *See*, Ryan, A.K. & Rosenfeld, M.G. *Genes Dev.* 11, 1207–1225 (1997). In some embodiments, variants have at least 85%, 90%, or 95% amino acid sequence identity across their whole sequence compared to a naturally occurring Oct polypeptide family member such as those listed above or such as listed in Genbank accession number NP_002692.2 (human Oct4) or NP_038661.1 15 (mouse Oct4). Oct polypeptides (*e.g.*, Oct3/4) can be from human, mouse, rat, bovine, porcine, or other animals. Generally, the same species of protein will be used with the species of cells being manipulated.

[0023] A "Klf polypeptide" refers to any of the naturally-occurring members of the family of Krüppel-like factors (Klfs), zinc-finger proteins that contain amino acid sequences similar 20 to those of the *Drosophila* embryonic pattern regulator Krüppel, or variants of the naturally-occurring members that maintain transcription factor activity, similar *e.g.*, within at least 50%, 80%, or 90% activity compared to the closest related naturally occurring family member, or polypeptides comprising at least the DNA-binding domain of the naturally occurring family member, and can further comprise a transcriptional activation domain. *See*, 25 Dang, D.T., Pevsner, J. & Yang, V.W. *Cell Biol.* 32, 1103-1121 (2000). Exemplary Klf family members include, Klf1, Klf2, Klf3, Klf-4, Klf5, Klf6, Klf7, Klf8, Klf9, Klf10, Klf11, Klf12, Klf13, Klf14, Klf15, Klf16, and Klf17. Klf2 and Klf-4 were found to be factors capable of generating iPS cells in mice, and related genes Klf1 and Klf5 did as well, although with reduced efficiency. *See*, Nakagawa, *et al.*, *Nature Biotechnology* 26:101-106 (2007). In 30 some embodiments, variants have at least 85%, 90%, or 95% amino acid sequence identity across their whole sequence compared to a naturally occurring Klf polypeptide family member such as those listed above or such as listed in Genbank accession number CAX16088 (mouse Klf4) or CAX14962 (human Klf4). Klf polypeptides (*e.g.*, Klf1, Klf4, and Klf5) can be from human, mouse, rat, bovine, porcine, or other animals. Generally, the

same species of protein will be used with the species of cells being manipulated. To the extent a Klf polypeptide is described herein, it can be replaced with an estrogen-related receptor beta (Essrb) polypeptide. Thus, it is intended that for each Klf polypeptide embodiment described herein, a corresponding embodiment using Essrb in the place of a Klf4
5 polypeptide is equally described.

[0024] A "Myc polypeptide" refers any of the naturally-occurring members of the Myc family (*see, e.g.,* Adhikary, S. & Eilers, M. *Nat. Rev. Mol. Cell Biol.* 6:635–645 (2005)), or variants thereof that maintain transcription factor activity, *e.g.,* within at least 50%, 80%, or 90% activity compared to the closest related naturally occurring family member, or
10 polypeptides comprising at least the DNA-binding domain of the naturally occurring family member, and can further comprise a transcriptional activation domain. Exemplary Myc polypeptides include, *e.g.,* c-Myc, N-Myc and L-Myc. In some embodiments, variants have at least 85%, 90%, or 95% amino acid sequence identity across their whole sequence compared to a naturally occurring Myc polypeptide family member, such as those listed
15 above or such as listed in Genbank accession number CAA25015 (human Myc). Myc polypeptides (*e.g.,* c-Myc) can be from human, mouse, rat, bovine, porcine, or other animals. Generally, the same species of protein will be used with the species of cells being manipulated. To the extent a Myc polypeptide is described herein, it can be replaced with a Wnt polypeptide, *e.g.,* Wnt 3A (*e.g.,* NP_149122.1), or agent that stimulates the Wnt
20 signaling pathway, *e.g.,* a glycogen synthase kinase alpha or beta inhibitor. Thus, it is intended that for each Myc polypeptide embodiment described herein, a corresponding embodiment using a Wnt polypeptide or agent that stimulates the Wnt signaling pathway in the place of a Myc polypeptide is equally described.

[0025] A "Sox polypeptide" refers to any of the naturally-occurring members of the SRY-related HMG-box (Sox) transcription factors, characterized by the presence of the high-mobility group (HMG) domain, or variants thereof that maintain transcription factor activity, *e.g.,* within at least 50%, 80%, or 90% activity compared to the closest related naturally occurring family member or polypeptides comprising at least the DNA-binding domain of the naturally occurring family member, and can further comprise a transcriptional activation
25 domain. *See, e.g.,* Dang, D.T., *et al., Int. J. Biochem. Cell Biol.* 32:1103–1121 (2000). Exemplary Sox polypeptides include, *e.g.,* Sox1, Sox-2, Sox3, Sox4, Sox5, Sox6, Sox7, Sox8, Sox9, Sox10, Sox11, Sox12, Sox13, Sox14, Sox15, Sox17, Sox18, Sox-21, and Sox30. Sox1 has been shown to yield iPS cells with a similar efficiency as Sox2, and genes Sox3, Sox15, and Sox18 have also been shown to generate iPS cells, although with somewhat less
30

efficiency than Sox2. *See, Nakagawa, et al., Nature Biotechnology 26:101 - 106 (2007).* In some embodiments, variants have at least 85%, 90%, or 95% amino acid sequence identity across their whole sequence compared to a naturally occurring Sox polypeptide family member such as those listed above or such as listed in Genbank accession number

5 CAA83435 (human Sox2). Sox polypeptides (*e.g.*, Sox1, Sox2, Sox3, Sox15, or Sox18) can be from human, mouse, rat, bovine, porcine, or other animals. Generally, the same species of protein will be used with the species of cells being manipulated.

[0026] An "exogenous transcription factor," as used herein, refers to a transcription factor that is not naturally (*i.e.*, endogenously) expressed in a cell of interest. Thus, an exogenous
10 transcription factor can be expressed from an introduced expression cassette (*e.g.*, under control of a promoter other than a native transcription factor promoter) or can be introduced as a protein from outside the cell. In some embodiments, the exogenous transcription factor comprises an Oct polypeptide (*e.g.*, Oct4), a Klf polypeptide (*e.g.*, Klf4), a Myc polypeptide (*e.g.*, c-Myc), or a Sox polypeptide (*e.g.*, Sox2).

15 [0027] "H3K9" refers to histone H3 lysine 9. H3K9 modifications associated with gene activity include H3K9 acetylation and H3K9 modifications associated with heterochromatin, include H3K9 di-methylation or tri-methylation. *See, e.g., Kubicek, et al., Mol. Cell 473-481 (2007).* "H3K4" refers to histone H3 lysine 4. *See, e.g., Ruthenburg et al., Mol. Cell 25:15-30 (2007).*

20 [0028] The term "pluripotent" or "pluripotency" refers to cells with the ability to give rise to progeny that can undergo differentiation, under the appropriate conditions, into cell types that collectively demonstrate characteristics associated with cell lineages from all of the three germinal layers (endoderm, mesoderm, and ectoderm). Pluripotent stem cells can contribute to many or all tissues of a prenatal, postnatal or adult animal. A standard art-accepted test,
25 such as the ability to form a teratoma in 8-12 week old SCID mice, can be used to establish the pluripotency of a cell population, however identification of various pluripotent stem cell characteristics can also be used to detect pluripotent cells.

[0029] "Pluripotent stem cell characteristics" refer to characteristics of a cell that distinguish pluripotent stem cells from other cells. The ability to give rise to progeny that can
30 undergo differentiation, under the appropriate conditions, into cell types that collectively demonstrate characteristics associated with cell lineages from all of the three germinal layers (endoderm, mesoderm, and ectoderm) is a pluripotent stem cell characteristic. Expression or non-expression of certain combinations of molecular markers are also pluripotent stem cell

characteristics. For example, human pluripotent stem cells express at least one, two, or three, and optionally all, of the markers from the following non-limiting list: SSEA-3, SSEA-4, TRA-1-60, TRA-1-81, TRA-2-49/6E, ALP, Sox2, E-cadherin, UTF-1, Oct4, Rex1, and Nanog. Cell morphologies associated with pluripotent stem cells are also pluripotent stem cell characteristics.

5 [0030] A "recombinant" polynucleotide is a polynucleotide that is not in its native state, *e.g.*, the polynucleotide comprises a nucleotide sequence not found in nature, or the polynucleotide is in a context other than that in which it is naturally found, *e.g.*, separated from nucleotide sequences with which it typically is in proximity in nature, or adjacent (or
10 contiguous with) nucleotide sequences with which it typically is not in proximity. For example, the sequence at issue can be cloned into a vector, or otherwise recombined with one or more additional nucleic acids.

[0031] "Expression cassette" refers to a polynucleotide comprising a promoter or other regulatory sequence operably linked to a sequence encoding a protein.

15 [0032] The terms "promoter" and "expression control sequence" are used herein to refer to an array of nucleic acid control sequences that direct transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase II type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located
20 as much as several thousand base pairs from the start site of transcription. Promoters include constitutive and inducible promoters. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence
25 (such as a promoter, or array of transcription factor binding sites) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

[0033] A "heterologous sequence" or a "heterologous nucleic acid", as used herein, is one that originates from a source foreign to the particular host cell, or, if from the same source, is
30 modified from its original form. Thus, a heterologous expression cassette in a cell is an expression cassette that is not endogenous to the particular host cell, for example by being linked to nucleotide sequences from an expression vector rather than chromosomal DNA, being linked to a heterologous promoter, being linked to a reporter gene, etc.

- 5 [0034] The terms "nucleic acid" and "polynucleotide" are used interchangeably herein to refer to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs).
- 10 [0035] Unless otherwise indicated, a particular nucleic acid sequence also encompasses conservatively modified variants thereof (*e.g.*, degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or
- 15 deoxyinosine residues (Batzer *et al.*, *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka *et al.*, *J. Biol. Chem.* 260:2605-2608 (1985); Rossolini *et al.*, *Mol. Cell. Probes* 8:91-98 (1994)).
- [0036] "Inhibitors," "activators," and "modulators" of expression or of activity are used to refer to inhibitory, activating, or modulating molecules, respectively, identified using *in vitro* and *in vivo* assays for expression or activity of a described target protein, *e.g.*, ligands,
- 20 agonists, antagonists, and their homologs and mimetics. The term "modulator" includes inhibitors and activators. Inhibitors are agents that, *e.g.*, inhibit expression or bind to, partially or totally block stimulation or protease inhibitor activity, decrease, prevent, delay activation, inactivate, desensitize, or down regulate the activity of the described target protein, *e.g.*, antagonists. Activators are agents that, *e.g.*, induce or activate the expression of
- 25 a described target protein or bind to, stimulate, increase, open, activate, facilitate, enhance activation or protease inhibitor activity, sensitize or up regulate the activity of described target protein (or encoding polynucleotide), *e.g.*, agonists. Modulators include naturally occurring and synthetic ligands, antagonists and agonists (*e.g.*, small chemical molecules, antibodies and the like that function as either agonists or antagonists). Such assays for
- 30 inhibitors and activators include, *e.g.*, applying putative modulator compounds to cells expressing the described target protein and then determining the functional effects on the described target protein activity, as described above. Samples or assays comprising described target protein that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of

effect. Control samples (untreated with modulators) are assigned a relative activity value of 100%. Inhibition of a described target protein is achieved when the activity value relative to the control is about 80%, optionally 50% or 25, 10%, 5% or 1%. Activation of the described target protein is achieved when the activity value relative to the control is 110%, optionally
5 150%, optionally 200, 300%, 400%, 500%, or 1000-3000% or more higher.

[0037] The term "allosteric" is used to refer to an effect that affects the activity of one part of an enzyme (such as an active site) by the binding of a molecule at a different site (regulatory site) at a different location on the enzyme. The binding of non-substrate molecules at allosteric sites effects the binding kinetics of the substrate-binding (active) site.
10 "Allosteric binding sites" are contained in many enzymes and receptors. As a result of binding to allosteric binding sites, the interaction with the normal ligand may be either enhanced or reduced. For example, an allosteric binding site in 3'-phosphoinositide-dependent kinase-1 (PDK1) is the PDK1 interacting fragment (PIF) binding pocket located between helix C, helix B and the -4 and -5 sheets (Pearl *et al.*, *Curr. Opin. Struct. Biol.* 12,
15 761–767 (2002); Biondi *et al.*, *Biochem. J.* 372, 1–13 (2003); Newton *et al.*, *Chem. Rev.* 101, 2353– 2364 (2001)).

[0038] As used herein, "promote" or "increase," or "promoting" or "increasing" are used interchangeably herein. These terms refer to the increase in a measured parameter (*e.g.*, activity, expression, glycolysis, glycolytic metabolism, glucose uptake, biosynthesis
20 downstream of glycolysis) in a treated cell (tissue or subject) in comparison to an untreated cell (tissue or subject). A comparison can also be made of the same cell or tissue or subject between before and after treatment. The increase is sufficient to be detectable. In some embodiments, the increase in the treated cell is at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 1-fold, 2-fold, 3-fold, 4-fold or more in comparison to an untreated
25 cell.

[0039] As used herein, "inhibit," "prevent" or "reduce," or "inhibiting," "preventing" or "reducing" are used interchangeably herein. These terms refer to the decrease in a measured parameter (*e.g.*, activity, expression, mitochondrial respiration, mitochondrial oxidation, oxidative phosphorylation) in a treated cell (tissue or subject) in comparison to an untreated
30 cell (tissue or subject). A comparison can also be made of the same cell or tissue or subject between before and after treatment. The decrease is sufficient to be detectable. In some embodiments, the decrease in the treated cell is at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or completely inhibited in comparison to an untreated cell. In some

embodiments the measured parameter is undetectable (*i.e.*, completely inhibited) in the treated cell in comparison to the untreated cell.

BRIEF DESCRIPTION OF THE DRAWINGS

- 5 [0040] **Figure 1. Generation of human induced pluripotent stem cells from primary keratinocytes by single gene, OCT4, and small molecules.** (a) Treatment with 0.5 μ M PD0325901 (PD) and 0.5 μ M A-83-01 (A83) significantly improved generation of iPSCs from primary human keratinocytes transduced with either 4TFs (4F, OKSM) or 3TFs (3F, OKS). NHEKs were seeded at a density of 100,000 transduced cells per 10 cm dish. (b) Further chemical screens identified PS48, NaB, and their combination that can substantially enhance reprogramming of primary human keratinocytes transduced with 2TFs (OK). NHEKs were seeded at a density of 100,000 transduced cells per 10 cm dish. (c) Experimental scheme for generation of human iPSCs from primary human keratinocytes transduced by a single reprogramming gene, *OCT4*. KCM, keratinocyte culture medium; hESCM, human ESC culture media. (d) Live immunostaining with TRA-1-81 of iPSC colonies that were generated from primary human keratinocytes transduced with 2TFs/OK or 1TF/OCT4 before picking-up of colonies. (e) The established human iPSC-OK and iPSC-O cells express typical pluripotency markers, including ALP (alkaline phosphatase), *OCT4*, *SOX2*, *NANOG*, *SSEA-4* and TRA-1-81. Nuclei were stained with DAPI.
- 10
- 15
- 20 [0041] **Figure 2. In depth characterizations of human iPSC-OK and iPSC-O cells.** (a) Expression analysis by RT-PCR of the endogenous pluripotency genes and exogenous *OCT4* and *KLF4*. *GAPDH* was used as an input control. (b) Methylation analysis of the *OCT4* and *NANOG* promoters by bisulfate genomic sequencing. Open circles and closed circles indicate unmethylated and methylated CpGs in the promoter regions, respectively. (c) Scatter plots comparing global gene expression patterns between iPSC-O cells and NHEKs, and hESCs. The positions of the pluripotency genes *OCT4*, *NANOG*, and *SOX2* are shown by arrows. Black lines indicate the linear equivalent and twofold changes in gene expression levels between the samples. (d) Human iPSC-OK and iPSC-O could effectively differentiate *in vitro* into cells in the three germ layers, including neural ectodermal cells (β III tubulin⁺), mesodermal cells (SMA⁺), and endodermal cells (AFP⁺) using EB method. (e) Quantitative PCR test of three germ layer markers from differentiated human iPSCs using EB method: ectoderm (*PAX6*, β III *TUBULIN*), mesoderm (*FOXF1*, *HAND1*) and endoderm (*AFP*, *GATA6*). Data denotes *GAPDH*- normalized fold changes relative to undifferentiated
- 25
- 30

parental human iPSCs. (f) Human iPSC-OK and iPSC-O could effectively produce full teratoma, which contains differentiated cells in the three germ layers, in SCID mice.

[0042] Figure 3. Generation and characterization of human induced pluripotent stem cells from human umbilical vein endothelial cells by single gene, OCT4, and small molecules.

(a) Experimental scheme for generation of human iPSCs from HUVECs transduced by OCT4. HCM, HUVEC culture medium; hESCM, human ESC culture media. (b) The established hiPSC-O cells from HUVECs express typical pluripotency markers, including NANOG and SSEA-4. Nuclei were stained with DAPI. (c) Expression analysis by RT-PCR of the endogenous pluripotency genes. GAPDH was used as an input control. (d) Methylation analysis of the OCT4 and NANOG promoters by bisulfate genomic sequencing. Open circles and closed circles indicate unmethylated and methylated CpGs in the promoter regions, respectively. (e) hiPSC-O cells from HUVECs could effectively differentiate *in vitro* into cells in the three germ layers, including neural ectodermal cells (β III tubulin⁺), mesodermal cells (SMA⁺), and endodermal cells (AFP⁺) using EB method. (f) hiPSC-O cells could effectively produce full teratoma, which contains differentiated cells in the three germ layers in SCID mice.

[0043] Figure 4. Characterization of human iPSC-O cells from AHEKs. (a) The established hiPSC-O cells from adult keratinocytes express typical pluripotency markers, including NANOG, SOX2 and SSEA-4. Nuclei were stained with DAPI. (b) These hiPSC-O cells could effectively differentiate *in vitro* into cells in the three germ layers, including neural ectodermal cells (β III tubulin⁺), mesodermal cells (SMA⁺), and endodermal cells (AFP⁺) using EB method.

[0044] Figure 5. Characterization of human iPSC-O cells from AFDCs. (a) The established hiPSC-O cells from amniotic fluid derived cells express typical pluripotency markers, including NANOG, SOX2 and SSEA-4. Nuclei were stained with DAPI. (b) These hiPSC-O cells could effectively differentiate *in vitro* into cells in the three germ layers, including neural ectodermal cells (β III tubulin⁺), mesodermal cells (SMA⁺), and endodermal cells (AFP⁺) using EB method.

[0045] Figure 6. Additional hiPSC cell lines express typical pluripotency markers. The other established hiPSC-O cell lines express typical pluripotency markers, including NANOG and SSEA-4. Nuclei were stained with DAPI.

[0046] Figure 7. Feeder-free culture of hiPSC cell lines. hiPSCs were split onto MatrigelTM/ECM-coated plates in chemically defined hESC medium as previously reported.

These hiPSCs could be maintained and expanded in a feeder-free environment. ICC showed the expression of pluripotency markers, OCT4 and SSEA4. Nuclei were stained with DAPI.

[0047] **Figure 8. Genotyping of hiPSCs.** RT-PCR analysis using genomic DNA shows that only OCT4 transgene integrated in the genome of hiPSC-O lines (hiPSC-O#1, hiPSC-O#3, hiPSC-O#21, hiPSC-O#26 and hiPSC-O#31). NHEKs (a) and HUVECs (b) were used as negative controls, while vectors were used as positive controls.

[0048] **Figure 9. Integration of the OCT4 transgene in hiPSCs.** Genomic DNA (10 µg) were digested with EcoRI and hybridized with the *OCT4* cDNA probe (an EcoRI/SpeI fragment of pSin-EF2-OCT4-Pur). Multiple transgenic integrations were detected.

10 [0049] **Figure 10. Karyotyping for hiPSC cell lines.** Metaphase spread of hiPSC-O#1 (a) and hiPSC-O#21 (b) show normal karyotype after passage 15.

[0050] **Figure 11. PS48 enhances reprogramming process by facilitating a metabolic switch toward glycolysis.** (a) PS48 treatment activated PDK1 activity. Western blotting analysis of phosphorylation of Akt (Thr-308) after PS48 (5 µM) or UCN-01(20 nM) treatment. (b) PS48 enhanced reprogramming of NHEKs, while UCN-01 (a PDK1 inhibitor) or 2-Deoxy-D-glucose (10 mM) (2-DG, a glycolysis inhibitor) inhibited reprogramming process. Three factor (Klf, Sox, and Oct)-transduced NHEKs were seeded at a density of 100,000 transduced cells per well, treated with compounds for four weeks, and then TRA 1 81 positive colonics were counted. (c) PS48 treatment facilitated/activated a metabolic switch to glycolysis, while treatment of UCN-01 or 2-DG inhibited glycolysis. NHEKs were treated with either PS48, PS48 and UCN-01, or PS48 and 2-DG for 8d and then lactate production in the medium was measured as a typical index of glycolysis by using the Lactate Assay Kit (BioVision, Mountain View, CA, USA). (d) PS48 treatment up-regulated the expression of several key glycolytic genes, including *GLUT1*, *HK2*, *PFK1* and *LDHA*. (e) Known small molecules that have been widely used to modulate mitochondrial respiration, glycolysis metabolism or HIF activation also showed corresponding consistent effects on reprogramming. Four factor (Klf, Sox, Myc, and Oct)-transduced HUVECs were seeded at a density of 20,000 transduced cells per well, treated with metabolism modulating compounds for three weeks and TRA-1-81 positive colonics were counted. F2,6P, 10 mM Fructose 2,6-bisphosphate; F6P, 10 mM Fructose 6-phosphate; 6-AN, 10 µM 6-aminonicotinamide; OA, 10 µM oxalate; DNP, 1 µM 2,4-dinitrophenol; NOG, 1 µM N-oxaloylglycine; QC, 1 µM Quercetin; 2-HA, 10 µM 2-Hydroxyglutaric acid; NA, 10 µM nicotinic acid; DMSO was used as a control.

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

5 [0051] The present invention is based on the surprising discovery that a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator greatly improves efficiency of induction of pluripotency in non-pluripotent mammalian cells. Accordingly, the present invention provides for methods of inducing pluripotency in non-pluripotent mammalian cells wherein the method comprises contacting the non-pluripotent cells with a PDK1 activator.

10 [0052] The present invention is also based on the surprising discovery that compounds that promote glycolytic metabolism as described herein greatly improve efficiency of induction of pluripotency in non-pluripotent mammalian cells. It was discovered in the present invention that compounds that promote glycolytic metabolism facilitate the metabolic reprogramming from mitochondrial oxidation (mainly used by adult somatic cells) to glycolysis (mainly used by embryonic stem cells (ESCs)), thereby inducing pluripotency in non-pluripotent
15 mammalian cells. Compounds that promote glycolysis or compounds that inhibit or impede mitochondrial respiration/oxidation are therefore useful in inducing pluripotency in non-pluripotent mammalian cells. Further, it was discovered that compounds that promote a process either upstream (*e.g.*, PDK1 pathway, hypoxia-inducible factor pathway, glucose uptake transporter pathway) or downstream of glycolysis (*e.g.*, fatty acids synthesis, lipids
20 synthesis, nucleotides synthesis, and amino acids synthesis) are useful in inducing pluripotency in non-pluripotent mammalian cells. Accordingly, the present invention provides for methods of inducing pluripotency in non-pluripotent mammalian cells wherein the method comprises contacting the non-pluripotent cells with one or more compounds that promote glycolytic metabolism as described herein.

25 [0053] To date, a large number of different methods and protocols have been established for inducing non-pluripotent mammalian cells into induced pluripotent stem cells (iPSCs). It is believed that the agents described herein can be used in combination with essentially any protocol for generating iPSCs and thereby improve the efficiency of the protocol. Thus, the present invention provides for incubation of non-pluripotent cells with at least a PDK1
30 activator, including but not limited to an allosteric PDK1 activator, in combination with any protocol for generating iPSCs. In other embodiments, the present invention provides for incubation of non-pluripotent cells with at least a compound that promotes glycolytic metabolism in combination with any protocol for generating iPSCs.

[0054] As used herein, "efficiency of induction," with respect to induction of a non-pluripotent cell into an induced pluripotent stem cell, refers to the number of non-pluripotent cells that can be converted into iPSCs in a defined time frame, or the amount of time it takes to convert a defined number of non-pluripotent cells into iPSCs, under conditions sufficient for inducing pluripotent stem cells. The improvement in efficiency of an iPSC generation protocol will depend on the protocol and which agents of the invention are used. In some embodiments, the efficiency is improved by at least 10%, 20%, 50%, 75%, 100%, 150%, 200%, 300% or more compared to the same protocol without inclusion of the agents of the invention (*e.g.*, a PDK1 activator, *e.g.*, an allosteric PDK1 activator, or a compound that promotes glycolytic metabolism, *e.g.*, PDK1 activators, glycolysis activators, glycolysis substrates, glycolytic intermediates and their metabolic precursors thereof, glucose uptake transporter activators, mitochondrial respiration modulators such as oxidative phosphorylation inhibitors, and hypoxia-inducible factor activators). In some embodiments, efficiency is measured with regard to improvement of the number of iPSCs generated in a particular time frame (*e.g.*, by comparing the number of iPSCs generated from non-pluripotent cells in a defined time frame under conditions comprising the introduction of one or more agents of the invention to the number of iPSCs generated from non-pluripotent cells in the defined time frame under conditions which do not comprise the introduction of one or more agents of the invention). In some embodiments, efficiency is measured with regard to improvement in the speed by which iPSCs are generated (*e.g.*, by comparing the length of time it takes to generate a defined number of iPSCs from non-pluripotent cells under conditions comprising the introduction of one or more agents of the invention to the length of time it takes to generate a defined number of iPSCs from non-pluripotent cells under conditions which do not comprise the introduction of one or more agents of the invention). In some embodiments, efficiency of induction is measured under conditions comprising transducing non-pluripotent cells (*e.g.*, normal human epidermal keratinocytes) with Oct4 and Klf4, then culturing the transduced cells in the absence or presence of one or more agents of the invention (*e.g.*, a PDK1 activator), as described in the Examples section below. Induction of iPSCs from non-pluripotent cells can be measured according to any method known in the art, including but not limited to marker analysis (*e.g.*, using pluripotency markers Tra-1-81 and/or OCT4).

[0055] According to the methods of the present invention, specific, context-dependent, treatment regimes can improve reprogramming efficiency. The effectiveness of a certain treatment regime can depend, in some embodiments, on cell types, cell passage numbers, and

exogenous transcription factors used. For example, in some embodiments, a more significant improvement in reprogramming efficiency by a specific treatment regime can be observed when reprogramming cells transduced with or in contact with fewer than four exogenous transcription factors, *i.e.*, with one, two, or three exogenous transcription factors, as compared to reprogramming cells transduced with or in contact with four exogenous transcription factors.

[0056] In general, human cells can take considerably longer (*e.g.*, 6-8 weeks) to be reprogrammed than mouse cells (*e.g.*, about 2 weeks). The effects of a specific treatment regime, in some embodiments, can be more exaggerated when reprogramming human cells as compared to mouse cells. Accordingly, when relatively longer periods (*e.g.*, at least 3, 4, 5, 6, or more weeks) are used in reprogramming, a treatment regime, *e.g.*, one that uses an epigenetic modifier, can be used to improve reprogramming efficiency.

[0057] The inventors have found that epigenetic modifiers can improve reprogramming efficiency. As defined herein, the term "epigenetic modifier" refers to a methylation modifying agent (*i.e.*, agents that induce methylation changes to DNA or histones) and/or an acetylation modifying agent (*i.e.*, agents that induce acetylation changes to DNA or histones). In some embodiments, the methylation modifying agent is a DNA methylation inhibitor (*e.g.*, a DNA methyltransferase (DNMT) inhibitor such as RG108), histone methylation inhibitor and/or histone demethylation inhibitor. In some embodiments, the acetylation modifying agent is a histone deacetylase (HDAC) inhibitor (*e.g.*, valproic acid (VPA), sodium butyrate (NaB), trichostatin A (TSA), or suberoylanilide hydroxamic acid (SAHA)), a histone acetyltransferase (HAT) inhibitor, histone deacetylase and histone acetyltransferase. In some embodiments, epigenetic modifiers are agents that inhibit methyltransferases or demethylases or agents that activate methyltransferases or demethylases. In some embodiments, the epigenetic modifier is an agent that inhibits histone H3K9 methylation or promotes H3K9 demethylation, *e.g.*, a G9a histone methyltransferase such as BIX01294.

[0058] Some epigenetic modifiers, however, may also induce cell differentiation. Accordingly, in some embodiments of the invention, epigenetic modifiers are used only in the earlier stage of the treatment, *e.g.*, in the first 1, 2, 3, or 4 weeks, in the first half, the first 1/3, the first quarter, or the first 1/5 of the treatment period. By omitting epigenetic modifiers in the later stage of the treatment, *e.g.*, in the last 1, 2, 3, or 4 weeks, in the last half, the last 1/3, the last quarter, or the last 1/5 of the treatment period, the side effects of inducing cell differentiation by these epigenetic modifiers can be, at least partially, avoided.

[0059] Alternatively, epigenetic modifiers that do not induce differentiation, or only minimally induce differentiation can be used. For example, when HDAC inhibitor is used in a treatment regime, a HDAC inhibitor that does not induce differentiation, or only minimally induces differentiation, *e.g.*, sodium butyrate, is used.


5 [0060] It is further discovered in the present invention that a treatment regime using MEK inhibitors can improve reprogramming efficiency. MEK inhibitors also support cell self-renewal of the induced pluripotent cells. Some MEK inhibitors, however, may inhibit cell proliferation. Accordingly, in some embodiments of the invention, MEK inhibitors are used only in the later stage of the treatment, *e.g.*, in the last 1, 2, 3, or 4 weeks, in the last half, the
10 last 1/3, the last quarter, or the last 1/5 of the treatment period. By omitting MEK inhibitors in the earlier stage of the treatment, *e.g.*, in the first 1, 2, 3, or 4 weeks, in the first half, the first 1/3, the first quarter, or the first 1/5 of the treatment period, cell proliferation is not inhibited in the earlier stage. For example, pluripotency can be induced by contacting a non-pluripotent mammalian cell with a PDK1 activator or with a compound that promotes
15 glycolytic metabolism (*e.g.*, a PDK1 activator) in the absence of a MEK inhibitor in the earlier stage, followed by contacting the non-pluripotent cell with a PDK1 activator or a compound that promotes glycolytic metabolism (*e.g.*, a PDK1 activator) and a MEK inhibitor in the later stage. In some embodiments, the method of inducing pluripotency comprises contacting the non-pluripotent cell with a PDK1 activator or with a compound that promotes
20 glycolytic metabolism (*e.g.*, a PDK1 activator), a TGF β receptor/ALK5 inhibitor, and a histone deacetylase (HDAC) inhibitor in the earlier stage, followed by contacting the non-pluripotent cell with a PDK1 activator or with a compound that promotes glycolytic metabolism (*e.g.*, a PDK1 activator), a TGF β receptor/ALK5 inhibitor, a histone deacetylase (HDAC) inhibitor and a MEK inhibitor in the later stage.

25 II. PDK1 Activators

[0061] 3'-phosphoinositide-dependent kinase-1 or "PDK1" is a master kinase associated with the activation of AKT/PKB and many other AGC kinases including PKC, S6K, SGK. An important role for PDK1 is in the signaling pathways activated by several growth factors and hormones including insulin signaling. The structure of PDK1 can be divided into two
30 domains; the kinase or catalytic domain and the PH domain. The PH domain functions mainly in the interaction of PDK1 with phosphatidylinositol (3,4)-bisphosphate and phosphatidylinositol (3,4,5)-trisphosphate which is important in localization and activation of some membrane associated PDK1 substrates including AKT. The kinase domain has three ligand binding sites; the substrate binding site, the ATP binding site, and the PIF binding

pocket. Several PDK1 substrates including S6K and Protein kinase C require binding at this PIF binding pocket. Small molecule allosteric activators of PDK1 were shown to selectively inhibit activation of substrates that require docking site interaction. These compounds do not bind to the active site and allow PDK1 to activate other substrates that do not require docking site interaction. PDK1 is constitutively active and at present, there are no known inhibitor proteins for PDK1. The activation of PDK1's main effector, AKT, is believed to require a proper orientation of the kinase and PH domains of PDK1 and AKT at the membrane. Phosphoinositide-dependent kinase-1 has been shown to interact with SGK, PRKACA, Sodium-hydrogen exchange regulatory cofactor 2, PRKCD, Protein kinase M ζ (PKMzeta), PKN2, PRKCI, Protein kinase N1, YWHAH and AKT1.

[0062] Exemplary PDK1 activators include sphingosine (King *et al.*, Journal of Biological Chemistry, 275:18108-18113, 2000). Exemplary allosteric activators of PDK1 include PS48 ((Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid), PS08 ((Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid) (Hindie *et al.*, Nature Chemical Biology, 5:758-764, 2009; Huse & Kuriyan, Cell 109: 275-282, 2002; Johnson & Lewis, Chem. Rev. 101:2209-2242, 2001), and compound 1 (2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid) (Engel *et al.*, EMBO J. 25: 5469-5480, 2006); 3,5-diphenylpent-2-enoic acids such as compound 12Z and compound 13Z (12Z: 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid; 13Z: (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid (Stroba *et al.*, J. Med. Chem. 52, 4683-4693 (2009)). PS48 has the following formula:



PS48

[0063] As shown in the Examples, inclusion of a PDK1 activator in cell reprogramming can increase efficiency greatly when used alone and results in even further efficiency increases when used in combination with an HDAC inhibitor. Additional inhibitors, including but not limited to an ALK5 inhibitor and/or a Mek inhibitor, as shown in the

Examples, can also be included in reprogramming, particularly where fewer than the four transcription factors (Oct4, Klf4, Sox2, and c-Myc) are introduced into the cell during reprogramming.

III. Compounds that Promote Glycolytic Metabolism

5 [0064] As defined herein, a metabolism modulating compound refers to a compound that modulates (*e.g.*, promotes or inhibits) metabolism of carbohydrate or other molecules. Metabolism modulating compounds include compounds that promote glycolytic metabolism. As defined herein, a compound that promotes glycolytic metabolism refers to a compound that facilitates cellular metabolic reprogramming from mitochondrial oxidation (mainly used
10 by adult somatic cells) to glycolysis (mainly used by ESCs). In some embodiments, a compound that promotes glycolytic metabolism is a compound that promotes glycolysis or a compound that promotes a process upstream of glycolysis (*e.g.*, PDK1 pathway, hypoxia-inducible factor pathway, glucose uptake transporter pathway). In some embodiments, a compound that promotes glycolytic metabolism is a compound that inhibits or impedes
15 mitochondrial respiration. In some embodiments, a compound that promotes glycolytic metabolism is a compound that promotes a process downstream of glycolysis (*e.g.*, fatty acids synthesis, lipids synthesis, nucleotides synthesis, and amino acids synthesis). Examples of compounds that promote glycolytic metabolism include PDK1 activators, glycolysis activators, glycolysis substrates, glycolytic intermediates and their metabolic precursors
20 thereof, glucose uptake transporter activators, mitochondrial respiration modulators such as oxidative phosphorylation inhibitors, and hypoxia-inducible factor activators. In some embodiments, a compound that promotes glycolytic metabolism is not a simple sugar (*e.g.*, a simple sugar commonly used in cell culture medium). Examples of simple sugars include aldoses such as D-glucose, D-mannose and D-galactose, and ketoses such as D-fructose.

25 I. Glycolysis activators

[0065] Glycolysis activators (*e.g.*, activators of the glycolytic pathway) are known in the art. Enzymes associated with glycolysis pathway are known in the art and include hexokinase, glucokinase, phosphoglucose isomerase, phosphofructokinase, aldolase, triose phosphate isomerase, glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase,
30 phosphoglyceromutase, enolase, pyruvate kinase and lactate dehydrogenase. In some embodiments, the glycolysis activator (*e.g.*, an activator of the glycolysis pathway) is an activator of an enzyme associated with the glycolytic pathway. In some embodiments, a

glycolysis activator is an activator of one of three particular enzymes uniquely associated with the glycolytic pathway: hexokinase, phosphofructokinase, and pyruvate kinase.

[0066] Examples of hexokinase activators include phosphate, citrate, D-malate, 3-phosphoglycerate, catecholamines and catecholamines derivatives. In some embodiments, the hexokinase activator is an allosteric activator. In some embodiments, the hexokinase activators do not include phosphate or citrate.

[0067] Examples of glucokinase activators include GKA1 (6-[(3-isobutoxy-5-isopropoxybenzoyl)amino]nicotinic acid; Brocklehurst *et al.*, Diabetes 53:535–541, 2004), GKA2 (5-({3-isopropoxy-5-[2-(3-thienyl)ethoxy]benzoyl}amino)-1,3,4-thiadiazole-2-carboxylic acid ; Brocklehurst *et al.*, Diabetes 53:535–541, 2004), RO-28-1675 (Grimsby *et al.*, Science 301:370–373, 2003), and compound A (N-Thiazol-2-yl-2-amino-4-fluoro-5-(1-methylimidazol-2-yl)thiobenzamide; Kamata *et al.*, Structure 12, 429–438, 2004), LY2121260 (2-(S)-cyclohexyl-1-(R)-(4-methanesulfonylphenyl)-cyclopropanecarboxylic acid thiazol-2-ylamide; Efanov *et al.*, Endocrinology, 146:3696-3701, 2005). In some embodiments, the glucokinase activator is an allosteric activator. Additional glucokinase activators are disclosed in WO 00/058293, WO 01/44216, WO 01/83465, WO 01/83478, WO 01/85706, WO 01/85707 and WO 02/08209, WO07/075847, WO07/061923, WO07/053345, WO07/104034, WO07/089512, WO08/079787, WO08/111473, WO09/106203, WO09/140624, WO09/140624, WO08/079787, WO02/046173, WO07/006814, WO07/006760, WO06/058923, WO02/048106, WO07/125103, WO07/125105, WO08/012227, WO08/074694, WO08/078674, WO08/084043, WO08/084044, WO09/127544, WO09/127546, WO07/125103, WO07/125105, WO02/014312, WO04/063179, WO07/006761, WO07/031739, WO08/091770, WO08/116107, WO08/118718, WO09/083553, WO04/052869, WO05/123132, WO04/072066, WO07/117381, WO07/115967, WO08/005964, WO08/154563, WO09/022179, WO09/046784, WO08/005964, WO/10/080333 , WO/03/095438, WO/06/016194, WO/05/066145, WO/07/115968, WO/07/143434, WO/08/005914, WO/08/149382, WO/09/018065, WO/09/047798, WO/09/046802, WO/10/029461, WO/08/005914, WO/08/149382, WO/07/143434, WO/10/103438, WO/03/047626, WO/05/095418, WO/08/104994, WO/09/082152, WO/09/082152, WO/05/049019, WO/07/048717, WO/09/042435, and WO/09/042435.

[0068] Examples of phosphofructokinase (or fructose-6-P kinase) activators include fructose 2,6-bisphosphate.

[0069] Examples of pyruvate kinase activators include xylulose 5-P, ceramide, an agonist of the A1 adenosine receptors such as N-6-cyclopentyladenosine. Additional pyruvate kinase activators are disclosed in are disclosed in WO10/042867, WO10/042867, WO99/048490, and WO09/025781.

- 5 [0070] Examples of phosphoglucosomerase activators, aldolase activators, glyceraldehyde 3P dehydrogenase activators, triose phosphate isomerase activators, phosphoglycerate kinase, enolase, phosphoglycerate mutase, and lactate dehydrogenase are known in the art.

2. Glycolysis substrates

- [0071] Examples of glycolysis substrates include glucose 6-phosphate, fructose 6-phosphate, fructose 1,6-bisphosphate, glyceraldehyde 3-phosphate, 1,3-bisphosphoglycerate, 3-phosphoglycerate, 2-phosphoglycerate, and phosphoenolpyruvate. In some embodiments, a compound that promotes glycolytic metabolism is not a simple sugar (*e.g.* glucose).

3. Glycolytic intermediates and the metabolic precursors thereof

- [0072] Glycolytic intermediates are all variously utilized as for biosynthesis of other important molecules such as fatty acids, lipids, nucleotides, and amino acids. Therefore, as defined herein, compounds that promote glycolytic metabolism include compounds that promote a process that is downstream of glycolysis (*e.g.*, fatty acids synthesis, lipids synthesis, nucleotides synthesis, and amino acids synthesis). In some embodiments, compounds that promote glycolytic metabolism include glycolytic intermediates, *e.g.*, glycolytic intermediates that were utilized in these downstream biosynthesis pathways. In some embodiments, compounds that promote glycolytic metabolism include metabolic precursors of glycolytic intermediates. As defined herein, the term "metabolic precursors" refers to compounds from which glycolytic intermediates are metabolically converted, *e.g.*, in a cell, a tissue, or human body.

- 25 [0073] Examples of glycolytic intermediates include glucose 6-phosphate, fructose 6-phosphate, fructose 1,6-bisphosphate, dihydroxyacetone phosphate, glyceraldehyde 3-phosphate, 1,3-bisphosphoglycerate, 3-phosphoglycerate, 2-phosphoglycerate, phosphoenolpyruvate, oxaloacetate, pyruvate, and metabolite precursors thereof. In some embodiments, the glycolytic intermediate is nicotinamide adenine dinucleotide (NADH). In some embodiments, the compound that promotes glycolytic metabolism is a metabolic precursor of NADH. In some embodiments, the compound that promotes glycolytic metabolism is nicotinic acid or nicotinamide.

4. Glucose uptake transporter activators

[0074] As defined herein, the term "glucose uptake transporter activator" refers to compounds that stimulate or otherwise promote the expression or activity of a glucose uptake transporter. As defined herein, the term "glucose transporter" refers to proteins that transport 5 compounds (whether glucose, glucose analogs, other sugars such as fructose or inositol, or non-sugars such as ascorbic acids) across a cell membrane and are members of the glucose transporter "family" based on structural similarity (*e.g.*, homology to other glucose transport proteins). As defined herein, glucose transporters also include transporter proteins that have a primary substrate other than glucose. For example, the glucose transporter GLUT5 is 10 primarily a transporter of fructose, and is reported to transport glucose itself with low affinity. Similarly, the primary substrate for the glucose transporter HMIT is myo-inositol (a sugar alcohol). As used herein, the term "glucose transporter," unless otherwise specified, includes transporters of fructose and inositol. Examples of glucose uptake transporters include a glucose transporter selected from the groups of GLUT1-12, HMIT and SGLT1-6 15 transporters.

[0075] Examples of glucose uptake transporter activators include insulin, pinitol (*see, e.g.*, WO/2000/071111), 8-bromo-cyclic AMP (*see, e.g.*, Ogura *et al.*, Journal of Endocrinology, 164:171-178, 2000), arachidonic acid (Fong *et al.*, Cellular Signalling, 8:179-183, 1996), phorbol esters such as 12-O-tetra-decanoyl-phorbol 13-acetate (*see, e.g.*, Molecular Brain 20 Research, 15:221-226, 1992).

5. Mitochondrial respiration modulators (oxidative phosphorylation inhibitors)

[0076] As defined herein, the term "mitochondrial respiration" or "mitochondrial oxidation" refers to the oxidation of substrate molecules (*e.g.*, sugars, organic acids, pyruvate, 25 etc.) inside mitochondria. In some embodiments, a compound that promotes glycolytic metabolism is a mitochondrial respiration modulator. A compound that can affect the degree of mitochondria respiration/oxidation is generally referred to herein as a "mitochondrial respiration modulator" or other similar term. In some embodiments, mitochondrial respiration modulator useful for the methods of the invention is a compound that inhibits or 30 impedes mitochondrial respiration or mitochondrial oxidation. In some embodiments, mitochondrial respiration modulator useful for the methods of the invention is an oxidative phosphorylation inhibitor.

[0077] An oxidative phosphorylation inhibitor of the invention can be any inhibitor of one or more enzymes of oxidative phosphorylation or an oxidative phosphorylation uncoupler.

The oxidative phosphorylation enzymes are known in the art and include enzyme complex I (NADH coenzyme Q reductase), II (succinate-coenzyme Q reductase), III (coenzyme Q cytochrome C reductase), IV (cytochrome oxydase), and V (F0-F1, ATP synthase).

[0078] Inhibitors of enzyme complex I are any known in the art and can include, but are not limited to any of the following: tritylthioalanine, carminomycin, and piperazinedione, rotenone, amytal, 1-methyl-4-phenylpyridinium (MPP+), paraquat, methylene blue, and ferricyanide (the latter 2 are electron acceptors). Inhibitors of enzyme complex II are any

known in the art. Inhibitors of coenzyme Q are any known in the art. Inhibitors of enzyme complex III are any known in the art and can include, but are not limited to myxothiazol, antimycin A, ubisemiquinone, cytochrome C, 4,6-diaminotriazine derivatives, methothrexate or electron acceptors such as phenazine methosulfate and 2,6-Dichlorophenol-indophenol.

Inhibitors of enzyme complex IV are any known in the art and can include, but are not limited to cyanide, hydrogen sulfide, azide, formate, phosphine, carbon monoxide and electron acceptor ferricyanide. Inhibitors of enzyme complex V are any known in the art and can

include, but are not limited to 2-hydroxyglutaric acid, VM-26 (4'-demethyl-epipodophyllotoxin thenylidene glucoside), tritylthioalanine, carminomycin, piperazinedione, dinitrophenol, dinitrocresol, 2-hydroxy-3-alkyl-1,4-naphthoquinones, apoptolidin aglycone,

oligomycin, ossamycin, cytovaricin, naphthoquinone derivatives (*e.g.*, dichloroallyl-lawsone and lapachol), rhodamine, rhodamine 123, rhodamine 6G, carbonyl cyanide p-trifluoromethoxyphenylhydrazone, rothenone, safranin O, cyhexatin, DDT, chlordecone, arsenate, pentachlorophenol, benzonitrile, thiadiazole herbicides, salicylate, cationic amphiphilic

drugs (amiodarone, perhexiline), gramicidin, calcimycin, pentachlorobutadienyl-cysteine (PCBD-cys), and trifluorocarbonyl cyanide phenylhydrazone (FCCP). Other inhibitors of oxidative phosphorylation may include atractyloside, DDT, free fatty acids,

lysophospholipids, n-ethylmaleimide, mersanyl, and p-benzoquinone.

[0079] Oxidative phosphorylation uncouplers refer to compounds that act as uncouplers of oxidative phosphorylation from electron transport. Examples of oxidative-phosphorylation

uncouplers include, but are not limited to, DNP, 5-chloro-3-tert-butyl-2'-chloro-4'-nitrosalicylanilide (S-13), sodium 2,3,4,5,6-pentachlorophenolate (PCP), 4,5,6,7-tetrachloro-2-(trifluoromethyl)-1H-benzimidazole (TTFB), Flufenamic acid (2-[3-

(trifluoromethyl)anilino]benzoic acid), 3,5-di-tert-butyl-4-hydroxy-benzylidenemalononitrile (SF6847), carbonyl cyanide m-chloro phenyl hydrazone (CCCP), Carbonyl cyanide p-

[trifluoromethoxy]-phenyl-hydrazone (FCCP), and alpha-(phenylhydrazono)phenylacetonitrile derivatives. phenylacetonitrile derivatives; and weak acids comprising: Weakly Acidic Phenols, benzimidazoles, N-phenylanthranilates, salicylanilides, phenylhydrazones, salicylic acids, acyldi-thiocarbazates, coumarins, and aromatic amines.

6. Hypoxia-inducible factor activators

[0080] Activators of hypoxia-inducible factor pathway are known in the art and include alkaloids and other amino acid derivatives, inhibitors of HIF asparaginyl hydroxylase (factor-inhibiting HIF, or FIH) and HIF prolyl hydroxylases (HPH or PHD), inhibitors of glycogen synthase kinase 3 β (GSK3 β), nitric oxide (NO) donors, microtubule-depolymerizing agents (MDA), phenolic compounds, terpenes/steroids, and prostaglandin E2 (PGE2). Examples of alkaloids and other amino acid derivatives include deferoxamine and desferri-exochelin DFE 722 SM, Ciclopirox olamine [Loprox[®], 6-cyclohexyl-1-hydroxy-4-methyl-2(1H)-pyridone 2-aminoethanol], and 8-methyl-pyridoxatin. Examples of inhibitors of glycogen synthase kinase 3 β (GSK3 β) include indirubin, derivatives of indirubin such as 5-iodoindirubin-3'-oxime and 5-methylindirubin-3'-oxime. Examples of nitric oxide (NO) donors include S-nitroso-N-acetyl-D,L-penicillamine (SNAP), 3-(hydroxy-1-(1-methylethyl)-2-nitrosohydrazino)-1-propanamine (NOC5), diazen-1-ium-1,2-diolate (NOC-18), S-nitrosoglutathione (GSNO), spermine NONOate (a complex of NO with the natural product spermine), diethylamine NONOate, and diethyltryamine NONOate. Examples of microtubule-depolymerizing agents (MDA) include plant alkaloids vinblastine, colchicine, and synthetic MDAs such as nocodazole. Examples of phenolic compounds include dibenzoylmethane (DBM), the flavonoid quercetin (3,3',4',5,7-pentahydroxyflavone), (-)-epicatechin-3-gallate (ECG), and (-)-epigallocatechin-3-gallate (EGCG). Examples of terpenes and steroids include sesquiterpene-tropolones (*e.g.*, pycnidione, epolone A and epolone B), 4-hydroxy estradiol (4-OHE2), dihydrotestosterone, methyltrienolone (R1881), and diterpene ester phorbol 12-O-myristate 13-acetate (PMA, also known as 12-O-tetradecanoylphorbol 13-acetate or TPA).

[0081] Examples of inhibitors of HIF asparaginyl hydroxylase (factor-inhibiting HIF, or FIH) and HIF prolyl hydroxylases (HPH or PHD) include analogues of 2-oxoglutarate (2OG) such as N-oxalylglycine (5, NOG), ester derivatives of NOG (*e.g.*, DMOG (dimethyl-oxalylglycine)), N-((3-hydroxy-6-chloroquinolin-2-yl)carbonyl)-glycine, 3-hydroxypyridine-2-carbonyl-glycine, 3,4-dihydroxybenzoate, pyridine-2,5-dicarboxylate, pyridine-2,4-dicarboxylate, N-oxalyl-2S-alanine, additional analogues of 2OG as described in Mole *et al.*,

Bioorg Med Chem Lett. 13:2677–80, 2003, alahopcin and dealanylalahopcin, dealanylalahopcin analogues 3-carboxymethylene N-hydroxy succinimide, 3-carboxy-N-hydroxy pyrrolidone, l-mimosine (L-Mim), ethyl 3,4-dihydroxybenzoate (3,4-DHB), and 6-chlor-3-hydroxyquinolin-2-carbonic acid-N-carboxymethylamid (S956711). Additional HIF
5 asparaginyl hydroxylase and HIF prolyl hydroxylases inhibitors are described in, e.g., Ivan *et al.*, *Proc Natl Acad Sci USA* 99: 13459-13464, 2002, WO03/049686, WO03/080566, and WO06/084210, WO10/056767.

[0082] Other activators of HIF pathway include iron chelators (*e.g.*, deferoxamine, 2,2'-pyridyl, 1,10-phenanthroline, Ca²⁺ chelator BAPTA (1,2-bis(2-aminophenoxy)ethane-N,N,N',N'-tetraacetic acid), transition metals (*e.g.*, cobalt, nickel, chromium (VI), and copper), the organomercurial compound mersalyl, and FG-0041 (a compound that is structurally related to 1,10-phenanthroline).

[0083] Additional activators of HIF pathway include proteins that up-regulate HIF-1 translation. Protein kinase C (PKC) increases the rate of HIF-1 α transcription and functions
15 in conjunction with the phosphatidylinositol 3-kinase (PI3K) pathway, which also enhances HIF-1 α translation. The PKC pathway activates expression of the S6 ribosomal protein, which specifically recognizes mRNA transcripts such as HIF-1 α . Via phosphorylation of the S6 protein in normoxic conditions, the rates of HIF-1 α mRNA translation can be greatly increased, effectively countering the effects of the proteasome degradation of this subunit and
20 increasing levels of the HIF-1 complex within the cell. The PI3K pathway has been identified as the primary means by which various mediators, such as lipopolysaccharides, affect activation of HIF-1 α in vascular smooth muscle cells and macrophages (Dery *et al.*, *Int J Biochem Cell Bio.* 37:535–540, 2004; Page *et al.*, *J Biol Chem.* 277:48403–48409, 2002).

[0084] The macrophage-derived peptide PR39 has been shown to stabilize HIF-1 α by
25 decreasing its degradation, resulting in accelerated formation of vascular structures in vitro and increased myocardial vasculature in mice (Li *et al.*, *Nat Med* 6: 49-55, 2000). Direct induction of HIF-1 has been achieved by using the N- or C-terminal of ODDD polypeptides that block VHL-mediated degradation (Maranchie *et al.*, *Cancer Cell* 1: 247-255, 2002).

[0085] HIF pathway activators further include other non-hypoxic physiological stimuli
30 such as growth factors, cytokines, and hormones. Examples of growth factors that activate HIF pathway include insulin-like growth factor (IGF)-1 and IGF-2, IGF-binding protein (IGFBP)-2 and IGFBP-3, EGF, basic fibroblast growth factor (bFGF), and heregulin. Examples of cytokines that activate HIF pathway include tumor necrosis factor alpha

(TNF α), interleukin-1 beta (IL-1 β), and IL-1. Examples of hormones that activate HIF pathway include the vascular hormones angiotensin II and thrombin, thyroid hormone and follicle-stimulating hormone. Other physiological factors such as the redox protein thioredoxin-1 (Trx-1) and oxidized low-density lipoprotein (oxLDL) can also induce HIF-1 α protein and activate HIF-1.

7. **PDK1 activators**

[0086] In some embodiments, compounds that promote glycolytic metabolism are PDK1 activators. Exemplary PDK1 activators are described herein in section II, supra.

IV. **HDAC Inhibitors**

[0087] Exemplary HDAC inhibitors can include antibodies that bind, dominant negative variants of, and siRNA and antisense nucleic acids that target HDAC. HDAC inhibitors include, but are not limited to, TSA (trichostatin A) (see, e.g., Adcock, *British Journal of Pharmacology* 150:829-831 (2007)), VPA (valproic acid) (see, e.g., Munster et al., *Journal of Clinical Oncology* 25:18S (2007): 1065), sodium butyrate (NaBu) (see, e.g., Han et al., *Immunology Letters* 108:143-150 (2007)), SAHA (suberoylanilide hydroxamic acid or vorinostat) (see, e.g., Kelly et al., *Nature Clinical Practice Oncology* 2:150-157 (2005)), sodium phenylbutyrate (see, e.g., Gore et al., *Cancer Research* 66:6361-6369 (2006)), depsipeptide (FR901228, FK228) (see, e.g., Zhu et al., *Current Medicinal Chemistry* 3(3):187-199 (2003)), trapoxin (TPX) (see, e.g., Furumai et al., *PNAS* 98(1):87-92 (2001)), cyclic hydroxamic acid-containing peptide 1 (CHAPI) (see, Furumai supra), MS-275 (see, e.g., Carninci et al., WO2008/126932), LBH589 (see, e.g., Goh et al., WO2008/108741) and PXD101 (see, Goh, supra). In general at the global level, pluripotent cells have more histone acetylation, and differentiated cells have less histone acetylation. Histone acetylation is also involved in histone and DNA methylation regulation. In some embodiments, HDAC inhibitors facilitate activation of silenced pluripotency genes.

V. **ALK5 Inhibitors**

[0088] TGF β receptor (e.g., ALK5) inhibitors can include antibodies to, dominant negative variants of, and antisense nucleic acids that suppress expression of, TGF β receptors (e.g., ALK5). Exemplary TGF β receptor/ALK5 inhibitors include, but are not limited to, SB431542 (see, e.g., Inman, et al., *Molecular Pharmacology* 62(1):65-74 (2002)), A-83-01, also known as 3-(6-Methyl-2-pyridinyl)-N-phenyl-4-(4-quinolinyl)-1H-pyrazole-1-carbothioamide (see, e.g., Tojo et al., *Cancer Science* 96(11):791-800 (2005), and commercially available from, e.g., Toicris Bioscience);

2-(3-(6-Methylpyridin-2-yl)-1H-pyrazol-4-yl)-1,5-naphthyridine, Wnt3a/BIO (see, e.g., Dalton et al., WO2008/094597, herein incorporated by reference), BMP4 (see, Dalton, supra), GW788388 (-{4-[3-(pyridin-2-yl)-1H-pyrazol-4-yl]pyridin-2-yl}-N-(tetrahydro-2H-pyran-4-yl)benzamide) (see, e.g., Gellibert et al., *Journal of Medicinal Chemistry* 49(7):2210-2221 (2006)), SM16 (see, e.g., Suzuki et al., *Cancer Research* 67(5):2351-2359 (2007)), IN-1130 (3-((5-(6-methylpyridin-2-yl)-4-(quinoxalin-6-yl)-1H-imidazol-2-yl)methyl)benzamide) (see, e.g., Kim et al., *Xenobiotica* 38(3):325-339 (2008)), GW6604 (2-phenyl-4-(3-pyridin-2-yl-1H-pyrazol-4-yl)pyridine) (see, e.g., de Gouville et al., *Drug News Perspective* 19(2):85-90 (2006)), SB-505124 (2-(5-benzo[1,3]dioxol-5-yl-2-tert-butyl-3H-imidazol-4-yl)-6-methylpyridine hydrochloride) (see, e.g., DaCosta et al., *Molecular Pharmacology* 65(3):744-752 (2004)) and pyrimidine derivatives (see, e.g., those listed in Stiefl et al., WO2008/006583), SU5416; 2-(5-benzo[1,3]dioxol-5-yl-2-tert-butyl-3H-imidazol-4-yl)-6-methylpyridine hydrochloride (SB-505124); lerdelimumb (CAT-152); metelimumab (CAT-192); GC-1008; ID11; AP-12009; AP-11014; LY550410; LY580276; LY364947; LY2109761; SB-505124; SB-431542; SD-208; SM16; NPC-30345; Ki26894; SB-203580; SD-093; Gleevec; 3,5,7,2',4'-pentahydroxyflavone (Morin); activin-M108A; P144; and soluble TBR2-Fc (see, e.g., Wrzesinski et al., *Clinical Cancer Research* 13(18):5262-5270 (2007); Kaminska et al., *Acta Biochimica Polonica* 52(2):329-337 (2005); and Chang et al., *Frontiers in Bioscience* 12:4393-4401 (2007)). Further, while "an ALK5 inhibitor" is not intended to encompass non-specific kinase inhibitors, an "ALK5 inhibitor" should be understood to encompass inhibitors that inhibit ALK4 and/or ALK7 in addition to ALK5, such as, for example, SB-431542 (see, e.g., Inman et al., *J. Mol. Pharmacol.* 62(1): 65-74 (2002)). Without intending to limit the scope of the invention, it is believed that ALK5 inhibitors affect the mesenchymal to epithelial conversion/transition (MET) process. TGF β /activin pathway is a driver for epithelial to mesenchymal transition (EMT). Therefore, inhibiting the TGF β /activin pathway can facilitate the MET (i.e., reprogramming) process.

[0089] In view of the data herein showing the effect of inhibiting ALK5, it is believed that inhibition of the TGF β /activin pathway will have similar effects. Thus, any inhibitor (e.g., upstream or downstream) of the TGF β /activin pathway can be used in combination with, or instead of, ALK5 inhibitors as described in each paragraph herein. Exemplary TGF β /activin pathway inhibitors include but are not limited to: TGF β receptor inhibitors, inhibitors of SMAD2/3 phosphorylation, inhibitors of the interaction of SMAD2/3 and SMAD4, and activators/agonists of SMAD6 and SMAD7. Furthermore, the categorizations described below are merely for

organizational purposes and one of skill in the art would know that compounds can affect one or more points within a pathway, and thus compounds may function in more than one of the defined categories.

[0090] Inhibitors of SMAD2/3 phosphorylation can include antibodies to, dominant negative variants of, and antisense nucleic acids that target SMAD2 or SMAD3. Specific examples of inhibitors include PD169316; SB203580; SB-431542; LY364947; A77-01; and 3,5,7,2',4'-pentahydroxyflavone (Morin). See, e.g., Wrzesinski, *supra*; Kaminska, *supra*; Shimanuki, et al., *Oncogene* 26:3311-3320 (2007); and Kataoka et al., EP1992360.

[0091] Inhibitors of the interaction of SMAD2/3 and SMAD4 can include antibodies to, dominant negative variants of, and antisense nucleic acids that target SMAD2, SMAD3 and/or SMAD4. Specific examples of inhibitors of the interaction of SMAD2/3 and SMAD4 include, but are not limited to, Trx-SARA, Trx-xFoxH1b and Trx-Lef1. (See, e.g., Cui et al., *Oncogene* 24:3864-3874 (2005) and Zhao et al., *Molecular Biology of the Cell*, 17:3819-3831 (2006).)

[0092] Activators/agonists of SMAD6 and SMAD7 include, but are not limited to, antibodies to, dominant negative variants of, and antisense nucleic acids that target SMAD 6 or SMAD 7. Specific examples of inhibitors include, but are not limited to, smad7-as PTO-oligonucleotides. See, e.g., Miyazono et al., US6534476, and Steinbrecher et al., US2005119203.

VI. MEK Inhibitors

[0093] Inhibitors of MEK can include antibodies to, dominant negative variants of, and siRNA and antisense nucleic acids that suppress expression of, MEK. Specific examples of MEK inhibitors include, but are not limited to, PD0325901, (see, e.g., Rinehart, et al., *Journal of Clinical Oncology* 22: 4456-4462 (2004)), PD98059 (available, e.g., from Cell Signaling Technology), U0126 (available, for example, from Cell Signaling Technology), SL 327 (available, e.g., from Sigma-Aldrich), ARRY-162 (available, e.g., from Array Biopharma), PD184161 (see, e.g., Klein et al., *Neoplasia* 8:1-8 (2006)), PD184352 (CI-1040) (see, e.g., Mattingly et al., *The Journal of Pharmacology and Experimental Therapeutics* 316:456-465 (2006)), sunitinib (see, e.g., Voss et al., US2008004287), sorafenib (see, Voss *supra*), Vandetanib (see, Voss *supra*), pazopanib (see, e.g., Voss *supra*), Axitinib (see, Voss *supra*) and PTK787 (see, Voss *supra*).

[0094] Currently, several MEK inhibitors are undergoing clinical trial evaluations. CI-1040 has been evaluated in Phase I and II clinical trials for cancer (*see, e.g., Rinehart et al., Journal of Clinical Oncology* 22(22):4456-4462 (2004)). Other MEK inhibitors being evaluated in clinical trials include PD184352 (*see, e.g., English et al., Trends in*
5 *Pharmaceutical Sciences* 23(1):40-45 (2002)), BAY 43-9006 (*see, e.g., Chow et al., Cytometry (Communications in Clinical Cytometry)* 46:72-78 (2001)), PD-325901 (also PD0325901), GSK1120212, ARRY-438162, RDEA119, AZD6244 (also ARRY-142886 or ARRY-886), RO5126766, XL518 and AZD8330 (also ARRY-704). *See, e.g.,* information from the National Institutes of Health located on the World Wide Web at clinicaltrials.gov as
10 well as information from the National Cancer Institute located on the World Wide Web at cancer.gov/clinicaltrials.

VII. Reprogramming

[0095] To date, a large number of different methods and protocols have been established for inducing non-pluripotent mammalian cells into induced pluripotent stem cells (iPSCs).
15 iPSCs are similar to ESCs in morphology, proliferation, and pluripotency, judged by teratoma formation and chimaera contribution. It is believed that PDK1 activators or compounds that promote glycolytic metabolism (*e.g.,* PDK1 activators), optionally in combination with an HDAC inhibitor, and optionally an ALK5 inhibitor and optionally a Mek inhibitor, will improve essentially any reprogramming protocol for generating iPSCs. Reprogramming
20 protocols that can be improved are believed to include those involving introduction of one or more reprogramming transcription factors selected from an Oct polypeptide (including but not limited to Oct 3/4), a Sox polypeptide (including but not limited to Sox2), a Klf polypeptide (including but not limited to Klf4) and/or a Myc polypeptide (including but not limited to c-Myc). Thus, in some embodiments, conditions sufficient to induce a cell to
25 become a pluripotent stem cell comprise conditions in which one or more reprogramming transcription factors selected from an Oct polypeptide (including but not limited to Oct 3/4), a Sox polypeptide (including but not limited to Sox2), a Klf polypeptide (including but not limited to Klf4) and/or a Myc polypeptide (including but not limited to c-Myc) are introduced into the cell. As noted in the Examples, PDK1 activators have been shown to improve
30 reprogramming with as few as one reprogramming transcription factor (*e.g.,* Oct4 alone). Thus, in some embodiments, conditions sufficient to induce a cell to become a pluripotent stem cell comprise conditions in which one reprogramming transcription factor (*e.g.,* Oct4 alone) is introduced into the cell.

[0096] In some embodiments, conditions sufficient to induce a cell to become a pluripotent stem cell comprise introducing reprogramming factors into the cells, for example, by expression from a recombinant expression cassette that has been introduced into the target cell, or by incubating the cells in the presence of exogenous reprogramming transcription factor polypeptides such that the polypeptides enter the cell.

[0097] Studies have shown that retroviral transduction of mouse fibroblasts with four transcription factors that are highly expressed in ESCs (Oct-3/4, Sox2, KLF4 and c-Myc) generate induced pluripotent stem (iPS) cells. See, Takahashi, K. & Yamanaka, S. *Cell* 126, 663-676 (2006); Okita, K., Ichisaka, T. & Yamanaka, S. *Nature* 448, 313-317 (2007); Wernig, M. et al. *Nature* 448, 318-324 (2007); Maherali, N. et al. *Cell Stem Cell* 1, 55-70 (2007); Meissner, A., Wernig, M. & Jaenisch, R. *Nature Biotechnol.* 25, 1177-1181 (2007); Takahashi, K. et al. *Cell* 131, 861-872 (2007); Yu, J. et al. *Science* 318, 1917-1920 (2007); Nakagawa, M. et al. *Nature Biotechnol.* 26, 101-106 (2007); Wernig, M., Meissner, A., Cassady, J. P. & Jaenisch, R. *Cell Stem Cell.* 2, 10-12 (2008). Studies have also demonstrated reprogramming of human somatic cells with transcription factors that are highly expressed in ESCs: Hockemeyer et al. *Cell Stem Cell.* 11;3(3):346-53 (2008); Lowry et al. *Proc Natl Acad Sci U S A.* 105(8):2883-8 (2008); Park et al. *Nature.* 10;451(7175):141-6 (2008); Nakagawa et al. *Nat Biotechnol.* Jan;26(1):101-6 (2008); Takahashi et al. *Cell.* 131(5):861-72 (2007); and Yu et al. *Science.* 318(5858):1917-20 (2007). Such methods are believed to be improved with the inclusion of a PDK1 activator or one of more compounds that promote glycolytic metabolism (e.g., a PDK1 activator) and optionally other agents as described herein.

[0098] To address the safety issues that arise from target cell genomes harboring integrated exogenous sequences, a number of modified genetic protocols have been further developed and can be used according to the present invention. These protocols produce iPS cells with potentially reduced risks, and include non-integrating adenoviruses to deliver reprogramming genes (Stadtfield, M., et al. (2008) *Science* 322, 945-949), transient transfection of reprogramming plasmids (Okita, K., et al. (2008) *Science* 322, 949-953), piggyBac transposition systems (Woltjen, K., et al. (2009). *Nature* 458, 766-770, Yusa et al. (2009) *Nat. Methods* 6:363-369, Kaji, K., et al. (2009) *Nature* 458, 771-775), Cre-excisable viruses (Soldner, F., et al. (2009) *Cell* 136, 964-977), and oriP/EBNA1-based episomal expression system (Yu, J., et al. (2009) *Science* DOI: 10.1126). Thus, in some embodiments, conditions sufficient to induce a cell to become a pluripotent stem cell comprise conditions in which

reprogramming factors are delivered by non-integrating adenoviruses, transient transfection of reprogramming plasmids, piggyBac transposition systems, re-excisable viruses (Soldner, F., *et al.* (2009) *Cell* 136, 964-977), and/or oriP/EBNA1-based episomal expression systems, according to any of the protocols described above. In some embodiments, a PDK1 activator
5 or one of more compounds that promote glycolytic metabolism (*e.g.*, a PDK1 activator) and optionally other agents as described herein are incubated with cells in any of the protocols described above.

[0099] As noted above, reprogramming can involve culturing target cells in the presence of one or more proteins under conditions to allow for introduction of the proteins into the cell.
10 *See, e.g.*, Zhou H *et al.*, *Cell Stem Cell*. 2009 May 8;4(5):381-4; WO/2009/117439. One can introduce an exogenous polypeptide (*i.e.*, a protein provided from outside the cell and/or that is not produced by the cell) into the cell by a number of different methods that do not involve introduction of a polynucleotide encoding the polypeptide. In some embodiments, conditions sufficient to induce a cell to become a pluripotent stem cell comprise introducing into the cell
15 one or more exogenous proteins, each exogenous protein comprising a transcription factor polypeptide of interest linked (*e.g.*, linked as a fusion protein or otherwise covalently or non-covalently linked) to a polypeptide that enhances the ability of the transcription factor to enter the cell (and in some embodiments the cell nucleus).

[0100] Examples of polypeptide sequences that enhance transport across membranes
20 include, but are not limited to, the *Drosophila* homeoprotein *antennapedia* transcription protein (AntHD) (Joliot *et al.*, *New Biol.* 3: 1121-34,1991; Joliot *et al.*, *Proc. Natl. Acad. Sci. USA*, 88: 1864-8,1991; Le Roux *et al.*, *Proc. Natl. Acad. Sci. USA*, 90: 9120-4, 1993), the herpes simplex virus structural protein VP22 (Elliott and O'Hare, *Cell* 88: 223-33, 1997); the HIV-1 transcriptional activator TAT protein (Green and Loewenstein, *Cell* 55: 1179-1188,
25 1988; Frankel and Pabo, *Cell* 55: 1 289-1193, 1988); Kaposi FGF signal sequence (kFGF); protein transduction domain-4 (PTD4); Penetratin, M918, Transportan-10; a nuclear localization sequence, a PEP-I peptide; an amphipathic peptide (*e.g.*, an MPG peptide); delivery enhancing transporters such as described in US Patent No. 6,730,293 (including but not limited to an peptide sequence comprising at least 5-25 or more contiguous arginines or
30 5-25 or more arginines in a contiguous set of 30, 40, or 50 amino acids; including but not limited to an peptide having sufficient, *e.g.*, at least 5, guanidino or amidino moieties); and commercially available Penetratin™ 1 peptide, and the Diatos Peptide Vectors ("DPVs") of the Vectocell® platform available from Daitos S.A. of Paris, France. *See also*, WO/2005/084158 and WO/2007/123667 and additional transporters described therein. Not

only can these proteins pass through the plasma membrane but the attachment of other proteins, such as the transcription factors described herein, is sufficient to stimulate the cellular uptake of these complexes. A number of polypeptides capable of mediating introduction of associated molecules into a cell have been described previously and can be adapted to the present invention. *See, e.g.,* Langel (2002) Cell Penetrating Peptides CRC Press, Pharmacology and Toxicology Series.

[0101] Exemplary polypeptide sequences that enhance transport across membranes include: **VP22**: G S P P T A P T R S K T P A Q G L A R K L H F S T A P P N P D A P W T P R V A G F N K R V F R F S P Q T A R R A T T T R I; **kFGF**: A G S G G A A V A L L P A V L L A L L A P G G E F A; **PTD4**: A G S G G Y A R A A A R Q A R A G G E F A; **PENETRATIN**: R Q I K I W F Q G R R M K W K K; **TAT**: Y G R K K R R Q R R R; **M918**: M V T V L F R R L R I R R A C G P P R V R V; **TRANSPORTAN-10**: A G Y L L G K I G L K A L A A L A K K I L.

[0102] In some embodiments, the polypeptide that enhances transport across membranes is a peptide sequence comprising at least 5 or more contiguous or non-contiguous arginines (*e.g.*, a 8-arginine peptide). In some embodiments, the polypeptide that enhances transport across membranes is a peptide sequence comprising at least 7 or more contiguous or non-contiguous arginines. For example, the polypeptide that enhances transport across membranes is a peptide sequence comprising 11 contiguous arginines, *e.g.*, ESGGGGSPGRRRRRRRRRRR. As noted above, the arginines in the transport enhancing sequence need not all be contiguous. In some embodiments, the polyarginine (*e.g.*, the contiguous or non-contiguous) region is at least 5, 8, 10, 12, 15, 20, or more amino acids long and has at least *e.g.*, 40%, 50%, 60%, 70%, 80%, 90%, or more arginines.

[0103] In some embodiments, conditions sufficient to induce a cell to become a pluripotent stem cell comprise conditions in which one or more exogenous polypeptides, *e.g.*, an Oct polypeptide (including but not limited to Oct 3/4), a Sox polypeptide (including but not limited to Sox2), a Klf polypeptide (including but not limited to Klf4) and/or a Myc polypeptide (including but not limited to c-Myc), is introduced into cells by traditional methods such as lipofection, electroporation, calcium phosphate precipitation, particle bombardment and/or microinjection, or can be introduced into cells by a protein delivery agent. For example, the exogenous polypeptide can be introduced into cells by covalently or noncovalently attached lipids, *e.g.*, by a covalently attached myristoyl group. Lipids used for lipofection are optionally excluded from cellular delivery modules in some embodiments. In some embodiments, the transcription factor polypeptides described herein are exogenously

introduced as part of a liposome, or lipid cocktail (such as commercially available Fugene6 and Lipofectamine). In another alternative, the transcription factor proteins can be microinjected or otherwise directly introduced into the target cell. In some embodiments, the transcription factor polypeptides are delivered into cells using Profect protein delivery reagents, *e.g.*, Profect-P1 and Profect-P2 (Targeting Systems, El Cajon, CA), or using Pro-
5 Ject® transfection reagents (Pierce, Rockford IL, USA). In some embodiments, the transcription factor polypeptides are delivered into cells using a single-wall nano tube (SWNT).

[0104] As discussed in the Examples of WO/2009/117439, incubation of cells with the transcription factor polypeptides of the invention for extended periods can be toxic to the
10 cells. Therefore, in some embodiments of the invention, conditions sufficient to induce the non-pluripotent mammalian cell to become a pluripotent stem cell comprise incubating a PDK1 activator or one of more compounds that promote glycolytic metabolism (*e.g.*, a PDK1 activator) and optionally an HDAC inhibitor, an ALK5 inhibitor and/or a Mck inhibitor, and
15 intermittently incubating the non-pluripotent mammalian cell with one or more of an Oct polypeptide (including but not limited to Oct 3/4), a Sox polypeptide (including but not limited to Sox2), a Klf polypeptide (including but not limited to Klf4) and/or a Myc polypeptide (including but not limited to c-Myc) with intervening periods of incubation of the cell in the absence of the one or more polypeptides. In some embodiments, the cycle of
20 incubation with and without the polypeptides can be repeated for 2, 3, 4, 5, 6, or more times and is performed for sufficient lengths of time (*i.e.*, the incubations with and without proteins) to achieve the development of pluripotent cells.

[0105] The various agents (*e.g.*, PDK1 activator or compounds that promote glycolytic metabolism, HDAC inhibitor, TGFβ receptor/ALK5 inhibitor, MEK/ERK pathway inhibitor, and/or Rho GTPase/ROCK inhibitor, etc.) can be contacted to non-pluripotent cells either
25 prior to, simultaneously with, or after delivery of, programming transcription factors (for example, delivered via expression cassette or as proteins). For convenience, the day the reprogramming factors are delivered is designated "day 1." In some embodiments, the inhibitors are contacted to cells in aggregate (*i.e.*, as a "cocktail") at about days 3-7 and
30 continued for 7-14 days. Alternatively, in some embodiments, the cocktail is contacted to the cells at day 0 (*i.e.*, a day before the preprogramming factors) and incubated for about 14-30 days.

[0106] The cell into which a protein of interest is introduced can be a mammalian cell. The cells can be human or non-human (*e.g.*, primate, rat, mouse, rabbit, bovine, dog, cat, pig, etc.). The cell can be, *e.g.*, in culture or in a tissue, fluid, etc. and/or from or in an organism. Cells that can be induced to pluripotency include, but are not limited to, keratinocyte cells, hair follicle cells, HUVEC (Human Umbilical Vein Endothelial Cells), cord blood cells, neural progenitor cells and fibroblasts.

[0107] In some embodiments, small molecules can improve the efficiency of a process for generating pluripotent cells (*e.g.*, iPS cells). For example, improved efficiency can be manifested by speeding the time to generate such pluripotent cells (*e.g.*, by shortening the time to development of pluripotent cells by at least a day compared to a similar or same process without the small molecule). Alternatively, or in combination, a small molecule can increase the number of pluripotent cells generated by a particular process (*e.g.*, increasing the number in a given time period by at least 10%, 30%, 50%, 100%, 200%, 500%, etc. compared to a similar or same process without the small molecule).

[0108] Optionally, or in addition, small molecules can "complement" or replace what is generally otherwise understood as a necessary expression of one of these proteins to result in pluripotent cells. By contacting a cell with an agent that functionally replaces one of the transcription factors, it is possible to generate pluripotent cells with all of the above-listed transcription factors except for the transcription factor replaced or complemented by the agent.

VIII. Transformation

[0109] This invention employs routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include Sambrook *et al.*, *Molecular Cloning, A Laboratory Manual* (3rd ed. 2001); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel *et al.*, eds., 1994)). In some embodiments, expression cassettes for expression of one or more reprogramming transcription factor is introduced into a cell.

[0110] In some embodiments, the species of cell and protein to be expressed is the same. For example, if a mouse cell is used, a mouse ortholog is introduced into the cell. If a human cell is used, a human ortholog is introduced into the cell.

[0111] It will be appreciated that where two or more proteins are to be expressed in a cell, one or multiple expression cassettes can be used. For example, where one expression cassette expresses multiple polypeptides, a polycistronic expression cassette can be used.

[0112] Any type of vector can be used to introduce an expression cassette of the invention into a cell. Exemplary vectors include but are not limited to plasmids and viral vectors. Exemplary viral vectors include, e.g., adenoviral vectors, AAV vectors, and retroviral (e.g., lentiviral) vectors.

[0113] Suitable methods for nucleic acid delivery for transformation of a cell, a tissue or an organism for use with the current invention are believed to include virtually any method by which a nucleic acid (e.g., DNA) can be introduced into a cell, a tissue or an organism, as described herein or as would be known to one of ordinary skill in the art (e.g., Stadtfeld and Hochedlinger, *Nature Methods* 6(5):329-330 (2009); Yusa et al., *Nat. Methods* 6:363-369 (2009); Woltjen, et al., *Nature* 458, 766-770 (9 April 2009)). Such methods include, but are not limited to, direct delivery of DNA such as by *ex vivo* transfection (Wilson et al., *Science*, 244:1344-1346, 1989, Nabel and Baltimore, *Nature* 326:711-713, 1987), optionally with Fugene6 (Roche) or Lipofectamine (Invitrogen), by injection (U.S. Patent Nos. 5,994,624, 5,981,274, 5,945,100, 5,780,448, 5,736,524, 5,702,932, 5,656,610, 5,589,466 and 5,580,859), including microinjection (Harland and Weintraub, *J. Cell Biol.*, 101:1094-1099, 1985; U.S. Pat. No. 5,789,215); by electroporation (U.S. Pat. No. 5,384,253; Tur-Kaspa et al., *Mol. Cell Biol.*, 6:716-718, 1986; Potter et al., *Proc. Nat'l Acad. Sci. USA*, 81:7161-7165, 1984); by calcium phosphate precipitation (Graham and Van Der Eb, *Virology*, 52:456-467, 1973; Chen and Okayama, *Mol. Cell Biol.*, 7(8):2745-2752, 1987; Rippe et al., *Mol. Cell Biol.*, 10:689-695, 1990); by using DEAE-dextran followed by polyethylene glycol (Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985); by direct sonic loading (Fechheimer et al., *Proc. Nat'l Acad. Sci. USA*, 84:8463-8467, 1987); by liposome mediated transfection (Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982; Fraley et al., *Proc. Nat'l Acad. Sci. USA*, 76:3348-3352, 1979; Nicolau et al., *Methods Enzymol.*, 149:157-176, 1987; Wong et al., *Gene*, 10:87-94, 1980; Kaneda et al., *Science*, 243:375-378, 1989; Kato et al., *J Biol. Chem.*, 266:3361-3364, 1991) and receptor-mediated transfection (Wu and Wu, *Biochemistry*, 27:887-892, 1988; Wu and Wu, *J. Biol. Chem.*, 262:4429-4432, 1987); and any combination of such methods.

IX. Mixtures

[0114] As discussed herein, the present invention provides for mammalian cells in a mixture with a PDK1 activator or a compound that promotes glycolytic metabolism, and one or more of (a) a TGF β receptor/ALK5 inhibitor; (b) a MEK inhibitor; (c) a histone deacetylase (HDAC) inhibitor; or (d) an exogenous polypeptide selected from an Oct

polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some
embodiments, the compound that promotes glycolytic metabolism is a PDK1 activator. In
some embodiments, the PDK1 activator is an allosteric PDK1 activator, *e.g.*, PS48. In some
embodiments, the compound that promotes glycolytic metabolism is a glycolysis activator,
5 *e.g.*, fructose 2,6-bisphosphate. In some embodiments, the compound that promotes
glycolytic metabolism is a substrate for glycolysis, *e.g.*, fructose 6-phosphate. In some
embodiments, the compound that promotes glycolytic metabolism is a glycolytic intermediate
or its metabolic precursors, *e.g.*, nicotinic acid, NADH, or fructose 6-phosphate. In some
embodiments, the compound that promotes glycolytic metabolism is a glucose uptake
10 transporter activator. In some embodiments, the compound that promotes glycolytic
metabolism is a mitochondrial respiration modulator. In some embodiments, the
mitochondrial respiration modulator is an oxidative phosphorylation inhibitor, *e.g.*, 2,4-
dinitrophenol, or 2-hydroxyglutaric acid. In some embodiments, the compound that promotes
glycolytic metabolism is a hypoxia-inducible factor activator, *e.g.*, N-oxaloylglycine, or
15 quercetin.

[0115] In some embodiments, the mixture further comprises a TGF β receptor/ALK5
inhibitor. TGF β receptor/ALK5 inhibitors include but are not limited to A-83-01. In some
embodiments, the mixture further comprises a MEK inhibitor. MEK inhibitors include but
are not limited to PD0325901. In some embodiments, the mixture further comprises a
20 histone deacetylase (HDAC) inhibitor. HDAC inhibitors include but are not limited to
sodium butyrate (NaB) and valproic acid (VPA). In some embodiments, the mixture further
comprises an exogenous transcription factor, *e.g.*, an exogenous transcription factor selected
from an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In
some embodiments, the exogenous transcription factor comprises an amino acid sequence
25 that enhances transport across cell membranes.

[0116] In some embodiments, the compound (*e.g.*, the PDK1 activator or the compound
that promotes glycolytic metabolism) is present in the mixture at a concentration sufficient to
induce or improve efficiency of induction to pluripotency. For example, in some
embodiments, the compounds are in a concentration of at least 0.1 nM, *e.g.*, at least 1, 10,
30 100, 1000, 10000, or 100000 nM, *e.g.*, between 0.1 nM and 100000 nM, *e.g.*, between 1 nM
and 10000 nM, *e.g.*, between 10 nM and 10000 nM. In some embodiments, the mixtures are
in a synthetic vessel (*e.g.*, a test tube, Petri dish, etc.). Thus, in some embodiments, the cells
are isolated cells (not part of an animal). In some embodiments, the cells are isolated from an
animal (human or non-human), placed into a vessel, contacted with one or more compound as

described herein. The cells can be subsequently cultured and optionally, inserted back into the same or a different animal, optionally after the cells have been stimulated to become a particular cell type or lineage. In some embodiments, the concentration of the inhibitors is sufficient to improve by at least 10%, 20%, 50%, 75%, 100%, 150%, 200%, 300% or more, the efficiency of induction of non-pluripotent cells in the mixture into induced pluripotent stem cells when the mixture is submitted to conditions sufficient to induce conversion of the cells into induced pluripotent stem cells.

[0117] As explained herein, in some embodiments, the cells comprise an expression cassette for heterologous expression of at least one or more of an Oct polypeptide, a Myc polypeptide, a Sox polypeptide and a Klf polypeptide. In some embodiments, the cells do not include an expression cassette to express one or more (including in some embodiments, any) of the Oct, Myc, Sox, or Klf polypeptides.

[0118] The cells according to the present invention can be human or non-human (*e.g.*, primate, rat, mouse, rabbit, bovine, dog, cat, pig, etc.). Examples of non-pluripotent cells include those described herein, including but not limited to, cells from a tissue selected from bone marrow, skin, skeletal muscle, fat tissue and peripheral blood. Exemplary cell types include, but are not limited to, fibroblasts, hepatocytes, myoblasts, neurons, osteoblasts, osteoclasts, T-cells, keratinocyte cells, hair follicle cells, human umbilical vein endothelial cells (HUVEC), cord blood cells, and neural progenitor cells. In some embodiments, at least 99% of the cells in the mixture are initially non-pluripotent cells. In some embodiments, essentially all of the cells in the mixture are initially non-pluripotent cells.

[0119] In some embodiments, at least 0.001%, at least 0.002%, at least 0.005%, at least 0.01%, at least 0.05%, at least 0.1%, at least 0.5%, least 1%, at least 2%, at least 3%, at least 4%, at least 5%, at least 6%, at least 7%, at least 8%, at least 9%, at least 10%, at least 15%, at least 20%, at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, or at least 90% of the cells in the mixture are induced into pluripotent cells. In some embodiments, at least 99% of the cells in the mixture are induced into pluripotent cells. In some embodiments, essentially all of the cells are induced into non-pluripotent cells.

X. Kits

[0120] The present invention provides a kit for inducing pluripotency in a non-pluripotent mammalian cell comprising a PDK1 activator or a compound that promotes glycolytic metabolism, and one or more of (a) a TGF β receptor/ALK5 inhibitor; (b) a MEK inhibitor; (c) a histone deacetylase (HDAC) inhibitor; or (d) an exogenous polypeptide selected from an

Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide.. In some embodiments, the compound that promotes glycolytic metabolism is a PDK1 activator. In some embodiments, the PDK1 activator is an allosteric PDK1 activator, *e.g.*, PS48. In some embodiments, the compound that promotes glycolytic metabolism is a glycolysis activator, *e.g.*, fructose 2,6-bisphosphate. In some embodiments, the compound that promotes glycolytic metabolism is a substrate for glycolysis, *e.g.*, fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glycolytic intermediate or its metabolic precursors, *e.g.*, nicotinic acid, NADH, or fructose 6-phosphate. In some embodiments, the compound that promotes glycolytic metabolism is a glucose uptake transporter activator. In some embodiments, the compound that promotes glycolytic metabolism is a mitochondrial respiration modulator. In some embodiments, the mitochondrial respiration modulator is an oxidative phosphorylation inhibitor, *e.g.*, 2,4-dinitrophenol, or 2-hydroxyglutaric acid. In some embodiments, the compound that promotes glycolytic metabolism is a hypoxia-inducible factor activator, *e.g.*, N-oxaloylglycine, or quercetin.

[0121] In some embodiments, the kit further comprises a TGF β receptor/ALK5 inhibitor, *e.g.*, A-83-01. In some embodiments, the kit further comprises a MEK inhibitor, *e.g.*, PD0325901. In some embodiments, the kit further comprises a histone deacetylase (HDAC) inhibitor, *e.g.*, sodium butyrate (NaB), or valproic acid (VPA). In some embodiments, the kit further comprises an exogenous transcription factor, *e.g.*, an exogenous transcription factor selected from an Oct polypeptide, a Klf polypeptide, a Myc polypeptide, and a Sox polypeptide. In some embodiments, the exogenous transcription factor comprises an amino acid sequence that enhances transport across cell membranes.

[0122] In some embodiments, the kits further comprise non-pluripotent cells. Examples of non-pluripotent cells include those described herein, including but not limited to, cells from a tissue selected from bone marrow, skin, skeletal muscle, fat tissue and peripheral blood. Exemplary cell types include, but are not limited to, fibroblasts, hepatocytes, myoblasts, neurons, osteoblasts, osteoclasts, T-cells, keratinocyte cells, hair follicle cells, human umbilical vein endothelial cells (HUVEC), cord blood cells, and neural progenitor cells.

30

EXAMPLES

[0123] The following examples are offered to illustrate, but not to limit the claimed invention.

Example 1: Reprogramming of human primary somatic cells by OCT4 and chemical compounds

[0124] Here we report a novel small molecule cocktail that enables reprogramming of human primary somatic cells to iPSCs with exogenous expression of only OCT4.

5 *Results*

[0125] Among several readily available primary human somatic cell types, keratinocytes that can be easily isolated from human skin or hair follicle represent an attractive cell source for reprogramming, because they endogenously express *KLF4* and *cMYC*, and were reported to be reprogrammed more efficiently using the conventional four TFs or three TFs (without
10 MYC) (Aasen, T. *et al.*, *Nat Biotechnol* 26:1276-1284 (2008); Maherali, N. *et al.*, *Cell Stem Cell* 3, 340-345(2008)). More recently, we reported that dual inhibition of TGF β and MAPK/ERK pathways using small molecules (*i.e.*, SB431542 and PD0325901, respectively) provides a drastically enhanced condition for reprogramming of human fibroblasts with four
15 exogenous TFs (*i.e.*, Oct4, Sox2, Klf4, and c-Myc TFs, or "OSKM") (Lin, T. *et al.*, *Nat Methods* 6:805-808 (2009)). Furthermore, we have shown that such dual pathway inhibition could also enhance reprogramming of human keratinocytes by two exogenous TFs (*i.e.*, Oct4 and Klf4, or "OK") with two small molecules, Parnate (an inhibitor of lysine-specific demethylase 1) and CHIR99021 (a GSK3 inhibitor) (Li, W. *et al.*, *Stem Cells* 27:2992-3000
20 (2009)). However, such a 2-TFs reprogramming process was very inefficient and complex (*e.g.*, involving two exogenous TFs and four chemicals), and reprogramming with even one less TF appeared daunting. Toward the OCT4 only reprogramming, we developed a step-wise strategy in refining reprogramming condition and identifying new reprogramming chemical entities.

[0126] We first attempted to further optimize the reprogramming process under the four or
25 three TFs (*i.e.*, OSKM or OSK) condition in neonatal human epidermal keratinocytes (NHEKs) by testing various inhibitors of TGF β and MAPK pathways at different concentrations using previously reported human iPSC characterization methods (Lin, T. *et al.*, *Nat Methods* 6:805-808 (2009)). We found that the combination of 0.5 μ M PD0325901 and 0.5 μ M A-83-01 (a more potent and selective TGF β receptor inhibitor) was more
30 effective in enhancing reprogramming of human keratinocytes transduced with OSKM or OSK (Figure 1a). Remarkably, when we further reduced viral transductions to only two factors/OK, we could still generate iPSCs from NHEKs when they were treated with 0.5 μ M PD0325901 and 0.5 μ M A-83-01, although with low efficiency. Then we began screening

additional small molecules from a collection of known bioactive compounds at various concentrations as previously reported. Among dozens of compounds tested so far, surprisingly we found that a small molecule activator of PDK1 (3'-phosphoinositide-dependent kinase-1), PS48 (5 μ M) that has never been reported in reprogramming, can significantly enhance the reprogramming efficiency about fifteen fold. Interestingly, we also found that 0.25 mM sodium butyrate (NaB, a histone deacetylase inhibitor) turned out to be much more reliable and efficient than the previously reported 0.5 mM VPA for the generation of iPSCs under OK condition (Figure 1b). Subsequent follow-up studies demonstrated that the combination of 5 μ M PS48 and 0.25 mM NaB could further enhance the reprogramming efficiency over twenty-five fold (Figure 1b and Table 3).

[0127] With such unprecedented efficiency in reprogramming NHEKs under only two TFs, we further explored the possibility of generating iPSCs with OCT4 alone by refining combinations of those small molecules during different treatment windows. Primary NHEKs were transduced with OCT4 and treated with chemicals (Figure 1c). Among various conditions, small iPSC colonies resembling hESCs (four to six colonies out of 1,000,000 seeded cells) appeared in OCT4 infected NHEKs that were treated with 0.25 mM NaB, 5 μ M PS48 and 0.5 μ M A-83-01 during the first four weeks, followed by treatment with 0.25 mM NaB, 5 μ M PS48, 0.5 μ M A-83-01 and 0.5 μ M PD0325901 for another four weeks (Figure 1c). Such TRA-1-81 positive iPSC colonies (Figure 1d) grew larger under conventional hESC culture media and could be serially passaged to yield stable iPSC clones that were further characterized (Figure 1e and 2). In addition, OCT4 only iPSCs could also be generated from human adult keratinocytes by addition of 2 μ M Pamate and 3 μ M CHIR99021 (which had been shown to improve reprogramming of NHEKs under OK condition) to this chemical cocktail. After the reliable reprogramming of primary keratinocytes to iPSCs by OCT4 and small molecules, we further applied the conditions to other human primary cell types, including HUVECs (differentiated mesoderm cells) and AFDCs (amniotic fluid derived cells). Similarly, TRA-1-81 positive iPSC colonies appeared in OCT4 infected HUVECs and AFDCs that were treated with chemicals. Remarkably, it appeared that reprogramming of HUVECs and AFDCs was more efficient and faster than reprogramming of NHEKs under the OCT4 and small molecule conditions (Table 3). Two clones of iPSCs from each cell type were long-term expanded for over 20 passages under conventional hESC culture condition and further characterized (Table 4).

[0128] These stably expanded hiPSC-OK and hiPSC-O cells are morphologically indistinguishable from hESCs, and could be cultured on ECM-coated surface under feeder-

free and chemically defined conditions (Figure 1e and Figure 6). They stained positive for alkaline phosphatase (ALP) and expressed typical pluripotency markers, including OCT4, SOX2, NANOG, TRA-1-81 and SSEA4, detected by immunocytochemistry/ICC (Figure 1e, 3b, Figures 4-5). In addition, RT-PCR analysis confirmed the expression of the endogenous human *OCT4*, *SOX2*, *NANOG*, *REX1*, *UTF1*, *TDGF2*, *FGF4* genes, and silencing of exogenous *OCT4* and *KLF4* (Figure 2a and 3c). Furthermore, bisulfite sequencing analysis revealed that the *OCT4* and *NANOG* promoters of hiPSC-OK and hiPSC-O cells are largely demethylated (Figure 2b and 3d). This result provides further evidence for reactivation of the pluripotency transcription program in the hiPSC-OK and hiPSC-O cells. Global gene expression analysis of hiPSC-O cells, NHEKs and hESCs showed that hiPSC-O cells are distinct from NHEKs (Pearson correlation value: 0.87) and most similar to hESCs (Pearson correlation value: 0.98) (Figure 2c). Genotyping analysis showed that hiPSC-O cells only contained the *OCT4* transgene without the contamination of transgene *KLF4* or *SOX2* (Figure 8). Southern blot analysis showed that there were multiple different integration sites of the *OCT4* transgene (Figure 9) among different clones. In addition, karyotyping result demonstrated that hiPSC-O maintained normal karyotype during the whole reprogramming and expansion process (Figure 10). Furthermore, DNA fingerprinting test excluded the possibility that these hiPSCs arose from hESC contamination in the laboratory (Table 5).

[0129] To examine the developmental potential of these hiPSC-O cells, they were differentiated *in vitro* by the standard embryoid body (EB) differentiation method. ICC analyses demonstrated that they could effectively differentiate into β III-tubulin⁺ characteristic neuronal cells (ectoderm), SMA⁺ mesodermal cells, and AFP⁺ endodermal cells (Figure 2d and 3e). Quantitative PCR analyses further confirmed the expression of these and additional lineage specific marker genes, including ectodermal cells (*β III-tubulin* and *NESTIN*), mesodermal cells (*MSX1* and *MLC2a*), and endodermal cells (*FOXA2* and *AFP*) (Figure 2e). Following EB protocol, these hiPSC-OK and hiPSC-O cells could also give rise to rhythmically beating cardiomyocytes. To test their *in vivo* pluripotency, they were transplanted into *SCID* mice. Four to six weeks later, these hiPSC-O cells effectively generated typical teratomas containing derivatives of all three germ layers (Figure 2f and 3f). Collectively, these *in vitro* and *in vivo* characterizations demonstrated that a single transcription factor, OCT4, combined with a defined small molecule cocktail is sufficient to reprogram several human primary somatic cells to iPSCs that are morphologically, molecularly and functionally similar to pluripotent hESCs.

Discussion

[0130] The studies presented above have a number of important implications: First, although fetal NSCs were shown to be reprogrammed to iPSCs by ectopic expression of OCT4 alone, there has been significant skepticism about whether exogenous OCT4 gene alone would be sufficient to reprogram other more practical human somatic cells that do not endogenously express SOX2 (one of the two master pluripotency genes in reprogramming), are at later developmental stages (*e.g.*, early embryonic/fetal vs. born/adult), and can be obtained without significant harms to the individual. To our knowledge, our study is the first demonstration that iPSCs can be practically derived from readily available primary human somatic cells (*e.g.*, keratinocytes) transduced with a single exogenous reprogramming gene, OCT4. In contrast to neural stem cells from the brain, keratinocytes are more accessible and can be easily obtained from born individuals with less invasive procedures. This further strengthens the strategy of exploiting various practically accessible human somatic cells for iPSC generation with safer approaches and/or better qualities. Thus, this new method and its further development would significantly facilitate production of patient-specific pluripotent stem cells for various applications.

[0131] Second, although small molecules and their combinations have been identified to replace only one or two reprogramming TFs, it becomes exponentially challenging to generate iPSCs when more exogenous reprogramming TFs are omitted together. The identification of this new small molecule cocktail, which functionally replaces three master transcription factors all together (*i.e.*, SOX2, KLF4 and MYC) in enabling generation of iPSCs with OCT4 alone, represents another major step toward the ultimate reprogramming with only small molecules, and further proved and solidified the chemical approach to iPSCs.

[0132] Third, this demonstrated single gene condition also has a significant implication for protein-induced pluripotent stem cell (piPSC) technology. A practical challenge for piPSC technology is large-scale and reliable production of the four transducible reprogramming proteins, each of which behaves differently in manufacture (*e.g.*, their expression, folding, stability etc.). Clearly, combining this small molecule cocktail with a single transducible protein would significantly simplify the piPSC technology and facilitate its applications.

[0133] Fourth, we identified a new small molecule, PS48, with a new target/mechanism in enhancing reprogramming. PS48 is an allosteric small molecule activator of PDK1 (Hindie, V. *et al.*, *Nat Chem Biol* 4:758-764 (2009)). One mechanism by which PS48 enhances reprogramming appears to be facilitating the metabolic reprogramming from mitochondrial

oxidation mainly used by adult somatic cells to glycolysis mainly used by ESCs (which is also known as the Warburg effect) (Manning, B. D. and Cantley, *Cell* 129:1261-1274 (2007); Kondoh, H. *et al.*, *Antioxid Redox Signal* 9:293-299 (2007); Heiden, M. G. V. *et al.*, *Science* 324:1029-1033 (2009)). Such differential use of glycolytic metabolism over mitochondrial

5 respiration by pluripotent stem cells would favor pluripotency by promoting proliferation/cell cycle transition with less oxidative stress. For highly proliferating cells, oxidative phosphorylation would not be able to meet the demand of providing macromolecular precursors for cell replication, but also generates significant amount of reactive oxygen species in mitochondria that could induce excessive oxidative damages. On the other hand,

10 glycolytic metabolism could more effectively generate macromolecular precursors, such as glycolytic intermediates for nonessential amino acids and acetyl-CoA for fatty acids, while provide sufficient energies to meet the needs of proliferating cells (Kondoh, H. *et al.*, *Antioxid Redox Signal* 9:293-299 (2007); Heiden, M. G. V. *et al.*, *Science* 324:1029-1033 (2009)). Interestingly, hypoxic condition and its effector HIF-1 α activation not only have

15 been closely linked to promoting glycolytic metabolism, but also were shown to enhance both mouse and human reprogramming (Yoshida, Y. *et al.*, *Cell Stem Cell* 5:237-241 (2009)). Mechanistically, growth factor signaling pathways, hypoxic condition/HIF-1 α and reprogramming factor Myc appear to regulate complementary aspects of cellular metabolism, including up-regulating glucose transporters and metabolic enzymes of glycolysis, such as

20 GLUT1, HK2 and PFK1 (Gordan, J. D. *et al.*, *Cancer Cell* 12:108-113 (2007); DeBerardinis, R. J. *et al.*, *Cell Metabolism* 7:11-20 (2008)). Those studies suggest that one potential conserved mechanism of Myc, hypoxic condition/HIF-1 α , and growth factors/ Akt pathway activation in enhancing reprogramming converge on their essential roles in regulating glycolytic metabolism. Supporting this notion, we found that treatment with PS48 activated

25 down-stream Akt/PKB (Figure 11a), and up-regulated expression of several key glycolytic genes (Figure 11d), facilitating the metabolic switch to glycolysis (Figure 11c). Conversely, we found that inactivation of PDK1 activity by UCN-01 (a PDK1 inhibitor) or inhibition of glycolysis by 2-Deoxy-D-glucose (a glycolysis inhibitor) not only attenuated glycolysis (Figure 11c) but also blocked reprogramming process (Figure 11b). Furthermore, several

30 known small molecules that have been widely used to modulate mitochondrial respiration (2,4-dinitrophenol), glycolytic metabolism (Fructose 2,6-bisphosphate and oxalate), or more specifically HIF pathway activation (N-oxaloylglycine and Quercetin) also showed corresponding consistent effects on reprogramming: *i.e.*, compounds facilitating glycolytic metabolism enhance reprogramming (such as 2,4-dinitrophenol and N-oxaloylglycine), while

35 compounds blocking glycolytic metabolism inhibit reprogramming (such as oxalate) (Figure

11e) (Hewitson, K. S. and Schofield, C. J., *Drug Discov Today* 9:704-711 (2004); Pelicano, H. *et al.*, *Oncogene* 25:4633-4646 (2006)). In conclusion, these results indicated that a metabolic switch to anaerobic glycolysis is critical for and facilitate reprogramming of somatic cells to pluripotent stem cells.

5 [0134] Finally, this new and powerful small molecule cocktail for reprogramming validated the step-wise chemical optimization and screening strategy presented here as a productive approach toward the ultimate purely chemical-induced pluripotent stem cells. Moreover, we found that different small molecules modulating the same target/mechanism could have significantly different effects on reprogramming in a different context, exemplified by A-83-10 01's and NaB's better reprogramming enhancing activities in human keratinocytes, suggests the importance of "individualized" optimization and treatment with different regimens for specific reprogramming context.

Methods

Cell Culture

15 [0135] Normal Human Epidermal Keratinocytes (Lonza) were maintained in Keratinocyte culturing medium (KCM, Lonza). Human Umbilical Vein Endothelial Cells (HUVECs, Millipore) were maintained in EndoGRO-VEGF Complete Medium (HCM, CHEMICON). Human ESCs and hiPSCs were cultured on MEF feeder cells in conventional human ESC culture media (hESCM: DMEM/F12, 15% Knockout serum replacement, 1% Glutamax, 1% 20 Non-essential amino acids, 1% penicillin/streptomycin, 0.1 mM β -mercaptoethanol and 10 ng/ml bFGF). All cell culture products were from Invitrogen/Gibco BRL except where mentioned.

Lentivirus Production

[0136] The lentivirus supernatants were produced and harvested as previously described 25 (Yu, J. *et al.*, *Science* 318:1917-1920 (2007)). The plasmids used for lentivirus production include pSin-EF2-Puro-hOCT4, pSin2-EF2-Puro-hSOX2, pLove-mKlf4, pLove-mMyc, the packaging plasmid psPAX2 and the envelop-coding plasmid pMD2.G (Yu, J. *et al.*, *Science* 318:1917-1920 (2007) and Li, W. *et al.*, *Stem Cells* 27:2992-3000 (2009)).

Reprogramming of NHEKs

30 [0137] NHEKs were cultured in a 100 mm tissue culture dish and transduced 3 times (3-4 hours each transduction) with freshly produced lentivirus supernatants. 1,000,000 transduced NHEKs were seeded on the irradiated x-ray inactivated CF1 MEF feeder cells in a 100-mm

dish and cultured in KCM and treated with 5 μ M PS48, 0.25 mM NaB (Stemgent) and 0.5 μ M A-83-01 (Stemgent) for 2 weeks, followed by changing half volume of media to hESCM and supplementing with 5 μ M PS48, 0.25 mM NaB and 0.5 μ M A-83-01 for another 2 weeks. Then cell culture media were changed to hESCM and supplemented with 5 μ M PS48, 0.25 mM NaB, 0.5 μ M A-83-01 and 0.5 μ M PD0325901 (Stemgent) for an additional four weeks. The same OCT4 infected keratinocytes cultured in media without chemicals were used as a control. The culture was split by Accutase (Millipore) and treated with 1 μ M Thiazovivin (Stemgent) in the first day after splitting. The iPSC colonies stained positive by Alexa Fluor 555 Mouse anti-Human TRA-1-81 antibody (BD Pharmingen) were picked up for expansion on feeder cells in hESCM and cultured routinely.

Reprogramming of HUVECs

[0138] HUVECs were cultured in a 100 mm tissue culture dish and transduced 2 times (4-6 hours each transduction) with freshly produced lentivirus supernatants. 200,000 transduced HUVECs were seeded on gelatin coated 100-mm dish, cultured in HCM, and treated with 5 μ M PS48, 0.25 mM NaB and 0.5 μ M A-83-01 for 2 weeks, followed by changing half volume of media to hESCM and supplementing with 5 μ M PS48, 0.25 mM NaB and 0.5 μ M A-83-01 for another 2 weeks. Then cell culture media were changed to hESCM and supplemented with 5 μ M PS48, 0.25 mM NaB, 0.5 μ M A-83-01 and 0.5 μ M PD0325901 for additional 1-2 weeks. The iPSC colonies stained positive by Alexa Fluor 555 Mouse anti-Human TRA-1-81 antibody were picked up for expansion on feeder cells in hESCM and cultured routinely. The culture was split by Accutase and treated with 1 μ M Thiazovivin in the first day after splitting.

Reprogramming of HUVECs using various metabolism modulating compounds

[0139] HUVECs were cultured in a 100-mm tissue culture dish and transduced 2 times (4-6 hours each transduction) with freshly produced lentivirus supernatants containing four reprogramming factors (Klf, Sox, Myc, and Oct). About 20,000 transduced HUVECs were seeded on gelatin coated 6-well plate, cultured in HCM, and treated with a metabolism modulating compound for 2 weeks. Then cell culture media were changed to hESCM and supplemented with a metabolism modulating compound for additional 1-2 weeks. The number of iPSC colonies stained positive by Alexa Fluor 555 Mouse anti-Human TRA-1-81 antibody was counted. Various metabolism modulating compounds have been tested, including 10 mM Fructose 2,6-bisphosphate (F2,6P), 10 mM Fructose 6-phosphate (F6P), 10 μ M 6-aminonicotinamide (6-AN), 10 μ M oxalate (OA), 1 μ M 2,4-dinitrophenol (DNP), 1 μ M

N-oxaloylglycine (NOG), 1 μ M Quercetin (QC), 10 μ M 2-Hydroxyglutaric acid (2-HA), or 10 μ M nicotinic acid (NA).

***In vitro* differentiation**

[0140] The *in vitro* differentiation of hiPSCs was carried out by the standard embryoid
5 body (EB) method. Briefly, the hiPSCs were dissociated by Accutase (Millipore), cultured in
ultra-low attachment 6-well plate for eight days and then transferred to Matrigel-coated 6-
well plate in differentiation medium. The cells were fixed for immunocytochemical analysis
or harvested for RT-PCR tests eight days later. Differentiation medium: DMEM/F12, 10%
FBS, 1% Glutamax, 1% Non-essential amino acids, 1% penicillin/streptomycin, 0.1 mM β -
10 mercaptoethanol.

Alkaline phosphatase staining and immunocytochemistry assay

[0141] Alkaline Phosphatase staining was performed according to the manufacturer's
protocol using the Alkaline Phosphatase Detection Kit (Stemgent). Standard
immunocytochemistry assay was carried out as previously reported (Li, W. *et al.*, *Stem Cells*
15 27:2992 3000 (2009)). Primary antibodies used can be found in the Table 2. Secondary
antibodies were Alexa Fluor 488 donkey anti-mouse or anti-rabbit IgG (1:1000) (Invitrogen).
Nuclei were visualized by DAPI (Sigma-Aldrich) staining. Images were captured using a
Nikon Eclipse TE2000-U microscope.

Gene expression analysis by RT-PCR and qRT-PCR

20 [0142] For RT-PCR and qRT-PCR analysis, total RNA was extracted from human iPSCs
using the RNeasy Plus Mini Kit in combination with QIAshredder (Qiagen). First strand
reverse transcription was performed with 2 μ g RNA using iScriptTM cDNA Synthesis Kit
(BioRad). The expression of pluripotency markers was analyzed by RT-PCR using Platinum
PCR SuperMix (Invitrogen). The expression of lineage specific markers after differentiation
25 was analyzed by qRT-PCR using iQ SYBR Green Supermix (Bio-Rad). The primers can be
found in the Table 1.

Microarray analysis

[0143] The Human Ref-8_v3 expression Beadchip (Illumina, CA, USA) was used for
microarray hybridizations to examine the global gene expression of NHEKs, hiPSC and hES
30 cells. Biotin-16-UTP-labeled cRNA was synthesized from 500 ng total RNA with the
Illumina TotalPrep RNA amplification kit (Ambion AMIL1791, Foster City, CA, USA). The
hybridization mix containing 750 ng of labeled amplified cRNA was prepared according to

the Illumina BeadStation 500× System Manual (Illumina, San Diego, CA, USA) using the supplied reagents and GE Healthcare Streptavidin-Cy3 staining solution. Hybridization to the Illumina Human Ref-8_v3 expression Beadchip was for 18 h at 55 °C on a BeadChip Hybridization Wheel. The array was scanned using the Illumina BeadArray Reader. All samples were prepared in two biological replicates. Processing and analysis of the microarray data were performed with the Illumina BeadStudio software. The data were subtracted for background and normalized using the rank invariant option.

Bisulfate genomic sequencing

[0144] Genomic DNAs were isolated using the Non Organic DNA Isolation Kit (Millipore) and then treated with the EZ DNA Methylation-Gold Kit (Zymo Research Corp., Orange, CA). The treated DNAs were then used as templates to amplify sequences of interest. Primers used for OCT4 and NANOG promoter fragment amplification are indicated in Table 1. The resulting fragments were cloned using the TOPO TA Cloning Kit for sequencing (Invitrogen) and sequenced.

15 Genotyping of hiPSCs

[0145] Genotyping of hiPSC lines was performed using RT-PCR of genomic DNA with specific primers (Table 1; Yu, J. *et al.*, *Science* 318:1917-1920 (2007) and Li, W. *et al.*, *Stem Cells* 27:2992-3000 (2009)).

Teratoma formation

20 [0146] The hiPSC lines were harvested by using 0.05 % Trypsin-EDTA. Five million cells were injected under the kidney capsule of *SCID* mice (n=3). After 4-6 weeks, well developed teratomas were harvested, fixed and then histologically analyzed at TSRI histology core facility.

Table 1. Primers used

Gene	Forward	Reverse
For RT-PCR		
<i>Endo-OCT4</i>	AGTTTGTGCCAGGGTTTTTG	ACTTCACCTTCCCTCCAACC
<i>Endo-SOX2</i>	CAAAAATGGCCATGCAGGTT	AGTTGGGATCGAACAAAAGCTATT
<i>Endo-NANOG</i>	TTTGAAGCTGCTGGGGAAG	GATGGGAGGAGGGGAGAGGA
<i>Endo-KLF4</i>	ACGATCGTGGCCCCGAAAAGGACC	GATTGTAGTGCTTTCTGGCTGGGCTCC
<i>Endo-cMYC</i>	GCGTCCTGGGAAGGGAGATCCGGAGC	TTGAGGGGCATCGTCGCGGGAGGCTG
<i>REX1</i>	CAGATCCTAAACAGCTCGCAGAAT	GCGTACGCAAATTAAGTCCAGA
<i>UTF1</i>	CCGTCGCTGAACACCGCCCTGCTG	CGCGCTGCCAGAATGAAGCCAC
<i>TDGF2</i>	CTGCTGCCTGAATGGGGGAACCTGC	GCCACGAGGTGCTCATCCATCACAAAGG

Gene	Forward	Reverse
<i>FGF4</i>	CTACAACGCCTACGAGTCTACA	GTTGCACCAGAAAAGTCAGAGTTG
<i>Exo-OCT4</i>	TGTCTCCGTCACCACTCTGG	ATGCATGCGGATCCTTCG
<i>PAX6</i>	TGTCCAACGGATGTGAGT	TTCCCAAGCAAAGATGGAC
<i>βIII TUBULIN</i>	CAACAGCACGGCCATCCAGG	CTTGGGGCCCTGGGCTCCGA
<i>FOXF1</i>	AAAGGAGCCACGAAGCAAGC	AGGCTGAAGCGAAGGAAGAGG
<i>HAND1</i>	TCCCTTTCCGCTTGCTCTC	CATCGCCTACCTGATGGACG
<i>AFP</i>	AGCAGCTTGGTGGTGGATGA	CCTGAGCTTGGCACAGATCCT
<i>GATA6</i>	TGTGCGTTCATGGAGAAGATCA	TTTGATAAGAGACCTCATGAACCGACT
<i>GAPDH</i>	GTGGACCTGACCTGCCGTCT	GGAGGAGTGGGTGTGCTGT
For bisulfate-sequencing		
<i>OCT4-1</i>	TTAGGAAAATGGGTAGTAGGGATTT	TACCCAAAAACAAATAAATTATAAAACCT
<i>OCT4-2</i>	GGATGTTATTAAGATGAAGATAGTTGG	CCTAAACTCCCCTTCAAAAATCTATT
<i>NANOG</i>	GAGTTAAAGAGTTTTGTTTTTAAAAAATTAT	TCCCAAATCTAATAATTTATCATATCTTTC
For genotyping		
<i>OCT4-Int</i>	CAGTGCCCGAAACCCACAC	AGAGGAACTGCTTCCTTCACGACA
<i>SOX2-Int</i>	TACCTCTTCCCTCCCACTCCA	AGAGGAACTGCTTCCTTCACGACA
<i>KLF4-Int</i>	CACCTGCCTTACACATGAAGAGG	CGTAGAATCGAGACCGAGGAGA

Table 2. Primary antibodies applied

Antibody	Species	Dilution	Vendor
Anti-OCT4 (1)	Mouse	1:500	Santa Cruz Biotechnology
Anti-OCT4 (2)	Rabbit	1:500	Stemgent
Anti-SOX2	Rabbit	1:1000	Chemicon
Anti-NANOG	Rabbit	1:500	Abcam
Anti-SSEA4	Mouse	1:500	Stemgent
Anti-TRA-1-81	Mouse	1:500	Stemgent
TUJ1 (Anti- β III TUBULIN)	Mouse	1:3000	Covance Research Products
Anti-SMA	Mouse	1:500	Sigma
Anti-AFP	Mouse	1:500	Sigma

Table 3. Summary of reprogramming experiments

Donor Cells	Induction factors	Chemicals	Experiments	TRA-1-81 positive colonies
	OCT4+KLF4+SOX2+MYC	DMSO	#1	17
			#2	20
			#3	23
		A83+PD	#1	72
			#2	104
			#3	91
	OCT4+KLF4+SOX2	DMSO	#1	2
			#2	3
			#3	8
		A83+PD	#1	26
			#2	35
			#3	44

Donor Cells	Induction factors	Chemicals	Experiments	TRA-1-81 positive colonies
NHEKs (lot number: 0000087940)	OCT4+KLF4	A83+PD	#1	1
			#2	2
			#3	0
		A83+PS48+PD	#1	15
			#2	18
			#3	5
		A83+VPA+PD	#1	6
			#2	0
			#3	3
		A83+NaB+PD	#1	20
			#2	17
			#3	18
	A83+PS48+NaB+PD	#1	21	
		#2	30	
		#3	27	
OCT4	A83+PS48+NaB+PD	#1	4	
		#2	0	
		#3	3	
NHEKs (lot number: 2F0661)	OCT4	A83+PS48+NaB+PD	#1	2
			#2	3
			#3	0
AHEKs	OCT4	A83+PS48+NaB+PD +Par+CHIR	#1	3
			#2	2
HUVECs	OCT4	A83+PS48+NaB+PD	#1	4
			#2	7
			#3	4
HUVECs	OCT4	A83+PS48+NaB+PD +Par+CHIR	#1	23
			#2	17
AFDCs	OCT4	A83+PS48+NaB+PD +Par+CHIR	#1	5
			#2	11

[0147] NHEKs, Neonatal Human Epidermal Keratinocytes; HUVECs, Human Umbilical Vein Endothelial Cells; AHEKs, Adult Human Epidermal Keratinocytes; AFDCs, Amniotic Fluid Derived Cells. Chemical concentration used: PD, 0.5 μ M PD0325901; A83, 0.5 μ M A-83-01; PS48, 5 μ M PS48; VPA, 0.5mM Valproic acid; NaB, 0.25 mM Sodium butyrate; Par, 2 μ M Parnate; CHIR, 3 μ M CHIR99021. For four-factor or three-factor induced reprogramming, NHEKs were seeded at a density of 100,000 transduced cells per 10 cm dish and positive colonies were counted four weeks later; For two-factor induced reprogramming, NHEKs were seeded at a density of 100,000 transduced cells per 10 cm dish and positive colonies were counted six weeks later; and for one-factor induced reprogramming, NHEKs and AHEKs were seeded at a density of 1,000,000 transduced cells per 10 cm dish and positive colonies were counted eight weeks later. HUVECs and AFDCs were seeded at a density of 200,000 transduced cells per 10 cm dish and positive colonies were counted six weeks later.

Table 4. Characterization of established human iPSC cell lines

hiPSC clone	Induction factors	Cell source	Marker expression	RT-PCR test	EB differentiation	Teratoma test
hiPSC-OK#1	OCT4+KLF4	NHEKs	√	√	√	√
hiPSC-OK#3			√	√	√	
hiPSC-O#1	OCT4	NHEKs	√	√	√	√
hiPSC-O#3			√	√	√	
hiPSC-O#4			√			
hiPSC-O#5			√			
2 more lines						
hiPSC-O#21	OCT4	HUVECs	√	√	√	√
hiPSC-O#22			√			
hiPSC-O#26			√	√	√	
hiPSC-O#31			√	√	√	√
7 more lines						
hiPSC-O#52	OCT4	AHEKs	√		√	
hiPSC-O#57			√			
hiPSC-O#63	OCT4	AFDCs	√		√	
hiPSC-O#65			√			

[0148] Those cell lines characterized were long-term expanded for over 20 passages under conventional hESC culture condition and further characterized for marker expression and pluripotency; while other cell lines established were stored at passage 5 or 6. Blank entries indicate "not determined."

Table 5. DNA fingerprint analysis on Oct4-induced iPSCs and parental cell lines

Genomic loci	NHEK (pooled)	hiPSC-O#1	HUVEC	hiPSC-O#21
Amelogenin	X, Y	X, Y	X	X
vWA	11, 15, 17, 18, 19	15, 18	15; 16	15; 16
D8S1179	10, 13, 16	13,	10; 13	10; 13
TPOX	8, 9, 11, 12	8	8	8
FGA	19, 22, 23, 24	19, 22	24; 27	24; 27
D3S1358	13, 14, 15, 17	17	14; 16	14; 16
THO1	6, 7, 9, 9.3	7, 9	6	6
D21S11	24.2, 29, 30.2, 35	24.2, 29	28; 30.2	28; 30.2
D18S51	13, 14, 16, 17, 18, 19	13, 17	13; 18	13; 18
Penta E	5, 8, 13, 14, 19	13, 19	12	12
D5S818	8, 11, 12, 13	11, 13	12; 13	12; 13
D13S317	8, 9, 11, 12, 13	9, 12	11; 14	11; 14
D7S820	8, 9, 10, 11	9, 10	11	11

Genomic loci	NHEK (pooled)	hiPSC-O#1	HUVEC	hiPSC-O#21
D16S539	9, 10, 11, 12, 13	9, 13	9; 11	9; 11
CSF1PO	10, 11, 12	11, 12	11; 12	11; 12
Penta D	2.2, 10, 12	10	12; 13	12; 13

[0149] Fifteen polymorphic short tandem repeat (STR) DNA loci and the sex chromosome marker amelogenin were investigated.

Example 2: Reprogramming of human umbilical vein endothelial cells

- 5 [0150] We tested the effects of the combination of a HDAC inhibitor, a PDK1 activator, a TGF β receptor inhibitor, and a MEK inhibitor on HUVECs that were lentivirally transduced with *Oct4* alone for their effects on reprogramming kinetics and efficiency.

Methods

- [0151] Human Umbilical Vein Endothelial Cells (HUVECs, Millipore) were maintained in
10 EndoGRO-VEGF Complete Medium (HCM, CHEMICON). HUVECs were cultured in a 100 mm tissue culture dish and transduced 2 times (4-6 hours/time) with freshly produced lentivirus supernatants. Then 200,000 transduced HUVECs were seeded on gelatin coated 100-mm dish and cultured in HCM and treated with PDK1 activator PS48 (5 μ M), HDAC inhibitor NaB (0.25 mM), and TGF β receptor inhibitor A-83-01(0.5 μ M) for 2 weeks,
15 followed by changing half volume of media to hESCM and supplementing with PDK1 activator PS48 (5 μ M), HDAC inhibitor NaB (0.25 mM), and TGF β receptor inhibitor A-83-01(0.5 μ M) for another 2 weeks. Then cell culture media were changed to hESCM and supplemented with PDK1 activator PS48 (5 μ M), HDAC inhibitor NaB (0.25 mM), and TGF β receptor inhibitor A-83-01(0.5 μ M) and MEK inhibitor PD0325901 (0.5 μ M) for
20 additional 2 weeks. The iPSC colonies were stained positive by Alexa Fluor 555 Mouse anti-Human TRA-1-81 antibody (BD Pharmingen). hESCM: DMEM/F12, 15% Knockout serum replacement, 1% Glutamax, 1% Non-essential amino acids, 1% penicillin/streptomycin, 0.1 mM β -mercaptoethanol and 10 ng/ml bFGF.

Results

- 25 [0152] For HUVECs transduced with *Oct4* alone, we tested the effects of the combination of a HDAC inhibitor, a PDK1 activator, a TGF β receptor inhibitor, a MEK inhibitor on reprogramming efficiency. We found that treatment with the combination of 5 μ M PS48, 0.25 mM NaB, 0.5 μ M A-83-01 and 0.5 μ M PD0325901 results in a ~0.0015% reprogramming efficiency.

[0153] The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims.

SEQUENCE LISTING IN ELECTRONIC FORM

This description contains a sequence listing in electronic form in ASCII text format. A copy of the sequence listing in electronic form is available from the Canadian Intellectual Property Office. The sequences in the sequence listing in electronic form are reproduced in the following Table.

SEQUENCE TABLE

SEQ ID NO: 1

```
Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg
 1           5           10           15
Arg Arg Arg Arg Arg Arg Arg Arg
          20           25
```

SEQ ID NO: 2

```
Gly Ser Pro Pro Thr Ala Pro Thr Arg Ser Lys Thr Pro Ala Gln Gly
 1           5           10           15
Leu Ala Arg Lys Leu His Phe Ser Thr Ala Pro Pro Asn Pro Asp Ala
          20           25           30
Pro Trp Thr Pro Arg Val Ala Gly Phe Asn Lys Arg Val Phe Arg Phe
          35           40           45
Ser Pro Gln Thr Ala Arg Arg Ala Thr Thr Thr Arg Ile
          50           55           60
```

SEQ ID NO: 3

```
Ala Gly Ser Gly Gly Ala Ala Val Ala Leu Leu Pro Ala Val Leu Leu
 1           5           10           15
Ala Leu Leu Ala Pro Gly Gly Glu Phe Ala
          20           25
```

SEQ ID NO: 4

```
Ala Gly Ser Gly Gly Tyr Ala Arg Ala Ala Ala Arg Gln Ala Arg Ala
 1           5           10           15
Gly Gly Glu Phe Ala
          20
```

SEQ ID NO: 5

```
Arg Gln Ile Lys Ile Trp Phe Gln Gly Arg Arg Met Lys Trp Lys Lys
 1           5           10           15
```

SEQ ID NO: 6

```
Tyr Gly Arg Lys Lys Arg Arg Gln Arg Arg Arg
 1           5           10
```

SEQ ID NO: 7

```
Met Val Thr Val Leu Phe Arg Arg Leu Arg Ile Arg Arg Ala Cys Gly
 1           5           10           15
Pro Pro Arg Val Arg Val
          20
```

SEQ ID NO: 8

Ala Gly Tyr Leu Leu Gly Lys Ile Gly Leu Lys Ala Leu Ala Ala Leu
 1 5 10 15
 Ala Lys Lys Ile Leu
 20

SEQ ID NO: 9

Arg Arg Arg Arg Arg Arg Arg Arg
 1 5

SEQ ID NO: 10

Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg Arg
 1 5 10

SEQ ID NO: 11

Glu Ser Gly Gly Gly Gly Ser Pro Gly Arg Arg Arg Arg Arg Arg Arg Arg
 1 5 10 15
 Arg Arg Arg Arg
 20

SEQ ID NO: 12

agtttgtgcc agggtttttg 20

SEQ ID NO: 13

acttcacctt cctccaacc 20

SEQ ID NO: 14

caaaaatggc catgcagggt 20

SEQ ID NO: 15

agttgggatc gaacaaaagc tatt 24

SEQ ID NO: 16

tttgaagct gctggggaag 20

SEQ ID NO: 17

gatgggagga ggggagagga 20

SEQ ID NO: 18

acgatcgtgg ccccgaaaa ggacc 25

SEQ ID NO: 19

gattgtagtg cttctgget gggetcc 27

SEQ ID NO: 20

gcgtcctggg aaggagatc cggagc 26

SEQ ID NO: 21

ttgaggggca tcgtcgcggg aggetg 26

SEQ ID NO: 22

cagatcctaa acagctcgca gaat 24

SEQ ID NO: 23

gcgtacgcaa attaaagtcc aga 23

SEQ ID NO: 24 ccgtcgctga acaccgccct gctg	24
SEQ ID NO: 25 cgcgctgccc agaatgaagc ccac	24
SEQ ID NO: 26 ctgctgcctg aatgggggaa cctgc	25
SEQ ID NO: 27 gccacgaggt gctcatccat cacaagg	27
SEQ ID NO: 28 ctacaacgcc tacgagtcct aca	23
SEQ ID NO: 29 gttgcaccag aaaagtcaga gttg	24
SEQ ID NO: 30 tgtctccgtc accactctgg	20
SEQ ID NO: 31 atgcatgcgg atccttcg	18
SEQ ID NO: 32 tgtccaacqg atgtgagt	18
SEQ ID NO: 33 tttccaagc aaagatggac	20
SEQ ID NO: 34 caacagcacy gccatccagg	20
SEQ ID NO: 35 cctggggccc tgggcctccg a	21
SEQ ID NO: 36 aaaggagcca cgaagcaagc	20
SEQ ID NO: 37 aggctgaagc gaaggaagag g	21
SEQ ID NO: 38 tccttttcc gcttgctctc	20
SEQ ID NO: 39 catgcctac ctgatggacg	20
SEQ ID NO: 40 agcagcttgg tggatgatga	20
SEQ ID NO: 41 cctgagcttg gcacagatcc t	21

SEQ ID NO: 42	tgtgcgttca tggagaagat ca	22
SEQ ID NO: 43	tttgataaga gacctcatga accgact	27
SEQ ID NO: 44	gtggacctga cctgccgtct	20
SEQ ID NO: 45	ggaggagtgg gtgtcgctgt	20
SEQ ID NO: 46	ttaggaaaat gggtagtagg gattt	25
SEQ ID NO: 47	tacccaaaaa acaaataaat tataaacct	30
SEQ ID NO: 48	ggatggttatt aagatgaaga tagttgg	27
SEQ ID NO: 49	cctaaactcc cttcaaaat ctatt	25
SEQ ID NO: 50	gagttaaaga gttttqtttt taaaaattat	30
SEQ ID NO: 51	tcccaaatct aataatttat catatctttc	30
SEQ ID NO: 52	cagtgcccgga aaccacac	19
SEQ ID NO: 53	agaggaactg cttccttcac gaca	24
SEQ ID NO: 54	tacctcttcc tcccactcca	20
SEQ ID NO: 55	agaggaactg cttccttcac gaca	24
SEQ ID NO: 56	caccttgect tacacatgaa gagg	24
SEQ ID NO: 57	cgtagaatcg agaccgagga ga	22

CLAIMS

1. An *in vitro* method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell, the method comprising:

(a) contacting the non-pluripotent cell with an Oct4 polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding an Oct4 polypeptide;

(b) contacting the non-pluripotent cell with (i) a TGF β receptor/ALK5 inhibitor or a MEK inhibitor, or (ii) a TGF β receptor/ALK5 inhibitor and a MEK inhibitor;

(c) contacting the non-pluripotent cell with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient to induce the cell to become a pluripotent stem cell, thereby inducing the non-pluripotent mammalian cell into an induced pluripotent stem cell; and

(d) one or more of (i) contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, (ii) contacting the non-pluripotent cell with a Klf polypeptide or (iii) introducing into the non-pluripotent cell a polynucleotide encoding a Klf polypeptide.

2. The method of claim 1, wherein the PDK1 activator is:

(a) an allosteric PDK1 activator; or

(b) (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid ("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z").

3. The method of claim 1 or 2, wherein step (d) comprises:

(i) contacting the non-pluripotent cell with the Klf polypeptide; or

(ii) introducing into the non-pluripotent cell the polynucleotide encoding the Klf polypeptide.

4. The method of claim 1 or 2, wherein step (d) comprises:

(i) contacting the non-pluripotent cell with the HDAC inhibitor and the Klf polypeptide;
or

(ii) contacting the non-pluripotent cell with the HDAC inhibitor and introducing into the non-pluripotent cell the polynucleotide encoding the Klf polypeptide.

5. The method of any one of claims 1 to 4, wherein the Klf polypeptide is Klf4.

6. The method of any one of claims 1 to 5, further comprising one or more of:

(a) contacting the non-pluripotent cell with a Sox polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding a Sox polypeptide; and

(b) contacting the non-pluripotent cell with a Myc polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding a Myc polypeptide.

7. The method of claim 6, wherein the Sox polypeptide is Sox2.

8. The method of any one of claims 1 to 7, wherein the Oct4 polypeptide contacted with the non-pluripotent cell comprises an amino acid sequence that enhances transport across cell membranes.

9. The method of claims 6 or 7, wherein one or more of the Oct4 polypeptide, Klf polypeptide, Sox polypeptide, and Myc polypeptide contacted with the non-pluripotent cell comprises an amino acid sequence that enhances transport across cell membranes.

10. The method of any one of claims 1 to 9, wherein the non-pluripotent cell is a human cell.

11. The method of any one of claims 1 to 10, wherein the method comprises:

(a) contacting the non-pluripotent cell with the PDK1 activator in the absence of the MEK inhibitor, followed by contacting the non-pluripotent cell with the PDK1 activator and the MEK inhibitor; or

(b) contacting the non-pluripotent cell with the PDK1 activator, the TGF β receptor/ALK5 inhibitor, and the histone deacetylase (HDAC) inhibitor in the absence of the MEK

inhibitor, followed by contacting the non-pluripotent cell with the PDK1 activator, the TGF β receptor/ ALK5 inhibitor, the histone deacetylase (HDAC) inhibitor and the MEK inhibitor.

12. The method of any one of claims 1 to 11, wherein the HDAC inhibitor is sodium butyrate or valproic acid.

13. The method of any one of claims 1 to 12, wherein the TGF β receptor/ ALK5 inhibitor is A-83-01.

14. The method of any one of claims 1 to 13, wherein the MEK inhibitor is PD0325901.

15. A mixture comprising: mammalian cells, a PDK1 activator, an Oct4 polypeptide, and:

(a) a TGF β receptor/ ALK5 inhibitor, a MEK inhibitor, or a combination thereof; and

(b) one or more of (i) a histone deacetylase (HDAC) inhibitor, (ii) a Klf polypeptide, or (iii) a polynucleotide encoding a Klf polypeptide.

16. The mixture of claim 15, wherein the PDK1 activator is:

(a) an allosteric PDK1 activator; or

(b) (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid ("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z").

17. The mixture of claim 15 or 16, wherein the mixture comprises:

(a) the Klf polypeptide; or

(b) the polynucleotide encoding the Klf polypeptide.

18. The mixture of claim 15 or 16, wherein the mixture comprises:

(a) the HDAC inhibitor and the Klf polypeptide, or

(b) the HDAC inhibitor and the polynucleotide encoding the Klf polypeptide.

19. The mixture of any one of claims 15 to 18, wherein the Klf polypeptide is Klf4.
20. The mixture of claim 17, 18, or 19, further comprising a Sox polypeptide, a Myc polypeptide, or a combination thereof.
21. The mixture of claim 20, wherein the Sox polypeptide is Sox2.
22. The mixture of any one of claims 15 to 21, wherein at least 99% of the cells are non-pluripotent cells.
23. The mixture of any one of claims 15 to 22, wherein the cells are human cells.
24. The mixture of any one of claims 15 to 23, wherein the mixture comprises the PDK1 activator, the TGF β receptor/ALK5 inhibitor, the MEK inhibitor, and the Oct4 polypeptide.
25. The mixture of any one of claims 15 to 24, wherein the HDAC inhibitor is sodium butyrate or valproic acid.
26. The mixture of any one of claims 15 to 25, wherein the TGF β receptor/ ALK5 inhibitor is A-83-01.
27. The mixture of any one of claims 15 to 26, wherein the MEK inhibitor is PD0325901.
28. An *in vitro* method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell, the method comprising:
 - (a) contacting the non-pluripotent cell with an Oct4 polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding an Oct4 polypeptide;
 - (b) contacting the non-pluripotent cell with a compound that promotes glycolytic metabolism, a small molecule TGF β receptor/ALK5 inhibitor, and a small molecule MEK inhibitor, thereby inducing the non-pluripotent mammalian cell to reprogram into an induced pluripotent stem cell,
wherein the compound that promotes glycolytic metabolism comprises F2,6P (fructose 2,6-bisphosphate), F6P (fructose 6-phosphate), DNP (2,4-dinitrophenol), NOG (N-

oxalylglycine), QC (quercetin), 2-HA (2-hydroxyglutaric acid), NA (nicotinic acid), or a PDK1 activator; and

(c) one or more of (i) contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, (ii) contacting the non-pluripotent cell with a Klf polypeptide or (iii) introducing into the non-pluripotent cell a polynucleotide encoding a Klf polypeptide.

29. The method of claim 28, wherein the PDK1 activator is:

(a) an allosteric PDK1 activator; or

(b) (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid ("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z").

30. The method of claim 29, wherein step (c) comprises:

(i) contacting the non-pluripotent cell with the Klf polypeptide; or

(ii) introducing into the non-pluripotent cell the polynucleotide encoding the Klf polypeptide.

31. The method of claim 28 or 29, wherein step (c) comprises

(i) contacting the non-pluripotent cell with the HDAC inhibitor and the Klf polypeptide;

or

(ii) contacting the non-pluripotent cell with the HDAC inhibitor and introducing into the non-pluripotent cell the polynucleotide encoding the Klf polypeptide.

32. The method of any one of claims 28 to 31, wherein the Klf polypeptide is Klf4.

33. The method of claim 30, 31, or 32, further comprising one or more of:

(a) contacting the non-pluripotent cell with a Sox polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding a Sox polypeptide; and

(b) contacting the non-pluripotent cell with a Myc polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding a Myc polypeptide.

34. The method of claim 33, wherein the Sox polypeptide is Sox2.

35. The method of any one of claims 28 to 34, wherein the Oct4 polypeptide contacted with the non-pluripotent cell comprises an amino acid sequence that enhances transport across cell membranes.

36. The method of claim 33 or 34, wherein one or more of the Oct4 polypeptide, Klf polypeptide, Sox polypeptide, and Myc polypeptide contacted with the non-pluripotent cell comprises an amino acid sequence that enhances transport across cell membranes.

37. The method of any one of claims 28 to 36, wherein the non-pluripotent cell is a human cell.

38. The method of any one of claims 28 to 37, wherein the method comprises:

(a) contacting the non-pluripotent cell with the compound that promotes glycolytic metabolism in the absence of the MEK inhibitor, followed by contacting the non-pluripotent cell with the compound that promotes glycolytic metabolism and the MEK inhibitor; or

(b) contacting the non-pluripotent cell with the compound that promotes glycolytic metabolism, the TGF β receptor/ALK5 inhibitor, and the HDAC inhibitor in the absence of the MEK inhibitor, followed by contacting the non-pluripotent cell with the compound that promotes glycolytic metabolism, the TGF β receptor/ALK5 inhibitor, the HDAC inhibitor and the MEK inhibitor.

39. The method of claim 37 or 38, wherein the HDAC inhibitor is sodium butyrate or valproic acid.

40. The method of any one of claims 28 to 39, wherein the TGF β receptor/ ALK5 inhibitor is A-83-01.

41. The method of any one of claims 28 to 40, wherein the MEK inhibitor is PD0325901.

42. A mixture comprising non-pluripotent mammalian cells, a compound that promotes glycolytic metabolism, one or more exogenous transcription factors, a small molecule TGF β receptor/ALK5 inhibitor, and a small molecule MEK inhibitor, wherein:

(a) the compound that promotes glycolytic metabolism comprises F2,6P (fructose 2,6-bisphosphate), F6P (fructose 6-phosphate), DNP (2,4-dinitrophenol), NOG (N-oxalylglycine), QC (quercetin), 2-HA (2-hydroxyglutaric acid), NA (nicotinic acid), or a PDK1 activator;

(b) the one or more exogenous transcription factors comprises an Oct4 polypeptide; and

(c) (i) the one or more exogenous transcription factors further comprises a Klf polypeptide; or (ii) the mixture comprises an HDAC inhibitor.

43. The mixture of claim 42, wherein the mixture comprises the HDAC inhibitor.

44. The mixture of claim 42 or 43, wherein the one or more exogenous transcription factors comprises the Oct4 polypeptide and the Klf polypeptide.

45. The mixture of claim 42, 43, or 44 wherein the Klf polypeptide is Klf4.

46. The mixture of claim 44 or 45, wherein the one or more exogenous transcription factors further comprise a Sox polypeptide, a Myc polypeptide, or a combination thereof.

47. The mixture of claim 46, wherein the Sox polypeptide is Sox2.

48. The mixture of any one of claims 42 to 47, wherein the non-pluripotent cells are human cells.

49. The mixture of any one of claims 42 to 48, wherein the HDAC inhibitor is sodium butyrate or valproic acid.

50. The mixture of any one of claims 42 to 49, wherein the TGF β receptor/ALK5 inhibitor is A-83-01.

51. The mixture of any one of claims 42 to 50, wherein the MEK inhibitor is PD0325901.

52. An *in vitro* method of inducing a non-pluripotent mammalian cell into an induced pluripotent stem cell, the method comprising:

(a) contacting the non-pluripotent cell with an Oct4 polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding an Oct4 polypeptide, wherein no c-Myc polypeptide and no polynucleotide encoding c-Myc are introduced into the non-pluripotent cell;

(b) contacting the non-pluripotent cell with a 3'-phosphoinositide-dependent kinase-1 (PDK1) activator under conditions sufficient to induce the cell to become a pluripotent stem cell, thereby inducing the non-pluripotent mammalian cell into an induced pluripotent stem cell; and

(c) one of more of (i) contacting the non-pluripotent cell with a histone deacetylase (HDAC) inhibitor, (ii) contacting the non-pluripotent cell with a Klf polypeptide, or (iii) introducing into the non-pluripotent cell a polynucleotide encoding a Klf polypeptide.

53. The method of claim 52, wherein the PDK1 activator is:

(a) an allosteric PDK1 activator; or

(b) (Z)-5-(4-Chlorophenyl)-3-phenylpent-2-enoic acid ("PS48"), (Z)-5-(4-Bromo-2-fluorophenyl)-3-phenylpent-2-enoic acid ("PS08"), 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ("12Z"), or (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ("13Z").

54. The method of claim 52 or 53, further comprising contacting the non-pluripotent cell with (a) a TGF β receptor/ALK5 inhibitor or a MEK inhibitor, or (b) a TGF β receptor/ALK5 inhibitor and a MEK inhibitor.

55. The method of claim 52 or 53, wherein the method comprises:

(a) contacting the non-pluripotent cell with the PDK1 activator in the absence of a MEK inhibitor, followed by contacting the non-pluripotent cell with the PDK1 activator and a MEK inhibitor; or

(b) contacting the non-pluripotent cell with the PDK1 activator, a TGF β receptor/ ALK5 inhibitor, and the histone deacetylase (HDAC) inhibitor in the absence of a MEK inhibitor, followed by contacting the non-pluripotent cell with the PDK1 activator, the TGF β receptor/ALK5 inhibitor, the histone deacetylase (HDAC) inhibitor and the MEK inhibitor.

56. The method of claim 54 or 55, wherein the TGF β receptor/ ALK5 inhibitor is A-83-01.

57. The method of claim 54, 55, or 56, wherein the MEK inhibitor is PD0325901.

58. The method of any one of claims 52 to 57, wherein step (c) comprises:

(i) contacting the non-pluripotent cell with the Klf polypeptide; or

(ii) introducing into the non-pluripotent cell the polynucleotide encoding the Klf polypeptide.

59. The method of any one of claims 52 to 57, wherein step (c) comprises:

(i) contacting the non-pluripotent cell with the HDAC inhibitor and the Klf polypeptide;

or

(ii) contacting the non-pluripotent cell with the HDAC inhibitor and introducing into the non-pluripotent cell the polynucleotide encoding the Klf polypeptide.

60. The method of any one of claims 52 to 59, wherein the Klf polypeptide is Klf4.

61. The method of claim 58, 59, or 60, further comprising contacting the non-pluripotent cell with a Sox polypeptide or introducing into the non-pluripotent cell a polynucleotide encoding a Sox polypeptide.

62. The method of claim 61, wherein the Sox polypeptide is Sox2.

63. The method of any one of claims 52 to 62, wherein the Oct4 polypeptide contacted with the non-pluripotent cell comprises an amino acid sequence that enhances transport across cell membranes.

64. The method of claim 61 or 62, wherein one or more of the Oct4 polypeptide, the Klf polypeptide, and the Sox polypeptide contacted with the non-pluripotent cell comprises an amino acid sequence that enhances transport across cell membranes.

65. The method of any one of claims 52 to 64, wherein the non-pluripotent cell is a human cell.

66. The method of any one of claims 52 to 65, wherein the HDAC inhibitor is sodium butyrate or valproic acid.

67. A small molecule for use in manufacturing a pluripotent stem cell from a non-pluripotent cell for application in regenerative medicine or tissue regeneration, wherein the small molecule enhances reprogramming of a mammalian cell, wherein the small molecule comprises a PDK1 (3'-phosphoinositide-dependent kinase-1) activator, and wherein the PDK1 activator comprises 2-(3-(4-Chlorophenyl)-3-oxo-1-phenylpropylthio)acetic acid, (Z)-5-(Naphthalen-2-yl)-3-phenylpent-2-enoic acid ('12Z'), (Z)-5-(1H-Indol-3-yl)-3-phenylpent-2-enoic acid ('13Z'), or a 3,5-diphenylpent-2-enoic acid.

68. A method of making a therapeutic composition comprising an induced pluripotent stem cell (iPSC) for tissue regeneration or regenerative medicine in a subject in need thereof, the method comprising:

(a) obtaining an iPSC induced according to a method as defined in any one of claims 1 to 14, 28 to 41 or 52 to 66, and

(b) making the therapeutic composition comprising the iPSC for tissue regeneration or regenerative medicine in the subject in need thereof.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

6/11




Fig. 6

7/11





Fig. 7

8/11

*Fig. 8*

9/11




Fig. 9

10/11





Fig. 10

11/11




Fig. 11

