



(86) Date de dépôt PCT/PCT Filing Date: 1998/02/24

(87) Date publication PCT/PCT Publication Date: 1998/09/03

(45) Date de délivrance/Issue Date: 2003/04/22

(85) Entrée phase nationale/National Entry: 1999/08/19

(86) N° demande PCT/PCT Application No.: EP 1998/001275

(87) N° publication PCT/PCT Publication No.: 1998/038186

(30) Priorités/Priorities: 1997/02/27 (PCT/EP97/00995) EP;
1997/07/25 (9715783.8) EP

(51) Cl.Int.⁶/Int.Cl.⁶ C07D 401/14, A61K 31/495

(72) Inventeurs/Inventors:

STOBIE, ALAN, GB;
GAUTIER, ELISABETH COLETTE LOUISE, FR;
WAITE, DAVID CHARLES, GB;
CROOK, ROBERT JAMES, GB


(73) Propriétaire/Owner:

PFIZER INC., US

(74) Agent: SMART & BIGGAR

(54) Titre : QUINOXALINEDIONES

(54) Title: QUINOXALINEDIONES



(I)


(57) Abrégé/Abstract:

The present invention provides a substantially pure compound of formula (I) (see formula I) or a pharmaceutically acceptable salt or solvate thereof, together with compositions containing, uses of, processes for the preparation of and intermediates used in the preparation of such compounds. The compounds are useful for treatment in a mammal of diseases by producing an antagonist effect at an NMDA receptor, for example acute neurodegenerative or chronic neurological disorders.

69387-272

ABSTRACT

The present invention provides a substantially pure compound of formula (I)



(I)

10 or a pharmaceutically acceptable salt or solvate thereof,
together with compositions containing, uses of, processes
for the preparation of and intermediates used in the
preparation of such compounds. The compounds are useful for
treatment in a mammal of diseases by producing an antagonist
15 effect at an NMDA receptor, for example acute
neurodegenerative or chronic neurological disorders.

QUINOXALINEDIONES

This invention relates to 2,3(1H,4H)-quinoxalinedione derivatives which
5 are selective antagonists of N-methyl-D-aspartate receptors. More particularly,
this invention relates to 5-triazolyl-2,3(1H,4H)-quinoxalinedione derivatives and
to the preparation of, compositions containing, and the uses of, such
derivatives.

L-Glutamic acid is an excitatory amino acid neurotransmitter whose
10 physiological role in the brain involves interaction with four receptors, three of
which are named after the selective agonists NMDA (N-methyl-D-aspartate),
AMPA (2-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid) and kainate.
The fourth receptor is termed the metabotropic receptor. In addition to a
binding site for glutamic acid, the NMDA receptor possesses high affinity
15 binding sites for dissociative anaesthetics (e.g. ketamine), polyamines (e.g.
spermine), glycine and certain metal ions (e.g. Mg^{2+} , Zn^{2+}). Since the NMDA
receptor has an absolute requirement to bind glycine for activation to occur,
glycine antagonists can act as functional NMDA antagonists.

In the region of a cerebral infarct, anoxia, for example, causes
20 abnormally high concentrations of glutamic acid to be released. This leads to
an over-stimulation of NMDA receptors resulting in the degeneration and death
of neurones. Thus, NMDA receptor antagonists, which have been shown to
block the neurotoxic effects of glutamic acid in vitro and in vivo, may be useful
in the treatment and/or prevention of any pathological condition in which NMDA
25 receptor activation is thought to be important. Examples of such conditions
include acute neurodegenerative disorders arising from events such as stroke,
transient ischaemic attack, peri-operative ischaemia, global ischaemia
(following cardiac arrest) and traumatic head injury to the brain or spinal cord.
In addition, NMDA antagonists may be of use in treating certain chronic
30 neurological disorders such as senile dementia, Parkinson's disease and
Alzheimer's disease. They may also have utility in conditions in which

-2-

peripheral nerve function has been impaired such as retinal and macular degeneration.

Furthermore, NMDA antagonists have been shown to possess anti-convulsant and anxiolytic activity and may therefore be used to treat epilepsy and anxiety. NMDA antagonists may also attenuate the effects of alcohol withdrawal from physically dependent animals (K.A. Grant *et al.*, J. Pharm.Exp.Ther., **260**, 1017 (1992)) and thus NMDA antagonists may be of use in the treatment of alcohol addiction and pain. NMDA antagonists may also be useful in the treatment of hearing disorders (e.g. tinnitus), migraine and psychiatric disorders.

EP-A-0572852 describes pyrrol-1-yl-substituted 2,3(1H,4H)-quinoxalinedione derivatives useful for the treatment of neurodegenerative illnesses and neurotoxic disorders of the central nervous system.


EP-A-0556393 discloses, *inter alia*, imidazolyl- or triazolyl-substituted 2,3(1H,4H)-quinoxalinedione derivatives with glutamate receptor antagonising activity, particularly NMDA-glycine receptor and AMPA receptor antagonising activities. However, no 5-triazolyl-substituted compounds are specifically described therein.

International Patent Application Publication no. WO 97/32873 discloses 5-heteroaryl-2,3-(1H,4H)-quinoxalinedione derivatives with NMDA receptor antagonist activity. Example 114 of that Application allegedly describes the preparation of (-)-6,7-dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione. However, further analysis of the product of Example 114 shows the stated title compound to be bound to a stoichiometric quantity of silica (see Reference Example 1 herein). This silica complex has been shown to have different properties compared with, and to be distinct, analytically, from, the stated title compound. Example 114 of that Application therefore discloses the preparation of a different compound to the alleged title compound although the skilled person, realising that a silica complex had been obtained, could readily apply common knowledge to prepare the stated title compound therefrom.

-3-

The present compounds are potent antagonists of the NMDA (glycine site) receptor. In addition, they are highly selective antagonists for the NMDA (glycine site) receptor in comparison to the AMPA receptor to which they have
 5 little, if any, affinity.

The present invention provides a novel, substantially pure compound of the formula:



(I)

10

or a pharmaceutically acceptable salt or solvate thereof.

The expression "substantially pure" means the compound preferably is at least of 90% w/w purity, more preferably is at least of 95% w/w purity and most preferably is at least of 98% w/w purity. For the purpose of pharmaceutical
 15 applications, the compound would normally be manufactured to at least 99% w/w purity.

The pharmaceutically acceptable salts of the compounds of the formula (I) include the acid addition and the base salts thereof.

Suitable acid addition salts are formed from acids which form non-toxic
 20 salts and examples are the hydrochloride, hydrobromide, hydroiodide, sulphate, hydrogen sulphate, nitrate, phosphate, hydrogen phosphate, acetate, maleate, fumarate, lactate, tartrate, citrate, gluconate, succinate, benzoate, methanesulphonate, benzenesulphonate and p-toluenesulphonate salts.

-4-

Suitable base salts are formed from bases which form non-toxic salts and examples are the calcium, lithium, magnesium, potassium, sodium, zinc, ethanolamine, diethanolamine and triethanolamine salts.


- 5 For a review on suitable salts see Berge *et al*, J.Pharm.Sci., 66, 1-19 (1977).

Suitable solvates include hydrates.

The compounds of the formula (I) are single stereoisomers known as atropisomers. Atropisomers are isomers that can be separated only because
10 rotation about single bonds is prevented or greatly slowed (see "Advanced Organic Chemistry", Third Edition, Jerry March, John Wiley and Sons (1985)). They may be prepared conventionally from a corresponding optically pure intermediate or by resolution of a racemic mixture containing the opposite stereoisomer. This can be achieved by H.P.L.C. of the corresponding
15 racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of the corresponding racemate with a suitable optically active acid or base.

The compounds of the formula (I) can be prepared by the following
20 methods.


- 1) The compounds of the formula (I) can be prepared by acidic or basic hydrolysis of a compound of the formula:



25

wherein R is group of the formula:

-5-



and R^1 and R^2 , either when taken alone or together, represent a group or groups that can be hydrolytically cleaved under acidic or basic conditions to provide a quinoxalinedione of the formula (I). Such group or groups are conventional and suitable examples will be well-known to the skilled person. Where R is a group of the formula (III), the reaction is followed by separation of the atropisomer of the formula (I) using conventional conditions.

Preferably R^1 and R^2 are either each independently selected from C_1 - C_4 alkyl (preferably methyl or ethyl) and benzyl, optionally ring-substituted by from 1 to 3 substituents each independently selected from C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halo, nitro and trifluoromethyl, or, when taken together, represent C_1 - C_6 alkylene, CH(phenyl), CH(4-methoxyphenyl) or CH(3,4-dimethoxyphenyl).

Preferably, the reaction is carried out by acidic hydrolysis of a compound of the formula (II).


In a typical procedure, a compound of the formula (II) is treated with an aqueous solution of a suitable acid, e.g. a mineral acid such as hydrochloric acid, optionally in the presence of a suitable organic co-solvent, e.g. 1,4-dioxane. The reaction is usually carried out by heating the mixture at up to the reflux temperature of the solvent(s).

The intermediates of the formula (II) can be prepared by conventional methods, for example,

-6-

a) by the route shown in Scheme I:

Scheme I



wherein R, R¹ and R² are as previously defined for a compound of the formula (II).

In a typical procedure, a 5-aminoquinoxaline of the formula (V) is reacted with a compound of the formula:




wherein X¹ is a suitable leaving group, e.g. chloro or bromo, in a suitable solvent, e.g. toluene or dichloromethane, and optionally in the presence of a suitable acid acceptor, e.g. pyridine, to provide an amide of the formula (VI).

An amide of the formula (VI) can be converted to a thioamide of the formula (VII) by treatment with 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulphide (Lawesson's reagent) in a suitable solvent, e.g. toluene or tetrahydrofuran.

-7-

A thioamide of the formula (VII) can be converted to a compound of the formula (II) by treatment with a compound of the formula:

5




in the presence of mercury (II) oxide, optionally a desiccant, e.g. 4A molecular sieves, and a suitable solvent, e.g. n-butanol.

10

A compound of the formula (II) where R is a group of the formula (III) may be resolved to provide a compound of the formula (II) where R is a group of the formula (IV) using conventional techniques, e.g. chiral H.P.L.C.; or


15 b) by using a similar method to that shown in Scheme I to prepare the corresponding pyridine compound of the formula:



(VIII)


20

wherein R³ is a group of the formula:



(IX)

or



(X)


-8-

and R^1 and R^2 are as previously defined for a compound of the formula (II), followed by N-oxidation thereof.


5 The N-oxidation can be performed using 3-chloroperoxybenzoic acid in a suitable solvent, e.g. aqueous methanol or acetone. Other suitable N-oxidation conditions include using hydrogen peroxide in acetic acid, dimethyldioxirane in acetone, monopero-phthalic acid in acetic acid/methanol, OXONE (trade mark, potassium peroxy-monosulphate) in a suitable solvent such as water, acetone or dichloromethane, and
10 sodium perborate in acetic acid.

Again, a compound of the formula (II) where R is a group of the formula (III) may be resolved to provide a compound of the formula (II) where R is a group of the formula (IV) as described in method (a) above.

15 2) The compounds of the formula (I) can also be prepared by N-oxidation of a compound of the formula:



20 where R^3 is a group of the formula:



25 The N-oxidation can be carried out using a suitable oxidising agent, e.g. 3-chloroperoxybenzoic acid, and a suitable solvent, e.g. methanol or

-9-

acetone. Other suitable N-oxidation conditions include using hydrogen peroxide in acetic acid, dimethyldioxirane in acetone, monoperphthalic acid in acetic acid/methanol, OXONE (trade mark, potassium
5 peroxymonosulphate) in a suitable solvent such as water, acetone or dichloromethane, and sodium perborate in acetic acid.

Where R^3 is a group of the formula (IX), the reaction is followed by separation of the atropisomer of the formula (I) using conventional
10 conditions.

The compounds of the formula (XI) may be prepared by acidic or basic hydrolysis of the compounds of the formula (VIII) using the conditions described in Method (1).

15

3) A compound of the formula (I) can be prepared from its corresponding silica complex by treating a solution of the complex in a suitable solvent, e.g. methanol, with a suitable acid, e.g. a mineral acid (e.g. hydrochloric acid) or acetic acid. This acid treatment degrades the silica complex and
20 liberates a compound of the formula (I).

During the preparation of a compound of the formula (I), the compound must not be treated with silica (e.g. during chromatography) otherwise it will become bound with a stoichiometric quantity thereof to form a different compound, i.e. a silica complex of the required
25 compound.

Accordingly, a compound of the formula (I) is preferably purified by reverse phase gel chromatography.

-10-

All of the above reactions and the preparations of novel starting materials used in the preceding methods are conventional and appropriate reagents and reaction conditions for their performance or preparation as well as procedures for isolating the desired products will be well known to those skilled in the art with reference to literature precedents and the Examples and Preparations hereto.

A pharmaceutically acceptable acid addition or base salt of a compound of the formula (I) may be readily prepared by mixing together solutions of a compound of the formula (I) and the desired acid or base, as appropriate. The salt may precipitate from solution and be collected by filtration or may be recovered by evaporation of the solvent.

The binding affinity of compound of the formula (I) for the glycine site of the NMDA receptor may be measured by testing its ability to displace a selective glycine site radioligand from rat brain membranes as described in Brit. J. Pharm., **104**, 74 (1991). In a variation of this method, thoroughly washed membrane protein is incubated with [³H]-L-689,560 (Mol. Pharmacol., **41**, 923 (1992)) for 90 minutes using tris-acetate buffer (pH 7.4). Displacement of the radioligand, using a range of test compound concentrations, is used to derive IC₅₀ (50% inhibitory concentration) values.

Functional *in vitro* glycine antagonism is demonstrated by the ability of the compounds to inhibit the depolarisations in rat cortical slices induced by NMDA by a similar method to that described in J. Med. Chem., **33**, 789 (1990) and Brit. J. Pharm., **84**, 381 (1985). In a variation of the procedure, the response to a standard concentration of NMDA is measured in the presence of a range of test compound concentrations and the results obtained are used to derive EC₅₀ (50% effective concentration) values.

The binding affinity of the compounds of the invention for the AMPA receptor may be measured by testing their ability to displace the radioligand [³H]-AMPA from rat brain membranes. Membrane homogenate is incubated

-11-

with radioligand (10 nM) in the presence or absence of test compounds at various concentrations at 4°C for 45 minutes. Free and bound radiolabel are separated by rapid filtration and radioactivity is measured by liquid scintillation
5 counting.

The compounds of the formula (I) can be administered to a subject to be treated alone, but will generally be administered in admixture with a pharmaceutically acceptable diluent or carrier selected with regard to the intended route of administration and standard pharmaceutical practice. For
10 example, they can be administered orally, including sublingually, in the form of tablets containing such excipients as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents. They can be injected parenterally, for example, intravenously, intramuscularly or subcutaneously.
15 For parenteral administration, they are best used in the form of a sterile aqueous solution which may contain other substances, for example, enough salts or glucose to make the solution isotonic with blood.

The compounds have potential for absorption through the gastrointestinal tract and thus administration by slow release formulations is
20 also possible.

In general, a therapeutically effective daily oral dose of the compounds of formula (I) is likely to range from 0.1 to 100 mg/kg body weight of the subject to be treated, preferably 1 to 20 mg/kg, and an intravenous or subcutaneous daily dose is likely to range from 0.01-20mg/kg body weight of subject to be
25 treated, preferably 0.1-20 mg/kg. The compounds of the formula (I) may also be administered by intravenous infusion at a dose which is likely to range from 0.01-10 mg/kg/hr.

Tablets or capsules of the compounds may be administered singly or two or more at a time, as appropriate.

-12-

The physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited and such are within the scope of this invention.

Alternatively, the compounds of the formula (I) can be administered by inhalation or in the form of a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. An alternative means of transdermal administration is by use of a skin patch. For example, they can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin. They can also be incorporated, at a concentration of between 1 and 10% by weight, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives as may be required.

It is to be appreciated that reference to treatment includes prophylaxis as well as the alleviation of established symptoms of the disease.

Thus the invention further provides:-

- i) a pharmaceutical composition comprising a compound of the formula (I), or a pharmaceutically acceptable salt or solvate thereof, together with a pharmaceutically acceptable diluent or carrier;
- ii) a compound of the formula (I), or a pharmaceutically acceptable salt, solvate or composition thereof, for use as a medicament;
- iii) the use of a compound of the formula (I), or of a pharmaceutically acceptable salt, solvate or composition thereof, for the manufacture of a medicament for the treatment of a disease by producing an antagonist effect at a NMDA receptor;
- iv) use as in (iii) where the disease is an acute neurodegenerative or a chronic neurological disorder;

-13-

- v) a method of treatment of a mammal to treat a disease by producing an antagonist effect at a NMDA receptor, which comprises treating said mammal with an effective amount of a compound of the formula (I) or
5 with a pharmaceutically acceptable salt, solvate or composition thereof;
- vi) a method as in (v) where the disease is an acute neurodegenerative or a chronic neurological disorder;
- vii) a compound of the formula (II) where R is a group of the formula (III) or (IV); and
- 10 viii) a compound of the formula (VIII) where R³ is a group of the formula (X).

-14-

The following Examples illustrate the preparation of the compounds of the formula (I) and a composition thereof.


- 5 Melting points were determined using a Buchi apparatus in glass capillary tubes and are uncorrected. Low Resolution Mass Spectroscopic (LRMS) data were recorded on a Fisons Trio 1000 Mass Spectrometer (thermospray using ammonium acetate in aqueous methanol as the carrier or atmospheric pressure chemical ionisation (APCI) using 97.5:2.5, by volume, methanol:acetic acid and gaseous nitrogen as the carrier). NMR data were recorded on a Varian Unity 300 or a Varian Inova 400 NMR instrument (300 and 400 MHz, respectively) and were consistent with the assigned structures. Proton NMR shifts are quoted in parts per million downfield from tetramethylsilane. The purity of the compounds was carefully assessed using analytical TLC and proton NMR and the latter technique was used to calculate the amount of solvent present in solvated samples. The term "residue" used in the microanalysis data indicates the residual material remaining following combustion, i.e. the non-flammable material.
- 10
- 15

-15-

EXAMPLE 1

(-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione hydrate

5



Concentrated hydrochloric acid (1ml) was added to a stirred solution of (-)-6,7-dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione, silica complex (See Reference Example 1) (2.3g, 4.64mmol) in methanol (40ml) and the mixture stirred for 2 hours. The solid precipitate was collected by filtration to afford the title compound as a white solid (1.4g, 65%). mp 264-265°C.

Found: C, 44.34; H, 3.21; N, 18.14; residue, 0.00.

15 $C_{17}H_{12}Cl_2N_6O_4 \cdot 1.5 H_2O$ requires C, 44.17; H, 3.21; N, 18.18; residue 0.00%.

1H -NMR (300 MHz, d_6 -DMSO): δ = 3.12 (3H, s), 4.36 (2H, m), 7.18 (1H, d, J = 9.5Hz), 7.36 (1H, dd, $J_1 = J_2 = 9.5$ Hz), 7.42 (1H, s), 8.24 (1H, d, J = 9.5Hz), 8.30 (1H, s), 12.22 (1H, s), 12.24 (1H, s).

20


m/z (thermospray): 435 (MH^+).

$[\alpha]_D^{25} -235^\circ$ (c=0.1, water)

-16-

EXAMPLE 2

(-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione hydrate



5

A solution of 3-chloroperoxybenzoic acid (50-55% w/w in water containing 3-chlorobenzoic acid impurity, 16.1g, 47mmol) in methanol (200ml) was added to a solution of (-)-6,7-dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (see Preparation 1) (13.8g, 31mmol) in methanol (400ml) at room temperature. The reaction mixture was stirred at room temperature for 3.5 days. The reaction mixture was pre-absorbed on reverse phase gel (MCI Gel CHP20P [trade mark], 75-100 μ) and purified by chromatography on reverse phase gel (MCI Gel CHP20P [trade mark], 75-100 μ) by gradient elution using water:methanol (3:1 changing to 2:1, by volume) as the eluent to give, after combination and concentration of the appropriate fractions, a light yellow solid which was recrystallised from methanol to give the title compound (7.6g, 54%) as a colourless solid. mp 265-267°C.

Found: C, 45.01; H, 3.08; N, 18.65. C₁₇H₁₂Cl₂N₆O₄.H₂O requires C, 45.05; H, 3.11; N, 18.54%.

20

-17-

¹H-NMR (300 MHz, d₆-DMSO): Identical spectrum to that obtained for the compound of Example 1.

5 m/z (thermospray): 435 (MH⁺)

[α]_D²⁵ -224° (c=0.1, water)

10

EXAMPLE 3

(-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione

15

(-)-6,7-Dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (see Preparation 1) (412.2g, 0.98 mol) and OXONE (trade mark) (1.44 kg, 2.3 mol) were slurried in water (4.13 L) and the mixture stirred at ambient temperature for 60 hours. Saturated aqueous sodium thiosulphate solution (2.2 L) was added and the slurry stirred for 1 hour before being filtered under reduced pressure. The filter cake was slurried at ambient temperature for 4 hours in 1:1, by volume, isopropyl alcohol: dichloromethane (111 L) and the solid collected by filtration. The filtrate was evaporated under reduced pressure to give the title compound as a colourless solid (366 g).

25

EXAMPLE 4

(-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione, sodium salt, hydrate

30

Sodium hydroxide (9.72ml of a 1 molar aqueous solution, 9.72mmol) was added to a stirred suspension of (-)-6,7-dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione hydrate (see Example 2) (4.406g, 9.72mmol) in water (60ml) and the mixture stirred for

-18-

5 minutes. The resulting solution was filtered and the filtrate freeze-dried to give the title compound (4.5g, 98%) as a pale yellow solid. mp 303°C (decomp.).

5

Found: C, 41.40; H, 3.05; N, 16.99. $C_{17}H_{11}Cl_2N_6NaO_4 \cdot 2H_2O$ requires C, 41.40; H, 3.07; N, 17.04%.

1H -NMR (400 MHz, d_6 -DMSO): δ = 3.08 (3H, s), 4.24 (2H, m), 7.22 (2H, m),
10 7.38 (1H, dd, $J_1 = J_2 = 9.5$ Hz), 8.02 (1H, s), 8.20 (1H, m), 11.66 (1H, s).

$[\alpha]_D^{25} -277^\circ$ (c=0.1, water)

EXAMPLE 5

15 Intravenous formulation of (-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione, sodium salt, hydrate

A formulation suitable for administering a 20mg/ml dose of the active
20 component by intravenous injection was prepared using (-)-6,7-dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione, sodium salt, $2H_2O$ (see Example 4) (22.7mg per unit dose), sodium chloride (9.0mg per unit dose) and water for injections (to 1.0ml).


25 To prepare the formulation, sodium chloride is dissolved in 75% of the total volume of water in a suitable vessel with mixing. (-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione, sodium salt, $2H_2O$ is then added and dissolved by mixing. The solution is then made up to volume with water and filtered through a
30 clarifying 0.2 micron filter. The filtrate is filled into sterile 10ml glass ampoules under aseptic conditions using a terminal clarifying filter and the ampoules sealed.

-19-

Part (i) of the following Reference Example 1 is a repeat preparation of the compound of Example 114 of International Patent Application Publication no. WO 97/32873. In Part (ii), the product obtained was recrystallised from aqueous acetone.

REFERENCE EXAMPLE 1

(-)-6,7-Dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione, silica complex



10

(i) A solution of 3-chloroperoxybenzoic acid (0.85g, 4.93 mmol) in acetone (20ml) was added in one portion to a suspension of (-)-6,7-dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (see Preparation 1) (1.0g, 2.24 mmol) in acetone (40ml) which caused all the solid to dissolve. The reaction was stirred at room temperature for 40 minutes after which time a white solid began to form. The reaction mixture was allowed to stir at room temperature for 3 days. The white solid was collected by filtration (this solid contained less than 90% w/w of the N-oxide product)¹ and subjected to flash chromatography on silica gel using dichloromethane:methanol:glacial acetic acid (90:10:1, by volume) as the eluant to give, after combination and concentration of the appropriate fractions, the title compound as a white solid, (0.16g). m.p. >310 °C.

15

20

25

¹H-NMR (300 MHz, d₆-DMSO): δ = 1.90 (s, acetic acid 0.3 eq), 3.10 (3H, s), 4.32 (2H, m), 7.22 (1H, m), 7.40 (2H, m), 8.10 (1H, m), 8.22 (1H, m).

-20-

m/z (thermospray): 435.

5 $[\alpha]_D^{25} -235^\circ$ (c=0.1, ethanol)*

(*It should be noted that a clerical error occurred when stating the $[\alpha]_D^{25}$ value in Example 114 of International Patent Application Publication no. WO 97/32873. The stated "c = 1.0" value is incorrect and this should
10 have read "c = 0.1").

(ii) Recrystallisation of this solid from aqueous acetone gave the title compound as a white solid. mp >310°C.

15 Found: C, 41.2; H, 3.1; N, 17.0; residue, 8.25.

$C_{17}H_{12}Cl_2N_6O_4 \cdot 0.5 H_2SiO_3 \cdot 1.2 H_2O$ requires: C, 41.18; H, 3.13; N, 16.95; residue 7.87%.

1H -NMR (300 MHz, d_6 -DMSO): δ = 3.05 (3H, s), 4.37 (2H, m), 7.16 (1H, d, J = 9.5Hz), 7.32 (1H, s), 7.32 (1H, m), 7.98 (1H, s), 8.18 (1H, d, J =
20 9.5Hz).

$[\alpha]_D^{25} -199^\circ$ (c=0.1, methanol)

25

Footnote

1. The method of Reference Example 1 (i) was repeated exactly and the precipitated white solid was collected by filtration (0.507 g).

This was found to contain 57.7% w/w (-)-6,7-dichloro-5-[3-methoxymethyl-5-(1-oxidopyridin-3-yl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione when analysed by high pressure liquid chromatography (HPLC) using a 15 cm x 0.46 cm i.d. Magellen (trade mark) C18 column and a gradient elution employing the following combinations of solvent A (acetonitrile)
30

-21-

and solvent B (8.3 mM phosphate buffer adjusted to pH3.7 using phosphoric acid):

5

Time (min.)	% (by volume) A	% (by volume) B	Duration (min.)
	2	98	(Initial)
0	98	2	30
35	2	98	1
45			(Finish)

at a flow rate of 1 ml/min. and at ambient temperature.

The components of the eluted mixture were detected at a wavelength of 220 nm and samples of the compounds of Examples 4 and Preparation 1 and
10 of 3-chloroperoxybenzoic acid were used as reference standards.


-22-

The following Preparations describe the manufacture of certain intermediates used in the preceding Examples and Reference Example.

5

PREPARATION 1

(±)-, (-)- and (+)-6,7-Dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione



10

- (a) Methoxyacetylchloride (27.3ml, 32.4g, 0.30mol) was added to a stirred mixture of 5-amino-6,7-dichloro-2,3-dimethoxyquinoxaline (Preparation 2) (73.8g, 0.27mol) and pyridine (26.4ml, 25.8g, 0.33mol) in dichloromethane (1.2 litres) at room temperature under nitrogen. After 18 hours stirring at room temperature, the mixture was washed with 2M aqueous hydrochloric acid solution followed by brine, then dried (MgSO₄) and concentrated under reduced pressure. The residue was triturated

15

-23-

with methanol and filtered to give 6,7-dichloro-2,3-dimethoxy-5-methoxyacetamidoquinoxaline (82.0g, 88%) as an off-white solid. mp 171-173°C.

5

Found: C, 44.97; H, 3.75; N, 12.03. $C_{13}H_{13}Cl_2N_3O_4$ requires C, 45.11; H, 3.79; N, 12.14%.

10 (b) 2,4-Bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulphide (Lawesson's reagent) (19.5g, 48.2mmol) was added to a solution of 6,7-dichloro-2,3-dimethoxy-5-methoxyacetamidoquinoxaline (27g, 78mmol) in tetrahydrofuran (480ml) and the mixture was stirred for 18 hours at room temperature, then evaporated under reduced pressure. The residue was purified by flash chromatography on silica gel by gradient
15 elution using hexane:dichloromethane (1:1 changing to 1:4, by volume) as the eluent to give 6,7-dichloro-2,3-dimethoxy-5-methoxythioacetamidoquinoxaline (29.1g, >100%) as a white solid, mp 198-200°C, containing a minor impurity.

20 Found: C, 43.06; H, 3.65; N, 11.59. $C_{13}H_{13}Cl_2N_3O_3S$ requires C, 43.11; H, 3.62; N, 11.60%.

(c) A mixture of 6,7-dichloro-2,3-dimethoxy-5-methoxythioacetamido-
quinoxaline (25.3g, 69.9mmol), nicotinic acid hydrazide (19.3g,
25 140.8mmol), mercury(II) oxide (15.1g, 69.7mmol) and 1,4-dioxane (600ml) was heated under reflux for 18 hours. After cooling, the mixture was filtered through ARBOCEL (trade mark) filter aid and the residue washed with dichloromethane. The filtrate was concentrated under

-24-

reduced pressure to afford a light brown solid which was partitioned between ethyl acetate and 2M aqueous hydrochloric acid solution. The layers were separated and the aqueous layer was extracted with
5 dichloromethane (2x500ml, 4x100ml). The combined dichloromethane extracts were dried (MgSO_4) and concentrated under reduced pressure. The residue was crystallised from ethyl acetate/methanol to give ($\pm$)-6,7-dichloro-2,3-dimethoxy-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]quinoxaline (11.6g, 37%) as a pale yellow solid. mp 189-
10 191°C.

Found: C, 50.10; H, 3.57; N, 18.53. $\text{C}_{19}\text{H}_{16}\text{Cl}_2\text{N}_6\text{O}_3 \cdot 0.5\text{H}_2\text{O}$ requires: C, 50.01; H, 3.76; N, 18.42%.

15 (d) A mixture of ($\pm$)-6,7-dichloro-2,3-dimethoxy-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]quinoxaline (3.0g, 6.7mmol), 2M aqueous hydrochloric acid solution (10ml) and 1,4-dioxane (50ml) was heated under reflux for 9 hours, cooled, and concentrated under reduced pressure. The residue was dissolved in 1M aqueous sodium hydroxide
20 solution and acidified to pH 4.5 with concentrated hydrochloric acid to afford a thick white precipitate. This was collected by filtration and washed with water to give ($\pm$)-6,7-dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (2.0g, 68%) as an off-white solid. mp 230-232°C.

25

Found: C, 46.23; H, 2.93; N, 19.00. $\text{C}_{17}\text{H}_{12}\text{Cl}_2\text{N}_6\text{O}_3 \cdot 1.25\text{H}_2\text{O}$ requires: C, 46.22; H, 3.31; N, 19.02%.

30 (e) (i) (-)-N-Methylephedrine (0.88g, 4.9mmol) and then methanol (66ml) were added to a stirred suspension of ($\pm$)-6,7-dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (1.9g,

-25-

4.3mmol) in ethyl acetate (400ml) at room temperature. The mixture was heated to its boiling point. The mixture was filtered, the filtrate concentrated to three quarters of its volume and then cooled to room temperature. The solid obtained was collected by filtration and washed with ethyl acetate. The solid was crystallised from ethyl acetate/methanol to give a single diastereoisomer of the quinoxalinedione starting material as the (-)-N-methylephedrine salt (1.28g, 43%). mp 162-164°C.

Found: C, 55.74; H, 5.38; N, 14.38. $C_{28}H_{29}Cl_2N_7O_4 \cdot CH_3CO_2C_2H_5$ requires: C, 55.98; H, 5.43; N, 14.28%.

$[\alpha]_D^{25} -135^\circ$ (c=0.1, ethanol).

(ii) A suspension of the (-)-N-methylephedrine salt (1.2g, 1.7mmol) from part (e)(i) in water (13ml) at room temperature was acidified to pH 5 with concentrated hydrochloric acid and the suspension was stirred for 1 hour. The solid obtained was collected by filtration, washed with water and crystallised from water/ethanol to give (-)-6,7-dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (0.48g, 62%) as a white solid. mp 220-222°C.

Found: C, 45.49; H, 3.21; N, 18.72. $C_{17}H_{12}Cl_2N_6O_3 \cdot 1.5H_2O$ requires C, 45.76; H, 3.39; N, 18.83%.

$[\alpha]_D^{25} -214^\circ$ (c=0.1, ethanol).

(iii) The combined filtrates from part (e)(i) were concentrated to dryness, the residue dissolved in water (20ml), acidified to pH 3 with concentrated

-26-

hydrochloric acid and the solid obtained was collected by filtration, washed with water and dried. (+)-N-Methylephedrine (0.37g, 2.06mmol) and then methanol (28ml) were added to a stirred suspension of this solid (0.80g, 1.87mmol) in ethyl acetate (170ml) at room temperature and the mixture was heated to its boiling point. The mixture was filtered, concentrated to three quarters of its volume and then cooled to room temperature. The solid obtained was collected by filtration and washed with ethyl acetate. The solid was crystallised from ethyl acetate/methanol to give a single diastereoisomer of the quinoxalinedione starting material as the (+)-N-methylephedrine salt (0.93g, 32%) as a white solid. mp 165-167°C.

Found: C, 55.88; H, 5.40; N, 14.31. $C_{28}H_{29}Cl_2N_7O_4 \cdot 0.8 CH_3CO_2C_2H_5$ requires: C, 56.01; H, 5.33; N, 14.66%.


$[\alpha]_D^{25} +127^\circ$ (c=0.1, ethanol).

- (iv) A suspension of the (+)-N-methylephedrine salt (0.90g, 1.35mmol) from part (e) (iii) in water (10ml) at room temperature was acidified to pH 5 with concentrated hydrochloric acid and the suspension was stirred for 1 hour. The solid was collected by filtration and washed with water to give (+)-6,7-dichloro-5-[3-methoxymethyl-5-(3-pyridyl)-4H-1,2,4-triazol-4-yl]-2,3(1H,4H)-quinoxalinedione (0.41g, 69%) as a white solid. mp 222-224°C.

Found: C, 46.44; H, 3.18; N, 19.01. $C_{17}H_{12}Cl_2N_6O_3 \cdot 1.25H_2O$ requires C, 46.22; H, 3.31; N, 19.02%.

$[\alpha]_D^{25} +212^\circ$ (c=0.1, ethanol).

-27-

PREPARATION 25-Amino-6,7-dichloro-2,3-dimethoxyquinoxaline

5

10

15

- (a) A mixture of 6,7-dichloro-5-nitro-2,3(1H,4H)-quinoxalinedione (Example 1 of WO-A-94/00124, 84 g, 0.34 mol), thionyl chloride (840ml) and dimethylformamide (0.5ml) was heated under reflux for 3 hours, cooled and concentrated under reduced pressure. Ethyl acetate (300ml) was added and removed by evaporation under reduced pressure and this procedure was then repeated with petroleum ether (bp 100-120°C). The solid residue was recrystallised from petroleum ether (bp 100-120°C) to give 2,3,6,7-tetrachloro-5-nitroquinoxaline (78g, 73%) as a light yellow solid.

¹H-NMR (300 MHz, CDCl₃): δ = 8.6 (1H, s).

20

- (b) Tin(II) chloride dihydrate (346.3g, 1.54mol) was added to a solution of 2,3,6,7-tetrachloro-5-nitroquinoxaline (96.2g, 0.31mol) in ethyl acetate (1.8 litres). The mixture was heated under reflux for 4 hours, cooled and poured cautiously into an excess of aqueous saturated sodium bicarbonate solution. The mixture was filtered through CELITE (trade

-28-

mark) filter aid washing well with ethyl acetate. The filter cake was macerated with further ethyl acetate and the solid material filtered off. The combined ethyl acetate phases were dried (MgSO_4) and concentrated under reduced pressure to give 5-amino-2,3,6,7-tetrachloroquinoxaline (73.4g, 84%) as a yellow solid.

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ = 5.45 (2H, br, s), 7.47 (1H, s).

m/z (thermospray): 385 (MH^+).

(In an alternative preparation, this reduction step was performed using iron filings in aqueous acetic acid).

(c) A solution of sodium methoxide (25% w/w solution in methanol, 274ml, 1.28mol) was added to a suspension of 5-amino-2,3,6,7-tetrachloroquinoxaline (72.4g, 0.256mol) in dry methanol (1 litre) and the resulting mixture was heated under reflux for 30 minutes. The mixture was cooled, concentrated under reduced pressure, and the residue partitioned between water and ethyl acetate (total of 8 litres). The organic extracts were dried (MgSO_4) and concentrated under reduced pressure. The crude product was triturated with methanol then dissolved in dichloromethane (2 litres) and filtered. The filtrate was concentrated under reduced pressure to give the title compound as a yellow solid (55.0g, 79%).

$^1\text{H-NMR}$ (300 MHz, CDCl_3): δ = 4.13 (3H, s), 4.14 (3H, s), 5.07 (2H, br s), 7.26 (1H, s).

m/z (thermospray): 274 (MH^+).

(In an alternative preparation, toluene was used as a co-solvent with methanol).

-29-

Solubility data

5 The compounds of Examples 1 and 2 and Reference Example 1 were tested for their solubility in water and methanol at ambient temperature.

The results are shown in the Table below.

Reference	Solubility in water at pH 7.3 (mg/ml)	Solubility in methanol (mg/ml)
Example 1 and 2	>20 mg/ml	<1 mg/ml
Reference Example 1	<1 mg/ml	ca. 15 mg/ml

10

Lipophilicity data

The lipophilicities of the compounds of Example 2 and Reference Example 1 were tested by the octanol/water partition method.

Reference	log D
Example 2	-1.7
Reference Example 1	-0.6

-30-

Pharmacological data

The binding affinities for the glycine site of the NMDA receptor and the functional in vitro glycine antagonism of the compounds of Example 2 and
5 Reference Example 1 were measured by the methods described on page 10.

The results were as follows:

Binding affinity	
Example 2	IC ₅₀ = 2.4 nm
Reference Example 1	IC ₅₀ = 3.8 nm


Functional <u>in vitro</u> glycine antagonism	
Example 2	IC ₅₀ = 140 nm
Reference Example 1	IC ₅₀ = 190 nm

69387-272

-31-

CLAIMS:

1. A compound of the formula:



(I)

or a pharmaceutically acceptable salt or solvate thereof.

2. A compound as claimed in claim 1 that is at least of 90% w/w purity.
3. A compound as claimed in claim 2 that is at least of 95% w/w purity.
4. A compound as claimed in claim 3 that is at least of 98% w/w purity.
5. A compound as claimed in any one of claims 1 to 4 wherein the pharmaceutically acceptable salt is a sodium salt.
6. A pharmaceutical composition comprising a compound of the formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 5, together with a pharmaceutically acceptable diluent or carrier.
7. A compound of the formula (I), or a pharmaceutically acceptable salt, solvate or composition thereof, as claimed in any one of claims 1 to 5 and 6, respectively, for use as a medicament.
8. The use of a compound of the formula (I), or of a pharmaceutically acceptable salt, solvate or composition thereof, as claimed in any one of claims

69387-272

-32-


1 to 5 and 6, respectively, for the manufacture of a medicament for the treatment of a disease by producing an antagonist effect at an NMDA receptor.

9. Use as claimed in claim 8 where the disease is an acute neurodegenerative or a chronic neurological disorder.


10. Use of a compound of the formula (I) or a pharmaceutically acceptable salt, solvate or composition thereof, as claimed in any one of claims 1 to 5 and 6, respectively, for treatment in a mammal of a disease by producing an antagonist effect at an NMDA receptor.

11. Use as claimed in claim 10 where the disease is an acute neurodegenerative or a chronic neurological disorder.

12. A compound of the formula:



wherein R is group of the formula:



and R¹ and R², either when taken alone or together, represent a group or groups that can be hydrolytically cleaved under acidic or basic conditions to provide the corresponding quinoxalinedione.

25


69387-272

-33-

13. A compound of the formula (II) as claimed in claim 12 wherein R^1 and R^2 are either each independently selected from C_1 - C_4 alkyl and benzyl, optionally ring-substituted by from 1 to 3 substituents each independently selected from C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halo, nitro and trifluoromethyl, or, when taken together, represent C_1 - C_6 alkylene, CH(phenyl), CH(4-methoxyphenyl) or CH(3,4-dimethoxyphenyl).


14. A compound as claimed in claim 13 wherein R^1 and R^2 are each independently selected from methyl, ethyl and benzyl, optionally ring-substituted by from 1 to 3 substituents each independently selected from C_1 - C_4 alkyl, C_1 - C_4 alkoxy, halo, nitro and trifluoromethyl.

15. A compound of the formula:



(VIII)

wherein R^3 is a group of the formula:



(X)


and R^1 and R^2 are as defined in claim 12, 13 or 14.

16. A process for the preparation of a compound of the formula (I), or a pharmaceutically acceptable salt or

69387-272

-33a-

solvate thereof, as claimed in claim 1 comprising acidic or basic hydrolysis of a compound of the formula:




5

wherein R is group of the formula:

69387-272

-34-




5 and R^1 and R^2 , either when taken alone or together, represent a group or groups that can be hydrolytically cleaved under acidic or basic conditions to provide the corresponding quinoxalinedione, said process being followed by:


10 (i) when a compound of the formula (II) wherein R is a group of the formula (III) is used, separation of the atropisomer of the formula (I); and/or

(ii) , optionally, conversion of a compound of the formula (I) to a pharmaceutically acceptable salt or solvate thereof.

15 17. A process for the preparation of a compound of the formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1 comprising N-oxidation of a compound of the formula:



where R^3 is a group of the formula:



20

69387-272

-35-

followed by work-up of the reaction under silica-free conditions, said process being followed by:


5 (i) when a compound of the formula (XI) wherein R is a group of the formula (IX) is used, separation of the atropisomer of the formula (I); and/or

(ii) , optionally, conversion of a compound of the formula (I) to a pharmaceutically acceptable salt or solvate thereof.

10 18. A process as claimed in claim 17 wherein the N-oxidation is carried out using OXONE (trade mark) in a reaction-inert solvent.

19. A process as claimed in claim 18 wherein the reaction-inert solvent is water.

15 20. A process for the preparation of a compound of the formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in claim 1 comprising acidic treatment of a silica complex of a compound of the formula (I), said process being optionally followed by conversion of a compound of the formula (I) to a pharmaceutically acceptable salt or solvate thereof.



(I)