



- (51) **International Patent Classification:**
C12N 5/075 (2010.01) C12N 15/877 (2010.01)
- (21) **International Application Number:**
PCT/US2014/067588
- (22) **International Filing Date:**
26 November 2014 (26.11.2014)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/963,246 27 November 2013 (27.11.2013) US
61/997,176 23 May 2014 (23.05.2014) US
- (71) **Applicant: THE CURATORS OF THE UNIVERSITY OF MISSOURI** [US/US]; 316 University Hall, Columbia, MO 65211 (US).
- (72) **Inventors: LEE, Kiho**; 316 University Hall, Columbia, MO 65211 (US). **PRATHER, Randall, S.**; 316 University Hall, Columbia, MO 65211 (US).
- (74) **Agent: HANSON, Robert, E.**; DENTONS US LLP, 233 South Wacker Drive, Suite 7800, Chicago, IL 60606-6306 (US).

- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** ARTIFICIAL OOCYTE ACTIVATION

(57) **Abstract:** The present invention provides novel methods for improving the efficiency of artificial activation of unfertilized mammalian oocytes by reducing the intracellular concentration of Zn^{2+} in the oocyte. The methods of the invention may additionally comprise a preceding step of increasing the intracellular concentration of Ca^{2+} in the oocyte prior to reduction of the intracellular Zn^{2+} concentration. The invention further provides unfertilized oocytes activated by the disclosed methods and viable mammalian animals produced from unfertilized oocytes activated by the disclosed methods.



WO 2015/081182 A1

ARTIFICIAL OOCYTE ACTIVATION

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of United States Provisional Application Nos. 61/963,246, filed November 27, 2013, and 61/997,176, filed May 23, 2014, herein incorporated
5 by reference in their entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

This invention was made with government support under grant number U42 RR018877 awarded by the National Swine Resource and Research Center. The government has certain rights in the invention.

FIELD OF THE INVENTION

The present invention relates to the fields of animal husbandry and biomedicine, more specifically, to a method for improving the cloning of mammals.

BACKGROUND OF THE INVENTION

Somatic cell nuclear transfer (SCNT), *i.e.*, cloning, can be used to generate genetically
15 engineered animals, preserve endangered species, and produce animals with a select genetic background. However, current SCNT technology is inefficient as only one percent of generated embryos are able to reach term development. One of the factors that contributes to the poor efficiency is the effectiveness of the artificial oocyte activation process, an essential procedure in SCNT. There is a great need for better approaches for artificial oocyte activation.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides a method of activating an unfertilized livestock or domestic mammalian oocyte comprising decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety. In one embodiment, the Zn^{2+} binding moiety comprises a concentration of between approximately 10 μM to approximately
25 500 μM , for instance, between approximately 100 μM to approximately 250 μM , such as approximately 200 μM . In another embodiment, the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 1 minute to approximately 5 hours, for

instance, between approximately 30 minutes to approximately 2.5 hours, such as approximately 30 minutes.

In further embodiments, the present method of activating an unfertilized livestock or domestic mammalian oocyte additionally comprises the step of increasing intracellular Ca^{2+} concentration of the oocyte prior to decreasing the intracellular Zn^{2+} concentration of the oocyte. In one embodiment, the intracellular Ca^{2+} concentration of the oocyte is not increased in an amount sufficient to induce oocyte activation, for instance, the intracellular Ca^{2+} concentration of the oocyte may be increased approximately ten times less than the amount sufficient to induce oocyte activation. In certain embodiments, in methods where the intracellular Ca^{2+} concentration of the oocyte is not increased in an amount sufficient to induce oocyte activation, the Zn^{2+} binding moiety may comprise a concentration of between approximately 1 μM to approximately 500 μM , for instance between approximately 100 μM to approximately 200 μM , such as approximately 200 μM . In other embodiments, in methods where the intracellular Ca^{2+} concentration of the oocyte is not increased in an amount sufficient to induce oocyte activation, the Zn^{2+} binding moiety may contact the oocyte for a period of time comprising between approximately 1 minutes to approximately 12 hours, for instance, between approximately 10 minutes to approximately 2 hours, such as approximately 30 minutes.

In still further embodiments, the intracellular Ca^{2+} concentration of the oocyte may be increased in an amount sufficient to induce oocyte activation. In certain embodiments, in methods where the intracellular Ca^{2+} concentration of the oocyte is increased in an amount sufficient to induce oocyte activation, the Zn^{2+} binding moiety may comprise a concentration of between approximately 0.1 μM to approximately 100 μM , for instance, between approximately 1 μM to approximately 10 μM , such as approximately 5 μM or approximately 10 μM . In other embodiments, in methods where the intracellular Ca^{2+} concentration of the oocyte is increased in an amount sufficient to induce oocyte activation, the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 1 minutes to approximately 5 hours, for instance, between approximately 10 minutes to approximately 30 minutes, such as approximately 30 minutes.

In certain embodiments, in methods of the present invention the intracellular Ca^{2+} concentration of the oocyte is increased by chemical or physical means or a combination thereof that induce calcium entry into the oocyte or release of internal calcium stores. In particular embodiments, the increase in the intracellular Ca^{2+} concentration of the oocyte by chemical means comprises treatment with a calcium containing medium, a calcium salt containing medium, thimerosal containing medium, ethanol containing medium, inositol trisphosphate containing medium, a calcium ionophore containing medium, a sodium-free medium or a combination thereof. For instance, In some embodiments, the chemical treatment comprises treatment with a medium comprising a Ca^{2+} concentration of between approximately 0.1 mM and approximately 250 mM, for instance, between approximately 0.1 mM and 1 mM. In other embodiments, in methods of the present invention, the increase in the intracellular Ca^{2+} concentration of the oocyte by physical means comprises application of an electrical stimulus.

In yet a further embodiment of the present invention, the Zn^{2+} binding moiety comprises TPEN (N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine). In other embodiments, the unfertilized livestock or domestic mammalian oocyte is a nuclear transfer oocyte. In additional embodiments, the unfertilized livestock or domestic mammalian oocyte is selected from the group consisting of a porcine oocyte, bovine oocyte, ovine oocyte, goat oocyte, horse oocyte, canine oocyte and feline oocyte.

In another aspect, provided herein is a parthenogenetic oocyte activated by the methods of the present invention. In still another aspect, provided herein is a nuclear transfer oocyte activated by the methods of the present invention.

In yet another aspect, the present invention provides a viable livestock or domestic mammalian animal produced from an unfertilized livestock or domestic mammalian oocyte activated by the methods of the invention, wherein the animal comprises the genomic DNA from a donor cell other than the oocyte and the mitochondrial DNA from the oocyte. In certain embodiments, the animal comprises epigenetic features different from the epigenetic features of the donor cell. In particular embodiments, the epigenetic features different from the epigenetic features of the donor cell are DNA methylation sites or histone modifications. In further

embodiments, the animal is selected from the group consisting of a porcine animal, bovine animal, ovine animal, goat, horse, dog and cat.

In a further aspect, the present invention provides a method of activating an unfertilized mammalian oocyte comprising increasing intracellular Ca^{2+} concentration of the oocyte by
5 contacting the oocyte with a medium comprising a calcium concentration of approximately 0.1 mM and decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety at a concentration of approximately 200 μM for a period of time comprising approximately 30 minutes.

In still a further aspect, the present invention provides a method of activating an
10 unfertilized mammalian oocyte comprising increasing intracellular Ca^{2+} concentration of the oocyte by contacting the oocyte with a medium comprising a calcium concentration of approximately 1.0 mM and decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety at a concentration of approximately 5 μM for a period of time comprising approximately 30 minutes.

In yet a further aspect, the present invention provides a method of activating an
15 unfertilized mammalian oocyte comprising increasing intracellular Ca^{2+} concentration of the oocyte by contacting the oocyte with a medium comprising a calcium concentration of approximately 1.0 mM and decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety at a concentration of approximately 10 μM for
20 a period of time comprising approximately 30 minutes.

In an additional aspect, the invention provides a method for activating an unfertilized mammalian oocyte, comprising the steps of (a) increasing Ca^{2+} concentration of said oocyte, and (b) subsequently decreasing Zn^{2+} concentration of said oocyte by treatment of Zn^{2+} removing agent. In one embodiment, the Zn^{2+} removing agent is N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine (TPEN). In another embodiment, the unfertilized mammalian
25 oocyte is a porcine somatic cell nuclear transfer embryo. In yet another embodiment, the TPEN concentration ranges between 0.1 μM to 100 μM . In still another embodiment, the TPEN concentration is 5 μM to 10 μM .

In another aspect, the invention provides a method for activating an unfertilized mammalian oocyte, comprising the steps of (a) providing a minor Ca^{2+} concentration increase of said oocyte, and (b) subsequently decreasing Zn^{2+} concentration of said oocyte by treatment of Zn^{2+} removing agent. In one embodiment, the Zn^{2+} removing agent is N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine (TPEN). In another embodiment, the unfertilized mammalian oocyte is a porcine somatic cell nuclear transfer embryo. In yet another embodiment, the TPEN concentration is higher than 10 μM . In still another embodiment, the TPEN concentration is 200 μM .

BRIEF DESCRIPTION OF THE DRAWINGS

For a more complete understanding of the features and advantages of the present invention, reference is now made to the detailed description of the invention along with the accompanying figures:

FIG. 1A - Graphical representation demonstrating that N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine (TPEN) does not induce a Ca^{2+} increase in Metaphase II (MII) oocytes. The graph demonstrates the Ca^{2+} increase after treating MII stage oocytes with thimerosal, an oocyte activating compound.

FIG. 1B - Graphical representation demonstrating that TPEN does not induce a Ca^{2+} increase in MII oocytes. The graph demonstrates the lack of Ca^{2+} increase from TPEN. No Ca^{2+} increase was detected when MII stage oocytes were exposed to various concentrations of TPEN. The arrow indicates when the MII oocytes were exposed to TPEN.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel methods for improving the efficiency of artificial oocyte activation of unfertilized mammalian oocytes. In particular, the invention provides methods for oocyte activation through reduction of intracellular Zn^{2+} . The methods of oocyte activation of the present invention by reduction of intracellular Zn^{2+} in the unfertilized oocyte may be either independent or dependent of conventional oocyte activation methods. For instance, in one embodiment, the reduction of intracellular Zn^{2+} is sufficient to activate an unfertilized oocyte without being coupled with a prior increase in intracellular Ca^{2+} . In another

embodiment, the reduction of intracellular Zn^{2+} is preceded by a Ca^{2+} increase in the unfertilized oocyte.

In certain embodiments the Ca^{2+} increase in the unfertilized oocyte may be a minor increase, for instance less than the amount that is sufficient to activate an unfertilized oocyte alone. In another embodiment, the Ca^{2+} increase in the unfertilized oocyte may be an amount
5 equal to or greater than the amount that is sufficient to activate an unfertilized oocyte alone.

Methods for artificial oocyte activation by increasing Ca^{2+} release in the oocyte have been studied with limited success, especially in parthenogenetic and somatic cell nuclear transfer (SCNT) porcine embryo development. Conventional methods to increase intracellular Ca^{2+} ,
10 including electroporation or thimerosal treatment followed by treatment with dithiothreitol (THI/DTT), are commonly employed in activating SCNT embryos, however, the efficiency and subsequent development is relatively low. The present invention thus provides methods to increase the developmental potential of artificially activated oocytes and increase the efficiency of oocyte activation and subsequent development during SCNT through a treatment with a Zn^{2+}
15 binding moiety and, in certain embodiments, combined with a Ca^{2+} stimulus. The present invention thus overcomes the difficulties in the art regarding artificial oocyte activation by conventional intracellular oocyte Ca^{2+} increase through the use of Zn^{2+} binding moieties, such as Zn^{2+} chelators, for the removal of Zn^{2+} .

In nature, penetration of the sperm head into the cytoplasm of the oocyte causes repetitive
20 calcium increases referred as Ca^{2+} oscillations. These oscillations result from the release of phospholipase C (PLC)-zeta from the sperm head. The Ca^{2+} oscillations activate Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII) and the CaMKII phosphorylates early mitotic inhibitor 2 (EMI2, also known as FBXO43) thus relieving anaphase-promoting complex/cyclosome (APC/C) from FBXO43-mediated inhibition. These events lead to the
25 degradation of cyclin B, a subunit of M-phase promoting factor (MPF).

Most conventional artificial activation methods induce an intracellular Ca^{2+} increase to mimic this Ca^{2+} signaling. However, because it is technically easier, most artificial activation methods produce a single Ca^{2+} increase, which is suboptimal for activating oocytes, as compared to the repetitive Ca^{2+} increases observed after sperm-induced oocyte activation.

Zn^{2+} is a key component which can maintain oocytes arrested at the metaphase II (MII) stage and is necessary for activating MPF, as Zn^{2+} regulates the activity of the cyclin-dependent kinase, CDC25. In addition, EMI2 (FBXO43), a zinc-binding protein, is required to maintain high MPF activity during the MII arrest, and the increase in total intracellular Zn^{2+} during oocyte maturation directly controls FBXO43 activity. Zn^{2+} is released from oocytes after fertilization indicating that removal of Zn^{2+} is a natural part of oocyte activation. By utilizing a Zn^{2+} specific binding moiety, such as the Zn^{2+} chelator TPEN, the natural process of oocyte activation can be stimulated thus increasing the developmental potential of activated oocytes.

Artificial oocyte activation is an essential process during SCNT. Unfortunately, current artificial activation methods used during SCNT are suboptimal as they cannot completely emulate the sperm-induced natural signaling pathway. Different strategies have been used to increase the efficiency of artificial oocyte activation by lowering the level of MPF during artificial oocyte activation. However, due to potential toxicity, they are not widely used in cloned animal production, such as pig production.

For the methods of the present invention, intracellular Ca^{2+} concentration may be increased by any conventional method known in the art. Such methods are known in the art would be fully appreciated and understood by one of skill in the art. One exemplary conventional artificial oocyte activation method comprises treating an unfertilized (or nuclear transferred) oocyte with an oocyte-modifying agent, such as thimerosal, followed by a reducing agent, such as dithiothreitol (DTT) (U.S. Patent No. 6,211,429). Additional methods include, without limitation, various chemical, physical or mechanical stimuli. For instance, the use of a medium comprising any calcium salt, such as $CaCl_2$, or a compound that dissociates calcium; calcium ionophores, such as ionomycin or A23187; use of a sodium-deficient or a sodium-free medium; by electric shock in the presence of Ca^{2+} ; ethanol treatment; or caged chelators which can release calcium inside the cell in response to specific wavelengths or combinations of one or more of these stimuli. Reference is made to U.S. Patent No. 6,211,429, which is incorporated herein by reference for a general discussion on increasing intracellular calcium in oocytes and the role of such an increase in artificial oocyte activation.

As used to herein "Zn²⁺ binding moieties" or "Zn²⁺ removing agents" refer to any molecule or composition that is capable of binding, removing or sequestering Zn²⁺. For instance, in certain embodiments, Zn²⁺ binding moieties of the present invention may include Zn²⁺ chelators. Such Zn²⁺ chelators include but are not limited to N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine (TPEN), known to have a high specificity toward Zn²⁺; clioquinol; and diethylenetriaminepentaacetic acid (DTPA). In particular embodiments, the concentration of Zn²⁺ binding moieties for use in the present invention can comprise approximately 0.01 μ M to approximately 1 mM. For instance, from approximately 0.1 μ M to approximately 500 μ M, approximately 1 μ M to approximately 250 μ M, approximately 10 μ M to approximately 100 μ M, approximately 200 μ M to approximately 500 μ M, approximately 200 μ M to approximately 250 μ M, approximately 100 μ M to approximately 500 μ M, approximately 100 μ M to approximately 250 μ M, approximately 100 μ M to approximately 200 μ M, approximately 100 μ M to approximately 150 μ M, approximately 50 μ M to approximately 500 μ M, approximately 50 μ M to approximately 250 μ M, approximately 50 μ M to approximately 200 μ M, approximately 200 μ M, approximately 50 μ M to approximately 150 μ M, approximately 50 μ M to approximately 100 μ M, approximately 10 μ M to approximately 500 μ M, approximately 10 μ M to approximately 250 μ M, approximately 10 μ M to approximately 200 μ M, approximately 10 μ M to approximately 150 μ M, approximately 10 μ M to approximately 50 μ M, approximately 10 μ M to approximately 20 μ M, approximately 5 μ M to approximately 200 μ M, approximately 5 μ M to approximately 150 μ M, approximately 5 μ M to approximately 100 μ M, approximately 5 μ M to approximately 50 μ M, approximately 5 μ M to approximately 20 μ M, approximately 5 μ M to approximately 10 μ M, approximately 1 μ M to approximately 500 μ M, approximately 1 μ M to approximately 100 μ M, approximately 1 μ M to approximately 50 μ M, approximately 1 μ M to approximately 20 μ M, approximately 1 μ M to approximately 10 μ M, approximately 1 μ M to approximately 5 μ M, approximately 0.1 μ M to approximately 100 μ M, approximately 0.1 μ M to approximately 50 μ M, approximately 0.1 μ M to approximately 20 μ M, approximately 0.1 μ M to approximately 10 μ M, approximately 0.1 μ M to approximately 5 μ M, approximately 0.1 μ M to approximately 1 μ M or a range equivalent thereto.

The concentration of the Zn²⁺ binding moieties may therefore comprise in particular embodiments approximately 0.01 μ M, approximately 0.05 μ M, approximately 0.1 μ M,

approximately 0.2 μM , approximately 0.5 μM , approximately 0.75 μM , approximately 1 μM ,
approximately 2 μM , approximately 3 μM , approximately 4 μM , approximately 5 μM ,
approximately 6 μM , approximately 7 μM , approximately 8 μM , approximately 9 μM ,
approximately 10 μM , approximately 11 μM , approximately 12 μM , approximately 13 μM ,
5 approximately 14 μM , approximately 15 μM , approximately 16 μM , approximately 17 μM ,
approximately 18 μM , approximately 19 μM , approximately 20 μM , approximately 25 μM ,
approximately 30 μM , approximately 35 μM , approximately 40 μM , approximately 45 μM ,
approximately 50 μM , approximately 55 μM , approximately 60 μM , approximately 65 μM ,
approximately 70 μM , approximately 75 μM , approximately 80 μM , approximately 85 μM ,
10 approximately 90 μM , approximately 95 μM , approximately 100 μM , approximately 110 μM ,
approximately 120 μM , approximately 130 μM , approximately 140 μM , approximately 150 μM ,
approximately 160 μM , approximately 170 μM , approximately 180 μM , approximately 190 μM ,
approximately 200 μM , approximately 210 μM , approximately 220 μM , approximately 230 μM ,
approximately 240 μM , approximately 250 μM , approximately 300 μM , approximately 350 μM ,
15 approximately 400 μM , approximately 450 μM , approximately 500 μM , approximately 550 μM ,
approximately 600 μM , approximately 650 μM , approximately 700 μM , approximately 750 μM ,
approximately 800 μM , approximately 850 μM , approximately 900 μM , approximately 950 μM ,
approximately 1000 μM or an amount equivalent thereto.

The period of time the oocytes are in contact with the Zn^{2+} binding moieties should be a
20 period effective to result in activation thereof either when preceded by an increase in intracellular
calcium levels or in the absence of intracellular calcium level increase. Such time periods can be
within the range of from approximately 1 minute to approximately 12 hours. For instance, from
approximately 1 minute to approximately 5 hours, approximately 1 minute to approximately 2
hours, approximately 1 minute to approximately 1.5 hours, approximately 1 minute to
25 approximately 1 hour, approximately 1 minute to approximately 45 minutes, approximately 1
minute to approximately 30 minutes, approximately 1 minute to approximately 20 minutes,
approximately 1 minute to approximately 15 minutes, approximately 1 minute to approximately
10 minutes, approximately 1 minute to approximately 5 minutes, approximately 5 minutes to
approximately 12 hours, approximately 5 minutes to approximately 5 hours, approximately 5
30 minutes to approximately 2 hours, approximately 5 minutes to approximately 1.5 hours,

approximately 5 minutes to approximately 1 hour, approximately 5 minutes to approximately 45 minutes, approximately 5 minutes to approximately 30 minutes, approximately 5 minutes to approximately 20 minutes, approximately 5 minutes to approximately 15 minutes, approximately 5 minutes to approximately 10 minutes, approximately 10 minutes to approximately 2.5 hours, approximately 10 minutes to approximately 2 hours, approximately 10 minutes to approximately 1.5 hours, approximately 10 minutes to approximately 1 hour, approximately 10 minutes to approximately 45 minutes, approximately 10 minutes to approximately 30 minutes, approximately 10 minutes to approximately 20 minutes, approximately 10 minutes to approximately 15 minutes, approximately 30 minutes to approximately 2.5 hours, approximately 30 minutes to approximately 2 hours, approximately 30 minutes to approximately 1.5 hours, approximately 30 minutes to approximately 1 hour, approximately 45 minutes to approximately 2.5 hours, approximately 45 minutes to approximately 2 hours, approximately 45 minutes to approximately 1.5 hours, approximately 45 minutes to approximately 1 hour, approximately 1 hour to approximately 2.5 hours, approximately 1 hour to approximately 2 hours, approximately 1 hour to approximately 1.5 hours, approximately 1.5 hours to approximately 2.5 hours, approximately 1.5 hours to approximately 2 hours, approximately 2 hours to approximately 2.5 hours or a range equivalent thereto.

The period of time the oocytes are in contact with the Zn^{2+} binding moieties may therefore comprise, approximately 1 minute, approximately 2 minutes, approximately 3 minutes, approximately 4 minutes, approximately 5 minutes, approximately 6 minutes, approximately 7 minutes, approximately 8 minutes, approximately 9 minutes, approximately 10 minutes, approximately 11 minutes, approximately 12 minutes, approximately 13 minutes, approximately 14 minutes, approximately 15 minutes, approximately 16 minutes, approximately 17 minutes, approximately 18 minutes, approximately 19 minutes, approximately 20 minutes, approximately 25 minutes, approximately 30 minutes, approximately 35 minutes, approximately 40 minutes, approximately 45 minutes, approximately 50 minutes, approximately 55 minutes, approximately 1 hour, approximately 1.25 hours, approximately 1.5 hours, approximately 1.75 hours, approximately 2 hours, approximately 2.25 hours, approximately 2.5 hours, approximately 2.75 hours, approximately 3 hours, approximately 3.5 hours, approximately 4 hours, approximately 4.5 hours, approximately 5 hours, approximately 5.5 hours, approximately 6 hours,

approximately 6.5 hours, approximately 7 hours, approximately 7.5 hours, approximately 8 hours, approximately 8.5 hours, approximately 9 hours, approximately 9.5 hours, approximately 10 hours, approximately 10.5 hours, approximately 11 hours, approximately 11.5 hours, approximately 12 hours, or a time equivalent thereto.

5 For those embodiments of the invention comprising decreasing the intracellular concentration of Zn^{2+} of the unfertilized oocytes preceded by an increase in intracellular Ca^{2+} of the unfertilized oocyte, the step of contacting the oocyte with a Zn^{2+} binding moieties after increasing the intracellular level of Ca^{2+} in the oocyte may occur substantially immediately after the step of increasing the intracellular level of Ca^{2+} or may occur after a period of time
10 comprising from about 5 seconds to about 10 minutes or a range equivalent thereto. The Zn^{2+} binding moiety may be added to the media in which the increase in intracellular Ca^{2+} occurred or the oocyte may be washed or transferred into new media after the increase in intracellular Ca^{2+} . In some embodiments, oocytes may be washed to remove the presence of the Zn^{2+} binding moiety after activation. In further embodiments, activation of oocytes may be determined
15 microscopically, by observing pronuclear formation, extrusion of a polar body, division of the oocytes to the 2-cell stage, or subsequent development to the blastocyst stage. Other methods for confirming oocyte activation would be known by one of skill in the art and would be suitable for use with the present method.

The methods described herein are generally applicable to unfertilized oocytes from a
20 wide array of mammalian animals, including, but not limited to, livestock mammals, domestic mammals, model animal mammals, and human or non-human mammals. As used to herein "livestock mammals" refer to any mammalian animal that is useful in an agricultural or livestock setting, such as pig (porcine), cattle (bovine), sheep (ovine), goat, horse and buffalo. "Domestic mammals" refer herein to any mammal that has been domesticated by humans such that they are
25 tame and depend upon man for survival, such as cat (feline), dog (canine), rabbit, guinea pig, and hamster. "Model animal mammals" refer herein to any mammal used for scientific and health related research, such as mice, and rats. In certain embodiments these categories of mammals may overlap, for instance domestic mammals such as dogs may also be classified as a model animal mammal.

In certain embodiments, the present methods of activating unfertilized mammalian oocytes disclosed herein facilitate parthenogenetic development of such eggs so that development thereof can proceed to the 2-cell stage, to at least the blastocyst stage, or anywhere in between, upon culturing *in vitro* or *in vivo*. In other embodiments, the present methods of oocyte activation may be employed in the field of mammalian cloning by nuclear transfer. In particular embodiments, the method of the present invention may be used in conjunction with conventional methods of mammalian cloning well known to those skilled in the art. Reference is made to First and Prather (*Differentiation* 48, 1991) and U.S. Patent Nos. 4,994,384 and 6,211,429, which are incorporated herein by reference for a general discussion on nuclear transfer techniques.

The term "nuclear transfer" as used herein refers to a process of transferring a nucleus of a cell of an adult, fetus, embryo, or cell line (referred to herein as "nuclear donor cell") into an enucleated oocyte (referred to herein as "recipient oocyte"). In certain embodiments, the enucleated oocyte referred to herein may comprise a MII oocyte from which the MII arrested chromosomes have been removed. A MII oocyte is referred to herein as "enucleated" despite comprising chromosomes arrested in metaphase and not comprising a membrane-bound nucleus. Oocytes for use in the present invention may be matured prior to nuclear transfer. Such maturation may be done either *in vitro* or *in vivo* using methods known in the art.

The resulting cell obtained by the nuclear transfer is referred to herein as a "nucleus-transferred oocyte," "nuclear transfer oocyte," "somatic cell nuclear transfer oocyte," or "somatic cell nuclear transfer embryo." The term nuclear donor cell may refer to any cell with a nucleus containing two sets of chromosomes (2n).

In certain embodiments, the methods of the present invention may involve the use of donor cells, for example those from a cell line, that have been genetically engineered using methods well known in the art (for instance as disclosed in Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons, Inc., 1989, and Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989) with a gene or other structural coding sequence of interest that may have been genetically modified prior to carrying out nuclear transfer. In this case, the product would be a

transgenic animal with a modified characteristic. Alternatively, the donor nucleus may have been genetically modified by using other methods, such as mega-nucleases, that do not necessarily leave a genetic footprint, such as a transgene or selectable marker. Reference is made to Whyte *et al.* (*Mol. Reprod. Dev.*, 78, 2011), Lee *et al.* (*PNAS*, 11, 2014), and Whitworth *et al.* (*Biology of Reproduction*, 91, 2014), for examples of mega-nucleases, such as zinc finger nucleases, TALENs, and CRISPR/Cas9 systems, respectively, the disclosure of which are incorporated herein.

Introduction of a nucleus from a donor cell into an enucleated recipient oocyte to form an oocyte containing the donor nucleus can be performed by fusing together the membrane of a membrane-bound nucleus from the donor cell with the membrane of the enucleated recipient oocyte to form an oocyte containing the nucleus from the donor cell. Alternatively, such introduction can be performed by microinjecting the nucleus from the donor cell into the enucleated recipient oocyte to form an oocyte containing the nucleus from the donor cell. As those of ordinary skill in the art are aware, further alternative means exist for introducing donor nuclear material into a recipient oocyte. For example, and not by way of limitation, one can introduce a donor cell (or nucleus) into the space under the zona pellucida or into the perivitelline space of the enucleated, recipient oocyte, and then subsequently carry out membrane fusion to produce an oocyte containing within its cytoplasm the donor nucleus. All means of introducing donor nuclear material into an enucleated recipient oocyte known to those of ordinary skill in the art are useful in the methods disclosed herein.

In certain embodiments the present invention provides mammalian animals produced through nuclear transfer, also referred to as cloned animals, that result from a nuclear transferred oocyte or nuclear transferred embryo activated by methods disclosed herein. Such animals are distinct from the animals from which the donor nucleus originated, referred to herein as a "donor animal." For instance, in certain embodiments although such cloned animals comprise the same genomic DNA as that of the donor nucleus, cell or animal from which the donor nucleus originated, the cloned animal may comprise distinct characteristics distinguishing it from the donor animal. In particular, embodiments, the cloned animal may comprise mitochondrial DNA or epigenetic traits or characteristics distinct from that of the donor animal. Such distinctions would be well understood and known by those of skill in the art.

Unlike nuclear DNA, mitochondrial DNA of an offspring, including animals produced through nuclear transfer, is inherited from the maternal parent, *i.e.* from the oocyte. A cloned animal would likely therefore comprise distinct mitochondrial DNA from that of a donor animal. Mitochondrial DNA drives the function of the cell's mitochondria and thus cellular respiration. Distinctions in mitochondrial DNA can therefore result in distinctions in mitochondrial function and cellular respiration.

Additionally, although cloned animals may comprise the same nuclear genomic sequence, the epigenetic markers or features associated with the nuclear genome may in certain embodiments be distinct from those of the donor animal. Although in certain embodiments some epigenetic markers or features may be inherited, referred to as transgenerational epigenetic markers, and thus remain unaltered between donor animal and cloned animal, many epigenetic markers or features are removed during development (referred to in the art as "reprogramming").

Such epigenetic markers or features that may differ between donor and cloned animal may comprise, but are not limited to, DNA methylation, histone modifications, imprinting, gene silencing, and X chromosome inactivation. In particular embodiments of the invention, the result of such epigenetic distinctions between a donor and cloned animal may include distinctions in gene or protein expression, thus may altering the phenotype of resulting animal. One example of such phenotypic distinction between cloned and donor animals can be seen in the original cloned cat, known in the art as "CopyCat." The nuclear donor for CopyCat was a calico cat but due to epigenetic changes between the donor and CopyCat (specifically X inactivation) CopyCat has a coloration distinct from that of the donor cat despite the two cats comprising the same nuclear genome. The genetic and epigenetic differences between cloned and donor animals, including but not limited to those discussed above, are known in the art.

EXAMPLES

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples, which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate

that many changes can be made in the specific embodiments, which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Oocyte maturation

5 Oocytes were obtained from either pre-pubertal gilt ovaries from an abattoir (Farmland Foods Inc., Milan, MO), or sow-derived oocytes that were purchased from Applied Reproductive Technology (Madison, WI). Immature oocytes from gilt ovaries were aspirated from medium size (3–6 mm) follicles by using an 18 gauge hypodermic needle attached to a 10 mL syringe. Oocytes with evenly dark cytoplasm and intact surrounding cumulus cells were then selected for
10 maturation. Around 50 cumulus oocyte complexes were placed in a well containing 500 μ L of maturation medium, TCM 199 (Invitrogen, Grand Island, NY) with 3.05 mM glucose, 0.91 mM sodium pyruvate, 0.57 mM cysteine, 10 ng/mL epidermal growth factor (EGF), 0.5 mg/mL luteinizing hormone (LH), 0.5 mg/mL follicle stimulating hormone (FSH), 10 ng/mL gentamicin (APP Pharm, Schaumburg, IL), and 0.1% polyvinyl alcohol (PVA) for 42–44 hr at 38.5°C, 5%
15 CO₂, in humidified air. The oocytes from sows were shipped overnight in maturation medium (TCM199 with 2.9 mM Hepes, 5 mg/mL insulin, 10 ng/mL EGF, 0.5 mg/mL p-FSH, 0.91 mM pyruvate, 0.5 mM cysteine, 10% porcine follicular fluid, 25 ng/mL gentamicin) and transferred into fresh medium at 24 hr. At the end of the maturation, the surrounding cumulus cells were removed from the oocytes by vortexing for 3 min in the presence of 0.1% hyaluronidase.
20 Oocytes with a visible polar body were selected in manipulation medium (TCM199 with 0.6 mM NaHCO₃, 2.9 mM Hepes, 30 mM NaCl, 10 ng/mL gentamicin, and 3 mg/mL bovine serum albumin [BSA]; and osmolarity of 305) and then used for the experiments.

Example 2

Various oocyte activation methods

25 Different oocyte activation methods were used for the experiments. As conventional activation methods, either electrical or chemical activation was used. For electrical activation, mature oocytes were activated in activation medium (0.3 M mannitol, 1.0 mM CaCl₂, 0.1 mM MgCl₂, and 0.5 mM Hepes) by two direct-current (DC) pulses (1-sec interval) at 1.2 kV/cm for 30 μ sec (using BTX Electro Cell Manipulator, Harvard Apparatus, Holliston, MA). For chemical

activation, mature oocytes were incubated in Hepes-buffered Tyrode's lactate (TL-Hepes) media in the presence of 200 mM thimerosal for 10 min in the dark followed by 8 mM dithiothreitol for 30 min. For TPEN mediated activation, mature oocytes were incubated in TL-Hepes media with various concentrations of TPEN for different durations.

Activated oocytes were washed three times in PZM3 then incubated at 38.5°C, 5% CO₂ until the embryos were examined on either day 2, 6 or 7. At the end of the culture, where applicable, the frequency of cleaved embryos and blastocysts, and total cell number in the blastocysts were recorded. Hoechst 33342 (1.2 mg/mL) was used to stain nuclei and the embryos were then evaluated by epi-fluorescence microscopy.

Example 3

Intracellular calcium measurement

The zonae pellucidae were removed after maturation and zona-free oocytes were loaded with the Ca²⁺ indicator dye, fura-2. For this purpose, they were incubated in the presence of 2 mM of the acetoxymethyl ester form of the dye and 0.02% pluronic F-127 for 40–50 min (both from Invitrogen). The dye loaded oocytes were transferred into a chamber with a cover-glass bottom and the chamber was then placed on the heated stage of an inverted microscope. Changes in the intracellular free Ca²⁺ concentration were recorded using InCytIm2, a dual-wavelength fluorescence imaging system (Intracellular Imaging, Inc.; Cincinnati, OH). During measurements the emitted fluorescence was detected at 510 nm after exciting the dye alternately at 340 and 380 nm. The ratio of the two emitted fluorescence intensities was calculated and the data are presented as fluorescence ratio values. Ratios of 1.0 and 5.0 correspond to about 100 and 1,200 nM Ca²⁺, respectively. In each treatment group the measurements were repeated at least 5 times using different oocytes.

Example 4

Parthenogenetic activation of porcine oocytes by TPEN

Metaphase II porcine oocytes were incubated *in vitro* with TPEN to determine if the concentration/duration of 100 µM of TPEN for 45 min, as described in the mouse study by Suzuki, *et al.* (*Development* 137, 2659-2669, 2010) had any effect on pig oocytes. The protocol

described by Suzuki, *et al.* could activate pig oocytes, however, the development was inferior to conventional artificial activation methods. Frequency of forming blastocysts on day 7 was comparable to the conventional oocyte activation method (13% vs. 15%, respectively) but the rate of blastocyst formation was slower. No blastocyst formation was observed on day 5 with the TPEN treatment whereas 10% of activated embryos reached blastocysts in the control group. In addition, no expanded blastocysts were derived from the TPEN activation group on day 7, indicating lower developmental competency of the embryos derived from the TPEN activation method.

A wide range of TPEN concentrations/durations was tested to identify an optimum concentration of TPEN that can activate pig oocytes; 100 μ M, 200 μ M, and 250 μ M and 30 min - 2.5 hours of duration were tested (Table 1). Optimum embryo development was achieved by using 200 μ M TPEN for 30 min. Although the frequency of blastocyst formation was not statistically superior to a conventional activation method, a numerical increase in blastocyst formation was observed. A lower total number of nuclei in blastocysts was observed in the embryos derived from the optimal TPEN activation group (200 μ M, 30 min), compared to the control (27.2 ± 1.92 vs. 32.6 ± 3.12 , respectively, $P < 0.01$). When oocytes were incubated in 250 μ M TPEN over 2.5 hours, a toxic effect of TPEN was observed. The 250 μ M TPEN treatment over 2.5 hours blocked embryo development as there was no blastocyst formation. An optimum concentration of TPEN which successfully induced development of activated oocytes was identified, but the total number of nuclei in these embryos was lower compared to the conventional activation method.

Table 1. Porcine oocytes activated with TPEN alone.

Treatment	Total number of oocytes	% blastocysts
Electroporation	98	11.2%
100 μ M, 2 hr	65	12.0%
200 μ M, 30 min	25	23.0%
200 μ M, 1 hr	79	20.0%

250 μ M, 2hr	30	10.0%
250 μ M, 2.5 hr	25	0%

Example 5

Treatment of activated oocytes with TPEN

Treatment of activated oocytes with TPEN was investigated for the potential to increase the efficiency of oocyte activation. When chemically activated oocytes were incubated with TPEN (100 μ M, 45 min), no blastocyst formation was observed suggesting the combination of the two methods is toxic to embryos. It was hypothesized that the detrimental effect was due to excess stimuli from Ca^{2+} signaling and TPEN. Therefore, lower concentrations of TPEN were introduced after activating oocytes with a conventional method. Incubating activated oocytes with a lower concentration of TPEN increased developmental potential of parthenogenetic embryos. Interestingly, when activated oocytes were incubated with a low concentration of TPEN (5-10 μ M) the TPEN treated group surprisingly showed higher developmental potential compared to the control group. Specifically, the average percent blastocyst formation of TPEN treated oocytes (5 μ M for 30 min) was $27.2 \pm 1.7\%$ but only $10.6 \pm 2.5\%$ developed to blastocyst in the control group (Table 2). Moreover, the average number of nuclei in blastocysts was higher in TPEN treated oocytes compared to the control group which was activated by thimerosal/DTT approach (Table 2).

Table 2. Incubation of TPEN can increase development potential of activated oocytes. Different letters indicate statistical difference ($p < 0.05$).

Treatment	Total number of embryos	% cleaved	% Blastocysts	Total cell number in blastocysts
THI/DTT	113	59.3 ^a	10.6 ^a	28.2 ± 2.1^a
THI/DTT + 5 μ M TPEN, 30 min	114	74.6 ^b	27.2 ^b	33.1 ± 2.6^b
THI/DTT + 10 μ M TPEN, 10 min	83	81.9 ^b	21.7 ^b	31.2 ± 2.6^b

Next, the optimal concentration and duration of TPEN (5 μ M, 30 min) was used to artificially activate reconstructed SCNT embryos. There was an increase in frequency of blastocyst formation when activated SCNT embryos were treated with TPEN (7.9% vs. 18.6%, respectively) (Table 3). This increase in development was comparable to the effect of Scriptaid, a histone deacetylase (HDAC) inhibitor, known to increase developmental potential of SCNT embryos. Thus lower concentrations of TPEN (5 – 10 μ M) can enhance embryo development when introduced after conventional oocyte activation.

Table 3. *In vitro* development of SCNT embryos treated with Scriptaid or TPEN alone.

Different letters indicate statistical difference ($p < 0.05$).

Treatment	Number of embryos	% Blastocysts
SCNT (control)	76	7.9% ^a
SCNT + Scriptaid	194	22.7% ^b
SCNT + 5 μ M TPEN, 30 min	102	18.6% ^b

Example 6

TPEN can serve as the main activating reagent

It was hypothesized that a proper combination of Ca^{2+} signal and TPEN can increase the developmental potential of activated oocytes. The ability of TPEN as the main activating reagent was therefore tested. First, oocytes received an intracellular Ca^{2+} increase through electroporation, although the concentration of Ca^{2+} (0.1 mM) in the electroporation media was not sufficient to induce oocyte activation. Then the oocytes were incubated with various concentrations (1 μ M, 5 μ M, 10 μ M, 100 μ M and 200 μ M) and durations (10 minutes, 20 minutes, 30 minutes, 1 hour, 2 hours, and overnight) of TPEN. The combination of a smaller Ca^{2+} increase and lower concentration of TPEN (10 μ M) resulted in 0 day 6 blastocysts. Surprisingly, oocytes activated by a smaller Ca^{2+} increase followed by TPEN incubation at higher concentration (100 μ M or above) could successfully activate oocytes and some

combinations resulted in a superior *in vitro* development compared to the oocytes activated by a conventional method (Table 4). Additionally, oocytes incubated in 1 μ M TPEN overnight presented an increased number of blastocysts at day 6 and 7, although the observed cell numbers in day 7 blastocysts were decreased (24.80 ± 7.84) as compared to oocytes incubated in 10 μ M TPEN for 1 hour (30.17 ± 5.97), 10 μ M TPEN for 2 hours (29.63 ± 10.47) or 100 μ M TPEN for 20 minutes (50.75 ± 17.94). Optimal oocyte activation was observed when oocytes were activated by a smaller Ca^{2+} signal followed by incubating in 200 μ M TPEN for 30 min.

Table 4. Effect of TPEN as a main oocyte activating reagent. Different letters indicate statistical difference ($p < 0.05$).

Treatment	Number of embryos	Day 6 Blastocysts (%)	Day 7 Blastocysts (%)
Control - electroporation	80	13.75 ^a	17.5 ^a
10 μ M, 30 min	40	0 ^b	5 ^b
10 μ M, 1 hour	40	15 ^a	22.5 ^a
10 μ M, 2 hours	40	15 ^a	25 ^a
100 μ M, 10 min	40	15 ^a	15 ^a
100 μ M, 20 min	40	25 ^a	30 ^a
200 μ M, 30 min	81	37.04 ^c	41.98 ^c
200 μ M, 1 hour	80	18.75 ^a	27.5 ^a

Additionally, the effect of the TPEN activation method alone on the development of SCNT embryos was tested. For this assay, SCNT embryos were activated using either the conventional thimerosal/DTT method or TPEN (TPEN as a major activating reagent). Although no statistical difference was observed, likely due to the small number of observations, a numerical increase was observed for embryos activated using TPEN as the major activating reagent as compared to those activated using the conventional thimerosal/DTT method during SCNT procedure (Table 5).

Table 5. Effect of TPEN activation method on the development of SCNT embryos.

Activation method	Day 6 blastocyst formation	Day 7 blastocyst formation
THI/DTT	23.3% (7/30)	26.7% (8/30)
TPEN (200 μ M, 30 min)	25.8% (8/31)	35.5% (11/31)

Example 7**No Ca^{2+} increase is observed in pig oocytes after exposure to TPEN**

5 To demonstrate that TPEN treatment is stimulating signaling pathways independent of Ca^{2+} signaling, oocytes were exposed to TPEN then the amount of intracellular Ca^{2+} was measured. As expected, no Ca^{2+} increase was detected when oocytes were exposed to a range of TPEN (5, 10, and 200 μ M), while oocytes incubated with thimerosal, serving as a positive control, could induce Ca^{2+} increase (Figure 1). This indicates that the benefit of TPEN observed
10 during oocyte activation experiments is independent of Ca^{2+} signaling pathway.

Example 8***In vivo* development competence of SCNT embryos activated by utilizing TPEN**

To address *in vivo* competency of SCNT embryos activated by utilizing TPEN, a series of embryo transfers were performed. For the study, at day 1, the SCNT-derived embryos were
15 surgically transferred into the ampullary-isthmic junction of a surrogate gilt at 0 or 1 days after observed estrus. At the end of gestation periods, piglets were recovered through C-section.

Six embryo transfers were performed for the study and two surrogates carried their pregnancy to term, each resulting in two healthy piglets (Table 6). One of the term development was from SCNT embryos that were not treated with a HDAC inhibitor, Scriptaid. The resulting
20 full term development indicates that TPEN is compatible with SCNT development to term and suggests that the TPEN technology is not toxic to the SCNT embryos. None of the resulting animals displayed any health related issues at birth as are sometime observed with cloned animals, and they have produced progeny.

Table 6. Result of embryo transfer.

Treatment	Number of embryos transferred	Result
SCNT + 5 μ M TPEN, 30 min + Scriptaid	207	2 normal piglets were delivered
SCNT + 5 μ M TPEN, 30 min + Scriptaid	245	Cycled
SCNT + 10 μ M TPEN, 10 min + Scriptaid	180	Cycled
SCNT + 5 μ M TPEN, 30 min	280	2 normal piglets were delivered
SCNT + 5 μ M TPEN, 30 min	230	Cycled
SCNT + 5 μ M TPEN, 30 min	280	Cycled

Example 9

Statistical analysis

Differences in the frequency of blastocyst formation was determined after analysis of variance (ANOVA) using the PROC MIX procedure of the Statistical Analysis System (SAS Institute, Cary, NC, USA) or Chi-square test. Percentage data was arcsin transformed prior to the ANOVA analysis. Average total number of nuclei in blastocysts was compared by using the Student's T-test. Differences with $P < 0.05$ were considered significant.

WHAT IS CLAIMED IS:

Claim 1. A method of activating an unfertilized livestock or domestic mammalian oocyte comprising decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety.

Claim 2. The method of claim 1, wherein the Zn^{2+} binding moiety comprises a concentration of between approximately 10 μM to approximately 500 μM .

Claim 3. The method of claim 2, wherein the Zn^{2+} binding moiety comprises a concentration of between approximately 100 μM to approximately 250 μM .

Claim 4. The method of claim 3, wherein the Zn^{2+} binding moiety comprises a concentration of approximately 200 μM .

Claim 5. The method of claim 1, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 1 minute to approximately 5 hours.

Claim 6. The method of claim 5, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 30 minutes to approximately 2.5 hours.

Claim 7. The method of claim 6, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising approximately 30 minutes.

Claim 8. The method of claim 1, further comprising the step of increasing intracellular Ca^{2+} concentration of the oocyte prior to decreasing the intracellular Zn^{2+} concentration of the oocyte.

Claim 9. The method of claim 8, wherein the intracellular Ca^{2+} concentration of the oocyte is not increased in an amount sufficient to induce oocyte activation.

Claim 10. The method of claim 9, wherein the intracellular Ca^{2+} concentration of the oocyte is increased approximately ten times less than the amount sufficient to induce oocyte activation.

Claim 11. The method of claim 9, wherein the Zn^{2+} binding moiety comprises a concentration of between approximately 1 μM to approximately 500 μM .

Claim 12. The method of claim 11, wherein the Zn^{2+} binding moiety comprises a concentration of between approximately 100 μM to approximately 200 μM .

Claim 13. The method of claim 12, wherein the Zn^{2+} binding moiety comprises a concentration of approximately 200 μM .

Claim 14. The method of claim 9, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 1 minutes to approximately 12 hours.

Claim 15. The method of claim 14, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 10 minutes to approximately 2 hours.

Claim 16. The method of claim 15, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising approximately 30 minutes.

Claim 17. The method of claim 8, wherein the intracellular Ca^{2+} concentration of the oocyte is increased in an amount sufficient to induce oocyte activation.

Claim 18. The method of claim 17, wherein the Zn^{2+} binding moiety comprises a concentration of between approximately 0.1 μM to approximately 100 μM .

Claim 19. The method of claim 18, wherein the Zn^{2+} binding moiety comprises a concentration of between approximately 1 μM to approximately 10 μM .

Claim 20. The method of claim 19, wherein the Zn^{2+} binding moiety comprises a concentration of approximately 5 μM .

Claim 21. The method of claim 19, wherein the Zn^{2+} binding moiety comprises a concentration of approximately 10 μM .

Claim 22. The method of claim 17, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 1 minutes to approximately 5 hours.

Claim 23. The method of claim 22, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising between approximately 10 minutes to approximately 30 minutes.

Claim 24. The method of claim 23, wherein the Zn^{2+} binding moiety contacts the oocyte for a period of time comprising approximately 30 minutes.

Claim 25. The method of claim 8, wherein the intracellular Ca^{2+} concentration of the oocyte is increased by chemical or physical means or a combination thereof that induce calcium entry into the oocyte or release of internal calcium stores.

Claim 26. The method of claim 25, wherein the increase in the intracellular Ca^{2+} concentration of the oocyte by chemical means comprises treatment with a calcium containing medium, a calcium salt containing medium, thimerosal containing medium, ethanol containing medium, inositol trisphosphate containing medium, a calcium ionophore containing medium, a sodium-free medium or a combination thereof.

Claim 27. The method of claim 26, wherein the chemical treatment comprises treatment with a medium comprising a Ca^{2+} concentration of between approximately 0.1 mM and approximately 250 mM.

Claim 28. The method of claim 27, wherein the chemical treatment comprises treatment with a medium comprising a Ca^{2+} concentration of between approximately 0.1 mM and 1 mM.

Claim 29. The method of claim 25, wherein the increase in the intracellular Ca^{2+} concentration of the oocyte by physical means comprises application of an electrical stimulus.

Claim 30. The method of claim 1, wherein the Zn^{2+} binding moiety comprises TPEN (N,N,N',N'-tetrakis(2-pyridylmethyl)ethane-1,2-diamine).

Claim 31. The method of claim 1, wherein the unfertilized livestock or domestic mammalian oocyte is a nuclear transfer oocyte.

Claim 32. The method of claim 1, wherein the unfertilized livestock or domestic mammalian oocyte is selected from the group consisting of a porcine oocyte, bovine oocyte, ovine oocyte, goat oocyte, horse oocyte, canine oocyte and feline oocyte.

Claim 33. A parthenogenetic oocyte activated by the method of claim 1.

Claim 34. A nuclear transfer oocyte activated by the method of claim 1.

Claim 35. A viable livestock or domestic mammalian animal produced from an unfertilized livestock or domestic mammalian oocyte activated by the method of claim 1, wherein said animal comprises the genomic DNA from a donor cell other than the oocyte and the mitochondrial DNA from the oocyte.

Claim 36. The viable livestock or domestic mammalian animal of claim 35, wherein the animal comprises epigenetic features different from the epigenetic features of the donor cell.

Claim 37. The viable livestock or domestic mammalian animal of claim 36, wherein the epigenetic features different from the epigenetic features of the donor cell are DNA methylation sites or histone modifications.

Claim 38. The viable livestock or domestic mammalian animal of claim 35, wherein the animal is selected from the group consisting of a porcine animal, bovine animal, ovine animal, goat, horse, dog and cat.

Claim 39. A method of activating an unfertilized mammalian oocyte comprising increasing intracellular Ca^{2+} concentration of the oocyte by contacting the oocyte with a medium comprising a calcium concentration of approximately 0.1 mM and decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety at a concentration of approximately 200 μM for a period of time comprising approximately 30 minutes.

Claim 40. A method of activating an unfertilized mammalian oocyte comprising increasing intracellular Ca^{2+} concentration of the oocyte by contacting the oocyte with a medium comprising a calcium concentration of approximately 1.0 mM and decreasing intracellular Zn^{2+}

concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety at a concentration of approximately 5 μM for a period of time comprising approximately 30 minutes.

Claim 41. A method of activating an unfertilized mammalian oocyte comprising increasing intracellular Ca^{2+} concentration of the oocyte by contacting the oocyte with a medium comprising a calcium concentration of approximately 1.0 mM and decreasing intracellular Zn^{2+} concentration of the oocyte by contacting the oocyte with a Zn^{2+} binding moiety at a concentration of approximately 10 μM for a period of time comprising approximately 30 minutes.




FIG. 1A

2/2




FIG.1B

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2014/067588

A. CLASSIFICATION OF SUBJECT MATTER INV. C12N5/075 C12N15/877 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) C12N		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, WPI Data, BIOSIS, EMBASE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	B. Y. KONG ET AL: "Zinc Maintains Prophase I Arrest in Mouse Oocytes Through Regulation of the MOS-MAPK Pathway", BIOLOGY OF REPRODUCTION, vol. 87, no. 1, 25 April 2012 (2012-04-25) , pages 11-11, XP055166423, ISSN: 0006-3363, DOI: 10.1095/biolreprod.112.099390 abstract page 2 - page 3	1-41
X	----- US 2013/017602 A1 (WOODRUFF TERESA K [US] ET AL) 17 January 2013 (2013-01-17) example 1 claims 1-5 ----- -/-	1-41
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier application or patent but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 24 February 2015		Date of mailing of the international search report 02/03/2015
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Loubradou-Bourges, N

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2014/067588

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	LAI L ET AL: "Production of alpha-1,3-galactosyltransferase knockout pigs by nuclear transfer cloning", SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, US, vol. 295, 8 February 2002 (2002-02-08), pages 1089-1092, XP002962717, ISSN: 0036-8075, DOI: 10.1126/SCIENCE.1068228 abstract	34-38
X	----- US 2003/066100 A1 (MACHATY ZOLTAN [US] ET AL) 3 April 2003 (2003-04-03)	33
A	paragraph [0051]	1-32, 34-41
X	----- US 5 496 720 A (SUSKO-PARRISH JOAN L [US] ET AL) 5 March 1996 (1996-03-05)	33
A	claim 1	1-32, 34-41
X,P	----- MING-HUI ZHAO ET AL: "Zinc Regulates Meiotic Resumption in Porcine Oocytes via a Protein Kinase C-Related Pathway", PLOS ONE, vol. 9, no. 7, 14 July 2014 (2014-07-14), page e102097, XP055166622, DOI: 10.1371/journal.pone.0102097 the whole document	1-41

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2014/067588

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2013017602 A1	17-01-2013	NONE	
US 2003066100 A1	03-04-2003	US 2003066100 A1 WO 03040354 A1	03-04-2003 15-05-2003
US 5496720 A	05-03-1996	US 5496720 A US 5843754 A US 6077710 A US 6194202 B1	05-03-1996 01-12-1998 20-06-2000 27-02-2001