



HU000024291T2



(19) **HU**

(11) Lajstromszám: **E 024 291**

(13) **T2**

**MAGYARORSZÁG**  
Szellemi Tulajdon Nemzeti Hivatala

## **EURÓPAI SZABADALOM** **SZÖVEGÉNEK FORDÍTÁSA**

(21) Magyar ügyszám: **E 08 865838**

(22) A bejelentés napja: **2008. 12. 23.**

(96) Az európai bejelentés bejelentési száma:  
**EP 20080865838**

(97) Az európai bejelentés közzétételi adatai:  
**EP 2242772 A1** **2009. 07. 02.**

(97) Az európai szabadalom megadásának meghirdetési adatai:  
**EP 2242772 B1** **2014. 11. 26.**

(51) Int. Cl.: **C07K 16/28**

(2006.01)  
**A61K 393/95**

(2006.01)  
**A61K 47/48**

(2006.01)  
**C07K 16/30**

(2006.01)  
**A61P 35/00**

(86) A nemzetközi (PCT) bejelentési szám:

**PCT/EP 08/068267**

(87) A nemzetközi közzétételi szám:

**WO 09080830**

(30) Elsőbbségi adatai:

16620 P	2007. 12. 26.	US
87466 P	2008. 08. 08.	US
87590 P	2008. 08. 08.	US

(73) Jogosult(ak):

**Biostest AG, 63303 Dreieich (DE)**  
**Immunogen, Inc., Waltham, MA 02451-1477**  
**(US)**

(72) Feltaláló(k):

KRAUS, Elmar, 61118 Bad Vilbel (DE)  
BRUECHER, Christoph, 65760 Eschborn (DE)  
DAELKEN, Benjamin, 60433 Frankfurt am Main (DE)  
ZENG, Steffen, 64839 Münster (DE)  
OSTERROTH, Frank, 63128 Dietzenbach (DE)  
UHEREK, Christoph, 63500 Seligenstadt (DE)  
AIGNER, Silke, 67227 Frankenthal (DE)  
GERMER, Matthias, 63225 Langen (DE)  
Schulz, Gregor, 79224 Umkirch (DE)  
HAEDER, Thomas, 63303 Dreieich (DE)

(74) Képviselő:

**Ráthonyi Zoltán, SBGK Szabadalmi Ügyvivői**  
**Iroda, Budapest**

(54)

**CD138-at megcélzó immunkonjugátumok és alkalmazásai**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

**EP 2 242 772 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention of the grant of the patent:  
**26.11.2014 Bulletin 2014/48**

(51) Int Cl.:  
**C07K 16/28 (2006.01)** **C07K 16/30 (2006.01)**  
**A61P 35/00 (2006.01)** **A61K 47/48 (2006.01)**  
**A61K 39/395 (2006.01)**

(21) Application number: **08865838.0**

(86) International application number:  
**PCT/EP2008/068267**

(22) Date of filing: **23.12.2008**

(87) International publication number:  
**WO 2009/080830 (02.07.2009 Gazette 2009/27)**

(54) **IMMUNOCONJUGATES TARGETING CD138 AND USES THEREOF**

AUF CD138 ZIELENDE IMMUNKONJUGATE UND ANWENDUNGEN DAVON

IMMUNOCONJUGUÉS CIBLANT CD138 ET UTILISATIONS DE CEUX-CI

(84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR**  
**HR HU IE IS IT LI LT LU LV MC MT NL NO PL PT**  
**RO SE SI SK TR**

(30) Priority: **26.12.2007 US 16620 P**  
**08.08.2008 US 87466 P**  
**08.08.2008 US 87590 P**

(43) Date of publication of application:  
**27.10.2010 Bulletin 2010/43**

(60) Divisional application:  
**14160780.4**

(73) Proprietors:  

- **Biotest AG**  
**63303 Dreieich (DE)**
- **Immunogen, Inc.**  
**Waltham, MA 02451-1477 (US)**

(72) Inventors:  

- **KRAUS, Elmar**  
**61118 Bad Vilbel (DE)**
- **BRUECHER, Christoph**  
**65760 Eschborn (DE)**
- **DAELKEN, Benjamin**  
**60433 Frankfurt am Main (DE)**
- **ZENG, Steffen**  
**64839 Münster (DE)**
- **OSTERROTH, Frank**  
**63128 Dietzenbach (DE)**
- **UHEREK, Christoph**  
**63500 Seligenstadt (DE)**

- **AIGNER, Silke**  
**67227 Frankenthal (DE)**
- **GERMER, Matthias**  
**63225 Langen (DE)**
- **SCHULZ, Gregor**  
**79224 Ummkirch (DE)**
- **HAEDER, Thomas**  
**63303 Dreieich (DE)**

(74) Representative: **Daniels, Jeffrey Nicholas**  
**Page White & Farrer**  
**Bedford House**  
**John Street**  
**London WC1N 2BF (GB)**

(56) References cited:  
**WO-A-2004/099379** **WO-A-2006/099875**  
**WO-A-2007/066109** **WO-A-2007/144046**  
**US-A1- 2005 272 128** **US-A1- 2007 183 971**

- **TASSONE PIERFRANCESCO ET AL: "Cytotoxic activity of the maytansinoid immunoconjugate B-B4-DM1 against CD138+ multiple myeloma cells."**  
**BLOOD 1 DEC 2004, vol. 104, no. 12, 1 December 2004 (2004-12-01), pages 3688-3696, XP002521112 ISSN: 0006-4971**
- **SHARKEY ROBERT M ET AL: "Targeted therapy of cancer: new prospects for antibodies and immunoconjugates." CA: A CANCER JOURNAL FOR CLINICIANS 2006 JUL-AUG, vol. 56, no. 4, July 2006 (2006-07), pages 226-243, XP002522041 ISSN: 0007-9235**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

- TURNER J HARVEY ET AL: "131I-Anti CD20 radioimmunotherapy of relapsed or refractory non-Hodgkins lymphoma: a phase II clinical trial of a nonmyeloablative dose regimen of chimeric rituximab radiolabeled in a hospital." *CANCER BIOTHERAPY & RADIOPHARMACEUTICALS* AUG 2003, vol. 18, no. 4, August 2003 (2003-08), pages 513-524, XP002522042 ISSN: 1084-9785
- ISRAEL L ET AL: "Plasmapheresis and immunological control of cancer." *LANCET* 18 SEP 1976, vol. 2, no. 7986, 18 September 1976 (1976-09-18), pages 642-643, XP002522043 ISSN: 0140-6736
- CORTESINI RAFFAELLO: "Pancreas cancer and the role of soluble immunoglobulin-like transcript 3 (ILT3)." *JOP : JOURNAL OF THE PANCREAS* 2007, vol. 8, no. 6, 1 November 2007 (2007-11-01), pages 697-703, XP002522044 ISSN: 1590-8577
- ROGUSKA M A ET AL: "HUMANIZATION OF MURINE MONOCLONAL ANTIBODIES THROUGH VARIABLE DOMAIN RESURFACING", *FASEB JOURNAL, FED. OF AMERICAN SOC. FOR EXPERIMENTAL BIOLOGY, US*, vol. 91, 1 February 1994 (1994-02-01), pages 969-973, XP002917393, ISSN: 0892-6638

**Description****FIELD OF THE INVENTION**

5 [0001] The present invention relates to immunoconjugates capable of targeting CD138 expressing cells, compositions comprising the immunoconjugates and methods employing them.

**BACKGROUND**

10 [0002] CD138, which acts as a receptor for the extracellular matrix, is overexpressed on multiple myeloma (MM) cells and has been shown to influence MM cell development and/or proliferation. CD138 is also expressed on cells of ovarian carcinoma, kidney carcinoma, gall bladder carcinoma, breast carcinoma, prostate cancer, lung cancer, colon carcinoma cells and cells of Hodgkin's and non-Hodgkin's lymphomas, chronic lymphocytic leukemia (CLL) to name just a few.

15 [0003] Tassone et al. (2004) have reported excellent binding of the murine IgG1 antibody B-B4 to the CD138 antigen expressed on the surface of MM cells. Tassone also reported high cytotoxic activity of the immunoconjugate B-B4-DM1, which comprises the mytansinoid DM1 as an effector molecule, against multiple myeloma cells (see also US Patent Publ. 20070183971).

20 [0004] While Tassone et al. have contributed to providing an effective treatment of MM and a composition of matter that may be employed in such a treatment, there remain a number of needs in the art.

25 [0005] There remains a need for immunoconjugates based on B-B4 that are devoid of certain properties and/or functions associated with B-B4. There is, in particular a need for a chimerized antibody based on B-B4 that binds the CD138 as effectively as B-B4 but can be administered to humans without *significant* side effects. There is also a need for such a B-B4 based immunoconjugate that shows one or more advantageous properties relative to its murine counterpart. Those properties include improved antigen binding, improved killing of tumor cells comprising, in particular of CD138 expressing tumor cells, and cells accessory thereto or more homogenous binding of the target.

**SUMMARY OF THE INVENTION**

30 [0006] The present invention is directed at an immunoconjugate capable of targeting CD138 expressing cells comprising:

(a) an engineered targeting antibody against CD138 comprising:

35 (i) an immunoglobulin heavy chain comprising a variable region comprising amino acid residues 31 to 35 (CDR1), 51 to 68 (CDR2) and 99 to 111 (CDR3) of SEQ ID NO: 1, and  
(ii) a light chain comprising a variable region comprising amino acid residues 24 to 34 (CDR1), 50 to 56 (CDR2) and 89 to 97 (CDR3) of SEQ ID NO: 2; and

40 (b) an effector molecule for inducing cell death,

45 wherein a constant region of said immunoglobulin heavy chain is an IgG4 isotype constant region.

[0007] The present invention also provides an immunoconjugate according to any one of claims 1 to 7 for use in treating multiple myeloma in a subject.

45 [0008] In addition the present invention provides an immunoconjugate according to any one of claims 1 to 7 for use in treating transitional cell bladder carcinoma in a subject.

[0009] The present invention further provides an immunoconjugate according to any one of claims 1 to 7 for use as a medicament.

50 [0010] Still further the present invention provides an immunoconjugate according to any one of claims 1 to 7 for use in inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient.

[0011] Moreover, the present invention provides an immunoconjugate according to any one of claims 1 to 7, for inhibiting, delaying and/or preventing the growth of a tumor and/or spread of malignant tumor cells comprising CD138 expressing cells in a patient, wherein the patient has been treated with one or more cytotoxic agents and/or radiation in an amount to reduce tumor load.

55 [0012] The present invention also provides an immunoconjugate according to any one of claims 1 to 7 for use in treating a subject having a condition that would benefit from the suppression of myeloma cell survival, wherein the immunoconjugate is capable of selectively decreasing survival or growth of said myeloma cells of said subject.

[0013] In addition, the present invention provides a pharmaceutical composition comprising the immunoconjugate of

any one of claims 1 to 7 and one or more pharmaceutically acceptable excipients, wherein the composition is suitable for the inhibition, delay and/or prevention of the growth of tumors and/or spread of tumor cells.

**[0014]** Further the present invention provides a kit comprising, in separate containers, pharmaceutical compositions for use in combination to inhibit, delay and/or prevent the growth of tumors and/or spread of tumor cells, wherein one container comprises an effective amount of the pharmaceutical composition of claim 16, and wherein, a separate container comprises a second pharmaceutical composition comprising an effective amount of a further agent, preferably a cytotoxic agent, for the inhibition, delay and/or prevention of the growth of tumors and/or spread of tumor cells, and one or more pharmaceutically acceptable excipients,

preferably wherein said agent in said second pharmaceutical composition is selected from the group consisting of mephalan, vincristine, doxorubicin, dexamethasone, cyclophosphamide, etoposide, cytarabine, cisplatin, thalidomide, prednisone, bortezomib, lenalidomide, sorafenib, romidepsin and combinations thereof or is antibody based.

**[0015]** Still further, the present invention provides use of an immunoconjugate according to any one of claims 1 to 7 for the manufacture of a medicament for inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient.

**[0016]** The engineered targeting antibody of the present description may

- (i) consist essentially of antigen binding region (ABR) against CD138 of a non-human antibody, or
- (ii) comprise an antigen binding region (ABR) against CD138, wherein said antigen binding region is of a non-human antibody, and

a further antibody region, wherein at least part of said further antibody region is of a human antibody.

**[0017]** The ABR of the description may comprise:

- (a) heavy chain variable region CDR3 comprising amino acid residues 99 to 111 of SEQ ID NO: 1, and
- (b) light chain variable region CDR3 comprising amino acid residues 89 to 97 of SEQ ID NO: 2, respectively.

**[0018]** The ABR of the description may further comprise:

- (a) heavy chain variable region CDR1 and CDR2 comprising amino acid residues 31 to 35 and 51 to 68 of SEQ ID NO: 1, and/or
- (b) light chain variable region CDR1 and CDR 2 comprising amino acid residues 24 to 34 and 50 to 56 of SEQ ID NO: 2, respectively.

**[0019]** The further antibody region may comprise:

- (a) amino acid residues 123 to 448 of SEQ ID NO: 1, and/or
- (b) amino acid residues 108 to 214 of SEQ ID NO: 2, respectively and mutations thereof that
  - (i) maintain or lower the antibody-dependent cytotoxicity and/or complement-dependent cytotoxicity of the engineered targeting antibody and/or
  - (ii) stabilize the engineered targeting antibody.

**[0020]** The effector molecule may be attached to said engineered targeting antibody via a linker. The linker may comprise a disulfide bond. The effector molecule (e.g., DM4) may provide sterical hindrance between the targeting antibody and the effector molecule. The effector molecule may be at least one maytansinoid (e.g., DM1, DM3, or DM4) taxane or a CC1065, or an analog thereof.

**[0021]** The immunoconjugate may bind CD138 with a targeting variation of less than 150%, 140%, 130%, 120%, 110%, 100%, 90%, 80%, 70%, 60% or 50%.

**[0022]** The present description is also directed at an immunoconjugate comprising:

- a targeting agent targeting CD138 comprising
- an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof, wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1. A constant region of said immunoglobulin heavy chain or said part thereof may be an IgG4 isotype constant region.

**[0023]** The present description is also directed at a method of treating MM in a subject, comprising:

- providing one of more of the immunoconjugates specified herein, and

administering to said subject said immunoconjugate in an amount effective to treat multiple myeloma.

5 [0024] The targeting agent of the immunoconjugate of the description may comprise a light chain sequence having at least about 70% sequence identity with SEQ ID NO:2. The targeting agent of the immunoconjugate of the description may also comprise a heavy chain sequence having at least about 70% sequence identity with SEQ ID NO:1.

[0025] The present description is also directed at a method for immunoconjugate mediated drug delivery comprising:

10 providing one or more of the immunoconjugates specified herein, and

15 administering said immunoconjugate in a therapeutically effective amount, wherein said IgG4 isotype alleviates ADCC, complement dependent cytotoxicity and/or Fc-mediated targeting of hepatic FcR.

[0026] The present description is also directed at a method for inhibiting, delaying and/or preventing the growth of tumor cells in a cell culture comprising

15 administering to said cell culture a growth of tumor cells inhibiting, delaying and/or preventing effective amount of one or more of the immunoconjugates specified herein. The effective amount may induce cell death or continuous cell cycle arrest in CD138 expressing tumor cells and, optionally, auxillary cells that do not express CD138, in particular tumor stroma cells. The cells in said cell culture may be obtained from a cancer patient and, after administration of said effective amount of said immunoconjugate, the cells of said cell culture may be reimplanted into said cancer patient.

20 [0027] The present description is also directed at a method for inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient in need thereof, comprising administering to said patient at least one or more of the immunoconjugates specified above in a growth of said tumor and/or spreading of said tumor cells inhibiting or reducing amount,

wherein said immunoconjugate inhibits, delays or prevents the growth and/or spread of said tumor cells.

25 [0028] The effector molecule of said immunoconjugate(s) may be a toxin, cytotoxic enzyme, low molecular weight cytotoxic drug, a pore-forming agent, biological response modifier, prodrug activating enzyme, an antibody, cytokine or a radionuclide.

[0029] Immunoconjugates of the present invention may be administered in a single dose of 5 mg/m<sup>2</sup> to about 300 mg/m<sup>2</sup>, optionally at hourly, daily, weekly intervals or combinations thereof.

30 [0030] Multiple dose regimes include, hourly, daily and weekly regimes are part of the present invention and include in particular administration at intervals of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 hours, 1, 2, 3, 4, 5, 6, 7 days, 1, 2, 3, 4, 5, 6, 7 or 8 weeks.

[0031] The present description is also directed at a method for inhibiting, delaying and/or preventing the growth of a tumor and/or spread of malignant tumor cells in a patient in need thereof, comprising

35 (a) administering to said patient one or more cytotoxic agents and/or radiation in an amount effective to reduce tumor load; and

(b) administering to said patient at least one of the immunconjugates specified herein in a growth of a tumor and/or spreading of tumor cells inhibiting, delaying or preventing amount,

40 wherein said immunoconjugate inhibits, delays or prevents the growth and/or spread of tumor cells comprising CD138 expressing cells.

[0032] The cytotoxic agent may, in particular, be mephalan, vincristine, doxorubicin, dexamethasone, cyclophosphamide, etoposide, cytarabine, cisplatin, thalidomide, prednisone, thalidomide, bortezomib, lenalidomide, sorafenib, romidepsin or combinations thereof or may be antibody based.

45 [0033] The present description is also directed at a method for treating a subject having a condition that would benefit from the suppression of myeloma cell survival, the method comprising:

(a) providing at least one of any of the immunoconjugates specified herein, and

50 (b) administering the immunoconjugate to the subject to selectively decrease survival or growth of said myeloma cells of said subject.

[0034] The present invention is also directed at a pharmaceutical composition comprising any of the immunoconjugates of the invention for the inhibition, delay and/or prevention of the growth of tumors and/or spread of tumor cells, and one or more pharmaceutically acceptable excipients.

55 [0035] The pharmaceutical composition may include cytotoxic agents as specified herein.

[0036] The present invention is also directed at a kit comprising, in separate containers, pharmaceutical compositions for use in combination to inhibit, delay and/or prevent the growth of tumors and/or spread of tumor cells, wherein one container comprises an effective amount of the above pharmaceutical composition, and wherein, a separate container

comprises a second pharmaceutical composition comprising an effective amount of an agent, preferably a cytotoxic agent, for the inhibition, delay and/or prevention of the growth of tumors and/or spread of tumor cells, and one or more pharmaceutically acceptable excipients.

5 [0037] The present description is also directed at a method for inhibiting, delaying and/or preventing growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a subject in need thereof, comprising

(a) providing an immunoconjugate comprising:

10 an engineered targeting antibody against CD138 attached to an effector molecule via a cleavable linker, wherein said effector molecule is sterically hindered, and

15 (b) administering to said subject the immunoconjugate of (a) in a growth of said tumor and/or spreading of said tumor cells inhibiting, delaying and/or preventing amount, wherein said immunoconjugate of (a) provides a growth of a tumor inhibiting activity that exceeds that of its unhindered counterpart by about 10%, about 20%, about 30%, about 40% or more.

[0038] A growth of a tumor inhibiting activity of an unhindered counterpart comprising a non-cleavable linker may exceed that of the growth of a tumor inhibiting activity of its unhindered counterpart comprising a cleavable linker, such as by at least about 5%, at least about 10%, up to about 15%.

20 [0039] Said engineered targeting antibody against CD138 may consist essentially of antigen binding region against CD138 of a non-human antibody or may comprise an antigen binding region against CD138 of a non-human antibody and a further antibody region, wherein at least part of said further antibody region is of a human antibody.

25 [0040] Said cleavable linker may comprise a disulfide bond. The effector molecule may be DM4. The immunoconjugate may be part of a pharmaceutical composition and may be administered to the subject in at least one dose in an amount from about 5 mg/m<sup>2</sup> to about 300 mg/m<sup>2</sup>.

[0041] The present description provides an immunoconjugate for use as a medicament wherein the immunoconjugate comprises:

30 (a) an engineered targeting antibody

35 (i) consisting essentially of antigen binding region against CD138 of a non-human antibody, or  
 (ii) comprising an antigen binding region against CD138, wherein said antigen binding region is of a non-human antibody,  
 a further antibody region, wherein at least part of said further antibody region is of a human antibody, and

35 (b) an effector molecule,

wherein said immunoconjugate homogenously binds to CD138.

40 [0042] The present description provides a further immunoconjugate for use as a medicament comprising:

45 a targeting agent targeting CD138 comprising  
 an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof, wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1.

50 [0043] In particular, in one aspect of the description the immunoconjugate of the above paragraph is for use in the treatment of multiple myeloma. In particular, the immunoconjugate can be used for the manufacture of a medicament for the treatment of multiple myeloma.

55 [0044] The present description further provides an immunoconjugate for use in immunoconjugate mediated drug delivery to a patient, in particular for alleviation of ADCC, complement-dependent cytotoxicity and/or Fc-mediated targeting of hepatic FcR, wherein the immunoconjugate comprises a targeting agent targeting CD138 comprising an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof, wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1, and wherein a constant region of said immunoglobulin heavy chain or part thereof is an IgG4 isotype constant region.

[0045] The present description also provides tumor cells for use in the treatment of cancer in a patient wherein the tumor cells have been treated in cell culture with an immunoconjugate comprising:

(a) an engineered targeting antibody

(i) consisting essentially of antigen binding region against CD138 of a non-human antibody, or  
(ii) comprising an antigen binding region against CD138, wherein said antigen binding region is of a non-human antibody,  
a further antibody region, wherein at least part of said further antibody region is of a human antibody, and

5

(b) an effector molecule,

wherein said immunoconjugate homogenously binds to CD138.

10 [0046] The present description also provides tumor cells for use in the treatment of cancer in a patient wherein the tumor cells have been treated in cell culture with an immunoconjugate comprising:

a targeting agent targeting CD138 comprising  
an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof,  
wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1.

15

[0047] The present description provides an immunoconjugate for use in inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient, wherein the immunoconjugate comprises:

20

(a) an engineered targeting antibody

(i) consisting essentially of antigen binding region against CD138 of a non-human antibody, or  
(ii) comprising an antigen binding region against CD138, wherein said antigen binding region is of a non-human antibody,

25

a further antibody region, wherein at least part of said further antibody region is of a human antibody, and

(b) an effector molecule,

wherein said immunoconjugate homogenously binds to CD138.

30 [0048] Alternatively, the present description provides an immunoconjugate for use in inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient, wherein the immunoconjugate comprises:

35

a targeting agent targeting CD138 comprising

an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof,  
wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1.

40

[0049] Still further, the present description provides a medicament comprising an immunoconjugate and one or more cancer drugs as a combined preparation for simultaneous, separate or sequential use in the treatment of tumor cells comprising CD138 expressing cells, wherein the immunoconjugate comprises:

(a) an engineered targeting antibody

(i) consisting essentially of antigen binding region against CD138 of a non-human antibody, or  
(ii) comprising an antigen binding region against CD138, wherein said antigen binding region is of a non-human antibody,

a further antibody region, wherein at least part of said further antibody region is of a human antibody, and

(b) an effector molecule,

50

wherein said immunoconjugate homogenously binds to CD138,

and wherein the one or more cancer drugs are capable of reducing the tumor load.

[0050] Alternatively, the present description provides a medicament comprising an immunoconjugate and one or more cancer drugs as a combined preparation for simultaneous, separate or sequential use in the treatment of tumor cells comprising CD138 expressing cells, wherein the immunoconjugate comprises:

a targeting agent targeting CD138 comprising

an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof,

wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1,

and wherein the one or more cancer drugs are capable of reducing the tumor load.

5 [0051] In a further aspect of the use of the above two paragraphs the combined preparation is to be administered to a patient who has been treated with radiation.

[0052] In an alternative aspect the present description provides the use of an immunoconjugate for the manufacture of a medicament for treating tumor cells in a patient comprising CD138 expressing cells, wherein the immunoconjugate comprises:

10 (a) an engineered targeting antibody

- (i) consisting essentially of antigen binding region against CD138 of a non-human antibody, or
- (ii) comprising an antigen binding region against CD138, wherein said antigen binding region is of a non-human antibody,

15 a further antibody region, wherein at least part of said further antibody region is of a human antibody, and

(b) an effector molecule,

wherein said immunoconjugate homogenously binds to CD138,

20 and wherein the medicament is to be administered to a patient treated with radiation to reduce the tumor load.

[0053] Still further the present description provides the use of an immunoconjugate for the manufacture of a medicament for treating tumor cells in a patient comprising CD138 expressing cells, wherein the immunoconjugate comprises:

a targeting agent targeting CD138 comprising

25 an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof, wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1,

and wherein the medicament is to be administered to a patient treated with radiation to reduce the tumor load.

[0054] In the above paragraphs, the medicament is capable of inhibiting, delaying and/or preventing the growth of a tumor and/or spread of malignant tumor cells in a patient.

[0055] Further the present description provides an immunoconjugate for suppression of myeloma cell survival in an individual wherein the immunoconjugate comprises:

30 (a) an engineered targeting antibody

35 (i) consisting essentially of antigen binding region against CD138 of a non-human antibody, or

- (ii) comprising an antigen binding region against CD138, wherein said antigen binding region is of a non-human antibody,

40 a further antibody region, wherein at least part of said further antibody region is of a human antibody, and

(b) an effector molecule,

wherein said immunoconjugate homogenously binds to CD138.

[0056] Still further the present description provides an immunoconjugate for suppression of myeloma cell survival in an individual wherein the immunoconjugate comprises:

45 a targeting agent targeting CD138 comprising

an isolated polypeptide comprising an amino acid sequence of an immunoglobulin heavy chain or part thereof, wherein said immunoglobulin heavy chain or part thereof has at least 70% sequence identity with SEQ ID NO:1.

50 [0057] In the above two paragraphs the immunoconjugate is, in particular, capable of selectively decreasing the survival or growth of said myeloma cells in the individual.

[0058] Further, the present description provides an immunoconjugate for use in inhibiting, delaying and/or preventing growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a subject wherein the immunoconjugate comprises an engineered targeting antibody against CD138 attached to an effector molecule via a cleavable linker, wherein said effector molecule is sterically hindered.

[0059] In the above paragraph, the immunoconjugate is, in particular, capable of providing a tumor growth inhibiting activity that exceeds that of its unhindered counterpart by about 10%, about 20%, about 30%, about 40% or more.

## BRIEF DESCRIPTION OF THE FIGURES

## [0060]

5 **FIG. 1** provides a schematic representation of nBT062 having effector molecules attached.  
**FIG. 2** is a chemical representation of BT062.  
**FIG. 3** shows the conversion of ansamitocin P-3 to maytansinol (stereochemistry is omitted for simplicity).  
**FIG. 4** shows a representative synthesis scheme of DM4.  
**FIG. 5** is a schematic representation of an antibody conjugation (nBT062 to DM4).  
10 **FIG. 6** shows an analysis of the binding of nBT062-SPDB-DM4, nBT062-SPP-DM1, nBT062-SMCC-DM1 and nBT062 antibody to OPM-2 cells. Different concentrations of nBT062 and conjugates were given to the cells and mean fluorescence was measured by FACS analysis.  
**FIG. 7(A)-(D)** depict *in vitro* cytotoxicity of nBT062-DMx conjugates towards MOLP-8 (CD138<sup>+</sup>) and BJAB (CD138<sup>-</sup>) cells. The cells were cultured in flat bottom plates and incubated with the indicated concentrations of immunoconjugates for 5 days. WST reagent was added for further 3 hours to assess cell viability. In (D) cytotoxic activity of nBT062-SPDB-DM4 was analyzed in the presence or absence of blocking antibody (1  $\mu$ M nBT062).  
15 **FIG. 8** shows tumor volumes for individual mice treated with (A) PBS, (B) nBT062 antibody, (C) free DM4 or (D) non-targeting conjugate huC242-DM4 over time (days) post-inoculation with MOLP-8 tumor cells.  
**FIG. 9** shows tumor volumes for individual mice treated with (A) PBS, (B) nBT062-SPDB-DM4, (C) B-B4-SPP-DM1 or (D) nBT062-SPP-DM1 over time (days) post-inoculation with MOLP-8 tumor cells.  
20 **FIG. 10** depicts mean tumor volume (+/- SD) of MOLP-8 human multiple myeloma xenografts in CB.17 SCID mice overtime (days) post-inoculation.  
**FIGS. 11A and B** show the anti-tumor activity of nBT062-DMx against CD138<sup>+</sup> MOLP-8 tumor cells in a bulky MOLP-8 tumor model in SCID mice. Tumor volume is given as mean (+/- SD) for each group.  
25 **FIG. 12** is a graph reflecting the anti-tumour efficacy of nBT062 containing DMx conjugates in the SCIDhu/INA-6 model towards multiple myeloma cells in the environment of human bone marrow. Soluble human IL-6 Receptor produced by multiple myeloma cells (shull-6R) was used as an indicator for tumor burden. Triangle: nBT062-SPP-DM1, Square: nBT062-SPDB-DM4; Diamond: vehicle control.  
30 **FIG. 13** shows nBT062-SPDB-DM4 mediated bystander killing *in vitro*. CD138 positive OPM2 cells and CD138 negative Namawla cells were cultured with nBT062-SPDB-DM4 at different concentrations and cell viability was measured. OD<sub>450</sub> values represent a measure for cell viability.

## DETAILED DESCRIPTION OF VARIOUS AND PREFERRED EMBODIMENTS OF THE INVENTION

35 **[0061]** The present invention relates to immunoconjugates comprising CD138 targeting antibodies and the delivery of the effector molecule(s) of the immunoconjugates to target sites and the site specific release of effector(s) molecule in, at or near target cells, tissues and organs. More particularly, the present invention relates to immunoconjugates comprising CD138 targeting antibodies and potent effector molecules that are attached to the targeting agent. The effector molecules may be activated by cleavage/dissociation from the targeting agent portion of the immunoconjugate at the target site.  
40 **[0062]** The immunoconjugates according to the present invention may be administered to a subject in need of therapeutic treatment. The effector molecule or molecules may be released from the immunoconjugate by cleavage/dissociation in, at or close to the target cell, tissue or organ.  
45 **[0063]** In one example, the immunoconjugate comprises the antibody nBT062, which targets CD138 expressing cells, and at least one highly cytotoxic drug or toxin as an effector molecule, is administered to a patient with cancer. In this example, a therapeutically effective amount of the immunoconjugate is administered intravenously to a patient so that it concentrates in the cancer cells. The effector molecule or molecules are then released from the antibody by natural means. After or during cleavage the effector molecule may be stabilized by alkylation and may diffuse to surrounding auxillary cells such as stroma cells that do not express CD138.  
50 **[0064]** In a second example, the immunoconjugate comprises the antibody nBT062, which targets CD 138 expressing cells, and at least one highly cytotoxic drug or toxin as an effector molecule, and an additional cytotoxic agent is administered to a patient with cancer. In this example, a therapeutically effective amount of the immunoconjugate and the cytotoxic agent are co-administered intravenously to a patient so that it concentrates in the cancer cells. The cytotoxic agent destroys more than 50% of the CD138 expressing cancer cells, but the immunconjugate attaches efficiently to further CD138 expressing cancer cells. The effector molecule or molecules are released from the antibody by natural means. After or during cleavage, the effector molecule may be stabilized by alkylation and may diffuse to surrounding auxillary cells such as stroma cells that do not express CD138.  
55 **[0065]** In a third example, the immunoconjugate comprises the antibody nBT062 and at least one highly cytotoxic drug

or toxin and is administered to a cell population isolated from a patient with cancer. In this example, a cell death or continuous cell cycle arrest inducing amount of the immunoconjugate is administered to the cell population so that it concentrates in the cancerous cells. The effector molecule or molecules are released from the targeting antibody by natural means or external means to induce cell death or continuous cell cycle arrest in the cancer cells.

5 [0066] In a fourth example, the immunoconjugate comprises the antibody nBT062 and at least one highly cytotoxic drug or toxin as an effector molecule and is administered to a patient with cancer. In this example, a therapeutically effective amount of the immunoconjugate is administered intravenously to a patient so that it concentrates in the cancerous cells. The effector molecule or molecules are released from the antibody target by an external means to induce cell death or continuous cell cycle arrest in the cancer cells.

10 [0067] CD138 or syndecan-1 (also described as SYND1; SYNDECAN; SDC; SCD1; CD138 ANTIGEN, SwissProt accession number: P18827 human) is a membrane glycoprotein that was originally described to be present on cells of epithelial origin, and subsequently found on hematopoietic cells (Sanderson, 1989). CD138 has a long extracellular domain that binds to soluble molecules (e.g., the growth factors EGF, FGF, HGF) and to insoluble molecules (e.g., to the extracellular matrix components collagen and fibronectin) through heparan sulfate chains (Langford, 1998; Yang, 15 2007) and acts as a receptor for the extracellular matrix. CD138 also mediates cell to cell adhesion through heparin-binding molecules expressed by adherent cells. It has been shown that CD138 has a role as a co-receptor for growth factors of myeloma cells (Bisping, 2006). Studies of plasma cell differentiation showed that CD138 must also be considered as a differentiation antigen (Bataille, 2006).

20 [0068] In malignant hematopoiesis, CD138 is highly expressed on the majority of MM cells, ovarian carcinoma, kidney carcinoma, gall bladder carcinoma, breast carcinoma, prostate cancer, lung cancer, colon carcinoma cells and cells of Hodgkin's and non-Hodgkin's lymphomas, chronic lymphocytic leukemia (CLL) (Horvathova, 1995), acute lymphoblastic leukemia (ALL), acute myeloblastic leukemia (AML) (Seftalioglu, 2003 (a); Seftalioglu, 2003 (b)), solid tissue sarcomas, colon carcinomas as well as other hematologic malignancies and solid tumors that express CD 138 (Carbone et al., 1999; Sebestyen et al., 1999; Han et al., 2004; Charnaux et al., 2004; O'Connell et al., 2004; Orosz and Kopper, 2001).

25 [0069] Other cancers that have been shown to be positive for CD138 expression are many ovarian adenocarcinomas, transitional cell bladder carcinomas, kidney clear cell carcinomas, squamous cell lung carcinomas; breast carcinomas and uterine cancers (see, for example, Davies et al., 2004; Barbareschi et al., 2003; Mennerich et al., 2004; Anttonen et al., 2001; Wijdenes, 2002).

30 [0070] In the normal human hematopoietic compartment, CD138 expression is restricted to plasma cells (Wijdenes, 1996; Chilosi, 1999) and CD138 is not expressed on peripheral blood lymphocytes, monocytes, granulocytes, and red blood cells. In particular, CD34<sup>+</sup> stem and progenitor cells do not express CD138 and anti-CD138 mAbs do not affect the number of colony forming units in hematopoietic stem cell cultures (Wijdenes, 1996). In non-hematopoietic compartments, CD138 is mainly expressed on simple and stratified epithelia within the lung, liver, skin, kidney and gut. Only a weak staining was seen on endothelial cells (Bernfield, 1992; Vooijs, 1996). It has been reported that CD138 exists in 35 polymorphic forms in human lymphoma cells (Gattei, 1999).

40 [0071] Monoclonal antibodies B-B4, BC/B-B4, B-B2, DL-101, 1 D4, MI15, 1.BB.210, 2Q1484, 5F7, 104-9, 281-2 in particular B-B4 have been reported to be specific to CD138. Of those B-B4, 1 D4 and MI15 recognized both the intact molecule and the core protein of CD138 and were shown to recognize either the same or closely related epitopes (Gattei, 1999). Previous studies reported that B-B4 did not recognize soluble CD138, but only CD138 in membrane bound form (Wijdenes, 2002).

45 [0072] B-B4, a murine IgG1 mAb, binds to a linear epitope between residues 90-95 of the core protein on human syndecan-1 (CD138) (Wijdenes, 1996; Dore, 1998). Consistent with the expression pattern of CD138, B-B4 was shown to strongly react with plasma cell line RPMI8226, but not to react with endothelial cells. Also consistent with the expression pattern of CD138, B-B4 also reacted with epithelial cell lines A431 (keratinocyte derived) and HepG2 (hepatocyte derived). An immunotoxin B-B4-saporin was also highly toxic towards the plasma cell line RPMI8226, in fact considerably more toxic than free saporin. However, from the two epithelial cell lines tested, B-B4-saporin showed only toxicity towards cell line A431, although in a clonogenic assay B-B4 saporin showed no inhibitory effect on the outgrowth of A431 cells (Vooijs, 1996). Other researchers reported lack of specificity of MM-associated antigens against tumors (Couturier, 1999).

50 [0073] An antibody/immunoconjugate "consisting essentially of" certain components means in the context of the present description that the antibody/immunoconjugate consists of the specified components and any additional materials or components that do not materially affect the basic characteristics of the antibody.

[0074] The present invention uses the term "tumor cell" to include cancer cells as well as pre-cancerous cells which may or may not form part of a solid tumor.

55 [0075] A "targeting agent" according to the present description is able to associate with a molecule expressed by a target cell and includes peptides and non-peptides. In particular, targeting agents according to the present description include targeting antibodies and non-immunoglobulin targeting molecules, which may be based on non-immunoglobulin proteins, including, but not limited to, AFFILIN® molecules, ANTICALINS® and AFFIBODIES®. Non-immunoglobulin targeting molecules also include non-peptidic targeting molecules such as targeting DNA and RNA oligonucleotides

(aptamers), but also physiological ligands, in particular ligands of the antigen in question, such as CD138.

**[0076]** A "targeting antibody" according to the present description is or is based on a natural antibody or is produced synthetically or by genetic engineering and binds to an antigen on a cell or cells (target cell(s)) of interest. A targeting antibody according to the present description includes a monoclonal antibody, a polyclonal antibody, a multispecific antibody (for example, a bispecific antibody), or an antibody fragment. The targeting antibody may be engineered to, for example, improve its affinity to the target cells (Ross, 2003) or diminish its immunogenicity. The targeting antibody may be attached to a liposomal formulation including effector molecules (Carter, 2003). An antibody fragment comprises a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments according to the present invention include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments, but also diabodies; domain antibodies (dAb) (Ward, 1989; United States Patent 6,005,079); linear antibodies; single-chain antibody molecules; and multispecific antibodies formed from antibody fragments. In a single chain variable fragment antibody (scFv) the heavy and light chains (VH and VL) can be linked by a short amino acid linker having, for example, the sequence (glycine<sub>4</sub>serine)<sub>n</sub>, which has sufficient flexibility to allow the two domains to assemble a functional antigen binding pocket. Addition of various signal sequences may allow for more precise targeting of the targeting antibody. Addition of the light chain constant region (CL) may allow dimerization via disulphide bonds, giving increased stability and avidity. Variable regions for constructing the scFv can, if a mAb against a target of interest is available, be obtained by RT-PCR which clones out the variable regions from mRNA extracted from the parent hybridoma. Alternatively, the scFv can be generated de novo by phage display technology (Smith, 2001). As used herein, the term "functional fragment", when used in reference to a targeting antibody, is intended to refer to a portion of the targeting antibody which is capable of specifically binding an antigen that is specifically bound by the antibody reference is made to. A bispecific antibody according to the present invention may, for example, have at least one arm that is reactive against a target tissue and one arm that is reactive against a linker moiety (United States Patent Publication 20020006379). A bispecific antibody according to the present description may also bind to more than one antigen on a target cell (Carter, 2003). An antibody according to the present invention may be modified by, for example, introducing cysteine residues to introduce thiol groups (Olafsen, 2004).

**[0077]** In accordance with the present description, the targeting antibody may be derived from any source and may be, but is not limited to, a camel antibody, a murine antibody, a chimeric human/mouse antibody or a chimeric human/monkey antibody, in particular, a chimeric human/mouse antibody such as nBT062.

**[0078]** Humanized antibodies are antibodies that contain sequences derived from a human-antibody and from a non-human antibody and are also within the scope of the present invention. Suitable methods for humanizing antibodies include CDR-grafting (complementarity determining region grafting) (EP 0 239 400; WO 91/09967; United States Patents 5,530,101; and 5,585,089), veneering or resurfacing (EP 0 592 106; EP 0 519 596; Padlan, 199; Studnicka et al., 1994; Roguska et al., 1994), chain shuffling (United States Patent 5,565,332) and Delmmunosation™ (Biovation, LTD). In CDR-grafting, the mouse complementarity-determining regions (CDRs) from, for example, mAb B-B4 are grafted into human variable frameworks, which are then joined to human constant regions, to create a human B-B4 antibody (hB-B4). Several antibodies humanized by CDR-grafting are now in clinical use, including MYLOTARG (Sievers et al., 2001) and HECEPTIN (Pegram et al., 1998).

**[0079]** The resurfacing technology uses a combination of molecular modeling, statistical analysis and mutagenesis to alter the non-CDR surfaces of antibody variable regions to resemble the surfaces of known antibodies of the target host. Strategies and methods for the resurfacing of antibodies, and other methods for reducing immunogenicity of antibodies within a different host, are disclosed, for example, in United States Patent 5,639,641. Human antibodies can be made by a variety of methods known in the art including phage display methods. See also United States Patents 4,444,887, 4,716,111, 5,545,806, and 5,814,318; and international patent application publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741.

**[0080]** Targeting antibodies that have undergone any non-natural modification such as chimeric human/mouse antibodies or a chimeric human/monkey antibodies, humanized antibodies or antibodies that were engineered to, for example, improve their affinity to the target cells or diminish their immunogenicity but also antibody fragments, in particular functional fragments of such targeting antibodies that have undergone any non-natural modification, diabodies; domain antibodies; linear antibodies; single-chain antibody molecules; and multispecific antibodies are referred to herein as **engineered targeting antibodies**.

**[0081]** Chimerized antibodies, maintain the antibody binding region (ABR or Fab region) of the non-human antibody, e.g., the murine antibody they are based on, while any constant regions may be provided for by, e.g., a human antibody. Generally, chimerization and/or the exchange of constant regions of an antibody will not affect the affinity of an antibody because the regions of the antibody which contribute to antigen binding are not affected by this exchange. In a preferred embodiment of the present invention, the engineered, in particular chimerized, antibody of the present invention, may have a higher binding affinity (as expressed by  $K_D$  values) than the respective non-human antibody it is based on. In particular, the nBT062 antibody and antibodies based thereon may have higher antibody affinity than the murine B-B4. In another preferred embodiment of the present invention, immunoconjugates comprising those engineered/chimerized antibodies also display this higher antibody affinity. These immunoconjugates may also display in certain embodiments

other advantageous properties, such as a higher reduction of tumor load than their B-B4 containing counterparts. In a preferred embodiment, the engineered, in particular chimerized targeting antibodies display binding affinities that are characterized by dissociation constants  $K_D$  (nM) of less than 1.6, less than 1.5 or about or less than 1.4, while their murine counterparts are characterized by dissociation constants  $K_D$  (nM) of about or more than 1.6. Immunoconjugates comprising targeting agents such as targeting antibodies may be characterized by dissociation constants of  $K_D$  (nM) of less than 2.6, less than 2.5, less than 2.4, less than 2.3, less than 2.2, less than 2.1, less than 2.0, less than or about 1.9 are preferred, while immunoconjugates comprising the murine counterpart antibodies may be characterized by dissociation constants  $K_D$  (nM) of about or more than 2.6 (compare Table 3, Materials and Methods).

**[0082]** Fully human antibodies may also be used. Those antibodies can be selected by the phage display approach, where CD138 or an antigenic determinant thereof is used to selectively bind phage expressing, for example, B-B4 variable regions (see, Krebs, 2001). This approach is advantageously coupled with an affinity maturation technique to improve the affinity of the antibody. All antibodies referred to herein are isolated antibodies.

**[0083]** In one embodiment, the targeting antibody is, in its unconjugated form, moderately or poorly internalized. Moderate internalization constitutes about 30% to about 75% internalization of antibody, poor internalization constitutes about 0.01% to up to about 30% internalization after 3 hours incubation at 37°C. In another preferred embodiment the targeting antibody binds to CD138, for example, antibodies B-B4, BC/B-B4, B-B2, DL-101, 1D4, MI15, 1.BB.210, 2Q1484, 5F7, 104-9, 281-2 in particular B-B4. Hybridoma cells, which were generated by hybridizing SP02/0 myeloma cells with spleen cells of Balb/c mice have been deposited with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1, D-38124 Braunschweig on December 11, 2007. The identification number of these B-B4 expressing hybridoma cells is DSM ACC2874. In another embodiment, the targeting antibody does not substantially bind non-cell-surface expressed CD138. When, in the context of the present invention, the name of a specific antibody is combined with the term "targeting antibody" such as "nBT062 targeting antibody," this means that this targeting antibody has the binding specificity of the antibody nBT062. If a targeting antibody is said to be "based on" a specified antibody, this means that this targeting antibody has the binding specificity of this antibody, but might take any form consistent with the above description of a targeting antibody. When, in the context of the present invention, the name of a specific antigen is combined with the term "targeting antibody" such as "CD138 targeting antibody," this means that this targeting antibody has the binding specificity for CD138. If, in the context of the present description, for example, a targeting antibody is said to do something "selectively" such as "selectively targeting cell-surface expressed CD138" or, to be "selective" for something, this means that there is a significant selectivity (i.e. a higher affinity towards CD138-positive cells compared with CD138-negative cells) for, in the case of the example provided, cell-surface expressed CD138, compared to any other antigens. Adverse side effects in a given environment are substantially reduced or even avoided due to this selectivity.

**[0084]** "Non-immunoglobulin targeting molecules" according to the present description include targeting molecules derived from non-immunoglobulin proteins as well as non-peptidic targeting molecules. Small non-immunoglobulin proteins which are included in this definition are designed to have specific affinities towards, in particular surface expressed CD138. These small non-immunoglobulin proteins include scaffold based engineered molecules such as Affilin® molecules that have a relatively low molecular weight such as between 10 kDa and 20 kDa. Appropriate scaffolds include, for example, gamma crystalline. Those molecules have, in their natural state, no specific binding activity towards the target molecules. By engineering the protein surfaces through locally defined randomization of solvent exposed amino acids, completely new binding sites are created. Former non-binding proteins are thereby transformed into specific binding proteins. Such molecules can be specifically designed to bind a target, such as CD138, and allow for specific delivery of one or more effector molecules (see, scil Proteins GmbH at [www.scilproteins.com](http://www.scilproteins.com), 2004). Another kind of non-immunoglobulin targeting molecules are derived from lipocalins, and include, for example ANTICALINS®, which resemble in structure somewhat immunoglobulins. However, lipocalins are composed of a single polypeptide chain with 160 to 180 amino acid residues. The binding pocket of lipocalins can be reshaped to recognize a molecule of interest with high affinity and specificity (see, for example, Beste et al., 1999). Artificial bacterial receptors such as those marketed under the trademark Affibody® (Affibody AB) are also within the scope of the present description. These artificial bacterial receptor molecules are small, simple proteins and may be composed of a three-helix bundle based on the scaffold of one of the IgG-binding domains of Protein A (*Staphylococcus aureus*). These molecules have binding properties similar to many immunoglobulins, but are substantially smaller, having a molecular weight often not exceeding 10kDa and are also comparatively stable. Suitable artificial bacterial receptor molecules are, for example, described in United States Patents 5,831,012; 6,534,628 and 6,740,734.

**[0085]** Other "non-immunoglobulin targeting molecules" are physiological ligands of the antigen in question. Physiological ligands of CD138 include for example, but not limited to, ADAMTS4 (aggrecanase-1), antithrombin-3, bFGF, cathepsin G, CCL5 (RANTES), CCL7, CCL11, CCL17, CD44, collagens (collagen type 1, collagen type 2, collagen type 3, collagen type 4, collagen type 5, collagen type 6), CXCL1, elastase, gp120, HGF [hepatocyte growth factor], laminin-1, laminin-2, laminin-5, midkine, MMP-7, neutrophil elastase, and pleiotrophin (HBNF, HBGF-8). Non-peptidic targeting molecules include, but are not limited to, to DNA and RNA oligonucleotides that bind to CD138 (aptamers).

[0086] An "effector molecule" according to the present invention is a molecule or a derivative, or an analogue thereof that is attached to an engineered targeting antibody, and that exerts a desired effect, for example, apoptosis, or another type of cell death, or a continuous cell cycle arrest on the target cell or cells. Effector molecules according to the present description include molecules that can exert desired effects in a target cell and include, but are not limited to, toxins, drugs, in particular low molecular weight cytotoxic drugs, radionuclides, biological response modifiers, pore-forming agents, ribonucleases, proteins of apoptotic signaling cascades with apoptosis-inducing activities, cytotoxic enzymes, prodrug activating enzymes, antisense oligonucleotides, antibodies or cytokines as well as functional derivatives or analogues/fragments thereof. Toxins may include bacterial toxins, such as, but not limited to, Diphtheria toxin or Exotoxin A, plant toxins, such as but not limited to, Ricin. Proteins of apoptotic signaling cascades with apoptosis-inducing activities, include, but are not limited to, Granzyme B, Granzyme A, Caspase-3, Caspase-7, Caspase-8, Caspase-9, truncated Bid (tBid), Bax and Bak.

[0087] In a preferred embodiment, the effector increases internal effector delivery of the immunoconjugate, in particular when the natural form of the antibody on which the targeting antibody of the immunoconjugate is based is poorly internalizable. In another preferred embodiment the effector is, in its native form, non-selective. In certain embodiments the effector has high non-selective toxicity, including systemic toxicity, when in its native form. The "native form" of an effector molecule of the present invention is an effector molecule before being attached to the targeting agent to form an immunoconjugate. In another preferred embodiment, the non-selective toxicity of the effector molecule is substantially eliminated upon conjugation to the targeting agent. In another preferred embodiment, the effector molecule causes, upon reaching the target cell, death or continuous cell cycle arrest in the target cell. A drug-effector molecule according to the present invention includes, but is not limited to, a drug including, for example, small highly cytotoxic drugs that act as inhibitors of tubulin polymerization such as maytansinoids, dolastatins, auristatin and cryptophycin; DNA alkylating agents like CC-1065 analogues or derivatives (United States Patents 5,475,092; 5,585,499; 6,716,821) and duocarmycin; enediyne antibiotics such as calicheamicin and esperamicin; and potent taxoid (taxane) drugs (Payne, 2003). Maytansinoids and calicheamicins are particularly preferred. An effector maytansinoid includes maytansinoids of any origin, including, but not limited to synthetic maytansinol and maytansinol analogue and derivative. Doxorubicin, daunomycin, methotrexate, vinblastine, neocarzinostatin, macromycin, trenimon and  $\alpha$ -amanitin are some other effector molecules within the scope of the present invention. Also within the scope of the present invention are antisense DNA molecules as effector molecules. When the name of, for example, a specific drug or class of drugs is combined herein with the term "effector" or "effector molecule," reference is made to an effector of an immunoconjugate according to the present invention that is based on the specified drug or class of drugs.

[0088] Maytansine is a natural product originally derived from the Ethiopian shrub *Maytenus serrata* (Remillard, 1975; United States Patent 3,896,111). This drug inhibits tubulin polymerization, resulting in mitotic block and cell death (Remillard, 1975; Bhattacharyya, 1977; Kupchan, 1978). The cytotoxicity of maytansine is 200-1000-fold higher than that of anti-cancer drugs in clinical use that affect tubulin polymerization, such as Vinca alkaloids or taxol. However, clinical trials of maytansine indicated that it lacked a therapeutic window due to its high systemic toxicity. Maytansine and maytansinoids are highly cytotoxic but their clinical use in cancer therapy has been greatly limited by their severe systemic side-effects primarily attributed to their poor selectivity for tumors. Clinical trials with maytansine showed serious adverse effects on the central nervous system and gastrointestinal system.

[0089] Maytansinoids have also been isolated from other plants including seed tissue of *Trewia nudiflora* (United States Patent 4,418,064)

[0090] Certain microbes also produce maytansinoids, such as maytansinol and C-3 maytansinol esters (United States Patent 4,151,042).

[0091] The present invention is directed to maytansinoids of any origin, including synthetic maytansinol and maytansinol analogues which are disclosed, for example, in United States Patents 4,137,230; 4,248,870; 4,256,746; 4,260,608; 4,265,814; 4,294,757; 4,307,016; 4,308,268; 4,308,269; 4,309,428; 4,313,946; 4,315,929; 4,317,821; 4,322,348; 4,331,598; 4,361,650; 4,362,663; 4,364,866; 4,371,533; 4,424,219 and 4,151,042.

[0092] In a preferred embodiment, the maytansinoid is a thiol-containing maytansinoid and is more preferably produced according to the processes disclosed in United States Patent 6,333,410 to Chari et al or in Chari et al.(Chari, 1992).

[0093] DM-1 (N<sup>2</sup>-deacetyl-N<sup>2</sup>-(3-mercaptopro-1-oxopropyl)-maytansine) is a preferred effector molecule in the context of the present invention. DM1 is 3- to 10-fold more cytotoxic than maytansine, and has been converted into a pro-drug by linking it via disulfide bond(s) to a monoclonal antibody directed towards a tumor-associated antigen. Certain of these conjugates (sometimes called "tumor activated prodrugs" (TAPs)) are not cytotoxic in the blood compartment, since they are activated upon associating with a target cells and internalized, thereby releasing the drug (Blartler, 2001). Several antibody-DM1 conjugates have been developed (Payne, 2003), and been evaluated in clinical trials. For example, huC242-DM1 treatment in colorectal cancer patients was well tolerated, did not induce any detectable immune response, and had a long circulation time (Tolcher, 2003).

[0094] Other particularly preferred maytansinoids comprise a side chain that contains a sterically hindered thiol bond such as, but not limited to, maytansinoids N<sup>2</sup>-deacetyl- N<sup>2</sup>-(4-mercaptopro-1-oxopentyl)-maytansine, also referred to as

"DM3," and N<sup>2</sup>'-deacetyl- N<sup>2</sup>'-(4-methyl-4-mercaptopentyl)-maytansine, also referred to as "DM4." The synthesis of DM4 is shown in FIGS. 3 and 4 and is described elsewhere herein. DM4 differs from DM1 and DM3 in that it bears methyl groups at its aC. This results in a sterical hindrance when DM4 is attached via a linker in particular, but not limited to, a linker comprising a disulfide bond, to a targeting agent such as nBT062. A wide variety of maytansinoids bearing a sterically hindered thiol group (possessing one or two substituents, in particular alkyls substituents, such as the methyl substituents of DM4) are disclosed U.S. Patent Publication 2004/0235840, published Nov. 25, 2004. The steric hindrance conferred by alkyl groups such as the methyl groups on the carbon adjacent to the sulfur atom of DM3 and DM4 may affect the rate of intracellular cleavage of the immunoconjugate. The variable alkyl unit may therefore affect potency, efficacy, and safety/toxicity *in vitro* and *in vivo*.

**[0095]** As reported by Goldmahker et al. in U.S. Patent Publication 2006/0233814, such a hindrance induces alkylation (e.g., methylation) of the free drug, once the drug is released at its target. The alkylation may increase the stability of the drug allowing for the so-called bystander effect. However, as the person skilled in the art will appreciate, other effector molecules comprising substituents such as alkyl groups at positions that result in a sterical hindrance when the effector is attached to a targeting agent via a linker are part of the present invention (U.S. Patent Publication 2004/0235840).

Preferably this hindrance induces a chemical modification such as alkylation of the free drug to increase its overall stability, which allows the drug to not only induce cell death or continuous cell cycle arrest in CD138 expressing tumor cells but, optionally, also to affect auxillary cells that, e.g., support or protect the tumor from drugs, in particular cells of the tumor stroma and the tumor vasculature and which generally do not express CD138 to diminish or lose their supporting or protecting function.

**[0096]** DNA alkylating agents are also particularly preferred as effector molecules and include, but are not limited to, CC-1065 analogues or derivatives. CC-1065 is a potent antitumor-antibiotic isolated from cultures of *Streptomyces zelensis* and has been shown to be exceptionally cytotoxic *in vitro* (United States Patent 4,169,888). Within the scope of the present invention are, for example the CC-1065 analogues or derivatives described in United States Patents 5,475,092, 5,585,499 and 5,739,350. As the person skilled in the art will readily appreciate, modified CC-1065 analogues or derivatives as described, for example, in United States Patent 6,756,397 are also within the scope of the present invention. In certain embodiments of the invention, CC-1065 analogues or derivatives may, for example, be synthesized as described in United States Patent 6,534,660.

**[0097]** Another group of compounds that make preferred effector molecules are taxanes, especially highly potent ones and those that contain thiol or disulfide groups. Taxanes are mitotic spindle poisons that inhibit the depolymerization of tubulin, resulting in an increase in the rate of microtubule assembly and cell death. Taxanes that are within the scope of the present invention are, for example, disclosed in United States Patents 6,436,931; 6,340,701; 6,706,708 and United States Patent Publications 20040087649; 20040024049 and 20030004210. Other taxanes are disclosed, for example, in United States Patent 6,002,023, United States Patent 5,998,656, United States Patent 5,892,063, United States Patent 5,763,477, United States Patent 5,705,508, United States Patent 5,703,247 and United States Patent 5,367,086. As the person skilled in the art will appreciate, PEGylated taxanes such as the ones described in United States Patent 6,596,757 are also within the scope of the present invention.

**[0098]** Calicheamicin effector molecules according to the present invention include gamma 1I, N-acetyl calicheamicin and other derivatives of calicheamicin. Calicheamicin binds in a sequence-specific manner to the minor groove of DNA, undergoes rearrangement and exposes free radicals, leading to breakage of double-stranded DNA, resulting in cell apoptosis and death. One example of a calicheamicin effector molecule that can be used in the context of the present invention is described in United States Patent 5,053,394.

**[0099]** An immunoconjugate according to the present description comprises at least one targeting agent, in particular targeting antibody and one effector molecule. The immunoconjugate might comprise further molecules for example for stabilization. For immunoconjugates, the term "conjugate" is generally used to define the operative association of the targeting agent with one or more effector molecules and is not intended to refer solely to any type of operative association, and is particularly not limited to chemical "conjugation". So long as the targeting agent is able to bind to the target site and the attached effector functions sufficiently as intended, particularly when delivered to the target site, any mode of attachment will be suitable. The conjugation methods according to the present description include, but are not limited to, direct attachment of the effector molecule to the targeting antibody, with or without prior modification of the effector molecule and/or the targeting antibody or attachment via linkers. Linkers can be categorized functionally into, for example, acid labile, photolabile linkers, enzyme cleavable linkers, such as linkers that can be cleaved by peptidases. Cleavable linkers are, in many embodiments of the invention preferred. Such cleavable linkers can be cleaved under conditions present in the cellular environment, in particular, an intracellular environment and that have no detrimental effect on the drug released upon cleavage. Low pHs such as pH of 4 to 5 as they exist in certain intracellular compartments will cleave acid labile linkers, while photolabile linkers can be cleaved by, e.g., infrared light. However, linkers that are cleaved by/under physiological conditions present in the majority of cells are preferred and are referred to herein as **physiologically cleavable linkers**. Accordingly, disulfide linkers are being preferred in many embodiments of the invention. These

linkers are cleavable through disulfide exchange, which can occur under physiological conditions. Preferred heterobifunctional disulfide linkers include, but are not limited to, N-succinimidyl 3-(2-pyridyldithio)propionate (SPDP) (see, e.g., Carlsson et al. (1978)), N-succinimidyl 4-(2-pyridyldithio)butanoate (SPDB) (see, e.g., U.S. Pat. No. 4,563,304), N-succinimidyl 4-(2-pyridyldithio)pentanoate (SPP) (see, e.g., CAS Registry number 341498-08-6), N-succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC) (see, e.g., Yoshitake et al., (1979)), and N-succinimidyl 4-methyl-4-[2-(5-nitro-pyridyl)-dithio]pentanoate (SMNP) (see, e.g., U.S. Pat. No. 4,563,304). The most preferred linker molecules for use in the inventive composition are SPP, SMCC, and SPDB.

**[0100]** Other suitable linkers may include "non-cleavable" bonds, such as, but not limited to Sulfosuccinimidyl maleimidomethyl cyclohexane carboxylate (SMCC), which is a heterobifunctional linker capable of linking compounds with SH-containing compounds. Bifunctional and heterobifunctional linker molecules, such as carbohydrate-directed heterobifunctional linker molecules, such as S-(2-thiopyridyl)-L-cysteine hydrazide (TPCH), are also within the scope of the present invention (Vogel, 2004). The effector molecule, such as a maytansinoid, may be conjugated to the targeting antibody via a two reaction step process, including as a first step modification of the targeting antibody with a cross-linking reagent such as N-succinimidyl pyridyldithiopropionate (SPDP) to introduce thiopyridyl groups into the targeting antibody. In a second step, a reactive maytansinoid having a thiol group, such as DM1, may be added to the modified antibody, resulting in the displacement of the thiopyridyl groups in the modified antibody, and the production of disulfide-linked cytotoxic maytansinoid/antibody conjugate (United States Patent 5,208,020). However, one-step conjugation processes such as the one disclosed in United States Patent Publication 20030055226 to Chari et al are also within the scope of the present invention. In one embodiment of the present invention multiple effector molecules of the same or different kind are attached to a targeting antibody. As discussed elsewhere herein, the nature of the linkers employed may influence bystander killing (Kovtun et al., 2006). See also discussion of Fig. 13.

**[0101]** CC-1065 analogues or derivatives may be conjugated to the targeting agent via for example PEG linking groups as described in United States Patent 6,716,821.

**[0102]** Calicheamicins may be conjugated to the targeting antibodies via linkers (United States Patent 5,877,296 and United States Patent 5,773,001) or according to the conjugation methods disclosed in United States Patent 5,712,374 and United States Patent 5,714,586. Another preferred method for preparing calicheamicin conjugates is disclosed in United States Patent Publication 20040082764. The immunoconjugates of the present description may take the form of recombinant fusion proteins.

**[0103]** The term "cytotoxic agents" comprises "cytotoxic/cancer drugs" including chemotherapeutic agents such as melphalan, cyclophosphamide, vincristine, doxorubicin and liposomal doxorubicin (DOXIL), cyclophosphamide, etoposide, cytarabine and cisplatin, corticosteroids such as prednisone and dexamethasone and agents such as thalidomide, bortezomib, lenalidomide, but also kinase inhibitor such as sorafenib or HDAC (histone deacetylase) inhibitors such as romidepsin as well as growth inhibitory agents, anti-hormonal agents, anti-angiogenic agents, cardioprotectants, immunostimulatory agents, immunosuppressive agents, angiogenesis inhibitors, protein tyrosine kinase (PTK) inhibitors. Also included in this definition are antibody based cytotoxic agents including immunoconjugates and antibodies that have an art recognized cytotoxic effect. Anti-CD40 is a preferred antibody. Other antibodies include, but are not limited to, e.g., AVASTIN (bevacizumab) or MYELOMACIDE (milatuzumab).

**[0104]** THALOMID ( $\alpha$ -(N-phthalimido) glutarimide; thalidomide), is an immunomodulatory agent. The empirical formula for thalidomide is  $C_{13}H_{10}N_2O_4$  and the gram molecular weight is 258.2. The CAS number of thalidomide is 50-35-1. It appears to have multiple actions, including the ability to inhibit the growth and survival of myeloma cells in various ways and to inhibit the growth of new blood vessels.

**[0105]** VELCADE is a proteasome inhibitor used to treat multiple myeloma. It is believed that VELCADE acts on myeloma cells to cause cell death, and/or acts indirectly to inhibit myeloma cell growth and survival by acting on the bone microenvironment. Without being limited to a specific theory or mode of action, VELCADE thus disrupts normal cellular processes, resulting in proteasome inhibition that promotes apoptosis.

**[0106]** REVIMID is an immunomodulatory agent. It is thought that REVIMID affects multiple pathways in myeloma cells, thereby inducing apoptosis, inhibiting myeloma cell growth, inhibiting vascular endothelial growth factor (VEGF) thereby inhibiting angiogenesis, and reducing adhesion of myeloma cells to bone marrow stromal cells.

**[0107]** Dexamethasone is a synthetic glucocorticoid steroid hormone that acts as an anti-inflammatory and immunosuppressant. When administered to cancer patients, dexamethasone can counteract side effects of cancer therapy. Dexamethasone can also be given alone or together with other anticancer agents, including thalidomide, adriamycin or vincristine.

**[0108]** The term "in combination with" is not limited to the administration at exactly the same time. Instead, the term encompassed administration of the immunoconjugate of the present invention and the other regime (e.g. radiotherapy) or agent, in particular the cytotoxic agents referred to above in a sequence and within a time interval such that they may act together to provide a benefit that is increased compared to treatment with only either the immunoconjugate of the present invention or, e.g., the other agent or agents. It is preferred that the immunoconjugate and the other agent or agents act additively, and especially preferred that they act synergistically. Such molecules are suitably provided in

amounts that are effective for the purpose intended. The skilled medical practitioner can determine empirically, or by considering the pharmacokinetics and modes of action of the agents, the appropriate dose or doses of each therapeutic agent, as well as the appropriate timings and methods of administration. As used in the context of the present description "co-administration" refers to administration at the same time as the immunoconjugate, often in a combined dosage form.

5 [0109] The term "sequence identity" refers to a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity", per se, has recognized meaning in the art and can be calculated using published techniques. (See, e.g.: Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heijne, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H. & Lipton, D., SIAM J Applied Math 48:1073 (1988)).

10 [0110] Whether any particular nucleic acid molecule is at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the nBT062 nucleic acid sequence, or a part thereof, can be determined conventionally using known computer programs such as DNAsis software (Hitachi Software, San Bruno, Calif.) for initial sequence alignment followed by ESEE version 3.0 DNA/protein sequence software (cabot@trog.mbb.sfu.ca) for multiple sequence alignments.

15 [0111] Whether the amino acid sequence is at least 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance SEQ ID NO:1 or SEQ ID NO:2, or a part thereof, can be determined conventionally using known computer programs such the BESTFIT program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, Wis. 53711). BESTFIT uses the local homology algorithm of Smith and Waterman, Advances in Applied Mathematics 2:482-489 (1981), to find the best segment of homology between two sequences.

20 [0112] When using DNAsis, ESEE, BESTFIT or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present description, the parameters are set such that the percentage of identity is calculated over the full length of the reference nucleic acid or amino acid sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

25 [0113] If, in the context of the present invention, reference is made to a certain sequence identity with a combination of residues of a particular sequence, this sequence identity relates to the sum of all the residues specified.

30 [0114] The basic antibody molecule is a bifunctional structure wherein the variable regions bind antigen while the remaining constant regions may elicit antigen independent responses. The major classes of antibodies, IgA, IgD, IgE, IgG and IgM, are determined by the constant regions. These classes may be further divided into subclasses (isotypes). For example, the IgG class has four isotypes, namely, IgG1, IgG2, IgG3, and IgG4 which are determined by the constant regions. Of the various human antibody classes, only human IgG1, IgG2, IgG3 and IgM are known to effectively activate the complement system. While the constant regions do not form the antigen binding sites, the arrangement of the constant regions and hinge region may confer segmental flexibility on the molecule which allows it to bind with the antigen.

35 [0115] Different IgG isotypes can bind to Fc receptors on cells such as monocytes, B cells and NK cells, thereby activating the cells to release cytokines. Different isotypes may also activate complement, resulting in local or systemic inflammation. In particular, the different IgG isotypes may bind Fc $\gamma$ R to different degrees. Fc $\gamma$ Rs are a group of surface glycoproteins belonging to the Ig superfamily and expressed mostly on leucocytes. The Fc $\gamma$ Rs are divided into three classes designated Fc $\gamma$ RI (CD64), Fc $\gamma$ RII (CD32) and Fc $\gamma$ RIII (CD16). While IgG1, IgG2 and IgG3 bind strongly to a variety of these classes of Fc $\gamma$ Rs, IgG4 display much weaker binding. In particular, IgG4 is an intermediate binder of Fc $\gamma$ RI, which results in relatively low or even no ADCC (antibody dependent cellular cytotoxicity), and does not bind to Fc $\gamma$ RIIIA or Fc $\gamma$ RIIA. IgG4 is also a weak binder of Fc $\gamma$ RIIB, which is an inhibitory receptor. Furthermore, IgG4 mediates only weak or no complement fixation and weak or no complement dependent cytotoxicity (CDC). In the context of the present invention, IgG4 is specifically employed to prevent Fc-mediated targeting of hepatic FcR as it displays no interaction with Fc $\gamma$ RII on LSECs (liver sinusoidal endothelial cells), no or weak interaction with Fc $\gamma$ RI-III on Kupffer cells (macrophages) and no interaction with Fc $\gamma$ RIII on hepatic NK cells. Certain mutations that further reduce any CDC are also part of the present invention. For example IgG4 residues at positions 327, 330 and 331 were shown to reduce ADCC (antibody dependent cellular cytotoxicity) and CDC (Amour, 1999; Shields, 2001). One or more mutations that stabilize the antibody are also part of the present invention (also referred to herein as "stabilizing mutations"). Those mutations include in particular, leucine-to-glutamic acid mutations in the CH2 region of IgG4 and serine-to-proline exchanges in the IgG4 hinge core. These mutations decrease, in certain embodiments of the invention, the amount of half-molecules to less than 10%, less than 5% and preferably less than 2% or 1%. Moreover, the in vivo half life of so stabilized antibodies might be increased several days, including 1, 2, 3, 4 or more than 5 days (Schuurman, 1999).

40 [0116] Targeting agents, including targeting antibodies disclosed herein may also be described or specified in terms

of their **binding affinity** to antigen, in particular to CD138. Preferred binding affinities of targeting agents such as targeting antibodies are characterized by dissociation constants  $K_D$  (nM) of less than 1.6, less than 1.5 or about or less than 1.4. For immunoconjugates comprising said targeting agents such as targeting antibodies dissociation constants  $K_D$  (nM) of less than 1.6, less than 1.5 or less than 2.5, less than 2.4, less than 2.3, less than 2.2, less than 2.1, less than 2.0, less than or about 1.9 are preferred.

[0117] An **antigen binding region** (ABR) according to the present description will vary based on the type of targeting antibody or engineered targeting antibody employed. In a naturally occurring antibody and in most chimeric and humanized antibodies, the antigen binding region is made up of a light chain and the first two domains of a heavy chain. However, in a heavy chain antibody devoid of light chains, the antigen binding region will be made up of, e.g., the first two domains of the heavy chain only, while in single chain antibodies (ScFv), which combine in a single polypeptide chain the light and heavy chain variable domains of an antibody molecule, the ABR is provided by only one polypeptide molecule. FAB fragments are usually obtained by papain digestion and have one light chain and part of a heavy chain and thus comprise an ABR with only one antigen combining site. On the other hand, diabodies are small antibody fragments with two antigen-binding regions. In the context of the present description, however, an antigen binding region of an targeting antibody or engineered targeting antibody is any region that primarily determines the **binding specificity** of the targeting antibody or engineered targeting antibody.

[0118] If an ABR or another targeting antibody region is said to be "**of a certain antibody**", e.g., a human or non-human antibody, this means in the context of the present description that the ABR is either identical to a corresponding naturally occurring ABR or is based thereon. An ABR is based on a naturally occurring ABR if it has the binding specificity of the naturally occurring ABR. However, such an ABR may comprise, e.g., point mutations, additions, deletions or posttranslational modification such as glycosylation. Such an ABR may in particular have more than 70%, more than 80%, more than 90%, preferably more than 95%, more than 98% or more than 99% sequence identity with the sequence of the naturally occurring ABR.

[0119] **Homogenous targeting** of a targeting agent such as a targeting antibody, but in particular an immunoconjugate comprising the same, in the context of the present description, is a measure of the variance associated with obtaining the desired result of said targeting with the targeting agent. In certain embodiments, the desired result is obtained by simple binding to the target. This is, for example, the case in embodiments in which a certain targeting agent provides a shield against subsequent binding. However, the homogeneity of a targeting agent can be readily assessed, e.g., via the efficacy of an immunoconjugate comprising said targeting agent. For example, the efficacy of said immunoconjugate against a tumor antigen such as CD138 that comprises an effector aimed at destroying tumor cells and/or arresting the growth of a tumor can be determined by the degree of growth suppression of a tumor comprising cells expressing the CD138 antigen. Such an immunoconjugate may display a high variance in its efficacy. It may, for example, arrest tumor growth sometimes with high efficacy, but other times with an efficacy that hardly exceeds the efficacy of the control. A low variance in the efficacy of an immunoconjugate, on the other hand, shows that the immunoconjugate and/or targeting agent, respectively, provide the desired result consistently. One way of quantifying the homogeneity of targeting is to calculate the **targeting variation**. In the context of tumor growth arrested by an immunoconjugate comprising a certain targeting agent, the targeting variation can be calculated by first determining the time for a tumor to reach a predetermined volume, e.g. 300mm<sup>3</sup>. Preferably, the predetermined volume is chosen so that any tumor growth before and after reaching said predetermined volume is steadily increasing at about the same rate. After such time has been determined for a group of subjects the mean of these times ( $T_m$ ) in the group of subjects (e.g., SCID mice or another suitable model displaying homogenous tumor growth) is calculated.  $T_m$  is then correlated to the observations made in the subject of the group showing the least efficacy in targeting and thus being associated with tumors that need the least time ( $T_f$ ) to reach said predetermined volume, and, on the other hand, the subject in the group showing the highest efficacy in targeting and thus being associated with tumors that need the most time ( $T_s$ ) to reach said predetermined volume by calculating the targeting variation for the predetermined volume according to the following formula:

**TARGETING VARIATION [%] =**

$$Ts-Tf/Tm \times 100$$

[0120] In a preferred embodiment, the targeting variation of the immunoconjugate comprising the engineered targeting antibody of the present description is less than 150%, less than 140%, less than 130%, less than 120%, less than 110%, less than 100%, less than 90%, less than 80%, less than 70%, less than 60%, or less than 50%, and in certain embodiments even less than 45%. Preferably, the targeting variation is between about 10% and about 150%, preferably between about 10% and about 100%, about 10% and about 80%, about 10% and about 70%, about 10% and about 60%, about 10% and about 50%.

[0121] The homogeneity of targeting can be also quantified by other means such as determining the tumor growth delay. Also, as the person skilled in the art will readily understand tumor volume of a certain size is only one parameter

on which basis targeting variation may be determined. Depending on the desired result, other parameters include time (for, e.g., measuring tumor growth delay) or % of binding may be employed. The person skilled in the art can readily determine such other parameters.

**[0122]** FIG. 9 shows in (C) and (D) the differences in homogeneity of targeting/binding between immunoconjugates comprising murine antibody BB4 (BB4-SPP-DM1; FIG. 9C) and the engineered targeting antibody nBT062 (nBT062-SPP-DM1; FIG. 9D) based thereon. As can be seen from these graphs, results obtained with the immunoconjugate comprising the engineered targeting antibody are substantially more homogenous than the ones obtained with the immunoconjugates comprising the murine antibody. This is particularly notable since the antibody binding region of BB4 was not modified in nBT062. Thus, the immunoconjugate comprising the antibody binding region of the murine antibody, but no other parts of the murine antibody, showed properties that far exceeded results the person skilled in the art would have expected.

**[0123]** nBT062 (see also FIG. 1) is a murine human chimeric IgG4 mAb a chimerized version of B-B4. This chimerized version of B-B4 was created to reduce the HAMA (Human Anti-Mouse Antibody) response, while maintaining the functionality of the antibody binding region of the B-B4 for CD138. Surprisingly, the results obtained using an immunoconjugate comprising the engineered targeting antibody were much more homogenous (the variance in the results was reduced). The protocol for producing nBT062 is specified below. Chinese hamster ovary cells expressing nBT062 have been deposited with the DSMZ-Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1, D-38124 Braunschweig on December 11, 2007. The identification number is DSM ACC2875. A CD138 specific chimeric antibody based on B-B4 is generically referred to herein as c-B-B4.

**[0124]** The amino acid sequence for both, the heavy and the light chains has been predicted from the translation of the nucleotide sequence for nBT062. The amino acid sequences predicted for the heavy chain and light chain are presented in **Table 1**. Predicted variable regions are bolded, predicted CDRs are underlined.

**Table 1.** Predicted Amino Acid Sequence for nBT062

- nBT062 heavy chain predicted sequence (SEQ ID NO:1):

1	<b>QVQLQQSGSE</b> <u>LMMPGASVKI</u> <b>SCKATGYTFS</b> <u>NYWIEWVKQR</u> <b>PGHGLEWIGE</b>
51	<u>ILPGTGRTIY</u> <b>NEKFKGKATF</b> <u>TADISSNTVQ</u> <b>MQLSSLTSED</b> <u>SAVYYCARRD</u>
101	<u>YYGNFYYAMD</u> <b>YWGQGTSVTV</b> <u>SSASTKGPSV</u> <b>FPLAPCSRST</b> <u>SESTAALGCL</u>
151	<b>VKDYFPEPV</b> T VSWNSGALTS GVHTFPAVLQ SSGLYSLSSV VTVPSSSLGT
201	KTYTCNVDHK PSNTKVDKRV ESKYGPPCPS CPAPEFLGGP SVFLFPPKPK
251	DTLMISRTPE VTCVVVDVVSQ EDPEVQFNWY VDGVEVHNAK TKPREEQFNS
301	TYRVVSVLTV LHQDWLNGKE YKCKVSNKGL PSSIEKTISK AKGQPREPQV
351	YTLPPSQEEM TKNQVSLTCL VKGFYPSDIA VEWESNGQPE NNYKTPPPV
401	DSDGSFFLYS RLTVDKSRWQ EGNVFSCSVM HEALHNHYTQKSLSLSLG (K)

The C-terminal lysine is prone to clipping and might be present due to incomplete clipping to a certain extent. The (K) in parenthesis is not part of SEQ ID NO:1.

- nBT062 light chain predicted sequence (SEQ ID NO:2):

1	<b>DIQMTQSTSS</b> <u>LSASLGDRVT</u> <b>ISCSASQGIN</b> <u>NYLNWYQQKP</u> <b>DGTVELLIYY</b>
51	<u>TSTLQSGVPS</u> <b>RFSGSGSGTD</b> <u>YSLTISNLEP</u> <b>EDIGTYYCQQ</b> <u>YSKLPRTFGG</u>
101	<b>GTKLEIKRTV</b> AAPSVFIFPP SDEQLKSGTA SVVCLLNNFY PREAKVQWKV
151	DNALQSGNSQ ESVTEQDSKD STYSLSSTLT LSKADYEKHK VYACEVTHQG
201	LSSPVTKSFN RGEc

**Table 2.** shows a comparison of the general CDR definitions of Krabat and Chothia and the predicted CDRs for BT062  
**Krabat CDR definition** **nBT062**

Light chain	CDR1: residues 24-34	<u>CDR1: residues 24-34</u>
	CDR2: residues 50-56	<u>CDR2: residues 50-56</u>

(continued)

<b>Kabat CDR definition</b>		<b>nBT062</b>
	CDR3: residues 89-97	<u>CDR3: residues 89-97</u>
5		
	Heavy chain	CDR1: residues 31-35 CDR2: residues 50-56 CDR3: residues 95-102
10	<b>Chothia CDR definition</b>	
	Light chain	CDR1: residues 26-32 CDR2: residues 50-52 CDR3: residues 91-96
15		
	Heavy chain	CDR1: residues 26-32 CDR2: residues 52-56 CDR3: residues 96-101

20 [0125] BT062 is an immunoconjugate comprising the CD138 targeting chimeric antibody nBT062 that is attached via a linker, here SPDB, to the cytostatic maytansinoid derivative DM4. A chemical representation of BT062 is provided in FIGS. 1 and 2. Immunoconjugates comprising nBT062 and a maytansinoid effector molecule are often characterized in terms of their linker and maytansinoid effector, e.g., nBT062-SMCC-DM1, is an immunoconjugate comprising nBT062, SMCC (a "noncleavable" linker containing a thioester bond) and DM1 as an effector. More generically, an immunoconjugate containing nBT062 and an effector molecule may also be described as nBT062-linker-effector or just as nBT062-effector (nBT062N, wherein N is any effector described herein).

25 [0126] Reference is made herein to a unhindered counterpart (**UI**: unhindered immunoconjugate) of an immunoconjugate comprising an engineered targeting antibody against CD138 attached to an effector molecule via a cleavable linker (CL) and is described herein as UICL, which is contrasted to an immunoconjugate in which said effector molecule is sterically hindered, and contains a cleavable linker (HICL -hindered immunoconjugate, cleavable linker). The UICL is an immunoconjugate equivalent to the HICL comprising an engineered targeting antibody in which the effector molecule is, however, not sterically hindered. Examples of a pair of HICL/UICL are BT062 and nBT062-SPP-DM1. An unhindered counterpart of such an immunoconjugate comprising a non-cleavable linker (**UNICL**) refers to the equivalent immunoconjugate comprising an engineered targeting antibody in which the effector molecule is not sterically hindered and comprises a noncleavable linker. For BT062, nBT062-SMCC-DM1 would constitute an example of such an unhindered counterpart comprising an non-cleavable linker (UNICL).

30 [0127] A growth of a tumor inhibiting activity (=tumor growth inhibiting activity) of an immunoconjugate is a relative measure. It describes the tumor growth inhibiting activity of a conjugate relative to the activity of the highest performing immunoconjugate whose activity is set as 100%. For example if the activity of the highest performing immunoconjugate, say, BT062, which causes a tumor growth delay (TGD) of 32 days, is set as 100%, the activity of, e.g., nBT062-DM1, which displays a tumor growth delay (TGD) of 18 days is calculated as follows:

35 **Tumor Growth Inhibiting Activity=**

40 
$$100 \times (TGD_{nBT062-DM1} / TGD_{BT062}),$$

45 more generically:

50 **Tumor Growth Inhibiting Activity=**

$$100 \times (TGD_{\text{Sample}} / TGD_{\text{Reference}}).$$

55 [0128] **Table 3** provides suitable examples from the results depicted in Fig. 11 B:

**Table 3:** Tumor growth delay (TGD) and % Activity of nBT062-DMx against MOLP-8 tumor xenografts in SCID mice based on treatment groups receiving a 450 µg/kg dose.

	TGD* (days)	% Activity**
PBS	0	0
nBT062-SMCC-DM1	18	56
BT062	32	100
nBT062-SPP-DM1	13	40

(\*) Tumor growth delay in days (TGD) as mean time in days for treatment group to reach a predetermined size (160 mm<sup>3</sup>) minus the mean time for the control group to reach this predetermined size.

(\*\*) Tumor Growth Inhibiting Activity = 100x(TGD<sub>Sample</sub>/TGD<sub>BT062</sub>). The activity of BT062 is defined to be 100%.

[0129] In the example provided in Table 2, BT062 provides a growth of a tumor inhibiting activity that exceeds that of its unhindered counterpart (nBT062-SPP-DM1) by 60%, and a growth of a tumor inhibiting activity that exceeds that of its unhindered counterpart immunoconjugate comprising a non-cleavable linker (nBT062-SMCC-DM1) by 44%.

[0130] It was previously reported that a cleavable linker in e.g., huC242-maytansinoid immunoconjugates may provide for a so called bystander effect. Goldmahker et al. (U.S. Patent Publication 2006/0233814) also disclose that the bystander effect is particularly pronounced when the effector molecule is subject to further modification, in particular alkylation, upon cleavage from the targeting agent. Goldmahker et al. also showed that UICL displayed better TGD than the respective UINCL (see, e.g., Fig. 6 of U.S. Patent Publication 2006/0233814).

[0131] However, the overall effectiveness of HICL/UICL/UINCL immunoconjugates appear to differ from immunoconjugate to immunoconjugate and/or target to target. For example the HICL trastuzumab-SPP-DM4 was clearly outperformed in its ability to reduce tumor size by the UINCL trastuzumab -SMCC-DM1, while performance of the UICL immunoconjugate trastuzumab -SPP-DM1 substantially resembled that of the corresponding HICL (see U.S. Patent Publication 2008/0171040 to Eberts et al.), thus rendering the results obtained a function of the immunoconjugate and the target.

[0132] Here, yet another relationship was found. While the HICL outperformed the UICL and UINCL, it was also surprisingly found that an UICL in a high single dosage regime (250 µg/kg) actually did not show any better results than the UINCL. In fact, the TGD in days that was observed in an UICL in such a regime was actually lower than that of the UINCL. This observation became more pronounced with an increase in dosage (450 µg/kg). In sum, as shown in Fig. 11A, HICL outperformed UICL in single dose experiments as well as multiple dose experiments (not shown), to an unexpected degree. In addition, the UICL was unexpectedly outperformed by UINCL at higher dosages.

[0133] The targeting agents, in particular targeting antibodies, and/or immunoconjugates disclosed herein can be administered by any route, including intravenously, parenterally, orally, intramuscularly, intrathecally or as an aerosol. The mode of delivery will depend on the desired effect. A skilled artisan will readily know the best route of administration for a particular treatment in accordance with the present invention. The appropriate dosage will depend on the route of administration and the treatment indicated, and can readily be determined by a skilled artisan in view of current treatment protocols.

[0134] Pharmaceutical compositions containing an unconjugated targeting agent, the immunoconjugate of the present invention and/or any further cytotoxic agent as active ingredients can be prepared according to conventional pharmaceutical compounding techniques. See, for example, Remington's Pharmaceutical Sciences, 17th Ed. (1985, Mack Publishing Co., Easton, Pa.). Typically, effective amounts of active ingredients will be admixed with a pharmaceutically acceptable carrier. The carrier may take a wide variety of forms depending on the form of preparation desired for administration, for example, intravenous, oral, parenteral, intrathecal, transdermal, or by aerosol.

[0135] For oral administration, the targeting agent and/or immunoconjugate and/or cytotoxic agent can be formulated into solid or liquid preparations such as capsules, pills, tablets, lozenges, melts, powders, suspensions or emulsions. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed, such as, for example, water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents, suspending agents, and the like in the case of oral liquid preparations (such as, for example, suspensions, elixirs and solutions); or carriers such as starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like in the case of oral solid preparations (such as, for example, powders, capsules and tablets). Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar-coated or enteric-coated by standard techniques. The active agent must be stable to passage through the gastrointestinal tract. If necessary, suitable agents for stable passage can be

used, and may include phospholipids or lecithin derivatives described in the literature, as well as liposomes, microparticles (including microspheres and macroospheres).

**[0136]** For parenteral administration, the targeting agent and/or the immunoconjugate and/or cytotoxic agent may be dissolved in a pharmaceutical carrier and administered as either a solution or a suspension. Illustrative of suitable carriers are water, saline, phosphate buffer solution (PBS), dextrose solutions, fructose solutions, ethanol, or oils of animal, vegetative or synthetic origin. The carrier may also contain other ingredients, for example, preservatives, suspending agents, solubilizing agents, buffers and the like. When the unconjugated targeting agent and/or immunoconjugate and/or cytotoxic agent are being administered intracerebroventricularly or intrathecally, they may also be dissolved in cerebrospinal fluid.

**[0137]** Dosages administered to a subject may be specified as amount, per surface area of the subject (which include humans as well as non-human animals). The dose may be administered to such a subject in amounts, preferably, but not exclusively from about 5 mg/m<sup>2</sup> to about 300 mg/m<sup>2</sup>, including about 20mg/m<sup>2</sup>, about 50mg/m<sup>2</sup>, about 100mg/m<sup>2</sup>, about 150mg/m<sup>2</sup>, about 200mg/m<sup>2</sup> and about 250mg/m<sup>2</sup>. The targeting agents/ immunoconjugates are suitably administered at one time or over a series of treatments. In a multiple dose regime these amounts may be administered once a day, once a week, once every two weeks, once every three weeks, once every four weeks, one every five weeks or once every six weeks. Loading doses with a single high dose or, alternatively, lower doses that are administered shortly after one another followed by dosages timed at longer intervals constitute a preferred embodiment of the present invention. In a preferred embodiment, the timing of the dosages are adjusted for a subject so that enough time has passed prior to a second and/or any subsequent treatment so that the previous dose has been metabolized substantially, but the amount of immunoconjugate present in the subject's system still inhibits, delays and/or prevents the growth of a tumor. An exemplary "repeated single dose" regime comprises administering an initial dose of immunoconjugate of about 200mg/m<sup>2</sup> once every three weeks. Alternatively, a high initial dose may be followed by a biweekly maintenance dose of about 150 $\mu$ g/m<sup>2</sup>. However, other dosage regimens may be useful. The progress of this therapy is easily monitored by known techniques and assays. Dosage may vary depending on whether they are administered for preventive or therapeutic purposes, the course of any previous therapy, the patient's clinical history and response to the targeting agent/immunoconjugate, and the discretion of the attending physician.

**[0138]** In one embodiment, the immunoconjugate is administered with one or more additional cytotoxic agents such as an relevant antibody or a fragment thereof, which is efficient in treating the same disease or an additional comorbidity; for example a CD138 specific immunoconjugate can be administered in combination with an antibody that recognizes another antigen that is overexpressed in the target type of cancer. An example of an antibody that can be administered in combination with a CD138 specific immunoconjugate is an anti-CD40 antibody (Tai et al., 2006).

**[0139]** Other anti-cancer antibodies and immunoconjugates that can be co-administered with the immunoconjugates disclosed herein, including but not limited to, e.g., AVASTIN (bevacizumab) or MYELOMACIDE (milatuzumab).

**[0140]** The immunoconjugate of the present invention may also particularly be administered in in treatment regimens with high-dose chemotherapy (preferably, melphalan, melphalan/prednisone (MP), vincristine/doxorubicin/dexamethasone (VAD), liposomal doxorubicin/vincristine, dexamethasone (DVd), cyclophosphamide, etoposide/dexamethasone/cytarabine, cisplatin (EDAP)), stem cell transplants (e.g., autologous stem cell transplantation or allogeneic stem cell transplantation, and/or mini-allogeneic (non-myeloablative) stem cell transplantation), steroids (e.g., corticosteroids, dexamethasone, thalidomide/dexamethasone, prednisone, melphalan/prednisone), supportive therapy (e.g., bisphosphonates, growth factors, antibiotics, intravenous immunoglobulin, low-dose radiotherapy, and/or orthopedic interventions), THALOMID (thalidomide, Celgene), VELCADE (bortezomib, Millennium), and/or REVOLIMID (lenalidomide) (Chugene Corporation) and/or other multiple myeloma treatments including radiation therapy.

**[0141]** If an immunoconjugate of the present invention is administered in combination with a cytotoxic agents, the above doses and regimes are often maintained. However, if the immunconjugate and the cytotoxic agent are co-administered, low dosages of each of these therapeutic components are, in certain embodiments, preferred. In such a situation, the immunoconjugate may be administered at doses from about 5 mg/m<sup>2</sup> to about 200 mg/m<sup>2</sup>, including about 20mg/m<sup>2</sup>, about 50mg/m<sup>2</sup>, about 100mg/m<sup>2</sup>, about 150mg/m<sup>2</sup>, while the cytotoxic agent is administered at doses that are below the recommendation when administered alone, such at about 80% to 20% of the recommended dose.

**[0142]** The experimental data obtained in the cell culture based (Fig. 7) and mouse experiments (Figs. 8 to 11), was further confirmed with experiments that further supported these finding.

**[0143]** The pathogenesis of multiple myeloma involves binding of myeloma cells, via cell-surface adhesion molecules, to bone marrow stroma cells (BMSCs) as well as the extracellular matrix (ECM). This binding triggers, and thus can be made ultimately responsible, for multiple myeloma cell growth, drug resistance, and migration of MM cells in the bone marrow milieu (Munshi et al. 2008). In particular, the adhesion of multiple myeloma cells to ECM via syndecan-1 (CD138) to type I collagen, induces the expression of matrix metalloproteinase 1, thus promoting bone resorption and tumour invasion (Hideshima et al. 2007). Interactions between multiple myeloma cells and the bone marrow microenvironment results in activation of a pleiotropic proliferative and anti-apoptotic cascade.

**[0144]** Following the homing of multiple myeloma cells to the bone marrow stromal compartment, adhesion between

multiple myeloma cells and BMSCs upregulates many cytokines like interleukin-6 (IL-6) and insulin like growth factor 1 (IGF-1) which have angiogenic and tumor growth promoting activities (Hideshima et al. 2007). The signalling cascades initiated by these cytokines eventually result in MM cell resistance to conventional therapeutics (Anderson et al. 2000; Hideshima et al. 2006).

5 [0145] The *in vivo* efficacy of nBT062-SPDB-DM4 and nBT062-SPP-DM1 against CD138-positive tumor cells in the presence of human bone marrow was analyzed in a mouse model and the results of this analysis are shown in Fig. 12. The Figure shows that both HICL and UICL perform well in this environment. The increase in the level of shuL-6R which can, in this model, be used as a parameter of MM cell growth, were both suppressed by the these immunoconjugates.

10 [0146] In accordance with the present invention, MM is treated as follows, with the use of BT062 as an example. This example is not intended to limit the present invention in any manner, and a skilled artisan could readily determine other immunoconjugate or nBT062 based systems that are within the scope of the present invention and other treatment regimes which could be utilized for the treatment of diseases such as MM.

15 [0147] Due to the selective expression of CD138 on patient MM cells on via the blood stream accessible cells, the specificity of nBT062 and the stability of BT062 in the bloodstream, BT062 removes the systemic toxicity of DM4 and provides an opportunity to target the delivery of the DM4-effector molecule(s) homogenously. The immunoconjugates of this invention provide a means for the effective administration of the effector molecules to cell sites where the effector molecules can be released from the immunoconjugates. This targeted delivery and release provides a significant advance in the treatment of multiple myeloma, for which current chemotherapy methods sometimes provide incomplete remission.

20 [0148] In accordance with the present invention multiple myeloma is also treated as follows: One or more cytotoxic agents are administered in the dosages and dosage forms and according to establish treatment protocols for these cytotoxic agents to an individual that is also treated with an immunoconjugate of the present invention.

25 [0149] In particular, a patient is subjected to a treatment regime using an oral dosage of melphalan according to the manufacturer's instruction (e.g. a pill traded under the trademark ALKERAN) and an appropriate dosage of BT062, e.g., 100mg/m<sup>2</sup> according to the present invention at certain intervals, e.g., at the beginning or end of a melphalan treatment session, to complement the effect of the melphalan treatment.

30 [0150] In accordance with the present invention, in particular solid tumors may also be treated as follows using BT062 as an example. This example is not intended to limit the present invention in any manner, and a skilled artisan could readily determine other immunoconjugates of the present invention and other treatment regimes which could be utilized for the treatment of solid tumors. The tumor is first treated to reduce the size of the tumor, for example chemotherapeutically, e.g., using liposomal doxorubicin, or radioactively. Subsequent administration of BT062 this invention provides a highly effective means for eliminating residual cancer cells. The administration of the immunoconjugate allows specific targeting of these residual cells and release of the effector molecules at the target site. The high efficiency of the immunoconjugate allows, in preferred embodiments, for a single dose regime. This targeted delivery and release provides a significant advance in the treatment of residual cancer cells of solid tumors, for which current chemotherapy methods sometimes provide incomplete remission.

35 [0151] The present invention is further described by reference to the following Examples, which are offered by way of illustration. Standard techniques well known in the art or the techniques specifically described below are utilized.

## Materials and Methods

### 40 Chimeric Antibody Construction (cB-B4: nBT062)

#### B-B4

45 [0152] Murine antibody B-B4 as previously characterized (Wijdenes et al., Br J Haematol., 94 (1996), 318) was used in these experiments.

#### Cloning and expression of B-B4 and cB-B4 / nBT062

50 [0153] Standard recombinant DNA techniques were performed as described in detail in text books, for example in J. Sambrook; Molecular Cloning, A Laboratory Manual; 2nd Ed. (1989), Cold Spring Harbor Laboratory Press, USA, or as recommended by the manufacturer's instruction in the cases when kits were used. PCR-cloning and modification of the mouse variable regions have been conducted using standard PCR methodology. Primers indicated in the respective results section have been used.

#### 55 Expression of cB-B4 / nB T062

[0154] Exponentially growing COS cells, cultured in DMEM supplemented with 10% FCS, 580 µg/ml L-glutamine, 50

5 Units/ml penicillin and 50 µg/ml streptomycin were harvested by trypsinisation and centrifugation and washed in PBS. Cells were resuspended in PBS to a final concentration of 1x10<sup>7</sup> cells/ml. 700 µl of COS cell suspension was transferred to a Gene Pulser cuvette and mixed with heavy and kappa light chain expression vector DNA (10 µg each or 13 µg of Suprvector). Cells were electroporated at 1900 V, 25 µF using a Bio-Rad Gene Pulser. Transformed cells were cultured in DMEM supplemented with 10% gamma-globulin free FBS, 580 µg/ml L-glutamine, 50 Units/ml penicillin and 50 µg/ml streptomycin for 72 h before antibody-containing cell culture supernatants were harvested.

*Capture ELISA to measure expression levels of cB-B4 / nBT062*

10 [0155] 96 well plates were coated with 100 µl aliquots of 0.4 µg/ml goat anti-human IgG antibody diluted in PBS (4°C, overnight). Plates were washed three times with 200 µl/well washing buffer (PBS+0.1% Tween-20). Wells were blocked with 0.2% BSA, 0.02% Tween-20 in PBS, before addition of 200 µl cell culture supernatants containing the secreted antibody (incubation at 37°C for one hour). The wells were washed six times with washing buffer, before detection of bound antibody with goat anti-human kappa light chain peroxidase conjugate.

15 *Purification of cB-B4 / nBT062 from cell culture supernatants*

20 [0156] The cB-B4 antibody was purified from supernatants of transformed COS 7 cells using the Protein A ImmunoPure Plus kit (Pierce, Rockford, IL), according to the manufacturer's recommendation.

25 *cB-B4 binding and competition assay*

[0157] Analysis of binding activity of B-B4 and cB-B4 to CD138 was performed using the Diaclone (Besançon, France) sCD138 kit according to the manufacturer's recommendation, considering the changes described in the results section.

30 *RNA preparation and cDNA synthesis*

35 [0158] Hybridoma B-B4 cells were grown and processed using the Qiagen Midi kit (Hilden, Germany) to isolate RNA following the manufacturer's protocol. About 5 µg of B-B4 RNA was subjected to reverse transcription to produce B-B4 cDNA using the Amersham Biosciences (Piscataway, NJ) 1st strand synthesis kit following the manufacturer's protocol.

*Cloning of B-B4 immunoglobulin cDNA*

40 [0159] Immunoglobulin heavy chain (IgH) cDNA was amplified by PCR using the IgH primer MHV7 (5'-ATGGGCAT-CAAGATGGAGTCACAGACCCAGG-3') [SEQ ID NO:3] and the IgG1 constant region primer MHCG1 (5'-CAGTGGATAGACAGATGGGG-3') [SEQ ID NO:4]. Similarly, immunoglobulin light chain (IgL) was amplified using the three different Igκ primers MKV2 (5'-ATGGAGACAGACACACTCCTGCTATGGGTG-3') [SEQ ID NO:5], MKV4 (5'-ATGAG-GGCCCTGCTCAGTTTTGGCTTCTG-3') [SEQ ID NO:6] and MKV9 (5'-ATGGTATCCACACCTCAGTTCTTG-3') [SEQ ID NO:7], each in combination with primer MKC (5'-ACTGGATGGTGGGAAGATGG-3') [SEQ ID NO:8]. All amplification products were directly ligated with the pCR2.1-TOPO vector using the TOPO-TA cloning kit (Invitrogen, Carlsbad, CA) according to the manufacturer's instruction.

45 [0160] *E. coli* TOP10 bacteria (Invitrogen) transformed with the ligated pCR2.1 vector constructs were selected on LB-ampicillin-Xgal agar plates. Small scale cultures were inoculated with single white colonies, grown overnight and plasmids were isolated using the QIAprep Spin Miniprep kit according to the manufacturer's instruction.

*cDNA sequence determination*

50 [0161] Plasmids were sequenced using the BigDye Termination v3.0 Cycle Sequencing Ready Reaction Kit (ABI, Foster City, CA). Each selected plasmid was sequenced in both directions using the 1210 and 1233 primers cycled on a GeneAmp9600 PCR machine. The electrophoretic sequence analysis was done on an ABI capillary sequencer.

[0162] The complete cycle of RT-PCR, cloning and DNA sequence analysis was repeated to obtain three completely independent sets of sequence information for each immunoglobulin chain.

55 *B-B4 Vκ DNA sequence*

[0163] 1st strand synthesis was performed in three independent reactions. The PCR products generated by using primers MKC and MKV2 (sequences given above) were ligated into pCR2.1-TOPO vectors according to the manufacturer's instruction. Clones from each independent set of RT-PCR reactions were sequenced in both directions. MKV2-

primed product sequence was highly similar to sterile kappa transcripts originating from the myeloma fusion partner such as MOPC-21, SP2 and Ag8 (Carroll et al., Mol Immunol., 25 (1988), 991; Cabilly et al., Gene, 40 (1985); 157) and was therefore disregarded.

5 [0164] The PCR products using MKC with MKV4 and MKV9 primers were similar to each other and differed only at the wobble positions within the leader sequence primer.

*B-B4 VH DNA sequence*

10 [0165] 1st strand synthesis was performed in three independent reactions and PCR products were cloned and sequenced from each 1st strand product. Five clones were sequenced from each 1st strand.

*Construction of chimeric cB-B4 expression vectors*

15 [0166] The construction of the chimeric expression vectors entails adding a suitable leader sequence to VH and V $\kappa$ , preceded by a *Bam*HI restriction site and a Kozak sequence. The Kozak consensus sequence is crucial for the efficient translation of a variable region sequence. It defines the correct AUG codon from which a ribosome can commence translation, and the single most critical base is the adenine (or less preferably, a guanine) at position -3, upstream of the AUG start. The leader sequence is selected as the most similar sequence in the Kabat database (Kabat et al., NIH National Technical Information Service, 1991). These additions are encoded within the forward (For) primers (both having 20 the sequence 5'-AGAGAAGCTTGGCCACCATGATT-GCCTCTGCTCAGTTCTGGTCTCC-3' [SEQ ID NO:9]; restriction site is underlined; Kozak sequence is in bold type). Furthermore, the construction of the chimeric expression vectors entails introducing a 5' fragment of the human gamma1 constant region, up to a natural *Apal* restriction site, contiguous with the 3' end of the J region of B-B4 and, for the light chain, adding a splice donor site and *Hind*III site. The splice donor sequence is important for the correct in-frame attachment of the variable region to its appropriate constant 25 region, thus splicing out the V:C intron. The kappa intron + CK are encoded in the expression construct downstream of the B-B4 V $\kappa$  sequence. Similarly, the gamma-4 CH is encoded in the expression construct downstream of the B-B4 VH sequence.

30 [0167] The B-B4 VH and V $\kappa$  genes were first carefully analyzed to identify any unwanted splice donor sites, splice acceptor sites, Kozak sequences and for the presence of any extra sub-cloning restriction sites which would later interfere with the subcloning and/or expression of functional whole antibody. An unwanted *Hind*III site was found in the V $\kappa$  sequence which necessarily was removed by site-directed mutagenesis via PCR without changing the amino acid sequence. For this reactions, oligonucleotide primers BT03 (5'-CAACAGTATAAGCTCCCTCGGACGTTGGTGG-3') [SEQ ID NO:10] and BT04 (5'-CCACCGAACGTCCGAGGGAGCTTACTATACTGTTG-3') [SEQ ID NO:11] were used and mutagenesis was performed according to the Stratagene (La Jolla, CA) Quickchange Mutagenesis Kit protocol.

35 *Kappa chain chimerization primers*

40 [0168] The non-ambiguous B-B4 V $\kappa$  leader sequence, independent of the PCR primer sequence, was aligned with murine leader sequences in the Kabat database. The nearest match for the B-B4 VH leader was VK-10 ARS-A (Sanz et al., PNAS, 84 (1987), 1085). This leader sequence is predicted to be cut correctly by the SignalP algorithm (Nielsen et al., Protein Eng, 10 (1997); 1). Primers CBB4Kfor (see above) and g2258 (5'-CGCGGGATCCACTCACGTTGATT-TCCAGCTTGGTGCCTCC-3' [SEQ ID NO:12]; Restriction site is underlined) were designed to generate a PCR product containing this complete leader, the B-B4 V $\kappa$  region, and *Hind*III and *Bam*HI terminal restriction sites, for cloning into the pKN100 expression vector. The forward primer, CBB4K introduces a *Hind*III restriction site, a Kozak translation initiation site and the VK-10 ARS-A leader sequence. The reverse primer g2258 introduces a splice donor site and a *Bam*HI restriction site. The resulting fragment was cloned into the *Hind*III/*Bam*HI restriction sites of pKN 100.

*Heavy chain chimerization primers*

50 [0169] The non-ambiguous B-B4 VH leader sequence, independent of the PCR primer sequence, was aligned with murine leader sequences in the Kabat database. The nearest match for the B-B4 VK leader was VH17-1A (Sun et al., PNAS, 84 (1987), 214). This leader sequence is predicted to be cut correctly by the SignalP algorithm. Primers cBB4Hfor (see above) and g22949 (5'-CGATGGGCCTTGGTGGAGGCTGAGGA-GACGGTGACTGAGGTTCC-3' [SEQ ID NO:13]; Restriction site is underlined) were designed to generate a PCR product containing VH17-1A leader, the B-B4 VH region, and terminal *Hind*III and *Apal* restriction sites, for cloning into the pG4D200 expression vector. The forward primer cBBHFor introduces a *Hind*III restriction site, a Kozak translation initiation site and the VH17-1A leader sequence. The reverse primer g22949 introduces the 5' end of the gamma4 C region and a natural *Apal* restriction site. The resulting fragment was cloned into the *Hind*III/*Apal* restriction sites of pG4D200, resulting in vector pG4D200cBB4.

*Production of cBB4 antibody*

5 [0170] One vial of COS 7 cells was thawed and grown in DMEM supplemented with 10% Fetal clone I serum with antibiotics. One week later, cells (0.7 ml at  $10^7$  cells/ml) were electroporated with pG4D200cBB4 plus pKN100cBB4 (10  $\mu$ g DNA each) or no DNA. The cells were plated in 8 ml growth medium for 4 days. Electroporation was repeated seven times.

*Detection of chimeric antibody*

10 [0171] A sandwich ELISA was used to measure antibody concentrations in COS 7 supernatants. Transiently transformed COS 7 cells secreted about 6956 ng/ml antibody (data not shown).

*Binding activity of cB-B4*

15 [0172] To assay the binding activity of cB-B4 in COS 7 culture supernatants, the Diaclone sCD138 kit has been used, a solid phase sandwich ELISA. A monoclonal antibody specific for sCD138 has been coated onto the wells of the microtiter strips provided. During the first incubation, sCD138 and biotinylated B-B4 (bio-B-B4) antibody are simultaneously incubated together with a dilution series of unlabeled test antibody (B-B4 or cB-B4).

20 [0173] The concentrations of bio-B-B4 in this assay have been reduced in order to obtain competition with low concentrations of unlabeled antibody (concentration of cB-B4 in COS 7 cell culture supernatants were otherwise too low to obtain sufficient competition). Results from this assay reveal that both antibodies have the same specificity for CD138 (data not shown).

*Purification of cB-B4*

25 [0174] Chimeric B-B4 was purified from COS 7 cell supernatants using the Protein A ImmunoPure Plus kit (Pierce), according to the manufacturer's recommendation (data not shown).

 **$K_D$ -determination: Comparison nBT062/ BB4**

30 *Purification of soluble CD138*

35 [0175] Soluble CD138 antigen from U-266 cell culture supernatant was purified by FPLC using a 1 mL "HiTrap NHS-activated HP" column coupled with B-B4. Cell culture supernatant was loaded in PBS-Buffer pH 7.4 onto the column and later on CD138 antigen was eluted with 50 mM tri-ethylamine pH 11 in 2 mL fractions. Eluted CD138 was immediately neutralised with 375  $\mu$ L 1 M Tris-HCl, pH 3 to prevent structural and/or functional damages.

*Biotinylation of CD138*

40 [0176] Sulfo-NHS-LC (Pierce) was used to label CD138. NHS-activated biotins react efficiently with primary amino groups like lysine residues in pH 7-9 buffers to form stable amide bonds.

[0177] For biotinylation of CD138, 50  $\mu$ l of CD138 were desalted using protein desalting spin columns (Pierce). The biotinylation reagent (EZ-Link Sulfo NHS-LC-Biotin, Pierce) was dissolved in ice-cooled deionised H<sub>2</sub>O to a final concentration of 0.5 mg/mL. Biotinylation reagent and capture reagent solution were mixed having a 12 times molar excess of biotinylation reagent compared to capture reagent (50 pmol CD138 to 600 pmol biotinylation reagent) and incubated 1 h at room temperature while shaking the vial gently. The unbound biotinylation reagent was removed using protein desalting columns.

*Immobilization of bCD138*

50 [0178] The sensorchip (SENSOR CHIP SA, BIACORE AB) used in the BIACORE assay is designed to bind biotinylated molecules for interaction analysis in BIACORE systems. The surface consists of a carboxymethylated dextran matrix pre-immobilized with streptavidin and ready for high-affinity capture of biotinylated ligands. Immobilization of bCD138 was performed on SENSOR CHIP SA using a flow rate of 10  $\mu$ L/min by manual injection. The chip surface was conditioned with three consecutive 1-minute injections of 1 M NaCl in 50 mM NaOH. Then biotinylated CD138 was injected for 1 minute.

*K<sub>D</sub>-Determination of different antibodies using BIACORE*

**[0179]** The software of BIACORE C uses pre-defined masks, so called "Wizards" for different experiments where only certain settings can be changed. As the BIACORE C was originally developed to measure concentrations, there is no wizard designed to carry out affinity measurements. However, with the adequate settings, the wizard for "non-specific binding" could be used to measure affinity rate constants and was therefore used for K<sub>D</sub>-determination. With this wizard, two flow cells were measured and the dissociation phase was set to 90 s by performing the "Regeneration 1" with BIACORE running buffer. "Regeneration 2" which is equivalent to the real regeneration was performed with 10 mM Glycine-HCl pH 2.5. After this step, the ligand CD138 was in its binding competent state again. During the whole procedure HBS-EP was used as running and dilution buffer. To determine binding of the different antibodies (-150 kDa) to CD138, association and dissociation was analysed at different concentrations (100, 50, 25, 12.5, 6.25 and 3.13 nM). The dissociation equilibrium constants were determined by calculating the rate constants k<sub>a</sub> and k<sub>d</sub>. Afterwards, the K<sub>D</sub>-values of the analytes were calculated by the quotient of k<sub>d</sub> and k<sub>a</sub> with the BIAscan evaluation software. The results are shown in **Table 4**.

15 Table 4: Comparative analysis of K<sub>D</sub> values of nBT062 and B-B4. Standard deviations are given for mean K<sub>D</sub> values.

Antibody	Affinity	
	K <sub>D</sub> (nM)	mean K <sub>D</sub> (nM)
<b>nBT062</b>	1.4	
	1.4	<b>1.4 +/- 0.06</b>
	1.5	
<b>B-B4</b>	1.7	
	1.7	<b>1.6 +/- 0.06</b>
	1.6	
<b>nBT062-SPDB-DM4</b>	1.9	
	1.9	<b>1.9 +/- 0.00</b>
	1.9	
<b>B-B4-SPP-DM1</b>	2.6	
	2.7	<b>2.6 +/- 0.06</b>
	2.6	

## 35 Discussion

**[0180]** Mean K<sub>D</sub> values for each antibody were calculated from three independent experiments. The results show that in all measurements nBT062 exhibits slightly decreased K<sub>D</sub> values compared to B-B4 (mean K<sub>D</sub> values were 1.4 and 1.6 nM, respectively).

40 **Preparation of Immunoconjugates***nBT062-DM1 and huC242-DM1*

**[0181]** The thiol-containing maytansinoid DM1 was synthesized from the microbial fermentation product ansamitocin P-3, as previously described by Chari (Chari et al., Cancer Res. 1 (1992), 127). Preparation of humanized C242 (huC242) (Roguska et al., PNAS, 91 (1994), 969) has been previously described. Antibody-drug conjugates were prepared as previously described (Liu et al., PNAS, 93 (1996), 8618). An average of 3.5 DM1 molecules was linked per antibody molecule.

50 *nBT062-DM4*

**[0182]** BT062 is an antibody-drug conjugate composed of the cytotoxic maytansinoid drug, DM4, linked via disulfide bonds through a linker to the nBT062 chimerized monoclonal antibody. Maytansinoids are anti-mitotics that inhibit tubulin polymerization and microtubule assembly (Remillard et al., Science 189 (1977), 1002). Chemical and schematic representations of BT062 (nBT062-DM4) are shown in **FIGS. 1 and 2**.

*Synthesis of DM4*

[0183] DM4 is prepared from the well known derivative maytansinol (Kupchan et al., J. Med. Chem., 21 (1978), 31). Maytansinol is prepared by reductive cleavage of the ester moiety of the microbial fermentation product, ansamitocin P3, with lithium trimethoxyaluminum hydride (see FIG. 3).

[0184] DM4 is synthesized by acylation of maytansinol with N-methyl-N-(4-methyldithiopentanoyl)-L-alanine (DM4 side chain) in the presence of dicyclohexylcarbodiimide (DCC) and zinc chloride to give the disulfide-containing maytansinoid DM4-SMe. The DM4-SMe is reduced with dithiothreitol (DTT) to give the desired thiol-containing maytansinoid DM4 (see FIG. 4 for the DM4 process flow diagram).

*Immunoconjugate BT062*

[0185] The procedure for the preparation of nBT062-DM4 is outlined in FIG. 5. The nBT062 antibody is modified with N-succinimidyl-4-(2-pyridyldithio) butyrate (SPDB linker) to introduce dithiopyridyl groups. DM4 is mixed with the modified antibody at a concentration in excess of the equivalents of dithiopyridyl groups. The BT062 conjugate forms by a disulfide exchange reaction between the thiol group of DM4 and the dithiopyridyl groups introduced into the antibody via the linker. Purification by chromatography and diafiltration removes the low molecular weight reactants (DM4) and reaction products (thiopyridine), as well as aggregates of conjugated antibody, to produce the bulk drug substance.

**20 FACS analysis and WST cytotoxicity assays***FACS analysis*

[0186] OPM-2 cells are plasma cell leukemia cell lines showing highly expressing CD138. OPM-2 cells were incubated with nBT062, nBT062-SPDB-DM4, nBT062-SPP-DM1 or nBT062-SMCC-DM1 at different concentrations (indicated in FIG. 6). The cells were washed and CD138-bound antibody or conjugates were detected using a fluorescence-labeled secondary antibody in FACS analysis. The mean fluorescence measured in these experiments was plotted against the antibody concentration.

*30 Cell viability assay*

[0187] CD138<sup>+</sup> MOLP-8 cells were seeded in flat bottom plates at 3000 cells/well. CD138-BJAB control cells were seeded at 1000 cells/well. The cells were treated with nBT062-SPDB-DM4, nBT062-SPP-DM1 or nBT062-SMCC-DM1 at different concentrations (indicated in FIG. 7) for five days. WST reagent (water-soluble tetrazolium salt, ROCHE) was added in order to measure cell viability according to the manufacturer's instruction (ROCHE). The reagent was incubated for 7.5 h on MOLP-8 cells and for 2 h on BJAB cells. The fraction of surviving cells was calculated based on the optical densities measured in a microplate reader using standard procedures.

*40 Discussion*

[0188] Binding of nBT062-SPDB-DM4, nBT062-SPP-DM1, nBT062-SMCC-DM1 or nBT062 was analyzed by FACS. CD138<sup>+</sup> OPM-2 as target cells were incubated with nBT062 or immunoconjugates and cell-bound molecules were detected using a fluorescence-labeled secondary antibody. In FIG. 6, the mean fluorescences as measure for the amount of cell bound antibody is plotted against different antibody or conjugate concentrations. The results show, that nBT062-SPDB-DM4, nBT062-SPP-DM1 and nBT062-SMCC-DM1 show very similar binding characteristics. In addition, the results strongly suggest that the binding characteristics of the unconjugated antibody is not affected by the conjugated toxins.

[0189] In cell viability assays, the cytotoxic activity of the antibody against CD138<sup>+</sup> MOLP-8 target cells and against CD138- BJAB B-lymphoblastoma control cells were analyzed. Both cell lines were seeded in flat-bottom plates and incubated with increasing concentrations of the immunoconjugates. Unconjugated antibody was used as a control. The cytotoxic activity was analyzed five days after addition of the immunoconjugates by using WST reagent in order to measure cell viability. In FIG. 7 (A)-(C), the fraction of surviving cells relative to control cells treated with vehicle control is plotted against increasing immunoconjugate concentrations. The results show that cytotoxic activity of nBT062-SPDB-DM4, nBT062-SPP-DM1 and nBT062-SMCC-DM1 against MOLP-8 cells is very similar. As expected, CD138- BJAB control cells were not killed by the immunoconjugates, indicating that all immunoconjugates act via cell specific binding to CD138. In competition experiments, in which MOLP-8 cells were preincubated with a molar excess of unconjugated nBT062. Preincubation substantially blocked the cytotoxicity of nBT062-SPDB-DM4, providing further evidence that the immunoconjugates kill the cells via specific binding to CD138 onto the cell surface (FIG. 7 (D)).

**Xenograft mouse experiments**

**[0190]** To evaluate the importance of CD138 targeting on the anti-tumor activity of antibody-maytansinoid conjugates of a human chimeric version of the B-B4 antibody, nBT062, xenograft mouse experiments were performed. Two versions of nBT062-maytansinoid conjugates were prepared that may differ in the chemical stability of their disulfide linkages (nBT062-SPP-DM1 and nBT062-SPDB-DM4). The anti-tumor activity of these antibody-drug conjugates was compared to the activity of the **B-B4-SPP-DM1** conjugate (comprising the murine parental antibody), as well as unconjugated free maytansinoid (DM4), native unmodified nBT062 antibody, and a non-targeting (irrelevant) IgG1-maytansinoid conjugate. The conjugates were evaluated in a CD138-positive xenograft model (MOLP-8) of human multiple myeloma in severe combined immunodeficient (SCID) mice.

**[0191]** In these mice, subcutaneous tumors were established (female CB.17 SCID mice) by inoculation with MOLP-8 cell suspensions. Treatment with a single bolus intravenous injection was conducted when tumor volumes reached an average 113 mm<sup>3</sup>. Changes in tumor volume and body weight were monitored twice per week. Experiments were carried out over 68 days after tumor cell inoculation.

**Xenograft mouse experiments A***Mice*

**[0192]** Female CB.17 SCID mice, five weeks old, were obtained from Charles River Laboratories.

*Human tumor cell lines*

**[0193]** MOLP-8, a human multiple myeloma cell line, was supplied from ATCC. MOLP-8 cells, which express the CD138 antigen on their cell surface and develop xenograft tumors in SCID mice, were maintained in RPMI-1640 medium supplemented with 4 mM L-glutamine (Biowhittaker, Walkersville, MD), 10% fetal bovine serum (Hyclone, Logan, Utah) and 1% streptomycin/penicillin, at 37°C in a humidified atmosphere that contained 5% CO<sub>2</sub>.

**PART I***Tumor growth in mice*

**[0194]** Each mouse was inoculated with 1x10<sup>7</sup> MOLP-8 cells subcutaneously into the area under the right shoulder. The total volume was 0.2 ml per mouse, in which the ratio of serum-free medium to matrigel (BD Bioscience, Bedford, MA) was 1/1 (v/v). Prior to treatment, the xenograft tumors were monitored daily and were allowed to become established. The tumor volume reached approximately 113 mm<sup>3</sup> about 11 days after tumor cell inoculation. Tumor take rate of CB.17 SCID mice was 100%.

**[0195]** Eleven days after tumor cell inoculation, 42 mice were selected based on tumor volumes and body weights. The tumor volume was in a range of 68.2 to 135.9 mm<sup>3</sup>. The forty-two mice were randomly divided into seven groups (A-G) of six animals each based on tumor volume.

**[0196]** Each of six mice in Group A received 200 µl of PBS as vehicle control. Each mouse in group B received 13.8 mg/kg of nBT062 naked antibody. This dose is equivalent to the amount of nBT062 antibody component in 250 µg/kg of linked maytansinoid. The ratio of molecular weights of maytansinoids to nBT062 antibody in a conjugate molecule is approximate 1/55. Each mouse in Group C received 250 µg/kg of DM4. Each mouse in Group D received 250 µg/kg of huC242-DM4. Mice in groups E, F and G received 250 µg/kg of nBT062-SPDB-DM4, **B-B4-SPP-DM1** and nBT062-SPP-DM1 each, respectively.

**[0197]** All agents were intravenously administered as a single bolus injection through a lateral tail vein with a 1 ml syringe fitted with a 27 gauge, ½ inch needle. Prior to administration, the stock solutions of nBT062 antibody, nBT062-SPDB-DM4 and nBT062-SPP-DM1 were diluted with sterile PBS to concentrations of 2 mg/ml, 28.1 µg/ml and 28.1 µg/ml, respectively, so that the injected volume for each mouse was between 120-220 µl.

**PART II**

**[0198]** In a second set of experiments, MOLP-8 cells (1.5x10<sup>7</sup> cells per mouse), suspended in a 50:50 mixture of serum free media and matrigel were injected subcutaneously in the area under the right shoulder in 100 µl. Tumor volumes reached about 80 mm<sup>3</sup> at day 11 and the mean of the controls was about 750 mm<sup>3</sup> at day 25, post cell inoculation. The tumor doubling time was estimated to be 4.58 days. Each mouse in the control group (n=6) received 0.2 ml of sterile PBS administered into the lateral tail vein (i.v.) in a bolus injection. All treatment doses were based on conjugated

maytansinoid. Nine groups (n=6) were treated with a single intravenous injection of nBT062-SMCC-DM1, nBT062-SPDB-DM4, or nBT062-SPP-DM1, each at doses of 450, 250 and 100  $\mu\text{g}/\text{kg}$ . An additional group (n=6) received 250  $\mu\text{g}/\text{kg}$  nBT062-SMCC-DM1 in a repeated dosing (weekly for five weeks). Mice were randomized into eleven groups (n=6) by tumor volume using the LabCat Program. The tumor volumes ranged from 40.0 to 152.5  $\text{mm}^3$ . The mice were dosed based on the individual body weight.

5 [0199] Tumor size was measured twice per week in three dimensions using the LabCat System (Tumor Measurement and Tracking, Innovative Programming Associated, Inc., Princeton, NJ). The tumor volume in  $\text{mm}^3$  was calculated using the methodology described in Tomayko et al., *Cancer Chemother. Pharmacol.*, 24 (1989), 148:

10 
$$\text{Volume} = \text{Length} \times \text{Width} \times \text{Height} \times \frac{1}{2}$$

Log<sub>10</sub> cell kill was calculated with the formula described in Bissery et al., *Cancer Res.*, 51 (1991), 4845:

15 
$$\text{Log}_{10} \text{ cell kill} = (\text{T}-\text{C}) / \text{T}_d \times 3.32$$

20 where (T-C) or tumor growth delay, is the median time in days required for the treatment group (T) and the control group (C) tumors, to reach a predetermined size (600  $\text{mm}^3$ ). T<sub>d</sub> is the tumor doubling time, based on the median tumor volume in the control mice, and 3.32 is the number of cell doublings per log of cell growth.

### Results

25 [0200] The tumor growth in individual mice is shown in FIGS. 8 and 9. The mean (+/- SD) tumor growth for each group is shown in FIG. 10.

[0201] As compared with tumor growth in the PBS-treated animals, treatment with nBT062 antibody, unconjugated free DM4 or the irrelevant non-targeting conjugate huC242-DM4 did not cause any significant inhibition of tumor growth.

30 [0202] All three CD138-targeting conjugates, nBT062-SPDB-DM4, **B-B4-SPP-DM1** and nBT062-SPP-DM1, at a dose of 250  $\mu\text{g}/\text{kg}$  caused marked delay in tumor growth. Based on the mean tumor volumes measured in the treatment groups, the DM4 conjugate nBT062-SPDB-DM4 was the most active one, while the nBT062-SPP-DM1 conjugate showed slightly increased activity as compared to its murine counterpart **B-B4-SPP-DM1** (FIG. 10). The results obtained in individual mice show in addition that the anti-tumor activity obtained with **B-B4-SPP-DM1** is more heterogeneously and therefore less predictable than that measure in mice treated with nBT062-SPP-DM1. In terms of homogeneity of anti tumor activity, the other conjugate that uses nBT062 as targeting antibody nBT062-SPDB-DM4 behaved similar to nBT062-SPP-DM1.

35 [0203] No body weight reduction was observed in any treatment group suggesting that the treatments were well tolerated.

### Discussion

40 [0204] The results of the analysis of three CD138-targeting conjugates in experimental animals demonstrate the importance of targeted delivery for the anti-tumor activity. While the maytansinoid conjugates of the human chimeric nBT062 and the murine B-B4 antibodies show significant activity as measured by log cell kill, there was no significant impact on tumor growth from treatment with unconjugated DM4, unmodified native huBT062 antibody, or a non-targeting control conjugate (huC242-DM4).

45 [0205] The immunoconjugate prepared from the human chimeric antibody, nBT062-SPP-DM1, gave slightly higher anti-tumor activity than the conjugate prepared from its murine counterpart, B-B4-SPP-DM1. In addition, treatment with nBT062-SPP-DM1 and nBT062-SPDB-DM4 resulted in more homogenous responses in individual mice as compared to treatment with B-B4-SPP-DM1. The high binding variation of **B-B4-SPP-DM1** explained that the measurement of the median tumor volume (+/- SD) of MOLP-8 human multiple myeloma xenografts in CB.17 SCID mice overtime (days) post-inoculation actually provided for relatively better results for **B-B4-SPP-DM1** than for nBT062-SPP-DM1 (data not shown). This feature of immunoconjugates using nBT062 as a targeting antibody seems to be beneficial especially for therapeutic use of the conjugates.

55 [0206] Lastly, the most potent of the maytansinoid conjugates, following single iv administration in the MOLP-8 xenograft models in SCID mice, was nBT062-SPDB-DM4.

*Bystander killing (cell viability assay)*

5 [0207] CD138<sup>+</sup> OPM2 cells and CD138<sup>-</sup> Namalwa cells were seeded in round bottom plates either in separate wells or in coculture. The cells were treated with nBT062-SPDB-DM4 at concentrations ranging from  $|x| \times 10^6$  to  $1 \times 10^9$  M. The fraction of viable cells was detected using WST reagent (water-soluble tetrazolium salt, ROCHE) according to the manufacturer's instruction (ROCHE). The fraction of surviving cells was calculated based on the optical densities measured in a microplate reader using standard procedures.

10 *Discussion*

10 [0208] Bystander killing of non-target cells in close proximity (as present in round bottom wells) to multiple myeloma cells upon nBT062-SPDB-DM4 treatment was analysed in an in vitro study in which CD138-positive OPM2 cells were cultured in coculture with CD138-negative Namalwa cells (FIG. 13). Generally, while CD138-positive cells are efficiently killed by nBT062-SPDB-DM4, CD138-negative cells were not affected by the conjugate. In the coculture in round bottom wells, however, nBT062-SPDB-DM4 also killed the antigen-negative cells in close proximity to the antigen-positive cells (an effect that is often referred to as bystander killing). Kovtun et al. (2006) discussed that bystander killing mediated by maytansinoid conjugates occurs only in close proximity to antigen-positive cells. Kovtun et al. (2006), also discusses the importance of the linker of the immunoconjugate. In vivo, bystander killing may contribute to 1) the eradication of tumour cells that heterogeneously express CD138, 2) the destruction of the tumour microenvironment by the killing of tumour stroma cells, and 3) the prevention of the selection of CD138-negative nBT062-SPDB-DM4-resistant cells.

15 [0209] The bystander effect is of particular importance if the activity of an immunoconjugate is impaired by a target antigen that is expressed in tumors in a heterogeneous fashion. If this is the case, a particular cell of a tumor expresses, if at all, the antigen not in amount that would allow effective direct targeting and killing of said cell by the respective immunoconjugate. The anti-tumor efficacy of nBT062-SPDB-DM4 on CD138-negative cells in coculture with CD138-positive cells clarified that the presence of target cells influences, under the appropriate circumstances, the cytotoxic activity of nBT062-SPDB-DM4 towards non-target cells.

**Xenograft mouse experiments B**

30 [0210] In this set of experiments, eighty-five mice were inoculated with MOLP-8 cells ( $1.5 \times 10^7$  cells/mouse) subcutaneously in the right shoulder. Tumor take rate was 100%. Sixty-six SCID mice bearing bulky MOLP-8 tumors with a mean tumor volume of about  $80 \text{ mm}^3$  were randomized into eleven treatment groups (n=6). Mice were treated with a single dose of one of three conjugates (nBT062-SMCC-DM1, nBT062-SPDB-DM4 or nBT062-SPP-DM1). An additional group received five weekly doses of nBT062-SMCC-DM1 and a control group received a single dose of PBS. Mean tumor volumes are shown in FIG. 11A. A dose response was established for each conjugate. A median tumor volume of  $750 \text{ mm}^3$  in the PBS-treated animals was reached on day 25. Tumor doubling time determined by the best-fit linear regression curve fit on a log-linear plot of control tumor growth was 4.58 days. Animals treated with nBT062-SPDB-DM4 at  $450 \mu\text{g/kg}$  had the highest log cell kill (LCK=2.89), followed by animals treated with nBT062-SMCC-DM1 at  $250 \mu\text{g/kg}$  weekly dosing (LCK=2.1; see Table 5). Comparison of the mean tumor growth curves for the treatment groups by repeated measures ANOVA performing Dunnett's Multiple Comparisopn Test showed a significant difference between the PBS control group and  $450 \mu\text{g/kg}$  nBT062-SPDB-DM4 ( $p < 0.01$ ),  $250 \mu\text{g/kg}$  nBT062-SPDB-DM4 ( $p < 0.05$ ) and five weekly doses of  $250 \mu\text{g/kg}$  nBT062-SMCC-DM1 ( $p < 0.05$ ). No partial or complete tumor regression in any of the treatment groups occurred with the exception of one animal receiving  $450 \mu\text{g/kg}$  nBT062-SPDB-DM4, which had partial regression of the tumor until day 85 post-inoculation.

45

**Table 5.** Log cell kill (LCK) values as measure for anti-tumor activity of different nBT062-DMx conjugates in different dosing schemes. Refer to the Materials and methods section for information on calculation of LCK values.

Test Material	Dose ( $\mu\text{g/kg}$ )	LCK	Dosing
PBS			single dose
nBT062-SMCC-DM1	450	0.85	single dose
nBT062-SMCC-DM1	250	0.53	single dose
nBT062-SMCC-DM1	100	0	single dose
nBT062-SPDB-DM4	450	2.89	single dose
nBT062-SPDB-DM4	250	1.05	sinle dose

(continued)

Test Material	Dose ( $\mu\text{g/kg}$ )	LCK	Dosing
5 nBT062-SPDB-DM4	100	0.39	single dose
10 nBT062-SPP-DM1	450	0.8	single dose
nBT062-SPP-DM1	250	0.39	single dose
nBT062-SPP-DM1	100	0.2	single dose
15 nBT062-SMCC-DM1	250	2.1	weekly for 5 weeks

***In vivo efficacy of nBT062-SPDB-DM4 and nBT062-SPP-DM1 in the bone marrow environment***15 *Preparation of SCID mice having human fetal bone implants*

5 [0211] Human fetal long bones (human fetal bone chips) were implanted into the upper body of CB17 SCID-mice (SCID-hu) as previously described (Urashima et al., 1997) and thus provided for a model in mouse for the homing of human MM cells to human BM cells.

20 *Treatment regime (SCID-hu/INA-6 mice)*

25 [0212] 4 weeks following bone implantation,  $2.5 \times 10^6$  INA-6 cells in a final volume of 100  $\mu\text{L}$  RPMI-1640 cell culture medium were injected directly into the human bone marrow cavity in the SCID-hu mice described above. An increase in the levels of soluble human IL-6 receptor (shuIL-6R), which is released by INA-6 cells, was used as a parameter of MM cell growth and disease burden.

30 [0213] Mice developed measurable serum shuIL-6R approximately 4 weeks following INA-6 cell injection and then received 0.176 mg conjugate or vehicle control via tail vein injection weekly for 7 weeks. After each treatment, blood samples were collected and measured for shuIL-6R levels by an enzyme-linked immunosorbent assay (ELISA, R&D Systems, Minneapolis, MN). The results are depicted in **Fig. 12**.

*Discussion*

35 [0214] Interleukin 6 (IL-6) is a growth and survival factor for multiple myeloma cells. INA-6 is an IL-6-dependent human myeloma cell line, which also requires bone marrow stromal cells (BMSC) to proliferate. INA-6 cell lines produce soluble IL-6 receptor (shuIL-6R). An increase in the levels of shuIL-6R can be used as a parameter of MM cell growth and disease burden.

40 [0215] Thus, the sCID-hu/INA-6 mice provide a model for multiple myeloma cells growing in their normal bone marrow environment. The tumor cells of this model, which directly interact with the human bone marrow, closely resemble the situation in patients, in which tumor cell growth is also promoted by the presence of stromal cells. As INA-6 cells release soluble human interleukin-6 receptor (shuIL-6R), serum concentrations of this protein can be used as a measure for tumor cell load in these mice. The *in vivo* potency of nBT062-SPDB-DM4 and nBT062-SPP-DM1 were tested in this environment.

45 [0216] Treatment of SCIDhu/INA-6 mice with weekly i.v. administrations of nBT062-SPDB-DM4 or nBT062-SPP-DM1 for seven weeks induced efficient tumour regression, as detected by a decrease in serum shuIL-6R levels relative to the control, indicating good efficacy of the conjugates even in the environment of human bone marrow, which reflect the relevant situation in patients (Fig. 12).

50 [0217] It will be appreciated that the methods and compositions of the instant invention can be incorporated in the form of a variety of embodiments, only a few of which are disclosed herein. Thus, the described embodiments are illustrative.

**Bibliography****[0218]**

55 Akkina RK, Rosenblatt JD, Campbell AG, Chen IS, Zack JA. Modeling human lymphoid precursor cell gene therapy in the SCID-hu mouse. *Blood*. 1994;84:1393-1398.

Armour KL, Clark MR, Hadley AG, et al. Recombinant human IgG molecules lacking Fc gamma receptor I binding and monocyte triggering activities. *Eur J Immunol*. 1999; 29(8):2613-24.

5 Anderson KC, Kyle RA, Dalton WS, Landowski T, Shain K, Jove R, Hazlehurst L, Berenson J. Multiple Myeloma: New Insights and Therapeutic Approaches. *Hematology* 2000;147-165.

Anttonen A, Heikkila P, Kajanti M, Jalkanen M, Joensuu H. High syndecan-1 expression is associated with favourable outcome in squamous cell lung carcinoma treated with radical surgery. *Lung Cancer*. 2001 Jun; 32(3):297-305.

10 Barbareschi M, Maisonneuve P, Aldovini D, Cangi MG, Pecciarini L, Angelo Mauri F, Veronese S, Caffo O, Lucenti A, Palma PD, Galligioni E, Doglioni C. High syndecan-1 expression in breast carcinoma is related to an aggressive phenotype and to poorer prognosis. *Cancer*. 2003 Aug 1;98(3):474-83.

15 Bataille R, Jégo G, Robillard N, Barillé-Nion S, Harousseau JL, Moreau P, Amiot M, Pellat-Deceunynck C. The phenotype of normal, reactive and malignant plasma cells. Identification of "many and multiple myelomas" and of new targets for myeloma therapy. *Haematologica*. 2006 Sep;91(9):1234-40. Review.

Bernfield M, Kokenyesi R, Kato M, Hinkes MT, Spring J, Gallo RL, Lose EJ. Biology of the syndecans: a family of transmembrane heparan sulfate proteoglycans. *Annu Rev Cell Biol*. 1992;8:365-393.

20 Beste G, Schmidt FS, Stibora T, Skerra A. Small antibody-like proteins with prescribed ligand specificities derived from the lipocalin fold. *Proc Natl Acad Sci USA*. 1999; 96, 1898-1903.

Bhattacharyya B, Wolff J. Maytansine binding to the vinblastine sites of tubulin. *FEBS Lett*. 1977;75:159-162.

25 Bisping G, Kropff M, Wenning D, Dreyer B, Bessonov S, Hilberg F, Roth GJ, Munzert G, Stefanic M, Stelljes M, Scheffold C, Müller-Tidow C, Liebisch P, Lang N, Tchinda J, Serve HL, Mesters RM, Berdel WE, Kienast J. Targeting receptor kinases by a novel indolinone derivative in multiple myeloma: abrogation of stroma-derived interleukin-6 secretion and induction of apoptosis in cytogenetically defined subgroups. *Blood*. 2006 Mar 1;107(5):2079-89. Epub 2005 Nov 8.

Blättler WA and Chari RVJ. Drugs to Enhance the Therapeutic Potency of Anticancer Antibodies: Antibody-Drug Conjugates as Tumor-Activated Prodrugs. In: Ojima, I., Vite, G.D. and Altmann, K.-H., Editors, 2001. *Anticancer Agents-Frontiers in Cancer Chemotherapy*, American Chemical Society, Washington, DC, pp. 317-338.

30 Bross PF, Beitz J, Chen G, Chen XH, Duffy E, Kieffer L, Roy S, Sridhara R, Rahman A, Williams G, Pazdur R. Approval summary: gemtuzumab ozogamicin in relapsed acute myeloid leukemia. *Clin Cancer Res*. 2001;7:1490-1496.

Carbone A, Gaidano G, Gloghini A, Ferlito A, Rinaldo A, Stein H. AIDS-related plasma- blastic lymphomas of the oral cavity and jaws: a diagnostic dilemma. *Ann. Otol. Rhinol. Laryngol.* 1999; 108: 95-99.

40 Carlsson J, Drevin H, Axen R. Protein thiolation and reversible protein-protein conjugation. N-succinimidyl-3-(2-pyridyldithio)propionate, a new heterobifunctional reagent. *Biochem J* 1978; 173: 723-737.

45 Carter P. Improving the efficacy of antibody-based cancer therapies. *Nat Rev Cancer*. 2001;1:118-129.

Chari RV, Martell BA, Gross JL, Cook SB, Shah SA, Blattler WA, McKenzie SJ, Goldmacher VS. Immunoconjugates containing novel maytansinoids: promising anticancer drugs. *Cancer Res*. 1992;52:127-131.

50 Chari RV, Jackel KA, Bourret LA, Derr SM, Tadayoni BM, Mattocks KM, Shah SA, Liu C, Blättler WA and Goldmacher VS. Enhancement of the selectivity and antitumor efficacy of a CC-1065 analogue through immunoconjugate formation. *Cancer Res*. 1995; 55: 4079-4084.

55 Charnaux N, Brûlé S, Chaigneau T, Saffar L, Sutton A, Hamon M, Prost C, Lievre N, Vita C, Gattegno L. RANTES (CCL5) induces a CCR5-dependent accelerated shedding of syndecan-1 (CD138) and syndecan-4 from HeLa cells and forms complexes with the shed ectodomains of these proteoglycans as well as with those of CD44. *Glycobiology*. 2004 Sep 8 [Epub ahead of print]

Chen BP, Galy A, Kyoizumi S, Namikawa R, Scarborough J, Webb S, Ford B, Cen DZ, Chen SC. Engraftment of human hematopoietic precursor cells with secondary transfer potential in SCID-hu mice. *Blood*. 1994;84:2497-2505.

5 Chilos M, Adami F, Lestani M, Montagna L, Cimarosta L, Semenzato G, Pizzolo G, Menestrina F. CD138/syndecan-1: a useful immunohistochemical marker of normal and neoplastic plasma cells on routine trephine bone marrow biopsies. *Mod Pathol*. 1999;12:1101-1106.

10 Clement C, Vooijs, W.C., Klein, B., and Wijdenes, J. In: al. SFSe, ed. *J. Leukocyte Typing V*. Oxford: Oxford University Press; 1995:714-715.

15 Couturier O, Faivre-Chauvet A ; Filippovich IV; Thedréz P, Sai-Maurel C; Bardiés M; Mishra AK; Gauvrit M; Blain G; Apostolidis C; Molinet R; Abbe JC; Bateille R; Wijdenes J; Chatal JF; Cherel M; Validation of 213Bi-alpha radioimmunotherapy for multiple myeloma. *Clinical Cancer Research* 5(10 Suppl.) (Oct 1999) 3165s-3170s.

20 Davies EJ et al. , Blackhall FH, Shanks JH, David G, McGown AT, Swindell R, Slade RJ, Martin-Hirsch P, Gallagher JT, Jayson GC. Distribution and Clinical Significance of Heparan Sulfate Proteoglycans in Ovarian Cancer *Clin Cancer Res*. 2004; 10(15):5178-86.

25 Dhodapkar MV, Abe E, Theus A, Lacy M, Langford JK, Barlogie B, Sanderson RD. Syndecan-1 is a multifunctional regulator of myeloma pathobiology: control of tumor cell survival, growth, and bone cell differentiation. *Blood*. 1998;91:2679-2688.

Dore JM, Morard F, Vita N, Wijdenes J. Identification and location on syndecan-1 core protein of the epitopes of B-B2 and B-B4 monoclonal antibodies. *FEBS Lett*. 1998;426:67-70.

30 Dowell JA, Korth-Bradley J, Liu H, King SP, Berger MS. Pharmacokinetics of gemtuzumab ozogamicin, an antibody-targeted chemotherapy agent for the treatment of patients with acute myeloid leukemia in first relapse. *J Clin Pharmacol*. 2001;41:1206-1214.

35 Edinger M, Sweeney TJ, Tucker AA, Olomu AB, Negrin RS, Contag CH. Noninvasive assessment of tumor cell proliferation in animal models. *Neoplasia*. 1999;1:303-310.

Gattei V, Godeas C, Degan M, Rossi FM, Aldinucci D, Pinto A. Characterization of Anti-CD138 monoclonal antibodies as tools for investigating the molecular polymorphism of syndecan-1 in human lymphoma cells. *Br J Haematol*. 1999;104:152-162.

40 Hamann PR, Hinman LM, Beyer CF, Lindh D, Upeslasis J, Flowers DA, Bernstein I. An anti-CD33 antibody-calicheamicin conjugate for treatment of acute myeloid leukemia. Choice of linker. *Bioconjug Chem*. 2002;13:40-46.

Han I, Park H, Oh ES. New insights into syndecan-2 expression and tumourigenic activity in colon carcinoma cells. *J Mol Histol*. 2004; 35(3):319-26.

45 Hideshima T, Catley L, Yasui H, Ishitsuka K, Raje N, Mitsiades C, Podar K, Munshi NC, Chauhan D, Richardson PG, Anderson KC. Perifosine, an oral bioactive novel alkylphospholipid, inhibits Akt and induces in vitro and in vivo cytotoxicity in human multiple myeloma cells. *Blood* 2006;107(10):4053-62.

Hideshima T, Mitsiades C, Tonon G, Richardson PG, Anderson KC. Understanding multiple myeloma pathogenesis in the bone marrow to identify new therapeutic targets. *Nat Rev Cancer* 2007;7(8):585-98.

50 Horvathova M, Gaillard, J.-P., Liutard, J., Duperray, C., Lavabre-Bertrand, T., Bourquard, P et al. In: al. SFSe, ed. *Leucocyte Typing V*. Oxford: Oxford University Press; 1995:713-714.

Kovtun YV, Audette CA, Ye Y, Xie H, Ruberti MF, Phinney SJ, et al. Antibody-drug conjugates designed to eradicate tumors with homogeneous and heterogeneous expression of the target antigen. *Cancer Res* 2006;66 (6):3214-21.

55 Krebs B, Rauchenberger R, Reiffert S, Rothe C, Tesar M, Thomassen E, Cao M, Dreier T, Fischer D, Hoss A et al. High-throughput generation and engineering of recombinant human antibodies. 2001. *J. Immunol. Methods* 254, pp. 67-84.

Kupchan SM, Sneden AT, Branfman AR, Howie GA, Rebhun LI, McIvor WE, Wang RW, Schnaitman TC. Structural requirements for antileukemic activity among the naturally occurring and semisynthetic maytansinoids. *J Med Chem.* 1978;21:31-37.

5 Kyoizumi S, Baum CM, Kaneshima H, McCune JM, Yee EJ, Namikawa R. Implantation and maintenance of functional human bone marrow in SCID-hu mice. *Blood.* 1992;79:1704-1711.

10 Kyoizumi S, Murray LJ, Namikawa R. Preclinical analysis of cytokine therapy in the SCID-hu mouse. *Blood.* 1993;81:1479-1488.

15 Langford JK, Stanley MJ, Cao D, Sanderson RD. Multiple heparan sulfate chains are required for optimal syndecan-1 function. *J Biol Chem.* 1998 Nov 6;273(45):29965-71.

20 Liu C, Tadayoni BM, Bourret LA, Mattocks KM, Derr SM, Widdison WC, Kedersha NL, Ariniello PD, Goldmacher VS, Lambert JM, Blattler WA, Chari RV. Eradication of large colon tumor xenografts by targeted delivery of maytansinoids. *Proc Natl Acad Sci USA.* 1996;93:8618-8623.

25 McCune JM, Namikawa R, Kaneshima H, Shultz LD, Lieberman M, Weissman IL. The SCID-hu mouse: murine model for the analysis of human hematolymphoid differentiation and function. *Science.* 1988;241:1632-1639.

30 Mennerich D, Vogel A, Klaman I, Dahl E, Lichtner RB, Rosenthal A, Pohlenz HD, Thierauch KH, Sommer A. Shift of syndecan-1 expression from epithelial to stromal cells during progression of solid tumours. *Eur J Cancer.* 2004 Jun; 40(9): 1373-82.

35 Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods.* 1983;65:55-63.

40 Munshi NC, Longo DL, Anderson KC. Plasma cell disorders. In: Braunwald E, Fauci AS, Kasper DL, Hauser SL, Longo DL, Jameson JL, editors. *Harrison's Principles of Internal Medicine.* 16th ed. New York : McGraw-Hill Medical Publishing Division; 2008. p. 700-7.

45 Namikawa R, Ueda R, Kyoizumi S. Growth of human myeloid leukemias in the human marrow environment of SCID-hu mice. *Blood.* 1993;82:2526-2536.

50 O'Connell FP, Pinkus JL, Pinkus GS. CD138 (Syndecan-1), a Plasma Cell Marker Immunohistochemical Profile in Hematopoietic and Nonhematopoietic Neoplasms. *Am J Clin Pathol* 2004; 121:254-263.

55 Ojima I, Geng X, Wu X, Qu C, Borella CP, Xie H, Wilhelm SD, Leece BA, Bartle LM, Goldmacher VS and Chari RV. Tumor-specific novel taxoid-monoclonal antibody conjugates. 2002. *J. Med. Chem.* 45, pp. 5620-5623.

60 Olafsen, T, Cheung, CC, Yazaki, PJ, Li L, Sundaresan G, Gambhir SS, Sherman, MA, Williams, LE, Shively, JE, Raubitschek, AA, and Wu, AM. Covalent disulfide-linked anti-CEA diabody allows site-specific conjugation and radiolabeling for tumor targeting applications. 2004; *Prot. Eng. Design & Selection* 17:1: 21-27.

65 Orosz Z, Kopper L. Syndecan-1 expression in different soft tissue tumours. *Anticancer Res.* 2001; 21(1B):733-7.

70 Padlan, EA. A possible procedure for reducing the immunogenicity of antibody variable domains while preserving their ligand-binding properties. *Mol. Immunol.* 1991; 28: 489-498.

75 Payne G. Progress in immunoconjugate cancer therapeutics. *Cancer Cell.* 2003;3:207-212.

80 Pegram MD, Lipton A, Hayes DF, Weber BL, Baselga JM, Tripathy D, Baly D, Baughman SA, Twaddell T, Glaspy JA and Slamon DJ. Phase II study of receptor-enhanced chemosensitivity using recombinant humanized anti-p185HER2/neu monoclonal antibody plus cisplatin in patients with HER2/neu-overexpressing metastatic breast cancer refractory to chemotherapy treatment. 1998. *J. Clin. Oncol.* 16, pp. 2659-2671.

85 Rawstron AC, Owen RG, Davies FE, Johnson RJ, Jones RA, Richards SJ, Evans PA, Child JA, Smith GM, Jack AS, Morgan GJ. Circulating plasma cells in multiple myeloma: characterization and correlation with disease stage.

Br J Haematol. 1997;97:46-55.

Remillard S, Rebhun LI, Howie GA, Kupchan SM. Antimitotic activity of the potent tumor inhibitor maytansine. Science. 1975;189:1002-1005.

Roguska MA, Pedersen JT, Keddy CA, Henry AH, Searle SJ, Lambert JM, Goldmacher VS, Blattler WA, Rees AR, Guild BC. Humanization of murine monoclonal antibodies through variable domain resurfacing. Proc Natl Acad Sci USA. 1994;91:969-973.

Ross S, Spencer SD, Holcomb I, Tan C, Hongo J, Devaux B, Rangell L, Keller GA, Schow P, Steeves RM, Lutz RJ, Frantz G, Hillan K, Peale F, Tobin P, Eberhard D, Rubin MA, Lasky LA, Koeppen H. Prostate stem cell antigen as therapy target: tissue expression and in vivo efficacy of an immunoconjugate. Cancer Res. 2002 May 1;62(9):2546-53.

Ross JS, Gray K, Gray G, Worland PJ, Rolfe M. Anticancer Antibodies, Am J Clin Path. (4/17/2003).

Sanderson RD, Lalor P, Bernfield M. B lymphocytes express and lose syndecan at specific stages of differentiation. Cell Regul. 1989;1:27-35.

Sandhu JS, Clark BR, Boynton EL, Atkins H, Messner H, Keating A, Hozumi N. Human hematopoiesis in SCID mice implanted with human adult cancellous bone. Blood. 1996;88:1973-1982.

Sasaki A, Boyce BF, Story B, Wright KR, Chapman M, Boyce R, Mundy GR, Yoneda T. Bisphosphonate risedronate reduces metastatic human breast cancer burden in bone in nude mice. Cancer Res. 1995;55:3551-3557.

Schneider U, van Lessen A, Huhn D, Serke S. Two subsets of peripheral blood plasma cells defined by differential expression of CD45 antigen. Br J Haematol. 1997;97:56-64.

Schuurman J, Van Ree R, G. J. Perdok GJ, Van Doorn HR, Tan KY, Aalberse RC, Normal human immunoglobulin G4 is bispecific: it has two different antigen-combining sites, Immunology 1999; 97:693-698.

Sebestyen A, Berczi L, Mihalik R, Paku S, Matolcsy A, Kopper L. Syndecan-1 (CD138) expression in human non-Hodgkin lymphomas. Br J Haematol. 1999; 104(2):412-9.

Seftalioglu A, Karakus S. Syndecan-1/CD138 expression in normal myeloid, acute lymphoblastic and myeloblastic leukemia cells. Acta Histochem. 2003;105:213-221.

Seftalioglu A, Karakus S, Dundar S, Can B, Erdemli E, Irmak MK, Oztas E, Korkmaz C, Yazar F, Cavusoglu I. Syndecan-1 (CD138) expression in acute myeloblastic leukemia cells--an immuno electron microscopic study. Acta Oncol. 2003;42:71-74.

Senter PD, Doronina S, Cerveny C, Chace D, Francisco J, Klussman K, Mendelsohn B, Meyer D, Siegall CB, Thompson J et al. (2002). Cures and regressions of established tumors with monoclonal antibody auristatin conjugates. Abstract #2062, American Association for Cancer Res. (San Francisco, CA: American Association for Cancer Res.), 414.

Shields RL, Namenuk AK, Hong K, Meng YG, Rae J, Briggs J, Xie D, Lai J, Stadlen A, Li B, Fox JA, Presta LG.. High resolution mapping of the binding site on human IgG1 for Fc gamma RI, Fc gamma RII, Fc gamma RIII, and FcRn and design of IgG1 variants with improved binding to the Fc gamma R. J Biol Chem. 2001; 276(9):6591-604.

Sievers EL, Larson R.A., Stadtmauer, E.A., Estey, E., Lowenberg, B., Dombret, H., Karanes, C., Theobald, M., Bennett, J.M., Sherman, M.L. et al. Efficacy and safety of gemtuzumab ozogamicin in patients with CD33-positive acute myeloid leukemia in first relapse. 2001. J. Clin. Oncol. 19, pp. 3244-3254.

Sievers EL and Linenberger M. Mylotarg: antibody-targeted chemotherapy comes of age. 2001. Curr. Opin. Oncol. 13, pp. 522-527.

Smith R., Single chain antibody variable region fragments; [www.stanford.edu/~smithr/scfv.html](http://www.stanford.edu/~smithr/scfv.html) (last up-

dated on May, 2001).

5 Studnicka GM, Soares S, Better M, Williams RE, Nadell R, Horwitz AH. Human-engineered monoclonal antibodies retain full specific binding activity by preserving non-CDR complementarity-modulating residues. *Protein Eng.* 1994; 7(6): 805-814.

10 Tai YT, Li XF, Catley L, Coffey R, Breitkreutz I, Bae J, Song W, Podar K, Hideshima T, Chauhan D, Schlossman R, Richardson P, Treon SP, Grewal IS, Munshi NC, Anderson KC. Immunomodulatory drug lenalidomide (CC-5013, IMiD3) augments anti-CD40 SGN-40-induced cytotoxicity in human multiple myeloma: clinical implications. *Cancer Res.* 2005 Dec 15; 65(24):11712-20.

15 Tassone P, Goldmacher VS, Neri P, Gozzini A, Shammas MA, Whiteman KA, Hylander-Gans LL, Carrasco DR, Hideshima T, Shringarpure R, Shi J, Allam CK, Wijdenes J, Venuta S, Munshi NC, Anderson KC, Cytotoxic activity of the maytansinoid immunoconjugate B-B4-DM1 against CD138+ multiple myeloma cells, *Blood*, 2004, 104 (12), pp. 3688-3696.

20 Tolcher AW, Ochoa L, Hammond LA, Patnaik A, Edwards T, Takimoto C, Smith L, de Bono J, Schwartz G, Mays T, Jonak ZL, Johnson R, DeWitte M, Martino H, Audette C, Maes K, Chari RV, Lambert JM, Rowinsky EK. Cantuzumab mertansine, a maytansinoid immunoconjugate directed to the CanAg antigen: a phase I, pharmacokinetic, and biologic correlative study. *J Clin Oncol.* 2003;21:211-222.

25 Urashima M, Chen BP, Chen S, Pinkus GS, Bronson RT, Dedera DA, Hoshi Y, Teoh G, Ogata A, Treon SP, Chauhan D, Anderson KC. The development of a model for the homing of multiple myeloma cells to human bone marrow. *Blood.* 1997;90:754-765.

30 Vooijs WC, Post J, Wijdenes J, Schuurman HJ, Bolognesi A, Polito L, Stirpe F, Bast EJ, de Gast GC. Efficacy and toxicity of plasma-cell-reactive monoclonal antibodies B-B2 and B-B4 and their immunotoxins. *Cancer Immunol Immunother.* 1996;42:319-328.

35 Ward, E.S., D. Gussow, A.D. Griffiths, P.T. Jones, and G. Winter. Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli*. *Nature.* 1989. 341:544-546.

35 Wargalla UC, Reisfeld RA. Rate of internalization of an immunotoxin correlates with cytotoxic activity against human tumor cells. *Proc. Natl. Acad. Sci. USA.* 1989;86:5146-5150.

40 Wijdenes J, Vooijs WC, Clement C, Post J, Morard F, Vita N, Laurent P, Sun RX, Klein B, Dore JM. A plasmocyte selective monoclonal antibody (B-B4) recognizes syndecan-1. *Br J Haematol.* 1996;94:318-323.

45 Wijdenes J, Dore JM, Clement C, Vermot-Desroches C. CD138, *J Biol Regul Homeost Agents.* 2002 Apr-Jun;16(2):152-5.

50 Witzig TE, Kimlinger TK, Ahmann GJ, Katzmman JA, Greipp PR. Detection of myeloma cells in the peripheral blood by flow cytometry. *Cytometry.* 1996;26:113-120.

55 Xie H, Audette C, Hoffee M, Lambert JM, Blättler W. Pharmacokinetics and biodistribution of the antitumor immunoconjugate, cantuzumab mertansine (huC242-DM1), and its two components in mice. *J Pharmacol Exp Ther.* 2004 Mar;308(3):1073-82.

Yang M, Jiang P, An Z, Baranov E, Li L, Hasegawa S, Al-Tuwaijri M, Chishima T, Shimada H, Moossa AR, Hoffman RM. Genetically fluorescent melanoma bone and organ metastasis models. *Clin Cancer Res.* 1999;5:3549-3559.

Yang M, Baranov E, Jiang P, Sun FX, Li XM, Li L, Hasegawa S, Bouvet M, Al-Tuwaijri M, Chishima T, Shimada H, Moossa AR, Penman S, Hoffman RM. Whole-body optical imaging of green fluorescent protein-expressing tumors and metastases. *Proc Natl Acad Sci USA.* 2000;97:1206-1211.

Yang Y, MacLeod V, Dai Y, Khotskaya-Sample Y, Shriner Z, Venkataraman G, Sasisekharan R, Naggi A, Torri G, Casu B, Vlodavsky I, Suva LJ, Epstein J, Yaccoby S, Shaughnessy JD Jr, Barlogie B, Sanderson RD. The syndecan-1 heparan sulfate proteoglycan is a viable target for myeloma therapy. *Blood*. 2007 Sep 15;110(6):2041-8. Epub 2007 May 29.

5

Yoshitake S, Yamada Y, Ishikawa E, Masseyeff R. Conjugation of glucose oxidase from *Aspergillus niger* and rabbit antibodies using N-hydroxysuccinimide ester of N-(4-carboxycyclohexylmethyl)-maleimide. *Eur J Biochem* 1979;101:395-399.

10 SEQUENCE LISTING

[0219]

15 <110> Biotest AG  
ImmunoGen, Inc.

<120> IMMUNOCOCONJUGATES TARGETING CD138 AND USES THEREOF

20 <130> 319935.WO/JND/CJS

25 <150> 61/016,620 <151> 2007-12-26

<150> 61/087,466 <151> 2008-08-08

30 <150> 61/087,590 <151> 2008-08-08

<160> 13

35 <170> PatentIn version 3.5

<210> 1

<211> 448

<212> PRT

35 <213> Artificial Sequence

<220>

<223> Amino Acid sequence (predicted) of heavy chain of chimeric human/mouse antibody

40 <220>

<221> CDR1

<222> (31)..(35)

45 <220>

<221> CDR2

<222> (51)..(68)

50 <220>

<221> CDR3

<222> (99) .. (111)

55 <220>

<221> region IgG4

<222> (123)..(448)

<400> 1

**EP 2 242 772 B1**

Gln Val Gln Leu Gln Gln Ser Gly Ser Glu Leu Met Met Pro Gly Ala  
1 5 10 15

5 Ser Val Lys Ile Ser Cys Lys Ala Thr Gly Tyr Thr Phe Ser Asn Tyr  
20 25 30

10 Trp Ile Glu Trp Val Lys Gln Arg Pro Gly His Gly Leu Glu Trp Ile  
35 40 45

15

20

25

30

35

40

45

50

55

**EP 2 242 772 B1**

Gly Glu Ile Leu Pro Gly Thr Gly Arg Thr Ile Tyr Asn Glu Lys Phe  
50 55 60

5 Lys Gly Lys Ala Thr Phe Thr Ala Asp Ile Ser Ser Asn Thr Val Gln  
65 70 75 80

10 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys  
85 90 95

15 Ala Arg Arg Asp Tyr Tyr Gly Asn Phe Tyr Tyr Ala Met Asp Tyr Trp  
100 105 110

20 Gly Gln Gly Thr Ser Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro  
115 120 125

25 Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu Ser Thr  
130 135 140

30 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr  
145 150 155 160

35 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro  
165 170 175

40 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr  
180 185 190

45 Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn Val Asp  
195 200 205

50 His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser Lys Tyr  
210 215 220

55 Gly Pro Pro Cys Pro Ser Cys Pro Ala Pro Glu Phe Leu Gly Gly Pro  
225 230 235 240

60 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser  
245 250 255

65 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Gln Glu Asp  
260 265 270

70 Pro Glu Val Gln Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn  
275 280 285

EP 2 242 772 B1

Ala Lys Thr Lys Pro Arg Glu Glu Gln Phe Asn Ser Thr Tyr Arg Val  
290 295 300

5 Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu  
305 310 315 320

10 Tyr Lys Cys Lys Val Ser Asn Lys Gly Leu Pro Ser Ser Ile Glu Lys  
325 330 335

15 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr  
340 345 350

Leu Pro Pro Ser Gln Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr  
355 360 365

20 Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu  
370 375 380

25 Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu  
385 390 395 400

30 Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Arg Leu Thr Val Asp Lys  
405 410 415

Ser Arg Trp Gln Glu Gly Asn Val Phe Ser Cys Ser Val Met His Glu  
420 425 430

35 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Leu Gly  
435 440 445

40 <210> 2

<211> 214

<212> PRT

<213> Artificial Sequence

45 <220>

<223> Amino Acid sequence (predicted) of light chain of chimeric human/mouse antibody

50 <220>

<221> CDR1

<222> (24) .. (34)

55 <220>

<221> CDR2

<222> (50)..(56)

55 <220>

<221> CDR3

<222> (89)..(97)

<220>  
 <221> region IgG4  
 <222> (108)..(214)

5 <400> 2

	Asp	Ile	Gln	Met	Thr	Gln	Ser	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu	Gly
1						5					10					15
10																
	Asp	Arg	Val	Thr	Ile	Ser	Cys	Ser	Ala	Ser	Gln	Gly	Ile	Asn	Asn	Tyr
											25					30
15																
	Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gly	Thr	Val	Glu	Leu	Leu	Ile
											40					45
20																
	Tyr	Tyr	Thr	Ser	Thr	Leu	Gln	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
											55					60
25																
	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu	Pro
											75					80
30																
	Glu	Asp	Ile	Gly	Thr	Tyr	Tyr	Cys	Gln	Gln	Tyr	Ser	Lys	Leu	Pro	Arg
											90					95
35																
	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	Thr	Val	Ala	Ala
											105					110
40																
	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys	Ser	Gly
											120					125
45																
	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg	Glu	Ala
											135					140
50																
	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn	Ser	Gln
											150					160
55																
	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser	Leu	Ser
											165					175
60																
	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys	Val	Tyr
											180					190
65																
	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr	Lys	Ser
											195					205
70																
	Phe	Asn	Arg	Gly	Glu	Cys										
											210					

5                   <210> 3  
                  <211> 31  
                  <212> DNA  
                  <213> Artificial Sequence  
10                  <220>  
                  <223> IgH primer MHV7  
                  <400> 3  
                  atgggcatca agatggagtc acagacccag g 31  
15                  <210> 4  
                  <211> 21  
                  <212> DNA  
                  <213> Artificial Sequence  
                  <220>  
                  <223> IgG1 constant region primer MHCG1  
20                  <400> 4  
                  cagtggatag acagatgggg g 21  
                  <210> 5  
                  <211> 30  
25                  <212> DNA  
                  <213> Artificial Sequence  
                  <220>  
                  <223> Ig kappa primer MKV2  
30                  <400> 5  
                  atggagacag acacactcct gctatgggtg 30  
                  <210> 6  
35                  <211> 33  
                  <212> DNA  
                  <213> Artificial Sequence  
                  <220>  
40                  <223> Ig kappa primer MKV4  
                  <400> 6  
                  atgaggggccc ctgctcagtt ttttggcttc ttg 33  
45                  <210> 7  
                  <211> 25  
                  <212> DNA  
                  <213> Artificial Sequence  
50                  <220>  
                  <223> Ig kappa primer MKV9  
                  <400> 7  
                  atggtatcca cacctcagtt ccttg 25  
55                  <210> 8  
                  <211> 20  
                  <212> DNA

<213> Artificial Sequence

<220>

5      <223> primer MKC

<400> 8  
actggatggt ggaaagatgg        20

10     <210> 9  
<211> 50  
<212> DNA  
<213> Artificial Sequence

15     <220>  
<223> forward (For) primer

<400> 9  
agagaagctt gccgccacca tgattgcctc tgctcagttc ctgggtctcc        50

20     <210> 10  
<211> 35  
<212> DNA  
<213> Artificial Sequence

25     <220>  
<223> oligonucleotide primer BT03

<400> 10  
caacagtata gtaagctccc tcggacgttc ggtgg 35

30     <210> 11  
<211> 35  
<212> DNA  
<213> Artificial Sequence

35     <220>  
<223> oligonucleotide primer BT04

<400> 11  
40     ccaccgaacg tccgagggag cttactatac tgg 35

<210> 12  
<211> 41  
<212> DNA  
45     <213> Artificial Sequence

<220>  
<223> primer g2258

50     <400> 12  
cgcgggatcc actcacgtt gatccacgtc ttgggtgcctc c        41

<210> 13  
<211> 45  
55     <212> DNA  
<213> Artificial Sequence

<220>

<223> Primer g22949

<400> 13  
cgatggcccttggggaggctgaggagacggtgactgaggttcc 45

5

### Claims

1. An immunoconjugate capable of targeting CD138 expressing cells comprising:

10 (a) an engineered targeting antibody against CD138 comprising:

(i) an immunoglobulin heavy chain comprising a variable region comprising amino acid residues 31 to 35 (CDR1), 51 to 68 (CDR2) and 99 to 111 (CDR3) of SEQ ID NO: 1, and  
15 (ii) a light chain comprising a variable region comprising amino acid residues 24 to 34 (CDR1), 50 to 56 (CDR2) and 89 to 97 (CDR3) of SEQ ID NO: 2; and

(b) an effector molecule for inducing cell death,

20 wherein a constant region of said immunoglobulin heavy chain is an IgG4 isotype constant region.

2. The immunoconjugate of claim 1, wherein said effector molecule is attached to said engineered targeting antibody via a linker, and preferably said linker is a cleavable linker, most preferably comprising a disulfide bond.

25 3. The immunoconjugate according to claim 1, wherein:

- said immunoconjugate has a  $K_D$  value of less than 2.6 nM.

30 4. The immunoconjugate according to any preceding claim wherein the effector molecule is at least one maytansinoid, taxane or a CC1065, or an analog thereof, preferably the effector molecule is at least one maytansinoid, and most preferably the at least one maytansinoid is DM1, DM3, or DM4.

5. The immunoconjugate of any preceding claim, wherein said engineered targeting antibody further comprises:

35 (a) amino acid residues 123 to 448 of SEQ ID NO: 1, and/or  
(b) amino acid residues 108 to 214 of SEQ ID NO: 2, respectively and mutations thereof that

40 (i) maintain or lower the antibody-dependent cytotoxicity and/or complement-dependent cytotoxicity of the engineered targeting antibody and/or  
(ii) stabilize the engineered targeting antibody.

6. The immunoconjugate of any preceding claim, wherein the light chain sequence has at least 70% sequence identity with SEQ ID NO:2.

45 7. The immunoconjugate of claim 6, wherein the engineered targeting antibody comprises an immunoglobulin heavy chain comprising SEQ ID NO: 1 and a light chain comprising SEQ ID NO: 2.

8. An immunoconjugate according to any one of claims 1 to 7 for use in treating multiple myeloma in a subject.

50 9. An immunoconjugate according to any one of claims 1 to 7 for use in treating transitional cell bladder carcinoma in a subject.

10. An immunoconjugate according to any one of claims 1 to 7 for use as a medicament.

55 11. An immunoconjugate according to any one of claims 1 to 7 for use in inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient.

12. The immunoconjugate for use according to claim 11, wherein:

- said patient suffers from a hematologic malignancy and/or a solid tumor comprising CD138 expressing cells, preferably wherein said patient suffers from one of the following: multiple myeloma, ovarian carcinoma, kidney carcinoma, gall bladder carcinoma, breast carcinoma, prostate cancer, lung cancer, colon carcinoma, Hodgkin's and non-Hodgkin's lymphoma, chronic lymphocytic leukemia (CLL), acute lymphoblastic leukemia (ALL), acute myeloblastic leukemia (AML), solid tissue sarcoma or colon carcinoma, and most preferably wherein the patient suffers from multiple myeloma;

- said effector molecule of said immunoconjugate is a toxin, cytotoxic enzyme, low molecular weight cytotoxic drug, a pore-forming agent, biological response modifier, prodrug activating enzyme, an antibody, cytokine or a radionuclide;

- said immunoconjugate is to be administered in a single dose of 5 mg/m<sup>2</sup> to about 300 mg/m<sup>2</sup>; or

- said immunoconjugate is to be administered in at least two doses of about 5 mg/m<sup>2</sup> to about 300 mg/m<sup>2</sup>, optionally at hourly, daily, weekly intervals or combinations thereof.

13. An immunoconjugate according to any one of claims 1 to 7, for inhibiting, delaying and/or preventing the growth of a tumor and/or spread of malignant tumor cells comprising CD138 expressing cells in a patient, wherein the patient has been treated with one or more cytotoxic agents and/or radiation in an amount to reduce tumor load.

14. The immunoconjugate for use according to claim 13 wherein the cytotoxic agent is mephalan, vincristine, doxorubicin, dexamethasone, cyclophosphamide, etoposide, cytarabine, cisplatin, thalidomide, prednisone, bortezomib, lenalidomide, sorafenib, romidepsin or combinations thereof, or is antibody based.

15. An immunoconjugate according to any one of claims 1 to 7 for use in treating a subject having a condition that would benefit from the suppression of myeloma cell survival, wherein the immunoconjugate is capable of selectively decreasing survival or growth of said myeloma cells of said subject.

16. A pharmaceutical composition comprising the immunoconjugate of any one of claims 1 to 7 and one or more pharmaceutically acceptable excipients, wherein the composition is suitable for the inhibition, delay and/or prevention of the growth of tumors and/or spread of tumor cells.

17. A kit comprising, in separate containers, pharmaceutical compositions for use in combination to inhibit, delay and/or prevent the growth of tumors and/or spread of tumor cells, wherein one container comprises an effective amount of the pharmaceutical composition of claim 16, and wherein, a separate container comprises a second pharmaceutical composition comprising an effective amount of a further agent, preferably a cytotoxic agent, for the inhibition, delay and/or prevention of the growth of tumors and/or spread of tumor cells, and one or more pharmaceutically acceptable excipients, preferably wherein said agent in said second pharmaceutical composition is selected from the group consisting of mephalan, vincristine, doxorubicin, dexamethasone, cyclophosphamide, etoposide, cytarabine, cisplatin, thalidomide, prednisone, bortezomib, lenalidomide sorafenib, romidepsin and combinations thereof or is antibody based.

18. Use of an immunoconjugate according to any one of claims 1 to 7 for the manufacture of a medicament for inhibiting, delaying and/or preventing the growth of a tumor comprising CD138 tumor cells and/or spread of tumor cells of such a tumor in a patient.

#### 45 Patentansprüche

1. Ein Immunkonjugat, das in der Lage ist, auf CD138-exprimierende Zellen abzuzielen, umfassend:

(a) einen gegen CD138 hergestellten Ziel- Antikörper, umfassend:

(i) eine schwere Immunglobulinkette, umfassend einen variablen Abschnitt, umfassend Aminosäurereste 31 bis 35 (CDR1), 51 bis 68 (CDR2) und 99 bis 111 (CDR3) von SEQ ID-Nr.: 1, und

(ii) eine leichte Kette, umfassend eine variablen Abschnitt, umfassend Aminosäurereste 24 bis 34 (CDR1), 50 bis 56 (CDR2) und 89 bis 97 (CDR3) von SEQ ID-Nr.: 2; und

(b) ein Effektmolekül zur Auslösung von Zelltod,

wobei ein konstanter Abschnitt der besagten schweren Immunglobulinkette ein IgG4-isotyper konstanter Abschnitt

ist.

5 2. Das Immunkonjugat nach Anspruch 1, wobei das besagte Effektormolekül mit dem hergestellten Ziel-Antikörper über einen Linker verbunden ist, und wobei solch ein Linker vorzugsweise ein spaltbarer Linker ist, am meisten bevorzugt mit einer Disulfidbindung.

10 3. Das Immunkonjugat nach Anspruch 1, wobei:

- das besagte Immunkonjugat einen  $K_D$ -Wert von weniger als 2,6 nM hat.

15 4. Das Immunkonjugat nach einem der vorangehenden Ansprüche, wobei das Effektormolekül zumindest ein Maytansinoid, Taxan oder ein CC1065 bzw. ein Analog davon ist, und wobei das Effektormolekül vorzugsweise mindestens ein Maytansinoid ist, und wobei am meisten bevorzugt zumindest ein Maytansinoid DM1, DM3, oder DM4 ist.

20 5. Das Immunkonjugat nach einem der vorstehenden Ansprüche, wobei der besagte hergestellte Ziel-Antikörper des Weiteren besteht aus:

(a) Aminosäureresten 123 bis 448 von SEQ ID-Nr.: 1, und/oder

(b) Aminosäureresten 108 bis 214 von SEQ ID-Nr.: 2 und Mutationen davon, welche

25 (i) die antikörperabhängige Zytotoxizität und/oder eine Komplement-abhängige Zytotoxizität des hergestellten Ziel-Antikörpers aufrechterhalten oder senken, und/oder

(ii) den hergestellten Ziel-Antikörper stabilisieren.

30 6. Das Immunkonjugat nach einem der vorstehenden Ansprüche, wobei die leichte Kettensequenz eine Sequenzidentität von mindestens 70 % mit der SEQ ID-Nr.: 2 hat.

35 7. Das Immunkonjugat nach Anspruch 6, wobei der hergestellte Ziel-Antikörper eine schwere Immunglobulinkette umfassend SEQ ID-Nr.: 1 und eine leichte Kette, umfassend SEQ ID-Nr.: 2, umfasst.

40 8. Ein Immunkonjugat nach einem der Ansprüche 1 bis 7 für die Verwendung in der Therapie von multiplen Myelom bei einem Probanden.

45 9. Ein Immunkonjugat nach einem der Ansprüche 1 bis 7 für die Verwendung in der Therapie eines Übergangszellkarzinoms der Blase bei einem Probanden.

10. Ein Immunkonjugat nach einem der Ansprüche 1 bis 7 für die Verwendung als ein Medikament.

50 11. Ein Immunkonjugat nach einem der Ansprüche 1 bis 7 für die Verwendung zur Hemmung, Verzögerung und/oder Prävention des Wachstums von Tumoren, die CD-138 Tumorzellen umfassen, und/oder der Ausbreitung von Tumorzellen eines solchen Tumors in einem Patienten.

12. Das Immunkonjugat zur Verwendung nach Anspruch 11, wobei:

55 - der besagte Patient an einer hämatologischen malignen Erkrankung und/oder einem soliden Tumor mit CD138-exprimierenden Zellen leidet, wobei der besagte Patient vorzugsweise an einer der folgenden Krankheiten leidet:

Multiples Myelom, Ovarialkarzinom, Nierenkarzinom, Gallenblasenkarzinom, Mammakarzinom, Prostatakrebs, Lungenkrebs, Kolonkarzinom, Morbus Hodgkin und Non-Hodgkin-Lymphom, chronische lymphatische Leukämie (CLL), akute lymphoblastische Leukämie (ALL), akute myeloische Leukämie (AML), Sarkom in festem Gewebe oder Kolonkarzinom, und wobei der Patient am meisten bevorzugt an einem multiplen Myelom leidet.

55 - das besagte Effektormolekül des besagten Immunkonjugats ein Toxin, zytotoxisches Enzym, niedermolekulares zytotoxisches Arzneimittel, porenbildendes Mittel, Biomodulator, Prodrugaktivierendes Enzym, Antikörper, Zytokin oder ein Radionuklid ist;

- das besagte Immunkonjugat in einer Einzeldosis von 5 mg/m<sup>2</sup> bis ca. 300 mg/m<sup>2</sup> verabreicht wird; oder

- das besagte Immunkonjugat in mindestens zwei Dosen von ca. 5 mg/m<sup>2</sup> bis ca. 300 mg/m<sup>2</sup>, optional ständiglich,

täglich, wöchentlich oder in Kombinationen davon, zu verabreichen ist.

5           13. Ein Immunkonjugat nach einem der Ansprüche 1 bis 7 zur Hemmung, Verzögerung und/oder Prävention des Wachstums eines Tumors und/oder der Ausbreitung von malignen Tumorzellen, die CD138-exprimierende Tumorzellen umfassen, in einem Patienten, wobei der Patient mit einem oder mehreren zytotoxischen Mitteln und/oder durch Bestrahlung mit einem Betrag zur Verringerung der Tumorlast behandelt wurde.

10          14. Das Immunkonjugat zur Verwendung nach Anspruch 13, wobei das zytotoxische Mittel Mephalan, Vincristine, Doxorubicin, Dexamethason, Cyclophosphamid, Etoposid, Cytarabin, Cisplatin, Thalidomid, Prednison, Bortezomib, Lenalidomid, Sorafenib, Romidepsin oder Kombinationen davon ist oder auf Antikörpern basiert.

15          15. Ein Immunkonjugat nach einem der Ansprüche 1 bis 7 für die Verwendung in der Behandlung eines Probanden der an einem Zustand leidet, der durch die Unterdrückung des Überlebens von Myelomzellen therapiert werden könnte, wobei das Immunkonjugat in der Lage ist, das Überleben oder Wachstum der besagten Myelomzellen des besagten Probanden gezielt einzuschränken.

20          16. Eine pharmazeutische Zusammensetzung, umfassend das Immunkonjugat von einem der Ansprüche 1 bis 7 und ein oder mehrere pharmazeutisch akzeptablen Bestandteile, wobei die Zusammensetzung für die Hemmung, Verzögerung und/oder Prävention des Wachstums von Tumoren und/oder der Ausbreitung von Tumorzellen geeignet ist.

25          17. Ein Set umfassend, in separaten Behältern, pharmazeutische Zusammensetzungen für die kombinierte Verwendung zur Hemmung, Verzögerung und/oder Prävention des Wachstums von Tumoren und/oder der Ausbreitung von Tumorzellen, wobei ein Behälter eine wirksame Menge der pharmazeutischen Zusammensetzung nach Anspruch 16 umfasst, und wobei ein separater Behälter eine zweite pharmazeutische Zusammensetzung umfasst, die eine wirksame Menge eines weiteren Agens umfasst, vorzugsweise eines zytotoxischen Wirkstoffes für die Hemmung, Verzögerung und/oder Prävention des Wachstums von Tumoren und/oder der Ausbreitung von Tumorzellen, sowie einen oder mehrere pharmazeutisch akzeptable Hilfsstoffe, wobei das besagte Agens in der besagten zweiten pharmazeutischen Zusammensetzung vorzugsweise aus der Gruppe stammt, die aus Mephalan, Vincristine, Doxorubicin, Dexamethason, Cyclophosphamid, Etoposid, Cytarabin, Cisplatin, Thalidomid, Prednison, Bortezomib, Lenalidomid, Sorafenib, Romidepsin oder Kombinationen davon besteht oder auf Antikörpern basiert.

30          18. Die Verwendung eines Immunkonjugats nach einem der Ansprüche 1 bis 7 für die Herstellung eines Medikaments zur Hemmung, Verzögerung und/oder Prävention des Wachstums von Tumoren, die CD-138 Tumorzellen umfassen, und/oder der Ausbreitung von Tumorzellen eines solchen Tumors in einem Patienten.

#### Revendications

40          1. Un immunoconjugué capable de cibler des cellules exprimant CD138 comprenant :

                 (a) un anticorps de ciblage produit par génie génétique contre CD138 comprenant :

                    (i) une chaîne lourde d'immunoglobuline comprenant une région variable comprenant des résidus d'acides aminés 31 à 35 (CDR1), 51 à 68 (CDR2) et 99 à 111 (CDR3) de SEQ ID N° : 1, et

                    (ii) une chaîne légère comprenant une région variable comprenant des résidus d'acides aminés 24 à 34 (CDR1), 50 à 56 (CDR2) et 89 à 97 (CDR3) de SEQ ID N° : 2, et

                    (b) une molécule effectrice pour induire la mort cellulaire,

50          dans lequel une région constante de ladite chaîne lourde d'immunoglobuline est une région constante d'isotype IgG4.

                 2. L'immunoconjugué de la revendication 1, dans lequel ladite molécule effectrice est attachée au dit anticorps de ciblage produit par génie génétique via un segment de liaison, et de préférence ledit segment de liaison est un segment de liaison pouvant être coupé, de manière très préférentielle comprenant un pont disulfure.

                 3. L'immunoconjugué selon la revendication 1, dans lequel :

- ledit immunoconjugué a une valeur  $K_D$  inférieure à 2,6 nM.

4. L'immunoconjugué selon l'une quelconque des revendications précédentes dans lequel la molécule effectrice est au moins un maytansinoïde, un taxane ou un CC1065, ou un analogue de ceux-ci, de préférence la molécule effectrice est au moins un maytansinoïde, et de manière très préférentielle le au moins un maytansinoïde est DM1, DM3, ou DM4.

5 5. L'immunoconjugué selon l'une quelconque des revendications précédentes, dans lequel ledit anticorps de ciblage produit par génie génétique comprend en outre :

10 (a) des résidus d'acides aminés 123 à 448 de SEQ ID N° : 1, et/ou  
 (b) des résidus d'acides aminés 108 à 214 de SEQ ID N° : 2, respectivement et des mutations de ceux-ci qui  
 15 (i) maintiennent ou diminuent la cytotoxicité dépendante des anticorps et/ou complémentent la cytotoxicité dépendante de l'anticorps de ciblage produit par génie génétique et/ou,  
 (ii) stabilisent l'anticorps de ciblage produit par génie génétique.

6. L'immunoconjugué selon l'une quelconque des revendications précédentes, dans lequel la séquence de chaîne légère a au moins 70 % d'identité de séquence avec SEQ ID N° : 2.

20 7. L'immunoconjugué de la revendication 6, dans lequel l'anticorps de ciblage produit par génie génétique comprend une chaîne lourde d'immunoglobuline comprenant SEQ ID N° : 1 et une chaîne légère comprenant SEQ ID N° : 2.

25 8. Un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour une utilisation dans le traitement du myélome multiple dans un sujet.

9. Un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour une utilisation dans le traitement du carcinome de cellules de type transitionnel de la vessie dans un sujet.

30 10. Un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour une utilisation comme un médicament.

11. Un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour une utilisation pour inhiber, retarder et/ou empêcher la croissance d'une tumeur comprenant des cellules tumorales CD138 et/ou la prolifération de cellules tumorales d'une telle tumeur dans un patient.

35 12. L'immunoconjugué pour une utilisation selon la revendication 11, dans lequel :

40 - ledit patient souffre d'affections hématologiques malignes et/ou d'une tumeur solide comprenant des cellules exprimant CD138, de préférence dans lequel ledit patient souffre d'une des affections suivantes : myélome multiple, carcinome de l'ovaire, carcinome du rein, carcinome de la vésicule biliaire, carcinome du sein, carcinome de la prostate, carcinome du poumon, carcinome du côlon, lymphome hodgkinien et non hodgkinien, leucémie lymphoïde chronique (LLC), leucémie lymphoblastique aigüe (LLA), leucémie myéloblastique aigüe (LMA), sarcome de tissu solide ou carcinome du côlon, et de manière très préférentielle dans lequel le patient souffre de myélome multiple ;

45 - ladite molécule effectrice dudit immunoconjugué est une toxine, un enzyme cytotoxique, un médicament cytotoxique à faible poids moléculaire, un agent formant des pores, un modificateur de la réponse biologique, un enzyme activant un précurseur de médicament, un anticorps, une cytokine ou un radionucléide ;

- ledit immunoconjugué doit être administré en une seule dose allant de 5 mg/m<sup>2</sup> à environ 300 mg/m<sup>2</sup> ; ou

50 - ledit immunoconjugué doit être administré en au moins deux doses allant de 5 mg/m<sup>2</sup> à environ 300 mg/m<sup>2</sup>, optionnellement à des intervalles d'une heure, d'un jour, d'une semaine ou de combinaisons de ceux-ci.

13. Un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour inhiber, retarder et/ou empêcher la croissance d'une tumeur et/ou la prolifération de cellules tumorales malignes comprenant des cellules exprimant CD138 dans un patient, dans lequel le patient a été traité avec un ou plusieurs agents cytotoxique et/ou par radiothérapie dans un montant pour réduire la charge tumorale.

55 14. L'immunoconjugué pour une utilisation selon la revendication 13, dans lequel l'agent cytotoxique est du melphalan, de la vincristine, de la doxorubicine, de la dexaméthasone, de la cyclophosphamide, de l'étoposide, de la cytarabine,

du cisplatine, de la thalidomide, de la prednisone, du bortézomib, de la lénalidomide, du sorafénib, de la romidepsine ou de combinaisons de ceux-ci, ou est à base d'anticorps.

- 5        15. Un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour une utilisation dans le traitement d'un sujet ayant une condition qui bénéficierait de la suppression de la survie de la cellule myélome, dans lequel l'immunoconjugué est capable de diminuer sélectivement la survie ou la croissance desdites cellules myélomes dudit sujet.
- 10        16. Une composition pharmaceutique comprenant l'immunoconjugué de l'une quelconque des revendications de 1 à 7 et un ou plusieurs excipients pharmaceutiquement acceptables, dans laquelle la composition est appropriée pour l'inhibition, le retard et/ou la prévention de la croissance de tumeurs et/ou la prolifération de cellules tumorales.
- 15        17. Un kit comprenant, dans des conteneurs séparés, des compositions pharmaceutiques pour une utilisation en combinaison pour inhiber, retarder et/ou empêcher la croissance de tumeurs et/ou la prolifération de cellules tumorales, dans lequel un conteneur comprend une quantité efficace de la composition pharmaceutique de la revendication 16, et dans lequel, un conteneur séparé comprend une seconde composition pharmaceutique comprenant une quantité efficace d'un autre agent, de préférence un agent cytotoxique, pour l'inhibition, le retard et/ou la prévention de la croissance de tumeurs et/ou la prolifération de cellules tumorales, et un ou plusieurs excipients pharmaceutiquement acceptables,
- 20        de préférence dans lequel ledit agent dans ladite seconde composition pharmaceutique est sélectionné du groupe consistant en du melphalan, de la vincristine, de la doxorubicine, de la dexaméthasone, de la cyclophosphamide, de l'étoposide, de la cytarabine, du cisplatine, de la thalidomide, de la prednisone, du bortézomib, de la lénalidomide, du sorafénib, de la romidepsine ou de combinaisons de ceux-ci, ou est à base d'anticorps.
- 25        18. Une utilisation d'un immunoconjugué selon l'une quelconque des revendications de 1 à 7 pour la fabrication d'un médicament pour inhiber, retarder et/ou empêcher la croissance d'une tumeur comprenant des cellules tumorales CD138 et/ou la prolifération de cellules tumorales d'une telle tumeur dans un patient.

30

35

40

45

50

55

FIG. 1





FIG. 2





FIG. 3



DCC: 1,3-dicyclohexylcarbodiimide  
 DTT: Dithiothreitol  
 DME: 1,2-Dimethoxyethane




FIG. 5




FIG. 6




FIG. 7A




FIG. 7B




FIG. 7C




FIG. 7D




FIG. 8A

FIG. 8B






FIG. 8C

FIG. 8D






FIG. 10




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13

## REFERENCES CITED IN THE DESCRIPTION

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

## Patent documents cited in the description

- US 20070183971 A [0003]
- US 6005079 A [0076]
- JP 20020006379 B [0076]
- EP 0239400 A [0078]
- WO 9109967 A [0078]
- US 5530101 A [0078]
- US 5585089 A [0078]
- EP 0592106 A [0078]
- EP 0519596 A [0078]
- US 5565332 A [0078]
- US 5639641 A [0079]
- US 4444887 A [0079]
- US 4716111 A [0079]
- US 5545806 A [0079]
- US 5814318 A [0079]
- WO 9846645 A [0079]
- WO 9850433 A [0079]
- WO 9824893 A [0079]
- WO 9816654 A [0079]
- WO 9634096 A [0079]
- WO 9633735 A [0079]
- WO 9110741 A [0079]
- US 5831012 A [0084]
- US 6534628 B [0084]
- US 6740734 B [0084]
- US 5475092 A [0087] [0096]
- US 5585499 A [0087] [0096]
- US 6716821 B [0087] [0101]
- US 3896111 A [0088]
- US 4418064 A [0089]
- US 4151042 A [0090] [0091]
- US 4137230 A [0091]
- US 4248870 A [0091]
- US 4256746 A [0091]
- US 4260608 A [0091]
- US 4265814 A [0091]
- US 4294757 A [0091]
- US 4307016 A [0091]
- US 4308268 A [0091]
- US 4308269 A [0091]
- US 4309428 A [0091]
- US 4313946 A [0091]
- US 4315929 A [0091]
- US 4317821 A [0091]
- US 4322348 A [0091]
- US 4331598 A [0091]
- US 4361650 A [0091]
- US 4362663 A [0091]
- US 4364866 A [0091]
- US 4371533 A [0091]
- US 4424219 A [0091]
- US 6333410 B, Chari [0092]
- US 20040235840 A [0094] [0095]
- US 20060233814 A, Goldmahker [0095] [0130]
- US 4169888 A [0096]
- US 5739350 A [0096]
- US 5846545 A [0096]
- US 6756397 B [0096]
- US 6534660 B [0096]
- US 6436931 B [0097]
- US 6340701 B [0097]
- US 6706708 B [0097]
- US 20040087649 A [0097]
- US 20040024049 A [0097]
- US 20030004210 A [0097]
- US 6002023 A [0097]
- US 5998656 A [0097]
- US 5892063 A [0097]
- US 5763477 A [0097]
- US 5705508 A [0097]
- US 5703247 A [0097]
- US 5367086 A [0097]
- US 6596757 B [0097]
- US 5053394 A [0098]
- US 4563304 A [0099]
- US 5208020 A [0100]
- US 20030055226 A, Chari [0100]
- US 5877296 A [0102]
- US 5773001 A [0102]
- US 5712374 A [0102]
- US 5714586 A [0102]
- US 20040082764 A [0102]
- US 20080171040 A, Eberts [0131]
- US 61016620 B [0219]
- US 61087466 B [0219]
- US 61087590 B [0219]

## Non-patent literature cited in the description

- Computational Molecular Biology. Oxford University Press, 1988 [0109]
- Biocomputing: Informatics and Genome Projects. Academic Press, 1993 [0109]

- Computer Analysis of Sequence Data. Humana Press, 1994 [0109]
- VON HEINJE, G. Sequence Analysis in Molecular Biology. Academic Press, 1987 [0109]
- Sequence Analysis Primer. M Stockton Press, 1991 [0109]
- CARILLO, H. ; LIPTON, D. SIAM J Applied Math. 1988, vol. 48, 1073 [0109]
- SMITH ; WATERMAN. *Advances in Applied Mathematics*, 1981, vol. 2, 482-489 [0111]
- Remington's Pharmaceutical Sciences. Mack Publishing Co, 1985 [0134]
- WIJDENES et al. *Br J Haematol.*, 1996, vol. 94, 318 [0152]
- J. SAMBROOK. Molecular Cloning, A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0153]
- CARROLL et al. *Mol Immunol.*, 1988, vol. 25, 991 [0163]
- CABILLY et al. *Gene*, 1985, vol. 40, 157 [0163]
- SUN et al. *PNAS*, 1987, vol. 84, 214 [0169]
- CHARI et al. *Cancer Res.*, 1992, vol. 1, 127 [0181]
- ROGUSKA et al. *PNAS*, 1994, vol. 91, 969 [0181]
- LIU et al. *PNAS*, 1996, vol. 93, 8618 [0181]
- REMILLARD et al. *Science*, 1977, vol. 189, 1002 [0182]
- KUPCHAN et al. *J. Med. Chem.*, 1978, vol. 21, 31 [0183]
- TOMAYKO et al. *Cancer Chemother. Pharmacol*, 1989, vol. 24, 148 [0199]
- BISSERY et al. *Cancer Res.*, 1991, vol. 51, 4845 [0199]
- AKKINA RK ; ROSENBLATT JD ; CAMPBELL AG ; CHENIS ; ZACK JA. Modeling human lymphoid precursor cell gene therapy in the SCID-hu mouse. *Blood*, 1994, vol. 84, 1393-1398 [0218]
- ARMOUR KL ; CLARK MR ; HADLEY AG et al. Recombinant human IgG molecules lacking Fc gamma receptor I binding and monocyte triggering activities. *Eur J Immunol.*, 1999, vol. 29 (8), 2613-24 [0218]
- ANDERSON KC ; KYLE RA ; DALTON WS ; LANDOWSKI T ; SHAIN K ; JOVE R ; HAZLEHURST L ; BERENSON J. Multiple Myeloma: New Insights and Therapeutic Approaches. *Hematology*, 2000, 147-165 [0218]
- ANTTONEN A ; HEIKKILA P ; KAJANTI M ; JALKANEN M ; JOENSUU H. High syndecan-1 expression is associated with favourable outcome in squamous cell lung carcinoma treated with radical surgery. *Lung Cancer*, June 2001, vol. 32 (3), 297-305 [0218]
- BARBARESCHI M ; MAISONNEUVE P ; ALDOVINI D ; CANGI MG ; PECCARINI L ; ANGELO MAURI F ; VERONESE S ; CAFFO O ; LUCENTI A ; PALMA PD. High syndecan-1 expression in breast carcinoma is related to an aggressive phenotype and to poorer prognosis. *Cancer*, 01 August 2003, vol. 98 (3), 474-83 [0218]
- BATAILLE R ; JÉGO G ; ROBILLARD N ; BARILLÉ-NION S ; HAROUSSEAU JL ; MOREAU P ; AMIOT M ; PELLAT-DECEUNYNCK C. The phenotype of normal, reactive and malignant plasma cells. Identification of "many and multiple myelomas" and of new targets for myeloma therapy. *Haematologica*, September 2006, vol. 91 (9), 1234-40 [0218]
- BERNFIELD M ; KOKENYESI R ; KATO M ; HINKES MT ; SPRING J ; GALLO RL ; LOSE EJ. Biology of the syndecans: a family of transmembrane heparan sulfate proteoglycans. *Annu Rev Cell Biol.*, 1992, vol. 8, 365-393 [0218]
- BESTE G ; SCHMIDT FS ; STIBORA T ; SKERRA A. Small antibody-like proteins with prescribed ligand specificities derived from the lipocalin fold. *Proc. Natl. Acad. Sci. USA.*, 1999, vol. 96, 1898-1903 [0218]
- BHATTACHARYYA B ; WOLFF J. Maytansine binding to the vinblastine sites of tubulin. *FEBS Lett.*, 1977, vol. 75, 159-162 [0218]
- BISPING G ; KROPFF M ; WENNING D ; DREYER B ; BESSONOV S ; HILBERG F ; ROTH GJ ; MUNZERT G ; STEFANIC M ; STELLJES M. Targeting receptor kinases by a novel indolinone derivative in multiple myeloma: abrogation of stroma-derived interleukin-6 secretion and induction of apoptosis in cytogenetically defined subgroups. *Blood*, 08 November 2005, vol. 107 (5), 2079-89 [0218]
- Drugs to Enhance the Therapeutic Potency of Anti-cancer Antibodies: Antibody-Drug Conjugates as Tumor-Activated Prodrugs. BLÄTTLER WA ; CHARI RVJ. Anticancer Agents-Frontiers in Cancer Chemotherapy. American Chemical Society, 2001, 317-338 [0218]
- BROSS PF ; BEITZ J ; CHEN G ; CHEN XH ; DUFFY E ; KIEFFER L ; ROY S ; SRIDHARA R ; RAHMAN A ; WILLIAMS G. Approval summary: gemtuzumab ozogamicin in relapsed acute myeloid leukemia. *Clin Cancer Res.*, 2001, vol. 7, 1490-1496 [0218]
- CARBONE A ; GAIDANO G ; GLOGHINI A ; FERLITO A ; RINALDO A ; STEIN H. AIDS-related plasma-blastic lymphomas of the oral cavity and jaws: a diagnostic dilemma. *Ann. Otol. Rhinol. Laryngol.*, 1999, vol. 108, 95-99 [0218]
- CARLSSON J ; DREVIN H ; AXEN R. Protein thiolation and reversible protein-protein conjugation. N-succinimidyl-3-(2-pyridyldithio)propionate, a new heterobifunctional reagent. *Biochem J*, 1978, vol. 173, 723-737 [0218]
- CARTER P. Improving the efficacy of antibody-based cancer therapies. *Nat Rev Cancer*, 2001, vol. 1, 118-129 [0218]
- CHARI RV ; MARTELL BA ; GROSS JL ; COOK SB ; SHAH SA ; BLATTNER WA ; MCKENZIE SJ ; GOLDMACHER VS. Immunoconjugates containing novel maytansinoids: promising anticancer drugs. *Cancer Res.*, 1992, vol. 52, 127-131 [0218]

- CHARI RV ; JACKEL KA ; BOURRET LA ; DERR SM ; TADAYONI BM ; MATTOCKS KM ; SHAH SA ; LIU C ; BLÄTTLER WA ; GOLDMACHER VS. Enhancement of the selectivity and antitumor efficacy of a CC-1065 analogue through immunoconjugate formation. *Cancer Res.*, 1995, vol. 55, 4079-4084 [0218]
- CHARNAUX N ; BRULE S ; CHAIGNEAU T ; SAF-FAR L ; SUTTON A ; HAMON M ; PROST C ; LIEVRE N ; VITA C ; GATTEGNO L. RANTES (CCL5) induces a CCR5-dependent accelerated shedding of syndecan-1 (CD138) and syndecan-4 from HeLa cells and forms complexes with the shed ectodomains of these proteoglycans as well as with those of CD44. *Glycobiology*, 08 September 2004 [0218]
- CHEN BP ; GALY A ; KYOIZUMI S ; NAMIKAWA R ; SCARBOROUGH J ; WEBB S ; FORD B ; CEN DZ ; CHEN SC. Engraftment of human hematopoietic precursor cells with secondary transfer potential in SCID-hu mice. *Blood*, 1994, vol. 84, 2497-2505 [0218]
- CHILOSI M ; ADAMI F ; LESTANI M ; MONTAGNA L ; CIMAROSTO L ; SEMENZATO G ; PIZZOLO G ; MENESTRINA F. CD138/syndecan-1: a useful immunohistochemical marker of normal and neoplastic plasma cells on routine trephine bone marrow biopsies. *Mod Pathol.*, 1999, vol. 12, 1101-1106 [0218]
- CLEMENT C ; VOOIJS, W.C. ; KLEIN, B. ; WIJDENES, J. *J. Leukocyte Typing V*. Oxford University Press, 1995, 714-715 [0218]
- COUTURIER O ; FAIVRE-CHAUVET A ; FILIP-POVICH IV ; THEDRÉZ P ; SAÏ-MAUREL C ; BARDIÉS M ; MISHRA AK ; GAUVRIT M ; BLAIN G ; APOSTOLIDIS C. Validation of 213Bi-alpha radioimmunotherapy for multiple myeloma. *Clinical Cancer Research*, October 1999, vol. 5 (10), 3165s-3170s [0218]
- DAVIES EJ ; BLACKHALL FH ; SHANKS JH ; DAVID G ; MCGOWN AT ; SWINDELL R ; SLADE RJ ; MARTIN-HIRSCH P ; GALLAGHER JT ; JAY-SON GC et al. Distribution and Clinical Significance of Heparan Sulfate Proteoglycans in Ovarian Cancer *Clin. Cancer Res.*, 2004, vol. 10 (15), 5178-86 [0218]
- DHODAPKAR MV ; ABE E ; THEUS A ; LACY M ; LANGFORD JK ; BARLOGIE B ; SANDERSON RD. Syndecan-1 is a multifunctional regulator of myeloma pathobiology: control of tumor cell survival, growth, and bone cell differentiation. *Blood*, 1998, vol. 91, 2679-2688 [0218]
- DORE JM ; MORARD F ; VITA N ; WIJDENES J. Identification and location on syndecan-1 core protein of the epitopes of B-B2 and B-B4 monoclonal antibodies. *FEBS Lett.*, 1998, vol. 426, 67-70 [0218]
- DOWELL JA ; KORTH-BRADLEY J ; LIU H ; KING SP ; BERGER MS. Pharmacokinetics of gemtuzumab ozogamicin, an antibody-targeted chemotherapy agent for the treatment of patients with acute myeloid leukemia in first relapse. *J Clin Pharmacol.*, 2001, vol. 41, 1206-1214 [0218]
- EDINGER M ; SWEENEY TJ ; TUCKER AA ; OLOMU AB ; NEGRIN RS ; CONTAG CH. Noninvasive assessment of tumor cell proliferation in animal models. *Neoplasia*, 1999, vol. 1, 303-310 [0218]
- GATTEI V ; GODEAS C ; DEGAN M ; ROSSI FM ; ALDINUCCI D ; PINTO A. Characterization of Anti-CD138 monoclonal antibodies as tools for investigating the molecular polymorphism of syndecan-1 in human lymphoma cells. *Br J Haematol.*, 1999, vol. 104, 152-162 [0218]
- HAMANN PR ; HINMAN LM ; BEYER CF ; LINDH D ; UPESLACIS J ; FLOWERS DA ; BERNSTEIN I. An anti-CD33 antibody-calicheamicin conjugate for treatment of acute myeloid leukemia. Choice of linker. *Bioconjug Chem.*, 2002, vol. 13, 40-46 [0218]
- HAN I ; PARK H ; OH ES. New insights into syndecan-2 expression and tumourigenic activity in colon carcinoma cells. *J Mol Histol.*, 2004, vol. 35 (3), 319-26 [0218]
- HIDESHIMA T ; CATLEY L ; YASUI H ; ISHITSUKA K ; RAJE N ; MITSIADES C ; PODAR K ; MUNSHI NC ; CHAUHAN D ; RICHARDSON PG. Perifosine, an oral bioactive novel alkylphospholipid, inhibits Akt and induces in vitro and in vivo cytotoxicity in human multiple myeloma cells. *Blood*, 2006, vol. 107 (10), 4053-62 [0218]
- HIDESHIMA T ; MITSIADES C ; TONON G ; RICHARDSON PG ; ANDERSON KC. Understanding multiple myeloma pathogenesis in the bone marrow to identify new therapeutic targets. *Nat Rev Cancer*, 2007, vol. 7 (8), 585-98 [0218]
- HORVATHOVA M ; GAILLARD, J.-P. ; LIUTARD, J. ; DUPERRAY, C. ; LAVABRE-BERTRAND, T. ; BOURQUARD, P et al. *Leucocyte Typing V*. Oxford University Press, 1995, 713-714 [0218]
- KOVTUN YV ; AUDETTE CA ; YE Y ; XIE H ; RUBERTI MF ; PHINNEY SJ et al. Antibody-drug conjugates designed to eradicate tumors with homogeneous and heterogeneous expression of the target antigen. *Cancer Res*, 2006, vol. 66 (6), 3214-21 [0218]
- KREBS B ; RAUCHENBERGER R ; REIFFERT S ; ROTHE C ; TESAR M ; THOMASSEN E ; CAO M ; DREIER T ; FISCHER D ; HOSS A et al. High-throughput generation and engineering of recombinant human antibodies. *J. Immunol. Methods*, 2001, vol. 254, 67-84 [0218]

- **KUPCHAN SM ; SNEDEN AT ; BRANFMAN AR ; HOWIE GA ; REBHUN LI ; MCLVOR WE ; WANG RW ; SCHNAITMAN TC.** Structural requirements for antileukemic activity among the naturally occurring and semisynthetic maytansinoids. *J Med Chem.*, 1978, vol. 21, 31-37 [0218]
- **KYOIZUMI S ; BAUM CM ; KANESHIMA H ; MCCUNE JM ; YEE EJ ; NAMIKAWA R.** Implantation and maintenance of functional human bone marrow in SCID-hu mice. *Blood*, 1992, vol. 79, 1704-1711 [0218]
- **KYOIZUMI S ; MURRAY LJ ; NAMIKAWA R.** Pre-clinical analysis of cytokine therapy in the SCID-hu mouse. *Blood*, 1993, vol. 81, 1479-1488 [0218]
- **LANGFORD JK ; STANLEY MJ ; CAO D ; SANDERSON RD.** Multiple heparan sulfate chains are required for optimal syndecan-1 function. *J Biol Chem.*, 06 November 1998, vol. 273 (45), 29965-71 [0218]
- **LIU C ; TADAYONI BM ; BOURRET LA ; MATTOCKS KM ; DERR SM ; WIDDISON WC ; KEDERSHA NL ; ARINIELLO PD ; GOLDMACHER VS ; LAMBERT JM.** Eradication of large colon tumor xenografts by targeted delivery of maytansinoids. *Proc Natl Acad Sci USA.*, 1996, vol. 93, 8618-8623 [0218]
- **MCCUNE JM ; NAMIKAWA R ; KANESHIMA H ; SHULTZ LD ; LIEBERMAN M ; WEISSMAN IL.** The SCID-hu mouse: murine model for the analysis of human hematolymphoid differentiation and function. *Science*, 1988, vol. 241, 1632-1639 [0218]
- **MENNERICH D ; VOGEL A ; Klaman I ; DAHL E ; LICHTNER RB ; ROSENTHAL A ; POHLENZ HD ; THIERAUCH KH ; SOMMER A.** Shift of syndecan-1 expression from epithelial to stromal cells during progression of solid tumours. *Eur J Cancer*, June 2004, vol. 40 (9), 1373-82 [0218]
- **MOSMANN T.** Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *J Immunol Methods*, 1983, vol. 65, 55-63 [0218]
- **Plasma cell disorders. MUNSHI NC ; LONGO DL ; ANDERSON KC.** Harrison's Principles of Internal Medicine. McGraw-Hill Medical Publishing Division, 2008, 700-7 [0218]
- **NAMIKAWA R ; UEDA R ; KYOIZUMI S.** Growth of human myeloid leukemias in the human marrow environment of SCID-hu mice. *Blood*, 1993, vol. 82, 2526-2536 [0218]
- **O'CONNELL FP ; PINKUS JL ; PINKUS GS.** CD138 (Syndecan-1), a Plasma Cell Marker Immunohistochemical Profile in Hematopoietic and Nonhematopoietic Neoplasms. *Am J Clin Pathol*, 2004, vol. 121, 254-263 [0218]
- **OJIMA I ; GENG X ; WU X ; QU C ; BORELLA CP ; XIE H ; WILHELM SD ; LEECE BA ; BARTLE LM ; GOLDMACHER VS.** Tumor-specific novel taxoid-monoclonal antibody conjugates. *J. Med. Chem.*, 2002, vol. 45, 5620-5623 [0218]
- **OLAFSEN T ; CHEUNG CC ; YAZAKI PJ ; LI L ; SUNDARESAN G ; GAMBHIR SS ; SHERMAN, MA ; WILLIAMS, LE ; SHIVELY, JE ; RAUBITSCHEK, AA.** Covalent disulfide-linked anti-CEA diabody allows site-specific conjugation and radiolabeling for tumor targeting applications. *Prot. Eng. Design & Selection*, 2004, vol. 17 (1), 21-27 [0218]
- **OROSZ Z ; KOPPER L.** Syndecan-1 expression in different soft tissue tumours. *Anticancer Res.*, 2001, vol. 21 (1B), 733-7 [0218]
- **PADLAN EA.** A possible procedure for reducing the immunogenicity of antibody variable domains while preserving their ligand-binding properties. *Mol. Immunol.*, 1991, vol. 28, 489-498 [0218]
- **PAYNE G.** Progress in immunoconjugate cancer therapeutics. *Cancer Cell*, 2003, vol. 3, 207-212 [0218]
- **PEGRAM MD ; LIPTON A ; HAYES DF ; WEBER BL ; BASELGA JM ; TRIPATHY D ; BALY D ; BAUGHMAN SA ; TWADDELL T ; GLASPY JA.** Phase II study of receptor-enhanced chemosensitivity using recombinant humanized anti-p185HER2/neu monoclonal antibody plus cisplatin in patients with HER2/neu-overexpressing metastatic breast cancer refractory to chemotherapy treatment. *J. Clin. Oncol.*, 1998, vol. 16, 2659-2671 [0218]
- **RAWSTRON AC ; OWEN RG ; DAVIES FE ; JOHN-SON RJ ; JONES RA ; RICHARDS SJ ; EVANS PA ; CHILD JA ; SMITH GM ; JACK AS.** Circulating plasma cells in multiple myeloma: characterization and correlation with disease stage. *Br J Haematol.*, 1997, vol. 97, 46-55 [0218]
- **REMILLARD S ; REBHUN LI ; HOWIE GA ; KUPCHAN SM.** Antimitotic activity of the potent tumor inhibitor maytansine. *Science*, 1975, vol. 189, 1002-1005 [0218]
- **ROGUSKA MA ; PEDERSEN JT ; KEDDY CA ; HENRY AH ; SEARLE SJ ; LAMBERT JM ; GOLDMACHER VS ; BLATTNER WA ; REES AR ; GUILD BC.** Humanization of murine monoclonal antibodies through variable domain resurfacing. *Proc Natl Acad Sci USA.*, 1994, vol. 91, 969-973 [0218]
- **ROSS S ; SPENCER SD ; HOLCOMB I ; TAN C ; HONGO J ; DEVAUX B ; RANGELL L ; KELLER GA ; SCHOW P ; STEEVES RM.** Prostate stem cell antigen as therapy target: tissue expression and in vivo efficacy of an immunoconjugate. *Cancer Res.*, 01 May 2002, vol. 62 (9), 2546-53 [0218]
- **ROSS JS ; GRAY K ; GRAY G ; WORLAND PJ ; ROLFE M.** Anticancer Antibodies. *Am J Clin Path.*, 17 April 2003 [0218]
- **SANDERSON RD ; LALOR P ; BERNFIELD M.** B lymphocytes express and lose syndecan at specific stages of differentiation. *Cell Regul.*, 1989, vol. 1, 27-35 [0218]

- **SANDHU JS ; CLARK BR ; BOYNTON EL ; ATKINS H ; MESSNER H ; KEATING A ; HOZUMI N.** Human hematopoiesis in SCID mice implanted with human adult cancellous bone. *Blood*, 1996, vol. 88, 1973-1982 [0218]
- **SASAKIA ; BOYCE BF ; STORY B ; WRIGHT KR ; CHAPMAN M ; BOYCE R ; MUNDY GR ; YONEDA T.** Bisphosphonate risedronate reduces metastatic human breast cancer burden in bone in nude mice. *Cancer Res.*, 1995, vol. 55, 3551-3557 [0218]
- **SCHNEIDER U ; VAN LESSEN A ; HUHN D ; SERKE S.** Two subsets of peripheral blood plasma cells defined by differential expression of CD45 antigen. *Br J Haematol.*, 1997, vol. 97, 56-64 [0218]
- **SCHUURMAN J ; VAN REE R ; G. J. PERDOK GJ ; VAN DOORN HR ; TAN KY ; AALBERSE RC.** Normal human immunoglobulin G4 is bispecific: it has two different antigen-combining sites. *Immunology*, 1999, vol. 97, 693-698 [0218]
- **SEBESTYEN A ; BERCZI L ; MIHALIK R ; PAKU S ; MATOLCSY A ; KOPPER L.** Syndecan-1 (CD138) expression in human non-Hodgkin lymphomas. *Br J Haematol.*, 1999, vol. 104 (2), 412-9 [0218]
- **SEFTALIOGLU A ; KARAKUS S.** Syndecan-1/CD138 expression in normal myeloid, acute lymphoblastic and myeloblastic leukemia cells. *Acta Histochem.*, 2003, vol. 105, 213-221 [0218]
- **SEFTALIOGLU A ; KARAKUS S ; DUNDAR S ; CAN B ; ERDEM LIE ; IRMAK MK ; OZTAS E ; KORKMAZ C ; YAZAR F ; CAVUSOGLU I.** Syndecan-1 (CD138) expression in acute myeloblastic leukemia cells--an immuno electron microscopic study. *Acta Oncol.*, 2003, vol. 42, 71-74 [0218]
- Cures and regressions of established tumors with monoclonal antibody auristatin conjugates. **SENTER PD ; DORONINA S ; CERVENY C ; CHACE D ; FRANCISCO J ; KLUSSMAN K ; MENDELSON B ; MEYER D ; SIEGALL CB ; THOMPSON J et al.** American Association for Cancer Res. American Association for Cancer Res, 2002, 414 [0218]
- **SHIELDS RL ; NAMENUK AK ; HONG K ; MENG YG ; RAE J ; BRIGGS J ; XIE D ; LAI J ; STADLEN A ; LI B.** High resolution mapping of the binding site on human IgG1 for Fc gamma RI, Fc gamma RII, Fc gamma RIII, and FcRn and design of IgG1 variants with improved binding to the Fc gamma R. *J Biol Chem.*, 2001, vol. 276 (9), 6591-604 [0218]
- **SIEVERS EL ; LARSON R.A. ; STADTMAUER, E.A. ; ESTEY, E. ; LOWENBERG, B. ; DOMBRET, H. ; KARANES, C. ; THEOBALD, M. ; BENNETT, J.M. ; SHERMAN, M.L. et al.** Efficacy and safety of gemtuzumab ozogamicin in patients with CD33-positive acute myeloid leukemia in first relapse. *J. Clin. Oncol.*, 2001, vol. 19, 3244-3254 [0218]
- **SIEVERS EL ; LINENBERGER M.** Mylotarg: antibody-targeted chemotherapy comes of age. *Curr. Opin. Oncol.*, May 2001, vol. 13, 522-527 [0218]
- **SMITH R.** Single chain antibody variable region fragments, May 2001, [www.stanford.edu/~smithr/science/scfv.html](http://www.stanford.edu/~smithr/science/scfv.html) [0218]
- **STUDNICKA GM ; SOARES S ; BETTER M ; WILLIAMS RE ; NADELL R ; HORWITZ AH.** Human-engineered monoclonal antibodies retain full specific binding activity by preserving non-CDR complementarity-modulating residues. *Protein Eng.*, 1994, vol. 7 (6), 805-814 [0218]
- **TAI YT ; LI XF ; CATLEY L ; COFFEY R ; BREITKREUTZ I ; BAE J ; SONG W ; PODAR K ; HIDESHIMA T ; CHAUHAN D.** Immunomodulatory drug lenalidomide (CC-5013, IMiD3) augments anti-CD40 SGN-40-induced cytotoxicity in human multiple myeloma: clinical implications. *Cancer Res.*, 15 December 2005, vol. 65 (24), 11712-20 [0218]
- **TASSONE P ; GOLDMACHER VS ; NERI P ; GOZZINIA ; SHAMMAS MA ; WHITEMAN KA ; HYLANDER-GANS LL ; CARRASCO DR ; HIDESHIMA T ; SHRINGARPURE R.** Cytotoxic activity of the maytansinoid immunoconjugate B-B4-DM1 against CD138+ multiple myeloma cells. *Blood*, 2004, vol. 104 (12), 3688-3696 [0218]
- **TOLCHER AW ; OCHOA L ; HAMMOND LA ; PATAIK A ; EDWARDS T ; TAKIMOTO C ; SMITH L ; DE BONO J ; SCHWARTZ G ; MAYS T.** Cantuzumab mertansine, a maytansinoid immunoconjugate directed to the CanAg antigen: a phase I, pharmacokinetic, and biologic correlative study. *J Clin Oncol.*, 2003, vol. 21, 211-222 [0218]
- **URASHIMA M ; CHEN BP ; CHEN S ; PINKUS GS ; BRONSON RT ; DEDERA DA ; HOSHI Y ; TEOH G ; OGATA A ; TREON SP.** The development of a model for the homing of multiple myeloma cells to human bone marrow. *Blood*, 1997, vol. 90, 754-765 [0218]
- **VOGEL CW.** Preparation of immunoconjugates using antibody oligosaccharide moieties. *Methods in Molecular Biology: Bioconjugation protocols strategies and methods*, 2004, vol. 283, 087-108 [0218]
- **VOOIJS WC ; POST J ; WIJDENES J ; SCHUURMAN HJ ; BOLOGNESI A ; POLITO L ; STRIPE F ; BAST EJ ; DE GAST GC.** Efficacy and toxicity of plasma-cell-reactive monoclonal antibodies B-B2 and B-B4 and their immunotoxins. *Cancer Immunol Immunother*, 1996, vol. 42, 319-328 [0218]
- **WARD, E.S. ; D. GUSSOW ; A.D. GRIFFITHS ; P.T. JONES ; G. WINTER.** Binding activities of a repertoire of single immunoglobulin variable domains secreted from *Escherichia coli*. *Nature*, 1989, vol. 341, 544-546 [0218]
- **WARGALLA UC ; REISFELD RA.** Rate of internalization of an immunotoxin correlates with cytotoxic activity against human tumor cells. *Proc. Natl. Acad. Sci. USA.*, 1989, vol. 86, 5146-5150 [0218]

- WIJDENES J ; VOOIJS WC ; CLEMENT C ; POST J ; MORARD F ; VITA N ; LAURENT P ; SUN RX ; KLEIN B ; DORE JM. A plasmocyte selective monoclonal antibody (B-B4) recognizes syndecan-1. *Br J Haematol.*, 1996, vol. 94, 318-323 [0218]
- WIJDENES J ; DORE JM ; CLEMENT C ; VERMOT-DESROCHES C. CD138. *J Biol Regul Homeost Agents*, April 2002, vol. 16 (2), 152-5 [0218]
- WITZIG TE ; KIMLINGER TK ; AHMANN GJ ; KATZMANN JA ; GREIPP PR. Detection of myeloma cells in the peripheral blood by flow cytometry. *Cytometry*, 1996, vol. 26, 113-120 [0218]
- XIE H ; AUDETTE C ; HOFFEE M ; LAMBERT JM ; BLÄTTLER W. Pharmacokinetics and biodistribution of the antitumor immunoconjugate, cantuzumab mertansine (huC242-DM1), and its two components in mice. *J Pharmacol Exp Ther.*, March 2004, vol. 308 (3), 1073-82 [0218]
- YANG M ; JIANG P ; AN Z ; BARANOV E ; LI L ; HASEGAWA S ; AL-TUWAIJRI M ; CHISHIMA T ; SHIMADA H ; MOOSA AR. Genetically fluorescent melanoma bone and organ metastasis models. *Clin Cancer Res.*, 1999, vol. 5, 3549-3559 [0218]
- YANG M ; BARANOV E ; JIANG P ; SUN FX ; LI XM ; LI L ; HASEGAWA S ; BOUVET M ; AL-TUWAIJRI M ; CHISHIMA T. Whole-body optical imaging of green fluorescent protein-expressing tumors and metastases. *Proc Natl Acad Sci USA.*, 2000, vol. 97, 1206-1211 [0218]
- YANG Y ; MACLEOD V ; DAI Y ; KHOT-SKAYA-SAMPLE Y ; SHRIVER Z ; VENKATARAMAN G ; SASISEKHARAN R ; NAGGI A ; TORRI G ; CASU B. The syndecan-1 heparan sulfate proteoglycan is a viable target for myeloma therapy. *Blood*, 29 May 2007, vol. 110 (6), 2041-8 [0218]
- YOSHITAKE S ; YAMADA Y ; ISHIKAWA E ; MASSEYEFF R. Conjugation of glucose oxidase from Aspergillus niger and rabbit antibodies using N-hydroxysuccinimide ester of N-(4-carboxycyclohexylmethyl)-maleimide. *Eur J Biochem*, 1979, vol. 101, 395-399 [0218]

## (CD138-at megcélzó immunkonjugátumok és alkalmazásai)

## Szabadalmi igénypontok

1. CD138-at kifejező sejtek megcélzására képes immunkonjugátum, amely magában foglal:

- (a) egy CD138 elleni manipulált célzó antitestet, amely magában foglal:
  - (i) egy, a SEQ ID NO: 1 31 – 35. (CDR1), 51 – 68. (CDR2) és 99 – 111. (CDR3) aminosavmaradékait magában foglaló immunglobulin nehéz láncot, és
  - (ii) egy, a SEQ ID NO: 2 24 – 34. (CDR1), 50 – 56. (CDR2) és 89 – 97. (CDR3) aminosavmaradékait magában foglaló variabilis régiót tartalmazó könnyű láncot, és
- (b) egy effektor molekulát sejthalál kivállására,

ahol az említett immunglobulin nehéz lánc egy IgG4 izotípusú konstans régió.

2. Az 1. igénypont szerinti immunkonjugátum, ahol az illető effektor molekula az illető manipulált célzó antitesthez egy linkerrel át csatlakozik, és az illető linker előnyösen egy hasítható linker, legelőnyösebben egy díszuifidkötést magában foglaló.

3. Az 1. igénypont szerinti immunkonjugátum, ahol:

- az illető immunkonjugátum 2,6 nM-nál kisebb  $K_D$  értékű.

4. Az előző igénypontok bármelyike szerinti immunkonjugátum, ahol az effektor molekula legalább egy maitanzinoid, és legelőnyösebben a legalább egy maitanzinoid DM1, DM3 vagy DM4.

5. Az előző igénypontok bármelyike szerinti immunkonjugátum, ahol az illető manipulált célzó antitest továbbá magában foglalja:

- (a) a SEQ ID NO: 1 123 – 448. aminosavmaradékait, és/vagy
- (b) a SEQ ID NO: 2 108 – 214. aminosavmaradékait és ezek mutációit, amelyek
  - (i) fenntarják vagy csökkentik az antitest-függő citotoxicitást és/vagy a manipulált célzó antitest komplement-függő citotoxicitását, és/vagy
  - (ii) stabilizálják a manipulált célzó antitestet.

6. Az előző igénypontok bármelyike szerinti immunkonjugátum, ahol a könnyű lánc szekvencia a SEQ ID NO: 2-vel legalább 70% szekvencia azonossággal rendelkezik.

7. A 6. igénypont szerinti immunkonjugátum, ahol a manipulált célzó antitest egy SEQ ID NO: 1-et magában foglaló immunglobulin nehéz láncot és egy SEQ ID NO: 2-t magában foglaló könnyű láncot foglal magában.

8. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum egy alanyban myeloma multiplex kezelésében való alkalmazásra.

9. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum egy alanyban húgyhólyag tranzisionális sejtes rákja kezelésében való alkalmazásra.

10. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum gyógyszerként való alkalmazásra.

11. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum egy paciensben egy CD138 tumorsejteket magában foglaló tumor növekedésének és/vagy egy ilyen tumor tumorsejjei szétterjedésének gátlására, késleltetésére és/vagy megelőzésére való alkalmazásra.

12. Immunkonjugátum a 11. igénypont szerinti alkalmazásra, ahol:

- az illető paciens rosszindulatú hematológiai betegségben és/vagy CD138-at kifejező sejteket magában foglaló szolid tumorról szenved, előnyösen ahol az illető paciens a következő betegségek egyikében szenved: myeloma multiplex, petefészekrák, veserák, epehólyagrák, mellrák, prosztatarák, tüdőrák, vastagbélrák, Hodgkin- és nem-Hodgkin-limfóma, krónikus limfocitás leukémia (CLL), akut limfoblasztikus leukémia (ALL), szolid szövetes szarkóma vagy vastagbél karcinóma, és legelőnyösebben ahol a paciens myeloma multiplexben szenved,
- az illető immunkonjugátum effektor molekulája egy toxin, citotoxikus enzim, kis molekulatömegű citotoxikus hatóanyag, egy pórus-képző szer, biológiai válasz módosító, prodrug aktiváló enzim, egy antitest, citokin vagy egy radionuklid,
- az illető immunkonjugátum  $5 \text{ mg/m}^2$ -től körülbelül  $300 \text{ mg/m}^2$ -ig terjedő egyszeri dózisban adandó be, vagy
- az illető immunkonjugátum körülbelül  $5 \text{ mg/m}^2$ -től körülbelül  $300 \text{ mg/m}^2$ -ig terjedő legalább két dózisban, adott esetben óránként, naponta, heti intervallumokban vagy ezek kombinációiban adandó be.

13. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum egy paciensben egy tumor növekedésének és/vagy CD138-at kifejező sejteket magukban foglaló malignus tumorsejtek szétterjedésének gátlására, késleltetésére és/vagy megelőzésére, ahol a pacienst már egy vagy több citotoxikus szemel és/vagy a tumorterhelést csökkentő mértékű besugárzással kezelték.

14. Immunkonjugátum a 13. igénypont szerinti alkalmazásra, ahol a citotoxikus szer mephalan, vínkrisztin, doxorubicin, dexametazon, ciklofoszfamid, etopozid, citarabin, ciszplatin, talidomid, prednizon, bortezomib, lenalidomid, szorafenib,

romidepszin vagy ezek kombinációja vagy antitest alapú.

15. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum egy, a myeloma sejt túlélés elnyomásától javulható betegségben szenvedő alany kezelésében való alkalmazásra, ahol az immunkonjugátum képes az illető alany említett myeloma sejtjei túlélésének vagy növekedésének szelektív csökkentésére.

16. Gyógyszerészeti kompozíció, amely magában foglal az 1 – 7. igénypontok bármelyike szerinti immunkonjugátumot és egy vagy több gyógyszerészeti leg elfogadható segédanyagot.

17. Készlet (kit), amely magában foglal, elkülönített tárolóedényekben, gyógyszerészeti kompozíciókat kombinációban tumorok növekedésének és/vagy tumorsejtek széttérjedésének gátlására, késleltetésére és/vagy megelőzésére való alkalmazásra, ahol egy tárolóedény a 16. igénypont szerinti gyógyszerészeti kompozíció hatásos mennyiségét foglalja magában, és ahol egy elkülönített tárolóedény egy további szernek, előnyösen egy citotoxikus szernek a tumorok növekedésének és/vagy tumorsejtek széttérjedésének gátlására, késleltetésére és/vagy megelőzésére hatásos mennyiségét és egy vagy több gyógyszerészeti segédanyagot magában foglaló második gyógyszerészeti kompozíciót foglal magában, ahol előnyösen az illető szer az illető második gyógyszerészeti kompozícióban a mephalan, vinkrisztin, doxorubicin, dexametazon, ciklofoszfamid, etopozid, citarabin, ciszplatin, talidomid, prednizon, bortezomib, lenalidomid, szorafenib, romidepszin vagy ezek kombinációja által alkotott csoportból kiválasztott vagy antitest alapú.

18. Egy 1 – 7. igénypontok bármelyike szerinti immunkonjugátum alkalmazása egy paciensben egy CD138 tumorsejteket magában foglaló tumor növekedésének és/vagy egy ilyen tumor tumorsejjei széttérjedésének gátlására, késleltetésére és/vagy megelőzésére szánt gyógyszer gyártására.

A meghatalmazott:



Ráthonyi Zoltán  
Kreatív Gyógyszergyártó  
szox Szabadalmi Ügyvivői Törzse  
H-1162 Budapest, Amassay út 113  
Telefon: +36 1 892 1000 / fax: +36 1 892 1099  
E-mail: [zoltan.rathonyi@szox.hu](mailto:zoltan.rathonyi@szox.hu)